

COMPOSITIONS FOR RNA-CHROMATIN INTERACTION ANALYSIS AND USES THEREOF

REFERENCE TO RELATED APPLICATION

This application claims priority to and the benefit of the filing date of U.S. Provisional Application No. 61/873,928, filed on September 5, 2013, the entire content of which is incorporated herein by reference.

BACKGROUND OF THE INVENTION

Noncoding RNAs (ncRNAs) are now believed to be transcribed pervasively in the genome, and large numbers of ncRNAs have been identified. However, disproportionately, still very little is known about their functional roles. Many of the known ncRNA functions were inferred by perturbation experiments, which lack the details of what specific target an ncRNA interact with. Technologies like CLIP/ RIP-Seq and ChiRP-Seq have provided tremendous insights of what the protein factors and chromatin loci for some ncRNAs to interact with. However, current methods are limited to examine ncRNA or interacting target one at a time. Thus it is desirable to have an unbiased genome-wide strategy to identify the functional targets for all ncRNAs.

SUMMARY OF THE INVENTION

One aspect of the invention provides a kit comprising: (1) an RNA linker comprising: (i) a first polynucleotide, and, (ii) a second polynucleotide, wherein the first and the second polynucleotides form a first double stranded region flanked by a first ligation compatible end, and a 3'-overhang at the 3'-end of the first polynucleotide, wherein the 3'-overhang comprises a random-sequence primer; and, (2) a DNA linker comprising: (iii) a third polynucleotide, and, (iv) a fourth polynucleotide, wherein the third and the fourth polynucleotides form a second double stranded region flanked by a blunt end and a second ligation compatible end, wherein the first and the second ligation compatible ends ligate to each other, or are adaptable to ligate to each other.

In certain embodiments, the first ligation compatible end is a 3'-overhang at the 3'-end of the second polynucleotide, and the second ligation compatible end is a 3'-overhang at the 3'-end of the third polynucleotide, wherein both 3'-overhangs anneal to each other for ligation.

In certain embodiments, the first double stranded region comprises a first recognition site for a first restriction enzyme (RE) that cleaves 3' to the random-sequence primer.

In certain embodiments, the second double stranded region comprises a second recognition site for a second restriction enzyme (RE) that cleaves 5' to the third polynucleotide.

In certain embodiments, one or more of said first, second, third, and fourth polynucleotides are DNA.

In certain embodiments, one or more of said first, second, third, and fourth polynucleotides comprise a modified nucleotide.

In certain embodiments, the modified nucleotide is a biotinylated T (Thymidine).

In certain embodiments, the first polynucleotide comprises a plurality of polynucleotides, each differing only at the random-sequence primer region.

In certain embodiments, the first polynucleotide comprises a homogeneous population of polynucleotides having identical random-sequence primer.

In certain embodiments, the random-sequence primer comprises 4, 5, 6, 7, 8, or more nucleotides.

In certain embodiments, the first double stranded region comprises a unique sequence that distinguishes the RNA linker from the DNA linker.

In certain embodiments, the second double stranded region comprises a unique sequence that distinguishes the RNA linker from the DNA linker.

In certain embodiments, the last nucleotide of the first recognition site is the last base-paired nucleotide 5' to the random-sequence primer.

In certain embodiments, the last nucleotide of the second recognition site is a base-paired nucleotide at the blunt end.

In certain embodiments, the first and the second restriction enzymes are the same.

In certain embodiments, the first or the second restriction enzyme is independently selected from: AarI, AceIII, AclI, BaeI, Bbr7I, BbvI, BbvII, BccI, Bce83I, BceAI, BceFI, BcgI, BciVI, BfiI, BinI, BplI, BsaXI, BscAI, BseMII, BseRI, BsgI, BsmI, BsmAI, BsmFI, Bsp24I, BspCNI, BspMI, BsrI, BsrDI, BstF5I, BtgZI, BtsI, CjeI, CjePI, EciI, Eco31I, Eco57I, Eco57MI, EcoP15I, Esp3I, Fall, FauI, FokI, GsuI, HaeIV, HgaI, Hin4I, HphI, HpyAV, Ksp632I, MboII, MlyI, MmeI, MnlI, PleI, PpiI, PsrI, RleAI, SapI, SfaNI, SspD5I, Sth132I, StsI, TaqII, TspDTI, TspGWI, TspRI or Tth111II.

In certain embodiments, the cleavage site of the first or the second restriction enzyme is at least about 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30 or more nucleotides 3' to the last nucleotide of the recognition site.

In certain embodiments, the first and the fourth polynucleotides are dephosphorylated.

In certain embodiments, the kit further comprises a reagent that cross-links protein and polynucleotide.

In certain embodiments, the reagent comprises formaldehyde.

In certain embodiments, the kit further comprises an affinity reagent (*e.g.*, an antibody, or a monoclonal antibody) that specifically or selectively binds a component of chromatin (*e.g.*, histone).

In certain embodiments, the kit further comprises an end-repairing mixture that converts DNA containing damaged or incompatible 5'- and/or 3'-protruding ends to 5'-phosphorylated, blunt-ended DNA.

In certain embodiments, the kit further comprises a DNA ligase (*e.g.*, T4 ligase).

In certain embodiments, the kit further comprises a reagent that reverses cross-linking of protein and polynucleotide (*e.g.*, Proteinase K).

In certain embodiments, the kit further comprises the first and/or the second restriction enzyme(s).

In certain embodiments, the kit further comprises a pair of concatenating adapters for PCR amplification of blunt-ended double stranded DNA.

In certain embodiments, the kit further comprises a Taq DNA polymerase.

In certain embodiments, the kit further comprises a reverse transcriptase.

Another aspect of the invention provides a paired-end tag (PET) polynucleotide comprising a central region comprising the first and second double stranded regions of the subject RNA and DNA linkers, said central region being flanked by: (1) at a site proximal to said first double stranded region, a sequence tag of a non-coding RNA (ncRNA); and (2) at a site proximal to said second double stranded region, a sequence tag of a genomic DNA.

In certain embodiments, the sequence tag of the non-coding RNA (ncRNA) has a free end resulting from digestion by said first restriction enzyme.

In certain embodiments, the sequence tag of the non-coding RNA (ncRNA) uniquely identifies a genomic region from which the ncRNA is transcribed.

In certain embodiments, the sequence tag of the non-coding RNA (ncRNA) is about 8-30 base pairs in length.

In certain embodiments, the sequence tag of the genomic DNA has a free end resulting from digestion by said second restriction enzyme.

In certain embodiments, the sequence tag of the genomic DNA uniquely identifies a genomic region at which the genomic DNA is located.

In certain embodiments, the sequence tag of the genomic DNA is about 8-30 base pairs in length.

Another aspect of the invention provides a paired-end tag (PET) library comprising two or more members of the subject PET polynucleotide, wherein each member of the PET library comprises the same said central region, and different said sequence tag of the subject non-coding RNA (ncRNA) or different said sequence tag of the subject genomic DNA or both.

Another aspect of the invention provides a vector comprising a subject PET polynucleotide.

In certain embodiments, the vector comprises a plurality of concatenated subject PET polynucleotide.

Another aspect of the invention provides a concatemer of two or more subject PET polynucleotides.

Another aspect of the invention provides a method of identifying functional interaction loci within a genome for non-coding RNAs (ncRNAs) of the genome, the method comprising: (1) providing chromatin fragments comprising cross-linked genomic DNA fragments and cross-linked ncRNAs; (2) using the RNA linker and the DNA linker of claim 1, ligating an end of a cross-linked genomic DNA fragment to an end of a cDNA of a cross-linked ncRNA, under a condition for proximity ligation, wherein said end of the cross-linked genomic DNA fragment is ligated to the DNA linker, and said end of the cDNA of the cross-linked ncRNA comprises the RNA linker; (3) isolating a PET polynucleotide of claim 29 for sequencing analysis; and, (4) mapping the sequence tag of the genomic DNA and the sequence tag of the ncRNA within each said PET polynucleotide to a reference genome, thereby identifying functional interaction loci within the reference genome for said non-coding RNAs (ncRNAs) of the reference genome.

In certain embodiments, the ncRNAs and the genomic DNA are cross-linked in live cells through formaldehyde-mediated cross-linking.

In certain embodiments, chromatin fragments are generated by sonication.

In certain embodiments, the cDNA of the cross-linked ncRNA comprises a first strand cDNA reverse transcribed from the random-sequence primer of the RNA linker, and the ncRNA template.

In certain embodiments, 2nd strand cDNA synthesis is carried out after proximity ligation but before step (3).

In certain embodiments, the method further comprises repairing the ends of the cross-linked genomic DNA fragments to 5'-phosphorylated, blunt-ended DNA prior to step (2).

In certain embodiments, the third polynucleotide of the DNA linker is dephosphorylated and the DNA linker does not self-ligate.

In certain embodiments, the method further comprises identifying clusters of two or more PET polynucleotides having overlapping sequence tags of the genomic DNA and overlapping sequence tags of the ncRNA.

In certain embodiments, the method further comprises excluding PET polynucleotides comprising sequence tags of rRNA.

In certain embodiments, the method further comprises isolating or enriching a subset of chromatin fragments prior to step (2).

In certain embodiments, the subset of chromatin fragments is isolated or enriched by immunoprecipitation using an antibody specific for a protein component of the subset of chromatin fragments.

In certain embodiments, the protein component is a histone, a transcription factor, a polycomb-group (PcG) family protein; a recombination involved factor; a chromatin insulator or chromatin waver; a methyl-CpG-binding protein; or an RNA binding protein.

It should be understood that any description disclosed for the purpose of carrying out one embodiment of this invention (such as embodiments only described in the example section only), including but not limited to any technique(s), reagents, experimental conditions, restrictions sites, enzymes, vectors, primers, and the like, may also be used in combination with other embodiments of the invention, including those embodiments described only in detail in one (but not any other) aspect of the invention. It will be evident

to any skilled person how to adapt techniques and material disclosed for the other embodiments to the present embodiment of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1A shows a schematic flow of a typical setting of the RICH-PET method using the RNA linker and DNA linker pair. The interactions of ncRNA to chromatin are captured by crosslinking, followed by sonication to break up the chromatin fibers into tethering complexes with RNA, DNA and protein components. The tethered RNA and DNA in each of the chromatin fragment complex are then connected by a set of ligation reactions mediated by specifically designed RNA linker and DNA linker oligonucleotides that may also have unique sequence barcodes for orientation and specificity. Within each of the chromatin complexes, the 3'-end of RNA is annealed to the random hexamer protruding portion of the RNA linker followed by reverse transcription for cDNA synthesis. Meanwhile, the DNA linker is added to the blunt ends of the tethered DNA fragments by ligation. After wash of the excess linker oligos, the attached RNA and DNA linkers are ligated to each other, thus linking the tethered RNA and DNA molecules. After reverse cross-linking, the hybrid ligation products are fragmented either by shearing or restriction digestion into desired sizes for further amplification, sequencing, and mapping analysis to identify the locations where the RNA was transcribed and where it interacted in the genome.

FIG. 1B shows a schematic flow of a typical setting of the RICH-PET method using the modified RNA linker.

FIG. 1C shows a schematic flow of a typical setting of the RICH-PET method using the direct RNA linker. "App" stands for the 5' adenylation at the 5'-end of the first polynucleotide.

FIGs. 2A-2C present statistics of selected RICH-PET libraries, and sequencing and mapping data. FIG. 2A shows proportions of singleton PET (no overlap with other PET sequences) and PET clusters. Using the PET cluster data, approximately 700 RNA loci and about 5000 DNA loci were identified. FIG. 2B shows RNA-Seq data intensity at the RNA and DNA loci identified by the RICH-PET data. FIG. 2C shows that most of the RICH-PET data defined ncRNA interactions were trans-acting and inter-chromosomal.

FIG. 3 demonstrates reproducibility and sensitivity of the subject method. The figure shows representative scatter plots showing the comparison of RNA interacting sites identified in technical and biological replicates. Known lncRNAs MALAT1 (PET count 174) and

NEAT1 (PET 18) were repeatedly detected in RICH-PET data (not shown). RNAPII ChIA-PET data also shows that these two lncRNAs are also connected spatially within the same RNAPII transcriptional complex possibly for co-regulation. In addition, RNA-Seq and RNA-PET data were used to assess the expression level of ncRNA genes in HeLa S3 (data not shown). Both data showed that MALAT1 was highly expressed, NEAT1 was expressed at middle level, and HOTAIR was expressed at very low level. RICH-PET mapping at the HOTAIR locus shows poor RICH-PET data in this region (data not shown).

FIGs. 4A-4B show data for validation of NEAT1 and MALAT1 RICH-PET data. FIG. 4A shows that both NEAT1 and MALAT1 are expressed in HeLa S3 cells, and are abundantly detected in RICH-PET data. NEAT1 is restricted only in cis-acting, in that both RNA and DNA tags were mapped in short distances within the same locus. MALAT1 is mostly trans-acting, in that most of the DNA tags were mapped in large distances in the same chromosome or in different chromosomes (inset). FIG. 4B shows RNA-FISH experiments in human A549 and HeLa S3. The NEAT1 probe generated few fluorescent spots (1-2 per nucleus in HeLa S3 cells), whereas the MALAT1 probe generated much more spots (13 per nucleus in HeLa S3 cells). Counts were based on 100 nuclei per probe per experiment.

FIGs. 5A-5B characterize the RICH-PET data. FIG. 5A shows pie charts of categories of RNA tag cluster locations in the genome, showing that the vast majority of RNA tags were found in putative ncRNA regions, only 3% were overlap with protein-coding exons. Many known ncRNAs were detected, and many new ones were identified. FIG. 5B shows pie charts of categories of DNA tag cluster locations in the genome, showing that the majority of DNA tag clusters were mapped to protein coding regions, mostly in either promoters or introns.

FIGs. 6A-6B show multi-targets and multi-functions by MALAT1 interactions. FIG. 6A is a connectivity map of MALAT1 interacting with 59 genomic loci. FIG. 6B is a box plot showing genes with MALAT1 presence at their promoter regions have higher RNA-seq reads than the genes with MALAT1 interactions at their intron regions. In an aggregation plot of RNAPII ChIP-Seq intensity (not shown), genes with MALAT1 presence at their promoter regions have higher RNA-seq reads than the genes with MALAT1 interactions at their intron regions.

FIG. 7 shows a schematic drawing of CCAT1 and its lncRNA transcript acting as transcription activator or co-activator for several target genes.

FIG. 8A shows the location on the human chromosome 8 of the CCAT1 genomic and cDNA sequences corresponding to SEQ ID NO: 9. FIG. 8B shows the location on the human

chromosome 8 of the eight additional CCAT1 genomic and cDNA sequences corresponding to SEQ ID NOs: 1-8 (CCAT1_JAX_1 to _8, respectively). Filled boxes represent exon sequences, while lines linking the exon sequences represent intron sequences.

FIG. 9A shows the counts of XIST (which specifically targets the X chromosome in female cells) measured by RNA-Seq data, in reads per kb per million reads (RPKM).

FIG. 9B shows the proportion of each chromosome covered by XIST binding.

DETAILED DESCRIPTION OF THE INVENTION

1. Overview

The invention described herein is partly based on the realization that, if an ncRNA had an epigenetic regulatory role in the nuclear space, it would have to either directly or indirectly interact with chromatin at certain locations in chromosomes, in which functions take place for modulating chromatin states and target gene activity. Hence, the invention described herein provides a new approach to globally map ncRNA-chromatin interactions through RNA-DNA ligation, followed by paired-end-tag sequencing (RICH-PET).

In brief, compositions described herein can be used in a method comprising three main parts: 1) chromatin crosslinking to capture (preferably all) molecular interaction events between RNA, DNA and proteins in a live cell (such as one cultured *in vitro*, or a primary cell obtained from a tissue sample); 2) ligation of the tethered interactive RNA and the chromatin DNA fragment (*e.g.*, through specifically designed linker, such as the RNA linker and DNA linker pairs, or through ligating RNA 3' end to 5' adenylated ssDNA or 5' adenylated overhang); and, 3) sequencing and mapping analysis of the RNA-DNA ligation products or tag sequences derived therefrom (*e.g.*, PET polynucleotide) to localize ncRNAs' transcription sites and their chromatin target sites in the genome.

Thus one aspect of the invention provides a method of identifying functional interaction loci within a genome for non-coding RNAs (ncRNAs) of the genome, the method comprising: (1) providing a chromatin fragment comprising a cross-linked genomic DNA fragment and a cross-linked ncRNA (or a fragment thereof); (2) ligating an end of the cross-linked genomic DNA fragment to an end of the cross-linked ncRNA, under a condition for proximity ligation; (3) isolating a paired-end tag (PET) polynucleotide for sequencing analysis, wherein the PET polynucleotide comprises a sequence tag of a non-coding RNA (ncRNA), and a sequence tag of a genomic DNA; and, (4) mapping the sequence tag of the genomic DNA and the sequence tag of the ncRNA to a reference genome, thereby identifying

functional interaction loci within the reference genome for the non-coding RNAs (ncRNAs) of the reference genome.

This RNA-DNA ligation approach not only applies to global study of all ncRNA-chromatin interactions, but can also be applied to studying RNA-protein interaction at specific chromatin locations. Thus, a chromosomal immunoprecipitation (ChIP)-based RICH-PET method could provide additional specificity of RNA-protein-chromatin interaction information.

The reagents and methods of the invention have a wide range of potential uses in research, development, drug target identification, drug screening, diagnosis, treatment / efficacy monitoring, prognosis, *etc.* For example, the reagents and methods of the invention can be used to comprehensively characterize ncRNA-chromatin interactomes for a number of established cell lines, stem cells, iPS cells, and cells from primary tissues, such as those derived from cancer and healthy tissue control; and to significantly increase our capability of investigating the immense complex world of RNA functions in regulating the output of the genome. The successful completion of the characterization of RNA-chromatin interactomes would provide a comprehensive chromatin address book for most (if not all) of the ncRNA species, which would add another dimension of genomic information to help understand how the genome functions in healthy and disease conditions.

Several specific embodiments of the invention are described in more details below.

a) RNA Linker and DNA Linker Pairs

In the first specific embodiment, the method of the invention can be carried out using an RNA linker and DNA linker pair to ligate the crosslinked RNA and chromosomal DNA in the same chromatin fragment.

Thus one aspect of the invention provides a kit comprising: (1) an RNA linker comprising: (i) a first polynucleotide, and, (ii) a second polynucleotide, wherein the first and the second polynucleotides form a first double stranded region flanked by a first ligation compatible end, and a 3'-overhang at the 3'-end of the first polynucleotide, wherein the 3'-overhang comprises a random-sequence primer; and, (2) a DNA linker comprising: (iii) a third polynucleotide, and, (iv) a fourth polynucleotide, wherein the third and the fourth polynucleotides form a second double stranded region flanked by a blunt end and a second ligation compatible end, wherein the first and the second ligation compatible ends ligate to each other, or are adaptable to ligate to each other.

In certain embodiments, the first ligation compatible end is a 3'-overhang at the 3'-end of the second polynucleotide, and the second ligation compatible end is a 3'-overhang at the 3'-end of the third polynucleotide, wherein both 3'-overhangs anneal to each other for ligation.

In certain embodiments, the first ligation compatible end is a 5'-overhang at the 5'-end of the first polynucleotide, and the second ligation compatible end is a 5'-overhang at the 5'-end of the fourth polynucleotide, wherein both 5'-overhangs anneal to each other for ligation.

In certain embodiments, the first and/or the second ligation compatible ends are adaptable for ligation. For example, instead of having the requisite 3' or 5' overhangs for ligation, the first and/or the second ligation compatible ends may comprise a restriction enzyme (RE) site, which can be cleaved by the RE to produce the requisite 3' or 5' overhangs required for ligation. Prior to cleavage by the restriction enzyme, however, the ligation compatible ends may be blunt ended (*e.g.*, dephosphorylated blunt end to prevent self-ligation), or have non-compatible overhang that prevents self-ligation or ligation with the other ligation compatible end.

In certain embodiments, the two 5'- or 3'-overhangs at the compatible ligation ends do not self-anneal and do not anneal with each other. This can be accomplished, for example, by designing the sequences of the overhangs such that the overhang sequences do not self-anneal or anneal with each other, at least when under the conditions the linkers are to be used.

This design may be advantageous in certain embodiments, in which, for example, a downstream step includes PCR amplification. One frequently observed type of non-specific amplification product is a template-independent artifact of amplification reactions referred to as "primer dimer," which is a double-stranded fragment whose length typically is close to the sum of the two primer lengths and appears to occur when one primer is extended over the other primer. The resulting extension product forms an undesired template which, because of its short length, is amplified efficiently.

Each of the first, second, third, and fourth polynucleotides may be provided in separate containers, such as synthesized polynucleotides, either in freeze dried, lyophilized form or in water or a suitable buffer solution. Alternatively, the first and the second polynucleotides may be combined in the same container (lyophilized or in solution), for example, in 1:1 molar ratio, such that they can be used as pre-annealed RNA linker. Similarly, the third and the fourth polynucleotides may be combined in the same container

(lyophilized or in solution), for example, in 1:1 molar ratio, such that they can be used as pre-annealed DNA linker.

The second, third, and fourth polynucleotides are substantially homogeneous or pure (*e.g.*, individual polynucleotide molecules within the same container are the same), while the 3'-end of the first polynucleotide in the 3'-overhang region comprises a random-sequence primer (*e.g.*, individual first polynucleotide molecules within the same container are the same except that each may have a different random sequence primer within the 3'-overhang region). Thus the first polynucleotide may be unique in that it is in fact a mixture of polynucleotides differing only at the random-sequence primer region of the individual polynucleotides.

In a related embodiment, however, when a specific ncRNA with a defined 3'-end sequence is of interest, and first polynucleotide of the invention may be homogeneously containing the same matching sequence at the random-sequence primer region, in order to initiate first strand cDNA synthesis specifically from the specific ncRNA with the defined 3'-end sequence.

The random-sequence primer generally has sufficient length (*e.g.*, hexamer), so as to be capable of directing 1st strand cDNA synthesis from the 3'-end of a non-coding RNA. Although hexamer random sequences can be used, other lengths, such as 4, 5, 7, 8, 9, 10, 11, 12 random sequence primers may also be used.

In certain embodiments, the most 3'-end of the random-sequence primer is not deoxythymidine (T) or Uridine (U), or other nucleotide analog that can base pair with adenine (A) in the poly A tail of mRNA. Such design may further help to avoid reverse transcription from the polyA tail of an mRNA.

The 5'- or 3'-overhangs at the 3'-end of the second and third polynucleotides (the first and second ligation compatible ends) are designed to be complementary such that they anneal to each other. The length of the overhang regions in the second and third polynucleotides can be the same, but need not be the same. In certain embodiments, about 2, 3, 4, 5, 6, 7, 8, or more nucleotides in the overhang regions of both polynucleotides are complementary and can form base pairs (Watson-Crick or wobble base pairs).

In certain embodiments, the length of the first double stranded region on the RNA linker is about 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 50, 60 or more base pairs.

In certain embodiments, the length of the second double stranded region on the DNA linker is about 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 50, 60 or more base pairs.

In certain embodiments, the total length of the first and the second double stranded regions, in the ligated RNA-DNA linker, is about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80 or more base pairs.

In certain embodiments, the first double stranded region may comprise a first recognition site for a first restriction enzyme, such as a Type II restriction enzyme (RE). The RE recognition site may be strategically placed such that, when the RE cleaves, it cleaves outside the RE site, 3' to the random-sequence primer. This allows the generation of an RNA tag linked to the RNA linker. For example, a MmeI recognition site may be placed at the end of the first double stranded region, distal to the other end of the first double stranded region (where the RNA linker and the DNA linker are linked via their respective 3'-overhang regions). The MmeI site is designed to be in the orientation such that when MmeI cuts, an RNA tag comprising a 18-bp fragment with a 2 bp overhang is generated in the cDNA derived from a linked ncRNA. However, the placement of the RE site does not need to be at the end of the first double stranded region. A more internal placement generates a correspondingly shorter RNA tag sequence.

In certain embodiments, the last nucleotide of the first recognition site (for the first (Type II) restriction enzyme) is the last base-paired nucleotide 5' to the random-sequence primer.

Likewise, in certain embodiments, the second double stranded region may comprise a second recognition site for a second restriction enzyme, such as a Type II restriction enzyme (RE), which may cleave 3' to the second RE recognition site and 5' to the third polynucleotide. The orientation of the RE recognition site is arranged in such a way that it generates a DNA tag based on the terminal sequence of a linked genomic DNA. In certain embodiments, the placement of the RE site does not need to be at the end of the second double stranded region. A more internal placement generates a correspondingly shorter DNA tag sequence.

In certain embodiments, the last nucleotide of the second recognition site (for the second (Type II) restriction enzyme) is a base-paired nucleotide at the blunt end.

In certain embodiments, the first and the second (Type II) restriction enzymes are the same. In other embodiments, the first and the second (Type II) restriction enzymes are different.

For RE that generates relatively long tag sequences, such as Type I or Type III RE, the orientation of the first and second RE recognition sequences may be reversed, such that

the RE site in the RNA linker directs the generation of a DNA tag, while the RE site in the DNA linker directs the generation of an RNA tag.

For RE that recognizes two recognition sites (such as Type IIB RE), one of the RE site may be in the RNA linker, and the other may be in the DNA linker, such that the RE only cleaves when the RNA and DNA linkers are correctly ligated as designed to reconstitute the full RE recognition site.

Suitable restriction enzymes that may be used according to the instant invention are described in more details below. In certain embodiments, the cleavage site of the first or the second restriction enzyme is at least about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more nucleotides 3' to the last nucleotide of the recognition site.

In certain embodiments, the RNA linker, the DNA linker, or both, does not have a restriction enzyme recognition site for generating the RNA tag or DNA tag.

In certain embodiments, one or more of the first, second, third, and fourth polynucleotides are DNA (*e.g.*, all are DNA), or comprise both DNA and RNA nucleotides. In other embodiments, any of them may be RNA.

In certain embodiments, one or more of the first, second, third, and fourth polynucleotides may comprise a modified nucleotide. The modified nucleotide may be at the 5'-end, 3'-end, and/or at an internal position.

In certain embodiments, the modified nucleotide is a biotinylated nucleotide, such as biotinylated dT (deoxy- thymidine). The presence of the biotinylated nucleotide allows affinity purification of the polynucleotide comprising one or more of such biotinylated nucleotides by, for example, using resins, agarose, nanoparticles, metal or magnetic beads conjugated to a biotin binding partner, such as avidin or streptavidin. Such beads can then be isolated by magnets. The biotinylated nucleotide may be present in the RNA linker, the DNA linker, or both. This technique may also be combined with high throughput next generation sequencing, such as single-molecule real-time sequencing (Pacific Bio); ion semiconductor (Ion Torrent sequencing); pyrosequencing (454); sequencing by synthesis (Illumina); sequencing by ligation (SOLiD sequencing); polony sequencing; massively parallel signature sequencing (MPSS); DNA nanoball sequencing; Heliscope single molecule sequencing, or be used with a Luminex-type system, using color beads or other antibodies for laser- or FACS-based sorting.

In certain embodiments, the modified nucleotide enhances the ability of the random sequence primer to synthesize first strand cDNA via reverse transcription, such as by enhancing the stability and/or specificity of the hybridization between the random primer with the 3'-end of the ncRNA.

In certain embodiments, the random priming sequence may include at least one nucleotide containing a sugar other than the conventional 2'-deoxy-D-ribose or D-ribose found in naturally occurring DNA and RNA, such as a nucleotide in which the sugar is modified by the addition or substitution of a side group, or in which the sugar is a stereoisomer of the conventional 2'-deoxy-D-ribose or D-ribose found in naturally occurring DNA and RNA, or both. See U.S. Pat. No. 6,794,142 (incorporated herein by reference). Such modified nucleotide may be at or near the 3'-end of the random priming sequence. In one embodiment, the modified random primer sequence consists essentially of an oligonucleotide in which at least one of the three 3' terminal nucleotides is a modified nucleotide selected from the group consisting of 2'-O-methyl-nucleotides, 2'-amino-nucleotides, and 2'-fluoro-nucleotides. In one embodiment, the modified primer sequence consists essentially of an oligonucleotide in which at least one of the three 3' terminal nucleotides is a modified nucleotide selected from the group consisting of 2'-O-methyl-ribonucleotides, 2'-deoxy-2'-amino-nucleotides, and 2'-deoxy-2'-fluoro-nucleotides. These modifications represent the addition of a moiety to the 2' OH, or the replacement of the 2'-OH by an alternative moiety.

In certain embodiments, the random priming sequence comprises one or more LNA or PNA. The presence of unusually thermodynamically stable structural fragments in RNAs, such as hairpins, can make it nearly impossible to carry out primer extension. Replacement of DNA primers with LNA-modified primers may overcome this limitation (see Fratzak *et al.*, *Biochemistry*, 48(3):514-6, 2009; Uppuladinne *et al.*, *Biomol. Struct. Dyn.*, 31(6):539-60, 2013).

Other modified nucleotide, such as thiophosphate (or phosphorothioate, a family of compounds and anions with the general chemical formula $PS_{4-x}O_x^{3-}$ ($x = 0, 1, 2, \text{ or } 3$)) modification that renders the internucleotide linkage resistant to nuclease degradation, morpholino oligonucleotides, 2' F-ANA, 2'-O-alkyl, etc., may also be incorporated to the linkers to enhance the stability and nuclease resistant ability of the linkers. See Verma & Eckstein, "Modified oligonucleotides: synthesis and strategy for users," *Annu. Rev. Biochem.*, 67:99-134, 1998 (incorporated herein by reference).

In certain embodiments, the RNA linker and/or the DNA linker may comprise a unique sequence (*e.g.*, a “bar code”) that distinguishes the RNA linker from the DNA linker, or the RNA/DNA linker from other RNA/DNA linker (*e.g.*, when two or more sets of RNA linkers are used together). For example, the first and/or the second double stranded region(s) may comprise a unique sequence that distinguishes the RNA linker from the DNA linker. Such bar code may simply be a small stretch of unique sequence, such as a 2-, 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-nucleotide sequence (or more). In certain embodiments, the difference in the sequence of the RNA linker and the DNA linker may be sufficient to distinguish the RNA linker from the DNA linker. In certain embodiments, only the RNA linker or only the DNA linker has the unique sequence / bar code. In certain embodiments, both the RNA linker and the DNA linker have their respective unique sequences / bar codes.

In certain embodiments, the first polynucleotide is dephosphorylated. In certain embodiments, the second polynucleotide is dephosphorylated. In certain embodiments, the third polynucleotide is dephosphorylated. In certain embodiments, the fourth polynucleotide is dephosphorylated. The dephosphorylation may help to avoid self-ligation of the polynucleotides or the DNA / RNA linkers, such as self-ligation through the blunt ends of two DNA linkers, each may be ligated to a chromosomal DNA fragment in the same chromatin fragment. In addition, if the linkers or the ligatable ends of the linkers are dephosphorylated, it is expected that the linkers are unlikely to ligate to form dimers or concatemers of linkers. Furthermore, it is expected that the DNA linker may ligate to the phosphorylated ends of the chromosomal DNA molecule but cannot ligate to link together the ends of the chromosomal DNA molecules until they are phosphorylated.

In an alternative embodiment, the first and the second polynucleotides may hybridize and form an RNA linker that has, at one end, the 3'-overhang comprising the random priming sequence of the first polynucleotide, and, at the other end, the first ligation compatible site comprising a recognition site for a restriction enzyme. Similarly, the third and the fourth polynucleotides may hybridize and form a DNA linker that has, at one end, the blunt end for ligating to a free end of a chromosomal fragment, and, at the other end, the second ligation compatible end comprising a recognition site for the same restriction enzyme, or a recognition site for a compatible restriction enzyme that generates a compatible ligatable end. Thus digestion by the restriction enzyme and/or its compatible RE produces the overhang (could be 3' or 5' overhang) that can be used to ligate the DNA and RNA linkers.

In this embodiment, prior to the restriction enzyme digestion, the ends of the DNA and RNA linkers may not be ligatable (for example, the RNA linker may have a 5' overhang and the DNA linker may have a blunt end or 3' overhang, or vice versa), and such ends may be further dephosphorylated. After the RE digestion, ligatable ends at the DNA and RNA linker ends are generated, with proper phosphorylation. The ligatable ends of the DNA and RNA linker(s) may then be ligated. The ligatable end after restriction may be a blunt end, or have a cohesive end with a 5' or 3' overhang. In particular, a restriction enzyme which cuts rarely may be used so as to reduce the possibility of cutting the nucleic acid material at unintended locations and/or to produce very short fragments.

The subject polynucleotides can be prepared by any suitable method, including direct chemical synthesis by a method such as the phosphotriester method of Narang *et al.*, 1979, *Meth. Enzymol.*, 68:90-99; the phosphodiester method of Brown *et al.*, 1979, *Meth. Enzymol.*, 68:109-151; the diethylphosphoramidite method of Beaucage *et al.*, 1981, *Tetrahedron Lett.*, 22:1859-1862; and the solid support method of U.S. Pat. No. 4,458,066, each incorporated herein by reference. A review of synthesis methods of conjugates of oligonucleotides and modified nucleotides is provided in Goodchild, 1990, *Bioconjugate Chemistry*, 1(3):165-187, incorporated herein by reference.

One or more additional reagents for carrying out the methods of the invention may also be included in the kit of the invention.

In certain embodiments, the kit further comprises a reagent that cross-links protein and polynucleotide, such as formaldehyde (*e.g.*, 1% formaldehyde).

In certain embodiments, the kit further comprises an affinity reagent that specifically or selectively binds a component of chromatin (*e.g.*, histone or a specific ncRNA of interest). For example, the affinity reagent may be an antibody (such as a monoclonal antibody), or any of the functional antigen-binding fragments or derivatives thereof. The affinity reagent may also be a polynucleotide (such as an antisense polynucleotide) that can hybridize / bind to the polynucleotide component of the chromatin. The antisense polynucleotide may be labeled to facilitate subsequent capture of the hybridization complex formed between the antisense polynucleotide and its complement target sequence. For example, the label can be a biotin label (such as biotinylated U or T) that can be captured by avidin or streptavidin coated beads. The antisense polynucleotide may also be immobilized on a solid support, such as on the surface of a microbeads or nanoparticles, which may be packed into a column or used in

batch mixture for affinity capture of the complement target sequence.

In certain embodiments, the kit further comprises an end-repairing mixture that converts DNA containing damaged or incompatible 5'- and/or 3'-protruding ends to 5'-phosphorylated, blunt-ended DNA. Such reagents are readily available commercially, such as the End-It™ DNA End-Repair Kit from Epicentre.

In certain embodiments, the kit further comprises a DNA ligase (*e.g.*, T4 DNA ligase from various commercial sources, such as New England Biolabs (NEB)).

In certain embodiments, the kit further comprises a reagent that reverses cross-linking of protein and polynucleotide (*e.g.*, Proteinase K from various commercial sources, such as New England Biolabs (NEB)).

In certain embodiments, the kit further comprises the first and/or the second restriction enzyme(s), and optionally any suitable buffers or cofactors required for RE digestion.

In certain embodiments, the kit further comprises a pair of concatenating adapters for PCR amplification of blunt-ended double stranded DNA. The adapters may comprise restriction enzyme sites useful for concatemerization, and may comprise PCR primer sequences suitable for PCR amplification.

In certain embodiments, the kit further comprises a Taq DNA polymerase for PCR amplification, or other DNA polymerases required for other forms of amplification (*e.g.*, rolling circle amplification).

In certain embodiments, the kit further comprises a reverse transcriptase for first strand cDNA synthesis.

Another aspect of the invention provides a paired-end tag (PET) polynucleotide comprising a central region comprising the first and second double stranded regions linked through the first and the second ligation compatible ends, said central region being flanked by: (1) at a site proximal to the first double stranded region, a sequence tag of a non-coding RNA (ncRNA); and (2) at a site proximal to the second double stranded region, a sequence tag of a genomic DNA.

Such PET polynucleotides comprise both the RNA tag and the DNA tag, each derived from the end sequence of the respective ncRNA and genomic DNA (paired-end tag). Together, the paired-end tag represents an observed event or incident where the ncRNA and the genomic DNA fragment are at close proximity of each other in a chromosomal fragment.

In certain embodiments, the sequence tag of the non-coding RNA (ncRNA) has a free end resulting from digestion by the first restriction enzyme.

The restriction enzyme may be any of the ones described above, such as a Type II RE (Type IIS, IIB, IIG, *etc.*), Type I RE, or a Type III RE, which may digest outside their recognition site. Alternatively, the free end may be generated by a naturally existing RE site on the cDNA corresponding to the ncRNA. Preferably, the RE is selected based on the sequence of the central region such that the RE does not cut inside the central region to disrupt the structure of the linked DNA linker and RNA linker.

In certain embodiments, the RNA sequence tag of the ncRNA or the DNA sequence tag of the genomic DNA has a free end resulting from physical shearing, such as shearing by sonication, hydroshearing, repeated drawing through a hypodermic syringe needle, *etc.*

In certain embodiments, the RNA sequence tag of the ncRNA or the DNA sequence tag of the genomic DNA has a free end resulting from limited digestion of a non-specific endonuclease, such as Micrococcal Nuclease (NEB Catalog M0247S), DNase I (NEB Catalog M0303S), or exonucleases that progressively digests from one end of a double stranded DNA, or a combination of endo- and exonucleases (*e.g.*, Exonuclease III and Mung Bean Nuclease) to reduce the average length of the cross-linked genomic DNA or cDNA of ncRNA. The extend of digestion may be controlled by limiting enzyme or substrate concentration, temperature and/or pH of digestion, availability of co-factors, or a combination thereof. Suitable digestion conditions may be pre-tested using standard substrates of defined length, and examining digestion products (by electrophoresis of CE (capillary electrophoresis), *etc.*) before and after digestion.

The length of the RNA or DNA sequence tags should be sufficient to uniquely identify a genomic region from which the ncRNA is transcribed, or at which the genomic DNA is located. For example, the RNA sequence tag of the non-coding RNA (ncRNA) and/or the DNA sequence tag may be about 10-100 base pairs in length (or 15-50 bp, 20-40 bp, 20-30 bp, 20-25 bp) for relatively complicated genomes of higher eukaryotes, but may be shorter (*e.g.*, 6-10 bp, 8-10 bp, 8-12 bp) for relatively simple genomes of bacteria, or lower eukaryotes.

In a related aspect, the invention provides a paired-end tag (PET) polynucleotide library comprising two or more members of the subject PET polynucleotides, wherein each member of the PET library comprises the same central region, and different RNA sequence

tag of the non-coding RNA (ncRNA), different DNA sequence tag of the genomic DNA, or both.

In yet another related aspect, the invention provides a vector or recombinant vector comprising the subject PET polynucleotides.

In certain embodiments, the vector comprises a plurality of concatenated subject PET polynucleotides.

Another aspect of the invention provides a method of identifying functional interaction loci within a genome for non-coding RNAs (ncRNAs) of the genome, the method comprising: (1) providing chromatin fragments comprising cross-linked genomic DNA fragments and cross-linked ncRNAs; (2) using the RNA linker and the DNA linker of the invention, ligating an end of a cross-linked genomic DNA fragment to an end of a cDNA of a cross-linked ncRNA, under a condition for proximity ligation, wherein the end of the cross-linked genomic DNA fragment is ligated to the DNA linker, and the end of the cDNA of the cross-linked ncRNA comprises the RNA linker; (3) isolating a PET polynucleotide of the invention for sequencing analysis; and, (4) mapping the sequence tag of the genomic DNA and the sequence tag of the ncRNA within each PET polynucleotides to a reference genome, thereby identifying functional interaction loci within the reference genome for the non-coding RNAs (ncRNAs) of the reference genome.

In certain embodiments, the methods of the invention are performed using live cells, such as tissue culture cells or cell isolated from freshly dissected tissues. In certain embodiments, the ncRNAs and the genomic DNA in live cells are cross-linked through formaldehyde- and/or EGS (Ethylene glycol bis[succinimidylsuccinate])-mediated cross-linking. Other similar bifunctional crosslinking reagents suitable for crosslinking protein-DNA, protein-RNA and/or protein-protein (*e.g.*, those having two or more reactive chemical groups suitable for reacting with the amide and/or thiol groups) may also be used. If EGS is used, the spacer region between the two NHS-ester may be a 12-atom spacer, although longer or shorter spacers (*e.g.*, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 atom spacers) may be used as well.

If formaldehyde or EGS (typically about 1-2 mM, or 1.5 mM) are used, EGS may be added first followed by (about 1%) formaldehyde. Reaction may be quenched by glycine. Alternatively, about 1% formaldehyde or about 1% glutaraldehyde may be used.

In other embodiments, the nucleic acids are cross-linked to the chromatin via UV cross-linking. For example, tissue culture cells may be UV-crosslinked at about 150 mJ/cm²

at 254 nm (e.g., by using a UV crosslinker, such as STRATALINKER[®] UV crosslinker).

For example, about $1 - 2 \times 10^8$ live tissue culture cells or isolated cells may be first collected and cross-linked with EGS with shaking for 40 min., then formaldehyde (final concentration of about 1%; Sigma) for 10 minutes at room temperature.

Proteinase inhibitor and/or RNase inhibitor may be added to prevent non-specific proteinase or RNase digestion.

The cells are then lysed in a suitable lysis buffer (e.g., 50 mM HEPES, 1 mM EDTA, 0.15 M NaCl, 1% SDS, 1% Triton X-100, 0.1% sodium deoxycholate, all from Ambion).

Once the crosslinking step is complete, various methods may be used to produce chromatin fragments comprising cross-linked genomic DNA and ncRNA.

For example, in certain embodiments, chromatin fragments are generated by physical shearing, such as sonication, hydroshearing, or repeated drawing through a hypodermic syringe needle. Sonication may be advantageous for breaking up the chromatin fibers into tethering complexes with RNA, DNA and protein components, while “shaking off” spurious, random, or weak ncRNA-chromatin-DNA interaction.

Alternatively, in certain embodiments, chromatin fragments may be generated by restriction enzyme digestion, or partial or limited endo- and/or exo-nuclease digestion under controlled conditions, in order to produce RNA and DNA tags of suitable length.

To generate chromatin fragments comprising cross-linked genomic DNA fragments and cross-linked ncRNAs, the chromatin can be solubilized by sonication (e.g., using a Branson 450 ultrasonic cell disruptor, operated at 20% duty power output, 30 second, 5 to 8 times; or using a probe sonicator operating at 35% power for 1.5 min, with 20sec on / 30sec off cycles).

Other commercially available instruments may be used for sonication. For example, the S220 Focused-ultrasonicator from Covaris, Inc. utilizes the Adaptive Focused Acoustics[™] (AFA) technology for DNA, RNA, and chromatin shearing. According to the manufacturer, its software incorporates various preset protocols for standard methods, such as DNA shearing to specific fragment lengths. Alternatively, the BIORUPTOR[®] UCD-200 (Life Technologies Corp.), a benchtop sonication device, may also be used for sonication shearing. The device consists of a high-power ultrasound generating element located below a water bath, and operates at a 20 kHz frequency (similar to a probe sonicator) to provide automated sonication steps suitable for standardized protocols such as ChIP, MeDIP, etc.

Once sheared, the chromatin is diluted (*e.g.*, 10 times) to lower the SDS concentration (*e.g.*, to about 0.1-0.5%). The extract is then cleared by centrifugation (*e.g.*, at 14,000 rpm for 10 minutes at 4°C). This extract can be stored at -80°C until use.

If immunoprecipitation is desired, about 2 µg of monoclonal antibody (specific for a chromatin component) can be bound to protein G sepharose (Pharmacia). The antibody coated beads are then incubated with the chromatin extract at 4°C for 16 hours. The beads are then washed (*e.g.*, with the following reagents from Sigma Chemical Company: Wash buffer 1 (50 mM HEPES, 1 mM EDTA, 0.15 M NaCl, 0.1% SDS, 1% Triton X-100, 0.1% sodium deoxycholate); 2 times Wash buffer 2 (50 mM HEPES, 1 mM EDTA, 0.5 M NaCl, 0.1% SDS, 1% Triton X-100, 0.1% sodium deoxycholate); 1 time Wash buffer 3 (20 mM Tris.HCl pH 8.0, 1 mM EDTA, 0.25 M LiCl, 0.5% NP40, 0.5% sodium deoxycholate); 1 time Wash buffer 4 (20 mM Tris.HCl pH 8.0, 1 mM EDTA). The protein-DNA complexes are then eluted from the beads with elution buffer (*e.g.*, 50 mM Tris.HCl pH 8.0, 1 mM EDTA, 1% SDS) for 20 min at 65°C. The eluent is then dialyzed in PBS (Ambion) to remove SDS (*e.g.*, for 3 hours at 4°C).

Optionally, the chromatin fragments may also be biotinylated (for example, by using EZlink Iodoacetyl-PEG2-Biotin (IPB) (Thermo Scientific, cat.21334)), and be isolated as streptavidin beads-bound chromatin fragments. For example, DYNABEADS® with streptavidin (DYNABEADS® MyOne™ Streptavidin C1/T1) may be used to enrich biotinylated chromatin fragments.

In addition, beads with silica like coating may be used to enrich the crosslinked nucleic acid on the chromatin fragments.

The chromatin fragments, after shearing or RE digestion, may have damaged ends or ends otherwise unsuitable for ligation with the DNA linker. Thus end-repair may be performed using, for example, the End-It kit from Epicentre or the T4 polymerase (Promega, R0191), according to the manufacture's suggestion.

First-strand cDNA synthesis can be performed using a reverse transcriptase and the RNA linker (or the modified RNA linker in the second specific embodiment below), such as the Superscript III First Strand Synthesis System (Life Technologies, cat.18080051).

The repaired chromatin DNA with 5' phosphorylation at its blunt end can then be used in ligation with the DNA linker. This can be carried out in the same container for reverse transcription using the RNA linker, provided that the proper buffer and other reaction conditions for DNA ligation are provided. A DNA ligase, such as the T4 DNA ligase, may

be used for this reaction. If necessary, the dephosphorylated DNA linker can then be phosphorylated (e.g., by T4 polynucleotide kinase).

In certain embodiments, first strand cDNA synthesis is performed (either before or after or concurrent with the DNA linker ligation) using the RNA linker.

In certain embodiments, the cDNA of the cross-linked ncRNA comprises a first strand cDNA reverse transcribed from the random-sequence primer of the RNA linker, and the ncRNA template. Due to the presence of the RNA linker, this first strand cDNA and ncRNA template hybrid molecule can be ligated to the DNA linker already ligated to the free end of the chromosomal DNA fragment.

Once the RNA linker and the DNA linker have been properly ligated to their respective ends of the target nucleic acid, proximity ligation can be performed to connect the DNA linker and the RNA linker on the same chromatin fragment. Proximity ligation is usually carried out at a diluted environment such that RNA and DNA linkers on the same chromatin fragment, due to their proximity to one another, are much more likely to be ligated as compared to RNA and DNA linkers on different chromatin fragments.

In certain embodiments, proximity ligation is carried out with about 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 15, 16, 70, 18, 19, 20-fold or more dilution with respect to the linker ligation steps.

In certain embodiments, proximity ligation is carried out in a total ligation volume of about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 mL or more, for each equivalent amount of captured chromatin fragments derived from about 1×10^8 human cells. Ligation volumes may be adjusted accordingly based on the type of cells (e.g., species of origin or genome size).

The proximity ligation conditions may be modified or adjusted, as required, so as to maximize the ligation of the DNA and RNA linkers. Any ligation condition may be modified or adjusted, including but not limited to increasing or decreasing the time for the ligation reaction and/or concentration of the reagents. In other words, the ligation reactions are adjusted or modified to maximize intermolecular ligation of separate nucleic acid molecules cross-linked to the same chromatin fragment. In particular, the ligation may be performed under very dilute conditions of the nucleic acid molecules to maximize the ligation of the ends of different nucleic acid molecules and to reduce the formation of circular multimers.

In certain embodiments, the method includes assessing the extent or frequency of undesired or false positive ligation events between genomic DNA and ncRNA crosslinked to

different chromatin fragments. Under ideal proximity ligation conditions, only genomic DNA and ncRNA crosslinked to the same chromatin fragment should be ligated.

For example, one set of DNA and RNA linkers (e.g., linker set A) can be used for ligating to the genomic DNA and RNA ends, respectively, in one reaction container. Meanwhile, a second set of DNA and RNA linkers (e.g., linker set B) can be used for ligating to the genomic DNA and RNA ends, respectively, in a second reaction container. The contents of the two reaction containers are then pooled for the proximity ligation. If the RNA linker in linker set A can be ligated to the DNA linkers of both linker sets (and the DNA linker in linker set A can be ligated to the RNA linkers of both linker sets), then proximity ligation condition is optimum if there is no or very infrequent ligation between linkers of sets A and B (e.g., RNA linker in set A ligate to DNA linker in set B). Conversely, proximity ligation condition is less than optimum if there is significant ligation between linkers of sets A and B.

In certain embodiments, the ratio of the RNA and DNA linkers in linker sets A and B can be further adjusted (e.g., not necessarily 1:1). For example, the molar ratio of RNA and DNA linkers in linker set A compared to that in linker set B may be 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, or vice versa.

In certain embodiments, the first, second, third, and/or the fourth polynucleotide of the invention is dephosphorylated and the DNA linker or RNA linker does not self-ligate.

Second strand cDNA synthesis can be completed either before or after the RNA linker - DNA linker ligation, using, for example, the Superscript Double-stranded cDNA Synthesis Kit (Life Technologies, cat.1197-020). In certain embodiments, 2nd strand cDNA synthesis is carried out after proximity ligation but before step (3).

In certain embodiments, a DNA polymerase, such as T4 DNA polymerase, may be added after the 2nd strand cDNA synthesis.

Next, the cross-linked nucleic acid and protein component of the chromatin fragment can be reverse cross-linked with proteinase K. In a typical reaction condition, for example, sample can be reverse cross-linked as 20 μ L aliquots by overnight incubation at 65°C in the presence of 15 μ L of 20 mg/ml proteinase K (Ambion) and optionally 0.3% SDS (Ambion). The following day, about 1 μ L of 10 mg/ml RNase A (Qiagen) may be added to degrade RNA (e.g., for 45 min at 37°C), followed by phenol extraction and ethanol precipitation of DNA.

Optionally, purification or enrichment of at least one linked and reverse cross-linked nucleic acid molecule may be performed using a binding system comprising at least two components, wherein at least one first component is coupled to the linker (*e.g.*, a biotinylated nucleotide incorporated into the RNA or DNA linker, for example), and at least a second component binds the first component. The components include but are not limited to streptavidin-biotin, avidin-biotin, protein-antibody and/or magnet/magnetic material.

In particular, biotinylated linker-ligated nucleic acid material may be purified using streptavidin beads, such as streptavidin-conjugated magnetic DYNABEADS™ (Life Technologies, cat.11206D-10ML). Only the nucleic acid material that contains biotinylated linkers will be immobilized on the streptavidin beads. If another component is bound to the linkers used, other systems of purifying the nucleic acid molecules suitable for the component may be used.

Alternatively, streptavidin columns may be used instead to capture the biotinylated beads. In yet another alternative, the beads may be color or fluorescently coated such that they can be sorted or collected by FACS, etc., on a flow-based detection instrument (*e.g.*, LUMINEX® 100™, LUMINEX® 200™ or BIO-RAD® BIO-PLEX® type analyzer).

The resulting released DNA can be used to produce PET polynucleotides having paired DNA and RNA tags through, for example, RE enzyme digestion. Optionally, the released PET polynucleotides may be further amplified by PCR before sequencing analysis. PCR adapters may be ligated to both ends of the PET polynucleotides (such as by T4 DNA ligase) before carrying out the PCR amplification. Only blunt ended, non-circularized nucleic acids can be ligated to the adapters. Self-ligated nucleic acid molecules and circular multimers cannot be ligated to the adapters.

The PCR adapters may also comprise modified nucleotides for PCR product purification. Similarly, streptavidin-biotin, avidin-biotin, protein-antibody and/or magnet/magnetic material may be used for this purpose.

The PET polynucleotides (with or without amplification) may be directly sequenced, such as according to the protocols for the various next-generation sequencing, such as 454 sequencing using the 454 multiplex sequencing machine (454 life sciences). The technique is taught in Margulies *et al* (2005) and US Application No. 20030068629 (both incorporated herein by reference). Any other high throughput or next-generation sequencing (NGS) methods may be used to determine the sequences of the PET polynucleotides.

Mapping of the obtained RNA / DNA tag sequences to their respective genomic locations can be performed using any of many commercially available tools, software, or services.

Once the RNA and DNA tags of the PET polynucleotides are sequenced and mapped to the reference genome, each linked RNA tag and DNA tag represents a putative ncRNA-chromatin interaction. The collection of all such observed interactions constitute the functional interaction loci within the reference genome for the non-coding RNAs (ncRNAs) of the reference genome.

In certain embodiments, the method further comprises identifying clusters of two or more PET polynucleotides having overlapping sequence tags of the genomic DNA and overlapping sequence tags of the ncRNA.

The PET clusters are considered as high confidence data, reflecting recurrent detection of more reliable events of ncRNA-chromatin interactions. In contrast, the singleton PETs with no overlap on both the RNA tag and the DNA tag with other PET sequences may represent weak linking signals, and may be indistinguishable from random background noises.

In certain embodiments, the method further comprises excluding PET polynucleotides comprising sequence tags of rRNA. Although some rRNA-chromatin-gDNA (genomic DNA) interactions may be of true biological significance, the presence of large amount (about 1/4 in some data set) of rRNA-chromatin-DNA interactions may obscure the other less abundant interactions. Thus such digital subtraction before further data analysis may be desirable for analyzing less frequent ncRNA-chromatin interactions.

In certain embodiments, the method further comprises isolating or enriching a subset of chromatin fragments prior to the proximity ligation step. For example, the subset of chromatin fragments can be isolated or enriched by immunoprecipitation using an antibody specific for a protein component of the subset of chromatin fragments, or by hybridization using a (labeled) polynucleotide specific for a nucleic acid component of the subset of chromatin fragments. This may be useful for identifying specific interactions between a known chromatin component and ncRNA.

In certain embodiments, the protein component is a histone, a transcription factor (such as a general transcription factor RNAPII, RNAPI, RNAPIII), a polycomb-group (PcG) family protein that remodels chromatin (such as EZH2, and others from insects, mammals, and plants); a recombination involved factor (such as PRDM9); a chromatin insulator or

chromatin waver (such as CTCF); a methyl-CpG-binding protein (such as MeCP2); or an RNA binding protein.

In a variation of the method, a specific labeled ncRNA (such as biotinylation) may be added to the cell before crosslinking. Such labeled ncRNA can be isolated or enriched by using magnetic beads coated with avidin or streptavidin.

In yet another variation of the method, complementary sequences to one or more specific ncRNAs of interest may be used to isolate or enrich such specific ncRNAs (using an array or column) cross-linked to chromatin fragments. Once isolated or enriched, such chromatin fragments can be subject to the remaining steps of the method to identify the regions of genomic DNA that interacts with the specific ncRNA.

In certain embodiments, the method further comprises validating one or more observed ncRNA- chromatin interaction by, for example, DNA/RNA FISH and immunofluorescence assays. For instance, if a specific ncRNA is linked to a particular genomic locus, DNA/RNA FISH and immunofluorescence assays may be performed using the ncRNA to confirm the observation (see, for example, Figure 4B).

b) Modified RNA Linker

In another / second specific embodiment, the method of the invention can be carried out using one modified RNA linker (and no DNA linker) to ligate the crosslinked RNA and chromosomal DNA in the same chromatin fragment.

Thus another aspect of the invention provides a modified RNA linker comprising: (i) a first polynucleotide, and, (ii) a second polynucleotide, wherein the first and the second polynucleotides form a double stranded region flanked by a genomic DNA ligation compatible end, and a 3'-overhang at the 3'-end of the first polynucleotide, wherein the 3'-overhang comprises a random-sequence primer.

According to this aspect of the invention, the 3'-overhang at the 3'-end of the first polynucleotide has a similar function as that of the RNA linker in the specific embodiment described in subsection a) (RNA and DNA linker pair), while the genomic DNA ligation compatible end can be used ligate blunt ended genomic DNA crosslinked to the same chromatin fragment.

In certain embodiments, the ligation compatible end may be blunt ended for direct ligation to the blunt end of the crosslinked genomic DNA fragment.

In another embodiment, the ligation compatible end may comprise a restriction enzyme site, which can be cleaved by the RE to produce the requisite blunt end required for ligation to the blunt end of the crosslinked genomic DNA fragment. Prior to cleavage by the restriction enzyme, however, the ligation compatible ends may be blunt ended (*e.g.*, dephosphorylated blunt end to prevent self-ligation), or have non-compatible overhang that prevents self-ligation.

In certain embodiments, the modified RNA linker does not self-ligate, either through its 3'-overhang or its ligation compatible end.

The first and second polynucleotides may be provided in separate containers, such as synthesized polynucleotides, either in freeze dried, lyophilized form or in water or a suitable buffer solution. Alternatively, the first and the second polynucleotides may be combined in the same container (lyophilized or in solution), for example, in 1:1 molar ratio, such that they can be used as pre-annealed modified RNA linker.

The second polynucleotide is substantially homogeneous or pure (*e.g.*, individual polynucleotide molecules within the same container are the same), while the 3'-end of the first polynucleotide in the 3'-overhang region comprises a random-sequence primer.

In a related embodiment, the first polynucleotide may be homogeneously containing the same matching sequence at the random-sequence primer region, in order to initiate first strand cDNA synthesis specifically from the specific ncRNA with the defined 3'-end sequence.

In certain embodiments, the double stranded region may comprise a first recognition site for a first restriction enzyme, such as a Type II restriction enzyme (RE). The RE recognition site may be strategically placed such that, when the RE cleaves, it cleaves outside the RE site, 3' to the random-sequence primer. This allows the generation of an RNA tag linked to the RNA linker. For example, a MmeI recognition site may be placed at the end of the double stranded region, proximal to the 3' overhang comprising the random-sequence primer. The MmeI site is designed to be in the orientation such that when MmeI cuts, an RNA tag comprising a 18-bp fragment with a 2 bp overhang is generated in the cDNA derived from a linked ncRNA. However, the placement of the RE site does not need to be at the end of the first double stranded region. A more internal placement generates a correspondingly shorter RNA tag sequence.

In certain embodiments, the last nucleotide of the first recognition site (for the first (Type II) restriction enzyme) is the last base-paired nucleotide 5' to the random-sequence

primer.

In certain embodiments, the double stranded region may comprise a second recognition site for a second restriction enzyme, such as a Type II restriction enzyme (RE), at or near the ligation compatible end. The RE may cleave 3' to the second RE recognition site and 5' to the first polynucleotide (e.g., into the ligated genomic DNA). The orientation of the RE recognition site is arranged in such a way that it generates a DNA tag based on the terminal sequence of a linked genomic DNA. In certain embodiments, the placement of the RE site does not need to be at the end of the double stranded region. A more internal placement generates a correspondingly shorter DNA tag sequence.

In certain embodiments, the last nucleotide of the second recognition site (for the second (Type II) restriction enzyme) is a base-paired nucleotide at the ligation compatible / blunt end.

In certain embodiments, the modified RNA linker does not have a restriction enzyme recognition site for generating the RNA tag or DNA tag.

In certain embodiments, the modified RNA linker may comprise a unique sequence (e.g., a "bar code") that distinguishes the modified RNA linker from other modified RNA linker(s).

In certain embodiments, the first and/or the second polynucleotide is dephosphorylated.

Another aspect of the invention provides a paired-end tag (PET) polynucleotide comprising a central region comprising the double stranded region (of the modified RNA linker) flanked by: (1) at a site proximal to the random-sequence primer, a sequence tag of a non-coding RNA (ncRNA); and (2) at a site proximal to the ligation compatible end, a sequence tag of a genomic DNA.

In a related aspect, the invention provides a paired-end tag (PET) polynucleotide library comprising two or more members of the subject PET polynucleotides, wherein each member of the PET library comprises the same central region, and different RNA sequence tag of the non-coding RNA (ncRNA), different DNA sequence tag of the genomic DNA, or both.

In yet another related aspect, the invention provides a vector or recombinant vector comprising the subject PET polynucleotides.

Another aspect of the invention provides a method of identifying functional interaction loci within a genome for non-coding RNAs (ncRNAs) of the genome, the method comprising: (1) providing chromatin fragments comprising cross-linked genomic DNA fragments and cross-linked ncRNAs; (2) using the modified RNA linker of the invention, ligating an end of a cross-linked genomic DNA fragment to an end of a cDNA of a cross-linked ncRNA, under a condition for proximity ligation, wherein the end of the cross-linked genomic DNA fragment is ligated to the ligation compatible end of the modified RNA linker, and the end of the cDNA of the cross-linked ncRNA comprises the modified RNA linker; (3) isolating a PET polynucleotide of the invention for sequencing analysis; and, (4) mapping the sequence tag of the genomic DNA and the sequence tag of the ncRNA within each PET polynucleotides to a reference genome, thereby identifying functional interaction loci within the reference genome for the non-coding RNAs (ncRNAs) of the reference genome.

In certain embodiments, the cDNA of the cross-linked ncRNA comprises a first strand cDNA reverse transcribed from the random-sequence primer of the modified RNA linker, and the ncRNA template. Due to the presence of the modified RNA linker, this first strand cDNA and ncRNA template hybrid molecule can be ligated to the free end of the chromosomal DNA fragment.

In certain embodiments, the length of the double stranded region on the modified RNA linker is about 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 50, 60 or more base pairs.

Other embodiments as described in the first specific embodiment described in subsection a) (RNA and DNA linker pair) are generally applicable, and are incorporated (but not reiterated) herein.

c) Direct RNA-DNA Ligation

In another / third specific embodiment, the method of the invention can be carried out using certain enzymes (such as the truncated RNA Ligase 2 or RNL2) that directly ligate the 3'-OH group of the ncRNA to a 5' adenylated single-stranded DNA (5' App-ssDNA), such as a ssDNA linker that is later hybridized to a complement polynucleotide, or a dsDNA with a 5' adenylated overhang that can serve as a substrate of the enzyme for direct ligation to the 3'-OH group of the ncRNA.

Thus the invention also provides an alternative way to ligate the 3'-end of the cross-linked ncRNA and a free end of a cross-linked genomic DNA fragment in the same chromatin fragment. According to this aspect of the invention, a single stranded DNA

oligonucleotide is provided with its 5' pre-Adenylated (5' App ssDNA). An RNA-DNA ligase (such as the Thermostable 5' AppDNA/RNA Ligase, NEB Catalog M0319S or M0319L) can then be used to directly link the 3'-OH of the ncRNA to the 5' App ssDNA.

According to the manufacture, the thermostable 5' App DNA/RNA Ligase is a point mutant of catalytic lysine of RNA ligase from *Methanobacterium thermoautotrophicum* (Zhelkovsky and McReynolds, *BMC Mol. Biol.*, 13:24, 2012). This enzyme is ATP independent, but requires a 5' pre-adenylated linker for ligation to the 3'-OH end of either RNA or single stranded DNA (ssDNA). The enzyme is also active in ligation of RNA with 2'-O-methylated 3' end to 5'-adenylated linkers (Zhelkovsky and McReynolds, *supra*). The mutant ligase is unable to adenylate the 5'-phosphate of RNA or ssDNA, which reduces the formation of undesired ligation products (concatemers and circles). The ability of the ligase to function at 65°C might further reduce the constraints of RNA secondary structure in RNA ligation reactions.

Another suitable ligase for this embodiment of the invention is RNA Ligase 2, such as the AIRTM RNA Ligase 2 (RNL2) from Bioo Scientific (Austin, TX), which specifically ligates the adenylated 5' end of an adapter to the 3' end of RNA. Similarly, the enzyme does not require ATP for ligation but does need an adenylated substrate, which dramatically reduces the amount of ligation between random RNA molecules. The ligase is a truncated version of T4 RNA Ligase 2. Unlike the full length RNA ligase 2, AIRTM Ligase does not ligate the phosphorylated 5' end of RNA or DNA without the adenylated substrate.

Alternatively, T4 RNA ligase 1 (NEB Cat. No. M0204S or M0204L) may be used to ligate the ncRNA 3'-OH to the 5' phosphoryl-terminated ssDNA.

Once the 3'-end of the ncRNA is ligated to the ssDNA, a complementary ssDNA can be anneal to the ligated ssDNA to initiate 2nd strand cDNA synthesis, and/or to form a blunt end suitable for ligation with the free end of a cross-linked genomic DNA fragment in the same chromatin fragment.

In an alternative embodiment, a dsDNA linker having a blunt end (or a ligation compatible end) at one end and a 5' adenylated overhang (that can serve as the single strand substrate for the various RNA ligases above) at the other end can first be ligated to the free end of the crosslinked genomic DNA fragment, before the protruding adenylated 5' end is directly ligated to the 3'-OH of the ncRNA.

Likewise, all the embodiments or variations described above for the ligated RNA linker - DNA linker or the modified RNA linker are generally applicable to the double

stranded region formed between the 5' App ssDNA and its complementary sequence.

For example, in certain embodiments, the double stranded region formed between the 5' App ssDNA and its complementary sequence may comprise one or more RE recognition sites to facilitate the generation of RNA and DNA tag sequences. Two MmeI sites can be situated at both ends of the double stranded region and direct the cleavage outside the double stranded region to generate 18-20 bp RNA and DNA tags flanking the double stranded region. Alternatively, one RE site may be used to generate the RNA tag (or the DNA tag), and the DNA tag (or the RNA tag) may be generated by physical shearing or limited non-specific enzyme digestion (see above).

Thus another aspect of the invention provides a direct RNA linker comprising: (i) a first polynucleotide, and, (ii) a second polynucleotide, wherein the first and the second polynucleotides form a double stranded region flanked by a genomic DNA ligation compatible end, and a 5'-overhang at the 5'-end of the first polynucleotide.

The 5'-overhang is optionally 5' adenylated, or can be adenylated by a suitable enzyme, such as the Mth RNA Ligase in the 5' DNA adenylation kit (Cat. No. E2610S or E2610L). If the RNA ligation is to be performed with the 5'-overhang, as opposed to the first polynucleotide as a ssDNA (before its annealing with the second polynucleotide), the 5'-overhang is of sufficient length (e.g., 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 bases or more) to be used as a substrate for the enzyme for direct RNA ligation.

In certain embodiments, the ligation compatible end may be blunt ended for direct ligation to the blunt end of the crosslinked genomic DNA fragment.

In another embodiment, the ligation compatible end may comprise a restriction enzyme site, which can be cleaved by the RE to produce the requisite blunt end required for ligation to the blunt end of the crosslinked genomic DNA fragment. Prior to cleavage by the restriction enzyme, however, the ligation compatible ends may be blunt ended (e.g., dephosphorylated blunt end to prevent self-ligation), or have non-compatible overhang that prevents self-ligation.

In certain embodiments, the direct RNA linker does not self-ligate. For example, the 3' end of the first polynucleotide may be blocked by a dideoxynucleotide or other modified nucleotide to prevent self-ligation (self-circularization) of the first polynucleotide. Upon completion of RNA-DNA ligation, the blocked 3' end of the first polynucleotide becomes part of the ligation compatible end, and may be cleaved off through RE digestion to create a blunt end for genomic DNA ligation.

In certain embodiments, the double stranded region may comprise a first recognition site for a first restriction enzyme, such as a Type II restriction enzyme (RE). The RE recognition site may be strategically placed such that, when the RE cleaves, it cleaves outside the RE site, 5' to the 5' adenylated end of the first polynucleotide. This allows the generation of an RNA tag linked to the direct RNA linker. For example, a MmeI recognition site may be placed at the end of the double stranded region, proximal to the 5' end of the 5'-overhang of the first polynucleotide. The MmeI site is designed to be in the orientation such that when MmeI cuts, an RNA tag comprising a 18-bp fragment with a 2 bp overhang is generated in the cDNA derived from a linked ncRNA. However, the placement of the RE site does not need to be at the end of the first polynucleotide. A more internal placement generates a correspondingly shorter RNA tag sequence. Longer RNA tag sequences can be generated if the first polynucleotide is used as a ssDNA substrate (as opposed to its 5'-overhang is used as substrate), since the RE site can be placed at the 5'-end of the first polynucleotide.

Thus in certain embodiments, the last nucleotide of the first recognition site (for the first (Type II) restriction enzyme) is the 5'-end of the first polynucleotide.

In certain embodiments, the double stranded region may comprise a second recognition site for a second restriction enzyme, such as a Type II restriction enzyme (RE), at or near the ligation compatible end. The RE may cleave 3' to the second RE recognition site and 3' to the first polynucleotide (e.g., into the ligated genomic DNA). The orientation of the RE recognition site is arranged in such a way that it generates a DNA tag based on the terminal sequence of a linked genomic DNA. In certain embodiments, the placement of the RE site does not need to be at the end of the double stranded region. A more internal placement generates a correspondingly shorter DNA tag sequence.

In certain embodiments, the last nucleotide of the second recognition site (for the second (Type II) restriction enzyme) is a base-paired nucleotide at the ligation compatible / blunt end.

In certain embodiments, the direct RNA linker does not have a restriction enzyme recognition site for generating the RNA tag or DNA tag.

In certain embodiments, the direct RNA linker may comprise a unique sequence (e.g., a "bar code") that distinguishes the direct RNA linker from other direct RNA linker(s).

In certain embodiments, the second polynucleotide is dephosphorylated.

The PET polynucleotides generated according to this aspect of the invention comprises a central region corresponding to the double stranded region formed between the

5' App ssDNA and its complementary sequence (i.e., the second polynucleotide). There is no specific sequence requirement for this region, and the length of the region is flexible (*e.g.*, as short as a few bp, sufficient long to support the substrate requirement of the RNA-DNA ligase, and that for the reverse transcriptase), although longer sequences may be used to incorporate any desired RE recognition sites, bar code sequences, or modified nucleotide (*e.g.*, biotinylated nucleotide for affinity purification).

Thus another aspect of the invention provides a paired-end tag (PET) polynucleotide comprising a central region comprising the double stranded region (of the direct RNA linker) flanked by: (1) at a site proximal to the 5' end of the first polynucleotide (either 5' adenylated or suitable to be 5' adenylated), a sequence tag of a non-coding RNA (ncRNA); and (2) at a site proximal to the ligation compatible end, a sequence tag of a genomic DNA.

In a related aspect, the invention provides a paired-end tag (PET) polynucleotide library comprising two or more members of the subject PET polynucleotides, wherein each member of the PET library comprises the same central region, and different RNA sequence tag of the non-coding RNA (ncRNA), different DNA sequence tag of the genomic DNA, or both.

In yet another related aspect, the invention provides a vector or recombinant vector comprising the subject PET polynucleotides.

Yet another aspect of the invention provides a method of identifying functional interaction loci within a genome for non-coding RNAs (ncRNAs) of the genome, the method comprising: (1) providing chromatin fragments comprising a cross-linked genomic DNA fragment and a cross-linked ncRNA; (2) ligating the 3'-OH of the ncRNA to a 5' pre-adenylated ssDNA; (3) providing a complement of the ssDNA to form a double stranded region between the ssDNA and the complement, (4) if necessary, producing a blunt end at the end of the double stranded region; (5) ligating the blunt end to an end of the cross-linked genomic DNA fragment under a condition for proximity ligation; (6) isolating a PET polynucleotide for sequencing analysis, wherein the PET polynucleotide comprises the double stranded region flanked by a DNA tag of the cross-linked genomic DNA fragment and an RNA tag of the ncRNA; and, (7) mapping the DNA tag and the RNA tag to a reference genome, thereby identifying functional interaction loci within the reference genome for the non-coding RNAs (ncRNAs) of the reference genome.

An alternative aspect of the invention provides a method of identifying functional interaction loci within a genome for non-coding RNAs (ncRNAs) of the genome, the method

comprising: (1) providing chromatin fragments comprising a cross-linked genomic DNA fragment and a cross-linked ncRNA; (2) ligating the 3'-OH of the ncRNA to a 5' pre-adenylated overhang of a dsDNA having a double stranded region, (4) if necessary, producing a blunt end at the end of the double stranded region distal to the 5' pre-adenylated overhang; (5) ligating the blunt end to an end of the cross-linked genomic DNA fragment under a condition for proximity ligation; (6) isolating a PET polynucleotide for sequencing analysis, wherein the PET polynucleotide comprises the double stranded region flanked by a DNA tag of the cross-linked genomic DNA fragment and an RNA tag of the ncRNA; and, (7) mapping the DNA tag and the RNA tag to a reference genome, thereby identifying functional interaction loci within the reference genome for the non-coding RNAs (ncRNAs) of the reference genome.

In certain embodiments, the complement of the ssDNA (i.e., the second polynucleotide) has the same length as the ssDNA. In certain embodiments, the complement is longer or shorter than the ssDNA, and forms a double stranded region with a protruding 3' or 5' end. In the latter case, the overhang can be filled-in by enzyme to generate a ligation suitable blunt end, or by cut off from the end by a restriction enzyme that generates a blunt end. The RE site can be engineered into the sequence of the ssDNA.

In certain embodiments, the length of the first polynucleotide of the direct RNA linker is about 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 50, 60 or more bases.

Other embodiments as described in the first and second specific embodiments described in subsection a) (RNA and DNA linker pair) and subsection b) (modified RNA linker), respectively, are generally applicable, and are incorporated (but not reiterated) herein.

With the general aspects of the invention so described, the following sections provide additional details and specific quantities and parameters relating to specific embodiments of the present invention. It shall be apparent to one of skill in the art that the invention may be practiced without such details or with minor modifications without departing from the general scope of the invention.

2. *Definitions*

“Non-coding RNA (ncRNA)” includes an RNA molecule that is not translated into a protein. Less frequently, it may also be referred to as non-protein-coding RNA (npcRNA), non-messenger RNA (nmRNA) and functional RNA (fRNA). It is usually a functional RNA having a function other than encoding protein, but some may be non-functional or without a

known function. Sometimes, the term small RNA (sRNA) is often used for short bacterial ncRNAs. The DNA sequence from which a non-coding RNA is transcribed is often called an RNA gene.

Non-coding RNA genes include highly abundant and functionally important RNAs, such as transfer RNA (tRNA) and ribosomal RNA (rRNA), as well as RNAs such as snoRNAs (including scRNA; for nucleotide modification of RNAs), snRNA (for splicing and other functions), gRNA (guide RNA; for mRNA nucleotide modification), RNase P (for tRNA maturation), RNase MRP (for rRNA maturation, and/or DNA replication), Y RNA (for RNA processing, and/or DNA replication), telomerase RNA (for Telomere synthesis), spliced leader RNA, SmY RNA (for mRNA trans-splicing), antisense RNA, cis-natural antisense transcript, microRNA (for gene regulation), siRNA (including trans-acting siRNA; for gene regulation), exRNAs, and piRNA (including repeat associated siRNA; for transposon defense, and maybe other functions), 7SK RNA (for negatively regulating CDK9/cyclin T complex), and the long ncRNAs that include examples such as Xist and HOTAIR. The number of ncRNAs encoded within the human genome is unknown, but recent transcriptomic and bioinformatic studies suggest the existence of thousands of ncRNAs. Since many of the newly identified ncRNAs have not been validated for their function, it is possible that many are non-functional.

In certain embodiments, ncRNA of the invention does not include any one or more of the above-referenced species. For example, in certain embodiments, ncRNA of the invention does not include rRNA. In certain embodiments, ncRNA of the invention does not include tRNA. In certain embodiments, ncRNA of the invention does not include tRNA.

“Restriction enzyme (RE)” and “restriction endonuclease” are used interchangeably herein to include an enzyme that cleaves double-stranded DNA. The enzyme typically makes two incisions, at, within, or near (*e.g.*, from about a few bases to about a few kilobases) specific recognition nucleotide sequences known as “restriction sites” or “RE recognition sites,” one through each of the phosphate backbones of the double helix without damaging the bases.

Restriction enzymes are commonly classified into three types, which differ in their structure and whether they cut their DNA substrate at their recognition site, or if the recognition and cleavage sites are separate from one another. Over 3000 restriction enzymes have been studied in detail so far, and more than 600 of these are available commercially, many of which are routinely used for DNA modification and manipulation in molecular

biology.

Type I restriction enzymes cut at a site that differs, and is a random distance (at least 1000 bp) away, from their recognition site. The Type I restriction enzyme recognition site is asymmetrical, and is composed of two specific portions - one containing 3-4 nucleotides, and another containing 4-5 nucleotides - separated by a non-specific spacer of about 6-8 nucleotides. These enzymes are multifunctional and are capable of both restriction and modification activities, depending upon the methylation status of the target DNA. Cofactors S-Adenosyl methionine (AdoMet), hydrolyzed adenosine triphosphate (ATP), and magnesium (Mg^{2+}) ions are required for their full activity.

Typical type II restriction enzymes are homodimers, with recognition sites that are usually undivided, palindromic, and 4-8 nucleotides in length. They recognize and cleave DNA at the same site, and they do not use ATP or AdoMet for their activity - they usually require only Mg^{2+} as a cofactor. Recently, new subfamily nomenclature (defined using a letter suffix) was developed to divide this large family into subcategories based on deviations from typical characteristics of type II enzymes. For example, Type IIB restriction enzymes (*e.g.*, BcgI and BpII) are multimers requiring both AdoMet and Mg^{2+} cofactors, and they cleave DNA on both sides of their recognition to cut out the recognition site. Type IIE restriction endonucleases (*e.g.*, NaeI) cleave DNA following interaction with two copies of their recognition sequence. One recognition site acts as the target for cleavage, while the other acts as an allosteric effector that speeds up or improves the efficiency of enzyme cleavage. Similar to type IIE enzymes, type IIF restriction endonucleases (*e.g.*, NgoMIV) interact with two copies of their recognition sequence but cleave both sequences at the same time. Type IIG restriction endonucleases (Eco57I) do have a single subunit, like classical Type II restriction enzymes, but require the cofactor AdoMet to be active. Type IIM restriction endonucleases, such as DpnI, are able to recognize and cut methylated DNA. Type IIS restriction endonucleases (*e.g.*, FokI) cleave DNA at a defined distance from their non-palindromic asymmetric recognition sites. That is, Type IIS enzymes cleave outside of their recognition sequence to one side. MmeI as well as most of the type IIS restriction enzymes produce variable end lengths. Dunn *et al.* (2002) showed that MmeI can cut 18/20 or 19/21 bases away in a rough proportion of 1:1. Therefore, when 18/20 is used to describe MmeI restriction cleavage site, 19/21 is also contemplated. Type IIT restriction enzymes (*e.g.*, Bpu10I and BslI) are composed of two different subunits. Some recognize palindromic sequences while others have asymmetric recognition sites.

Type III restriction enzymes (*e.g.*, EcoP15) recognize two separate non-palindromic sequences that are inversely oriented. They cut DNA about 20-30 base pairs after the recognition site. These enzymes contain more than one subunit and require AdoMet and ATP cofactors for their roles in DNA methylation and restriction, respectively. Type III enzymes recognize short 5-6 bp long asymmetric DNA sequences and cleave 25-27 bp downstream to leave short, single-stranded 5' protrusions. They require the presence of two inversely oriented unmethylated recognition sites for restriction to occur.

Restriction enzyme cleavage products may be blunt-ended or have sticky ends with 5' or 3' overhangs, which sticky-end fragment can be ligated not only to the fragment from which it was originally cleaved, but also to any other fragment with a compatible cohesive or sticky end.

"Nucleotide" as used herein includes a phosphoric ester of nucleoside - the basic structural unit of nucleic acids (DNA or RNA). Short strands of two or more nucleotides (*e.g.*, 2-30, 5-25, 10-15 nucleotides) are sometimes referred to as "oligonucleotides," while longer strands are referred to as polynucleotides, although there is no definitive length limitation between the two terms. The term nucleotide may be used interchangeably with the term "nucleic acid." A polynucleotide may be either single-stranded, or double-stranded with each strand having a 5' end and a 3' end. The end regions of a *stretch* of nucleic acid may be referred to as the 5' terminus and the 3' terminus respectively. The nucleotides in a polynucleotide may be natural nucleotides (deoxyribonucleotides A, T, C, or G for DNA, and ribonucleotides A, U, C, G for RNA), or may include modified nucleotides, which may be incorporated into a polynucleotide by, for example, chemical synthesis. Such modified nucleotides may confer additional desirable properties absent or lacking in the natural nucleotides, and polynucleotides comprising modified nucleotides may be used in the compositions and methods of the invention.

The term "primer" or "priming sequence" refers to an oligonucleotide capable of acting as a point of initiation of DNA synthesis under conditions in which synthesis of a primer extension product complementary to a nucleic acid strand is induced, *i.e.*, either in the presence of four different nucleoside triphosphates and an agent for extension (*e.g.*, a DNA polymerase or reverse transcriptase) in an appropriate buffer and at a suitable temperature. A primer may be a single-stranded DNA. The appropriate length of a primer depends on the intended use of the primer but typically ranges from 10 to 50 nucleotides, such as from 15-35 nucleotides. Short primer molecules generally require lower temperatures to form

sufficiently stable hybrid complexes with the template. A primer need not reflect the exact sequence of the template nucleic acid, but must be sufficiently complementary to hybridize with the template. The design of suitable primers for the amplification of a given target sequence is well known in the art and described in, for example, the literature cited herein.

A “probe” generally refers to a nucleic acid molecule or a sequence complementary therewith, used to detect the presence of at least a portion of the cDNA or an mRNA of a target sequence, such as the CCAT1 ncRNA sequence or cDNA thereof. The detection may be carried out by identification of hybridization complexes between the probe and the assayed target sequence. The probe can be attached to a solid support or to a detectable label. The probe will generally be single stranded. The probe(s) typically comprise 10 to 200 nucleotides. The particular properties of a probe will depend upon the particular use and are within the competence of one of ordinary skill in the art to determine. Generally, the probe will hybridize to at least a portion of the target cDNA or RNA under conditions of high stringency hybridization.

“Adapter” refers to an oligonucleotide molecule to be ligated or is ligated to an end of a nucleic acid molecule. Adapters may be used for amplification (PCR adapter having PCR primer sequences), sequencing (having sequencing primer sequences), and/or inserting a nucleic acid fragment into a vector (having suitable cloning sequences, such as RE recognition sites).

“Concatemer” is usually composed of at least two nucleotide monomer sequences linked end to end, optionally separated by a linker or spacer. The monomers may or may not be the same in sequence, but may have similar structural elements (such as the RNA and DNA linkers of the invention). The monomers may also be in the same or different orientation (*e.g.*, monomers within a concatemer may be linked to one another head-to-head, head-to-tail, or a mixture of both). A concatemer of the invention comprises at least two oligonucleotides (*e.g.*, PET polynucleotides) prepared according to the method of the invention.

“Library” includes a collection of like nucleic acid sequences, oligonucleotides, or polynucleotides, with each member of the library sharing one or more defining characteristics. For example, a library of PET polynucleotide of the invention comprises two or more (*e.g.*, tens of thousands, hundreds of thousands, millions, tens of millions, *etc.*) PET polynucleotides of the invention, with each PET polynucleotide sharing a similar or identical structure but having different DNA and/or RNA tag sequences.

“Vector” or “recombinant vector” is an art-recognized term referring to a bacteriophage, plasmid, or other agent that is capable of transferring or amplifying a genetic material contained within (*e.g.*, a cloned genetic information or cloned DNA) from one cell to another. Such vectors, depending on specific nature and characteristics, may be introduced into different host cells by transfection and/or transformation, such as lipofection, calcium phosphate precipitation, retroviral deliver, electroporation, and biolistic transformation, and any other molecular biology techniques available in the art.

Suitable vectors may include a plasmid, a viral vector or other vehicle known in the art that has been manipulated by insertion or incorporation of heterologous genetic sequences. Such vectors may contain a replication origin for suitable host amplification, a promoter sequence that may facilitate the efficient transcription of the cloned sequences, flanking PCR primers for direct amplification of the cloned sequences. The vector may also comprise specific genes that allow phenotypic selection of the transformed cells. Vectors suitable for use in the present invention include for example, pBlueScript (Stratagene, La Jolla, Calif.); pBC, pZErO-1 (Invitrogen, Carlsbad, CA) and pGEM3z (Promega, Madison, WI) or modified vectors thereof as well as other similar vectors known to those of skill in the art. See, for example, the pGEM vectors disclosed in U.S. Pat. No. 4,766,072, herein incorporated by reference.

“Chromatin” is used to describe a complex of nucleic acids and proteins, primarily histones, in the cell nucleus that stains readily with basic dyes and condenses to form chromosomes during cell division. Chromatin is an example of a nucleic acid-protein complex.

“Tag” as used herein includes an identifiable stretch of sequence of nucleic acids that may uniquely identify the origin of the sequence within a reference genome. The tag may be of sufficient length (usually 18-20 bp, but can be shorter depending on the sequence composition and reference genome size and complexity, *etc.*) that uniquely or unambiguously maps the tag to one or several locations (such as duplicate copies of one gene or related genes with high sequence identity) in a reference genome. A DNA tag of the invention originates from a genomic DNA sequence. It may be linked to an ncRNA, or a cDNA of the ncRNA, through, for example, the DNA linker and RNA linker of the invention (or the modified RNA linker of the invention, or the direct RNA linker of the invention). An RNA tag of the invention originates from an ncRNA, or a cDNA reverse transcribed from the ncRNA. The RNA tag may be linked to genomic DNA through, for example, the DNA linker and RNA

linker of the invention (or the modified RNA linker of the invention, or the direct RNA linker of the invention).

The RNA or DNA tags of the invention can be of any size, but needs to be meaningful and advantageous over the size of the parental sequence from which it is derived. In certain embodiments, the size of a DNA or RNA tag is determined by genome complexity. For a bacterial genome, a tag from about 8 bp to about 16 bp may be sufficient, whereas for a complex genome like the human genome, a 16-20 bp tag may be considered.

“Linker” is usually an artificial sequence of nucleic acids designed for a specific purpose, such as linking two polynucleotides together. The “RNA linker” of the invention is designed to be linked to the DNA linker of the invention and to the cDNA synthesized from the free 3'-end of an RNA, such as a cross-linked non-coding RNA. The “DNA linker” of the invention is designed to be linked to the RNA linker of the invention and to a free end of a DNA, such as a chromosomal DNA cross-linked to a chromatin fragment. The “modified RNA linker” of the invention is designed to be linked to a genomic DNA fragment at one end (e.g., a blunt end or a ligation compatible end capable of generating a blunt end), and to the cDNA synthesized from the free 3'-end of an RNA, such as a cross-linked non-coding RNA, at the other end. The “direct RNA linker” of the invention is designed to be directly linked to the 3'-OH of ncRNA through a pre-adenylated 5'-end, and to be linked to genomic DNA fragment at the other end (e.g., a blunt end or a ligation compatible end capable of generating a blunt end).

“Sequencing” refers to the various methods used to determine the order of constituents in a biopolymer, in this case, a nucleic acid. Suitable sequencing techniques that can be used with the instant invention includes the traditional chain termination Sanger method, as well as the so-called next-generation (high throughput) sequencing available from a number of commercial sources, such as massively parallel signature sequencing (or MPSS, by Lynx Therapeutics / Solexa / Illumina), polony sequencing (Life Technologies), pyrosequencing or “454 sequencing” (454 Life Sciences / Roche Diagnostics), sequencing by ligation (SOLiD sequencing, by Applied Biosystems / Life Technologies), sequencing by synthesis (Solexa / Illumina), DNA nanoball sequencing, heliscope sequencing (Helicos Biosciences), ion semiconductor or Ion Torrent sequencing (Ion Torrent Systems Inc. / Life Technologies), and single-molecule real-time (SMRT) sequencing (Pacific Bio), *etc.* Numerous other high throughput sequencing methods are still being developed or perfected, with may also be used to sequence the PET polynucleotides of the invention, including

nanopore DNA sequencing, sequencing by hybridization, sequencing with mass spectrometry, microfluidic Sanger sequencing, transmission electron microscopy DNA sequencing, RNAP sequencing, and *In vitro* virus high-throughput sequencing, *etc.*

In certain embodiments, the sequencing method is capable of sequencing tags from both sides of the subject PET polynucleotides, thus providing paired end tag information. In certain embodiments, the sequencing method is capable of performing reads on long DNA fragments of variable length, such as concatemers of the subject PET polynucleotides.

“Reference genome” refers to the genome of the organism of interest, or the genome from which the ncRNA and genomic DNA originates. The method and compositions of the invention apply to any reference genomes for which a complete or substantially complete sequence is available, including numerous archaeal or eubacterial, protist, fungi (e.g., *S. cerevisiae* or *S. pombe*), plant, animal genomes. For example, the genome sequences of human, mouse and numerous other mammals and non-mammalian species are now readily available in the public domain. See, for example, Venter *et al.*, “The Sequence of the Human Genome,” *Science*, 291(5507):1304-1351, 2001. Other non-limiting reference genomes include those for numerous non-human primates, mammals, rodents (rats, mice, hamsters, rabbits, *etc.*), livestock animals (cattle, pigs, horses, sheep, goat), birds (chickens), reptiles, amphibians (*Xenopus*), fish (zebrafish (*Danio rerio*), puffer fish), insects (*Drosophila*, mosquito), nematodes, parasites, fungi (e.g., yeast, such as *S. cerevisiae* or *S. pombe*), various plants, virus (such as those integrated into a host genome), *etc.*

Locked nucleic acid (LNA) is a modified RNA nucleotide in which the ribose moiety of an LNA nucleotide is modified with an extra bridge connecting the 2' oxygen and 4' carbon. The bridge “locks” the ribose in the 3'-endo conformation. LNA nucleotides can be mixed with DNA or RNA residues in the oligonucleotide whenever desired. Such oligomers are synthesized chemically and are commercially available. The locked ribose conformation enhances base stacking and backbone pre-organization. This significantly increases the hybridization properties (melting temperature) of oligonucleotides.

Peptide nucleic acid (PNA) is an artificially synthesized polymer similar to DNA or RNA. PNA oligomers show greater specificity in binding to complementary DNAs, with a PNA/DNA base mismatch being more destabilizing than a similar mismatch in a DNA/DNA duplex. This binding strength and specificity also applies to PNA/RNA duplexes.

A “paired-end tag (PET) polynucleotide” of the invention is a polynucleotide that, at or near one end, an RNA tag originating from a ncRNA, and at or near the other end, a DNA

tag originating from a genomic DNA, wherein the ncRNA and the genomic DNA preferably are crosslinked to the same chromatin fragment. In that sense, the RNA and DNA tags at the two ends of the PET polynucleotide are paired, and reflects an event of physical proximity between the ncRNA and the genomic DNA at the time of crosslinking.

“Proximity ligation condition” refers to a condition for polynucleotide ligation reaction under which ligatable polynucleotide ends in close proximity, such as those genomic DNA and ncRNA crosslinked to the same chromatin fragment, are ligated preferentially. Meanwhile, ligatable polynucleotide ends not in close proximity, such as those genomic DNA and ncRNA crosslinked to different chromatin fragments, are not ligated or substantially not ligated. Such ligation condition include large volume ligation, such that ligatable ends on the same chromatin fragment, due to their physical proximity to one another, are much more likely to be ligated than ligation between ligatable ends on different chromatin fragments.

“Mapping (a sequence tag to a genome)” includes the identification of the genomic location of the sequence tag in the genome.

A “bifunctional crosslinking agent / reagent” or “crosslinking agent / reagent” includes modifying agents that possess two or more reactive groups, each is capable of reacting with one moiety (such as a DNA, an RNA, or a protein), thus crosslinking the two moieties together when the two moieties represent separate molecules. Such bifunctional crosslinkers are well known in the art (see, for example, Isalm and Dent in *Bioconjugation*, Chapter 5, pp. 218-363, Groves Dictionaries Inc., New York, 1999). For example, formaldehyde, glutaraldehyde or other similar reagents having aldehyde reactive groups may cross-link primary amino groups in proteins with other nearby nitrogen atoms in protein or DNA through a methylene (-CH₂-) linkage. Other bifunctional crosslinking agents that enable linkage via a thioether bond include *N*-succinimidyl-4-(*N*-maleimidomethyl)-cyclohexane-1-carboxylate (SMCC) to introduce maleimido groups, or with *N*-succinimidyl-4-(iodoacetyl)-aminobenzoate (SIAB) to introduce iodoacetyl groups. Other bifunctional crosslinking agents that introduce maleimido groups or haloacetyl groups on to a polypeptide are well known in the art (see US Patent Applications 2008/0050310, 2005/0169933, available from Pierce Biotechnology Inc. P.O. Box 117, Rockland, IL 61105, USA) and include, but not limited to, bis-maleimidopolyethyleneglycol (BMPEO), BM(PEO)₂, BM(PEO)₃, *N*-(β-maleimidopropoxy)succinimide ester (BMPS), γ-maleimidobutyric acid *N*-succinimidyl ester (GMBS), ε-maleimidocaproic acid *N*-hydroxysuccinimide ester

(EMCS), 5-maleimidovaleric acid NHS, HBVS, N-succinimidyl-4-(N-maleimidomethyl)-cyclohexane-1-carboxy-(6-amidocaproate), which is a “long chain” analog of SMCC (LC-SMCC), m-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS), 4-(4-N-maleimidophenyl)-butyric acid hydrazide or HCl salt (MPBH), N-succinimidyl 3-(bromoacetamido)propionate (SBAP), N-succinimidyl iodoacetate (SIA), κ -maleimidoundecanoic acid N-succinimidyl ester (KMUA), N-succinimidyl 4-(p-maleimidophenyl)-butyrate (SMPB), succinimidyl-6-(β -maleimidopropionamido)hexanoate (SMPH), succinimidyl-(4-vinylsulfonyl)benzoate (SVSB), dithiobis-maleimidoethane (DTME), 1,4-bis-maleimidobutane (BMB), 1,4 bismaleimidyl-2,3-dihydroxybutane (BMDDB), bis-maleimidoethane (BMH), bis-maleimidoethane (BMOE), sulfosuccinimidyl 4-(N-maleimido-methyl)cyclohexane-1-carboxylate (sulfo-SMCC), sulfosuccinimidyl(4-iodoacetyl)aminobenzoate (sulfo-SIAB), m-maleimidobenzoyl-N-hydroxysulfosuccinimide ester (sulfo-MBS), N-(γ -maleimidobutyloxy)sulfosuccinimide ester (sulfo-GMBS), N-(ϵ -maleimidocaproyloxy)sulfosuccinimide ester (sulfo-EMCS), N-(κ -maleimidoundecanoyloxy)sulfosuccinimide ester (sulfo-KMUS), and sulfosuccinimidyl 4-(p-maleimidophenyl)butyrate (sulfo-SMPB).

Heterobifunctional crosslinking agents that may be used for crosslinking may contain an amine-reactive *N*-hydroxysuccinimide group (NHS group), and/or a carbonyl-reactive hydrazine group. Examples of such commercially available heterobifunctional crosslinking agents include succinimidyl 6-hydrazinonicotinamide acetone hydrazone (SANH), succinimidyl 4-hydrazidoterephthalate hydrochloride (SHTH) and succinimidyl hydrazinium nicotinate hydrochloride (SHNH). Conjugates bearing an acid-labile linkage can also be prepared using a hydrazine-bearing benzodiazepine derivative of the present invention. Examples of bifunctional crosslinking agents that can be used include succinimidyl-p-formyl benzoate (SFB) and succinimidyl-p-formylphenoxyacetate (SFPA).

Other bifunctional crosslinking agents that enable crosslinking via disulfide bonds are known in the art and include *N*-succinimidyl-3-(2-pyridyldithio)propionate (SPDP), *N*-succinimidyl-4-(2-pyridyldithio)pentanoate (SPP), *N*-succinimidyl-4-(2-pyridyldithio)butanoate (SPDB), *N*-succinimidyl-4-(2-pyridyldithio)2-sulfo butanoate (sulfo-SPDB) to introduce dithiopyridyl groups. Other bifunctional crosslinking agents that can be used to introduce disulfide groups are known in the art and are disclosed in U.S. Patents 6,913,748, 6,716,821 and US Patent Publications 2009/0274713 and 2010/0129314, all of which are incorporated herein by reference. Alternatively, crosslinking agents such as 2-

iminothiolane, homocysteine thiolactone or S-acetylsuccinic anhydride that introduce thiol groups can also be used.

Two or more of the above bifunctional crosslinking reagents may be used together to crosslink DNA, RNA, and protein in a chromatin fragment.

3. *Restriction Enzymes*

It is not required that the DNA and/or RNA linkers of the invention comprise restriction enzyme recognition sites. Indeed, in certain embodiments, it may even be desired that the DNA and/or RNA linkers of the invention comprise no restriction enzyme recognition sites. However, in certain embodiments, the DNA and/or the RNA linkers of the invention may comprise at least one RE recognition site, such as a Type II RE recognition site (*e.g.*, Type IIS RE site).

In general, any RE and their recognition sites known in the art may be used, if the result of the RE cleavage produces a DNA or RNA tag of desired length, such as 10-20 bp. Such restriction enzymes recognizing at least one recognition site within the nucleic acid molecule and which may be used with the instant invention will be evident to those skilled in the art, particularly in view of the guidance provided herein and the illustrative examples. See, for example, *Current Protocols in Molecular Biology*, Vol. 2, 1995, Ed. Ausubel, *et al.*, Greene Publish. Assoc. & Wiley Interscience, Unit 3.1.15; and the most up-to-date New England Biolabs Catalog or website information, 2005 and beyond.

A non-exclusive list of possible restriction enzyme recognition sites and the corresponding restriction enzymes recognizing the same is reported below.

As an example, a Type IIS RE, such as MmeI, may be used to generate a fixed length DNA or RNA tag that flanks the ligated RNA-DNA linkers. In particular, an MmeI recognition site may be placed at the end of the double-stranded region of the RNA or DNA linker, such that, upon MmeI cleavage, an 17-21 bp tag sequence originating from the RNA or DNA sequence is linked to the now ligated RNA linker and DNA linker. If one MmeI site appears in each of the RNA and DNA linkers, the two generated tags - one being a DNA tag, another being an RNA tag - flanks the now ligated RNA linker and DNA linker. The two tags may be additionally processed by blunting such that further downstream operation, such as PCR amplification, concatenation, or sequencing, may be performed.

Examples of some non-exhaustive Type II restriction enzymes that may be used with

the instant invention include: AarI, AceIII, AoiI, BaeI, Bbr7I, BbvI, BbvII, BccI, Bce83I, BceAI, BceFI, BcgI, BciVI, BfiI, BinI, BplI, BsaXI, BscAI, BseMII, BseRI, BsgI, BsmI, BsmAI, BsmFI, Bsp24I, BspCNI, BspMI, BsrI, BsrDI, BstF5I, BtgZI, BtsI, CjeI, CjePI, EciI, Eco31I, Eco57I, Eco57MI, Esp3I, Fall, FauI, FokI, GsuI, HaeIV, HgaI, Hin4I, HphI, HpyAV, Ksp632I, MboII, MlyI, MmeI, MnlI, PleI, PpiI, PsrI, RleAI, SapI, SfaNI, SspD5I, Sth132I, StsI, TaqII, TspDTI, TspGWI, TspRI and Tth111II (see the list in the web site of Rebase Enzymes: [rebase dot neb dot com slash cgi-bin slash outsidelist](http://rebase.neb.com/cgi-bin/outsidelist); see also Szybalski, W., 1985, *Gene*, 40:169). Other suitable RE enzymes known in the art or those later discovered, which have the similar property of being able to generate a tag sequence of desired length (*e.g.*, 10-25 bp to hundreds of bps) may also be used to practice the present invention.

In certain embodiments, the restriction enzyme is a Type IIS enzyme. In certain embodiments, the RE produces a DNA or an RNA tag sequence of about 10-25 bp or 15-20 bp. In certain embodiments, the RE is MmeI or GsuI.

Other examples of recognition sites and cleavage sites of several class II restriction enzymes include (into parenthesis are the recognition site and the cleavage site): BbvI (GCAGC 8/12), HgaI (GACGC 5/10), BsmFI (GGGAC 10/14) SfaNI (GCATC 5/9), and Bsp I (ACCTGC 4/8).

Artificial restriction endonucleases may also be used. These endonucleases may be prepared by protein engineering. For example, the endonuclease FokI has been engineered by insertions so that it cleaves one nucleotide further away from its recognition site on both strands of the DNA substrates. See Li and Chandrasegaran, *Proc. Nat. Acad. Sciences USA*, 90:2764-8, 1993. Such techniques may be applied to prepare restriction endonucleases with desirable recognition sequences and desirable distances from recognition site to cleavage site.

Thus in certain embodiments, the RE enzymes that may be useful for the composition and methods of the invention includes artificial restriction endonucleases, such as those capable of generating Type IIS type cleavage fragments outside the recognition sites. In certain other embodiments, however, the RE enzymes that may be useful for the composition and methods of the invention excludes artificial restriction endonucleases.

In certain embodiments, Type IIB restriction enzyme recognition sites may be incorporated into the design DNA and/or RNA linkers. Type IIB restriction enzymes (*e.g.*, BcgI and BplI) are multimers requiring both AdoMet and Mg^{2+} cofactors, and they cleave DNA on both sides of their recognition to cut out the recognition site. Thus, a Type IIB RE

site may be engineered to span or straddle the linked RNA and DNA linkers (*e.g.*, part of the RE site is on the RNA linker, and the remaining part of the RE site is on the DNA linker, such that the ligated DNA and RNA linkers reconstitute a complete Type IIB RE site), or completely within the RNA linker or the DNA linker. Upon digestion with the Type IIB RE, both the RNA and DNA tags can be generated.

In certain embodiments, Type IIG RE (such as *AcuI*) recognition sites may be used instead of the Type IIS RE sites. Such Type IIG RE recognize continuous sequences and cleave on just one side (*AcuI*).

A list of all suitable Type II RE recognition sites, *e.g.*, Type II RE that cleaves outside its recognition sequence on one or both sides, may be obtained from various sources. See, for example, *Restriction Endonucleases (Nucleic Acids and Molecular Biology)*, edited by A. Pingoud, Springer; 2004 edition (December 1, 2004), incorporated herein by reference. Also see, New England Biolabs' 2010 catalog and subsequent updates (incorporated herein by reference).

In certain embodiments, Type I restriction enzymes may also be used to generate RNA or DNA tags, particularly DNA tags. For example, the Type I RE recognition sites may be included in the DNA linker such that the RE cuts at a random distance in the linked chromosomal DNA.

In certain embodiments, Type III RE recognition sites (*e.g.*, *EcoP15I* site) may be used in the RNA and/or DNA linkers. Type III RE enzymes cleave outside of their recognition sequences and require two such sequences in opposite orientations within the same DNA molecule to accomplish cleavage. The two required recognition site for each cleavage may be contained completely within the DNA linker, or completely within the RNA linker, or in both linkers (such that only correctly linked RNA-DNA linkers regenerate the RE recognition site).

Examples of Type III restriction site(s) and Type III enzyme(s) have been described in, for example, Matsumura *et al.*, *SuperSAGE, Proc. Natl. Acad. Sci., USA* 100(26):15718-23 (Dec. 2003; Moencke-Buchner *et al.*, *J. Biotechnol.*, 114: 99-106, 2004; Mucke *et al.*, *J. Mol. Biol.*, 312: 687-698, 2001; Rao *et al.*, *J. Mol. Biol.*, 209: 599-606, 1989; Hadi *et al.*, *J. Mol. Biol.*, 134: 655-666, 1979, all incorporated herein by reference. Type III restriction enzymes can also be purchased from New England Biolabs (NEB). In particular, an exemplary Type III RE for carrying out an embodiment of the present invention is the type III enzyme *EcoP15I*. The recognition site(s) of *EcoP15I* is CAGCAG (25/27).

Any of the above restriction sites may be used together in the DNA or RNA linkers. For example, the RNA linker may comprise a Type IIS RE site, and the corresponding DNA linker may have no RE site, a Type IIG site, or a Type III RE site, *etc.*

4. *Concatemers and Libraries*

In certain embodiments, the isolated PET polynucleotides of the present invention may be joined or concatenated with other isolated PET polynucleotides to form a concatemer of PET polynucleotides. Any number of PET polynucleotides may be joined together for the purposes of sequencing or for cloning into a suitable plasmid or vector.

Accordingly, in another aspect, the present invention provides a concatemer of PET polynucleotides comprising at least two PET polynucleotides, each comprising at least a DNA tag and at least one RNA tag, wherein the DNA tag is obtained from a chromosomal or genomic DNA and the RNA tag obtained from a cDNA of a ncRNA, wherein the DNA and the cDNA of the ncRNA are obtained from a cross-linked nucleic acid-protein complex, using the RNA/DNA linkers and methods of the invention.

Each PET polynucleotide of the concatemer of PET polynucleotides may thus have the general structure of RNA tag-RNA linker-DNA linker-DNA tag (or the reverse orientation).

The concatemers may be formed by any of many art recognized methods. In particular, the length controlled concatenation method (Ruan *et al.*, U.S. patent application publication US 2008/0124707 A1, incorporated herein by reference) may be used. In another example, the isolated PET polynucleotides may be polished at both ends, if necessary, before the ends are linked to one or more adapter oligonucleotide(s) that can be digested by a (Type II) restriction enzyme. The digestion products may have compatible sticky ends that can facilitate the concatemerization of the individual PET polynucleotides. If the RE sites are the same for all the adapters linked to the ends of the PET polynucleotides, all sticky ends are compatible for ligation and concatemerization, and the individual PET polynucleotides may be independently linked together either in head-to-tail manner or head-to-head manner. If the adapters are different, for example, a first adapter having a first RE site may be linked to the RNA tag, while a second adapter having a second (different) RE site may be linked to the DNA tag. Upon concatemerization, all PET polynucleotides will be linked in a head-to-head manner.

Thus each PET polynucleotide of the concatemer of PET polynucleotides may be independently linked to one (for the terminal PET polynucleotides) or two (for the internal PET polynucleotides) another PET polynucleotide in a head-to-tail or head-to-head manner. In certain embodiments, all PET polynucleotides within the concatemer are linked in a head-to-head manner.

The DNA and/or RNA linkers of the PET polynucleotides may comprise at least one restriction enzyme recognition site, such as an RE recognition site for a type IIS restriction enzyme (*e.g.*, MmeI or GsuI).

The concatemer of PET polynucleotides may be inserted into or cloned in a vector or a cell; the cell may be a bacterial cell. The cloned concatemer of PET polynucleotides may be digested by RE and isolated individually if desired.

It will be apparent that the number of PET polynucleotides of the present invention that may be concatenated depends on the length of the PET polynucleotides, which may be readily determined by those of skilled in the art without undue experimentation. After formation of concatemers, multiple tags may be cloned into a vector for sequence analysis, or the concatemers may be directly sequenced without cloning by methods known to those of skill in the art, such as by any of the so called next-generation high throughput sequencing methods described herein or known in the art, including single molecule sequencing methods. Hence, the concatenation of the PET polynucleotides allows an efficient analysis of the nucleic acid molecules in a serial manner by sequencing multiple PET polynucleotides within a single vector or clone.

In a related aspect, the present invention provides a library of PET polynucleotides comprising at least two PET polynucleotides, each comprising at least a DNA tag and at least one RNA tag, wherein the DNA tag is obtained from a chromosomal or genomic DNA and the RNA tag obtained from a cDNA of a ncRNA, wherein the DNA and the cDNA of the ncRNA are obtained from a cross-linked nucleic acid-protein complex, using the RNA/DNA linkers and methods of the invention.

In certain embodiments, the library may comprise up to 10 million PET polynucleotides, or up to 1 million, 100 thousand, 10 thousand, 1 thousand, 100 hundred, or 10 PET polynucleotides.

In certain embodiments, the library has not been through any amplification, such as PCR amplification.

In certain embodiments, the library has been amplified, such that at least two members within the library originate from amplification, such as PCR amplification, rolling circle amplification, biological amplification of cloned genetic materials, or any other known amplification methods. The PCR primers and the probes sequences may be prepared based on the information of the PCR adapters linked to the end of the PET polynucleotides, or based on the primer sequences on the cloning vector flanking the cloned PET polynucleotides or concatemers thereof.

The PCR or other amplification products that contain the PET polynucleotides may then be isolated with an enzyme recognizing the flanking RE restriction site (inside the adaptors) to give rise to the amplified library, which may be used for any of many downstream analysis.

In certain embodiments, the PET polynucleotide concatemers, before or after amplification, may be selected for suitable sizes, by any standard method, including gel electrophoresis and gel excision. The main considerations in selection for the appropriate sizes are that the sizes should be above the size of primer dimers and unannealed adapters and below the sizes of certain long linear multimers. In particular, concatemers with sizes of approximately 100-1000 bp, or 200-500 bp may be selected. Accordingly, with size selection, an advantage is that long linear multimers may be eliminated as their sizes will be above the size range. Similarly, fragments that are too short, unannealed adapters and primer dimers may also be eliminated.

5. *Chromatin Immunoprecipitation (ChIP)*

In certain embodiments, the methods of the invention may be used to identify specific ncRNA-chromatin/protein-DNA interaction. For example, in certain embodiments, it may be of interest to determine any ncRNA-DNA-chromatin interaction associated with a particular chromatin component or protein. The methods of the invention may further comprising using ChIP to immunoprecipitate the protein of interest.

ChIP has been used to enrich and thereby allow the identification of genomic regions associated with specific proteins such as histones and other proteins binding to nucleic acids in nucleic-acid protein complexes (reviewed in Taverner *et al.*, *Genome Biol.*, 2004, 5(3):210). The aim is to cross-link proteins with DNA at their sites of interaction.

This may be accomplished quickly and efficiently by adding a suitable fixative such as formaldehyde, paraformaldehyde, glutaraldehyde, acetone, methanol, or other bifunctional

crosslinking reagents (or mixtures thereof) directly to living cells in culture. Crude extracts of these fixed cells are then prepared, and the chromatin fragmented according to the methods of the invention. For example, fragmentation may be achieved either by physical sheering (*e.g.*, shearing by sonication, hydroshearing, repeated drawing through a hypodermic syringe needle), or by enzymatic digestion (such as restriction enzyme digestion, or digestion with endonuclease with controlled timing, enzyme concentration, temperature, pH, *etc.*) so as to achieve a desired average size (*e.g.*, usually about 1 kb). The cross-linked and sheered chromatin fragments are then used in immunoprecipitation reactions with antibodies raised against the specific protein of interest (*e.g.* transcription factors or histones). Crosslinked ncRNA and DNA fragments enriched in each immunoprecipitation are subsequently linked using the DNA and RNA linkers of the invention through proximity ligation, then de-linked or reverse cross-linked from the protein components (*e.g.*, through heat and/or Protease K digestion), and purified to allow their identification by the methods of the invention.

The advantage of using ChIP is that this approach is able to “freeze” the ncRNA or gene regulatory network in live cells, as such interactions exist in their natural states, by rapid cross-linking of chromatin and other non-histone proteins, thereby in theory representing a “true” picture of the specific ncRNA or gene regulatory system at any point in time, free of potential artifacts imposed by heterologous expression, for instance.

6. *Applications*

The methods and compositions of the invention allow one to identify interaction between ncRNA and genomic loci, either at a non-biased global level, or at the level of specific ncRNA or specific chromatin components of interest. Information obtained using the instant methods can be used in a wide variety of research and development settings.

For example, the invention provides a method to identify chromatin targets of a specific ncRNA, which may previously have unknown or incompletely understood function, the method comprising determining interaction between the specific ncRNA and its genomic target sequences using the methods and compositions of the invention. The identified genomic target sequences represent candidate targets upon which the ncRNA exerts its biological function.

In a related aspect, the invention provides a method to identify ncRNAs that interact with a specific gene or genomic region, such as gene or genomic region harboring a tumor suppressor gene or an oncogene, the method comprising determining interaction between the

specific gene or genomic region and ncRNAs of the genome using the methods and compositions of the invention. The identified ncRNAs represent candidate modulators (e.g., suppressors, enhancers or co-activators) of gene function.

In certain embodiments, the method further comprises comparing the presence / absence or the extent of the interaction between the ncRNA and the gene / genomic region, among two or more samples. Such comparison may help to further decipher the biological significance of the interaction and any observed differences between the samples.

For example, one of the samples may be a healthy control sample, and the other samples may be disease samples, such as disease samples from animal models (e.g., mouse or rat models); disease samples before and after a particular treatment; disease samples over different stages of treatment; disease samples from patients who have responded to a particular treatment, or patients who are resistant to a treatment, or patients who has relapsed after a treatment.

In certain embodiments, one of the samples is a stem cell or induced pluripotent stem (iPS) cell derived from the patient, and, optionally, the other samples may be cell lines differentiated from such stem cells or iPS cells. Here, a specific ncRNA-chromatin interaction may be associated with the initiation of a developmental or differentiation program.

In certain embodiments, the sample(s) may be from a human, a non-human primate / mammal, a livestock animal (cattle, horse, pig, sheep, goat, chicken, camel, donkey, cat, and dog), a mammalian model organism (mouse, rat, hamster, guinea pig, rabbit or other rodents), an amphibian (e.g., *Xenopus*), fish (e.g., zebrafish), an insect (*Drosophila*), a nematode (e.g., *C. elegans*), a plant, an algae, a fungus (yeast, such as *S. cerevisiae* or *S. pombe*). The sample(s) may be a tissue culture of established cell lines, cultured primary cells, tissue biopsies (freshly dissected or frozen), etc.

As shown in Example 9, the methods of the invention identified an ncRNA - CCAT1 (Colon Cancer Associated Transcript 1) - as having a very complicated transcript isoform structures in this locus. The RICH-PET data provides important insights of potential function and underlying mechanism of CCAT1. Specifically, it was found that CCAT1 locus itself has significant enhancer features, that CCAT1 locus is highly transcribed in cervical cancer cell line HeLa cells, and the RICH-PET data shows that the transcribed product from this locus targets other enhancer and promoter regions. For example, for 122 loci targeted by CCAT1 ncRNA transcript (each with ≥ 3 RNA tags), 88 loci are enhancer regions, including

six enhancer loci with RNAPII interaction. Another 34 loci are within promoter regions. This is consistent with the observation that CCAT1 target genes on average are more highly expressed than randomly selected groups of genes. Thus the lncRNA CCAT1 may act as a transcription co-factor to activate a network of genes, including the oncogene *c-myc*.

Thus another aspect of the invention provides a method to treat a cancer expressing CCAT1, the method comprising administering an antagonist of the CCAT1-encoded lncRNA.

In a related aspect, the invention provides a method to disrupt transcription activation or co-activation mediated by a gene product of CCAT1 (e.g., a transcribed lncRNA), comprising contacting the gene product with an antagonist of the CCAT1-encoded lncRNA. In certain embodiments, the transcription activation or co-activation occurs in a cancer cell. In certain embodiments, the transcription activation or co-activation is for *c-myc*, FAN84B, and/or SNX14. In certain embodiments, the transcription activation or co-activation is effected by bringing the CCAT1 genomic locus to physical proximity of a target gene locus.

In certain embodiments, the cancer is a colon cancer (e.g., adenocarcinoma of the colon), a rectal cancer, a cervical cancer, a lung cancer, a gastric carcinoma, a liver cancer, and a metastase thereof. In certain embodiments, the cancer expresses CCAT1 transcript at a level that is 2-fold, 3-, 5-, 10-, 15-, 20-, 30-, 40-, 50-, 60-, 70-, 80-, 90-, 100-, 120-, 150-, 175-, 200-, 250-, 300-, 500-, 1000-fold higher compared to a matching or control sample.

In certain embodiments, the antagonist is an antisense polynucleotide that may optionally comprise modified nucleotides to, for example, improve serum stability, pharmacological or pharmacokinetic properties, etc. The modified nucleotide may comprise PNA, LNA, 2'-O-alkyl or other 2' modifications, and/or modifications on the sugar-phosphate backbone.

In certain embodiments, the antagonist is an siRNA or miRNA construct that targets the encoded CCAT1 lncRNA.

The invention also provides an antagonist of the CCAT1 lncRNA (antisense, siRNA, miRNA, or vector encoding / expressing the same).

In another aspect, the invention provides a method for drug screening, the method comprises establishing a statistically significant association or correlation between drug efficacy and a specific observed ncRNA-chromatin interaction identified by the methods of

the invention (such as an interaction identified in a responsive patient but not in a resistant patient), determining the effect of a plurality of candidate drugs on the statistically significant association or correlation, and identifying candidate drugs that promote the statistically significant association or correlation.

In certain embodiments, effects of the candidate drugs are tested using samples from the resistant patient. This may allow the identification of candidate drugs that restores the statistically significant association in the resistant patient.

In another aspect, the invention provide a method to identify a target gene for treating a disease, the method comprising: (1) using the methods of the invention, identifying (from among the observed ncRNA - genomic DNA interactions), a statistically significantly association between the efficacy of a drug and a particular ncRNA- genomic DNA (gene) interaction (*e.g.*, whenever efficacy is observed in a patient responsive to treatment, a particular ncRNA-genomic DNA (gene) interaction(s) is observed; whenever efficacy is not observed in a patient not responsive to treatment, the particular ncRNA-genomic DNA (gene) interaction(s) is not observed), (2) determining the expression level of the involved ncRNA and/or the DNA (gene); wherein the DNA (gene) is identified as a potential target gene for treating the disease when drug efficacy is associated with increased ncRNA expression and inhibition of DNA (gene) expression.

The compositions and methods of the invention can also be used to identify as yet unknown ncRNA in a particular genome, since the method of the invention is an unbiased approach for identifying such ncRNAs. If a cluster of PET polynucleotides consistently identify a cluster of RNA tags in one region of genome that does not encode any protein, and consistently link these RNA tags to a (remote, *e.g.*, an interchromosomal) locus that is represented by the corresponding DNA tags, it is likely that the RNA tags reveal an ncRNA.

Any candidate therapeutic reagents or target genes identified by the screening methods of the invention can be validated *in vitro* and/or *in vivo*, using well-known experimental models correlating to a disease or condition. For example, if a particular ncRNA is identified as promoting the expression of an oncogene (or as inhibiting the expression of a tumor suppressor gene), thus becoming a candidate drug target, a potential therapy using antagonists of the ncRNA, such as siRNA, miRNA, antisense, *etc.*, may be further validated *in vitro* and/or *in vivo*, the latter may be carried out in an established cancer

model, e.g., in a model animal, such as a mouse model of the cancer to be treated.

The mouse is a well-established model for drug discovery and development, with many different strains available. For example, a large number of useful models for studying cancer can be found at Mouse Models of Human Cancers Consortium, which has developed several databases, e.g., Emice (emice dot nci dot nih dot gov), Cancer Models Database (cancermodels dot nci dot nih dot gov) and Cancer Images Database (cancerimages dot nci dot nih dot gov), or other resource such as cancer research models distributed via The Jackson Laboratory (see jaxmice dot jax dot org slash list slash rax3 dot html). Further xenograft models, using either primary cancer biopsies or cell lines, are useful to investigate cancer.

For example, to develop a lung cancer model in which the efficacy of a potential antagonist against a candidate ncRNA can be verified, six to eight 8-week-old female immunodeficient mice, such as CB17-SCID beige mice (Taconic, cat. no. CBSCBG) or NOD/SCID (The Jackson Laboratory cat. 001303) or NOD SCID Gamma mice, also known as NSG (The Jackson Laboratory cat. 5557) are injected either subcutaneously or transthoracically (orthotopic; 10^4 /sup cells/25 μ L) via the left lung with human lung carcinoma A549 cells (ATCC® CCL-185). Tumor-bearing mice are intraperitoneally injected with neutralizing anti-CXCL12 or preimmune serum, or receive no treatment. Alternatively, tumor-bearing mice may be treated with Platinol (Cisplatin) or Abitrexate (Methotrexate) or Paclitaxel, or other compounds. Tumors are isolated at various time points, treated and untreated. Noncoding RNAs are identified according to the method described previously.

7. *CCAT1 Transcripts, Antagonists, and Uses Thereof*

In another aspect, the invention provides various CCAT1 transcripts identified by the methods of the invention, their cDNA sequences (both strands), antagonist (e.g., antisense sequences, siRNA or miRNA constructs that antagonizes the function of these CCAT1 ncRNA transcripts.

The eight identified cDNA sequences representing different isoforms of the CCAT1 ncRNA are provided below in SEQ ID NOs: 1-8.

```
>CCAT1_JAX_1 Transcript sequence; Genomic location:  
chr8:128128655-128241571 strand:-  
<128241571>
```

3' -

TATAGGTATAACCAATATATATGTATAACATATATATGTCCATATATATGTATAACCAAACC
ACAGGTGTTTTTTTTTGGAAAGTCATATTATACAGGGAGTTGACAGAGGTGTGAGCTGGACTTTA
AGAAGCTGCACATAAGATGCTAGTATGATCAAGCTGGAATGGACTTAGACAATTTGAAACAA
CTTTTCTCAGTTTTTCAGATGAGGAACTGACGGGTACCAAGCTTAAATGACTTGACGAAGCT
CATAGAAGATTAGCAGGTAGTAGAATAATGACTGCTGACTCCTAATTCAGTGGATCTTCCCT
GGCCACCGTTTTTGTATTGAGCTGCAATGCTTCCTTGACTGTTCTCCATGCCAGATTCTTATC
AATGATCTTTTACCTAAGAAACAGCAAAGATTCTGGCAAGCACACGATCTAGAGATACATCT
TATTGCGATTTTTTCACAAAAATCAAAAGAAGAAAGAAGGCTTAGCTGGTGTTAATTATTGT
TATTTTTTTTCAATAGGGAAATCTGTACACAATGATTTATCTCCAGTGATTTGCCATTGATCA
ATTTTTTTTCTCATTTTCATTTTCTATTTTTTTGTTTTTTGTTTTTCTTTATTTTTTATTTTT
TCTCCTTTTTCTTTTTTTTAAATTTTCTGTTTATCACAAATGATCATGTAATTATATGTTAAT
ACTATGTAACCCCAAGTGTTTTCAACTGTTTGTGATTCAATGTTACCCAGTTTTCTTTTTCTTA
ATTTTAAATAAATTTGAAAAATTATCTTAGAGTGTTTTGAGCCTGTGTTGGTACATTTAGTT
CTAGTTCATTGTGGTAAATCCACTTCAGTTTCTAAGTTTTCCACCCTTTAGTAAAGACATATT
TCTAAATTTGGTTTTATATCCTCAGTTACAAAAGATTCTAACTGCTAGTTTTTGTGACAGCAAT
CACACATACCATCAAGGGATGGGGCAGGCAGTTTTGGAATGTGCTGATGCTAGCATTTTTTAT
AAGCCTATGGCTTTTTATAGTCTAAATTGTTCTTATTTCTATTAATGCTTCAATTTTTTGACAA
ACACACAACCATAGAAACAAATAATAAATTTGTTTTTTGGGAATTATCCAGGATTCCTGGTT
AAGTGGGGAATTTAGGCTTTGACAGCATAAAGGATCACGGATAAGTATTTTTTACGGTGGCT
CTAATTATAATGCTGAGCATGTGGCAGGCACAGAAATATTTACTCATTGACTGAATATAGCA
CATCGTAATGTTGATTTTTTTTCCAACATAATTTTAGAGCTAGGCATATTGTATTCTATTACA
CTAGACTATATATCATTCTTAAATAGAACCAGCCTTGCTAGATAACACATGTTGGAGGAGAG
GCCCTTCTTCTTAGCCCTCAGTGTTTCCATCTATGGGGAAGAAGTTCCACCATACTAACATT
ACTATCGTCTCTCCACCTGCTCACTCACTTCTCCCCAAGGGAGGGGTGTTTCGATATGGTTTC
TGAGCTTGGAAGAAACTCAGGCATGTGTAACATGGTTCCTTCAGTCCCATGACCCACTGT
CCACAAATGGGCTGCTCACAGAGTGCATGCCCTTACCCTTGTTCTTGGCCATGCAGGAAATT
GTATGAAACAGTCCTAGCTGAAGCCTGAGATTTTCTGCTGATTGCCTAGTCCTGGTGGGTATC
TGTCTACTCCTGGAGTTTGGATTGGAAAGTCCACATGCCTGAAGGTATAAACCTATTCTACA
AAGGGGTGTTTTCTAGAATGAAGGTAATATTTTTATCTTACATTTGCAGAAAGAGACAGAAC
AATGTTATAGGTGAGTGCATGGACAATGACCTCAAACAGCTAAGATTCAAACCCCTGCGTTG
AATGATTGAATTGAAATGATTCAATGAGCTAATGTACATAAAGCATCCAGAATGTTGCCTGG
CACAAGGGACTGTATTGTCTGCTAGACCATTTATTCAAAGTGGGAGGATGATGTTCTAAAAG
CCAATGATAAAGCTCATGGCAATGCAGGGTATATCTGATGGCATGGAATGCTTTAGGATGGC
CAAGATTGCCCATCAAATGCCAAGTCACCGAAGGGTTCTGGGAAATAAGAAATCATTAGAAA
AAGATTTCTGCCTTCTAGAAGTACACAGTCTAATGGTGAGATAGGCAGGTTATTAATGGCTC
TCCTACTAGGAGCCTGACATCATGGTGAGCATCGAGAAAGGAATAACCTAAGCTGAAGACAC
GCCTTTTTCAGGAGGGCCAAGTTCCACGTTCTGTGCATGCTTTTGGCGAAAGTCAGGTAAAGCC
TCCAGAAATGCATGGTCTATTCTCTCGGACCATATGGCTGTGGGCAAACCTCTGGCTTCTTGG
AGCCTCCATTAGCCACCTAGGGAAAGGTAATTGGCTTTATGTTTGGCTCCATCACTGCTGGC
GACAAGCTCCACTGCATTCTCAAGCAGTAGATGAAATACAACCTGTGCTCCTGAATGCAGCAA
AGGAAAGAGAACCCAACTGGACCAGTGTGAAATTGGGACCTTTGGAGGCAAGAATGCACATA
TTTATAAATGAATAAACATAATTTTTCTCTCTTCTTCCCATTGGCAGCATATCTGAAGGCCC
TGACTCAACAGTAGTGGGTCTTAATGCTTGTGCTTGTCCACCCTTCTCGGCAGTGATTGATT
TCATTGTTGGCTTGTTTTTCATGACTTGAATGTGCGATCTTCAGAGGGCCTAATTCCTTGCAA
GTTCCATTTGAGGGTGAAGAGCTAGAACAAAGCTATAACCAGTAGACAACCTCCGTGACTCAG
GAGCTTAAGCATGTGACTAATTAGTAAGAAAAAATGTGGTGAAGATTTGTAGTTAATAAGAA
GGAAAGAAGAATCACTGGGGCTAGAATTATGCAAGCTTTTGTTCCTTTTGGGACTATATCA
GAACTATGAGAGAAGAAAGGCTACCTTTTACCTTTGAGGAATTTTCAAAGCCTTTTTTTTTT
TTTTTGGTTGGTTTGTGACTGACAAAGGGCACAAATTTCAACACCTCAGAAAAATGCCTCATC
ATTTCTCTTGTGAAATCTGGGTGCTTCTGAATGAATCCATGTTAGGAATGTGTACTTCCAT

CCATTAAAGTCAATGTCCAGTCTCATT TTTGGGCCAGAGGCTGTTACTAAGTTGTAGTACTGG
 TGAGAAAAGGACAGAGCATTTACCTTCCCTGGGTATCCTGTGACCTACATGTGTCCTTGCAA
 GGCAGGGAAATGTTACTAATTAAGAGCATAGCCTTTTCGTGAGATGGGTTTCAGATCCTAACAC
 ATCCTCTTTCTGGTTACAGGACAGTCTCTCTGAGCCTCAGTTTTCTCATCTCTAAGATGAG
 GCTAATTATACCTACTTCCCAGGGGGGTACTAAGGATAAAGTAACAACACAAGAAAATTTTT
 TATCCTTGATTTTTCTTAGCTGGAAAATTAGGCTAGAAAGACCTCCATCCTTGGGGTATTTAA
 AATAAAAGGACAAACATATGATGAGTCTAAGTGATCAATACATTGTGGCTTTTGTACTTACT
 ATTCTGAAACATGGGTGTGGCCTAGATACTTTCCAAAATTCTGCCCTCTCCACCTAGCAAAA
 TGACAACCAGACTTACAGATATGCCAACAAAGCTAGTGTTAATAATGGTGTTTGATGATAAA
 TGGCATTCTTTTCGACATTTGTCTCTTTTTTAAAACCTTGGTGCTCTTAAATGCATCATTGGA
 TAAGTGATGACTGTTCCCATTCGCAAGAAGAGACCTGGAATCTAAGCATGAAGGACCTGTCC
 TGATGTTGAGAAGTGTTGATCACATTTATTTGTAACTTAGCTTCTTGCTCGTCTCATGGTT
 GCTTTTTTTTTCTTTTTCTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTGAGAGG
 GAGTCTCACTCTGTCACCCAGGCTGGAGTGCAGTGGTGATCCCGGCTCACTGCAACCTCC
 ACCTCCCGGTTTCAAGCAATTCTCCTGCTTCAGCCTCCCCAGAAGCTGGGATTACAGGCACG
 TGCCAAGGATTTAATGGCAAGATGCCATTATAGACAAGAACAGGATTTCAGACACTTTTCGTGT
 TATGTGTTCTTGTCCAAATACTGTGTCACTCTGCCTGGCATCCCAATCCCACCAACACCTT
 TAGCAGGAACCTCCTGTTTCAATAACATTTCTCAATACTCTACCTGATTGCTTAGAATCCAT
 GAGACAGCCATATTCTCCATGTCTAGGTCCCTATCTTATTTTTGTTGAGATGGTGTTAAGAC
 TTTATTTGTGAGGCTTCTGGGGAAGAGGAGTAAGGTATTGATCCCACTGACTGGATAATTTG
 GGTCTCAAAATGGATAATAAATAAGCATTACATATTTTGACCACTTCTTGGAGGAGAACTT
 CTTGGAATGTGCACCATGTCCGCTGCACTTTTTTTTGCACAGATATCTAAGTTGGAGAAACAT
 ACTACTAGATAAATCAATTTGTTCTCTTAGTACTCATGATATGGTTCCTGGGAACCTTCTGAT
 TCACCAAATTAATCTTGGCCAGGTACATACCTGGCAGGAATCCAAAAATCCCCAAGTCTCC
 TTGAAGTTTCAGGATCATCATTCTTAATAAATACACCGAGGGAAAAACCATGGAGAGTTTGTC
 CCAGATGCTGTGAATCTGGCCCGGGGTACATGAAGAAGTCCTTAATTGCAGTCATTTACATG
 GTAGATTCTCTATAATCATTTAATTTGCTATAGGTCTATGATTTTTTAGTCCTTCTTCTCTAA
 ATGATTGAACATGTATAATTTCCCATTTCAATCATATTACCTGGATGAACAAAAGTAACGCTA
 GACTCATTCATGCATTCTGGTTGCCAAGGAAAAGGAAAAAAAAACAAAACAATCAACAGGAT
 GTTTAACTGTCTTAGGGCAACTTCAGGCCATAGTCACTGGTGTTCTTGCAGACTATGAGAT
 ATTTTACATTCTGATAAGGGATAAAAATTCGTGCCTCACATGGCTCCCATCACACTAAGATC
 TTGCAACAATAACACTACTGATTTCAGACATTAATCTTAAGTATCCAGGGAGCCCTAAAACAT
 TGTATCCCACTAGCAAGGACCATGGTAATTGCCACGTAAATCCCCTCCATTATGTGGCCCTT
 ATTATGACCAGCCAGCCAAGGCTTGCCTTTAAATCATACCAATTGAACCGAGCCTTGTAGAA
 ACACTATCACCTACGCATACCTCTGCTTCTTTTCATTAACCTGCTATCCTCTTTACAAATGG
 GATTCTTCACCCACTCCCTTCTTCTAGATTAGCAATGCCCTGTAAAGTAAACGAACACGAAA
 TTCAAAGGGAAACAGGAGCAATCATCATTACCAGCTGCCGTGTAAAGCATTGCGAAAACGCT
 CACGATTTCAGAAAAATCCATGCTGTTCTTTGAAGGCATTCAAGCCTTAATAGCTAGCTGG
 ATGAATGTTTAACTTCTAGGCCAGGCACTACTCTGTCCCAACAATAAGCCCTGTACATTGGG
 AAAGGTGCCGAGACATGAACTTTGGTCTTCTCTGCAATCCATCTGGAGCATTCACTGACAAC
 ATCGACTTTGAAGTTGCACTGACCTGGCCAGCCCTGCCACTTACCAGGTTGGGCTCTGTATG
 GCTAAGCGTTTTCTCCTAAAAATCCCTTGAAAACCTGTGAGAAGACCATAAGAAGATCATATCT
 TTAATTCTATTTTACAAGTCACACAATATTCCAATCAAATACAGATGGTTGAGAAAAGTCAT
 CCATCTTCCCTCCCCACCCTCCCACAGCCCTCAACCCTGCCCTGAAACTTATATGCTGTT
 ATCCGCAGCTCCATCTGGAGCATCACAGCTACTGTCAACCCTGACGCTCTTTCTGAAAAAAC
 ACCGGATGGACATCAGAACTATTTCTTTAAGGATGTTACTGAGCCACACAGGAAAACCTTGCC
 TTATGATTTTGAATGCACGGATCTGATTTGACTAAACATGATAACTAGAGAATCACCCAATC
 TACTCCCATTTTCAACTCTAAATCATCAGAGTGTCTCAAATCCAAAGCACACACAGACCAGC
 CTGGCCAACACGGTGAAACTCCACCCCTACTAAAAGTATAAAAAATTATCCAGGTGTGGTGGC
 GGGCGCCTGTAATCCAAGCTACTTGGGAGTCTGGAGGCAGGAGAATCCCTTGAACCTGGGAG
 ATGGAGGTTGCAGTGAGCAGAGATCACACCACCGCACTCTAGCCTGGGCCACAAATCAACAA

CAACAACAACAACAAAAACAAAGCGCACACAGAGACTGAGGTCCTCTTTGGCATTGAGAAG
ATGGCTATGCAAGTCCCAACTAGCAAGTGCAAACCTCCCAGCTTCACTTCTGCCAGTGTCCC
TTCACCCCTTCTCAACCCCACTGGGAGGCAGGAGGGTGCTTGACAATAACAGCCTTGGCATC
ACTCTGCCAGGGTGTAATAGGAAGTGTTACAATTCTGAGATTCTGTGTAAGCACTGGCCTTT
CTGCCTAGAAATGCCTTCTCCTCTCTTTTTTAAGTGCATGCTCCTATTTATCTTTCAAAGCCC
GGAAAAAATAACACTGCACACGGGAAATGCTCCCTTCTACTGCAGTCATTTAGATGACTCT
ATGCCATTCCATTCAATTTCTCTTTCTACCACAGAAGTGCTTTGAGATTTTGGAGTCAGACT
GCTTGAAGTTGAATCCTGGCCCTCTCATCAGAGACTTGACTTATTTTAGGCAAGTTATATAA
CCAATTTTACCTCAGTTCCTTACCCATAAAAATGGGTCTAATGAGAGTACCTACCACACAGAA
TTTTGATGAAAAGTGAATGAGATGAAGGCCTTTAAGGCAGTGGTCCCCAACCTGGGGACAC
AGACAGGTACCATTTTGTGGCCTGTTAGGAACTGGGCCACACAGCAGGAGGTGAGCAGTGGG
TGAGTGAGATCAGCGTTATTTACAGCTGCTCCCCATTGCTCACCTTACTGCCTGAGCTCCAC
CTCCTGTCAGATCAGCAGTGGCATTAAATTCTCATAGCAGCACAAACCCTGTCATGAACTGC
ACATGCGAGGGATCTAGGTTGTGCGCTCCTTATGAGAATCTAATGCCTAATGACCTGTCACC
GTCTCCCATCACCCCTAGATGGGAGTGTCTAGTTGCAGGAAACAAGCTCAGGGCTTCCACTG
ATTCTACATTATGGTGAGTTGTATAATTATTTCAATTATATAATAACAATGTAATAATAATAGA
AACACAGTGCACAACAAATGTAATGTGCTTGAATCATCCCCAAACCATCCAGTCCACGGTC
TTCCACATTTTGTCTTTTACAAAAATTGTCTTCCACAAAAGTGGTCCCTGGTGCCAAAAAGG
CTTGGGACCACTGCTTTAAAGCCTTTCATAGTGCTTAGAATTGAGGGGGAAAAAAGAAACA
AAAACAATGTAGCTAGTTGCTACAATCACTATATTGGTGAGTTTCAAAGGAAAAAGAATTCT
GTCCCATTTATGCTTGAGCCTTGAGTTGCTAACCAAGCCTGACACAAAATTACTGTTGAAGG
GATGTGTGAGTCCTAATTGAAATGAGGCCTCTTAAGGGAATTGTGGACCAAACCCCAAGCAG
GCAGAAAGCCGTATCTTAATTATTGCAAGTATTTAGGCAAGGTGTGGATGGCCATTTGAAT
TCAAGCAGACTAGGACCTGGGATGAGAAAGAAGGTGTGTACGTGACTTGATCTTTGAACCTT
AGCTCACCATCTGGAAGAAGGCTGAGTATTCTCTGCACTCACATAGTAGCTAATGCCTACTC
CCCAGCCACCCACAATTCTTTCTGTAGGAAGGCTCGCTAGAATACTTTGTGATATTGGATAT
TAGTTCCATATTCTACTGTGTATCTTAGTTCAACCAAATTGTAATCATCTGATATTTATTTT
TTTTAATATAAATATAAGTATATTAAGTCTTGGCATGCTTGCTCAGTCTCTCTCTCTCTCCC
ATTCCTCCCCGCTCCCCTCTCTCTTTCCCAACAGGCTTGGAAAGCAGGCATCACCATGCCTA
TTTAACAGTTGGGGTCCCTTGGCCACCAGGTGCTGGAGTAGGAATCTGAGCCCGGACATGCC
TGATCTGTAAATTTTGTGTTTTCCCACTGTGCTGGGCAGATCACAGCTATCAGCGCCAAAT
TCATAGAAGGGGCGCCCCCTGTGGTCAATTGAGGGATTGTGTTTTGAGGTAGATCTCAAGAA
GGAATGGGTGGGGAAGTTAGCCTAGGACAGAGCAGAAAGGAGCCCTCACTCCCCAAGCACCA
ACGGCCTCAGTCCTTCTGCTGACTCCAGCCTCTAGCTCTCACCCAGACTATCTGCATCCTT
CTCTCCACCACGCTCCTTTGGAACCTGCGTAAAACACAGATTAAAGGAATTCGCGCTTACTT
CCCTTTCCGCATTATGACCAAATGGTTTTTACACTATCATTGAACAGTTTAGTACAAAACATG
CCACCTTTTAATCTATTCAATTCATTTAACAAATACTTTGGAGTGTTTACCATGTGCCAAGTG
CTGTTCTAATAGACATAAGCTGTGAGGTTATGCTTATCTGATTCTCACAGCAACAGCTTTTCG
AGATATGAATTGGTATACTCATTTGACAGATGAGGAAATTGAATTCATGTAGTGAGAAGGAA
GAGCTGCAATTCAGGGTTACTGGTTTTCTCCTGCACTAAGCACTGAGCCACACTAGAAGAGAA
GGCATGAGGAAGACAAAAGTGAGGCTGTGGCTTGCCTTTTCACTTCTTGTGTCCTGTTTTAAG
AAATATCTGCTATCCCAAGGTAAGATACTATACTTTTTTTAACATGTTATTTTGTGTTTTACCT
TTCACATTTGGTGTATACTACATTTGTAATTAATTTGTCCATATTATATGACATATAGCCAA
GATTTATTTTTTACCATACAGATACTCAATATTGCCATTTACGTAGAACATCGTTCTCTTCC
TACTCAATTGCCTTGGCACCTTTGTAATAAATCAGATGATCGTGTATGTGTAGTTCAAGTTTC
TGGACTCTGTCTTCTGTTTTCTTTAGTCTATTTGCCTATTCTTGTACCAATATGCACTGTGTT
AATAATCGTAGCTTTGTAGTAGGTCTCGAAATCTGACAGTGTAATCTTTTAGTTTTCTTTCT
TCTGCAAATTTTCTTTAGCTGTTTTACGTCCTTTGCATTTGTATATAAACTTCAGAAATCAGC
TTGTCTATTCCAAAAACAACAACAACAAATGAAAGTTTCAGAAACTTTAACTGAAATTGTAT
TGAATCTGTAGACAATTTTGGAGTAAATTGCATCTTAGCAATGTAGAGTCTTTCGAACCATA
ACCATGGTAAGTCTCTCCATTTGCTTAAATCTTCTTTAATTTATTTCAACAATGGCTTCCAA

TTTCCAGCGGGAGCTCTTGAAACTACAATTTACATGAACTTCTAATTTGATATTTTTTCGGT
GTCATTATAAACATTGTTGTTTTAAAAGTTGTCTTCAATTTTTTGTGTGCAGGCACAGAAAT
ACAATTATTGATAACATTTATATATAAACTGTATCCAGTGATCTTGCTAGATTCAGTGATGA
GTCTGATGTTGTAGATTCTTTGGGATTTTTCTCCGTACATAATCATATCCTTTCTGAATAAGA
TAGTTTTACTTCTTCATTTTTAATATCTATGCCTTTTATTTATTTTTCTTTTGAACTTTTTG
CTGACTTCATTATTCACTCTCATGTTTTTCTCTTTCATTAGACTATGACTCCTCGATGGTAG
CAATTTGTAGTAATCAAATTTTTGTATTTTATTTTAGCATCTGGCATCTTTCTTGACATATG
TAGCAGTTGCTTTTTTGACAGCTTGCTTCTTCAGTGAATGAATAAATTAATAAAGAGAAATGT
GATGTTTCAGTGATCCATTTTGCAGGTAAGAAAACCTGAGGCAGACAGAGGATGTTAGCAAGCA
AGAGGCCTTGGCCTACAATTTAGATCACTGGACTCTTACTCCAGATGCAATCTGCAGAACCC
ACATACTTTTTAATTAGTCCCTTTGTCTATGTTCTGCCACTGTCACTTCTAAGGAAGGTGTGT
CATCCCAAATGGGGTAGTATCTTATTGGTAGACCTAAATCTGCTGTGTTCCGCCATCTCACCT
ACATGAGTATCTATGTGTAGCATTCTGCATATTCATCTTTTCCACCTTCTGGAGGTTTTGTC
TTTTTATAGGCAGCATGTGAATAACAATGGGGCCAACTGGGGACCAGAAGGGGCCATTTTC
TAGTTCTGAACATAGATAAGCATCACTAACTTTTCCCTCCTGGCAGTAATGGCCTCAAAAAGT
TCCAACCTTAGGAGAAAAAGGCCAAAACCGTCTGCCAAAGTGTGTGAAAAGTTAGAGCAAACCT
TGGTTTTACCAAGAACCTGTGTCCCTCTTATGGAAATTCACACTTTCACACTTTTAGACAAA
TATTAAATGTGTGACATTCTATTACGTACAGTGCCTGGCACATCTAGACACGCAGCACACTT
TAGCCCCCTTCTTTCTTCTTCTAACTCCAAGTTCTAACTAGAAAAAGCCCCACTTGAGTCT
GAGATTTGCTTTTTTGAAGTAGTTTATTTTCTAGATTGTAATCATGCTATCTAGGGTGTGACAG
TGTTTGCTATTTCTAGGGCACCTTGACCTGATTCTTTTTTGCACAGGAACTTGTTCTACCCCT
TTTGCCCACTTCACTGAAGTGAGGACTGAGACAGAGAAGGATTAAGTCACTTTTTTATTTAAC
AAATATTTGTTCCACATTCTCAGCATTTTATTAAATACTGGTGCATAATGATGGAATAAATT
TTATACCGTAAGGATAAACCAGTATTCTGGACTAAGCCAACGTGGGAGACCCTAGGAGGCCT
GTTTGAGAAAGTGACATTTAAATTGAGACTTGGCGGTGGCTGTGGCTACATATCTAGTAAGT
GGCTGGGTGGGATTTGAAGTCTATGCCTGCTTAGCTCTAAAGATGATGCTTTTTGGCTTTGTA
CTCTGCTCTCTCTAGACAACTCTGGTCCAAAATCGTTAAAGCTAACATTTATCCCTGCC
CAACTGGAATTGTCATGTTATGACAAATGGCTCTGTGGTCTCAGATGCCCAGCAGACCCATT
AGTGGAATTCTATGTGCTACAGACCTGGGCAAAATGCCAGAGCCTTATACACCCATCACATT
TCGTCTGGCAAAGGTCTTCAACAAAGAGAAGTAATTACAGCAATGAAAAGCAACAGGTCCAG
CAACACCATAAGAACAAAATAATTAATTTCCCTAAAATAGAAGAAACCATTTATAGAGTAAG
AGCCGATACAATCAATAATTGGAAGAAATAGAAGAGGCTTTAGTATTCTAGCCTTCTTTATT
TGTAGATGTAAATGTGAGCCTCAGAGAAGTTATATATCTAATTAGTGTCACTCAGGTAGAT
ACAACAGAAATTAGGATTAGAACTTAATTCTTATGACTCCCAGAGCAGGGAAAAGACAGGAT
GAAGTCCCAAAACATTGCGTGTGAAGTCAATCTGACTCTGAATTGAGAGTCTGCTATTTAC
TCCCTATGTGACCAGAATCCTTCAGAGCCCATGAGATTCCCTGTCATAGAGTAGATTTTGAT
CCACACTAGGCATTTTTACCTCTTGCTCTTTGAGTTGGTGCCCCATGTTTACTCAGAAATAT
TCCAAAGGTGTTACATCTATTGGTTTTACATGTTGAGCACAGATCATTATAAGACAAATTGA
AATGAAAACATCAACAAGTCTCATTTCATTGTCTAACTTACGCTGAGCAATATTTAATAACTA
GAATATCAAGAGAGTCCAAAGTGTTTGCCCATCCCCTCAAGACCAATGTAATGGAATTTTAC
TCTTATCACCTGCTCAGGGTAGTGGCAATTACAGGATACAGAGGACAGAAATAAAGAATCATG
ACACACAATCCACAGAATTCACAGATGCCAAACATCTACCCTTCTCTGTCACCACACATTGG
ACTCACATGGTGGAAATAGGCAACACAAGCAGAGAGGTGGCTTAACCTTTTCATAATTTTTCA
ACGACCTATGGGAAGAGAGTTTTCTTGTTCAAATCCCAGCTTAGCCACACAGAGTGTGGTA
ATATTGGGCAAGTCAACCAAGCTCTCTGTGCTTCAGTTTCCTCATTATTAAAATGGGGGAAA
TAATAGTGCCTGCATCAGAGGGTTGTTGTGAGAACTAAACGAAATAATTTATCTGAGCTTTA
GAACCCACACCATATTAGTTAAAAATTCATGCATTTTCTTTTATTATATTTCTCTACCTTAG
ACTGCAAACCTCTAAGAGGAAAGGCCGGACTGTTATATTCATAAAGCATTACAGGAACAGTAA
TTAGAACTAGGAGCTTTTCAATGGCCTGCCTGAAATCTGAAAAATAGGTATATTATTTGAAA
TTTTGAAAAAATCAAATAATTAAAAATTAATAGATGTTAATAAAATATCTGTAATATGTAAT
ATCAAGGTCAACTCAACTCTTAATTGTTTATATAAAATATAGTGAAGTTTAAATTGCAAAAT

CTTACAGAAAATGTGCTATTAAAACTCAAAAGTATAATTCTTTCTAATATGTATATATATGT
ATATAGTTTTATTTTAAAGTTCAGGGGCACATGTGCAAGTTTGTTACCTAGGGAAACTCACGT
CACAGGACTTTGTTATACAGATACTTTTCATCACCCAGGTATTAAGCCTAGTGCTCATTAGTT
ATTTTTCTGATCCTCTCCCTCCTCCCAACCTCCACCCTCAGGTAGGCCTCAGTATCTCTTG
TTCCCCTCTATGTGTTTCATGAGTTCTTATCATTTAGCTCCCACTTACAAGTGAGAACATGTG
GCATTTGGTTTTCTGTTCTGCTTAGTTTGCTAAGGAAAGCACTGTGGCAATTTCTCAAAG
GACTAAAAACGGAATTACCATTCAACCCAGCAATCCCATTACTAGTTACGTACTCAAATATT
TTTAAGGCAAAACAAAGCTGCAACCAGAACACCTGGACTCCCTGAAACCCCTTCCACTGATG
TTGTTGTTGTTGTTTTCTTTTTCCCAGCTTCTCAGGCCAAAATACTGGATCATCTTGGGCAC
TGTTCTCTCTGCCCACCCTTTCCCATATGCAGAGTGTTGTCACCTTCTCTCTGCTTCCACTG
CTAACTCCCTGGTCCAAGCCGCTGCACCACTTTTCGTGATTATTGCCACAGTCTCCTCACTG
GTTCCCTGCCCCCACTCTTGCTCTGAACCTATCCAGTTAAACCTGAATTAGATCATCTCATC
CTCATCTCAGAGCTTTCTCGTGGCTCCTCTGCCCTCTCAGGAAAAAATCTAAATTCTAGATG
ACCTAAAAATCCCTTGCTCTTACTGTTTATCTGACCTCATTTACTACCACCTTTTTCTTTG
ATCATTCTGTTCCAGCCACACTGGCCTCCTTACCCTCCTCAAATATGCCAAGCACAGCCCC
CACCCCCCAGGGCTTTGAACTGGCTGATCCCCCTTCCCTGGAATGCCTTACCCCCAAATATCAA
CTTAGCCAACCTCCCTCCTCTCCTCCAAGTGTCTGTTTAAACATGGCTTCAGTAGGAGCTGTC
TTAACATCCTATTAAATATTGTAATTCCTCTCATGACACTTTACACCCCTTCCCTGATATGC
TTTCCATATACCATGCAATATCTGCTGAGATAATATATAATTCACTTATTTTTCTTTATTGTC
ATTTCAAAGAGGGCGGTGTGTTCTGTGTTTTATTTAGTGCCAAAATACTTGCTGATGAAGAG
AGTTCCCTGCCACATAGTAGGTGCTCAATATGTGCTTGTTGAATAAATGTGTCAATGTTTGAT
GTACAGACCTTTTATTATGTTTTGATTTGCTGCCAGTGCTGCCTCCAAACACAGGAGTGCTTC
ATGAGATGTTCAAAAAGCTCTTAAAATATTCCACAAAAATCTTAAAATATTTTCATGAGTTT
TCTTTCTGTATTTTTATAGCAGCATCTGGAATTTAGCCTGCATAGGACCCTCTGTAAGCTG
ACCCTGTTTATCTATTTCAGCTTTACTTCTCCCCTCTCTCCACTTTGTATTTTTATTCTCTACT
ACTTCCAACCTGATTGTAATTTGACCAGACTCCAGACTATCTTATGCCTCTTTGCTTTTGTTT
ACCTGTTACTTCTCTCTGGAATTCCTGCCCCTTCTTAATTTTTCTGGCCAATTCTCACTCT
CTAGGACTCAGAGGTTTTCTCCTCAGGAGACTTCCATGAGTCTCATGTTGAGTTAGGTGACCC
CAATCCTCTGTTCTTCATAGTCATTTCGCGCATTTATCTAGCTCAGCATTTGCCATACTACAT
TGAAATTATTTCTTATGTGCCCATCACTCCCCGTAGATTGCAAACCTCCTAGAGAAGGGCTC
AACAGTGAGTGCTGAGGCTGCACAGAGGAGGAAGGCAGCACAAATGATGGAAGGCTTCCTAAA
GAGGTATGTTCCAAGAGCCCCCACTTCCCTTTCATGGGAGACTCATGCTGTTACACCTAGACT
ATCTAGGGATACATCTAATGTAGTCGTGGAAAGAACAGAGGACTTGAGTACTAGACTGACGT
GATTTTGAATCCTGGCTCCCTATTGACCAGATGTGTGTCTTGAACAAGTCCCTGAGCCTCAG
TGTCTTCATCTGCACAGTGAGGATAATGATACCACACTGCATATATGAGGTATCCGGCACAT
GTAAATGTCCACTACATGCTGATTTCTTCACCCGCTACTCACCCCTGGGAAAGAAGTAGACT
CACCTACTCTTGGTACCCATTTCATTCCCCCTCAGTTGGAAGCATGAGGTGTGCAGCTGCCTG
ACCTGGGGGAAGGGCTGCAAGCAGTAGGTGTTGTGATGTGGTGGAGCTTGTTGACTTCCT
CCCAGGGGCCAGCTAACAACCTGCCTCTGTTCCCTTGATAGTCAAGTTCAACTTTCACTTCT
TAGCACCACAGGAAGTTGACTGAACATTAAGTGAAGTCTCTCTCAAACAGGAGACATCTTTG
CCAGGTCCCTGTACTTCCTAGCCTCATTCCTGCTCTCCCTAGTGAGCAGGCTGCCCTCCCTT
CTCGCCCCAGCACCCTGATAGGCAAGGGTACTCAGAACTACTACCTTGTGGGCCATGTGAT
GTGCCAGGAGCTGCACCCAGGACTTTAATACAGTAGTTGGCTCCCACTGAATGTTTCATTGTT
ACCCCAGGATAAAAAGGGGACACTGTGATCATTTTTCTATTTTGCTGTGATCAGGCTTGTTGA
GCTAAAGTCACCTACCTTCCCAGTCTCTACTAATAGAAGTCATGGATCAGTCCTATTGGTTT
TTCTGTTACAAGGATTGAGAAATTCATAATCATGGAGCTGCATTTACAGGCAGAAGTTTCTTT
CATAGTTTTCTAAGTGTTTCTTTTAGCAACAATGGAGAAAATCAAAGAGGGCAAAGGTGAGG
GGAGAAAATAACATTTCCCTTTCTGTCTTTGCTCTTGTAGTCTTTTGCTTTAGTTTCTTTA
CTATGACTGTGAGGGTGAAACTAGTGATCAGAGTGGTCCAGAATGGGTTTGATGAATCTGAT
TCTGGTGACACAAGATGAATTGGGTATATGTTTCCCTAAAGATAGAGAGACAATATAACATA
GTCTTTACATTAATAGACTCTGGAGCCAATTTTTTTTAGGTTCACTCTCTTTCCCTTTCATGTG

TGTTGATTTTCAACAAACATCTTGCACTCGAATTCCATCTCACTGTTTTATTTTCAAAAAAT
TTAATTTGAGAAAGTTAGCTGTATTAATTTTTTCTTTTTCTAAAATTCTTTTACTAATTGCA
ATTATTTCCATTGATGCTATTCCATTGAAACCATTTTAACATGGACTCAATAACTTCATTGT
ATGTTAATGTTTAATTTTCATTTCTTTACCTTCTTGGTTTTCTAGCTGTGTTAATGTGGTT
GACCACTCATTCTTTGAAGCTCTATTCCCTCTGGCTACTACAGTATGACACATTTTGTCTCCT
TCTTCAGTCTCTGTCTTCTCCACAGTCTTCTCTTCTTTTATATACCTTTAAATATTAATGT
TTCCCAGAGATATTTTCTTAACTCACTTCTCTACTGATTCTAGGTACTTTTCTTGATCCAAC
TCTTCTGATTTTACCCATCTCGATGATTCTATAATTTGTATTTTCTGTTTTGATCTCTTTTC
AGTCTTCCAGACCTAAATATCCAAATGCCTGATGGATAGTGCTTTCTTTTTTACTACCAAGC
CCTCAAAGGCACTATGTTCAAAAGGAATTTGTCATCAGTCTCACGGCACATAAGCTTCCTCT
TGTGTTCAATCTGGAGACTTGAGAGTCTTCCTGTTCCCTTCTTCTCCTTATTTTCTCCATAA
TCAATCACAAAGTCATGTGGATTTTGCTCCTAAATATGTTAACTTCTTTCCTCTCTCACTTT
ATATCCCTTACATCTAGGTATTTTCAAGCCCTCAGTCTCTCTCACATAGACTTTGGCAATAAC
CTTCTAATATCAGTCAACCTGACCAATAGGCCACCAGTGCTTCATGTAGAATCTGGACAATG
TAGAGCACTGAGAATGCTCACACTGGTCATATATGTATGAGTTGGTATGACATCTAGGGAAG
TTGAAGACTTACATAGCCTTTGGCCCAGCAATATACACCATAATACATTAGAGAACTCTAG
CATGTGTACACAGTGATATACACACAAGAATGTTTCACAATGCCATTATTTTAATAGCAAAAT
TGTGGAAACAACACAAATGTTTATCAATAACAGAATGGATAAGTGAGCCATGGCATAGTCAT
ACAATGAAAAATAATATAATAGTCAAAATGAATGATCTGAAGAGATATCATTATTGGCAATC
TTATAAAAGACTGAGTTAAAAATGCAATTTGTGAAAAATTTTAAATTATTTGATATTATTTAA
TGCAAAGTTTTAGAACATGCAAACAACCTGTATATATTATTTATGTATATATGCAAATTCAGC
AATAGCATTTAATCATGCCTGGGAATGATAAGTATCAAAGTCAGAAAGTGGTTACCTTGGG
AAGAGAGGTATGTATCAGCGGTGGGGCACATAGGATGTTGCAGCCATATCTGTAATGTTTCT
TTGCTTTAAAAAATTTGAATCAAGCTTGGCAAAGTGTGACATTTGATTAAGCAGGATAGTGA
GTGCATATCTGTTACTTATATTGTTCTTTATAATTTTCTCTATGCTAAAGCATTTTGTAATT
TAAAAAACCTGACAGTGTTACTCCCATGCTTAAAATATGCCAGTGGTCAAACCAAATCCAGC
AGCACATCAAAAAGCTTATCCACCATGATCAAGTGGGCTTCATCCCTGGGATGCAAGGCTGG
TTCAATATAAGCAAATCAATAAATGTAATCCAGCATATAAACAGAACCAGCAAAAACCA
CATGATTATCTCAATAGATGCAGAAAAGGCCTTTGACAAAATTCAACAACCTCTTCATGCCAA
AACTCTCAATAAATTAAGTATTGATGGGACGTATCTCAAAATAATAAGAGTTATCTATGAA
AAACCCACAGCCAATATCATACTGAATGGGCAAAAACCTGGAAGCATTCCCTTTGAAAACCTGG
CACAAGACAGGGATGCCCTCTCTCACCCTCCTATTCAACATGGTGTGTTGGAAGTTCTGGCCA
GGGCAATTAGGCAGGAGAAGGAAATAAAGGGTATTCAATTAGGAAAAGAGGAAATCAAATTG
TCCCTGTTTGCAGATGACATGTATATCTAGAAAACCCCATTTGTCTCAGCCCAAAATCTCCTT
AAGCTGATAAGCAACTTCAGCAAAGTCTCAGGATATAAAAATCAATGTACAAAATCAGAAG
CATTCTTATACACCAACAACAGACAAACAGAGAGCCAAATCATGAGTGAACCTCCATTACACA
ATTGCTTCAAAGAGAATAAAAATACCTAGGAATCCAACCTTACAAGGGACATGAAGGAACTCTT
CAAGGAGAACTACAAACCACTGCTCAATGAAATAAAAAGAGGATACAAACAAATGGAAGAACA
TTCCATGCTCATGGGTAGGAAGAATCAATATCGTGAAAATGGCCATACTGCCCAAGGTAATT
TATAGATTCAATGCCATCCCCATCAAGCTACCAATGACTTTCTTCACAGAATTGGAAAAAAC
TGCTTTAAAGTTTCAATATGGCACCAAAAAAGAGCCCGCATCACCAAGTCAATCCTAAGCCAAA
AGAACAAAGCTGGAGGCATCACACTACCTGACTTCAAACCTATACTACAAGGCTACAGTAACC
CAAACAGCATGGTACTGGTACCAAAAACAGAGATATAGATCAATGGAACAGAACAGAGCCCTC
AGAAATAACGCCACATATCTACAACCTCTCTGATCTTTGACAAACCTGAGAAAAACAAGCAAT
GGGGAAAGGATTCCCTATTTAATAAATGGTGCTGGGAAAACCTGGCTAGCCATATGGAGAAAG
CTGAAACTGGATCCCTTCTTACACCTTATACAAAATTAATTCAAGATGGATTAAAGACTT
AAATGTTAGACCTAAAACCATAAAAACCCTAGAAGAAAACCTAGGCATTACCATTCAAGGACA
TAGGCATGGGCAAGGACTTCATGTCTAAAACACCAAAAGCAATGGCAACAAAAGACAAAATT
GACAAAGGGGATCTAATTAACTGAAGAGCTTCTGCACAGCAAAAAGAACTACCATCAGAGT
GAACAGGCAACCTACAAAATGGGAGAAAATTTTCACAACCTACTCATCTGACAAAGGGCTAA
TATCCAGAATCTACAATGAACTCAAACAAATTTACAAGAAAAAACAACAACCCCATCAAA

AAGTGGGCAAAGGACATGAACAGACACTTCTCAAAAAGAAGACATTTATGCAGCCAAAAAACA
CATGAAAAAATGCTCATCATCACTGGCCATCAGAGAAATGCAAATCAAAACCACAATGAGAT
ACCATCTCACACCAGTTAGAATGGCAATCATTAAAAAGTCAGGAAACAACAGGTGCTCGAGA
GGATGTGGAGAAATAGGAACACTTTTACACTGTTAGTGGGACTGTAAACTAGTTCAACCATT
GTGGAAGTCAGTGTGGCGATTCTCAGGGATCTAGAACTAGAAATACCATTTGACCCAGCCA
TCCCATTACTGGGTATATACCCAAAGGACTATAAATCATGCTGCTATAAAGACACATGCATA
CGTATGTTTATTGTGGCACTATTCACAATAGCAAAGACTTGGAACCAAGCCAAATGTCCAAC
AATGATAGACTGGATTAAGAGAATGTGGCACATATACACCATGGAATACTATGCAGCCATAA
AAAATGATGAGTTTCATGTCCTTTGTAGGGACATGGATGAAAATTGGAAATCATCATTCTCAGT
AAACTATCGCAAGGACAAAAACCAACACCGCATGTTCTCACTCATAGGTGGGAACTGAACA
ATGAAAACACATGGACACAGGAAGGGGAACATCACACTCTGGGGACTGTTGTGGGGTGGGGG
GAGGGGGGAGGGATAGCATTAGGAGATATACCTAATGCTAAATGATGAGTTAATGGGTGCAG
CACACCAGCATGGCACACGTATACATATGTAATAACCTGCACATTGTGCACATGTACCCTA
AAACTTAAAGTATAATAATAATTAATAAACCATAAGTTTATGAAACCCCCCCCCAAAAA
TATATGCCAGTGGCCTCCAGTTGCCACCAGGTAGCATCCACATTCTTAAATGGAAAGCCCT
TCCTTGCTTCGAACCTGCCAACTGGGATTGGACATTTGTAGTTGCATTTCTAAGAACTGTTT
CCTTTTGTCAATGGAGCCTGATTTCCACTTGGATATCTGGGTGATTTAGGGAACTGACCTC
AAAACCAATTCTACATTTTCGACCATGTGACCTTGGCTTAATCAATTCACGCATCTTTTTCC
CTCACCTCAGGGGATGATCATATGAACTAAGCCAGTTGCAATAGAGTAAACCTCATGTTCTT
AATGAGAAATCCAGAACAAAATGCTTTATTTTCTTCAATTTTTTATTAGGTATCTCCTGA
ATCAATTAAAAAACCACAGTGACAACAAAATAAAAAATATGAAGAAGCTGAAACAT
GAAAGCTCTGCCAACTGCAATATGTAGCTGCTAAGGTTGCTGTATTTATTGGAATCAAGCAA
GTGTTCCAGTAAAGAGCACAGAAGATGTGTCTGGGAGCCTTTATGTGTAGGTCTGCAAGTGG
TGGATATCACTACTACTCACACGCCATTGGCTAGAAGTGTGATGATGATACACCTAATTG
TAAAGAGAGGCTGGGAAATAGAGACTATTGTGCCAGAAAGAAGAGAAAATTCATTTATGGAA
GAGGTAGCTAGTCTCTCACAGCCATGAAAAGAGAAGTGTTTAGCTAATTGAAGTGAATAGCA
GCCATCTTGGGTCCCTAAGGCAAGTTAGACTAATATTGAAGTGGAAACCATGAGGAAAGCAG
TGATACTGAAAGTAACCGCATCTTTGAGAACATGCATTCATTTCTTACAACATGAATTTATT
GAGGACCTACCTTAATACAGGCAGCGTGCTAGACACCAAGAGAACTGATGTCCTCTTCCTTC
CTGCCTGCCTGGAGCCTGTATTCTGGAGGGGACAGAGCTAGCAGATCAGACCTAACTGGAAA
TCTGCTGTGCCAGTATATATTTTCAGTGATGTGAGCCAATATATCCCCTTGATTGCTCAAAGT
AGTTTGGTCGATATATTTTGTGCTTTAAATTGAACACATTCTTATGTACAGCCTCTGTCTC
CTCATCTCCAACCAAGCAAAATAGCTTGTTCTCTTTATGCAGGGACACATGACATTTCCCAC
GTGGCTTTGTGCATATCTCCACCTCAATTTAAAATGCCTTCCAATCCCTGCTCAAAGTCAA
CAGCTTAATAATGGTAGACATAGAATTTGACTTATTCTAATAATAGGTCTTTTAAACAATGC
CTTCTTCTCTTCAATCTTTCTTCTTAGAGTGGGTATTCTTTCTGGTGCATCATGTAAAGGA
AGGTAACATCATGCATGTAATGATGAGAATATTTATATGTATTTATGATTATCACAAAAA
CAAAGATTCTACCATTCAAGAGGAACATTTATTTTATTTTTTTTATTTGAGAAAAGTATAATT
TTATTTATTTATTTATTTGTGCAAATTTATGGGGTACTTGAGAAAATGTGTTACATGTATAT
AATGTGTAGTGATCCAATCAGGATACTAAGGGTGTCCATCACCTGAGTGTATTACATTTTTG
TTAAGTATAATCATCCTACTCCAGGAGAACATTTTAAAACTGTTCTGTAGAGATACTACTC
AAATTAAGTTCTCAGTCCTGAAACATCAGATCAGCTAGGAATCTGACAAAAATGCAAGTTCT
CAGATGACAGATGAGACCACTTCAATCAGAATTTCTGGAGTGGAGCCCACACATTTGTATTT
TTGCAACCTTTCCAATGATACTTATGTACATGCTCAAGCTTGAAAACCACTTTCCCTAGGACA
TTAGTTCCCTCGACAAGATTTGTGAGTAACCTTGTTTCATGAAAAAGTGTTTAGGAGATACTG
ATTCAATAAAAACTAATCAGGCTTTTATTGTTTGCAGGGCTTTCAAACCTTGCAATAGGCCA
CTGTGCATTGTAAATTTCTAAGAGGAAGATGCTTATGTCCTCAATGAATATCTTCCCACCAT
GAAGTACTCTTCTTCCCCCACTTTTTTAAACAATTACTAACACCTGGCAGAAGTAGGCAGAC
AGCTTACAGCTTAGAAAAAGTTGGCCTAAGATAATGGCTAATTTTCATACATTATTTATTG
TCATCATGCTTATCTTCTCTCTCTAAATTGTATATTTTCATCTCTGTGATCACAGATTGAGC
CTCATATTTTCATATCTGCCCCCTGGCCTAATGGTTGTTTACAGAATGAGCTCAATGAATATTG

TTAAGTGAGTAGGATTTAATTTATTTGATAAATAGATAACCTTAAGTTTTAAACGGTGGATT
TCACATGAGGACATTTACTTACTATTGTTGAGCTGTAATTAATTTTTTAATACTGTTTAGGTA
CTCATAATAAAGAACAGGATATTTGGAGAAGGAAGACAGTATCATTCTGGTTCTTAGTCTT
ACCAGCTTATTGATCATGAGTATATAACCTCTCTGTGGCTCAGTGCCTTTCTCTGTAAATG
GGAACACAGTGATGTTTACCTCACAGGATTGATTTGTAAAAGGGCTGGATAAGGTTATGAGA
ATGTTTTGCAAAGTGATATCGAAAGATTAATTGCAAACCTCATTGAATCTTAAATTGTTTG
AGATAGGTCATGCTATGAATCAACTATGAAGTGCAGATATTGTCAAGATTCAATATTTCTTT
CCCAAGAGCTGAGAGGAGGGGCTGCTTGTGTTGTTTCTTTCTTTTGTAGAAACATGCCAG
GACAGGCTCATTTTTCGGGTTTTCTCTCACTTGCTCATCTTACTTTTTCTTTAGTTTTCTCTA
TTCATTAGGATACAGTACTGTAAAGCTTTATGGCATTTTTTATTTTGTGGGAGATGAATCTGA
ATAAGAATTACAGTTAAATCATTGCTAAGTTTGATGAATGAGCACCAAAGAACTCTTCAAG
ATGTCATTTTTTAAAGTTTTGTAAATGATTGGCTTTCAGTGGTTTCCTCTAAGGAATTTTAAT
TTTGAATAATGCATAGAAAAATGTGCGCACACACAAATCATTCAAGTATCCACCTCGAAGGGA
AATCAAAGTGCCTGTGAAGTGAACCTTTACCTTTCTATATCACCAGCTTCCTGTTAGAGCAG
ACTTTTTCTTTGCTCAAAGTCTAAGCATTGAAGAACTTCTTTTTAGTAGGTAGATTTTTGTG
TTTTTTTGTGTTGTTTTTGAGACGGAGTCTCACTCTGTGCGCCAGGCTGTGGTGCAGTGGCAC
GATCTCGGCCCCACTGCAAGCTCTGCCTCCCGGGTTCACACCATTCTCCTGCCTCAGCCTCCC
AAGTAGCTGGGACTACAGGTGCCAGCCACCACGCCTGGCTAATTTTTTTGCATTTTTTAGTA
GAGATGGGGTTTTATCGTGTTAGCCAGGATGGTCTTGATCTCCTGACCTCATGATCCACCCG
CCTTGGCCTGCCAGAGTGCTGGAATTACAGGCGTGATTTAGTAGGTAGTTTTGAGTAGGGAG
TATACATTTAAATGCTGAACTCAGTTAAGGAATAATCTAATACTGTATTCAACTGAACT
CAGTTGAGAAATTTCTTTCCAATAATAAAGGAAAATCAACTGCAGTAATGAGGGAGATGATT
TTGCTGCTAATTACAACAAATATTTACTACAGACCTGTTATGCACCAGGAAGTGTGCTAAAT
GTTTTATACATATAACTTTATCTTGCTCCAACAACTTATTACATAGAAATTGCTATTATT
CCCATTTTCTAGATTAATAAATTGGTTTTAGAGGGGTCGTATAGGTGAAACAACCTCACTCAAT
ATCACAAGCTGTTATGTGGTGAAGTTTGCATGATCAGTACAGGGTTCTGGTCATCCCACTCA
TTGAGTGGTGCTAGTCAAGATCTGGAAGCTCTTCTGGTCTTAGTTTTCTCTAGCCGTGAAGTG
ACAATGATTAGGTCTAATCATAGAACATGAGAGTACATGTGAAAAAATGCCTTTTTTAAAGAG
TATGAAAACTTGAGTTGTAAATTTTCTTGTTGGATAATTTATTATTGCTTTTCTTTTTTAG
ATAACACTAACAAAGTTGACCTTAGAATTGGAGTGCCTGGGTAGAACCTGCTGGTACCAC
CTGCTTACTGGCATGCTTCTGATGTGAGTTCAGGAGAAGACACTGGCAAGGACAGCAAAGAA
CAGGAGAACACTCTAGCTTCCCTGATAGCATTCAAGGTGCTGTCCAAAGCTGACTGTGATGG
CACCTCCAGACAGACAGCGATGCCACATGTTCAAGATGGCAGAATCACTATCAGCTTCAAT
TCCTGAATGACTGCAGAGCAAAATTTCTTACCTGCAACATACACTCTATTTTCAGCCTCCCT
GGACTGTTACATAATGATACATAAAAAATATTTCTTGTTGAGGCATCCCAAATTTGATTTA
TTTGTATCACAGTCTATCCTATGAATATATTTCTGATCCAAATAATGCTAGATTCATGAGA
GTTTATAGTCCAGAGATTTTTTTTTTTTTTTTGGAGACAGAATCTGTCTCCAGGCTGGGGTG
CAGTGGCATAATCTTAGCTCACTGCAACCTCTGCCTCCCGGGCTCAAGTGATCCTCCCTCCT
TAGCCTCCATGAGTAGCTGGGACCACAGGTGTGCACCACCACACCCCGCTAATTATTTTGTA
TTTTTTGCAGAGGGGGGTTTACCACATTGCCCAAGCTGGTCTCAAACCTCCTAGACTTAAGG
GATCTGCCTGCCTTGGCCTCCCAAAATGCTGAAATTACACTTGTTAGCCATAGTAATAGTTC
CTAGCCATAGTCCGGAGATATTTTAAAAACTTTAGTGTGAGTTATTTCTCTTTGCTATAATT
CTCTCATGCATTTTGAAAACCTTTTCTGCCATACAGTGGTTTTTTGGGAAGTCCTTTTGACA
GACATATCACAGGATGGAGAACTAAAAAGAGAGAATGCATTAAAGAAAAAATAGTACTTCG
TATTTTCAGAAATTACAAAAGGGTGTGATGCCCCACAGCATGGTGGGCAGTCATTTGTAACCC
ACAGAACATTGAGTTTCTAAAATTTGAGTGTTTTTTAACTTACGAAGGCCAACTTTACTGATG
AAGTGATTACCCCAAATATGTGAAATCCATATTTCAAGTACAATAGTAGGAAATTGGAAATG
GGGCTCTTGAGTTTCTGCTTTCAAGTGACCCTCAGAAATTCCTCCATTCACTGCAGAGGTTT
TGTTTCTCCTTGCTTTGTTCTGACTTTACGGCAGAACTAAGCTAATGAGTTAGTTACTATGG
GTTATCACTTGGATTTGAAGAACCATCATTTCTAGGCATTGCTGCTCAGAGTGGGGTCTGCC
CTCAGGGGAAGCTGTCTAACCAGAGCCTAACCCTAGGGTTTTATCAGAACCTAACTGACC

TGAGGAAAAGAAATACTCAACTCCAGCAAGCTCCAGCCTTCCAAATGGAGGAGGAGAAAACC
CAACTCCAGCCCACCTTCACTATCTTGTCCCACATAAAGGGGAAGTGAAGAAACCTGAGAA
GCAGGCATGAAGTTCACATGTATTCCACACCATTTCATATTCCATCGCATTCCATTCTACAC
AATTCCACATCTCCCCGGCCACATCAGCCTGAGAGTAATAAACAGACCTTGCCATCGTATCC
CCTTTTGGAGACACAAGCCAGGAAGGAAGCTCTTCAGTGCAGCTGTGAATAGAGAAATGCTG
GCTGAGGTTTTGGAGACACTGGGTGATCATTCCCAGTGAAATTTTTAGATCCCTGGGGACAT
GAGCTGCTTTGTCTTTTCCAAGGTCAGGATAAACAGAATAACTTCAGGCTTCTCTACCCAGA
AAGAACATGTGGCATAAATATCAACTGCAGAATAAATATGATTAATCTGGTACATGGACGAA
GATGTTTTCTAGGAGATGCTTATCCTGGGATGAGAGCTTTCATAAGCATTGATATTTACATG
ACTCTTACCGTGTGTCAGGAAGTGTCTAATTGTTTTACCTATGTCAATTCATTTGATAGTC
ACAACCACTGAAAGGAGTAAGTACTCTTATTACTTCCATTTTACAGATAGGGAAACTGAGGT
ATAGAGAAGTCAAGTGACTTGCCCGAGGTTATTAACTACTTAATGTCAACACCAGGATTTG
AACCCAGATCATTGTGTTCTGAAGTACATGCTCACAATCACTGCATTACTGATACACTGTTT
TGTCTTTGCATACTTAAGTGGTCATAACTTAGTCTGAAACACTTTGTGAGAGCAGGAAGCAA
ACTGTCCCCTTATTAGGTGGACCAGTATAGTGATAATACAAAAGTGTATTGCATTTGAATTA
CTTGCTAATATCTTCTAATTGAGGCAATTTTGAACAGGAATATACATATCTAGCCTCTATTT
TTCTAGCTAGAAGTTCTGAAATCCCTGGGCTTAATATTGTATGGCAACAATTGGCTGGAGTT
GAGTTGCTGCCACTCTCTTTTAACTGAGCCATGCTCTCTCTAGTTTGCTACAGGCCCCACCA
CTCCCTATTGCCTCTCCAATACCAGGTCATTTGGCATCTTACTCAGCTCATTTCATGCACAT
GTGTTCCACAATTGGTAACATAACCCTAAAAGTATTTGAGTTTGTGGCTTCTGCTCTTGTGA
CAGAAGACTTTTCTCGAATTCCAAGGTCAACATATACCATATTGACTCTGGGCCACATTTTT
TAATGTGCTCAAGCTCAGTTTTTCTCCTTAAAAAATGAGGGGTAAAAAATAACACTTCGTAG
TTGCCTCATCTGTGGATTGGAAGAATGAATGCCTGTCAATTTCTAGAGTTGTAGTAAGGGCCA
GTTGGGGCAGTGCCTGTGAATATACGCAATGGGCCATCAAGCAATCTCAGGGCTTCAGGCAA
TGCTGGGGTTTTATAAAGCACTTTATGTTTTAAGTTCACTTTTATTTCTAAAGTCTCATTGAC
TGCTCTGAAAATCTCTCAAGTAAAGTGGGCACTAAAGGCTTTATACTCTCTCCAATTATACC
TTCCATTGTATAGATTTGGAAGCTGGGGTCCAAATGTGTTAAATGACTTGCCTAACATGGTC
CATTACTGTAAGTACAGAAACGGAATGAGACCCCAAATCTGCTTCATGGAGGGACACTCTTC
TAAGATACTGTGATGCTTCTTCCCAAGTAATCCGTCTTCAGACTTCAAGGTCTCAATTCTGA
ATGACAATTCAATATAGGACTTTCATAATCTTAAAAGCAACCTGACAGTCATTACAGTGGGC
TGTGAGAAATACTAACGCAGAGCCACATACTCTGGGCTTCATGCTAGGTTTTGCCACTCACT
ATCTTTGCTGAAAAAGTTTTGGAAGCCCTCCTAAGCAGGTGCCAGACCCTTTTTTGGCCAGA
GGACAGGATCTTACGCTGTTGCCAGGCTGGAGTGCAATAGTGTCAATAGTGTCAATAGTGTG
CCTGGAAGTGTGAGGCTCAAGGGATCCTCCCAGCTCAGCCTCCCAAGTAGCTAGGAGTAAAG
GTACATGCCATCATGCCTAGCTTTTTTTTTTTTTTAAATTTTTGGAGAGATAAGTTCCCATCAT
GTTGCCAGGCTGGTCTCTAACTCTTGACTCAAGTGGTCCTCCCACATCAGCCTCTCAAAG
TGCTGAGATTGCAGGTGTTAGCCACTGCCTTGGCCTGGCCCCAGTATTCTTTGGGATCTGG
AGTTTGCTGTTGAATGAGAAGGCAAGATGAAATCCATGTAGTCAGACTCCTACGCTGCTGT
TCTAAGCAGGGTTGGGCCTGATTAGTAGGTTATAGATGATGTTTTTCTGTGGTGCTATCTGG
ACCTAGTGCTCTTTGGCATCTGGGAAGGTATGGCCTTTAAAAAGCAAACCTGCCATGAGAACT
GCTCTACCCCAAATTTTGGTTCACAGCCTTCATTTGATTATGTATTGGGGCAAAAATAGTTT
AGCCATGTGAACCTGTTTGTAAACTGGTGAGTTTCTATTGCTATTTTCATAGCTAAAGTTTTG
AGGTAAATGCTATTGGATCTTTGTGTCTGTGTGTGTATACATATTTAGATTTTTTTTTTTTT
TTTTTTTTTGGTTCTTTGAAACATTGCTGATTCTTTTTGTTTTGTTTTTTCAGAGTCTGGAGA
ACACTTTTTCTTTTGGAGCTGTTTACAACCTTTAGCAATTGAGTAGAGTGACTCTTGTCAAT
AGAATTTGAAGCACATTTCTCTCTCTGCCTGATTTCTGTAGAATTTGTAAACTATTTGTGAA
TATTCTTAATTTATGGCAATATGGTTGCTTACATAAGTTCAATAATAATCTGTTTTCTTTTA
CAATGAGACACAGTTGGAGGAAGTGGTTATTTTCCCAGGGCTTTGACTGAAATGGCCTTGTG
AATGGTTCCAGGAAAGCCAATTTTGGAGACCCTATGTGGATGATGATGCTTGCTGTACTTTC
TGTGGGTAATCGGGCCAAGTATATGGGACTGAAGCTAATTTTGCAGGGCAACATAGAGAGAC
TTGAGTTCCAGGGGGAAAGTTTTAGGATGGAGAGAAGCCCTCACTGGACTGTGATGTGGGGT

GAGAAAATGGAGGTCCAGAGAGAAAAGTGACTCACCCAAGCTGAAAACAGTTGCAGGAAGAA
AAGCCAGGACAGAACTGGGCTGTTAGAAACCTTGGTAGTGCACTTTTGGATTTCACTCTTAAA
ATGCTCAAATGTCTTTTTCTGACAACATGTAGGGAGACCTGAGGTTGGACCTAAAGACACCA
TATATTGGATTGAGTGCTTCACACAAGTTAATCTAAATTATGAAGGAGTACACTAAAGATGA
GGAAGTGAAGTACCGCAATGCCAAATGTAACACGCTGTGGGTCAAATTTGGATGGTTGCGT
TGGAAATTATTCTGAAGCCACAGACAGGGCTAAAAGGAGTCTAGAATGTCCTACTAGCTGTT
GGAACCTGCTGGAGTTCTAAAGGCATGCGATGTCCTGAATGTGCTTGACCAATGGACACAGC
AGGTATTTGAAGAGTTTTGATGCTTCCTTTCTGTTTTTCAGTGCTATTTTTTTGCTTTATTTT
TTTTAAAAAATTTCCCCACAGAGGCTATTTATTACACAGCAGCAGCCAGCATGCACCAAGC
AGAAGTGTCTTTTCATAATAAACCTATTTATCTTCTGGCATGTCAGAAATTTCTCAAATTGAT
ACTTATAATAAATATGATTAATAACACATTGTATTTTCAGTTGTTGCCGTAATAAATTGGGA
ATAAATATCGTATTGTGATGTTGACAGTGTGAGGTTGGTTGGTGCCAAACAGCGGGAGCAGC
CAACTGTATTTTCATTAAACAGCCACATTGCTACAATGCTATTAAAGTTTTTCATAATCTCTTCT
GCTCAAGGTTGATCATTCTGTTAGACAAATTCTTGCAACAATCACACTGGAAGTAAAT
CTGCCAGCGACAATCTCAGCAAGATGGGTTCTCAC-5'

<128128655> (SEQ ID NO: 1)

>CCAT1_JAX_2 Transcript sequence; Genomic location:

chr8:128128655-128232653 strand:-

<128232653>

3'-

AAGGATTTAATGGCAAGATGCCATTATAGACAAGAACAGGATTCAGACACTTTTCGTGTTATG
TGTTCTTGTCCAAATACTGTGTCACCTCTGCACTGGCATCCCAATCCCACCAACACCTTTAGC
AGGAACCTCCTGTTTCAATAACATTTCTCAATACTCTACCTGATTGCTTAGAATCCATGAGA
CAGCCATATTTCTCCATGTCTAGGTCCCTATCTTATTTTTGTTGAGATGGTGTAAAGACTTTA
TTTGTGAGGCTTCTGGGGAAGAGGAGTAAGGTATTGATCCCACTGACTGGATAATTTGGGTC
TCAAATGGATAATAAATAAGCATTACATATTTTGACCACTTCCTTGGAGGAGAAGCTTCTTG
GAATGTGCACCATGTCCGCTGCACTTTTTTTTGACAGATATCTAAGTTGGAGAAACATACTA
CTAGATAAATCAATTTGTTCTCTTAGTACTCATGATATGGTTCCCTGGGAAGCTTCTGATTAC
CAAATTAATCTTGCCAGGTACATACCTGGCAGGAATCCAAAAATCCCCAAGTCTCCTTGA
AGTTCAGGATCATCATTCTTAATAAATACACCGAGGGAAAAACCATGGAGAGTTTGTCCCAG
ATGCTGTGAATCTGGCCCGGGGTACATGAAGAAGTCCTTAATTGCAGTCATTTACATGGTAG
ATTCTCTATAATCATTTAATTTGCTATAGGTCTATGATTTTTAGTCCTTCTTCTCTAAATGA
TTGAACATGTATAATTCCCATTTCAATCATATTACCTGGATGAACAAAAGTAACGCTAGACT
CATTCATGCATTCTGGTTGCCAAGGAAAAGGAAAAAAACAAAACAATCAACAGGATGTTT
AAACTGTCTTAGGGCAACTTCAGGCCATAGTCACTGGTGTCTTGCAGACTATGAGATATTT
TACATTCTGATAAGGGATAAAAATTCGTGCCTCACATGGCTCCCATCACACTAAGATCTTGC
AACAAATAACACTACTGATTTCAGACATTAATCTTAAGTATCCAGGGAGCCCTAAAACATTGTA
TCCCCTAGCAAGGACCATGGTAATTGCCACGTAAATCCCCTCCATTATGTGGCCCTTATTA
TGACCAGCCAGCCAAGGCTTGCCCTTAAATCATACCAATTGAACCGAGCCTTGTAGAAACAC
TATCACCTACGCATACCTCTGCTTCTTTTCATTAACCTGCTATCCTCTTTACAAATGGGATT
CTTCACCCACTCCCTTCTTCTAGATTAGCAATGCCCTGTAAAGTAAACGAACACGAAATTCA
AAGGGAAACAGGAGCAATCATCATTACCAGCTGCCGTGTTAAGCATTGCGAAAACGCTCACG
ATTCACAGAAAAATCCATGCTGTTCTTTGAAGGCATTCAAGCCTTAATAGCTAGCTGGATGA
ATGTTTAACTTCTAGGCCAGGCACTACTCTGTCCCAACAATAAGCCCTGTACATTGGGAAAG
GTGCCGAGACATGAACCTTGGTCTTCTCTGCAATCCATCTGGAGCATTCACTGACAACATCG
ACTTTGAAGTTGCACTGACCTGGCCAGCCCTGCCACTTACCAGGTTGGGCTCTGTATGGCTA
AGCGTTTTCTCCTAAAATCCCTTGAAAATGTGAGAAGACCATAAGAAGATCATATCTTTAA

TTCTATTTTACACAAGTCACACAATATTCCAATCAAATACAGATGGTTGAGAAAAGTCATCCAT
CTTCCCTCCCCACCCTCCCACAGCCCCTCAACCACTGCCCTGAAACTTATATGCTGTTATCC
GCAGCTCCATCTGGAGCATCACAGCTACTGTCAACCCTGACGCTCTTTCTGAAAAAACACCG
GATGGACATCAGAACTATTTCTTTAAGGATGTTACTGAGCCACACAGGAAAACCTGCCTTAT
GATTTTGAATGCACGGATCTGATTTGACTAAACATGATAACTAGAGAATCACCCAATCTACT
CCCATTTTCAACTCTAAATCATCAGAGTGTCTCAAATCCAAAGCACACACAGACCAGCCTGG
CCAACACGGTGAAGCTCCACCCCTACTAAAAGTATAAAAAATTATCCAGGTGTGGTGGCGGGC
GCCTGTAATCCAAGCTACTTGGGAGTCTGGAGGCAGGAGAATCCCTTGAACCTGGGAGATGG
AGGTTGCAGTGAGCAGAGATCACACCACCGCACTCTAGCCTGGGCCACAAATCAACAACAAC
AACAACAACAAAAACAAAGCGCACACAGAGACTGAGGTCTCTTTGGCATTGAGAAGATGG
CTATGCAAGTCCCAACTAGCAAGTGCAAACTTCCCAGCTTCACTTCTGCCAGTGTCCCTTCA
CCCCTTCTCAACCCCACTGGGAGGCAGGAGGGTGCTTGACAATAACAGCCTTGGCATCACTC
TGCCAGGGTGTAAATAGGAAGTGTACAATTCTGAGATTCTGTGTAAGCACTGGCCTTTCTGC
CTAGAATGCCTTCTCCTCTCTTTTTTAACTGCATGCTCCTATTTATCTTTCAAAGCCCGGAA
AAAATAACACTGCACACGGGAAATGCTCCCTTCTACTGCAGTCATTTAGATGACTCTATGC
CATTCCATTCAATTTCTCTTTTCTACCACAGAAGTGCTTTGAGATTTTGGAGTCAGACTGCTT
GAACTTGAATCCTGGCCCTCTCATCAGAGACTTGACTTATTTTAGGCAAGTTATATAACCAA
TTTTACCTCAGTTCCTTACCCATAAAATGGGTCTAATGAGAGTACCTACCACACAGAATTTT
GATGAAAAGTGAATGAGATGAAGGCCTTTAAGGCAGTGGTCCCCAACCTGGGGACACAGAC
AGGTACCATTTTGTGGCCTGTTAGGAAGTGGGCCACACAGCAGGAGGTGAGCAGTGGGTGAG
TGAGATCAGCGTTATTTACAGCTGCTCCCCATTGCTCACCTTACTGCCTGAGCTCCACCTCC
TGTCAGATCAGCAGTGGCATTAAATTCTCATAGCAGCACAAACCCTGTCATGAACTGCACAT
GCGAGGGATCTAGGTTGTGCGCTCCTTATGAGAATCTAATGCCTAATGACCTGTCACCGTCT
CCCATCACCCCTAGATGGGAGTGTCTAGTTGCAGGAAACAAGCTCAGGGCTTCCACTGATTC
TACATTATGGTGAGTTGTATAATTATTTTATTATATAATACAATGTAATAATAATAGAAACA
CAGTGCACAACAAATGTAATGTGCTTGAATCATCCCCAAACCATCCCAGTCCACGGTCTTCC
ACATTTTGTCTTTTACAAAAATTGTCTTCCACAAAAGTGGTCCCTGGTGCCAAAAAGGCTTG
GGACCACTGCTTTAAAGCCTTTGCATAGTGCTTAGAATTGAGGGGGAAAAAAAACAAAAA
CAATGTAGCTAGTTGCTACAATCACTATATTGGTGAGTTTCAAAGGAAAAGAATTCTGTCC
CATTTATGCTTGAGCCTTGAGTTGCTAACCAAGCCTGACACAAAATTACTGTTGAAGGGATG
TGTGAGTCCTAATTGAAATGAGGCCTCTTAAGGGAATTGTGGACCAACCCCAAGCAGGCAG
AAAGCCGTATCTTAATTATTGCAAGTATTTTCAAGGCAAGGTGTGGATGGCCATTTGAATTCAA
GCAGACTAGGACCTGGGATGAGAAAGAAGGTGTGTACGTGACTTGATCTTTGAACTTTAGCT
CACCATCTGGAAGAAGGCTGAGTATTCTCTGCACTCACATAGTAGCTAATGCCTACTCCCCA
GCCACCCACAATTCTTTCTGTAGGAAGGCTCGCTAGAATACTTTGTGATATTGGATATTAGT
TCCATATTCTACTGTGTATCTTAGTTCAACCAAATTGTAATCATCTGATATTTATTTCTTTT
AATATAAATATAAGTATATTAAGTCTTGGCATGCTTGCTCAGTCTCTCTCTCTCTCCCATTC
CTCCCCGCTCCCCCTCTCTCTTTCCCAACAGGCTTGGAAGCAGGCATCACCATGCCTATTTA
ACAGTTGGGGTCCCTTGCCACCAGGTGCTGGAGTAGGAATCTGAGCCCGACATGCCTGAT
CTGTAAATTTTGTGTTTTCCCCACTGTGCTGGGCAGATCACAGCTATCAGCGCCAAATTCAT
AGAAGGGGCGCCCCCTGTGGTCAATTGAGGGATTTGTGTTTGAGGTAGATCTCAAGAAGGAA
TGGGTGGGGAACCTAGCCTAGGACAGAGCAGAAAGGAGCCCTCACTCCCCAAGCACCAACGG
CCTCAGTCTTCCCTGCTGACTCCAGCCTCTAGCTCTCACCCAGACTATCTGCATCCTTCTCT
CCACCACGCTCCTTTGGAACCTGCGTAAAACACAGATTAAAGGAATTCCGCCTTACTTCCCT
TTCCGCATTATGACCAAATGGTTTTTACACTATCATTGAACAGTTTAGTACAAAACATGCCAC
CTTTTAATCTATTTCATTCAATTAACAAATACTTTGGAGTGTTTACCATGTGCCAAGTGCTGT
TCTAATAGACATAAGCTGTGAGGTTATGCTTATCTGATTCTCACAGCAACAGCTTTTCGAGAT
ATGAATTGGTATACTCATTTGACAGATGAGGAAATTGAATTCATGTAGTGAGAAGGAAGAGC
TGCAATTCAGGGTTACTGGTTTCTCCTGCACTAAGCACTGAGCCACACTAGAAGAGAAGGCA
TGAGGAAGACAAAAGTGAGGCTGTGGCTTGCCCTTTTCACTTCTTGTGTCCTGTTTTAAGAAAT
ATCTGCTATCCCAAGGTAAGATACTATACTTTTTTTAACATGTTATTTTGTTTTACCTTTCA

CATTGTGTATACTACATTTGTAATTAATTTGTCCATATTATATGACATATAGCCAAGATT
 TATTTTTTACCATACAGATACTCAATATTGCCATTTACGTAGAACATCGTTCTCTTCTACT
 CAATTGCCTTGGCACCTTTGTAATAAATCAGATGATCGTGTATGTGTAGTTTCAGTTTCTGGA
 CTCTGTCTTCTGTTTTCTTTAGTCTATTTGCCTATTCTTGTACCAATATGCACTGTGTTAATA
 ATCGTAGCTTTGTAGTAGGTCTCGAAATCTGACAGTGTAAATCTTTTAGTTTCTTTCTTCTG
 CAAATTTTCTTTAGCTGTTTTACGTCCCTTGCATTTGTATATAAACTTCAGAATCAGCTTGT
 CTATTCCAAAAACAACAACAACAAATGAAAGTTTCAGAACTTTAACTGAAATTGTATTGAA
 TCTGTAGACAATTTTGGAGTAAATTGCATCTTAGCAATGTAGAGTCTTTCGAACCATAACCA
 TGGTAAGTCTCTCCATTTGCTTAAATCTTCTTTAATTTATTTCAACAATGGCTTCCAATTTT
 CAGCGGGAGCTCTTGGAACTACAATTTACATGAACTTCTAATTTGATATTTTTCGGTGTCA
 TTATAAACATTGTTGTTTTAAAAGTTGTCTTCAATTTTTTGTGTGTCAGGCACAGAAATACAA
 TTATTGATAACATTTATATATAAACTGTATCCAGTGATCTTGCTAGATTCACTGATGAGTCT
 GATGTTGTAGATTCTTTGGGATTTTCTCCGTACATAATCATATCCTTTCTGAATAAGATAGT
 TTTACTTCTTCATTTTTAATATCTATGCCTTTTATTTATTTTTCTTTTGAACCTTTTGTCTGA
 CTTCAATTATTCATCTCATGTTTTTCTCTTTCATTAGACTATGACTCCTCGATGGTAGCAAT
 TTGTAGTAATCAAATTTTTGTATTTTTATTTTAGCATCTGGCATCTTTCTTGACATATGTAGC
 AGTTGCTTTTTGACAGCTTGCTTCTTCAGTGAATGAATAAATTAATAAAGAGAAATGTGATG
 TTCAGTGATCCATTTTGCAGGTAAGAAAACCTGAGGCAGACAGAGGATGTTAGCAAGCAAGAG
 GCCTTGGCCTACAATTTAGATCACTGGACTCTTACTCCAGATGCAATCTGCAGAACCCACAT
 ACTTTTAATTAGTCCCTTTGTCTATGTTCTGCCACTGTCACCTTCTAAGGAAGGTGTGTCATC
 CCAAATGGGGTAGTATCTTATTGGTAGACCTAAATCTGCTGTGTTTCGCCATCTCACCTACAT
 GAGTATCTATGTGTAGCATTCTGCATATTCATCTTTTCCACCTTCTGGAGGTTTTGTCTTTT
 TATAGGCAGCATGTGAATAACAATGGGGCCAACTGGGGACCAGAAGGGGCCATTTTCTAGT
 TCTGAACATAGATAAGCATCACTAACTTTTCCCTCCTGGCAGTAATGGCCTCAAAAGTTCCA
 ACTTAGGAGAAAAAGGCAAAACCGTCTGCCAAAGTGTGTGAAAAGTTAGAGCAAACCTTGGT
 TTTACCAAGAACCTGTGTCCCTCTTATGGAAATTCACACTTTTACACTTTTAGACAAATATT
 AAATGTGTGACATTCTATTACGTACAGTGCCTGGCACATCTAGACACGCAGCACACTTTAGC
 CCCCTTCTTTCTTCTTCTAACTCCAAGTTCTAACTAGAAAAAGCCCCACTTGAGTCTGAGA
 TTTGCTTTTTTGAAGTAGTTTATTTAGATTGTAATCATGCTATCTAGGGTTGTGACAGTGT
 TGCTATTTCTAGGGCACTTTGACCTGATTCTTTTTGCACAGGAACTTGTCTACCCTTTTG
 CCCACTTCACTGAAGTGAGGACTGAGACAGAGAAGGATTAAGTCACTTTTTATTTAAACAAT
 ATTTGTTCCACATTCCTCAGCATTTATTAATACTGGTGCATAATGATGGAATAAATTTTAT
 ACCGTAAGGATAAACCAGTATCTTGACTAAGCCAACGTGGGAGACCCTAGGAGGCCTGTTT
 GAGAAAGTGACATTTAAATTGAGACTTGGCGGTGGCTGTGGCTACATATCTAGTAAGTGGCT
 GGGTTGGGATTTGAACTCATGCCTGCTTAGCTCTAAAGATGATGCTTTTGGCTTTGTACTCT
 GCTCTCTCTCTAGACAACTCTGGTCCAAAATCGTTAAAGCTAACATTTATCCCTGCCAAC
 TGGAATTGTGATGTTATGACAAATGGCTCTGTGGTCTCAGATGCCAGCAGACCCATTAGTG
 GAATTCTATGTGCTACAGACCTGGGCAAAATGCCAGAGCCTTATACACCCATCACATTTCTG
 CTGGCAAAGGTCTTCAACAAAGAGAAGTAATTACAGCAATGAAAAGCAACAGGTCCAGCAAC
 ACCATAAGAACAATAATTAATTTCCCTAAAATAGAAGAAACCATTTATAGAGTAAGAGCC
 GATACAATCAATAATTGGAAGAAATAGAAGAGGCTTTAGTATTCTAGCCTTCTTTATTTGTA
 GATGTAAATGTCGAGCCTCAGAGAAGTTATATATCTAATTAGTGTCACTCAGGTAGATAACA
 ACAGAATTAGGATTAGAACTTAATTCTTATGACTCCCAGAGCAGGGAAAAGACAGGATGAAG
 TCCCAAAACATTGCGTGTGAACTCACATCTGACTCTGAATTGAGAGTCTGCTATTTACTCCC
 TATGTGACCAGAATCCTTCAGAGCCCATGAGATTCCCTGTCATAGAGTAGATTTTGATCCAC
 ACTAGGCATTTTTACCTCTTGCTCTTTGAGTTGGTGCCCCATGTTTACTCAGAAATATTCCA
 AAGGTGTTACATCTATTGGTTTTACATGTTGAGCACAGATCATTATAAGACAAATTGAAATG
 AAAACATCAACAAGTCTCATTGTTGTTGTTGTTGTTGTTGTTGTTGTTGTTGTTGTTGTTG
 ATCAAGAGAGTCCAAAGTGTGTTGTTGTTGTTGTTGTTGTTGTTGTTGTTGTTGTTGTTG
 ATCACCTGCTCAGGGTAGTGGCAATTCAGGATACAGAGGACAGAAATAAGAATCATGACAC
 ACAATCCACAGAATTCACAGATGCCAAACATCTACCCTTCTCTGTCAACACACATTGGACTC

ACATGGTGGAAATAGGCAACACAAGCAGAGAGGTGGCTTAACCTTTTCATAATTTTTCAACGA
CCTATGGGAAGAGAGTTTTCTTGTTCAAATCCCAGCTTAGCCACACAGAGTGTGGTAATAT
TGGGCAAGTCAACCAAGCTCTCTGTGCTTCAGTTTCCTCATTATTAAAATGGGGGAAATAAT
AGTGCCTGCATCAGAGGGTTGTTGTGAGAATAAACGAAATAATTTATCTGAGCTTTAGAAC
CCACACCATATTAGTTAAAAATTCATGCATTTTCTTTTATTATATTTCTCTACCTTAGACTG
CAAACCTCTAAGAGGAAAGGCCGACTGTTATATTCATAAAGCATTACAGGAACAGTAATTAG
AACTAGGAGCTTTTCAATGGCCTGCCTGAAATCTGAAAAATAGGTATATTATTTGAAATTTT
GAAAAAATCAAATAATTAAAAATTAATAGATGTTAATAAAAAATCTGTAATATGTAATATCA
AGGTCAACTCAACTCTTAATTGTTTATATAAAATATAGTGAAGTTTAAATTGCAAAATCTTA
CAGAAAATGTGCTATTAAAACTCAAAAGTATAATTCTTTCTAATATGTATATATATGTATAT
AGTTTTTATTTTAAAGTTCAGGGGCACATGTGCAAGTTTGTTACCTAGGGAAACTCACGTCACA
GGACTTTGTTATACAGATACTTTCATCACCCAGGTATTAAGCCTAGTGCTCATTAGTTATTT
TTCCTGATCCTCTCCCTCCTCCCAACCTCCACCCTCAGGTAGGCCTCAGTATCTCTTGTTCC
CCTCTATGTGTTTCATGAGTTCTTATCATTTAGCTCCCACTTACAAGTGAGAACATGTGGCAT
TTGGTTTTCTGTTCTGCTTAGTTTGCTAAGGAAAGCACTGTGGCAATTTCTCAAAGGACT
AAAAACGGAATTACCATTCAACCCAGCAATCCCATTAAGTTACGTACTCAAATATTTTTTA
AGGCAAAACAAAGCTGCAACCAGAACACCTGGACTCCCTGAAACCCCTTCCACTGATGTTGT
TGTTGTTGTTTCTTTTTTCCCCAGCTTCTCAGGCCAAAATACTGGATCATCTTGGGCACTGTT
CTCTCCTGCCCCACCTTTCCCATATGCAGAGTGTTGTCACCTTCTCTGCTTCCACTGCTAA
CTCCCTGGTCCAAGCCGCTGCACCACTTTTCGTGATTATTGCCACAGTCTCCTCACTGGTTC
CCTGCCCCCACTCTTGCTCTGAACATCCAGTTAAAACCTGAATTAGATCATCTCATCCTCA
TCTCAGAGCTTTCTCGTGGCTCCTCTGCCCTCTCAGGAAAAAATCTAAATTCTAGATGACCT
AAAAATCCCTTGCTCTCTTACTGTTTATCTGACCTCATTTACTACCACCTTTTTCTTTGATCA
TTCTGTTCCAGCCACACTGGCCTCCTTACCCTCCTCAAATATGCCAAGCACAGCCCCCACC
CCCCAGGGCTTTGAACTGGCTGATCCCCCTTCTGGAATGCCTTACCCCAAATATCAACTTA
GCCAACTCCCTCCTCTCCTCCAAGTGCTGTTTAAACATGGCTTCAGTAGGAGCTGTCTTAA
CATCCTATTAATATTGTAATTCTCTCATGACACTTTACACCCCTTCCCTGATATGCTTTC
CATATACCATGCAATATCTGCTGAGATAATATATAATTCACTTATTTTCTTTATTGTCATTT
CAAAGAGGGCGGTGTGTTCTGTGTTTTATTTAGTGCCAAAATACTTGCTGATGAAGAGAGTT
CCTGCCACATAGTAGGTGCTCAATATGTGCTTGTTGAATAAATGTGTCAATGTTTGATGTAC
AGACCTTTTATTATGTTTGATTTGCTGCCAGTGCTGCCCTCCAAACACAGGAGTGCTTCATGA
GATGTTTCAAAAAGCTCTTAAAATATTCACAAAAATCTTAAAATATTTTCATGAGTTTTCTT
TCCTGTATTTTTATAGCAGCATCTGGAATTTAGCCTGCATAGGACCCTCTGTAAGCTGACCC
TGTTTATCTATTTCAGCTTTACTTCTCCCCTCTCTCCACTTTGTATTTTATTCTCTACTACTT
CCAACCTGATTGTAATTTGACCAGACTCCAGACTATCTTATGCCTCTTTGCTTTTGTTTACCT
GTTACTTCTCTCTGGAATTCCTGCCCTTCTTAATTTTTCTGGCCAATTCTCACTCTCTAG
GACTCAGAGGTTTCTCCTCAGGAGACTTCCATGAGTCTCATGTTGAGTTAGGTGACCCCAAT
CCTCTGTTCTTCATAGTCATTTCGCGCATTTATCTAGCTCAGCATTGCCATACTACATTGAA
ATTATTTTCCTTATGTGCCCATCACTCCCCGTAGATTGCAAACCTCCTAGAGAAGGGCTCAACA
GTGAGTGCTGAGGCTGCACAGAGGAGGAAGGCAGCACAAATGATGGAAGGCTTCCTAAAGAGG
TATGTTCCAAGAGCCCCCACTTCCTTTCATGGGAGACTCATGCTGTTACACCTAGACTATCT
AGGGATACATCTAATGTAGTCGTGGAAGAACAAGGAGCTTGAGTACTAGACTGACGTGATT
TTGAATCCTGGCTCCCTATTGACCAGATGTGTGCTTGAACAAGTCCCTGAGCCTCAGTGTC
TTCATCTGCACAGTGAGGATAATGATACCACACTGCATATATGAGGTATCCGGGCACATGTAA
ATGTCCACTACATGCTGATTTCTTCACCCGCTACTCACCCCTGGGAAAGAAGTAGACTCACC
TACTCTTGTTACCCATTTCATTCCCCCTCAGTTGGAAGCATGAGGTGTGCAGCTGCCTGACCT
GGGGGAAGGGCTGCAAGCAGTAGGTGTTGTGATGTTGAGGCTTGTTGACTTCCTCCCA
GGGGCCCAGCTAACAACCTGCCTCTGTTTCTTGATAGTCAAGTTCAACTTTCACTTCTTAGC
ACCACAGGAAGTTGACTGAACATTAACTGAAGTCTCTCTCAAACAGGAGACATCTTTGCCAG
GTCCCTGTACTTCCTAGCCTCATTCCTGCTCTCCCTAGTGAGCAGGCTGCCCTCCCTTCTCG
CCCCAGCACCACTGATAGGCAAGGGTACTCAGAACTACTACCTTGTGGGCCATGTCATGTGC

CAGGAGCTGCACCCAGGACTTTAATACAGTAGTTGGCTCCCACTGAATGTTTATTGTTACCC
CAGGATAAAAAGGGGACACTGTGATCATTTTTCTATTTTGCTGTGATCAGGCTTGGTGAGCTA
AAGTCACCTACCTTCCCAGTCTCTACTAATAGAAGTCATGGATCAGTCCTATTGGTTCTTCT
GTTACAAGGATTGAGAAATTCATAATCATGGAGCTGCATTTACAGGCAGAAGTTTCTTTTATA
GTTTTCTAAGTGTTTCTTTTAGCAACAATGGAGAAAATCAAAGAGGGCAAAGGTGAGGGGAG
AAAATAACATTTCCCTTTCTGTCTTTGCTCTTGTAGTCTTTTGCTTTAGTTTCTTTACTAT
GACTGTGAGGGTGAACTAGTGATCAGAGTGGTCCAGAATGGGTTTGATGAATCTGATTCTG
GTGACACAAGATGAATTGGGTATATGTTTTCCCTAAAGATAGAGAGACAATATAACATAGTCT
TTACATTAATAGACTCTGGAGCCAATTTTTTTAGGTTCACTCTCTTTTCTTTTCTGTTGTT
GATTTTCAACAAACATCTTGCACTCGAATTCATCTCACTGTTTTATTTTCAAAAAATTTAA
TTTGAGAAAGTTAGCTGTATTAATTTTTTCTTTTTTCTAAAAATCTTTTACTAATTGCAATTA
TTTCCATTGATGCTATTCCATTGAAACCATTTTAAACATGGACTCAATAACTTCATTGTATGT
TAATGTTTAAATTTTCAATTTCTTTACCTTCTTGGTTTTCTAGCTGTGTTAATGTGGTTGACC
ACTCATTCTTTGAAGCTCTATTCTCTGGCTACTACAGTATGACACATTTTGTCTCCTTCTT
CAGTCTCTGTCTTCTCCACAGTCTTCTCTTCTTTTATATACCTTTAAATATTAATGTTTCC
CAGAGATATTTTCTTAACTCACTTCTCTACTGATTCTAGGTACTTTTTCTTGATCCAACCTCTT
CTGATTTTACCCATCTCGATGATTCTATAATTTGTATTTTCTGTTTTGATCTCTTTTTCAGTC
TTCCAGACCTAAATATCCAAATGCCTGATGGATAGTGCTTTCTTTTTTACTACCAAGCCCTC
AAAGGCACTATGTTCAAAAGGAATTTGTTCATCAGTCTCACGGCACATAAGCTTCCCTCTGTG
TTCAATCTGGAGACTTGAGAGTCTTCTGTTCCTTCTTCTCTCTTATTTTCTCCATAATCAA
TCACAAAGTCATGTGGATTTTGCTCCTAAATATGTTAACTTCTTTCTCTCTCACTTTTATAT
CCCTTACATCTAGGTATTTTACAGACCCTCAGTCTCTCTCACATAGACTTTGGCAATAACCTTC
TAATATCAGTCAACCTGACCAATAGGCCACCAGTGCTTCATGTAGAATCTGGACAATGTAGA
GCACTGAGAATGCTCACACTGGTCATATATGTATGAGTTGGTATGACATCTAGGGAAGTTGA
AGACTTACATAGCCTTTGGCCCAGCAATATACACCATAATACATTAGAGAACTCTAGCATG
TGTACACAGTGATATACACACAAGAATGTTTCACAATGCCATTATTTTAAATAGCAAAATTGTG
GAAACAACACAAATGTTTATCAATAACAGAATGGATAAGTGAGCCATGGCATAGTCATACAA
TGAAAAATAATATAATAGTCAAAATGAATGATCTGAAGAGATATCATTATTGGCAATCTTAT
AAAAGACTGAGTTAAAAATGCAATTTGTGAAAATTTTTAATTATTTGATATTATTTAATGCA
AAGTTTTAGAACATGCAACAACCTGTATATATTATTTATGTATATATGCAAATTCAGCAATA
GCATTTAATCATGCCTGGGAATGATAAGTATCAAAGTCAGAAAGTGGTTACCCTTGGGAAGA
GAGGTATGTATCAGCGGTGGGGCACATAGGATGTTGCAGCCATATCTGTAATGTTTCTTTGC
TTTAAAAAATTTGAATCAAGCTTGGCAAAGTGTGACATTTGATTAAGCAGGATAGTGAGTGC
ATATCTGTTACTTATATTGTTCTTTATAATTTTCTCTATGCTAAAGCATTTTGTAAATTTAAA
AAACCTGACAGTGTTACTCCCATGCTTAAATATGCCAGTGGTCAAACCAAATCCAGCAGCA
CATCAAAAAGCTTATCCACCATGATCAAGTGGGCTTCATCCCTGGGATGCAAGGCTGGTTCA
ATATAAGCAAATCAATAAATGTAATCCAGCATATAAACAGAACCAAGACAAAAACCACATG
ATTATCTCAATAGATGCAGAAAAGGCCTTTGACAAAATTCACCAACTCTTCATGCCAAAAAC
TCTCAATAAATTAAGTATTGATGGGACGTATCTCAAAATAATAAGAGTTATCTATGAAAAAC
CCACAGCCAATATCATACTGAATGGGCAAAAACCTGGAAGCATTCCCTTTGAAAACCTGGCACA
AGACAGGGATGCCCTCTCTCACCCTCTTATCAACATGGTGTGGAAGTTCTGGCCAGGGC
AATTAGGCAGGAGAAGGAAATAAAGGGTATTCAATTAGGAAAAGAGGAAATCAAATTGTCCC
TGTTTGCAGATGACATGTATATCTAGAAAACCCCATTTGTCTCAGCCCAAAATCTCCTTAAGC
TGATAAGCAACTTCAGCAAAAGTCTCAGGATATAAAAAATCAATGTACAAAAATCAGAAGCATT
CTTATACACCAACAACAGACAAAACAGAGAGCCAAATCATGAGTGAACCTCCCATTCACAATTG
CTTCAAAGAGAATAAAATACCTAGGAATCCAACCTTACAAGGGACATGAAGGAACCTTTCAAG
GAGAACTACAAACCACTGCTCAATGAAATAAAAGAGGATACAAACAAATGGAAGAACATTCC
ATGCTCATGGGTAGGAAGAATCAATATCGTGAAAATGGCCATACTGCCCAAGGTAATTTATA
GATTCAATGCCATCCCCATCAAGCTACCAATGACTTTCTTACAGAATTGGAAAAAACTGCT
TTAAAGTTCATATGGCACCAAAAAAGAGCCCGCATCACCAGTCAATCCTAAGCCAAAAGAA
CAAAGCTGGAGGCATCACACTACCTGACTTCAAACCTATACTACAAGGCTACAGTAACCCAAA

CAGCATGGTACTGGTACCAAAACAGAGATATAGATCAATGGAACAGAACAGAGCCCTCAGAA
ATAACGCCACATATCTACAACCTCTCTGATCTTTGACAAACCTGAGAAAAACAAGCAATGGGG
AAAGGATTCCCTATTTAATAAATGGTGCTGGGAAAACCTGGCTAGCCATATGGAGAAAGCTGA
AACTGGATCCCTTCCCTTACACCTTATACAAAAATTAATTCAAGATGGATTAAAGACTTAAAT
GTTAGACCTAAAACCATAAAAACCCTAGAAGAAAACCTAGGCATTACCATTTCAGGACATAGG
CATGGGCAAGGACTTCATGTCTAAAACACCAAAAGCAATGGCAACAAAAGACAAAATTGACA
AAGGGGATCTAATTAACTGAAGAGCTTCTGCACAGCAAAAGAACTACCATCAGAGTGAAC
AGGCAACCTACAAAATGGGAGAAAATTTTCACAACCTACTCATCTGACAAAGGGCTAATATC
CAGAATCTACAATGAACTCAAACAAATTTACAAGAAAAAAACAAACAACCCCATCAAAAAAGT
GGGCAAAGGACATGAACAGACACTTCTCAAAAGAAGACATTTATGCAGCCAAAAAACACATG
AAAAAATGCTCATCATCACTGGCCATCAGAGAAATGCAAATCAAACCACAATGAGATACCA
TCTCACACCAGTTAGAATGGCAATCATTAAAAAGTCAGGAAACAACAGGTGCTCGAGAGGAT
GTGGAGAAATAGGAACACTTTTACACTGTTAGTGGGACTGTAACTAGTTCAACCATTGTGG
AAGTCAGTGTGGCGATTCCCTCAGGGATCTAGAAGTAAATACCATTTGACCCAGCCATCCC
ATTACTGGGTATATACCCAAAGGACTATAAATCATGCTGCTATAAAGACACATGCATACGTA
TGTTTTATTGTGGCACTATTACAAATAGCAAAGACTTGAACCAAGCCAAATGTCCAACAATG
ATAGACTGGATTAAGAGAATGTGGCACATATACACCATGGAATACTATGCAGCCATAAAAAA
TGATGAGTTCATGTCCTTTGTAGGGACATGGATGAAATTGGAAATCATCATTCTCAGTAAAC
TATCGCAAGGACAAAAACCAAACACCGCATGTTCTCACTCATAGGTGGGAACTGAACAATGA
AAACACATGGACACAGGAAGGGGAACATCACACTCTGGGGACTGTTGTGGGGTGGGGGGAGG
GGGGAGGGATAGCATTAGGAGATATACCTAATGCTAAATGATGAGTTAATGGGTGCAGCACA
CCAGCATGGCACACGTATACATATGTAACCTAACCTGCACATTGTGCACATGTACCCTAAAAC
TTAAAGTATAATAATAATTAAAAAAACCAATAGTTTATGAAACCCCCCCCCAAAAAAATATA
TGCCAGTGGCCTCCAGTTGCCACCAGGTAGCATCCACATTCTTTAATGGAAAGCCCTTCCT
TGCTTCGAACCTTGCCAACTGGGATTGGACATTTGTAGTTGCATTTCTAAGAACTGTTCCCTT
TTGTCAATGGAGCCTGATTTCCACTTGGATATCTGGGTGATTTAGGGAACTGACCTCAAAA
CCCAATTCTACATTTTCGACCATGTGACCTTGGCTTAATCAATTCACGCATCTTTTTCCCTCA
CCTCAGGGGATGATCATATGAACCTAAGCCAGTTGCAATAGAGTAAACCTCATGTTCCCTAATG
AGAAATCCAGAACAAAATGCTTTATTTTTCTTCAATTTTTTATTAGGTCATCTCCTGAATCA
ATTAATAAAAAAAACCAACAGTGACAACAAAACCTAAAAATATGAAGAAGCTGAAACATGAAA
GCTCTGCCAACTGCAATATGTAGCTGCTAAGGTTGCTGTATTTATTGGAATCAAGCAAGTGT
TCCAGTAAAGAGCACAGAAGATGTGTCTGGGAGCCTTTATGTGTAGGTCTGCAAGTGGTGG
TATCACTACTACTCACACGCCATTGGCTAGAAGTGAAGTTGCATGGATACACCTAATTGTAAA
AGAGGCTGGGAAATAGAGACTATTGTGCCCAGAAAGAAGAGAAAATTCATTTATGGAAGAGG
TAGCTAGTCTCTCACAGCCATGAAAAGAGAAGTGTTTAGCTAATTGAAGTGAATAGCAGCCA
TCTTGGGTCCCTAAGGCAAGTTAGACTAATATTGAAGTGGAAACCATGAGGAAAGCAGTGAT
ACTGAAAGTAACCGCATCTTTGAGAACATGCATTTCATTTCCCTACAACATGAATTTATTGAGG
ACCTACCTTAATACAGGCAGCGTGCTAGACACCAAGAGAACTGATGTCTCTCTCCTTCCTGC
CTGCCTGGAGCCTGTATTCTGGAGGGGACAGAGCTAGCAGATCAGACCTAACTGGAAATCTG
CTGTGCCAGTATATATTTTCAAGTGTGTGAGCCAATATATCCCCTTGATTGCTCAAAGTAGTT
TGGTCGATATATTTTGTGCTTTAAATTGAACACATTCTTATGTACAGCCTCTGTCTCCTCA
TCTCCAACCAAGCAAAATAGCTTGTTCTCTTTATGCAGGGACACATGACATTTCCACAGTGG
CTTTGTGCATATCTCCACCTCAATTTAAATGCCTTCCAATCCCTGCTCAAAAGTCAAAACAGC
TTAATAATGGTAGACATAGAATTTGACTTATTCTAATAATAGGTCTTTTAAACAATGCCTTC
TTCTCTTCATTCTTTCTTCTTAGAGTGGGTATTCTTTCTGGTGCATCATGTAAAGGAAGGT
AACTACATGCATGTAATGATGAGAATATTTATATGTATTTATGATTATCACAAAAAACAAA
GATTCTACCATTCAAGAGGAACATTTATTTTATTTTTTTATTTTGAGAAAAGTATAATTTTAT
TTATTTATTTATTTGTGCAATTTATGGGGTACTTGAGAAAATGTGTTACATGTATATAATG
TGTAGTGATCCAATCAGGATACTAAGGGTGTCCATCACCTGAGTGTATTACATTTTTTGTAA
GTATAATCATCTACTCCAGGAGAACATTTTAAAACTGTTCTGTAGAGATACTACTCAAAT
TAAGTTCTCAGTCTGAAACATCAGATCAGCTAGGAATCTGACAAAAATGCAAGTTCTCAGA

TGACAGATGAGACCACTTCAATCAGAATTTCTGGAGTGGAGCCCACACATTTGTATTTTGC
AACCTTTCCAATGATACTTATGTACATGCTCAAGCTTGAAAACCACTTTCCTAGGACATTAG
TTCTCGACAAGATTTGTGAGTAACCTTGTTTCATGAAAAAGTGTTTAGGAGATACTGATTC
AATAAAAACTAATCAGGCTTTTATTGTTTGCAGGGCTTTCAAACTTGCAATAGGCCACTGT
GCATTGTTAATTTCTAAGAGGAAGATGCTTATGTCCTCAATGAATATCTTCCCACCATGAAG
TACTCTTCTTCCCCCACTTTTTAAACAATTACTAACACCTGGCAGAAGTAGGCAGACAGCT
TACAGCTTAGAAAAAGTTGGCCTAAGATAATGGCTAATTTTCATACATTATTTATTTGTCAT
CATGCTTATCTTTCTCTCTCTAAATTGTATATTTTCATCTCTGTGATCACAGATTGAGCCTCA
TATTTTCATATCTGCCCCCTGGCCTAATGGTTGTTTACAGAATGAGCTCAATGAATATTGTAA
GTGAGTAGGATTTAATTTATTTGATAAATAGATAACCTTAAGTTTTAAACGGTGGATTTTAC
ATGAGGACATTTACTTACTATTGTTGAGCTGTAATTAATTTTAATACTGTTTAGGTACTCA
TAATAAAGAACAGGATATTTGGAGAAGGAAGACAGTATCATTCTGGTTCTTAGTCTTACCA
GCTTATTGATCATGAGTATATAACCTCTCTGTGGCTCAGTGCCTTTCTCTGTAAAATGGGAA
CACAGTGATGTTACCTCACAGGATTGATTTGTAAAAGGGCTGGATAAGGTTATGAGAATGT
TTTGCAAAGTGATATCGAAAGATTAATTGCAAACCTTCATTTGAATCTTAAATTGTTTGAGAT
AGGTCATGCTATGAATCAACTATGAAGTGCAGATATTGTCAAGATTCAATATTTCTTTCCCA
AGAGCTGAGAGGAGGGGCTGCTTGTTTGTGTTGTTCTTTCTTTTAGAAACATGCCAGGACA
GGCTCATTTTCGGGTTTTCTCTCACTTGCTCATCTTACTTTTTCTTTAGTTTTCTCTATTCA
TTAGGATACAGTACTGTAAAGCTTTATGGCATTTTTTATTTTGTGGGAGATGAATCTGAATAA
AGAATTACAGTTAAATCATTGCTAAGTTTGATGAATGAGCACCAAAGAACTCTTCAAGATGT
CATTTTTAAAGTTTTGTAAATGATTGGCTTTTCTAGTGGTTTTCTCTAAGGAATTTTAATTTTG
AATAATGCATAGAAAAATGTGCGCACACACAAATCATTCAAGTATCCACCTCGAAGGGAAATC
AAAGTGCCTGTGAAGTGAACTTTACCTTTCTATATCACCAGCTTCCTGTTAGAGCAGACTT
TTTCTTTGCTCAAAGTCTAAGCATTGAAGAACTTCTTTTTAGTAGGTAGATTTTTGTGTTTT
TTTGTGTTGTTTTGAGACGGAGTCTCACTCTGTCGCCCAGGCTGTGGTGCAGTGGCACGATC
TCGGCCCACTGCAAGCTCTGCCTCCCGGGTTCACACCATTCTCCTGCCTCAGCCTCCCAAGT
AGCTGGGACTACAGGTGCCAGCCACCACGCCTGGCTAATTTTTTTTGCATTTTTTTAGTAGAGA
TGGGGTTTTCATCGTGTAGCCAGGATGGTCTTGATCTCCTGACCTCATGATCCACCCGCCTT
GGCCTGCCAGAGTGCTGGAATTACAGGCGTGATTTAGTAGGTAGTTTTGAGTAGGGAGTATA
CATTTAAATGCTGAAACTCAGTTAAGGAATAATCTAATACTGTATTCAACTGAACTCAGT
TGAGAAATTTCTTTCCAATAATAAAGGAAAAATCAACTGCAGTAATGAGGGAGATGATTTTGC
TGCTAATTACAACAAATATTTACTACAGACCTGTTATGCACCAGGAAGTGTGCTAAATGTTT
TATACATATAACTTTATCTTGTGCTCCAACAACCTATTACATAGAAATTGCTATTATTCCCA
TTTTCTAGATTAATAAATTGGTTTAGAGGGGTCGTATAGGTGAAACAACCTCACTCAATATCA
CAAGCTGTTATGTGGTGAAGTTTGCATGATCAGTACAGGGTTCTGGTTCATCCCACTCATTGA
GTGGTGTAGTCAAGATCTGGAAGCTCTTCTGGTCTTAGTTTTCTCTAGCCGTGAAGTGACAA
TGATTAGGTCTAATCATAGAACATGAGAGTACATGTGAAAAAATGCCTTTTTTAAAGAGTATG
AAAACTTGAGTTGTAAATTTTCTTGTGGATAATTTATTATTGCTTTTTCTTTTTTAGATAA
CACTAACAAAGTTGACCTTAGAATTGGAGTGCCTGGGTAGAACCTGCTGGTACCACCTGC
TTACTGGCATGCTTCTGATGTGAGTTCAGGAGAAGACACTGGCAAGGACAGCAAAGAACAGG
AGAACTCTAGCTTCCCTGATAGCATTCAAGGTGCTGTCCAAAGCTGACTGTGATGGCACC
CTCCAGACAGACAGCGATGCCACATGTTCAAGATGGCAGAATCACTATCAGCTTCAATTCTT
GAATGACTGCAGAGCAAAATTTCTTACCTGCAACATACACTCTATTTTCAGCCTCCCTGGAC
TGTTACATAATGATACATAAAAAATATTTCTTGTGTTGAGGCATCCCAAATTTGATTTATTTG
TCATCACAGTCTATCCTATGAATATATTTCTGATCCAAATAATGCTAGATTTCATGAGAGTTT
ATAGTCCAGAGATTTTTTTTTTTTTTTTGGAGACAGAATCTGTCTCCAGGCTGGGGTGCAGT
GGCATAATCTTAGCTCACTGCAACCTCTGCCTCCCGGGCTCAAGTGATCCTCCCTCCTTAGC
CTCCATGAGTAGCTGGGACCACAGGTGTGCACCACCACACCCCGCTAATTATTTTGTATTTT
TTGCAGAGGGGGGTTTTACCACATTGCCCAAGCTGGTCTCAAACCTCCTAGACTTAAGGGATC
TGCCTGCCTTGGCCTCCCAAAATGCTGAAATTACACTTGTTAGCCATAGTAATAGTTCCTAG
CCATAGTCCGGAGATATTTTAAAACTTTAGTGTGAGTTATTTCTCTTTGCTATAATTCTCT

CATGCATTTTGTAAAACCTTTTCTGCCATACAGTGGTTTTTGGGAAGTCCTTTTGGACAGACA
TATCACAGGATGGAGAACTAAAAAGAGAGAATGCATTAAAGAAAAAATAGTACTTCGTATT
TCAGAAATTACAAAAGGGTGTTCATGCCCCACAGCATGGTGGGCAGTCATTTGTAACCCACAG
AACATTGAGTTTCTAAAATTTGAGTGTTTTTAACTTACGAAGGCCAACTTTACTGATGAAGT
GATTACCCCAAATATGTGAAATCCATATTTCAAGTACAATAGTAGGAAATTGGAAATGGGGC
TCTTGAGTTTCTGCTTTCAAGTGACCCTCAGAAATTCCTCCATTCACTGCAGAGGTTCTGTT
TCTCCTTGCTTTGTTCTGACTTTACGGCAGAACTAAGCTAATGAGTTAGTTACTATGGGTTA
TCACTTGGATTTGAAGAACCATCATTTCTAGGCATTGCTGCTCAGAGTGGGGTCTGCCCTCA
GGGGAAGCTGTCTAACCAGAGCCTAACCCACTAGGGTTTTATCAGAACCTAACTGACCTGAG
GAAAAGAAATACTCAACTCCAGCAAGCTCCAGCCTTCCAAATGGAGGAGGAGAAAACCCAAC
TCCAGCCCACTTCAGCTATCTTGTCCCACATAAAGGGGAAGTGAAGAAACCTGAGAAGCAG
GCATGAAGTTCACATGTATTCCACACCATTTTCATATTCATCGCATTCCATTCTACACAATT
CCACATCTCCCCGGCCACATCAGCCTGAGAGTAATAAACAGACCTTGCCATCGTATCCCCTT
TTGGAGACACAAGCCAGGAAGGAAGCTCTTCAGTGCAGCTGTGAATAGAGAAATGCTGGCTG
AGGTTTTGGAGACACTGGGTGATCATTTCCAGTGAAATTTTTAGATCCCTGGGGACATGAGC
TGCTTTGTCTTTTCCAAGGTCAGGATAAACAGAATAACTTCAGGCTTCTCTACCCAGAAAGA
ACATGTGGCATAAATATCAACTGCAGAATAAATATGATTAATCTGGTACATGGACGAAGATG
TTTTCTAGGAGATGCTTATCCTGGGATGAGAGCTTTCATAAGCATTGATATTTACATGACTC
TTACCGTGTGTCAGGAAGTCTTAATTGTTTTACCTATGTCAATTCATTTGATAGTCACAA
CCACTGAAAGGAGTAAGTACTCTTATTACTTCCATTTTACAGATAGGGAACTGAGGTATAG
AGAAGTCAAGTGACTTGCCCGAGGTTATTAACTACTTAATGTCAACACCAGGATTTGAACC
CAGATCATTTGTTTTCTGAAGTACATGCTCACAATCACTGCATTACTGATACACTGTTTTGTC
TTTGCATACTTAAGTGGTCATAACTTAGTCTGAAACACTTTGTGAGAGCAGGAAGCAAACCTG
TCCCCTTATTAGGTGGACCAGTATAGTGATAATACAAAAGTGTATTGCATTTGAATTACTTG
CTAATATCTTCTAATTGAGGCAATTTTGAACAGGAATATACATATCTAGCCTCTATTTTTCT
AGCTAGAAGTTCTGAAATCCCTGGGCTTAATATTGTATGGCAACAATTGGCTGGAGTTGAGT
TGCTGCCACTCTCTTTTAACTGAGCCATGCTCTCTCTAGTTTTGCTACAGGCCCCACCACTCC
CTATTGCCTCTCCAATACCAGGTCATTTGGCATCTTACTCAGCTCATTTTCATGCACATGTGT
TCCACAATTGGTAACATAACCCTAAAAGTATTTGAGTTTGTGGCTTCTGCTCTTGTGACAGA
AGACTTTTCTCGAATTCCAAGGTCAACATATACCATATTGACTCTGGGCCACATTTTTTAAAT
GTGCTCAAGCTCAGTTTTTCTCCTTAAAAAATGAGGGGTAAAAATAACACTTCGTAGTTGC
CTCATCTGTGGATTGGAAGAATGAATGCCTGTCAATTTCTAGAGTTGTAGTAAGGGCCAGTTG
GGGCAGTGCCTGTGAATATACGCAATGGGCCATCAAGCAATCTCAGGGCTTCAGGCAATGCT
GGGTTTTATAAAGCACTTTATGTTTTAAGTTCATTTTATTTCTAAAGTCTCATTGACTGCT
CTGAAAATCTCTCAAGTAAAGTGGGCACTAAAGGCTTTATACTCTCTCCAATTATACCTTCC
ATTGTATAGATTTGGAAGCTGGGGTCCAAATGTGTTAAATGACTTGCCTAACATGGTCCATT
ACTGTAAGTACAGAAACGGAATGAGACCCCAAATCTGCTTCATGGAGGGACACTCTTCTAAG
ATACTGTGATGCTTCTTCCCAAGTAATTCCGTCTTCAGACTTCAAGGTCTCAATTCTGAATGA
CAATTCAATATAGGACTTTTCATAATCTTAAAAGCAACCTGACAGTCATTACAGTGGGCTGTG
AGAAATACTAACGCAGAGCCACATACTCTGGGCTTCATGCTAGGTTTTGCCACTCACTATCT
TTGCTGAAAAAGTTTTGGAAGCCCTCCTAAGCAGGTGCCAGACCCTTTTTTGGCCAGAGGAC
AGGATCTTACGCTGTTGCCAGGCTGGAGTGCAATAGTGTATCATGGCTCACTGTAGCCTG
GAACTGCTGGGCTCAAGGGATCCTCCCAGCTCAGCCTCCCAAGTAGCTAGGAGTAAAGGTAC
ATGCCATCATGCCTAGCTTTTTTTTTTTTTTAAATTTTTTGGAGAGATAAGTTCCCATCATGTTG
CCCAGGCTGGTCTCTAACTCTTGGACTCAAGTGGTCTCCTCCACATCAGCCTCTCAAAGTGCT
GAGATTGCAGGTGTTAGCCACTGCACTTGGCCTGGCCCCAGTATTCTTTGGGATCTGGAGTT
TGCTGTTGAATGAGAAGGCAAGATGAAATTCCATGTAGTCAGACTCCTACGCTGCTGTTCTA
AGCAGGGTTGGGCTGATTAGTAGGTTATAGATGATGTTTTTCTGTGGTGCTATCTGGACCT
AGTGCTCTTTGGCATCTGGGAAGGTATGGCCTTTAAAAAGCAAACCTGCCATGAGAACTGCTC
TACCCCAAATTTTTGGTTCACAGCCTTCATTTGATTATGTATTGGGGCAAAAATAGTTTAGCC
ATGTGAACCTGTTTGTAACCTGGTGAGTTTCTATTGCTATTTTCATAGCTAAAGTTTTGAGGT

AAATGCTATTGGATCTTTGTGTCTGTGTGTGTATACATATTTAGATTTTTTTTTTTTTTTTTT
TTTTTGGTTCTTTTGAACATTGCTGATTCTTTTTGTGTTTTGTTTTTCAGAGTCTGGAGAACAC
TTTTTCTTTTGAGCTGTTTACAACCTTTAGCAATTGAGTAGAGTGACTCTTGTCAATAGAA
TTTGAAGCACATTTCTCTCTCTGCCTGATTTCTGTAGAATTTGTAAACTATTTGTGAATATT
CTTAATTTATGGCAATATGGTTGCTTACATAAGTTCAATAATAATCTGTTTTCTTTTACAAT
GAGACACAGTTGGAGGAAGTGGTTATTTTCCCAGGGCTTTGACTGAAATGGCCTTGTGAATG
GTTCCAGGAAAGCCAATTTTGGAGACCCTATGTGGATGATGATGCTTGCTGTACTTTCTGTG
GGTAATCGGGCCAAGTATATGGGACTGAAGCTAATTTTGCAGGGCAACATAGAGAGACTTGA
GTTCCAGGGGGAAAGTTTTAGGATGGAGAGAAGCCCTCACTGGACTGTGATGTGGGGTGAGA
AAATGGAGGTCCAGAGAGAAAAGTGAAGTCAACCAAGCTGAAAACAGTTGCAGGAAGAAAAGC
CAGGACAGAACTGGGCTGTTAGAAACCTTGGTAGTGCACCTTTTGATTTCACTCTTAAAATGC
TCAATATGTCTTTTTCTGACAACATGTAGGGAGACCTGAGGTTGGACCTAAAGACACCATATA
TTGGATTCAAGTCTTACACAAGTTAATCTAAATTATGAAGGAGTACACTAAAGATGAGGAA
CTGAGTCACCGCAATGCCAAATGTAACACGCTGTGGGTCAAAAATTTGGATGGTTGCGTTGGA
AATTATTCTGAAGCCACAGACAGGGCTAAAAGGAGTCTAGAATGTCCTACTAGCTGTTGGAA
CCTGCTGGAGTTCTAAAGGCATGCGATGTCCTGAATGTGCTTGACCAATGGACACAGCAGGT
ATTTGAAGAGTTTTGATGCTTCCCTTTCTGTTTTTCAAGTCTATTTTTTTGCTTTATTTCTTTT
AAAAAATTTTCCCCACAGAGGCTATTTATTACACAGCAGCAGCCAGCATGCACCAAGCAGAA
CTGCTTTTTCATAATAAACCTATTTATCTTCTGGCATGTCAGAAATTTCTCAAATTGATACTT
ATAATAAATATGATTAATAACACATTGTATTTTCAAGTGTGCGTAAAATAATTGGGAATAA
ATATCGTATTGTGATGTTGACAGTGTGAGGTTGGTTGGTGCCAAACAGCGGGAGCAGCCAAC
TGTATTTTCATTAACAGCCACATTGCTACAATGCTATTAAGTTTTTTCATAATCTCTTCTGCTC
AAGGTTGATCATTCACTTCTGTTAGACAAATCTTGCAACAATCACACTGGAAGTAAATCTGC
CAGCGACAATCTCAGCAAGATGGGTTCTCAC-5'

<128128655>(SEQ ID NO: 2)

>CCAT1_JAX_3 Transcript sequence; Genomic location:

chr8:128152989-128231094 strand:-

<128231094>

3'-

GTTGCACTGACCTGGCCAGCCCTGCCACTTACCAGGTTGGGGCTTTGACTGAAATGGCCTTG
TGAATGGTTCCAGGAAAGCCAATTTTGGAGACCCTATGTGGATGATGATGCTTGCTGTACTT
TCTGTGGGTAATCGGGCCAAGTATATGGGACTGAAGCTAATTTTGCAGGCAACATAGAGAGA
CTTGAGTTCCAGGGGGAAAGTTTTAGGATGGAGAGAAGCCCTCACTGGACTGTGATGTGGGT
GAGAAAATGGAGGTCCAGAGAGAAAAGTGAAGTCAACCAAGCTGAAAACAGTTGCAGGAAGAA
AAGCCAGGACAGAACTGGGCTGTTAGAAACCTTGATGAAGGAGTACACTAAAGATGAGGAAC
TGAGTCACCGCAATGCCAAATGTAACACGCTGTGGGTCAAAAATTTGGATGGTTGCGTTGGAA
ATTATTCTGAAGCCACAGACAGGGCTAAAAG-5'<128152989>(SEQ ID NO: 3)

>CCAT1_JAX_4 Transcript sequence; Genomic location:

chr8:128160497-128232653 strand:-

<128232653>

3'-

AAGGATTTAATGGCAAGATGCCATTATAGACAAGAACAGGATTCAGACACTTTCGTGTTATG
TGTTCTTGTCCAAATACTGTGTCACTCTGCACTGGCATCCCAATCCCACCAACACCTTTAGC
AGGAACTTCCTGTTTCAATAACATTTCTCAATACTCTACCTGATTGCTTAGAATCCATGAGA
CAGCCATATTCTCCATGTCTAGGTCCTATCTTATTTTTGTTGAGATGGTGTTAAGACTTTA

TTTGTGAGGCTTCTGGGGAAGAGGAGTAAGGTATTGATCCCACTGACTGGATAATTTGGGTC
 TCAAAATGGATAATAAATAAGCATTACATATTTTGACCACTTCCTTGAGGAGAACTTCTTG
 GAATGTGCACCATGTCCGCTGCACTTTTTTTGCACAGATATCTAAGTTGGAGAAACATACTA
 CTAGATAAATCAATTTGTTCTCTTAGTACTCATGATATGGTTCCCTGGGAACCTTCTGATTAC
 CAAATTAATCTTGGCCAGGTACATACCTGGCAGGAATCCAAAAATTCCCCAAGTCTCCTTGA
 AGTTCAGGATCATCATTCTTAATAAATACACCGAGGGAAAAACCATGGAGAGTTTGTCCCAG
 ATGCTGTGAATCTGGCCCGGGGTACATGAAGAAGTCCTTAATTGCAGTCATTTACATGGTAG
 ATTCTCTATAATCATTTAATTTGCTATAGGTCTATGATTTTTAGTCCTTCTTCTCTAAATGA
 TTGAACATGTATAATTCCCATTTCATCATATTACCTGGATGAACAAAAGTAACGCTAGACT
 CATTTCATGCATTCTGGTTGCCAAGGAAAAAGGAAAAAACAACAAATCAACAGGATGTTTA
 AACTGTCTTAGGGCAACTTCAGGCCATAGTCACTGGTGTCTTGCAGACTATGAGATATTTT
 ACATTCTGATAAGGGATAAAAATTCGTGCCTCACATGGCTCCCATCACACTAAGATCTTGCA
 ACAATAACACTACTGATTGAGACATTAATCTTAAGTATCCAGGGAGCCCTAAAACATTGTAT
 CCCACTAGCAAGGACCATGGTAATTGCCACGTAAATCCCCTCCATTATGTGGCCCTTATTAT
 GACCAGCCAGCCAAGGCTTGCCTTTAAATCATACCATTGAACCGAGCCTTGTAGAAACACTA
 TCACCTACGCATACCTCTGCTTCTTTTCATTAACCTGCTATCCTCTTTACAAATGGGATTCT
 TCACCCACTCCCTTCTTCTAGATTAGCAATGCCCTGTAAAGTAAACGAACACGAAATTCAAA
 GGGAAACAGGAGCAATCATCATTACCAGCTGCCGTGTAAAGCATTGCGAAAACGCTCACGAT
 TCACAGAAAAATCCATGCTGTTCTTTGAAGGCATTCAAGCCTTAATAGCTAGCTGGATGAAT
 GTTTAACTTCTAGGCCAGGCACTACTCTGTCCCAACAATAAGCCCTGTACATTGGGAAAGGT
 GCCGAGACATGAACTTTGGTCTTCTCTGCAATCCATCTGGAGCATTCACTGACAACATCGAC
 TTTGAAGTTGCACTGACCTGGCCAGCCCTGCCACTTACCAGGTTGGGCTGTGGCTTGCCTTT
 TCACCTTCTTGTGTCCTGTTTAAGAAATATCTGCTATCCCAAGGTAAGATACTATACTTTTTT
 TAACATGTTATTTTGTTTTACCTTTCACATTTGGTGTATACTACATTTGTAATTAATTTGTC
 CATATTATATGACATATAGCCAAGATTTATTTTTTACCATACAGATACTCAATATTGCCATT
 TACGTAGAACATCGTTCTCTTCTACTCAATTGCCTTGGCACCTTTGTAATAAATCAGATGA
 TCGTGTATGTGTAGTTTCTGACTCTGTCTTCTGTTTCTTTAGTCTATTTGCCTATT
 CTTGTACCAATATGCACTGTGTTAATAATCGTAGCTTTGTAGTAGGTCTCGAAATCTGACAG
 TGTAATTCTTTTAGTTTCTTCTTCTGCAAATTTCTTTAGCTGTTTTACGTCCTTTGCATT
 TGTATATAAACTTCAGAATCAGCTTGTCTATTCCAAAAACAACAACAACAAATGAAAGTTTC
 AGAACTTTAACTGAAATTGTATTGAATCTGTAGACAATTTTGGAGTAAATTCATCTTAGC
 AATGTAGAGTCTTTTGAACCATACCATGGTAAGTCTCTCCATTTGCTTAAATCTTCTTTAA
 TTTATTTTCAACAATGGCTTCCAATTTCCAGCGGGAGCTCTTGGAACTACAATTTACATGAA
 CTTCTAATTTGATATTTTTTCGGTGTCAATTATAAACATTGTTGTTTTAAAGTTGTCTTCAAT
 TTTTTGTTGTGAGGCACAGAAATACAATTATTGATAACATTTATATATAAACTGTATCCAGT
 GATCTTGCTAGATTCACTGATGAGTCTGATGTTGTAGATTCTTTGGGATTTTCTCCGTACAT
 AATCATATCCTTTCTGAATAAGATAGTTTTACTTCTTCATTTTTAATATCTATGCCTTTTAT
 TTATTTTTCTTTTGAACCTTTTGGCTGACTTCATTATTCACCTCTCATGTTTTTCTCTTCATT
 AGACTATGACTCCTCGATGGTAGCAATTTGTAGTAATCAAATTTTTGTATTTTTATTTTAGCA
 TCTGGCATCTTTCTTGACATATGTAGCAGTTGCTTTTTGACAGCTTGCTTCTTCAGTGAATG
 AATAAATTAATAAAGAGAAATGTGATGTTTCAGTGATCCATTTTGCAGGTAAGAAAACCTGAGG
 CAGACAGAGGATGTTAGCAAGCAAGAGGCCTTGGCCTACAATTTAGATCACTGGACTCTTAC
 TCCAGATGCAATCTGCAGAACCCACATACTTTTAATTAGTCCCTTTGTCTATGTTCTGCCAC
 TGTCACCTTCTAAGGAAGGTGTGTATCCCAAATGGGGTAGTATCTTATTGGTAGACCTAAAT
 CTGCTGTGTTTCGCCATCTCACCTACATGAGTATCTATGTGTAGCATTCTGCATATTCATCTT
 TTCCACCTTCTGGAGGTTTTGTCTTTTTATAGGCAGCATGTGAATAACAATGGGGCCAACT
 GGGGACCAGAAGGGGCCATTTTCTAGTTCTGAACATAGATAAGCATCACTAACTTTTCCCTC
 CTGGCAGTAATGGCCTCAAAAGTTCCAACCTTAGGAGAAAAAGGCAAAACCGTCTGCCAAAGT
 GTGTGAAAAGTTAGAGCAAACCTTGGTTTTACCAAGAACCTGTGTCCCTCTTATGGAAATTC
 ACACTTTTCACACTTTTGTAGACAAATATTAAATGTGTGACATTCTATTACGTACAGTGCCTGGC
 ACATCTAGACACGCAGCACACTTTAGCCCCCTTCTTTCTTCTTCTAACTCCAAGTTCTAAAC

TAGAAAAAGCCCCACTTGAGTCTGAGATTTGCTTTTTTGAAGTAGTTTATTTTCAGATTGTAAT
CATGCTATCTAGGGTTGTGACAGTGTTTGCTATTTCTAGGGCACTTTGACCTGATTCTTTTT
GCACAGGAACTTGTTCTACCCTTTTGCCCACTTCACTGAAGTGAGGACTGAGACAGAGAAG
GATTAAGTCACTTTTTATTTAACAAATATTTGTTCCACATTCTCAGCATTTATTAATACT
GGTGCATAATGATGGAATAAATTTTATACCGTAAGGATAAACCAGTATTCTGGACTAAGCCA
ACGTGGGAGACCCTAGGAGGCCTGTTTGAGAAAGTGACATTTAAATTGAGACTTGGCGGTGG
CTGTGGCTACATATCTAGTAAGTGGCTGGGTGGGATTTGAACTCATGCCTGCTTAGCTCTA
AAGATGATGCTTTTGGCTTTGTACTCTGCTCTCTCTCTAGACAACTCTGGTCCAAAATCGT
TAAAGCTAACATTTATCCCTGCCCCAACTGGAATTGTCATGTTATGACAAATGGCTCTGTGGT
CTCAGATGCCCAGCAGACCCATTAGTGGAATTCTATGTGCTACAGACCTGGGCAAAATGCCA
GAGCCTTATACACCCATCACATTTCTGCTGGCAAAGGTCTTCAACAAAGAGAAGTAATTACA
GCAATGAAAAGCAACAGGTCCAGCAACACCATAAGAACAAAATAATTAATTTCCCTAAAATA
GAAGAAACCATTTATAGAGTAAGAGCCGATACAATCAATAATTGGAAGAAATAGAAGAGGCT
TTAGTATTCTAGCCTTCTTTATTTGTAGATGTAAATGTCGAGCCTCAGAGAAGTTATATATC
TAATTAGTGTCACTCAGGTAGATAACAACAGAATTAGGATTAGAAGTTAATTCTTATGACTC
CCAGAGCAGGGAAAAGACAGGATGAAGTCCCAAAACATTGCGTGTGAACTCACATCTGACTC
TGAATTGAGAGTCTGCTATTTACTCCCTATGTGACCAGAATCCTTCAGAGCCCATGAGATTC
CCTGTCATAGAGTAGATTTTGATCCACACTAGGCATTTTTACCTCTTGCTCTTTGAGTTGGT
GCCCCATGTTTACTCAGAAATATTCCAAAGGTGTTACATCTATTGGTTTTACATGTTGAGCA
CAGATCATTATAAGACAAATTGAAATGAAAACATCAACAAGTCTCATTCAATTGTCTAACTTA
CGCTGAGCAATATTTAATAACTAGAAATATCAAGAGAGTCCAAAGTGTTTGCCCATCCCCTCA
AGACCAATGTAATGGAATTTTACTCTTATCACCTGCTCAGGGTAGTGGCAATTCAGGATACA
GAGGACAGAAATAAAGAATCATGACACACAATCCACAGAATTCACAGATGCCAAACATCTAC
CCTTCTCTGTCAACCACACATTGGACTCACATGGTGGAAATAGGCAACACAAGCAGAGAGGTG
GCTTAACCTTTTCATAATTTTTCAACGACCTATGGGAAGAGAGTTTTCTTGTTTCAAATCCCA
GCTTAGCCACACAGAGTGTGGTAATATTGGGCAAGTCAACCAAGCTCTCTGTGCTTCAGTTT
CCTCATTATTAATAATGGGGGAAATAATAGTGCCTGCATCAGAGGGTTGTTGTGAGAACTAAA
CGAAATAATTTATCTGAGCTTTAGAACCCACACCATATTAGTTAAAAATTCATGCATTTTCT
TTTATTATATTTCTCTACCTTAGACTGCAAACTCTAAGAGGAAAGGCCGGACTGTTATATTC
ATAAAGCATTACAGGAACAGTAATTAGAAGTAGGAGCTTTTCAATGGCCTGCCTGAAATCTG
AAAAATAGGTATATTTTGAATTTTGAAAAAATCAAATAATTAATAATTAATAGATGTTA
ATAAAATATCTGTAATATGTAATATCAAGGTCAACTCAACTCTTAATTGTTTATATAAAATA
TAGTGAAGTTTAAATTGCAAAATCTTACAGAAAATGTGCTATTAAAACTCAAAGTATAATT
CTTTCTAATATGTATATATATGTATATAGTTTTATTTTAAGTTCAGGGGCACATGTGCAAGT
TTGTTACCTAGGGAACTCACGTCACAGGACTTTGTTATACAGATACTTTTCATCACCCAGGT
ATTAAGCCTAGTGCTCATTAGTTATTTTTCTGATCCTCTCCCTCCTCCCAACCTCCACCCT
CAGGTAGGCCTCAGTATCTCTTGTTCCCCTCTATGTGTTTCATGAGTTCTTATCATTTAGCTC
CCACTTACAAGTGAGAACATGTGGCATTGTTGTTTTCTGTTTCTGCCTTAGTTTGCTAAGGAA
AGCACTGTGGCAATTTCTCAAAGGACTAAAAACGGAATTACCATTCAACCCAGCAATCCCAT
TACTAGTTACGTACTCAAATATTTTTAAGGCAAAACAAAGCTGCAACCAGAACACCTGGACT
CCCTGAAACCCCTTCCACTGATGTTGTTGTTGTTGTTTCTTTTTCCCAGCTTCTCAGGCCA
AAATACTGGATCATCTTGGGCACTGTTCTCTCCTGCCCCACCCTTTCCCATATGCAGAGTGTT
GTCATTTCTCTCTGCTTCCACTGCTAACTCCCTGGTCCAAGCCGCTGCACCACTTTTCGTGA
TTATTGCCACAGTCTCCTCACTGGTTCCCTGCCCCCACTCTTGCTCTGAACTATCCAGTTAA
AACCTGAATTAGATCATCTCATCCTCATCTCAGAGCTTTCTCGTGGCTCCTCTGCCCTCTCA
GGAAAAAATCTAAATTCTAGATGACCTAAAAATCCCTTGCTCTTACTGTTTATCTGACCTC
ATTTACTACCACCTTTTTCTTTGATCATTCTGTTCCAGCCACACTGGCCTCCTTACCCTCC
TCAAATATGCCAAGCACAGCCCCCAGGGCTTTGAACTGGCTGATCCCCCTTCCCTG
GAATGCCTTACCCCAAATATCAACTTAGCCAACTCCCTCCTCTCCTCCAAGTGTCTGTTTAA
ACATGGCTTCAGTAGGAGCTGTCTTAACATCCTATTAATATTGTAATTCCTCTCATGACACT
TTACACCCCTTCCCTGATATGCTTTCCATATACCATGCAATATCTGCTGAGATAATATATA

ATTCAC TTATTTTCTTTATTGTCATTTCAAAGAGGGCGGTGTGTTCTGTGTTTTATTTAGTG
 CCAAAATACTTGCTGATGAAGAGAGTTCTTGCCACATAGTAGGTGCTCAATATGTGCTTGTT
 GAATAAATGTGTCAATGTTTGATGTACAGACCTTTTATTATGTTTGATTTGCTGCCAGTGCT
 GCCTCCAAACACAGGAGTGCTTCATGAGATGTTTACAAAAGCTCTTAAAATATTCACAAAA
 ATCTTAAAATATTTTCATGAGTTTTCTTTCCTGTATTTTTATAGCAGCATCTGGAATTTAGCC
 TGCATAGGACCCTCTGTAAGCTGACCCTGTTTATCTATTACGCTTTACTTCTCCCCTCTCTC
 CACTTTGTATTTTATTCTCTACTACTTCCAAGTGATTGTAATTTGACCAGACTCCAGACTAT
 CTTATGCCTCTTTGCTTTTGTTTACCTGTTACTTCTCTCTGGAATTCCTGCCCTTCTTAA
 TTTTTCTGGCCAATTCTCACTCTCTAGGACTCAGAGGTTTCTCCTCAGGAGACTTCCATGAG
 TCTCATGTTGAGTTAGGTGACCCCAATCCTCTGTTCTTCATAGTCATTGCGCATTTTATCTA
 GCTCAGCATTTGCCATACTACATTGAAATTATTTCTTATGTGCCCATCACTCCCCGTAGAT
 TGCAAACCTCTAGAGAAGGGCTCAACAGTGAGTGCTGAGGCTGCACAGAGGAGGAAGGCAGC
 ACAATGATGGAAGGCTTCTTAAAGAGGCTGAAAAAGTTTTGGAAGCCCTCTAAGCAGGTGC
 CAGACCCTTTTTTGGCCAGAGACAGGATCTTACGCTGTTGCCCAGGCTGGAGTGCAATAGTG
 TCATCATGGCTCACTGTAGCCTGGAAGTCTGCTGGGCTCAAGGGATCCTCCCAGCTCAGCCTCC
 CAAGTAGCTAGGAGTAAAGGTACATGCCATCATGCCTAGCTTTTTTTTTTTTTTAAATTTTGG
 AGAGATAAGTTCCCATCATGTTGCCCAGGCTGGTCTCTAACTCTTGGACTCAAGTGGTCCTC
 CCACATCAGCCTCTCAAAGTGCTGAGATTGCAGGTGTTAGCCACTGCACTTGGCCTGGCCCC
 AGTATTCTTTGGGATCTGGAGTTTGCTGTTGAATGAGAAGGCAAGATGAAATTCCATGTAGT
 CAGACTCCTACGCTGCTGTTCTAAGCAGGGTTGGGCCTGATTAGTAGGTTATAGATGATGTT
 TTTCTGTGGTGCTATCTGGACCTAGTGCTCTTTGGCATCTGGGAAGGTATGGCCTTTAAAAA
 GCAAACCTGCCATGAGAAGTCTCTACCCCAAATTTTGGTTTACAGCCTTCATTTGATTATGT
 ATTGGGGCAAAAATAGTTTAGCCATGTGAACCTGTTTGTAAGTGGTGAGTTTCTATTGCTA
 TTTTCATAGCTAAAGTTTTGAGGTAAATGCTATTGGATCTTTGTGTCTGTGTGTATACATA
 TTTAGA-5'

<128160497>(SEQ ID NO: 4)

>CCAT1_JAX_5 Transcript sequence; Genomic location:

chr8:128172634-128231094 strand:-

<128231094>

3'-

GTTGCAC TGACCTGGCCAGCCCTGCCACTTACCAGGTTGGTGTATTCCACACCATTTTCATAT
 TCCATCGCATTCCATTCTACACAATTCCACATCTCCCCGGCCACATCAGCCTGAGAGTAATA
 AACAACCTTGCCATCGTATCCCCTTTTGGAGACACAAGCCAGGAAGGAAGCTCTTCAGTGCA
 GCTGTGAATAGAGAAATGCTGGCTGAGGTTTTTGGAGACACTGGGTGATCATTCCCAGTGAAA
 TTTTATAGATCCCTGGGGACATGAGCTGCTTTGTCTTTTCCAAGGTCAGGATAAACAGAATAA
 CTTCAGGCTTCTCTACCCAGAAAGAACATGTGGCATAAATATCAACTGCAGAATAAATATGA
 TTAATCTGGTACATGGACGAAGATGTTTTCTAGGAGATGCTTATCCTGGGATGAGAGCTTTC
 ATAAGCATTGATATTTACATGACTCTTACCGTGTGTGTCAGGAAGTGTCTAATTGTTTTACCT
 ATGTCAATTCATTTGATAGTCACAACCACTGAAAGGAGTAAGTACTCTTATTACTTCCATTT
 TACAGATAGGGAAACTGAGGTATAGAGAAGTCAAGTGACTTGCCCCGAGGTTATTAACTACT
 TAATGTCAACACCAGGATTTGAACCCAGATCATTTGTTTCTGAAGTACATGCTCACAATCAC
 TGCATTACTGATACACTGTTTTGTCTTTGCATACTTAAGTGGTCATAACTTAGTCTGAAACA
 CTTTGTGAGAGCAGGAAGCAAACCTGTCCCCTTATTAGGTGGACCAGTATAGTGATAATACAA
 AAGTGTATTGCATTTGAATTACTTGCTAATATCTTCTAATTGAGGCAATTTTGAACAGGAAT
 ATACATATCTAGCCTCTATTTTTCTAGCTAGAAGTCTGAAATCCCTGGGCTTAATATTGTA
 TGGCAACAATTGGCTGGAGTTGAGTTGCTGCCACTCTCTTTTAACTGAGCCATGCTCTCTCT
 AGTTTGCTACAGGCCCCACCCTCCCTATTGCCTCTCCAATACCAGGTCATTTGGCATCTTA
 CTCAGCTCATTTTCATGCACATGTGTTCCACAATTGGTAACATAACCCTAAAAGTATTGAGT

TTGTGGCTTCTGCTCTTGTGACAGAAGACTTTTCTCGAATTCCAAGGTCAACATATACCATA
 TTGACTCTGGGCCACATTTTTTAATGTGCTCAAGCTCAGTTTTTCTCCTTAAAAAATGAGGG
 GTTAAAAATAACACTTCGTAGTTGCCTCATCTGTGGATTGGAAGAATGAATGCCTGTCATTT
 CTAGAGTTGTAGTAAGGGCCAGTTGGGGCAGTGCCTGTGAATATACGCAATGGGCCATCAAG
 CAATCTCAGGGCTTCAGGCAATGCTGGGGTTTATAAAGCACTTTATGTTTTAAGTTCACCTT
 TATTTCTAAAGTCTCATTGACTGCTCTGAAAATCTCTCAAGTAAAGTGGGCACTAAAGGCTT
 TATACTCTCTCCAATTATACCTTCCATTGTATAGATTTGGAAGCTGGGGTCCAAATGTGTTA
 AATGACTTGCCTAACATGGTCCATTACTGTAAGTACAGAAACGGAATGAGACCCCAAATCTG
 CTTCATGGAGGGACACTCTTCTAAGATACTGTGATGCTTCTTCCCAAGTAATTCCGTCTTCA
 GACTTCAAGGTCTCAATTCGAATGACAATTCAATATAGGACTTTCATAATCTTAAAAGCAAC
 CTGACAGTCATTACAGTGGGCTGTGAGAAATACTAACGCAGAGCCACATACTCTGGGCTTCA
 TGCTAGGTTTTGCCACTCACTATCTT-5'

<128172634>(SEQ ID NO: 5)

>_CCAT1_JAX_6 Transcript sequence; Genomic location:

chr8:128197810-128240377 strand:-

<128240377>

3'-

GCATGTGGCAGGCACAGAAATATTTACTCATTGACTGAATATAGCACATCGTAATGTTGATT
 TTTTTCCAACATAATTTTAGAGCTAGGCATATTGTATTCTATTACACTAGACTATATATCAT
 TCTTAAATAGAACCGCCTTGCTAGATAACACATGTTGGAGGAGAGGCCCTTCTTCTTAGCC
 CTCAGTGTTTCCATCTATGGGGAAGAAGTTCCACCATACTAACATTACTATCGTCTCTCCAC
 CTGCTCACTCACTTCTCCCCAAGGGAGGGGTGTTTCGATATGGTTTCTGAGCTTGGAAAGAAA
 ACTCAGGCATGTGTAACATGGTTCCTTCAGTCCCATGACCCACTGTCCACAAATGGGCTGCT
 CACAGAGTGCATGCCTTCACCCTTGTTCTTGCCATGCAGGAAATTGTATGAAACAGTCCTA
 GCTGAAGCCTGAGATTTTCTGCAATTGCCTAGTCCTGGTGGGTATCTGTCTACTCCTGGAGT
 TTGGATTGGAAAGTCCACATGCCTGAAGGTATAAACCTATTCTACAAAGGGGTGTTTTCTAG
 AATGAAGGTAATATTTTTATCTTACATTTGCAGAAAGAGACAGAACAAATGTTATAGGTGAGT
 GCATGGACAATGACCTCAAACAGCTAAGATTCAAACCCCTGCGTTGAATGATTGAATTGAAA
 TGATTCAATGAGCTAATGTACATAAAGCATCCAGAATGTTGCCTGGCACAAGGACTGTATTG
 TCTGCTAGACCATTTATTCAAAGTGGGAGGATGATGTTCTAAAAGCCAATGATAAAGCTCAT
 GGCAATGCAGGGTATATCTGATGGCATGGAATGCTTTAGGATGGCCAAGATTGCCCATCAAA
 TGCCAAGTCACCGAAGTTATTAATGGCTCTCCTACTAGGAGCCTGACATCATGGTGAGCATC
 GAGAAAGGAATAACCTAAGCTGAAGACACGCCTTTTCAGGAGGCCAAGTTCCACGTTCTGTG
 CATGCTTTTGGCGAAAGTCAGGCCAGGCACTACTCTGTCCCAACAATAAGCCCTGTACATTG
 GGAAAGGTGCCGAGACATGAACCTTTGGTCTTCTCTGCAATCCATCTGGAGCATTCACTGACA
 ACATCGACTTGAAGTTGCACTGACCTGGCCAGCCCTGCCACTTACCAGGTTGGGCTGTGGCT
 TGCCTTTTCACTTCTTGTGTCCTGTTTAAGAAATATCTGCTATCCCAAGATTGCAAACTCCT
 AGAGAAGGGCTCAACAGTGAGTGCTGAGGCTGCACAGAGGAGGAAGGCAGCACAAATGATGGA
 AGGCTTCCCTAAAGAGGTATGTTCCAAGAGCCCCACTTCCTTTTCATGGGAGACTCATGCTGT
 TACACCTAGACTATCTAGGGATACATCTAATGTAGTCGTGGAAGAAGAGAGGACTTGAGTA
 CTAGACTGACGTGATTTTGAATCCTGGCTCCCTATTGACCAGATGTGTGTCTTGAACAAGTC
 CCTGAGCCTCAGTGTCTTCATCTGCACAGTGAGGATAATGATACCACACTGCATATATGAGG
 TATCCGGCACATGTAAATGTCCACTACATGCTGATTTCTTACCCGCTACTCACCCCTGGGA
 AAGAAGTAGACTCACCTACTCTTGGTACCCATTCAATCCCCCTCAGTTGGAAGCATGAGGTG
 TGCAGCTGCCTGACCTGGGGGAAGGGCTGCAAGCAGTAGGTGTTGTCAGATGTGGTGGAGCT
 TGTTGACTTCCTCCAGGGGCCAGCTAACAACCTGCCTCTGTTTCTTGTATAGTCAAGTTCA
 ACTTTCACTTCTTAGCACCACAGGAAGTTGACTGAACATTAAGTGAAGTCTCTCTCAAACAG
 GAGACATCTTTGCCAGGTCCCTGTACTTCCTAGCCTCATTCCTGCTCTCCCTAGTGAGCAGG

CTGCCCTCCCTTCTCGCCCCAGCACCCTGATAGGCAAGGGTACTCAGAACTACTACCTTGT
GGGCCATGTGATGTGCCAGGAGCTGCACCCAGGACTTTAATACAGTAGTTGGCTCCCACTGA
ATGTTTATTGTTACCCCAGGATAAAAAGGGGACACTGTGATCATTCTTCTATTTTGTGTGAT
CAGGCTTGGTGAGCTAAAGTCACCTACCTTCCCAGTCTCTACTAATAGAAGTCATGGATCAG
TCCTATTGGTTCTTCTGTGTACAAGGATTGAGAATTCATAATCATGGAGCTGCATTTACAGGC
AGAAGTTTCTTTTCATAGTTTTCTAAGTGTTCCCTTTTAGCAACAATGGAGAAAATCAAAGAGG
GCAAAGGTGAGGGGAGAAAATAACATTTCCCTTTCTGTCTTTGCTCTTGTAGTCTTTTGCT
TTAGTTTCTTTACTATGACTGTGAGGGTGAACTAGTGATCAGAGTGGTCCAGAATGGGTTT
GATGAATCTGATTCTGGTGACACAAGATGAATTGGGTATATGTTTCCCTAAAGATAGAGAGA
CAATATAACATAGTCTTTACATTAATAGACTCTGGAGCCAATTTTTTTTAGGTTCACTCTCTT
TCCTTTTCATGTGTGTTGATTTTCAACAAACATCTTGCACCTCGAATTCCATCTCACTGTTTTA
TTTTCAAAAATTTAATTTGAGAAAGTTAGCTGTATTAATTTTTTCTTTTTCTAAAATTCTT
TTACTAATTGCAATTATTTCCATTGATGCTATTCCATTGAAACCATTTTAACATGGACTCAA
TAACTTCATTGTATGTTAATGTTTAATTTTCATTTCTTTACCTTCTTGGTTTTCTAGCTGTG
TTTAATGTGGTTGACCCTCATTCTTTGAAGCTCTATTCCCTCTGGCTACTACAGTATGACAC
ATTTTGTCTCCTTCTTCAGTCTCTGTCTTCTCCACAGTCTTCTCTTCCTTTTATATACCTTT
AAATATTAATGTTTCCCAGAGATATTTTCTTAACTCACTTCTCTACTGATTCTAGGTACTTT
TCTTGATCCAACCTCTTCTGATTTTACCCATCTCGATGATTCTATAATTTGTATTTTCTGTTT
TGATCTCTTTTTCAGTCTTCCAGACCTAAATATCCAAATGCCTGATGGATAGTGCTTTCTTTT
TTACTACCAAGCCCTCAAAGGCACTATGTTCAAAGGAATTTGTCATCAGTCTCACGGCACA
TAAGCTTCCTCTTGTGTTCAATCTGGAGACTTGAGAGTCTTCCGTGTTCCCTTCTTCTCCTTA
TTTTCTCCATAATCAATCACAAAGTCATGTGGATTTTGCTCCTAAATATGTTAACTTCTTTC
CTCTCTCACTTTATATCCCTTACATCTAGGTATTTTCCAGACCCTCAGTCTCTCTCACATAGAC
TTTGGCAATAACCTTCTAATATCAGTCAACCTGACCAATAGGCCACCAGTGCTTCATGTAGA
ATCTGGACAATGTAGAGCACTGAGAATGCTCACACTGGTCATATATGTATGAGTTGGTATGA
CATCTAGGGAAGTTGAAGACTTACATAGCCTTTGGCCCAGCAATATACACCATAATACATTA
GAGAAACTCTAGCATGTGTACACAGTGATATACACACAAGAATGTTTCACAATGCCATTATTT
TAATAGCAAAATTTGTGGAAACAACACAAATGTTTATCAATAACAGAATGGATAAGTGAGCCA
TGGCATAGTCATACAATGAAAAATAATATAATAGTCAAAATGAATGATCTGAAGAGATATCA
TTATTGGCAATCTTATAAAAAGACTGAGTTAAAAATGCAATTTGTGAAAATTTTTAATTATTT
GATATTATTTAATGCAAAGTTTTAGAACATGCAAACAACCTGTATATATTATTTATGTATATA
TGCAAATTCAGCAATAGCATTTAATCATGCCTGGGAATGATAAGTATCAAAGTCAGAAAGTG
GTTACCTTGGGAAGAGAGGTATGTATCAGCGGTGGGGCACATAGGATGTTGCAGCCATATC
TGTAATGTTTCTTTGCTTTAAAAAATTTGAATCAAGCTTGGCAAAGTGTGACATTTGATTAA
GCAGGATAGTGAGTGATATCTGTTACTTATATTGTTCTTTATAATTTTCTCTATGCTAAAG
CATTTTGTAAATTTAAAAAACCTGACAGTGTTACTCCCATGCTTAAAATATGCCAGTGGTCAA
ACCAAATCCAGCAGCACATCAAAAAGCTTATCCACCATGATCAAGTGGGCTTCATCCCTGGG
ATGCAAGGCTGGTTCAATATAAGCAAATCAATAAATGTAATCCAGCATATAAACAGAACCAA
AGACAAAAACCACATGATTATCTCAATAGATGCAGAAAAGGCCTTTGACAAAATTCACAAC
TCTTCATGCCAAAACTCTCAATAAATTAAGTATTGATGGGACGTATCTCAAAATAATAAGA
GTTATCTATGAAAAACCCACAGCCAATATCATACTGAATGGGCAAAAACCTGGAAGCATTCCC
TTTGAAAACCTGGCACAAGACAGGGATGCCCTCTCTCACCCTCCTATTCAACATGGTGTGG
AAGTTCTGGCCAGGGCAATTAGGCAGGAGAAGGAAATAAAGGGTATTCAATTAGGAAAAGAG
GAAATCAAATTTGTCCCTGTTTGCAGATGACATGTATATCTAGAAAACCCCATTTGTCTCAGCC
CAAAATCTCCTTAAGCTGATAAGCAACTTCAGCAAAGTCTCAGGATATAAAAATCAATGTAC
AAAAATCAGAAGCATTCTTATACACCAACAACAGACAAACAGAGAGCCAAATCATGAGTGAA
CTCCCATTACAAATTGCTTCAAAGAGAATAAAAATACCTAGGAATCCAACCTACAAGGGACAT
GAAGGAACTCTTCAAGGAGAACTACAAACCACTGCTCAATGAAATAAAAGAGGATACAAACA
AATGGAAGAACATTCCATGCTCATGGGTAGGAAGAATCAATATCGTGAAAATGGCCATACTG
CCCAAGGTAAATTTATAGATTCAATGCCATCCCCATCAAGCTACCAATGACTTTCTTCACAGA
ATTGGAAAAAATGCTTTAAAGTTCATATGGCACCAAAAAAGAGCCCGCATCACCAAGTCAA

TCCTAAGCCAAAAGAACAAAGCTGGAGGCATCACACTACCTGACTTCAAACCTATACTACAAG
GCTACAGTAACCCAAACAGCATGGTACTGGTACCAAACAGAGATATAGATCAATGGAACAG
AACAGAGCCCTCAGAAATAACGCCACATATCTACAACCTCTCTGATCTTTGACAAACCTGAGA
AAAACAAGCAATGGGGAAAGGATTCCCTATTTAATAAATGGTGCTGGGAAAACCTGGCTAGCC
ATATGGAGAAAGCTGAAACTGGATCCCTTCCTTACACCTTATACAAAAATTAATTCAAGATG
GATTAAAGACTTAAATGTTAGACCTAAAACCATAAAAACCCTAGAAGAAAACCTAGGCATTA
CCATTCAAGGACATAGGCATGGGCAAGGACTTCATGTCTAAAACACCAAAAAGCAATGGCAACA
AAAGACAAAATTGACAAAGGGGATCTAATTAAACTGAAGAGCTTCTGCACAGCAAAAAGAAAC
TACCATCAGAGTGAACAGGCAACCTACAAAATGGGAGAAAAATTTTACAAACCTACTCATCTG
ACAAAGGGCTAATATCCAGAATCTACAATGAACTCAAACAAATTTACAAGAAAAAAACAAAC
AACCCCATCAAAAAGTGGGCAAAGGACATGAACAGACACTTCTCAAAAAGAAGACATTTATGC
AGCCAAAAAACACATGAAAAAATGCTCATCATCACTGGCCATCAGAGAAATGCAATCAAAA
CCACAATGAGATACCATCTCACACCAGTTAGAATGGCAATCATTAAAAAGTCAGGAAACAAC
AGGTGCTCGAGAGGATGTGGAGAAATAGGAACACTTTTACACTGTTAGTGGGACTGTAAACT
AGTTCAACCATTTGTGGAAGTCAGTGTGGCGATTTCCTCAGGGATCTAGAACTAGAAATACCAT
TTGACCCAGCCATCCCATTACTGGGTATATACCCAAAGGACTATAAATCATGCTGCTATAAA
GACACATGCATACGTATGTTTATTGTGGCACTATTCACAATAGCAAAGACTTGGAAACCAAGC
CAAATGTCCAACAATGATAGACTGGATTAAAGAGAATGTGGCACATATACACCATGGAATACT
ATGCAGCCATAAAAAATGATGAGTTTCATGTCCTTTGTAGGGACATGGATGAAATTGGAAATC
ATCATTTCTCAGTAAACTATCGCAAGGACAAAAACCAACACCGCATGTTCTCACTCATAGGT
GGGAACTGAACAATGAAAACACATGGACACAGGAAGGGGAACATCACACTCTGGGGACTGTT
GTGGGGTGGGGGGAGGGGGGAGGGATAGCATTAGGAGATATACCTAATGCTAAATGATGAGT
TAATGGGTGCAGCACACCAGCATGGCACACGTATACATATGTAACCTAACCTGCACATTGTGC
ACATGTACCCTAAAACCTTAAAGTATAATAATAATTAAAAAAACCAATAGTTTATGAAACCCC
CCCCAAAAAAATATATGCCAGTGGCCTCCAGTTGCCACCAGGTAGCATCCACATTCTTTA
ATGGAAAGCCCTTCCTTGCTTCGAACTTGCCAACCTGGGATTGGACATTTGTAGTTGCATTTT
TAAGAACTGTTCCCTTTTGTCAATGGAGCCTGATTTCCACTTGGATATCTGGGTGATTTAGG
GAACTGACCTCAAAACCCAATTCTACATTTTCGACCATGTGACCTTGGCTTAATCAATTCAC
GCATCTTTTTCCCTCACCTCAGGGGATGATCATATGAACTAAGCCAGTTGCAATAGAGTAAA
CCTCATGTTCCCTAATGAGAAATCCAGAACAAAATGCTTTATTTTTCTTCAATTTTTTATTAG
GTCATCTCCTGAATCAATTAAAAAAAACCAACAGTGACAACAAAACCTAAAAATATGAAG
AAGCTGAAACATGAAAGCTCTGCCAACTGCAATATGTAGCTGCTAAGGTTGCTGTATTTATT
GGAATCAAGCAAGTGTTCAGTAAAGAGCACAGAAGATGTGTCTGGGAGCCTTTATGTGTAG
GTCTGCAAGTGGTGGATATCACTACTACTCACACGCCATTGGCTAGAACTGAGTTGCATGGA
TACACCTAATTGTAAGAGAGGCTGGGAAATAGAGACTATTGTGCCCAGAAAGAAGAGAAAAT
TCATTTATGGAAGAGGTAGCTAGTCTCTCACAGCCATGAAAAGAGAAGTGTTTAGCTAATTG
AAGTGAATAGCAGCCATCTTGGGTCCCTAAGGCAAGTTAGACTAATATTGAAGTGGAAACCA
TGAGGAAAGCAGTGATACTGAAAGTAACCGCATCTTTGAGAACATGCATTTCATTTCTTACAA
CATGAATTTATTGAGGACCTACCTTAATACAGGCAGCGTGCTAGACACCAAGAGAAGTATG
TCCTCTTCCTTCCTGCCTGCCTGGAGCCTGTATTCTGGAGGGGACAGAGCTAGCAGATCAGA
CCTAACTGGAAATCTGCTGTGCCAGTATATATTTTCAGTGATGTGAGCCAATATATCCCCTTG
ATTGCTCAAAGTAGTTTGGTCGATATATTTTGTGCTTTAAATTGAACACATTCTTATGTAC
AGCCTCTGTCTCCTCATCTCCAACCAAGCAAAAATAGCTTGTTCTCTTTATGCAGGGACACAT
GACATTTCCCACGTGGCTTTGTGCATATCTCCACCTCAATTTAAAATGCCTTCCAATCCCTG
CTCAAAGTCAAACAGCTTAATAATGGTAGACATAGAATTTGACTTATTCTAATAATAGGTCT
TTTAAACAATGCCTTCTTCTCTTCATTCTTTCTTCTTAGAGTGGGTATTCTTTCTGGTGCA
TCATGTAAAGGAAGGTAACCTACATGCATGTAATGATGAGAATATTTATATGTATTTATGATT
ATCACAAAAAACAAAGATTCTACCATTCAAGAGGAACATTTATTTTATTTTTTTATTTGAG
AAAAGTATAATTTTTATTTATTTATTTATTTGTGCAAATTTATGGGGTACTTGAGAAAATGTG
TTACATGTATATAATGTGTAGTGATCCAATCAGGATACTAAGGGTGTCCATCACCTGAGTGT
ATTACATTTTTGTAAAGTATAATCATCCTACTCCAGGAGAACATTTTAAAAACTGTTCTGTA

GAGATACTACTCAAATTAAGTTCTCAGTCCTGAAACATCAGATCAGCTAGGAATCTGACAAA
AATGCAAGTTCTCAGATGACAGATGAGACCACTTCAATCAGAATTTCTGGAGTGGAGCCCAC
ACATTTGTATTTTTTGCAACCTTTCCAATGATACTTATGTACATGCTCAAGCTTGAAAACCAC
TTTCCTAGGACATTAGTTCTCTCGACAAGATTTGTGAGTAACCTTGTTTCATGAAAAAGTGTT
TAGGAGATACTGATTCAATAAAAACTAATCAGGCTTTTATTGTTTGCAGGGCTTTCAAACCT
TGCAATAGGCCACTGTGCATTGTTAATTTCTAAGAGGAAGATGCTTATGTCCTCAATGAATA
TCTTCCCACCATGAAGTACTCTTCTTCCCCCACTTTTAAAAACAATTACTAACACCTGGCAG
AAGTAGGCAGACAGCTTACAGCTTAGAAAAAGTTGGCCTAAGATAATGGCTAATTTTTCATAC
ATTATTTATTTGTTCATCATGCTTATCTTTCTCTCTCTAAATTGTATATTTTCATCTCTGTGAT
CACAGATTGAGCCTCATATTTTCATATCTGCCCTGGCCTAATGGTTGTTTACAGAATGAGCT
CAATGAATATTGTTAAGTGAGTAGGATTTAATTTATTTGATAAATAGATAACCTTAAGTTTT
AAACGGTGGATTTACATGAGGACATTTACTTACTATTGTTGAGCTGTAATTAATTTTTTAAT
ACTGTTTAGGTACTCATAATAAAGAACAGGATATTTGGAGAAGGAAGACAGTATCATTCTCTG
GTTCTTAGTCTTACCAGCTTATTGATCATGAGTATATAACCTCTCTGTGGCTCAGTGCCTTT
CTCTGTAATAATGGGAACACAGTGATGTTTACCTCACAGGATTGATTTGTAAAAGGGCTGGAT
AAGGTTATGAGAATGTTTTGCAAAGTGATATCGAAAGATTAATTGCAAACCTTCATTTGAATC
TTAAATTGTTTGAAGATAGGTCATGCTATGAATCAACTATGAAGTGCAGATATTGTCAAGATT
CAATATTTCTTTCCCAAGAGCTGAGAGGAGGGGCTGCTTGTTTGTGTTTCTTTCTTTTCT
GAAACATGCCAGGACAGGCTCATTTTCGGGTTTTCTCTCACTTGCTCATCTTACTTTTTCT
TTAGTTTCTCTATTTCATTAGGATACAGTACTGTAAAGCTTTATGGCATTATTTATTTGTGGG
AGATGAATCTGAATAAAGAATTACAGTTAAATCATTGCTAAGTTTGATGAATGAGCACCAAA
GAACTCTTCAAGATGTCATTTTTTAAAGTTTTGTAAATGATTGGCTTTCAGTGGTTTCTCTA
AGGAATTTTAAATTTTGAATAATGCATAGAAAAATGTGCGCACACACAAATCATTAGTATCC
ACCTCGAAGGGAAATCAAAGTGCTGTGAAGTGAACTTTACCTTTCTATATCACCAGCTTC
CTGTTAGAGCAGACTTTTTCTTTGCTCAAAGTCTAAGCATTGAAGAACTTCTTTTGTAGTAGG
TAGATTTTTGTGTTTTTTTGTGTTTTGTGAGACGGAGTCTCACTCTGTCGCCCAGGCTGTG
GTGCAAGTGGCAGATCTCGGCCCCACTGCAAGCTCTGCCTCCCGGGTTCACACCATTCTCTCTG
CCTCAGCCTCCCAAGTAGCTGGGACTACAGGTGCCAGCCACCACGCCTGGCTAATTTTTTTTG
CATTTTTTAGTAGAGATGGGGTTTCATCGTGTTAGCCAGGATGGTCTTGATCTCCTGACCTC
ATGATCCACCCGCCCTTGGCCTGCCAGAGTGCTGGAATTACAGGCGTGATTTAGTAGGTAGTT
TTGAGTAGGGAGTATACATTTAAATGCTGAACTCAGTTAAGGAATAATCTAATACTGTAT
TCAACTGAACTCAGTTGAGAAATTTCTTTCCAATAATAAAGGAAAATCAACTGCAGTAATG
AGGGAGATGATTTTGCTGCTAATTACAACAAATATTTACTACAGACCTGTTATGCACCAGGA
ACTGTGCTAAATGTTTTATACATATAACTTTATCTTGTGCTCCAACAACCTTATTACATAGAA
ATTGCTATTATTCCCATTTTCTAGATTAATAAATTGGTTTAGAGGGGTCGTATAGGTGAAAC
AACTCACTCAATATCACAAGCTGTTATGTGGTGAAGTTTGCATGATCAGTACAGGGTTCTGG
TCATCCCCTCATTGAGTGGTGCTAGTCAAGATCTGGAAGCTCTTCTGGTCTTAGTTTCTCT
AGCCGTGAAGTGACAATGATTAGGTCTAATCATAGAACATGAGAGTACATGTGAAAAAATGC
CTTTTTTAAAGAGTATGAAAACTTGAGTTGTAAAATTTTCTTGTGGATAATTTATTATTGCT
TTTCTTTTTTAGATAACACTAACAAAGTTGACCTTAGAATTGGAGTGCCTGGGTTAGAACCC
TGCTGGTACCACCTGCTTACTGCATGCTTCTGATGTGAGTTTCAGGAGAAGACACTGGCAAGG
ACAGCAAAGAACAGGAGAACACTCTAGCTTCCCTGATAGCATTCAAGGTGCTGTCCAACTG
ACTGTGATGGCACCCTCCAGACAGACAGCGATGCCACATGTTCAAGATGGCAGAATCACTAT
CAGCTTCAATTCCCTGAATGACTGCAGAGCAAAAATTTCTTACCTGCAACATACACTCTATTTT
CAGCCTCCCTGGACTGTTACATAATGATACATAAAAAATATTTCTTGTGTTGAGGCATCCCAA
ATTTGATTTATTTGTCA-5'

<128197810>(SEQ ID NO: 6)

>CCAT1_JAX_7 Transcript sequence; Genomic location:

chr8:128186443-128240377 strand:-

<128240377>

3' -

GTAATGTTGATTTTTTTTCCAACATAATTTTAGAGCTAGGCATATTGTATTCTATTACACTAG
ACTATATATCATTTCTTAAATAGAACCAGCCTTGCTAGATAACACATGTTGGAGGAGAGGCC
TTCTTCTTAGCCCTCAGTGTTTCCATCTATGGGGAAGAAGTTCCACCATACTAACATTACTA
TCGTCTCTCCACCTGCTCACTCACTTCTCCCCAAGGGAGGGGTGTTTCGATATGGTTTCTGAG
CTTGGAAGAAAACCTCAGGCATGTGTAACATGGTTCCTTCAGTCCCATGACCCACTGTCCAC
AAATGGGCTGCTCACAGAGTGCATGCCTTCACCCTTGTTCCCTGGCCATGCAGGAAATTGTAT
GAAACAGTCCTAGCTGAAGCCTGAGATTTTCTGCATTGCCTAGTCCTGGTGGGTATCTGTC
TACTCCTGGAGTTTGGATTGGAAAGTCCACATGCCTGAAGGTATAAACCTATTCTACAAAGG
GGTGTCTTCTAGAATGAAGGTAATATTTTATCTTACATTTGCAGAAAGAGACAGAACAATG
TTATAGGACTGTATTGTCTGCTAGACCATTATTTCAAAGTGGGAGGATGATGTTCTAAAAGC
CAATGATAAAGCTCATGGCAATGCAGGGTATATCTGATGGCATGGAATGCTTTAGGATGGCC
AAGATTGCCCATCAAATGCCAAGTCACCGAAGTTATTAATGGCTCTCCTACTAGGAGCCTGA
CATCATGGTGAGCATCGAGAAAGGAATAACCTAAGCTGAAGACACGCCTTTTCAGGAGGCCA
AGTTCCACGTTCTGTGCATGCTTTTGGCGAAAGTCAGGCCAGGCACTACTCTGTCCCAACAA
TAAGCCCTGTACATTGGGAAAGGTGCCGAGACATGAACTTTGGTCTTCTCTGCAATCCATCT
GGAGCATTCACTGACAACATCGACTTGAAGTTGCACTGACCTGGCCAGCCCTGCCACTTACC
AGGTTGGGCTCTTGAGTTTCTGCTTTCAAGTGACCTCAGAAATTCTCCTCATTCACTGCAGA
GGTTCTGTTTCTCCTTGCTTTGTTCTGACTTTACGGCAGAACTAAGCTAATGAGTTAGTTAC
TATGGGTTATCACTTGGATTTGAAGAACCATCATTTCTAGGCATTGCTGC-5'

<128186443>(SEQ ID NO: 7)

>CCAT1_JAX_8 Transcript sequence; Genomic location:

chr8:128218833-128240377 strand=-

<128240377>

3' -

GCATGTGGCAGGCACAGAAATATTTACTCATTGACTGAATATAGCACATCGTAATGTTGATT
TTTTTCCAACATAATTTTAGAGCTAGGCATATTGTATTCTATTACACTAGACTATATATCAT
TCTTAAATAGAACCAGCCTTGCTAGATAACACATGTTGGAGGAGAGGCCCTTCTTCTTAGCC
CTCAGTGTTTCCATCTATGGGGAAGAAGTTCCACCATACTAACATTACTATCGTCTCTCCAC
CTGCTCACTCACTTCTCCCCAAGGGAGGGGTGTTTCGATATGGTTTCTGAGCTTGGAAAGAAA
ACTCAGGCATGTGTAACATGGTTCCTTCAGTCCCATGACCCACTGTCCACAAATGGGCTGCT
CACAGAGTGCATGCCTTCACCCTTGTTCCCTGGCCATGCAGGAAATTGTATGAAACAGTCCTA
GCTGAAGCCTGAGATTTTCTGCATTGCCTAGTCCTGGTGGGTATCTGTCTACTCCTGGAGT
TTGGATTGGAAAGTCCACATGCCTGAAGGTATAAACCTATTCTACAAAGGGGTGTTTTCTAG
AATGAAGGTAATATTTTTTATCTTACATTTGCAGAAAGAGACAGAACAATGTTATAGGTTCTG
GGAAATAAGAAATCATTAGAAAAAGATTTCTGCCTTCTAGAAGTACACAGTCTAATGGTGAG
ATAGGCAGTTATTAATGGCTCTCCTACTAGGAGCCTGACATCATGGTGAGCATCGAGAAAGG
AATAACCTAAGCTGAAGACACGCCTTTTCAGGAGGCCAAGTTCCACGTTCTGTGCATGCTTT
TGGCGAAAGTCAGGCCAGGCACTACTCTGTCCCAACAATAAGCCCTGTACATTGGGAAAGGT
GCCGAGACATGAACTTTGGTCTTCTCTGCAATCCATCTGGAGCATTCACTGACAACATCGAC
TTGAAGTTGCACTGACCTGGCCAGCCCTGCCACTTACCAGGTTGGCTCTGTATGGCTAAGCG
TTTTCTCCTAAAATCCCTTGAAAACCTGTGAGAAGACCATAAGAAGATCATATCTTTAATTCT
ATTTACAAAGTCACACAATATTTCAATCAAATACAGATGGTTGAGAAAAGTCATCCATCTTC
CTCCCCACCCTCCCACAGCCCCTCAACCACTGCCCTGAACTTATATGCTGTTATCCGCAG
CTCCATCTGGAGCATCACAGCTACTGTCAACCCTGACGCTCTTTCTGAAAAAACACCGGATG
GACATCAGAACTATTTCTTTAAGGATGTTACTGAGCCACACAGGAAAACCTTGCCTTATGATT

TTGAATGCACGGATCTGATTTGACTAAACATGATAACTAGAGAATCACCCAATCTACTCCCA
 TTTTCAACTCTAAATCATCAGAGTGTCTCAAATCCAAAGCACACACAGACCAGCCTGGCCAA
 CACGGTGAAACTCCACCCCTACTAAAAGTATAAAAAATTATCCAGGTGTGGTGGCGGGCGCCT
 GTAATCCAAGCTACTTGGGAGTCTGGAGGCAGGAGAATCCCTTGAACCTGGGAGATGGAGGT
 TGCAGTGAGCAGAGATCACACCACCGCACTCTAGCCTGGGCCACAAATCAACAACAACAACA
 ACAACAAAAAACAAGCGCACACAGAGACTGAGGTCCTCTTTGGCATTGAGAAGATGGCTAT
 GCAAGTCCCAACTAGCAAGTGCAAACCTCCCAGCTTCACTTCTGCCAGTGTCCCTTCACCCC
 TTCTCAACCCCACTGGGAGGCAGGAGGGTGCTTGACAATAACAGCCTTGGCATCACTCTGCC
 AGGGTGTAATAGGAAGTGTACAAATTCTGAGATTCTGTGTAAGCACTGGCCTTTCTGCCTAG
 AATGCCTTCTCCTCTCTTTTTTAAGTGCATGCTCCTATTTATCTTTCAAAGCCCGGAAAAAA
 TAACACTGCACACGGGAAATGCTCCCTTCCTACTGCAGTCATTTAGATGACTCTATGCCATT
 CCATTCACTTCTCTTTTCTACCACAGAAGTGCTTTGAGATTTTGGAGTCAGACTGCTTGAAC
 TTGAATCCTGGCCCTCTCATCAGAGACTTGACTTATTTTAGGCAAGTTATATAACCAATTTT
 ACCTCAGTTTCTTACCCATAAAATGGGTCTAATGAGAGTACCTACCACACAGAATTTTGATG
 AAAACTGAATGAGATGAAGGCCTTTAAGGCAGTGGTCCCCAACCCCTGGGGACACAGACAGGT
 ACCATTTTGTGGCCTGTTAGGAACTGGGCCACACAGCAGGAGGTGAGCAGTGGGTGAGTGAG
 ATCAGCGTTATTTACAGCTGCTCCCCATTGCTCACCTTACTGCCTGAGCTCCACCTCCTGTC
 AGATCAGCAGTGGCATTAAATTCTCATAGCAGCACAAACCCTGTCATGAACTGCACATGCGA
 GGGATCTAGGTTGTGCGCTCCTTATGAGAATCTAATGCCTAATGACCTGTCACCGTCTCCCA
 TCACCCCTAGATGGGAGTGTCTAGTTGCAGGAAACAAGCTCAGGGCTTCCACTGATTCTACA
 TTATGGTGAGTTGTATAATTATTTCACTATATAATACAATGTAATAATAATAGAAACACAGT
 GCACAACAAATGTAATGTGCTTGAATCATCCCCAAACCATCCCAGTCCACGGTCTTCCACAT
 TTTGTCTTTTACAAAAATTGTCTTCCACAAAACCTGGTCCCTGGTGCCAAAAAGGCTTGGGAC
 CACTGCTTTAAAGCCTTTGCATAGTGCTTAGAATTGAGGGGGGAAAAAAAAAACAACAAAT
 GTAGCTAGTTGCTACAATCACTATATTGGTGAGTTTCAAAGGAAAAGAATTCTGTCCCAT
 TATGCTTGAGCCTTGAGTTGCTAACCAAGCCTGACACAAAATTACTGTTGAAGGGATGTGTG
 AGTCCTAATTGAAATGAGGCCTCTTAAGGGAATTGTGGACCAAACCCCAAGCAGGCAGAAAG
 CCGTATCTTAATTATTGCAAGTATTTCAAGCAAGGTGTGGATGGCCATTTGAATTCAAGCAG
 ACTAGGACCTGGGATGAGAAAGAAGGTGTGTACGTGACTTGATCTTTGAACTTTAGCTCACC
 ATCTGGAAGAAGGCTGAGTATTCTCTGCACTCACATAGTAGCTAATGCCTACTCCCCAGCCA
 CCCACAATTCTTTCTGTAGGAAGGCTCGCTAGAATACTTTGTGATATTGGATATTAGTTCCA
 TATTCTACTGTGTATCTTAGTTCAACCAAAATTGTAATCATCTGATATTTATTTCTTTTAATA
 TAAATATAAGTATATTAAGTCTTGGCATGCTTGCTCAGTCTCTCTCTCTCTCCCATTCCTCC
 CCGCTCCCCTCTCTCTTTTCCCAACAGGCTTGGAAGCAGGCATCACCATGCCTATTTAACAG
 TTGGGGTCCCTTGCCACCAGGTGCTGGAGTAGGAATCTGAGCCCGGACATGCCTGATCTGT
 AAATTTTGTGTTTTTCCCCACTGTGCTGGGCAGATCACAGCTATCAGCGCCAAATTCATAGAA
 GGGGCGCCCCCTGTGGTCAATTGAGGGATTTGTGTTTGGAGTAGATCTCAAGAAGGAATGGG
 TGGGGAACCTTAGCCTAGGACAGAGCAGAAAGGAGCCCTCACTCCCCAAGCACCAACGGCCTC
 AGTCCTTCCTGCTGACTCCAGCCTCTAGCTCTCACCCAGACTATCTGCATCCTTCTCTCCAC
 CACGCTCCTTTGGAACCTGCGTAAACACAGATTAAAGGAATTCCGCCTTACTTCCCTTTCC
 GCATTATGACCAAAATGGTTTTACACTATCATTGAACAGTTTAGTACAAAACATGCCACCTTT
 TAATCTATTCACTTCACTTAACAAATACTTTGGAGTGTTTACCATGTGCCAAGTGCTGTTCTA
 ATAGACATAAGCTGTGAGGTATGCTTATCTGATTCTCACAGCAACAGCTTTCGAGATATGA
 ATTGGTATACTCATTTGACAGATGAGGAAATTGAATTCATGTAGTGAAAGGAAGAGCTGCAA
 TTCAGGGTACTGGTTTCTCCTGCACTAAGCACTGAGCCACACTAGAAGAGAAGGCATGAGG
 AAGACAAAAGT-5'

<128218833>(SEQ ID NO: 8)

For each of SEQ ID NOs: 1-8, the cDNA sequence “-” strand having the same sequence (except that the U’s in RNA are replaced with T’s in cDNA) as the respective

CCAT1 ncRNA transcript isoform is shown, from 3' end to 5' end. In addition, the first and the last nucleotides of each cDNA “-” strand, as they are mapped to the corresponding nucleotides on the genomic sequence, are also shown (*e.g.*, in SEQ ID NO:1, the first cDNA nucleotide C at the 5' end corresponds to nucleotide 128128655 on Chromosome 8 of the human genome, and the last cDNA nucleotide T at the 5' end corresponds to nucleotide 128241571 on Chromosome 8 of the human genome).

Furthermore, the following table lists additional information for the 8 transcripts, CCAT1_JAX_1 to CCAT1_JAX_8 (SEQ ID NOs: 1-8, respectively), including the start and end nucleotide positions for each exon of each CCAT1 transcript as represented by the nucleotide positions on human chromosome 8., the length of each exon, and the corresponding genomic sequence spans.

Name	Feature	Start	End	Genomic span	Transcript length
CCAT1_JAX_1	Transcript	128128655	128241571	112917	29299
CCAT1_JAX_1	Exon1	128128655	128129210	556	556
CCAT1_JAX_1	Exon2	128152988	128153109	122	122
CCAT1_JAX_1	Exon3	128153590	128153816	227	227
CCAT1_JAX_1	Exon4	128155104	128155178	75	75
CCAT1_JAX_1	Exon5	128156007	128156437	431	431
CCAT1_JAX_1	Exon6	128160496	128161163	668	668
CCAT1_JAX_1	Exon7	128161860	128161917	58	58
CCAT1_JAX_1	Exon8	128172633	128174329	1697	1697
CCAT1_JAX_1	Exon9	128176683	128176771	89	89
CCAT1_JAX_1	Exon10	128181151	128181362	212	212
CCAT1_JAX_1	Exon11	128186434	128186609	176	176
CCAT1_JAX_1	Exon12	128197071	128198015	945	945
CCAT1_JAX_1	Exon13	128200029	128200129	101	101
CCAT1_JAX_1	Exon14	128200289	128215467	15179	15179
CCAT1_JAX_1	Exon15	128218832	128218920	89	89
CCAT1_JAX_1	Exon16	128218922	128221962	3041	3041
CCAT1_JAX_1	Exon17	128231054	128231498	445	445
CCAT1_JAX_1	Exon18	128231499	128231806	308	308
CCAT1_JAX_1	Exon19	128231808	128232653	846	846
CCAT1_JAX_1	Exon20	128234035	128235911	1877	1877
CCAT1_JAX_1	Exon21	128236644	128236720	77	77
CCAT1_JAX_1	Exon22	128236779	128236929	151	151
CCAT1_JAX_1	Exon23	128239643	128241571	1929	1929
CCAT1_JAX_2	Transcript	128128655	128232653	103999	25265
CCAT1_JAX_2	Exon1	128128655	128129210	556	556
CCAT1_JAX_2	Exon2	128152988	128153109	122	122

CCAT1_JAX_2	Exon3	128153590	128153816	227	227
CCAT1_JAX_2	Exon4	128155104	128155178	75	75
CCAT1_JAX_2	Exon5	128156007	128156437	431	431
CCAT1_JAX_2	Exon6	128160496	128161163	668	668
CCAT1_JAX_2	Exon7	128161860	128161917	58	58
CCAT1_JAX_2	Exon8	128172633	128174329	1697	1697
CCAT1_JAX_2	Exon9	128176683	128176771	89	89
CCAT1_JAX_2	Exon10	128181151	128181362	212	212
CCAT1_JAX_2	Exon11	128186434	128186609	176	176
CCAT1_JAX_2	Exon12	128197071	128198015	945	945
CCAT1_JAX_2	Exon13	128200029	128200129	101	101
CCAT1_JAX_2	Exon14	128200289	128215467	15179	15179
CCAT1_JAX_2	Exon15	128218832	128218920	89	89
CCAT1_JAX_2	Exon16	128218922	128221962	3041	3041
CCAT1_JAX_2	Exon17	128231054	128231498	445	445
CCAT1_JAX_2	Exon18	128231499	128231806	308	308
CCAT1_JAX_2	Exon19	128231808	128232653	846	846
CCAT1_JAX_3	Transcript	128152989	128231094	78106	465
CCAT1_JAX_3	Exon1	128152989	128153109	121	121
CCAT1_JAX_3	Exon2	128153719	128153816	98	98
CCAT1_JAX_3	Exon3	128155105	128155178	74	74
CCAT1_JAX_3	Exon4	128156008	128156139	132	132
CCAT1_JAX_3	Exon5	128231055	128231094	40	40
CCAT1_JAX_4	Transcript	128160497	128232653	72157	8066
CCAT1_JAX_4	Exon1	128160497	128161163	667	667
CCAT1_JAX_4	Exon2	128161861	128161917	57	57
CCAT1_JAX_4	Exon3	128209720	128215465	5746	5746
CCAT1_JAX_4	Exon4	128231055	128231498	444	444
CCAT1_JAX_4	Exon5	128231500	128231806	307	307
CCAT1_JAX_4	Exon6	128231809	128232653	845	845
CCAT1_JAX_5	Transcript	128172634	128231094	58461	1824
CCAT1_JAX_5	Exon1	128172634	128174329	1696	1696
CCAT1_JAX_5	Exon2	128176684	128176771	88	88
CCAT1_JAX_5	Exon3	128231055	128231094	40	40
CCAT1_JAX_6	Transcript	128197810	128240377	42568	11053
CCAT1_JAX_6	Exon1	128197810	128198015	206	206
CCAT1_JAX_6	Exon2	128200030	128200129	100	100
CCAT1_JAX_6	Exon3	128200290	128209809	9520	9520
CCAT1_JAX_6	Exon4	128215408	128215465	58	58
CCAT1_JAX_6	Exon5	128231055	128231098	44	44
CCAT1_JAX_6	Exon6	128231100	128231211	112	112
CCAT1_JAX_6	Exon7	128235783	128235911	129	129
CCAT1_JAX_6	Exon8	128236780	128236929	150	150
CCAT1_JAX_6	Exon9	128239644	128240377	734	734

CCAT1_JAX_7	Transcript	128186443	128240377	53935	1216
CCAT1_JAX_7	Exon1	128186443	128186609	167	167
CCAT1_JAX_7	Exon2	128231055	128231098	44	44
CCAT1_JAX_7	Exon3	128231100	128231212	113	113
CCAT1_JAX_7	Exon4	128235784	128235911	128	128
CCAT1_JAX_7	Exon5	128236780	128236929	150	150
CCAT1_JAX_7	Exon6	128239764	128240377	614	614
CCAT1_JAX_8	Transcript	128218833	128240377	21545	4103
CCAT1_JAX_8	Exon1	128218833	128218920	88	88
CCAT1_JAX_8	Exon2	128218923	128221962	3040	3040
CCAT1_JAX_8	Exon3	128231055	128231098	44	44
CCAT1_JAX_8	Exon4	128231100	128231211	112	112
CCAT1_JAX_8	Exon5	128235783	128235911	129	129
CCAT1_JAX_8	Exon6	128236645	128236720	76	76
CCAT1_JAX_8	Exon7	128239764	128240377	614	614

These CCAT1 transcripts are different from the CCAT1 transcript described below in NCBI Reference Sequence: XR_133500.3:

```

1 TCATCATTAC CAGCTGCCGT GTTAAGCATT GCGAAAACGC TCACGATTCA CAGAAAAATC
61 CAIGCTGTTT TTTGAAGGCA TTCAAGCCTT AATAGCTAGC TGGATGAATG TTTAACTTCT
121 AGGCCAGGCA CTACTCTGTC CCAACAATAA GCCCTGTACA TTGGGAAAGG TGCCGAGACA
181 TGAACCTTGG TCTTCTCTGC AATCCATCTG GAGCATTAC TGACAACATC GACTTTGAAG
241 TTGCACTGAC CTGGCCAGCC CTGCCACTTA CCAGGTTGGC TCTGIATGGC TAAGCGTTTT
301 CTCCTAAAAT CCCTTGAAAA CTGTGAGAAG ACCATAAGAA GATCATATCT TTAATTCTAT
361 TTCACAAGTC ACACAATATT CCAATCAAAT ACAGATGGTT GAGAAAAGTC ATCCATCTTC
421 CCTCCCCACC CTCCCACAGC CCTCAACCA CTGCCCTGAA ACTTATATGC TGTATCCGC
481 AGCTCCATCT GGAGCATCAC AGCTACTGTC AACCTTGACG CTCTTCTGA AAAAACACCG
541 GATGGACATC AGAACTATTT CTTAAGGAT GTTACTGAGC CACACAGGAA AACTTGCCTT
601 ATGATTTTGA ATGCACGGAT CTGATTTGAC TAAACATGAT AACTAGAGGA TCACCCAATC
661 TACTCCCATT TTCAACTCTA AATCATCAGA GTGTCTCAAA TCCAAAGCAC ACACAGACCA
721 GCCTGGCCAA CGCGGTGAAA CTCCACCCCT ACTAAAAGTA TAAAAATTAT CCAGGTGTGG
781 TGGCGGGGCG CTGTAATCCA AGCTACTTGG GAGTCTGAGG CAGGAGAATC CCTTGAACCT
841 GGGAGATGGA GGTTCAGTG AGCAGAGATC ACACCACCGC ACTTAGCCT GGGCCACAAA
901 TCAACAACAA CAACAACAAC AAAAAACAAA GCGCACACAG AGACTGAGGT CCTCTTTGGC
961 ATTGAGAAGA TGGCTATGCA AGTCCCAACT AGCAAGTGCA AACTTCCCAG CTTCACTTCT
1021 GCCAGTGTCC CTTACCCCT TCTCAACCC ACTGGGAGGC AGGAGGGTGC TTGACAATAA
1081 CAGCCTTGGC ATCACTCTGC CAGGGTGTA TAGGAAGTGT TACAATTCTG AGATTCTGTG
1141 TAAGCACTGG CCTTCTGCC TAGAATGCCT TCTCCTCTCT TTTTAACTG CATGCTCCTA
1201 TTTATCTTTC AAAGCCCGGA AAAAATAACA CTGCACACGG GAAATGCTCC CTTCTACTG
1261 CAGTCATTTA GATGACTCTA TGCCATTCCA TTCATTTCTC TTTCTACCA CAGAAGTGCT
1321 TTGAGATTTT GGAGTCAGAC TGCTTGAAC TGAATCCTGG CCTCTCATC AGAGACTTGA
1381 CTTATTTTAG GCAAGTTATA TAACCAATTT TACCTCAGT CTTACCCAT AAAATGGGTC
1441 TAATGAGAGT ACCTACCACA CAGAATTTTG ATGAAAACG AATGAGATGA AGGCCTTTAA
1501 GGCAGTGGTC CCCAACCCTG GGGACACAGA CAGGTACCAT TTTGTGGCCT GTTAGGAAC
1561 GGGCCACACA GCAGGAGGTG AGCAGTGGGT GAGTGAGATC AGCGTTATTT ACAGCTGCTC
1621 CCCATTGCTC ACCTTACTGC CTGAGCTCCA CCTCCTGTCA GATCAGCAGT GGCATTAAAT
1681 TCTCATAGCA GCACAAACCC TGTCATGAAC TGCACATGCG AGGGATCTAG GTTGTGCGCT
1741 CCTTATGAGA ATCTAATGCC TAATGACCTG TCACCGTCTC CCATCACCCC TAGATGGGAG
1801 TGTCTAGTTG CAGGAAACAA GCTCAGGGCT TCCACTGATT CTACATTATG GTGAGTTGTA
1861 TAATTATTTT ATTATATAAT ACAATGTAAT AATAATAGAA ACACAGTGCA CAACAAATGT
1921 AATGTGCTTG AATCATCCCC AAACCATCCC AGTCCACGGT CTCCACATT TTGTCTTTTC
1981 ACAAATTTGT CTTCCACAAA ACTGGTCCCT GGTGCCAAA AGGCTTGGGA CCACTGCTTT

```

```

2041 AAAGCCTTTG CATAGTGCTT AGAATTGAGG GGGAAAAAAA AAACAAAAAC AATGTAGCTA
2101 GTTGCTACAA TCACTATATT GGTGAGTTTC AAAAGGAAAA GAATTCTGTC CCATTTATGC
2161 TTGAGCCTTG AGTTGCTAAC CAAGCCTGAC ACAAATTAC TGTTGAAGGG ATGTGTGAGT
2221 CCTAATTGAA ATGAGGCCTC TTAAGGGAAT TGTGGACCAA ACCCCAAGCA GGCAGAAAGC
2281 CGTATCTTAA TTATTGCAAG TATTCAGGC AAGGTGTGGA TGGCCATTG AATTCAAGCA
2341 GACTAGGACC TGGGATGAGA AAGAAGGTGT GTACGTGACT TGATCTTTGA ACTTTAGCTC
2401 ACCATCTGGA AGAAGGCTGA GTATTCTCTG CACTCACATA GTAGCTAATG CCTACTCCCC
2461 AGCCACCCAC AATTCTTTCT GTAGGAAGGC TCGCTAGAAT ACTTTGTGAT ATTGGATATT
2521 AGTTCCATAT TCTACTGTGT ATCTTAGTTC AACCAAATTG TAATCATCTG ATATTTATTT
2581 CTTTAAATAT AAATATAAGT ATATTAAGTC TT (SEQ ID NO: 9)

```

Thus in one aspect, the invention provides cDNA sequences of the CCAT1 ncRNA transcripts, wherein the cDNA sequences are represented by a sequence selected from the group consisting of SEQ ID NOs: 1-8.

In a related aspect, the invention provides an antagonist sequence of a CCAT1 ncRNA, wherein the antagonist sequence antagonizes a function of the CCAT1 ncRNA.

In certain embodiments, the antagonizing sequence does not antagonize a function of the CCAT1 ncRNA corresponding to SEQ ID NO: 9.

In certain embodiments, the antagonist sequence is an antisense sequence to any one of the “-” strand cDNA sequences shown in SEQ ID NOs: 1-8.

In certain embodiments, the antisense sequence hybridizes to any one of the “-” strand cDNA sequences shown in SEQ ID NOs: 1-8 (but not SEQ ID NO: 9), under physiological conditions (*e.g.*, in the nucleus of a cell), or under a high stringency hybridization condition, such as one described in *Molecular Cloning: A Laboratory Manual* by Sambrook and Russell, Third Edition, 2001, published by Cold Spring Harbor Laboratory Press (incorporated herein by reference). One such high stringency hybridization condition may include 6 × sodium chloride/sodium citrate (SSC) at approximately 45°C, followed by one or more washes in 0.2×SSC and 0.1% SDS at 50°C, at 55°C, or at about 60°C, or about 65°C or more.

In certain embodiments, the antisense sequence is at least about 40%, 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 99% or more identical to any one of the “-” strand cDNA sequences shown in SEQ ID NOs: 1-8, at least in a region to which the antisense sequence hybridizes with the cDNA sequence. In certain embodiments, the antisense sequence is no more than about 50%, 40%, 30%, 20% identical to SEQ ID NO: 9.

In certain embodiments, the antisense sequence is about 10, 12, 14, 16, 20, 22, 24, 26, 28, 30 or more nucleotides in length.

In certain embodiments, the antagonist sequence is an siRNA or miRNA sequence that targets the destruction of any one or more of the CCAT1 ncRNA isoforms represented by the “-” strand cDNA sequences shown in SEQ ID NOs: 1-8 (but not SEQ ID NO: 9).

In certain embodiments, the antagonist sequence is a vector that encodes the siRNA / miRNA, or a dsRNA substrate for RNase III (such as Dicer) that can be processed to the siRNA or miRNA.

In certain embodiments, the siRNA or miRNA comprises a guide sequence of about 20-25 nucleotides that targets the destruction of the CCAT1 ncRNA isoforms.

In a related aspect, the invention provides a method of diagnosing cancer or precancerous lesions, comprising measuring the level of expression of any one of SEQ ID NOs: 1-8 or a fragment thereof in a biological sample, wherein expression of any one of SEQ ID NOs: 1-8 or a fragment thereof in the biological sample is indicative of cancer or a precancerous lesion. In certain embodiments, the fragment is not a fragment of SEQ ID NO: 9.

In certain embodiments, the method further comprises comparing the expression level measured in the biological sample with a standard, wherein a higher level of expression of any one of SEQ ID NOs: 1-8 or a fragment thereof in the biological sample is indicative of cancer or a precancerous lesion. In certain embodiments, the fragment is not a fragment of SEQ ID NO: 9.

In certain embodiments, the method comprising: (a) isolating nucleic acids from a biological sample obtained from a subject; (b) hybridizing a probe capable of recognizing to any one of SEQ ID NOs: 1-8 with the nucleic acids, under conditions allowing the formation of hybridization complexes; and (c) comparing hybridization complex formation with a standard; wherein a higher level of hybridization complexes in the biological sample is indicative of cancer or a precancerous lesion. In certain embodiments, the probe does not hybridize to SEQ ID NO:9.

In certain embodiments, the method comprising: (a) isolating nucleic acids from a biological sample obtained from a subject; (b) amplifying any one of SEQ ID NOs: 1-8 or any fragment thereof in the isolated nucleic acids; (c) visualizing the amplified CCAT1 product; and (d) comparing the amount of the CCAT1 amplification product with a standard; wherein the presence of a higher level of a CCAT-1 amplification product is indicative of cancer or a precancerous lesion. In certain embodiments, the fragment is not a fragment of SEQ ID NO: 9.

In certain embodiments, the amplification is performed by PCR (such as real-time quantitative PCR) using a probe specific for one or more of SEQ ID NOs: 1-8.

In certain embodiments, the standard is determined by measuring the level of expression of CCAT-1 in a subject not afflicted with cancer. In a related embodiment, the standard is determined by measuring the level of expression of CCAT-1 in a non-cancerous tissue of the same subject.

In certain embodiments, the cancer is selected from the group consisting of: colon cancer (*e.g.*, adenocarcinoma of the colon), rectal cancer, cervical cancer, lung cancer, gastric carcinoma, liver cancer and, metastases thereof.

In certain embodiments, the precancerous lesion is an adenomatous polyp.

In certain embodiments, the biological sample is selected from the group consisting of tissue, blood, saliva, urine, stool, and bone marrow samples.

A related aspect of the invention provides an oligonucleotide comprising at least 8 contiguous nucleotides of any one of SEQ ID NOs: 1-8 or a complement thereof, useful as a probe or a primer. In certain embodiments, the oligonucleotide does not hybridize to SEQ ID NO: 9.

A related aspect of the invention provides a method for detecting the expression of CCAT-1 in a biological sample, the method comprising: (a) isolating nucleic acids from the biological sample; (b) hybridizing the CCAT1 oligonucleotide probe of the invention to the nucleic acids under conditions allowing the formation of hybridization complexes; and (c) comparing hybridization complex formation with a standard, wherein a higher level of hybridization complexes in the biological sample indicates expression of CCAT-1 in the sample.

Another related aspect of the invention provides a vector comprising a cDNA or a fragment thereof, wherein the cDNA is selected from the group consisting of SEQ ID NOs: 1-8. In certain embodiments, the cDNA fragment does not hybridize to SEQ ID NO: 9.

Another related aspect of the invention provides a host cell comprising the subject vector.

Another related aspect of the invention provides a method of imaging cancer or precancerous lesions, comprising: (a) administering to a subject a CCAT1 probe of the invention; wherein the probe is conjugated to an indicator molecule; and (b) detecting the indicator molecule (*e.g.*, a radio-isotope, a fluorescent dye, a visible dye or a nano-particle)

conjugated to the probe by an imaging device.

A further related aspect of the invention provides a method to antagonize the function of a CCAT1 ncRNA transcript represented by any one or more of SEQ ID NOs: 1-8, comprising contacting the CCAT1 ncRNA with a subject antagonist sequence of CCAT1 (*e.g.*, antisense, miRNA or siRNA).

In certain embodiments, the method is carried out *in vitro*, and the CCAT1 ncRNA transcript is present in cells from a tissue culture sample.

In certain embodiments, the method is carried out *in vivo*, comprising administering to a subject in need thereof the subject antagonist sequence of CCAT1 (*e.g.*, antisense, miRNA or siRNA).

Yet another related aspect of the invention provides a pharmaceutical composition comprising a subject antagonist sequence of CCAT1 (*e.g.*, antisense, miRNA or siRNA), and a pharmaceutically acceptable excipient and/or carrier.

It should be understood that any embodiments described in the application, including embodiments only described under one aspect of the invention, can be combined with other embodiments of other aspects of the invention.

A person of ordinary skill in the art will appreciate that techniques not specifically taught herein may be found in standard molecular biology reference books, such as *Molecular Cloning: A Laboratory Manual* by Sambrook and Russell, Third Edition, 2001, published by Cold Spring Harbor Laboratory Press; *Oligonucleotide Synthesis* (M. J. Gait, ed., 1984); *Nucleic Acid Hybridization* (B. D. Hames and S. J. Higgins. eds., 1984); *PCR Technology - principles and applications for DNA amplification*, 1989, (ed. H. A. Erlich) Stockton Press, New York; *PCR Protocols: A Guide to Methods and Applications*, 1990, (ed. M. A. Innis *et al.*) Academic Press, San Diego; and *PCR Strategies*, 1995, (ed. M. A. Innis *et al.*) Academic Press, San Diego; all of which are incorporated herein by reference.

EXAMPLES

The invention generally described above will be more readily understood through reference to the following illustrative examples that are for illustration only, and are not intended to be limiting in any respect.

Example 1 General RICH-PET Methodology

Using RNA-DNA ligation followed by paired-end-tag sequencing (RICH-PET), Applicants have developed an exemplary method described below to study ncRNA (non-coding RNA) and chromatin interactions in an unbiased and genome-wide manner.

A principal concept behind the method is based on the realization that most of the ncRNA regulatory functions, particularly those adopted by long ncRNAs (lncRNAs), likely have direct or indirect contacts in specific chromatin loci through any combinations of RNA-protein, RNA-DNA, and/or RNA-RNA interactions. Therefore, a comprehensive collection of ncRNA contact addresses of chromatin locations in the entire genome would provide a large structural framework and the detail contents of genomic elements in order to understand the global impact as well as specific functions mediated by individual and/or collective ncRNAs.

Through crosslinking, RNA-chromatin interactions can be captured. After fragmentation of chromatin fibers by sonication, ncRNA and DNA fragments tethered together via protein bindings in each chromatin complex are then subjective for RNA-DNA ligation using the subject RNA and DNA linkers, in order to establish an artificial connectivity relationship of the RNA molecules and the DNA fragments for high throughput analysis with specificity.

The RNA linker of the invention may comprise a random oligonucleotide sequence, *e.g.*, random hexonucleotides, for annealing to the 3'-end of any tethered RNA molecules, and as the primer for reverse transcription to convert the RNA templates into first-strand cDNA molecules. Meanwhile, the DNA linker of the invention is ligated to the blunt-ended chromatin DNA fragments. The RNA linker and the DNA linker each has a sticky end complementary to each other but not to itself. Hence, once the linkers are attached accordingly to their intended targets, the RNA and DNA fragments can be covalently connected through ligation. The hybrid ligation products are then subjective for paired end tag (PET) library construction and subsequent high throughput sequencing analysis. A schematic drawing for this method is depicted in Fig 1A.

Alternatively, a modified RNA linker may be used to carry out the RNA-DNA ligation step. A schematic drawing for this method is depicted in Fig 1B.

Additionally, a direct RNA linker may be used to carry out the RNA-DNA ligation step by taking advantage of certain enzymes (such as the truncated RNL2) that can directly link RNA 3'-end to 5' adenylated ssDNA or 5' adenylated overhang, a schematic drawing for the latter method is depicted in Fig 1C.

To further distinguish the tag sequences from their original nature as RNA or DNA, specific nucleotide bar codes may be incorporated into RNA and/or DNA linker sequence designs, which then allow accurate calling of the paired RNA-tag and DNA-tag in a RICH-PET library dataset. The processed RNA-tag and DNA-tag sequences are then mapped to reference genome (*e.g.*, a reference human genome for human originated sequences) to identify ncRNAs and their chromatin target loci (data not shown).

Certain experimental details are provided below for illustration purpose.

I. Cell Culture and Crosslinking

HeLa S3 cells were grown in Ham's F-12 Nutrient Mix (Life Technologies, cat.11765-054) supplemented with 5% Fetal Bovine Serum (FBS) (Life Technologies, cat. 10082147). For each batch of crosslinked cells, EGS (spacer Arm: 16.1A; Thermo Scientific, cat.21565) and formaldehyde (spacer Arm: 2.0A; Merck – Calbiochem, cat. 344198-250ML) were used to treat the cells for dual-crosslinking of protein-DNA, protein-RNA and protein-protein, which could provide better connectivity than using only formaldehyde.

Around 1×10^8 cells in 245 mm square plate (Corning, cat.431110) were crosslinked with 45 ml of 1.5 mM EGS in pre-warmed DPBS (Life Technologies, cat.14190250), first shaking at 75 rpm for 40 min, then adding 1% formaldehyde (Merck – Calbiochem, cat.344198-250ML), and keep shaking for 20 min, followed by quenching with 0.125 M glycine (Promega, cat. H5071) for 10 min, then washing with ice-cold DPBS twice. Then 3 to 5 ml of ice-cold DPBS containing proteinase inhibitor (Roche, cat.11873580001) and RNase inhibitor (such as the SUPERase• In™ RNase Inhibitor, Life Technologies, cat. AM2696) were added, and then cells were scraping and transferred to 15 ml-Falcon tube (Life Technologies, cat.AM1250). This process was repeated as necessary to ensure that all cells were collected. Cells were spun down at 2000 rpm for 5min at 4°C, then the cell pellet were stored at -80°C until use.

II. Cell Lysis and Chromatin Biotinylation

Cell lysis was performed as described previously (Goh *et al.*, *J. Vis. Exp.*, (62), e3770, doi:10.3791/3770, 2012; Fullwood *et al.*, *Nature*, 462:58-64, 2009, both incorporate herein

by reference). Briefly, the nuclei pellet was washed twice with an ice-cold wash buffer (50 mM Tris-HCl pH= 8.0, 150 mM NaCl, 1 mM EDTA, 1% TritonX-100, 0.1% SDS), and suspended in 1 mL of the same buffer. Chromatin was sheared to fragments with average size of about 500 bp by, for example, sonication. SDS was then added to the final concentration of about 0.5% to the shearing chromatin, and the mixture was then incubated at 37°C for 15 min, before mixing with EZlink Iodoacetyl-PEG2-Biotin (IPB) (Thermo Scientific, cat.21334) and rotating at room temperature for 60 min as described previously (Kalhor et al., *Nat. Biotechnol.*, 30:90–98, 2012, incorporate herein by reference). The streptavidin beads-bound chromatin was then subjected to RICH-PET library construction.

III. RICH-PET Library Construction

The DNA fragments present in streptavidin beads-bound chromatin were end-repaired using T4 polymerase (Promega, R0191), followed by first-strand cDNA synthesis using Superscript III First Strand Synthesis System (Life Technologies, cat.18080051).

Briefly, 1 µg of biotinylated RNA linker a (tube 1) and RNA linker b (tube 2) containing a flanking MmeI site (IDT), were added to two tubes containing annealing mixture (5 µl 10 mM dNTPs, 40 µl DEPC-treated water), respectively, and incubated at 65°C for 5 min, then placed on ice for at least about 1 min, then mixed with cDNA synthesis mixture (10 µl 10× RT (reverse transcription) buffer, 20 µl 25 mM MgCl₂, 10 µl 0.1 M DTT, 5 µl RNaseOUT, 5 µl SuperScript III RT) for incubation for 10 min at 25°C, followed by 30 min at 50°C.

Overnight ligation was performed using 1 µg of DNA linker A (tube 1) and DNA linker B (tube 2), respectively, in ligation mixture (140 µl 5× T4 DNA ligase buffer with PEG, 3.5 µl RNase inhibitor, 546.5 µl nuclease free water) using 5 µl of T4 DNA ligase at 16°C. The linker-added DNA fragments were then phosphorylated with 14 µl of T4 polynucleotide kinase (NEB) in PNK master mix buffers (70 µl 10× T4 DNA ligase buffer, 3.5 µl RNase inhibitor, 612.5 µl Nuclease free water), followed by the two tubes proximity ligation with 34 µl of T4 DNA ligase in reaction buffer (1000 µl 10× T4 DNA ligase buffer, 50 µl RNase, 8916 µl Nuclease free water) overnight at 16°C.

Chromatin DNA fragments with linkers were subjected to second-strand cDNA synthesis with Superscript Double-stranded cDNA Synthesis Kit (Life Technologies, cat.1197-020). Specifically, chromatin fragments were mixed with second-strand cDNA mixture (111 µl DEPC-treated water, 30 µl 5× Second-strand reaction buffer, 3 µl 10 mM

dNTP mix, 1 μ l *E. coli* DNA ligase, 4 μ l *E. coli* DNA Polymerase I, 1 μ l *E. coli* RNase H), and were incubated at 16°C for 2 hours. Following the reaction, 2 μ L of T4 DNA polymerase was added for continued incubation at 16°C for 5 min.

The crosslinks in DNA/RNA/protein complexes were then reversed by incubation at 65°C overnight with 0.3% SDS (Ambion) and proteinase K (Ambion). The cDNA-DNA fragments were purified by phenol/chloroform isopropanol precipitation. The purified cDNA-DNA was then digested by 1 μ l of *MmeI* (NEB) in suitable buffer (5 μ l 10 $\times$ NEBuffer 4, 5 μ l Half linker non-Biotinylated to quench excess *MmeI*, 5 μ l 10 $\times$ SAM) for at least 2 hrs at 37°C to release the cDNA tag-RNA linker-DNA linker -DNA tag structure (paired end tag, PET).

The biotinylated PETs were then immobilized on streptavidin-conjugated magnetic Dynabeads (Life Technologies, cat.11206D-10ML) in 50 μ l of 2 $\times$ B&W buffer (10 mM Tris-HCl pH7.5, 1 mM EDTA, 1 M NaCl), rocked at room temperature for 45 min. The ends of each PET structure were then ligated to an adaptor by 1 μ l of T4 DNA ligase (Thermo Scientific, cat. EL0013) in Adaptor ligation buffer (4 μ l Adaptor A, 4 μ l Adaptor B, 5 μ l 10 $\times$ T4 DNA ligase buffer, 36 μ l Nuclease free water) at 16°C overnight with mixing. The beads were then washed three times with 1 $\times$ B&W buffer (5 mM Tris-HCl pH7.5, 0.5 mM EDTA, 1 M NaCl).

Nick translation was performed with 4 μ l of *E. coli* DNA polymerase I in a reaction mixture (38.5 μ l Nuclease free water, 10 $\times$ NEBuffer 2, 2.5 μ l 10 mM dNTPs), which was incubated at room temperature for 2 hours with rotation on an Intelli-Mixer (F8, 30 rpm, U=50, u=60; ELMI Ltd., Riga, Latvia). This was followed by 16 rounds of PCR to amplify the PETs. RICH-PET libraries were sequenced on an Illumina HiSeq2000 (2 $\times$ 36 bp reads).

All steps were performed in buffer with protease inhibitor and RNase-inhibitors to prevent or minimize protein and RNA degradation.

The various polynucleotides or primers used herein are listed below:

Polynucleotides	Name	Sequences
DNA linker A2	Rb-top-6	5'-Phos-GTTGGACTTGTACGATAGCTCTC-3'
	Rb-bot-6	5'-OH-GCTA/iBIOdT/CGTACAAGTCCAAC NNNNNV-3'
DNA linker B2	DB-top-6	5'-OH-GCGATATCACTGTTCCAAC-3'

Polynucleotides	Name	Sequences
	DB-bot-6	5'-OH-GTTGGAACAGTGATATCGCGAGA-3'
linker without biotin for sequencing access MmeI	top	5'-GGCCGCGATATCGGATCCAAC -3'
	bottom	5'- GTTGATCCGATATCGC -3'
Adaptor A	top	5'-CCATCTCATCCCTGCGTGTCCTCATC TGTTCCCTCCCTGTCTCAGNN-3'
	bottom	5'-CTGAGACAGGGAGGGAACAGAT GGGACACGCAGGGATGAGATGG-3'
Adaptor B	top	5'-CTGAGACACGCAACAGGGGAT AGGCAAGGCACACAGGGGATAGG-3'
	bottom	5'-CCTATCCCCTGTGTGCCTTGC CTATCCCCTGTTGCGTGTCCTCAGNN-3'
PCR primer 1		5'-AATGATACGGCGACCACCGAGAT CTACACCCTATCCCCTGTGTGCCTTG-3'
PCR primer 2		5'-CAAGCAGAAGACGGCATAACGAGA TCGGTCCATCTCATCCCTGCGTGTC-3'
sequencing primer 1		5'-GTGCCTTGCCTATCCCCTGTT GCGTGTCCTCAG-3'
sequencing primer 2		5'-TGCGTGTCCCATCTGTTCCCT CCCTGTCTCAG-3'

Example 2 RICH-PET Library Statistics

Three RICH-PET library datasets were generated using technical and biological replicates from HeLa S3 cells.

HeLa S3 RICH-PET data mapping results

Libraries	Replicates	Reads	Unique PET	Cluster ($\geq$ PET2)
CHH2430	1 (Tec)	52,254,130	2,367,898	5,371
JCHH2430	2 (Tec)	211,837,204	2,920,369	9,089
JCHH2431	3 (Bio)	83,143,999	2,049,942	3,128

RICH-PET data is classified either as singleton PET (*i.e.*, no overlap on both RNA-tag and DNA-tag with other PET sequences) or as PET cluster (*i.e.*, both of the paired RNA-tag and DNA-tag sequences overlap with other PETs) with 2 and more PET sequences. The PET clusters are considered to be more reliable, or as high confidence data reflecting recurrent detection of more reliable events of ncRNA-chromatin interactions, whereas the singleton PETs may represent weak linking signals, but are indistinguishable from random background noises. Using the clustering criterion, approximately 700 putative RNA loci that are connected to about 5000 chromatin loci were identified (Fig 2A).

As a quick verification, RNA-seq signals for these RNA and DNA loci were checked, and it was found that the RNA loci indeed had significantly higher RNA counts than the DNA loci, suggesting that the RICH-PET data are as expected (Fig 2B).

About one fifth (about 22%) of the obtained RNA-DNA connectivity data can be considered as *cis*-acting in nature (*i.e.*, <100kb from the RNA to DNA mapping sites), while the majority of the RNA-DNA connectivity data is *trans*-acting (Figs. 2C).

One concern was that the chromatin RNA-DNA ligation approach may capture mostly the nascent mRNA when transcription is still in process. Surprisingly, the data shows that most nascent mRNA transcripts appear to have their 3'-ends hidden within the center of the RNA polymerase complex, such that the method of the invention, which is partly based on using the supposedly free 3'-ends of ncRNA molecules, largely avoids the interference from nascent mRNA.

Specifically, mapping paired RICH-PET data reveals the distance between the paired RNA and DNA tags, thus suggesting possible mode of interacting action, *cis* or *trans*. The

mapping results showed that only a small set of the data was cis-acting, and the majority was trans-acting and inter-chromosomal, indicating that the likelihood of capturing nascent transcripts in the RICH-PET protocol is low.

Further annotation analysis of the RNA tag clusters (see below) showed that only 3% of the RNA tags mapped to mRNA exons, while the vast majority mapped to ncRNAs.

Another concern was the abundance of rRNAs in cells, which is a general issue for RNA related analysis because in some cells, rRNA could be as high as 80% of total RNAs.

One strategy to deal with rRNAs includes the avoidance approach, such as the polyA+ selection approach for mRNA and subtractive depletion of rRNA, used prior to the start of specific analysis. We assessed the abundance level of rRNA sequences in one of the RICH-PET libraries, and found that rRNA sequences constitute about 26% of the total RNA tags. In contrast, almost none (0.23%) of the DNA tags correspond to rRNA sequences. Thus a digital depletion approach may be used to remove all rRNA sequences before any further analysis to reduce data noise due to rRNA.

RICH-PET	Total	Non-rRNA	rRNA
RNA Tag	2308959	1699014 (73.58%)	609945 (26.42%)
DNA Tag	2308959	2303550 (99.77%)	5409 (0.23%)

Example 3 Reproducibility and Sensitivity of the RICH-PET Method

To assess the reproducibility of the RICH-PET data, two technical replicates (same cell preparations split into two aliquots for parallel library construction and sequencing analysis) and two biological replicates (different cell preparations collected at different times for use in library construction using nearly identical procedures with slight modifications) were performed. The resulting replicate results showed genuine reproducibility (Fig. 3). For example, the two well studied lncRNAs NEAT1 and MALAT1, known to be involved in cancers, were reproducibly detected in all three libraries (data not shown).

It is noteworthy that the two lncRNA genes were found to be spatially organized in an extensive chromatin interaction loop structure mediated by RNA polymerase II (RNAPII or RNA Pol2), indicating that their expressions are most likely co-regulated under a common transcription complex of machinery.

In RICH-PET data obtained herein, both MALAT1 and NEAT1 were highly expressed in HeLa S3 cells, and were abundantly detected in all three RICH-PET datasets. Specifically, NEAT1 was expressed relatively less compared to MALAT1 in the cells, thus the RICH-PET data counts to NEAT1 was less than that to MALAT1 (data not shown). As a control, HOTAIR is another known lncRNA expressed in low level in HeLa S3 cells, and it was not detected in the obtained RICH-PET data (data not shown).

Thus it appeared that the detection of ncRNA in RICH-PET data was well correlated with ncRNA expression levels.

Example 4 Validation of the RICH-PET Data

Based on the obtained RICH-PET mapping data, it is intriguing that even though these two ncRNAs are co-transcribed in the same transcription factory, their interaction properties are very different. Specifically, NEAT1 RNA is restrictively in *cis*, binding only to where it was transcribed; whereas MALAT1 is mostly out going in *trans*, interacting with many loci in the genome (Fig 4A).

To validate this observation, RNA-FISH experiments were conducted using NEAT1 and MALAT1 RNAs as fluorescent probes to examine HeLa nuclei (Fig 4B). As expected, the NEAT1 probe yielded only 1 or 2 spots per nucleus, whereas the MALAT1 probe spotted all over the nuclear space, consistent to what were observed in the RICH-PET data. Similar RNA-FISH result for NEAT1 and MALAT1 in A549 cells was also obtained. This validation suggests that RICH-PET data is qualitative and accurate in detecting and distinguishing authentic *cis* and *trans* interactions.

Example 5 Characterization of the RICH-PET Data

The RNA and DNA tag clusters were characterized based on Genecode V14 annotation of the human genome. Only 3% of the RNA tag clusters overlapped with protein-coding exons, and the vast majority of the RNA tag clusters were mapped to non-coding regions, many of which are previously known ncRNAs (172, 24%). The rest are potentially novel ncRNAs located in protein-coding intron regions, antisense, and inter-genetic regions (Fig 5A).

All putative ncRNAs identified in the RICH-PET data have RNA-Seq data support, indicating that they are actively transcribed in HeLa cells. In contrary, the DNA tag clusters

of the RICH-PET data mapped mostly to protein-coding genes, and a significant portion to gene promoters (Fig 5B).

A set of chromatin activity marks around the RNA and DNA tag clusters were subjected to further analysis. It is interesting to note that the center of RNA tag clusters are off the peaks of transcription activity defined by the signals of RNA Pol2 and DHS for open chromatin state, and such “off-center” property is strand specific (data not shown). This strand-specific “off-center” property is consistent with the RICH-PET methodology, as it is designed to capture the 3’-end of RNAs. Therefore, the RNA tag clusters are expected to be downstream of transcription start sites. In contrast, the chromatin activity signals are symmetrically peaked around the center of DNA tag clusters (data not shown), reflecting the random shearing of chromatin fibers by sonication.

Example 6 MALAT1 interacts with many genomic features and may function for both gene activation and gene repression

Using all RICH-PET data connected to MALAT1 (including singleton PETs), Applicants generated the chromosome-wide and genome-wide MALAT1-interaction profile, showing that MALAT1 has a potential to interact with a large territory in the genome (data not shown). Of more than 50 high-confidence interactions (PET cluster with tag counts ≥ 2) sites, about half are located in promoters and a quarter in intron regions of known genes (Fig 6A). RNA-Seq and RNA Pol2 ChIP-seq data from the same cells showed that genes with MALAT1 presence in their promoters have significantly higher transcriptional activities than those with MALAT1 interacting at their intron regions (Figure 6B; data not shown). It had been reported that MALAT1 was involved in modulating splicing functions through interacting with a number of splicing factors including SRSF2 (Tripathi *et al.*, 2011).

Applicants also found that MALAT1 RNA may be directly involved in modulating the expression of SRSF2 by interacting with its promoter (data not shown). These observations suggest that MALAT1 may have multiple functional roles in regulating gene activation and repression.

Example 7 XIST’s Function Beyond X-Chromosome

The most well characterized lncRNA is XIST, which is transcribed from one copy of the X chromosome and binds (cis-acting) to the same site in the other copy of the X

chromosome, and further extends to coat the entire chromosome for inactivation (not shown). The RICH-PET mapping data indeed showed that the DNA tags paired with the RNA tags of XIST were highly enriched in the X chromosome, while the background noise was scattered throughout the genome, indicating that XIST is specifically bound to the X chromosome as expected.

Interestingly, it also appeared that there was some level of XIST-binding enrichment in one non-X chromosome, and somehow depleted to another non-X chromosome. More data and further analysis are being obtained to further validate this observation.

Example 8 Complex interaction networks by ncRNAs

The RICH-PET data presented here provided a first glimpse into the complex systems of ncRNA interaction networks. In addition to the classic view that one ncRNA may have multiple targets in the genome (MALAT1), it had been found that many putative ncRNA loci have “in-and-out” RICH-PET data, in that a locus was found to be interacted by an ncRNA and from where an interactive ncRNA was also detected to interact with another locus.

In many sense, this ncRNA interaction network is similar to transcription factor (TF) binding networks, in which many TFs bind to each other's genes for transcriptional modulation. More data will help to further illustrate how ncRNAs function, and how the ncRNA interaction networks impact the genome system.

Example 9 lncRNA Encoded by CCAT1 Is a Transcription Co-Activator

The RICH-PET method was used to identify global ncRNA - genomic DNA interaction. Among the identified interactions, one ncRNA - the Colon Cancer Associated Transcript 1 - was of particular interest.

Colon Cancer Associated Transcript 1 (CCAT1) is a 2628 nucleotide-long, non-coding RNA recently discovered using Representational Difference Analysis (RDA), cDNA cloning, and rapid amplification of cDNA ends (RACE) (Nissan *et al.*, “Colon cancer associated transcript-1: A novel RNA expressed in malignant and pre-malignant human tissues,” *Int. J. Cancer*, 13:1598–1606, 2012). It is recently found to be over-expressed in colon cancer (CC), but not in normal tissues, thereby making it a potential disease-specific biomarker (Nissan *et al.*, *Int. J. Cancer*, 130(7):1598-606, 2012; Alaiyan *et al.*, *BMC Cancer*, 13:196, 2013).

Careful analysis based on the RICH-PET data revealed a new complex model of isoform transcripts in this locus (data not shown). In addition, CCAT1 is highly transcribed in the cervical cancer cell line HeLa cells.

RICH-PET data also revealed that the CCAT1 lncRNA transcripts targets many other loci in the genome (data not shown), including all the human chromosomes except for chromosomes 15, 16, 20, X and Y.

Among the CCAT1 chromatin targets having at least 2 CCAT1 tags, many show the strongest lncRNA-genomic DNA association in enhancers or promoters (data not shown). For example, for the 122 CCAT1 genomic target loci associated with at least 3 CCAT1 RNA tags, 88 target loci are in the enhancer region, including 6 of the enhancer loci with RNAPII interaction. Another 34 genomic target loci of CCAT1 are in promoters.

These CCAT1 target genes have an average expression level several folds higher than randomly selected collections of control genes, suggesting that CCAT1 lncRNA promotes target gene expression.

One of these CCAT1 target genes is c-myc, an oncogene overexpressed in a wide variety of human cancers, including about 80% of breast cancers, 70% of colon cancers, 90% of gynecological cancers, 50% of hepatocellular carcinomas, and a variety of hematological tumors (such as Burkitt's lymphoma) possessing abnormal myc expression. Additional data suggests that the CCAT1 lncRNA functions by binding to the CCAT1 locus itself as well as the myc locus, thus bringing the CCAT1 and myc loci to close physical proximity, and allowing the enhancers in the CCAT1 locus to stimulate myc transcription. In addition, the CCAT1 transcribed lncRNA may bind to protein factors and serve as transcription co-activators, thus directly enhancing transcription of myc, as well as other CCAT1 target genes such as FAM84B and SNX14.

Example 10 Additional Applications in Human B-lymphoblastoid Cells GM12878 and *Drosophila* S2 Cells

Using substantially the same RICH-PET methods described above, Applicants obtained additional data from the human B-lymphoblastoid cells GM12878 and the *Drosophila* S2 cells to further support the general applicability of the RICH-PET methods.

Specifically, the human GM12878 cells were used for RICH-PET analysis because the ncRNA gene XIST is highly expressed in this cell line, while the previous HeLa cells used

for RICH-PET analysis have low level of XIST expression, and HCT116 is derived from a male, thus having no XIST expression. Hence, GM12878 is a much better cell type for RICH-PET analysis when using XIST as a model to evaluate the performance of RICH-PET analysis to detect ncRNA interaction with chromatin.

As previously described, XIST specifically or preferentially binds to the X-chromosome. See FIG. 9A, which shows the counts of XIST measured by RNA-Seq data, in reads per kb per million reads (RPKM); and FIG. 9B, which shows the proportion of each chromosome covered by XIST binding. In GM12878 cells, most of the chromosomes were only covered by XIST in 10-20% of the total chromosome space, whereas the X-chromosome was covered by nearly 90% by XIST. This coverage represents almost 6-fold (5.9-fold) specificity of the XIST to its targeting chromosome over the other non-specific chromosomes. In contrast, in HeLa cells, the coverage represented about 3.4-fold specificity of the XIST to its targeting chromosome over the other non-specific chromosomes, and in HCT116 cells, there was no observed X-chromosome enrichment, as expected.

Similarly, in *Drosophila* S2 cells, the ncRNA gene *rox2* - an equivalent to XIST of human - showed similar enrichment of *rox2* binding to X-chromosome: 5-fold over the other chromosomes (data not shown). Specifically, *rox2* binding data in whole *Drosophila* genome was obtained. More than 80% of the *rox2*-linked DNA-tags bind to the X-chromosome, representing a 5-fold enrichment to the X-chromosome. There was a reasonably strong correlation value (0.6) observed between *rox2* mapping on X-chromosome by CHART-seq, and by the RICH-PET method, demonstrating the suitability of the RICH-PET method.

The majority of the RNA-tags of the RICH-PET data mapped to non-coding regions, while only about 26% are in coding regions, indicating the method has enrichment for ncRNA (data not shown). Comparison of the RNA-tags of RICH-PET data with the RNA-seq data from *Drosophila* S2 cells showed significant enrichment for known ncRNAs (data not shown).

In summary, data presented in the examples above demonstrates that the method of the invention (*e.g.*, RICH-PET method) works as designed. The vast majority of the RNA tags in the RICH-PET data were mapped to non-coding regions, and some of them mapped to known lncRNAs such as MALAT1 and NEAT1. This is a strong indication that this method performed as expected. More importantly, through the RNA-DNA connectivity mapping data, Applicants are able to identify potential ncRNA-chromatin interaction loci genome-

wide. Several lines of preliminary validations done so far have suggested that RICH-PET identified ncRNA interactions are *bona fide*.

Claims:

1. A kit comprising:

(1) an RNA linker comprising:

(i) a first polynucleotide, and,

(ii) a second polynucleotide,

wherein the first and the second polynucleotides form a first double stranded region flanked by a first ligation compatible end, and a 3'-overhang at the 3'-end of the first polynucleotide, wherein the 3'-overhang comprises a random-sequence primer; and,

(2) a DNA linker comprising:

(iii) a third polynucleotide, and,

(iv) a fourth polynucleotide,

wherein the third and the fourth polynucleotides form a second double stranded region flanked by a blunt end and a second ligation compatible end, wherein the first and the second ligation compatible ends ligate to each other, or are adaptable to ligate to each other.

2. The kit of claim 1, wherein the first ligation compatible end is a 3'-overhang at the 3'-end of the second polynucleotide, and the second ligation compatible end is a 3'-overhang at the 3'-end of the third polynucleotide, wherein both 3'-overhangs anneal to each other for ligation.

3. The kit of claim 1, wherein the first double stranded region comprises a first recognition site for a first restriction enzyme (RE) that cleaves 3' to the random-sequence primer, or, wherein the second double stranded region comprises a second recognition site for a second restriction enzyme (RE) that cleaves 5' to the third polynucleotide.

4. The kit of claim 1, wherein one or more of said first, second, third, and fourth polynucleotides are DNA, or, wherein one or more of said first, second, third, and fourth polynucleotides comprise a modified nucleotide, such as a biotinylated T (Thymidine).

5. The kit of claim 1, wherein said first polynucleotide comprises a plurality of polynucleotides, each differing only at the random-sequence primer region, or, wherein said first polynucleotide

comprises a homogeneous population of polynucleotides having identical random-sequence primer, or, wherein said random-sequence primer comprises 4, 5, 6, 7, 8, or more nucleotides.

6. The kit of claim 1, wherein the first double stranded region comprises a unique sequence that distinguishes the RNA linker from the DNA linker, or, wherein the second double stranded region comprises a unique sequence that distinguishes the RNA linker from the DNA linker, or, wherein the last nucleotide of the first recognition site is the last base-paired nucleotide 5' to the random sequence primer, or, wherein the last nucleotide of the second recognition site is a base-paired nucleotide at the blunt end.

7. The kit of claim 1, wherein the first and the second restriction enzymes are the same, or, wherein the first or the second restriction enzyme is independently selected from:

AarI, AceIII, AfoI, BaeI, Bbr7I, BbvI, BbvII, BccI, Bce83I, BceAI, BceFI, BcgI, BciVI, BfiI, BinI, BpII, BsaXI, BscAI, BseMII, BseRI, BsgI, BsmI, BsmAI, BsmFI, Bsp24I, BspCNI, BspMI, BsrI, BsrDI, BstF5I, BtgZI, BtsI, CjeI, CjePI, EciI, Eco31I, Eco57I, Eco57MI, EcoP15I, Esp3I, FaeI, FauI, FokI, GsuI, HaeIV, HgaI, Hin4I, HphI, HpyAV, Ksp632I, MboII, MlyI, MmeI, MnlI, PfiI, PpiI, PstI, RleAI, SapI, SfaNI, SspD5I, Sth132I, StsI, TaqII, TspDTI, TspGWI, TspRI or Tth111II, or,

wherein the cleavage site of the first or the second restriction enzyme is at least about 10, 12, 14, 16, 18, 20, or more nucleotides 3' to the last nucleotide of the recognition site, or, wherein the first and the fourth polynucleotides are dephosphorylated.

8. The kit of claim 1, further comprising a reagent that cross-links protein and polynucleotide, such as a reagent comprising formaldehyde, or, further comprising an affinity reagent (*e.g.*, an antibody, or a monoclonal antibody) that specifically or selectively bind a component of chromatin (*e.g.*, histone), or, further comprising an end-repairing mixture that converts DNA containing damaged or incompatible 5'- and/or 3'-protruding ends to 5'-phosphorylated, blunt-ended DNA, or, further comprising a DNA ligase (*e.g.*, T4 ligase), or, further comprising a reagent that reverses cross-linking of protein and polynucleotide (*e.g.*, Proteinase K), or, further comprising the first and/or the second restriction enzyme(s), or, further comprising a pair of concatenating adapters for PCR amplification of blunt-ended double stranded DNA,

or, further comprising a Taq DNA polymerase, or, further comprising a reverse transcriptase.

9. A paired-end tag (PET) polynucleotide comprising a central region comprising the first and second double stranded regions of claim 1, said central region being flanked by:

(1) at a site proximal to said first double stranded region, a sequence tag of a non-coding RNA (ncRNA); and

(2) at a site proximal to said second double stranded region, a sequence tag of a genomic DNA.

10. The PET polynucleotide of claim 9, wherein the sequence tag of the non-coding RNA (ncRNA) has a free end resulting from digestion by said first restriction enzyme, or, wherein the sequence tag of the non-coding RNA (ncRNA) uniquely identifies a genomic region from which the ncRNA is transcribed, or, wherein said sequence tag of the non-coding RNA (ncRNA) is about 8-30 base pairs in length, or, wherein the sequence tag of the genomic DNA has a free end resulting from digestion by said second restriction enzyme, or, wherein the sequence tag of the genomic DNA uniquely identifies a genomic region at which the genomic DNA is located, or, wherein said sequence tag of the genomic DNA is about 8-30 base pairs in length.

11. A paired-end tag (PET) library comprising two or more members of the PET polynucleotide of claim 9, wherein each member of the PET library comprises the same said central region, and different said sequence tag of the noncoding RNA (ncRNA) of claim 9 or different said sequence tag of the genomic DNA of claim 9 or both.

12. A vector comprising a PET polynucleotide of claim 9, optionally further comprising a plurality of concatenated PET polynucleotide of claim 9.

13. A concatemer of two or more PET polynucleotides of claim 9.

14. A method of identifying functional interaction loci within a genome for non-coding RNAs (ncRNAs) of the genome, the method comprising:

(1) providing chromatin fragments comprising cross-linked genomic DNA fragments and cross-linked ncRNAs;

(2) using the RNA linker and the DNA linker of claim 1, ligating an end of a cross-linked genomic DNA fragment to an end of a cDNA of a cross-linked ncRNA, under a condition for proximity ligation, wherein said end of the cross-linked genomic DNA fragment is ligated to the DNA linker, and said end of the cDNA of the cross-linked ncRNA comprises the RNA linker;

(3) isolating a PET polynucleotide of claim 9 for sequencing analysis; and,

(4) mapping the sequence tag of the genomic DNA and the sequence tag of the ncRNA within each said PET polynucleotide to a reference genome, thereby identifying functional interaction loci within the reference genome for said non-coding RNAs (ncRNAs) of the reference genome.

15. The method of claim 14, wherein the ncRNAs and the genomic DNA are cross-linked in live cells through formaldehyde-mediated cross-linking, or, wherein chromatin fragments are generated by sonication, or, wherein the cDNA of the cross-linked ncRNA comprises a first strand cDNA reverse transcribed from the random sequence primer of the RNA linker, and the ncRNA template, or, wherein 2nd strand cDNA synthesis is carried out after proximity ligation but before step (3), or, further comprising repairing the ends of the cross-linked genomic DNA fragments to 5'-phosphorylated, blunt ended DNA prior to step (2), or, wherein the third polynucleotide of the DNA linker is dephosphorylated and the DNA linker does not self-ligate, or, further comprising identifying clusters of two or more PET polynucleotides having overlapping sequence tags of the genomic DNA and overlapping sequence tags of the ncRNA, optionally, further comprising excluding PET polynucleotides comprising sequence tags of rRNA, or, further comprising isolating or enriching a subset of chromatin fragments prior to step (2), optionally, the subset of chromatin fragments is isolated or enriched by immunoprecipitation using an antibody specific for a protein component of the subset of chromatin fragments and further optionally, wherein the protein component is a histone, a transcription factor, a polycomb-group (PcG) family protein; a recombination involved factor; a chromatin insulator or chromatin waver; a methyl-CpG-binding protein; or an RNA binding protein.