

PHARMACEUTICAL COMPOSITIONS FOR COMBINATION THERAPY

The present invention relates to a combination therapy of inflammatory, metabolic, fibrotic and cholestatic diseases.

5

[2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl]ethanoate, (or nitazoxanide, or NTZ), first described in 1975 (Rossignol and Cavier 1975), was shown to be highly effective against anaerobic protozoa, helminths, and a wide spectrum of microbes including both anaerobic and aerobic bacteria (Rossignol and Maisonneuve 1984; Dubreuil, Houcke et al. 1996; 10 Megraud, Occhialini et al. 1998; Fox and Saravolatz 2005; Pankuch and Appelbaum 2006; Finegold, Molitoris et al. 2009). It was first studied in humans for the treatment of intestinal cestodes (Rossignol and Maisonneuve 1984) and it is now licensed in the United States (Alinia®, Romark laboratories) for the treatment of diarrhea caused by the protozoan parasites *Cryptosporidium parvum* and *Giardia intestinalis*. NTZ has also been widely 15 commercialized in Latin America and in India where it is indicated for treating a broad spectrum of intestinal parasitic infections (Hemphill, Mueller et al. 2006). The proposed mechanism of action by which NTZ exerts its antiparasitic activity is through the inhibition of pyruvate:ferredoxin oxidoreductase (PFOR) enzyme-dependent electron transfer reactions that are essential for anaerobic metabolism (Hoffman, Sisson et al. 2007). NTZ also 20 exhibited activity against *Mycobacterium tuberculosis*, which does not possess a homolog of PFOR, thus suggesting an alternative mechanism of action. Indeed, the authors showed that NTZ can also act as an uncoupler disrupting membrane potential and intra-organism pH homeostasis (de Carvalho, Darby et al. 2011).

The pharmacological effects of NTZ are not restricted to its antiparasitic or antibacterial 25 activities and in recent years, several studies revealed that NTZ can also confer antiviral activity (Di Santo and Ehrisman 2014; Rossignol 2014). NTZ interferes with the viral replication by diverse ways including a blockade in the maturation of hemagglutinin (influenza) or VP7 (rotavirus) proteins, or the activation of the protein PKR involved in the innate immune response (for a review, see (Rossignol 2014)). NTZ was also shown to have 30 broad anticancer properties by interfering with crucial metabolic and prodeath signaling pathways (Di Santo and Ehrisman 2014).

The present invention describes novel combinations of (i) NTZ or an analogue thereof, and (ii) an anti-NASH, anti-fibrotic or anti-cholestatic agent, and their use in therapy, in particular 35 in the treatment of inflammatory, metabolic, fibrotic and cholestatic diseases.


The inventors found that NTZ, a synthetic antiprotozoal agent or its derivatives or its metabolites in combination with an anti-NASH, anti-fibrotic or anti-cholestatic show therapeutic activities that are useful in therapy, in particular for the treatment of immune, inflammatory, metabolic, fibrotic or cholestatic diseases.

5

Therefore, the present invention relates to a combination product comprising:

- (i) a compound selected from NTZ and an analogue thereof ; and
- (ii) an anti-NASH, anti-fibrotic or anti-cholestatic agent.

- 10 In a particular embodiment, the compound of component (i) of the combination product is a compound of formula (I):




or a pharmaceutical acceptable salt thereof, in which :


- 15 R1 represents a hydrogen atom (H), a deuterium atom (D), a halogen atom, a (C6-C14)aryl group, a heterocyclic group, a (C3-C14)cycloalkyl group, a (C1-C6)alkyl group, a sulfonyl group, a sulfoxide group, a (C1-C6)alkylcarbonyl group, a (C1-C6)alkyloxy, a carboxylic group, a carboxylate group, a nitro group (NO₂), an amino group (NH₂), a (C1-C6)alkylamino group, an amido group, a (C1-C6)alkylamido group, a (C1-C6)dialkylamido group.
- 20 R2 represents a hydrogen atom, a deuterium atom, a NO₂ group, a (C6-C14)aryl group, a heterocyclic group, a halogen atom, a (C1-C6)alkyl group, a (C3-C14)cycloalkyl group, a (C2-C6)alkynyl group, a (C1-C6)alkyloxy group, a (C1-C6)alkylthio group, a (C1-C6)alkylcarbonyl group, a (C1-C6)alkylcarbonylamino group, a (C6-C14)arylcarbonylamino group, a carboxylic or carboxylate group, an amido group, a (C1-C6)alkylamido group, a (C1-C6)dialkylamido group, a NH₂ group, a (C1-C6)alkylamino group,
- 25 or R1 and R2, together with the carbon atoms to which they are attached, form a substituted or unsubstituted 5- to 8- membered cycloalkyl, heterocyclic or aryl group,
- R3, R4, R5, R6, and R7, identical or different, represent a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a (C1-C6)alkylcarbonyl group, an (C1-C6)alkyl group, an

(C1-C6)alkyloxy group, an (C1-C6)alkylthio group, an (C1-C6)alkylcarbonyloxy group, an (C6-C14)aryloxy group, a (C6-C14)aryl group, a heterocyclic group, a (C3-C14)cycloalkyl group, a NO₂ group, a sulfonylaminoalkyle group, an NH₂ group, an amino(C1-C6)alkyl group, an (C1-C6)alkylcarbonylamino group, a carboxylic group, a carboxylate group, or a
 5 R₉ group;

R₉ represents a O-R₈ group or an amino acid selected from the group consisting of alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, or a moiety of formula (A):



wherein R' represents an (C1-C6)alkyl group, an (C2-C6)alkenyl group, an (C2-C6)alkynyl group, a (C3-C14)cycloalkyl group, (C3-C14)cycloalkylalkyl group, a (C3-C14)cycloalkyl(C2-C6)alkenyl group, a (C3-C14)cycloalkenyl group, a (C3-C14)cycloalkenyl(C1-C6)alkyl group, a (C3-C14)cycloalkenyl(C2-C6)alkenyl group, a (C3-C14)cycloalkenyl(C2-C6)alkynyl group;
 15 R'' and R''', independently, represent hydrogen atom, an (C1-C6)alkyl group, or a nitrogen protecting group.




R₈ represents a hydrogen atom, a deuterium atom, a glucuronidyl group, or a group wherein, R_{8a}, R_{8b} and R_{8c}, identical or different, represent a hydrogen atom or a deuterium atom.

20


In a further particular embodiment, the compound of component (i) of the combination product is a compound of formula (I), or a pharmaceutical acceptable salt thereof, in which:
 R₁ represents a hydrogen atom (H), a deuterium atom (D), a halogen atom, a (C6-C14)aryl group, a heterocyclic group, a (C3-C14)cycloalkyl group, a (C1-C6)alkyl group, a sulfonyl group, a sulfoxyde group, a (C1-C6)alkylcarbonyl group, a (C1-C6)alkyloxy, a carboxylic group, a carboxylate group, a NO₂ group, a NH₂ group, a (C1-C6)alkylamino group, an amido group, a (C1-C6)alkylamido group, a (C1-C6)dialkylamido group.

R₂ represents a hydrogen atom, a deuterium atom, a NO₂ group, a (C6-C14)aryl group, a heterocyclic group, a halogen atom, a (C1-C6)alkyl group, a (C3-C14)cycloalkyl group, a
 30 (C2-C6)alkynyl group, a (C1-C6)alkyloxy group, a (C1-C6)alkylthio group, a (C1-

C6)alkylcarbonyl group, a (C1-C6)alkylcarbonylamino group, a (C6-C14)arylcarbonylamino group, a carboxylic or carboxylate group, an amido group, a (C1-C6)alkylamido group, a (C1-C6)dialkylamido group, a NH₂ group, a (C1-C6)alkylamino group,
 or R1 and R2, together with the carbon atoms to which they are attached, form a substituted
 5 or unsubstituted 5- to 8- membered cycloalkyl, heterocyclic or aryl group,
 R3 represents a hydrogen atom, a deuterium atom, a halogen atom, a O-R8 group, a (C1-C6)alkylcarbonyl group, an (C1-C6)alkyl group, an (C1-C6)alkyloxy group, an (C1-C6)alkylthio group, an (C1-C6)alkylcarbonyloxy group, an (C6-C14)aryloxy group, a (C6-C14)aryl group, a heterocyclic group, a (C3-C14)cycloalkyl group a NO₂, a
 10 sulfonylaminoalkyle group, an NH₂ group, an amino(C1-C6)alkyl group, an (C1-C6)alkylcarbonylamino group, a carboxylic group, a carboxylate group, an amino acid selected from the group consisting of alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, or a moiety of formula
 15 (A):




wherein R' represents an (C1-C6)alkyl group, an (C2-C6)alkenyl group, an (C2-C6)alkynyl group, a (C3-C14)cycloalkyl group, (C3-C14)cycloalkylalkyl group, a (C3-C14)cycloalkyl(C2-C6)alkenyl group, a (C3-C14)cycloalkenyl group, a (C3-C14)cycloalkenyl(C1-C6)alkyl group,
 20 a (C3-C14)cycloalkenyl(C2-C6)alkenyl group, a (C3-C14)cycloalkenyl(C2-C6)alkynyl group;
 R'' and R''', independently, represent hydrogen atom, an (C1-C6)alkyl group, or a nitrogen protecting group.



R8 represents a hydrogen atom, a deuterium atom, or a group wherein, R8a, R8b and R8c, identical or different, represent a hydrogen atom or a deuterium atom.
 25 R4, R5, R6, and R7, identical or different, represent a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, an (C1-C6)alkylcarbonyl group, an (C1-C6)alkyl group, an (C1-C6)alkyloxy group, an (C1-C6)alkylthio group, an (C1-C6)alkylcarbonyloxy group, an (C6-C14)aryloxy group, an (C6-C14)aryl group, a heterocyclic group, a (C3-C14)cycloalkyl group, a NO₂, a sulfonylamino(C1-C6)alkyl group, an NH₂ group, an amino(C1-C6)alkyl
 30 group, an (C1-C6)alkylcarbonylamino group, a carboxylic group, a carboxylate group, an

amino acid selected from the group consisting of alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, or a moiety of formula (A):




wherein R' represents an (C1-C6)alkyl group, an (C2-C6)alkenyl group, an (C2-C6)alkynyl group, a (C3-C14)cycloalkyl group, (C3-C14)cycloalkyl(C1-C6)alkyl group, a (C3-C14)cycloalkyl(C1-C6)alkenyl group, a (C3-C14)cycloalkenyl group, a (C3-C14)cycloalkenyl(C1-C6)alkyl group, a (C3-C14)cycloalkenyl(C2-C6)alkenyl group, a (C3-C14)cycloalkenyl(C2-C6)alkynyl group; R'' and R''', independently, represent a hydrogen atom, an (C1-C6)alkyl group, or a nitrogen protecting group.

In a particular embodiment, in the compound of formula (I) of the present invention:

- an alkyl group may be a substituted or unsubstituted (C1-C6)alkyl group, in particular a substituted or unsubstituted (C1-C4)alkyl group;
- an alkynyl group may be a substituted or unsubstituted (C2-C6)alkynyl group;
- a cycloalkyl group may be a substituted or unsubstituted (C3-C14)cycloalkyl group
- an alkyloxy group may be a substituted or unsubstituted (C1-C6)alkyloxy group, such as a substituted or unsubstituted (C1-C4)alkyloxy group;
- an alkylthio group may be a substituted or unsubstituted (C1-C6)alkylthio group, such as a substituted or unsubstituted (C1-C4)alkylthio group;
- an alkylamino group may be a (C1-C6)alkylamino group, such as a (C1-C4)alkylamino group;
- a dialkylamino group may be a (C1-C6)dialkylamino group, such as a (C1-C4)dialkylamino group;
- an aryl group may be a substituted or unsubstituted (C6-C14)aryl group, such as a substituted or unsubstituted (C6-C14)aryl group;
- a heterocyclic group may be a substituted or unsubstituted heterocycloalkyl or heteroaryl group.


In a particular embodiment, the compound of component (i) of the combination product is a compound of formula (II)

6



(II),

in which R₉ represents a hydrogen atom, a deuterium atom, a O-R₈ group (R₈ being as defined above), or an amino acid selected from the group consisting of alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, or a moiety of formula (A):




(A);

wherein R' represents an (C1-C6)alkyl group, an (C2-C6)alkenyl group, an (C2-C6)alkynyl group, a (C3-C14)cycloalkyl group, (C3-C14)cycloalkyl(C1-C6)alkyl group, a (C3-C14)cycloalkyl(C1-C6)alkenyl group, a (C3-C14)cycloalkenyl group, a (C3-C14)cycloalkenyl(C1-C6)alkyl group, a (C3-C14)cycloalkenyl(C2-C6)alkenyl group, a (C3-C14)cycloalkenyl(C2-C6)alkynyl group; R'' and R''', independently, represent a hydrogen atom, an (C1-C6)alkyl group, or a nitrogen protecting group or a pharmaceutically acceptable salt thereof.

In a particular embodiment, component (i) of the combination product of the invention is selected from:


- NTZ:




;

- tizoxanide (TZ):

7



- tizoxanide glucuronide (TZG):



- 5 In another embodiment, component (i) of the combination product is a compound of formula (II) wherein R8a, R8b and R8c, identical or different, represent a hydrogen atom or a deuterium atom; and/or

wherein R1, R3, R4, R5, and R6, identical or different, represent a hydrogen atom or a deuterium atom with the proviso that R1, R2, R8a, R8b, R8c, R3, R4, R5, and R6 are not
10 simultaneously a hydrogen atom.


In a particular embodiment, component (i) is [(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d3)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d2) ethanoate; or 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d1) ethanoate.

15


In another particular embodiment, component (i) is 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate, in particular (S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate, or a pharmaceutically acceptable salt thereof such as its hydrochloride salt (RM5061) of formula:

20

8




In another particular embodiment, component (i) is 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate, in particular (2S,3S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate, or a pharmaceutically acceptable salt thereof such as its hydrochloride salt (RM5066) of formula:




10

In another particular embodiment, component (i) is 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate, in particular (S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate, or a pharmaceutically acceptable salt thereof such as its hydrochloride salt (RM5064) of formula:


15



9



In another particular embodiment, component (i) is -2-(5-chlorothiazol-2-ylcarbamoyl)phenyl
2-amino-3-methylpentanoate, in particular (2S,3S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl
5 2-amino-3-methylpentanoate, or a pharmaceutically acceptable salt thereof such as its
hydrochloride salt (RM5065) of formula:



10

In a particular embodiment, component (i) is NTZ or a pharmaceutically acceptable salt thereof.

In the present invention, component (ii) is selected from an anti-NASH agent, an anti-fibrotic
15 agent or an anti-cholestatic agent, with the proviso that the anti-NASH, anti-fibrotic or anti-cholestatic agent.

In a particular embodiment, the anti-NASH, anti-fibrotic or anti-cholestatic agent is not a
PPAR agonist.

20

In a further particular embodiment of the invention, component (ii) is an anti-NASH agent. Illustrative, non-limiting, anti-NASH agents useful in the practice of the present invention include:

- Acetyl-CoA Carboxylase inhibitors;
- 5 - Anti-LPS antibodies;
- Apical sodium-codependent bile acid transporter inhibitors;
- bioactive lipids;
- Cannabinoid CB1 receptor antagonists;
- Caspase inhibitors;
- 10 - Cathepsin inhibitors;
- CCR antagonists;
- Diacylglycerol-O-acyltransferase (DGAT) inhibitors;
- Dipeptidyl peptidase IV (DPP4) inhibitors;
- Dual NOX (NADPH oxidase) 1&4 inhibitors;
- 15 - Extracellular matrix protein modulators;
- Stearoyl CoA desaturase-1 inhibitors/fatty acid bile acid conjugates (FABAC);
- Fatty Acid Synthase (FAS) inhibitors;
- Fibroblast Growth Factor 19 (FGF-19) recombinants;
- Fibroblast Growth Factor 21 (FGF-21) agonists;
- 20 - Farnesoid X Receptor (FXR) agonists;
- Galectin 3 inhibitors;
- Glucagon-Like Peptide-1 (GLP-1) analogs and GLP-1 receptor agonists;
- G-Protein Coupled Receptor (GPCR) modulators;
- Integrin inhibitors;
- 25 - Leukotriene (LT)/Phosphodiesterase (PDE)/Lipoxygenase (LO) inhibitors;
- Macrolides;
- miRNA antagonists;
- Monoclonal antibodies;
- Myeloperoxidase inhibitors;
- 30 - mTOR modulators;
- nuclear receptor ligands;
- P2Y13 protein agonists;
- Protease-Activated Receptor (PAR)-2 antagonists;
- Protein kinase modulators
- 35 - Rho-associated protein kinase 2 (ROCK2) inhibitors;
- Sodium-GLucose Transport (SGLT) 1 and 2 inhibitors;

- signal-regulating kinase 1 (ASK1) inhibitors;
- Thyroid Hormone Receptor β (THR β) agonists;
- Toll Like Receptor 2 and 4 (TLR-2 and 4) antagonists;
- Tyrosine Kinase Receptor (RTK) modulators;
- 5 - Vascular Adhesion Protein-1 (VAP-1) inhibitors; and
- Vitamin D receptor (VDR) agonists.

In a further particular embodiment, component (ii) is an anti-NASH agent. Illustrative, non-limiting, anti-NASH agents useful in the practice of the present invention include:

- 10 - Acetyl-CoA Carboxylase inhibitors;
- Anti-LPS antibodies;
- Apical sodium-codependent bile acid transporter inhibitors;
- bioactive lipids;
- Cannabinoid CB1 receptor antagonists;
- 15 - Caspase inhibitors;
- Cathepsin inhibitors;
- CCR antagonists;
- Diacylglycerol-O-acyltransferase (DGAT) inhibitors;
- Dipeptidyl peptidase IV (DPP4) inhibitors;
- 20 - Dual NOX (NADPH oxidase) 1&4 inhibitors;
- Extracellular matrix protein modulators;
- Stearoyl CoA desaturase-1 inhibitors/fatty acid bile acid conjugates (FABAC);
- Fibroblast Growth Factor 19 (FGF-19) recombinants;
- Fibroblast Growth Factor 21 (FGF-21) agonists;
- 25 - Farnesoid X Receptor (FXR) agonists;
- Galectin 3 inhibitors;
- Glucagon-Like Peptide-1 (GLP-1) analogs;
- G-Protein Coupled Receptor (GPCR) modulators;
- Integrin inhibitors;
- 30 - Leukotriene (LT)/Phosphodiesterase (PDE)/Lipoxygenase (LO) inhibitors;
- Macrolides;
- miRNA antagonists ;
- Monoclonal antibodies;
- mTOR modulators;
- 35 - nuclear receptor ligands;
- P2Y13 protein agonists;

- Protease-Activated Receptor (PAR)-2 antagonists;
- Protein kinase modulators;
- Rho-associated protein kinase 2 (ROCK2) inhibitors;
- Sodium-GLucose Transport (SGLT) 2 inhibitors;
- 5 - signal-regulating kinase 1 (ASK1) inhibitors;
- Thyroid Hormone Receptor β (THR β) agonists;
- Toll Like Receptor 4 (TLR- 4) antagonists;
- Tyrosine Kinase Receptor (RTK) modulators;
- Vascular Adhesion Protein-1 (VAP-1) inhibitors and
- 10 - Vitamin D receptor (VDR) agonists.

Other anti-NASH agents include KB-GE-001, NGM-386, NGM-395, NC-10, and TCM-606F.

In a particular embodiment of the invention, component (ii) is an anti-fibrotic agent.
 15 Illustrative, non-limiting, anti-fibrotic agents useful in the practice of the present invention include:

- Angiotensin II receptor blockers
- antisense oligonucleotides targeting Transforming Growth Factor beta 2 (TGF- β 2);
- Bioactive lipids;
- 20 - Caspase inhibitors;
- Dual Farnesoid X receptor (FXR) / TGR5 agonists;
- NOX (NADPH oxidase) inhibitors, such as Dual NOX (NADPH oxidase) 1&4 inhibitors;
- Galectin 3 inhibitors;
- 25 - Immunomodulators;
- Integrin inhibitors;
- Macrophage mannose receptor modulators;
- Metalloprotease-9 (MMP-9) stimulators;
- Monoclonal antibodies;
- 30 - NF-kappa B inhibitors;
- Non-Steroid Anti-Inflammatory Drugs (NSAIDs); and
- PDGFR modulators.

In a further particular embodiment of the invention, component (ii) is an anti-fibrotic agent.
 35 Illustrative, non-limiting, anti-fibrotic agents useful in the practice of the present invention include:

- antisense oligonucleotides targeting Transforming Growth Factor beta 2 (TGF- β 2);
- Bioactive lipids;
- Caspase inhibitors;
- Dual Farnesoid X receptor (FXR) / TGR5 agonists;
- 5 - Dual NOX (NADPH oxidase) 1&4 inhibitors;
- Galectin 3 inhibitors;
- Immunomodulators;
- Integrin inhibitors;
- Macrophage mannose receptor modulators;
- 10 - Metalloprotease-9 (MMP-9) stimulators;
- Monoclonal antibodies;
- NF-kappa B inhibitors;
- Non-Steroid Anti-Inflammatory Drugs (NSAIDs); and
- PDGFR modulators.

15

Other anti-fibrotic agents include HEC-585, INV-240, RNAi therapeutic (e.g. Silence Therapeutics) and SAMiRNA program (Bioneer Corp).

Other illustrative antifibrotic agents include pirfenidone or receptor tyrosine kinase inhibitors (RTKIs) such as Nintedanib, Sorafenib and other RTKIs, or angiotensin II (AT1) receptor blockers, or CTGF inhibitor, or any antifibrotic compound susceptible to interfere with the TGF β and BMP-activated pathways including activators of the latent TGF β complex such as MMP2, MMP9, THBS1 or cell-surface integrins, TGF β receptors type I (TGFBRI) or type II (TGFBRII) and their ligands such as TGF β , Activin, inhibin, Nodal, anti-Müllerian hormone, 25 GDFs or BMPs, auxiliary co-receptors (also known as type III receptors), or components of the SMAD-dependent canonical pathway including regulatory or inhibitory SMAD proteins, or members of the SMAD-independent or non-canonical pathways including various branches of MAPK signaling, TAK1, Rho-like GTPase signaling pathways, phosphatidylinositol-3 kinase/AKT pathways, TGF β -induced EMT process, or canonical and non-canonical 30 Hedgehog signaling pathways including Hh ligands or target genes, or any members of the WNT, or Notch pathways which are susceptible to influence TGF β .

According to a particular embodiment, the anti-fibrotic agent is not an agent selected from: pirfenidone or receptor tyrosine kinase inhibitors (RTKIs) such as Nintedanib, Sorafenib and other RTKIs, or angiotensin II (AT1) receptor blockers, or CTGF inhibitor, or any antifibrotic 35 compound susceptible to interfere with the TGF β and BMP-activated pathways including activators of the latent TGF β complex such as MMP2, MMP9, THBS1 or cell-surface

integrins, TGF β receptors type I (TGFBRI) or type II (TGFBRII) and their ligands such as TGF β , Activin, inhibin, Nodal, anti-Müllerian hormone, GDFs or BMPs, auxiliary co-receptors (also known as type III receptors), or components of the SMAD-dependent canonical pathway including regulatory or inhibitory SMAD proteins, or members of the SMAD-
5 independent or non-canonical pathways including various branches of MAPK signaling, TAK1, Rho-like GTPase signaling pathways, phosphatidylinositol-3 kinase/AKT pathways, TGF β -induced EMT process, or canonical and non-canonical Hedgehog signaling pathways including Hh ligands or target genes, or any members of the WNT, or Notch pathways which are susceptible to influence TGF β .

10

In a particular embodiment of the invention, component (ii) is an anti-cholestatic agent. Illustrative, non-limiting, anti-cholestatic agents useful in the practice of the present invention include:

- Apical Sodium-codependent Bile acid Transporter inhibitors (ASBTi);
- 15 - Bile acids;
- cathepsin inhibitors;
- CCR antagonists;
- CD40 inhibitors;
- CD80 inhibitors;
- 20 - NOX (NADPH oxidase) inhibitors, such as dual NOX (NADPH oxidase) 1&4 inhibitors;
- Farnesoid X receptor (FXR) agonists;
- Fibroblast Growth Factor (FGF) 19 recombinant;
- Fractalkine ligand inhibitors;
- 25 - ileal sodium bile acid cotransporter inhibitors; and
- Monoclonal antibodies.

In a further particular embodiment of the invention, component (ii) is an anti-cholestatic agent. Illustrative, non-limiting, anti-cholestatic agents useful in the practice of the present
30 invention include:

- Apical Sodium-codependent Bile acid Transporter inhibitors (ASBTi);
- Bile acids;
- cathepsin inhibitors;
- CCR antagonists;
- 35 - CD40 inhibitors;
- CD80 inhibitors;

- dual NOX (NADPH oxidase) 1&4 inhibitors;
- Farnesoid X receptor (FXR) agonists;
- Fibroblast Growth Factor (FGF) 19 recombinant;
- Fractalkine ligand inhibitors;
- 5 - ileal sodium bile acid cotransporter inhibitors; and
- Monoclonal antibodies.

According to the invention, the term “acetyl-CoA carboxylase inhibitors” as used herein includes, but is not limited to GS-0976, ND-654, AC-8632, and PF05175157.

10 According to the invention, the term “angiotensin type 1 receptor antagonists” as used herein includes, but is not limited to, Irbesartan.

According to the invention, the term “anti-LPS antibodies” as used herein includes, but is not limited to IMM-124-E.

15 According to the invention, the term “antisense oligonucleotide targeting transforming growth factor beta 2” as used herein includes, but is not limited to ASPH-0047, IMC-TR1, and ISTH-0047.

According to the invention, the term “apical sodium-codependent bile acid transporter inhibitor” as used herein includes, but is not limited to A-4250, volixibat, maralixibat formerly SHP-625, GSK-2330672, elobixibat, and CJ-14199.

20 According to the invention, the term “bile acids” as used herein includes, but is not limited to obeticholic acid, UDCA, norursodeoxycholic acid, and ursodiol.

According to the invention, the term “bioactive lipids” as used herein includes, but is not limited to 5-hydroxyeicosapentaenoic acid (15-HEPE, DS-102). In a further particular embodiment, the bioactive lipid may be selected from unsaturated fatty acids such as
25 arachidonic acid, eicosapentanoic acid, and docosahexaenoic acid.

According to the invention, the term “cannabinoid CB1 receptor antagonists” as used herein includes, but is not limited to nacamizumab, GRC-10801, MRI-1569, MRI-1867, DBPR-211, AM-6527, AM-6545, NESS-11-SM, CXB-029, GCC-2680, TM-38837, Org-50189, PF-514273, BMS-812204, ZYO-1, AZD-2207, AZD-1175, otenabant, ibipinabant,
30 surinabant, rimonabant, drinabant, SLV-326, V-24343, and O-2093.

According to the invention, the term “caspase inhibitors” as used herein includes, but is not limited to emricasan, belnacasan, nivocasan, IDN-7314, F-573, VX-166, YJP-60107, MX-1122, IDN-6734, TLC-144, SB-234470, IDN-1965, VX-799, SDZ-220-976, and L-709049.

According to the invention, the term “cathepsin inhibitors” as used herein includes, but
35 is not limited to VBY-376, VBY-825, VBY-036, VBY-129, VBY-285, Org-219517, LY3000328, RG-7236, and BF/PC-18.

According to the invention, the term “CCR antagonists” as used herein includes, but is not limited to cenicriviroc (CCR2/5 antagonist), PG-092, RAP-310, INCB-10820, RAP-103, PF-04634817, and CCX-872.

According to the invention, the term “CD40 inhibitors” as used herein includes, but is not limited to FFp-104, xl-050, DOM-0800, XmAb-5485, KGY-15, FFP-106, TDI-0028, and ABI-793.

According to the invention, the term “CD80 inhibitors” as used herein includes, but is not limited to RhuDex, FPT-155, Tolerimab, galiximab, SCH-212394, IGM-001, ASP-2408, and SCH-204698.

According to the invention, the term “diacylglycerol-O-acyltransferase inhibitors” as used herein includes, but is not limited to IONIS-DGAT2Rx formerly ISIS-DGAT2Rx, LY-3202328, BH-03004, KR-69530, OT-13540, AZD-7687, and ABT-046.






According to the invention, the term “dipeptidyl peptidase IV inhibitors” as used herein includes, but is not limited to evogliptin, vidagliptin, fotagliptin, alogliptin, saxagliptin, tilagliptin, anagliptin, sitagliptin, retagliptin, melogliptin, gosogliptin, trelagliptin, teneligliptin, dutogliptin, linagliptin, gemigliptin, yogliptin, betagliptin, imigliptin, omarigliptin, vidagliptin, and denagliptin.






According to the invention, the term “dual Farnesoid X receptor (FXR) / TGR5 agonists” as used herein includes, but is not limited to INT-767.


According to the invention, the term “dual NOX (NADPH oxidase) 1&4 inhibitors” as used herein includes, but is not limited to GKT-831 (2-(2-chlorophenyl)-4-[3-(dimethylamino)phenyl]-5-methyl-1H-pyrazolo[4,3-c]pyridine-3,6(2H,5H)-dione), formerly GKT137831, and GKT-901.

According to the invention, the term “extracellular matrix protein modulators” as used herein includes, but is not limited to CNX-024, CNX-025, and SB-030.


According to the invention, the term “Fatty Acid Synthase (FAS) inhibitors” as used herein include, but is not limited to TVB-2640; TVB-3664; TVB-3166 ; TVB-3150 ; TVB-3199 ; TVB-3693; BZL-101 ; 2-octadecynoic acid ; MDX-2 ; Fasnall ; MT-061 ; G28UCM ; MG-28 ; HS-160 ; GSK-2194069 ; KD-023 ; cilostazol. In a particular embodiment, the FAS inhibitor is a compound selected in the following list of compounds:








In another particular embodiment, the FAS inhibitor is selected from:



5



In a particular embodiment, the FAS inhibitor is TVB-2640.

10 According to the invention, the term “Farnesoid X receptor (FXR) agonists” as used herein includes, but is not limited to obeticholic acid, GS-9674, LJM-452, LJM-763, LMB763, EDP-305, AKN-083, INT-767, GNF-5120, LY2562175, INV-33, NTX-023-1, EP-024297, EPY-001, Px-103, and SR-45023.

15 According to the invention, the term “Fatty Acid Bile Acid Conjugates, FABAC)” as used herein includes but is not limited to Aramchol.

According to the invention, the term “Fractalkine ligand inhibitors” as used herein includes, but is not limited to E-6011, and KAN-0440567.

According to the invention, the term “Fibroblast Growth Factor 19 (FGF-19) recombinants” as used herein includes, but is not limited to NGM-282.

According to the invention, the term “Fibroblast Growth Factor 21 (FGF-21) agonists” as used includes, but is not limited to PEG-FGF21 formerly BMS-986036, YH-25348, BMS-986171, YH-25723, LY-3025876, and NNC-0194-0499.

According to the invention, the term “Galectin 3 inhibitors” as used herein includes, but is not limited to GR-MD-02, TD-139, ANG-4021, Galectin-3C, LJPC-201, TFD-100, GR-MD-03, GR-MD-04, GM-MD-01, GM-CT-01, GM-CT-02, Gal-100, and Gal-200.

According to the invention, the term “Glucagon-like peptide-1 (GLP-1) analogs” as used herein includes, but is not limited to semaglutide, liraglutide, exenatide, albiglutide, dulaglutide, lixisenatide, loxenatide, efpeglenatide, taspoglutide, MKC-253, DLP-205, and ORMD-0901.

According to the invention, the term “Glucagon-like peptide-1 (GLP-1) receptor agonist” as used herein includes, but is not limited to LY-3305677, and Oxyntomodulin long acting.

According to the invention, the term “G-protein coupled receptor (GPCR) modulators” as used herein includes, but is not limited to CNX-023.

According to the invention, the term “ileal sodium bile acid cotransporter inhibitors” as used herein includes, but is not limited to A-4250, GSK-2330672, volixibat, CJ-14199, and elobixibat.

According to the invention, the term “immunomodulators” as used herein includes, but is not limited to PBI-4050, and AIC-649.

According to the invention, the term “integrin inhibitors” as used herein includes, but is not limited to integrin inhibitors of Pliant Therapeutic, integrin inhibitors of Indalo Therapeutics, integrin inhibitors of St Louis University, ProAgio, and GSK-3008348.

According to the invention, the term “leukotriene/phosphodiesterase /lipoxygenase inhibitors” as used includes, but is not limited to tielukast (formerly MN-001), tomelukast, sulukast, masilukast, zafirlukast, pranlukast, montelukast, gemilukast, verlukast, aklukast, pobilikast, cinalukast, and iralukast.

According to the invention, the term “macrolides” as used herein includes, but is not limited to solithromycin, azithromycin, and erythromycin .

According to the invention, the term “macrophage mannose receptor modulators” as used herein includes, but is not limited to AB-0023, MT-1001, [18F]FB18mHSA, Xemys,technetium Tc 99m tilmanocept, and CDX-1307.

According to the invention, the term “miRNA antagonists” as used herein includes, but is not limited to RG-125 formerly AZD4076, RGLS-5040, RG-101, MGN-5804, and MRG-201.

According to the invention, the term “metalloprotease-9 (MMP-9) stimulators” as used herein includes, but is not limited to MMP-9 stimulator of Elastomics Ab.

According to the invention, the term “myeloperoxidase inhibitor” as used herein includes, but is not limited to PF-06667272.

According to the invention, the term “monoclonal antibodies” as used includes, but is not limited to bertilimumab, NGM-313, IL-20 targeting mAbs, fresolimumab (antiTGFβ) 5 formerly GC1008, timolumab formerly BTT-1023, namacizumab, omalizumab, ranibizumab, bevacizumab, lebrikizumab, epratuzumab, felvizumab, matuzumab, monalizumab, reslizumab, inebilizumab, foralumab (NI-0401, anti-CD3), simtizumab (GS-6624) mAb against LOXL2, and ustekinumab an anti-TNF antibody.

In a particular embodiment, the monoclonal antibody is selected in the group consisting 10 of anti-IL20 mAbs, anti-TGFβ antibodies, anti-CD3 antibodies, anti-LOXL2 antibodies and anti-TNF antibodies.

According to the invention, the term “mTOR modulators” as used herein includes, but is not limited to MSDC-0602, AAV gene therapy co-administered with SVP-sirolimus.

According to the invention, the term “NF-kappa B inhibitors” as used herein includes, 15 but is not limited to LC-280126.

According to the invention, the term “non-steroid anti-inflammatory drugs (NSAIDs)” as used herein includes, but is not limited to F-351, salicylates (aspirin), acetaminophen, propionic acid derivatives (ibuprofen, naproxen), acetic acid derivatives (indomethacin, diclofenac), enolic acid derivatives (piroxicam, phenylbutazone), anthranilic acid derivatives 20 (meclofenalmic acid, flufenamic acid), selective COX-2 inhibitors (celecoxib, parecoxib), and sulfonanilides (nimesulide).

According to the invention, the term “nuclear receptor ligands” as used herein includes, but is not limited to DUR-928 formerly DV 928.

According to the invention, the term “P2Y13 protein agonists” as used herein includes, 25 but is not limited to CER-209.

According to the invention, the term “PDGFR modulators” as used herein includes, but is not limited to BOT-501, and BOT-191.

According to the invention, the term “protein kinase modulators” as used herein includes, but not limited to, CNX-014, MB-11055, ALF-1, mangiferin, amlexanox, GS- 30 444217, REG-101, and valine.

According to the invention, the term “Rho-associated protein kinase 2 (ROCK2) inhibitors” as used herein includes, but is not limited to KD-025, TRX-101, BA-1049, LYC- 53976, INS-117548, and RKI-1447.

According to the invention, the term “signal-regulating kinase 1 (ASK1) inhibitors” as 35 used herein includes, but is not limited to selonsertib (formerly GS-4997).

According to the invention, the term “sodium-glucose transport (SGLT) 1 inhibitors” as used herein includes, but is not limited to LIK-066 (Licogliflozin).

According to the invention, the term “sodium-glucose transport (SGLT) 2 inhibitors” as used herein includes, but is not limited to remogliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin, ipragliflozin, tianagliflozin, canagliflozin, tofogliflozin, janagliflozin, bexagliflozin, luseogliflozin, sergliflozin, HEC-44616, AST-1935, and PLD-101.

According to the invention, the term “stearoyl CoA desaturase-1 inhibitors/fatty acid bile acid conjugates” as used herein includes, but is not limited to aramchol, GRC-9332, steamchol, TSN-2998, GSK-1940029, and XEN-801.

According to the invention, the term “Thyroid Hormone Receptor β (THR β) agonists” as used herein includes, but is not limited to VK-2809 ; MGL-3196 ; MGL-3745 ; SKL-14763 ; sobetirome ; BCT-304 ; ZYT-1 ; MB-07811 ; and eprotirome.

According to the invention, the term “Toll Like Receptor 4 (TLR-4) antagonists” as used herein includes, but is not limited to naltrexone, JKB-121 also known as Nalmefene, M-62812, resatorvid, dendrophilin, CS-4771, AyuV-1, AyuV-25, NI-0101, EDA-HPVE7, and eritoran.

According to the invention, the term “Toll Like Receptor 2 and 4 (TLR-2) antagonists” as used herein includes, but is not limited to CI-201 also known as VB-201.

According to the invention, the term “tyrosine kinase receptor (RTK) modulators” as used herein includes, but is not limited to, RTK inhibitors. In a particular embodiment of the invention, the RTK modulators include CNX-025, and KBP-7018. In a particular embodiment, the RTK modulator is not a PPAR agonist.

According to the invention, the term “vascular adhesion protein-1 (VAP-1) inhibitors” as used herein includes, but is not limited to PXS-4728A, CP-664511, PRX-167700, ASP-8232, RTU-1096, RTU-007, and BTT-1023.

According to the invention, the term “vitamin D receptor (VDR) agonists” as used herein includes, but is not limited to calciferol, alfacalcidol, 1,25-dihydroxyvitamin D3, Vitamin D2, Vitamin D3, calcitriol, Vitamin D4, Vitamin D5, dihydrotachysterol, calcipotriol, tacalcitol 1,24- dihydroxyvitamin D3 and paricalcitol.

In a particular embodiment of the combination product of the invention:

- component (ii) is an anti-NASH, anti-fibrotic or anti-cholestatic; and
- component (i) is selected from NTZ, TZ, TZG, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d3)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d2)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d1) ethanoate, 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-nitrothiazol-2-

ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), and 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt).

In a particular variant of this embodiment, component (ii) is not a PPAR agonist.

10

In a particular embodiment, component (i) is selected from NTZ, TZ, TZG, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d3)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d2) ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d1) ethanoate, 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), and 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), component (i) being more particularly NTZ or a pharmaceutically acceptable salt thereof; and

component (ii) is an anti-NASH agent selected in the group consisting of:

- 25 - Acetyl-CoA Carboxylase inhibitors;
- Anti-LPS antibodies;
- Apical sodium-codependent bile acid transporter inhibitors;
- bioactive lipids;
- Cannabinoid CB1 receptor antagonists;
- 30 - Caspase inhibitors;
- Cathepsin inhibitors;
- CCR antagonists;
- Diacylglycerol-O-acyltransferase (DGAT) inhibitors;
- Dipeptidyl peptidase IV (DPP4) inhibitors;
- 35 - Dual NOX (NADPH oxidase) 1&4 inhibitors;
- Extracellular matrix protein modulators;

- Stearoyl CoA desaturase-1 inhibitors/fatty acid bile acid conjugates (FABAC);
- Fatty Acid Synthase (FAS) inhibitors;
- Fibroblast Growth Factor 19 (FGF-19) recombinants;
- Fibroblast Growth Factor 21 (FGF-21) agonists;
- 5 - Farnesoid X Receptor (FXR) agonists;
- Galectin 3 inhibitors;
- Glucagon-Like Peptide-1 (GLP-1) analogs and GLP-1 receptor agonists;
- G-Protein Coupled Receptor (GPCR) modulators;
- Integrin inhibitors;
- 10 - Leukotriene (LT)/Phosphodiesterase (PDE)/Lipoxygenase (LO) inhibitors;
- Macrolides;
- miRNA antagonists;
- Monoclonal antibodies;
- Myeloperoxidase inhibitors;
- 15 - mTOR modulators;
- nuclear receptor ligands;
- P2Y₁₃ protein agonists;
- Protease-Activated Receptor (PAR)-2 antagonists;
- Protein kinase modulators
- 20 - Rho-associated protein kinase 2 (ROCK2) inhibitors;
- Sodium-GLucose Transport (SGLT) 1 and 2 inhibitors;
- signal-regulating kinase 1 (ASK1) inhibitors;
- Thyroid Hormone Receptor β (THR β) agonists;
- Toll Like Receptor 2 and 4 (TLR-2 and 4) antagonists;
- 25 - Tyrosine Kinase Receptor (RTK) modulators;
- Vascular Adhesion Protein-1 (VAP-1) inhibitors;
- Vitamin D receptor (VDR) agonists;
- KB-GE-001;
- NGM-386;
- 30 - NGM-395;
- NC-10; and
- TCM-606F.

In a particular embodiment, component (i) is selected from NTZ, TZ, TZG, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d3)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d2) ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d1) ethanoate, 2-(5-nitrothiazol-

2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), and 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), component (i) being more particularly NTZ or a pharmaceutically acceptable salt thereof; and

component (ii) is an anti-NASH agent selected in the group consisting of:

- Acetyl-CoA Carboxylase inhibitors;
- Anti-LPS antibodies;
- Apical sodium-codependent bile acid transporter inhibitors;
- 15 - bioactive lipids;
- Cannabinoid CB1 receptor antagonists;
- Caspase inhibitors;
- Cathepsin inhibitors;
- CCR antagonists;
- 20 - Diacylglycerol-O-acyltransferase (DGAT) inhibitors;
- Dipeptidyl peptidase IV (DPP4) inhibitors;
- Dual NOX (NADPH oxidase) 1&4 inhibitors;
- Extracellular matrix protein modulators;
- Stearoyl CoA desaturase-1 inhibitors/fatty acid bile acid conjugates (FABAC);
- 25 - Fibroblast Growth Factor 19 (FGF-19) recombinants;
- Fibroblast Growth Factor 21 (FGF-21) agonists;
- Farnesoid X Receptor (FXR) agonists;
- Galectin 3 inhibitors;
- Glucagon-Like Peptide-1 (GLP-1) analogs;
- 30 - G-Protein Coupled Receptor (GPCR) modulators;
- Integrin inhibitors;
- Leukotriene (LT)/Phosphodiesterase (PDE)/Lipoxygenase (LO) inhibitors;
- Macrolides;
- miRNA antagonists ;
- 35 - Monoclonal antibodies;
- mTOR modulators;

- nuclear receptor ligands;
- P2Y₁₃ protein agonists;
- Protease-Activated Receptor (PAR)-2 antagonists;
- Protein kinase modulators;
- 5 - Rho-associated protein kinase 2 (ROCK2) inhibitors;
- Sodium-GLucose Transport (SGLT) 2 inhibitors;
- signal-regulating kinase 1 (ASK1) inhibitors;
- Thyroid Hormone Receptor β (THR β) agonists;
- Toll Like Receptor 4 (TLR- 4) antagonists;
- 10 - Tyrosine Kinase Receptor (RTK) modulators;
- Vascular Adhesion Protein-1 (VAP-1) inhibitors;
- Vitamin D receptor (VDR) agonists;
- KB-GE-001;
- NGM-386;
- 15 - NGM-395;
- NC-10; and
- TCM-606F.

In a particular embodiment, component (i) is selected from NTZ, TZ, TZG, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d3)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d2) ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d1) ethanoate, 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), and 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), component (i) being more particularly NTZ or a pharmaceutically acceptable salt thereof; and

component (ii) is an anti-fibrotic agent selected in the group consisting of:

- Angiotensin II receptor blockers
- antisense oligonucleotides targeting Transforming Growth Factor beta 2 (TGF- β 2);
- 35 - Bioactive lipids;
- Caspase inhibitors;

- Dual Farnesoid X receptor (FXR) / TGR5 agonists;
- NOX (NADPH oxidase) inhibitors, such as Dual NOX (NADPH oxidase) 1&4 inhibitors;
- Galectin 3 inhibitors;
- 5 - Immunomodulators;
- Integrin inhibitors;
- Macrophage mannose receptor modulators;
- Metalloprotease-9 (MMP-9) stimulators;
- Monoclonal antibodies;
- 10 - NF-kappa B inhibitors;
- Non-Steroid Anti-Inflammatory Drugs (NSAIDs);
- PDGFR modulators;
- HEC-585,
- INV-240,
- 15 - RNAi therapeutic (e.g. Silence Therapeutics); and
- the SAMiRNA program (Bioneer Corp).

In a particular embodiment, component (i) is selected from NTZ, TZ, TZG, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d3)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d2) ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d1) ethanoate, 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), and 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), component (i) being more particularly NTZ or a pharmaceutically acceptable salt thereof; and

30 component (ii) is an anti-fibrotic agent selected in the group consisting of:

- antisense oligonucleotides targeting Transforming Growth Factor beta 2 (TGF- β 2);
- Bioactive lipids;
- Caspase inhibitors;
- 35 - Dual Farnesoid X receptor (FXR) / TGR5 agonists;
- Dual NOX (NADPH oxidase) 1&4 inhibitors;

- Galectin 3 inhibitors;
 - Immunomodulators;
 - Integrin inhibitors;
 - Macrophage mannose receptor modulators;
 - 5 - Metalloprotease-9 (MMP-9) stimulators;
 - Monoclonal antibodies;
 - NF-kappa B inhibitors;
 - Non-Steroid Anti-Inflammatory Drugs (NSAIDs);
 - PDGFR modulators;
 - 10 - HEC-585,
 - INV-240,
 - RNAi therapeutic (e.g. Silence Therapeutics); and
 - the SAMiRNA program (Bioneer Corp).
- 15 In a particular embodiment, component (i) is selected from NTZ, TZ, TZG, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d3)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d2) ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d1) ethanoate, 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), and 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), component (i) being more particularly NTZ or a pharmaceutically acceptable salt thereof; and
- 20 component (ii) is an anti-cholestatic agent selected in the group consisting of:
- Apical Sodium-codependent Bile acid Transporter inhibitors (ASBTi);
 - 30 - Bile acids;
 - cathepsin inhibitors;
 - CCR antagonists;
 - CD40 inhibitors;
 - CD80 inhibitors;
 - 35 - NOX (NADPH oxidase) inhibitors, such as dual NOX (NADPH oxidase) 1&4 inhibitors;

- Farnesoid X receptor (FXR) agonists;
- Fibroblast Growth Factor (FGF) 19 recombinant;
- Fractalkine ligand inhibitors;
- ileal sodium bile acid cotransporter inhibitors;
- 5 - Monoclonal antibodies.

- In a particular embodiment, component (i) is selected from NTZ, TZ, TZG, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d3)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d2) ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d1) ethanoate, 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), and 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), component (i) being more particularly NTZ or a pharmaceutically acceptable salt thereof; and
- 20 component (ii) is an anti-cholestatic agent selected in the group consisting of:
- Apical Sodium-codependent Bile acid Transporter inhibitors (ASBTi);
 - Bile acids;
 - cathepsin inhibitors;
 - CCR antagonists;
 - 25 - CD40 inhibitors;
 - CD80 inhibitors;
 - dual NOX (NADPH oxidase) 1&4 inhibitors;
 - Farnesoid X receptor (FXR) agonists;
 - Fibroblast Growth Factor (FGF) 19 recombinant;
 - 30 - Fractalkine ligand inhibitors;
 - ileal sodium bile acid cotransporter inhibitors; and
 - Monoclonal antibodies.

- In a particular embodiment, component (i) is selected from NTZ, TZ, TZG, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d3)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d2) ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d1) ethanoate, 2-(5-nitrothiazol-
- 35

2-ylcarbamoylethyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-nitrothiazol-2-ylcarbamoylethyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), 2-(5-nitrothiazol-2-ylcarbamoylethyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-nitrothiazol-2-ylcarbamoylethyl 2-amino-3-methylpentanoate or its hydrochloride salt), 2-(5-chlorothiazol-2-ylcarbamoylethyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-chlorothiazol-2-ylcarbamoylethyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), and 2-(5-chlorothiazol-2-ylcarbamoylethyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-chlorothiazol-2-ylcarbamoylethyl 2-amino-3-methylpentanoate or its hydrochloride salt), component (i) being more particularly NTZ or a pharmaceutically acceptable salt thereof; and
 component (ii) is a FAS inhibitor.

In a particular embodiment of the combination product of the invention:

- component (ii) is a ROCK2 inhibitor (in particular KD-025), an ACC inhibitor (in particular (GS-0976), an ASK1 inhibitor (in particular Selonsertib), a dual NOX1 and NOX4 inhibitor (in particular GKT-831, formerly GKT137831), a FABAC (in particular Aramchol), a FXR agonist (in particular Tropicifexor) or a CCR antagonist (in particular CVC); and

- component (i) is selected from NTZ, TZ, TZG, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoylethyl (d3)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoylethyl (d2)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoylethyl (d1) ethanoate, 2-(5-nitrothiazol-2-ylcarbamoylethyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-nitrothiazol-2-ylcarbamoylethyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), 2-(5-nitrothiazol-2-ylcarbamoylethyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-nitrothiazol-2-ylcarbamoylethyl 2-amino-3-methylpentanoate or its hydrochloride salt), 2-(5-chlorothiazol-2-ylcarbamoylethyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-chlorothiazol-2-ylcarbamoylethyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), and 2-(5-chlorothiazol-2-ylcarbamoylethyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-chlorothiazol-2-ylcarbamoylethyl 2-amino-3-methylpentanoate or its hydrochloride salt).

30

In a particular embodiment of the combination product of the invention:

- component (ii) is a ROCK2 inhibitor (in particular KD-025), an ACC inhibitor (in particular (GS-0976), an ASK1 inhibitor (in particular Selonsertib), a dual NOX1 and NOX4 inhibitor (in particular GKT-831, formerly GKT137831), a FABAC (in particular Aramchol), a FXR agonist (in particular Tropicifexor) or a CCR antagonist (in particular CVC); and

35

- component (i) is selected from NTZ, TZ, TZG, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d3)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d2)ethanoate, 2-[(5-nitro-1,3-thiazol-2-yl)carbamoyl]phenyl (d1) ethanoate, 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), 2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-nitrothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt), 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate (such as (S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3,3-dimethylbutanoate or its hydrochloride salt), and 2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate (such as (2S,3S)-2-(5-chlorothiazol-2-ylcarbamoyl)phenyl 2-amino-3-methylpentanoate or its hydrochloride salt).

In a particular embodiment, component (i) is NTZ or a pharmaceutically acceptable salt thereof, and component (ii) is a ROCK2 inhibitor (in particular KD-025), an ACC inhibitor (in particular (GS-0976), an ASK1 inhibitor (in particular Selonsertib), a dual NOX1 and NOX4 inhibitor (in particular GKT-831, formerly GKT137831), a FABAC (in particular Aramchol), a FXR agonist (in particular Tropifexor) or a CCR antagonist (in particular CVC).

In a particular embodiment, component (i) is NTZ or a pharmaceutically acceptable salt thereof, and component (ii) is a ROCK2 inhibitor (in particular KD-025)

In a particular embodiment, component (i) is NTZ or a pharmaceutically acceptable salt thereof, and component (ii) is an ACC inhibitor (in particular (GS-0976),

In a particular embodiment, component (i) is NTZ or a pharmaceutically acceptable salt thereof, and component (ii) is an ASK1 inhibitor (in particular Selonsertib)

In a particular embodiment, component (i) is NTZ or a pharmaceutically acceptable salt thereof, and component (ii) is or a dual NOX1 and NOX4 inhibitor (in particular GKT-831, formerly GKT137831).

In a particular embodiment, component (i) is NTZ or a pharmaceutically acceptable salt thereof, and component (ii) a FABAC (in particular Aramchol).

In a particular embodiment, component (i) is NTZ or a pharmaceutically acceptable salt thereof, and component (ii) a FXR agonist (in particular Tropicamorex).

5 In a particular embodiment, component (i) is NTZ or a pharmaceutically acceptable salt thereof, and component (ii) a CCR antagonist (in particular CCR).

In a particular embodiment, the combination product of the invention further comprises at least one other therapeutically active agent selected from JAK/STAT inhibitors and other anti-inflammatory agent and/or an immunosuppressant agent.

10

Illustrative anti-inflammatory and/or immunosuppressant agents are selected in the group consisting of glucocorticoids, NSAIDs, cyclophosphamide, nitrosoureas, folic acid analogs, purine analogs, pyrimidine analogs, methotrexate, azathioprine, mercaptopurine, ciclosporin, myriocin, tacrolimus, sirolimus, mycophenolic acid derivatives, fingolimod and other
15 sphingosine-1-phosphate receptor modulators, monoclonal and/or polyclonal antibodies against such targets as proinflammatory cytokines and proinflammatory cytokine receptors, T-cell receptor and integrins.

20

In another particular embodiment the synergistic combination of the invention may further comprise at least one therapeutically active agent with known antifibrotic activity selected from pirfenidone or receptor tyrosine kinase inhibitors (RTKIs) such as Nintedanib, Sorafenib and other RTKIs, or angiotensin II (AT1) receptor blockers, or CTGF inhibitor, or any antifibrotic compound susceptible to interfere with the TGF β and BMP-activated pathways including activators of the latent TGF β complex such as MMP2, MMP9, THBS1 or cell-
25 surface integrins, TGF β receptors type I (TGFBR1) or type II (TGFBR2) and their ligands such as TGF β , Activin, inhibin, Nodal, anti-Müllerian hormone, GDFs or BMPs, auxiliary co-receptors(also known as type III receptors), or components of the SMAD-dependent canonical pathway including regulatory or inhibitory SMAD proteins, or members of the SMAD-independent or non-canonical pathways including various branches of MAPK
30 signaling, TAK1, Rho-like GTPase signaling pathways, phosphatidylinositol-3 kinase/AKT pathways, TGF β -induced EMT process, or canonical and non-canonical Hedgehog signaling pathways including Hh ligands or target genes, or any members of the WNT, or Notch pathways which are susceptible to influence TGF β signaling.

35

In a specific embodiment of the invention, the other therapeutically active agent is a PPAR agonist.

In another particular embodiment, the PPAR agonist is a PPAR-alpha agonist, a PPAR-gamma agonist, a PPAR-delta agonist, a PPAR-alpha/gamma dual agonist, a PPAR alpha/delta dual agonist, a PPAR gamma/delta dual agonist or PPAR alpha/gamma/delta pan agonist.

In a particular embodiment, the other therapeutically active agent is:

- at least one PPAR-alpha agonist;
- at least one PPAR-gamma agonist;
- 10 - at least one PPAR-delta agonist;
- at least one PPAR-alpha/delta dual agonist;
- at least one PPAR-alpha agonist and at least one PPAR delta agonist;
- at least one PPAR-alpha/gamma dual agonist;
- at least one PPAR-alpha agonist and at least one PPAR gamma agonist;
- 15 - at least one PPAR-gamma/delta dual agonist;
- at least one PPAR-gamma agonist and at least one PPAR delta agonist;
- at least one PPAR-alpha/gamma/delta pan agonist; and
- at least one PPAR-alpha agonist, at least one PPAR-gamma agonist and at least one PPAR-delta agonist.

20

According to the present invention, the term "PPAR(s) agonists" refers the Peroxisome Proliferator Activated Receptor agonists, which are a class of drugs which plays a central role in lipid and glucose homeostasis. PPAR α mainly influences fatty acid metabolism and its activation lowers lipid levels, while PPAR γ is mostly involved in the regulation of the adipogenesis, energy balance, and lipid biosynthesis. PPAR δ participates in fatty acid oxidation, mostly in skeletal and cardiac muscles, but it also regulates blood glucose and cholesterol levels.

PPAR agonist may be in the form of a salt, hydrate, solvate, polymorph, or a co-crystal. PPAR agonist may also be in the form of a hydrate, solvate, polymorph, or a co-crystal of a salt.

According to the invention, the term "PPAR alpha agonists" as used herein includes, but is not limited to fenofibrate, ciprofibrate, pemafibrate, gemfibrozil, clofibrate, binifibrate, clinofibrate, clofibric acid, nicofibrate, pirifibrate, plafibrade, ronifibrate, theofibrate, tocofibrate, SR10171;

According to the invention, the term "PPAR gamma agonists" as used herein includes, but is not limited to Pioglitazone, deuterated pioglitazone, Rosiglitazone, efatutazone, ATx08-

001, OMS-405, CHS-131, THR-0921, SER-150-DN, KDT-501, GED-0507-34-Levo, CLC-3001, ALL-4.

According to the invention, the term "PPAR delta agonists" as used herein includes, but is not limited to GW501516 (Endurabol or ({4-[(4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl]sulfanyl}-2-methylphenoxy}acetic acid)) or MBX8025 (Seladelpar or {2-methyl-4-[5-methyl-2-(4-trifluoromethyl-phenyl)-2H-[1,2,3]triazol-4-ylmethylsulfanyl]-phenoxy}-acetic acid) or GW0742 ([4-[[[2-[3-fluoro-4-(trifluoromethyl)phenyl]-4-methyl-5-thiazolyl]methyl]thio]-2-methyl phenoxy]acetic acid) or L165041 or HPP-593 or NCP-1046.


According to the invention, the term "PPAR alpha/gamma agonists" (also named glitazars) used herein includes, but is not limited to Saroglitazar, Aleglitazar, Muraglitazar, Tesaglitazar, DSP-8658.

According to the invention, the term "PPAR alpha/delta agonists" used herein includes, but is not limited to Elafibranor (GFT505) or T913659.

According to the invention, the term "PPAR gamma/delta agonist" used herein includes, but is not limited to a conjugated linoleic acid (CLA), T3D-959.

According to the invention, the term "PPAR alpha/gamma/delta agonists" or "PPARpan agonists" used herein includes, but is not limited to IVA337 or TTA (tetradecylthioacetic acid) or Bavachinin or GW4148 or GW9135, or Bezafibrate or Lobeglitazone, or CS038. According to the invention, the term "protease-activated receptor (PAR)-2 antagonists" as used herein includes, but is not limited to PZ-235; NP-003.

In a more particular embodiment, the PPAR agonist is a compound of formula (III), or a pharmaceutically acceptable salt thereof:



in which:

Y1 represents a halogen, a Ra, or Ga-Ra group;

A represents a CH=CH or a CH2-CH2 group;

Y2 represents a Gb-Rb group;

Ga and Gb, identical or different, represent an atom of oxygen or sulfur;

Ra represents a hydrogen atom, an unsubstituted (C1-C6)alkyl group, a (C6-C14)aryl group or a (C1-C6)alkyl group that is substituted by one or more halogen atoms, a (C1-

C6)alkoxy or a (C1-C6)alkylthio group, (C3-C14)cycloalkyl groups, (C3-C14)cycloalkylthio groups or heterocyclic groups;

Rb represents a (C1-C6)alkyl group substituted by at least a –COORc group, wherein Rc represents a hydrogen atom, or a (C1-C6)alkyl group that is substituted or not by one or more halogen atoms, (C3-C14)cycloalkyl groups, or heterocyclic groups; and Y4 and Y5, identical or different, representing a (C1-C6)alkyl group that is substituted or not by one or more halogen atoms, (C3-C14)cycloalkyl groups or heterocyclic groups.

In a particular embodiment of the compound of formula (III):

Y1 represents a halogen, a Ra, or a Ga-Ra group;

A represents a CH=CH group;

Y2 represents a Gb-Rb group;

Ga and Gb, identical or different, represent an atom of oxygen or sulfur;

Ra represents a (C1-C6)alkyl or (C3-C14)cycloalkyl group, in particular a (C1-C7)alkyl or (C3-C14)cycloalkyl group substituted or not by one or more halogen atoms;

Rb represents a (C1-C6)alkyl group substituted by a –COOR3 group, wherein Rc represents a hydrogen atom or an alkyl group having from one to four carbon atoms; and

Y4 and Y5 independently represent a (C1-C4)alkyl group.

In a particular embodiment of the compound of formula (III):

Y1 represents a Ra or Ga-Ra group;

A represents a CH2-CH2 group;

Y2 represents a Gb-Rb group;

Ga represents an atom of oxygen or sulfur and Gb represents an atom of oxygen;

Ra represents a (C1-C6)alkyl or (C3-C7)cycloalkyl group;

Rb represents a (C1-C6)alkyl group substituted by at least a –COORc group, wherein Rc represents a hydrogen atom or (C1-C4)alkyl group; and

Y4 and Y5 independently represent a (C1-C4)alkyl group.

In a particular embodiment of the compound of formula (III):

Y1 represents a halogen atom or a Ra or Ga-Ra group;

A represents a CH2-CH2 group;

Y2 represents a Gb-Rb group;

Ga represents an atom of oxygen or sulfur and Gb represents an atom of oxygen;

Ra represents a (C1-C6)alkyl or (C3-C14)cycloalkyl group that is substituted by one or more halogen atoms;

Rb represents a (C1-C6)alkyl group substituted or not by one or more halogen atoms and substituted by at least a –COORc group, wherein Rc represents a hydrogen atom or a (C1-C4)alkyl group; and

Y4 and Y5 represent a (C1-C4)alkyl group.

5

In a particular embodiment of the compound of formula (III), Gb is an oxygen atom and Rb is (C1-C6)alkyl group substituted by a –COORc group, wherein Rc represents a hydrogen atom or an unsubstituted linear or branched (C1-C4)alkyl group.

10 In a particular embodiment of the compound of formula (III), Y1 is a (C1-C6)alkylthio group that comprises a (C1-C6)alkyl group that is linear or branched that is substituted or not by one or more halogen atoms.

In a particular embodiment, the compound of formula (III) is selected in the group
15 consisting of 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxyphenyl]prop-2-en-1-one (Elafibranor or GFT505), 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-isopropoxy carbonyldimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyldimethylmethoxyphenyl] prop-2-en-1-one, 1-[4-trifluoromethylphenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyl dimethylmethoxyphenyl]prop-
20 2-en-1-one, 1-[4-trifluoromethylphenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-trifluoromethyl oxyphenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyldimethylmethoxy phenyl] prop-2-en-1-one, 1-[4-trifluoromethoxyphenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxyphenyl]prop-2-en-1-one, 2-[2,6-dimethyl-4-[3-[4-(methylthio)phenyl]-3-oxo-propyl] phenoxy]-2-methylpropanoic
25 acid, and 2-[2,6-dimethyl-4-[3-[4-(methylthio) phenyl]-3-oxo-propyl]phenoxy]-2-methylpropanoic acid isopropyl ester.

In a more particular embodiment, the PPAR agonist is 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxy phenyl]prop-2-en-1-one (or Elafibranor – GFT505), or
30 a pharmaceutically acceptable salt thereof.

In a particular embodiment, the combination product of the invention is a composition comprising components (i) and (ii) as described above, and a pharmaceutically acceptable carrier.

35

In a particular embodiment, the combination product is a kit of parts comprising components (i) and (ii) as described above, for sequential, separate or simultaneous use.

5 In a further embodiment, components i) and ii) are formulated in an injectable suspension, a gel, an oil, a pill, a tablet, a suppository, a powder, a capsule, an aerosol, an ointment, a cream, a patch, or means of galenic forms for a prolonged and/or slow release.

In another embodiment, components (i) and (ii) of the combination product are for use in the treatment of any of the diseases mentioned below.

10

The present invention also relates to the combination product according to the invention, for use as a medicament.

15 The invention also relates to the combination product herein disclosed, for use in a method for the treatment of a disease. In another embodiment, the invention relates to a method for the treatment of a disease, comprising administering to a subject in need thereof a therapeutically effective amount of the combination product herein disclosed. In another embodiment, it is provided the use of a combination product according to the invention, for the manufacture of a medicament for the treatment of a disease.

20

In particular, the combination product of the present invention is useful for the treatment of diseases such as immune, inflammatory, metabolic, fibrotic and cholestatic diseases. In a particular embodiment, the disease is selected in the group consisting of metabolic liver diseases, non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH),
25 drug-induced liver diseases, alcohol-induced liver diseases, infectious agent induced liver diseases, inflammatory liver diseases, immune system dysfunction-mediated liver diseases, dyslipidemia, cardiovascular diseases, restenosis, syndrome X, metabolic syndrome, diabetes, obesity, hypertension, chronic cholangiopathies such as Primary Sclerosing Cholangitis (PSC), Primary Biliary Cholangitis (PBC), biliary atresia, familial intrahepatic
30 cholestasis type 3 (PFIC3), inflammatory bowel diseases, Crohn's disease, ulcerative colitis, keloid, old myocardial infarction, scleroderma/systemic sclerosis, inflammatory diseases, neurodegenerative diseases, cancers, liver cancer, hepatocellular carcinoma, gastrointestinal cancer, gastric cancer, meningioma associated with neurofibromatosis, pancreatic neuroendocrine tumors, pancreatic exocrine tumors, leukemia,

- myeloproliferative/myelodysplastic diseases, mastocytosis, dermatofibrosarcoma, solid tumors including breast, lung, thyroid or colorectal cancer, a prostate cancer, liver fibrosis or cirrhosis of any origin, metabolic disease-induced liver fibrosis or cirrhosis, NAFLD-induced fibrosis or cirrhosis, NASH-induced fibrosis or cirrhosis, alcohol-induced liver fibrosis or cirrhosis, drug-induced liver fibrosis or cirrhosis, infectious agent-induced liver fibrosis or cirrhosis, parasite infection-induced liver fibrosis or cirrhosis, bacterial infection-induced liver fibrosis or cirrhosis, viral infection-induced fibrosis or cirrhosis, HBV-infection induced liver fibrosis or cirrhosis, HCV-infection induced liver fibrosis or cirrhosis, HIV-infection induced liver fibrosis or cirrhosis, dual HCV and HIV-infection induced liver fibrosis or cirrhosis, radiation- or chemotherapy-induced fibrosis or cirrhosis, biliary tract fibrosis, liver fibrosis or cirrhosis due to any chronic cholestatic disease, gut fibrosis of any etiology, Crohn's disease-induced fibrosis, ulcerative colitis-induced fibrosis, intestine (e.g. small intestine) fibrosis, colon fibrosis, stomach fibrosis, skin fibrosis, epidermis fibrosis, endodermis fibrosis, skin fibrosis due to scleroderma/systemic sclerosis, lung fibrosis, lung fibrosis consecutive to chronic inflammatory airway diseases, such as COPD, asthma, emphysema, smoker's lung, tuberculosis, pulmonary fibrosis, idiopathic pulmonary fibrosis (IPF), heart fibrosis, kidney fibrosis, nephrogenic systemic fibrosis, muscle fibrosis, soft tissue (e.g. mediastinum or retroperitoneum) fibrosis, bone marrow fibrosis, joint fibrosis, tendon fibrosis, cartilage fibrosis, pancreas fibrosis, uterus fibrosis, nervous system fibrosis, testis fibrosis, ovary fibrosis, adrenal gland fibrosis, artery fibrosis, vein fibrosis, eye fibrosis, endomyocardial fibrosis, mediastinal fibrosis, myelofibrosis, retroperitoneal fibrosis, progressive massive fibrosis (a complication of coal workers' pneumoconiosis), proliferative fibrosis, neoplastic fibrosis, peri-implantational fibrosis and asbestosis, arthrofibrosis, adhesive capsulitis.
- In a most preferred embodiment, the disease is selected in the group consisting of metabolic liver diseases, non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), drug-induced liver diseases, alcohol-induced liver diseases, infectious agent induced liver diseases, inflammatory liver diseases, immune system dysfunction-mediated liver diseases, dyslipidemia, cardiovascular diseases, restenosis, syndrome X, metabolic syndrome, diabetes, obesity, hypertension, chronic cholangiopathies such as Primary Sclerosing Cholangitis (PSC), Primary Biliary Cholangitis (PBC), biliary atresia, familial intrahepatic cholestasis type 3 (PFIC3), inflammatory bowel diseases, Crohn's disease, ulcerative colitis, liver cancer, hepatocellular carcinoma, gastrointestinal cancer, gastric cancer, colorectal cancer, metabolic disease-induced liver fibrosis or cirrhosis, NAFLD-induced fibrosis or cirrhosis, NASH-induced fibrosis or cirrhosis, alcohol-induced liver fibrosis or cirrhosis, drug-induced liver fibrosis or cirrhosis, infectious agent-induced liver fibrosis or

cirrhosis, parasite infection-induced liver fibrosis or cirrhosis, bacterial infection-induced liver fibrosis or cirrhosis, viral infection-induced fibrosis or cirrhosis, HBV-infection induced liver fibrosis or cirrhosis, HCV-infection induced liver fibrosis or cirrhosis, HIV-infection induced liver fibrosis or cirrhosis, dual HCV and HIV-infection induced liver fibrosis or cirrhosis, radiation- or chemotherapy-induced fibrosis or cirrhosis, biliary tract fibrosis, liver fibrosis or cirrhosis due to any chronic cholestatic disease, gut fibrosis of any etiology, Crohn's disease-induced fibrosis, ulcerative colitis-induced fibrosis, intestine (e.g. small intestine) fibrosis, colon fibrosis, stomach fibrosis, lung fibrosis, lung fibrosis consecutive to chronic inflammatory airway diseases, such as COPD, asthma, emphysema, smoker's lung, tuberculosis, pulmonary fibrosis, idiopathic pulmonary fibrosis (IPF),

In a further aspect, the invention relates to the combination of the invention, for use in the inhibition of proliferation and/or activation of fibroblasts responsible for the production of collagen fibers and/or responsible for the production of the extracellular matrix.

According to the present invention, the term "autoimmune diseases" is used to designate a condition that arises from an abnormal immune response of the body against substances and tissues normally present in the body. The disease may be restricted to certain organs (e.g. in type I diabetes or autoimmune thyroiditis) or involve a particular tissue in different places (e.g. in Goodpasture's disease, affection of the basement membrane in the lung and the kidney).

The term "inflammation" is used to designate a condition that arise from a protective response involving host cells, blood vessels, and proteins and other mediators which may serve to eliminate the cause of cell/tissue injury, as well as the necrotic cells/tissues resulting from the original insult, and to initiate the process of repair. The inflammatory reaction may be manifested by pain, heat, redness, swelling, blood vessels dilatation, blood flow increase and loss of function.

According to the present invention, the terms "fibrosis", "fibrotic disease", "fibrotic disorder" and declinations thereof denote a pathological condition of excessive deposition of fibrous connective tissue in an organ or tissue. More specifically, fibrosis is a pathological process defined by a persistent fibrotic scar formation and overproduction of extracellular matrix by the connective tissue, as a response to tissue damage. Physiologically, the deposit of connective tissue can obliterate the architecture and function of the underlying organ or tissue.

According to the present invention, the fibrosis or fibrotic disorder may be associated with any organ or tissue fibrosis. Illustrative, non-limiting examples of particular organ fibrosis include liver, gut, kidney, skin, epidermis, endodermis, muscle, tendon, cartilage, heart, pancreas, lung, uterus, nervous system, testis, penis, ovary, adrenal gland, artery, vein,

colon, intestine (e.g. small intestine), biliary tract, soft tissue (e.g. mediastinum or retroperitoneum), bone marrow, joint or stomach fibrosis, in particular liver, kidney, skin, epidermis, endodermis, muscle, tendon, cartilage, heart, pancreas, lung, uterus, nervous system, testis, ovary, adrenal gland, artery, vein, colon, intestine (e.g. small intestine), biliary tract, soft tissue (e.g. mediastinum or retroperitoneum), bone marrow, joint, eye or stomach fibrosis.

According to the present invention, the terms “cholestasis” or “cholestatic disease”, or “cholestatic disorder” and declinations thereof denote a pathological condition defined by a decrease in bile flow due to impaired secretion by hepatocytes or to obstruction of bile flow through intra-or extrahepatic bile ducts. Therefore, the clinical definition of cholestasis is any condition in which substances normally excreted into bile are retained.

In a particular embodiment, the fibrotic disorder is selected in the group consisting of a liver, gut, lung, heart, kidney, muscle, skin, soft tissue (e.g. mediastinum or retroperitoneum), bone marrow, intestinal, and joint (e.g. knee, shoulder or other joints) fibrosis.

In a preferred embodiment, the fibrotic disorder is selected in the group consisting of liver, lung, skin, kidney and intestinal fibrosis.

In a more preferred embodiment of the present invention, treated fibrotic disorder is selected in the group consisting of the following non exhaustive list of fibrotic disorders: non-alcoholic steatohepatitis (NASH), pulmonary fibrosis, idiopathic pulmonary fibrosis, skin fibrosis, eye fibrosis (such as capsular fibrosis), endomyocardial fibrosis, mediastinal fibrosis, myelofibrosis, retroperitoneal fibrosis, progressive massive fibrosis (a complication of coal workers' pneumoconiosis), proliferative fibrosis, neoplastic fibrosis, lung fibrosis consecutive to chronic inflammatory airway disease (COPD, asthma, emphysema, smoker's lung, tuberculosis), alcohol or drug-induced liver fibrosis, liver cirrhosis, infection-induced liver fibrosis, radiation or chemotherapeutic-induced fibrosis, nephrogenic systemic fibrosis, Crohn's disease, ulcerative colitis, keloïd, old myocardial infarction, scleroderma/systemic sclerosis, arthrofibrosis, some forms of adhesive capsulitis, chronic fibrosing cholangiopathies such as Primary Sclerosing Cholangitis (PSC) and Primary Biliary Cholangitis (PBC), biliary atresia, familial intrahepatic cholestasis type 3 (PFIC3), peri-implantational fibrosis and asbestosis.

Cholestasis is defined as a decrease in bile flow due to impaired secretion by hepatocytes (hepato-cellular cholestasis) or to obstruction of bile flow through intra-or extrahepatic bile ducts (obstructive cholestasis). In clinical practice, cholestasis is any condition in which the flow of bile from the liver is slowed or blocked. According to a particular embodiment of the invention, the cholestestatic disease is selected in the group consisting of primary biliary cholangitis (PBC), primary sclerosing cholangitis (PSC),

Intrahepatic Cholestasis of Pregnancy, Progressive Familial Intrahepatic Cholestasis, Biliary atresia, Cholelithiasis, Infectious Cholangitis, Cholangitis associated with Langerhans cell histiocytosis, Alagille syndrome, Nonsyndromic ductal paucity, Drug-induced cholestasis, and Total parenteral nutrition-associated cholestasis. In a preferred embodiment, the cholestatic disease is PBC or PSC, in particular PBC.

Examples of inflammatory diseases, fibrotic diseases, metabolic diseases and cholestatic diseases include metabolic liver diseases, non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), drug-induced liver diseases, alcohol-induced liver diseases, infectious agent induced liver diseases, inflammatory liver diseases, immune system dysfunction-mediated liver diseases, dyslipidemia, cardiovascular diseases, restenosis, syndrome X, metabolic syndrome, diabetes, obesity, hypertension, chronic cholangiopathies such as Primary Sclerosing Cholangitis (PSC), Primary Biliary Cholangitis (PBC), biliary atresia, familial intrahepatic cholestasis type 3 (PFIC3), inflammatory bowel diseases, Crohn's disease, ulcerative colitis, keloid, old myocardial infarction, scleroderma/systemic sclerosis, inflammatory diseases, neurodegenerative diseases, cancers, liver cancer, hepatocellular carcinoma, gastrointestinal cancer, gastric cancer, meningioma associated with neurofibromatosis, pancreatic neuroendocrine tumors, pancreatic exocrine tumors, leukemia, myeloproliferative/myelodysplastic diseases, mastocytosis, dermatofibrosarcoma, solid tumors including breast, lung, thyroid or colorectal cancer, a prostate cancer, liver fibrosis or cirrhosis of any origin, metabolic disease-induced liver fibrosis or cirrhosis, NAFLD-induced fibrosis or cirrhosis, NASH-induced fibrosis or cirrhosis, alcohol-induced liver fibrosis or cirrhosis, drug-induced liver fibrosis or cirrhosis, infectious agent-induced liver fibrosis or cirrhosis, parasite infection-induced liver fibrosis or cirrhosis, bacterial infection-induced liver fibrosis or cirrhosis, viral infection-induced fibrosis or cirrhosis, HBV-infection induced liver fibrosis or cirrhosis, HCV-infection induced liver fibrosis or cirrhosis, HIV-infection induced liver fibrosis or cirrhosis, dual HCV and HIV-infection induced liver fibrosis or cirrhosis, radiation- or chemotherapy-induced fibrosis or cirrhosis, biliary tract fibrosis, liver fibrosis or cirrhosis due to any chronic cholestatic disease, gut fibrosis of any etiology, Crohn's disease-induced fibrosis, ulcerative colitis-induced fibrosis, intestine (e.g. small intestine) fibrosis, colon fibrosis, stomach fibrosis, skin fibrosis, epidermis fibrosis, endodermis fibrosis, skin fibrosis due to scleroderma/systemic sclerosis, lung fibrosis, lung fibrosis consecutive to chronic inflammatory airway diseases, such as COPD, asthma, emphysema, smoker's lung, tuberculosis, pulmonary fibrosis, idiopathic pulmonary fibrosis (IPF), heart fibrosis, kidney fibrosis, nephrogenic systemic fibrosis, muscle fibrosis, soft tissue (e.g. mediastinum or retroperitoneum) fibrosis, bone marrow fibrosis, joint

fibrosis, tendon fibrosis, cartilage fibrosis, pancreas fibrosis, uterus fibrosis, nervous system fibrosis, testis fibrosis, ovary fibrosis, adrenal gland fibrosis, artery fibrosis, vein fibrosis, eye fibrosis, endomyocardial fibrosis, mediastinal fibrosis, myelofibrosis, retroperitoneal fibrosis, progressive massive fibrosis (a complication of coal workers' pneumoconiosis), proliferative
5 fibrosis, neoplastic fibrosis, peri-implantational fibrosis and asbestosis, arthrofibrosis, adhesive capsulitis.

Preferably, the disease is selected in the group consisting of metabolic liver diseases, non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), drug-induced
10 liver diseases, alcohol-induced liver diseases, infectious agent induced liver diseases, inflammatory liver diseases, immune system dysfunction-mediated liver diseases, dyslipidemia, cardiovascular diseases, restenosis, syndrome X, metabolic syndrome, diabetes, obesity, hypertension, chronic cholangiopathies such as Primary Sclerosing Cholangitis (PSC), Primary Biliary Cholangitis (PBC), biliary atresia, familial intrahepatic
15 cholestasis type 3 (PFIC3), inflammatory bowel diseases, Crohn's disease, ulcerative colitis, liver cancer, hepatocellular carcinoma, gastrointestinal cancer, gastric cancer, colorectal cancer, metabolic disease-induced liver fibrosis or cirrhosis, NAFLD-induced fibrosis or cirrhosis, NASH-induced fibrosis or cirrhosis, alcohol-induced liver fibrosis or cirrhosis, drug-induced liver fibrosis or cirrhosis, infectious agent-induced liver fibrosis or cirrhosis, parasite
20 infection-induced liver fibrosis or cirrhosis, bacterial infection-induced liver fibrosis or cirrhosis, viral infection-induced fibrosis or cirrhosis, HBV-infection induced liver fibrosis or cirrhosis, HCV-infection induced liver fibrosis or cirrhosis, HIV-infection induced liver fibrosis or cirrhosis, dual HCV and HIV-infection induced liver fibrosis or cirrhosis, radiation- or chemotherapy-induced fibrosis or cirrhosis, biliary tract fibrosis, liver fibrosis or cirrhosis due
25 to any chronic cholestatic disease, gut fibrosis of any etiology, Crohn's disease-induced fibrosis, ulcerative colitis-induced fibrosis, intestine (e.g. small intestine) fibrosis, colon fibrosis, stomach fibrosis, lung fibrosis, lung fibrosis consecutive to chronic inflammatory airway diseases, such as COPD, asthma, emphysema, smoker's lung, tuberculosis, pulmonary fibrosis, idiopathic pulmonary fibrosis (IPF),

30 .

The term "treatment" or "treating" refers to the curative or preventive treatment of a disorder in a subject in need thereof. The treatment involves the administration of the compound, in particular comprised in a pharmaceutical composition, to a subject having a declared disorder, i.e. to a patient, to cure, delay, reverse, or slow down the progression of
35 the disorder, improving thereby the condition of the subject. The treatment may also result in an alleviation of symptoms associated with a disorder, or halt of further progression or

worsening of those symptoms. A treatment may also be administered to a subject that is healthy or at risk of developing a cholestatic or fibrotic disorder to prevent or delay the disorder.

Therefore, according to the invention, the treatment of a immune, inflammatory,
5 metabolic, fibrotic and cholestatic disease involves the administration of the combination of the present invention, for example in the form of a pharmaceutical composition containing components (i) and (ii) of the combination, to a subject having a declared disorder to cure, delay, reverse or slow down the progression of the disorder, thus improving the condition of the patient or to a healthy subject, in particular a subject who is at risk of developing such
10 disease.

The treatment involves the administration of the combination of the invention to a patient having a declared disorder to cure, delay, or slow down the progress, thus improving the condition of the patient or to a healthy subject, in particular a subject who is at risk of developing an inflammatory, metabolic, fibrotic and cholestatic disease.

15 The subject to be treated is a mammal, preferably a human. The subject to be treated according to the invention can be selected on the basis of several criteria associated to fibrotic diseases such as previous drug treatments, associated pathologies, genotype, exposure to risk factors, viral infection, as well as on the basis of the detection of any
20 relevant biomarker that can be evaluated by means of imaging methods and immunological, biochemical, enzymatic, chemical, or nucleic acid detection methods.

The subjects to be treated according to the invention can be selected on the basis of several criteria associated to inflammatory, metabolic, fibrotic and cholestatic diseases such
25 as previous drug treatments, associated pathologies, genotype, exposure to risk factors, viral infection, as well as any other relevant biomarker that can be evaluated by means of imaging methods and immunological, biochemical, enzymatic, chemical, or nucleic acid detection method.

Synthesis of NTZ or analogues can be for example carried out as described in
30 (Rossignol and Cavier 1975), or by any other way of synthesis known by a person skilled in the art.

In a particular embodiment, the treatment of an inflammatory, metabolic, fibrotic and cholestatic disease may comprise the administration of a composition comprising at least two compounds selected from NTZ and NTZ analogues. In this embodiment, the administered
35 component (ii) is provided in the same composition as the two compounds, or in a separate form, such as in a different composition.

In another embodiment, the combination of the invention is for simultaneous, sequential or separate administration in therapy, therefore being possibly included in different compositions. In case of sequential administration, NTZ and/or NTZ analogue(s) may be administrated prior to component (ii), or component (ii) is (are) administrated prior to NTZ and/or NTZ analogue(s).

NTZ and NTZ analogue(s) can be formulated as pharmaceutically acceptable salts, particularly acid or base salts compatible with pharmaceutical use. Salts of NTZ and NTZ analogue(s) include pharmaceutically acceptable acid addition salts, pharmaceutically acceptable base addition salts, pharmaceutically acceptable metal salts, ammonium and alkylated ammonium salts. These salts can be obtained during the final purification step of the compound or by incorporating the salt into the previously purified compound.

Combination of a compound of Formula (I) or (II) with one or more component(s) (ii) (for example a ROCK2 inhibitor, an ACC inhibitor, an ASK1 inhibitor, a dual NOX1 and NOX4 inhibitor, a Fatty Acid Bile Acid Conjugate, a FXR agonist, or a CCR antagonist) can be formulated as pharmaceutically acceptable non-toxic salts obtained from organic or inorganic bases or acids of compound of Formula (I) or (II) or component(s) (ii) (such as a ROCK2 inhibitor, an ACC inhibitor, an ASK1 inhibitor, a dual NOX1 and NOX4 inhibitor, a Fatty Acid Bile Acid Conjugate, a FXR agonist, or a CCR antagonist). These salts can be obtained during the final purification step of the compound or by incorporating the salt into the previously purified compound.

The pharmaceutical compositions of the present invention comprising a compound of Formula (I) or (II) and one or more component(s) (ii), such as a ROCK2 inhibitor, an ACC inhibitor, an ASK1 inhibitor, a dual NOX1 and NOX4 inhibitor, a Fatty Acid Bile Acid Conjugate, a FXR agonist, or a CCR antagonist, can also comprise one or several excipients or vehicles, acceptable within a pharmaceutical context (e.g. saline solutions, physiological solutions, isotonic solutions, etc., compatible with pharmaceutical usage and well-known by one of ordinary skill in the art).

These compositions can also comprise one or several agents or vehicles chosen among dispersants, solubilisers, stabilisers, preservatives, etc. Agents or vehicles useful for these formulations (liquid and/or injectable and/or solid) are particularly methylcellulose, hydroxymethylcellulose, carboxymethylcellulose, polysorbate 80, mannitol, gelatin, lactose, vegetable oils, acacia, liposomes, etc.

These compositions can be formulated in the form of injectable suspensions, gels, oils, ointments, pills, tablets, suppositories, powders, gel caps, capsules, aerosols, etc., eventually by means of galenic forms or devices assuring a prolonged and/or slow release.

For this kind of formulation, agents such as cellulose, carbonates or starches can be advantageously used.

The pharmaceutical compositions of the present invention comprising a compound of Formula (I) or (II) and one or more component(s) (ii), such as a ROCK2 inhibitor, an ACC inhibitor, an ASK1 inhibitor, a dual NOX1 and NOX4 inhibitor, a Fatty Acid Bile Acid Conjugate, a FXR agonist, or a CCR antagonist may be administered by different routes and in different forms. For example, the compound(s) may be administered via a systemic way, per os, parenterally, by inhalation, by nasal spray, by nasal instillation, or by injection, such as for example intravenously, by intra-muscular route, by subcutaneous route, by transdermal route, by topical route, by intra-arterial route, etc.

Of course, the route of administration will be adapted to the form of NTZ and/or NTZ analogue(s) in combination with one or more component(s) (ii), such as a ROCK2 inhibitor, an ACC inhibitor, an ASK1 inhibitor, a dual NOX1 and NOX4 inhibitor, a Fatty Acid Bile Acid Conjugate, a FXR agonist, or a CCR antagonist, according to procedures well known by those skilled in the art.

NTZ and/or NTZ analogue(s) in combination with one or more component(s) (ii), such as a ROCK2 inhibitor, an ACC inhibitor, an ASK1 inhibitor, a dual NOX1 and NOX4 inhibitor, a Fatty Acid Bile Acid Conjugate, a FXR agonist, or a CCR antagonist is administered in a therapeutically effective amount. Within the context of the invention, the term "effective amount" refers to an amount of the compound sufficient to produce the desired therapeutic result.

The frequency and/or dose relative to the administration can be adapted by one of ordinary skill in the art, in function of the patient, the pathology, the form of administration, etc. Typically, the combination (such as in the form of a pharmaceutical composition or a kit-of-parts) of the present invention can be administered for the treatment of a fibrotic disease at a dose for NTZ, or a NTZ analogue or component (ii), such as a ROCK2 inhibitor, an ACC inhibitor, an ASK1 inhibitor, a dual NOX1 and NOX4 inhibitor, a Fatty Acid Bile Acid Conjugate, a FXR agonist, or a CCR antagonist, such as a compound of Formula (I) or (II), comprised between 0.01 mg/day to 4000 mg/day, such as from 50 mg/day to 2000 mg/day, and particularly from 100 mg/day to 1000 mg/day.

In another preferred embodiment, the active ingredients are administered as one or more pharmaceutical composition(s) in the form of a pill intended for an oral ingestion.

Administration can be performed daily or even several times per day, if necessary.

The invention is further described with reference to the following, non-limiting, examples.

DESCRIPTION OF THE FIGURES AND TABLES

Figure 1: Antifibrotic effect of Elafibranor and Nitazoxanide in TGF β -induced hHSC

Serum-deprived HSC were preincubated for 1 hour with Elafibranor (A) or Nitazoxanide (B),
5 or Bezafibrate (C) before the activation with the profibrogenic cytokine TGF β 1 (1 ng/ml). After
48 hours of incubation, the expression of α -SMA was measured by ELISA. The obtained
values were transformed into percentage inhibition over TGF β 1 control. Data are presented
as mean (triplicates) $\pm$ standard deviation (SD). The curve fitting and the calculation of half
maximal inhibitory concentration (IC₅₀) were performed with XLFit software 5.3.1.3.

Figure 2: Combination of Elafibranor with Nitazoxanide synergistically inhibits α -SMA in
TGF β 1-induced hHSC

Combinations were tested in a dose-response matrix format and analyzed according to the
excess over Bliss (EOB) additivism model. Dilution series of Elafibranor (row) and
15 Nitazoxanide (column) were prepared, including their respective DMSO controls. The
resulting mixes were added to serum-deprived HSC, 1 hour prior to the activation with the
profibrogenic cytokine TGF β 1 (1 ng/ml). (A) Percentage of α -SMA inhibition over the TGF β 1
control for all combination pairs. Data are presented as mean of quadruplicates. (B) EOB
scores were calculated as described in Materials and Methods. Any compound pair with
20 positive EOB value were considered synergistic (colored from light grey to black). The total
EOB score including all combinations was also calculated. (C) Data values derived from a
synergistic combination pair were plotted in a bar graph representation. Data are presented
as mean (quadruplicates) $\pm$ standard deviation (SD). EOHSA model was used as described
in the Materials and Methods section to confirm the synergism of the selected NTZ/ELA
25 combination pairs.

Figure 3: Hepatic collagen content

6 week-old C57BL/6 mice were fed a control (CSAA) diet, CDAA + 1% CHOL (CDAAc) diet,
or CDAAc diet supplemented with NTZ (30 mg/kg/day or 100 mg/kg/day for 12 weeks) or
30 Elafibranor (1 mg/kg/day or 3 mg/kg/day) or a combination of Elafibranor and NTZ
(respectively 1+30 mg/kg/day, 1 +100 mg/kg/day, 3 + 30 mg/kg/day and 3 + 100 mg/kg/day) .
For each graph the exact amount of exposure doses was indicated.
After the sacrifice, the hepatic collagen content was determined.

Figure 4: Hepatic fibrosis percentage

6 week-old C57BL/6 mice were fed a control (CSAA) diet, CDAA + 1% CHOL (CDAAc) diet, or CDAAc diet supplemented with NTZ (30 mg/kg/day or 100 mg/kg/day for 12 weeks) or Elafibranor (1 mg/kg/day or 3 mg/kg/day) or a combination of Elafibranor and NTZ (respectively 1+30 mg/kg/day, 1 +100 mg/kg/day, 3 + 30 mg/kg/day and 3 + 100 mg/kg/day) .

- 5 For each graph the exact amount of exposure doses was indicated.
After the sacrifice, the hepatic fibrosis area was determined.

Figure 5: Hepatic α SMA gene expression

- 10 Figure 6: Hepatic CCR2 gene expression

Figure 7: Hepatic CCR5 gene expression

Figure 8: Hepatic Col1a2 gene expression

15

Figure 9: Hepatic MMP2 gene expression

Figure 10: Hepatic TIMP2 gene expression

- 20 Figure 11: Hepatic TGF β 1 gene expression

Figure 12: Combination of GS-0976 (A), Selonsertib (B), GKT-831 (C), KD-025 (D), Cenicriviroc (CVC) (E), Tropifexor (LJN-452) (F), and Aramchol (G) with Nitazoxanide synergistically inhibits α -SMA in TGF β 1-induced hHSC

- 25 Combinations were tested in a dose-response matrix format and analyzed according to the excess over Bliss (EOB) additivism model. Dilution series of GS-0976, Selonsertib, GKT-831, KD-025, CVC, Tropifexor (LJN-452), and Aramchol (row) and Nitazoxanide (column) were prepared, including their respective DMSO controls. The resulting mixes were added to serum-deprived HSC, 1 hour prior to the activation with the profibrogenic cytokine TGF β 1 (1
30 ng/ml). EOB scores were calculated as described in Materials and Methods. Any compound pair with positive EOB value were considered synergistic (colored from light grey to black). Data values derived from a synergistic combination pair were plotted in a bar graph representation. Data are presented as mean (quadruplicates) $\pm$ standard deviation (SD). EOHSA model was used as described in the Materials and Methods section to confirm the
35 synergism of the selected NTZ/ GS-0976 (A), Selonsertib (B), GKT-831 (C), KD-025 (D), Cenicriviroc (E), Tropifexor (F), and Aramchol (G) combination pairs.

Abbreviations used in the figures, in the tables, and in the text:

	AP-1	Activator Protein 1
	ASBTi	Apical Sodium-codependent Bile acid Transporter Inhibitor
5	ASK1	signal-regulating kinase 1
	AT1	AngioTensin 1
	COPD	Chronic Obstructive Pulmonary Disease
	CTGF	Connective Tissue Growth Factor
	DGAT	DiacylGlycerol-O-AcylTransferase
10	DMSO	DiMethylSulfOxyde
	DNA	DesoxyriboNucleic Acid
	DPP4	DiPeptidyl Peptidase
	ELISA	Enzyme-Linked Immuno Assay
	EOB	Excess Over Bliss
15	FABAC	Fatty Acid Bile Acid Conjugates
	FBS	Fetal Bovine Serum
	FGF	Fibroblast Growth Factor
	FXR	Farnesoid X Receptor
	GDF	Growth Differentiation Factor
20	GLP-1	Glucagon-Like Peptide-1
	GPCR	G-Protein Coupled Receptor
	HBV	Hepatitis B Virus
	HCV	Hepatitis C Virus
	15-HEPE	5-HydroxyEicosaPentaEnoic acid
25	HIV	Human Immunodeficiency Virus
	HSC	Hepatic Stellate Cell
	IC50	half maximal inhibitory concentration
	iNOS	inducible Nitric Oxide Synthase
	IPF	Idiopathic Pulmonary Fibrosis
30	LBD	Ligand Binding Domain
	LPS	LipoPolySaccharide
	LT	LeukoTriene
	MAPK	Mitogen-Activated Protein Kinases
	MMPase	Matrix MetalloProtease
35	NADPH	Nicotinamide Adenine Dinucleotide PHosphate
	NAFLD	Non-Alcoholic Fatty Liver Disease

	NASH	Non-Alcoholic SteatoHepatitis
	NF-κB	Nuclear Factor-kappa B
	NOX	NADPH oxidase
	NSAIDs	Non-Steroid Anti-Inflammatory Drugs
5	NTZ	NiTaZoxanide
	PAR	Protease-Activated Receptor
	PBC	Primary Biliary Cholangitis
	PDE	PhosphoDiEsterase
	PDGF	Platelet-Derived Growth Factor
10	PFIC3	Progressive Familial Intrahepatic Cholestasis type 3
	PFOR	Pyruvate:Ferredoxin OxidoReductase
	PPAR	Peroxisome proliferator activated receptor
	PPRE	PPAR Response Elements
	PSC	Primary Sclerosing Cholangitis
15	ROCK	Rho-associated protein kinase
	RTK	Receptor Tyrosine Kinase
	SD	Standard Deviation
	SGLT	Sodium-GLucose Transport
	STAT	Signal Transducers and Activators of Transcription
20	TGFβ	Transforming Growth Factor β
	TGFBRI	TGFβ Receptors type I
	TGFBRII	TGFβ Receptors type II
	THBS1	THromBoSpondin 1
	THR β	Thyroid Hormone Receptor β
25	TIMP	Tissue Inhibitor of MetalloProtease
	TLR-4	Toll Like Receptor 4
	TZ	TiZoxanide
	TZG	TiZoxanide Glucuronide
	VAP-1	Vascular Adhesion Protein-1
30		

EXAMPLES

MATERIALS AND METHODS

- 35 Compounds were dissolved in dimethyl sulfoxide (DMSO, Fluka cat# 41640). Lanifibranor (ARK PHARM cat# AK689102), Nitazoxanide (INTERCHIM cat#RQ550U), GKT-831

(cat#S7171, Absource Diagnostica GmbH), KD-025 (cat#S7936, Absource Diagnostica GmbH), GS-0976 (cat#HY-16901, Haoyuan Chemexpress Co), GS-4997 or Selonsertib (cat#S8292, Absource Diagnostica GmbH), Aramchol (cat#A765150, Toronto Research Chemical (TRC)), Tropifexor (LJN-452, cat# HY-107418 Haoyuan Chemexpress Co),
5 Cenicriviroc (PB0001630-239-01/PBLJ6483, Pharmablock), Nalmefene (cat#N284400, TRC), and PBI-4050 (cat#ACDS-072808, Acadechem) were obtained commercially.

hHSC culture

10 The human primary hepatic stellate cells (hHSC) (Innoprot) were cultured in STeCM medium (ScienCell cat# 5301) that was supplemented with 2% fetal bovine serum (FBS, ScienCell cat# 0010), 1% penicillin / streptomycin (ScienCell cat# 0503) and stellate cell growth supplement (SteCGS; ScienCell cat# 5352). Cell-culture flasks were coated with Poly-L
15 Lysine (Sigma cat# P4707) for a better adherence.

Preparation of compositions

2 components combination matrix (NTZ/component (ii))

For these experiments, a checkerboard matrix was generated. NTZ and selected component
(ii) stocks were serially diluted in DMSO in 5-points series in a row (component (ii)) and a 6-
20 points series in a column (NTZ) of a 96-well plate. Subsequently, the 5X6 combination matrix was generated by 1:1 mixing of all single agent concentrations. The test concentrations for each compound were chosen based on the respective IC₅₀ of each compound as single agent obtained by measuring α -SMA content in the HSC model stimulated with TGF- β 1.

Then, 2-fold and 4-fold higher and lower concentrations were selected.

Activation of hHSC with TGF- β 1 and compound treatment

The human primary hepatic stellate cells (hHSC) (Innoprot) were cultured under standard conditions, as described above. The cells were subsequently plated at a density of 2×10^4 cells/well into 96-well plates for the measure of α -SMA by ELISA. The next day, cell-
30 culture medium was removed, and cells were washed with PBS (Invitrogen cat# 14190). hHSC were deprived for 24 hours in serum-free and SteCGS-free medium. For the treatments with NTZ, Elafibranor, component (ii) and the respective NTZ/component (ii) or Elafibranor combinations, the serum-deprived hHSC were preincubated for 1 hour with the compounds followed by addition of the profibrogenic stimuli TGF- β 1 (PeproTech cat# 100-
35 21, 1 ng/mL) in serum-free and SteCGS-free medium for an additional 48 hour period.

α-SMA ELISA

The level of α-SMA was measured using a Sandwich ELISA. Briefly, the wells of an ELISA plate were first coated with the capture antibody (mouse monoclonal anti-ACTA2, Abnova) at 4°C overnight. After 3 washes in PBS + 0,2% Tween 20, a blocking solution consisting of PBS +0.2% BSA was added for one hour followed by another washing cycle. The cell lysates were transferred into the wells for binding to the capture antibody for a period of 2h at room temperature. After the washing procedure, the detection antibody (biotinylated mouse monoclonal anti-ACTA2, Abnova) was added for 2 hours at room temperature followed by 3 washes. For the detection, an HRP-conjugated Streptavidin (R&D Systems cat# DY998) was first applied for 30 min at room temperature. After washing, the HRP substrate TMB (BD,#555214) was added and incubated for 7min at room temperature in the dark. Upon oxidation, TMB forms a water-soluble blue reaction product that becomes yellow with addition of sulfuric acid (solution stop), enabling accurate measurement of the intensity at 450nm using a spectrophotometer. The developed color is directly proportional to the amount of α-SMA present in the lysate.

Determination of synergism by Excess Over Bliss (EOB) method and confirmation by EOHSA (Excess over Highest Single Agent)

The values obtained in the αSMA ELISA assays were first transformed into percentage inhibitions over TGF-β1 control. Then, using these percentage inhibitions, EOB (Excess Over Bliss) was determined to define the synergistic effects of drug combinations. Expected Bliss additivism score (E) was firstly determined by the equation:

$$E = (A + B) - (A \times B)$$
 where A and B are the percentage inhibition of NTZ (A) and component (ii) or Elafibranor (B) at a given dose. The difference between the Bliss expectation and the observed inhibition of the combined NTZ/component (ii) or Elafibranor at the same dose is the 'Excess over Bliss' score.

- Excess over Bliss score = 0 indicates that the combination treatment is additive (as expected for independent pathway effects);
- Excess over Bliss score >0 indicates activity greater than additive (synergy); and
- Excess over Bliss score <0 indicates the combination is less than additive (antagonism).

For the combination NTZ + component (ii) or Elafibranor, an additional total Bliss score was calculated by summation of all EOB.

EOHSA is a standard measure of synergy used by the FDA for evaluation of drug combinations and is calculated as the difference of the effect produced by the drug combination and the greatest effect produced by each of the combination's single agents at

the same concentrations as when combined (Borisý et al., 2003). For synergistic combinations identified by the EOB method, the experimental % inhibition were plotted in a bar graph and the significance of the observed differences between NTZ/component (ii) or Elafibranor and single agent were estimated by one-way ANOVA and uncorrected Fisher's LSD *post-hoc* (*: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$).

Evaluation of Elafibranor, Nitazoxanide and the combination Elafibranor + Nitazoxanide in a chronic CDAA + 1% cholesterol model of fibrosing NASH (12 weeks)

Experimental design

The choline-deficient and L-amino acid-defined (CDAA) diet lacks choline, which is essential for hepatic β -oxidation and very low density lipoprotein production, and is believed to induce hepatocytes to store fat and subsequently cause cellular damage. The CDAA diet-induced rodent model develops fibrosis within a relatively short period of time, ideal for rapidly studying the reversibility of NASH pathology, particularly fibrosis.

Increased cholesterol intake accelerates liver fibrosis in several mouse models of NASH. The exacerbation of liver fibrosis mainly involves free cholesterol accumulation in hepatic stellate cells, that sensitizes the cells to transforming growth factor β (TGF β) and subsequently aggravates liver fibrosis.

In the current study, we examined the effects of nitazoxanide on liver fibrosis in C57Bl/6J mice fed a CDAA diet supplemented with 1% cholesterol.

The preventive effects of alone, NTZ alone and the combination of both were assessed in a fibrosing NASH -model of mice fed a CDAA + 1% cholesterol diet. 6 week-old male C57Bl/6J mice were fed a control (CSAA) diet, CDAA + 1% cholesterol diet, or CDAA + 1% cholesterol diet supplemented with 1, and 3 mg/kg/day, NTZ 30 and 100 mg/kg/day or combined drugs (1 and 3 mg/kg/day combined to NTZ 30 and 100 mg/kg/day) for 12 weeks.

The body weight and the food intake were monitored twice per week. On the last day of treatment, mice were sacrificed after a 6h fasting period. The liver was rapidly excised for biochemical and histological studies.

All animal procedures were performed according to standard protocols and in accordance with the standard recommendations for the proper care and use of laboratory animals.

6 weeks-old C57BL/6 male mice were fed for 12 weeks according to an experimental plan detailed in the table 1:

Diet	Compound	Dose mg/kg/day	Number of mice
CSAA			8
			12
		1	8
		3	8
CDA + 1% Chol	NTZ	30	8
		100	8
	+ NTZ	1+30	8
		1+100	8
		3+30	8
		3+100	8

Table 1: Experimental plan

The control was the CSAA diet.

Some mice were fed with the CDAAc diet.

- 5 Some mice were fed with CDAAc diet supplemented with Elafibranor at 1 or 3 mg/kg/day.

Some mice were fed with CDAAc diet supplemented with NTZ at 30 or 100 mg/kg/day.

Some mice were fed with CDAAc diet supplemented with Elafibranor + NTZ combination at different ratios: 1 +30, 1 +100, 3 + 30 and 3 +100 mg/kg/day.

- 10 The group corresponding to CDAAc diet supplemented with NTZ at 30mg/kg is also named C57Bl/6J mice fed CDAA diet + nitazoxanide 0.02% (wt/wt) corresponding to the theoretical dose of 30 mg/kg/day.

The group corresponding to CDAAc diet supplemented with NTZ at 100mg/kg is also named C57Bl/6J mice fed CDAA diet + nitazoxanide 0.0667% (wt/wt) corresponding to the

- 15 theoretical dose of 100 mg/kg/day.

The food was purchased from Ssniff® company (Soest, Germany).

Nitazoxanide (cf reference in Table 1) was incorporated by Ssniff® into CDAA + 1% chol diet in powder form to the required dose.

The reference and batch number of nitazoxanide are summarized in the table Table 2:

- 20 Table 2: Nitazoxanide references

Compound (INN)	Laboratory code		Supplier	Reference	Batch
	External ID	Genfit ID			
Nitazoxanide	GFE 50455	GSL022597.08	Interchim	RQ550	1501

For each dose a calculation of the exact doses was done according to the following example. This allows taking in count the exact dose of each product that was exactly consumed by each group of mice.

Calculation of the actual treatment doses: example with nitazoxanide 0.02% (wt/wt)

- Food intake is expressed in grams of food / grams of animal / day
- 0.02% of nitazoxanide in diet
 - 0.02 g of cpd /100 g of food=0.2 mg of cpd/g of food
- Actual dose of cpd :
 - 0.2mg of cpd/gram of food/gram of animal/day
 - (0.2mg of cpd/gram of food/gram of animal/day)X1000= (0.2mg of cpd/gram of food/kg of animal/day)=200 mg of cpd/gram of food/kg of animal per day

Consequently, multiplying by 200 the food intake value expressed in grams of food / grams of animal / day; the obtained value corresponds to the actual administered dose expressed in mg of NTZ/kg of animal/day.

In the same manner:

- For the dose of NTZ 0.00667% wt/wt, the actual treatment dose was obtained by multiplying the food intake value (grams of food / grams of animal / day) by 66.7.
- For the dose of NTZ 0.0667% wt/wt, the actual treatment dose was obtained by multiplying the food intake value (grams of food / grams of animal / day) by 667.

According to the calculation the calculated dose versus estimated dose are given in the following tables 3 and 4.

Groups	GFT505		NTZ	
Estimated doses	1mpk	3mpk	30mpk	100mpk
Calculated doses	1mpk	2.8mpk	26.3mpk	78.1mpk

Table 3: estimated and calculated doses for each compound a different doses

Groups	Combo 1		Combo 2		Combo 3		Combo 4	
	GFT505	NTZ	GFT505	NTZ	GFT505	NTZ	GFT505	NTZ
Estimated doses	1mpk	30mpk	1mpk	100mpk	3mpk	30mpk	3mpk	100mpk
Calculated doses	0.9mpk	26.4mpk	0.9mpk	79.4mpk	2.9mpk	27.4mpk	2.8mpk	77.7mpk

Table 4: estimated and calculated doses for each combinations at different doses

The body weight and food intake were recorded twice a week throughout the study.

- 5 At the end of the treatment period, animals were anesthetized with isoflurane and blood samples were taken as described below. Animals were then sacrificed by cervical dislocation and beheaded for brain excision and weighing. The liver was also collected and weighed. Part of the liver was fixed in 4% formalin, embedded in paraffin and used for histological analyses. The remaining liver was snap frozen in liquid nitrogen and kept at -80°C until use
- 10 for further analyses.

Blood sampling was performed at sacrifice following a 6-hour fasting period. Blood samples were withdrawn under anesthesia by retro orbital puncture. Heparin tubes containing blood were rapidly centrifuged (15 minutes at 4,000 rpm/4°C) and the plasma fraction was collected. Plasma aliquots were stored at -20°C until further analyses.

15

Plasma biochemistry

Alanine Amino Transferase (ALT)

- The plasmatic concentration of ALT was determined using the appropriate Randox kit for Daytona automate (Randox, cat# AL 3801). Briefly, the ALT within the plasma sample
- 20 enzymatically transforms α -oxoglutarate and L-alanine into L-glutamate and pyruvate. In the presence of NADH, the generated pyruvate is converted by lactate dehydrogenase to form L-lactate and NAD⁺. The kinetics of the reaction is studied and allows the plasmatic level of ALT to be calculated. Results are expressed in U/L.

Aspartate Amino Transferase (AST)

The plasmatic concentration of AST was determined using the appropriate Randox kit for Daytona automate (Randox, cat# AS 3804). Briefly, the AST within the plasma sample enzymatically transforms α -oxoglutarate and L-aspartate into L-glutamate and oxaloacetate. In the presence of NADH, the generated oxaloacetate is converted by malate

dehydrogenase to form L-malate and NAD⁺. The kinetics of the reaction is studied and allows the plasmatic levels of AST to be calculated. Results are expressed in U/L.

Histology

- 5 At sacrifice, liver samples were processed for histological analysis and examined as follows.

Tissue embedding and sectioning

- The liver slices were first fixed for 12 hours in formalin 4% solution. Then, the liver pieces were washed 30 minutes in PBS, and dehydrated in ethanol solutions (successive baths at 10 70, 80, 95 and 100% ethanol). The liver pieces were incubated in three different baths of Xylene (Sigma-Aldrich cat# 534056), followed by two baths in liquid paraffin (60°C). Liver pieces were then put into racks that were gently filled with Histowax® to completely cover the tissue.

- The paraffin blocks containing the tissue pieces were removed from the racks and stored at 15 room temperature. The liver blocks were cut into 3 µm slices.

Picrosirius red staining

- Liver sections were deparaffinized, rehydrated and incubated for 15 minutes in a solution of Fast Green FCF 0.04% (Sigma-Aldrich, cat# F7258) before rinsing in a bath of 0.5% acetic acid (Sigma-Aldrich, cat# 695092). Then, the liver sections were rinsed in water and 20 incubated 30 minutes in a solution of Fast Green FCF 0.04% - 0.1% sirius red (Direct Red 80, Fluka cat# 43665) in saturated aqueous picric acid (Sigma-Aldrich cat# P6744). Sections were then dehydrated, and mounted using the CV Mount medium (Leica, cat #14046430011).

25

Histological examinations

- A technician blinded to the source of each liver specimen performed histological examinations. Virtual slides were generated using the Pannoramic 250 scanner from 3D Histech. Using Quant Center software (3D Histech, including Pattern Quant and Histo Quant 30 modules), collagen-stained areas were quantified. Briefly, Pattern Quant was used to detect the tissue and measure its surface. Then, Histo Quant was used to detect the stained collagen content and measure its surface, based on a color threshold method. The fibrosis area was then expressed as the percentage of the collagen surface to the whole tissue for each animal.

- 35 Hepatic fibrosis staging was blind-evaluated, using CRN fibrosis criteria.

Details on the parameters, quantification/counting, and number of fields considered are provided in the following table 5.

Table 5: CRN's criteria for fibrosis

Parameter	Points in score	Description (the entire section was considered)
Fibrosis	0	No fibrosis
	1	Centrilobular perisinusoidal/pericellular fibrosis or portal/periportal fibrosis
	2	Centrilobular perisinusoidal/pericellular fibrosis and portal/periportal fibrosis
	3	Centrilobular perisinusoidal/pericellular fibrosis and/or portal fibrosis with focal or extensive bridging fibrosis
	4	Cirrhosis

Statistical analysis

5 Experimental results were expressed as mean $\pm$ standard deviation (SD) and plotted as bar graphs or curves. Statistical analyses were performed as follows:

- For body weight and Food intake parameters:

On the one hand, CSAA diet and CDAA + 1% chol groups were compared by two-way ANOVA measures on factor Time followed by a Bonferroni's test, using Sigma Plot 11.0 software (\$: $p < 0.05$; \$\$: $p < 0.01$; \$\$\$ $p < 0.001$); on the other hand, treatment groups were compared to CDAA + 1% chol diet with the same statistical analyses and software (*: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$).

15 For the measures performed after sacrifice, CSAA vs CDAA + 1% chol groups were compared by a Student t-test using Sigma Plot 11.0 software (#: $p < 0.05$; ##: $p < 0.01$; ###: $p < 0.001$) and treatment groups were compared to CDAA + 1 % chol diet by one-way ANOVA followed by a Bonferroni's test, using Sigma Plot 11.0 software (*: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$).

Measurement of hepatic collagen content

20 The hepatic collagen content was determined using the appropriate QuickZyme kit (Total collagen assay, cat# QZB-totcol5). The assay is based on the detection of hydroxyproline, which is a non-proteinogenic amino acid mainly found in the triple helix of collagen. Thus,

hydroxyproline in tissue hydrolysates can be used as a direct measure of the amount of collagen present in the tissue (without discrimination between procollagen, mature collagen and collagen degradation products).

Complete hydrolysis of tissue samples in 6M HCl at 95°C is required before dosing the hydroxyproline. The assay results in the generation of a chromogen with a maximum absorbance at 570 nm. Results are expressed as mg of collagen / g of liver.

Procollagen III N-terminal Propeptide (PIIINP)

The plasmatic concentration of PIIINP was determined using an ELISA assay from Cloud-Clone Corp (cat# SEA573Ra), according to the manufacturer's instructions. The microtiter plate is pre-coated with an antibody specific to PIIINP. Standards or samples are added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to PIIINP. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain PIIINP, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of PIIINP in the samples is then determined by comparing the OD of the samples to the standard curve. Results are expressed in pg/mL.

Gene expression

RNA extraction

Hepatic Total RNA was isolated using Nucleospin® 96 Kit (Macherey Nagel) following manufacturer's instructions. 150ng of total RNA were reverse transcribed in cDNA using M-MLV-RT (Moloney Murine Leukemia Virus Reverse Transcriptase) (Invitrogen cat# 28025) in presence of RT buffer 1x (Invitrogen cat#P/NY02321), 1mM DTT (Invitrogen cat#P/NY00147), 0.5mM dNTPs (Promega), 200ng pdN6 (Roche cat#11034731001) and 40U of Ribonuclease inhibitor (Promega cat#N2515).

Quantitative PCR was then carried out using the CFX96 Touch™ Real-Time PCR Detection System (Biorad). Briefly, PCR reactions were performed in 96 well plates on 5µl of 5x diluted reverse transcription mix using the iQ SYBR Green Supermix kit (Biorad cat#170887). The experimental conditions were: 20 µL of volume reaction and 0.5µL each of reverse and forward primers (10pMol).

Primer name	Sequence ID	Sequence (5'→3')
αSMA forward	1	CTGACAGAGGCACCACTGAA
αSMA reverse	2	CATCTCCAGAGTCCAGCACA
Col1α1 forward	3	AGGCGAACAAGGTGACAGAG
Col1α1 reverse	4	GCCAGGAGAACCAGCAGAG
Col1α2 forward	5	ATTGGAAGCCGAGGTCCCAG
Col1α2 reverse	6	TTTGCCCCCAGGTATGCCAG
TGFβ1 forward	7	TTGCTTCAGCTCCACAGAGA
TGFβ1 reverse	8	TGGTTGTAGAGGGCAAGGAC
TIMP1 forward	9	ATTCAAGGCTGTGGGAAATG
TIMP1 reverse	10	CTCAGAGTACGCCAGGGAAC
TIMP2 forward	11	GCATCACCCAGAAGAAGAGC
TIMP2 reverse	12	GGGTCCTCGATGTCAAGAAA
MMP2 forward	13	TCCCTAAGCTCATCGCAGAC
MMP2 reverse	14	GCTTCCAACTTCACGCTCT
MMP7 forward	15	TAATTGGCTTCGCAAGGAGA
MMP7 reverse	16	AAGGCATGACCTAGAGTGTTCC
CCR2 forward	17	TAATATGTTACCTCAGTTCATCCACGG
CCR2 reverse	18	TGCTCTTCAGCTTTTTACAGCCTATC
CCR5 forward	19	ATTCTCCACACCCTGTTTCG
CCR5 reverse	20	GAATTCCTGGAAGGTGGTCA
GAPDH forward	21	TATGACTCCACTCACGGCAA
GAPDH reverse	22	TCCACGACATACTCAGCACC

Expression levels were normalized using the expression of GAPDH gene as reference.

For each gene, the standard curve was drawn by selecting the best points (at least three points) in order to have PCR reaction efficiency close to 100% and a correlation coefficient close to 1. Expression levels were determined using the standard curve equation for both the housekeeping gene and the target gene (taking into account the specific PCR efficiency of each target gene).

Results and conclusions:

The abnormal persistence of differentiated myofibroblasts is a characteristic of many fibrotic diseases.

Following liver injury, quiescent HSCs undergo a process of activation that is characterized by a differentiation into (α -SMA)-positive myofibroblasts.

NTZ as a single agent was shown to confer an anti-fibrotic activity in TGF β -induced hHSC (Fig.1B). Since it is known that NTZ is rapidly hydrolyzed into its active metabolite tizoxanide (TZ) (Broekhuysen, Stockis et al. 2000), this metabolite was also evaluated for its antifibrotic activity in HSC. TZ showed a profile similar to the parent drug (data not shown). On the other hand, some PPAR agonists like Elafibranor also revealed an antifibrotic profile in the TGF β induced HSC model (Fig. 1A). Other PPAR agonists like bezafibrate revealed a weak activity suggesting that PPAR agonists are not equivalent regarding their antifibrotic properties (Fig. 1C).

In order to evaluate if a combination of Elafibranor with NTZ could reduce fibrosis in a synergistic manner, combination matrix experiments were performed in TGF β -induced HSCs. Briefly, NTZ and Elafibranor solutions were serially diluted in a checkerboard format generating a 42 combinations matrix covering a large panel of Elafibranor/NTZ ratios. Synergy was first determined by calculating Excess Over Bliss scores. These experiments revealed that NTZ could synergize with Elafibranor to reduce α -SMA production in activated HSCs. Several combination pairs revealed an EOB score over 10, which is indicative of a synergism (Fig.2B). To validate the synergism, the experimental values corresponding to top EOB score were plotted in a bar graph (Fig. 2C). These graphs illustrate that the combination of NTZ with Elafibranor shows a superior antifibrotic effect that is statistically significant compared to the highest single agent (NTZ or Elafibranor). The most impressive example is represented with the combination pair NTZ at 0,6 μ M and Elafibranor at 5 μ M. Although NTZ shows almost no antifibrotic activity at 0,6 μ M, the addition of Elafibranor at 5 μ M results in a strong reduction of α SMA of 55%, which is much stronger than the effect observed with the single agents alone. In conclusion, the applicant has discovered unexpected antifibrotic activities for a combination of a compound of Formula (I) with specific PPAR agonist(s). These results suggest that a combination of a compound of Formula (I) with a PPAR agonist can be synergistic and can provide therapeutic benefits in multiple types of fibrotic diseases.

The administration of a choline-deficient and L-amino acid-deficient (CDAA) + 1 % cholesterol diet to mice causes progressive fibrosing steatohepatitis that is pathologically similar to human non-alcoholic steatohepatitis (NASH).

The CDAA +1 % cholesterol diet notably induces a significant increase in the hepatic collagen, as shown in the figure 3.

This figure also shows that administration of Elafibranor or NTZ alone decreases the hepatic collagen content. The decrease in collagen is proportional to the dose of elafibranor or NTZ administered.

When a combination of Elafibranor and NTZ is administered, the decrease in collagen produced is greater than the decrease observed for each compound taken separately.

There is therefore a synergistic effect of the combination of elafibranor and NTZ on the decrease in collagen production. In other words, there is a noticed anti-fibrotic effect when Elafibranor and NTZ are combined.

The better effect is observed for the combination comprising Elafibranor at 2.8 mpk (mg per kilogram) and NTZ at 77,7 mpK, expressed in calculated doses.

The figure 4 represents the results obtained with the histology, i.e. the determination of the fibrosis area which was expressed as the percentage of the collagen surface to the whole tissue for each animal.

The CDAA +1 % cholesterol diet notably induces a significant increase in the fibrosis percentage.

The figure 4 also shows that administration of Elafibranor or NTZ alone decreases the fibrosis percentage. The decrease is proportional to the dose of elafibranor or NTZ administered.

When a combination of Elafibranor and NTZ is administered, the decrease in fibrosis percentage is greater than the decrease observed for each compound taken separately.

There is therefore a synergistic effect of the combination of elafibranor and NTZ on the decrease in the fibrosis percent. In other words, there is a noticed anti-fibrotic effect when Elafibranor and NTZ are combined.

The better effect is observed for the combination comprising Elafibranor at 2.8 mpk (mg per kilogram) and NTZ at 77,7 mpK, expressed in calculated doses.

The figures 5 to 11 represent the gene expression of different hepatic markers of fibrosis. For all markers, the CDAA +1 % cholesterol diet notably induces a significant increase in the gene expressions.

It can also be noted that that administration of Elafibranor or NTZ alone decreases the level of the different gene expression. The decrease is proportional to the dose of elafibranor or NTZ administered.

When a combination of Elafibranor and NTZ is administered, the decrease in the gene expression is greater than the decrease observed for each compound taken separately.

There is therefore a synergistic effect of the combination of elafibranor and NTZ on the decrease in the gene expression of different hepatic markers of fibrosis. In other words, there is a noticed anti-fibrotic effect when Elafibranor and NTZ are combined.

- 5 To assess the synergism with the combination of NTZ with component (ii), GS-0976, Selonsertib (GS-4997), GKT-831, KD-025, Cenicriviroc (CVC), Tropifexor (LJN-452), and Aramchol were validated on anti-fibrotic activity. The combination of NTZ with GS-0976, Selonsertib (GS-4997), GKT-831, KD-025, Cenicriviroc, Tropifexor, and Aramchol revealed an EOB score > 0, which is indicative of a synergism.
- 10 The combination of GS-0976 at 10 μ M and NTZ at 1.25 μ M resulted also in a significant and strong reduction of α SMA of 64%, which was much stronger than the effect observed with the single agents alone (Fig. 12A). The combination of Selonsertib (GS-4997) at 10 μ M and NTZ at 2.5 μ M also resulted in a significant and strong reduction of α SMA of 77%, which was much stronger than the effect observed with the single agents alone (Fig. 12B). The
- 15 synergism was even more pronounced with the combination of GKT-831 0.3 μ M and NTZ at 0.3 μ M (Fig. 12C) and the combination of KD-025 at 1 μ M and NTZ at 0.3 μ M (Fig. 12D). The combination of Cenicriviroc at 3 μ M and NTZ at 2.5 μ M resulted also in a significant and strong reduction of α SMA of 47%, which was much stronger than the effect observed with the single agents alone (Fig. 12E). The synergism was to the same extent with the
- 20 combination of Tropifexor 3 μ M and NTZ at 1.3 μ M (Fig. 12F). The combination with Aramchol at 3 μ M and NTZ at 0.625 μ M reached a reduction of α SMA of 52%.

- By contrast, no synergism could be observed with the combination of NTZ with Nalmefene (JKB-121), originally known as nalmetrene, an opioid antagonist and TLR-4 antagonist used
- 25 primarily in the management of alcohol dependence, or PBI-4050, a potential treatment for idiopathic pulmonary fibrosis, acute lung injury, cystic fibrosis, diabetic.

- In conclusion, the applicant has discovered unexpected antifibrotic activities for a combination of a compound of Formula (I) with specific ROCK2 inhibitor(s), ACC inhibitor(s),
- 30 dual NOX1 and NOX4 inhibitors, ASK1 inhibitor(s), Fatty Acid Bile Acid conjugates (FABAC), FXR agonists, and CCR antagonists.

- These results suggest that a combination of a compound of Formula (I) or Formula (II) with a ROCK2 inhibitor, an ACC inhibitor, a dual NOX1 and NOX4 inhibitor, an ASK1 inhibitor, a Fatty Acid Bile Acid Conjugate, an FXR agonist, and a CCR antagonist can be synergistic
- 35 and/or can have additional effect and can also provide therapeutic benefits in multiple types of fibrotic diseases.

REFERENCES

- 5 Broekhuysen, J., A. Stockis, et al. (2000). "Nitazoxanide: pharmacokinetics and metabolism in man." Int J Clin Pharmacol Ther **38**(8): 387-394.
- de Carvalho, L. P. S., C. M. Darby, et al. (2011). "Nitazoxanide disrupts membrane potential and intrabacterial pH homeostasis of Mycobacterium tuberculosis." ACS Med. Chem. Lett. **2**(Copyright (C) 2015 American Chemical Society (ACS). All Rights Reserved.): 849-854.
- 10 Di Santo, N. and J. Ehrisman (2014). "A functional perspective of nitazoxanide as a potential anticancer drug." Mutat. Res., Fundam. Mol. Mech. Mutagen. **768**(Copyright (C) 2015 American Chemical Society (ACS). All Rights Reserved.): 16-21.
- 15 Dubreuil, L., I. Houcke, et al. (1996). "In vitro evaluation of activities of nitazoxanide and tizoxanide against anaerobes and aerobic organisms." Antimicrob. Agents Chemother. **40**(Copyright (C) 2015 American Chemical Society (ACS). All Rights Reserved.): 2266-2270.
- 20 Finegold, S. M., D. Molitoris, et al. (2009). "Study of the in vitro activities of rifaximin and comparator agents against 536 anaerobic intestinal bacteria from the perspective of potential utility in pathology involving bowel flora." Antimicrob. Agents Chemother. **53**(Copyright (C) 2015 American Chemical Society (ACS). All Rights Reserved.): 281-286.
- Fox, L. M. and L. D. Saravolatz (2005). "Nitazoxanide: a new thiazolide antiparasitic agent." Clin. Infect. Dis. **40**(Copyright (C) 2015 American Chemical Society (ACS). All Rights Reserved.): 1173-1180.
- 25 Hemphill, A., J. Mueller, et al. (2006). "Nitazoxanide, a broad-spectrum thiazolide anti-infective agent for the treatment of gastrointestinal infections." Expert Opin. Pharmacother. **7**(Copyright (C) 2015 American Chemical Society (ACS). All Rights Reserved.): 953-964.
- 30 Hoffman, P. S., G. Sisson, et al. (2007). "Antiparasitic drug nitazoxanide inhibits the pyruvate oxidoreductases of Helicobacter pylori, selected anaerobic bacteria and parasites, and Campylobacter jejuni." Antimicrob. Agents Chemother. **51**(Copyright (C) 2015 American Chemical Society (ACS). All Rights Reserved.): 868-876.
- 35 Megraud, F., A. Occhialini, et al. (1998). "Nitazoxanide, a potential drug for eradication of Helicobacter pylori with no cross-resistance to metronidazole." Antimicrob. Agents Chemother. **42**(Copyright (C) 2015 American Chemical Society (ACS). All Rights Reserved.): 2836-2840.
- 40 Pankuch, G. A. and P. C. Appelbaum (2006). "Activities of tizoxanide and nitazoxanide compared to those of five other thiazolides and three other agents against anaerobic species." Antimicrob. Agents Chemother. **50**(Copyright (C) 2015 American Chemical Society (ACS). All Rights Reserved.): 1112-1117.
- Rossignol, J.-F. (2014). "Nitazoxanide: A first-in-class broad-spectrum antiviral agent." Antiviral Res. **110**(Copyright (C) 2015 American Chemical Society (ACS). All Rights Reserved.): 94-103.

Rossignol, J. F. and R. Cavier (1975). 2-Benzamido-5-nitrothiazoles, S.P.R.L. Phavic, Belg. . 11 pp.

Rossignol, J. F. and H. Maisonneuve (1984). "Nitazoxanide in the treatment of *Taenia saginata* and *Hymenolepis nana* infections." Am J Trop Med Hyg **33**(Copyright (C) 2015 U.S. National Library of Medicine.): 511-512.