

PERFUSION AND/OR PRESERVATION SOLUTION FOR ORGANS**FIELD OF THE INVENTION**

The present invention relates to a solution for preservation, perfusion, and/or
5 reperfusion of an organ, especially the heart, for transplantation. The solution
contains peptide inhibitor(s) of protein kinase C beta II (PKC β II) and/or of protein
kinase C zeta (PKC ζ) and/or peptide activator(s) of protein kinase C delta (PKC δ).

BACKGROUND OF THE INVENTION

10 Successful organ transplantation is often limited due to ischemic/reperfusion
injury. Isolated human hearts deprived of oxygen for more than four hours
progressively lose vigor and often do not survive in recipient hosts. Other organs
such as the kidney, liver, pancreas and lung are also subject to tissue and cellular
damage when removed from their hosts prior to transplantation. This damage is due
15 to hypoxic conditions and a lack of circulation, which normally delivers physiological
concentrations of oxygen and nutrients, and removes toxic compounds produced by
an organ's cells. Organ transplants have a higher frequency of success when
performed immediately after excision from their hosts.

Recent advances have increased the rate of successful organ transplants and
20 organ surgery, such as coronary bypass surgery. The first includes organ preservation
and organ perfusion solutions. The second is improved methods and devices for the
delivery of organ perfusion solutions to an organ.

Short-term myocardial preservation is currently provided by cold storage after
cardioplegic arrest. A variety of processes exist however differing by the composition
25 of the solution used, the preservation temperature and the administration protocol.

Different solutions for arresting and preserving the heart have been developed to protect the myocardium in cardiac surgery. Examples of these solutions include Krebs-Henseleit solution, UW solution, St. Thomas II solution, Collins solution and Stanford solution. (See, for example, U.S. Patent Nos. 4,798,824 and 4,938,961; Southard and Belzer, *Ann. Rev. Med.* 46:235-247 (1995); and Donnelly and Djuric, *Am. J. Hosp. Pharm.* 48:2444-2460 (1991)). Nevertheless, organ rejections still remains due to deterioration in the condition of the transplanted organ between the time of removal and the restoration of blood flow in the recipient.

Restoration of blood flow is the primary objective for treatment of organ tissue experiencing prolonged ischemia, e.g., during transplant. However, reperfusion of blood flow induces endothelium and myocyte injury, resulting in organ dysfunction (Buerke et al., *Am J Physiol* 266: H128-136, 1994; Lucchesi and Mullane, *Ann Rev Pharmacol Toxicol* 26: 2011-2024, 1986; and Lucchesi et al., *J Mol Cell Cardiol* 21: 1241-1251, 1989). The sequential events associated with reperfusion injury are initiated by endothelial dysfunction which is characterized by a reduction of the basal endothelial cell release of nitric oxide (NO) within the first 2.5-5 min post-reperfusion (Tsao and Lefer, *Am J Physiol* 259: H1660-1666, 1990). The decrease in endothelial derived NO is associated with adhesion molecule up-regulation on endothelial and polymorphonuclear (PMN) leukocyte cell membranes (Ma et al., *Circ Res* 72: 403-412, 1993; and Weyrich et al., *J Leuko Biol* 57: 45-55, 1995). This event promotes PMN/endothelial interaction, which occurs by 10 to 20 min post-reperfusion, and subsequent PMN infiltration into the myocardium is observed by 30 min post reperfusion (Lefer and Hayward, In *The Role of Nitric Oxide in Ischemia-Reperfusion: Contemporary Cardiology*, Loscalzo et al. (Eds.), Humana Press, Totowa, NJ, pp. 357-380, 2000; Lefer and Lefer, *Cardiovasc Res* 32: 743-751, 1996;

Tsao et al., *Circulation* 82: 1402-1412, 1990; and Weyrich et al., *J Leuko Biol* 57: 45-55, 1995).

Chemotactic substances released from reperfused tissue and plasma factors activate PMNs that augment PMN release of cytotoxic substances (i.e. superoxide anion) and contribute to organ dysfunction following ischemia/reperfusion (Lucchesi et al. *J Mol Cell Cardiol* 21: 1241-1251, 1989; Ma et al., *Circ Res* 69: 95-106, 1991; Tsao et al., *Circulation* 82: 1402-1412, 1990; and Tsao et al., *Am Heart J* 123: 1464-1471, 1992). Superoxide combines with NO to produce peroxynitrite anion thus reducing the bioavailability of NO and promotes endothelial dysfunction and PMN infiltration after myocardial ischemia/reperfusion (Clancey et al., *J Clin Invest* 90: 1116-1121, 1992; Hansen, *Circulation* 91: 1872-85, 1995; Lucchesi et al., *J Mol Cell Cardiol* 21: 1241-1251, 1989; Rubanyi and Vanhoutte, *Am J Physiol* 250: H815-821, 1986; Tsao et al., *Am Heart J* 123: 1464-1471, 1992; and Weiss, *New Eng J Med* 320: 365-375, 1989).

Therefore, there remains a need for a solution of improved quality that can extend the preservation time of an organ for transplantation and protect the organ from reperfusion injury after ischemia, so that the organ can resume proper function after restoration of blood flow.

SUMMARY OF THE INVENTION

In one aspect, there is provided a solution for preservation, perfusion, and/or reperfusion of an organ, especially the heart, containing peptide inhibitor(s) of protein kinase C β II (PKC β II) and/or of protein kinase C ζ (PKC ζ) and/or peptide activator(s) of protein kinase C δ (PKC δ). The solution protects organ tissues and cells from damage while the organ is isolated from the circulatory system or is experiencing decreased blood flow (ischemia). The present inventor has discovered that the peptide inhibitors of PKC β II and/or PKC ζ , and/or peptide activator(s) of PKC δ enhance NO release or inhibit endothelial/PMN superoxide release, which can exert protective effect in organs undergoing ischemia/reperfusion.

In another aspect, there is provided a method for preserving an organ for transplantation comprising the step of perfusing the organ with the solution of the first aspect.

In still another aspect, there is provided a method for protecting an ischemic organ from damage comprising the step of perfusing the organ with the solution of the first aspect.

In yet another aspect, there is provided a method for attenuating organ dysfunction after ischemia comprising the step of perfusing the organ with the solution of the first aspect.

In a further aspect, there is provided a method for maintaining nitric oxide release in an ischemic organ comprising the step of perfusing the organ with the solution of the first aspect.

In another aspect, there is provided a method for protecting an organ from damage after isolation from the circulatory system comprising the step of perfusing the organ with the solution of the first aspect.

In an embodiment, the solution contains about 5-10 μM of the peptide inhibitor of PKC βII and/or about 2.5-5 μM of the peptide inhibitor of PKC ζ and/or a 5-10 μM of the peptide activator of PKC δ dissolved in a saline solution.

The solution of the present disclosure can be used as a perfusion solution or a preservation solution. As a perfusion solution, it is pumped into the vasculature of the organ to protect the organ tissues and cells. As a preservation solution, it serves as a bathing solution into which the organ is submerged. Preferably, the organ is perfused with and submerged in the present solution. Further, the present solution also serves as a reperfusion solution upon restoration of blood flow to the organ after ischemia.

Also disclosed herein are methods of using the solution of the present disclosure. These include methods for preserving an organ for transplantation, for protecting an ischemic organ from damage, for attenuating organ dysfunction after ischemia, for maintaining nitric oxide release in an ischemic organ, and for protecting an organ from damage when isolated from the circulatory system.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Time course of LVDP (left ventricular developed pressure = left ventricular end systolic pressure – left ventricular end diastolic pressure) in sham, I/R, I/R+PMNs and I/R+PMN+ PKC ζ peptide inhibitor (5 μ M) perfused rat hearts. LVDP data at initial (baseline) and reperfusion from 0 to 45 min following 20 min ischemia. The sham group (n=6) maintained the same LVDP throughout the 80 min. protocol. The I/R (n=6) group recovered to initial baseline values. I/R+PMN group (n=6) exhibited a significant and sustained reduction in LVDP compared to and I/R+PMN+ PKC ζ peptide inhibitor (n=6) group. All values are expressed as mean $\pm$ SEM.

*p<0.05 and **p<0.01, I/R+PMN+ PKC ζ peptide inhibitor group from I/R+PMNs.

Figure 2. Initial and final LVDP expressed in mmHg from isolated perfused rat hearts before ischemia (I) (initial) and after 45 min post reperfusion (R) (final). Hearts were perfused in the presence or absence of PMNs. PMNs induced a significant contractile dysfunction, which was attenuated by the PKC ζ peptide inhibitor, but was significantly blocked by N^G-nitro-L-arginine methyl ester (L-NAME). All values are expressed as mean $\pm$ SEM. Numbers of hearts examined are at the bottom of the bars. **p<0.01, from final I/R+PMNs; NS=not significant.

Figure 3. Initial and final maximal rate of LVDP (+dP/dt max) expressed in mmHg/s in isolated perfused rat hearts before ischemia (I) and after reperfusion (R). Hearts were perfused in the presence or absence of PMNs. PMNs induced a significant contractile dysfunction, which was attenuated by a PKC ζ peptide inhibitor, but was blocked by L-NAME. All values are expressed as means $\pm$ SEM. Numbers of hearts examined are at the bottom of the bars. **P<0.01, from final I/R + PMNs; NS=not significant.

Figure 4a. Histological assessment of total intravascular and infiltrated PMNs in

isolated perfused rat heart samples taken from 3 rats per group and 10 areas per heart. The numbers of total intravascular and infiltrated PMNs in

- 5 post-reperfusion cardiac tissue and adhering to coronary vasculature was significantly attenuated by the PKC ζ peptide inhibitor. Hatched boxes represent non-PMN perfused hearts and black boxes represent PMN-perfused hearts. **P<0.01, from I/R+PMNs.

Figure 4b. Histological assessment of intravascular PMNs that adhered to the
10 coronary vasculature in isolated perfused rat heart samples taken from 3 rats per group

and 10 areas per heart. The numbers of PMNs adhering to the coronary vasculature was not significantly different from I/R+PMNs. Hatched boxes represent non-PMN perfused hearts and black boxes represent PMN-perfused hearts. All values are

- 15 mean numbers of PMNs/mm² of heart area $\pm$ SEM.

Figure 5. Measurement of NO release from rat aortic segments. Endothelial NO release was significantly increased from basal NO release in PKC ζ peptide inhibitor treated segments (2.5-15 μ M), as well as acetylcholine (Ach, 200nM). NO release was significantly reduced in both groups given 400 μ M L-NAME, and in
20 endothelium removed (denuded) segments. All values are expressed as means $\pm$ SEM. Numbers at bottom of bars are numbers of separate experiments per group. *p<0.05, **p<0.01, from basal values.

Figure 6. Superoxide release from rat PMNs. Superoxide release was measured from 5 X 10⁶ PMNs after phorbol-12-myristate-13-acetate (PMA) (15 nM)
25 stimulation. Superoxide dismutase (SOD) (10 μ g/ml) was employed as a positive

control. The change in absorbance (Δ) was measured 360 sec after PMA addition (peak response). Superoxide release was significantly inhibited by the PKC ζ peptide inhibitor (** $p < 0.01$, 2.5, 5 and 15 μ M). All values are means $\pm$ SEM. Numbers at bottom of bars show the numbers of separate experiments per group.

5 Figure 7. Time course of LVDP in Sham I/R, I/R, I/R+PMNs and I/R+PMN+ β II peptide inhibitor (10 μ M) perfused rat hearts. LVDP data at initial (baseline) and reperfusion from 0 to 45 min following 20 min ischemia. The sham group (n=6) maintained the same LVDP throughout the 80 min. protocol. The I/R+PMN group (n=9) exhibited a significant and sustained reduction in LVDP
10 compared to I/R (n=6) and I/R+PMN+ β II peptide inhibitor (n=7) groups. All values are expressed as mean $\pm$ SEM. * $p < 0.05$ and ** $p < 0.01$, from I/R+PMNs.

Figure 8. Initial and final LVDP expressed in mmHg from isolated perfused rat hearts before ischemia (I) (initial) and after 45 min post reperfusion (R) (final). Hearts were perfused in the presence or absence of PMNs. PMNs induced a
15 significant contractile dysfunction, which was attenuated by PKC β II peptide inhibitor, but was significantly blocked by the presence of L-NAME. All values are expressed as mean $\pm$ SEM. Numbers of hearts examined are at the bottom of the bars. * $p < 0.05$ and ** $p < 0.01$, from final I/R+PMNs; NS=not significant.

Figure 9. Initial and final +dP/dt max expressed in mmHg/s in isolated
20 perfused rat hearts before ischemia (I) and after reperfusion (R). Hearts were perfused in the presence or absence of PMNs. PMNs induced a significant contractile dysfunction, which was attenuated by PKC β II peptide inhibitor, but was blocked by L-NAME. All values are expressed as means $\pm$ SEM. Numbers of hearts examined are at the bottom of the bars. * $p < 0.05$ and ** $p < 0.01$, from final I/R + PMNs; NS=not
25 significant.

Figure 10. Measurement of NO release from rat aortic segments. Endothelial NO release was significantly increased from basal NO release in PKC β II peptide inhibitor treated segments (1, 2.5, 5 and 10 μ M), as well as acetylcholine (Ach, 500nM). NO release was significantly reduced in both groups given 400 μ M L-NAME. All values are expressed as means $\pm$ SEM. Numbers at bottom of bars are numbers of separate experiments per group. * p <0.05, ** p <0.01, from basal values.

Figure 11. Superoxide release from rat PMNs. Superoxide release was measured from 5×10^6 PMNs after formyl-methionyl-leucyl-phenylalanine (fMLP) (200 nM) stimulation. SOD (10 μ g/ml) was employed as a positive control. The change in absorbance (Δ) was measured 90 sec after fMLP addition (peak response). Superoxide release was significantly inhibited by the PKC β II peptide inhibitor (** p <0.01, 5, 10 and 20 μ M). All values are means $\pm$ SEM. Numbers at bottom of bars are numbers of separate experiments per group.

Figure 12. Time course of LVDP in sham, I/R, I/R+PMNs and I/R+PMN+ PKC β II (10 μ M)+PKC ζ (5 μ M) peptide inhibitors perfused rat hearts. LVDP data at initial (baseline) and reperfusion from 0 to 45 min following 20 min ischemia. The sham group (n=6) maintained the same LVDP throughout the 80 min. protocol. The I/R (n=6) group partially recovered toward initial baseline values. I/R+PMN group (n=11) exhibited a significant and sustained reduction in LVDP compared to and I/R+PMN+ PKC β II (10 μ M)+PKC ζ (5 μ M) peptide inhibitors (n=7) group. All values are expressed as mean $\pm$ SEM. * p <0.05 from I/R + PMNs.

Figure 13. Initial and final LVDP expressed in mmHg from isolated perfused rat hearts before ischemia (I) (initial) and after 45 min post reperfusion (R) (final). Hearts were perfused in the presence or absence of PMNs. PMNs induced a significant contractile dysfunction, which was attenuated by the presence of PKC β II

and PKC ζ peptide inhibitors. This protective effect was blocked by L-NAME. All values are expressed as mean $\pm$ SEM. Numbers of hearts examined are at the bottom of the bars. * $p < 0.05$ from final I/R + PMNs; NS=not significant.

Figure 14. Initial and final +dP/dt max expressed in mmHg/s in isolated perfused rat hearts before ischemia (I) and after reperfusion (R). Hearts were perfused in the presence or absence of PMNs. PMNs induced a significant contractile dysfunction, which was attenuated by PKC β II and PKC ζ peptide inhibitors. This protective effect was blocked by L-NAME. All values are expressed as means $\pm$ SEM. Numbers of hearts examined are at the bottom of the bars. * $P < 0.05$ from final I/R + PMNs; NS=not significant.

Figure 15. Histological assessment of total intravascular and infiltrated PMNs in isolated perfused rat heart samples taken from 3 rats per group and 10 areas per heart. The numbers of total intravascular and infiltrated PMNs in post-reperfusion cardiac tissue and adhering to coronary vasculature was significantly attenuated by the PKC β II and PKC ζ peptide inhibitors. Hatched boxes represent non-PMN perfused hearts and black boxes represent PMN-perfused hearts. ** $P < 0.01$, from I/R+PMNs.

Figure 16. Histological assessment of intravascular PMNs that adhered to the coronary vasculature in isolated perfused rat heart. The numbers of PMNs adhering to the coronary vasculature in hearts treated with PKC β II and PKC ζ peptide inhibitors was lower than I/R+PMN hearts. Hatched boxes represent non-PMN perfused hearts and black boxes represent PMN-perfused hearts. All values are mean numbers of PMNs/mm² of heart area $\pm$ SEM. ** $P < 0.01$, from I/R+PMNs.

Figure 17. Measurement of NO release from rat aortic segments. Endothelial NO release was significantly increased from basal NO release in treated segments treated with PKC β II and PKC ζ peptide inhibitors, as well as acetylcholine (Ach, 500nM). NO release was significantly reduced in both groups given 400 μ M L-NAME. All values are expressed as means $\pm$ SEM. Numbers at bottom of bars are numbers of separate experiments per group. **p<0.01, from basal values.

Figure 18. Superoxide release from rat PMNs. Superoxide release was measured from 5×10^6 PMNs after PMA (15 nM) stimulation. SOD (10 μ g/ml) was employed as a positive control. The change in absorbance (Δ) was measured 360 sec after PMA addition (peak response). Superoxide release was significantly inhibited by the presence of PKC β II and PKC ζ peptide inhibitors (*p<0.05, **p<0.01). All values are means $\pm$ SEM. Numbers at bottom of bars show the numbers of separate experiments per group.

Figure 19. Time course of LVDP in sham, I/R, I/R+PMNs and I/R+PMN+ PKC δ peptide activator (10 μ M) perfused rat hearts. LVDP data at initial (baseline) and reperfusion from 0 to 45 min following 20 min ischemia. The sham group (n=6) maintained the same LVDP throughout the 80 min. protocol. The I/R (n=6) group recovered to initial baseline values. I/R+PMN group (n=6) exhibited a significant and sustained reduction in LVDP compared to and I/R+PMN+ PKC δ peptide activator (n=6) group. All values are expressed as mean $\pm$ SEM. **p<0.01, from I/R+PMNs.

Figure 20. Initial and final LVDP expressed in mmHg from isolated perfused rat hearts before ischemia (I) (initial) and after 45 min post reperfusion (R) (final). Hearts were perfused in the presence or absence of PMNs. PMNs induced a significant contractile dysfunction, which was attenuated by the PKC δ peptide activator. All values are expressed as mean $\pm$ SEM. Numbers of hearts examined are

at the bottom of the bars. $\overline{p}<0.05$ and $\overline{p}<0.01$, from final I/R+PMNs; NS=not significant.

Figure 21. Initial and final +dP/dt max expressed in mmHg/s in isolated perfused rat hearts before ischemia (I) and after reperfusion (R). Hearts were
5 perfused in the presence or absence of PMNs. PMNs induced a significant contractile dysfunction, which was attenuated by a PKC δ peptide activator. All values are expressed as means $\pm$ SEM. Numbers of hearts examined are at the bottom of the bars. $\overline{P}<0.01$, from final I/R + PMNs; NS=not significant.

Figure 22. Superoxide release from rat PMNs. Superoxide release was
10 measured from 5×10^6 PMNs after PMA (15 nM) stimulation. The change in absorbance (Δ) was measured 360 sec after PMA addition (peak response). Superoxide release was significantly inhibited by the PKC δ peptide activator ($\overline{p}<0.01$, 5 and 10 μ M). All values are means $\pm$ SEM. Numbers at bottom of bars show the numbers of separate experiments per group.

15 Figure 23a. Histological assessment of total intravascular and infiltrated PMNs in isolated perfused rat heart samples taken from 3 rats per group and 10 areas per heart. The numbers of total intravascular and infiltrated PMNs in post-reperfusion cardiac tissue and adhering to coronary vasculature was attenuated by the PKC δ peptide
20 activator. Hatched boxes represent non-PMN perfused hearts and black boxes represent PMN-perfused hearts.

Figure 23b. Histological assessment of intravascular PMNs that adhered to the coronary vasculature in isolated perfused rat heart samples taken from 3 rats per
group
25 and 10 areas per heart. The numbers of PMNs adhering to the coronary vasculature

was attenuated by the PKC δ peptide activator. Hatched boxes represent non-PMN perfused hearts and black boxes represent PMN-perfused hearts. All values are mean numbers of PMNs/mm² of heart area $\pm$ SEM.

Figure 24. Time course of left ventricular end diastolic pressure (LVEDP) in
5 sham, I/R, I/R+PMNs and I/R+PMN+ PKC δ peptide activator (10 μ M) perfused rat
hearts. LVEDP data at initial (baseline) and reperfusion from 0 to 45 min following
20 min ischemia. The sham group (n=6) maintained the same LVEDP throughout the
80 min. protocol. The I/R (n=6) group partially recovered to initial baseline values.
I/R+PMN group (n=6) exhibited a significant and sustained elevation in LVEDP
10 compared to and I/R+PMN+ PKC δ peptide activator (n=6) group. All values are
expressed as mean $\pm$ SEM. *p<0.05 and **p<0.01, from I/R+PMNs.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

The present invention provides a solution for the preservation, perfusion, and/or reperfusion of an organ, especially the heart. The solution contains peptide inhibitor(s) of protein kinase C β II (PKC β II) and/or of protein kinase C ζ (PKC ζ) and/or peptide activator(s) of PKC δ . Preferably, the peptide inhibitor of PKC β II or the peptide activator of PKC δ is present in the solution in an amount of about 5-10 μ M; and the peptide inhibitor of PKC ζ is present in an amount of about 2.5-5 μ M.

In a preferred embodiment, the peptide inhibitor of PKC β II has an amino acid sequence of SEQ ID NO: 1; the peptide inhibitor of PKC ζ has an amino acid sequence of SEQ ID NO: 2; and the peptide activator has an amino acid sequence of SEQ ID NO: 3. Also, in other embodiments, it is preferred that the peptide inhibitor/activator is myristoylated to facilitate absorption into the cells of the organ.

In a preferred embodiment, the peptide inhibitor(s) or peptide activator (s) are dissolved in a saline solution, preferably normal saline (0.9 % NaCl). The peptide inhibitor(s) can also be dissolved in known preservation solution, such as Krebs-Henseleit solution, UW solution, St. Thomas II solution, Collins solution, Stanford solution, and the like. The solution may also contain one or more of sodium (Na^+), potassium (K^+), calcium (Ca^{2+}), magnesium (Mg^{2+}), glutamate, arginine, adenosine, manitol, allopurinol, glutathione, raffinose, and lactobionic acid in concentrations of about 4-7 mM, about 0.2-0.3 mM, about 108-132 mM, about 13-16 mM, about 18-22 mM, about 2-4 mM, about 0.5-1 mM, about 27-33 mM, about 0.9-1.1 mM, about 2.7-3.3 mM, about 25-35 mM, and about 80-120 mM, respectively. Na^+ can be in the form of NaOH; K^+ can be in the form of KCl and/or KH_2PO_4 , most preferably at ratio of about 2-3.5 mM KCl and about 2-3.5 mM KH_2PO_4 ; Ca^{2+} can be in the form of

'CaCl₂ and Mg²⁺' can be in the form of MgCl₂. The solution is preferably maintained at physiological pH of about 7.2-7.4.

The solution of the present invention can be used during all phases of an organ, especially the heart, transplant, including, but are not limited to, 1) isolating of
5 the organ from the donor (cardioplegic solution); 2) preserving the organ (hypothermic storage/transport); and 3) re-implanting the organ in the recipient (reperfusion solution).

During perfusion or reperfusion, especially for the heart, it is preferred that the organ be perfused at a rate of about 1 mL/min for about 5 min. The perfusion rate can be
10 varied, but it should not exceed about 25 mL/min. Overall, the perfusion rate should not be so high as to impose undue pressure on the vasculature of the organ.

The solution of the present invention can be prepared by 1) dissolving and diluting the peptide inhibitor(s) and the different constituents in distilled water; 2) adjusting the pH to about 7.2-7.4, e.g. with NaOH; and 3) sterilizing the solution, e.g.,
15 by filtering with a 0.2 µm filter. The sterilized solution is then kept isolated from contaminants in the environment.

Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods.
20 The following example is given to illustrate the present invention. It should be understood that the invention is not to be limited to the specific conditions or details described in this example.

Example 1 – Effects of Peptide Inhibitor of PKC ζ

Male Sprague Dawley rats (275-325 g, Ace Animals, Boyertown, PA) were anesthetized with 60 mg/kg pentobarbital sodium intraperitoneally (i.p.). Sodium heparin (1,000 U) was also administered i.p. The hearts were rapidly excised, the ascending aortas were cannulated, and retrograde perfusion of the heart was initiated
5 with a modified Krebs buffer maintained at 37°C at a constant pressure of 80 mmHg. The Krebs buffer had the following composition (in mmol/l): 17 dextrose, 120 NaCl, 25 NaHCO₃, 2.5 CaCl₂, 0.5 EDTA, 5.9 KCl, and 1.2 MgCl₂. The perfusate was aerated with 95% O₂ and 5% CO₂ and equilibrated at a pH of 7.3-7.4. The two side arms in the perfusion line proximal to the heart inflow cannula allowed PMNs, plasma
10 without PKC ζ peptide inhibitor (control hearts) or plasma containing different concentrations of PKC ζ peptide inhibitor (1, 2.5 or 5 μ M) to be directly infused into the coronary inflow line. Coronary flow was monitored by a flow meter (T106, Transonic System, Inc., Ithaca, NY). LVDP and $+dP/dt_{\max}$ were monitored using a pressure transducer (SPR-524, Millar Instruments, Inc., Houston, TX), which was
15 positioned in the left ventricular cavity. Hearts were immersed in a water-jacketed reservoir containing 160 mL of Krebs buffer maintained at 37°C. Coronary flow, LVDP and $+dP/dt_{\max}$ were recorded using a Powerlab Station acquisition system (ADInstruments, Grand Junction, CO) in conjunction with a computer.

LVDP, $+dP/dt_{\max}$, and coronary flow were measured every 5 min for 15 min to
20 equilibrate the hearts and obtain a baseline measurement. LVDP was defined as left ventricular end-systolic pressure minus left ventricular end-diastolic pressure. After 15 min, the flow of the Krebs buffer was reduced to zero for 20 min to induce global ischemia. At reperfusion, hearts were infused for 5 min with 200×10^6 PMN resuspended in 5 mL of Krebs buffer plus 5 mL of plasma at a rate of 1 mL/min. In
25 some experiments, PKC ζ peptide inhibitor (Genemed Synthesis, Inc., San Francisco,

"CA") was added to plasma at a final concentration of 1, 2.5 or 5 μ M. Sham I/R hearts were not subjected to ischemia and were not perfused with PMNs.

The following groups of isolated perfused rat hearts were used:

- Group 1: Sham Ischemia/Reperfusion (I/R) hearts was not subjected to ischemia and not perfused with PMNs, but was perfused with 5 mL of plasma (1 mL/min) at 35 minutes into perfusion (the same time point that I/R hearts would be given 5 mL of plasma, 15 minutes of baseline recordings plus 20 minutes ischemia). These hearts represented a control group to determine if the isolated rat heart can maintain LVDP and $+dP/dt_{\max}$ throughout the 80-minute protocol (n=6).
- Group 2: Sham I/R + PKC ζ peptide inhibitor (5 μ M) hearts were not subjected to ischemia and not perfused with PMNs. These hearts were administered the PKC ζ peptide inhibitor (5 μ M, dissolved in plasma from a 5 mM stock in H₂O) 35 minutes into perfusion. This group was employed to determine if the PKC ζ peptide inhibitor causes a cardiotonic or cardiodepressant effect (n=6).
- Group 3: I/R hearts were subjected to 20 min of ischemia and perfused with 5 mL of plasma (1 mL/min) during the first 5 min of reperfusion, but were not perfused with PMNs. These hearts represented a control group to determine if 20 min of ischemia followed by reperfusion stuns the heart, but LVDP and $+dP/dt_{\max}$ will recover to baseline values (initial) by the end of the 45-minute reperfusion period (n=6).
- Group 4: I/R + PKC ζ peptide inhibitor (5 μ M, dissolved in plasma) hearts were subjected to 20 min. of ischemia and not perfused with PMNs. These hearts were perfused with 5 mL of plasma + PKC ζ inhibitor during the first 5 min of reperfusion. This group was employed to determine if the PKC ζ peptide inhibitor caused a cardiodepressant effect in the setting of I/R without PMNs (n=6).

Group 5: I/R + PMNs hearts were subjected to 20 min of ischemia and perfused with 5 mL of plasma (1 mL/min) and PMNs (resuspended in 5 mL Krebs buffer) during the first 5 min of reperfusion. These hearts represented a control group to determine if 20 min of ischemia followed by 45 min reperfusion in the presence of PMNs (200×10^6) resulted in a sustained cardiac contractile dysfunction throughout the 45 min reperfusion period compared to initial baseline values ($n=6$).

Group 6: I/R + PMNs + PKC ζ peptide inhibitor (1 μ M) hearts were subjected to 20 min of ischemia and perfused with 1 μ M PKC ζ peptide inhibitor (dissolved in plasma) and PMNs (200×10^6) during the first 5 minutes of reperfusion. These hearts represented a group to determine the effect of PKC ζ inhibition in attenuating PMN-induced cardiac contractile dysfunction ($n=6$).

Group 7: I/R + PMNs + PKC ζ peptide inhibitor (2.5 μ M) hearts were subjected to 20 min of ischemia and perfused with 2.5 μ M PKC ζ peptide inhibitor (dissolved in plasma) and PMNs (200×10^6) during the first 5 minutes of reperfusion. These hearts represented a group to determine the effect of PKC ζ inhibition in attenuating PMN-induced cardiac contractile dysfunction ($n=6$).

Group 8: I/R + PMNs + PKC ζ peptide inhibitor (5 μ M) hearts were subjected to 20 min of ischemia and are perfused with 5 μ M PKC ζ peptide inhibitor (dissolved in plasma) and PMNs (200×10^6) during the first 5 minutes of reperfusion. These hearts represented a group to determine the effect of PKC ζ inhibition at a higher concentration of the PKC ζ peptide inhibitor in attenuating PMN-induced cardiac contractile dysfunction ($n=6$).

Group 9: I/R + PMNs + PKC ζ peptide inhibitor (5 μ M) + N^G -nitro-L-arginine methyl ester (L-NAME, 50 μ M) hearts were subjected to 20 min of ischemia and perfused with 5 μ M PKC ζ peptide inhibitor (dissolved in 5 mL plasma) and 50

μM L-NAME (dissolved in Krebs buffer from 50 mM stock in H_2O) and PMNs (200×10^6) during the first 5 minutes of reperfusion. The L-NAME (50 μM) was continually infused into the heart throughout the 45 min reperfusion period. These hearts represent a group to determine if the cardioprotective effect of PKC ζ peptide inhibition could be blocked with a nitric oxide synthase inhibitor (L-NAME) (n=5).

Data were recorded every 5 min for 45 min post-reperfusion. After each experiment, the left ventricle was isolated, fixed in 4% paraformaldehyde and stored at 4°C for later histological analysis.

Figure 1 showed the time course of cardiac contractile function (i.e., LVDP).

The data from the sham I/R, I/R, I/R+PMN+ PKC ζ peptide inhibitor (5 μM) and I/R+PMN groups illustrated the relative changes in LVDP during the 80 min perfusion period. As shown, the sham I/R remained near or greater than 100% of initial baseline values of LVDP for the entire perfusion period. The I/R hearts experienced a depression in LVDP at the beginning of reperfusion, but recovered to 95 $\pm$ 7% of initial baseline values by the end of reperfusion. In contrast, the I/R+PMN hearts suffered severe cardiac contractile dysfunction, recovering to only 47 $\pm$ 7% of initial baseline values by 45 min post-reperfusion. Conversely, the I/R+PMN+ PKC ζ peptide inhibitor (5 μM) hearts recovered to 84 $\pm$ 4% at 45 min post-reperfusion.

To determine whether PKC ζ peptide inhibitor produced direct inotropic effects on cardiac contractile function, non-ischemic sham I/R hearts were perfused with PKC ζ peptide inhibitor (5 μM). Treatment of Sham I/R hearts with PKC ζ peptide inhibitor did not result in any significant change in LVDP (Fig. 2) or $+\text{dP}/\text{dt}_{\text{max}}$ (Fig. 3) during the 80 min perfusion period, demonstrating the PKC ζ peptide inhibitor at 5 μM exerts no direct effect on cardiac contractile function. A 15 μM PKC ζ peptide inhibitor concentration was initially tested, since this concentration

corresponded with an 82% inhibition of PMN superoxide release. However, 15 μ M produced a cardiodepressant effect (50% reduction in LVDP Sham I/R hearts) and could not be used in these experiments (data not shown).

Figures 2 and 3 showed the initial and final values for LVDP and $+dP/dt_{\max}$ from isolated perfused rat hearts. The initial baselines were similar for all groups. However, the final LVDP and $+dP/dt_{\max}$ (45 min post-reperfusion) was significantly decreased ($p < 0.01$) by $47 \pm 7\%$ and $41 \pm 7\%$ respectively for the I/R hearts reperfused with PMNs compared to its initial baseline. The PKC ζ peptide inhibitor (2.5 μ M and 5 μ M concentrations) significantly attenuated the decrease in LVDP and $+dP/dt_{\max}$ associated with post-ischemic reperfusion with PMNs. In the group receiving 5 μ M of drug, the hearts recovered to $84 \pm 4\%$ and $76 \pm 5\%$ for final LVDP and $+dP/dt_{\max}$ compared to its initial baseline. The lowest effective dose was observed at 2.5 μ M; and these hearts recovered to $83 \pm 4\%$ for LVDP and $75 \pm 5\%$ for $+dP/dt_{\max}$ compared to initial baseline values. The cardioprotective effects of the PKC ζ peptide inhibitor (5 μ M) were blocked in the presence of L-NAME (50 μ M). These hearts only recovered to $58 \pm 7\%$ and $54 \pm 9\%$ for LVDP and $+dP/dt_{\max}$, respectively, at 45 min post-reperfusion compared to its initial baseline. These hearts were similar to the IR + PMN group ($47 \pm 7\%$ and $41 \pm 7\%$ LVDP, $+dP/dt_{\max}$). At 1 μ M, PKC ζ peptide inhibitor treated hearts exposed to I/R + PMNs only recovered to $57 \pm 10\%$ and $46 \pm 6\%$ for LVDP and $+dP/dt_{\max}$, respectively, at 45 min post-reperfusion compared to its initial baseline. These hearts were not significantly different from control I/R + PMN hearts at 45 min post-reperfusion at this lower dose.

The cardiac injury associated with I/R in this model was closely correlated with the substantial number of PMNs infiltrating the myocardium within the 45 min reperfusion period. During reperfusion, a significant number of PMNs transmigrated

into the myocardium, increasing from less than 25 PMN/mm² in Sham I/R hearts to more than 180 PMN/mm² in I/R +PMN hearts at the end of the reperfusion period (Fig. 4a). In contrast, I/R+PMN+PKC ζ peptide inhibitor treated hearts experienced a 20 $\pm$ 6%, 46 $\pm$ 4 % and 48 $\pm$ 3% significant reduction in PMN infiltration into the post-reperfused cardiac tissue at 1, 2.5 and 5 μ M (p<0.01), respectively; and this effect was blocked in the presence of L-NAME. Furthermore, the 2.5 and 5 μ M treated hearts had significantly fewer infiltrated PMNs compared to 1 μ M treated hearts (p<0.01) (Fig. 4a).

PMN adherence to coronary vascular endothelium was also evaluated within the assessment of total intravascular and infiltrated PMNs. As seen in Figure 4b, the number of adherent PMNs to the coronary endothelium was not significantly reduced in I/R+PMN+PKC ζ peptide inhibitor hearts (5 μ M) (43 $\pm$ 10%, p<0.06) (Fig. 4b). NO release from rat aortic endothelium was measured to determine if PKC ζ peptide inhibitor provides cardioprotection by a mechanism involving increased endothelial NO release. In Figure 5, PKC ζ peptide inhibitor-treated endothelium generated significantly more NO by 47 $\pm$ 2% (2.5 μ M, p<0.05), 54 $\pm$ 5% (5 μ M, p<0.01), and 91 $\pm$ 15% (15 μ M, p<0.01) compared to basal NO release. At 1 μ M, NO release was not significantly different from basal NO release. Acetylcholine (200 nM) was used as a positive control in the NO assay, and significantly increased NO release by 67 $\pm$ 4% (p<0.01) compared to basal NO release. The eNOS inhibitor, L-NAME, was used as another control in order to decrease basal release of NO to zero. Both the acetylcholine and the PKC ζ peptide inhibitor-induced production of NO were completely inhibited by treating the endothelium with L-NAME (400 μ M). To attribute the source of the NO to the endothelium, experiments with endothelium

removed (denuded) rat aortic segments were incubated with PKC ζ peptide inhibitor (5 μ M) and were not different from L-NAME –treated segments.

Another mechanism of the cardioprotective effects of PKC ζ peptide inhibitor may be related to inhibition of superoxide release. PKC ζ peptide inhibitor significantly inhibited superoxide release by 33-82% (2.5 - 15 μ M, $p < 0.01$), except at 1 μ M where there was no difference from suspensions of PMA-stimulated rat PMNs (Fig. 6). SOD (10 μ g /mL) was used as a positive control in the superoxide assays, and degraded superoxide release produced by the PMA-stimulated rat PMNs by 99% ($p < 0.01$; Fig. 6).

10

Example 2 – Effects of Peptide Inhibitor of PKC β II

Experiments with PKC β II peptide inhibitors were performed substantially as described in Example 1 for PKC ζ peptide inhibitors.

The following groups of isolated perfused rat hearts were used:

15 Group 1: Sham I/R hearts were not subjected to ischemia and were not perfused with PMNs, but were perfused with 5 mL of plasma (1 mL/min) at 35 minutes into perfusion (the same time point that I/R hearts would be given 5 mL of plasma, 15 minutes of baseline recordings plus 20 minutes ischemia). These hearts represented a control group to determine if the isolated rat heart can maintain LVDP and $+dP/dt_{\max}$ throughout the 80-minute protocol ($n=6$).

Group 2: Sham I/R + PKC β II peptide inhibitor (10 μ M) hearts were not subjected to ischemia and not perfused with PMNs. These hearts were administered the PKC β II peptide inhibitor (10 μ M, dissolved in plasma from a 5 mM stock in H_2O) 35 minutes into perfusion. This group was employed to determine if the PKC β II peptide inhibitor causes a cardiotonic or cardiodepressant effect ($n=6$).

25

Group 3: I/R hearts were subjected to 20 min of ischemia and perfused with 5 mL of plasma (1 mL/min) during the first 5 min of reperfusion, but were not perfused with PMNs. These hearts represented a control group to determine if 20 min of ischemia followed by reperfusion stunned the heart, but LVDP and $+dP/dt_{max}$ will
5 recover to baseline values (initial) by the end of the 45-minute reperfusion period (n=6).

Group 4: I/R + PKC β II peptide inhibitor (10 μ M, dissolved in plasma) hearts were subjected to 20 min of ischemia and not perfused with PMNs. These hearts were perfused with 5 mL of plasma + PKC β II peptide inhibitor during the first 5 min
10 of reperfusion. This group was employed to determine if the PKC β II peptide inhibitor causes a cardiodepressant effect in the setting of I/R without PMNs (n=6).

Group 5: I/R + PMNs hearts were subjected to 20 min of ischemia and perfused with 5 mL of plasma (1 mL/min) and PMNs (resuspended in 5 mL Krebs buffer) during the first 5 min of reperfusion. These hearts represented a control group
15 to determine if 20 min of ischemia followed by 45 min reperfusion in the presence of PMNs (200×10^6) resulted in a sustained cardiac contractile dysfunction throughout the 45 min reperfusion period compared to initial baseline values (n=9).

Group 6: I/R + PMNs + PKC β II peptide inhibitor (1 μ M) hearts were subjected to 20 min of ischemia and perfused with 1 μ M PKC β II peptide inhibitor
20 (dissolved in plasma) and PMNs (200×10^6) during the first 5 minutes of reperfusion. These hearts represented a group to determine the effect of PKC β II inhibition in attenuating PMN-induced cardiac contractile dysfunction (n=6).

Group 7: I/R + PMNs + PKC β II peptide inhibitor (5 μ M) hearts were subjected to 20 min of ischemia and perfused with 5 μ M PKC β II peptide inhibitor
25 (dissolved in plasma) and PMNs (200×10^6) during the first 5 minutes of reperfusion.

These hearts represented a group to determine the effect of PKC β II inhibition at a higher concentration of the PKC β II peptide inhibitor in attenuating PMN-induced cardiac contractile dysfunction (n=7).

Group 8: I/R + PMNs + PKC β II peptide inhibitor (10 μ M) hearts were
5 subjected to 20 min of ischemia and perfused with 10 μ M PKC β II peptide inhibitor (dissolved in plasma) and PMNs (200×10^6) during the first 5 minutes of reperfusion. These hearts represented a group to determine the effect of PKC β II inhibition at a higher concentration of the PKC β II peptide inhibitor in attenuating PMN-induced cardiac contractile dysfunction (n=7).

10 Group 9: I/R + PMNs + PKC β II peptide inhibitor (10 μ M) + N^G -nitro-L-arginine methyl ester (L-NAME, 50 μ M) hearts were subjected to 20 min of ischemia and perfused with 10 μ M PKC β II peptide inhibitor (dissolved in 5 mL plasma) and 50 μ M L-NAME (dissolved in Krebs buffer from 50 mM stock in H_2O) and PMNs (200×10^6) during the first 5 minutes of reperfusion. The L-NAME (50 μ M) was
15 continually infused into the heart throughout the 45 min reperfusion period. These hearts represented a group to determine if the cardioprotective effect of PKC β II peptide inhibition can be blocked with a nitric oxide synthase inhibitor (L-NAME) (n=6).

Previous studies showed that sham I/R hearts given PMNs exhibited no
20 changes from initial control values (Lefer et al., *Circulation* 100: 178-184, 1999). Data were recorded every 5 min for 45 min post-reperfusion. After each experiment, the left ventricle was isolated, fixed in 4% paraformaldehyde and stored at 4°C for later histological analysis.

Figure 7 showed the time course of cardiac contractile function (LVDP) for
25 the sham I/R, I/R, I/R+PMN+PKC β II peptide inhibitor (10 μ M) and I/R+PMN

groups, and illustrated the changes in LVDP during the 80 min perfusion period. The hearts in the Sham I/R group remained at $103 \pm 4\%$ of initial baseline values of LVDP for the entire duration of the perfusion period. Hearts in the I/R group experienced a depression in LVDP during the initial stages of reperfusion, but by the end of

5 reperfusion they had recovered to $94 \pm 6\%$ of initial baseline values. However, the hearts in the I/R+PMN group exhibited severe cardiac contractile dysfunction, only recovering to $43 \pm 5\%$ of initial baseline values by the end of reperfusion. By contrast, the hearts in the I/R+PMN+PKC β II peptide inhibitor ($10 \mu\text{M}$), although initially showing a depression in LVDP of $61 \pm 10\%$ of initial baseline values at 15 min into

10 reperfusion, recovered to $82 \pm 9\%$ of baseline.

In order to establish whether the PKC β II peptide inhibitor produced any direct inotropic effects on cardiac contractile function, Sham I/R hearts were perfused with PKC β II peptide inhibitor ($10 \mu\text{M}$). This group served as one of the controls for the study. These hearts did not show any significant change in LVDP (Fig. 8) or

15 $+dP/dt_{\text{max}}$ (Fig. 9) at the end of the 80 min reperfusion period, thus, indicating that at this dose the PKC β II peptide inhibitor had no direct effect on cardiac contractile function. A $20 \mu\text{M}$ dose of PKC β II peptide inhibitor was tested on a Sham I/R heart and there were no cardiodepressant effects noticed at this dose.

Figs. 8 and 9 showed the initial and final values for LVDP and $+dP/dt_{\text{max}}$ from

20 isolated perfused hearts respectively. There was no significant difference between the initial baseline values of all the groups studied. There was also no significant difference between the initial and final values of LVDP and $+dP/dt_{\text{max}}$ for the Sham I/R, I/R, Sham I/R+PKC β II peptide inhibitor ($5 \mu\text{M}$), and I/R+PKC β II peptide inhibitor ($10 \mu\text{M}$) groups. However, there was a significant difference between the

25 initial and final values of LVDP and $+dP/dt_{\text{max}}$ for the I/R+PMN group. A significant

decrease ($p<0.01$) from initial baseline of $43\pm5\%$ in LVDP and $42\pm6\%$ in $+dP/dt_{\max}$ at 45 min post-reperfusion was observed.

The presence of the PKC β II peptide inhibitor at a 5 μ M and 10 μ M dose attenuated the decrease in LVDP and $+dP/dt_{\max}$ associated with the post-ischemic perfusion with PMNs. The 10 μ M dose was the most cardioprotective as the hearts in the I/R+PMN+PKC β II peptide inhibitor (10 μ M) recovered to $82\pm9\%$ and $79\pm10\%$ of initial baseline at 45 min post-reperfusion for LVDP and $+dP/dt_{\max}$, respectively. These values were significantly different from I/R+PMN at 45 min post-reperfusion ($p<0.01$). The 5 μ M dose was also cardioprotective, although not to the same extent as the 10 μ M dose, as the hearts in the I/R+PMN+PKC β II peptide inhibitor (5 μ M) recovered to $69\pm7\%$ and $63\pm7\%$ for LVDP and $+dP/dt_{\max}$ of initial baseline at 45 min post-reperfusion, respectively. The LVDP values for the 5 μ M dose were significantly different from I/R+PMN at 45 min post-reperfusion ($p<0.05$). The 1 μ M dose of PKC β II peptide inhibitor was not cardioprotective as the hearts in the I/R+PMN+PKC β II peptide inhibitor (1 μ M) group only recovered to $57\pm4\%$ and $53\pm6\%$ for LVDP and $+dP/dt_{\max}$ respectively. The final values of LVDP and $+dP/dt_{\max}$ at the 1 μ M dose group were not significantly different from the final values of the I/R+PMN group.

The cardioprotective effects of the PKC β II peptide inhibitor (10 μ M) were blocked by the presence of L-NAME (50 μ M) in the IR+PMN+PKC β II peptide inhibitor (10 μ M)+L-NAME (50 μ M) group, as the LVDP and $+dP/dt_{\max}$ values at the end of the 45 min reperfusion period were only $56\pm2\%$ and $53\pm5\%$ of the initial baseline values, respectively, and were not significantly different from the final values of the IR+PMN group (Figs. 3 and 4).

NO release from rat aortic endothelium was measured to determine if PKC β II peptide inhibitor provides cardioprotection by a mechanism involving increased endothelial NO release. Fig. 10 showed that segments of the endothelium treated with PKC β II peptide inhibitor generated significantly more NO when compared to the basal NO release at 5 μ M ($p<0.05$) and 10 μ M ($p<0.01$). The basal value of NO release was measured at 1.85 ± 0.18 pmoles NO/mg tissue. There was a definite dose-response effect of stimulating the endothelium with PKC β II peptide inhibitor as the 1 μ M, 2.5 μ M, 5 μ M and 10 μ M produced an increase in NO release above basal of 0.75 ± 0.19 , 1.91 ± 0.44 , 2.54 ± 0.29 , and 3.49 ± 0.62 pmoles NO/mg tissue, respectively.

Acetylcholine (Ach, 500 nM) was used as a positive control in this assay and stimulated the endothelium causing an increase of 3.75 ± 0.58 pmoles NO/mg tissue, above the baseline basal value. L-NAME was used as another control in order to decrease basal release of NO to zero. Both the acetylcholine and PKC β II peptide inhibitor production of NO were completely inhibited by treating the endothelium with L-NAME (400 μ M).

Another mechanism that may contribute to the cardioprotective effects (i.e. LVDP) of the PKC β II peptide inhibitor may be inhibition of PMN superoxide release. PKC β II peptide inhibitor significantly inhibited superoxide release (i.e. absorbance) from suspensions of fMLP-stimulated rat PMNS from 0.13 ± 0.01 to 0.05 ± 0.009 ($p<0.01$), 0.02 ± 0.004 ($p<0.01$), and 0.02 ± 0.007 ($p<0.01$) for 5 μ M, 10 μ M and 20 μ M respectively (Fig. 11). There was no significant inhibition of superoxide at the 1 μ M dose. SOD (10 μ g/ml) was used as a positive control and it scavenged the superoxide released by the fMLP-stimulated rat PMNs reducing the response to 0.0016 ± 0.0006 .

Example 3 – Effects of Peptide Inhibitors of PKC β II and PKC ζ in Combination

Experiments with PKC β II and PKC ζ peptide inhibitors were performed substantially as described in Examples 1 and 2.

The following groups of isolated perfused rat hearts were used:

5 Group 1: Sham I/R hearts were not subjected to ischemia and were not perfused with PMNs, but were perfused with 5 mL of plasma (1 mL/min) at 35 minutes into perfusion (the same time point that I/R hearts would be given 5 mL of plasma, 15 minutes of baseline recordings plus 20 minutes ischemia). These hearts represented a control group to determine if the isolated rat heart can maintain LVDP
10 and $+dP/dt_{\max}$ throughout the 80-minute protocol (n=6).

 Group 2: Sham I/R + PKC β II (10 μ M) + PKC ζ (5 μ M) peptide inhibitors hearts were not subjected to ischemia and not perfused with PMNs. These hearts were administered the PKC β II and PKC ζ peptide inhibitor (10 μ M and 5 μ M, respectively, dissolved in plasma from a 5 mM stock in H₂O) 35 minutes into
15 perfusion. This group was employed to determine if the peptide inhibitors cause a cardiotoxic or cardiodepressant effect (n=6).

 Group 3: I/R hearts were subjected to 20 min of ischemia and perfused with 5 mL of plasma (1 mL/min) during the first 5 min of reperfusion, but were not perfused with PMNs. These hearts represented a control group to determine if 20 min of
20 ischemia followed by reperfusion stunned the heart, but LVDP and $+dP/dt_{\max}$ will recover to baseline values (initial) by the end of the 45-minute reperfusion period (n=6).

 Group 4: I/R + PKC β II (10 μ M) + PKC ζ (5 μ M) peptide inhibitors (dissolved in plasma) hearts were subjected to 20 min of ischemia and not perfused
25 with PMNs. These hearts were perfused with 5 mL of plasma + PKC β II + PKC ζ

peptide inhibitors during the first 5 min of reperfusion. This group was employed to determine if the peptide inhibitors cause a cardiodepressant effect in the setting of I/R without PMNs (n=6).

Group 5: I/R + PMNs hearts were subjected to 20 min of ischemia and perfused with 5 mL of plasma (1 mL/min) and PMNs (resuspended in 5 mL Krebs buffer) during the first 5 min of reperfusion. These hearts represented a control group to determine if 20 min of ischemia followed by 45 min reperfusion in the presence of PMNs (200×10^6) resulted in a sustained cardiac contractile dysfunction throughout the 45 min reperfusion period compared to initial baseline values (n=11).

Group 6: I/R + PMNs + PKC β II (5 μ M) + PKC ζ (2.5 μ M) peptide inhibitors hearts were subjected to 20 min of ischemia and perfused with 1 μ M PKC β II peptide inhibitor (dissolved in plasma) and PMNs (200×10^6) during the first 5 minutes of reperfusion. These hearts represented a group to determine the effect of PKC β II inhibition in attenuating PMN-induced cardiac contractile dysfunction (n=7).

Group 7: I/R + PMNs + PKC β II (10 μ M) + PKC ζ (2.5 μ M) peptide inhibitors hearts were subjected to 20 min of ischemia and perfused with 5 μ M PKC β II peptide inhibitor (dissolved in plasma) and PMNs (200×10^6) during the first 5 minutes of reperfusion. These hearts represented a group to determine the effect of PKC β II inhibition at a higher concentration of the PKC β II peptide inhibitor in attenuating PMN-induced cardiac contractile dysfunction (n=6).

Group 8: I/R + PMNs + PKC β II (10 μ M) + PKC ζ (5 μ M) peptide inhibitors hearts were subjected to 20 min of ischemia and perfused with 10 μ M PKC β II peptide inhibitor (dissolved in plasma) and PMNs (200×10^6) during the first 5 minutes of reperfusion. These hearts represented a group to determine the effect of

PKC β II inhibition at a higher concentration of the PKC β II peptide inhibitor in attenuating PMN-induced cardiac contractile dysfunction (n=7).

Group 9: I/R + PMNs + PKC β II (10 μ M) + PKC ζ (5 μ M) peptide inhibitors + *N*^G-nitro-L-arginine methyl ester (L-NAME, 50 μ M) hearts were subjected to 20 min of ischemia and perfused with 10 μ M PKC β II and 5 μ M PKC ζ peptide inhibitor (dissolved in 5 mL plasma) and 50 μ M L-NAME (dissolved in Krebs buffer from 50 mM stock in H₂O) and PMNs (200 x 10⁶) during the first 5 minutes of reperfusion. The L-NAME (50 μ M) was continually infused into the heart throughout the 45 min reperfusion period. These hearts represented a group to determine if the cardioprotective effect of the peptide inhibitors can be blocked with a nitric oxide synthase inhibitor (L-NAME) (n=5).

From the time course of Fig. 12, the combination of PKC β II (10 μ M) and PKC ζ (5 μ M) peptide inhibitors resulted in a recovery rate matching that of the I/R heart. This result was not observed when either PKC β II (10 μ M) or PKC ζ (5 μ M) peptide inhibitor was used alone (Examples 1 and 2) where the recovery rate of the peptide inhibitor treated heart was slower than that of the I/R heart.

Figs. 13 and 14 showed the initial and final values for LVDP and $+dP/dt_{\max}$ from isolated perfused hearts respectively. As expected and similar to Examples 1 and 2, the initial and final values of LVDP and $+dP/dt_{\max}$ for the I/R+PMN group change significantly, decreasing (p<0.01) from initial baseline by about 50% in LVDP and by about 45% in $+dP/dt_{\max}$ at 45 min post-reperfusion.

The presence of the PKC β II and PKC ζ peptide inhibitors attenuated the decrease in LVDP and $+dP/dt_{\max}$ associated with the post-ischemic perfusion with PMNs. The 10 μ M PKC β II and 5 μ M PKC ζ peptide inhibitors dose was the most cardioprotective. Increasing the PKC β II peptide inhibitors dosage from 5 μ M to 10

μM did not result in a significant change in the cardioprotective effect. On the other hand, increasing the PKC ζ peptide inhibitors dosage from 2.5 μM to 5 μM resulted in a marginal increase in the cardioprotective effect.

The cardioprotective effects of the combination of PKC βII and PKC ζ peptide inhibitors were blocked by the presence of L-NAME (50 μM) in the I/R + PMNs + PKC βII (10 μM) + PKC ζ (5 μM) peptide inhibitors+L-NAME (50 μM) group, as the LVDP and $+\text{dP}/\text{dt}_{\text{max}}$ values at the end of the 45 min reperfusion period were only about 50% and 40% of the initial baseline values, respectively, and were not significantly different from the final values of the IR+PMN group (Figs. 13 and 14).

The cardiac injury associated with I/R in this model was closely correlated with the substantial number of PMNs infiltrating the myocardium within the 45 min reperfusion period. During reperfusion, a significant number of PMNs transmigrated into the myocardium, increasing from less than 25 PMN/ mm^2 in Sham I/R hearts to more than 175 PMN/ mm^2 in I/R +PMN hearts at the end of the reperfusion period (Fig. 15). In contrast, I/R+PMN+PKC βII + PKC ζ peptide inhibitors treated hearts experienced a significant reduction in PMN infiltration into the post-reperfused cardiac tissue. This effect was blocked in the presence of L-NAME. Furthermore, doubling the PKC βII peptide inhibitor dosage from 5 μM to 10 μM , while maintaining the PKC ζ peptide inhibitor dosage at 2.5 μM , did not significantly decrease the PMN infiltration. On the other hand, doubling the PKC ζ peptide inhibitor dosage from 2.5 μM to 5 μM , while maintaining the PKC βII peptide inhibitor dosage at 10 μM , marginally decrease the PMN infiltration (Fig. 15).

PMN adherence to coronary vascular endothelium was also evaluated within the assessment of total intravascular and infiltrated PMNs. As seen in Figure 16, the number of adherent PMNs to the coronary endothelium was reduced in

I/R+PMN+PKC β II+PKC ζ peptide inhibitors hearts. This protective effect was reversed in the presence of L-NAME.

In Figure 17, endothelium of hearts treated with PKC β II and PKC ζ peptide inhibitors generated significantly more NO release, particularly at a dosage of 10 μ M PKC β II and 5 μ M PKC ζ peptide inhibitors, compared to basal NO release. Acetylcholine (500 nM) was used as a positive control in the NO assay, and significantly increased NO release compared to basal NO release. L-NAME was used as another control in order to decrease basal release of NO to zero. Both the acetylcholine and the peptide inhibitors-induced production of NO were completely inhibited by treating the endothelium with L-NAME (400 μ M).

Another mechanism of the cardioprotective effects of PKC β II and PKC ζ peptide inhibitors may be related to inhibition of superoxide release. As seen from Fig. 18, the combination of PKC β II and PKC ζ peptide inhibitors significantly inhibited superoxide release when compared to suspensions of PMA-stimulated rat PMNs or when compared to PKC β II or PKC ζ peptide inhibitor used alone. SOD (10 μ g/mL) was used as a positive control in the superoxide assays, and degraded superoxide release produced by the PMA-stimulated rat PMNs by 99% (Fig. 18).

The Examples all showed that the presence of sufficient amount of PKC β II peptide inhibitor and/or PKC ζ peptide inhibitor provided significant cardioprotective effect by attenuating PMN-induced cardiac dysfunction.

Example 4 – Effects of Peptide Activator of PKC δ

Experiments with PKC δ peptide activators were performed substantially as described in Example 1 for PKC ζ peptide inhibitor.

The following groups of isolated perfused rat hearts were used:

Group 1: Sham I/R hearts were not subjected to ischemia and were not perfused with PMNs, but were perfused with 5 mL of plasma (1 mL/min) at 35 minutes into perfusion (the same time point that I/R hearts would be given 5 mL of plasma, 15 minutes of baseline recordings plus 20 minutes ischemia). These hearts represented a control group to determine if the isolated rat heart can maintain LVDP and $+dP/dt_{\max}$ throughout the 80-minute protocol (n=6).

Group 2: Sham I/R + PKC δ peptide activator (10 μ M) hearts were not subjected to ischemia and not perfused with PMNs. These hearts were administered the PKC δ peptide activator (10 μ M, dissolved in plasma from a 5 mM stock in H₂O) 35 minutes into perfusion. This group was employed to determine if the PKC δ peptide activator causes a cardiotonic or cardiodepressant effect (n=6).

Group 3: I/R hearts were subjected to 20 min of ischemia and perfused with 5 mL of plasma (1 mL/min) during the first 5 min of reperfusion, but were not perfused with PMNs. These hearts represented a control group to determine if 20 min of ischemia followed by reperfusion stunned the heart, but LVDP and $+dP/dt_{\max}$ will recover to baseline values (initial) by the end of the 45-minute reperfusion period (n=6).

Group 4: I/R + PKC δ peptide activator (10 μ M, dissolved in plasma) hearts were subjected to 20 min of ischemia and not perfused with PMNs. These hearts were perfused with 5 mL of plasma + PKC δ peptide activator during the first 5 min of reperfusion. This group was employed to determine if the PKC δ peptide activator causes a cardiodepressant effect in the setting of I/R without PMNs (n=6).

Group 5: I/R + PMNs hearts were subjected to 20 min of ischemia and perfused with 5 mL of plasma (1 mL/min) and PMNs (resuspended in 5 mL Krebs buffer) during the first 5 min of reperfusion. These hearts represented a control group

to determine if 20 min of ischemia followed by 45 min reperfusion in the presence of PMNs (200×10^6) resulted in a sustained cardiac contractile dysfunction throughout the 45 min reperfusion period compared to initial baseline values ($n=10$).

Group 6: I/R + PMNs + PKC δ peptide activator ($1 \mu\text{M}$) hearts were
5 subjected to 20 min of ischemia and perfused with $1 \mu\text{M}$ PKC βII peptide inhibitor (dissolved in plasma) and PMNs (200×10^6) during the first 5 minutes of reperfusion. These hearts represented a group to determine the effect of PKC δ activation in attenuating PMN-induced cardiac contractile dysfunction ($n=6$).

Group 7: I/R + PMNs + PKC δ peptide activator ($5 \mu\text{M}$) hearts were
10 subjected to 20 min of ischemia and perfused with $5 \mu\text{M}$ PKC βII peptide inhibitor (dissolved in plasma) and PMNs (200×10^6) during the first 5 minutes of reperfusion. These hearts represented a group to determine the effect of PKC δ activation at a higher concentration of the PKC δ peptide activator in attenuating PMN-induced cardiac contractile dysfunction ($n=6$).

Group 8: I/R + PMNs + PKC δ peptide activator ($10 \mu\text{M}$) hearts were
15 subjected to 20 min of ischemia and perfused with $10 \mu\text{M}$ PKC δ peptide activator (dissolved in plasma) and PMNs (200×10^6) during the first 5 minutes of reperfusion. These hearts represented a group to determine the effect of PKC δ activation at a higher concentration of the PKC δ peptide activator in attenuating PMN-induced
20 cardiac contractile dysfunction ($n=6$).

Previous studies showed that sham I/R hearts given PMNs exhibited no changes from initial control values (Lefer et al., *Circulation* 100: 178-184, 1999). Data were recorded every 5 min for 45 min post-reperfusion. After each experiment, the left ventricle was isolated, fixed in 4% paraformaldehyde and stored at 4°C for
25 later histological analysis.

Figure 19 showed the time course of cardiac contractile function (LVDP) for the sham I/R, I/R, I/R+PMN+PKC δ peptide activator (10 μ M) and I/R+PMN groups, and illustrated the changes in LVDP during the 80 min perfusion period. The hearts in the Sham I/R group remained at $103 \pm 4\%$ of initial baseline values of LVDP for the entire duration of the perfusion period. Hearts in the I/R group experienced a depression in LVDP during the initial stages of reperfusion, but by the end of reperfusion they had recovered to $82 \pm 3\%$ of initial baseline values. However, the hearts in the I/R+PMN group exhibited severe cardiac contractile dysfunction, only recovering to $46 \pm 9\%$ of initial baseline values by the end of reperfusion. By contrast, the hearts in the I/R+PMN+PKC δ peptide activator (10 μ M), although initially showing a depression in LVDP of $58 \pm 8\%$ of initial baseline values at 15 min into reperfusion, recovered to $83 \pm 3\%$ of baseline.

In order to establish whether the PKC δ peptide activator produced any direct inotropic effects on cardiac contractile function, Sham I/R hearts were perfused with PKC δ peptide activator (10 μ M). This group served as one of the controls for the study. These hearts did not show any significant change in LVDP (Fig. 20) or $+dP/dt_{\max}$ (Fig. 21) at the end of the 80 min reperfusion period, thus, indicating that at this dose the PKC δ peptide activator had no direct effect on cardiac contractile function.

Figs. 20 and 21 showed the initial and final values for LVDP and $+dP/dt_{\max}$ from isolated perfused hearts respectively. There was no significant difference between the initial baseline values of all the groups studied. There was also no significant difference between the initial and final values of LVDP and $+dP/dt_{\max}$ for the Sham I/R, I/R, Sham I/R+PKC δ peptide activator and I/R+PKC δ peptide activator (10 μ M) groups. However, there was a significant difference between the

initial and final values of LVDP and $+dP/dt_{\max}$ for the I/R+PMN group. A significant decrease ($p<0.01$) from initial baseline of $46\pm9\%$ in LVDP and $45\pm8\%$ in $+dP/dt_{\max}$ at 45 min post-reperfusion was observed.

The presence of the PKC δ peptide activator at a 5 μM and 10 μM dose
5 attenuated the decrease in LVDP and $+dP/dt_{\max}$ associated with the post-ischemic perfusion with PMNs. The 10 μM dose was the most cardioprotective as the hearts in the I/R+PMN+PKC δ peptide activator (10 μM) recovered to $83\pm3\%$ and $79\pm5\%$ of initial baseline at 45 min post-reperfusion for LVDP and $+dP/dt_{\max}$, respectively. These values were significantly different from I/R+PMN at 45 min
10 post-reperfusion ($p<0.01$). The 5 μM dose was also cardioprotective, although not to the same extent as the 10 μM dose, as the hearts in the I/R+PMN+PKC δ peptide activator (5 μM) recovered to $80\pm9\%$ and $67\pm7\%$ for LVDP and $+dP/dt_{\max}$ of initial baseline at 45 min post-reperfusion, respectively. The LVDP values for the 5 μM dose were significantly different from I/R+PMN at 45 min post-reperfusion ($p<0.05$).
15 The 1 μM dose of PKC δ peptide activator was not cardioprotective as the hearts in the I/R+PMN+PKC δ peptide activator (1 μM) group only recovered to $60\pm13\%$ and $51\pm5\%$ for LVDP and $+dP/dt_{\max}$ respectively. The final values of LVDP and $+dP/dt_{\max}$ at the 1 μM dose group were not significantly different from the final values of the I/R+PMN group.

20 A mechanism that may contribute to the cardioprotective effects (i.e. LVDP) of the PKC δ peptide activator may be inhibition of PMN superoxide release. PKC δ peptide activator significantly inhibited superoxide release (i.e. absorbance) from suspensions of PMA-stimulated rat PMNS from 0.49 ± 0.03 to 0.32 ± 0.03 ($p<0.01$), 0.32 ± 0.02 ($p<0.01$), for 5 μM and 10 μM respectively (Fig. 22). There was no
25 significant inhibition of superoxide at the 1 μM dose.

Although certain presently preferred embodiments of the invention have been specifically described herein, it will be apparent to those skilled in the art to which the invention pertains that variations and modifications of the various embodiments shown and described herein may be made without departing from the spirit and scope
5 of the invention. Accordingly, it is intended that the invention be limited only to the extent required by the appended claims and the applicable rules of law.

What is claimed is

1. A solution for perfusion, preservation, and/or re-perfusion for organ preservation comprising at least one peptide inhibitor of protein kinase C β II (PKC β II) and/or at least one peptide inhibitor of protein kinase C ζ (PKC ζ) and/or at least one peptide activator of PKC δ .
2. The solution of claim 1, wherein the peptide inhibitors are dissolved in saline solution.
3. The solution of claim 1, further comprising potassium chloride.
4. The solution of claim 1, wherein the at least one peptide inhibitor of PKC β II is SEQ ID NO: 1.
5. The solution of claim 1, wherein the at least one peptide activator of PKC δ is SEQ ID NO: 3.
6. The solution of claim 1, wherein the at least one peptide inhibitor of PKC ζ is selected from the group consisting of SEQ ID NO: 2 and G86983.
7. The solution of claim 1, wherein the concentration of the at least one peptide inhibitor of PKC β II is about 5-10 μ M.
8. The solution of claim 1, wherein the concentration of the at least one peptide activator of PKC δ is about 5-10 μ M.

9. The solution of claim 1, wherein the concentration of the at least one peptide inhibitor of PKC ζ is about 2.5-5 μ M.
- 5
10. The solution of claim 1, wherein the organ is a heart.
11. The solution of claim 1, wherein the organ is a mammalian organ.
- 10 12. The solution of claim 11, wherein the mammal is human.
13. The solution of claim 1, wherein the organ is preserved for transplantation.
14. The solution of claim 1, wherein the peptide inhibitors/activator are
- 15 myristolated.
15. A method for preserving an organ for transplantation comprising the step of perfusing the organ with the solution of claim 1.
- 20 16. The method of claim 15, wherein the peptide inhibitors/activator are dissolved in saline solution.
17. The method of claim 15, further comprising potassium chloride.

18. The method of claim 15, wherein the at least one peptide inhibitor of PKC β II is SEQ ID NO: 1.
19. The method of claim 15, wherein the at least one peptide activator of PKC δ is
5 SEQ ID NO: 3.
20. The method of claim 15, wherein the at least one peptide inhibitor of PKC ζ is selected from the group consisting of SEQ ID NO: 2 and G66983.
- 10 21. The method of claim 15, wherein the concentration of the at least one peptide inhibitor of PKC β II is about 5-10 μ M.
22. The method of claim 15, wherein the concentration of the at least one peptide activator of PKC δ is about 5-10 μ M.
- 15 23. The method of claim 15, wherein the concentration of the at least one peptide inhibitor of PKC ζ is about 2.5-5 μ M.
24. The method of claim 15, wherein the organ is a heart.
- 20 25. The method of claim 15, wherein the organ is a mammalian organ.
26. The method of claim 25, wherein the mammal is human.
- 25 27. The method of claim 15, wherein the organ is preserved for transplantation.

28. The method of claim 15, wherein the wherein the peptide inhibitors/activators are myristolated.

5 29. The method of claim 15, further comprising the step of submerging the organ in the solution of claim 1.

30. The method of claim 15, wherein the perfusing step takes place at a rate of less than about 20 mL/minute.

10

31. The method of claim 15, wherein the perfusing step takes place at a rate of about 1 mL/minute.

32. The method of claim 15, wherein the perfusing step is a retrograde perfusion.

15

33. The method of claim 15, wherein the perfusing step lasts about 5 minutes.

34. A method for protecting an ischemic organ from damage comprising the step of perfusing the organ with the solution of claim 1.

20

35. The method of claim 34, wherein the peptide inhibitors/activators are dissolved in saline solution.

36. The method of claim 34, further comprising potassium chloride.

25

37. The method of claim 34, wherein the at least one peptide inhibitor of PKC β II is SEQ ID NO: 1.
38. The method of claim 34, wherein the at least one peptide activator of PKC δ is
5 SEQ ID NO: 3.
39. The method of claim 34, wherein the at least one peptide inhibitor of PKC ζ is selected from the group consisting of SEQ ID NO: 2 and G66983.
- 10 40. The method of claim 34, wherein the concentration of the at least one peptide inhibitor of PKC β II is about 5-10 μ M.
41. The method of claim 34, wherein the concentration of the at least one peptide activator of PKC δ is about 5-10 μ M.
- 15 42. The method of claim 34, wherein the concentration of the at least one peptide inhibitor of PKC ζ is about 2.5-5 μ M.
43. The method of claim 34, wherein the organ is a heart.
- 20 44. The method of claim 34, wherein the organ is a mammalian organ.
45. The method of claim 44, wherein the mammal is human.
- 25 46. The method of claim 34, wherein the organ is preserved for transplantation.

47. The method of claim 34, wherein the wherein the peptide inhibitors/activators are myristolated.

5 48. The method of claim 34, further comprising the step of submerging the organ in the solution of claim 1.

49. The method of claim 34, wherein the perfusing step takes place at a rate of less than about 20 mL/minute.

10

50. The method of claim 34, wherein the perfusing step takes place at a rate of about 1 mL/minute.

51. The method of claim 34, wherein the perfusing step is a retrograde perfusion.

15

52. The method of claim 34, wherein the perfusing step lasts about 5 minutes.

53. A method for attenuating organ dysfunction after ischemia comprising the step of perfusing the organ with the solution of claim 1.

20

54. The method of claim 53, wherein the peptide inhibitors/activator are dissolved in saline solution.

55. The method of claim 53, further comprising potassium chloride.

25

56. The method of claim 53, wherein the at least one peptide inhibitor of PKC β II is SEQ ID NO: 1.
57. The method of claim 53, wherein the at least one peptide activator of PKC δ is
5 SEQ ID NO: 3.
58. The method of claim 53, wherein the at least one peptide inhibitor of PKC ζ is selected from the group consisting of SEQ ID NO: 2 and G66983.
- 10 59. The method of claim 53, wherein the concentration of the at least one peptide inhibitor of PKC β II is about 5-10 μ M.
60. The method of claim 53, wherein the concentration of the at least one peptide activator of PKC δ is about 5-10 μ M.
- 15 61. The method of claim 53, wherein the concentration of the at least one peptide inhibitor of PKC ζ is about 2.5-5 μ M.
62. The method of claim 53, wherein the organ is a heart.
- 20 63. The method of claim 53, wherein the organ is a mammalian organ.
64. The method of claim 63, wherein the mammal is human.
- 25 65. The method of claim 53, wherein the organ is preserved for transplantation.

66. The method of claim 53, wherein the wherein the peptide inhibitors/activator are myristolated.

5 67. The method of claim 53, further comprising the step of submerging the organ in the solution of claim 1.

68. The method of claim 53, wherein the perfusing step takes place at a rate of less than about 20 mL/minute.

10

69. The method of claim 53, wherein the perfusing step takes place at a rate of about 1 mL/minute.

70. The method of claim 53, wherein the perfusing step is a retrograde perfusion.

15

71. The method of claim 53, wherein the perfusing step lasts about 5 minutes.

72. A method for maintaining nitric oxide release in an ischemic organ comprising the step of perfusing the organ with the solution of claim 1.

20

73. The method of claim 72, wherein the peptide inhibitors/activator are dissolved in saline solution.

74. The method of claim 72, further comprising potassium chloride.

25

75. The method of claim 72, wherein the at least one peptide inhibitor of PKC β II is SEQ ID NO: 1.
76. The method of claim 72, wherein the at least one peptide inhibitor of PKC δ is
5 SEQ ID NO: 3.
77. The method of claim 72, wherein the at least one peptide inhibitor of PKC ζ is selected from the group consisting of SEQ ID NO: 2 and G66983.
- 10 78. The method of claim 72, wherein the concentration of the at least one peptide inhibitor of PKC β II is about 5-10 μ M.
79. The method of claim 72, wherein the concentration of the at least one peptide activator of PKC δ is about 5-10 μ M.
- 15 80. The method of claim 72, wherein the concentration of the at least one peptide inhibitor of PKC ζ is about 2.5-5 μ M.
81. The method of claim 72, wherein the organ is a heart.
- 20 82. The method of claim 72, wherein the organ is a mammalian organ.
83. The method of claim 82, wherein the mammal is human.
- 25 84. The method of claim 72, wherein the organ is preserved for transplantation.

85. The method of claim 72, wherein the wherein the peptide inhibitors/activator are myristolated.
- 5 86. The method of claim 72, further comprising the step of submerging the organ in the solution of claim 1.
87. The method of claim 72, wherein the perfusing step takes place at a rate of less than about 20 mL/minute.
- 10 88. The method of claim 72, wherein the perfusing step takes place at a rate of about 1 mL/minute.
89. The method of claim 72, wherein the perfusing step is a retrograde perfusion.
- 15 90. The method of claim 72, wherein the perfusing step lasts about 5 minutes.
91. A method for protecting an organ from damage after isolation from the circulatory system comprising the step of perfusing the organ with the solution of
- 20 claim 1.
92. The method of claim 91, wherein the peptide inhibitors/activator are dissolved in saline solution.
- 25 93. The method of claim 91, further comprising potassium chloride.

94. The method of claim 91, wherein the at least one peptide inhibitor of PKC β II is SEQ ID NO: 1.
- 5 95. The method of claim 91, wherein the at least one peptide activator of PKC δ is SEQ ID NO: 3.
96. The method of claim 91, wherein the at least one peptide inhibitor of PKC ζ is selected from the group consisting of SEQ ID NO: 2 and G66983.
- 10 97. The method of claim 91, wherein the concentration of the at least one peptide inhibitor of PKC β II is about 5-10 μ M.
98. The method of claim 91, wherein the concentration of the at least one peptide
15 activator of PKC δ is about 5-10 μ M.
99. The method of claim 91, wherein the concentration of the at least one peptide inhibitor of PKC ζ is about 2.5-5 μ M.
- 20 100. The method of claim 91, wherein the organ is a heart.
101. The method of claim 91, wherein the organ is a mammalian organ.
102. The method of claim 101, wherein the mammal is human.

103. The method of claim 91, wherein the organ is preserved for transplantation.

104. The method of claim 91, wherein the wherein the peptide inhibitors/activator are myristolated.

5

105. The method of claim 91, further comprising the step of submerging the organ in the solution of claim 1.

106. The method of claim 91, wherein the perfusing step takes place at a rate of less
10 than about 20 mL/minute.

107. The method of claim 91, wherein the perfusing step takes place at a rate of about 1 mL/minute.

15 108. The method of claim 91, wherein the perfusing step is a retrograde perfusion.

109. The method of claim 91, wherein the perfusing step lasts about 5 minutes.

SEQUENCE LISTING

<110> Young, Lindon

<120> PERFUSION AND/OR PRESERVATION SOLUTION FOR ORGANS

<130> 072689-00107

<160> 3

<170> PatentIn version 3.3

<210> 1

<211> 9

<212> PRT

<213> Homo sapiens

<400> 1

Ser Leu Asn Pro Glu Trp Asn Glu Thr
1 5

<210> 2

<211> 13

<212> PRT

<213> Homo sapiens

<400> 2

Ser Ile Tyr Arg Arg Gly Ala Arg Arg Trp Arg Lys Leu
1 5 10

<210> 3

<211> 8

<212> PRT

<213> Homo sapiens

<400> 3

Met Arg Ala Ala Glu Asp Pro Met
1 5