

5

TERPHENYL DERIVATIVES FOR TREATMENT OF ALZHEIMER'S DISEASE

The present invention relates to compounds having the general formula (I) with the definitions of X, R₁-R₄ given below, and/or a salt or ester thereof.

10 Furthermore, the invention relates to the use of said compounds for the treatment of Alzheimer's disease and their use for the modulation of γ -secretase activity.

Alzheimer's Disease (AD) is a progressive neurodegenerative disorder marked by loss of memory, cognition, and behavioural stability. AD afflicts 6-10% of the population over
15 age 65 and up to 50% over age 85. It is the leading cause of dementia and the third leading cause of death after cardiovascular disease and cancer. There is currently no effective treatment for AD. The total net cost related to AD in the U.S. exceeds \$100 billion annually.

20 AD does not have a simple etiology, however, it has been associated with certain risk factors including (1) age, (2) family history (3) and head trauma; other factors include environmental toxins and low level of education. Specific neuropathological lesions in the limbic and cerebral cortices include intracellular neurofibrillary tangles consisting of hyperphosphorylated tau protein and the extracellular deposition of fibrillar aggregates of
25 amyloid beta peptides (amyloid plaques). The major component of amyloid plaques are the amyloid beta (A-beta, Abeta or A β) peptides of various lengths. A variant thereof, which is the A β 1-42-peptide (Abeta-42), is believed to be the major causative agent for amyloid formation. Another variant is the A β 1-40-peptide (Abeta-40). Amyloid beta is the proteolytic product of a precursor protein, beta amyloid precursor protein (beta-APP or APP).
30 or APP).

Familial, early onset autosomal dominant forms of AD have been linked to missense mutations in the β -amyloid precursor protein (β -APP or APP) and in the presenilin

5 proteins 1 and 2. In some patients, late onset forms of AD have been correlated with a specific allele of the apolipoprotein E (ApoE) gene, and, more recently, the finding of a mutation in alpha2-macroglobulin, which may be linked to at least 30% of the AD population. Despite this heterogeneity, all forms of AD exhibit similar pathological findings. Genetic analysis has provided the best clues for a logical therapeutic approach
10 to AD. All mutations, found to date, affect the quantitative or qualitative production of the amyloidogenic peptides known as Abeta-peptides ($A\beta$), specifically $A\beta_{42}$, and have given strong support to the "amyloid cascade hypothesis" of AD (Tanzi and Bertram, 2005, Cell 120, 545). The likely link between $A\beta$ peptide generation and AD pathology emphasizes the need for a better understanding of the mechanisms of $A\beta$ production and
15 strongly warrants a therapeutic approach at modulating $A\beta$ levels.

The release of $A\beta$ peptides is modulated by at least two proteolytic activities referred to as β - and γ - secretase cleaving at the N-terminus (Met-Asp bond) and the C-terminus (residues 37-42) of the $A\beta$ peptide, respectively. In the secretory pathway, there is
20 evidence that β -secretase cleaves first, leading to the secretion of s-APP β ($s\beta$) and the retention of a 11 kDa membrane-bound carboxy terminal fragment (CTF). The latter is believed to give rise to $A\beta$ peptides following cleavage by γ -secretase. The amount of the longer isoform, $A\beta_{42}$, is selectively increased in patients carrying certain mutations in a particular protein (presenilin), and these mutations have been correlated with early-onset familial Alzheimer's disease. Therefore, $A\beta_{42}$ is believed by many researchers to
25 be the main culprit of the pathogenesis of Alzheimer's disease.

It has now become clear that the γ -secretase activity cannot be ascribed to a single particular protein, but is in fact associated with an assembly of different proteins.

The gamma-secretase activity resides within a multiprotein complex containing at least
30 four components: the presenilin (PS) heterodimer, nicastrin, aph-1 and pen-2. The PS heterodimer consists of the amino- and carboxyterminal PS fragments generated by endoproteolysis of the precursor protein. The two aspartates of the catalytic site are at the interface of this heterodimer. It has recently been suggested that nicastrin serves as a gamma-secretase-substrate receptor. The functions of the other members of gamma-

5 secretase are unknown, but they are all required for activity (Steiner, 2004. Curr. Alzheimer Research 1(3): 175-181).

Thus, although the molecular mechanism of the second cleavage-step has remained elusive until present, the γ -secretase-complex has become one of the prime targets in the search for compounds for the treatment of Alzheimer's disease.


10 Various strategies have been proposed for targeting gamma-secretase in Alzheimer's disease, ranging from targeting the catalytic site directly, developing substrate-specific inhibitors and modulators of gamma-secretase activity (Marjaux et al., 2004. Drug Discovery Today: Therapeutic Strategies, Volume 1, 1-6). Accordingly, a variety of compounds were described that have secretases as targets (Larner, 2004. Secretases as
15 therapeutics targets in Alzheimer's disease: patents 2000 – 2004. Expert Opin. Ther. Patents 14, 1403-1420.)

Indeed, this finding was recently supported by biochemical studies in which an effect of certain NSAIDs on γ -secretase was shown (Weggen et al (2001) Nature 414, 6860, 212 and WO 01/78721 and US 2002/0128319; Morihara et al (2002) J. Neurochem. 83, 1009;
20 Eriksen (2003) J. Clin. Invest. 112 , 440). Potential limitations for the use of NSAIDs to prevent or treat AD are their inhibition activity of Cox enzymes, which can lead to unwanted side effects, and their low CNS penetration (Peretto et al., 2005, J. Med. Chem. 48, 5705-5720).

Thus, there is a strong need for novel compounds which modulate γ -secretase activity
25 thereby opening new avenues for the treatment of Alzheimer's disease.

The object of the present invention is to provide such compounds.

The invention is directed to novel compounds of Formula I wherein:



Formula I

5

X is a bond or a group $-CR_5R_6$ wherein R_5 and R_6 are, independently of each other, selected from the group consisting of H; alkyl selected from the group CH_3 , C_2H_5 , $i-C_3H_7$, $n-C_3H_7$, $i-C_4H_9$, $n-C_4H_9$, $sec-C_4H_9$, $tert-C_4H_9$; alkenyl selected from C_2H_3 , $i-C_3H_5$, $n-C_3H_5$, $n-C_4H_7$, $i-C_4H_7$, $sec-C_4H_7$; wherein in any of the alkyl or alkenyl groups one or more H atoms optionally can be substituted with one or more substituents independently selected from the group consisting of OH, F, Cl, Br, I and CF_3 ; or R_5 and R_6 being part of a ring, either saturated or unsaturated, substituted or unsubstituted, having 3 to 6 C-atoms, and which may contain in the ring one or more heteroatoms from the group N, S or O, and which heteroatom may be identical or different if more than one heteroatom is present;

15

R_1 , R_2 , R_3 and R_4 are independently selected from the group consisting of H; F; Cl; Br; I; CN; OH; $C(O)N(R_7R_8)$; $S(O)_2R_7$; $SO_2N(R_7R_8)$; $S(O)N(R_7R_8)$; $N(R_7)S(O)_2R_8$; $N(R_8)S(O)R_8$; $S(O)_2R_7$; $N(R_7)S(O)_2N(R_8R_{8a})$; SR_7 ; $N(R_7R_8)$; $N(R_7)C(O)R_8$; $N(R_7)C(O)N(R_8R_{8a})$; $N(R_7)C(O)OR_8$; $OC(O)N(R_7R_8)$; $C(O)R_7$; substituted and unsubstituted C_1 - C_4 -alkyl and substituted and unsubstituted C_1 - C_4 -alkoxy, and wherein the substituents of both groups C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy are selected from OH, F, Cl, Br, I, CF_3 ;

25

R_7 , R_8 , R_{8a} are independently selected from the group consisting of H; C_1 - C_4 -alkyl; heterocyclyl; and C_{3-7} cycloalkyl, wherein C_1 - C_4 -alkyl; heterocyclyl; and C_{3-7} cycloalkyl

- 5 optionally can be substituted with one or more substituents independently selected from the group consisting of OH, F, Cl, Br, I and CF₃;

R₉, and R₁₀ are independently H, F, or CF₃;

- 10 Y is a carboxy group –C(O)OH or a substituted or unsubstituted tetrazole group and solvates, hydrates, esters, and pharmaceutically acceptable salts thereof.

The term “substituted” as used herein includes both part and full substitution. Substituents can be either saturated or unsaturated.

15

In case R₅ and R₆ are part of a ring, the ring can be substituted by C₁-C₄-alkyl or OH, F, Cl, Br, I and CF₃

- Esters are those according to formula (I) in which H of the carboxy group is replaced by
20 an organic residue R_{7a}. Suitable organic residues are known to a person skilled in the art. Preferred R_{7a} include the following: an unsubstituted or at least monosubstituted alkyl, preferably a C₁-C₁₀ alkyl, an alkenyl, preferably C₂-C₁₀-alkenyl, an alkynyl, preferably C₃-C₁₀-alkynyl, and an unsubstituted or at least monosubstituted, saturated or unsaturated,
25 one or more heteroatoms from the group N, S or O, and which heteroatom may be identical or different if more than one heteroatom is present. Said substituents being selected from the group consisting of halogen, alkyl, alkenyl, alkynyl, N, S, O, carboxy, sulphonyl, and the like and which can be further substituted.

- 5 Examples for current aromatic groups include aryl groups, for example phenyl groups, and heteroaryl groups, which aryl and heteroaryl groups may be substituted, preferably by the substituents given above.

In another embodiment of the invention:

- X is group $-CR_5R_6$ wherein R_5 and R_6 are, independently of each other, selected from the
10 group consisting of H; alkyl selected from the group CH_3 , C_2H_5 , $i-C_3H_7$, $n-C_3H_7$, $i-C_4H_9$, $n-C_4H_9$, $sec-C_4H_9$, $tert-C_4H_9$; wherein in the all named alkyl groups one or more H atoms optionally can be substituted with one or more substituents independently selected from the group consisting of OH, F, Cl, Br and I; or R_5 , R_6 jointly form together with the carbon atom to which they are attached a cyclopropyl ring; and/or
- 15 R_1 , R_2 , R_3 and R_4 are independently selected from the group consisting of H, OH, $C_{(1-4)}$ alkyl, $C_{(1-4)}$ alkoxy, $-N(CH_3)_2$, $-SO_2CH_3$, CN, OCF_3 , $-C(O)CH_3$, OCH_3 , CF_3 , F, and Cl; wherein said $C_{(1-4)}$ alkyl and $C_{(1-4)}$ alkoxy optionally can be independently substituted with one, two, or three substituents selected from the group consisting of OH, I, Br, F, and Cl; and/or
- 20 R_5 and R_6 being H; or R_5 being H and R_6 being CH_3 , C_2H_5 , C_3H_7 or C_4H_9 or isomers thereof; or R_1 and R_2 being CH_3 or R_5 , R_6 jointly form together with the carbon atom to which they are attached a cyclopropyl ring; and/or

R_9 , and R_{10} are independently H, F, or CF_3 ;

Y is a carboxy group

- 25 and solvates, hydrates, esters, and pharmaceutically acceptable salts thereof.

In another embodiment of the invention:

X is a group $-CR_5R_6$ with R_5 and R_6 being H; or R_5 being H and R_6 being CH_3 , C_2H_5 , C_3H_7 or C_4H_9 or isomers thereof; and/or

- R_1 , R_2 , R_3 and R_4 are independently selected from the group consisting of H, OH, $C_{(1-4)}$ alkyl, $C_{(1-4)}$ alkoxy, $-N(CH_3)_2$, $-SO_2CH_3$, CN, OCF_3 , $-C(O)CH_3$, OCH_3 , CF_3 , F, and Cl;
30 wherein said $C_{(1-4)}$ alkyl and $C_{(1-4)}$ alkoxy can optionally be independently substituted with one, two, or three substituents selected from the group consisting of OH, I, Br, F, and Cl;

5 R₉, and R₁₀ are independently H, F, or CF₃; and/or

and/or

Y is a carboxy group;

and solvates, hydrates, esters, and pharmaceutically acceptable salts thereof.

In another embodiment of the invention:

10 X is a group -CR₅R₆, with R₅ and R₆ being H; or R₅ being H and R₆ being CH₃, C₂H₅, C₃H₇ or C₄H₉ or isomers thereof;


Y is a carboxy group;

R₉, and R₁₀ are independently H, F, or CF₃;

15 R₁, R₂, R₃ and R₄ are independently selected from the group consisting of H, OH, C₍₁₋₄₎alkyl, -N(CH₃)₂, -SO₂CH₃, CN, OCF₃, -C(O)CH₃, OCH₃, CF₃, F, and Cl;

and solvates, hydrates, esters, and pharmaceutically acceptable salts thereof.

Another embodiment of the invention is directed to compounds of Formula I*



Formula I*

20

X is a bond or a group -CR₅R₆ wherein R₅ and R₆ are, independently of each other, selected from the group consisting of H; alkyl selected from the group CH₃, C₂H₅, i-C₃H₇, n-C₃H₇, i-C₄H₉, n-C₄H₉, sec-C₄H₉, tert-C₄H₉; alkenyl selected from C₂H₃, i-C₃H₅, n-C₃H₅, n-C₄H₇, i-C₄H₇, sec-C₄H₇; wherein in the alkyl or alkenyl groups one or more H

- 5 atoms optionally can be substituted with one or more substituents independently selected from the group consisting of OH, F, Cl, Br, I and CF₃; or R₅ and R₆ being part of a ring, either saturated or unsaturated, substituted or unsubstituted, having 3 to 6 C-atoms, and which may contain in the ring one or more heteroatoms from the group N, S or O, and which heteroatom may be identical or different if more than one heteroatom is present;
- 10 R₁, R₂, R₃ and R₄ are independently selected from the group consisting of H; F; Cl; Br; I; CN; OH; C(O)N(R₇R₈); S(O)₂R₇; SO₂N(R₇R₈); S(O)N(R₇R₈); N(R₇)S(O)₂R₈; N(R₈)S(O)R₈; S(O)₂R₇; N(R₇)S(O)₂N(R₈R_{8a}); SR₇; N(R₇R₈); N(R₇)C(O)R₈; N(R₇)C(O)N(R₈R_{8a}); N(R₇)C(O)OR₈; OC(O)N(R₇R₈); C(O)R₇; substituted and unsubstituted C₁-C₄-alkyl and substituted and unsubstituted C₁-C₄-alkoxy, and wherein
- 15 the substituents of both groups C₁-C₄-alkyl and C₁-C₄-alkoxy are selected from OH, F, Cl, Br, I, CF₃;

- R₇, R₈, R_{8a} are independently selected from the group consisting of H; C₁-C₄-alkyl; heterocyclyl; and C₃₋₇ cycloalkyl, wherein C₁-C₄-alkyl; heterocyclyl; and C₃₋₇ cycloalkyl optionally can be substituted with one or more substituents independently selected from
- 20 the group consisting of OH, F, Cl, Br, I and CF₃;

Y is a carboxy group -C(O)OH or a substituted or unsubstituted tetrazole group and/or a salt or ester thereof.

- The term "substituted" as used herein includes both part and full substitution.
- 25 Substituents can be either saturated or unsaturated.

In case R₅ and R₆ are part of a ring, the ring can be substituted by C₁-C₄-alkyl or OH F, Cl, Br, I and CF₃

- 30 Esters are those according to formula (I) in which H of the carboxy group is replaced by an organic residue R_{7a}. Suitable organic residues are known to a person skilled in the art. Preferred R_{7a} include the following:

5

An unsubstituted or at least monosubstituted alkyl, preferably a C₁-C₁₀ alkyl, an alkenyl, preferably C₂-C₁₀-alkenyl, an alkynyl, preferably C₃-C₁₀-alkynyl, and an unsubstituted or at least monosubstituted, saturated or unsaturated, non-aromatic or aromatic ring having 3 to 6 C-atoms, and which may contain in the ring one or more heteroatoms from the group N, S or O, and which heteroatom may be identical or different if more than one heteroatom is present. Said substituents being selected from the group consisting of halogen, alkyl, alkenyl, alkynyl, N, S, O, carboxy, sulphonyl, and the like and which can be further substituted.

15 Examples for current aromatic groups include aryl groups, for example phenyl groups, and heteroaryl groups, which aryl and heteroaryl groups may be substituted, preferably by the substituents given above.

The term "C₁-C₄-alkyl" refers to methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and tert-butyl.

"C₃₋₇ cycloalkyl" or "C₃₋₇ cycloalkyl ring" means a cyclic alkyl chain having 3 - 7 carbon atoms, e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexenyl, cycloheptyl. Each hydrogen of a cycloalkyl carbon may be replaced by a substituent.

25

"Heterocyclyl" or "heterocycle" means a cyclopentane, cyclohexane or cycloheptane ring that may contain up to the maximum number of double bonds (aromatic or non-aromatic ring which is fully, partially or un-saturated) wherein at least one carbon atom up to 4 carbon atoms are replaced by a heteroatom selected from the group consisting of sulfur (including -S(O)-, -S(O)₂-), oxygen and nitrogen (including =N(O)-) and wherein the ring is linked to the rest of the molecule via a carbon or nitrogen atom. Examples for a heterocycle include but are not restricted to furan, thiophene, pyrrole, pyrroline,

5 imidazole, imidazoline, pyrazole, pyrazoline, oxazole, oxazoline, isoxazole, isoxazoline, thiazole, thiazoline, isothiazole, isothiazoline, thiadiazole, thiadiazoline, tetrahydrofuran, tetrahydrothiophene, pyrrolidine, imidazolidine, pyrazolidine, oxazolidine, isoxazolidine, thiazolidine, isothiazolidine, thiadiazolidine, sulfolane, pyran, dihydropyran, tetrahydropyran, imidazolidine, pyridine, pyridazine, pyrazine, pyrimidine, piperazine,
 10 piperidine, morpholine, tetrazole, triazole, triazolidine, tetrazolidine, azepine or homopiperazine. "Heterocycle" means also azetidine.

In preferred embodiments, the invention relates to a compound having the general formula (I) wherein X; Y; R₁ and R₂; and R₃, R₄, R₅ and R₆ independently of each other have the following meanings:

- 15 X is a group -CR₅R₆ wherein R₅ and R₆ are, independently of each other, selected from the group consisting of H; alkyl selected from the group CH₃, C₂H₅, i-C₃H₇, n-C₃H₇, i-C₄H₉, n-C₄H₉, sec-C₄H₉, tert-C₄H₉; wherein in the all named alkyl groups one or more H atoms optionally can be substituted with one or more substituents independently selected from the group consisting of OH, F, Cl, Br and I; and/or
- 20 R₁, R₂, R₃ and R₄ are independently selected from the group consisting of H; OH; C₁-C₄-alkyl or C₁-C₄-alkoxy, substituted partly or fully by OH, F, Cl, Br, I; and/or
- R₅ and R₆ being H; or R₅ being H and R₆ being CH₃, C₂H₅, C₃H₇ or C₄H₉ or isomers thereof; or R₁ and R₂ being CH₃ or R₅, R₆ jointly form together with the carbon atom to which they are attached a cyclopropyl ring; and/or
- 25 Y is a carboxy group
- and/or a salt or ester thereof.

Within this group of embodiments, it is even more preferred if all the groups X; Y; R₁, R₂, R₃, R₄, R₅ and R₆ have the meanings defined beforehand.

It is even more preferred if X; Y; R₁ and R₂; and R₃, R₄, R₅ and R₆ independently of each
 30 other have the following meanings:

5 X is a group $-\text{CR}_5\text{R}_6$ with R_5 and R_6 being H; or R_5 being H and R_6 being CH_3 , C_2H_5 , C_3H_7 or C_4H_9 or isomers thereof; or R_5 and R_6 being CH_3 or R_5 , R_6 jointly form together with the carbon atom to which they are attached a cyclopropyl ring; and/or

R_1 , R_2 , R_3 and R_4 are independently selected from the group consisting of H; OH; C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, substituted partly or fully by OH, F, Cl, Br, I; and/or

10 and/or

Y is a carboxy group

and/or a salt or ester thereof.

Within this group of embodiments, it is even more preferred if all the groups X; Y; R_1 , R_2 , R_3 , R_4 , R_5 and R_6 have the meanings defined beforehand.

15

It is still more preferred if X; Y; R_1 and R_2 ; and R_3 , R_4 , R_5 and R_6 independently of each other have the following meanings:

X is a group $-\text{CR}_5\text{R}_6$, with R_5 and R_6 being H; or R_5 being H and R_6 being CH_3 , C_2H_5 , C_3H_7 or C_4H_9 or isomers thereof;

20 Y is a carboxy group

R_1 , R_2 , R_3 and R_4 are independently selected from the group consisting of H, OH, CH_3 , OCH_3 , CF_3 , F, and Cl; and/or

and/or a salt or ester thereof.

25 Within this group of embodiments, it is even more preferred if all the groups X; Y; R_1 and R_2 ; and R_3 , R_4 , R_5 and R_6 have the meanings defined beforehand.

In an even more preferred embodiment, the invention relates to compounds selected from the group consisting of

30 (i) 4''-Chloro-4-trifluoromethyl-[1,1';3,1'']terphenyl-5'-yl)-acetic acid






- 5 (ii) (4"-Trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid
- (iii) (3-Chloro-4"-trifluoromethyl [1,1';3',1'']terphenyl-5'-yl)-acetic acid
- (iv) (4-Hydroxy-4"-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid
- (v) (4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid
- (vi) [1,1';3',1'']Terphenyl-5'-yl-acetic acid
- 10 (vii) (4,4"-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid
- (viii) (4,4"-Difluoro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid
- (ix) (3,3"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid
- (x) (3,3"-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid
- (xi) (4,4"-Dimethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid
- 15 (xii) (4,4"-Dimethoxy-[1,1';3',1'']terphenyl-5'-yl)-acetic acid
- (xiii) 2-(4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
- (xiv) (R)-2-(4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
- (xv) (S)-2-(4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
- (xvi) 4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-carboxylic acid
- 20 (xvii) 5-(4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-1H-tetrazole






and solvates, hydrates, esters, and pharmaceutically acceptable salts thereof.

Some of the compounds of the inventions and/or salts or esters thereof will exist in different stereoisomeric forms. All of these forms are subjects of the invention.

25






In another embodiment, the invention relates to compounds selected from the group consisting of

Compound #	Structure	Chemical Name
1		2-(3,5-Difluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methylpentanoic acid
2		2-(2,4-Difluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methylpentanoic acid
3		2-(4-Chloro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methylpentanoic acid
4		2-(4-Isopropyl-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methylpentanoic acid
5		2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methylpentanoic acid





6		2-(2,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
7		4-Methyl-2-(3,5,4''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
8		2-(2-Fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
9		2-(3-Fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
10		2-(4-Fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid

11		4-Methyl-2-(4-methyl-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
12		2-(3,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
13		2-(4-Chloro-3-fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
14		4-Methyl-2-(3,4,5-trifluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
15		2-(3,5-Dichloro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid






16		2-(4-Cyano-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methylpentanoic acid
17		2-(3-Cyano-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methylpentanoic acid
18		2-(4-Cyano-3-fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methylpentanoic acid
19		2-(3-Cyano-4-fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methylpentanoic acid
20		4-Methyl-2-(4-trifluoromethoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid





21		2-(4-Methanesulfonyl-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
22		2-(4-Chloro-3,4''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
23		2-(2,5-Dichloro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
24		2-(4-Methoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
25		2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pent-4-enoic acid




26		2-(3-Fluoro-4-trifluoromethoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
27		2-(3-Fluoro-4-trifluoromethoxy-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
28		2-(4-Chloro-3-fluoro-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
29		2-(3''-Fluoro-3,5,5''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
30		2-(4''-Chloro-3,5,3''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid

31		2-(3,5-Difluoro-4''-chloro-3''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
32		4-Methyl-2-(3,5,3''-trifluoro-5''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
33		2-(3,5-Difluoro-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
34		2-(3-Cyano-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl pentanoic acid.

35		4-Methyl-2-(3,5,4''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
36		2-(5-Cyano-3-fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methylpentanoic acid
37		2-(4-Cyano-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methylpentanoic acid.
38		4-Methyl-2-(3,5,3'',5''-tetrafluoro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
39		(R)-2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methylpentanoic acid

40		(S)-2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
41		(S)-2-(3,5-Difluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
42		(R)-2-(3,5-Difluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
43		(S)-4-Methyl-2-(3,5,4''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
44		(R)-4-Methyl-2-(3,5,4''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid

45		(R)-2-(4-Chloro-4''-trifluoromethyl-[1,1';3'1'']terphenyl-5'-yl)-4-methylpentanoic acid
46		(R)-2-(4-Isopropyl-4''-trifluoromethyl-[1,1';3'1'']terphenyl-5'-yl)-4-methylpentanoic acid
47		(R)-2-(4-Chloro-3-fluoro-4''-trifluoromethyl-[1,1';3'1'']terphenyl-5'-yl)-4-methylpentanoic acid
48		2-(4-Acetyl-4''-trifluoromethyl-[1,1';3'1'']terphenyl-5'-yl)-4-methylpentanoic acid

49		2-(3-Dimethylamino-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
50		(S)-2-(4-Chloro-3-fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid
51		(R)-4-Methyl-2-(4-trifluoromethoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
52		(S)-4-Methyl-2-(4-trifluoromethoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid

5

and solvates, hydrates, esters, and pharmaceutically acceptable salts thereof.

Described below are exemplary salts of the compounds according to the invention which are included herein. The list of the different salts stated below is not meant to be complete and limiting.

10

Compounds according to the invention which contain one or more acidic groups can be used according to the invention, e.g. as their alkali metal salts, alkaline earth metal salts

5 or ammonium salts. More precise examples of such salts include sodium salts, potassium salts, calcium salts, magnesium salts or salts with ammonia or organic amines such as, e.g. ethylamine, ethanolamine, triethanolamine or amino acids.

Compounds according to the invention which contain one or more basic groups, i.e.
10 groups which can be protonated, can be used according to the invention in the form of their addition salts with inorganic or organic acids.

Examples for suitable acids include hydrogen chloride, hydrogen bromide, phosphoric acid, sulfuric acid, nitric acid, methanesulfonic acid, p-toluenesulfonic acid,
15 naphthalenedisulfonic acid, oxalic acid, acetic acid, tartaric acid, lactic acid, salicylic acid, benzoic acid, formic acid, propionic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, malic acid, sulfamic acid, phenylpropionic acid, gluconic acid, ascorbic acid, isonicotinic acid, citric acid, adipic acid and other acids known to a person skilled in the art.

20

The term “pharmaceutically acceptable” means approved by a regulatory agency such as the EMEA (Europe) and/or the FDA (US) and/or any other national regulatory agency for use in animals, preferably in humans.

25 Compounds according to the invention which contain several basic groups can simultaneously form different salts.

If a compound according to the invention simultaneously contains acidic and basic groups in the molecule, the invention also includes, in addition to the salt forms
30 mentioned, inner salts or betaines.

- 5 The respective salts of the compounds according to the invention can be obtained by customary methods which are known to the person skilled in the art, for example by contacting these with an organic or inorganic acid or base in a solvent or dispersant, or by anion exchange or cation exchange with other salts.
- 10 Furthermore, the invention includes all salts of the compounds according to the invention which, owing to low physiological compatibility, are not directly suitable for use in pharmaceuticals but which can be used, for example, as intermediates for chemical reactions or for the preparation of pharmaceutically acceptable salts or which might be suitable for studying γ -secretase modulating activity of a compound according of the
- 15 invention in any suitable manner, such as any suitable in vitro assay.

The present invention furthermore includes all solvates of the compounds according to the invention.

- 20 The present invention furthermore includes derivatives/prodrugs (including the salts thereof) of the compounds according to the invention which contain physiologically tolerable and cleavable groups and which are metabolized in animals, preferably mammals, most preferably humans into a compound according to the invention.
- 25 The present invention furthermore includes the metabolites of the compounds according to the invention.

The term "metabolites" refers to all molecules derived from any of the compounds according to the invention in a cell or organism, preferably mammal.

- 5 Preferably the term “metabolites” relates to molecules which differ from any molecule which is present in any such cell or organism under physiological conditions.

The structure of the metabolites of the compounds according to the invention will be obvious to any person skilled in the art, using the various appropriate methods.

10

The compounds according to general formula (I) can be prepared according to methods published in the literature or by analogous methods.

- 15 Depending on the circumstances of the individual case, in order to avoid side reactions during the synthesis of a compound of the general formula (I), it can be necessary or advantageous to temporarily block functional groups by introducing protective groups and to deprotect them in a later stage of the synthesis, or to introduce functional groups in the form of precursor groups and at a later stage to convert them into the desired functional groups. Suitable synthetic strategies, protective groups and precursor groups
20 are known to the person skilled in the art.

- If desired, the compounds of the formula (I) can be purified by customary purification procedures, for example by recrystallization or chromatography. The starting materials for the preparation of the compounds of the formula (I) are commercially available or can
25 be prepared according to or analogously to literature procedures.

These can serve as a basis for the preparation of the other compounds according to the invention by several methods well known to the person skilled in the art.

- 30 The invention also relates to a compound of the invention for use as a medicament. The compounds are as defined above, furthermore with respect to the medicament the

5 embodiments as described below with respect to the use of the invention, e.g. formulation, application and combination, also apply to this aspect of the invention.

In particular the compounds according to the invention are suitable for the treatment of Alzheimer's disease.

10

Details relating to said use are further disclosed below.

The compounds can be used for modulation of γ -secretase activity.

15 As used herein, the term "modulation of γ -secretase activity" refers to an effect on the processing of APP by the γ -secretase-complex. Preferably it refers to an effect in which the overall rate of processing of APP remains essentially as without the application of said compounds, but in which the relative quantities of the processed products are changed, more preferably in such a way that the amount of the A β 42-peptide produced is
20 reduced. For example a different Abeta species can be produced (e.g. Abeta-38 or other Abeta peptide species of shorter amino acid sequence instead of Abeta-42) or the relative quantities of the products are different (e.g. the ratio of Abeta-40 to Abeta-42 is changed, preferably increased).

Gamma secretase activity can e.g. be measured by determining APP processing, e.g. by
25 determining the levels of Abeta peptide species produced, most importantly levels of Abeta-42 (see Example section, *infra*).

It has been previously shown that the γ -secretase complex is also involved in the processing of the Notch-protein. Notch is a signalling protein which plays a crucial role
30 in developmental processes (e.g. reviewed in Schweisguth F (2004) Curr. Biol. 14, R129).

5 With respect to the use of said compounds for the modulation of γ -secretase activity in therapy, it seems particularly advantageous not to interfere with the Notch-processing activity of the γ -secretase activity in order to avoid putative undesired side-effects.

Thus, compounds are preferred which do not show an effect on the Notch-processing activity of the γ -secretase-complex.

10

Within the meaning of the invention, "effect on the Notch processing activity" includes both an inhibition or an activation of the Notch-processing activity by a certain factor.

A compound is defined as not having an effect on the Notch processing activity, if said factor is smaller than 20, preferably smaller than 10, more preferably smaller than 5, most
15 preferably smaller than 2 in the respective assay as described in Shimizu et al (2000) Mol. Cell. Biol, 20: 6913 at a concentration of 30 μ M.

Such a γ -secretase modulation can be carried out, e.g. in animals such as mammals. Exemplary mammals are mice, rats, guinea pigs, monkeys, dogs, cats. The modulation can also be carried out in humans. In a particular embodiment of the invention, said
20 modulation is performed in vitro or in cell culture. As known to the person skilled in the art, several in vitro and cell culture assays are available.

Exemplary assays useful for measuring the production of C-terminal APP fragments in cell lines or transgenic animals by Western blot analysis include but are not limited to
25 those described in Yan et al., 1999, Nature 402, 533-537.

An example of an in vitro γ -secretase assay is described in WO-03/008635. In this assay a suitable peptide substrate is contacted with a γ -secretase preparation and the ability to cleave the substrate is measured.

30

5 Concentrations of the various products of the γ -secretase cleavage (the A β -peptides) can be determined by various methods known to a person skilled in the art. Examples for such methods include determination of the peptides by mass-spectrometry or detection by antibodies.

10 Exemplary assays useful for the characterization of the profile of soluble Abeta peptides in cultured cell media and biological fluids include but are not limited to those described by Wang et al., 1996, J. Biol. Chem. 271, 31894-31902. In this assay a combination of immunoprecipitation of Abeta-peptides with specific antibodies and detection and quantification of the peptide species with matrix-assisted laser desorption ionization time-
15 of-flight mass spectrometry is used.

Exemplary assays useful for measuring the production of Abeta-40 and Abeta-42 peptides by ELISA include but are not limited to those described in Vassar et al, 1999, Science 286, 735-741. Further information is disclosed for example in N. Ida et al. (1996)
20 J. Biol. Chem. 271, 22908, and M. Jensen et al. (2000) Mol. Med. 6, 291. Suitable antibodies are available for example from The Genetics Company, Inc., Switzerland. Antibody-based kits are also available from Innogenetics, Belgium.

Cells which can be employed in such assays include cells which endogenously express
25 the γ -secretase complex and transfected cells which transiently or stably express some or all interactors of the γ -secretase complex. Numerous available cell lines suitable for such assays are known to the skilled person. Cells and cell lines of neuronal or glial origin are particularly suitable. Furthermore, cells and tissues of the brain as well as homogenates and membrane preparations thereof may be used (Xia et al., 1998, Biochemistry 37,
30 16465-16471).

- 5 Such assays might be carried out for example to study the effect of the compounds according to the invention in different experimental conditions and configurations.

Furthermore, such assays might be carried out as part of functional studies on the γ -secretase complex.

- 10 For example, either one or more interactors (either in their wild-type form or carrying certain mutations and/or modifications) of the γ -secretase complex of an animal, preferably a mammal, more preferably humans, might be expressed in certain cell lines and the effect of the compounds according to the invention might be studied.

- 15 Mutated forms of the interactor(s) used can either be mutated forms which have been described in certain animals, preferably mammals, more preferably humans or mutated forms which have not previously been described in said animals.

- 20 Modifications of the interactors of the γ -secretase complex include both any physiological modification of said interactors and other modifications which have been described as modifications of proteins in a biological system.

Examples of such modifications include, but are not limited to, glycosylation, phosphorylation, prenylation, myristylation and farnesylation.

25

Furthermore, the compounds according to the invention can be used for the preparation of a medicament for the modulation of γ -secretase activity.

- 30 The invention further relates to the use of said compounds for the preparation of a medicament for the modulation of γ -secretase activity.

5

The activity of the γ -secretase can be modulated in different ways, i.e. resulting in different profiles of the various A β -peptides.

10 Uses of a compound for the modulation of γ -secretase activity resulting in a decrease in the relative amount of A β 42-peptides produced are preferred.

Respective dosages, routes of administration, formulations etc are disclosed further below.

15 The invention further relates to the use of the compounds according to the invention for the treatment of a disease associated with an elevated level of A β 42-production. The disease with elevated levels of Abeta peptide production and deposition in the brain is typically Alzheimer's disease (AD), cerebral amyloid angiopathy, multi-infarct dementia, dementia pugilistica or Down syndrome, preferably AD.

20 As used herein, the term "treatment" is intended to refer to all processes, wherein there may be a slowing, interrupting, arresting, or stopping of the progression of a disease, but does not necessarily indicate a total elimination of all symptoms.

As used herein, the term "elevated level of A β 42-production" refers to a condition in which the rate of production of A β 42-peptide is increased due to an overall increase in
25 the processing of APP or, preferably, it refers to a condition in which the production of the A β 42 peptide is increased due to a modification of the APP-processing profile in comparison to the wild-type APP and non-pathological situation.

30 As outlined above, such an elevated A β 42-level is a hallmark of patients developing or suffering from Alzheimer's disease.

- 5 One advantage of the compounds or a part of the compounds of the present invention may lie in their enhanced CNS-penetration.

Furthermore the invention relates to a pharmaceutical composition comprising a compound according to the invention in a mixture with an inert carrier.

10

In a preferred embodiment, the invention relates to a pharmaceutical composition comprising a compound according to the invention in a mixture with an inert carrier, where said inert carrier is a pharmaceutical carrier.

- 15 The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the compound is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, including but not limited to peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered orally.
- 20 Saline and aqueous dextrose are preferred carriers when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions are preferably employed as liquid carriers for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatine, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride,
- 25 dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsions, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers
- 30 such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such

5 compositions will contain a therapeutically effective amount of the compound, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

10 Furthermore, the invention relates to methods for the preparation of a compound according to the invention.

In one embodiment for the preparation of a compound according to the present invention, a dibromofluorobenzene can be treated with a benzyl alcohol in the presence of an alkali metal hydride, typically sodium hydride, in a suitable aprotic solvent such as tetrahydrofuran. The product can be treated with a suitable malonic acid derivative, such as malonic acid tert-butyl ester ethyl ester in the presence of an alkali metal hydride, typically sodium hydride and a metal halide, typically a copper halide, preferably copper bromide. Further treatment in an acidic solvent such as acetic acid at elevated temperature provides a benzyloxy-bromophenylacetic acid ester. This can be coupled to a boronic acid under the variety of conditions known to those skilled in the art for such Suzuki coupling, typically using solvents such as 1,2-dimethoxyethane and water, an alkali metal carbonate such as potassium carbonate, and a palladium compound such as tetrakis(triphenylphosphine)palladium (0).

Removal of the benzyl ether protecting group can be achieved under the variety of conditions known to those skilled in the art for such deprotections, typically using a palladium catalyst such as 10% palladium on charcoal in a suitable solvent, such as ethanol, and under an atmosphere of hydrogen.

The resultant hydroxy compound can be converted to a triflate using, e.g., trifluoromethanesulphonic anhydride, an organic base such as pyridine and in a suitable solvent such as dichloromethane. This triflate can then be coupled to a boronic acid under the variety of conditions known to those skilled in the art for such Suzuki coupling, typically using solvents such as 1,2-dimethoxyethane and water, an alkali metal carbonate

5 such as potassium carbonate, and a palladium compound such as bis(tri-tert-butylphosphine)palladium (0).

If required the triphenyl carboxylic acid can be alkylated by treatment in a suitable aprotic solvent such as tetrahydrofuran with a suitable base such as a metal alkylamide, typically LDA, and the appropriate halide at a suitable temperature, typically -78°C

10 Conversion of the ester to the acid can be done using a base such as an alkali metal hydroxide, typically sodium hydroxide in the presence of water and other suitable solvents such as ethanol.

In another embodiment a dihydroxyphenylacetic acid derivative can be converted to a bis-triflate using eg trifluoromethanesulphonic anhydride, an organic base such as
15 pyridine and in a suitable solvent such as dichloromethane. This triflate can then be coupled to a boronic acid under the variety of conditions known to those skilled in the art for such Suzuki coupling, typically using solvents such as 1,2-dimethoxyethane and water, an alkali metal carbonate such as potassium carbonate, and a palladium compound such as bis(tri-tert-butylphosphine)palladium (0).

20 If required the triphenyl carboxylic acid can be alkylated by treatment in a suitable aprotic solvent such as tetrahydrofuran with a suitable base such as a metal alkylamide, typically LDA, and the appropriate halide at a suitable temperature, typically -78°C

Conversion of the ester to the acid can be done using a base such as an alkali metal hydroxide, typically sodium hydroxide in the presence of water and other suitable
25 solvents such as ethanol.

In another embodiment a dihydroxybenzonitrile can be converted to a bis-triflate using eg trifluoromethanesulphonic anhydride, an organic base such as pyridine and in a suitable solvent such as dichloromethane. This triflate can then be coupled to a boronic acid under the variety of conditions known to those skilled in the art for such Suzuki coupling,
30 typically using solvents such as 1,2-dimethoxyethane and water, an alkali metal carbonate such as potassium carbonate, and a palladium compound such as bis(tri-tert-butylphosphine)palladium (0).

5 Hydrolysis of the nitrile to the acid can be done using a base such as an alkali metal hydroxide, typically sodium hydroxide in the presence of water and other suitable solvents such as ethanol at elevated temperature.

 Alternatively the nitrile can be converted to a tetrazole by treatment with an alkali metal azide, typically sodium azide, an ammonium halide such as ammonium chloride and in a
10 suitable solvent such as DMF at an elevated temperature.

 When compounds of the invention are produced as racemates, these can be separated into their enantiomers by methods known to those skilled in the art, typically by using a chiral HPLC column.

15 Furthermore, the invention relates to a method for the preparation of a medicament comprising the steps of:

- a) preparing a compound according to the invention
- b) formulation of a medicament containing said compound.

20 The compounds according to the invention and their pharmaceutically acceptable salts, optionally in combination with other pharmaceutically active compounds are suitable to treat or prevent Alzheimer's disease or the symptoms thereof. Such additional compounds include cognition-enhancing drugs such as acetylcholinesterase inhibitors (e.g. Donepezil, Tacrine, Galantamine, Rivastigmin), NMDA antagonists (e.g.
25 Memantine) PDE4 inhibitors (e.g. Ariflo) or any other drug known to a person skilled in the art suitable to treat or prevent Alzheimer's disease. Such compounds also include cholesterol-lowering drugs such as statins (e.g. simvastatin). These compounds can be administered to animals, preferably to mammals, and in particular humans, as pharmaceuticals by themselves, in mixtures with one another or in the form of
30 pharmaceutical preparations.

5 Various delivery systems are known and can be used to administer a compound of the invention for the treatment of Alzheimer's disease or for the modulation of the γ -secretase activity, e.g. encapsulation in liposomes, microparticles, and microcapsules:

If not delivered directly to the central nervous system, preferably the brain, it is advantageous to select and/or modify methods of administration in such a way as to allow
10 the pharmaceutical compound to cross the blood-brain barrier.

Methods of introduction include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes.

15 The compounds may be administered by any convenient route, for example by infusion, by bolus injection, by absorption through epithelial or mucocutaneous linings and may be administered together with other biologically active agents.

Administration can be systemic or local. In addition, it may be desirable to introduce the
20 pharmaceutical compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, e.g. by use of an inhaler or nebulizer, and formulation with an aerosolizing
25 agent.

In another embodiment, the compound can be delivered in a vesicle, in particular a liposome (Langer (1990) Science 249, 1527; Treat et al. (1989) Liposomes in the Therapy of Infectious Disease and Cancer, Lopez-Berestein and Fidler, eds., Liss, New
30 York, 353; Lopez-Berestein, *ibid.*, 317)

5 In yet another embodiment, the compound can be delivered via a controlled release system. In one embodiment, a pump may be used (Sefton (1987) CRC Crit. Ref. Biomed. Eng. 14, 201; Buchwald et al. (1980) Surgery 88, 507; Saudek et al. (1989) N. Engl. J. Med. 321, 574). In another embodiment, polymeric materials can be used (Ranger and Peppas (1983) Macromol. Sci. Rev. Macromol. Chem. 23, 61; Levy et al. 10 (1985) Science 228, 190; During et al. (1989) Ann. Neurol. 25, 351; Howard et al. (1989) J. Neurosurg. 71, 858). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (e.g. Goodson, 1984, In: Medical Applications of Controlled Release, supra, Vol. 2, 115). Other controlled release systems are discussed in the review 15 by Langer (1990, Science 249, 1527).

In order to select an appropriate way of administration, the person skilled in the art will also consider routes of administration which have been selected for other known Anti-Alzheimer-drugs.

20 For example, Aricept/Donepezil and Cognex/Tacrine (all acetylcholinesterase-inhibitors) are being taken orally, Axura/Memantine (an NMDA-receptor antagonist) has been launched both as tablets/liquid and as an i.v.-solution.

Furthermore, the skilled person in the art will take into account the available data with respect to routes of administration of members of the NSAID-family in clinical trials and 25 other studies investigating their effect on Alzheimer's disease.

In order to select the appropriate dosage, the person skilled in the art will choose a dosage which has been shown to be not toxic in preclinical and/or clinical studies and which can be in accordance with the values given beforehand, or which may deviate from these.

30 The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. However, suitable dosage ranges for intravenous administration are generally about 20-500

5 micrograms of active compound per kilogram body weight. Suitable dosage ranges for intranasal administration are generally about 0.01 mg/kg body weight to 1 mg/kg body weight. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems.

10 An exemplary animal model is the transgenic mouse strain "Tg2576" containing an APP695-form with the double mutation KM670/671NL. For reference see e.g. patent US5877399 and Hsiao et al. (1996) Science 274, 99 and also Kawarabayashi T (2001) J. Neurosci. 21, 372; Frautschy et al. (1998) Am. J. Pathol. 152, 307; Irizarry et al. (1997) J. Neuropathol. Exp. Neurol. 56, 965; Lehman et al. (2003) Neurobiol. Aging 24, 645.

15 Substantial data from several studies are available to the skilled person in the art which are instructive to the skilled person to select the appropriate dosage for the chosen therapeutic regimen.

The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a
20 tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

Wherein the present invention is directed to co-therapy or combination therapy, comprising administration of one or more compound(s) "therapeutically effective
25 amount" shall mean that amount of the combination of agents taken together so that the combined effect elicits the desired biological or medicinal response. Further, it will be recognized by one skilled in the art that in the case of co-therapy the amount of each component of the combination if used by itself may or may not be a therapeutically effective amount.

30 Numerous studies have been published in which the effects of molecules on the γ -secretase activity are described. Exemplary studies are Lim et al. (2001) Neurobiol. Aging 22, 983; Lim et al. (2000) J Neurosci. 20, 5709; Weggen et al. (2001) Nature 414, 212; Eriksen et al. (2003) J Clin Invest. 112, 440; Yan et al. (2003) J Neurosci. 23, 7504.

5 As used herein, unless otherwise noted, the term “isolated form” shall mean that the compound is present in a form which is separate from any solid mixture with another compound(s), solvent system or biological environment.

As used herein, unless otherwise noted, the term “substantially pure base” shall mean that
10 the mole percent of impurities in the isolated base is less than about 5 mole percent, preferably less than about 2 mole percent, more preferably, less than about 0.5 mole percent, most preferably, less than about 0.1 mole percent.


As used herein, unless otherwise noted, the term “substantially free of a corresponding
15 salt form(s)” when used to described the compound of formula (I) shall mean that mole percent of the corresponding salt form(s) in the isolated base of formula (I) is less than about 5 mole percent, preferably less than about 2 mole percent, more preferably, less than about 0.5 mole percent, most preferably less than about 0.1 mole percent.

The term “pharmaceutically acceptable” means a non-toxic material that does not
20 interfere with the effectiveness of the biological activity of the active ingredients. Such preparations may routinely contain pharmaceutically acceptable concentrations of salts, buffering agents, preservatives, compatible carriers and optionally other therapeutic agents.

25 **General Synthetic Description**

The following general description is for illustrative purposes only and is in no way meant to limit the invention.

Compounds of Formula I wherein X, R¹, R², R³, R⁴, R⁹, and R¹⁰ are defined as in
Formula I, and Y is CO₂H, may be obtained by hydrolysis of ester II under standard
30 acidic or basic hydrolysis conditions, including reaction with NaOH, at room temperature, for several hours, in an appropriate solvent mixture, such as water, tetrahydrofuran (THF), and methanol. For illustrative purposes, ester II is shown with X as CHR⁵, but those skilled in the art will recognize that ester hydrolysis will work for all X as defined in Formula I.



5

Compound II may be obtained by coupling either compound IIIa or IIIb with an aryl boronic acid or an aryl pinacol boronate ester under typical Suzuki conditions, e.g. in aqueous dimethoxyethane (DME) or THF in the presence of sodium carbonate and $\text{Pd(PPh}_3)_4$. Compound II may be also obtained coupling reaction of compound IIIa or


10 IIIb with aryltin bromide or arylzinc bromide under Stille coupling conditions, e.g. in THF or DME with $\text{Pd(PPh}_3)_4$. Compound IIIa may be obtained from the reaction of phenol IIIc with trifluoromethanesulfonic anhydride ($\text{ Tf}_2\text{O}$) in dichloromethane (DCM) in the presence of an amine such as triethylamine at 0 °C. Intermediate IIIb can be obtained from the reaction of phenol IIIc with a concentrate of HCl, or HBr, HI at elevated

15 temperature (25-150 °C). Alternatively, compound IIIb can be obtained under mild conditions by treatment of the corresponding triflates IIIc with pinacolborane in dioxane in the presence of triethylamine catalyzed with PdCl_2 to give an aryl pinacolboronate ester which is then were treated with copper (II) halide in the methanol-water procedure described by Nesmejanow et al. (Chem Ber. **1960**, 2729). Alternatively, compound IIIb

20 where Q is iodine can be obtained by reacting the aforementioned pinacolboronate ester with NaI in aqueous THF in the presence of chloramines-T, as described by J. W. Huffman et. al.(Synthesis , **2005**, 547).

Compound IIIc may be prepared by debenzylation of compound IV by hydrogenation in alcohol, e.g. MeOH or EtOH in the presence of Pd-C. Debenzylation can also be


25 achieved with other methods, such as BBr_3 in DCM, NaCN in DMSO/ 120-200 °C or LiCN in DMF/ 120-200 °C.



5


Compound IV may be prepared from alkylation of compound V with either an alkyl or alkenyl halide. Treatment of compound V in THF or another aprotic solvent with a base, e.g. lithium bis(trimethylsilyl) amide, sodium bis(trimethylsilyl) amide, or lithium diisopropylamide at -78°C , followed by the addition of an electrophile, e.g. an alkyl or

10 alkenyl halide, yields alkylated compound IV.




Compound V may be prepared from compound VI through a coupling reaction with an arylboronic acid under Suzuki conditions of aqueous sodium carbonate in DME in the presence of $\text{Pd}(\text{PPh}_3)_4$. Similarly, the triflates can be converted to boronate esters under

15 the conditions described above and then can be coupled with aryl bromides or aryl chlorides to give compound V.




V

Intermediate compound VI may be prepared from compound VII with trifluoromethanesulfonic anhydride in DCM in the presence of one equivalent of pyridine at 0 °C.




VI

Intermediate compound VII can be prepared from mono-debenzylation of compound VIII. Selective mono-debenzylation of compound VIII can be achieved by treatment with 1.1 equivalents of base, e.g. sodium hydroxide or potassium hydroxide, in ethanol or methanol solution in the presence of Pd-C catalyst under hydrogen atmosphere in a Parr shaker.




VII

- 5 Intermediate VIII can be easily prepared from reaction of 3,5-dihydroxyphenyl acetic acid methyl ester (commercially available) with benzyl bromide and potassium carbonate in DMF at room temperature.




VIII


- Compound I has a chiral center α to the carboxylic group, and can exist as one of two enantiomers (or a mixture thereof, wherein an enantiomeric excess may or may not be present). The enantiomers Ia (R enantiomer) and Ib (S enantiomer) are shown. The pure enantiomers Ia and Ib may be obtained by chiral separation using a chiral column. The enantiomers Ia and Ib may also be separated by resolutions through forming chiral amine salts by fractional recrystallizations. The enantiomers Ia and Ib also may be obtained from kinetic resolution of the racemate of corresponding esters using lipase enzymes, e.g. AmanoAk, Amano lipase PS, Amano lipaseA, Amano lipase M, Amano lipase F-15 Amano lipase G (from Biocatalytics Inc) in aqueous organic solvents, e.g. aqueous DMF, DMSO, t-butylethyl ether or triton X-100 aqueous solutions.



- 20 Both enantiomers of compound I may be prepared from chiral syntheses. Compounds Ia and Ib may be obtained from the removal of the chiral auxiliary groups from compounds IXa and IXb respectively with lithium hydroxide in aqueous THF in the presence of hydrogen peroxide.




IXa




IXb

Compounds IXa and IXb may be obtained by coupling compounds Xa or Xb respectively with an aryl boronic acid or aryl pinacol boronate ester under Suzuki conditions, e.g. in aqueous DME or THF in the presence of sodium carbonate and Pd(PPh₃)₄. Compounds IXa and IXb may be also obtained by coupling compounds Xa or Xb respectively with an aryltin bromide or arylzinc bromide under Stille coupling conditions, e.g. in THF or DME with Pd(PPh₃)₄.




Xa




Xb

Compounds Xa and Xb may be obtained from reaction of phenols XIa and XIb respectively with trifluoromethanesulfonyl anhydride in DCM in the presence of an amine, such as triethylamine, at 0 °C.





XIa



XIb



5

Compounds XIa and XIb may be prepared from debenzylation of compounds XIIa and XIIb respectively by hydrogenation in an alcohol solvent, e.g. MeOH or EtOH, in the presence of Pd-C.



10


Compounds XIIa and XIIb may be prepared from the alkylation of compounds XIIIa and XIIIb respectively with an alkyl halide or alkenyl halide. Treatment of compounds XIIIa and XIIIb in THF or other aprotic solvents with bases, e.g. lithium bis(trimethylsilyl) amide, sodium bis(trimethylsilyl) amide, or lithium diisopropylamide at -78°C ,
 15 followed by the addition of electrophiles, e.g. alkyl halides or alkenyl halides, gives alkylated compounds XIIa and XIIb respectively.



20

Compounds XIIIa and XIIIb may be prepared from the common intermediate XIV by coupling with either R-isomer of 4-benzyl-oxazolidin-one XVa or S-isomer of 4-benzyl -
 oxazolidin-one XVb by Evans's procedures. Intermediate XIV may be reacted with pivaloyl chloride, oxalyl chloride or isopropyl chloroformate in THF in the presence of a

- 5 base, e.g. triethylamine or N-methylmorpholine, to mixed anhydrides or acid chlorides which then were reacted with the lithium salt of XVa or XVb in THF.



Intermediate compound XIV may be obtained from ester hydrolysis of compound V with bases in aqueous alcohol solution, e.g. LiOH or NaOH in aqueous methanol solution.

10 Synthetic Procedures

All reactions were carried out under inert atmosphere unless otherwise stated. NMR spectra were obtained on a Bruker dp400. LCMS was carried out on an Agilent 1100 using a ZORBAX[®] SB-C18, 4.6 x 75 mm, 3.5 micron column for method A. Column flow was 1ml/min and solvents used were water and acetonitrile (0.1%TFA) with an injection volume of 10ul. Wavelengths were 254 and 210nm. Methods are described below:

Method	Flow Rate	Solvent
A	1ml/min	0-1.5 min 30-95%MeCN 1.5-4.5min 95% MeCN 4.5-5 min 95%-5% MeCN

20 Abbreviations

Ac	Acetyl
d	Doublet
DCM	Dichloromethane
DME	1,2-dimethoxyethane
DMF	N,N-dimethylformamide
DMSO	Dimethyl sulfoxide
e.e.	enantiomeric excess
eq	Equivalents
Et	Ethyl
EtOAc	ethyl acetate
g	Gram
h	Hour
HPLC	high pressure liquid chromatography
K ₂ CO ₃	Potassium carbonate
l	Litre
LCMS	liquid chromatography – mass spectrometry
LDA	lithium diisopropylamide
M	Molar
m	Multiplet
Me	Methyl
min	Minute
Mol	Mole
NMR	nuclear magnetic resonance

q	Quartet
RT	Retention time
s	Singlet
Sat	Saturated
t	Triplet
TFA	Trifluoroacetic acid
THF	Tetrahydrofuran

5 **EXAMPLES:****Example (i) Preparation of (4''-Chloro-4-trifluoromethyl-[1,1';3,1'']terphenyl-5'-yl)-acetic acid**Preparation of 1-benzyloxy-3,5-dibromobenzene

10 Benzylalcohol (9.7 mL, 94 mmol) was added dropwise to a suspension of NaH (4.0 g of a 60% suspension in mineral oil, 100 mmol) in THF (150 mL) at room temperature and the mixture was stirred at room temperature for 1h before 1,3-dibromo-5-fluorobenzene (15.9 g, 62.5 mmol) was added. The reaction was stirred at room temperature for 12h. Water was added carefully and the THF was evaporated under reduced pressure. The residue was extracted with iso-hexane (x3) and the combined organic extracts were washed with
15 NaOH solution (1M aq), water, brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash column chromatography (EtOAc : petroleum ether) to give 1-benzyloxy-3,5-dibromobenzene as a colourless liquid, 14.7 g 69 % yield. ¹H NMR (CDCl₃) δ 7.45-7.33 (m, 5H), 7.30-7.28 (m, 1H), 7.10-7.08 (m, 2H), 5.02 (s, 2H).

20

Preparation of (3-benzyloxy-5-bromo-phenyl)-acetic acid ethyl ester

Malonic acid *tert*-butyl ester ethyl ester (10.2 mL, 53.8mmol) was added dropwise to a suspension of NaH (2.2 g of a 60% suspension in mineral oil, 53.8 mmol) in dioxane (200 mL) at room temperature and the mixture was stirred at this temperature for 1h before
25 CuBr (7.7 g, 53.8 mmol) and 1-benzyloxy-3,5-dibromobenzene (9.2 g, 26.9 mmol) were added. The reaction mixture was heated to reflux for 5h. HCl solution (1M aq, 100 mL) was carefully added and the mixture was extracted with iso-hexane (x3). The combined organic extracts were washed with HCl solution (1M aq), water, brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash
30 column chromatography (EtOAc : petroleum ether) to give, in order of elution, recovered 1-benzyloxy-3,5-dibromobenzene (3.2 g, 9.4 mmol) in 35% yield and 2-(3-benzyloxy-5-bromo-phenyl)-malonic acid *tert*-butyl ester ethyl ester (7.2 g, contains 1.4 equivalent malonic acid *tert*-butyl ester ethyl ester, 10 mmol) as a colourless liquid in 37% yield.

5 2-(3-Benzyloxy-5-bromophenyl)malonic acid *tert*-butyl ester ethyl ester (7.2 g, contains 1.4 equivalent malonic acid *tert*-butyl ester ethyl ester, 10 mmol) was dissolved in glacial AcOH (50 mL) and heated to reflux for 12h. The AcOH was removed under reduced pressure. The residue was poured into Na₂CO₃ solution (sat aq) and the mixture was extracted with EtOAc (x3). The combined organic extracts were washed with water,
10 brine, dried (MgSO₄), filtered and concentrated under reduced pressure to give (3-benzyloxy-5-bromo-phenyl)-acetic acid ethyl ester as a yellow liquid yield 6.8 g (97%).
¹H NMR (CDCl₃) δ 7.44-7.30 (m, 5H), 7.07-7.03 (m, 2H), 6.87-6.84 (m, 1H), 5.03 (s, 2H), 4.15 (q, 2H), 3.54 (s, 2H), 1.26 (t, 3H).

15 (5-Benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid ethyl ester

(3-Benzyloxy-5-bromo-phenyl)-acetic acid ethyl ester (2.50 g, 7.2 mmol) was added to a solution of 4-(trifluoromethyl)phenyl boronic acid (1.5 g, 8.0 mmol) and K₂CO₃ (14.4 mmol, 2 M aq.) in DME (25 mL). Nitrogen was bubbled through the reaction mixture for 10 minutes before addition of tetrakis(triphenylphosphine)palladium(0) (10 % wt) and the
20 resultant mixture was heated to 80 °C for 4 hours under inert atmosphere. The reaction mixture was diluted with water and extracted with EtOAc (x3). The combined organic extracts were washed with sat. Na₂CO₃, brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash column chromatography (EtOAc : petroleum ether) to give (5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic
25 acid ethyl ester (2.2g) as a colourless gum in 74% yield. ¹H NMR (CDCl₃) δ 7.59-7.54 (m, 2H), 7.48-7.30 (m, 8H), 7.13-7.11 (m, 2H), 6.94-6.91 (m, 1H), 5.12 (s, 2H), 4.16 (q, 2H), 3.64 (s, 2H), 1.27 (t, 3H).

Preparation of (5-hydroxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid ethyl ester

30 To a solution of (5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid (2.5 g, 5.5 mmol) in EtOH (50 mL) was added 10% Pd/C (5% wt) and the resultant black suspension stirred under an atmosphere of H₂ for 5 hours. The resultant mixture was filtered through celite and evaporated to dryness. The residue was purified by flash column

- 5 chromatography (EtOAc : petroleum ether) to give (5-hydroxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid ethyl ester as a white solid, yield 2.3 g (93 %).

Preparation of (5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid ethyl ester.

- 10 Trifluoromethanesulphonic anhydride (570 mg, 2.0 mmol) was added drop wise to a solution of (5-hydroxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid ethyl ester (546 mg, 1.69 mmol) and pyridine (0.41 mL, 5.0 mmol) in DCM (10 mL) at 0 °C. The temperature was maintained at 0 °C for 15mins before warming to room temperature and the mixture stirred for 18 hrs. The reaction was diluted with DCM, washed with H₂O, Na₂CO₃, dil.
15 HCl, brine, dried (MgSO₄) and evaporated under reduced pressure to give a yellow oil. The residue was purified by flash column chromatography (EtOAc : petroleum ether), yield 695 mg (96 %). ¹H NMR (CDCl₃) δ 7.72 (d, 2H), 7.66 (d, 2H), 7.54 (t, 1H), 7.39 (t, 1H), 7.28 (t, 1H), 4.19 (q, 2H), 3.73 (s, 2H), 1.28 (t, 3H).

- 20 Preparation of (4''-chloro-4-trifluoromethyl-[1,1';3'1'']terphenyl-5'-yl)-acetic acid methyl ester.

- A solution of (5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid ethyl ester (100 mg, 0.2 mmol), 4-chlorophenyl boronic acid (41 mg, 0.24 mmol), K₂CO₃ (2 M solution in H₂O, 220 μL, 0.4 mmol) in DME (2.0 mL) was heated to 80 °C
25 in the presence of bis(tri-t-Butylphosphine) palladium (0) (cat) for 2 hrs. The mixture was cooled to room temperature filtered, diluted with EtOAc, washed with Na₂CO₃, dil HCl, brine, dried (MgSO₄) and evaporated under reduced pressure to give an off white solid. The residue was purified by flash column chromatography (EtOAc : petroleum ether), ¹H NMR (CDCl₃) δ 7.71 (s, 4H), 7.65-7.68 (m, 1H), 7.42 (d, 2H), 7.52-7.48 (m, 2H), 7.36
30 (d, 2H), 3.77 (s, 2H), 3.74 (s, 3H)

Preparation of (4''-Chloro-4-trifluoromethyl-[1,1';3,1'']terphenyl-5'-yl)-acetic acid.

- 5 NaOH solution (1 ml, 1M aq) was added to a solution of (5-benzyloxy-biphenyl-3-yl)-acetic acid ethyl ester (50 mg) in EtOH (2 ml) and the mixture was stirred at room temperature for 12h. The reaction mixture was diluted with HCl solution (2M aq) and extracted with EtOAc (x3). The combined organic extracts were washed with water, brine, dried (MgSO₄), filtered and concentrated under reduced pressure to give colourless
 10 solid in 90% yield. ¹H NMR (CDCl₃) δ 7.71 (s, 4H), 7.65-7.68 (m, 1H), 7.42 (d, 2H), 7.52-7.48 (m, 2H), 7.36 (d, 2H), 3.79 (s, 2H). LCMS R.T. 3.4 min (389 M -H)

In an analogous fashion, using the appropriate boronic acid, the following were prepared:

Example No	Name	LC method	retention time (min)
(ii)	(4"-Trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid (II)	A	3.3 (M-H 355)
(iii)	(3-Chloro-4"-trifluoromethyl [1,1';3',1'']terphenyl-5'-yl)-acetic acid (III)	A	3.5 (M-H 389)
(iv)	(4-Hydroxy-4"-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid (IV)	A	2.9 (M-H 372)

15

Example (v) Preparation of (4,4''-dichloro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid

Preparation of (3,5-bis-trifluoromethanesulfonyloxy-phenyl)-acetic acid methyl ester

- Trifluoromethanesulphonic anhydride (7.05 g, 25.0 mmol) was added dropwise to a solution of 3,5-dihydroxyphenyl acetic acid methyl ester (1.80 g, 10.0 mmol) and
 20 pyridine (4.9 mL, 60.0 mmol) in DCM (20 mL) at 0 °C. The temperature was maintained

5 at 0 °C for 15mins before warming to room temperature and the mixture stirred for 18 hrs. The reaction was diluted with DCM washed with H₂O, Na₂CO₃, dil. HCl, brine, dried (MgSO₄) and evaporated under reduced pressure to give a yellow oil. The residue was purified by flash column chromatography (EtOAc : petroleum ether), yield 4.0 g (89 %), ¹H NMR (CDCl₃) δ 7.31 (d, 2H), 7.17 (t, 1H), 3.74 (s, 3H), 7.73 (s, 2H).

10

Preparation of (4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid methyl ester

A solution of (3,5-bis-trifluoromethanesulfonyloxy-phenyl)-acetic acid methyl ester (250 mg, 0.56 mmol), 4-chlorophenyl boronic acid (219 mg, 1.4 mmol), K₂CO₃ (2 M solution in H₂O, 1.1 mL, 2.24 mmol) in DME (4.0 mL) and heated to 80 °C in the presence of
 15 bis(tri-*t*-butylphosphine) palladium (0) (cat) for 4 hrs. The mixture was cooled to room temperature filtered, diluted with EtOAc, washed with Na₂CO₃, dil HCl, brine, dried (MgSO₄) and evaporated under reduced pressure to give an off white solid. The residue was purified by flash column chromatography (EtOAc : petroleum ether), ¹H NMR (CDCl₃) δ 7.61 (t, 1H), 7.55 (d, 4H), 7.46 (d, 2H), 7.42 (d, 4H), 3.75 (s, 2H), 3.73 (s, 3H).

20

Preparation of (4,4"-dichloro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid

(4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid methyl ester was hydrolysed under conditions previously described to give (4,4"-dichloro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid as a clear oil. ¹H NMR (CDCl₃) δ 7.61-7.63 (m, 1H), 7.54 (d, 4H), 7.45-7.62 (m,
 25 2H), 7.42 (d, 2H), 3.78 (s, 2H), LCMS Method A R.T. 3.5 min

In an analogous fashion, using the appropriate boronic acid, the following were prepared:

Example No	Name	LC method	retention time (min)
(vi)	[1,1';3',1'']Terphenyl-5'-yl-acetic acid (VI)	A	3.1

(vii)	(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid (VII)	A	3.4
(viii)	(4,4''-Difluoro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid (VIII)	A	3.1
(ix)	(3,3''-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid (IX)	A	3.5
(x)	(3,3''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid (X)	A	3.4
(xi)	(4,4''-Dimethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid (XI)	A	3.2
(xii)	(4,4''-Dimethoxy-[1,1';3',1'']terphenyl-5'-yl)-acetic acid (XII)	A	3.1

5

Example (xiii) Preparation of 2-(4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid

Preparation of 2-(4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid methyl ester

10 A solution of LDA (0.18 mL of 1.8 M in THF, 0.31 mmol) was added dropwise to a stirred solution of (4,4''-dichloro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid methyl ester (100 mg, 0.26 mmol) in THF (10 mL) at -78 °C. The reaction mixture was stirred for 30 minutes at -78 °C before iodopropane (0.035 mL, 0.31 mmol) was added dropwise. The reaction mixture was allowed to warm to room temperature overnight. A saturated

aqueous solution of ammonium chloride (10 mL) was carefully added and the residue was partitioned between EtOAc and water. The aqueous layers were extracted with EtOAc (x3). The combined organic layers were washed with water, brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash column chromatography (EtOAc : petroleum ether) yield 70 mg (66 %) ¹H NMR (CDCl₃) δ 7.60 (t, 1H), 7.55 (d, 4H), 7.48 (d, 2H), 7.42 (d, 4H), 3.65-3.72 (m, 4H), 2.11-2.23 (m, 1H), 1.78-1.89 (m, 1H), 1.28-1.40 (m, 4H), 0.94 (t, 3H)

Preparation of 2-(4,4''-dichloro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid

2-(4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid methyl ester was hydrolysed under conditions previously described to afford 2-(4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid as a clear oil.

Examples (xiv) and (xv) Preparation of (R)-2-(4,4''-Dichloro-[1,1';3',1'']

terphenyl-5'-yl)-pentanoic acid and (S)-2-(4,4''-Dichloro-[1,1';3',1'']

terphenyl-5'-yl)-pentanoic acid

The enantiomers of 2-(4,4''-dichloro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid were separated on a 80 mm I.D. Dynamic axial compression column filled with 500 grams of 20 micrometer Chiralpak AD [3,5 dimethyl phenyl carbamate of amylose (Daicel)] of packing bed length: 21 cm with ethanol and 0.1 % TFA as the eluent at a flow rate of 80 ml/min at ambient temperature. The first peak off the column at 18.25 min was designated R* and the 2nd peak at 24.75 min was designated S*.

Example (xvi) Preparation of 4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-carboxylic acid

Preparation of 4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-carbonitrile

4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-carbonitrile was prepared in an analogous fashion to example 5 replacing 3,5-dihydroxyphenyl acetic acid methyl ester by 3,5-dihydroxybenzonitrile. ¹H NMR (CDCl₃) δ 7.91 (t, 1H), 7.80 (d, 2H), 7.52-7.52 (m, 4H), 7.45-7.49 (m, 4H).

5

Preparation of 4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-carboxylic acid

A suspension of 4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-carbonitrile (40 mg, 0.12 mmol) in EtOH (1 mL) and NaOH (25% aq., 0.5 mL) was heated at reflux for 3 h. The resultant brown solution was evaporated to dryness, acidified with dil. HCl and the resultant ppt
10 extracted with EtOAc. The organic layer was washed with brine, dried (MgSO₄), and evaporated to give an off white solid. Trituration with petrol/EtOAc gave a beige solid yield 23 mg (55%). ¹H NMR (MDOD) δ 8.24 (t, 1H), 8.06 (d, 2H), 7.71-7.74 (m, 4H), 7.48-7.52 (m, 4H). LCMS Method A R.T. 3.6 min


15 **Example (xvii) Preparation of 5-(4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-1H-tetrazole**

A mixture of 4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-carbonitrile (50 mg, 0.15 mmol), sodium azide (20 mg, 0.31 mmol) and ammonium chloride (16 mg, 0.31 mmol) in DMF (1 mL) was heated at 100 °C for 16h. The reaction mixture was acidified with 1M HCl
20 solution poured into water and extracted with ethyl acetate. The organic layer was washed with brine, dried (Na₂SO₄) and the solvent was removed in vacuo to give 5-(4,4''-dichloro-[1,1';3',1'']terphenyl-5'-yl)-1H-tetrazole yield 36mg (64%) as an off-white solid. ¹H NMR (DMSO) δ 7.49 (d, 2H), 7.23 (t, 1H), 7.17 (s br, 1H), 6.96-6.99 (m, 4H), 6.71-6.74 (m, 4H), LCMS method A R.T. 3.5

25


Example 1

2-(3,5-Difluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



5

a) (3,5-Bis-benzyloxy-phenyl)-acetic acid methyl ester



10

A mixture of (3,5-dihydroxy-phenyl)-acetic acid methyl ester (from Aldrich, 70g, 0.385 mole), benzylbromide (137mL, 1.16mole), potassium carbonate (160g, 1.16mole) and DMF (1.5L) under N₂ was mechanically stirred at room temperature overnight. The resulting reaction mixture was poured into a mixture of 1.5L of ice-water with stirring.


15 The precipitate was obtained by filtration and washed with heptane successively to remove benzyl bromide to give the title compounds (123.7g) as a brown solid which was air dried for the next reaction.

¹H-NMR(CDCl₃): δ 3.60 (s, 2H), 3.71(s,3H), 5.05 (s, 4H), 6.60 (s, 3H), 7.35-7.50 (m, 10H); Calcd for C₂₃H₂₂O₄ (M+H) 363.15, Found 363.

20


b) 3-Benzyloxy-5-hydroxy-phenyl)-acetic acid ethyl ester

5



A solution of 50 grams (1.38moles) of 3,5-Bis-benzyloxy-phenyl)-acetic acid methyl ester and NaOH (6.6 g, 1.65 moles) in 1 L of EtOH in the presence of 10 % of Pd-C was hydrogenated in a Parr shaker till one equivalent of hydrogen was consumed. The mixture was acidified with concentrated HCl and then the catalyst and solvent were removed to give an oil residue. The crude was purified by ISCO silica gel column chromatography using EtOAc-heptane as eluents (gradient from 10% to 75% of EtOAc) to give 25 grams (65% yield) the title compound. ¹H-NMR(CDCl₃): δ 1.15-1.20 (t, 3H), 3.4-(s,2H), 4.05-4.1 (q, 2H),4.9(s, 2H), 5.5(s, 1H), 6.4(s, 2H), 6.5(s, 1H), 7.207.35(m, 5H); Calcd for C₁₇H₁₈O₄ (M+H) 287.3, Found 287.

c) (3-Benzyloxy-5-trifluoromethanesulfonyloxy-phenyl)-acetic acid ethyl ester




20

To a solution of 3-(benzyloxy-5-hydroxy-phenyl)-acetic acid ethyl ester (74.4 g, 0.26 mol) in dichloromethane (700 mL) was added pyridine (62.5 mL, 0.78 mol). The mixture was cooled to 0 °C. To this cold solution was added trifluoromethanesulfonic anhydride (65.6 mL, 0.39 mol), over 1.5 h, maintaining the internal temperature below 5 °C and stirred for further 0.5 h at 0 °C. This reaction mixture was poured to a mixture of 1 N HCl (420 mL), and wet-ice (105 g) and stirred for 0.5 h. The aqueous layer was extracted with dichloromethane (2 x 100 mL). Combined fractions were washed with water (2 x 100 mL), saturated aqueous NaHCO₃ solution (2 x 100 mL), and brine (2 x 100 mL). The

5 organics were dried (MgSO_4) and concentrated in vacuo to receive a reddish liquid (108 g) which was carried on to the next step without further purification.

Calcd for $\text{C}_{18}\text{H}_{17}\text{F}_3\text{O}_6\text{S}$ (M+H) 419.07, Found 419.1

d) (5-Benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid ethyl ester




10

A mixture of (3-benzyloxy-5-trifluoromethanesulfonyloxy-phenyl)-acetic acid ethyl ester (108 g, 0.26 mol), 4-(trifluoromethyl)phenylboronic acid (55.6 g, 0.29 mol), 1,2-dimethoxyethane (1.1 L) and aqueous Na_2CO_3 (2 M, 129 mL, 0.26 mol) was mechanically stirred while purging N_2 at room temperature for 10 min. To this system
 15 was added $\text{Pd}(\text{Ph}_3)_4$ (480 mg, 0.42 mmol) and heated to reflux (95 °C) for 2.5 h. The red-brown mixture was diluted with EtOAc (0.5 L) and washed with saturated aqueous NaHCO_3 solution (3 x 200 mL) and brine (2 x 200 mL). The organic fraction was dried (Na_2SO_4) and concentrated in vacuo. The crude mixture was purified by ISCO silica gel column chromatography to obtain (5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid ethyl ester (107 g, 100%).
 20

$^1\text{H-NMR}$ (CDCl_3): δ 1.26 (t, 3H), 3.66 (s, 2H), 4.17 (q, 2H), 5.12 (s, 2H), 6.99 (s, 1H), 7.12 (s, 2H), 7.34-7.49 (m, 5H), 7.67 (s, 4H); Calcd for $\text{C}_{24}\text{H}_{21}\text{F}_3\text{O}_3$ (M+H) 415.14, Found 415.2.

e) 2-(5-Benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pent-4-enoic acid ethyl ester

25



5


To a solution of compound **1d** (4.9g, 11.8mmole) in THF (50 mL) at -78°C was added $\text{Li}[\text{N}(\text{SiMe}_3)_2]$ (1N in THF, 14.2mL, 14.2mmol) dropwise. The reaction mixture was stirred for 1 h at -78°C and then 3-bromo-2-methyl-propene (1.25mL, 12.4mmole) was added dropwise. The solution was slowly warmed up to -35°C and stirred at -35°C for 0.5 h. The reaction was quenched with NH_4Cl saturated solution and extracted with EtOAc. The organic extracts was dried (Na_2SO_4), concentrated and purified by column chromatography give the title compound (5.1g, 92%) as a clear oil; ^1H NMR (400 MHz, CHLOROFORM- D) δ ppm 1.19 - 1.29 (m, 3 H), 1.74 (s, 3 H), 2.47 (m, 1 H), 2.85 (m, 1 H), 3.83 (m, 1 H), 4.11 (m, 2 H), 4.72 (s, 1 H), 4.77 (s, 1 H), 5.12 (s, 2 H), 7.03 (s, 1 H), 7.10 (s, 1 H), 7.15 (s, 1 H), 7.35 - 7.48 (m, 5 H), 7.67 (s, 4 H); Calcd for $\text{C}_{28}\text{H}_{27}\text{F}_3\text{O}_3$ (M+H) 469.19, Found 469.

10

15

f) 2-(5-Hydroxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoic acid ethyl ester


20



5 A mixture of compound **1e** (5.1g, 10.9mmole), 10% Pd/C (500mg) in EtOH (50mL) was hydrogenated under H₂ (40psi) in par-shaker for 20h. The resulting reaction mixture was filtered through celite and the filtrate was concentrated to give the title compound (4.2g, 100%) as a clear oil; ¹H NMR (300 MHz, CHLOROFORM-D) δ ppm 0.92 (d, *J*=6.6 Hz, 6 H), 1.25 (m, 3 H), 1.49 – 1.61 (m, 1 H), 1.65 – 1.70 (m, 1 H), 1.95 – 2.05 (m, 1 H), 3.67
 10 (t, *J*=7.7 Hz, 1 H), 4.10 – 4.29 (m, 2 H), 6.91 (s, 1 H), 6.97 (t, *J*=2.0 Hz, 1 H), 7.08 (s, 1 H), 7.65 (s, 4 H); Calcd for C₂₁H₂₃F₃O₃ (M+H) 381.16, Found 381.

g) 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester

15



To a solution of compound **1f** (2.8g, 7.36mmol) and N-phenyl-bis-
 (trifluoromethanesulfonylimide) (3.16g, 8.83mmol) in THF (30mL) under N₂ was added
 20 Et₃N (2.05mL, 14.7mmol). The reaction mixture was heated to reflux overnight. After cooling to room temperature, the solution was concentrated and purified by column chromatography to give the title compound (3.7g, 98%) as a colorless thick oil; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 0.94 (dd, *J*=6.60, 1.47 Hz, 6 H), 1.22 - 1.28 (m, 3 H), 1.46 - 1.52 (m, 1 H), 1.69 (ddd, *J*=13.82, 7.09, 6.97 Hz, 1 H), 1.98 - 2.06 (m, 1 H),
 25 3.75 (t, *J*=7.83 Hz, 1 H), 4.10 - 4.21 (m, 2 H), 7.31 (s, 1 H), 7.38 (s, 1 H), 7.57 (s, 1 H), 7.65 - 7.75 (m, 4 H); Calcd for C₂₂H₂₂F₆O₅S (M+H) 513.11, Found 513.

5

h) 2-(3,5-Difluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid


A mixture of compound **1g** (50mg, 0.098mmol), 3, 5-difluorobenzeneboronic acid (23mg, 0.146mmol), Pd(PPh₃)₄ (23mg, 0.0196mmol) and Na₂CO₃ (2N in H₂O, 0.098mL, 0.196mmol) in DME (1mL) was heated at 85 °C for 3h. After cooling to room temperature, the solution was partitioned between EtOAc and H₂O. The organic layer was dried (Na₂SO₄), concentrated and purified by column chromatography to give an ethyl ester intermediate.

A mixture of the above intermediate and NaOH (2N in H₂O, 0.147mL, 0.294mmol) in THF-MeOH (0.6mL-0.6mL) was stirred for 18h and concentrated. CH₂Cl₂ and water were added, and the mixture was acidified with 1N HCl. The organic phase was separated and the aqueous phase was extracted with CH₂Cl₂. The combined organic layers were dried, concentrated, and purified by column chromatography to give 30mg (69%, 2 steps) of the title compound as a white solid; ¹H NMR (400 MHz, MeOD) δ ppm 0.88 (dd, *J*=6.60, 3.18 Hz, 6 H), 1.43 - 1.50 (m, 1 H), 1.66 (ddd, *J*=13.82, 7.09, 6.97 Hz, 1 H), 1.92 - 1.98 (m, 1 H), 3.76 (t, *J*=7.83 Hz, 1 H), 6.87 (tt, *J*=9.08, 2.29 Hz, 1 H), 7.21 - 7.26 (m, 2 H), 7.55 (d, *J*=1.47 Hz, 1 H), 7.58 - 7.60 (m, 1 H), 7.66 - 7.72 (m, 3 H), 7.79 (d, *J*=8.07 Hz, 2 H).

25

Example 2

2-(2,4-Difluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



5

The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 2,4-difluorophenylboronic acid under the conditions described in **Example 1**.


10

¹H NMR (400 MHz, MeOD) δ ppm 0.87 (dd, $J=6.60, 2.45$ Hz, 6 H), 1.46 (dt, $J=13.39, 6.63$ Hz, 1 H), 1.64 (ddd, $J=13.82, 7.09, 6.97$ Hz, 1 H), 1.93 (dt, $J=13.63, 7.61$ Hz, 1 H), 3.72 (t, $J=7.83$ Hz, 1 H), 6.96 - 7.02 (m, 2 H), 7.42 - 7.52 (m, 2 H), 7.55 - 7.60 (m, 2 H), 7.64 - 7.70 (m, 2 H), 7.73 - 7.77 (m, 2 H).

15

Example 3

2-(4-Chloro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid




20

The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester

- 5 (intermediate **Example 1g**) with 4-chlorophenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, MeOD) δ ppm 0.87 (dd, *J*=6.60, 3.18 Hz, 6 H), 1.41 - 1.51 (m, 1 H), 1.65 (ddd, *J*=13.69, 7.21, 6.97 Hz, 1 H), 1.95 (ddd, *J*=13.57, 7.70, 7.58 Hz, 1 H), 3.74 (t, *J*=7.83 Hz, 1 H), 7.33 - 7.40 (m, 2 H), 7.50 - 7.59 (m, 4 H), 7.65 (s, 1 H), 7.67 (d, *J*=1.47 Hz, 2 H), 7.75 (d, *J*=8.31 Hz, 2 H); Calcd for
10 C₂₅H₂₂ClF₃O₂ (M+H) 447.13, Found 447.

Example 4

2-(4-Isopropyl-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid




15

- The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 4-isopropyl-phenylboronic acid under the conditions
20 described in **Example 1**; ¹H NMR (400 MHz, MeOD) δ ppm 0.88 (dd, *J*=6.60, 3.42 Hz, 6 H), 1.20 (d, *J*=6.85 Hz, 6 H), 1.44 - 1.49 (m, 1 H), 1.64 - 1.67 (m, 1 H), 1.91 - 1.96 (m, 1 H), 2.83 - 2.88 (m, 1 H), 3.73 (t, *J*=7.70 Hz, 1 H), 7.25 (d, *J*=8.07 Hz, 2 H), 7.51 (dd, *J*=8.44, 1.83 Hz, 4 H), 7.67 (m, 3 H), 7.77 (d, *J*=8.07 Hz, 2 H); Calcd for
C₂₈H₂₉ClF₃O₂ (M+H) 455.5, Found 455.

25

Example 5

2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



5

The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 4-trifluoromethyl-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm


0.93 - 1.02 (m, 6 H), 1.59 (dt, $J=13.39$, 6.63 Hz, 1 H), 1.79 (ddd, $J=13.76$, 7.34, 7.03 Hz, 1 H), 2.08 (dt, $J=13.69$, 7.58 Hz, 1 H), 3.84 (t, $J=7.83$ Hz, 1 H), 7.26 (s, 2 H), 7.58 (d, $J=1.47$ Hz, 2 H), 7.68 - 7.75 (m, 7 H).

10

Example 6

2-(2,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid

15




The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 2-trifluoromethyl-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm

20

- 5 0.86 - 0.97 (m, 6 H), 1.54 (dt, $J=13.39, 6.63$ Hz, 1 H), 1.75 - 1.85 (m, 1 H), 2.01 (ddd, $J=13.94, 7.95, 7.70$ Hz, 1 H), 3.78 (t, $J=7.83$ Hz, 1 H), 7.36 (s, 1 H), 7.38 (d, $J=7.34$ Hz, 1 H), 7.45 - 7.53 (m, 2 H), 7.56 - 7.62 (m, 2 H), 7.67 - 7.78 (m, 5 H); Calcd for $C_{26}H_{22}ClF_6O_2$ (M+Na) 503.44, Found 503.

10 Example 7


4-Methyl-2-(3,5,4''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid



- 15 The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 3,5-bis-trifluoromethyl-phenylboronic acid under the conditions described in **Example 1**; 1H NMR (400 MHz, CHLOROFORM-D) δ ppm
- 20 0.94 - 1.01 (m, 6 H), 1.54 - 1.65 (m, 1 H), 1.79 (ddd, $J=13.82, 7.09, 6.97$ Hz, 1 H), 2.04 - 2.15 (m, 1 H), 3.85 (t, $J=7.70$ Hz, 1 H), 7.56 (d, $J=1.71$ Hz, 1 H), 7.64 (s, 1 H), 7.67 (t, $J=1.59$ Hz, 1 H), 7.73 (s, 4 H), 7.90 (s, 1 H), 8.02 (s, 2 H).

Example 8

2-(2-Fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid




5

The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 2-fluoro-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 0.94 (td, $J=19.93, 6.60$ Hz, 6 H), 1.58 (dt, $J=13.39, 6.63$ Hz, 1 H), 1.78 (ddd, $J=13.69, 7.21, 6.97$ Hz, 1 H), 2.01 - 2.11 (m, 1 H), 3.80 (t, $J=7.83$ Hz, 1 H), 7.16 - 7.27 (m, 2 H), 7.29 - 7.40 (m, 1 H), 7.47 (td, $J=7.70, 1.71$ Hz, 1 H), 7.52 - 7.61 (m, 2 H), 7.66 - 7.76 (m, 5 H); Calcd for C₂₅H₂₂F₄O₂ (M+Na) 453.16, Found 453.

15

Example 9

2-(3-Fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid




The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester

- 5 (intermediate **Example 1g**) with 3-fluoro-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 0.89 - 1.00 (m, 6 H), 1.58 (dt, $J=13.39$, 6.63 Hz, 1 H), 1.78 (ddd, $J=13.76$, 7.34, 7.03 Hz, 1 H), 2.01 - 2.13 (m, 1 H), 3.82 (t, $J=7.83$ Hz, 1 H), 7.05 - 7.13 (m, 1 H), 7.32 (dd, $J=10.03$, 1.47 Hz, 1 H), 7.36 - 7.47 (m, 2 H), 7.52 - 7.61 (m, 2 H), 7.66 - 7.76 (m, 5 H); Calcd for
- 10 C₂₅H₂₂F₄O₂ (M+Na) 453.16, Found 453.

Example 10


2-(4-Fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



- 15 The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 4-fluoro-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 0.88 - 0.99
- (m, 6 H), 1.57 (ddd, $J=13.33$, 6.85, 6.72 Hz, 1 H), 1.77 (ddd, $J=13.88$, 7.21, 7.03 Hz, 1
- 20 H), 2.02 - 2.11 (m, 1 H), 3.80 (t, $J=7.70$ Hz, 1 H), 7.11 - 7.20 (m, 3 H), 7.51 - 7.65 (m, 4 H), 7.68 - 7.73 (m, 4 H); Calcd for C₂₅H₂₂F₄O₂ (M+Na) 453.16, Found 453.

Example 11

4-Methyl-2-(4-methyl-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid




5

The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 4-methyl-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 0.94 (t, *J*=6.60 Hz, 6 H), 1.50 – 1.62 (m, 1 H), 1.6 (s, 1 H), 2.03 - 2.13 (m, 4 H), 3.81 (t, *J*=7.80 Hz, 1 H), 7.25 - 7.33 (m, 3 H) 7.49 - 7.60 (m, 4 H), 7.67 - 7.77 (m, 4 H); Calcd for C₂₆H₂₅F₃O₂ (M+H) 427.18, Found 427.

15 **Example 12**

2-(3,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid




The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 3-trifluoromethyl-phenylboronic acid under the

20

- 5 conditions described in **Example 1**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 0.88 - 0.99 (m, 6 H), 1.53 - 1.63 (m, 1 H), 1.78 (ddd, $J=13.82, 7.09, 6.97$ Hz, 1 H), 2.0 - 2.15 (m, 1 H), 3.83 (t, $J=7.70$ Hz, 1 H), 7.56 - 7.61 (m, 3 H) 7.63 - 7.74 (m, 6 H) 7.74 - 7.86 (m, 2 H); Calcd for C₂₆H₂₂F₆O₂ (M+H) 481.44, Found 481.

10 **Example 13**


2-(4-Chloro-3-fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



- 15 The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 4-chloro-3-fluoro-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 0.88 - 0.99 (m, 6 H), 1.58 (dt, $J=13.21, 6.60$ Hz, 1 H), 1.77 (ddd, $J=13.69, 7.21, 6.97$ Hz, 1 H), 2.07 (ddd, $J=13.57, 7.83, 7.70$ Hz, 1 H), 3.82 (t, $J=7.83$ Hz, 1 H), 7.33 - 7.43 (m, 2 H), 7.45 - 7.50 (m, 1 H), 7.52 - 7.58 (m, 2 H), 7.61 - 7.69 (m, 1 H), 7.71 (s, 4 H); Calcd for C₂₅H₂₁ClF₄O₂ (M+H) 465.88, Found 465.
- 20

Example 14

- 25 **4-Methyl-2-(3,4,5-trifluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid**




5

The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 3,4,5-trifluoro-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 0.92 (d, *J*=6.60 Hz, 6 H), 1.50 - 1.58 (m, 1 H), 1.70 - 1.78 (m, 1 H), 2.01 - 2.15 (m, 1 H), 3.75 - 3.81 (m, 1 H), 7.18 - 7.25 (m, 2 H), 7.46 (s, 1 H), 7.52 - 7.58 (m, 2 H), 7.63 - 7.73 (m, 4 H); Calcd for C₂₅H₂₀F₆O₂ (M+Na) 489.42, Found 488.4.

10

15 **Example 15**

2-(3,5-Dichloro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid




20

The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester

- 5 (intermediate **Example 1g**) with 3,5-dichloro-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 0.96 (d, *J*=6.60 Hz, 6 H), 1.58 (dt, *J*=13.45, 6.72 Hz, 1 H), 1.77 (ddd, *J*=13.76, 7.34, 7.03 Hz, 1 H), 2.09 (ddd, *J*=13.45, 7.83, 7.58 Hz, 1 H), 3.83 (t, *J*=7.70 Hz, 1 H), 7.38 (t, *J*=1.83 Hz, 1 H), 7.48 (d, *J*=1.71 Hz, 2 H), 7.52 (s, 1 H), 7.57 - 7.67 (m, 2 H), 7.71 - 7.77 (m, 4 H);
- 10 Calcd for C₂₅H₂₁Cl₂F₃O₂ (M+H) 482.33, Found 482.

Example 16

2-(4-Cyano-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid




15

- The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 4-cyano-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, MeOD) δ ppm 0.88 (dd, *J*=6.60, 3.18 Hz, 6 H) 1.47 (dt, *J*=13.45, 6.72 Hz, 1 H) 1.66 (ddd, *J*=13.63, 7.09, 6.91 Hz, 1 H) 1.91 - 2.00 (m, 1 H) 3.77 (t, *J*=7.83 Hz, 1 H) 7.61 (dd, *J*=4.16, 1.71 Hz, 2 H) 7.68 (d, *J*=8.31 Hz, 2 H) 7.73 - 7.81 (m, 7 H).
- 20

Example 17

- 25 **2-(3-Cyano-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid**




5

The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 3-cyano-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, MeOD) δ ppm 0.88 (dd, $J=6.60, 3.18$ Hz, 6 H), 1.47 (dt, $J=13.45, 6.72$ Hz, 1 H), 1.66 (ddd, $J=13.63, 7.09, 6.91$ Hz, 1 H), 1.91 - 2.00 (m, 1 H), 3.77 (t, $J=7.83$ Hz, 1 H), 7.61 (dd, $J=4.16, 1.71$ Hz, 2 H), 7.68 (d, $J=8.31$ Hz, 2 H), 7.73 - 7.81 (m, 7 H).

10

15 **Example 18**

2-(4-Cyano-3-fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid




20


- 5 The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 4-cyano-3-fluoro-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, MeOD) δ ppm 0.88 (dd, $J=6.60, 2.93$ Hz, 6 H), 1.46 (dt, $J=13.45, 6.72$ Hz, 1 H), 1.66 (ddd, $J=13.69, 7.21, 6.97$ Hz, 1 H), 1.96 (ddd, $J=13.27, 7.95, 7.64$ Hz, 1 H), 3.78 (t, $J=7.83$ Hz, 1 H), 7.59 - 7.70 (m, 6 H) 7.72 - 7.82 (m, 4 H); Calcd for C₂₆H₂₁F₄NO₂ (M+H) 456.44, Found 456.

Example 19


- 15 **2-(3-Cyano-4-fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid**



- 20 The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 3-cyano-4-fluoro-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, MeOD) δ ppm 0.87 (dd, $J=6.48, 2.81$ Hz, 6 H), 1.47 (dt, $J=13.39, 6.63$ Hz, 1 H), 1.65 (ddd, $J=13.63, 7.09, 6.91$ Hz, 1 H), 1.92 - 2.00 (m, 1 H), 3.71 - 3.77 (m, 1 H), 7.54 (td, $J=7.70, 1.47$ Hz, 1 H), 7.59 - 7.61 (m, 1 H), 7.65 - 7.72 (m, 3 H), 7.77 - 7.81 (m, 2 H), 7.86 - 7.89 (m, 1 H), 7.96 (ddd, $J=8.80, 5.14, 2.45$ Hz, 1 H), 8.02 (dd, $J=5.99, 2.32$ Hz, 1 H); Calcd for C₂₆H₂₁F₄NO₂ (M+H) 456.44, Found 456.

5 **Example 20****4-Methyl-2-(4-trifluoromethoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid**

- 10 The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 4-trifluoromethoxy-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, MeOD) δ ppm 0.87 - 0.98 (m, 6 H), 1.56 (dt, $J=13.39$, 6.63 Hz, 1 H), 1.75 (dt, $J=13.76$, 6.94 Hz, 1 H), 1.98 - 2.08 (m, 1
- 15 H), 3.84 (t, $J=7.83$ Hz, 1 H), 7.36 (t, $J=8.56$ Hz, 2 H), 7.64 (d, $J=5.62$ Hz, 2 H), 7.70 - 7.80 (m, 5 H), 7.84 - 7.89 (m, 2 H).


Example 2120 **2-(4-Methanesulfonyl-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid**

5


The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 4-methylsulfonyl-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, MeOD) δ ppm 0.98 (dd, $J=6.60, 3.18$ Hz, 6 H), 1.57 (dt, $J=13.45, 6.72$ Hz, 1 H), 1.77 (ddd, $J=13.69, 7.21, 6.97$ Hz, 1 H), 2.06 (ddd, $J=13.57, 7.70, 7.58$ Hz, 1 H), 3.17 (s, 3 H), 3.87 (t, $J=7.83$ Hz, 1 H), 7.72 (d, $J=1.71$ Hz, 2 H), 7.78 (d, $J=8.56$ Hz, 2 H), 7.87 - 7.91 (m, 3 H), 7.95 - 7.98 (m, 2 H), 8.05 - 8.08 (m, 2 H).

15 **Example 22**


2-(4-Chloro-3,4''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



20 The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 4-chloro-3-trifluoromethyl-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, MeOD) δ ppm 0.85 - 0.96 (m, 6 H), 1.42 - 1.54 (m, 1 H), 1.65 (ddd, $J=13.82, 7.09, 6.97$ Hz, 1 H), 1.96 (dd, $J=13.57, 7.21$ Hz, 1 H), 3.76 (t, $J=7.70$ Hz, 1 H), 7.56 - 7.92 (m, 10 H); Calcd for C₂₆H₂₁ClF₆O₂ (M+Na) 537.11, Found 537.

5 **Example 23****2-(2,5-Dichloro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid**

- 10 The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 2,5-dichloro-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, MeOD) δ ppm 0.97 - 1.08 (m, 6 H), 1.59 (dd, *J*=13.33, 6.72 Hz, 1 H), 1.78 (dt, *J*=13.76, 6.94 Hz, 1 H), 2.04 (dd, *J*=13.45, 7.34 Hz, 1 H), 3.85 (t, *J*=7.70 Hz, 1 H), 7.39 - 7.87 (m, 10 H); Calcd for C₂₅H₂₁Cl₂F₃O₂ (M+Na) 503.09, Found 503.
- 15


Example 24**2-(4-Methoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid**

5


The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 4-methoxy-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (400 MHz, MeOD) δ ppm 0.79 - 0.90 (m, 6 H), 1.46 (ddd, $J=13.33$, 6.85, 6.72 Hz, 1 H), 1.64 (ddd, $J=13.63$, 7.09, 6.91 Hz, 1 H), 1.87 - 1.98 (m, 1 H), 3.67 - 3.77 (m, 4 H), 6.87 - 6.95 (m, 2 H), 7.43 - 7.54 (m, 4 H), 7.59 - 7.69 (m, 3 H), 7.73 (d, $J=8.07$ Hz, 2 H); Calcd for C₂₆H₂₅F₃O₃ (M+Na) 465.18, Found 465.

Example 25

15 **2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pent-4-enoic acid**



a) **(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid methyl ester**




20

5 A mixture of (3,5-dihydroxy-phenyl)-acetic acid methyl ester (2g, 11 mmol), N-phenyl-bis-(trifluoromethanesulfonimide) (8.6g, 24.2mmol) in THF (100mL) under N₂ was added Et₃N (6.1mL, 44mmol). The reaction mixture was heated at 50 °C for 48h. After cooling to room temperature, the solution was concentrated and purified by column chromatography to give bis-triflate intermediate.

10 The bis-triflate intermediate was under the same Suzuki coupling procedure as in the preparation of **Example 1h** gave the title compound **25a**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 3.74 (s, 3 H), 3.79 (s, 2 H), 7.55 (s, 2 H), 7.69 - 7.78 (m, 9 H); Calcd for C₂₃H₁₆F₆O₂ (M+H) 439.11, Found 439.

15 **b) 2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pent-4-enoic acid methyl ester**



20 The above intermediate **25a** was subjected to the same alkylation procedure as described in the preparation of **Example 1e** to give the title compound **25b**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 1.68 (s, 3 H), 2.46 (dd, *J*=14.67, 6.36 Hz, 1 H), 2.86 (dd, *J*=14.55, 9.17 Hz, 1 H), 3.60 (s, 3 H), 3.90 (dd, *J*=9.17, 6.48 Hz, 1 H), 4.69 (d, *J*=16.38 Hz, 2 H), 7.51 (d, *J*=1.47 Hz, 2 H), 7.58 - 7.68 (m, 9 H); Calcd for C₂₇H₂₂F₆O₂ (M+H) 493.15, Found 493.


25

c) 2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pent-4-enoic acid

- 5 A mixture of intermediate **25b** (40mg, 0.081mmol) and NaOH (2N in H₂O, 0.121mL, 0.243mmol) in THF-MeOH (0.6mL-0.6mL) was stirred for 18h and concentrated. CH₂Cl₂ and water were added, and the mixture was acidified with 1N HCl. The organic phase was separated and the aqueous phase was extracted with CH₂Cl₂. The combined organic layers were dried, concentrated, and purified by column chromatography to give 33mg
 10 (85%) of the title compound as a white solid; ¹H NMR (400 MHz, MeOD) δ ppm 1.77 (s, 3 H), 2.59 (dd, *J*=14.43, 7.09 Hz, 1 H), 2.91 (dd, *J*=14.55, 8.44 Hz, 1 H), 4.01 (t, *J*=7.83 Hz, 1 H), 4.76 (s, 2 H), 7.70 (d, *J*=1.47 Hz, 2 H), 7.77 (d, *J*=8.31 Hz, 4 H), 7.82 - 7.90 (m, 5 H); Calcd for C₂₆H₂₀F₆O₂ (M+H) 479.14, Found 479.


15 **Example 26**

2-(3-Fluoro-4-trifluoromethoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



20


a) **2-(3,5-Bis-benzyloxy-phenyl)-4-methyl-pent-4-enoic acid methyl ester**



5

A 2M solution of LDA in THF-heptane-ethylbenzene (21.5 mL, 43.0 mmol) was added dropwise over 12 min to a stirred solution of (3,5-bis-benzyloxyphenyl)acetic acid methyl ester (prepared in **Example 1**, step (a)) (13.0 g, 35.9 mmol) in THF (80 mL) at -78 °C under a nitrogen atmosphere. The temperature was maintained below -70 °C for an additional 50 min, then 3-bromo-2-methylpropene (4.0 mL, 39.7 mmol) was added in one portion and the reaction mixture was warmed to 0 °C. After 2 h the mixture was concentrated *in vacuo*, diluted with sat. aq. NH₄Cl (100 mL) and extracted with EtOAc (100 mL). The organic layer was washed with brine (100 mL), dried (MgSO₄), concentrated *in vacuo* and purified by flash chromatography (silica, 0-10% EtOAc in petroleum ether) to afford the title product as a yellow oil (14.1g, 94 %). ¹H-NMR (400 MHz, CDCl₃): δ 7.42-7.25 (m, 10H), 6.58 (s, 2H), 6.52 (s, 1H), 5.02 (s, 4H), 4.74 (s, 1H), 4.66 (s, 1H), 3.74 (t, 1H), 3.64 (s, 3H), 2.79 (dd, 1H), 2.38 (dd, 1H), 1.70 (s, 3H).

b) 2-(3-Benzyloxy-5-hydroxy-phenyl)-4-methyl-pent-4-enoic acid methyl ester




5

10% Pd/C (Aldrich cat no 205699, 0.55 g) was added to a stirred solution of 2-(3,5-bis-benzyloxyphenyl)-4-methyl-pent-4-enoic acid methyl ester (14.1 g, 33.8 mmol) and NaOH (1.50 g, 37.5 mmol) in MeOH (180 mL) at room temperature. Stirring was continued for 1 h under H₂ (1 atm.) then the mixture was filtered through Celite, concentrated *in vacuo*, suspended in water (100 mL) and adjusted to pH 2 with 1M HCl. The mixture was extracted with EtOAc (2 x 180 mL); the combined organic layer was washed with brine (50 mL), dried (MgSO₄), concentrated *in vacuo* and purified by flash chromatography (silica, 0-30% EtOAc in petroleum ether) to give the title product as a yellow oil (7.40 g, 67 %) ¹H-NMR (400 MHz, CDCl₃): δ 7.41-7.25 (m, 5H), 6.54 (s, 1H), 6.43 (s, 1H), 6.38 (s, 1H), 5.01 (s, 3H), 4.74 (s, 1H), 4.67 (s, 1H), 3.70 (t, 1H), 3.65 (s, 3H), 2.78 (dd, 1H), 2.38 (dd, 1H), 1.71 (s, 3H); Mass spectrum (ESI, m/z) 325 (M-1)

c) 2-(3-Benzyloxy-5-trifluoromethanesulfonyloxy-phenyl)-4-methyl-pent-4-enoic acid methyl ester


20



Trifluoromethanesulfonic anhydride (3.3 mL, 10.1 mmol) was added dropwise to a stirred solution of 2-(3-benzyloxy-5-hydroxyphenyl)-4-methyl-pent-4-enoic acid methyl ester (4.5 g, 13.8 mmol) and pyridine (3.0 mL, 38.7 mmol) in DCM (80 mL) at 0 °C then warmed to room temperature. After 1 h, the mixture was washed with 1M HCl (50 mL), dried (MgSO₄) and concentrated *in vacuo* to afford the title product as an orange oil (6.10 g, 96 %). ¹H-NMR (400 MHz, CDCl₃): δ 7.42-7.30 (m, 5H), 6.97 (s, 1H), 6.85 (s, 1H),

5 6.78 (s, 1H), 5.05 (s, 3H), 4.75 (s, 1H), 4.64 (s, 1H), 3.77 (t, 1H), 3.66 (s, 3H), 2.77 (dd, 1H), 2.40 (dd, 1H), 1.69 (s, 3H).

d) 2-(5-Benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pent-4-enoic acid methyl ester




10

A mixture of 2-(3-benzyloxy-5-trifluoromethanesulfonyloxy-phenyl)-4-methyl-pent-4-enoic acid methyl ester (4.3 g, 9.4 mmol), 4-trifluoromethylphenylboronic acid (2.6 g, 13.7 mmol), K₂CO₃ solution (2M, 9.4 mL) and DME (50 mL) was purged with N₂ three times before adding Pd(PPh₃)₄ (400 mg, 0.3 mmol). The mixture was heated to 95 °C for 5 h (followed by HPLC). The reaction was diluted with EtOAc (200 mL) and then was washed successively with NaHCO₃ solution and brine. The organic layer was dried (MgSO₄), filtered and concentrated *in vacuo* to give the title compound as an oil. The residue was used crude in the next step.

20

e) 2-(5-Hydroxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoic acid methyl ester




5

10% Pd/C (Aldrich cat no 205699, 0.30 g) was added to a stirred solution of 2-(5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pent-4-enoic acid methyl ester (2.71 g, 5.96 mmol) in MeOH (75 mL) at room temperature. Stirring was continued for 2 days under H₂ (1 atm.). Then the mixture was filtered through Celite, concentrated *in vacuo* to give the compound (1.83 g, 84%) as a yellow oil.

¹H-NMR (400 MHz, CD₃Cl): δ 7.70-7.60 (m, 4H), 7.08 (t, 1H), 7.00-6.95 (m, 1H), 6.90-6.87 (m, 1H), 5.56 (br. s, 1H), 3.73-3.65 min (m, 4H), 2.04-1.92 (m, 1H), 1.76-1.64 (m, 1H), 1.56-1.42 (m, 1H), 0.92 (d, 6H).

15

f) 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid methyl ester




20

- 5 The title compound was prepared in 95 % yield from 2-(5-hydroxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoic acid methyl ester under the conditions described in step (c).

¹H-NMR (400 MHz, CD₃Cl): δ 7.77-7.64 (m, 4H), 7.57-7.54 (m, 1H), 7.40-7.35 (m, 1H), 7.31-7.28 (m, 1H), 3.77 (t, 1H), 3.70 (s, 3H), 3.07-1.95 (m, 1H), 1.75-1.65 (m, 1H),
10 1.54-1.40 (m, 1H), 0.93 (dd, 6H). Mass spectrum (ESI, m/z): 365 (M-H)

g) 2-(3-Fluoro-4-trifluoromethoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-ethyl-pentanoic acid methyl ester




15

- The title compound was prepared in 53 % yield from 4-methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid methyl ester and 3-fluoro-4-trifluoromethoxyphenylboronic acid under the conditions described
20 in step (d).

¹H-NMR (400 MHz, CD₃Cl): δ ¹H-NMR (400 MHz, CDCl₃): δ 7.73 (br, s, 4H), 7.63 (t, 1H), 7.57 (t, 1H), 7.51 (t, 1H), 7.45 (m, 1H), 7.40 (m, 2H), 3.80 (m, 1H), 3.70 (s, 3H), 2.07 (m, 1H), 1.75 (m, 1H), 1.55 (m, 1H), 0.95 (d, 6H).

- 25 **h) 2-(3-Fluoro-4-trifluoromethoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-ethyl-pentanoic acid**

5




A mixture of 2-(3-fluoro-4-trifluoromethoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-ethyl-pentanoic acid methyl ester (2 mg) THF (1 mL), 10% LiOH in MeOH (0.3 mL) and H₂O (0.3 mL) was stirred at 30 °C for 3 h. The solution was concentrated *in vacuo* and the residue was diluted with H₂O and then acidified with concentrated HCl. The aqueous solution was extracted with DCM and filtered through polytetrafluoroethylene filter. The solution was concentrated *in vacuo* to give a solid residue. The solid was purified using reverse phase preparative HPLC (MeCN, H₂O) to give the title product (11 mg, 44 %).

¹H-NMR (CD₃Cl; 400 MHz): δ 7.70 (br. s, 4H), 7.61 (t, 1H), 7.55 (t, 1H), 7.51 (t, 1H), 7.43 (m, 1H), 7.38 (m, 2H), 3.79 (m, 1H), 2.05 (m, 1H), 1.76 (m, 1H), 1.55 (m, 1H), 0.93 (d, 6H).

20 Intermediate A

4-Methyl-2-(5-(2-(3-fluoro-4-(trifluoromethanesulfonyloxy-3',5'-bis-trifluoromethyl-biphenyl-3-yl)-pentanoic acid methyl ester



5

The title compound was prepared in 71 % yield from 2-(3-benzyloxy-5-trifluoromethanesulfonyloxy-phenyl)-4-methyl-pent-4-enoic acid methyl ester (prepared in **Example 26**, step (c)) under the conditions described in **Example 26**, step (d-f)) using 3,5-*bis*-trifluoromethylphenylboronic acid in step (d).


10

¹H-NMR (400 MHz, CD₃Cl): δ 7.96 (s, 2H), 7.94 (s, 1H), 7.55 (m, 1H), 7.38 (m, 2H), 3.80 (t, 1H), 3.71 (s, 3H), 2.01 (m, 1H), 1.71 (m, 1H), 1.49 (m, 1H), 0.94 (d, 6H).

Intermedaite B

15

2-(3'-Fluoro-5-trifluoromethanesulfonyloxy-5'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoic acid methyl ester




- 5 The title compound was prepared in 66% yield from 2-(3-benzyloxy-5-trifluoromethanesulfonyloxy-phenyl)-4-methyl-pent-4-enoic acid methyl ester (prepared in **Example 26**, step (c)) under the conditions described in **Example 26**, step (d-f)) using 3-fluoro-5-trifluoromethylphenylboronic acid in step (d).

¹H-NMR (400 MHz, CD₃Cl): δ ¹H-NMR (400 MHz, CDCl₃): δ 7.58 (s, 1H), 7.53 (m, 1H), 7.44 (dm, 1H, J= 9.1Hz) 7.39 (m, 1H), 7.36 (m, 1H), 7.33 (m, 1H), 3.78 (m, 1H),
 10 3.70 (s, 3H), 2.03 (m, 1H), 1.71 (m, 1H), 1.49 (m, 1H), 0.94 (d, 6H, J=6.8Hz).

Intermediate C


- 15 **2-(4'-Chloro-5-trifluoromethanesulfonyloxy-3'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoic acid methyl ester**




- 20 The title compound was prepared in 70 % yield from 2-(3-benzyloxy-5-trifluoromethanesulfonyloxy-phenyl)-4-methyl-pent-4-enoic acid methyl ester (prepared in **Example 26**, step (c)) under the conditions described in **Example 26**, step (d-f)) using 4-chloro-5-trifluoromethylphenylboronic acid in step (d).

Example 27

- 5 **2-(3-Fluoro-4-trifluoromethoxy-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid**



- 10 **a) 2-(3-Fluoro-4-trifluoromethoxy-3'',5''-bis-trifluoromethyl [1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester**



15

The title compound was prepared in 73 % yield from 4-methyl-2-(5-trifluoromethanesulfonyloxy-3',5'-bis-trifluoromethyl-biphenyl-3-yl)-pentanoic acid methyl ester (prepared in **Intermediate A**) and 3-fluoro-4-trifluoromethoxyphenylboronic acid under the conditions described in **Example 26**, step

20 (d).

5

b) 2-(3-Fluoro-4-trifluoromethoxy-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid


The title compound was prepared in 45 % yield from 2-(3-fluoro-4-trifluoromethoxy-3'',5''-bis-trifluoromethyl [1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester under the conditions described in **Example 26**, step (h).

¹H-NMR (400 MHz, CDCl₃): δ 7.99 (m, 2H), 7.89 (m, 2H), 7.67 (t, 1H), 7.57 (t, 1H), 7.54 (t, 1H), 7.43 (m, 1H), 7.39 (m, 2H), 3.81 (m, 1H), 2.07 (m, 1H), 1.75 (m, 1H), 1.55 (m, 1H), 0.96 (d, 6H).

15


Example 28

2-(3-Fluoro-4-chloro-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



20

a) 2-(3-Fluoro-4-chloro-3'',5''-bis-trifluoromethyl [1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester



5

The title compound was prepared in 65 % yield from 4-methyl-2-(5-trifluoromethanesulfonyloxy-3',5'-bis-trifluoromethyl-biphenyl-3-yl)-pentanoic acid methyl ester (prepared in **Intermediate A**) and 3-fluoro-4-chlorophenylboronic acid under the conditions described in **Example 26**, step (d).

10

b) 2-(3-Fluoro-4-chloro-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid

The title compound was prepared in 37 % yield from 2-(3-fluoro-4-chloro-3'',5''-bis-trifluoromethyl [1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester under the conditions described in **Example 26**, step (h).


15

¹H-NMR (400 MHz, CDCl₃): δ 8.00 (m, 2H), 7.90 (m, 2H), 7.62 (t, 1H), 7.58 (t, 1H), 7.54 (t, 1H), 7.49 (dd, 1H), 7.41 (dd, 1H), 7.34 (dd, 1H), 3.84 (m, 1H), 2.10 (m, 1H), 1.78 (m, 1H), 1.59 (m, 1H), 0.97 (d, 6H)

20


Example 29

2-(3''-Fluoro-3,5,5''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



5

a) 2-(3''-Fluoro-3,5,5''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester



10

The title compound was prepared in 91 % yield from 2-(3'-fluoro-5-trifluoromethanesulfonyloxy-5'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoic acid methyl ester (prepared in **Intermediate B**) and 3,5-bis-trifluoromethylphenylboronic acid under the conditions described in **Example 26**, step (d).

15


b) 2-(3''-Fluoro-3,5,5''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid

- 5 The title compound was prepared in 33 %yield from 2-(3''-Fluoro-3,5,5''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester under the conditions described in **Example 26**, step (h).

¹H-NMR (400 MHz, CDCl₃): δ 8.01 (br.s, 1H), 7.91 (br. s, 1H), 7.63 (br. d, 2H, J=6.1Hz), 7.59 (br. s, 2H), 7.50 (br. d, 1H, J=8.3Hz), 7.37 (br. d, 1H, J=8.1Hz), 3.90 (br.
10 s, 1H), 2.14 (br. s, 1H), 1.80 (br. s, 1H), 1.60 (br. s, 1H), 0.97 (br. s, 6H)


Example 30

2-(3,5-bis-trifluoromethyl-4''-chloro-3''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



15

a) 2-(3,5-bis-trifluoromethyl-4''-chloro-3''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester



20

- 5 The title compound was prepared in 50 % yield from 2-(4'-chloro-5-trifluoromethanesulfonyloxy-3'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoic acid methyl ester (prepared in **Intermediate C**) and 3,5-*bis*-trifluoromethylphenylboronic acid under the conditions described in **Example 26**, step (d).


10 **b) 2-(3,5-bis-trifluoromethyl-4''-chloro-3''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid**

The title compound was prepared in 61 % yield from 2-(3,5-*bis*-trifluoromethyl-4''-chloro-3''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester under the conditions described in **Example 26**, step (h).

- 15 ¹H-NMR (400 MHz, CDCl₃): δ 7.98 (s, 2H), 7.88 (1H), 7.85 (1H), 6.65 (d, 1H), 7.56-7.50 (m, 3H), 7.25 (1H), 3.73 (t, 1H), 1.97 (m, 1H), 1.69 (m, 1H), 1.47 (m, 1H), 0.86 (m, 6H).


Example 31

20 **2-(3,5-Difluoro-4''-chloro-3''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid**



25 **a) 2-(3,5-Difluoro-4''-chloro-3''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester**

5




The title compound was prepared in 40 % yield from 2-(4'-chloro-5-trifluoromethanesulfonyloxy-3'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoic acid methyl ester (prepared in **Intermedaite C**) and 3,5-difluorophenylboronic acid under the conditions described in **Example 26**, step (d). ¹H-NMR (400 MHz, CDCl₃): δ 7.90 (s, 1H), 7.72 (d, 1H), 7.60 (d, 1H), 7.58 (s, 1H), 7.52 (s, 2H), 7.15 (d, 2H), 6.83 (t, 1H), 3.80 (t, 1H), 3.70 (s, 3H), 2.04 (m, 1H), 1.73 (m, 1H), 0.95 (d, 6H).

b) 2-(3,5-Difluoro-4''-chloro-3''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid

The title compound was prepared in 99 % yield from 2-(3,5-difluoro-4''-chloro-3''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester under the conditions described in **Example 26**, step (h). ¹H-NMR (400 MHz, CDCl₃): δ 7.89 (d, 2H), 7.70 (dd, 1H), 7.61 (s, 1H), 7.59 (m, 1H), 7.53 (m, 2H), 7.13 (m, 2H), 6.84 (tt, 1H), 3.84 (t, 1H), 2.10 (m, 1H), 1.77 (m, 1H), 1.58 (m, 1H), 0.97 (d, 6H).


Example 32

4-Methyl-2-(3,5,3''-trifluoro-5''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid



5

a) 4-Methyl-2-(3,5,3''-trifluoro-5''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid methyl ester



10

The title compound was prepared in 95 % yield from 2-(3'-fluoro-5'-trifluoromethanesulfonyloxy-5'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoic acid methyl ester (prepared in **Intermediate B**) and 3,5-difluorophenylboronic acid under the conditions described in **Example 26**, step (d).

15

b) 4-Methyl-2-(3,5,3''-trifluoro-5''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid


- 5 The title compound was prepared in 36 % yield from 4-methyl-2-(3,5,3''-trifluoro-5''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid methyl ester

under the conditions described in **Example 26**, step (h).

- ¹H-NMR (400 MHz, CDCl₃): δ 7.63 (br. s, 1H), 7.60 (br. s, 1H), 7.55 (br. s, 1H), 7.49 (br. d, 1H, J=9.1Hz), 7.13 (dm, 2H, J=6.3Hz), 6.84 (tm, 1H, J=8.8Hz), 3.87 (br. s, 1H),
10 2.10 (br. s, 1H), 1.80 (br. s, 1H), 1.58 (br. s, 1H), 0.96 (d, 6H, J= 6.1Hz).


Example 33

2-(3,5-Difluoro-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



15

a) 2-(3,5-Difluoro-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester



20

5

The title compound was prepared in 72 % yield from 4-methyl-2-(5-trifluoromethanesulfonyloxy-3',5'-bis-trifluoromethyl-biphenyl-3-yl)-pentanoic acid methyl ester (prepared in **Intermedaite A**) under the conditions described in **Example 26**, step (d).

10

b) 2-(3,5-Difluoro-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid


The title compound was prepared in 61 % yield from 2-(3,5-difluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester under the conditions described in **Example 26**, step (h).

15

¹H-NMR (400 MHz, CD₃Cl): δ 8.00 (m, 2H), 7.90 (m, 1H), 7.62 (m, 1H), 7.58 (m, 1H), 7.55 (m 1H), 7.14 (m, 2H), 6.84 (m, 1H), 3.87 (m, 1H), 2.10 (m, 1H), 1.80 (m, 1H), 1.59 (m, 1H), 0.97 (d, 6H).


20 **Example 34**

2-(3-Cyano-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl pentanoic acid.



25

- 5 a) 2-(3-Cyano-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester



The title compound was prepared in 48 % yield from 4-methyl-2-(5-trifluoromethanesulfonyloxy-3',5'-bis-trifluoromethyl-biphenyl-3-yl)-pentanoic acid methyl ester (prepared in **Intermedaite A**) and 3-cyanophenylboronic acid under the conditions described in **Example 26**, step (d).

- 15 b) 2-(3-Cyano-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl pentanoic acid.


The title compound was prepared in 68 % yield from 2-(3-cyano-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester under the conditions described in **Example 26**, step (h).

20 ¹H-NMR (400 MHz, CDCl₃): δ 8.02 (m, 2H), 7.91 (m, 2H), 7.85 (m, 1H), 7.69 (m, 1H), 7.5-7.7 (m, 4H), 3.86 (m, 1H), 2.10 (m, 1H), 1.80 (m, 1H), 1.57 (m, 1H), 0.97 (d, 6H).

Example 35


4-Methyl-2-(3,5,4''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid

25



5

a) **4-Methyl-2-(3,5,4''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid methyl ester**



10

The title compound was prepared in 68 % yield from 4-methyl-2-(5-trifluoromethanesulfonyloxy-3',5'-bis-trifluoromethyl-biphenyl-3-yl)-pentanoic acid methyl ester (prepared in **Intermediate A**) and 3,5-bis-trifluoromethylphenylboronic acid under the conditions described in **Example 26**, step (d).

15

b) **2-(3,5-Difluoro-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid.**


- 5 The title compound was prepared in 65 % yield from 4-methyl-2-(3,5,4''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid methyl ester under the conditions described in **Example 26**, step (h).

¹H-NMR (400 MHz, CDCl₃): δ 8.00 (m, 4H), 7.92 (m, 2H), 7.65 (m, 1H), 7.61 (m, 2H), 3.88 (t, 1H), 2.13 (m, 1H), 1.79 (m, 1H), 1.61 (m, 1H), 0.97 (d, 6H).

10


Example 36

2-(5-Cyano-3-fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



15

a) 2-(5-Cyano-3-fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester



20

- 5 The title compound was prepared in 21 % yield from 4-methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid methyl ester (prepared in **Example 26**, step(f)) and 3-fluoro-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzonitrile under the conditions described in **Example 26**, step (d).

10

b) 2-(5-Cyano-3-fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid


- The title compound was prepared in 71 % yield from 2-(5-cyano-3-fluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester under the conditions described in **Example 26**, step (h).

¹H-NMR (400 MHz, CD₃Cl): δ 7.64 (q, 4H), 7.10 (s, 1H), 6.97 (s, 1H), 6.94-6.92 (m, 2H), 6.89 (s, 1H), 6.82 (d, 1H), 3.70 (t, 1H), 2.01-1.93 (m, 1H), 1.75-1.68 (m, 1H), 1.56-1.51 (m, 1H), 0.92 (d, 6H).

20

Example 37


2-(4-Cyano-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid.



25

5

a) 2-(4-Cyano-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester



10

The title compound was prepared in 57 % yield from 4-methyl-2-(5-trifluoromethanesulfonyloxy-3',5'-bis-trifluoromethyl-biphenyl-3-yl)-pentanoic acid methyl ester (prepared in **Intermediate A**) and 4-cyanophenylboronic acid under the conditions described in **Example 26**, step (d).

15

b) 2-(3-Cyano-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid

The title compound was prepared in 37 % yield from 2-(4-cyano-3'',5''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid methyl ester under the conditions described in **Example 26**, step (h).


20

¹H-NMR (400 MHz, CDCl₃): δ 8.00 (m, 2H), 7.91 (m, 2H), 7.77 (d, 2H), 7.72 (d, 2H), 7.66 (t, 1H), 7.63 (t, 1H), 7.58 (t, 1H), 3.86 (m, 1H), 2.11 (m, 1H), 1.78 (m, 1H), 1.58 (m, 1H), 0.96 (d, 6H).

Example 38


25

5 **4-Methyl-2-(3,5,3'',5''-tetrafluoro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid**



a) **2-(3,5-Dihydroxy-phenyl)-4-methyl-pentanoic acid methyl ester**


10



10% Pd/C (Aldrich cat no 205699, 214 mg) was added to a stirred solution of 2-(3,5-*bis*-benzyloxyphenyl)-4-methyl-pent-4-enoic acid methyl ester (2.14 g, 5.1 mmol) and NaOH (225 mg, 5.6 mmol) in MeOH (30 mL) at room temperature. Stirring was continued for 2 h under H₂ (1 atm.) then the mixture was filtered through Celite, concentrated *in vacuo*, suspended in water (15 mL) and adjusted to pH 2 with 1M HCl. The mixture was extracted with DCM (3 x 60 mL); the combined organic layer was washed with brine (20 mL), dried (MgSO₄), concentrated *in vacuo* and purified by flash chromatography (silica, 0-20% EtOAc in petroleum ether) to give the title product as a white syrup (1.1 g, 90 %).


b) **2-(3,5-Bis-trifluoromethanesulfonyloxy-phenyl)-4-methyl-pentanoic acid methyl ester**

5



Trifluoromethanesulfonic anhydride (208 μ L, 1.26 mmol) was added dropwise to a stirred solution of 2-(3,5-dihydroxy-phenyl)-4-methyl-pentanoic acid methyl ester (50 mg, 0.21 mmol) and pyridine (195 μ L, 2.52 mmol) in DCM (5 mL) at 0 °C then warmed to room temperature. After 1 h, the mixture was washed with 1M HCl (2 mL), dried (MgSO₄) and concentrated *in vacuo* to afford the title product as an orange oil (104 mg, 99 %). The product was used without further purification in the next step

c) **4-Methyl-2-(3,5,3'',5''-tetrafluoro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid methyl ester**



A mixture of 2-(3,5-*bis*-trifluoromethanesulfonyloxy-phenyl)-4-methyl-pentanoic acid methyl ester (104 mg, 0.21 mmol), 3,5-difluorophenylboronic acid (82 mg, 0.52 mmol), K₂CO₃ solution (2M, 310 μ L) and DME (2 mL) was purged with N₂ three times before adding Pd(PPh₃)₄ (10 mg, 0.01 mmol). The mixture was heated at 95 °C overnight

5 (followed by HPLC). The reaction was diluted with EtOAc (5 mL) and then was washed successively with NaHCO₃ solution and brine. The organic layer was dried (MgSO₄), filtered and concentrated *in vacuo* to give the title compound as an oil. The residue was purified by flash chromatography (0-15% EtOAc in Petroleum ether) to give the title compound (75 mg, 84%) as a yellow oil.

10 ¹H-NMR (400 MHz, CDCl₃): δ 7.58 (br. s, 1H), 7.52 (br. s, 2H), 7.26, (br. s, H), 7.15-7.05 (m, 3H), 6.83, (m, 2H), 3.79 (br. s, 1H), 3.70 (s, 3H), 2.06 (br. s, 1H), 1.73 (br. s, 1H), 1.52 (br. s, 1H), 0.94 (m, 6H).


b) 4-Methyl-2-(3,5,3'',5''-tetrafluoro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid

15 The title compound was prepared in 74% yield from 4-methyl-2-(3,5,3'',5''-tetrafluoro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid methyl ester under the conditions described in **Example 26** step (h).

20 ¹H-NMR (400 MHz, CDCl₃): δ 7.58 (br. s, 1H), 7.52 (br. s, 2H), 7.25 (m, 1H), 7.19 (br. s, 3H), 6.82 (m, 2H), 3.80 (m, 1H), 2.06 (m, 1H), 1.76 (m, 1H), 1.55 (m, 1H), 0.95 (m, 6H).


Example 39

(R)-2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



5


a) 5-Benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid



- 10 To a solution of (5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid ethyl ester (120 g, 0.29 mol) in THF (1.2 L) was added water (240 mL), LiOH·H₂O (16 g, 0.32 mol) and stirred at room temperature for 16 h. The solution was filtered and concentrated in vacuo to remove THF. The resulting thick liquid was acidified to pH 2 by adding 2N aqueous HCl solution and the white suspension was mechanically stirred for 1h at room
- 15 temperature. The wet white product was recovered after filtration and dissolved in EtOAc (500 mL). The organic layer was separated from water, dried (MgSO₄) and concentrated in vacuo to obtain (5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid (105 g, 94%).

- ¹H-NMR (d₆-DMSO): δ 3.64 (s, 2H), 5.18 (s, 2H), 7.02 (s, 1H), 7.24 (d, 2H), 7.34-7.50 (m, 5H), 7.81 (d, 2H), 7.89 (d, 2H), 12.25 (bs, 0.6H); Calcd for C₂₂H₁₇F₃O₃ (M+H) 387.11, Found 387.1.
- 20

5 **b) 4-Benzyl-3-[2-(5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetyl]-oxazolidin-2-one**



10 To a mechanically stirred solution of (5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid (20 g, 52 mmol) in THF (104 mL) at -78°C was added N-methyl morpholine (6.3 mL, 57 mmol) and trimethylacetyl chloride (7.0 mL, 57 mmol) maintaining the internal temperature below -70°C . This mixture was stirred at -78°C for 15 minute and 0°C at 1h. The white solid was filtered off to receive the anhydride in the filtrate which

15 was cooled back to -78°C . In a separate flask, to a solution of (R)-(+)-4-benzyl-2-oxazolidinone (9.6 g, 54.4 mmol) in THF (109 mL) at -78°C was added nBuLi (1.6M in hexanes, 34 mL, 54.4 mol), drop-wise, maintaining the internal temperature below -70°C and stirred at -78°C for 45 min. This metalated chiral auxiliary was cannulated to the anhydride at -78°C and warmed to 0°C over 1.5h. The resulting mixture was stirred


20 further at 0°C for 30 minutes and quenched by adding excess saturated aqueous NH_4Cl solution. The solution was diluted with EtOAc (200 mL) and the organic phase was washed with saturated aqueous NaHCO_3 solution (3 x 100 mL) and brine (2 x 100 mL). The solution was dried over MgSO_4 and the solvent was removed in vacuo. The crude material was purified by ISCO silica gel column chromatography to receive 20.3 g (72%)

25 of 4-benzyl-3-[2-(5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetyl]-oxazolidin-2-one as a white solid.

$^1\text{H-NMR}$ (CDCl_3): δ 2.76 (dd, 1H), 3.26 (dd, 1H), 4.19 (m, 2H), 4.35 (q, 2H), 4.69 (m, 1H), 5.13 (s, 2H), 7.04-7.46 (m, 13H), 7.67 (s, 4H); Calcd for $\text{C}_{32}\text{H}_{26}\text{F}_3\text{NO}_4$ (M+H) 546.18, Found 546.3.

5

c) 4-Benzyl-3-[2-(5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pent-4-enoyl]-oxazolidin-2-one




- 10 To a colorless solution of 4-benzyl-3-[2-(5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetyl]-oxazolidin-2-one (6.0 g, 11.00 mmol) in dry THF (22 mL) at -78°C was added NaHMDS (1 M in THF solution, 12.11 mL, 12.11 mmol), drop-wise, maintaining the internal temperature below -75°C . The resulting red solution was stirred at -78°C for 30 minutes. To this was added 3-bromo-2-methyl propene (4.44 mL, 44 mmol) maintaining
- 15 the temperature below -75°C . When the addition was at near completion, the system turned to green. At this point the dry-ice bath was quickly removed and replaced with wet-ice bath and completed the addition. The reaction mixture was stirred further at 0°C for 30 min and quenched with saturated aqueous NH_4Cl solution. The system was diluted with EtOAc (100 mL) and the organic phase was washed with saturated aqueous
- 20 NaHCO_3 solution (3 x 50 mL) and dried (MgSO_4). Solvent was removed in vacuo and the crude mixture was purified by ISCO silica gel column to yield 4-benzyl-3-[2-(5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pent-4-enoyl]-oxazolidin-2-one (6.3 g, 95 %).

$^1\text{H-NMR}$ (CDCl_3): δ 1.80 (s, 3H), 2.46 (dd, 1H), 2.75 (dd, 1H), 3.05 (dd, 1H), 3.32 (dd, 1H), 4.08 (m, 2H), 4.59 (m, 1H), 4.80 (d, 2H), 5.13 (s, 2H), 5.48 (dd, 1H), 7.11 (d, 2H), 7.21-7.49 (m, 11H), 7.67 (s, 4H); Calcd for $\text{C}_{36}\text{H}_{32}\text{F}_3\text{NO}_4$ ($\text{M}+\text{H}$) 600.23, Found 600.3.

25

5 **d) 4-Benzyl-3-[2-(5-hydroxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoyl]-oxazolidin-2-one**




To a solution of 4-benzyl-3-[2-(5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pent-4-enoyl]-oxazolidin-2-one (6.7 g, 11.2 mmol) in MeOH (150 mL) was added 10% Pd/C (670 mg, 10 w %). The black suspension was hydrogenated at 45-45 psi overnight. The mixture was filtered through celite and the solvent was

15 removed in vacuo to obtain relatively pure 4-benzyl-3-[2-(5-hydroxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoyl]-oxazolidin-2-one (5.4 g, 93 %).

¹H-NMR (CDCl₃): δ 0.94 (d, 3H), 0.98 (d, 3H), 1.54 (m, 1H), 1.74 (m, 1H), 2.12 (m, 1H), 2.79 (dd, 1H), 3.36 (dd, 1H), 4.11 (m, 2H), 4.62 (m, 1H), 5.25 (t, 1H), 6.97 (m, 2H), 7.21-7.37 (m, 6H), 7.67 (s, 4H); Calcd for C₂₉H₂₈F₃NO₄ (M+H) 512.20, Found 512.3.

20

e) Trifluoro-methanesulfonic acid 5-[1-(4-benzyl-2-oxo-oxazolidine-3-carbonyl)-3-methyl-butyl]-4'-trifluoromethyl-biphenyl-3-yl ester




25

To a solution of 4-benzyl-3-[2-(5-hydroxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoyl]-oxazolidin-2-one (32 g, 62.6 mmol) in dichloromethane (170 mL) was added pyridine (15.0 mL). The system was cooled to 0 °C. To this cold solution was added

5 trifluoromethanesulfonic anhydride (16 mL, 94 mmol) maintaining the internal temperature below 5 °C and stirred for further 0.5 h at 0 °C. This reaction mixture was poured to a mixture of 1 N HCl (100 mL), and wet-ice (25 g) and stirred for 0.5 h. The aqueous layer was extracted with dichloromethane (2 x 100 mL). Combined fractions were washed with water (2 x 100 mL), saturated aqueous NaHCO₃ solution (2 x 100 mL),
 10 and brine (2 x 100 mL). The organics were dried (MgSO₄) and concentrated in vacuo to receive a reddish liquid which was purified by ISCO column chromatography to receive trifluoro-methanesulfonic acid 5-[1-(4-benzyl-2-oxo-oxazolidine-3-carbonyl)-3-methyl-butyl]-4'-trifluoromethyl-biphenyl-3-yl ester (34 g, 84%).

¹H-NMR (CDCl₃): δ 0.96 (d, 3H), 0.98 (d, 3H), 1.52 (m, 1H), 1.77 (m, 1H), 2.13 (m, 1H), 2.79 (dd, 1H), 3.37 (dd, 1H), 4.14 (m, 2H), 4.67 (m, 1H), 5.33 (t, 1H), 7.20-7.38 (m, 7H), 7.70 (m, 5H); Calcd for C₃₀H₂₇F₆NO₆S (M+H) 644.15, Found 644.2.

f) 4-Benzyl-3-[2-(4,4''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoyl]-oxazolidin-2-one



20

A mixture of trifluoro-methanesulfonic acid 5-[1-(4-benzyl-2-oxo-oxazolidine-3-carbonyl)-3-methyl-butyl]-4'-trifluoromethyl-biphenyl-3-yl ester (4.03 g, 6.27 mmol), 4-(trifluoromethyl)phenylboronic acid (1.34 g, 7.05 mmol), 1,2-dimethoxyethane (24 mL) and aqueous Na₂CO₃ (2 M, 3.2 mL, 6.4 mmol) was stirred while purging N₂ at room temperature for 10 min. To this system was added Pd(Ph₃)₄ (1.45 g, 1.25 mmol) and heated to reflux (95 °C) for 1 h. The red-brown mixture was diluted with EtOAc (50 mL) and washed with saturated aqueous NaHCO₃ solution (3 x 50 mL) and brine (2 x 50 mL). The organic fraction was dried (Na₂SO₄) and concentrated

in vacuo. The crude mixture was purified by ISCO column chromatography to obtain 4-benzyl-3-[2-(4,4''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoyl]-oxazolidin-2-one (3.2 g, 79%).

¹H-NMR (CDCl₃): δ 0.97 (d, 3H), 0.99 (d, 3H), 1.58 (m, 1H), 1.80 (m, 1H), 2.17 (m, 1H), 2.79 (dd, 1H), 3.39 (dd, 1H), 4.12 (m, 2H), 4.65 (m, 1H), 5.35 (t, 1H), 7.22-7.37 (m, 5H), 7.68-7.76 (m, 11H); Calcd for C₃₆H₃₁F₆NO₃ (M+H) 640.22, Found 640.3.

g) (R)-2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid


To a solution of 4-benzyl-3-[2-(4,4''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoyl]-oxazolidin-2-one (3.66 g, 5.7 mmol) in THF (24 mL) was added water (8 mL). The system was cooled to 0 °C. To this cold solution was added LiOH·H₂O (240 mg, 5.7 mmol) and 30% H₂O₂ (1.95 mL, 17.2 mmol,) and stirred at 0 °C for 15 min. The excess H₂O₂ was quenched by adding 1.5 M aqueous Na₂SO₃ solution (11.5 mL, 17.2 mmol) and stirred at room temperature for 10 min. The organic solvent was removed in vacuo. The resulting liquid was acidified to pH = 2 by adding 1 N aqueous HCl solution.

The aqueous layer was extracted with EtOAc (3 x 50 mL) and dried (MgSO₄). The mixture was concentrated in vacuo to receive a crude mixture which was purified by ISCO silica gel column chromatography to yield (R)-2-(4,4''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid (2.5 g, 92 %).

¹H-NMR (CDCl₃): δ 0.96 (d, 6H), 1.59 (m, 1H), 1.79 (m, 1H), 2.08 (m, 1H), 3.83 (t, 1H), 7.58 (d, 2H), 7.69 (t, 1H), 7.72 (s, 8H); Calcd for C₂₆H₂₂F₆O₂ (M+H) 481.15, Found 481.2.

Example 40


(S)-2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



5

a) 4-Benzyl-3-[2-(5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetyl]-oxazolidin-2-one

10




The title compound was prepared from (5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetic acid and (S)-(-)-4-benzyl-2-oxazolidinone following the same procedure as for the synthesis of compound **39a**.

15

$^1\text{H-NMR}$ (CDCl_3): δ 2.76 (dd, 1H), 3.26 (dd, 1H), 4.19 (m, 2H), 4.35 (q, 2H), 4.69 (m, 1H), 5.13 (s, 2H), 7.04-7.46 (m, 13H), 7.67 (s, 4H); Calcd for $\text{C}_{32}\text{H}_{26}\text{F}_3\text{NO}_4$ (M+H) 546.18, Found 546.3.

b) 4-Benzyl-3-[2-(5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pent-4-enoyl]-oxazolidin-2-one

20




113

5

The title compound was prepared from 4-benzyl-3-[2-(5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-acetyl]-oxazolidin-2-one (**40a**) following the same procedure as for the synthesis of compound **39b**.

¹H-NMR (CDCl₃): δ 1.80 (s, 3H), 2.46 (dd, 1H), 2.75 (dd, 1H), 3.05 (dd, 1H), 3.32 (dd, 1H), 4.08 (m, 2H), 4.59 (m, 1H), 4.80 (d, 2H), 5.13 (s, 2H), 5.48 (dd, 1H), 7.11 (d, 2H), 7.21-7.49 (m, 11H), 7.67 (s, 4H); Calcd for C₃₆H₃₂F₃NO₄ (M+H) 600.23, Found 600.3.

c) 4-Benzyl-3-[2-(5-hydroxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pentanoyl]-oxazolidin-2-one




15

The title compound was prepared from 4-benzyl-3-[2-(5-benzyloxy-4'-trifluoromethyl-biphenyl-3-yl)-4-methyl-pent-4-enoyl]-oxazolidin-2-one (**40b**) following the same procedure as for the synthesis of compound **39c**.

¹H-NMR (CDCl₃): δ 0.94 (d, 3H), 0.98 (d, 3H), 1.54 (m, 1H), 1.74 (m, 1H), 2.12 (m, 1H), 2.79 (dd, 1H), 3.36 (dd, 1H), 4.11 (m, 2H), 4.62 (m, 1H), 5.25 (t, 1H), 6.97 (m, 2H), 7.21-7.37 (m, 6H), 7.67 (s, 4H); Calcd for C₂₉H₂₈F₃NO₄ (M+H) 512.20, Found 512.3.

d) Trifluoro-methanesulfonic acid 5-[1-(4-benzyl-2-oxo-oxazolidine-3-carbonyl)-3-methyl-butyl]-4'-trifluoromethyl-biphenyl-3-yl ester

25




5

The title compound was prepared from 4-benzyl-3-[2-(5-hydroxy-4'-trifluoromethyl-
 10 biphenyl-3-yl)-4-methyl-pentanoyl]-oxazolidin-2-one (**40c**) following the same procedure
 as for the synthesis of compound **39d**.

¹H-NMR (CDCl₃): δ 0.96 (d, 3H), 0.98 (d, 3H), 1.52 (m, 1H), 1.77 (m, 1H), 2.13 (m,
 1H), 2.79 (dd, 1H), 3.37 (dd, 1H), 4.14 (m, 2H), 4.67 (m, 1H), 5.33 (t, 1H), 7.20-7.38 (m,
 7H), 7.70 (m, 5H); Calcd for C₃₀H₂₇F₆NO₆S (M+H) 644.15, Found 644.2.

15

**e) 4-Benzyl-3-[2-(4,4''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-
 pentanoyl]-oxazolidin-2-one**



20 The title compound was prepared from trifluoro-methanesulfonic acid 5-[1-(4-benzyl-2-
 oxo-oxazolidine-3-carbonyl)-3-methyl-butyl]-4'-trifluoromethyl-biphenyl-3-yl ester (**40d**)
 following the same procedure as for the synthesis of compound **39e**.

- 5 $^1\text{H-NMR}$ (CDCl_3): δ 0.97 (d, 3H), 0.99 (d, 3H), 1.58 (m, 1H), 1.80 (m, 1H), 2.17 (m, 1H), 2.79 (dd, 1H), 3.39 (dd, 1H), 4.12 (m, 2H), 4.65 (m, 1H), 5.35 (t, 1H), 7.22-7.37 (m, 5H), 7.68-7.76 (m, 11H); Calcd for $\text{C}_{36}\text{H}_{31}\text{F}_6\text{NO}_3$ (M+H) 640.22, Found 640.3.


10 **f) (S)-2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid**

The title compound was prepared from 4-benzyl-3-[2-(4,4''-bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoyl]-oxazolidin-2-one (**40e**) following the same procedure as for the synthesis of compound **39**.

- 15 $^1\text{H-NMR}$ (CDCl_3): δ 0.96 (d, 6H), 1.59 (m, 1H), 1.79 (m, 1H), 2.08 (m, 1H), 3.83 (t, 1H), 7.58 (d, 2H), 7.69 (t, 1H), 7.72 (s, 8H); Calcd for $\text{C}_{26}\text{H}_{22}\text{F}_6\text{O}_2$ (M+H) 481.15, Found 481.2.

Example 41

20 **(S)-2-(3,5-Difluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid**




- 25 The title compound was prepared from a Suzuki coupling of trifluoro-methanesulfonic acid 5-[1-(4-benzyl-2-oxo-oxazolidine-3-carbonyl)-3-methyl-butyl]-4'-trifluoromethyl-biphenyl-3-yl ester(intermediate **Example 40c**) with 3,5-difluoro-phenyl-boronic acid under the conditions described in **Example 40**; $^1\text{H NMR}$ (400 MHz, CHLOROFORM-D) δ ppm 0.96 (d, $J=6.60$ Hz, 6 H), 1.53 - 1.64 (m, 1 H), 1.78 (ddd, $J=13.88$, 7.21, 7.03 Hz, 1 H), 2.04 - 2.12 (m, 1 H), 3.83 (t, $J=7.70$ Hz, 1 H), 6.80 - 6.86 (m, 1 H), 7.11 - 7.16 (m, 2 H), 7.54 (d, $J=1.47$ Hz, 1 H), 7.58 (s, 1 H), 7.64 (t, $J=1.59$ Hz, 1 H), 7.71 (s, 4 H).

5

Example 42


(R)-2-(3,5-Difluoro-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



- 10 The title compound was prepared from a Suzuki coupling of trifluoro-methanesulfonic acid 5-[1-(4-benzyl-2-oxo-oxazolidine-3-carbonyl)-3-methyl-butyl]-4'-trifluoromethyl-biphenyl-3-yl ester (intermediate **Example 39d**) with 3,5-difluoro-phenyl-boronic acid under the conditions described in **Example 39**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 0.96 (d, *J*=6.60 Hz, 6 H), 1.53 - 1.80 (m, 2 H), 1.78 (dt, *J*=14.00, 7.06 Hz, 1 H),
 15 2.08 (dd, *J*=14.55, 7.46 Hz, 1 H), 3.83 (t, *J*=7.70 Hz, 1 H), 6.80 - 6.86 (m, 1 H), 7.13 (s, 1 H), 7.14 - 7.16 (m, 1 H), 7.54 (d, *J*=1.22 Hz, 1 H), 7.58 (s, 1 H), 7.64 (d, *J*=1.47 Hz, 1 H), 7.72 (s, 4 H).

Example 43

- 20 **(S)-4-Methyl-2-(3,5,4''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid**




5

The title compound was prepared from a Suzuki coupling of trifluoro-methanesulfonic acid 5-[1-(4-benzyl-2-oxo-oxazolidine-3-carbonyl)-3-methyl-butyl]-4'-trifluoromethyl-biphenyl-3-yl ester (intermediate **Example 40c**) with 3,5-bis-trifluoromethyl-phenyl-

10 boronic acid under the conditions described in **Example 40**; ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 0.94 - 0.99 (m, 6 H), 1.60 (dt, $J=13.45$, 6.72 Hz, 1 H), 1.79 (ddd, $J=13.82$, 7.09, 6.97 Hz, 1 H), 2.11 (dt, $J=13.76$, 7.67 Hz, 1 H), 3.86 (t, $J=7.83$ Hz, 1 H), 7.56 (d, $J=1.71$ Hz, 1 H), 7.63 - 7.68 (m, 2 H), 7.73 (s, 4 H), 7.90 (s, 1 H), 8.02 (s, 2 H).

15 **Example 44**

(R)-4-Methyl-2-(3,5,4''-tris-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid




20 The title compound was prepared from a Suzuki coupling of trifluoro-methanesulfonic acid 5-[1-(4-benzyl-2-oxo-oxazolidine-3-carbonyl)-3-methyl-butyl]-4'-trifluoromethyl-

- 5 biphenyl-3-yl ester (intermediate **Example 39d**) with 3,5-bis-trifluoromethyl-phenyl-boronic acid under the conditions described in **Example 39**; ^1H NMR (400 MHz, CHLOROFORM-D) δ ppm 0.97 (d, $J=6.60$ Hz, 6 H), 1.60 (ddd, $J=13.33, 6.60, 6.48$ Hz, 1 H), 1.79 (ddd, $J=13.88, 7.21, 7.03$ Hz, 1 H), 2.07 - 2.15 (m, 1 H), 3.86 (t, $J=7.70$ Hz, 1 H), 7.56 (s, 1 H), 7.65 (d, $J=13.69$ Hz, 2 H), 7.73 (s, 4 H), 7.90 (s, 1 H), 8.02 (s, 2 H).

10

Example 47

(R)-2-(4-Chloro-4''-trifluoromethyl-[1,1';3'1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



15

The title compound was synthesized using a similar procedure as described for the preparation of compound **39** using 4-chlorophenylboronic acid in **Example 39e**.


^1H -NMR (CDCl_3): δ 0.95 (d, 6H), 1.58 (m, 1H), 1.78 (m, 1H), 2.06 (m, 1H), 3.81 (t, 1H), 7.43 (m, 2H), 7.54 (m, 4H), 7.64 (t, 1H), 7.71 (s, 4H); Calcd for $\text{C}_{25}\text{H}_{22}\text{ClF}_3\text{O}_2$ (M+H)

20 447.13, Found 447.

Example 46

(R)-2-(4-Isopropyl-4''-trifluoromethyl-[1,1';3'1'']terphenyl-5'-yl)-4-methyl-pentanoic acid

25



5

The title compound was synthesized using a similar procedure as described for the preparation of compound **39** using 4-isopropylphenylboronic acid in **Example 39e**.


¹H-NMR (CDCl₃): δ 0.94 (d, 6H), 1.30 (d, 6H), 1.58 (m, 1H), 1.78 (m, 1H), 2.05 (m, 1H), 2.97 (m, 1H), 3.80 (t, 1H), 7.32 (d, 2H), 7.54 (m, 4H), 7.70 (m, 5H); Calcd for C₂₈H₂₉F₃O₂ (M+H) 455.21, Found 455.3.

10

Example 47

(R)-2-(4-Chloro-3-fluoro-4''-trifluoromethyl-[1,1';3'1'']terphenyl-5'-yl)-4-methyl-
pentanoic acid

15




The title compound was synthesized using a similar procedure as described for the preparation of compound **39** using 4-chloro-3-fluorophenylboronic acid in **Example 39e**.

¹H-NMR (CDCl₃): δ 0.96 (d, 6H), 1.58 (m, 1H), 1.77 (m, 1H), 2.07 (m, 1H), 3.83 (t, 1H), 7.33-7.64 (m, 6H), 7.71 (s, 4H); Calcd for C₂₅H₂₁ClF₄O₂ (M+H) 465.12, Found 465.

20

Example 48

5 **2-(4-Acetyl-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid**



The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester (intermediate **Example 1g**) with 4-acetyl-phenylboronic acid under the conditions


10 described in **Example 1**; ¹H NMR (300 MHz, MeOD) δ ppm 0.88 (dd, *J*=6.78, 2.26 Hz, 6 H), 1.42 - 1.52 (m, 1 H), 1.67 (ddd, *J*=13.75, 7.16, 6.97 Hz, 1 H), 1.91 - 2.01 (m, 1 H), 2.55 (s, 3 H), 3.76 (t, *J*=7.72 Hz, 1 H), 7.59 (t, *J*=5.84 Hz, 2 H), 7.66 - 7.81 (m, 7 H), 8.01 (d, *J*=8.29 Hz, 2 H);

Calcd for C₂₇H₂₅F₃O₃ (M+H) 455.18, Found 455.

15

Example 49

2-(3-Dimethylamino-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid




20 The title compound was prepared from a Suzuki coupling of 4-Methyl-2-(5-trifluoromethanesulfonyloxy-4'-trifluoromethyl-biphenyl-3-yl)-pentanoic acid ethyl ester

(intermediate **Example 1g**) with 3-dimethylamino-phenylboronic acid under the conditions described in **Example 1**; ¹H NMR (300 MHz, MeOD) δ ppm 0.99 (dd, *J*=6.59, 2.45 Hz, 6 H), 1.58 (dt, *J*=13.28, 6.73 Hz, 1 H), 1.78 (ddd, *J*=13.75, 7.16, 6.97 Hz, 1 H), 2.03 - 2.13 (m, 1 H), 3.32 (s, 6 H), 3.88 (t, *J*=7.72 Hz, 1 H), 7.49 (d, *J*=7.91 Hz, 1 H), 7.68 - 7.83 (m, 7 H), 7.86 - 7.94 (m, 3 H); Calcd for C₂₇H₂₈F₃NO₂ (M+H) 456.21, Found 456.

Example 50


(S)-2-(4-Chloro-3-fluoro-4''-trifluoromethyl-[1,1';3'1'']terphenyl-5'-yl)-4-methyl-pentanoic acid



The title compound was synthesized using a similar procedure as described for the preparation of compound **40** using 4-chloro-3-fluorophenylboronic acid in **Example 40c**. ¹H-NMR (CDCl₃): δ 0.96 (d, 6H), 1.58 (m, 1H), 1.77 (m, 1H), 2.07 (m, 1H), 3.83 (t, 1H), 7.33-7.64 (m, 6H), 7.71 (s, 4H); Calcd for C₂₅H₂₁ClF₄O₂ (M+H) 465.12, Found 465.

Example 51

(R)-4-Methyl-2-(4-trifluoromethoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid




5

The title compound was synthesized using a similar procedure as described for the preparation of compound **39** using 4-(trifluoromethoxy)phenylboronic acid and the intermediate in **Example 39e**.

¹H-NMR (CDCl₃): δ 0.96 (d, 6H), 1.59 (m, 1H), 1.79 (m, 1H), 2.08 (m, 1H), 3.82 (t, 1H), 7.30 (d, 2H), 7.55 (d, 2H), 7.64 (m, 3H), 7.71 (s, 4H); Calcd for C₂₆H₂₂F₆O₃ (M+H) 497.15, Found 497.2.

Example 52

(S)-4-Methyl-2-(4-trifluoromethoxy-4''-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid



The title compound was prepared from a Suzuki coupling of trifluoro-methanesulfonic acid 5-[1-(4-benzyl-2-oxo-oxazolidine-3-carbonyl)-3-methyl-butyl]-4'-trifluoromethyl-biphenyl-3-yl ester (intermediate **Example 40c**) with 4-trifluoromethoxy-phenyl-boronic acid under the conditions described in **Example 40**; ¹H NMR (400 MHz, MeOD) δ ppm 0.87 - 0.98 (m, 6 H), 1.56 (dt, *J*=13.39, 6.63 Hz, 1 H), 1.75 (dt, *J*=13.76, 6.94 Hz, 1 H),

- 5 1.98 - 2.08 (m, 1 H), 3.84 (t, $J=7.83$ Hz, 1 H), 7.36 (t, $J=8.56$ Hz, 2 H), 7.64 (d, $J=5.62$ Hz, 2 H), 7.70 - 7.80 (m, 5 H), 7.84 - 7.89 (m, 2 H).

Determination of the effect of the compounds according to the invention on cyclooxygenase-1 and cyclooxygenase-2 (Cox-1, Cox-2)

10

Inhibition of Cox-1 and Cox-2 was determined using the Colorimetric Cox inhibitor screening assay provided by Cayman Chemical Company, Ann Arbor, MI, USA. (Cat. No. 760111) according to manufacturer's instructions.

- 15 Compounds of the invention will show <50% inhibition at 100micromolar.

Screening of the compounds of the invention for γ -secretase-modulating activity

- 20 Screening was carried out using SKNBE2 cells carrying the APP 695 – wild type, grown in DMEM/NUT-mix F12 (HAM) provided by Gibco (cat no. 31330-38) containing 5% Serum/Fe supplemented with 1% non-essential amino acids.

Cells were grown to near confluency.

The screening was performed using the assay as described in Citron et al (1997) Nature Medicine 3: 67.

25

IC50-values of selected compounds of the invention for γ -secretase activity.

Activity range: 1-10uM

(4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid; (example v)

4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-carboxylic acid; (example xvi)

- 30 5-(4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-1H-tetrazole; (example xvii)

5

Additional IC₅₀-values of selected compounds of the invention for γ -secretase activity

Compound #		WTAPP SKNBE2 A β 42 EC ₅₀ , μ M	WTAPP SKNBE2 A β 42 % Inhibition at 3 μ M
1		0.19	
2		-	53
3		0.44	
4		0.55	
5		0.19	
6		-	65
7		0.08	
8		-	-1
9		0.95	
10		0.99	
11		-	68
12		-	86
13		0.17	
14		0.19	
15		0.14	
16		0.36	
17		0.30	
18		0.17	
19		0.26	
20		0.25	
21		-	46
22		0.15	
23		-	84
24		-	53
25		-	82
26		0.25	

27		0.09	
28		0.28	
29		0.12	
30		0.08	
31		0.16	
32		-	80
33		0.21	
34		-	73
35		0.14	
36		-	0
37		-	84
38		-	53
39		0.14	
40		0.09	
41		0.51	
42		0.33	
43		0.21	
44		0.15	
45		0.22	
46		0.45	
47		0.18	
48		-	18
49		-	-17
50		0.25	
51		0.38	
52		0.24	

5

Demonstration of in vivo efficacy

A β 42 lowering agents of the invention can be used to treat AD in mammals such as humans or alternatively in a validated animal model such as the mouse, rat, or guinea pig.

5 The mammal may not be diagnosed with AD, or may not have a genetic predisposition for AD, but may be transgenic such that it overproduces and eventually deposits A β in a manner similar to that seen in humans afflicted with AD.

A β 42 lowering agents can be administered in any standard form using any standard method. For example, but not limited to, A β 42 lowering agents can be in the form of
10 liquid, tablets or capsules that are taken orally or by injection. A β 42 lowering agents can be administered at any dose that is sufficient to significantly reduce levels of A β 42 in the blood, blood plasma, serum, cerebrospinal fluid (CSF), or brain.

To determine whether acute administration of an A β 42 lowering agent would reduce A β 42 levels in vivo, non-transgenic rodents, e.g. mice or rats can be used. Alternatively,
15 two to three month old Tg2576 mice expressing APP695 containing the "Swedish" variant can be used or a transgenic mouse model developed by Dr. Fred Van Leuven (K.U.Leuven, Belgium) and co-workers, with neuron-specific expression of a clinical mutant of the human amyloid precursor protein [V717I] (Moechars et al., 1999 J. Biol. Chem. 274, 6483). The single transgenic mouse displays spontaneous, progressive
20 accumulation of β -amyloid (A β) in the brain, eventually resulting in amyloid plaques within subiculum, hippocampus and cortex. Animals of this age have high levels of A β in the brain but no detectable A β deposition. Mice treated with the A β 42 lowering agent will be examined and compared to those untreated or treated with vehicle and brain levels of soluble A β 42 and total A β would be quantitated by standard techniques, for example,
25 using ELISA. Treatment periods may vary from hours to days and will be adjusted based on the results of the A β 42 lowering once a time course of onset of effect can be established.

A typical protocol for measuring A β 42 lowering in vivo is shown but it is only one of many variations that could be used to optimize the levels of detectable A β . For example,
30 aliquots of compounds can be dissolved in DMSO (volume equal to 1/10th of the final formulation volume), vortexed and further diluted (1:10) with a 10 % (w/v)

- 5 hydroxypropyl β cyclodextrin (HBC, Aldrich, Ref N° 33,260-7) solution in PBS, where after they are sonicated for 20 seconds.

A β 42 lowering agents may be administered as a single oral dose given three to four hours before sacrifice and analysis or alternatively could be given over a course of days and the animals sacrificed three to four hours after the final dose is given.

- 10 Blood is collected at sacrifice. The blood collection is performed via a heart puncture during anesthesia with a mixture of Ketalar (Ketamin), Rompun (Xylazin 2%) and Atropin (2:1:1) and collected in EDTA treated collection tubes. Blood is centrifuged at 4000 g for 5 minutes at 4 °C and the plasma recovered for analysis.

- The mice are anaesthetized with a mixture of Ketalar (Ketamin), Rompun (Xylazin 2%)
15 and Atropin (2:1:1) and flushed trans-cardially with physiological serum at 4 °C.

The brain is removed from the cranium and hindbrain and forebrain are separated with a cut in the coronal/frontal plane. The cerebellum is removed. The forebrain is divided evenly into left and right hemisphere by using a midline sagital cut.

- One hemisphere is immediately immersed in liquid nitrogen and stored at -70 °C until
20 homogenization for biochemical assays.

- Brains are homogenized using a Potter, a glass tube (detergent free, 2 cm³) and a mechanical homogenizer (650 rpm). A volume of 6,5 x 1/2 brain weight of freshly prepared 20 mM Tris/HCl buffer (pH 8,5) with Proteinase Inhibitors (1 tablet per 50 ml Tris/HCl buffer, CompleteTM, Roche, Mannheim, Germany) is used as homogenization
25 buffer.

Samples are transferred from -70 °C into a sample holder with liquid nitrogen and each individual sample is pre-warmed by incubation on the bench for a few seconds prior to homogenization. The homogenates are collected in Beckman centrifuge tubes TLX and collected on ice prior to centrifugation. Between two samples, the Potter and the glass

5 tube are rinsed carefully with distilled water without detergents and dried with absorption paper.

Samples are centrifuged in a pre-cooled ultracentrifuge (Beckman, Mannheim, Germany) for 1 hour and 20 minutes at 48000 rpm (135.000 x g) at 4°C. The supernatant (soluble fraction containing secreted APP and amyloid peptides) is separated from the pellet
10 (membrane fraction containing membrane-bound APP-fragments and plaque-associated amyloid peptides in case of aged mice).

Small reversed phase columns (C18-Sep-Pack Vac 3cc cartridges, Waters, Massachusetts, MA) are mounted on a vacuum system and washed with 80% acetonitrile in 0,1% Trifluoroacetic acid (A-TFA) followed with 0,1% TFA twice. Then the samples
15 are applied and the columns are washed successively with 5% and 25% A-TFA. Amyloid peptides are eluted with 75% A-TFA and the eluates are collected in 2 ml tubes on ice. Eluates are freeze-dried in a speedvac concentrator (Savant, Farmingdale, NY) overnight and resolved in 240 µl of the sample diluent furnished with the ELISA kits.

To quantify the amount of human Aβ-42 in the soluble fraction of the brain homogenates,
20 commercially available Enzyme-Linked-Immunosorbent-Assay (ELISA) kits are used (h Amyloid β42 ELISA high sensitive, The Genetics Company, Zurich, Switzerland). The ELISA is performed according to the manufacturer's protocol. Briefly, the standard (a dilution of synthetic Aβ1-42) and samples are prepared in a 96-well polypropylene plate without protein binding capacity (Greiner bio-one, Frickenhausen, Germany). The
25 standard dilutions with final concentrations of 1000, 500, 250, 125, 62.5, 31.3 and 15.6 pg/ml and the samples are prepared in the sample diluent, furnished with the ELISA kit, to a final volume of 60 µl. Samples, standards and blancs (50 µl) are added to the anti-Aβ-coated polystyrol plate (capture antibody selectively recognizes the C-terminal end of the antigen) in addition with a selective anti-Aβ-antibody conjugate (biotinylated
30 detection antibody) and incubated overnight at 4 °C in order to allow formation of the antibody-Amyloid-antibody-complex. The following day, a Streptavidine-Peroxidase-Conjugate is added, followed 30 minutes later by an addition of TMB/peroxide mixture, resulting in the conversion of the substrate into a colored product. This reaction is

5 stopped by the addition of sulfuric acid (1M) and the color intensity is measured by means of photometry with an ELISA-reader with a 450 nm filter. Quantification of the Abeta content of the samples is obtained by comparing absorbance to a standard curve made with synthetic A β 1-42.

10 In such a model at least 20% A β 42 lowering compared to untreated animals would be advantageous.

In Vivo Data Oral dose 30 mpk at 4 hr time point


Compound #		Mouse Efficacy % Lowering A β 42	Rat Efficacy % Lowering A β 42
4		30	na
5		48	29
14		58	20
16		10	na
20		50	20
39		50	27
40		40	40
41		58	na
42		48	na
43		40	na
44		40	-10
47		45	na
50		48	28
51		50	na
52		48	na

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.

All publications disclosed in the above specification are hereby incorporated by reference in full.

5 **Claims**

1. A compound having the general formula (I)



10 wherein

X is a bond or a group $-CR_5R_6$ wherein R_5 and R_6 are, independently of each other, selected from the group consisting of H; alkyl selected from the group CH_3 , C_2H_5 , $i-C_3H_7$, $n-C_3H_7$, $i-C_4H_9$, $n-C_4H_9$, $sec-C_4H_9$, $tert-C_4H_9$; alkenyl selected from C_2H_3 , $i-C_3H_5$, $n-C_3H_5$, $n-C_4H_7$, $i-C_4H_7$, $sec-C_4H_7$; wherein in any of the alkyl or alkenyl groups one or

15 more H atoms optionally can be substituted with one or more substituents independently selected from the group consisting of OH, F, Cl, Br, I and CF_3 ; or R_5 and R_6 being part of a ring, either saturated or unsaturated, substituted or unsubstituted, having 3 to 6 C-atoms, and which may contain in the ring one or more heteroatoms from the group N, S or O, and which heteroatom may be identical or different if more than one heteroatom is

20 present;

R_1 , R_2 , R_3 and R_4 are independently selected from the group consisting of H; F; Cl; Br; I; CN; OH; $C(O)N(R_7R_8)$; $S(O)_2R_7$; $SO_2N(R_7R_8)$; $S(O)N(R_7R_8)$; $N(R_7)S(O)_2R_8$; $N(R_8)S(O)R_8$; $S(O)_2R_7$; $N(R_7)S(O)_2N(R_8R_{8a})$; SR_7 ; $N(R_7R_8)$; $N(R_7)C(O)R_8$; $N(R_7)C(O)N(R_8R_{8a})$; $N(R_7)C(O)OR_8$; $OC(O)N(R_7R_8)$; $C(O)R_7$; substituted and

25 unsubstituted C_1 - C_4 -alkyl and substituted and unsubstituted C_1 - C_4 -alkoxy, and wherein the substituents of both groups C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy are selected from OH, F, Cl, Br, I, CF_3 ;

5

R₇, R₈, R_{8a} are independently selected from the group consisting of H; C₁-C₄-alkyl; heterocyclyl; and C₃₋₇ cycloalkyl, wherein C₁-C₄-alkyl; heterocyclyl; and C₃₋₇ cycloalkyl optionally can be substituted with one or more substituents independently selected from the group consisting of OH, F, Cl, Br, I and CF₃;

10 R₉, and R₁₀ are independently H, F, or CF₃;

Y is a carboxy group -C(O)OH or a substituted or unsubstituted tetrazole group and solvates, hydrates, esters, and pharmaceutically acceptable salts thereof..

2. A compound according to claim 1, wherein:

15 X is group -CR₅R₆ wherein R₅ and R₆ are, independently of each other, selected from the group consisting of H; alkyl selected from the group CH₃, C₂H₅, i-C₃H₇, n-C₃H₇, i-C₄H₉, n-C₄H₉, sec-C₄H₉, tert-C₄H₉; wherein in any of the alkyl groups one or more H atoms optionally can be substituted with one or more substituents independently selected from the group consisting of OH, F, Cl, Br and I; or R₅, R₆ jointly form together with the
20 carbon atom to which they are attached a cyclopropyl ring;

R₁, R₂, R₃ and R₄ are independently selected from the group consisting of H, OH, C₍₁₋₄₎alkyl, C₍₁₋₄₎alkoxy, -N(CH₃)₂, -SO₂CH₃, CN, OCF₃, -C(O)CH₃, OCH₃, CF₃, F, and Cl; wherein said C₍₁₋₄₎alkyl and C₍₁₋₄₎alkoxy optionally can be independently substituted with one, two, or three substituents selected from the group consisting of OH, I, Br, F, and Cl;

25 Y is a carboxy group
and solvates, hydrates, esters, and pharmaceutically acceptable salts thereof.

3. A compound according to claim 2, wherein:

X is a group -CR₅R₆ with R₅ and R₆ being H; or R₅ being H and R₆ being CH₃, C₂H₅,
30 C₃H₇ or C₄H₉ or isomers thereof;

5

and solvates, hydrates, esters, and pharmaceutically acceptable salts thereof..

4. A compound according to claim 3, wherein:

X is a group $-CR_5R_6$ with R_5 and R_6 being H; or R_5 being H and R_6 being CH_3 , C_2H_5 ,
10 C_3H_7 or C_4H_9 or isomers thereof;

R_1 , R_2 , R_3 and R_4 are independently selected from the group consisting of H, OH, $C_{(1-4)}$ alkyl, $-N(CH_3)_2$, $-SO_2CH_3$, CN, OCF_3 , $-C(O)CH_3$, OCH_3 , CF_3 , F, and Cl;

and solvates, hydrates, esters, and pharmaceutically acceptable salts thereof.

15 5. A compound selected from the group consisting of

(i) 4''-Chloro-4-trifluoromethyl-[1,1';3,1'']terphenyl-5'-yl)-acetic acid

(ii) (4"-Trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid

(iii) (3-Chloro-4"-trifluoromethyl [1,1';3',1'']terphenyl-5'-yl)-acetic acid

(iv) (4-Hydroxy-4"-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid

20 (v) (4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid

(vi) [1,1';3',1'']Terphenyl-5'-yl)-acetic acid

(vii) (4,4"-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid

(viii) (4,4"-Difluoro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid

(ix) (3,3"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-acetic acid

25 (x) (3,3"-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid

(xi) (4,4"-Dimethyl-[1,1';3',1'']terphenyl-5'-yl)-acetic acid


(xii) (4,4"-Dimethoxy-[1,1';3',1'']terphenyl-5'-yl)-acetic acid

(xiii) 2-(4,4"-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid






- 5 (xiv) (R)-2-(4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
 (xv) (S)-2-(4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-pentanoic acid
 (xvi) 4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-carboxylic acid
 (xvii) 5-(4,4''-Dichloro-[1,1';3',1'']terphenyl-5'-yl)-1H-tetrazole
 (xviii) 2-(4,4''-Bis-trifluoromethyl-[1,1';3',1'']terphenyl-5'-yl)-4-methyl-pentanoic acid






10 and solvates, hydrates, esters, and pharmaceutically acceptable salts thereof.






6. A compound selected from the group consisting of









15

















5




10




5


;








;








;




;









5




;




;



;



;



;

10

- 5 and solvates, hydrates, esters, and pharmaceutically acceptable salts thereof.
7. A compound according to any of claims 1 to 6 for use as a medicament.
8. Use of a compound according to any of claims 1 to 6 for the preparation of a medicament for the modulation of γ -secretase.
- 10 9. Use of a compound according to any of claims 1 to 6 for the preparation of a medicament for the treatment of a disease associated with an elevated level of A β 42-production.
10. Use of a compound according to any of claims 1 to 6 for the preparation of a medicament for the treatment of Alzheimer's disease.
- 15 11. A pharmaceutical composition comprising a compound according to any of claims 1 to 6 in admixture with an inert carrier.
12. A process for the preparation of a compound according to any of claims 1 to 6, comprising the following steps:
- 20 a) treating a dihalidefluorobenzene compound, preferably dibromofluorobenzene, with a benzyl alcohol in the presence of an alkali metal hydride;
- b) treating the product with a suitable malonic ester derivative in the presence of an alkali metal hydride and a metal halide;
- 25 c) treatment in an acidic solvent;
- d) coupling to a boronic acid derivative;
- e) removal of the benzyl ether protecting group;
- f) converting the resulting hydroxycompound to a triflate and coupling to a boronic acid;
- 30

- 5 g) optionally alkylating the resulting triphenyl compound;
- h) conversion of the ester to the acid;
13. A process for the preparation of a compound according to any of claims 1 to 6, comprising the following steps:
- a) converting a dihydroxyphenylacetic acid derivative to a bis-triflate;
- 10 b) coupling the bis-triflate to a boronic acid;
- c) optionally alkylating the resulting triphenyl compound;
- d) converting the ester to the acid;
- e) resolution of racemic mixture into enantiomers.
- 15 14. A process for the preparation of a compound according to any of claims 1 to 6, comprising the following steps:
- a) converting a dihydroxybenzonitrile to a bis-triflate;
- b) coupling the bis-triflate to a boronic acid;
- c) hydrolysis of the nitrile; or
- 20 d) converting the nitrile to tetrazole.
15. Method for the preparation of a medicament comprising the steps of:
- a) preparing a compound according to claim 1; and
- b) formulation of a medicament containing said compound.
- 25
16. A method of treating a mammal for the modulation of γ -secretase, which method comprises administering to said mammal a therapeutically effective amount of a compound according to any of claims 1 to 6.

- 5 **17.** A method of treating a disease in a mammal associated with an elevated level of
 A β 42-production, which method comprises administering to said mammal a
 therapeutically effective amount of a compound according to any of claims 1 to 6.
- 18.** A method of treating Alzheimer's disease in a mammal, which method comprises
10 administering to said mammal a therapeutically effective amount of a compound
 according to any of claims 1 to 6.
- 19.** A compound according to claim 1 as a substantially pure base.
- 20.** A compound according to claim 1 in isolated form.