CLAIMS

1. Use of a composition in the manufacture of a medicament for the treatment and/or prevention of a mitochondrial dysfunction-related disease, wherein the mitochondrial dysfunction-related disease is obesity, the composition comprising an active agent, said active agent containing the amino acids leucine, isoleucine, valine, threonine, lysine and citric acid, succinic acid, malic acid.

2. The use according to claim 1, wherein the weight ratio between the sum of citric acid, malic acid, succinic acid and the sum of the branched chain amino acids leucine, isoleucine, valine plus lysine and threonine is comprised between 0.05 and 0.3, preferably between 0.1 and 0.25.

3. The use according to any one of the preceding claims, wherein the weight ratio between the overall amount of citric acid, malic acid, succinic acid and the overall amount of the branched chain amino acids leucine, isoleucine, valine is comprised between 0.1 and 0.4, preferably between 0.15 and 0.35.

4. The use according to any one of the preceding claims, wherein the weight ratio between citric acid and the sum of malic acid and succinic acid is comprised between 1.0 and 4.0, preferably between 1.5 and 2.5.

5. The use according to any one of the preceding claims, wherein the citric acid:malic acid:succinic acid weight ratio is comprised between 10:1:1 and 2:1.5:1.5, preferably between 7:1:1 and 1.5:1:1, more

preferably between 5:1:1 and 3:1:1.

6. The use according to any one of the preceding claims, wherein said active agent further comprises at least one amino acid selected in the group consisting of histidine, phenylalanine, methionine, tryptophan, tyrosine, cysteine.

7. The use according to any one of the preceding claims, wherein said active agent further comprises histidine, phenylalanine, methionine, tryptophan, cysteine and optionally tyrosine.

8. The use according to any one of the preceding claims, the ratio between the overall molar amount of citric acid, malic acid, succinic acid and the overall molar amount of methionine, phenylalanine, histidine and tryptophan is higher than 1.35.

9. The use according to any one of the preceding claims, wherein the ratio between the overall molar amount of the three acids citric acid, succinic acid, malic acid and the overall molar amount of lysine and threonine is comprised between 0.1 and 0.7, preferably between 0.15 and 0.55.

10. The use according to any one of the preceding claims, wherein the weight or molar amount of citric acid is higher than the overall weight or molar amount of malic acid and succinic acid.

11. The use according to any one of the preceding claims, wherein the weight ratio between leucine and citric acid is comprised between 5 and 1, preferably between 2.50 and 3.50.

12. The use according to any one of the preceding claims, wherein said active agent is free of arginine.

13. The use according to any one of the preceding claims, wherein the composition further comprises one or more vitamins, preferably selected in the group of vitamins B, more preferably vitamin B1 and/or vitamin B6.