

- 1 -

COMBINATION THERAPY TO TREAT BRAIN CANCER

TECHNICAL FIELD

The present invention relates to combination therapies and methods for treating brain cancer.

BACKGROUND

Despite advances in therapy, glioblastoma (GBM) remains one of the deadliest cancers. The current standard treatment for GBM is surgery, followed by concurrent radiation therapy (RT) and temozolomide (TMZ) chemotherapy administered daily during RT and then for 6-12 maintenance (adjuvant) cycles following the completion of RT for select patients [Stupp R, Mason WP, van den Bent MJ, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. N Engl J Med 2005, 352:987-996].

Checkpoint inhibitors, such as programmed cell death-1 (PD-1) inhibitors, have increased response rates in many cancers, but have not yet shown clinical benefit in GBM.

Accordingly, a need exists for the identification and development of methods for the treatment and prognosis of GBM to facilitate clinical management and progression of disease.

SUMMARY

Provided herein are methods of treating brain cancer in a subject, comprising: (a) measuring an expression level of a miR-331-3p miRNA comprising SEQ ID NO: 30 or isomiRs thereof and a miR-1537-3p miRNA comprising SEQ ID NO: 31 or isomiRs thereof in a biological sample from a subject who has brain cancer; and (b) administering interleukin-12 (IL-12); an immunogenic composition of human telomerase reverse transcriptase (hTERT), Wilms Tumor-1 (WT-1), and prostate specific membrane antigen (PSMA); and an anti-programmed cell death receptor 1 (PD-1) antibody to said subject if the subject has an increased expression level of the miR-331-3p miRNA or isomiRs

- 2 -

thereof and the miR-1537-3p miRNA or isomiRs thereof relative to a control population of subjects. In certain embodiments, the presence of the miRNA is determined by RNA sequencing. In certain embodiments, the biological sample is a plasma sample. In certain embodiments, the biological sample is taken from the subject prior to administration of the IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and the anti-PD-1 antibody.

Also provided herein are methods of treating brain cancer in a subject, comprising: (a) measuring an expression level of a miR-331-3p miRNA comprising SEQ ID NO: 30 or isomiRs thereof and a miR-1537-3p miRNA comprising SEQ ID NO: 31 or isomiRs thereof in a biological sample from a subject who has brain cancer; and (b) administering interleukin-12 (IL-12); an immunogenic composition of human telomerase reverse transcriptase (hTERT), Wilms Tumor-1 (WT-1), and prostate specific membrane antigen (PSMA); and an anti-programmed cell death receptor 1 (PD-1) antibody to said subject if the subject has an increased expression level of the miR-331-3p miRNA or isomiRs thereof and the miR-1537-3p miRNA or isomiRs thereof relative to a control population of subjects. In certain embodiments, the biological sample is a primary tumor sample. In certain embodiments, the biological sample is taken from the subject prior to administration of the IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and the anti-PD-1 antibody.

In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in survival at 18 months post administration. In certain embodiments, the control population of subjects is deceased at 18 months post administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody. In certain embodiments, the subject has an unmethylated MGMT gene promoter.

In certain embodiments, IL-12 is encoded by a DNA plasmid, for example, INO-9012 or a biosimilar or bioequivalent thereof. In certain embodiments, hTERT, WT-1, and PSMA are encoded by one or more DNA plasmids, for example, INO-5401 or a biosimilar or bioequivalent thereof. In certain embodiments, the anti-PD-1 antibody is cemiplimab or a biosimilar or bioequivalent thereof. In certain embodiments, the methods further

- 3 -

comprise administering radiation therapy and/or a chemotherapeutic agent, for example, temozolomide or a bioequivalent thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

The summary, as well as the following detailed description, is further understood when read in conjunction with the appended drawings. For the purpose of illustrating the disclosed methods, there are shown in the drawings exemplary embodiments of the thereof; however, the methods are not limited to the specific embodiments disclosed. In the drawings:

FIG. 1 illustrates the study design for the examples.

FIG. 2A and FIG. 2B show the study population demographics for the example for the 18-month (**FIG. 2A**) and 42-month (**FIG. 2B**) analysis timepoints.

FIG. 3 shows representative MRI Images from two patients demonstrating increase in MRI signal at timepoints following first dose of INO-5401 + INO-9012 and cemiplimab-rwlc, suggestive of edema or tumor. Biopsy on several patients shows treatment-related changes with necrosis and mixed inflammation; absence of mitotic activity; and no evidence of viable tumor. The subject represented by the MRI images in the lower panel showed evidence of disease progression at Week 9 but resolution at Week 21. Subjects with similar findings on MRI who were resected showed only immune infiltrate with an absence of viable tumor.

FIG. 4 demonstrates ELISpot results supporting the combination of INO-5401 and cemiplimab-rwlc as immunogenic- with IFN-g magnitudes above baseline to all 3 antigens in 5/11 subjects and to at least one antigen in 9 subjects obtained at the 12-month data cut-off.

FIG. 5A, FIG. 5B, and FIG. 5C show the lytic granule loading assay results demonstrating frequencies of live, antigen-specific, activated (CD38+) CD3+CD8+ T cells with lytic potential (expressing Granzyme A, Perforin) obtained at the 12-month data cut-off. **FIG. 5A** shows the frequencies of live, antigen specific, activated (CD38+) CD3+CD8+ T cells with lytic potential (expressing Granzyme A, Perforin) from before treatment (pre) and the highest magnitude (peak) after treatment with INO-5401 and

- 4 -

cemiplimab-rwlc. Each subject is represented by an open circle, bars represent the mean. The difference from pre to peak, delta, is shown for each antigen graph as well as together for 8 subjects assayed (**FIG. 5B**) and for the 5 subjects with sample available to week 12 (**FIG. 5C**). INO-5401 is the sum of WT1, PSMA and hTERT. Box plots extend from the 25th to 75th percentile, with a horizontal line at the median, and “+” at the mean.

FIG. 6 shows the visual representation of the Kaplan-Meier estimator of the progression-free survival at six months (PFS6) for Cohort A, patients with the O6-methylguanine methyltransferase gene promoter unmethylated in their tumor cells. The curve shows the probability of an event at a certain time interval. The probability of the event is represented numerically on the y-axis, and the time interval on the x-axis. The event shown is progression-free survival. Progression-free survival is the absence of progression of disease at a given time point for a given subject.

FIG. 7 shows the visual representation of the Kaplan-Meier estimator of the progression-free survival at six months (PFS6) for Cohort B, patients with the O6-methylguanine methyltransferase gene promoter methylated in their tumor cells. The curve shows the probability of an event at a certain time interval. The probability of the event is represented numerically on the y-axis, and the time interval on the x-axis. The event shown is progression-free survival. Progression-free survival is the absence of progression of disease at a given time point for a given subject.

FIG. 8 shows the visual representation of the Kaplan-Meier estimator of the progression-free survival at six months (PFS6) for Cohort A and Cohort B, patients with the O6-methylguanine methyltransferase gene promoter unmethylated or methylated in their tumor cells. The curve shows the probability of an event at a certain time interval. The probability of the event is represented numerically on the y-axis, and the time interval on the x-axis. The event shown is progression-free survival. Progression-free survival is the absence of progression of disease at a given time point for a given subject.

FIG. 9 shows the tabular representation of the Kaplan-Meier estimator of the progression-free survival at six months (PFS6) for Cohort A, Cohort B, and both cohorts combined. The total number of subjects per cohort, number of events, estimation of the

event (PFS6), and the 95% confidence interval (CI) in which the numerical estimate of the event (PFS6) exists are all provided.

FIG. 10A shows the visual representation of the Kaplan-Meier estimator of the overall survival probability over twelve months for Cohort A, for patients with the O6-methylguanine methyltransferase gene promoter unmethylated in their tumor cells. The stepwise curve shows the probability of surviving up to and beyond a specific time point. The survival probability is represented numerically on the y-axis, and survival time in days on the x-axis. **FIG. 10B** shows the visual representation of the Kaplan-Meier estimator of the overall survival probability over eighteen months for Cohort A, for patients with the O6-methylguanine methyltransferase gene promoter unmethylated in their tumor cells. The stepwise curve shows the probability of surviving up to and beyond a specific time point. The survival probability is represented numerically on the y-axis, and survival time in days on the x-axis. Median follow-up in Cohort A is 17.8 months. mITT includes any subject who received ≥ 1 dose of study therapy. Shading represents confidence band on point estimate for survival at that timepoint. **FIG. 10C** shows the visual representation of the Kaplan-Meier analysis of the overall survival for MGMT unmethylated (miTT) patients up to 42 months (cohort A).

FIG. 11A shows the visual representation of the Kaplan-Meier estimator of the overall survival probability over twelve months for Cohort B, for patients with the O6-methylguanine methyltransferase gene promoter methylated in their tumor cells. The stepwise curve shows the probability of surviving up to and beyond a specific time point. The survival probability is represented numerically on the y-axis, and survival time in days on the x-axis. **FIG. 11B** shows the visual representation of the Kaplan-Meier estimator of the overall survival probability over eighteen months for Cohort B, for patients with the O6-methylguanine methyltransferase gene promoter methylated in their tumor cells. The stepwise curve shows the probability of surviving up to and beyond a specific time point. The survival probability is represented numerically on the y-axis, and survival time in days on the x-axis. Median follow-up in Cohort B is 15.6 months. Censored; two subjects in Cohort B withdrew consent for follow-up at Week 3. mITT includes any subject who received ≥ 1 dose of study therapy. Shading represents confidence bands on point estimate for survival at that timepoint.

FIG. 11C shows the visual representation of the Kaplan-Meier analysis of the overall survival for MGMT methylated (miTT) patients up to 42 months (cohort B).

FIG. 12 shows the visual representation of the Kaplan-Meier estimator of the overall survival probability over twelve months for Cohorts A + B combined. The stepwise curve shows the probability of surviving up to and beyond a specific time point. The survival probability is represented numerically on the y-axis, and survival time in days on the x-axis.

FIG. 13 shows the efficacy data of the overall survival at 12 months and 18 months for Cohort A, for Cohort B, and combined. The figure shows the total number of subjects who were reported alive at 12 months and at 18 months. The total number of subjects, estimation of the event (OS12 or OS18), and the 95% confidence interval (CI) in which the numerical estimate of the event (OS12 or OS18) exists are all provided. The 95% CI were calculated using the exact Clopper-Pearson method.

FIG. 14 illustrates all Adverse Events as defined by the clinical study protocol $\geq$ NCI CTCAE Grade 3 from the example at the 18-month analysis timepoint.

FIG. 15 illustrates Immune Related Adverse Events as defined by the clinical study protocol from the example at the 18-month analysis timepoint.

FIG. 16A and **FIG. 16B** provide ELISpot results by Cohort at the 18-month data cut-off. In Cohort A, 19/22 (86%) subjects tested to date had an IFN-g magnitude above baseline to one or more of the antigens INO-5401 (**FIG. 16A**). In Cohort B, 16/17 (94%) subjects tested to date had an IFN-g magnitude above baseline to one or more of the antigens in INO-5401 (**FIG. 16B**). Baseline values from the peak timepoint following treatment are plotted. Samples collected Q3 weeks x 4 and then Q12 weeks.

FIG. 17A and **FIG. 17B** provide results of assessment of post-INO-5401 peripheral immune responses by Cohort by flow cytometry (the expansion of antigen specific CD8+ T cells with lytic potential) at the 18-month data cut-off. In Cohort A, 13/19 (68%) subjects tested to date had a frequency of CD38+GrzA+Prf+ CD8+T cells above baseline to one or more of the antigens in INO-5401 (**FIG. 17A**). In Cohort B, 8/10 (80%) subjects tested to date had a frequency of CD38+GrzA+Prf+ CD8+T cells above baseline to one or more of the antigens in INO-5401 (**FIG. 17B**). Baseline values from the peak timepoint following treatment are plotted. Samples were collected Q3 weeks x 4 and then Q12 weeks.

- 7 -

FIG. 18A and **FIG. 18B** show the normalized transcripts for miR-1537-3p (**FIG. 18A**) and miR-331-3p (**FIG. 18B**) for subjects deceased at 18 months post tumor resection versus subject alive at 18 months post tumor resection.

FIG. 19 shows the dual axis miRNA segregation data for miR-1537-3p (x-axis) miR-331-3p (y-axis) for subjects deceased at 18 months post tumor resection versus subject alive at 18 months post tumor resection.

FIG. 20 provides the transcriptome analysis of primary tumors from Cohort A (MGMT Unmethylated) patients grouped according to OS18 assessment (n=31 patients) (15 alive, 16 deceased). Each subject is represented by an open circle. Box plots extend from the 25th to 75th percentile, with a horizontal line at the median, and “+” at the mean.

FIG. 21 shows changes in gene expression levels for tumor antigen targets and markers for immune cell modulators and subsets from six patients in Cohort A (MGMT Unmethylated) in progression (Post-) versus primary (Pre-) tumor tissue is expressed as log2 (fold change); red indicates an increase and green indicates a decrease in gene expression levels.

DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

The disclosed nucleic acid molecules, proteins, vaccines, and methods may be understood more readily by reference to the following detailed description taken in connection with the accompanying figures, which form a part of this disclosure. It is to be understood that the disclosed nucleic acid molecules, proteins, vaccines, and methods are not limited to the specific nucleic acid molecules, proteins, vaccines, and methods described and/or shown herein, and that the terminology used herein is for the purpose of describing particular embodiments by way of example only and is not intended to be limiting of the claimed nucleic acid molecules, proteins, vaccines, and methods.

Unless specifically stated otherwise, any description as to a possible mechanism or mode of action or reason for improvement is meant to be illustrative only, and the disclosed nucleic acid molecules, proteins, vaccines, and methods are not to be constrained by the

- 8 -

correctness or incorrectness of any such suggested mechanism or mode of action or reason for improvement.

For recitation of numeric ranges herein, each intervening number therebetween with the same degree of precision is explicitly contemplated. For example, for the range of 6-9, the numbers 7 and 8 are contemplated in addition to 6 and 9, and for the range 6.0-7.0, the numbers 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, and 7.0 are explicitly contemplated.

It is to be appreciated that certain features of the disclosed nucleic acid molecules, proteins, vaccines, and methods which are, for clarity, described herein in the context of separate embodiments, may also be provided in combination in a single embodiment.

Conversely, various features of the disclosed nucleic acid molecules, proteins, vaccines, and methods that are, for brevity, described in the context of a single embodiment, may also be provided separately or in any subcombination.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art. In case of conflict, the present document, including definitions, will control. Exemplary methods and materials are described below, although methods and materials similar or equivalent to those described herein can be used in practice or testing of the present invention. All publications, patent applications, patents and other references mentioned herein are incorporated by reference in their entirety. The materials, methods, and examples disclosed herein are illustrative only and not intended to be limiting. The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

Throughout this text, the descriptions refer to compositions and methods of using said compositions. Where the disclosure describes or claims a feature or embodiment associated with a composition, such a feature or embodiment is equally applicable to the methods of using said composition. Likewise, where the disclosure describes or claims a feature or embodiment associated with a method of using a composition, such a feature or embodiment is equally applicable to the composition.

Certain Terminology

The terms “comprise(s),” “include(s),” “having,” “has,” “can,” “contain(s),” and variants thereof, as used herein, are intended to be open-ended transitional phrases, terms, or words that do not preclude the possibility of additional acts or structures. The singular forms “a,” “and” and “the” include plural references unless the context clearly dictates otherwise. The present disclosure also contemplates other embodiments “comprising,” “consisting of,” and “consisting essentially of” the embodiments or elements presented herein, whether explicitly set forth or not.

Some of the quantitative expressions given herein are not qualified with the term “about”. It is understood that, whether the term “about” is used explicitly or not, every quantity given is intended to refer to the actual given value, and it is also meant to refer to the approximation to such given value that would reasonably be inferred based on the ordinary skill in the art, including approximations due to the experimental and/or measurement conditions for such value.

“Adjuvant” as used herein means any molecule added to the immunogenic compositions described herein to enhance the immunogenicity of the antigens encoded by the nucleic acid molecules and the encoding nucleic acid sequences described hereinafter.

“Biosimilar” (of an approved reference product/biological drug, *i.e.*, reference listed drug) refers to a biological product that is highly similar to the reference product notwithstanding minor differences in clinically inactive components with no clinically meaningful differences between the biosimilar and the reference product in terms of safety, purity and potency, based upon data derived from (a) analytical studies that demonstrate that the biological product is highly similar to the reference product notwithstanding minor differences in clinically inactive components; (b) animal studies (including the assessment of toxicity); and/or (c) a clinical study or studies (including the assessment of immunogenicity and pharmacokinetics or pharmacodynamics) that are sufficient to demonstrate safety, purity, and potency in one or more appropriate conditions of use for which the reference product is licensed and intended to be used and for which licensure is sought for the biosimilar. The biosimilar may be an interchangeable product that may be substituted for the reference product at the pharmacy without the intervention of the prescribing healthcare professional.

- 10 -

To meet the additional standard of "interchangeability," the biosimilar is to be expected to produce the same clinical result as the reference product in any given patient and, if the biosimilar is administered more than once to an individual, the risk in terms of safety or diminished efficacy of alternating or switching between the use of the biosimilar and the reference product is not greater than the risk of using the reference product without such alternation or switch. The biosimilar utilizes the same mechanisms of action for the proposed conditions of use to the extent the mechanisms are known for the reference product. The condition or conditions of use prescribed, recommended, or suggested in the labeling proposed for the biosimilar have been previously approved for the reference product. The route of administration, the dosage form, and/or the strength of the biosimilar are the same as those of the reference product and the biosimilar is manufactured, processed, packed or held in a facility that meets standards designed to assure that the biosimilar continues to be safe, pure and potent. The biosimilar may include minor modifications in the amino acid sequence when compared to the reference product, such as N- or C-terminal truncations that are not expected to change the biosimilar performance.

The term "antibody," as used herein, includes immunoglobulin molecules comprising four polypeptide chains, two heavy (H) chains and two light (L) chains interconnected by disulfide bonds, as well as multimers thereof (e.g., IgM). In a typical antibody, each heavy chain comprises a heavy chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region comprises three domains, CH1, CH2 and CH3. Each light chain comprises a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region comprises one domain (CL1). The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDRs), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. In different embodiments of the invention, the FRs of the antibody (or antigen-binding portion thereof) may be identical to the human germline

- 11 -

sequences or may be naturally or artificially modified. An amino acid consensus sequence may be defined based on a side-by-side analysis of two or more CDRs.

The term “antibody,” as used herein, also includes antigen-binding fragments of full antibody molecules. The terms “antigen-binding portion” of an antibody, “antigen-binding fragment” of an antibody, and the like, as used herein, include any naturally occurring, enzymatically obtainable, synthetic, or genetically engineered polypeptide or glycoprotein that specifically binds an antigen to form a complex. Antigen-binding fragments of an antibody may be derived, e.g., from full antibody molecules using any suitable standard techniques such as proteolytic digestion or recombinant genetic engineering techniques involving the manipulation and expression of DNA encoding antibody variable and optionally constant domains. Such DNA is known and/or is readily available from, e.g., commercial sources, DNA libraries (including, e.g., phage-antibody libraries), or can be synthesized. The DNA may be sequenced and manipulated chemically or by using molecular biology techniques, for example, to arrange one or more variable and/or constant domains into a suitable configuration, or to introduce codons, create cysteine residues, modify, add or delete amino acids, etc.

Non-limiting examples of antigen-binding fragments include: (i) Fab fragments; (ii) F(ab')₂ fragments; (iii) Fd fragments; (iv) Fv fragments; (v) single-chain Fv (scFv) molecules; (vi) dAb fragments; and (vii) minimal recognition units consisting of the amino acid residues that mimic the hypervariable region of an antibody (e.g., an isolated complementarity determining region (CDR) such as a CDR3 peptide), or a constrained FR3- CDR3-FR4 peptide. Other engineered molecules, such as domain-specific antibodies, single domain antibodies, domain-deleted antibodies, chimeric antibodies, CDR-grafted antibodies, diabodies, triabodies, tetrabodies, minibodies, nanobodies (e.g. monovalent nanobodies, bivalent nanobodies, etc.), small modular immunopharmaceuticals (SMIPs), and shark variable IgNAR domains, are also encompassed within the expression “antigen-binding fragment,” as used herein.

An antigen-binding fragment of an antibody will typically comprise at least one variable domain. The variable domain may be of any size or amino acid composition and will generally comprise at least one CDR which is adjacent to or in frame with one or

- 12 -

more framework sequences. In antigen-binding fragments having a VH domain associated with a VL domain, the VH and VL domains may be situated relative to one another in any suitable arrangement. For example, the variable region may be dimeric and contain VH-VH, VH-VL or VL-VL dimers. Alternatively, the antigen-binding fragment of an antibody may contain a monomeric VH or VL domain.

In certain embodiments, an antigen-binding fragment of an antibody may contain at least one variable domain covalently linked to at least one constant domain. Non-limiting, exemplary configurations of variable and constant domains that may be found within an antigen-binding fragment of an antibody of the present invention include: (i) VH-CH1; (ii) VH-CH2; (iii) VH-CH3; (iv) VH-CH1-CH2; (v) VH-CH1-CH2-CH3; (vi) VH-CH2-CH3; (vii) VH-CL; (viii) VL-CH1; (ix) VL-CH2; (x) VL-CH3; (xi) VL-CH1-CH2; (xii) VL-CH2-CH3; (xiii) VL-CH2-CH3; and (xiv) VL-CL. In any configuration of variable and constant domains, including any of the exemplary configurations listed above, the variable and constant domains may be either directly linked to one another or may be linked by a full or partial hinge or linker region. A hinge region may consist of at least 2 (e.g., 5, 10, 15, 20, 40, 60 or more) amino acids which result in a flexible or semi-flexible linkage between adjacent variable and/or constant domains in a single polypeptide molecule. Moreover, an antigen-binding fragment of an antibody of the present invention may comprise a homo-dimer or hetero-dimer (or other multimer) of any of the variable and constant domain configurations listed above in non-covalent association with one another and/or with one or more monomeric VH or VL domain (e.g., by disulfide bond(s)).

“Coding sequence” or “encoding nucleic acid” as used herein means the nucleic acids (RNA or DNA molecule) that comprise a nucleotide sequence which encodes a protein. The coding sequence can further include initiation and termination signals operably linked to regulatory elements including a promoter and polyadenylation signal capable of directing expression in the cells of an individual or mammal to which the nucleic acid is administered.

“Complement” or “complementary” as used herein means a nucleic acid can mean Watson-Crick (e.g., A-T/U and C-G) or Hoogsteen base pairing between nucleotides or nucleotide analogs of nucleic acid molecules.

- 13 -

“Consensus” or “consensus sequence” as used herein means a polypeptide sequence based on analysis of an alignment of multiple sequences for the same gene from different organisms. Nucleic acid sequences that encode a consensus polypeptide sequence can be prepared. Immunogenic compositions comprising proteins that comprise consensus sequences and/or nucleic acid molecules that encode such proteins can be used to induce broad immunity against an antigen.

“Electroporation,” “electro-permeabilization,” or “electro-kinetic enhancement” (“EP”) as used interchangeably herein means the use of a transmembrane electric field pulse to induce microscopic pathways (pores) in a bio-membrane; their presence allows biomolecules such as plasmids, oligonucleotides, siRNA, drugs, ions, and water to pass from one side of the cellular membrane to the other.

“Fragment” as used herein with respect to nucleic acid sequences means a nucleic acid sequence or a portion thereof, that encodes a polypeptide capable of eliciting an immune response in a mammal that cross reacts with an antigen disclosed herein. The fragments can be DNA fragments selected from at least one of the various nucleotide sequences that encode protein fragments set forth below. Fragments can comprise at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 95% of one or more of the nucleic acid sequences set forth below. In some embodiments, fragments can comprise at least 20 nucleotides or more, at least 30 nucleotides or more, at least 40 nucleotides or more, at least 50 nucleotides or more, at least 60 nucleotides or more, at least 70 nucleotides or more, at least 80 nucleotides or more, at least 90 nucleotides or more, at least 100 nucleotides or more, at least 150 nucleotides or more, at least 200 nucleotides or more, at least 250 nucleotides or more, at least 300 nucleotides or more, at least 350 nucleotides or more, at least 400 nucleotides or more, at least 450 nucleotides or more, at least 500 nucleotides or more, at least 550 nucleotides or more, at least 600 nucleotides or more, at least 650 nucleotides or more, at least 700 nucleotides or more, at least 750 nucleotides or more, at least 800 nucleotides or more, at least 850 nucleotides or more, at least 900 nucleotides or more, at least 950 nucleotides or more, or at least 1000 nucleotides or more of at least one of the nucleic acid sequences set forth below.

- 14 -

“Fragment” or “immunogenic fragment” with respect to polypeptide sequences means a polypeptide capable of eliciting an immune response in a mammal that cross reacts with an antigen disclosed herein. The fragments can be polypeptide fragments selected from at least one of the various amino acid sequences below. Fragments of consensus proteins can comprise at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90% or at least 95% of a consensus protein. In some embodiments, fragments of consensus proteins can comprise at least 20 amino acids or more, at least 30 amino acids or more, at least 40 amino acids or more, at least 50 amino acids or more, at least 60 amino acids or more, at least 70 amino acids or more, at least 80 amino acids or more, at least 90 amino acids or more, at least 100 amino acids or more, at least 110 amino acids or more, at least 120 amino acids or more, at least 130 amino acids or more, at least 140 amino acids or more, at least 150 amino acids or more, at least 160 amino acids or more, at least 170 amino acids or more, at least 180 amino acids or more of a protein sequence disclosed herein.

As used herein, the term “genetic construct” refers to the DNA or RNA molecules that comprise a nucleotide sequence which encodes a protein. The coding sequence includes initiation and termination signals operably linked to regulatory elements including a promoter and polyadenylation signal capable of directing expression in the cells of the individual to whom the nucleic acid molecule is administered. As used herein, the term “expressible form” refers to gene constructs that contain the necessary regulatory elements operably linked to a coding sequence that encodes a protein such that when present in the cell of the individual, the coding sequence will be expressed.

The term “homology,” as used herein, refers to a degree of complementarity. There can be partial homology or complete homology (i.e., identity). A partially complementary sequence that at least partially inhibits a completely complementary sequence from hybridizing to a target nucleic acid is referred to using the functional term “substantially homologous.” When used in reference to a double-stranded nucleic acid sequence such as a cDNA or genomic clone, the term “substantially homologous,” as used herein, refers to a probe that can hybridize to a strand of the double-stranded nucleic acid sequence under conditions of low stringency. When used in reference to a single-stranded nucleic acid

- 15 -

sequence, the term "substantially homologous," as used herein, refers to a probe that can hybridize to (i.e., is the complement of) the single-stranded nucleic acid template sequence under conditions of low stringency.

"Identical" or "identity" as used herein in the context of two or more nucleic acids or polypeptide sequences means that the sequences have a specified percentage of residues that are the same over a specified region. The percentage can be calculated by optimally aligning the two sequences, comparing the two sequences over the specified region, determining the number of positions at which the identical residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the specified region, and multiplying the result by 100 to yield the percentage of sequence identity. In cases where the two sequences are of different lengths or the alignment produces one or more staggered ends and the specified region of comparison includes only a single sequence, the residues of single sequence are included in the denominator but not the numerator of the calculation. When comparing DNA and RNA, thymine (T) and uracil (U) can be considered equivalent. Identity can be performed manually or by using a computer sequence algorithm such as BLAST or BLAST 2.0.

"Substantially complementary" as used herein means that a first sequence is at least 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the complement of a second sequence over a region of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 180, 270, 360, 450, 540, or more nucleotides or amino acids, or that the two sequences hybridize under stringent hybridization conditions.

"Substantially identical" as used herein means that a first and second sequence are at least 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical over a region of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 180, 270, 360, 450, 540 or more nucleotides or amino acids, or

- 16 -

with respect to nucleic acids, if the first sequence is substantially complementary to the complement of the second sequence.

The term “therapeutically effective amount” refers to a therapeutically effective amount of a biologic, compound, or composition that can produce a therapeutic effect in a human subject. A therapeutically effective amount is an amount that can treat, ameliorate, or prevent an identified disease or condition, or to exhibit a detectable therapeutic effect. A therapeutically effective amount is an amount that results in one or more of: (a) a reduction in the severity or duration of a symptom or an indication of a cancer, e.g., glioblastoma; (b) inhibition of tumor growth, or an increase in tumor necrosis, tumor shrinkage and/or tumor disappearance; (c) delay in tumor growth and development; (d) inhibition of tumor metastasis; (e) prevention of recurrence of tumor growth; (f) increase in survival of a subject with a cancer; and/or (g) a reduction in the use or need for conventional anti-cancer therapy (e.g., reduced or eliminated use of chemotherapeutic or cytotoxic agents) as compared to an untreated subject or a subject administered the anti-cancer therapy as monotherapy. The precise effective amount for a subject will depend upon the subject's body weight, size, and health; the nature and extent of the condition; and the therapeutic selected for administration. Therapeutically effective amounts for a given situation can be determined by routine experimentation that is within the skill and judgment of the clinician.

As used herein, “therapeutic effect” is a consequence of a medical treatment of any kind, the results of which are judged to be desirable and beneficial. This is true whether the result was expected, unexpected, or even an unintended consequence of the treatment. A therapeutic effect may also be an objectively identifiable improvement as noted by the clinician or other qualified observer.

“Variant” used herein with respect to a nucleic acid means (i) a portion or fragment of a referenced nucleotide sequence; (ii) the complement of a referenced nucleotide sequence or portion thereof; (iii) a nucleic acid that is substantially identical to a referenced nucleic acid or the complement thereof; or (iv) a nucleic acid that hybridizes under stringent conditions to the referenced nucleic acid, complement thereof, or a sequence substantially identical thereto. A variant may be a nucleic acid sequence that is substantially identical over the full length of the full gene sequence or a fragment thereof.

- 17 -

The nucleic acid sequence may be 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical over the full length of the gene sequence or a fragment thereof.

“Variant” with respect to a polypeptide is one that differs in amino acid sequence by the insertion, deletion, or conservative substitution of amino acids, but retains at least one biological activity of the reference polypeptide. Variant can also mean a protein with an amino acid sequence that is substantially identical to a referenced protein with an amino acid sequence that retains at least one biological activity. A variant may be an amino acid sequence that is substantially identical over the full length of the amino acid sequence or fragment thereof. The amino acid sequence may be 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical over the full length of the amino acid sequence or a fragment thereof.

“Vector” as used herein means a nucleic acid sequence containing an origin of replication. A vector can be a viral vector, bacteriophage, bacterial artificial chromosome, or yeast artificial chromosome. A vector can be a DNA or RNA vector. A vector can be a self-replicating extrachromosomal vector, and in one embodiment, is an expression plasmid. The vector can contain or include one or more heterologous nucleic acid sequences.

“Immune response” as used herein means the activation of a host’s immune system, e.g., that of a mammal, in response to the introduction of antigen. The immune response can be in the form of a cellular or humoral response, or both.

“Nucleic acid” or “oligonucleotide” or “polynucleotide” as used herein means at least two nucleotides covalently linked together. The depiction of a single strand also defines the sequence of the complementary strand. Thus, a nucleic acid also encompasses the complementary strand of a depicted single strand. Many variants of a nucleic acid can be used for the same purpose as a given nucleic acid. Thus, a nucleic acid also encompasses substantially identical nucleic acids and complements thereof. A single strand provides a probe that can hybridize to a target sequence under stringent hybridization conditions. Thus, a nucleic acid also encompasses a probe that hybridizes under stringent hybridization conditions.

- 18 -

Nucleic acids can be single stranded or double-stranded or can contain portions of both double-stranded and single-stranded sequence. The nucleic acid can be DNA, both genomic and cDNA, RNA, or a hybrid, where the nucleic acid can contain combinations of deoxyribo- and ribo-nucleotides, and combinations of bases including uracil, adenine, thymine, cytosine, guanine, inosine, xanthine hypoxanthine, isocytosine and isoguanine. Nucleic acids can be obtained by chemical synthesis methods or by recombinant methods.

“Operably linked” as used herein means that expression of a gene is under the control of a promoter with which it is spatially connected. A promoter can be positioned 5' (upstream) or 3' (downstream) of a gene under its control. The distance between the promoter and a gene can be approximately the same as the distance between that promoter and the gene it controls in the gene from which the promoter is derived. As is known in the art, variation in this distance can be accommodated without loss of promoter function.

A “peptide,” “protein,” or “polypeptide” as used herein can mean a linked sequence of amino acids and can be natural, synthetic, or a modification or combination of natural and synthetic.

“Promoter” as used herein means a synthetic or naturally-derived molecule which is capable of conferring, activating, or enhancing expression of a nucleic acid in a cell. A promoter can comprise one or more specific transcriptional regulatory sequences to further enhance expression and/or to alter the spatial expression and/or temporal expression of same. A promoter can also comprise distal enhancer or repressor elements, which can be located as much as several thousand base pairs from the start site of transcription. A promoter can be derived from sources including viral, bacterial, fungal, plants, insects, and animals. A promoter can regulate the expression of a gene component constitutively, or differentially with respect to cell, the tissue or organ in which expression occurs or, with respect to the developmental stage at which expression occurs, or in response to external stimuli such as physiological stresses, pathogens, metal ions, or inducing agents. Representative examples of promoters include the bacteriophage T7 promoter, bacteriophage T3 promoter, SP6 promoter, lac operator-promoter, tac promoter, SV40 late promoter, SV40 early promoter, RSV-LTR promoter, CMV IE promoter, SV40 early promoter or SV40 late promoter and the CMV IE promoter.

- 19 -

“Signal peptide” and “leader sequence” are used interchangeably herein and refer to an amino acid sequence that can be linked at the amino terminus of a protein set forth herein. Signal peptides/leader sequences typically direct localization of a protein. Signal peptides/leader sequences used herein can facilitate secretion of the protein from the cell in which it is produced. Signal peptides/leader sequences are often cleaved from the remainder of the protein, often referred to as the mature protein, upon secretion from the cell. Signal peptides/leader sequences are linked at the amino terminus (i.e., N terminus) of the protein.

As used herein, the expression "a subject in need thereof" means a human or non-human mammal that exhibits one or more symptoms or indications of brain cancer, and/or who has been diagnosed with brain cancer, including for example glioblastoma, and who needs treatment for the same. In many embodiments, the term "subject" may be interchangeably used with the term "patient". For example, a human subject may be diagnosed with a primary or a metastatic tumor and/or with one or more symptoms or indications including, but not limited to, unexplained weight loss, general weakness, persistent fatigue, loss of appetite, fever, night sweats, bone pain, shortness of breath, swollen abdomen, chest pain/pressure, enlargement of spleen, and elevation in the level of a cancer-related biomarker (e.g., CA125). The expression includes subjects with primary or established tumors. The term includes subjects with primary or metastatic tumors (advanced malignancies). For example, the expression includes subjects who have been newly diagnosed. In some embodiment, the expression includes subjects for whom treatment in accordance with the disclosed methods is an initial treatment (e.g., “first line” treatment, wherein the patient has not received prior systemic treatment for the cancer). In certain embodiments, the expression includes subjects for whom treatment in accordance with the disclosed methods is “second-line” treatment, wherein the patient has been previously treated with “standard-of-care” therapy including, but not limited to chemotherapy, surgery and radiation.

As used herein, the term "treat", "treating", or the like, means to alleviate symptoms, eliminate the causation of symptoms either on a temporary or permanent basis, to delay or inhibit tumor growth, to reduce tumor cell load or tumor burden, to promote tumor

- 20 -

regression, to cause tumor shrinkage, necrosis and/or disappearance, to prevent tumor recurrence, to prevent or inhibit metastasis, to inhibit metastatic tumor growth, and/or to increase duration of survival of the subject.

As used herein, the phrase "in combination with" means that the cancer antigens hTERT, PSMA, and WT-1 are administered to the subject at the same time as, just before, or just after administration of the adjuvant, the PD-1 checkpoint inhibitor, radiation therapy, and/or chemotherapeutic agent. In certain embodiments, the cancer antigens are administered as a co-formulation with the adjuvant.

As used herein, unless otherwise noted, the term "clinically proven" (used independently or to modify the terms "safe" and/or "effective") shall mean that it has been proven by a clinical trial wherein the clinical trial has met the approval standards of U.S. Food and Drug Administration, EMA or a corresponding national regulatory agency. For example, proof may be provided by the clinical trial described in the example provided herein.

The term "clinically proven safe", as it relates to a dose, dosage regimen, treatment or method with cancer antigens hTERT, PSMA, WT1 (for example, administered as INO-5401 or a biosimilar or bioequivalent thereof) in combination with the adjuvant, such as IL-12 (for example, administered as INO-9012 or a biosimilar or bioequivalent thereof) and a PD-1 checkpoint inhibitor, such as an anti-PD-1 antibody (e.g., the anti-PD-1 antibody REGN2810 or a biosimilar or bioequivalent thereof), refers to a favorable risk:benefit ratio with an acceptable frequency and/or acceptable severity of treatment-emergent adverse events (referred to as AEs or TEAEs) compared to the standard of care or to another comparator. An adverse event is an untoward medical occurrence in a patient administered a medicinal product. One index of safety is the National Cancer Institute (NCI) incidence of adverse events (AE) graded per Common Toxicity Criteria for Adverse Events CTCAE v4.03.

The terms "clinically proven efficacy" and "clinically proven effective" as used herein in the context of a dose, dosage regimen, treatment or method refer to the effectiveness of a particular dose, dosage, or treatment regimen. Efficacy can be measured based on change in the course of the disease in response to an agent of the

- 21 -

present invention. For example, a combination of cancer antigens hTERT, PSMA, WT1, and adjuvant, (for example, INO-5401 or a biosimilar or bioequivalent thereof in combination with INO-9012 or a biosimilar or bioequivalent thereof) with a PD-1 checkpoint inhibitor, such as an anti-PD-1 antibody (e.g., the anti-PD-1 antibody cemiplimab or a biosimilar or bioequivalent thereof), is administered to a patient in an amount and for a time sufficient to induce an improvement, preferably a sustained improvement, in at least one indicator that reflects the severity of the disorder that is being treated. Various indicators that reflect the extent of the subject's illness, disease or condition may be assessed for determining whether the amount and time of the treatment is sufficient. Such indicators include, for example, clinically recognized indicators of disease severity, symptoms, or manifestations of the disorder in question. The degree of improvement generally is determined by a physician, who may make this determination based on signs, symptoms, biopsies, or other test results, and who may also employ questionnaires that are administered to the subject, such as quality-of-life questionnaires developed for a given disease. For example, the combination of cancer antigens hTERT, PSMA, WT1, and adjuvant, (for example, INO-5401 or a biosimilar or bioequivalent thereof in combination with INO-9012 or a biosimilar or bioequivalent thereof) with an anti-PD-1 antibody (e.g., the anti-PD-1 antibody cemiplimab or a biosimilar or bioequivalent thereof), may be administered to achieve an improvement in a patient's condition related to brain cancer, such as glioblastoma (GBM). Improvement may be indicated by an improvement in an index of disease activity, by amelioration of clinical symptoms or by any other measure of disease activity.

As used herein, "INO-5401" refers to an immunologic composition of three DNA plasmids: a DNA plasmid comprising an insert encoding hTERT operably controlled by a promoter, a DNA plasmid comprising an insert encoding WT1 operably controlled by a promoter, and a DNA plasmid comprising an insert encoding PSMA operably controlled by a promoter.

As used herein, the term "radiation therapy", also referred to as "XRT," means using ionizing radiation to kill cancer cells, generally as part of anti-cancer therapy. X-rays,

- 22 -

gamma rays or charged particles (e.g., protons or electrons) are used to generate ionizing radiation.

Radiation therapy may be delivered by a machine placed outside the patient's body (external- beam radiation therapy), or by a source placed inside a patient's body (internal radiation therapy or brachytherapy), or through systemic radioisotopes delivered intravenously or orally (systemic radioisotope therapy). Radiation therapy may be planned and administered in conjunction with imaging-based techniques such as a computed tomography (CT), magnetic resonance imaging (MRI) to accurately determine the dose and location of radiation to be administered. In various embodiments, radiation therapy is selected from the group consisting of total all-body radiation therapy, conventional external beam radiation therapy, stereotactic radiosurgery, stereotactic body radiation therapy, 3-D conformal radiation therapy, intensity- modulated radiation therapy, image-guided radiation therapy, tomotherapy, brachytherapy, and systemic radiation therapy. Depending upon the intent, in certain embodiments, radiation therapy is curative, adjuvanting or palliative. In specific embodiments, the term “radiation therapy” refers to hypofractionated radiation therapy. Hypofractionated radiation therapy refers to a radiation treatment schedule in which the total dose of radiation is divided into large doses and treatments are given once a day or less often. Hypofractionated radiotherapy may provide more radiation per dose in fewer doses than standard radiotherapy. In various embodiments, each fraction comprises 2-20 Gy. For example, a radiation dose of 50 Gy may be split up into 10 fractions, each comprising 5 Gy. In certain embodiments, the 2 or more fractions are administered on consecutive or sequential days. In certain other embodiments, the 2 or more fractions are administered once in 2 days, once in 3 days, once in 4 days, once in 5 days, once in 6 days, once in 7 days, or in a combination thereof.

As used herein, the term “isomiR” refers to isoforms of canonical (mature) miRNAs. IsomiRs may have the following variations from the canonical miRNA: (1) addition and/or deletion of nucleotide(s) at the 5'-end of the canonical miRNA; (2) addition and/or deletion of nucleotide(s) at the 3'-end of canonical miRNA; (3) addition and/or deletion of nucleotide(s) at both 5'- and 3'-ends of the canonical miRNA; and (4) nucleotide substitution within the sequence of the canonical miRNA.

- 23 -

As used herein, the term “sensitivity” refers to the proportion of treated subjects with brain cancer who do not meet the defined TMM threshold.

As used herein, the term “specificity” refers to the proportion of treated subjects with brain cancer who meet the defined TMM threshold.

As used herein, the term “negative predictive value” refers to the ability to accurately select for subjects who would respond clinically to a disclosed prior to treatment with the disclosed therapy.

As used herein, the term “positive predictive value” refers to the ability to accurately predict for lack of response to treatment with a disclosed therapy.

As used herein, the term “defined TMM threshold” or “defined threshold” refers to a TMM value that causes both negative and positive predictive values as well as sensitivity and specificity to exceed 70% according to the Classification and Regression Tree (CART) analysis described in the Examples.

Methods of Treatment, Uses and Diagnosis

Provided herein are vaccines and methods of their use to prevent or treat cancer. Also provided herein are methods of selecting a subject for brain cancer therapy. The cancer can be brain cancer, for example, glioblastoma. The vaccine preferably includes at least three cancer antigens, hTERT, WT-1, and PSMA. In certain embodiments, the vaccine also includes an adjuvant, such as IL-12, and an anti-PD-1 antibody. The methods involve administering cancer antigens hTERT, WT-1, and PSMA, an adjuvant, and a PD-1 checkpoint inhibitor, such as an anti-PD-1 antibody to a subject in need thereof. In some embodiments, the methods prevent tumor growth. In some embodiments, the methods can reduce tumor growth and/or mass. In some embodiments, the methods can prevent metastasis of tumor cells. In some embodiments, the methods can increase a cellular immune response in the subject. In some embodiments, the methods increase tumor-free survival, progression-free survival, overall survival, or any combination thereof, of the subject.

In particular embodiments, the disclosed methods and uses can mediate clearance or prevent growth of tumor cells by inducing (1) humoral immunity via B cell responses to

- 24 -

generate antibodies that block monocyte chemoattractant protein-1 (MCP-1) production, thereby retarding myeloid derived suppressor cells (MDSCs) and suppressing tumor growth; (2) increase cytotoxic T lymphocyte such as CD8⁺ (CTL) to attack and kill tumor cells; (3) increase T helper cell responses; (4) and increase inflammatory responses via IFN- γ and TFN- α ; or (5) any combination of the aforementioned. The methods can increase progression-free survival by 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, and 45%. The methods can reduce tumor mass by 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, and 60% after immunization. The methods can prevent and block increases in monocyte chemoattractant protein 1 (MCP- 1), a cytokine secreted by myeloid derived suppressor cells. The methods can increase tumor survival by 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, and 60%.

The disclosed methods and uses can increase a cellular immune response in a subject by about 50-fold to about 6000-fold, about 50-fold to about 5500-fold, about 50-fold to about 5000- fold, about 50-fold to about 4500-fold, about 100-fold to about 6000-fold, about 150-fold to about 6000-fold, about 200-fold to about 6000-fold, about 250-fold to about 6000-fold, or about 300-fold to about 6000-fold as compared to a cellular immune response in a subject not administered the method or administered a standard-of-care treatment method. In some embodiments the methods can increase the cellular immune response in the subject by about 50-fold, 100-fold, 150-fold, 200-fold, 250-fold, 300-fold, 350-fold, 400-fold, 450-fold, 500- fold, 550-fold, 600-fold, 650-fold, 700-fold, 750-fold, 800-fold, 850-fold, 900-fold, 950-fold, 1000-fold, 1100-fold, 1200-fold, 1300-fold, 1400-fold, 1500-fold, 1600-fold, 1700-fold, 1800-fold, 1900-fold, 2000-fold, 2100-fold, 2200-fold, 2300-fold, 2400-fold, 2500-fold, 2600-fold, 2700-fold, 2800-fold, 2900-fold, 3000-fold, 3100-fold, 3200-fold, 3300-fold, 3400-fold, 3500-fold, 3600-fold, 3700-fold, 3800-fold, 3900-fold, 4000-fold, 4100-fold, 4200-fold, 4300-fold, 4400-fold, 4500-fold, 4600-fold, 4700-fold, 4800-fold, 4900-fold, 5000-fold, 5100-fold, 5200-fold, 5300-fold, 5400-fold, 5500-fold, 5600-fold, 5700-fold, 5800-fold, 5900-fold, or 6000-fold as compared to

- 25 -

the cellular immune response in the subject not administered the method or administered a standard-of-care treatment method.

In some embodiments, the disclosed methods and uses can increase tumor-free survival, reduce tumor mass, increase progression-free survival, increase overall survival, or a combination thereof in the subject. The methods can increase tumor-free survival by 20%, 21%, 22%, 23%, 24%, 25%, 26%, 27%, 28%, 29%, 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, and 60% in the subject. The methods can reduce tumor mass by 20%, 21%, 22%, 23%, 24%, 25%, 26%, 27%, 28%, 29%, 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, and 70% in the subject. The methods can increase progression-free survival by 20%, 21%, 22%, 23%, 24%, 25%, 26%, 27%, 28%, 29%, 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, and 60% in the subject. The methods can increase overall survival by 20%, 21%, 22%, 23%, 24%, 25%, 26%, 27%, 28%, 29%, 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, and 60% in the subject.

In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in survival or improved overall survival at 1 month, 2 months, 3 months, 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, 12 months, 13 months, 14 months, 15 months, 16 months, 17 months, 18 months, 19 months, 20 months, 21 months, 22 months, 23 months, 24 months, 25 months, 26 months, 27 months, 28 months, 29 months, 30 months, 31 months, 32 months, 33 months, 34 months, 35 months, 36 months, 37 months, 38 months, 39 months, 40 months, 41 months, or 42 months post administration.

In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in survival at 18 months post administration. In further embodiments, administration of IL-12; an immunogenic

- 26 -

composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in improved overall survival at 18 months post administration. In further embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in improved overall survival at 18 months post administration relative to a patient population with brain cancer that has been administered a placebo. In certain embodiments, said administration or of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in improved overall survival at 18 months post administration relative to a patient population with brain cancer that has not been administered an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody. In certain embodiments, said administration or of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in improved overall survival at 18 months post administration relative to a patient population with brain cancer that is deceased at 18 months post administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody.

In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in survival at 12 months post administration. In further embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in improved overall survival at 12 months post administration. In further embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in improved overall survival at 12 months post administration relative to a patient population with brain cancer that has been administered a placebo. In certain embodiments, said administration or of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in improved overall survival at 12 months post administration relative to a patient population with brain cancer that has not been administered an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody. In certain embodiments, said administration or of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in improved overall survival at 12 months post administration relative to a patient population with brain cancer that is deceased at 12 months post administration of IL-

- 27 -

12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody.

In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in improved median overall survival. In further embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in improved median overall survival relative to a patient population with brain cancer that has been administered a placebo. In certain embodiments, said administration of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in improved median overall survival relative to a patient population with brain cancer that has not been administered an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody. In certain embodiments, said administration of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in improved median overall survival relative to a patient population with brain cancer that is deceased post administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody. In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in median overall survival of about 19.5 months. In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in median overall survival in subjects having an unmethylated MGMT promoter of about 17.9 months. In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in median overall survival in subjects having a methylated MGMT promoter of about 32.5 months.

The recombinant cancer antigens can induce antigen-specific T cell and/or high titer antibody responses, thereby inducing or eliciting an immune response that is directed to or reactive against the cancer or tumor expressing the antigen. In some embodiments, the induced or elicited immune response can be a cellular, humoral, or both cellular and humoral immune responses. In some embodiments, the induced or elicited cellular immune response can include induction or secretion of interferon-gamma (IFN- γ) and/or tumor necrosis factor alpha (TNF- α). In other embodiments, the induced or elicited immune

- 28 -

response can reduce or inhibit one or more immune suppression factors that promote growth of the tumor or cancer expressing the antigen, for example, but not limited to, factors that down regulate MHC presentation, factors that up regulate antigen-specific regulatory T cells (Tregs), PD-L1, FasL, cytokines such as IL-10 and TFG- β , tumor associated macrophages, tumor associated fibroblasts, soluble factors produced by immune suppressor cells, CTLA-4, PD-1, MDSCs, MCP-1, and an immune checkpoint molecule.

In certain embodiments, IL-12 is encoded by a DNA plasmid, for example, INO-9012 or a biosimilar or bioequivalent thereof. In certain embodiments, hTERT, WT-1, and PSMA are encoded by one or more DNA plasmids, for example, INO-5401 or a biosimilar or bioequivalent thereof. In further embodiments, the hTERT is encoded by a DNA plasmid, WT-1 is encoded by a DNA plasmid, and/or PSMA is encoded by a DNA plasmid. In still further embodiments, the hTERT, WT-1, and PSMA are encoded by the same DNA plasmid; two of hTERT, WT-1, and PSMA are encoded by the same DNA plasmid; or wherein hTERT, WT-1, and PSMA are each encoded by a different DNA plasmid.

In certain embodiments, the anti-PD-1 antibody is cemiplimab or a biosimilar or bioequivalent thereof. In certain embodiments, the methods further comprise administering radiation therapy and/or a chemotherapeutic agent, for example, temozolomide or a bioequivalent thereof. In certain embodiments, the anti-PD-1 antibody: comprises the heavy chain complementarity determining regions (HCDR1, HCDR2 and HCDR3) of a heavy chain variable region (HCVR) comprising the amino acid sequence of SEQ ID NO: 1 and three light chain complementarity determining regions (LCDR1, LCDR2 and LCDR3) of a light chain variable region (LCVR) comprising the amino acid sequence of SEQ ID NO: 2; comprises three HCDRs (HCDR1, HCDR2 and HCDR3) and three LCDRs (LCDR1, LCDR2 and LCDR3), wherein HCDR1 comprises the amino acid sequence of SEQ ID NO: 3, HCDR2 comprises the amino acid sequence of SEQ ID NO: 4, HCDR3 comprises the amino acid sequence of SEQ ID NO: 5, LCDR1 comprises the amino acid sequence of SEQ ID NO: 6, LCDR2 comprises the amino acid sequence of SEQ ID NO: 7, and LCDR3 comprises the amino acid sequence of SEQ ID NO: 8;

- 29 -

comprises a HCVR with 90% sequence identity to SEQ ID NO: 1; comprises a LCVR with 90% sequence identity to SEQ ID NO: 2; comprises a HCVR with 90% sequence identity to SEQ ID NO: 1 and a LCVR with 90% sequence identity to SEQ ID NO: 2; comprises a HCVR comprising the amino acid sequence of SEQ ID NO: 1 and a LCVR comprising the amino acid sequence of SEQ ID NO: 2; comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 9 and a light chain comprising the amino acid sequence of SEQ ID NO: 10; is an IgG4 antibody; or is REGN2810 or a biosimilar or bioequivalent thereof.

In certain embodiments, the IL12 p35 subunit comprises the amino acid sequence of SEQ ID NO: 23; the IL12 p40 subunit comprises the amino acid sequence of SEQ ID NO: 25; the IL12 p35 subunit comprises the amino acid sequence of SEQ ID NO: 23 and the IL12 p40 subunit comprises the amino acid sequence of SEQ ID NO: 25; the IL12 p35 subunit is encoded by the nucleic acid sequence of SEQ ID NO: 22; the IL12 p40 subunit is encoded by the nucleic acid sequence of SEQ ID NO: 24; or the IL12 p35 subunit is encoded by the nucleic acid sequence of SEQ ID NO: 22 and the IL12 p40 subunit is encoded by the nucleic acid sequence of SEQ ID NO: 24.

In certain embodiments, the hTERT comprises the amino acid sequence of SEQ ID NO: 20 or is encoded by the nucleic acid sequence of SEQ ID NO: 19; the WT-1 comprises the amino acid sequence of SEQ ID NO: 26 or is encoded by the nucleic acid sequence of SEQ ID NO: 27; and/or the PSMA comprises the amino acid sequence of SEQ ID NO: 28 or is encoded by the nucleic acid sequence of SEQ ID NO: 29.

The cancer can be brain cancer, for example, glioblastoma. In certain embodiments, the brain cancer is MGMT-methylated glioblastoma. In certain embodiments, the brain cancer is MGMT-unmethylated glioblastoma.

- 30 -

MicroRNA (miRNA) and IsomiRs

MicroRNAs (miRNAs) are a class of endogenous small noncoding RNA, typically ranging from 19 to 24 nucleotides. When loaded into the microribonucleoprotein (miRNP) or RNA induced silencing complex (RISC), miRNAs bind to miRNA recognition elements (MREs) on mRNA targets, exerting gene regulation through post-transcriptional and/translational mechanisms. To date, there is no known miRNA or group of miRNAs that have been used in a fashion that allows for the prediction of a response to immunotherapy for a given disease state.

Thought to be the products of altered miRNA biogenesis, isomiRs may have additions or deletions at the 5'-, 3'- or both ends of the canonical miRNA. In some rare cases, isomiRs may have some nucleotides substituted within the canonical sequence (polymorphic isomiRs). These isomiRs, originally mistaken to be sequencing artifacts, have recently been confirmed by multiple, parallel sequencing and bioinformatic studies to be bona fide miRNA species, greatly expanding the miRNA repertoire.

Recent studies have increasingly shown that these isomiRs may be conserved across species and produced in a regulated cell-specific manner or at a particular developmental stage. IsomiRs can also be loaded into miRNPs/RISCs and the variations in length and sequences would potentially alter target selection, miRNA loading and stability, possibly with important and different biological functions.

With the advancement of molecular biological techniques, miRNAs can be detected using multiple platforms. Most current miRNA studies are reported using quantitative reverse transcription polymerase chain reaction (qRT-PCR), microarrays and high-throughput sequencing. The use of next generation sequencing has led to the discovery of miRNA isoforms (isomiRs), which are variants of the canonical mature miRNA sequences.

Provided herein are methods of treating brain cancer in a subject, comprising, consisting of, or consisting essentially of: (a) measuring an expression level of a miR-331-3p miRNA comprising SEQ ID NO: 30 or isomiRs thereof and a miR-1537-3p miRNA comprising SEQ ID NO: 31 or isomiRs thereof in a biological sample from a subject who has brain cancer; and (b) administering IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody to said subject if the subject has

- 31 -

an increased expression level of the miR-331-3p miRNA or isomiRs thereof and the miR-1537-3p miRNA or isomiRs thereof relative to a control population of subjects.

Also provided herein are methods for selecting a subject for brain cancer therapy comprising, consisting of, or consisting essentially of: (a) evaluating one or more biological samples from a subject who has brain cancer for the presence of a miR-331-3p miRNA comprising SEQ ID NO: 30 or isomiRs thereof and a miR-1537-3p miRNA comprising SEQ ID NO: 31 or isomiRs thereof; and (b) characterizing the subject as a candidate for brain cancer therapy with IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody if the subject has an increased expression level of the miR-331-3p miRNA or isomiRs thereof and the miR-1537-3p miRNA or isomiRs thereof relative to a control population of subjects.

Further provided herein are vaccines for use in treating brain cancer in a subject who has brain cancer and who harbors an miR-331-3p miRNA comprising SEQ ID NO: 30 or isomiRs thereof and a miR-1537-3p miRNA comprising SEQ ID NO: 31 or isomiRs thereof, wherein the expression level of the miR-331-3p miRNA and the miR-1537-3p miRNA is increased relative to a control population of subjects, and wherein the vaccine comprises an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody.

Still further provided herein are uses of a vaccine in the manufacture of a medicament for treating brain cancer in a subject who has brain cancer and who harbors an miR-331-3p miRNA comprising SEQ ID NO: 30 or isomiRs thereof and a miR-1537-3p miRNA comprising SEQ ID NO: 31 or isomiRs thereof, wherein the expression level of the miR-331-3p miRNA and the miR-1537-3p miRNA is increased relative to a control population of subjects, and wherein the vaccine comprises an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody.

In certain embodiments, the biological sample is taken from the subject at baseline, e.g. prior to administration of the therapeutic composition. In certain embodiments, the increase relative to a control population of subjects is measured at baseline for both the subject and the control population of subjects, e.g. prior to administration of the therapeutic composition.

- 32 -

In certain embodiments, the biological sample is taken from the subject prior to administration of the IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and the anti-PD-1 antibody. In certain embodiments, the biological sample is taken from the control population of subjects prior to administration of the IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and the anti-PD-1 antibody. In certain embodiments, the biological sample is taken from the subject and from the control population of subjects prior to administration of the IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and the anti-PD-1 antibody.

In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in survival or improved overall survival at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 or 18 months post administration. In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in survival or improved overall survival at 18 months post administration. In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in survival or improved overall survival at 12 months post administration.

In further embodiments, the control population of subjects is deceased at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 or 18 months months post administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody. In certain embodiments, the control population of subjects is deceased at 18 months post administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody. In certain embodiments, the control population of subjects is deceased at 12 months post administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody.

In some aspects of the present disclosure, the biological sample is a peripheral blood sample. In certain embodiments, the biological sample is a plasma sample.

In some aspects of the present disclosure, the methods comprise evaluating one or more biological sample. In certain embodiments, the methods comprising evaluating one biological sample. In certain embodiments, the methods comprising evaluating two biological

- 33 -

samples. In certain embodiments, the methods comprising evaluating three biological sample. In certain embodiments, the methods comprising evaluating four biological samples. In certain embodiments, the methods comprising evaluating five biological samples.

In certain embodiments, the subject has an unmethylated MGMT gene promoter. In further embodiments, the subject has a methylated MGMT gene promoter.

Messenger RNA (mRNA)

Provided herein are methods of treating brain cancer in a subject, comprising, consisting of, or consisting essentially of: (a) measuring an expression level of at least one mRNA biomarker selected from SYNGR3, OTX1, GABBR2, LHX1, CADM3, MLLT11, MNX1, GRB14, SLC34A2, PHYHIP, WNT10B, SLC17A6, CRLF1, HOXD13, TGFβR3, UBA7, SFRP4, or any combination thereof, in a tumor sample from a subject who has brain cancer; and (b) administering IL-12; an immunogenic composition hTERT, WT-1, and PSMA; and an anti-PD-1 antibody to said subject if the expression level of SYNGR3, OTX1, GABBR2, LHX1, CADM3, MLLT11, MNX1, GRB14, SLC34A2, PHYHIP, WNT10B, SLC17A6, CRLF1 and HOXD13 is decreased or if the expression level of TGFβR3, UBA7, SFRP4 is increased relative to a control population of subjects.

Also provided herein are methods for selecting a subject for brain cancer therapy comprising, consisting of, or consisting essentially of: (a) measuring an expression level of at least one mRNA biomarker selected from SYNGR3, OTX1, GABBR2, LHX1, CADM3, MLLT11, MNX1, GRB14, SLC34A2, PHYHIP, WNT10B, SLC17A6, CRLF1, HOXD13, TGFβR3, UBA7, SFRP4, or any combination thereof, in a tumor sample from a subject who has brain cancer; and (b) characterizing the subject as a candidate for brain cancer therapy with IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody if the expression level of SYNGR3, OTX1, GABBR2, LHX1, CADM3, MLLT11, MNX1, GRB14, SLC34A2, PHYHIP, WNT10B, SLC17A6, CRLF1 and HOXD13 is decreased or if the expression level of TGFβR3, UBA7, SFRP4 is increased relative to a control population of subjects.

Further provided herein are vaccines for use in treating brain cancer in a subject who has brain cancer, who has an unmethylated MGMT gene promoter, and who harbors at least

- 34 -

one mRNA biomarker selected from SYNGR3, OTX1, GABBR2, LHX1, CADM3, MLLT11, MNX1, GRB14, SLC34A2, PHYHIP, WNT10B, SLC17A6, CRLF1, HOXD13, TGF β R3, UBA7, SFRP4, or any combination thereof, wherein the expression level of SYNGR3, OTX1, GABBR2, LHX1, CADM3, MLLT11, MNX1, GRB14, SLC34A2, PHYHIP, WNT10B, SLC17A6, CRLF1 and HOXD13 is decreased or the expression level of TGF β R3, UBA7, SFRP4 is increased relative to a control population of subjects.

Still further provided herein are uses of a vaccine in the manufacture of a medicament for treating brain cancer in a subject who has brain cancer, who has an unmethylated MGMT gene promoter, and who harbors at least one mRNA biomarker selected from SYNGR3, OTX1, GABBR2, LHX1, CADM3, MLLT11, MNX1, GRB14, SLC34A2, PHYHIP, WNT10B, SLC17A6, CRLF1, HOXD13, TGF β R3, UBA7, SFRP4, or any combination thereof, wherein the expression level of SYNGR3, OTX1, GABBR2, LHX1, CADM3, MLLT11, MNX1, GRB14, SLC34A2, PHYHIP, WNT10B, SLC17A6, CRLF1 and HOXD13 is decreased or the expression level of TGF β R3, UBA7, SFRP4 is increased relative to a control population of subjects.

In certain embodiments, the biological sample is taken from the subject at baseline, e.g. prior to administration of the therapeutic composition. In certain embodiments, the increase relative to a control population of subjects is measured at baseline for both the subject and the control population of subjects, e.g. prior to administration of the therapeutic composition.

In certain embodiments, the biological sample is taken from the subject prior to administration of the IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and the anti-PD-1 antibody. In certain embodiments, the biological sample is taken from the control population of subjects prior to administration of the IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and the anti-PD-1 antibody. In certain embodiments, the biological sample is taken from the subject and from the control population of subjects prior to administration of the IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and the anti-PD-1 antibody.

In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in survival or improved

- 35 -

overall survival at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 or 18 months post administration. In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in survival or improved overall survival at 18 months post administration. In certain embodiments, administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in survival or improved overall survival at 12 months post administration.

In further embodiments, the control population of subjects is deceased at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 or 18 months post administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody. In certain embodiments, the control population of subjects is deceased at 18 months post administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody. In certain embodiments, the control population of subjects is deceased at 12 months post administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody.

In certain embodiments, the biological sample is a primary tumor sample.

In certain embodiments, the subject has an unmethylated MGMT gene promoter. In further embodiments, the subject has a methylated MGMT gene promoter.

Therapeutic Compositions

Disclosed herein are optimized consensus sequences of cancer antigens hTERT, WT-1, and PSMA. In one embodiment, the antigen encoded by the optimized consensus sequence is capable of eliciting an immune response in a mammal. In one embodiment, the antigen encoded by the optimized consensus sequence can comprise an epitope(s) that makes it particularly effective as an immunogen against which an immune response can be induced.

In one embodiment is provided an optimized consensus PSMA designed to break tolerance to native human PSMA. In one embodiment, a human optimized consensus PSMA encoding sequence is as set forth in SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 21, or SEQ ID NO: 28. In one embodiment, a human optimized consensus PSMA encoded

- 36 -

antigen has an amino acid sequence as set forth in SEQ ID NO:13, SEQ ID NO:14, or SEQ ID NO: 28.

In one embodiment, an optimized consensus WT-1 is designed to break tolerance to native human WT-1. In one embodiment, a human optimized consensus WT-1 encoding sequence is as set forth in SEQ ID NO:15 or SEQ ID NO: 27. In one embodiment, a human optimized consensus WT-1 encoded antigen has an amino acid sequence as set forth in SEQ ID NO:16 or SEQ ID NO: 26.

In one embodiment, an optimized consensus TERT is designed to break tolerance to native human TERT. In one embodiment, a human optimized consensus TERT encoding sequence is as set forth in SEQ ID NO:17 or SEQ ID NO:19. In one embodiment, a human optimized consensus TERT encoded antigen has an amino acid sequence as set forth in SEQ ID NO:18 and SEQ ID NO:20.

The disclosed vaccines may further comprise an adjuvant. In certain embodiments, the disclosed methods of treatment further comprise administering to the subject an adjuvant. In certain embodiments, the adjuvant is IL12. IL12 may be included in a vaccine in the form of its p35 and p40 subunits. The adjuvant IL12 may be administered to the subject as its p35 and p40 subunits. The IL12 p35 and p40 subunits may be encoded by the same expression vector or by separate expression vectors. In one embodiment, the IL12 p35 encoding sequence is as set forth in SEQ ID NO:22. In one embodiment, the IL12 p35 subunit has an amino acid sequence as set forth in SEQ ID NO:23. In one embodiment, the IL12 p40 encoding sequence is as set forth in SEQ ID NO:24. In one embodiment, the IL12 p40 subunit has an amino acid sequence as set forth in SEQ ID NO:25.

The cancer antigens TERT, WT-1, PSMA, and/or adjuvant can be present in the vaccine or administered to the subject as the polypeptide, fragment thereof, variant thereof, nucleic acid sequence encoding the polypeptide, fragment or variant thereof, or any combination thereof. The cancer antigen can be any form that induces an immune response in a subject. The nucleic acid sequence can be DNA, RNA, cDNA, a variant thereof, a fragment thereof, or a combination thereof. The nucleic acid sequence can also include additional sequences that encode linker or tag sequences that are linked to the

- 37 -

antigen by a peptide bond. The amino acid sequence can be a protein, a peptide, a variant thereof, a fragment thereof, or a combination thereof.

The cancer antigens TERT, WT-1, PSMA, and/or adjuvant can be included in a vaccine or administered to the subject as the polypeptide, fragment thereof, variant thereof, nucleic acid sequence encoding the polypeptide, fragment or variant thereof, or any combination thereof. The cancer antigen can be any form that induces an immune response in a subject. The nucleic acid sequence can be DNA, RNA, cDNA, a variant thereof, a fragment thereof, or a combination thereof. The nucleic acid sequence can also include additional sequences that encode linker or tag sequences that are linked to the antigen by a peptide bond. The amino acid sequence can be a protein, a peptide, a variant thereof, a fragment thereof, or a combination thereof.

The cancer antigens TERT, WT-1, PSMA and/or IL-12 can be included in a vaccine or administered as one or more nucleic acid molecules, for example but not limited to, an expression vector(s). An expression vector can be a circular plasmid or a linear nucleic acid. An expression vector is capable of directing expression of a particular nucleotide sequence in an appropriate subject cell. An expression vector can have a promoter operably linked to the antigen-encoding nucleotide sequence, which may be operably linked to termination signals. An expression vector can also contain sequences required for proper translation of the nucleotide sequence. The expression vector comprising the nucleotide sequence of interest may be chimeric, meaning that at least one of its components is heterologous with respect to at least one of its other components. The expression of the nucleotide sequence in the expression cassette may be under the control of a constitutive promoter or of an inducible promoter, which initiates transcription only when the host cell is exposed to some particular external stimulus. In the case of a multicellular organism, the promoter can also be specific to a particular tissue or organ or stage of development.

In one embodiment, the nucleic acid is an RNA molecule. Accordingly, in one embodiment, the invention provides an RNA molecule encoding one or more polypeptides of interest. The RNA may be plus-stranded. Accordingly, in some embodiments, the RNA molecule can be translated by cells without needing any intervening replication steps such as reverse transcription. An RNA molecule useful with the invention may have a 5'

- 38 -

cap (e.g. a 7- methylguanosine). This cap can enhance in vivo translation of the RNA. The 5' nucleotide of an RNA molecule useful with the invention may have a 5' triphosphate group. In a capped RNA this may be linked to a 7-methylguanosine via a 5'-to-5' bridge. An RNA molecule may have a 3' poly-A tail. It may also include a poly-A polymerase recognition sequence (e.g., AAUAAA) near its 3' end. An RNA molecule useful with the invention may be single- stranded. In some embodiments, the RNA molecule is a naked RNA molecule. In one embodiment, the RNA molecule is comprised within a vector.

In one embodiment, the RNA has 5' and 3' UTRs. In one embodiment, the 5' UTR is between zero and 3000 nucleotides in length. The length of 5' and 3' UTR sequences to be added to the coding region can be altered by different methods, including, but not limited to, designing primers for PCR that anneal to different regions of the UTRs. Using this approach, one of ordinary skill in the art can modify the 5' and 3' UTR lengths required to achieve optimal translation efficiency following transfection of the transcribed RNA.

The 5' and 3' UTRs can be the naturally occurring, endogenous 5' and 3' UTRs for the gene of interest. Alternatively, UTR sequences that are not endogenous to the gene of interest can be added by incorporating the UTR sequences into the forward and reverse primers or by any other modifications of the template. The use of UTR sequences that are not endogenous to the gene of interest can be useful for modifying the stability and/or translation efficiency of the RNA. For example, it is known that AU-rich elements in 3' UTR sequences can decrease the stability of RNA. Therefore, 3' UTRs can be selected or designed to increase the stability of the transcribed RNA based on properties of UTRs that are well known in the art.

In one embodiment, the 5' UTR can contain the Kozak sequence of the endogenous gene. Alternatively, when a 5' UTR that is not endogenous to the gene of interest is being added by PCR as described above, a consensus Kozak sequence can be redesigned by adding the 5' UTR sequence. Kozak sequences can increase the efficiency of translation of some RNA transcripts, but does not appear to be required for all RNAs to enable efficient translation. The requirement for Kozak sequences for many RNAs is known in the art. In other embodiments, the 5' UTR can be derived from an RNA virus

- 39 -

whose RNA genome is stable in cells. In other embodiments, various nucleotide analogues can be used in the 3' or 5' UTR to impede exonuclease degradation of the RNA.

In one embodiment, the RNA has both a cap on the 5' end and a 3' poly(A) tail which determine ribosome binding, initiation of translation and stability of RNA in the cell.

In one embodiment, the RNA is a nucleoside-modified RNA. Nucleoside-modified RNA have particular advantages over non-modified RNA, including for example, increased stability, low or absent innate immunogenicity, and enhanced translation.

The expression vector may be a circular plasmid, which may transform a target cell by integration into the cellular genome or exist extrachromosomally (e.g., autonomous replicating plasmid with an origin of replication). The vector can be pVAX, pcDNA3.0, or provax, or any other expression vector capable of expressing DNA encoding the antigen and enabling a cell to translate the sequence to an antigen that is recognized by the immune system.

Also provided herein is a linear nucleic acid immunogenic composition, or linear expression cassette ("LEC"), that is capable of being efficiently delivered to a subject via electroporation and expressing one or more desired antigens. The LEC may be any linear DNA devoid of any phosphate backbone. The DNA may encode one or more antigens. The LEC may contain a promoter, an intron, a stop codon, and/or a polyadenylation signal. The expression of the antigen may be controlled by the promoter. The LEC may not contain any antibiotic resistance genes and/or a phosphate backbone. The LEC may not contain other nucleotide sequences unrelated to the desired antigen gene expression. The LEC may be derived from any plasmid capable of being linearized. The plasmid may be capable of expressing the antigen. The plasmid can be pNP (Puerto Rico/34) or pM2 (New Caledonia/99). The plasmid may be WLV009, pVAX, pcDNA3.0, or provax, or any other expression vector capable of expressing DNA encoding the antigen and enabling a cell to translate the sequence to an antigen that is recognized by the immune system. The LEC can be pcrM2. The LEC can be pcrNP. pcrNP and pcrMR can be derived from pNP (Puerto Rico/34) and pM2 (New Caledonia/99), respectively.

- 40 -

The vector can comprise heterologous nucleic acid encoding the above described antigens and can further comprise an initiation codon, which can be upstream of the one or more cancer antigen coding sequence(s), and a stop codon, which can be downstream of the coding sequence(s) of the above described antigens.

The vector may have a promoter. A promoter may be any promoter that is capable of driving gene expression and regulating expression of the isolated nucleic acid. Such a promoter is a cis-acting sequence element required for transcription via a DNA dependent RNA polymerase, which transcribes the antigen sequence described herein. Selection of the promoter used to direct expression of a heterologous nucleic acid depends on the particular application. The promoter may be positioned about the same distance from the transcription start in the vector as it is from the transcription start site in its natural setting. However, variation in this distance may be accommodated without loss of promoter function.

The initiation and termination codon can be in frame with the coding sequence(s) of the above described antigens. The vector can also comprise a promoter that is operably linked to the coding sequence(s) of the above described antigens. The promoter operably linked to the coding sequence(s) of the above described antigens can be a promoter from simian virus 40 (SV40), a mouse mammary tumor virus (MMTV) promoter, a human immunodeficiency virus (HIV) promoter such as the bovine immunodeficiency virus (BIV) long terminal repeat (LTR) promoter, a Moloney virus promoter, an avian leukosis virus (ALV) promoter, a cytomegalovirus (CMV) promoter such as the CMV immediate early promoter, Epstein Barr virus (EBV) promoter, or a Rous sarcoma virus (RSV) promoter. The promoter can also be a promoter from a human gene such as human actin, human myosin, human hemoglobin, human muscle creatine, or human metallothionein. The promoter can also be a tissue specific promoter, such as a muscle or skin specific promoter, natural or synthetic. Examples of such promoters are described in US patent application publication no. US20040175727, the contents of which are incorporated herein in its entirety.

The vector can also comprise a polyadenylation signal, which can be downstream of the coding sequence(s) of the above described antigens and/or antibodies. The polyadenylation signal can be a SV40 polyadenylation signal, LTR polyadenylation

- 41 -

signal, bovine growth hormone (bGH) polyadenylation signal, human growth hormone (hGH) polyadenylation signal, or human β -globin polyadenylation signal. The SV40 polyadenylation signal can be a polyadenylation signal from a pCEP4 vector (Invitrogen, San Diego, CA).

The vector can also comprise an enhancer upstream of the above described antigens.

The enhancer can be necessary for expression. The enhancer can be human actin, human myosin, human hemoglobin, human muscle creatine or a viral enhancer such as one from CMV, HA, RSV or EBV.

The vector may include an enhancer and an intron with functional splice donor and acceptor sites. The vector may contain a transcription termination region downstream of the structural gene to provide for efficient termination. The termination region may be obtained from the same gene as the promoter sequence or may be obtained from different genes.

The disclosed methods may comprise administration of a plurality of copies of a single nucleic acid molecule such as a single plasmid, or a plurality of copies of two or more different nucleic acid molecules such as two or more different plasmids. For example, the methods may comprise administration of two, three, four, five, six, seven, eight, nine or ten or more different nucleic acid molecules.

The nucleic acid molecules used in accordance with the disclosed methods, such as plasmids, may collectively contain coding sequence for a single antigen or for multiple antigens. As an example, in one embodiment, the antigens are multiple antigens selected from TERT and one or more additional cancer antigens. In one exemplary embodiment, the antigens are TERT and WT-1. In one exemplary embodiment, the antigens are TERT and PSMA. In one exemplary embodiment, the antigens are PSMA and one or more additional cancer antigens. In one exemplary embodiment, the antigens are PSMA and WT-1. In another exemplary embodiment, the antigens are TERT, WT-1 and PSMA.

The vector can further comprise elements or reagents that inhibit it from integrating into the chromosome. The vector can comprise a mammalian origin of replication in order to maintain the vector extrachromosomally and produce multiple copies of the vector in a cell. The vector can be pVAX1, pCEP4 or pREP4 from

- 42 -

Invitrogen (San Diego, CA), which can comprise the Epstein Barr virus origin of replication and nuclear antigen EBNA-1 coding region, which can produce high copy episomal replication without integration. The vector can be pVAX1 or a pVax1 variant with changes such as the variant plasmid described herein. The variant pVax1 plasmid is a 2998 base pair variant of the backbone vector plasmid pVAX1 (Invitrogen, Carlsbad CA). The CMV promoter is located at bases 137-724. The T7 promoter/priming site is at bases 664-683. Multiple cloning sites are at bases 696-811.

Bovine GH polyadenylation signal is at bases 829-1053. The Kanamycin resistance gene is at bases 1226-2020. The pUC origin is at bases 2320-2993.

Based upon the sequence of pVAX1 available from Invitrogen, the following mutations were found in the sequence of pVAX1 that was used as the backbone for plasmids 1-6 set forth herein:

C>G241 in CMV promoter

C>T 1942 backbone, downstream of the bovine growth hormone polyadenylation signal (bGHpolyA)

A> - 2876 backbone, downstream of the Kanamycin gene

C>T 3277 in pUC origin of replication (Ori) high copy number mutation (see Nucleic Acid Research 1985)

G>C 3753 in very end of pUC Ori upstream of RNaseH site

Base pairs 2, 3 and 4 are changed from ACT to CTG in backbone, upstream of CMV promoter.

The backbone of the vector can be pAV0242. The vector can be a replication defective adenovirus type 5 (Ad5) vector.

The vector can also comprise a regulatory sequence, which can be well suited for gene expression in a mammalian or human cell into which the vector is administered. The one or more cancer antigen sequences disclosed herein can comprise a codon, which can allow more efficient transcription of the coding sequence in the host cell.

The vector can be pSE420 (Invitrogen, San Diego, Calif.), which can be used for protein production in *Escherichia coli* (*E. coli*). The vector can also be pYES2 (Invitrogen, San Diego, Calif.), which can be used for protein production in *Saccharomyces cerevisiae*

- 43 -

strains of yeast. The vector can also be of the MAXBAC™ complete baculovirus expression system (Invitrogen, San Diego, Calif.), which can be used for protein production in insect cells. The vector can also be pcDNA I or pcDNA3 (Invitrogen, San Diego, Calif.), which may be used for protein production in mammalian cells such as Chinese hamster ovary (CHO) cells. The vector can be expression vectors or systems to produce protein by routine techniques and readily available starting materials including Sambrook et al., Molecular Cloning and Laboratory Manual, Second Ed., Cold Spring Harbor (1989), incorporated fully herein by reference.

Exemplary DNA plasmids encoding the cancer antigens hTERT, WT-1, and/or PSMA are disclosed in U.S. Application No. 62/899,543, filed September 12, 2019, the entire contents of which are disclosed herein by reference.

Dosing and Routes of Administration

In accordance with the disclosed methods and uses, the subject may be administered about 5 nanograms to about 20 mg of a nucleic acid molecule(s) encoding an antigen or antigens. In some embodiments, the subject may be administered about 5 mg to about 15 mg of a nucleic acid molecule(s) encoding an antigen or antigens. In some embodiments, the subject may be administered about 9 mg to about 11 mg of a nucleic acid molecule(s) encoding an antigen or antigens. In some embodiments, the subject may be administered about 10 mg of a nucleic acid molecule(s) encoding an antigen or antigens.

In certain embodiments, the disclosed methods and uses comprise administering to the subject 3 mg of the DNA plasmid encoding hTERT, 3 mg of the DNA plasmid encoding PSMA, 3 mg of the DNA plasmid encoding WT-1, and 1 mg of the plasmid encoding IL-12.

In certain embodiments, the IL-12 and the immunogenic composition are co-administered by intramuscular injection every three weeks for four doses and then every nine weeks. In further embodiments, the disclosed methods and uses further comprise electroporation following each intramuscular injection.

The DNA plasmid(s) can be delivered via a variety of routes. Typical delivery routes include parenteral administration, e.g., intradermal, intramuscular, or subcutaneous

- 44 -

delivery. Other routes include oral administration, intranasal, and intravaginal routes. For the DNA of the vaccine in particular, the vaccine can be delivered to the interstitial spaces of tissues of an individual (Felgner et al., U.S. Pat. Nos. 5,580,859 and 5,703,055, the contents of all of which are incorporated herein by reference in their entirety). The DNA plasmid(s) can also be administered to muscle, or can be administered via intradermal or subcutaneous injections, or transdermally, such as by iontophoresis. Epidermal administration of the DNA plasmid(s) can also be employed. Epidermal administration can involve mechanically or chemically irritating the outermost layer of epidermis to stimulate an immune response to the irritant (Carson et al., U.S. Pat. No. 5,679,647, the contents of which are incorporated herein by reference in its entirety).

The DNA plasmid(s) can be a liquid preparation such as a suspension, syrup, or elixir. The vaccine can also be a preparation for parenteral, subcutaneous, intradermal, intramuscular, or intravenous administration (e.g., injectable administration), such as a sterile suspension or emulsion.

The DNA plasmid(s) can be incorporated into liposomes, microspheres, or other polymer matrices (Felgner et al., U.S. Pat. No. 5,703,055; Gregoriadis, *Liposome Technology*, Vols. I to III (2nd ed. 1993), the contents of which are incorporated herein by reference in their entirety). Liposomes can consist of phospholipids or other lipids, and can be nontoxic, physiologically acceptable and metabolizable carriers that are relatively simple to make and administer.

The DNA plasmid(s) can be administered via electroporation, such as by a method described in U.S. Pat. No. 7,664,545, the contents of which are incorporated herein by reference. The electroporation can be by a method and/or apparatus described in U.S. Pat. Nos. 6,302,874; 5,676,646; 6,241,701; 6,233,482; 6,216,034; 6,208,893; 6,192,270; 6,181,964; 6,150,148; 6,120,493; 6,096,020; 6,068,650; and 5,702,359, the contents of which are incorporated herein by reference in their entirety. The electroporation may be carried out via a minimally invasive device.

The minimally invasive electroporation device ("MID") may be an apparatus for injecting the vaccine described above and associated fluid into body tissue. The device may comprise a hollow needle, DNA cassette, and fluid delivery means, wherein the

- 45 -

device is adapted to actuate the fluid delivery means in use so as to concurrently (for example, automatically) inject DNA into body tissue during insertion of the needle into the said body tissue. This has the advantage that the ability to inject the DNA and associated fluid gradually while the needle is being inserted leads to a more even distribution of the fluid through the body tissue. The pain experienced during injection may be reduced due to the distribution of the DNA being injected over a larger area.

The MID may inject the DNA plasmid(s) into tissue without the use of a needle. The MID may inject the vaccine as a small stream or jet with such force that the vaccine pierces the surface of the tissue and enters the underlying tissue and/or muscle. The force behind the small stream or jet may be provided by expansion of a compressed gas, such as carbon dioxide through a micro-orifice within a fraction of a second. Examples of minimally invasive electroporation devices, and methods of using them, are described in published U.S. Patent Application No. 20080234655; U.S. Pat. Nos. 6,520,950; 7,171,264; 6,208,893; 6,009,347; 6,120,493; 7,245,963; 7,328,064; and 6,763,264, the contents of each of which are herein incorporated by reference.

The MID may comprise an injector that creates a high-speed jet of liquid that painlessly pierces the tissue. Such needle-free injectors are commercially available. Examples of needle-free injectors that can be utilized herein include those described in U.S. Pat. Nos. 3,805,783; 4,447,223; 5,505,697; and 4,342,310, the contents of each of which are herein incorporated by reference.

A desired vaccine in a form suitable for direct or indirect electrotransport may be introduced (e.g., injected) using a needle-free injector into the tissue to be treated, usually by contacting the tissue surface with the injector so as to actuate delivery of a jet of the agent, with sufficient force to cause penetration of the vaccine into the tissue. For example, if the tissue to be treated is mucosa, skin or muscle, the agent is projected towards the mucosal or skin surface with sufficient force to cause the agent to penetrate through the stratum corneum and into dermal layers, or into underlying tissue and muscle, respectively.

Needle-free injectors are well suited to deliver DNA plasmid(s) to all types of tissues, particularly to skin and mucosa. In some embodiments, a needle-free injector

- 46 -

may be used to propel a liquid that contains the DNA plasmid(s) to the surface and into the subject's skin or mucosa. Representative examples of the various types of tissues that can be treated using the invention methods include pancreas, larynx, nasopharynx, hypopharynx, oropharynx, lip, throat, lung, heart, kidney, muscle, breast, colon, prostate, thymus, testis, skin, mucosal tissue, ovary, blood vessels, or any combination thereof.

The MID may have needle electrodes that electroporate the tissue. By pulsing between multiple pairs of electrodes in a multiple electrode array, for example set up in rectangular or square patterns, provides improved results over that of pulsing between a pair of electrodes. Disclosed, for example, in U.S. Pat. No. 5,702,359 entitled "Needle Electrodes for Mediated Delivery of Drugs and Genes" is an array of needles wherein a plurality of pairs of needles may be pulsed during the therapeutic treatment. In that application, which is incorporated herein by reference as though fully set forth, needles were disposed in a circular array, but have connectors and switching apparatus enabling a pulsing between opposing pairs of needle electrodes. A pair of needle electrodes for delivering recombinant expression vectors to cells may be used. Such a device and system are described in U.S. Pat. No. 6,763,264, the contents of which are herein incorporated by reference. Alternatively, a single needle device may be used that allows injection of the DNA and electroporation with a single needle resembling a normal injection needle and applies pulses of lower voltage than those delivered by presently used devices, thus reducing the electrical sensation experienced by the patient.

The MID may comprise one or more electrode arrays. The arrays may comprise two or more needles of the same diameter or different diameters. The needles may be evenly or unevenly spaced apart. The needles may be between 0.005 inches and 0.03 inches, between 0.01 inches and 0.025 inches; or between 0.015 inches and 0.020 inches. The needle may be 0.0175 inches in diameter. The needles may be 0.5 mm, 1.0 mm, 1.5 mm, 2.0 mm, 2.5 mm, 3.0 mm, 3.5 mm, 4.0 mm, or more spaced apart.

The MID may consist of a pulse generator and a two or more-needle injectors that deliver the DNA plasmid(s) and electroporation pulses in a single step. The pulse generator may allow for flexible programming of pulse and injection parameters via a flash card operated personal computer, as well as comprehensive recording and storage of

- 47 -

electroporation and patient data. The pulse generator may deliver a variety of volt pulses during short periods of time. For example, the pulse generator may deliver three 15-volt pulses of 100 ms in duration. An example of such a MID is the Elgen 1000 system by Inovio Biomedical Corporation, which is described in U.S. Pat. No. 7,328,064, the contents of which are herein incorporated by reference.

The MID may be a CELLECTRA® (Inovio Pharmaceuticals, Blue Bell Pa.) device and system, which is a modular electrode system, that facilitates the introduction of a macromolecule, such as a DNA, into cells of a selected tissue in a body or plant. The modular electrode system may comprise a plurality of needle electrodes; a hypodermic needle; an electrical connector that provides a conductive link from a programmable constant-current pulse controller to the plurality of needle electrodes; and a power source. An operator can grasp the plurality of needle electrodes that are mounted on a support structure and firmly insert them into the selected tissue in a body or plant. The macromolecules are then delivered via the hypodermic needle into the selected tissue. The programmable constant-current pulse controller is activated, and constant-current electrical pulse is applied to the plurality of needle electrodes. The applied constant-current electrical pulse facilitates the introduction of the macromolecule into the cell between the plurality of electrodes. Cell death due to overheating of cells is minimized by limiting the power dissipation in the tissue by virtue of constant-current pulses. The CELLECTRA® device and system is described in U.S. Pat. No. 7,245,963, the contents of which are herein incorporated by reference.

The MID may be an Elgen 1000 system (Inovio Pharmaceuticals). The Elgen 1000 system may comprise device that provides a hollow needle; and fluid delivery means, wherein the apparatus is adapted to actuate the fluid delivery means in use so as to concurrently (for example automatically) inject fluid, the described DNA plasmid(s) herein, into body tissue during insertion of the needle into the said body tissue. The advantage is the ability to inject the fluid gradually while the needle is being inserted leads to a more even distribution of the fluid through the body tissue. It is also believed that the pain experienced during injection is reduced due to the distribution of the volume of fluid being injected over a larger area.

- 48 -

In addition, the automatic injection of fluid facilitates automatic monitoring and registration of an actual dose of fluid injected. This data can be stored by a control unit for documentation purposes if desired.

It will be appreciated that the rate of injection could be either linear or non-linear and that the injection may be carried out after the needles have been inserted through the skin of the subject to be treated and while they are inserted further into the body tissue.

Suitable tissues into which fluid may be injected by the apparatus of the present invention include tumor tissue, skin or liver tissue but may be muscle tissue.

The apparatus further comprises needle insertion means for guiding insertion of the needle into the body tissue. The rate of fluid injection is controlled by the rate of needle insertion. This has the advantage that both the needle insertion and injection of fluid can be controlled such that the rate of insertion can be matched to the rate of injection as desired. It also makes the apparatus easier for a user to operate. If desired means for automatically inserting the needle into body tissue could be provided.

A user could choose when to commence injection of fluid. Ideally however, injection is commenced when the tip of the needle has reached muscle tissue and the apparatus may include means for sensing when the needle has been inserted to a sufficient depth for injection of the fluid to commence. This means that injection of fluid can be prompted to commence automatically when the needle has reached a desired depth (which will normally be the depth at which muscle tissue begins). The depth at which muscle tissue begins could for example be taken to be a preset needle insertion depth such as a value of 4 mm which would be deemed sufficient for the needle to get through the skin layer.

The sensing means may comprise an ultrasound probe. The sensing means may comprise a means for sensing a change in impedance or resistance. In this case, the means may not as such record the depth of the needle in the body tissue but will rather be adapted to sense a change in impedance or resistance as the needle moves from a different type of body tissue into muscle. Either of these alternatives provides a relatively accurate and simple to operate means of sensing that injection may commence. The depth of insertion of the needle can further be recorded if desired and could be used to control

- 49 -

injection of fluid such that the volume of fluid to be injected is determined as the depth of needle insertion is being recorded.

The apparatus may further comprise: a base for supporting the needle; and a housing for receiving the base therein, wherein the base is moveable relative to the housing such that the needle is retracted within the housing when the base is in a first rearward position relative to the housing and the needle extends out of the housing when the base is in a second forward position within the housing. This is advantageous for a user as the housing can be lined up on the skin of a patient, and the needles can then be inserted into the patient's skin by moving the housing relative to the base.

As stated above, it is desirable to achieve a controlled rate of fluid injection such that the fluid is evenly distributed over the length of the needle as it is inserted into the skin. The fluid delivery means may comprise piston driving means adapted to inject fluid at a controlled rate. The piston driving means could for example be activated by a servo motor. However, the piston driving means may be actuated by the base being moved in the axial direction relative to the housing. It will be appreciated that alternative means for fluid delivery could be provided. Thus, for example, a closed container which can be squeezed for fluid delivery at a controlled or non-controlled rate could be provided in the place of a syringe and piston system.

The apparatus described above could be used for any type of injection. It is however envisaged to be particularly useful in the field of electroporation and so it may further comprise means for applying a voltage to the needle. This allows the needle to be used not only for injection but also as an electrode during, electroporation. This is particularly advantageous as it means that the electric field is applied to the same area as the injected fluid. There has traditionally been a problem with electroporation in that it is very difficult to accurately align an electrode with previously injected fluid and so users have tended to inject a larger volume of fluid than is required over a larger area and to apply an electric field over a higher area to attempt to guarantee an overlap between the injected substance and the electric field. Using the present invention, both the volume of fluid injected, and the size of electric field applied may be reduced while achieving a good fit between the electric field and the fluid.

- 50 -

Upon administration of nucleic acid molecule(s) encoding cancer antigens hTERT, PSMA, and WT-1 to the subject, the transfected cells will express and secrete one or more of the cancer antigens. These secreted proteins, or synthetic antigens, will be recognized as foreign by the immune system, which will mount an immune response that can include antibodies made against the one or more cancer antigens, and T-cell response specifically against the one or more cancer antigens. In some examples, a mammal administered the immunogenic composition discussed herein will have a primed immune system and when challenged with the one or more cancer antigens as disclosed herein, the primed immune system will allow for rapid clearing of subsequent cancer antigens as disclosed herein, whether through the humoral, cellular, or both cellular and humoral immune responses.

The recombinant cancer antigen can induce antigen-specific T cell and/or high titer antibody responses, thereby inducing or eliciting an immune response that is directed to or reactive against the cancer or tumor expressing the antigen. In some embodiments, the induced or elicited immune response can be a cellular, humoral, or both cellular and humoral immune responses. In some embodiments, the induced or elicited cellular immune response can include induction or secretion of interferon-gamma (IFN- γ) and/or tumor necrosis factor alpha (TNF- α). In other embodiments, the induced or elicited immune response can reduce or inhibit one or more immune suppression factors that promote growth of the tumor or cancer expressing the antigen, for example, but not limited to, factors that down regulate MHC presentation, factors that up regulate antigen-specific regulatory T cells (Tregs), PD-L1, FasL, cytokines such as IL-10 and TGF- β , tumor associated macrophages, tumor associated fibroblasts, soluble factors produced by immune suppressor cells, CTLA-4, PD-1, MDSCs, MCP-1, and an immune checkpoint molecule.

Combination with anti-PD-1 antibody

The disclosed vaccines may further comprise an anti-PD-1 antibody. The disclosed methods of treatment may further comprise administering to the subject an anti-PD-1 antibody. According to certain embodiments of the present invention, the anti-PD-1

- 51 -

antibody comprises a heavy chain variable region (HCVR), light chain variable region (LCVR), and/or complementarity determining regions (CDRs) comprising the amino acid sequences of any of the anti-PD-1 antibodies as set forth in US Patent Publication No. 20150203579, hereby incorporated in its entirety. In certain exemplary embodiments, the anti-PD-1 antibody that can be used in the context of the disclosed methods comprises the heavy chain complementarity determining regions (HCDRs) of a heavy chain variable region (HCVR) comprising the amino acid sequence of SEQ ID NO: 1 and the light chain complementarity determining regions (LCDRs) of a light chain variable region (LCVR) comprising the amino acid sequence of SEQ ID NO: 2. According to certain embodiments, the anti-PD-1 antibody comprises three HCDRs (HCDR1, HCDR2 and HCDR3) and three LCDRs (LCDR1, LCDR2 and LCDR3), wherein the HCDR1 comprises the amino acid sequence of SEQ ID NO: 3; the HCDR2 comprises the amino acid sequence of SEQ ID NO: 4; the HCDR3 comprises the amino acid sequence of SEQ ID NO: 5; the LCDR1 comprises the amino acid sequence of SEQ ID NO: 6; the LCDR2 comprises the amino acid sequence of SEQ ID NO: 7; and the LCDR3 comprises the amino acid sequence of SEQ ID NO: 8. In yet other embodiments, the anti-PD-1 antibody comprises an HCVR comprising SEQ ID NO: 1 and an LCVR comprising SEQ ID NO: 2. In certain embodiments, the methods of the present invention comprise the use of an anti-PD-1 antibody, wherein the antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 9. In some embodiments, the anti-PD-1 antibody comprises a light chain comprising the amino acid sequence of SEQ ID NO: 10. An exemplary antibody comprising a heavy chain comprising the amino acid sequence of SEQ ID NO: 9 and a light chain comprising the amino acid sequence of SEQ ID NO: 10 is the fully human anti-PD-1 antibody known as REGN2810 and also known as cemiplimab or cemiplimab-rwlc.

According to certain exemplary embodiments, the methods of the present invention comprise the use of REGN2810, or a biosimilar or bioequivalent thereof. The term “bioequivalent”, as used herein, refers to anti-PD-1 antibodies or PD-1-binding proteins or fragments thereof that are pharmaceutical equivalents or pharmaceutical alternatives whose rate and/or extent of absorption do not show a significant difference with that of

- 52 -

REGN2810 when administered at the same molar dose under similar experimental conditions, either single dose or multiple dose. In the context of the invention, the term refers to antigen-binding proteins that bind to PD-1 which do not have clinically meaningful differences with REGN2810 in their safety, purity and/or potency.

According to certain embodiments of the present invention, the anti-human PD-1 antibody comprises a HCVR having 90%, 95%, 98% or 99% sequence identity to SEQ ID NO: 1.

According to certain embodiments of the present invention, the anti-human PD-1 antibody comprises a LCVR having 90%, 95%, 98% or 99% sequence identity to SEQ ID NO: 2.

According to certain embodiments of the present invention, the anti-human PD-1 antibody comprises a HCVR comprising an amino acid sequence of SEQ ID NO: 1 having no more than 5 amino acid substitutions. According to certain embodiments of the present invention, the anti-human PD-1 antibody comprises a LCVR comprising an amino acid sequence of SEQ ID NO: 2 having no more than 2 amino acid substitutions.

Sequence identity may be measured by any method known in the art (e.g., GAP, BESTFIT, and BLAST).

The present invention also includes use of anti-PD-1 antibodies in methods to treat cancer, wherein the anti-PD-1 antibodies comprise variants of any of the HCVR, LCVR and/or CDR amino acid sequences disclosed herein having one or more conservative amino acid substitutions. For example, the present invention includes use of anti-PD-1 antibodies having HCVR, LCVR and/or CDR amino acid sequences with, e.g., 10 or fewer, 8 or fewer, 6 or fewer, 4 or fewer, etc. conservative amino acid substitutions relative to any of the HCVR, LCVR and/or CDR amino acid sequences disclosed herein.

The amount of anti-PD-1 antibody administered to a subject according to the disclosed methods can be a therapeutically effective amount. As used herein, the phrase "therapeutically effective amount" of anti-PD-1 antibody is an amount that results in one or more of: (a) a reduction in the severity or duration of a symptom or an indication of a cancer, e.g., glioblastoma; (b) inhibition of tumor growth, or an increase in tumor necrosis, tumor shrinkage and/or tumor disappearance; (c) delay in tumor growth and development;

- 53 -

(d) inhibition of tumor metastasis; (e) prevention of recurrence of tumor growth; (f) increase in survival of a subject with a cancer; and/or (g) a reduction in the use or need for conventional anti-cancer therapy (e.g., reduced or eliminated use of chemotherapeutic or cytotoxic agents) as compared to an untreated subject or a subject administered the antibody as monotherapy.

In the case of an anti-PD-1 antibody or antigen-binding fragment thereof, a therapeutically effective amount can be from about 0.05 mg to about 600 mg, from about 1 mg to about 500 mg, from about 10 mg to about 450 mg, from about 50 mg to about 400 mg, from about 75 mg to about 350 mg, or from about 100 mg to about 300 mg of the antibody. For example, in various embodiments, the amount of the anti-PD-1 antibody is about 0.05 mg, about 0.1 mg, about 1.0 mg, about 1.5 mg, about 2.0 mg, about 10 mg, about 20 mg, about 30 mg, about 40 mg, about 50 mg, about 60 mg, about 70 mg, about 80 mg, about 90 mg, about 100 mg, about 110 mg, about 120 mg, about 130 mg, about 140 mg, about 150 mg, about 160 mg, about 170 mg, about 180 mg, about 190 mg, about 200 mg, about 210 mg, about 220 mg, about 230 mg, about 240 mg, about 250 mg, about 260 mg, about 270 mg, about 280 mg, about 290 mg, about 300 mg, about 310 mg, about 320 mg, about 330 mg, about 340 mg, about 350 mg, about 360 mg, about 370 mg, about 380 mg, about 390 mg, about 400 mg, about 410 mg, about 420 mg, about 430 mg, about 440 mg, about 450 mg, about 460 mg, about 470 mg, about 480 mg, about 490 mg, about 500 mg, about 510 mg, about 520 mg, about 530 mg, about 540 mg, about 550 mg, about 560 mg, about 570 mg, about 580 mg, about 590 mg, or about 600 mg, of the anti-PD-1 antibody. In one embodiment, 250 mg of an anti-PD-1 antibody is administered according to the methods of the present invention. In one embodiment, 200 mg of an anti-PD-1 antibody is administered according to the methods of the present invention. In one embodiment, 350 mg of an anti-PD-1 antibody is administered according to the methods of the present invention.

The anti-PD-1 antibody may be administered to the subject in multiple doses, e.g., as part of a specific therapeutic dosing regimen. For example, the therapeutic dosing regimen may comprise administering one or more doses of an anti-PD-1 antibody to the subject at a frequency of about once a day, once every two days, once every three days,

- 54 -

once every four days, once every five days, once every six days, once a week, once every two weeks, once every three weeks, once every four weeks, once a month, once every two months, once every three months, once every four months, or less frequently.

In some embodiments, the anti-PD-1 antibody is contained within a pharmaceutical composition. The pharmaceutical compositions of the invention may be formulated with suitable carriers, excipients, and other agents that provide suitable transfer, delivery, tolerance, and the like. A multitude of appropriate formulations can be found in the formulary known to all pharmaceutical chemists: Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles (such as LIPOFECTIN™), DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax. See also Powell et al. "Compendium of excipients for parenteral formulations" PDA (1998) J Pharm Sci Technol 52:238-311.

Various delivery systems are known and can be used to administer the anti-PD-1 antibody, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the mutant viruses, receptor mediated endocytosis (see, e.g., Wu et al., 1987, J. Biol. Chem. 262: 4429-4432). Methods of administration include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The composition may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents.

The anti-PD-1 antibody can be delivered subcutaneously or intravenously with a standard needle and syringe. In addition, with respect to subcutaneous delivery, a pen delivery device readily has applications in delivering the anti-PD-1 antibody. Such a pen delivery device can be reusable or disposable. A reusable pen delivery device generally utilizes a replaceable cartridge that contains a pharmaceutical composition of the anti-PD-1 antibody. Once all of the pharmaceutical composition within the cartridge has been

- 55 -

administered and the cartridge is empty, the empty cartridge can readily be discarded and replaced with a new cartridge that contains the pharmaceutical composition. The pen delivery device can then be reused. In a disposable pen delivery device, there is no replaceable cartridge. Rather, the disposable pen delivery device comes prefilled with the pharmaceutical composition held in a reservoir within the device. Once the reservoir is emptied of the pharmaceutical composition, the entire device is discarded.

In certain situations, the anti-PD-1 antibody can be delivered in a controlled release system. In one embodiment, a pump may be used. In another embodiment, polymeric materials can be used; see, *Medical Applications of Controlled Release*, Langer and Wise (eds.), 1974, CRC Pres., Boca Raton, Fla. In yet another embodiment, a controlled release system can be placed in proximity of the target, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, 1984, in *Medical Applications of Controlled Release*, supra, vol. 2, pp. 115-138). Other controlled release systems are discussed in the review by Langer, 1990, *Science* 249:1527-1533.

Injectable preparations of the anti-PD-1 antibody may include dosage forms for intravenous, subcutaneous, intracutaneous and intramuscular injections, drip infusions, etc. These injectable preparations may be prepared by known methods. For example, the injectable preparations may be prepared, e.g., by dissolving, suspending, or emulsifying the antibody or its salt described above in a sterile aqueous medium or an oily medium conventionally used for injections. As the aqueous medium for injections, there are, for example, physiological saline, an isotonic solution containing glucose and other auxiliary agents, etc., which may be used in combination with an appropriate solubilizing agent such as an alcohol (e.g., ethanol), a polyalcohol (e.g., propylene glycol, polyethylene glycol), a nonionic surfactant [e.g., polysorbate 80, HCO-50 (polyoxyethylene (50 mol) adduct of hydrogenated castor oil)], etc. As the oily medium, there are employed, e.g., sesame oil, soybean oil, etc., which may be used in combination with a solubilizing agent such as benzyl benzoate, benzyl alcohol, etc. The injection thus prepared is preferably filled in an appropriate ampoule.

In certain embodiments, the anti-PD-1 antibody is formulated in a pharmaceutical composition for use in intravenous administration.

- 56 -

Combination with Radiation Therapy or Chemotherapy

In certain embodiments, the methods further comprise administering radiation therapy to the subject. In certain embodiments, the one or more doses of radiation therapy are administered to the subject at a frequency of about once a day, once every two days, once every three days, once every four days, once every five days, once every six days, once a week, once every two weeks, once every three weeks, once every four weeks, once a month, once every two months, once every three months, once every four months, or less frequently.

In certain embodiments, the radiation therapy is hypofractionated radiation therapy. In some embodiments, each dose of radiation therapy comprises 20-50 Gy. In some embodiments, the subject is administered 20-50 Gy in 2-20 fractions. In some embodiments, each dose of radiation therapy comprises 20-60 Gy. In some embodiments, the subject is administered 20-60 Gy in 2-20 fractions. In certain embodiments, the fractionated radiation therapy comprises 40 Gy in 15 fractions.

In certain embodiments, the hypofractionated radiation therapy comprises 15 fractions. In certain embodiments, the 15 fractions are administered on 15-25 consecutive days. In certain embodiments, the 15 fractions are administered on 21 consecutive days.

In certain embodiments, the methods further comprise administering a chemotherapeutic agent to the subject, for example, temozolomide (TMZ). The chemotherapeutic agent can be administered with the radiation therapy. For example, TMZ is administered at a daily dose of 75 mg/m^2 concomitant with hypofractionated radiation therapy. In some embodiments, subjects having a tumor with a methylated MGMT promoter will be administered maintenance therapy of the chemotherapeutic agent. For example, following radiation therapy, subjects having a tumor with a methylated MGMT promoter may receive TMZ at a starting dose of $150 \text{ mg/m}^2/\text{day}$ for 6 cycles on the first 5 days of a 28-day cycle (5 days “on,” 23 days “off”) with increased each maintenance cycle by $50 \text{ mg/m}^2/\text{dose}$ to a maximum of $200 \text{ mg/m}^2/\text{dose}$, in the absence of hematologic toxicity. In some embodiments, the maintenance therapy will start

- 57 -

approximately three to five weeks, preferably about 4 weeks, after the last dose of radiation therapy.

EXAMPLES

The examples and embodiments described herein are for illustrative purposes only and various modifications or changes suggested to persons skilled in the art are to be included within the spirit and purview of this application and scope of the appended claims.

EXAMPLE 1: A Study of INO-5401 and INO-9012 Delivered by Electroporation (EP) in Combination With Cemiplimab (REGN2810) in Newly-Diagnosed Glioblastoma (GBM)

A non-limiting example of a phase 1/2 trial to evaluate safety, immunogenicity, and preliminary efficacy of INO-5401 and INO-9012 in combination with cemiplimab (REGN2810), with radiation and chemotherapy, in subjects with newly-diagnosed glioblastoma (GBM) is provided herein.

Objectives

Primary Objective: To evaluate the safety and tolerability of INO-5401 and INO-9012 delivered by intramuscular (IM) injection followed by EP with CELLECTRA® 2000 in combination with cemiplimab-rwlc in adult subjects with newly-diagnosed GBM.

Primary Endpoint(s) and Assessments:

- Incidence of adverse events (AE) graded per Common Toxicity Criteria for Adverse Events (CTCAE) v4.03, classified by system organ class, preferred term, severity, and relationship to trial treatment.
- Clinically significant changes in safety laboratory parameters from baseline.

Secondary Objectives:

- To evaluate preliminary clinical efficacy and immunogenicity of INO-5401 and INO-9012 delivered by IM injection followed by EP with CELLECTRA® 2000 in combination with cemiplimab-rwlc in adult subjects with newly-diagnosed GBM.
- To evaluate preliminary immunogenicity of INO-5401 and INO-9012

- 58 -

delivered by IM injection followed by EP with CELLECTRA® 2000 in combination with REGN2810 in adult subjects with newly-diagnosed GBM.

Secondary Endpoint(s) and Assessments:

- Overall survival at 18 months (OS18);
- Antigen-specific cellular immune responses assessed by:
 - Interferon- γ secreting T lymphocytes in peripheral blood mononuclear cells (PBMC) by ELISpot;
 - T-cell phenotype (e.g. activation and cytolytic cell, myeloid derived suppressor cell frequency (MDSC)) in PBMC by Flow Cytometry;
 - T cell receptor (TCR) sequencing from PBMCs to assess diversity and putative antigen specificity;
- Antigen-specific humoral responses (e.g. B cell activation/antibody secretion).

Exploratory Objective(s):

- To explore correlative association between clinical efficacy and tumor genetics and/or biomarkers.
- To further evaluate efficacy of INO-5401 and INO-9012 delivered by IM injection followed by EP with CELLECTRA® 2000 in combination with REGN2810 and hypofractionated radiation therapy in adult subjects with newly-diagnosed GBM. Exploratory Endpoint(s):
- Tumor infiltrating lymphocytes (TILs) and immunosuppressive elements, where feasible;
- Expression of tumor oncoproteins, including but not limited to tumor expression of hTERT, WT1, and PSMA by IHC, immunofluorescence (IF) or genome sequencing;
- MicroRNA signatures in blood plasma and/or sera;
- Circulating tumor cells, circulating endothelial cells, and/or circulating cancer associated macrophage-like cells from peripheral blood where feasible;

- 59 -

- Assessment of tumor-associated antigen (TAA)-specific peripheral T cells by RNAseq;
- Assessment of cytokine profiles from plasma and/or sera;
- Progression-Free Survival, as assessed by RANO (Response Assessment in Neurooncology) criteria and Immunotherapy Response Assessment in Neuro- oncology (iRANO) criteria;
- Overall Survival (OS).

Study Design

The study described in this example corresponds to ClinicalTrials.gov identifier NCT03491683. The data presented herein as associated with this study reflects the state of this study as of the time of this filing. In this study, antigen-specific T cell-generating therapy, INO-5401, combined with INO-9012, followed by electroporation with the CELLECTRA® 2000 device, together with a PD-1 checkpoint inhibitor, cemiplimab-rwlc, was given to patients with newly-diagnosed GBM, together with radiation and temozolomide, in order to evaluate tolerability, immunogenicity and anti-tumor activity of the combination. Ethics Approval by NYU Ethics Board; approval number 17-00764.

This is a Phase 1/2, open-label, multi-center trial to evaluate the safety, immunogenicity, and preliminary efficacy of INO-5401 and INO-9012 in combination with cemiplimab (also known as REGN2810) in subjects with newly-diagnosed GBM. All patients provided written informed consent.

Subjects started immunotherapy with REGN2810 upon definitive histopathological diagnosis of GBM and adequate recovery from surgical intervention. Subjects were assigned to a cohort based on the results of the MGMT gene methylation assay performed in a CLIA-certified laboratory, which was available prior to the completion of RT. The start of immunotherapy is designated as Day 0. REGN2810 was administered intravenously (IV) every three weeks until disease progression defined by iRANO (Immune Response Assessment in Neuro-Oncology), unacceptable toxicity, withdrawal of consent, or death.

On Day 0, subjects received INO-5401 and INO-9012 intramuscularly (IM) followed by electroporation (EP). INO-5401 and INO-9012 were administered, followed

- 60 -

by EP, every three weeks for four doses, and then every 9 weeks until disease progression defined by iRANO, unacceptable toxicity, withdrawal of consent, or death.

Temozolomide was administered to all subjects both with and without MGMT promoter methylation, unless clinically contraindicated, during radiation therapy. Radiation therapy (RT) began no later than 42 days after surgical intervention. Radiation therapy started approximately 1 to 2 weeks after Day 0, and was continue(d) for approximately three weeks. Temozolomide (TMZ) was given daily during radiation therapy (TMZ/RT). Subjects with MGMT promoter methylation received maintenance (adjuvant) TMZ for 6 cycles, following recovery from TMZ/RT. Maintenance (adjuvant) TMZ was administered for the first 5 days of a 28-day cycle. This study had two cohorts: Cohort A, consisting of subjects with an unmethylated MGMT promoter, and Cohort B consisting of subjects with a methylated MGMT promoter.

Study Population

Each potential subject satisfied all of the following criteria to be enrolled in the study: Overview of Patient Eligibility

Adults with newly-diagnosed GBM who are post-definitive surgery, and are able to receive standard therapy. Estimated Number of Subjects: 52. Cohort A: MGMT promoter unmethylated (N=32 for 30 evaluable subjects). Cohort B: MGMT promoter methylated (N=20 for 19 evaluable subjects).

Inclusion Criteria:

- Subjects must provide written IRB approved informed consent in accordance with institutional guidelines;
- Be 18 years of age or older on the day of signing the informed consent, and able and willing to comply with all trial procedures;
- Newly-diagnosed brain cancer with histopathological diagnosis of glioblastoma (GBM);
- Karnofsky Performance Status (KPS) rating of ≥ 70 at baseline;
- Receipt of dexamethasone equivalent dose ≤ 2 mg per day, stable or decreased for $\geq$ three days prior to Day 0;
- Recovery from the effects of prior GBM surgery as defined by the Investigator;

- 61 -

- ECG with no clinically significant findings as assessed by the Investigator performed within 28 days of signing the informed consent form (ICF);
- Adequate organ function as demonstrated by hematological, renal, hepatic parameters as defined in Table 1 below, obtained within 28 days prior to the first trial treatment;

Table 1.

Absolute neutrophil count (ANC)	$\geq 1500/\text{mm}^3$
Platelets	$\geq 100,000/\text{mm}^3$
Hemoglobin	$\geq 9 \text{ g/dL}$
Creatinine	$\leq 1.5 \times \text{upper limit of normal (ULN)}$
OR	OR
Measured or calculated	$\geq 50 \text{ mL/min}$ for subject with creatinine level > 1.5
creatinine clearance or GFR*	$\geq \text{institutional ULN}$
Total bilirubin	$\leq 1.5 \times \text{ULN}$
AST (SGOT) and ALT (SGPT)	$\leq 2.5 \times \text{ULN}$

*Creatinine clearance should be calculated per Cockcroft-Gault equation

- Agree that, during the trial, men will not father a child, and women cannot be or become pregnant if they are of child-bearing potential. Subjects must be of non-child bearing potential (≥ 12 months of non-therapy-induced amenorrhea, confirmed by follicle stimulating hormone [FSH], if not on hormone replacement); or surgically sterile (vasectomy in males or absence of ovaries and/or uterus in females); or agree to use one highly effective or combined contraceptive methods that result in a failure rate of $< 1\%$ per year during the treatment period and at least through week 12 after last dose. Periodic abstinence (e.g., calendar, ovulation, symptothermal, or post-ovulation methods) and withdrawal are not acceptable methods of contraception. Examples of contraceptive methods with an expected failure rate of $< 1\%$ per year include male sterilization and hormonal implants. Alternatively, proper use of combined oral or injected hormonal contraceptives and certain intrauterine devices (IUDs) or two methods (e.g., two barrier methods such as a condom and a cervical cap) may be combined to achieve a failure rate of $< 1\%$ per year (barrier methods must always be supplemented with the use of a spermicide);

- Ability to tolerate magnetic resonance imaging (MRI).

Exclusion Criteria:

- Presence of greater than 1 cm x 1 cm residual tumor enhancement on post-

- 62 -

operative MRI;

- Multifocal disease or leptomeningeal disease (LM) disease on post-operative

MRI;

- Are not able to start radiation within 42 days of surgical resection of their tumor;
- Receive dexamethasone equivalent dose >2 mg per day;
- Prior treatment with an agent that blocks the PD-1/PD-L1 pathway at any point

in the past;

- Receipt of previous approved or investigative immune modulatory agent (for example, anti-TNF, therapeutic anti-cancer vaccines, cytokine treatments (other than G-CSF or erythropoietin), or agents that target cytotoxic T-lymphocyte antigen 4 (CTLA-4), 4-1BB (CD137), PI3K-delta, or OX-40) within 28 days of receiving the first dose of treatment;

- Have received prior treatment with idelalisib at any point in the past;

- Past, current, or planned treatment with tumor treatment fields (Optune;

NovoTTF); oncolytic viral treatment; or prior exposure to an investigational agent or device, including Gliadel wafer (Carmustine) implant for chemotherapy; within 28 days of receiving the first dose of treatment;

- Allergy or hypersensitivity to REGN2810 or to any of its excipients;
- History of documented allergic reactions or acute hypersensitivity reaction

attributed to antibody treatments;

- Ongoing or recent (within 5 years) evidence of autoimmune disease that required treatment with systemic immunosuppressive treatments, which may suggest risk for immune-related adverse events (irAEs), with the exception of: vitiligo, childhood asthma that has resolved, type 1 diabetes, residual hypothyroidism that required only hormone replacement, or psoriasis that does not require systemic treatment;

- Diagnosis of immunodeficiency or treatment with systemic immunosuppressive therapy within 28 days prior to the first dose of trial treatment, other than dexamethasone for the underlying disease under investigation, as noted in the inclusion criteria;

- Positive serological test for human immunodeficiency virus (HIV), or a history

- 63 -

of HIV infection; or positive tests for hepatitis B virus surface antigen (HBV sAg) or hepatitis C virus ribonucleic acid (HCV RNA) indicating active or chronic infection, as these infections may interfere with the ability to mount an appropriate immune response to vaccination;

- Current malignancy at another site, with the exception of adequately treated basal or squamous cell skin cancers, or carcinoma of the cervix in situ, with no evidence of disease within 3 years. Cancer survivors who have undergone curative therapy for a prior malignancy, have no evidence of disease for 3 years and are deemed at low risk for recurrence are eligible for the trial;

- Receipt of any vaccine within 4 weeks prior to first dose of trial treatment with the exception of the inactivated influenza vaccine, which may be given up to 2 weeks prior;

- History of clinically significant, medically unstable disease which, in the judgment of the investigator, would jeopardize the safety of the subject, interfere with trial assessments or endpoint evaluation, or otherwise impact the validity of the trial results (e.g. chronic renal failure, angina, myocardial ischemia or infarction, New York Heart Association (NYHA) class III/ IV cardiac disease); or any cardiac pre-excitation syndromes (such as Wolff- Parkinson-White; cardiomyopathy, or clinically significant arrhythmias);

- History of pneumonitis within the last 5 years;

- Acute or chronic bleeding or clotting disorder that would contraindicate IM injections or use of blood thinners (e.g. anticoagulants or antiplatelet drugs, excluding over-the-counter aspirin or non-steroidal anti-inflammatory drugs, such as ibuprofen) within 2 weeks of Day 0;

- Fewer than two acceptable sites available for IM injection considering the deltoid and anterolateral quadriceps muscles. The following are unacceptable sites:

- Tattoos, keloids, or hypertrophic scars located within 2 cm of intended treatment site;
 - Cardioverter-defibrillator or pacemaker (to prevent a life-threatening arrhythmia) that is located ipsilateral to the deltoid injection site (unless deemed acceptable by a cardiologist);

- 64 -

- Metal implants or implantable medical device within the intended treatment site (i.e. EP area);
- Active drug or alcohol use or dependence that, in the opinion of the Investigator, would interfere with adherence to trial requirements;
- Imprisonment, or compulsory detainment (involuntary incarceration) for treatment of either a psychiatric or physical (i.e. infectious disease) illness;
- Pregnant or current breastfeeding;
- As determined by the Investigator, any medical or psychological or non-medical condition that might interfere with the subject's ability to participate or affect the safety of the subject.

Dosage and Administration

Investigational Drug Products are provided in Table 2.

Table 2.

Product	Formulation	Unit
INO-5401	10.0 ± 0.5 mg/mL total plasmid (pGX1108, pGX1404, pGX1434; 1:1:1 w/w) in 165 mM sodium chloride and 16.5 mM sodium citrate	1.4 mL/ 2-mL vial
INO-9012	10.0 ± 0.5 mg/mL pGX6001 in water for injection	0.2 mL/ 2-mL vial
REGN2810	REGN2810 is supplied as a sterile liquid solution of 5.5 mL in a 10 mL glass vial (50 mg/mL) for IV administration. REGN2810 may also be supplied as a sterile liquid solution of 7.44 mL in a 10 mL glass vial (50 mg/mL) for IV administration.	5.5 mL/ 10-mL vial Or 7.44 mL/ 10-mL vial

The active pharmaceutical ingredients (APIs) in INO-5401 are DNA plasmid sequences that were designed and constructed using proprietary synthetic consensus (SynCon®) technology. This process involves synthetically deriving consensus genes across multiple strains and optimizing DNA inserts at the genetic level to allow high expression in human cells. The INO-5401 plasmids are as follows:

- pGX1108, a plasmid for expression of prostate-specific membrane antigen (PSMA; SEQ ID NO: 28). 3 mg of pGX1108 will be present in each 10 mg dose of trial treatment (INO-5401 + INO-9012).

- 65 -

- pGX1404, a plasmid for expression of Wilms' tumor gene-1 (WT1) antigen (SEQ ID NO: 26). 3 mg of pGX1404 will be present in each 10 mg dose of trial treatment.
- pGX1434, a plasmid for expression of human telomerase reverse transcriptase (hTERT) (SEQ ID NO: 20). 3 mg of pGX1434 will be present in each 10 mg dose of trial treatment.

The API in drug product INO-9012 is pGX6001, a DNA plasmid for expression of human IL-12 p35 (SEQ ID NO: 22) and p40 (SEQ ID NO: 25) subunit proteins. 1 mg of pGX6001 will be present in each 10 mg dose of trial treatment. Both DNA plasmid products INO-5401 and INO-9012, are administered using a syringe and the investigational CELLECTRA® 2000 electroporation (EP) device.

Cemiplimab-rwlc (REGN2810) is a covalent heterotetramer consisting of two disulfide-linked human heavy chains, each of which is covalently bonded through disulfide linkages to a human kappa light chain. The antibody possesses an approximate molecular weight of 143.6 kDa based on the primary sequence. There is a single N-linked glycosylation site on each heavy chain, located within the constant region in Fc portion of the molecule.

The REGN2810 heavy chain possesses an IgG4 isotype-constant region. The variable domains of the heavy and light chains combine to form PD-1 binding site within the antibody. Antibody generation by VelocImmune® mice is carried out using standard techniques after immunization with PD-1. The genes encoding the heavy and light chains of REGN2810 were introduced into CHO cells, and a stable expression cell line with a higher titer (Cell Line 2) was developed for this antibody. For both cell lines, the recombinant CHO cells were grown in suspension culture and chemically induced to initiate antibody expression and secretion into the cell culture medium. Antibody is harvested via filtration and purified through a series of preparative column chromatographic and filtration steps to generate drug substance. Drug substance is then formulated and sterile-filtered to produce the final drug product.

REGN2810 (50 mg/mL) is formulated in an aqueous buffered solution at pH 6.0 containing 10 mM histidine, 5% (w/v) sucrose, 1.5% (w/v) L-proline, and 0.2% (w/v)

- 66 -

polysorbate 80. REGN2810 is supplied as a sterile liquid solution of 5.5 mL in a 10 or 20 mL glass vial for IV administration. A maximum volume of 5.0 mL can be withdrawn from each vial containing 250 mg of REGN2810. Seven mLs are needed to provide a 350 mg dose of REGN2810, thus 2 vials must be used when supplied with the 5 mL vials. REGN2810 (50 mg/mL) may also be supplied as a sterile liquid solution of 7.44 mL in a 10 or 20 mL glass vial for IV administration. A maximum volume of 7.0 mL can be withdrawn from each vial containing 350 mg of REGN2810.

Treatment

Subjects who meet all of the inclusion criteria and none of the exclusion criteria started immunotherapy with REGN2810 and INO-5401 + INO-9012 on Day 0. REGN2810 was administered IV every three weeks at a dose of 350 mg per dose, in the absence of dose holding, until disease progression as defined by iRANO, unacceptable toxicity, withdrawal of consent, or death. INO-5401 and INO-9012 IM followed by EP was administered every three weeks for four doses, and then every 9 weeks, at a dose of 10 mg/DNA per dose, in the absence of dose holding, until disease progression as defined by iRANO, unacceptable toxicity, withdrawal of consent, or death. RT began no later than 42 days after surgical intervention and approximately 1 to 2 weeks after Day 0. RT continued for approximately three weeks. The total dose of RT was 40 Gy given over 3 weeks.

Daily TMZ with radiation therapy (TMZ/RT) began no later than 42 days after surgical intervention and approximately 2 weeks after Day 0. TMZ/RT continued for approximately three weeks. TMZ was given at a dose of 75 mg/m²/dose, in the absence of dose reduction. Subjects should then receive maintenance (adjuvant) TMZ for an additional 6 cycles. Cohort B received TMZ following radiotherapy for up to six cycles. Maintenance (adjuvant) TMZ was administered to subjects in Cohort B for the first 5 days of a 28-day cycle at 150-200 mg/m²/dose, following peripheral blood count recovery from TMZ/RT per standard guidelines TMZ treatment.

Day 0 (first dose of INO 5401, INO 9012 and REGN2810) was at least 14 days after completion of resection of primary tumor and the subject has recovered from surgery, but no later than post-operative day 28.

- 67 -

For subjects who discontinued one therapy (either INO-5401 + INO-9012 or REGN2810) for reasons other than progression, the other therapy was allowed to continue after consultation with the Medical Monitor.

FIG. 1 illustrates the trial design for Cohorts A and B.

- INO-5401 (3 mg each of hTERT, WT-1 and PSMA plasmids) combined with 1 mg INO-9012 (IL-12), for a total of 10 mg of DNA, administered via intramuscular (IM) injection followed by electroporation (EP) with CELLECTRA® 2000 device, and delivered every three weeks for four doses, then every 9 weeks.
- Chemoradiation: Radiation (RT) given in a hypofractionated schedule (40 Gy over three weeks)
- Temozolomide (TMZ) concurrent with radiation (Cohorts A and B), followed by six maintenance (adjuvant) cycles (Cohort B only)
- Cemiplimab-rwlc (REGN2810) was administered IV at a dose of 350 mg every three weeks (Q3W) over approximately 30 minutes, starting at Day 0, and continued until disease progression as defined by iRANO, unacceptable toxicity, withdrawal of consent, or death.

INO-5401 is a mixture of three separate synthetic plasmids that target WT1, PSMA and hTERT proteins. Each plasmid was dosed at 3 mg DNA, for a total of 9 mg DNA per dose of INO-5401. INO-5401 was administered IM at Day 0, Week 3, Week 6, and Week 9, and then every 9 weeks thereafter, and continued until disease progression as defined by iRANO, unacceptable toxicity, withdrawal of consent, or death. INO-9012 is a synthetic plasmid that expresses human IL-12, and is dosed at 1 mg DNA, and was administered IM together with INO-5401. The total dose of DNA in each dose of INO-5401 + INO-9012 when mixed and administered together was 10 mg. INO-5401 (3 mg each of hTERT, WT-1 and PSMA plasmids) combined with 1 mg INO-9012 (IL-12), for a total of 10 mg of DNA, was administered via intramuscular (IM) injection followed by electroporation (EP) with CELLECTRA® 2000 device, and delivered every three weeks for four doses, then every 9 weeks.

All subjects received a total of 40 Gy in 15 fractions (three weeks).

- 68 -

Hypofractionated radiation therapy (hfRT) began no later than 42 days after surgery. Radiotherapy was given for three weeks.

All patients received TMZ, regardless of MGMT methylation status, unless clinically contraindicated, during radiotherapy. TMZ was administered at 75 mg/m² daily by mouth for 21 days concomitant (7 days a week for three weeks) with hfRT therapy.

Following radiation therapy, subjects with MGMT promoter methylation (Cohort B) continued TMZ maintenance therapy at a starting dose of 150 mg/m²/day for 6 cycles on the first 5 days of a 28-day cycle (5 days “on,” 23 days “off”), and increased each maintenance cycle by 50 mg/m²/dose to a maximum of 200 mg/m²/dose, in the absence of hematologic toxicity. Maintenance (adjuvant) TMZ started approximately four weeks after the last dose of RT ($\pm$ 3 days) and following peripheral blood count recovery, per TMZ treatment guidelines. The dose was determined using actual body surface area (BSA) as calculated in square meters at the beginning of each treatment cycle.

The daily dose was rounded to the nearest 5 mg.

For detailed information on efficacy evaluations/endpoints ELISpot, lytic granule loading, safety assessments, medical and clinical assessments, physical assessments, medical post-treatment reaction assessment, vital signs, weight, height, and body mass index, 12-lead ECG, pregnancy test, laboratory evaluations, peripheral blood immunogenicity assessments, tissue immunology, clinical assessments, documentation of overall survival and progression, rano and irano (immunotherapy) response assessment in neuro-oncology), nanoscale neurological assessment in neuro-oncology, statistical and analytical plan, statistical hypothesis, analysis populations and datasets, subject disposition, and safety data are described in WO 2021/092019, the entirety of which is incorporated herein by reference.

Exploratory Analyses

Exploratory tumor genetics and/or biomarker responses were presented using descriptive statistics at each time point and as changes from baseline for the mITT population and per protocol population.

Results

Demographics of enrolled patients at the 18-month analysis timepoint are shown in **FIG. 2A** (Demographics Table). Updated patient demographics as assessed at the 42-

- 69 -

month analysis timepoint are provided in **FIG. 2B**. Assessment of gene transcripts from tumors at diagnosis confirmed expression of antigens encoded by INO-5401 and a diverse immune gene profile. 5 of 47 subjects (811%) with available tissue for assessment exhibited transcript expression of one or more Tumor Associated Antigens encoded by INO-5401 (WT1, PSMA and hTERT). 43 of 47 subjects (89%) with available tissue for assessment exhibited transcript expression of two (2) or more Tumor Associated Antigens encoded by INO-5401 (WT1, PSMA and hTERT). 19 of 47 subjects (40%) exhibited transcript expression of all three Tumor Associated Antigens. No subject exhibited PD-1 expression without concomitant PD-L1 expression. 27 of 47 subjects (57%) exhibited PD-L1 expression with no concomitant PD-1 expression. 20 of 47 subjects (43%) showed co-expression of PD-1 and PD-L1. Normalized transcript read counts >1 were considered “positive.”

MRI Imaging

Several patients have experienced pseudo-progression, with radiographic evidence of progression on MRI without evidence of tumor on repeat biopsy. Images from example patients demonstrate increase in MRI signal at timepoints following first dose of INO-5401 + INO-9012 and cemiplimab-rwlc, suggestive of edema or tumor. Biopsy on several patients shows treatment-related changes with necrosis and mixed inflammation; absence of mitotic activity; and no evidence of viable tumor. Representative images from two patients are presented in **FIG. 3**.

ELISpot

ELISpot is/was employed to give a qualitative measure of whether antigen specific T cells are present in a peripheral blood mononuclear cell (PBMC) sample. PBMCs are collected from subjects at study weeks 0, 3, 6, 9, 12, and 24 and assayed by IFN-g ELISpot. At the 12-month data cut-off, antigen specific IFN-g spot forming units (SFU) per million PBMCs are shown from before (pre) and the highest magnitude (peak) after treatment with INO-5401 and cemiplimab-rwlc from 8 subjects with sample to week 24 (**Figure 4**). Each subject is represented by an open circle, bars represent the mean. The difference from pre to peak, delta, is shown for each antigen graph as well as together for 11 subjects assayed and for the 8 with sample available to week 24. INO- 5401 is the sum of WT1, PSMA and

- 70 -

hTERT. Box plots extend from the 25th to 75th percentile, with a horizontal line at the median, and “+” at the mean. ELISpot results support the combination of INO-5401 and cemiplimab-rwlc are immunogenic- with IFN-g magnitudes above baseline to all 3 antigens in 5/11 subjects and to at least one antigen in 9 subjects as shown in **FIG. 4**.

Assessment of post-INO-5401 peripheral immune responses by Cohort at the 18-month data cut-off revealed antigen-specific T cell responses by Interferon gamma ELISpot (cytokine production in response to each component of INO-5401). Results of assessment of post-INO-5401 peripheral immune responses by ELISpot at by Cohort are provided in **FIG. 16A and FIG. 16B**. Baseline values from the peak timepoint following treatment are plotted. In Cohort A, 19/22 (86%) subjects tested to date had an IFN-g magnitude above baseline to one or more of the INO-5401 antigens (FIG. 16A). In Cohort B, 16/17 (94%) subjects tested to date had an IFN-g magnitude above baseline to one or more of the INO-5401 antigens (FIG. 16B).

Lytic Granule Loading

A lytic granule loading assay was performed to explore the activation status and lytic potential of antigen specific T cells present in PBMC samples collected from subjects at study weeks 0, 3, 6, 9, 12, and 24. PBMCs were stimulated with overlapping peptide libraries for INO-5401 antigens (hTERT, PSMA and WT1) or relevant controls in the absence of any exogenous cytokines. After 5 days, cells were stained with antibodies and assessed by flow cytometry. Frequencies of live, antigen-specific, activated (CD38+) CD3+CD8+ T cells with lytic potential (expressing Granzyme A, Perforin) are shown from before treatment at baseline (pre) and the highest magnitude (peak) after treatment with INO-5401 and cemiplimab-rwlc from 8 subjects for each antigen (**FIG. 5A**). Each subject is represented by an open circle, bars represent the mean. The difference from pre to peak, delta, is shown for each antigen as well as INO-5401 at the 12-month data cut-off for 8 subjects assayed (**FIG. 5B**) and for the 5/8 subjects with sample available to week 12 (**FIG. 5C**). INO-5401 is the sum of WT1, PSMA and hTERT. Box plots extend from the 25th to 75th percentile, with a horizontal line at the median, and “+” at the mean. Five subjects had a frequency of activated CD8+T cells with lytic potential (CD38+Prf+GrzA+) above baseline (pre) to more than

one antigen; three subjects had a frequency of activated CD8+T cells with lytic potential (CD38+Prf+GrzA+) above baseline to all three antigens. Three subjects did not have a response above baseline to any antigen at any time.

Assessment of post-INO-5401 peripheral immune responses by Cohort at the 18-month data cut-off revealed antigen-specific T cell responses by flow cytometry (the expansion of antigen specific CD8+ T cells with lytic potential). In Cohort A, 13/19 (68%) subjects tested to date had a frequency of CD38+GrzA+Prf+ CD8+T cells above baseline to one or more of the INO-5401 antigens (**FIG. 17A**). In Cohort B, 8/10 (80%) subjects tested to date had a frequency of CD38+GrzA+Prf+ CD8+T cells above baseline to one or more of the INO-5401 antigens (**FIG. 17B**). Samples were collected Q3 weeks x 4 and then Q12 weeks. Baseline values from the peak timepoint following treatment are plotted.

Progression-Free Survival

FIG. 6 shows the visual representation of the Kaplan-Meier estimator of the progression-free survival at six months (PFS6) for Cohort A, patients with the O6-methylguanine methyltransferase gene promoter unmethylated in their tumor cells. The curve shows the probability of an event at a certain time interval. The probability of the event is represented numerically on the y-axis, and the time interval on the x-axis. The event shown is progression-free survival. Progression-free survival is the absence of progression of disease at a given time point for a given subject.

FIG. 7 shows the visual representation of the Kaplan-Meier estimator of the progression-free survival at six months (PFS6) for Cohort B, patients with the O6-methylguanine methyltransferase gene promoter methylated in their tumor cells. The curve shows the probability of an event at a certain time interval. The probability of the event is represented numerically on the y-axis, and the time interval on the x-axis. The event shown is progression-free survival. Progression-free survival is the absence of progression of disease at a given time point for a given subject.

FIG. 8 shows the visual representation of the Kaplan-Meier estimator of the progression-free survival at six months (PFS6) for Cohort A and Cohort B, patients with the O6-methylguanine methyltransferase gene promoter unmethylated or methylated in their tumor cells. The curve shows the probability of an event at a certain time interval. The

- 72 -

probability of the event is represented numerically on the y-axis, and the time interval on the x-axis. The event shown is progression-free survival. Progression-free survival is the absence of progression of disease at a given time point for a given subject.

FIG. 9 shows the tabular representation of the Kaplan-Meier estimator of the progression-free survival at six months (PFS6) for Cohort A, Cohort B, and both cohorts combined. The total number of subjects per cohort, number of events, estimation of the event (PFS6), and the 95% confidence interval (CI) in which the numerical estimate of the event (PFS6) exists are all provided.

Confirmed progressive disease (PD) is determined by confirmation by consecutive PD scan ≥ 4 weeks from original PD event, or progressed according to biopsy surgery. Subjects who terminated for any reason prior to 6 months other than PD included as confirmed progressive events, including two (2) subjects in Cohort B who came off-study at week three (3), and declined long-term follow-up.

Overall Survival

All efficacy analyses (OS12, OS18, & Kaplan-Meier) were analyzed on subjects in the modified intent-to-treat (mITT) population which is defined as all subjects who received at least one dose of planned treatment. Overall Survival at twelve months (OS12) was tabulated as the proportion of subjects alive at 12 months out of all subjects at risk of death at the start of the study. Subjects who dropped out were considered failures (that is deaths). All subjects in Cohort A who dropped out prior to 12 months also died before 12 months of follow-up. The 95% confidence intervals (CIs) are calculated using the Clopper-Pearson exact method. Overall Survival at eighteen months (OS18) was tabulated as the proportion of subjects alive at 18 months out of all subjects at risk of death at the start of the study. The 95% confidence intervals (CIs) are calculated using the Clopper-Pearson exact method.

FIG. 10A shows the visual representation of the Kaplan-Meier estimator of the overall survival probability over twelve months for Cohort A, for patients with the O6-methylguanine methyltransferase gene promoter unmethylated in their tumor cells. The stepwise curve shows the probability of surviving up to and beyond a specific time point. The survival probability is represented numerically on the y-axis, and survival time in days on the x-axis. **FIG. 10B** shows the visual representation of the Kaplan-Meier estimator of the overall survival

probability over eighteen months for Cohort A, for patients with the O6-methylguanine methyltransferase gene promoter unmethylated in their tumor cells. The stepwise curve shows the probability of surviving up to and beyond a specific time point. The survival probability is represented numerically on the y-axis, and survival time in days on the x-axis. Median follow-up in Cohort A is 17.8 months. mITT includes any subject who received ≥ 1 dose of study therapy. Shading represents confidence band on point estimate for survival at that timepoint.

FIG. 10C shows the visual representation of the Kaplan-Meier analysis of the overall survival for MGMT unmethylated (miTT) patients up to 42 months (cohort A). At risk = number who have not met OS endpoint or have not been censored by a given timepoint. NR = not reached. **FIG. 11A** shows the visual representation of the Kaplan-Meier estimator of the overall survival probability over twelve months for Cohort B, for patients with the O6-methylguanine methyltransferase gene promoter methylated in their tumor cells. The stepwise curve shows the probability of surviving up to and beyond a specific time point. The survival probability is represented numerically on the y-axis, and survival time in days on the x-axis.

FIG. 11B shows the visual representation of the Kaplan-Meier estimator of the overall survival probability over eighteen months for Cohort B, for patients with the O6-methylguanine methyltransferase gene promoter methylated in their tumor cells. The stepwise curve shows the probability of surviving up to and beyond a specific time point. The survival probability is represented numerically on the y-axis, and survival time in days on the x-axis. Median follow-up in Cohort B is 15.6 months. Censored; two subjects in Cohort B withdrew consent for follow-up at Week 3. mITT includes any subject who received ≥ 1 dose of study therapy. Shading represents confidence bands on point estimate for survival at that timepoint.

FIG. 11C shows the visual representation of the Kaplan-Meier analysis of the overall survival for MGMT methylated (miTT) patients up to 42 months (cohort B). At risk = number who have not met OS endpoint or have not been censored by a given timepoint. NR = not reached.

FIG. 12 shows the visual representation of the Kaplan-Meier estimator of the overall survival probability over twelve months for Cohorts A + B combined. The stepwise curve shows the probability of surviving up to and beyond a specific time point. The survival probability is represented numerically on the y-axis, and survival time in days on the x-axis.

- 74 -

FIG. 13 shows the efficacy data of the overall survival at 12 months and 18 months for Cohort A, for Cohort B, and combined. The figure shows the total number of subjects who were reported alive at 12 months and at 18 months. The total number of subjects, estimation of the event (OS12 or OS18), and the 95% confidence interval (CI) in which the numerical estimate of the event (OS12 or OS18) exists are all provided. The 95% CI were calculated using the exact Clopper-Pearson method.

Median overall survival values for Cohorts A and B for over a period of 42 months are provided in Table A.

Table A

Median OS; unmethylated (A)	17.9 months (14.5-19.8)	Historical 14.6-16 months
Media OS; methylated (B)	32.5 months (18.4-NR)	Historical 23.2-25 months
Median OS; combined (A+B)	19.5 months (16.9-23.3)	-

Conclusions

In patients with newly diagnosed GBM, INO-5401 + INO-9012 in combination with cemiplimab-rwlc, given with radiation and temozolomide, has an acceptable safety profile, is immunogenic and is potentially efficacious in patients with newly diagnosed GBM. Common AEs included injection site administration events; $\geq$ Grade 3 AEs were primarily due to TMZ or radiation, and immune-related AEs were consistent with the profile of cemiplimab-rwlc. SAEs were consistent with those seen in patients with GBM (seizure).

Antigen-specific T cell responses were seen to one or more of the antigens included in INO- 5401 in almost all patients tested to date. PFS6 exceeds that of historical controls in this study, in patients with and without MGMT promoter methylation, and OS12 exceeds that of historical controls in patients without MGMT promoter methylation [Stupp R, Mason WP, van den Bent MJ, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. N Engl J Med 2005, 352:987-996]. Treatment of subjects without methylation of the promoter region of the MGMT gene with INO-5401 resulted in an Overall Survival rate of 50% at 18 months following resection of tumor.

EXAMPLE 2: Biomarker Analysis

Biomarker assessments were performed from peripheral and tissue samples referenced in the phase 1/2 trial described in Example 1, above, and collected per lab manual instructions. Immunohistochemical assessment of the expression of the hTERT, WT1 and/or PSMA proteins within tissue samples from enrolled subjects occurred contingent on the presence of sufficient sample quantity and continued relevance as supported by available data. IDH-1 status was performed on tumor tissue if available. MicroRNA signatures in blood plasma and/or sera were assessed in order to determine disease and/or therapy specific signatures that predict disease course and/or response to treatment with INO-5401 as well as INO-5401 driven changes. RNAseq may be used for this method.

MicroRNA Signature

miRNA as a novel prognostic marker of clinical response to INO-5401 for treatment of glioblastoma

Plasma samples taken from Cohort A in trial described in Example 1 were assessed in a post-hoc fashion for the presence of miRNAs using an RNA sequencing platform for the purpose of exploration of dynamic regulation of miRNAs during the treatment process. However, and surprisingly, analysis of miRNA data taken prior to treatment with INO-5401 suggested that the presence and amount of particular miRNAs could be prognostic for response to treatment in the form of survival out to 18 months post tumor resection.

Final selection of specific prognostic miRNA species for response to INO-5401, including graphical representations of quantitative differences and tabular outputs:

Final selection of miRNA species for use as a pre-treatment indicator overall survival at 18 months during treatment with INO-5401 using post-hoc data was achieved by requiring both Negative and Positive Predictive Values as well as Sensitivity and Specificity to exceed 70%. Once miRNA species meeting these requirements were identified, TMM values for the group exhibiting each clinical outcome were individually graphed along with data outputted in tabular form. These single miRNA species, when put together into a “composite” signature, result in a high prognostic value. Graphs of miR-1537-3p normalized transcripts

- 76 -

and miR-33-3p normalized transcripts are provided in **FIG. 18A** and **FIG. 18B**, respectively. A graph of dual axis miRNA segregation for miR-331-3p and miR-1537-3p showing relative levels of each miRNA species based on survival to 18 months is provided in **FIG. 19**. For both miRNA species, subjects that survived at least 18 months had higher levels of a given miRNA species than those that were deceased. Table 3 provide the metrics and outcomes for the dual axis miRNA segregation for miR-331-3p and miR-1537-3p

Table 3.

Metric	Outcome
Accuracy	96.43%
Negative Predictive Value	93.33%
Positive Predictive Value	100.00%
Sensitivity	92.86%
Specificity	100.00%

Transcriptome Analysis

RNA Sequencing

RNA Sequencing was performed on tumor samples from Cohort A collected in the phase 1/2 trial described in Example 1, above, to evaluate gene expression levels. Formalin-fixed, paraffin-embedded (FFPE) tissue taken during initial resection of the primary tumor and second/third resection upon suspected progression was processed for RNA Sequencing. Briefly, one section was H&E stained and annotated by a pathologist for tumor content. H&E annotations were transferred to adjacent sections and macro-dissection was performed where possible to remove surrounding non-tumor tissue. Dual isolation of nucleic acids was performed utilizing the AllPrep DNA/RNA FFPE Tissue kit (Qiagen), followed by quality control (QC) of extracted nucleic acids using Agilent Tape Station for size distribution assessment and fluorometric method for quantification. RNA Sequencing libraries were prepared using the Stranded RNA Sequencing kit (Kapa Biosystems) and followed by QC metrics. RNA Sequencing was performed on NovaSeq™ 6000 (Illumina, San Diego, CA, USA) sequencers with paired-end 2 x 150 bp read lengths and Illumina's proprietary reversible terminator-based method. Tumor specimens were

- 77 -

sequenced to an average output of 100 million paired end reads (total of 200 M reads). Reads that mapped to the preferred transcript of each gene were counted to measure expression levels for genes from each tumor sample. Raw strand-specific counts per gene were generated by the STAR aligner. Counts per Million mapped reads (CPM) were calculated and globally normalized across samples using trimmed mean of M values (TMM). Transcriptome analysis was performed.

Transcriptome Analysis- Primary tumor samples

Patients in Cohort A (MGMT Unmethylated) were divided into two groups, those who were alive (n=15) or deceased (n=16) at the OS18 assessment. Gene transcripts were compared on a group basis and statistically significant differences in gene expression were assessed using the Wilcoxon rank sum (Mann-Whitney U) test. Differentially expressed genes (DEGs) were interrogated for function and relevance and a gene expression signature that may be associated with survival at OS18 in response to treatment with INO-5401 and INO-9012, in combination with cemiplimab, was identified. Patients who were alive at OS18 displayed reduced expression of *SYNGR3*, *OTX1*, *GABBR2*, *LHX1*, *CADM3*, *MLLT11*, *MNX1*, *GRB14*, *SLC34A2*, *PHYHIP*, *WNT10B*, *SLC17A6*, *CRLF1* and *HOXD13*, and enhanced expression of *TGF β R3*, *UBA7*, *SFRP4*, in comparison to patients who were deceased at OS18 (**FIG. 20**).

Transcriptome Analysis- Paired tumor samples

Paired samples (primary and progression tissue) were analyzed from six patients in Cohort A (MGMT Unmethylated) and gene transcripts were compared on a per subject basis. Gene expression was assessed for tumor antigen targets and markers for immune cell modulators and subsets (**FIG. 21**).

Six patients had 'treatment-related changes' noted in pathology reports for their second/third resections. Each patient displayed altered gene expression of at least two tumor antigen targets encoded by INO-5401 in progression vs. primary tumor tissue. Extent of treatment-related changes (extensive to minimal) noted in pathology reports from second/third resections correlated with changes in expression levels of immune-related genes, especially those for T cell markers, in progression vs. primary tumor tissue: Three patients (60010003, 60010006, 60010010) displayed overall

- 78 -

enhanced expression, one patient (60010009) displayed mildly enhanced or relatively unchanged expression, and two patients (60010005, 60010008) displayed overall reduced expression of immune cell modulators and subsets in progression vs. primary tumor tissue.

It is understood that the foregoing detailed description and accompanying examples are merely illustrative and are not to be taken as limitations upon the scope of the invention, which is defined solely by the appended claims and their equivalents.

Various changes and modifications to the disclosed embodiments will be apparent to those skilled in the art. Such changes and modifications, including without limitation those relating to the chemical structures, substituents, derivatives, intermediates, syntheses, compositions, formulations, or methods of use of the invention, may be made without departing from the spirit and scope thereof.

- 79 -

SEQUENCE LISTING

SEQ ID NO: 1 R2810 HCVR

Glu ₁	Val ₁	Gln ₁	Leu ₅	Leu ₅	Glu ₅	Ser ₅	Gly ₅	Gly ₁₀	Val ₁₀	Leu ₁₀	Val ₁₀	Gln ₁₅	Pro ₁₅	Gly ₁₅	Gly ₁₅
Ser ₂₀	Leu ₂₀	Arg ₂₀	Leu ₂₀	Ser ₂₀	Cys ₂₀	Ala ₂₀	Ala ₂₀	Ser ₂₅	Gly ₂₅	Phen ₂₅	Thr ₂₅	Phen ₃₀	Ser ₃₀	Asn ₃₀	Phen ₃₀
Gly ₃₅	Met ₃₅	Thr ₃₅	Trp ₃₅	Val ₃₅	Arg ₃₅	Gln ₄₀	Ala ₄₀	Pro ₄₀	Gly ₄₀	Lys ₄₀	Gly ₄₀	Leu ₄₅	Glu ₄₅	Trp ₄₅	Val ₄₅
Ser ₅₀	Gly ₅₀	Ile ₅₀	Ser ₅₀	Gly ₅₀	Gly ₅₀	Gly ₅₅	Arg ₅₅	Asp ₅₅	Thr ₅₅	Tyr ₅₅	Phen ₆₀	Ala ₆₀	Asp ₆₀	Ser ₆₀	Val ₆₀
Lys ₆₅	Gly ₆₅	Arg ₆₅	Phen ₆₅	Thr ₆₅	Ile ₇₀	Ser ₇₀	Arg ₇₀	Asp ₇₀	Asn ₇₀	Ser ₇₅	Lys ₇₅	Asn ₇₅	Thr ₇₅	Leu ₇₅	Tyr ₈₀
Leu ₈₅	Glu ₈₅	Met ₈₅	Asn ₈₅	Ser ₈₅	Leu ₈₅	Lys ₈₅	Gly ₈₅	Glu ₉₀	Asp ₉₀	Thr ₉₀	Ala ₉₀	Val ₉₀	Tyr ₉₅	Tyr ₉₅	Cys ₉₅
Val ₁₀₀	Lys ₁₀₀	Trp ₁₀₀	Gly ₁₀₀	Asn ₁₀₀	Ile ₁₀₀	Tyr ₁₀₀	Phen ₁₀₀	Asp ₁₀₅	Tyr ₁₀₅	Trp ₁₀₅	Gly ₁₀₅	Glu ₁₁₀	Gly ₁₁₀	Thr ₁₁₀	Leu ₁₁₀
Val ₁₁₅	Thr ₁₁₅	Val ₁₁₅	Ser ₁₁₅	Ser ₁₁₅											

SEQ ID NO: 2 R2810 LCVR

Asp ₁	Ile ₁	Gln ₁	Met ₅	Thr ₅	Gln ₅	Ser ₅	Pro ₅	Ser ₁₀	Ser ₁₀	Leu ₁₀	Ser ₁₀	Ala ₁₀	Ser ₁₅	Val ₁₅	Gly ₁₅
Asp ₂₀	Ser ₂₀	Ile ₂₀	Thr ₂₀	Ile ₂₀	Thr ₂₀	Cys ₂₀	Arg ₂₀	Ala ₂₅	Ser ₂₅	Leu ₂₅	Ser ₂₅	Ile ₃₀	Asn ₃₀	Thr ₃₀	Phen ₃₀
Leu ₃₅	Asn ₃₅	Trp ₃₅	Tyr ₃₅	Gln ₃₅	Gln ₃₅	Lys ₄₀	Pro ₄₀	Gly ₄₀	Lys ₄₀	Ala ₄₅	Pro ₄₅	Asn ₄₅	Leu ₄₅	Leu ₄₅	Ile ₄₅
Tyr ₅₀	Ala ₅₀	Ala ₅₀	Ser ₅₀	Ser ₅₀	Leu ₅₀	His ₅₅	Gly ₅₅	Gly ₅₅	Val ₅₅	Pro ₆₀	Ser ₆₀	Arg ₆₀	Phen ₆₀	Ser ₆₀	Gly ₆₀

- 80 -

Se	Gl	Se	Gl	Th	As	Ph	Th	Le	Th	Il	Ar	Th	Le	Gl	Pr
r	y	r	y	r	p	e	r	u	r	e	g	r	u	n	o
65					7					7					8
					0					5					0
Gl	As	Ph	Al	Th	Ty	Ty	Cy	Gl	Gl	Se	Se	As	Th	Pr	Ph
u	p	e	a	r	r	r	s	n	n	r	r	n	r	o	e
				85					90					95	
Th	Ph	Gl	Pr	Gl	Th	Va	Va	As	Ph	Ar					
r	e	y	o	y	r	l	l	p	e	g					
			10					10							
			0					5							

SEQ ID NO: 3 R2810 HCDR1

Gly Phe Thr Phe Ser Asn
 Phe Gly 1 5

SEQ ID NO: 4 R2810 HCDR2

Ile Ser Gly Gly Gly Arg
 Asp Thr 1 5

SEQ ID NO: 5 R2810 HCDR3

Val Lys Trp Gly Asn Ile Tyr Phe
 Asp Tyr 1 5 10

- 81 -

SEQ ID NO: 6 R2810 LCDR1

Leu Ser Ile Asn
 Thr Phe 1 5

SEQ ID NO: 7 R2810 LCDR2

Ala Ala
 Ser 1

SEQ ID NO: 8 R2810 LCDR3

Gln Gln Ser Ser Asn Thr Pro
 Phe Thr 1 5

SEQ ID NO: 9 R2810 HC

Glu	Val	Gln	Leu	Leu	Glu	Ser	Glu	Glu	Val	Leu	Val	Gln	Pro	Glu	Glu
1	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Glu	Phen	Thr	Phen	Ser	Asn	Phen
16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
Glu	Meth	Thr	Trp	Val	Arg	Glu	Ala	Pro	Glu	Lys	Glu	Leu	Glu	Trp	Val
32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47
Ser	Glu	Ile	Ser	Glu	Glu	Glu	Arg	Asp	Thr	Tyr	Phen	Ala	Asp	Ser	Val
48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63
Lys	Glu	Arg	Phen	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79
Leu	Glu	Meth	Asn	Ser	Leu	Lys	Glu	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95
Val	Lys	Trp	Glu	Asn	Ile	Tyr	Phen	Asp	Tyr	Trp	Glu	Glu	Glu	Thr	Leu
96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Glu	Pro	Ser	Val	Phen	Pro	Leu
112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127

- 83 -

Va₁₃₀₅ Le_u Hi_s Gl_n As_p Tr_{p310} Le_u As_n Gl_y Ly_s Gl_{u315} Ty_r Ly_s Cy_s Ly_s Va₁₃₂₀
 Se_r As_n Ly_s Gl_y Le_{u325} Pr_o Se_r Se_r Il_e Gl_{u330} Ly_s Th_r Il_e Se_r Ly_{s335} Al_a
 Ly_s Gl_y Gl_n Pr_{o340} Ar_g Gl_u Pr_o Gl_n Va₁₃₄₅ Ty_r Th_r Le_u Pr_o Pr_{o350} Se_r Gl_n
 Gl_u Gl_u Me_{t355} Th_r Ly_s As_n Gl_n Va₁₃₆₀ Se_r Le_u Th_r Cy_s Le_{u365} Va_l Ly_s Gl_y
 Ph_e Ty_{r370} Pr_o Se_r As_p Il_e Al_{a375} Va_l Gl_u Tr_p Gl_u Se_{r380} As_n Gl_y Gl_n Pr_o
 Gl_{u385} As_n As_n Ty_r Ly_s Th_{r390} Th_r Pr_o Pr_o Va_l Le_{u395} As_p Se_r As_p Gl_y Se_{r400}
 Ph_e Ph_e Le_u Ty_r Se_{r405} Ar_g Le_u Th_r Va_l As_{p410} Ly_s Se_r Ar_g Tr_p Gl_{n415} Gl_u
 Gl_y As_n Va_l Ph_{e420} Se_r Cy_s Se_r Va_l Me_{t425} Hi_s Gl_u Al_a Le_u Hi_{s430} As_n Hi_s
 Ty_r Th_r Gl_{n435} Ly_s Se_r Le_u Se_r Le_{u440} Se_r Le_u Gl_y Ly_s

SEQ ID NO: 10 R2810 LC

As_{p1} Il_e Gl_n Me_{t5} Th_r Gl_n Se_r Pr_o Se_r Se_{r10} Le_u Se_r Al_a Se_r Va_{l15} Gl_y
 As_p Se_r Il_e Th_r Il_e Th_r Cy_s Ar_g Al_{a25} Se_r Le_u Se_r Il_e As_{n30} Th_r Ph_e
 Le_u As_n Tr_{p35} Ty_r Gl_n Gl_n Ly_{s40} Pr_o Gl_y Ly_s Al_a Pr_o As_{n45} Le_u Le_u Il_e
 Ty_r Al_{a50} Al_a Se_r Se_r Le_u Hi_{s55} Gl_y Gl_y Va_l Pr_{o60} Se_r Ar_g Ph_e Se_r Gl_y
 Se_r Gl_y Se_r Gl_y Th_r As_p Ph_e Th_r Le_u Th_r Il_e Ar_g Th_r Le_u Gl_n Pr_o

- 84 -

65						70						75						80
Gl u	As p	Ph e	Al a	Th r 85	Ty r	Ty r	Cy s	Gl n	Gl n 90	Se r	Se r	As n	Th r	Pr o 95	Ph e			
Th r	Ph e	Gl y	Pr o 100	Gl y	Th r	Va l	Va l	As p 105	Ph e	Ar g	Ar g	Th r	Va l 110	Al a	Al a			
Pr o	Se r	Va l 115	Ph e	Il e	Ph e	Pr o	Pr o 120	Se r	As p	Gl u	Gl n	Le u 125	Ly s	Se r	Gl y			
Th r	Al a 130	Se r	Va l	Va l	Cy s	Le u 135	Le u	As n	As n	Ph e	Ty r 140	Pr o	Ar g	Gl u	Al a			
Ly s 145	Va l	Gl n	Tr p	Ly s	Va l 150	As p	As n	Al a	Le u	Gl n 155	Se r	Gl y	As n	Se r	Gl n 160			
Gl u	Se r	Va l	Th r	Gl u 165	Gl n	As p	Se r	Ly s	As p 170	Se r	Th r	Ty r	Se r	Le u 175	Se r			
Se r	Th r	Le u	Th r 180	Le u	Se r	Ly s	Al a	As p 185	Ty r	Gl u	Ly s	Hi s	Ly s 190	Va l	Ty r			
Al a	Cy s	Gl u 195	Va l	Th r	Hi s	Gl n	Gl y 200	Le u	Se r	Se r	Pr o	Va l 205	Th r	Ly s	Se r			
Ph e	As n 210	Ar g	Gl y	Gl u	Cy s													

- 85 -

SEQ ID NO: 11 PSMA with no IgE leader sequence and with
HA tag atgtggaacg cactgcatga gactgattct gctgtcgcac
tgggacggag accccggtgg 60 ctgtgcgctg gagcactggt
gctggccggc gggggattcc tgctgggatt cctgtttggc 120
tggtttatca aaagctccag cgaggctacc aatattaccc ctaagcacia
taagaaagca 180 ttcttgatg aactgaaagc cgagaacatc
aagaaattcc tgtacaactt cacaagaatt 240 ccacatctgg
ctggcactga gcagaacttc cagctggcaa aacagatcca gactcagtgg
300 aaggaatttg ggctggactc agtggagctg acccactacg
atgtcctgct gtcctatcca 360 aataagactc atcccaacta
catctctatc attaacgaag acggaatga gattttcaac 420
acctctctgt ttgaaccccc tccaccggc tatgagaatg tcagtgcgt
ggccctcca 480 ttctcagcct tcagcccca ggggatgcct
gaggagatc tgggtgacgt caattatgct 540 agaacagaag
acttcttta gctggagagg gatatgaaa tcaactgttc cggcaagatc
600 gtgattgccc ggtacgggaa ggtgttcaga ggaaataagg
tcaaaaacgc tcagctggcc 660 ggagctaccg gcgtgatcct
gtacagcgac cccgctgatt attttgacc tggcgtgaag 720
tcctatccag acggatggaa tctgccggc gggggagtgc agaggggaaa
catcctgaac 780 ctgaatggag ccggcgatcc tctgactcca
ggataccccc ccaacgaata cgcttatcgc 840 cggggaattg
cagaggccgt gggcctgcct agcatcccag tccatcccat tggctattac
900 gatgccaga agctgctgga gaaaatgggc gggagcgtc
cccctgactc tagttggaag 960 ggctccctga aagtgcctta
caatgtcggg ccaggattca ctgggaactt ttctaccag 1020
aaggtgaaaa tgcacatcca tagtaccagc gaggtgacac gaatctaca
cgctattggc 1080 accctgagag gcgccgtgga gcctgatcgc
tatgtcattc tgggaggcca cagagactca 1140 tgggtgttcg
ggggaatcga tccacagagc ggagcagctg tggccatga aattgtgcgc
1200 agctttggga ccctgaagaa agagggatgg cgaccaggc
gcacaatcct gttcgcatcc 1260 tgggacgccg aggaatttgg
gctgctgggc agcacagaat gggccgagga aaattctcgc 1320
ctgctgcagg agcaggggt ggcttacatc aatgcagact caagcattga
aggaaactat 1380 accctgcggg tggattgcac acccctgatg
tacagtctgg tctataacct gacaaaggag 1440 ctgaaatcac
ctgacgaggg ctgcgaaggg aaaagcctgt acgaatcctg gactgagaag
1500 agcccatccc ccgaattcag cggcatgcct aggatctcta
agctgggcag tgggaacgat 1560 tttaggtgt tctttcagcg
cctgggaatt gcctctggcc gagctcggta cacaaaaaat 1620
tgggagacta acaagttctc ctcttaccga ctgtatcaca gcgtgtacga
gacttatgaa 1680 ctggtcgaga aattctacga cccactttt
aagtatcatc tgaccgtggc acaggtcagg 1740 ggcgggatgg

- 86 -

tgttcgaact ggccaatagc atcgctcctgc catttgactg tcgagattac
 1800 gctgtggtcc tgcggaagta cgcagacaag atctataaca
 tctccatgaa gcacccccag 1860 gagatgaagg cctatttctgt
 gagtttcgat tccctgtttt ctgccgtcaa aaatttcacc 1920
 gaaatcgcta gtaagttttc agagcgcctg caggacctgg ataagtccaa
 tcccatcctg 1980 ctgcggatta tgaacgatca gctgatgttc
 ctggaaagag cttttatcga ccctctgggc 2040 ctgcctgata
 gaccattcta caggcacgtg atctacgcac ctagttcaca taacaagtac
 2100 gccggcgagt ctttcccagg gatctatgac gctctgtttg
 atattgaatc aaaggtggac 2160 ccagcaaag catggggcga
 ggtcaagaga cagatcagca ttgcagcctt tacagtgcag 2220
 gccgcccgcg aaaccctgtc cgaagtcgct taccatacag atgtccccga
 ttacgcatga 2280 taa

SEQ ID NO: 12 PSMA with IgE leader sequence (underlined)
 and with HA tag atggactgga catggattct gttcctggtc
gccgcccga ctcgctgca ttcctggaac 60 gcactgcatg agactgattc
 tgctgtcgca ctgggacgga gaccccgggtg gctgtgcgct 120
 ggagcactgg tgctggccgg cgggggattc ctgctgggat tctgttttg
 ctggtttatc 180 aaaagctcca gcgaggctac caatattacc
 cctaagcaca ataagaaagc attcctggat 240 gaactgaaag
 ccgagaacat caagaaatc ctgtacaact tcacaagaat tccacatctg
 300 gctggcactg agcagaactt ccagctggca aacagatcc
 agagtcagtg gaaggaattt 360 gggctggact cagtggagct
 gaccactac gatgtcctgc tgtcctatcc aaataagact 420
 catcccaact acatctctat cattaacgaa gacggaaatg agattttcaa
 cacctctctg 480 tttgaacccc ctccaccgg ctatgagaat
 gtcagtgcg tggtccctcc attctcagcc 540 ttcagcccc
 aggggatgcc tgaggagat ctggtgtacg tcaattatgc tagaacagaa
 600 gacttcttta agctggagag ggatatgaaa atcaactgtt
 ccggcaagat cgtgattgcc 660 cggtagcgga aggtgttcag
 aggaaataag gtcaaaaacg ctacagctggc cggagctacc 720
 ggctgatcc tgtacagcga ccccgctgat tattttgcac ctggcgtgaa
 gtcctatcca 780 gacggatgga atctgcccgg cgggggagtg
 cagaggggaa acatcctgaa cctgaatgga 840 gccggcgatc
 ctctgactcc aggatacccc gccaacgaat acgcttatcg ccggggaatt
 900 gcagaggccg tgggcctgcc tagcatccca gtccatccca
 ttggctatta cgatgccag 960 aagctgctgg agaaaatggg
 cgggagcgct cccctgact ctagttgga gggctccctg 1020

- 87 -

```

aaagtgcctt acaatgtcgg gccaggattc actgggaact tttctaccca
gaaggtgaaa 1080 atgcacatcc atagtaccag cgaggtgaca
cgaatctaca acgtcattgg caccctgaga 1140 ggcgccgtgg
agcctgatcg ctatgtcatt ctgggaggcc acagagactc atgggtgttc
1200 gggggaatcg atccacagag cggagcagct gtggtccatg
aaattgtgcg cagctttggg 1260 accctgaaga aagagggatg
gcgacccagg cgcacaatcc tgttcgcatc ctgggacgcc 1320
gaggaatttg ggctgctggg cagcacagaa tgggccgagg aaaattctcg
cctgctgcag 1380 gagcgagggg tggcttacat caatgcagac
tcaagcattg aaggaaacta taccctgcgg 1440 gtggattgca
caccctgat gtacagtctg gtctataacc tgacaaagga gctgaaatca
1500 cctgacgagg gcttcgaagg gaaaagcctg tacgaatcct
ggactgagaa gagcccatcc 1560 cccgaattca gcggcatgcc
taggatctct aagctgggca gtgggaacga ttttgagggtg 1620
ttctttcagc gcctgggaat tgctctggc cgagctcggg acacaaaaaa
ttgggagact 1680 aacaagttct cctcttacc actgtatcac
agcgtgtacg agacttatga actggtcgag 1740 aaattctacg
acccacttt taagtatcat ctgaccgtgg cacaggtcag ggcgggatg
1800 gtgttcgaac tggccaatag catcgtcctg ccatgtgact
gtcgagatta cgctgtggtc 1860 ctgcggaagt acgcagacaa
gatctataac atctccatga agcaccacca ggagatgaag 1920
gcctattctg tgagtttcga ttccctgttt tctgccgtca aaaatttcac
cgaaatcgct 1980 agtaagtttt cagagcgct gcaggacctg
gataagtcca atcccatcct gctgcggatt 2040 atgaacgatc
agctgatgtt cctggaaaga gcctttatcg accctctggg cctgcctgat
2100 agaccattct acaggcacgt gatctacgca cctagttcac
ataacaagta cgccggcgag 2160 tctttcccag ggatctatga
cgctctgttt gatattgaat caaagtgga cccagcaaa 2220
gcatggggcg aggtcaagag acagatcagc attgcagcct ttacagtgca
ggccgccgcc 2280 gaaaccctgt ccgaagtcgc ttaccatac
gatgtccccg attacgatg ataa

```

SEQ ID NO: 13 PSMA with no IgE leader sequence
Met Trp Asn Ala Leu His Glu Thr Asp Ser Ala Val Ala
Leu Gly Arg Arg Pro Arg Trp Leu Cys Ala Gly Ala Leu
Val Leu Ala Gly Gly Gly Phe Leu Leu Gly Phe Leu Phe
Gly Trp Phe Ile Lys Ser Ser Ser Glu Ala Thr Asn Ile
Thr Pro Lys His Asn Lys Lys Ala Phe Leu Asp Glu Leu
Lys Ala Glu Asn Ile Lys Lys Phe Leu Tyr Asn Phe Thr
Arg Ile Pro His Leu Ala Gly Thr Glu Gln Asn Phe Gln
Leu Ala Lys Gln Ile Gln Ser Gln Trp Lys Glu Phe Gly
Leu Asp Ser Val Glu Leu Thr His Tyr Asp Val Leu Leu

- 88 -

Ser Tyr Pro Asn Lys Thr His Pro Asn Tyr Ile Ser Ile
Ile Asn Glu Asp Gly Asn Glu Ile Phe Asn Thr Ser Leu
Phe Glu Pro Pro Pro Pro Gly Tyr Glu Asn Val Ser Asp
Val Val Pro Pro Phe Ser Ala Phe Ser Pro Gln Gly Met
Pro Glu Gly Asp Leu Val Tyr Val Asn Tyr Ala Arg Thr
Glu Asp Phe Phe Lys Leu Glu Arg Asp Met Lys Ile Asn
Cys Ser Gly Lys Ile Val Ile Ala Arg Tyr Gly Lys Val
Phe Arg Gly Asn Lys Val Lys Asn Ala Gln Leu Ala Gly
Ala Thr Gly Val Ile Leu Tyr Ser Asp Pro Ala Asp Tyr
Phe Ala Pro Gly Val Lys Ser Tyr Pro Asp Gly Trp Asn
Leu Pro Gly Gly Gly Val Gln Arg Gly Asn Ile Leu Asn
Leu Asn Gly Ala Gly Asp Pro Leu Thr Pro Gly Tyr Pro
Ala Asn Glu Tyr Ala Tyr Arg Arg Gly Ile Ala Glu Ala
Val Gly Leu Pro Ser Ile Pro Val His Pro Ile Gly Tyr
Tyr Asp Ala Gln Lys Leu Leu Glu Lys Met Gly Gly Ser
Ala Pro Pro Asp Ser Ser Trp Lys Gly Ser Leu Lys Val
Pro Tyr Asn Val Gly Pro Gly Phe Thr Gly Asn Phe Ser
Thr Gln Lys Val Lys Met His Ile His Ser Thr Ser Glu
Val Thr Arg Ile Tyr Asn Val Ile Gly Thr Leu Arg Gly
Ala Val Glu Pro Asp Arg Tyr Val Ile Leu Gly Gly His
Arg Asp Ser Trp Val Phe Gly Gly Ile Asp Pro Gln Ser
Gly Ala Ala Val Val His Glu Ile Val Arg Ser Phe Gly
Thr Leu Lys Lys Glu Gly Trp Arg Pro Arg Arg Thr Ile
Leu Phe Ala Ser Trp Asp Ala Glu Glu Phe Gly Leu Leu
Gly Ser Thr Glu Trp Ala Glu Glu Asn Ser Arg Leu Leu
Gln Glu Arg Gly Val Ala Tyr Ile Asn Ala Asp Ser Ser
Ile Glu Gly Asn Tyr Thr Leu Arg Val Asp Cys Thr Pro
Leu Met Tyr Ser Leu Val Tyr Asn Leu Thr Lys Glu Leu
Lys Ser Pro Asp Glu Gly Phe Glu Gly Lys Ser Leu Tyr
Glu Ser Trp Thr Glu Lys Ser Pro Ser Pro Glu Phe Ser
Gly Met Pro Arg Ile Ser Lys Leu Gly Ser Gly Asn Asp
Phe Glu Val Phe Phe Gln Arg Leu Gly Ile Ala Ser Gly
Arg Ala Arg Tyr Thr Lys Asn Trp Glu Thr Asn

- 89 -

Lys Phe Ser Ser Tyr Pro Leu Tyr His Ser Val Tyr Glu
 Thr Tyr Glu Leu Val Glu Lys Phe Tyr Asp Pro Thr Phe
 Lys Tyr His Leu Thr Val Ala Gln Val Arg Gly Gly Met
 Val Phe Glu Leu Ala Asn Ser Ile Val Leu Pro Phe Asp
 Cys Arg Asp Tyr Ala Val Val Leu Arg Lys Tyr Ala Asp
 Lys Ile Tyr Asn Ile Ser Met Lys His Pro Gln Glu Met
 Lys Ala Tyr Ser Val Ser Phe Asp Ser Leu Phe Ser Ala
 Val Lys Asn Phe Thr Glu Ile Ala Ser Lys Phe Ser Glu
 Arg Leu Gln Asp Leu Asp Lys Ser Asn Pro Ile Leu Leu
 Arg Ile Met Asn Asp Gln Leu Met Phe Leu Glu Arg Ala
 Phe Ile Asp Pro Leu Gly Leu Pro Asp Arg Pro Phe Tyr
 Arg His Val Ile Tyr Ala Pro Ser Ser His Asn Lys Tyr
 Ala Gly Glu Ser Phe Pro Gly Ile Tyr Asp Ala Leu Phe
 Asp Ile Glu Ser Lys Val Asp Pro Ser Lys Ala Trp Gly
 Glu Val Lys Arg Gln Ile Ser Ile Ala Ala Phe Thr Val
 Gln Ala Ala Ala Glu Thr Leu Ser Glu Val Ala

SEQ ID NO: 14 PSMA with IgE leader sequence (underlined)

Met Asp Trp Thr Trp Ile Leu Phe Leu Val Ala Ala Ala
Thr Arg Val His Ser Trp Asn Ala Leu His Glu Thr Asp
 Ser Ala Val Ala Leu Gly Arg Arg Pro Arg Trp Leu Cys
 Ala Gly Ala Leu Val Leu Ala Gly Gly Gly Phe Leu Leu
 Gly Phe Leu Phe Gly Trp Phe Ile Lys Ser Ser Ser Glu
 Ala Thr Asn Ile Thr Pro Lys His Asn Lys Lys Ala Phe
 Leu Asp Glu Leu Lys Ala Glu Asn Ile Lys Lys Phe Leu
 Tyr Asn Phe Thr Arg Ile Pro His Leu Ala Gly Thr Glu
 Gln Asn Phe Gln Leu Ala Lys Gln Ile Gln Ser Gln Trp
 Lys Glu Phe Gly Leu Asp Ser Val Glu Leu Thr His Tyr
 Asp Val Leu Leu Ser Tyr Pro Asn Lys Thr His Pro Asn
 Tyr Ile Ser Ile Ile Asn Glu Asp Gly Asn Glu Ile Phe
 Asn Thr Ser Leu Phe Glu Pro Pro Pro Gly Tyr Glu
 Asn Val Ser Asp Val Val Pro Pro Phe Ser Ala Phe Ser
 Pro Gln Gly Met Pro Glu Gly Asp Leu Val Tyr Val Asn
 Tyr Ala Arg Thr Glu Asp Phe Phe Lys Leu Glu Arg Asp
 Met Lys Ile Asn Cys Ser Gly Lys Ile Val Ile Ala Arg
 Tyr Gly Lys Val Phe Arg Gly Asn Lys Val Lys Asn Ala
 Gln Leu Ala Gly Ala Thr Gly Val Ile Leu Tyr Ser Asp
 Pro Ala Asp Tyr Phe Ala Pro Gly Val Lys Ser Tyr Pro
 Asp Gly Trp Asn Leu Pro Gly Gly Gly Val Gln Arg Gly
 Asn Ile Leu Asn Leu Asn Gly Ala Gly Asp Pro Leu Thr
 Pro Gly Tyr Pro Ala Asn Glu Tyr Ala Tyr Arg Arg Gly
 Ile Ala Glu Ala Val Gly Leu Pro Ser Ile Pro Val His
 Pro Ile Gly Tyr Tyr Asp Ala Gln Lys Leu Leu Glu Lys

- 90 -

Met Gly Gly Ser Ala Pro Pro Asp Ser Ser Trp Lys Gly
Ser Leu Lys Val Pro Tyr Asn Val Gly Pro Gly Phe Thr
Gly Asn Phe Ser Thr Gln Lys Val Lys Met His Ile His
Ser Thr Ser Glu Val Thr Arg Ile Tyr Asn Val Ile Gly
Thr Leu Arg Gly Ala Val Glu Pro Asp Arg Tyr Val Ile
Leu Gly Gly His Arg Asp Ser Trp Val Phe Gly Gly Ile
Asp Pro Gln Ser Gly Ala Ala Val Val His Glu Ile Val
Arg Ser Phe Gly Thr Leu Lys Lys Glu Gly Trp Arg Pro
Arg Arg Thr Ile Leu Phe Ala Ser Trp Asp Ala Glu Glu
Phe Gly Leu Leu Gly Ser Thr Glu Trp Ala Glu Glu Asn
Ser Arg Leu Leu Gln Glu Arg Gly Val Ala Tyr Ile Asn
Ala Asp Ser Ser Ile Glu Gly Asn Tyr Thr Leu Arg Val
Asp Cys Thr Pro Leu Met Tyr Ser Leu Val Tyr Asn Leu
Thr Lys Glu Leu Lys Ser Pro Asp Glu Gly Phe Glu Gly
Lys Ser Leu Tyr Glu Ser Trp Thr Glu Lys Ser Pro Ser
Pro Glu Phe Ser Gly Met Pro Arg Ile Ser Lys Leu Gly
Ser Gly Asn Asp Phe Glu Val Phe Phe Gln Arg Leu Gly
Ile Ala Ser Gly Arg Ala Arg Tyr Thr Lys Asn Trp Glu
Thr Asn Lys Phe Ser Ser Tyr Pro Leu Tyr His Ser Val
Tyr Glu Thr Tyr Glu Leu Val Glu Lys Phe Tyr Asp Pro
Thr Phe Lys Tyr His Leu Thr Val Ala Gln Val Arg Gly
Gly Met Val Phe Glu Leu Ala Asn Ser Ile Val Leu Pro
Phe Asp Cys Arg Asp Tyr Ala Val Val Leu Arg Lys Tyr
Ala Asp Lys Ile Tyr Asn Ile Ser Met Lys His Pro Gln
Glu Met Lys Ala Tyr Ser Val Ser Phe Asp Ser Leu Phe
Ser Ala Val Lys Asn Phe Thr Glu Ile Ala Ser Lys Phe
Ser Glu Arg Leu Gln Asp Leu Asp Lys Ser Asn Pro Ile
Leu Leu Arg Ile Met Asn Asp Gln Leu Met Phe Leu Glu
Arg Ala Phe Ile Asp Pro Leu Gly Leu Pro Asp Arg Pro
Phe Tyr

- 91 -

Arg His Val Ile Tyr Ala Pro Ser Ser His Asn Lys Tyr
 Ala Gly Glu Ser Phe Pro Gly Ile Tyr Asp Ala Leu Phe
 Asp Ile Glu Ser Lys Val Asp Pro Ser Lys Ala Trp Gly
 Glu Val Lys Arg Gln Ile Ser Ile Ala Ala Phe Thr Val
 Gln Ala Ala Ala Glu Thr Leu Ser Glu Val Ala

SEQ ID NO: 15 Consensus WT1-L with modified Zinc
 Fingers nucleic acid sequence plus IgE leader
ggatccgccca ccatggactg gacctggatt ctgttcctgg tcgccgccgc
aacacgggtg 60 catagtggga gtgatgtgag agacctgaac
 gccctgctgc cagcagtgcc atccctgcct 120 ggcgggggag
 gctgcgctct gccagtctct ggagcagctc agtgggctcc cgtgctggac
 180 tttgcacccc ctgcagcccc ttacggaagt ctgggcggcc
 cacactcatt catcaaacag 240 gagccaagct ggggcggggc
 agatcctcat gaggaacagt gcctgtcagc cttcacagtc 300
 cacttttagcg ggcagttcac tggaaccgca ggagcttgta gatacggacc
 ctttgagca 360 ccaccccctt cccaggcacc ttctggacag
 gcacgcatgt tcccaaacgc tccctatctg 420 cctaattgtc
 tggaaagcca gcccgctatt aggaaccagg gctactccac agtggcattt
 480 gacgggactc ctgctatgg acatacccca tcccaccatg
 ctgcacagtt tcctaatacac 540 tccttcaagc atgaggaccc
 catgggacag caggggtccc tgggagaaca gcagtactct 600
 gtgccccctc ccgtgtacgg atgccacaca ccaactgaca gttgtacagg
 ctacagggcc 660 ctgctgctgc gaactccata caacagtgat
 aatctgtatc agatgacctc acagctggag 720 tgcattgacat
 ggaaccagat gaatctgggc agcacactga aaggccatgc cactgggtac
 780 gaatctgaca accacaccac acctatgctg tacagttgtg
 gagccagta tagaatccac 840 actcatggag tcttcagagg
 cattcaggat gtgcggagag tcccaggagt ggcaccaact 900
 atcgtgcgga gcgcctccga gaccaacgaa aagcgcccct ttatgggcgc
 ctaccctgga 960 ggcaataagc ggtatttcaa actgtctcac
 ctgcagatgg ggagtagaaa ggggaccgga 1020 gagaaacctt
 atcagggcga ctttaaagat ggggaaaggc gcttctctcg cagtgaccag
 1080 ctgaagcgag gacagcgacg aggaaccggg gtgaagccat
 ttcagtgcaa aacatgtcag 1140 agaaagttct caaggagcga
 tcacctgaag acccatacaa gaactcacac cggcaagacc 1200
 agcgagaaac cattttcctg ccgatggccc tcttgtcaga agaaattcgc
 ccgctccgac 1260 gaactgggtcc gacaccacaa tatgcatcag
 agaaatatga caaaactgca gctggctctg 1320 tgataactcg ag

SEQ ID NO: 16 Con WT1-L with modified Zinc Fingers
 protein sequence plus IgE leader (underlined)

- 92 -

Met Asp Trp Thr Trp Ile Leu Phe Leu Val Ala Ala Ala
Thr Arg Val His Ser Gly Ser Asp Val Arg Asp Leu Asn
Ala Leu Leu Pro Ala Val Pro Ser Leu Pro Gly Gly Gly
Gly Cys Ala Leu Pro Val Ser Gly Ala Ala Gln Trp Ala
Pro Val Leu Asp Phe Ala Pro Pro Ala Ala Pro Tyr Gly
Ser Leu Gly Gly Pro His Ser Phe Ile Lys Gln Glu Pro
Ser Trp Gly Gly Ala Asp Pro His Glu Glu Gln Cys Leu
Ser Ala Phe Thr Val His Phe Ser Gly Gln Phe Thr Gly
Thr Ala Gly Ala Cys Arg Tyr Gly Pro Phe Gly Ala Pro
Pro Pro Ser Gln Ala Pro Ser Gly Gln Ala Arg Met Phe
Pro Asn Ala Pro Tyr Leu Pro Asn Cys Leu Glu Ser Gln
Pro Ala Ile Arg Asn Gln Gly Tyr Ser Thr Val Ala Phe
Asp Gly Thr Pro Ser Tyr Gly His Thr Pro Ser His His
Ala Ala Gln Phe Pro Asn His Ser Phe Lys His Glu Asp
Pro Met Gly Gln Gln Gly Ser Leu Gly Glu Gln Gln Tyr
Ser Val Pro Pro Pro Val Tyr Gly Cys His Thr Pro Thr
Asp Ser Cys Thr Gly Ser Gln Ala Leu Leu Leu Arg Thr
Pro Tyr Asn Ser Asp Asn Leu Tyr Gln Met Thr Ser Gln
Leu Glu Cys Met Thr Trp Asn Gln Met Asn Leu Gly Ser
Thr Leu Lys Gly His Ala Thr Gly Tyr Glu Ser Asp Asn
His Thr Thr Pro Met Leu Tyr Ser Cys Gly Ala Gln Tyr
Arg Ile His Thr His Gly Val Phe Arg Gly Ile Gln Asp
Val Arg Arg Val Pro Gly Val Ala Pro Thr Ile Val Arg
Ser Ala Ser Glu Thr Asn Glu Lys Arg Pro Phe Met Gly
Ala Tyr Pro Gly Gly Asn Lys Arg Tyr Phe Lys Leu Ser
His Leu Gln Met Gly Ser Arg Lys Gly Thr Gly Glu Lys
Pro Tyr Gln Gly Asp Phe Lys Asp Gly Glu Arg Arg Phe
Ser Arg Ser Asp Gln Leu Lys Arg Gly Gln Arg Arg Gly
Thr Gly Val Lys Pro Phe Gln Cys Lys Thr Cys Gln Arg
Lys Phe Ser Arg Ser Asp His Leu Lys Thr His Thr Arg
Thr His Thr Gly Lys Thr Ser Glu Lys Pro Phe Ser Cys
Arg Trp Pro Ser Cys Gln Lys Lys Phe Ala Arg Ser Asp

- 93 -

Glu Leu Val Arg His His Asn Met His Gln Arg Asn
Met Thr Lys Leu Gln Leu Ala Leu

SEQ ID NO: 17 Synthetic Consensus hTERT nucleic acid
sequence operably linked to IgE (underlined)
atggattgga catggattct gttcctgggtc gcagccgcca cagagtgca
tagccctaga 60 gcccacgggt gtagagcagt ccgcagcctg
ctgcgcagcc gataccggga agtgctgcct 120 ctggccacct
ttgtccggag actgggacca cagggcaggc gcctgggtgca gcgcggcgac
180 cccgcagctt tccgagcact ggtggcacag tgcctgggtg
gcgtgccatg ggatgcacgg 240 cccctccag cagcccctag
ctttagacag gtgtcctgcc tgaaagaact ggtcgcaagg 300
gtgggtccagc ggctgtgcga gagaggcgcc aggaacgtgc tggcattcgg
ctttgcaactg 360 ctggacggag ctaggggcgg gcccctgag
gcattcacca caagcgtgcg ctccctacctg 420 ccaaatacag
tcaactgatac cctgcgaggg tccggagcat ggggactgct gctgcgacgg
480 gtgggggacg atgtgctggt ccacctgctg gctagatgcg
cactgtatgt gctggctcgt 540 ccctcttgcg cataccaggt
gtgcggacca cccctgtatg acctgggcgc tgcaaccag 600
gcaagacctc caccacgc ctctggcact agaaggggac tgggcaccga
acaggcatgg 660 aaccatagtg tcagggaggc aggagtcca
ctgggactgc cagcacctgg ggctcgccga 720 cggagaggga
gtgccggacg gtcactgcca ctggctaaga gaccaaggcg cggagccgct
780 ccagaaccag agaggacacc tgtgggacag ggaagctggg
cacaccctgg aagaactagg 840 gggccaagt ataggggctt
ctgcgtgggtc tcaccagcac gaccagcaga ggaagctact 900
tctctggagg gagctctgag tggcaccgg cactctcatc ctagtgtggg
aagacagcac 960 catgcaggcc ctccaagcac cagccggcct
ccccggccat gggacactcc ttgtccaccc 1020 gtgtacgctg
aaaccaaaca ctttctgtat agctccggag ataaggagca gctgcggccc
1080 tctttcctgc tgtctagtct gagacctagt ctgaccggag
cacgacggct ggtggaaaca 1140 atctttctgg ggtcccggcc
ttggatgcca ggaaccccca gaaggacacc tcgactgcca 1200
cagcgggtact ggcagatgcg gccactgttc ctggagctgc tgggcaatca
cgctcagtgc 1260 ccctatgggg cactgctgcg aacacattgt
cctctgcggg cagccgtgac tccagctgca 1320 ggagtctgcg
ccagggaaaa gccacagggc agcgtggcag ctccctgagga agaggacacc
1380 gatccacgcc gactggtgca gctgctgaga cagcactcaa
gcccctggca ggtgtacgga 1440 tttctgaggg cctgtctgcg
gagactggtg cctccaggac tgtggggggtc caggcacaac 1500
gaaaggcgct ttctgcgcaa tactaagaaa ttcacagcc tgggcaagca
tgctaaactg 1560 tccctgcagg agctgacctg gaaaatgagt

- 94 -

gtgcgcgact gcgcatggct gcgacgggtca 1620 ccaggagtcg
ggtgcgtgcc tgcagccgag caccgcctgc gagaagagat tctggccaag
1680 tttctgcatt ggctgatgtc agtgtacgtg gtcgaactgc
tgccggagctt cttttatgtg 1740 acagagacta cttccagaa
aaactacctg ttcttttatt gcaagtcagt gtggagcaaa 1800
ctgcagtcaa tcggcattcg gcagcacctg aagagagtgc agctgaggga
actgagtga 1860 gccgagggtcc ggcagcatag agaggcaagg
cctgccctgc tgacctcccg gctgagattc 1920 ctgcctaagc
cagacgggct gagaccaatc gtgaacatgg attacgtggc cggagcacgg
1980 accttccgga gggaaaaacg cgctgagcga ctgacatccc
gcgtgaagac tctgttctct 2040 gtcctgaatt atgagcgcgc
tcgccgaccc ggactgctgg gagcatctgt gctgggactg 2100
gacgatattc accgggcttg gagagcattt gtcctgaggg tgcgcgcaca
ggaccctccc 2160 ccagaactgt acttcgtgaa agtcgccgtg
accggggctt atgacacaat ccctcaggat 2220 cggctgactg
aagtgatcgc ctccatcatt aagccacaga atacctactg cgtgcggaga
2280 tatgctgtgg tcaggcgcgc tgcacacggc catgtgagga
agagcttcaa gcgccacgtc 2340 agcacactga ctgatctgca
gccctacatg agacagttcg tggtcatctc gcaggagacc 2400
agccctctga gggacgcagt ggtcatcgaa cagtcctcta gtcgaacga
ggcatcaagc 2460 gggtgttcg atgtctttct gcggttcgtg
tgccaccatg ccgtcagaat tggaggcaaa 2520 tcttacgtgc
agtgtcaggg catccccag ggcagcattc tgtctaccct gctgtgcagc
2580 ctgtgctatg gcgacatgga aaataagctg tttgccggaa
tccgacggga tggcctgctg 2640 ctgagactgg tggccgcttt
tctgctggtc actccacacc tgacctatgc caaagctttc 2700
ctgcgcacac tggtcgagg ggtgccagag tacggatgcg tggtaacct
gaggaagacc 2760 gtggtcaatt tcccagtgga agacgaggcc
ctgggcggca cagcatttgt ccagctgcca 2820 gcacacggac
tgttcccatg gtgtggactg ctgctggaca cccgcacact ggaggtgcag
2880 tccgattact cctcttatgc ccggacaagc atcagagctt
ccctgacttt taacagaggc 2940 ttcaaggccg ggaggaatat
gagaaggaaa ctgtttggcg tgctgcgcct gaagtgccat 3000
tccctgttcc tgtatctgca ggtgaactct ctgcagactg tctgtaccaa
cgtgtacaaa 3060 atttttctgc tgcaggccta tcggttccac
gcttgctgctg tgcagctgcc attccatcag 3120 caggtcagga
agaacccac cttctttctg cgcgtgatct ctgatacagc tagtctgtgc
3180 tactcaattc tgaaggccaa aaatgctggc atgagcctgg
gagcaaaagg agcagcagga 3240

- 95 -

ccatttcctt ccgaggctgc acagtggctg tgccaccagg cattcctgct
 gaagctggcc 3300 cgacatcggt tgacatatag gtgcctgctg
 ggcgcactgc gaacagcaca gactcagctg 3360 tgcagaaagc
 tgcccggggc cactctgggt gccctggaag ccgctgccga ccctgccctg
 3420 acctccgatt tcaagactat tctggac

SEQ ID NO: 18 Synthetic consensus hTERT amino acid sequence
 operably linked to IgE (underlined)

Met Asp Trp Thr Trp Ile Leu Phe Leu Val Ala Ala Ala
Thr Arg Val His Ser Pro Arg Ala Pro Arg Cys Arg Ala
 Val Arg Ser Leu Leu Arg Ser Arg Tyr Arg Glu Val Leu
 Pro Leu Ala Thr Phe Val Arg Arg Leu Gly Pro Gln Gly
 Arg Arg Leu Val Gln Arg Gly Asp Pro Ala Ala Phe Arg
 Ala Leu Val Ala Gln Cys Leu Val Cys Val Pro Trp Asp
 Ala Arg Pro Pro Pro Ala Ala Pro Ser Phe Arg Gln Val
 Ser Cys Leu Lys Glu Leu Val Ala Arg Val Val Gln Arg
 Leu Cys Glu Arg Gly Ala Arg Asn Val Leu Ala Phe Gly
 Phe Ala Leu Leu Asp Gly Ala Arg Gly Gly Pro Pro Glu
 Ala Phe Thr Thr Ser Val Arg Ser Tyr Leu Pro Asn Thr
 Val Thr Asp Thr Leu Arg Gly Ser Gly Ala Trp Gly Leu
 Leu Leu Arg Arg Val Gly Asp Asp Val Leu Val His Leu
 Leu Ala Arg Cys Ala Leu Tyr Val Leu Val Ala Pro Ser
 Cys Ala Tyr Gln Val Cys Gly Pro Pro Leu Tyr Asp Leu
 Gly Ala Ala Thr Gln Ala Arg Pro Pro Pro His Ala Ser
 Gly Thr Arg Arg Gly Leu Gly Thr Glu Gln Ala Trp Asn
 His Ser Val Arg Glu Ala Gly Val Pro Leu Gly Leu Pro
 Ala Pro Gly Ala Arg Arg Arg Arg Gly Ser Ala Gly Arg
 Ser Leu Pro Leu Ala Lys Arg Pro Arg Arg Gly Ala Ala
 Pro Glu Pro Glu Arg Thr Pro Val Gly Gln Gly Ser Trp
 Ala His Pro Gly Arg Thr Arg Gly Pro Ser Asp Arg Gly
 Phe Cys Val Val Ser Pro Ala Arg Pro Ala Glu Glu Ala
 Thr Ser Leu Glu Gly Ala Leu Ser Gly Thr Arg His Ser
 His Pro Ser Val Gly Arg Gln His His Ala Gly Pro Pro
 Ser Thr Ser Arg Pro Pro Arg Pro Trp Asp Thr Pro Cys
 Pro Pro Val Tyr Ala Glu Thr Lys His Phe Leu Tyr Ser
 Ser Gly Asp Lys Glu Gln Leu Arg Pro Ser Phe Leu Leu
 Ser Ser Leu Arg Pro Ser Leu Thr Gly Ala Arg Arg Leu
 Val Glu Thr Ile Phe Leu Gly Ser Arg Pro Trp Met Pro
 Gly Thr Pro Arg Arg Thr Pro Arg Leu Pro Gln Arg Tyr
 Trp Gln Met Arg Pro Leu Phe Leu Glu Leu Leu Gly Asn
 His Ala Gln Cys Pro Tyr Gly Ala Leu Leu Arg Thr His
 Cys Pro Leu Arg Ala Ala Val Thr Pro Ala Ala Gly Val

- 96 -

Cys Ala Arg Glu Lys Pro Gln Gly Ser Val Ala Ala Pro
Glu Glu Glu Asp Thr Asp Pro Arg Arg Leu Val Gln Leu
Leu Arg Gln His Ser Ser Pro Trp Gln Val Tyr Gly Phe
Leu Arg Ala Cys Leu Arg Arg Leu Val Pro Pro Gly Leu
Trp Gly Ser Arg His Asn Glu Arg Arg Phe Leu Arg Asn
Thr Lys Lys Phe Ile Ser Leu Gly Lys His Ala Lys Leu
Ser Leu Gln Glu Leu Thr Trp Lys Met Ser Val Arg Asp
Cys Ala Trp Leu Arg Arg Ser Pro Gly Val Gly Cys Val
Pro Ala Ala Glu His Arg Leu Arg Glu Glu Ile Leu Ala
Lys Phe Leu His Trp Leu Met Ser Val Tyr Val Val Glu
Leu Leu Arg Ser Phe Phe Tyr Val Thr Glu Thr Thr Phe
Gln Lys Asn Tyr Leu Phe Phe Tyr Arg Lys Ser Val Trp
Ser Lys Leu Gln Ser Ile Gly Ile Arg Gln His Leu Lys
Arg Val Gln Leu Arg Glu Leu Ser Glu Ala Glu Val Arg
Gln His Arg Glu Ala Arg Pro Ala Leu Leu Thr Ser Arg
Leu Arg Phe Leu Pro Lys Pro Asp Gly Leu Arg Pro Ile
Val Asn Met Asp Tyr Val Val Gly Ala Arg Thr Phe Arg
Arg Glu Lys Arg Ala Glu Arg Leu Thr Ser Arg Val Lys
Thr Leu Phe Ser Val Leu Asn Tyr Glu Arg Ala Arg Arg
Pro Gly Leu Leu Gly Ala Ser Val Leu Gly Leu Asp Asp
Ile His Arg Ala Trp Arg Ala Phe Val Leu Arg Val Arg
Ala Gln Asp Pro Pro Pro Glu Leu Tyr Phe Val Lys Val
Ala Val Thr Gly Ala Tyr Asp Thr Ile Pro Gln Asp Arg
Leu Thr Glu Val Ile Ala Ser Ile Ile Lys Pro Gln Asn
Thr Tyr Cys Val Arg Arg Tyr Ala Val Val Arg Arg Ala
Ala His Gly His Val Arg Lys Ser Phe Lys Arg His Val
Ser Thr Leu Thr Asp Leu Gln Pro Tyr Met Arg Gln Phe
Val Ala His Leu Gln Glu Thr Ser Pro Leu Arg Asp Ala
Val Val Ile Glu Gln Ser Ser Ser Leu Asn Glu Ala Ser
Ser Gly Leu Phe Asp Val Phe Leu Arg Phe Val Cys His

- 97 -

His Ala Val Arg Ile Gly Gly Lys Ser Tyr Val Gln Cys
 Gln Gly Ile Pro Gln Gly Ser Ile Leu Ser Thr Leu Leu
 Cys Ser Leu Cys Tyr Gly Asp Met Glu Asn Lys Leu Phe
 Ala Gly Ile Arg Arg Asp Gly Leu Leu Leu Arg Leu Val
 Ala Ala Phe Leu Leu Val Thr Pro His Leu Thr His Ala
 Lys Ala Phe Leu Arg Thr Leu Val Arg Gly Val Pro Glu
 Tyr Gly Cys Val Val Asn Leu Arg Lys Thr Val Val Asn
 Phe Pro Val Glu Asp Glu Ala Leu Gly Gly Thr Ala Phe
 Val Gln Leu Pro Ala His Gly Leu Phe Pro Trp Cys Gly
 Leu Leu Leu Asp Thr Arg Thr Leu Glu Val Gln Ser Asp
 Tyr Ser Ser Tyr Ala Arg Thr Ser Ile Arg Ala Ser Leu
 Thr Phe Asn Arg Gly Phe Lys Ala Gly Arg Asn Met Arg
 Arg Lys Leu Phe Gly Val Leu Arg Leu Lys Cys His Ser
 Leu Phe Leu Tyr Leu Gln Val Asn Ser Leu Gln Thr Val
 Cys Thr Asn Val Tyr Lys Ile Phe Leu Leu Gln Ala Tyr
 Arg Phe His Ala Cys Val Leu Gln Leu Pro Phe His Gln
 Gln Val Arg Lys Asn Pro Thr Phe Phe Leu Arg Val Ile
 Ser Asp Thr Ala Ser Leu Cys Tyr Ser Ile Leu Lys Ala
 Lys Asn Ala Gly Met Ser Leu Gly Ala Lys Gly Ala Ala
 Gly Pro Phe Pro Ser Glu Ala Ala Gln Trp Leu Cys His
 Gln Ala Phe Leu Leu Lys Leu Ala Arg His Arg Val Thr
 Tyr Arg Cys Leu Leu Gly Ala Leu Arg Thr Ala Gln Thr
 Gln Leu Cys Arg Lys Leu Pro Gly Ala Thr Leu Ala Ala
 Leu Glu Ala Ala Ala Asp Pro Ala Leu Thr Ser Asp Phe
 Lys Thr Ile Leu Asp

SEQ ID NO: 19 Synthetic Consensus hTERT nucleic acid sequence
 CCTAGAGCCCCACGGTGTAGAGCAGTCCGCAGCCTGCTGCGCAGCCGATACCGGGAAGTGCTGCC
 TCTGGCCACC
 TTTGTCCGGAGACTGGGACCACAGGGCAGGCGCCTGGTGCAGCGCGGCGACCCCGCAGCTTTCCG
 AGCACTGGTG
 GCACAGTGCCTGGTGTGCGTGCCATGGGATGCACGGCCCCCTCCAGCAGCCCCTAGCTTTAGACA
 GGTGTCCTGC
 CTGAAAGAACTGGTGCAGAGGGTGGTCCAGCGGCTGTGCGAGAGAGGCGCCAGGAACGTGCTGGC
 ATTCGGCTTT
 GCACTGCTGGACGGAGCTAGGGGCGGGCCCCCTGAGGCATTACCAACAAGCGTGCGCTCCTACCT
 GCCAAATACA
 GTCAGTATACCCTGCGAGGCTCCGGAGCATGGGGACTGCTGCTGCGACGGGTGGGGGACGATGT
 GCTGGTCCAC
 CTGCTGGCTAGATGCGCACTGTATGTGCTGGTGCCTCCCTCTTGCGCATACCAGGTGTGCGGACC
 ACCCTGTAT
 GACCTGGGCGCTGCAACCCAGGCAAGACCTCCACCCACGCCTCTGGCACTAGAAGGGGACTGGG

- 98 -

CACCGAACAG
GCATGGAACCATAGTGTCAGGGAGGCAGGAGTGCCACTGGGACTGCCAGCACCTGGGGCTCGCCG
ACGGAGAGGG
AGTGCCGGACGGTCACTGCCACTGGCTAAGAGACCAAGGCGGGAGCCGCTCCAGAACCAGAGAG
GACACCTGTG
GGACAGGGAAGCTGGGCACACCCTGGAAGAACTAGGGGGCCAAGTGATAGGGGCTTCTGCGTGGT
CTCACCAGCA
CGACCAGCAGAGGAAGCTACTTCTCTGGAGGGAGCTCTGAGTGGCACCCGGCACTCTCATCCTAG
TGTGGGAAGA
CAGCACCATGCAGGCCCTCCAAGCACCCAGCCGGCCTCCCCGGCCATGGGACACTCCTTGTCCACC
CGTGTACGCT
GAAACCAAACACTTTCTGTATAGCTCCGGAGATAAGGAGCAGCTGCGGCCCTCTTTCCTGCTGTC
TAGTCTGAGA
CCTAGTCTGACCGGAGCACGACGGCTGGTGGAAACAATCTTTCTGGGGTCCCCGCCCTTGGATGCC
AGGAACCCCC
AGAAGGACACCTCGACTGCCACAGCGGTACTGGCAGATGCGGCCACTGTTCTTGGAGCTGCTGGG
CAATCACGCT
CAGTGCCCTATGGGGCACTGCTGCGAACACATTGTCCTCTGCGGGCAGCCGTGACTCCAGCTGC
AGGAGTCTGC
GCCAGGGAAGCCACAGGGCAGCGTGGCAGCTCCTGAGGAAGAGGACACCGATCCACGCCGACT
GGTGCAGCTG
CTGAGACAGCACTCAAGCCCCTGGCAGGTGTACGGATTTCTGAGGGCCTGTCTGCGGAGACTGGT
GCCTCCAGGA
CTGTGGGGGTCCAGGCACAACGAAAGGCGCTTTCTGCGCAATACTAAGAAATTCATCAGCCTGGG
CAAGCATGCT
AAACTGTCCCTGCAGGAGCTGACCTGGAAAATGAGTGTGCGCGACTGCGCATGGCTGCGACGGTC
ACCAGGAGTC
GGGTGCGTGCCTGCAGCCGAGCACCGCCTGCGAGAAGAGATTCTGGCCAAGTTTCTGCATTGGCT
GATGTCACTG
TACGTGGTCGAACTGCTGCGGAGCTTCTTTTATGTGACAGAGACTACCTTCCAGAAAACTACCT
GTTCTTTTAT
CGCAAGTCAGTGTGGAGCAAACCTGCAGTCAATCGGCATTTCGGCAGCACCTGAAGAGAGTGCAGCT
GAGGGAACCTG
AGTGAAGCCGAGGTCCGGCAGCATAGAGAGGCAAGGCCTGCCCTGCTGACCTCCCGGCTGAGATT
CCTGCCTAAG
CCAGACGGGCTGAGACCAATCGTGAACATGGATTACGTGGTTCGGAGCACGGACCTTCCGGAGGGA
AAAACGCGCT
GAGCGACTGACATCCCGCGTGAAGACTCTGTTCTCTGTCTGAATTATGAGCGAGCTCGCCGACC
CGGACTGCTG
GGAGCATCTGTGCTGGGACTGGACGATATTACCGGGCTTGGAGAGCATTTGTCTGAGGGTGCG
CGCACAGGAC
CCTCCCCAGAAGTGTACTTCGTGAAAGTCGCCGTGACCGGGGCTTATGACACAATCCCTCAGGA
TCGGCTGACT
GAAGTGATCGCCTCCATCATTAAGCCACAGAATACCTACTGCGTGCGGAGATATGCTGTGGTCAG

- 99 -

GCGCGCTGCA
CACGGCCATGTGAGGAAGAGCTTCAAGCGCCACGTCAGCACACTGACTGATCTGCAGCCCTACAT
GAGACAGTTC
GTGGCTCATCTGCAGGAGACCAGCCCTCTGAGGGACGCAGTGGTCATCGAACAGTCCTCTAGTCT
GAACGAGGCA
TCAAGCGGGCTGTTTCGATGTCTTTCTGCGGTTTCGTGTGCCACCATGCCGTCAGAATTGGAGGCAA
ATCTTACGTG
CAGTGTGAGGGCATCCCCAGGGCAGCATTCTGTCTACCCTGCTGTGCAGCCTGTGCTATGGCGA
CATGGAAAAT
AAGCTGTTTGCCGGAATCCGACGGGATGGCCTGCTGCTGAGACTGGTGGCCGCTTTTCTGCTGGT
CACTCCACAC
CTGACCCATGCCAAAGCTTTCCTGCGCACACTGGTCCGAGGGGTGCCAGAGTACGGATGCGTGGT
CAACCTGAGG
AAGACCGTGGTCAATTTCCAGTGGAAGACGAGGCCCTGGGCGGCACAGCATTTGTCCAGCTGCC
AGCACACGGA

- 100 -

CTGTTCCCATGGTGTGGACTGCTGCTGGACACCCGCACACTGGAGGTGCAGTCCGATTACTC
CTCTTATGCCCCG
ACAAGCATCAGAGCTTCCCTGACTTTTAAACAGAGGCTTCAAGGCCGGGAGGAATATGAGAAG
GAAACTGTTTGGC
GTGCTGCGCCTGAAGTGCCATTCCCTGTTCTGTATCTGCAGGTGAACTCTCTGCAGACTGT
CTGTACCAACGTG
TACAAAATTTTTCTGCTGCAGGCCTATCGGTTCCACGCTTGCGTGCTGCAGCTGCCATTCCA
TCAGCAGGTCAGG
AAGAACCCACCTTCTTTCTGCGCGTGATCTCTGATACAGCTAGTCTGTGCTACTCAATTCT
GAAGGCCAAAAAT
GCTGGCATGAGCCTGGGAGCAAAGGAGCAGCAGGACCATTTCCTTCCGAGGCTGCACAGTG
GCTGTGCCACCAG
GCATTCCTGCTGAAGCTGGCCCGACATCGGGTGACATATAGGTGCCTGCTGGGCGCACTGCG
AACAGCACAGACT
CAGCTGTGCAGAAAGCTGCCCCGGGGCCACTCTGGCTGCCCTGGAAGCCGCTGCCGACCCTGC
CCTGACCTCCGAT TTCAAGACTATTCTGGACTGATAA

SEQ ID NO: 20 Synthetic Consensus hTERT amino acid sequence
PRAPRCRAVRSLLRSRYREVLPLATFVRLGPGQRRLVQRGDPAAFRALVAQCLVCPWDARPPP
AAPSFRQVSC
LKELVARVVQRLCERGARNVLAFCFALLDGARGGPPEAFTTSVRSYLPNTVTDTLRGSGAWGLLL
RRVGDDVLVH
LLARCALYVLVAPSCAYQVCGPPLYDLGAATQARPPPHASGTRRGLGTEQAWNHSVREAGVPLGL
PAPGARRRRG
SAGRSLPLAKRPRRGAAPEPERTPVGQGSWAHPGRTRGPSDRGFCVVSPARPAEEATSLEGALSG
TRHSHPSVGR
QHHAGPPSTSRPPRPWDTPCPPVYAETKHFLYSSGDKEQLRPSFLLSSLRPSLTGARRLVETIFL
GSRPWMPGTP
RRTPRLPQRYWQMRPLFLELLGNHAQCPYGALLRTHCPLRAAVTPAAGVCAREKPQGSVAAPEEE
DTDPRRLVQL
LRQHSSPWQVYGFRLACLRLVPPGLWGSRHNERFLRNTKKFISLGKHAKLSLQELTWKMSVRD
CAWLRRSPGV
GCVPAAEHRLREEILAKFLHWLMSVYVVELLRSFFYVTETTFQKNYLFFYRKSVWSKLQSIGIRQ
HLKRVQLREL
SEAEVRQHREARPALLTSRLRFLPKPDGLRPIVNMDYVVGARTFRREKRAERLTSRVKTLFSVLN
YERARRPGLL
GASVLGLDDIHRAWRAFLRVRAQDPPELYFVKVAVTGAYDTIPQDRLTEVIASIIKPQNTYCV
RRYAVVRRAA
HGHVRKSFKRHVSTLTDLPYMRQFVAHLQETSPLRDAVVIEQSSSLNEASSGLFDVFLRFVCHH
AVRIGGKSYV
QCQGI PQGSILSTLLCSLCYGD MENKLFAGIRRDGLLLRLVAAFLLVTPHLTHAKAFLRTLVRGV
PEYGCVVNLR

- 101 -

KTVVNFPVEDEALGGTAFVQLPAHGLFPWCGLLLDTRTLEVQSDYSSYARTSIRASLTFNRGFKA
GRNMRRKLFG
VLRLLKCHSLFLYLQVNSLQTVCTNVYKIFLLQAYRFHACVLQLPFHQQVRKNPTFFLRVISDTAS
LCYSILKAKN
AGMSLGAKGAAGFPFPSEAAQWLCHQAFLLKLARHRVTYRCLLGALRTAQTQLCRKLPGATLAALE
AAADPALTSDFKTILD**

SEQ ID NO: 21 nucleic acid sequence encoding PSMA with IgE
leader sequence
atggactggacatggattctgttcctggctgcgcgcgcgcaactcgcgtgcattcctggaacgcact
gcatgagact
gattctgctgtcgcactgggacggagacccccgggtggctgtgcgcgtggagcactgggtgctggccgg
cgggggattc
ctgctgggattcctgtttggctggtttatcaaaagctccagcgaggctaccaatattaccacctaa
gcacaataag
aaagcattcctggatgaactgaaagccgagaacatcaagaaattcctgtacaacttcacaagaat
tccacatctg
gctggcactgagcagaacttcagctggcaaacagatccagagtcagtggaaggaatttgggct
ggactcagtg
gagctgaccactacgatgtcctgctgtcctatccaaataagactcatcccaactacatctctat
cattaacgaa
gacggaaatgagattttcaacacctctctgtttgaacccccctccacccggctatgagaatgtcag
tgacgtggtc
ctccattctcagccttcagccccaggggatgcctgaggagatctgggtgtacgtcaattatgc
tagaacagaa
gacttctttaagctggagagggatatgaaaatcaactgttcgggaagatcgtgattgcccggtg
cgggaaggtg
ttcagaggaaataaggtcaaaaacgctcagctggccggagctaccggcgatcctgtacagcga
ccccgctgat
tattttgcacctggcgtgaagtcctatccagacggatggaatctgcccggcgggggagtgacag
gggaaacatc
ctgaacctgaatggagccggcgatcctctgactccaggataccccgccaaacgaatacgcttatcg
ccggggaatt
gcagaggccgtgggcctgcctagcatcccagtcctatccattggctattacgatgcccagaagct
gctggagaaa
atgggcgggagcgctccccctgactctagttggaagggtccctgaaagtgccttacaatgtcgg
gccaggattc
actgggaacttttctaccagaaggtgaaaatgcacatccatagtagcagcgaggtgacacgaat
ctacaacgtc
attggcaccctgagagggcgccgtggagcctgatcgctatgtcattctgggaggccacagagactc
atgggtgttc
gggggaatcgaatccacagagcggagcagctgtgggtccatgaaattgtgcgcagctttgggaccct
gaagaaagag
ggatggcgacccaggcgcaaatcctgttcgcatcctgggacgcgcgaggaatttgggctgctggg

- 102 -

cagcacagaa
tgggccgaggaaaattctcgctgctgcaggagcgaggggtggcttacatcaatgcagactcaag
cattgaagga
aactataccctgcgggtggattgcacacccctgatgtacagtctgggtctataacctgacaaagga
gctgaaatca
cctgacgagggccttgaagggaaaagcctgtacgaatcctggactgagaagagcccatccccga
attcagcggc
atgcctaggatctctaagctgggcagtggggaacgattttgaggtgttctttcagcgccctgggaat
tgcctctggc
cgagctcggtacacaaaaaattgggagactaacaagttctcctcttaccactgtatcacagcgt
gtacgagact
tatgaactggctcgagaaattctacgacccacttttaagtatcatctgaccgtggcacagggtcag
ggcggggatg
gtgttcgaactggccaatagcatcgctcctgccatttgactgtcgagattacgctgtggtcctgcg
gaagtacgca
gacaagatctataacatctccatgaagcacccccaggagatgaaggcctattctgtgagtttcga
ttccctgttt
tctgccgtcaaaaatttcaccgaaatcgctagtaagttttcagagcgccctgcaggacctggataa
gtccaatccc
atcctgctgcggattatgaacgatcagctgatgttcctggaaagagcctttatcgaccctctggg
cctgcctgat
agaccattctacaggcacgtgatctacgcacctagttcacataacaagtacgccggcgagtttt
cccagggatc

- 103 -

tatgacgctctgtttgatattgaatcaaaggtggaccccagcaaagcatggggcgagggtcaagag
acagatcagc
attgcagcctttacagtgcaggccgcccgaaccctgtccgaagtcgcttgataa

SEQ ID NO: 22 nucleic acid sequence encoding IL12 p35
(pGX6001)

atgtgtccagcgcgcagcctcctccttgtggctaccctggctcctcctggaccacctcagtttggc
cagaaacctc
cccgtggccactccagacccaggaatgttcccatgccttcaccactcccaaacctgctgagggc
cgtcagcaac
atgctccagaaggccagacaaactctagaatttacccttgcaattctgaagagattgatcatga
agatatcaca
aaagataaaaccagcacagtggaggcctgtttaccattggaattaaccaagaatgagagttgcct
aaattccaga
gagacctctttcataactaatgggagttgcctggcctccagaaagacctcttttatgatggccct
gtgccttagt
agtatttatgaagacttgaagatgtaccaggtggagttcaagaccatgaatgcaaagcttctgat
ggatcctaag
aggcagatctttctagatcaaaacatgctggcagttattgatgagctgatgcaggccctgaattt
caacagtgag
actgtgccacaaaaatcctcccttgaagaaccggatttttataaaaactaaaatcaagctctgcat
acttcttcat
gctttcagaattcgggcagtgactattgatagagtgatgagctatctgaatgcttcctaa

SEQ ID NO: 23 IL12 p35 amino acid sequence (pGX6001)

MCPARSLLLVATLVLLDHLNLPVATPDGPMFCLHHSQNLLRAVSNNMLQKARQTLEFYFCT
SEEDHEDIT
KDKTSTVEACLPLELTKNESCLNSRETSFITNGSCLSRKTSFMMALCLSSIEDLKMYQVEFKT
MNAKLLMDPK
RQIFLDQNMLAVIDELMQALNFNSETVPQKSSLEEPDFYKTKIKLCILLHAFRIRAVTIDR
VMSYLNAS

SEQ ID NO: 24 nucleic acid sequence encoding IL12
p40 (pGX6001)

atgtgtcaccagcagttggtcatctcttgggttttccctgggttttctggcatctcccctcgtggc
catatgggaa
ctgaagaaagatgtttatgtcgtagaattggattggtatccggatgcccctggagaaatggtggt
cctcacctgt
gacaccttgaagaagatggtatcacctggaccttggaccagagcagtgagggtcttaggctctgg
caaacacctg
accatccaagtcaaagagtttggagatgctggccagtacacctgtcacaaggaggcgaggttct
aagccattcg

- 104 -

ctcctgctgcttcacaaaaaggaagatggaatttggtccactgatatttttaaggaccagaaaga
acccaaaaat
aagaccttttctaagatgcgaggccaagaattattctggacgtttcacctgctggtggctgacgac
aatcagtact
gatttgacattcagtggtcaaaagcagcagaggctcttctgaccccccaaggggtgacgtgaggagc
tgctacactc
tctgcagagagagtcagaggggacaacaaggagtatgagtactcagtgaggagtgccaggaggacag
tgctgcccc
gctgctgaggagagtcctgcccattgaggtcatggtggatgccgttcacaagctcaagtatgaaaa
ctacaccagc
agcttcttcatcaggacatcatcaaacctgacccaccaagaacttgacagctgaagccattaaa
gaattctcgg
caggtggaggtcagctgggagtaccctgacacctggagtactccacattcctacttctccctgac
attctgcgtt
caggtccagggaagagcaagagagaaaaagaaagatagagtcttcacggacaagacctcagccac
ggtcatctgc
cgcaaaaatgccagcattagcgtgcgggcccaggaccgctactatagctcatcttgagcgcaatg
ggcatctgtg ccctgcagttag

SEQ ID NO: 25 IL12 p40 amino acid sequence (pGX6001)
MCHQQLVISWFLVFLASPLVAIWELKKDVYVVELDWYPDAPGEMVVLTCDTPEEDGITWTLDQS
SEVLGSGKTL
TIQVKEFGDAGQYTCHKGGEVLSHSLLLLHKKEDGIWSTDILKDQKEPKNKTFLRCEAKNYSGRF
TCWWLTITIST
DLTFSVKSSRGSSDPQGVTCGAATLSAERVGRDNKEYEYSVEQCQEDSACPAAEESLPIEVMVDAV
HKLKYENYTS
SFFIRDIKPDPPKNLQLKPLKNSRQVEVSWEYPDTWSTPHSYFSLTFCVQVQGKSKREKKDRVF
TDKTSATVIC RKNASISVRAQDRYYSSSWSEWASVPCS

SEQ ID NO: 26 Con WT1-L with modified Zinc Fingers protein
GSDVRDLNALLPAVPSLPGGGGALPVSGAAQWAPVLDFAPPAAPYGLGGPHSFIKQEPSWGGA
DPHEEQCLSA
FTVHFSGQFTGTAGACRYGPFAPPPSQAPSGQARMFPNAPYLPNCLESQPAIRNQGYSTVAFDG
TPSYGHTPSH
HAAQFPNHSFKHEDPMGQQGSLGEQQYSVPPPVYGCHTPTDCTGSQALLLRTPYNSDNLYQMTS
QLECMTNQM
NLGSTLKGHATGYESDNHTTPMLYSCGAQYRIHTHGVRGIQDVRRVPGVAPTIVRSASETNEKR
PFMGAYPGGN
KRYFKLSHLQMGRKGTGEKPYQGDFKDGERRFSRSDQLKRGQRRGTGVKPFQCKTCQRKFSRSD
HLKTHTRHT GKTSEKPFSCRWPSCQKKFARSDELVRHHNMHQRNMTKLQLAL**

SEQ ID NO: 27 Consensus WT1-L with modified Zinc Fingers
nucleic acid sequence
GGGAGTGATGTGAGAGACCTGAACGCCCTGCTGCCAGCAGTGCCATCCCTGCCTGGCGGGGG

- 105 -

AGGCTGCGCTCTG
CCAGTCTCTGGAGCAGCTCAGTGGGCTCCCGTGCTGGACTTTGCACCCCCTGCAGCCCCTTA
CGGAAGTCTGGGC
GGCCCACACTCATTTCATCAACAGGAGCCAAGCTGGGGCGGGGCAGATCCTCATGAGGAACA
GTGCCTGTCAGCC
TTCACAGTCCACTTTAGCGGGCAGTTCACTGGAACCGCAGGAGCTTGTAGATACGGACCCTT
TGGAGCACCACCC
CCTTCCCAGGCACCTTCTGGACAGGCACGCATGTTCCCAAACGCTCCCTATCTGCCTAATTG
TCTGGAAAGCCAG
CCCGCTATTAGGAACCAGGGCTACTCCACAGTGGCATTGTGACGGGACTCCTAGCTATGGACA
TACCCCATCCCAC
CATGCTGCACAGTTTCCTAATCACTCCTTCAAGCATGAGGACCCCATGGGACAGCAGGGGTC
CCTGGGAGAACAG

- 106 -

CAGTACTCTGTGCCCCCTCCCGTGTACGGATGCCACACACCAACTGACAGTTGTACAGGCTC
ACAGGCCCTGCTG
CTGCGAACTCCATACAACAGTGATAATCTGTATCAGATGACCTCACAGCTGGAGTGCATGAC
ATGGAACCAGATG
AATCTGGGCAGCACACTGAAAGGCCATGCCACTGGGTACGAATCTGACAACCACACCACACC
TATGCTGTACAGT
TGTGGAGCCCAGTATAGAATCCACACTCATGGAGTCTTCAGAGGCATTGAGGATGTGCGGAG
AGTCCCAGGAGTG
GCACCAACTATCGTGCAGGAGCGCCTCCGAGACCAACGAAAAGCGCCCCCTTTATGGGCGCCTA
CCCTGGAGGCAAT
AAGCGGTATTTCAAACGTCTCTCACCTGCAGATGGGGAGTAGAAAGGGGACCGGAGAGAAAACC
TTATCAGGGCGAC
TTTAAAGATGGGGAAAGGCGCTTCTCTCGCAGTGACCAGCTGAAGCGAGGACAGCGACGAGG
AACCAGGGTGAAG
CCATTTCACTGCAAAACATGTCTCAGAGAAAGTTCTCAAGGAGCGATCACCTGAAGACCCATAC
AAGAACTCACACC
GGCAAGACCAGCGAGAAACCATTTTCTGCGGATGGCCCTCTTGTCAGAAGAAATTCGCCCC
CTCCGACGAACTG
GTCCGACACCACAATATGCATCAGAGAAATATGACAAAACCTGCAGCTGGCTCTGTGATA
A

SEQ ID NO: 28 PSMA

WNALHETDSVALGRRPRWLCAGALVLAGGGFLLGFLFGWFIKSSSEATNITPKHNKKAFLDELK
AENIKKFLYN
FTRIPHLAGTEQNFLAKQIQSQWKEFGLDSVELTHYDVLLSYPNKTHPNYISIIINEDGNEIFNT
SLFEPPPPGY
ENVSDVVPFSAFSPQGMPEGDLVYVNYARTEDFFKLERDMKINCSGKIVIARYGKVFRGNKVKN
AQLAGATGVI
LYSDPADYFAPGVKSYPDGWNLPGGGVQGRNINLNLNGAGDPLTPGYPANAYARRGIAEAVGLPS
IPVHPIGYYD
AQKLEKMGGSAAPPDSSWKSLKVPYNVPGPGFTGNFSTQKVKMHIHSTSEVTRIYNVIGTLRGAV
EPDRYVILGG
HRDSWVFGGIDPQSGAAVVHEIVRSFGTLKKEGWRPRRTILFASWDAEEFGLLGSTEWAEENSRL
LQERGVAYIN
ADSSIEGNYTLRVDCTPLMYSLVYNLTKEKSPDEGFEGKSLYESWTEKSPSPEFSGMPRISKLG
SGNDFEVFFQ
RLGASGRARYTKNWETNKFSYPLYHSVYETYELVEKFYDPTFKYHLTVAQVRGGMVFELANSI
VLPFDCRDYA
VVLRYADKIYNISMKHPQEMKAYSVSFDSLFSVKNFTEIASKFSERLQDLDKSNPILLRIMND
QLMFLERAFT
DPLGLPDRPFYRHVIYAPSSHNKYAGESFPGIYDALFDIESKVDPSKAWGEVKRQISIAAFTVQA
AAETLSEVA*

*

- 107 -

SEQ ID NO: 29 nucleic acid sequence encoding PSMA
TGGAACGCACTGCATGAGACTGATTCTGCTGTCGCACTGGGACGGAGACCCCGGTGGCTGT
GCGCTGGAGCACTG
GTGCTGGCCGGCGGGGATTCCCTGCTGGGATTCCCTGTTTGGCTGGTTTATCAAAAGCTCCA
GCGAGGCTACCAAT
ATTACCCCTAAGCACAATAAGAAAGCATTCCTGGATGAACTGAAAGCCGAGAACATCAAGA
AATTCCTGTACAAC
TTCACAAGAATTCCACATCTGGCTGGCACTGAGCAGAACTTCCAGCTGGCAAAACAGATCC
AGAGTCAGTGGAAAG
GAATTTGGGCTGGACTCAGTGGAGCTGACCCACTACGATGTCCTGCTGTCTATCCAAATA
AGACTCATCCCAAC
TACATCTCTATCATTAACGAAGACGGAAATGAGATTTTCAACACCTCTCTGTTTGAACCCC
CTCCACCCGGCTAT
GAGAATGTCAGTGACGTGGTCCCTCCATTCTCAGCCTTCAGCCCCAGGGGATGCCTGAGG
GAGATCTGGTGTAC
GTCAATTATGCTAGAACAGAAGACTTCTTTAAGCTGGAGAGGGATATGAAAATCAACTGTT
CCGGCAAGATCGTG
ATTGCCCGGTACGGGAAGGTGTTTCCAGAGGAAATAAGGTCAAAAACGCTCAGCTGGCCGGAG
CTACCGGCGTGATC
CTGTACAGCGACCCCGCTGATTATTTTGCACCTGGCGTGAAGTCTATCCAGACGGATGGA
ATCTGCCCCGGCGGG
GGAGTGCAGAGGGGAAACATCCTGAACCTGAATGGAGCCGGCGATCCTCTGACTCCAGGAT
ACCCCGCCAACGAA
TACGCTTATCGCCGGGAATTGCAGAGGCCGTGGGCCTGCCTAGCATCCCAGTCCATCCCA
TTGGCTATTACGAT
GCCCAGAAGCTGCTGGAGAAAATGGGCGGGAGCGCTCCCCCTGACTCTAGTTGGAAGGGCT
CCCTGAAAGTGCCT
TACAATGTCGGGCCAGGATTCCTGGGAACCTTTTCTACCCAGAAGGTGAAAATGCACATCC
ATAGTACCAGCGAG
GTGACACGAATCTACAACGTCATTGGCACCCCTGAGAGGCGCCGTGGAGCCTGATCGCTATG
TCATTCTGGGAGGC
CACAGAGACTCATGGGTGTTTCGGGGGAATCGATCCACAGAGCGGAGCAGCTGTGGTCCATG
AAATTGTGCGCAGC
TTTGGGACCCCTGAAGAAAGAGGGATGGCGACCCAGGCGCACAAATCCTGTTGCGATCCTGGG
ACGCCGAGGAATTT
GGGCTGCTGGGCAGCACAGAATGGGCCGAGGAAAATTCCTCGCCTGCTGCAGGAGCGAGGGG
TGGCTTACATCAAT
GCAGACTCAAGCATTGAAGGAACTATACCCTGCGGGTGGATTGCACACCCCTGATGTACA
GTCTGGTCTATAAC
CTGACAAAGGAGCTGAAATCACCTGACGAGGGCTTCGAAGGGAAAAGCCTGTACGAATCCT
GGACTGAGAAGAGC
CCATCCCCGAATTCAGCGGCATGCCTAGGATCTCTAAGCTGGGCAGTGGGAACGATTTTG
AGGTGTTCTTTTCAG

- 108 -

CGCCTGGGAATTGCCTCTGGCCGAGCTCGGTACACAAAAAATTGGGAGACTAACAAGTTCT
CCTCTTACCCACTG
TATCACAGCGTGTACGAGACTTATGAACTGGTCGAGAAATTCTACGACCCCACTTTTAAGT
ATCATCTGACCGTG
GCACAGGTCAAGGGCGGGATGGTGTTCGAACTGGCCAATAGCATCGTCCTGCCATTTGACT
GTCGAGATTACGCT
GTGGTCCTGCGGAAGTACGCAGACAAGATCTATAACATCTCCATGAAGCACCCCCAGGAGA
TGAAGGCCTATTCT
GTGAGTTTCGATTCCCTGTTTTCTGCCGTCAAAAATTTACCGAAATCGCTAGTAAGTTTT
CAGAGCGCCTGCAG
GACCTGGATAAGTCCAATCCCATCCTGCTGCGGATTATGAACGATCAGCTGATGTTCTTG
AAAGAGCCTTTATC
GACCCTCTGGGCCTGCCTGATAGACCATTCTACAGGCACGTGATCTACGCACCTAGTTCAC
ATAACAAGTACGCC
GGCGAGTCTTTCCAGGGATCTATGACGCTCTGTTTGATATTGAATCAAAGGTGGACCCCA
GCAAAGCATGGGGC
GAGGTCAAGAGACAGATCAGCATTGCAGCCTTTACAGTGCAGGCCGCCGCCGAAACCCTGT
CCGAAGTCGCTTGA
TAA

SEQ ID NO: 30 miR-331-3p (miRbase accession #
MIMAT0000760)
GCCCCUGGGCCUAUCCUAGAA

SEQ ID NO: 31 miR-1537-3p (miRbase accession #
MIMAT0007399)
AAAACCGUCUAGUUACAGUUGU

- 109 -

What is claimed:

1. A method of treating brain cancer in a subject, comprising:
 - (a) measuring an expression level of a miR-331-3p miRNA comprising SEQ ID NO: 30 or isomiRs thereof and a miR-1537-3p miRNA comprising SEQ ID NO: 31 or isomiRs thereof in a biological sample from a subject who has brain cancer; and
 - (b) administering interleukin-12 (IL-12); an immunogenic composition of human telomerase reverse transcriptase (hTERT), Wilms Tumor-1 (WT-1), and prostate specific membrane antigen (PSMA); and an anti-programmed cell death receptor 1 (PD-1) antibody to said subject if the subject has an increased expression level of the miR-331-3p miRNA or isomiRs thereof and the miR-1537-3p miRNA or isomiRs thereof relative to a control population of subjects.
2. The method of claim 1 or 2, wherein the presence of the miRNA is determined by RNA sequencing.
3. The method of any one of the preceding claims, wherein the biological sample is a plasma sample.
4. The method of any one of the preceding claims, wherein the biological sample is taken from the subject prior to administration of the IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and the anti-PD-1 antibody.
5. The method of any one of the preceding claims, wherein administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody results in survival at 18 months post administration.
6. The method of any one of the preceding claims, wherein the control population of subjects is deceased at 18 months post administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody.

- 110 -

7. The method of any one of the preceding claims, wherein the subject has an unmethylated O6-methylguanine methyltransferase (MGMT) gene promoter.
8. A method of treating brain cancer in a subject, comprising:
 - (a) measuring an expression level of at least one mRNA biomarker selected from SYNGR3, OTX1, GABBR2, LHX1, CADM3, MLLT11, MNX1, GRB14, SLC34A2, PHYHIP, WNT10B, SLC17A6, CRLF1, HOXD13, TGF β 3, UBA7, SFRP4, or any combination thereof, in a tumor sample from a subject who has brain cancer; and
 - (b) administering interleukin-12 (IL-12); an immunogenic composition of human telomerase reverse transcriptase (hTERT), Wilms Tumor-1 (WT-1), and prostate specific membrane antigen (PSMA); and an anti-programmed cell death receptor 1 (PD-1) antibody to said subject if the expression level of SYNGR3, OTX1, GABBR2, LHX1, CADM3, MLLT11, MNX1, GRB14, SLC34A2, PHYHIP, WNT10B, SLC17A6, CRLF1 and HOXD13 is decreased or if the expression level of TGF β 3, UBA7, SFRP4 is increased relative to a control population of subjects.
8. The method of claim 8, wherein the subject has an unmethylated O6-methylguanine methyltransferase (MGMT) gene promoter.
9. The method of claim 7, wherein the biological sample is a primary tumor sample.
10. The method of any one of the preceding claims, wherein the biological sample is taken from the subject prior to administration of the IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and the anti-PD-1 antibody.
11. The method of claim 7 or 8, wherein administration of IL-12; an immunogenic composition of hTERT, WT-1, and PSMA; and an anti anti-PD-1 antibody results in survival at 18 months post administration.
12. The method of any one of the preceding claims, wherein the control population of subjects is deceased at 18 months post administration of IL-12; an immunogenic

- 111 -

composition of hTERT, WT-1, and PSMA; and an anti-PD-1 antibody.

13. The method of any one of the preceding claims, wherein IL-12 is encoded by a DNA plasmid.

14. The method of any one of the preceding claims, wherein hTERT is encoded by a DNA plasmid, WT-1 is encoded by a DNA plasmid, and/or PSMA is encoded by a DNA plasmid.

15. The method of any one of the preceding claims, wherein hTERT, WT-1, and PSMA are encoded by the same DNA plasmid; two of hTERT, WT-1, and PSMA are encoded by the same DNA plasmid; or wherein hTERT, WT-1, and PSMA are each encoded by a different DNA plasmid.

16. The method of any one of the preceding claims, wherein the anti-PD-1 antibody:
comprises the heavy chain complementarity determining regions (HCDR1, HCDR2 and HCDR3) of a heavy chain variable region (HCVR) comprising the amino acid sequence of SEQ ID NO: 1 and three light chain complementarity determining regions (LCDR1, LCDR2 and LCDR3) of a light chain variable region (LCVR) comprising the amino acid sequence of SEQ ID NO: 2;

comprises three HCDRs (HCDR1, HCDR2 and HCDR3) and three LCDRs (LCDR1, LCDR2 and LCDR3), wherein HCDR1 comprises the amino acid sequence of SEQ ID NO: 3, HCDR2 comprises the amino acid sequence of SEQ ID NO: 4, HCDR3 comprises the amino acid sequence of SEQ ID NO: 5, LCDR1 comprises the amino acid sequence of SEQ ID NO: 6, LCDR2 comprises the amino acid sequence of SEQ ID NO: 7, and LCDR3 comprises the amino acid sequence of SEQ ID NO: 8;

comprises a HCVR with 90% sequence identity to SEQ ID NO: 1;

comprises a LCVR with 90% sequence identity to SEQ ID NO: 2;

comprises a HCVR with 90% sequence identity to SEQ ID NO: 1 and a LCVR with 90% sequence identity to SEQ ID NO: 2;

- 112 -

comprises a HCVR comprising the amino acid sequence of SEQ ID NO: 1 and a LCVR comprising the amino acid sequence of SEQ ID NO: 2;

comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 9 and a light chain comprising the amino acid sequence of SEQ ID NO: 10;

is an IgG4 antibody; or

is REGN2810 or a biosimilar or bioequivalent thereof.

17. The method of any one of the preceding claims, wherein the anti-PD-1 antibody is administered intravenously or subcutaneously.

18. The method of any one of the preceding claims, wherein 350 mg of the anti-PD-1 antibody is administered every three weeks.

19. The method of any one of the preceding claims, wherein:

the IL12 p35 subunit comprises the amino acid sequence of SEQ ID NO: 23;

the IL12 p40 subunit comprises the amino acid sequence of SEQ ID NO: 25;

the IL12 p35 subunit comprises the amino acid sequence of SEQ ID NO: 23 and the IL12 p40 subunit comprises the amino acid sequence of SEQ ID NO: 25;

the IL12 p35 subunit is encoded by the nucleic acid sequence of SEQ ID NO: 22;

the IL12 p40 subunit is encoded by the nucleic acid sequence of SEQ ID NO: 24; or

the IL12 p35 subunit is encoded by the nucleic acid sequence of SEQ ID NO: 22 and the IL12 p40 subunit is encoded by the nucleic acid sequence of SEQ ID NO: 24.

20. The method of any one of the preceding claims, wherein

the hTERT comprises the amino acid sequence of SEQ ID NO: 20 or is encoded by the nucleic acid sequence of SEQ ID NO: 19;

the WT-1 comprises the amino acid sequence of SEQ ID NO: 26 or is encoded by the nucleic acid sequence of SEQ ID NO: 27; and/or

the PSMA comprises the amino acid sequence of SEQ ID NO: 28 or is encoded by

- 113 -

the nucleic acid sequence of SEQ ID NO: 29.

21. The method of claim 19, wherein the method comprises administering to the subject 3 mg of the DNA plasmid encoding hTERT, 3 mg of the DNA plasmid encoding PSMA, 3 mg of the DNA plasmid encoding WT-1, and 1 mg of the plasmid encoding IL-12.
22. The method of any one of the preceding claims, wherein the IL-12 and the immunogenic composition are co-administered by intramuscular injection every three weeks for four doses and then every nine weeks.
23. The method of claim 21, further comprising electroporation following each intramuscular injection.
24. The method of any one of the preceding claims, further comprising administering to the subject one or more doses of radiation therapy.
25. The method of claim 23, wherein each dose of radiation therapy comprises 20-50 Gy.
26. The method of claim 24, wherein the radiation therapy is fractionated radiation therapy.
27. The method of claim 25, wherein the fractionated radiation therapy comprises 2-20 fractions.
28. The method of claim 26, wherein the fractionated radiation therapy comprises 40 Gy in 15 fractions.
29. The method of claim 27, wherein the fractionated radiotherapy is given over 21 consecutive days.

- 114 -

30. The method of any one of the preceding claims, further comprising administering to the subject one or more doses of a chemotherapeutic agent; optionally wherein the chemotherapeutic agent is temozolomide.
31. The method of claim 29, comprising administering 75 mg/m^2 temozolomide to the subject daily for 21 consecutive days with the fractionated radiotherapy.
32. The method of claim 29 or 30, further comprising administering temozolomide maintenance therapy to the subject if the subject has a tumor having a methylated MGMT promoter.
33. The method of any one of the preceding claims, wherein the brain cancer is glioblastoma.
34. The method of claim 32, wherein the brain cancer is MGMT-methylated glioblastoma.
35. The method of claim 32, wherein the brain cancer is MGMT-unmethylated glioblastoma.