

ANTI-CD71 ANTIBODIES, ACTIVATABLE ANTI-CD71 ANTIBODIES, AND METHODS OF USE THEREOF

Related Applications

[0001] This application claims the benefit under of U.S. Provisional Application Nos. 62/156,838, filed May 4, 2015; 62/257,321, filed November 19, 2015; 62/257,484, filed November 19, 2015; 62/277,775, filed January 12, 2016; 62/310,553, filed March 18, 2016; and 62/315,276, filed March 30, 2016, the contents of each of which are incorporated herein by reference in their entirety.

Field of the Invention

[0002] The invention relates generally to antibodies that bind CD71, activatable antibodies that specifically bind to CD71 and methods of making and using these anti-CD71 antibodies and anti-CD71 activatable antibodies in a variety of therapeutic, diagnostic and prophylactic indications.

Background of the Invention

[0003] Antibody-based therapies have proven effective treatments for several diseases but in some cases, toxicities due to broad target expression have limited their therapeutic effectiveness. In addition, antibody-based therapeutics have exhibited other limitations such as rapid clearance from the circulation following administration.

[0004] In the realm of small molecule therapeutics, strategies have been developed to provide prodrugs of an active chemical entity. Such prodrugs are administered in a relatively inactive (or significantly less active) form. Once administered, the prodrug is metabolized *in vivo* into the active compound. Such prodrug strategies can provide for increased selectivity of the drug for its intended target and for a reduction of adverse effects.

[0005] Accordingly, there is a continued need in the field of antibody-based therapeutics for antibodies that mimic the desirable characteristics of the small molecule prodrug.

Summary of the Invention

[0006] The disclosure provides antibodies or antigen-binding fragments thereof that specifically bind CD71, also known as transferrin receptor protein 1 (TfR1).

[0007] In some embodiments, the antibody includes an antibody or antigen-binding fragment thereof that specifically binds CD71. In some embodiments, the antibody or antigen-binding fragment thereof that binds CD71 is a monoclonal antibody, domain antibody, single chain, Fab fragment, a F(ab')₂ fragment, a scFv, a scAb, a dAb, a single domain heavy chain antibody, or a single domain light chain antibody. In some embodiments, such an antibody or antigen-binding fragment thereof that binds CD71 is a mouse, other rodent, chimeric, humanized or fully human monoclonal antibody.

[0008] In some embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1, and 3-5. In some embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 3-5.

[0009] In some embodiments, the antibody or antigen-binding fragment thereof comprises a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8. In some embodiments, the antibody or antigen-binding fragment thereof comprises a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00010] In some embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a light chain variable region amino acid sequence or antigen-binding fragment thereof selected from the group consisting of SEQ ID NO: 2 and 6-8.

[00011] In some embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 3-5, and a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00012] In some embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5. In some embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 3-5.

[00013] In some embodiments, the antibody or antigen-binding fragment thereof comprises a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8. In some embodiments, the antibody or antigen-binding fragment thereof comprises a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00014] In some embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[00015] In some embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 3-5, and a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00016] In some embodiments, the antibody or antigen-binding fragment thereof comprises a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3) sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2) sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one complementarity determining region (CDR) sequence is selected from the group consisting of a VH CDR1 sequence comprising the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); a VH CDR2 sequence comprising the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); a VH

CDR3 sequence comprising the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); a VL CDR1 sequence comprising the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); a VL CDR2 sequence comprising the amino acid sequence STSNLAS (SEQ ID NO: 14); and a VL CDR3 sequence comprising the amino acid sequence QQRRNYPYT (SEQ ID NO: 15).

[00017] In some embodiments, the antibody or antigen-binding fragment thereof comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR1 sequence comprising the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); a VH CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR2 sequence comprising the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); a VH CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR3 sequence comprising the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); a VL CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR1 sequence comprising the amino acid sequence ASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); a VL CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR2 sequence comprising the amino acid sequence STSNLAS (SEQ ID NO: 14); and a VL CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR3 sequence comprising the amino acid sequence QQRRNYPYT (SEQ ID NO: 15).

[00018] In some embodiments, the antibody or antigen-binding fragment thereof comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR1 sequence comprises the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence comprises the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence comprises the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence comprises the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2

sequence comprises the amino acid sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence comprises the amino acid sequence QQRRNYPYT (SEQ ID NO: 15).

[00019] In some embodiments, the antibody or antigen-binding fragment thereof comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR1 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to comprises the amino acid sequence QQRRNYPYT (SEQ ID NO: 15).

[00020] In some embodiments, the antibody or antigen-binding fragment thereof is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5. In some embodiments, the antibody or antigen-binding fragment thereof is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 3-5.

[00021] In some embodiments, the antibody or antigen-binding fragment thereof is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8. In some embodiments, the antibody or antigen-binding fragment thereof is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00022] In some embodiments, the antibody or antigen-binding fragment thereof is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a nucleic acid sequence encoding a light chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[00023] In some embodiments, the antibody or antigen-binding fragment thereof is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 3-5, and a nucleic acid sequence encoding a light chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00024] In some embodiments, the antibody or antigen-binding fragment thereof is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5. In some embodiments, the antibody or antigen-binding fragment thereof is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 3-5.

[00025] In some embodiments, the antibody or antigen-binding fragment thereof is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a light chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8. In some embodiments, the antibody or antigen-binding fragment thereof is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a light chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00026] In some embodiments, the antibody or antigen-binding fragment thereof is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a light chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[00027] In some embodiments, the antibody or antigen-binding fragment thereof is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 3-5, and a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a light chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00028] The disclosure also provides methods for producing an antibody of the disclosure by culturing a cell under conditions that lead to expression of the antibody, wherein the cell comprises a nucleic acid molecule of the disclosure or a vector of the disclosure.

[00029] In some embodiments, the antibody or antigen-binding fragment thereof is incorporated in a multispecific antibody or antigen-binding fragment thereof, where at least one arm of the multispecific antibody or antigen-binding fragment thereof specifically binds CD71. In some embodiments, the antibody or antigen-binding fragment thereof is incorporated in a bispecific antibody or antigen-binding fragment thereof, where at least one arm of the bispecific antibody or antigen-binding fragment thereof specifically binds CD71.

[00030] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5. In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 3-5.

[00031] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8. In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00032] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[00033] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain variable region amino acid sequence c selected from the group consisting of SEQ ID NO: 3-5, and a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00034] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5. In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 3-5.

[00035] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8. In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific

antibody or antigen-binding fragment thereof, comprises a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00036] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[00037] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 3-5, and a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00038] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3) sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2) sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one complementarity determining region (CDR) sequence is selected from the group consisting of a VH CDR1 sequence comprising the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); a VH CDR2 sequence comprising the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); a VH

CDR3 sequence comprising the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); a VL CDR1 sequence comprising the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); a VL CDR2 sequence comprising the amino acid sequence STSNLAS (SEQ ID NO: 14); and a VL CDR3 sequence comprising the amino acid sequence QQRRNYPYT (SEQ ID NO: 15).

[00039] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR1 sequence comprising the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); a VH CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR2 sequence comprising the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); a VH CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR3 sequence comprising the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); a VL CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR1 sequence comprising the amino acid sequence ASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); a VL CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR2 sequence comprising the amino acid sequence STSNLAS (SEQ ID NO: 14); and a VL CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR3 sequence comprising the amino acid sequence QQRRNYPYT (SEQ ID NO: 15).

[00040] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR1 sequence comprises the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence comprises the amino acid sequence

AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence comprises the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence comprises the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence comprises the amino acid sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence comprises the amino acid sequence QQRRNYPYT (SEQ ID NO: 15).

[00041] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR1 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence QQRRNYPYT (SEQ ID NO: 15).

[00042] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain or a heavy chain variable region that comprises or is derived from a heavy chain amino acid sequence or heavy chain variable region amino acid sequence shown in Table 12. In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a light chain or a light chain variable region that comprises or is derived from a light chain amino acid sequence or light chain variable region amino acid sequence shown in Table 12. In some embodiments, at least one arm of the multispecific

antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain or a heavy chain variable region amino acid sequence that comprises or is derived from a heavy chain amino acid sequence or heavy chain variable region amino acid sequence shown in Table 12 and a light chain or a light chain variable region amino acid sequence that comprises or is derived from a light chain amino acid sequence or light chain variable region amino acid sequence shown in Table 12.

[00043] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the heavy chain variable region sequences shown in Table 12. In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence that is selected from the group consisting of the light chain variable region sequences shown in Table 12. In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the heavy chain variable region sequences shown in Table 12 and a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the light chain variable region sequences shown in Table 12.

[00044] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3) sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2)

sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence shown in Table 13; a VH CDR2 sequence shown in Table 13; a VH CDR3 sequence shown in Table 13; a VL CDR1 sequence shown in Table 13; a VL CDR2 sequence shown in Table 13; and a VL CDR3 sequence shown in Table 13.

[00045] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR1 sequence shown in Table 13; a VH CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR2 sequence shown in Table 13; a VH CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR3 sequence shown in Table 13; a VL CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR1 sequence shown in Table 13; a VL CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR2 sequence shown in Table 13; and a VL CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR3 sequence shown in Table 13.

[00046] In some embodiments at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 13.

[00047] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a light chain variable region that comprise a combination of a VL CDR1

sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the three light chain CDR sequences (VL CDR1, VL CDR2, VL CDR3) shown in a single row in Table 13.

[00048] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain variable region that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein the combination is a combination of the three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 13.

[00049] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 13.

[00050] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a heavy chain variable region that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 13.

[00051] In some embodiments, at least one arm of the multispecific antibody or antigen-binding fragment thereof, *e.g.*, a bispecific antibody or antigen-binding fragment thereof, comprises a light chain variable region that comprise a combination of a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of three light chain CDR sequences (VL CDR1, VL CDR2, VL CDR3) shown in a single row in Table 13.

[00052] Suitable anti-CD71 antibodies of the disclosure also include an antibody or antigen binding fragment thereof that binds to the same epitope on human CD71 and/or cynomolgus monkey CD71 as an anti-CD71 antibody comprising a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[00053] Suitable anti-CD71 antibodies of the disclosure also include an antibody or antigen binding fragment thereof that binds to the same epitope on human CD71 and/or cynomolgus monkey CD71 as an anti-CD71 antibody comprising a VH CDR1 sequence comprising the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); a VH CDR2 sequence comprising the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); a VH CDR3 sequence comprising the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); a VL CDR1 sequence comprising the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); a VL CDR2 sequence comprising the amino acid sequence STSNLAS (SEQ ID NO: 14); and a VL CDR3 sequence comprising the amino acid sequence QRRNYPYT (SEQ ID NO: 15).

[00054] Suitable anti-CD71 antibodies of the disclosure also include an antibody or antigen binding fragment thereof that cross-competes for binding to human CD71 and/or cynomolgus monkey CD71 to an anti-CD71 antibody comprising a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[00055] Suitable anti-CD71 antibodies of the disclosure also include an antibody or antigen binding fragment thereof that cross-competes for binding to human CD71 and/or cynomolgus monkey CD71 to an anti-CD71 antibody comprising a VH CDR1 sequence comprising the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); a VH CDR2 sequence comprising the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); a VH CDR3 sequence comprising the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); a VL CDR1 sequence comprising the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); a VL CDR2 sequence comprising the amino acid sequence STSNLAS (SEQ ID NO: 14); and a VL CDR3 sequence comprising the amino acid sequence QRRNYPYT (SEQ ID NO: 15).

[00056] In some embodiments, the anti-CD71 antibody of the disclosure comprises an isolated antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian CD71, wherein the AB specifically binds human CD71 and cynomolgus monkey CD71. In some embodiments, the antibody or antigen binding fragment thereof comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QRRNYPYT (SEQ ID NO: 15). In some embodiments, the antibody or antigen binding fragment thereof comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8.

[00057] In some embodiments, the isolated antibody or antigen binding fragment thereof binds to the same epitope on human CD71 and/or cynomolgus monkey CD71 as an isolated antibody that comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QRRNYPYT (SEQ ID NO: 15). In some embodiments, the isolated antibody or antigen binding fragment thereof binds to the same epitope on human CD71 and/or cynomolgus monkey CD71 as an isolated antibody that comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8. In some embodiments, the antigen binding fragment thereof is selected from the group consisting of a Fab fragment, a F(ab')₂ fragment, a scFv, a scAb, a dAb, a single domain heavy chain antibody, and a single domain light chain antibody. In some embodiments, the antibody or antigen-binding fragment thereof specifically binds human CD71.

[00058] In some embodiments, the isolated antibody or antigen binding fragment thereof cross-competes with an isolated antibody that comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence

SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QQRRNYPYT (SEQ ID NO: 15). In some embodiments, the isolated antibody or antigen binding fragment thereof cross-competes with an isolated antibody that comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8.

[00059] In some embodiments, antibody or antigen binding fragment thereof is conjugated to an agent. In some embodiments, the isolated antibody or antigen binding fragment thereof cross-competes with an isolated antibody that comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QQRRNYPYT (SEQ ID NO: 15). In some embodiments, the isolated antibody or antigen binding fragment thereof cross-competes with an isolated antibody that comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8. In some embodiments, the agent is a toxin or fragment thereof. In some embodiments, the agent is a microtubule inhibitor. In some embodiments, the agent is a nucleic acid damaging agent. In some embodiments, the agent is selected from the group consisting of a dolastatin or a derivative thereof, an auristatin or a derivative thereof, a maytansinoid or a derivative thereof, a duocarmycin or a derivative thereof, a calicheamicin or a derivative thereof, and a pyrrolobenzodiazepine or a derivative thereof. In some embodiments, the agent is auristatin E or a derivative thereof. In some embodiments, the agent is monomethyl auristatin E (MMAE). In some embodiments, the agent is monomethyl auristatin D (MMAD). In some embodiments, the agent is a maytansinoid selected from the group consisting of DM1 and DM4. In some embodiments, the agent is maytansinoid DM4. In some embodiments, the agent is duocarmycin. In some embodiments, the agent is conjugated to the AB via a linker. In some embodiments, the linker with which the agent is conjugated to the AB comprises an SPDB moiety, a vc moiety, or a PEG2-vc moiety. In some embodiments, the linker and toxin conjugated to the AB comprises an SPDB-DM4 moiety, a vc-MMAD moiety, a vc-MMAE moiety, vc-

duocarmycin, or a PEG2-vc-MMAD moiety. In some embodiments, the linker is a cleavable linker. In some embodiments, the linker is a non-cleavable linker. In some embodiments, the agent is a detectable moiety. In some embodiments, the detectable moiety is a diagnostic agent.

[00060] In some embodiments, the conjugated antibody comprises a conjugated antibody comprising: (a) an antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian CD71, wherein the AB comprises: (i) the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QQRRNYPYT (SEQ ID NO: 15), or (ii) a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8; (b) an agent conjugated to the AB, wherein the agent is selected from the group consisting of auristatin E, monomethyl auristatin F (MMAF), monomethyl auristatin E (MMAE), monomethyl auristatin D (MMAD), maytansinoid DM4, maytansinoid DM1, a pyrrolobenzodiazepine, a pyrrolobenzodiazepine dimer, and a duocarmycin.

[00061] The disclosure also provides activatable antibodies that include an antibody or antigen-binding fragment thereof that specifically binds CD71 coupled to a masking moiety (MM), such that coupling of the MM reduces the ability of the antibody or antigen-binding fragment thereof to bind CD71. In some embodiments, the MM is coupled via a sequence that includes a substrate for a protease, for example, a protease that is active in diseased tissue and/or a protease that is co-localized with CD71 at a treatment site in a subject. The activatable anti-CD71 antibodies provided herein, also referred to herein interchangeably as anti-CD71 activatable antibodies or CD71 activatable antibodies, are stable in circulation, activated at intended sites of therapy and/or diagnosis but not in normal, *e.g.*, healthy tissue or other tissue not targeted for treatment and/or diagnosis, and, when activated, exhibit binding to CD71 that is at least comparable to the corresponding, unmodified antibody, also referred to herein as the parental antibody.

[00062] The invention also provides methods of treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with aberrant expression and/or activity of CD71 in a subject using activatable antibodies that bind CD71,

particularly activatable antibodies that bind and neutralize or otherwise inhibit at least one biological activity of CD71 and/or CD71-mediated signaling.

[00063] The invention also provides methods of treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with the presence, growth, proliferation, metastasis, and/or activity of cells which are expressing CD71 or aberrantly expressing CD71 in a subject using activatable antibodies that bind CD71, particularly activatable antibodies that bind, target, neutralize, kill, or otherwise inhibit at least one biological activity of cells which are expressing or aberrantly expressing CD71.

[00064] The invention also provides methods of treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with the presence, growth, proliferation, metastasis, and/or activity of cells which are expressing CD71 in a subject using activatable antibodies that bind CD71, particularly activatable antibodies that bind, target, neutralize, kill, or otherwise inhibit at least one biological activity of cells which are expressing CD71.

[00065] The invention also provides methods of treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with the presence, growth, proliferation, metastasis, and/or activity of cells which are aberrantly expressing CD71 in a subject using activatable antibodies that bind CD71, particularly activatable antibodies that bind, target, neutralize, kill, or otherwise inhibit at least one biological activity of cells which are aberrantly expressing CD71.

[00066] The activatable antibodies in an activated state bind CD71 and include (i) an antibody or an antigen binding fragment thereof (AB) that specifically binds to CD71; (ii) a masking moiety (MM) that, when the activatable antibody is in an uncleaved state, inhibits the binding of the AB to CD71; and (c) a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease.

[00067] In some embodiments, the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM.

[00068] In some embodiments, the activatable antibody comprises a linking peptide between the MM and the CM.

[00069] In some embodiments, the activatable antibody comprises a linking peptide between the CM and the AB.

[00070] In some embodiments, the activatable antibody comprises a first linking peptide (LP1) and a second linking peptide (LP2), and wherein the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-LP1-CM-LP2-AB or AB-LP2-CM-LP1-MM. In some embodiments, the two linking peptides need not be identical to each other.

[00071] In some embodiments, at least one of LP1 or LP2 comprises an amino acid sequence selected from the group consisting of (GS)_n, (GGS)_n, (GSGGS)_n (SEQ ID NO: 339) and (GGGS)_n (SEQ ID NO: 340), where n is an integer of at least one.

[00072] In some embodiments, at least one of LP1 or LP2 comprises an amino acid sequence selected from the group consisting of GGSG (SEQ ID NO: 341), GGSGG (SEQ ID NO: 342), GSGSG (SEQ ID NO: 343), GSGGG (SEQ ID NO: 344), GGGSG (SEQ ID NO: 345), and GSSSG (SEQ ID NO: 346).

[00073] In some embodiments, LP1 comprises the amino acid sequence GSSGGSGGSGGSG (SEQ ID NO: 347), GSSGGSGGSGG (SEQ ID NO: 348), GSSGGSGGSGGS (SEQ ID NO: 349), GSSGGSGGSGGSGGGS (SEQ ID NO: 350), GSSGGSGGSG (SEQ ID NO: 351), or GSSGGSGGSGS (SEQ ID NO: 352).

[00074] In some embodiments, LP2 comprises the amino acid sequence GSS, GGS, GGGS (SEQ ID NO: 353), GSSGT (SEQ ID NO: 354) or GSSG (SEQ ID NO: 355).

[00075] In some embodiments, the AB has a dissociation constant of about 100 nM or less for binding to mammalian CD71. In some embodiments, the AB has a dissociation constant of about 10 nM or less for binding to mammalian CD71. In some embodiments, the AB has a dissociation constant of about 5 nM or less for binding to CD71. In some embodiments, the AB has a dissociation constant of about 1 nM or less for binding to CD71. In some embodiments, the AB has a dissociation constant of about 0.5 nM or less for binding to CD71. In some embodiments, the AB has a dissociation constant of about 0.1 nM or less for binding to CD71. In some embodiments, the AB has a dissociation constant of 0.01 nM to 100 nM, 0.01 nM to 10 nM, 0.01 nM to 5 nM, 0.01 nM to 1 nM, 0.01 to 0.5 nM, 0.01 nm to 0.1 nM, 0.01 nm to 0.05 nM, 0.05 nM to 100 nM, 0.05 nM to 10 nM, 0.05 nM to 5 nM, 0.05 nM to 1 nM, 0.05 to 0.5 nM, 0.05 nm to 0.1 nM, 0.1 nM to 100 nM, 0.1 nM to 10 nM, 0.1 nM to 5 nM, 0.1 nM to 1 nM, 0.1 to 0.5 nM, 0.5 nM to 100 nM, 0.5 nM to 10 nM, 0.5 nM to 5 nM, 0.5 nM to 1 nM, 1 nM to 100 nM, 1 nM to 10 nM, 1 nM to 5 nM, 5 nM to 100 nM, 5 nM to 10 nM, or 10 nM to 100 nM, for binding to mammalian CD71.

[00076] In some embodiments, the activatable antibody includes an antibody or antigen-binding fragment thereof (AB) that specifically binds CD71. In some embodiments, the antibody or antigen-binding fragment thereof that binds CD71 is a monoclonal antibody, domain antibody, single chain, Fab fragment, a $F(ab')_2$ fragment, a scFv, a scAb, a dAb, a single domain heavy chain antibody, or a single domain light chain antibody. In some embodiments, such an antibody or antigen-binding fragment thereof that binds CD71 is a mouse, other rodent, chimeric, humanized or fully human monoclonal antibody.

[00077] In some embodiments, the activatable antibody in an uncleaved state specifically binds to the mammalian CD71 with a dissociation constant less than or equal to 1 nM, less than or equal to 5 nM, less than or equal to 10 nM, less than or equal to 15 nM, less than or equal to 20 nM, less than or equal to 25 nM, less than or equal to 50 nM, less than or equal to 100 nM, less than or equal to 150 nM, less than or equal to 250 nM, less than or equal to 500 nM, less than or equal to 750 nM, less than or equal to 1000 nM, and/or less than or equal to 2000 nM.

[00078] In some embodiments, the activatable antibody in an uncleaved state specifically binds to the mammalian CD71 with a dissociation constant in the range of 1 nM to 2000 nM, 1 nM to 1000 nM, 1 nM to 750 nM, 1 nM to 500 nM, 1 nM to 250 nM, 1 nM to 150 nM, 1 nM to 100 nM, 1 nM to 50 nM, 1 nM to 25 nM, 1 nM to 15 nM, 1 nM to 10 nM, 1 nM to 5 nM, 5 nM to 2000 nM, 5 nM to 1000 nM, 5 nM to 750 nM, 5 nM to 500 nM, 5 nM to 250 nM, 5 nM to 150 nM, 5 nM to 100 nM, 5 nM to 50 nM, 5 nM to 25 nM, 5 nM to 15 nM, 5 nM to 10 nM, 10 nM to 2000 nM, 10 nM to 1000 nM, 10 nM to 750 nM, 10 nM to 500 nM, 10 nM to 250 nM, 10 nM to 150 nM, 10 nM to 100 nM, 10 nM to 50 nM, 10 nM to 25 nM, 10 nM to 15 nM, 15 nM to 2000 nM, 15 nM to 1000 nM, 15 nM to 750 nM, 15 nM to 500 nM, 15 nM to 250 nM, 15 nM to 150 nM, 15 nM to 100 nM, 15 nM to 50 nM, 15 nM to 25 nM, 25 nM to 2000 nM, 25 nM to 1000 nM, 25 nM to 750 nM, 25 nM to 500 nM, 25 nM to 250 nM, 25 nM to 150 nM, 25 nM to 100 nM, 25 nM to 50 nM, 50 nM to 2000 nM, 50 nM to 1000 nM, 50 nM to 750 nM, 50 nM to 500 nM, 50 nM to 250 nM, 50 nM to 150 nM, 50 nM to 100 nM, 100 nM to 2000 nM, 100 nM to 1000 nM, 100 nM to 750 nM, 100 nM to 500 nM, 100 nM to 250 nM, 100 nM to 150 nM, 150 nM to 2000 nM, 150 nM to 1000 nM, 150 nM to 750 nM, 150 nM to 500 nM, 150 nM to 250 nM, 250 nM to 2000 nM, 250 nM to 1000 nM, 250 nM to 750 nM, 250 nM to 500 nM, 500 nM to 2000 nM, 500 nM to 1000 nM, 500 nM to 750 nM, 500 nM to 500 nM, 500 nM to 250 nM,

500 nM to 150 nM, 500 nM to 100 nM, 500 nM to 50 nM, 750 nM to 2000 nM, 750 nM to 1000 nM, or 1000 nM to 2000 nM.

[00079] In some embodiments, the activatable antibody in an activated state specifically binds to the mammalian CD71 with a dissociation constant is less than or equal to 0.01 nM, 0.05 nM, 0.1 nM, 0.5 nM, 1 nM, 5 nM, or 10 nM.

[00080] In some embodiments, the activatable antibody in an activated state specifically binds to the mammalian CD71 with a dissociation constant in the range of 0.01 nM to 100 nM, 0.01 nM to 10 nM, 0.01 nM to 5 nM, 0.01 nM to 1 nM, 0.01 to 0.5 nM, 0.01 nm to 0.1 nM, 0.01 nm to 0.05 nM, 0.05 nM to 100 nM, 0.05 nM to 10 nM, 0.05 nM to 5 nM, 0.05 nM to 1 nM, 0.05 to 0.5 nM, 0.05 nm to 0.1 nM, 0.1 nM to 100 nM, 0.1 nM to 10 nM, 0.1 nM to 5 nM, 0.1 nM to 1 nM, 0.1 to 0.5 nM, 0.5 nM to 100 nM, 0.5 nM to 10 nM, 0.5 nM to 5 nM, 0.5 nM to 1 nM, 1 nM to 100 nM, 1 nM to 10 nM, 1 nM to 5 nM, 5 nM to 100 nM, 5 nM to 10 nM, or 10 nM to 100 nM.

[00081] In some embodiments, the mammalian CD71 is selected from the group consisting of a human CD71, a murine CD71, a rat CD71, and a cynomolgus monkey CD71. In some embodiments, the AB specifically binds to human CD71, murine CD71 or cynomolgus monkey CD71 with a dissociation constant of less than 1 nM. In some embodiments, the mammalian CD71 is a human CD71.

[00082] In some embodiments, the AB has one or more of the following characteristics: (a) the AB specifically binds to human CD71; and (b) the AB specifically binds to human CD71 and cynomolgus monkey CD71.

[00083] In some embodiments, the AB has one or more of the following characteristics: (a) the AB specifically binds human CD71 and cynomolgus monkey CD71; (b) the AB inhibits binding of transferrin to mammalian CD71; (c) the AB inhibits binding of human transferrin to human CD71; and (d) the AB inhibits binding of cynomolgus monkey transferrin to cynomolgus monkey CD71.

[00084] In some embodiments, the AB blocks the ability of a natural ligand to bind to the mammalian CD71 with an EC_{50} less than or equal to 5 nM, less than or equal to 10 nM, less than or equal to 50 nM, less than or equal to 100 nM, less than or equal to 500 nM, and/or less than or equal to 1000 nM. In some embodiments, the AB blocks the ability of a transferrin to bind to the mammalian CD71 with an EC_{50} less than or equal to 5 nM, less than or equal to 10 nM, less than or equal to 50 nM, less than or equal to 100 nM, less than

or equal to 500 nM, and/or less than or equal to 1000 nM. In some embodiments, the natural ligand of CD71 is transferrin.

[00085] In some embodiments, the AB blocks the ability of a natural ligand to bind to the mammalian CD71 with an EC₅₀ of 5 nM to 1000 nM, 5 nM to 500 nM, 5 nM to 100 nM, 5 nM to 50 nM, 5 nM to 10 nM, 10 nM to 1000 nM, 10 nM to 500 nM, 10 nM to 100 nM, 10 nM to 50 nM, 50 nM to 1000 nM, 50 nM to 500 nM, 50 nM to 100 nM, 100 nM to 1000 nM, 100 nM to 500 nM, 500 nM to 1000 nM. In some embodiments, the AB blocks the ability of a transferrin to bind to the mammalian CD71 with an EC₅₀ of 5 nM to 1000 nM, 5 nM to 500 nM, 5 nM to 100 nM, 5 nM to 50 nM, 5 nM to 10 nM, 10 nM to 1000 nM, 10 nM to 500 nM, 10 nM to 100 nM, 10 nM to 50 nM, 50 nM to 1000 nM, 50 nM to 500 nM, 50 nM to 100 nM, 100 nM to 1000 nM, 100 nM to 500 nM, 500 nM to 1000 nM. In some embodiments, the natural ligand of CD71 is transferrin.

[00086] In some embodiments, the AB of the present disclosure inhibits or reduces the growth, proliferation, and/or metastasis of cells expressing mammalian CD71. Without intending to be bound by any theory, the AB of the present disclosure may inhibit or reduce the growth, proliferation, and/or metastasis of cells expressing mammalian CD71 by specifically binding to CD71 and inhibiting, blocking, and/or preventing the binding of a natural ligand to mammalian CD71. In some embodiments, the natural ligand of mammalian CD71 is transferrin.

[00087] In some embodiments, the activatable antibody comprises a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1, and 3-5. In some embodiments, the activatable antibody comprises a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 3-5.

[00088] In some embodiments, the activatable antibody comprises a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8. In some embodiments, the activatable antibody comprises a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00089] In some embodiments, the activatable antibody comprises a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[00090] In some embodiments, the activatable antibody comprises a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 3-5,

and a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00091] In some embodiments, the activatable antibody comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5. In some embodiments, the activatable antibody comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 3-5.

[00092] In some embodiments, the activatable antibody comprises a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8. In some embodiments, the activatable antibody comprises a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00093] In some embodiments, the activatable antibody comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[00094] In some embodiments, the activatable antibody comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 3-5, and a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[00095] In some embodiments, the activatable antibody comprises a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3)

sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2) sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence comprising the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); a VH CDR2 sequence comprising the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); a VH CDR3 sequence comprising the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); a VL CDR1 sequence comprising the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); a VL CDR2 sequence comprising the amino acid sequence STSNLAS (SEQ ID NO: 14); and a VL CDR3 sequence comprising the amino acid sequence QRRNYPYT (SEQ ID NO: 15).

[00096] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR1 sequence comprising the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); a VH CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR2 sequence comprising the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); a VH CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR3 sequence comprising the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); a VL CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR1 sequence comprising the amino acid sequence ASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); a VL CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR2 sequence comprising the amino acid sequence STSNLAS (SEQ ID NO: 14); and a VL CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR3 sequence comprising the amino acid sequence QRRNYPYT (SEQ ID NO: 15).

[00097] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR1 sequence comprises the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence comprises the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence comprises the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence comprises the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence comprises the amino acid sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence comprises the amino acid sequence QQRRNYPYT (SEQ ID NO: 15).

[00098] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR1 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence QQRRNYPYT (SEQ ID NO: 15).

[00099] In some embodiments, the AB of the activatable anti-CD71 antibody comprises a heavy chain variable region amino acid sequence selected from the group consisting of the heavy chain variable region sequences shown in Table 12. In some embodiments, the AB of the activatable anti-CD71 antibody comprises a light chain variable region amino acid sequence selected from the group consisting of the light chain variable region sequences shown in Table 12. In some embodiments, the AB of the

activatable anti-CD71 antibody comprises a heavy chain variable region amino acid sequence selected from the group consisting of the heavy chain variable region sequences shown in Table 12 and a light chain variable region amino acid sequence selected from the group consisting of the light chain variable region sequences shown in Table 12.

[000100] In some embodiments, the AB of the activatable anti-CD71 antibody comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the heavy chain variable region sequences shown in Table 12. In some embodiments, the AB of the activatable anti-CD71 antibody comprises a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the light chain variable region sequences shown in Table 12. In some embodiments, the AB of the activatable anti-CD71 antibody comprises a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the heavy chain variable region sequences shown in Table 12 and a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the light chain variable region sequences shown in Table 12.

[000101] In some embodiments, the activatable antibody comprises a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3) sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2) sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence shown in Table 13; a VH CDR2 sequence shown in Table 13; a VH CDR3 sequence shown in Table 13; a VL CDR1 sequence shown in Table 13; a VL CDR2 sequence shown in Table 13; and a VL CDR3 sequence shown in Table 13.

[000102] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR1 sequence shown in Table 13; a VH CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR2 sequence shown in Table 13; a VH CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR3 sequence shown in Table 13; a VL CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR1 sequence shown in Table 13; a VL CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR2 sequence shown in Table 13; and a VL CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR3 sequence shown in Table 13.

[000103] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 13.

[000104] In some embodiments, the activatable antibody comprises a heavy chain that comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein the combination is a combination of the three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 13.

[000105] In some embodiments, the activatable antibody comprises a light chain that comprises a combination of a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the three light chain CDR sequences (VL CDR1, VL CDR2, VL CDR3) shown in a single row in Table 13.

[000106] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%,

97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 13.

[000107] In some embodiments, the activatable antibody comprises a heavy chain variable region that comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 13.

[000108] In some embodiments, the activatable antibody comprises a light chain variable region that comprises a combination of a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of three light chain CDR sequences (VL CDR1, VL CDR2, VL CDR3) shown in a single row in Table 13.

[000109] In some embodiments, the MM has a dissociation constant for binding to the AB which is greater than the dissociation constant of the AB to CD71.

[000110] In some embodiments, the MM has a dissociation constant for binding to the AB which is no more than the dissociation constant of the AB to CD71.

[000111] In some embodiments, the MM has a dissociation constant for binding to the AB which is less than the dissociation constant of the AB to CD71.

[000112] In some embodiments, the dissociation constant (K_d) of the MM towards the AB is no more than 2, 3, 4, 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 50,000, 100,000, 500,000, 1,000,000, 5,000,000, 10,000,000, 50,000,000 times or greater, or between 1-5, 5-10, 10-100, 10-1,000, 10-10,000, 10-100,000, 10-1,000,000, 10-10,000,000, 100-1,000, 100-10,000, 100-100,000, 100-1,000,000, 100-10,000,000, 1,000-10,000, 1,000-100,000, 1,000-1,000,000, 1,000-10,000,000, 10,000-100,000, 10,000-1,000,000, 10,000-10,000,000, 100,000-1,000,000, or 100,000-10,000,000 times or greater than the dissociation constant of the AB towards the target.

[000113] In some embodiments, the MM does not interfere or compete with the AB for binding to CD71 when the activatable antibody is in a cleaved state.

[000114] In some embodiments, the MM is a polypeptide of about 2 to 40 amino acids in length. In some embodiments, the MM is a polypeptide of up to about 40 amino acids in length.

[000115] In some embodiments, the MM polypeptide sequence is different from that of CD71. In some embodiments, the MM polypeptide sequence is no more than 50% identical to any natural binding partner of the AB. In some embodiments, the MM polypeptide sequence is different from that of CD71 and is no more than 40%, 30%, 25%, 20%, 15%, or 10% identical to any natural binding partner of the AB.

[000116] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind CD71 such that the dissociation constant (K_d) of the AB when coupled to the MM towards CD71 is at least two times greater than the K_d of the AB when not coupled to the MM towards CD71.

[000117] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind CD71 such that the dissociation constant (K_d) of the AB when coupled to the MM towards CD71 is at least five times greater than the K_d of the AB when not coupled to the MM towards CD71.

[000118] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind CD71 such that the dissociation constant (K_d) of the AB when coupled to the MM towards CD71 is at least 10 times greater than the K_d of the AB when not coupled to the MM towards CD71.

[000119] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind CD71 such that the dissociation constant (K_d) of the AB when coupled to the MM towards CD71 is at least 20 times greater than the K_d of the AB when not coupled to the MM towards CD71.

[000120] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind CD71 such that the dissociation constant (K_d) of the AB when coupled to the MM towards CD71 is at least 40 times greater than the K_d of the AB when not coupled to the MM towards CD71.

[000121] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind CD71 such that the dissociation constant (K_d) of the AB when coupled to the MM towards CD71 is at least 100 times greater than the K_d of the AB when not coupled to the MM towards CD71.

[000122] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind CD71 such that the dissociation constant (K_d) of the AB when coupled to the MM towards CD71 is at least 1000 times greater than the K_d of the AB when not coupled to the MM towards CD71.

[000123] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind CD71 such that the dissociation constant (K_d) of the AB when coupled to the MM towards CD71 is at least 10,000 times greater than the K_d of the AB when not coupled to the MM towards CD71.

[000124] In some embodiments, in the presence of CD71, the MM reduces the ability of the AB to bind CD71 by at least 90% when the CM is uncleaved, as compared to when the CM is cleaved when assayed *in vitro* using a target displacement assay such as, for example, the assay described in PCT Publication No. WO 2010/081173, the contents of which are hereby incorporated by reference in their entirety.

[000125] In some embodiments, MM comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 16-295 and 297-314.

[000126] In some embodiments, the protease that cleaves the CM is active, e.g., up-regulated or otherwise unregulated, in diseased tissue, and the protease cleaves the CM in the activatable antibody when the activatable antibody is exposed to the protease.

[000127] In some embodiments, the protease is co-localized with CD71 in a tissue, and the protease cleaves the CM in the activatable antibody when the activatable antibody is exposed to the protease.

[000128] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to CD71 is reduced to occur with a dissociation constant that is at least twofold greater than the dissociation constant of an unmodified AB binding to CD71, whereas in the cleaved state (i.e., when the activatable antibody is in the cleaved state), the AB binds CD71.

[000129] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to CD71 is reduced to occur with a dissociation constant that is at least fivefold greater than the dissociation constant of an unmodified AB binding to CD71, whereas in the cleaved state (i.e., when the activatable antibody is in the cleaved state), the AB binds CD71.

[000130] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to CD71 is reduced to occur with a dissociation constant that is at least 10-fold greater than the dissociation constant of an unmodified AB binding to CD71, whereas in the cleaved state (i.e., when the activatable antibody is in the cleaved state), the AB binds CD71.

[000131] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to CD71 is reduced to occur with a dissociation constant that is at least 20-fold greater than the dissociation constant of an unmodified AB binding to CD71, whereas in the cleaved state (i.e., when the activatable antibody is in the cleaved state), the AB binds CD71.

[000132] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to CD71 is reduced to occur with a dissociation constant that is at least 40-fold greater than the dissociation constant of an unmodified AB binding to CD71, whereas in the cleaved state, the AB binds CD71.

[000133] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to CD71 is reduced to occur with a dissociation constant that is at least 50-fold greater than the dissociation constant of an unmodified AB binding to CD71, whereas in the cleaved state, the AB binds CD71.

[000134] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to CD71 is reduced to occur with a dissociation constant that is at least 100-fold greater than the dissociation constant of an unmodified AB binding to CD71, whereas in the cleaved state, the AB binds CD71.

[000135] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to CD71 is reduced to occur with a dissociation constant that is at least 200-fold greater than the dissociation constant of an unmodified AB binding to CD71, whereas in the cleaved state, the AB binds CD71.

[000136] In some embodiments, the CM is a polypeptide of up to 15 amino acids in length.

[000137] In some embodiments, the CM is a polypeptide that includes a first cleavable moiety (CM1) that is a substrate for at least one matrix metalloprotease (MMP) and a second cleavable moiety (CM2) that is a substrate for at least one serine protease (SP). In some embodiments, each of the CM1 substrate sequence and the CM2 substrate sequence of the CM1-CM2 substrate is independently a polypeptide of up to 15 amino acids in length.

[000138] In some embodiments, the CM is a substrate for at least one protease that is or is believed to be up-regulated or otherwise unregulated in cancer.

[000139] In some embodiments, the CM is a substrate for at least one protease selected from the group consisting of a matrix metalloprotease (MMP), thrombin, a neutrophil elastase, a cysteine protease, legumain, and a serine protease, such as matriptase (MT-SP1), and urokinase (uPA). Without being bound by theory, it is believed that these proteases are up-regulated or otherwise unregulated in at least one of cancer.

[000140] Exemplary substrates include but are not limited to substrates cleavable by one or more of the following enzymes or proteases listed in Table 4.

[000141] In some embodiments, the CM is selected for use with a specific protease, for example a protease that is known to be co-localized with the target of the activatable antibody.

[000142] In some embodiments, the CM is a substrate for at least one MMP. Examples of MMPs include the MMPs listed in the Table 4. In some embodiments, the CM is a substrate for a protease selected from the group consisting of MMP 9, MMP14, MMP1, MMP3, MMP13, MMP17, MMP11, and MMP19. In some embodiments the CM is a substrate for MMP9. In some embodiments, the CM is a substrate for MMP14.

[000143] In some embodiments, the CM is a substrate that includes the sequence TGRGPSWV (SEQ ID NO: 356); SARGPSRW (SEQ ID NO: 357); TARGPSFK (SEQ ID NO: 358); LSGRSDNH (SEQ ID NO: 359); GGWHTGRN (SEQ ID NO: 360); HTGRSGAL (SEQ ID NO: 361); PLTGRSGG (SEQ ID NO: 362); AARGPAIH (SEQ ID NO: 363); RGPAFNPM (SEQ ID NO: 364); SSRGPAYL (SEQ ID NO: 365); RGPATPIM (SEQ ID NO: 366); RGPA (SEQ ID NO: 367); GGQPSGMWGW (SEQ ID NO: 368); FPRPLGITGL (SEQ ID NO: 369); VHMPLGFLGP (SEQ ID NO: 370); SPLTGRSG (SEQ ID NO: 371); SAGFSLPA (SEQ ID NO: 372); LAPLGLQRR (SEQ ID NO: 373); SGGPLGVR (SEQ ID NO: 374); PLGL (SEQ ID NO: 375); LSGRSGNH (SEQ ID NO:

789); SGRSANPRG (SEQ ID NO: 790); LSGRSDDH (SEQ ID NO: 791); LSGRSDIH (SEQ ID NO: 792); LSGRSDQH (SEQ ID NO: 793); LSGRSDTH (SEQ ID NO: 794); LSGRSDYH (SEQ ID NO: 795); LSGRSDNP (SEQ ID NO: 796); LSGRSANP (SEQ ID NO: 797); LSGRSANI (SEQ ID NO: 798); LSGRSDNI (SEQ ID NO: 799); MIAPVAYR (SEQ ID NO: 800); RPSPMWAY (SEQ ID NO: 801); WATPRPMR (SEQ ID NO: 802); FRLLDWQW (SEQ ID NO: 803); ISSGL (SEQ ID NO: 804); ISSGLLS (SEQ ID NO: 805); and/or ISSGLL (SEQ ID NO: 806).

[000144] In some embodiments, the CM comprises the amino acid sequence LSGRSDNH (SEQ ID NO: 359). In some embodiments, the CM comprises the amino acid sequence TGRGPSWV (SEQ ID NO: 356). In some embodiments, the CM comprises the amino acid sequence PLTGRSGG (SEQ ID NO: 362). In some embodiments, the CM comprises the amino acid sequence GGQPSGMWGW (SEQ ID NO: 368). In some embodiments, the CM comprises the amino acid sequence FPRPLGITGL (SEQ ID NO: 369). In some embodiments, the CM comprises the amino acid sequence VHMPGLGFLGP (SEQ ID NO: 370). In some embodiments, the CM comprises the amino acid sequence PLGL (SEQ ID NO: 375). In some embodiments, the CM comprises the amino acid sequence SARGPSRW (SEQ ID NO: 357). In some embodiments, the CM comprises the amino acid sequence TARGPSFK (SEQ ID NO: 358). In some embodiments, the CM comprises the amino acid sequence GGWHTGRN (SEQ ID NO: 360). In some embodiments, the CM comprises the amino acid sequence HTGRSGAL (SEQ ID NO: 361). In some embodiments, the CM comprises the amino acid sequence AARGPAIH (SEQ ID NO: 363). In some embodiments, the CM comprises the amino acid sequence RGPAFNPM (SEQ ID NO: 364). In some embodiments, the CM comprises the amino acid sequence SSRGPAYL (SEQ ID NO: 365). In some embodiments, the CM comprises the amino acid sequence RGPATPIM (SEQ ID NO: 366). In some embodiments, the CM comprises the amino acid sequence RGPA (SEQ ID NO: 367). In some embodiments, the CM comprises the amino acid sequence LSGRSGNH (SEQ ID NO: 789). In some embodiments, the CM comprises the amino acid sequence SGRSANPRG (SEQ ID NO: 790). In some embodiments, the CM comprises the amino acid sequence LSGRSDDH (SEQ ID NO: 791). In some embodiments, the CM comprises the amino acid sequence LSGRSDIH (SEQ ID NO: 792). In some embodiments, the CM comprises the amino acid sequence LSGRSDQH (SEQ ID NO: 793). In some embodiments, the CM comprises the amino acid sequence LSGRSDTH (SEQ ID NO: 794). In some embodiments, the CM comprises the amino acid

sequence LSGRSDYH (SEQ ID NO: 795). In some embodiments, the CM comprises the amino acid sequence LSGRSDNP (SEQ ID NO: 796). In some embodiments, the CM comprises the amino acid sequence LSGRSANP (SEQ ID NO: 797). In some embodiments, the CM comprises the amino acid sequence LSGRSANI (SEQ ID NO: 798). In some embodiments, the CM comprises the amino acid sequence LSGRSDNI (SEQ ID NO: 799). In some embodiments, the CM comprises the amino acid sequence MIAPVAYR (SEQ ID NO: 800). In some embodiments, the CM comprises the amino acid sequence RPSPMWAY (SEQ ID NO: 801). In some embodiments, the CM comprises the amino acid sequence WATPRPMR (SEQ ID NO: 802). In some embodiments, the CM comprises the amino acid sequence FRLLDWQW (SEQ ID NO: 803). In some embodiments, the CM comprises the amino acid sequence ISSGL (SEQ ID NO: 804). In some embodiments, the CM comprises the amino acid sequence ISSGLLS (SEQ ID NO: 805). In some embodiments, the CM comprises the amino acid sequence and/or ISSGLL (SEQ ID NO: 806).

[000145] In some embodiments, the CM is a substrate for an MMP and includes the sequence ISSGLSS (SEQ ID NO: 376); QNQALRMA (SEQ ID NO: 377); AQNLLGMV (SEQ ID NO: 378); STFPFGMF (SEQ ID NO: 379); PVGYTSSL (SEQ ID NO: 380); DWLYWPGI (SEQ ID NO: 381); ISSGLLSS (SEQ ID NO: 382); LKAAPRWA (SEQ ID NO: 383); GPSHLVLT (SEQ ID NO: 384); LPGGLSPW (SEQ ID NO: 385); MGLFSEAG (SEQ ID NO: 386); SPLPLRVP (SEQ ID NO: 387); RMHLRSLG (SEQ ID NO: 388); LAAPLGLL (SEQ ID NO: 389); AVGLLAPP (SEQ ID NO: 390); LLAPSHRA (SEQ ID NO: 391); and/or PAGLWLDP (SEQ ID NO: 392).

[000146] In some embodiments, the CM comprises the amino acid sequence ISSGLSS (SEQ ID NO: 376). In some embodiments, the CM comprises the amino acid sequence QNQALRMA (SEQ ID NO: 377). In some embodiments, the CM comprises the amino acid sequence AQNLLGMV (SEQ ID NO: 378). In some embodiments, the CM comprises the amino acid sequence STFPFGMF (SEQ ID NO: 379). In some embodiments, the CM comprises the amino acid sequence PVGYTSSL (SEQ ID NO: 380). In some embodiments, the CM comprises the amino acid sequence DWLYWPGI (SEQ ID NO: 381). In some embodiments, the CM comprises the amino acid sequence ISSGLLSS (SEQ ID NO: 382). In some embodiments, the CM comprises the amino acid sequence LKAAPRWA (SEQ ID NO: 383). In some embodiments, the CM comprises the amino acid sequence GPSHLVLT (SEQ ID NO: 384). In some embodiments, the CM comprises the amino acid sequence LPGGLSPW (SEQ ID NO: 385). In some embodiments, the CM comprises the amino acid

sequence MGLFSEAG (SEQ ID NO: 386). In some embodiments, the CM comprises the amino acid sequence SPLPLRVP (SEQ ID NO: 387). In some embodiments, the CM comprises the amino acid sequence RMHLRSLG (SEQ ID NO: 388). In some embodiments, the CM comprises the amino acid sequence LAAPLGLL (SEQ ID NO: 389). In some embodiments, the CM comprises the amino acid sequence AVGLLAPP (SEQ ID NO: 390). In some embodiments, the CM comprises the amino acid sequence LLAPSHRA (SEQ ID NO: 391). In some embodiments, the CM comprises the amino acid sequence PAGLWLDP (SEQ ID NO: 392).

[000147] In some embodiments, the CM is a substrate for thrombin. In some embodiments, the CM is a substrate for thrombin and includes the sequence GPRSFGL (SEQ ID NO: 393) or GPRSFG (SEQ ID NO: 394). In some embodiments, the CM comprises the amino acid sequence GPRSFGL (SEQ ID NO: 393). In some embodiments, the CM comprises the amino acid sequence GPRSFG (SEQ ID NO: 394).

[000148] In some embodiments, the CM comprises an amino acid sequence selected from the group consisting of NTLSGRSENHSG (SEQ ID NO: 395); NTLSGRSGNHGS (SEQ ID NO: 396); TSTSGRSANPRG (SEQ ID NO: 397); TSGRSANP (SEQ ID NO: 398); VAGRSMRP (SEQ ID NO: 399); VVPEGRRS (SEQ ID NO: 400); ILPRSPAF (SEQ ID NO: 401); MVLGRSLL (SEQ ID NO: 402); QGRAITFI (SEQ ID NO: 403); SPRSIMLA (SEQ ID NO: 404); and SMLRSMPL (SEQ ID NO: 405).

[000149] In some embodiments, the CM comprises the amino acid sequence NTLSGRSENHSG (SEQ ID NO: 395). In some embodiments, the CM comprises the amino acid sequence NTLSGRSGNHGS (SEQ ID NO: 396). In some embodiments, the CM comprises the amino acid sequence TSTSGRSANPRG (SEQ ID NO: 397). In some embodiments, the CM comprises the amino acid sequence TSGRSANP (SEQ ID NO: 398). In some embodiments, the CM comprises the amino acid sequence VAGRSMRP (SEQ ID NO: 399). In some embodiments, the CM comprises the amino acid sequence VVPEGRRS (SEQ ID NO: 400). In some embodiments, the CM comprises the amino acid sequence ILPRSPAF (SEQ ID NO: 401). In some embodiments, the CM comprises the amino acid sequence MVLGRSLL (SEQ ID NO: 402). In some embodiments, the CM comprises the amino acid sequence QGRAITFI (SEQ ID NO: 403). In some embodiments, the CM comprises the amino acid sequence SPRSIMLA (SEQ ID NO: 404). In some embodiments, the CM comprises the amino acid sequence SMLRSMPL (SEQ ID NO: 405).

[000150] In some embodiments, the CM is a substrate for a neutrophil elastase. In some embodiments, the CM is a substrate for a serine protease. In some embodiments, the CM is a substrate for uPA. In some embodiments, the CM is a substrate for legumain. In some embodiments, the CM is a substrate for matriptase. In some embodiments, the CM is a substrate for a cysteine protease. In some embodiments, the CM is a substrate for a cysteine protease, such as a cathepsin.

[000151] In some embodiments, the CM is a CM1-CM2 substrate and includes the sequence ISSGLLSGRSDNH (SEQ ID NO: 406); ISSGLLSSGGSGGSLSGRSDNH (SEQ ID NO: 407); AVGLLAPPGGTSTSGRSANPRG (SEQ ID NO: 408); TSTSGRSANPRGGGAVGLLAPP (SEQ ID NO: 409); VHMPGLGFLPGGTSTSGRSANPRG (SEQ ID NO: 410); TSTSGRSANPRGGGVHMPGLGFLGP (SEQ ID NO: 411); AVGLLAPPGGLSGRSDNH (SEQ ID NO: 412); LSGRSDNHGGAVGLLAPP (SEQ ID NO: 413); VHMPGLGFLPGGLSGRSDNH (SEQ ID NO: 414); LSGRSDNHGGVHMPGLGFLGP (SEQ ID NO: 415); LSGRSDNHGGSGGSISSGLLSS (SEQ ID NO: 416); LSGRSGNHGGSGGSISSGLLSS (SEQ ID NO: 417); ISSGLLSSGGSGGSLSGRSGNH (SEQ ID NO: 418); LSGRSDNHGGSGGSQLNRMA (SEQ ID NO: 419); QNQALRMAGGSGGSLSGRSDNH (SEQ ID NO: 420); LSGRSGNHGGSGGSQLNRMA (SEQ ID NO: 421); QNQALRMAGGSGGSLSGRSGNH (SEQ ID NO: 422); ISSGLLSGRSGNH (SEQ ID NO: 423); ISSGLLSGRSANPRG (SEQ ID NO: 680); AVGLLAPPTSGRSANPRG (SEQ ID NO: 681); AVGLLAPPSGRSANPRG (SEQ ID NO: 682); ISSGLLSGRSDDH (SEQ ID NO: 683); ISSGLLSGRSDIH (SEQ ID NO: 684); ISSGLLSGRSDQH (SEQ ID NO: 685); ISSGLLSGRSDTH (SEQ ID NO: 686); ISSGLLSGRSDYH (SEQ ID NO: 687); ISSGLLSGRSDNP (SEQ ID NO: 688); ISSGLLSGRSANP (SEQ ID NO: 689); ISSGLLSGRSANI (SEQ ID NO: 690); AVGLLAPPGGLSGRSDDH (SEQ ID NO: 691); AVGLLAPPGGLSGRSDIH (SEQ ID NO: 692); AVGLLAPPGGLSGRSDQH (SEQ ID NO: 693); AVGLLAPPGGLSGRSDTH (SEQ ID NO: 694); AVGLLAPPGGLSGRSDYH (SEQ ID NO: 695); AVGLLAPPGGLSGRSDNP (SEQ ID NO: 696); AVGLLAPPGGLSGRSANP (SEQ ID NO: 697); AVGLLAPPGGLSGRSANI (SEQ ID NO: 698); ISSGLLSGRSDNI (SEQ ID NO: 713); AVGLLAPPGGLSGRSDNI (SEQ ID NO: 714); GLSGRSDNHGGAVGLLAPP (SEQ ID NO: 807); and/or GLSGRSDNHGGVHMPGLGFLGP (SEQ ID NO: 808).

[000152] In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDNH (SEQ ID NO: 406), which is also referred to herein as substrate 2001. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLSSGGSGGSLSGRSDNH (SEQ ID NO: 407), which is also referred to herein as substrate 1001/LP'/0001, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 1037). In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGTSTSGRSANPRG (SEQ ID NO: 408), which is also referred to herein as substrate 2015 and/or substrate 1004/LP'/0003, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence TSTSGRSANPRGGGAVGLLAPP (SEQ ID NO: 409), which is also referred to herein as substrate 0003/LP'/1004, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence VHMPLGFLGPGGTSTSGRSANPRG (SEQ ID NO: 410), which is also referred to herein as substrate 1003/LP'/0003, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence TSTSGRSANPRGGGVHMPLGFLGP (SEQ ID NO: 411), which is also referred to herein as substrate 0003/LP'/1003, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDNH (SEQ ID NO: 412), which is also referred to herein as substrate 3001 and/or substrate 1004/LP'/0001, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence LSGRSDNHGGAVGLLAPP (SEQ ID NO: 413), which is also referred to herein as substrate 0001/LP'/1004, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence VHMPLGFLGPGGLSGRSDNH (SEQ ID NO: 414), which is also referred to herein as substrate 1003/LP'/0001, wherein LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence LSGRSDNHGGVHMPLGFLGP (SEQ ID NO: 415), which is also referred to herein as substrate 0001/LP'/1003, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence LSGRSDNHGGSGGSISSGLLSS (SEQ ID NO: 416), which is also referred to herein as substrate 0001/LP'/1001, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 1037). In some embodiments,

the CM1-CM2 substrate includes the sequence LSGRSGNHGGSGGSISSGLLSS (SEQ ID NO: 417), which is also referred to herein as substrate 0002/LP'/1001, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 1037). In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSSGGSGGSLSGRSGNH (SEQ ID NO: 418), which is also referred to herein as substrate 1001/LP'/0002, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 1037). In some embodiments, the CM1-CM2 substrate includes the sequence LSGRSDNHGGSGGSGQNQALRMA (SEQ ID NO: 419), which is also referred to herein as substrate 0001/LP'/1002, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 1037). In some embodiments, the CM1-CM2 substrate includes the sequence QNQALRMAGGSGGSLSGRSDNH (SEQ ID NO: 420), which is also referred to herein as substrate 1002/LP'/0001, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 1037). In some embodiments, the CM1-CM2 substrate includes the sequence LSGRSGNHGGSGGSGQNQALRMA (SEQ ID NO: 421), which is also referred to herein as substrate 0002/LP'/1002, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 1037). In some embodiments, the CM1-CM2 substrate includes the sequence QNQALRMAGGSGGSLSGRSGNH (SEQ ID NO: 422), which is also referred to herein as substrate 1002/LP'/0002, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 1037). In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSGNH (SEQ ID NO: 423), which is also referred to herein as substrate 2002. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSANPRG (SEQ ID NO: 680), which is also referred to herein as substrate 2003. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPTSGRSANPRG (SEQ ID NO: 681), which is also referred to herein as substrate 2004. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPSGRSANPRG (SEQ ID NO: 682), which is also referred to herein as substrate 2005. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDDH (SEQ ID NO: 683), which is also referred to herein as substrate 2006. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDIH (SEQ ID NO: 684), which is also referred to herein as substrate 2007. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDQH (SEQ ID NO: 685), which is also referred to herein as substrate 2008.

In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDTH (SEQ ID NO: 686), which is also referred to herein as substrate 2009. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDYH (SEQ ID NO: 687), which is also referred to herein as substrate 2010. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDNP (SEQ ID NO: 688), which is also referred to herein as substrate 2011. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSANP (SEQ ID NO: 689), which is also referred to herein as substrate 2012. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSANI (SEQ ID NO: 690), which is also referred to herein as substrate 2013. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDDH (SEQ ID NO: 691), which is also referred to herein as substrate 3006. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDIH (SEQ ID NO: 692), which is also referred to herein as substrate 3007. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDQH (SEQ ID NO: 693), which is also referred to herein as substrate 3008. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDTH (SEQ ID NO: 694), which is also referred to herein as substrate 3009. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDYH (SEQ ID NO: 695), which is also referred to herein as substrate 3010. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDNP (SEQ ID NO: 696), which is also referred to herein as substrate 3011. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSANP (SEQ ID NO: 697), which is also referred to herein as substrate 3012. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSANI (SEQ ID NO: 698), which is also referred to herein as substrate 3013. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDNI (SEQ ID NO: 713), which is also referred to herein as substrate 2014. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDNI (SEQ ID NO: 714), which is also referred to herein as substrate 3014. In some embodiments, the CM1-CM2 substrate includes the sequence GLSGRSDNHGGAVGLLAPP (SEQ ID NO: 807), which is also referred to herein as substrate 0001/LP'/1004, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence

GLSGRSDNHGGVHMPGLGFLGP (SEQ ID NO: 808), which is also referred to herein as substrate 0001/LP'/1003, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG.

[000153] In some embodiments, the CM is a substrate for at least two proteases. In some embodiments, each protease is selected from the group consisting of those shown in Table 4. In some embodiments, the CM is a substrate for at least two proteases, wherein one of the proteases is selected from the group consisting of a MMP, thrombin, a neutrophil elastase, a cysteine protease, uPA, legumain and matriptase and the other protease is selected from the group consisting of those shown in Table 4. In some embodiments, the CM is a substrate for at least two proteases selected from the group consisting of a MMP, thrombin, a neutrophil elastase, a cysteine protease, uPA, legumain and matriptase.

[000154] In some embodiments, the activatable antibody includes at least a first CM and a second CM. In some embodiments, the first CM and the second CM are each polypeptides of no more than 15 amino acids long. In some embodiments, the first CM and the second CM in the activatable antibody in the uncleaved state have the structural arrangement from N-terminus to C-terminus as follows: MM-CM1-CM2-AB or AB-CM2-CM1-MM. In some embodiments, at least one of the first CM and the second CM is a polypeptide that functions as a substrate for a protease selected from the group consisting of a MMP, thrombin, a neutrophil elastase, a cysteine protease, uPA, legumain, and matriptase. In some embodiments, the first CM is cleaved by a first cleaving agent selected from the group consisting of a MMP, thrombin, a neutrophil elastase, a cysteine protease, uPA, legumain, and matriptase in a target tissue and the second CM is cleaved by a second cleaving agent in a target tissue. In some embodiments, the other protease is selected from the group consisting of those shown in Table 4. In some embodiments, the first cleaving agent and the second cleaving agent are the same protease selected from the group consisting of a MMP, thrombin, a neutrophil elastase, a cysteine protease, uPA, legumain, and matriptase, and the first CM and the second CM are different substrates for the enzyme. In some embodiments, the first cleaving agent and the second cleaving agent are the same protease selected from the group consisting of those shown in Table 4. In some embodiments, the first cleaving agent and the second cleaving agent are different proteases. In some embodiments, the first cleaving agent and the second cleaving agent are co-localized in the target tissue. In some embodiments, the first CM and the second CM are cleaved by at least one cleaving agent in the target tissue.

[000155] In some embodiments, the activatable antibody is exposed to and cleaved by a protease such that, in the activated or cleaved state, the activated antibody includes a light chain amino acid sequence that includes at least a portion of LP2 and/or CM sequence after the protease has cleaved the CM.

[000156] Suitable activatable anti-CD71 antibodies of the disclosure also include an antibody or antigen binding fragment thereof that binds to the same epitope on human CD71 and/or cynomolgus monkey CD71 as an anti-CD71 antibody comprising a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[000157] Suitable activatable anti-CD71 antibodies of the disclosure also include an antibody or antigen binding fragment thereof that binds to the same epitope on human CD71 and/or cynomolgus monkey CD71 as an anti-CD71 antibody comprising a VH CDR1 sequence comprising the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); a VH CDR2 sequence comprising the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); a VH CDR3 sequence comprising the amino acid sequence ENWDPGFAP (SEQ ID NO: 11); a VL CDR1 sequence comprising the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); a VL CDR2 sequence comprising the amino acid sequence STSNLAS (SEQ ID NO: 14); and a VL CDR3 sequence comprising the amino acid sequence QRRNYPYT (SEQ ID NO: 15).

[000158] Suitable activatable anti-CD71 antibodies of the disclosure also include an antibody or antigen-binding fragment thereof that binds to the same epitope on human CD71 and/or cynomolgus monkey CD71 as an anti-CD71 antibody comprising a heavy chain variable region amino acid sequence selected from the group consisting of the heavy chain variable region sequences shown in Table 12 and a light chain variable region amino acid sequence selected from the group consisting of the light chain variable region sequences shown in Table 12.

[000159] Suitable activatable anti-CD71 antibodies of the disclosure also include an antibody or antigen-binding fragment thereof that binds to the same epitope on human CD71 and/or cynomolgus monkey CD71 as an anti-CD71 antibody comprising a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the

combination is a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 13.

[000160] Suitable activatable anti-CD71 antibodies of the disclosure also include an antibody or antigen binding fragment thereof that cross-competes for binding to human CD71 and/or cynomolgus monkey CD71 to an anti-CD71 antibody comprising a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[000161] Suitable activatable anti-CD71 antibodies of the disclosure also include an antibody or antigen binding fragment thereof that cross-competes for binding to human CD71 and/or cynomolgus monkey CD71 to an anti-CD71 antibody comprising a VH CDR1 sequence comprising the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); a VH CDR2 sequence comprising the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); a VH CDR3 sequence comprising the amino acid sequence ENWDPGF AF (SEQ ID NO: 11); a VL CDR1 sequence comprising the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); a VL CDR2 sequence comprising the amino acid sequence STSNLAS (SEQ ID NO: 14); and a VL CDR3 sequence comprising the amino acid sequence QQR RNYPYT (SEQ ID NO: 15).

[000162] Suitable activatable anti-CD71 antibodies of the disclosure also include an antibody or antigen-binding fragment thereof that cross-competes for binding to human CD71 and/or cynomolgus monkey CD71 as an anti-CD71 antibody comprising a heavy chain variable region amino acid sequence selected from the group consisting of the heavy chain variable region sequences shown in Table 12 and a light chain variable region amino acid sequence selected from the group consisting of the light chain variable region sequences shown in Table 12.

[000163] Suitable activatable anti-CD71 antibodies of the disclosure also include an antibody or antigen-binding fragment thereof that cross-competes for binding to human CD71 and/or cynomolgus monkey CD71 as an anti-CD71 antibody comprising a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 13.

[000164] In some embodiments, the activatable anti-CD71 antibody is an activatable antibody that, in an activated state, binds CD71 comprising: an antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian CD71, wherein the AB specifically binds human CD71 and cynomolgus monkey CD71; a masking moiety (MM) that inhibits the binding of the AB to CD71 when the activatable antibody is in an uncleaved state; and a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease.

[000165] In some embodiments, the MM has a dissociation constant for binding to the AB that is greater than the dissociation constant of the AB to CD71. In some embodiments, the MM does not interfere or compete with the AB for binding to CD71 when the activatable antibody is in a cleaved state. In some embodiments, the MM is a polypeptide of no more than 40 amino acids in length. In some embodiments, the MM polypeptide sequence is different from that of human CD71. In some embodiments, the MM polypeptide sequence is no more than 50% identical to any natural binding partner of the AB. In some embodiments, the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 16-295 and 297-314. In some embodiments, the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 16, 17, and 297-314.

[000166] In some embodiments, the CM is a substrate for a protease that is active in diseased tissue. In some embodiments, the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 356-423, 680-698, 713, 714, and 789-808. In some embodiments, the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 406-423, 680-698, 713, 714, and 807-808.

[000167] In some embodiments, the activatable antibody comprises an antigen binding fragment thereof is selected from the group consisting of a Fab fragment, a F(ab')₂ fragment, a scFv, a scAb, a dAb, a single domain heavy chain antibody, and a single domain light chain antibody. In some embodiments, the AB of the activatable antibody specifically binds human CD71. In some embodiments, the AB comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QQRNYPYT (SEQ ID NO: 15). In some embodiments, the AB comprises a heavy chain variable region

comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8 and 809-908.

[000168] In some embodiments, the AB is linked to the CM. In some embodiments, the AB is linked directly to the CM. In some embodiments, the AB is linked to the CM via a linking peptide. In some embodiments, the MM is linked to the CM such that the activatable antibody in an uncleaved state comprises the structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM. In some embodiments, the activatable antibody comprises a linking peptide between the MM and the CM. In some embodiments, the activatable antibody comprises a linking peptide between the CM and the AB. In some embodiments, the activatable antibody comprises a first linking peptide (LP1) and a second linking peptide (LP2), and wherein the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-LP1-CM-LP2-AB or AB-LP2-CM-LP1-MM. In some embodiments, the two linking peptides need not be identical to each other. In some embodiments, each of LP1 and LP2 is a peptide of about 1 to 20 amino acids in length.

[000169] In some embodiments, the activatable antibody comprises the heavy chain sequence of SEQ ID NO: 325 or 699 and a light chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 327, 329, 331, 333, 335, 337, 650, 652, 654, 656, 658, 660, 670-673, 701-712, 721-788, 809-836, and 841-908.

[000170] In some embodiments, the activatable antibody comprises a combination of amino acid sequences, wherein the combination of amino acid sequences is selected from a single row in Table D, wherein for a given combination, (a) the heavy chain of the AB comprises the amino acid sequences of the VH CDR sequences corresponding to the given combination in the single row listed in Table D, (b) the light chain of the AB comprises the amino acid sequences of the VL CDR sequences corresponding to the given combination in the single row listed in Table D, (c) the MM comprises the amino acid sequence of the mask sequence (MM) corresponding to the given combination in the single row listed in Table D, and (d) the CM comprises the amino acid sequence of the substrate sequence (CM) corresponding to the given combination in the single row listed in Table D.

[000171] In some embodiments, the activatable antibody comprises a combination of amino acid sequences, wherein for a given combination of amino acid sequences, (a) the heavy chain of the AB comprises the amino acid sequences of the VH sequence or VH CDR

sequences selected from the group consisting of: the VH sequence or VH CDR sequences listed in the corresponding column of Table E, (b) the light chain of the AB comprises the amino acid sequences of the VL sequence or VL CDR sequences selected from the group consisting of: the VL sequence or VL CDR sequences listed in the corresponding column of Table E, (c) the MM comprises the amino acid sequence of the mask sequence (MM) selected from the group consisting of: the MM sequences listed in the corresponding column of Table E, and (d) the CM comprises the amino acid sequence of the substrate sequence (CM) selected from the group consisting of: the CM sequences listed in the corresponding column of Table E.

[000172] In some embodiments, the activatable anti-CD71 antibody comprises an antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian CD71, a MM, and a CM, wherein the activatable antibody comprises: a heavy chain sequence of SEQ ID NOS: 325 or 699; and a light chain comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 327, 329, 331, 333, 335, 337, 650, 652, 654, 656, 658, 660, 670-673, 701-712, 721-788, 809-836, and 841-908. In some embodiments, the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 16, 17, and 297-314, and the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 406-423, 680-698, 713, 714, and 789-808. In some embodiments, the AB comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QQRRNYPYT (SEQ ID NO: 15). In some embodiments, the AB comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 6-8 and 809-908.

[000173] In some embodiments, the activatable anti-CD71 antibody comprises an antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian CD71, a MM comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 16-295 and 297-314, and a CM comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 356-423, 680-698, 713, 714, and 789-808. In some embodiments, the MM comprises an amino acid sequence selected from the group

consisting of SEQ ID NOs: 16, 17, and 297-314, and the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 406-423, 680-698, 713, 714, and 807-808. In some embodiments, the AB comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QRRNYPYT (SEQ ID NO: 15). In some embodiments, the AB comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8 and 809-908.

[000174] In some embodiments, the activatable anti-CD71 antibody comprises an antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian CD71, wherein the AB specifically binds to the same epitope on human CD71 and/or cynomolgus monkey CD71 as an isolated antibody of the disclosure; a masking moiety (MM) that inhibits the binding of the AB to CD71 when the activatable antibody is in an uncleaved state; and a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease.

[000175] In some embodiments, the anti-CD71 activatable antibody of the disclosure comprises an isolated antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian CD71, wherein the AB specifically binds human CD71 and cynomolgus monkey CD71. In some embodiments, the antibody or antigen binding fragment thereof comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QRRNYPYT (SEQ ID NO: 15). In some embodiments, the activatable antibody or antigen binding fragment thereof comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8.

[000176] In some embodiments, the activatable anti-CD71 antibody comprises an antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian

CD71, wherein the AB specifically cross-competes with an isolated antibody of the disclosure for binding to human CD71 and/or cynomolgus monkey CD71; a masking moiety (MM) that inhibits the binding of the AB to CD71 when the activatable antibody is in an uncleaved state; and a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease.

[000177] In some embodiments, the anti-CD71 activatable antibody of the disclosure comprises an isolated antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian CD71, wherein the AB specifically binds human CD71 and cynomolgus monkey CD71. In some embodiments, the antibody or antigen binding fragment thereof comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QQRRNYPYT (SEQ ID NO: 15). In some embodiments, the activatable antibody or antigen binding fragment thereof comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8.

[000178] In some embodiments, the activatable antibody also includes an agent conjugated to the AB. In some embodiments, the agent conjugated to the AB or the AB of an activatable antibody is a therapeutic agent. In some embodiments, the agent is an antineoplastic agent. In some embodiments, the agent is a toxin or fragment thereof. As used herein, a fragment of a toxin is a fragment that retains toxic activity. In some embodiments, the agent is conjugated to the AB via a cleavable linker. In some embodiments, the agent is conjugated to the AB via a linker that includes at least one CM1-CM2 substrate sequence. In some embodiments, the agent is conjugated to the AB via a noncleavable linker. In some embodiments, the agent is conjugated to the AB via a linker that is cleavable in an intracellular or lysosomal environment. In some embodiments, the agent is a microtubule inhibitor. In some embodiments, the agent is a nucleic acid damaging agent, such as a DNA alkylator, a DNA cleaving agent, a DNA cross-linker, a DNA intercalator, or other DNA damaging agent. In some embodiments, the agent is an agent selected from the group listed in Table 5. In some embodiments, the agent is a dolastatin. In some embodiments, the agent is an auristatin or derivative thereof. In some embodiments,

the agent is auristatin E or a derivative thereof. In some embodiments, the agent is monomethyl auristatin E (MMAE). In some embodiments, the agent is monomethyl auristatin D (MMAD). In some embodiments, the agent is a maytansinoid or maytansinoid derivative. In some embodiments, the agent is DM1 or DM4. In some embodiments, the agent is a duocarmycin or derivative thereof. In some embodiments, the agent is a calicheamicin or derivative thereof. In some embodiments, the agent is a pyrrolobenzodiazepine. In some embodiments, the agent is a pyrrolobenzodiazepine dimer.

[000179] In some embodiments, the activatable antibody is conjugated to one or more equivalents of an agent. In some embodiments, the activatable antibody is conjugated to one equivalent of the agent. In some embodiments, the activatable antibody is conjugated to two, three, four, five, six, seven, eight, nine, ten, or greater than ten equivalents of the agent. In some embodiments, the activatable antibody is part of a mixture of activatable antibodies having a homogeneous number of equivalents of conjugated agents. In some embodiments, the activatable antibody is part of a mixture of activatable antibodies having a heterogeneous number of equivalents of conjugated agents. In some embodiments, the mixture of activatable antibodies is such that the average number of agents conjugated to each activatable antibody is between zero to one, between one to two, between two and three, between three and four, between four and five, between five and six, between six and seven, between seven and eight, between eight and nine, between nine and ten, and ten and greater. In some embodiments, the mixture of activatable antibodies is such that the average number of agents conjugated to each activatable antibody is one, two, three, four, five, six, seven, eight, nine, ten, or greater.

[000180] In some embodiments, the activatable antibody comprises one or more site-specific amino acid sequence modifications such that the number of lysine and/or cysteine residues is increased or decreased with respect to the original amino acid sequence of the activatable antibody, thus in some embodiments correspondingly increasing or decreasing the number of agents that can be conjugated to the activatable antibody, or in some embodiments limiting the conjugation of the agents to the activatable antibody in a site-specific manner. In some embodiments, the modified activatable antibody is modified with one or more non-natural amino acids in a site-specific manner, thus in some embodiments limiting the conjugation of the agents to only the sites of the non-natural amino acids.

[000181] In some embodiments, the agent is an anti-inflammatory agent.

[000182] In some embodiments, the activatable antibody also includes a detectable moiety. In some embodiments, the detectable moiety is a diagnostic agent.

[000183] In some embodiments, the activatable antibody also includes a signal peptide. In some embodiments, the signal peptide is conjugated to the activatable antibody via a spacer. In some embodiments, the spacer is conjugated to the activatable antibody in the absence of a signal peptide. In some embodiments, the spacer is joined directly to the MM of the activatable antibody. In some embodiments, the spacer is joined directly to the MM of the activatable antibody in the structural arrangement from N-terminus to C-terminus of spacer-MM-CM-AB. An example of a spacer joined directly to the N-terminus of MM of the activatable antibody is QGQSGQ (SEQ ID NO: 424). Other examples of a spacer joined directly to the N-terminus of MM of the activatable antibody include QGQSGQG (SEQ ID NO: 645), QGQSG (SEQ ID NO: 646), QGQS (SEQ ID NO: 647), QGQ (SEQ ID NO: 648), QG (SEQ ID NO: 649), and Q. Other examples of a spacer joined directly to the N-terminus of MM of the activatable antibody include GQSGQG (SEQ ID NO: 666), QSGQG (SEQ ID NO: 667), SGQG (SEQ ID NO: 668), GQG (SEQ ID NO: 669), and G. In some embodiments, no spacer is joined to the N-terminus of the MM. In some embodiments, the spacer includes at least the amino acid sequence QGQSGQ (SEQ ID NO: 424). In some embodiments, the spacer includes at least the amino acid sequence QGQSGQG (SEQ ID NO: 645). In some embodiments, the spacer includes at least the amino acid sequence QGQSG (SEQ ID NO: 646). In some embodiments, the spacer includes at least the amino acid sequence QGQS (SEQ ID NO: 647). In some embodiments, the spacer includes at least the amino acid sequence QGQ (SEQ ID NO: 648). In some embodiments, the spacer includes at least the amino acid sequence QG (SEQ ID NO: 649). In some embodiments, the spacer includes at least the amino acid residue Q. In some embodiments, the spacer includes at least the amino acid sequence GQSGQG (SEQ ID NO: 666). In some embodiments, the spacer includes at least the amino acid sequence QSGQG (SEQ ID NO: 667). In some embodiments, the spacer includes at least the amino acid sequence SGQG (SEQ ID NO: 668). In some embodiments, the spacer includes at least the amino acid sequence GQG (SEQ ID NO: 669). In some embodiments, the spacer includes at least the amino acid sequence G. In some embodiments, the spacer is absent.

[000184] In some embodiments, the AB of the activatable antibody naturally contains one or more disulfide bonds. In some embodiments, the AB can be engineered to include one or more disulfide bonds.

[000185] In some embodiments, activatable antibody or antigen binding fragment thereof is conjugated to an agent. In some embodiments, the activatable antibody comprises an antibody or antigen binding fragment thereof cross-competes with an isolated antibody that comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QRRNYPYT (SEQ ID NO: 15). In some embodiments, the activatable antibody comprises an antibody or antigen binding fragment thereof cross-competes with an isolated antibody that comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8. In some embodiments, the agent is a toxin or fragment thereof. In some embodiments, the agent is a microtubule inhibitor. In some embodiments, the agent is a nucleic acid damaging agent. In some embodiments, the agent is selected from the group consisting of a dolastatin or a derivative thereof, an auristatin or a derivative thereof, a maytansinoid or a derivative thereof, a duocarmycin or a derivative thereof, a calicheamicin or a derivative thereof, and a pyrrolobenzodiazepine or a derivative thereof. In some embodiments, the agent is auristatin E or a derivative thereof. In some embodiments, the agent is monomethyl auristatin E (MMAE). In some embodiments, the agent is monomethyl auristatin D (MMAD). In some embodiments, the agent is a maytansinoid selected from the group consisting of DM1 and DM4. In some embodiments, the agent is maytansinoid DM4. In some embodiments, the agent is duocarmycin. In some embodiments, the agent is conjugated to the AB via a linker. In some embodiments, the linker with which the agent is conjugated to the AB comprises an SPDB moiety, a vc moiety, or a PEG2-vc moiety. In some embodiments, the linker and toxin conjugated to the AB comprises an SPDB-DM4 moiety, a vc-MMAD moiety, a vc-MMAE moiety, vc-duocarmycin, or a PEG2-vc-MMAD moiety. In some embodiments, the linker is a cleavable linker. In some embodiments, the linker is a non-cleavable linker. In some embodiments, the agent is a detectable moiety. In some embodiments, the detectable moiety is a diagnostic agent.

[000186] In some embodiments, the conjugated activatable antibody comprises a conjugated activatable antibody that, in an activated state, binds CD71 comprising: an antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian

CD71, wherein the AB specifically binds human CD71 and cynomolgus monkey CD71; a masking moiety (MM) that inhibits the binding of the AB to CD71 when the activatable antibody is in an uncleaved state; a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease; and an agent conjugated to the AB. In some embodiments, the agent is selected from the group consisting of a dolastatin or a derivative thereof, an auristatin or a derivative thereof, a maytansinoid or a derivative thereof, a duocarmycin or a derivative thereof, a calicheamicin or a derivative thereof, and a pyrrolobenzodiazepine or a derivative thereof. In some embodiments, the agent is selected from the group consisting of auristatin E, monomethyl auristatin F (MMAF), monomethyl auristatin E (MMAE), monomethyl auristatin D (MMAD), maytansinoid DM4, maytansinoid DM1, a duocarmycin, a pyrrolobenzodiazepine, and a pyrrolobenzodiazepine dimer. In some embodiments, the agent is conjugated to the AB via a linker. In some embodiments, the linker with which the agent is conjugated to the AB comprises an SPDB moiety, a vc moiety, or a PEG2-vc moiety. In some embodiments, the linker and toxin conjugated to the AB comprises an SPDB-DM4 moiety, a vc-MMAD moiety, a vc-MMAE moiety, vc-duocarmycin, or a PEG2-vc-MMAD moiety. In some embodiments, the AB of the conjugated activatable antibody or antigen binding fragment thereof comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAP (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QRRNYPYT (SEQ ID NO: 15). In some embodiments, the AB of the conjugated activatable antibody or antigen binding fragment thereof comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8 and 809-908. In some embodiments, the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 16-295 and 297-314. In some embodiments, the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 16, 17, and 297-314. In some embodiments, the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 356-423, 680-698, 713, 714, and 789-808. In some embodiments, the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 406-423, 680-698, 713, 714, and 807-808. In some embodiments, the

activatable antibody comprises a combination of amino acid sequences, wherein the combination of amino acid sequences is selected from a single row in Table D, wherein for a given combination, (a) the heavy chain of the AB comprises the amino acid sequences of the VH CDR sequences corresponding to the given combination in the single row listed in Table D, (b) the light chain of the AB comprises the amino acid sequences of the VL CDR sequences corresponding to the given combination in the single row listed in Table D, (c) the MM comprises the amino acid sequence of the mask sequence (MM) corresponding to the given combination in the single row listed in Table D, and (d) the CM comprises the amino acid sequence of the substrate sequence (CM) corresponding to the given combination in the single row listed in Table D. In some embodiments, the activatable antibody comprises a combination of amino acid sequences, wherein for a given combination of amino acid sequences, (a) the heavy chain of the AB comprises the amino acid sequences of the VH sequence or VH CDR sequences selected from the group consisting of: the VH sequence or VH CDR sequences listed in the corresponding column of Table E, (b) the light chain of the AB comprises the amino acid sequences of the VL sequence or VL CDR sequences selected from the group consisting of: the VL sequence or VL CDR sequences listed in the corresponding column of Table E, (c) the MM comprises the amino acid sequence of the mask sequence (MM) selected from the group consisting of: the MM sequences listed in the corresponding column of Table E, and (d) the CM comprises the amino acid sequence of the substrate sequence (CM) selected from the group consisting of: the CM sequences listed in the corresponding column of Table E. In some embodiments, the activatable antibody comprises: a heavy chain comprising the amino acid sequence of SEQ ID NOS: 325 or 699; and a light chain comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 327, 329, 331, 333, 335, 337, 650, 652, 654, 656, 658, 660, 670-673, 701-712, 721-788, 809-836, and 841-908.

[000187] In some embodiments, the conjugated activatable antibody comprises a conjugated activatable antibody that, in an activated state, binds to CD71, comprising: an antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian CD71, wherein the AB specifically binds human CD71 and cynomolgus monkey CD71; a masking moiety (MM) that inhibits the binding of the AB to CD71 when the activatable antibody is in an uncleaved state; a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease; and an agent conjugated to the AB, wherein the AB comprises: (i) the VH CDR1 sequence GYTFTSYWMH (SEQ ID

NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QQRRNYPYT (SEQ ID NO: 15), or (ii) a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 6-8, or (iii) a heavy chain comprising the amino acid sequence of SEQ ID NOS: 325 or 699, and a light chain comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 327, 329, 331, 333, 335, 337, 650, 652, 654, 656, 658, 660, 670-673, 701-712, 721-788, 809-836, and 841-908; and wherein the agent is selected from the group consisting of auristatin E, monomethyl auristatin F (MMAF), monomethyl auristatin E (MMAE), monomethyl auristatin D (MMAD), maytansinoid DM4, maytansinoid DM1, a pyrrolobenzodiazepine, a pyrrolobenzodiazepine dimer, and a duocarmycin. In some embodiments, the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 16-295 and 297-314. In some embodiments, the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 16, 17, and 297-314. In some embodiments, the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 356-423, 680-698, 713, 714, and 789-808. In some embodiments, the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 406-423, 680-698, 713, 714, and 807-808. In some embodiments, the agent is conjugated to the AB via a linker, and wherein the linker to which the agent is conjugated to the AB comprises an SPDB moiety, a vc moiety, or a PEG2-vc moiety. In some embodiments, the linker and toxin conjugated to the AB comprises an SPDB-DM4 moiety, a vc-MMAD moiety, a vc-MMAE moiety, vc-duocarmycin, or a PEG2-vc-MMAD moiety.

[000188] In some embodiments, the conjugated activatable antibody comprises a conjugated activatable antibody or conjugated antibody comprising: an antibody or antigen binding fragment thereof (AB) that, in an activated state, binds CD71; and a toxin conjugated to the AB via a linker, wherein the conjugated activatable antibody or the conjugated antibody comprises amino acid sequences, a linker, and a toxin selected from a single row in Table F, wherein for the given combination: (a) the AB comprises a heavy chain comprising the amino acid sequence of the heavy chain sequence or heavy chain variable domain sequence corresponding to the given combination in the single row listed in

Table F, (b) the AB comprises a light chain comprising the amino acid sequence of the light chain sequence or light chain variable domain sequence corresponding to the given combination in the single row listed in Table F, and (c) the linker and the toxin comprise the linker and the toxin corresponding to the given combination in the single row listed in Table F.

[000189] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 3-5.

[000190] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[000191] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a nucleic acid sequence encoding a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[000192] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 3-5, and a nucleic acid sequence encoding a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[000193] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%,

95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a heavy chain variable region amino acid sequence comprising selected from the group consisting of SEQ ID NO: 3-5.

[000194] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a light variable region chain amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[000195] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 1 and 3-5, and a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 2 and 6-8.

[000196] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 3-5, and a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence encoding a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NO: 6-8.

[000197] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain variable region amino acid sequence selected from the group consisting of the heavy chain variable region sequences shown in Table 12. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain variable region amino acid sequence selected from the group consisting of the light chain variable region sequences shown in Table 12. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence

encoding a heavy chain variable region amino acid sequence selected from the group consisting of the heavy chain variable region sequences shown in Table 12 and a nucleic acid sequence encoding a light chain variable region amino acid sequence selected from the group consisting of the light chain variable region sequences shown in Table 12.

[000198] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the heavy chain variable region sequences shown in Table 12. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the light chain variable region sequences shown in Table 12. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the heavy chain variable region sequences shown in Table 12 and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain variable region amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the light chain variable region sequences shown in Table 12.

[000199] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3) sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2) sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one CDR sequence is selected from the group consisting of a VH

CDR1 sequence shown in Table 13; a VH CDR2 sequence shown in Table 13; a VH CDR3 sequence shown in Table 13; a VL CDR1 sequence shown in Table 13; a VL CDR2 sequence shown in Table 13; and a VL CDR3 sequence shown in Table 13.

[000200] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR1 sequence shown in Table 13; a VH CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR2 sequence shown in Table 13; a VH CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR3 sequence shown in Table 13; a VL CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR1 sequence shown in Table 13; a VL CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR2 sequence shown in Table 13; and a VL CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR3 sequence shown in Table 13.

[000201] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 13.

[000202] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain variable region that comprise a combination of a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the three light chain CDR sequences (VL CDR1, VL CDR2, VL CDR3) shown in a single row in Table 13.

[000203] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain variable region

that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein the combination is a combination of the three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 13.

[000204] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 13.

[000205] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain variable region that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 13.

[000206] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain variable region that comprise a combination of a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of three light chain CDR sequences (VL CDR1, VL CDR2, VL CDR3) shown in a single row in Table 13.

[000207] The disclosure also provides methods for producing an activatable antibody of the disclosure by culturing a cell under conditions that lead to expression of the activatable antibody, wherein the cell comprises a nucleic acid molecule of the disclosure or a vector of the disclosure.

[000208] The disclosure also provides methods of manufacturing an activatable antibody that, in an activated state, binds CD71, the method comprising: (a) culturing a cell comprising a nucleic acid construct that encodes the activatable antibody under conditions that lead to expression of the activatable antibody, wherein the activatable antibody

comprises an activatable antibody of the disclosure; and (b) recovering the activatable antibody.

[000209] In some embodiments, the activatable antibody includes one or more polypeptides that include the combination of sequences in a given row of Table D or any combination of a mask sequence (MM), a substrate sequence (CM), a light chain variable domain sequence or light chain variable domain CDR sequences, and a heavy chain variable domain sequence or heavy chain variable domain CDR sequences of Table E.

Table D: Anti-CD71 Activatable Antibody Combinations

Comb. No.	Mask Sequence (MM)	Substrate Sequence (CM)	VL CDRs SEQ ID NOs	VH CDRs SEQ ID NOs
1	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	LSGRSDNH (SEQ ID NO: 359)	12, 14, 15	9, 10, 11
2	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSS (SEQ ID NO: 382)	12, 14, 15	9, 10, 11
3	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	LSGRSGNH (SEQ ID NO: 789)	12, 14, 15	9, 10, 11
4	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPP (SEQ ID NO: 390)	12, 14, 15	9, 10, 11
5	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	VHMPLGFLGP (SEQ ID NO: 370)	12, 14, 15	9, 10, 11
6	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	TSTSGRSANPRG (SEQ ID NO: 397)	12, 14, 15	9, 10, 11
7	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	QNQALRMA (SEQ ID NO: 377)	12, 14, 15	9, 10, 11
8	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSGRSDNH (SEQ ID NO: 406)	12, 14, 15	9, 10, 11
9	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSGRSGNH (SEQ ID NO: 423)	12, 14, 15	9, 10, 11
10	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSGRSANPRG (SEQ ID NO: 680)	12, 14, 15	9, 10, 11
11	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPPTSGRSANPRG (SEQ ID NO: 681)	12, 14, 15	9, 10, 11
12	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPPSGRSANPRG (SEQ ID NO: 682)	12, 14, 15	9, 10, 11
13	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSGRSDDH (SEQ ID NO: 683)	12, 14, 15	9, 10, 11
14	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSGRSDIH (SEQ ID NO: 684)	12, 14, 15	9, 10, 11
15	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSGRSDQH (SEQ ID NO: 685)	12, 14, 15	9, 10, 11
16	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSGRSDTH (SEQ ID NO: 686)	12, 14, 15	9, 10, 11
17	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSGRSDYH (SEQ ID NO: 687)	12, 14, 15	9, 10, 11

18	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSGRSDNP (SEQ ID NO: 688)	12, 14, 15	9, 10, 11
19	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSGRSANP (SEQ ID NO: 689)	12, 14, 15	9, 10, 11
20	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSGRSANI (SEQ ID NO: 690)	12, 14, 15	9, 10, 11
21	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSGRSDNI (SEQ ID NO: 713)	12, 14, 15	9, 10, 11
22	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPPGGLSGRSDNH (SEQ ID NO: 412)	12, 14, 15	9, 10, 11
23	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPPGGLSGRSDDH (SEQ ID NO: 691)	12, 14, 15	9, 10, 11
24	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPPGGLSGRSDIH (SEQ ID NO: 692)	12, 14, 15	9, 10, 11
25	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPPGGLSGRSDQH (SEQ ID NO: 693)	12, 14, 15	9, 10, 11
26	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPPGGLSGRSDTH (SEQ ID NO: 694)	12, 14, 15	9, 10, 11
27	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPPGGLSGRSDYH (SEQ ID NO: 695)	12, 14, 15	9, 10, 11
28	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPPGGLSGRSDNP (SEQ ID NO: 696)	12, 14, 15	9, 10, 11
29	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPPGGLSGRSANP (SEQ ID NO: 697)	12, 14, 15	9, 10, 11
30	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPPGGLSGRSANI (SEQ ID NO: 698)	12, 14, 15	9, 10, 11
31	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	AVGLLAPPGGLSGRSDNI (SEQ ID NO: 714)	12, 14, 15	9, 10, 11
32	QFCPWSYYLIGDCDI (SEQ ID NO: 16)	ISSGLLSSGGSGGLSGRSDNH (SEQ ID NO: 407)	12, 14, 15	9, 10, 11
33	NLCTEHSFALDCRSY (SEQ ID NO: 17)	LSGRSDNH (SEQ ID NO: 359)	12, 14, 15	9, 10, 11
34	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSS (SEQ ID NO: 382)	12, 14, 15	9, 10, 11
35	NLCTEHSFALDCRSY (SEQ ID NO: 17)	LSGRSGNH (SEQ ID NO: 789)	12, 14, 15	9, 10, 11
36	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPP (SEQ ID NO: 390)	12, 14, 15	9, 10, 11
37	NLCTEHSFALDCRSY (SEQ ID NO: 17)	VHMPLGFLGP (SEQ ID NO: 370)	12, 14, 15	9, 10, 11
38	NLCTEHSFALDCRSY (SEQ ID NO: 17)	TSTSGRSANPRG (SEQ ID NO: 397)	12, 14, 15	9, 10, 11
39	NLCTEHSFALDCRSY (SEQ ID NO: 17)	QNQALRMA (SEQ ID NO: 377)	12, 14, 15	9, 10, 11
40	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSGRSDNH (SEQ ID NO: 406)	12, 14, 15	9, 10, 11
41	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSGRSGNH (SEQ ID NO: 423)	12, 14, 15	9, 10, 11
42	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSGRSANPRG (SEQ ID NO: 680)	12, 14, 15	9, 10, 11
43	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPPTSGRSANPRG (SEQ ID NO: 681)	12, 14, 15	9, 10, 11
44	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPPSGRSANPRG (SEQ ID NO: 682)	12, 14, 15	9, 10, 11

45	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSGRSDDH (SEQ ID NO: 683)	12, 14, 15	9, 10, 11
46	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSGRSDIH (SEQ ID NO: 684)	12, 14, 15	9, 10, 11
47	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSGRSDQH (SEQ ID NO: 685)	12, 14, 15	9, 10, 11
48	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSGRSDTH (SEQ ID NO: 686)	12, 14, 15	9, 10, 11
49	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSGRSDYH (SEQ ID NO: 687)	12, 14, 15	9, 10, 11
50	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSGRSDNP (SEQ ID NO: 688)	12, 14, 15	9, 10, 11
51	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSGRSANP (SEQ ID NO: 689)	12, 14, 15	9, 10, 11
52	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSGRSANI (SEQ ID NO: 690)	12, 14, 15	9, 10, 11
53	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLLSGRSDNI (SEQ ID NO: 713)	12, 14, 15	9, 10, 11
54	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPPGGLSGRSDNH (SEQ ID NO: 412)	12, 14, 15	9, 10, 11
55	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPPGGLSGRSDDH (SEQ ID NO: 691)	12, 14, 15	9, 10, 11
56	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPPGGLSGRSDIH (SEQ ID NO: 692)	12, 14, 15	9, 10, 11
57	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPPGGLSGRSDQH (SEQ ID NO: 693)	12, 14, 15	9, 10, 11
58	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPPGGLSGRSDTH (SEQ ID NO: 694)	12, 14, 15	9, 10, 11
59	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPPGGLSGRSDYH (SEQ ID NO: 695)	12, 14, 15	9, 10, 11
60	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPPGGLSGRSDNP (SEQ ID NO: 696)	12, 14, 15	9, 10, 11
61	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPPGGLSGRSANP (SEQ ID NO: 697)	12, 14, 15	9, 10, 11
62	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPPGGLSGRSANI (SEQ ID NO: 698)	12, 14, 15	9, 10, 11
63	NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPPGGLSGRSDNI (SEQ ID NO: 714)	12, 14, 15	9, 10, 11
64	NLCTEHSFALDCRSY (SEQ ID NO: 17)	ISSGLSSGGSGGSLSGRSDNH (SEQ ID NO: 407)	12, 14, 15	9, 10, 11
65	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	LSGRSDNH (SEQ ID NO: 359)	12, 14, 15	9, 10, 11
66	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLSS (SEQ ID NO: 382)	12, 14, 15	9, 10, 11
67	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	LSGRSGNH (SEQ ID NO: 789)	12, 14, 15	9, 10, 11
68	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPP (SEQ ID NO: 390)	12, 14, 15	9, 10, 11
69	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	VHMPGLGFLGP (SEQ ID NO: 370)	12, 14, 15	9, 10, 11
70	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	TSTSGRSANPRG (SEQ ID NO: 397)	12, 14, 15	9, 10, 11
71	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	QNQALRMA (SEQ ID NO: 377)	12, 14, 15	9, 10, 11

72	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSGRSDNH (SEQ ID NO: 406)	12, 14, 15	9, 10, 11
73	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSGRSGNH (SEQ ID NO: 423)	12, 14, 15	9, 10, 11
74	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSGRSANPRG (SEQ ID NO: 680)	12, 14, 15	9, 10, 11
75	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPPTSGRSANPRG (SEQ ID NO: 681)	12, 14, 15	9, 10, 11
76	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPPSGRSANPRG (SEQ ID NO: 682)	12, 14, 15	9, 10, 11
77	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSGRSDDH (SEQ ID NO: 683)	12, 14, 15	9, 10, 11
78	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSGRSDIH (SEQ ID NO: 684)	12, 14, 15	9, 10, 11
79	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSGRSDQH (SEQ ID NO: 685)	12, 14, 15	9, 10, 11
80	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSGRSDTH (SEQ ID NO: 686)	12, 14, 15	9, 10, 11
81	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSGRSDYH (SEQ ID NO: 687)	12, 14, 15	9, 10, 11
82	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSGRSDNP (SEQ ID NO: 688)	12, 14, 15	9, 10, 11
83	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSGRSANP (SEQ ID NO: 689)	12, 14, 15	9, 10, 11
84	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSGRSANI (SEQ ID NO: 690)	12, 14, 15	9, 10, 11
85	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSGRSDNI (SEQ ID NO: 713)	12, 14, 15	9, 10, 11
86	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPPGGLSGRSDNH (SEQ ID NO: 412)	12, 14, 15	9, 10, 11
87	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPPGGLSGRSDDH (SEQ ID NO: 691)	12, 14, 15	9, 10, 11
88	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPPGGLSGRSDIH (SEQ ID NO: 692)	12, 14, 15	9, 10, 11
89	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPPGGLSGRSDQH (SEQ ID NO: 693)	12, 14, 15	9, 10, 11
90	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPPGGLSGRSDTH (SEQ ID NO: 694)	12, 14, 15	9, 10, 11
91	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPPGGLSGRSDYH (SEQ ID NO: 695)	12, 14, 15	9, 10, 11
92	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPPGGLSGRSDNP (SEQ ID NO: 696)	12, 14, 15	9, 10, 11
93	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPPGGLSGRSANP (SEQ ID NO: 697)	12, 14, 15	9, 10, 11
94	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPPGGLSGRSANI (SEQ ID NO: 698)	12, 14, 15	9, 10, 11
95	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	AVGLLAPPGGLSGRSDNI (SEQ ID NO: 714)	12, 14, 15	9, 10, 11
96	NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSSGGSGGLSGRSDNH (SEQ ID NO: 407)	12, 14, 15	9, 10, 11
97	CTEHSFALDC (SEQ ID NO: 314)	LSGRSDNH (SEQ ID NO: 359)	12, 14, 15	9, 10, 11
98	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSS (SEQ ID NO: 382)	12, 14, 15	9, 10, 11

99	CTEHSFALDC (SEQ ID NO: 314)	LSGRSGNH (SEQ ID NO: 789)	12, 14, 15	9, 10, 11
100	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPP (SEQ ID NO: 390)	12, 14, 15	9, 10, 11
101	CTEHSFALDC (SEQ ID NO: 314)	VHMLPLGFLGP (SEQ ID NO: 370)	12, 14, 15	9, 10, 11
102	CTEHSFALDC (SEQ ID NO: 314)	TSTSGRSANPRG (SEQ ID NO: 397)	12, 14, 15	9, 10, 11
103	CTEHSFALDC (SEQ ID NO: 314)	QNQALRMA (SEQ ID NO: 377)	12, 14, 15	9, 10, 11
104	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSGRSDNH (SEQ ID NO: 406)	12, 14, 15	9, 10, 11
105	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSGRSGNH (SEQ ID NO: 423)	12, 14, 15	9, 10, 11
106	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSGRSANPRG (SEQ ID NO: 680)	12, 14, 15	9, 10, 11
107	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPPTSGRSANPRG (SEQ ID NO: 681)	12, 14, 15	9, 10, 11
108	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPPSGRSANPRG (SEQ ID NO: 682)	12, 14, 15	9, 10, 11
109	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSGRSDDH (SEQ ID NO: 683)	12, 14, 15	9, 10, 11
110	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSGRSDIH (SEQ ID NO: 684)	12, 14, 15	9, 10, 11
111	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSGRSDQH (SEQ ID NO: 685)	12, 14, 15	9, 10, 11
112	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSGRSDTH (SEQ ID NO: 686)	12, 14, 15	9, 10, 11
113	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSGRSDYH (SEQ ID NO: 687)	12, 14, 15	9, 10, 11
114	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSGRSDNP (SEQ ID NO: 688)	12, 14, 15	9, 10, 11
115	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSGRSANP (SEQ ID NO: 689)	12, 14, 15	9, 10, 11
116	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSGRSANI (SEQ ID NO: 690)	12, 14, 15	9, 10, 11
117	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSGRSDNI (SEQ ID NO: 713)	12, 14, 15	9, 10, 11
118	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPPGGLSGRSDNH (SEQ ID NO: 412)	12, 14, 15	9, 10, 11
119	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPPGGLSGRSDDH (SEQ ID NO: 691)	12, 14, 15	9, 10, 11
120	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPPGGLSGRSDIH (SEQ ID NO: 692)	12, 14, 15	9, 10, 11
121	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPPGGLSGRSDQH (SEQ ID NO: 693)	12, 14, 15	9, 10, 11
122	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPPGGLSGRSDTH (SEQ ID NO: 694)	12, 14, 15	9, 10, 11
123	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPPGGLSGRSDYH (SEQ ID NO: 695)	12, 14, 15	9, 10, 11
124	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPPGGLSGRSDNP (SEQ ID NO: 696)	12, 14, 15	9, 10, 11
125	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPPGGLSGRSANP (SEQ ID NO: 697)	12, 14, 15	9, 10, 11

126	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPPGGLSGRSANI (SEQ ID NO: 698)	12, 14, 15	9, 10, 11
127	CTEHSFALDC (SEQ ID NO: 314)	AVGLLAPPGGLSGRSDNI (SEQ ID NO: 714)	12, 14, 15	9, 10, 11
128	CTEHSFALDC (SEQ ID NO: 314)	ISSGLLSSGGSGGSLSGRSDNH (SEQ ID NO: 407)	12, 14, 15	9, 10, 11

Table E: Anti-CD71 Activatable Antibody Components

Mask Sequence (MM)	Substrate Sequence (CM)	VL or VL CDRs	VH or VH CDRs
QFCPWSYYLIGDCDI (SEQ ID NO: 16)	LSGRSDNH (SEQ ID NO: 359)	SEQ ID NOS: 12, 14, 15	SEQ ID NOS: 9, 10, 11
QFCAWSYYLIGDCDI (SEQ ID NO: 297)	TGRGPSWV (SEQ ID NO: 356)	SEQ ID NO: 6	SEQ ID NO: 3
QFCPASYYLIGDCDI (SEQ ID NO: 298)	PLTGRSGG (SEQ ID NO: 362)	SEQ ID NO: 7	SEQ ID NO: 4
QFCPWAYYYLIGDCDI (SEQ ID NO: 299)	TARGPSFK (SEQ ID NO: 358)	SEQ ID NO: 8	SEQ ID NO: 5
QFCPWSAYLIGDCDI (SEQ ID NO: 300)	NTLSGRSENHSG (SEQ ID NO: 395)	SEQ ID NOS: 13, 14, 15	
QFCPWSYALIGDCDI (SEQ ID NO: 301)	NTLSGRSGNHGS (SEQ ID NO: 396)		
QFCPWSYYAIGDCDI (SEQ ID NO: 302)	TSTSGRSANPRG (SEQ ID NO: 397)		
QFCPWSYYLAGDCDI (SEQ ID NO: 303)	TSGRSANP (SEQ ID NO: 398)		
QFCPWSYYLIGACDI (SEQ ID NO: 304)	VHMPLGFLGP (SEQ ID NO: 370)		
NLCTEHSFALDCRSY (SEQ ID NO: 17)	AVGLLAPP (SEQ ID NO: 390)		
NLCAEHSFALDCRSY (SEQ ID NO: 305)	AQNLLGMV (SEQ ID NO: 378)		
NLCTAHSFALDCRSY (SEQ ID NO: 306)	QNQALRMA (SEQ ID NO: 377)		
NLCTEASFALDCRSY (SEQ ID NO: 307)	LAAPLGLL (SEQ ID NO: 389)		
NLCTEHAFALDCRSY (SEQ ID NO: 308)	STFPFGMF (SEQ ID NO: 379)		
NLCTEHSAAALDCRSY (SEQ ID NO: 309)	ISSGLLSS (SEQ ID NO: 382)		
NLCTEHSFAALDCRSY (SEQ ID NO: 310)	PAGLWLDP (SEQ ID NO: 392)		
NLCTEHSFALACRSY (SEQ ID NO: 311)	VAGRSMRP (SEQ ID NO: 399)		
NLCTEHSFALDCASY (SEQ ID NO: 312)	VVPEGRRS (SEQ ID NO: 400)		
CTEHSFALDCRSY (SEQ ID NO: 313)	ILPRSPAF (SEQ ID NO: 401)		

CTEHSFALDC (SEQ ID NO: 314)	MVLGRSLL (SEQ ID NO: 402)		
	QGRAITFI (SEQ ID NO: 403)		
	SPRSIMLA (SEQ ID NO: 404)		
	SMLRSMPL (SEQ ID NO: 405)		
	ISSGLLSGRSDNH (SEQ ID NO: 406)		
	AVGLLAPPGGLSGRSDNH (SEQ ID NO: 412)		
	ISSGLLSSGGSGGSLSGRSDNH (SEQ ID NO: 407)		
	LSGRSGNH (SEQ ID NO: 789)		
	SGRSANPRG (SEQ ID NO: 790)		
	LSGRSDDH (SEQ ID NO: 791)		
	LSGRSDIH (SEQ ID NO: 792)		
	LSGRSDQH (SEQ ID NO: 793)		
	LSGRSDTH (SEQ ID NO: 794)		
	LSGRSDYH (SEQ ID NO: 795)		
	LSGRSDNP (SEQ ID NO: 796)		
	LSGRSANP (SEQ ID NO: 797)		
	LSGRSANI (SEQ ID NO: 798)		
	LSGRSDNI (SEQ ID NO: 799)		
	MIAPVAYR (SEQ ID NO: 800)		
	RPSPMWAY (SEQ ID NO: 801)		
	WATPRPMR (SEQ ID NO: 802)		
	FRLLDWQW (SEQ ID NO: 803)		
	ISSGL (SEQ ID NO: 804)		
	ISSGLLS (SEQ ID NO: 805)		
	ISSGLL (SEQ ID NO: 806)		
	ISSGLLSGRSANPRG (SEQ ID NO: 680)		
	AVGLLAPPTSGRSANPRG (SEQ ID NO: 681)		

	AVGLLAPPSGRSANPRG (SEQ ID NO: 682)		
	ISSGLLSGRSDDH (SEQ ID NO: 683)		
	ISSGLLSGRSDIH (SEQ ID NO: 684)		
	ISSGLLSGRSDQH (SEQ ID NO: 685)		
	ISSGLLSGRSDTH (SEQ ID NO: 686)		
	ISSGLLSGRSDYH (SEQ ID NO: 687)		
	ISSGLLSGRSDNP (SEQ ID NO: 688)		
	ISSGLLSGRSANP (SEQ ID NO: 689)		
	ISSGLLSGRSANI (SEQ ID NO: 690)		
	AVGLLAPPGGLSGRSDDH (SEQ ID NO: 691)		
	AVGLLAPPGGLSGRSDIH (SEQ ID NO: 692)		
	AVGLLAPPGGLSGRSDQH (SEQ ID NO: 693)		
	AVGLLAPPGGLSGRSDTH (SEQ ID NO: 694)		
	AVGLLAPPGGLSGRSDYH (SEQ ID NO: 695)		
	AVGLLAPPGGLSGRSDNP (SEQ ID NO: 696)		
	AVGLLAPPGGLSGRSANP (SEQ ID NO: 697)		
	AVGLLAPPGGLSGRSANI (SEQ ID NO: 698)		
	ISSGLLSGRSDNI (SEQ ID NO: 713)		
	AVGLLAPPGGLSGRSDNI (SEQ ID NO: 714)		
	GLSGRSDNHGGAVGLLAPP (SEQ ID NO: 807)		
	GLSGRSDNHGGVHMPGLGFLGP (SEQ ID NO: 808)		

[000210] In some embodiments, an activatable antibody of the present disclosure includes one or more polypeptides that include the combination of sequences selected from Table D or Table E, where the polypeptide includes a combination of a masking sequence selected from the column titled “Mask Sequence (MM)” of Table D or Table E, a substrate sequence from the column titled “Substrate Sequence (CM)” of Table D or Table E, a light chain variable domain or light chain CDRs from the column titled “VL or VL CDRs” or

“VL CDRs SEQ ID NOs” of Table D or Table E, and a heavy chain variable domain or heavy chain CDRs from the column titled “VH or VH CDRs” or “VH CDRs SEQ ID Nos” of Table D or Table E. For example, an activatable antibody of the present disclosure may include the amino acid sequences of combination no. 54, which includes the masking sequence of SEQ ID NO: 17, the substrate sequence of SEQ ID NO: 412, a light chain variable domain that includes the VL CDR sequences of SEQ ID NOS: 12, 14, and 15, and a heavy chain variable domain that includes the VH CDR sequences of 9, 10, and 11. Therefore, an activatable antibody that includes at least the combination of sequences in any given row of Table D is described herein. Similarly, any combination of a mask sequence (MM), a substrate sequence (CM), a light chain variable domain sequence or light chain variable domain CDR sequences, and a heavy chain variable domain sequence or heavy chain variable domain CDR sequences of Table E is described herein. An activatable antibody that includes at least any combination of a masking sequence, a substrate sequence, a variable heavy chain or variable heavy chain CDRs, and a variable light chain or variable light chain CDRs selected from the corresponding columns Table D or Table E is also described herein. In some exemplary embodiments, an activatable antibody that includes at least the combination of sequences in any given row of Table D or any combination of a mask sequence (MM), a substrate sequence (CM), a light chain variable domain sequence or light chain variable domain CDR sequences, and a heavy chain variable domain sequence or heavy chain variable domain CDR sequences of Table E can be combined with one or more toxins, including a dolastatin or a derivative thereof, an auristatin or a derivative thereof, a maytansinoid or a derivative thereof, a duocarmycin or a derivative thereof, a calicheamicin or a derivative thereof, or a pyrrolobenzodiazepine or a derivative thereof. In some exemplary embodiments, an activatable antibody that includes at least the combination of sequences in any given row of Table D or any combination of a mask sequence (MM), a substrate sequence (CM), a light chain variable domain sequence or light chain variable domain CDR sequences, and a heavy chain variable domain sequence or heavy chain variable domain CDR sequences of Table E can be combined with one or more toxins, including auristatin E, monomethyl auristatin F (MMAF), monomethyl auristatin E (MMAE), monomethyl auristatin D (MMAD), maytansinoid DM4, maytansinoid DM1, a pyrrolobenzodiazepine, a pyrrolobenzodiazepine dimer, and/or a duocarmycin.

[000211] Any of the combinations in Table D or Table E as described above can be combined with human immunoglobulin constant regions to result in fully human IgGs

including IgG1, IgG2, IgG4 or mutated constant regions to result in human IgGs with altered functions such as IgG1 N297A, IgG1 N297Q, or IgG4 S228P. The combinations described in Table D or Table E are not limited by the particular combinations shown in any given row, and thus may include any mask sequence from column 2 of Table D (or column 1 of Table E) combined with any substrate sequence from column 3 of Table D (or column 2 of Table E) combined with any VL sequence or set of VL CDR sequences from column 4 of Table D (or column 3 of Table E) combined with any VH sequence or set of VH CDR sequences from column 5 of Table D (or column 4 of Table E). In addition to the mask sequences disclosed in column 2 of Table D or column 1 of Table E, any mask sequence disclosed herein can be used in a combination. In addition to the substrate sequences disclosed in column 3 of Table D or column 2 of Table E, any CM disclosed herein can be used in a combination. In addition to the light chain variable region sequence or light chain CDR sequences disclosed in column 4 of Table D or column 3 of Table E, any light chain variable region sequence or light chain CDR sequences disclosed herein can be used in a combination. In addition to the heavy chain variable region sequence or heavy chain CDR sequences disclosed in column 5 of Table D or column 4 of Table E, any heavy chain variable region sequence or heavy chain CDR sequences disclosed herein can be used in a combination.

[000212] In some embodiments, the antibody drug conjugates (ADCs) and activatable antibody drug conjugates (AADCs) can include one or more polypeptides that include the combination of a light chain sequence or a light chain variable domain sequence, and a heavy chain sequence or a heavy chain variable domain sequence, a linker, and a toxin in a given row of Table F or any combination of a light chain sequence or a light chain variable domain sequence, and a heavy chain sequence or a heavy chain variable domain sequence, a linker, and a toxin of Table F.

Table F: Anti-CD71 ADC and Anti-CD71 Activatable ADC Combinations

Comb. No.	Heavy Chain (HC) or HC Variable Region SEQ ID NO.	Light Chain (LC) or LC Variable Region SEQ ID NO.	Linker	Toxin
1	5	7	vc	MMAD
2	5	7	PEG2-vc	MMAD
3	5	7	vc	MMAE
4	5	7	vc	duocarmycin
5	5	7	spdb	DM4

6	325	323	vc	MMAD
7	325	323	PEG2-vc	MMAD
8	325	323	vc	MMAE
9	325	323	vc	duocarmycin
10	325	323	spdb	DM4
11	325	327	vc	MMAD
12	325	327	PEG2-vc	MMAD
13	325	327	vc	MMAE
14	325	327	vc	duocarmycin
15	325	327	spdb	DM4
16	5	810	vc	MMAD
17	5	810	PEG2-vc	MMAD
18	5	810	vc	MMAE
19	5	810	vc	duocarmycin
20	5	810	spdb	DM4
21	325	329	vc	MMAD
22	325	329	PEG2-vc	MMAD
23	325	329	vc	MMAE
24	325	329	vc	duocarmycin
25	325	329	spdb	DM4
26	5	812	vc	MMAD
27	5	812	PEG2-vc	MMAD
28	5	812	vc	MMAE
29	5	812	vc	duocarmycin
30	5	812	spdb	DM4
31	325	331	vc	MMAD
32	325	331	PEG2-vc	MMAD
33	325	331	vc	MMAE
34	325	331	vc	duocarmycin
35	325	331	spdb	DM4
36	5	814	vc	MMAD
37	5	814	PEG2-vc	MMAD
38	5	814	vc	MMAE
39	5	814	vc	duocarmycin
40	5	814	spdb	DM4
41	325	333	vc	MMAD
42	325	333	PEG2-vc	MMAD
43	325	333	vc	MMAE
44	325	333	vc	duocarmycin
45	325	333	spdb	DM4
46	5	816	vc	MMAD
47	5	816	PEG2-vc	MMAD
48	5	816	vc	MMAE
49	5	816	vc	duocarmycin
50	5	816	spdb	DM4
51	325	335	vc	MMAD
52	325	335	PEG2-vc	MMAD
53	325	335	vc	MMAE
54	325	335	vc	duocarmycin

55	325	335	spdb	DM4
56	5	818	vc	MMAD
57	5	818	PEG2-vc	MMAD
58	5	818	vc	MMAE
59	5	818	vc	duocarmycin
60	5	818	spdb	DM4
61	325	337	vc	MMAD
62	325	337	PEG2-vc	MMAD
63	325	337	vc	MMAE
64	325	337	vc	duocarmycin
65	325	337	spdb	DM4
66	5	820	vc	MMAD
67	5	820	PEG2-vc	MMAD
68	5	820	vc	MMAE
69	5	820	vc	duocarmycin
70	5	820	spdb	DM4
71	325	673	vc	MMAD
72	325	673	PEG2-vc	MMAD
73	325	673	vc	MMAE
74	325	673	vc	duocarmycin
75	325	673	spdb	DM4
76	5	824	vc	MMAD
77	5	824	PEG2-vc	MMAD
78	5	824	vc	MMAE
79	5	824	vc	duocarmycin
80	5	824	spdb	DM4
81	325	702	vc	MMAD
82	325	702	PEG2-vc	MMAD
83	325	702	vc	MMAE
84	325	702	vc	duocarmycin
85	325	702	spdb	DM4
86	5	826	vc	MMAD
87	5	826	PEG2-vc	MMAD
88	5	826	vc	MMAE
89	5	826	vc	duocarmycin
90	5	826	spdb	DM4
91	325	671	vc	MMAD
92	325	671	PEG2-vc	MMAD
93	325	671	vc	MMAE
94	325	671	vc	duocarmycin
95	325	671	spdb	DM4
96	5	822	vc	MMAD
97	5	822	PEG2-vc	MMAD
98	5	822	vc	MMAE
99	5	822	vc	duocarmycin
100	5	822	spdb	DM4
101	325	704	vc	MMAD
102	325	704	PEG2-vc	MMAD
103	325	704	vc	MMAE

104	325	704	vc	duocarmycin
105	325	704	spdb	DM4
106	5	828	vc	MMAD
107	5	828	PEG2-vc	MMAD
108	5	828	vc	MMAE
109	5	828	vc	duocarmycin
110	5	828	spdb	DM4
111	325	706	vc	MMAD
112	325	706	PEG2-vc	MMAD
113	325	706	vc	MMAE
114	325	706	vc	duocarmycin
115	325	706	spdb	DM4
116	5	830	vc	MMAD
117	5	830	PEG2-vc	MMAD
118	5	830	vc	MMAE
119	5	830	vc	duocarmycin
120	5	830	spdb	DM4
121	325	708	vc	MMAD
122	325	708	PEG2-vc	MMAD
123	325	708	vc	MMAE
124	325	708	vc	duocarmycin
125	325	708	spdb	DM4
126	5	832	vc	MMAD
127	5	832	PEG2-vc	MMAD
128	5	832	vc	MMAE
129	5	832	vc	duocarmycin
130	5	832	spdb	DM4
131	325	710	vc	MMAD
132	325	710	PEG2-vc	MMAD
133	325	710	vc	MMAE
134	325	710	vc	duocarmycin
135	325	710	spdb	DM4
136	5	834	vc	MMAD
137	5	834	PEG2-vc	MMAD
138	5	834	vc	MMAE
139	5	834	vc	duocarmycin
140	5	834	spdb	DM4
141	325	712	vc	MMAD
142	325	712	PEG2-vc	MMAD
143	325	712	vc	MMAE
144	325	712	vc	duocarmycin
145	325	712	spdb	DM4
146	5	836	vc	MMAD
147	5	836	PEG2-vc	MMAD
148	5	836	vc	MMAE
149	5	836	vc	duocarmycin
150	5	836	spdb	DM4
151	325	323	vc	MMAD
152	325	323	PEG2-vc	MMAD

153	325	323	vc	MMAE
154	325	323	vc	duocarmycin
155	325	323	spdb	DM4
156	325	650	vc	MMAD
157	325	650	PEG2-vc	MMAD
158	325	650	vc	MMAE
159	325	650	vc	duocarmycin
160	325	650	spdb	DM4
161	5	809	vc	MMAD
162	5	809	PEG2-vc	MMAD
163	5	809	vc	MMAE
164	5	809	vc	duocarmycin
165	5	809	spdb	DM4
166	325	652	vc	MMAD
167	325	652	PEG2-vc	MMAD
168	325	652	vc	MMAE
169	325	652	vc	duocarmycin
170	325	652	spdb	DM4
171	5	811	vc	MMAD
172	5	811	PEG2-vc	MMAD
173	5	811	vc	MMAE
174	5	811	vc	duocarmycin
175	5	811	spdb	DM4
176	325	654	vc	MMAD
177	325	654	PEG2-vc	MMAD
178	325	654	vc	MMAE
179	325	654	vc	duocarmycin
180	325	654	spdb	DM4
181	5	813	vc	MMAD
182	5	813	PEG2-vc	MMAD
183	5	813	vc	MMAE
184	5	813	vc	duocarmycin
185	5	813	spdb	DM4
186	325	656	vc	MMAD
187	325	656	PEG2-vc	MMAD
188	325	656	vc	MMAE
189	325	656	vc	duocarmycin
190	325	656	spdb	DM4
191	5	815	vc	MMAD
192	5	815	PEG2-vc	MMAD
193	5	815	vc	MMAE
194	5	815	vc	duocarmycin
195	5	815	spdb	DM4
196	325	658	vc	MMAD
197	325	658	PEG2-vc	MMAD
198	325	658	vc	MMAE
199	325	658	vc	duocarmycin
200	325	658	spdb	DM4
201	5	817	vc	MMAD

202	5	817	PEG2-vc	MMAD
203	5	817	vc	MMAE
204	5	817	vc	duocarmycin
205	5	817	spdb	DM4
206	325	660	vc	MMAD
207	325	660	PEG2-vc	MMAD
208	325	660	vc	MMAE
209	325	660	vc	duocarmycin
210	325	660	spdb	DM4
211	5	819	vc	MMAD
212	5	819	PEG2-vc	MMAD
213	5	819	vc	MMAE
214	5	819	vc	duocarmycin
215	5	819	spdb	DM4
216	325	672	vc	MMAD
217	325	672	PEG2-vc	MMAD
218	325	672	vc	MMAE
219	325	672	vc	duocarmycin
220	325	672	spdb	DM4
221	5	823	vc	MMAD
222	5	823	PEG2-vc	MMAD
223	5	823	vc	MMAE
224	5	823	vc	duocarmycin
225	5	823	spdb	DM4
226	325	701	vc	MMAD
227	325	701	PEG2-vc	MMAD
228	325	701	vc	MMAE
229	325	701	vc	duocarmycin
230	325	701	spdb	DM4
231	5	825	vc	MMAD
232	5	825	PEG2-vc	MMAD
233	5	825	vc	MMAE
234	5	825	vc	duocarmycin
235	5	825	spdb	DM4
236	325	670	vc	MMAD
237	325	670	PEG2-vc	MMAD
238	325	670	vc	MMAE
239	325	670	vc	duocarmycin
240	325	670	spdb	DM4
241	5	821	vc	MMAD
242	5	821	PEG2-vc	MMAD
243	5	821	vc	MMAE
244	5	821	vc	duocarmycin
245	5	821	spdb	DM4
246	325	703	vc	MMAD
247	325	703	PEG2-vc	MMAD
248	325	703	vc	MMAE
249	325	703	vc	duocarmycin
250	325	703	spdb	DM4

251	5	827	vc	MMAD
252	5	827	PEG2-vc	MMAD
253	5	827	vc	MMAE
254	5	827	vc	duocarmycin
255	5	827	spdb	DM4
256	325	705	vc	MMAD
257	325	705	PEG2-vc	MMAD
258	325	705	vc	MMAE
259	325	705	vc	duocarmycin
260	325	705	spdb	DM4
261	5	829	vc	MMAD
262	5	829	PEG2-vc	MMAD
263	5	829	vc	MMAE
264	5	829	vc	duocarmycin
265	5	829	spdb	DM4
266	325	707	vc	MMAD
267	325	707	PEG2-vc	MMAD
268	325	707	vc	MMAE
269	325	707	vc	duocarmycin
270	325	707	spdb	DM4
271	5	831	vc	MMAD
272	5	831	PEG2-vc	MMAD
273	5	831	vc	MMAE
274	5	831	vc	duocarmycin
275	5	831	spdb	DM4
276	325	709	vc	MMAD
277	325	709	PEG2-vc	MMAD
278	325	709	vc	MMAE
279	325	709	vc	duocarmycin
280	325	709	spdb	DM4
281	5	833	vc	MMAD
282	5	833	PEG2-vc	MMAD
283	5	833	vc	MMAE
284	5	833	vc	duocarmycin
285	5	833	spdb	DM4
286	325	711	vc	MMAD
287	325	711	PEG2-vc	MMAD
288	325	711	vc	MMAE
289	325	711	vc	duocarmycin
290	325	711	spdb	DM4
291	5	835	vc	MMAD
292	5	835	PEG2-vc	MMAD
293	5	835	vc	MMAE
294	5	835	vc	duocarmycin
295	5	835	spdb	DM4

[000213] An antibody drug conjugate (ADC) of the present disclosure or activatable antibody drug conjugate (AADC) of the present disclosure may include one or more

polypeptides that include the combination of amino acid sequences, a linker, and a toxin listed in a given row of Table F. Therefore, an activatable antibody drug conjugate (ADC) of the present disclosure or activatable antibody drug conjugate (AADC) of the present disclosure that includes the combination of amino acid sequences, a linker, and a toxin listed in a given row or provided as a specific combination is described herein. For example, an activatable antibody drug conjugate of the present disclosure may include the amino acid sequences of combination no. 45, which includes a heavy chain comprising the amino acid sequence of SEQ ID NO: 325, a light chain comprising the amino acid sequence of SEQ ID NO: 333, and a spdb-DM4 linker-toxin. In another example of the AADCs disclosed and described herein, an activatable antibody drug conjugate of the present disclosure may include the amino acid sequences of combination no. 33, which includes a heavy chain comprising the amino acid sequence of SEQ ID NO: 325, a light chain comprising the amino acid sequence of SEQ ID NO: 331, and a vc-MMAE linker-toxin.

[000214] Any of the combinations in Table F that list a heavy chain and light chain variable region can be combined with human immunoglobulin constant regions to result in fully human IgGs including IgG1, IgG2, IgG4 or mutated constant regions to result in human IgGs with altered functions such as IgG1 N297A, IgG1 N297Q, or IgG4 S228P. The combinations described in Table F are not limited by the particular combinations shown in any given row, and thus can include any heavy chain sequence or heavy chain variable region sequence from column 2 of Table F combined with any light chain sequence or light chain variable region sequence from column 3 of Table F combined with any linker from column 4 combined with any toxin from column 5. In addition to the heavy chain sequences or heavy chain variable region sequences listed in column 2, any heavy chain sequence or heavy chain variable region sequence disclosed herein can be used in a combination. In addition to the light chain sequences or light chain variable region sequences listed in column 3, any light chain sequence or light chain variable region sequence disclosed herein can be used in a combination. In addition to the linkers listed in column 4, any linker disclosed herein can be used in a combination. In addition to the toxins listed in column 5, any toxin disclosed herein can be used in a combination.

[000215] In some embodiments, the serum half-life of the activatable antibody is longer than that of the corresponding antibody; e.g., the pK of the activatable antibody is longer than that of the corresponding antibody. In some embodiments, the serum half-life of the activatable antibody is similar to that of the corresponding antibody. In some

[illegible]

[000216] In some embodiments, the activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody is monospecific. In some embodiments, the activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody is multispecific, *e.g.*, by way of non-limiting example, bispecific or trifunctional. In some embodiments, the activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody is formulated as part of a pro-Bispecific T Cell Engager (BITE) molecule. In some embodiments, the activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody is formulated as part of a pro-Chimeric Antigen Receptor (CAR) modified T cell or other engineered receptor.

[000217] In some embodiments, the activatable antibody or antigen-binding fragment thereof is incorporated in a multispecific activatable antibody or antigen-binding fragment thereof, where at least one arm of the multispecific activatable antibody specifically binds CD71. In some embodiments, the activatable antibody or antigen-binding fragment thereof is incorporated in a bispecific antibody or antigen-binding fragment thereof, where at least one arm of the bispecific activatable antibody specifically binds CD71.

[000218] In some embodiments, the anti-CD71 antibodies, conjugated anti-CD71 antibodies, activatable anti-CD71 antibodies and/or conjugated activatable anti-CD71 antibodies described herein are used in conjunction with one or more additional agents or a combination of additional agents. Suitable additional agents include current pharmaceutical and/or surgical therapies for an intended application, such as, for example, cancer. For example, the anti-CD71 antibodies, conjugated anti-CD71 antibodies, activatable anti-CD71 antibodies and/or conjugated activatable anti-CD71 antibodies can be used in conjunction with an additional chemotherapeutic or anti-neoplastic agent.

[000219] In some embodiments, the additional agent(s) is a chemotherapeutic agent, such as a chemotherapeutic agent selected from the group consisting of docetaxel, paclitaxel, abraxane (*i.e.*, albumin-conjugated paclitaxel), doxorubicin, oxaliplatin, carboplatin, cisplatin, irinotecan, and gemcitabine.

[000220] In some embodiments, the additional agent(s) is a checkpoint inhibitor, a kinase inhibitor, an agent targeting inhibitors in the tumor microenvironment, and/or a T cell or NK agonist. In some embodiments, the additional agent(s) is radiation therapy, alone or in combination with another additional agent(s) such as a chemotherapeutic or anti-neoplastic agent. In some embodiments, the additional agent(s) is a vaccine, an oncovirus, and/or a DC-activating agent such as, by way of non-limiting example, a toll-like receptor

(TLR) agonist and/or α -CD40. In some embodiments, the additional agent(s) is a tumor-targeted antibody designed to kill the tumor via ADCC or via direct conjugation to a toxin (e.g., an antibody drug conjugate (ADC)).

[000221] In some embodiments, the checkpoint inhibitor is an inhibitor of a target selected from the group consisting of CTLA-4, LAG-3, PD-1, CD71, TIGIT, TIM-3, B7H4, and Vista. In some embodiments, the kinase inhibitor is selected from the group consisting of B-RAFi, MEKi, and Btk inhibitors, such as ibrutinib. In some embodiments, the kinase inhibitor is crizotinib. In some embodiments, the tumor microenvironment inhibitor is selected from the group consisting of an IDO inhibitor, an α -CSF1R inhibitor, an α -CCR4 inhibitor, a TGF-beta, a myeloid-derived suppressor cell, or a T-regulatory cell. In some embodiments, the agonist is selected from the group consisting of Ox40, GITR, CD137, ICOS, CD27, and HVEM.

[000222] In some embodiments, the inhibitor is a CTLA-4 inhibitor. In some embodiments, the inhibitor is a LAG-3 inhibitor. In some embodiments, the inhibitor is a PD-1 inhibitor. In some embodiments, the inhibitor is a CD71 inhibitor. In some embodiments, the inhibitor is a TIGIT inhibitor. In some embodiments, the inhibitor is a TIM-3 inhibitor. In some embodiments, the inhibitor is a B7H4 inhibitor. In some embodiments, the inhibitor is a Vista inhibitor. In some embodiments, the inhibitor is a B-RAFi inhibitor. In some embodiments, the inhibitor is a MEKi inhibitor. In some embodiments, the inhibitor is a Btk inhibitor. In some embodiments, the inhibitor is ibrutinib. In some embodiments, the inhibitor is crizotinib. In some embodiments, the inhibitor is an IDO inhibitor. In some embodiments, the inhibitor is an α -CSF1R inhibitor. In some embodiments, the inhibitor is an α -CCR4 inhibitor. In some embodiments, the inhibitor is a TGF-beta. In some embodiments, the inhibitor is a myeloid-derived suppressor cell. In some embodiments, the inhibitor is a T-regulatory cell.

[000223] In some embodiments, the agonist is Ox40. In some embodiments, the agonist is GITR. In some embodiments, the agonist is CD137. In some embodiments, the agonist is ICOS. In some embodiments, the agonist is CD27. In some embodiments, the agonist is HVEM.

[000224] In some embodiments, the anti-CD71 antibody, conjugated antibody, activatable antibody and/or conjugated activatable antibody is administered during and/or after treatment in combination with one or more additional agents such as, for example, a chemotherapeutic agent, an anti-inflammatory agent, and/or an immunosuppressive agent.

In some embodiments, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and the additional agent are formulated into a single therapeutic composition, and the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and additional agent are administered simultaneously. Alternatively, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and additional agent are separate from each other, *e.g.*, each is formulated into a separate therapeutic composition, and the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and the additional agent are administered simultaneously, or the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and the additional agent are administered at different times during a treatment regimen. For example, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody is administered prior to the administration of the additional agent, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody is administered subsequent to the administration of the additional agent, or the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and the additional agent are administered in an alternating fashion. As described herein, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and additional agent are administered in single doses or in multiple doses.

[000225] In some embodiments, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and the additional agent(s) are administered simultaneously. For example, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and the additional agent(s) can be formulated in a single composition or administered as two or more separate compositions. In some embodiments, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and the additional agent(s) are administered sequentially, or the anti-CD71 antibody, conjugated anti-CD71

antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and the additional agent are administered at different times during a treatment regimen.

[000226] In some embodiments, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody is administered during and/or after treatment in combination with one or more additional agents such as, by way of non-limiting example, a chemotherapeutic agent, an anti-inflammatory agent, and/or an immunosuppressive agent, such as an alkylating agent, an anti-metabolite, an anti-microtubule agent, a topoisomerase inhibitor, a cytotoxic antibiotic, and/or any other nucleic acid damaging agent. In some embodiments, the additional agent is a taxane, such as paclitaxel (e.g., Abraxane®). In some embodiments, the additional agent is an anti-metabolite, such as gemcitabine. In some embodiments, the additional agent is an alkylating agent, such as platinum-based chemotherapy, such as carboplatin or cisplatin. In some embodiments, the additional agent is a targeted agent, such as a kinase inhibitor, e.g., sorafenib or erlotinib. In some embodiments, the additional agent is a targeted agent, such as another antibody, e.g., a monoclonal antibody (e.g., bevacizumab), a bispecific antibody, or a multispecific antibody. In some embodiments, the additional agent is a proteasome inhibitor, such as bortezomib or carfilzomib. In some embodiments, the additional agent is an immune modulating agent, such as lenolidomide or IL-2. In some embodiments, the additional agent is radiation. In some embodiments, the additional agent is an agent considered standard of care by those skilled in the art. In some embodiments, the additional agent is a chemotherapeutic agent well known to those skilled in the art.

[000227] In some embodiments, the additional agent is another antibody or antigen-binding fragment thereof, another conjugated antibody or antigen-binding fragment thereof, another activatable antibody or antigen-binding fragment thereof and/or another conjugated activatable antibody or antigen-binding fragment thereof. In some embodiments the additional agent is another antibody or antigen-binding fragment thereof, another conjugated antibody or antigen-binding fragment thereof, another activatable antibody or antigen-binding fragment thereof and/or another conjugated activatable antibody or antigen-binding fragment thereof against the same target as the first antibody or antigen-binding fragment thereof, the first conjugated antibody or antigen-binding fragment thereof, activatable antibody or antigen-binding fragment thereof and/or a conjugated activatable antibody or antigen-binding fragment thereof, e.g., against CD71. In some embodiments the additional agent is another antibody or antigen-binding fragment thereof, another

conjugated antibody or antigen-binding fragment thereof, another activatable antibody or antigen-binding fragment thereof and/or another conjugated activatable antibody or antigen-binding fragment thereof against a target different than the target of the first antibody or antigen-binding fragment thereof, the first conjugated antibody or antigen-binding fragment thereof, activatable antibody or antigen-binding fragment thereof and/or a conjugated activatable antibody or antigen-binding fragment thereof.

[000228] As a non-limiting example, the antibody or antigen-binding fragment and/or the AB of an activatable antibody is a binding partner for any target listed in Table 1.

Table 1: Exemplary Targets

1-92-LFA-3	CD52	DL44	HVEM	LIF-R	STEAP1
Alpha-4 integrin	CD56	DLK1	Hyaluronidase	Lewis X	STEAP2
Alpha-V integrin	CD64	DLL4	ICOS	LIGHT	TAG-72
alpha4beta1 integrin	CD70	DPP-4	IFNalpha	LRP4	TAPA1
alpha4beta7 integrin	CD71	DSG1	IFNbeta	LRRC26	TGFbeta
AGR2	CD74	EGFR	IFNgamma	MCSP	TIGIT
Anti-Lewis-Y		EGFRviii	IgE	Mesothelin	TIM-3
Apelin J receptor	CD80	Endothelin B receptor (ETBR)	IgE Receptor (FcεRI)	MRP4	TLR2
APRIL	CD81	ENPP3	IGF	MUC1	TLR4
B7-H4	CD86	EpCAM	IGF1R	Mucin-16 (MUC16, CA-125)	TLR6
BAFF	CD95	EPHA2	IL1B	Na/K ATPase	TLR7
BTLA	CD117	EPHB2	IL1R	Neutrophil elastase	TLR8
C5 complement	CD125	ERBB3	IL2	NGF	TLR9
C-242	CD132 (IL-2RG)	F protein of RSV	IL11	Nicastrin	TMEM31
CA9	CD133	FAP	IL12	Notch Receptors	TNFAalpha
CA19-9 (Lewis a)	CD137	FGF-2	IL12p40	Notch 1	TNFR
Carbonic anhydrase 9	CD138	FGF8	IL-12R, IL-12Rbeta1	Notch 2	TNFRS12 A
CD2	CD166	FGFR1	IL13	Notch 3	TRAIL-R1
CD3	CD172A	FGFR2	IL13R	Notch 4	TRAIL-R2
CD6	CD248	FGFR3	IL15	NOV	Transferrin

CD9	CDH6	FGFR4	IL17	OSM-R	Transferrin receptor
CD11a	CEACAM5 (CEA)	Folate receptor	IL18	OX-40	TRK-A
CD19	CEACAM6 (NCA-90)	GAL3ST1	IL21	PAR2	TRK-B
CD20	CLAUDIN-3	G-CSF	IL23	PDGF-AA	uPAR
CD22	CLAUDIN-4	G-CSFR	IL23R	PDGF-BB	VAP1
CD24	cMet	GD2	IL27/IL27R (wsx1)	PDGFRalpha	VCAM-1
CD25	Collagen	GITR	IL29	PDGFRbeta	VEGF
CD27	Cripto	GLUT1	IL-31R	PD-1	VEGF-A
CD28	CSFR	GLUT4	IL31/IL31R	PD-L1	VEGF-B
CD30	CSFR-1	GM-CSF	IL2R	PD-L2	VEGF-C
CD33	CTLA-4	GM-CSFR	IL4	Phosphatidyl-serine	VEGF-D
CD38	CTGF	GP IIb/IIIa receptors	IL4R	P1GF	VEGFR1
CD40	CXCL10	Gp130	IL6, IL6R	PSCA	VEGFR2
CD40L	CXCL13	GPIIB/IIIA	Insulin Receptor	PSMA	VEGFR3
CD41	CXCR1	GPNMB	Jagged Ligands	RAAG12	VISTA
CD44	CXCR2	GRP78	Jagged 1	RAGE	WISP-1
CD44v6		HER2/neu	Jagged 2	SLC44A4	WISP-2
CD47	CXCR4	HGF	LAG-3	Sphingosine 1 Phosphate	WISP-3
CD51	CYR61	hGH			

[000229] As a non-limiting example, the antibody or antigen-binding fragment and/or the AB of an activatable antibody is or is derived from an antibody listed in Table 2.

Table 2: Exemplary sources for Abs

Antibody Trade Name (antibody name)	Target
Avastin™ (bevacizumab)	VEGF
Lucentis™ (ranibizumab)	VEGF
Erbix™ (cetuximab)	EGFR
Vectibix™ (panitumumab)	EGFR
Remicade™ (infliximab)	TNF α
Humira™ (adalimumab)	TNF α
Tysabri™ (natalizumab)	Integrin α 4
Simulect™ (basiliximab)	IL2R
Soliris™ (eculizumab)	Complement C5
Raptiva™ (efalizumab)	CD11a
Bexxar™ (tositumomab)	CD20
Zevalin™ (ibritumomab tiuxetan)	CD20
Rituxan™ (rituximab)	CD20

Ocrelizumab	CD20
Arzerra™ (ofatumumab)	CD20
Gazyva™ (obinutuzumab)	CD20
Zenapax™ (daclizumab)	CD25
Adcetris™ (brentuximab vedotin)	CD30
Myelotarg™ (gemtuzumab)	CD33
Mylotarg™ (gemtuzumab ozogamicin)	CD33
Campath™ (alemtuzumab)	CD52
ReoPro™ (abixicimab)	Glycoprotein receptor IIb/IIIa
Xolair™ (omalizumab)	IgE
Herceptin™ (trastuzumab)	Her2
Kadcyla™ (trastuzumab emtansine)	Her2
Synagis™ (palivizumab)	F protein of RSV
(ipilimumab)	CTLA-4
(tremelimumab)	CTLA-4
Hu5c8	CD40L
(pertuzumab)	Her2-neu
(ertumaxomab)	CD3/Her2-neu
Orencia™ (abatacept)	CTLA-4
(tanezumab)	NGF
(bavituximab)	Phosphatidylserine
(zalutumumab)	EGFR
(mapatumumab)	EGFR
(matuzumab)	EGFR
(nimotuzumab)	EGFR
ICR62	EGFR
mAb 528	EGFR
CH806	EGFR
MDX-447	EGFR/CD64
(edrecolomab)	EpCAM
RAV12	RAAG12
huJ591	PSMA
Enbrel™ (etanercept)	TNF-R
Amevive™ (alefacept)	1-92-LFA-3
Antril™, Kineret™ (ankinra)	IL-1Ra
GC1008	TGFbeta
	Notch, e.g., Notch 1
	Jagged 1 or Jagged 2
(adecatumumab)	EpCAM
(figitumumab)	IGF1R
(tocilizumab)	IL-6 receptor
Stelara™ (ustekinumab)	IL-12/IL-23
Prolia™ (denosumab)	RANKL

[000230] In some embodiments, the additional antibody or antigen binding fragment thereof, conjugated antibody or antigen binding fragment thereof, activatable antibody or antigen binding fragment thereof, and/or conjugated activatable antibody or antigen binding

fragment thereof is a monoclonal antibody, domain antibody, single chain, Fab fragment, a $F(ab')_2$ fragment, a scFv, a scAb, a dAb, a single domain heavy chain antibody, or a single domain light chain antibody. In some embodiments, the additional antibody or antigen binding fragment thereof, conjugated antibody or antigen binding fragment thereof, activatable antibody or antigen binding fragment thereof, and/or conjugated activatable antibody or antigen binding fragment thereof is a mouse, other rodent, chimeric, humanized or fully human monoclonal antibody.

[000231] The disclosure also provides methods of producing an anti-CD71 antibody and/or activatable anti-CD71 antibody polypeptide by culturing a cell under conditions that lead to expression of the polypeptide, wherein the cell comprises an isolated nucleic acid molecule encoding an antibody and/or an activatable antibody described herein, and/or vectors that include these isolated nucleic acid sequences. The disclosure provides methods of producing an antibody and/or activatable antibody by culturing a cell under conditions that lead to expression of the antibody and/or activatable antibody, wherein the cell comprises an isolated nucleic acid molecule encoding an antibody and/or an activatable antibody described herein, and/or vectors that include these isolated nucleic acid sequences.

[000232] The invention also provides a method of manufacturing activatable antibodies that in an activated state binds CD71 by (a) culturing a cell comprising a nucleic acid construct that encodes the activatable antibody under conditions that lead to expression of the activatable antibody, wherein the activatable antibody comprises a masking moiety (MM), a cleavable moiety (CM), and an antibody or an antigen binding fragment thereof (AB) that specifically binds CD71, (i) wherein the CM is a polypeptide that functions as a substrate for a protease; and (ii) wherein the CM is positioned in the activatable antibody such that, when the activatable antibody is in an uncleaved state, the MM interferes with specific binding of the AB to CD71 and in a cleaved state the MM does not interfere or compete with specific binding of the AB to CD71; and (b) recovering the activatable antibody. Suitable AB, MM, and/or CM include any of the AB, MM, and/or CM disclosed herein.

[000233] In some embodiments, the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM. In some embodiments, the activatable antibody comprises a linking peptide between the MM and the CM. In some embodiments, the activatable antibody comprises a linking peptide between the CM and the AB. In some embodiments, the activatable antibody

comprises a first linking peptide (LP1) and a second linking peptide (LP2), and wherein the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-LP1-CM-LP2-AB or AB-LP2-CM-LP1-MM. In some embodiments, the two linking peptides need not be identical to each other. In some embodiments, the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: spacer-MM-LP1-CM-LP2-AB or AB-LP2-CM-LP1-MM-spacer.

[000234] In some embodiments, at least one of LP1 or LP2 comprises an amino acid sequence selected from the group consisting of (GS)_n, (GGS)_n, (GSGGS)_n (SEQ ID NO: 339) and (GGGS)_n (SEQ ID NO: 340), where n is an integer of at least one.

[000235] In some embodiments, at least one of LP1 or LP2 comprises an amino acid sequence selected from the group consisting of GGSG (SEQ ID NO: 341), GGSGG (SEQ ID NO: 342), GSGSG (SEQ ID NO: 343), GSGGG (SEQ ID NO: 344), GGGSG (SEQ ID NO: 345), and GSSSG (SEQ ID NO: 346).

[000236] In some embodiments, LP1 comprises the amino acid sequence GSSGGSGGSGGSG (SEQ ID NO: 347), GSSGGSGGSGG (SEQ ID NO: 348), GSSGGSGGSGGS (SEQ ID NO: 349), GSSGGSGGSGGSGGGS (SEQ ID NO: 350), GSSGGSGGSG (SEQ ID NO: 351), or GSSGGSGGSGS (SEQ ID NO: 352).

[000237] In some embodiments, LP2 comprises the amino acid sequence GSS, GGS, GGGS (SEQ ID NO: 353), GSSGT (SEQ ID NO: 354) or GSSG (SEQ ID NO: 356).

[000238] The invention provides methods of preventing, delaying the progression of, treating, alleviating a symptom of, or otherwise ameliorating an CD71 mediated disease in a subject by administering a therapeutically effective amount of an anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody described herein to a subject in need thereof.

[000239] The invention also provides methods of preventing, delaying the progression of, treating, alleviating a symptom of, or otherwise ameliorating cancer in a subject by administering a therapeutically effective amount of an anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody described herein to a subject in need thereof. CD71 is known to be expressed in a variety of cancers, such as, by way of non-limiting example, adenocarcinoma, bile duct (biliary) cancer, bladder cancer, breast cancer, e.g., triple-negative breast cancer and Her2-negative breast cancer; carcinoid cancer; cervical cancer; cholangiocarcinoma; colorectal;

endometrial; glioma; head and neck cancer, e.g., head and neck squamous cell cancer; leukemia; liver cancer; lung cancer, e.g., NSCLC, SCLC; lymphoma; melanoma; oropharyngeal cancer; ovarian cancer; pancreatic cancer; prostate cancer, e.g., metastatic castration-resistant prostate carcinoma; renal cancer; skin cancer; squamous cell cancer, stomach cancer; testis cancer; thyroid cancer; and urothelial cancer.

[000240] In some embodiments, the cancer is associated with a CD71-expressing tumor. In some embodiments, the cancer is due to a CD71-expressing tumor.

[000241] An anti-CD71 antibody, a conjugated anti-CD71 antibody, an activatable anti-CD71 antibody and/or a conjugated activatable anti-CD71 antibody used in any of the embodiments of these methods and uses can be administered at any stage of the disease. For example, such an anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody can be administered to a patient suffering cancer of any stage, from early to metastatic. The terms subject and patient are used interchangeably herein.

[000242] In some embodiments, the subject is a mammal, such as a human, non-human primate, companion animal (e.g., cat, dog, horse), farm animal, work animal, or zoo animal. In some embodiments, the subject is a human. In some embodiments, the subject is a companion animal. In some embodiments, the subject is an animal in the care of a veterinarian.

[000243] The anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and therapeutic formulations thereof are administered to a subject suffering from or susceptible to a disease or disorder associated with aberrant CD71 expression and/or activity. A subject suffering from or susceptible to a disease or disorder associated with aberrant CD71 expression and/or activity is identified using any of a variety of methods known in the art. For example, subjects suffering from cancer or other neoplastic condition are identified using any of a variety of clinical and/or laboratory tests such as, physical examination and blood, urine and/or stool analysis to evaluate health status. For example, subjects suffering from inflammation and/or an inflammatory disorder are identified using any of a variety of clinical and/or laboratory tests such as physical examination and/or bodily fluid analysis, e.g., blood, urine and/or stool analysis, to evaluate health status.

[000244] Administration of an anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody to a

patient suffering from a disease or disorder associated with aberrant CD71 expression and/or activity is considered successful if any of a variety of laboratory or clinical objectives is achieved. For example, administration of an anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody to a patient suffering from a disease or disorder associated with aberrant CD71 expression and/or activity is considered successful if one or more of the symptoms associated with the disease or disorder is alleviated, reduced, inhibited or does not progress to a further, *i.e.*, worse, state. Administration of an anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody to a patient suffering from a disease or disorder associated with aberrant CD71 expression and/or activity is considered successful if the disease or disorder enters remission or does not progress to a further, *i.e.*, worse, state.

[000245] In some embodiments, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and therapeutic formulations thereof are administered to a subject suffering from or susceptible to a disease or disorder, such as subjects suffering from cancer or other neoplastic condition, wherein the subject's diseased cells are expressing CD71. In some embodiments, the diseased cells are associated with aberrant CD71 expression and/or activity. In some embodiments, the diseased cells are associated with normal CD71 expression and/or activity. A subject suffering from or susceptible to a disease or disorder wherein the subject's diseased cells express CD71 is identified using any of a variety of methods known in the art. For example, subjects suffering from cancer or other neoplastic condition are identified using any of a variety of clinical and/or laboratory tests such as, physical examination and blood, urine and/or stool analysis to evaluate health status. For example, subjects suffering from inflammation and/or an inflammatory disorder are identified using any of a variety of clinical and/or laboratory tests such as physical examination and/or bodily fluid analysis, *e.g.*, blood, urine and/or stool analysis, to evaluate health status.

[000246] In some embodiments, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and therapeutic formulations thereof are administered to a subject suffering from or susceptible to a disease or disorder associated with cells expressing CD71 or the presence, growth, proliferation, metastasis, and/or activity of such cells, such as subjects suffering

from cancer or other neoplastic conditions. In some embodiments, the cells are associated with aberrant CD71 expression and/or activity. In some embodiments, the cells are associated with normal CD71 expression and/or activity. A subject suffering from or susceptible to a disease or disorder associated with cells that express CD71 is identified using any of a variety of methods known in the art. For example, subjects suffering from cancer or other neoplastic condition are identified using any of a variety of clinical and/or laboratory tests such as, physical examination and blood, urine and/or stool analysis to evaluate health status. For example, subjects suffering from inflammation and/or an inflammatory disorder are identified using any of a variety of clinical and/or laboratory tests such as physical examination and/or bodily fluid analysis, e.g., blood, urine and/or stool analysis, to evaluate health status.

[000247] Administration of an anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody to a patient suffering from a disease or disorder associated with cells expressing CD71 is considered successful if any of a variety of laboratory or clinical objectives is achieved. For example, administration of an anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody to a patient suffering from a disease or disorder associated with cells expressing CD71 is considered successful if one or more of the symptoms associated with the disease or disorder is alleviated, reduced, inhibited or does not progress to a further, *i.e.*, worse, state. Administration of an anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody to a patient suffering from a disease or disorder associated with cells expressing CD71 is considered successful if the disease or disorder enters remission or does not progress to a further, *i.e.*, worse, state.

[000248] In some embodiments, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody is administered during and/or after treatment in combination with one or more additional agents such as, for example, a chemotherapeutic agent, an anti-inflammatory agent, and/or an immunosuppressive agent. In some embodiments, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and the additional agent(s) are administered simultaneously. For example, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and the additional agent(s) can be

formulated in a single composition or administered as two or more separate compositions. In some embodiments, the anti-CD71 antibody, conjugated anti-CD71 antibody, activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody and the additional agent(s) are administered sequentially.

[000249] The disclosure also provides methods of treating, alleviating a symptom of, or delaying the progression of a disorder or disease in which diseased cells express CD71 comprising administering a therapeutically effective amount of an antibody of the disclosure or a conjugated antibody of the disclosure or a pharmaceutical composition comprising an antibody of the disclosure or a pharmaceutical composition comprising a conjugated antibody of the disclosure to a subject in need thereof. In some embodiments, the disorder or disease is cancer.

[000250] The disclosure also provides methods of treating, alleviating a symptom of, or delaying the progression of a disorder or disease associated with cells expressing CD71 comprising administering a therapeutically effective amount of an antibody of the disclosure or a conjugated antibody of the disclosure or a pharmaceutical composition comprising an antibody of the disclosure or a pharmaceutical composition comprising a conjugated antibody of the disclosure to a subject in need thereof. In some embodiments, the disorder or disease associated with cells expressing CD71 is cancer. In some embodiments, the cancer is an adenocarcinoma, a bile duct (biliary) cancer, a bladder cancer, a bone cancer, a breast cancer, a triple-negative breast cancer, a Her2-negative breast cancer, a carcinoid cancer, a cervical cancer, a cholangiocarcinoma, a colorectal cancer, a colon cancer, an endometrial cancer, a glioma, a head and neck cancer, a head and neck squamous cell cancer, a leukemia, a liver cancer, a lung cancer, a non-small cell lung cancer, a small cell lung cancer, a lymphoma, a melanoma, an oropharyngeal cancer, an ovarian cancer, a pancreatic cancer, a prostate cancer, a metastatic castration-resistant prostate carcinoma, a renal cancer, a sarcoma, a skin cancer, a squamous cell cancer, a stomach cancer, a testis cancer, a thyroid cancer, a urogenital cancer, or a urothelial cancer. In some embodiments, the natural ligand is transferrin. In some embodiments, the expression and/or activity of the mammalian CD71 is aberrant. In some embodiments, the method comprises administering an additional agent. In some embodiments, the additional agent is a therapeutic agent.

[000251] The disclosure also provides methods of inhibiting or reducing the growth, proliferation, or metastasis of cells expressing mammalian CD71 comprising administering a therapeutically effective amount of an antibody of the disclosure or a conjugated antibody

of the disclosure or a pharmaceutical composition comprising an antibody of the disclosure or a pharmaceutical composition comprising a conjugated antibody of the disclosure to a subject in need thereof. In some embodiments, the natural ligand is transferrin. In some embodiments, the expression and/or activity of the mammalian CD71 is aberrant. In some embodiments, the method comprises administering an additional agent. In some embodiments, the additional agent is a therapeutic agent.

[000252] The disclosure also provides methods of inhibiting, blocking, or preventing the binding of a natural ligand to mammalian CD71, comprising administering a therapeutically effective amount of an antibody of the disclosure or a conjugated antibody of the disclosure or a pharmaceutical composition comprising an antibody of the disclosure or a pharmaceutical composition comprising a conjugated antibody of the disclosure to a subject in need thereof. In some embodiments, the natural ligand is transferrin. In some embodiments, the expression and/or activity of the mammalian CD71 is aberrant. In some embodiments, the method comprises administering an additional agent. In some embodiments, the additional agent is a therapeutic agent.

[000253] The disclosure also provides methods of treating, alleviating a symptom of, or delaying the progression of a disorder or disease in which diseased cells express CD71 comprising administering a therapeutically effective amount of an activatable antibody of the disclosure or a conjugated activatable antibody of the disclosure or a pharmaceutical composition comprising an activatable antibody of the disclosure or a pharmaceutical composition comprising a conjugated activatable antibody of the disclosure to a subject in need thereof. In some embodiments, the disorder or disease is cancer.

[000254] The disclosure also provides methods of treating, alleviating a symptom of, or delaying the progression of a disorder or disease associated with cells expressing CD71 comprising administering a therapeutically effective amount of an activatable antibody of the disclosure or a conjugated activatable antibody of the disclosure or a pharmaceutical composition comprising an activatable antibody of the disclosure or a pharmaceutical composition comprising a conjugated activatable antibody of the disclosure to a subject in need thereof. In some embodiments, the disorder or disease associated with cells expressing CD71 is cancer. In some embodiments, the cancer is an adenocarcinoma, a bile duct (biliary) cancer, a bladder cancer, a bone cancer, a breast cancer, a triple-negative breast cancer, a Her2-negative breast cancer, a carcinoid cancer, a cervical cancer, a cholangiocarcinoma, a colorectal cancer, a colon cancer, an endometrial cancer, a glioma, a

head and neck cancer, a head and neck squamous cell cancer, a leukemia, a liver cancer, a lung cancer, a non-small cell lung cancer, a small cell lung cancer, a lymphoma, a melanoma, an oropharyngeal cancer, an ovarian cancer, a pancreatic cancer, a prostate cancer, a metastatic castration-resistant prostate carcinoma, a renal cancer, a sarcoma, a skin cancer, a squamous cell cancer, a stomach cancer, a testis cancer, a thyroid cancer, a urogenital cancer, or a urothelial cancer. In some embodiments, the natural ligand is transferrin. In some embodiments, the expression and/or activity of the mammalian CD71 is aberrant. In some embodiments, the method comprises administering an additional agent. In some embodiments, the additional agent is a therapeutic agent.

[000255] The disclosure also provides methods of inhibiting or reducing the growth, proliferation, or metastasis of cells expressing mammalian CD71 comprising administering a therapeutically effective amount of an activatable antibody of the disclosure or a conjugated activatable antibody of the disclosure or a pharmaceutical composition comprising an activatable antibody of the disclosure or a pharmaceutical composition comprising a conjugated activatable antibody of the disclosure to a subject in need thereof. In some embodiments, the natural ligand is transferrin. In some embodiments, the expression and/or activity of the mammalian CD71 is aberrant. In some embodiments, the method comprises administering an additional agent. In some embodiments, the additional agent is a therapeutic agent.

[000256] The disclosure also provides methods of inhibiting, blocking, or preventing the binding of a natural ligand to mammalian CD71, comprising administering a therapeutically effective amount of an activatable antibody of the disclosure or a conjugated activatable antibody of the disclosure or a pharmaceutical composition comprising an activatable antibody of the disclosure or a pharmaceutical composition comprising a conjugated activatable antibody of the disclosure to a subject in need thereof. In some embodiments, the natural ligand is transferrin. In some embodiments, the expression and/or activity of the mammalian CD71 is aberrant. In some embodiments, the method comprises administering an additional agent. In some embodiments, the additional agent is a therapeutic agent.

[000257] The invention also provides methods and kits for using the activatable anti-CD71 antibodies and/or conjugated activatable anti-CD71 antibodies in a variety of diagnostic and/or prophylactic indications. For example, the invention provides methods and kits for detecting the presence or absence of a cleaving agent and a target of interest in a

subject or a sample by (i) contacting a subject or sample with an anti-CD71 activatable antibody, wherein the anti-CD71 activatable antibody comprises a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, and an antigen binding domain or fragment thereof (AB) that specifically binds the target of interest, wherein the anti-CD71 activatable antibody in an uncleaved, non-activated state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to CD71, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB and is not a modified form of a natural binding partner of the AB; and (b) wherein, when the AB is in an uncleaved, non-activated state, the MM interferes with specific binding of the AB to CD71, and when the AB is in a cleaved, activated state the MM does not interfere or compete with specific binding of the AB to CD71; and (ii) measuring a level of activated anti-CD71 activatable antibody in the subject or sample, wherein a detectable level of activated anti-CD71 activatable antibody in the subject or sample indicates that the cleaving agent and CD71 are present in the subject or sample and wherein no detectable level of activated anti-CD71 activatable antibody in the subject or sample indicates that the cleaving agent, CD71 or both the cleaving agent and CD71 are absent in the subject or sample.

[000258] In some embodiments, the activatable anti-CD71 antibody is an activatable anti-CD71 antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable anti-CD71 antibody is not conjugated to an agent. In some embodiments, the activatable anti-CD71 antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable anti-CD71 antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000259] In some embodiments of these methods and kits, the activatable anti-CD71 antibody includes a detectable label. In some embodiments of these methods and kits, the detectable label includes an imaging agent, a contrasting agent, an enzyme, a fluorescent label, a chromophore, a dye, one or more metal ions, or a ligand-based label. In some embodiments of these methods and kits, the imaging agent comprises a radioisotope. In some embodiments of these methods and kits, the radioisotope is indium or technetium. In some embodiments of these methods and kits, the contrasting agent comprises iodine,

gadolinium or iron oxide. In some embodiments of these methods and kits, the enzyme comprises horseradish peroxidase, alkaline phosphatase, or β -galactosidase. In some embodiments of these methods and kits, the fluorescent label comprises yellow fluorescent protein (YFP), cyan fluorescent protein (CFP), green fluorescent protein (GFP), modified red fluorescent protein (mRFP), red fluorescent protein tdimer2 (RFP tdimer2), HCRED, or a europium derivative. In some embodiments of these methods and kits, the luminescent label comprises an N-methylacrydium derivative. In some embodiments of these methods, the label comprises an Alexa Fluor[®] label, such as Alex Fluor[®] 680 or Alexa Fluor[®] 750. In some embodiments of these methods and kits, the ligand-based label comprises biotin, avidin, streptavidin or one or more haptens.

[000260] In some embodiments of these methods and kits, the subject is a mammal. In some embodiments of these methods, the subject is a human. In some embodiments, the subject is a non-human mammal, such as a non-human primate, companion animal (e.g., cat, dog, horse), farm animal, work animal, or zoo animal. In some embodiments, the subject is a rodent.

[000261] In some embodiments of these methods and kits, the method is an *in vivo* method. In some embodiments of these methods, the method is an *in situ* method. In some embodiments of these methods, the method is an *ex vivo* method. In some embodiments of these methods, the method is an *in vitro* method.

[000262] In some embodiments of the methods and kits, the method is used to identify or otherwise refine a patient population suitable for treatment with an anti-CD71 activatable antibody of the disclosure, followed by treatment by administering that activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody to a subject in need thereof. For example, patients that test positive for both the target (e.g., CD71) and a protease that cleaves the substrate in the cleavable moiety (CM) of the anti-CD71 activatable antibody being tested in these methods are identified as suitable candidates for treatment with such an anti-CD71 activatable antibody comprising such a CM, and the patient is then administered a therapeutically effective amount of the activatable anti-CD71 antibody and/or conjugated activatable anti-CD71 antibody that was tested. Likewise, patients that test negative for either or both of the target (e.g., CD71) and the protease that cleaves the substrate in the CM in the activatable antibody being tested using these methods might be identified as suitable candidates for another form of therapy. In some embodiments, such patients can be tested with other anti-CD71 activatable antibodies until a

suitable anti-CD71 activatable antibody for treatment is identified (e.g., an anti-CD71 activatable antibody comprising a CM that is cleaved by the patient at the site of disease). In some embodiments, the patient is then administered a therapeutically effective amount of the activatable anti-CD71 antibody and/or conjugated for which the patient tested positive. Suitable AB, MM, and/or CM include any of the AB, MM, and/or CM disclosed herein.

[000263] Pharmaceutical compositions according to the invention can include an antibody of the invention and a carrier. These pharmaceutical compositions can be included in kits, such as, for example, diagnostic kits.

[000264] In some embodiments, the pharmaceutical composition comprises an antibody of the disclosure, an activatable antibody of the disclosure, a conjugated antibody of the disclosure, and/or a conjugated activatable antibody of the disclosure, and a carrier. In some embodiments, the pharmaceutical composition comprises an additional agent. In some embodiments, the additional agent is a therapeutic agent.

Brief Description of the Drawings

[000265] Figure 1 is a graph depicting the ability of anti-CD71 antibodies of the disclosure to bind human CD71.

[000266] Figure 2 is a graph depicting the ability of various activatable anti-CD71 antibodies of the disclosure to bind human CD71.

[000267] Figure 3A is a graph depicting the ability of unconjugated and conjugated anti-CD71 activatable antibodies of the disclosure to bind human CD71 when the activatable antibody is intact or proteolytically activated. Figure 3B is a graph depicting the *in vitro* cytotoxicity of conjugated anti-CD71 activatable antibodies and activated conjugated anti-CD71 activatable antibodies of the disclosure.

[000268] Figure 4 is a graph depicting the efficacy of a conjugated activatable anti-CD71 antibody of the disclosure in NCI-H292 xenograft tumors.

[000269] Figure 5 is a graph depicting the efficacy of a conjugated activatable anti-CD71 antibody of the disclosure in breast cancer HCC1806 xenograft tumors.

[000270] Figure 6A is a graph depicting that an anti-CD71 antibody of the disclosure binds to cynomolgus monkey primary kidney epithelial cells, Figure 6B is a graph depicting that the anti-CD71 antibody of the disclosure binds to cynomolgus monkey CD71 but does not bind to mouse CD71, and Figure 6C is a graph depicting that the murine and humanized anti-CD71 antibodies of the disclosure bind to the human BxPC3 cell line.

[000271] Figure 7 is a graph depicting that a single dose of an activatable anti-CD71 antibody of the disclosure demonstrates a prolonged half-life compared to the CD71 parental antibody in cynomolgus monkeys.

[000272] Figure 8 is a graph depicting the pharmacokinetics of a conjugated activatable anti-CD71 antibody of the disclosure when administered to cynomolgus monkeys.

[000273] Figure 9 is a graph depicting the tolerability of a conjugated activatable anti-CD71 antibody of the disclosure when administered to cynomolgus monkeys.

[000274] Figure 10 is a graph depicting an exemplary assay of the binding between recombinant CD71 and its ligand transferrin.

[000275] Figures 11A is a graph depicting a competitive binding assay to CD71 between an anti-CD71 antibody of the present disclosure and transferrin. Figure 11B is a graph depicting a binding assay of the binding of transferrin to CD71 demonstrating the inhibition of the binding by an anti-CD71 antibody of the present disclosure.

[000276] Figures 12A, 12B, and 12C depict exemplary immunohistochemical (IHC) assays to determine levels of CD71 expression in various primary and metastatic cancer tissue types.

[000277] Figure 13A, 13B, and 13C depict exemplary efficacy studies of activatable anti-CD71 conjugated antibodies (AADCs) of the present disclosure using non-Hodgkin's lymphoma tumor xenografts in mice.

[000278] Figures 14A to 14G depict exemplary tolerability studies of anti-CD71 conjugated activatable antibodies (AADCs) of the present disclosure in cynomolgus monkeys.

[000279] Figure 15 depicts exemplary CD71 expression levels in various cell lines using an anti-CD71 antibody, as well as cytotoxicity studies in the same cell lines using an anti-CD71 antibody of the present disclosure with a drug-conjugated secondary antibody.

[000280] Figures 16A to 16D depict exemplary studies of the ability of various anti-CD71 activatable antibodies of the present disclosure to bind human CD71 on various human-derived cell lines.

[000281] Figures 17A to 17D depict exemplary studies of the cytotoxicity of anti-CD71 antibody drug conjugates of the present disclosure on various cell lines.

[000282] Figure 18 depicts exemplary studies of the cytotoxicity of various activatable anti-CD71 antibody drug conjugates of the present disclosure on a non-Hodgkin's lymphoma-derived Raji cell line.

[000283] Figure 19 depicts an exemplary study of the ability of various anti-CD71 activatable antibodies of the disclosure to bind human CD71.

[000284] Figures 20A and 20B depicts an exemplary study of the efficacy of anti-CD71 antibody drug conjugates (ADC) and anti-CD71 activatable antibody drug conjugates (AADC) of the present disclosure in a patient-derived xenograft tumor model.

[000285] Figures 21A – 21H depict exemplary studies of the cytotoxicity of various anti-CD71 antibody drug conjugates of the present disclosure on a variety of colorectal cancer-derived cell lines.

[000286] Figures 22A and 22B depicts exemplary studies of the cytotoxicity of various anti-CD71 antibody drug conjugates and activatable drug conjugates of the present disclosure on a HT29 colorectal cancer-derived cell line.

[000287] Figures 23A and 23B depict exemplary efficacy studies of activatable anti-CD71 conjugated antibodies (AADCs) of the present disclosure using non-Hodgkin's lymphoma tumor xenografts in mice.

[000288] Figures 24A and 24B depict exemplary efficacy studies of activatable anti-CD71 conjugated antibodies (AADCs) of the present disclosure using non-small cell lung cancer (NCI-H292) tumor xenografts in mice.

[000289] Figure 25 depicts exemplary efficacy studies of activatable anti-CD71 conjugated antibodies (AADCs) of the present disclosure using pancreatic cancer (BxPC3) tumor xenografts in mice.

[000290] Figures 26A and 26B depict exemplary efficacy studies of activatable anti-CD71 conjugated antibodies (AADCs) of the present disclosure using patient-derived tumor xenografts (non-Hodgkin's lymphoma) in mice.

[000291] Figures 27A and 27B depict exemplary efficacy studies of activatable anti-CD71 conjugated antibodies (AADCs) of the present disclosure using patient-derived tumor xenografts in mice.

[000292] Figure 28 depicts an exemplary study of the ability of various anti-CD71 antibodies and activatable antibodies of the disclosure to bind human CD71.

[000293] Figures 29A to 29C depict exemplary studies of the ability of various anti-CD71 activatable antibodies and activatable antibody drug conjugates of the present disclosure to bind human CD71 on various human-derived cell lines.

[000294] Figures 30A to 30D depict exemplary studies of the ability of various anti-CD71 activatable antibodies of the present disclosure to bind human CD71 *in vitro* and on various human-derived cell lines.

[000295] Figures 31A to 31J depicts exemplary studies of the cytotoxicity of various anti-CD71 activatable drug conjugates of the present disclosure on various cancer-derived cell lines.

[000296] Figures 32A and 32B depict exemplary studies of the ability of anti-CD71 antibodies and antibody drug conjugates of the present disclosure to bind human CD71 on various human-derived small cell lung cancer cell lines.

[000297] Figures 33A and 33B depict exemplary studies of the anti-proliferative effect of anti-CD71 antibodies of the present disclosure against various cancer-derived cell lines.

[000298] Figure 34 depicts exemplary studies of the expression level of CD71 in multiple patient-derived metastatic cancer samples.

[000299] Figure 35 depicts exemplary studies of *in vivo* binding of anti-CD71 antibodies of the present disclosure to tumors in a pancreatic cancer mouse model.

[000300] Figure 36 depicts exemplary studies of *in vivo* binding of anti-CD71 antibodies of the present disclosure to tumors in a metastatic breast cancer mouse model.

[000301] Figure 37 depicts exemplary studies of *in situ* binding of anti-CD71 antibodies of the present disclosure in a lung cancer xenograft model.

Detailed Description of the Invention

[000302] The present disclosure provides monoclonal antibodies (mAbs) and activatable monoclonal antibodies that specifically bind CD71, also known as transferrin receptor protein 1 (TfR1). In some embodiments, the monoclonal antibodies and activatable monoclonal antibodies are internalized by CD71-containing cells. The use of the term “CD71” is intended to cover any variation thereof, such as, by way of non-limiting example, CD-71 and/or CD 71, and all variations are used herein interchangeably.

[000303] CD71 is a transmembrane glycoprotein that primarily binds transferrin. CD71 is essential for cell homeostasis. CD71 is continuously recycled through ligand-

mediated endocytosis, where the main ligand is transferrin. CD71 is also known to be ubiquitously expressed on dividing cells.

[000304] Aberrant expression and/or activity of CD71 and CD71-related signaling have been implicated in the pathogenesis of many diseases and disorders, such as cancer. CD71 is overexpressed in many cancers, including both solid and hematological cancers. CD71 has broad cell surface expression. CD71 in malignant cells mediates higher iron uptake required for cell division. CD71 is also associated with poor prognosis in leukemias.

[000305] CD71 is desirable target because it is prevalent across multiple cancer indications.

[000306] The disclosure provides anti-CD71 antibodies, conjugated anti-CD71 antibodies, activatable anti-CD71 antibodies, and/or conjugated activatable anti-CD71 antibodies that are useful in methods of treating, preventing, delaying the progression of, ameliorating and/or alleviating a symptom of a disease or disorder associated with aberrant CD71 expression and/or activity. For example, the activatable anti-CD71 antibodies are used in methods of treating, preventing, delaying the progression of, ameliorating and/or alleviating a symptom of a cancer or other neoplastic condition.

[000307] The disclosure provides anti-CD71 antibodies, conjugated anti-CD71 antibodies, activatable anti-CD71 antibodies, and/or conjugated activatable anti-CD71 antibodies that are useful in methods of treating, preventing, delaying the progression of, ameliorating and/or alleviating a symptom of a disease or disorder associated with cells expressing CD71. In some embodiments, the cells are associated with aberrant CD71 expression and/or activity. In some embodiments, the cells are associated with normal CD71 expression and/or activity. For example, the activatable anti-CD71 antibodies are used in methods of treating, preventing, delaying the progression of, ameliorating and/or alleviating a symptom of a cancer or other neoplastic condition.

[000308] The disclosure provides anti-CD71 antibodies, conjugated anti-CD71 antibodies, activatable anti-CD71 antibodies, and/or conjugated activatable anti-CD71 antibodies that are useful in methods of treating, preventing, delaying the progression of, ameliorating and/or alleviating a symptom of a disease or disorder in which diseased cells express CD71. In some embodiments, the diseased cells are associated with aberrant CD71 expression and/or activity. In some embodiments, the diseased cells are associated with normal CD71 expression and/or activity. For example, the activatable anti-CD71 antibodies

are used in methods of treating, preventing, delaying the progression of, ameliorating and/or alleviating a symptom of a cancer or other neoplastic condition.

[000309] The activatable anti-CD71 antibodies and/or conjugated activatable anti-CD71 antibodies include an antibody or antigen-binding fragment thereof that specifically binds CD71 coupled to a masking moiety (MM), such that coupling of the MM reduces the ability of the antibody or antigen-binding fragment thereof to bind CD71. In some embodiments, the MM is coupled via a sequence that includes a substrate for a protease, for example, a protease that is co-localized with CD71 at a treatment site in a subject.

[000310] Exemplary activatable anti-CD71 antibodies of the invention include, for example, activatable antibodies that include a heavy chain and a light chain that are, or are derived from, the heavy chain variable and light chain variable sequences shown below (CDR sequences are shown in bold and underline):

muM21 VH:

EVQLQESGTVLARPGASVKMSCKAS**GYTF****TSYWMH**WVKQRPQGLEWIG**AIYPGNSETG**YNQNFKGK
AKLTAVTSASTAYMDLSSLTNEDSAVYYCT**ENWD****PGFAF**WGQGLITVSA (SEQ ID NO: 1)

muM21 VL:

DIVMTQTPAIMSASPGEKVTITC**SASSSVYYMY**WFQKPGTSPKLWIY**STSNLAS**GVPVRFSGSGSG
TSYSLTISRMEAEDAATYYC**QQRNYPYT**FGGGTKLEIKRA (SEQ ID NO: 2)

hu2vHa variable heavy chain

QVQLVQSGAEVKKPGASVKVSCCKAS**GYTF****TSYWMH**WVRQAPQGQLEWMG**AIYPGNSETG**YAQKFQGR
VTMTRDTSTSTVYMESSLRSEDVAVYYCAR**ENWD****PGFAF**WGQGLTVTVSS (SEQ ID NO: 3)

hu2vHb variable heavy chain

QVQLVQSGAEVKKPGASVKMSCKAS**GYTF****TSYWMH**WVRQAPQGQLEWIG**AIYPGNSETG**YAQKFQGR
ATLTADTSTSTAYMESSLRSEDVAVYYCT**ENWD****PGFAF**WGQGLTVTVSS (SEQ ID NO: 4)

hu2vHc variable heavy chain

QVQLVQSGAEVKKPGASVKMSCKAS**GYTF****TSYWMH**WVRQAPQGQLEWIG**AIYPGNSETG**YAQKFQGR
ATLTADTSTSTAYMESSLRSEDVAVYYCT**ENWD****PGFAF**WGQGLITVSS (SEQ ID NO: 5)

hu2lvKa variable light chain

DIQMTQSPSSLSASVGRVTITC**SASSSVYYMY**WYQKPGKAPKLLIY**STSNLAS**GVPSRFSGSGSG
TDFTLTISSLQPEDFATYYC**QQRNYPYT**FGGQTKLEIK (SEQ ID NO: 6)

hu21vKb variable light chain

DIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSG
TDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIK (SEQ ID NO: 7)

hu21vKc variable light chain

DIQMTQSPSSLSASVGDRVTITCCRASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSG
TDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIK (SEQ ID NO: 8)

[000311] Exemplary activatable anti-CD71 antibodies of the invention include, for example, activatable antibodies that include a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3) sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2) sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence comprising the amino acid sequence GYTFTSYWMH (SEQ ID NO: 9); a VH CDR2 sequence comprising the amino acid sequence AIYPGNSETG (SEQ ID NO: 10); a VH CDR3 sequence comprising the amino acid sequence ENWDPGFAF (SEQ ID NO: 11); a VL CDR1 sequence comprising the amino acid sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); a VL CDR2 sequence comprising the amino acid sequence STSNLAS (SEQ ID NO: 14); and a VL CDR3 sequence comprising the amino acid sequence QQRNYPYT (SEQ ID NO: 15).

[000312] In some embodiments, the activatable anti-CD71 antibody includes a heavy chain that comprises or is derived from a heavy chain amino acid sequence shown in PCT Publication Nos. WO 2014/144060, WO 2014/189973, WO 2014/020140, in US Patent Nos. 8,663,598; 8,129,503; 7,736,647; 7,572,895; 4,434,156; in US Patent Application Publication Nos. US2014114054, US20140212423, US2013177579, US2013045206, US20130216476, US20120282176, and/or in Chinese Patent No. CN101245107B, the contents of each of which are hereby incorporated by reference in their entirety.

[000313] In some embodiments, the activatable anti-CD71 antibody includes a heavy chain that comprises or is derived from a heavy chain amino acid sequence shown in Table 12. In some embodiments, the activatable anti-CD71 antibody includes a light chain that comprises or is derived from a heavy chain amino acid sequence shown in Table 12. In some embodiments, the activatable anti-CD71 antibody includes a heavy chain that comprises or is derived from a heavy chain amino acid sequence shown in Table 12, and a light chain that comprises or is derived from a light chain amino acid sequence shown in Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of heavy chain variable region and light chain variable region sequences from the combinations shown in Group A in Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of heavy chain variable region and light chain variable region sequences from the sequences shown in Group B in Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of heavy chain variable region and light chain variable region sequences from the sequences shown in Group C in Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of heavy chain variable region and light chain variable region sequences from the sequences shown in Group D in Table 12. In some embodiments, the activatable anti-CD71 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group E in Table 12. In some embodiments, the activatable anti-CD71 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group F in Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of heavy chain variable region and light chain variable region sequences from the sequences shown in Group G in Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of heavy chain variable region and light chain variable region sequences from the sequences shown in Group H in Table 12. In some embodiments, the activatable anti-CD71 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group I in Table 12. In some embodiments, the activatable anti-CD71 antibody includes the heavy chain variable region sequence shown in Group J in Table 12. In some embodiments, the activatable anti-CD71 antibody includes the heavy chain variable region sequence shown in Group J in Table 12, or the combination of heavy chain variable region and light chain variable region sequences shown in Group K in Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of heavy chain

variable region and light chain variable region sequences from the sequences shown in Group L in Table 12.

[000314] In some embodiments, the activatable anti-CD71 antibody includes a combination of the complementarity determining region (CDR) sequences of a heavy chain sequence from the heavy chain sequences shown in Group A Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a light chain sequence from the light chain sequences shown in Group A Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group A Table 12 and the CDRs of a light chain sequence from the heavy chain sequences shown in Group A Table 12.

[000315] In some embodiments, the activatable anti-CD71 antibody includes a combination of CDRs of a heavy chain sequence from the heavy chain sequences shown in Group B Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a light chain sequence from the light chain sequences shown in Group B Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group B Table 12 and the CDRs of a light chain sequence from the heavy chain sequences shown in Group B Table 12.

[000316] In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group C Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a light chain sequence from the light chain sequences shown in Group C Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group C Table 12 and the CDRs of a light chain sequence from the heavy chain sequences shown in Group C Table 12.

[000317] In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group D Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a light chain sequence from the light chain sequences shown in Group D Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown

in Group D Table 12 and the CDRs of a light chain sequence from the heavy chain sequences shown in Group D Table 12.

[000318] In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group E Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a light chain sequence from the light chain sequences shown in Group E Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group E Table 12 and the CDRs of a light chain sequence from the heavy chain sequences shown in Group E Table 12.

[000319] In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group F Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a light chain sequence from the light chain sequences shown in Group F Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group F Table 12 and the CDRs of a light chain sequence from the heavy chain sequences shown in Group F Table 12.

[000320] In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group G Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a light chain sequence from the light chain sequences shown in Group G Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group G Table 12 and the CDRs of a light chain sequence from the heavy chain sequences shown in Group G Table 12.

[000321] In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group H Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a light chain sequence from the light chain sequences shown in Group H Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown

in Group H Table 12 and the CDRs of a light chain sequence from the heavy chain sequences shown in Group H Table 12.

[000322] In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group I Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a light chain sequence from the light chain sequences shown in Group I Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group I Table 12 and the CDRs of a light chain sequence from the heavy chain sequences shown in Group I Table 12.

[000323] In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group J Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a light chain sequence from the light chain sequences shown in Group J Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group J Table 12 and the CDRs of a light chain sequence from the heavy chain sequences shown in Group J Table 12.

[000324] In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group K Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a light chain sequence from the light chain sequences shown in Group K Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group K Table 12 and the CDRs of a light chain sequence from the heavy chain sequences shown in Group K Table 12.

[000325] In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown in Group L Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a light chain sequence from the light chain sequences shown in Group L Table 12. In some embodiments, the activatable anti-CD71 antibody includes a combination of the CDRs of a heavy chain sequence from the heavy chain sequences shown

in Group L Table 12 and the CDRs of a light chain sequence from the heavy chain sequences shown in Group L Table 12.

Table 12. Variable Heavy Chain Region (VH) and Variable Light Chain Region (VL) Sequences for Activatable Antibodies that Bind CD71

Group A	
VH	VTLKVC ^{CGP} GILQPSQTLGLACTFSGISLSTSGMGLSWLRKPSGKALEWLASIWNNDNYNPSLKSRLTISKE TSNNQVFLKLTSVDTADSTTYFC ^{AWR} ERTMVTTSM ^{LWT} TGVKEPQSPS (SEQ ID NO: 425)
VL	DI ^{LMT} QSPASLSASVGENVTITCRASENIYSYLAWYQQKQKSPQLLLYKEKTLAEGVSSRFSGSGSGTQFS LRINSLQPEDFGSY ^{YCQH} HYGIPWTFGGGTKLEIKR (SEQ ID NO: 426)
Group B	
VH	EVQLVQSGAEVKKPGASVKVSC ^{KAS} GYTFTSYWMHWVRQAPGQRL ^{EWIG} EINPTNGRTNYIEKFKSRATLTV DKSASTAYMELSSLRSEDTAVYYCARGTRAYHYWGQGTMTVTVSS (SEQ ID NO: 427)
VH	EVQLVQSGAEVKKPGASVKVSC ^{KSG} YTF ^{TDY} AMHWVRQAPGQGLEWMGGISTYFGR ^{TNYN} QKFKGRVTMTV DTSISTAYMELSLRLSDDTAVYYCARGLSGNYVMDYWGQGTMTVTVSS (SEQ ID NO: 428)
VH	EVQLVQSGAEVKKPGASVKVSC ^{KAS} GYTFTSYWMHWVRQAPGQGLEWIGNIYPGSGSTKYDERFKSRVTITV DTSTSTAYLELSSLRSEDTAVYYCTRGGYDSRAWFAYWGQGTMTVTVSS (SEQ ID NO: 429)
VH	EVQLVESGPGLVKPS ^{ETLS} LTCTVSGNSITSEYAWN ^{WIRQ} PPGKLEWIGYISYSGTTSYNPSLKS ^{RV} TISR DTSKNQLSLKLSSVTAADTAVYYCARYGYGNPATRYFDVWGQGTMTVTVSS (SEQ ID NO: 430)
VH	EVQLVQSGAEVKKPGASVKVSC ^{KAS} GYTFTSYWMHWVRQAPGQRL ^{EWIG} EIAPTNGRTNYIEKFKSRATLTV DKSASTAYMELSSLRSEDTAVYYCARGTRAYHYWGQGTMTVTVSS (SEQ ID NO: 431)
VH	EVQLVQSGAEVKKPGASVKVSC ^{KAS} GYTFTSYWMHWVRQAPGQRL ^{EWIG} EINPANGRTNYIEKFKSRATLTV DKSASTAYMELSSLRSEDTAVYYCARGTRAYHYWGQGTMTVTVSS (SEQ ID NO: 432)
VL	DIQMTQSPSSLSASVGD ^{RV} TITCRASDNLYSNLAWYQQKPGKSPKLLVYDATNLADGVPSRFSGSGSGTDYT LTISLQPEDFATYYCQHFWGTPLTFGQGTKEIK (SEQ ID NO: 433)
VL	DIVMTQSPDSLAVSLGERATINCRASEVDSYGNSEFMHWYQQKPGQPPKLLIYRASNL ^{ESG} VPDRFSGSGSR TDFTLTISLQAEDVAVYYCQQSNEAPPTFGQGTKEIK (SEQ ID NO: 434)
VL	DIVMTQSPDSLAVSLGERATINCRARQSVSTSSYSFMHWYQQPAGQPPKLLIKYASIQ ^{ESG} VPDRFSGSGSG TDFTLTISLQAEDVAVYYCQHTWEIPPTFGQGTKEIK (SEQ ID NO: 435)
VL	DIQMTQSPSSLSASVGD ^{RV} TITCRASKSISKYLAWYQQKPGKTNKLLYSGSTLQSGVPSRFSGSGSGTDYT LTISLQPEDFATYYCQQHNEYPWTFGQGTKEIK (SEQ ID NO: 436)
VL	DIQMTQSPSSLSASVGD ^{RV} TITCRASDNLYSNLAWYQQKPGKSPKLLVYDATNLADGVPSRFSGSGSGTDYT LTISLQPEDFATYYCQHFA ^{GT} PLTFGQGTKEIK (SEQ ID NO: 437)
Group C	
VH	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNYGMHWIRQAPGKLEWIAMIIYDSSKMNYADTVKGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCAVPTSHYVVDVWGQGTMTVTVSS (SEQ ID NO: 438)

VH	QVQLVQSGAEVKKPGASVKVSCKASGFTFSNYGMHWIRQAPGGGLEWIAMIIYYDSSKMNYADTVKGRFTITR DNSTNTLYMELSSLRSEDTAVYYCAVPTSHYVVDVWGQGTITVTVSS (SEQ ID NO: 440)
VH	EVQLVESGGGLVQPNSLTLSCLVASGFTFSNYGMHWIRQAPKKGLEWIAMIIYYDSSKMNYADTVKGRFTISR DNSKNTLYLEMNSLRSEDTAMYYCAVPTSHYVVDVWGQGVSVTVSS (SEQ ID NO: 442)
VH	QVQLVQSGAEVKKPGSSVKVSCKASGYTFTNYDIHWVRQAPGGGLEWMGWIYPGDGSTKYNEKFKGRVTITA DESTSTAYMELSSLRSEDTAVYYCAREWAYWGQGTITVTVSS (SEQ ID NO: 443)
VH	EVQLQQSGAVLVKPGASVKLSCPASGFNIKDTYIHWVIQRPEQGLEWIGRIDPANGDTKCDPKFQVKATITA DTSSNTAYLQLSSLTSEDTAVYFCVRDYLPPYFDWFGQGTTLTVSS (SEQ ID NO: 988)
VH	QSMEESSGGRLVTPGTPLTLCTVSGFSLSSYAMSWVRQAPGKGLEWIGYIWSGGSTDYASWAKGRFTISKTS TTVDLKITSPTTEDTATYFCARRYGTSYPDYGDANGFDPWGPGLTVTVSS (SEQ ID NO: 989)
VL	DIQMTQSPASLSASLEEIIVTITCQASQDIGNWLAWYQQKPGKSPQLLIYGATSLADGVPSRFSGSRSGTQFS LKISRVQVEDIGIYYCLQAYNTPWTFGGGTKLELK (SEQ ID NO: 985)
VL	DIQMTQSPASLSASLEEIIVTITCQASQDIGNWLAWYQQKPGKSPQLLIYGATSLADGVPSRFSGSRSGTQFS LKISRVQVEDIGIYYCLQAYNTPWTFGGGTKVEIK (SEQ ID NO: 986)
VL	KIVMTQSPKSMSSVGERVTLNCRASESVDTYVSWYQQKPEQSPPELLIYGASNRYTGVPDRFTGSGSATDFT LTISSVQAEDLADYYCGQTNYNPLTFGAGTKLELKR (SEQ ID NO: 987)
VL	AYDMTQTPASVEVAVGGTVTIKCQASQSISSYLSWYQQKPGQRPKLLIYRASTLASGVSSRFKSGSGTQFT LTISGVECAADAATYYCQQCYSSSNVDNTFGGGTEVVVKR (SEQ ID NO: 990)
Group D	
VH	MEWSWIFLFLLSGTAGVLSEVELQQFGIEMVKPGASVKISCKASGYIFTDYHMDWVRQSHGKSLEWIGDIDP KYDRVTYNQKFKGKASLTADKSSSTAYMELRSLTSEDTAVYYCAKTGAYGDYLAAYWGQGTITVTVSA (SEQ ID NO: 444)
VH	MGWSYIILFLVATATGVHSQVQLQQPGAELVKPGTSVKLSCKASGYNFTSYWINWVKLRPGQGLEWIGDIYP GSGSTNYNEKFKSKATLTVDTSSSTAYMQLSSSLASEDSALYYCARSAYRYDWFAYWGQGTITVTVSA (SEQ ID NO: 445)
VH	MLLGLKWVFFVVFYQGVHCEVQLVESGGGLVQPKGSLKLSAASGFTFNTYAMNWVRQAPGKLEWVARIRS KSNNYATYYADSVKDRFTISRDDSQSMYLYQMNNLKTEDTAMYYCVAYGSRNYWGQGTITVTVSS (SEQ ID NO: 446)
VH	MGWSWIFLFLLSGTAGVHSEVQLQQSGPEVVKPGASMKMSCKTSGYKFTGYMDWVKQSLGASFEWIGRVIP SNGDTRYNQKFEGKATLTVDRSSSTAYMELNSLTSEDSAVYYCARKPLSGNAADYWGQGTITVTVST (SEQ ID NO: 447)
HC	EVQLQQSGPEVVKPGASMKMSCKTSGYKFTGYMDWVKQSLGASFEWIGRVIPSNGDTRYNQKFEGKATLTV DRSSSTAYMELNSLTSEDSAVYYCARKPLSGNAADYWGQGTITVTVSTASTKGPSVFPLAPSSKSTSGGTAAL GCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKK VEPKSCDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHE ALHNHYTQKSLSLSPGK (SEQ ID NO: 448)

VL	MKLPVRLVLVLMFWIPASSSDLVMTQTPLSLPVSLGDQASISCRSSQSLVHSDGNTYFYWYLQKPGQSPKLLI YKVSNRFSGVPDRFSAGGSGTYFTLKI SRVEAEDLG VYFCSQTTHFPPTFGGGTKLEIKR (SEQ ID NO: 449)
VL	MSVPTQVLGLLLLLWLT DARC DIQMTQSPASLSVSVGETVTITCRASENIYSNLAWYQQKQKSPQLLVYAAT NLADGVPSRFSGSGSGTQYSLKINSIQSEDFGSYYCQHFWGTPYTFGGGKLEIKR (SEQ ID NO: 450)
VL	MESQTQVLSLLFWVSGTCGDIVMTQSPSSLTVTAGEKVTMSCKSSQSLNLSGNQKNYLTWYQQKPGQPPKL LIYWASTRESGVPDRFTGSGSGTDFTLTISSVQAEDLAVYYCQNDYSYPYTFGGGKLEIKR (SEQ ID NO: 451)
VL	MTMLSLAPLLSLLLLCVSDSRAETTVTQSPASLSVATGEKVTIRCITSTDDIDDMNHYQQKPGEPKLLISD GNTLRPGVPSRFSGSGYGTDFVFTIENTLSEDI TDYYCMQSDNMPFTFGSGTKLEIKR (SEQ ID NO: 452)
LC	ETTVTQSPASLSVATGEKVTIRCITSTDDIDDMNHYQQKPGEPKLLISDGNTLRPGVPSRFSGSGYGTDFV FTIENTLSEDI TDYYCMQSDNMPFTFGSGTKLEIKRTVAAPSVFI FPPSDEQLKSGTASVVCLLNNFYPREA KVQWKVDNALQSGNSQESVTEQDSKDSYSTLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNREGC (SEQ ID NO: 453)
Group E	
VH	MEWSVVMFLFLLSGTAGVRSEVQLQQSGPELVKPGASMKISCKASGYSFTGYTMNVVKQSHGENLEWIGRINP HNGGTDYNQKFKDKAPLTVDKSSNTAYMELLSLTSEDSAVYYCARGYYYYSLDYWGQGTSTVTVSS (SEQ ID NO: 454)
VL	MDFQVQIFSFLLISASVILSRGQIVLTQSPAIMASAPGEKVTMTCSASSSIDYIHWYQQKSGTSPKRWIYDT SKLASGVPARFSGSGSGTSYSLTISSMEPEDAATYYCHQRNSYPWTFGGGTRLEIR (SEQ ID NO: 455)
Group F	
VH	MAQVQLLESGGGLVQPGGSLRLSCAASGFI FNTEYMAWVRQAPGKGLEWVSAIKEQSGSTYYADSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCAAQMHEAEVKFWGQGT LVTVSS (SEQ ID NO: 456)
VH	MAQVQLLESGGGLVQPGGSLRLSCAASGFKFSAEYMSWVRQAPGKGLEWVSTIKMNGSTYYADSVKGRFTI SRDNSKHTLYLQMNSLRAEDTAVYYCARPMAWRGNVVR AENLRFWGQGT LVTVSS (SEQ ID NO: 457)
Group G	
VH	MEFGLSWLFLVAILKGVQCEVQLQQSGTVLARP GASVKMSCKASGYSFTIYWIHWVKQRPQG LEWIATIYP GNSDIIYNQKFKGKAKLTAVTSASTAYMELSSLTNEASAVYYCTRQGYDYAMDYWGQGTSTVTVSS (SEQ ID NO: 458)
VL	MDMRVPAQLLGLLLLWLPGARCDVQITQSPSYLAASPGETII INCRASKSISKYLAWYQEKPGKTNKLLIYS GSTLQSGIPSRFSGSGSGTDFTLTISSLEPQDFAMYYCQHNEYPWTFGGGKLEIK (SEQ ID NO: 459)
VL	NIVMTQSPKSMSSVGERVTLTCKASENVVTVSWYQQKPEQSPKLLIYGASNRYTGVPDRFTGSGSATDFT LTISSVQAEDLADYHCGQGYSPYTFGGGKLEIKR (SEQ ID NO: 460)

VL	MDMRVPAQLLGLLLLWLPGARCDVQITQSPSYLAASPGETIIINCRASKSISKYLAWYQEKPGKTNKLLIYS GSTLQSGIPSRFSGSGSGTDFTLTISSELPQDFAMYYCQQHNEYPWTFGGGTKLEIKR (SEQ ID NO: 461)
VL	MDMRVPAQLLGLLLLWLPGARCDVQITQSPSYLAASPGETIIINCRASKSISKYLAWYQEKPGKTNKLLIYS GSTLQSGIPSRFSGSGSGTDFTLTISSELPQDFAMYYCQQHNEYPWTFGGGTKLQIK (SEQ ID NO: 462)
VL	NIVMTQSPKSMMSVGERVTLTCKASENVVTVSWYQKPEQSPKLLIYGASNRYTGVPDRFTGSGSATDFT LTISSEVQAEDLADYHCGQGYSPYTFGGGTKLQIKR (SEQ ID NO: 463)
VL	MDMRVPAQLLGLLLLWLPGARCDVQITQSPSYLAASPGETIIINCRASKSISKYLAWYQEKPGKTNKLLIYS GSTLQSGIPSRFSGSGSGTDFTLTISSELPQDFAMYYCQQHNEYPWTFGGGTKLQIKR (SEQ ID NO: 464)
Group H	
VH	QVQLQESGGGLVQPGGSLRLSCAASGFTSFNTYTMHWVRQAPGKGLEWVADIAYDGSTKYYADSVKGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCARD AVAGEGYFDLWGRGTLTVS (SEQ ID NO: 465)
VH	QVQLQESGGGLVQPGGSLRLSCAASEFTFSASGMHWVRQAPGKGLEWMAFIAYDGNQKFYADSVKGRFTISR DNSKNTLYLQMDSLRLGEDTAVYYCAKEMQREGYFDYWGQGTTLTVS (SEQ ID NO: 466)
VH	QVQLAESGGGLVQPGGSLRLSCAASGFTFS DYYMSWIRQAPGKGLEWVS YISTSGSSIYYVDSVKGRFTISR DNAKNSLYLQMDSLRLDDDTAVYYCARDLHGDYAFDSWGQGTTLTVS (SEQ ID NO: 467)
VH	QVQLQESGGGLVQPGGSLRLSCAASGFTFSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTISR DNSKNTLYLQMNSLRAEDTAVYYCAK VSSWSHF YWGQGTTLTVS (SEQ ID NO: 468)
VH	QVQLVESGGGLVPEPGGSLRLSCAASGFTFSNYAINWVRQAPGKGLEWVANIHHDGNGKYYVDSVEGRFTISR DNAKNSLYLQMDSLRAEDTAIYYCARDGYGGYLDLWGQGTTLTVS (SEQ ID NO: 469)
VH	QVQLQESGGGLVQPGGSLRLSCAASRTFSSYAMHWVRQAPGKGLEWVAVISYDGSNKYYADSVKGRFTISR DNSKNTLYLQMNSLRAEDTAVYYCARDLSGYGDYDPDYWGQGTTLTVS (SEQ ID NO: 470)
VL	SNFMLTQDPAVSVALGQTVRITCQGDSLRSYASWYQKPGQAPVLVIYGKNNRPSGIPDRFSGSKSGNSAS LDISGLQSEDEADYYCATWDDNLSGPIFGGGTKVTVLG (SEQ ID NO: 471)
VL	SQSALTQDPAVSVALGQTVRITCQGDSLRSYASWYQQLPGTAPKLLIYRNNQRPSGVPDRFSGSKSGTSAS LAISGLRSEDEADYYCAAWDDSLSAWVFGGGTKLTVLGA (SEQ ID NO: 472)
VL	SSELTQDPAVSVALGQTVRITCQGDSLRSYASWYQKPGQAPVLVIYGKNNRPSGIPDRFSGSKSGNSASL DISGLQSEDEADYYCATWDDNLSGPIFGGGTKVTVLG (SEQ ID NO: 473)
VL	SDVMTQSPSTLSASVGRVTITCRASQYISNWLAWYQKPGKAPKLLIYKASSLES GVPSRFSGSGSGTEF TLTISSELPEDFATYYCQESYNTPLFTFGPGTKLEIKR (SEQ ID NO: 474)
VL	SSELTQDPAVSVALGQTVRITCQGDSLRSYASWYQKPGQAPVLVIYGKNNRPSGIPDRFSGSGSGNTASL TITGAQAEDEADYYCAAWDDSLSGPVFGGGTKVTVLG (SEQ ID NO: 475)
VL	SSELTQDPAVSVALGQTVRITCQGDSLRSYASWYQKPGQAPVLVYGRNERPSGVPDRFSGSKSGTSASL AISGLQPEDEANYYCAGWDDSLTGPVFGGGTKLTVLG (SEQ ID NO: 476)

VH / VL	QVQLQESGGGLVQPGGSLRLSCAASGFSFNTYTMHWVRQAPGKGLEWVADIAVDGSKYYADSVKGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCARD <u>AVAGEGYFDLWGRGTLVTVSSGGGGSGGGSGGGGSQ</u> SALTQDPA VSVALGQTVRITCQGD <u>SLRSYYASWYQQ</u> LPGTAPKLLIYRNNQRP <u>SGVPDR</u> FSGSKSGTSASLAISGLRSED EADYYCAAWDDSLSAWVFGGGTKLTVLG (SEQ ID NO: 477)
VH / VL	QVQLQQSGGGVVPQGGSLRLSCAASEFTFSASGMHWVRQAPGKGLEWMAFIAYDGNQKFYADSVKGRFTISR DNSKNTLYLQMDSLRLGEDTAVYYCAKEMQREGYFDYWGQGT <u>LVTVSSGGGGSGGGSGGGGSNF</u> MLTQDPAV SVALGQTVRITCQGD <u>SLRSYYASWYQQ</u> KPGQAPVLVIY <u>GKNNRPSGI</u> PDRFSGSKSGNSASLDISGLQSEDE ADYYCATWDDNLSGPIFGGGTKVTVLG (SEQ ID NO: 478)
VH / VL	QVQLAESGGGLVKPQGGSLRLSCAASGFTFSDYIMSWIRQAPGKGLEWVSISTSGSSIIYYVDSVKGRFTISR DNAKNSLYLQMDSLRLDDDTAVYYCARDLHGDYAFDSWGQGT <u>LVTVSSGGGGSGGGSGGGGSSELT</u> QDPAVS VALGQTVRITCQGD <u>SLRSYYASWYQQ</u> KPGQAPVLVIY <u>GKNNRPSGI</u> PDRFSGSKSGNSASLDISGLQSEDEA DYCATWDDNLSGPIFGGGTKVTVLG (SEQ ID NO: 479)
VH / VL	QVQLQESGGGLVQPGGSLRLSCAASGFTFSYAMS WVRQ APGKGLEWVSAISGSGGSTYYADSVKGRFTISR DNSKNTLYLQMNSLRAEDTAVYYCAK <u>VSSWSHFDYWGQGT</u> LVTVSSGGGGSGGGSGGGGSDVMTQSPST LSASVGDRVTITCRASQYISNWLAWYQQKPGKAPKLLIYKASSLES <u>GVPSRF</u> SGSGSGTEFTLTISLQPED FATYYCQESYNTPLFTFGPGTKLEIKR (SEQ ID NO: 480)
VH / VL	QVQLVESGGGLVPEPQGGSLRLSCAASGFTFSNYAINWVRQAPGKGLEWVANIHHDGNGKYYVDSVEGRFTISR DNAKNSLYLQMDSLRAEDTAIYYCARDGYGGYLDLWGQGT <u>LVTVSSGGGGSGGGSGGGGSSELT</u> QDPAVSV ALGQTVRITCQGD <u>SLRSYYASWYQQ</u> KPGQAPVLVIY <u>GKNNRPSGI</u> PDRFSGSGSGNTASLTITGAQAEDEAD YYCAAWDDSLSGPVFGGGTKVTVLG (SEQ ID NO: 481)
VH / VL	QVQLQESGGGVVPGRSLRLSCAASRFTFSYAMHWVRQAPGKGLEWVAVISYDGSNKYYADSVKGRFTISR DNSKNTLYLQMNSLRAEDTAVYYCARDLSGYGDYDPDYWGQGT <u>LVTVSSGGGGSGGGSGGGGSSELT</u> QDPAV SVALGQTVRITCQGD <u>SLRSYYASWYQQ</u> KPGQAPVLVYGRNERPSGV <u>PDRFS</u> GSKSGTSASLAISGLQPEDE ANYYCAGWDDSLTGPVFGGGTKLTVLG (SEQ ID NO: 482)
Group I	
VH	MEFGLSWLFLVAILKGVQCEVQLQQSGTVLARPGASVKMSCKASGYSTIYWIHWVKQRPQGQGLEWIAIYIP GNSDIIYNQKFKGAKLTAVTSASTAYMELSSLTNEASAVYYCTRQGYDYAMDYWGQGT <u>SVTVSS</u> (SEQ ID NO: 483)
VL	MDMRVPAQLLGLLLLWLPGARCDVQITQSPSYLAASPGETIIINCRASKSISKYLAWYQEKPGKTNKLLIYS GSTLQSGIPSRFSGSGSGTDFTLTISSELPQDFAMYYCQQHNEYPWTFGGGTKLEIKR (SEQ ID NO: 484)
Group J	
VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFAHETMVWVRQAPGKGLEWVSHIPVGGQDPFYADSVKGRFTISR DNSKNTLYLQMNSLRAEDTAVYYCALLPKRGPWFYWGQGT <u>LVTVSS</u> (SEQ ID NO: 485)
Group K	
VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFAHETMVWVRQAPGKGLEWVSHIPVGGQDPFYADSVKGRFTISR DNSKNTLYLQMNSLRAEDTAVYYCALLPKRGPWFYWGQGT <u>LVTVSS</u> (SEQ ID NO: 485)

VH / VL	MASYELTQPPSVSVAPGQTARITCSGDALGNKYASWYQQKPGQAPVLVIYEDSKRPSGIPERFSGSNSGNTA TLTISGTQAEDEADYYCSSGDSPCRAFGGGTKLTVLGSGGSTITSYNVYYTKLSSSGSEVQLVESGGGLVQP GGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTISRDNKNTLYLQMNSL RAEDTAVYYCARHSIYRCFFAVWGQGTLLTVSS (SEQ ID NO: 487)
Group L	
VH	DVQLVESGGGLVQPGGSLRLSCAASGFPFKSYGMQWVRQAPGKGLEWVAVISFDGSSRYADSVKGRFTISR DNKNTLYLQMNSLRAEDTAVYYCARDGALWGGYYSPVDVWGQGTLLTVS (SEQ ID NO: 999)
VH	DVQLVESGGGLVQPGGSLRLSCAASGFPFKSYAMHWVRQAPGKGLEWVAVISYDGSNKYYADSVKGRFTISR DNKNTLYLQMNSLRAEDTAVYYCARDGALWGGYYSPVDVWGQGTLLTVSS (SEQ ID NO: 1000)
VH	DVQLVESGGGLVQPGGSLRLSCAASGFPFKSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVKGRFTISR DNKNTLYLQMNSLRAEDTAVYYCARDGALWGGYYSPVDVWGQGTLLTVSS (SEQ ID NO: 1001)
VH	DVQLVESGGGLVQPGGSLRLSCAASGFPFKSYAMHWVRQAPGKGLEWVAVISFDGSSRYADSVKGRFTISR DNKNTLYLQMNSLRAEDTAVYYCARDGALWGGYYSPVDVWGQGTLLTVSS (SEQ ID NO: 1002)
VH	DVQLVESGGGLVQPGGSLRLSCAASGFPFKSYGMQWVRQAPGKGLEWVAVISFDGSSRYADSVKGRFTISR DNKNTLYLQMNSLRAEDTAVYYCARDGALWGGYYSPVDVWGQGTLLTVSS (SEQ ID NO: 1003)
VH	DVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMQWVRQAPGKGLEWVAVISFDGSSRYADSVKGRFTISR DNKNTLYLQMNSLRAEDTAVYYCARDGALWGGYYSPVDVWGQGTLLTVSS (SEQ ID NO: 1004)
VH	DVQLVESGGGLVQPGGSLRLSCAASGFTFSSYGMQWVRQAPGKGLEWVAVISFDGSSRYADSVKGRFTISR DNKNTLYLQMNSLRAEDTAVYYCARDGALWGGYYSPVDVWGQGTLLTVSS (SEQ ID NO: 1005)
VH	DVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMQWVRQAPGKGLEWVVISFDGGNRYADSVKGRFTISR DNKNTLYLQMNSLRAEDTAVYYCARDGALWGGYYSPVDVWGQGTLLTVSS (SEQ ID NO: 1006)
VH	DVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMQWVRQAPGKGLEWVISVFDGGNRYADSIGRFTISR DNKNTLYLQMNSLRAEDTAVYYCARDGALWGGYYSPIDVWGQGTLLTVSS (SEQ ID NO: 1007)
VL	DFMLTQPHSVSESPGKTVTISCTRSSGSIASNYVQWYQQRPGSSPTTVIYEDNQRPSPGVDRFSGSIDSSN SASLTISGLKTEDEADYYCQSYDSSNHWFVGGGTKLAVL (SEQ ID NO: 1008)
VL	DFMLTQPQSVSESPGKTVTISCTRSSGSIASNYVQWYQQRPGSSPTTVIYEDNQRPSPGVDRFSGSIDSSN SASLTISGLKTEDEADYYCQSYDSSNQWVFGGGLKLAFL (SEQ ID NO: 1009)
VL	QSXLTPPPSVSGSPGQSVTISCTGSSSNIASXNYVSWYQQXPGTAPKLMYENNRKPSGPVDRFSGSKXXSG NTASLTISGLQAEDEADYYCSDSSLSX (SEQ ID NO: 1010)
VL	DSALTQPPSVSGSPGQSVTISCTGSSSNIASNSVQWYQQLPGTAPKTVIYEDTQRPSPGVDRFSGSKDSSG NTASLTISGLQAEDEADYYCQSYDSAYHWVFGGGLKLAFL (SEQ ID NO: 1011)
VL	DSALTQPPSVSGSPGQSVTISCTGSSSNIASNSVQWYQQLPGTAPKTVIYENTQRPSPGVDRFSGSKDSSG NTASLTISGLQAEDEADYYCSDSAYHWVFGGGLKLAFL (SEQ ID NO: 1012)
VL	DFMLTQPHSVSESPGKTVIIISCTRS DGTIAGYYVQWYQQRPGRAPTTVIFEDTQRPSPGVDRFSGSIDRSSN SASLTISGLQTEDEADYYCQSYDRDHWVFGGGLKLTVLG (SEQ ID NO: 1013)
VL	DFMLTQPHSVSESPGKTVIIISCTRS DGTIAGYYVQWYQQRPGRAPTTVIFEDTQRPSPGVDRFSGSIDRSSN SASLTISGLQTEDEADYYCQSYDSRDHWVFGGGLKLTVL (SEQ ID NO: 1014)
VL	DFMLTQPQSVSESPGKTVIIISCTRS DGTIAGYYVQWYQQRPGRAPTTVIFDETQRPSPGVDRFSGSIDRSSN SASLTISGLQTEDEADYYCQSYDSRDQWVFGGGLKLTVL (SEQ ID NO: 1015)

VL	DIQMTQSPSSLSASVGDRVTITCRASQISNYLAWYQQKPGKAPKLLIYAASSLESQVPSRFSGSGSGTDFT LTISSLQPEDFATYYCQQYNLSPWTFGQGTKVEIK (SEQ ID NO: 1016)
VL	DIQMTQSPSSLSASVGDRVTITCRASQIASNSVQWYQQKPGKAPKTVIYEDTQLESQVPSRFSGSGSGTDFT LTISSLQPEDFATYYCQSYDSAYHWVFGQGTKVEIK (SEQ ID NO: 1017)
VL	DIQMTQSPSTLSASVGDRVTITCRASQISSWLAWYQQKPGKAPKLLIYDASSLESQVPSRFSGSGSGTEFT LTISSLQPDDFATYYCQQYNSYS (SEQ ID NO: 1018)
VL	DIQMTQSPSTLSASVGDRVTITCRASQIASNSVQWYQQKPGKAPKTVIYEDTQLESQVPSRFSGSGSGTEFT LTISSLQPDDFATYYCQSYDSAYHWVFGQGTKVEIK (SEQ ID NO: 1019)
VL	DIQMTQSPSTLSASVGDRVTITCRASQIASNSVQWYQQKPGKAPKTVIYEDTQLESQVPSRFSGSGSGTEFT LTISSLQPDDFATYYCQSYNSAYHWVFGQGTKVEIK (SEQ ID NO: 1020)
VL	DIQMTQSPSTLSASVGDRVTITCRASQIASNSVQWYQQKPGKAPKTVIYEDTQLESQVPSRFSGSGSGTEFT LTISSLQPDDFATYYCQSYNSAYQWVFGQGTKVEIK (SEQ ID NO: 1021)
VL	DIQMTQSPSSLSASVGDRVTITCRASQIRNDLTWYQQKPGTAPKRLIYGATSLQSGVPSRFSGSGSGTEFT LTINSLQPEDFATYYCLQYSSFPWTFGQGTKVEIK (SEQ ID NO: 1022)
VL	DIQMTQSPSSLSASVGDRVTITCRASQIASNSVQWYQQKPGTAPKTVIYEDTQLQSGVPSRFSGSGSGTEFT LTINSLQPEDFATYYCQSYDSAYHWVFGQGTKVEIK (SEQ ID NO: 1023)
VL	DIQMTQSPSSLSASVGDRVTITCRASQIASNSVQWYQQKPGTAPKTVIYEDTQLQSGVPSRFSGSGSGTEFT LTINSLQPEDFATYYCQSYNSAYHWVFGQGTKVEIK (SEQ ID NO: 1024)
VL	DIQMTQSPSSLSASVGDRVTITCRASQIASNSVQWYQQKPGTAPKTVIYEDTQLQSGVPSRFSGSGSGTEFT LTINSLQPEDFATYYCQSYNSAYQWVFGQGTKVEIK (SEQ ID NO: 1025)

(VH/VL = VH linked to VL)

[000326] In some embodiments, the activatable anti-CD71 antibody includes a CDR sequence shown in Table 13, a combination of VL CDR sequences (VL CDR1, VL CDR2, VL CDR3) selected from the group consisting of those combinations shown in a single row Table 13, a combination of VH CDR sequences (VH CDR1, VH CDR2, VH CDR3) selected from the group consisting of those combinations shown in Table 13, or a combination of VL CDR and VH CDR sequences (VL CDR1, VL CDR2, VL CDR3, VH CDR1, VH CDR2, VH CDR3) selected the group consisting of those combinations shown in Table 13.

Table 13. CDR Sequences for Antibodies and Activatable Antibodies that Bind CD71

Group A					
VH			VL		
CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)	CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)
TSGMGL (488)	ASIWNNDNYNPS LKS (489)	AWRERTMVTTS MLWT (490)	RASENIYSYL A (491)	KEKTLAE (492)	QHGYGIPWT (493)

	GISTYFGRNTYNQ KFKG (494)	GLSGNYVMDY (495)
	VIS (F/P) YSGKT NYSQNFMG (501)	GLSGNFVVDF (502)
	EINPTNGRTNYIE KFKS (506)	YGYGNPATRYF DV (507)
	EINP (I/S) NGRT NY (N/S) E (N/T) FKK (511)	GLSGNYVMDY (495)
	NIYPGSGSTKYDE RFKS (516)	GLSGNYVVDF (517)
	(S/G) I (S/L) NG GDNTYY (P/N) D (K/T) VKG (521)	GLSGNFVMDY (522)
	VISPYSGRNTYNQ NFKG (525)	GTRAYHY (526)
	EIAPTNGRTNYIE KFKS (528)	GTRAYHF (529)
	VISFYSGKTNYNQ KFMG (531)	GTRAYHY (526)
	VISPYSGKTNYSQ KFKG (534)	GTRAYHF (529)
	EINPTNGRTNYIE KFK (537)	GGYDSRAWFAY (538)
	EINPTNGRTNYNE NFKS (540)	GGYDSRAWFAH (541)
	EINPINGRTNYSE KFKK (543)	QGALYDGYRG AMDY (544)
	YISYSGTTSYNPS LKS (546)	RGYGYDGEFA Y (547)
	NIYPGSGSTKYDE RFKS (516)	
	NIYPGSGSTKYDE KFKS (551)	
	SISNGGDNTYYPD TVK (553)	
	EILPGSGSTKYNE KFKG (555)	

Group B

VH		
CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)
GFTFSNYGMH (558)	MIYYDSSKM NYADTVKG (559)	PTSHYVVDV (560)
GFTFSNYGMH (561)	MIYYDSSKM NYADTVKG (559)	PTSHYVVDV (563)
GYTFTNYDIH (564)	MIYYDSSK MNYADTVK G (559)	PTSHYVVDV (566)
	WIYPGDGST KYNEKFKG (567)	YWQGGTTV (578)

Group C

RASESVDSYG NSFMH (496)	RASNLES (497)	QQSNEAPPT (500)
RASESVDDYP NSFMH (503)	DATNLAD (504)	QHSNE (G/D) PPT (505)
RASENLYSNL A (508)	DATDLAD (509)	QHFWSGTPLT (510)
RASDNLYSNL A (513)	YASIQES (514)	GGYDSRAWFRGAMA Y (515)
RARQSVSTSS YSFMH (518)	STS (N/R) L (A/H) S (519)	(Q/R) G (A/G) (L/Y) (Y/G) (D/Y) (G/D) (Y/G) (F or Absent) DH (520)
TT (S/G) S (V/D) (P/I) (S/T) NY (F/L) N (523)		QQSNEAPPT (524)
		QQSNEGPPPT (527)
		GISTYFGRNTYNQK FK (530)
		QHSNEDPPT (533)
		QHFWSGTPLT (510)
		QHFWSGTPLM (539)
		QQHNEYPTT (542)
		QHTWEIPPT (545)
		HQYHRSPTT (548)
		QQANTLPYT (550)
		QHTWEIPPT (552)
		HQ (Y/A) (H/N) (R/T) (S/L) PYT (554)
		QQSNEAPPT (556)
		QHFAGTPLT (557)

VH			VL		
CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)	CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)
TYTMH (579)	DIAYDGGSTKYY ADSVKG (580)	DAVAGEGYFDL (581)	QGDSLRSYYA S (582)	RNNQRPS (583)	AAWDDSLSAVV (584)
ASGMH (585)	FIAYDGNQKFY ADSVKG (586)	EMQREGYFDY (587)	QGDSLRSYYA S (582)	GKNNRPS (589)	ATWDDNLSGPI (590)
DYYMS (591)	YISTSGSSIYY VDSVKG (592)	DLHGDYAFDS (593)	QGDSLRSYYA S (582)	GKNNRPS (589)	ATWDDNLSGPI (590)
SYAMS (597)	AISGSGGSTYY ADSVKG (598)	VSSSWSHFDY (599)	RASQYISNWL A (600)	KASSLES (601)	QESYNTPLFT (602)
NYAIN (603)	NIHHDGNGKYY VDSVEG (604)	DGYGGYLDL (605)	QGDSLRSYYA S (582)	GKNNRPS (589)	AAWDDSLSGPV (608)
SYAMH (609)	VISYDGSNKYY ADSVKG (610)	DLSGYGDYPDY (611)	QGDSLRSYYA S (582)	GRNERPS (613)	AGWDDSLTGPV (614)

Group D

VH			VL		
CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)	CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)
Any of the VH CDR combinations shown in paragraphs [0060], [00150] and/or [00151] of US Patent Application Publication No. 2014/0114054			Any of the VL CDR combinations shown in paragraph [0060], [00150] and/or [00151] of US Patent Application Publication No. 2014/0114054		
TSGVGVG (439)	LIYWDDDKHYS PSLKS (441)	NGDYGIEFDY (486)	GGNNIGSKSV H (512)	YSDRPS (516)	QVWDSSSDHVV (532)
SYSMN (535)	SISSSSYIYY ADSVKG (536)	ARESVDAFDI (562)	QGDSLRSYDA S (565)	GLSDRPS (588)	ISRDSGGNPH (594)
SYAMS (597)	AISGSGGSTYY ADSVKG (598)	GYGGSNYGYMD V (595)	SGSSSNIGSN YVY (596)	RNNQRPS (583)	AAWDDSLSGPV (608)
DFVFS (606)	WISAHGNTNY AQKLQD (607)	DTFTNLLGDYSY DAMDV (612)	GSSTGAVTSG HYPY (909)	DTTEKHS (910)	LLSSGDGRAV (911)
NYGMS (912)	WISAYNGNTNY GEKLQG (913)	DDYYGSGVDAFD I (914)	GGNKIGSKSV H (915)	YDRDRPS (916)	QVWDSSSDVV (917)
SYGMH (918)	VISFDGSSKYY ADSVKG (919)	DSNFWSGYYSPV DV (920)	TRSSGSIASN SVQ (921)	YEDTQRPS (922)	QSYDSAYHWV (923)
SYWLS (924)	KIDPSDSYTQY SPSFEG (925)	HGYDAFHV (926)	SGSSSNIGNN AVN (927)	YDDLPS (928)	AAWDDSLNGWV (929)
DYAMH (930)	GISWNSGSIGY ADSVKG (931)	DQHREFYYYGMD V (932)	SGSSSNIGSN YVY (596)	RNNQRPS (583)	AAWDDSLSGPV (608)
SYWIG (933)	IIPGDSDTTRY SPSFQG (934)	QGTNWGVGDAFD I (935)	GGNNIGSKSV H (512)	DDSDRPS (936)	QVWDISSDHVV (937)
SYAMS (597)	AISGSGGSTYY ADSVKG (598)	DRYYYGSGSYD AFDI (938)	QGDSLRSYYA S (582)	GKNNRPS (589)	NSRDSSGNHVV (939)
SYSMN (535)	VISYDGSNKYY ADSVKG (610)	VDPGDRGWYFDL (940)	SGSSSNIGSN TVN (941)	SNNQRPS (942)	AAWDDSLNGWV (929)
SSPYWG (943)	SVYYSGNTYYN PSLTR (944)	HSWGINDAFDV (945)	SGSSSNIGNN YVS (946)	DNNKRPS (947)	GTWDDSSLVWV (948)

DYAMH (930)	GISWNSGSIDY ADSVKG (949)	ENLAVAGLDY (950)	QGDSLIRGYA S (951)	DKNTRPS (952)	QSRDNSGEMVV (953)
ELSMH (954)	GFDPEDGETIY AQKFQG (955)	DAYYGS SPRDA FDI (956)	GGDNVGGKSL H (957)	DDRDRPS (958)	QVWDDISRLVI (959)
SYIIH (960)	IINPRGGGTDF AQKFQG (961)	GDCTNGVCYSGG LDV (962)	SGSSSNIGNN YVS (946)	DNDKRPS (963)	GTWDNSLSGV (964)
DYAMH (930)	GISWNSGSIGY ADSVKG (931)	DVDLWFGEYYFD Y (965)	SGSSSNIGNN YVS (946)	DNNKRPS (947)	GTWDSSLSAPYV (966)
DYAMY (967)	GINWNSAIGY ADSVKG (968)	EALYYSAFFDS (969)	SGSSSNIGNN YVS (946)	DNNKRPS (947)	GTWDSSLSAWV (970)
DYAMH (930)	GINWNGGSTDY ADSVKG (971)	DYADLGSGSDY (972)	SGSRNIGSN YVH (973)	RNDQRPS (974)	ASWDDKMSGRL (975)
SYEMN (976)	YISSSGSTIYY ADSVKG (977)	HSNYDILTGYST DAFDI (978)	TGTSSDIGFY DSVS (979)	DVSNRPS (980)	TSNTKTNTLYV (981)
RGNYWWT (982)	SVHYSGSTNYN PSLKS (983)	DSYDGYDFDY (984)	QGDSLRSYYA S (582)	GKNNRPS (589)	NSRDSSGNHV (939)

Group E

VH			VL		
CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)	CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)
SYGMH (918)	VISFDGSSKYY ADSVK (919)	DGNFWSGYYSPV DV (1026)	TRSSGSIASN SVQ (921)	YEDTQRPS (922)	QSYDSAYHWV (923)
SYGMH (918)	VISFDGSSKYY ADSVK (919)	DSAFWSGYYSPV DV (991)			
SYGMH (918)	VISFDGSSKYY ADSVK (919)	DSNLWSGYYSPV DV (992)			
SYGMH (918)	VISFDGSSKYY ADSVK (919)	DSNFWGGYYSPV DV (993)			
SYGMH (918)	VISFDGSSKYY ADSVK (919)	DSALWGGYYSPV DV (994)			
SYGMH (918)	VISFDGSSKYY ADSVK (919)	DGNLWGGYYSPV DV (995)			
SYGMH (918)	VISFDGSSKYY ADSVK (919)	DGAFWGGYYSPV DV (996)			
SYGMH (918)	VISFDGSSKYY ADSVK (919)	DGALWSGYYSPV DV (997)			
SYGMH (918)	VISFDGSSKYY ADSVK (919)	DGALWGGYYSPV DV (998)			
SYGMH (918)	VISYDGSNKYY ADSVKG (1027)	DSNFWSGYYSPV DV (920)			
SYGMH (918)	VISYDGSNKYY ADSVKG (610)	DSNFWSGYYSPV DV (920)			

Group F

VH			VL		
CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)	CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)
FSLSSY (1028)	WS	GTSYPDYGDANG FD (1029)	SQSISSY (1030)	RAS (1031)	CYSSSNVDN (1032)
GFNIKDT (1033)	ANG	YLYPPYFD (1034)	SESVDTY (1035)	GAS	TYNYPL (1036)

[000327] In some embodiments, the activatable anti-CD71 antibody comprises or is derived from an antibody that is manufactured, secreted or otherwise produced by a

hybridoma, such as, for example, the hybridoma(s) designated BA120 as disclosed in US Patent No. 7,736,647 and deposited at the Collection Nationale de Cultures de Microorganismes (CNCM) (Institut Pasteur, Paris, France, 25, Rue du Docteur Roux, F-75724, Paris, Cedex 15) on Jun. 14, 2005, under number CNCM I-3449; the hybridoma(s) disclosed in US Patent No. 7,572,895 and deposited at the ATCC under PTA-6055; the hybridoma(s) disclosed in PCT Publication No. WO 2014/020140 and WO 2005/111082 and deposited with CNCM on May 10 2001, under number 1-2665; the hybridoma(s) disclosed in US Patent No. 4,434,156 and deposited at the ATCC under HB-8094; the hybridoma(s) disclosed in US Patent No. 5,648,469 and deposited at the ATCC under HB-11011 and HB-11010.

[000328] The anti-CD71 antibodies and the ABs in the activatable antibodies of the disclosure specifically bind a CD71 target, such as, for example, mammalian CD71, and/or human CD71. Also included in the disclosure are anti-CD71 antibodies and ABs that bind to the same CD71 epitope as an antibody of the disclosure and/or an activated activatable antibody described herein. Also included in the disclosure are anti-CD71 antibodies and ABs that compete with an anti-CD71 antibody and/or an activated anti-CD71 activatable antibody described herein for binding to a CD71 target, e.g., human CD71. Also included in the disclosure are anti-CD71 antibodies and ABs that cross-compete with an anti-CD71 antibody and/or an activated anti-CD71 activatable antibody described herein for binding to a CD71 target, e.g., human CD71.

[000329] The activatable anti-CD71 antibodies provided herein include a masking moiety. In some embodiments, the masking moiety is an amino acid sequence that is coupled or otherwise attached to the anti-CD71 antibody and is positioned within the activatable anti-CD71 antibody construct such that the masking moiety reduces the ability of the anti-CD71 antibody to specifically bind CD71. Suitable masking moieties are identified using any of a variety of known techniques. For example, peptide masking moieties are identified using the methods described in PCT Publication No. WO 2009/025846 by Daugherty et al., the contents of which are hereby incorporated by reference in their entirety.

[000330] The activatable anti-CD71 antibodies provided herein include a cleavable moiety. In some embodiments, the cleavable moiety includes an amino acid sequence that is a substrate for a protease, usually an extracellular protease. Suitable substrates are identified using any of a variety of known techniques. For example, peptide substrates are identified

using the methods described in U.S. Patent No. 7,666,817 by Daugherty et al.; in U.S. Patent No. 8,563,269 by Stagliano et al.; and in PCT Publication No. WO 2014/026136 by La Porte et al., the contents of each of which are hereby incorporated by reference in their entirety. (See also Boulware et al. "Evolutionary optimization of peptide substrates for proteases that exhibit rapid hydrolysis kinetics." Biotechnol Bioeng. 106.3 (2010): 339-46).

[000331] Exemplary substrates include but are not limited to substrates cleavable by one or more of the following enzymes or proteases listed in Table 4.

Table 4: Exemplary Proteases and/or Enzymes

ADAMS, ADAMTS, <i>e.g.</i> ADAM8 ADAM9 ADAM10 ADAM12 ADAM15 ADAM17/TACE ADAMDEC1 ADAMTS1 ADAMTS4 ADAMTS5	Cysteine proteinases, <i>e.g.</i> , Cruzipain Legumain Otubain-2	Serine proteases, <i>e.g.</i> , activated protein C Cathepsin A Cathepsin G Chymase coagulation factor proteases (<i>e.g.</i> , FVIIa, FIXa, FXa, FXIa, FXIIa)
		Elastase
		Granzyme B
		Guanidinobenzoatase
		HtrA1
Aspartate proteases, <i>e.g.</i> , BACE Renin	KLKs, <i>e.g.</i> , KLK4 KLK5 KLK6 KLK7 KLK8 KLK10 KLK11 KLK13 KLK14	Human Neutrophil Elastase Lactoferrin
		Marapsin NS3/4A PACE4 Plasmin PSA tPA Thrombin Trypsin uPA
Aspartic cathepsins, <i>e.g.</i> , Cathepsin D Cathepsin E	Metallo proteinases, <i>e.g.</i> , Meprin Neprilysin PSMA BMP-1	Type II Transmembrane Serine Proteases (TTSPs), <i>e.g.</i> , DESC1 DPP-4 FAP Hepsin Matriptase-2 MT-SP1/Matriptase TMPRSS2 TMPRSS3 TMPRSS4
Caspases, <i>e.g.</i> , Caspase 1 Caspase 2 Caspase 3 Caspase 4 Caspase 5 Caspase 6 Caspase 7 Caspase 8 Caspase 9 Caspase 10 Caspase 14	MMPs, <i>e.g.</i> , MMP1 MMP2 MMP3 MMP7 MMP8 MMP9 MMP10 MMP11 MMP12 MMP13 MMP14 MMP15 MMP16	
Cysteine cathepsins, <i>e.g.</i> , Cathepsin B Cathepsin C		

Cathepsin K	MMP17
Cathepsin L	MMP19
Cathepsin S	MMP20
Cathepsin V/L2	MMP23
Cathepsin X/Z/P	MMP24
	MMP26
	MMP27

[000332] The activatable anti-CD71 antibodies described herein overcome a limitation of antibody therapeutics, particularly antibody therapeutics that are known to be toxic to at least some degree *in vivo*. Target-mediated toxicity constitutes a major limitation for the development of therapeutic antibodies. The activatable anti-CD71 antibodies provided herein are designed to address the toxicity associated with the inhibition of the target in normal tissues by traditional therapeutic antibodies. These activatable anti-CD71 antibodies remain masked until proteolytically activated at the site of disease. Starting with an anti-CD71 antibody as a parental therapeutic antibody, the activatable anti-CD71 antibodies of the invention were engineered by coupling the antibody to an inhibitory mask through a linker that incorporates a protease substrate.

[000333] When the AB is modified with a MM and is in the presence of the target, specific binding of the AB to its target is reduced or inhibited, as compared to the specific binding of the AB not modified with an MM or the specific binding of the parental AB to the target.

[000334] The K_d of the AB modified with a MM towards the target is at least 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 50,000, 100,000, 500,000, 1,000,000, 5,000,000, 10,000,000, 50,000,000 or greater, or between 5-10, 10-100, 10-1,000, 10-10,000, 10-100,000, 10-1,000,000, 10-10,000,000, 100-1,000, 100-10,000, 100-100,000, 100-1,000,000, 100-10,000,000, 1,000-10,000, 1,000-100,000, 1,000-1,000,000, 1000-10,000,000, 10,000-100,000, 10,000-1,000,000, 10,000-10,000,000, 100,000-1,000,000, or 100,000-10,000,000 times greater than the K_d of the AB not modified with an MM or of the parental AB towards the target. Conversely, the binding affinity of the AB modified with a MM towards the target is at least 2, 3, 4, 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 50,000, 100,000, 500,000, 1,000,000, 5,000,000, 10,000,000, 50,000,000 or greater, or between 5-10, 10-100, 10-1,000, 10-10,000, 10-100,000, 10-1,000,000, 10-10,000,000, 100-1,000, 100-10,000, 100-100,000, 100-1,000,000, 100-10,000,000, 1,000-10,000, 1,000-100,000, 1,000-1,000,000, 1000-10,000,000, 10,000-100,000, 10,000-1,000,000, 10,000-10,000,000, 100,000-1,000,000, 100,000-10,000,000 times greater than the K_d of the AB not modified with an MM or of the parental AB towards the target.

10,000,000, 100,000-1,000,000, or 100,000-10,000,000 times lower than the binding affinity of the AB not modified with an MM or of the parental AB towards the target.

[000335] The dissociation constant (K_d) of the MM towards the AB is generally greater than the K_d of the AB towards the target. The K_d of the MM towards the AB can be at least 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 100,000, 1,000,000 or even 10,000,000 times greater than the K_d of the AB towards the target. Conversely, the binding affinity of the MM towards the AB is generally lower than the binding affinity of the AB towards the target. The binding affinity of MM towards the AB can be at least 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 100,000, 1,000,000 or even 10,000,000 times lower than the binding affinity of the AB towards the target.

[000336] In some embodiments, the dissociation constant (K_d) of the MM towards the AB is approximately equal to the K_d of the AB towards the target. In some embodiments, the dissociation constant (K_d) of the MM towards the AB is no more than the dissociation constant of the AB towards the target.

[000337] In some embodiments, the dissociation constant (K_d) of the MM towards the AB is less than the dissociation constant of the AB towards the target.

[000338] In some embodiments, the dissociation constant (K_d) of the MM towards the AB is greater than the dissociation constant of the AB towards the target.

[000339] In some embodiments, the MM has a K_d for binding to the AB that is no more than the K_d for binding of the AB to the target.

[000340] In some embodiments, the MM has a K_d for binding to the AB that is no less than the K_d for binding of the AB to the target.

[000341] In some embodiments, the MM has a K_d for binding to the AB that is approximately equal to the K_d for binding of the AB to the target.

[000342] In some embodiments, the MM has a K_d for binding to the AB that is less than the K_d for binding of the AB to the target.

[000343] In some embodiments, the MM has a K_d for binding to the AB that is greater than the K_d for binding of the AB to the target.

[000344] In some embodiments, the MM has a K_d for binding to the AB that is no more than 2, 3, 4, 5, 10, 25, 50, 100, 250, 500, or 1,000 fold greater than the K_d for binding of the AB to the target. In some embodiments, the MM has a K_d for binding to the AB that is between 1-5, 2-5, 2-10, 5-10, 5-20, 5-50, 5-100, 10-100, 10-1,000, 20-100, 20-1000, or 100-1,000 fold greater than the K_d for binding of the AB to the target.

[000345] In some embodiments, the MM has an affinity for binding to the AB that is less than the affinity of binding of the AB to the target.

[000346] In some embodiments, the MM has an affinity for binding to the AB that is no more than the affinity of binding of the AB to the target.

[000347] In some embodiments, the MM has an affinity for binding to the AB that is approximately equal of the affinity of binding of the AB to the target.

[000348] In some embodiments, the MM has an affinity for binding to the AB that is no less than the affinity of binding of the AB to the target.

[000349] In some embodiments, the MM has an affinity for binding to the AB that is greater than the affinity of binding of the AB to the target.

[000350] In some embodiments, the MM has an affinity for binding to the AB that is 2, 3, 4, 5, 10, 25, 50, 100, 250, 500, or 1,000 less than the affinity of binding of the AB to the target. In some embodiments, the MM has an affinity for binding to the AB that is between 1-5, 2-5, 2-10, 5-10, 5-20, 5-50, 5-100, 10-100, 10-1,000, 20-100, 20-1000, or 100-1,000 fold less than the affinity of binding of the AB to the target. In some embodiments, the MM has an affinity for binding to the AB that is 2 to 20 fold less than the affinity of binding of the AB to the target. In some embodiments, a MM not covalently linked to the AB and at equimolar concentration to the AB does not inhibit the binding of the AB to the target.

[000351] When the AB is modified with a MM and is in the presence of the target specific binding of the AB to its target is reduced or inhibited, as compared to the specific binding of the AB not modified with an MM or the specific binding of the parental AB to the target. When compared to the binding of the AB not modified with an MM or the binding of the parental AB to the target the AB's ability to bind the target when modified with an MM can be reduced by at least 50%, 60%, 70%, 80%, 90%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% and even 100% for at least 2, 4, 6, 8, 12, 28, 24, 30, 36, 48, 60, 72, 84, or 96 hours, or 5, 10, 15, 30, 45, 60, 90, 120, 150, or 180 days, or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months or more when measured *in vivo* or in an *in vitro* assay.

[000352] The MM inhibits the binding of the AB to the target. The MM binds the antigen binding domain of the AB and inhibits binding of the AB to the target. The MM can sterically inhibit the binding of the AB to the target. The MM can allosterically inhibit the binding of the AB to its target. In these embodiments when the AB is modified or coupled to a MM and in the presence of target there is no binding or substantially no binding of the

AB to the target, or no more than 0.001%, 0.01%, 0.1%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, or 50% binding of the AB to the target, as compared to the binding of the AB not modified with an MM, the parental AB, or the AB not coupled to an MM to the target, for at least 2, 4, 6, 8, 12, 28, 24, 30, 36, 48, 60, 72, 84, or 96 hours, or 5, 10, 15, 30, 45, 60, 90, 120, 150, or 180 days, or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months or longer when measured *in vivo* or in an *in vitro* assay.

[000353] When an AB is coupled to or modified by a MM, the MM ‘masks’ or reduces or otherwise inhibits the specific binding of the AB to the target. When an AB is coupled to or modified by a MM, such coupling or modification can effect a structural change that reduces or inhibits the ability of the AB to specifically bind its target.

[000354] An AB coupled to or modified with an MM can be represented by the following formulae (in order from an amino (N) terminal region to carboxyl (C) terminal region:

(MM)-(AB)

(AB)-(MM)

(MM)-L-(AB)

(AB)-L-(MM)

where MM is a masking moiety, the AB is an antibody or antibody fragment thereof, and the L is a linker. In many embodiments, it can be desirable to insert one or more linkers, *e.g.*, flexible linkers, into the composition so as to provide for flexibility.

[000355] In certain embodiments, the MM is not a natural binding partner of the AB. In some embodiments, the MM contains no or substantially no homology to any natural binding partner of the AB. In some embodiments, the MM is no more than 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% similar to any natural binding partner of the AB. In some embodiments, the MM is no more than 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 25% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 50% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 20% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 10% identical to any natural binding partner of the AB.

[000356] In some embodiments, the activatable antibodies include an AB that is modified by an MM and also includes one or more cleavable moieties (CM). Such activatable antibodies exhibit activatable/switchable binding, to the AB's target. Activatable antibodies generally include an antibody or antibody fragment (AB), modified by or coupled to a masking moiety (MM) and a modifiable or cleavable moiety (CM). In some embodiments, the CM contains an amino acid sequence that serves as a substrate for at least one protease.

[000357] The elements of the activatable antibodies are arranged so that the MM and CM are positioned such that in a cleaved (or relatively active) state and in the presence of a target, the AB binds a target while the activatable antibody is in an uncleaved (or relatively inactive) state in the presence of the target, specific binding of the AB to its target is reduced or inhibited. The specific binding of the AB to its target can be reduced due to the inhibition or masking of the AB's ability to specifically bind its target by the MM.

[000358] The K_d of the AB modified with a MM and a CM towards the target is at least 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 50,000, 100,000, 500,000, 1,000,000, 5,000,000, 10,000,000, 50,000,000 or greater, or between 5-10, 10-100, 10-1,000, 10-10,000, 10-100,000, 10-1,000,000, 10-10,000,000, 100-1,000, 100-10,000, 100-100,000, 100-1,000,000, 100-10,000,000, 1,000-10,000, 1,000-100,000, 1,000-1,000,000, 1000-10,000,000, 10,000-100,000, 10,000-1,000,000, 10,000-10,000,000, 100,000-1,000,000, or 100,000-10,000,000 times greater than the K_d of the AB not modified with an MM and a CM or of the parental AB towards the target. Conversely, the binding affinity of the AB modified with a MM and a CM towards the target is at least 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 50,000, 100,000, 500,000, 1,000,000, 5,000,000, 10,000,000, 50,000,000 or greater, or between 5-10, 10-100, 10-1,000, 10-10,000, 10-100,000, 10-1,000,000, 10-10,000,000, 100-1,000, 100-10,000, 100-100,000, 100-1,000,000, 100-10,000,000, 1,000-10,000, 1,000-100,000, 1,000-1,000,000, 1000-10,000,000, 10,000-100,000, 10,000-1,000,000, 10,000-10,000,000, 100,000-1,000,000, or 100,000-10,000,000 times lower than the binding affinity of the AB not modified with an MM and a CM or of the parental AB towards the target.

[000359] When the AB is modified with a MM and a CM and is in the presence of the target but not in the presence of a modifying agent (for example at least one protease), specific binding of the AB to its target is reduced or inhibited, as compared to the specific binding of the AB not modified with an MM and a CM or of the parental AB to the target.

When compared to the binding of the parental AB or the binding of an AB not modified with an MM and a CM to its target, the AB's ability to bind the target when modified with an MM and a CM can be reduced by at least 50%, 60%, 70%, 80%, 90%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% and even 100% for at least 2, 4, 6, 8, 12, 28, 24, 30, 36, 48, 60, 72, 84, or 96 hours or 5, 10, 15, 30, 45, 60, 90, 120, 150, or 180 days, or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months or longer when measured *in vivo* or in an *in vitro* assay.

[000360] As used herein, the term cleaved state refers to the condition of the activatable antibodies following modification of the CM by at least one protease. The term uncleaved state, as used herein, refers to the condition of the activatable antibodies in the absence of cleavage of the CM by a protease. As discussed above, the term “activatable antibodies” is used herein to refer to an activatable antibody in both its uncleaved (native) state, as well as in its cleaved state. It will be apparent to the ordinarily skilled artisan that in some embodiments a cleaved activatable antibody may lack an MM due to cleavage of the CM by protease, resulting in release of at least the MM (*e.g.*, where the MM is not joined to the activatable antibodies by a covalent bond (*e.g.*, a disulfide bond between cysteine residues).

[000361] By activatable or switchable is meant that the activatable antibody exhibits a first level of binding to a target when the activatable antibody is in a inhibited, masked or uncleaved state (*i.e.*, a first conformation), and a second level of binding to the target in the uninhibited, unmasked and/or cleaved state (*i.e.*, a second conformation), where the second level of target binding is greater than the first level of binding. In general, the access of target to the AB of the activatable antibody is greater in the presence of a cleaving agent capable of cleaving the CM, *i.e.*, a protease, than in the absence of such a cleaving agent. Thus, when the activatable antibody is in the uncleaved state, the AB is inhibited from target binding and can be masked from target binding (*i.e.*, the first conformation is such the AB cannot bind the target), and in the cleaved state the AB is not inhibited or is unmasked to target binding.

[000362] The CM and AB of the activatable antibodies are selected so that the AB represents a binding moiety for a given target, and the CM represents a substrate for a protease. In some embodiments, the protease is co-localized with the target at a treatment site or diagnostic site in a subject. As used herein, co-localized refers to being at the same site or relatively close nearby. In some embodiments, a protease cleaves a CM yielding an activated antibody that binds to a target located nearby the cleavage site. The activatable

antibodies disclosed herein find particular use where, for example, a protease capable of cleaving a site in the CM, i.e., a protease, is present at relatively higher levels in target-containing tissue of a treatment site or diagnostic site than in tissue of non-treatment sites (for example in healthy tissue). In some embodiments, a CM of the disclosure is also cleaved by one or more other proteases. In some embodiments, it is the one or more other proteases that is co-localized with the target and that is responsible for cleavage of the CM *in vivo*.

[000363] In some embodiments activatable antibodies provide for reduced toxicity and/or adverse side effects that could otherwise result from binding of the AB at non-treatment sites if the AB were not masked or otherwise inhibited from binding to the target.

[000364] In general, an activatable antibody can be designed by selecting an AB of interest and constructing the remainder of the activatable antibody so that, when conformationally constrained, the MM provides for masking of the AB or reduction of binding of the AB to its target. Structural design criteria can be taken into account to provide for this functional feature.

[000365] Activatable antibodies exhibiting a switchable phenotype of a desired dynamic range for target binding in an inhibited versus an uninhibited conformation are provided. Dynamic range generally refers to a ratio of (a) a maximum detected level of a parameter under a first set of conditions to (b) a minimum detected value of that parameter under a second set of conditions. For example, in the context of an activatable antibody, the dynamic range refers to the ratio of (a) a maximum detected level of target protein binding to an activatable antibody in the presence of at least one protease capable of cleaving the CM of the activatable antibodies to (b) a minimum detected level of target protein binding to an activatable antibody in the absence of the protease. The dynamic range of an activatable antibody can be calculated as the ratio of the dissociation constant of an activatable antibody cleaving agent (*e.g.*, enzyme) treatment to the dissociation constant of the activatable antibodies cleaving agent treatment. The greater the dynamic range of an activatable antibody, the better the switchable phenotype of the activatable antibody. Activatable antibodies having relatively higher dynamic range values (*e.g.*, greater than 1) exhibit more desirable switching phenotypes such that target protein binding by the activatable antibodies occurs to a greater extent (*e.g.*, predominantly occurs) in the presence of a cleaving agent (*e.g.*, enzyme) capable of cleaving the CM of the activatable antibodies than in the absence of a cleaving agent.

[000366] Activatable antibodies can be provided in a variety of structural configurations. Exemplary formulae for activatable antibodies are provided below. It is specifically contemplated that the N- to C-terminal order of the AB, MM and CM can be reversed within an activatable antibody. It is also specifically contemplated that the CM and MM may overlap in amino acid sequence, *e.g.*, such that the CM is contained within the MM.

[000367] For example, activatable antibodies can be represented by the following formula (in order from an amino (N) terminal region to carboxyl (C) terminal region:

(MM)-(CM)-(AB)

(AB)-(CM)-(MM)

where MM is a masking moiety, CM is a cleavable moiety, and AB is an antibody or fragment thereof. It should be noted that although MM and CM are indicated as distinct components in the formulae above, in all exemplary embodiments (including formulae) disclosed herein it is contemplated that the amino acid sequences of the MM and the CM could overlap, *e.g.*, such that the CM is completely or partially contained within the MM. In addition, the formulae above provide for additional amino acid sequences that can be positioned N-terminal or C-terminal to the activatable antibodies elements.

[000368] In certain embodiments, the MM is not a natural binding partner of the AB. In some embodiments, the MM contains no or substantially no homology to any natural binding partner of the AB. In some embodiments, the MM is no more than 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% similar to any natural binding partner of the AB. In some embodiments, the MM is no more than 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 50% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 25% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 20% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 10% identical to any natural binding partner of the AB.

[000369] In many embodiments it may be desirable to insert one or more linkers, *e.g.*, flexible linkers, into the activatable antibody construct so as to provide for flexibility at one or more of the MM-CM junction, the CM-AB junction, or both. For example, the AB, MM, and/or CM may not contain a sufficient number of residues (*e.g.*, Gly, Ser, Asp, Asn,

especially Gly and Ser, particularly Gly) to provide the desired flexibility. As such, the switchable phenotype of such activatable antibody constructs may benefit from introduction of one or more amino acids to provide for a flexible linker. In addition, as described below, where the activatable antibody is provided as a conformationally constrained construct, a flexible linker can be operably inserted to facilitate formation and maintenance of a cyclic structure in the uncleaved activatable antibody.

[000370] For example, in certain embodiments an activatable antibody comprises one of the following formulae (where the formula below represent an amino acid sequence in either N- to C-terminal direction or C- to N-terminal direction):

(MM)-L1-(CM)-(AB)

(MM)-(CM)-L2-(AB)

(MM)-L1-(CM)-L2-(AB)

wherein MM, CM, and AB are as defined above; wherein L1 and L2 are each independently and optionally present or absent, are the same or different flexible linkers that include at least 1 flexible amino acid (*e.g.*, Gly). In addition, the formulae above provide for additional amino acid sequences that can be positioned N-terminal or C-terminal to the activatable antibodies elements. Examples include, but are not limited to, targeting moieties (*e.g.*, a ligand for a receptor of a cell present in a target tissue) and serum half-life extending moieties (*e.g.*, polypeptides that bind serum proteins, such as immunoglobulin (*e.g.*, IgG) or serum albumin (*e.g.*, human serum albumin (HAS)).

[000371] The CM is specifically cleaved by at least one protease at a rate of about $0.001\text{--}1500 \times 10^4 \text{ M}^{-1}\text{S}^{-1}$ or at least 0.001, 0.005, 0.01, 0.05, 0.1, 0.5, 1, 2.5, 5, 7.5, 10, 15, 20, 25, 50, 75, 100, 125, 150, 200, 250, 500, 750, 1000, 1250, or $1500 \times 10^4 \text{ M}^{-1}\text{S}^{-1}$. In some embodiments, the CM is specifically cleaved at a rate of about $100,000 \text{ M}^{-1}\text{S}^{-1}$. In some embodiments, the CM is specifically cleaved at a rate from about 1×10^2 to about $1 \times 10^6 \text{ M}^{-1}\text{S}^{-1}$ (*i.e.*, from about 1×10^2 to about $1 \times 10^6 \text{ M}^{-1}\text{S}^{-1}$).

[000372] For specific cleavage by an enzyme, contact between the enzyme and CM is made. When the activatable antibody comprising an AB coupled to a MM and a CM is in the presence of target and sufficient enzyme activity, the CM can be cleaved. Sufficient enzyme activity can refer to the ability of the enzyme to make contact with the CM and effect cleavage. It can readily be envisioned that an enzyme may be in the vicinity of the CM but unable to cleave because of other cellular factors or protein modification of the enzyme.

[000373] Linkers suitable for use in compositions described herein are generally ones that provide flexibility of the modified AB or the activatable antibodies to facilitate the inhibition of the binding of the AB to the target. Such linkers are generally referred to as flexible linkers. Suitable linkers can be readily selected and can be of any of a suitable of different lengths, such as from 1 amino acid (*e.g.*, Gly) to 20 amino acids, from 2 amino acids to 15 amino acids, from 3 amino acids to 12 amino acids, including 4 amino acids to 10 amino acids, 5 amino acids to 9 amino acids, 6 amino acids to 8 amino acids, or 7 amino acids to 8 amino acids, and can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acids in length.

[000374] Exemplary flexible linkers include glycine polymers (G)_n, glycine-serine polymers (including, for example, (GS)_n, (GSGGS)_n (SEQ ID NO: 339) and (GGGS)_n (SEQ ID NO: 340), where n is an integer of at least one), glycine-alanine polymers, alanine-serine polymers, and other flexible linkers known in the art. Glycine and glycine-serine polymers are relatively unstructured, and therefore may be able to serve as a neutral tether between components. Glycine accesses significantly more phi-psi space than even alanine, and is much less restricted than residues with longer side chains (see Scheraga, *Rev. Computational Chem.* 11173-142 (1992)). Exemplary flexible linkers include, but are not limited to Gly-Gly-Ser-Gly (SEQ ID NO: 341), Gly-Gly-Ser-Gly-Gly (SEQ ID NO: 342), Gly-Ser-Gly-Ser-Gly (SEQ ID NO: 343), Gly-Ser-Gly-Gly-Gly (SEQ ID NO: 344), Gly-Gly-Gly-Ser-Gly (SEQ ID NO: 345), Gly-Ser-Ser-Ser-Gly (SEQ ID NO: 346), and the like. The ordinarily skilled artisan will recognize that design of an activatable antibodies can include linkers that are all or partially flexible, such that the linker can include a flexible linker as well as one or more portions that confer less flexible structure to provide for a desired activatable antibodies structure.

[000375] The disclosure also provides compositions and methods that include an activatable anti-CD71 antibody that includes an antibody or antibody fragment (AB) that specifically binds CD71, where the AB is coupled to a masking moiety (MM) that decreases the ability of the AB to bind its target. In some embodiments, the activatable anti-CD71 antibody further includes a cleavable moiety (CM) that is a substrate for a protease. The compositions and methods provided herein enable the attachment of one or more agents to one or more cysteine residues in the AB without compromising the activity (*e.g.*, the masking, activating or binding activity) of the activatable anti-CD71 antibody. In some embodiments, the compositions and methods provided herein enable the attachment of one

or more agents to one or more cysteine residues in the AB without reducing or otherwise disturbing one or more disulfide bonds within the MM. The compositions and methods provided herein produce an activatable anti-CD71 antibody that is conjugated to one or more agents, *e.g.*, any of a variety of therapeutic, diagnostic and/or prophylactic agents, for example, in some embodiments, without any of the agent(s) being conjugated to the MM of the activatable anti-CD71 antibody. The compositions and methods provided herein produce conjugated activatable anti-CD71 antibodies in which the MM retains the ability to effectively and efficiently mask the AB of the activatable antibody in an uncleaved state. The compositions and methods provided herein produce conjugated activatable anti-CD71 antibodies in which the activatable antibody is still activated, *i.e.*, cleaved, in the presence of a protease that can cleave the CM.

[000376] The activatable anti-CD71 antibodies have at least one point of conjugation for an agent, but in the methods and compositions provided herein less than all possible points of conjugation are available for conjugation to an agent. In some embodiments, the one or more points of conjugation are sulfur atoms involved in disulfide bonds. In some embodiments, the one or more points of conjugation are sulfur atoms involved in interchain disulfide bonds. In some embodiments, the one or more points of conjugation are sulfur atoms involved in interchain sulfide bonds, but not sulfur atoms involved in intrachain disulfide bonds. In some embodiments, the one or more points of conjugation are sulfur atoms of cysteine or other amino acid residues containing a sulfur atom. Such residues may occur naturally in the antibody structure or can be incorporated into the antibody by site-directed mutagenesis, chemical conversion, or mis-incorporation of non-natural amino acids.

[000377] Also provided are methods of preparing a conjugate of an activatable anti-CD71 antibody having one or more interchain disulfide bonds in the AB and one or more intrachain disulfide bonds in the MM, and a drug reactive with free thiols is provided. The method generally includes partially reducing interchain disulfide bonds in the activatable antibody with a reducing agent, such as, for example, TCEP; and conjugating the drug reactive with free thiols to the partially reduced activatable antibody. As used herein, the term partial reduction refers to situations where an activatable anti-CD71 antibody is contacted with a reducing agent and less than all disulfide bonds, *e.g.*, less than all possible sites of conjugation are reduced. In some embodiments, less than 99%, 98%, 97%, 96%,

95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10% or less than 5% of all possible sites of conjugation are reduced.

[000378] In yet other embodiments, a method of reducing and conjugating an agent, e.g., a drug, to an activatable anti-CD71 antibody resulting in selectivity in the placement of the agent is provided. The method generally includes partially reducing the activatable anti-CD71 antibody with a reducing agent such that any conjugation sites in the masking moiety or other non-AB portion of the activatable antibody are not reduced, and conjugating the agent to interchain thiols in the AB. The conjugation site(s) are selected so as to allow desired placement of an agent to allow conjugation to occur at a desired site. The reducing agent is, for example, TCEP. The reduction reaction conditions such as, for example, the ratio of reducing agent to activatable antibody, the length of incubation, the temperature during the incubation, the pH of the reducing reaction solution, etc., are determined by identifying the conditions that produce a conjugated activatable antibody in which the MM retains the ability to effectively and efficiently mask the AB of the activatable antibody in an uncleaved state. The ratio of reduction agent to activatable anti-CD71 antibody will vary depending on the activatable antibody. In some embodiments, the ratio of reducing agent to activatable anti-CD71 antibody will be in a range from about 20:1 to 1:1, from about 10:1 to 1:1, from about 9:1 to 1:1, from about 8:1 to 1:1, from about 7:1 to 1:1, from about 6:1 to 1:1, from about 5:1 to 1:1, from about 4:1 to 1:1, from about 3:1 to 1:1, from about 2:1 to 1:1, from about 20:1 to 1:1.5, from about 10:1 to 1:1.5, from about 9:1 to 1:1.5, from about 8:1 to 1:1.5, from about 7:1 to 1:1.5, from about 6:1 to 1:1.5, from about 5:1 to 1:1.5, from about 4:1 to 1:1.5, from about 3:1 to 1:1.5, from about 2:1 to 1:1.5, from about 1.5:1 to 1:1.5, or from about 1:1 to 1:1.5. In some embodiments, the ratio is in a range of from about 5:1 to 1:1. In some embodiments, the ratio is in a range of from about 5:1 to 1.5:1. In some embodiments, the ratio is in a range of from about 4:1 to 1:1. In some embodiments, the ratio is in a range from about 4:1 to 1.5:1. In some embodiments, the ratio is in a range from about 8:1 to about 1:1. In some embodiments, the ratio is in a range of from about 2.5:1 to 1:1.

[000379] In some embodiments, a method of reducing interchain disulfide bonds in the AB of an activatable anti-CD71 antibody and conjugating an agent, e.g., a thiol-containing agent such as a drug, to the resulting interchain thiols to selectively locate agent(s) on the AB is provided. The method generally includes partially reducing the AB with a reducing agent to form at least two interchain thiols without forming all possible interchain thiols in

the activatable antibody; and conjugating the agent to the interchain thiols of the partially reduced AB. For example, the AB of the activatable antibody is partially reduced for about 1 hour at about 37°C at a desired ratio of reducing agent:activatable antibody. In some embodiments, the ratio of reducing agent to activatable antibody will be in a range from about 20:1 to 1:1, from about 10:1 to 1:1, from about 9:1 to 1:1, from about 8:1 to 1:1, from about 7:1 to 1:1, from about 6:1 to 1:1, from about 5:1 to 1:1, from about 4:1 to 1:1, from about 3:1 to 1:1, from about 2:1 to 1:1, from about 20:1 to 1:1.5, from about 10:1 to 1:1.5, from about 9:1 to 1:1.5, from about 8:1 to 1:1.5, from about 7:1 to 1:1.5, from about 6:1 to 1:1.5, from about 5:1 to 1:1.5, from about 4:1 to 1:1.5, from about 3:1 to 1:1.5, from about 2:1 to 1:1.5, from about 1.5:1 to 1:1.5, or from about 1:1 to 1:1.5. In some embodiments, the ratio is in a range of from about 5:1 to 1:1. In some embodiments, the ratio is in a range of from about 5:1 to 1.5:1. In some embodiments, the ratio is in a range of from about 4:1 to 1:1. In some embodiments, the ratio is in a range from about 4:1 to 1.5:1. In some embodiments, the ratio is in a range from about 8:1 to about 1:1. In some embodiments, the ratio is in a range of from about 2.5:1 to 1:1.

[000380] The thiol-containing reagent can be, for example, cysteine or N-acetyl cysteine. The reducing agent can be, for example, TCEP. In some embodiments, the reduced activatable antibody can be purified prior to conjugation, using for example, column chromatography, dialysis, or diafiltration. Alternatively, the reduced antibody is not purified after partial reduction and prior to conjugation.

[000381] The invention also provides partially reduced activatable anti-CD71 antibodies in which at least one interchain disulfide bond in the activatable antibody has been reduced with a reducing agent without disturbing any intrachain disulfide bonds in the activatable antibody, wherein the activatable antibody includes an antibody or an antigen binding fragment thereof (AB) that specifically binds to CD71, a masking moiety (MM) that inhibits the binding of the AB of the activatable antibody in an uncleaved state to the CD71 target, and a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease. In some embodiments the MM is coupled to the AB via the CM. In some embodiments, one or more intrachain disulfide bond(s) of the activatable antibody is not disturbed by the reducing agent. In some embodiments, one or more intrachain disulfide bond(s) of the MM within the activatable antibody is not disturbed by the reducing agent. In some embodiments, the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-

terminus as follows: MM-CM-AB or AB-CM-MM. In some embodiments, reducing agent is TCEP.

[000382] In yet other embodiments, a method of reducing and conjugating an agent, e.g., a drug, to an activatable anti-CD71 antibody resulting in selectivity in the placement of the agent by providing an activatable anti-CD71 antibody with a defined number and positions of lysine and/or cysteine residues. In some embodiments, the defined number of lysine and/or cysteine residues is higher or lower than the number of corresponding residues in the amino acid sequence of the parent antibody or activatable antibody. In some embodiments, the defined number of lysine and/or cysteine residues may result in a defined number of agent equivalents that can be conjugated to the anti-CD71 antibody or activatable anti-CD71 antibody. In some embodiments, the defined number of lysine and/or cysteine residues may result in a defined number of agent equivalents that can be conjugated to the anti-CD71 antibody or activatable anti-CD71 antibody in a site-specific manner. In some embodiments, the modified activatable antibody is modified with one or more non-natural amino acids in a site-specific manner, thus in some embodiments limiting the conjugation of the agents to only the sites of the non-natural amino acids. In some embodiments, the anti-CD71 antibody or activatable anti-CD71 antibody with a defined number and positions of lysine and/or cysteine residues can be partially reduced with a reducing agent as discussed herein such that any conjugation sites in the masking moiety or other non-AB portion of the activatable antibody are not reduced, and conjugating the agent to interchain thiols in the AB.

[000383] The disclosure also provides partially reduced activatable antibodies in which at least one interchain disulfide bond in the activatable antibody has been reduced with a reducing agent without disturbing any intrachain disulfide bonds in the activatable antibody, wherein the activatable antibody includes an antibody or an antigen binding fragment thereof (AB) that specifically binds to the target, e.g., CD71, a masking moiety (MM) that inhibits the binding of the AB of the activatable antibody in an uncleaved state to the target, and a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for at least one protease. In some embodiments, the MM is coupled to the AB via the CM. In some embodiments, one or more intrachain disulfide bond(s) of the activatable antibody is not disturbed by the reducing agent. In some embodiments, one or more intrachain disulfide bond(s) of the MM within the activatable antibody is not disturbed by the reducing agent. In some embodiments, the activatable

antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM. In some embodiments, reducing agent is TCEP.

[000384] In some embodiments, the activatable antibodies described herein also include an agent conjugated to the activatable antibody. In some embodiments, the conjugated agent is a therapeutic agent, such as an anti-inflammatory and/or an antineoplastic agent. In such embodiments, the agent is conjugated to a carbohydrate moiety of the activatable antibody, for example, in some embodiments, where the carbohydrate moiety is located outside the antigen-binding region of the antibody or antigen-binding fragment in the activatable antibody. In some embodiments, the agent is conjugated to a sulfhydryl group of the antibody or antigen-binding fragment in the activatable antibody.

[000385] In some embodiments, the agent is a cytotoxic agent such as a toxin (*e.g.*, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (*i.e.*, a radioconjugate).


[000386] In some embodiments, the agent is a detectable moiety such as, for example, a label or other marker. For example, the agent is or includes a radiolabeled amino acid, one or more biotinyl moieties that can be detected by marked avidin (*e.g.*, streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or calorimetric methods), one or more radioisotopes or radionuclides, one or more fluorescent labels, one or more enzymatic labels, and/or one or more chemiluminescent agents. In some embodiments, detectable moieties are attached by spacer molecules.

[000387] The disclosure also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a toxin (*e.g.*, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (*i.e.*, a radioconjugate). Suitable cytotoxic agents include, for example, dolastatins and derivatives thereof (*e.g.* auristatin E, AFP, MMAF, MMAE, MMAD, DMAF, DMAE). For example, the agent is monomethyl auristatin E (MMAE) or monomethyl auristatin D (MMAD). In some embodiments, the agent is an agent selected from the group listed in Table 5. In some embodiments, the agent is a dolastatin. In some embodiments, the agent is an auristatin or derivative thereof. In some embodiments, the agent is auristatin E or a derivative thereof. In some embodiments, the agent is monomethyl auristatin E (MMAE). In some embodiments, the agent is monomethyl auristatin D (MMAD). In some embodiments, the agent is a maytansinoid or maytansinoid derivative. In some embodiments, the agent is


DM1 or DM4. In some embodiments, the agent is a duocarmycin or derivative thereof. In some embodiments, the agent is a calicheamicin or derivative thereof. In some embodiments, the agent is a pyrrolobenzodiazepine. In some embodiments, the agent is a pyrrolobenzodiazepine dimer.

[000388] In some embodiments, the agent is linked to the AB using a maleimide caproyl-valine-citrulline linker or a maleimide PEG-valine-citrulline linker. In some embodiments, the agent is linked to the AB using a maleimide caproyl-valine-citrulline linker. In some embodiments, the agent is linked to the AB using a maleimide PEG-valine-citrulline linker. In some embodiments, the agent is monomethyl auristatin D (MMAD) linked to the AB using a maleimide PEG-valine-citrulline-para-aminobenzyloxycarbonyl linker, and this linker payload construct is referred to herein as “vc-MMAD.” In some embodiments, the agent is monomethyl auristatin E (MMAE) linked to the AB using a maleimide PEG-valine-citrulline-para-aminobenzyloxycarbonyl linker, and this linker payload construct is referred to herein as “vc-MMAE.” In some embodiments, the agent is linked to the AB using a maleimide PEG-valine-citrulline linker. In some embodiments, the agent is monomethyl auristatin D (MMAD) linked to the AB using a maleimide bis-PEG-valine-citrulline-para-aminobenzyloxycarbonyl linker, and this linker payload construct is referred to herein as “PEG2-vc-MMAD.” The structures of vc-MMAD, vc-MMAE, and PEG2-vc-MMAD are shown below:


vc-MMAD:



vc-MMAE:



PEG2-vc-MMAD:



[000389] The disclosure also provides conjugated activatable antibodies that include an activatable antibody linked to monomethyl auristatin D (MMAD) payload, wherein the activatable antibody includes an antibody or an antigen binding fragment thereof (AB) that specifically binds to a target, a masking moiety (MM) that inhibits the binding of the AB of the activatable antibody in an uncleaved state to the target, and cleavable moiety (CM) coupled to the AB, and the CM is a polypeptide that functions as a substrate for at least one MMP protease.

[000390] In some embodiments, the MMAD-conjugated activatable antibody can be conjugated using any of several methods for attaching agents to ABs: (a) attachment to the carbohydrate moieties of the AB, or (b) attachment to sulfhydryl groups of the AB, or (c) attachment to amino groups of the AB, or (d) attachment to carboxylate groups of the AB.

[000391] In some embodiments, the MMAD payload is conjugated to the AB via a linker. In some embodiments, the MMAD payload is conjugated to a cysteine in the AB via a linker. In some embodiments, the MMAD payload is conjugated to a lysine in the AB via a linker. In some embodiments, the MMAD payload is conjugated to another residue of the AB via a linker, such as those residues disclosed herein. In some embodiments, the linker is a thiol-containing linker. In some embodiments, the linker is a cleavable linker. In some embodiments, the linker is a non-cleavable linker. In some embodiments, the linker is selected from the group consisting of the linkers shown in Tables 6 and 7. In some embodiments, the activatable antibody and the MMAD payload are linked via a maleimide caproyl-valine-citrulline linker. In some embodiments, the activatable antibody and the MMAD payload are linked via a maleimide PEG-valine-citrulline linker. In some embodiments, the activatable antibody and the MMAD payload are linked via a maleimide caproyl-valine-citrulline-para-aminobenzyloxycarbonyl linker. In some embodiments, the

activatable antibody and the MMAD payload are linked via a maleimide PEG-valine-citrulline-para-aminobenzyloxycarbonyl linker. In some embodiments, the MMAD payload is conjugated to the AB using the partial reduction and conjugation technology disclosed herein.

[000392] In some embodiments, the polyethylene glycol (PEG) component of a linker of the present disclosure is formed from 2 ethylene glycol monomers, 3 ethylene glycol monomers, 4 ethylene glycol monomers, 5 ethylene glycol monomers, 6 ethylene glycol monomers, 7 ethylene glycol monomers, 8 ethylene glycol monomers, 9 ethylene glycol monomers, or at least 10 ethylene glycol monomers. In some embodiments of the present disclosure, the PEG component is a branched polymer. In some embodiments of the present disclosure, the PEG component is an unbranched polymer. In some embodiments, the PEG polymer component is functionalized with an amino group or derivative thereof, a carboxyl group or derivative thereof, or both an amino group or derivative thereof and a carboxyl group or derivative thereof.

[000393] In some embodiments, the PEG component of a linker of the present disclosure is an amino-tetra-ethylene glycol-carboxyl group or derivative thereof. In some embodiments, the PEG component of a linker of the present disclosure is an amino-tri-ethylene glycol-carboxyl group or derivative thereof. In some embodiments, the PEG component of a linker of the present disclosure is an amino-di-ethylene glycol-carboxyl group or derivative thereof. In some embodiments, an amino derivative is the formation of an amide bond between the amino group and a carboxyl group to which it is conjugated. In some embodiments, a carboxyl derivative is the formation of an amide bond between the carboxyl group and an amino group to which it is conjugated. In some embodiments, a carboxyl derivative is the formation of an ester bond between the carboxyl group and an hydroxyl group to which it is conjugated.

[000394] Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolacca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcun, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re .

[000395] Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., Science 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. (See WO94/11026).

[000396] Table 5 lists some of the exemplary pharmaceutical agents that can be employed in the herein described disclosure but in no way is meant to be an exhaustive list.

Table 5: Exemplary Pharmaceutical Agents for Conjugation

CYTOTOXIC AGENTS

Auristatins	Turbostatin
Auristatin E	Phenstatins
Monomethyl auristatin D (MMAD)	Hydroxyphenstatin
Monomethyl auristatin E (MMAE)	Spongistatin 5
Desmethyl auristatin E (DMAE)	Spongistatin 7
Auristatin F	Halistatin 1
Monomethyl auristatin F (MMAF)	Halistatin 2
Desmethyl auristatin F (DMAF)	Halistatin 3
Auristatin derivatives, <i>e.g.</i> , amides thereof	Modified Bryostatins
Auristatin tyramine	Halocomstatins
Auristatin quinoline	Pyrrolobenzimidazoles (PBI)
Dolastatins	Cibrostatin6
Dolastatin derivatives	Doxaliform
Dolastatin 16 DmJ	Anthracyclins analogues
Dolastatin 16 Dpv	
Maytansinoids, <i>e.g.</i> DM-1; DM-4	
Maytansinoid derivatives	Cemadotin analogue (CemCH2-SH)
Duocarmycin	Pseudomonas toxin A (PE38) variant
Duocarmycin derivatives	Pseudomonas toxin A (ZZ-PE38) variant
Alpha-amanitin	ZJ-101
Anthracyclines	OSW-1
Doxorubicin	4-Nitrobenzyloxycarbonyl Derivatives of
	O6-Benzylguanine
Daunorubicin	Topoisomerase inhibitors
Bryostatins	Hemiasterlin
Camptothecin	Cephalotaxine

Camptothecin derivatives
 7-substituted Camptothecin
 10, 11-
 Difluoromethylenedioxy camptothecin
 Combretastatins
 Debromoaplysiatoxin

Kahalalide-F
 Discodermolide
 Ecteinascidins

ANTIVIRALS

Acyclovir
 Vira A
 Symmetrel

ANTIFUNGALS

Nystatin

ADDITIONAL ANTI-NEOPLASTICS

Adriamycin
 Cerubidine
 Bleomycin
 Alkeran
 Velban
 Oncovin
 Fluorouracil
 Methotrexate
 Thiotepa
 Bisantrene
 Novantrone
 Thioguanine
 Procarabazine
 Cytarabine

ANTI-BACTERIALS

Aminoglycosides
 Streptomycin
 Neomycin
 Kanamycin
 Amikacin
 Gentamicin
 Tobramycin
 Streptomycin B
 Spectinomycin
 Ampicillin
 Sulfanilamide
 Polymyxin

Homoharringtonine
 Pyrrolobenzodiazepine dimers (PBDs)
 Pyrrolobenzodiazepenes

Functionalized pyrrolobenzodiazepenes
 Functionalized pyrrolobenzodiazepene
 dimers
 Calicheamicins
 Podophyllotoxins
 Taxanes
 Vinca alkaloids

CONJUGATABLE DETECTION REAGENTS

Fluorescein and derivatives thereof
 Fluorescein isothiocyanate (FITC)

RADIOPHARMACEUTICALS

¹²⁵I
¹³¹I
⁸⁹Zr
¹¹¹In
¹²³I
¹³¹I
^{99m}Tc
²⁰¹Tl
¹³³Xe
¹¹C
⁶²Cu
¹⁸F
⁶⁸Ga
¹³N
¹⁵O
³⁸K
⁸²Rb
^{99m}Tc (Technetium)

HEAVY METALS

Barium
 Gold
 Platinum

ANTI-MYCOPLASMAS

Tylosine
 Spectinomycin

Chloramphenicol

[000397] Those of ordinary skill in the art will recognize that a large variety of possible moieties can be coupled to the resultant antibodies of the disclosure. (*See, for example, "Conjugate Vaccines", Contributions to Microbiology and Immunology, J. M. Cruse and R. E. Lewis, Jr (eds), Carger Press, New York, (1989), the entire contents of which are incorporated herein by reference.*)

[000398] Coupling can be accomplished by any chemical reaction that will bind the two molecules so long as the antibody and the other moiety retain their respective activities. This linkage can include many chemical mechanisms, for instance covalent binding, affinity binding, intercalation, coordinate binding and complexation. In some embodiments, the binding is, however, covalent binding. Covalent binding can be achieved either by direct condensation of existing side chains or by the incorporation of external bridging molecules. Many bivalent or polyvalent linking agents are useful in coupling protein molecules, such as the antibodies of the present disclosure, to other molecules. For example, representative coupling agents can include organic compounds such as thioesters, carbodiimides, succinimide esters, diisocyanates, glutaraldehyde, diazobenzenes and hexamethylene diamines. This listing is not intended to be exhaustive of the various classes of coupling agents known in the art but, rather, is exemplary of the more common coupling agents. (*See Killen and Lindstrom, Jour. Immun. 133:1335-2549 (1984); Jansen et al., Immunological Reviews 62:185-216 (1982); and Vitetta et al., Science 238:1098 (1987).*)

[000399] In some embodiments, in addition to the compositions and methods provided herein, the conjugated activatable antibody can also be modified for site-specific conjugation through modified amino acid sequences inserted or otherwise included in the activatable antibody sequence. These modified amino acid sequences are designed to allow for controlled placement and/or dosage of the conjugated agent within a conjugated activatable antibody. For example, the activatable antibody can be engineered to include cysteine substitutions at positions on light and heavy chains that provide reactive thiol groups and do not negatively impact protein folding and assembly, nor alter antigen binding. In some embodiments, the activatable antibody can be engineered to include or otherwise introduce one or more non-natural amino acid residues within the activatable antibody to provide suitable sites for conjugation. In some embodiments, the activatable

antibody can be engineered to include or otherwise introduce enzymatically activatable peptide sequences within the activatable antibody sequence.

[000400] Suitable linkers are described in the literature. (*See, for example*, Ramakrishnan, S. et al., Cancer Res. 44:201-208 (1984) describing use of MBS (M-maleimidobenzoyl-N-hydroxysuccinimide ester). *See also*, U.S. Patent No. 5,030,719, describing use of halogenated acetyl hydrazide derivative coupled to an antibody by way of an oligopeptide linker. In some embodiments, suitable linkers include: (i) EDC (1-ethyl-3-(3-dimethylamino-propyl) carbodiimide hydrochloride); (ii) SMPT (4-succinimidylloxycarbonyl-alpha-methyl-alpha-(2-pyridyl-dithio)-toluene (Pierce Chem. Co., Cat. (21558G); (iii) SPDP (succinimidyl-6 [3-(2-pyridyldithio) propionamido]hexanoate (Pierce Chem. Co., Cat #21651G); (iv) Sulfo-LC-SPDP (sulfosuccinimidyl 6 [3-(2-pyridyldithio)-propionamide] hexanoate (Pierce Chem. Co. Cat. #2165-G); and (v) sulfo-NHS (N-hydroxysulfo-succinimide: Pierce Chem. Co., Cat. #24510) conjugated to EDC. Additional linkers include, but are not limited to, SMCC ((succinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate), sulfo-SMCC (sulfosuccinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate), SPDB (N-succinimidyl-4-(2-pyridyldithio) butanoate), or sulfo-SPDB (N-succinimidyl-4-(2-pyridyldithio)-2-sulfo butanoate).

[000401] The linkers described above contain components that have different attributes, thus leading to conjugates with differing physio-chemical properties. For example, sulfo-NHS esters of alkyl carboxylates are more stable than sulfo-NHS esters of aromatic carboxylates. NHS-ester containing linkers are less soluble than sulfo-NHS esters. Further, the linker SMPT contains a sterically hindered disulfide bond, and can form conjugates with increased stability. Disulfide linkages, are in general, less stable than other linkages because the disulfide linkage is cleaved *in vitro*, resulting in less conjugate available. Sulfo-NHS, in particular, can enhance the stability of carbodimide couplings. Carbodimide couplings (such as EDC) when used in conjunction with sulfo-NHS, forms esters that are more resistant to hydrolysis than the carbodimide coupling reaction alone.

[000402] In some embodiments, the linkers are cleavable. In some embodiments, the linkers are non-cleavable. In some embodiments, two or more linkers are present. The two or more linkers are all the same, *i.e.*, cleavable or non-cleavable, or the two or more linkers are different, *i.e.*, at least one cleavable and at least one non-cleavable.

[000403] The present disclosure utilizes several methods for attaching agents to ABs: (a) attachment to the carbohydrate moieties of the AB, or (b) attachment to sulfhydryl

groups of the AB, or (c) attachment to amino groups of the AB, or (d) attachment to carboxylate groups of the AB. According to the disclosure, ABs can be covalently attached to an agent through an intermediate linker having at least two reactive groups, one to react with AB and one to react with the agent. The linker, which may include any compatible organic compound, can be chosen such that the reaction with AB (or agent) does not adversely affect AB reactivity and selectivity. Furthermore, the attachment of linker to agent might not destroy the activity of the agent. Suitable linkers for reaction with oxidized antibodies or oxidized antibody fragments include those containing an amine selected from the group consisting of primary amine, secondary amine, hydrazine, hydrazide, hydroxylamine, phenylhydrazine, semicarbazide and thiosemicarbazide groups. Such reactive functional groups may exist as part of the structure of the linker, or can be introduced by suitable chemical modification of linkers not containing such groups.

[000404] According to the present disclosure, suitable linkers for attachment to reduced ABs include those having certain reactive groups capable of reaction with a sulfhydryl group of a reduced antibody or fragment. Such reactive groups include, but are not limited to: reactive haloalkyl groups (including, for example, haloacetyl groups), p-mercuribenzoate groups and groups capable of Michael-type addition reactions (including, for example, maleimides and groups of the type described by Mitra and Lawton, 1979, J. Amer. Chem. Soc. 101: 3097-3110).

[000405] According to the present disclosure, suitable linkers for attachment to neither oxidized nor reduced Abs include those having certain functional groups capable of reaction with the primary amino groups present in unmodified lysine residues in the Ab. Such reactive groups include, but are not limited to, NHS carboxylic or carbonic esters, sulfo-NHS carboxylic or carbonic esters, 4-nitrophenyl carboxylic or carbonic esters, pentafluorophenyl carboxylic or carbonic esters, acyl imidazoles, isocyanates, and isothiocyanates.

[000406] According to the present disclosure, suitable linkers for attachment to neither oxidized nor reduced Abs include those having certain functional groups capable of reaction with the carboxylic acid groups present in aspartate or glutamate residues in the Ab, which have been activated with suitable reagents. Suitable activating reagents include EDC, with or without added NHS or sulfo-NHS, and other dehydrating agents utilized for carboxamide formation. In these instances, the functional groups present in the suitable linkers would include primary and secondary amines, hydrazines, hydroxylamines, and hydrazides.

[000407] The agent can be attached to the linker before or after the linker is attached to the AB. In certain applications it may be desirable to first produce an AB-linker intermediate in which the linker is free of an associated agent. Depending upon the particular application, a specific agent may then be covalently attached to the linker. In some embodiments, the AB is first attached to the MM, CM and associated linkers and then attached to the linker for conjugation purposes.

[000408] *Branched Linkers:* In specific embodiments, branched linkers that have multiple sites for attachment of agents are utilized. For multiple site linkers, a single covalent attachment to an AB would result in an AB-linker intermediate capable of binding an agent at a number of sites. The sites can be aldehyde or sulfhydryl groups or any chemical site to which agents can be attached.

[000409] In some embodiments, higher specific activity (or higher ratio of agents to AB) can be achieved by attachment of a single site linker at a plurality of sites on the AB. This plurality of sites can be introduced into the AB by either of two methods. First, one may generate multiple aldehyde groups and/or sulfhydryl groups in the same AB. Second, one may attach to an aldehyde or sulfhydryl of the AB a "branched linker" having multiple functional sites for subsequent attachment to linkers. The functional sites of the branched linker or multiple site linker can be aldehyde or sulfhydryl groups, or can be any chemical site to which linkers can be attached. Still higher specific activities can be obtained by combining these two approaches, that is, attaching multiple site linkers at several sites on the AB.

[000410] *Cleavable Linkers:* Peptide linkers that are susceptible to cleavage by enzymes of the complement system, such as but not limited to u-plasminogen activator, tissue plasminogen activator, trypsin, plasmin, or another enzyme having proteolytic activity can be used in one embodiment of the present disclosure. According to one method of the present disclosure, an agent is attached via a linker susceptible to cleavage by complement. The antibody is selected from a class that can activate complement. The antibody-agent conjugate, thus, activates the complement cascade and releases the agent at the target site. According to another method of the present disclosure, an agent is attached via a linker susceptible to cleavage by enzymes having a proteolytic activity such as a u-plasminogen activator, a tissue plasminogen activator, plasmin, or trypsin. These cleavable linkers are useful in conjugated activatable antibodies that include an extracellular toxin, e.g., by way of non-limiting example, any of the extracellular toxins shown in Table 5.

[000411] Non-limiting examples of cleavable linker sequences are provided in Table 6.

Table 6: Exemplary Linker Sequences for Conjugation

Types of Cleavable Sequences	Amino Acid Sequence
<u>Plasmin cleavable sequences</u>	
Pro-urokinase	PRFKIIGG (SEQ ID NO: 615)
	PRFRIIGG (SEQ ID NO: 616)
TGFβ	SSRHRRALD (SEQ ID NO: 617)
Plasminogen	RKSSIIIRMRDVVL (SEQ ID NO: 618)
Staphylokinase	SSSFDKGKYKKGDDA (SEQ ID NO: 619)
	SSSFDKGKYKRGDDA (SEQ ID NO: 620)
<u>Factor Xa cleavable sequences</u>	
	IEGR (SEQ ID NO: 621)
	IDGR (SEQ ID NO: 622)
	GGSIDGR (SEQ ID NO: 623)
<u>MMP cleavable sequences</u>	
Gelatinase A	PLGLWA (SEQ ID NO: 624)
<u>Collagenase cleavable sequences</u>	
Calf skin collagen (α1(I) chain)	GPQGIAGQ (SEQ ID NO: 625)
Calf skin collagen (α2(I) chain)	GPQGLLGA (SEQ ID NO: 626)
Bovine cartilage collagen (α1(II) chain)	GIAGQ (SEQ ID NO: 627)
Human liver collagen (α1(III) chain)	GPLGIAGI (SEQ ID NO: 628)
Human α ₂ M	GPEGLRVG (SEQ ID NO: 629)
Human PZP	YGAGLGVV (SEQ ID NO: 630)
	AGLGVVER (SEQ ID NO: 631)
	AGLGISST (SEQ ID NO: 632)
Rat α ₁ M	EPQALAMS (SEQ ID NO: 633)
	QALAMSAI (SEQ ID NO: 634)
Rat α ₂ M	AAYHLVSQ (SEQ ID NO: 635)
	MDAFLESS (SEQ ID NO: 636)
Rat α ₁ I ₃ (2J)	ESLPVVAV (SEQ ID NO: 637)
Rat α ₁ I ₃ (27J)	SAPAVESE (SEQ ID NO: 638)
Human fibroblast collagenase	DVAQFVLT (SEQ ID NO: 639)
<u>(autolytic cleavages)</u>	VAQFVLTE (SEQ ID NO: 640)
	AQFVLTEG (SEQ ID NO: 641)
	PVQPIGPQ (SEQ ID NO: 642)

[000412] In addition, agents can be attached via disulfide bonds (for example, the disulfide bonds on a cysteine molecule) to the AB. Since many tumors naturally release high levels of glutathione (a reducing agent) this can reduce the disulfide bonds with subsequent release of the agent at the site of delivery. In some embodiments, the reducing

agent that would modify a CM would also modify the linker of the conjugated activatable antibody.

[000413] *Spacers and Cleavable Elements:* In some embodiments, it may be necessary to construct the linker in such a way as to optimize the spacing between the agent and the AB of the activatable antibody. This can be accomplished by use of a linker of the general structure:



wherein

W is either --NH--CH₂-- or --CH₂--;

Q is an amino acid, peptide; and

n is an integer from 0 to 20.

[000414] In some embodiments, the linker may comprise a spacer element and a cleavable element. The spacer element serves to position the cleavable element away from the core of the AB such that the cleavable element is more accessible to the enzyme responsible for cleavage. Certain of the branched linkers described above may serve as spacer elements.

[000415] Throughout this discussion, it should be understood that the attachment of linker to agent (or of spacer element to cleavable element, or cleavable element to agent) need not be particular mode of attachment or reaction. Any reaction providing a product of suitable stability and biological compatibility is acceptable.

[000416] *Serum Complement and Selection of Linkers:* According to one method of the present disclosure, when release of an agent is desired, an AB that is an antibody of a class that can activate complement is used. The resulting conjugate retains both the ability to bind antigen and activate the complement cascade. Thus, according to this embodiment of the present disclosure, an agent is joined to one end of the cleavable linker or cleavable element and the other end of the linker group is attached to a specific site on the AB. For example, if the agent has an hydroxy group or an amino group, it can be attached to the carboxy terminus of a peptide, amino acid or other suitably chosen linker via an ester or amide bond, respectively. For example, such agents can be attached to the linker peptide via a carbodimide reaction. If the agent contains functional groups that would interfere with attachment to the linker, these interfering functional groups can be blocked before attachment and deblocked once the product conjugate or intermediate is made. The opposite

or amino terminus of the linker is then used either directly or after further modification for binding to an AB that is capable of activating complement.

[000417] Linkers (or spacer elements of linkers) can be of any desired length, one end of which can be covalently attached to specific sites on the AB of the activatable antibody. The other end of the linker or spacer element can be attached to an amino acid or peptide linker.

[000418] Thus when these conjugates bind to antigen in the presence of complement the amide or ester bond that attaches the agent to the linker will be cleaved, resulting in release of the agent in its active form. These conjugates, when administered to a subject, will accomplish delivery and release of the agent at the target site, and are particularly effective for the *in vivo* delivery of pharmaceutical agents, antibiotics, antimetabolites, antiproliferative agents and the like as presented in but not limited to those in Table 5.

[000419] *Linkers for Release without Complement Activation:* In yet another application of targeted delivery, release of the agent without complement activation is desired since activation of the complement cascade will ultimately lyse the target cell. Hence, this approach is useful when delivery and release of the agent should be accomplished without killing the target cell. Such is the goal when delivery of cell mediators such as hormones, enzymes, corticosteroids, neurotransmitters, genes or enzymes to target cells is desired. These conjugates can be prepared by attaching the agent to an AB that is not capable of activating complement via a linker that is mildly susceptible to cleavage by serum proteases. When this conjugate is administered to an individual, antigen-antibody complexes will form quickly whereas cleavage of the agent will occur slowly, thus resulting in release of the compound at the target site.

[000420] *Biochemical Cross Linkers:* In some embodiments, the activatable antibody can be conjugated to one or more therapeutic agents using certain biochemical cross-linkers. Cross-linking reagents form molecular bridges that tie together functional groups of two different molecules. To link two different proteins in a step-wise manner, hetero-bifunctional cross-linkers can be used that eliminate unwanted homopolymer formation.

[000421] Peptidyl linkers cleavable by lysosomal proteases are also useful, for example, Val-Cit, Val-Ala or other dipeptides. In addition, acid-labile linkers cleavable in the low-pH environment of the lysosome can be used, for example: bis-sialyl ether. Other suitable linkers include cathepsin-labile substrates, particularly those that show optimal function at an acidic pH.

[000422] Exemplary hetero-bifunctional cross-linkers are referenced in Table 7.

Table 7: Exemplary Hetero-Bifunctional Cross Linkers

<u>HETERO-BIFUNCTIONAL CROSS-LINKERS</u>			
Linker	Reactive Toward	Advantages and Applications	Spacer Arm Length after cross-linking (Angstroms)
SMPT	Primary amines Sulfhydryls	Greater stability	11.2 Å
SPDP	Primary amines Sulfhydryls	Thiolation	6.8 Å
LC-SPDP	Primary amines Sulfhydryls	Cleavable cross-linking Extended spacer arm	15.6 Å
Sulfo-LC-SPDP	Primary amines Sulfhydryls	Extender spacer arm	15.6 Å
SMCC	Primary amines Sulfhydryls	Water-soluble Stable maleimide reactive group Enzyme-antibody conjugation Hapten-carrier protein conjugation	11.6 Å
Sulfo-SMCC	Primary amines Sulfhydryls	Stable maleimide reactive group Water-soluble Enzyme-antibody conjugation	11.6 Å
MBS	Primary amines Sulfhydryls	Enzyme-antibody conjugation Hapten-carrier protein conjugation	9.9 Å
Sulfo-MBS	Primary amines Sulfhydryls	Water-soluble	9.9 Å
SIAB	Primary amines Sulfhydryls	Enzyme-antibody conjugation	10.6 Å
Sulfo-SIAB	Primary amines Sulfhydryls	Water-soluble	10.6 Å
SMPB	Primary amines Sulfhydryls	Extended spacer arm Enzyme-antibody conjugation	14.5 Å
Sulfo-SMPB	Primary amines Sulfhydryls	Extended spacer arm Water-soluble	14.5 Å
EDE/Sulfo-NHS	Primary amines Carboxyl groups	Hapten-Carrier conjugation	0
ABH	Carbohydrates Nonselective	Reacts with sugar groups	11.9 Å

[000423] *Non-Cleavable Linkers or Direct Attachment:* In some embodiments of the disclosure, the conjugate can be designed so that the agent is delivered to the target but not released. This can be accomplished by attaching an agent to an AB either directly or via a non-cleavable linker.

[000424] These non-cleavable linkers may include amino acids, peptides, D-amino acids or other organic compounds that can be modified to include functional groups that can subsequently be utilized in attachment to ABs by the methods described herein. A general formula for such an organic linker could be



wherein

W is either --NH--CH₂-- or --CH₂--;

Q is an amino acid, peptide; and

n is an integer from 0 to 20.

[000425] *Non-Cleavable Conjugates:* In some embodiments, a compound can be attached to ABs that do not activate complement. When using ABs that are incapable of complement activation, this attachment can be accomplished using linkers that are susceptible to cleavage by activated complement or using linkers that are not susceptible to cleavage by activated complement.

[000426] The antibodies disclosed herein can also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang et al., Proc. Natl Acad. Sci. USA, 77: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

[000427] Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present disclosure can be conjugated to the liposomes as described in Martin et al., J. Biol. Chem., 257: 286-288 (1982) via a disulfide-interchange reaction.

Definitions:

[000428] Unless otherwise defined, scientific and technical terms used in connection with the present disclosure shall have the meanings that are commonly understood by those of ordinary skill in the art. The term "a" entity or "an" entity refers to one or more of that entity. For example, a compound refers to one or more compounds. As such, the terms "a", "an", "one or more" and "at least one" can be used interchangeably. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall

include the singular. Generally, nomenclatures utilized in connection with, and techniques of, cell and tissue culture, molecular biology, and protein and oligo- or polynucleotide chemistry and hybridization described herein are those well-known and commonly used in the art. Standard techniques are used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (*e.g.*, electroporation, lipofection). Enzymatic reactions and purification techniques are performed according to manufacturer's specifications or as commonly accomplished in the art or as described herein. The foregoing techniques and procedures are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. *See e.g.*, Sambrook *et al.* Molecular Cloning: A Laboratory Manual (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)). The nomenclatures utilized in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well-known and commonly used in the art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[000429] As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

[000430] As used herein, the term "antibody" refers to immunoglobulin molecules and immunologically active portions of immunoglobulin (Ig) molecules, *i.e.*, molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen. By "specifically bind" or "immunoreacts with" or "immunospecifically bind" is meant that the antibody reacts with one or more antigenic determinants of the desired antigen and does not react with other polypeptides or binds at much lower affinity ($K_d > 10^{-6}$). Antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, domain antibody, single chain, Fab, and F(ab')₂ fragments, scFvs, and an Fab expression library.

[000431] The basic antibody structural unit is known to comprise a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. In general, antibody molecules obtained from humans relate to any of the classes IgG, IgM, IgA, IgE and IgD,

which differ from one another by the nature of the heavy chain present in the molecule. Certain classes have subclasses as well, such as IgG₁, IgG₂, and others. Furthermore, in humans, the light chain can be a kappa chain or a lambda chain.

[000432] The term “monoclonal antibody” (mAb) or “monoclonal antibody composition”, as used herein, refers to a population of antibody molecules that contain only one molecular species of antibody molecule consisting of a unique light chain gene product and a unique heavy chain gene product. In particular, the complementarity determining regions (CDRs) of the monoclonal antibody are identical in all the molecules of the population. MAbs contain an antigen binding site capable of immunoreacting with a particular epitope of the antigen characterized by a unique binding affinity for it.

[000433] The term “antigen-binding site” or “binding portion” refers to the part of the immunoglobulin molecule that participates in antigen binding. The antigen binding site is formed by amino acid residues of the N-terminal variable (“V”) regions of the heavy (“H”) and light (“L”) chains. Three highly divergent stretches within the V regions of the heavy and light chains, referred to as “hypervariable regions,” are interposed between more conserved flanking stretches known as “framework regions,” or “FRs”. Thus, the term “FR” refers to amino acid sequences that are naturally found between, and adjacent to, hypervariable regions in immunoglobulins. In an antibody molecule, the three hypervariable regions of a light chain and the three hypervariable regions of a heavy chain are disposed relative to each other in three dimensional space to form an antigen-binding surface. The antigen-binding surface is complementary to the three-dimensional surface of a bound antigen, and the three hypervariable regions of each of the heavy and light chains are referred to as “complementarity-determining regions,” or “CDRs.” The assignment of amino acids to each domain is in accordance with the definitions of Kabat Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md. (1987 and 1991)), or Chothia & Lesk J. Mol. Biol. 196:901-917 (1987), Chothia *et al.* Nature 342:878-883 (1989).

[000434] As used herein, the term “epitope” includes any protein determinant capable of specific binding to an immunoglobulin, an scFv, or a T-cell receptor. The term “epitope” includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor. Epitopic determinants usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics. For

example, antibodies can be raised against N-terminal or C-terminal peptides of a polypeptide. An antibody is said to specifically bind an antigen when the dissociation constant is $\leq 1 \mu\text{M}$; in some embodiments, $\leq 100 \text{ nM}$ and in some embodiments, $\leq 10 \text{ nM}$.

[000435] As used herein, the terms “specific binding,” “immunological binding,” and “immunological binding properties” refer to the non-covalent interactions of the type which occur between an immunoglobulin molecule and an antigen for which the immunoglobulin is specific. The strength, or affinity of immunological binding interactions can be expressed in terms of the dissociation constant (K_d) of the interaction, wherein a smaller K_d represents a greater affinity. Immunological binding properties of selected polypeptides can be quantified using methods well known in the art. One such method entails measuring the rates of antigen-binding site/antigen complex formation and dissociation, wherein those rates depend on the concentrations of the complex partners, the affinity of the interaction, and geometric parameters that equally influence the rate in both directions. Thus, both the “on rate constant” (K_{on}) and the “off rate constant” (K_{off}) can be determined by calculation of the concentrations and the actual rates of association and dissociation. (*See* Nature 361:186-87 (1993)). The ratio of K_{off}/K_{on} enables the cancellation of all parameters not related to affinity, and is equal to the dissociation constant K_d . (*See, generally*, Davies et al. (1990) Annual Rev Biochem 59:439-473). An antibody of the present disclosure is said to specifically bind to the target, when the binding constant (K_d) is $\leq 1 \mu\text{M}$, in some embodiments $\leq 100 \text{ nM}$, in some embodiments $\leq 10 \text{ nM}$, and in some embodiments $\leq 100 \text{ pM}$ to about 1 pM , as measured by assays such as radioligand binding assays or similar assays known to those skilled in the art.

[000436] The term “isolated polynucleotide” as used herein shall mean a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, which by virtue of its origin the “isolated polynucleotide” (1) is not associated with all or a portion of a polynucleotide in which the “isolated polynucleotide” is found in nature, (2) is operably linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence. Polynucleotides in accordance with the disclosure include the nucleic acid molecules encoding the heavy chain immunoglobulin molecules shown herein, and nucleic acid molecules encoding the light chain immunoglobulin molecules shown herein.

[000437] The term “isolated protein” referred to herein means a protein of cDNA, recombinant RNA, or synthetic origin or some combination thereof, which by virtue of its

origin, or source of derivation, the “isolated protein” (1) is not associated with proteins found in nature, (2) is free of other proteins from the same source, *e.g.*, free of murine proteins, (3) is expressed by a cell from a different species, or (4) does not occur in nature.

[000438] The term “polypeptide” is used herein as a generic term to refer to native protein, fragments, or analogs of a polypeptide sequence. Hence, native protein fragments, and analogs are species of the polypeptide genus. Polypeptides in accordance with the disclosure comprise the heavy chain immunoglobulin molecules shown herein, and the light chain immunoglobulin molecules shown herein, as well as antibody molecules formed by combinations comprising the heavy chain immunoglobulin molecules with light chain immunoglobulin molecules, such as kappa light chain immunoglobulin molecules, and vice versa, as well as fragments and analogs thereof.

[000439] The term “naturally-occurring” as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and that has not been intentionally modified by man in the laboratory or otherwise is naturally-occurring.

[000440] The term “operably linked” as used herein refers to positions of components so described are in a relationship permitting them to function in their intended manner. A control sequence “operably linked” to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

[000441] The term “control sequence” as used herein refers to polynucleotide sequences that are necessary to effect the expression and processing of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and transcription termination sequence in eukaryotes, generally, such control sequences include promoters and transcription termination sequence. The term “control sequences” is intended to include, at a minimum, all components whose presence is essential for expression and processing, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences. The term “polynucleotide” as referred to herein means nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA.

[000442] The term oligonucleotide referred to herein includes naturally occurring, and modified nucleotides linked together by naturally occurring, and non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide subset generally comprising a length of 200 bases or fewer. In some embodiments, oligonucleotides are 10 to 60 bases in length and in some embodiments, 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides are usually single stranded, *e.g.*, for probes, although oligonucleotides may be double stranded, *e.g.*, for use in the construction of a gene mutant. Oligonucleotides of the disclosure are either sense or antisense oligonucleotides.

[000443] The term “naturally occurring nucleotides” referred to herein includes deoxyribonucleotides and ribonucleotides. The term “modified nucleotides” referred to herein includes nucleotides with modified or substituted sugar groups and the like. The term “oligonucleotide linkages” referred to herein includes oligonucleotide linkages such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoronmidate, and the like. *See e.g.*, LaPlanche *et al.* Nucl. Acids Res. 14:9081 (1986); Stec *et al.* J. Am. Chem. Soc. 106:6077 (1984), Stein *et al.* Nucl. Acids Res. 16:3209 (1988), Zon *et al.* Anti Cancer Drug Design 6:539 (1991); Zon *et al.* Oligonucleotides and Analogues: A Practical Approach, pp. 87-108 (F. Eckstein, Ed., Oxford University Press, Oxford England (1991)); Stec *et al.* U.S. Patent No. 5,151,510; Uhlmann and Peyman Chemical Reviews 90:543 (1990). An oligonucleotide can include a label for detection, if desired.

[000444] As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. *See* Immunology - A Synthesis (2nd Edition, E.S. Golub and D.R. Green, Eds., Sinauer Associates, Sunderland, Mass. (1991)). Stereoisomers (*e.g.*, D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as α -, α -disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids may also be suitable components for polypeptides of the present disclosure. Examples of unconventional amino acids include: 4 hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, σ -N-methylarginine, and other similar amino acids and imino acids (*e.g.*, 4-hydroxyproline). In the polypeptide notation used herein, the left-hand direction is the amino terminal direction and the right-hand direction is the carboxy-terminal direction, in accordance with standard usage and convention.

[000445] Similarly, unless specified otherwise, the left-hand end of single-stranded polynucleotide sequences is the 5' end and the left-hand direction of double-stranded polynucleotide sequences is referred to as the 5' direction. The direction of 5' to 3' addition of nascent RNA transcripts is referred to as the transcription direction sequence regions on the DNA strand having the same sequence as the RNA and that are 5' to the 5' end of the RNA transcript are referred to as "upstream sequences", sequence regions on the DNA strand having the same sequence as the RNA and that are 3' to the 3' end of the RNA transcript are referred to as "downstream sequences".

[000446] As applied to polypeptides, the term "substantial identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 80 percent sequence identity, in some embodiments, at least 90 percent sequence identity, in some embodiments, at least 95 percent sequence identity, and in some embodiments, at least 99 percent sequence identity.

[000447] In some embodiments, residue positions that are not identical differ by conservative amino acid substitutions.

[000448] As discussed herein, minor variations in the amino acid sequences of antibodies or immunoglobulin molecules are contemplated as being encompassed by the present disclosure, providing that the variations in the amino acid sequence maintain at least 75%, in some embodiments, at least 80%, 90%, 95%, and in some embodiments, 99%. In particular, conservative amino acid replacements are contemplated. Conservative replacements are those that take place within a family of amino acids that are related in their side chains. Genetically encoded amino acids are generally divided into families: (1) acidic amino acids are aspartate, glutamate; (2) basic amino acids are lysine, arginine, histidine; (3) non-polar amino acids are alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan, and (4) uncharged polar amino acids are glycine, asparagine, glutamine, cysteine, serine, threonine, tyrosine. The hydrophilic amino acids include arginine, asparagine, aspartate, glutamine, glutamate, histidine, lysine, serine, and threonine. The hydrophobic amino acids include alanine, cysteine, isoleucine, leucine, methionine, phenylalanine, proline, tryptophan, tyrosine and valine. Other families of amino acids include (i) serine and threonine, which are the aliphatic-hydroxy family; (ii) asparagine and glutamine, which are the amide containing family; (iii) alanine, valine, leucine and isoleucine, which are the aliphatic family; and (iv) phenylalanine, tryptophan, and tyrosine, which are the aromatic family. For example, it is reasonable to expect that an isolated

replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the binding or properties of the resulting molecule, especially if the replacement does not involve an amino acid within a framework site. Whether an amino acid change results in a functional peptide can readily be determined by assaying the specific activity of the polypeptide derivative. Assays are described in detail herein. Fragments or analogs of antibodies or immunoglobulin molecules can be readily prepared by those of ordinary skill in the art. Suitable amino- and carboxy-termini of fragments or analogs occur near boundaries of functional domains. Structural and functional domains can be identified by comparison of the nucleotide and/or amino acid sequence data to public or proprietary sequence databases. In some embodiments, computerized comparison methods are used to identify sequence motifs or predicted protein conformation domains that occur in other proteins of known structure and/or function. Methods to identify protein sequences that fold into a known three-dimensional structure are known. Bowie *et al.* Science 253:164 (1991). Thus, the foregoing examples demonstrate that those of skill in the art can recognize sequence motifs and structural conformations that can be used to define structural and functional domains in accordance with the disclosure.

[000449] Suitable amino acid substitutions are those that: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinities, and (5) confer or modify other physicochemical or functional properties of such analogs. Analogs can include various muteins of a sequence other than the naturally-occurring peptide sequence. For example, single or multiple amino acid substitutions (for example, conservative amino acid substitutions) can be made in the naturally-occurring sequence (for example, in the portion of the polypeptide outside the domain(s) forming intermolecular contacts. A conservative amino acid substitution should not substantially change the structural characteristics of the parent sequence (*e.g.*, a replacement amino acid should not tend to break a helix that occurs in the parent sequence, or disrupt other types of secondary structure that characterizes the parent sequence). Examples of art-recognized polypeptide secondary and tertiary structures are described in Proteins, Structures and Molecular Principles (Creighton, Ed., W. H. Freeman and Company, New York (1984)); Introduction to Protein Structure (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)); and Thornton et al. Nature 354:105 (1991).

[000450] The term “polypeptide fragment” as used herein refers to a polypeptide that has an amino terminal and/or carboxy-terminal deletion and/or one or more internal deletion(s), but where the remaining amino acid sequence is identical to the corresponding positions in the naturally-occurring sequence deduced, for example, from a full length cDNA sequence. Fragments typically are at least 5, 6, 8 or 10 amino acids long, in some embodiments, at least 14 amino acids long, in some embodiments, at least 20 amino acids long, usually at least 50 amino acids long, and in some embodiments, at least 70 amino acids long. The term “analog” as used herein refers to polypeptides that are comprised of a segment of at least 25 amino acids that has substantial identity to a portion of a deduced amino acid sequence and that has specific binding to the target, under suitable binding conditions. Typically, polypeptide analogs comprise a conservative amino acid substitution (or addition or deletion) with respect to the naturally-occurring sequence. Analog typically are at least 20 amino acids long, in some embodiments, at least 50 amino acids long or longer, and can often be as long as a full-length naturally-occurring polypeptide.

[000451] The term “agent” is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials.

[000452] As used herein, the terms “label” or “labeled” refers to incorporation of a detectable marker, *e.g.*, by incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotinyl moieties that can be detected by marked avidin (*e.g.*, streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or calorimetric methods). In certain situations, the label or marker can also be therapeutic. Various methods of labeling polypeptides and glycoproteins are known in the art and can be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (*e.g.*, ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I), fluorescent labels (*e.g.*, FITC, rhodamine, lanthanide phosphors), enzymatic labels (*e.g.*, horseradish peroxidase, p-galactosidase, luciferase, alkaline phosphatase), chemiluminescent, biotinyl groups, predetermined polypeptide epitopes recognized by a secondary reporter (*e.g.*, leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance. The term “pharmaceutical agent or drug” as used herein refers to a chemical compound or composition capable of inducing a desired therapeutic effect when properly administered to a patient.

[000453] Other chemistry terms herein are used according to conventional usage in the art, as exemplified by The McGraw-Hill Dictionary of Chemical Terms (Parker, S., Ed., McGraw-Hill, San Francisco (1985)).

[000454] As used herein, “substantially pure” means an object species is the predominant species present (*i.e.*, on a molar basis it is more abundant than any other individual species in the composition), and in some embodiments, a substantially purified fraction is a composition wherein the object species comprises at least about 50 percent (on a molar basis) of all macromolecular species present.

[000455] Generally, a substantially pure composition will comprise more than about 80 percent of all macromolecular species present in the composition, in some embodiments, more than about 85%, 90%, 95%, and 99%. In some embodiments, the object species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species.

[000456] The term patient includes human and veterinary subjects.

[000457] Antibodies and/or activatable antibodies of the disclosure specifically bind a given target, *e.g.*, a human target protein such as human CD71. Also included in the disclosure are antibodies and/or activatable antibodies that bind to the same epitope as the antibodies and/or activatable antibodies described herein. Also included in the disclosure are antibodies and/or antibodies activatable antibodies that compete with an anti-CD71 antibody and/or an anti-CD71 activatable antibody described herein for binding to CD71, *e.g.*, human CD71. Also included in the disclosure are antibodies and/or antibodies activatable antibodies that cross-compete with an anti-CD71 antibody and/or an anti-CD71 activatable antibody described herein for binding to CD71, *e.g.*, human CD71.

[000458] Those skilled in the art will recognize that it is possible to determine, without undue experimentation, if a monoclonal antibody (*e.g.*, a murine monoclonal or humanized antibody) has the same specificity as a monoclonal antibody used in the methods described herein by ascertaining whether the former prevents the latter from binding to the target. If the monoclonal antibody being tested competes with the monoclonal antibody of the disclosure, as shown by a decrease in binding by the monoclonal antibody of the disclosure, then the two monoclonal antibodies bind to the same, or a closely related, epitope. An alternative method for determining whether a monoclonal antibody has the specificity of a monoclonal antibody of the disclosure is to pre-incubate the monoclonal antibody of the

disclosure with the target and then add the monoclonal antibody being tested to determine if the monoclonal antibody being tested is inhibited in its ability to bind the target. If the monoclonal antibody being tested is inhibited then, in all likelihood, it has the same, or functionally equivalent, epitopic specificity as the monoclonal antibody of the disclosure.

Multispecific Activatable Antibodies

[000459] The disclosure also provides multispecific anti-CD71 activatable antibodies. The multispecific activatable antibodies provided herein are multispecific antibodies that recognize CD71 and at least one or more different antigens or epitopes and that include at least one masking moiety (MM) linked to at least one antigen- or epitope-binding domain of the multispecific antibody such that coupling of the MM reduces the ability of the antigen- or epitope-binding domain to bind its target. In some embodiments, the MM is coupled to the antigen- or epitope-binding domain of the multispecific antibody via a cleavable moiety (CM) that functions as a substrate for at least one protease. The activatable multispecific antibodies provided herein are stable in circulation, activated at intended sites of therapy and/or diagnosis but not in normal, *i.e.*, healthy tissue, and, when activated, exhibit binding to a target that is at least comparable to the corresponding, unmodified multispecific antibody.

[000460] In some embodiments, the multispecific activatable antibodies are designed to engage immune effector cells, also referred to herein as immune-effector cell engaging multispecific activatable antibodies. In some embodiments, the multispecific activatable antibodies are designed to engage leukocytes, also referred to herein as leukocyte engaging multispecific activatable antibodies. In some embodiments, the multispecific activatable antibodies are designed to engage T cells, also referred to herein as T-cell engaging multispecific activatable antibodies. In some embodiments, the multispecific activatable antibodies engage a surface antigen on a leukocyte, such as on a T cell, on a natural killer (NK) cell, on a myeloid mononuclear cell, on a macrophage, and/or on another immune effector cell. In some embodiments, the immune effector cell is a leukocyte. In some embodiments, the immune effector cell is a T cell. In some embodiments, the immune effector cell is a NK cell. In some embodiments, the immune effector cell is a mononuclear cell, such as a myeloid mononuclear cell. In some embodiments, the multispecific activatable antibodies are designed to bind or otherwise interact with more than one target and/or more than one epitope, also referred to herein as multi-antigen targeting activatable antibodies. As used herein, the terms “target” and “antigen” are used interchangeably.

[000461] In some embodiments, immune effector cell engaging multispecific activatable antibodies of the disclosure include a targeting antibody or antigen-binding fragment thereof that binds CD71 and an immune effector cell engaging antibody or antigen-binding portion thereof, where at least one of the targeting antibody or antigen-binding fragment thereof and/or the immune effector cell engaging antibody or antigen-binding portion thereof is masked. In some embodiments, the immune effector cell engaging antibody or antigen binding fragment thereof includes a first antibody or antigen-binding fragment thereof (AB1) that binds a first, immune effector cell engaging target, where the AB1 is attached to a masking moiety (MM1) such that coupling of the MM1 reduces the ability of the AB1 to bind the first target. In some embodiments, the targeting antibody or antigen-binding fragment thereof includes a second antibody or fragment thereof that includes a second antibody or antigen-binding fragment thereof (AB2) that binds CD71, where the AB2 is attached to a masking moiety (MM2) such that coupling of the MM2 reduces the ability of the AB2 to bind CD71. In some embodiments, the immune effector cell engaging antibody or antigen binding fragment thereof includes a first antibody or antigen-binding fragment thereof (AB1) that binds a first, immune effector cell engaging target, where the AB1 is attached to a masking moiety (MM1) such that coupling of the MM1 reduces the ability of the AB1 to bind the first target, and the targeting antibody or antigen-binding fragment thereof includes a second antibody or fragment thereof that includes a second antibody or antigen-binding fragment thereof (AB2) that binds CD71, where the AB2 is attached to a masking moiety (MM2) such that coupling of the MM2 reduces the ability of the AB2 to bind CD71. In some embodiments, the non-immune effector cell engaging antibody is a cancer targeting antibody. In some embodiments the non-immune cell effector antibody is an IgG. In some embodiments the immune effector cell engaging antibody is a scFv. In some embodiments the CD71-targeting antibody (e.g., non-immune cell effector antibody) is an IgG and the immune effector cell engaging antibody is a scFv. In some embodiments, the immune effector cell is a leukocyte. In some embodiments, the immune effector cell is a T cell. In some embodiments, the immune effector cell is a NK cell. In some embodiments, the immune effector cell is a myeloid mononuclear cell.

[000462] In some embodiments, T-cell engaging multispecific activatable antibodies of the disclosure include a CD71-targeting antibody or antigen-binding fragment thereof and a T-cell engaging antibody or antigen-binding portion thereof, where at least one of the

CD71-targeting antibody or antigen-binding fragment thereof and/or the T-cell engaging antibody or antigen-binding portion thereof is masked. In some embodiments, the T-cell engaging antibody or antigen binding fragment thereof includes a first antibody or antigen-binding fragment thereof (AB1) that binds a first, T-cell engaging target, where the AB1 is attached to a masking moiety (MM1) such that coupling of the MM1 reduces the ability of the AB1 to bind the first target. In some embodiments, the targeting antibody or antigen-binding fragment thereof includes a second antibody or fragment thereof that includes a second antibody or antigen-binding fragment thereof (AB2) that binds CD71, where the AB2 is attached to a masking moiety (MM2) such that coupling of the MM2 reduces the ability of the AB2 to bind CD71. In some embodiments, the T-cell engaging antibody or antigen binding fragment thereof includes a first antibody or antigen-binding fragment thereof (AB1) that binds a first, T-cell engaging target, where the AB1 is attached to a masking moiety (MM1) such that coupling of the MM1 reduces the ability of the AB1 to bind the first target, and the targeting antibody or antigen-binding fragment thereof includes a second antibody or fragment thereof that includes a second antibody or antigen-binding fragment thereof (AB2) that binds CD71, where the AB2 is attached to a masking moiety (MM2) such that coupling of the MM2 reduces the ability of the AB2 to bind CD71.

[000463] In some embodiments of an immune effector cell engaging multispecific activatable antibody, one antigen is CD71, and another antigen is typically a stimulatory or inhibitory receptor present on the surface of a T-cell, natural killer (NK) cell, myeloid mononuclear cell, macrophage, and/or other immune effector cell, such as, but not limited to, B7-H4, BTLA, CD3, CD4, CD8, CD16a, CD25, CD27, CD28, CD32, CD56, CD137, CTLA-4, GITR, HVEM, ICOS, LAG3, NKG2D, OX40, PD-1, TIGIT, TIM3, or VISTA. In some embodiments, the antigen is a stimulatory receptor present on the surface of a T cell or NK cell; examples of such stimulatory receptors include, but are not limited to, CD3, CD27, CD28, CD137 (also referred to as 4-1BB), GITR, HVEM, ICOS, NKG2D, and OX40. In some embodiments, the antigen is an inhibitory receptor present on the surface of a T-cell; examples of such inhibitory receptors include, but are not limited to, BTLA, CTLA-4, LAG3, PD-1, TIGIT, TIM3, and NK-expressed KIRs. The antibody domain conferring specificity to the T-cell surface antigen may also be substituted by a ligand or ligand domain that binds to a T-cell receptor, a NK-cell receptor, a macrophage receptor, and/or other immune effector cell receptor, such as, but not limited to, B7-1, B7-2, B7H3, PDL1, PDL2, or TNFSF9.

[000464] In some embodiments, the T-cell engaging multispecific activatable antibody includes an anti-CD3 epsilon (CD3 ϵ , also referred to herein as CD3e and CD3) scFv and a targeting antibody or antigen-binding fragment thereof, where at least one of the anti-CD3 ϵ scFv and/or the targeting antibody or antigen-binding portion thereof is masked. In some embodiments, the CD3 ϵ scFv includes a first antibody or antigen-binding fragment thereof (AB1) that binds CD3 ϵ , where the AB1 is attached to a masking moiety (MM1) such that coupling of the MM1 reduces the ability of the AB1 to bind CD3 ϵ . In some embodiments, the targeting antibody or antigen-binding fragment thereof includes a second antibody or fragment thereof that includes a second antibody or antigen-binding fragment thereof (AB2) that binds CD71, where the AB2 is attached to a masking moiety (MM2) such that coupling of the MM2 reduces the ability of the AB2 to bind CD71. In some embodiments, the CD3 ϵ scFv includes a first antibody or antigen-binding fragment thereof (AB1) that binds CD3 ϵ , where the AB1 is attached to a masking moiety (MM1) such that coupling of the MM1 reduces the ability of the AB1 to bind CD3 ϵ , and the targeting antibody or antigen-binding fragment thereof includes a second antibody or fragment thereof that includes a second antibody or antigen-binding fragment thereof (AB2) that binds CD71, where the AB2 is attached to a masking moiety (MM2) such that coupling of the MM2 reduces the ability of the AB2 to bind CD71.

[000465] In some embodiments, the multi-antigen targeting antibodies and/or multi-antigen targeting activatable antibodies include at least a first antibody or antigen-binding fragment thereof that binds a first target and/or first epitope and a second antibody or antigen-binding fragment thereof that binds a second target and/or a second epitope. In some embodiments, the multi-antigen targeting antibodies and/or multi-antigen targeting activatable antibodies bind two or more different targets. In some embodiments, the multi-antigen targeting antibodies and/or multi-antigen targeting activatable antibodies bind two or more different epitopes on the same target. In some embodiments, the multi-antigen targeting antibodies and/or multi-antigen targeting activatable antibodies bind a combination of two or more different targets and two or more different epitopes on the same target.

[000466] In some embodiments, a multispecific activatable antibody comprising an IgG has the IgG variable domains masked. In some embodiments, a multispecific activatable antibody comprising a scFv has the scFv domains masked. In some embodiments, a multispecific activatable antibody has both IgG variable domains and scFv

domains, where at least one of the IgG variable domains is coupled to a masking moiety. In some embodiments, a multispecific activatable antibody has both IgG variable domains and scFv domains, where at least one of the scFv domains is coupled to a masking moiety. In some embodiments, a multispecific activatable antibody has both IgG variable domains and scFv domains, where at least one of the IgG variable domains is coupled to a masking moiety and at least one of the scFv domains is coupled to a masking moiety. In some embodiments, a multispecific activatable antibody has both IgG variable domains and scFv domains, where each of the IgG variable domains and the scFv domains is coupled to its own masking moiety. In some embodiments, one antibody domain of a multispecific activatable antibody has specificity for a target antigen and another antibody domain has specificity for a T-cell surface antigen. In some embodiments, one antibody domain of a multispecific activatable antibody has specificity for a target antigen and another antibody domain has specificity for another target antigen. In some embodiments, one antibody domain of a multispecific activatable antibody has specificity for an epitope of a target antigen and another antibody domain has specificity for another epitope of the target antigen.

[000467] In a multispecific activatable antibody, a scFv can be fused to the carboxyl terminus of the heavy chain of an IgG activatable antibody, to the carboxyl terminus of the light chain of an IgG activatable antibody, or to the carboxyl termini of both the heavy and light chains of an IgG activatable antibody. In a multispecific activatable antibody, a scFv can be fused to the amino terminus of the heavy chain of an IgG activatable antibody, to the amino terminus of the light chain of an IgG activatable antibody, or to the amino termini of both the heavy and light chains of an IgG activatable antibody. In a multispecific activatable antibody, a scFv can be fused to any combination of one or more carboxyl termini and one or more amino termini of an IgG activatable antibody. In some embodiments, a masking moiety (MM) linked to a cleavable moiety (CM) is attached to and masks an antigen binding domain of the IgG. In some embodiments, a masking moiety (MM) linked to a cleavable moiety (CM) is attached to and masks an antigen binding domain of at least one scFv. In some embodiments, a masking moiety (MM) linked to a cleavable moiety (CM) is attached to and masks an antigen binding domain of an IgG and a masking moiety (MM) linked to a cleavable moiety (CM) is attached to and masks an antigen binding domain of at least one scFv.

[000468] The disclosure provides examples of multispecific activatable antibody structures which include, but are not limited to, the following: (VL-CL)₂:(VH-CH1-CH2-CH3-L4-VH*-L3-VL*-L2-CM-L1-MM)₂; (VL-CL)₂:(VH-CH1-CH2-CH3-L4-VL*-L3-VH*-L2-CM-L1-MM)₂; (MM-L1-CM-L2-VL-CL)₂:(VH-CH1-CH2-CH3-L4-VH*-L3-VL*)₂; (MM-L1-CM-L2-VL-CL)₂:(VH-CH1-CH2-CH3-L4-VL*-L3-VH*)₂; (VL-CL)₂:(MM-L1-CM-L2-VL*-L3-VH*-L4-VH-CH1-CH2-CH3)₂; (VL-CL)₂:(MM-L1-CM-L2-VH*-L3-VL*-L4-VH-CH1-CH2-CH3)₂; (MM-L1-CM-L2-VL-CL)₂:(VL*-L3-VH*-L4-VH-CH1-CH2-CH3)₂; (MM-L1-CM-L2-VL-CL)₂:(VH*-L3-VL*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VH*-L3-VL*-L2-CM-L1-MM)₂:(VH-CH1-CH2-CH3)₂; (VL-CL-L4-VL*-L3-VH*-L2-CM-L1-MM)₂:(VH-CH1-CH2-CH3)₂; (MM-L1-CM-L2-VL*-L3-VH*-L4-VL-CL)₂:(VH-CH1-CH2-CH3)₂; (MM-L1-CM-L2-VH*-L3-VL*-L4-VL-CL)₂:(VH-CH1-CH2-CH3)₂; (VL-CL-L4-VH*-L3-VL*-L2-CM-L1-MM)₂; (MM-L1-CM-L2-VL*-L3-VH*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VH*-L3-VL*-L2-CM-L1-MM)₂; (MM-L1-CM-L2-VL*-L3-VH*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VH*-L3-VL*-L2-CM-L1-MM)₂; (MM-L1-CM-L2-VH*-L3-VL*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VL*-L3-VH*-L2-CM-L1-MM)₂; (MM-L1-CM-L2-VL*-L3-VH*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VL*-L3-VH*)₂; (MM-L1-CM-L2-VH*-L3-VL*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VH*-L3-VL*)₂; (MM-L1-CM-L2-VL*-L3-VH*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VH*-L3-VL*)₂; (MM-L1-CM-L2-VH*-L3-VL*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VL*-L3-VH*)₂; (MM-L1-CM-L2-VL*-L3-VH*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VL*-L3-VH*)₂; (MM-L1-CM-L2-VH*-L3-VL*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VH*-L3-VL*-L2-CM-L1-MM)₂; (VL*-L3-VH*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VH*-L3-VL*-L2-CM-L1-MM)₂; (VH*-L3-VL*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VL*-L3-VH*-L2-CM-L1-MM)₂; (VL*-L3-VH*-L4-VH-CH1-CH2-CH3)₂; or (VL-CL-L4-VL*-L3-VH*-L2-CM-L1-MM)₂; (VH*-L3-VL*-L4-VH-CH1-CH2-CH3)₂, wherein: VL and VH represent the light and heavy variable domains of the first specificity, contained in the IgG; VL* and VH* represent the variable domains of the second specificity, contained in the scFv; L1 is a linker peptide connecting the masking moiety (MM) and the cleavable moiety (CM); L2 is a linker peptide connecting the cleavable moiety (CM), and the antibody; L3 is a linker peptide connecting the variable domains of the scFv; L4 is a linker peptide connecting the antibody of the first specificity to the antibody of the second specificity; CL is the light-chain constant domain; and CH1, CH2, CH3 are the heavy chain constant domains. The first and second specificities can be toward any antigen or epitope.

[000469] In some embodiments of a T-cell engaging multispecific activatable antibody, one antigen is CD71, and another antigen is typically a stimulatory (also referred to herein as activating) or inhibitory receptor present on the surface of a T-cell, natural killer (NK) cell, myeloid mononuclear cell, macrophage, and/or other immune effector cell, such as, but not limited to, B7-H4, BTLA, CD3, CD4, CD8, CD16a, CD25, CD27, CD28, CD32, CD56, CD137 (also referred to as TNFRSF9), CTLA-4, GITR, HVEM, ICOS, LAG3, NKG2D, OX40, PD-1, TIGIT, TIM3, or VISTA. The antibody domain conferring specificity to the T-cell surface antigen may also be substituted by a ligand or ligand domain that binds to a T-cell receptor, a NK-cell receptor, a macrophage receptor, and/or other immune effector cell receptor.

[000470] In some embodiments, the targeting antibody is an anti-CD71 antibody disclosed herein. In some embodiments, the targeting antibody can be in the form an activatable antibody. In some embodiments, the scFv(s) can be in the form of a Pro-scFv (see, e.g., WO 2009/025846, WO 2010/081173).

[000471] In some embodiments, the scFv is specific for binding CD3 ϵ , and comprises or is derived from an antibody or fragment thereof that binds CD3 ϵ , e.g., CH2527, FN18, H2C, OKT3, 2C11, UCHT1, or V9. In some embodiments, the scFv is specific for binding CTLA-4 (also referred to herein as CTLA and CTLA4).

[000472] In some embodiments, the anti-CTLA-4 scFv includes the amino acid sequence:

```
GGGSGGGSGSGGGSGGGSGGGEIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQ
QKPGQAPRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYYCQQYGSSPLTF
GGGTKVEIKRSGGSTITSYNVYYTKLSSSGTQVQLVQTGGGVVQPGRSLRLSCAASGSTFS
SYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDT
AVYYCATNSLYWYFDLWGRGTLTVSSAS (SEQ ID NO: 643)
```

[000473] In some embodiments, the anti-CTLA-4 scFv includes the amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence of SEQ ID NO: 643.

[000474] In some embodiments, the anti-CD3 ϵ scFv includes the amino acid sequence:

```
GGGSGGGSGSGGGSGGGSGGGQVQLQQSGAELARPGASVKMSCKASGYTFTRYTMHWVK
QRPQGQLEWIGYINPSRGYTNYNQKFKDKATLTDDKSSSTAYMQLSSLTSEDSAVYYCARY
YDDHYCLDYWGQGTTLTVSSGGGGSGGGSGGGGSQIVLTQSPAIMSASPGKVTMTCSAS
```

SSVSYMNWYQQKSGTSPKRWIYDTSKLGVPAPFRGSGSGTSYSLTISGMEAEDAATYYC
QQWSSNPFTFGSGTKLEINR (SEQ ID NO: 644)

[000475] In some embodiments, the anti-CD3 ϵ scFv includes the amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence of SEQ ID NO: 644.

[000476] In some embodiments, the scFv is specific for binding one or more T-cells, one or more NK-cells and/or one or more macrophages. In some embodiments, the scFv is specific for binding a target selected from the group consisting of B7-H4, BTLA, CD3, CD4, CD8, CD16a, CD25, CD27, CD28, CD32, CD56, CD137, CTLA-4, GITR, HVEM, ICOS, LAG3, NKG2D, OX40, PD-1, TIGIT, TIM3, or VISTA.

[000477] In some embodiments, the multispecific activatable antibody also includes an agent conjugated to the AB. In some embodiments, the agent is a therapeutic agent. In some embodiments, the agent is an antineoplastic agent. In some embodiments, the agent is a toxin or fragment thereof. In some embodiments, the agent is conjugated to the multispecific activatable antibody via a linker. In some embodiments, the agent is conjugated to the AB via a cleavable linker. In some embodiments, the linker is a non-cleavable linker. In some embodiments, the agent is a microtubule inhibitor. In some embodiments, the agent is a nucleic acid damaging agent, such as a DNA alkylator or DNA intercalator, or other DNA damaging agent. In some embodiments, the linker is a cleavable linker. In some embodiments, the agent is an agent selected from the group listed in Table 5. In some embodiments, the agent is a dolastatin. In some embodiments, the agent is an auristatin or derivative thereof. In some embodiments, the agent is auristatin E or a derivative thereof. In some embodiments, the agent is monomethyl auristatin E (MMAE). In some embodiments, the agent is monomethyl auristatin D (MMAD). In some embodiments, the agent is a maytansinoid or maytansinoid derivative. In some embodiments, the agent is DM1 or DM4. In some embodiments, the agent is a duocarmycin or derivative thereof. In some embodiments, the agent is a calicheamicin or derivative thereof. In some embodiments, the agent is a pyrrolobenzodiazepine. In some embodiments, the agent is a pyrrolobenzodiazepine dimer.

[000478] In some embodiments, the multispecific activatable antibody also includes a detectable moiety. In some embodiments, the detectable moiety is a diagnostic agent.

[000479] In some embodiments, the multispecific activatable antibody naturally contains one or more disulfide bonds. In some embodiments, the multispecific activatable antibody can be engineered to include one or more disulfide bonds.

[000480] The disclosure also provides an isolated nucleic acid molecule encoding a multispecific activatable antibody described herein, as well as vectors that include these isolated nucleic acid sequences. The disclosure provides methods of producing a multispecific activatable antibody by culturing a cell under conditions that lead to expression of the activatable antibody, wherein the cell comprises such a nucleic acid molecule. In some embodiments, the cell comprises such a vector.

[000481] The disclosure also provides a method of manufacturing multispecific activatable antibodies of the disclosure by (a) culturing a cell comprising a nucleic acid construct that encodes the multispecific activatable antibody under conditions that lead to expression of the multispecific activatable, and (b) recovering the multispecific activatable antibody. Suitable AB, MM, and/or CM include any of the AB, MM, and/or CM disclosed herein.

[000482] The disclosure also provides multispecific activatable antibodies and/or multispecific activatable antibody compositions that include at least a first antibody or antigen-binding fragment thereof (AB1) that specifically binds a first target or first epitope and a second antibody or antigen-binding fragment thereof (AB2) that binds a second target or a second epitope, where at least AB1 is coupled or otherwise attached to a masking moiety (MM1), such that coupling of the MM1 reduces the ability of AB1 to bind its target. In some embodiments, the MM1 is coupled to AB1 via a first cleavable moiety (CM1) sequence that includes a substrate for a protease, for example, a protease that is co-localized with the target of AB1 at a treatment site or a diagnostic site in a subject. The multispecific activatable antibodies provided herein are stable in circulation, activated at intended sites of therapy and/or diagnosis but not in normal, *i.e.*, healthy tissue, and, when activated, exhibit binding to the target of AB1 that is at least comparable to the corresponding, unmodified multispecific antibody. Suitable AB, MM, and/or CM include any of the AB, MM, and/or CM disclosed herein.

[000483] The disclosure also provides compositions and methods that include a multispecific activatable antibody that includes at least a first antibody or antibody fragment (AB1) that specifically binds a target and a second antibody or antibody fragment (AB2), where at least the first AB in the multispecific activatable antibody is coupled to a masking

moiety (MM1) that decreases the ability of AB1 to bind its target. In some embodiments, each AB is coupled to a MM that decreases the ability of its corresponding AB to each target. For example, in bispecific activatable antibody embodiments, AB1 is coupled to a first masking moiety (MM1) that decreases the ability of AB1 to bind its target, and AB2 is coupled to a second masking moiety (MM2) that decreases the ability of AB2 to bind its target. In some embodiments, the multispecific activatable antibody comprises more than two AB regions; in such embodiments, AB1 is coupled to a first masking moiety (MM1) that decreases the ability of AB1 to bind its target, AB2 is coupled to a second masking moiety (MM2) that decreases the ability of AB2 to bind its target, AB3 is coupled to a third masking moiety (MM3) that decreases the ability of AB3 to bind its target, and so on for each AB in the multispecific activatable antibody. Suitable AB, MM, and/or CM include any of the AB, MM, and/or CM disclosed herein.

[000484] In some embodiments, the multispecific activatable antibody further includes at least one cleavable moiety (CM) that is a substrate for a protease, where the CM links a MM to an AB. For example, in some embodiments, the multispecific activatable antibody includes at least a first antibody or antibody fragment (AB1) that specifically binds a target and a second antibody or antibody fragment (AB2), where at least the first AB in the multispecific activatable antibody is coupled via a first cleavable moiety (CM1) to a masking moiety (MM1) that decreases the ability of AB1 to bind its target. In some bispecific activatable antibody embodiments, AB1 is coupled via CM1 to MM1, and AB2 is coupled via a second cleavable moiety (CM2) to a second masking moiety (MM2) that decreases the ability of AB2 to bind its target. In some embodiments, the multispecific activatable antibody comprises more than two AB regions; in some of these embodiments, AB1 is coupled via CM1 to MM1, AB2 is coupled via CM2 to MM2, and AB3 is coupled via a third cleavable moiety (CM3) to a third masking moiety (MM3) that decreases the ability of AB3 to bind its target, and so on for each AB in the multispecific activatable antibody. Suitable AB, MM, and/or CM include any of the AB, MM, and/or CM disclosed herein.

Activatable antibodies Having Non-Binding Steric Moieties or Binding Partners for Non-Binding Steric Moieties

[000485] The disclosure also provides activatable antibodies that include non-binding steric moieties (NB) or binding partners (BP) for non-binding steric moieties, where the BP recruits or otherwise attracts the NB to the activatable antibody. The activatable antibodies

provided herein include, for example, an activatable antibody that includes a non-binding steric moiety (NB), a cleavable linker (CL) and antibody or antibody fragment (AB) that binds a target; an activatable antibody that includes a binding partner for a non-binding steric moiety (BP), a CL and an AB; and an activatable antibody that includes a BP to which an NB has been recruited, a CL and an AB that binds the target. Activatable antibodies in which the NB is covalently linked to the CL and AB of the activatable antibody or is associated by interaction with a BP that is covalently linked to the CL and AB of the activatable antibody are referred to herein as “NB-containing activatable antibodies.” By activatable or switchable is meant that the activatable antibody exhibits a first level of binding to a target when the activatable antibody is in an inhibited, masked or uncleaved state (*i.e.*, a first conformation), and a second level of binding to the target when the activatable antibody is in an uninhibited, unmasked and/or cleaved state (*i.e.*, a second conformation, *i.e.*, activated antibody), where the second level of target binding is greater than the first level of target binding. The activatable antibody compositions can exhibit increased bioavailability and more favorable biodistribution compared to conventional antibody therapeutics.

[000486] In some embodiments, activatable antibodies provide for reduced toxicity and/or adverse side effects that could otherwise result from binding of the at non-treatment sites and/or non-diagnostic sites if the AB were not masked or otherwise inhibited from binding to such a site.

[000487] Anti-CD71 activatable antibodies that include a non-binding steric moiety (NB) can be made using the methods set forth in PCT Publication No. WO 2013/192546, the contents of which are hereby incorporated by reference in their entirety.

Use Of Antibodies, Conjugated Antibodies, Activatable Antibodies, and Conjugated Activatable Antibodies

[000488] It will be appreciated that administration of therapeutic entities in accordance with the disclosure will be administered with suitable carriers, excipients, and other agents that are incorporated into formulations to provide improved transfer, delivery, tolerance, and the like. A multitude of appropriate formulations can be found in the formulary known to all pharmaceutical chemists: Remington’s Pharmaceutical Sciences (15th ed, Mack Publishing Company, Easton, PA (1975)), particularly Chapter 87 by Blaug, Seymour, therein. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles (such as Lipofectin™), DNA

conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax. Any of the foregoing mixtures may be appropriate in treatments and therapies in accordance with the present disclosure, provided that the active ingredient in the formulation is not inactivated by the formulation and the formulation is physiologically compatible and tolerable with the route of administration. *See also* Baldrick P. "Pharmaceutical excipient development: the need for preclinical guidance." Regul. Toxicol Pharmacol. 32(2):210-8 (2000), Wang W. "Lyophilization and development of solid protein pharmaceuticals." Int. J. Pharm. 203(1-2):1-60 (2000), Charman WN "Lipids, lipophilic drugs, and oral drug delivery-some emerging concepts." J Pharm Sci. 89(8):967-78 (2000), Powell *et al.* "Compendium of excipients for parenteral formulations" PDA J Pharm Sci Technol. 52:238-311 (1998) and the citations therein for additional information related to formulations, excipients and carriers well known to pharmaceutical chemists.

[000489] Therapeutic formulations of the disclosure, which include an anti-CD71 antibody and/or activatable anti-CD71 antibody, such as by way of non-limiting example, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody, are used to prevent, treat or otherwise ameliorate a disease or disorder associated with aberrant target expression and/or activity. For example, therapeutic formulations of the disclosure, which include an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody, are used to treat or otherwise ameliorate a cancer or other neoplastic condition, inflammation, an inflammatory disorder, and/or an autoimmune disease. In some embodiments, the cancer is a solid tumor or a hematologic malignancy where the target is expressed. In some embodiments, the cancer is a solid tumor where the target is expressed. In some embodiments, the cancer is a hematologic malignancy where the target is expressed. In some embodiments, the target is expressed on parenchyma (e.g., in cancer, the portion of an organ or tissue that often carries out function(s) of the organ or tissue). In some embodiments, the target is expressed on a cell, tissue, or organ. In some embodiments, the target is expressed on stroma (i.e., the connective supportive framework of a cell, tissue, or organ). In some embodiments, the target is expressed on an osteoblast. In some embodiments, the target is expressed on the endothelium (vasculature). In some embodiments, the target is expressed on a cancer stem cell. In some embodiments, the agent to which the antibody and/or the activatable antibody

is conjugated is a microtubule inhibitor. In some embodiments, the agent to which the antibody and/or the activatable antibody is conjugated is a nucleic acid damaging agent.

[000490] Efficaciousness of prevention, amelioration or treatment is determined in association with any known method for diagnosing or treating the disease or disorder associated with target expression and/or activity, such as, for example, aberrant target expression and/or activity. Prolonging the survival of a subject or otherwise delaying the progression of the disease or disorder associated with target expression and/or activity, e.g., aberrant target expression and/or activity, in a subject indicates that the antibody, conjugated antibody, activatable antibody and/or conjugated activatable antibody confers a clinical benefit.

[000491] An antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody can be administered in the form of pharmaceutical compositions. Principles and considerations involved in preparing such compositions, as well as guidance in the choice of components are provided, for example, in Remington : The Science And Practice Of Pharmacy 19th ed. (Alfonso R. Gennaro, et al., editors) Mack Pub. Co., Easton, Pa.: 1995; Drug Absorption Enhancement: Concepts, Possibilities, Limitations, And Trends, Harwood Academic Publishers, Langhorne, Pa., 1994; and Peptide And Protein Drug Delivery (Advances In Parenteral Sciences, Vol. 4), 1991, M. Dekker, New York.

[000492] In some embodiments where antibody fragments are used, the smallest fragment that specifically binds to the binding domain of the target protein is selected. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. (*See, e.g.,* Marasco et al., Proc. Natl. Acad. Sci. USA, 90: 7889-7893 (1993)). The formulation can also contain more than one active compounds as necessary for the particular indication being treated, for example, in some embodiments, those with complementary activities that do not adversely affect each other. In some embodiments, or in addition, the composition can comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[000493] The active ingredients can also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example,

hydroxymethylcellulose or gelatin-microcapsules and poly-(methylemethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions.

[000494] The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

[000495] Sustained-release preparations can be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOTTM (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods.

[000496] In some embodiments, the antibody, the conjugated antibody, activatable antibody and/or conjugated activatable antibody contains a detectable label. An intact antibody, or a fragment thereof (*e.g.*, Fab, scFv, or F(ab)₂) is used. The term “labeled”, with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently-labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently-labeled streptavidin. The term “biological sample” is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. Included within the usage of the term “biological sample”, therefore, is blood and a fraction or component of blood including blood serum, blood plasma, or lymph. That is, the detection method of the disclosure can be used to detect an analyte mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of an analyte mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro*

techniques for detection of an analyte protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations, immunochemical staining, and immunofluorescence. *In vitro* techniques for detection of an analyte genomic DNA include Southern hybridizations. Procedures for conducting immunoassays are described, for example in “ELISA: Theory and Practice: Methods in Molecular Biology”, Vol. 42, J. R. Crowther (Ed.) Human Press, Totowa, NJ, 1995; “Immunoassay”, E. Diamandis and T. Christopoulos, Academic Press, Inc., San Diego, CA, 1996; and “Practice and Theory of Enzyme Immunoassays”, P. Tijssen, Elsevier Science Publishers, Amsterdam, 1985. Furthermore, *in vivo* techniques for detection of an analyte protein include introducing into a subject a labeled anti-analyte protein antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

[000497] The antibodies, conjugated antibodies, activatable antibodies and/or conjugated activatable antibodies of the disclosure are also useful in a variety of diagnostic and prophylactic formulations. In one embodiment, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody is administered to patients that are at risk of developing one or more of the aforementioned disorders. A patient's or organ's predisposition to one or more of the aforementioned disorders can be determined using genotypic, serological or biochemical markers.

[000498] In some embodiments of the disclosure, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody is administered to human individuals diagnosed with a clinical indication associated with one or more of the aforementioned disorders. Upon diagnosis, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody is administered to mitigate or reverse the effects of the clinical indication.

[000499] An antibody, a conjugated antibody, an activatable antibody, and/or a conjugated activatable antibody of the disclosure is also useful in the detection of a target in patient samples and accordingly are useful as diagnostics. For example, the antibodies and/or activatable antibodies, and conjugated versions thereof, of the disclosure are used in *in vitro* assays, *e.g.*, ELISA, to detect target levels in a patient sample.

[000500] In one embodiment, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody of the disclosure is immobilized on a solid support (*e.g.*, the well(s) of a microtiter plate). The immobilized antibody, conjugated

antibody, activatable antibody and/or conjugated activatable antibody serves as a capture antibody for any target that may be present in a test sample. Prior to contacting the immobilized antibody and/or activatable antibody, and/or conjugated versions thereof, with a patient sample, the solid support is rinsed and treated with a blocking agent such as milk protein or albumin to prevent nonspecific adsorption of the analyte.

[000501] Subsequently the wells are treated with a test sample suspected of containing the antigen, or with a solution containing a standard amount of the antigen. Such a sample is, *e.g.*, a serum sample from a subject suspected of having levels of circulating antigen considered to be diagnostic of a pathology. After rinsing away the test sample or standard, the solid support is treated with a second antibody that is detectably labeled. The labeled second antibody serves as a detecting antibody. The level of detectable label is measured, and the concentration of target antigen in the test sample is determined by comparison with a standard curve developed from the standard samples.

[000502] It will be appreciated that based on the results obtained using the antibodies and activatable antibodies of the disclosure, and conjugated versions thereof, in an *in vitro* diagnostic assay, it is possible to stage a disease in a subject based on expression levels of the target antigen. For a given disease, samples of blood are taken from subjects diagnosed as being at various stages in the progression of the disease, and/or at various points in the therapeutic treatment of the disease. Using a population of samples that provides statistically significant results for each stage of progression or therapy, a range of concentrations of the antigen that may be considered characteristic of each stage is designated.

[000503] An antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody can also be used in diagnostic and/or imaging methods. In some embodiments, such methods are *in vitro* methods. In some embodiments, such methods are *in vivo* methods. In some embodiments, such methods are *in situ* methods. In some embodiments, such methods are *ex vivo* methods. For example, activatable antibodies having an enzymatically cleavable CM can be used to detect the presence or absence of an enzyme that is capable of cleaving the CM. Such activatable antibodies can be used in diagnostics, which can include *in vivo* detection (*e.g.*, qualitative or quantitative) of enzyme activity (or, in some embodiments, an environment of increased reduction potential such as that which can provide for reduction of a disulfide bond) through measured accumulation of activated antibodies (*i.e.*, antibodies resulting from cleavage of an activatable antibody) in a

given cell or tissue of a given host organism. Such accumulation of activated antibodies indicates not only that the tissue expresses enzymatic activity (or an increased reduction potential depending on the nature of the CM) but also that the tissue expresses target to which the activated antibody binds.

[000504] For example, the CM can be selected to be substrate for at least one protease found at the site of a tumor, at the site of a viral or bacterial infection at a biologically confined site (*e.g.*, such as in an abscess, in an organ, and the like), and the like. The AB can be one that binds a target antigen. Using methods as disclosed herein, or when appropriate, methods familiar to one skilled in the art, a detectable label (*e.g.*, a fluorescent label or radioactive label or radiotracer) can be conjugated to an AB or other region of an antibody and/or activatable antibody. Suitable detectable labels are discussed in the context of the above screening methods and additional specific examples are provided below. Using an AB specific to a protein or peptide of the disease state, along with at least one protease whose activity is elevated in the disease tissue of interest, activatable antibodies will exhibit an increased rate of binding to disease tissue relative to tissues where the CM specific enzyme is not present at a detectable level or is present at a lower level than in disease tissue or is inactive (*e.g.*, in zymogen form or in complex with an inhibitor). Since small proteins and peptides are rapidly cleared from the blood by the renal filtration system, and because the enzyme specific for the CM is not present at a detectable level (or is present at lower levels in non-disease tissues or is present in inactive conformation), accumulation of activated antibodies in the disease tissue is enhanced relative to non-disease tissues.

[000505] In another example, activatable antibodies can be used to detect the presence or absence of a cleaving agent in a sample. For example, where the activatable antibodies contain a CM susceptible to cleavage by an enzyme, the activatable antibodies can be used to detect (either qualitatively or quantitatively) the presence of an enzyme in the sample. In another example, where the activatable antibodies contain a CM susceptible to cleavage by reducing agent, the activatable antibodies can be used to detect (either qualitatively or quantitatively) the presence of reducing conditions in a sample. To facilitate analysis in these methods, the activatable antibodies can be detectably labeled, and can be bound to a support (*e.g.*, a solid support, such as a slide or bead). The detectable label can be positioned on a portion of the activatable antibody that is not released following cleavage, for example, the detectable label can be a quenched fluorescent label or other label that is not detectable until cleavage has occurred. The assay can be conducted by, for example, contacting the

immobilized, detectably labeled activatable antibodies with a sample suspected of containing an enzyme and/or reducing agent for a time sufficient for cleavage to occur, then washing to remove excess sample and contaminants. The presence or absence of the cleaving agent (*e.g.*, enzyme or reducing agent) in the sample is then assessed by a change in detectable signal of the activatable antibodies prior to contacting with the sample *e.g.*, the presence of and/or an increase in detectable signal due to cleavage of the activatable antibody by the cleaving agent in the sample.

[000506] Such detection methods can be adapted to also provide for detection of the presence or absence of a target that is capable of binding the AB of the activatable antibodies when cleaved. Thus, the assays can be adapted to assess the presence or absence of a cleaving agent and the presence or absence of a target of interest. The presence or absence of the cleaving agent can be detected by the presence of and/or an increase in detectable label of the activatable antibodies as described above, and the presence or absence of the target can be detected by detection of a target-AB complex *e.g.*, by use of a detectably labeled anti-target antibody.

[000507] Activatable antibodies are also useful in *in situ* imaging for the validation of activatable antibody activation, *e.g.*, by protease cleavage, and binding to a particular target. *In situ* imaging is a technique that enables localization of proteolytic activity and target in biological samples such as cell cultures or tissue sections. Using this technique, it is possible to confirm both binding to a given target and proteolytic activity based on the presence of a detectable label (*e.g.*, a fluorescent label).

[000508] These techniques are useful with any frozen cells or tissue derived from a disease site (*e.g.* tumor tissue) or healthy tissues. These techniques are also useful with fresh cell or tissue samples.

[000509] In these techniques, an activatable antibody is labeled with a detectable label. The detectable label can be a fluorescent dye, (*e.g.* a fluorophore, Fluorescein Isothiocyanate (FITC), Rhodamine Isothiocyanate (TRITC), an Alexa Fluor® label), a near infrared (NIR) dye (*e.g.*, Qdot® nanocrystals), a colloidal metal, a hapten, a radioactive marker, biotin and an amplification reagent such as streptavidin, or an enzyme (*e.g.* horseradish peroxidase or alkaline phosphatase).

[000510] Detection of the label in a sample that has been incubated with the labeled, activatable antibody indicates that the sample contains the target and contains a protease that is specific for the CM of the activatable antibody. In some embodiments, the presence

of the protease can be confirmed using broad spectrum protease inhibitors such as those described herein, and/or by using an agent that is specific for the protease, for example, an antibody such as A11, which is specific for the protease matriptase and inhibits the proteolytic activity of matriptase; see e.g., International Publication Number WO 2010/129609, published 11 November 2010. The same approach of using broad spectrum protease inhibitors such as those described herein, and/or by using a more selective inhibitory agent can be used to identify a protease that is specific for the CM of the activatable antibody. In some embodiments, the presence of the target can be confirmed using an agent that is specific for the target, e.g., another antibody, or the detectable label can be competed with unlabeled target. In some embodiments, unlabeled activatable antibody could be used, with detection by a labeled secondary antibody or more complex detection system.

[000511] Similar techniques are also useful for *in vivo* imaging where detection of the fluorescent signal in a subject, e.g., a mammal, including a human, indicates that the disease site contains the target and contains a protease that is specific for the CM of the activatable antibody.

[000512] These techniques are also useful in kits and/or as reagents for the detection, identification or characterization of protease activity in a variety of cells, tissues, and organisms based on the protease-specific CM in the activatable antibody.

[000513] The disclosure provides methods of using the antibodies and/or activatable antibodies in a variety of diagnostic and/or prophylactic indications. For example, the disclosure provides methods of detecting presence or absence of a cleaving agent and a target of interest in a subject or a sample by (i) contacting a subject or sample with an activatable antibody, wherein the activatable antibody comprises a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, e.g., a protease, and an antigen binding domain or fragment thereof (AB) that specifically binds the target of interest, wherein the activatable antibody in an uncleaved, non-activated state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB and is not a modified form of a natural binding partner of the AB; and (b) wherein, in an uncleaved, non-activated state, the MM interferes with specific binding of the AB to the target, and in a cleaved, activated state the MM does not interfere or compete with specific

binding of the AB to the target; and (ii) measuring a level of activated activatable antibody in the subject or sample, wherein a detectable level of activated activatable antibody in the subject or sample indicates that the cleaving agent and the target are present in the subject or sample and wherein no detectable level of activated activatable antibody in the subject or sample indicates that the cleaving agent, the target or both the cleaving agent and the target are absent and/or not sufficiently present in the subject or sample. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000514] The disclosure also provides methods of detecting presence or absence of a cleaving agent in a subject or a sample by (i) contacting a subject or sample with an activatable antibody in the presence of a target of interest, *e.g.*, the target, wherein the activatable antibody comprises a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, *e.g.*, a protease, and an antigen binding domain or fragment thereof (AB) that specifically binds the target of interest, wherein the activatable antibody in an uncleaved, non-activated state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB and is not a modified form of a natural binding partner of the AB; and (b) wherein, in an uncleaved, non-activated state, the MM interferes with specific binding of the AB to the target, and in a cleaved, activated state the MM does not interfere or compete with specific binding of the AB to the target; and (ii) measuring a level of activated activatable antibody in the subject or sample, wherein a detectable level of activated activatable antibody in the subject or sample indicates that the cleaving agent is present in the subject or sample and wherein no detectable level of activated activatable antibody in the subject or sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or sample. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In

some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000515] The disclosure also provides kits for use in methods of detecting presence or absence of a cleaving agent and the target in a subject or a sample, where the kits include at least an activatable antibody comprises a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, e.g., a protease, and an antigen binding domain or fragment thereof (AB) that specifically binds the target of interest, wherein the activatable antibody in an uncleaved, non-activated state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB and is not a modified form of a natural binding partner of the AB; and (b) wherein, in an uncleaved, non-activated state, the MM interferes with specific binding of the AB to the target, and in a cleaved, activated state the MM does not interfere or compete with specific binding of the AB to the target; and (ii) measuring a level of activated activatable antibody in the subject or sample, wherein a detectable level of activated activatable antibody in the subject or sample indicates that the cleaving agent is present in the subject or sample and wherein no detectable level of activated activatable antibody in the subject or sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or sample. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000516] The disclosure also provides methods of detecting presence or absence of a cleaving agent in a subject or a sample by (i) contacting a subject or sample with an activatable antibody, wherein the activatable antibody comprises a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, e.g., a protease, an antigen binding domain (AB) that specifically binds the target, and a detectable label, wherein the activatable antibody in an uncleaved, non-activated state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB and is not a modified form of a natural binding partner of the AB; wherein, in an uncleaved, non-activated state, the MM interferes with specific binding of the AB to the target, and in a cleaved, activated state the MM does not interfere or compete with specific binding of the AB to the target; and wherein the detectable label is positioned on a portion of the activatable antibody that is released following cleavage of the CM; and (ii) measuring a level of detectable label in the subject or sample, wherein a detectable level of the detectable label in the subject or sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or sample and wherein no detectable level of the detectable label in the subject or sample indicates that the cleaving agent is present in the subject or sample. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000517] The disclosure also provides kits for use in methods of detecting presence or absence of a cleaving agent and the target in a subject or a sample, where the kits include at least an activatable antibody and/or conjugated activatable antibody (e.g., an activatable antibody to which a therapeutic agent is conjugated) described herein for use in contacting a subject or biological sample and means for detecting the level of activated activatable antibody and/or conjugated activatable antibody in the subject or biological sample, wherein a detectable level of activated activatable antibody in the subject or biological sample

indicates that the cleaving agent and the target are present in the subject or biological sample and wherein no detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent, the target or both the cleaving agent and the target are absent and/or not sufficiently present in the subject or biological sample, such that the target binding and/or protease cleavage of the activatable antibody cannot be detected in the subject or biological sample.

[000518] The disclosure also provides methods of detecting presence or absence of a cleaving agent in a subject or a sample by (i) contacting a subject or biological sample with an activatable antibody in the presence of the target, and (ii) measuring a level of activated activatable antibody in the subject or biological sample, wherein a detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent is present in the subject or biological sample and wherein no detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or biological sample at a detectable level, such that protease cleavage of the activatable antibody cannot be detected in the subject or biological sample. Such an activatable antibody includes a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, e.g., a protease, and an antigen binding domain or fragment thereof (AB) that specifically binds the target, wherein the activatable antibody in an uncleaved (*i.e.*, non-activated) state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB; and (b) wherein the MM of the activatable antibody in an uncleaved state interferes with specific binding of the AB to the target, and wherein the MM of an activatable antibody in a cleaved (*i.e.*, activated) state does not interfere or compete with specific binding of the AB to the target. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the detectable label is attached to the masking moiety. In some embodiments, the detectable label is attached to the cleavable moiety N-terminal to the protease cleavage site. In some embodiments, a single antigen binding site of the AB is masked. In some embodiments wherein an antibody of the disclosure has at least two antigen binding sites, at least one antigen binding site is masked and at least one antigen binding site is not masked. In some

embodiments all antigen binding sites are masked. In some embodiments, the measuring step includes use of a secondary reagent comprising a detectable label.

[000519] The disclosure also provides kits for use in methods of detecting presence or absence of a cleaving agent and the target in a subject or a sample, where the kits include at least an activatable antibody and/or conjugated activatable antibody described herein for use in contacting a subject or biological sample with an activatable antibody in the presence of the target, and measuring a level of activated activatable antibody in the subject or biological sample, wherein a detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent is present in the subject or biological sample and wherein no detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or biological sample at a detectable level, such that protease cleavage of the activatable antibody cannot be detected in the subject or biological sample. Such an activatable antibody includes a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, e.g., a protease, and an antigen binding domain or fragment thereof (AB) that specifically binds the target, wherein the activatable antibody in an uncleaved (*i.e.*, non-activated) state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB; and (b) wherein the MM of the activatable antibody in an uncleaved state interferes with specific binding of the AB to the target, and wherein the MM of an activatable antibody in a cleaved (*i.e.*, activated) state does not interfere or compete with specific binding of the AB to the target. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the detectable label is attached to the masking moiety. In some embodiments, the detectable label is attached to the cleavable moiety N-terminal to the protease cleavage site. In some embodiments, a single antigen binding site of the AB is masked. In some embodiments wherein an antibody of the disclosure has at least two antigen binding sites, at least one antigen binding site is masked and at least one antigen binding site is not masked. In some embodiments all antigen binding sites are masked. In some embodiments, the measuring step includes use of a secondary reagent comprising a detectable label.

[000520] The disclosure also provides kits for use in methods of detecting presence or absence of a cleaving agent in a subject or a sample, where the kits include at least an activatable antibody and/or conjugated activatable antibody described herein for use in contacting a subject or biological sample and means for detecting the level of activated activatable antibody and/or conjugated activatable antibody in the subject or biological sample, wherein the activatable antibody includes a detectable label that is positioned on a portion of the activatable antibody that is released following cleavage of the CM, wherein a detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or biological sample such that the target binding and/or protease cleavage of the activatable antibody cannot be detected in the subject or biological sample, and wherein no detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent is present in the subject or biological sample at a detectable level.

[000521] The disclosure provides methods of detecting presence or absence of a cleaving agent and the target in a subject or a sample by (i) contacting a subject or biological sample with an activatable antibody, wherein the activatable antibody includes a detectable label that is positioned on a portion of the activatable antibody that is released following cleavage of the CM and (ii) measuring a level of activated activatable antibody in the subject or biological sample, wherein a detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent, the target or both the cleaving agent and the target are absent and/or not sufficiently present in the subject or biological sample, such that the target binding and/or protease cleavage of the activatable antibody cannot be detected in the subject or biological sample, and wherein a reduced detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent and the target are present in the subject or biological sample. A reduced level of detectable label is, for example, a reduction of about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95% and/or about 100%. Such an activatable antibody includes a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, and an antigen binding domain or fragment thereof (AB) that specifically binds the target, wherein the activatable antibody in an uncleaved (*i.e.*, non-activated) state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-

MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB; and (b) wherein the MM of the activatable antibody in an uncleaved state interferes with specific binding of the AB to the target, and wherein the MM of an activatable antibody in a cleaved (*i.e.*, activated) state does not interfere or compete with specific binding of the AB to the target. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000522] The disclosure also provides kits for use in methods of detecting presence or absence of a cleaving agent and the target in a subject or a sample, where the kits include at least an activatable antibody and/or conjugated activatable antibody described herein for use in contacting a subject or biological sample and means for detecting the level of activated activatable antibody and/or conjugated activatable antibody in the subject or biological sample, wherein a detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent, the target or both the cleaving agent and the target are absent and/or not sufficiently present in the subject or biological sample, such that the target binding and/or protease cleavage of the activatable antibody cannot be detected in the subject or biological sample, and wherein a reduced detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent and the target are present in the subject or biological sample. A reduced level of detectable label is, for example, a reduction of about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95% and/or about 100%.

[000523] The disclosure also provides methods of detecting presence or absence of a cleaving agent in a subject or a sample by (i) contacting a subject or biological sample with an activatable antibody, wherein the activatable antibody includes a detectable label that is positioned on a portion of the activatable antibody that is released following cleavage of the

CM; and (ii) measuring a level of detectable label in the subject or biological sample, wherein a detectable level of the detectable label in the subject or biological sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or biological sample at a detectable level, such that protease cleavage of the activatable antibody cannot be detected in the subject or biological sample, and wherein a reduced detectable level of the detectable label in the subject or biological sample indicates that the cleaving agent is present in the subject or biological sample. A reduced level of detectable label is, for example, a reduction of about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95% and/or about 100%. Such an activatable antibody includes a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, and an antigen binding domain or fragment thereof (AB) that specifically binds the target, wherein the activatable antibody in an uncleaved (*i.e.*, non-activated) state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB; and (b) wherein the MM of the activatable antibody in an uncleaved state interferes with specific binding of the AB to the target, and wherein the MM of an activatable antibody in a cleaved (*i.e.*, activated) state does not interfere or compete with specific binding of the AB to the target. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000524] The disclosure also provides kits for use in methods of detecting presence or absence of a cleaving agent of interest in a subject or a sample, where the kits include at least an activatable antibody and/or conjugated activatable antibody described herein for use in contacting a subject or biological sample and means for detecting the level of activated activatable antibody and/or conjugated activatable antibody in the subject or biological

sample, wherein the activatable antibody includes a detectable label that is positioned on a portion of the activatable antibody that is released following cleavage of the CM, wherein a detectable level of the detectable label in the subject or biological sample indicates that the cleaving agent, the target, or both the cleaving agent and the target are absent and/or not sufficiently present in the subject or biological sample, such that the target binding and/or protease cleavage of the activatable antibody cannot be detected in the subject or biological sample, and wherein a reduced detectable level of the detectable label in the subject or biological sample indicates that the cleaving agent and the target are present in the subject or biological sample. A reduced level of detectable label is, for example, a reduction of about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95% and/or about 100%.

[000525] In some embodiments of these methods and kits, the activatable antibody includes a detectable label. In some embodiments of these methods and kits, the detectable label includes an imaging agent, a contrasting agent, an enzyme, a fluorescent label, a chromophore, a dye, one or more metal ions, or a ligand-based label. In some embodiments of these methods and kits, the imaging agent comprises a radioisotope. In some embodiments of these methods and kits, the radioisotope is indium or technetium. In some embodiments of these methods and kits, the contrasting agent comprises iodine, gadolinium or iron oxide. In some embodiments of these methods and kits, the enzyme comprises horseradish peroxidase, alkaline phosphatase, or β -galactosidase. In some embodiments of these methods and kits, the fluorescent label comprises yellow fluorescent protein (YFP), cyan fluorescent protein (CFP), green fluorescent protein (GFP), modified red fluorescent protein (mRFP), red fluorescent protein tdimer2 (RFP tdimer2), HCRED, or a europium derivative. In some embodiments of these methods and kits, the luminescent label comprises an N-methylacrydium derivative. In some embodiments of these methods, the label comprises an Alexa Fluor[®] label, such as Alex Fluor[®] 680 or Alexa Fluor[®] 750. In some embodiments of these methods and kits, the ligand-based label comprises biotin, avidin, streptavidin or one or more haptens.

[000526] In some embodiments of these methods and kits, the subject is a mammal. In some embodiments of these methods and kits, the subject is a human. In some embodiments, the subject is a non-human mammal, such as a non-human primate,

companion animal (e.g., cat, dog, horse), farm animal, work animal, or zoo animal. In some embodiments, the subject is a rodent.

[000527] In some embodiments of these methods, the method is an *in vivo* method. In some embodiments of these methods, the method is an *in situ* method. In some embodiments of these methods, the method is an *ex vivo* method. In some embodiments of these methods, the method is an *in vitro* method.

[000528] In some embodiments, *in situ* imaging and/or *in vivo* imaging are useful in methods to identify which patients to treat. For example, in *in situ* imaging, the activatable antibodies are used to screen patient samples to identify those patients having the appropriate protease(s) and target(s) at the appropriate location, e.g., at a tumor site.

[000529] In some embodiments *in situ* imaging is used to identify or otherwise refine a patient population suitable for treatment with an activatable antibody of the disclosure. For example, patients that test positive for both the target (e.g., the target) and a protease that cleaves the substrate in the cleavable moiety (CM) of the activatable antibody being tested (e.g., accumulate activated antibodies at the disease site) are identified as suitable candidates for treatment with such an activatable antibody comprising such a CM. Likewise, patients that test negative for either or both of the target (e.g., the target) and the protease that cleaves the substrate in the CM in the activatable antibody being tested using these methods might be identified as suitable candidates for another form of therapy. In some embodiments, such patients that test negative with respect to a first activatable antibody can be tested with other activatable antibodies comprising different CMs until a suitable activatable antibody for treatment is identified (e.g., an activatable antibody comprising a CM that is cleaved by the patient at the site of disease). In some embodiments, the patient is then administered a therapeutically effective amount of the activatable antibody for which the patient tested positive.

[000530] In some embodiments *in vivo* imaging is used to identify or otherwise refine a patient population suitable for treatment with an activatable antibody of the disclosure. For example, patients that test positive for both the target (e.g., the target) and a protease that cleaves the substrate in the cleavable moiety (CM) of the activatable antibody being tested (e.g., accumulate activated antibodies at the disease site) are identified as suitable candidates for treatment with such an activatable antibody comprising such a CM. Likewise, patients that test negative might be identified as suitable candidates for another form of therapy. In some embodiments, such patients that test negative with respect to a first

activatable antibody can be tested with other activatable antibodies comprising different CMs until a suitable activatable antibody for treatment is identified (e.g., an activatable antibody comprising a CM that is cleaved by the patient at the site of disease). In some embodiments, the patient is then administered a therapeutically effective amount of the activatable antibody for which the patient tested positive.

[000531] In some embodiments of the methods and kits, the method or kit is used to identify or otherwise refine a patient population suitable for treatment with an activatable antibody of the disclosure. For example, patients that test positive for both the target (e.g., the target) and a protease that cleaves the substrate in the cleavable moiety (CM) of the activatable antibody being tested in these methods are identified as suitable candidates for treatment with such an activatable antibody comprising such a CM. Likewise, patients that test negative for both of the targets (e.g., the target) and the protease that cleaves the substrate in the CM in the activatable antibody being tested using these methods might be identified as suitable candidates for another form of therapy. In some embodiments, such patients can be tested with other activatable antibodies until a suitable activatable antibody for treatment is identified (e.g., an activatable antibody comprising a CM that is cleaved by the patient at the site of disease). In some embodiments, patients that test negative for either of the target (e.g., the target) are identified as suitable candidates for treatment with such an activatable antibody comprising such a CM. In some embodiments, patients that test negative for either of the target (e.g., the target) are identified as not being suitable candidates for treatment with such an activatable antibody comprising such a CM. In some embodiments, such patients can be tested with other activatable antibodies until a suitable activatable antibody for treatment is identified (e.g., an activatable antibody comprising a CM that is cleaved by the patient at the site of disease). In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000532] In some embodiments, a method or kit is used to identify or otherwise refine a patient population suitable for treatment with an anti-the target activatable antibody and/or conjugated activatable antibody (e.g., activatable antibody to which a therapeutic agent is conjugated) of the disclosure, followed by treatment by administering that activatable antibody and/or conjugated activatable antibody to a subject in need thereof. For example, patients that test positive for both the targets (e.g., the target) and a protease that cleaves the substrate in the cleavable moiety (CM) of the activatable antibody and/or conjugated activatable antibody being tested in these methods are identified as suitable candidates for treatment with such antibody and/or such a conjugated activatable antibody comprising such a CM, and the patient is then administered a therapeutically effective amount of the activatable antibody and/or conjugated activatable antibody that was tested. Likewise, patients that test negative for either or both of the target (e.g., the target) and the protease that cleaves the substrate in the CM in the activatable antibody being tested using these methods might be identified as suitable candidates for another form of therapy. In some embodiments, such patients can be tested with other antibody and/or conjugated activatable antibody until a suitable antibody and/or conjugated activatable antibody for treatment is identified (e.g., an activatable antibody and/or conjugated activatable antibody comprising a CM that is cleaved by the patient at the site of disease). In some embodiments, the patient is then administered a therapeutically effective amount of the activatable antibody and/or conjugated activatable antibody for which the patient tested positive.

[000533] In some embodiments of these methods and kits, the MM is a peptide having a length from about 4 to 40 amino acids. In some embodiments of these methods and kits, the activatable antibody comprises a linker peptide, wherein the linker peptide is positioned between the MM and the CM. In some embodiments of these methods and kits, the activatable antibody comprises a linker peptide, where the linker peptide is positioned between the AB and the CM. In some embodiments of these methods and kits, the activatable antibody comprises a first linker peptide (L1) and a second linker peptide (L2), wherein the first linker peptide is positioned between the MM and the CM and the second linker peptide is positioned between the AB and the CM. In some embodiments of these methods and kits, each of L1 and L2 is a peptide of about 1 to 20 amino acids in length, and wherein each of L1 and L2 need not be the same linker. In some embodiments of these methods and kits, one or both of L1 and L2 comprises a glycine-serine polymer. In some embodiments of these methods and kits, at least one of L1 and L2 comprises an amino acid

sequence selected from the group consisting of (GS)_n, (GSGGS)_n (SEQ ID NO: 339) and (GGGS)_n (SEQ ID NO: 340), where n is an integer of at least one. In some embodiments of these methods and kits, at least one of L1 and L2 comprises an amino acid sequence having the formula (GGS)_n, where n is an integer of at least one. In some embodiments of these methods and kits, at least one of L1 and L2 comprises an amino acid sequence selected from the group consisting of Gly-Gly-Ser-Gly (SEQ ID NO: 341, Gly-Gly-Ser-Gly-Gly (SEQ ID NO: 342), Gly-Ser-Gly-Ser-Gly (SEQ ID NO: 343, Gly-Ser-Gly-Gly-Gly (SEQ ID NO: 344), Gly-Gly-Gly-Ser-Gly (SEQ ID NO: 345), and Gly-Ser-Ser-Ser-Gly (SEQ ID NO: 346).

[000534] In some embodiments of these methods and kits, the AB comprises an antibody or antibody fragment sequence selected from the cross-reactive antibody sequences presented herein. In some embodiments of these methods and kits, the AB comprises a Fab fragment, a scFv or a single chain antibody (scAb).

[000535] In some embodiments of these methods and kits, the cleaving agent is a protease that is co-localized in the subject or sample with the target and the CM is a polypeptide that functions as a substrate for the protease, wherein the protease cleaves the CM in the activatable antibody when the activatable antibody is exposed to the protease. In some embodiments of these methods and kits, the CM is a polypeptide of up to 15 amino acids in length. In some embodiments of these methods and kits, the CM is coupled to the N-terminus of the AB. In some embodiments of these methods and kits, the CM is coupled to the C-terminus of the AB. In some embodiments of these methods and kits, the CM is coupled to the N-terminus of a VL chain of the AB.

[000536] The antibodies, conjugated antibodies, activatable antibodies and/or conjugated activatable antibodies of the disclosure are used in diagnostic and prophylactic formulations. In one embodiment, an activatable antibody is administered to patients that are at risk of developing one or more of the aforementioned inflammation, inflammatory disorders, cancer or other disorders.

[000537] A patient's or organ's predisposition to one or more of the aforementioned disorders can be determined using genotypic, serological or biochemical markers.

[000538] In some embodiments of the disclosure, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody is administered to human individuals diagnosed with a clinical indication associated with one or more of the aforementioned disorders. Upon diagnosis, an antibody, a conjugated antibody, an

activatable antibody and/or a conjugated activatable antibody is administered to mitigate or reverse the effects of the clinical indication.

[000539] Antibodies, conjugated antibodies, activatable antibodies and/or conjugated activatable antibodies of the disclosure are also useful in the detection of the target in patient samples and accordingly are useful as diagnostics. For example, the antibodies, conjugated antibodies, the activatable antibodies and/or conjugated activatable antibodies of the disclosure are used in *in vitro* assays, *e.g.*, ELISA, to detect target levels in a patient sample.

[000540] In one embodiment, an antibody and/or activatable antibody of the disclosure is immobilized on a solid support (*e.g.*, the well(s) of a microtiter plate). The immobilized antibody and/or activatable antibody serves as a capture antibody for any target that may be present in a test sample. Prior to contacting the immobilized antibody and/or activatable antibody with a patient sample, the solid support is rinsed and treated with a blocking agent such as milk protein or albumin to prevent nonspecific adsorption of the analyte.

[000541] Subsequently the wells are treated with a test sample suspected of containing the antigen, or with a solution containing a standard amount of the antigen. Such a sample is, *e.g.*, a serum sample from a subject suspected of having levels of circulating antigen considered to be diagnostic of a pathology. After rinsing away the test sample or standard, the solid support is treated with a second antibody that is detectably labeled. The labeled second antibody serves as a detecting antibody. The level of detectable label is measured, and the concentration of target antigen in the test sample is determined by comparison with a standard curve developed from the standard samples.

[000542] It will be appreciated that based on the results obtained using the antibodies and/or activatable antibodies of the disclosure in an *in vitro* diagnostic assay, it is possible to stage a disease in a subject based on expression levels of the Target antigen. For a given disease, samples of blood are taken from subjects diagnosed as being at various stages in the progression of the disease, and/or at various points in the therapeutic treatment of the disease. Using a population of samples that provides statistically significant results for each stage of progression or therapy, a range of concentrations of the antigen that may be considered characteristic of each stage is designated.

[000543] Antibodies, conjugated antibodies, activatable antibodies and/or conjugated activatable antibodies can also be used in diagnostic and/or imaging methods. In some embodiments, such methods are *in vitro* methods. In some embodiments, such methods are

in vivo methods. In some embodiments, such methods are *in situ* methods. In some embodiments, such methods are *ex vivo* methods. For example, activatable antibodies having an enzymatically cleavable CM can be used to detect the presence or absence of an enzyme that is capable of cleaving the CM. Such activatable antibodies can be used in diagnostics, which can include *in vivo* detection (*e.g.*, qualitative or quantitative) of enzyme activity (or, in some embodiments, an environment of increased reduction potential such as that which can provide for reduction of a disulfide bond) through measured accumulation of activated antibodies (*i.e.*, antibodies resulting from cleavage of an activatable antibody) in a given cell or tissue of a given host organism. Such accumulation of activated antibodies indicates not only that the tissue expresses enzymatic activity (or an increased reduction potential depending on the nature of the CM) but also that the tissue expresses target to which the activated antibody binds.

[000544] For example, the CM can be selected to be a protease substrate for a protease found at the site of a tumor, at the site of a viral or bacterial infection at a biologically confined site (*e.g.*, such as in an abscess, in an organ, and the like), and the like. The AB can be one that binds a target antigen. Using methods familiar to one skilled in the art, a detectable label (*e.g.*, a fluorescent label or radioactive label or radiotracer) can be conjugated to an AB or other region of an activatable antibody. Suitable detectable labels are discussed in the context of the above screening methods and additional specific examples are provided below. Using an AB specific to a protein or peptide of the disease state, along with a protease whose activity is elevated in the disease tissue of interest, activatable antibodies will exhibit an increased rate of binding to disease tissue relative to tissues where the CM specific enzyme is not present at a detectable level or is present at a lower level than in disease tissue or is inactive (*e.g.*, in zymogen form or in complex with an inhibitor). Since small proteins and peptides are rapidly cleared from the blood by the renal filtration system, and because the enzyme specific for the CM is not present at a detectable level (or is present at lower levels in non-disease tissues or is present in inactive conformation), accumulation of activated antibodies in the disease tissue is enhanced relative to non-disease tissues.

[000545] In another example, activatable antibodies can be used to detect the presence or absence of a cleaving agent in a sample. For example, where the activatable antibodies contain a CM susceptible to cleavage by an enzyme, the activatable antibodies can be used to detect (either qualitatively or quantitatively) the presence of an enzyme in the sample. In

another example, where the activatable antibodies contain a CM susceptible to cleavage by reducing agent, the activatable antibodies can be used to detect (either qualitatively or quantitatively) the presence of reducing conditions in a sample. To facilitate analysis in these methods, the activatable antibodies can be detectably labeled, and can be bound to a support (*e.g.*, a solid support, such as a slide or bead). The detectable label can be positioned on a portion of the activatable antibody that is not released following cleavage, for example, the detectable label can be a quenched fluorescent label or other label that is not detectable until cleavage has occurred. The assay can be conducted by, for example, contacting the immobilized, detectably labeled activatable antibodies with a sample suspected of containing an enzyme and/or reducing agent for a time sufficient for cleavage to occur, then washing to remove excess sample and contaminants. The presence or absence of the cleaving agent (*e.g.*, enzyme or reducing agent) in the sample is then assessed by a change in detectable signal of the activatable antibodies prior to contacting with the sample *e.g.*, the presence of and/or an increase in detectable signal due to cleavage of the activatable antibody by the cleaving agent in the sample.

[000546] Such detection methods can be adapted to also provide for detection of the presence or absence of a target that is capable of binding the AB of the activatable antibodies when cleaved. Thus, the assays can be adapted to assess the presence or absence of a cleaving agent and the presence or absence of a target of interest. The presence or absence of the cleaving agent can be detected by the presence of and/or an increase in detectable label of the activatable antibodies as described above, and the presence or absence of the target can be detected by detection of a target-AB complex *e.g.*, by use of a detectably labeled anti-target antibody.

[000547] Activatable antibodies are also useful in *in situ* imaging for the validation of activatable antibody activation, *e.g.*, by protease cleavage, and binding to a particular target. *In situ* imaging is a technique that enables localization of proteolytic activity and target in biological samples such as cell cultures or tissue sections. Using this technique, it is possible to confirm both binding to a given target and proteolytic activity based on the presence of a detectable label (*e.g.*, a fluorescent label).

[000548] These techniques are useful with any frozen cells or tissue derived from a disease site (*e.g.* tumor tissue) or healthy tissues. These techniques are also useful with fresh cell or tissue samples.

[000549] In these techniques, an activatable antibody is labeled with a detectable label. The detectable label can be a fluorescent dye, (e.g. Fluorescein Isothiocyanate (FITC), Rhodamine Isothiocyanate (TRITC), a near infrared (NIR) dye (e.g., Qdot® nanocrystals), a colloidal metal, a hapten, a radioactive marker, biotin and an amplification reagent such as streptavidin, or an enzyme (e.g. horseradish peroxidase or alkaline phosphatase).

[000550] Detection of the label in a sample that has been incubated with the labeled, activatable antibody indicates that the sample contains the target and contains a protease that is specific for the CM of the activatable antibody. In some embodiments, the presence of the protease can be confirmed using broad spectrum protease inhibitors such as those described herein, and/or by using an agent that is specific for the protease, for example, an antibody such as A11, which is specific for the protease matriptase and inhibits the proteolytic activity of matriptase; see e.g., International Publication Number WO 2010/129609, published 11 November 2010. The same approach of using broad spectrum protease inhibitors such as those described herein, and/or by using a more selective inhibitory agent can be used to identify a protease or class of proteases specific for the CM of the activatable antibody. In some embodiments, the presence of the target can be confirmed using an agent that is specific for the target, e.g., another antibody, or the detectable label can be competed with unlabeled target. In some embodiments, unlabeled activatable antibody could be used, with detection by a labeled secondary antibody or more complex detection system.

[000551] Similar techniques are also useful for *in vivo* imaging where detection of the fluorescent signal in a subject, e.g., a mammal, including a human, indicates that the disease site contains the target and contains a protease that is specific for the CM of the activatable antibody.

[000552] These techniques are also useful in kits and/or as reagents for the detection, identification or characterization of protease activity in a variety of cells, tissues, and organisms based on the protease-specific CM in the activatable antibody.

[000553] In some embodiments, *in situ* imaging and/or *in vivo* imaging are useful in methods to identify which patients to treat. For example, in *in situ* imaging, the activatable antibodies are used to screen patient samples to identify those patients having the appropriate protease(s) and target(s) at the appropriate location, e.g., at a tumor site.

[000554] In some embodiments *in situ* imaging is used to identify or otherwise refine a patient population suitable for treatment with an activatable antibody of the disclosure. For

example, patients that test positive for both the target and a protease that cleaves the substrate in the cleavable moiety (CM) of the activatable antibody being tested (e.g., accumulate activated antibodies at the disease site) are identified as suitable candidates for treatment with such an activatable antibody comprising such a CM. Likewise, patients that test negative for either or both of the target and the protease that cleaves the substrate in the CM in the activatable antibody being tested using these methods are identified as suitable candidates for another form of therapy (*i.e.*, not suitable for treatment with the activatable antibody being tested). In some embodiments, such patients that test negative with respect to a first activatable antibody can be tested with other activatable antibodies comprising different CMs until a suitable activatable antibody for treatment is identified (e.g., an activatable antibody comprising a CM that is cleaved by the patient at the site of disease).

[000555] In some embodiments *in vivo* imaging is used to identify or otherwise refine a patient population suitable for treatment with an activatable antibody of the disclosure. For example, patients that test positive for both the target and a protease that cleaves the substrate in the cleavable moiety (CM) of the activatable antibody being tested (e.g., accumulate activated antibodies at the disease site) are identified as suitable candidates for treatment with such an activatable antibody comprising such a CM. Likewise, patients that test negative are identified as suitable candidates for another form of therapy (*i.e.*, not suitable for treatment with the activatable antibody being tested). In some embodiments, such patients that test negative with respect to a first activatable antibody can be tested with other activatable antibodies comprising different CMs until a suitable activatable antibody for treatment is identified (e.g., an activatable antibody comprising a CM that is cleaved by the patient at the site of disease).

Pharmaceutical compositions

[000556] The antibodies, conjugated antibodies, activatable antibodies and/or conjugated activatable antibodies of the disclosure (also referred to herein as “active compounds”), and derivatives, fragments, analogs and homologs thereof, can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the antibody, the conjugated antibody, activatable antibody and/or conjugated activatable antibody and a pharmaceutically acceptable carrier. As used herein, the term “pharmaceutically acceptable carrier” is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and

absorption delaying agents, and the like, compatible with pharmaceutical administration. Suitable carriers are described in the most recent edition of Remington's Pharmaceutical Sciences, a standard reference text in the field, which is incorporated herein by reference. Suitable examples of such carriers or diluents include, but are not limited to, water, saline, Ringer's solutions, dextrose solution, and 5% human serum albumin. Liposomes and non-aqueous vehicles such as fixed oils may also be used. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

[000557] A pharmaceutical composition of the disclosure is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (*i.e.*, topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[000558] Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringeability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the

like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In some embodiments, it will be desirable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, aluminum monostearate and gelatin.

[000559] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[000560] Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

[000561] For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser that contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

[000562] Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

[000563] The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

[000564] In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

[000565] It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the disclosure are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

[000566] The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

[000567] The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

Examples

EXAMPLE 1. Characterization of Anti-CD71 Antibodies

[000568] The studies provided herein were designed to evaluate binding of anti-CD71 antibodies of the disclosure.

[000569] The anti-CD71 M21 monoclonal antibody of the present disclosure was obtained using mouse hybridoma technology in accordance with techniques known in the art. Mice were immunized with human CD71 extracellular domain (ECD) and subsequent hybridomas were screened using a cytotoxicity piggyback assay, and cytotoxicity positive clones from this assay were confirmed by ELISA to bind the human CD71 ECD polypeptide and confirmed to bind cell surfaces by FACS. The anti-CD71 M21 monoclonal antibody of the present disclosure includes a heavy chain variable region (VH) of SEQ ID NO: 1, and a light chain variable region (VL) of SEQ ID NO: 2, and was used where described herein as a positive control.

[000570] The following humanized anti-CD71 antibodies, which were based on the anti-CD71 mouse monoclonal antibody M21, were tested: Ab21.10 LcB:HcA (VH of SEQ ID NO: 3 and VL of SEQ ID NO: 7), Ab21.11 LcB:HcB (VH of SEQ ID NO: 4 and VL of SEQ ID NO: 7), Ab21.12 LcB:HcC (VH of SEQ ID NO: 5 and VL of SEQ ID NO: 7), and M21 (VH of SEQ ID NO: 1 and VL of SEQ ID NO: 2). Binding of various anti-CD71 antibodies of the disclosure was confirmed by FACS (Figure 1).

[000571] As shown in Figure 1, all of the humanized anti-CD71 antibodies showed binding to human CD71 that was comparable to the binding demonstrated by the CD71 M21 mouse antibody. Binding of the humanized anti-CD71 antibodies was confirmed on the BxPC3 cell line by FACS. Briefly, BxPC3 cells were labeled with mouse monoclonal Mab21 or huCD71 (Ab21.10, Ab21.11, and Ab21.12) antibody at the indicated concentrations and subsequently detected with an Alexa Fluor 647 labeled goat anti-mouse or anti-human IgG Alexa Fluor 647, respectively.

EXAMPLE 2. Mask Discovery

[000572] The studies provided herein were designed to identify and characterize masking moieties for use in activatable anti-CD71 antibodies of the disclosure.

[000573] Anti-CD71 21.12 antibody, comprising a VH of SEQ ID NO: 5 and a VL of SEQ ID NO: 7, was used to screen a random X₁₅ peptide library with a total diversity of 6x10¹⁰, where X is any amino acid, using a method similar to that described in PCT International Publication Number WO 2010/081173, published 15 July 2010. The screening consisted of one round of MACS and five rounds of FACS sorting. The initial MACS sorting was done with protein-A Dynabeads (Invitrogen) and the anti-CD71 21.12 antibody at a concentration of 200 nM. For MACS, approximately 1x10¹² cells were screened for binding and 1x10⁷ cells were collected. Anti-CD71 21.12 was conjugated with DyLight-488 (ThermoFisher), CD71 binding activity was confirmed and anti-CD71 21.12-488 was used as a fluorescent probe for all FACS rounds. Bacterial cells were stained and positive clones were collected as follows: 20nM anti-CD71 21.12-488 with 1x10⁶ cells collected in FACS round 1, 5nM anti-CD71 21.12-488 with 6.2x10⁴ cells collected in FACS round 2 and 5nM anti-CD71 21.12-488 with 5x10³ cells and 1nM anti-CD71 21.12-488 with 5x10² cells collected in FACS round 3, 1nM anti-CD71 21.12-488 with > 2x10² cells collected in FACS rounds 4 and 5. The positive population from the second FACS round was verified to inhibit binding of the anti-CD71 21.12-488 antibody to recombinant CD71 protein. Individual peptide clones were identified by sequence analysis from the 5nM binders from FACS round 3 and the 1nM binders from FACS rounds 3, 4 and 5.

[000574] The sequences of the anti-CD71 masking moieties are listed in Table A.

Table A. Anti-CD71 masking moieties (MM)

MM	Amino Acid Sequence	SEQ ID
TF01	QFCPWSYYLIGDCDI	16
TF02	NLCTEHSFALDCRSY	17
TF03	APMPCTSPGCWLTLR	18
TF04	QSACIFPMTNSCTYS	19
TF05	ASQFCTANPECNYAG	20
TF06	THRCLPMQNFCHNPF	21
TF07	ICSFESWHQFSNCNP	22

TF08	RYS PACYSTCNSINW	23
TF09	NRWCAPMQNYCHHST	24
TF10	AHCVQMSNHPYCNHG	25
TF11	LSHVCLITPMCNAMEQ	26
TF12	AWWCAPMQNACQHYQ	27
TF13	RSPCAVPMSNSCYII	28
TF14	STRCLPMQNYCHFSD	29
TF15	PNCLPMQNAQRCHSP	30
TF16	NTAFCTHNPFENRPT	31
TF17	MYS PYCYSMCENNIYP	32
TF18	IMRSCTSPGCYLYNT	33
TF19	TRCFPMSNNPNCMNY	34
TF20	NCGNFYYTMMANCNY	35
TF21	NPNCWAPMMNSCNAF	36
TF22	WNCENVNLYLPACMQ	37
TF23	TWCPMSNHPTCTLK	38
TF24	GYPCVPMQNRCAWKS	39
TF25	LLSPFCTQMCNPKLQ	40
TF26	MSRFCTHNPECNFTY	41
TF27	MSTCWTQMTNRCTYN	42
TF28	RKSPNCYSMCNYYFN	43
TF29	TGQCWAPMQNCTHNK	44
TF30	TKMGCSKANLCNGAM	45
TF31	AHWCAPMQNRCTNAP	46
TF32	ASCPLMSNKPTCRQP	47
TF33	DQCWAPMMNMFHCNK	48
TF34	EYCEYTRPVPMGCWG	49
TF35	GMCPMSNNPNCNNY	50
TF36	GSCPLMSNRPSCHSS	51
TF37	HGDCAPMTNRCTNTT	52
TF38	KLPCWTLHQNSCTEY	53
TF39	LASCAPMQNRCYGLN	54
TF40	MDRFGCLQMCNWPKN	55

TF41	MMHCAMMQNRCSEYH	56
TF42	MRSCHSPMMNQCNHI	57
TF43	MTHCLPMQNRCTTYM	58
TF44	QSKCWTTMMNGCHIN	59
TF45	RECMQNNPACHMY	60
TF46	RPSPACKIMCNMYQH	61
TF47	SNRCWAPMTNMCTYN	62
TF48	SRKCFPMENRCNHHT	63
TF49	TPFCLPMQNSCKHTS	64
TF50	TTHCMQMKCNRPN	65
TF51	TTSCIMPMTNRCTIH	66
TF52	VYSPLCHNMCNPRNT	67
TF53	WCNNRYIAPMSNCAS	68
TF54	WEYCHNTEFIGCTYK	69
TF55	WTSFCTHNIFCNMNG	70
TF56	YLRGCYEYNMCNGYY	71
TF57	YSCAPMQNHPQCNQT	72
TF58	ACVNSNYPFDSNCMY	73
TF59	ADCQIFAMTNRCMNN	74
TF60	AECSPMSNRPQCHIH	75
TF61	AISCMPMQNSCYHTY	76
TF62	AKCWTEYMNHPECRI	77
TF63	ALHVLFCYNFCNGPTT	78
TF64	AMCVGCSRECCWL GK	79
TF65	AQPCAPMQNSCHQSV	80
TF66	ARLGCTKFNMCFYS	81
TF67	ASCPLMSNKPQCLKH	82
TF68	ASTFCTHNPACAYIP	83
TF69	ATCPLMSNRPSCISA	84
TF70	ATSMYCYSMCNTNNY	85
TF71	AYRCMPMQNSCYSTN	86
TF72	AYTGCTANNMCNAAY	87
TF73	CASLCKNQPCSSNRN	88

TF74	DCPMSYYLPMVNCMH	89
TF75	DCWRYLGNMNNWCYL	90
TF76	DLTFCTANPFCSREF	91
TF77	DNSFCTHNYCMLNK	92
TF78	DSRCWTGHQNRNTI	93
TF79	EACMHSLHWRSWCKV	94
TF80	ECENLYKAPMENCLR	95
TF81	ECRLRNQPMMSNCSM	96
TF82	EHPCMPMENRCYTLN	97
TF83	ESVCMQPMQNVCTWTT	98
TF84	EYCNMIPTNYCSYN	99
TF85	EYLWCELMVSCSRLM	100
TF86	FKHFCTHNPECKYNT	101
TF87	FPQCSPMQNKCGEGR	102
TF88	FRSQYCQSMCNYYNN	103
TF89	FRWCAPMQNYCHHST	104
TF90	FSCTAMSYAEDCWSW	105
TF91	FSQGCFHDNMCNRYQ	106
TF92	GCFQLCETCCIHSFD	107
TF93	GGHCAPMQNNCYYS	108
TF94	GKYCKMPMVNRCYIM	109
TF95	GMYCTRMQNSCAHNN	110
TF96	GPFFCTHSPECQFDK	111
TF97	GQCFCSSLFCRPLKP	112
TF98	GQYCNLPMVNSCNYL	113
TF99	GRSCWTSFMNRCTND	114
TF100	GTCSWGSPGGQLCGR	115
TF101	GVCLLGLETYLLCFS	116
TF102	GWSYCINIYCRIDDN	117
TF103	HCANHHPTPMQNCKT	118
TF104	HCKEAPWTLYMNCNT	119
TF105	HCWWPQNI PMNNCIS	120
TF106	HFCQNHQMVNACTA	121

TF107	HGQCMQPMQNRCSNSY	122
TF108	HISPYCQQMCNKWYN	123
TF109	HLMECSYENMCNRY	124
TF110	HMYCAPMQNACTKNT	125
TF111	HNCWTMMMNKPYCSN	126
TF112	HNYCWAPMQNNCMIE	127
TF113	HPPCMQPMQNRVSDH	128
TF114	HPVFCTHNPCHHNY	129
TF115	HRGCMQPMQNSCSTKN	130
TF116	HSSCWAPHTNRCHMG	131
TF117	HSTFCTHNRNCNNHS	132
TF118	HTCHYNTPMTNCHYS	133
TF119	HVSPSCTMMCNSYNF	134
TF120	IAAECTKHNMCCNAM	135
TF121	ICDSSAAPMTNRCMN	136
TF122	INVCMPMQNRCIGQN	137
TF123	IPCHGQWNFSNCKYQ	138
TF124	IPTCVPMQNHCHYSA	139
TF125	IVSKHCTSMCNPNWY	140
TF126	IYCPMSNNIACQNN	141
TF127	KCALSNQPLMSNCLM	142
TF128	KCHDNNFCRNNMCNC	143
TF129	KCSLKHTLPMSNCMA	144
TF130	KCTNAFRMPMENCMS	145
TF131	KFCQQPQTPMQNCSS	146
TF132	KGCYTWTMMNCIPK	147
TF133	KHSPYCFSMCNLQIS	148
TF134	KPRCSPMQNMCYNKQ	149
TF135	KTPCWTSMMNGCNHY	150
TF136	KYCPLMSNKPPCKPK	151
TF137	KYSPMCFHCNSQIK	152
TF138	LCELSASTWSELCCR	153
TF139	LCGVYEMPMTNACAA	154

TF140	LDTFCTHNFNCSKHN	155
TF141	LEKNCTYNNMCNGYH	156
TF142	LESFCTHSPACVKHR	157
TF143	LGTCLPMQNGCTGSR	158
TF144	LHDCRSPAWD CFSLG	159
TF145	LHGFCTHNYSCSLEK	160
TF146	LKCLAEQISLDCRST	161
TF147	LKCQQPYTLMQNCAI	162
TF148	LMCTQLPMLNGPCHT	163
TF149	LQRFCTHNMLCQHNS	164
TF150	LRSPYCMGMCNFMTY	165
TF151	LSSFCTSNPFCNPYH	166
TF152	LTSPACRIMCNMDMS	167
TF153	LYNFCTHNNQCNTF	168
TF154	MHDCRMPMTNSCTYP	169
TF155	MKSPACKSMCNLYIN	170
TF156	MKYFCTHNYNCNNH	171
TF157	MPCHLMPMHNKCQST	172
TF158	MPGCMQPMQNGCKHYN	173
TF159	MTEFCTHNRNCMMIS	174
TF160	MTICAPMQNYCPNAN	175
TF161	MYAPTCQQMCNPVSK	176
TF162	MYCWTAMMNKPCRFN	177
TF163	MYSSYCQIMCNPVPK	178
TF164	NCTMDQILPMSNCNM	179
TF165	NCVNIHYTMSNCNF	180
TF166	NDYCLPMQNKCNMLS	181
TF167	NFCSIPMSNHPTCNN	182
TF168	NFNCWYNPNSDCNYY	183
TF169	NPACMPMQNSCSHYD	184
TF170	NRPCWTDMMNACNHG	185
TF171	NSNFCTANYNCNWIN	186
TF172	PCHTGTYPMPMNCHT	187

TF173	PECPPMSNNPHCNKL	188
TF174	PMCWEFYMMNKCIPY	189
TF175	PSCPPMSNQPACNRT	190
TF176	PSRCFVPMQNYCHNY	191
TF177	PTCREHFAMHNRCHD	192
TF178	PTHCAPMQNACTPHI	193
TF179	PTNCLPMQNRCKMNH	194
TF180	QASFCTHNYNCRTNN	195
TF181	QCGFLPTDKFSNCKN	196
TF182	QNHCVTTMINGCSWT	197
TF183	QQSFCTHSPACIASY	198
TF184	QSICWTYMTNGCMNY	199
TF185	QSQCAPMQNSCAQKH	200
TF186	QSYCYYWTMHCNDRY	201
TF187	QWYCSPMQNGCSNDN	202
TF188	QYCAVMSNNPYCRIN	203
TF189	QYSPYCYSMCNGHKN	204
TF190	RANCSQLGWSHCNIP	205
TF191	RCGNTYYTQMANCNN	206
TF192	RCPAYQHMAHVNCAN	207
TF193	RCSNLVNTPMQNCNM	208
TF194	REPCNVPMTNSCMRN	209
TF195	RGPCAVPMTNHCYSL	210
TF196	RISPGCTLMCNHYMY	211
TF197	RLCPPMSNKTACNNR	212
TF198	RMSCMPMQNSCHNTT	213
TF199	RNICLPMQNYCNNNN	214
TF200	RPCVPMQNDPTCTHI	215
TF201	RPMPCTSPGCTIGVH	216
TF202	RPSFCTHNKNCNHNR	217
TF203	RSTFCTHNPYCTQKL	218
TF204	RTATCSSFGTCSSLW	219
TF205	RTCQLMQNHDPCTMK	220

TF206	RTHCMPMQNHCMDSK	221
TF207	RTWVGCENHCYMLEK	222
TF208	RYGCVPMQNACGGPW	223
TF209	SAQCAPMQNSCHSIK	224
TF210	SCLNAHYQPMSNCNG	225
TF211	SCTPVDDNYFNGCSK	226
TF212	SDYCFPMQNCAMWS	227
TF213	SFSEKCQNMCPFNQ	228
TF214	SGCLPMSNNPVCNNR	229
TF215	SIGCLLPMTNGCNYK	230
TF216	SMCLPMQNNPYCNHK	231
TF217	SMFCFWHPDVCHDQE	232
TF218	SMHCSPMQNFCSNNF	233
TF219	SPPCDNMRNWCHPNM	234
TF220	SRECWAMMQNCSRNM	235
TF221	SRVCLPMQNYCKMNH	236
TF222	SSFCSPMQNFCAISI	237
TF223	SSLCMPMQNYCMNHK	238
TF224	STNACNMRNYCMWHQ	239
TF225	STNCIMPMTNGCRLN	240
TF226	STRCNFQAQMHCNYM	241
TF227	STSPACNHMCNPWNG	242
TF228	SVCSRSSPGGVLCRE	243
TF229	SVSPVCMSLCNRYHH	244
TF230	SWTCVPMQNACMHRT	245
TF231	SYCLPMTNKPMCNYM	246
TF232	SYCPHMAMAESCNEW	247
TF233	SYSSSCTQMCNHSYY	248
TF234	TAMPCTSPGCAMEPN	249
TF235	TAMRNCFDNMTN	250
TF236	TASRSCFTMCNLYNH	251
TF237	TCHFPMYYQFANCIP	252
TF238	TCHSNYGPMQNSCYM	253

TF239	TCMHPNTGFYENCTI	254
TF240	TCSIDNYTSSKNC.M	255
TF241	TDNCYGSRWIKCPQT	256
TF242	TEACSPMQNKCTHIY	257
TF243	TFDCVPMQNWCSNN	258
TF244	TGRCQSPMTNGCHYK	259
TF245	THCSPMQNHPSCLHQ	260
TF246	TMQCMQPMQNKCNAN	261
TF247	TNPCIMPMVNHCHPL	262
TF248	TPWCAPMQNACPKGQ	263
TF249	TQMFCTHNPECQINL	264
TF250	TRCLPMSNHPRCAMP	265
TF251	TRFCAPMQNHCMGHN	266
TF252	TRKPCHSPGQCIAMY	267
TF253	TTCPMWAPMTNCTKS	268
TF254	TTHCSPMQNGCTINR	269
TF255	TWSPYCLSMCNLRYP	270
TF256	VAFCLPMTNKCAQV	271
TF257	VATWCTVGPACAIKG	272
TF258	VCSNTQNFPMMNCNY	273
TF259	VGGCYVDDLGCARMY	274
TF260	VHLCAPMQNGCMNTQ	275
TF261	VKSPFCFSTCNMRMN	276
TF262	VSCSHGSPDGLMCRG	277
TF263	VSRECTHLNHCNRLS	278
TF264	VTCPMSNRNPACNYH	279
TF265	VTSPACQNM CNTYNN	280
TF266	VVGCGFWDFGCQRHF	281
TF267	VVSPSCFSFCNTHWV	282
TF268	VWCPRMSNNPHCASM	283
TF269	VYSPLCKSFCNPIYY	284
TF270	WGKCSPMQNKCTNNS	285
TF271	WRGGCFSPGSCGLLL	286

TF272	WSQCQFPMVNQCMVR	287
TF273	WTSPNCMSMCNNWIR	288
TF274	WWGCVPM SNRCEGGR	289
TF275	YGNRCNMINSCMNNY	290
TF276	YKAFCTHNYNCISK	291
TF277	YLFPTCIEFCNSSRQ	292
TF278	YQCPPMSNHPHCIVT	293
TF279	YQYCPPMSNRCAIKA	294
TF280	YSAYCNYSNMCNRNS	295

[000575] The TF01 and TF02 masks were truncated and alanine scanned to generate families of activatable antibodies with different masking efficiencies:

Table B. Truncation and Alanine Scanning of Masking Peptides

<u>TF01</u>	<u>QFCPWSYYLIGDCDI</u>	(SEQ ID NO: 16)
01 TF01.01	QFCaWSYYLIGDCDI	(SEQ ID NO: 297)
02 TF01.02	QFCPaSYYLIGDCDI	(SEQ ID NO: 298)
03 TF01.03	QFCPWaYYLIGDCDI	(SEQ ID NO: 299)
04 TF01.04	QFCPWSaYLIGDCDI	(SEQ ID NO: 300)
05 TF01.05	QFCPWSYaLIGDCDI	(SEQ ID NO: 301)
06 TF01.06	QFCPWSYYaIGDCDI	(SEQ ID NO: 302)
07 TF01.07	QFCPWSYYLaGDCDI	(SEQ ID NO: 303)
08 TF01.08	QFCPWSYYLIGaCDI	(SEQ ID NO: 304)
<u>TF02</u>	<u>NLCTEHSFALDCRSY</u>	(SEQ ID NO: 17)
09 TF02.09	NLCaEHSFALDCRSY	(SEQ ID NO: 305)
10 TF02.10	NLCaHSFALDCRSY	(SEQ ID NO: 306)
11 TF02.11	NLCTEaSFALDCRSY	(SEQ ID NO: 307)
12 TF02.12	NLCTEHaSFALDCRSY	(SEQ ID NO: 308)
13 TF02.13	NLCTEHSaALDCRSY	(SEQ ID NO: 309)
14 TF02.14	NLCTEHSFAaDCRSY	(SEQ ID NO: 310)
15 TF02.15	NLCTEHSFALaCRSY	(SEQ ID NO: 311)
16 TF02.16	NLCTEHSFALDCaSY	(SEQ ID NO: 312)
17 TF02.17	CTEHSFALDCRSY	(SEQ ID NO: 313)
18 TF02.18	CTEHSFALDC	(SEQ ID NO: 314)

[000576] These masking peptides were used to generate anti-CD71 activatable antibodies of the disclosure. The sequences for certain of these anti-CD71 activatable antibodies are shown below in Table C. In some embodiments, these anti-CD71 activatable antibodies include cleavable moiety 2001 (ISSGLLSGRSDNH; SEQ ID NO: 406), cleavable moiety 3001 (AVGLLAPPGGLSGRSDNH; SEQ ID NO: 412), cleavable moiety 2007 (ISSGLLSGRSDIH; SEQ ID NO: 684), cleavable moiety 2008 (ISSGLLSGRSDQH; SEQ ID NO: 685), cleavable moiety 2011 (ISSGLLSGRSDNP; SEQ ID NO: 688), cleavable moiety 2012 (ISSGLLSGRSANP; SEQ ID NO: 689), cleavable moiety 2013 (ISSGLLSGRSANI; SEQ ID NO: 690), cleavable moiety 3007 (AVGLLAPPGGLSGRSDIH; SEQ ID NO: 692), cleavable moiety 3008 (AVGLLAPPGGLSGRSDQH; SEQ ID NO: 693), cleavable moiety 3011 (AVGLLAPPGGLSGRSDNP; SEQ ID NO: 696), cleavable moiety 3012 (AVGLLAPPGGLSGRSANP; SEQ ID NO: 697), or cleavable moiety 3013 (AVGLLAPPGGLSGRSANI; SEQ ID NO: 698), as indicated.

[000577] While certain sequences shown below include the spacer sequence of SEQ ID NO: 645, those of ordinary skill in the art appreciate that the activatable anti-CD71 antibodies of the disclosure can include any suitable spacer sequence, such as, for example, a spacer sequence selected from the group consisting of QGQSGQG (SEQ ID NO: 645), QGQSGQ (SEQ ID NO: 424), QGQSG (SEQ ID NO: 646), QGQS (SEQ ID NO: 647), QGQ (SEQ ID NO: 648), QG (SEQ ID NO: 649), GQSGQG (SEQ ID NO: 666), QSGQG (SEQ ID NO: 667), SGQG (SEQ ID NO: 668), GQG (SEQ ID NO: 669), G, or Q. In some embodiments, the activatable anti-CD71 antibodies of the disclosure can have no spacer sequence joined to its N-terminus.

Table C. Anti-CD71 Activatable Antibody Sequences

HuCD71_HcC-des

Amino Acid Sequence

QVQLVQSGAEVKKPGASVKMSCKASGYTFTSYWMHWVRQAPGQGLEWIGAIYPGNSETGYAQKFQGR
ATLTADTSTSTAYMELSSLRSEDTAVYYCTRENWDPGFQAFWGQGTLLITVSSASTKGPSVFPLAPSSK
STSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYIC
NVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVS
HEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK

TISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDS
GSFFLYSKLTVDKSRWQQGNVFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 699)

HuCD71_HcC-des

Nucleotide sequence

CAGGTGCAGCTGGTGCAGTCTGGCGCCGAAGTGAAGAAACCTGGCGCCTCCGTGAAGATGTCCTGCA
AGGCCTCCGGCTACACCTTCACCAGCTACTGGATGCACTGGGTGCGACAGGCTCCAGGCCAGGGCCT
CGAATGGATCGGCGCCATCTACCCCGGCAACTCCGAGACAGGCTACGCCCAGAAGTTCCAGGGCAGA
GCCACCCTGACCGCCGACACCTCCACCTCCACCGCCTACATGGAAGTGTCCAGCCTGCGGAGCGAGG
ACACCGCCGTGTACTACTGCACCAGAGAGAACTGGGACCCCGGCTTCGCCTTCTGGGGCCAGGGCAC
CCTGATCACCGTGTCTCCGCCAGCACCAAGGGCCCCCTCCGTGTTCCCTCTGGCCCCCTTCCAGCAAG
TCCACCTCTGGCGGCACAGCTGCCCTGGGCTGCCTGGTGAAAGACTACTTCCCCGAGCCCGTGACCG
TGTCTGGAAGTCTGGCGCCCTGACCAGCGGAGTGCACACCTTCCCTGCCGTGCTGCAGTCTCCGG
CCTGTACTCCCTGTCTCCGTGGTGACAGTGCCTCCTCCAGCCTGGGCACCCAGACCTACATCTGC
AACGTGAACCACAAGCCCTCCAACACCAAGGTGGACAAGAAGTGGAACCCAAGTCTGCGACAAGA
CCCACACCTGTCCTCCCTGCCCTGCCCCCTGAACTGCTGGGCGGACCTTCCGTGTTTCTGTTCCCCC
AAAGCCCAAGGACACCCTGATGATCTCCCGGACCCCGAAGTGACCTGCGTGGTGGTGGACGTGTCC
CACGAGGACCCTGAAGTGAAGTTCAATTGGTACGTGGACGGCGTGGAAGTGACAACGCCAAGACCA
AGCCCAGAGAGGAACAGTACAACCTCCACCTACCGGGTGGTGTCCGTGCTGACCGTGTGACCAGGA
CTGGCTGAACGGCAAAGAGTACAAGTGCAAGGTGTCCAACAAGGCCCTGCCTGCCCCCATCGAAAAG
ACCATCTCCAAGGCCAAGGGCCAGCCCCGCGAGCCCCAGGTGTACACACTGCCACCTAGCCGGGAAG
AGATGACCAAGAACCAGGTGTCCCTGACCTGTCTGGTGAAAGGCTTCTACCCCTCCGATATCGCCGT
GGAATGGGAGAGCAACGGCCAGCCCCGAGAACAACCTACAAGACCACCCACCTGTGCTGGACTCCGAC
GGCTCATTCCTTCTGTACTCCAAGCTGACCGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCT
CCTGCAGCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGAGCCCCGG
C (SEQ ID NO: 700)

HuCD71_HcC

Amino Acid Sequence

QVQLVQSGAEVKKPGASVKMSCKASGYTFTSYWMHWVRQAPGQGLEWIGAIYPGNSETGYAQKFQGR
ATLTADTSTSTAYMELSSLRSEDTAVYYCTRENWDPGFAFWGQGLITVSSASTKGPSVFPLAPSSK
STSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYIC
NVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVS
HEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK
TISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDS
GSFFLYSKLTVDKSRWQQGNVFSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 325)

HuCD71_HcC

Nucleotide sequence

CAGGTGCAGCTGGTGCAGTCTGGCGCCGAAGTGAAGAAACCTGGCGCCTCCGTGAAGATGTCCTGCA
AGGCCTCCGGCTACACCTTCACCAGCTACTGGATGCACTGGGTGCGACAGGCTCCAGGCCAGGGCCT
CGAATGGATCGGCGCCATCTACCCCGGCAACTCCGAGACAGGCTACGCCCAGAAGTTCCAGGGCAGA
GCCACCCTGACCGCCGACACCTCCACCTCCACCGCCTACATGGAACTGTCCAGCCTGCGGAGCGAGG
ACACCGCCGTGTACTACTGCACCAGAGAGAACTGGGACCCCGGCTTCGCCTTCTGGGGCCAGGGCAC
CCTGATCACCGTGTCTCCCGCCAGCACCAAGGGCCCCCTCCGTGTTCCCTCTGGCCCCCTTCCAGCAAG
TCCACCTCTGGCGGCACAGCTGCCCTGGGCTGCCTGGTGAAAGACTACTTCCCCGAGCCCGTGACCG
TGTCCTGGAACCTCTGGCGCCCTGACCAGCGGAGTGACACCTTCCCTGCCGTGCTGCAGTCTCCGG
CCTGTACTCCCTGTCTCCGTGGTGACAGTGCCCTCCTCCAGCCTGGGCACCCAGACCTACATCTGC
AACGTGAACCACAAGCCCTCCAACACCAAGGTGGACAAGAAGGTGGAACCCAAGTCTCGACAAGA
CCCACACCTGTCTCTCCCTGCCCTGCCCCTGAAGTGTCTGGGCGGACCTTCCGTGTTTCTGTTCCCCC
AAAGCCCAAGGACACCCCTGATGATCTCCCGGACCCCCGAAGTGACCTGCGTGGTGGTGGACGTGTCC
CACGAGGACCCTGAAGTGAAGTTCAATTGGTACGTGGACGGCGTGGAAGTGCACAACGCCAAGACCA
AGCCCAGAGAGGAACAGTACAACCTCCACCTACCGGGTGGTGTCCGTGCTGACCGTGCTGCACCAGGA
CTGGCTGAACGGCAAAGAGTACAAGTGCAAGGTGTCCAACAAGGCCCTGCCTGCCCCCATCGAAAAG
ACCATCTCCAAGGCCAAGGGCCAGCCCCGCGAGCCCCAGGTGTACACACTGCCACCTAGCCGGGAAG
AGATGACCAAGAACCAGGTGTCCCTGACCTGTCTGGTGAAAGGCTTCTACCCCTCCGATATCGCCGT
GGAATGGGAGAGCAACGGCCAGCCCGAGAACAACCTACAAGACCACCCACCTGTGCTGGACTCCGAC
GGCTCATTCCTTCTGTACTCCAAGCTGACCGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCT
CCTGCAGCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGAGCCCCGG
CAAG (SEQ ID NO: 326)

HuCD71_LcB

Amino Acid Sequence

DIQMTQSPSSLSASVGDRTITCSASSSVYYMYWFQKPKGAPKLWIYSTSNLASGVPSRFSGSGSG
TDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCL
LNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLS
SPVTKSFNRGEC (SEQ ID NO: 323)

HuCD71_LcB

Nucleotide sequence

GACATCCAGATGACCCAGTCCCCATCCAGCCTGTCCGCCTCCGTGGGCGACAGAGTGACAATCACCT
GTTCCGCCAGCTCCTCCGTGTACTACATGTACTGGTTCCAGCAGAAGCCCGGCAAGGCCCCCAAGCT
GTGGATCTACTCCACCTCCAACCTGGCCTCCGGCGTGCCCTCCAGATTCTCCGGCTCTGGCTCCGGC
ACCGACTACACCCTGACCATCTCCAGCATGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGC

GGCGGAACCTACCCCTACACCTTCGGCCAGGGCACCAAGCTGGAAATCAAGCGGACCGTGGCCGCTCC
 CAGCGTGTTCATCTTCCCACCCCTCCGACGAGCAGCTGAAGTCCGGCACC GCCAGCGTCGTGTGCCTG
 CTGAACAACCTTCTACCCCCGCGAGGCCAAGGTGCAGTGGAAAGTGGACAACGCCCTGCAGTCCGGCA
 ACTCCCAGGAATCCGTACCGAGCAGGACTCCAAGGACAGCACCTACTCCCTGTCTCCACCCTGAC
 CCTGTCCAAGGCCGACTACGAGAAGCACAAGGTGTACGCCTGCGAAGTGACCCACCAGGGCCTGTCC
 AGCCCCGTGACCAAGTCCTTCAACCGCGGCGAGTGC (SEQ ID NO: 324)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_2001 (SEQ ID NO: 650)]

Amino Acid Sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGSISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGD
 RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
 YYCQQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCCLNNFYPREAKVQWKVDN
 ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
 (SEQ ID NO: 327)

[spacer (SEQ ID NO: 662)] [huCD71Lc_TF01_2001 (SEQ ID NO: 651)]

Nucleotide Sequence

[CAGGGCCAGTCCGGCCAGGGA] [CAGTTCTGCCCTTGGTCCTACTACCTGATCGGCGACTGCGACA
 TCGGCGGAGGCTCCTCCGGCGGCTCCATCTCCTCTGGCCTGCTGTCCGGCAGATCCGACAACCACGG
 CGGTGGCAGCGACATCCAGATGACCCAGTCCCCATCCAGCCTGTCCGCCTCCGTGGGCGACAGAGTG
 ACAATCACCTGTTCCGCCAGCTCCTCCGTGTACTACATGTACTGGTTCAGCAGAAGCCCGGCAAGG
 CCCCCAAGCTGTGGATCTACTCCACCTCCAACCTGGCCTCCGGCGTGCCCTCCAGATTCTCCGGCTC
 TGGCTCCGGCACCGACTACACCCTGACCATCTCCAGCATGCAGCCCGAGGACTTCGCCACCTACTAC
 TGCCAGCAGCGGCGGAACCTACCCCTACACCTTCGGCCAGGGCACCAAGCTGGAAATCAAGCGGACCG
 TGGCCGCTCCCAGCGTGTTCATCTTCCCACCCCTCCGACGAGCAGCTGAAGTCCGGCACC GCCAGCGT
 CGTGTGCCTGCTGAACAACCTTCTACCCCCGCGAGGCCAAGGTGCAGTGGAAAGTGGACAACGCCCTG
 CAGTCCGGCAACTCCCAGGAATCCGTACCGAGCAGGACTCCAAGGACAGCACCTACTCCCTGTCTCT
 CCACCCTGACCCTGTCCAAGGCCGACTACGAGAAGCACAAGGTGTACGCCTGCGAAGTGACCCACCA
 GGCCTGTCCAGCCCCGTGACCAAGTCCTTCAACCGCGGCGAGTGC] (SEQ ID NO: 328)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_3001 (SEQ ID NO: 652)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS AVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSLSA
 SVGDRVITITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
 DFATYYCQQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCCLNNFYPREAKVQW
 KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
 (SEQ ID NO: 329)

[spacer (SEQ ID NO: 662)] [huCD71Lc_TF01_3001 (SEQ ID NO: 653)]

Nucleotide sequence

[CAGGGCCAGTCCGGCCAGGGA] [CAGTTCTGCCCTTGGTCCTACTACCTGATCGGCGACTGCGACA
TCGGCGGAGGCTCCTCTGGCGGCTCTGCTGTGGGCCTGCTGGCTCCACCTGGCGGCCTGTCCGGCAG
ATCTGACAACCACGGCGGCTCCGACATCCAGATGACCCAGTCCCCCTCCAGCCTGTCCGCCTCCGTG
GGCGACAGAGTGACAATCACCTGTTCCGCCAGCTCCTCCGTGTACTACATGTACTGGTTCCAGCAGA
AGCCCGGAAAGGCCCCCAAGCTGTGGATCTACTCCACCTCCAACCTGGCCTCCGGCGTGCCCTCCAG
ATTCTCCGGCTCTGGCTCCGGCACCGACTACACCCTGACCATCTCCAGCATGCAGCCCGAGGACTTC
GCCACCTACTACTGCCAGCAGCGGCGGAACCTACCCCTACACCTTCGGCCAGGGCACCAAGCTGGAAA
TCAAGCGGACCGTGGCCGCTCCTTCCGTGTTTCATCTTCCACCCCTCCGACGAGCAGCTGAAGAGCGG
CACCGCCAGCGTGGTCTGCCTGCTGAACAACTTCTACCCCCGCGAGGCCAAGGTGCAGTGGAAGGTG
GACAACGCCCTGCAGTCCGGCAACTCCCAGGAATCCGTCACCGAGCAGGACTCCAAGGACAGCACCT
ACTCCCTGTCTCCACCTGACCCTGTCCAAGGCCGACTACGAGAAGCACAAGGTGTACGCCTGCGA
AGTGACCCACCAGGGCCTGTCCAGCCCCGTGACCAAGTCCTTCAACCGGGGCGAGTGC] (SEQ ID
NO: 330)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_2001 (SEQ ID NO: 654)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAIDCRSYGGGSSGGSISSGLLSGRSDNHGGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQKPKGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVCLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 331)

[spacer (SEQ ID NO: 663)] [huCD71Lc_TF02.13_2001 (SEQ ID NO: 655)]

Nucleotide sequence

[CAGGGACAGTCTGGCCAGGGC] [AACCTGTGCACCGAGCACTCTGCCGCTCTGGACTGCAGATCCT
ACGGCGGAGGCTCCTCCGGCGGCTCCATCTCCTCTGGCCTGCTGTCCGGCAGATCCGACAACCATGG
CGGCGGATCCGACATCCAGATGACCCAGTCCCCCTCCAGCCTGTCCGCCTCCGTGGGCGACAGAGTG
ACAATCACCTGTTCCGCCAGCTCCTCCGTGTACTACATGTACTGGTTCCAGCAGAAGCCCGGCAAGG
CCCCCAAGCTGTGGATCTACTCCACCAGCAACCTGGCCTCCGGCGTGCCCTCCAGATTCTCCGGCTC
TGGCTCCGGCACCGACTACACCCTGACCATCTCCAGCATGCAGCCCGAGGACTTCGCCACCTACTAC
TGCCAGCAGCGGCGGAACCTACCCCTACACCTTCGGACAGGGCACCAAGCTGGAAATCAAGCGGACCG
TGGCCGCTCCCAGCGTGTTCATCTTCCACCCCTCCGACGAGCAGCTGAAGTCCGGCACCGCCAGCGT
GGTCTGCCTGCTGAACAACTTCTACCCCCGCGAGGCCAAGGTGCAGTGGAAGGTGGACAACGCCCTG
CAGTCCGGCAACTCCCAGGAATCCGTCACCGAGCAGGACTCCAAGGACAGCACCTACTCCCTGTCTCT

CCACCCTGACCCTGTCCAAGGCCGACTACGAGAAGCACAAGGTGTACGCCTGCGAAGTGACCCACCA
GGGCCTGTCCAGCCCCGTGACCAAGTCCTTCAACCGGGGCGAGTGC] (SEQ ID NO: 332)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_3001 (SEQ ID NO: 656)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAIDCRSYGGSSGSAVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 333)

[spacer (SEQ ID NO: 664)] [huCD71Lc_TF02.13_3001 (SEQ ID NO: 657)]

Nucleotide sequence

[CAGGGCCAGTCTGGACAGGGC] [AACCTGTGCACCGAGCACTCTGCCGCTCTGGACTGCAGATCCT
ACGGCGGAGGCTCCTCTGGCGGCTCTGCTGTGGGCCTGCTGGCTCCACCTGGCGGCCTGTCCGGCAG
ATCTGACAACCACGGCGGCTCCGACATCCAGATGACCCAGTCCCCCTCCAGCCTGTCTGCCTCCGTG
GGCGACAGAGTGACAATCACCTGTTCCGCCAGCTCCTCCGTGTACTACATGTACTGGTTCCAGCAGA
AGCCCGGCAAGGCCCCCAAGCTGTGGATCTACTCCACCTCCAACCTGGCCTCCGGCGTGCCCTCCAG
ATTCTCCGGCTCTGGCTCCGGCACCGACTACACCCTGACCATCTCCAGCATGCAGCCCCGAGGACTTC
GCCACCTACTACTGCCAGCAGCGGCGGAACCTACCCCTACACCTTCGGACAGGGCACCAAGCTGGAAA
TCAAGCGGACCGTGGCCGCTCCCTCCGTGTTTCATCTTCCACCCCTCCGACGAGCAGCTGAAGTCCGG
CACCGCCAGCGTGGTCTGCCTGCTGAACAACTTCTACCCCCGCGAGGCCAAGGTGCAGTGGAAGGTG
GACAACGCCCTGCAGTCCGGCAACTCCCAGGAATCCGTACCGAGCAGGACTCCAAGGACAGCACCT
ACTCCCTGTCCTCCACCCTGACCCTGTCCAAGGCCGACTACGAGAAGCACAAGGTGTACGCCTGCGA
AGTGACCCACCAGGGCCTGTCCAGCCCCGTGACCAAGTCCTTCAACCGGGGCGAGTGC] (SEQ ID
NO: 334)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_2001 (SEQ ID NO: 658)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGSISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVTIT
CSASSSVYYMYWFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQ
RRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSG
NSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC] (SEQ ID
NO: 335)

[spacer (SEQ ID NO: 665)] [huCD71Lc_TF02.18_2001 (SEQ ID NO: 659)]

Nucleotide sequence

[CAGGGCCAGTCTGGCCAGGGC] [TGCACCGAGCACAGCTTCGCCCTGGACTGTGGCGGCGGATCCT
CCGGCGGCTCCATCTCCTCTGGCCTGCTGTCCGGCAGATCCGACAACCACGGCGGAGGCTCCGACAT
CCAGATGACCCAGTCCCCCTCCAGCCTGTCCGCCTCCGTGGGCGACAGAGTGACAATCACCTGTTCC
GCCAGCTCCTCCGTGTACTACATGTACTGGTTCCAGCAGAAGCCCGGCAAGGCCCCCAAGCTGTGGA
TCTACTCCACCTCCAACCTGGCCTCCGGCGTGCCCTCCAGATTCTCCGGCTCTGGCTCCGGCACCGA
CTACACCCTGACCATCTCCAGCATGCAGCCCCGAGGACTTCGCCACCTACTACTGCCAGCAGCGGCGG
AACTACCCCTACACCTTCGGCCAGGGCACCAAGCTGGAAATCAAGCGGACCGTGGCCGCTCCCTCCG
TGTTTCATCTTCCCACCCTCCGACGAGCAGCTGAAGTCCGGCACCGCCAGCGTGGTGTGCCTGCTGAA
CAACTTCTACCCCCGCGAGGCCAAGGTGCAGTGGAAGGTGGACAACGCCCTGCAGTCCGGCAACTCC
CAGGAATCCGTGACCGAGCAGGACTCCAAGGACAGCACCTACTCCCTGTCTCTCCACCCTGACCCTGT
CCAAGGCCGACTACGAGAAGCACAAGGTGTACGCCTGCGAAGTGACCCACCAGGGCCTGTCCAGCCC
CGTGACCAAGTCCTTCAACCGGGGCGAGTGC] (SEQ ID NO: 336)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_3001 (SEQ ID NO: 660)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSVAVGLLAPPGGLSGRSDNHGGSIDIQMTQSPSSLSASVGDR
VTITCSASSSVYYMYWFQQKPKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATY
YCQQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNA
LQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC] (SEQ
ID NO: 337)

[spacer (SEQ ID NO: 665)] [huCD71Lc_TF02.18_3001 (SEQ ID NO: 661)]

Nucleotide sequence

[CAGGGCCAGTCTGGCCAGGGC] [TGCACCGAGCACAGCTTCGCCCTGGACTGTGGCGGCGGATCTT
CTGGCGGCTCTGCTGTGGGCCTGCTGGCTCCTCCTGGCGGCCTGTCCGGCAGATCTGACAACCACGG
CGGCTCCGACATCCAGATGACCCAGTCCCCCTCCAGCCTGTCCGCCTCCGTGGGCGACAGAGTGACA
ATCACCTGTTCCGCCAGCTCCTCCGTGTACTACATGTACTGGTTCCAGCAGAAGCCCGCAAGGCCC
CCAAGCTGTGGATCTACTCCACCTCCAACCTGGCCTCCGGCGTGCCCTCCAGATTCTCCGGCTCTGG
CTCCGGCACCGACTACACCCTGACCATCTCCAGCATGCAGCCCCGAGGACTTCGCCACCTACTACTGC
CAGCAGCGGCGGAACTACCCCTACACCTTCGGCCAGGGCACCAAGCTGGAAATCAAGCGGACCGTGG
CCGCTCCCTCCGTGTTTCATCTTCCCACCCTCCGACGAGCAGCTGAAGTCCGGCACCGCCAGCGTGGT
GTGCCTGCTGAACAACCTTCTACCCCCGCGAGGCCAAGGTGCAGTGGAAGGTGGACAACGCCCTGCAG
TCCGGCAACTCCCAGGAATCCGTGACCGAGCAGGACTCCAAGGACAGCACCTACTCCCTGTCTCTCCA
CCCTGACCCTGTCCAAGGCCGACTACGAGAAGCACAAGGTGTACGCCTGCGAAGTGACCCACCAGGG
CCTGTCCAGCCCCGTGACCAAGTCCTTCAACCGGGGCGAGTGC] (SEQ ID NO: 338)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_2012 (SEQ ID NO: 670)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAIDCRSYGGGSSGGSSISSGLLSGRSANPGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 671)

[spacer (SEQ ID NO: 674)] [huCD71Lc_TF02.13_2012 (SEQ ID NO: 675)]

Nucleotide sequence

[CAGGGACAGTCAGGCCAGGGC] [AATCTCTGCACGGAGCATAGCGCCGCACTTGACTGTGATCTT
ACGGCGGCGGTTCTCTGGAGGCTCTATATCATCCGACTCCTCTCAGGCAGAAGCGCTAATCCTGG
CGGCGGATCTGATATACAAATGACTCAGTCACCAAGCTCCCTGAGTGCGTCAGTTGGTGATAGGGTG
ACGATCACTTGTAGTGCGAGCTCATCTGTTTATTATATGTACTGGTTTCAACAGAAACCCGGAAAAG
CACCTAAGTTGTGGATCTACAGTACCTCCAATCTGGCTTCCGGCGTCCCAGCCGGTTTTCCGGCTC
TGGAAGCGGAACGGATTACACGCTCACCATATCCTCTATGCAACCTGAAGATTCGCAACTTACTAC
TGTCAGCAACGCAGGAATTATCCATATACATTTGGTCAAGGGACTAAGCTCGAAATCAAGCGTACGG
TGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGT
TGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAGGTGGATAACGCCCTC
CAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTCAGCA
GCACCCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCATCA
GGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT] (SEQ ID NO: 676)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_3011 (SEQ ID NO: 672)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAIDCRSYGGGSSGGSSAVGLLAPPGLSGRSDNPGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 673)

[spacer (SEQ ID NO: 677)] [huCD71Lc_TF02.13_3011 (SEQ ID NO: 678)]

Nucleotide sequence

[CAAGGGCAGTCCGGTCAAGGG] [AACTTGTGTACAGAGCATCTGCCGCCCTTGACTGCAGGTCTT
ACGGCGGAGGGAGTAGTGGCGGGAGCGCGGTGGGACTTCTGGCACCACTGGTGGGTGTGTCAGGCAG
GAGCGACAATCCAGGGGGGTGAGACATCCAGATGACACAAAGTCCGAGTAGTCTCTCAGCTAGTGTG

GGCGATAGAGTCACAATTACATGTAGTGCGTCCAGTAGCGTGACTACATGTACTGGTTTCAGCAGA
 AGCCGGGCAAAGCACCGAAACTGTGGATTTACAGTACCAGCAACCTCGCCAGCGGTGTTCCCTCTCG
 ATTTTCAGGGAGTGGGAGTGGGACCGACTACACGCTCACCATCTCAAGTATGCAGCCAGAAGATTTTC
 GCTACCTACTATTGCCAGCAGCGGCGGAATTATCCCTACACGTTCCGGTCAAGGCACAAAACCTGGAAA
 TCAAACGTACGGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGG
 AACTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCAGAGAGGCCAAAGTACAGTGGAAGGTG
 GATAACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCT
 ACAGCCTCAGCAGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGA
 AGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT] (SEQ ID
 NO: 679)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_2011 (SEQ ID NO: 701)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSISSGLLSGRSDNPGGSDIQMTQSPSSLSASVGD
 RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
 YYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
 ALQSGNSQESVTEQDSKDSTYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
 (SEQ ID NO: 702)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_3012 (SEQ ID NO: 703)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSAVGLLAPPGGLSGRSANPGGSDIQMTQSPSSLSA
 SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
 DFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
 KVDNALQSGNSQESVTEQDSKDSTYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
 (SEQ ID NO: 704)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_2011 (SEQ ID NO: 705)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSISSGLLSGRSDNPGGSDIQMTQSPSSLSASVGD
 RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQ
 QRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSG
 NSQESVTEQDSKDSTYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC] (SEQ ID
 NO: 706)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_3012 (SEQ ID NO: 707)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSAVGLLAPPGGLSGRSANPGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC] (SEQ ID NO: 708)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_2012 (SEQ ID NO: 709)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSISSGLLSGRSANPGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC] (SEQ ID NO: 710)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_3011 (SEQ ID NO: 711)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSAVGLLAPPGGLSGRSDNPGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC] (SEQ ID NO: 712)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_NSUB (SEQ ID NO: 715)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSGGSGGGSGGGSGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC] (SEQ ID NO: 716)

[spacer (SEQ ID NO: 717)] [huCD71Lc_TF02.13_NSUB (SEQ ID NO: 718)]

Nucleotide sequence

[CAAGGCCAGTCTGGCCAGGGT] [AATTTGTGCACGGAGCATAGTGCAGCTCTGGATTGCCGGAGTTATGGAGGTGGCTCGAGCGGAGGCAGCGGAGGTT CAGGCGGGAGCGGTGGGGGGTCAGGAGGTGGCTC

TGGAGGCTCAGACATCCAGATGACCCAGTCCCCCTCTTCCCTCTCTGCCAGCGTGGGTGATCGAGTG
 ACAATTACATGTTCCGCCTCTTCTAGCGTATACTATATGTACTGGTTTCAGCAGAAACCTGGAAAAG
 CCCCCAAACTGTGGATCTATTCTACTAGCAACCTGGCCTCCGGAGTCCCATCCCGGTTCTCTGGCAG
 CGGTTCTGGAACCGACTACACTCTGACCATCTCTTCTATGCAACCAGAGGACTTTGCTACTTACTAC
 TGTCAACAGAGAAGGAACTATCCTTATACTTTCCGGTCAGGGAACTAAGCTGGAAATCAAGCGTACGG
 TGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGT
 TGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAAGTGGATAACGCCCTC
 CAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTCAGCA
 GCACCCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCCAAGTCACCCATCA
 GGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT] (SEQ ID NO: 719)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_NSUB (SEQ ID NO: 720)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSGGSGGSGGSGGSGGSDIQMTQSPSSLSASVGDRVITIT
 CSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQ
 RRNYPYTFQGQTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNMFYPREAKVQWKVDNALQSG
 NSQESVTEQDSKDYSLSSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC] (SEQ ID
 NO: 315)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_2007 (SEQ ID NO: 721)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSISSGLLSGRSDIHGGGSDIQMTQSPSSLSASVGD
 RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
 YYCQQRRNYPYTFQGQTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNMFYPREAKVQWKVDN
 ALQSGNSQESVTEQDSKDYSLSSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
 (SEQ ID NO: 722)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_2008 (SEQ ID NO: 723)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSISSGLLSGRSDQHGGGSDIQMTQSPSSLSASVGD
 RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
 YYCQQRRNYPYTFQGQTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNMFYPREAKVQWKVDN
 ALQSGNSQESVTEQDSKDYSLSSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
 (SEQ ID NO: 724)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_2013 (SEQ ID NO: 725)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSSISSGLLSGRSANIGGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 726)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_3007 (SEQ ID NO: 727)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSSAVGLLAPPGGLSGRSDIHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 728)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_3008 (SEQ ID NO: 729)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSSAVGLLAPPGGLSGRSDQHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 730)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_3013 (SEQ ID NO: 731)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSSAVGLLAPPGGLSGRSANIGGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 732)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_2007 (SEQ ID NO: 733)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSSISSGLLSGRSDIHGGGSDIQMTQSPSSLSASVGDRVTIT
CSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQ
RRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSG

NSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNREGC] (SEQ ID NO: 734)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_2008 (SEQ ID NO: 735)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSISSGLLSGRSDQHGGGSDIQMTQSPSSLSASVGDRVITITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNREGC] (SEQ ID NO: 736)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_2013 (SEQ ID NO: 737)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSISSGLLSGRSANIGGGSDIQMTQSPSSLSASVGDRVITITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNREGC] (SEQ ID NO: 738)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_3007 (SEQ ID NO: 739)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSAGVLLAPPGGLSGRSDIHGGSDIQMTQSPSSLSASVGDRVITITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNREGC] (SEQ ID NO: 740)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_3008 (SEQ ID NO: 741)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSAGVLLAPPGGLSGRSDQHGGSDIQMTQSPSSLSASVGDRVITITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNREGC] (SEQ ID NO: 742)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_3013 (SEQ ID NO: 743)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSAVGLLAPPGGLSGRSANIGGSDIQMTQSPSSLSASVGDR
VTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATY
YCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNA
LQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC] (SEQ
ID NO: 744)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_2007 (SEQ ID NO: 745)]

Amino Acid Sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGSISSGLLSGRSDIHGGSDIQMTQSPSSLSASVGDR
VVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 746)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_3007 (SEQ ID NO: 747)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGSAVGLLAPPGGLSGRSDIHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 748)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_2008 (SEQ ID NO: 749)]

Amino Acid Sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGSISSGLLSGRSDQHGGSDIQMTQSPSSLSASVGDR
VVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 750)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_3008 (SEQ ID NO: 751)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS AVGLLAPPGGLSGRSDQHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 752)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_2011 (SEQ ID NO: 753)]

Amino Acid Sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS ISSGLLSGRSDNPGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 754)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_3011 (SEQ ID NO: 755)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS AVGLLAPPGGLSGRSDNPGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 756)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_2012 (SEQ ID NO: 757)]

Amino Acid Sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS ISSGLLSGRSANPGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 758)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_3012 (SEQ ID NO: 759)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS AVGLLAPPGGLSGRSANPGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE

DFATYYCQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 760)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_2013 (SEQ ID NO: 761)]

Amino Acid Sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGSISSGLLSGRSANIGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 762)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_3013 (SEQ ID NO: 763)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGSASVGLLAPPGGLSGRSANIGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
KVDNALQSGNSQESVTEQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 764)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_2001 (SEQ ID NO: 765)]

Amino Acid Sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 766)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_3001 (SEQ ID NO: 767)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSASVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
KVDNALQSGNSQESVTEQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 768)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_2007 (SEQ ID NO: 769)]

Amino Acid Sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSISSGLLSGRSDIHGGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 770)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_3007 (SEQ ID NO: 771)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSAVGLLAPPGGLSGRSDIHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 772)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_2008 (SEQ ID NO: 773)]

Amino Acid Sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSISSGLLSGRSDQHGGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 774)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_3008 (SEQ ID NO: 775)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSAVGLLAPPGGLSGRSDQHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 776)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_2011 (SEQ ID NO: 777)]

Amino Acid Sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSISSGLLSGRSDNPGGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT

YYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 778)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_3011 (SEQ ID NO: 779)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGS AVGLLAPPGGLSGRSDNPGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 780)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_2012 (SEQ ID NO: 781)]

Amino Acid Sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGS ISSGLLSGRSANPGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 782)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_3012 (SEQ ID NO: 783)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGS AVGLLAPPGGLSGRSANPGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 784)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_2013 (SEQ ID NO: 785)]

Amino Acid Sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGS ISSGLLSGRSANIGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 786)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_3013 (SEQ ID NO: 787)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGS AVGLLAPPGGLSGRSANIGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKDSSTLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 788)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_2001 VL domain (SEQ ID NO: 809)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS ISSGLLSGRSDNHGGGSDIQMTQSPSSL
ASVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLT
ISSMQPEDFATYYCQRRNYPYTFGQGTKLEIK] (SEQ ID NO: 810)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_3001 VL domain (SEQ ID NO: 811)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS AVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSL
ASVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIK] (SEQ ID NO: 812)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_2001 VL domain (SEQ ID NO: 813)]

Amino Acid sequence

[QGQSGQG] [NLCTEHS AALDCRSYGGGSSGGS ISSGLLSGRSDNHGGGSDIQMTQSPSSL
SASVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIK] (SEQ ID NO: 814)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_3001 VL domain (SEQ ID NO: 815)]

Amino Acid sequence

[QGQSGQG] [NLCTEHS AALDCRSYGGGSSGGS AVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSL
SASVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIK] (SEQ ID NO: 816)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_2001 VL domain (SEQ ID NO: 817)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSSISGLLSGRSDNHGGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 818)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_3001 VL domain (SEQ ID NO: 819)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSSAVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 820)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_2012 VL domain (SEQ ID NO: 821)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSSISGLLSGRSANPGGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 822)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_3011 VL domain (SEQ ID NO: 823)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSSAVGLLAPPGGLSGRSDNPGGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 824)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_2011 VL domain (SEQ ID NO: 825)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSSISGLLSGRSDNPGGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 826)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_3012 VL domain (SEQ ID NO: 827)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGS AVGLLAPPGGLSGRSANPGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 828)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_2011 VL domain (SEQ ID NO: 829)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGS ISSGLLSGRSDNPGGSDIQMTQSPSSLSASVGDRVTIT
CSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQ
QRNYPYTFGQGTKLEIK] (SEQ ID NO: 830)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_3012 VL domain (SEQ ID NO: 831)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGS AVGLLAPPGGLSGRSANPGGSDIQMTQSPSSLSASVGDR
VTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATY
YCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 832)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_2012 VL domain (SEQ ID NO: 833)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGS ISSGLLSGRSANPGGSDIQMTQSPSSLSASVGDRVTIT
CSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQ
QRNYPYTFGQGTKLEIK] (SEQ ID NO: 834)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_3011 VL domain (SEQ ID NO: 835)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGS AVGLLAPPGGLSGRSDNPGGSDIQMTQSPSSLSASVGDR
VTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATY
YCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 836)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_NSUB VL domain (SEQ ID NO: 837)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSGGSGGSGGSGGSGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 838)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_NSUB VL domain (SEQ ID NO: 839)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSGGSGGSGGSGGSGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 840)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_2007 VL domain (SEQ ID NO: 841)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSISSGLLSGRSDIHGGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 842)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_2008 VL domain (SEQ ID NO: 843)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSISSGLLSGRSDQHGGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 844)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_2013 VL domain (SEQ ID NO: 845)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGSISSGLLSGRSANIGGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 846)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_3007 VL domain (SEQ ID NO: 847)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGS AVGLLAPPGGLSGRSDIHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 848)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_3008 VL domain (SEQ ID NO: 849)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGS AVGLLAPPGGLSGRSDQHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 850)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.13_3013 VL domain (SEQ ID NO: 851)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSAAALDCRSYGGGSSGGS AVGLLAPPGGLSGRSANIGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 852)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_2007 VL domain (SEQ ID NO: 853)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGS ISSGLLSGRSDIHGGSDIQMTQSPSSLSASVGDRVTIT
CSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQ
RRNYPYTFGQGTKLEIK] (SEQ ID NO: 854)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_2008 VL domain (SEQ ID NO: 855)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGS ISSGLLSGRSDQHGGSDIQMTQSPSSLSASVGDRVTIT
CSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQ
RRNYPYTFGQGTKLEIK] (SEQ ID NO: 856)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_2013 VL domain (SEQ ID NO: 857)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSSISGLLSGRSANIGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 858)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_3007 VL domain (SEQ ID NO: 859)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSSAVGLLAPPGGLSGRSDIHGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 860)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_3008 VL domain (SEQ ID NO: 861)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSSAVGLLAPPGGLSGRSDQHGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 862)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02.18_3013 VL domain (SEQ ID NO: 863)]

Amino Acid sequence

[QGQSGQG] [CTEHSFALDCGGGSSGGSSAVGLLAPPGGLSGRSANIGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 864)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_2007 VL domain (SEQ ID NO: 865)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGSSISGLLSGRSDIHGGSDIQMTQSPSSLSASVGDRVTITCSASSSVYYMYWFFQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 866)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_3007 VL domain (SEQ ID NO: 867)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS AVGLLAPP GGLSGRSDIHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIK] (SEQ ID NO: 868)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_2008 VL domain (SEQ ID NO: 869)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS ISSGLLSGRSDQHGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQRRNYPYTFGQGTKLEIK] (SEQ ID NO: 870)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_3008 VL domain (SEQ ID NO: 871)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS AVGLLAPP GGLSGRSDQHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIK] (SEQ ID NO: 872)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_2011 VL domain (SEQ ID NO: 873)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS ISSGLLSGRSDNPGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQRRNYPYTFGQGTKLEIK] (SEQ ID NO: 874)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_3011 VL domain (SEQ ID NO: 875)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS AVGLLAPP GGLSGRSDNPGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIK] (SEQ ID NO: 876)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_2012 VL domain (SEQ ID NO: 877)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGSISSGLLSGRSANPGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 878)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_3012 VL domain (SEQ ID NO: 879)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS AVGLLAPPGGLSGRSANPGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 880)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_2013 VL domain (SEQ ID NO: 881)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGSISSGLLSGRSANIGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 882)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF01_3013 VL domain (SEQ ID NO: 883)]

Amino Acid sequence

[QGQSGQG] [QFCPWSYYLIGDCDIGGGSSGGS AVGLLAPPGGLSGRSANIGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 884)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_2001 VL domain (SEQ ID NO: 885)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 886)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_3001 VL domain (SEQ ID NO: 887)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGS AVGLLAPP GGLSGRSDNHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 888)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_2007 VL domain (SEQ ID NO: 889)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGS ISSGLLSGRSDIHGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 890)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_3007 VL domain (SEQ ID NO: 891)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGS AVGLLAPP GGLSGRSDIHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 892)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_2008 VL domain (SEQ ID NO: 893)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGS ISSGLLSGRSDQHGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 894)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_3008 VL domain (SEQ ID NO: 895)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGS AVGLLAPP GGLSGRSDQHGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 896)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_2011 VL domain (SEQ ID NO: 897)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSISSGLLSGRSDNPGGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 898)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_3011 VL domain (SEQ ID NO: 899)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSAVGLLAPPGGLSGRSDNPGGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 900)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_2012 VL domain (SEQ ID NO: 901)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSISSGLLSGRSANPGGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 902)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_3012 VL domain (SEQ ID NO: 903)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSAVGLLAPPGGLSGRSANPGGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 904)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_2013 VL domain (SEQ ID NO: 905)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGSISSGLLSGRSANIGGGSDIQMTQSPSSLSASVGD
RVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPEDFAT
YYCQQRNYPYTFGQGTKLEIK] (SEQ ID NO: 906)

[spacer (SEQ ID NO: 645)] [huCD71Lc_TF02_3013 VL domain (SEQ ID NO: 907)]

Amino Acid sequence

[QGQSGQG] [NLCTEHSFALDCRSYGGGSSGGS AVGLLAPPGLSGRSANIGGSDIQMTQSPSSLSA
SVGDRVTITCSASSSVYYMYWFQQKPGKAPKLWIYSTSNLASGVPSRFSGSGSGTDYTLTISSMQPE
DFATYYCQRRNYPYTFGQGTKLEIK] (SEQ ID NO: 908)

EXAMPLE 3. Generation and Characterization of Activatable Anti-CD71 Antibodies

[000578] The studies provided herein were designed to generate activatable anti-CD71 antibodies of the disclosure.

[000579] Anti-CD71 activatable antibodies were generated with different masking efficiencies (i.e., a measurement of the ability of the MM of the activatable antibody to block binding of the AB of the activatable antibody to its target). The peptides TF01 and TF02 were mutated by truncation and alanine scanning as described in Example 2, and these masking peptide variants were used to generate families of anti-CD71 activatable antibodies of the present disclosure with a range of masking efficiencies.

[000580] Binding of anti-CD71 activatable antibodies of the present disclosure to the NCI H292 (also referred to herein as H292) cell line was evaluated using FACS. Briefly, cells were labeled with huCD71 antibody or activatable antibody at the indicated concentrations and subsequently detected with an Alexa Fluor 647 labeled goat anti-human IgG secondary antibody. As shown in Figure 2, anti-CD71 activatable antibodies of the present disclosure show a range of masking efficiencies compared to the parental anti-CD71 antibody (huCD71 21.12 Ab).

[000581] As shown in Figure 3A, huCD71 activatable antibody drug conjugates of the present disclosure behave similarly to un-conjugated activatable antibodies and recover huCD71 antibody binding activity upon proteolytic activation. This study was done using the activatable anti-CD71 antibody referred to herein as huCD71 TF01-2001 conjugated to MMAD via a val-cit linker (the linker-toxin is referred to herein as vc-MMAD), and this activatable antibody construct includes the VH of SEQ ID NO: 5, the VL of SEQ ID NO: 7, the MM of SEQ ID NO: 16, and the CM comprising the sequence ISSGLLSGRSDNH (SEQ ID NO: 406). The binding of uPA treated or intact huCD71 TF01-2001 activatable antibody -MMAD conjugated to H292 cells was evaluated by FACS using a standard FACS labeling protocol. An anti-CD71 activatable antibody drug conjugate, also referred to herein as "AADC," was activated by an overnight incubation with uPA. Figure 3B shows the

ability of activated anti-CD71 activatable antibody huCD71 TF01-2001 conjugated to MMAD via a val-cit linker (the linker-toxin is referred to herein as v-MMAD or vc-MMAD), to kill cells *in vitro* in a manner similar to anti-CD71 antibody 21.12-MMAD. Similar results were observed when an activatable antibody of the embodiments conjugated to a nucleic acid damaging agent was tested in such cell killing assays.

[000582] H292 xenograft tumors treated with isotype-DM4 control, huCD71-DM4 ADC, and the conjugated anti-CD71 activatable antibody huCD71 TF02.13_2001-DM4 AADC, which has the VH of SEQ ID NO: 5, the VL of SEQ ID NO: 7, the MM of SEQ ID NO: 309 (NLCTEHSALDCRSY), and the CM comprising the sequence ISSGLLSGRSDNH (SEQ ID NO: 406). Tumors were grown to an average of 150 mm³; then the mice were randomized into groups of eight and dosed on day 0 and 7 with the indicated test articles. The results of this study are shown in Figure 4. Mean tumor volume $\pm$ SEM for each group is plotted. Both the ADC and AADC induce tumor regressions. All DM4-conjugated activatable antibodies disclosed herein were produced by TCRS (The Chemistry Research Solution).

[000583] Figure 5 is a graph depicting the results of breast cancer HCC1806 xenograft tumors treated with PBS as a control and with two different doses, 1 mg/kg and 3 mg/kg, of the conjugated activatable anti-CD71 antibody huCD71 TF02.13-2001-DM4 AADC, which has the VH of SEQ ID NO: 5, the VL of SEQ ID NO: 7, the MM of SEQ ID NO: 309 (NLCTEHSALDCRSY), and the CM1 comprising the sequence ISSGLLSGRSDNH (SEQ ID NO: 406). Mean tumor volume $\pm$ SEM for each group is plotted. Figure 5 demonstrates that the conjugated activatable anti-CD71 antibodies of the disclosure induce complete tumor regressions in breast HCC1806 xenograft tumors at lower than clinically relevant doses.

[000584] Figure 6A is a graph depicting the ability of the anti-CD71 antibody of the present disclosure (VH of SEQ ID NO: 5 and the VL of SEQ ID NO: 7) to bind cynomolgus monkey primary kidney epithelial cells, with an observed K_d of about 3 nM in this exemplary study. In this study, the binding of the antibody of the present disclosure to the cells were performed using a standard FACS labelling method. Briefly, cells were labeled with the anti-CD71 antibody or isotype (palivizumab) control of the present disclosure at the indicated concentrations and subsequently detected with an Alexa Fluor 647 labeled goat anti-human IgG secondary antibody.

[000585] Figure 6B demonstrates that the anti-CD71 antibody huCD71 (VH of SEQ ID NO: 5 and the VL of SEQ ID NO: 7) binds to human and cynomolgus monkey CD71 but does not bind to mouse CD71. In this assay, recombinant CD71 of the indicated type (1 µg/mL) was absorbed to ELISA plates and subsequently incubated with the anti-human CD71 21.12 antibody at the indicated concentration, followed and subsequently detected with goat anti-human IgG-HRP secondary antibody and Ultra TMB (Thermo Fisher Scientific) detection using an assay and in accordance with methods known to those skilled in the art.

[000586] Figure 6C is a graph depicting the ability of the murine anti-CD71 antibody muM21 (“CD71 ms Ab”) of the present disclosure (VH of SEQ ID NO: 1 and the VL of SEQ ID NO: 2) and the humanized anti-CD71 antibody Ab21.12 (“CD71 Hu Ab”) of the present disclosure (VH of SEQ ID NO: 5 and the VL of SEQ ID NO: 7) to bind the human pancreatic cancer-derived BxPC3 cell line, both with an observed K_d of about 3 nM in this exemplary study. In this study, the binding of the antibodies of the present disclosure to the cells were performed using a standard FACS labelling method. Briefly, cells were labeled with the anti-CD71 antibody or human IgG1 isotype control antibody (palivizumab, “Isotype”) of the present disclosure at the indicated concentrations and subsequently detected with an Alexa Fluor 647 labeled goat anti-human or goat anti-mouse IgG secondary antibody.

[000587] Figure 7 is a graph depicting that a single dose of an anti-CD71 activatable antibody of the present disclosure demonstrates a prolonged half-life compared to the CD71 parental antibody when administered to cynomolgus monkeys. In this study, monkeys were injected with a single dose on day 0 at 5 mg/kg of the indicated antibody, where the “CD71 antibody” was anti-human CD71 21.12 antibody of the present disclosure, and the “CD71 activatable antibody” was anti-human CD71 TF01.2001 activatable antibody of the present disclosure. The amount of total human IgG in the serum of each monkey was assayed on the indicated days using an anti-human IgG sandwich ELISA.

[000588] The pharmacokinetics and tolerability of the huCD71 TF02.13-2001 DM4 drug conjugate in two cynomolgus monkeys was evaluated after a single 5 mg/kg dose (Figure 8). Total serum levels of human IgG were measured using an anti-human IgG sandwich ELISA.

[000589] The studies depicted in Figures 7, 8, and 9 were conducted as part of a 3 week tolerability study (non-terminal); single dose n=2. The prolonged PK of intact

activatable antibody that was observed was consistent with a decrease in normal tissue binding. Low circulating levels of activated conjugated activatable antibody were detected. The conjugated activatable antibody was well tolerated at 5 mg/mg. No evidence of on or off-target toxicity was observed. There were no clinical signs, weight loss, clinical chemistry or hematologic findings.

[000590] Figure 9 shows that two cynomolgus monkeys that received the conjugated activatable anti-CD71 antibody of the disclosure (anti-huCD71 TF02.13-2001 DM4) showed no evidence of neutropenia (based on neutrophil counts) or liver toxicity (based on levels of alanine transaminase (ALT) or aspartate transaminase (AST)).

EXAMPLE 4: Characterization of the Binding and Inhibitory Activity of the CD71

Antibody

[000591] This Example shows the ability of an anti-CD71 antibody of the present disclosure to inhibit binding of recombinant CD71 to its ligand, transferrin.

[000592] Figure 10 shows the binding of recombinant human transferrin (Tf) ligand to recombinant human CD71 in the absence of anti-CD71 antibody using a solid-phase ELISA binding assay. In this study, a dissociation constant (K_d) of about 1.6 nM of transferrin for human CD71 was determined in this example. In this assay, recombinant human CD71 (1 μ g/ml) was absorbed to ELISA plates and subsequently incubated with the indicated concentration of biotinylated transferrin ligand, followed by detection of bound biotinylated transferrin using streptavidin-horse radish peroxidase (HRP) and Ultra TMB detection (Thermo Fisher Scientific) using an assay and in accordance with methods known to those skilled in the art.

[000593] Figure 11A shows the effect on binding between recombinant human transferrin (Tf) ligand and recombinant human CD71 under various conditions using a solid-phase ELISA binding assay, including binding in the presence of an anti-CD71 antibody of the present disclosure. In particular as shown in Figure 11A, this study showed that a humanized anti-CD71 antibody of the present disclosure ("CD71 Ab21.12") inhibited the binding between the transferrin ligand and its CD71 receptor as compared to an isotype control. The results also showed that this inhibition was similar to the binding inhibition demonstrated by a different, commercially-available mouse anti-human CD71 monoclonal antibody ("OKT9"). The results also showed that neither a commercially-available rabbit anti-human CD71 polyclonal antibody (LS-C334364, LS Bio) nor an isotype control

significantly inhibited transferrin binding to CD71, and that binding activity was abolished by the addition of an excess amount of cold transferrin. In this assay, recombinant human CD71 protein (1 µg/ml) was absorbed to ELISA plates and subsequently incubated with 0.6 nM of biotinylated-transferrin in the presence of the indicated concentrations of antibody or transferrin (where the anti-CD71 21.12 of the present disclosure having VH of SEQ ID NO: 5 and VL of SEQ ID NO: 7) as part of a titration assay. The bound transferrin was detected with streptavidin-conjugated HRP and Ultra TMB detection using an assay and in accordance with methods known to those skilled in the art.

[000594] Figure 11B shows the extent of inhibition of transferrin binding to CD71 by a known anti-CD71 monoclonal antibody (OKT9) is significantly less than that of the extent of inhibition by anti-CD71 21.12 antibody of the present disclosure. In this assay, recombinant human CD71 protein (1 µg/ml) was absorbed to ELISA plates and subsequently incubated with 0.6 nM of biotinylated-transferrin in the presence of the indicated concentrations of OKT9 anti-CD71 antibody or anti-CD71 21.12 antibody of the present disclosure (anti-CD71 21.12 having VH of SEQ ID NO: 5 and VL of SEQ ID NO: 7) as part of a titration assay. The bound transferrin was detected with streptavidin-conjugated HRP and Ultra TMB detection using an assay and in accordance with methods known to those skilled in the art.

EXAMPLE 5: CD71 Expression in Multiple Primary and Metastatic Tumors

[000595] This Example shows that CD71 is expressed in a large variety of primary and metastatic tumor types by immunohistochemical (IHC) staining using an anti-CD71 antibody.

[000596] Figures 12A, 12B, and 12C show that CD71 is highly expressed in a large number of primary and metastatic tumor samples, using IHC staining with a commercially-purchased anti-CD71 antibody on multiple primary tumors and metastatic tissue microarrays (TMA). Figure 12A shows an IHC staining of CD71 in cervical cancer (1), head and neck cancer (2), an H292 xenograft (3), non-Hodgkin's lymphoma (NHL) in lymph nodes (4), NHL in colon (5), and NHL in stomach (6). Figure 12B shows an IHC staining of CD71 in a TMA consisting of cores from metastatic tumors demonstrated a moderate to high level of expression of CD71 in the majority of the cores. Figure 12C shows a summary of the level of IHC staining of CD71 of one exemplary TMA shows that a large number of metastasis cores derived from multiple tissue sources showed a strong CD71 signal.

EXAMPLE 6: Activatable Anti-CD71-AADC *in vivo* Efficacy in Raji Xenograft Model

[000597] This Example shows that anti-human CD71 activatable antibodies with conjugated toxins (AADCs) of the present disclosure are efficacious in a non-Hodgkin's lymphoma (NHL) mouse xenograft model. These efficacies are specific to anti-human CD71 antibodies and are comparable or equivalent to the efficacy demonstrated by a parental anti-human CD71 drug conjugate.

[000598] Figure 13A shows the efficacy of various anti-CD71 activatable antibody drug conjugates in a mouse NHL xenograft model, where the activatable antibody drug conjugates of the present disclosure (CD71 TF02.18-2001-spdb-DM4, CD71 TF02.13-2001-spdb-DM4, and CD71 TF02.13.3001-spdb-DM4) showed significantly higher efficacy than an isotype control (palivizumab-spdb-DM4). The figure also shows that the activatable antibody drug conjugates of the present disclosure demonstrated an efficacy comparable to an unmasked anti-CD71 antibody drug conjugate (CD71 21.12-spdb-DM4). In this study, non-Hodgkin's lymphoma (NHL) Raji xenograft tumors in mice were grown to an average volume of 150 mm³. The mice were then randomized into groups of eight and dosed on days 1 and 8 with 5 mg/kg of each indicated test article. The mean tumor volume $\pm$ SEM was plotted for each time point.

[000599] Figure 13B shows the efficacy of various anti-CD71 activatable antibody drug conjugates in a mouse NHL xenograft model, where the activatable antibody drug conjugates of the present disclosure (CD71 TF02.18-2001-vc-MMAE and CD71 TF02.13-3001-vc-MMAE) showed significantly higher efficacy than an isotype control (palivizumab-vc-MMAE). The figure also shows that the activatable antibody drug conjugates of the present disclosure demonstrated an efficacy comparable to an unmasked anti-CD71 antibody drug conjugate (CD71 21.12-vc-MMAE). In this study, non-Hodgkin's lymphoma (NHL) Raji xenograft tumors in mice were grown to an average volume of 150 mm³. The mice were then randomized into groups of eight and dosed on days 1 and 8 with 3 mg/kg of each indicated test article. The mean tumor volume $\pm$ SEM was plotted for each time point.

[000600] Figure 13C shows the efficacy of various anti-CD71 activatable antibody drug conjugates in a mouse NHL xenograft model, where the activatable antibody drug conjugates of the present disclosure (CD71 TF02.18-2001-vc-MMAD and CD71 TF02.13-3001-vc-MMAD) showed significantly higher efficacy than an isotype control

(palivizumab-vc-MMAD). The figure also shows that the activatable antibody drug conjugates of the present disclosure demonstrated an efficacy comparable to an unmasked anti-CD71 antibody drug conjugate (CD71 21.12-vc-MMAD). In this study, non-Hodgkin's lymphoma (NHL) Raji xenograft tumors in mice were grown to an average volume of 150 mm³. The mice were then randomized into groups of eight and dosed on days 1 and 8 with 0.5 or 1 mg/kg of each indicated test article. The mean tumor volume $\pm$ SEM was plotted for each time point.

EXAMPLE 7: Cynomolgus Tolerability of Anti-CD71-ADC and Activatable-Anti-CD71-AADC

[000601] This Example shows that activatable anti-human CD71 antibodies with conjugated toxins (AADCs) of the present disclosure are well-tolerated in cynomolgus monkeys compared to the corresponding parental anti-CD71 ADCs based on one or more hematology readouts.

[000602] Figures 14A-14G show that anti-CD71 activatable antibody drug conjugates (AADCs) of the present disclosure demonstrate a higher tolerance in cynomolgus monkeys based on an absence of marked changes in each hematology readout, as compared to monkeys treated with high or low doses of the corresponding parental anti-CD71 ADC. The following hematology results were observed upon treatment of the monkeys with 5 mg/kg of the parental anti-CD71 ADC: a marked reduction in total white blood cells (WBC) with high dose ADC treatment (Figure 14A), severe neutropenia by days 8 to 11 with high and low dose ADC treatment (Figure 14B), a marked reduction in lymphocytes with high dose ADC treatment (Figure 14C), decreased monocytes with both high and low dose ADC treatment, with monocyte recovery in the low-dose-treated animal on day 11 (Figure 14D), decreased red blood cell count with both high and low dose ADC treatment (RBC) (Figure 14E), a marked reduction of hemoglobin count (HGB) with both high and low dose ADC treatment (Figure 14F), and a marked reduction of reticulocytes with both high and low dose ADC treatment (Figure 14G). Monkeys treated with anti-CD71 ADC showed a normal platelet count (data not shown). In comparison, animals treated with activatable anti-CD71 AADCs of the present disclosure showed no marked changes in all of these hematological readouts (Figures 14A-14F). In this study, a single dose of activatable anti-CD71 TF02.13-2001-spdb-DM4 ("AADC") of the present disclosure having a drug-antibody ratio (DAR) of about 3.4 was dosed in cynomolgus monkeys at 5 mg/kg and anti-CD71 21.12-spdb-

DM4 (“ADC”) of the present disclosure was dosed at 5 mg/kg (low dose) and 7.5 mg/kg (high dose). The hematological results were obtained from each monkey at the indicated days of the study using techniques and methods known in the art.

EXAMPLE 8: CD71 Expression and Sensitivity to Anti-CD71-Mediated Cytotoxicity In Multiple Cell Lines

[000603] This Example shows that CD71 is expressed in high levels in many tumor-derived cell lines, and that many of these cell lines demonstrated sensitivity to anti-CD71-targeted cytotoxicity.

[000604] Figure 15 shows the maximum binding or relative expression of CD71 in the indicated cell lines by FACS analysis. FACS staining was carried out using a commercially available anti-CD71 mouse monoclonal antibody, OKT9, followed by an Alexa Fluor 647 conjugated goat anti mouse secondary antibody, with the height of the bar for a given cell line corresponding to the relative amount of CD71-derived signal for that cell line. The color of the bar for a given cell line shows the relative sensitivity of the corresponding cell line in an *in vitro* cytotoxicity assay where the cell line was treated with anti-CD71 21.12 antibody of the present disclosure in the presence of an anti-human secondary antibody conjugated to MMAE toxin. A cell line was categorized as “sensitive” if cytotoxicity was observed with a less than 1 nM EC50. A cell line was categorized as “moderately sensitive” if maximum killing was less than 50% and EC50 was greater than 1 nM. A cell line was categorized as “resistant” if no to little cytotoxicity was observed.

EXAMPLE 9: CD71 Expression on HT29, BxPc3, Fadu and MDA MB 231 Cell Lines

[000605] This Example shows that activatable anti-human CD71 antibodies of the present disclosure bind CD71 on multiple cell lines with a higher dissociation constant than that of the unmasked anti-human CD7 antibody of the present disclosure, thus showing the effect of the mask in reducing binding prior to activation.

[000606] Figures 16A to 16D shows the amount of binding of anti-CD71 activatable and parental antibodies of the present disclosure to the cell lines HT29 (Figure 16A), BxPc3 (Figure 16B), FaDu (Figure 16C), and MDA MB 231 (Figure 16D). In this study, the binding of the antibodies of the present disclosure to the indicated cell lines were performed using a standard FACS labelling method. Briefly, cells were labeled with the indicated antibodies of the present disclosure: anti-human CD71 antibody (anti-human CD71

Ab21.12 antibody, “CD71-Ab”) or one of two anti-human CD71 activatable antibodies (anti-human CD71 TF02.13-3001, “CD71-ActAb 1,” or anti-human CD71 TF02.18-2001, “CD71-ActAb 2”) at the indicated concentrations and subsequently detected with an Alexa Fluor 647 labeled goat anti-human IgG secondary antibody. Table 14 below shows the equilibrium dissociation constants based on the binding curves depicted in Figures 14A to 14D. These results show that anti-human CD71 21.12 antibody (CD71-Ab) binds all cell lines with similar K_d (0.39 to 0.81 nM), while the binding of anti-human CD71 TF02.13-3001 (CD71-ActAb 1) and anti-human CD71 TF02.18-2001 (CD71-ActAb 2) to the cell lines were significantly shifted to the right (38 to 46 fold), which is indicative of the masking efficiency of the masking moiety.

Table 14: Exemplary Observed CD71 Binding Activity of Activatable Anti-CD71

Cell Line	CD71-Ab K _d (nM)	CD71-ActAb 1 K _d (nM)	ME*	CD71-ActAb 2 K _d (nM)	ME*	B _{max} (MFI)
FaDu	0.39	14.74	38	15.36	39	1000
HT29	0.51	23.02	45	19.74	39	700
MB231	0.84	38.84	46	31.99	38	500
BxPC3	0.81	35.18	43	36.35	43	400

*ME = masking efficiency

EXAMPLE 10: Anti-CD71 ADC Cytotoxicity on HT29, BxPc3, FaDu and MDA MB

231 Cell Lines

[000607] This Example shows that anti-human CD71 antibody drug conjugates of the present disclosure demonstrate a higher cytotoxicity against multiple cell lines compared to an isotype control ADC.

[000608] Figures 17A to 17D show that anti-human CD71 antibody of the present disclosure conjugated to various toxins demonstrate higher *in vitro* cytotoxicity against various CD71-expressing cell lines as compared to an isotype control ADC. In this study, anti-human CD71 21.12-ADC of the present disclosure (conjugated to either spdb-DM4, vc-MMAE or vc-MMAD) or an isotype control (palivizumab-ADC, conjugated to either spdb-DM4, vc-MMAE or vc-MMAD) as applied at the indicated concentrations to HT29 (Figure 17A), BxPc3 (Figure 17B), FaDu (Figure 17C), and MDA MB 231 (Figure 17D). The cytotoxicity was determined as a percentage of a population of untreated cells that were used as a control.

EXAMPLE 11: Activatable Anti-CD71 Cytotoxicity on NHL Raji Cells

[000609] This Example shows that anti-human CD71 antibody drug conjugates (ADCs) of the present disclosure demonstrate a higher cytotoxicity against a Raji (non-Hodgkin's lymphoma) cell line as compared to anti-human CD71 activatable antibody drug conjugates (AADCs) of the present disclosure.

[000610] Figure 18 shows that anti-human CD71 21.12-ADC drug conjugates (ADCs) of the present disclosure (conjugated to either spdb-DM4, vc-MMAE or vc-MMAD) showed a higher cytotoxicity against Raji non-Hodgkin's lymphoma-derived cells than that of either their corresponding anti-human CD71 TF02.18-2001-AADC activatable antibody drug conjugates (conjugated to either spdb-DM4, vc-MMAE or vc-MMAD) or anti-human CD71 TF02.13-3001-AADC activatable antibody drug conjugates (conjugated to either spdb-DM4, vc-MMAE or vc-MMAD). The anti-human CD71 ADCs demonstrated a similar cytotoxic efficacy to each other (showing an IC₅₀ of ~ 0.1 nM based on comparison to a control sample) against the Raji cell line. In comparison, the anti-human CD71 activatable antibody drug conjugate (AADC) showed a cytotoxic effect that is similar or comparable to that of the isotype-ADC control. In this study, the various anti-human CD71 antibody drug conjugates and activatable antibody drug conjugates of the present disclosure were applied at the indicated concentrations to Raji cells. The cytotoxicity was determined as a percentage of a population of untreated cells that were used as a control.

EXAMPLE 12: CD71 Binding Assay

[000611] This Example show that anti-human CD71 activatable antibodies of the present disclosure demonstrated a shifted binding affinity to recombinant CD71 protein compared to the parental anti-CD71 antibody of the present disclosure.

[000612] As shown by the examples depicted in Figure 19, a solid-phase binding assay was used to demonstrate the binding of anti-human CD71 antibodies of the present disclosure. In these examples, recombinant human CD71 protein (R&D Systems) was coated on nickel-plated ELISA plates at a concentration of 1 µg/mL, and then incubated with the indicated concentration of anti-CD71 antibody ("CD71 21.12") or activatable anti-CD71 antibodies ("CD71.TF02.13-2001" or "CD71.TF02.18-2001"), where in the activatable antibodies were assayed in their non-proteolytically-activated form. The amount of bound antibody was detected by incubation and detection by goat anti-human antibody conjugated to horseradish peroxidase and Ultra TMB (Thermo Fisher Scientific) detection.

[000613] As depicted in Figure 19, the exemplary assay demonstrated that the anti-human CD71 21.12 antibody bound with a K_d of ~ 0.3 nM. The binding constants of anti-human CD71 TF02.18-2001 and human CD71 TF02.13-2001 activatable antibodies are shifted to the right as a result of their respective masking efficiencies (9 and 42 fold respectively).

Table 15: Exemplary Observed CD71 Binding Activity of Activatable Anti-CD71 Antibodies

Antibody	K_d (nM) 1 μ g/mL CD71 antigen	ME*
CD71 21.12	0.3	-
CD71 TF02.13-2001	13	42
CD71 TF02.18-2001	3.0	9

*ME = masking efficiency

EXAMPLE 13: Activatable Anti-CD71-AADC *in vivo* Efficacy in a Patient-Derived Xenograft Model of Non-Hodgkin's Lymphoma

[000614] This Example shows that an anti-human CD71 activatable antibody with conjugated toxins of the present disclosure (AADC) is efficacious in a non-Hodgkin's lymphoma (NHL) patient-derived xenograft model, and that this efficacy is comparable to or better than that observed with a parental anti-human CD71 antibody drug conjugate (ADC).

[000615] Figures 20A and 20B show that using a patient-derived xenograft mouse model of non-Hodgkin's lymphoma (NHL) primary tumor, mice treated with an activatable anti-huCD71-AADC of the present disclosure showed over time efficacies that were similar to or higher than that of their respective anti-huCD71-ADC of the present disclosure. In this study, two patient-derived NHL xenografts (labeled as LY2214 and LY0257, Crown Bioscience) were each treated with an isotype-ADC control (palivizumab conjugated to spdb-DM4, "Isotype-ADC"); activatable anti-huCD71 21.12-ADC of the present disclosure conjugated to spdb-DM4 ("CD71-ADC") of the present disclosure; or activatable anti-huCD71 TF02.13.3001 of the present disclosure conjugated to spdb-DM4 ("CD71-

AADC”). In each of the examples, tumor xenografts were grown to an average volume of 150 mm³; then the mice were randomized into groups of 4 and dosed on days 1 and 7 with 5 mg/kg of the indicated test articles. The mean tumor volume $\pm$ SEM is plotted for each time point.

EXAMPLE 14: Anti-CD71-ADC Cytotoxicity on Multiple Colorectal Cancer-Derived Cell Lines

[000616] This Example shows that anti-human CD71 antibody drug conjugates (ADCs) of the present disclosure demonstrate a higher cytotoxicity against multiple colorectal cancer (CRC)-derived cell lines as compared to isotype-ADC controls.

[000617] Figures 21A – 21H show that anti-human CD71 21.12 antibody drug conjugates (ADCs) of the present disclosure conjugated to either spdb-DM4 or vc-MMAE showed a higher cytotoxicity against a variety of CRC-derived cell lines than that of an isotype antibody (chKTI, a chimeric human IgG1 anti-soybean trypsin inhibitor antibody) conjugated to either spdb-DM4 or vc-MMAE. The CRC-derived cell lines that were treated with the ADCs were SW1417, SW48, Ls411N, Lovo, HCT116, DLD1, Ls174T, and SW480. The anti-human CD71 ADCs demonstrated a similar cytotoxic efficacy to each other against the cell lines. In this study, the various anti-human CD71 antibody drug conjugates of the present disclosure and isotype drug conjugates were applied at the indicated concentrations to the cells. The cytotoxicity was determined as a percentage of a population of untreated cells that were used as a control.

[000618] Figures 22A and 22B shows that anti-human CD71 21.12-ADC drug conjugates (ADCs) of the present disclosure conjugated to vc-MMAE or vc-MMAD (CD71 21.12 -vc-MMAE, CD71 21.12 -vc-MMAD) showed a higher cytotoxicity against a HT29 CRC-derived cells than that of either their corresponding anti-human CD71 TF02.18-2001-AADC activatable antibody drug conjugate conjugated to vc-MMAE or vc-MMAD (CD71 TF02.18-2001-vc-MMAE, CD71 TF02.18-2001-vc-MMAD), an anti-human CD71 TF02.13-3001-AADC activatable antibody drug conjugate conjugated to vc-MMAE or vc-MMAD (CD71 TF02.13-3001-vc-MMAE, CD71 TF02.13-3001-vc-MMAD), or an isotype control (palivizumab-ADC) conjugated to vc-MMAE or vc-MMAD (Isotype-vc-MMAE, Isotype-vc-MMAD). In this study, the various anti-human CD71 antibody drug conjugates and activatable antibody drug conjugates of the present disclosure, as well as the isotype-

ADC, were applied at the indicated concentrations to HT29 cells. The cytotoxicity was determined as a percentage of a population of untreated cells that were used as a control.

EXAMPLE 15: Activatable Anti-CD71-AADC *in vivo* Efficacy in a Xenograft Model of Non-Hodgkin's Lymphoma

[000619] This Example shows that anti-human CD71 activatable antibodies with conjugated toxins (AADCs) of the present disclosure are efficacious in a non-Hodgkin's lymphoma (NHL) mouse xenograft model. These efficacies are specific to anti-human CD71 antibodies and are comparable or equivalent to the efficacy demonstrated by a parental anti-human CD71 drug conjugates.

[000620] Figure 23A shows the efficacy of various anti-CD71 activatable antibody drug conjugates in a mouse NHL (Raji) xenograft model, where the activatable antibody drug conjugates of the present disclosure (CD71 TF02.18-2001-spdb-DM4, CD71 TF02.13-2001-spdb-DM4, and CD71 TF02.13.3001-spdb-DM4) showed significantly higher efficacy than an isotype control (palivizumab-spdb-DM4). The figure also shows that the activatable antibody drug conjugates of the present disclosure demonstrated an efficacy comparable to an unmasked anti-CD71 antibody drug conjugate (CD71 21.12-spdb-DM4). In this study, non-Hodgkin's lymphoma (NHL) Raji xenograft tumors in mice were grown to an average volume of 150 mm³. The mice were then randomized into groups of eight and dosed on days 1 and 8 with 5 mg/kg of each indicated test article. The mean tumor volume $\pm$ SEM was plotted for each time point.

[000621] Figure 23B shows the efficacy of various anti-CD71 activatable antibody drug conjugates in a mouse NHL (Raji) xenograft model, where the activatable antibody drug conjugates of the present disclosure (CD71 TF02.18-2001-vc-MMAE and CD71 TF02.13-3001-vc-MMAE) showed significantly higher efficacy than an isotype control (palivizumab-vc-MMAE). The figure also shows that the activatable antibody drug conjugates of the present disclosure demonstrated an efficacy comparable to an unmasked anti-CD71 antibody drug conjugate (CD71 21.12-vc-MMAE). In this study, non-Hodgkin's lymphoma (NHL) Raji xenograft tumors in mice were grown to an average volume of 150 mm³. The mice were then randomized into groups of eight and dosed on days 1 and 8 with 3 mg/kg of each indicated test article. The mean tumor volume $\pm$ SEM was plotted for each time point.

EXAMPLE 16: Activatable Anti-CD71-AADC *in vivo* Efficacy in a Xenograft Model of Non-Small Cell Lung Carcinoma

[000622] This Example shows that anti-human CD71 activatable antibodies with conjugated toxins (AADCs) of the present disclosure are efficacious in a non-small cell lung carcinoma (NSCLC) mouse xenograft model over a range of dosages. These efficacies are specific to anti-human CD71 antibodies and are comparable or equivalent to the efficacy demonstrated by a parental anti-human CD71 drug conjugate.

[000623] Figures 24A and 24B show the efficacy of anti-CD71 activatable antibody drug conjugates in a mouse NSCLC xenograft model, where the activatable antibody drug conjugates of the present disclosure (CD71 TF02.13-3001-spdb-DM4, CD71 TF02.13-2001-spdb-DM4, CD71 TF01-2001-PEG2-vc-MMAD) at the indicated dosages (5 mg/kg, 3 mg/kg, and 1 mg/kg) showed significantly higher efficacy than an isotype control (palivizumab-spdb-DM4 or palivizumab-PEG2-vc-MMAD) administered at a 5 mg/kg dosage. The figure also shows that the activatable antibody drug conjugates of the present disclosure demonstrated an efficacy comparable to an unmasked anti-CD71 antibody drug conjugate (CD71 21.12-spdb-DM4 or CD71 21.12-PEG2-vc-MMAD) at 5 mg/kg dosage. In this study, NSCLC H292 xenograft tumors in mice were grown to an average volume of 150 mm³. The mice were then randomized into groups of eight and dosed on days 1 and 8 with the indicated amount of each indicated test article. The mean tumor volume $\pm$ SEM was plotted for each time point.

EXAMPLE 17: Activatable Anti-CD71-AADC *in vivo* Efficacy in a Xenograft Model of Pancreatic Adenocarcinoma

[000624] This Example shows that anti-human CD71 activatable antibodies with conjugated toxins (AADCs) of the present disclosure are efficacious in a BxPC3 pancreatic adenocarcinoma mouse xenograft model over a range of dosages. These efficacies are specific to anti-human CD71 antibodies and are comparable or equivalent to the efficacy demonstrated by a parental anti-human CD71 drug conjugate, and/or higher than the efficacy of an isotype control ADC.

[000625] Figure 25 shows the efficacy of various anti-CD71 activatable antibody drug conjugates in a BxPC3 mouse xenograft model, where the activatable antibody drug conjugates of the present disclosure (CD71 TF02.13-2001-spdb-DM4, CD71 TF02.13-3001-spdb-DM4, and CD71 TF02.13-3001-vc-MMAE) each at the indicated dosages

showed significantly higher efficacy than an isotype control (palivizumab-spdb-DM4 at 5 mg/kg and palivizumab-vc-MMAE at 3 mg/kg). The figure also shows that the activatable antibody drug conjugates of the present disclosure demonstrated an efficacy comparable to an unmasked anti-CD71 antibody drug conjugate (CD71 21.12-spdb-DM4 at 5 mg/kg and CD71 21.12-vc-MMAE at 3 mg/kg). In this study, BxPC3 xenograft tumors in mice were grown to an average volume of 150 mm³. The mice were then randomized into groups of eight and dosed on day 1 with the indicated amount of each indicated test article. The mean tumor volume $\pm$ SEM was plotted for each time point.

EXAMPLE 18: Activatable Anti-CD71-AADC *in vivo* Efficacy in a Patient-Derived Xenograft Model of Non-Hodgkin's Lymphoma

[000626] This Example shows that an anti-human CD71 activatable antibody with conjugated toxins of the present disclosure (AADC) is efficacious in a non-Hodgkin's lymphoma (NHL) patient-derived xenograft model, and that this efficacy is comparable to or better than that observed with a parental anti-human CD71 antibody drug conjugate (ADC).

[000627] Figures 26A and 26B show that using a patient-derived xenograft mouse model of non-Hodgkin's lymphoma (NHL) primary tumor, mice treated with an activatable anti-hu CD71-AADC (CD71 TF02.13-3001-spdb-DM4 and CD71 TF02.13-3001-vc-MMAD) of the present disclosure showed over time efficacies that were similar to or higher than that of their respective anti-huCD71-ADC of the present disclosure. In this study, two patient-derived NHL xenografts (labeled as LY2214 and LY0257, Crown Bioscience) were each treated with a palivizumab isotype-ADC control conjugated to spdb-DM4 ("Isotype-spdb-DM4") or vc-MMAD ("Isotype-vc-MMAD"); antibody drug conjugates of the present disclosure anti-huCD71 21.12-spdb-DM4 or anti-huCD71 21.12-vc-MMAD; or activatable antibodies of the present disclosure anti-hu CD71 TF02.13-3001-spdb-DM4 and anti-huCD71 TF02.13-3001-vc-MMAD. In each of the examples, tumor xenografts were grown to an average volume of 150 mm³; then the mice were randomized into groups of 4 and dosed on days 1 and 7 of the indicated test articles at the indicated dosages. The mean tumor volume $\pm$ SEM is plotted for each time point.

EXAMPLE 19: Activatable Anti-CD71-AADC *in vivo* Efficacy in a Patient-Derived Xenograft Models

[000628] This Example shows that an anti-human CD71 activatable antibody with conjugated toxins of the present disclosure (AADC) is efficacious in various patient-derived xenograft models, and that this efficacy is higher than that of an isotype control ADC.

[000629] Figure 27A shows that using a patient-derived xenograft mouse model of a lung tumor (CTG-0166), mice treated with an activatable anti-hu CD71-AADC (CD71 TF02.13-3001-spdb-DM4) of the present disclosure showed over time efficacies that were higher than that of their respective isotype-ADC control (chKTI, a chimeric human IgG1 anti-soybean trypsin inhibitor antibody, conjugated to spdb-DM4, “Isotype-spdb-DM4”). Figure 27B shows that using a patient-derived xenograft mouse model of a endometrial tumor (CTG-0774), mice treated with an activatable anti-hu CD71-AADC (CD71 TF02.13-3001-spdb-DM4) of the present disclosure showed over time efficacies that were higher than that of their respective isotype-ADC control (chKTI conjugated to spdb-DM4, “Isotype-spdb-DM4”). In another study using a patient-derived xenograft mouse model of a cholangiocarcinoma tumor (bile duct cancer) (CTG-1941), mice treated with an activatable anti-hu CD71-AADC (CD71 TF02.13-3001-vc-MMAE) of the present disclosure showed over time efficacies that were higher than that of an isotype-ADC control (chKTI, a chimeric human IgG1 anti-soybean trypsin inhibitor antibody, conjugated to spdb-DM4). In each of the examples, tumor xenografts were grown to an average volume of 150 mm³; then the mice were randomized into groups of 4 and dosed on day 1 with the indicated amount for each of the indicated test articles. The mean tumor volume $\pm$ SEM is plotted for each time point.

EXAMPLE 20: CD71 Binding Assay

[000630] This Example show that anti-human CD71 activatable antibodies of the present disclosure demonstrated a shifted binding affinity to recombinant CD71 protein compared to the parental anti-CD71 antibody of the present disclosure.

[000631] As shown by the examples depicted in Figure 28, a solid-phase binding assay was used to demonstrate the binding of anti-human CD71 antibodies of the present disclosure. In these examples, recombinant human CD71 protein (R&D Systems) was coated on copper or nickel-plated ELISA plates at a concentration of 1 μ g/mL, and then incubated with the indicated concentration of anti-CD71 antibody (“CD71 21.12”) or

activatable anti-CD71 antibodies (“CD71.TF01-2001”, “CD71.TF02-2001”, “CD71.TF02.13-2001” or “CD71.TF02.18-2001”), where in the activatable antibodies were assayed in their uncleaved form. The amount of bound antibody was detected by incubation and detection by goat anti-human antibody conjugated to horseradish peroxidase and Ultra TMB (Thermo Fisher Scientific) detection.

[000632] As depicted in Figure 28, the exemplary assay demonstrated that the anti-human CD71 21.12 antibody bound with a K_d of ~0.14 nM. The binding constants of activatable anti-CD71 antibodies are shifted to the right as a result of their respective masking efficiencies.

Table 16: Exemplary Observed CD71 Binding Activity of Activatable Anti-CD71 Antibodies

Antibody	K_d (nM) 1 $\mu\text{g/mL}$ CD71 antigen	ME*
CD71 21.12	0.14	-
CD71 TF01-2001	69.99	500
CD71 TF02-2001	21.45	153
CD71 TF02.13-2001	9.14	65
CD71 TF02.18-2001	1.59	11

*ME = masking efficiency

EXAMPLE 21: CD71 Expression in Normal Human and Cynomolgus Tissues

[000633] This Example shows that CD71 is expressed in a large variety of normal human and cynomolgus tissues by immunohistochemical (IHC) staining using an anti-CD71 antibody.

[000634] Table 17 shows that CD71 is expressed in certain normal human and cynomolgus tissue samples, using IHC staining with a commercially-purchased anti-CD71 antibody (rabbit anti-CD71 mAb IgG; D7G9X, Cell Signaling Tech.). Table 17 shows a summary of the relative level of IHC staining of CD71 of a variety of formalin-fixed

paraffin-embedded normal human and cynomolgus tissues, where each plus sign (“+”) corresponds to a higher level of staining, and each minus sign (“-”).

Table 17: IHC Assay of CD71 Expression in FFPE Normal Tissues

Tissue Type	Cynomolgus	Human
Bone	+	++
Breast	-	-
Brain	+	+
Colon	-	-
Esophagus	-	-
Heart	-	-
Kidney	+	- / +
Liver	-	-
Lung	-	++ (in a few cells)
Nerve	+++	++
Ovary	+	+
Pancreas	-	-
Prostate	-	-
Skin	N/A	-/+
Small Intestine	+	-
Spleen	-	-
Stomach	- / +	+
Striated / Skeletal Muscle	-	-
Testis	-	+
Uterus	-	- / +

EXAMPLE 22: CD71 Expression in Multiple Primary and Metastatic Tumors

[000635] This Example shows that CD71 is expressed in a large number and variety of patient-derived tumors by immunohistochemical (IHC) staining using an anti-CD71 antibody.

[000636] Table 18 shows that CD71 is moderately or highly expressed in a large number and variety of patient-derived tumor samples, using IHC staining with a commercially-purchased anti-CD71 antibody on multiple patient-derived tumor tissue microarrays (TMA). Table 18 shows a summary of the level of IHC staining of CD71 of the TMAs shows that a large number of cores derived from multiple patient-derived samples showed a strong CD71 signal. The number in parentheses following each type of cancer indicate the number of individual patient-derived samples.

Table 18: IHC Assay of CD71 Expression In Patient-Derived Cancers

Cancer Type (Total No. of Samples)	Samples with IHC Score (%)		
	0	1	2/3
Colorectal Cancer (228)	1 (0.4)	20 (8.8)	207 (90.8)
Pancreatic Cancer (1060)	0 (0)	26 (24.5)	90 (75.5)
Small Cell Lung Cancer (8)	2 (25)	4 (50)	2 (25)
Non-Small Cell Lung Cancer (42)	2 (3.2)	26 (41.9)	34 (54.8)
Ovarian Cancer (53)	3 (7.1)	19 (45.2)	20 (47.6)
Head and Neck Squamous Cell Carcinoma (53)	3 (5.7)	20 (38)	30 (57)
Breast Cancer (41)	1 (2.4)	15 (36.6)	57 (61.0)
Gastric Cancer (17)	0 (0)	6 (35.3)	11 (64.7)
Esophageal Cancer (14)	0 (0)	7 (50)	7 (50)

EXAMPLE 23: CD71 Fab Binding Assay

[000637] This Example shows that an anti-human CD71 antigen-binding fragment (Fab) of the present disclosure, which is derived from anti-human CD71 antibody of the present disclosure, demonstrated a binding affinity to recombinant human and cynomolgus CD71 protein.

[000638] A Fab antigen-binding fragment was generated by digestion of anti-CD71 21.12 antibody of the present disclosure with the papain enzyme in accordance with known protocols to generate an anti-CD71 Fab fragment. The anti-CD71 Fab fragment of the present disclosure was assayed for binding to recombinant human or cynomolgus CD71 protein by measurement of the kinetic on- and off-rates of a 1:3 dilution series of the Fab fragment to a substrate-immobilized recombinant CD71 protein (Octet system, ForteBio). The recombinant CD71 proteins included a hexa-histidine (His₆) peptide tag, by which the protein was immobilized to a Ni-NTA (nitrilotriacetic acid)-containing substrate. The results are shown in Table 19.

Table 19: Binding Kinetics of Anti-CD71 Fab to Human and Cynomolgus CD71

Target	k_{on} (1/Msec)	k_{dis} (sec ⁻¹)	K_d (nM)
Human CD71-His ₆	6.64×10^5	3.98×10^{-3}	14
Cyno CD71-His ₆	3.90×10^5	5.59×10^{-3}	6

EXAMPLE 24: Histopathology Of Cynomolgus Monkeys After ADC Treatment

[000639] This Example describes a study in which a cynomolgus monkey treated with an anti-human CD71 antibody drug conjugate (ADC) of the present disclosure demonstrated histopathological abnormalities as a result of the toxicity of the ADC.

[000640] Two male cynomolgus monkeys were each given a single intravenous dose of an ADC of the present disclosure (CD71 21.12-spdb-DM4) at either 5 mg/kg or 7.5 mg/kg dosage. The specimen given the 7.5 mg/kg dose was subsequently euthanized on day 11 of the study due to moribundity, and its tissues were examined. Light microscopic examination of a limited selection of tissues revealed prominent, interrelated intestinal lesions (*e.g.*, ulceration, inflammation, hemorrhage, bacterial overgrowth, and/or fibrino-necrotic exudate in the colon, duodenum, and jejunum). The specimen also exhibited lung hemorrhage, necrosis, inflammation, and bacteria, suggestive of secondary bacteremia. Ulceration and bacteria in the tongue were also observed. Marked splenic lymphoid hyperplasia was also noted.

[000641] The moribundity of the specimen may be attributable to, for example, the fulminant and extensive intestinal lesions, as well as the secondary systemic lesions (*e.g.*, in the lung). The development of the intestinal and systemic lesions may be attributable to, for example, observed marked femoral bone marrow myeloid depletion and/or clinically observed neutropenia.

EXAMPLE 25: CD71 Expression on H292, HCC1806, and MDA MB 231 Cell Lines

[000642] This Example shows that anti-human CD71 activatable antibodies of the present disclosure and anti-human CD71 activatable antibody drug conjugates of the present disclosure bind CD71 on multiple cell lines with a higher dissociation constant than that of the unmasked anti-human CD7 antibody of the present disclosure, thus showing the effect of the mask in reducing binding prior to activation.

[000643] Figures 29A, 29B, and 29C shows the amount of binding of anti-CD71 activatable antibodies, activatable antibody drug conjugates, and parental antibodies of the present disclosure to the cell lines MDA MB 231 (Figure 29A), H292 (Figure 29B), and HCC1806 (Figure 29C). In this study, the binding of the antibodies of the present disclosure to the indicated cell lines were performed using a standard FACS labelling method. Briefly, cells were labeled with the indicated antibodies or activatable antibodies of the present disclosure: an isotype control (palivizumab), human CD71 antibody (anti-human CD71 Ab 21.12 antibody), an anti-human CD71 activatable antibodies (anti-human CD71 TF02.13-2001 or anti-human CD71 TF02.13-3001). In addition, cell were also labeled with an isotype ADC control (palivizumab-vc-MMAE), an anti-human CD71 antibody drug conjugate (ADC) of the present disclosure (CD71 Ab21.12-PEG2-vc-MMAD, CD71 Ab 21.12-vc-MMAE, or CD71 Ab 21.12-spdb-DM4), or an activatable antibody drug conjugate (AADC) of the present disclosure: CD71 TF02.13-3001-PEG2-vc-MMAD, CD71 TF02.13-3001-vc-MMAE, CD71 TF02.13-2001-spdb-DM4 or CD71 TF02.13-3001-spdb-DM4) at the indicated concentrations and subsequently detected with an Alexa Fluor 647 labeled goat anti-human IgG secondary antibody.

[000644] Table 20 below shows the equilibrium dissociation constants and Bmax values based on the binding curves depicted in Figures 29A to 29C. These results show that anti-human CD71 Ab 21.12 antibody bound all cell lines with similar Kd, while the binding of anti-human CD71 activatable antibodies and activatable antibody drug conjugates of the present disclosure to the cell lines demonstrated a significantly higher Kd, which is indicative of the masking efficiency of the masking moiety in the activatable antibodies of the present disclosure.

Table 20: Exemplary Observed CD71 Binding Activity of Anti-CD71 Binders

Article	MDA MB231 Kd (nM)	H292 Kd (nM)	H292 Bmax (MFI)	HCC1806 Kd (nM)
Ab 21.12	1.032	1.533	1824	5.0
Ab 21.12- PEG2-vc- MMAD	1.471	-	-	-
Ab 21.12- vc-MMAE	-	1.328	1597	6.4
Ab 21.12- spdb-DM4	1.238	-	-	-
TF02.13-	-	92.32	2208	~200

2001				
TF02.13-				
2001-				
PEG2-vc-	-	-	-	-
MMAD				
TF02.13-				
2001-vc-	-	-	-	-
MMAE				
TF02.13-				
2001-spdb-	-	-	-	~200
DM4				
TF02.13-	30.60	52.12	1821	~200
3001				
TF02.13-				
3001-	52.58	-	-	-
PEG2-vc-				
MMAD				
TF02.13-				
3001-vc-	-	53.78	1848	-
MMAE				
TF02.13-				
3001-spdb-	46.92	-	-	~200
DM4				
Isotype	-	-	-	-
Isotype-vc-	-	57.05	10.64	-
MMAE				

EXAMPLE 26: Binding of Anti-CD71 Activatable Antibodies With Modified

Substrates

[000645] This Example shows that anti-human CD71 activatable antibodies of the present disclosure bind CD71 on a cell line and *in vitro* with a higher dissociation constant than that of the unmasked anti-human CD71 antibody of the present disclosure, thus showing the effect of the mask in reducing binding prior to activation.

[000646] Figures 30A and 30B shows the amount of binding of anti-CD71 activatable antibodies and parental antibodies of the present disclosure to the H292 cell line. In this study, the binding of the antibodies of the present disclosure to the indicated cell lines were performed using a standard FACS labelling method. Briefly, cells were labeled with the indicated antibodies or activatable antibodies of the present disclosure: an isotype control (palivizumab), human CD71 antibody (anti-human CD71 Ab 21.12), an anti-human CD71 activatable antibodies (anti-human CD71 TF02.13-3001, TF02.13-3007, TF02.13-3008, TF02.13-3011, TF02.13-3012, TF02.13-3013, TF02.13-2001, TF02.13-2007, TF02.13-2008, TF02.13-2011, TF02.13-2012, TF02.13-2013), where in the activatable antibodies

were assayed in their uncleaved form. These activatable antibodies include the light chain of the correspondingly named sequence listed above in Table C. The cells were labeled with the indicated test article at the indicated concentrations and subsequently detected with an Alexa Fluor 647 labeled goat anti-human IgG secondary antibody.

[000647] Figures 30C and 30C shows the amount of binding of anti-CD71 activatable antibodies and parental antibodies of the present disclosure to human CD71 in a solid-phase ELISA binding assay. In these examples, recombinant human CD71 protein (R&D Systems) was coated on copper or nickel-plated ELISA plates at a concentration of 1 µg/mL, and then incubated with the indicated concentration of an anti-CD71 antibody (anti-human CD71 Ab 21.12) or an activatable anti-CD71 antibody (anti-human CD71 TF02.13-3001, TF02.13-3007, TF02.13-3008, TF02.13-3011, TF02.13-3013, TF02.13-2001, TF02.13-2007, TF02.13-2008, TF02.13-2011, TF02.13-2012, TF02.13-2013), where in the activatable antibodies were assayed in their uncleaved form. The amount of bound antibody was detected by incubation and detection by goat anti-human antibody conjugated to horseradish peroxidase and Ultra TMB (Thermo Fisher Scientific) detection.

[000648] Table 21 below shows the equilibrium dissociation constants based on the binding curves depicted in Figures 30A to 30D. These results show that anti-human CD71 activatable antibodies and activatable antibody drug conjugates bind to the cell line and to the solid-phase bound CD71 at a significantly lower affinity than that of the parent anti-CD71 antibody, which is indicative of the masking efficiency of the masking moiety.

Table 21: Exemplary Observed CD71 Binding Activity of Anti-CD71 Binders

Article	H292 Kd (nM)	ELISA Kd (nM)
Ab 21.12	0.4412	0.2553 0.1624
Isotype	~22544	-
TF02.13-3001	10.17	0.8132
TF02.13-3007	86.66	2.210
TF02.13-3008	97.79	2.613
TF02.13-3011	67.36	0.7778
TF02.13-3013	78.27	4.912
TF02.13-2001	27.50	2.529
TF02.13-	70.27	-

2007		
TF02.13-2008	89.99	15.12
TF02.13-2011	57.15	4.274
TF02.13-2012	39.15	1.636
TF02.13-2013	70.27	3.248

EXAMPLE 27: Anti-CD71-AADC and ADC Cytotoxicity on Skin and Lung Cancer-Derived Cell Lines

[000649] This Example shows that anti-human CD71 antibody drug conjugates (ADCs) and activatable antibody drug conjugates (AADCs) of the present disclosure demonstrate a higher cytotoxicity against a lung adenocarcinoma and a skin squamous cell carcinoma-derived cell lines as compared to isotype-ADC controls.

[000650] Figures 31A – 31J show that activatable antibody drug conjugates (AADCs) of the present disclosure conjugated to either PEG2-vc-MMAD, or vc-MMAE showed a lower cytotoxicity against a variety of cancer-derived cell lines than that of a correspondingly conjugated parental anti-CD71 antibody drug conjugate. The cancer-derived cell lines that were treated with the ADCs or AADCs were derived from a skin squamous cell carcinoma (A431, Figure 31A and 31B), a lung adenocarcinoma (A549, Figure 31C and 31D), a hypopharyngeal carcinoma (FaDu, Figure 31E and 31F), a non-small cell lung cancer (NCI-H292, Figure 31G and 31H), and a pancreatic adenocarcinoma (BxPC3, Figure 31I and 31J).

[000651] The anti-human CD71 ADCs demonstrated a similar cytotoxic efficacy to each other against the cell lines. The anti-human CD71 AADCs demonstrated a lower efficacy than the parental anti-CD71 ADC, which may be attributable to the masking effect of the masking moiety. In this study, the various anti-human CD71 ADCs or AADCs (TF02.13-3001 or TF02.18-2001 conjugated to the indicated drug) of the present disclosure and isotype drug conjugates were applied at the indicated concentrations to the cells. The cytotoxicity was determined as a percentage of a population of untreated cells that were used as a control.

EXAMPLE 28: Anti-CD71 ADC Cytotoxicity on Various Cancer-Derived Cell Lines

[000652] This Example shows that anti-human CD71 antibody drug conjugates (ADCs) of the present disclosure demonstrate significant cytotoxicity against a variety of cancer-derived cell lines.

[000653] Table 23 shows the observed EC50s of an anti-CD71 antibody drug conjugate (ADCs) of the present disclosure conjugated to vc-MMAE. In this study, the anti-human CD71 Ab 21-12-vc-MMAE of the present disclosure was applied over a range of concentrations to the indicated cells. The cytotoxicity was determined as a percentage of a population of untreated cells that were used as a control, and an EC50 was calculated based on the observed cytotoxicity curve.

Table 23: Cytotoxicity EC50 of Anti-CD71-vc-MMAE

Cancer Origin	Cell Line	EC50 (nM)
Pancreatic Cancer	Miapaca2	0.2
	HPAF 2	0.2
	BxPC3	0.37
Colorectal Cancer	SW480	0.11
	SW1417	0.17
	HT29	0.18
	Ls411N	0.2
	SW48	0.2
	Ls174T	0.23
	Lovo	0.27
	HCT-116	0.54
	DLD1	1.76
	Fadu	0.40
Head-Neck Small Cell Carcinoma	A253	0.83
	SCC9	0.69
	KYSE 70	0.6
	SCC1	0.51
	KYSE 150	0.47
	SAS	0.33
	BHY	0.25
	SCC25	0.24
	A431	0.22
	NCI H292	0.3
Lung Cancer	Detroit 562	0.3
	A549	0.49
	NCI H2141	0.5
	NCI-H69	0.57
	NCI H526	0.7
	HS766T	1.02
	NCI H727	1.29

	NCI H889	1.3
Non-Hodgkin's Lymphoma	Ramos	0.2
	Raji	0.3
Ovarian Cancer	OVCAR3	0.20
	ET201	0.49
Breast Cancer	HCC1806	0.2
	MDA MB231	1.1

EXAMPLE 29: CD71 Expression on Small Cell Lung Cancer Cell Lines

[000654] This Example shows that CD71 antibody and antibody drug conjugates of the present disclosure bind CD71 on multiple small lung cancer cell lines (SCLC) essentially equivalently and specifically as shown by lack of binding of an isotype control or isotype control-ADC.

[000655] Figures 32A and 32B and Table 24 show the amount of binding of anti-CD71 antibodies and ADCs of the present disclosure to the SCLC cell lines DMS 79 and DMS 153. In this study, the binding of the antibodies and ADCs of the present disclosure to the indicated cell lines were performed using a standard FACS labelling method. Briefly, cells were labeled at the indicated concentrations with the indicated antibodies of the present disclosure: anti-human CD71 antibody (anti-human CD71 Ab21.12) or anti-human CD71 antibody conjugated to duocarmycin (anti-human CD71 Ab21.12-Duoc) or the corresponding isotype controls (pavilizumab) and subsequently detected with an Alexa Fluor 647 labeled goat anti-human IgG secondary antibody. Table 24 below shows the equilibrium dissociation constants based on the observed binding curves. These results show that anti-human CD71 21.12 antibody (CD71-Ab) and its corresponding ADC bound with essentially equivalent affinity.

Table 24: Exemplary Observed CD71 Binding Activity of Activatable Anti-CD71

Test Article	DMS 79 Kd (nM)	DMS 153 Kd (nM)
Ab 21.12	0.8983	4.471
Ab 21-12-Duoc	1.087	2.298

EXAMPLE 30: Anti-Proliferative Effect of Anti-CD71 Antibodies

[000656] This Example shows that anti-human CD71 antibodies of the present disclosure that bind CD71 on cancer cell lines demonstrate an anti-proliferative effect compared to an isotype control.

[000657] Figures 33A and 33B show the inhibition of cell proliferation by application of anti-CD71 Ab 21.12 antibody of the present disclosure to the H292 and HCC1806 cell lines, as compared to an isotype control. In this study, the indicated cells were plated at a density of 1000 cells per well in a 96 well plate and left to adhere overnight. An equal volume of test article at twice the indicated concentration was added and the plates were allowed to incubate for 5 days. Viability was subsequently measured using Cell Titer Glo reagent (Promega), and inhibition of proliferation was determined as a percentage of a population of untreated cells that were used as a control. The results indicate that anti-CD71 antibodies of the present disclosure showed significant inhibition of proliferation (EC₅₀ of ~ 4 nM in H292 cells, and ~ 10 nM in HCC1806 cells)

EXAMPLE 31: CD71 Expression In Multiple Metastatic Cancer Samples

[000658] This Example shows that CD71 is expressed in a large number and variety of patient-derived metastatic tumors by immunohistochemical (IHC) staining using an anti-CD71 antibody.

[000659] Figure 34 shows that CD71 is moderately or highly expressed in a large number and variety of patient-derived metastatic tumor samples, using IHC staining with a commercially-purchased anti-CD71 antibody on multiple patient-derived tumor tissue microarrays (TMA). Figure 34 shows a summary of the level of IHC staining of CD71 of the TMAs shows that a large number of cores derived from multiple patient-derived metastatic samples showed a strong CD71 signal.

EXAMPLE 32: *In Vivo* Imaging of Activatable Anti-CD71 Antibodies in a Mouse Pancreatic Cancer Xenograft Model

[000660] This Example shows that activatable anti-CD71 antibodies of the present disclosure demonstrate tumor-associated protease-dependent *in vivo* activation and binding to CD71 expressed in a pancreatic cancer mouse xenograft model by *in vivo* fluorescent imaging.

[000661] Figure 35 shows *in vivo* imaging of live mice with pancreatic tumor xenografts (BxPC3 cells) using fluorescently-conjugated anti-CD71 and activatable anti-CD71 antibodies of the present disclosure. In this study, 7-8 week-old *nu/nu* female mice were implanted subcutaneously in the right hind flank with 5×10^6 BxPC3 cells, a human pancreatic cancer-derived cell line. After the tumors grew to 300 to 380 mm³, an anti-CD71

antibody Ab 21.12 of the present disclosure (“anti-CD71”), an activatable anti-CD71 antibody of the present disclosure (“CD71 TF02.13-2012”), or a masked anti-CD71 antibody of the present disclosure lacking a CM domain (“CD71 TF02.13-NSUB”) were administered to each of the mice at a dose of 1 mg/kg. The administered antibodies were labeled by conjugation to the fluorescent label AlexFluor 750. The mice were subjected to *in vivo* fluorescent imaging 96 hours after administration of the antibodies, using a 745 nm excitation signal and detecting an 800 nm emission signal. The scale shows the relative magnitude of the detected fluorescent signal.

[000662] The results of this exemplary study showed that fluorescent signals from the labeled unmasked anti-CD71 antibody of the present disclosure and the labeled activatable anti-CD71 antibody of the present disclosure accumulated in the xenograft. In contrast, a correspondingly masked anti-CD71 antibody but which lacked a protease cleavage site (CM) did not detectably accumulate in its tumor xenograft site. Without being bound by any particular theory, this exemplary study demonstrated that activatable anti-CD71 antibodies of the present disclosure can be activated *in vivo* via tumor-associated protease cleavage, thus allowing the activated activatable anti-CD71 antibody to bind CD71 in the xenograft tumor to an extent comparable to the unmasked anti-CD71 antibody of the present disclosure. The masked anti-CD71 antibody lacking a protease cleavage domain (CM) of the present disclosure was not activatable in the same manner, and thus did not appreciably bind to the tumor xenograft.

EXAMPLE 33: *In Vivo* Imaging of Activatable Anti-CD71 Antibodies in a Mouse Model of Breast Cancer Metastasis

[000663] This Example shows that activatable anti-CD71 antibodies of the present disclosure demonstrate tumor-associated protease-dependent *in vivo* activation and binding to CD71 expressed in a mouse model of breast cancer metastasis by *in vivo* imaging.

[000664] Figure 36 shows *in vivo* imaging of live mice with a human-derived breast cancer metastatic cancer model, using fluorescently-conjugated anti-CD71 and activatable anti-CD71 antibodies of the present disclosure, and in which the cancer cells also express a bioluminescent signal. In this study, 7-8 week-old *nu/nu* female mice were injected into the left ventricle with 5×10^5 MDA-MB-231-luc2-D3LN cells, a human breast cancer-derived cell line that expresses the bioluminescent enzyme luciferase. Mice with detectable cancer cells (as determined by detecting a bioluminescent signal via *in vivo* imaging) received a 5

mg/kg intravenous injection of an isotype control antibody palivizumab (“isotype”), an anti-CD71 antibody Ab21.12 of the present disclosure (“anti-CD71”), or an activatable antibody of the present disclosure (“CD71 TF02.13-3001”). The administered antibodies were labeled by conjugation to the fluorescent label AlexFluor 750. The mice were subjected to *in vivo* fluorescent imaging 48 hours after administration of the antibodies, using a 745 nm excitation signal and detecting an 800 nm emission signal. Following fluorescent imaging of the antibodies, the mice were injected intraperitoneally with 3 mg luciferin, and then subjected to *in vivo* bioluminescent imaging 10 minutes post-injection by blocked excitation and open emission.

[000665] The results of this exemplary study showed that fluorescent signals from the labeled unmasked anti-CD71 antibody of the present disclosure and the labeled activatable anti-CD71 antibody of the present disclosure accumulated in the same location as the tumor location, as determined by the overlapping bioluminescent and fluorescent signals. In contrast, a correspondingly labeled isotype control antibody did not detectably accumulate at or near any tumor site. Without being bound by any particular theory, this exemplary study demonstrated that activatable anti-CD71 antibodies of the present disclosure can be activated *in vivo* via tumor-associated protease cleavage, thus allowing the activated activatable anti-CD71 antibody to bind CD71 in the metastatic tumor model to an extent comparable to the unmasked anti-CD71 antibody of the present disclosure.

EXAMPLE 34: *In Situ* Imaging of Activatable Anti-CD71 Antibodies

[000666] This Example shows that activatable anti-CD71 antibodies of the present disclosure demonstrate tumor-dependent *in situ* activation and binding to CD71 expressed in a non-small cell lung cancer (NSCLC) mouse xenograft model by immunohistochemical (IHC) staining.

[000667] Figure 37 shows an IHC staining assay of tissues sections of H292 NSCLC xenograft tumors using anti-CD71 and activatable anti-CD71 antibodies of the present disclosure. In this study, tissue sections from H292 xenograft tumors were pre-incubated either in the presence (+BSPI) or absence (-BSPI) of a broad-spectrum protease inhibitor cocktail (Protease Inhibitor Cocktail Set III, EDTA-Free (Cat. No. 539134, EMD Millipore) at a 1:15 dilution and 20 mM EDTA in Tris buffer). The tissue sections were then incubated for 2 hours at room temperature with either anti-CD71 antibody Ab 21.12 of the present disclosure (“anti-CD71”) or activatable anti-CD71 antibodies of the present disclosure

(“CD71 TF02.13-3001”, “CD71 TF02.13-2001”, or “CD71 TF02.13-2012”). After extensive washing to remove non-bound material, the presence of bound anti-CD71 antibodies or activated anti-CD71 antibodies of the present disclosure were detected with donkey anti-human IgG secondary antibody labeled with AlexaFluor 647 (Cat. No. 709-605-149, Jackson ImmunoResearch). Without being bound by any particular theory, this exemplary study shows that activatable anti-CD71 antibodies of the present disclosure demonstrated *in situ* activation and binding to CD71 in the xenograft tissue sections, which was inhibited by the pre-incubation of the tissues with a broad spectrum of protease inhibitors. The unmasked anti-CD71 antibody of the present disclosure demonstrated *in situ* binding to CD71 in the xenograft tissue sections irrespective of the presence or absence of protease-inhibitor pre-incubation.

Other Embodiments

[000668] While the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following.

CLAIMS

1. An isolated antibody or an antigen binding fragment thereof (AB) that specifically binds human CD71 and cynomolgus monkey CD71,
wherein the AB comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QRRNYPYT (SEQ ID NO: 15).
2. The AB of claim 1, wherein the AB comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8.
3. An activatable antibody that, in an activated state, specifically binds to human CD71 and cynomolgus CD71 comprising:
an antibody or antigen binding fragment thereof (AB) that specifically binds human CD71 and cynomolgus monkey CD71;
a masking moiety (MM) coupled to the AB that inhibits the binding of the AB to CD71 when the activatable antibody is in an uncleaved state; and
a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease,
wherein the AB comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QRRNYPYT (SEQ ID NO: 15).
4. The activatable antibody of claim 3, wherein the MM has one or more of the characteristics selected from the group consisting of:
(i) the MM has a dissociation constant for binding to the AB that is greater than the dissociation constant of the AB to CD71;

(ii) the MM does not interfere or compete with the AB for binding to CD71 when the activatable antibody is in a cleaved state;

(iii) the MM is a polypeptide of no more than 40 amino acids in length;

(iv) the MM polypeptide sequence is different that of human CD71;

(v) the MM polypeptide sequence is no more than 50% identical to any natural binding partner of the AB; and

(vi) the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 16-295 and 297-314.

5. The activatable antibody of claim 3, wherein the CM has one or more of the characteristics selected from the group consisting of:

(i) the CM is a substrate for a protease that is active in diseased tissue; and

(ii) the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 356-423, 680-698, 713, 714, and 789-808.

6. The activatable antibody of claim 3, wherein the antigen binding fragment thereof is selected from the group consisting of a Fab fragment, a F(ab')₂ fragment, a scFv, and a scAb.

7. The activatable antibody of claim 3, wherein the AB comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8.
8. The activatable antibody of claim 3, wherein the AB is linked to the CM.
9. The activatable antibody of claim 3, wherein the AB is linked directly to the CM.
10. The activatable antibody of claim 3, wherein the AB is linked to the CM via a linking peptide.
11. The activatable antibody of claim 3, wherein the MM is linked to the CM such that the activatable antibody in an uncleaved state comprises the structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM.
12. The activatable antibody of claim 3, wherein the activatable antibody comprises a linking peptide between the MM and the CM, a linking peptide between the CM and the AB, or both a linking peptide between the MM and the CM and a linking peptide between the CM and the AB.
13. The activatable antibody of claim 3, wherein the activatable antibody comprises a first linking peptide (LP1) and a second linking peptide (LP2), and wherein the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-LP1-CM-LP2-AB or AB-LP2-CM-LP1-MM.
14. The activatable antibody of claim 13, wherein the two linking peptides are not identical to each other.

15. The activatable antibody of claim 13, wherein each of LP1 and LP2 is a peptide of about 1 to 20 amino acids in length.

16. The activatable antibody of claim 3, wherein the activatable antibody has one or more of the characteristics selected from the group consisting of:

(a) the activatable antibody comprises the heavy chain sequence of SEQ ID NO: 325 or 699 and a light chain comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 327, 329, 331, 333, 335, 337, 650, 652, 654, 656, 658, 660, 670-673, 701-712, and 721-788;

(b) the activatable antibody comprises a combination of amino acid sequences, wherein the combination of amino acid sequences is selected from a single row in Table D, wherein for a given combination,

(i) the heavy chain of the AB comprises the amino acid sequences of the VH CDR sequences corresponding to the given combination in the single row listed in Table D,

(ii) the light chain of the AB comprises the amino acid sequences of the VL CDR sequences corresponding to the given combination in the single row listed in Table D,

(iii) the MM comprises the amino acid sequence of the mask sequence (MM) corresponding to the given combination in the single row listed in Table D, and

(iv) the CM comprises the amino acid sequence of the substrate sequence (CM) corresponding to the given combination in the single row listed in Table D;

(c) the activatable antibody comprises a combination of amino acid sequences, wherein for a given combination of amino acid sequences,

(i) the heavy chain of the AB comprises the amino acid sequences of the VH sequence or VH CDR sequences selected from the group consisting of: the VH sequence or VH CDR sequences listed in the corresponding column of Table E,

(ii) the light chain of the AB comprises the amino acid sequences of the VL sequence or VL CDR sequences selected from the group consisting of: the VL sequence or VL CDR sequences listed in the corresponding column of Table E,

(iii) the MM comprises the amino acid sequence of the mask sequence (MM) selected from the group consisting of: the MM sequences listed in the corresponding column of Table E, and

(iv) the CM comprises the amino acid sequence of the substrate sequence (CM) selected from the group consisting of: the CM sequences listed in the corresponding column of Table E; and

(d) the activatable antibody comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8, 809-836, and 841-908.

17. An activatable antibody comprising an antibody or an antigen binding fragment thereof (AB) that specifically binds to human CD71 and cynomolgus CD71, a masking moiety (MM), and a cleavable moiety (CM),

wherein the AB comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9), the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10), the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11), the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13), the VL CDR2 sequence STSNLAS (SEQ ID NO: 14), and the VL CDR3 sequence QQRRNYPYT (SEQ ID NO: 15); and

wherein the activatable antibody has one or more of the following characteristics selected from the group consisting of:

(i) the activatable antibody comprises a heavy chain sequence of SEQ ID NOs: 325 or 699; and a light chain comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 327, 329, 331, 333, 335, 337, 650, 652, 654, 656, 658, 660, 670-673, 701-712, and 721-788;

(ii) the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 16-295 and 297-314, and the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 356-423, 680-698, 713, 714, and 789-808; and

(iii) the activatable antibody comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8, 809-836, and 841-908.

18. A conjugated activatable antibody comprising the activatable antibody of claim 3 conjugated to an agent.

19. The conjugated activatable antibody of claim 18, wherein the agent has one or more of the characteristics selected from the group consisting of:

- (a) the agent is a toxin or fragment thereof;
- (b) the agent is a microtubule inhibitor;
- (c) the agent is a nucleic acid damaging agent;
- (d) the agent is a dolastatin or a derivative thereof;
- (e) the agent is an auristatin or a derivative thereof;
- (f) the agent is a maytansinoid or a derivative thereof;
- (g) the agent is a duocarmycin or a derivative thereof;
- (h) the agent is a calicheamicin or a derivative thereof;
- (i) the agent is a pyrrolobenzodiazepine or a derivative thereof;
- (j) the agent is auristatin E or a derivative thereof;
- (k) the agent is monomethyl auristatin E (MMAE);
- (l) the agent is monomethyl auristatin D (MMAD);
- (m) the agent is the maytansinoid DM1;
- (n) the agent is the maytansinoid DM4;
- (o) the agent is a detectable moiety; and
- (p) the agent is a diagnostic agent.

20. The conjugated activatable antibody of claim 18, wherein the agent is conjugated to the AB via a linker.

21. The conjugated activatable antibody of claim 20, wherein the linker with which the agent is conjugated to the AB comprises an SPDB moiety, a vc moiety, or a PEG2-vc moiety.

22. The conjugated activatable antibody of claim 18, wherein the agent is a toxin conjugated to the AB via a linker, and wherein the linker and the toxin conjugated to the AB comprise a moiety selected from the group consisting of: an SPDB-DM4 moiety, a vc-MMAD moiety, a vc-MMAE moiety, a vc-duocarmycin moiety, or a PEG2-vc-MMAD moiety.

23. The conjugated activatable antibody of claim 20, wherein the linker is a cleavable linker.
24. The conjugated activatable antibody of claim 20, wherein the linker is a non-cleavable linker.
25. A conjugated activatable antibody that, in an activated state, binds human CD71 and cynomolgus CD71 comprising:
- (a) an activatable antibody comprising:
 - (i) an antibody or an antigen binding fragment thereof (AB) that specifically binds to human CD71 and cynomolgus CD71, wherein the AB specifically binds human CD71 and cynomolgus monkey CD71, wherein the AB comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9), the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10), the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11), the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13), the VL CDR2 sequence STSNLAS (SEQ ID NO: 14), and the VL CDR3 sequence QQRRNYPYT (SEQ ID NO. 15);
 - (ii) a masking moiety (MM) coupled to the AB that inhibits the binding of the AB to CD71 when the activatable antibody is in an uncleaved state; and
 - (iii) a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease; and
 - (b) an agent conjugated to the AB.
26. The conjugated activatable antibody of claim 25, wherein the agent is selected from the group consisting of a dolastatin or a derivative thereof, an auristatin or a derivative thereof, a maytansinoid or a derivative thereof, a duocarmycin or a derivative thereof, a

calicheamicin or a derivative thereof, a pyrrolobenzodiazepine or a derivative thereof, auristatin E, monomethyl auristatin F (MMAF), monomethyl auristatin E (MMAE), monomethyl auristatin D (MMAD), maytansinoid DM4, maytansinoid DM1, a duocarmycin, a calicheamicin, a pyrrolobenzodiazepine, and a pyrrolobenzodiazepine dimer.

27. The conjugated activatable antibody of claim 25, wherein the agent is conjugated to the AB via a linker.

28. The conjugated activatable antibody of claim 27, wherein the linker with which the agent is conjugated to the AB comprises an SPDB moiety, a vc moiety, or a PEG2-vc moiety.

29. The conjugated activatable antibody of claim 25, wherein the agent is a toxin conjugated to the AB via a linker, and wherein the linker and the toxin conjugated to the AB comprise a moiety selected from the group consisting of: an SPDB-DM4 moiety, a vc-MMAD moiety, a vc-MMAE moiety, a vc-duocarmycin moiety, or a PEG2-vc-MMAD moiety.

30. The conjugated activatable antibody of claim 25, wherein the activatable antibody has one or more of the characteristics selected from the group consisting of:

(i) the activatable antibody comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light

chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8, 809-836, and 841-908; and

(ii) the activatable antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NOs: 325 or 699, and a light chain comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 327, 329, 331, 333, 335, 337, 650, 652, 654, 656, 658, 660, 670-673, 701-712, and 721-788.

31. The conjugated activatable antibody of claim 25, wherein the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 16-295 and 297-314.

32. The conjugated activatable antibody of claim 25, wherein the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 356-423, 680-698, 713, 714, and 789-808.

33. The conjugated activatable antibody of claim 25, wherein the activatable antibody has one or more of the following characteristics selected from the group consisting of:

(i) the activatable antibody comprises a combination of amino acid sequences, wherein the combination of amino acid sequences is selected from a single row in Table D, wherein for a given combination,

(a) the heavy chain of the AB comprises the amino acid sequences of the VH CDR sequences corresponding to the given combination in the single row listed in Table D,

(b) the light chain of the AB comprises the amino acid sequences of the VL CDR sequences corresponding to the given combination in the single row listed in Table D,

(c) the MM comprises the amino acid sequence of the mask sequence (MM) corresponding to the given combination in the single row listed in Table D, and

(d) the CM comprises the amino acid sequence of the substrate sequence (CM) corresponding to the given combination in the single row listed in Table D; and

(ii) the activatable antibody comprises a combination of amino acid sequences, wherein for a given combination of amino acid sequences,

(a) the heavy chain of the AB comprises the amino acid sequences of the VH sequence or VH CDR sequences selected from the group consisting of: the VH sequence or VH CDR sequences listed in the corresponding column of Table E,

(b) the light chain of the AB comprises the amino acid sequences of the VL sequence or VL CDR sequences selected from the group consisting of: the VL sequence or VL CDR sequences listed in the corresponding column of Table E,

(c) the MM comprises the amino acid sequence of the mask sequence (MM) selected from the group consisting of: the MM sequences listed in the corresponding column of Table E, and

(d) the CM comprises the amino acid sequence of the substrate sequence (CM) selected from the group consisting of: the CM sequences listed in the corresponding column of Table E.

34. A conjugated antibody comprising:

(a) an antibody or an antigen binding fragment thereof (AB) that specifically binds to human CD71 and cynomolgus CD71, wherein the AB comprises:

(i) the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QQRRNYPYT (SEQ ID NO: 15), or

(ii) a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8; and

(b) an agent conjugated to the AB, wherein the agent is selected from the group consisting of auristatin E, monomethyl auristatin F (MMAF), monomethyl auristatin E (MMAE), monomethyl auristatin D (MMAD), maytansinoid DM4, maytansinoid DM1, a calicheamicin, a pyrrolobenzodiazepine, a pyrrolobenzodiazepine dimer, and a duocarmycin.

35. A conjugated activatable antibody that, in an activated state, specifically binds to human CD71 and cynomolgus CD71, comprising:

(a) an activatable antibody comprising:

(i) an antibody or an antigen binding fragment thereof (AB) that specifically binds human CD71 and cynomolgus monkey CD71, wherein the AB comprises the VH CDR1 sequence GYTFTSYWMH (SEQ ID NO: 9); the VH CDR2 sequence AIYPGNSETG (SEQ ID NO: 10); the VH CDR3 sequence ENWDPGFAF (SEQ ID NO: 11); the VL CDR1 sequence SASSSVYYMY (SEQ ID NO: 12) or CRASSSVYYMY (SEQ ID NO: 13); the VL CDR2 sequence STSNLAS (SEQ ID NO: 14); and the VL CDR3 sequence QQRRNYPYT (SEQ ID NO: 15);

(ii) a masking moiety (MM) coupled to the AB that inhibits the binding of the AB to CD71 when the activatable antibody is in an uncleaved state;

(iii) a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease; and

(b) an agent conjugated to the AB, wherein the activatable antibody comprises:

(i) a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3-5, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6-8, 809-836, and 841-908, or

(ii) a heavy chain comprising the amino acid sequence of SEQ ID NOs: 325 or 699, and a light chain comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 327, 329, 331, 333, 335, 337, 650, 652, 654, 656, 658, 660, 670-673, 701-712, and 721-788; and

wherein the agent is selected from the group consisting of auristatin E, monomethyl auristatin F (MMAF), monomethyl auristatin E (MMAE), monomethyl auristatin D (MMAD), maytansinoid DM4, maytansinoid DM1, a calicheamicin, a pyrrolobenzodiazepine, a pyrrolobenzodiazepine dimer, and a duocarmycin.

36. The conjugated activatable antibody of claim 35, wherein the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 16-295 and 297-314.

37. The conjugated activatable antibody of claim 35, wherein the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 356-423, 680-698, 713, 714, and 789-808.

38. The conjugated activatable antibody of claim 35, wherein the agent is conjugated to the AB via a linker, and wherein the linker to which the agent is conjugated to the AB comprises an SPDB moiety, a vc moiety, or a PEG2-vc moiety.

39. The conjugated activatable antibody of claim 35, wherein the agent is a toxin conjugated to the AB via a linker, and wherein the linker and the toxin conjugated to the AB comprise a moiety selected from the group consisting of: an SPDB-DM4 moiety, a vc-MMAD moiety, a vc-MMAE moiety, a vc-duocarmycin moiety, or a PEG2-vc-MMAD moiety.

40. A conjugated activatable antibody or conjugated antibody comprising:
an antibody or antigen binding fragment thereof (AB) that, in an activated state,
binds human CD71 and cynomolgus CD71; and
a toxin conjugated to the AB via a linker,
wherein the conjugated activatable antibody or the conjugated antibody comprises
amino acid sequences, a linker, and a toxin selected from a single row in Table F, wherein
for the given combination:
- (a) the AB comprises a heavy chain comprising the amino acid sequence of the
heavy chain sequence or heavy chain variable domain sequence corresponding to the
given combination in the single row listed in Table F,
 - (b) the AB comprises a light chain comprising the amino acid sequence of the light
chain sequence or light chain variable domain sequence corresponding to the given
combination in the single row listed in Table F, and
 - (c) the linker and the toxin comprise the linker and the toxin corresponding to the
given combination in the single row listed in Table F.
41. A pharmaceutical composition comprising the activatable antibody of claim 3; and a
carrier.
42. The pharmaceutical composition of claim 41 comprising an additional agent.

43. The pharmaceutical composition of claim 42, wherein the additional agent is a therapeutic agent.
44. A pharmaceutical composition comprising the conjugated activatable antibody of claim 18; and a carrier.
45. The pharmaceutical composition of claim 44 comprising an additional agent.
46. The activatable antibody of claim 3, wherein the MM is linked to the CM.

47. The activatable antibody of claim 3, wherein the MM is linked directly to the CM.
48. The activatable antibody of claim 3, wherein the MM is linked to the CM via a linking peptide.