

Description

Title of Invention:

PEPTIDE TAG AND NUCLEIC ACID ENCODING SAME

Technical Field

[0001]

The present disclosure relates to a peptide tag, and a nucleic acid encoding the peptide tag.

Background Art

[0002]

An antibody functioning in a cell, namely, an intrabody (intracellular antibody) can affect the function of the cell by recognizing and binding to an antigen (target molecule) in the cell of a higher organism. Such an antigen can be a significant intracellular therapeutic target that can be inactivated by binding to an intracellular antibody. As a research method, use of an intracellular antibody attracts attention as means for specifically inhibiting the function of a protein directly by binding to the antibody in the cell.

[0003]

In case of an intracellular antibody, a hybridoma producing a monoclonal antibody recognizing an antigen is first produced by a standard method, and from the cDNA

thereof, an intracellular expression vector containing a DNA encoding a single chain antibody (single chain Fv: scFv) is constructed to obtain a complex of a heavy chain (VH) and a light chain (VL) as an intracellular antibody. Recently, a phage library for presenting a human scFv from an antibody isolated from a human B cell is produced, and is used for isolation of a scFv binding to an intracellular antigen in some cases.

[0004]

An antibody usually moves around in an extracellular space such as blood in a body, and recognizes an extracellular antigen to function, and hence works in the extracellular space as a premise. Accordingly, if an antibody is expressed in the cytoplasm, there arise problems of reduction of the expression level, folding causing limitation of a half-life of an antibody domain, and stability. The problem of stability of an intracellular antibody in the cytoplasm can lead to formation of an aggregation of the intracellular antibody in the cytoplasm. The formation of the aggregation can lead to reduction of a production amount of the intracellular antibody, and inhibition of expression of normal function. The same applies to a protein except for the intracellular antibody. The intracellular antibody has a characteristic of easily aggregating in particular, but a protein except for the intracellular

antibody also can form an aggregation in the cytoplasm when produced in the cytoplasm.

[0005]

By contrast, it has been shown that a peptide tag having an amino acid sequence containing 45% or more of acidic amino acids improves stability of an intracellular antibody (Patent Literature 1, and Non Patent Literature 1). In proposing the effectiveness of a peptide tag having an amino acid sequence containing 45% or more of acidic amino acids, Patent Literature 1 and Non Patent Literature 1 point out, as a design guideline for the peptide tag, significance of designing the peptide tag in such a manner that a charge value and a pI value are sufficiently low based not on the pH environment of the cytoplasm but on the pH environment on the surface of an endosome on the side of the cytoplasm. In Non Patent Literature 3, a membrane localization signal of HRAS is added to a heavy chain variable region of an antibody.

Citation List

Patent Literature

[0006]

Patent Literature 1: WO2019/004213

Non Patent Literature

[0007]

Non Patent Literature 1: Kabayama et al., 2020, Nature Communication, 11, 336

Non Patent Literature 2: Shubhada et al., 2012,
Biochemical genetics, Vol. 50, No. 7-8, pp. 625-41

Non Patent Literature 3: Tanaka et al., 2007, EMBO
Journal, 26:3250-3259

Summary of Invention

[0008]

The present disclosure provides a peptide tag, and a nucleic acid encoding the peptide tag. The peptide tag of the present disclosure can reduce the aggregation property of a protein in a cell.

[0009]

The present inventors made earnest studies on peptide tags having various amino acid sequences, resulting in finding a peptide tag having an effect of reducing an aggregation property of a protein in a cell.

[0010]

The present disclosure provides the following inventions:

[1] A peptide having an amino acid sequence with a length of, for example, 600 amino acids or less, for example, 10 to 200 amino acids (for example, 10 to 90 amino acids),

wherein (a) 5% or more and less than 45% of amino acids contained in the amino acid sequence are acidic amino acids, and

(b) 20% or more, and preferably 30% or more of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A, and

the peptide is preferably capable of reducing an aggregation property in a cell of a protein linked to the peptide, wherein 10% or 15% or more of the amino acids contained in the amino acid sequence are N or P.

[2] The peptide according to [1] above, wherein 30% or less, preferably 20% or less, more preferably 15% or less, and further preferably 10% or less of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of M, T, W, C, I, V, and L.

[3] The peptide according to [1] or [2] above, wherein each of A and G constitutes less than 10% of the amino acids contained in the amino acid sequence thereof.

[4] The peptide according to any one of [1] to [3] above, wherein

(a) 20% or more and less than 45% of the amino acids contained in the amino acid sequence are acidic amino acids,

(b) 30% or more and less than 70% of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A,

(c) 20% or less of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of M, T, W, C, I, V, and L, and

(d) each of A and G constitutes less than 10% of the amino acids contained in the amino acid sequence.

[5] A peptide having an amino acid sequence set forth in any one of SEQ ID NOs: 2 to 11.

[6] A nucleic acid encoding the peptide according to any one of [1] to [5] above.

[7] A protein expression vector comprising: the nucleic acid according to [6] above operably linked to a regulatory sequence; and a nucleic acid encoding a protein of interest in-frame to the nucleic acid according to [6] above.

[8] The protein expression vector according to [7] above, wherein the protein of interest is an antibody, or an antigen-binding fragment of an antibody.

[9] The protein expression vector according to [8] above, wherein the antigen-binding fragment of the antibody is a single chain Fv (scFv).

[10] A fusion protein of the peptide according to any one of [1] to [5] above and a protein of interest.

[11] The fusion protein according to [10] above, wherein the protein of interest is an antibody, or an antigen-binding fragment of an antibody.

[12] The fusion protein according to [11] above, wherein the antigen-binding fragment of the antibody is a single chain Fv (scFv).

[13] A protein-producing cell comprising: the nucleic acid according to [6] above operably linked to a regulatory sequence; and a nucleic acid encoding a protein of interest in-frame to the nucleic acid according to [6] above.

[14] A method for selecting or identifying an amino acid sequence having a length of 600 amino acids or less, for example, 10 to 200 amino acids (for example, 10 to 90 amino acids), the method comprising:

acquiring, from an amino acid sequence (group) (that can include an amino acid sequence (group) having a length of 10 to 200 amino acids (for example, 10 to 90 amino acids)), an amino acid sequence (group) in which (a) 5% or more and less than 45% of amino acids contained in the amino acid sequence are acidic amino acids, and (b) 20% or more of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A;

selecting or identifying an amino acid sequence of a peptide tag that, when the fusion protein of a peptide tag having the selected or identified amino acid sequence and a reference protein is expressed in a mammal cell (preferably in a human cell), provides reduction of a proportion of cells in which the fusion protein forms an

aggregation (for example, the proportion which is not more than a predetermined value); and

obtaining a peptide tag having the amino acid sequence, or a nucleic acid encoding the peptide tag.

[15] The method according to claim 14, wherein the amino acid sequence group to be acquired is the peptide according to any one of [1] to [5] above.

[16] The method according to [14] or [15] above, wherein the amino acid sequence group to be acquired is a group of amino acid sequences encoded by coding regions of human genome.

[17] The method according to any one of [14] to [16] above, wherein the amino acid sequence to be acquired contains a neo-antigen.

[0011]

[18] A peptide satisfying one or more selected from the group consisting of (a) to (h) described below, and capable of reducing an aggregation property in a cell of a protein linked to the peptide.

[19] A peptide selected from the group consisting of (A) to (AE) and (AF) to (AU) described below, and capable of reducing an aggregation property in a cell of a protein linked to the peptide.

[0012]

[20] A nucleic acid encoding the peptide according to [18] above.

[21] A protein expression vector comprising: the nucleic acid according to [20] above operably linked to a regulatory sequence; and a nucleic acid encoding a protein of interest in-frame to the nucleic acid according to [20] above.

[22] A fusion protein of the peptide according to [18] above and a protein of interest.

[23] The fusion protein according to [22] above, wherein the protein of interest is an antibody, or an antigen-binding fragment of an antibody.

[24] The fusion protein according to [23] above, wherein the antigen-binding fragment of the antibody is a single chain Fv (scFv).

[0013]

[25] A nucleic acid encoding the peptide according to [19] above.

[26] A protein expression vector comprising: the nucleic acid according to [25] above operably linked to a regulatory sequence; and a nucleic acid encoding a protein of interest in-frame to the nucleic acid according to [25] above.

[27] A fusion protein of the peptide according to [25] above, and a protein of interest.

[28] The fusion protein according to [26] above, wherein the protein of interest is an antibody, or an antigen-binding fragment of an antibody.

[29] The fusion protein according to [27] above, wherein the antigen-binding fragment of the antibody is a single chain Fv (scFv).

[0014]

[30] The method according to [14] above, wherein the amino acid sequence (group) includes the peptide according to [18] above.

[31] The method according to [14] above, wherein the amino acid sequence (group) includes the peptide according to [19] above.

[0015]

[32] The method according to [14] above, wherein the reference protein is a scFv, and the predetermined value is a value of 30% or less.

[33] The method according to [14] above, wherein the reference protein is a scFv, and the predetermined value is a value of 20% or less.

[34] The method according to [14] above, wherein the reference protein is a scFv, and the predetermined value is a value of 15% or less.

[35] The method according to [14] above, wherein the reference protein is a scFv, and the predetermined value is a value of 10% or less.

[36] The method according to [14] above, wherein the reference protein is a scFv, and the predetermined value is a value of 5% or less.

[0016]

[37] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (A) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (A) described below.

[38] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (B) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (B) described below.

[39] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (C) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (C) described below.

[40] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (D) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (D) described below.

[41] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (E) described

below, and the amino acid sequence to be acquired preferably further satisfies a condition of (E) described below.

[42] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (F) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (F) described below.

[43] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (G) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (G) described below.

[44] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (H) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (H) described below.

[45] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (I) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (I) described below.

[46] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (J) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (J) described below.

[47] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (K) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (K) described below.

[48] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (L) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (K) described below.

[49] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (M) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (M) described below.

[50] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (N) described

below, and the amino acid sequence to be acquired preferably further satisfies a condition of (N) described below.

[51] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (O) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (O) described below.

[52] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (P) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (P) described below.

[53] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (Q) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (Q) described below.

[54] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (R) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (R) described below.

[55] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (S) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (S) described below.

[56] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (T) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (T) described below.

[57] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (U) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (U) described below.

[58] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (V) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (V) described below.

[59] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (W) described

below, and the amino acid sequence to be acquired preferably further satisfies a condition of (W) described below.

[60] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (X) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (X) described below.

[61] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (Y) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (Y) described below.

[62] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (Z) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (Z) described below.

[63] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AA) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AA) described below.

[64] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AB) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AB) described below.

[65] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AC) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AC) described below.

[66] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AD) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AD) described below.

[67] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AE) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AE) described below.

[67A] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AF) described

below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AF) described below.

[67B] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AG) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AG) described below.

[67C] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AH) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AH) described below.

[67D] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AI) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AI) described below.

[67E] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AJ) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AJ) described below.

[67F] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AK) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AK) described below.

[67G] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AL) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AL) described below.

[67H] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AM) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AM) described below.

[67I] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AN) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AN) described below.

[67J] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AO) described

below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AO) described below.

[67K] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AP) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AP) described below.

[67L] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AQ) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AQ) described below.

[67M] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AR) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AR) described below.

[67N] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AS) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AS) described below.

[67O] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AT) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AT) described below.

[67P] The method according to [14] above, wherein the reference protein is a scFv, the amino acid sequence group includes the peptide according to (AU) described below, and the amino acid sequence to be acquired preferably further satisfies a condition of (AU) described below.

[0017]

[68] The method according to any one of [37] to [67] and [67A] to [67N] above, wherein the reference protein is a scFv, and the predetermined value is a value of 15% or less.

[69] The method according to any one of [37] to [67] and [67A] to [67N] above, wherein the reference protein is a scFv, and the predetermined value is a value of 10% or less.

[70] The method according to any one of [37] to [67] and [67A] to [67N] above, wherein the reference protein is a scFv, and the predetermined value is a value of 5% or less.

[0018]

[71] The method according to any one of [37] to [70] above, wherein a proportion of cells in which the reference protein forms an aggregation is a value more than 30%.

[72] The method according to any one of [37] to [70] above, wherein a proportion of cells in which the reference protein forms an aggregation is a value in a range of 30 to 40%.

[73] The method according to any one of [37] to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 40 to 50%.

[74] The method according to any one of [37] to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 50 to 60%.

[75] The method according to any one of [37] to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 60 to 70%.

[76] The method according to any one of [37] to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 70 to 80%.

[77] The method according to any one of [37] to [70] above, wherein a rate of cells in which the reference

protein forms an aggregation is a value in a range of 80 to 90%.

[78] The method according to any one of [37] to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 90 to 95%.

[79] The method according to any one of [37] to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 95 to 99%.

[80] The method according to any one of [37] to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 99 to 99.9%.

[81] The method according to any one of [37] to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 99.9 to 100%.

[0019]

[82] The method according to [69] above, wherein a proportion of cells in which the reference protein forms an aggregation is a value more than 30%.

[83] The method according to [69] above, wherein a proportion of cells in which the reference protein forms an aggregation is a value in a range of 30 to 40%.

[84] The method according to [69] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 40 to 50%.

[85] The method according to [69] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 50 to 60%.

[86] The method according to [69] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 60 to 70%.

[87] The method according to [69] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 70 to 80%.

[88] The method according to [69] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 80 to 90%.

[89] The method according to [69] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 90 to 95%.

[91] The method according to [69] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 99 to 99.9%.

[92] The method according to [69] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 99.9 to 100%.

[0020]

[93] The method according to [70] above, wherein a proportion of cells in which the reference protein forms an aggregation is a value more than 30%.

[94] The method according to [70] above, wherein a proportion of cells in which the reference protein forms an aggregation is a value in a range of 30 to 40%.

[95] The method according to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 40 to 50%.

[96] The method according to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 50 to 60%.

[97] The method according to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 60 to 70%.

[98] The method according to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 70 to 80%.

[99] The method according to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 80 to 90%.

[100] The method according to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 90 to 95%.

[101] The method according to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 95 to 99%.

[102] The method according to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 99 to 99.9%.

[103] The method according to [70] above, wherein a rate of cells in which the reference protein forms an aggregation is a value in a range of 99.9 to 100%.

[0021]

[104] The peptide according to any one of [1] to [5] above, wherein a peptide tag is capable of reducing an aggregation property of a scFv having at least an amino acid sequence set forth in SEQ ID NO: 1.

[105] The nucleic acid according to [6] above, wherein a peptide tag is capable of reducing an aggregation property of a scFv having at least an amino acid sequence set forth in SEQ ID NO: 1.

[106] The protein expression vector according to any one of [7] to [9] above, wherein a peptide tag is capable of reducing an aggregation property of a scFv having at least an amino acid sequence set forth in SEQ ID NO: 1.

[0022]

[107] The protein expression vector according to any one of [7] to [9] above, wherein the protein expression vector is a virus vector.

[108] The protein expression vector according to [107] above, wherein the virus vector is selected from the group consisting of a retrovirus vector, a lentivirus vector, an adenovirus vector, an adeno-associated virus

vector, a herpes simplex virus vector, a vaccinia virus vector, a Sendai virus vector, and a vesicular stomatitis virus vector.

[0023]

[109] The nucleic acid according to [6] above, wherein the nucleic acid is an mRNA.

[110] The nucleic acid according to [109] above, wherein the nucleic acid has a cap structure at the 5' end, and a poly A chain at the 3' UTR.

[111] The nucleic acid according to [109] or [110] above, wherein the nucleic acid contains pseudouridine as U.

[112] A nanoparticle, comprising the nucleic acid according to any one of [109] to [111] above.

[113] The nanoparticle according to [112], wherein the nanoparticle is a lipid nanoparticle.

[114] The method according to any one of [14] to [17] and [30] to [103] above, wherein the reference protein has an amino acid sequence set forth in SEQ ID NO: 1.

[115] The method, the peptide, the fusion protein, the nucleic acid, or the vector according to any one of those described above, wherein the cell is a eukaryotic cell.

[116] The method, the peptide, the fusion protein, the nucleic acid, or the vector according to any one of those described above, wherein the cell is a human cell.

[117] The method, the peptide, the fusion protein, the nucleic acid, or the vector according to any one of those described above, wherein a peptide tag does not prevent free localization of a protein of interest.

[0024]

A peptide tag of the present disclosure can cause an intracellular stability of a tagged protein. Accordingly, the peptide tag of the present disclosure can be a more highly biocompatible peptide tag.

Brief Description of Drawings

[0025]

[Figure 1] Figure 1 illustrates an effect of a peptide tag Tag4-1 on an aggregation property of a single chain Fv (scFv) in a cell.

[Figure 2A] Figure 2A illustrates a scheme for constructing a model of an intracellular accumulation of α -synuclein, that is, an amyloid.

[Figure 2B] Figure 2B illustrates fluorescence microscope images showing influence on intracellular synuclein fibril caused by intracellular expression of scFv-E6-CMA peptide fusion protein having Tag18-1, that is, one of peptide tags of the present disclosure.

[Figure 2C] Figure 2C illustrates an effect of removing synuclein fibril by intracellular expression of scFv-E6-CMA peptide fusion protein having Tag4-8 or Tag18-1, that is, one of peptide tags of the present disclosure.

[Figure 3A] Figure 3A illustrates fluorescence microscope images showing intracellular localization of a scFv-C2 itself having Tag18-1, that is, one of peptide tags of the present disclosure expressed in the cell.

[Figure 3B] Figure 3B illustrates a stabilizing action of the scFv-C2 having Tag18-1, that is, one of peptide tags of the present disclosure.

Description of Embodiments

[0026]

In the present invention, the term "subject" is a vertebrate, examples include birds and mammals, and specific examples include mammals such as a mouse, a rat, a hamster, a guinea pig, a horse, a cow, a pig, a goat, sheep, a donkey, a dog, and a cat, and primates such as a monkey, a chimpanzee, a gorilla, an orangutan, a bonobo, and a human, and particularly a human. Herein, the term "subject" is used in the meaning including a human as described above, and when a human is excluded, the term "non-human" is used.

[0027]

Herein, the term "antibody" means an immunoglobulin, and refers to a protein having a structure in which two heavy chains (H chains) and two light chains (L chains) stabilized through a disulfide bond are associated with each other. The heavy chain contains a heavy chain variable region VH, heavy chain constant regions CH1,

CH2, and CH3, and a hinge region positioned between the CH1 and the CH2, and the light chain contains a light chain variable region VL (wherein VL can be VK or V λ), and a light chain constant region CL. Among these regions, a variable region fragment (Fv) consisting of the VH and the VL is a region directly involved in an antigen bond, and imparting variety to the antibody. An antigen binding region consisting of the VL, the CL, the VH, and the CH1 is designated as a Fab region, and a region consisting of the hinge region, the CH2 and the CH3 is designated as a Fc region.

Among the variable regions, a region directly contacting an antigen is particularly largely changed, and is designated as a complementarity-determining region (CDR). A portion except for the CDRs that is comparatively less mutated is designated as a framework region (FR). There are three CDRs in each variable region of the heavy chain and the light chain, and these are designated, successively from the N terminal side, heavy chain CDR1 to CDR3, and light chain CDR1 to CDR3, respectively. Each CDR is incorporated into the framework regions. The heavy chain variable region of the antibody includes, from the N terminal side to the C terminal side, a heavy chain framework region 1, the heavy chain CDR1, a heavy chain framework region 2, the heavy chain CDR2, a heavy chain framework region 3, the heavy chain CDR3, and a heavy chain framework region 4 in

the stated order. The light chain variable region of the antibody includes, from the N terminal side to the C terminal side, a light chain framework region 1, the light chain CDR1, a light chain framework region 2, the light chain CDR2, a light chain framework region 3, the light chain CDR3, and a light chain framework region 4 in the stated order. The antibody may be a recombinant protein (recombinant antibody), and can be produced in an animal cell such as a Chinese hamster ovarian cell (CHO cell). The derivation of the antibody is not especially limited, and examples include an antibody of a non-human animal, an antibody of a non-human mammal (such as a mouse antibody, a rat antibody, or a camel antibody), and a human antibody. The antibody may be a chimeric antibody, a humanized antibody, or a fully humanized antibody. The antibody may be a polyclonal antibody or a monoclonal antibody, and is preferably a monoclonal antibody. A "chimeric antibody" refers to an antibody in which a heavy chain variable region and a light chain variable region are respectively linked to a heavy chain constant region and a light chain constant region of different species. A humanized antibody means an antibody in which an amino acid sequence characteristic to a non-human-derived antibody is substituted in the corresponding position of a human antibody, and an example includes an antibody having heavy chain CDR1 to CDR3 and light chain CDR1 to CDR3 of an antibody produced

by immunizing a mouse or a rat, and having the other regions including four framework regions (FR) each of the heavy chain and the light chain all derived from a human antibody. Such an antibody is designated as a CDR-grafted antibody in some cases. A "humanized antibody" encompasses a human chimeric antibody in some cases. A "human chimeric antibody" refers to a non-human-derived antibody in which a constant region of the non-human-derived antibody is substituted with a constant region of a human antibody. The antibody can be an isolated antibody, or a purified antibody. The antibody can be, for example, an IgG.

[0028]

A variable region of an immunoglobulin chain generally has the same entire structure including relatively preserved framework regions (FR) linked through three hypervariable regions (more frequently designated as "complementarity-determining regions" or CDRs). The CDRs obtained from the two chains of each heavy chain/light chain pair are typically arranged parallel by the framework region for forming a structure specifically binding to a specific epitope on a protein of interest (such as PCSK9). Light chain and heavy chain variable regions present in nature all usually have these elements in the following order from the N terminal to the C terminal: FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. In order to assign numbers to amino acids positioned in

these respective domains, a numbering system has been devised. This numbering system is defined in "Kabat Sequences of Proteins of Immunological Interest (1987 and 1991, NIH, Bethesda, MD)", or "Chothia & Lesk, 1987, J. Mol. Biol. 196: 901-917; Chothia et al., 1989, Nature, 342: 878-883".

[0029]

Herein, the antibody encompasses an antigen-binding fragment of an antibody. Herein, an antibody not fragmented may be referred to as a full length antibody. A full length antibody can contain the full length of the antibody excluding a signal sequence.

[0030]

Herein, the term "antigen-binding fragment" means a part of an antibody maintaining a binding property to an antigen. The antigen-binding fragment can contain either or both of a heavy chain variable region and a light chain variable region of the antibody of the present disclosure. The antigen-binding fragment may be chimerized or humanized. Examples of the antigen-binding fragment include Fab, Fab', F(ab')₂, and Fv. The antigen-binding fragment may contain a bonded product or functional equivalent produced by recombination (for example, a part of another antibody in the form of a scFv (single chain Fv), a diabody, a scDb, a tandem scFv, a leucine zipper type, or a sc(Fv)₂ (single chain (Fv)₂)). Such an antigen-binding fragment of an antibody can be

obtained, for example, by treating the antibody with an enzyme, although not especially limited. For example, when an antibody is digested with papain, a Fab can be obtained. Alternatively, when an antibody is digested with pepsin, a $F(ab')_2$ can be obtained, and when this is further reduced, a Fab' can be obtained. Herein, such an antigen-binding fragment of the antibody can be used. In an scFv, the VL and the VH are linked via an artificial polypeptide linker, and thus, the same antigen specificity as that of the original antibody can be maintained. The VL and the VH can be linked in the order of the VH and the VL, or the VL and the VH from the N terminal side. The linker can have a length of about 10 to 25 amino acids. The linker may contain glycine in a large amount, and may contain an amino acid such as serine or threonine for purpose of increasing water solubility.

[0031]

Herein, the term "intracellular antibody" (intrabody) refers to an antibody expressed in a cell (for example, in the cytoplasm or in the nucleus). Although an antibody is extracellularly secreted to function, an intracellular antibody is different in that it is designed to be expressed in a cell to function. The intracellular antibody can affect the function of an intracellular protein, and can inhibit the function thereof in the cytoplasm, the nucleus, or the secretory

pathway. A cancer gene product can be a target of the intracellular antibody (Biocca, S., Pierandrei-Amaldi, P., and Cattaneo, A. (1993), *Biochem Biophys Res Commun*, Vol. 197, p. 422 to 427; Biocca, S., Pierandrei-Amaldi, P., Campioni, N., and Cattaneo, A. (1994), *Biotechnology (NY)*, Vol. 12, p. 396 to 399; Cochet, O. et al., (1998), *Cancer Res*, Vol. 58, p. 1170 to 1176). The intracellular antibody directly binds to a protein for purpose of inhibiting the protein function. The bond may directly inhibit the function of the protein in some cases, and may inhibit the protein from binding to another protein in other cases.

[0032]

Examples of the intracellular antibody include various antibodies and antigen-binding fragments thereof, and although not especially limited, a scFv, a tandem scFv, a VHH antibody (nanobody), a minibody, and a diabody can be preferably used. A scFv is typically an antibody fragment having a heavy chain variable region and a light chain variable region of an antibody, and the heavy chain variable region and the light chain variable region are linked via a linker. A tandem scFv is typically an antibody fragment having two scFvs having different antigen specificities, and these are linked via a linker. A diabody is typically a dimer of a scFv. Diabodies are roughly divided into bivalent monospecific diabodies and bispecific diabodies. A minibody is

typically dimerized two fusion proteins each of a dimerized domain and a scFv via the dimerized domain. A VHH antibody is an antibody fragment containing a heavy chain variable domain of a heavy chain antibody. The VHH antibody is typically a heavy chain variable domain of a heavy chain antibody derived from a camelid (such as a camel, a llama, or an alpaca). Although a general antibody is extracellularly expressed, and hence can be caused to function only extracellularly, the intracellular antibody is superior because it can be caused to exhibit the antibody function in a cell. The intracellular antibody can be used in various applications in a cell such as activation and inactivation of a target protein, and neutralization and block of protein-protein interaction. A scFv tends to exhibit an aggregation property when expressed in a cell. Accordingly, in such a case, it is useful to reduce the aggregation property by obtaining a fusion protein by linking a peptide tag of the present disclosure to the intracellular antibody. When the aggregation property of a protein is reduced, the protein can be caused to exhibit functions inherent to the protein in the cell.

[0033]

Herein, the term "peptide tag" refers to one that labels a protein of interest, or changes a biochemical property of the protein of interest when fused with the protein of interest. Examples of the peptide tag include

various tags such as a FLAG tag, a 3×FLAG tag, a Myc tag, an HA tag, T7, a 6×His tag, a PA tag, an S tag, an E tag, VSV-G, Glu-Glu, Strep-tag II, a HSV tag, a Chitin Binding Domain (CBD) tag, a Calmodulin Binding Peptide (CBP) tag, a V5 tag, a GST tag, a maltose binding protein (MBP) tag, a thioredoxin (Trx) tag, and a mini-AID tag. These can be used for affinity purification of a protein of interest by utilizing affinity for the tag, or for detection of the protein of interest with an antibody to the tag produced. An antibody recognizing a tag is generally designated as a tag antibody, and a tag sequence corresponding to an epitope of the tag antibody is designated as an epitope tag. A tag can have a polypeptide chain generally with a length of several amino acids to several tens amino acids.

[0034]

Herein, the term "protein of interest" refers to a protein to be expressed in a cell. The protein of interest may be an aggregating protein or a non-aggregating protein. In either case, when the peptide tag of the present disclosure is added thereto, the stability is further increased, and robustness against formation of aggregation can be obtained. Even when added to an aggregating protein, however, the peptide tag of the present disclosure can reduce the aggregation property thereof in a cell, and therefore, the protein of interest can be preferably an aggregating protein. Even

when the protein of interest is a secretory protein, aggregation may be formed in a cell before the secretory protein is secreted extracellularly in some cases. The peptide tag of the present disclosure can be advantageously used also for a secretory protein, preferably a secretory protein having an aggregation property.

[0035]

Herein, the term "aggregating protein" refers to a protein that forms aggregation (particularly, an insoluble aggregation) in a cell. Herein, the term "aggregation property" means a property of forming aggregation, and the term "non-aggregation property" means a property of not forming aggregation. Attenuation of the aggregation property can be promotion of the non-aggregation property, and promotion of the aggregation property can be attenuation of the non-aggregation property. Herein, the term "non-aggregation property" is used interchangeably with the term "stability". Aggregation can be observed, for example, as a bright point under a microscope by immunocytochemistry (IC). An aggregation rate can be calculated, for example, as a proportion of cells exhibiting aggregation in cells forcedly expressing a protein. Reduction of the aggregation rate thus calculated means increase of cells that forcedly express a protein and are not affected by the aggregation, and therefore can be an index of

physiological favorability. Reduction of the aggregation property (for example, reduction of the aggregation rate) and increase of solubility are different indexes. The increase of solubility means increase of a concentration in an aqueous solution of available protein, and does not directly lead to the number of aggregations, or a proportion of cells having the aggregations. Accordingly, the increase of solubility does not always mean the reduction of the aggregation property (for example, the reduction of the aggregation rate).

[0036]

Herein, an amino acid sequence is described by one letter amino acid code. Specifically, A denotes alanine, R denotes arginine, N denotes asparagine, D denotes aspartic acid, C denotes cysteine, Q denotes glutamine, E denotes glutamic acid, G denotes glycine, H denotes histidine, I denotes isoleucine, L denotes leucine, K denotes lysine, M denotes methionine, F denotes phenylalanine, P denotes proline, S denotes serine, T denotes threonine, W denotes tryptophan, Y denotes tyrosine, and V denotes valine. Amino acids are usually 20 types of L-amino acids mentioned above.

[0037]

Herein, the term "regulatory sequence" refers to a sequence having activity of driving a gene operably linked thereto to transcribe RNA from the gene. The regulatory sequence is, for example, a promoter.

Examples of the promoter include a class I promoter (usable for transcription of an rRNA precursor), a class II promoter (containing a core promoter and an upstream promoter element, and usable for transcription of an mRNA), and a class III promoter (further roughly divided into type I, type II, and type III).

[0038]

The present invention provides a peptide tag that reduces aggregation tendency of an aggregating protein. The present invention provides a protein expression vector operably linked to a regulatory sequence, and containing a gene encoding the peptide tag. The present invention provides a protein of interest fused with the peptide tag. The present invention provides a protein expression vector operably linked to a regulatory sequence, and containing a gene encoding a protein of interest fused with the peptide tag. The protein of interest can be an intracellular protein in one embodiment. The protein of interest can be an intracellular antibody in one embodiment. The protein of interest can be an scFv in one embodiment.

[0039]

Hereinafter, the peptide tag of the present disclosure that reduces aggregation tendency of an aggregating protein will be described in detail. The peptide tag of the present disclosure can reduce aggregation tendency of a protein of interest in a

eukaryotic cell, particularly, in a human cell. In examination of pharmaceutical application and the like, it can be useful to reduce the aggregation tendency of a protein of interest in a human cell.

[0040]

The length of the peptide tag of the present disclosure is not especially limited, and can be, for example, 600 amino acids or less, 500 amino acids or less, 400 amino acids or less, 300 amino acids or less, or 200 amino acids or less, and for example, 5 amino acids to 100 amino acids, such as 10 amino acids to 90 amino acids, 20 amino acids to 80 amino acids, 30 amino acids to 70 amino acids, 40 amino acids to 60 amino acids, 10 amino acids to 50 amino acids, 10 amino acids to 40 amino acids, or 10 amino acids to 30 amino acids. In this embodiment, the lower limit of the length of the peptide tag of the present disclosure can be 5 amino acids or more, 10 amino acids or more, 15 amino acids or more, 20 amino acids or more, 30 amino acids or more, 40 amino acids or more, 50 amino acids or more, 60 amino acids or more, 70 amino acids or more, or 80 amino acids or more, and/or the upper limit can be 100 amino acids or less, 90 amino acids or less, 80 amino acids or less, 70 amino acids or less, 60 amino acids or less, 50 amino acids or less, 40 amino acids or less, 30 amino acids or less, or 20 amino acids or less.

[0041]

(a) The peptide tag of the present disclosure can contain acidic amino acids (amino acids belonging to Element 1) in the following ratio.

In the peptide tag of the present disclosure, 45% or more of amino acids contained in the amino acid sequence thereof can be acidic amino acids.

In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids. In a preferable embodiment, in the peptide tag of the present disclosure, 5% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, more preferably, 10% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, further preferably, 20% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, further preferably, 30% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, still further preferably, 35% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, and particularly preferably, 40% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids. The acidic amino acids are D or E. For example, an acidic amino acid content in the peptide tag of the

present disclosure can be 44% or less, 43.5% or less, 43% or less, 42.5% or less, 42% or less, 41.5% or less, 41% or less, 40% or less, 35% or less, 30% or less, 25% or less, or 20% or less. Thus, in one embodiment, a risk of occurrence of unexpected interaction with an intracellular molecule or the like having a positive charge based on a high acidic amino acid ratio in the peptide tag can be reduced.

[0042]

(b) The peptide tag of the present disclosure can contain basic amino acids (amino acids belonging to Element 2) in the following ratio.

The peptide tag of the present disclosure can contain basic amino acids in a rate of preferably 25% or less or 20% or less, and more preferably can contain basic amino acids in a rate of 15% or less of amino acids, can contain basic amino acids further preferably in a rate of 10% or less, can contain basic amino acids further preferably in a rate of 5% or less, and can contain basic amino acids particularly preferably in a rate less than 3%, less than 2%, or less than 1%. In a most preferable embodiment, the peptide tag of the present disclosure does not contain a basic amino acid in the amino acid sequence thereof. The basic amino acids are K, R, or H.

[0043]

(c) The peptide tag of the present disclosure can contain amino acids belonging to Element 3 in the following ratio.

The amino acids belonging to the Element 3 can be F, P, Y, G, S, Q, N, and A.

In the peptide tag of the present disclosure, 10% or more, preferably 20% or more, more preferably 30% or more, or 40% or more of amino acids contained in the amino acid sequence thereof can be preferably the amino acids of the Element 3. In the peptide tag of the present disclosure, 50% or more, 60% or more, or 70% or more of amino acids contained in the amino acid sequence thereof can be amino acids of the Element 3. In the peptide tag of the present disclosure, preferably 80% or less, more preferably 70% or less, and further preferably 60% or less of amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3. 50% or less, 40% or less, 30% or less, or 20% or less of amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3. In a preferable embodiment, in the peptide tag of the present disclosure, 20% or more and 80% or less, 30% or more and 70% or less, 30% or more and 60% or less, 30% or more and 50% or less, 30% or more and 40% or less, 40% or more and 70% or less, 40% or more and 60% or less, 40% or more and 50% or less, 50% or more and 70% or less, 50% or more and 60% or less, 60% or more and 80% or less, or 60% or more and 70% or

less of amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3.

[0044]

In a particularly preferable embodiment, in the peptide tag of the present disclosure, 5% or more, 10% or more, 15% or more, or 20% or more (preferably 21% or more, 25% or more, or 30% or more) of amino acids contained in the amino acid sequence thereof are either N or P. In the peptide tag of the present disclosure, for example, 90% or less, 80% or less, 70% or less, 60% or less, 50% or less, 45% or less, 40% or less, 35% or less, 30% or less, or 25% or less of amino acids contained in the amino acid sequence thereof can be either N or P. In one embodiment, in the peptide tag of the present disclosure, 10% or more and 20% or less of amino acids contained in the amino acid sequence thereof can be either N or P. In one embodiment, in the peptide tag of the present disclosure, 55% or more and 90% or less of amino acids contained in the amino acid sequence thereof can be either N or P. In one embodiment, in the peptide tag of the present disclosure, more than 10% and 20% or less of amino acids contained in the amino acid sequence thereof can be either N or P. In one embodiment, in the peptide tag of the present disclosure, more than 20% and 30% or less of amino acids contained in the amino acid sequence thereof can be either N or P. In one embodiment, in the peptide tag of the present disclosure,

more than 30% and 40% or less of amino acids contained in the amino acid sequence thereof can be either N or P. In one embodiment, in the peptide tag of the present disclosure, more than 40% and 50% or less of amino acids contained in the amino acid sequence thereof can be either N or P. In one embodiment, in the peptide tag of the present disclosure, more than 50% and 60% or less of amino acids contained in the amino acid sequence thereof can be either N or P.

[0045]

For example, in the peptide tag of the present disclosure, 5% or more, 10% or more, 15% or more, or 20% or more (preferably 21% or more, 25% or more, or 30% or more) of amino acids contained in the amino acid sequence thereof are N. In the peptide tag of the present disclosure, for example, 90% or less, 80% or less, 70% or less, 60% or less, 50% or less, 45% or less, 40% or less, 35% or less, 30% or less, or 25% or less of amino acids contained in the amino acid sequence thereof can be N. In one embodiment, in the peptide tag of the present disclosure, 10% or more and 20% or less of amino acids contained in the amino acid sequence thereof can be N. In one embodiment, in the peptide tag of the present disclosure, 55% or more and 90% or less of amino acids contained in the amino acid sequence thereof can be N. In one embodiment, in the peptide tag of the present disclosure, more than 10% and 20% or less of amino acids

contained in the amino acid sequence thereof can be N.

In one embodiment, in the peptide tag of the present disclosure, more than 20% and 30% or less of amino acids contained in the amino acid sequence thereof can be N.

In one embodiment, in the peptide tag of the present disclosure, more than 30% and 40% or less of amino acids contained in the amino acid sequence thereof can be N.

In one embodiment, in the peptide tag of the present disclosure, more than 40% and 50% or less of amino acids contained in the amino acid sequence thereof can be N.

In one embodiment, in the peptide tag of the present disclosure, more than 50% and 60% or less of amino acids contained in the amino acid sequence thereof can be N.

[0046]

For example, in the peptide tag of the present disclosure, 5% or more, 6% or more, 7% or more, 8% or more, 9% or more, 10% or more, 15% or more, or 20% or more (preferably 21% or more, 25% or more, or 30% or more) of amino acids contained in the amino acid sequence thereof are P. In the peptide tag of the present disclosure, for example, 90% or less, 80% or less, 70% or less, 60% or less, 50% or less, 45% or less, 40% or less, 35% or less, 30% or less, or 25% or less of amino acids contained in the amino acid sequence thereof can be P.

In one embodiment, in the peptide tag of the present disclosure, 55% or more and 90% or less of amino acids contained in the amino acid sequence thereof can be P.

In one embodiment, in the peptide tag of the present disclosure, 10% or more and 20% or less of amino acids contained in the amino acid sequence thereof can be P.

In one embodiment, in the peptide tag of the present disclosure, more than 10% and 20% or less of amino acids contained in the amino acid sequence thereof can be P.

In one embodiment, in the peptide tag of the present disclosure, more than 20% and 30% or less of amino acids contained in the amino acid sequence thereof can be P.

In one embodiment, in the peptide tag of the present disclosure, more than 30% and 40% or less of amino acids contained in the amino acid sequence thereof can be P.

In one embodiment, in the peptide tag of the present disclosure, more than 40% and 50% or less of amino acids contained in the amino acid sequence thereof can be P.

In one embodiment, in the peptide tag of the present disclosure, more than 50% and 60% or less of amino acids contained in the amino acid sequence thereof can be P.

[0047]

In a particularly preferable embodiment, in the peptide tag of the present disclosure, 5% or less, 10% or less, 15% or less, or 20% or less of amino acids contained in the amino acid sequence thereof are F or Y.

In a particularly preferable embodiment, in the peptide tag of the present disclosure, 5% or less, 10% or less, 15% or less, or 20% or less of amino acids contained in the amino acid sequence thereof are F and/or Y. In a

particularly preferable embodiment, in the peptide tag of the present disclosure, 5% or less, 10% or less, 15% or less, or 20% or less of amino acids contained in the amino acid sequence thereof are F and Y. In a particularly preferable embodiment, 5% or more, 10% or more, 15% or more, or 20% or more of amino acids contained in the amino acid sequence thereof are either N or P, and 5% or less, 10% or less, 15% or less, or 20% or less thereof are F and/or Y.

[0048]

(d) The peptide tag of the present disclosure can contain amino acids belonging to Element 4 in the following ratio.

The amino acids belonging to the Element 4 can be amino acids that are none of an acidic amino acid, a basic amino acid, and the amino acids of the Element 3. The amino acids of the Element 4 can be, for example, M, T, W, C, I, V, and L.

In the peptide tag of the present disclosure, 80% or less, 70% or less, 60% or less, 50% or less, 40% or less, 30% or less, 20% or less, 15% or less, 10% or less, or 5% or less of amino acids contained in the amino acid sequence thereof can be preferably the amino acids of the Element 4. In a preferable embodiment, the peptide tag of the present disclosure does not contain the amino acids of the Element 4.

[0049]

(e) In the peptide tag of the present disclosure, 40% or less, 30% or less, 20% or less, 15% or less, 10% or less, or 5% or less of amino acids contained in the amino acid sequence thereof can be preferably G. In one embodiment, the peptide tag of the present disclosure does not contain G.

[0050]

(f) In the peptide tag of the present disclosure, 40% or less, 30% or less, 20% or less, 15% or less, 10% or less, or 5% or less of amino acids contained in the amino acid sequence thereof can be preferably A. In one embodiment, the peptide tag of the present disclosure does not contain A.

[0051]

(g) In the peptide tag of the present disclosure, 40% or less, 30% or less, 20% or less, 15% or less, 10% or less, or 5% or less of amino acids contained in the amino acid sequence thereof can be preferably G, and 40% or less, 30% or less, 20% or less, 15% or less, 10% or less, or 5% or less of amino acids contained in the amino acid sequence thereof can be A.

[0052]

(h) The peptide tag of the present disclosure can preferably contain S. The peptide tag of the present disclosure preferably does not contain S. The peptide tag of the present disclosure can contain S in a rate of 10% or more, 20% or more, 30% or more, 40% or more, or

50% or more of amino acids contained in the amino acid sequence thereof. The peptide tag of the present disclosure can contain S in a rate of 60% or less, 50% or less, 40% or less, 30% or less, 20% or less, 15% or less, 10% or less, or 5% or less of amino acids contained in the amino acid sequence thereof.

[0053]

(A) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, and 20% or more and 80% or less (preferably 30% or more and 70% or less) of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3.

[0054]

(B) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 20% or more and 80% or less (preferably 30% or more and 70% or less) of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A.

[0055]

(C) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 20% or more and 80% or less (preferably 30% or more and 70% or less)

of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, and 10% or less of the amino acids contained in the amino acid sequence thereof can be G.

[0056]

(D) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 20% or more and 80% or less (preferably 30% or more and 70% or less) of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 10% or less of the amino acids contained in the amino acid sequence thereof can be A, and 10% or less of the amino acids contained in the amino acid sequence thereof can be G.

[0057]

(E) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 20% or more and 80% or less (preferably 30% or more and 70% or less) of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, and 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4.

[0058]

(F) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid

sequence thereof can be acidic amino acids, 20% or more and 80% or less (preferably 30% or more and 70% or less) of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A.

[0059]

(G) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 20% or more and 80% or less (preferably 30% or more and 70% or less) of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, and 10% or less of the amino acids contained in the amino acid sequence thereof can be G.

[0060]

(H) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 20% or more and 80% or less (preferably 30% or more and 70% or less) of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 30% or less of the amino acids contained in the amino acid

sequence thereof can be the amino acids of the Element 4, 10% or less of the amino acids contained in the amino acid sequence thereof can be A, and 10% or less of the amino acids contained in the amino acid sequence thereof can be G.

[0061]

(I) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, and 30% or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3.

[0062]

(J) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 30% or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A.

[0063]

(K) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 30% or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the

Element 3, and 10% or less of the amino acids contained in the amino acid sequence thereof can be G.

[0064]

(L) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 30% or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 10% or less of the amino acids contained in the amino acid sequence thereof can be A, and 10% or less of the amino acids contained in the amino acid sequence thereof can be G.

[0065]

(M) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 30% or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, and 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4.

[0066]

(N) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 30% or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the

Element 3, 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A.
[0067]

(O) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 30% or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, and 10% or less of the amino acids contained in the amino acid sequence thereof can be G.
[0068]

(P) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 30% or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, 10% or less of the amino acids contained in the amino acid sequence thereof can be A, and 10% or less of the amino acids contained in the amino acid sequence thereof can be G.
[0069]

(Q) In the above-described peptide tag, preferably 20% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, more preferably 30% or more and less than 45% of the amino acids contained in the amino acid sequence thereof can be acidic amino acids, further preferably 35% or more and less than 45% of the amino acids contained in the amino acid sequence thereof can be acidic amino acids, and particularly preferably 40% or more and less than 45% of the amino acids contained in the amino acid sequence thereof can be acidic amino acids.

[0070]

(R) In the above-described peptide tag, 10% or less of the amino acids contained in the amino acid sequence thereof are preferably basic amino acids. The peptide tag can contain preferably 5% or less of basic amino acids, and particularly preferably less than 3%, less than 2%, or less than 1% of basic amino acids. Alternatively, the peptide tag does not contain a basic amino acid in a preferable embodiment.

[0071]

(S) In the above-described peptide tag, 40% or more and 60% or less of the amino acids contained in the amino acid sequence thereof are preferably the amino acids of the Element 3.

[0072]

(T) In the above-described peptide tag, 20% or less, 15% or less, 10% or less, or 5% or less of the amino acids contained in the amino acid sequence thereof are preferably the amino acids of the Element 4. In a preferable embodiment, the above-described peptide tag does not contain the amino acids of the Element 4.

[0073]

(U) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 10% or less of the amino acids contained in the amino acid sequence thereof can be basic amino acids, 20% or more and 80% or less (preferably 30% or more and 70% or less) of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, 10% or less of the amino acids contained in the amino acid sequence thereof can be G, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A. In this embodiment, preferably 5% or more and less than 45% of the amino acids contained in the amino acid sequence thereof can be acidic amino acids, more preferably 10% or more and less than 45% of the amino acids contained in the amino acid sequence thereof can be acidic amino acids, further preferably 20% or more and less than 45% of the amino acids contained in the amino

acid sequence thereof can be acidic amino acids, further preferably 30% or more and less than 45% of the amino acids contained in the amino acid sequence thereof can be acidic amino acids, still further preferably 35% or more and less than 45% of the amino acids contained in the amino acid sequence thereof can be acidic amino acids, and particularly preferably 40% or more and less than 45% of the amino acids contained in the amino acid sequence thereof can be acidic amino acids.

[0074]

(V) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 5% or less of the amino acids contained in the amino acid sequence thereof can be basic amino acids, 20% or more and 80% or less (preferably 30% or more and 70% or less) of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, 10% or less of the amino acids contained in the amino acid sequence thereof can be G, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A. In this embodiment, in the peptide tag of the present disclosure, it is preferable that less than 3%, less than 2%, or less than 1% of the amino acids contained in the

amino acid sequence thereof can be basic amino acids, or that it does not contain a basic amino acid.

[0075]

(W) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 10% or less of the amino acids contained in the amino acid sequence thereof can be basic amino acids, 30% or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, 10% or less of the amino acids contained in the amino acid sequence thereof can be G, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A. In this embodiment, in the peptide tag of the present disclosure, 40% or more and 60% or less of the amino acids contained in the amino acid sequence thereof can be preferably the amino acids of the Element 3.

[0076]

(X) In the peptide tag of the present disclosure, less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 10% or less of the amino acids contained in the amino acid sequence thereof can be basic amino acids, 20% or more and 80% or less (preferably 30% or more and 70% or less) of the

amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, 10% or less of the amino acids contained in the amino acid sequence thereof can be G, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A. In this embodiment, in the peptide tag of the present disclosure, 20% or less, 15% or less, 10% or less, or 5% or less of the amino acids contained in the amino acid sequence thereof can be preferably the amino acids of the Element 4. In a preferable embodiment, the peptide tag of the present disclosure does not contain the amino acids of the Element 4.

[0077]

(Y) In the peptide tag of the present disclosure, 20% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 10% or less of the amino acids contained in the amino acid sequence thereof can be basic amino acids, 20% or more and 80% or less (preferably 30% or more and 70% or less) of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, 10% or less of the amino acids contained in the amino acid sequence thereof can be G, and 10% or less

of the amino acids contained in the amino acid sequence thereof can be A. In this embodiment, preferably 30% or more and less than 45% of the amino acids contained in the amino acid sequence thereof are acidic amino acids, more preferably 35% or more and less than 45% of the amino acids of the amino acid sequence are acidic amino acids, and further preferably 40% or more and less than 45% of the amino acids contained in the amino acid sequence thereof are acidic amino acids.

[0078]

(Z) In the peptide tag of the present disclosure, 20% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 10% or less of the amino acids contained in the amino acid sequence thereof can be basic amino acids, 30% or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, 10% or less of the amino acids contained in the amino acid sequence thereof can be G, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A. In this embodiment, preferably 40% or more and 60% or less of the amino acids contained in the amino acid sequence thereof are the amino acids of the Element 3.

[0079]

(AA) In the peptide tag of the present disclosure, 20% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 10% or less of the amino acids contained in the amino acid sequence thereof can be basic amino acids, 20% or more and 80% or less (preferably 30% or more and 70% or less) of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 20% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, 10% or less of the amino acids contained in the amino acid sequence thereof can be G, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A. In this embodiment, it is preferable that 15% or less of the amino acids contained in the amino acid sequence thereof are the amino acids of the Element 4, that 10% or less of the amino acids contained in the amino acid sequence thereof are the amino acids of the Element 4, and that 5% or less of the amino acids contained in the amino acid sequence thereof are the amino acids of the Element 4.

[0080]

(AB) In the peptide tag of the present disclosure, 30% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 10% or less of the amino acids contained in the amino acid sequence thereof can be basic amino acids, 30%

or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 30% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, 10% or less of the amino acids contained in the amino acid sequence thereof can be G, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A. In this embodiment, preferably 40% or more and 60% or less of the amino acids contained in the amino acid sequence thereof are the amino acids of the Element 3. In this embodiment, more preferably 35% or more and less than 45% of the amino acids contained in the amino acid sequence thereof can be acidic amino acids, and further preferably 40% or more and less than 45% of the amino acids contained in the amino acid sequence thereof can be acidic amino acids. In this embodiment, preferably 40% or more and 60% or less of the amino acids contained in the amino acid sequence thereof are the amino acids of the Element 3, and 35% or more and less than 45% of the amino acids contained in the amino acid sequence thereof are acidic amino acids, and further preferably 40% or more and 60% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, and 40% or more and less than 45% of the amino acids contained in the amino acid sequence thereof can be acidic amino acids.

[0081]

(AC) In the peptide tag of the present disclosure, 30% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 10% or less of the amino acids contained in the amino acid sequence thereof can be basic amino acids, 30% or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 20% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 4, 10% or less of the amino acids contained in the amino acid sequence thereof can be G, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A. In this embodiment, preferably 15% or less, 10% or less, or 5% or less of the amino acids contained in the amino acid sequence thereof can be G. In one embodiment, the peptide tag of the present disclosure does not contain G.

[0082]

(AD) In the peptide tag of the present disclosure, 35% or more and less than 45% of amino acids contained in the amino acid sequence thereof can be acidic amino acids, 10% or less of the amino acids contained in the amino acid sequence thereof can be basic amino acids, 30% or more and 70% or less of the amino acids contained in the amino acid sequence thereof can be the amino acids of the Element 3, 10% or less of the amino acids contained

in the amino acid sequence thereof can be the amino acids of the Element 4, 10% or less of the amino acids contained in the amino acid sequence thereof can be G, and 10% or less of the amino acids contained in the amino acid sequence thereof can be A.

[0083]

(AE) The peptide tag of the present disclosure can have, for example, an amino acid sequence set forth in any of SEQ ID NOs: 2 to 11. The peptide tag of the present disclosure can have preferably an amino acid sequence of SEQ ID NO: 5.

[0084]

(AF) The peptide tag of the present disclosure can have, for example, any one of amino acid sequences shown in Tables 1 to 11, Table 12-1, Table 12-2, Table 13-1, Table 13-2, Table 14-1, and Table 14-2.

[0085]

(AG) The peptide tag of the present disclosure may have one or more selected from the group consisting of addition and insertion of one or more amino acids selected from the group consisting of N and P, and substitution with the amino acids (for example, substitution of one to about 30% of amino acids, such as substitution with 1 to 20, 10, or several amino acids) in any amino acids of any amino acid sequences of (AE) and (AF) described above (for example, the amino acids of the Element 1, the Element 2, the Element 3, or the Element

4, or for example, the amino acids of the Element 2, the Element 3, or the Element 4). The peptide tag of the present disclosure may have addition and insertion of S in any amino acids of any amino acid sequences of (AF) and (AG) described above (for example, the amino acids of the Element 1, the Element 2, the Element 3, or the Element 4, or for example, amino acids of the Element 2, the Element 3, or the Element 4), and substitution of S with the amino acids (for example, substitution of one to about 30% of amino acids, such as substitution with 1 to 20, 10, or several amino acids). The peptide tag of the present disclosure may have deletion of an arbitrary amino acid of any one of the amino acid sequences of (AF) and (AG) described above (for example, the amino acids of the Element 1, the Element 2, the Element 3, or the Element 4, for example, the amino acids of the Element 2, the Element 3 (particularly, F and/or Y), and the Element 4 (A or G).

[0086]

(AH) In the peptide tag of the present disclosure, preferably 5% or more, 6% or more, 7% or more, 8% or more, 9% or more, 10% or more, 11% or more, 12% or more, 13% or more, 14% or more, 15% or more, 16% or more, 17% or more, 18% or more, 19% or more, or 20% or more of amino acids contained in the amino acid sequence thereof can be either N or P, or N and P. In the peptide tag of the present disclosure, preferably 5% or more, 6% or

more, 7% or more, 8% or more, 9% or more, 10% or more, 11% or more, 12% or more, 13% or more, 14% or more, 15% or more, 16% or more, 17% or more, 18% or more, 19% or more, 20% or more, 21% or more, 25% or more, or 30% or more of the amino acids contained in the amino acid sequence thereof can be P. In the peptide tag of the present disclosure, preferably 5% or more, 6% or more, 7% or more, 8% or more, 9% or more, 10% or more, 11% or more, 12% or more, 13% or more, 14% or more, 15% or more, 16% or more, 17% or more, 18% or more, 19% or more, 20% or more, 21% or more, 25% or more, or 30% or more of the amino acids contained in the amino acid sequence thereof can be N.

[0087]

(AI) In the peptide tag of the present disclosure, 45% or more of amino acids contained in the amino acid sequence thereof can be acidic amino acids, and 5% or more, 6% or more, 7% or more, 8% or more, 9% or more, 10% or more, 11% or more, 12% or more, 13% or more, 14% or more, 15% or more, 16% or more, 17% or more, 18% or more, 19% or more, 20% or more, 21% or more, 25% or more, or 30% or more thereof can be N or P. In this embodiment, less than 10% (preferably less than 5%, and more preferably 0%) of the amino acids contained in the amino acid sequence thereof can be G, less than 10% (preferably less than 5%, and more preferably 0%) thereof can be A,

and/or less than 10% (preferably less than 5%, and more preferably 0%) thereof can be F and Y.

[0088]

(AJ) In the peptide tag of the present disclosure,

(a) 5% or more and less than 45% of amino acids contained in the amino acid sequence can be acidic amino acids,

(b) 30% or more and less than 70% of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A, and 60% or less of the amino acids contained in the amino acid sequence can be S,

(c) 10% or less of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of M, T, W, C, I, V, and L, and

(d) each of A and G can constitute less than 10% of the amino acids contained in the amino acid sequence. In this embodiment, the amino acid sequence can have a length of 10 to 200 amino acids (such as 10 to 90 amino acids).

[0089]

(AK) In the peptide tag of the present disclosure,

(a) 5% or more and less than 45% of amino acids contained in the amino acid sequence can be acidic amino acids,

(b) 30% or more and less than 70% of the amino acids contained in the amino acid sequence can be amino acids

selected from the group consisting of F, P, Y, G, S, Q, N, and A, 60% or less of the amino acids contained in the amino acid sequence can be S, and 10% or more of the amino acids contained in the amino acid sequence can be N or P,

(c) 10% or less of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of M, T, W, C, I, V, and L, and

(d) each of A and G can constitute less than 10% of the amino acids contained in the amino acid sequence. In this embodiment, the amino acid sequence can have a length of 10 to 200 amino acids (such as 10 to 90 amino acids).

[0090]

(AJ) In the peptide tag of the present disclosure,

(a) 5% or more and less than 45% of amino acids contained in the amino acid sequence can be acidic amino acids,

(b) 30% or more and less than 70% of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A, 60% or less of the amino acids contained in the amino acid sequence can be S, and 10% or less of the amino acids contained in the amino acid sequence can be F and/or Y,

(c) 10% or less of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of M, T, W, C, I, V, and L, and

(d) each of A and G can constitute less than 10% of the amino acids contained in the amino acid sequence.

In this embodiment, the amino acid sequence can have a length of 10 to 200 amino acids (such as 10 to 90 amino acids).

[0091]

(AM) In the peptide tag of the present disclosure,

(a) 5% or more and less than 45% of amino acids contained in the amino acid sequence can be acidic amino acids,

(b) 30% or more and less than 70% of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A, 60% or less of the amino acids contained in the amino acid sequence can be S, 10% or more of the amino acids contained in the amino acid sequence can be N or P, and 10% or less of the amino acids contained in the amino acid sequence can be F and/or Y,

(c) 10% or less of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of M, T, W, C, I, V, and L, and

(d) each of A and G can constitute less than 10% of the amino acids contained in the amino acid sequence.

In this embodiment, the amino acid sequence can have a length of 10 to 200 amino acids (such as 10 to 90 amino acids).

[0092]

(AN) In the peptide tag of the present disclosure,

(a) 5% or more and less than 45% of amino acids contained in the amino acid sequence can be acidic amino acids,

(b) 30% or more and less than 65% of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A, 60% or less of the amino acids contained in the amino acid sequence can be S, and 10% or less of the amino acids contained in the amino acid sequence can be F and/or Y,

(c) 10% or less of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of M, T, W, C, I, V, and L, and

(d) each of A and G can constitute less than 10% of the amino acids contained in the amino acid sequence.

In this embodiment, the amino acid sequence can have a length of 10 to 200 amino acids (such as 10 to 90 amino acids).

[0093]

(AN) In the peptide tag of the present disclosure,

(a) 5% or more and less than 45% of amino acids contained in the amino acid sequence can be acidic amino acids,

(b) 30% or more and less than 65% of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A, 60% or less of the amino acids contained in the amino acid sequence can be S, 10% or more of the amino acids contained in the amino acid sequence can be N or P,

(c) 10% or less of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of M, T, W, C, I, V, and L, and

(d) each of A and G can constitute less than 10% of the amino acids contained in the amino acid sequence. In this embodiment, the amino acid sequence can have a length of 10 to 200 amino acids (such as 10 to 90 amino acids).

[0094]

(AO) In the peptide tag of the present disclosure,

(a) 5% or more and less than 45% of amino acids contained in the amino acid sequence can be acidic amino acids,

(b) 30% or more and less than 65% of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A, 60% or less of the amino acids contained in the amino acid sequence can be S, 10% or more of the amino

acids contained in the amino acid sequence can be N or P, and 10% or less of the amino acids contained in the amino acid sequence can be F and/or Y,

(c) 10% or less of the amino acids contained in the amino acid sequence can be amino acids selected from the group consisting of M, T, W, C, I, V, and L, and

(d) each of A and G can constitute less than 10% of the amino acids contained in the amino acid sequence.

In this embodiment, the amino acid sequence can have a length of 10 to 200 amino acids (such as 10 to 90 amino acids).

[0095]

(AP) In the peptide tag of the present disclosure,

(a) 5% or more and less than 45% of amino acids contained in the amino acid sequence can be acidic amino acids,

(b1) 55% or more and less than 90% of the amino acids contained in the amino acid sequence can be either of N and P, and

the rest of the amino acids contained in the amino acid sequence can be other amino acids. In this embodiment, 20% or less (preferably 15% or less, 10% or less, 5% or less, 4% or less, 3% or less, 2% or less, 1% or less, or 0%) of the amino acids contained in the amino acid sequence are neither an acidic amino acid nor N and P.

[0096]

(AQ) In the peptide tag of the present disclosure,

(a) 45% or more of amino acids contained in the amino acid sequence can be acidic amino acids,

(b1) 21% or more of the amino acids contained in the amino acid sequence can be N, and/or 7% or more thereof can be P, and

20% or less (preferably 15% or less, 10% or less, 5% or less, 4% or less, 3% or less, 2% or less, 1% or less, or 0%) of the amino acids contained in the amino acid sequence can be other amino acids.

[0097]

(AR) In the (AK) to (AQ) above, a rate of the Element 4 can be preferably 0%.

[0098]

(AS) In the (AK) to (AQ) above, a rate of A or G can be 0%. In the (AK) to (AO) above, a rate of A and G is preferably 0%.

[0099]

(AT) In the (AK) to (AQ) above, preferably, a rate of the Element 4 is 0%, and a rate of A and G is 0%.

[0100]

(AU) In the (AR) to (AT) above, a rate of the Element 2 is preferably 0%.

[0101]

In the (a) above, preferably 10% or more and less than 45% of the amino acids contained in the amino acid sequence are acidic amino acids, more preferably 15% or

more and less than 45% thereof are acidic amino acids, further preferably 20% or more and less than 45% thereof are acidic amino acids, still further preferably 25% or more and less than 45% thereof are acidic amino acids, and particularly preferably 30% or more and less than 45% thereof are acidic amino acids. In these examples, the upper limit of the acidic amino acid content can be, for example, less than 45%, 40% or less, or 35% or less.

[0102]

The peptide tag of the present disclosure can be added to a protein (such as an intracellular aggregating protein). Accordingly, the present disclosure provides a fusion protein of the peptide tag of the present disclosure and an intracellular aggregating protein. The peptide tag of the present disclosure may be added to an intracellular non-aggregating protein. When the tag is added thereto, toughness of the non-aggregating protein against a non-aggregation property can be increased. The protein can be, for example, an intracellular antibody. The intracellular antibody can be an antigen-binding fragment of an antibody. The peptide tag of the present disclosure may be added to a fusion protein of an intracellular antibody and a degradation-inducing sequence. Thus, selective degradation of a target to which the intracellular antibody binds can be induced. Examples of the intracellular antibody include the above-described antibody fragments. Other examples of the

intracellular antibody include antibodies that bind to α -synuclein, LRRK2, Tau, β -amyloid, amyloid precursor protein (APP), C9orf72, superoxide dismutase 1 (SOD1), TAR DNA-binding protein 43 (TDP43), Fused in Sarcoma (FUS), and a prion protein, and pathological forms thereof. Another example of the intracellular antibody includes an antibody inhibiting protein-protein interaction (PPI). Still another example of the intracellular antibody includes one in the form of a fusion protein with a degradation-inducing sequence that binds to a target. Other examples of the intracellular antibody include intracellular antibodies described in Molecular Therapy, 29(2): 859-872, 2021 (such as CP13 iB, PHF1 iB, and Tau5 iB), and intracellular antibodies each having all CDR sequences of these intracellular antibodies. "iB" is an abbreviation of "intrabody", and specifically means an intracellular antibody. The present disclosure provides a fusion protein of, for example, such an intracellular antibody and the peptide tag of the present disclosure. The intracellular antibody may preferably further include a degradation-inducing sequence. Other examples of the intracellular antibody include intracellular antibodies described in Molecular Therapy, 30(4): 1484-1499, 2022 (such as VHH E4-1, and VHHZ70), and intracellular antibodies each having all CDR sequences of these intracellular antibodies. The intracellular antibody may preferably

further include a degradation-inducing sequence. Still other examples of the intracellular antibody include an intracellular antibody described in J. Biol. Chem., 295(31): 10662-10676, 2020 (such as M204-scFv), and an intracellular antibody having all CDR sequences of this intracellular antibody. The intracellular antibody may preferably further include a degradation-inducing sequence. Still other examples of the intracellular antibody include an intracellular antibody described in WO2018/231254 (such as BIIB092 antibody), an intracellular antibody described in WO2016/207245, an antibody described in WO2018/011073 (such as C10-2), intracellular antibodies described in WO2015/114538 (such as VHH tau A2, VHH tau A2-SH, and VHH tau A2var-SH), intracellular antibodies described in WO2014/059442 (such as F9T, D11C, D4G, G12C, H2A and H7T), and JP2020/515233 (such as IE4, 9B11, 3A9, 10F10, 11F11, AC8, AE8, AA9, DG5, AD2, AD7, DG11, DG8, and DA9) and intracellular antibodies each having all CDR sequences of these intracellular antibodies. The intracellular antibody may preferably further include a degradation-inducing sequence. Examples of the intracellular antibody further include an intracellular antibodies capable of degrading and removing abnormal TDP-43 (such as SEQ ID NOs: 21 to 24) described in WO2019177138, and an intracellular antibody having all CDR sequences of this intracellular antibody. The intracellular antibody may preferably

further include a degradation-inducing sequence. Thus, an intracellular antibody (such as a scFv or VHH) binding to tau, an intracellular antibody (such as a scFv or VHH) binding to α -synuclein, and other intracellular antibodies (such as a scFv or VHH) against amyloid causing cytotoxicity in a cell are preferred, and can be linked to the tag to form a fusion protein with the tag. [0103]

In one embodiment, the peptide tag of the present disclosure does not have a CAAX motif (such as SEQ ID NO: 58: KLNPPDESGPGCMSCKCVLS). In one embodiment, the peptide of the present disclosure does not have a membrane localization signal. In one embodiment, the peptide tag of the present disclosure does not have a signal peptide sequence for extracellular secretion from the viewpoint of expressing a protein of interest in a cell. In one embodiment, the peptide tag of the present disclosure may have a signal sequence for extracellular secretion from the viewpoint of promoting extracellular secretion of a protein of interest. When a secretory protein has an aggregation property, the peptide tag of the present disclosure containing a signal sequence in the sequence thereof, or the peptide tag of the present disclosure linked tandem to the signal sequence can be advantageous. In one embodiment, the peptide tag of the present disclosure can contain a nuclear localization signal. In one embodiment, the peptide tag of the

present disclosure does not have a sequence preventing protein localization in the cytoplasm. In one embodiment, the peptide tag of the present disclosure promotes free distribution in a cell of the protein of interest. In one embodiment, the peptide tag of the present disclosure can promote intracellular bond of the protein of interest to an original binding partner, and co-localization with the binding partner. In one embodiment, the peptide tag of the present disclosure can have a sequence that imposes unique constraints on the distribution in a cell (or a sequence that prevents free distribution) of the protein of interest, but is possible not to have such a sequence.

[0104]

In any embodiment, the peptide tag of the present disclosure does not have the following sequence (Enzymol. 326, 362-267 (2000)): S-tag: KETAAAKFERQHMDS (SEQ ID NO: 14). In one embodiment, the peptide tag of the present disclosure can have a sequence in which a rate of the Element 2 is 10% or less, and/or a rate of A is 10% or less.

[0105]

In any embodiment, the peptide tag of the present disclosure does not have KLNPPDESGPGCMSCKCVLS (SEQ ID NO: 15) (Tanaka et al., 2007, EMBO Journal, 26: 3250-3259). In one embodiment, the peptide tag of the present

disclosure does not have a sequence having 90% or more sequence identity to this sequence.

[0106]

In any embodiment, the peptide tag of the present disclosure does not have EFGGAPEFPPKSTPPGSSGL (SEQ ID NO: 16), and a sequence having 90% or more sequence identity to this sequence (Paolo et al., 2003, Clinical Cancer Research, 9: 2837-2848). In one embodiment, the peptide tag of the present disclosure does not have a sequence having 90% or more sequence identity to this sequence.

[0107]

In any embodiment, the peptide tag of the present disclosure does not have any one of the following sequences (Arimori et al., 2017, Structures, 25: 1611-1622):

hMst1: DYEFLKSWTVEDLQKRLALDPMMEQEIEEIRQKYQSKRQPILDAIEAK
(SEQ ID NO: 17);

hMST2: DFDFLKNLSLEELQMRLKALDPMMEIEELRQRYTAKRQPILDAMDAK
(SEQ ID NO: 18);

hRaf1: GEVNWDAFSMPFLHNFLRILQREEEHLRQILQKYSYSRQKIQEALHAS
(SEQ ID NO: 19);

hRaf5: GEVEWDAFSIPELQNFLTILEKEEQDKIQVQKKYDKFRQKLEELRES
(SEQ ID NO: 20);

hSAV1: HILKWELFQLADLDITYQGMLKLLFMKELEQIVKMYEAYRQALLTELENR
(SEQ ID NO: 21). In one embodiment, the peptide tag of the present disclosure does not have a sequence having

90% or more sequence identity to any one of these sequences.

[0108]

In any embodiment, the peptide tag of the present disclosure have none of the following (Zhang et al., 2004, Protein Expression and Purification, 36(2): 207-216):

T7C:

LEDPFQSGVMLGVASTVAASPEEASVTSTEETLTPAQEAARTRAANKARKEAELAAATAEQ (SEQ ID NO: 22);

T7B: LEDPEEASVTSTEETLTPAQEAARTRAANKARKEAELAAATAEQ (SEQ ID NO: 23);

T7B1: LEDPEEASVTSTEETLTPAQEAARTRPPNKARKEAELAAATAEQ (SEQ ID NO: 24);

T7B2: LEDPEEASVTSTEETLTPAQEAARTRGGNKARKEAELAAATAEQ (SEQ ID NO: 25);

T7B3: LEDPEEASVTSTEETLTPAQEAARTRAANKARKEAELTAEQ (SEQ ID NO: 26);

T7B4: LEDPEEASVTSTEETLTPAQEAARTRAANKARKEAELEAETAEQ (SEQ ID NO: 27);

T7B5: LEDPEEASVTSTEETLTPAQEAARTRAAAKARKEAELAAATAEQ (SEQ ID NO: 28);

T7B6: LEDPEEASVTSTEETLTPAQEAARTRKARKEAELAAATAEQ (SEQ ID NO: 29);

T7B7: LEDPEEASVTSTEETLTPAQEAARTRAANKARKEAELAAATAEQ (SEQ ID NO: 30);

T7B8: LEDPEEASVTSTEETLTPAQEAARTRAANKARKEAELAAA (SEQ ID NO: 31);

T7B9: LEDPEEASVTSTEETLTPAQEAETEAAANKARKEAELEAETAEQ (SEQ ID NO: 32);

T7B10: LEDPTPAQEAARTRAANKARKEAELAAATAEQ (SEQ ID NO: 33);

T7A: LEDPAANKARKEAELAAATAEQ (SEQ ID NO: 34);

T7A1: LEDPERNKERKEAELAAATAEQ (SEQ ID NO: 35);

T7A2: LEDPERNKERKEAELEAATAEQ (SEQ ID NO: 36);

T7A3: LEDPERNKERKEAELEAETAEQ (SEQ ID NO: 37);

T3: LEDPAVWEAGKVVAKGVGTADITATTSNGLIASCKVIVNAATS (SEQ ID NO: 38);

T3A: LEDPAVWEAGKVVAKGVGTADITATTSNGLIASSEEADNAATS (SEQ ID NO: 39). In one embodiment, the peptide tag of the present disclosure does not have a sequence having 90% or more sequence identity to any one of these sequences.

[0109]

In any embodiment, the peptide tag of the present disclosure have none of the following (Japanese Patent Laid-Open No. 2015-97519):

Zif628:

ERPYPVESCRRFRSDELTRHIRIHTGQKPFQCRICMRNFSRSDHLTTHIRHT
GEKPFACDICGRKFARSDEKRHTKIHLRQKD (SEQ ID NO: 40);

HinR:

GRPRAITKHEQEQISRLLKGGHPRQQLAIIFGIGVSTLYRYFPASSIKKRMN (SEQ ID NO: 41); and

TrpR:

MAQQSPYSAAMAEQRHXXQEWLRFVDLLKNAYQNXXDLHLPLLNMLTPDERXXEAL

GTRVRIVEELLRGEMSQRELKNELGAGIATITRGSNSLKAAPVELRQWLEEVLLKSD
(SEQ ID NO: 42).

[0110]

In a preferable embodiment, the peptide tag of the present disclosure can be a natural sequence found in a non-human living thing. In a preferable embodiment, the peptide tag of the present disclosure can be a non-natural sequence or a part thereof. In either embodiment, the peptide tag of the present disclosure is none of the following (WO2010/034183):

NE-1: TKENPRSNQEESYDDNES (SEQ ID NO: 43);

NE-8: TKENPRTNQEESYDDNES (SEQ ID NO: 44);

NE-9: TKENPRSNQDESYDDNES (SEQ ID NO: 45);

NE-10: TKENPRSNQPPSYDDNES (SEQ ID NO: 46).

[0111]

In one embodiment, the peptide tag of the present disclosure is none of the following (WO2011/034605):

ACID.P1: GGSAQLEKELQALEKENAQLEWELQALEKELAQQGAT (SEQ ID NO: 50).

[0112]

In one embodiment, the peptide tag of the present disclosure is none of the following (WO2009/023270):

rPEG_K288-GFP: (GEGEGGGEG)₃₂ (SEQ ID NO: 51).

[0113]

In one embodiment, the peptide tag of the present disclosure is none of the following (WO2020/059228):

Hero7: MTRGNQRELARQKNMKKQSDSVKGKRRDDGLSAAARKQRDSEIMQQKQKK
ANEKKEPK (SEQ ID NO: 1038);

Hero9:

MSGPNGDLGMPVEAGAEGEEDGFGEAEYAAINSMLDQINSCLDHLEEKNDHLHARLQ
ELLESNRQTRLEFQQQLGEAPSDASP (SEQ ID NO: 1039);

Hero11:

MAQGQRKFQAHKPAKSKTAAAASEKNRGPRKGGRVIAPKKARVVQQQKLKKNLEVGI
RKKIEHDVVMKASSSLPKKLALLKAPAKKKGAAAATSSKTPS (SEQ ID NO:
1040).

[0114]

In one embodiment, the peptide tag of the present disclosure is none of the following (Protein Engineering, Design & Selection, 26(8): 490-501, 2013):

PAS#1: ASPAAPAPASPAAPAPSAPAA (SEQ ID NO: 52);

1P2: ASAAAPAAASAAASAPSAAAA (SEQ ID NO: 53);

PAS#5: AASPAAPSAPPAAASPAAPSAPPAA (SEQ ID NO: 54);

and repeated sequences of these (the number of repetition being, for example, 200 ± 20 times, 400 ± 40 times, or 600 ± 60 times).

[0115]

In one embodiment, the peptide tag of the present disclosure is none of the following (Protein Engineering, Design & Selection, 17(11): 779-786, 2004):

Z(W):

VDNKFNFKEQQNAFYEILHLPNLNEEQRNAFIQSLKDDPSQSANLLAEAKKLNDQAAP
K (SEQ ID NO: 55);

Z(a1):

VDNKFNKEQQNAEYEIEHLPNLNNEEQENAFIQSLEDDPSQSANLLAEAKKLNDQAAP

K (SEQ ID NO: 56);

Z(a2):

VDNKFNKEEEEEAEIEHLPNLNNEEQEEAFIESLEDDPSQSANLLAEAKKLNDQAAP

K (SEQ ID NO: 57)

[0116]

In one embodiment, the peptide tag of the present disclosure may have a mutation selected from the group consisting of substitution, insertion, deletion, addition, and elimination of one or more, preferably two or more amino acids in any one of the amino acid sequences of SEQ ID NOs: 43 to 46 and 47 to 58. In one embodiment, the peptide tag of the present disclosure can have less than 90%, 85% or less, 80% or less, 75% or less, 70% or less, 60% or less, 50% or less, 40% or less, 30% or less, 20% or less, or 10% or less sequence identity to any one of the amino acid sequences of SEQ ID NOs: 44 to 47 and 47 to 58.

[0117]

In any embodiment, the peptide tag of the present disclosure does not contain the following sequences (Protein Science (2019) 28, 823-836):

PA12-tag: GVAMPGAEDDVV (SEQ ID NO: 47);

PA14-tag: EGGVAMPGAEDDVV (SEQ ID NO: 48).

In any embodiment, the peptide tag of the present disclosure does not contain the following sequences:

DYKDDDDVEAEESDNVDSADAEEDSDVWWGGADTDYADGSEDKVVEVAEEEEVAEV
EEEEADDDDEDEDGDEVEEEAEOPYEEATERTTSIATTTTTTTESVEEVYPGQVGYP
GQVGYPGQV (SEQ ID NO: 49).

[0118]

In one embodiment, the peptide tag of the present disclosure has less than 90%, 80% or less, 70% or less, 60% or less, 50% or less, 40% or less, 30% or less, 20% or less, or 10% or less sequence identity to any one of SEQ ID NOs: 14 to 48 and 49 to 58.

[0119]

In one embodiment, the peptide tag of the present disclosure can have a sequence satisfying one, two or all of the following (i) to (iii): (i) a rate of the Element 2 is 10% or less, (ii) a rate of A is 10% or less, and (iii) a rate of G is 10% or less.

[0120]

In a preferable embodiment, the peptide tag of the present disclosure does not have a sequence consecutively containing 5 or more As. In a preferable embodiment, the peptide tag of the present disclosure does not have a sequence consecutively containing 5 or more Qs. In a preferable embodiment, the peptide tag of the present disclosure does not have a sequence consecutively containing 5 or more Ss. In a preferable embodiment, the peptide tag of the present disclosure does not have a sequence consecutively containing 5 or more Ns. In a preferable embodiment, the content of a specific single

amino acid in the amino acid sequence of the peptide tag does not exceed 50%, 40%, 35%, 30%, 25%, or 20%. In a preferable embodiment, the peptide tag of the present disclosure does not contain an amino acid sequence having a length of 3 to 8 amino acids described in Table 1 in WO2002/092132, or does not contain a consecutive repeat (for example, consecutive repeat of 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, or 10 or more times) of the amino acid sequence.

[0121]

For example, the peptide tag of the present disclosure can inhibit an intracellular antibody from forming aggregation to promote uniform distribution in a cell of the antibody, and/or inhibit aggregation formation to promote bond of the intracellular antibody to an antigen in a cell, and co-localization with the antigen. In one embodiment, the peptide tag of the present disclosure is possible not to have a sequence that imposes unique constraints on the distribution in a cell (or a sequence that prevents free distribution) of the intracellular antibody, and/or is possible not to have bond in a cell of the intracellular antibody to an antigen, and co-localization with the antigen.

[0122]

All the peptide tags of the present disclosure can mitigate, inhibit, or improve aggregation tendency; increase, promote, or improve a non-aggregation property;

or increase, promote, or improve stability of a tagged protein. The peptide tag of the present disclosure is possible not to have a sequence that imposes unique constraints on the distribution in a cell (or a sequence that prevents free distribution) of the intracellular antibody, and/or is possible not to have bond in a cell of the intracellular antibody to an antigen and co-localization with the antigen. When the protein of interest is an antigen-binding fragment of an antibody, the peptide tag of the present disclosure can promote co-localization with an antigen through bond of the protein to the antigen.

[0123]

The peptide tag of the present disclosure can be, for example, a gene product encoded by a gene of a living thing, or a fragment thereof, and here, can be a gene product encoded by a gene of a non-human living thing (for example, a microorganism such as a bacteria, an alga, or a fungus, an animal such as a mammal, a bird, or fish, or a plant), or a fragment thereof. Alternatively, in a preferable embodiment, the peptide tag of the present disclosure can be a gene product encoded by a gene of a human, or a fragment thereof.

[0124]

When fused with an aggregating protein, the peptide tag of the present disclosure can mitigate, inhibit or improve the aggregation tendency of the aggregating

protein, or can increase, promote, or improve the non-aggregation property of the aggregating protein. Aggregation of a protein can adversely affect a cell in which the protein is expressed, and in addition, can adversely affect the protein production amount and functionality by the aggregation. Accordingly, the mitigation, inhibition, or improvement of the aggregation tendency reduces the influence of the aggregation on the cell, and can lead to reduction of the influence on the protein production amount and the functionality. In this manner, the peptide tag of the present disclosure can be beneficial in improvement of the expression level of an aggregating protein expressed in a cell and/or improvement of the functionality, and accordingly, can be useful for forced expression of the aggregating protein *in vivo*.

[0125]

Accordingly, the peptide tag of the present disclosure may be fused with an aggregating protein. The aggregating protein can be a protein that forms aggregation in 30% or more, 40% or more, 50% or more, 60% or more, 70% or more, 80% or more, or 90% or more of cells when intracellularly expressed. The aggregating protein may be a protein that forms aggregation in 90% or less, 80% or less, 70% or less, 60% or more, or 50% or less of cells when intracellularly expressed. The fusion can be performed, for example, on the N terminal and/or

the C terminal (preferably, both the N terminal and the C terminal) of the aggregating protein. The fusion can be conducted by, for example, linking a nucleic acid encoding the peptide tag of the present disclosure and a nucleic acid encoding the aggregating protein, in-frame (in such a manner as to match the reading frames of codons). The peptide tag of the present disclosure may be fused with a non-aggregating protein. Thus, the non-aggregation property of the non-aggregating protein can be further increased. The non-aggregating protein may be a protein that forms aggregation in 20% or less, 15% or less, 10% or less, 9% or less, 8% or less, 7% or less, 6% or less, 5% or less, 4% or less, 3% or less, 2% or less, or 1% or less of cells or a protein that does not form aggregation, when intracellularly expressed.

[0126]

When fused with a protein, the peptide tag of the present disclosure can increase, promote, or improve the stability under an intracellular environment of the protein. The stability under an intracellular environment of a protein is beneficial in both an aggregating protein and a non-aggregating protein.

[0127]

In one embodiment, the protein of interest may be linked to a second peptide tag. The second peptide tag can be added to the protein of interest for purpose of, for example, detection or purification. In this case,

the peptide tag of the present disclosure can be used for reducing, inhibiting, or improving aggregation tendency; increasing, promoting, or improving a non-aggregation property; or increasing, promoting, or improving stability of a fusion protein of the protein of interest and the second peptide tag. As the second peptide tag, a usual peptide tag, such as an HA tag, can be used.

[0128]

The aggregating protein is not especially limited, and can be, for example, an antigen-binding fragment of an antibody. The aggregating protein can be preferably a single chain Fv (scFv) or a VHH antibody. A scFv is a fusion protein containing a heavy chain variable region and a light chain variable region of an antibody in which the heavy chain variable region and the light chain variable region are linked via a linker (preferably, a flexible linker). For example, there is an undruggable therapeutic target in a cell. This is conspicuous, for example, when a site for binding to a low molecular weight compound cannot be found in a therapeutic target. An antibody can bind to the target with strong binding affinity with different principles from the low molecular weight compound, and hence can effectively work on the therapeutic target regarded as undruggable with the low molecular weight compound. An antibody is, however, usually extracellularly secreted, and extracellularly functions. Therefore, in order to express, in a cell, a

secretory protein (protein extracellularly secreted), a gene can be designed to express a secretory protein (intracellular antibody) in a cell. For example, in order to express a secretory protein (intracellular antibody), a signal sequence of the protein can be disrupted, preferably removed or the like. In particular, a scFv can exhibit an aggregation property in a cell. Accordingly, the tag of the present disclosure can be fused with an aggregating protein, particularly a secretory protein, particularly an antigen-binding fragment of an antibody, and with preferably a scFv. A secretory protein exhibits an aggregation property in a cell in some cases. For stabilizing such a secretory protein before secretion in a cell, the tag of the present disclosure can be effective.

[0129]

The antibody, or the antigen-binding fragment of the antibody can have binding affinity (KD) to an antigen thereof of, for example, 10^{-5} M or less, 10^{-6} M or less, 10^{-7} M or less, 10^{-8} M or less, 10^{-9} M or less, 10^{-10} M or less, 10^{-11} M or less, or 10^{-12} M or less. A test of the binding property and a test of the binding affinity can be performed, for example, in a buffered saline.

[0130]

An example of the antigen includes an intracellular antigen such as an intracellular protein. Examples of the intracellular protein include an intracytoplasmic

protein (such as an intracellular extravesicular cytoplasmic protein), a nuclear protein (in this case, the peptide tag or the fusion protein may contain a nuclear localization signal), a nuclear transcription factor, a protein binding to a transcription factor, a protein binding to a genomic DNA, a protein binding to a protein binding to a genomic DNA, a constituent protein of chromatin, a protein binding to chromatin, an intracellular cell skeleton, and a protein binding to an intracellular cell skeleton. The intracellular protein is not especially limited, and other examples include a gene product of a cancer driver gene, a protein in an activated signal cascade (particularly in an activated immune cell of a patient having a cancer cell or an immune-related disease), and a gene product of a tumor suppressor gene (particularly under regulation of a binding partner for negative regulation thereof). In one embodiment, the antigen can be Kras.

[0131]

Introduction of the protein into a cell can be conducted by introducing, into the cell, a protein expression vector containing a nucleic acid encoding a fusion protein (a fusion protein of the peptide tag and the protein of interest) operably linked to a regulatory sequence. The protein expression vector is introduced into a protein-producing cell or a mammal cell, and can express the fusion protein in the cell.

[0132]

The regulatory sequence can be a promoter capable of transcribing an mRNA, and for example, various types of pol II promoters can be used. The pol II promoters are not especially limited, and examples include a CMV promoter, an EF1 promoter (EF1 α promoter), an SV40 promoter, an MSCV promoter, an hTERT promoter, a β actin promoter, a CAG promoter, and a CBh promoter. Further, a promoter driving bacteriophage-derived RNA polymerase, such as a T7 promoter, a T3 promoter, or an SP6 promoter, and a pol III promoter such as a U6 promoter can be used as the promoter. For a cyclic DNA, the T7 promoter can be preferably used, and for a linear DNA, the SP6 promoter can be preferably used. The promoter may be a promoter of a virus. Alternatively, the promoter may be an inducible promoter. The inducible promoter is a promoter capable of inducing expression of a polynucleotide functionally linked to the promoter only in the presence of an inducer driving the promoter. An example of the inducible promoter includes a promoter inducing gene expression by heat such as a heat shock promoter. Another example of the inducible promoter includes a promoter using a drug as the inducer driving the promoter. Examples of such a drug inducible promoter include a Cumate operator sequence, a λ operator sequence (such as 12 $\times$ λ Op), and a tetracycline-based inducible promoter. An example of the tetracycline-based inducible

promoter includes a promoter driving gene expression in the presence of tetracycline or a derivative thereof (such as doxycycline), or a reverse tetracycline controlled transactivator (rtTA). An example of the tetracycline-based inducible promoter includes a TRE3G promoter.

[0133]

The protein expression vector is not especially limited, and can be a virus vector or a plasmid vector. The virus vector is not especially limited, and examples include a retrovirus vector, a lentivirus vector, an adenovirus vector, an adeno-associated virus vector, a herpes simplex virus vector, a vaccinia virus vector, a Sendai virus vector, and a vesicular stomatitis virus vector. From the viewpoint of changing infectiveness to a cell, these vectors may be of pseudo type. These vectors may be derived from attenuated strains. Such a vector can be appropriately prepared by known technique.

[0134]

From the viewpoint of convenience in production of the protein expression vector, the protein expression vector may contain a nucleic acid encoding the regulatory sequence and the peptide tag of the present disclosure operably linked to the regulatory sequence, and have, on the downstream of the nucleic acid, a cloning site of a nucleic acid encoding the protein of interest. The cloning site has a restriction enzyme cleavage site

uniquely present in the vector, and is suitable for introducing a fragment of a gene encoding the protein of interest. A gene encoding the fusion protein of the peptide tag and the protein of interest is obtained by linking a gene encoding the protein of interest in-frame to a gene encoding the peptide tag. Accordingly, the present disclosure provides a protein expression vector containing a nucleic acid encoding the regulatory sequence and the peptide tag of the present disclosure operably linked to the regulatory sequence. In one preferable embodiment, this vector has, on the downstream of the nucleic acid, a cloning site of a nucleic acid encoding the protein of interest. The present disclosure provides a protein expression vector containing: a nucleic acid encoding the regulatory sequence and the peptide tag of the present disclosure operably linked to the regulatory sequence; and a nucleic acid encoding the protein of interest linked in-frame to the former nucleic acid. In this manner, the fusion protein of the peptide tag and the protein of interest can be expressed in a cell.

[0135]

The present disclosure provides a messenger RNA (mRNA) containing a nucleic acid encoding the peptide tag of the present disclosure. The mRNA further contains a nucleic acid encoding the protein of interest. The nucleic acid encoding the protein of interest is linked

in-frame to the nucleic acid encoding the peptide tag. In one embodiment, at least one or more uridines may be changed to pseudouridines in the mRNA. The pseudouridine can be 1-methyl-pseudouridine. The mRNA can be one transcribed from a cDNA, namely, may not have an intron. The mRNA may have a cap structure at the 5' end (Furuichi Y. & Miura K., *Nature*, 1975; 253 (5490): 374-5). As the cap structure, a Cap0 structure can be added to the mRNA by Anti-Reverse Cap Analogues (ARCA) method using a cap analogue (Stepinski J. et al., *RNA*, 2001 Oct; 7(10): 1486-95). When 2'-O methyltransferase treatment is further performed, the Cap0 structure of the mRNA can be converted to a Cap1 structure. Such an operation can be performed by an ordinary method, and can be practiced using, for example, a commercially available kit, such as ScriptCap m7G Capping System, ScriptCap 2'-O-Methyltransferase Kit, or T7 mScript Standard mRNA Production System (AR Brown Co., Ltd.). The mRNA may have a poly A chain. The addition of a poly A chain can be performed by an ordinary method, and can be performed, for example, with A-Plus Poly(A) Polymerase Tailing Kit (AR Brown Co., Ltd.). Accordingly, in one embodiment, the mRNA can be an mRNA that has a cap structure at the 5' end, has a poly A chain at the 3' end, and preferably has pseudouridine (preferably 1-methyl-pseudouridine) as at least a part of uridines. The mRNA can be an isolated mRNA or a synthesized mRNA.

[0136]

The mRNA can be encapsulated in a nanoparticle, such as a lipid nanoparticle (LNP). Thus, degradation of the mRNA in a living body is prevented, and efficiency of delivering the mRNA into a cell is improved.

Accordingly, in one embodiment, the mRNA can be an mRNA that has a cap structure at the 5' end, has a poly A chain at the 3' end, and preferably has pseudouridine (preferably 1-methyl-pseudouridine) as at least a part of uridines. Such a lipid nanoparticle encapsulating the mRNA is also provided. The lipid nanoparticle is not especially limited, and lipid nanoparticles described in, for example, US9364435B, US8822668B, US8802644B, and US8058069B2 can be used. Alternatively, the mRNA may be encapsulated in a polyion complex micelle, or a polyion complex polymersome (Miyata et al., Chem. Soc. Rev., 2012, 41, 2562-2574). A nanoparticle refers to a particle having a diameter (for example, a hydrodynamic diameter) less than 1 μm .

[0137]

Accordingly, the present disclosure provides a nanovesicle (such as a lipid nanovesicle, or a polyion complex polymersome) containing an mRNA at least containing a nucleic acid encoding the peptide tag of the present disclosure. The mRNA can contain a nucleic acid encoding the fusion protein of the peptide tag and the protein of interest.

[0138]

The present disclosure provides a method for reducing, inhibiting, or improving aggregation tendency of a protein, including fusing, to the protein, the peptide tag of the present disclosure that reduces, inhibits, or improves the aggregation tendency. In this embodiment, the protein can be an aggregating protein. The aggregation tendency can be aggregation tendency under an intracellular environment. The fusion is conducted usually on the N terminal and/or the C terminal of the protein. The method can be an *in vitro* method.

[0139]

The present disclosure provides a method for increasing, promoting, or improving a non-aggregation property a protein, including fusing, to the protein, the peptide tag of the present disclosure that reduces, inhibits, or improves the aggregation tendency. In this embodiment, the protein can be an aggregating protein. The non-aggregation property can be a non-aggregation property under an intracellular environment. The method can be an *in vitro* method.

[0140]

The present disclosure provides a method for increasing, promoting, or improving stability of a protein, including fusing, to the protein, the peptide tag of the present disclosure that reduces, inhibits, or improves aggregation tendency. In this embodiment, the

protein can be an aggregating protein. In this embodiment, the protein can be a non-aggregation property. The stability can be stability under an intracellular environment. The method can be an *in vitro* method.

[0141]

The present disclosure provides use of the peptide tag of the present disclosure for reducing, inhibiting, or improving aggregation tendency of a protein. The present disclosure also provides use of the peptide tag of the present disclosure for increasing, promoting, or improving a non-aggregation property of a protein. The present disclosure also provides use of the peptide tag of the present disclosure for increasing, promoting, or improving stability of a protein. The use can be use *in vitro*.

[0142]

It can be tested by *in vitro* assay how strongly the peptide tag of the present disclosure can reduce, inhibit, or improve the aggregation tendency of a protein; increase, promote, or improve the non-aggregation property; or increase, promote, or improve the stability. For example, a gene encoding the peptide tag of the present disclosure is fused, for introduction into a cell, to the N terminal or the C terminal of a gene encoding an aggregating protein such as a scFv having an amino acid sequence of SEQ ID NO: 1, and thus,

the aggregating protein fused with the peptide tag of the present disclosure can be expressed in the cell. An aggregation formed by the aggregating protein is observed with an antibody against the aggregating protein, and thus, a rate (%) of cells having aggregations to cells expressing the aggregating protein can be calculated. This rate can be used for evaluating influence of the peptide tag of the present disclosure on the aggregation tendency, the non-aggregation tendency, and the stability of the protein.

[0143]

The present disclosure provides a composition containing the peptide tag of the present disclosure. The peptide tag of the present disclosure can be linked to the protein of interest in a reaction solution by, for example, a click reaction. The click reaction can be a Huisgen reaction. One of the peptide tag and the protein of interest is modified with an alkyne and the other is modified with an azide compound, and thus, a 1,2,3-triazole ring is formed to obtain the link therebetween.

[0144]

The present disclosure provides a composition containing the fusion protein of the peptide tag of the present invention and the protein of interest. The protein of interest can be an aggregating protein in one embodiment. The protein of interest can be an antigen-

binding fragment of an antibody in one embodiment. The protein of interest can be a scFv in one embodiment.

[0145]

The present disclosure provides an mRNA containing a nucleic acid encoding the fusion protein of the peptide tag of the present disclosure and the protein of interest, and a composition containing the mRNA. The present disclosure provides a vesicle containing an mRNA containing a nucleic acid encoding the fusion protein of the peptide tag of the present disclosure and the protein of interest, and a composition containing the vesicle.

[0146]

The present disclosure provides a protein expression vector containing a nucleic acid encoding the fusion protein of the peptide tag of the present disclosure operably linked to a regulatory sequence and the protein of interest, and a composition containing the protein expression vector.

[0147]

In one embodiment, the composition may further contain a pharmaceutically acceptable carrier, excipient, and/or additive. The composition can be a pharmaceutical composition in one embodiment.

[0148]

In all embodiments of the present disclosure, the fusion protein having the peptide tag of the present disclosure linked thereto can be a non-natural protein.

[0149]

In all embodiments of the present disclosure, the peptide and the protein can be respectively recombinant peptide and protein.

[0150]

In all embodiments of the present disclosure, regarding a scFv having at least an amino acid sequence of SEQ ID NO: 1, the peptide tag of the present disclosure can reduce, inhibit, or improve aggregation tendency of the protein; increase, promote, or improve a non-aggregation property; or increase, promote, or improve stability.

[0151]

A peptide, a protein, and a nucleic acid can be isolated, concentrated, or purified. Isolation means that one or more components of a system are separated from a given component. Purification means that a relative concentration of a given component is increased as compared with a concentration of one or more other components of a system. Concentration means that a concentration of a given component is increased.

[0152]

One aspect of the present disclosure provides:

a method for acquiring (or selecting or identifying) an amino acid sequence (or a nucleic acid encoding the amino acid sequence), including acquiring an amino acid sequence in which:

(a) 5% or more and less than 45% of amino acids contained in the amino acid sequence are acidic amino acids; and

(b) 20% or more of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A.

[0153]

One aspect of the present disclosure provides:

a method for acquiring (or selecting or identifying) an amino acid sequence (or a nucleic acid encoding the amino acid sequence) including:

acquiring an amino acid sequence satisfying any one of conditions (A) to (AE) and (AF) to (AU) described above, or any combination of these conditions.

[0154]

In one aspect of the present disclosure, the method for acquiring an amino acid sequence may further include:

selecting or identifying an amino acid sequence of a peptide tag that, when a fusion protein of a peptide having the selected or identified amino acid sequence and a reference protein is expressed in a mammal cell (preferably in a human cell), provides reduction in a proportion of cells in which the fusion protein forms an aggregation (aggregation rate) (or the proportion which is not more than a predetermined value).

[0155]

In one aspect of the present disclosure, the method for acquiring may further include obtaining a peptide having the amino acid sequence, or a nucleic acid encoding the peptide.

[0156]

In the method for acquiring, the amino acid to be obtained may have a length of 10 to 200 amino acids (for example, a length of 10 to 90 amino acids).

[0157]

One aspect of the present disclosure provides a method for acquiring (or selecting or identifying) an amino acid sequence (or a nucleic acid encoding the amino acid sequence) having a length of 10 to 200 amino acids (10 to 90 amino acids), including:

acquiring an amino acid sequence in which:

(a) 5% or more and less than 45% of amino acids contained in the amino acid sequence are acidic amino acids; and

(b) 20% or more of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A;

Selecting or identifying an amino acid sequence of a peptide tag that, when the fusion protein of a peptide having the selected or identified amino acid sequence and a reference protein (reference protein) is expressed in a mammal cell (preferably in a human cell), provides reduction of a proportion of cells in which a fusion

protein forms an aggregation (aggregation rate) (or the proportion which is not more than a predetermined value); and

obtaining a peptide having the amino acid, or a nucleic acid encoding the peptide.

In this method, a peptide having a particularly excellent stabilizing action can be selected from peptides having a stabilizing action by increasing the extent of reduction or by reducing the predetermined value. In one aspect, the present method may further include expressing, in a mammal cell (preferably in a human cell), the fusion protein of the peptide having the selected or identified amino acid sequence and the reference protein. The present method may further include obtaining a nucleic acid encoding the selected or identified amino acid sequence. The predetermined value can be a numerical value based on a rate (%) of cells having an aggregation to cells expressing the aggregating protein. For example, the predetermined value can be a value of 30% or less, a value of 20% or less, a value of 15% or less, a value of 10% or less, a value of 5% or less, a value of 3% or less, a value of 2% or less, or a value of 1% or less, or 0%. The predetermined value can be a value in a range of 0% to 10%, a value in a range of 10% to 20%, or a value in a range of 20 to 30%. When a peptide having a higher effect is desired to be acquired, the predetermined value is preferably smaller. In this

embodiment, a peptide tag exhibiting a stronger effect can be selected from, for example, the above-described peptide tags of the present disclosure (for example, any one of the peptide tags of (A) to (Z), (AA) to (AE), and (AF) to (AU) described above).

[0158]

The reference protein can be, for example, a protein (aggregating protein, such as a scFv) in which the proportion of cells having an aggregation formed therein by the fusion protein is more than 30%, 40% to 50% or less, 50% to 60%, 60% to 70%, 70% to 80%, 80% to 90%, 90% to 95%, 95% to 99%, 99% to 99.9%, or 99.9% or more. The reference protein needs not have special functionality or binding property to an antigen but is used simply for evaluating the reduction of the aggregation rate, and therefore, the CDR sequence thereof may be any sequence. The aggregation rate of the scFv may be varied depending on the amino acid sequence of the CDR. It is possible to search for an amino acid sequence of a peptide tag that provides the aggregation rate not more than the predetermined value in the scFv having the varied aggregation rate. In one preferable embodiment, the reference protein can be a protein having an amino acid sequence of SEQ ID NO: 1.

[0159]

In one embodiment, the amino acid sequence to be obtained further satisfies the condition defined in (B)

described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (D) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (E) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (F) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (G) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (H) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (I) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (I) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (J) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (K) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (L) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (M) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (N)

described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (O) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (P) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (Q) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (R) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (S) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (T) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (U) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (V) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (W) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (X) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (Y) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (Z)

described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AA) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AB) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AC) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AD) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AE) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AF) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AG) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AH) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AI) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AJ) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AK) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AL)

described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AM) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AN) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AO) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AP) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AQ) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AR) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AS) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AT) described above. In one embodiment, the amino acid sequence to be acquired further satisfies the condition defined in (AU) described above. In one embodiment, the amino acid sequence to be obtained satisfies the condition described in any one or more of (A) to (Z), (AA) to (AE), and (AF) to (AU) described above.

[0160]

In one embodiment, the amino acid sequence to be acquired can be selected from an amino acid sequence group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (A) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (B) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (C) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (D) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (E) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (F) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (G) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (H) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (I) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (J) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (K) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (L) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (M) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (N) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (O) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (P) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (Q) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (R) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (S) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (T) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (U) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (V) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (W) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (X) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (Y) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (Z) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AA) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AB) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AC) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AD) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AE) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AF) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AG) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AH) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AI) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AJ) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AK) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AL) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AM) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AN) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AO) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AP) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AQ) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AR) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AS) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AT) above, or can include this group.

In one embodiment, the amino acid sequence group can be a group of the amino acid sequences defined in (AU) above, or can include this group.

[0161]

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the aggregation rate of the fusion protein is lower than this aggregation rate.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 1% or more.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 2% or more.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the aggregation rate of the fusion protein is not more than a

predetermined value, which is lower than this aggregation rate by 3% or more.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 4% or more.

In one preferable embodiment, the aggregation rate of the reference protein is 6 to 10%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 5% or more.

In one preferable embodiment, the aggregation rate of the reference protein is 7 to 10%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 6% or more.

In one preferable embodiment, the aggregation rate of the reference protein is 8 to 10%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 7% or more.

In one preferable embodiment, the aggregation rate of the reference protein is 9 to 10%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 8% or more.

[0162]

In a preferable embodiment, a ratio of the aggregation rate of the fusion protein to the aggregation rate of the reference protein is not more than a predetermined value, which is 0.9 or less, 0.8 or less, 0.7 or less, 0.6 or less, 0.5 or less, 0.4 or less, 0.3 or less, 0.2 or less, or 0.1 or less.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.9 or less.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.8 or less.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.7 or less.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.6 or less.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.5 or less.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.4 or less.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.3 or less.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.2 or less.

In one preferable embodiment, the aggregation rate of the reference protein is 5 to 10%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.1 or less.

[0163]

In one embodiment, the aggregation rate of the reference protein is less than 5%, and the aggregation rate of the fusion protein is lower than this aggregation rate.

In one preferable embodiment, the aggregation rate of the reference protein is less than 5%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.9 or less.

In one preferable embodiment, the aggregation rate of the reference protein is less than 5%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.8 or less.

In one preferable embodiment, the aggregation rate of the reference protein is less than 5%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.7 or less.

In one preferable embodiment, the aggregation rate of the reference protein is less than 5%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.6 or less.

In one preferable embodiment, the aggregation rate of the reference protein is less than 5%, and the ratio of the aggregation rate of the fusion protein to this

aggregation rate is not more than a predetermined value, which is 0.5 or less.

In one preferable embodiment, the aggregation rate of the reference protein is less than 5%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.4 or less.

In one preferable embodiment, the aggregation rate of the reference protein is less than 5%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.3 or less.

In one preferable embodiment, the aggregation rate of the reference protein is less than 5%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.2 or less.

In one preferable embodiment, the aggregation rate of the reference protein is less than 5%, and the ratio of the aggregation rate of the fusion protein to this aggregation rate is not more than a predetermined value, which is 0.1 or less.

[0164]

In one preferable embodiment, the aggregation rate of the reference protein is more than 10%, and the aggregation rate of the fusion protein is not more than a

predetermined value, which is lower than this aggregation rate by 10% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 20%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 10% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 20%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 20% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 30%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 10% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 30%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 20% or more.

[0165]

In one preferable embodiment, the aggregation rate of the reference protein is more than 30%, and the aggregation rate of the fusion protein is not more than a

predetermined value, which is lower than this aggregation rate by 30% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 40%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 10% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 40%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 20% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 40%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 30% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 40%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 40% or more.

[0166]

In one preferable embodiment, the aggregation rate of the reference protein is more than 50%, and the aggregation rate of the fusion protein is not more than a

predetermined value, which is lower than this aggregation rate by 10% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 50%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 20% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 50%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 30% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 50%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 40% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 50%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 50% or more.

[0167]

In one preferable embodiment, the aggregation rate of the reference protein is more than 60%, and the aggregation rate of the fusion protein is not more than a

predetermined value, which is lower than this aggregation rate by 10% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 60%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 20% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 60%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 30% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 60%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 40% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 60%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 50% or more.

[0168]

In one preferable embodiment, the aggregation rate of the reference protein is more than 60%, and the aggregation rate of the fusion protein is not more than a

predetermined value, which is lower than this aggregation rate by 60% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 70%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 10% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 70%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 20% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 70%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 30% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 70%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 40% or more.

[0169]

In one preferable embodiment, the aggregation rate of the reference protein is more than 70%, and the aggregation rate of the fusion protein is not more than a

predetermined value, which is lower than this aggregation rate by 50% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 70%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 60% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 70%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 70% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 80%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 10% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 80%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 20% or more.

[0170]

In one preferable embodiment, the aggregation rate of the reference protein is more than 80%, and the aggregation rate of the fusion protein is not more than a

predetermined value, which is lower than this aggregation rate by 30% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 80%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 40% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 80%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 50% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 80%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 60% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 80%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 70% or more.

[0171]

In one preferable embodiment, the aggregation rate of the reference protein is more than 80%, and the aggregation rate of the fusion protein is not more than a

predetermined value, which is lower than this aggregation rate by 80% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 90%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 10% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 90%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 20% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 90%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 30% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 90%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 40% or more.

[0172]

In one preferable embodiment, the aggregation rate of the reference protein is more than 90%, and the aggregation rate of the fusion protein is not more than a

predetermined value, which is lower than this aggregation rate by 50% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 90%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 60% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 90%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 70% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 90%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 80% or more.

In one preferable embodiment, the aggregation rate of the reference protein is more than 90%, and the aggregation rate of the fusion protein is not more than a predetermined value, which is lower than this aggregation rate by 90% or more.

[0173]

A nucleic acid encoding the obtained peptide tag is linked in-frame to a nucleic acid encoding the protein of interest, and thus, a nucleic acid encoding the fusion protein of the peptide tag and the protein of interest

can be obtained. From the nucleic acid encoding the fusion protein, the fusion protein can be expressed. A protein expression vector containing a nucleic acid encoding a fusion protein operably linked to a regulatory sequence may be prepared.

[0174]

In one aspect of the present disclosure, the method can be employed for determining whether or not a peptide tag having an amino acid sequence with a length of 10 to 200 amino acids (for example, 10 to 90 amino acids) has an effect of improving an aggregation property not less than a predetermined intensity against a tagged protein. In other words, in one aspect of the present disclosure, a method for determining whether or not a peptide tag having an amino acid sequence with a length of 10 to 90 amino acids has an effect of improving an aggregation property not less than a predetermined intensity against a tagged protein, and the method including:

acquiring an amino acid sequence in which:

(α) 5% or more and less than 45% of amino acids contained in the amino acid sequence are acidic amino acids; and

(β) 20% or more of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A;

determining that an amino acid sequence of a peptide tag that, when the fusion protein of a peptide having the

selected or identified amino acid sequence and a reference protein is expressed in a mammal cell (preferably in a human cell), provides reduction of a proportion of cells in which a fusion protein forms an aggregation (aggregation rate) (or the proportion which is not more than a predetermined value), has the effect of improving an aggregation property (for example, an aggregation property not less than a predetermined intensity).

In one embodiment, the method may further include expressing, in a mammal cell (preferably in a human cell), the fusion protein of the peptide having the selected or identified amino acid sequence and the reference protein. In one embodiment, the amino acid sequence to be acquired can be selected from an amino acid sequence group. The details of the amino acid group are the same as those described above. In one aspect, the amino acid sequence to be acquired satisfies another one or more conditions. The conditions are the same as those described above. In one embodiment, the aggregation rate of the reference protein and the predetermined value are the same as those described above.

[0175]

In one embodiment, in the method for selecting or identifying an amino acid sequence having a length of 10 to 200 amino acids (for example, 10 to 90 amino acids),

the amino acid sequence group having a length of 10 to 200 amino acids (for example, 10 to 90 amino acids) can be a group of amino acid sequences encoded by the coding region of the human genome. In one embodiment, the amino acid sequence group having a length of 10 to 90 amino acids can be a group of amino acid sequences encoded by the coding region of the genome of a non-human living thing.

[0176]

In one embodiment, the amino acid sequence having a length of 10 to 200 amino acids (for example, 10 to 90 amino acids) can be a neo-antigen. A neo-antigen was discovered as a mutant antigen newly caused by gene mutation peculiar to a cancer cell. The neo-antigen is not expressed in a non-cancer cell. It is expected that immunity can be induced specifically to cancer by inducing immunity to a neo-antigen by administering a peptide containing the neo-antigen. The neo-antigen can be different among cancer cells. The neo-antigen can be used, for example, for tagging a protein of interest to be expressed in a cell having the neo-antigen, and can be thus usable because it is a peptide originally expressed, and hence the cell is not or little adversely affected. The neo-antigen can be a naturally occurring mutant. The neo-antigen has one or more mutations selected from the group consisting of addition, insertion, deletion, and substitution in, for example, a wild type sequence

thereof. For example, a neo-antigen of a human typically has one or more (for example, 1 to 10, for example, 1 to several, 1 to 5, 1 to 4, 1 to 3, 2 or 1) mutations selected from the group consisting of addition, insertion, deletion, and substitution in a wild type sequence of a human. The neo-antigen of a human can have, for example, 80% or more identity, 85% or more identity, 90% or more identity, or 95% or more identity to the wild type sequence of a human.

[0177]

In one preferable embodiment, the reference protein is a scFv. In one embodiment, the scFv has the amino acid sequence of SEQ ID NO: 1.

[0178]

Still another aspect of the present invention provides a method for modifying an amino acid sequence of a peptide tag, including:

preparing a peptide tag (that may be any one of the peptide tags disclosed herein) for producing a fusion protein; and

obtaining a modified amino acid sequence by substituting, with either of P and N, one or more (preferably, a plurality of) amino acids of the Element 2, 3, or 4 (preferably, any one of the Element 2, the Element 4, and A, G, Y, and F).

The method of this aspect may further include:

determining that an amino acid sequence of a peptide tag that, when the fusion protein of a peptide having a selected or identified amino acid sequence and a reference protein is expressed in a mammal cell (preferably in a human cell), provides reduction of a proportion of cells in which a fusion protein forms an aggregation (aggregation rate) (or the proportion which is not more than a predetermined value), has the effect of improving an aggregation property (for example, an aggregation property not less than a predetermined intensity).

In this manner, modification having particularly strong aggregation reducing action can be performed.

Examples

[0179]

Example 1

[Method]

- Construction of Gene Expression Vector

A gene fragment encoding a fusion protein containing a peptide tag and an aggregating protein was produced by Eurofins Genomics K.K. or VectorBuilder Japan, Inc. The thus synthesized gene fragment was cloned into a pEF-BOS vector (Mizushima and Nagata, Nucleic Acids Res. 1990 Sep 11; 18(17): 5322). As the aggregating protein, a protein (specifically, a scFv) having an amino acid sequence set forth in SEQ ID NO: 1 was used. This aggregating protein

aggregates in the cytoplasm when expressed in a cell.
The sequences of tags used here and SEQ ID NOs thereof
were as shown in Table 1 below.

[0180]

[Table 1]

Table 1: List of Tags used in Experiment

Tag Name	Amino Acid Sequence	SEQ ID NO:
Tag-1-1	AHSSSAESESTSDSDSSSDSESESSSSDSEGS	2
Tag-1-2	AHSLSAELESTIDSDCSDWESELSSSDSEGS	3
Tag-1-3	AQSSSAEESGSDSDSSSDSESESSSSDSEGS	4
Tag-4-1	NEGYREAFDEYEQQDEDFAEQDPDGNEAFEGEYDGPNQDEYPDEAQNF	5
Tag-2-1	DEAGSSGAPADEAGSSGAPADEAGSSGAPADEAGSSGAPADEAGSSGAPAGS	6
Tag-2-2	DEVGISLAPTDEVGISLAPTDEVGISLAPTDEVGISLAPTDEVGISLAPTGS	7
Tag-2-3	DEVMISLWPTDEVMISLWPTDEVMISLWPTDEVMISLWPTDEVMISLWPTGS	8
Tag-3-1	DFAGSSGAPADFAGSSGAPADFAGSSGAPADFAGSSGAPADFAGSSGAPAGS	9
Tag-3-2	DTAVSSIAPLDTAVSSIAPLDTAVSSIAPLDTAVSSIAPLDTAVSSIAPLGS	10
Tag-3-3	DTWVSLIAILDWVSLIAILDWVSLIAILDWVSLIAILDWVSLIAILGS	11
Myc	EQKLISEEDL	12

[0181]

Specific content ratios of amino acids were as follows.

[0182]

[Table 2]

Table 2: Amino Acid Content Ratios in Each Peptide and Influence of Tag Addition on Intracellular Aggregation Formation of Protein

Tag Name	Element 1 (%)	Element 2 (%)	Element 3 (%)	Element 4 (%)	S (%)	G (%)	A (%)	Aggregation Rate (%)
Myc (control)	40.0	10.0	20.0	30.0	10.0	0	0	40.1
Tag-1-1	28.1	3.1	65.6	3.1	56.3	3.1	6.3	11.4
Tag-1-2	28.1	3.1	46.9	21.9	37.5	3.1	6.3	20.2
Tag-1-3	28.1	0.0	71.9	0.0	56.3	6.3	6.3	15.7
Tag-4-1	42.0	2.0	56.0	0.0	0.0	8.0	8.0	4.0
Tag-2-1	19.2	0.0	80.8	0.0	21.2	21.2	28.8	29.7
Tag-2-2	19.2	0.0	42.3	38.5	11.5	11.5	9.6	22.5
Tag-2-3	19.2	0.0	23.1	57.7	11.5	1.9	0.0	41.2
Tag-3-1	9.6	0.0	90.4	0.0	21.2	21.2	28.8	43.4
Tag-3-2	9.6	0.0	51.9	38.5	21.2	1.9	19.2	26.1
Tag-3-3	9.6	0.0	23.1	67.3	11.5	1.9	9.6	65.0

[0183]

In Table 2, Element 1 refers to D and E, Element 2 refers to H, K, and R, Element 3 refers to F, P, Y, G, S, Q, N, and A, and Element 4 refers to the other amino acids.

[0184]

The peptide tag was fused to the N terminal of an antibody fragment. The antibody fragment fused with the peptide tag was linked, for detection, to an HA tag of SEQ ID NO: 13 at the C terminal of the antibody fragment.

[0185]

- Measurement of Intracellular Aggregation Rate of Intracellular Antibody
(Cell Culture)

A HeLa cell derived from human cervical epithelial cancer was prepared. The HeLa cell was purchased from JCRB Cell Bank, National Institutes of Biomedical Innovation, Health and Nutrition (JCRB9004), and cultured in DMEM (D-MEM, FUJIFILM Wako Pure Chemical Corporation, 4548995066251) containing 10% FBS.

(Transfection)

In a 35 mm glass bottom dish (IWAKI 3911-035, glass hole, inner diameter: 12 mm) coated with poly-L-lysine (Sigma-Aldrich P1399 Poly-L-lysine hydrobromide mol. Wt. 150000-300000), 4×10^5 HeLa cells were plated, and after 24 hours, the antibody fragment gene described above was introduced into the HeLa cells with Lipofectamine 3000 (Invitrogen, L3000-008) based on a use method provided by the manufacturer, and thus, a tagged antibody fragment was expressed in the cells. A culture fluid was removed 24 hours after the transfection, and the resultant cells were fixed with 4% PFA. 20 minutes after the fixation, the resultant cells were washed with PBS(-), and thereafter, the antibody expressed in the cells (hereinafter, simply referred to as the "intracellular antibody") was observed by immunostaining to measure the aggregation rate thereof.

(Immunostaining)

The immunostaining was performed by a standard method. The cells were treated with 0.3% Triton X-100/PBS(-) for 2 minutes, and kept for 1 hour at room

temperature in a blocking solution (1% BSA, 0.1% Triton X-100/PBS(-)). The cells were kept for 2 hours at room temperature in an anti-HA antibody (rabbit anti-Ha antibody: Sigma-Aldrich, H6908) diluted with the blocking solution, the resultant cells were washed with the blocking solution, and then the resultant cells were kept at room temperature in Alexa Fluor 488 Goat anti-rabbit IgG (H+L) (Invitrogen, A11034) for 2 hours. The resultant cells were washed with the blocking solution, kept for 15 minutes at room temperature in a nuclear staining probe (NucBlue Fixed cell stain ReadyProbes, Invitrogen, R37606), washed with PBS(-), and stored at 4°C until fluorescence imaging.

(Fluorescence Imaging)

The fluorescence imaging was performed with Keyence BZ

-X700 or BZ-X800 using a 40× objective lens. An image including 200 or more cells per dish was acquired to count the number of cells having aggregation of the intracellular antibody therein (intracellular antibody aggregating cells). The number of the intracellular antibody aggregating cells was normalized with the total number of cells expressing the intracellular antibody to quantify an intracellular aggregation rate of the intracellular antibody.

[0186]

[Results]

The peptides having the amino acid sequences shown in Table 1 were fused with antigen-binding fragments (specifically, scFvs) of antibodies, and the thus obtained fusion proteins were expressed in the cytoplasm of the HeLa cell. The amino acid ratio in each peptide tag are shown in Table 2. The intracellular aggregation rates obtained based on fluorescence images were as shown in Table 2. In Table 2, a molecule provided with a Myc tag was used as a control, and the effect of each peptide tag was evaluated based on a difference in the aggregation rate from that of the Myc tag.

[0187]

As shown in Table 2,

- the aggregation rate tended to be reduced when the rate of the Element 3 was in a range of 40 to 75%;
- the aggregation rate tended to be reduced as the rate of the Element 4 was lower;
- the aggregation rate tended to be increased when the rate of alanine (A) was more than 10%;
- the aggregation rate tended to be increased when the rate of glycine (G) was more than 10%; and
- the intracellular aggregation rate tended to be reduced as the rate of the Element 1 was higher.

[0188]

Serine (S) may not be present (see Tag-4-1), but did not adversely affect the reduction of the aggregation

rate even present in a large amount (see Tag-1-1 to Tag-1-3).

[0189]

In either case, a peptide tag having an acidic amino acid content less than 45% did not exhibit non-specific adsorption (particularly, non-specific adsorption due to negative charge of the peptide tag) to an intracellular protein or the like. It was suggested that the peptide tag of the present disclosure is useful from the viewpoint that free intracellular localization of a protein of interest is not restricted peculiarly to the tag.

[0190]

It is understood, through comparison between the Myc tag (control) and Tag-4-1, that reduction of the Element 4 with increase of the Element 3 makes a strong contribution to the aggregation rate reducing action of the peptide tag. Similarly, it is understood, also through comparison between Tag-1-2 and Tag-1-1, that the reduction of the Element 4 with increase of the Element 3 makes a contribution to the aggregation rate reducing action of the peptide tag.

[0191]

In particular, when the rate of the Element 1 was 20% or more and less than 45%, that of the Element 2 was 10% or less, that of the Element 3 was 40 to 75%, that of the Element 4 was 10% or less, that of alanine was 10% or

less, and that of glycine was 10% or less, the aggregation rate was favorably reduced.

[0192]

The HeLa cell expressing the fusion protein of the aggregating protein and the tag (Tag4-1) was observed under a fluorescence microscope. As a negative control, a HeLa cell expressing an aggregating protein without Tag4-1 was observed under a fluorescence microscope. Results were as illustrated in Figure 1. As illustrated in Figure 1, when tagged with Tag4-1 (fused with Tag-4-1), the aggregating protein homogeneously distributed in the cytoplasm, but the aggregating protein without Tag4-1 formed an aggregation in the cytoplasm of most of cells.

[0193]

Example 2: Experiment of Adding Amino Acid to Tag

Amino acid contents in a tag and the aggregation rate of a tagged protein were further analyzed. Serine (S) was randomly inserted into or added to Tag4-1 or Tag11-1, and the effect of the thus obtained modified tags on the protein aggregation rate was examined. The experiment was conducted in the same manner as described in Example above except that the tags were different. Results were as shown in Table 3. In Table 3, added or inserted amino acids are underlined.

[0194]

[Table 3]

Table 3: Sequences, S Contents, D/E Contents, and Aggregation Rates of Modified Tags

Tag	sequence	Aggregation Rate	S (%)	D,E(%)
Tag4-1 (SEQ ID NO: 59)	NEGYREAFD <u>ED</u> YEQQDEDFAEQDPDGNEA FEGEYDGNPQDEYPDEAQNFE	4.57	0	42
Tag4-1-S10 (SEQ ID NO: 60)	NEGSYREAFD <u>SE</u> DYEQSQD <u>SE</u> DFSAEQDPD GN <u>SE</u> AEFG <u>SE</u> YSDGPN <u>SE</u> QDEYPDEAQN <u>SE</u>	5.06	16.7	35
Tag4-1-S20 (SEQ ID NO: 61)	NEGSYRE <u>SA</u> FD <u>SE</u> DYEQSQD <u>SS</u> EDFSAEQDS SPDGNSSEAFEGSSEYSDGPN <u>SE</u> QDEYPDEA <u>SS</u> QNFS <u>SS</u> SE	6.76	28.6	30
Tag11-1 (SEQ ID NO: 62)	YDNPYFEPQYGFPPPEDEDE	14.48	0	15
Tag11-1-S10 (SEQ ID NO: 63)	SYD <u>SN</u> P <u>SY</u> F <u>SE</u> PSQYGSFPP <u>SE</u> SE <u>SE</u> DE <u>SE</u>	11.29	33.3	10
Tag11-1-S20 (SEQ ID NO: 64)	<u>SS</u> YSD <u>SN</u> P <u>SY</u> F <u>SE</u> PS <u>SS</u> QYGSFPP <u>SE</u> SE <u>SS</u> <u>ES</u> D <u>SS</u> SE	18.77	50	7.5

[0195]

As shown in Table 3 as results, increase of S in a tag did not largely affect the aggregation rate of the tagged peptide. Although it is known that increase of acidic amino acids in a tag largely affects reduction of the aggregation rate of the tagged peptide (US2020/0157210A), on the contrary, the reduction of the rate of acidic amino acids did not largely reduce the aggregation rate in this example.

[0196]

Similarly, amino acid contents except for serine were changed to examine the influence of tagged peptides on the aggregation rate. In Tables 4 to 8, added or inserted amino acids are underlined.

[0197]

[Table 4]

Table 4: Sequences, Q Contents, D/E Contents, and Aggregation Rates of Modified Tags

Tag	sequence	Aggregation Rate	Q (%)	D,E(%)
Tag 4-1	NEGYREAFDEYEQQDEDFAEQDPDGNEAF EGEYDGNQDEYQDEAQNFE	4.88	10	42
Tag4-1-Q10 (SEQ ID NO: 65)	NEGQYREAFDQEDYEQQDQEDFQAEQDP DGNQAEFEGQYDGNQDEYQDEAQNFE QE	8.31	25	35
Tag4-1-Q20 (SEQ ID NO: 66)	NEGQYREAFDQEDYEQQDQEDFQAEQ DQQPDGNQAEFEGQYDGNQDEY PDEAQQQNFQQQE	8.40	35.7	30
Tag 11-1	YDNPYFEPQYGFPPPEEDEDE	16.59	5	15
Tag11-1-Q10 (SEQ ID NO: 67)	QYDQNPQYFQEPQQYGFPPQEQEDQEDQ E	26.70	36.7	10
Tag11-1-Q20 (SEQ ID NO: 68)	QQYQDQNPQYFQEPQQQYGFPPQEQE QEDQQEQDQQQE	17.27	52.5	7.5

[0198]

[Table 5]

Table 5: Sequences, N Contents, D/E Contents, and Aggregation Rates of Modified Tags

Tag	sequence	Aggregation Rate	N (%)	D,E(%)
Tag 4-1	NEGYREAFDEYEQQDEDFAEQDPDGNEAF EGEYDGNQDEYQDEAQNFE	4.88	8.00	42
Tag4-1-N10 (SEQ ID NO: 69)	NEGNYREAFDNEDYEQQDNEDFNAEQDP DGNNEAFEGNEYNDGNQDEYQDEAQNFE NE	6.12	23.33	35
Tag4-1-N20 (SEQ ID NO: 70)	NEGNYREAFDNEDYEQQDNEDFNAEQ DNNPDGNNEAFEGNNEYNDGNQDEYQ DEANNQNFNNNE	6.86	34.29	30
Tag 11-1	YDNPYFEPQYGFPPPEEDEDE	16.59	5.00	15
Tag11-1-N10 (SEQ ID NO: 71)	NYDNNPNYFNEPNQYGNFPPNEDNEDNE	18.56	36.67	10
Tag11-1-N20 (SEQ ID NO: 72)	NNYNDNNPNYFNEPNQNNYGNFPPNEN EDNNENDNNNE	17.68	52.50	7.5

[0199]

[Table 6]

Table 6: Sequences, P Contents, D/E Contents, and Aggregation Rates of Modified Tags

Tag	sequence	Aggregation Rate	P (%)	D,E(%)
Tag 4-1	NEGYREAFDEDEYEQQDEDFAEQDPDGNEAF EGEYDGPNQDEYPDEAQNFE	4.49	6.00	42
Tag4-1-P10 (SEQ ID NO: 73)	NEGPYREAFDEDEYEQQDEDFAEQDPDG NPEAFEGPEYEDGPNQDEYPDEAQNFE	8.84	21.67	35
Tag4-1-P20 (SEQ ID NO: 74)	NEGPYREAFDEDEYEQQDEDFAEQDPDG PPDGNPPEAFEGPEYEDGPNQDEYPDEAP EQNFPPPE	6.44	32.86	30
Tag 11-1	YDNPYFEPQYGFPPPEEDEDE	17.75	20.00	15
Tag11-1-P10 (SEQ ID NO: 75)	EYDNPPEYFPEPEQYGFPPPEDEDEDE	15.57	46.67	10
Tag11-1-P20 (SEQ ID NO: 76)	PPYEDNPPEYFPEPEQYGFPPPEDEDE PEPDPPPE	18.12	60.00	7.5

[0200]

[Table 7]

Table 7: Sequences, F Contents, D/E Contents, and Aggregation Rates of Modified Tags

Tag	sequence	Aggregation Rate	F (%)	D,E(%)
Tag 4-1	NEGYREAFDEDEYEQQDEDFAEQDPDGNEAF EGEYDGPNQDEYPDEAQNFE	4.90	8.00	42
Tag4-1-F10 (SEQ ID NO: 77)	NEGEYREAFDEDEYEQQDEDFAEQDPDG NEEAFEGEEYEDGPNQDEYPDEAQNFE	35.13	34.29	35
Tag4-1-F20 (SEQ ID NO: 78)	NEGEYREAFDEDEYEQQDEDFAEQDFF PDGNFEEAFEGEEYEDGPNQDEYPDEAEE QNFEEEE	38.76	23.33	30
Tag 11-1	YDNPYFEPQYGFPPPEEDEDE	16.59	10.00	15
Tag11-1-F10 (SEQ ID NO: 79)	EYDENPEYFEEPEQYGFPPPEEEDFEEDE	43.58	55.00	10
Tag11-1-F20 (SEQ ID NO: 80)	FFYEDENPEYFEEPEEQYGFPPPEEEDFEE EDFEE	27.43	40.00	7.5

[0201]

[Table 8]

Table 8: Sequences, Y Contents, D/E Contents, and Aggregation Rates of Modified Tags

Tag	sequence	Aggregation Rate	Y (%)	D,E(%)
Tag 4-1	NEGYREAFDEYEQQDEDFAEQDPDGNEAF EGEYDGPNQDEYPDEAQNFE	4.94	8.00	42
Tag4-1-Y10 (SEQ ID NO: 81)	NEGYYREAFDYEDYEQYQDYEDFYAEQDPDG NYEAFEGYFYDGPNYQDEYPDEAQNFE	29.59	23.33	35
Tag4-1-Y20 (SEQ ID NO: 82)	NEGYYREYAFDYEDYEQYQDYEDFYAEQDY PDGNYEAFEGYFYDGPNYQDEYPDEAY QNFYYE	37.93	34.29	30
Tag 11-1	YDNPYFEPQYGFPPPEDEDE	17.75	15.00	15
Tag11-1-Y10 (SEQ ID NO: 83)	YYDYNPYFYFEPYQYGYFPPYFYEDYEDYE	17.28	43.33	10
Tag11-1-Y20 (SEQ ID NO: 84)	YYYDYNPYFYFEPYQYGYGYFPPYFYEDYY EYDYYE	30.53	57.50	7.5

[0202]

As described above, when the influence on the protein aggregation rate of tags obtained by randomly adding or inserting 10 to 20 specific amino acids was examined, the addition or insertion of F and Y tended to worsen the aggregation rate of the tagged proteins. The adverse effect of the addition or insertion of the other amino acids on the aggregation rates of tagged proteins was restrictive even if D and E contents were reduced.

[0203]

Example 3: Effect on Aggregation Rate of Protein by Amino Acid Content Change by Substitution of Amino Acids in Tag

N, P, S, Q, F, and Y were respectively substituted with other amino acids, and thus, attempts were made to specify, among these amino acids, an amino acid

exhibiting the effect on the protein aggregation rate.

In Tables 9 and 10, substituted amino acids are underlined.

[0204]

[Table 9]

Table 9: Sequences, N Contents, D/E Contents, and Aggregation Rates of Modified Tags

N-terminus	scFv	C-terminus	Cell	sequence	Aggregation Rate	N (%)	D, E (%)
Tag 9-1	Y13-259	HA	HeLa	GNNQSSSDNEADEASDDDEDNDGN (SEQ ID NO: 85)	8.14	20.00	44.00
Tag 4-8	Y13-259	HA	HeLa	NEGNREASDEDSEQQDEDNAEQDPDGNEANE GESDGPNQDENPDEAQNNE(SEQ ID NO: 86)	3.67	16.00	42.00
Tag 4-9	Y13-259	HA	HeLa	NEGNREANDEDNEQQDEDNAEQDPDGNEANE GENDGPNQDENPDEAQNNE(SEQ ID NO: 87)	4.10	24.00	42.00
Tag 4-10	Y13-259	HA	HeLa	NEGNREANDEDNEQQDEDNAEQDPDGNEANE GENDGPNQDENPDEAQNNE(SEQ ID NO: 88)	5.51	32.00	42.00
Tag 4-11	Y13-259	HA	HeLa	NEGNREANDEDNEQQDEDNAEQDPDGNEANE GENDGPNQDENPDEAQNNE(SEQ ID NO: 89)	4.44	32.00	42.00
Tag 18-1	Y13-259	HA	HeLa	DNNESADDNNENPEDNNKNTDDNEENPNNNEN (SEQ ID NO: 90)	4.82	43.75	37.50
Tag 4-8	6E	Tag 4-8, CMA, HA	SHSY5Y	NEGNREASDEDSEQQDEDNAEQDPDGNEANE GESDGPNQDENPDEAQNNE	0.96	16.00	42.00
Tag 18-1	6E	Tag 18-1, CMA, HA	SHSY5Y	DNNESADDNNENPEDNNKNTDDNEENPNNNEN	1.14	43.75	37.50

[0205]

[Table 10]

Table 10: Sequences, P Contents, N Contents, and Aggregation Rates of Modified Tags

		Aggregation Rate	N (%)	P (%)
Tag 18-1	DNNESADDNNENPEDNNKNTDDNEENPNNNEN	4.76	43.75	6.25
Tag 18-1-NS7 (SEQ ID NO: 91)	DNSESADDNSENPEDNSKNTDDSEENPSNSES	6.37	21.88	6.25
Tag 18-1-NS14 (SEQ ID NO: 92)	DSSESADDSSSPEDSSKSTDDSEESPSSSES	5.72	0.00	6.25
Tag 18-NQ7 (SEQ ID NO: 93)	DNQESADDNQENPEDNQKNTDDQEENPQNQEQ	6.60	21.88	6.25
Tag 18-NQ14 (SEQ ID NO: 94)	DQGESADDQEQPEDQKQKTDQEEQPQQQEQ	6.03	0.00	6.25
Tag 18-1-NF7 (SEQ ID NO: 95)	DNFESADDNFENPEDNFKNTDDFEENPFNFEF	13.50	21.88	6.25
Tag 18-1-NF14 (SEQ ID NO: 96)	DNFESADDNFEENPEDNFKNTDDFEENPFNFEF	13.60	0.00	6.25
Tag 18-1-NP7 (SEQ ID NO: 97)	DNPEESADDNPEENPEDNPKNTDDPEENPPNPEP	4.48	21.88	28.13
Tag 18-1-NP14 (SEQ ID NO: 98)	DNPEESADDNPEENPEDNPKNTDDPEENPPNPEP	3.18	0.00	50.00
Tag 18-1-NY7 (SEQ ID NO: 99)	DNYESADDNYENPEDNYKNTDDYEENPYNYEY	11.60	21.88	6.25
Tag 18-1-NY14 (SEQ ID NO: 100)	DYYESADDYEEYPEDYYKYTDDYEEYPYYEY	15.70	0.00	6.25
Tag 11-1	YDNPYFEPQYGFPPPEEDEDE	17.75	5.00	20.00
Tag 11-1-FYPN (SEQ ID NO: 101)	NDNNNNENQNGNNEEDEDE	12.46	50.00	0.00
Tag 11-1-FYN (SEQ ID NO: 102)	NDNPNNEPQNGNPPPEEDEDE	5.79	30.00	20.00
45Tag1 (SEQ ID NO: 1041)	NDEYSDFEDSFDGDYKDSDEYKDDSENFDGFE	13.91	3.71	0
45Tag1-m1 (SEQ ID NO: 1042)	PDEPPDPEDPDPPDPKDPDEDPKDDPEPPDDPPE	5.70	0	42.86

[0206]

It was found, based on Tables 9 and 10, that the aggregation rate reducing action of a tagged protein is the greatest when the content of P or N in the tag was higher, and subsequently, the aggregation rate reducing action was exhibited in the order that the content of S and Q was higher, and the content of F and Y was higher. 45Tag1-m1 peptide tag, which was produced by substituting, with P, all of the Element 3 in 45Tag1 having an acidic amino acid content of about 51%, largely improved the aggregation inhibiting action thereof through the amino acid substitution with P. In this manner, it was revealed that the increase of the aggregation rate reducing action by adding P and N does

not depend on the acidic amino acid content. This result shows that most of constituent amino acids can be acidic amino acids, and N or P.

[0207]

Example 4: Sequence Shuffle

Two tags, Tag4-8 (Scr1) and Tag4-8 (Scr2), were synthesized by randomly shuffling the amino acid sequence of Tag4-8, and were tested for the aggregation rate reducing action against the scFv in the same manner as described above. Results were as shown in Table 11.

[0208]

[Table 11]

Table 11: Sequence Shuffling

	Sequence	Aggregation Rate
Tag4-8	NEGNREASDEDSEQQDEDNAEQDPD GNEANEGESDGPNQDENPDEAQNSE	3.93
Tag4-8 (Scr1) (SEQ ID NO: 103)	ENESEDNDEEPPNQNADDGDPNANP AAEQGGDSSDDEEGQENDQSRENEQ	3.25
Tag4-8 (Scr2) (SEQ ID NO: 104)	QNEENGGDDQDQEEGSEEQQGESDRS NEESDNEPAADDNAPEGAEEPDDNN	4.52

[0209]

As shown in Table 11, Tag4-8, Tag4-8 (Scr1) and Tag4-8 (Scr2) exhibited equivalent aggregation rate reducing actions. Even when only the order of amino acids was changed without changing the content ratios and the lengths of the amino acids, the aggregation rate reducing action was not affected.

[0210]

Example 5: Use as Tag of Human-derived Peptide

From human proteome database (Proteome ID: UP000005640), peptides having specific amino acid content ratios were all extracted, these peptides were randomly selected to be used as tags, and thus, the aggregation rate reducing action against a tagged protein was examined.

[0211]

Extraction Condition 1:

length: 20 to 70 amino acids

group [D, E]: content of [30] or more

group [D, E]: content of less than [45]

group [H, K, R]: content of [5] or less

group [C, T, V, L, I, W, M]: content of [5] or less

group [G]: content of less than [10]

group [A]: content of less than [10]

group [F, Y]: content of [5] or less

group [N]: content of [15] or more

* In the above-described extraction condition, the unit of each content is %. The amino acids are described by one letter codes.

[0212]

[Table 12]

Table 12: Aggregation Rates of scFvs having Tags Extracted under Extraction Condition 1 Added

Tag Name	Sequence	Aggregation Rate %	DE Rate	N Rate	HKR Rate	CTVLIWM Rate	G Rate	A Rate	FyRate
Tag 18-1(465-1)	DNNESADDNNENPEDNNKNTDDNEENPNNNEN	4.68	37.50	43.75	3.13	3.13	0	3.13	0
Tag 465-2	GNNQDSSDSDNEADEASDDDEDNDGN	5.04	44.00	20	0	0	8	8	0
Tag 465-3	SHGNNQDSSDSDNEADEASDDDEDNDGNEGDNE	3.98	43.75	18.75	3.13	0	9.38	6.25	0
Tag 465-4	ESADDNNENPEDNNKNTDDNE	9.99	42.86	33.33	4.76	4.76	0	4.76	0
Tag 465-5	NNENPEDNNKNTDDNEENPNNN	13.94	33.33	47.62	4.76	4.76	0	0	0
Tag 465-6	TDNNESADDNNENPEDNNKNT	8.95	35.00	40	5	5	0	5	0
Tag 465-7	DNNESADDNNENPEDNNKNT	7.77	35.00	40	5	5	0	5	0
Tag 465-8	NNESADDNNENPEDNNKNT	7.39	35.00	40	5	5	0	5	0
Tag 465-9	FWGSHNNQDSSDSDNEADEASDDDEDNDGNE	7.36	38.71	16.13	3.23	3.23	9.68	6.45	3.23
Tag 1121-1	KPNNSNAPNEDQEEIIQQSE	14.73	30	20	5	5	0	5	0
Tag 1121-2	NNSNAPNEDQEEIIQQSEGH	13.76	30	20	5	5	0	5	0
Tag 1121-3	SEGEQQLKPNNSNAPNEDQEEI	14.15	31.82	18.18	4.55	4.55	4.55	4.55	0
Tag 2408-1	EQLNFSDDDEQGSNSPKENNSDQ	7.56	33.33	16.67	4.17	4.17	4.17	0	4.17
Tag 2408-2	EQLNFSDDDEQGSNSPKENNSDQ	7.53	32.00	16	4	4	8	0	4
Tag 2408-3	EQLNFSDDDEQGSNSPKENNSDQGS	9.42	30.77	15.38	3.85	3.85	7.69	0	3.85
Tag 6301-1	EKENDENSLSSSDSDSED	12.66	40.00	15	5	5	0	0	0
Tag 6301-2	NENDENSLSSSDSDSEDKDE	8.85	40.00	15	5	5	0	0	0
Tag 6626-1	KETNNNAQNPSEEEGEGQDE	12.50	33.33	19.05	4.76	4.76	9.52	4.76	0
Tag 6626-2	ETNNNAQNPSEEEGEGQDED	8.04	38.10	19.05	0	4.76	9.52	4.76	0
Tag 6626-3	KETNNNAQNPSEEEGEGQDED	13.56	36.36	18.18	4.55	4.55	9.09	4.55	0
Tag 6915-1	ENANDSSDSDSEETDESFPN	9.52	40.00	15	0	5	5	5	5
Tag 7128-1	DDNESNESAENGWDSGSNFSEE	12.10	34.78	17.39	0	4.35	8.7	4.35	4.35
Tag 7128-2	SDDNESNESAENGWDSGSNFSEE	8.21	33.33	16.67	0	4.17	8.33	4.17	4.17
Tag 7128-3	SDDNESNESAENGWDSGSNFSEE	8.06	32.00	16	0	4	8	4	4
Tag 7315-1	EENASSGDSEENTNSDHESE	10.74	40.00	15	5	5	5	5	0
Tag 7315-2	SEENASSGDSEENTNSDHESE	11.72	35.00	15	5	5	5	5	0
Tag 7315-3	ENASSGDSEENTNSDHESEQ	10.07	35.00	15	5	5	5	5	0
Tag 8482-1	DESENENWRNYPEEESDSDG	8.17	42.86	19.05	4.76	4.76	4.76	0	4.76
Tag 8482-2	ENSENNWRNYPEEESDSDGDE	7.11	42.86	19.05	4.76	4.76	4.76	0	4.76
Tag 8482-3	NSENNWRNYPEEESDSDGDESD	5.08	40.91	18.18	4.55	4.55	4.55	0	4.55
Tag 8974-1	EQQNEASEENNDDQQSQEVPE	7.84	35.00	15	0	5	0	5	0
Tag 8974-2	QQNEASEENNDDQQSQEVPEK	6.39	30.00	15	5	5	0	5	0
Tag 9333-1	MQEDEFDQGNQGEQEDNSNAE	15.37	40.00	15	0	5	5	5	5
Tag 9333-2	SKMQEDEFDQGNQGEQEDNSN	9.62	35.00	15	5	5	5	0	5
Tag 9333-3	QDEFDQGNQGEQEDNSNAEENASN	7.13	40.74	18.52	0	3.7	3.7	7.41	3.7
Tag 10381-1	PSENNQSSEDVGGDNDSEN	5.59	33.33	19.05	0	4.76	9.52	0	0
Tag 11717-1	EVEENPSAKEDSNPNSSGE	10.48	30.00	15	5	5	5	5	0
Tag 11717-2	VEENPSAKEDSNPNSSGED	12.10	30.00	15	5	5	5	5	0
Tag 12237-1	KEENSEPLNENSDSEYSEE	9.98	40.00	15	5	5	0	0	5
Tag 12809-1	QPGPNHEEDADSYENMDNPD	28.15	35.00	15	5	5	5	5	5
Tag 12809-2	PNHEEDADSYENMDNPDGPD	20.94	40.00	15	5	5	5	5	5
Tag 12809-3	NHEEDADSYENMDNPDGPD	23.41	40.00	15	5	5	5	5	5
Tag 12885-1	DDPNSSDESNGNDANSESD	14.62	40.00	20	0	0	5	5	0
Tag 12885-2	NSSDESNGNDANSESDNNS	17.00	30.00	30	0	0	5	5	0
Tag 12885-3	DESNGNDANSESDNNSSSRGD	13.93	31.82	22.73	4.55	0	9.09	4.55	0
Tag 12968-1	DNNEAGEDGDNDFSPSDEEL	13.42	42.86	19.05	0	4.76	9.52	4.76	4.76
Tag 12968-2	AEEEDDNNENAGEDGDNDFSPS	10.18	43.48	17.39	0	4.35	8.7	8.7	4.35
Tag 12968-3	DNNEAGEDGDNDFSPSDEELAN	14.63	39.13	21.74	0	4.35	8.7	8.7	4.35
Tag 13648-1	NPADDPNNQGEDEFEEAEQVREEN	15.03	41.67	16.67	4.17	4.17	4.17	8.33	4.17
Tag 14056-1	NEENTEPGAESSENADDPNKD	13.22	38.10	19.05	4.76	4.76	4.76	9.52	0
Tag 14056-2	SSENADDPNKDTSENADGQSDEN	11.45	34.78	17.39	4.35	4.35	4.35	8.7	0
Tag 14056-3	ESSENADDPNKDTSENADGQSDEN	10.38	37.50	16.67	4.17	4.17	4.17	8.33	0
Tag 14681-1	DRDPEMENEQPSSENDSON	5.63	40.00	15	5	5	0	0	0
Tag 14681-2	PEMENEQPSSENDSONQSG	11.50	30.00	15	0	5	5	0	0
Tag 14681-3	ENEQPSSENDSONQSGEQI	17.15	30.00	15	0	5	5	0	0
Tag 14844-1	DSESANVSDKEAGSNENDQGN	12.13	33.33	19.05	4.76	4.76	4.76	9.52	0
Tag 15481-1	NYNDGSGEDRDWQDDQSDNQ	9.92	35.00	15	5	5	5	0	5
Tag 16043-1	RENTNEASSEGNSDDSEDE	13.45	40.00	15	5	5	5	5	0
Tag 16043-2	ENTNEASSEGNSDDSEDER	11.56	40.00	15	5	5	5	5	0
Tag 16400-1	QENDNGNETESEQPKESNENQ	15.9	33.33	23.81	4.76	4.76	4.76	0	0
Tag 16400-2	ENDNGNETESEQPKESNENQE	18.1	38.1	23.81	4.76	4.76	4.76	0	0
Tag 16400-3	QENDNGNETESEQPKESNENQE	8.84	36.36	22.73	4.55	4.55	4.55	0	0
Tag 16417-1	QNEENPGDEEAKNQVNSSDSDSEE	12.18	40.00	10	4	4	4	4	0
Tag 16417-2	NEENPGDEEAKNQVNSSDSDSEES	11.00	40.00	16	4	4	4	4	0
Tag 16417-3	QNEENPGDEEAKNQVNSSDSDSEES	12.37	38.46	15.38	3.85	3.85	3.85	3.85	0
Tag 18137-1	YNGGNANPRPANNEEEDEEYD	7.47	40.91	22.73	4.55	0	9.09	9.09	4.55
Tag 18137-2	NGGNANPRPANNEEEDEEYD	8.17	40.91	22.73	4.55	0	9.09	9.09	4.55
Tag 18137-3	NGGNANPRPANNEEEDEEYD	9.86	43.48	21.74	4.35	0	8.7	8.7	4.35
Tag 18347-1	GASENEEDDDYKPLDPNS	16.06	40.00	15	5	5	5	5	5
Tag 18478-1	LQNKAEAEPPGPDSENSQEN	16.82	30.00	15	5	5	5	5	0
Tag 18478-2	QNKAEAEPPGPDSENSQENP	14.77	30.00	15	5	0	5	5	0
Tag 18478-3	NQKAEAEPPGPDSENSQENPP	12.74	30.00	15	5	0	5	5	0
Tag 20166-1	KESVSPENNEEGGNDNODNEN	16.95	33.33	28.57	4.76	4.76	9.52	0	0
Tag 20166-2	ESVSPENNEEGGNDNODNENP	13.67	33.33	28.57	0	4.76	9.52	0	0
Tag 20166-3	KESVSPENNEEGGNDNODNENP	26.56	31.82	27.27	4.55	4.55	9.09	0	0
Tag 41693-1	ASPPPREPSDDENSDNSNEC	7.83	30.00	15	5	5	0	5	0
Tag 55443-1	NNSQDEDGFQELNENGNAKDE	19.76	33.33	23.81	4.76	4.76	9.52	4.76	4.76
Tag 55443-2	NSQDEDGFQELNENGNAKDEN	22.25	33.33	23.81	4.76	4.76	9.52	4.76	4.76
Tag 55443-3	NNSQDEDGFQELNENGNAKDEN	18.69	31.82	27.27	4.55	4.55	9.09	4.55	4.55

[0213]

In Table 12, the N content ratio in the amino acid sequences of the extracted tags is 15% or more, and the content ratios of the other amino acids are as described above in Extraction Condition 1. In order to confirm that the aggregation inhibiting action does not depend on a specific amino acid sequence, tags were randomly selected from the tags extracted under Extraction Condition 1. As for some tags, an amino acid sequence satisfying the extraction condition was additionally selected from another portion of the same protein. The aggregation rates of scFvs tagged with the selected amino acid sequences were tested, and results as shown in Table 12 were obtained. As shown in Table 12, the aggregation rates of the tagged scFvs were low as a whole. Accordingly, it is obvious that the aggregation inhibiting action of a peptide tag largely depends on the amino acid content ratios thereof, and depends merely weakly on the specific amino acid sequence itself.

[0214]

It is noted that human-derived amino acid sequences that can be extracted under Extraction Condition 1 were as follows.

[Table 12-2]

Table 12-2: Examples of human-derived amino acid sequences that can be extracted under

Extraction Condition 1

SEQ ID NO:	Sequence
105	TDNNEASADDNNENPEDNNKN (Tag465-6)
106	DNNEASADDNNENPEDNNKNT (Tag465-7)
107	NNESADDNNENPEDNNKNTD (Tag465-8)
108	NESADDNNENPEDNNKNTDD
109	ESADDNNENPEDNNKNTDDN
110	SADDNNENPEDNNKNTDDNE
111	DNNEPEDNNKNTDDNEENP
112	NNENPEDNNKNTDDNEENPN
113	NENPEDNNKNTDDNEENPNN
114	ENPEDNNKNTDDNEENPNNN
115	NPEDNNKNTDDNEENPNNNE
116	PEDNNKNTDDNEENPNNNEN
117	DNNEASADDNNENPEDNNKNTD
118	NNESADDNNENPEDNNKNTDD
119	NESADDNNENPEDNNKNTDDN
120	ESADDNNENPEDNNKNTDDNE (Tag 465-4)
121	SADDNNENPEDNNKNTDDNEE
122	ADDNNENPEDNNKNTDDNEEN
123	DDNNENPEDNNKNTDDNEENP
124	DNNEPEDNNKNTDDNEENPN
125	NNENPEDNNKNTDDNEENPNN (Tag 465-5)
126	NENPEDNNKNTDDNEENPNNN
127	ENPEDNNKNTDDNEENPNNNE
128	NPEDNNKNTDDNEENPNNNEN
129	DNNEASADDNNENPEDNNKNTDD
130	NNESADDNNENPEDNNKNTDDN
131	NESADDNNENPEDNNKNTDDNE
132	SADDNNENPEDNNKNTDDNEEN
133	ADDNNENPEDNNKNTDDNEENP
134	DDNNENPEDNNKNTDDNEENPN
135	DNNEPEDNNKNTDDNEENPNN
136	NNENPEDNNKNTDDNEENPNNN
137	NENPEDNNKNTDDNEENPNNNE
138	ENPEDNNKNTDDNEENPNNNEN
139	DNNEASADDNNENPEDNNKNTDDN
140	NNESADDNNENPEDNNKNTDDNE
141	NESADDNNENPEDNNKNTDDNEE
142	ESADDNNENPEDNNKNTDDNEEN
143	SADDNNENPEDNNKNTDDNEENP
144	ADDNNENPEDNNKNTDDNEENPN
145	DDNNENPEDNNKNTDDNEENPNN
146	DNNEPEDNNKNTDDNEENPNNN
147	NNENPEDNNKNTDDNEENPNNNE
148	NENPEDNNKNTDDNEENPNNNEN
149	HGNNQDSSSDNEADEASDDEDN
150	DNNEASADDNNENPEDNNKNTDDNE
151	NNESADDNNENPEDNNKNTDDNEE
152	NESADDNNENPEDNNKNTDDNEEN
153	ESADDNNENPEDNNKNTDDNEENP
154	SADDNNENPEDNNKNTDDNEENPN
155	ADDNNENPEDNNKNTDDNEENPNN
156	DDNNENPEDNNKNTDDNEENPNNN
157	DNNEPEDNNKNTDDNEENPNNNE
158	NNENPEDNNKNTDDNEENPNNNEN
159	SHGNNQDSSSDNEADEASDDEDN
160	DNNEASADDNNENPEDNNKNTDDNEE
161	NNESADDNNENPEDNNKNTDDNEEN
162	NESADDNNENPEDNNKNTDDNEENP
163	ESADDNNENPEDNNKNTDDNEENPN
164	SADDNNENPEDNNKNTDDNEENPNN
165	ADDNNENPEDNNKNTDDNEENPNNN
166	DDNNENPEDNNKNTDDNEENPNNNE
167	DNNEPEDNNKNTDDNEENPNNNEN
168	GSHGNNQDSSSDNEADEASDDEDN
169	SHGNNQDSSSDNEADEASDDEDND
170	HGNNQDSSSDNEADEASDDEDNDG
171	GNNQDSSSDNEADEASDDEDNDGN (Tag 465-2)
172	DNNEASADDNNENPEDNNKNTDDNEEN
173	NNESADDNNENPEDNNKNTDDNEENP
174	NESADDNNENPEDNNKNTDDNEENPN
175	ESADDNNENPEDNNKNTDDNEENPNN
176	SADDNNENPEDNNKNTDDNEENPNNN
177	ADDNNENPEDNNKNTDDNEENPNNNE
178	DDNNENPEDNNKNTDDNEENPNNNEN
179	WGSNGNNQDSSSDNEADEASDDEDN
180	GSHGNNQDSSSDNEADEASDDEDND
181	SHGNNQDSSSDNEADEASDDEDNDG
182	HGNNQDSSSDNEADEASDDEDNDGN
183	DNNEASADDNNENPEDNNKNTDDNEENP
184	NNESADDNNENPEDNNKNTDDNEENPN
185	NESADDNNENPEDNNKNTDDNEENPNN
186	ESADDNNENPEDNNKNTDDNEENPNNN
187	SADDNNENPEDNNKNTDDNEENPNNNE
188	ADDNNENPEDNNKNTDDNEENPNNNEN
189	SHGNNQDSSSDNEADEASDDEDNDGN
190	HGNNQDSSSDNEADEASDDEDNDGNE
191	DNNEASADDNNENPEDNNKNTDDNEENPN
192	NNESADDNNENPEDNNKNTDDNEENPNN
193	NESADDNNENPEDNNKNTDDNEENPNNN
194	ESADDNNENPEDNNKNTDDNEENPNNNE
195	SADDNNENPEDNNKNTDDNEENPNNNEN
196	SHGNNQDSSSDNEADEASDDEDNDGNE
197	DNNEASADDNNENPEDNNKNTDDNEENPN N
198	NNESADDNNENPEDNNKNTDDNEENPNN N

199	NESADDNNENPEDNNKNTDDNEENPNNN E
200	ESADDNNENPEDNNKNTDDNEENPNNNE N
201	DNNESADDNNENPEDNNKNTDDNEENPN NN
202	NNESADDNNENPEDNNKNTDDNEENPNN NE
203	NESADDNNENPEDNNKNTDDNEENPNNN EN
204	DNNESADDNNENPEDNNKNTDDNEENPN NNE
205	NNESADDNNENPEDNNKNTDDNEENPNN NEN
206	FWGSHGNNQDSSSDNEADEASDDEDN DGNE (Tag465-9)
207	SHGNNQDSSSDNEADEASDDEDNDGNE GDN
208	DNNESADDNNENPEDNNKNTDDNEENPN NNEN (18-1)
209	SHGNNQDSSSDNEADEASDDEDNDGNE GDNE (Tag 465-3)
210	LEDNNEEPRDPQSPDPNE (Tag656-1)
211	EDNNEEPRDPQSPDPNEF (Tag656-2)
212	DNNEEPRDPQSPDPNEFG (Tag656-3)
213	EGEQQLKPNNSNAPNEDQEE
214	GEQQLKPNNSNAPNEDQEEE
215	KPNNSNAPNEDQEEIQQSE (Tag1121-1)
216	PNNSNAPNEDQEEIQQSEQ
217	NNSNAPNEDQEEIQQSEQH (Tag1121-2)
218	EGEQQLKPNNSNAPNEDQEEE
219	SEGEQQLKPNNSNAPNEDQEEE (Tag1121-3)
220	EQLNFSDDDEQGSNSPKENN
221	LNFSDDDEQGSNSPKENNSE
222	NFSDDDEQGSNSPKENNSED
223	FSDDDEQGSNSPKENNSEDQ
224	LNFSDDDEQGSNSPKENNSED
225	NFSDDDEQGSNSPKENNSEDQ
226	EQLNFSDDDEQGSNSPKENNSE
227	QLNFSDDDEQGSNSPKENNSED
228	LNFSDDDEQGSNSPKENNSEDQ
229	NFSDDDEQGSNSPKENNSEDQG
230	EQLNFSDDDEQGSNSPKENNSED
231	QLNFSDDDEQGSNSPKENNSEDQ
232	LNFSDDDEQGSNSPKENNSEDQG
233	NFSDDDEQGSNSPKENNSEDQGS
234	EQLNFSDDDEQGSNSPKENNSEDQ (Tag2408-1)
235	EQLNFSDDDEQGSNSPKENNSEDQG (Tag2408-2)
236	EQLNFSDDDEQGSNSPKENNSEDQGS (Tag2408-3)
237	EEKNENDENSLSSSSDSSD (Tag6301-1)
238	NENDENSLSSSSDSSDKDE (Tag6301-1)
239	KETNNSNAQNPSEEEEGQDE (Tag6626- 1)

240	ETNNSNAQNPSEEEEGQDE (Tag6626- 2)
241	KETNNSNAQNPSEEEEGQDE (Tag6626-3)
242	ENANDSSDSDGEETDESFP (Tag6915-1)
243	DDNESNSESANGWDSGSNFSE
244	DNESNSESANGWDSGSNFSEE
245	SDDNESNSESANGWDSGSNFSE
246	DDNESNSESANGWDSGSNFSEE (Tag7128-1)
247	SDDNESNSESANGWDSGSNFSEE (Tag7128-2)
248	SSDDNESNSESANGWDSGSNFSEE (Tag7128-3)
249	SEENASSGDSEENTNSDHES (Tag7135-2)
250	EENASSGDSEENTNSDHESE (Tag 7315-1)
251	ENASSGDSEENTNSDHESEQ (Tag7135-3)
252	DENSENNWRNEYPEEESSDG
253	ENSENNWRNEYPEEESSGD
254	NSENNWRNEYPEEESSGDGE
255	DDENSENNWRNEYPEEESSDG (Tag8482- 1)
256	DENSENNWRNEYPEEESSGD
257	ENSENNWRNEYPEEESSGDGE (Tag8482- 2)
258	NSENNWRNEYPEEESSGDGED
259	NSENNWRNEYPEEESSGDGEDS (Tag8482-3)
260	ENSENNWRNEYPEEESSGDGEDS
261	EQQNEASEENNDQQSQEVPE (Tag8974-1)
262	QQNEASEENNDQQSQEVPEK (Tag8974-2)
263	SKMQEDEFDQGNQEEDNSN (Tag9333- 2)
264	KMQEDEFDQGNQEEDNSNA
265	MQEDEFDQGNQEEDNSNAE (Tag 9333- 1)
266	QEDEFDQGNQEEDNSNAEM
267	FDQGNQEEDNSNAEMEEEN
268	EFDQGNQEEDNSNAEMEEEN
269	FDQGNQEEDNSNAEMEEENA
270	DQGNQEEDNSNAEMEEENAS
271	QGNQEEDNSNAEMEEENASN
272	EFDQGNQEEDNSNAEMEEENA
273	FDQGNQEEDNSNAEMEEENAS
274	DQGNQEEDNSNAEMEEENASN
275	DEFDQGNQEEDNSNAEMEEENA
276	EFDQGNQEEDNSNAEMEEENAS
277	FDQGNQEEDNSNAEMEEENASN
278	DEFDQGNQEEDNSNAEMEEENAS
279	EFDQGNQEEDNSNAEMEEENASN
280	QEDEFDQGNQEEDNSNAEMEEENA
281	EDEFDQGNQEEDNSNAEMEEENAS
282	DEFDQGNQEEDNSNAEMEEENASN
283	QEDEFDQGNQEEDNSNAEMEEENAS
284	EDEFDQGNQEEDNSNAEMEEENASN
285	QEDEFDQGNQEEDNSNAEMEEENASN (Tag9333-3)

286	PSEENENSQSEDSVGGDNSEN (Tag10381-1)
287	EVEESNPSEKEDSNPNSSGE (Tag11717-1)
288	VEESNPSEKEDSNPNSSGED (Tag11717-2)
289	KEENSESPLNENSDESYSEE (Tag12237-1)
290	QPGPNHEEDADSYENMDNPD (Tag12809-1)
291	PNHEEDADSYENMDNPDGPD (Tag12809-2)
292	NHEEDADSYENMDNPDGPD (Tag12809-3)
293	DDPNSSDESNGNDDANSESD (Tag 12885-1)
294	DPNSSDESNGNDDANSESDN
295	PNSSDESNGNDDANSESDNN
296	NSSDESNGNDDANSESDNNS (Tag 12885-2)
297	SSDESNGNDDANSESDNNS
298	SDESNGNDDANSESDNNS
299	DESNGNDDANSESDNNS
300	MQGDDPNSSDESNGNDDANSE
301	QGDDPNSSDESNGNDDANSES
302	GDDPNSSDESNGNDDANSESD
303	DDPNSSDESNGNDDANSESDN
304	DPNSSDESNGNDDANSESDNN
305	DKSMQGDDPNSSDESNGNDDAN
306	SMQGDDPNSSDESNGNDDANSE
307	MQGDDPNSSDESNGNDDANSES
308	QGDDPNSSDESNGNDDANSESD
309	GDDPNSSDESNGNDDANSESDN
310	DDPNSSDESNGNDDANSESDNN
311	DPNSSDESNGNDDANSESDNNS
312	DESNGNDDANSESDNNS
313	DDKSMQGDDPNSSDESNGNDDAN
314	DKSMQGDDPNSSDESNGNDDANS
315	KSMQGDDPNSSDESNGNDDANSE
316	SMQGDDPNSSDESNGNDDANSES
317	MQGDDPNSSDESNGNDDANSESD
318	QGDDPNSSDESNGNDDANSESDN
319	GDDPNSSDESNGNDDANSESDNN
320	DDPNSSDESNGNDDANSESDNNS
321	DPNSSDESNGNDDANSESDNNS
322	SDESNGNDDANSESDNNS
323	DESNGNDDANSESDNNS
324	DDANSESDNNS
325	FDDKSMQGDDPNSSDESNGNDDAN
326	DDKSMQGDDPNSSDESNGNDDANS
327	DKSMQGDDPNSSDESNGNDDANSE
328	SMQGDDPNSSDESNGNDDANSESD
329	MQGDDPNSSDESNGNDDANSESDN
330	QGDDPNSSDESNGNDDANSESDNN
331	GDDPNSSDESNGNDDANSESDNNS
332	DDPNSSDESNGNDDANSESDNNS
333	DFDDKSMQGDDPNSSDESNGNDDAN
334	FDDKSMQGDDPNSSDESNGNDDANS
335	DDKSMQGDDPNSSDESNGNDDANSE

336	DKSMQGDDPNSSDESNGNDDANSES
337	KSMQGDDPNSSDESNGNDDANSESD
338	SMQGDDPNSSDESNGNDDANSESDN
339	MQGDDPNSSDESNGNDDANSESDNN
340	QGDDPNSSDESNGNDDANSESDNNS
341	GDDPNSSDESNGNDDANSESDNNS
342	DDPNSSDESNGNDDANSESDNNS
343	DFDDKSMQGDDPNSSDESNGNDDANS
344	FDDKSMQGDDPNSSDESNGNDDANSE
345	DDKSMQGDDPNSSDESNGNDDANSES
346	DKSMQGDDPNSSDESNGNDDANSESD
347	KSMQGDDPNSSDESNGNDDANSESDN
348	SMQGDDPNSSDESNGNDDANSESDNN
349	MQGDDPNSSDESNGNDDANSESDNNS
350	QGDDPNSSDESNGNDDANSESDNNS
351	GDDPNSSDESNGNDDANSESDNNS
352	DDPNSSDESNGNDDANSESDNNS
353	DKSMQGDDPNSSDESNGNDDANSESDN
354	DDKSMQGDDPNSSDESNGNDDANSESD N
355	DKSMQGDDPNSSDESNGNDDANSESDN N
356	DDPNSSDESNGNDDANSESDNNS
357	FDDKSMQGDDPNSSDESNGNDDANSESD N
358	DDKSMQGDDPNSSDESNGNDDANSESD NN
359	DKSMQGDDPNSSDESNGNDDANSESDN NS
360	DDPNSSDESNGNDDANSESDNNS
361	DESNGNDDANSESDNNS
362	DFDDKSMQGDDPNSSDESNGNDDANSES DN
363	FDDKSMQGDDPNSSDESNGNDDANSESD NN
364	DDKSMQGDDPNSSDESNGNDDANSESD NNS
365	DKSMQGDDPNSSDESNGNDDANSESDN NSS
366	DDPNSSDESNGNDDANSESDNNS
367	SDESNGNDDANSESDNNS
368	DESNGNDDANSESDNNS
369	DFDDKSMQGDDPNSSDESNGNDDANSES DNN
370	FDDKSMQGDDPNSSDESNGNDDANSESD NNS
371	DDKSMQGDDPNSSDESNGNDDANSESD NNS
372	DFDDKSMQGDDPNSSDESNGNDDANSES DNN
373	FDDKSMQGDDPNSSDESNGNDDANSESD NNS

374	DDKSMQGDDPNSSDESNGNDDANSESD NNSSS
375	DFDDKSMQGDDPNSSDESNGNDDANSES DNNSS
376	FDDKSMQGDDPNSSDESNGNDDANSESD NNSSS
377	DFDDKSMQGDDPNSSDESNGNDDANSES DNNSSS
378	DDPNSSDESNGNDDANSESDNNSSSRGD ASYNSDE
379	GDDPNSSDESNGNDDANSESDNNSSSRG DASYNSDE
380	DDPNSSDESNGNDDANSESDNNSSSRGD ASYNSDES
381	DSYDFDDKSMQGDDPNSSDESNGNDDA NSESDNNSSSRGD
382	SYDFDDKSMQGDDPNSSDESNGNDDANS ESDNNSSSRGDA
383	YDFDDKSMQGDDPNSSDESNGNDDANSE SDNNSSSRGDAS
384	DFDDKSMQGDDPNSSDESNGNDDANSES DNNSSSRGDASY
385	QGDDPNSSDESNGNDDANSESDNNSSSR GDASYNSDESKD
386	GDDPNSSDESNGNDDANSESDNNSSSRG DASYNSDESKDN
387	DDPNSSDESNGNDDANSESDNNSSSRGD ASYNSDESKDNG
388	DDKSMQGDDPNSSDESNGNDDANSESD NNSSSRGDASYNSDE
389	DFDDKSMQGDDPNSSDESNGNDDANSES DNNSSSRGDASYNSD
390	FDDKSMQGDDPNSSDESNGNDDANSESD NNSSSRGDASYNSDE
391	DDKSMQGDDPNSSDESNGNDDANSESD NNSSSRGDASYNSDES
392	DFDDKSMQGDDPNSSDESNGNDDANSES DNNSSSRGDASYNSDE
393	DFDDKSMQGDDPNSSDESNGNDDANSES DNNSSSRGDASYNSDES
394	DDPNSSDESNGNDDANSESDNNSSSRGD ASYNSDESKDNGNGSDSKGAEDDDSDST SDTN
395	DPNSSDESNGNDDANSESDNNSSSRGDA SYNSDESKDNGNGSDSKGAEDDDSDSTS DTNN
396	NSSDESNGNDDANSESDNNSSSRGDASY NSDESKDNGNGSDSKGAEDDDSDSTS DTNNSD
397	SSDESNGNDDANSESDNNSSSRGDASYN SDESKDNGNGSDSKGAEDDDSDSTS DTNNSDS
398	SDESNGNDDANSESDNNSSSRGDASYNS DESKDNGNGSDSKGAEDDDSDSTS DTNNSDSN
399	AEDDDSDSTS DTNNSDSN

400	EDDDSDSTS DTNNSDSN
401	DDSDSTS DTNNSDSN
402	GDDPNSSDESNGNDDANSESDNNSSSRG DASYNSDESKDNGNGSDSKGAEDDDSDS TSDTN
403	DDPNSSDESNGNDDANSESDNNSSSRGD ASYNSDESKDNGNGSDSKGAEDDDSDST SDTN
404	QGDDPNSSDESNGNDDANSESDNNSSSR GDASYNSDESKDNGNGSDSKGAEDDDSD TSDTN
405	GDDPNSSDESNGNDDANSESDNNSSSRG DASYNSDESKDNGNGSDSKGAEDDDSDS TSDTN
406	DDPNSSDESNGNDDANSESDNNSSSRGD ASYNSDESKDNGNGSDSKGAEDDDSDST SDTN
407	DPNSSDESNGNDDANSESDNNSSSRGDA SYNSDESKDNGNGSDSKGAEDDDSDSTS DTNNSD
408	MQGDDPNSSDESNGNDDANSESDNNS SRGDASYNSDESKDNGNGSDSKGAEDDD SDSTS DTNNSD
409	QGDDPNSSDESNGNDDANSESDNNSSSR GDASYNSDESKDNGNGSDSKGAEDDDSD TSDTN
410	GDDPNSSDESNGNDDANSESDNNSSSRG DASYNSDESKDNGNGSDSKGAEDDDSDS TSDTN
411	DDPNSSDESNGNDDANSESDNNSSSRGD ASYNSDESKDNGNGSDSKGAEDDDSDST SDTN
412	DPNSSDESNGNDDANSESDNNSSSRGDA SYNSDESKDNGNGSDSKGAEDDDSDSTS DTNNSDS
413	GDDPNSSDESNGNDDANSESDNNSSSRG DASYNSDESKDNGNGSDSKGAEDDDSDS TSDTN
414	DDPNSSDESNGNDDANSESDNNSSSRGD ASYNSDESKDNGNGSDSKGAEDDDSDST SDTN
415	QGDDPNSSDESNGNDDANSESDNNSSSR GDASYNSDESKDNGNGSDSKGAEDDDSD TSDTN
416	GDDPNSSDESNGNDDANSESDNNSSSRG DASYNSDESKDNGNGSDSKGAEDDDSDS TSDTN
417	DDPNSSDESNGNDDANSESDNNSSSRGD ASYNSDESKDNGNGSDSKGAEDDDSDST SDTN
418	EDDDSDSTS DTNNSDSN

419	MQGDDPNSSDESNGNDDANSESDNNSS SRGDASYNSDESKDNGNGSDSKGAEDDD SDSTSDTNNSD
420	QGDDPNSSDESNGNDDANSESDNNSSSR GDASYNSDESKDNGNGSDSKGAEDDDSD STSDTNNSDS
421	GDDPNSSDESNGNDDANSESDNNSSSRG DASYNSDESKDNGNGSDSKGAEDDDSDS TSDTNNSDSN
422	DDPNSSDESNGNDDANSESDNNSSSRGD ASYNSDESKDNGNGSDSKGAEDDDSDST SDTNNSDSNG
423	AEDDDSDSTSDTNNSDSNGNGNNGNDDN DKSDSGKGKSDSSSDSSSSNSSDSSD SSSDSSSDSN
424	EDDDSDSTSDTNNSDSNGNGNNGNDDN DKSDSGKGKSDSSSDSSSDSSNSSDSSD SSSDSSSDSN
425	LEEDDNNENAGEDGDNDFPS
426	DNNENAGEDGDNDFPSDEEL (Tag 12968-1)
427	NNENAGEDGDNDFPSDEELA
428	NENAGEDGDNDFPSDEELAN
429	DNNENAGEDGDNDFPSDEELA
430	NNENAGEDGDNDFPSDEELAN
431	AEEEDDNNENAGEDGDNDFPS (Tag 12968-2)
432	DDNNENAGEDGDNDFPSDEELA
433	DNNENAGEDGDNDFPSDEELAN (Tag 12968-3)
434	DDNNENAGEDGDNDFPSDEELAN
435	EDDNNENAGEDGDNDFPSDEELAN
436	NPADDPNQGEDEFEEAEQVREEN (Tag 13648-1)
437	NEENTEPGAESSENADDPNKD (Tag 14056-1)
438	ENADDPNKDTSENADGQSDEN
439	SENADDPNKDTSENADGQSDEN
440	SSENADDPNKDTSENADGQSDEN (Tag 14056-2)
441	ESSENADDPNKDTSENADGQSDEN (Tag 14056-3)
442	DRDPEMENEEQPSSENDSQN (Tag 14681- 1)
443	RDPEMENEEQPSSENDSQNQ
444	DPEMENEEQPSSENDSQNQS
445	PEMENEEQPSSENDSQNQSG (Tag 14681- 2)
446	EMENEEQPSSENDSQNQSGE
447	MENEEQPSSENDSQNQSGEQ
448	ENEEQPSSENDSQNQSGEQI (Tag 14681- 3)
449	DSESANVSDKEAGSNENDQDN (Tag 14844-1)
450	NYNDGSQEDRDWQDDQSDNQ (Tag 15481-1)
451	RENTNEASSEGNSDDSEDE (Tag 16043- 1)

452	ENTNEASSEGNSDDSEDER (Tag 16043- 2)
453	QENDNGNETESEQPKESNEN
454	ENDNGNETESEQPKESNENQ
455	NDNGNETESEQPKESNENQE
456	QENDNGNETESEQPKESNENQ (Tag 16400-1)
457	ENDNGNETESEQPKESNENQE (Tag 16400-2)
458	QENDNGNETESEQPKESNENQE (Tag 16400-3)
459	QNEENPGDEEAKNQVNSESD
460	NEENPGDEEAKNQVNSESDS
461	EENPGDEEAKNQVNSESDSD
462	ENPGDEEAKNQVNSESDSDS
463	NPGDEEAKNQVNSESDSDSE
464	QNEENPGDEEAKNQVNSESDS
465	NEENPGDEEAKNQVNSESDSD
466	QNEENPGDEEAKNQVNSESDSD
467	NEENPGDEEAKNQVNSESDSDS
468	QNEENPGDEEAKNQVNSESDSDS
469	NEENPGDEEAKNQVNSESDSDSE
470	QNEENPGDEEAKNQVNSESDSDSE
471	NEENPGDEEAKNQVNSESDSDSEE
472	QNEENPGDEEAKNQVNSESDSDSEE (Tag 16417-1)
473	NEENPGDEEAKNQVNSESDSDSEES (Tag 16417-2)
474	QNEENPGDEEAKNQVNSESDSDSEES (Tag 16417-3)
475	YNGGNANPRPANNEEEDEED
476	NGGNANPRPANNEEEDEEDE
477	GGNANPRPANNEEEDEEDEY
478	YNGGNANPRPANNEEEDEEDE (Tag 18137-1)
479	NGGNANPRPANNEEEDEEDEY (Tag 18137-2)
480	NGGNANPRPANNEEEDEEDEYD (Tag 18137-3)
481	GASENEEDDDYNKPLDPNS (Tag 18347- 1)
482	LQNQKEAEEPGPDSENSQEN (Tag 18478- 1)
483	QNQKEAEEPGPDSENSQENP (Tag 18478- 2)
484	NQKEAEEPGPDSENSQENPP (Tag 18478- 3)
485	KESVSPENNEEGGNDNQDNEN (Tag 20166-1)
486	ESVSPENNEEGGNDNQDNENP (Tag 20166-2)
487	KESVSPENNEEGGNDNQDNENP (Tag 20166-3)
488	ASPQPREPSDDENS DNSNEC (Tag 41693- 1)
489	NNSQDEDFGQELNENGNAKDE (Tag 55443-1)

490	NSQDEDDGFQELNENGNAKDEN (Tag 55443-2)
491	NNSQDEDDGFQELNENGNAKDEN (Tag 55443-3)

[0215]

Extraction Condition 2:

length: 20 to 70 amino acids

group [D, E]: content of [30] or more

group [D, E]: content of less than [45]

group [P]: content of [10] or more

group [H, K, R]: content of [5] or less

group [G]: content of less than [10]

group [A]: content of less than [10]

group [C, T, V, L, I, M, W]: content of [0]

group [F, Y]: content of [0]

* In the above-described extraction condition, the unit of each content is %. The amino acids are described by one letter codes.

[0216]

[Table 13]

Table 13: Aggregation Rates of scFvs having Tags Extracted under Extraction Condition 2 Added

Tag Name	Sequence	Aggregation Rate %	DE Rate	P Rate	H K R Rate	G Rate	A Rate
tag47-1	SSDSPKQSPPEDSGESEAD	9.48	35	15	5	5	5
tag1784-1	GPEPEPEPEPEPAPEPEPE	15.91	42.86	47.62	0	4.76	4.76
tag1784-2	PEPEPEPEPEPAPEPEPEP	14.03	42.86	52.38	0	0	4.76
tag1784-3	EPEPEPEPEPAPEPEPEPK	23.67	42.86	47.62	4.76	0	4.76
tag2257-1	PEEDGSPDEPSPPEPKPS	22.33	35	40	5	5	0
tag2257-2	EDPEEDGSPDEPSPPEPKP	14.76	42.86	38.1	4.76	4.76	0
tag2257-3	DPEEDGSPDEPSPPEPKPS	17.21	38.1	38.1	4.76	4.76	0
tag4398-1	PDDGSDSDSPSPSPAIESEP	35.71	33.33	23.81	0	4.76	9.52
tag4398-2	DDDGSDSDSPSPSPAIESEPQ	18.64	33.33	19.05	0	4.76	9.52
tag4398-3	PDDGSDSDSPSPSPAIESEPQ	18.58	31.82	22.73	0	4.55	9.09
tag4898-1	PPPEQGQGDAPPQHEDEEPA	9.9	33.33	28.57	4.76	9.52	9.52
tag5533-1	GAPSPAPSPDSDSDSDGEE	19.43	33.33	19.05	0	9.52	9.52
tag5533-2	APSPAPSPDSDSDSDGEE	16.43	38.1	19.05	0	4.76	9.52
tag5533-3	PSAPSPDSDSDSDGEEEE	8.76	42.86	19.05	0	4.76	4.76
tag5601-1	PAPPPPPPEEDPEQDSGPE	33.03	30	50	0	5	5
tag5601-2	APPPPPPEEDPEQDSGPED	20.10	35	45	0	5	5
tag5601-3	PAPPPPPPEEDPEQDSGPED	24.34	33.33	47.62	0	4.76	4.76
tag6354-1	PPPPQAPPEEENESEPEEPS	7.87	33.33	42.86	0	0	4.76
tag6354-2	PPPPQAPPEEENESEPEEPSG	10.68	33.33	38.1	0	4.76	4.76
tag6354-3	PPPPQAPPEEENESEPEEPSG	7.98	31.82	40.91	0	4.55	4.55
tag6681-1	PQDSSSKSPSPAESPDND	8.07	30	20	5	0	5
tag7124-1	QSSDNSEDEEEPPDNADSKS	5.56	40	10	5	0	5
tag7702-1	EEEEQPGKAPDPQDPQDAESD	13.15	42.86	19.05	4.76	4.76	9.52
tag7702-2	EEEEQPGKAPDPQDPQDAESDS	17.29	38.1	19.05	4.76	4.76	9.52
tag7702-3	EEEEQPGKAPDPQDPQDAESDS	17.57	40.91	18.18	4.55	4.55	9.09
tag8341-1	PPPEEEGPEEPKASPESE	23.65	35	35	5	5	5
tag8341-2	PPPEEEGPEEPKASPESE	23.58	33.33	38.1	4.76	4.76	4.76
tag8341-3	PPPEEEGPEEPKASPESEA	24.91	33.33	33.33	4.76	4.76	9.52
tag10102-1	DDAEEPPSPPPPSPSPEP	19.00	30	45	5	0	5
tag11508-1	SGEASSSEEEPPSPDDKENQA	14.91	33.33	14.29	4.76	4.76	9.52
tag11508-2	GEASSSEEEPPSPDDKENQAP	14.49	33.33	19.05	4.76	4.76	9.52
tag11508-3	SGEASSSEEEPPSPDDKENQAP	13.03	31.82	18.18	4.55	4.55	9.09
tag13088-1	PPPPPPPESSDSEPEAEPG	14.98	30	45	0	5	5
tag13088-2	PPPPPESSDSEPEAEPGSP	16.42	30	40	0	5	5
tag13088-3	PPPESSDSEPEAEPGSPQK	20.58	30	30	5	5	5
tag13619-1	SDPEPPDAGEDSKSENGENAP	11.52	33.33	19.05	4.76	9.52	9.52
tag14205-1	SDSEEDPPRNQASDSENEE	10.73	40	10	5	0	5
tag14666-1	PGGEDAEPEEDPQSEDSEAPS	10.21	42.86	19.05	0	9.52	9.52
tag14666-2	PGEDAEPEEDPQSEDSEAPSS	10.04	42.86	19.05	0	4.76	9.52
tag14666-3	PGGEDAEPEEDPQSEDSEAPSS	12.08	40.91	18.18	0	9.09	9.09
tag15430-1	ESESSSSDSEANEPSSASPEPE	14.06	30.43	13.04	0	0	8.7
tag15430-2	SDSESSSSDSEANEPSSASPEPE	10.87	30.77	11.54	0	0	7.69
tag15430-3	DSESSSSDSEANEPSSASPEPEP	7.54	30.77	15.38	0	0	7.69
tag16604-1	HQEDSEEESEEEAEGASEPPPP	12.72	43.48	17.39	4.35	4.35	8.7
tag16604-2	GDHQEDSEEESEEEAEGASEPPPP	12.95	44	16	4	8	8
tag17053-1	PSQPPEEPPEDEAESSPDQ	15.35	35	35	0	0	5
tag17053-2	PAPSPQPEEPPEDEAESSPD	16.53	33.33	38.1	0	0	9.52
tag17053-3	DPAPSPQPEEPPEDEAESSPD	14.03	36.36	36.36	0	0	9.09
tag17170	PGSQPQASSGPEAEFEEDDE	11.11	42.86	14.29	0	9.52	9.52
tag17514-1	SPDSQEEQKGESSASSPEEP	13.45	30	15	5	5	5
tag17514-2	PADSPDSQEEQKGESSASSPEEP	15.02	30.43	17.39	4.35	4.35	8.7
tag17514-3	ADSPDSQEEQKGESSASSPEEP	16.14	34.78	13.04	4.35	4.35	8.7
tag17603-1	PSPEDESSSSSSSSSEDEE	6.01	35	10	0	0	0
tag17603-2	PRSPSPDESSSSSSSSSEDEE	5.96	30.43	13.04	4.35	0	0
tag17603-3	PRSPSPDESSSSSSSSSEDEE	7.22	33.33	12.5	4.17	0	0
tag18253-1	SSSDSDSDSSEDEAPSKP	3.59	35	10	5	0	5
tag18453-1	PSPGSPRGPQDQDDDDDEE	6.12	42.86	19.05	4.76	9.52	0
tag18467-1	PAGDGEAGPQQAEDHPQNPPDPNQPDPEDD	9.87	32.26	25.81	3.23	9.68	9.68
tag18467-2	AGDGEAGPQQAEDHPQNPPDPNQPDPEDDS	5.73	32.26	22.58	3.23	9.68	9.68
tag18467-3	PAGDGEAGPQQAEDHPQNPPDPNQPDPEDDS	8.73	31.25	25	3.13	9.38	9.38
tag18478-1	QNQKEAEPPGPDSENSQENP	14.94	30	15	5	5	5
tag18478-2	NQKEAEPPGPDSENSQENPP	11.47	30	20	5	5	5
tag19033-1	DQNESQSPQPEEGPSDEDDKA	12.69	38.1	14.29	4.76	4.76	4.76
tag19033-2	QNESQSPQPEEGPSDEDDKAE	12.4	38.1	14.29	4.76	4.76	4.76
tag19033-3	NESQSPQPEEGPSDEDDKAE	12.28	38.1	14.29	4.76	9.52	4.76
tag29487-1	PASSSSNPEEGPEEDREAESE	12.77	38.1	14.29	4.76	4.76	9.52
tag34831-1	DKPEEDDEAQPPQSQSGPEAE	7.58	43.48	17.39	4.35	4.35	8.7
tag34831-2	KPEEDDEAQPPQSQSGPEAE	8.02	43.48	17.39	4.35	4.35	8.7
tag34831-3	PEEDDEAQPPQSQSGPEAE	9.14	43.48	17.39	0	8.7	8.7
tag34858-1	PEEEHAPGEDESSPQSPQS	17.52	30	25	5	5	5

In Table 13, the P content ratio in the amino acid sequences of the extracted tags is 15% or more, and the content ratios of the other amino acids are as described above in Extraction Condition 2. In order to confirm that the aggregation inhibiting action does not depend on a specific amino acid sequence, tags were randomly selected from the tags extracted under Extraction Condition 2. As for some tags, an amino acid sequence satisfying the extraction condition was additionally selected from another portion of the same protein. The aggregation rates of scFvs tagged with the selected amino acid sequences were tested, and results as shown in Table 13 were obtained. As shown in Table 13, the aggregation rates of the tagged scFvs were low as a whole.

[0218]

It is noted that human-derived amino acid sequences that can be extracted under Extraction Condition 2 were as follows.

[Table 13-2]

Table 13-2: Examples of human-derived amino acid sequences that can be extracted under Extraction Condition 2

SEQ ID NO:	Sequence
492	SSDSPKQSPPEDSGESEAD (tag47-1)
493	PGPEPEPEPEPEPEPEPEPE
494	GPEPEPEPEPEPEPEPEPEPE
495	PEPEPEPEPEPEPEPEPEPK
496	EPEPEPEPEPEPEPEPEPKP
497	PEPEPEPEPEPEPEPEPKPG
498	AGPGPEPEPEPEPEPEPEPEPE
499	GPGPEPEPEPEPEPEPEPEPE
500	PGPEPEPEPEPEPEPEPEPEPE
501	GPEPEPEPEPEPEPEPEPEPE (tag1784-1)
502	PEPEPEPEPEPEPEPEPEPE (tag1784-2)
503	EPEPEPEPEPEPEPEPEPEPK (tag1784-3)
504	PEPEPEPEPEPEPEPEPEPKP
505	EPEPEPEPEPEPEPEPEPKPG
506	PEPEPEPEPEPEPEPEPKPGA
507	EPEPEPEPEPEPEPEPKPGAG
508	AGPGPEPEPEPEPEPEPEPEPE
509	GPGPEPEPEPEPEPEPEPEPEPE
510	PGPEPEPEPEPEPEPEPEPEPE
511	GPEPEPEPEPEPEPEPEPEPEPE
512	PEPEPEPEPEPEPEPEPEPEPK
513	EPEPEPEPEPEPEPEPEPEPKP
514	PEPEPEPEPEPEPEPEPEPKPG
515	EPEPEPEPEPEPEPEPEPKPGA
516	PEPEPEPEPEPEPEPEPKPGAG
517	AGPGPEPEPEPEPEPEPEPEPEPE
518	GPGPEPEPEPEPEPEPEPEPEPE
519	PGPEPEPEPEPEPEPEPEPEPEPE
520	GPEPEPEPEPEPEPEPEPEPEPK
521	PEPEPEPEPEPEPEPEPEPEPKP
522	EPEPEPEPEPEPEPEPEPEPKPG
523	PEPEPEPEPEPEPEPEPEPKPGA
524	EPEPEPEPEPEPEPEPEPKPGAG
525	AGPGPEPEPEPEPEPEPEPEPEPE
526	GPGPEPEPEPEPEPEPEPEPEPEPE
527	PGPEPEPEPEPEPEPEPEPEPEPK
528	GPEPEPEPEPEPEPEPEPEPEPKP
529	PEPEPEPEPEPEPEPEPEPEPKPG
530	EPEPEPEPEPEPEPEPEPEPKPGA
531	PEPEPEPEPEPEPEPEPEPKPGAG
532	AGPGPEPEPEPEPEPEPEPEPEPEPE
533	GPGPEPEPEPEPEPEPEPEPEPEPK
534	PGPEPEPEPEPEPEPEPEPEPEPKP
535	GPEPEPEPEPEPEPEPEPEPEPKPG
536	PEPEPEPEPEPEPEPEPEPEPKPGA
537	EPEPEPEPEPEPEPEPEPEPKPGAG
538	AGPGPEPEPEPEPEPEPEPEPEPEPK
539	GPGPEPEPEPEPEPEPEPEPEPEPKP
540	PGPEPEPEPEPEPEPEPEPEPEPKPG
541	GPEPEPEPEPEPEPEPEPEPEPKPGA
542	PEPEPEPEPEPEPEPEPEPEPKPGAG
543	AGPGPEPEPEPEPEPEPEPEPEPKP
544	PGPEPEPEPEPEPEPEPEPEPKPGA
545	DPEEDGSPDPEPSPEPEPKP
546	PEEDGSPDPEPSPEPEPKPS (tag2257-1)
547	EDPEEDGSPDPEPSPEPEPKP (tag2257-2)
548	DPEEDGSPDPEPSPEPEPKPS (tag2257-3)
549	EDPEEDGSPDPEPSPEPEPKPS
550	DEDPEEDGSPDPEPSPEPEPKPS
551	PDDGSDSSPPSASPAESEP (tag4398-1)
552	DDGSDSSPPSASPAESEPQ (tag4398-2)
553	PDDGSDSSPPSASPAESEPQ (tag4398-3)
554	PPPEEQGGDAPPQHEDEEPA (tag4898-1)
555	PSPAPSPDSDSDSDSDGEEE
556	GAPSPAPSPDSDSDSDSDGEE (tag5533-1)
557	APSPAPSPDSDSDSDSDGEEE (tag5533-2)
558	PSPAPSPDSDSDSDSDGEEEE (tag5533-3)
559	NGAPSPAPSPDSDSDSDSDGEE
560	GAPSPAPSPDSDSDSDSDGEEE
561	APSPAPSPDSDSDSDSDGEEEE
562	QNGAPSPAPSPDSDSDSDSDGEE
563	NGAPSPAPSPDSDSDSDSDGEEE
564	GAPSPAPSPDSDSDSDSDGEEEE
565	APSPAPSPDSDSDSDSDGEEEE
566	QNGAPSPAPSPDSDSDSDSDGEEE
567	NGAPSPAPSPDSDSDSDSDGEEEE
568	GAPSPAPSPDSDSDSDSDGEEEE
569	QNGAPSPAPSPDSDSDSDSDGEEEE
570	NGAPSPAPSPDSDSDSDSDGEEEE
571	GAPSPAPSPDSDSDSDSDGEEEE
572	QNGAPSPAPSPDSDSDSDSDGEEEE
573	NGAPSPAPSPDSDSDSDSDGEEEE
574	QNGAPSPAPSPDSDSDSDSDGEEEE
575	NGAPSPAPSPDSDSDSDSDGEEEE
576	QNGAPSPAPSPDSDSDSDSDGEEEE
577	QNGAPSPAPSPDSDSDSDSDGEEEE GER
578	QNGAPSPAPSPDSDSDSDSDGEEEE GERD
579	EPPAPPPPPPEEDPEQDSG
580	PAPPPPPPPPEEDPEQDSGPE (tag5601-1)
581	APPPPPPPPEEDPEQDSGPE (tag5601-2)
582	PAPPPPPPPPEEDPEQDSGPED (tag5601-3)
583	EPPAPPPPPPPPEEDPEQDSGPE
584	PPAPPPPPPPPEEDPEQDSGPED

585	AEPPAPPPPPPEEDPEQDSGPE
586	EPPAPPPPPPEEDPEQDSGPED
587	AEPPAPPPPPPEEDPEQDSGPED
588	PPPPQAPPEEENESEPEEP
589	PPPPQAPPEEENESEPEEPS
590	PPPQAPPEEENESEPEEPSG
591	PPPPQAPPEEENESEPEEPS (tag6354-1)
592	PPPPQAPPEEENESEPEEPSG (tag6354-2)
593	PPPPQAPPEEENESEPEEPSG (tag6354-3)
594	PQDSSSKSPEPSADESPDND (tag6681-1)
595	QSSDNSEDEEPPDNADSKS (tag7124-1)
596	EEEEQPGKAPDPQDPQDAES
597	EEEEQPGKAPDPQDPQDAESD (tag7702-1)
598	EEEQPGKAPDPQDPQDAESDS (tag7702-2)
599	EEEEQPGKAPDPQDPQDAESDS (tag7702-3)
600	EEEEQPGKAPDPQDPQDAESDS
601	PPPPSEEEGPEEPPKASPE
602	PPPPSEEEGPEEPPKASPES
603	PPPSSEEEGPEEPPKASPESE (tag8341-1)
604	PPPPSEEEGPEEPPKASPESE (tag8341-2)
605	PPPSSEEEGPEEPPKASPESEA (tag8341-3)
606	PPPPSEEEGPEEPPKASPESE
607	PPPPSEEEGPEEPPKASPESEA
608	PPPPSEEEGPEEPPKASPESEA
609	DDAEEPESPPPPRSPSEPE (tag10102-1)
610	SGEASSSEEEPPSPDDKENQ
611	SGEASSSEEEPPSPDDKENQA (tag11508-1)
612	GEASSSEEEPPSPDDKENQAP (tag11508-2)
613	SGEASSSEEEPPSPDDKENQAP (tag11508-3)
614	PQPPPPPPPEESSDSEPEAE
615	QPPPPPPPEESSDSEPEAEP
616	PPPPPPPEESSDSEPEAEPG (tag13088-1)
617	PPPPPEESSDSEPEAEPGS
618	PPPPPEESSDSEPEAEPGSP (tag13088-2)
619	PPPPEESSDSEPEAEPGSPQ
620	PPPEESSDSEPEAEPGSPQK (tag13088-3)
621	SDPEPPDAGEDSKSENGENAP (tag13619-1)
622	SDSESDPPRNQASDSENEE (tag14205-1)
623	GPGEDAEPDEDQSEDSEAPS (tag14666-1)
624	PGEDAEPDEDQSEDSEAPSS (tag14666-2)
625	GPGEDAEPDEDQSEDSEAPSS (tag14666-3)
626	ESESSSSDSEANEPSQSASPEPE (tag15430-1)
627	DSESESSSSDSEANEPSQSASPEPE
628	SDSESESSSSDSEANEPSQSASPEPE (tag15430-2)

629	DSESESSSSDSEANEPSQSASPEPEP (tag15430-3)
630	HQEDSEEEESQEEEAEGASEPPPP (tag16604-1)
631	GDHQEDSEEEESQEEEAEGASEPPPP (tag16604-2)
632	PSQPPEEPEPDEAESSDPQ (tag17053-1)
633	DPAPSQPPEEPEPDEAESSPD
634	PAPSQPPEEPEPDEAESSDP (tag17053-2)
635	APSQPPEEPEPDEAESSDPQ
636	PSQPPEEPEPDEAESSDPQA
637	DPAPSQPPEEPEPDEAESSDP (tag17053-3)
638	PAPSQPPEEPEPDEAESSDPQ
639	DPAPSQPPEEPEPDEAESSDPQ
640	PGSQPQASSGPEAEEDDE (tag17170)
641	DSPDSQEEQKGEASSASPEE
642	SPDSQEEQKGEASSASPEEP (tag17514-1)
643	PDSQEEQKGEASSASPEEPE
644	DSQEEQKGEASSASPEEPEE
645	DSPDSQEEQKGEASSASPEEP
646	SPDSQEEQKGEASSASPEEPE
647	PDSQEEQKGEASSASPEEPEE
648	PADSPDSQEEQKGEASSASPEE
649	ADSPDSQEEQKGEASSASPEEP
650	DSPDSQEEQKGEASSASPEEPE
651	SPDSQEEQKGEASSASPEEPEE
652	PADSPDSQEEQKGEASSASPEEP (tag17514-2)
653	ADSPDSQEEQKGEASSASPEEPE (tag17514-3)
654	DSPDSQEEQKGEASSASPEEPEE
655	PADSPDSQEEQKGEASSASPEEPE
656	ADSPDSQEEQKGEASSASPEEPEE
657	PADSPDSQEEQKGEASSASPEEPEE
658	SPSPDESSSSSSSSSEDE
659	PSPDESSSSSSSSSEDEE (tag17603-1)
660	PRSPSPDESSSSSSSSSEDEE (tag17603-2)
661	PRSPSPDESSSSSSSSSEDEEE (tag17603-3)
662	SSSDSSDSDSEDDEAPSKP (tag18253-1)
663	PSPGSPRGQPQDQDDDEDEE (tag18453-1)
664	QQAEDHPQNPPEDPNQDPPE
665	QAEDHPQNPPEDPNQDPPE
666	AEDHPQNPPEDPNQDPPEDD
667	EDHPQNPPEDPNQDPPEDDS
668	QQAEDHPQNPPEDPNQDPPE
669	QAEDHPQNPPEDPNQDPPEDD
670	AEDHPQNPPEDPNQDPPEDDS
671	PQQAEDHPQNPPEDPNQDPPE
672	QQAEDHPQNPPEDPNQDPPEDD
673	QAEDHPQNPPEDPNQDPPEDDS
674	DGEAGPQQAEDHPQNPPEDPNQD

675	GPQQAEDHPQNPPEDPNQDPPED
676	PQQAEDHPQNPPEDPNQDPPEDD
677	QQAEDHPQNPPEDPNQDPPEDDS
678	GPQQAEDHPQNPPEDPNQDPPEDD
679	PQQAEDHPQNPPEDPNQDPPEDDS
680	EAGPQQAEDHPQNPPEDPNQDPPED
681	AGPQQAEDHPQNPPEDPNQDPPEDD
682	GPQQAEDHPQNPPEDPNQDPPEDDS
683	DGEAGPQQAEDHPQNPPEDPNQDPPE
684	GEAGPQQAEDHPQNPPEDPNQDPPED
685	EAGPQQAEDHPQNPPEDPNQDPPEDD
686	AGPQQAEDHPQNPPEDPNQDPPEDDS
687	DGEAGPQQAEDHPQNPPEDPNQDPPED
688	GEAGPQQAEDHPQNPPEDPNQDPPEDD
689	EAGPQQAEDHPQNPPEDPNQDPPEDDS
690	DGEAGPQQAEDHPQNPPEDPNQDPPED D
691	GEAGPQQAEDHPQNPPEDPNQDPPEDD S
692	DGEAGPQQAEDHPQNPPEDPNQDPPED DS
693	PAGDGEAGPQQAEDHPQNPPEDPNQDP PEDD (tag18467-1)
694	AGDGEAGPQQAEDHPQNPPEDPNQDPP EDDS (tag18467-2)
695	PAGDGEAGPQQAEDHPQNPPEDPNQDP PEDDS (tag18467-3)
696	QNQKEAEEPGPDSSENSQENP (tag18478- 1)
697	NQKEAEEPGPDSSENSQENPP (tag18478-2)
698	DQNESQSPQEPEEGPSEDDK
699	QNESQSPQEPEEGPSEDDKA
700	NESQSPQEPEEGPSEDDKAE
701	DQNESQSPQEPEEGPSEDDKA (tag19033- 1)
702	QNESQSPQEPEEGPSEDDKAE (tag19033- 2)
703	NESQSPQEPEEGPSEDDKAE (tag19033- 3)
704	ESQSPQEPEEGPSEDDKAEGE
705	SQSPQEPEEGPSEDDKAEGEE
706	DQNESQSPQEPEEGPSEDDKAE
707	QNESQSPQEPEEGPSEDDKAEG
708	NESQSPQEPEEGPSEDDKAEGE
709	DQNESQSPQEPEEGPSEDDKAEG
710	QNESQSPQEPEEGPSEDDKAEGE
711	NESQSPQEPEEGPSEDDKAEGEE
712	DQNESQSPQEPEEGPSEDDKAEGE

713	QNESQSPQEPEEGPSEDDKAEGEE
714	DQNESQSPQEPEEGPSEDDKAEGEE
715	QNESQSPQEPEEGPSEDDKAEGEEE
716	XEASSSEEEPPSPDDKENQAP (tag25919- 1)
717	PASSSSNP EEGPSEEDREAEE (tag29487- 1)
718	DKPEEEDDEAQQPQPQSGPE
719	KPEEEDDEAQQPQPQSGPEE
720	AGEGDKPEEEDDEAQQPQPQSGP
721	EGDKPEEEDDEAQQPQPQSGP
722	GDKPEEEDDEAQQPQPQSGPE
723	DKPEEEDDEAQQPQPQSGPEE
724	KPEEEDDEAQQPQPQSGPEEA
725	PEEEDDEAQQPQPQSGPEEAE
726	EEDDEAQQPQPQSGPEEAEEG
727	EDDEAQQPQPQSGPEEAEEGE
728	DDEAQQPQPQSGPEEAEEGEE
729	DEAQQPQPQSGPEEAEEGEEE
730	EAQQPQPQSGPEEAEEGEEEE
731	QQPQPQSGPEEAEEGEEEEAE
732	QPQPQSGPEEAEEGEEEEAER
733	EGDKPEEEDDEAQQPQPQSGPE
734	GDKPEEEDDEAQQPQPQSGPEE
735	DKPEEEDDEAQQPQPQSGPEEA
736	KPEEEDDEAQQPQPQSGPEEAE
737	QQPQPQSGPEEAEEGEEEEAER
738	EGDKPEEEDDEAQQPQPQSGPEE
739	GDKPEEEDDEAQQPQPQSGPEEA
740	DKPEEEDDEAQQPQPQSGPEEAE (tag34831-1)
741	KPEEEDDEAQQPQPQSGPEEAE (tag34831-2)
742	PEEEDDEAQQPQPQSGPEEAE (tag34831-3)
743	EGDKPEEEDDEAQQPQPQSGPEEA
744	GDKPEEEDDEAQQPQPQSGPEEAE
745	KPEEEDDEAQQPQPQSGPEEAE
746	EGDKPEEEDDEAQQPQPQSGPEEAE
747	GDKPEEEDDEAQQPQPQSGPEEAE
748	DKPEEEDDEAQQPQPQSGPEEAE
749	KPEEEDDEAQQPQPQSGPEEAE
750	PEEEHAPGEDESSPQSPQPS (tag34858-1)
751	XPPPEESSDSEPEAEPGSPQ (tag)
752	PPPEESSDSEPEAEPGSPQK (tag)

[0219]

Extraction Condition 3:

length: 20 to 70 amino acids

group [D, E]: content of [45] or more

group [G]: content of less than [10]

group [A]: content of less than [10]

group [F, Y]: content of [0]

group [C, M, L, I, W, T, V]: content of [0]

group [P]: content of [15] or more

[0220]

[Table 14]

Table 14: Aggregation Rates of scFvs having Tags Extracted under Extraction Condition 3 Added

Tag Name	Sequence	Aggregation Rate %	DE Rate	P Rate	G Rate	A Rate
Tag 167-1	KEPKKEKKDDDEEAPKPSSD	8.02	45	15	0	5
Tag 1034-1	SEEEKPPEEDKEEEEEKKAP	8.91	55	15	0	5
Tag 1409-1	PAEEEDDDPEQEKEAGEPGRP	5.17	47.62	19.05	9.52	9.52
Tag 1784-1	EPEPEPEPEPEPAPEPEPEP	12.97	45	50	0	5
Tag 2257-1	EDPEEDGSPDPEPSPEPEPK	12.61	45	35	5	0
Tag 2257-2	DEDPEEDGSPDPEPSPEPEPK	11.50	47.62	33.33	4.76	0
Tag 2257-3	DEDPEEDGSPDPEPSPEPEPKP	8.62	45.45	36.36	4.55	0
Tag 2740-1	KPEDKDPDPPEESKEPKKEEK	13.22	45	20	0	0
Tag 2740-2	PEDKDPDPPEESKEPKKEEKQ	11.68	45	20	0	0
Tag 2740-3	EDKDPDPPEESKEPKKEEKQR	8.79	45	15	0	0
Tag 3227-1	KRNDSEEEERERDEEQEPPP	20.63	50	15	0	0
Tag 4898-1	PEEPDDQDAPDEHEPSPSED	10.71	52.38	23.81	0	4.76
Tag 4898-2	EEEPDDQDAPDEHEPSPSEDA	7.01	52.38	19.05	0	9.52
Tag 4898-3	EEPDQDAPDEHEPSPSEDA	19.90	47.62	23.81	0	9.52
Tag 5533-1	PSPAPSPDSDSDSDSGEEEEEEE	1.89	50	16.67	4.17	4.17
Tag 5533-2	APSPAPSPDSDSDSDSGEEEEEEE	1.00	48	16	4	8
Tag 5533-3	PSPAPSPDSDSDSDSGEEEEEEEG	0.90	48	16	8	4
Tag 6236-1	EKNDDEPQKPEDKGDPEGPE	10.19	47.62	19.05	9.52	0
Tag 6236-2	EKNDDEPQKPEDKGDPEGPEA	8.40	45.45	18.18	9.09	4.55
Tag 6755-1	EDEEEEEEEEEDEGPAPP	4.10	75	15	5	5
Tag 6755-2	DEEEEEEEEEDEGPAPPS	2.38	70	15	5	5
Tag 7167-1	GEREPDPPDDRDASDGEDEKP	12.37	47.62	19.05	9.52	4.76
Tag 7167-2	EREPPDPPDDRDASDGEDEKPP	8.77	47.62	23.81	4.76	4.76
Tag 7167-3	EGEREPPDPPDDRDASDGEDEKP	9.97	50	18.18	9.09	4.55
Tag 7702-1	EEEEEQPGKAPDPQDPQDAESD	13.30	45.45	18.18	4.55	9.09
Tag 8243-1	EPEEKQEPEEKQEPEEKQKPE	11.90	47.62	19.05	0	0
Tag 8243-2	QEPEEKQEPEEKQEPEEKQKPE	12.44	45.45	18.18	0	0
Tag 8243-3	EPEEKQEPEEKQEPEEKQKPEA	12.60	45.45	18.18	0	4.55
Tag 8818-1	DDDDDDSDPDSPDSDSDSD	2.58	65	15	0	0
Tag 8818-2	DDDDDDSDPDSPDSDSDSDSD	2.19	60	15	0	0
Tag 8818-3	DDSDPDSPDSDSDSDSDSEK	8.33	50	15	0	0
Tag 9050-1	DPDQPREDPAAEEKEEKDAPE	15.60	52.38	19.05	0	9.52
Tag 9166-1	PENESEPKHEEPPKPEEKPEEE	17.13	50	22.73	0	0
Tag 9166-2	NESEPKHEEPPKPEEKPEEEK	7.62	50	18.18	0	0
Tag 9166-3	PENESEPKHEEPPKPEEKPEEEE	13.00	52.17	21.74	0	0
Tag 9590-1	PPEEDPEEQAEENPEGEQPE	13.99	50	25	5	5
Tag 9590-2	KPPEEDPEEQAEENPEGEQPE	10.18	47.62	23.81	4.76	4.76
Tag 9590-3	KPPEEDPEEQAEENPEGEQPEE	9.67	50	22.73	4.55	4.55
Tag 9704-1	PDDDDSESDHDDPDNAHESP	7.65	55	15	0	5
Tag 9749-1	PEPEPEPEPEPESEPEPEPE	15.74	50	45	0	0
Tag 9749-2	GGEPEPEPEPEPEPESEPEPE	12.24	47.83	39.13	8.7	0
Tag 9749-3	GGEPEPEPEPEPEPESEPEPEPE	14.29	48	40	8	0
Tag 10346-1	PEEPDDQDAPDEHESPPPE	8.16	50	30	0	5
Tag 10346-2	EEEPDDQDAPDEHESPPPEDA	6.88	52.38	23.81	0	9.52
Tag 10346-3	PEEPDDQDAPDEHESPPPEDAP	8.97	47.83	30.43	0	8.7
Tag 11099-1	GPSSDDENEESKPEKEDEP	7.21	50	15	5	0
Tag 11099-2	PSSDDENEESKPEKEDEPQ	5.09	50	15	0	0

Tag 12127-1	SDDSDSEKRRPEEQEEEEQP	17.51	45	15	0	0
Tag 12127-2	DDSDSEKRRPEEQEEEEQPR	17.44	45	15	0	0
Tag 13036-1	PEEEDEEPGDPREGEEEEDEPD	3.17	68	20	8	0
Tag 13036-2	PEEEDEEPGDPREGEEEEDEPDPEAP	5.60	64.29	21.43	7.14	3.57
Tag 13036-3	AAPEEEDEEPGDPREGEEEEDEPDPEAPENG	6.19	55.88	17.65	8.82	8.82
Tag 14128-1	PPPSEGSDDEEEEEDEE	2.43	70	15	5	0
Tag 14128-2	KPPPSEGSDDEEEEEDEEERKP	10.86	60	16	4	0
Tag 14128-3	KPPPSEGSDDEEEEEDEEERKPQ	9.76	57.69	15.38	3.85	0
Tag 16549-1	EEEEEEEEEEEEEAAPPP	4.95	75	20	0	5
Tag 16549-2	EEEEEEEEEEEEEAAPPPR	3.01	73.91	17.39	0	4.35
Tag 16549-3	PDDDEEEEEEEEEEEEEEAAPPP	0.89	81.82	15.15	0	3.03
Tag 16604-1	EDSEESQEEEAEGASEPPPP	6.00	47.62	19.05	4.76	9.52
Tag 16604-2	QEDSEESQEEEAEGASEPPPP	7.39	45.45	18.18	4.55	9.09
Tag 16604-3	DHQEDSEESQEEEAEGASEPPPP	10.59	45.83	16.67	4.17	8.33
Tag 16741-1	DQSEEEEEKHPKPAKPE	10.62	45	20	0	5
Tag 16991-1	PAPAHRRPEDEGEENEGEED	14.97	47.62	19.05	9.52	9.52
Tag 16991-2	PAPAHRRPEDEGEENEGEED	7.12	50	18.18	9.09	9.09
Tag 17199-1	QENGQREEEEEKEPEAEP	9.09	50	15	5	5
Tag 17199-2	PAEQENGQREEEEEKEPEAEP	10.29	45.83	16.67	8.33	8.33
Tag 17936-1	EQEPEPEPEPEPEPEPE	6.37	50	45	0	0
Tag 17936-2	EPEPEPEPEPEPEPEPEQ	10.94	50	45	0	0
Tag 17936-3	EQEPEPEPEPEPEPEPEQ	16.61	50	40.91	0	0
Tag 18132-1	PEEEEEEEEEEPASPPERK	3.36	60	20	0	5
Tag 18453-1	PSPGSPRGQPQDQDDDEDEED	4.31	45.45	18.18	9.09	0
Tag 18453-2	RPSGSPRGQPQDQDDDEDEEDE	13.65	45.83	16.67	8.33	0
Tag 18453-3	PSPGSPRGQPQDQDDDEDEDEA	6.54	45.83	16.67	8.33	4.17
Tag 18866-1	AEDDDEEEEEEEEPDPP	2.24	80	15	0	5
Tag 18866-2	EDDDEEEEEEEEPDPP	2.29	85	15	0	0
Tag 19350-1	KQEPDPEDKEENKDDAS	14.73	45	15	0	5
Tag 19350-2	QEPDPEDKEENKDDAS	14.12	45	15	0	5
Tag 19511-1	EDEDEDESSEEDSEDEPP	1.70	70	15	0	0
Tag 19511-2	PDDSRDEDEDESSEEDSEDEPP	4.37	65.38	15.38	0	0
Tag 19511-3	PKKEPDDSRDEDEDESSEEDSEDEPPPKRR	3.87	54.55	15.15	0	0
Tag 22900-1	PEEEAAEEEEEEERPKPSRP	10.93	52.38	19.05	0	9.52
Tag 22900-2	EQPEEEAAEEEEEEERPKPSRP	6.90	52.17	17.39	0	8.7
Tag 22900-3	EEQPEEEAAEEEEEEERPKPSRP	9.67	56	16	0	8
Tag 34831-1	PEEEDDEAQQPQSGPEEAE	10.16	45.45	18.18	4.55	9.09
Tag 34831-2	EGDKPEEEDDEAQQPQSGPEEAE	10.78	46.15	15.38	7.69	7.69
Tag 34831-3	KPEEEDDEAQQPQSGPEEAEAEGERP	8.33	45.45	15.15	9.09	9.09
Tag 39056-1	NNSEEEEDDDDEEPPDKPP	0.44	65	15	0	0
Tag 39056-2	NSEEEEDDDDEEPPDKPPA	0.99	65	15	0	5
Tag 39056-3	SEEEEDDDDEEPPDKPPAN	1.23	65	15	0	5

[0221]

In Table 14, the acidic amino acid ratio in the amino acid sequences of the extracted tags is 45% or more, the P content ratio is 15% or more, and the content ratios of the other amino acids are as described above in Extraction Condition 3. In order to confirm that the aggregation inhibiting action does not depend on a specific amino acid sequence, tags were randomly selected from the tags extracted under Extraction Condition 3. As for some tags, an amino acid sequence satisfying the

extraction condition was additionally selected from another portion of the same protein. The aggregation rates of scFvs tagged with the selected amino acid sequences were tested, and results as shown in Table 14 were obtained. As shown in Table 14, the aggregation rates of the tagged scFvs were low as a whole. It is understood, from Table 14-1, that a strategy of reducing the G, A, F, Y, C, M, L, I, W, T, and V contents and increasing the P content works on a peptide tag having a high acidic amino acid content.

[0222]

It is noted that human-derived amino acid sequences that can be extracted under Extraction Condition 3 were as follows.

[Table 14-2]

Table 14-2: Examples of human-derived amino acid sequences that can be extracted under Extraction Condition 3

SEQ ID NO:	Sequence
753	KEPKKEKKDDDEEAPKPSSD (Tag 167-1)
754	SEEEKPPEEDKEEEEEKKAP (Tag 1034-1)
755	PAEEDDDPEQEKEAGEPGRP (Tag 1409-1)
756	PEPEPEPEPEPEPAPEPEPE
757	EPEPEPEPEPEPAPEPEPEP (Tag 1784-1)
758	DEDPEEDGSPDPEPSPEPEP
759	EDPEEDGSPDPEPSPEPEPK (Tag 2257-1)
760	DEDPEEDGSPDPEPSPEPEPK (Tag 2257-2)
761	DEDPEEDGSPDPEPSPEPEPK (Tag 2257-3)
762	KPEDKDPDRDPEESKEPKKEK (Tag 2740-1)
763	PEDKDPDRDPEESKEPKKEKQ (Tag 2740-2)
764	EDKDPDRDPEESKEPKKEKQR (Tag 2740-3)
765	KRNDSEEEERERDEEQEPPP (Tag 3227-1)
766	PEEPPDDQDAPDEHEPSPSE
767	EEPPDDQDAPDEHEPSPSED
768	PEEPPDDQDAPDEHEPSPSED (Tag 4898-1)
769	EEPPDDQDAPDEHEPSPSEDA (Tag 4898-2)
770	EEPPDDQDAPDEHEPSPSEDAP (Tag 4898-3)
771	PEEPPDDQDAPDEHEPSPSEDA
772	EEPPDDQDAPDEHEPSPSEDAP
773	PEEPPDDQDAPDEHEPSPSEDAP
774	SPAPSPDSDSDSDSGEEEE
775	PAPSPDSDSDSDSGEEEE
776	PSPAPSPDSDSDSDSGEEEE
777	PSPAPSPDSDSDSDSGEEEE
778	APSPAPSPDSDSDSDSGEEEE
779	PSPAPSPDSDSDSDSGEEEE (Tag 5533-1)
780	APSPAPSPDSDSDSDSGEEEE (Tag 5533-2)
781	PSPAPSPDSDSDSDSGEEEEEG (Tag 5533-3)
782	GAPSPAPSPDSDSDSDSGEEEE
783	APSPAPSPDSDSDSDSGEEEEEG
784	PSPAPSPDSDSDSDSGEEEEEGE
785	EKNDEDEPQKPEDKGDPEGPE (Tag 6236-1)
786	EKNDEDEPQKPEDKGDPEGPEA (Tag 6236-2)
787	EDEEEEEEEEEDEGPAPP (Tag 6755-1)
788	DEEEEEEEEEDEGPAPPS (Tag 6755-2)
789	EREPPPPDDRDASDGEDEKP
790	REPDPDDRDASDGEDEKPP
791	GEREPDPDDRDASDGEDEKP (Tag 7167-1)
792	EREPPPPDDRDASDGEDEKPP (Tag 7167-2)
793	EGEREPPPPDDRDASDGEDEKP (Tag 7167-3)
794	GEREPDPDDRDASDGEDEKPP
795	EGEREPPPPDDRDASDGEDEKPP
796	EEEEEQPGKAPDPQDPQDAESD (Tag 7702-1)
797	QEPEEKQEPEEKQEPEEKQK
798	EPEEKQEPEEKQEPEEKQKP
799	PEEKQEPEEKQEPEEKQKPE
800	EEKQEPEEKQEPEEKQKPEA
801	EPEEKQEPEEKQEPEEKQKPE (Tag 8243-1)
802	QEPEEKQEPEEKQEPEEKQKPE (Tag 8243-2)
803	EPEEKQEPEEKQEPEEKQKPEA (Tag 8243-3)
804	AGDDDDDDDDSDPESPDDSD
805	GDDDDDDDDSDPESPDDSE
806	DDDDDDDDSDPESPDDSES
807	DDDDDDSDPESPDDSESD (Tag 8818-1)
808	DDDDDDSDPESPDDSESDS (Tag 8818-2)
809	DDDDSDPESPDDSESDSE
810	DDDDSDPESPDDSESDSES
811	DDSDPESPDDSESDSESE
812	DDSDPESPDDSESDSESEK (Tag 8818-3)
813	DSPDPESPDDSESDSESEKE
814	SPDPESPDDSESDSESEKEE
815	PDPEPDDSESDSESEKEES
816	DPDQPREDPAAAAKEEKDAPE (Tag 9050-1)
817	EGKPENESEPKHEEPPKPEE
818	PENESEPKHEEPPKPEEKPE
819	ENESEPKHEEPPKPEEKPEE
820	NESEPKHEEPPKPEEKPEEE
821	ESEPKHEEPPKPEEKPEEEE
822	SEPKHEEPPKPEEKPEEEEE
823	PENESEPKHEEPPKPEEKPEE
824	ENESEPKHEEPPKPEEKPEEE

825	NESEPKHEEHPKPEEKPEEEE
826	ESEPKHEEHPKPEEKPEEEEK
827	KPENESEPKHEEHPKPEEKPEE
828	PENESEPKHEEHPKPEEKPEEE (Tag 9166-1)
829	ENESEPKHEEHPKPEEKPEEEE
830	NESEPKHEEHPKPEEKPEEEEK (Tag 9166-2)
831	KPENESEPKHEEHPKPEEKPEEE
832	PENESEPKHEEHPKPEEKPEEEE (Tag 9166-3)
833	ENESEPKHEEHPKPEEKPEEEEK
834	EGKPENESEPKHEEHPKPEEKPEE
835	GKPENESEPKHEEHPKPEEKPEEE
836	KPENESEPKHEEHPKPEEKPEEEE
837	PENESEPKHEEHPKPEEKPEEEEK
838	EGKPENESEPKHEEHPKPEEKPEEE
839	GKPENESEPKHEEHPKPEEKPEEEE
840	KPENESEPKHEEHPKPEEKPEEEEK
841	EGKPENESEPKHEEHPKPEEKPEEEE
842	GKPENESEPKHEEHPKPEEKPEEEEK
843	EGKPENESEPKHEEHPKPEEKPEEEEK
844	KPPEEDPEEQAEENPEGEQP
845	PPEEDPEEQAEENPEGEQPE (Tag 9590-1)
846	PEEDPEEQAEENPEGEQPEE
847	KPPEEDPEEQAEENPEGEQPE (Tag 9590-2)
848	PPEEDPEEQAEENPEGEQPEE
849	KPPEEDPEEQAEENPEGEQPEE (Tag 9590-3)
850	PDDDDSESDHDDPDNAHESP (Tag 9704-1)
851	GEPEPEPEPEPEPESEPE
852	EPEPEPEPEPEPESEPEP
853	PEPEPEPEPEPEPESEPEPE
854	EPEPEPEPEPEPESEPEPEP
855	PEPEPEPEPEPEPESEPEPE (Tag 9749-1)
856	GGEPEPEPEPEPEPESEPE
857	GEPEPEPEPEPEPESEPEP
858	EPEPEPEPEPEPESEPEPE
859	PEPEPEPEPEPEPESEPEPEP
860	EPEPEPEPEPEPESEPEPEPE
861	GGEPEPEPEPEPEPESEPEP
862	GEPEPEPEPEPEPESEPEPE
863	EPEPEPEPEPEPESEPEPEP
864	PEPEPEPEPEPEPESEPEPE
865	GGEPEPEPEPEPEPESEPEPE (Tag 9749-2)
866	GEPEPEPEPEPEPESEPEPEP
867	EPEPEPEPEPEPESEPEPEPE
868	GGEPEPEPEPEPEPESEPEPEP
869	GEPEPEPEPEPEPESEPEPEPE
870	GGEPEPEPEPEPEPESEPEPEPE (Tag 9749-3)

871	PEEEPDDQDAPDEHESPPPE (Tag 10346-1)
872	EEEEPDDQDAPDEHESPPPED
873	PEEEPDDQDAPDEHESPPPED
874	EEEEPDDQDAPDEHESPPPEDA (Tag 10346-2)
875	EEPDDQDAPDEHESPPPEDAP
876	PEEEPDDQDAPDEHESPPPEDA
877	EEEEPDDQDAPDEHESPPPEDAP
878	PEEEPDDQDAPDEHESPPPEDAP (Tag 10346-3)
879	GPSSDDENEEESKPEKEDEP (Tag 11099-1)
880	PSSDDENEEESKPEKEDEPQ (Tag 11099-2)
881	SDDSDSEKRRPEEQEEEPQP (Tag 12127-1)
882	DDSDSEKRRPEEQEEEPQPR (Tag 12127-2)
883	DPREGEEEEEEDEPDPEAPE
884	PREGEEEEEEDEPDPEAPEN
885	DEEPGDPREGEEEEEEDEPDPE
886	EEPGDPREGEEEEEEDEPDPE
887	EPGDPREGEEEEEEDEPDPEA
888	PGDPREGEEEEEEDEPDPEAP
889	GDPREGEEEEEEDEPDPEAPE
890	DPREGEEEEEEDEPDPEAPEN
891	PREGEEEEEEDEPDPEAPENG
892	EDEEPGDPREGEEEEEEDEPDPE
893	DEEPGDPREGEEEEEEDEPDPE
894	EEPGDPREGEEEEEEDEPDPEA
895	EPGDPREGEEEEEEDEPDPEAP
896	PGDPREGEEEEEEDEPDPEAPE
897	GDPREGEEEEEEDEPDPEAPEN
898	DPREGEEEEEEDEPDPEAPENG
899	PREGEEEEEEDEPDPEAPENG
900	PEEEDEEPGDPREGEEEEEEDEP
901	EEDEEPGDPREGEEEEEEDEPDPE
902	EDEEPGDPREGEEEEEEDEPDPE
903	DEEPGDPREGEEEEEEDEPDPEA
904	EEPGDPREGEEEEEEDEPDPEAP
905	EPGDPREGEEEEEEDEPDPEAPE
906	PGDPREGEEEEEEDEPDPEAPEN
907	DPREGEEEEEEDEPDPEAPENG
908	APEEEDEEPGDPREGEEEEEEDEP
909	PEEEDEEPGDPREGEEEEEEDEPD
910	EEDEEPGDPREGEEEEEEDEPDPE
911	EEDEEPGDPREGEEEEEEDEPDPE
912	EDEEPGDPREGEEEEEEDEPDPEA
913	DEEPGDPREGEEEEEEDEPDPEAP
914	EEPGDPREGEEEEEEDEPDPEAPE
915	EPGDPREGEEEEEEDEPDPEAPEN
916	AAPEEEDEEPGDPREGEEEEEEDEP
917	APEEEDEEPGDPREGEEEEEEDEPD
918	PEEEDEEPGDPREGEEEEEEDEPDPE (Tag 13036-1)

919	EEDEEPPGDPRGEEEEEEDEPDPE
920	EEDEEPPGDPRGEEEEEEDEPDPEA
921	EDEEPPGDPRGEEEEEEDEPDPEAP
922	DEEPPGDPRGEEEEEEDEPDPEAPE
923	EEPPGDPRGEEEEEEDEPDPEAPEN
924	AAPEEEDEEPPGDPRGEEEEEEDEPD
925	APEEEEDEEPPGDPRGEEEEEEDEPDP
926	PEEEDEEPPGDPRGEEEEEEDEPDPE
927	EEDEEPPGDPRGEEEEEEDEPDPEA
928	EEDEEPPGDPRGEEEEEEDEPDPEAP
929	EDEEPPGDPRGEEEEEEDEPDPEAPE
930	DEEPPGDPRGEEEEEEDEPDPEAPEN
931	AAPEEEDEEPPGDPRGEEEEEEDEPDP
932	APEEEEDEEPPGDPRGEEEEEEDEPDPE
933	PEEEDEEPPGDPRGEEEEEEDEPDPEA
934	EEDEEPPGDPRGEEEEEEDEPDPEAP
935	EEDEEPPGDPRGEEEEEEDEPDPEAPE
936	EDEEPPGDPRGEEEEEEDEPDPEAPEN
937	AAPEEEDEEPPGDPRGEEEEEEDEPDPE
938	APEEEEDEEPPGDPRGEEEEEEDEPDPEA
939	PEEEDEEPPGDPRGEEEEEEDEPDPEAP (Tag 13036-2)
940	EEDEEPPGDPRGEEEEEEDEPDPEAPE
941	EEDEEPPGDPRGEEEEEEDEPDPEAPEN
942	APEEEEDEEPPGDPRGEEEEEEDEPDPEA P
943	PEEEDEEPPGDPRGEEEEEEDEPDPEAP E
944	EEDEEPPGDPRGEEEEEEDEPDPEAPE N
945	APEEEEDEEPPGDPRGEEEEEEDEPDPEA PE
946	PEEEDEEPPGDPRGEEEEEEDEPDPEAP EN
947	PRGAAPEEEDEEPPGDPRGEEEEEEDE EPDP
948	RGAAPEEEDEEPPGDPRGEEEEEEDE PDPE
949	AAPEEEDEEPPGDPRGEEEEEEDEPDPE APE
950	APEEEEDEEPPGDPRGEEEEEEDEPDPEA PEN
951	PEEEDEEPPGDPRGEEEEEEDEPDPEAP ENG
952	EEDEEPPGDPRGEEEEEEDEPDPEAPE NGS
953	PRGAAPEEEDEEPPGDPRGEEEEEEDE EPDPE
954	AAPEEEDEEPPGDPRGEEEEEEDEPDPE APEN
955	APEEEEDEEPPGDPRGEEEEEEDEPDPEA PENG
956	PEEEDEEPPGDPRGEEEEEEDEPDPEAP ENG
957	AAPEEEDEEPPGDPRGEEEEEEDEPDPE APENG

958	APEEEEDEEPPGDPRGEEEEEEDEPDPEA PENG
959	AAPEEEDEEPPGDPRGEEEEEEDEPDPE APENG (Tag 13036-3)
960	KPPPSEGSDDEEEEEDEEDE
961	PPPSEGSDDEEEEEDEEDEE (Tag 14128- 1)
962	PPPSEGSDDEEEEEDEEDEERKP
963	KPPPSEGSDDEEEEEDEEDEERKP (Tag 14128-2)
964	PPPSEGSDDEEEEEDEEDEERKPKQ
965	KPPPSEGSDDEEEEEDEEDEERKPKQ (Tag 14128-3)
966	EEEEEEEEEEEEEEEEAPPP
967	EEEEEEEEEEEEEEEEAPPPP (Tag 16549- 1)
968	EEEEEEEEEEEEEEEEAPPPPR
969	EEEEEEEEEEEEEEEEAPPPP
970	EEEEEEEEEEEEEEEEAPPPPR
971	EEEEEEEEEEEEEEEEAPPPP
972	EEEEEEEEEEEEEEEEAPPPPR
973	EEEEEEEEEEEEEEEEAPPPP
974	EEEEEEEEEEEEEEEEAPPPPR (Tag 16549-2)
975	EEEEEEEEEEEEEEEEAPPPP
976	EEEEEEEEEEEEEEEEAPPPPR
977	EEEEEEEEEEEEEEEEAPPPP
978	EEEEEEEEEEEEEEEEAPPPPR
979	EEEEEEEEEEEEEEEEAPPPP
980	EEEEEEEEEEEEEEEEAPPPPR
981	PDDDEEEEEEEEEEEEEEEEEEEEE APPPP (Tag 16549-3)
982	EDSEEEESQEEEAEGASEPPPP (Tag 16604-1)
983	QEDSEEEESQEEEAEGASEPPPP (Tag 16604-2)
984	DHQEDSEEEESQEEEAEGASEPPPP (Tag 16604-3)
985	DQSEEEEEEEKHPPKPAKPE (Tag 16741- 1)
986	PAPAHRRPPEDEGEENEGEEDE (Tag16991-1)
987	PAPAHRRPPEDEGEENEGEEDEE (Tag16991-2)
988	QENGQREEEEEKEPEAEPP (Tag 17199- 1)
989	PAEQENGQREEEEEKEPEAEPP (Tag 17199-2)
990	EQEPEPEPEPEPEPEPEPEPE (Tag 17936- 1)
991	QEPEPEPEPEPEPEPEPEPE
992	EPEPEPEPEPEPEPEPEPEQ (Tag 17936- 2)
993	EQEPEPEPEPEPEPEPEPEPE
994	QEPEPEPEPEPEPEPEPEPEQ
995	EQEPEPEPEPEPEPEPEPEPEQ (Tag 17936-3)

996	PEEEEEEEEEEPASPPERK (Tag 18132-1)	1014	PEEEAEEEEEEEEERPKPSRP (Tag 22900-1)
997	PSPGSPRGQPQDQDDDEDDEED (Tag 18453-1)	1015	QPEEEAEEEEEEEEERPKPSRP
998	PSPGSPRGQPQDQDDDEDDEEDE	1016	EQPEEEAEEEEEEEEERPKPSRP (Tag 22900-2)
999	RSPGSPRGQPQDQDDDEDDEEDE (Tag 18453-2)	1017	EEQPEEEAEEEEEEEEERPKPSRP
1000	PSPGSPRGQPQDQDDDEDDEEDEA (Tag 18453-3)	1018	EEEQPEEEAEEEEEEEEERPKPSRP (Tag 22900-3)
1001	AEDDDEEDEEEEEEPDPP (Tag 18866-1)	1019	PEEEDDEAQQPQPQSGPEEAEE (Tag 34831-1)
1002	EDDDEEDEEEEEEPDPP (Tag 18866-2)	1020	DKPEEEDDEAQQPQPQSGPEEAEE
1003	KQEPPDPEEDKEENKDDSAS (Tag 19350-1)	1021	PEEEDDEAQQPQPQSGPEEAEEGE
1004	QEPPDPEEDKEENKDDSASK (Tag 19350-2)	1022	PEEEDDEAQQPQPQSGPEEAEEGEE
1005	EDEDEDESSEEDSEDEEPPP (Tag 19511-1)	1023	EGDKPEEEDDEAQQPQPQSGPEEAEE (Ta 34831-2)
1006	DEDEDESSEEDSEDEEPPPK	1024	DKPEEEDDEAQQPQPQSGPEEAEEGE
1007	EDEDESSEEDSEDEEPPPKR	1025	KPEEEDDEAQQPQPQSGPEEAEEGEE
1008	DEDESSEEDSEDEEPPPKRR	1026	PEEEDDEAQQPQPQSGPEEAEEGEEE
1009	PDDSRDEDEDESSEEDSEDEEPPP (Tag 19511-2)	1027	PEEEDDEAQQPQPQSGPEEAEEGEEEE AERGP
1010	PKKEPDDSRDEDEDESSEEDSEDEEPP	1028	KPEEEDDEAQQPQPQSGPEEAEEGEEEE EAERGP (Tag 34831-3)
1011	PKKEPDDSRDEDEDESSEEDSEDEEPPK	1029	NNSEEEEDDDDEEEEPDKPP (Tag 39056-1)
1012	PKKEPDDSRDEDEDESSEEDSEDEEPPKR	1030	NSEEEEDDDDEEEEPDKPPA (Tag 39056-2)
1013	PKKEPDDSRDEDEDESSEEDSEDEEPPKRR (Tag 19511-3)	1031	SEEEEDDDDEEEEPDKPPAN (Tag 39056-3)

[0223]

As shown by these examples, the protein aggregation rate reducing action of a peptide tag had low dependency on a specific amino acid sequence, also had low dependency on a protein from which it is derived, but had high dependency on amino acid contents.

[0224]

Example 6: Addition to Various Proteins and Aggregation Inhibiting Action

In the above Example, Y13-259 was used as the scFv. In this example, a VHH antibody (a heavy chain variable

domain of a heavy chain antibody) was used. As the VHH antibody, iDab#6 binding to Ras was used. As the tag, Tag4-8 was used. The other conditions were the same as those employed in Example 1. As a result, the aggregation rate of the VHH antibody not having the tag was 57.89%, the aggregation rate of the VHH antibody having Tag4-8 at the C terminal was 8.77%, and thus, a strong aggregation inhibiting action was exhibited by the tag addition.

[0225]

When SHSY5Y (human dopamine-like cell) that is a neuroblastoma cell line was used as the cell, scFv-6E (6E) was used as the scFv, and Tag4-8 or Tag18-1 was added as the tag to the C terminal and the N terminal, the aggregation rate of the Tag4-8 added scFv was 0.96%, the aggregation rate of the Tag18-1 added scFv was 1.14%, and thus, it was revealed that the tag addition makes a contribution to the low aggregation rate.

[0226]

When D4 binding to botulinum toxin type A (SEQ ID NO: 1032:
QVQLQQSGGGLVQPGGSLRLSCAASGFTLDYYAIGWFRQAPGKEREGVLCISSSGGS
TNYADSVKGRFTISRDNAKNTVYLQMNSLPEDTAVYYCAADDLRCSNWSSYFRGS
WGQGTQVTVSS) was used as the VHH antibody, and the Tag4-8 was added as the tag to the C terminal of the VHH antibody, the aggregation rate of the Tag4-8 added D4 was 7.8%, and the aggregation rate of D4 not having the tag

was 81.5%. This result reveals that the tag addition makes a contribution to the reduction of the aggregation rate of the VHH antibody.

[0227]

Example 7: Action Enhancement by Stabilization of Intracellular Antibody (Effect of Enhancement of Antibody Action on Amyloid Accumulation)

It is known that a central nervous system disease is caused by accumulation of amyloid in a nerve cell. When human α -synuclein fibril is extracellularly introduced into a nerve cell, synuclein fibril is formed with synuclein having a normal structure in the cell involved. When GFP-tagged synuclein has been expressed in the cell, the synuclein forms synuclein fibril together with the introduced human α -synuclein fibril, and the thus formed synuclein fibril can be observed with fluorescence of GFP. In this example, GFP-tagged synuclein was expressed in SHSY-5Y cell, and α -synuclein fibril (Cosmo Bio Co., Ltd., SYNO3) was extracellularly introduced into the SHSY-5Y cell. In this example, the ability of an antibody to reduce the synuclein fibril was tested. Specifically, as the antibody, scFv-6E binding to fibrilized synuclein (SEQ ID NO: 1033:

AEVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSYIASGGD
TTNYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKGASAFDYWGQGT
LVTVSSGGGGSGGGGSGGGGSTDIQMTQSPSSLSASVGDRVTITCRASQSISSYL
NWYQQKPGKAPKLLIYAASYLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQ
QSSNDP

YTFGQG~~TK~~VEIKR) was used. The scFv-6E is an antibody not having a significant binding property to a monomer or oligomer of synuclein, and such an antibody selective or specific to synuclein fibril is suitable for selectively removing synuclein fibril. To the scFv-6E, a tag having the aggregation rate reducing action (Tag4-8 or Tag18-1) and a degradation-inducing sequence (CMA (SEQ ID NO: 1034): MARVKKDQAEPLHRKFERQPPG) were added. An expression plasmid vector for the protein and the synuclein fibril were introduced into the cell respectively with X-trem GENE9 and Multifectam (Merck). The antibody was provided with an HA tag or a myc tag, and detected with an anti-HA tag antibody. The synuclein fibril was detected with an anti-phosphorylated α -synuclein antibody. These antibodies were specifically detected with Alexa 555 labeled antibody and Alexa 633 labeled antibody. Fluorescent stained cells were observed with Keyence BZ-X800.

[0228]

The tag was added to the N terminal and the C terminal. In consideration that the aggregation rate of one having Tag4-8 added to the N terminal and the C terminal was 0.96% in the SHSY-5Y cell, that the aggregation rate of one having TAG18-1 added to the N terminal and the C terminal was 1.14% in the SHSY-5 cell, and that the aggregation rate of the scFv-E6 having no

tag added was 41.6% in the HeLa cell, it seems that favorable aggregation inhibition was exhibited.

[0229]

When a scFv binds to synuclein fibril, the lysosome is caused to target the synuclein fibril for a degradation-inducing sequence to degrade the synuclein fibril, and as a result, the amount of synuclein fibril in the cell is expected to be reduced. It was evaluated whether or not the reduction amount of the synuclein fibril is increased when the aggregation rate of the scFv was reduced by using the tag to increase the amount of functional scFv in the cell.

[0230]

The antibody was expressed in the cell where the synuclein fibril was formed, and the number of synuclein fibril-positive cells was counted. A positive rate of the synuclein fibril was compared between a cell in which the antibody was expressed and a cell in which it was not expressed. The result was obtained as a rate of synuclein fibril-positive cells in antibody-positive cells/a rate of synuclein fibril-positive cells in antibody-negative cells (P/N).

[0231]

As illustrated in Figure 2, phosphorylated synuclein was lost in the cell into which the tagged scFv-E6 was introduced. The P/N was as illustrated in Figure 3. As illustrated in Figure 3, both the scFv-E6 tagged with

Tag4-8 and the scFv-E6 tagged with Tag18-1 largely reduced the rate of synuclein-positive cells.

[0232]

Example 8: Enhancement of Action by Stabilization of Intracellular Antibody (Effect of Enhancement of Antibody Action for Recovering Function of CFTR)

The cystic fibrosis transmembrane conductance regulator (CFTR; UniprotKB/Swiss-Prot: P13569.3) is a negative ion channel expressed in epithelial membrane cells of the whole body, and abnormality thereof causes cystic fibrosis. F508 deletion mutation of CFTR (CFTR Δ F508) is known as the most common mutation of CFTR, in which the 508th phenylalanine is deleted due to deletion of three nucleotides. As a result, CFTR Δ F508 cannot be normally folded, strongly tends to form an aggregation, and is deemed to move onto the membrane in a smaller amount than the wild type. The scFv-C2 (SEQ ID NO: 1035:

EVQLLES GGGGLVQP GGS LRLSCAASGFT FSSYAMSWVRQAPGKGLEWVSAISGSGGS
TYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKMRLGLFDYWGQGT LVT
VSSGGGGSGGGSGGGGEIVLTQSPGTL SLSPGERATLSCRASQSVSSSYLAWYQQK
PGQAPRL LIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYYCQQRGDVPP
TFGQGTKVEIKAAA) binds to the NBD1 domain of CFTR, and thus can inhibit formation of an aggregation by the NBD1 (Lovato et al., Protein Engineering, Design and Selection, 20(12): 607-614, 2007). The scFv-C2 exhibits an effect of increasing the amount of CFTR Δ F508 moving

onto the membrane. In this example, it was evaluated whether or not aggregation of the mutant $\Delta F508$ in the NBD1 domain in the HeLa cell can be inhibited by adding a tag (Tag18-1) to the scFv-C2. Plasmid vectors for expressing these proteins were introduced into HeLa cell with Lipo3000(TM). An antibody was tagged with a myc tag to be detected with a rabbit anti-myc tag antibody, and the NBD1 was tagged with a His tag to be detected with a mouse anti-His tag antibody. The mouse anti-His tag antibody was detected with Alexa 633-labeled anti-mouse IgG antibody, and the rabbit anti-myc tag antibody was detected with Alexa 488-labeled anti-rabbit IgG antibody. Fluorescent stained cells were observed with Keyence BZ-X800.

[0233]

First, the aggregation rate of the scFv-C2 in HeLa cell was 82%. On the contrary, the aggregation rate of the scFv-C2 having Tag18-1 at the N terminal was 31%, and the aggregation rate of the scFv-C2 having Tag18-1 at the N terminal and the C terminal was 4.6% (see Figure 3A and Figure 3B). The aggregation rate of wild type NBD1 domain was 74% in a cell expressing scFv-C2, and the aggregation rate of the $\Delta F508$ mutant of the NBD1 was 85% in a cell expressing scFv-C2. On the contrary, in a cell expressing scFv-C2 having Tag18-1 at the N terminal and the C terminal, the aggregation rate of the wild type NBD1 domain was 32%, and the aggregation rate of the

ΔF508 mutant of NBD1 was 43%. In this manner, aggregation formation of the scFv-C2 itself could be inhibited by tagging the scFv-C2, and thus, the formation of an aggregation by the NBD1 could be inhibited. The inhibition of the aggregation of the NBD1 is expected to make a contribution to improvement of expression level of the NBD1 in cell membrane.

[0234]

Example 9: Effect of Amino Acid Substitution in Existing Tag

Through Examples described above, it was revealed that amino acid substitution for satisfying the condition needed for the peptide tag of the present disclosure, particularly amino acid substitution with P or N increases the aggregation rate reducing action of the tag. In this example, with some amino acids of PEST sequence substituted with N, the aggregation rates of the scFvs (Y14-259) having tags before and after the substitution were examined.

[0235]

[Table 15]

Table 15: Comparison of sequence between before and after substitution

PEST(before substitution) SEQ ID NO: 1036	YPYDVPDYAGSPQPVEDGEDEFCTPMACEANIQSGDSAAPMSAVHRHRL
PEST(after substitution) SEQ ID NO: 1037	NNYDVPDNAGSPQPQEDGEDEFNNPQANEANQQSGDSNNPNSAVNRHNN

[0236]

After these scFvs were expressed in a cell, the aggregation rates of the scFvs were evaluated in the same manner as in Example 1, and the aggregation rate of the scFv having the tag before the substitution was 50.3%, but the aggregation rate of the scFv having the tag after the substitution was 18.0%. In this manner, it was revealed that the amino acid substitution for satisfying the condition needed for the peptide tag of the present disclosure, particularly the amino acid substitution with P or N increases the effect of inhibiting protein aggregation of the tag.

[0237]

As described so far, various proteins including antibodies form aggregation in a cell, and thus, the functions can be partially or entirely impaired. A protein tag for inhibiting the formation of an aggregation can inhibit the aggregation formation of these proteins, and thus, can cause the proteins to exhibit their actions to be originally exhibited. A tag in which an acidic amino acid ratio is relatively low can be helpful in a scene where a tag having a high acidic amino acid ratio is difficult to use.

Claims

[Claim 1]

A peptide, wherein

(a) 5% or more and less than 45% of amino acids contained in an amino acid sequence thereof are acidic amino acids, and

(b) 20% or more, and preferably 30% or more of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A, and

the peptide is capable of reducing an aggregation property in a cell of a protein linked to the peptide.

[Claim 2]

The peptide according to claim 1, wherein

(c) 30% or less, preferably 20% or less, more preferably 15% or less, and further preferably 10% or less of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of M, T, W, C, I, V, and L.

[Claim 3]

The peptide according to claim 1 or 2, wherein

(d) each of A and G constitutes less than 10% of the amino acids contained in the amino acid sequence.

[Claim 4]

The peptide according to any one of claims 1 to 3, wherein

(a) 20% or more and less than 45% of the amino acids contained in the amino acid sequence are acidic amino acids,

(b) 30% or more and less than 70% of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A,

(c) 20% or less of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of M, T, W, C, I, V, and L, and

(d) each A and G constitutes less than 10% of the amino acids contained in the amino acid sequence.

[Claim 5]

A peptide having an amino acid sequence set forth in any one of SEQ ID NOs: 2 to 11.

[Claim 6]

A nucleic acid encoding the peptide according to any one of claims 1 to 5.

[Claim 7]

A protein expression vector comprising: the nucleic acid according to claim 6 operably linked to a regulatory sequence; and a nucleic acid encoding a protein of interest in-frame to the nucleic acid according to claim 6.

[Claim 8]

The protein expression vector according to claim 7, wherein the protein of interest is an antibody, or an antigen-binding fragment of an antibody.

[Claim 9]

The protein expression vector according to claim 8, wherein the antigen-binding fragment of the antibody is a single chain Fv (scFv).

[Claim 10]

A fusion protein of the peptide according to any one of claims 1 to 5 and a protein of interest.

[Claim 11]

The fusion protein according to claim 10, wherein the protein of interest is an antibody, or an antigen-binding fragment of an antibody.

[Claim 12]

The fusion protein according to claim 11, wherein the antigen-binding fragment of the antibody is a single chain Fv (scFv).

[Claim 13]

A protein-producing cell comprising: the nucleic acid according to claim 6 operably linked to a regulatory sequence; and a nucleic acid encoding a protein of interest in-frame to the nucleic acid according to claim 6.

[Claim 14]

A method for selecting or identifying an amino acid sequence, comprising:

acquiring an amino acid sequence in which:

(a) 5% or more and less than 45% of amino acids contained in the amino acid sequence are acidic amino acids; and

(b) 20% or more of the amino acids contained in the amino acid sequence are amino acids selected from the group consisting of F, P, Y, G, S, Q, N, and A;

selecting or identifying an amino acid sequence of a peptide tag that, when a fusion protein of the peptide tag having the selected or identified amino acid sequence and a reference protein is expressed in a mammal cell (preferably in a human cell), provides reduction of a proportion of cells in which the fusion protein forms an aggregation, or the proportion which is not more than a predetermined value; and

obtaining the peptide tag having the amino acid sequence or a nucleic acid encoding the peptide tag.

[Claim 15]

The method according to claim 14, wherein the amino acid sequence to be acquired is the peptide according to any one of claims 1 to 5.

[Claim 16]

The method according to claim 14 or 15, wherein the amino acid sequence to be acquired is a group of amino acid sequences encoded by coding regions of human genome.

[Claim 17]

The method according to any one of claims 14 to 16,
wherein the amino acid sequence to be acquired contains a
neo-antigen.