ANTI-JAGGED1 ANTIBODIES AND METHODS OF USE

FIELD OF THE INVENTION

The present invention relates to anti-Jagged antibodies and methods of using the same.

BACKGROUND

The Notch signaling pathway regulates a diverse array of cell functions (Kopan et al., *Cell* 137, 216–233 (2009)). Four Notch receptors have been identified in mammals, i.e., Notch 1-4, that share basic structural elements that include an extracellular domain, a transmembrane domain, and an intracellular domain. Similarly, the canonical ligands of Notch share certain structural similarities but a number of non-canonical ligands of Notch have also been identified (Kopan et al., *Cell* 137, 216–233 (2009)). The five canonical ligands in mammals are Delta-like 1, Delta-like 3, Delta-like 4, Jagged1 and Jagged2. Binding of a Notch ligand to the extracellular domain of a Notch receptor sets a signaling cascade in motion that begins with proteolytic cleavage at the extracellular S2 site by an alpha secretase of the ADAM (a disintegrin and metalloprotease) family. Cleavage at S2 is followed by proteolytic cleavage by a gamma secretase at the intracellular S3 site, which results in release of the intracellular domain and downstream events that ultimately activate Notch-dependent transcription factors such as Hes1 and Hey.

Aberrant Notch expression and signaling has been implicated in a number of diseases, including cancer (Koch et al., *Cell. Mol. Life Sci.* 64, 2746–2762 (2007)). It is clear that there continues to be a need for agents that have clinical attributes that are optimal for development as therapeutic agents. The invention described herein meets this need and provides other benefits.

SUMMARY

The ~~invention~~ disclosure provides anti-Jagged1 antibodies and methods of using the same.

The present inventors unexpectedly found that anti-Jagged1 antibody A-1 (*see* Figure 4 and PCT Publication No. 2014/028446) is cleaved in the heavy chain following heat treatment and/or freeze-thaw conditions. The poor stability of the antibody potentially reduces its value as a therapeutic. Analysis of the cleavage site revealed no known protease cleavage motifs. It was therefore not known whether changes to the antibody sequence could reduce the observed

cleavage. Further, because the cleavage site is in a heavy chain HVR, even if amino acid change(s) would reduce cleavage, it was not known whether such changes could be made without significantly reducing the antibody's affinity for Jagged1. The inventors found, surprisingly, that mutation of the amino acid at position 101 in HVR-H3 reduced cleavage with only a slight loss of affinity.

In ~~some embodiments~~one aspect, an isolated antibody that binds to human Jagged1 is provided, wherein the antibody comprises an HVR-H3 comprising the amino acid sequence of SEQ ID NO: 55 or 59, wherein X is any amino acid other than S. In some embodiments, the antibody comprises at least one, at least two, at least three, at least four, or five HVRs selected from HVR-H1 comprising the amino acid sequence of SEQ ID NO: 35 or 78; HVR-H2 comprising the amino acid sequence of SEQ ID NO: 28 or 36; HVR-L1 comprising the amino acid sequence of SEQ ID NO: 38; HVR-L2 comprising the amino acid sequence of SEQ ID NO: 39; and HVR-L3 comprising the amino acid sequence of SEQ ID NO: 16 or 40. In some embodiments, the antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 78; HVR-H2 comprising the amino acid sequence of SEQ ID NO: 36; and HVR-H3 comprising the amino acid sequence of SEQ ID NO: 55; or (b) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 78; HVR-H2 comprising the amino acid sequence of SEQ ID NO: 36; and HVR-H3 comprising the amino acid sequence of SEQ ID NO: 59; or (c) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 78; HVR-H2 comprising the amino acid sequence of SEQ ID NO: 28; and HVR-H3 comprising the amino acid sequence of SEQ ID NO: 59. In any of the embodiments described herein, the antibody may comprise (a) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 38; HVR-L2 comprising the amino acid sequence of SEQ ID NO: 39; and HVR-L3 comprising the amino acid sequence of SEQ ID NO: 40; or (b) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 38; HVR-L2 comprising the amino acid sequence of SEQ ID NO: 39; and HVR-L3 comprising the amino acid sequence of SEQ ID NO: 16; or (c) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 38; HVR-L2 comprising the amino acid sequence of SEQ ID NO: 39; and HVR-L3 comprising the amino acid sequence of SEQ ID NO: 16.

In some embodiments, an isolated antibody that binds to Jagged1 comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 78; HVR-H2 comprising the amino acid sequence of SEQ ID NO: 36; HVR-H3 comprising the amino acid sequence of SEQ ID NO: 55; HVR-L1 comprising the amino acid sequence of SEQ ID NO: 38; HVR-L2 comprising the amino acid sequence of SEQ ID NO: 39; and HVR-L3 comprising the amino acid sequence of SEQ ID NO: 40; or (b) HVR-H1 comprising the amino acid sequence of SEQ

ID NO: 78; HVR-H2 comprising the amino acid sequence of SEQ ID NO: 36; HVR-H3 comprising the amino acid sequence of SEQ ID NO: 59; HVR-L1 comprising the amino acid sequence of SEQ ID NO: 38; HVR-L2 comprising the amino acid sequence of SEQ ID NO: 39; and HVR-L3 comprising the amino acid sequence of SEQ ID NO: 16; or (c) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 78; HVR-H2 comprising the amino acid sequence of SEQ ID NO: 28; HVR-H3 comprising the amino acid sequence of SEQ ID NO: 59; HVR-L1 comprising the amino acid sequence of SEQ ID NO: 38; HVR-L2 comprising the amino acid sequence of SEQ ID NO: 39; and HVR-L3 comprising the amino acid sequence of SEQ ID NO: 16.

In some embodiments, an isolated antibody that binds to Jagged1 comprises a VH sequence having at least 95% identity to the amino acid sequence of SEQ ID NO: 54, 58, or 62. In some embodiments, an isolated antibody that binds to Jagged1 comprises a VL sequence having at least 95% identity to the amino acid sequence of SEQ ID NO: 10, 26, or 34. In some embodiments, the antibody comprises (a) a VH sequence having at least 95% identity to the amino acid sequence of SEQ ID NO: 54 and a VL sequence having at least 95% identity to the amino acid sequence of SEQ ID NO: 34; or (b) a VH sequence having at least 95% identity to the amino acid sequence of SEQ ID NO: 58 and a VL sequence having at least 95% identity to the amino acid sequence of SEQ ID NO: 10; or (c) a VH sequence having at least 95% identity to the amino acid sequence of SEQ ID NO: 62 and a VL sequence having at least 95% identity to the amino acid sequence of SEQ ID NO: 26.

In ~~some embodiments~~one aspect, an isolated antibody that binds to human Jagged1 is provided, comprising ~~comprises~~ (a) a VH sequence of SEQ ID NO: 54, wherein X is any amino acid other than S, and a VL sequence of SEQ ID NO: 34; or (b) a VH sequence of SEQ ID NO: 58, wherein X is any amino acid other than S, and a VL sequence of SEQ ID NO: 10; or (c) a VH sequence of SEQ ID NO: 62, wherein X is any amino acid other than S, and a VL sequence of SEQ ID NO: 26. In some embodiments, the antibody comprises a VH sequence of SEQ ID NO: 54 and a VL sequence of SEQ ID NO: 34. In some embodiments, the heavy chain comprises the amino acid sequence of SEQ ID NO: 56 and the light chain comprises the amino acid sequence of SEQ ID NO: 53. In some embodiments, the heavy chain comprises the amino acid sequence of SEQ ID NO: 57 and the light chain comprises the amino acid sequence of SEQ ID NO: 53.

In any of the embodiments described herein, X may be any amino acid other than S or H. In any of the embodiments described herein, X may be selected from A, D, E, G, I, K, L, N, Q, R T, and V. In any of the embodiments described herein, X may be T.

In ~~some embodiments~~one aspect, an isolated antibody that binds human Jagged1 is provided, comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 78; HVR-H2 comprising the amino acid sequence of SEQ ID NO: 36; HVR-H3 comprising the amino acid sequence of SEQ ID NO: 37; HVR-L1 comprising the amino acid sequence of SEQ ID NO: 38; HVR-L2 comprising the amino acid sequence of SEQ ID NO: 39; and HVR-L3 comprising the amino acid sequence of SEQ ID NO: 40; or (b) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 78; HVR-H2 comprising the amino acid sequence of SEQ ID NO: 36; HVR-H3 comprising the amino acid sequence of SEQ ID NO: 64; HVR-L1 comprising the amino acid sequence of SEQ ID NO: 38; HVR-L2 comprising the amino acid sequence of SEQ ID NO: 39; and HVR-L3 comprising the amino acid sequence of SEQ ID NO: 16; or (c) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 78; HVR-H2 comprising the amino acid sequence of SEQ ID NO: 28; HVR-H3 comprising the amino acid sequence of SEQ ID NO: 64; HVR-L1 comprising the amino acid sequence of SEQ ID NO: 38; HVR-L2 comprising the amino acid sequence of SEQ ID NO: 39; and HVR-L3 comprising the amino acid sequence of SEQ ID NO: 16.

In some embodiments, an isolated antibody that binds human Jagged1 is provided, comprising (a) a VH sequence of SEQ ID NO: 33 and a VL sequence of SEQ ID NO: 34; or (b) a VH sequence of SEQ ID NO: 65 and a VL sequence of SEQ ID NO: 10; or (c) a VH sequence of SEQ ID NO: 66 and a VL sequence of SEQ ID NO: 26. In some embodiments, the antibody comprises a VH sequence of SEQ ID NO: 33 and a VL sequence of SEQ ID NO: 34.

In any of the above embodiments described herein, the antibody may be a monoclonal antibody. In certain embodiments, the antibody is a human, humanized, or chimeric antibody. In any of the above embodiments described herein, the antibody may be an antibody fragment.

Any of the above embodiments may be a full-length IgG1 antibody. In some embodiments, the antibody is an IgG1 antibody lacking effector function. In some embodiments, the antibody is an IgG1 antibody comprising an N297G or N297A mutation. In some embodiments, the antibody is an IgG1 antibody comprising an N297G mutation.

In some embodiments, an isolated antibody that binds to Jagged1 is provided, wherein the heavy chain comprises the amino acid sequence of SEQ ID NO: 51 and the light chain comprises the amino acid sequence of SEQ ID NO: 53. In some embodiments, the heavy chain comprises the amino acid sequence of SEQ ID NO: 52 and the light chain comprises the amino acid sequence of SEQ ID NO: 53. In some embodiments, the antibody comprises (a) a heavy chain comprising the amino acid sequence of SEQ ID NO: 69 and a light chain comprising the amino acid sequence of SEQ ID NO: 75; or (b) a heavy chain comprising the amino acid

stronger, or 0.9 nM or stronger, 0.8 nM or stronger, 0.7 nM or stronger, or 0.6 nM or stronger, or 0.5 nM or stronger (i.e., less than 1.5 nM, less than 1 nM, less than 0.9 nM, less than 0.8 nM, less than 0.7 nM, less than 0.6 nM, or less than 0.5 nM). In some embodiments, affinity (Kd) is measured using surface plasmon resonance.

In some embodiments, the antibody binds human Jagged1 with an association constant (k_{on}) of at least 1.0E+04/Ms, or at least 1.5E+04/Ms, or at least 2.0E+04/Ms. In some embodiments, the antibody binds human Jagged1 with a dissociation constant (k_{off}) of less than 10.0E-04/s, or less than 9.0E-04/s, or less than 8.0E-04/s, or less than 7.0E-04/s. In some embodiments, the antibody binds murine Jagged1 with an association constant (k_{on}) of at least 1.0E+04/Ms, or at least 1.5E+04/Ms, or at least 2.0E+04/Ms, or at least 3.0E+04/Ms, or at least 4.0E+04/Ms, or at least 5.0E+04/Ms, or at least 6.0E+04/Ms, or at least 7.0E+04/Ms. In some embodiments, the antibody binds murine Jagged1 with a dissociation constant (k_{off}) of less than 10.0E-04/s, or less than 9.0E-04/s, or less than 8.0E-04/s, or less than 7.0E-04/s. In some embodiments, association and dissociation constants are measured using surface plasmon resonance.

In some embodiments, the antibody binds native, folded Jagged1, but does not bind denatured Jagged1. In some embodiments, the antibody binds Jagged1 in an enzyme-linked immunosorbent assay (ELISA) but does not bind Jagged1 on a Western blot. In some embodiments, the antibody binds folded Jagged1 in an enzyme-linked immunosorbent assay (ELISA) but does not bind denatured Jagged1 on a Western blot. In some embodiments, the antibody binds folded Jagged1 under physiological conditions but does not bind denatured Jagged1.

In some embodiments, the antibody reduces tumor growth in a mouse xenograft model without causing weight loss. In some embodiments, the mouse xenograft model is a liver cancer xenograft model. In some embodiments, tumor growth is reduced by at least 50, at least 60, at least 70, at least 80, or at least 90 AUC/day TGI%.

In some embodiments, use of the Jagged1 antibody reduces goblet cell metaplasia in lungs in a mouse model of airway hyperresponsiveness. In some embodiments, administration of the antibody reduces the number of goblet cells in the lungs by at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90%.In another ~~aspect~~embodiment, the ~~invention~~ disclosure provides an isolated antibody that competes with any of the above embodiments for specific binding to Jagged1. In some embodiments, an antibody is that competes for binding to human Jagged1 with an antibody comprising a heavy chain variable region comprising the

sequence of SEQ ID NO: 33 and a light chain variable region comprising the sequence of SEQ ID NO: 34, wherein the antibody is not antibody A, antibody A-1, or antibody A-2.

In another aspect, ~~the invention provides~~ an isolated nucleic acid encoding an isolated antibody ~~of the above embodiments~~described above is provided. In a further aspect, ~~the invention provides~~ a host cell comprising the isolated nucleic acid ~~encoding the antibody~~described above is provided. In a further aspect, ~~the invention provides~~ a method of producing an antibody is provided, comprising culturing the host cell described above so that the antibody is produced.

In another aspect, ~~the invention provides~~ an immunoconjugate is provided, comprising any of the antibodies described~~an antibody of any of the~~ above ~~embodiments~~ and a cytotoxic agent.

In another aspect, ~~the invention provides~~ a pharmaceutical formulation is provided, comprising any of the antibodies described ~~an antibody of any of the~~ above ~~embodiments~~ and a pharmaceutically acceptable carrier. In another aspect, a pharmaceutical formulation is provided, comprising an immunoconjugate described above and a pharmaceutically acceptable carrier.

In another aspect, any of the antibodies or immunoconjugates described~~an antibody of any of the~~ above ~~embodiments~~ is provided for use as a medicament. In ~~some embodiments~~another aspect, any of the antibodies or immunoconjugates described~~an antibody of any of the~~ above ~~embodiments~~ is provided for use in treating a cancer. In ~~some embodiments~~another aspect, any of the antibodies or immunoconjugates described ~~an antibody of any of the~~ above ~~embodiments~~ is provided for use in reducing cancer cell growth.

In ~~another aspect~~one embodiment, a method of inhibiting Jagged1-mediated signaling is provided. In one embodiment, a method of inhibiting Jagged1-mediated signaling *in vitro* is provided. In one embodiment, a method of inhibiting Jagged1-mediated signaling *in vivo* is provided.

In one aspect, use of any of the antibodies or immunoconjugates described above in the manufacture of a medicament is provided. In ~~another aspect~~one embodiment, a method of treating an individual having a cancer comprising administering to the individual an effective amount of an antibody of any of the above embodiments is provided. In one embodiment, the cancer is selected from the group consisting of: breast cancer, lung cancer, brain cancer, cervical cancer, colon cancer, liver cancer, bile duct cancer, pancreatic cancer, skin cancer, B-cell malignancies, and T-cell malignancies.

The present inventors discovered that treatment with Jagged1-antibody biases cell fate in the airways away from a secretory cell (including goblet cell) fate and toward a ciliated cell fate. Jagged1 signaling is important for maintaining the secretory cell fate, and inhibition of Jagged1 signaling prevented goblet cell metaplasia. The present inventors also showed that the club cell-to-ciliated cell conversion is direct and did not involve cell division (data not shown). This transdifferentiation of one cell type to another occurred in the adult lung and is distinct from cell fate choices that involve progenitor cell division, such as after damage or during development. Goblet cell metaplasia or excess mucus is a hallmark of several airway diseases, such as asthma, cystic fibrosis, COPD and Barrett's esophagus. These Jagged inhibition results provide the basis for therapeutic applications involving use of Jagged1 or Jagged2 inhibitors for prevention or reversal of goblet cell metaplasia and for treatment of conditions

with goblet cell metaplasia. In some embodiments, the disease is selected from asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis, and Barrett's esophagus.

In some embodiments, an antibody that binds to human Jagged1 (including, for example, any of the anti-Jagged1 antibodies described herein) is provided for use in treating a disease selected from allergy, asthma, autoimmune disease, diseases associated with goblet cell metaplasia (e.g., in lung) and diseases associated with excess mucus is provided, comprising administering to an individual with cancer an effective amount of an antibody of any of the embodiments described herein, wherein the antibody that binds to human Jagged1 increases ciliated cell number in the adult human airway (e.g., lung). In some embodiments, the disease is associated with goblet cell metaplasia. In some embodiments, the disease is selected from asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis, and Barrett's esophagus.

In some embodiments, an antibody that binds to human Jagged1 (including, for example, any of the anti-Jagged1 antibodies described herein) is provided for use in treating a disease selected from allergy, asthma, autoimmune disease, diseases associated with goblet cell metaplasia (e.g., in lung) and diseases associated with excess mucus is provided, comprising administering to an individual with cancer an effective amount of an antibody of any of the embodiments described herein, wherein the antibody that binds to human Jagged1 increases formation of ciliated cells in the adult human airway (e.g., lung). In some embodiments, the disease is associated with goblet cell metaplasia. In some embodiments, the disease is selected from asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis, and Barrett's esophagus.

~~In another aspect,~~Also provided are methods of (a) converting a ciliated cell to a club cell (e.g., wherein the ciliated cell is found in the adult human airway, e.g., lung), (b) increasing mucus (e.g., airway mucus), (c) decreasing ciliated cell number (e.g., human airway ciliated cell), comprising administering an agonist of Jagged signaling to an individual.

In some embodiments, a method of treating a disease selected from allergy, asthma, autoimmune disease, diseases associated with goblet cell metaplasia (e.g., in lung) and diseases associated with excess mucus is provided, comprising administering to an individual with cancer an effective amount of antibody that binds to human Jagged1 (including, for example, any of the antibodies that bind to human Jagged1 described herein). In some embodiments, the disease is associated with goblet cell metaplasia (e.g., in the lung). In some embodiments, the disease is selected from asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis, and Barrett's esophagus.

In any of the embodiments described herein, wherein the Jagged1 antibody reduces tumor growth in a mouse xenograft model without causing weight loss.

In ~~any of the embodiments described herein~~one aspect, ~~the~~ an anti-Jagged1 antibody as described above ~~may be~~ conjugated to a label is provided. In some embodiments, the label is a positron emitter. In some embodiments, the positron emitter is ^{89}Zr. In ~~some embodiments~~one aspect, a method of detecting human Jagged1 in a biological sample is provided, comprising contacting the biological sample with an antibody described herein under conditions permissive for binding of the antibody to a naturally occurring human Jagged1, and detecting whether a complex is formed between the antibody and a naturally occurring human Jagged1 in the biological sample. In some embodiments, the biological sample is selected from breast cancer, lung cancer, brain cancer, cervical cancer, colon cancer, liver cancer, bile duct cancer, pancreatic cancer, skin cancer, B-cell malignancies, and T-cell malignancies.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 shows exemplary amino acid sequences of human and murine Jagged1 protein.

FIG. 2 shows exemplary amino acid sequences of human and murine Jagged2 protein.

FIGS. 3A-D show the amino acid sequences of peptides used for phage antibody library screening and selection. All proteins were expressed as a secreted protein in BEVS cells and their sequences are listed in the N-terminal to C-terminal direction. (A) Amino acid sequence of expressed protein murine Jagged 1-DSL-EGF1-4 (Q34-D377). The bold font at the N-terminus represents a short linker sequence (ADLGS) (SEQ ID NO: 82). The bold font at the C-terminus represents a short linker sequence (EFG), a thrombin cleavage site (LVPRGS) (SEQ ID NO: 83), a G spacer and the 6-His tag. (B) Amino acid sequence of expressed protein human Jag1-DSL-EGF1-4. Only the Jag1 sequence is shown although the antigen also contained a TEV protease cleavage site and 6-His tag at the C-terminus. (C) Amino acid sequence of expressed protein murine Jag2-DSL-EGF1-4 (M27-E388). The bold font at the N-terminus represents a short linker sequence (ADLGS) (SEQ ID NO: 82). The bold font at the C-terminus represents a short linker sequence (EFG), a thrombin cleavage site (LVPRGS) (SEQ ID NO: 83), a G spacer and the 6-His tag. (D) Amino acid sequence of expressed protein human Jag2-DSL-EGF1-4 (R2-E388). The bold font at the C-terminus represents a short linker sequence (EFG), a thrombin cleavage site (LVPRGS) (SEQ ID NO: 83), a G spacer and the 6-His tag.

FIGS. 4A-1-B-2 show an alignment of the amino acid sequences for the heavy (FIG. 4A-1 and FIG. 4A-2) and light (FIG. 4B-1 and FIG. 4B-2) chain variable domains of anti-

41. The antibody of any one of claims 1 to 33 or the immunoconjugate of claim 36 for reducing cancer cell growth.

42. Use of the antibody of any one of claims 1 to 33 or the immunoconjugate of claim 36 in the manufacture of a medicament.

43. The use of claim 42, wherein the medicament is for treating cancer.

44. The use of claim 42, wherein the medicament is for reducing cancer cell growth.

45. The use of claim 42, wherein the medicament is for treating allergy, asthma, autoimmune disease, diseases associated with goblet cell metaplasia (e.g., in lung) and/or excess mucus.

46. The use of claim 45, wherein the medicament is for treating a disease associated with goblet cell metaplasia.

47. The use of claim 46, wherein the medicament is for treating asthma, cystic fibrosis, chronic obstructive pulmonary disease, or Barrett's esophagus.

48. A method of treating cancer comprising administering to an individual with cancer an effective amount of the antibody of any one of claims 1 to 33 or the immunoconjugate of claim 36.

49. The method of claim 48, wherein the cancer is selected from breast cancer, lung cancer, brain cancer, cervical cancer, colon cancer, liver cancer, bile duct cancer, pancreatic cancer, skin cancer, B-cell malignancies, and T-cell malignancies.

50. A method of treating a disease selected from allergy, asthma, autoimmune disease, diseases associated with goblet cell metaplasia (e.g., in lung) and excess mucus, comprising administering to an individual with cancer an effective amount of the antibody of any one of claims 1 to 33 or the immunoconjugate of claim 36.

51. The method of claim 50, wherein the disease is associated with goblet cell metaplasia.

52. The method of claim 51, wherein the disease is selected from asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis, and Barrett's esophagus.

53. The isolated antibody of any one of claims 1 to 33 conjugated to a label.

54. The isolated antibody of claim 53, wherein the label is a positron emitter.

55. The isolated antibody of claim 54, wherein the positron emitter is ^{89}Zr.

56. A method of detecting human Jagged1 in a biological sample comprising contacting the biological sample with the antibody of any one of claims 1 to 33 and 53 to 55 under conditions permissive for binding of the antibody to a naturally occurring human

Jagged1, and detecting whether a complex is formed between the antibody and a naturally occurring human Jagged1 in the biological sample.

57. The method of claim 56, wherein the biological sample is selected from breast cancer, lung cancer, brain cancer, cervical cancer, colon cancer, liver cancer, bile duct cancer, pancreatic cancer, skin cancer, B-cell malignancies, and T-cell malignancies.

41. The antibody of any one of claims 1 to 33 or the immunoconjugate of claim 36 for reducing cancer cell growth.

42. Use of the antibody of any one of claims 1 to 33 or the immunoconjugate of claim 36 in the manufacture of a medicament.

43. The use of claim 42, wherein the medicament is for treating cancer.

44. The use of claim 42, wherein the medicament is for reducing cancer cell growth.

45. The use of claim 42, wherein the medicament is for treating allergy, asthma, autoimmune disease, diseases associated with goblet cell metaplasia (e.g., in lung) and/or excess mucus.

46. The use of claim 45, wherein the medicament is for treating a disease associated with goblet cell metaplasia.

47. The use of claim 46, wherein the medicament is for treating asthma, cystic fibrosis, chronic obstructive pulmonary disease, or Barrett's esophagus.

48. The use of claim 43, wherein the cancer is selected from breast cancer, lung cancer, brain cancer, cervical cancer, colon cancer, liver cancer, bile duct cancer, pancreatic cancer, skin cancer, B-cell malignancies, and T-cell malignancies.

49. The isolated antibody of any one of claims 1 to 33 conjugated to a label.

50. The isolated antibody of claim 49, wherein the label is a positron emitter.

51. The isolated antibody of claim 50, wherein the positron emitter is ^{89}Zr.

52. A method of detecting human Jagged1 in a biological sample comprising contacting the biological sample with the antibody of any one of claims 1 to 33 and 49 to 51 under conditions permissive for binding of the antibody to a naturally occurring human Jagged1, and detecting whether a complex is formed between the antibody and a naturally occurring human Jagged1 in the biological sample.

53. The method of claim 52, wherein the biological sample is selected from breast cancer, lung cancer, brain cancer, cervical cancer, colon cancer, liver cancer, bile duct cancer, pancreatic cancer, skin cancer, B-cell malignancies, and T-cell malignancies.