

PYRIMIDOOXAZOCINE DERIVATIVES AS MTOR - INHIBITORS

The present invention relates to bicyclic heterocyclic derivatives, and to the
5 preparation and therapeutic use thereof.

mTOR (mammalian target of rapamycin), also known as FRAP (FKBP12 and rapamycin associated protein), is a 289-kDa serine/threonine kinase of the family of PIKKs (phosphoinositide 3-kinase-like kinases), although mTOR does not phosphorylate phospholipids.

10 The protein contains several domains, including a C-terminal kinase domain, an FKBP12-rapamycin binding domain, 20 N-terminal HEAT repetitions involved in protein-protein interactions, a FAT (FRAP-ATM-TRRAP) domain and a C-terminal FAT domain, which is also present in other PIKKs (Wullschleger et al. (2006) Cell, 124, 471-484).

The kinase mTOR is a central regulator of cell growth and proliferation, but also
15 plays an important role in cell metabolism and angiogenesis. mTOR is activated by the PI3K/Akt axis and in turn phosphorylates the downstream effectors of the PI3K/Akt signalling pathway, in particular the ribosomal protein S6 kinase (S6K1) and the eukaryotic initiation factor 4E binding protein (4E-BP1), which are two main regulators of the cell protein translation machinery (the mTOR signalling pathway is described in
20 Zoncu et al. (2011) Nature Rev. Mol. Cell Biol. 12, 21-35).

The mTOR signalling pathway is mutated and deregulated in a variety of human cancers. Mutations of the protein kinase Akt, of the lipid kinase PI3K and/or inactivation of the tumour suppressants PTEN and TSC2, but also amplifications and/or mutations affecting the growth factor receptors, are a few of the events upstream of mTOR that
25 lead to constitutive inactivation of the PI3K/Akt/mTOR pathway and uncontrolled cell proliferation (for a review of the role of the protein mTOR in cancer, see Guertin and Sabatini (2007) Cancer Cell 12, 9-22).

Genetic mutations and amplifications affecting the mTOR pathway have been identified in the formation of glioblastomas, prostate cancers, tuberous sclerosis, lung
30 cancers (NSCLC), breast cancer, ovarian cancer, endometrial cancer, bowel cancer, pancreatic cancer, cancer of the head and neck, skin cancer and hepatocellular carcinomas (Yuan and Cantley (2008) Oncogene 27, 5497-5510; Whittaker et al. (2010) Oncogene 29, 4989-5005).

mTOR recruits several partners to form two multi-protein complexes that are
35 essential for tumour growth. By phosphorylating the proteins S6K and 4E-BP1, the mTORC1 complex makes the connection between oncogene signalling and protein synthesis, glycolysis and lipid biosynthesis (Yecies and Manning (2011) J. Mol. Med. 89,

221-228). The mTORC2 complex has recently been identified as the kinase that phosphorylates Akt on the Ser-473 residue, thus acting as an essential activator of the kinase Akt. The role of the complex mTORC2 has recently been specifically linked to cell transformation (Sparks and Guertin (2010) *Oncogene* 29, 3733-3744).

5 Rapamycin and analogues thereof, the rapalogues, via their association with the protein FKBP12, are allosteric inhibitors of the mTORC1 complex. Several of them are in clinical development for the treatment of certain cancers. Everolimus (Novartis) and Temsirolimus (Wyeth) have recently been approved for the treatment of kidney cancer (renal cell carcinoma or RCC). However, the efficacy of the rapalogues in cancer
10 treatment has remained below expectations, despite certain promising results, and appears to be limited to a subset of cancers. These limitations have been attributed to the fact that rapalogues do not interact with the mTORC2 complex, and that certain aspects of the activity of the mTORC1 complex, and in particular the phosphorylating of 4E-BP1, are resistant to rapamycin and analogues thereof (Benjamin et al. (2011) *Nature*
15 *Reviews Drug Discovery* 10, 868-880).

On the other hand, inhibitors of the kinase site of mTOR do not suffer from this drawback (Feldman et al. (2009) *PLoS Biology* 7, 371-383) and are considered as a novel generation of modulators of the mTOR pathway with, as advantages, a potential increase in antitumour efficacy and a broadened field of therapeutic applications. Several
20 of them have now entered into clinical phases (Garcia-Echeverria (2011) *Biochem. Soc. Trans.* 39, 451-455; Richard et al. (2010) *Curr. Drug Opinion Disc. Dev.* 13, 428-440).


Other potential therapeutic indications have been proposed for mTOR inhibitors (Tsang et al. (2007) *Drug Discovery Today* 12, 112-124). mTOR inhibitors may have a neuroprotective role in neurodegenerative diseases such as Alzheimer's disease,
25 Parkinson's disease and Huntington's disease (Bové et al. (2011) *Nature Reviews Neuroscience* 12, 437-452). Moreover, mTOR hyperactivity has been correlated with age-related pathologies (Harrison et al. (2009) *Nature* 460, 392). Other applications include kidney, lung and liver fibrosis (Mehrad et al. (2009) *Int. J. Biochem. Cell Biol.* 41, 1708-1718; Lieberthal and Levine (2009); Shouval (2011)), inflammation and
30 autoimmune disorders (Bhonde et al. (2008) *Am. J. Physiol. Gastrointest. Liver Physiol.* 295, G1237-G1245; Young and Nickerson-Nutter (2005) *Curr. Opinion Pharmacol.* 5, 418-423), diabetes, obesity (Dann et al. (2007) *Trends Mol. Medicine* 13, 252-259), Pompe's disease (Ashe et al. (2010) *Molecular Genetics and Metabolism* 100, 309-315), cardiovascular diseases (Hagenmueller et al. (2010). *FEBS Lett.* 584, 74-80), ocular
35 diseases (Blumenkranz (2007) *Retina Today* 24-26), epilepsy (Huang et al. (2010) *Neurobiology of Disease* 40, 193-199), and parasitic diseases (Wang et al. (2010) 240th ACS National Meeting Boston, MEDI-152).

mTOR inhibition also activates autophagocytosis and a certain number of diseases should be sensitive to its inhibition, in particular proteinopathies, bacterial and viral infections and cancer (review in Rubinsztein et al. (2007) Nature 6, 304-312).

The development of compounds targeting the mTOR kinase site and inhibiting the activity of the two compounds mTORC1 and mTORC2 is thus important.

[COMPOUNDS]

According to a first aspect, a subject of the present invention is the compounds corresponding to formula (I):



(I)

10 in which:

- each of the radicals R_1 independently represents a group (C₁-C₆)-alkyl, it being understood that two groups R_1 borne by two different carbon atoms of the morpholinyl ring may be linked together to form, with the morpholinyl ring, a bicyclic heterocyclic structure,

15 - R_2 and R_3 independently represent:

- H, or
- a group (C₁-C₆)-alkyl,

- R_4 represents:

- H,

20 - a group (C₁-C₆)-alkyl, optionally substituted with a hydroxyl group,

- a group (C₁-C₆)-alkoxy,

- a group -L₁-R₁₀, in which:

- L₁ represents a group (C₁-C₆)-alkylene,

- R₁₀ represents a group chosen from -COOR₁₁, -CO-R₁₂, -OR₁₃,

25 -CONR₁₄R₁₅ and a 5- or 6-atom heterocycle optionally substituted with a group -R₁₆, -COOR₁₇, -CO-R₁₈, -OR₁₉ or -NR₂₀R₂₁, or

- a 5- or 6-atom heterocycle optionally substituted with a group $-R_{22}$, $-COOR_{23}$, $-CO-R_{24}$, $-OR_{25}$ or $-CONR_{26}R_{27}$,
- G_1 represents a divalent phenyl or 5- to 6-atom heteroaryl radical, optionally substituted with 1 to 4 groups R_5 independently chosen from a halogen atom, a group $-OR_{30}$ and a group $-(C_1-C_6)$ -alkyl optionally substituted with a hydroxyl,
- X_1 represents a group $-O-$ or $-NR_{40}-$,
- X_2 represents a single bond or a group $-CONR_{50}-$, $-CONR_{51}-O-$, $-COO-$, $-CO-$ or $-SO_2-$,
- R_6 represents:
 - H,
 - a group $-L_2-R_7$, in which:
 - $-L_2-$ represents a group (C_1-C_6) -alkylene or (C_3-C_6) -cycloalkylene,
 - R_7 represents a group chosen from:
 - H,
 - OR_{60} ,
 - a halogen atom,
 - a group (C_1-C_6) -haloalkyl,
 - a 5- to 6-atom heterocycle optionally substituted with one or more groups chosen from a group (C_1-C_6) -alkyl, a group (C_1-C_6) -alkoxy and a group $=O$, or
 - a group $-G_2-X_3-G_3$, in which:
 - X_3 represents a single bond or a group $-O-$, $-CO-$ or $-CH_2-$,
 - G_2 represents a single bond or a 5- to 6-atom divalent cyclic radical optionally substituted with one or more groups R_{80} ,
 - G_3 represents a 4- to 8-atom ring optionally substituted with one or more groups R_{81} ,
 - or $-G_2-X_3-G_3$ together form a 7- to 10-atom fused bicycle,
 - each radical R_{80} and each radical R_{81} is independently chosen from:
 - a halogen atom,
 - a group $-COOR_{70}$, $-CO-R_{71}$, $-OR_{72}$, $-NR_{73}R_{74}$, $-CONR_{75}R_{76}$, $-CN$ or $-S(O)_p-R_{77}$,
 - a group (C_1-C_6) -alkyl optionally substituted with one or more groups R_{100} and/or optionally interrupted with one or more oxygen atoms,
 - a group (C_1-C_6) -alkoxy optionally substituted with one or more groups R_{101} and/or optionally interrupted with one or more oxygen atoms,
- n represents 0, 1 or 2,

- m represents 0 or 1,
 - p represents 0, 1 or 2,
 - R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, R₂₄, R₂₅, R₂₆, R₂₇, R₃₀, R₄₀, R₅₀, R₅₁, R₆₀, R₇₀, R₇₁, R₇₂, R₇₃, R₇₄, R₇₅, R₇₆ and R₇₇ independently represent H or a group (C₁-C₆)-alkyl optionally substituted with a group R₉₀ and/or optionally interrupted with one or more oxygen atoms,
 - R₉₀ represents a group –OR₉₁ or –NR₉₂R₉₃,
 - R₉₁, R₉₂ and R₉₃ independently represent H or a group (C₁-C₆)-alkyl,
 - R₁₀₀ and R₁₀₁ are independently chosen from a halogen atom and a hydroxyl
- 10 in the form of the base or of an acid-addition salt.

The compounds of formula (I) can comprise one or more asymmetric carbon atoms. They may therefore exist in the form of enantiomers or diastereoisomers. These enantiomers, diastereoisomers, and also mixtures thereof, including racemic mixtures, form part of the invention.

- 15 The pure enantiomers of the compounds of the invention may be obtained from enantiomerically pure precursors or alternatively by chromatography on chiral phases or alternatively, when the compounds comprise acid or amine functions, by selective crystallization of diastereoisomeric salts obtained by reacting the compounds (I) with, respectively, chiral amines or acids.

- 20 The compounds of formula (I) may exist in the form of tautomers. These tautomeric forms form part of the invention.

By virtue of their structure, the compounds of general formula (I) may also exist in the form of enantiomers of rotamer or atropoisomer type.

- 25 The compounds of formula (I) may exist in the form of bases or acid-addition salts. Such addition salts form part of the invention.

These salts may be advantageously prepared with pharmaceutically acceptable acids, but salts of other acids that are of use, for example, for purifying or isolating the compounds of formula (I) also form part of the invention.

In the context of the present invention, the following definitions apply:

- 30 - a halogen atom: a fluorine atom, a chlorine atom, a bromine atom or an iodine atom, especially a fluorine atom;
- an alkyl group: a saturated, linear or branched hydrocarbon-based aliphatic group


comprising, unless otherwise mentioned, from 1 to 6, especially from 1 to 4 and in particular from 1 to 2 carbon atoms, or a cyclic hydrocarbon-based aliphatic group, also be noted as cycloalkyl and defined hereinbelow. Examples that may be mentioned include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, etc. groups;

- a cycloalkyl group: a cyclic alkyl group comprising, unless otherwise mentioned, from 3 to 6 carbon atoms. Examples that may be mentioned include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc. groups;
- an alkylene group: a linear or branched saturated hydrocarbon-based divalent aliphatic group, comprising, unless otherwise mentioned, from 1 to 6 carbon atoms;
- a cycloalkylene group: a cyclic or carbocyclic hydrocarbon-based divalent aliphatic group, comprising, unless otherwise mentioned, from 3 to 6 carbon atoms;
- an alkoxy group: a radical -O-alkyl in which the alkyl group is as defined previously. Examples that may be mentioned include the groups -O-(C₁-C₆)alkyl or -(C₁-C₆)-alkoxy, and in particular as (i) group -O-C₁alkyl, the group -O-methyl, as (ii) group -O-C₂alkyl, the group -O-ethyl, as (iii) group -O-C₃alkyl, the group -O-propyl, -O-isopropyl, as (iv) group -O-C₄alkyl, the group -O-butyl, -O-isobutyl, -O-tert-butyl, as (v) group -O-C₅alkyl, the group -O-pentyl, -O-isopentyl; as (vi) group -O-C₆alkyl, the group -O-hexyl;
- a ring: a saturated, unsaturated or aromatic (aryl or heteroaryl) carbocycle or heterocycle;
- a carbocycle: a ring composed of carbon atoms, which is saturated (the saturated carbocycles especially being a cycloalkyl, such as a cyclopropyl, a cyclopentyl, cyclohexyl or an adamantyl (example of a bridge tricycle)), unsaturated (for example a cyclohexene) or aromatic (i.e. an aryl). When the carbocycle comprises several substituents, they may be borne by the same atom or different atoms,
- an aryl: a monocyclic or bicyclic aromatic hydrocarbon-based group comprising, unless otherwise mentioned, between 5 and 10 carbon atoms. Examples of aryl groups that may be mentioned include phenyl and naphthyl, in particular phenyl,
- a heterocycle: a monocyclic or bicyclic group comprising, unless otherwise mentioned, from 3 to 10 atoms, and comprising one or more heteroatoms chosen from O, N and/or S. The said heterocycle may be saturated or partially unsaturated and may comprise one or more double bonds. It is then referred to as a heterocycloalkyl group. It may also be aromatic, comprising, unless otherwise mentioned, from 5 to 10 atoms and then represent a heteroaryl group. When the heterocycle is substituted, the substitution(s) may be on one (or more) carbon atom(s) and/or on the heteroatom(s). When the heterocycle comprises several

substituents, they may be borne by the same atom or different atoms.


Non-aromatic heterocycles or heterocycloalkyls that may be mentioned include epoxyethyl, oxiranyl, aziridinyl, tetrahydrofuryl, dioxolanyl, pyrrolidinyl, pyrazolidinyl, imidazolidinyl, tetrahydrothiophenyl, dithiolanyl, thiazolidinyl, 5 tetrahydropyranyl, dioxanyl, morpholinyl, piperidyl, piperazinyl, tetrahydrothiopyranyl, dithianyl, thiomorpholinyl, dihydrofuryl, 2-imidazolinyl, 2,-3-pyrrolinyl, pyrazolinyl, dihydrothiophenyl, dihydropyranyl, pyranal, tetrahydropyridyl, dihydropyridyl, tetrahydropyridinyl, dihydrothiopyranyl, isoxazolidinyl and the corresponding groups derived from fusion with a phenyl 10 nucleus, and more particularly morpholinyl, dioxalanyl, benzothiazolidinyl, pyrrolidinyl and benzopyrrolidinyl rings.

Bridged heterocycles that may be mentioned include rings of bridged morpholinyl type such as:



15 Heteroaryls that may especially be mentioned include the following representative groups: benzimidazolyl, benzothiazolyl, furyl, furazanyl, imidazolyl, indolyl, indoliziny, isoxazolyl, isoquinolinyl, isothiazolyl, oxadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, pyrazinyl, pyridazinyl, pyrazolyl, pyridyl, pyrimidinyl, pyrrolyl, quinazolinyl, quinolinyl, 1,3,4-thiadiazolyl, 1,2,3-thiadiazolyl, thiazolyl, 20 thienyl, triazolyl, 1,2,3-triazolyl and 1,2,4-triazolyl;

- a fused bicycle comprises two rings as defined above, the fusion of the two rings possibly taking place by the sharing of a sequence of atoms (branch and bridgehead) (for example norbornane) or by sharing a bond between two atoms (for example isoquinoline or quinoline);
- 25 - the ring of a cyclic divalent radical is as defined above. The ring of the cyclic divalent radical may be a carbocycle or a heterocycle and it may be saturated, unsaturated or aromatic, especially aromatic. In particular, the ring of a cyclic divalent radical may be an aryl, such as a phenyl, or a heteroaryl, in which the heteroatom(s) are especially nitrogen atoms. An example of a cyclic divalent radical of 5 atoms that may be mentioned is a divalent triazole radical, and examples of cyclic divalent radicals of 6 30 atoms that may be mentioned are divalent radicals of phenyl (phenylene), of pyridine and of pyrazine. For example, a cyclic divalent radical of 5 or 6 atoms may have one of the following formulae, it being understood that the atoms of the rings may be substituted (with one or more groups R_{80}):




In one embodiment, a subject of the present invention is the compounds corresponding to formula (I) in which:

- 5 - each of the radicals R_1 independently represents a group (C_1-C_6) -alkyl, advantageously a group (C_1-C_3) -alkyl, it being understood that two groups R_1 borne by two different carbon atoms of the morpholinyl ring may be linked together to form, with the morpholinyl ring, a bicyclic heterocyclic structure,
- R_2 and R_3 independently represent H or a group (C_1-C_2) -alkyl,
- 10 - R_4 represents:
 - H,
 - a group (C_1-C_6) -alkyl, advantageously a group (C_1-C_3) -alkyl optionally substituted with a hydroxyl group,
 - a group $-L_1-R_{10}$, in which:
 - 15 - L_1 represents a group (C_1-C_2) -alkylene,
 - R_{10} represents a group chosen from $-COOR_{11}$, $-OR_{13}$, $-CONR_{14}R_{15}$ or a heterocycle of 5 atoms optionally substituted with a group $-COOR_{17}$, $-CO-R_{18}$, or
 - a heterocycle of 6 atoms optionally substituted with a group $-R_{22}$,
- 20 - G_1 represents a divalent phenyl or 5- to 6-atom heteroaryl radical, optionally substituted with 1 to 4 groups R_5 independently chosen from a halogen atom and a group $-(C_1-C_6)$ -alkyl optionally substituted with a hydroxyl,
- X_1 represents a group $-O-$ or $-NR_{40}-$,
- X_2 represents a single bond or a group $-CONR_{50}-$, $-CONR_{51}-O-$, $-COO-$, $-CO-$ or $-$
- 25 SO_2- ,
- R_6 represents:
 - H,
 - a group $-L_2-R_7$, in which:
 - $-L_2-$ represents a group (C_1-C_4) -alkylene or (C_3-C_6) -cycloalkylene,
 - 30 - $-R_7$ represents a group chosen from:
 - H,
 - OR_{60} ,
 - a halogen atom,


- a 5- to 6-atom heterocycle optionally substituted with one or more groups chosen from a group (C₁-C₆)-alkyl and a group =O, or
- a group -G₂-X₃-G₃, in which:
 - X₃ represents a single bond or a group -O-, -CO- or -CH₂-,
 - 5 - G₂ represents a single bond or a cyclic divalent radical of 5 to 6 atoms,
 - G₃ represents a 4- to 8-atom ring optionally substituted with one or more groups R₈₁,
 - or -G₂-X₃-G₃ together form a 7- to 10-atom fused bicycle,
- 10 - each radical R₈₁ is independently chosen from:
 - a halogen atom,
 - a group -COOR₇₀, -OR₇₂, -NR₇₃R₇₄, -CONR₇₅R₇₆, -CN or -S(O)_p-R₇₇,
 - a group (C₁-C₂)-alkyl optionally substituted with one or more groups R₁₀₀,
 - a group (C₁-C₂)-alkoxy optionally substituted with one or more groups R₁₀₁,
- 15 - n represents 0, 1 or 2,
- m represents 0 or 1,
- p represents 2,
- R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, R₂₄, R₂₅, R₂₆, R₂₇, R₃₀, R₄₀,
- 20 R₅₀, R₅₁, R₆₀, R₇₀, R₇₁, R₇₂, R₇₃, R₇₄, R₇₅, R₇₆ and R₇₇ independently represent H or a group (C₁-C₂)-alkyl optionally substituted with a group R₉₀,
- R₉₀ represents a group -OR₉₁ or -NR₉₂R₉₃,
- R₉₁, R₉₂ and R₉₃ represent a group (C₁-C₂)-alkyl,
- R₁₀₀ and R₁₀₁ represent a halogen atom,
- 25 in the form of the base or of an acid-addition salt.

In one embodiment, in formula (I) above, m represents 0 and R₂ and R₃ represent a methyl, the compounds then having the formula (II) below:



in which n , R_1 , R_4 , G_1 , X_1 , X_2 and R_6 are as defined above, in the form of the base or of an acid-addition salt.

In one embodiment, in formula (II) above, G_1 represents an optionally substituted phenyl group, X_1 represents $-\text{NH}-$ and the compounds have the formula (III) below:



5


in which:

- R_4 represents a group $(\text{C}_1\text{-C}_6)\text{-alkyl}$, in particular an ethyl group or a methyl group,
- R'_5 and R''_5 represent $-\text{H}$ or $-\text{F}$, it being understood that when one of the radicals R'_5 and R''_5 represents $-\text{F}$, then the other represents $-\text{H}$,
- 10 - X_2 represents $-\text{CONR}_{50}-$ or $-\text{COO}-$, especially $-\text{CONH}-$ or $-\text{COO}-$, in particular $-\text{CONH}-$,
- R_6 represents:
 - a group $-\text{L}_2\text{-R}_7$, in which:
 - $-\text{L}_2-$ represents a group $(\text{C}_1\text{-C}_6)\text{-alkylene}$ or $(\text{C}_3\text{-C}_6)\text{-cycloalkylene}$,
 - 15 - R_7 represents a group chosen from:
 - H , or
 - OR_{60} ,
 - a group $-\text{G}_2\text{-X}_3\text{-G}_3$, in which:
 - X_3 represents a single bond or a group $-\text{O}-$,
 - 20 - G_2 represents a single bond or a cyclic divalent radical of 6 atoms,
 - G_3 represents a 4- to 8-atom ring optionally substituted with one or more (especially 1 or 2) groups R_{81} ,
- n , R_1 , R_{50} , R_{60} and R_{81} are as defined above, in the form of the base or of an acid-addition salt.


In the abovementioned formulae, the following embodiments may be used and independently combined with each other:

- m represents 0,

5 - the group



is especially chosen from the following groups:



10 - R_2 and R_3 represent a methyl group,

- R_4 represents:

- H,

- a group (C_1-C_6) -alkyl, (especially (C_1-C_2) -alkyl), optionally substituted with a hydroxyl group,

15 - a group $-L_1-R_{10}$, in which:

- L_1 represents a group (C_1-C_6) -alkylene,

- R_{10} represents a group chosen from $-COOR_{11}$, $-OR_{13}$, $-CONR_{14}R_{15}$ and a 5- or 6-atom heterocycle optionally substituted with a group $-COOR_{17}$ or $-CO-R_{18}$, or

- a 5- or 6-atom heterocycle optionally substituted with a group $-R_{22}$,

20 - R_4 especially represents a group (C_1-C_6) -alkyl, such as a methyl group or an ethyl group,

- G_1 represents a divalent phenyl or pyridyl radical optionally substituted with a group R_5 chosen from halogen atoms, especially fluorine,

25 - G_1 represents a divalent pyridyl radical or a divalent phenyl radical optionally substituted with a group R_5 chosen from halogen atoms, especially fluorine,

- G_1 especially represents a divalent phenyl radical optionally substituted with a fluorine group,

30 - when X_1 represents O, X_2 represents a single bond and R_6 represents a group (C_1-C_6) -alkyl (i.e. a group $-L_2-R_7$ in which $-L_2-$ represents a (C_1-C_6) -alkylene and R_7 represents $-H$),

- X_1 represents a group $-NR_{40}-$, especially $-NH-$,
 - X_2 represents a single bond or a group $-CONH-$, $-COO-$, $-CO-$ or $-SO_2-$, especially $-CONH-$ or $-COO-$,
 - X_2 represents $-CONR_{50}-$ or $-COO-$, especially $-CONH-$ or $-COO-$,
- 5 - When R_7 represents a 5- to 6-atom heterocycle optionally substituted with one or more groups $=O$, the said heterocycle is typically a heterocycle bearing an oxidized sulfur atom, such as tetramethylene sulfone,
- when $-G_2-X_3-G_3$ together form a 7- to 10-atom fused bicycle, the said ring is a benzimidazolyl or a 7-azaindolyl,
- 10 - G_2 is a 6-atom cyclic divalent radical,
- G_2 is a 6-atom cyclic divalent radical and the groups X_2 and X_3 are in the para position on this ring,
 - G_2 is optionally substituted with a group R_{80} ,
 - G_3 is optionally substituted with 1, 2 or 3, especially 1 or 2, groups R_{80} ,
- 15 - G_2 is an aromatic ring,
- G_2 is an aromatic ring and G_3 is a saturated ring,
 - each radical R_{80} and each radical R_{81} is independently chosen from:
 - a halogen atom, especially fluorine,
 - a group $-COOR_{70}$, $-OR_{72}$, $-NR_{73}R_{74}$, $-CONR_{75}R_{76}$, $-CN$ or $-S(O)_p-R_{77}$,
- 20 - a group (C_1-C_6) -alkyl optionally substituted with one or more groups R_{100} ,
- a group (C_1-C_6) -alkoxy optionally substituted with one or more groups R_{101} and/or optionally interrupted with one or more oxygen atoms,
- in the form of the base or of an acid-addition salt.

Among the compounds of formula (I) that are subjects of the invention, mention
 25 may be made especially of the compounds in Table 1 below:

1	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea
2	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea
3	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea

4	1-{4-[8,8-Dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(S)-1-pyrrolidin-2-ylmethyl-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-ethylurea
5	(S)-2-[4-[4-(3-Ethylureido)-3-fluorophenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-ylmethyl]pyrrolidine-1-carboxylic acid ethyl ester
6	1-{4-[6-((S)-1-Acetylpyrrolidin-2-ylmethyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-ethylurea
7	3-[4-[4-(3-Ethylureido)phenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl]propionic acid
8	3-[4-[4-(3-Ethylureido)phenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl]-N,N-dimethylpropionamide
9	1-Ethyl-3-{4-[6-(3-hydroxypropyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
10	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6,7,8,9-tetrahydro-5H-10-oxa-1,3,6-triazabenzocycloocten-4-yl]phenyl}urea
11	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-methylurea
12	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-pyridin-4-ylurea
13	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[4-(4-methylpiperazin-1-yl)phenyl]urea
14	1-{4-[8,8-Dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-ethylurea
15	1-{4-[8,8-Dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methylisoxazol-3-yl)urea
16	1-{4-[8,8-Dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(1-methyl-1H-pyrazol-3-yl)urea

17	1-Ethyl-3-{5-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]pyridin-2-yl}urea
18	1-(3-Difluoromethoxyphenyl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
19	1-(2,4-Difluorophenyl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
20	{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid (S)-2-hydroxypropyl ester
21	1-(4-Methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
22	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(1-methylpiperidin-4-yloxy)pyridin-3-yl]urea
23	1-[4-(4-Dimethylaminopiperidine-1-carbonyl)phenyl]-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
24	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-hydroxy-2-methylpropyl)urea
25	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-pyrrolidin-1-ylpyridin-2-yl)urea
26	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-methylphenyl}-3-isoxazol-3-ylurea
27	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-((R)-2-hydroxy-1-methylethyl)urea
28	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methoxypyrazin-2-yl)urea

29	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((S)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
30	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((R)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
31	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)urea
32	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-pyrrolidin-1-ylpyridin-3-yl)urea
33	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-2-ylurea
34	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methylpyridin-2-yl)urea
35	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(4-methoxypyridin-2-yl)urea
36	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methylpyridin-2-yl)urea
37	1-{4-[6-Ethyl-2-((S)-3-ethylmorpholin-4-yl)-8,8-dimethyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methylpyridin-3-yl)urea
38	1-Methyl-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
39	1-Pyridin-2-yl-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
40	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[5-(2-methoxyethoxy)pyridin-2-yl]urea
41	1-Pyridin-2-yl-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea

42	1-(5-Methylpyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
43	1-Methyl-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
44	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methoxypyridin-2-yl)urea
45	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-methylurea
46	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-methylurea
47	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-methylurea
48	1-(5-Methylpyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
49	1-(5-Methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
50	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-3-ylurea
51	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-4-ylurea
52	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-4-ylurea
53	{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid (1S,2R)-2-hydroxycyclopentyl ester
54	1-(5-Methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
55	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-methylpyridin-4-yl)urea

56	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-methylpyridin-4-yl)urea
57	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-((S)-2-hydroxy-1-methylethyl)urea
58	1-(2,6-Dimethylpyridin-4-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
59	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-methylpyridin-4-yl)urea
60	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-methylpyridin-4-yl)urea
61	{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid (S)-2-hydroxy-1-methylethyl ester
62	{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid (R)-2-hydroxy-1-methylethyl ester
63	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-2-ylurea
64	1-(5-Dimethylaminopyridin-2-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
65	{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid 2-hydroxyethyl ester
66	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methylpyridin-3-yl)urea
67	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methylpyridin-3-yl)urea

68	1-Pyridin-4-yl-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
69	1-Pyridin-4-yl-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
70	1-(4-Methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
71	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-pyridin-4-ylurea
72	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[5-(3-hydroxypyrrolidin-1-yl)pyridin-2-yl]urea
73	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-((S)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
74	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-((S)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
75	1-[4-((1S,5R)-6-Ethyl-8,8-dimethyl-2-3-oxa-8-azabicyclo[3.2.1]oct-8-yl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)phenyl]-3-(3-fluoropyridin-4-yl)urea
76	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(3-fluoropyridin-4-yl)urea
77	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[4-(tetrahydrofuran-3-yloxy)phenyl]urea
78	1-(2-Dimethylaminopyridin-4-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
79	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(4-methoxypyridin-2-yl)urea
80	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[2-(4-methanesulfonylpiperazin-1-yl)pyridin-4-yl]urea

81	4-[5-(3-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}ureido)pyridin-2-yl]piperazine-1-carboxylic acid methyl ester
82	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-4-ylurea
83	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-4-ylurea
84	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-pyridin-2-ylurea
85	1-(3-Fluoropyridin-4-yl)-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
86	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(1H-pyrrolo[2,3-b]pyridin-4-yl)urea
87	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-((R)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
88	{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid (S)-2-hydroxypropyl ester
89	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-(4-methoxypyridin-2-yl)urea
90	1-(2,6-Dimethylpyrimidin-4-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
91	1-(2,6-Dimethylpyrimidin-4-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
92	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyrimidin-4-ylurea
93	1-(2,6-Dimethylpyrimidin-4-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea

94	1-[4-(8,8-Dimethyl-2-morpholin-4-yl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)phenyl]-3-ethylurea
95	4-(6-Aminopyridin-3-yl)-8,8-dimethyl-2-morpholin-4-yl-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one
96	1-Ethyl-3-[4-(6-ethyl-8,8-dimethyl-2-morpholin-4-yl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)phenyl]urea
97	1-Ethyl-3-[4-(6-ethyl-2-morpholin-4-yl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)phenyl]urea
98	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
99	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
100	6-Ethyl-4-(3-hydroxymethyl-4-methoxyphenyl)-8,8-dimethyl-2-morpholin-4-yl-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one
101	6-Ethyl-4-(3-hydroxymethyl-4-methoxyphenyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one
102	4-(4-Aminophenyl)-6-ethyl-8,8-dimethyl-2-morpholin-4-yl-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one
103	[4-(6-Ethyl-8,8-dimethyl-2-morpholin-4-yl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)phenyl]carbamic acid methyl ester
104	1-Methoxy-3-[4-(6-ethyl-8,8-dimethyl-2-morpholin-4-yl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)phenyl]urea
105	{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid methyl ester
106	{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid ethyl ester
107	4-(6-Aminopyridin-3-yl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one
108	N-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}propionamide
109	1-{4-[2-((2S,6R)-2,6-Dimethylmorpholin-4-yl)-6-ethyl-8,8-dimethyl-5-oxo-6,7,8,9-tetrahydro-5H-10-oxa-1,3,6-triazabenzocycloocten-4-yl]phenyl}-3-ethylurea

110	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
111	Ethanesulfonic acid {4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}amide
112	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-3-ylurea
113	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
114	1-{4-[2-((2S,6R)-2,6-Dimethylmorpholin-4-yl)-6-ethyl-8,8-dimethyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-ethylurea
115	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-methyl-2H-pyrazol-3-yl)urea
116	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methylisoxazol-3-yl)urea
117	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-hydroxyethyl)urea
118	4-[4-(1H-Benzimidazol-2-ylamino)phenyl]-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one
119	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(3-hydroxypropyl)urea
120	3-[4-[4-(3-Ethylureido)phenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl]propionic acid tert-butyl ester
121	6-Ethyl-8,8-dimethyl-4-[4-(5-methyl-[1,3,4]thiadiazol-2-ylamino)phenyl]-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one
122	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(1-methyl-1H-pyrazol-3-yl)urea
123	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-fluoroethyl)urea

124	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridin-3-yl)urea
125	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyrazin-2-ylurea
126	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(1-methyl-1H-pyrazol-4-yl)urea
127	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(3-methoxyphenyl)urea
128	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(tetrahydrofuran-2-ylmethyl)urea
129	1-(1,1-Dioxotetrahydro-1λ6-thiophen-3-ylmethyl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
130	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(1-methylpiperidin-4-ylmethyl)urea
131	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-methoxyethyl)urea
132	1-{4-[8,8-Dimethyl-6-(1-methylpiperidin-4-yl)-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-ethylurea
133	1-Cyclopropyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
134	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(tetrahydropyran-4-ylmethyl)urea
135	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-isoxazol-3-ylurea

136	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-methoxypyrimidin-5-yl)urea
137	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-oxetan-3-ylurea
138	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(4-fluoro-3-methoxyphenyl)urea
139	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyrazin-2-yl)urea
140	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridazin-3-yl)urea
141	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridin-3-yl)urea
142	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridin-2-yl)urea
143	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(1-methyl-1H-pyrazol-4-yl)urea
144	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-methoxypyrimidin-4-yl)urea
145	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridin-3-ylmethyl)urea
146	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methoxypyridazin-3-yl)urea
147	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(3-trifluoromethylphenyl)urea

148	1-{4-[8,8-Dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methoxy-pyridin-3-yl)urea
149	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2,2,2-trifluoroethyl)urea
150	1-Cyclopropyl-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
151	1-Ethyl-3-{4-[6-ethyl-2-((S)-3-ethylmorpholin-4-yl)-8,8-dimethyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
152	6-Ethyl-4-[3-fluoro-4-(5-methyl-4H-[1,2,4]triazol-3-ylamino)phenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one
153	1-{4-[8,8-Dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-isoxazol-3-ylurea
154	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-fluoro-4-trifluoromethylphenyl)urea
155	1-(2,4-Difluoro-5-methoxyphenyl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
156	1-{4-[8,8-Dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-isoxazol-3-ylurea
157	1-{4-[8,8-Dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-isoxazol-3-ylurea
158	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-2-ylurea
159	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-isoxazol-3-ylurea

160	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyrazin-2-ylurea
161	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methylpyridin-3-yl)urea
162	1-(2,4-Difluorophenyl)-3-{4-[6-(2-methoxyethyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
163	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(5-methoxypyridin-2-yl)urea
164	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(5-methylpyridin-2-yl)urea
165	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-methylphenyl}-3-isoxazol-3-ylurea
166	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(1-methyl-1H-pyrazol-3-yl)urea
167	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(5-methylpyrazin-2-yl)urea
168	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methylpyridin-2-yl)urea
169	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methylpyridin-2-yl)urea
170	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(5-methyl-[1,3,4]oxadiazol-2-yl)urea
171	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]urea

172	1-Isoxazol-3-yl-3-{4-[6-(2-methoxyethyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
173	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methoxypyridin-2-yl)urea
174	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]urea
175	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-methylphenyl}-3-(6-methoxypyridin-3-yl)urea
176	N-(2-Dimethylaminoethyl)-4-(3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}ureido)-N-methylbenzamide
177	1-[4-(4-Dimethylaminopiperidine-1-carbonyl)phenyl]-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
178	1-{2-Fluoro-4-[6-(2-methoxyethyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methoxypyridin-3-yl)urea
179	1-{2-Fluoro-4-[6-(2-methoxyethyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-isoxazol-3-ylurea
180	1-Isoxazol-3-yl-3-{4-[6-(2-methoxyethyl)-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
181	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(4-methoxypyridin-2-yl)urea
182	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(1-methyl-1H-imidazol-4-yl)urea
183	6-Ethyl-4-{3-fluoro-4-[5-(6-methoxypyridin-3-yl)-4H-[1,2,4]triazol-3-ylamino]phenyl}-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one

184	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(4-methoxypyridin-2-yl)urea
185	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methylpyridin-3-yl)urea
186	{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid 2-hydroxyethyl ester
187	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-morpholin-4-ylpyridin-3-yl)urea
188	1-(2,4-Difluorophenyl)-3-{4-[6-(2-methoxyethyl)-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
189	1-{4-[8,8-Dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridin-3-yl)urea
190	4-(3-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}ureido)-N-(2-hydroxy-2-methylpropyl)benzamide
191	1-(2,4-Difluorophenyl)-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
192	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-2-ylurea
193	{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid 3-methoxyphenyl ester
194	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(1-methyl-1H-imidazol-4-yl)urea
195	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]urea

196	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-methylphenyl}-3-(6-methoxypyridin-3-yl)urea
197	1-{4-[6-(2-Methoxyethyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methoxypyridin-3-yl)urea
198	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methoxypyrazin-2-yl)urea
199	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-methylpyridin-4-yl)urea
200	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-morpholin-4-ylpyridin-3-yl)urea
201	1-{4-[6-(2-Methoxyethyl)-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(1-methyl-1H-pyrazol-4-yl)urea
202	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-fluorophenyl)urea
203	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(4-piperidin-1-ylmethylphenyl)urea
204	1-(3-Difluoromethoxyphenyl)-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
205	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[4-(4-methylpiperazin-1-yl)phenyl]urea
206	1-[4-((1S,4S)-6-Ethyl-8,8-dimethyl-2-2-oxa-5-azabicyclo[2.2.1]hept-5-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl]-3-methylurea
207	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea

208	1-(2,4-Difluoro-5-methoxyphenyl)-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
209	{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid 2-hydroxyethyl ester
210	1-{2-Fluoro-4-[6-(2-methoxyethyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-methylurea
211	1-Ethyl-3-{2-fluoro-4-[6-(2-methoxyethyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
212	1-(3-Difluoromethoxyphenyl)-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
213	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-fluoro-5-methoxyphenyl)urea
214	{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid 2-hydroxy-2-methylpropyl ester
215	N-(2-Dimethylamino-ethyl)-4-(3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}ureido)-N-methylbenzamide
216	1-(3-Difluoromethoxyphenyl)-3-{4-[6-(2-methoxyethyl)-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
217	1-Ethyl-3-{4-[6-(2-methoxyethyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
218	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[5-(4-methylpiperazin-1-yl)pyridin-2-yl]urea
219	4-(3-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}ureido)-N-(2-hydroxy-2-methylpropyl)benzamide

220	1-(3-Difluoromethoxyphenyl)-3-{2-fluoro-4-[6-(2-methoxyethyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
221	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-2-ylurea
222	{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid 2-hydroxyethyl ester
223	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-3-ylurea
224	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-hydroxy-1,1-dimethylethyl)urea
225	1-{2-Fluoro-4-[6-(2-methoxyethyl)-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-methylurea
226	1-{2-Fluoro-4-[6-(2-methoxyethyl)-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-hydroxy-2-methylpropyl)urea
227	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(4-morpholin-4-ylphenyl)urea
228	1-{4-[6-Ethyl-2-((S)-3-ethylmorpholin-4-yl)-8,8-dimethyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea
229	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methylpyridin-3-yl)urea
230	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-trifluoromethylpyridin-3-yl)urea
231	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-4-ylurea

232	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-morpholin-4-ylpyrazin-2-yl)urea
233	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(5-morpholin-4-ylpyrazin-2-yl)urea
234	1-(5-Dimethylaminopyrazin-2-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
235	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[4-(4-methanesulfonylpiperazin-1-yl)phenyl]urea
236	4-[4-(3-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}ureido)phenyl]piperazine-1-carboxylic acid methyl ester
237	1-[4-(1,1-Dioxo-1λ6-thiomorpholin-4-yl)phenyl]-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
238	{4-[6,8,8-Trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid (S)-2-hydroxy-1-methylethyl ester
239	{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid (R)-2-hydroxypropyl ester
240	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridin-3-yl)urea
241	1-(2,6-Dimethylpyridin-4-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
242	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-(4-methoxypyridin-2-yl)urea
243	{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}carbamic acid 2-hydroxyethyl ester

244	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-3-yl]urea
245	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)pyridin-3-yl]urea
246	{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}carbamic acid 2-hydroxyethyl ester
247	1-(6-Cyanopyridin-3-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
248	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((S)-3-ethylmorpholin-4-yl)pyridin-3-yl]urea
249	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((S)-3-ethylmorpholin-4-yl)pyridin-3-yl]urea
250	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-morpholin-4-ylpyridin-2-yl)urea
251	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[5-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-2-yl]urea
252	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[(1S,4S)-6-(2-oxa-5-azabicyclo[2.2.1]hept-5-yl)pyridin-3-yl]urea
253	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[(1S,4S)-6-(2-oxa-5-azabicyclo[2.2.1]hept-5-yl)pyridin-3-yl]urea
254	1-Cyclopropyl-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
255	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea

256	1-[6-(3,3-Dimethylmorpholin-4-yl)pyridin-3-yl]-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
257	1-[6-(3,3-Dimethylmorpholin-4-yl)pyridin-3-yl]-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
258	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-3-yl]urea
259	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-3-yl]urea
260	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-4-yl]urea
261	1-(4,4-Difluoro-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4'-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea

in the form of the base or of an acid-addition salt.

Table 1: Compounds according to the invention - IUPAC nomenclature

It should be noted that the numbers attributed to the compounds correspond to those of the examples and remain the same throughout the patent application.

- 5 Among the compounds of formula (III) that are subjects of the invention, mention may be made especially of the compounds of Table 2 below:

1	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea
2	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea
3	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
12	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-pyridin-4-ylurea

13	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[4-(4-methylpiperazin-1-yl)phenyl]urea
21	1-(4-Methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
22	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(1-methylpiperidin-4-yloxy)pyridin-3-yl]urea
29	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((S)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
30	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((R)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
31	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)urea
32	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-pyrrolidin-1-ylpyridin-3-yl)urea
37	1-{4-[6-Ethyl-2-((S)-3-ethylmorpholin-4-yl)-8,8-dimethyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methylpyridin-3-yl)urea
38	1-Methyl-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
40	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[5-(2-methoxyethoxy)pyridin-2-yl]urea
43	1-Methyl-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
45	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-methylurea
46	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-methylurea
49	1-(5-Methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea

50	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-3-ylurea
51	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-4-ylurea
52	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-4-ylurea
54	1-(5-Methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
55	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-methylpyridin-4-yl)urea
56	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-methylpyridin-4-yl)urea
58	1-(2,6-Dimethylpyridin-4-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
59	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-methylpyridin-4-yl)urea
60	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-methylpyridin-4-yl)urea
64	1-(5-Dimethylaminopyridin-2-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
66	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methylpyridin-3-yl)urea
67	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methylpyridin-3-yl)urea

68	1-Pyridin-4-yl-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
69	1-Pyridin-4-yl-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
70	1-(4-Methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
71	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-pyridin-4-ylurea
73	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-((S)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
74	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-((S)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
75	1-[4-((1S,5R)-6-Ethyl-8,8-dimethyl-2-3-oxa-8-azabicyclo[3.2.1]oct-8-yl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)phenyl]-3-(3-fluoropyridin-4-yl)urea
76	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(3-fluoropyridin-4-yl)urea
77	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[4-(tetrahydrofuran-3-yloxy)phenyl]urea
78	1-(2-Dimethylaminopyridin-4-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
79	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(4-methoxypyridin-2-yl)urea
80	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[2-(4-methanesulfonylpiperazin-1-yl)pyridin-4-yl]urea
81	4-[5-(3-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}ureido)pyridin-2-yl]piperazine-1-carboxylic acid methyl ester

82	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-4-ylurea
83	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-4-ylurea
87	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-((R)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
98	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
99	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
110	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
112	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-3-ylurea
116	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methylisoxazol-3-yl)urea
124	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridin-3-yl)urea
133	1-Cyclopropyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
135	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-isoxazol-3-ylurea
141	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridin-3-yl)urea
150	1-Cyclopropyl-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
159	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-isoxazol-3-ylurea
161	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methylpyridin-3-yl)urea

168	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methylpyridin-2-yl)urea
171	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]urea
174	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]urea
184	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(4-methoxypyridin-2-yl)urea
185	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methylpyridin-3-yl)urea
186	{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid 2-hydroxyethyl ester
187	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-morpholin-4-ylpyridin-3-yl)urea
195	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]urea
199	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-methylpyridin-4-yl)urea
200	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-morpholin-4-ylpyridin-3-yl)urea
202	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-fluorophenyl)urea
205	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[4-(4-methylpiperazin-1-yl)phenyl]urea
207	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea

218	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[5-(4-methylpiperazin-1-yl)pyridin-2-yl]urea
221	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-2-ylurea
223	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-3-ylurea
227	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(4-morpholin-4-ylphenyl)urea
229	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methylpyridin-3-yl)urea
231	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-4-ylurea
235	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[4-(4-methanesulfonylpiperazin-1-yl)phenyl]urea
236	4-[4-(3-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}ureido)phenyl]piperazine-1-carboxylic acid methyl ester
237	1-[4-(1,1-Dioxo-1λ6-thiomorpholin-4-yl)phenyl]-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
240	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridin-3-yl)urea
241	1-(2,6-Dimethylpyridin-4-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
243	{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}carbamic acid 2-hydroxyethyl ester

244	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-3-yl]urea
245	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)pyridin-3-yl]urea
247	1-(6-Cyanopyridin-3-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
248	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((S)-3-ethylmorpholin-4-yl)pyridin-3-yl]urea
249	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((S)-3-ethylmorpholin-4-yl)pyridin-3-yl]urea
250	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-morpholin-4-ylpyridin-2-yl)urea
251	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[5-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-2-yl]urea
252	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[(1S,4S)-6-(2-oxa-5-azabicyclo[2.2.1]hept-5-yl)pyridin-3-yl]urea
253	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[(1S,4S)-6-(2-oxa-5-azabicyclo[2.2.1]hept-5-yl)pyridin-3-yl]urea
254	1-Cyclopropyl-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
255	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
256	1-[6-(3,3-Dimethylmorpholin-4-yl)pyridin-3-yl]-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea

257	1-[6-(3,3-Dimethylmorpholin-4-yl)pyridin-3-yl]-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
258	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-3-yl]urea
259	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-3-yl]urea
260	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-4-yl]urea
261	1-(4,4-Difluoro-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4'-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea

in the form of the base or of an acid-addition salt.

Table 2: Compounds according to the invention - IUPAC nomenclature

Combinations of the groups of compounds according to the invention also form part of the invention as an embodiment.

5

[PREPARATION PROCESS]

In the text hereinbelow, the term "PG" (protecting group) means a group that can, firstly, protect a reactive function such as a hydroxyl or an amine during the synthesis and, secondly, regenerate the intact reactive function at the end of the synthesis.

10 Examples of protecting groups and of protection and deprotection methods are given in Protective Groups in Organic Chemistry, J.F.W. McOmie, Plenum Press, 1973, in Greene's Protective Groups in Organic Synthesis, by Theodora W. Greene published by John Wiley & Sons Inc., 2006 or in Protecting Groups, Kocienski P.J., 1994, Georg Thieme Verlag. Examples that may be mentioned, as temporary protecting groups for

15 amines: benzyls, carbamates (such as tert-butyloxycarbonyl, which may be cleaved in acidic medium, or benzyloxycarbonyl, which may be cleaved by hydrogenolysis), as temporary protecting groups for carboxylic acids: alkyl esters (such as methyl, ethyl or

tert-butyl, which may be hydrolysed in basic or acidic medium) and benzyl esters, which may be hydrogenolysed, temporary protecting groups for alcohols or phenols such as tetrahydropyranyl, methyloxymethyl, methylethoxymethyl, tert-butyl, benzyl and tert-butyl dimethylsilyl ethers, temporary protecting groups for carbonyl derivatives such as linear or cyclic acetals, for instance 1,3-dioxan-2-yl or 1,3-dioxolan-2-yl; and reference may be made to the well-known general methods described in these publications.

The processes for protecting and deprotecting the protecting groups are typically those described in these publications.

Protection/deprotection steps for the various groups present on the compounds and reaction intermediates may be added before or after each of the steps mentioned below (or between two steps). Determining whether or not it is necessary to protect the groups, and the nature of the protecting groups to be used, are standard approaches for a person skilled in the art, who knows, in the light of his general knowledge, whether a group is liable to react during subsequent steps and whether or not it should be protected.


In the text hereinbelow, the term "LG" (leaving group) means a group that can be readily cleaved from a molecule by breaking a heterolytic bond, with loss of an electron pair. This group may thus be readily replaced with another group, for example during a substitution reaction. Such leaving groups are, for example, halogen atoms or an activated hydroxyl group such as a methanesulfonate, benzenesulfonate, p-toluenesulfonate, triflate, acetate, etc. Examples of leaving groups and also references for their preparation are given in *Advances in Organic Chemistry*, J. March, 3rd Edition, Wiley Interscience, pp. 310-316.

The intermediates described below may comprise groups that are precursors of other functions which are generated subsequently in one or more other steps.

In accordance with the invention, the compounds of general formula (I) may be prepared according to the processes that follow.


In a first embodiment, the process for preparing the compound of formula (I) may comprise step m) below:

m) grafting a group $-X_2-R_6$ onto a compound of formula (XLII) below:



in which R_1 , n , m , R_2 , R_3 , R_4 , G_1 and X_1 are as defined above,
to obtain the compound of formula (I).


The reaction scheme for the preparation of the compound of formula (I) via step
5 m) is represented in scheme 1 below.



Scheme 1: reaction scheme for preparing the compound of formula (I) from the
compound of formula (XLII).

When the process performed requires the protection of X_1 (more precisely of the
10 hydroxyl function when X_1 represents -O- or of the amine function when X_1 represents
-NR₄₀-), the process for preparing the compound of formula (XLII) used in the
abovementioned step m) generally comprises step l) below:

l) deprotection of a compound of formula (XLI) below:




15 in which:

- R_1 , n , m , R_2 , R_3 , R_4 , G_1 and X_1 are as defined above, and
- PG_1 is a protecting group for X_1 (protecting group for a hydroxyl function when

X_1 represents -O- or for an amine when X_1 represents $-NR_{40-}$,
to obtain the compound of formula (XLII).

The reaction scheme for the preparation of the compound of formula (XLII) via
step I) is represented in scheme 2 below.

5



Scheme 2: Reaction scheme for the preparation of the compound of formula (XLII) from
the compound of formula (XLI), when PG_1 is a protecting group for X_1 .


Step I) is not necessary when the process performed does not require protection
of X_1 . In this case, in the steps that follow, PG_1 represents H and the compound of
formula (XLII) corresponds to the compound of formula (XLI) in which PG_1 represents H.

Thus, in the steps described below (for preparing the compounds of formula (I),
(XLI) or (XLII)), PG_1 represents H or a protecting group for X_1 .

Two alternative embodiments have been developed for preparing the
abovementioned compounds of formula (XLI).


In a first alternative, the process for preparing the compound of formula (XLI) may
comprise the following steps:

c) Suzuki reaction of a compound of formula (XIII) below:




in which m , R_2 , R_3 and R_4 are as defined above and LG_2 is a leaving group, for
example a halogen atom such as chlorine, with a compound of formula $PG_1-X_1-G_1-$
 R_{300} , in which X_1 , and G_1 are as defined above, PG_1 is H or a protecting group for X_1

and R_{300} is a boronic acid or ester radical, typically with a compound of formula (CII) below:




in which X_1 , PG_1 and G_1 are as defined above,

5 to obtain a compound of formula (XXI) below:



in which m , R_2 , R_3 , R_4 , X_1 , PG_1 and G_1 are as defined above,


d) oxidation of the compound of formula (XXI) obtained in step c) to obtain a compound of formula (XXII) below:



10

in which m , R_2 , R_3 , R_4 , X_1 , PG_1 and G_1 are as defined above,

e) nucleophilic attack of a compound of formula (CI) below:




in which n and R_1 are as defined above,

15 on the compound of formula (XXII) obtained in step d),
to obtain the compound of formula (XLI).

When PG_1 represents H, the compound of formula (XLI) obtained at the end of step e) corresponds to the compound of formula (XLII), which may be engaged directly in step m), without the deprotection step l) being necessary.


The reaction scheme for the preparation of the compound of formula (XLI) via steps c) to e) is represented in scheme 3 below (in the embodiment in which step c) uses the compound of formula (CII)).



Scheme 3: reaction scheme for the preparation of the compound of formula (XLI) starting with the compound of formula (XIII) via steps c) to e).

10 According to a second alternative, the process for preparing the compound of formula (XLI) may comprise the following steps:


f) nucleophilic attack of a compound of formula PG_2-OH , in which PG_2 is a protecting group for the hydroxyl function, for example a benzyl, on a compound of formula (XIII) below:



15


in which m , R_2 , R_3 and R_4 are as defined above and LG_2 is a leaving group, for example a halogen atom such as chlorine,

to obtain a compound of formula (XXXI) below:




in which PG₂, m, R₂, R₃ and R₄ are as defined above,

- g) oxidation of the compound of formula (XXXI) obtained in step f),
 5 to obtain a compound of formula (XXXII) below:




in which m, R₂, R₃, R₄ and PG₂ are as defined above,

- h) nucleophilic attack of a compound of formula (CI) below:




- 10 in which n and R₁ are as defined above,
 on the compound of formula (XXXII) obtained in step g),
 to obtain the compound of formula (XXXIII) below:




in which n , R_1 , m , R_2 , R_3 , R_4 and PG_2 are as defined above,

- i) deprotection of the compound of formula (XXXIII) obtained in step h),
to obtain a compound of formula (XXXIV) below:



- 5 in which n , R_1 , m , R_2 , R_3 and R_4 are as defined above,


- j) grafting of a leaving group LG_5 onto the compound of formula (XXXIV) obtained in step i),
to obtain the compound of formula (XXXV) below:



- 10 in which n , R_1 , m , R_2 , R_3 and R_4 are as defined above and LG_5 is a leaving group, for example a halogen atom or a triflate,

- k) Suzuki reaction of the compound of formula (XXXV) obtained in step j) with a compound of formula $PG_1-X_1-G_1-R_{300}$, in which X_1 and G_1 are as defined above, PG_1 is H or a protecting group for X_1 and R_{300} is a boronic acid or ester radical,

- 15 typically with a compound of formula (CII) below:




in which X_1 , PG_1 and G_1 are as defined above,

to obtain the compound of formula (XLI).

When PG_1 represents H, the compound of formula (XLI) obtained at the end of step k) corresponds to the compound of formula (XLII), which may be engaged directly in step m), without the deprotection step l) being necessary.


- 5 The reaction scheme for the preparation of the compound of formula (XLI) via steps f) to k) is represented in scheme 4 below (in the embodiment in which step k) uses the compound of formula (CII)).



- Scheme 4: reaction scheme for the preparation of the compound of formula (XLI) starting with the compound of formula (XIII) via steps f) to k).


The process for the preparation of the abovementioned compound of formula (XIII), performed in steps c) or f) described above, typically comprises the following steps:

- a) nucleophilic attack of a compound of formula (XI) below:




(XI)

in which m, R₂, R₃ and R₄ are as defined above,
on a compound of formula (X) below:



(X)

- 5 in which LG₂ is as defined above and LG₁ and LG₄ are independently leaving groups, for example halogen atoms such as chlorine atoms, to obtain a compound of formula (XII) below:




(XII)

in which m, R₂, R₃, R₄, LG₁ and LG₂ are as defined above,

- 10 b) cyclization of the compound of formula (XII) obtained in step a),
to obtain the compound of formula (XIII).


The reaction scheme for the preparation of the compound of formula (XIII) via steps a) and b) is represented in scheme 5 below.



Scheme 5: reaction scheme for the preparation of the compound of formula (XIII) starting with the compound of formula (X) via steps a) and b).

In a second embodiment, when, in formula (I), X₁ represents -NR₄₀⁻ and X₂ represents -CONR₅₀⁻, the process for preparing the compound of formula (I) may comprise the following step:


n) Suzuki reaction of the compound of formula (XXXV):



10 in which n, R₁, m, R₂, R₃ and R₄ are as defined above and LG₅ is a leaving group, for example a halogen atom or a triflate,

with a compound of formula R₆R₅₀N-(C=O)-NR₄₀-G₁-R₃₀₀, in which G₁, R₆, R₅₀ and R₄₀ are as defined above and R₃₀₀ is a boronic acid or ester radical, typically with a compound of formula (CIII) below:


52



in which G_1 , R_6 , R_{50} and R_{40} are as defined above,
to obtain the compound of formula (I) in which X_1 represents $-NR_{40}-$ and X_2 represents $-CONR_{50}-$.

- 5 The process for the preparation of the compound of formula (XXXV) performed in step n) typically comprises the abovementioned steps f), g), h), i) and j).

The reaction scheme for the preparation of the compound of formula (I) via step n) is represented in scheme 6 below (in the embodiment in which step n) uses the compound of formula (CIII)).





10

Scheme 6: reaction scheme for preparing the compound of formula (I) from the compound of formula (XXXV) via a step n).

In a third embodiment, the process for preparing the compound of formula (I) may comprise step q) below:

- 15 q) nucleophilic attack of a compound of formula (CI) below:





Chemical reaction scheme showing the conversion of compound (LII) to compound (I) using reagent (CI).

Compound (LII) is a pyrimidine derivative with a sulfonamide group, a morpholine ring, and a side chain with substituents R_6 , X_2 , X_1 , G_1 , R_4 , R_3 , R_2 , and a repeating unit $(CH_2)_m$.

Reagent (CI) is a morpholine derivative with substituent R_1 .

The reaction arrow is labeled 'q)'.

Compound (I) is a pyrimidine derivative where the sulfonamide group has been converted to a morpholine ring, and the side chain remains unchanged.


The process for the preparation of the compound of formula (LII) performed in the abovementioned step q) typically comprises the following steps:

CSC1=NC2=C(N1)C(=C(C2=O)C(=O)N(R4)CC(R2)CC(R3)O)C3=CC=CC=C3LG2

(XIII)


in which m , R_2 , R_3 , R_4 and LG_2 are as defined above,

with a compound of formula $R_6-X_2-X_1-G_1-R_{300}$, in which R_6 , X_2 , X_1 and G_1 are as defined above and R_{300} is a boronic acid or ester radical, typically with a compound of formula (CIV) below:



(CIV)

- 5 in which R_6 , X_2 , X_1 and G_1 are as defined above, to obtain the compound of formula (LI) below:




(LI)

in which R_6 , X_2 , X_1 , G_1 , m , R_2 , R_3 and R_4 are as defined above,

- p) oxidation of the compound of formula (LI) obtained in step o),
10 to obtain the compound of formula (LII).

The process for the preparation of the compound of formula (XIII) performed in step o) typically comprises the abovementioned steps a) and b).

- The reaction scheme for the preparation of the compound of formula (LII) via steps o) and p) is represented in scheme 8 below (in the embodiment in which step o)
15 uses the compound of formula (CIV)):



Scheme 8: reaction scheme for the preparation of the compound of formula (LII) starting with the compound of formula (XIII) via steps o) and p).

Embodiments for the abovementioned steps a) to q) are described below.

- 5 Step m) may be performed according to the procedures described in *Advanced Organic Chemistry*, J. March, 3rd Edition, Wiley Interscience, and the references cited therein.

For example, when X₂ represents a single bond and R₆ represents a group -G₂-X₃-G₃ in which:

- 10 - G₂ represents a divalent heteroaryl radical optionally substituted with one or more groups R₈₀, X₃ and G₃ being as defined above, or
- G₂ and X₃ represent single bonds and G₃ represents heteroaryl optionally substituted with one or more groups R₈₁,

- step m) may consist of a Buchwald-Hartwig reaction, as described in Topics in Current Chemistry Chemical Review 2002, 219, 131-209, by reaction of the compound of formula
- 15 (XLII) with a compound of formula LG₇-G₂-X₃-G₃ in which:

- G₂ represents a divalent heteroaryl radical optionally substituted with one or more groups R₈₀, or
- G₂ and X₃ represent single bonds and G₃ represents heteroaryl optionally substituted
- 20 with one or more groups R₈₁, and

- LG₇ represents a leaving group, for example a halogen atom,
 in the presence of a palladium catalyst, a base such as potassium carbonate, potassium
 fluoride, potassium tert-butoxide or potassium phosphate, and optionally a phosphine, in
 a solvent typically chosen from dimethylformamide, dioxane, ethylene glycol dimethyl
 5 ether, toluene, tert-butanol, tetrahydrofuran and water, or a mixture thereof, at a
 temperature generally ranging from room temperature to 150°C.

Four embodiments were developed for performing step m) when X₂ represents
 -CONR₅₀-, -CONR₅₁-O- or -COO-.

In a first embodiment, step m) comprises the following steps:

- 10 m1-1) reaction of the compound of formula (XLII) with phosgene or triphosgene in the
 presence of a base,
 m1-2) reaction of a compound of formula HNR₅₀R₆, HNR₅₁-O-R₆ or HO-R₆ in which R₆,
 R₅₀ and R₅₁ are as defined above with the compound obtained in step m1-1), to
 obtain the compound of formula (I),
 15 reactions m1-1) and m1-2) possibly being performed as a one-pot reaction.

In a second embodiment, step m) comprises the following steps:

- m2-1) reaction of a compound of formula HNR₅₀R₆, HNR₅₁-O-R₆ or HO-R₆ in which R₆,
 R₅₀ and R₅₁ are as defined above with phosgene or triphosgene in the presence
 of a base,
 20 m2-2) reaction of the compound of formula (XLII) with the compound obtained in step
 m2-1), to obtain the compound of formula (I),
 reactions m2-1) and m2-2) possibly being performed as a one-pot reaction.

Steps m1-1), m1-2), m2-1) and m2-2) are typically performed in dichloromethane,
 dioxane or a mixture of the two. Steps m1-1) and m2-2) are generally performed in the
 25 presence of an amine base such as triethylamine or diisopropylethylamine.

In a third embodiment, step m) comprises the following steps:

- m3-1) reaction of the compound of formula (XLII) with an alkyl or aryl chloroformate in
 the presence of a base,
 m3-2) reaction of a compound of formula HNR₅₀R₆, HNR₅₁-O-R₆ or HO-R₆ in which R₆,
 30 R₅₀ and R₅₁ are as defined above with the compound obtained in step m3-1), to
 obtain the compound of formula (I),
 reactions m3-1) and m3-2) possibly being performed as a one-pot reaction.

In a fourth embodiment, step m) comprises the following steps:

- m4-1) reaction of a compound of formula HNR₅₀R₆, HNR₅₁-O-R₆ or HO-R₆ in which R₆,
 35 R₅₀ and R₅₁ are as defined above with an alkyl or aryl chloroformate in the

presence of a base,
 m4-2) reaction of the compound of formula (XLII) with the compound obtained in step
 m4-1), to obtain the compound of formula (I),
 reactions m4-1) and m4-2) possibly being performed as a one-pot reaction.

5 Steps m3-1), m3-2), m4-1) and m4-2) are typically performed in a polar aprotic solvent such as dimethylformamide, tetrahydrofuran or a mixture of the two. Steps m3-1) and m4-1) are typically performed with phenyl chloroformate, generally at room temperature. Steps m3-2) and m4-2) are typically performed between 80 and 120°C and may be performed by microwave.

10 Step a) is typically performed in the presence of a base, especially an amine base such as triethylamine, generally in a polar aprotic solvent such as tetrahydrofuran, at a temperature especially of -78°C.

Step b) is typically performed in the presence of a base, especially an amine base such as 1,8-diazabicyclo[5.4.0]undec-7-ene, generally in a polar aprotic solvent such as
 15 dimethyl sulfoxide, at a temperature especially between room temperature (25°C) and 60°C.

Steps c), n) and o), which correspond to Suzuki coupling reactions, are typically performed in the presence of a mineral base such as sodium or potassium carbonate, potassium fluoride, potassium tert-butoxide or tripotassium phosphate and of a palladium
 20 complex such as tetrakis(triphenylphosphine)palladium or bis(triphenylphosphine)palladium (II) chloride, generally in a solvent chosen from dimethylformamide, dioxane, ethylene glycol dimethyl ether, toluene, tetrahydrofuran and water or a mixture thereof, typically in a dioxane/water mixture, typically at a temperature from room temperature to 150°C. The conditions described in Chemical Review 2007,
 25 107, 133-173 may especially be followed. These steps may be performed by microwave.

Steps d), g) and p) are typically performed in the presence of an oxidizing agent such as magnesium monoperoxyphthalate hexahydrate, generally in an acetonitrile/alcohol mixture such as ethanol, typically a 3/1 to 3/2 mixture, generally at a temperature from room temperature to 70°C.

30 Steps e), h) and q) are generally performed in the presence of a base, especially an amine base such as triethylamine, typically in a polar aprotic solvent such as dimethyl sulfoxide or dioxane, at a temperature of between 70 and 110°C. These steps may be performed by microwave.

The conditions of step I) vary as a function of the nature of the protecting group

PG₁. PG₁ is especially a Boc group. When PG₁ is a Boc group, step l) is typically performed in a solvent generally chosen from dichloromethane, diethyl ether, tetrahydrofuran and dioxane, or a mixture thereof, typically dichloromethane, in the presence of an acid such as trifluoroacetic acid or hydrochloric acid, generally at room temperature.

The conditions of step f) vary according to the nature of the protecting group PG₂. PG₂ is especially a benzyl group. When PG₂ is a benzyl group, step f) is typically performed by placing the compound of formula (XIII) in contact with benzoate (generally obtained by reacting benzyl alcohol and a base such as sodium hydride), typically in an anhydrous polar aprotic solvent, such as anhydrous tetrahydrofuran, generally at a temperature of between -10 and 10°C.

The conditions of step i) vary according to the nature of the protecting group PG₂. PG₂ is especially a benzyl group. When PG₂ is a benzyl group, step i) may be performed:

- by hydrogenolysis, for example in the presence of hydrogen and palladium-on-charcoal, or
- by reaction with ammonium formate in the presence of a palladium catalyst, especially in methanol at a temperature between room temperature and the reflux point.

The conditions of step j) vary according to the nature of the protecting group LG₅. LG₅ is especially a triflate group. When LG₅ is a triflate group, step i) may be performed by reacting the compound of formula (XXXIV) with triflic anhydride, typically in the presence of a base, especially an amine base such as triethylamine, for example dichloromethane, at a temperature generally between -10 and 10°C.

In the abovementioned reaction schemes, the starting compounds and the reagents, when their preparation method is not described, are commercially available or described in the literature, or alternatively may be prepared according to methods that are described therein or that are known to those skilled in the art.

The examples that follow describe the preparation of certain compounds in accordance with the invention. These examples are not limiting and serve merely to illustrate the present invention. The numbers of the compounds exemplified refer to those given in Table 3 below, which illustrates the chemical structures and physical properties of a number of compounds according to the invention.

A person skilled in the art can adapt without difficulty the teaching below to the compounds of general formula (I). He will be capable of selecting, in the light of his knowledge and of the literature, the appropriate protecting groups enabling the introduction of all the groups or functions described in the present invention.

In the procedures and examples below:

- the proton magnetic resonance spectra (^1H NMR), as described below, are recorded at a temperature of 300 K (exchangeable protons not recorded) at 300, 400 or 600 MHz in DMSO- d_6 , using the DMSO- d_6 peak as reference. The chemical shifts δ are expressed in parts per million (ppm). The signals observed are expressed as follows:
s = singlet, d = doublet, m = multiplet, bs = broad signal, t = triplet, q = quartet.
- The LC/MS characteristics as described below indicate successively the high-performance liquid chromatography analytical method used and detailed below, the peak $\text{M}+\text{H}^+$ identified by mass spectrometry and the retention time (T_r) of the compound expressed in minutes.

* Method A

Instrument: Acquity UPLC chain (Waters); LCT mass spectrometer (Waters)

Column: BEH C18 50 x 2.1 mm 1.7 μm , $T^\circ = 40^\circ\text{C}$

Solvent A: H_2O + 0.05% TFA; Solvent B: acetonitrile + 0.035% TFA

15 Flow rate: 1.0 ml/min

Gradient A/B: t 0 min 2% B, t 1.6 min 100% B, t 2.1 min 100% B, t 2.5 min 2% B, t 3.0 min 2% B

Detection: UV 220 nm

Ionization: electrospray positive mode

20 * Method B

Instrument: Alliance HPLC chain (Waters); ZQ mass spectrometer (Waters)

Column: Kromasil C18 50 x 2.1 mm, 3.5 μm , $T^\circ = 40^\circ\text{C}$

Solvent A: Ammonium acetate 5 mM + 3% acetonitrile; Solvent B: acetonitrile

Flow rate: 0.8 ml/min

25 Gradient A/B: t 0 min 0% B, t 5.5 min 100% B, t 7.0 min 100% B, t 7.1 min 0% B, t 10.0 min 0% B

Detection: UV 220 nm

Ionization: electrospray positive mode

* Method C

30 Instrument: Acquity UPLC type HPLC chain (Waters); SQD mass spectrometer (Waters)

Column: Ascentis Express C18 50 x 2.1 mm 2.7 μm , $T^\circ = 55^\circ\text{C}$

Solvent A: H_2O + 0.02 % TFA; Solvent B: acetonitrile + 0.014 % TFA

Flow rate: 1 ml/min

Gradient A/B: t 0 min 2% B, t 1.2 min 98% B, t 1.5 min 2% B

35 Detection: UV 220 nm

Ionization: electrospray positive mode

*** Method D**

Instrument: Acquity UPLC type HPLC chain (Waters); SQD mass spectrometer (Waters)

Column: BEH C18 100 x 2.1 mm 1.7 μ m, T° = 70°C

- 5 Solvent A: H₂O + 0.02 % TFA; Solvent B: acetonitrile + 0.014 % TFA

Flow rate: 0.8 ml/min

Gradient A/B: t 0 min 2% B, t 5.0 min 98% B, t 5.3 min 98% B, t 5.33 min 2% B

Detection: UV 220 nm

Ionization: electrospray positive mode

10 *** Method E**

Instrument: Alliance HPLC chain (Waters); ZQ mass spectrometer (Waters)

Column: Xbridge C18 30 x 2.1 mm 2.5 μ m, T° = 55°C

Solvent A: H₂O + 0.02% TFA; Solvent B: MeOH

Flow rate: 0.7 ml/min

- 15 Gradient A/B: t 0 min 2% B, t 3.0 min 100% B, t 3.5 min 100% B, t 3.6 min 2% B

Detection: UV 220 nm

Ionization: electrospray positive mode

*** Method F**

Instrument: Alliance HPLC chain (Waters); ZQ mass spectrometer (Waters)

- 20 Column: Waters Select CSH C18 30 x 7.5 mm, 3.5 μ m, T° = 60°C

Solvent A: H₂O + 0.1% formic acid; Solvent B: acetonitrile + 0.1% formic acid

Flow rate: 1.1 ml/min

Gradient A/B: t 0 min 6% B, t 0.8 min 6% B, t 4.7 min 100% B, t 4.8 min 100% B, t 5.0 min 6% B, t 6.0 min 6% B

- 25 Detection: UV 220 nm

Ionization: electrospray positive mode

The solvent mixtures are quantified in volumetric ratios.

The NMR spectra and mass spectra confirm the structures of the compounds obtained according to the examples below.

- 30 In the examples that follow, the following abbreviations and empirical formulae are used:

DBU: 1,8-diazabicyclo[5.4.0]undec-7-ene

DCM: dichloromethane


DIEA: N,N-diisopropylethylamine

- 35 DMF: N,N-dimethylformamide

- DMSO: dimethyl sulfoxide
 Et₂O: diethyl ether
 EtOH: ethanol
 HPLC: high performance liquid chromatography
 5 K₂CO₃: potassium carbonate
 LC/MS: liquid chromatography/mass spectrometry
 MeOH: methanol
 MgSO₄: magnesium sulfate
 MHz: MegaHertz
 10 Min: minute(s)
 Mmol: milimoles
 Na₂CO₃: sodium carbonate
 NaCl: sodium chloride
 NaHCO₃: sodium hydrogen carbonate
 15 Na₂SO₄: sodium sulfate
 Psi: pounds per square inch, with 1 psi = 0.069 bar
 TFA: trifluoroacetic acid
 THF: tetrahydrofuran
 °C: degrees Celsius
 20 Tr: retention time


Example 1:

- Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea**



Step 1.1: (preparation of a compound of formula (X))


4,6-Dichloro-2-methylsulfanylpurine-5-carbonyl chloride



10 g of 4,6-dichloro-2-methylsulfanylpurine-5-carboxylic acid (41.83 mmol) are placed in 61 ml of thionyl chloride (836.54 mmol) with stirring. After 18 hours at 80°C, the thionyl chloride is evaporated off and the residue is taken up twice with 15 ml of toluene and concentrated to give 10.77 g of 4,6-dichloro-2-methylsulfanylpurine-5-carbonyl chloride.

Step 1.2: (which corresponds to step a))

4,6-Dichloro-2-methylsulfanylpurine-5-carboxylic acid ethyl-(2-hydroxy-2-methylpropyl)amide




10

4.9 g of 1-(ethylamino)-2-methylpropan-2-ol (41.82 mmol) and 17.5 ml of triethylamine (125.46 mmol) are placed in 30 ml of anhydrous THF with stirring. After cooling the reaction medium with a bath of cardice in acetone to -70°C, 10.77 g and (41.82 mmol) of the compound described in step 1.1 diluted in 80 ml of THF are added dropwise. After stirring for 4 hours at -70°C, the reaction medium is poured into ice-water and then extracted twice with DCM. The organic phase is washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness. 13.54 g of 4,6-dichloro-2-methylsulfanylpurine-5-carboxylic acid ethyl-(2-hydroxy-2-methylpropyl)amide are obtained.

20 LC/MS (method F): M+H⁺ = 338; Tr = 3.70 and 3.80 min (conformers)

Step 1.3: (which corresponds to step b))

4-Chloro-6-ethyl-8,8-dimethyl-2-methylsulfanyl-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one




13 g (38.43 mmol) of the compound described in step **1.2** are placed in 200 ml of DMSO, with stirring. 6.3 ml of DBU (42.28 mmol) are added. The reaction medium is stirred for 5 hours at 55°C, and 500 ml of ice-water are then added. A white precipitate forms, which is filtered off and rinsed twice with water. The solid is then taken up in
 5 acetonitrile and filtered off to give 3.6 g of 4-chloro-6-ethyl-8,8-dimethyl-2-methylsulfanyl-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one.

The mother liquors are recovered and extracted with 3 times 300 ml of DCM. The organic phases are combined, washed with brine, dried over Na₂SO₄ and then filtered. The solution is evaporated to dryness. The residue is then chromatographed on silica gel
 10 with a gradient of ethyl acetate in DCM ranging from 0% to 10%. 3 g of 4-chloro-6-ethyl-8,8-dimethyl-2-methylsulfanyl-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one are obtained.

LC/MS (method F): M+H⁺ = 302; Tr = 3.68 min

Step 1.4: (which corresponds to step c))

15 **[4-(6-Ethyl-8,8-dimethyl-2-methylsulfanyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)-2-fluorophenyl]carbamic acid tert-butyl ester**




8 g (26.51 mmol) of the compound described in step **1.3** are placed in 150 ml of dioxane. 7.67 g (29.16 mmol) of 4-N-Boc-amino-3-fluorophenylboronic acid, 66.27 ml
 20 (132.5 mmol) of aqueous 1N Na₂CO₃ solution and 1.53 g (1.33 mmol) of tetrakis(triphenylphosphine)palladium are successively added, with stirring. The mixture is stirred at 85°C for 2 hours and is then poured into water. The precipitate obtained is then filtered off and rinsed with water. The solid obtained is taken up in MeOH, filtered off and rinsed with pentane. 11.2 g of [4-(6-ethyl-8,8-dimethyl-2-methylsulfanyl-5-oxo-
 25 5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)-2-fluorophenyl]carbamic acid tert-butyl ester are obtained.

LC/MS (method F): M+H⁺ = 477; Tr = 4.55 min

Step 1.5: (which corresponds to step d))

[4-(6-Ethyl-2-methanesulfonyl-8,8-dimethyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-

triazabenzocyclohepten-4-yl)-2-fluorophenyl]carbamic acid tert-butyl ester




21 g (44.06 mmol) of the compound described in step **1.4** and 34.1 g of magnesium monoperoxyphthalate hexahydrate (88.13 mmol) are placed in 300 ml of a 2/1 acetonitrile/EtOH mixture. After stirring for 18 hours at room temperature, the solvents are evaporated off. The residue is taken up in 500 ml of water. The product obtained is filtered off and rinsed twice with EtOH to give 22 g of [4-(6-ethyl-2-methanesulfonyl-8,8-dimethyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-

LC/MS (method F): $M+H^+ = 509$; Tr = 4.08 min

Step 1.6: (which corresponds to step e))

{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid tert-butyl ester



15


5 g (9.83 mmol) of the compound described in step **1.5** are placed in 100 ml of DMSO. 5.67 ml of triethylamine (39.33 mmol) and 2.94 g of 8-oxa-3-azabicyclo[3.2.1]octane hydrochloride (19.66 mmol) are successively added. After stirring for 1 hour 30 minutes at 90°C, the DMSO is evaporated off. The residue is taken up in 200 ml of water and the solid formed is filtered off, rinsed with water, with acetonitrile and then with pentane to give 5 g of {4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid tert-butyl ester.

20

LC/MS (method F): $M+H^+ = 542$; Tr = 4.49 min

Step 1.7: (which corresponds to step l))

4-(4-Amino-3-fluorophenyl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one




5

5 g (9.23 mmol) of the compound described in step **1.6** are placed in 90 ml of DCM. 21.3 ml of TFA are added dropwise and the reaction medium is stirred at room temperature for 3 hours. Saturated aqueous NaHCO_3 solution is added to pH 10, and the reaction medium is extracted twice with DCM. The organic phase is dried over Na_2SO_4 , filtered and evaporated. The solid is taken up in acetonitrile and filtered off to give 3.25 g of 4-(4-amino-3-fluorophenyl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one.

LC/MS (method F): $M+H^+ = 442$; Tr = 3.71 min

Step 1.8: (which corresponds to step m) (steps m1-1) and m1-2))

15 **1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea**



400 mg (0.91 mmol) of the compound described in step **1.7** and 0.24 ml of DIEA (1.36 mmol) are placed in 10 ml of dioxane. 0.715 ml of 1.9 N phosgene in toluene (1.6 mmol) is then added. After stirring for 30 minutes at 50°C, 3.17 ml of 2 N methylamine in

20

THF (6.34 mmol) are added. After stirring for 1 hour at room temperature, the reaction mixture is filtered. The solid is rinsed with water, with ethyl acetate and then with pentane. 343 mg of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-


5 methylurea are obtained.

LC/MS (method A): $M+H^+$ = 499; Tr = 0.90 min

^1H NMR (400 MHz, DMSO- d_6) δ ppm 8.48 (m, 1H); 8.14 (dd, 1H); 7.35 (dd, 1H); 7.24 (dd, 1H); 6.53 (q, 1H); 4.41(m, 2H); 4.22 (m, 2H); 3.66 (s, 2H); 3.51 (m, 2H); 3.10 (d, 2H); 2.67 (d, 3H); 1.81 (m, 2H); 1.64 (m, 2H); 1.34 (s, 6H); 1.14 (t, 3H)


10 **Example 2:**

Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea



Step 2.1: (which corresponds to step e))

15 **{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid tert-butyl ester**




2.5 g of [4-(6-ethyl-2-methanesulfonyl-8,8-dimethyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)-2-fluorophenyl]carbamic acid tert-butyl ester

(19.66 mmol) in 20 ml of dioxane are placed in 4 microwave tubes, respectively. One quarter of (S)-3-methylmorpholine (98.32 mmol) is added to each tube, respectively. The tubes are heated for 2 hours at 100°C in a Biotage microwave machine. The reaction media are filtered and rinsed with MeOH. The solid is washed with water, with acetonitrile and then with pentane. 3.7 g of {4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid tert-butyl ester are obtained. The DMSO/MeOH phase is evaporated to dryness. The residue is taken up in water and filtered, rinsed with acetonitrile and then with pentane. This product is chromatographed on silica gel with a gradient of MeOH in DCM ranging from 0 to 5%. 1.7 g of {4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid tert-butyl ester are obtained.

LC/MS (method F): $M+H^+ = 530$; Tr = 4.52 min

Step 2.2: (which corresponds to step l))

15 **4-(4-Amino-3-fluorophenyl)-6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one**




5.4 g (10.2 mmol) of the compound described in step 2.1 are placed in 100 ml of DCM. 23.57 ml of TFA (305.89 mmol) are added dropwise. After stirring for 2 hours at room temperature, the medium is concentrated, taken up in water and basified with saturated NaHCO_3 solution. The aqueous phase is extracted with DCM and the organic phase is washed with brine, dried over Na_2SO_4 and then concentrated. 4.2 g of 4-(4-amino-3-fluorophenyl)-6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one are obtained.

25 LC/MS (method F): $M+H^+ = 430$; Tr = 3.75 min

Step 2.3: (which corresponds to step m) (steps m1-1) and m1-2))

1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea




400 mg (0.93 μmol) of the compound described in step **2.2** and 0.23 ml of DIEA (1.30 mmol) are placed in 20 ml of dioxane. 0.686 ml of 1.9 N phosgene in toluene (1.3 mmol) is added. After stirring for 30 minutes at 50°C, 2.33 ml of 2 N methylamine in THF
 5 (4.66 mmol) are added. After stirring for 1 hour at room temperature, the reaction mixture is washed with water, dried over Na_2SO_4 , filtered and concentrated. The residue is chromatographed on silica gel with a gradient of a 95/5 ethyl acetate/MeOH mixture in DCM ranging from 0% to 100%. 268 mg of 1-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea are obtained.

LC/MS (method A): $\text{M}+\text{H}^+ = 487$; $\text{Tr} = 0.91$ min

^1H NMR (400 MHz, DMSO-d_6) δ ppm 8.48 (m, 1H); 8.14 (dd, 1H); 7.34 (dd, 1H); 7.25 (dd, 1H); 6.54 (q, 1H); 4.65 (m, 1H); 4.34 (m, 1H); 3.91 (m, 1H); 3.72-3.67 (m, 3H); 3.59-3.51 (m, 3H); 3.45-3.36 (m, 1H); 3.22-3.15 (m, 1H); 2.67 (d, 3H); 1.35 (s, 6H); 1.21 (d,
 15 3H); 1.15 (t, 3H)


Example 3:

Synthesis of 1-ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea



Step 3.1: (which corresponds to step f))

4-Benzyloxy-6-ethyl-8,8-dimethyl-2-methylsulfonyl-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one



5


6.20 ml of benzyl alcohol (59.64 mmol) are placed in 400 ml of anhydrous THF. The solution is cooled to 4°C in an ice bath and 2.39 g of 60% sodium hydride (59.64 mmol) are added portionwise. The reaction mixture is stirred at 4°C for 15 minutes. 15 g of 4-chloro-6-ethyl-8,8-dimethyl-2-methylsulfonyl-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one (49.70 mmol) dissolved in 400 ml of anhydrous THF are added dropwise. After stirring for 2 hours at 4°C, 400 ml of water are added and the THF is evaporated off. The aqueous phase is extracted twice with 300 ml of DCM, and the organic phases are combined and washed with 400 ml of water and then 400 ml of brine. The organic phase is dried over Na₂SO₄, filtered and concentrated. The residue is chromatographed on silica gel with a gradient of ethyl acetate in heptane ranging from 30% to 50%. 18.6 g of 4-benzyloxy-6-ethyl-8,8-dimethyl-2-methylsulfonyl-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one are obtained.

LC/MS (method F): M+H⁺ = 374; Tr = 4.21 min

Step 3.2: (which corresponds to step g))

4-Benzyloxy-6-ethyl-2-methanesulfonyl-8,8-dimethyl-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one

20




18.6 g (49.80 mmol) of the compound described in step **3.1** and 49.27 g of magnesium monoperoxyphthalate hexahydrate (99.61 mmol) are placed in 375 ml of a 2/1 acetonitrile/EtOH mixture. After stirring for 2 hours at room temperature, the solvents are evaporated off. The residue is taken up in 400 ml of water. The mixture is filtered and rinsed twice with water. The solid obtained is taken up in DCM, dried over Na₂SO₄, filtered and evaporated to give 15.9 g of 4-benzyloxy-6-ethyl-2-methanesulfonyl-8,8-dimethyl-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one.

LC/MS (method F): M+H⁺ = 406; Tr = 3.70 min

10 **Step 3.3:** (which corresponds to step h))

4-Benzyloxy-6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one



11.23 g (27.7 mmol) of the compound described in step **3.2** are placed in 120 ml of DMSO. 5.79 ml of triethylamine (41.54 mmol) and 8.4 g of (S)-3-methylmorpholine (83.09 mmol) are successively added. The medium is dried with anhydrous MgSO₄ with stirring for 30 minutes. The mixture is filtered and divided among six 20 ml microwave tubes. The tubes are heated for 30 minutes at 100°C in a Biotage microwave machine.


The reaction media are poured into water and extracted three times with ethyl acetate. The organic phase is washed three times with water, dried over Na₂SO₄, filtered and concentrated. The residue is chromatographed on silica gel with a gradient of ethyl

acetate in DCM to give 6.91 g of 4-benzyloxy-6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one.

LC/MS (method F): $M+H^+ = 427$; Tr = 4.16 min

Step 3.4: (which corresponds to step i))

- 5 **6-Ethyl-4-hydroxy-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one**




- 6.9 g (16.18 mmol) of the compound described in step **3.3** and 160 ml of EtOH are placed in a Parr bottle. 0.17 g of 50% hydrated 10% palladium-on-charcoal (1.62 mmol) is added and the reaction medium is stirred at 40 psi of hydrogen for 1 hour at 25°C. The reaction medium is filtered and concentrated to give 4.81 g of 6-ethyl-4-hydroxy-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one.

LC/MS (method F): $M+H^+ = 337$; Tr = 3.40 min

- 15 **Step 3.5:** (which corresponds to step j))

Trifluoromethanesulfonic acid 6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl ester




- 4.6 g (13.67 mmol) of the compound described in step **3.4** are placed in 140 ml of DCM. 2.86 ml of triethylamine (20.51 mmol) are added. The solution is cooled to 4°C in an ice bath and 3.47 ml of trifluoromethanesulfonic anhydride (20.51 mmol) are then added. After stirring the mixture at 4°C for 3 hours, 300 ml of water are added. The organic phase is extracted and washed with saturated aqueous NaHCO_3 solution, dried over Na_2SO_4 , filtered and concentrated. The residue is chromatographed on silica gel with a gradient of from 20% to 50% of ethyl acetate in heptane to give 5.27 g of trifluoromethanesulfonic acid 6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-

5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl ester.

LC/MS (method F): $M+H^+ = 469$; Tr = 4.35 min

Step 3.6: (preparation of a compound of formula (CIII))

1-Ethyl-3-[2-fluoro-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]urea




5

5 g of 4-amino-3-fluorophenylboronic acid pinacol ester (21.08 mmol) are placed in 100 ml of DCM. 3.23 ml of triethylamine (23.20 mmol) are added. The reaction mixture is cooled in an ice bath to 4°C. After stirring for 5 minutes, 2.21 g of triphosgene (7.38 mmol) are added portionwise, and the ice bath is then removed. After stirring for 1 hour
 10 30 minutes at room temperature, 8.77 g of ethylamine hydrochloride (105.45 mmol) and then 14.57 g of K_2CO_3 (105.45 mmol) are added portionwise. After 2 hours, the medium is cooled in an ice bath and then acidified with 1 N hydrochloric acid solution. The aqueous phase is extracted 3 times with DCM. The organic phases are combined, washed with brine, dried over $MgSO_4$, filtered and concentrated. The residue is
 15 chromatographed on silica gel with a gradient of from 0% to 5% MeOH in DCM to give 4.65 g of 1-ethyl-3-[2-fluoro-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]urea. LC/MS (method F): $M+H^+ = 309$; Tr = 4.11 min

Step 3.7: (which corresponds to step n))

**1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-
 20 tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl]urea**




420 mg (0.89 mmol) of the compound described in step **3.5** are placed in 10 ml of dioxane in a 20 ml microwave tube. 276 mg (0.89 mmol) of the compound described in step **3.6**, 380 mg of tribasic potassium phosphate (1.79 mmol) and 0.5 ml of water are successively added. The solution is degassed with argon, and 63 mg of bis(triphenylphosphine)palladium (II) chloride (0.09 mmol) are then added. The tube is heated at 100°C for 20 minutes in a Biotage microwave machine. The reaction medium is diluted in DCM, and washed twice with water and once with brine. The organic phase is dried over Na₂SO₄, filtered and concentrated. The residue is chromatographed on silica gel with a gradient of a 5/1 ethyl acetate/MeOH mixture in DCM ranging from 0% to 30%. The product is taken up in Et₂O and then filtered off to give 222 mg of 1-ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea.

LC/MS (method A): M+H⁺ = 501; Tr = 1.02 min

¹H NMR (400 MHz, DMSO-d₆) δ ppm 8.42 (m, 1H); 8.15 (dd, 1H); 7.34 (dd, 1H); 7.24 (dd, 1H); 6.66 (q, 1H); 4.65 (m, 1H); 4.32 (m, 1H); 3.91 (m, 1H); 3.72-3.67 (m, 3H); 3.59-3.37 (m, 4H); 3.21-3.09 (m, 3H); 1.35 (s, 6H); 1.21 (d, 3H); 1.15 (t, 3H); 1.07 (t, 3H)


Example 4:

Synthesis of 1-{4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(S)-1-pyrrolidin-2-ylmethyl-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-ethylurea



Step 4.1: (preparation of a compound of formula (XI))

(S)-2-[(2-Hydroxy-2-methylpropylamino)methyl]pyrrolidine-1-carboxylic acid tert-butyl ester




5

4.34 g of 1-amino-2-methylpropan-2-ol (48.68 mmol) are placed in 50 ml of MeOH. 2.79 ml of acetic acid (48.68 mmol) and then 4.72 ml of N-Boc-L-prolinal (24.34 mmol), diluted beforehand in 20 ml of MeOH, are successively added dropwise. The reaction mixture is stirred at room temperature for 30 minutes and 3.22 g of sodium cyanoborohydride (48.68 mmol) are then added portionwise. After stirring at room temperature for 18 hours, the medium is poured into 400 ml of saturated aqueous NaHCO₃ solution. The aqueous phase is extracted with 3 times 150 ml of DCM. The organic phases are combined, washed with 200 ml of water, with 200 ml of brine, dried over MgSO₄, filtered and evaporated. 7.19 g of (S)-2-[(2-hydroxy-2-methylpropylamino)methyl]pyrrolidine-1-carboxylic acid tert-butyl ester are obtained.

LC/MS (method F): M+H⁺ = 273; Tr = 2.55 min

Step 4.2: (which corresponds to step a))


(S)-2-[[[(4,6-Dichloro-2-methylsulfanylpurimidine-5-carbonyl)-(2-hydroxy-2-methylpropyl)amino]methyl]pyrrolidine-1-carboxylic acid tert-butyl ester



6.63 g (24.34 mmol) of the compound described in step **4.1** are placed in 17 ml of anhydrous THF. The reaction mixture is cooled in a bath of cardice in acetone at -70°C , followed by successive dropwise addition of 10.18 ml of triethylamine (73.02 mmol) and 5 6.46 g of 4,6-dichloro-2-methylsulfanylpirimidine-5-carbonyl chloride (25.08 mmol) predissolved in 50 ml of THF. After stirring for 3 hours at -70°C , the reaction mixture is poured into ice-water and then extracted 4 times with 150 ml of DCM. The organic phases are combined and washed with 200 ml of water and then with 200 ml of brine. The organic phase is dried over Na_2SO_4 , filtered and evaporated. The residue is 10 chromatographed on silica gel with a gradient of from 6% to 60% ethyl acetate in heptane to give 10.56 g of (S)-2-[[[(4,6-dichloro-2-methylsulfanylpirimidine-5-carbonyl)-(2-hydroxy-2-methylpropyl)amino]methyl]pyrrolidine-1-carboxylic acid tert-butyl ester. LC/MS (method F): $\text{M}+\text{H}^+ = 493$; $\text{Tr} = 4.57$ min

Step 4.3: (which corresponds to step b))


15 **(S)-2-(4-Chloro-8,8-dimethyl-2-methylsulfanyl-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-ylmethyl)pyrrolidine-1-carboxylic acid tert-butyl ester**



10.56 g (21.40 mmol) of the compound described in step **4.2** are placed in 107 ml of DMSO. 3.55 ml of DBU (23.54 mmol) are added dropwise and the reaction mixture is 20 stirred at 55°C for 5 hours. The medium is then poured into water and extracted with DCM. The organic phase is washed with brine, dried over MgSO_4 , filtered and evaporated. The residue is chromatographed on silica gel with a gradient of from 5% to 50% ethyl acetate in heptane to give 3.55 g of (S)-2-(4-chloro-8,8-dimethyl-2-methylsulfanyl-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-ylmethyl)pyrrolidine-1-carboxylic 25 acid tert-butyl ester. LC/MS (method F): $\text{M}+\text{H}^+ = 457$; $\text{Tr} = 4.47$ min

Step 4.4: (which corresponds to step o))

(S)-2-{4-[4-(3-Ethylureido)-3-fluorophenyl]-8,8-dimethyl-2-methylsulfanyl-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-ylmethyl}pyrrolidine-1-carboxylic acid tert-butyl ester



5


1 g (2.19 mmol) of the compound described in step **4.3** is placed in 10 ml of dioxane in a 20 ml microwave tube. 809 mg (2.63 mmol) of 1-ethyl-3-[2-fluoro-4-(4,4,5,5-tetramethyl-[1,3,2] dioxaborolan-2-yl)phenyl]urea and 6.02 ml of 2 N Na₂CO₃ solution (12.04 mmol) are successively added. The solution is degassed with argon, followed by addition of 126 mg of tetrakis(triphenylphosphine)palladium (109.41 μmol). The tube is heated at 110°C for 30 minutes in a Biotage microwave machine. The medium is poured into 300 ml of water and extracted twice with 300 ml of ethyl acetate. The organic phases are combined, washed with water and then with brine, dried over MgSO₄, filtered and evaporated. The residue is chromatographed on silica gel with a gradient of ethyl acetate in heptane ranging from 7% to 70% to give 1.9 g of (S)-2-{4-[4-(3-ethylureido)-3-fluorophenyl]-8,8-dimethyl-2-methylsulfanyl-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-ylmethyl}pyrrolidine-1-carboxylic acid tert-butyl ester. LC/MS (method F): M+H⁺ = 603; Tr = 4.42 min

Step 4.5: (which corresponds to step p))

(S)-2-{4-[4-(3-Ethylureido)-3-fluorophenyl]-2-methanesulfonyl-8,8-dimethyl-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-ylmethyl}pyrrolidine-1-carboxylic acid tert-butyl ester

20


77



- 1.9 g (3.15 mmol) of the compound described in step 4.4 and 3.9 g of magnesium monoperoxyphthalate hexahydrate (6.3 mmol) are placed in 25 ml of a 3/2 acetonitrile/EtOH mixture. After stirring for 3 hours at room temperature, the medium is poured into water and extracted 3 times with DCM. The organic phases are combined, washed with water and then with brine, dried over MgSO_4 , filtered and evaporated. 2.5 g of (S)-2-{4-[4-(3-ethylureido)-3-fluorophenyl]-2-methanesulfonyl-8,8-dimethyl-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-ylmethyl}pyrrolidine-1-carboxylic acid tert-butyl ester are obtained.
- 10 LC/MS (method F): $\text{M}+\text{H}^+ = 635$; Tr = 4.08 min

Step 4.6: (which corresponds to step q))

(S)-2-[4-[4-(3-Ethylureido)-3-fluorophenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-ylmethyl]pyrrolidine-1-carboxylic acid tert-butyl ester



15


Half of 1.93 g (3.04 mmol) of the compound described in step 4.5 is placed in half of 26 ml of DMSO in two 20 ml microwave tubes, respectively. Half of each amount of reagents of the 1 g of 8-oxa-3-azadicyclo[3.2.1]octane hydrochloride (6.35 mmol) and 1.78 ml of triethylamine (12.68 mmol) are successively and respectively added. The

tubes are heated for 30 minutes at 80°C in a Biotage microwave machine. The media are poured into water and extracted 3 times with DCM. The organic phases are combined, washed with brine, dried over MgSO_4 , filtered and evaporated. The residue is chromatographed on silica gel with a gradient of MeOH in DCM ranging from 1% to 10% to give 1.46 g of (S)-2-[4-[4-(3-ethylureido)-3-fluorophenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-ylmethyl]pyrrolidine-1-carboxylic acid tert-butyl ester.

LC/MS (method F): $\text{M}+\text{H}^+ = 668$; Tr = 4.37 min

Step 4.7: (deprotection)

- 10 **1-{4-[8,8-Dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(S)-1-pyrrolidin-2-ylmethyl-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-ethylurea**




- 15 1.17 g (1.75 mmol) of the compound described in step 4.6 are placed in 20 ml of DCM. 4 ml of TFA (51.92 mmol) are added dropwise. After stirring for 30 minutes at room temperature, the reaction medium is concentrated to dryness. The residue is diluted in DCM and washed with saturated aqueous NaHCO_3 solution and then with brine. The organic phase is dried over MgSO_4 , filtered and evaporated to give 1.15 g of 1-{4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(S)-1-pyrrolidin-2-ylmethyl-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-ethylurea.

LC/MS (method A): $\text{M}+\text{H}^+ = 568$; Tr = 0.76 min

- 25 ^1H NMR (400 MHz, DMSO-d_6) δ ppm 8.43 (m, 1H); 8.14 (dd, 1H); 7.43 (dd, 1H); 7.35 (dd, 1H); 6.64 (t, 1H); 4.41 (m, 2H); 4.29 (d, 2H); 3.75 (d, 2H); 3.69-3.43 (m, 3H); 3.21-3.12 (m, 4H); 3.17-3.08 (m, 2H); 1.95-1.69 (m, 8H); 1.51 (m, 1H); 1.39 (d, 6H); 1.11 (t, 3H)

Example 5:

Synthesis of (S)-2-[4-[4-(3-ethylureido)-3-fluorophenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-ylmethyl]pyrrolidine-1-carboxylic acid ethyl ester



5

400 mg (704.65 μmol) of 1-{4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(S)-1-pyrrolidin-2-ylmethyl-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-ethylurea are placed in 5 ml of DCM. 0.15 ml of triethylamine (1.06 mmol) and 5 mg of 4-dimethylaminopyridine (40.93 μmol) are successively added. The reaction mixture is cooled in an ice bath to 4°C and 89 μL of ethyl chloroformate (916.05 μmol) are added dropwise. The ice bath is removed and, after 4 hours, the reaction medium is poured into 100 ml of saturated aqueous NaHCO_3 solution. The organic phase is extracted, washed with water, with brine, dried over MgSO_4 , filtered and evaporated. The residue is chromatographed on silica gel with a gradient of a 5/1 ethyl acetate/MeOH mixture in DCM ranging from 0% to 45%. The solid obtained is taken up in Et_2O and filtered off to give 310 mg of (S)-2-[4-[4-(3-ethylureido)-3-fluorophenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-ylmethyl]pyrrolidine-1-carboxylic acid ethyl ester. LC/MS (method A): $\text{M}+\text{H}^+ = 640$; $\text{Tr} = 1.08$ min.


^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 8.43 (m, 1H); 8.15 (m, 1H); 7.39 (dd, 2H); 7.30 (dd, 1H); 6.63 (q, 1H); 4.40 (m, 2H); 4.08-3.99 (m, 4H); 3.77-3.59 (m, 3H); 3.41-3.31 (m, 3H); 3.17-3.08 (m, 4H); 1.90-1.81 (m, 6H); 1.65 (m, 2H); 1.35 (d, 6H); 1.19 (t, 3H); 1.06 (t, 3H)

Example 6:

Synthesis of 1-{4-[6-((S)-1-acetylpyrrolidin-2-ylmethyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-ethylurea

25

80




- 400 mg (704.65 μmol) of 1-{4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(S)-1-pyrrolidin-2-ylmethyl-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-ethylurea are placed in 5 ml of DCM. 0.19 ml of triethylamine (1.41 mmol) and 0.1 ml of acetic anhydride (1.06 mmol) are successively added dropwise. After stirring for 3 hours at room temperature, the reaction medium is poured into saturated aqueous NaHCO_3 solution and extracted twice with DCM. The organic phases are combined, washed with water and then with brine, dried over MgSO_4 , filtered and evaporated. The residue is chromatographed on silica gel with a gradient of a 5/1 ethyl acetate/MeOH mixture in DCM ranging from 0% to 45%. The solid obtained is taken up in Et_2O and filtered off to give 277 mg of 1-{4-[6-((S)-1-acetylpyrrolidin-2-ylmethyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-ethylurea. LC/MS (method A): $\text{M}+\text{H}^+ = 610$; Tr = 0.91 min
- ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 8.10-8.04 (m, 2H); 7.41-7.30 (m, 2H); 6.46 (m, 1H); 4.38 (m, 2H); 4.29-4.22 (m, 2H); 3.38 (m, 3H); 3.43 (m, 3H); 3.14-3.10 (m, 5H); 1.97-1.81 (m, 9H); 1.68 (m, 2H); 1.36 (d, 6H); 1.08 (t, 3H)

Example 7:


- Synthesis of 3-[4-[4-(3-ethylureido)phenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl]propionic acid**

81



Step 7.1: (preparation of a compound of formula (XI))


3-(2-Hydroxy-2-methylpropylamino)propionic acid tert-butyl ester



5 1 g of 1-aminomethylpropan-2-ol (11.22 mmol) is placed in 10 ml of DMF. 1.58 ml of triethylamine (11.22 mmol) and 1.87 ml of tert-butyl 3-bromopropionate (11.22 mmol) prediluted in 5 ml of DMF are successively added dropwise. After stirring for 18 hours at room temperature, the solvent is evaporated off. The residue is chromatographed on silica gel with a gradient of from 0% to 20% MeOH in DCM to give 2 g of 3-(2-hydroxy-2-
10 methylpropylamino)propionic acid tert-butyl ester.

Step 7.2: (which corresponds to step a))

3-[(4,6-Dichloro-2-methylsulfanylpuridine-5-carbonyl)-(2-hydroxy-2-methylpropyl)amino]propionic acid tert-butyl ester



15 1 g (4.58 mmol) of the compound described in step 7.1 is placed in 5 ml of anhydrous THF. 0.35 ml of triethylamine (4.58 mmol) is added and the mixture is cooled to -70°C in a bath of cardice in acetone. 1.18 g of 4,6-dichloro-2-methylsulfanylpuridine-5-carbonyl chloride (4.58 mmol) prediluted in 10 ml of anhydrous THF are then added dropwise. After stirring for 2 hours at -70°C, 10 ml of


water are added and the reaction medium is extracted with DCM. The organic phase is washed with brine, dried and concentrated to dryness. The residue is chromatographed on silica gel with a gradient of ethyl acetate in DCM ranging from 0% to 10% to give 1.3 g of

3-[(4,6-dichloro-2-methylsulfanylpyrimidine-5-carbonyl)-(2-hydroxy-2-methylpropyl)amino]propionic acid tert-butyl ester.

LC/MS (method F): $M+H^+ = 438$; Tr = 4.34 min

Step 7.3: (which corresponds to step b))

3-(4-Chloro-8,8-dimethyl-2-methylsulfanyl-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl)propionic acid tert-butyl ester




10

800 mg (1.82 mmol) of the compound described in step 7.2 are placed in 10 ml of DMSO. 0.3 ml of DBU (2.01 mmol) is added and the reaction mixture is stirred at 55°C for 2 hours. The reaction medium is poured into water and extracted with ethyl acetate. The organic phase is washed with brine, dried over $MgSO_4$, filtered and evaporated. The residue is chromatographed on silica gel with a gradient of ethyl acetate in DCM ranging from 0% to 10% to give 460 mg of 3-(4-chloro-8,8-dimethyl-2-methylsulfanyl-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl)propionic acid tert-butyl ester.

LC/MS (method F): $M+H^+ = 402m$; Tr = 4.32 min

Step 7.4: (which corresponds to step o))

20 3-{4-[4-(3-Ethylureido)phenyl]-8,8-dimethyl-2-methylsulfanyl-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl}propionic acid tert-butyl ester




410 mg (1.02 mmol) of the compound described in step **7.3** are placed in 15 ml of dioxane in a 20 ml microwave tube. 350 mg (1.22 mmol) of 1-ethyl-3-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]urea and 2.8 ml of 2N Na₂CO₃ solution (5.61 mmol) are successively added. The solution is degassed with argon, followed by addition
 5 of 60 mg of tetrakis(triphenylphosphine)palladium (51 μmol). The tube is heated at 110°C for 30 minutes in a Biotage microwave machine. The medium is poured into 75 ml of water and the solution is filtered. The solid is rinsed with acetonitrile and then with pentane to give 500 mg of 3-{4-[4-(3-ethylureido)phenyl]-8,8-dimethyl-2-methylsulfonyl-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl}propionic acid tert-butyl
 10 ester.

LC/MS (method F): M+H⁺ = 603; Tr = 4.42 min

Step 7.5: (which corresponds to step p))

3-{4-[4-(3-Ethylureido)phenyl]-2-methanesulfonyl-8,8-dimethyl-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl}propionic acid tert-butyl ester



15

450 mg (0.85 mmol) of the compound described in step **7.4** and 1.05 g of magnesium monoperoxyphthalate hexahydrate (1.7 mmol) are placed in 6 ml of a 2/1 acetonitrile/EtOH mixture. After stirring for 18 hours at room temperature, the solvents are evaporated off. The residue is taken up in 20 ml of water. The product obtained is
 20 filtered off and rinsed with acetonitrile, with pentane and then with Et₂O to give 410 mg of 3-{4-[4-(3-ethylureido)phenyl]-8,8-dimethyl-2-methanesulfonyl-8,8-dimethyl-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl}propionic acid tert-butyl ester.


LC/MS (method F): M+H⁺ = 562; Tr = 3.81 min

Step 7.6: (which corresponds to step q))

3-[4-[4-(3-Ethylureido)phenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl]propionic acid tert-butyl ester

25

84




410 mg (0.73 mmol) of the compound described in step **7.5** are placed in 10 ml of DMSO in a 20 ml microwave tube. 345 mg of 8-oxa-3-azabicyclo[3.2.1]octane hydrochloride (2.19 mmol) and 0.6 ml of triethylamine (4.38 mmol) are successively added. The tube is heated at 80°C for 45 minutes in a Biotage microwave machine. The medium is poured into water and extracted twice with DCM. The organic phases are combined, washed with brine, dried over MgSO_4 , filtered and evaporated. The solid is rinsed with acetonitrile and with pentane to give 420 mg of 3-[4-[4-(3-ethylureido)phenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl]propionic acid tert-butyl ester.

LC/MS (method A): $\text{M}+\text{H}^+ = 595$; $\text{Tr} = 1.14$ min

Step 7.7: (deprotection))

3-[4-[4-(3-Ethylureido)phenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl]propionic acid



15

300 mg (0.5 mmol) of the compound described in step **7.6** are placed in 5 ml of DCM. 0.5 ml of a 4M solution of hydrogen chloride in dioxane (2 mmol) is added slowly to


the medium. After stirring for 3 hours at room temperature, the solvent is evaporated off and the residue is chromatographed on silica gel with a gradient of MeOH containing 10% aqueous ammonia in DCM ranging from 0% to 10%. 130 mg of 3-[4-[4-(3-ethylureido)phenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl]propionic acid are obtained.

LC/MS (method A): $M+H^+$ = 539; Tr = 0.86 min

^1H NMR (400 MHz, DMSO- d_6) δ ppm 9.48 (m, 1H); 8.53 (m, 1H); 7.45-7.35 (m, 4H); 6.15 (t, 1H); 4.38 (m, 2H); 4.26 (m, 2H); 3.75-3.63 (m, 4H); 3.15-3.03 (m, 4H); 2.55 (t, 2H); 1.81 (m, 2H); 1.66 (m, 2H); 1.35 (s, 6H); 1.06 (t, 3H)

10 **Example 8:**

Synthesis of 3-[4-[4-(3-ethylureido)phenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl]-N,N-dimethylpropionamide




15 70 mg (0.13 mmol) of 3-[4-[4-(3-ethylureido)phenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl]propionic acid are placed in 3 ml of DCM. 0.66 ml of DIEA (0.39 mmol), 12 mg of dimethylamine hydrochloride (0.14 mmol), 21 mg of hydroxybenzotriazole (0.14 mmol) and 27 mg of 1-ethyl-3-(dimethylaminopropyl)carbodiimide hydrochloride (0.14 mmol) are
20 successively added, with stirring. After stirring for 3 days at room temperature, the DCM is evaporated off. The residue is taken up in water and extracted with ethyl acetate. The organic phase is washed with saturated aqueous $NaHCO_3$ solution and then with brine, dried over $MgSO_4$, filtered and evaporated. The solid obtained is taken up in acetonitrile, filtered off and rinsed with pentane to give 34 mg of 3-[4-[4-(3-ethylureido)phenyl]-8,8-
25 dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl]-N,N-dimethylpropionamide.

LC/MS (method A): $M+H^+$ = 566; Tr = 0.86 min

^1H NMR (400 MHz, DMSO- d_6) δ ppm 8.64 (m, 1H); 7.38 (m, 4H); 6.15 (t, 1H); 4.40 (m, 2H); 4.23 (m, 2H); 3.72 (s, 2H); 3.66 (m, 2H); 3.15-3.05 (m, 4H); 3.01 (s, 3H); 2.88 (s, 3H); 2.68 (m, 2H); 1.81 (m, 2H); 1.65 (m, 2H); 1.34 (s, 6H); 1.05 (t, 3H)

5 **Example 9:**

Synthesis of 1-ethyl-3-{4-[6-(3-hydroxypropyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea




10 13 mg of lithium aluminium hydride (0.34 mmol) are placed in 5 ml of anhydrous THF. A solution containing 90 mg (0.17 mmol) of 3-[4-[4-(3-ethylureido)phenyl]-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-7,8-dihydro-5H-9-oxa-1,3,6-triazabenzocyclohepten-6-yl]propionic acid in 10 ml of THF is added dropwise. The reaction mixture is heated at 66°C for 18 hours. 72 mg of lithium aluminium hydride (1.89
15 mmol) are added and the mixture is heated for a further 24 hours at 66°C. The medium is then cooled in an ice-water bath to 4°C and basified with saturated aqueous Na_2SO_4 solution. Ethyl acetate is added and the mixture is filtered through Celite, rinsed with ethyl acetate and concentrated. The residue is chromatographed on silica gel with a gradient of MeOH containing 10% aqueous ammonia DCM ranging from 0% to 10%. The solid is
20 rinsed with acetonitrile and then with pentane to give 22 mg of 1-ethyl-3-{4-[6-(3-hydroxypropyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea.

LC/MS (method A): $M+H^+$ = 525; Tr = 0.82 min


^1H NMR (400 MHz, DMSO- d_6) δ ppm 8.29 (m, 1H); 7.44-7.34 (m, 4H); 5.95 (t, 1H); 4.38
25 (m, 2H); 4.27 (d, 2H); 3.62 (s, 2H); 3.58-3.52 (m, 4H); 3.15-3.09 (m, 5H); 1.83-1.66 (m, 6H); 1.36 (s, 6H); 1.07 (t, 3H)

Example 10:

Synthesis of 1-ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6,7,8,9-tetrahydro-5H-10-oxa-1,3,6-triazabenzocycloocten-4-yl]phenyl}urea

5 **Step 10.1:**


3-(tert-Butyldimethylsilyloxy)-2,2-dimethylpropylamine



2 g of 3-amino-2,2-dimethylpropan-1-ol (19.39 mmol) are placed in 40 ml of DCM. 2.56 g of imidazole (38.77 mmol) are added and the solution is cooled to +4°C in an ice bath, followed by dropwise addition of 3.36 ml of tert-butyl(chloro)dimethylsilane (19.39 mmol). The medium is stirred for 30 minutes at +4°C, and the ice bath is then removed and the mixture is stirred at room temperature for 18 hours. The medium is then washed with water, and then with brine. The organic phase is dried over Na₂SO₄, filtered and concentrated to give 4.02 g of 3-(tert-butyldimethylsilyloxy)-2,2-dimethylpropylamine.

15 **Step 10.2:** (preparation of a compound of formula (XI) whose hydroxyl function is protected)

[3-(tert-Butyldimethylsilyloxy)-2,2-dimethylpropyl]ethylamine




4.02 g (18.49 mmol) of the compound described in step 10.1 are placed in 40 ml of anhydrous THF. 0.81 g of sodium hydride (20.34 mmol) is added portionwise. After 30 minutes, 1.63 ml of iodoethane (20.34 mmol) are added and the mixture is heated at 55°C for 18 hours. The reaction medium is poured into water and extracted with ethyl acetate. The organic phase is washed with brine, dried over Na₂SO₄, filtered and

concentrated to give 4.47 g of [3-(tert-butyldimethylsilyloxy)-2,2-dimethylpropyl]ethylamine.

Step 10.3: (which corresponds to step a), with the hydroxyl function protected)

**4,6-Dichloro-2-methylsulfanylpurimidine-5-carboxylic acid [3-(tert-butyl-
5 dimethylsilyloxy)-2,2-dimethylpropyl]ethylamide**




4.47 g (18.21 mmol) of the compound described in step **10.2** are placed in 400 ml of THF. 7.57 ml of triethylamine (5.53 mmol) are added and the mixture is cooled to -70°C in a bath of cardice in acetone. 4.69 g of 4,6-dichloro-2-methylsulfanylpurimidine-5-carbonyl chloride (18.21 mmol) prediluted in 20 ml of anhydrous THF are then added dropwise. After stirring for 4 hours at -70°C, aqueous 1N hydrochloric acid solution is added and the reaction medium is extracted with ethyl acetate, dried over Na₂SO₄ and concentrated to dryness. The residue is taken up in acetonitrile and filtered. The filtrate is evaporated to give 4.1 g of 4,6-dichloro-2-methylsulfanylpurimidine-5-carboxylic acid [3-(tert-butyl-5 dimethylsilyloxy)-2,2-dimethylpropyl]ethylamide.

LC/MS (method F): M+H⁺ = 466; Tr = 5.77 min

Step 10.4: (deprotection of the alcohol function)

4,6-Dichloro-2-methylsulfanylpurimidine-5-carboxylic acid ethyl(3-hydroxy-2,2-dimethylpropyl)amide




20

4.1 g (8.79 mmol) of the compound described in step **10.3** are placed in 40 ml of MeOH. 1.83 ml of 12N hydrochloric acid (21.98 mmol) are added. After 18 hours, the MeOH is evaporated off. The residue is taken up in ethyl acetate and washed with saturated NaHCO₃ solution. The organic phase is dried over Na₂SO₄, filtered and concentrated to give 3.1 g of 4,6-dichloro-2-methylsulfanylpurimidine-5-carboxylic acid ethyl(3-hydroxy-2,2-dimethylpropyl)amide.

LC/MS (method F): M+H⁺ = 352; Tr = 4.08 min

Step 10.5: (which corresponds to step b))

4-Chloro-6-ethyl-8,8-dimethyl-2-methylsulfanyl-6,7,8,9-tetrahydro-10-oxa-1,3,6-triazabenzocycloocten-5-one




2.95 g (8.37 mmol) of the compound described in step **10.4** are placed in 55 ml of DMSO with stirring. 1.38 ml of DBU (9.21 mmol) are added. The reaction medium is stirred for 3 hours at 55°C, and 40 ml of ice-water are then added. The medium is extracted with ethyl acetate, washed with brine, dried over Na₂SO₄ and then filtered. The solution is evaporated to dryness. The residue is then chromatographed on silica gel with a gradient of ethyl acetate in DCM ranging from 0% to 10%. 1.18 g of 4-chloro-6-ethyl-8,8-dimethyl-2-methylsulfanyl-6,7,8,9-tetrahydro-10-oxa-1,3,6-triazabenzocycloocten-5-one are obtained.

LC/MS (method F): M+H⁺ = 316; Tr = 3.91 min

Step 10.6: (which corresponds to step o))

1-Ethyl-3-[4-(6-ethyl-8,8-dimethyl-2-methylsulfanyl-5-oxo-6,7,8,9-tetrahydro-5H-10-oxa-1,3,6-triazabenzocycloocten-4-yl)phenyl]urea




0.3 g (0.95 mmol) of the compound described in step **10.5** is placed in 8 ml of dioxane in a 20 ml microwave tube. 330 mg (1.14 mmol) of 1-ethyl-3-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]urea and 2.61 ml of 2N Na₂CO₃ solution (5.22 mmol) are successively added. The solution is degassed with argon, followed by addition of 55 mg of tetrakis(triphenylphosphine)palladium (50 μmol). The tube is heated at 110°C for 30 minutes in a Biotage microwave machine. The medium is poured into water and filtered to give 330 mg of 1-ethyl-3-[4-(6-ethyl-8,8-dimethyl-2-methylsulfanyl-5-oxo-6,7,8,9-tetrahydro-5H-10-oxa-1,3,6-triazabenzocycloocten-4-yl)phenyl]urea.

LC/MS (method F): M+H⁺ = 444; Tr = 3.70 min

Step 10.7: (which corresponds to step p))

1-Ethyl-3-[4-(6-ethyl-2-methanesulfonyl-8,8-dimethyl-5-oxo-6,7,8,9-tetrahydro-5H-10-oxa-1,3,6-triazabenzocycloocten-4-yl)phenyl]urea




- 5 0.292 g (0.66 mmol) of the compound described in step **10.6** and 0.65 g of magnesium monoperoxyphthalate hexahydrate (1.32 mmol) are placed in 20 ml of a 2/1 acetonitrile/EtOH mixture. After stirring for 4 hours at room temperature, the solvents are evaporated off. The residue is taken up in water. The product obtained is filtered off and rinsed twice with water to give 0.25 g of 1-ethyl-3-[4-(6-ethyl-2-methanesulfonyl-8,8-
- 10 dimethyl-5-oxo-6,7,8,9-tetrahydro-5H-10-oxa-1,3,6-triazabenzocycloocten-4-yl)phenyl]urea.

LC/MS (method F): $M+H^+ = 476$; Tr = 3.34 min

Step 10.8: (which corresponds to step q))

- 1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6,7,8,9-**
- 15 **tetrahydro-5H-10-oxa-1,3,6-triazabenzocycloocten-4-yl]phenyl}urea**



410 mg (0.26 mmol) of the compound described in step **10.7** are placed in 7 ml of DMSO in a 20 ml microwave tube. 118 mg of 8-oxa-3-azabicyclo[3.2.1]octane hydrochloride (0.79 mmol) and 0.22 ml of triethylamine (1.58 mmol) are successively


added. The tube is heated at 100°C for 1 hour 30 minutes in a Biotage microwave machine. The medium is poured into water and extracted twice with ethyl acetate. The organic phases are combined, washed with brine, dried over MgSO_4 , filtered and evaporated. The residue is chromatographed on silica gel with a gradient of MeOH containing 10% aqueous ammonia in DCM to give 48 mg of 1-ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6,7,8,9-tetrahydro-5H-10-oxa-1,3,6-triazabenzocycloocten-4-yl]phenyl}urea.

LC/MS (method A): $\text{M}+\text{H}^+ = 509$; $\text{Tr} = 0.96$ min

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 8.60 (m, 1H); 7.36 (m, 4H); 6.13 (t, 1H); 4.39-3.92 (m, 5H); 3.72-3.62 (m, 2H); 3.14-3.03 (m, 5H); 2.93 (m, 1H); 1.81-1.61 (m, 5H); 1.1-1.03 (m, 9H); 0.93 (m, 3H)

Example 11:


Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-methylurea



15

Step 11.1: (which corresponds to step c))


[4-(6-Ethyl-8,8-dimethyl-2-methylsulfanyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)-3-fluorophenyl]carbamic acid tert-butyl ester



1.77 g (5.85 mmol) of 4-chloro-6-ethyl-8,8-dimethyl-2-methylsulfanyl-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one, 2.27 g (6.73 mmol) of [3-fluoro-4-(4,4,5,5-tetramethyl[1,3,2]dioxaborolan-2-yl)phenyl]carbamic acid tert-butyl ester and 14.63 ml of a 2 M solution of Na_2CO_3 in 35 ml dioxane are mixed together, followed by degassing 5 times with argon. 339 mg (293 μmol) of tetrakis(triphenylphosphine)palladium (0) are added and the mixture is degassed a further 3 times with argon and then heated at 90°C for 2 hours 30 minutes. The resulting mixture is allowed to cool. It is poured into water. The product is filtered off, washed with water, dissolved in DCM, dried and concentrated. It is taken up in Et_2O , and the mixture is stirred, filtered and dried. 2.94 g of [4-(6-ethyl-8,8-dimethyl-2-methylsulfanyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)-3-fluorophenyl]carbamic acid tert-butyl ester are recovered. LC/MS (method E): $\text{M}+\text{H}^+ = 477$; $\text{Tr} = 2.60$ min

Step 11.2: (which corresponds to step d))


[4-(6-Ethyl-2-methanesulfonyl-8,8-dimethyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)-3-fluorophenyl]carbamic acid tert-butyl ester



2.79 g (5.85 mmol) of the compound described in step 11.1 are suspended in 30 ml of acetonitrile and 10 ml of EtOH , followed by addition of 7.23 g of magnesium monoperoxyphthalate hexahydrate (11.70 mmol). The mixture is stirred at 60°C for 2 hours and is then stirred at room temperature overnight. 1 equivalent of magnesium monoperoxyphthalate hexahydrate is added. The mixture is heated at 60°C for 2 hours. It is allowed to cool, diluted with water, stirred for 10 minutes and then filtered. The product is washed with water, dissolved in DCM, dried and concentrated to give 3.05 g of [4-(6-ethyl-2-methanesulfonyl-8,8-dimethyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl)-3-fluorophenyl]carbamic acid tert-butyl ester. LC/MS (method E): $\text{M}+\text{H}^+ = 509$; $\text{Tr} = 2.20$ min

Step 11.3: (which corresponds to step e))


{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}carbamic acid tert-butyl ester



- 5 To 1.02 g (2.0 mmol) of the compound described in step **11.2** suspended in 5 ml of anhydrous dioxane under argon are added 2.02 g (20 mmol) of (S)-3-methylmorpholine, followed by heating at 95°C for 3 hours. The mixture is allowed to cool and is partly concentrated, water is added and the resulting mixture is then filtered. The residue is taken up in DCM and the solution is dried and concentrated. The solid
- 10 obtained (1 g) is used without purification in the following step.
LC/MS (method E): $M+H^+ = 530$; Tr = 2.64 min

Step 11.4: (which corresponds to step I))

4-(4-Amino-2-fluorophenyl)-6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one



15


The compound described in step **11.3** is suspended in 5 ml of DCM and 500 μ L of water. 10 ml of TFA are added. The mixture is reacted at room temperature for 2 hours. The resulting mixture is concentrated, taken up in a DCM/water mixture and basified with K_2CO_3 . The phases are separated by settling and the organic phase is dried and

concentrated. The solid is taken up in Et₂O. The product is stirred, filtered off and air-dried to give 690 mg of 4-(4-amino-2-fluorophenyl)-6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one.

LC/MS (method E): M+H⁺ = 430; Tr = 2.04 min

5 **Step 11.5:** (which corresponds to step m) (steps m1-1) and m1-2))

1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-methylurea



200 mg (466 μmol) of the compound described in step **11.4** are suspended in 5
 10 ml of anhydrous dioxane under argon. The mixture is heated to dissolve the suspension. The resulting mixture is allowed to cool and 123 μl (699 μmol) of DIEA are added, followed by 368 μl (699 μmol) of a 1.9M solution of phosgene in toluene, after which the mixture is heated at 50°C for 30 minutes. It is allowed to cool and 1.63 ml of 2M methylamine in THF (3.26 mmol) are added. The resulting mixture is reacted overnight.
 15 is poured into water and filtered. The product is dissolved in DCM, dried over Na₂SO₄ and then concentrated under reduced pressure. The resulting product is taken up in Et₂O, stirred, filtered off and dried to give 191 mg of 1-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-methylurea.


20 LC/MS (method C): M+H⁺ = 487; Tr = 0.64 min

¹H NMR (600 MHz, DMSO-d₆) δ ppm 8.95 (s, 1H); 7.49 (d, 1H); 7.45 (d, 1H); 7.05 (d, 1H); 6.17 (s, 1H); 4.63 (s, 1H); 4.29 (d, 1H); 3.90 (dd, 1H); 3.70 (d, 1H); 3.57 (dd, 1H); 3.52 (s, 2H); 3.36-3.50 (bs, 3H); 3.18 (bs, 1H); 2.65 (d, 3H); 1.35 (d, 6H); 1.21 (d, 3H); 1.08 (t, 3H)

25 **Example 12:**

Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-

tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl]-3-pyridin-4-ylurea (which corresponds to step m) (steps m1-1) and m1-2))




150 mg (394 μ mol) of 4-(4-amino-2-fluorophenyl)-6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one are suspended in 4 ml anhydrous dioxane under argon. 92 μ l (524 μ mol) of DIEA are added, followed by 276 μ l of a 1.9M solution of phosgene in toluene, after which the mixture is heated at 50°C for 30 minutes. The resulting mixture is allowed to cool, followed by addition of 202 mg (2.10 mmol) of 4-aminopyridine dissolved in 2 ml of anhydrous DMF. The resulting mixture is reacted for 3 hours. It is poured into water, stirred, filtered and washed with water. The product is dissolved in DCM, dried over Na₂SO₄ and then concentrated under reduced pressure. It is taken up in Et₂O, and the mixture is stirred and filtered. The product is dissolved in 5 ml of DCM with a small amount of MeOH, and 20 ml of ethyl acetate are added. The product is placed in a refrigerator and the solid is then filtered off to give 146 mg of 1-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl]-3-pyridin-4-ylurea.

LC/MS (method C): M+H⁺ = 550; Tr = 0.57 min

¹H NMR (600 MHz, DMSO-d₆) δ ppm 9.30 (s, 2H); 8.38 (d, 2H); 7.57 (t, 1H); 7.48 (dd, 1H); 7.45 (d, 2H); 7.17 (dd, 1H); 4.64 (s, 1H); 4.30 (d, 1H); 3.91 (dd, 1H); 3.71 (d, 1H); 3.58 (dd, 1H); 3.54 (s, 2H); 3.39-3.53 (m, 3H); 3.19 (dt, 1H); 1.36 (d, 6H); 1.22 (d, 3H); 1.09 (t, 3H)

Example 13:

Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl]-3-[4-(4-methylpiperazin-1-yl)phenyl]urea (which corresponds to step m) (steps m2-1) and m2-2))




136.5 mg (460 μ mol) of trisphosgene are dissolved in 2 ml of DCM. A mixture of 90.7 mg (460 μ mol) of 4-(4-methylpiperazin-1-yl)phenylamine and 389 μ l (2.76 mmol) of triethylamine in 3 ml of DCM is added in a single portion. The resulting mixture is stirred for 30 minutes at room temperature. 100 mg (226.5 μ mol) of 4-(4-amino-3-fluorophenyl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one are added. The resulting mixture is reacted for 15 hours at room temperature. The reaction mixture is diluted in 50 ml of DCM and the organic phase is washed with twice 20 ml of saturated NaHCO_3 solution. The organic phase is dried over Na_2SO_4 , filtered and evaporated. The residue is chromatographed on silica gel, eluting with an MeOH/DCM mixture (0/100 v/v) up to (10/90 v/v) to give 75 mg of a white powder. The powder obtained is washed again with 2 ml of MeOH and then 2 ml of ethyl acetate. The resulting product is dried. 68 mg of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[4-(4-methylpiperazin-1-yl)phenyl]urea are recovered.

LC/MS (method C): $\text{M}+\text{H}^+ = 659$; $\text{Tr} = 0.53$ min

^1H NMR (600 MHz, $\text{DMSO}-d_6$) δ ppm 8.91 (1H); 8.67 (s, 1H); 8.19 (t, 1H); 7.42 (d, 1H); 7.30 (t, 3H); 6.91 (d, 2H); 4.43 (s, 2H); 4.1-4.3 (bs, 2H); 3.70 (s, 2H); 3.4-3.6 (bs, 2H); 3.11 (d, 2H); 3.06 (t, 4H); 2.46 (t, 4H); 2.22 (s, 3H); 1.8-1.9 (bs, 2H); 1.6-1.7 (bs, 2H); 1.36 (s, 6H); 1.17 (t, 3H)


Example 14:

Synthesis of 1-{4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-ethylurea



Step 14.1: (preparation of a compound of formula (XI))


2-Methyl-1-((tetrahydro-2H-pyran-4-yl)amino)propan-2-ol



- 5 2.75 g of 1-amino-2-methylpropan-2-ol are dissolved in 20 ml of MeOH. 1.71 ml of glacial acetic acid are added dropwise under an inert atmosphere. 2 g of tetrahydropyran-4-one in 10 ml of MeOH are added, the mixture is stirred for 30 minutes at room temperature and 1.98 g of sodium cyanoborohydride are added. The resulting mixture is stirred under an inert atmosphere for 15 hours. The reaction medium is poured
- 10 into saturated aqueous NaHCO₃ solution (100 ml). The resulting mixture is stirred for 30 minutes and then extracted with 4 times 100 ml of DCM. The organic phase is dried over Na₂SO₄, filtered and evaporated without heating. The MeOH contained in the aqueous phase is evaporated off and the extraction, drying, filtration and evaporation operation is repeated 3 times. 2.8 g of 2-methyl-1-((tetrahydro-2H-pyran-4-yl)amino)propan-2-ol are
- 15 recovered in total.

Step 14.2(which corresponds to step a))

4,6-Dichloro-2-methylsulfanylpurimidine-5-carboxylic acid (2-hydroxy-2-methylpropyl)(tetrahydropyran-4-yl)amide




This compound is prepared according to the procedure described in step **1.2** starting with 4,6-dichloro-2-methylsulfanylpurimidine-5-carbonyl chloride and the compound described in step **14.1**.

LC/MS (method E): $M+H^+ = 395$; Tr = 2.13 min

5 **Step 14.3:** (which corresponds to step b))

4-Chloro-8,8-dimethyl-2-methylsulfanyl-6-(tetrahydropyran-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one




10 This compound is prepared according to the procedure described in step **1.3** starting with the compound described in step **14.2**.

LC/MS (method E): $M+H^+ = 360$; Tr = 1.90 min

Step 14.4: (which corresponds to step c))

{4-[8,8-Dimethyl-2-methylsulfanyl-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid tert-butyl ester



15


This compound is prepared according to the procedure described in step **1.4** starting with the compound described in step **14.3** and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]carbamic acid tert-butyl ester.

LC/MS (method E): $M+H^+ = 515$; Tr = 2.44 min

20 **Step 14.5:** (which corresponds to step d))

{4-[2-Methanesulfonyl-8,8-dimethyl-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-

tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid tert-butyl ester




- This compound is prepared according to the procedure described in step **1.5**
 5 starting with the compound described in step **14.4**.

LC/MS (method E): $M+H^+ = 547$; Tr = 2.03 min

Step 14.6: (which corresponds to step e))

- {4-[8,8-Dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic**
 10 **acid tert-butyl ester**




- This compound is prepared according to the procedure described in step **1.6**
 starting with the compound described in step **14.5** and 8-oxa-3-azabicyclo[3.2.1]octane.

LC/MS (method E): $M+H^+ = 580$; Tr = 2.43 min

- 15 **Step 14.7:** (which corresponds to step l))

4-(4-Aminophenyl)-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-6-
(tetrahydropyran-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one

100




This compound is prepared according to the procedure described in step **1.7** starting with the compound described in step **14.6**.

LC/MS (method E): $M+H^+ = 480$; Tr = 1.74 min

5 **Step 14.8:** (which corresponds to step m) (step m3-1))

{4-[8,8-Dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid phenyl ester




10 200 mg (417 μmol) of the compound described in step **14.7** are suspended in THF, and 152 μl (917.5 μmol) of DIEA are added, followed by addition of 66 mg (417 μmol) of phenyl chloroformate. The mixture is stirred at room temperature for 10 minutes. The solid is filtered off, rinsed with MeOH and dried. 171 mg of {4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-
15 triazabenzocyclohepten-4-yl]phenyl}carbamic acid phenyl ester are recovered.

LC/MS (method E): $M+H^+ = 600$; Tr = 2.40 min

Step 14.9: (which corresponds to step m) (step m3-2))

1-{4-[8,8-Dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-ethylurea




100 mg (166.8 μmol) of the compound described in step **14.8** suspended in 2 ml of DMF are placed in a microwave reactor, and 1 ml of a 2M solution of ethylamine (2.0 mmol) in THF is added. The mixture is heated in a Biotage microwave machine for 1 hour at 100°C. The solvents and the excess ethylamine are evaporated off. The residue is chromatographed on silica gel, eluting with an MeOH/DCM mixture from (0/100 v/v) up to (5/95 v/v) to give 66 mg of 1-{4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-ethylurea.

10 LC/MS (method C): $M+H^+ = 551$; Tr = 0.61 min

^1H NMR (600 MHz, DMSO- d_6) δ ppm 8.61 (s, 1H); 7.38 (dd, 4H); 6.14 (t, 1H); 4.41 (m, 4H); 4.0-4.3 (bs, 1H); 3.95 (dd, 2H); 3.61 (s, 2H); 3.38 (t, 2H); 3.09-3.14 (m, 4H); 1.85-1.94 (m, 2H); 1.78-1.84 (m, 2H); 1.60-1.70 (m, 4H); 1.34 (s, 6H); 1.07 (t, 3H)

Example 15: (which corresponds to step m) (step m3-2))


15 **Synthesis of 1-{4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methylisoxazol-3-yl)urea**



- 50 mg (83.38 μmol) of {4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid phenyl ester are placed in a microwave reactor and 25 mg (0.25 mmol) of 5-methylisoxazol-3-ylamine are added. 2 ml of DMF and 1 ml of THF are added. The reaction mixture is heated in a Biotage microwave machine for 30 minutes at 100°C. 100 mg (1.0 mmol) of 5-methylisoxazol-3-ylamine are added and the mixture is heated for 2 hours at 110°C. A further 300 mg (3 mmol) of 5-methylisoxazol-3-ylamine are added and the mixture is heated for 2 hours at 110°C. The solvents are evaporated off. The residue is taken up in water and the insoluble material is filtered off, washed three times with water, taken up in MeOH/DCM and dried. The residue is chromatographed on silica gel, eluting with an MeOH/DCM mixture from (0/100 v/v) up to (5/95 v/v) to give 6 mg of 1-{4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methylisoxazol-3-yl)urea.
- 15 LC/MS (method C): $M+H^+ = 604$; $T_r = 0.68$ min
- ^1H NMR (600 MHz, $\text{DMSO}-d_6$) δ ppm 9.52 (s, 1H); 9.03 (s, 1H); 7.44 (dd, 4H); 6.55 (s, 1H); 4.41 (m, 4H); 4.0-4.3 (bs, 1H); 3.95 (dd, 2H); 3.61 (s, 2H); 3.38 (t, 2H); 3.11 (d, 2H); 2.36 (s, 3H); 1.85-1.94 (m, 2H); 1.78-1.84 (m, 2H); 1.60-1.70 (m, 4H); 1.36 (s, 6H)

Example 16: (which corresponds to step m) (step m3-2))

- 20 **Synthesis of 1-{4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(1-methyl-1H-pyrazol-3-yl)urea**



- 50 mg (83.4 μmol) of {4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid phenyl ester are placed in a microwave reactor and 25 mg (26 μmol) of 1-methyl-1H-pyrazol-3-ylamine are added. 1 ml of DMF and 2 ml of THF are


added. The reactor is heated in a Biotage microwave machine for 30 minutes at 100°C. 100 mg (100 µmol) of 1-methyl-1H-pyrazol-3-ylamine are added and the mixture is heated for 2 hours at 110°C. The solvents are evaporated off. The residue is chromatographed on silica gel, eluting with an MeOH/DCM mixture from (0/100 v/v) up to 5 (5/95 v/v) to give a solid, which is then washed with water and dried. 23 mg of 1-{4-[8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-6-(tetrahydropyran-4-yl)-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(1-methyl-1H-pyrazol-3-yl)urea are recovered.

LC/MS (method C): $M+H^+ = 603$; Tr = 0.63 min

10 ^1H NMR (600 MHz, DMSO- d_6) δ ppm 9.12 (s, 1H); 8.93 (s, 1H); 7.54 (s, 1H); 7.44 (dd, 4H); 6.24 (s, 1H); 4.41 (m, 4H); 4.0-4.3 (bs, 1H); 3.95 (dd, 2H); 3.74 (s, 3H); 3.61 (s, 2H); 3.38 (t, 2H); 3.11 (d, 2H); 1.85-1.94 (m, 2H); 1.78-1.84 (m, 2H); 1.60-1.70 (m, 4H); 1.36 (s, 6H)


Example 17:

15 **Synthesis of 1-ethyl-3-{5-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]pyridin-2-yl}urea**



Step 17.1: (which corresponds to step k))

20 **4-(6-Aminopyridin-3-yl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one**




170 mg (353.83 μmol) of trifluoromethanesulfonic acid 6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl ester and 90.16 mg (389.21 μmol) of 5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)pyridin-2-ylamine are dissolved in 3 ml of dioxane in a microwave tube. 150.21 mg (707.66 μmol) of tripotassium phosphate and 1 ml of water are added. The solution is purged with argon. Finally, 24.84 mg (35.38 μmol) of bis(triphenylphosphine)palladium(II) chloride are added and the mixture is heated in a Biotage microwave machine for 15 minutes at 100°C. The reaction mixture is diluted in ethyl acetate and washed with water and then with saturated NaCl solution. The precipitate present in the organic phase is filtered off to give 120 mg of 4-(6-aminopyridin-3-yl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one.

^1H NMR (300 MHz, DMSO-d_6) δ ppm 8.08 (d, 1H); 7.58 (dd, 1H); 6.37-6.45 (m, 3H); 6.37-6.45 (m, 3H); 4.06-4.46 (bs, 4H); (bs, 4H); 3.42-3.68 (bs, 4H); 3.03-3.13 (bs, 2H); 1.73-1.86 (bs, 2H); 1.60-1.69 (bs, 2H); 1.33 (bs, 6H); 1.14 (t, 3H)

Step 17.2: (which corresponds to step m) (steps m3-1) and m3-2))


1-Ethyl-3-{5-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]pyridin-2-yl}urea



- 100 mg (235.57 μmol) of the compound described in step **17.1**, 30.52 μl (235.57 μmol) of phenyl chloroformate and 90.52 μl (518.26 μmol) of DIEA are placed in 2.5 ml of THF in a microwave tube. The resulting mixture is stirred for 30 minutes at room temperature. 235.57 μl (471.15 μmol) of 2N ethylamine are added in THF and the mixture is heated for 1 hour at 100°C in a Biotage microwave machine. The reaction medium is evaporated under reduced pressure. The residue is chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to (95/5 v/v) to give 56 mg of 1-ethyl-3-{5-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]pyridin-2-yl}urea.
- 10 LC/MS (method C): $M+H^+ = 496$; $T_r = 0.59$ min
 ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ ppm 9.39 (s, 1H); 8.34 (d, 1H); 7.92 (bs, 1H); 7.79 (dd, 1H); 7.41 (d, 1H); 4.05-4.46 (bs, 4H); 3.69 (bs, 2H); 3.51 (bs, 2H); 3.20 (m, 2H); 3.10 (bs, 2H); 1.82 (bs, 2H); 1.65 (bs, 2H); 1.35 (bs, 6H); 1.05-1.18 (m, 6H)

Example 18: (which corresponds to step m) (steps m4-1) and m4-2))

- 15 **Synthesis of 1-(3-difluoromethoxyphenyl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea**



- 96.84 mg of 3-difluoromethoxyphenylamine (590.31 μmol), 206.20 μl of DIEA (1.18 mmol) and 30.81 μl of phenyl chloroformate (237.83 μmol) are added to 4 ml of THF in a microwave tube. The mixture is stirred at room temperature for 20 minutes. 250 mg of 4-(4-aminophenyl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one (590.31 μmol) are added and the resulting mixture is heated for 1 hour at 120°C in a Biotage microwave machine. The reaction medium is evaporated under reduced pressure. The residue is chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to (98/2 v/v) to give 180 mg of 1-(3-difluoromethoxyphenyl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-

azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea.


LC/MS (method C): $M+H^+ = 609$; Tr = 0.81 min

^1H NMR (300 MHz, DMSO- d_6) δ ppm 9.23 (bs, 2H); 7.44-7.52 (bs, 5H); 7.27-7.36 (m, 1H); 6.92-7.25 (m, 2H); 6.73-6.81 (m, 1H); 4.41 (bs, 2H); 3.95-4.36 (bs, 2H); 3.67 (s, 2H); 3.37-3.63 (bs, 2H); 3.10 (m, 2H); 1.82 (bs, 2H); 1.65 (bs, 2H); 1.35 (bs, 6H); 1.15 (t, 3H)

Example 19: (which corresponds to step m) (steps m1-1) and m1-2))

Synthesis of 1-(2,4-difluorophenyl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-

10 triazabenzocyclohepten-4-yl]phenyl}urea



To a solution of 350.35 mg (1.18 mmol) of triphosgene in 6 ml of DCM is added in a single portion a suspension of 500 mg (1.18 mmol) of 4-(4-aminophenyl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one and 987.33 μ l (7.08 mmol) of triethylamine in 6 ml of DCM. The resulting mixture is stirred for 20 minutes at room temperature. 607.12 μ l (5.90 mmol) of 2,4-difluoroaniline are added. The precipitate obtained is filtered off, rinsed twice with MeOH and dried under reduced pressure to give 575 mg of 1-(2,4-difluorophenyl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea.


LC/MS (method C): $M+H^+ = 579$; Tr = 0.80 min

^1H NMR (300 MHz, DMSO- d_6) δ ppm 9.22 (bs, 1H); 8.56 (bs, 1H); 8.08 (dt, 1H); 7.42-7.52 (m, 4H); 7.33 (m, 1H); 7.06 (m, 1H); 4.41 (bs, 2H); 4.11 (q, 2H); 3.67 (bs, 2H); 3.40-3.62 (bs, 2H); 3.10 (bs, 2H); 1.82 (bs, 2H); 1.65 (bs, 2H); 1.35 (bs, 6H); 1.15 (t, 3H)

Example 20: (which corresponds to step m) (steps m1-1) and m1-2))

{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-

**tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid
(S)-2-hydroxypropyl ester**




A mixture of 134.43 mg (0.45 mmol) of triphosgene, 200 mg (0.45 mmol) of 4-(4-
5 amino-3-fluorophenyl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-
dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one and 275.04 mg (2.72 mmol) of
triethylamine in 50 ml of DCM is stirred at room temperature for 30 minutes. 134 mg
(1.81 mmol) of (S)-propane-1,2-diol are added and the mixture is stirred for 2 hours at
room temperature. The resulting mixture is hydrolysed with 5 ml of water and then
10 evaporated under reduced pressure. The product is precipitated by addition of ethyl
acetate/water. The product is filtered off and dried under vacuum to give 120 mg {4-[6-
ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-
1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}carbamic acid (S)-2-hydroxypropyl
ester.

15 LC/MS (method C): $M+H^+$ = 544; Tr = 0.64 min

^1H NMR (600 MHz, DMSO- d_6) δ ppm 9.45 (bs, 1H); 7.7 (t, 1H); 7.32 (t, 2H); 4.8 (d, 1H);
4.42 (bs, 2H); 4.3 (m, 2H); 3.95 (m, 2H); 3.87 (m, 1H); 3.7 (bs, 2H); 3.5 (bs, 2H); 3.09
(bs, 1H); 1.82 (bs, 1H); 1.66 (bs, 1H); 1.35 (bs, 6H); 1.12 (t, 3H); 1.1 (t, 3H)

Example 21: (which corresponds to step m) (steps m1-1) and m1-2))

20 **Synthesis of 1-(4-methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-
azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-
triazabenzocyclohepten-4-yl]phenyl}urea**





- 210 mg (0.51 mmol) of 4-(4-aminophenyl)-6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one and 314.5 mg (3 mmol) of triethylamine are added to 3 ml of DCM. This mixture is then added
- 5 under nitrogen to a solution containing 152.2 mg (0.51 mmol) of triphosgene in 3 ml of DCM. The mixture is stirred at room temperature for 30 minutes, and 268.1 mg (2 mmol) of 2-amino-4-methoxypyridine are then added. This mixture is stirred for 2 hours at room temperature, extracted with DCM, washed once with water and then with saturated NaCl solution, and the organic phase is dried over Na₂SO₄. The resulting phase is filtered and
- 10 the filtrate is evaporated under reduced pressure. The residue is chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to (100/4 v/v). The solid obtained is then taken up in a DCM/MeOH/pentane mixture, filtered off and dried to give 85 mg of 1-(4-methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea.
- 15 LC/MS (method C): M+H⁺ = 560; Tr = 0.55 min


¹H NMR (600 MHz, DMSO-d₆) δ ppm 9.42 (s, 1H); 8.11 (d, 1H); 7.50-7.54 (m, 4H); 7.10 (s, 1H); 6.64 (d, 1H); 4.41 (s, 2H); 4.10-4.40 (bs, 2H); 3.81 (s, 3H); 3.70 (s, 2H); 3.07-3.13 (m, 5H); 1.80-1.84 (m, 2H); 1.62-1.68 (m, 2H); 1.35 (s, 6H)

Example 22:

- 20 **Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(1-methylpiperidin-4-yloxy)pyridin-3-yl]urea**

**Step 22.1:****2-(1-Methylpiperidin-4-yloxy)-5-nitropyridine**

- 5 557 mg of NaH (14 mmol) suspended in 10 ml of THF are placed under nitrogen. The mixture is cooled to 0°C and a solution of 1.5 g (9.3 mmol) of 2-chloro-5-nitropyridine and 1.28 g (11.1 mmol) of 4-hydroxy-1-methylpiperidine in 15 ml of THF is added dropwise. The reaction mixture is refluxed for 4 hours. After hydrolysing the excess NaH, the solvent is evaporated off under reduced pressure. The residue is taken up in ethyl
- 10 acetate and washed with water and then with saturated NaCl solution. It is then dried over Na₂SO₄ and filtered, and the filtrate is evaporated under reduced pressure. The residue is chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to 100/4 v/v to give 1.1 g of 2-(1-methylpiperidin-4-yloxy)-5-nitropyridine.
- LC/MS (method E): M+H⁺ = 238; Tr = 0.79 min


15 **Step 22.2:****6-(1-Methylpiperidin-4-yloxy)pyridin-3-ylamine**

- 1.05 g of the compound obtained in step **20.1** are dissolved in 90 ml of MeOH and then hydrogenated at atmospheric pressure at 35°C over 10% Pd/C. The medium is
- 20 filtered and then evaporated under reduced pressure to give 860 mg of 6-(1-

methylpiperidin-4-yloxy)pyridin-3-ylamine.

Step 22.3: (which corresponds to step m) (steps m1-1) and m1-2))

1-**{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(1-methylpiperidin-4-yloxy)pyridin-3-yl]urea**



130 mg (0.29 mmol) of 4-(4-amino-3-fluorophenyl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one and 180.6 mg (1.8 mmol) of triethylamine are placed in 3 ml of DCM. This mixture is then added under nitrogen to a solution containing 87.4 mg (0.29 mmol) of triphosgene in 3 ml of DCM. The resulting mixture is stirred at room temperature for 30 minutes, followed by addition of 244.1 mg (1.18 mmol) of the compound described in step **21.2**. The resulting mixture is stirred for 2 hours at room temperature. This mixture is extracted with DCM, washed once with water and then with saturated NaCl solution, and the organic phase is dried over Na₂SO₄. The resulting phase is filtered and the filtrate is evaporated under reduced pressure. The residue is chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to (100/20 v/v). The solid obtained is then taken up in a DCM/pentane mixture, filtered off and dried to give 90 mg of 1-**{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(1-methylpiperidin-4-yloxy)pyridin-3-yl]urea**.


LC/MS (method C): M+H⁺ = 675; Tr = 0.53 min

¹H NMR (300 MHz, DMSO-d₆) δ ppm 9.03 (s, 1H); 8.80 (d, 1H); 8.15 (d, 1H); 8.13 (t, 1H); 7.84; (dd, 1H); 7.40 (dd, 1H); 7.29 (dd, 1H); 6.75 (d, 1H); 4.89 (m, 1H); 4.02-4.47 (m, 4H); 3.68 (s, 2H); 3.53 (bs, 2H); 3.08 (m, 2H); 2.63 (m, 2H); 2.17 (s, 3H); 2.12 (m, 2H); 1.93 (m, 2H); 1.81 (m, 2H); 1.65 (m, 4H); 1.34 (s, 6H); 1.14 (t, 3H)

Example 23:

Synthesis of 1-[4-(4-dimethylaminopiperidine-1-carbonyl)phenyl]-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea (which corresponds to step m) (steps m1-1)

5 and m1-2))




130 mg (316 μ mol) of 4-(4-aminophenyl)-6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one suspended in 3 ml of dioxane are heated at 50°C for 30 minutes in the presence of 233 μ l of a 1.9 M solution of phosgene in toluene (442.3 mmol) and 75 μ l (442.3 mmol) of DIEA. 391 mg (1.58 mmol) of (4-aminophenyl)-(4-dimethylaminopiperidin-1-yl)methanone are then added. The medium is stirred for 16 hours at room temperature. The mixture is evaporated under reduced pressure and the residue is taken up in water and extracted with DCM. The organic phase is washed twice with water, dried over MgSO_4 and then filtered. The filtrate is evaporated under reduced pressure and the residue is then chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to (93/7 v/v). The product obtained is dissolved in a minimum amount of DCM and then reprecipitated from Et_2O to give 61 mg of 1-[4-(4-dimethylaminopiperidine-1-carbonyl)phenyl]-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea.

LC/MS (method C): $\text{M}+\text{H}^+$ = 685; Tr = 0.52 min

^1H NMR (600 MHz, DMSO-d_6) δ ppm 8.94 (s, 2H); 7.51 (d, 2H); 7.48 (m, 4H); 7.34 (d, 2H); 4.66 (m, 1H); 4.35 (m, 2H); 3.91 (m, 1H); 3.71 (m, 1H); 3.67 (m, 2H); 3.57 (m, 1H); 3.52 (m, 2H); 3.42 (m, 1H); 3.19 (m, 1H); 2.89 (bs, 2H); 2.32 (m, 1H); 2.18 (s, 6H); 1.75 (m, 2H); 1.33 (m, 8H); 1.22 (m, 3H); 1.16 (t, 3H)

Example 24: (which corresponds to step m) (steps m1-1) and m1-2))

Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-hydroxy-2-methylpropyl)urea





5

100 mg (226.5 μ mol) of 4-(4-amino-3-fluorophenyl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one suspended in 2 ml of dioxane are heated at 50°C for 30 minutes in the presence of 179 μ l of a 1.9 M solution of phosgene (339.75 μ mol) in toluene and 58 μ l (339.5 μ mol) of DIEA. 101 mg (1.13 mmol) of 2-hydroxy-2-methylpropylamine dissolved in 1 ml of dioxane are then added. The reaction medium is stirred for 16 hours at room temperature and then diluted with water. The precipitate formed is filtered off and washed with water and then with ether. The solid obtained is dissolved in a minimum amount of DCM and then reprecipitated from ether to give 66 mg of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-hydroxy-2-methylpropyl)urea.


LC/MS (method C): $M+H^+$ = 557; Tr = 0.62 min
 ^1H NMR (600 MHz, DMSO- d_6) δ ppm 8.66 (s, 1H); 8.16 (t, 1H); 7.35 (d, 1H); 7.23 (d, 1H); 6.80 (t, 1H); 4.49 (m, 1H); 4.41 (m, 2H); 4.2 (bs, 2H); 3.66 (m, 2H); 3.52 (bs, 2H); 3.09 (m, 2H); 3.05 (m, 2H); 1.81 (m, 2H); 1.65 (m, 2H); 1.34 (s, 6H); 1.14 (t, 3H); 1.09 (s, 6H)

Example 25:
Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-pyrrolidin-1-ylpyridin-2-yl)urea

25

**Step 25.1:****2-Nitro-5-pyrrolidin-1-ylpyridine**


- 5 A mixture of 600 mg (2.93 mmol) of 5-bromo-2-nitropyridine and 1.05 g (14.63 mmol) of pyrrolidine is heated in a Biotage microwave machine for 1 hour at 120°C. The reaction medium is diluted with water and then extracted with ethyl acetate. The organic phase is washed twice with water, dried over MgSO₄ and then filtered. The filtrate is evaporated under reduced pressure and then chromatographed on silica gel, eluting with
- 10 a heptane/ethyl acetate mixture from (100/0 v/v) up to (50/50 v/v) to give 410 mg of 2-nitro-5-pyrrolidin-1-ylpyridine.

Step 25.2:**5-Pyrrolidin-1-ylpyridin-2-ylamine**

- 15 410 mg (2.12 mmol) of the product obtained in step **25.1** are hydrogenated in the presence of 10% Pd/C at atmospheric pressure at 35°C. The solution obtained is evaporated under reduced pressure to give 321 mg of 5-pyrrolidin-1-ylpyridin-2-ylamine.

Step 25.3: (which corresponds to step m) (steps m1-1) and m1-2))

1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-pyrrolidin-1-ylpyridin-2-yl)urea




5 120 mg (223 μ mol) of 4-(4-aminophenyl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one 2,2,2-trifluoroacetic salt are extracted in carbonate medium with DCM and the free base obtained is then heated in dioxane at 50°C for 30 minutes in the presence of 177 μ l of a 1.9 M solution of phosgene (334.9 μ mol) in toluene and 56 μ l (893 μ mol) of DIEA. 146
10 mg (893 μ mol) of 5-pyrrolidin-1-ylpyridin-2-ylamine obtained in step **25.2** are then added and the reaction medium is stirred for 16 hours at room temperature. The medium is evaporated under reduced pressure and the residue is taken up in water and extracted with DCM. The organic phase is washed twice with water, dried over MgSO_4 and then filtered. The filtrate is evaporated under reduced pressure and the residue is then
15 chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to (95/5 v/v) to give 68 mg of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-pyrrolidin-1-ylpyridin-2-yl)urea.

LC/MS (method C): $\text{M}+\text{H}^+$ = 613; Tr = 0.63 min

20 ^1H NMR (600 MHz, DMSO-d_6) δ ppm 10.42 (bs, 1H); 9.02 (bs, 1H); 7.64 (m, 1H); 7.44 (m, 4H); 7.39 (m, 1H); 7.05 (m, 1H); 4.30 (bs, 4H); 3.66 (s, 2H); 3.50 (bs, 2H); 3.22 (m, 4H); 3.10 (m, 2H); 1.95 (m, 4H); 1.82 (m, 2H); 1.66 (m, 2H); 1.35 (s, 6H); 1.16 (t, 3H)


Example 26:

Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-methylphenyl}-3-isoxazol-3-ylurea



5 **Step 26.1:** (which corresponds to step k))

{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-methylphenyl}carbamic acid tert-butyl ester




- 10 A mixture of 1 g (2.13 mmol) of trifluoromethanesulfonic acid 6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl ester, 711.35 mg (2.13 mmol) of [2-methyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]carbamic acid tert-butyl ester, 906.25 mg (4.27 mmol) of tribasic potassium phosphate, 149.83 mg (213.47 μ mol) of
- 15 bis(triphenylphosphine)palladium(II) chloride in 10 ml of dioxane and 3 ml of water is heated in a Biotage microwave machine at 100°C for 30 minutes. The reaction medium is diluted in 250 ml of ethyl acetate and filtered through Celite. The organic phase is dried over MgSO_4 . The filtrate is evaporated under reduced pressure. The residue is taken up in 30 ml of heptane, brought to reflux and filtered while hot to give 1.24 g of {4-[6-ethyl-

8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-methylphenyl}carbamic acid tert-butyl ester.

Step 26.2: (which corresponds to step l))


4-(4-Amino-3-methylphenyl)-6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one



A mixture of 1.24 g (2.36 mmol) of the compound described in step **26.1** and 5 ml (64.90 mmol) of TFA in 15 ml of DCM is stirred for 45 minutes at room temperature. The reaction medium is evaporated under reduced pressure, taken up in 10% and Na₂CO₃ solution and filtered. The precipitate is purified by chromatography on silica gel, eluting with a DCM/ethyl acetate mixture from (100/0 v/v) up to (95/5 v/v) to give 720 mg of 4-(4-amino-3-methylphenyl)-6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one.

Step 26.3: (which corresponds to step m) (steps m1-1) and m1-2))

1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-methylphenyl}-3-isoxazol-3-ylurea



160.40 mg (540.51 μmol) of triphosgene are added to a mixture of 230 mg


(540.51 μmol) of the compound described in step **26.2** and 452.02 μl (3.24 mmol) of triethylamine in 10 ml of DCM, under argon. The mixture is stirred for 45 minutes at room temperature, and 199.67 μl (2.70 mmol) of isoxazol-3-ylamine are then added. This mixture is stirred overnight at room temperature and the reaction medium is then poured
 5 into water. The resulting mixture is extracted with DCM and washed with saturated NaCl solution. The organic phase is dried over MgSO_4 , filtered and evaporated under reduced pressure. The residue is purified by chromatography on silica gel, eluting with a DCM/EtOAc mixture from (100/0 v/v) up to 95/5 v/v) to give 39 mg of 1-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-
 10 triazabenzocyclohepten-4-yl]-2-methylphenyl}-3-isoxazol-3-ylurea.

LC/MS (method C): $\text{M}+\text{H}^+ = 536$; $\text{Tr} = 0.70$ min

^1H NMR (600 MHz, $\text{DMSO}-d_6$) δ ppm 10.03 (s, 1 H); 9.34 (s, 1H); 9.35 (s, 1H); 7.94 (m, 1H); 7.34-7.39 (m, 2H); 6.83 (d, 1H); 4.65 (s, 1H); 4.30-4.35 (m, 1H); 3.92 (dd, 1H); 3.65-3.73 (m, 3H); 3.58 (dd, 1H); 3.52 (m, 2H); 3.40-3.44 (m, 1H); 3.16-3.22 (m, 1H); 2.26 (s,
 15 3H); 1.36 (s, 6H); 1.22 (d, 3H); 1.16 (t, 3H)

Example 27:

Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-((R)-2-hydroxy-1-methylethyl)urea (which corresponds to step m) (steps m1-1) and m1-2))



20

A mixture of 134.43 mg (0.45 mmol) of triphosgene, 200 mg (0.45 mmol) of 4-(4-amino-3-fluorophenyl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one and 275.04 mg (2.72 mmol) of triethylamine in 50 ml of DCM is stirred at room temperature for 30 minutes. 138.88 mg
 25 (1.81 mmol) of (R)-2-aminopropan-1-ol are then added and the mixture is stirred at room temperature for 2 hours. The resulting mixture is hydrolysed with 5 ml of water and the solvent is then evaporated off under reduced pressure. The product is precipitated by


addition of ethyl acetate/water. The product is filtered off and solvents are evaporated off under reduced pressure to give 68 mg of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-((R)-2-hydroxy-1-methylethyl)urea.

- 5 LC/MS (method C): $M+H^+ = 544$; Tr = 0.64 min

^1H NMR (600 MHz, DMSO- d_6) δ ppm 8.49 (bs, 1H); 8.16 (t, 1H); 7.35 (d, 1H); 7.24 (d, 1H); 6.70 (d, 1H); 3.78 (t, 1H); 4.42 (bs, 4H); 3.70 (m, 1H); 3.68 (s, 2H); 3.5 (bs, 2H); 3.37 (m, 2H); 3.1 (bs, 2H); 1.72 (bs, 2H); 1.66 (bs, 2H); 1.35 (bs, 6H); 1.15 (t, 3H); 1.08 (d, 3H)


10 **Example 28:**

Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methoxypyrazin-2-yl)urea



- 15 **Step 28.1:** (preparation of a compound of formula (CIII))

1-(6-Methoxypyrazin-2-yl)-3-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]urea




A mixture of 350 mg (1.60 mmol) of 4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenylamine and 1.35 ml (9.59 mmol) of triethylamine in 10 ml of DCM is added, under nitrogen, to a solution of 474.07 mg (1.60 mmol) of triphosgene in 30 ml of DCM. The mixture is stirred for 30 minutes at room temperature, followed by addition of 1 g (7.99 mmol) of 6-methoxypyrazin-2-amine. The resulting mixture is stirred for 2 hours at room temperature. Water is added and the product is extracted with DCM. The solution is dried over MgSO_4 and then filtered. The filtrate is evaporated under reduced pressure and the residue is then chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to (96/4 v/v). The solvent is evaporated off under reduced pressure. The residue is taken up in ethyl acetate and washed three times with 1N hydrochloric acid solution and then with saturated NaCl solution. The resulting solution is dried over MgSO_4 . 167 mg of 1-(6-methoxypyrazin-2-yl)-3-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]urea are obtained.

LC/MS (method E): $\text{M}+\text{H}^+ = 371$; $\text{Tr} = 2.46$ min

15 **Step 28.2:** (which corresponds to step n))

1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methoxypyrazin-2-yl)urea



20 A mixture of 167 mg (0.451 mmol) of the compound described in step **28.1**, 216 mg (0.451 mmol) of trifluoromethanesulfonic acid 6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl ester, 197 mg (0.902 mmol) of tripotassium phosphate and 32 mg (0.045 mmol) of bis(triphenylphosphine)palladium (II) chloride in 5 ml of water and 15 ml of dioxane is

25 heated in a Biotage microwave machine at 100°C for 20 minutes. The reaction mixture is evaporated under reduced pressure and the residue is then chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to (96/4 v/v). The product is


taken up in EtOH and then filtered through a Millipore filter to give 85 mg of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methoxypyrazin-2-yl)urea.

LC/MS (method C): $M+H^+$ = 575; Tr = 0.70 min

- 5 ^1H NMR (600 MHz, $\text{DMSO}-d_6$) δ ppm 9.47 (s, 1H) 9.37 (s, 1H); 8.68 (s, 1H); 7.92 (s, 1H); 7.5 (m, 4H); 4.15-4.4 (bs, 4H); 3.95 (s, 3H); 3.65 (bs, 4H); 3.1 (d, 2H); 1.82 (bs, 2H); 1.65 (bs, 2H); 1.37 (s, 6H); 1.17 (t, 3H)


Example 29:

- Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((S)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
- 10



Step 29.1:

(S)-3-Methyl-4-(5-nitropyridin-2-yl)morpholine



15


- A mixture of 1 g (6.18 mmol) of 2-chloro-5-nitropyridine and 1.31 g (12.36 mmol) of (S)-3-methylmorpholine is heated in a Biotage microwave machine for 1 hour at 150°C. The reaction medium is taken up in MeOH and then evaporated under reduced pressure. The residue is chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to (95/5 v/v) to give 1.35 g of (S)-3-methyl-4-(5-nitropyridin-2-
- 20

yl)morpholine.

LC/MS (method E): $M+H^+ = 224$; Tr = 1.72 min

Step 29.2:

6-((S)-3-Methylmorpholin-4-yl)pyridin-3-ylamine




5

1.35 g (6 mmol) of the product obtained in step **29.1** are hydrogenated in the presence of 10% Pd/C at atmospheric pressure at 50°C. The solution obtained is evaporated under reduced pressure to give 1.17 g of 6-((S)-3-Methylmorpholin-4-yl)pyridin-3-ylamine.

10 LC/MS (method E): $M+H^+ = 194$; Tr = 0.45 min

Step 29.3: (which corresponds to step m) (steps m1-1) and m1-2))

1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((S)-3-methylmorpholin-4-yl)pyridin-3-yl]urea



15

150 mg (0.34 mmol) of 4-(4-amino-3-fluorophenyl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one and 287.20 μ l (2.04 mmol) of triethylamine are placed in 3 ml of DCM. This mixture is then added under nitrogen to a solution containing 100 mg (0.34 mmol) of triphosgene in 20 3 ml of DCM. The resulting mixture is stirred at room temperature for 30 minutes,

followed by addition of 262.62 mg (1.36 mmol) of the compound described in step **29.2**. This mixture is stirred for 2 hours at room temperature and is then extracted with DCM. The organic phase is washed once with water and with saturated NaCl solution. The organic phase is dried over Na₂SO₄, filtered and evaporated under reduced pressure.


- 5 The residue is chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to (98/2 v/v). The solid obtained is then washed twice with EtOH and taken up in an ethyl acetate/pentane mixture (10/90), filtered and dried to give 72 mg of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((S)-3-methylmorpholin-4-yl)pyridin-3-yl]urea.

LC/MS (method C): M+H⁺ = 661; Tr = 0.59 min

- ¹H NMR (600 MHz, DMSO-d₆) δ ppm 8.9 (s, 1H); 8.73 (s, 1H); 8.15 (m, 2H); 7.72 (dd, 1H); 7.4 (d, 1H); 7.28 (d, 1H); 6.75 (d, 1H); 4.4 (bs, 4H); 4.25 (m, 1H); 3.92 (dd, 1H); 3.68 (m, 5H); 3.5 (dt, 3H); 3.1 (m, 2H); 3.02 (m, 1H); 1.82 (bs, 2H); 1.65 (bs, 2H); 1.35 (s, 6H);
- 15 1.15 (t, 4H); 1.08 (d, 3H)

Example 30:


Synthesis of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((R)-3-methylmorpholin-4-yl)pyridin-3-yl]urea



20

Step 30.1:

(R)-3-Methyl-4-(5-nitropyridin-2-yl)morpholine




A mixture of 1 g (6.18 mmol) of 2-chloro-5-nitropyridine and 1.31 g (12.36 mmol) of (R)-3-methylmorpholine is heated in a Biotage microwave machine for 1 hour at 150°C. The reaction medium is taken up in MeOH and then evaporated under reduced pressure. The residue is chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to (95/5 v/v) to give 1.07 g of (R)-3-methyl-4-(5-nitro-pyridin-2-yl)-morpholine.

LC/MS (method E): $M+H^+ = 224$; Tr = 1.72 min

Step 30.2:

10 6-((R)-3-Methylmorpholin-4-yl)pyridin-3-ylamine




1.07 g (4.79 mmol) of the product obtained in step **30.1** are hydrogenated in the presence of 10% Pd/C at atmospheric pressure at 45°C. The solution obtained is evaporated under reduced pressure to give 840 mg of 6-((R)-3-methylmorpholin-4-yl)pyridin-3-ylamine.

LC/MS (method E): $M+H^+ = 194$; Tr = 0.45 min

Step 30.3: (which corresponds to step m) (steps m1-1) and m1-2))

1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((R)-3-methylmorpholin-4-yl)pyridin-3-yl]urea










200 mg (453 μ mol) of 4-(4-amino-3-fluorophenyl)-6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-7,8-dihydro-6H-9-oxa-1,3,6-triazabenzocyclohepten-5-one and 287.20 μ l (2.72 mmol) of triethylamine are placed in 3 ml of DCM. This mixture is then added under nitrogen to a solution containing 134 mg (453 μ mol) of triphosgene in 3 ml of DCM. The resulting mixture is stirred at room temperature for 30 minutes, followed by addition of 350.16 mg (1.81 mmol) of the compound obtained in step **30.2**. The resulting mixture is stirred for 2 hours at room temperature. The product is taken up in DCM and washed once with water and then with saturated NaCl solution. The organic phase is dried over Na₂SO₄. The resulting phase is filtered and the filtrate is evaporated under reduced pressure. The residue is chromatographed on silica gel, eluting with a DCM/MeOH mixture from (100/0 v/v) up to (98/2 v/v). The solid obtained is then washed twice with EtOH and taken up in an ethyl acetate/pentane mixture (10/90), filtered and dried over MgSO₄ to give 40 mg of 1-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((R)-3-methylmorpholin-4-yl)pyridin-3-yl]urea.





LC/MS (method C): M+H⁺ = 661; Tr = 0.59 min




¹H NMR (600 MHz, DMSO-d₆) δ ppm 8.9 (s, 1H); 8.73 (s, 1H); 8.15 (m, 2H); 7.72 (dd, 1H); 7.4 (d, 1H); 7.28 (d, 1H); 6.75 (d, 1H); 4.4 (bs, 4H); 4.25 (m, 1H); 3.92 (dd, 1H); 3.68 (m, 5H); 3.5 (dt, 3H); 3.1 (m, 2H); 3.02 (m, 1H); 1.82 (bs, 2H); 1.65 (bs, 2H); 1.35 (s, 6H); 1.15 (t, 4H); 1.08 (d, 3H)




Table 3 below illustrates the chemical structures and physical properties of a number of examples of compounds according to the invention.




Exptl	Structure	Mass	Tr (min)	LC/MS method
1		499	0.9	A
2		487	0.91	A
3		501	1.02	A
4		568	0.76	F



Expt	Structure	Mass	Tr (min)	LC/MS method
5		640	1.08	A
6		610	0.91	A
7		539	0.86	A

Exptl	Structure	Mass	Tr (min)	LC/MS method
8		566	0.86	A
9		525	0.82	A
10		509	0.96	A
11		487	0.64	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
12		550	0.57	C
13		659	0.53	C
14		551	0.61	C



Expt	Structure	Mass	Tr (min)	LC/MS method
15		604	0.68	C
16		603	0.63	C
17		496	0.59	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
18		609	0.81	C
19		579	0.8	C
20		544	0.64	C




Expl	Structure	Mass	Tr (min)	LC/MS method
21		560	0.55	C
22		675	0.53	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
23		685	0.52	C
24		557	0.62	C
25		613	0.63	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
26		536	0.7	C
27		543	0.6	C
28		575	0.7	C




Expl	Structure	Mass	Tr (min)	LC/MS method
29		661	0.59	C
30		661	0.59	C




Expl	Structure	Mass	Tr (min)	LC/MS method
31		645	0.57	C
32		631	0.54	C
33		544	0.61	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
34		558	0.62	C
35		574	0.55	C
36		546	0.93	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
37		578	0.91	A
38		467	0.54	C
39		530	0.57	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
40		618	0.68	C
41		530	0.58	C
42		544	0.58	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
43	 <chem>CNC(=O)Nc1ccc(cc1)c2nc3c(ncn3C45CCOCC45)C(=O)N(C)CC6C(C)OC62</chem>	467	0.54	C
44	 <chem>COc1ccc(cc1)NC(=O)Nc2ccc(cc2)c3nc4c(ncn4C56CCOCC56)C(=O)N(CC)CC7C(C)OC73</chem>	574	0.68	C
45	 <chem>CNC(=O)Nc1ccc(cc1F)c2nc3c(ncn3C45CCOCC45)C(=O)N(C)CC6C(C)OC62</chem>	485	0.57	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
46		485	0.58	C
47		499	0.62	C
48		544	0.59	C




Expl	Structure	Mass	Tr (min)	LC/MS method
49		560	0.64	C
50		562	0.81	A
51		562	0.82	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
52		562	0.82	A
53		540	1.03	A
54		560	0.64	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
55		576	0.85	A
56		576	0.84	A
57		542	0.6	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
58		590	0.81	A
59		562	0.5	C
60		562	0.5	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
61		544	0.65	C
62		544	0.65	C
63		544	0.88	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
64		587	0.58	C
65		512	0.88	A
66		562	0.53	C




Expl	Structure	Mass	Tr (min)	LC/MS method
67		562	0.53	C
68		530	0.47	C
69		530	0.74	C




Expl	Structure	Mass	Tr (min)	LC/MS method
70		560	0.56	C
71		562	0.57	C
72		629	0.56	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
73		647	0.57	C
74		647	0.57	C
75		562	0.81	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
76		580	0.86	A
77		647	1.08	A
78		605	0.81	A





Exptl	Structure	Mass	Tr (min)	LC/MS method
79		574	0.84	A
80		724	0.82	A
81		704	0.86	A




Expl	Structure	Mass	Tr (min)	LC/MS method
82		548	0.52	C
83		548	0.52	C
84		562	0.69	C




Expl	Structure	Mass	Tr (min)	LC/MS method
85		548	0.53	C
86		601	0.59	C
87		647	0.56	C




Expl	Structure	Mass	Tr (min)	LC/MS method
88		526	0.63	C
89		592	0.62	C
90		591	0.9	A




Expl	Structure	Mass	Tr (min)	LC/MS method
91		561	0.84	A
92		545	0.89	A
93		573	0.82	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
94		441	3.8	B
95		371	0.67	A
96		469	3.27	B
97		441	0.77	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
98		483	0.96	A
99		495	0.88	A
100		443	0.88	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
101		469	0.88	A
102		398	0.67	A
103		456	0.93	A




Expl	Structure	Mass	Tr (min)	LC/MS method
104		471	0.88	A
105		482	1.01	A
106		496	1.08	A

Expl	Structure	Mass	Tr (min)	LC/MS method
107		425	0.72	A
108		480	0.97	A
109		511	1.07	A



Exptl	Structure	Mass	Tr (min)	LC/MS method
110		513	1	A
111		516	0.99	A
112		544	0.81	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
113		481	0.75	A
114		497	1.03	A
115		547	0.91	A




Expl	Structure	Mass	Tr (min)	LC/MS method
116		548	1.06	A
117		511	0.82	A
118		540	0.91	A



Expt	Structure	Mass	Tr (min)	LC/MS method
119		525	0.85	A
120		595	1.14	A
121		522	0.96	A




Expl	Structure	Mass	Tr (min)	LC/MS method
122	 <chem>CCN(CC)C(=O)O[C@H]1C(C)(C)O[C@H]1C2=NC(=C(N3C(=O)N(C)C=C3)C(=O)N4C(=O)C=C(C4)C5=CC=CC=C5N6C(=O)O[C@H]6C7C8C9C10C9C8C7C6)C2=NC(=O)N1</chem>	526	0.97	A
123	 <chem>CCN(CC)C(=O)O[C@H]1C(C)(C)O[C@H]1C2=NC(=C(N3C(=O)N(C)C=C3)C(=O)N4C(=O)C=C(C4)C5=CC=CC=C5N6C(=O)O[C@H]6C7C8C9C10C9C8C7C6)C2=NC(=O)N1</chem>	513	0.93	A
124	 <chem>CCN(CC)C(=O)O[C@H]1C(C)(C)O[C@H]1C2=NC(=C(N3C(=O)N(C)C=C3)C(=O)N4C(=O)C=C(C4)C5=CC=CC=C5N6C(=O)O[C@H]6C7C8C9C10C9C8C7C6)C2=NC(=O)N1</chem>	592	1.2	A



Expl	Structure	Mass	Tr (min)	LC/MS method
125		547	1.07	A
126		547	0.89	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
127		573	1.26	A
128		569	0.68	C
129		617	0.93	A



Exptl	Structure	Mass	Tr (min)	LC/MS method
130		596	0.49	C
131		543	0.96	A
132		564	3.23	B



Exptl	Structure	Mass	Tr (min)	LC/MS method
133		525	1.01	A
134		583	0.67	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
135		552	1.04	A
136		593	0.98	A
137		541	3.4	B




Expl	Structure	Mass	Tr (min)	LC/MS method
138		609	1.19	A
139		593	0.75	C




Expl	Structure	Mass	Tr (min)	LC/MS method
140		593	0.72	C
141		580	1.08	A
142		592	0.81	C

Expl	Structure	Mass	Tr (min)	LC/MS method
143	 <chem>CCN(CC1OC2C(C1)C3=NC4=C(N2)N=CN(C4)N5C(=O)N(C5)C6=CC=C(C=C6)NC(=O)N7C=CC(=N7)C</chem>	565	0.93	A
144	 <chem>CCN(CC1OC2C(C1)C3=NC4=C(N2)N=CN(C4)N5C(=O)N(C5)C6=CC=C(C=C6)NC(=O)N7C=CC(=C7)OC</chem>	593	1.05	A




Expl	Structure	Mass	Tr (min)	LC/MS method
145		606	1.02	A
146		575	0.67	C




Expl	Structure	Mass	Tr (min)	LC/MS method
147		611	0.86	C
148		630	0.66	C
149		555	1.1	A




Expl	Structure	Mass	Tr (min)	LC/MS method
150		513	1.01	A
151		515	1.07	A
152		523	0.84	A





Exptl	Structure	Mass	Tr (min)	LC/MS method
153		608	0.69	C
154		629	0.9	C
155		609	0.8	C

Expl	Structure	Mass	Tr (min)	LC/MS method
156		596	0.7	C
157		590	0.65	C
158		562	1.09	A



Exptl	Structure	Mass	Tr (min)	LC/MS method
159		540	1.03	A
160		551	1.04	A
161		564	0.79	A



Expl	Structure	Mass	Tr (min)	LC/MS method
162		609	0.79	C
163		580	1.15	A
164		564	1.15	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
165		548	0.68	C
166		553	1.01	A
167		565	1.07	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
168		546	0.9	A
169		546	1.01	A
170		555	0.95	A
171		648	0.81	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
172		564	0.66	C
173		562	1.06	A
174		630	0.76	A




Expl	Structure	Mass	Tr (min)	LC/MS method
175		588	0.69	C
176		671	0.51	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
177		697	0.5	C
178		622	0.76	C




Expl	Structure	Mass	Tr (min)	LC/MS method
179		582	0.7	C
180		552	0.67	C
181		580	0.99	A




Expl	Structure	Mass	Tr (min)	LC/MS method
182		553	0.8	A
183		616	1.04	A
184		562	0.87	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
185		576	0.78	A
186		500	0.88	A
187		635	0.86	A




Expl	Structure	Mass	Tr (min)	LC/MS method
188		597	3.13	D
189		636	4.38	D
190		658	2.49	D

Expl	Structure	Mass	Tr (min)	LC/MS method
191		567	0.81	C
192		550	1.1	A
193		562	1.19	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
194		565	0.75	A
195		660	0.78	A
196		576	0.71	C

Exptl	Structure	Mass	Tr (min)	LC/MS method
197		604	0.67	C
198		575	0.72	C
199		564	0.81	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
200		647	0.85	A
201		565	0.6	C
202		549	0.8	C




Expl	Structure	Mass	Tr (min)	LC/MS method
203		640	0.54	C
204		597	0.82	C
205		641	0.51	C




Expl	Structure	Mass	Tr (min)	LC/MS method
206		485	0.77	A
207		499	0.89	A
208		597	0.81	C
209		518	0.92	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
210		529	0.61	C
211		543	0.66	C
212		615	0.86	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
213	 <chem>CCN(CC)C(=O)c1c2nc(NC(=O)c3ccc(NC(=O)c4cc(OC)c(F)cc4)cc3)nc2c1OCC(C)(C)O</chem>	579	0.81	C
214	 <chem>CCN(CC)C(=O)c1c2nc(NC(=O)c3ccc(NC(=O)OC(C)(C)O)cc3)cc2c1OCC(C)(C)O</chem>	528	0.98	A
215	 <chem>CCN(CC)C(=O)c1c2nc(NC(=O)c3ccc(NC(=O)c4ccc(NC(C)C)cc4)cc3)cc2c1OCC(C)(C)O</chem>	659	0.52	C

Exptl	Structure	Mass	Tr (min)	LC/MS method
216		627	0.82	C
217		525	0.62	C
218		630	0.75	A



Exptl	Structure	Mass	Tr (min)	LC/MS method
219		646	0.64	C
220		657	0.84	C
221		532	1.01	A




Expl	Structure	Mass	Tr (min)	LC/MS method
222	 <chem>CCN(CC)C(=O)c1c2c(c3c1nc(NC(=O)OCC)cc(F)c3)nc4c2OCC5(C)CC6C(C5)OC6</chem>	530	0.88	A
223	 <chem>CCN(CC)C(=O)c1c2c(c3c1nc(NC(=O)Nc4ccncc4)cc(F)c3)nc4c2OCC5(C)CC6C(C5)OC6</chem>	562	0.8	A
224	 <chem>CCN(CC)C(=O)c1c2c(c3c1nc(NC(=O)OC(C)(C)CO)cc(F)c3)nc4c2OCC5(C)CC6C(C5)OC6</chem>	557	0.66	C




Exptl	Structure	Mass	Tr (min)	LC/MS method
225		517	0.63	C
226		575	0.64	C
227		634	0.97	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
228		501	0.98	A
229		576	0.84	A
230		630	1.2	A




Expl	Structure	Mass	Tr (min)	LC/MS method
231		550	0.82	A
232		618	0.71	C
233		648	0.74	C




Expl	Structure	Mass	Tr (min)	LC/MS method
234		606	0.76	C
235		723	1.04	A

Expl	Structure	Mass	Tr (min)	LC/MS method
236		703	0.97	A
237		694	1.01	A
238		512	0.6	C




Expl	Structure	Mass	Tr (min)	LC/MS method
239		544	0.66	C
240		592	1.03	A
241		590	0.80	A




Exptl	Structure	Mass	Tr (min)	LC/MS method
242		580	0.64	C
243		530	0.64	C
244		659	0.56	C

Exptl	Structure	Mass	Tr (min)	LC/MS method
245		659	0.56	C
246		530	0.64	C
247		587	0.75	C

Expl	Structure	Mass	Tr (min)	LC/MS method
248		663	1.28	A
249		675	0.92	A
250		629	0.63	C

[illegible]

Exptl	Structure	Mass	Tr (min)	LC/MS method
254		525	0.98	A
255		513	0.97	A
256		663	0.96	A

Exptl	Structure	Mass	Tr (min)	LC/MS method
257		675	0.95	A
258		661	0.87	A
259		673	0.85	A

[illegible]

Table 3: Chemical structures and physical properties of a number of examples of compounds according to the invention.

[THERAPEUTIC APPLICATION]

5 The compounds according to the invention underwent pharmacological trials to determine their inhibitory effect on mTOR and the effect of the two complexes mTORC1 and mTORC2.

The trials consisted in measuring the *in vitro* activity of the compounds of the invention on mTOR and also on PI3K α , Akt-pS473 and on the proliferation of U87MG 10 cells. The inhibitory activities are given by the concentration that inhibits 50% of the activity of mTOR, PI3K α , Akt-pS473 or of the proliferation of U87MG cells.

Model 1: mTOR enzymatic test by TR-FRET (time-resolved fluorescence resonance energy transfer).

The test detects the phosphorylation of the substrate GFP-4EBP1 using a terbium-labelled antibody that is specific for the phospho-Thr46 epitope. Binding of the
 5 Tb-antibody to the phosphorylated GFP peptide allows the transfer of energy from Tb, excited at 340 nm, to GFP with an increase in the fluorescence emission of the GFP at 520 nm. The test is performed in 96-well format (Corning/Costar 96 black flat-bottomed half-wells plate, ref. 3694) in a total volume of 30 µl. To 1 µl of inhibitor in 100% DMSO are added (final concentrations) 400 nM of GFP-4EBP1 (Invitrogen PV4759), 8 µM of
 10 ATP, 200 ng/ml of enzyme (human recombinant aa mTOR 1360-2549, Invitrogen PV4753) in a buffer of HEPES 50 mM pH 7.5, EGTA 1 mM, MnCl₂ 10 mM, BSA 0.1 mg/ml, glycerol 0.5%, NV-10 0.1 mg/ml, DTT 2 mM. The mixture is incubated for 30 minutes at room temperature. 10 µl of solution are taken up and the reaction is quenched by adding 10 µl of Tb-antibody solution (Invitrogen PV4757)/EDTA (2 nM Tb-Ac and 10
 15 mM EDTA final). After 60 minutes at room temperature, the fluorescence at 490 nm, from the Tb emission, and at 520 nm, from the GFP emission, is measured in a BMG PHERAstar reader equipped with a LanthascreenTM filter (excitation at 340 nm, emission at 490 and 520 nm). The results are given in the form of the 520/490 ratio.

The IC₅₀ values are determined by the preparation of successive threefold
 20 dilutions on at least a scale of 10 000.

Model 2: PI3K α enzymatic test by measuring the consumption of ATP.

The test uses a luciferin/luciferase system to measure the concentration of ATP and its consumption during the enzymatic reaction. The test is performed in 96-well format (Corning/Costar 96 black flat-bottomed half-wells plate, ref.3694) in a total volume
 25 of 30 µl.

To 1 µl of inhibitor in 100% DMSO are added (final concentrations) 50 µM of the substrate PIP2 ((L- α -phosphatidyl-D-myoinositol 4,5-bisphosphate, Echelon 117P-4516-0500), 2 µM of ATP and 1.7 µg/ml of PI3K α (p110 α /p85 α , Invitrogen PV4788) in a buffer of Tris/HCl 50 mM pH 7.5, EGTA 1 mM, MgCl₂ 10 mM, Chaps 0.03%, 1 mM DTT). After
 30 90 minutes, the reaction is quenched by adding 20 µl/well of KinaseGlo reagent (Promega V6713). After 10 minutes in the dark, the luminescence is read on the PHERAStar microplate reader (reading at 0.8 sec/well).

The IC₅₀ values are determined by the preparation of successive threefold dilutions on at least a scale of 10 000.

Model 3: Cellular test of phosphorylation of the epitope S473 of the Akt protein on the U87MG line via an ELISA test

5 Principle

Starting with a cell lysate, the phosphorylation of Serine-473 of AKT kinase is measured via an ELISA sandwich test using an anti-AKT uptake antibody and an anti-AKT Phospho Serine-473 detection antibody. An electrochemiluminescence signal is generated by a ruthenium-labelled anti-rabbit antibody.

10 Cellular test

The cells are seeded at 12×10^4 cells/well/100 µl of culture medium [90% of DMEM 4.5 g/l of glucose and L-glutamine (Gibco 41965), 10% of heat-inactivated foetal calf serum, 1 mM of sodium pyruvate (Gibco 11360-039), penicillin-streptomycin (10 000 IU-10 mg/ml, Gibco 15140-023, dil. 200x), and non-essential amino acids (Gibco 11140-035, dil. 100x)] in a type I collagen-coated 96-well culture plate (Becton Dickinson 356407). After culturing for 16-20 hours (the cells are at 100% of confluence) at 37°C under 5% CO₂, 1 µl of the chemical compounds in 100% DMSO (1% DMSO final concentration on the cells) is added. The cells are incubated with the compounds for 3 hours at 37°C under 5% CO₂.

20 After addition of 90 µl of lysis buffer [Hepes 50 mM (Sigma H-0887), NaCl 150 mM, Triton X-100 1%, protease inhibitor cocktail (Complete Mini: Roche 11836 153 001, extemporaneous addition), phosphatase inhibitor cocktail (Sigma P5726 and P0044, extemporaneous addition), AEBSF hydrochloride 2 µM (Calbiochem 101500), NaF 10 mM (Fluka 67414)], the mixture is incubated for 1 hour on ice with agitation.

25 Preparation of the ELISA plates

Standard 96-well plates (MSD L11XA-6) are coated with an anti-goat Akt antibody (Santa Cruz sc1618g, without gelatin, without BSA, 2 mg/ml, diluted 30-40 fold in a buffer of HEPES 50 mM pH 7.2, 0.0075% Triton X-100) 2.5 µl/well overnight at room temperature. The free sites are blocked with 150 µl/well of a buffer of 50 mM Tris pH 7.5, 0.15 M NaCl, 0.02% Tween 20, 3% BSA with stirring at 450 rpm for 1 hour at room temperature, followed by a step of washing with 3×300 µl of a buffer of 50 mM Tris pH 7.5, 0.15 M NaCl, 0.02% Tween 20.

ELISA

15 μ l of cell lysate are mixed with 15 μ l of a 50 mM buffer of Tris pH 7.5, 0.15 M NaCl, 0.02% Tween 20, 1% BSA in a precoated ELISA plate (see above), and incubated for 1 hour at room temperature with stirring, followed by a step of washing with 3 $\times$ 300 μ l of a buffer of 50 mM Tris pH 7.5, 0.15 M NaCl, 0.02% Tween 20. The pS473 epitope of Akt is detected with a rabbit polyclonal antibody (Cell Signaling, SC9271), diluted 100-fold in 50 mM Tris pH 7.5, 0.15 M NaCl, 0.02% Tween 20, 1% BSA, 25 μ l/well (incubation for 1 hour at room temperature with stirring at 450 rpm).

After washing (see above), revelation is performed with a ruthenium-labelled anti-rabbit antibody (MSD R32AB1), diluted 3000-fold in a 50 mM buffer of Tris pH 7.5, 0.15 M NaCl, 0.02% Tween 20, 1% BSA, 25 μ l/well, followed by incubation for 1 hour at room temperature with stirring at 450 rpm, and the plates are covered with aluminium.

After a further washing step, the plates are emptied and 150 μ l of MSD Read Buffer T diluted fourfold with water are added. The electrochemiluminescence is measured on the MSD Sector Imager 6000.

15 **Model 4: Test of *in vitro* proliferation of U87MG tumour cells.**

U87MG tumour cells are incubated with the chemical compounds for 72 hours at 37°C under 5% CO₂. The cell growth is determined with the reagent CellTiter 96 AQueous One Solution Cell Proliferation Assay from Promega (Cat. No. G3581).

Seeding is performed at 2500 cells/well/100 μ l of culture medium [90% DMEM 4.5 g/l of glucose and L-glutamine (Gibco 41965), 10% heat-inactivated foetal calf serum, 1 mM sodium pyruvate (Gibco 11360-039), penicillin-streptomycin (10 000 IU-10 mg/ml, Gibco 15140-023, diluted 200-fold), and non-essential amino acids (Gibco 11140-035, diluted 100-fold)], in a type I collagen-coated transparent 96-well culture plate (BD Biosciences Cat No. 356407).

25 After adhesion overnight, 1 μ l of the chemical compounds in 100% DMSO (1% DMSO final concentration) is added. After 72 hours at 37°C under 5% CO₂, 20 μ l of the reagent CellTiter 96 AQueous One Solution Cell Proliferation Assay from Promega (Cat. No. G3581) are added per well. The mixture is incubated for 1 hour 30 minutes at 37 °C under 5% CO₂, and an absorbance reading at 490 nm is then taken (for example on the Victor reader from Wallac/Perkin-Elmer). A reading at T = 0 (before the 72 hours of incubation) is taken, corresponding to 100% inhibition.

The pharmacological data are given in Table 4 below:

Expl	mTOR	PI3Ka	U87	U87
------	------	-------	-----	-----

	IC50, nM	IC50, nM	Akt-pS473 % Inhibition or IC50, nM	Cell Proliferation IC50, nM
1	23	8841	55	197
2	22	547	31	142
3	18	2110	48	242
4	721	>10000	>3000	>10000
5	905	>10000	NT	>10000
6	1070	>10000	NT	7150
7	17	>10000	>3000	>10000
8	44	>10000	537	3810
9	65	>10000	NT	2500
10	344	>10000	611	8770
11	13	2300	61	271
12	17	640	43	121
13	132	6470	13	32
14	41	>10000	458	1420
15	12	5300	76	218
16	19	>10000	269	527
17	2487	>10000	NT	>10000
18	25	7570	29	63
19	32	6219	28	122
20	286	>10000	>1000	7590
21	15	>10000	51	549
22	76	3070	28	68
23	51	79	62	30
24	1072	NT	>1000	>10000
25	130	>10000	124	196
26	112	3860	>1000	3570
27	394	NT	>1000	9150
28	59	>10000	42	508

29	38	>10000	32	165
30	21	>10000	15	46
31	65	>10000	17	35
32	84	>10000	21	115
33	24	9240	42	226
34	43	>10000	70	320
35	37	>10000	75	336
36	55	4400	129	686
37	122	8630	54	869
38	22	3450	23	218
39	26	5870	34	207
40	56	>10000	59	274
41	35	>10000	31	122
42	43	>10000	52	350
43	27	4730	20	82
44	34	>10000	40	220
45	24	4000	32	85
46	26	8880	24	90
47	34	>10000	151	564
48	47	>10000	86	185
49	32	>10000	21	155
50	39	2870	7	40
51	20	1900	14	12
52	18	1710	16	29
53	222	>10000	555	2160
54	32	3000	41	161
55	66	4810	27	54
56	41	3230	26	57
57	239	>10000	>1000	7580
58	28	9470	20	71

59	67	4200	36	42
60	58	NT	17	46
61	993	NT	>1000	4670
62	667	>10000	>1000	>10000
63	31	>10000	31	190
64	24	>10000	37	270
65	27	22400	27	257
66	16	2950	21	71
67	15	>10000	20	69
68	12	1360	16	82
69	6	472	35	79
70	27	>10000	40	288
71	21	>10000	59	209
72	23	>10000	58	498
73	71	5910	48	119
74	64	>10000	31	126
75	40	5570	8	47
76	28	2360	29	367
77	56	4490	16	53
78	90	6260	28	83
79	53	>10000	41	383
80	72	>10000	48	151
81	124	2100	17	108
82	21	2240	13	29
83	20	1460	16	41
84	23	>10000	255	760
85	24	1950	28	77
86	377	>10000	144	490
87	57	>10000	20	59
88	198	>10000	65% at 1000 nM	1900

89	84	>10000	260	221
90	450	NT	64% at 1000 nM	2900
91	53	4640	47	410
92	40	>10000	12	99
93	72	>10000	65	410
94	112	7320	>10000	NT
95	624	>10000	NT	NT
96	72	1444	107	1469
97	800	2159	NT	>10000
98	26	1138	20	184
99	31	>10000	40	239
100	7285	7440	NT	>10000
101	4045	NoVal	NT	>10000
102	927	3610	NT	9830
103	194	6590	425	3510
104	1494	3750	NT	>10000
105	180	>10000	360	2690
106	442	>10000	767	6560
107	2648	9380	NT	>10000
108	472	>10000	NT	4450
109	7770	2046	NT	NT
110	45	>10000	89	506
111	6349	>10000	NT	NT
112	20	2660	24	154
113	3960	>10000	NT	>10000
114	1220	>10000	NT	>10000
115	147	>10000	NT	5000
116	42	4310	98	282
117	25	7410	812	1699
118	58	5040	32	233

221

119	104	>10000	1060	>3000
120	351	>10000	1180	2258
121	66	>10000	130	802
122	19	4200	140	340
123	22	3627	150	221
124	20	>10000	61	115
125	16	>10000	86	260
126	13	2963	68	225
127	16	3646	24	79
128	3475	>10000	NT	>10000
129	1384	>10000	>3000	>10000
130	2120	>10000	>3000	>10000
131	383	>10000	1990	5900
132	245	4320	>3000	>10000
133	33	>10000	92	219
134	3165	>10000	>3000	>10000
135	11	6525	26	112
136	34	>10000	37	239
137	29	>10000	241	1330
138	299	>10000	105	331
139	251	>10000	721	3010
140	45	>10000	507	472
141	34	598	19	94
142	495	>10000	NT	3100
143	19	3180	62	295
144	105	>10000	142	2570
145	2977	>10000	NT	>10000
146	35	>10000	82	441
147	182	>10000	28	297
148	68	>10000	21	490

149	113	>10000	154	854
150	40	5800	73	201
151	116	>10000	82	531
152	348	>10000	257	2680
153	31	>10000	116	346
154	182	>10000	54	97
155	79	>10000	59	142
156	27	1140	92	395
157	27	>10000	67	260
158	64	>10000	1600	1700
159	18	561	21	98
160	41	4900	114	744
161	38	321	9	60
162	147	>10000	110	486
163	55	3870	130	1020
164	85	3800	227	872
165	180	>10000	1530	4200
166	86	2300	278	1270
167	62	3520	158	500
168	18	1385	48	231
169	31	1450	26	299
170	106	824	94	557
171	37	308	41	91
172	20	>10000	98	198
173	12	1602	42	242
174	22	326	36	125
175	585	>10000	NT	3050
176	69	100	314	83
177	84	87	435	57
178	45	>10000	104	252

179	58	>10000	331	1330
180	11	604	65	146
181	74	7650	285	1320
182	26	159	407	2930
183	128	>10000	165	752
184	12	12978	65	368
185	24	5680	20	73
186	25	3940	60	266
187	37	472	22	67
188	63	2330	45	505
189	62	1890	56	451
190	40	927	69% at 3000 nM	734
191	24	570	26	105
192	29	1310	82% at 3000 nM	1770
193	970	9440	>1000	4660
194	27	1230	62% at 3000 nM	2130
195	61	6800	45	57
196	276	7470	60% at 3000 nM	5050
197	117	>10000	150	523
198	57	>10000	44	318
199	39	174	10	32
200	52	>10000	9	108
201	48	813	230	898
202	19	213	5	73
203	96	9610	111	43
204	61	468	18	74
205	63	3740	13	22
206	145	>10000	87% at 1000 nM	1340
207	18	>10000	19	74
208	81	1240	26	181

209	155	>10000	69	1050
210	41	>10000	174	625
211	87	>10000	270	894
212	141	1540	8	71
213	48	991	20	103
214	2448	>10000	>1000	>10000
215	48	37	33	32
216	125	1830	NT	514
217	59	>10000	80% at 3000 nM	1760
218	58	1060	17	211
219	38	98	28	436
220	348	1860	30	285
221	13	372	11	137
222	123	>10000	213	1680
223	20	2070	4	61
224	1120	>10000	NT	NT
225	36	1360	91	365
226	1280	>10000	NT	NT
227	76	484	7	29
228	70	6600	72	229
229	43	>10000	19	86
230	77	>10000	18	58
231	16	103	12	29
232	32	5630	26	95
233	68	>10000	87	908
234	67	>10000	118	2290
235	89	>7200	10	29
236	112	>7200	20	29
237	39	>7200	80	49
238	678	>10000	>1000	1700

239	1070	NT	38% at 1000 nM	>10000
240	61	>10000	29	75
241	165	>10000	43	59
242	89	>10000	770	2300
243	63	3870	160	610
244	60	>10000	29	77
245	63	>10000	67	NT
246	82	>10000	240	1600
247	26	7200	8	62
248	68	>10000	29	96
249	365	>10000	27	29
250	36	>10000	38	100
251	64	>10000	80	340
252	68	1360	48	282
253	72	>10000	68	167
254	68	>10000	68	240
255	72	>10000	68	210
256	196	199	42	83
257	142	>10000	46	63
258	53	2940	20	125
259	87	>10000	16	87
260	174	>10000	24	95
261	387	>10000	20	160

Table 4: Pharmacological data for a number of examples of compounds according to the invention (NT means not tested)

It thus appears that the compounds according to the invention have selective inhibitory activity on the kinase mTOR.

- 5 The compounds according to the invention may thus be used for the preparation of medicaments, in particular medicaments for inhibiting mTOR.

A subject of the present invention is also the use of a compound according to the invention in the treatment of diseases requiring inhibition of the kinase mTOR.

According to another of its aspects, a subject of the present invention is more particularly the use of a compound as defined above for the preparation of a medicament
5 that is useful for treating or preventing:

- cancers, especially:
 - malignant haematological tumours, such as leukaemia, multiple myeloma, lymphomas such as Hodgkin's disease, non-Hodgkin lymphomas (including mantle-cell lymphoma) and myelodysplastic syndromes,
 - 10 - solid tumours and metastases thereof such as breast cancer and lung cancer (non-small cell lung cancer, NSCL; small cell lung cancer, SCLC), squamous cell carcinoma and cancer of the endometrium,
 - central nervous system tumours, such as gliomas, neuro-dysembryoplastic tumours, glioblastoma multiforme, mixed gliomas, medulloblastoma,
 - 15 - retinoblastoma, neuroblastoma, germinoma and teratoma,
 - cancers of the gastrointestinal tract, such as stomach cancer, oesophageal cancer, hepatocellular carcinoma (liver), cholangiocarcinomas, bowel and rectal carcinomas, bowel cancer and pancreatic cancers,
 - skin cancers such as melanomas (in particular metastatic melanoma), thyroid
 - 20 - cancers, cancers of the head and neck, cancers of the salivary glands, of the prostate, of the testicles, of the ovaries, of the cervix, of the vulva, of the bladder, of the kidneys (including clear-cell renal carcinoma and renal oncocytoma), squamous cell carcinomas, sarcomas such as osteosarcoma, chondrosarcoma, leiomyosarcoma, soft-tissue sarcomas, Ewing's sarcomas,
 - 25 - gastrointestinal stromal tumours (GIST), Kaposi's sarcoma, and paediatric cancers such as rhabdomyosarcomas and neuroblastomas,
 - inflammatory and immune diseases, such as rheumatoid arthritis,
 - diabetes,
 - obesity,
 - 30 - obstructive diseases of the respiratory pathways,
 - fibroses (including lung, kidney and liver fibrosis),
 - Pompe's disease (Cori's type II glycogenosis),
 - cardiovascular diseases, or
 - age-related diseases such as neurodegenerative diseases (including Alzheimer's
 - 35 - disease, Huntington's diseases and Parkinson's disease), loss of hearing and ocular diseases, including age-related macular degeneration (ARMD),

- epilepsy,
- parasitic diseases.

Thus, according to another of its aspects, a subject of the invention is
 5 medicaments that comprise a compound of formula (I), or a pharmaceutically acceptable acid-addition salt thereof.

These medicaments find their use in therapeutics, especially in the treatment or prevention of:

- cancers, especially:
 - 10 - malignant haematological tumours, such as leukaemia, multiple myeloma, lymphomas such as Hodgkin's disease, non-Hodgkin lymphomas (including mantle-cell lymphoma) and myelodysplastic syndromes,
 - solid tumours and metastases thereof such as breast cancer and lung cancer (non-small cell lung cancer, NSCL; small cell lung cancer, SCLC), squamous
 15 cell carcinoma and cancer of the endometrium,
 - central nervous system tumours, such as gliomas, neuro-dysembryoplastic tumours, glioblastoma multiforme, mixed gliomas, medulloblastoma, retinoblastoma, neuroblastoma, germinoma and teratoma,
 - cancers of the gastrointestinal tract, such as stomach cancer, oesophageal
 20 cancer, hepatocellular carcinoma (liver), cholangiocarcinomas, bowel and rectal carcinomas, bowel cancer and pancreatic cancers,
 - skin cancers such as melanomas (in particular metastatic melanoma), thyroid cancers, cancers of the head and neck, cancers of the salivary glands, of the prostate, of the testicles, of the ovaries, of the cervix, of the vulva, of the
 25 bladder, of the kidneys (including clear-cell renal carcinoma and renal oncocytoma), squamous cell carcinomas, sarcomas such as osteosarcoma, chondrosarcoma, leiomyosarcoma, soft-tissue sarcomas, Ewing's sarcomas, gastrointestinal stromal tumours (GIST), Kaposi's sarcoma, and paediatric cancers such as rhabdomyosarcomas and neuroblastomas,
- 30 - inflammatory and immune diseases, such as rheumatoid arthritis,
- diabetes,
- obesity,
- obstructive diseases of the respiratory pathways,
- fibroses (including lung, kidney and liver fibrosis),
- 35 - Pompe's disease (Cori's type II glycogenosis),
- cardiovascular diseases, or

- age-related diseases such as neurodegenerative diseases (including Alzheimer's disease, Huntington's diseases and Parkinson's disease), loss of hearing and ocular diseases, including age-related macular degeneration (ARMD),
- epilepsy,
- 5 - parasitic diseases.

According to another of its aspects, the present invention relates to pharmaceutical compositions comprising, as active principle, a compound according to the invention. These pharmaceutical compositions contain an effective dose of at least one compound according to the invention, or a pharmaceutically acceptable salt of the
10 said compound, and also at least one pharmaceutically acceptable excipient.

The said excipients are chosen, according to the pharmaceutical form and the mode of administration desired, from the usual excipients which are known to those skilled in the art.

In the pharmaceutical compositions of the present invention for oral, sublingual,
15 subcutaneous, intramuscular, intravenous, topical, local, intratracheal, intranasal, transdermal or rectal administration, the active principle of formula (I) above, or the salt thereof, may be administered in unit administration form, as a mixture with standard pharmaceutical excipients, to man and animals for the treatment or prevention of the above disorders or diseases.

20 The appropriate unit administration forms include oral-route forms such as tablets, soft or hard gel capsules, powders, granules and oral solutions or suspensions, sublingual, buccal, intratracheal, intraocular and intranasal administration forms, inhalation forms, topical, transdermal, subcutaneous, intramuscular or intravenous administration forms, rectal administration forms and implants. For topical application, the
25 compounds according to the invention may be used in creams, gels, pomades or lotions.


By way of example, a unit administration form of a compound according to the invention in tablet form may comprise the following constituents:

	Compound according to the invention	50.0 mg
	Mannitol	223.75 mg
30	Sodium croscarmellose	6.0 mg
	Corn starch	15.0 mg
	Hydroxypropylmethylcellulose	2.25 mg
	Magnesium stearate	3.0 mg

According to another of its aspects, the present invention also relates to a method for treating or preventing the pathologies indicated above, which comprises the administration, to a patient, of an effective dose of a compound according to the invention, or a pharmaceutically acceptable salt thereof.

CLAIMS

1. A compound of formula (I):




wherein:

- each R_1 is independently (C_1-C_6) -alkyl, wherein two R_1 groups on different carbon atoms of the morpholinyl ring may be linked together to form, with the morpholinyl ring, a bicyclic heterocyclic structure,
- R_2 and R_3 are independently hydrogen or (C_1-C_6) alkyl;
- R_4 is hydrogen, (C_1-C_6) -alkyl optionally substituted with a hydroxyl group, (C_1-C_6) -alkoxy or $-L_1-R_{10}$, wherein
 - L_1 is (C_1-C_6) -alkylene, and
 - R_{10} is $-COOR_{11}$, $-CO-R_{12}$, $-OR_{13}$, $-CONR_{14}R_{15}$, a 5- or 6-atom heterocycle optionally substituted with $-R_{16}$, $-COOR_{17}$, $-CO-R_{18}$, $-OR_{19}$ or $-NR_{20}R_{21}$, or a 5- or 6-atom heterocycle optionally substituted with $-R_{22}$, $-COOR_{23}$, $-CO-R_{24}$, $-OR_{25}$ or $-CONR_{26}R_{27}$, wherein each 5- to 6-atom heterocycle is saturated, partially unsaturated or aromatic, and includes one or more heteroatoms selected from O, N and S;
- G_1 is phenylene or a 5- to 6-atom heteroarylene, optionally substituted with 1 to 4 R_5 groups, each R_5 group being independently chosen from halogen, $-OR_{30}$ and $-(C_1-C_6)$ -alkyl optionally substituted with hydroxyl;
- X_1 is $-O-$ or $-NR_{40}-$;
- X_2 is a single bond, $-CONR_{50}-$, $-CONR_{51}-O-$, $-COO-$, $-CO-$ or $-SO_2-$;
- R_6 is hydrogen or L_2-R_7 , wherein
 - L_2- is (C_1-C_6) -alkylene or (C_3-C_6) -cycloalkylene;

- R_7 is hydrogen, $-OR_{60}$, halogen, (C_1-C_6) -haloalkyl, a 5- to 6-atom heterocycle optionally substituted with one or more groups chosen from (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy and $=O$, or $-G_2-X_3-G_3$, wherein
 - X_3 is a single bond or $-O-$, $-CO-$ or $-CH_2-$;
- 5 - G_2 is a single bond or a 5- to 6-atom divalent cyclic radical optionally substituted with one or more groups R_{80} ,
- G_3 is a 4- to 8-atom ring optionally substituted with one or more groups R_{81} ,
- or $-G_2-X_3-G_3$ together form a 7- to 10-atom fused bicyclic, wherein
 - each R_{80} and R_{81} is independently halogen, $-COOR_{70}$, $-CO-R_{71}$, $-OR_{72}$, -
 - 10 $NR_{73}R_{74}$, $-CONR_{75}R_{76}$, $-CN$ or $-S(O)_p-R_{77}$,
 - (C_1-C_6) -alkyl optionally substituted with one or more R_{100} groups, and/or optionally interrupted with one or more oxygen atoms, and
 - (C_1-C_6) -alkoxy optionally substituted with one or more R_{101} groups, and/or optionally interrupted with one or more oxygen atoms,
- 15 - n is 0, 1 or 2;
- m is 0 or 1;
- p is 0, 1 or 2;
- R_{11} , R_{12} , R_{13} , R_{14} , R_{15} , R_{16} , R_{17} , R_{18} , R_{19} , R_{20} , R_{21} , R_{22} , R_{23} , R_{24} , R_{25} , R_{26} , R_{27} , R_{30} , R_{40} , R_{50} , R_{51} , R_{60} , R_{70} , R_{71} , R_{72} , R_{73} , R_{74} , R_{75} , R_{76} and R_{77} are independently H or (C_1-C_6) -
- 20 alkyl optionally substituted with a R_{90} group, and/or optionally interrupted with one or more oxygen atoms;
- R_{90} is $-OR_{91}$ or $-NR_{92}R_{93}$;
- R_{91} , R_{92} and R_{93} are independently hydrogen or (C_1-C_6) -alkyl; and
- R_{100} and R_{101} are independently halogen or hydroxyl;
- 25 or a pharmaceutically acceptable salt thereof.

2. The compound of claim 1, of formula (II):



30 or a pharmaceutically acceptable salt thereof.

3. The compound of claim 1, wherein

- R_4 is hydrogen, (C_1-C_6) -alkyl, optionally substituted with a hydroxyl group, or $-L_1-R_{10}$, wherein

5 - L_1 is (C_1-C_6) -alkylene, and


- R_{10} is $-COOR_{11}$, $-OR_{13}$, $-CONR_{14}R_{15}$, a 5- or 6-atom heterocycle optionally substituted with $-COOR_{17}$ or $-CO-R_{18}$, or a 5- or 6-atom heterocycle optionally substituted with $-R_{22}$.

10 4. The compound of claim 1, wherein G_1 is phenyl or pyridyl, each optionally substituted with an R_5 group wherein R_5 is halogen.

5. The compound of claim 1, wherein X_1 is $-NR_{40}-$.

15 6. The compound of claim 1, wherein X_2 is $-CONR_{50}-$ or $-COO-$.

7. The compound of claim 1, of formula (III):



20

wherein

- R_4 is (C_1-C_6) -alkyl;

25 - R'_5 and R''_5 are independently hydrogen or fluoro, provided that when one of R'_5 and R''_5 is fluoro, the other is hydrogen;

- X_2 is $-CONR_{50}-$ or $-COO-$;

- R_6 is $-L_2-R_7$, wherein

- $-L_2-$ is (C_1-C_6) -alkylene or (C_3-C_6) -cycloalkylene, and
- R_7 is hydrogen, $-OR_{60}$, or $-G_2-X_3-G_3$, wherein
- X_3 is a single bond or $-O-$,
- G_2 is a single bond or a cyclic divalent radical of 6 atoms,
- 5 - G_3 is a 4- to 8-atom ring optionally substituted with one or more groups R_{81} ,

or a pharmaceutically acceptable salt thereof.

8. The compound of claim 1, wherein R_4 is methyl or ethyl.

10

9. The compound of claim 1, wherein the compound is:

1	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea
2	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea
3	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
12	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-pyridin-4-ylurea
13	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[4-(4-methylpiperazin-1-yl)phenyl]urea
21	1-(4-Methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
22	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(1-methylpiperidin-4-yloxy)pyridin-3-yl]urea
29	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((S)-3-methylmorpholin-4-yl)pyridin-3-yl]urea

30	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((R)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
31	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)urea
32	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-pyrrolidin-1-ylpyridin-3-yl)urea
37	1-{4-[6-Ethyl-2-((S)-3-ethylmorpholin-4-yl)-8,8-dimethyl-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methylpyridin-3-yl)urea
38	1-Methyl-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
40	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[5-(2-methoxyethoxy)pyridin-2-yl]urea
43	1-Methyl-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
45	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-methylurea
46	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-methylurea
49	1-(5-Methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
50	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-3-ylurea
51	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-4-ylurea

52	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-4-ylurea
54	1-(5-Methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
55	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-methylpyridin-4-yl)urea
56	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-methylpyridin-4-yl)urea
58	1-(2,6-Dimethylpyridin-4-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
59	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-methylpyridin-4-yl)urea
60	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-methylpyridin-4-yl)urea
64	1-(5-Dimethylaminopyridin-2-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
66	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methylpyridin-3-yl)urea
67	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(6-methylpyridin-3-yl)urea
68	1-Pyridin-4-yl-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
69	1-Pyridin-4-yl-3-{4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea

70	1-(4-Methoxypyridin-2-yl)-3-{4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
71	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}-3-pyridin-4-ylurea
73	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-((S)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
74	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-((S)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
75	1-[4-((1S,5R)-6-Ethyl-8,8-dimethyl-2-3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl]-3-(3-fluoropyridin-4-yl)urea
76	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(3-fluoropyridin-4-yl)urea
77	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[4-(tetrahydrofuran-3-yloxy)phenyl]urea
78	1-(2-Dimethylaminopyridin-4-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
79	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(4-methoxypyridin-2-yl)urea
80	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[2-(4-methanesulfonylpiperazin-1-yl)pyridin-4-yl]urea
81	4-[5-(3-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}ureido)pyridin-2-yl]piperazine-1-carboxylic acid methyl ester
82	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-4-ylurea

83	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-4-ylurea
87	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-((R)-3-methylmorpholin-4-yl)pyridin-3-yl]urea
98	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
99	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}urea
110	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
112	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-3-ylurea
116	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methylisoxazol-3-yl)urea
124	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridin-3-yl)urea
133	1-Cyclopropyl-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
135	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-isoxazol-3-ylurea
141	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridin-3-yl)urea
150	1-Cyclopropyl-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea

159	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-isoxazol-3-ylurea
161	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methylpyridin-3-yl)urea
168	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-methylpyridin-2-yl)urea
171	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]urea
174	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]urea
184	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(4-methoxypyridin-2-yl)urea
185	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methylpyridin-3-yl)urea
186	{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}carbamic acid 2-hydroxyethyl ester
187	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-morpholin-4-ylpyridin-3-yl)urea
195	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]urea
199	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(2-methylpyridin-4-yl)urea
200	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-morpholin-4-ylpyridin-3-yl)urea


202	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(2-fluorophenyl)urea
205	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[4-(4-methylpiperazin-1-yl)phenyl]urea
207	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-methylurea
218	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[5-(4-methylpiperazin-1-yl)pyridin-2-yl]urea
221	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-pyridin-2-ylurea
223	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-3-ylurea
227	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(4-morpholin-4-ylphenyl)urea
229	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methylpyridin-3-yl)urea
231	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-pyridin-4-ylurea
235	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[4-(4-methanesulfonylpiperazin-1-yl)phenyl]urea
236	4-[4-(3-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}ureido)phenyl]piperazine-1-carboxylic acid methyl ester
237	1-[4-(1,1-Dioxo-1λ6-thiomorpholin-4-yl)phenyl]-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea

240	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-(6-methoxypyridin-3-yl)urea
241	1-(2,6-Dimethylpyridin-4-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
243	{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-3-fluorophenyl}carbamic acid 2-hydroxyethyl ester
244	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-3-yl]urea
245	1-{2-Fluoro-4-[6,8,8-trimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[6-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)pyridin-3-yl]urea
247	1-(6-Cyanopyridin-3-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
248	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((S)-3-ethylmorpholin-4-yl)pyridin-3-yl]urea
249	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-((S)-3-ethylmorpholin-4-yl)pyridin-3-yl]urea
250	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-(5-morpholin-4-ylpyridin-2-yl)urea
251	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]phenyl}-3-[5-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-2-yl]urea
252	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[(1S,4S)-6-(2-oxa-5-azabicyclo[2.2.1]hept-5-yl)pyridin-3-yl]urea

253	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[(1S,4S)-6-(2-oxa-5-azabicyclo[2.2.1]hept-5-yl)pyridin-3-yl]urea
254	1-Cyclopropyl-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
255	1-Ethyl-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
256	1-[6-(3,3-Dimethylmorpholin-4-yl)pyridin-3-yl]-3-{4-[6-ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
257	1-[6-(3,3-Dimethylmorpholin-4-yl)pyridin-3-yl]-3-{4-[6-ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea
258	1-{4-[6-Ethyl-8,8-dimethyl-2-((S)-3-methylmorpholin-4-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-3-yl]urea
259	1-{4-[6-Ethyl-8,8-dimethyl-2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[6-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-3-yl]urea
260	1-{4-[6-Ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}-3-[2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)pyridin-4-yl]urea
261	1-(4,4-Difluoro-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4'-yl)-3-{4-[6-ethyl-8,8-dimethyl-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-5,6,7,8-tetrahydro-9-oxa-1,3,6-triazabenzocyclohepten-4-yl]-2-fluorophenyl}urea


or a pharmaceutically acceptable salt thereof.

10. A process for preparing a compound of formula (I) according to claim 1, the
5 process comprising
grafting a group -X₂-R₆ onto a compound of formula (XLII) below:



to obtain the compound of formula (I).


- 11.** A process for preparing a compound of formula (I) according to claim 1, in which X_1 represents $-NR_{40}^-$ and X_2 is $-CONR_{50}^-$, the process comprising Suzuki reaction of the compound of formula (XXXV):




in which LG_5 is a leaving group,

- with a compound of formula $R_6R_{50}N-(C=O)-NR_{40}-G_1-R_{300}$, in which R_{300} is a boronic acid or boronic ester
- to obtain the compound of formula (I) wherein X_1 is $-NR_{40}^-$ and X_2 represents $-CONR_{50}^-$.

- 12.** A process for preparing a compound of formula (I) according to claim 1, the process comprising nucleophilic attack of a compound of formula (CI):



on a compound of formula (LII):



to obtain the compound of formula (I).

- 13.** A pharmaceutical composition, comprising a compound according to claim 1,
 5 or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable excipient.

