

ENGINEERED ACID ALPHA-GLUCOSIDASE VARIANTS

[0001] The present application claims priority to US Prov. Appln. Ser. No. 62/951,625, filed December 20, 2019, hereby incorporated by reference in its entirety for all purposes.

FIELD OF THE INVENTION

[0002] The present invention provides engineered acid alpha-glucosidase (GAA) polypeptides and compositions thereof. In some embodiments, the engineered GAA polypeptides have been optimized to provide increased expression, stability at neutral and acidic pH, increased uptake into cells, and activity in cell lysates. The invention also provides methods for utilization of the compositions comprising the engineered GAA polypeptides for therapeutic and other purposes.

REFERENCE TO SEQUENCE LISTING, TABLE OR COMPUTER PROGRAM

[0003] The official copy of the Sequence Listing is submitted concurrently with the specification as an ASCII formatted text file via EFS-Web, with a file name of "CX7-199WO2_ST25.txt", a creation date of December 17, 2020, and a size of 18.2 megabytes. The Sequence Listing filed via EFS-Web is part of the specification and is incorporated in its entirety by reference herein.

BACKGROUND OF THE INVENTION

[0004] Pompe disease is an autosomal recessive lysosomal storage disorder that is caused by mutations in the gene encoding acid alpha-glucosidase. This genetic defect leads to reduction of or absence of GAA in the body tissues. The resulting accumulation of glycogen in the lysosomes results in lysosomal swelling and rupture, which can lead to cell damage, organelle dysfunction, and other cellular defects. There are two forms of Pompe disease, including the classical infantile form and late-onset (childhood or adulthood) form. Disease severity is related to the amount of enzyme activity present in the cells of affected individuals. The infantile form is the most severe and rapidly progressive form, typically with GAA activity that is less than 1%, resulting in marked accumulation of glycogen in skeletal muscle, as well as heart and other tissues (See e.g., Hahn and Schänzer, *Ann. Transl. Med.*, 7:283 [2019]). In these patients, there is multi-system storage of accumulated lysosomal and non-lysosomal bound-glycogen in the heart, skeletal muscle, and brain tissue (See, Schoser, *Ann. Transl. Med.*, 7:292 [2019]). Patients present with elevated creatinine kinase levels, hypertrophic cardiomyopathy, failure to thrive, muscular hypotonia, and axial muscle weakness. If untreated, patients typically die within the first year of life due to cardiorespiratory insufficiency. Survival beyond 18 months of age is exceptional. This form is distinguished from non-classic or late-infantile Pompe disease, in which patients present with much less severe cardiac hypertrophy. Patients with late-onset Pompe disease typically experience progressive limb-girdle myopathy and respiratory dysfunction. These patients present with predominant, but not exclusive, muscle

involvement. The patients eventually become wheelchair and/or ventilator-dependent. Respiratory insufficiency is the leading cause of death in these patients. Some patients may synthesize a non-functional form of GAA, but others are incapable of producing any type of native enzyme. The human *GAA* gene encoding GAA has been localized to chromosome 17q25.2-q25.3 and has been cloned and sequenced (See, Peruzzo et al., *Ann. Transl. Med.*, 7:278-287 [2019]; and Martiniuk et al., *DNA Cell. Biol.*, 10:283-292 [1991]). Although numerous mutations in the gene have been reported, the pathological mechanisms that lead to the wide range of phenotypes observed in affected patients remains unknown. Despite the availability of enzyme replacement therapy (ERT) utilizing recombinant GAA, there remains the need for better treatment and management options for affected patients.

SUMMARY OF THE INVENTION

[0005] The present invention provides engineered acid alpha-glucosidase (GAA) polypeptides and compositions thereof. In some embodiments, the engineered GAA polypeptides have been optimized to provide increased expression, stability at neutral and acidic pH, increased uptake into cells, and activity in cell lysates. The invention also provides methods for utilization of the compositions comprising the engineered GAA polypeptides for therapeutic and other purposes.

[0006] In some embodiments, the present invention provides engineered GAA polypeptides (also referred to herein as “recombinant GAA polypeptides”) and biologically active fragments and analogs thereof having improved properties when compared to a wild-type GAA enzyme and/or a reference GAA polypeptide under essentially the same conditions. The invention is further directed to methods of using the engineered GAA polypeptides and biologically active fragments and analogs thereof in therapeutic and/or other compositions.

[0007] The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% sequence identity to SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104. In some embodiments, the present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104. In some embodiments, the present invention also provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104. In some embodiments, the

present invention further provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence consisting of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104. In some embodiments, the recombinant acid alpha-glucosidase sequences include a signal peptide sequence (e.g., SEQ ID NO: 3382 or 3384) encoded by the polynucleotides set forth in SEQ ID NO: 3381 and 3383, respectively. In some embodiments, the recombinant polynucleotides encoding the recombinant acid alpha-glucosidases of the present invention comprise a 57 base pair sequence that encodes a signal peptide. In some embodiments, the polypeptides of the recombinant acid alpha-glucosidases of the present invention comprise a 19 amino acid signal peptide. In some alternative embodiments, the recombinant polynucleotides encoding recombinant acid alpha-glucosidases do not include a sequence encoding a signal peptide. In some additional embodiments, the recombinant polypeptides comprising recombinant acid alpha-glucosidases do not include a signal peptide. It is not intended that the present invention be limited to recombinant acid alpha-glucosidase polynucleotide or polypeptide sequences comprising the signal peptide nucleotide or polypeptide sequence. It is also not intended that the present invention be limited to recombinant acid alpha-glucosidase polynucleotide or polypeptide sequences that do not comprise a signal peptide nucleotide or polypeptide sequence.

[0008] The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 2. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 27, 27/944, 28, 29/478, 30, 88, 107, 109, 109/842, 110, 113, 135, 137, 138, 148, 150, 247, 274, 276, 278, 375, 403, 414, 418, 418/499, 421, 426, 437, 444, 455, 463, 471, 471/478, 476, 489, 527, 547, 581, 610, 642, 668, 670, 692, 725/732, 750, 753, 786, 820, 862, 871, 895, 897, 930, 934, and 944, wherein the positions are numbered with reference to SEQ ID NO: 2. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from 27P, 27P/944W, 27R, 28P, 28R, 28S, 29T/478T, 30G, 30K, 30T, 88G, 88S, 107G, 107P, 109G/842E, 109P, 110G, 110L, 113S, 135A, 135Q, 137P, 138A, 148G, 148Y, 150G, 247R, 274G, 276F, 276Y, 278A, 278G, 375E, 403W, 414P, 418E/499R, 418R, 421S, 426R, 437S, 444T, 455V, 463A, 471Q/478S, 471S, 476A, 476H, 489R, 527R, 547G, 581G, 581T, 610A, 610G, 610S, 642M, 642Q, 642S, 668H, 670N, 692Q, 725N/732I, 750P, 753T, 786P, 786Y, 820E, 862G, 871E, 895R, 897V, 930R, 934R, 944G, and 944R, wherein the positions are numbered with reference to SEQ ID NO: 2. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from F27P, F27P/C944W, F27R, L28P, L28R, L28S, L29T/A478T, V30G, V30K, V30T, K88G, K88S, Q107G, Q107P, L109G/G842E, L109P, Q110G, Q110L, Q113S, S135A, S135Q,

E137P, M138A, T148G, T148Y, T150G, Q247R, D274G, A276F, A276Y, T278A, T278G, I375E, R403W, R414P, A418E/H499R, A418R, Q421S, G426R, A437S, A444T, R455V, E463A, K471Q/A478S, K471S, S476A, S476H, A489R, N527R, A547G, K581G, K581T, W610A, W610G, W610S, L642M, L642Q, L642S, S668H, L670N, T692Q, K725N/V732I, A750P, A753T, R786P, R786Y, G820E, R862G, L871E, K895R, T897V, C930R, L934R, C944G, and C944R, wherein the positions are numbered with reference to SEQ ID NO: 2. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 29/218/240/668/700/744/869, 29/218/240/700/869, 29/240/596/668/700/744/869, 29/240/596/668/869, 36/106/150/218/527/750/883/894, 106/112/150/218/414/527/793/883, 106/150/169/218/414/486/527/750/894, 106/150/169/218/414/486/527/894, 106/150/169/218/414/486/749/793/883/894, 106/150/169/218/414/486/750/793/883/894, 106/150/169/218/414/486/793/883, 106/150/169/218/414/486/894, 106/150/169/218/414/749/750/793/883, 106/150/169/218/414/749/793, 106/150/169/218/414/749/793/883, 106/150/169/218/486/527/749/793/894, 106/150/169/218/486/749/883, 106/150/169/218/486/883, 106/150/169/218/749/800, 106/150/169/414/486/749/750/883, 106/150/169/527/749/793/883, 106/150/169/749/793/883/894, 106/150/218/331/414/486/527/733/749/793, 106/150/218/414/486/642/750/793/883, 106/150/218/414/486/750/793/894, 106/150/218/414/527/749/750/883, 106/150/218/414/527/749/793/883/894, 106/150/218/414/749/750/793/883/894, 106/150/218/414/749/793/883, 106/150/218/486/527/749/894, 106/150/218/486/793/883, 106/150/218/527/749/750/793, 106/150/218/527/793/894, 106/150/218/749/750/793, 106/150/218/793, 106/150/218/793/894, 106/150/245/793/883/894, 106/150/414/749/750/793/894, 106/150/414/749/793/894, 106/150/486/527/750/793, 106/150/486/749/793/883/894, 106/150/749/793/883, 106/169/185/218/414/749/750/793, 106/191/280/402/414/444/727, 106/191/414/444/522/928/944, 106/191/414/489/928/944, 106/280/402/414/444/489/727/944, 150/169/218/414/527/793, 150/218/414/486/749/750, 150/218/414/486/750/793, 150/218/414/486/750/793/883, 150/218/414/749/750/793/894, 150/218/414/749/793, 150/218/527/749/793, 150/218/749/750/793, 150/218/749/793, 150/414/486/527/750/894, 150/414/486/749/750/793, 150/486/750/883/894, 169/486/750/793/883, 180/275/402/518/547/610/638/669/671, 180/402/431/507/547/610/669/671/793, 180/402/507/547/610/671, 191/280/402/414/444/465/842/928, 191/280/402/414/444/489/500/944, 191/280/414/444/489/500/522/842/928/944, 191/280/414/444/489/522/727/944, 191/280/414/489/842/928/944, 191/280/414/944, 191/414/522/842/944, 196/402/431/547/610/638, 218/668/700/869, 224/402/507/518/547/638/668, 269/275/431/518/547/638/668/669, 275/281/402/431/507/518/610/668, 275/281/402/431/518/547/610/669/671, 275/281/402/507/518/547/638/669/671, 275/281/402/518/547/610/638/671, 275/281/402/518/547/610/668/669/887, 275/281/402/547/610/638/669/671,

275/281/431/518/547/638/669/671, 275/281/507/547/669/671, 275/281/610/638/668/669, 275/281/671, 275/377/402/507/518/669/671/715, 275/402/431/507/547/671, 275/402/431/518/610/638/669/671/922, 275/402/507/547/610/638/668/669, 275/402/507/547/610/638/669/671, 275/402/507/547/610/671, 275/402/547/610/638/669/671, 275/402/547/638/669/671, 275/402/638/669/671, 275/431/507/518/547/668/669/671, 275/431/507/518/610/669/671, 275/431/507/547/610/638/671, 275/431/518/547/638/668, 275/431/518/610/638/669/671, 275/431/638, 275/507/518/547/610/638/668/669, 275/507/518/547/638/669/671, 275/507/547/610/638/669/671, 275/507/547/668/669/671, 275/518/671, 280/402/536/928, 281/402/507/518/547/610/638/669/671, 281/402/507/547/638/669/671, 281/402/518/547/610/638/668/669, 281/402/518/547/668, 281/431/507/518/547/610/638/668, 402/431/518/547/610/668, 402/431/518/547/671, 402/431/518/610, 402/431/547/638/671, 431/507/518/541/547/638/669/671, 431/507/518/669/671, 507/547/610, 507/547/638/669/671, 547/610/638/671, and 547/638/668, wherein the positions are numbered with reference to SEQ ID NO: 2. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from 29Q/218S/240I/668D/700F/744V/869L, 29Q/240I/596P/668D/869L, 29Q/240I/596S/668D/700F/744V/869T, 29V/218S/240I/700F/869T, 36R/106P/150S/218S/527D/750P/883H/894R, 106P/112S/150S/218S/414G/527D/793K/883H, 106P/150S/169S/218S/414G/486E/527D/750P/894R, 106P/150S/169S/218S/414G/486E/527D/894R, 106P/150S/169S/218S/414G/486E/749E/793K/883H/894R, 106P/150S/169S/218S/414G/486E/750P/793K/883H/894R, 106P/150S/169S/218S/414G/486E/793K/883H, 106P/150S/169S/218S/414G/486E/894R, 106P/150S/169S/218S/414G/749E/750P/793K/883H, 106P/150S/169S/218S/414G/749E/793K, 106P/150S/169S/218S/414G/749E/793K/883H, 106P/150S/169S/218S/486E/527D/749E/793K/894R, 106P/150S/169S/218S/486E/749E/883H, 106P/150S/169S/218S/486E/883H, 106P/150S/169S/218S/749E/800A, 106P/150S/169S/414G/486E/749E/750P/883H, 106P/150S/169S/527D/749E/793K/883H, 106P/150S/169S/749E/793K/883H/894R, 106P/150S/218S/331A/414G/486E/527D/733E/749E/793K, 106P/150S/218S/414G/486E/642F/750P/793K/883H, 106P/150S/218S/414G/486E/750P/793K/894R, 106P/150S/218S/414G/527D/749E/750P/883H, 106P/150S/218S/414G/527D/749E/793K/883H/894G, 106P/150S/218S/414G/749E/750P/793K/883H/894R, 106P/150S/218S/414G/749E/793K/883H, 106P/150S/218S/486E/527D/749E/894R, 106P/150S/218S/486E/793K/883H, 106P/150S/218S/527D/749E/750P/793K, 106P/150S/218S/527D/793K/894G, 106P/150S/218S/749E/750P/793K, 106P/150S/218S/793K, 106P/150S/218S/793K/894R, 106P/150S/245S/793K/883H/894R, 106P/150S/414G/749E/750P/793K/894R, 106P/150S/414G/749E/793K/894R, 106P/150S/486E/527D/750P/793K,

106P/150S/486E/749E/793K/883H/894G, 106P/150S/749E/793K/883H,
106P/169S/185G/218S/414G/749E/750P/793K, 106P/191R/280D/402A/414G/444P/727P,
106P/191R/414G/444P/522V/928T/944S, 106P/191R/414G/489D/928T/944S,
106P/280D/402A/414G/444P/489D/727P/944S, 150S/169S/218S/414G/527D/793K,
150S/218S/414G/486A/750P/793K, 150S/218S/414G/486E/749E/750P,
150S/218S/414G/486E/750P/793K/883H, 150S/218S/414G/749E/750P/793K/894R,
150S/218S/414G/749E/793K, 150S/218S/527D/749E/793K, 150S/218S/749E/750P/793K,
150S/218S/749E/793K, 150S/414G/486E/527D/750P/894R, 150S/414G/486E/749E/750P/793K,
150S/486E/750P/883H/894G, 169S/486E/750P/793K/883H,
180H/275M/402A/518V/547G/610R/638I/669H/671N,
180H/402A/431V/507L/547G/610R/669H/671N/793G, 180H/402A/507L/547G/610R/671N,
191R/280D/402A/414G/444P/465E/842S/928T, 191R/280D/402A/414G/444P/489D/500A/944S,
191R/280D/414G/444P/489D/500A/522V/842S/928T/944S,
191R/280D/414G/444P/489D/522V/727P/944S, 191R/280D/414G/489D/842S/928T/944S,
191R/280D/414G/944S, 191R/414G/522V/842S/944S, 196V/402A/431V/547G/610R/638I,
218S/668D/700F/869T, 224F/402A/507L/518V/547G/638I/668D,
269N/275M/431V/518V/547G/638I/668D/669H, 275M/281V/402A/431V/507L/518V/610R/668D,
275M/281V/402A/507L/518V/547G/638I/669H/671N,
275M/281V/402A/518V/547G/610R/638I/671N,
275M/281V/402A/518V/547G/610R/668D/669H/887D,
275M/281V/402A/547G/610R/638I/669H/671N, 275M/281V/507L/547G/669H/671N,
275M/281V/610R/638I/668D/669H, 275M/402A/431V/507L/547G/671N,
275M/402A/507L/547G/610R/671N, 275M/402A/547G/638I/669H/671N,
275M/431V/518V/547G/638I/668D, 275M/431V/518V/610R/638I/669H/671N, 275M/431V/638I,
275M/507L/547G/668D/669H/671N, 275V/281V/402A/431V/518V/547G/610R/669H/671N,
275V/281V/431V/518V/547G/638I/669H/671N, 275V/281V/671N,
275V/377K/402A/507L/518V/669H/671N/715G,
275V/402A/431V/518V/610R/638I/669H/671N/922L,
275V/402A/507L/547G/610R/638I/668D/669H, 275V/402A/507L/547G/610R/638I/669H/671N,
275V/402A/547G/610R/638I/669H/671N, 275V/402A/638I/669H/671N,
275V/431V/507L/518V/547G/668D/669H/671N, 275V/431V/507L/518V/610R/669H/671N,
275V/431V/507L/547G/610R/638I/671N, 275V/507L/518V/547G/610R/638I/668D/669H,
275V/507L/518V/547G/638I/669H/671N, 275V/507L/547G/610R/638I/669H/671N,
275V/518V/671N, 280D/402A/536I/928T, 281V/402A/507L/518V/547G/610R/638I/669H/671N,
281V/402A/507L/547G/638I/669H/671N, 281V/402A/518V/547G/610R/638I/668D/669H,
281V/402A/518V/547G/668D, 281V/431V/507L/518V/547G/610R/638I/668D,
402A/431V/518V/547G/610R/668D, 402A/431V/518V/547G/671N, 402A/431V/518V/610R,

402A/431V/547G/638I/671N, 431V/507L/518V/541E/547G/638I/669H/671N, 431V/507L/518V/669H/671N, 507L/547G/610R, 507L/547G/638I/669H/671N, 547G/610R/638I/671N, and 547G/638I/668D, wherein the positions are numbered with reference to SEQ ID NO: 2. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from L29Q/L218S/L240I/S668D/H700F/I744V/I869L, L29Q/L240I/A596P/S668D/I869L, L29Q/L240I/A596S/S668D/H700F/I744V/I869T, L29V/L218S/L240I/H700F/I869T, G36R/K106P/T150S/L218S/N527D/A750P/R883H/Q894R, K106P/A112S/T150S/L218S/R414G/N527D/E793K/R883H, K106P/T150S/N169S/L218S/R414G/T486E/N527D/A750P/Q894R, K106P/T150S/N169S/L218S/R414G/T486E/N527D/Q894R, K106P/T150S/N169S/L218S/R414G/T486E/Q749E/E793K/R883H/Q894R, K106P/T150S/N169S/L218S/R414G/T486E/A750P/E793K/R883H/Q894R, K106P/T150S/N169S/L218S/R414G/T486E/E793K/R883H, K106P/T150S/N169S/L218S/R414G/T486E/Q894R, K106P/T150S/N169S/L218S/R414G/Q749E/A750P/E793K/R883H, K106P/T150S/N169S/L218S/R414G/Q749E/E793K, K106P/T150S/N169S/L218S/R414G/Q749E/E793K/R883H, K106P/T150S/N169S/L218S/T486E/N527D/Q749E/E793K/Q894R, K106P/T150S/N169S/L218S/T486E/Q749E/R883H, K106P/T150S/N169S/L218S/T486E/R883H, K106P/T150S/N169S/L218S/Q749E/P800A, K106P/T150S/N169S/R414G/T486E/Q749E/A750P/R883H, K106P/T150S/N169S/N527D/Q749E/E793K/R883H, K106P/T150S/N169S/Q749E/E793K/R883H/Q894R, K106P/T150S/L218S/V331A/R414G/T486E/N527D/D733E/Q749E/E793K, K106P/T150S/L218S/R414G/T486E/L642F/A750P/E793K/R883H, K106P/T150S/L218S/R414G/T486E/A750P/E793K/Q894R, K106P/T150S/L218S/R414G/N527D/Q749E/A750P/R883H, K106P/T150S/L218S/R414G/N527D/Q749E/E793K/R883H/Q894G, K106P/T150S/L218S/R414G/Q749E/A750P/E793K/R883H/Q894R, K106P/T150S/L218S/R414G/Q749E/E793K/R883H, K106P/T150S/L218S/T486E/N527D/Q749E/Q894R, K106P/T150S/L218S/T486E/E793K/R883H, K106P/T150S/L218S/N527D/Q749E/A750P/E793K, K106P/T150S/L218S/N527D/E793K/Q894G, K106P/T150S/L218S/Q749E/A750P/E793K, K106P/T150S/L218S/E793K, K106P/T150S/L218S/E793K/Q894R, K106P/T150S/P245S/E793K/R883H/Q894R, K106P/T150S/R414G/Q749E/A750P/E793K/Q894R, K106P/T150S/R414G/Q749E/E793K/Q894R, K106P/T150S/T486E/N527D/A750P/E793K, K106P/T150S/T486E/Q749E/E793K/R883H/Q894G,

K106P/T150S/Q749E/E793K/R883H, K106P/N169S/V185G/L218S/R414G/Q749E/A750P/E793K,
K106P/H191R/G280D/S402A/R414G/A444P/S727P,
K106P/H191R/R414G/A444P/E522V/D928T/C944S, K106P/H191R/R414G/A489D/D928T/C944S,
K106P/G280D/S402A/R414G/A444P/A489D/S727P/C944S,
T150S/N169S/L218S/R414G/N527D/E793K, T150S/L218S/R414G/T486A/A750P/E793K,
T150S/L218S/R414G/T486E/Q749E/A750P, T150S/L218S/R414G/T486E/A750P/E793K/R883H,
T150S/L218S/R414G/Q749E/A750P/E793K/Q894R, T150S/L218S/R414G/Q749E/E793K,
T150S/L218S/N527D/Q749E/E793K, T150S/L218S/Q749E/A750P/E793K,
T150S/L218S/Q749E/E793K, T150S/R414G/T486E/N527D/A750P/Q894R,
T150S/R414G/T486E/Q749E/A750P/E793K, T150S/T486E/A750P/R883H/Q894G,
N169S/T486E/A750P/E793K/R883H,
N180H/L275M/S402A/I518V/A547G/W610R/V638I/L669H/S671N,
N180H/S402A/M431V/M507L/A547G/W610R/L669H/S671N/E793G,
N180H/S402A/M507L/A547G/W610R/S671N,
H191R/G280D/S402A/R414G/A444P/G465E/G842S/D928T,
H191R/G280D/S402A/R414G/A444P/A489D/D500A/C944S,
H191R/G280D/R414G/A444P/A489D/D500A/E522V/G842S/D928T/C944S,
H191R/G280D/R414G/A444P/A489D/E522V/S727P/C944S,
H191R/G280D/R414G/A489D/G842S/D928T/C944S, H191R/G280D/R414G/C944S,
H191R/R414G/E522V/G842S/C944S, A196V/S402A/M431V/A547G/W610R/V638I,
L218S/S668D/H700F/I869T, L224F/S402A/M507L/I518V/A547G/V638I/S668D,
T269N/L275M/M431V/I518V/A547G/V638I/S668D/L669H,
L275M/A281V/S402A/M431V/M507L/I518V/W610R/S668D,
L275M/A281V/S402A/M507L/I518V/A547G/V638I/L669H/S671N,
L275M/A281V/S402A/I518V/A547G/W610R/V638I/S671N,
L275M/A281V/S402A/I518V/A547G/W610R/S668D/L669H/E887D,
L275M/A281V/S402A/A547G/W610R/V638I/L669H/S671N,
L275M/A281V/M507L/A547G/L669H/S671N, L275M/A281V/W610R/V638I/S668D/L669H,
L275M/S402A/M431V/M507L/A547G/S671N, L275M/S402A/M507L/A547G/W610R/S671N,
L275M/S402A/A547G/V638I/L669H/S671N, L275M/M431V/I518V/A547G/V638I/S668D,
L275M/M431V/I518V/W610R/V638I/L669H/S671N, L275M/M431V/V638I,
L275M/M507L/A547G/S668D/L669H/S671N,
L275V/A281V/S402A/M431V/I518V/A547G/W610R/L669H/S671N,
L275V/A281V/M431V/I518V/A547G/V638I/L669H/S671N, L275V/A281V/S671N,
L275V/R377K/S402A/M507L/I518V/L669H/S671N/V715G,
L275V/S402A/M431V/I518V/W610R/V638I/L669H/S671N/P922L,
L275V/S402A/M507L/A547G/W610R/V638I/S668D/L669H,

L275V/S402A/M507L/A547G/W610R/V638I/L669H/S671N,
L275V/S402A/A547G/W610R/V638I/L669H/S671N, L275V/S402A/V638I/L669H/S671N,
L275V/M431V/M507L/I518V/A547G/S668D/L669H/S671N,
L275V/M431V/M507L/I518V/W610R/L669H/S671N,
L275V/M431V/M507L/A547G/W610R/V638I/S671N,
L275V/M507L/I518V/A547G/W610R/V638I/S668D/L669H,
L275V/M507L/I518V/A547G/V638I/L669H/S671N,
L275V/M507L/A547G/W610R/V638I/L669H/S671N, L275V/I518V/S671N,
G280D/S402A/V536I/D928T, A281V/S402A/M507L/I518V/A547G/W610R/V638I/L669H/S671N,
A281V/S402A/M507L/A547G/V638I/L669H/S671N,
A281V/S402A/I518V/A547G/W610R/V638I/S668D/L669H, A281V/S402A/I518V/A547G/S668D,
A281V/M431V/M507L/I518V/A547G/W610R/V638I/S668D,
S402A/M431V/I518V/A547G/W610R/S668D, S402A/M431V/I518V/A547G/S671N,
S402A/M431V/I518V/W610R, S402A/M431V/A547G/V638I/S671N,
M431V/M507L/I518V/G541E/A547G/V638I/L669H/S671N, M431V/M507L/I518V/L669H/S671N,
M507L/A547G/W610R, M507L/A547G/V638I/L669H/S671N, A547G/W610R/V638I/S671N, and
A547G/V638I/S668D, wherein the positions are numbered with reference to SEQ ID NO: 2. In some
embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one
or more positions selected from 27, 27/944, 28, 29/218/240/668/700/744/869, 29/218/240/700/869,
29/240/596/668/700/744/869, 29/240/596/668/869, 29/478, 30, 36/106/150/218/527/750/883/894,
106/112/150/218/414/527/793/883, 106/150/169/218/414/486/527/750/894,
106/150/169/218/414/486/527/894, 106/150/169/218/414/486/749/793/883/894,
106/150/169/218/414/486/750/793/883/894, 106/150/169/218/414/486/793/883,
106/150/169/218/414/486/894, 106/150/169/218/414/749/750/793/883,
106/150/169/218/414/749/793, 106/150/169/218/414/749/793/883, 106/150/169/218/486/749/883,
106/150/169/218/486/883, 106/150/169/414/486/749/750/883, 106/150/169/527/749/793/883,
106/150/169/749/793/883/894, 106/150/218/331/414/486/527/733/749/793,
106/150/218/414/486/642/750/793/883, 106/150/218/414/486/750/793/894,
106/150/218/414/527/749/750/883, 106/150/218/414/527/749/793/883/894,
106/150/218/414/749/750/793/883/894, 106/150/218/414/749/793/883,
106/150/218/486/527/749/894, 106/150/218/486/793/883, 106/150/218/527/793/894,
106/150/245/793/883/894, 106/150/414/749/750/793/894, 106/150/414/749/793/894,
106/150/486/527/750/793, 106/150/486/749/793/883/894, 106/150/749/793/883,
106/169/185/218/414/749/750/793, 106/191/280/402/414/444/727,
106/280/402/414/444/489/727/944, 107, 109, 109/842, 110, 135, 138, 148, 150,
150/218/414/486/749/750, 150/218/414/486/750/793, 150/218/414/486/750/793/883,
150/218/414/749/750/793/894, 150/218/414/749/793, 150/414/486/527/750/894,

150/414/486/749/750/793, 150/486/750/883/894, 169/486/750/793/883,
180/275/402/518/547/610/638/669/671, 180/402/431/507/547/610/669/671/793,
180/402/507/547/610/671, 191/280/402/414/444/489/500/944, 191/280/414/444/489/522/727/944,
191/280/414/944, 191/414/522/842/944, 196/402/431/547/610/638, 218/668/700/869,
224/402/507/518/547/638/668, 269/275/431/518/547/638/668/669, 274,
275/281/402/431/507/518/610/668, 275/281/402/431/518/547/610/669/671,
275/281/402/507/518/547/638/669/671, 275/281/402/518/547/610/638/671,
275/281/402/518/547/610/668/669/887, 275/281/402/547/610/638/669/671,
275/281/431/518/547/638/669/671, 275/281/507/547/669/671, 275/281/610/638/668/669,
275/377/402/507/518/669/671/715, 275/402/431/507/547/671,
275/402/431/518/610/638/669/671/922, 275/402/507/547/610/638/668/669,
275/402/507/547/610/638/669/671, 275/402/507/547/610/671, 275/402/547/610/638/669/671,
275/402/547/638/669/671, 275/402/638/669/671, 275/431/507/518/547/668/669/671,
275/431/507/518/610/669/671, 275/431/507/547/610/638/671, 275/431/518/547/638/668,
275/431/518/610/638/669/671, 275/431/638, 275/507/518/547/610/638/668/669,
275/507/518/547/638/669/671, 275/507/547/610/638/669/671, 275/507/547/668/669/671, 276,
281/402/507/518/547/610/638/669/671, 281/402/507/547/638/669/671,
281/402/518/547/610/638/668/669, 281/402/518/547/668, 281/431/507/518/547/610/638/668, 375,
402/431/518/547/610/668, 402/431/518/547/671, 402/431/518/610, 402/431/547/638/671, 403, 414,
418/499, 431/507/518/541/547/638/669/671, 431/507/518/669/671, 437, 471/478, 507/547/610,
507/547/638/669/671, 547, 547/610/638/671, 547/638/668, 581, 642, 670, 692, 750, 753, 820, 871,
and 944, wherein the positions are numbered with reference to SEQ ID NO: 2. In some embodiments,
the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more
positions selected from 27P, 27P/944W, 27R, 28P, 28S, 29Q/218S/240I/668D/700F/744V/869L,
29Q/240I/596P/668D/869L, 29Q/240I/596S/668D/700F/744V/869T, 29T/478T,
29V/218S/240I/700F/869T, 30G, 30K, 30T, 36R/106P/150S/218S/527D/750P/883H/894R,
106P/112S/150S/218S/414G/527D/793K/883H,
106P/150S/169S/218S/414G/486E/527D/750P/894R, 106P/150S/169S/218S/414G/486E/527D/894R,
106P/150S/169S/218S/414G/486E/749E/793K/883H/894R,
106P/150S/169S/218S/414G/486E/750P/793K/883H/894R,
106P/150S/169S/218S/414G/486E/793K/883H, 106P/150S/169S/218S/414G/486E/894R,
106P/150S/169S/218S/414G/749E/750P/793K/883H, 106P/150S/169S/218S/414G/749E/793K,
106P/150S/169S/218S/414G/749E/793K/883H, 106P/150S/169S/218S/486E/749E/883H,
106P/150S/169S/218S/486E/883H, 106P/150S/169S/414G/486E/749E/750P/883H,
106P/150S/169S/527D/749E/793K/883H, 106P/150S/169S/749E/793K/883H/894R,
106P/150S/218S/331A/414G/486E/527D/733E/749E/793K,
106P/150S/218S/414G/486E/642F/750P/793K/883H, 106P/150S/218S/414G/486E/750P/793K/894R,

106P/150S/218S/414G/527D/749E/750P/883H,
106P/150S/218S/414G/527D/749E/793K/883H/894G,
106P/150S/218S/414G/749E/750P/793K/883H/894R, 106P/150S/218S/414G/749E/793K/883H,
106P/150S/218S/486E/527D/749E/894R, 106P/150S/218S/486E/793K/883H,
106P/150S/218S/527D/793K/894G, 106P/150S/245S/793K/883H/894R,
106P/150S/414G/749E/750P/793K/894R, 106P/150S/414G/749E/793K/894R,
106P/150S/486E/527D/750P/793K, 106P/150S/486E/749E/793K/883H/894G,
106P/150S/749E/793K/883H, 106P/169S/185G/218S/414G/749E/750P/793K,
106P/191R/280D/402A/414G/444P/727P, 106P/280D/402A/414G/444P/489D/727P/944S, 107G,
109G/842E, 109P, 110G, 110L, 135Q, 138A, 148G, 148Y, 150G,
150S/218S/414G/486A/750P/793K, 150S/218S/414G/486E/749E/750P,
150S/218S/414G/486E/750P/793K/883H, 150S/218S/414G/749E/750P/793K/894R,
150S/218S/414G/749E/793K, 150S/414G/486E/527D/750P/894R,
150S/414G/486E/749E/750P/793K, 150S/486E/750P/883H/894G, 169S/486E/750P/793K/883H,
180H/275M/402A/518V/547G/610R/638I/669H/671N,
180H/402A/431V/507L/547G/610R/669H/671N/793G, 180H/402A/507L/547G/610R/671N,
191R/280D/402A/414G/444P/489D/500A/944S, 191R/280D/414G/444P/489D/522V/727P/944S,
191R/280D/414G/944S, 191R/414G/522V/842S/944S, 196V/402A/431V/547G/610R/638I,
218S/668D/700F/869T, 224F/402A/507L/518V/547G/638I/668D,
269N/275M/431V/518V/547G/638I/668D/669H, 274G,
275M/281V/402A/431V/507L/518V/610R/668D,
275M/281V/402A/507L/518V/547G/638I/669H/671N,
275M/281V/402A/518V/547G/610R/638I/671N,
275M/281V/402A/518V/547G/610R/668D/669H/887D,
275M/281V/402A/547G/610R/638I/669H/671N, 275M/281V/507L/547G/669H/671N,
275M/281V/610R/638I/668D/669H, 275M/402A/431V/507L/547G/671N,
275M/402A/507L/547G/610R/671N, 275M/402A/547G/638I/669H/671N,
275M/431V/518V/547G/638I/668D, 275M/431V/518V/610R/638I/669H/671N, 275M/431V/638I,
275M/507L/547G/668D/669H/671N, 275V/281V/402A/431V/518V/547G/610R/669H/671N,
275V/281V/431V/518V/547G/638I/669H/671N, 275V/377K/402A/507L/518V/669H/671N/715G,
275V/402A/431V/518V/610R/638I/669H/671N/922L,
275V/402A/507L/547G/610R/638I/668D/669H, 275V/402A/507L/547G/610R/638I/669H/671N,
275V/402A/547G/610R/638I/669H/671N, 275V/402A/638I/669H/671N,
275V/431V/507L/518V/547G/668D/669H/671N, 275V/431V/507L/518V/610R/669H/671N,
275V/431V/507L/547G/610R/638I/671N, 275V/507L/518V/547G/610R/638I/668D/669H,
275V/507L/518V/547G/638I/669H/671N, 275V/507L/547G/610R/638I/669H/671N, 276Y,
281V/402A/507L/518V/547G/610R/638I/669H/671N, 281V/402A/507L/547G/638I/669H/671N,

281V/402A/518V/547G/610R/638I/668D/669H, 281V/402A/518V/547G/668D,
281V/431V/507L/518V/547G/610R/638I/668D, 375E, 402A/431V/518V/547G/610R/668D,
402A/431V/518V/547G/671N, 402A/431V/518V/610R, 402A/431V/547G/638I/671N, 403W, 414P,
418E/499R, 431V/507L/518V/541E/547G/638I/669H/671N, 431V/507L/518V/669H/671N, 437S,
471Q/478S, 507L/547G/610R, 507L/547G/638I/669H/671N, 547G, 547G/610R/638I/671N,
547G/638I/668D, 581G, 581T, 642Q, 642S, 670N, 692Q, 750P, 753T, 820E, 871E, and 944G,
wherein the positions are numbered with reference to SEQ ID NO: 2. In some embodiments, the acid
alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions
selected from F27P, F27P/C944W, F27R, L28P, L28S,
L29Q/L218S/L240I/S668D/H700F/I744V/I869L, L29Q/L240I/A596P/S668D/I869L,
L29Q/L240I/A596S/S668D/H700F/I744V/I869T, L29T/A478T, L29V/L218S/L240I/H700F/I869T,
V30G, V30K, V30T, G36R/K106P/T150S/L218S/N527D/A750P/R883H/Q894R,
K106P/A112S/T150S/L218S/R414G/N527D/E793K/R883H,
K106P/T150S/N169S/L218S/R414G/T486E/N527D/A750P/Q894R,
K106P/T150S/N169S/L218S/R414G/T486E/N527D/Q894R,
K106P/T150S/N169S/L218S/R414G/T486E/Q749E/E793K/R883H/Q894R,
K106P/T150S/N169S/L218S/R414G/T486E/A750P/E793K/R883H/Q894R,
K106P/T150S/N169S/L218S/R414G/T486E/E793K/R883H,
K106P/T150S/N169S/L218S/R414G/T486E/Q894R,
K106P/T150S/N169S/L218S/R414G/Q749E/A750P/E793K/R883H,
K106P/T150S/N169S/L218S/R414G/Q749E/E793K,
K106P/T150S/N169S/L218S/R414G/Q749E/E793K/R883H,
K106P/T150S/N169S/L218S/T486E/Q749E/R883H, K106P/T150S/N169S/L218S/T486E/R883H,
K106P/T150S/N169S/R414G/T486E/Q749E/A750P/R883H,
K106P/T150S/N169S/N527D/Q749E/E793K/R883H,
K106P/T150S/N169S/Q749E/E793K/R883H/Q894R,
K106P/T150S/L218S/V331A/R414G/T486E/N527D/D733E/Q749E/E793K,
K106P/T150S/L218S/R414G/T486E/L642F/A750P/E793K/R883H,
K106P/T150S/L218S/R414G/T486E/A750P/E793K/Q894R,
K106P/T150S/L218S/R414G/N527D/Q749E/A750P/R883H,
K106P/T150S/L218S/R414G/N527D/Q749E/E793K/R883H/Q894G,
K106P/T150S/L218S/R414G/Q749E/A750P/E793K/R883H/Q894R,
K106P/T150S/L218S/R414G/Q749E/E793K/R883H,
K106P/T150S/L218S/T486E/N527D/Q749E/Q894R, K106P/T150S/L218S/T486E/E793K/R883H,
K106P/T150S/L218S/N527D/E793K/Q894G, K106P/T150S/P245S/E793K/R883H/Q894R,
K106P/T150S/R414G/Q749E/A750P/E793K/Q894R, K106P/T150S/R414G/Q749E/E793K/Q894R,
K106P/T150S/T486E/N527D/A750P/E793K, K106P/T150S/T486E/Q749E/E793K/R883H/Q894G,

K106P/T150S/Q749E/E793K/R883H, K106P/N169S/V185G/L218S/R414G/Q749E/A750P/E793K,
K106P/H191R/G280D/S402A/R414G/A444P/S727P,
K106P/G280D/S402A/R414G/A444P/A489D/S727P/C944S, Q107G, L109G/G842E, L109P,
Q110G, Q110L, S135Q, M138A, T148G, T148Y, T150G,
T150S/L218S/R414G/T486A/A750P/E793K, T150S/L218S/R414G/T486E/Q749E/A750P,
T150S/L218S/R414G/T486E/A750P/E793K/R883H,
T150S/L218S/R414G/Q749E/A750P/E793K/Q894R, T150S/L218S/R414G/Q749E/E793K,
T150S/R414G/T486E/N527D/A750P/Q894R, T150S/R414G/T486E/Q749E/A750P/E793K,
T150S/T486E/A750P/R883H/Q894G, N169S/T486E/A750P/E793K/R883H,
N180H/L275M/S402A/I518V/A547G/W610R/V638I/L669H/S671N,
N180H/S402A/M431V/M507L/A547G/W610R/L669H/S671N/E793G,
N180H/S402A/M507L/A547G/W610R/S671N,
H191R/G280D/S402A/R414G/A444P/A489D/D500A/C944S,
H191R/G280D/R414G/A444P/A489D/E522V/S727P/C944S, H191R/G280D/R414G/C944S,
H191R/R414G/E522V/G842S/C944S, A196V/S402A/M431V/A547G/W610R/V638I,
L218S/S668D/H700F/I869T, L224F/S402A/M507L/I518V/A547G/V638I/S668D,
T269N/L275M/M431V/I518V/A547G/V638I/S668D/L669H, D274G,
L275M/A281V/S402A/M431V/M507L/I518V/W610R/S668D,
L275M/A281V/S402A/M507L/I518V/A547G/V638I/L669H/S671N,
L275M/A281V/S402A/I518V/A547G/W610R/V638I/S671N,
L275M/A281V/S402A/I518V/A547G/W610R/S668D/L669H/E887D,
L275M/A281V/S402A/A547G/W610R/V638I/L669H/S671N,
L275M/A281V/M507L/A547G/L669H/S671N, L275M/A281V/W610R/V638I/S668D/L669H,
L275M/S402A/M431V/M507L/A547G/S671N, L275M/S402A/M507L/A547G/W610R/S671N,
L275M/S402A/A547G/V638I/L669H/S671N, L275M/M431V/I518V/A547G/V638I/S668D,
L275M/M431V/I518V/W610R/V638I/L669H/S671N, L275M/M431V/V638I,
L275M/M507L/A547G/S668D/L669H/S671N,
L275V/A281V/S402A/M431V/I518V/A547G/W610R/L669H/S671N,
L275V/A281V/M431V/I518V/A547G/V638I/L669H/S671N,
L275V/R377K/S402A/M507L/I518V/L669H/S671N/V715G,
L275V/S402A/M431V/I518V/W610R/V638I/L669H/S671N/P922L,
L275V/S402A/M507L/A547G/W610R/V638I/S668D/L669H,
L275V/S402A/M507L/A547G/W610R/V638I/L669H/S671N,
L275V/S402A/A547G/W610R/V638I/L669H/S671N, L275V/S402A/V638I/L669H/S671N,
L275V/M431V/M507L/I518V/A547G/S668D/L669H/S671N,
L275V/M431V/M507L/I518V/W610R/L669H/S671N,
L275V/M431V/M507L/A547G/W610R/V638I/S671N,

L275V/M507L/I518V/A547G/W610R/V638I/S668D/L669H,
 L275V/M507L/I518V/A547G/V638I/L669H/S671N,
 L275V/M507L/A547G/W610R/V638I/L669H/S671N, A276Y,
 A281V/S402A/M507L/I518V/A547G/W610R/V638I/L669H/S671N,
 A281V/S402A/M507L/A547G/V638I/L669H/S671N,
 A281V/S402A/I518V/A547G/W610R/V638I/S668D/L669H, A281V/S402A/I518V/A547G/S668D,
 A281V/M431V/M507L/I518V/A547G/W610R/V638I/S668D, I375E,
 S402A/M431V/I518V/A547G/W610R/S668D, S402A/M431V/I518V/A547G/S671N,
 S402A/M431V/I518V/W610R, S402A/M431V/A547G/V638I/S671N, R403W, R414P,
 A418E/H499R, M431V/M507L/I518V/G541E/A547G/V638I/L669H/S671N,
 M431V/M507L/I518V/L669H/S671N, A437S, K471Q/A478S, M507L/A547G/W610R,
 M507L/A547G/V638I/L669H/S671N, A547G, A547G/W610R/V638I/S671N, A547G/V638I/S668D,
 K581G, K581T, L642Q, L642S, L670N, T692Q, A750P, A753T, G820E, L871E, and C944G,
 wherein the positions are numbered with reference to SEQ ID NO: 2.

[0009] The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 6. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 27, 27/944, 28, 29/218/240/668/700/744/869, 29/218/240/700/869, 29/240/596/668/700/744/869, 29/240/596/668/869, 29/478, 30, 36/106/150/218/527/750/883/894, 88, 106/112/150/218/414/527/793/883, 106/150/169/218/414/486/527/750/894, 106/150/169/218/414/486/527/894, 106/150/169/218/414/486/749/793/883/894, 106/150/169/218/414/486/750/793/883/894, 106/150/169/218/414/486/793/883, 106/150/169/218/414/486/894, 106/150/169/218/414/749/750/793/883, 106/150/169/218/414/749/793, 106/150/169/218/414/749/793/883, 106/150/169/218/486/527/749/793/894, 106/150/169/218/486/749/883, 106/150/169/218/486/883, 106/150/169/218/749/800, 106/150/169/414/486/749/750/883, 106/150/169/527/749/793/883, 106/150/169/749/793/883/894, 106/150/218/331/414/486/527/733/749/793, 106/150/218/414/486/642/750/793/883, 106/150/218/414/486/750/793/894, 106/150/218/414/527/749/750/883, 106/150/218/414/527/749/793/883/894, 106/150/218/414/749/750/793/883/894, 106/150/218/486/527/749/894, 106/150/218/486/793/883, 106/150/218/527/749/750/793, 106/150/218/527/793/894, 106/150/218/749/750/793, 106/150/218/793, 106/150/218/793/894, 106/150/245/793/883/894, 106/150/414/749/750/793/894, 106/150/414/749/793/894, 106/150/486/527/750/793, 106/150/486/749/793/883/894, 106/150/749/793/883, 106/169/185/218/414/749/750/793, 106/191/280/402/414/444/727,

106/191/414/444/522/928/944, 106/191/414/489/928/944, 106/280/402/414/444/489/727/944, 107, 109, 109/842, 110, 113, 135, 137, 138, 148, 150, 150/169/218/414/527/793, 150/218/414/486/749/750, 150/218/414/486/750/793, 150/218/414/486/750/793/883, 150/218/414/749/750/793/894, 150/218/414/749/793, 150/218/527/749/793, 150/218/749/750/793, 150/218/749/793, 150/414/486/527/750/894, 150/414/486/749/750/793, 150/486/750/883/894, 169/486/750/793/883, 180/275/402/518/547/610/638/669/671, 180/402/431/507/547/610/669/671/793, 180/402/507/547/610/671, 191/280/402/414/444/465/842/928, 191/280/402/414/444/489/500/944, 191/280/414/444/489/500/522/842/928/944, 191/280/414/444/489/522/727/944, 191/280/414/489/842/928/944, 191/280/414/944, 191/414/522/842/944, 196/402/431/547/610/638, 218/668/700/869, 224/402/507/518/547/638/668, 247, 269/275/431/518/547/638/668/669, 274, 275/281/402/431/507/518/610/668, 275/281/402/431/518/547/610/669/671, 275/281/402/507/518/547/638/669/671, 275/281/402/518/547/610/638/671, 275/281/402/518/547/610/668/669/887, 275/281/402/547/610/638/669/671, 275/281/431/518/547/638/669/671, 275/281/507/547/669/671, 275/281/610/638/668/669, 275/281/671, 275/377/402/507/518/669/671/715, 275/402/431/507/547/671, 275/402/431/518/610/638/669/671/922, 275/402/507/547/610/638/668/669, 275/402/507/547/610/638/669/671, 275/402/507/547/610/671, 275/402/547/610/638/669/671, 275/402/547/638/669/671, 275/402/638/669/671, 275/431/507/518/547/668/669/671, 275/431/507/518/610/669/671, 275/431/507/547/610/638/671, 275/431/518/547/638/668, 275/431/518/610/638/669/671, 275/431/638, 275/507/518/547/610/638/668/669, 275/507/518/547/638/669/671, 275/507/547/610/638/669/671, 275/507/547/668/669/671, 275/518/671, 276, 278, 280/402/536/928, 281/402/507/518/547/610/638/669/671, 281/402/507/547/638/669/671, 281/402/518/547/610/638/668/669, 281/402/518/547/668, 281/431/507/518/547/610/638/668, 375, 402/431/518/547/610/668, 402/431/518/547/671, 402/431/518/610, 402/431/547/638/671, 403, 414, 418, 418/499, 421, 426, 431/507/518/541/547/638/669/671, 431/507/518/669/671, 437, 444, 455, 463, 471, 471/478, 476, 489, 507/547/610, 507/547/638/669/671, 527, 547, 547/610/638/671, 547/638/668, 581, 610, 642, 668, 670, 692, 725/732, 750, 753, 786, 820, 862, 871, 895, 897, 934, and 944, wherein the positions are numbered with reference to SEQ ID NO: 6. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 27P, 27P/944W, 27R, 28P, 28R, 28S, 29Q/218S/240I/668D/700F/744V/869L, 29Q/240I/596P/668D/869L, 29Q/240I/596S/668D/700F/744V/869T, 29T/478T, 29V/218S/240I/700F/869T, 30G, 30K, 30T, 36R/106P/150S/218S/527D/750P/883H/894R, 88G, 88S, 106P/112S/150S/218S/414G/527D/793K/883H, 106P/150S/169S/218S/414G/486E/527D/750P/894R, 106P/150S/169S/218S/414G/486E/527D/894R, 106P/150S/169S/218S/414G/486E/749E/793K/883H/894R,

106P/150S/169S/218S/414G/486E/750P/793K/883H/894R,
106P/150S/169S/218S/414G/486E/793K/883H, 106P/150S/169S/218S/414G/486E/894R,
106P/150S/169S/218S/414G/749E/750P/793K/883H, 106P/150S/169S/218S/414G/749E/793K,
106P/150S/169S/218S/414G/749E/793K/883H, 106P/150S/169S/218S/486E/527D/749E/793K/894R,
106P/150S/169S/218S/486E/749E/883H, 106P/150S/169S/218S/486E/883H,
106P/150S/169S/218S/749E/800A, 106P/150S/169S/414G/486E/749E/750P/883H,
106P/150S/169S/527D/749E/793K/883H, 106P/150S/169S/749E/793K/883H/894R,
106P/150S/218S/331A/414G/486E/527D/733E/749E/793K,
106P/150S/218S/414G/486E/642F/750P/793K/883H, 106P/150S/218S/414G/486E/750P/793K/894R,
106P/150S/218S/414G/527D/749E/750P/883H,
106P/150S/218S/414G/527D/749E/793K/883H/894G,
106P/150S/218S/414G/749E/750P/793K/883H/894R, 106P/150S/218S/414G/749E/793K/883H,
106P/150S/218S/486E/527D/749E/894R, 106P/150S/218S/486E/793K/883H,
106P/150S/218S/527D/749E/750P/793K, 106P/150S/218S/527D/793K/894G,
106P/150S/218S/749E/750P/793K, 106P/150S/218S/793K, 106P/150S/218S/793K/894R,
106P/150S/245S/793K/883H/894R, 106P/150S/414G/749E/750P/793K/894R,
106P/150S/414G/749E/793K/894R, 106P/150S/486E/527D/750P/793K,
106P/150S/486E/749E/793K/883H/894G, 106P/150S/749E/793K/883H,
106P/169S/185G/218S/414G/749E/750P/793K, 106P/191R/280D/402A/414G/444P/727P,
106P/191R/414G/444P/522V/928T/944S, 106P/191R/414G/489D/928T/944S,
106P/280D/402A/414G/444P/489D/727P/944S, 107G, 107P, 109G/842E, 109P, 110G, 110L, 113S,
135A, 135Q, 137P, 138A, 148G, 148Y, 150G, 150S/169S/218S/414G/527D/793K,
150S/218S/414G/486A/750P/793K, 150S/218S/414G/486E/749E/750P,
150S/218S/414G/486E/750P/793K/883H, 150S/218S/414G/749E/750P/793K/894R,
150S/218S/414G/749E/793K, 150S/218S/527D/749E/793K, 150S/218S/749E/750P/793K,
150S/218S/749E/793K, 150S/414G/486E/527D/750P/894R, 150S/414G/486E/749E/750P/793K,
150S/486E/750P/883H/894G, 169S/486E/750P/793K/883H,
180H/275M/402A/518V/547G/610R/638I/669H/671N,
180H/402A/431V/507L/547G/610R/669H/671N/793G, 180H/402A/507L/547G/610R/671N,
191R/280D/402A/414G/444P/465E/842S/928T, 191R/280D/402A/414G/444P/489D/500A/944S,
191R/280D/414G/444P/489D/500A/522V/842S/928T/944S,
191R/280D/414G/444P/489D/522V/727P/944S, 191R/280D/414G/489D/842S/928T/944S,
191R/280D/414G/944S, 191R/414G/522V/842S/944S, 196V/402A/431V/547G/610R/638I,
218S/668D/700F/869T, 224F/402A/507L/518V/547G/638I/668D, 247R,
269N/275M/431V/518V/547G/638I/668D/669H, 274G,
275M/281V/402A/431V/507L/518V/610R/668D,
275M/281V/402A/507L/518V/547G/638I/669H/671N,

275M/281V/402A/518V/547G/610R/638I/671N,
275M/281V/402A/518V/547G/610R/668D/669H/887D,
275M/281V/402A/547G/610R/638I/669H/671N, 275M/281V/507L/547G/669H/671N,
275M/281V/610R/638I/668D/669H, 275M/402A/431V/507L/547G/671N,
275M/402A/507L/547G/610R/671N, 275M/402A/547G/638I/669H/671N,
275M/431V/518V/547G/638I/668D, 275M/431V/518V/610R/638I/669H/671N, 275M/431V/638I,
275M/507L/547G/668D/669H/671N, 275V/281V/402A/431V/518V/547G/610R/669H/671N,
275V/281V/431V/518V/547G/638I/669H/671N, 275V/281V/671N,
275V/377K/402A/507L/518V/669H/671N/715G,
275V/402A/431V/518V/610R/638I/669H/671N/922L,
275V/402A/507L/547G/610R/638I/668D/669H, 275V/402A/507L/547G/610R/638I/669H/671N,
275V/402A/547G/610R/638I/669H/671N, 275V/402A/638I/669H/671N,
275V/431V/507L/518V/547G/668D/669H/671N, 275V/431V/507L/518V/610R/669H/671N,
275V/431V/507L/547G/610R/638I/671N, 275V/507L/518V/547G/610R/638I/668D/669H,
275V/507L/518V/547G/638I/669H/671N, 275V/507L/547G/610R/638I/669H/671N,
275V/518V/671N, 276F, 276Y, 278A, 278G, 280D/402A/536I/928T,
281V/402A/507L/518V/547G/610R/638I/669H/671N, 281V/402A/507L/547G/638I/669H/671N,
281V/402A/518V/547G/610R/638I/668D/669H, 281V/402A/518V/547G/668D,
281V/431V/507L/518V/547G/610R/638I/668D, 375E, 402A/431V/518V/547G/610R/668D,
402A/431V/518V/547G/671N, 402A/431V/518V/610R, 402A/431V/547G/638I/671N, 403W, 414P,
418E/499R, 418R, 421S, 426R, 431V/507L/518V/541E/547G/638I/669H/671N,
431V/507L/518V/669H/671N, 437S, 444T, 455V, 463A, 471Q/478S, 471S, 476A, 476H, 489R,
507L/547G/610R, 507L/547G/638I/669H/671N, 527R, 547G, 547G/610R/638I/671N,
547G/638I/668D, 581G, 581T, 610A, 610G, 610S, 642M, 642Q, 642S, 668H, 670N, 692Q,
725N/732I, 750P, 753T, 786P, 786Y, 820E, 862G, 871E, 895R, 897V, 934R, 944G, and 944R,
wherein the positions are numbered with reference to SEQ ID NO: 6. In some embodiments, the acid
alpha-glucosidase comprises at least one substitution at position or set of positions selected from
F27P, F27P/C944W, F27R, L28P, L28R, L28S, L29Q/L218S/L240I/S668D/H700F/I744V/I869L,
L29Q/L240I/A596P/S668D/I869L, L29Q/L240I/A596S/S668D/H700F/I744V/I869T, L29T/A478T,
L29V/L218S/L240I/H700F/I869T, V30G, V30K, V30T,
G36R/K106P/T150S/L218S/N527D/A750P/R883H/Q894R, K88G, K88S,
K106P/A112S/T150S/L218S/R414G/N527D/E793K/R883H,
K106P/T150S/N169S/L218S/R414G/T486E/N527D/A750P/Q894R,
K106P/T150S/N169S/L218S/R414G/T486E/N527D/Q894R,
K106P/T150S/N169S/L218S/R414G/T486E/Q749E/E793K/R883H/Q894R,
K106P/T150S/N169S/L218S/R414G/T486E/A750P/E793K/R883H/Q894R,
K106P/T150S/N169S/L218S/R414G/T486E/E793K/R883H,

K106P/T150S/N169S/L218S/R414G/T486E/Q894R,
K106P/T150S/N169S/L218S/R414G/Q749E/A750P/E793K/R883H,
K106P/T150S/N169S/L218S/R414G/Q749E/E793K,
K106P/T150S/N169S/L218S/R414G/Q749E/E793K/R883H,
K106P/T150S/N169S/L218S/T486E/N527D/Q749E/E793K/Q894R,
K106P/T150S/N169S/L218S/T486E/Q749E/R883H, K106P/T150S/N169S/L218S/T486E/R883H,
K106P/T150S/N169S/L218S/Q749E/P800A,
K106P/T150S/N169S/R414G/T486E/Q749E/A750P/R883H,
K106P/T150S/N169S/N527D/Q749E/E793K/R883H,
K106P/T150S/N169S/Q749E/E793K/R883H/Q894R,
K106P/T150S/L218S/V331A/R414G/T486E/N527D/D733E/Q749E/E793K,
K106P/T150S/L218S/R414G/T486E/L642F/A750P/E793K/R883H,
K106P/T150S/L218S/R414G/T486E/A750P/E793K/Q894R,
K106P/T150S/L218S/R414G/N527D/Q749E/A750P/R883H,
K106P/T150S/L218S/R414G/N527D/Q749E/E793K/R883H/Q894G,
K106P/T150S/L218S/R414G/Q749E/A750P/E793K/R883H/Q894R,
K106P/T150S/L218S/R414G/Q749E/E793K/R883H,
K106P/T150S/L218S/T486E/N527D/Q749E/Q894R, K106P/T150S/L218S/T486E/E793K/R883H,
K106P/T150S/L218S/N527D/Q749E/A750P/E793K, K106P/T150S/L218S/N527D/E793K/Q894G,
K106P/T150S/L218S/Q749E/A750P/E793K, K106P/T150S/L218S/E793K,
K106P/T150S/L218S/E793K/Q894R, K106P/T150S/P245S/E793K/R883H/Q894R,
K106P/T150S/R414G/Q749E/A750P/E793K/Q894R, K106P/T150S/R414G/Q749E/E793K/Q894R,
K106P/T150S/T486E/N527D/A750P/E793K, K106P/T150S/T486E/Q749E/E793K/R883H/Q894G,
K106P/T150S/Q749E/E793K/R883H, K106P/N169S/V185G/L218S/R414G/Q749E/A750P/E793K,
K106P/H191R/G280D/S402A/R414G/A444P/S727P,
K106P/H191R/R414G/A444P/E522V/D928T/C944S, K106P/H191R/R414G/A489D/D928T/C944S,
K106P/G280D/S402A/R414G/A444P/A489D/S727P/C944S, Q107G, Q107P, L109G/G842E, L109P,
Q110G, Q110L, Q113S, S135A, S135Q, E137P, M138A, T148G, T148Y, T150G,
T150S/N169S/L218S/R414G/N527D/E793K, T150S/L218S/R414G/T486A/A750P/E793K,
T150S/L218S/R414G/T486E/Q749E/A750P, T150S/L218S/R414G/T486E/A750P/E793K/R883H,
T150S/L218S/R414G/Q749E/A750P/E793K/Q894R, T150S/L218S/R414G/Q749E/E793K,
T150S/L218S/N527D/Q749E/E793K, T150S/L218S/Q749E/A750P/E793K,
T150S/L218S/Q749E/E793K, T150S/R414G/T486E/N527D/A750P/Q894R,
T150S/R414G/T486E/Q749E/A750P/E793K, T150S/T486E/A750P/R883H/Q894G,
N169S/T486E/A750P/E793K/R883H,
N180H/L275M/S402A/I518V/A547G/W610R/V638I/L669H/S671N,
N180H/S402A/M431V/M507L/A547G/W610R/L669H/S671N/E793G,

N180H/S402A/M507L/A547G/W610R/S671N,
H191R/G280D/S402A/R414G/A444P/G465E/G842S/D928T,
H191R/G280D/S402A/R414G/A444P/A489D/D500A/C944S,
H191R/G280D/R414G/A444P/A489D/D500A/E522V/G842S/D928T/C944S,
H191R/G280D/R414G/A444P/A489D/E522V/S727P/C944S,
H191R/G280D/R414G/A489D/G842S/D928T/C944S, H191R/G280D/R414G/C944S,
H191R/R414G/E522V/G842S/C944S, A196V/S402A/M431V/A547G/W610R/V638I,
L218S/S668D/H700F/I869T, L224F/S402A/M507L/I518V/A547G/V638I/S668D, Q247R,
T269N/L275M/M431V/I518V/A547G/V638I/S668D/L669H, D274G,
L275M/A281V/S402A/M431V/M507L/I518V/W610R/S668D,
L275M/A281V/S402A/M507L/I518V/A547G/V638I/L669H/S671N,
L275M/A281V/S402A/I518V/A547G/W610R/V638I/S671N,
L275M/A281V/S402A/I518V/A547G/W610R/S668D/L669H/E887D,
L275M/A281V/S402A/A547G/W610R/V638I/L669H/S671N,
L275M/A281V/M507L/A547G/L669H/S671N, L275M/A281V/W610R/V638I/S668D/L669H,
L275M/S402A/M431V/M507L/A547G/S671N, L275M/S402A/M507L/A547G/W610R/S671N,
L275M/S402A/A547G/V638I/L669H/S671N, L275M/M431V/I518V/A547G/V638I/S668D,
L275M/M431V/I518V/W610R/V638I/L669H/S671N, L275M/M431V/V638I,
L275M/M507L/A547G/S668D/L669H/S671N,
L275V/A281V/S402A/M431V/I518V/A547G/W610R/L669H/S671N,
L275V/A281V/M431V/I518V/A547G/V638I/L669H/S671N, L275V/A281V/S671N,
L275V/R377K/S402A/M507L/I518V/L669H/S671N/V715G,
L275V/S402A/M431V/I518V/W610R/V638I/L669H/S671N/P922L,
L275V/S402A/M507L/A547G/W610R/V638I/S668D/L669H,
L275V/S402A/M507L/A547G/W610R/V638I/L669H/S671N,
L275V/S402A/A547G/W610R/V638I/L669H/S671N, L275V/S402A/V638I/L669H/S671N,
L275V/M431V/M507L/I518V/A547G/S668D/L669H/S671N,
L275V/M431V/M507L/I518V/W610R/L669H/S671N,
L275V/M431V/M507L/A547G/W610R/V638I/S671N,
L275V/M507L/I518V/A547G/W610R/V638I/S668D/L669H,
L275V/M507L/I518V/A547G/V638I/L669H/S671N,
L275V/M507L/A547G/W610R/V638I/L669H/S671N, L275V/I518V/S671N, A276F, A276Y,
T278A, T278G, G280D/S402A/V536I/D928T,
A281V/S402A/M507L/I518V/A547G/W610R/V638I/L669H/S671N,
A281V/S402A/M507L/A547G/V638I/L669H/S671N,
A281V/S402A/I518V/A547G/W610R/V638I/S668D/L669H, A281V/S402A/I518V/A547G/S668D,
A281V/M431V/M507L/I518V/A547G/W610R/V638I/S668D, I375E,

S402A/M431V/I518V/A547G/W610R/S668D, S402A/M431V/I518V/A547G/S671N, S402A/M431V/I518V/W610R, S402A/M431V/A547G/V638I/S671N, R403W, R414P, A418E/H499R, A418R, Q421S, G426R, M431V/M507L/I518V/G541E/A547G/V638I/L669H/S671N, M431V/M507L/I518V/L669H/S671N, A437S, A444T, R455V, E463A, K471Q/A478S, K471S, S476A, S476H, A489R, M507L/A547G/W610R, M507L/A547G/V638I/L669H/S671N, N527R, A547G, A547G/W610R/V638I/S671N, A547G/V638I/S668D, K581G, K581T, W610A, W610G, W610S, L642M, L642Q, L642S, S668H, L670N, T692Q, K725N/V732I, A750P, A753T, R786P, R786Y, G820E, R862G, L871E, K895R, T897V, L934R, C944G, and C944R, wherein the positions are numbered with reference to SEQ ID NO: 6.

[0010] The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 8. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 4, 27, 27/28/489, 27/418/478, 28, 28/29, 28/29/113/135/138, 28/29/113/135/418, 28/29/135, 28/29/418, 29/113/126/135/193, 29/113/135, 29/113/135/455, 29/113/138, 29/148, 29/478, 106, 106/138/218/431/671/749, 106/218/281, 106/218/455, 106/218/455/507/749, 106/489/671, 106/638, 106/671/934, 113, 113/135/418, 113/418/455/478/581, 113/418/478/489/581, 135, 135/148/150/418, 135/478/489/581, 135/489, 135/944, 138/218/668/671, 138/218/749/934, 138/671/749/934, 157, 218, 218/281, 218/281/431, 218/281/671, 218/431, 218/431/489/507/749/934, 218/455, 218/507/749, 218/507/934, 218/638/671, 218/749, 281/431/489/668, 345/934, 418, 418/489, 431/668/671, 489/638/934, 489/671/934, 489/749, 489/934, 507/668, 507/671/934, 671/749, 671/934, and 749/784, wherein the positions are numbered with reference to SEQ ID NO: 8. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from 4H, 27P/28S/489R, 27P/418E/478T, 27R, 28S, 28S/29T, 28S/29T/113S/135Q/138A, 28S/29T/113S/135Q/418E, 28S/29T/135Q, 28S/29T/418E, 29T/113S/126Q/135Q/193Q, 29T/113S/135Q, 29T/113S/135Q/455V, 29T/113S/138A, 29T/148G, 29T/478T, 106P, 106P/138A/218S/431V/671N/749E, 106P/218S/281V, 106P/218S/455V, 106P/218S/455V/507L/749E, 106P/489R/671N, 106P/638I, 106P/671N/934R, 113S, 113S/135Q/418E, 113S/418E/455V/478T/581T, 113S/418E/478T/489R/581T, 135P/944Y, 135Q, 135Q/148G/150G/418E, 135Q/478T/489R/581T, 135Q/489R, 138A/218S/668D/671N, 138A/218S/749E/934R, 138A/671N/749E/934R, 157M, 218S, 218S/281V, 218S/281V/431V, 218S/281V/671N, 218S/431V, 218S/431V/489R/507L/749E/934R, 218S/455V, 218S/507L/749E, 218S/507L/934R, 218S/638I/671N, 218S/749E, 281V/431V/489R/668D, 345K/934R, 418E, 418E/489R, 431V/668D/671N, 489R/638I/934R, 489R/671N/934R, 489R/749E, 489R/934R, 507L/668D, 507L/671N/934R, 671N/749E, 671N/934R, and 749E/784T, wherein the positions are

numbered with reference to SEQ ID NO: 8. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from P4H, F27P/L28S/A489R, F27P/A418E/A478T, F27R, L28S, L28S/L29T, L28S/L29T/Q113S/S135Q/M138A, L28S/L29T/Q113S/S135Q/A418E, L28S/L29T/S135Q, L28S/L29T/A418E, L29T/Q113S/P126Q/S135Q/H193Q, L29T/Q113S/S135Q, L29T/Q113S/S135Q/R455V, L29T/Q113S/M138A, L29T/T148G, L29T/A478T, K106P, K106P/M138A/L218S/M431V/S671N/Q749E, K106P/L218S/A281V, K106P/L218S/R455V, K106P/L218S/R455V/M507L/Q749E, K106P/A489R/S671N, K106P/V638I, K106P/S671N/L934R, Q113S, Q113S/S135Q/A418E, Q113S/A418E/R455V/A478T/K581T, Q113S/A418E/A478T/A489R/K581T, S135P/C944Y, S135Q, S135Q/T148G/S150G/A418E, S135Q/A478T/A489R/K581T, S135Q/A489R, M138A/L218S/S668D/S671N, M138A/L218S/Q749E/L934R, M138A/S671N/Q749E/L934R, L157M, L218S, L218S/A281V, L218S/A281V/M431V, L218S/A281V/S671N, L218S/M431V, L218S/M431V/A489R/M507L/Q749E/L934R, L218S/R455V, L218S/M507L/Q749E, L218S/M507L/L934R, L218S/V638I/S671N, L218S/Q749E, A281V/M431V/A489R/S668D, Q345K/L934R, A418E, A418E/A489R, M431V/S668D/S671N, A489R/V638I/L934R, A489R/S671N/L934R, A489R/Q749E, A489R/L934R, M507L/S668D, M507L/S671N/L934R, S671N/Q749E, S671N/L934R, and Q749E/A784, wherein the positions are numbered with reference to SEQ ID NO: 8. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from 27, 27/418/478, 28, 28/29, 28/29/113/135/138, 28/29/113/135/418, 28/29/135, 28/29/418, 29/113/126/135/193, 29/113/135, 29/113/135/455, 29/113/138, 29/148, 29/478, 106, 106/138/218/431/671/749, 106/489/671, 106/638, 113/135/418, 113/418/455/478/581, 113/418/478/489/581, 135, 135/148/150/418, 135/478/489/581, 135/944, 138/218/668/671, 157, 218/638/671, 418, 418/489, 431/668/671, 507/668, and 671/749, wherein the positions are numbered with reference to SEQ ID NO: 8. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from 27P/418E/478T, 27R, 28S, 28S/29T, 28S/29T/113S/135Q/138A, 28S/29T/113S/135Q/418E, 28S/29T/135Q, 28S/29T/418E, 29T/113S/126Q/135Q/193Q, 29T/113S/135Q, 29T/113S/135Q/455V, 29T/113S/138A, 29T/148G, 29T/478T, 106P, 106P/138A/218S/431V/671N/749E, 106P/489R/671N, 106P/638I, 113S/135Q/418E, 113S/418E/455V/478T/581T, 113S/418E/478T/489R/581T, 135P/944Y, 135Q, 135Q/148G/150G/418E, 135Q/478T/489R/581T, 138A/218S/668D/671N, 157M, 218S/638I/671N, 418E, 418E/489R, 431V/668D/671N, 507L/668D, and 671N/749E, wherein the positions are numbered with reference to SEQ ID NO: 8. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from F27P/A418E/A478T, F27R, L28S, L28S/L29T, L28S/L29T/Q113S/S135Q/M138A, L28S/L29T/Q113S/S135Q/A418E, L28S/L29T/S135Q, L28S/L29T/A418E,

L29T/Q113S/P126Q/S135Q/H193Q, L29T/Q113S/S135Q, L29T/Q113S/S135Q/R455V, L29T/Q113S/M138A, L29T/T148G, L29T/A478T, K106P, K106P/M138A/L218S/M431V/S671N/Q749E, K106P/A489R/S671N, K106P/V638I, Q113S/S135Q/A418E, Q113S/A418E/R455V/A478T/K581T, Q113S/A418E/A478T/A489R/K581T, S135P/C944Y, S135Q, S135Q/T148G/S150G/A418E, S135Q/A478T/A489R/K581T, M138A/L218S/S668D/S671N, L157M, L218S/V638I/S671N, A418E, A418E/A489R, M431V/S668D/S671N, M507L/S668D, and S671N/Q749E, wherein the positions are numbered with reference to SEQ ID NO: 8.

[0011] The present invention also provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 12. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 4, 27, 27/28/489, 27/418/478, 28, 28/29, 28/29/113/135/138, 28/29/113/135/418, 28/29/135, 28/29/418, 29/113/126/135/193, 29/113/135, 29/113/135/455, 29/113/138, 29/148, 29/478, 106, 106/138/218/431/671/749, 106/218/281, 106/218/455, 106/218/455/507/749, 106/489/671, 106/638, 106/671/934, 113, 113/135/418, 113/418/455/478/581, 113/418/478/489/581, 135, 135/148/150/418, 135/478/489/581, 135/489, 135/944, 138/218/668/671, 138/218/749/934, 138/671/749/934, 157, 218, 218/281/431, 218/281/671, 218/431, 218/455, 218/507/749, 218/638/671, 218/749, 281/431/489/668, 345/934, 418, 418/489, 431/668/671, 489/638/934, 489/671/934, 489/749, 489/934, 507/668, 507/671/934, 671/749, 671/934, and 749/784, wherein the positions are numbered with reference to SEQ ID NO: 12. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 4H, 27P/28S/489R, 27P/418E/478T, 27R, 28S, 28S/29T, 28S/29T/113S/135Q/138A, 28S/29T/113S/135Q/418E, 28S/29T/135Q, 28S/29T/418E, 29T/113S/126Q/135Q/193Q, 29T/113S/135Q, 29T/113S/135Q/455V, 29T/113S/138A, 29T/148G, 29T/478T, 106P, 106P/138A/218S/431V/671N/749E, 106P/218S/281V, 106P/218S/455V, 106P/218S/455V/507L/749E, 106P/489R/671N, 106P/638I, 106P/671N/934R, 113S, 113S/135Q/418E, 113S/418E/455V/478T/581T, 113S/418E/478T/489R/581T, 135P/944Y, 135Q, 135Q/148G/150G/418E, 135Q/478T/489R/581T, 135Q/489R, 138A/218S/668D/671N, 138A/218S/749E/934R, 138A/671N/749E/934R, 157M, 218S, 218S/281V/431V, 218S/281V/671N, 218S/431V, 218S/455V, 218S/507L/749E, 218S/638I/671N, 218S/749E, 281V/431V/489R/668D, 345K/934R, 418E, 418E/489R, 431V/668D/671N, 489R/638I/934R, 489R/671N/934R, 489R/749E, 489R/934R, 507L/668D, 507L/671N/934R, 671N/749E, 671N/934R, and 749E/784T, wherein the positions are numbered with reference to SEQ ID NO: 12. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from P4H, F27P/L28S/A489R, F27P/A418E/A478T, F27R, L28S, L28S/L29T,

L28S/L29T/Q113S/S135Q/M138A, L28S/L29T/Q113S/S135Q/A418E, L28S/L29T/S135Q, L28S/L29T/A418E, L29T/Q113S/P126Q/S135Q/H193Q, L29T/Q113S/S135Q, L29T/Q113S/S135Q/R455V, L29T/Q113S/M138A, L29T/T148G, L29T/A478T, K106P, K106P/M138A/L218S/M431V/S671N/Q749E, K106P/L218S/A281V, K106P/L218S/R455V, K106P/L218S/R455V/M507L/Q749E, K106P/A489R/S671N, K106P/V638I, K106P/S671N/L934R, Q113S, Q113S/S135Q/A418E, Q113S/A418E/R455V/A478T/K581T, Q113S/A418E/A478T/A489R/K581T, S135P/C944Y, S135Q, S135Q/T148G/S150G/A418E, S135Q/A478T/A489R/K581T, S135Q/A489R, M138A/L218S/S668D/S671N, M138A/L218S/Q749E/L934R, M138A/S671N/Q749E/L934R, L157M, L218S, L218S/A281V/M431V, L218S/A281V/S671N, L218S/M431V, L218S/R455V, L218S/M507L/Q749E, L218S/V638I/S671N, L218S/Q749E, A281V/M431V/A489R/S668D, Q345K/L934R, A418E, A418E/A489R, M431V/S668D/S671N, A489R/V638I/L934R, A489R/S671N/L934R, A489R/Q749E, A489R/L934R, M507L/S668D, M507L/S671N/L934R, S671N/Q749E, S671N/L934R, and Q749E/A784T, wherein the positions are numbered with reference to SEQ ID NO: 12.

[0012] The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 14. The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 16. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 22, 24, 27, 27/165, 30, 33, 34, 37/62, 37/62/79/196/696/862, 37/62/523, 37/62/523/793, 37/64/66/79/154/523/681/793/862, 37/79/154/793, 37/196, 37/528/696/793, 37/528/790, 37/528/790/793/862, 37/790/793, 39, 39/58/489/725/830/842/930/944, 39/70/109/830/842, 39/70/489/612, 39/70/725, 39/267, 39/267/489/522/612/830/842, 39/267/489/830/944, 39/489/500/612, 39/500/612, 40, 44/157, 47, 49, 50, 55, 60/500/612, 62/79/154/862, 62/79/196/681/862, 62/79/523/528/790, 62/79/790/793, 62/79/862, 62/92, 62/92/790/793, 62/106/523/528/696/793/862, 62/154/696/793/862, 62/793/862, 68, 70, 70/267/725/944, 70/267/930/944, 70/489/930, 70/725/830/860/930/944, 77, 79/154/681, 79/154/793/862, 79/862, 89, 97, 106/154, 107, 109, 109/522/612/725, 109/522/830/944, 109/612, 118, 149, 157, 158, 178, 179, 196/528/681/790/793, 207, 208, 217, 267/489/500/725/830/930, 267/522/725, 352, 385, 424, 448, 463, 489/830/944, 500, 500/612/830/860, 500/860/930, 500/930/944, 522/725, 523, 523/790/793, 528/681, 528/793, 528/862, 672, 673, 725, 734, 740, 753, 774, 778, 793, 830, 844, 862, 875, 880, 892, 902, 922, 925, 930, 932, 934, 938, and 944, wherein the

positions are numbered with reference to SEQ ID NO: 14 and/or 16. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from 22R, 24E, 24R, 24W, 27A, 27G, 27G/165I, 27K, 27R, 27S, 27V, 27W, 30D, 30L, 33G, 33P, 34D, 34M, 34T, 37F/62E, 37F/62E/79S/196T/696S/862Q, 37F/62E/523N, 37F/62E/523N/793K, 37F/64Q/66G/79S/154R/523N/681Q/793K/862Q, 37F/79S/154R/793K, 37F/196T, 37F/528S/696S/793K, 37F/528S/790V, 37F/528S/790V/793K/862Q, 37F/790V/793K, 39D, 39H, 39Q, 39Q/58L/489D/725E/830K/842S/930P/944S, 39Q/70A/109P/830K/842S, 39Q/70A/489D/612D, 39Q/70A/725E, 39Q/267K, 39Q/267K/489D/522V/612D/830K/842S, 39Q/267K/489D/830K/944S, 39Q/489D/500A/612D, 39Q/500A/612D, 40W, 44I/157V, 47G, 47R, 49A, 49G, 50G, 50L, 50V, 55C, 55L, 60V/500A/612D, 62E/79S/154R/862Q, 62E/79S/196T/681Q/862Q, 62E/79S/523N/528S/790V, 62E/79S/790V/793K, 62E/79S/862Q, 62E/92R, 62E/92R/790V/793K, 62E/106R/523N/528S/696S/793K/862Q, 62E/154R/696S/793K/862Q, 62E/793K/862Q, 68N, 68S, 68W, 70A/267K/725E/944S, 70A/267K/930P/944S, 70A/489D/930P, 70A/725E/830K/860F/930P/944S, 70Q, 77W, 79S/154R/681Q, 79S/154R/793K/862Q, 79S/862Q, 89R, 97D, 97G, 106R/154R, 107G, 109D, 109P/522V/612D/725E, 109P/522V/830K/944S, 109P/612D, 118F, 149R, 157Q, 158E, 158F, 178G, 178V, 179L, 196T/528S/681Q/790V/793K, 207R, 207Y, 208G, 208I, 217A, 217D, 267K/489D/500A/725E/830K/930P, 267K/522V/725E, 352K, 352V, 385G, 424K, 448L, 463A, 489D/830K/944S, 500A, 500A/612D/830K/860F, 500A/860F/930P, 500A/930P/944S, 522V/725E, 523N, 523N/790V/793K, 528S/681Q, 528S/793K, 528S/862Q, 672E, 672K, 673N, 673R, 725F, 725V, 734K, 740G, 740Q, 753S, 774G, 774S, 778Q, 793K, 830V, 844R, 862Q, 875D, 880R, 892L, 902L, 922E, 925A, 925W, 930P, 932A, 934F, 938A, 938P, 944R, and 944S, wherein the positions are numbered with reference to SEQ ID NO: 14 and/or 16. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from I22R, L24E, L24R, L24W, F27A, F27G, F27G/M165I, F27K, F27R, F27S, F27V, F27W, V30D, V30L, E33G, E33P, L34D, L34M, L34T, S37F/A62E, S37F/A62E/N79S/A196T/A696S/R862Q, S37F/A62E/D523N, S37F/A62E/D523N/E793K, S37F/P64Q/R66G/N79S/K154R/D523N/E681Q/E793K/R862Q, S37F/N79S/K154R/E793K, S37F/A196T, S37F/N528S/A696S/E793K, S37F/N528S/I790V, S37F/N528S/I790V/E793K/R862Q, S37F/I790V/E793K, P39D, P39H, P39Q, P39Q/R58L/A489D/K725E/Q830K/G842S/C930P/C944S, P39Q/V70A/L109P/Q830K/G842S, P39Q/V70A/A489D/S612D, P39Q/V70A/K725E, P39Q/R267K, P39Q/R267K/A489D/E522V/S612D/Q830K/G842S, P39Q/R267K/A489D/Q830K/C944S, P39Q/A489D/D500A/S612D, P39Q/D500A/S612D, V40W, T44I/L157V, A47G, A47R, Q49A, Q49G, Q50G, Q50L, Q50V, P55C, P55L, A60V/D500A/S612D, A62E/N79S/K154R/R862Q, A62E/N79S/A196T/E681Q/R862Q, A62E/N79S/D523N/N528S/I790V, A62E/N79S/I790V/E793K, A62E/N79S/R862Q, A62E/Q92R, A62E/Q92R/I790V/E793K, A62E/K106R/D523N/N528S/A696S/E793K/R862Q, A62E/K154R/A696S/E793K/R862Q,

A62E/E793K/R862Q, R68N, R68S, R68W, V70A/R267K/K725E/C944S, V70A/R267K/C930P/C944S, V70A/A489D/C930P, V70A/K725E/Q830K/L860F/C930P/C944S, V70Q, P77W, N79S/K154R/E681Q, N79S/K154R/E793K/R862Q, N79S/R862Q, A89R, A97D, A97G, K106R/K154R, Q107G, L109D, L109P/E522V/S612D/K725E, L109P/E522V/Q830K/C944S, L109P/S612D, W118F, P149R, L157Q, T158E, T158F, P178G, P178V, A179L, A196T/N528S/E681Q/I790V/E793K, E207R, E207Y, E208G, E208I, Q217A, Q217D, R267K/A489D/D500A/K725E/Q830K/C930P, R267K/E522V/K725E, Y352K, Y352V, R385G, H424K, R448L, E463A, A489D/Q830K/C944S, D500A, D500A/S612D/Q830K/L860F, D500A/L860F/C930P, D500A/C930P/C944S, E522V/K725E, D523N, D523N/I790V/E793K, N528S/E681Q, N528S/E793K, N528S/R862Q, L672E, L672K, P673N, P673R, K725F, K725V, H734K, E740G, E740Q, A753S, A774G, A774S, L778Q, E793K, Q830V, E844R, R862Q, N875D, E880R, Q892L, A902L, P922E, K925A, K925W, C930P, S932A, L934F, Q938A, Q938P, C944R, and C944S, wherein the positions are numbered with reference to SEQ ID NO: 14 and/or 16.

[0013] The present invention also provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 18, wherein the positions are numbered with reference to SEQ ID NO: 18. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 22, 24, 27, 27/165, 30, 33, 34, 37/62, 37/62/79/196/696/862, 37/62/523, 37/196, 37/528/790, 39, 39/70/109/830/842, 39/70/725, 39/267, 39/267/489/522/612/830/842, 39/267/489/830/944, 40, 70/267/725/944, 70/267/930/944, 70/489/930, 107, 109, 109/522/830/944, 217, 267/489/500/725/830/930, 267/522/725, 352, 385, 500/930/944, 673, 734, 774, 778, 875, 930, 932, and 934, wherein the positions are numbered with reference to SEQ ID NO: 18. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 22R, 24E, 24R, 24W, 27G, 27G/165I, 27K, 27R, 27W, 30D, 33G, 34D, 34M, 34T, 37F/62E, 37F/62E/79S/196T/696S/862Q, 37F/62E/523N, 37F/196T, 37F/528S/790V, 39D, 39Q, 39Q/70A/109P/830K/842S, 39Q/70A/725E, 39Q/267K, 39Q/267K/489D/522V/612D/830K/842S, 39Q/267K/489D/830K/944S, 40W, 70A/267K/725E/944S, 70A/267K/930P/944S, 70A/489D/930P, 107G, 109D, 109P/522V/830K/944S, 217D, 267K/489D/500A/725E/830K/930P, 267K/522V/725E, 352K, 352V, 385G, 500A/930P/944S, 673N, 734K, 774G, 778Q, 875D, 930P, 932A, and 934F, wherein the positions are numbered with reference to SEQ ID NO: 18. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from I22R, L24E, L24R, L24W, F27G, F27G/M165I, F27K, F27R, F27W, V30D, E33G, L34D, L34M, L34T, S37F/A62E, S37F/A62E/N79S/A196T/A696S/R862Q, S37F/A62E/D523N, S37F/A196T, S37F/N528S/I790V, P39D, P39Q, P39Q/V70A/L109P/Q830K/G842S, P39Q/V70A/K725E, P39Q/R267K, P39Q/R267K/A489D/E522V/S612D/Q830K/G842S,

P39Q/R267K/A489D/Q830K/C944S, V40W, V70A/R267K/K725E/C944S, V70A/R267K/C930P/C944S, V70A/A489D/C930P, Q107G, L109D, L109P/E522V/Q830K/C944S, Q217D, R267K/A489D/D500A/K725E/Q830K/C930P, R267K/E522V/K725E, Y352K, Y352V, R385G, D500A/C930P/C944S, P673N, H734K, A774G, L778Q, N875D, C930P, S932A, and L934F, wherein the positions are numbered with reference to SEQ ID NO: 18.

[0014] The present invention also provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 20, wherein the positions are numbered with reference to SEQ ID NO: 20. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 22/24, 22/24/27/50/107/109/489/612/725, 22/24/27/489, 22/24/27/489/612/774, 22/24/27/612/944, 22/24/50/109/267/489/774/944, 22/24/50/267/612/922/944, 22/24/107/267/489/922, 22/24/489, 22/24/612/725/944, 22/50/107/267/489/612/944, 22/50/109/267/489, 22/267/489/612, 24, 24/27/50/107/267/774/944, 24/27/89/500/842, 24/27/107/267/612/944, 24/27/267/944, 24/27/500/842, 24/27/500/842/932, 24/27/944, 24/39/49/89/97/842/932, 24/39/68/89/107/500/842, 24/39/89/97/842/932, 24/39/842/932, 24/50/489/944, 24/50/612, 24/70/107/109/489/612/725, 24/70/267/774, 24/89/500, 24/107/109/267/489/612/725/774, 24/109/612, 24/109/944, 24/267/725/944, 24/489/944, 24/725, 24/842/932, 24/944, 27/39/49/97/500/842, 27/49/68/500/842, 34/39/500/932, 39/89/97/500, 42, 48, 50/109/489/612, 50/489/774, 50/612/944, 57, 62, 68, 68/89/97/932, 71, 88, 89/97/107, 89/97/500, 89/842, 107/109, 107/500/842, 108, 109/612/774/944, 112, 123, 124, 148, 188, 193, 197, 204, 253, 264, 305, 312, 333, 381, 402, 402/781, 489, 489/944, 500/842, 500/932, 523, 527, 612, 612/725/944, 612/922, 614, 727, 742, 748, 820, 823, 832, 842/932, 858, 862, 911, 913, 914, 916, 923, 937, and 940, wherein the positions are numbered with reference to SEQ ID NO: 20. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 22R/24W, 22R/24W/27A/50V/107G/109D/489A/612S/725E, 22R/24W/27A/489A, 22R/24W/27A/489A/612S/774S, 22R/24W/27A/612S/944R, 22R/24W/50V/109D/267R/489A/774S/944R, 22R/24W/50V/267R/612S/922E/944R, 22R/24W/107G/267R/489A/922E, 22R/24W/489A, 22R/24W/612S/725E/944R, 22R/50V/107G/267R/489A/612S/944S, 22R/50V/109D/267R/489A, 22R/267R/489A/612S, 24R, 24R/27G/89R/500A/842G, 24R/27G/500A/842G, 24R/27G/500A/842G/932A, 24R/39D/68S/89R/107G/500A/842G, 24R/39H/49G/89R/97G/842G/932A, 24R/39H/89R/97D/842G/932A, 24R/39H/842G/932A, 24R/89R/500A, 24R/842G/932A, 24W, 24W/27A/50V/107G/267R/774S/944S, 24W/27A/107G/267R/612S/944S, 24W/27A/267R/944R, 24W/27A/944R, 24W/50V/489A/944S, 24W/50V/612S, 24W/70A/107G/109D/489A/612S/725E, 24W/70A/267R/774S, 24W/107G/109D/267R/489A/612S/725E/774S, 24W/109D/612S,

24W/109D/944S, 24W/267R/725E/944S, 24W/489A/944R, 24W/725E, 24W/944S, 27G/39H/49G/97G/500A/842G, 27G/49G/68S/500A/842G, 34T/39D/500A/932A, 39D/89R/97G/500A, 42G, 48Q, 48V, 48W, 50V/109D/489A/612S, 50V/489A/774S, 50V/612S/944S, 57F, 57L, 57M, 62F, 62L, 62W, 68N/89R/97G/932A, 68S, 71G, 71L, 71V, 71W, 71Y, 88L, 88R, 89R/97G/107G, 89R/97G/500A, 89R/842G, 107G/109D, 107G/500A/842G, 108R, 109D/612S/774S/944S, 112H, 123L, 123V, 124G, 124M, 124V, 148K, 148R, 188R, 188W, 193E, 193P, 197G, 204A, 253M, 264M, 305F, 312A, 333L, 381R, 381V, 381W, 402N, 402V/781Q, 489A, 489A/944R, 500A/842G, 500A/932A, 523E, 527R, 527V, 612S, 612S/725E/944S, 612S/922E, 614Q, 614R, 614W, 727W, 742V, 748V, 820A, 820V, 823F, 823V, 832A, 832R, 842G/932A, 858C, 858W, 862I, 862M, 862Q, 862Y, 911G, 911R, 913G, 913R, 913W, 914G, 914I, 914K, 914Q, 914R, 914S, 914T, 916G, 916H, 916R, 923L, 923V, 923W, 937K, and 940Q, wherein the positions are numbered with reference to SEQ ID NO: 20. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from I22R/L24W, I22R/L24W/F27A/Q50V/Q107G/L109D/D489A/D612S/K725E, I22R/L24W/F27A/D489A, I22R/L24W/F27A/D489A/D612S/A774S, I22R/L24W/F27A/D612S/C944R, I22R/L24W/Q50V/L109D/K267R/D489A/A774S/C944R, I22R/L24W/Q50V/K267R/D612S/P922E/C944R, I22R/L24W/Q107G/K267R/D489A/P922E, I22R/L24W/D489A, I22R/L24W/D612S/K725E/C944R, I22R/Q50V/Q107G/K267R/D489A/D612S/C944S, I22R/Q50V/L109D/K267R/D489A, I22R/K267R/D489A/D612S, L24R, L24R/F27G/A89R/D500A/S842G, L24R/F27G/D500A/S842G, L24R/F27G/D500A/S842G/S932A, L24R/Q39D/R68S/A89R/Q107G/D500A/S842G, L24R/Q39H/Q49G/A89R/A97G/S842G/S932A, L24R/Q39H/A89R/A97D/S842G/S932A, L24R/Q39H/S842G/S932A, L24R/A89R/D500A, L24R/S842G/S932A, L24W, L24W/F27A/Q50V/Q107G/K267R/A774S/C944S, L24W/F27A/Q107G/K267R/D612S/C944S, L24W/F27A/K267R/C944R, L24W/F27A/C944R, L24W/Q50V/D489A/C944S, L24W/Q50V/D612S, L24W/V70A/Q107G/L109D/D489A/D612S/K725E, L24W/V70A/K267R/A774S, L24W/Q107G/L109D/K267R/D489A/D612S/K725E/A774S, L24W/L109D/D612S, L24W/L109D/C944S, L24W/K267R/K725E/C944S, L24W/D489A/C944R, L24W/K725E, L24W/C944S, F27G/Q39H/Q49G/A97G/D500A/S842G, F27G/Q49G/R68S/D500A/S842G, L34T/Q39D/D500A/S932A, Q39D/A89R/A97G/D500A, E42G, H48Q, H48V, H48W, Q50V/L109D/D489A/D612S, Q50V/D489A/A774S, Q50V/D612S/C944S, P57F, P57L, P57M, A62F, A62L, A62W, R68N/A89R/A97G/S932A, R68S, P71G, P71L, P71V, P71W, P71Y, K88L, K88R, A89R/A97G/Q107G, A89R/A97G/D500A, A89R/S842G, Q107G/L109D, Q107G/D500A/S842G, G108R, L109D/D612S/A774S/C944S, A112H, P123L, P123V, S124G, S124M, S124V, T148K, T148R, E188R, E188W, H193E, H193P, P197G, E204A, A253M, S264M, L305F, V312A, I333L, E381R, E381V, E381W, S402N, S402V/P781Q, D489A, D489A/C944R, D500A/S842G, D500A/S932A, D523E, N527R, N527V, D612S,

D612S/K725E/C944S, D612S/P922E, E614Q, E614R, E614W, S727W, L742V, L748V, G820A, G820V, L823F, L823V, P832A, P832R, S842G/S932A, E858C, E858W, R862I, R862M, R862Q, R862Y, N911G, N911R, V913G, V913R, V913W, P914G, P914I, P914K, P914Q, P914R, P914S, P914T, S916G, S916H, S916R, D923L, D923V, D923W, E937K, and L940Q, wherein the positions are numbered with reference to SEQ ID NO: 20. In some additional embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 22/24, 22/24/27/50/107/109/489/612/725, 22/24/27/489, 22/24/27/489/612/774, 22/24/27/612/944, 22/24/50/109/267/489/774/944, 22/24/50/267/612/922/944, 22/24/107/267/489/922, 22/24/489, 22/24/612/725/944, 22/50/107/267/489/612/944, 22/50/109/267/489, 22/267/489/612, 24, 24/27/50/107/267/774/944, 24/27/89/500/842, 24/27/107/267/612/944, 24/27/267/944, 24/27/500/842, 24/27/500/842/932, 24/27/944, 24/39/49/89/97/842/932, 24/39/68/89/107/500/842, 24/39/89/97/842/932, 24/39/842/932, 24/50/489/944, 24/50/612, 24/70/107/109/489/612/725, 24/70/267/774, 24/89/500, 24/107/109/267/489/612/725/774, 24/109/612, 24/109/944, 24/267/725/944, 24/489/944, 24/725, 24/842/932, 24/944, 27/39/49/97/500/842, 27/49/68/500/842, 34/39/500/932, 50/109/489/612, 50/612/944, 68/89/97/932, 89/97/107, 89/842, 106, 107/109, 107/500/842, 108, 109/612/774/944, 112, 148, 148/772, 188/377, 238, 240, 240/374, 243, 244, 246, 248, 249/777, 252, 253, 259, 260, 261, 262, 264, 279, 305, 309, 312, 319, 320, 329, 333, 387, 402, 421, 432, 500/842, 500/932, 556, 612, 612/725/944, 612/922, 727, 736, 737, 741, 742, 748, 815, 816, 818, 823, 832, 842/932, 911, 913, 914, 916, 923, 937, and 940, wherein the positions are numbered with reference to SEQ ID NO: 20. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 22R/24W, 22R/24W/27A/50V/107G/109D/489A/612S/725E, 22R/24W/27A/489A, 22R/24W/27A/489A/612S/774S, 22R/24W/27A/612S/944R, 22R/24W/50V/109D/267R/489A/774S/944R, 22R/24W/50V/267R/612S/922E/944R, 22R/24W/107G/267R/489A/922E, 22R/24W/489A, 22R/24W/612S/725E/944R, 22R/50V/107G/267R/489A/612S/944S, 22R/50V/109D/267R/489A, 22R/267R/489A/612S, 24R, 24R/27G/89R/500A/842G, 24R/27G/500A/842G, 24R/27G/500A/842G/932A, 24R/39D/68S/89R/107G/500A/842G, 24R/39H/49G/89R/97G/842G/932A, 24R/39H/89R/97D/842G/932A, 24R/39H/842G/932A, 24R/89R/500A, 24R/842G/932A, 24W, 24W/27A/50V/107G/267R/774S/944S, 24W/27A/107G/267R/612S/944S, 24W/27A/267R/944R, 24W/27A/944R, 24W/50V/489A/944S, 24W/50V/612S, 24W/70A/107G/109D/489A/612S/725E, 24W/70A/267R/774S, 24W/107G/109D/267R/489A/612S/725E/774S, 24W/109D/612S, 24W/109D/944S, 24W/267R/725E/944S, 24W/489A/944R, 24W/725E, 24W/944S, 27G/39H/49G/97G/500A/842G, 27G/49G/68S/500A/842G, 34T/39D/500A/932A, 50V/109D/489A/612S, 50V/612S/944S, 68N/89R/97G/932A, 89R/97G/107G, 89R/842G, 106A, 106G, 106N, 106T, 107G/109D, 107G/500A/842G, 108H, 108N, 108R, 108S, 108V, 109D/612S/774S/944S, 112H, 112P, 148E, 148G, 148H, 148K, 148R/772I, 188Q/377Q, 238Q, 240I,

240W/374T, 240Y, 243E, 243G, 243R, 243V, 244I, 244V, 246A, 246G, 248A, 248R, 248V, 249V/777N, 252V, 253G, 253P, 259G, 259N, 259S, 260W, 261E, 262P, 264C, 279E, 305F, 305G, 305R, 305V, 305Y, 309C, 309G, 312A, 319F, 320M, 329F, 333L, 333V, 387L, 402G, 402N, 421P, 432C, 500A/842G, 500A/932A, 556H, 556R, 556S, 556Y, 612S, 612S/725E/944S, 612S/922E, 727G, 727Q, 727T, 727W, 736M, 736V, 736W, 737M, 741C, 741D, 741E, 741G, 741T, 742V, 748I, 748T, 748V, 815A, 815M, 816V, 818T, 818V, 823A, 823F, 823G, 823R, 832E, 832G, 842G/932A, 911G, 913A, 913E, 913G, 913H, 913L, 913Q, 913R, 913W, 914E, 914G, 914H, 914K, 914Q, 914R, 914S, 914T, 916A, 916G, 916H, 916I, 916R, 916V, 923W, 937Q, 940G, 940Q, 940T, and 940W, wherein the positions are numbered with reference to SEQ ID NO: 20. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from I22R/L24W, I22R/L24W/F27A/Q50V/Q107G/L109D/D489A/D612S/K725E, I22R/L24W/F27A/D489A, I22R/L24W/F27A/D489A/D612S/A774S, I22R/L24W/F27A/D612S/C944R, I22R/L24W/Q50V/L109D/K267R/D489A/A774S/C944R, I22R/L24W/Q50V/K267R/D612S/P922E/C944R, I22R/L24W/Q107G/K267R/D489A/P922E, I22R/L24W/D489A, I22R/L24W/D612S/K725E/C944R, I22R/Q50V/Q107G/K267R/D489A/D612S/C944S, I22R/Q50V/L109D/K267R/D489A, I22R/K267R/D489A/D612S, L24R, L24R/F27G/A89R/D500A/S842G, L24R/F27G/D500A/S842G, L24R/F27G/D500A/S842G/S932A, L24R/Q39D/R68S/A89R/Q107G/D500A/S842G, L24R/Q39H/Q49G/A89R/A97G/S842G/S932A, L24R/Q39H/A89R/A97D/S842G/S932A, L24R/Q39H/S842G/S932A, L24R/A89R/D500A, L24R/S842G/S932A, L24W, L24W/F27A/Q50V/Q107G/K267R/A774S/C944S, L24W/F27A/Q107G/K267R/D612S/C944S, L24W/F27A/K267R/C944R, L24W/F27A/C944R, L24W/Q50V/D489A/C944S, L24W/Q50V/D612S, L24W/V70A/Q107G/L109D/D489A/D612S/K725E, L24W/V70A/K267R/A774S, L24W/Q107G/L109D/K267R/D489A/D612S/K725E/A774S, L24W/L109D/D612S, L24W/L109D/C944S, L24W/K267R/K725E/C944S, L24W/D489A/C944R, L24W/K725E, L24W/C944S, F27G/Q39H/Q49G/A97G/D500A/S842G, F27G/Q49G/R68S/D500A/S842G, L34T/Q39D/D500A/S932A, Q50V/L109D/D489A/D612S, Q50V/D612S/C944S, R68N/A89R/A97G/S932A, A89R/A97G/Q107G, A89R/S842G, K106A, K106G, K106N, K106T, Q107G/L109D, Q107G/D500A/S842G, G108H, G108N, G108R, G108S, G108V, L109D/D612S/A774S/C944S, A112H, A112P, T148E, T148G, T148H, T148K, T148R/V772I, E188Q/R377Q, L238Q, L240I, L240W/A374T, L240Y, S243E, S243G, S243R, S243V, L244I, L244V, S246A, S246G, Y248A, Y248R, Y248V, I249V/S777N, L252V, A253G, A253P, L259G, L259N, L259S, M260W, L261E, S262P, S264C, P279E, L305F, L305G, L305R, L305V, L305Y, A309C, A309G, V312A, A319F, L320M, L329F, I333L, I333V, H387L, S402G, S402N, Q421P, M432C, D500A/S842G, D500A/S932A, F556H, F556R, F556S, F556Y, D612S, D612S/K725E/C944S, D612S/P922E, S727G, S727Q, S727T, S727W, L736M, L736V, L736W, L737M, A741C, A741D, A741E, A741G, A741T, L742V, L748I, L748T, L748V, I815A, I815M,

I816V, L818T, L818V, L823A, L823F, L823G, L823R, P832E, P832G, S842G/S932A, N911G, V913A, V913E, V913G, V913H, V913L, V913Q, V913R, V913W, P914E, P914G, P914H, P914K, P914Q, P914R, P914S, P914T, S916A, S916G, S916H, S916I, S916R, S916V, D923W, E937Q, L940G, L940Q, L940T, and L940W, wherein the positions are numbered with reference to SEQ ID NO: 20.

[0015] The present invention further provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 946, wherein the positions are numbered with reference to SEQ ID NO: 946. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 19/124/149/381/727, 24/39/489/862, 24/57/62/89/489/823/862, 24/57/823/862, 24/62/89/188/823/842/862, 24/89/489, 24/89/489/727/862, 24/489/500/842, 39/57/62/188/500/842, 39/57/500/862, 57, 57/62/120/527/913/916, 57/62/305/437/500/614/727/916, 57/62/305/437/500/727/913/916, 57/62/305/437/614/683/913/916/932, 57/62/305/489/907/913/916, 57/62/305/489/913/916, 57/62/305/500/913/916, 57/62/305/913, 57/62/305/916, 57/62/437/500/761/914/916, 57/62/437/527/727, 57/62/437/913/916, 57/62/913/916/932, 57/62/916, 57/188/489/823/862, 57/305, 57/305/437/916, 57/437/500/527/727/916, 57/437/500/614/727/914, 57/437/913/914, 57/489/527/914/916, 57/614/916/932, 62/89, 62/89/124/148/381/858, 62/89/124/381/858, 62/89/148/381/614/858, 62/89/148/923, 62/89/149/381, 62/89/149/381/832, 62/89/188/489/500/727/823, 62/89/381, 62/89/381/858, 62/89/381/923, 62/89/858, 62/96/614, 62/124/148/149/381/614, 62/124/149/381/832/858/937, 62/124/188/823/842/862, 62/124/381/832, 62/148/149/381/858/937, 62/148/381/614/937, 62/148/381/727, 62/148/381/858, 62/149/381/614/937, 62/149/381/858/937, 62/149/727, 62/305/437/500/727/913, 62/305/727, 62/381, 62/437/489/527/727/913/932, 62/437/489/614/727/913, 62/437/527/727, 62/437/527/916/932, 62/437/913/916, 62/489/500/932, 62/489/527/916/932, 62/489/614/916, 62/500, 62/527, 62/527/727/916, 62/614, 62/727, 62/916, 89/148/149, 89/148/149/381, 89/381, 124/148/381/727/858/937, 124/381/614, 124/500/842/862, 124/832/937, 148/832/858/937, 381, 381/614/832, 381/858/937, 437, 437/489/914/916, 437/727/914, 437/914/916, 489/500, 489/614/916, 500/727/913/916, 500/914/916, and 923, wherein the positions are numbered with reference to SEQ ID NO: 946. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 19T/124V/149R/381V/727W, 24R/39H/489A/862Q, 24R/57L/62W/89R/489A/823F/862Q, 24R/57L/823F/862Q, 24R/62W/89R/188W/823V/842G/862Q, 24R/89R/489A, 24R/89R/489A/727W/862Q, 24R/489A/500A/842G, 39H/57L/62W/188W/500A/842G, 39H/57L/500A/862Q, 57F/62L/305F/437G/500A/614Q/727W/916R, 57F/62L/305F/437G/614Q/683S/913R/916R/932A,

57F/62L/305F/500A/913R/916G, 57F/62L/305F/913R, 57F/62L/437G/500A/761F/914K/916R,
 57F/62L/437G/527R/727W, 57F/62L/913R/916R/932A, 57F/62L/916G,
 57F/62W/120I/527R/913R/916R, 57F/62W/305F/437G/500A/727W/913R/916R,
 57F/62W/305F/489A/907K/913R/916G, 57F/62W/305F/489A/913R/916G, 57F/62W/305F/916R,
 57F/62W/437G/913R/916G, 57F/305F, 57F/305F/437G/916G, 57F/437G/500A/527R/727W/916R,
 57F/437G/500A/614Q/727W/914R, 57F/437G/913R/914R, 57F/489A/527R/914R/916G,
 57F/614Q/916G/932A, 57L, 57L/188W/489A/823F/862Q, 62F/89R,
 62F/89R/124V/148R/381W/858W, 62F/89R/148R/381V/614R/858W, 62F/89R/148R/923W,
 62F/89R/149R/381W/832R, 62F/89R/381V/923W, 62F/89R/858C, 62F/96K/614R,
 62F/124V/381W/832R, 62F/149R/381V/858W/937K, 62F/149R/727W, 62F/381V, 62F/614R,
 62L/305F/437G/500A/727W/913R, 62L/305F/727W, 62L/437G/489A/527R/727W/913R/932A,
 62L/437G/527R/727W, 62L/437G/527R/916G/932A, 62L/437G/913R/916R, 62L/489A/500A/932A,
 62L/489A/614Q/916R, 62L/527R, 62L/527R/727W/916G, 62W/89R/124V/381W/858C,
 62W/89R/149R/381W, 62W/89R/188W/489A/500A/727W/823F, 62W/89R/381V,
 62W/89R/381W/858C, 62W/124V/148R/149R/381W/614R,
 62W/124V/149R/381V/832R/858C/937K, 62W/124V/188W/823F/842G/862Q,
 62W/148R/149R/381V/858C/937K, 62W/148R/381W/614R/937K, 62W/148R/381W/727W,
 62W/148R/381W/858C, 62W/149R/381W/614R/937K, 62W/381V,
 62W/437G/489A/614Q/727W/913R, 62W/489A/527R/916R/932A, 62W/500A, 62W/727W,
 62W/916G, 89R/148R/149R, 89R/148R/149R/381W, 89R/381W,
 124V/148R/381W/727W/858W/937K, 124V/381W/614R, 124V/500A/842G/862Q,
 124V/832R/937K, 148R/832R/858W/937K, 381V, 381V/614R/832R, 381W/858C/937K, 437G,
 437G/489A/914R/916R, 437G/727W/914K, 437G/914R/916G, 489A/500A, 489A/614Q/916G,
 500A/727W/913R/916R, 500A/914R/916G, and 923W, wherein the positions are numbered with
 reference to SEQ ID NO: 946. In some embodiments, the acid alpha-glucosidase comprises at least
 one substitution at position or set of positions selected from S19T/S124V/P149R/E381V/S727W,
 W24R/Q39H/D489A/R862Q, W24R/P57L/A62W/A89R/D489A/L823F/R862Q,
 W24R/P57L/L823F/R862Q, W24R/A62W/A89R/E188W/L823V/S842G/R862Q,
 W24R/A89R/D489A, W24R/A89R/D489A/S727W/R862Q, W24R/D489A/D500A/S842G,
 Q39H/P57L/A62W/E188W/D500A/S842G, Q39H/P57L/D500A/R862Q,
 P57F/A62L/L305F/A437G/D500A/E614Q/S727W/S916R,
 P57F/A62L/L305F/A437G/E614Q/A683S/V913R/S916R/S932A,
 P57F/A62L/L305F/D500A/V913R/S916G, P57F/A62L/L305F/V913R,
 P57F/A62L/A437G/D500A/L761F/P914K/S916R, P57F/A62L/A437G/N527R/S727W,
 P57F/A62L/V913R/S916R/S932A, P57F/A62L/S916G, P57F/A62W/F120I/N527R/V913R/S916R,
 P57F/A62W/L305F/A437G/D500A/S727W/V913R/S916R,
 P57F/A62W/L305F/D489A/Q907K/V913R/S916G, P57F/A62W/L305F/D489A/V913R/S916G,

P57F/A62W/L305F/S916R, P57F/A62W/A437G/V913R/S916G, P57F/L305F, P57F/L305F/A437G/S916G, P57F/A437G/D500A/N527R/S727W/S916R, P57F/A437G/D500A/E614Q/S727W/P914R, P57F/A437G/V913R/P914R, P57F/D489A/N527R/P914R/S916G, P57F/E614Q/S916G/S932A, P57L, P57L/E188W/D489A/L823F/R862Q, A62F/A89R, A62F/A89R/S124V/T148R/E381W/E858W, A62F/A89R/T148R/E381V/E614R/E858W, A62F/A89R/T148R/D923W, A62F/A89R/P149R/E381W/P832R, A62F/A89R/E381V/D923W, A62F/A89R/E858C, A62F/E96K/E614R, A62F/S124V/E381W/P832R, A62F/P149R/E381V/E858W/E937K, A62F/P149R/S727W, A62F/E381V, A62F/E614R, A62L/L305F/A437G/D500A/S727W/V913R, A62L/L305F/S727W, A62L/A437G/D489A/N527R/S727W/V913R/S932A, A62L/A437G/N527R/S727W, A62L/A437G/N527R/S916G/S932A, A62L/A437G/V913R/S916R, A62L/D489A/D500A/S932A, A62L/D489A/E614Q/S916R, A62L/N527R, A62L/N527R/S727W/S916G, A62W/A89R/S124V/E381W/E858C, A62W/A89R/P149R/E381W, A62W/A89R/E188W/D489A/D500A/S727W/L823F, A62W/A89R/E381V, A62W/A89R/E381W/E858C, A62W/S124V/T148R/P149R/E381W/E614R, A62W/S124V/P149R/E381V/P832R/E858C/E937K, A62W/S124V/E188W/L823F/S842G/R862Q, A62W/T148R/P149R/E381V/E858C/E937K, A62W/T148R/E381W/E614R/E937K, A62W/T148R/E381W/S727W, A62W/T148R/E381W/E858C, A62W/P149R/E381W/E614R/E937K, A62W/E381V, A62W/A437G/D489A/E614Q/S727W/V913R, A62W/D489A/N527R/S916R/S932A, A62W/D500A, A62W/S727W, A62W/S916G, A89R/T148R/P149R, A89R/T148R/P149R/E381W, A89R/E381W, S124V/T148R/E381W/S727W/E858W/E937K, S124V/E381W/E614R, S124V/D500A/S842G/R862Q, S124V/P832R/E937K, T148R/P832R/E858W/E937K, E381V, E381V/E614R/P832R, E381W/E858C/E937K, A437G, A437G/D489A/P914R/S916R, A437G/S727W/P914R, A437G/P914R/S916G, D489A/D500A, D489A/E614Q/S916G, D500A/S727W/V913R/S916R, D500A/P914R/S916G, and D923W, wherein the positions are numbered with reference to SEQ ID NO: 946. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 57/62/120/527/913/916, 57/62/305/437/500/614/727/916, 57/62/305/437/500/727/913/916, 57/62/305/437/614/683/913/916/932, 57/62/305/489/907/913/916, 57/62/305/489/913/916, 57/62/305/500/913/916, 57/62/305/913, 57/62/305/916, 57/62/437/500/761/914/916, 57/62/437/527/727, 57/62/437/913/916, 57/62/913/916/932, 57/62/916, 57/188/489/823/862, 57/305, 57/305/437/916, 57/437/500/527/727/916, 57/437/500/614/727/914, 57/437/913/914, 57/489/527/914/916, 57/614/916/932, 62/89/188/489/500/727/823, 62/124/188/823/842/862, 62/305/437/500/727/913, 62/305/727, 62/437/489/527/727/913/932, 62/437/489/614/727/913, 62/437/527/727, 62/437/527/916/932, 62/437/913/916, 62/489/500/932, 62/489/527/916/932, 62/527/727/916, 62/727, 62/916, 124/500/842/862, 437, 437/489/914/916, 437/727/914, 437/914/916, 489/614/916, 500/727/913/916, 500/914/916, and 923, wherein the positions are numbered with

reference to SEQ ID NO: 946. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from

57F/62L/305F/437G/500A/614Q/727W/916R, 57F/62L/305F/437G/614Q/683S/913R/916R/932A, 57F/62L/305F/500A/913R/916G, 57F/62L/305F/913R, 57F/62L/437G/500A/761F/914K/916R, 57F/62L/437G/527R/727W, 57F/62L/913R/916R/932A, 57F/62L/916G, 57F/62W/120I/527R/913R/916R, 57F/62W/305F/437G/500A/727W/913R/916R, 57F/62W/305F/489A/907K/913R/916G, 57F/62W/305F/489A/913R/916G, 57F/62W/305F/916R, 57F/62W/437G/913R/916G, 57F/305F, 57F/305F/437G/916G, 57F/437G/500A/527R/727W/916R, 57F/437G/500A/614Q/727W/914R, 57F/437G/913R/914R, 57F/489A/527R/914R/916G, 57F/614Q/916G/932A, 57L/188W/489A/823F/862Q, 62L/305F/437G/500A/727W/913R, 62L/305F/727W, 62L/437G/489A/527R/727W/913R/932A, 62L/437G/527R/727W, 62L/437G/527R/916G/932A, 62L/437G/913R/916R, 62L/489A/500A/932A, 62L/527R/727W/916G, 62W/89R/188W/489A/500A/727W/823F, 62W/124V/188W/823F/842G/862Q, 62W/437G/489A/614Q/727W/913R, 62W/489A/527R/916R/932A, 62W/727W, 62W/916G, 124V/500A/842G/862Q, 437G, 437G/489A/914R/916R, 437G/727W/914K, 437G/914R/916G, 489A/614Q/916G, 500A/727W/913R/916R, 500A/914R/916G, and 923W, wherein the positions are numbered with reference to SEQ ID NO: 946. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from

P57F/A62L/L305F/A437G/D500A/E614Q/S727W/S916R,
P57F/A62L/L305F/A437G/E614Q/A683S/V913R/S916R/S932A,
P57F/A62L/L305F/D500A/V913R/S916G, P57F/A62L/L305F/V913R,
P57F/A62L/A437G/D500A/L761F/P914K/S916R, P57F/A62L/A437G/N527R/S727W,
P57F/A62L/V913R/S916R/S932A, P57F/A62L/S916G, P57F/A62W/F120I/N527R/V913R/S916R,
P57F/A62W/L305F/A437G/D500A/S727W/V913R/S916R,
P57F/A62W/L305F/D489A/Q907K/V913R/S916G, P57F/A62W/L305F/D489A/V913R/S916G,
P57F/A62W/L305F/S916R, P57F/A62W/A437G/V913R/S916G, P57F/L305F,
P57F/L305F/A437G/S916G, P57F/A437G/D500A/N527R/S727W/S916R,
P57F/A437G/D500A/E614Q/S727W/P914R, P57F/A437G/V913R/P914R,
P57F/D489A/N527R/P914R/S916G, P57F/E614Q/S916G/S932A,
P57L/E188W/D489A/L823F/R862Q, A62L/L305F/A437G/D500A/S727W/V913R,
A62L/L305F/S727W, A62L/A437G/D489A/N527R/S727W/V913R/S932A,
A62L/A437G/N527R/S727W, A62L/A437G/N527R/S916G/S932A, A62L/A437G/V913R/S916R,
A62L/D489A/D500A/S932A, A62L/N527R/S727W/S916G,
A62W/A89R/E188W/D489A/D500A/S727W/L823F, A62W/S124V/E188W/L823F/S842G/R862Q,
A62W/A437G/D489A/E614Q/S727W/V913R, A62W/D489A/N527R/S916R/S932A, A62W/S727W,
A62W/S916G, S124V/D500A/S842G/R862Q, A437G, A437G/D489A/P914R/S916R,
A437G/S727W/P914K, A437G/P914R/S916G, D489A/E614Q/S916G,

D500A/S727W/V913R/S916R, D500A/P914R/S916G, and D923W, wherein the positions are numbered with reference to SEQ ID NO: 946.

[0016] The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 1956, wherein the positions are numbered with reference to SEQ ID NO: 1956. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from , 3/569, 41/53, 44, 44/347, 56, 65, 78, 78/87/176/266/536/615, 78/87/266, 78/87/266/372/386/777, 78/87/266/372/536, 78/87/266/483/924, 78/87/483/777, 78/87/536, 78/266/483/536/615, 78/266/483/795, 78/266/763, 78/372/390, 78/390, 78/536/615, 87, 87/266, 87/266/372/483, 87/266/483, 87/266/924, 87/372/777, 87/536/777, 87/615, 87/795, 105, 136, 141, 145, 154/588, 156, 157, 199, 202, 222, 225, 227, 229, 266, 266/372/536/615/763/777, 266/372/924, 266/536/615/795, 344, 348, 390/615, 412, 423, 425/678/894, 430, 446, 484, 488, 496, 499/711, 503, 530, 543, 569, 572, 573, 574, 577, 578, 579, 580, 581, 583, 585, 588, 589/663, 615, 628, 629, 631, 633, 656, 663, 669, 670, 671, 678, 679, 687, 690, 691, 692, 693, 705, 706, 708, 709, 710, 711, 726, 768, 773, 777, 779, 795, 797, 816, 826, 834, 857, 859, 868, 869, 871, 873, 877, 878, and 909, wherein the positions are numbered with reference to SEQ ID NO: 1956. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 3L/569F, 41I/53M, 44E, 44G, 44L, 44P/347I, 44R, 56A, 56L, 56R, 65L, 65V, 78E, 78E/87E/176T/266N/536T/615D, 78E/87E/266N, 78E/87E/266N/372T/386Y/777G, 78E/87E/266N/372T/536T, 78E/87E/266N/483S/924N, 78E/87E/483S/777G, 78E/87E/536T, 78E/266N/483S/536T/615D, 78E/266N/483S/795E, 78E/266N/763L, 78E/372T/390Q, 78E/390Q, 78E/536T/615D, 87E, 87E/266N, 87E/266N/372T/483S, 87E/266N/483S, 87E/266N/924N, 87E/372T/777G, 87E/536T/777G, 87E/615D, 87E/795E, 105T, 136G, 141S, 141W, 145I, 145R, 154R/588L, 156L, 157S, 199V, 202K, 202L, 202N, 202R, 202T, 222C, 222P, 225D, 227A, 229C, 266N, 266N/372T/536T/615D/763L/777G, 266N/372T/924N, 266N/536T/615D/795E, 344G, 344M, 348G, 390Q/615D, 412Y, 423V, 425R/678I/894C, 430F, 446T, 484L, 488G, 488K, 488M, 496G, 499Y/711F, 503S, 503T, 530V, 543C, 543Q, 543S, 543V, 569H, 569I, 569Q, 569S, 569T, 569V, 569Y, 572G, 572S, 573C, 573D, 573H, 573M, 573Q, 574S, 577A, 577D, 577E, 577M, 577T, 577V, 578S, 579V, 580E, 580G, 580I, 580L, 580W, 580Y, 581F, 581G, 581H, 581L, 581S, 581T, 581V, 581Y, 583C, 583G, 583K, 583L, 585F, 585L, 585M, 585Q, 585V, 588L, 588V, 589I/663F, 615G, 628I, 628M, 628V, 629A, 629C, 629G, 629I, 631I, 631L, 631M, 633V, 656M, 656V, 663A, 663F, 669R, 670E, 670F, 670I, 670Q, 670R, 670S, 670T, 670V, 670W, 671A, 671G, 671M, 671T, 678H, 678L, 678T, 678Y, 679W, 687L, 690V, 691F, 691V, 692C, 692F, 692G, 692I, 692L, 692R, 692S, 692V, 692Y, 693F, 693I, 693Y, 705M, 706F, 706M, 706V, 708C, 709S, 710K, 710L, 710M, 710N,

710S, 711C, 711F, 711G, 711H, 711L, 711R, 711W, 726E, 768S, 773V, 777G, 777I, 777R, 779E, 779H, 779R, 795E, 797L, 797M, 816L, 826G, 834H, 857T, 859T, 859Y, 868I, 868L, 869L, 869S, 871E, 871K, 871R, 873A, 873F, 873Y, 877L, 877V, 878A, 878F, 878G, 878K, 878L, 878Q, 878R, 878S, 878W, and 909F, wherein the positions are numbered with reference to SEQ ID NO: 1956. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from W3L/L569F, L41I/S53M, T44E, T44G, T44L, T44P/L347I, T44R, G56A, G56L, G56R, G65L, G65V, P78E, P78E/D87E/K176T/T266N/V536T/Q615D, P78E/D87E/T266N, P78E/D87E/T266N/S372T/A386Y/S777G, P78E/D87E/T266N/S372T/V536T, P78E/D87E/T266N/T483S/T924N, P78E/D87E/T483S/S777G, P78E/D87E/V536T, P78E/T266N/T483S/V536T/Q615D, P78E/T266N/T483S/Q795E, P78E/T266N/T763L, P78E/S372T/L390Q, P78E/L390Q, P78E/V536T/Q615D, D87E, D87E/T266N, D87E/T266N/S372T/T483S, D87E/T266N/T483S, D87E/T266N/T924N, D87E/S372T/S777G, D87E/V536T/S777G, D87E/Q615D, D87E/Q795E, A105T, S136G, T141S, T141W, T145I, T145R, K154R/F588L, I156L, L157S, P199V, S202K, S202L, S202N, S202R, S202T, V222C, V222P, N225D, T227A, A229C, T266N, T266N/S372T/V536T/Q615D/T763L/S777G, T266N/S372T/T924N, T266N/V536T/Q615D/Q795E, Q344G, Q344M, D348G, L390Q/Q615D, G412Y, L423V, Q425R/S678I/G894C, Y430F, S446T, N484L, L488G, L488K, L488M, A496G, H499Y/A711F, P503S, P503T, L530V, T543C, T543Q, T543S, T543V, L569H, L569I, L569Q, L569S, L569T, L569V, L569Y, A572G, A572S, I573C, I573D, I573H, I573M, I573Q, A574S, R577A, R577D, R577E, R577M, R577T, R577V, A578S, L579V, V580E, V580G, V580I, V580L, V580W, V580Y, K581F, K581G, K581H, K581L, K581S, K581T, K581V, K581Y, R583C, R583G, R583K, R583L, T585F, T585L, T585M, T585Q, T585V, F588L, F588V, V589I/M663F, Q615G, L628I, L628M, L628V, L629A, L629C, L629G, L629I, V631I, V631L, V631M, L633V, L656M, L656V, M663A, M663F, L669R, L670E, L670F, L670I, L670Q, L670R, L670S, L670T, L670V, L670W, S671A, S671G, S671M, S671T, S678H, S678L, S678T, S678Y, F679W, M687L, A690V, L691F, L691V, T692C, T692F, T692G, T692I, T692L, T692R, T692S, T692V, T692Y, L693F, L693I, L693Y, F705M, H706F, H706M, H706V, A708C, H709S, V710K, V710L, V710M, V710N, V710S, A711C, A711F, A711G, A711H, A711L, A711R, A711W, D726E, Q768S, E773V, S777G, S777I, S777R, P779E, P779H, P779R, Q795E, V797L, V797M, I816L, T826G, A834H, L857T, V859T, V859Y, V868I, V868L, I869L, I869S, L871E, L871K, L871R, R873A, R873F, R873Y, I877L, I877V, V878A, V878F, V878G, V878K, V878L, V878Q, V878R, V878S, V878W, and L909F, wherein the positions are numbered with reference to SEQ ID NO: 1956. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position selected from 44, 53, 56, 63, 65, 105, 125, 129, 136, 139, 141, 142, 145, 152, 156, 162, 176, 177, 185, 186, 187, 199, 199/775, 202, 265, 267, 337, 344, 348, 350, 354, 372, 373, 401, 412, 446, 469, 484, 488, 493, 496, 499, 503, 526, 543, 612, 615, 649, 677, 678, 679, 730, 752, 765, 768, 773, 777, 779, 788, 797, 822, 826, 834, 855, 856, 857, 859, 860, 924, 926, 931, and 936, wherein the positions are numbered

with reference to SEQ ID NO: 1956. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position selected from 44A, 44F, 44V, 44W, 44Y, 53I, 56S, 56W, 63N, 65A, 65F, 65R, 65Y, 105V, 105W, 125H, 125W, 129E, 129I, 129S, 129T, 129V, 129W, 136K, 136R, 136V, 139E, 141K, 141R, 142G, 145A, 145L, 152L, 152S, 152W, 156C, 156K, 156R, 156S, 162T, 176R, 177Q, 185L, 186H, 187I, 199A, 199G, 199I, 199R, 199T, 199V/775I, 199W, 202A, 202D, 202G, 202H, 202Q, 202Y, 265D, 265F, 265H, 267E, 267G, 267R, 337H, 344C, 348E, 348W, 350F, 350I, 354L, 354S, 372D, 373A, 373S, 401G, 401S, 412R, 412S, 412W, 446C, 446D, 446G, 446I, 446K, 469M, 469T, 469V, 484A, 484K, 484R, 488C, 488E, 488S, 493L, 496M, 496W, 499A, 499E, 499I, 499M, 499Q, 499V, 503C, 503H, 503N, 526L, 526V, 543G, 543H, 543K, 543L, 543R, 612G, 612L, 612R, 612T, 615M, 615S, 649M, 677T, 678Q, 678R, 678V, 678W, 679Y, 730K, 730L, 730R, 752F, 752G, 752L, 752N, 752S, 752W, 765W, 768I, 768K, 768V, 773P, 777M, 777W, 779I, 779M, 779S, 788A, 788H, 788I, 788L, 788N, 788Q, 788S, 788T, 788Y, 797E, 797F, 797I, 797R, 797W, 822R, 826I, 826M, 834G, 834S, 834V, 834W, 855G, 855L, 856A, 856G, 857A, 857E, 857R, 857S, 857V, 859A, 859G, 860S, 924A, 926M, 926T, 931L, 936N, 936R, and 936S, wherein the positions are numbered with reference to SEQ ID NO: 1956. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position selected from T44A, T44F, T44V, T44W, T44Y, S53I, G56S, G56W, H63N, G65A, G65F, G65R, G65Y, A105V, A105W, Y125H, Y125W, K129E, K129I, K129S, K129T, K129V, K129W, S136K, S136R, S136V, G139E, T141K, T141R, A142G, T145A, T145L, F152L, F152S, F152W, I156C, I156K, I156R, I156S, D162T, K176R, D177Q, V185L, P186H, L187I, P199A, P199G, P199I, P199R, P199T, P199V/L775I, P199W, S202A, S202D, S202G, S202H, S202Q, S202Y, W265D, W265F, W265H, K267E, K267G, K267R, P337H, Q344C, D348E, D348W, V350F, V350I, F354L, F354S, S372D, T373A, T373S, D401G, D401S, G412R, G412S, G412W, S446C, S446D, S446G, S446I, S446K, I469M, I469T, I469V, N484A, N484K, N484R, L488C, L488E, L488S, D493L, A496M, A496W, H499A, H499E, H499I, H499M, H499Q, H499V, P503C, P503H, P503N, P526L, P526V, T543G, T543H, T543K, T543L, T543R, S612G, S612L, S612R, S612T, Q615M, Q615S, L649M, Y677T, S678Q, S678R, S678V, S678W, F679Y, W730K, W730L, W730R, K752F, K752G, K752L, K752N, K752S, K752W, Y765W, Q768I, Q768K, Q768V, E773P, S777M, S777W, P779I, P779M, P779S, P788A, P788H, P788I, P788L, P788N, P788Q, P788S, P788T, P788Y, V797E, V797F, V797I, V797R, V797W, G822R, T826I, T826M, A834G, A834S, A834V, A834W, E855G, E855L, S856A, S856G, L857A, L857E, L857R, L857S, L857V, V859A, V859G, L860S, T924A, V926M, V926T, V931L, G936N, G936R, and G936S, wherein the positions are numbered with reference to SEQ ID NO: 1956. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 78/87/176/266/536/615, 78/87/266/372/386/777, 78/87/266/372/536, 78/266/763, 78/372/390, 87/266/372/483, 87/372/777, 105, 125, 129, 136, 139, 141, 142, 152, 154/588, 156, 222, 225, 227, 229, 266/372/536/615/763/777, 266/372/924, 267, 372, 401, 493, 496, 499, 569, 572, 573, 574, 577, 579, 580, 581, 583, 585, 588, 589/663, 628, 629, 631, 663, 669, 670,

671, 691, 692, 693, 706, 708, 710, 711, 765, 768, 779, 797, 826, 834, 855, 856, 857, 869, 871, 873, 878, 909, 924, and 926, wherein the positions are numbered with reference to SEQ ID NO: 1956. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 78E/87E/176T/266N/536T/615D, 78E/87E/266N/372T/386Y/777G, 78E/87E/266N/372T/536T, 78E/266N/763L, 78E/372T/390Q, 87E/266N/372T/483S, 87E/372T/777G, 105T, 105W, 125W, 129E, 129S, 136G, 136K, 136V, 139E, 141S, 142G, 152S, 152W, 154R/588L, 156C, 156L, 222C, 222P, 225D, 227A, 229C, 266N/372T/536T/615D/763L/777G, 266N/372T/924N, 267E, 372D, 401G, 493L, 496G, 499E, 499I, 499M, 499Q, 569H, 569Q, 569S, 569T, 569V, 569Y, 572G, 572S, 573C, 573D, 573H, 573M, 573Q, 574S, 577A, 577D, 577E, 577T, 579V, 580E, 580G, 580W, 581G, 581H, 581T, 583C, 583G, 585Q, 588L, 588V, 589I/663F, 628V, 629A, 629C, 629G, 629I, 631I, 631L, 631M, 663A, 663F, 669R, 670E, 670F, 670I, 670Q, 670R, 670S, 670T, 670V, 670W, 671A, 671G, 671T, 691V, 692C, 692F, 692G, 692I, 692L, 692Y, 693F, 693I, 693Y, 706F, 708C, 710K, 710L, 710M, 710N, 710S, 711C, 711G, 711H, 711W, 765W, 768I, 779E, 797E, 797F, 797I, 797L, 797M, 797W, 826G, 834G, 834H, 834S, 834W, 855G, 856G, 857A, 857E, 857S, 857T, 857V, 869L, 869S, 871E, 871K, 873A, 873F, 873Y, 878A, 878G, 878K, 878Q, 878S, 878W, 909F, 924A, 926M, and 926T, wherein the positions are numbered with reference to SEQ ID NO: 1956. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from P78E/D87E/K176T/T266N/V536T/Q615D, P78E/D87E/T266N/S372T/A386Y/S777G, P78E/D87E/T266N/S372T/V536T, P78E/T266N/T763L, P78E/S372T/L390Q, D87E/T266N/S372T/T483S, D87E/S372T/S777G, A105T, A105W, Y125W, K129E, K129S, S136G, S136K, S136V, G139E, T141S, A142G, F152S, F152W, K154R/F588L, I156C, I156L, V222C, V222P, N225D, T227A, A229C, T266N/S372T/V536T/Q615D/T763L/S777G, T266N/S372T/T924N, K267E, S372D, D401G, D493L, A496G, H499E, H499I, H499M, H499Q, L569H, L569Q, L569S, L569T, L569V, L569Y, A572G, A572S, I573C, I573D, I573H, I573M, I573Q, A574S, R577A, R577D, R577E, R577T, L579V, V580E, V580G, V580W, K581G, K581H, K581T, R583C, R583G, T585Q, F588L, F588V, V589I/M663F, L628V, L629A, L629C, L629G, L629I, V631I, V631L, V631M, M663A, M663F, L669R, L670E, L670F, L670I, L670Q, L670R, L670S, L670T, L670V, L670W, S671A, S671G, S671T, L691V, T692C, T692F, T692G, T692I, T692L, T692Y, L693F, L693I, L693Y, H706F, A708C, V710K, V710L, V710M, V710N, V710S, A711C, A711G, A711H, A711W, Y765W, Q768I, P779E, V797E, V797F, V797I, V797L, V797M, V797W, T826G, A834G, A834H, A834S, A834W, E855G, S856G, L857A, L857E, L857S, L857T, L857V, I869L, I869S, L871E, L871K, R873A, R873F, R873Y, V878A, V878G, V878K, V878Q, V878S, V878W, L909F, T924A, V926M, and V926T, wherein the positions are numbered with reference to SEQ ID NO: 1956.

[0017] The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising

at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 2496, wherein the positions are numbered with reference to SEQ ID NO: 2496. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 246, 304, 313, 569, 569/588, 569/588/589/628/629/692/711, 569/588/678/692, 569/588/711/869/871/878, 569/588/878, 569/589/628/670/678/692/711/795/871, 569/589/628/670/692/711, 569/589/628/692/711/795, 569/589/670, 569/589/670/678/692/711/795, 569/589/670/871, 569/589/678/871/878, 569/589/692/795/871/878, 569/589/711/871, 569/589/871, 569/628, 569/628/670, 569/628/670/678, 569/628/670/692/711/871, 569/628/670/711, 569/628/678/711, 569/628/692, 569/670, 569/670/678, 569/670/678/692/871, 569/670/692, 569/670/711, 569/670/711/871, 569/678/692/795, 569/678/869/878, 569/678/878, 569/692, 569/692/711, 569/692/711/869/871/878, 569/711, 569/711/795/871/878, 569/711/869/878, 569/711/871, 569/795, 569/871, 572/588/678/692/869/878, 572/588/795, 572/692/869/878, 572/692/878, 582, 584, 585, 588/589/628/678, 589, 589/670/692/795/871, 589/670/795/871, 589/871, 628/629/692/871/878, 628/670, 628/670/692/711/795, 628/711/795, 628/871, 628/878, 629/869/878, 670/678, 670/678/692/871, 670/692/871, 678/692/711/869, 678/692/795/869, 678/692/869, 678/795/871/878, 692, 692/711, 692/711/795/869, 692/711/795/869/871/878, 692/711/869/878, 692/711/871/878, 692/869, 692/869/871/878/916, 692/871, 711, 711/795/869/878, 711/869/878, 711/871, 795/878, 812, 871, and 878, wherein the positions are numbered with reference to SEQ ID NO: 2496. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 246T, 304M, 313I, 313L, 569H/589I/628M/670T/678T/692G/711H/795E/871S, 569H/628M, 569H/628M/670F/678T, 569H/628M/670T, 569H/628M/670T/692Y/711H/871K, 569H/670F/678T/692G/871S, 569H/670T/692G, 569H/678T/692G/795E, 569H/692G, 569H/692Y, 569H/711H/795E/871S/878S, 569H/711H/869S/878S, 569H/711H/871K, 569T, 569T/588L, 569T/588L/589I/628M/629I/692Y/711H, 569T/588L/678T/692Y, 569T/588L/711H/869L/871K/878S, 569T/588L/878S, 569T/589I/628M/670F/692G/711H, 569T/589I/678T/871K/878S, 569T/589I/692G/795E/871K/878S, 569T/589I/871S, 569T/628M/678T/711H, 569T/628M/692Y, 569T/670T/678T/692G/871K, 569T/670T/711H/871E, 569T/678T/869S/878S, 569T/678T/878S, 569T/692Y, 569T/692Y/711H/869L/871K/878A, 569T/711H, 569T/795E, 569T/871K, 569T/871S, 569Y/589I/628M/670T/692Y/711H, 569Y/589I/628M/692G/711H/795E, 569Y/589I/670F, 569Y/589I/670T/678T/692G/711H/795E, 569Y/589I/670T/871K, 569Y/589I/711H/871K, 569Y/628M/670T/692Y/711H/871S, 569Y/628M/670T/711H, 569Y/670F/678T, 569Y/670T, 569Y/670T/711H, 569Y/692Y/711H, 569Y/711H, 569Y/871E, 572S/588L/678T/692G/869L/878A, 572S/588L/795E, 572S/692G/869S/878S, 572S/692G/878S, 572S/692L/869S/878S, 582T, 584E, 585K, 588L/589I/628M/678T, 589I, 589I/670T/692G/795E/871K, 589I/670T/795E/871S, 589I/871E,

628M/629I/692Y/871S/878S, 628M/670F, 628M/670T/692G/711H/795E, 628M/711H/795E, 628M/871S, 628M/878S, 629I/869L/878S, 670T/678T, 670T/678T/692Y/871S, 670T/692G/871K, 678T/692G/711H/869S, 678T/692G/795E/869S, 678T/692G/869S, 678T/795E/871K/878A, 692G/711H, 692G/711H/795E/869L/871K/878A, 692G/711H/795E/869S, 692G/711H/869L/878S, 692G/711H/869S/878A, 692G/711H/871S/878A, 692G/869L/871K/878S/916R, 692G/869S, 692G/871K, 692Y, 711H, 711H/795E/869S/878S, 711H/869S/878S, 711H/871K, 795E/878S, 812E, 871K, 871S, and 878S, wherein the positions are numbered with reference to SEQ ID NO: 2496. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from S246T, L304M, V313I, V313L, L569H/V589I/L628M/L670T/S678T/T692G/A711H/Q795E/L871S, L569H/L628M, L569H/L628M/L670F/S678T, L569H/L628M/L670T, L569H/L628M/L670T/T692Y/A711H/L871K, L569H/L670F/S678T/T692G/L871S, L569H/L670T/T692G, L569H/S678T/T692G/Q795E, L569H/T692G, L569H/T692Y, L569H/A711H/Q795E/L871S/V878S, L569H/A711H/I869S/V878S, L569H/A711H/L871K, L569T, L569T/F588L, L569T/F588L/V589I/L628M/L629I/T692Y/A711H, L569T/F588L/S678T/T692Y, L569T/F588L/A711H/I869L/L871K/V878S, L569T/F588L/V878S, L569T/V589I/L628M/L670F/T692G/A711H, L569T/V589I/S678T/L871K/V878S, L569T/V589I/T692G/Q795E/L871K/V878S, L569T/V589I/L871S, L569T/L628M/S678T/A711H, L569T/L628M/T692Y, L569T/L670T/S678T/T692G/L871K, L569T/L670T/A711H/L871E, L569T/S678T/I869S/V878S, L569T/S678T/V878S, L569T/T692Y, L569T/T692Y/A711H/I869L/L871K/V878A, L569T/A711H, L569T/Q795E, L569T/L871K, L569T/L871S, L569Y/V589I/L628M/L670T/T692Y/A711H, L569Y/V589I/L628M/T692G/A711H/Q795E, L569Y/V589I/L670F, L569Y/V589I/L670T/S678T/T692G/A711H/Q795E, L569Y/V589I/L670T/L871K, L569Y/V589I/A711H/L871K, L569Y/L628M/L670T/T692Y/A711H/L871S, L569Y/L628M/L670T/A711H, L569Y/L670F/S678T, L569Y/L670T, L569Y/L670T/A711H, L569Y/T692Y/A711H, L569Y/A711H, L569Y/L871E, A572S/F588L/S678T/T692G/I869L/V878A, A572S/F588L/Q795E, A572S/T692G/I869S/V878S, A572S/T692G/V878S, A572S/T692L/I869S/V878S, A582T, G584E, T585K, F588L/V589I/L628M/S678T, V589I, V589I/L670T/T692G/Q795E/L871K, V589I/L670T/Q795E/L871S, V589I/L871E, L628M/L629I/T692Y/L871S/V878S, L628M/L670F, L628M/L670T/T692G/A711H/Q795E, L628M/A711H/Q795E, L628M/L871S, L628M/V878S, L629I/I869L/V878S, L670T/S678T, L670T/S678T/T692Y/L871S, L670T/T692G/L871K, S678T/T692G/A711H/I869S, S678T/T692G/Q795E/I869S, S678T/T692G/I869S, S678T/Q795E/L871K/V878A, T692G/A711H, T692G/A711H/Q795E/I869L/L871K/V878A, T692G/A711H/Q795E/I869S, T692G/A711H/I869L/V878S, T692G/A711H/I869S/V878A, T692G/A711H/L871S/V878A, T692G/I869L/L871K/V878S/S916R, T692G/I869S, T692G/L871K, T692Y, A711H, A711H/Q795E/I869S/V878S, A711H/I869S/V878S, A711H/L871K, Q795E/V878S, A812E, L871K,

L871S, and V878S, wherein the positions are numbered with reference to SEQ ID NO: 2496. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 60/589, 307, 313, 584, and 810, wherein the positions are numbered with reference to SEQ ID NO: 2496. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 60V/589A, 307T, 313T, 584C, and 810V, wherein the positions are numbered with reference to SEQ ID NO: 2496. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from A60V/V589A, S307T, V313T, G584C, and L810V, wherein the positions are numbered with reference to SEQ ID NO: 2496. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 60/589, 246, 304, 307, 313, 569, 569/588, 569/588/589/628/629/692/711, 569/588/678/692, 569/588/711/869/871/878, 569/588/878, 569/589/628/670/678/692/711/795/871, 569/589/628/670/692/711, 569/589/628/692/711/795, 569/589/670, 569/589/670/678/692/711/795, 569/589/670/871, 569/589/678/871/878, 569/589/692/795/871/878, 569/589/711/871, 569/589/871, 569/628, 569/628/670, 569/628/670/678, 569/628/670/692/711/871, 569/628/670/711, 569/628/678/711, 569/628/692, 569/670, 569/670/678, 569/670/678/692/871, 569/670/692, 569/670/711, 569/670/711/871, 569/678/692/795, 569/678/869/878, 569/678/878, 569/692, 569/692/711, 569/692/711/869/871/878, 569/711, 569/711/795/871/878, 569/711/869/878, 569/711/871, 569/795, 569/871, 572/588/678/692/869/878, 572/588/795, 572/692/869/878, 572/692/878, 582, 584, 585, 588/589/628/678, 589, 589/670/692/795/871, 589/670/795/871, 589/871, 628/629/692/871/878, 628/670, 628/670/692/711/795, 628/711/795, 628/871, 628/878, 629/869/878, 670/678, 670/678/692/871, 670/692/871, 678/692/711/869, 678/692/795/869, 678/692/869, 678/795/871/878, 692, 692/711, 692/711/795/869, 692/711/795/869/871/878, 692/711/869/878, 692/711/871/878, 692/869, 692/869/871/878/916, 692/871, 711, 711/795/869/878, 711/869/878, 711/871, 795/878, 810, 812, 871, and 878, wherein the positions are numbered with reference to SEQ ID NO: 2496. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 60V/589A, 246T, 304M, 307T, 313I, 313L, 313T, 569H/589I/628M/670T/678T/692G/711H/795E/871S, 569H/628M, 569H/628M/670F/678T, 569H/628M/670T, 569H/628M/670T/692Y/711H/871K, 569H/670F/678T/692G/871S, 569H/670T/692G, 569H/678T/692G/795E, 569H/692G, 569H/692Y, 569H/711H/795E/871S/878S, 569H/711H/869S/878S, 569H/711H/871K, 569T, 569T/588L, 569T/588L/589I/628M/629I/692Y/711H, 569T/588L/678T/692Y, 569T/588L/711H/869L/871K/878S, 569T/588L/878S, 569T/589I/628M/670F/692G/711H, 569T/589I/678T/871K/878S, 569T/589I/692G/795E/871K/878S, 569T/589I/871S, 569T/628M/678T/711H, 569T/628M/692Y, 569T/670T/678T/692G/871K, 569T/670T/711H/871E, 569T/678T/869S/878S, 569T/678T/878S, 569T/692Y, 569T/692Y/711H/869L/871K/878A, 569T/711H, 569T/795E, 569T/871K, 569T/871S, 569Y/589I/628M/670T/692Y/711H,

569Y/589I/628M/692G/711H/795E, 569Y/589I/670F, 569Y/589I/670T/678T/692G/711H/795E, 569Y/589I/670T/871K, 569Y/589I/711H/871K, 569Y/628M/670T/692Y/711H/871S, 569Y/628M/670T/711H, 569Y/670F/678T, 569Y/670T, 569Y/670T/711H, 569Y/692Y/711H, 569Y/711H, 569Y/871E, 572S/588L/678T/692G/869L/878A, 572S/588L/795E, 572S/692G/869S/878S, 572S/692G/878S, 572S/692L/869S/878S, 582T, 584C, 584E, 585K, 588L/589I/628M/678T, 589I, 589I/670T/692G/795E/871K, 589I/670T/795E/871S, 589I/871E, 628M/629I/692Y/871S/878S, 628M/670F, 628M/670T/692G/711H/795E, 628M/711H/795E, 628M/871S, 628M/878S, 629I/869L/878S, 670T/678T, 670T/678T/692Y/871S, 670T/692G/871K, 678T/692G/711H/869S, 678T/692G/795E/869S, 678T/692G/869S, 678T/795E/871K/878A, 692G/711H, 692G/711H/795E/869L/871K/878A, 692G/711H/795E/869S, 692G/711H/869L/878S, 692G/711H/869S/878A, 692G/711H/871S/878A, 692G/869L/871K/878S/916R, 692G/869S, 692G/871K, 692Y, 711H, 711H/795E/869S/878S, 711H/869S/878S, 711H/871K, 795E/878S, 810V, 812E, 871K, 871S, and 878S, wherein the positions are numbered with reference to SEQ ID NO: 2496. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from A60V/V589A, S246T, L304M, S307T, V313I, V313L, V313T, L569H/V589I/L628M/L670T/S678T/T692G/A711H/Q795E/L871S, L569H/L628M, L569H/L628M/L670F/S678T, L569H/L628M/L670T, L569H/L628M/L670T/T692Y/A711H/L871K, L569H/L670F/S678T/T692G/L871S, L569H/L670T/T692G, L569H/S678T/T692G/Q795E, L569H/T692G, L569H/T692Y, L569H/A711H/Q795E/L871S/V878S, L569H/A711H/I869S/V878S, L569H/A711H/L871K, L569T, L569T/F588L, L569T/F588L/V589I/L628M/L629I/T692Y/A711H, L569T/F588L/S678T/T692Y, L569T/F588L/A711H/I869L/L871K/V878S, L569T/F588L/V878S, L569T/V589I/L628M/L670F/T692G/A711H, L569T/V589I/S678T/L871K/V878S, L569T/V589I/T692G/Q795E/L871K/V878S, L569T/V589I/L871S, L569T/L628M/S678T/A711H, L569T/L628M/T692Y, L569T/L670T/S678T/T692G/L871K, L569T/L670T/A711H/L871E, L569T/S678T/I869S/V878S, L569T/S678T/V878S, L569T/T692Y, L569T/T692Y/A711H/I869L/L871K/V878A, L569T/A711H, L569T/Q795E, L569T/L871K, L569T/L871S, L569Y/V589I/L628M/L670T/T692Y/A711H, L569Y/V589I/L628M/T692G/A711H/Q795E, L569Y/V589I/L670F, L569Y/V589I/L670T/S678T/T692G/A711H/Q795E, L569Y/V589I/L670T/L871K, L569Y/V589I/A711H/L871K, L569Y/L628M/L670T/T692Y/A711H/L871S, L569Y/L628M/L670T/A711H, L569Y/L670F/S678T, L569Y/L670T, L569Y/L670T/A711H, L569Y/T692Y/A711H, L569Y/A711H, L569Y/L871E, A572S/F588L/S678T/T692G/I869L/V878A, A572S/F588L/Q795E, A572S/T692G/I869S/V878S, A572S/T692G/V878S, A572S/T692L/I869S/V878S, A582T, G584C, G584E, T585K, F588L/V589I/L628M/S678T, V589I, V589I/L670T/T692G/Q795E/L871K, V589I/L670T/Q795E/L871S, V589I/L871E, L628M/L629I/T692Y/L871S/V878S, L628M/L670F, L628M/L670T/T692G/A711H/Q795E, L628M/A711H/Q795E, L628M/L871S, L628M/V878S, L629I/I869L/V878S, L670T/S678T,

L670T/S678T/T692Y/L871S, L670T/T692G/L871K, S678T/T692G/A711H/I869S, S678T/T692G/Q795E/I869S, S678T/T692G/I869S, S678T/Q795E/L871K/V878A, T692G/A711H, T692G/A711H/Q795E/I869L/L871K/V878A, T692G/A711H/Q795E/I869S, T692G/A711H/I869L/V878S, T692G/A711H/I869S/V878A, T692G/A711H/L871S/V878A, T692G/I869L/L871K/V878S/S916R, T692G/I869S, T692G/L871K, T692Y, A711H, A711H/Q795E/I869S/V878S, A711H/I869S/V878S, A711H/L871K, Q795E/V878S, L810V, A812E, L871K, L871S, and V878S, wherein the positions are numbered with reference to SEQ ID NO: 2496.

[0018] The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 2880, wherein the positions are numbered with reference to SEQ ID NO: 2880. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from

24/28/29/39/50/62/78/87/135/150/266/267/522/527/551/670/727/750/830/842/871/883/894/932, 24/28/39/50/62/78/87/135/150/266/267/522/527/551/569/727/830/842/871/883/894/913, 24/28/50/135/150/437/522/527/871/883/894/932, 24/28/62/522/569/932, 24/28/437/486/527, 24/29/39/50/62/78/87/135/150/267/437/486/522/527/551/711/727/750/830/842/871/883/894/913/932, 24/50/78/87/135/150/267/486/522/527/551/670/727/750/830/842/871/883/894/913/932, 24/50/486/527/711/727, 24/62/87/486/727, 24/62/727/830/932, 24/87/135/522/670/711/830/842/913, 24/150/522/527/727/883/894, 24/527/727/842/871/883/913/932, 24/670/727/750/842/871, 28, 28/50/78/87/135/266/267/437/486/527/551/670/727/750/830/842/871/883/894/913/932, 28/50/522/527/711/727/871, 28/62, 28/62/267/932, 28/437/527/871, 28/522/527/569/711/830/894, 28/727, 28/727/871, 29/39/50/62/65/78/87/135/150/437/551/569/670/727/750/830/842/883/894/932, 29/62/437/527, 29/78/87/150/527/727, 29/78/135/727/830, 29/87, 29/135/150/527/670/727/883, 29/150/267/727/750/871/883/932, 29/150/437/727, 29/522/670/711/871, 29/670/932, 39/50, 39/727/750/932, 50/135/150/932, 50/437/522/527, 50/711, 50/727/750/883/894, 62, 62/87/150, 62/87/150/727, 62/135/522/711/727/750/842/871/894, 62/437, 62/437/727, 78/87/486/527/670/727/750/830/842/871/913/932, 87/750, 89/109/527/678/727/842, 89/109/678/727/736/812/878, 89/109/727/932, 89/109/932, 89/527, 89/527/678/692/736/842/878/932, 89/527/678/932, 89/527/727/812/860, 89/678/692/736/932, 89/678/812/878, 89/842/878, 109/527/678/812, 109/678/692/842/860/878/932, 109/678/727/860, 109/678/736/812/878, 109/678/812, 109/678/842/878, 109/692/727/736/812, 109/692/727/812/842/860, 109/727/860/878, 109/736/932, 109/812, 109/842, 109/932, 135/670/727, 135/711/750/932, 150/527/842/871/913, 150/871/932, 150/883/932, 267/527/727, 403/527/678/692/736/812/842/860, 437/522/527/670/871, 437/750/830/932, 522, 522/527/569/727, 522/830, 527, 527/678/692/727/736/878, 527/678/692/812/932, 527/692/727/736/812, 527/692/727/736/842/860/878, 527/727/736,

527/736/932, 527/812, 670/711/871, 670/830/871, 678/692/727/812/842, 678/692/812, 678/812, 678/860/878, 678/913, 678/932, 692/727/736/842/913, 692/812, 727, 727/932, 871, and 878/932, wherein the positions are numbered with reference to SEQ ID NO: 2880. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from wherein the positions are numbered with reference to SEQ ID NO: 2880. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 24L/28L/29L/39P/50Q/62A/78P/87D/135S/150T/266T/267R/522E/527N/551V/670L/727S/750A/830Q/842G/871L/883R/894Q/932S, 24L/28L/39P/50Q/62A/78P/87D/135S/150T/266T/267R/522E/527N/551V/569L/727S/830Q/842G/871L/883R/894Q/913V, 24L/28L/50Q/135S/150T/437A/522E/527N/871L/883R/894Q/932S, 24L/28L/62A/522E/569L/932S, 24L/28L/437A/486T/527N, 24L/29L/39P/50Q/62A/78P/87D/135S/150T/267R/437A/486T/522E/527N/551V/711A/727S/750A/830Q/842G/871L/883R/894Q/913V/932S, 24L/50Q/78P/87D/135S/150T/267R/486T/522E/527N/551V/670L/727S/750A/830Q/842G/871L/883R/894Q/913V/932S, 24L/50Q/486T/527N/711A/727S, 24L/62A/87D/486T/727S, 24L/62A/727S/830Q/932S, 24L/87D/135S/522E/670L/711A/830Q/842G/913V, 24L/150T/522E/527N/727S/883R/894Q, 24L/527N/727S/842G/871L/883R/913V/932S, 24L/670L/727S/750A/842G/871L, 28L, 28L/50Q/78P/87D/135S/266T/267R/437A/486T/527N/551V/670L/727S/750A/830Q/842G/871L/883R/894Q/913V/932S, 28L/50Q/522E/527N/711A/727S/871L, 28L/62A, 28L/62A/267R/932S, 28L/437A/527N/871L, 28L/522E/527N/569L/711A/830Q/894Q, 28L/727S, 28L/727S/871L, 29L/39P/50Q/62A/65R/78P/87D/135S/150T/437A/551V/569L/670L/727S/750A/830Q/842G/883R/894Q/932S, 29L/62A/437A/527N, 29L/78P/87D/150T/527N/727S, 29L/78P/135S/727S/830Q, 29L/87D, 29L/135S/150T/527N/670L/727S/883R, 29L/150T/267R/727S/750A/871L/883R/932S, 29L/150T/437A/727S, 29L/522E/670L/711A/871L, 29L/670L/932S, 39P/50Q, 39P/727S/750A/932S, 50Q/135S/150T/932S, 50Q/437A/522E/527N, 50Q/711A, 50Q/727S/750A/883R/894Q, 62A, 62A/87D/150T, 62A/87D/150T/727S, 62A/135S/522E/711A/727S/750A/842G/871L/894Q, 62A/437A, 62A/437A/727S, 78P/87D/486T/527N/670L/727S/750A/830Q/842G/871L/913V/932S, 87D/750A, 89R/109D/527N/678T/727S/842G, 89R/109D/678T/727S/736M/812E/878S, 89R/109D/727S/932S, 89R/109D/932S, 89R/527N, 89R/527N/678T/692G/736M/842G/878S/932S, 89R/527N/678T/932S, 89R/527N/727S/812E/860F, 89R/678T/692G/736M/932S, 89R/678T/812E/878S, 89R/842G/878S, 109D/527N/678T/812E, 109D/678T/692G/842G/860F/878S/932S, 109D/678T/727S/860F, 109D/678T/736M/812E/878S, 109D/678T/812E, 109D/678T/842G/878S, 109D/692G/727S/736M/812E, 109D/692G/727S/812E/842G/860F, 109D/727S/860F/878S, 109D/736M/932S, 109D/812E, 109D/842G, 109D/932S, 135S/670L/727S, 135S/711A/750A/932S, 150T/527N/842G/871L/913V, 150T/871L/932S, 150T/883R/932S, 267R/527N/727S,

403H/527N/678T/692G/736M/812E/842G/860F, 437A/522E/527N/670L/871L, 437A/750A/830Q/932S, 522E, 522E/527N/569L/727S, 522E/830Q, 527N, 527N/678T/692G/727S/736M/878S, 527N/678T/692G/812E/932S, 527N/692G/727S/736M/812E, 527N/692G/727S/736M/842G/860F/878S, 527N/727S/736M, 527N/736M/932S, 527N/812E, 670L/711A/871L, 670L/830Q/871L, 678T/692G/727S/812E/842G, 678T/692G/812E, 678T/812E, 678T/860F/878S, 678T/913V, 678T/932S, 692G/727S/736M/842G/913V, 692G/812E, 727S, 727S/932S, 871L, and 878S/932S, wherein the positions are numbered with reference to SEQ ID NO: 2880. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from

W24L/S28L/T29L/Q39P/V50Q/L62A/E78P/E87D/Q135S/S150T/N266T/K267R/V522E/R527N/A551V/T670L/W727S/P750A/K830Q/S842G/E871L/H883R/G894Q/A932S, W24L/S28L/Q39P/V50Q/L62A/E78P/E87D/Q135S/S150T/N266T/K267R/V522E/R527N/A551V/T569L/W727S/K830Q/S842G/E871L/H883R/G894Q/R913V, W24L/S28L/V50Q/Q135S/S150T/G437A/V522E/R527N/E871L/H883R/G894Q/A932S, W24L/S28L/L62A/V522E/T569L/A932S, W24L/S28L/G437A/E486T/R527N, W24L/T29L/Q39P/V50Q/L62A/E78P/E87D/Q135S/S150T/K267R/G437A/E486T/V522E/R527N/A551V/H711A/W727S/P750A/K830Q/S842G/E871L/H883R/G894Q/R913V/A932S, W24L/V50Q/E78P/E87D/Q135S/S150T/K267R/E486T/V522E/R527N/A551V/T670L/W727S/P750A/K830Q/S842G/E871L/H883R/G894Q/R913V/A932S, W24L/V50Q/E486T/R527N/H711A/W727S, W24L/L62A/E87D/E486T/W727S, W24L/L62A/W727S/K830Q/A932S, W24L/E87D/Q135S/V522E/T670L/H711A/K830Q/S842G/R913V, W24L/S150T/V522E/R527N/W727S/H883R/G894Q, W24L/R527N/W727S/S842G/E871L/H883R/R913V/A932S, W24L/T670L/W727S/P750A/S842G/E871L, S28L, S28L/V50Q/E78P/E87D/Q135S/N266T/K267R/G437A/E486T/R527N/A551V/T670L/W727S/P750A/K830Q/S842G/E871L/H883R/G894Q/R913V/A932S, S28L/V50Q/V522E/R527N/H711A/W727S/E871L, S28L/L62A, S28L/L62A/K267R/A932S, S28L/G437A/R527N/E871L, S28L/V522E/R527N/T569L/H711A/K830Q/G894Q, S28L/W727S, S28L/W727S/E871L, T29L/Q39P/V50Q/L62A/G65R/E78P/E87D/Q135S/S150T/G437A/A551V/T569L/T670L/W727S/P750A/K830Q/S842G/H883R/G894Q/A932S, T29L/L62A/G437A/R527N, T29L/E78P/E87D/S150T/R527N/W727S, T29L/E78P/Q135S/W727S/K830Q, T29L/E87D, T29L/Q135S/S150T/R527N/T670L/W727S/H883R, T29L/S150T/K267R/W727S/P750A/E871L/H883R/A932S, T29L/S150T/G437A/W727S, T29L/V522E/T670L/H711A/E871L, T29L/T670L/A932S, Q39P/V50Q, Q39P/W727S/P750A/A932S, V50Q/Q135S/S150T/A932S, V50Q/G437A/V522E/R527N,

V50Q/H711A, V50Q/W727S/P750A/H883R/G894Q, L62A, L62A/E87D/S150T, L62A/E87D/S150T/W727S, L62A/Q135S/V522E/H711A/W727S/P750A/S842G/E871L/G894Q, L62A/G437A, L62A/G437A/W727S, E78P/E87D/E486T/R527N/T670L/W727S/P750A/K830Q/S842G/E871L/R913V/A932S, E87D/P750A, A89R/L109D/R527N/S678T/W727S/S842G, A89R/L109D/S678T/W727S/L736M/A812E/V878S, A89R/L109D/W727S/A932S, A89R/L109D/A932S, A89R/R527N, A89R/R527N/S678T/T692G/L736M/S842G/V878S/A932S, A89R/R527N/S678T/A932S, A89R/R527N/W727S/A812E/L860F, A89R/S678T/T692G/L736M/A932S, A89R/S678T/A812E/V878S, A89R/S842G/V878S, L109D/R527N/S678T/A812E, L109D/S678T/T692G/S842G/L860F/V878S/A932S, L109D/S678T/W727S/L860F, L109D/S678T/L736M/A812E/V878S, L109D/S678T/A812E, L109D/S678T/S842G/V878S, L109D/T692G/W727S/L736M/A812E, L109D/T692G/W727S/A812E/S842G/L860F, L109D/W727S/L860F/V878S, L109D/L736M/A932S, L109D/A812E, L109D/S842G, L109D/A932S, Q135S/T670L/W727S, Q135S/H711A/P750A/A932S, S150T/R527N/S842G/E871L/R913V, S150T/E871L/A932S, S150T/H883R/A932S, K267R/R527N/W727S, R403H/R527N/S678T/T692G/L736M/A812E/S842G/L860F, G437A/V522E/R527N/T670L/E871L, G437A/P750A/K830Q/A932S, V522E, V522E/R527N/T569L/W727S, V522E/K830Q, R527N, R527N/S678T/T692G/W727S/L736M/V878S, R527N/S678T/T692G/A812E/A932S, R527N/T692G/W727S/L736M/A812E, R527N/T692G/W727S/L736M/S842G/L860F/V878S, R527N/W727S/L736M, R527N/L736M/A932S, R527N/A812E, T670L/H711A/E871L, T670L/K830Q/E871L, S678T/T692G/W727S/A812E/S842G, S678T/T692G/A812E, S678T/A812E, S678T/L860F/V878S, S678T/R913V, S678T/A932S, T692G/W727S/L736M/S842G/R913V, T692G/A812E, W727S, W727S/A932S, E871L, and V878S/A932S, wherein the positions are numbered with reference to SEQ ID NO: 2880. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 62/87/150, 89/109/527/678/727/842, 89/109/678/727/736/812/878, 89/109/932, 89/527/678/692/736/842/878/932, 89/527/727/812/860, 89/678/692/736/932, 89/678/812/878, 109/527/678/812, 109/678/692/842/860/878/932, 109/678/736/812/878, 109/678/812, 109/692/727/736/812, 109/692/727/812/842/860, 109/736/932, 109/812, 109/842, 109/932, 403/527/678/692/736/812/842/860, 522/830, 527/678/692/727/736/878, 527/678/692/812/932, 527/692/727/736/812, 527/692/727/736/842/860/878, 527/727/736, 527/736/932, 527/812, 678/692/727/812/842, 678/692/812, 678/812, 692/727/736/842/913, and 692/812, wherein the positions are numbered with reference to SEQ ID NO: 2880. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 62A/87D/150T, 89R/109D/527N/678T/727S/842G, 89R/109D/678T/727S/736M/812E/878S, 89R/109D/932S, 89R/527N/678T/692G/736M/842G/878S/932S, 89R/527N/727S/812E/860F,

89R/678T/692G/736M/932S, 89R/678T/812E/878S, 109D/527N/678T/812E, 109D/678T/692G/842G/860F/878S/932S, 109D/678T/736M/812E/878S, 109D/678T/812E, 109D/692G/727S/736M/812E, 109D/692G/727S/812E/842G/860F, 109D/736M/932S, 109D/812E, 109D/842G, 109D/932S, 403H/527N/678T/692G/736M/812E/842G/860F, 522E/830Q, 527N/678T/692G/727S/736M/878S, 527N/678T/692G/812E/932S, 527N/692G/727S/736M/812E, 527N/692G/727S/736M/842G/860F/878S, 527N/727S/736M, 527N/736M/932S, 527N/812E, 678T/692G/727S/812E/842G, 678T/692G/812E, 678T/812E, 692G/727S/736M/842G/913V, and 692G/812E, wherein the positions are numbered with reference to SEQ ID NO: 2880. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from L62A/E87D/S150T, A89R/L109D/R527N/S678T/W727S/S842G, A89R/L109D/S678T/W727S/L736M/A812E/V878S, A89R/L109D/A932S, A89R/R527N/S678T/T692G/L736M/S842G/V878S/A932S, A89R/R527N/W727S/A812E/L860F, A89R/S678T/T692G/L736M/A932S, A89R/S678T/A812E/V878S, L109D/R527N/S678T/A812E, L109D/S678T/T692G/S842G/L860F/V878S/A932S, L109D/S678T/L736M/A812E/V878S, L109D/S678T/A812E, L109D/T692G/W727S/L736M/A812E, L109D/T692G/W727S/A812E/S842G/L860F, L109D/L736M/A932S, L109D/A812E, L109D/S842G, L109D/A932S, R403H/R527N/S678T/T692G/L736M/A812E/S842G/L860F, V522E/K830Q, R527N/S678T/T692G/W727S/L736M/V878S, R527N/S678T/T692G/A812E/A932S, R527N/T692G/W727S/L736M/A812E, R527N/T692G/W727S/L736M/S842G/L860F/V878S, R527N/W727S/L736M, R527N/L736M/A932S, R527N/A812E, S678T/T692G/W727S/A812E/S842G, S678T/T692G/A812E, S678T/A812E, T692G/W727S/L736M/S842G/R913V, and T692G/A812E, wherein the positions are numbered with reference to SEQ ID NO: 2880.

[0019] The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 3104, wherein the positions are numbered with reference to SEQ ID NO: 3104. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 62, 62/89/830, 62/248/678/830/878/932, 62/678, 62/678/785, 62/678/830, 62/678/830/860, 62/678/830/860/871/878/932, 62/678/830/860/878, 62/678/830/860/878/932, 62/678/830/860/932, 62/678/830/871, 62/678/830/871/932, 62/678/830/878/932, 62/678/830/932, 62/678/860, 62/678/860/878, 62/678/860/932, 62/678/871, 62/678/871/932, 62/678/878/932, 62/678/932, 62/830, 62/830/860, 62/830/860/871/873, 62/830/860/878/932, 62/830/860/932, 62/830/871/932, 62/830/878, 62/830/932, 62/833/860/932, 62/860, 62/860/871, 62/860/871/878, 62/860/871/932, 62/860/878/932, 62/860/932, 62/871/878/932, 62/871/932, 62/878, 62/878/932, 62/932, 678, 678/830/932, 678/932, and 860/932, wherein the

positions are numbered with reference to SEQ ID NO: 3104. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 62A, 62A/89D/830Q, 62A/248H/678T/830Q/878S/932S, 62A/678T, 62A/678T/785Q, 62A/678T/830Q, 62A/678T/830Q/860F, 62A/678T/830Q/860F/871L/878S/932S, 62A/678T/830Q/860F/878S, 62A/678T/830Q/860F/878S/932S, 62A/678T/830Q/860F/932S, 62A/678T/830Q/871L, 62A/678T/830Q/871L/932S, 62A/678T/830Q/878S/932S, 62A/678T/830Q/932S, 62A/678T/860F, 62A/678T/860F/878S, 62A/678T/860F/932S, 62A/678T/871L, 62A/678T/871L/932S, 62A/678T/878S/932S, 62A/678T/878S/932T, 62A/678T/932S, 62A/830Q, 62A/830Q/860F, 62A/830Q/860F/871L/873H, 62A/830Q/860F/878S/932S, 62A/830Q/860F/932S, 62A/830Q/871L/932S, 62A/830Q/878S, 62A/830Q/932S, 62A/833I/860F/932S, 62A/860F, 62A/860F/871L, 62A/860F/871L/878S, 62A/860F/871L/932S, 62A/860F/878S/932S, 62A/860F/932S, 62A/871L/878S/932S, 62A/871L/932S, 62A/878S, 62A/878S/932S, 62A/932S, 678T, 678T/830Q/932S, 678T/932S, and 860F/932S, wherein the positions are numbered with reference to SEQ ID NO: 3104. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from L62A, L62A/A89D/K830Q, L62A/Y248H/S678T/K830Q/V878S/A932S, L62A/S678T, L62A/S678T/P785Q, L62A/S678T/K830Q, L62A/S678T/K830Q/L860F, L62A/S678T/K830Q/L860F/E871L/V878S/A932S, L62A/S678T/K830Q/L860F/V878S, L62A/S678T/K830Q/L860F/V878S/A932S, L62A/S678T/K830Q/L860F/A932S, L62A/S678T/K830Q/E871L, L62A/S678T/K830Q/E871L/A932S, L62A/S678T/K830Q/V878S/A932S, L62A/S678T/K830Q/A932S, L62A/S678T/L860F, L62A/S678T/L860F/V878S, L62A/S678T/L860F/A932S, L62A/S678T/E871L, L62A/S678T/E871L/A932S, L62A/S678T/V878S/A932S, L62A/S678T/V878S/A932T, L62A/S678T/A932S, L62A/K830Q, L62A/K830Q/L860F, L62A/K830Q/L860F/E871L/R873H, L62A/K830Q/L860F/V878S/A932S, L62A/K830Q/L860F/A932S, L62A/K830Q/E871L/A932S, L62A/K830Q/V878S, L62A/K830Q/A932S, L62A/M833I/L860F/A932S, L62A/L860F, L62A/L860F/E871L, L62A/L860F/E871L/V878S, L62A/L860F/E871L/A932S, L62A/L860F/V878S/A932S, L62A/L860F/A932S, L62A/E871L/V878S/A932S, L62A/E871L/A932S, L62A/V878S, L62A/V878S/A932S, L62A/A932S, S678T, S678T/K830Q/A932S, S678T/A932S, and L860F/A932S, wherein the positions are numbered with reference to SEQ ID NO: 3104.

[0020] In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 62/89/830, 62/830, 62/830/860, 62/830/860/932, and 62/830/932, wherein the positions are numbered with reference to SEQ ID NO: 3104. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 62A/89D/830Q, 62A/830Q, 62A/830Q/860F, 62A/830Q/860F/932S, and 62A/830Q/932S, wherein the positions are numbered with reference to SEQ ID NO: 3104. In some

embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from L62A/A89D/K830Q, L62A/K830Q, L62A/K830Q/L860F, L62A/K830Q/L860F/A932S, and L62A/K830Q/A932S, wherein the positions are numbered with reference to SEQ ID NO: 3104.

[0021] In some embodiments, the recombinant acid alpha glucosidase comprises at least one mutation in at least one position as provided in Tables 3-1, 3-2, 4-1, 6-1, 10-1, 10-2, 12-1, 13-1, 13-2, 14-1, 14-2, 15-1, 16-1, 17-1, 17-2, 17-3, 17-4, 17-5, 17-6, 17-7, 17-8, and/or 17-9. In some additional embodiments, the recombinant acid alpha glucosidase is derived from a human acid alpha glucosidase. In yet some additional embodiments, the recombinant acid alpha glucosidase comprising the polypeptide sequence of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

[0022] In some additional embodiments, the recombinant acid alpha-glucosidase provided herein is thermostable. In some further embodiments, the recombinant acid alpha-glucosidase is resistant to proteolysis. In yet some additional embodiments, the recombinant acid alpha-glucosidase is resistant to at least one digestive tract protease. In some embodiments, the digestive tract protease is selected from chymotrypsin, trypsin, carboxypeptidases, and elastases. In some further embodiments, the recombinant acid alpha-glucosidase is acid stable. In some additional embodiments, the recombinant acid alpha-glucosidase is stable to acidic pH and neutral pH. In yet some additional embodiments, the recombinant acid alpha-glucosidase is purified. In some further embodiments, the recombinant acid alpha-glucosidase exhibits at least one improved property selected from: i) enhanced catalytic activity; ii) increased expression; iii) increased stability at neutral pH levels; iv) increased stability at acidic pH levels; v) enhanced activity in cell lysates; and vi) decreased immunogenicity; or a combination of any of i), ii), iii), iv), v), and/or vi), as compared to a reference sequence. In some further embodiments, the recombinant acid alpha glucosidase exhibits at least one improved property selected from: i) enhanced catalytic activity; ii) increased tolerance to pH 7; iii) increased tolerance to pH 4; iv) increased expression; v) increased uptake into cells; vi) increased enzymatic activity in cell lysates; vii) decreased immunogenicity, or a combination of any of i), ii), iii), iv), v), vi), and/or vii), as compared to a reference sequence. In some embodiments, the reference sequence is selected from SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104. In some further embodiments, the recombinant acid alpha glucosidase is more stable at pH 7 than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104. In yet some other embodiments, the recombinant acid alpha glucosidase is more stable at pH 4 than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104. In some further embodiments, the recombinant acid alpha glucosidase exhibits increased expression than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104. In still some additional embodiments, the recombinant acid alpha glucosidase is more lysosomally stable than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946,

1956, 2496, 2880, and/or 3104. In some further embodiments, the recombinant acid alpha glucosidase is more readily taken up by cells than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104. In some additional embodiments, the recombinant acid alpha glucosidase exhibits greater enzymatic activity in cell lysates than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104. In some additional embodiments, the recombinant acid alpha glucosidase exhibits reduced or decreased immunogenicity than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104. In some embodiments, the recombinant acid alpha glucosidase is purified.

[0023] In some further embodiments, the recombinant acid alpha-glucosidase comprises a polypeptide sequence comprising at least about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to at least one of the even-numbered sequences of SEQ ID NOS: 8-3378. In some further embodiments, the recombinant acid alpha-glucosidase comprises a polypeptide sequence comprising at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to at least one of the even-numbered sequences of SEQ ID NOS: 8-3378. In some embodiments the recombinant acid alpha glucosidase comprises a polypeptide sequence at least 90% identical to at least one of the even-numbered sequences of SEQ ID NOS: 8-3378. In some further embodiments, the recombinant acid alpha-glucosidase comprises a polypeptide sequence comprises at least one of the even-numbered sequences of SEQ ID NOS: 2-3378. In some further embodiments, the recombinant acid alpha-glucosidase consists of a polypeptide sequence comprising at least one of the even-numbered sequences of SEQ ID NOS: 8-3378.

[0024] The present invention also provides compositions comprising at least one recombinant acid alpha-glucosidase provided herein. In some embodiments, the compositions comprise one recombinant acid alpha-glucosidase provided herein.

[0025] The present invention also provides recombinant polynucleotide sequences encoding at least one recombinant acid alpha-glucosidase provided herein. In some embodiment, a recombinant polynucleotide sequence encodes one recombinant acid alpha-glucosidase. In some embodiments, the recombinant polynucleotide sequence is selected from DNA, RNA, and mRNA. In some embodiments, the polynucleotide sequence is codon-optimized. In some further embodiments, the recombinant polynucleotide sequence encodes a recombinant acid alpha-glucosidase comprising a polypeptide sequence comprising at least about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to at least one of the even-numbered sequences of SEQ ID NOS: 8-3378. In some further embodiments, the recombinant acid alpha-glucosidase encoded by the polynucleotide sequence comprises a polypeptide sequence comprising at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to at least one of the even-numbered sequences of SEQ ID NOS: 8-3378.

[0026] In some further embodiments, the recombinant polynucleotide sequence is at least about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to at least one of the odd-numbered sequences of SEQ ID NOS: 7-3377. In some further embodiments, the recombinant polynucleotide sequence is at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%, identical to at least one of the odd-numbered sequences of SEQ ID NOS: 7-3377. In some additional embodiments, the recombinant polynucleotide sequence comprises a sequence comprising at least 90% sequence identity to at least one of the odd-numbered sequences of SEQ ID NOS: 7-3377. In yet some further embodiments, the recombinant polynucleotide sequence comprises an odd-numbered sequence of SEQ ID NOS: 7-3377.

[0027] The present invention also provides expression vectors comprising the recombinant polynucleotide sequence encoding recombinant acid alpha-glucosidase. In some embodiments, the recombinant polynucleotide sequence is operably linked to a control sequence. In some further embodiments, the control sequence is a promoter. In some additional embodiments, the promoter is a heterologous promoter. The present invention also provides the expression vector referred to herein as pDH. In some embodiments, the pDH vector comprises at least one polynucleotide sequence encoding an acid alpha glucosidase. In some additional embodiments, the pDH vector comprises at least one polynucleotide sequence encoding an acid alpha glucosidase provided herein. In some further embodiments, the pDH vector comprises at least one polynucleotide sequence selected from the odd-numbered sequences of SEQ ID NOS: 1-3377. In some additional embodiments, the pDH vector comprises at least one polynucleotide sequence encoding an acid alpha glucosidase selected from the even-numbered sequences of SEQ ID NOS: 2-3378. In some embodiments, the pDH vector comprises SEQ ID NO: 3379, while in some other embodiments, the pDH vector comprises SEQ ID NO: 3380. In some additional embodiments, SEQ ID NO:1, which is included in SEQ ID NO: 3379 is replaced with another polynucleotide sequence. In some embodiments, the SEQ ID NO: 1 in SEQ ID NO: 3379 is replaced by at least one polynucleotide sequence selected from the odd-numbered sequences of SEQ ID NOS: 1-3377. In some additional embodiments, the pDH vector comprises the plasmid provided in the map at Figure 9, while in some other embodiments, the pDH vector comprises the plasmid the map at Figure 10. In some embodiments, the “stuffer sequence” (i.e., the bla sequence) in the plasmid shown in Figure 10 is replaced by a gene of interest. In some embodiments, the “stuffer gene” ATG start codon to the last codon of the gene is numbered from base pair 724 to 1581 (for a total of 858 base pairs). As used herein, the terms “stuffer gene” and “stuffer sequence” refer to a sequence within a plasmid vector that is replaced by a gene of interest. As used herein, the term “gene of interest” refers to a gene that encodes a desired polypeptide (e.g., a “polypeptide of interest”). In some additional embodiments, the stuffer sequence is replaced by a gene of interest (i.e., a gene that is desired for expression, such that a polypeptide of interest, such as a variant acid alpha glucosidase is produced).

[0028] The present invention also provides host cells comprising at least one expression vector provided herein. In some embodiments, the expression vector provided in the host cells is pDH. In some embodiments, the host cell is selected from eukaryotes and prokaryotes. In some further embodiments, the host cell is a mammalian cell.

[0029] The present invention also provides methods of producing a recombinant acid alpha glucosidase variant, comprising culturing at least one host cell provided herein, under conditions that the acid alpha glucosidase encoded by the recombinant polynucleotide is produced. In some embodiments, the methods further comprise the step of recovering the acid alpha glucosidase. In yet some further embodiments, the methods further comprise the step of purifying the acid alpha glucosidase. The present invention also provides recombinant acid alpha glucosidase variants produced according to a method provided herein.

[0030] The present invention also provides compositions comprising at least one recombinant acid alpha glucosidase provided herein. The present invention also provides for use of the compositions provided herein. In some embodiments, the present invention provides pharmaceutical compositions for the treatment of Pompe disease, comprising at least one composition provided herein. In some additional embodiments, the pharmaceutical composition further comprises at least one pharmaceutically acceptable carrier and/or excipient. In some further embodiments, the pharmaceutical composition is suitable for parenteral injection or infusion to a human. In yet some additional embodiments, the present invention provides pharmaceutical compositions comprising at least one recombinant polynucleotide provided herein. In yet some further embodiments, the present invention provides pharmaceutical compositions comprising at least one recombinant polypeptide provided herein. In yet some additional embodiments, the present invention provides compositions comprising at least one recombinant polynucleotide and at least one recombinant polypeptide provided herein.

[0031] The present invention also provides methods for treating and/or preventing the symptoms of Pompe disease in a subject, comprising providing a subject having Pompe disease and at least one pharmaceutical composition provided herein, and administering the pharmaceutical composition to the subject. In some embodiments, the symptoms of Pompe disease are ameliorated. In some additional embodiments, the subject is an infant or child. In yet some further embodiments, the subject is an adult or young adult.

DESCRIPTION OF THE DRAWINGS

[0032] Figure 1 provides a graph showing the kinetic assay results for three GAA variants, as described in Example 9.

[0033] Figure 2 provides a graph showing the glycogen hydrolysis to glucose activity of four GAA variants, as described in Example 9.

[0034] Figure 3 provides a graph showing the 4-MuGlu hydrolysis levels of four GAA variants, as described in Example 9.

[0035] Figure 4 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 5 hours incubation at 37 °C with cultured Pompe patient fibroblasts, followed by 24 hr incubation.

[0036] Figure 5 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 24 hours incubation at 37 °C with cultured Pompe patient fibroblasts.

[0037] Figure 6 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 5 hours incubation at 37 °C with cultured C2C12 GAA-/- myoblasts, followed by 24 hr incubation.

[0038] Figure 7 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 24 hours incubation at 37 °C with cultured C2C12 GAA-/- myoblasts.

[0039] Figure 8 provides a graph showing the cellular uptake of different purified GAA variants, expressed as relative activity, after 1-7 days incubation at 37 °C with cultured Pompe patient fibroblasts.

[0040] Figure 9 provides a plasmid map of the pDH vector containing WT GAA.

[0041] Figure 10 provides a plasmid map showing the pDH vector comprising the stuffer sequence bla (i.e., beta-lactamase).

[0042] Figure 11 provides a graph showing the stability at neutral pH and 37 °C, expressed as normalized residual RFU activity, of eight GAA variants.

[0043] Figure 12 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 24 hours incubation at 37 °C with cultured Pompe patient fibroblasts.

[0044] Figure 13 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 72 hours incubation at 37 °C with cultured Pompe patient fibroblasts.

[0045] Figure 14 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 24 hours incubation at 37 °C with cultured C2C12 GAA-/- myoblasts.

[0046] Figure 15 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 72 hours incubation at 37 °C with cultured C2C12 GAA-/- myoblasts.

[0047] Figure 16 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 4 hours incubation at 37 °C with cultured Pompe patient fibroblasts, followed by cell washing and an additional 20 hr incubation.

[0048] Figure 17 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 24 hours incubation at 37 °C with cultured Pompe patient fibroblasts.

[0049] Figure 18 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 72 hours incubation at 37 °C with cultured Pompe patient fibroblasts.

[0050] Figure 19 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 4 hours incubation at 37 °C with cultured C2C12 GAA-/- myoblasts, followed by cell washing and an additional 20 hr incubation.

[0051] Figure 20 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 24 hours incubation at 37 °C with cultured C2C12 GAA-/- myoblasts.

[0052] Figure 21 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 72 hours incubation at 37 °C with cultured C2C12 GAA-/- myoblasts.

[0053] Figure 22 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 96 hours incubation at 37 °C with cultured Pompe patient fibroblasts.

[0054] Figure 23 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 96 hours incubation at 37 °C with cultured C2C12 GAA-/- myoblasts.

[0055] Figure 24 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 48 hours incubation at 37 °C with cultured Pompe patient fibroblasts.

[0056] Figure 25 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 48 hours incubation at 37 °C with cultured C2C12 GAA-/- myoblasts.

[0057] Figure 26 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 1 hour incubation at 37 °C with cultured Pompe patient fibroblasts, followed by cell washing and an additional 71 hr incubation.

[0058] Figure 27 provides a graph showing the cellular uptake of different purified GAA variants, expressed as RFU activity, after 1 hour incubation at 37 °C with cultured C2C12 GAA-/- myoblasts, followed by cell washing and an additional 71 hr incubation.

DESCRIPTION OF THE INVENTION

[0059] The present invention provides engineered acid alpha-glucosidase (GAA) polypeptides and compositions thereof. In some embodiments, the engineered GAA polypeptides have been optimized to provide enhanced catalytic activity and enhanced acid stability, while reducing sensitivity to proteolysis. The invention also provides methods for utilization of the compositions comprising the engineered GAA polypeptides for therapeutic and other purposes.

Abbreviations and Definitions:

[0060] Unless defined otherwise, all technical and scientific terms used herein generally have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. Generally, the nomenclature used herein and the laboratory procedures of cell culture, molecular genetics, microbiology, organic chemistry, analytical chemistry and nucleic acid chemistry described below are those well-known and commonly employed in the art. Such techniques are well-known and described in numerous texts and reference works well known to those of skill in the art. Standard techniques, or modifications thereof, are used for chemical syntheses and chemical analyses. All patents, patent applications, articles and publications mentioned herein, both *supra* and *infra*, are hereby expressly incorporated herein by reference.

[0061] Although any suitable methods and materials similar or equivalent to those described herein find use in the practice of the present invention, some methods and materials are described herein. It is to be understood that this invention is not limited to the particular methodology, protocols, and reagents described, as these may vary, depending upon the context they are used by those of skill in the art. Accordingly, the terms defined immediately below are more fully described by reference to the application as a whole. All patents, patent applications, articles and publications mentioned herein, both *supra* and *infra*, are hereby expressly incorporated herein by reference.

[0062] Also, as used herein, the singular "a", "an," and "the" include the plural references, unless the context clearly indicates otherwise.

[0063] Numeric ranges are inclusive of the numbers defining the range. Thus, every numerical range disclosed herein is intended to encompass every narrower numerical range that falls within such broader numerical range, as if such narrower numerical ranges were all expressly written herein. It is also intended that every maximum (or minimum) numerical limitation disclosed herein includes every lower (or higher) numerical limitation, as if such lower (or higher) numerical limitations were expressly written herein.

[0064] The term "about" means an acceptable error for a particular value. In some instances "about" means within 0.05%, 0.5%, 1.0%, or 2.0%, of a given value range. In some instances, "about" means within 1, 2, 3, or 4 standard deviations of a given value. In some instances, "about" encompasses values that are within 2.5%, 3%, 3.5%, 4%, 4.5%, 5%, 5.5%, 6%, 6.5%, 7%, 7.5%, 8%, 8.5%, 9%, 9.5%, or 10% of a given value.

[0065] Furthermore, the headings provided herein are not limitations of the various aspects or embodiments of the invention which can be had by reference to the application as a whole. Accordingly, the terms defined immediately below are more fully defined by reference to the application as a whole. Nonetheless, in order to facilitate understanding of the invention, a number of terms are defined below.

[0066] Unless otherwise indicated, nucleic acids are written left to right in 5' to 3' orientation; amino acid sequences are written left to right in amino to carboxy orientation, respectively.

[0067] As used herein, the term “comprising” and its cognates are used in their inclusive sense (i.e., equivalent to the term “including” and its corresponding cognates).

[0068] “EC” number refers to the Enzyme Nomenclature of the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology (NC-IUBMB). The IUBMB biochemical classification is a numerical classification system for enzymes based on the chemical reactions they catalyze.

[0069] “ATCC” refers to the American Type Culture Collection whose biorepository collection includes genes and strains.

[0070] “NCBI” refers to National Center for Biological Information and the sequence databases provided therein.

[0071] As used herein, the terms “acid alpha-glucosidase,” “acid α -glucosidase,” “acid alpha-glucosidase polypeptide” “lysosomal alpha-glucosidase,” and “GAA” refer to enzymes within a family (EC 3.2.1.20) of enzymes that break down glycogen present in lysosomes. The enzyme is also sometimes referred to as “alpha-1,4-glucosidase,” α -1,4-glucosidase,” “acid maltase,” “glucoinvertase,” “glucosidosucrase,” “lysosomal alpha-glucosidase,” “lysosomal α -glucosidase,” “maltase,” or “maltase-glucoamylase.” One reaction catalyzed by the enzyme is the hydrolysis of terminal, non-reducing (1 to 4) linked alpha-D-glucose residues with release of alpha-D-glucose.

[0072] As used herein, “Pompe disease” refers to a glycogen storage disease type II, which is an autosomal recessive genetic disorder that results in a metabolic disorder characterized by lysosomal accumulation of glycogen in skeletal muscle and other tissues. It is characterized based on age of onset, organ involvement, severity, and rate of progression. The more severe form is infantile-onset Pompe disease (IOPD), which occurs in infants. The other form, referred to as “late-onset Pompe disease” (LOPD), occurs in individuals with an onset of disease before 12 months of age, but without the cardiomyopathy associated with IOPD, and all individuals with an onset of disease after 12 months of age. Synonyms for Pompe disease include “acid alpha-glucosidase deficiency,” acid maltase deficiency,” “GAA deficiency,” “glycogen storage disease type II,” “GSD II,” “GSD2,” and “glycogenosis type II.”

[0073] “Protein,” “polypeptide,” and “peptide” are used interchangeably herein to denote a polymer of at least two amino acids covalently linked by an amide bond, regardless of length or post-translational modification (e.g., glycosylation or phosphorylation).

[0074] “Amino acids” are referred to herein by either their commonly known three-letter symbols or by the one-letter symbols recommended by IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single letter codes.

[0075] The terms “engineered,” “recombinant,” “non-naturally occurring,” and “variant,” when used with reference to a cell, a polynucleotide or a polypeptide refers to a material or a material corresponding to the natural or native form of the material that has been modified in a manner that

would not otherwise exist in nature or is identical thereto but produced or derived from synthetic materials and/or by manipulation using recombinant techniques.

[0076] As used herein, “wild-type” and “naturally-occurring” refer to the form found in nature. For example, a wild-type polypeptide or polynucleotide sequence is a sequence present in an organism that can be isolated from a source in nature and which has not been intentionally modified by human manipulation.

[0077] “Coding sequence” refers to that part of a nucleic acid (e.g., a gene) that encodes an amino acid sequence of a protein.

[0078] The term “percent (%) sequence identity” is used herein to refer to comparisons among polynucleotides and polypeptides, and are determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence for optimal alignment of the two sequences. The percentage may be calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Alternatively, the percentage may be calculated by determining the number of positions at which either the identical nucleic acid base or amino acid residue occurs in both sequences or a nucleic acid base or amino acid residue is aligned with a gap to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Those of skill in the art appreciate that there are many established algorithms available to align two sequences. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith and Waterman (Smith and Waterman, *Adv. Appl. Math.*, 2:482 [1981]), by the homology alignment algorithm of Needleman and Wunsch (Needleman and Wunsch, *J. Mol. Biol.*, 48:443 [1970]), by the search for similarity method of Pearson and Lipman (Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444 [1988]), by computerized implementations of these algorithms (e.g., GAP, BESTFIT, FASTA, and TFASTA in the GCG Wisconsin Software Package), or by visual inspection, as known in the art. Examples of algorithms that are suitable for determining percent sequence identity and sequence similarity include, but are not limited to the BLAST and BLAST 2.0 algorithms, which are described by Altschul et al. (See, Altschul et al., *J. Mol. Biol.*, 215: 403-410 [1990]; and Altschul et al., 1977, *Nucleic Acids Res.*, 3389-3402 [1977], respectively). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information website. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as, the neighborhood word score threshold (See, Altschul et

al., supra). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (See, Henikoff and Henikoff, Proc. Natl. Acad. Sci. USA 89:10915 [1989]). Exemplary determination of sequence alignment and % sequence identity can employ the BESTFIT or GAP programs in the GCG Wisconsin Software package (Accelrys, Madison WI), using default parameters provided.

[0079] As used herein, the term “reference sequence” refers to a defined sequence used as a basis for a sequence comparison. A reference sequence may be a subset of a larger sequence, for example, a segment of a full-length gene or polypeptide sequence. Generally, a reference sequence is at least 20 nucleotide or amino acid residues in length, at least 25 residues in length, at least 50 residues in length, at least 100 residues in length or the full length of the nucleic acid or polypeptide. Since two polynucleotides or polypeptides may each (1) comprise a sequence (i.e., a portion of the complete sequence) that is similar between the two sequences, and (2) may further comprise a sequence that is divergent between the two sequences, sequence comparisons between two (or more) polynucleotides or polypeptide are typically performed by comparing sequences of the two polynucleotides or polypeptides over a “comparison window” to identify and compare local regions of sequence similarity. In some embodiments, a “reference sequence” can be based on a primary amino acid sequence, where the reference sequence is a sequence that can have one or more changes in the primary sequence. “Comparison window” refers to a conceptual segment of at least about 20 contiguous nucleotide positions or amino acids residues wherein a sequence may be compared to a reference sequence of at least 20 contiguous nucleotides or amino acids and wherein the portion of the sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The comparison window can be longer than 20 contiguous residues, and includes, optionally 30, 40, 50, 100, or longer windows.

[0080] “Corresponding to”, “reference to” or “relative to” when used in the context of the numbering of a given amino acid or polynucleotide sequence refers to the numbering of the residues of a

specified reference sequence when the given amino acid or polynucleotide sequence is compared to the reference sequence. In other words, the residue number or residue position of a given polymer is designated with respect to the reference sequence rather than by the actual numerical position of the residue within the given amino acid or polynucleotide sequence. For example, a given amino acid sequence, such as that of an engineered GAA, can be aligned to a reference sequence by introducing gaps to optimize residue matches between the two sequences. In these cases, although the gaps are present, the numbering of the residue in the given amino acid or polynucleotide sequence is made with respect to the reference sequence to which it has been aligned.

[0081] “Amino acid difference” or “residue difference” refers to a difference in the amino acid residue at a position of a polypeptide sequence relative to the amino acid residue at a corresponding position in a reference sequence. The positions of amino acid differences generally are referred to herein as “X_n,” where n refers to the corresponding position in the reference sequence upon which the residue difference is based. For example, a “residue difference at position X₂₇ as compared to SEQ ID NO: 2” refers to a difference of the amino acid residue at the polypeptide position corresponding to position 27 of SEQ ID NO: 2. Thus, if the reference polypeptide of SEQ ID NO: 2 has a phenylalanine at position 2, then a “residue difference at position X₂₇ as compared to SEQ ID NO: 2” is an amino acid substitution of any residue other than phenylalanine at the position of the polypeptide corresponding to position 27 of SEQ ID NO: 2. In most instances herein, the specific amino acid residue difference at a position is indicated as “X_nY” where “X_n” specified the corresponding position as described above, and “Y” is the single letter identifier of the amino acid found in the engineered polypeptide (i.e., the different residue than in the reference polypeptide). In some instances (e.g., as shown in in Tables 3-1, 3-2, 4-1, 6-1, 10-1, 10-2, 12-1, 13-1, 13-2, 14-1, 14-2, 15-1, 16-1, 17-1, 17-2, 17-3, 17-4, 17-5, 17-6, 17-7, 17-8, and/or 17-9), the present disclosure also provides specific amino acid differences denoted by the conventional notation “A_nB”, where A is the single letter identifier of the residue in the reference sequence, “n” is the number of the residue position in the reference sequence, and B is the single letter identifier of the residue substitution in the sequence of the engineered polypeptide. In some instances, a polypeptide of the present disclosure can include one or more amino acid residue differences relative to a reference sequence, which is indicated by a list of the specified positions where residue differences are present relative to the reference sequence. In some embodiments, where more than one amino acid can be used in a specific residue position of a polypeptide, the various amino acid residues that can be used are separated by a “/” (e.g., X₂₇P/X₂₇R or X₂₇P/R). In some embodiments, the enzyme variants comprise more than one substitution. These substitutions are separated by a slash for ease in reading (e.g., F₂₇P/C₉₄₄W). In some cases, the “X” does not precede the position number in the present application. The present application includes engineered polypeptide sequences comprising one or more amino acid differences that include either/or both conservative and non-conservative amino acid substitutions.

[0082] “Conservative amino acid substitution” refers to a substitution of a residue with a different residue having a similar side chain, and thus typically involves substitution of the amino acid in the polypeptide with amino acids within the same or similar defined class of amino acids. By way of example and not limitation, an amino acid with an aliphatic side chain may be substituted with another aliphatic amino acid (e.g., alanine, valine, leucine, and isoleucine); an amino acid with hydroxyl side chain is substituted with another amino acid with a hydroxyl side chain (e.g., serine and threonine); an amino acids having aromatic side chains is substituted with another amino acid having an aromatic side chain (e.g., phenylalanine, tyrosine, tryptophan, and histidine); an amino acid with a basic side chain is substituted with another amino acid with a basis side chain (e.g., lysine and arginine); an amino acid with an acidic side chain is substituted with another amino acid with an acidic side chain (e.g., aspartic acid or glutamic acid); and/or a hydrophobic or hydrophilic amino acid is replaced with another hydrophobic or hydrophilic amino acid, respectively.

[0083] “Non-conservative substitution” refers to substitution of an amino acid in the polypeptide with an amino acid with significantly differing side chain properties. Non-conservative substitutions may use amino acids between, rather than within, the defined groups and affects (a) the structure of the peptide backbone in the area of the substitution (e.g., proline for glycine) (b) the charge or hydrophobicity, or (c) the bulk of the side chain. By way of example and not limitation, an exemplary non-conservative substitution can be an acidic amino acid substituted with a basic or aliphatic amino acid; an aromatic amino acid substituted with a small amino acid; and a hydrophilic amino acid substituted with a hydrophobic amino acid.

[0084] “Deletion” refers to modification to the polypeptide by removal of one or more amino acids from the reference polypeptide. Deletions can comprise removal of 1 or more amino acids, 2 or more amino acids, 5 or more amino acids, 10 or more amino acids, 15 or more amino acids, or 20 or more amino acids, up to 10% of the total number of amino acids, or up to 20% of the total number of amino acids making up the reference enzyme while retaining enzymatic activity and/or retaining the improved properties of an engineered enzyme. Deletions can be directed to the internal portions and/or terminal portions of the polypeptide. In various embodiments, the deletion can comprise a continuous segment or can be discontinuous.

[0085] “Insertion” refers to modification to the polypeptide by addition of one or more amino acids from the reference polypeptide. Insertions can be in the internal portions of the polypeptide, or to the carboxy or amino terminus. Insertions as used herein include fusion proteins as is known in the art. The insertion can be a contiguous segment of amino acids or separated by one or more of the amino acids in the naturally occurring polypeptide.

[0086] A “functional fragment” or a “biologically active fragment” used interchangeably herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion(s) and/or internal deletions, but where the remaining amino acid sequence is identical to the corresponding positions in

the sequence to which it is being compared (e.g., a full-length engineered GAA of the present invention) and that retains substantially all of the activity of the full-length polypeptide.

[0087] “Isolated polypeptide” refers to a polypeptide which is substantially separated from other contaminants that naturally accompany it (e.g., protein, lipids, and polynucleotides). The term embraces polypeptides which have been removed or purified from their naturally-occurring environment or expression system (e.g., host cell or *in vitro* synthesis). The recombinant GAA polypeptides may be present within a cell, present in the cellular medium, or prepared in various forms, such as lysates or isolated preparations. As such, in some embodiments, the recombinant GAA polypeptides can be an isolated polypeptide.

[0088] “Substantially pure polypeptide” refers to a composition in which the polypeptide species is the predominant species present (i.e., on a molar or weight basis it is more abundant than any other individual macromolecular species in the composition), and is generally a substantially purified composition when the object species comprises at least about 50 percent of the macromolecular species present by mole or % weight. Generally, a substantially pure GAA composition comprises about 60% or more, about 70% or more, about 80% or more, about 90% or more, about 95% or more, and about 98% or more of all macromolecular species by mole or % weight present in the composition. In some embodiments, the object species is purified to essential homogeneity (i.e., contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species. Solvent species, small molecules (<500 Daltons), and elemental ion species are not considered macromolecular species. In some embodiments, the isolated recombinant GAA polypeptides are substantially pure polypeptide compositions.

[0089] “Improved enzyme property” refers to an engineered GAA polypeptide that exhibits an improvement in any enzyme property as compared to a reference GAA polypeptide and/or as a wild-type GAA polypeptide or another engineered GAA polypeptide. Improved properties include but are not limited to such properties as increased protein expression, increased thermoactivity, increased thermostability, increased pH activity, increased stability, increased enzymatic activity, increased substrate specificity or affinity, increased specific activity, increased resistance to substrate or end-product inhibition, increased chemical stability, improved chemoselectivity, improved solvent stability, increased tolerance to acidic, neutral, or basic pH, increased tolerance to proteolytic activity (i.e., reduced sensitivity to proteolysis), reduced aggregation, increased solubility, reduced immunogenicity, improved post-translational modification (e.g., glycosylation), altered temperature profile, increased lysosomal stability, etc.

[0090] “Increased enzymatic activity” or “enhanced catalytic activity” refers to an improved property of the engineered GAA polypeptides, which can be represented by an increase in specific activity (e.g., product produced/time/weight protein) or an increase in percent conversion of the substrate to the product (e.g., percent conversion of starting amount of substrate to product in a specified time

period using a specified amount of GAA) as compared to the reference GAA enzyme. Exemplary methods to determine enzyme activity are provided in the Examples. Any property relating to enzyme activity may be affected, including the classical enzyme properties of K_m , V_{max} or k_{cat} , changes of which can lead to increased enzymatic activity. Improvements in enzyme activity can be from about 1.1 fold the enzymatic activity of the corresponding wild-type enzyme, to as much as 2-fold, 5-fold, 10-fold, 20-fold, 25-fold, 50-fold, 75-fold, 100-fold, 150-fold, 200-fold or more enzymatic activity than the naturally occurring GAA or another engineered GAA from which the GAA polypeptides were derived.

[0091] In some embodiments, the engineered GAA polypeptides have a k_{cat} of at least 0.1/sec, at least 0.5/sec, at least 1.0/sec, at least 5.0/sec, at least 10.0/sec and in some preferred embodiments greater than 10.0/sec. In some embodiments, the K_m is in the range of about 1 μ M to about 5mM; in the range of about 5 μ M to about 10mM; in the range of about 30 μ M to about 30mM; or in the range of about 50 μ M to about 50mM. In some specific embodiments, the engineered GAA enzyme exhibits improved enzymatic activity after exposure to certain conditions in the range of 1.5 to 10 fold, 1.5 to 25 fold, 1.5 to 50 fold, 1.5 to 100 fold or greater than that of a reference GAA enzyme (e.g., a wild-type GAA or any other reference GAA, such as SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104).

[0092] GAA activity can be measured by any suitable method known in the art (e.g., standard assays, such as monitoring changes in spectrophotometric properties of reactants or products). In some embodiments, the amount of products produced can be measured by High-Performance Liquid Chromatography (HPLC) separation combined with UV absorbance or fluorescent detection. In some embodiments, the amount of product produced can be measured by monitoring fluorescence (Ex. 355 nm, Em. 460 nm) after hydrolysis of a 4-methylumbelliferyl-alpha-D-glucopyranoside (4-MUGlu) molecule. Comparisons of enzyme activities are made using a defined preparation of enzyme, a defined assay under a set condition, and one or more defined substrates, as further described in detail herein. Generally, when lysates are compared, the numbers of cells and the amount of protein assayed are determined as well as use of identical expression systems and identical host cells to minimize variations in amount of enzyme produced by the host cells and present in the lysates.

[0093] The term “improved tolerance to acidic pH” means that a recombinant GAA according to the invention will have increased stability (higher retained activity at about pH 4.8, after exposure to an acidic pH for a specified period of time (e.g., 1 hour, up to 24 hours)) as compared to a reference GAA or another enzyme.

[0094] The term “improved tolerance to neutral pH” means that a recombinant GAA according to the invention will have increased stability (higher retained activity at about pH 7, after exposure to a neutral pH for a specified period of time (e.g., 1 hour, up to 24 hours)) as compared to a reference GAA or another enzyme.

[0095] The term “improved cellular uptake” means that a recombinant GAA provided herein exhibits increased endocytosis into cells, as compared to a reference GAA (including wild-type GAA) or another enzyme. In some embodiments, the cells are cultured Pompe patient cells (higher retained intracellular activity after incubation with cultured cells over a specified period of time, as compared to a reference GAA or another enzyme). In some additional embodiments, the recombinant GAA provided herein exhibits greater retained intracellular activity with cultured cells over a specific period of time as compared to a reference GAA (including wild-type GAA) or another enzyme. In some additional embodiments, the time period is about 4 hours, while in some other embodiments, the time period is less than 4 hours (e.g., 1, 2, or 3 hours), and in some alternative embodiments, the time period is more than 4 hours (e.g., 5, 6, 7, 8, or more hours).

[0096] The terms “reduced immunogenicity” and “decreased immunogenicity” mean that a recombinant GAA provided herein induces a reduced immune response as compared to a wild-type or another reference GAA.

[0097] “Physiological pH” as used herein means the pH range generally found in a subject’s (e.g., human) blood.

[0098] The term “basic pH” (e.g., used with reference to improved stability to basic pH conditions or increased tolerance to basic pH) means a pH range of about 7 to 11.

[0099] The term “acidic pH” (e.g., used with reference to improved stability to acidic pH conditions or increased tolerance to acidic pH) means a pH range of about 1.5 to 4.5.

[0100] “Conversion” refers to the enzymatic conversion (or biotransformation) of a substrate(s) to the corresponding product(s). “Percent conversion” refers to the percent of the substrate that is converted to the product within a period of time under specified conditions. Thus, the “enzymatic activity” or “activity” of a GAA polypeptide can be expressed as “percent conversion” of the substrate to the product in a specific period of time.

[0101] “Hybridization stringency” relates to hybridization conditions, such as washing conditions, in the hybridization of nucleic acids. Generally, hybridization reactions are performed under conditions of lower stringency, followed by washes of varying but higher stringency. The term “moderately stringent hybridization” refers to conditions that permit target-DNA to bind a complementary nucleic acid that has about 60% identity, preferably about 75% identity, about 85% identity to the target DNA, with greater than about 90% identity to target-polynucleotide. Exemplary moderately stringent conditions are conditions equivalent to hybridization in 50% formamide, 5× Denhart’s solution, 5×SSPE, 0.2% SDS at 42°C, followed by washing in 0.2×SSPE, 0.2% SDS, at 42°C. “High stringency hybridization” refers generally to conditions that are about 10°C or less from the thermal melting temperature T_m as determined under the solution condition for a defined polynucleotide sequence. In some embodiments, a high stringency condition refers to conditions that permit hybridization of only those nucleic acid sequences that form stable hybrids in 0.018M NaCl at 65°C (i.e., if a hybrid is not stable in 0.018M NaCl at 65°C, it will not be stable under high stringency

conditions, as contemplated herein). High stringency conditions can be provided, for example, by hybridization in conditions equivalent to 50% formamide, 5× Denhart's solution, 5×SSPE, 0.2% SDS at 42°C, followed by washing in 0.1×SSPE, and 0.1% SDS at 65°C. Another high stringency condition is hybridizing in conditions equivalent to hybridizing in 5X SSC containing 0.1% (w:v) SDS at 65°C and washing in 0.1x SSC containing 0.1% SDS at 65°C. Other high stringency hybridization conditions, as well as moderately stringent conditions, are described in the references cited above.

[0102] “Codon optimized” refers to changes in the codons of the polynucleotide encoding a protein to those preferentially used in a particular organism such that the encoded protein is more efficiently expressed in the organism of interest. Although the genetic code is degenerate in that most amino acids are represented by several codons, called “synonyms” or “synonymous” codons, it is well known that codon usage by particular organisms is nonrandom and biased towards particular codon triplets. This codon usage bias may be higher in reference to a given gene, genes of common function or ancestral origin, highly expressed proteins versus low copy number proteins, and the aggregate protein coding regions of an organism's genome. In some embodiments, the polynucleotides encoding the GAA enzymes may be codon optimized for optimal production from the host organism selected for expression.

[0103] “Control sequence” refers herein to include all components, which are necessary or advantageous for the expression of a polynucleotide and/or polypeptide of the present application. Each control sequence may be native or foreign to the nucleic acid sequence encoding the polypeptide. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter sequence, signal peptide sequence, initiation sequence and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the nucleic acid sequence encoding a polypeptide.

[0104] “Operably linked” is defined herein as a configuration in which a control sequence is appropriately placed (i.e., in a functional relationship) at a position relative to a polynucleotide of interest such that the control sequence directs or regulates the expression of the polynucleotide and/or polypeptide of interest.

[0105] “Promoter sequence” refers to a nucleic acid sequence that is recognized by a host cell for expression of a polynucleotide of interest, such as a coding sequence. The promoter sequence contains transcriptional control sequences, which mediate the expression of a polynucleotide of interest. The promoter may be any nucleic acid sequence which shows transcriptional activity in the host cell of choice including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

[0106] “Suitable reaction conditions” refers to those conditions in the enzymatic conversion reaction solution (e.g., ranges of enzyme loading, substrate loading, temperature, pH, buffers, co-solvents, etc.) under which a GAA polypeptide of the present application is capable of converting a substrate to the desired product compound. Exemplary “suitable reaction conditions” are provided in the present application and illustrated by the Examples. “Loading”, such as in “compound loading” or “enzyme loading” refers to the concentration or amount of a component in a reaction mixture at the start of the reaction. “Substrate” in the context of an enzymatic conversion reaction process refers to the compound or molecule acted on by the GAA polypeptide. “Product” in the context of an enzymatic conversion process refers to the compound or molecule resulting from the action of the GAA polypeptide on a substrate.

[0107] As used herein the term “culturing” refers to the growing of a population of microbial cells under any suitable conditions (e.g., using a liquid, gel or solid medium).

[0108] Recombinant polypeptides can be produced using any suitable methods known the art. Genes encoding the wild-type polypeptide of interest can be cloned in vectors, such as plasmids, and expressed in desired hosts, such as *E. coli*, *S. cerevisiae*, etc. Variants of recombinant polypeptides can be generated by various methods known in the art. Indeed, there is a wide variety of different mutagenesis techniques well known to those skilled in the art. In addition, mutagenesis kits are also available from many commercial molecular biology suppliers. Methods are available to make specific substitutions at defined amino acids (site-directed), specific or random mutations in a localized region of the gene (regio-specific), or random mutagenesis over the entire gene (e.g., saturation mutagenesis). Numerous suitable methods are known to those in the art to generate enzyme variants, including but not limited to site-directed mutagenesis of single-stranded DNA or double-stranded DNA using PCR, cassette mutagenesis, gene synthesis, error-prone PCR, shuffling, and chemical saturation mutagenesis, or any other suitable method known in the art. Non-limiting examples of methods used for DNA and protein engineering are provided in the following patents: US Pat. No. 6,117,679; US Pat. No. 6,420,175; US Pat. No. 6,376,246; US Pat. No. 6,586,182; US Pat. No. 7,747,391; US Pat. No. 7,747,393; US Pat. No. 7,783,428; and US Pat. No. 8,383,346. After the variants are produced, they can be screened for any desired property (e.g., high or increased activity, or low or reduced activity, increased thermal activity, increased thermal stability, and/or acidic pH stability, etc.). In some embodiments, “recombinant GAA polypeptides” (also referred to herein as “engineered GAA polypeptides,” “variant GAA enzymes,” and “GAA variants”) find use.

[0109] As used herein, a “vector” is a DNA construct for introducing a DNA sequence into a cell. In some embodiments, the vector is an expression vector that is operably linked to a suitable control sequence capable of effecting the expression in a suitable host of the polypeptide encoded in the DNA sequence. In some embodiments, an “expression vector” has a promoter sequence operably linked to the DNA sequence (e.g., transgene) to drive expression in a host cell, and in some embodiments, also

comprises a transcription terminator sequence. In some preferred embodiments, the pDH vector provided herein finds use.

[0110] As used herein, the term “gene therapy vector” refers to vehicles or carriers suitable for delivery of polynucleotide sequences to cells. In some embodiments, the vectors encapsulate genes (e.g., therapeutic genes) or polynucleotide sequences for delivery to cells or tissues, including but not limited to adenovirus (AV), adeno-associated virus (AAV), lentivirus (LV), and non-viral vectors, such as liposomes. It is not intended that the present invention be limited to any specific gene therapy vector, as any vehicle suitable for a given setting finds use. The gene therapy vector may be designed to deliver genes to a specific species or host, or may find more general applicability.

[0111] As used herein, the term “expression” includes any step involved in the production of the polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, and post-translational modification. In some embodiments, the term also encompasses secretion of the polypeptide from a cell.

[0112] As used herein, the term “produces” refers to the production of proteins and/or other compounds by cells. It is intended that the term encompass any step involved in the production of polypeptides including, but not limited to, transcription, post-transcriptional modification, translation, and post-translational modification. In some embodiments, the term also encompasses secretion of the polypeptide from a cell.

[0113] As used herein, an amino acid or nucleotide sequence (e.g., a promoter sequence, signal peptide, terminator sequence, etc.) is “heterologous” to another sequence with which it is operably linked if the two sequences are not associated in nature.

[0114] As used herein, the terms “host cell” and “host strain” refer to suitable hosts for expression vectors comprising DNA provided herein (e.g., the polynucleotides encoding the GAA variants). In some embodiments, the host cells are prokaryotic or eukaryotic cells that have been transformed or transfected with vectors constructed using recombinant DNA techniques as known in the art.

[0115] The term “analogue” means a polypeptide having more than 70% sequence identity but less than 100% sequence identity (e.g., more than 75%, 78%, 80%, 83%, 85%, 88%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% sequence identity) with a reference polypeptide. In some embodiments, the term analogue refers to polypeptides that contain one or more non-naturally occurring amino acid residues including, but not limited, to homoarginine, ornithine and norvaline, as well as naturally occurring amino acids. In some embodiments, analogues also include one or more D-amino acid residues and non-peptide linkages between two or more amino acid residues.

[0116] The term “therapeutic” refers to a compound administered to a subject who shows signs or symptoms of pathology having beneficial or desirable medical effects.

[0117] The term “pharmaceutical composition” refers to a composition suitable for pharmaceutical use in a mammalian subject (e.g., human) comprising a pharmaceutically effective amount of an engineered GAA polypeptide encompassed by the invention and an acceptable carrier.

[0118] The term “gene therapy” refers to the delivery of a gene, polydeoxyribonucleotide, or polynucleotide sequence(s) with a gene therapy vector to cells or tissues for the modification of those cells or tissues for the treatment or prevention of a disease. Gene therapy may include replacing a mutated gene that causes disease with a healthy copy of the gene, or inactivating, or “knocking out,” a mutated gene that is functioning improperly. In some embodiments, gene therapy is used in the treatment of disease in patients.

[0119] The term “mRNA therapy” refers to the delivery of an mRNA polyribonucleotide sequence to cells or tissues for the modification of those cells or tissues for the treatment or prevention of a disease. In some embodiments, the mRNA polynucleotide sequences for delivery to cells or tissue, are formulated, for instance, but not limited to, in liposomes. In some embodiments, mRNA therapy is used in the treatment of disease in patients.

[0120] The term “cell therapy” refers to the delivery of living cells that have been modified exogenously to patients to provide a missing gene for the treatment or prevention of a disease. The modified cells are then reintroduced into the body.

[0121] The term “effective amount” means an amount sufficient to produce the desired result. One of general skill in the art may determine what the effective amount by using routine experimentation.

[0122] The terms “isolated” and “purified” are used to refer to a molecule (e.g., an isolated nucleic acid, polypeptide, etc.) or other component that is removed from at least one other component with which it is naturally associated. The term “purified” does not require absolute purity, rather it is intended as a relative definition.

[0123] The term “subject” encompasses mammals such as humans, non-human primates, livestock, companion animals, and laboratory animals (e.g., rodents and lagomorphs). It is intended that the term encompass females as well as males.

[0124] As used herein, the term “patient” means any subject that is being assessed for, treated for, or is experiencing disease.

[0125] The term “infant” refers to a child in the period of the first month after birth to approximately one (1) year of age. As used herein, the term “newborn” refers to child in the period from birth to the 28th day of life. The term “premature infant” refers to an infant born after the twentieth completed week of gestation, yet before full term, generally weighing ~500 to ~2499 grams at birth. A “very low birth weight infant” is an infant weighing less than 1500 g at birth.

[0126] As used herein, the term “child” refers to a person who has not attained the legal age for consent to treatment or research procedures. In some embodiments, the term refers to a person between the time of birth and adolescence.

[0127] As used herein, the term “adult” refers to a person who has attained legal age for the relevant jurisdiction (e.g., 18 years of age in the United States). In some embodiments, the term refers to any fully grown, mature organism. In some embodiments, the term “young adult” refers to a person less than 18 years of age, but who has reached sexual maturity.

[0128] As used herein, “composition” and “formulation” encompass products comprising at least one engineered GAA of the present invention, intended for any suitable use (e.g., pharmaceutical compositions, dietary/nutritional supplements, feed, etc.).

[0129] The terms “administration” and “administering” a composition mean providing a composition of the present invention to a subject (e.g., to a person suffering from the effects of Pompe disease).

[0130] The term “carrier” when used in reference to a pharmaceutical composition means any of the standard pharmaceutical carrier, buffers, and excipients, such as stabilizers, preservatives, and adjuvants.

[0131] The term “pharmaceutically acceptable” means a material that can be administered to a subject without causing any undesirable biological effects or interacting in a deleterious manner with any of the components in which it is contained and that possesses the desired biological activity.

[0132] As used herein, the term “excipient” refers to any pharmaceutically acceptable additive, carrier, diluent, adjuvant, or other ingredient, other than the active pharmaceutical ingredient (API; e.g., the engineered GAA polypeptides of the present invention). Excipients are typically included for formulation and/or administration purposes.

[0133] The term “therapeutically effective amount” when used in reference to symptoms of disease/condition refers to the amount and/or concentration of a compound (e.g., engineered GAA polypeptides) that ameliorates, attenuates, or eliminates one or more symptom of a disease/condition or prevents or delays the onset of symptom(s).

[0134] The term “therapeutically effective amount” when used in reference to a disease/condition refers to the amount and/or concentration of a composition (e.g., engineered GAA polypeptides) that ameliorates, attenuates, or eliminates the disease/condition. In some embodiments, the term is used in reference to the amount of a composition that elicits the biological (e.g., medical) response by a tissue, system, or animal subject that is sought by the researcher, physician, veterinarian, or other clinician.

[0135] It is intended that the terms “treating,” “treat” and “treatment” encompass preventative (e.g., prophylactic), as well as palliative treatment.

Engineered GAA Polypeptides:

[0136] In some embodiments, engineered GAA polypeptides are produced by cultivating a microorganism comprising at least one polynucleotide sequence encoding at least one engineered GAA polypeptide under conditions which are conducive for producing the engineered GAA polypeptide(s). In some embodiments, the engineered GAA polypeptide is recovered from the resulting culture medium and/or cells.

[0137] The present invention provides exemplary engineered GAA polypeptides having GAA activity. The Examples provide Tables showing sequence structural information correlating specific amino acid sequence features with the functional activity of the engineered GAA polypeptides. This

structure-function correlation information is provided in the form of specific amino acid residues differences relative to a reference engineered polypeptide, as indicated in the Examples. The Examples further provide experimentally determined activity data for the exemplary engineered GAA polypeptides.

Polynucleotides Encoding Engineered Polypeptides, Expression Vectors and Host Cells:

[0138] The present invention provides polynucleotides encoding the engineered GAA polypeptides described herein. In some embodiments, the polynucleotides are operatively linked to one or more heterologous regulatory sequences that control gene expression to create a recombinant polynucleotide capable of expressing the polypeptide. Expression constructs containing a heterologous polynucleotide encoding the engineered GAA polypeptides can be introduced into appropriate host cells to express the corresponding GAA polypeptide.

[0139] As will be apparent to the skilled artisan, availability of a protein sequence and the knowledge of the codons corresponding to the various amino acids provide a description of all the polynucleotides capable of encoding the subject polypeptides. The degeneracy of the genetic code, where the same amino acids are encoded by alternative or synonymous codons, allows an extremely large number of nucleic acids to be made, all of which encode the engineered GAA polypeptide. Thus, having knowledge of a particular amino acid sequence, those skilled in the art could make any number of different nucleic acids by simply modifying the sequence of one or more codons in a way which does not change the amino acid sequence of the protein. In this regard, the present invention specifically contemplates each and every possible variation of polynucleotides that could be made encoding the polypeptides described herein by selecting combinations based on the possible codon choices, and all such variations are to be considered specifically disclosed for any polypeptide described herein, including the variants provided in Tables 3-1, 3-2, 4-1, 6-1, 10-1, 10-2, 12-1, 13-1, 13-2, 14-1, 14-2, 15-1, 16-1, 17-1, 17-2, 17-3, 17-4, 17-5, 17-6, 17-7, 17-8, and/or 17-9, as well as SEQ ID NOS: 2, 8, and/or 14.

[0140] In various embodiments, the codons are preferably selected to fit the host cell in which the protein is being produced. For example, preferred codons used in bacteria are used for expression in bacteria. Consequently, codon optimized polynucleotides encoding the engineered GAA polypeptides contain preferred codons at about 40%, 50%, 60%, 70%, 80%, or greater than 90% of codon positions of the full-length coding region.

[0141] In some embodiments, as described above, the polynucleotide encodes an engineered polypeptide having GAA activity with the properties disclosed herein, wherein the polypeptide comprises an amino acid sequence having at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identity to a reference sequence selected from SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104, or the amino acid sequence of any variant as disclosed in Tables 3-1, 3-2, 4-1, 6-1, 10-1, 10-2, 12-1, 13-1, 13-2, 14-1, 14-2, 15-1,

16-1, 17-1, 17-2, 17-3, 17-4, 17-5, 17-6, 17-7, 17-8, and/or 17-9, and one or more residue differences as compared to the reference polypeptide of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104, or the amino acid sequence of any variant as disclosed in Table 3-1, 3-2, 4-1, 6-1, 10-1, 10-2, 12-1, 13-1, 13-2, 14-1, 14-2, 15-1, 16-1, 17-1, 17-2, 17-3, 17-4, 17-5, 17-6, 17-7, 17-8, and/or 17-9 (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acid residue positions). In some embodiments, the reference sequence is selected from SEQ ID NOS: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

[0142] The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% sequence identity to SEQ ID NO: 2. The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 2. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 27, 27/944, 28, 29/478, 30, 88, 107, 109, 109/842, 110, 113, 135, 137, 138, 148, 150, 247, 274, 276, 278, 375, 403, 414, 418, 418/499, 421, 426, 437, 444, 455, 463, 471, 471/478, 476, 489, 527, 547, 581, 610, 642, 668, 670, 692, 725/732, 750, 753, 786, 820, 862, 871, 895, 897, 930, 934, and 944, wherein the positions are numbered with reference to SEQ ID NO: 2. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from 27P, 27P/944W, 27R, 28P, 28R, 28S, 29T/478T, 30G, 30K, 30T, 88G, 88S, 107G, 107P, 109G/842E, 109P, 110G, 110L, 113S, 135A, 135Q, 137P, 138A, 148G, 148Y, 150G, 247R, 274G, 276F, 276Y, 278A, 278G, 375E, 403W, 414P, 418E/499R, 418R, 421S, 426R, 437S, 444T, 455V, 463A, 471Q/478S, 471S, 476A, 476H, 489R, 527R, 547G, 581G, 581T, 610A, 610G, 610S, 642M, 642Q, 642S, 668H, 670N, 692Q, 725N/732I, 750P, 753T, 786P, 786Y, 820E, 862G, 871E, 895R, 897V, 930R, 934R, 944G, and 944R, wherein the positions are numbered with reference to SEQ ID NO: 2. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from F27P, F27P/C944W, F27R, L28P, L28R, L28S, L29T/A478T, V30G, V30K, V30T, K88G, K88S, Q107G, Q107P, L109G/G842E, L109P, Q110G, Q110L, Q113S, S135A, S135Q, E137P, M138A, T148G, T148Y, T150G, Q247R, D274G, A276F, A276Y, T278A, T278G, I375E, R403W, R414P, A418E/H499R, A418R, Q421S, G426R, A437S, A444T, R455V, E463A, K471Q/A478S, K471S, S476A, S476H, A489R, N527R, A547G, K581G, K581T, W610A, W610G, W610S, L642M, L642Q, L642S, S668H, L670N, T692Q, K725N/V732I, A750P, A753T, R786P, R786Y, G820E, R862G, L871E, K895R, T897V, C930R, L934R, C944G,

and C944R, wherein the positions are numbered with reference to SEQ ID NO: 2. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 29/218/240/668/700/744/869, 29/218/240/700/869, 29/240/596/668/700/744/869, 29/240/596/668/869, 36/106/150/218/527/750/883/894, 106/112/150/218/414/527/793/883, 106/150/169/218/414/486/527/750/894, 106/150/169/218/414/486/527/894, 106/150/169/218/414/486/749/793/883/894, 106/150/169/218/414/486/750/793/883/894, 106/150/169/218/414/486/793/883, 106/150/169/218/414/486/894, 106/150/169/218/414/749/750/793/883, 106/150/169/218/414/749/793, 106/150/169/218/414/749/793/883, 106/150/169/218/486/527/749/793/894, 106/150/169/218/486/749/883, 106/150/169/218/486/883, 106/150/169/218/749/800, 106/150/169/414/486/749/750/883, 106/150/169/527/749/793/883, 106/150/169/749/793/883/894, 106/150/218/331/414/486/527/733/749/793, 106/150/218/414/486/642/750/793/883, 106/150/218/414/486/750/793/894, 106/150/218/414/527/749/750/883, 106/150/218/414/527/749/793/883/894, 106/150/218/414/749/750/793/883/894, 106/150/218/414/749/793/883, 106/150/218/486/527/749/894, 106/150/218/486/793/883, 106/150/218/527/749/750/793, 106/150/218/527/793/894, 106/150/218/749/750/793, 106/150/218/793, 106/150/218/793/894, 106/150/245/793/883/894, 106/150/414/749/750/793/894, 106/150/414/749/793/894, 106/150/486/527/750/793, 106/150/486/749/793/883/894, 106/150/749/793/883, 106/169/185/218/414/749/750/793, 106/191/280/402/414/444/727, 106/191/414/444/522/928/944, 106/191/414/489/928/944, 106/280/402/414/444/489/727/944, 150/169/218/414/527/793, 150/218/414/486/749/750, 150/218/414/486/750/793, 150/218/414/486/750/793/883, 150/218/414/749/750/793/894, 150/218/414/749/793, 150/218/527/749/793, 150/218/749/750/793, 150/218/749/793, 150/414/486/527/750/894, 150/414/486/749/750/793, 150/486/750/883/894, 169/486/750/793/883, 180/275/402/518/547/610/638/669/671, 180/402/431/507/547/610/669/671/793, 180/402/507/547/610/671, 191/280/402/414/444/465/842/928, 191/280/402/414/444/489/500/944, 191/280/414/444/489/500/522/842/928/944, 191/280/414/444/489/522/727/944, 191/280/414/489/842/928/944, 191/280/414/944, 191/414/522/842/944, 196/402/431/547/610/638, 218/668/700/869, 224/402/507/518/547/638/668, 269/275/431/518/547/638/668/669, 275/281/402/431/507/518/610/668, 275/281/402/431/518/547/610/669/671, 275/281/402/507/518/547/638/669/671, 275/281/402/518/547/610/638/671, 275/281/402/518/547/610/668/669/887, 275/281/402/547/610/638/669/671, 275/281/431/518/547/638/669/671, 275/281/507/547/669/671, 275/281/610/638/668/669, 275/281/671, 275/377/402/507/518/669/671/715, 275/402/431/507/547/671, 275/402/431/518/610/638/669/671/922, 275/402/507/547/610/638/668/669, 275/402/507/547/610/638/669/671, 275/402/507/547/610/671, 275/402/547/610/638/669/671,

275/402/547/638/669/671, 275/402/638/669/671, 275/431/507/518/547/668/669/671,
275/431/507/518/610/669/671, 275/431/507/547/610/638/671, 275/431/518/547/638/668,
275/431/518/610/638/669/671, 275/431/638, 275/507/518/547/610/638/668/669,
275/507/518/547/638/669/671, 275/507/547/610/638/669/671, 275/507/547/668/669/671,
275/518/671, 280/402/536/928, 281/402/507/518/547/610/638/669/671,
281/402/507/547/638/669/671, 281/402/518/547/610/638/668/669, 281/402/518/547/668,
281/431/507/518/547/610/638/668, 402/431/518/547/610/668, 402/431/518/547/671,
402/431/518/610, 402/431/547/638/671, 431/507/518/541/547/638/669/671, 431/507/518/669/671,
507/547/610, 507/547/638/669/671, 547/610/638/671, and 547/638/668, wherein the positions are
numbered with reference to SEQ ID NO: 2. In some embodiments, the acid alpha-glucosidase
comprises at least one substitution or substitution sets at one or more positions selected from
29Q/218S/240I/668D/700F/744V/869L, 29Q/240I/596P/668D/869L,
29Q/240I/596S/668D/700F/744V/869T, 29V/218S/240I/700F/869T,
36R/106P/150S/218S/527D/750P/883H/894R, 106P/112S/150S/218S/414G/527D/793K/883H,
106P/150S/169S/218S/414G/486E/527D/750P/894R, 106P/150S/169S/218S/414G/486E/527D/894R,
106P/150S/169S/218S/414G/486E/749E/793K/883H/894R,
106P/150S/169S/218S/414G/486E/750P/793K/883H/894R,
106P/150S/169S/218S/414G/486E/793K/883H, 106P/150S/169S/218S/414G/486E/894R,
106P/150S/169S/218S/414G/749E/750P/793K/883H, 106P/150S/169S/218S/414G/749E/793K,
106P/150S/169S/218S/414G/749E/793K/883H, 106P/150S/169S/218S/486E/527D/749E/793K/894R,
106P/150S/169S/218S/486E/749E/883H, 106P/150S/169S/218S/486E/883H,
106P/150S/169S/218S/749E/800A, 106P/150S/169S/414G/486E/749E/750P/883H,
106P/150S/169S/527D/749E/793K/883H, 106P/150S/169S/749E/793K/883H/894R,
106P/150S/218S/331A/414G/486E/527D/733E/749E/793K,
106P/150S/218S/414G/486E/642F/750P/793K/883H, 106P/150S/218S/414G/486E/750P/793K/894R,
106P/150S/218S/414G/527D/749E/750P/883H,
106P/150S/218S/414G/527D/749E/793K/883H/894G,
106P/150S/218S/414G/749E/750P/793K/883H/894R, 106P/150S/218S/414G/749E/793K/883H,
106P/150S/218S/486E/527D/749E/894R, 106P/150S/218S/486E/793K/883H,
106P/150S/218S/527D/749E/750P/793K, 106P/150S/218S/527D/793K/894G,
106P/150S/218S/749E/750P/793K, 106P/150S/218S/793K, 106P/150S/218S/793K/894R,
106P/150S/245S/793K/883H/894R, 106P/150S/414G/749E/750P/793K/894R,
106P/150S/414G/749E/793K/894R, 106P/150S/486E/527D/750P/793K,
106P/150S/486E/749E/793K/883H/894G, 106P/150S/749E/793K/883H,
106P/169S/185G/218S/414G/749E/750P/793K, 106P/191R/280D/402A/414G/444P/727P,
106P/191R/414G/444P/522V/928T/944S, 106P/191R/414G/489D/928T/944S,
106P/280D/402A/414G/444P/489D/727P/944S, 150S/169S/218S/414G/527D/793K,

150S/218S/414G/486A/750P/793K, 150S/218S/414G/486E/749E/750P,
150S/218S/414G/486E/750P/793K/883H, 150S/218S/414G/749E/750P/793K/894R,
150S/218S/414G/749E/793K, 150S/218S/527D/749E/793K, 150S/218S/749E/750P/793K,
150S/218S/749E/793K, 150S/414G/486E/527D/750P/894R, 150S/414G/486E/749E/750P/793K,
150S/486E/750P/883H/894G, 169S/486E/750P/793K/883H,
180H/275M/402A/518V/547G/610R/638I/669H/671N,
180H/402A/431V/507L/547G/610R/669H/671N/793G, 180H/402A/507L/547G/610R/671N,
191R/280D/402A/414G/444P/465E/842S/928T, 191R/280D/402A/414G/444P/489D/500A/944S,
191R/280D/414G/444P/489D/500A/522V/842S/928T/944S,
191R/280D/414G/444P/489D/522V/727P/944S, 191R/280D/414G/489D/842S/928T/944S,
191R/280D/414G/944S, 191R/414G/522V/842S/944S, 196V/402A/431V/547G/610R/638I,
218S/668D/700F/869T, 224F/402A/507L/518V/547G/638I/668D,
269N/275M/431V/518V/547G/638I/668D/669H, 275M/281V/402A/431V/507L/518V/610R/668D,
275M/281V/402A/507L/518V/547G/638I/669H/671N,
275M/281V/402A/518V/547G/610R/638I/671N,
275M/281V/402A/518V/547G/610R/668D/669H/887D,
275M/281V/402A/547G/610R/638I/669H/671N, 275M/281V/507L/547G/669H/671N,
275M/281V/610R/638I/668D/669H, 275M/402A/431V/507L/547G/671N,
275M/402A/507L/547G/610R/671N, 275M/402A/547G/638I/669H/671N,
275M/431V/518V/547G/638I/668D, 275M/431V/518V/610R/638I/669H/671N, 275M/431V/638I,
275M/507L/547G/668D/669H/671N, 275V/281V/402A/431V/518V/547G/610R/669H/671N,
275V/281V/431V/518V/547G/638I/669H/671N, 275V/281V/671N,
275V/377K/402A/507L/518V/669H/671N/715G,
275V/402A/431V/518V/610R/638I/669H/671N/922L,
275V/402A/507L/547G/610R/638I/668D/669H, 275V/402A/507L/547G/610R/638I/669H/671N,
275V/402A/547G/610R/638I/669H/671N, 275V/402A/638I/669H/671N,
275V/431V/507L/518V/547G/668D/669H/671N, 275V/431V/507L/518V/610R/669H/671N,
275V/431V/507L/547G/610R/638I/671N, 275V/507L/518V/547G/610R/638I/668D/669H,
275V/507L/518V/547G/638I/669H/671N, 275V/507L/547G/610R/638I/669H/671N,
275V/518V/671N, 280D/402A/536I/928T, 281V/402A/507L/518V/547G/610R/638I/669H/671N,
281V/402A/507L/547G/638I/669H/671N, 281V/402A/518V/547G/610R/638I/668D/669H,
281V/402A/518V/547G/668D, 281V/431V/507L/518V/547G/610R/638I/668D,
402A/431V/518V/547G/610R/668D, 402A/431V/518V/547G/671N, 402A/431V/518V/610R,
402A/431V/547G/638I/671N, 431V/507L/518V/541E/547G/638I/669H/671N,
431V/507L/518V/669H/671N, 507L/547G/610R, 507L/547G/638I/669H/671N,
547G/610R/638I/671N, and 547G/638I/668D, wherein the positions are numbered with reference to
SEQ ID NO: 2. In some embodiments, the acid alpha-glucosidase comprises at least one substitution

or substitution sets at one or more positions selected from

L29Q/L218S/L240I/S668D/H700F/I744V/I869L, L29Q/L240I/A596P/S668D/I869L,
L29Q/L240I/A596S/S668D/H700F/I744V/I869T, L29V/L218S/L240I/H700F/I869T,
G36R/K106P/T150S/L218S/N527D/A750P/R883H/Q894R,
K106P/A112S/T150S/L218S/R414G/N527D/E793K/R883H,
K106P/T150S/N169S/L218S/R414G/T486E/N527D/A750P/Q894R,
K106P/T150S/N169S/L218S/R414G/T486E/N527D/Q894R,
K106P/T150S/N169S/L218S/R414G/T486E/Q749E/E793K/R883H/Q894R,
K106P/T150S/N169S/L218S/R414G/T486E/A750P/E793K/R883H/Q894R,
K106P/T150S/N169S/L218S/R414G/T486E/E793K/R883H,
K106P/T150S/N169S/L218S/R414G/T486E/Q894R,
K106P/T150S/N169S/L218S/R414G/Q749E/A750P/E793K/R883H,
K106P/T150S/N169S/L218S/R414G/Q749E/E793K,
K106P/T150S/N169S/L218S/R414G/Q749E/E793K/R883H,
K106P/T150S/N169S/L218S/T486E/N527D/Q749E/E793K/Q894R,
K106P/T150S/N169S/L218S/T486E/Q749E/R883H, K106P/T150S/N169S/L218S/T486E/R883H,
K106P/T150S/N169S/L218S/Q749E/P800A,
K106P/T150S/N169S/R414G/T486E/Q749E/A750P/R883H,
K106P/T150S/N169S/N527D/Q749E/E793K/R883H,
K106P/T150S/N169S/Q749E/E793K/R883H/Q894R,
K106P/T150S/L218S/V331A/R414G/T486E/N527D/D733E/Q749E/E793K,
K106P/T150S/L218S/R414G/T486E/L642F/A750P/E793K/R883H,
K106P/T150S/L218S/R414G/T486E/A750P/E793K/Q894R,
K106P/T150S/L218S/R414G/N527D/Q749E/A750P/R883H,
K106P/T150S/L218S/R414G/N527D/Q749E/E793K/R883H/Q894G,
K106P/T150S/L218S/R414G/Q749E/A750P/E793K/R883H/Q894R,
K106P/T150S/L218S/R414G/Q749E/E793K/R883H,
K106P/T150S/L218S/T486E/N527D/Q749E/Q894R, K106P/T150S/L218S/T486E/E793K/R883H,
K106P/T150S/L218S/N527D/Q749E/A750P/E793K, K106P/T150S/L218S/N527D/E793K/Q894G,
K106P/T150S/L218S/Q749E/A750P/E793K, K106P/T150S/L218S/E793K,
K106P/T150S/L218S/E793K/Q894R, K106P/T150S/P245S/E793K/R883H/Q894R,
K106P/T150S/R414G/Q749E/A750P/E793K/Q894R, K106P/T150S/R414G/Q749E/E793K/Q894R,
K106P/T150S/T486E/N527D/A750P/E793K, K106P/T150S/T486E/Q749E/E793K/R883H/Q894G,
K106P/T150S/Q749E/E793K/R883H, K106P/N169S/V185G/L218S/R414G/Q749E/A750P/E793K,
K106P/H191R/G280D/S402A/R414G/A444P/S727P,
K106P/H191R/R414G/A444P/E522V/D928T/C944S, K106P/H191R/R414G/A489D/D928T/C944S,
K106P/G280D/S402A/R414G/A444P/A489D/S727P/C944S,

T150S/N169S/L218S/R414G/N527D/E793K, T150S/L218S/R414G/T486A/A750P/E793K,
T150S/L218S/R414G/T486E/Q749E/A750P, T150S/L218S/R414G/T486E/A750P/E793K/R883H,
T150S/L218S/R414G/Q749E/A750P/E793K/Q894R, T150S/L218S/R414G/Q749E/E793K,
T150S/L218S/N527D/Q749E/E793K, T150S/L218S/Q749E/A750P/E793K,
T150S/L218S/Q749E/E793K, T150S/R414G/T486E/N527D/A750P/Q894R,
T150S/R414G/T486E/Q749E/A750P/E793K, T150S/T486E/A750P/R883H/Q894G,
N169S/T486E/A750P/E793K/R883H,
N180H/L275M/S402A/I518V/A547G/W610R/V638I/L669H/S671N,
N180H/S402A/M431V/M507L/A547G/W610R/L669H/S671N/E793G,
N180H/S402A/M507L/A547G/W610R/S671N,
H191R/G280D/S402A/R414G/A444P/G465E/G842S/D928T,
H191R/G280D/S402A/R414G/A444P/A489D/D500A/C944S,
H191R/G280D/R414G/A444P/A489D/D500A/E522V/G842S/D928T/C944S,
H191R/G280D/R414G/A444P/A489D/E522V/S727P/C944S,
H191R/G280D/R414G/A489D/G842S/D928T/C944S, H191R/G280D/R414G/C944S,
H191R/R414G/E522V/G842S/C944S, A196V/S402A/M431V/A547G/W610R/V638I,
L218S/S668D/H700F/I869T, L224F/S402A/M507L/I518V/A547G/V638I/S668D,
T269N/L275M/M431V/I518V/A547G/V638I/S668D/L669H,
L275M/A281V/S402A/M431V/M507L/I518V/W610R/S668D,
L275M/A281V/S402A/M507L/I518V/A547G/V638I/L669H/S671N,
L275M/A281V/S402A/I518V/A547G/W610R/V638I/S671N,
L275M/A281V/S402A/I518V/A547G/W610R/S668D/L669H/E887D,
L275M/A281V/S402A/A547G/W610R/V638I/L669H/S671N,
L275M/A281V/M507L/A547G/L669H/S671N, L275M/A281V/W610R/V638I/S668D/L669H,
L275M/S402A/M431V/M507L/A547G/S671N, L275M/S402A/M507L/A547G/W610R/S671N,
L275M/S402A/A547G/V638I/L669H/S671N, L275M/M431V/I518V/A547G/V638I/S668D,
L275M/M431V/I518V/W610R/V638I/L669H/S671N, L275M/M431V/V638I,
L275M/M507L/A547G/S668D/L669H/S671N,
L275V/A281V/S402A/M431V/I518V/A547G/W610R/L669H/S671N,
L275V/A281V/M431V/I518V/A547G/V638I/L669H/S671N, L275V/A281V/S671N,
L275V/R377K/S402A/M507L/I518V/L669H/S671N/V715G,
L275V/S402A/M431V/I518V/W610R/V638I/L669H/S671N/P922L,
L275V/S402A/M507L/A547G/W610R/V638I/S668D/L669H,
L275V/S402A/M507L/A547G/W610R/V638I/L669H/S671N,
L275V/S402A/A547G/W610R/V638I/L669H/S671N, L275V/S402A/V638I/L669H/S671N,
L275V/M431V/M507L/I518V/A547G/S668D/L669H/S671N,
L275V/M431V/M507L/I518V/W610R/L669H/S671N,

L275V/M431V/M507L/A547G/W610R/V638I/S671N,
 L275V/M507L/I518V/A547G/W610R/V638I/S668D/L669H,
 L275V/M507L/I518V/A547G/V638I/L669H/S671N,
 L275V/M507L/A547G/W610R/V638I/L669H/S671N, L275V/I518V/S671N,
 G280D/S402A/V536I/D928T, A281V/S402A/M507L/I518V/A547G/W610R/V638I/L669H/S671N,
 A281V/S402A/M507L/A547G/V638I/L669H/S671N,
 A281V/S402A/I518V/A547G/W610R/V638I/S668D/L669H, A281V/S402A/I518V/A547G/S668D,
 A281V/M431V/M507L/I518V/A547G/W610R/V638I/S668D,
 S402A/M431V/I518V/A547G/W610R/S668D, S402A/M431V/I518V/A547G/S671N,
 S402A/M431V/I518V/W610R, S402A/M431V/A547G/V638I/S671N,
 M431V/M507L/I518V/G541E/A547G/V638I/L669H/S671N, M431V/M507L/I518V/L669H/S671N,
 M507L/A547G/W610R, M507L/A547G/V638I/L669H/S671N, A547G/W610R/V638I/S671N, and
 A547G/V638I/S668D, wherein the positions are numbered with reference to SEQ ID NO: 2.

[0143] The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% sequence identity to SEQ ID NO: 8. The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 8. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 4, 27, 27/28/489, 27/418/478, 28, 28/29, 28/29/113/135/138, 28/29/113/135/418, 28/29/135, 28/29/418, 29/113/126/135/193, 29/113/135, 29/113/135/455, 29/113/138, 29/148, 29/478, 106, 106/138/218/431/671/749, 106/218/281, 106/218/455, 106/218/455/507/749, 106/489/671, 106/638, 106/671/934, 113, 113/135/418, 113/418/455/478/581, 113/418/478/489/581, 135, 135/148/150/418, 135/478/489/581, 135/489, 135/944, 138/218/668/671, 138/218/749/934, 138/671/749/934, 157, 218, 218/281, 218/281/431, 218/281/671, 218/431, 218/431/489/507/749/934, 218/455, 218/507/749, 218/507/934, 218/638/671, 218/749, 281/431/489/668, 345/934, 418, 418/489, 431/668/671, 489/638/934, 489/671/934, 489/749, 489/934, 507/668, 507/671/934, 671/749, 671/934, and 749/784, wherein the positions are numbered with reference to SEQ ID NO: 8. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from 4H, 27P/28S/489R, 27P/418E/478T, 27R, 28S, 28S/29T, 28S/29T/113S/135Q/138A, 28S/29T/113S/135Q/418E, 28S/29T/135Q, 28S/29T/418E, 29T/113S/126Q/135Q/193Q, 29T/113S/135Q, 29T/113S/135Q/455V, 29T/113S/138A, 29T/148G, 29T/478T, 106P,

106P/138A/218S/431V/671N/749E, 106P/218S/281V, 106P/218S/455V, 106P/218S/455V/507L/749E, 106P/489R/671N, 106P/638I, 106P/671N/934R, 113S, 113S/135Q/418E, 113S/418E/455V/478T/581T, 113S/418E/478T/489R/581T, 135P/944Y, 135Q, 135Q/148G/150G/418E, 135Q/478T/489R/581T, 135Q/489R, 138A/218S/668D/671N, 138A/218S/749E/934R, 138A/671N/749E/934R, 157M, 218S, 218S/281V, 218S/281V/431V, 218S/281V/671N, 218S/431V, 218S/431V/489R/507L/749E/934R, 218S/455V, 218S/507L/749E, 218S/507L/934R, 218S/638I/671N, 218S/749E, 281V/431V/489R/668D, 345K/934R, 418E, 418E/489R, 431V/668D/671N, 489R/638I/934R, 489R/671N/934R, 489R/749E, 489R/934R, 507L/668D, 507L/671N/934R, 671N/749E, 671N/934R, and 749E/784T, wherein the positions are numbered with reference to SEQ ID NO: 8. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from P4H, F27P/L28S/A489R, F27P/A418E/A478T, F27R, L28S, L28S/L29T, L28S/L29T/Q113S/S135Q/M138A, L28S/L29T/Q113S/S135Q/A418E, L28S/L29T/S135Q, L28S/L29T/A418E, L29T/Q113S/P126Q/S135Q/H193Q, L29T/Q113S/S135Q, L29T/Q113S/S135Q/R455V, L29T/Q113S/M138A, L29T/T148G, L29T/A478T, K106P, K106P/M138A/L218S/M431V/S671N/Q749E, K106P/L218S/A281V, K106P/L218S/R455V, K106P/L218S/R455V/M507L/Q749E, K106P/A489R/S671N, K106P/V638I, K106P/S671N/L934R, Q113S, Q113S/S135Q/A418E, Q113S/A418E/R455V/A478T/K581T, Q113S/A418E/A478T/A489R/K581T, S135P/C944Y, S135Q, S135Q/T148G/S150G/A418E, S135Q/A478T/A489R/K581T, S135Q/A489R, M138A/L218S/S668D/S671N, M138A/L218S/Q749E/L934R, M138A/S671N/Q749E/L934R, L157M, L218S, L218S/A281V, L218S/A281V/M431V, L218S/A281V/S671N, L218S/M431V, L218S/M431V/A489R/M507L/Q749E/L934R, L218S/R455V, L218S/M507L/Q749E, L218S/M507L/L934R, L218S/V638I/S671N, L218S/Q749E, A281V/M431V/A489R/S668D, Q345K/L934R, A418E, A418E/A489R, M431V/S668D/S671N, A489R/V638I/L934R, A489R/S671N/L934R, A489R/Q749E, A489R/L934R, M507L/S668D, M507L/S671N/L934R, S671N/Q749E, S671N/L934R, and Q749E/A784, wherein the positions are numbered with reference to SEQ ID NO: 8.

[0144] The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% sequence identity to SEQ ID NO: 14. The present invention provides recombinant acid alpha-glucosidases and/or biologically active recombinant acid alpha-glucosidase fragments comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%

sequence identity to SEQ ID NO: 14. In some embodiments, the acid alpha-glucosidase comprises at least one substitution at position or set of positions selected from 22, 24, 27, 27/165, 30, 33, 34, 37/62, 37/62/79/196/696/862, 37/62/523, 37/62/523/793, 37/64/66/79/154/523/681/793/862, 37/79/154/793, 37/196, 37/528/696/793, 37/528/790, 37/528/790/793/862, 37/790/793, 39, 39/58/489/725/830/842/930/944, 39/70/109/830/842, 39/70/489/612, 39/70/725, 39/267, 39/267/489/522/612/830/842, 39/267/489/830/944, 39/489/500/612, 39/500/612, 40, 44/157, 47, 49, 50, 55, 60/500/612, 62/79/154/862, 62/79/196/681/862, 62/79/523/528/790, 62/79/790/793, 62/79/862, 62/92, 62/92/790/793, 62/106/523/528/696/793/862, 62/154/696/793/862, 62/793/862, 68, 70, 70/267/725/944, 70/267/930/944, 70/489/930, 70/725/830/860/930/944, 77, 79/154/681, 79/154/793/862, 79/862, 89, 97, 106/154, 107, 109, 109/522/612/725, 109/522/830/944, 109/612, 118, 149, 157, 158, 178, 179, 196/528/681/790/793, 207, 208, 217, 267/489/500/725/830/930, 267/522/725, 352, 385, 424, 448, 463, 489/830/944, 500, 500/612/830/860, 500/860/930, 500/930/944, 522/725, 523, 523/790/793, 528/681, 528/793, 528/862, 672, 673, 725, 734, 740, 753, 774, 778, 793, 830, 844, 862, 875, 880, 892, 902, 922, 925, 930, 932, 934, 938, and 944, wherein the positions are numbered with reference to SEQ ID NO: 14. In some embodiments, the acid alpha-glucosidase comprises at least one substitution or substitution sets at one or more positions selected from 22R, 24E, 24R, 24W, 27A, 27G, 27G/165I, 27K, 27R, 27S, 27V, 27W, 30D, 30L, 33G, 33P, 34D, 34M, 34T, 37F/62E, 37F/62E/79S/196T/696S/862Q, 37F/62E/523N, 37F/62E/523N/793K, 37F/64Q/66G/79S/154R/523N/681Q/793K/862Q, 37F/79S/154R/793K, 37F/196T, 37F/528S/696S/793K, 37F/528S/790V, 37F/528S/790V/793K/862Q, 37F/790V/793K, 39D, 39H, 39Q, 39Q/58L/489D/725E/830K/842S/930P/944S, 39Q/70A/109P/830K/842S, 39Q/70A/489D/612D, 39Q/70A/725E, 39Q/267K, 39Q/267K/489D/522V/612D/830K/842S, 39Q/267K/489D/830K/944S, 39Q/489D/500A/612D, 39Q/500A/612D, 40W, 44I/157V, 47G, 47R, 49A, 49G, 50G, 50L, 50V, 55C, 55L, 60V/500A/612D, 62E/79S/154R/862Q, 62E/79S/196T/681Q/862Q, 62E/79S/523N/528S/790V, 62E/79S/790V/793K, 62E/79S/862Q, 62E/92R, 62E/92R/790V/793K, 62E/106R/523N/528S/696S/793K/862Q, 62E/154R/696S/793K/862Q, 62E/793K/862Q, 68N, 68S, 68W, 70A/267K/725E/944S, 70A/267K/930P/944S, 70A/489D/930P, 70A/725E/830K/860F/930P/944S, 70Q, 77W, 79S/154R/681Q, 79S/154R/793K/862Q, 79S/862Q, 89R, 97D, 97G, 106R/154R, 107G, 109D, 109P/522V/612D/725E, 109P/522V/830K/944S, 109P/612D, 118F, 149R, 157Q, 158E, 158F, 178G, 178V, 179L, 196T/528S/681Q/790V/793K, 207R, 207Y, 208G, 208I, 217A, 217D, 267K/489D/500A/725E/830K/930P, 267K/522V/725E, 352K, 352V, 385G, 424K, 448L, 463A, 489D/830K/944S, 500A, 500A/612D/830K/860F, 500A/860F/930P, 500A/930P/944S, 522V/725E, 523N, 523N/790V/793K, 528S/681Q, 528S/793K, 528S/862Q, 672E, 672K, 673N, 673R, 725F, 725V, 734K, 740G, 740Q, 753S, 774G, 774S, 778Q, 793K, 830V, 844R, 862Q, 875D, 880R, 892L, 902L, 922E, 925A, 925W, 930P, 932A, 934F, 938A, 938P, 944R, and 944S, wherein the positions are numbered with reference to SEQ ID NO: 14. In some embodiments, the acid alpha-glucosidase

comprises at least one substitution or substitution sets at one or more positions selected from I22R, L24E, L24R, L24W, F27A, F27G, F27G/M165I, F27K, F27R, F27S, F27V, F27W, V30D, V30L, E33G, E33P, L34D, L34M, L34T, S37F/A62E, S37F/A62E/N79S/A196T/A696S/R862Q, S37F/A62E/D523N, S37F/A62E/D523N/E793K, S37F/P64Q/R66G/N79S/K154R/D523N/E681Q/E793K/R862Q, S37F/N79S/K154R/E793K, S37F/A196T, S37F/N528S/A696S/E793K, S37F/N528S/I790V, S37F/N528S/I790V/E793K/R862Q, S37F/I790V/E793K, P39D, P39H, P39Q, P39Q/R58L/A489D/K725E/Q830K/G842S/C930P/C944S, P39Q/V70A/L109P/Q830K/G842S, P39Q/V70A/A489D/S612D, P39Q/V70A/K725E, P39Q/R267K, P39Q/R267K/A489D/E522V/S612D/Q830K/G842S, P39Q/R267K/A489D/Q830K/C944S, P39Q/A489D/D500A/S612D, P39Q/D500A/S612D, V40W, T44I/L157V, A47G, A47R, Q49A, Q49G, Q50G, Q50L, Q50V, P55C, P55L, A60V/D500A/S612D, A62E/N79S/K154R/R862Q, A62E/N79S/A196T/E681Q/R862Q, A62E/N79S/D523N/N528S/I790V, A62E/N79S/I790V/E793K, A62E/N79S/R862Q, A62E/Q92R, A62E/Q92R/I790V/E793K, A62E/K106R/D523N/N528S/A696S/E793K/R862Q, A62E/K154R/A696S/E793K/R862Q, A62E/E793K/R862Q, R68N, R68S, R68W, V70A/R267K/K725E/C944S, V70A/R267K/C930P/C944S, V70A/A489D/C930P, V70A/K725E/Q830K/L860F/C930P/C944S, V70Q, P77W, N79S/K154R/E681Q, N79S/K154R/E793K/R862Q, N79S/R862Q, A89R, A97D, A97G, K106R/K154R, Q107G, L109D, L109P/E522V/S612D/K725E, L109P/E522V/Q830K/C944S, L109P/S612D, W118F, P149R, L157Q, T158E, T158F, P178G, P178V, A179L, A196T/N528S/E681Q/I790V/E793K, E207R, E207Y, E208G, E208I, Q217A, Q217D, R267K/A489D/D500A/K725E/Q830K/C930P, R267K/E522V/K725E, Y352K, Y352V, R385G, H424K, R448L, E463A, A489D/Q830K/C944S, D500A, D500A/S612D/Q830K/L860F, D500A/L860F/C930P, D500A/C930P/C944S, E522V/K725E, D523N, D523N/I790V/E793K, N528S/E681Q, N528S/E793K, N528S/R862Q, L672E, L672K, P673N, P673R, K725F, K725V, H734K, E740G, E740Q, A753S, A774G, A774S, L778Q, E793K, Q830V, E844R, R862Q, N875D, E880R, Q892L, A902L, P922E, K925A, K925W, C930P, S932A, L934F, Q938A, Q938P, C944R, and C944S, wherein the positions are numbered with reference to SEQ ID NO: 14.

[0145] In some embodiments, the recombinant acid alpha glucosidase comprises at least one mutation in at least one position as provided in Tables 3-1, 3-2, 4-1, 6-1, 10-1, 10-2, 12-1, 13-1, 13-2, 14-1, 14-2, 15-1, 16-1, 17-1, 17-2, 17-3, 17-4, 17-5, 17-6, 17-7, 17-8, and/or 17-9. In some additional embodiments, the recombinant acid alpha glucosidase is derived from a human acid alpha glucosidase. In yet some additional embodiments, the recombinant acid alpha glucosidase comprising the polypeptide sequence of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

[0146] In some embodiments, the polynucleotide encoding the engineered GAA polypeptides comprises a polynucleotide sequence selected from a polynucleotide sequence encoding SEQ ID NO: 1, 5, 7, 11, 13, 15, 17, 19, 945, 1955, 2495, 2879, and/or 3103. In some embodiments, the

polynucleotide encoding an engineered GAA polypeptide has at least about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% nucleotide residue identity to SEQ ID NO: 1, 5, 7, 11, 13, 15, 17, 19, 945, 1955, 2495, 2879, and/or 3103. In some embodiments, the polynucleotide encoding an engineered GAA polypeptide has at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% nucleotide residue identity to SEQ ID NO: 1, 5, 7, 11, 13, 15, 17, 19, 945, 1955, 2495, 2879, and/or 3103. In some embodiments, the polynucleotide encoding an engineered GAA polypeptide comprises SEQ ID NO: 1, 5, 7, 11, 13, 15, 17, 19, 945, 1955, 2495, 2879, and/or 3103. In some embodiments, the polynucleotide encoding an engineered GAA polypeptide consists of SEQ ID NO: 1, 5, 7, 11, 13, 15, 17, 19, 945, 1955, 2495, 2879, and/or 3103. In some embodiments, the polynucleotides are capable of hybridizing under highly stringent conditions to a reference polynucleotide sequence selected from SEQ ID NO: 1, 5, 7, 11, 13, 15, 17, 19, 945, 1955, 2495, 2879, and/or 3103, or a complement thereof, or a polynucleotide sequence encoding any of the variant GAA polypeptides provided herein.

[0147] In some embodiments, an isolated polynucleotide encoding any of the engineered GAA polypeptides provided herein is manipulated in a variety of ways to provide for expression of the polypeptide. In some embodiments, the polynucleotides encoding the polypeptides are provided as expression vectors where one or more control sequences is present to regulate the expression of the polynucleotides and/or polypeptides. Manipulation of the isolated polynucleotide prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotides and nucleic acid sequences utilizing recombinant DNA methods are well known in the art.

[0148] In some embodiments, the control sequences include among other sequences, promoters, Kozak sequence, leader sequences, polyadenylation sequences, propeptide sequences, signal peptide sequences, DNA based regulatory elements for gene therapy retention and transcription terminators. As known in the art, suitable promoters can be selected based on the host cells used. For bacterial host cells, suitable promoters for directing transcription of the nucleic acid constructs of the present application, include, but are not limited to the promoters obtained from the *E. coli* lac operon, *Streptomyces coelicolor* agarase gene (*dagA*), *Bacillus subtilis* levansucrase gene (*sacB*), *Bacillus licheniformis* alpha-amylase gene (*amyL*), *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), *Bacillus amyloliquefaciens* alpha-amylase gene (*amyQ*), *Bacillus licheniformis* penicillinase gene (*penP*), *Bacillus subtilis* *xylA* and *xylB* genes, and prokaryotic beta-lactamase gene (See e.g., Villa-Kamaroff et al., Proc. Natl Acad. Sci. USA 75: 3727-3731 [1978]), as well as the *tac* promoter (See e.g., DeBoer et al., Proc. Natl Acad. Sci. USA 80: 21-25 [1983]). Exemplary promoters for filamentous fungal host cells, include promoters obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase

(glaA), *Rhizomucor miehei* lipase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Aspergillus nidulans* acetamidase, and *Fusarium oxysporum* trypsin-like protease (See e.g., WO 96/00787), as well as the NA2-tpi promoter (a hybrid of the promoters from the genes for *Aspergillus niger* neutral alpha-amylase and *Aspergillus oryzae* triose phosphate isomerase), and mutant, truncated, and hybrid promoters thereof. Exemplary yeast cell promoters can be from the genes can be from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* galactokinase (GAL1), *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP), and *Saccharomyces cerevisiae* 3-phosphoglycerate kinase. Other useful promoters for yeast host cells are known in the art (See e.g., Romanos et al., Yeast 8:423-488 [1992]). Exemplary promoters for use in mammalian cells include, but are not limited to those from cytomegalovirus (CMV), chicken β -actin promoter fused with the CMV enhancer, Simian vacuolating virus 40 (SV40), from *Homo sapiens* phosphoglycerate kinase, beta actin, elongation factor-1a or glyceraldehyde-3-phosphate dehydrogenase, or from *Gallus gallus* β -actin.

[0149] In some embodiments, the control sequence is a suitable transcription terminator sequence, a sequence recognized by a host cell to terminate transcription. The terminator sequence is operably linked to the 3' terminus of the nucleic acid sequence encoding the polypeptide. Any terminator which is functional in the host cell of choice finds use in the present invention. For example, exemplary transcription terminators for filamentous fungal host cells can be obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* alpha-glucosidase, and *Fusarium oxysporum* trypsin-like protease. Exemplary terminators for yeast host cells can be obtained from the genes for *Saccharomyces cerevisiae* enolase, *Saccharomyces cerevisiae* cytochrome C (CYC1), and *Saccharomyces cerevisiae* glyceraldehyde-3-phosphate dehydrogenase. Other useful terminators for yeast host cells are known in the art (See e.g., Romanos et al., *supra*). Exemplary terminators for mammalian cells include, but are not limited to those from cytomegalovirus (CMV), Simian vacuolating virus 40 (SV40), from *Homo sapiens* growth hormone hGH, from bovine growth hormone BGH, and from human or rabbit beta globulin.

[0150] In some embodiments, the control sequence is a suitable leader sequence, 5'-cap modification, 5' UTR, etc. In some embodiments, these regulatory sequence elements mediate binding to molecules involved in mRNA trafficking and translation, inhibit 5'-exonucleolytic degradation and confer resistance to de-capping. The leader sequence is operably linked to the 5' terminus of the nucleic acid sequence encoding the polypeptide. Any leader sequence that is functional in the host cell of choice may be used. Exemplary leaders for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase and *Aspergillus nidulans* triose phosphate isomerase. Suitable leaders for yeast host cells include, but are not limited to those obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* 3-phosphoglycerate kinase,

Saccharomyces cerevisiae alpha-factor, and *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP). Suitable leaders for mammalian host cells include but are not limited to the 5'-UTR element present in orthopoxvirus mRNA.

[0151] In some embodiments, the control sequence comprises a 3' untranslated nucleic acid region and polyadenylation tail nucleic acid sequence, sequences operably linked to the 3' terminus of the protein coding nucleic acid sequence and which mediate binding to proteins involved in mRNA trafficking and translation and mRNA half-life. Any polyadenylation sequence and 3' UTR which is functional in the host cell of choice may be used in the present invention. Exemplary polyadenylation sequences for filamentous fungal host cells include, but are not limited to those from the genes for *Aspergillus oryzae* TKA amylose, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Fusarium oxysporum* trypsin-like protease, and *Aspergillus niger* alpha-glucosidase. Useful polyadenylation sequences for yeast host cells are also known in the art (See e.g., Guo and Sherman, Mol. Cell. Biol., 15:5983-5990 [1995]). Useful polyadenylation and 3' UTR sequences for mammalian host cells include, but are not limited to the 3'-UTRs of α - and β -globin mRNAs that harbor several sequence elements that increase the stability and translation of mRNA.

[0152] In some embodiments, the control sequence is a signal peptide coding region that codes for an amino acid sequence linked to the amino terminus of a polypeptide and directs the encoded polypeptide into the cell's secretory pathway. The 5' end of the coding sequence of the nucleic acid sequence may inherently contain a signal peptide coding region naturally linked in translation reading frame with the segment of the coding region that encodes the secreted polypeptide. Alternatively, the 5' end of the coding sequence may contain a signal peptide coding region that is foreign to the coding sequence. Any signal peptide coding region that directs the expressed polypeptide into the secretory pathway of a host cell of choice finds use for expression of the engineered GAA polypeptides provided herein. Effective signal peptide coding regions for bacterial host cells include, but are not limited to the signal peptide coding regions obtained from the genes for *Bacillus* NCIB 11837 maltogenic amylase, *Bacillus stearothermophilus* alpha-amylase, *Bacillus licheniformis* subtilisin, *Bacillus licheniformis* beta-lactamase, *Bacillus stearothermophilus* neutral proteases (nprT, nprS, nprM), and *Bacillus subtilis* prsA. Further signal peptides are known in the art (See e.g., Simonen and Palva, Microbiol. Rev., 57:109-137 [1993]). Effective signal peptide coding regions for filamentous fungal host cells include, but are not limited to the signal peptide coding regions obtained from the genes for *Aspergillus oryzae* TKA amylose, *Aspergillus niger* neutral amylase, *Aspergillus niger* glucoamylase, *Rhizomucor miehei* aspartic proteinase, *Humicola insolens* cellulase, and *Humicola lanuginosa* lipase. Useful signal peptides for yeast host cells include, but are not limited to those from the genes for *Saccharomyces cerevisiae* alpha-factor and *Saccharomyces cerevisiae* invertase. In some embodiments, the *S. cerevisiae* a-mating factor prepro peptide (Mfalpa) finds use (e.g., SEQ ID NOS: 3383 and 3384). Useful signal peptides for mammalian host cells include, but are not

limited to those from the genes for immunoglobulin gamma (IgG). Additional signal peptides useful for mammalian hosts include mouse signal peptides. In some embodiments, a synthetic mouse IG signal peptide finds use (e.g., SEQ ID NOS: 3381 and 3382).

[0153] In some embodiments, the control sequence is a propeptide coding region that codes for an amino acid sequence positioned at the amino terminus of a polypeptide. The resultant polypeptide is referred to as a “proenzyme,” “propolypeptide,” or “zymogen,” in some cases). A propolypeptide can be converted to a mature active polypeptide by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide. The propeptide coding region includes, but is not limited to the genes for *Bacillus subtilis* alkaline protease (aprE), *Bacillus subtilis* neutral protease (nprT), *Saccharomyces cerevisiae* alpha-factor, *Rhizomucor miehei* aspartic proteinase, and *Myceliophthora thermophila* lactase (See e.g., WO 95/33836). Where both signal peptide and propeptide regions are present at the amino terminus of a polypeptide, the propeptide region is positioned next to the amino terminus of a polypeptide and the signal peptide region is positioned next to the amino terminus of the propeptide region.

[0154] In some embodiments, regulatory sequences are also utilized. These sequences facilitate the regulation of the expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those which cause the expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. In prokaryotic host cells, suitable regulatory sequences include, but are not limited to the *lac*, *tac*, and *trp* operator systems. In yeast host cells, suitable regulatory systems include, but are not limited to the ADH2 system or GAL1 system. In filamentous fungi, suitable regulatory sequences include, but are not limited to the TAKA alpha-amylase promoter, *Aspergillus niger* glucoamylase promoter, and *Aspergillus oryzae* glucoamylase promoter.

[0155] In another aspect, the present invention also provides a recombinant expression vector comprising a polynucleotide encoding an engineered GAA polypeptide, and one or more expression regulating regions such as a promoter and a terminator, a replication origin, etc., depending on the type of hosts into which they are to be introduced. In some embodiments, the various nucleic acid and control sequences described above are joined together to produce a recombinant expression vector which includes one or more convenient restriction sites to allow for insertion or substitution of the nucleic acid sequence encoding the variant GAA polypeptide at such sites. Alternatively, the polynucleotide sequence(s) of the present invention are expressed by inserting the polynucleotide sequence or a nucleic acid construct comprising the polynucleotide sequence into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression.

[0156] The recombinant expression vector may be any suitable vector (e.g., a plasmid or virus including but not limited to adenovirus (AV), adeno-associated virus (AAV), lentivirus (LV), and non-viral vectors, such as liposomes), that can be conveniently subjected to recombinant DNA

procedures and can result in the expression of the variant GAA polynucleotide sequence. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vectors may be linear or closed circular plasmids. Indeed, it is not intended that the present invention be limited to any specific vector.

[0157] In some embodiments, the expression vector is an autonomously replicating vector (i.e., a vector that exists as an extra-chromosomal entity, the replication of which is independent of chromosomal replication, such as a plasmid, an extra-chromosomal element, a minichromosome, or an artificial chromosome). The vector may contain any means for assuring self-replication. In some alternative embodiments, the vector may be one which, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. Furthermore, a single vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the host cell, or a transposon may be used.

[0158] In some embodiments, the expression vector is pDH. A plasmid map of this vector containing SEQ ID NO:1 is provided in Figure 9. The sequence of this plasmid is provided as SEQ ID NO: 3379. A plasmid map of this vector containing a beta-lactamase stuffer sequence is provided in Figure 10. The sequence of this plasmid is provided as SEQ ID NO: 3380. It is contemplated that the pDH vector will find use in expression of various genes, including but not limited to the polynucleotide sequences encoding the acid alpha-glucosidases provided herein. Indeed, it is contemplated that the stuffer sequence (or SEQ ID NO: 1 present in SEQ ID NO: 3379) will be substituted with any suitable gene of interest.

[0159] In some embodiments, the expression vector preferably contains one or more selectable markers, which permit easy selection of transformed cells. A “selectable marker” is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like. Examples of bacterial selectable markers include, but are not limited to the *dal* genes from *Bacillus subtilis* or *Bacillus licheniformis*, or markers, which confer antibiotic resistance such as ampicillin, kanamycin, chloramphenicol or tetracycline resistance. Suitable markers for yeast host cells include, but are not limited to ADE2, HIS3, LEU2, LYS2, MET3, TRP1, and URA3. Selectable markers for use in a filamentous fungal host cell include, but are not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferases), *bar* (phosphinothricin acetyltransferase), *hph* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase), *sC* (sulfate adenylyltransferase), and *trpC* (anthranilate synthase), as well as equivalents thereof. In another aspect, the present invention provides a host cell comprising a polynucleotide encoding at least one engineered GAA polypeptide of the present application, the polynucleotide being operatively linked to one or more control sequences for expression of the engineered GAA enzyme(s) in the host cell. Host cells for use in expressing the polypeptides encoded by the expression vectors of the present invention are well known in the art and include but are not limited to, bacterial cells, such as *E. coli*, *Vibrio fluvialis*, *Streptomyces* and *Salmonella typhimurium* cells, fungal cells,

such as yeast cells (e.g., *Saccharomyces cerevisiae* and *Pichia pastoris* [ATCC Accession No. 201178]); insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells (e.g., CHO, CHO-K1, COS, and BHK), and human cells (e.g., HEK293T, human fibroblast, THP-1, Jurkat and Bowes melanoma cell lines); and plant cells.

[0160] Accordingly, in another aspect, the present invention provides methods for producing the engineered GAA polypeptides, where the methods comprise culturing a host cell capable of expressing a polynucleotide encoding the engineered GAA polypeptide under conditions suitable for expression of the polypeptide. In some embodiments, the methods further comprise the steps of isolating and/or purifying the GAA polypeptides, as described herein.

[0161] Appropriate culture media and growth conditions for the above-described host cells are well known in the art. Polynucleotides for expression of the GAA polypeptides may be introduced into cells by various methods known in the art. Techniques include, among others, electroporation, biolistic particle bombardment, liposome mediated transfection, calcium chloride transfection, and protoplast fusion.

[0162] The engineered GAA with the properties disclosed herein can be obtained by subjecting the polynucleotide encoding the naturally occurring or engineered GAA polypeptide to mutagenesis and/or directed evolution methods known in the art, and as described herein. An exemplary directed evolution technique is mutagenesis and/or DNA shuffling (See e.g., Stemmer, Proc. Natl. Acad. Sci. USA 91:10747-10751 [1994]; WO 95/22625; WO 97/0078; WO 97/35966; WO 98/27230; WO 00/42651; WO 01/75767 and U.S. Pat. 6,537,746). Other directed evolution procedures that can be used include, among others, staggered extension process (StEP), in vitro recombination (See e.g., Zhao et al., Nat. Biotechnol., 16:258–261 [1998]), mutagenic PCR (See e.g., Caldwell et al., PCR Methods Appl., 3:S136-S140 [1994]), and cassette mutagenesis (See e.g., Black et al., Proc. Natl. Acad. Sci. USA 93:3525-3529 [1996]).

[0163] For example, mutagenesis and directed evolution methods can be readily applied to polynucleotides to generate variant libraries that can be expressed, screened, and assayed.

Mutagenesis and directed evolution methods are well known in the art (See e.g., US Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, 5,837,458, 5,928,905, 6,096,548, 6,117,679, 6,132,970, 6,165,793, 6,180,406, 6,251,674, 6,265,201, 6,277,638, 6,287,861, 6,287,862, 6,291,242, 6,297,053, 6,303,344, 6,309,883, 6,319,713, 6,319,714, 6,323,030, 6,326,204, 6,335,160, 6,335,198, 6,344,356, 6,352,859, 6,355,484, 6,358,740, 6,358,742, 6,365,377, 6,365,408, 6,368,861, 6,372,497, 6,337,186, 6,376,246, 6,379,964, 6,387,702, 6,391,552, 6,391,640, 6,395,547, 6,406,855, 6,406,910, 6,413,745, 6,413,774, 6,420,175, 6,423,542, 6,426,224, 6,436,675, 6,444,468, 6,455,253, 6,479,652, 6,482,647, 6,483,011, 6,484,105, 6,489,146, 6,500,617, 6,500,639, 6,506,602, 6,506,603, 6,518,065, 6,519,065, 6,521,453, 6,528,311, 6,537,746, 6,573,098, 6,576,467, 6,579,678, 6,586,182, 6,602,986, 6,605,430, 6,613,514, 6,653,072, 6,686,515, 6,703,240, 6,716,631, 6,825,001, 6,902,922, 6,917,882, 6,946,296, 6,961,664, 6,995,017, 7,024,312, 7,058,515, 7,105,297, 7,148,054, 7,220,566, 7,288,375, 7,384,387,

7,421,347, 7,430,477, 7,462,469, 7,534,564, 7,620,500, 7,620,502, 7,629,170, 7,702,464, 7,747,391, 7,747,393, 7,751,986, 7,776,598, 7,783,428, 7,795,030, 7,853,410, 7,868,138, 7,783,428, 7,873,477, 7,873,499, 7,904,249, 7,957,912, 7,981,614, 8,014,961, 8,029,988, 8,048,674, 8,058,001, 8,076,138, 8,108,150, 8,170,806, 8,224,580, 8,377,681, 8,383,346, 8,457,903, 8,504,498, 8,589,085, 8,762,066, 8,768,871, 9,593,326, 9,684,771, 9,665,694; and WO 95/22625; WO 97/0078; WO 97/35966; WO 98/27230; WO 00/42651; WO 01/75767; WO 2009/152336, WO 2013/138339, WO 2015/048572, and WO 2015/048573; and all related US and non-US counterparts of these listed patents and applications; Ling et al., *Anal. Biochem.*, 254:157-78 [1997]; Dale et al., *Meth. Mol. Biol.*, 57:369-74 [1996]; Smith, *Ann. Rev. Genet.*, 19:423-462 [1985]; Botstein et al., *Science*, 229:1193-1201 [1985]; Carter, *Biochem. J.*, 237:1-7 [1986]; Kramer et al., *Cell*, 38:879-887 [1984]; Wells et al., *Gene*, 34:315-323 [1985]; Minshull et al., *Curr. Op. Chem. Biol.*, 3:284-290 [1999]; Christians et al., *Nat. Biotechnol.*, 17:259-264 [1999]; Crameri et al., *Nature*, 391:288-291 [1998]; Crameri, et al., *Nat. Biotechnol.*, 15:436-438 [1997]; Zhang et al., *Proc. Nat. Acad. Sci. U.S.A.*, 94:4504-4509 [1997]; Crameri et al., *Nat. Biotechnol.*, 14:315-319 [1996]; Stemmer, *Nature*, 370:389-391 [1994]; and Stemmer, *Proc. Nat. Acad. Sci. USA*, 91:10747-10751 [1994]; all of which are incorporated herein by reference).

[0164] In some embodiments, the enzyme clones obtained following mutagenesis treatment are screened by subjecting the enzymes to a defined temperature (or other assay conditions) and measuring the amount of enzyme activity remaining after heat treatments or other assay conditions. Clones containing a polynucleotide encoding a GAA polypeptide are then isolated from the gene, sequenced to identify the nucleotide sequence changes (if any), and used to express the enzyme in a host cell. Measuring enzyme activity from the expression libraries can be performed using any suitable method known in the art (e.g., standard biochemistry techniques, such as HPLC analysis).

[0165] For engineered polypeptides of known sequence, the polynucleotides encoding the enzyme can be prepared by standard solid-phase methods, according to known synthetic methods. In some embodiments, fragments of up to about 100 bases can be individually synthesized, then joined (e.g., by enzymatic or chemical ligation methods, or polymerase mediated methods) to form any desired continuous sequence. For example, polynucleotides and oligonucleotides disclosed herein can be prepared by chemical synthesis using the classical phosphoramidite method (See e.g., Beaucage et al., *Tetra. Lett.*, 22:1859-69 [1981]; and Matthes et al., *EMBO J.*, 3:801-05 [1984]), as it is typically practiced in automated synthetic methods. According to the phosphoramidite method, oligonucleotides are synthesized (e.g., in an automatic DNA synthesizer), purified, annealed, ligated and cloned in appropriate vectors.

[0166] Accordingly, in some embodiments, a method for preparing the engineered GAA polypeptide can comprise: (a) synthesizing a polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the amino acid sequence of any variant provided in Table 3-1, 3-2, 4-1, 6-1, 10-1, 10-2, 12-1, 13-1, 13-2, 14-1, 14-2, 15-1, 16-1, 17-1, 17-2, 17-3, 17-4, 17-5, 17-6, 17-7, 17-8,

and/or 17-9, as well as SEQ ID NOS: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104, and (b) expressing the GAA polypeptide encoded by the polynucleotide. In some embodiments of the method, the amino acid sequence encoded by the polynucleotide can optionally have one or several (e.g., up to 3, 4, 5, or up to 10) amino acid residue deletions, insertions and/or substitutions. In some embodiments, the amino acid sequence has optionally 1-2, 1-3, 1-4, 1-5, 1-6, 1-7, 1-8, 1-9, 1-10, 1-15, 1-20, 1-21, 1-22, 1-23, 1-24, 1-25, 1-30, 1-35, 1-40, 1-45, or 1-50 amino acid residue deletions, insertions and/or substitutions. In some embodiments, the amino acid sequence has optionally 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, or 50 amino acid residue deletions, insertions and/or substitutions. In some embodiments, the amino acid sequence has optionally 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, 20, 21, 22, 23, 24, or 25 amino acid residue deletions, insertions and/or substitutions. In some embodiments, the substitutions can be conservative or non-conservative substitutions.

[0167] The expressed engineered GAA polypeptide can be measured for any desired improved property (e.g., activity, selectivity, stability, acid tolerance, protease sensitivity, etc.), using any suitable assay known in the art, including but not limited to the assays and conditions described herein.

[0168] In some embodiments, any, some, or all of the engineered GAA polypeptides expressed in a host cell are recovered from the cells and/or the culture medium using any one or more of the well-known techniques for protein purification, including, among others, lysozyme treatment, sonication, filtration, salting-out, ultra-centrifugation, and chromatography.

[0169] Chromatographic techniques for isolation of the GAA polypeptides include, among others, reverse phase chromatography high performance liquid chromatography, ion exchange chromatography, hydrophobic interaction chromatography, gel electrophoresis, and affinity chromatography. Conditions for purifying a particular enzyme depends, in part, on factors such as net charge, hydrophobicity, hydrophilicity, molecular weight, molecular shape, etc., and will be apparent to those having skill in the art. In some embodiments, affinity techniques may be used to isolate the improved variant GAA enzymes. In some embodiments utilizing affinity chromatography purification, any antibody which specifically binds the variant GAA polypeptide finds use. For the production of antibodies, various host animals, including but not limited to rabbits, mice, rats, etc., are immunized by injection with a GAA polypeptide (e.g., a GAA variant), or a fragment thereof. In some embodiments, the GAA polypeptide or fragment is attached to a suitable carrier, such as BSA, by means of a side chain functional group or linkers attached to a side chain functional group.

[0170] In some embodiments, the engineered GAA polypeptide is produced in a host cell by a method comprising culturing a host cell (e.g., *S. cerevisiae*, *Daucus carota*, *Nicotiana tabacum*, *H. sapiens* [e.g., HEK293T], or *Cricetulus griseus* [e.g., CHO]) comprising a polynucleotide sequence encoding an engineered GAA polypeptide as described herein under conditions conducive to the

production of the engineered GAA polypeptide and recovering the engineered GAA polypeptide from the cells and/or culture medium.

[0171] In some embodiments, the engineered GAA polypeptide is produced in a host cell by a method comprising culturing a host cell comprising a polynucleotide sequence encoding an engineered GAA polypeptide as described herein under conditions conducive to the production of the engineered GAA polypeptide and recovering the engineered GAA polypeptide from the cells and/or culture medium.

[0172] In some preferred embodiments, the invention encompasses a method of producing an engineered GAA polypeptide comprising culturing a recombinant bacterial cell comprising a polynucleotide sequence encoding an engineered GAA polypeptide having at least 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to reference sequences SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104, and one or more amino acid residue differences as compared to SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104, and/or combinations thereof when aligned, under suitable culture conditions to allow the production of the engineered GAA polypeptide and optionally recovering the engineered GAA polypeptide from the culture and/or cultured bacterial cells.

[0173] In some embodiments, once the engineered GAA polypeptides are recovered from the recombinant host cells or cell culture and they are further purified by any suitable method(s) known in the art. In some additional embodiments, the purified GAA polypeptides are combined with other ingredients and compounds to provide compositions and formulations comprising the engineered GAA polypeptide as appropriate for different applications and uses (e.g., pharmaceutical compositions).

Compositions:

[0174] The present invention provides engineered GAA polypeptides suitable for use in pharmaceutical and other compositions, such as dietary/nutritional supplements, as well as for other purposes.

Pharmaceutical Compositions:

[0175] Depending on the mode of administration, the compositions comprising a therapeutically effective amount of an engineered GAA according to the present invention are in the form of a solid, semi-solid, gel, or liquid. In some embodiments, the compositions include other pharmaceutically acceptable components such as diluents, buffers, excipients, salts, emulsifiers, preservatives, stabilizers, fillers, and other ingredients. Details on techniques for formulation and administration are well known in the art and described in the literature.

[0176] In some embodiments, the engineered GAA polypeptides are formulated for use in oral pharmaceutical compositions. Any suitable format for use in delivering the engineered GAA polypeptides find use in the present invention, including but not limited to pills, tablets, gel tabs,

capsules, lozenges, dragees, powders, soft gels, sol-gels, gels, emulsions, implants, patches, sprays, ointments, liniments, creams, pastes, jellies, paints, aerosols, chewing gums, demulcents, sticks, suspensions (including but not limited to oil-based suspensions, oil-in water emulsions, etc.), slurries, syrups, controlled release formulations, suppositories, etc. In some embodiments, the engineered GAA polypeptides are provided in a format suitable for injection (i.e., in an injectable formulation). In some embodiments, the engineered GAA polypeptides are provided in biocompatible matrices such as sol-gels, including silica-based (e.g., oxysilane) sol-gels. In some embodiments, the engineered GAA polypeptides are encapsulated. In some alternative embodiments, the engineered GAA polypeptides are encapsulated in nanostructures (e.g., nanotubes, nanotubules, nanocapsules, or microcapsules, microspheres, liposomes, etc.). Indeed, it is not intended that the present invention be limited to any particular delivery formulation and/or means of delivery. It is intended that the engineered GAA polypeptides be administered by any suitable means known in the art, including but not limited to parenteral, oral, topical, transdermal, intranasal, intraocular, intrathecal, via implants, etc.

[0177] In some embodiments, the engineered GAA polypeptides are chemically modified by glycosylation, pegylation (i.e., modified with polyethylene glycol [PEG] or activated PEG, etc.) or other compounds (See e.g., Ikeda, Amino Acids 29:283-287 [2005]; US Pat. Nos. 7,531,341, 7,534,595, 7,560,263, and 7,53,653; US Pat. Appln. Publ. Nos. 2013/0039898, 2012/0177722, etc.). Indeed, it is not intended that the present invention be limited to any particular delivery method and/or mechanism.

[0178] In some additional embodiments, the engineered GAA polypeptides are provided for delivery to cells or tissues via gene therapy, including viral delivery vectors, including but not limited to adenovirus (AV), adeno-associated virus (AAV), lentivirus (LV), or non-viral vectors (e.g., liposomes). In some embodiments, the engineered GAA polypeptides are provided for delivery to cells or tissues via mRNA therapy following formulation of polyribonucleotide sequences in a encapsulated delivery, such as liposomes. In some additional embodiments, the engineered GAA polypeptides are provided for delivery to cells or tissues via cell therapy, where the polynucleotide sequence encoding the engineered GAA polypeptides is introduced into exogenous cell and that cell (or cells) are introduced into a recipient (e.g., a patient exhibiting or at risk for developing Pompe disease).

[0179] In some additional embodiments, the engineered GAA polypeptides are provided in formulations comprising matrix-stabilized enzyme crystals. In some embodiments, the formulation comprises a cross-linked crystalline engineered GAA enzyme and a polymer with a reactive moiety that adheres to the enzyme crystals. The present invention also provides engineered GAA polypeptides in polymers.

[0180] In some embodiments, compositions comprising the engineered GAA polypeptides of the present invention include one or more commonly used carrier compounds, including but not limited to

sugars (e.g., lactose, sucrose, mannitol, and/or sorbitol), starches (e.g., corn, wheat, rice, potato, or other plant starch), cellulose (e.g., methyl cellulose, hydroxypropylmethyl cellulose, sodium carboxymethylcellulose), gums (e.g., arabic, tragacanth, guar, etc.), and/or proteins (e.g., gelatin, collagen, etc.). Additional components in oral formulations may include coloring and or sweetening agents (e.g., glucose, sucrose, and mannitol) and lubricating agents (e.g., magnesium stearate), as well as enteric coatings (e.g., methacrylate polymers, hydroxyl propyl methyl cellulose phthalate, and/or any other suitable enteric coating known in the art). In some embodiments, disintegrating or solubilizing agents are included (e.g., cross-linked polyvinyl pyrrolidone, agar, alginic acid or salts thereof, such as sodium alginate). In some embodiments, the engineered GAA polypeptide are be combined with various additional components, including but not limited to preservatives, suspending agents, thickening agents, wetting agents, alcohols, fatty acids, and/or emulsifiers, particularly in liquid formulations. In some embodiments, the engineered GAA polypeptides are administered to subjects in combination with other compounds, molecules, and/or materials used in the treatment of Pompe disease, including but not limited to pharmacological chaperones, as well as any other suitable compounds. In some additional embodiments, the pharmaceutical composition is suitable for parenteral injection into a human. In some embodiments, the pharmaceutical composition comprises a pill, tablet, capsule, or gelcap that further comprises an enteric coating.

[0181] In some embodiments, the present invention provides engineered GAA polypeptides suitable for use in decreasing the concentration of glycogen in tissues. The dosages of engineered GAA polypeptide(s) administered to an animal depend upon the condition or disease, the general condition of the animal, and other factors known to those in the art. In some embodiments, the compositions are intended for single or multiple administrations to an animal. In some embodiments, it is contemplated that the concentration of engineered GAA polypeptide(s) in the composition(s) administered to an animal (e.g., a human with Pompe disease) is sufficient to effectively treat, ameliorate and/or prevent the symptoms of disease (e.g., Pompe disease and/or Pompe disease-related conditions, diseases and/or symptoms). In some embodiments, the engineered GAA polypeptides are administered in combination with other pharmaceutical and/or dietary compositions.

[0182] The foregoing and other aspects of the invention may be better understood in connection with the following non-limiting examples. The examples are provided for illustrative purposes only and are not intended to limit the scope of the present invention in any way.

EXPERIMENTAL

[0183] The following Examples, including experiments and results achieved, are provided for illustrative purposes only and are not to be construed as limiting the present invention.

[0184] In the experimental disclosure below, the following abbreviations apply: ppm (parts per million); M (molar); mM (millimolar), uM and μ M (micromolar); nM (nanomolar); mol (moles); gm and g (gram); mg (milligrams); ug and μ g (micrograms); L and l (liter); ml and mL (milliliter); cm

(centimeters); mm (millimeters); μm and μm (micrometers); sec. (seconds); min(s) (minute(s)); h(s) and hr(s) (hour(s)); U (units); MW (molecular weight); rpm (rotations per minute); °C (degrees Centigrade); CDS (coding sequence); DNA (deoxyribonucleic acid); RNA (ribonucleic acid); *E. coli* W3110 (commonly used laboratory *E. coli* strain, available from the Coli Genetic Stock Center [CGSC], New Haven, CT); DPBS (Dulbecco's phosphate buffered saline); LB (Luria-Bertani); TB (terrific broth); 4-MUGlu (4-methylumbelliferyl α -D-glucopyranoside); SD-Ura (single drop out medium without uracil); HPLC (high pressure liquid chromatography); SDS-PAGE (sodium dodecyl sulfate polyacrylamide gel electrophoresis); MU-Glu (4-methylumbelliferyl α -D-glucopyranoside); IPTG (isopropyl β -D-1-thiogalactopyranoside); PMBS (polymyxin B sulfate); FIOPC (fold improvements over positive control); PBMC (peripheral blood mononuclear cells); LB (Luria broth); MeOH (methanol); Axygen (Axygen, Inc., Union City, CA); Athens Research (Athens Research Technology, Athens, GA); ProSpec (ProSpec Tany Technogene, East Brunswick, NJ); Sigma-Aldrich (Sigma-Aldrich, St. Louis, MO); Ram Scientific (Ram Scientific, Inc., Yonkers, NY); Pall Corp. (Pall, Corp., Pt. Washington, NY); Millipore (Millipore, Corp., Billerica MA); Difco (Difco Laboratories, BD Diagnostic Systems, Detroit, MI); Molecular Devices (Molecular Devices, LLC, Sunnyvale, CA); Kuhner (Adolf Kuhner, AG, Basel, Switzerland); Microfluidics (Microfluidics Corp., Westwood, MA); Thermotron (Thermotron, Inc., Holland, MI); Cambridge Isotope Laboratories, (Cambridge Isotope Laboratories, Inc., Tewksbury, MA); Applied Biosystems (Applied Biosystems, part of Life Technologies, Corp., Grand Island, NY); Greiner Bio-One (Greiner Bio-One North America, Monroe, NC); Agilent (Agilent Technologies, Inc., Santa Clara, CA); Thermo Scientific (part of Thermo Fisher Scientific, Waltham, MA); Corning (Corning, Inc., Palo Alto, CA); Megazyme (Megazyme International, Wicklow, Ireland); Enzo (Enzo Life Sciences, Inc., Farmingdale, NY); GE Healthcare (GE Healthcare Bio-Sciences, Piscataway, NJ); Pierce (Pierce Biotechnology (now part of Thermo Fisher Scientific), Rockford, IL); Phenomenex (Phenomenex, Inc., Torrance, CA); Optimal (Optimal Biotech Group, Belmont, CA); and Bio-Rad (Bio-Rad Laboratories, Hercules, CA).

EXAMPLE 1

GAA Gene Acquisition and Construction of Expression Vectors

[0185] In this Example, GAA gene acquisition and expression vector construction are described. A synthetic gene coding for a WT human GAA (Uniprot ID P10253) with the native signal peptide removed was designed for optimized gene expression in *Saccharomyces cerevisiae* and fused to the yeast MFA signal peptide sequence (SEQ ID NO: 3383) to generate the gene sequence represented by SEQ ID NO:2, which was cloned into the yeast expression vector pYT-72, as previously described (See e.g. US Pat. Appln. Publn. No. 2017/0360900 A1). Recombination cloning and gene expression were performed in *S. cerevisiae* strain INVSc1. Directed evolution techniques were used to generate

libraries of gene variants from this plasmid construct (See e.g., U.S. Pat. No. 8,383,346 and WO2010/144103).

[0186] For secreted expression and transient transfection in mammalian cells, a chimeric GAA expression construct encoding the synthetic mouse IG signal peptide (residues 1-19 of Uniprot accession number: A0N1R5; SEQ ID NO: 3381) fused to a synthetic gene coding for the different GAA variants was generated as follows. In some embodiments, the synthetic GAA gene variants are based on a GAA sequence that is codon-optimized for yeast (SEQ ID NO: 3), while in some alternative embodiments, the synthetic GAA variants are based on a GAA sequence that is codon-optimized for mammalian (SEQ ID NO: 5) expression. Oligonucleotides containing restriction enzyme flanks to enable cloning into either the *Bam*HI/*Xho*I site or *Hind*III/*Xho*I site were used to amplify a fragment coding for the synthetic mouse IG signal peptide (SEQ ID NOS: 3381 and 3382) and the coding sequence for the mature form of GAA. For mammalian expression, the PCR product was ligated into the linearized vector pcDNA3.1(+) either at the *Bam*HI/*Xho*I site or *Hind*III/*Xho*I site (Invitrogen). Directed evolution was used to generate specific gene variants derived from SEQ ID NOS: 18 to 828, within the pcDNA3.1(+) plasmid construct (See e.g., US Pat. No. 8,383,346 and WO2010/144103, each of which is incorporated by reference in its entirety).

[0187] The pDH vector was generated to optimize for vector copy number, desired selection drug, and base pair length, to enable compatibility with library variant generation. For mammalian expression, the PCR product was ligated into the linearized vector pDH either at the *Bam*HI/*Xho*I site or *Hind*III/*Xho*I site. Directed evolution was used to generate specific gene variants (SEQ ID NOS: 829 to 3378) derived from SEQ ID NO: 20, within the pDH plasmid construct. A plasmid map for pDH containing WT GAA is shown in Figure 9 and the plasmid sequence is provided as SEQ ID NO: 3379. In addition, a plasmid map for pDH containing a beta-lactamase (bla) stuffer sequence is provided in Figure 10 and the sequence is provided as SEQ ID NO: 3380.

[0188] In some experiments, expression of GAA variants was performed using a linear PCR amplification product of the expression cassettes described above (i.e., a chimeric expression construct composed of the synthetic mouse IG signal peptide (residues 1-19 of Uniprot accession number: A0N1R5; SEQ ID NO: 3381) fused to a synthetic gene coding for the different GAA variants) in pcDNA3.1(+) or pDH. PCR amplification was performed with optimizations generally known by those skilled in the art with primer pair (SEQ ID NOS: 3387 and 3388) for pcDNA3.1(+) and primer pair (SEQ ID NOS: 3385 and 3386; or SEQ ID NOS: 3387 and 3388) for pDH. In some cases, phosphorothioate primers were used.

EXAMPLE 2

High-Throughput Growth of *Saccharomyces cerevisiae* and GAA Assays

[0189] In this Example, experiments involving high-throughput growth of cells producing GAA variants, and assays to determine GAA activity are described.

High-Throughput (HTP) Growth of *S. cerevisiae*

[0190] Yeast (INVSc1) cells transformed with vectors expressing GAA and GAA variants using the lithium acetate method as known in the art, were selected on SD-Ura agar plates. After 72 h incubation at 30° C, colonies were placed into the wells of AXYGEN® 1.1 ml 96-well deep well plates (Axygen) filled with 200 µl/well SD-Ura broth (2 g/L SD-Ura, 6.8 g/L yeast nitrogen base without amino acids [Sigma Aldrich]), 3.06 g/L sodium dihydrogen phosphate, 0.804 g/L disodium hydrogen phosphate, pH 6.0, supplemented with 6% glucose. The cells were grown for 20-24 hours in a Kuhner shaker (250 rpm, 30° C, and 85% relative humidity). Overnight culture samples (20 µL) were transferred into COSTAR® 96-well deep plates (Corning), filled with 380 µL of SD-ura broth supplemented with 2% glucose. The plates were incubated for 66-84 h in a Kuhner shaker (250 rpm, 30° C, and 85% relative humidity). The cells were then pelleted (4000 rpm×20 min), and the supernatants (conditioned media) stored at 4° C until analyzed.

HTP-Analysis of Supernatants

[0191] GAA variant activity was determined by measuring the hydrolysis of 4-methylumbelliferyl α-D-glucopyranoside (4-MUGlu). For the unchallenged assay, 50 µL of SD-URA conditioned media produced as described above was mixed with 50 µL of 1 mM 4-MUGlu in McIlvaine Buffer (McIlvaine, J. Biol. Chem., 49:183-186 [1921]), pH 4.5, in a 96-well, black, opaque bottom plate. The reactions were mixed briefly and incubated at 37 °C for 4-24 hours, prior to quenching with 100 µL of 0.5 M sodium carbonate pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX® M2 microplate reader (Molecular Devices), by monitoring fluorescence (Ex. 355 nm, Em. 460 nm).

HTP-Analysis of Supernatants Challenged with Neutral Buffer

[0192] GAA variants were challenged with neutral to near neutral (pH 6.5-7.5) buffer to simulate the pH that the variants encounter in the blood following their administration to a patient. First, 50 µL of GAA variants in SD-URA conditioned media and 50 µL of McIlvaine buffer (pH 6.5-7.4) were added to the wells of a 96-well round bottom plate. The plates were sealed and incubated at 37 °C for 1 h. Then, 50 µL of each challenged sample were mixed with 50 µL of 1 mM 4-MUGlu in McIlvaine buffer pH 4.4. The reactions were mixed briefly and incubated at 37 °C for 4-24 hrs, prior to quenching with 100 µL of 0.5 M sodium carbonate, pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX® M2 microplate reader (Molecular Devices) by monitoring fluorescence (Ex. 355 nm, Em. 460 nm).

HTP-Analysis of Supernatants Pretreated with Acid

[0193] GAA variants were challenged with acidic (pH 3) buffer to simulate the pH that the variants encounter in the lysosome following their administration to a patient. First, 50 µL of GAA variants in

SD-URA conditioned media and 50 μ L of McIlvaine buffer (pH 3) were added to the wells of a 96-well round bottom plate. The plates were sealed and incubated at 37 °C for 1 h. Then, 50 μ L of each challenged sample were mixed with 50 μ L of 1 mM 4-MUGlu in McIlvaine buffer pH 4.4. The reactions were mixed briefly and incubated at 37 °C for 4-24 hrs, prior to quenching with 100 μ L of 0.5 M sodium carbonate pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX® M2 microplate reader (Molecular Devices) monitoring fluorescence (Ex. 355 nm, Em. 460 nm).

HTP-Analysis of Supernatants with Glycogen Substrate

[0194] The hydrolytic activity of the GAA variants was determined by measuring the hydrolysis of glycogen to glucose. For the unchallenged assay, 50 μ L of SD-URA conditioned media produced as described above was mixed with 50 μ L of 5mM glycogen in pH 4.5 McIlvaine buffer (7.71 mL of 0.2 M Na₂HPO₄ and 12.29 of 0.1 M citric acid) in a 96-well, black, opaque bottom plate. The reactions were mixed briefly and incubated at 37 °C for 24 hours. After incubation, 20uL of the glycogen hydrolysis reaction mixtures were mixed with 80uL of AMPLEX® Red Glucose Assay Kit (Sigma) to determine the glucose content of each reaction. Hydrolysis was analyzed using a SPECTRAMAX® M2 microplate reader (Molecular Devices) by monitoring fluorescence (Ex. 544 nm, Em. 585 nm)

EXAMPLE 3

GAA and GAA Variants

[0195] GAA and GAA variant activity was determined by assaying the enzyme activity after a series of independent challenges. The results for all of the variants and the substitutions in each of the polypeptide sequences are reported in reference to SEQ ID NO: 2. These variants were tested for GAA 4-MUGlu activity (Unchallenged Activity FIOPC), after pH 6.5 incubation (pH 6.5 Stability Activity FIOPC), and for glycogen hydrolysis (Glycogen Activity FIOPC), as described in Example 2. Tables 3-1 and 3-2 provide the results of these assays.

Table 3-1. Activity of GAA Variants Under Various Conditions¹				
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	Unchallenged Activity FIOPC	pH 6.5 Stability Activity FIOPC	Glycogen Activity FIOPC
21/22	A276F	++	+++	+
23/24	A276Y	+++	+++	+
25/26	A418E/H499R			++
27/28	A418R		++	+
29/30	A437S	+	+	
31/32	A444T		++	+
33/34	A489R	++	++	+++
35/36	A547G	+	++	
37/38	A750P	+	+	+

Table 3-1. Activity of GAA Variants Under Various Conditions¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	Unchallenged Activity FIOPC	pH 6.5 Stability Activity FIOPC	Glycogen Activity FIOPC
39/40	A753T	+	+	+
41/42	C930R	+	+	+
43/44	C944G	+	+	+
45/46	C944R	+	+	+
47/48	D274G		++	
49/50	E137P	+	++	+
51/52	E463A		++	+
53/54	F27P	++	++	+
55/56	F27P/C944W	++	++	++
57/58	F27R		+++	++
59/60	G426R		++	+
61/62	G820E		++	+
63/64	I375E	++	++	+
65/66	K471Q/A478S	+	+	+
67/68	K471S	+	+	
69/70	K581G	+		+
71/72	K581T	++	++	++
73/74	K725N/V732I	+	++	
75/76	K88G	++	++	
77/78	K88S		++	
79/80	K895R		++	++
81/82	L109G/G842E	++	+++	+
83/84	L109P	+	+	+
85/86	L28P	++	+	+
87/88	L28R	++	+	++
89/90	L28S	++	++	++
91/92	L29T/A478T	+	++	++
93/94	L642M	++	++	+
95/96	L642Q	++	++	+
97/98	L642S	++	++	+
99/100	L670N		++	
101/102	L871E		++	+
103/104	L934R	++	++	++
105/106	M138A	+	+	++
107/108	N527R		++	+
109/110	Q107G		++	+
111/112	Q107P			+
113/114	Q110G		+++	+
115/116	Q110L		++	+
117/118	Q113S	+	+	++
119/120	Q247R		+	+
121/122	Q421S			+
123/124	R403W	+	+	+

Table 3-1. Activity of GAA Variants Under Various Conditions¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	Unchallenged Activity FIOPC	pH 6.5 Stability Activity FIOPC	Glycogen Activity FIOPC
125/126	R414P	+	+	++
127/128	R455V			++
129/130	R786P		++	+
131/132	R786Y		++	
133/134	R862G		++	+
135/136	S135A		++	+
137/138	S135Q	+	+++	++
139/140	S476A		++	+
141/142	S476H		++	
143/144	S668H	++	++++	
145/146	T148G	++	++	+
147/148	T148Y	+	+	
149/150	T150G	++	++	++
151/152	T278A	+	+	
153/154	T278G		++	+
155/156	T692Q	+	+	+
157/158	T897V		++	
159/160	V30G		+++	++
161/162	V30K	++	++	+
163/164	V30T	++	++	+
165/166	W610A	++	+	
167/168	W610G	++	++	
169/170	W610S	+	++	+

¹All activities were determined relative to the reference polypeptide of SEQ ID NO: 2. Levels of increased activity are defined as follows: "+" = 0.9 to 1.1; "++" > 1.1; "+++" > 2; and "++++" > 3.5.

Table 3-2. Activity of GAA Variants Relative to SEQ ID NO: 2¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	Unchallenged Activity FIOPC	pH 6.5 Stability Activity FIOPC	Glycogen Activity FIOPC
171/172	G280D/S402A/V536I/D928T			++
173/174	H191R/G280D/S402A/R414G/A444P/G465E/G842S/D928T	+	+	++
175/176	L275V/A281V/S671N	++	+	++
177/178	K106P/H191R/R414G/A444P/E522V/D928T/C944S	++	+	+
179/180	H191R/G280D/R414G/A444P/A489D/D500A/E522V/G842S/D928T/C944S	++	++	++
181/182	K106P/H191R/R414G/A489D/D928T/C944S	++	++	

Table 3-2. Activity of GAA Variants Relative to SEQ ID NO: 2¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	Unchallenged Activity FIOPC	pH 6.5 Stability Activity FIOPC	Glycogen Activity FIOPC
183/184	L275M/A281V/S402A/M431V/M507L/I518V/W610R/S668D	++	+	++
185/186	L275M/M507L/A547G/S668D/L669H/S671N	++	++	+
187/188	L275M/M431V/V638I	++	++	++
189/190	A281V/S402A/I518V/A547G/S668D	++	++	+
191/192	L275V/I518V/S671N	++	++	+
193/194	L275V/M431V/M507L/I518V/A547G/S668D/L669H/S671N	++	++	+
195/196	S402A/M431V/I518V/W610R	++	++	+
197/198	K106P/H191R/G280D/S402A/R414G/A444P/S727P	++	++	+
199/200	M431V/M507L/I518V/L669H/S671N	++	++	+
201/202	L275V/R377K/S402A/M507L/I518V/L669H/S671N/V715G	++	++	+
203/204	H191R/R414G/E522V/G842S/C944S	++	++	+
205/206	H191R/G280D/R414G/A489D/G842S/D928T/C944S	++	++	+
207/208	L275M/A281V/W610R/V638I/S668D/L669H	++	++	+
209/210	A196V/S402A/M431V/A547G/W610R/V638I	++	++	++
211/212	H191R/G280D/S402A/R414G/A444P/A489D/D500A/C944S	++	++	+
213/214	L275V/S402A/V638I/L669H/S671N	++	++	+
215/216	L29Q/L240I/A596P/S668D/I869L	++	++	++
217/218	K106P/G280D/S402A/R414G/A444P/A489D/S727P/C944S	++	++	++
219/220	L29Q/L240I/A596S/S668D/H700F/I744V/I869T	++	++	+
221/222	L218S/S668D/H700F/I869T	++	++	+
223/224	M507L/A547G/W610R	++	++	
225/226	A281V/M431V/M507L/I518V/A547G/W610R/V638I/S668D	++	++	++
227/228	H191R/G280D/R414G/A444P/A489D/E522V/S727P/C944S	++	++	+
229/230	L275V/S402A/M431V/I518V/W610R/V638I/L669H/S671N/P922L	++	++	
231/232	L29V/L218S/L240I/H700F/I869T	++	++	+
233/234	A547G/V638I/S668D	++	++	+
235/236	H191R/G280D/R414G/C944S	++	++	+
237/238	L275V/M431V/M507L/I518V/W610R/L669H/S671N	++	++	
239/240	L275M/S402A/M431V/M507L/A547G/S671N	++	++	
241/242	S402A/M431V/A547G/V638I/S671N	++	++	+
243/244	A281V/S402A/M507L/A547G/V638I/L669H/S671N	++	++	+

Table 3-2. Activity of GAA Variants Relative to SEQ ID NO: 2¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	Unchallenged Activity FIOPC	pH 6.5 Stability Activity FIOPC	Glycogen Activity FIOPC
245/246	L275M/A281V/M507L/A547G/L669H/S671N	++	++	
247/248	L275M/M431V/I518V/A547G/V638I/S668D	++	++	
249/250	A547G/W610R/V638I/S671N	++	++	+
251/252	L275V/M431V/M507L/A547G/W610R/V638I/S671N	++	++	+
253/254	L275M/S402A/M507L/A547G/W610R/S671N	++	++	
255/256	L275M/A281V/S402A/I518V/A547G/W610R/V638I/S671N	++	++	
257/258	A281V/S402A/I518V/A547G/W610R/V638I/S668D/L669H	++	++	
259/260	L275M/A281V/S402A/A547G/W610R/V638I/L669H/S671N	++	++	
261/262	L275M/M431V/I518V/W610R/V638I/L669H/S671N	++	++	+
263/264	L29Q/L218S/L240I/S668D/H700F/I744V/I869L	++	++	++
265/266	A281V/S402A/M507L/I518V/A547G/W610R/V638I/L669H/S671N	++	++	+
267/268	S402A/M431V/I518V/A547G/S671N	++	++	
269/270	L275V/A281V/S402A/M431V/I518V/A547G/W610R/L669H/S671N	++	++	
271/272	L224F/S402A/M507L/I518V/A547G/V638I/S668D	++	++	++
273/274	N180H/S402A/M507L/A547G/W610R/S671N	++	++	+
275/276	L275M/A281V/S402A/M507L/I518V/A547G/V638I/L669H/S671N	++	++	
277/278	K106P/T150S/T486E/Q749E/E793K/R883H/Q894G	++	+++	++
279/280	L275V/A281V/M431V/I518V/A547G/V638I/L669H/S671N	++	++	
281/282	L275V/S402A/A547G/W610R/V638I/L669H/S671N	++	++	
283/284	L275M/S402A/A547G/V638I/L669H/S671N	++	++	+
285/286	K106P/T150S/T486E/N527D/A750P/E793K	++	+++	++
287/288	S402A/M431V/I518V/A547G/W610R/S668D	++	++	
289/290	T150S/L218S/N527D/Q749E/E793K	++	+++	++
291/292	L275V/M507L/I518V/A547G/V638I/L669H/S671N	++	++	+
293/294	N180H/S402A/M431V/M507L/A547G/W610R/L669H/S671N/E793G	++	++	
295/296	L275V/M507L/I518V/A547G/W610R/V638I/S668D/L669H	++	++	+

Table 3-2. Activity of GAA Variants Relative to SEQ ID NO: 2¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	Unchallenged Activity FIOPC	pH 6.5 Stability Activity FIOPC	Glycogen Activity FIOPC
297/298	L275V/S402A/M507L/A547G/W610R/V638I/S668D/L669H	++	++	+
299/300	M507L/A547G/V638I/L669H/S671N	+++	++	+
301/302	K106P/T150S/N169S/L218S/Q749E/P800A	+++	+++	++
303/304	T150S/R414G/T486E/Q749E/A750P/E793K	+++	+++	++
305/306	K106P/T150S/L218S/R414G/T486E/L642F/A750P/E793K/R883H	+++	+++	++
307/308	N180H/L275M/S402A/I518V/A547G/W610R/V638I/L669H/S671N	+++	++	
309/310	T150S/L218S/R414G/Q749E/E793K	+++	+++	+
311/312	L275V/M507L/A547G/W610R/V638I/L669H/S671N	+++	++	+
313/314	T150S/R414G/T486E/N527D/A750P/Q894R	+++	+++	++
315/316	K106P/T150S/L218S/N527D/E793K/Q894G	+++	+++	++
317/318	T150S/N169S/L218S/R414G/N527D/E793K	+++	+++	++
319/320	K106P/T150S/L218S/E793K	+++	+++	++
321/322	K106P/T150S/R414G/Q749E/A750P/E793K/Q894R	+++	+++	++
323/324	K106P/T150S/N169S/N527D/Q749E/E793K/R883H	+++	+++	++
325/326	L275V/S402A/M507L/A547G/W610R/V638I/L669H/S671N	+++	++	+
327/328	K106P/T150S/L218S/Q749E/A750P/E793K	+++	+++	++
329/330	M431V/M507L/I518V/G541E/A547G/V638I/L669H/S671N	+++	+++	+
331/332	T150S/L218S/R414G/T486A/A750P/E793K	+++	+++	++
333/334	T150S/L218S/Q749E/A750P/E793K	+++	+++	++
335/336	K106P/T150S/N169S/L218S/R414G/T486E/Q894R	+++	+++	++
337/338	T150S/L218S/R414G/T486E/A750P/E793K/R883H	+++	+++	++
339/340	K106P/T150S/N169S/L218S/R414G/Q749E/E793K	+++	+++	++
341/342	T269N/L275M/M431V/I518V/A547G/V638I/S668D/L669H	+++	+++	
343/344	K106P/T150S/R414G/Q749E/E793K/Q894R	+++	+++	++
345/346	K106P/T150S/L218S/N527D/Q749E/A750P/E793K	+++	+++	++
347/348	K106P/T150S/N169S/Q749E/E793K/R883H/Q894R	+++	+++	++
349/350	K106P/T150S/L218S/E793K/Q894R	+++	+++	++
351/352	K106P/T150S/N169S/L218S/T486E/N527D/Q749E/E793K/Q894R	+++	+++	++
353/354	K106P/T150S/L218S/R414G/T486E/A750P/E793K/Q894R	+++	+++	++

Table 3-2. Activity of GAA Variants Relative to SEQ ID NO: 2¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	Unchallenged Activity FIOPC	pH 6.5 Stability Activity FIOPC	Glycogen Activity FIOPC
355/356	K106P/T150S/N169S/L218S/R414G/T486E/Q749E/E793K/R883H/Q894R	+++	+++	++
357/358	K106P/T150S/N169S/L218S/T486E/R883H	+++	+++	++
359/360	K106P/T150S/N169S/L218S/R414G/T486E/A750P/E793K/R883H/Q894R	+++	+++	++
361/362	G36R/K106P/T150S/L218S/N527D/A750P/R883H/Q894R	+++	+++	++
363/364	K106P/T150S/N169S/L218S/T486E/Q749E/R883H	+++	+++	++
365/366	K106P/N169S/V185G/L218S/R414G/Q749E/A750P/E793K	+++	+++	++
367/368	K106P/T150S/P245S/E793K/R883H/Q894R	+++	+++	+++
369/370	K106P/T150S/N169S/L218S/R414G/Q749E/A750P/E793K/R883H	+++	+++	++
373/374	T150S/L218S/Q749E/E793K	+++	+++	++
375/376	T150S/L218S/R414G/Q749E/A750P/E793K/Q894R	+++	+++	++
377/378	K106P/T150S/N169S/R414G/T486E/Q749E/A750P/R883H	+++	+++	++
379/380	K106P/T150S/N169S/L218S/R414G/T486E/E793K/R883H	+++	+++	++
381/382	K106P/T150S/L218S/T486E/N527D/Q749E/Q894R	+++	+++	++
383/384	K106P/T150S/N169S/L218S/R414G/T486E/N527D/Q894R	+++	+++	++
385/386	T150S/L218S/R414G/T486E/Q749E/A750P	+++	+++	++
387/388	K106P/T150S/L218S/R414G/Q749E/E793K/R883H	+++	+++	++
389/390	K106P/T150S/N169S/L218S/R414G/T486E/N527D/A750P/Q894R	+++	+++	++
391/392	N169S/T486E/A750P/E793K/R883H	+++	+++	++
7/8	T150S/T486E/A750P/R883H/Q894G	+++	++++	+++
393/394	K106P/T150S/L218S/R414G/N527D/Q749E/A750P/R883H	+++	++++	++
395/396	K106P/T150S/L218S/R414G/Q749E/A750P/E793K/R883H/Q894R	+++	+++	++
397/398	K106P/T150S/N169S/L218S/R414G/Q749E/E793K/R883H	+++	++++	++
399/400	K106P/T150S/L218S/T486E/E793K/R883H	+++	+++	++
401/402	K106P/T150S/L218S/V331A/R414G/T486E/N527D/D733E/Q749E/E793K	+++	++++	++
403/404	L275M/A281V/S402A/I518V/A547G/W610R/S668D/L669H/E887D	+++	+++	++
405/406	K106P/T150S/L218S/R414G/N527D/Q749E/E793K/R883H/Q894G	+++	+++	++
407/408	K106P/T150S/Q749E/E793K/R883H	+++	++++	+++
409/410	K106P/A112S/T150S/L218S/R414G/N527D/E793K/R883H	++++	++++	+++

Table 3-2. Activity of GAA Variants Relative to SEQ ID NO: 2 ¹				
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	Unchallenged Activity FIOPC	pH 6.5 Stability Activity FIOPC	Glycogen Activity FIOPC
¹ All activities were determined relative to the reference polypeptide of SEQ ID NO: 2. Levels of increased activity are defined as follows: ""+"" = 0.9 to 1.1; ""++"" > 1.1; ""+++"" > 2; and ""++++"" > 3.5				

EXAMPLE 4

GAA variants of SEQ ID NO: 8

[0196] In this Example, analysis of GAA variants derived from SEQ ID NO: 8 for improved GAA activity after a series of challenges are described. Directed evolution of the GAA encoded by SEQ ID NO: 8 was carried out by constructing libraries of variant genes. These libraries were then plated, grown, and screened for GAA 4-MUGlu activity after in the unchallenged, no pre-incubation, activity assay (Unchallenged Activity FIOPC), after pH 7 incubation (pH 7 Stability Activity FIOPC), or after pH 3 incubation (pH 3 Stability Activity FIOPC), as described in Example 2. The results are presented in Table 4-1.

Table 4-1. Activity of GAA Variants Relative to SEQ ID NO: 8 ¹				
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 8)	Unchallenged Activity FIOPC	pH 3 Stability Activity FIOPC	pH 7 Stability Activity FIOPC
411/412	A281V/M431V/A489R/S668D	+++	+++	+++
413/414	A418E	++++	++++	++++
415/416	A418E/A489R	+	++	+
417/418	A489R/L934R	+	+	
419/420	A489R/Q749E	+	+	
421/422	A489R/S671N/L934R	+	+	
423/424	A489R/V638I/L934R	++	++	++
425/426	F27P/A418E/A478T	+	+	+++
427/428	F27P/L28S/A489R	+	+++	++
429/430	F27R	+	+	++
431/432	K106P	++	+++	++
433/434	K106P/A489R/S671N	+++	++	+++
435/436	K106P/L218S/A281V	++	+++	++
437/438	K106P/L218S/R455V	+++	+++	+++
439/440	K106P/L218S/R455V/M507L/Q749E	+++	+++	+++
441/442	K106P/M138A/L218S/M431V/S671N/Q749E	+++		++
443/444	K106P/S671N/L934R	+		+
445/446	K106P/V638I	+	+++	+
447/448	L157M	+++	+++	+++
449/450	L218S	+	++	+
451/452	L218S/A281V	++	+++	++

Table 4-1. Activity of GAA Variants Relative to SEQ ID NO: 8 ¹				
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 8)	Unchallenged Activity FIOPC	pH 3 Stability Activity FIOPC	pH 7 Stability Activity FIOPC
453/454	L218S/A281V/M431V	+++	+++	+++
455/456	L218S/A281V/S671N	+		
457/458	L218S/M431V	+	+	
459/460	L218S/M431V/A489R/M507L/Q749E/L934R	+++	++	+++
461/462	L218S/M507L/L934R	+	+	+
463/464	L218S/M507L/Q749E	+++	++	+++
465/466	L218S/Q749E	+++	+++	+++
467/468	L218S/R455V	+	+	+
469/470	L218S/V638I/S671N	+++	+++	+++
471/472	L28S	+	++	+
473/474	L28S/L29T	+	+	+
475/476	L28S/L29T/A418E	++	+++	++
477/478	L28S/L29T/Q113S/S135Q/A418E	+	++++	+++
479/480	L28S/L29T/Q113S/S135Q/M138A	+	+	+
13/14	L28S/L29T/S135Q	+++	++++	+++
481/482	L29T/A478T	+		+
483/484	L29T/Q113S/M138A	+	+	+
485/486	L29T/Q113S/P126Q/S135Q/H193Q	+	+++	++
487/488	L29T/Q113S/S135Q	++	+++	++
489/490	L29T/Q113S/S135Q/R455V	+	+	+
491/492	L29T/T148G	+++	+++	+++
493/494	M138A/L218S/Q749E/L934R	+	+	
495/496	M138A/L218S/S668D/S671N	+++	++	+++
497/498	M138A/S671N/Q749E/L934R	+++	++	+++
499/500	M431V/S668D/S671N	+	+	+
501/502	M507L/S668D	+	+	+
503/504	M507L/S671N/L934R	+++	+++	+++
505/506	P4H	+	++	+
507/508	Q113S	+	++	+
509/510	Q113S/A418E/A478T/A489R/K581T	+	+	+
511/512	Q113S/A418E/R455V/A478T/K581T	+	+	+
513/514	Q113S/S135Q/A418E	++	+++	+++
515/516	Q345K/L934R	+	++	
517/518	Q749E/A784T	+++	+++	++
519/520	S135P/C944Y	+++	+++	+++
521/522	S135Q	+	++	+
523/524	S135Q/A478T/A489R/K581T	+		+
525/526	S135Q/A489R	+	+++	++
527/528	S135Q/T148G/S150G/A418E	+	+++	+
529/530	S671N/L934R	+++	++	+++

Table 4-1. Activity of GAA Variants Relative to SEQ ID NO: 8 ¹				
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 8)	Unchallenged Activity FIOPC	pH 3 Stability Activity FIOPC	pH 7 Stability Activity FIOPC
531/532	S671N/Q749E	+++	+++	+++
¹ All activities were determined relative to the reference polypeptide of SEQ ID NO: 8. Levels of increased activity are defined as follows: """" = 0.5-0.9; """" 0.9 to 1.1; """" > 1.1; """" > 2; and """" > 3.5.				

EXAMPLE 5

High-Throughput Growth of Adherent Mammalian Cells and GAA Assays Obtained Through Adherent Mammalian Expression

High-Throughput (HTP) Growth of GAA and GAA Variants in Adherent Mammalian Cells (HEK293T)

[0197] HEK 293T cells were transfected with a pcDNA 3.1(+) vector (ThermoFisher Scientific), pDH vector, or PCR-amplified linear DNA (as described in Example 1) encoding a synthetic mouse IG signal peptide (SEQ ID NOS: 3381 and 3382) fusion to wild-type GAA or GAA variants using the lipofection method with LIPOFECTAMINE[®] 3000 Reagent (ThermoFisher Scientific). HEK 293T cells were cultured in standard complete growth medium (DMEM with 10% fetal bovine serum [both from Corning]) and seeded into NUNC[®] Edge 2.0 96-well plate (ThermoFisher Scientific), at densities of 0.5×10^5 cells/well/250 μ L to adhere and grow for 24 hours at 37 °C, and in the presence of 5% CO₂ prior to lipofection-mediated transfection. Following transfection, cells were incubated for 24-96 hours to allow for expression and secretion of GAA variants into the conditioned media. Conditioned media (20-100 μ L) from the HEK293T transfection was then transferred into new 96-well plates for activity, stability or uptake into cell analysis.

HTP-Analysis of Supernatants

[0198] GAA variant activity was determined by measuring the hydrolysis of 4-methylumbelliferyl α -D-glucopyranoside (4-MUGlu). For the unchallenged assay, 20 μ L of HEK 293T conditioned media produced as described above was mixed with 50 μ L of 1.5 mM 4-MUGlu in McIlvaine Buffer (McIlvaine, J. Biol. Chem., 49:183-186 [1921]), pH 4.4, in a 96-well, black, opaque bottom plate. The reactions were incubated at 25 - 37 °C for 30 - 60 minutes with agitation at 400 rpm, prior to quenching with 100 μ L of 0.5 M sodium carbonate pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX[®] M2 microplate reader (Molecular Devices) or an ENVISION[®] microplate reader (Perkin Elmer) monitoring fluorescence (Ex. 355 nm, Em. 460 nm).

HTP-Analysis of Supernatants Challenged with Neutral Buffer

[0199] GAA variants were challenged with neutral buffer to simulate the pH that the variants encounter in the blood following their administration to a patient. First, 20 μ L of conditioned media

containing GAA variants from HEK 293T expression were combined with 100 μ L of DPBS buffer (pH 7.4) in a 96-well plate. The plates were sealed and incubated at 37 °C for 24-96 h. Next, 20 μ L of neutral-pH-challenged sample were mixed with 50 μ L of 1.5 mM 4-MUGlu in McIlvaine buffer, pH 4.4. The reactions were incubated at 37 °C for 180 minutes with agitation at 400 rpm, prior to quenching with 100 μ L of 0.5 M sodium carbonate pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX® M2 microplate reader or an ENVISION® microplate reader (Perkin Elmer) monitoring fluorescence (Ex. 355 nm, Em. 460 nm).

HTP-Analysis of Supernatants Challenged with Plasma

[0200] GAA variants were challenged with plasma to simulate the conditions that the variants encounter in the blood following their administration to a patient. First, 30 μ L of conditioned media containing GAA variants from HEK 293T expression were combined with 30 μ L of plasma (Innovative Research, Innovative Grade US Origin Monkey Cynomolgus Plasma K2 EDTA) in a 96-well plate. The plates were sealed and incubated at 37 °C for 2-4 h. Next, 10 μ L of plasma-challenged sample were mixed with 50 μ L of 1.5 mM 4-MUGlu in McIlvaine buffer, pH 4.4. The reactions were incubated at 25-37 °C for 15-60 minutes with agitation at 400 rpm, prior to quenching with 100 μ L of 0.5 M sodium carbonate pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX® M2 microplate reader or an ENVISION® microplate reader (Perkin Elmer) monitoring fluorescence (Ex. 355 nm, Em. 460 nm).

HTP-Analysis of GAA Activity in Lysates of Pompe Fibroblasts and C2C12 GAA Knockout Myoblasts

[0201] GAA variants produced in HTP were incubated with target cells and assayed for residual intracellular activity after 24 hours. For these experiments, mammalian cells lacking functional GAA activity were used, namely Pompe patient-derived fibroblasts (Coriell Institute for Medical Research # GM00248) and C2C12 myoblasts whose native GAA gene had been knocked out using Crispr-Cas9 editing. In these experiments, Pompe fibroblasts or C2C12 GAA knockout myoblasts were seeded into black, clear bottom, tissue culture treated COSTAR® 96-well plates (Corning, 3904) and allowed to grow to confluency in standard complete growth medium. Upon confluency, complete growth culture media was removed from the plates using an automated BIOMEK® i5 liquid handling robot. Conditioned media produced by HEK293T cells transiently transfected as described above, were transferred to Pompe patient-derived fibroblasts and C2C12 myoblasts, and allowed to incubate for 4-24 hours at 37°C, 5% CO₂. Medium was removed from the cultures using an automated BIOMEK® i5 liquid handling robot. The cells were briefly washed with 150 μ L 1xDPBS /well, and DPBS was removed using an automated BIOMEK® i5 liquid handling robot. Then, 200 μ L standard complete growth culture medium was added to each well, and the plates were returned to the incubator for 0-72 hours. At the conclusion of incubation, standard complete growth media was removed using an

automated BIOMEK® i5 liquid handling robot. The cells were washed with 150 µL 1xDPBS /well, and the DPBS removed using an automated BIOMEK® i5 liquid handling. The cells were lysed via addition of 50 µL of McIlvaine buffer, pH 4.4, supplemented with 0.2-0.5% TRITON X-100™ non-ionic surfactant (Sigma #93443) and agitation at room temperature for 30 minutes. Activity was assessed by addition of 50 µL of 1.5 mM 4-MUGlu in McIlvaine buffer, pH 4.4. The plates were sealed, incubated at 37 °C for 300-360 minutes with agitation at 400 rpm, prior to quenching with 100 µL of 0.5 M sodium carbonate, pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX® M2 microplate reader or an ENVISION® microplate reader (Perkin Elmer) monitoring fluorescence (Ex. 355 nm, Em. 460 nm). Cellular uptake FIOPC was calculated by dividing normalized GAA variant intracellular activity by the activity of the reference polypeptide with the indicated SEQ ID NO.

EXAMPLE 6

GAA Variants of SEQ ID NO: 16

[0202] In this Example, experiments for evolution and screening of GAA variants derived from SEQ ID NO: 16 for improved GAA activity after a series of challenges are described. The GAA synthetic gene coding for SEQ ID NO: 13 was fused to the synthetic mouse IG signal peptide polynucleotide (SEQ ID NO: 3381) (as described in Example 1) to generate the synthetic gene coding for SEQ ID NO: 15. Directed evolution of the GAA encoded by SEQ ID NO: 16 was carried out by constructing libraries of variant genes. These libraries were then plated, grown, and screened for GAA MU-Glu activity (“Unchallenged Activity FIOPC”), as well as after pH 7.4 pre-incubation (“pH 7.4 Stability and Activity FIOPC”), as described in Example 5. Variants were also tested for 4-MUGlu activity after lysis of Pompe fibroblasts (“Activity from Pompe Fibroblast Lysate FIOPC”) or GAA^{-/-} C2C12 cells (“Activity from C2C12 GAA^{-/-} Lysate FIOPC”), as described in Example 5. The results of these assays are presented in Table 6-1.

Table 6-1. Activity of GAA Variants Relative to SEQ ID NO: 16 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 16)	Unchallenged Activity FIOPC	pH 7.4 Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC	Activity from C2C12 GAA ^{-/-} Lysate FIOPC
533/534	V70A/R267K/K725E/C944S	+++	+++	+++	+++
535/536	R267K/A489D/D500A/K725E/Q830K/C930P	+++	+++	++	++
537/538	L109P/E522V/Q830K/C944S	+++	+++	+++	+++
539/540	V70A/R267K/C930P/C944S	+++	+++	+++	++
541/542	V70A/K725E/Q830K/L860F/C930P/C944S	+++	+++	+++	+++

Table 6-1. Activity of GAA Variants Relative to SEQ ID NO: 16¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 16)	Unchallenged Activity FIOPC	pH 7.4 Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC	Activity from C2C12 GAA-/- Lysate FIOPC
545/546	P39Q/R58L/A489D/K725E/Q830K/G842S/C930P/C944S	++	+++	++	++
547/548	A60V/D500A/S612D	+	+	+	+
549/550	P39Q/D500A/S612D	+++	+++	+++	+++
551/552	E522V/K725E	+++	+++	+++	+++
553/554	P39Q/V70A/L109P/Q830K/G842S	+++	+++	+++	+++
555/556	P39Q/V70A/K725E	+++	+++	+++	+++
557/558	P39Q/R267K/A489D/Q830K/C944S	+++	+++	++	+++
559/560	C930P	+++	+++	+++	+++
561/562	D500A/C930P/C944S	+++	+++	+++	+++
563/564	C944S	++	+++	++	++
565/566	L109P/E522V/S612D/K725E	+++	+++	++	++
19/20	P39Q/R267K/A489D/E522V/S612D/Q830K/G842S	+++	+++	+++	+++
567/568	P39Q/R267K	+++	+	+++	++
569/570	P39Q/V70A/A489D/S612D	+++	+++	+++	+++
571/572	L109P/S612D	++	+	+	+
573/574	D500A/L860F/C930P	+++	+++	++	++
575/576	R267K/E522V/K725E	++	++	+++	+
577/578	V70A/A489D/C930P	+++	+++	+++	+++
579/580	A489D/Q830K/C944S	+++	+++	+++	+++
581/582	D500A/S612D/Q830K/L860F	+++	+	+++	+++
583/584	P39Q/A489D/D500A/S612D	+++	+	++	+
585/586	P39Q	+++	+++	+++	+++
587/588	N528S/E793K	+		++	
589/590	K106R/K154R	+++	+		++
591/592	A62E/K106R/D523N/N528S/A696S/E793K/R862Q	+		+	
593/594	A196T/N528S/E681Q/I790V/E793K	+			
595/596	A62E/K154R/A696S/E793K/R862Q	+++	+	+	++
597/598	A62E/Q92R/I790V/E793K	++	+		++
599/600	E793K	+++	++	+	+++
601/602	S37F/N528S/I790V	+			
603/604	N528S/E681Q	+			
605/606	D523N/I790V/E793K	+++	++	+	++
607/608	D523N	++	+	+	++
609/610	N528S/R862Q	++			

Table 6-1. Activity of GAA Variants Relative to SEQ ID NO: 16 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 16)	Unchallenged Activity FIOPC	pH 7.4 Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC	Activity from C2C12 GAA-/- Lysate FIOPC
611/612	S37F/A62E	++	+	+	++
613/614	A753S	+++	+++		++
615/616	A62E/N79S/K154R/R862Q	+++			++
617/618	A62E/N79S/I790V/E793K	+++	+		+++
619/620	A62E/N79S/R862Q	+++	+	+++	+++
621/622	A62E/N79S/D523N/N528S/I790V	+++		+	
623/624	A62E/N79S/A196T/E681Q/R862Q	++		+	+
625/626	S37F/A196T	+			+
627/628	S37F/A62E/D523N	+++	+	++	++
629/630	S37F/N79S/K154R/E793K	+++	++		++
631/632	S37F/P64Q/R66G/N79S/K154R/D523N/E681Q/E793K/R862Q	+		+	
633/634	S37F/A62E/N79S/A196T/A696S/R862Q	+++	+	+	+
635/636	S37F/I790V/E793K	+			+
637/638	N79S/R862Q	+++	+	++	++
639/640	N79S/K154R/E681Q	+++		++	+
641/642	S37F/A62E/D523N/E793K	++	+	+	+
643/644	A62E/E793K/R862Q	+++	+++	+	+++
645/646	S37F/N528S/A696S/E793K	+			
647/648	S37F/N528S/I790V/E793K/R862Q	++			
649/650	A62E/Q92R	++	+	+	++
651/652	N79S/K154R/E793K/R862Q	+++	+	+	+
653/654	R862Q	++	+		++
655/656	L34D	+++	+++	+++	+++
657/658	Y352K	++	++	++	++
659/660	E207R	++	++	++	++
661/662	Q50L	+	+	++	+
663/664	L672K	++	+	++	+
665/666	Q938P	+++	+++	+++	+++
667/668	N875D	++	+	++	++
669/670	Q830V	+++	+	++	++
671/672	P178G	++		+	+
673/674	I22R	+++	++	+++	++
675/676	L24W	+++	++	++	++
677/678	L778Q	+++	+	+	+
679/680	E208I	++	++	++	+
681/682	K925A	++	+	++	++

Table 6-1. Activity of GAA Variants Relative to SEQ ID NO: 16 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 16)	Unchallenged Activity FIOPC	pH 7.4 Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC	Activity from C2C12 GAA-/- Lysate FIOPC
683/684	A47G	++	+	++	+
685/686	Y352V	++			
687/688	F27S	++	+	++	++
689/690	P149R	+++	+++	+++	+++
691/692	F27K	++	++	++	++
693/694	L672E	+++	+	++	+
695/696	S932A	++	++	++	++
697/698	L24R	+++	+++	+++	+++
699/700	L24E	++	++	++	+
701/702	C944R	+++	+++	++	+++
703/704	E740G	+++	++	+++	++
705/706	P55C	+	+	++	++
707/708	P673R	++			
709/710	F27G	++++	+++	+++	+++
711/712	Q49A	+	+	++	++
713/714	L34M	+++	++	++	++
715/716	E207Y	+	+	++	++
717/718	R68S	+++	++	+++	+++
719/720	P55L	++	++	++	+
721/722	V70Q	+++	+++	+++	+++
723/724	P39H	+++	+++	+++	+++
725/726	P922E	+++	++	+++	++
727/728	R68W	++	++	++	++
729/730	P77W	++	+	++	++
731/732	A774G	++	+	++	++
733/734	Q50G	++	++	++	+++
735/736	F27W	++	+	+	++
737/738	W118F	++	+	++	++
739/740	F27V	+++	+++	+++	+++
741/742	P178V	++	+	+	+
743/744	H424K	+++			
745/746	R385G	+++	+	+	+
747/748	F27G/M165I	+++	+++	+++	+++
749/750	V30D	+++	+++	+++	+++
751/752	V30L	+	++	+	+
753/754	V40W	++	+	++	+++
755/756	R68N	+++	+++	+++	+++
757/758	A179L	++	++	++	++
759/760	A774S	+++	+++	+++	+++
761/762	E33G	++	++	++	++
763/764	A902L	++	++	++	++
765/766	L109D	+++	+++	+++	++
767/768	E463A	+	+	+	+
769/770	F27A	+++	+++	+++	+++
771/772	E880R	++	+	++	++
773/774	T44I/L157V	+++	++	++	++

Table 6-1. Activity of GAA Variants Relative to SEQ ID NO: 16 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 16)	Unchallenged Activity FIOPC	pH 7.4 Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC	Activity from C2C12 GAA-/- Lysate FIOPC
775/776	T158E	+++	++	++	++
777/778	L157Q	++	++	++	++
779/780	K725V	++	+	++	+
781/782	Q217A	++	+++	+	+
783/784	Q938A	+++	++	++	+
785/786	D500A	+	+	++	+++
787/788	A89R	+++	+++	+++	+++
789/790	E844R	++	+	++	+
791/792	R448L	+			
793/794	Q107G	+++	+++	+++	+++
795/796	E208G	+	+	+	+
797/798	A47R	+++	+++	+++	+++
799/800	E740Q	++	+	++	++
801/802	F27R	++	++	++	++
803/804	L934F	+	+	+	+
805/806	K725F	++	+++	+	+
807/808	K925W	+	++	++	++
809/810	Q49G	+++	+++	+++	+++
811/812	Q50V	+++	++	+++	+++
813/814	Q217D	++	++	++	+++
815/816	Q892L	++	+++	++	++
817/818	L34T	+++	+++	+++	+++
819/820	A97G	+++	+++	+++	+++
821/822	P673N	+			
823/824	T158F	++	+	+	++
825/826	A97D	+++	+++	+++	+++
827/828	E33P	++	++	+++	++
371/372	P39D	+++	+++	+++	+++
543/544	H734K	+	+	++	+

¹All activities were determined relative to the reference polypeptide of SEQ ID NO: 16. Levels of increased activity are defined as follows: ""+"" = 0.5-0.9; ""++"" = 0.9 to 1.1; ""+++"" > 1.1; and ""++++"" > 2.

EXAMPLE 7

Production of GAA Variants

[0203] In this Example, production of GAA variants is described.

Production of GAA in EXPI293F™ Cells

[0204] Milligram-scale production of GAA variants was achieved by transient transfection of EXPI293™ cells (ThermoFisher Scientific) using the lipofection method with EXPIFECTAMINE™ 293 Reagent (ThermoFisher Scientific) in EXPI293™ Expression Medium (ThermoFisher Scientific). GAA variants fused to an N-terminal synthetic mammalian signal peptide (SEQ ID NO: 3381) were

subcloned into the mammalian expression vectors pLEV113, pcDNA 3.1(+), or pDH as described in Example 1. EXPI293™ cells were transfected with plasmid DNA and grown in suspension for 4-7 days. Conditioned media was then harvested, clarified by centrifugation and filtration and stored at 4 °C until analysis.

EXAMPLE 8

Purification of GAA Variants

[0205] In this Example, methods of purifying GAA variants are described.

Purification of GAA Variants From Mammalian Cell Supernatants

[0206] GAA variants (SEQ ID NOS: 4, 6, 10, 12, 16, 18, 20, 946, 1894, 1924, 1950, 1956, 1984, 2034, 2054, 2066, 2074, 2178, 2202, and 2496) produced in EXPI293F™ cells as described in Example 7, were purified from mammalian culture supernatant as described in the literature (Yasuda et al., Prot. Exp. Pur., 37:499-506 [2004]). Concanavalin A resin (Sigma Aldrich) was equilibrated with 0.1 M sodium acetate, 0.1 M NaCl, 1 mM MgCl₂, CaCl₂, and MnCl₂, pH 6.0 (Concanavalin A binding buffer). Supernatant was sterile-filtered with a 0.2µm bottle-top filter before it was loaded onto the column. After loading, the column was washed with 10 column volumes of Concanavalin A binding buffer and the bound protein was eluted with Concanavalin A binding buffer supplemented with 0.9 M methyl- α -D-mannopyranoside and 0.9 M methyl- α -D-glucopyranoside. Eluted protein was concentrated, and the buffer exchanged into storage buffer (20 mM sodium phosphate, 150 mM sodium chloride, 185 µM TWEEN®-20 non-ionic detergent, pH 6.0) using an AMICON® Ultra 15mL filtration unit with a 50 kDa molecular weight cut off membrane (Millipore). The GAA in storage buffer was sterile filtered through ANOTOP® 0.2µm syringe filters (Whatman), and stored at -80°C. Based on BCA quantitation (described below), the purification process produced 2.4-50 ng of purified protein/ml of culture supernatant.

Protein Quantification by BCA assay

[0207] A bicinchoninic acid (BCA) protein assay (Sigma Aldrich) was used to quantify purified GAA variants produced as described above. In a microtiter plate, 25 µL of protein standards and purified GAA with proper dilution were mixed with 200 µL of working reagent containing 50 parts of BCA reagent A and 1 part of BCA reagent B. The plate was thoroughly mixed on a plate shaker for 30 seconds and incubated at 37°C for 30 minutes. Absorbance was measured at 562 nm on a plate reader.

EXAMPLE 9

In vitro Characterization of GAA Variants

[0208] In this Example, experiments conducted to characterize GAA variants produced as indicated herein are described.

Kinetic Characterization of rhGAA and GAA variants

[0209] GAA variant activity was determined by measuring the hydrolysis of 4-methylumbelliferyl α -D-glucopyranoside (4-MUGlu). Purified enzyme produced as described above was mixed with 50 μ L of 0-2.5 mM 4-MUGlu in McIlvaine Buffer (McIlvaine, J. Biol. Chem., 49:183-186 [1921]), pH 4.5, in a 96-well, black, opaque bottom plate. The reactions were mixed briefly and incubated at 37 °C for 15-30 min, prior to quenching with 100 μ L of 0.5 M sodium carbonate pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX® M2 microplate reader monitoring fluorescence (Ex. 355 nm, Em. 460 nm), plotted and analyzed using the Michaelis-Menten equation. The results from this assay are presented in Figure 1.

Stability at Neutral pH of rhGAA and GAA variants

[0210] GAA variant stability to neutral pH was determined by incubating variants in 100 μ L of MEM complete growth medium (pH 7.4) in a 96-well plate. The plates were sealed and incubated at 37 °C for up to 48 h. Next, 10 μ L of neutral-pH-challenged sample were mixed with 50 μ L of 1.5 mM 4-MUGlu in McIlvaine buffer, pH 4.4 in a 96-well, black, opaque bottom plate. The reactions were incubated at room temperature for 30 minutes with agitation, prior to quenching with 100 μ L of 0.5 M sodium carbonate pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX® M2 microplate reader or an ENVISION® microplate reader (Perkin Elmer) monitoring fluorescence (Ex. 355 nm, Em. 460 nm) measuring the hydrolysis of 4-methylumbelliferyl α -D-glucopyranoside (4-MUGlu). The results from this assay are presented in Figure 11.

Cellular Uptake in Pompe Fibroblasts or C2C12 GAA Knockout Myoblasts of Purified GAA Variants Expressed in Expi293F Cells

[0211] The cellular uptake of GAA variants as compared to reference enzymes (SEQ ID NOS: 4 and 6), was determined to assess their overall ability to be endocytosed into cultured cells. Pompe fibroblasts (GM00248, Coriell Institute for Medical Research) or C2C12 GAA knockout cells were seeded into a black walled, clear bottom 96-well plate (Costar, #3603) in standard complete growth medium and allowed to reach confluency (2-3 days at 37 °C, 5% CO₂). After reaching confluency, standard complete growth medium was removed using an automated BIOMEK® i5 liquid handling robot. Enzymes purified as described in Example 8, were added to cells at 0-10 μ g GAA/mL in standard complete growth media and allowed to incubate for acute (1-6 hours) or extended (16-96 hours) treatments at 37 °C, 5% CO₂. Media containing GAA variants were aspirated using an automated BIOMEK® i5 liquid handling robot. The cells were briefly washed with 150 μ L 1xDPBS /well, and the DPBS was removed by an automated BIOMEK® i5 liquid handling robot. Then, 200 μ L

standard complete growth medium was added to each well, and the plates were returned to the incubator for 0-72 hours. At the conclusion of incubation, MEM complete growth media was removed using an automated BIOMEK[®] i5 liquid handling robot. The cells were washed with 150 μ L 1xDPBS /well, and the DPBS was removed using an automated BIOMEK[®] i5 liquid handling robot. Cells were lysed via addition of 50 μ L of McIlvaine buffer, pH 4.4, supplemented with 0.2-0.5% TRITON X-100[™] non-ionic surfactant (Sigma #93443) and agitation at room temperature for 30 minutes. GAA activity was assessed by addition of 50 μ L of 1.5 mM 4-MUGlu in McIlvaine buffer, pH 4.4. The plates were sealed, incubated at 37 °C for 300-360 minutes with agitation at 400 rpm, prior to quenching with 100 μ L of 0.5 M sodium carbonate pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX[®] M2 microplate reader monitoring fluorescence (Ex. 355 nm, Em. 460 nm). Cellular uptake FIOPC was calculated by dividing the normalized GAA variant intracellular activity by the control (WT) activity. Figures 4, 5, 8, 12, 13, 16-18, 22, 24, and 26, provide graphs showing the cellular uptake in Pompe fibroblasts of purified GAA variants after treatments of 1 to 96 hours. Figure 6, 7, 14, 15, 19-21, 23, 25, and 27 provide graphs showing the cellular uptake in C2C12 GAA KO myoblasts of purified GAA variants after treatments of 1 to 96 hours. Figure 8 provides a graph showing the cellular uptake and stability of GAA variants after 1-7 days.

Glycogen Degradation Activity of GAA Variants Expressed in EXP1293F[™] Cells

[0212] The relative glycogen hydrolysis activity of GAA variants was assessed using an AMPLEX[®] Red Glucose/Glucose Oxidase Kit (Invitrogen, #A22189) following the manufacturer's instructions with the modifications described herein. Briefly, GAA variants were diluted to an appropriate concentration range (0 – 2 μ g/mL) in 50 μ L reactions with 10 mM glycogen in McIlvaine buffer pH 4.4, in a COSTAR[®] black walled, clear bottom 96-well plate (#3603, Corning). The plates were sealed and gently shaken at room temperature for 30 minutes. Reactions were neutralized by addition of 25 μ L of 0.5 M sodium carbonate, pH 10.5. 50 μ L of the AMPLEX[®] Red/Horseradish Peroxidase/Glucose oxidase solution prepared as per manufacturer's instructions, was added to each well. The plates were sealed and gently shaken at room temperature for 30 minutes. Reactions were quantified to a standard curve of free glucose using a SPECTRAMAX[®] M2 microplate reader monitoring fluorescence (Ex. 530 nm, Em. 590 nm). The results are shown in Figure 2.

EXAMPLE 10

GAA Variants of SEQ ID NO: 20

[0213] In this Example, experiments for evolution and screening of GAA variants derived from SEQ ID NO: 20 for improved GAA activity after a series of challenges are described. Libraries of variant genes GAA encoded based off of by SEQ ID NO: 20 were constructed, plated, grown, and screened for GAA MU-Glu activity ("Unchallenged Activity FIOPC"), as well as after plasma challenge ("Plasma Stability and Activity FIOPC"), as described in Example 5. Variants were also tested for 4-

MUGlu activity after lysis of Pompe fibroblasts (“Activity from Pompe Fibroblast Lysate FIOPC”) or GAA^{-/-} C2C12 cells (“Activity from C2C12 GAA^{-/-} Lysate FIOPC”), as described in Example 5. The results of these assays are presented in Table 10-1 and 10-2.

Table 10-1. Activity of GAA Variants Relative to SEQ ID NO: 20 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC	Activity from C2C12 GAA ^{-/-} Lysate FIOPC
19/20		+	+	+	+
829/830	L24R/F27G/A89R/D500A/S842G		++	++	++
831/832	L24R/F27G/D500A/S842G	+	++	++	+
833/834	D500A/S932A		+	++	+
835/836	L24R/Q39H/S842G/S932A		+	+	++
837/838	Q39D/A89R/A97G/D500A		+	++	++
839/840	D500A/S842G		++	++	+
841/842	S842G/S932A			+	++
843/844	A89R/A97G/Q107G		++	++	++
845/846	A89R/A97G/D500A			++	++
847/848	F27G/Q39H/Q49G/A97G/D500A/S842G		++	++	++
849/850	R68N/A89R/A97G/S932A		+	++	+
851/852	A89R/S842G		+	++	+
853/854	L24R/Q39D/R68S/A89R/Q107G/D500A/S842G	+	++	++	++
855/856	L24R/F27G/D500A/S842G/S932A			++	++
857/858	L24R/Q39H/Q49G/A89R/A97G/S842G/S932A			++	++
859/860	L24R/S842G/S932A	+	+	++	++
861/862	L24R/A89R/D500A		+	++	++
863/864	L24R/Q39H/A89R/A97D/S842G/S932A			++	++
865/866	L24R	+	+	++	++
867/868	F27G/Q49G/R68S/D500A/S842G		+	++	++
869/870	Q107G/D500A/S842G		+	++	++
871/872	L34T/Q39D/D500A/S932A	+	+	++	++
873/874	R68S	+	+	++	+
875/876	L24W/L109D/D612S		+	++	++

Table 10-1. Activity of GAA Variants Relative to SEQ ID NO: 20 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC	Activity from C2C12 GAA-/- Lysate FIOPC
877/878	L24W/V70A/K267R/A774S		+	++	++
879/880	I22R/L24W/D612S/K725E/C944R	+	+	++	++
881/882	L24W			++	+
883/884	I22R/L24W/F27A/D489A		+	++	+
885/886	D489A			+	+
887/888	I22R/L24W/F27A/Q50V/Q107G/L109D/D489A/D612S/K725E		+	++	+
889/890	Q50V/D612S/C944S	+		++	+
891/892	D612S/K725E/C944S	++	++	++	
893/894	L109D/D612S/A774S/C944S	+	++	+	++
895/896	I22R/L24W/F27A/D612S/C944R	+	+	++	+
897/898	Q50V/D489A/A774S			++	++
899/900	Q50V/L109D/D489A/D612S			+	+
901/902	I22R/L24W/Q50V/L109D/K267R/D489A/A774S/C944R		+	++	++
903/904	I22R/L24W	+		++	++
905/906	L24W/Q50V/D489A/C944S		+	++	++
907/908	I22R/Q50V/L109D/K267R/D489A	+	++	++	+
909/910	L24W/D489A/C944R			++	++
911/912	L24W/K725E			++	+
913/914	L24W/L109D/C944S	+	+	++	+
915/916	D612S/P922E		+	+	+
917/918	L24W/K267R/K725E/C944S	+	+	++	++
919/920	Q107G/L109D				++
921/922	D489A/C944R	+	+	+	+
923/924	I22R/L24W/Q50V/K267R/D612S/P922E/C944R			++	++
925/926	I22R/L24W/F27A/D489A/D612S/A774S	+		+++	+

Table 10-1. Activity of GAA Variants Relative to SEQ ID NO: 20 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC	Activity from C2C12 GAA-/- Lysate FIOPC
927/928	L24W/F27A/Q107G/K267R/D612S/C944S	+		++	+
929/930	L24W/F27A/K267R/C944R			+	+
931/932	I22R/L24W/D489A			++	++
933/934	I22R/Q50V/Q107G/K267R/D489A/D612S/C944S	+	+	++	+
935/936	L24W/V70A/Q107G/L109D/D489A/D612S/K725E	+	+	++	++
937/938	I22R/L24W/Q107G/K267R/D489A/P922E	+		+++	++
939/940	L24W/Q107G/L109D/K267R/D489A/D612S/K725E/A774S	+	+	++	++
941/942	L24W/F27A/C944R	+	+	++	+
943/944	D612S			+	++
945/946	L24W/Q50V/D612S	+		++	++
947/948	I22R/K267R/D489A/D612S		+	++	++
949/950	L24W/F27A/Q50V/Q107G/K267R/A774S/C944S	+	+	++	+
951/952	L24W/C944S	+		++	++
953/954	E614W			++	+
955/956	N527R	+	+	++	+
957/958	E381W			++	+
959/960	R862Y	+		++	+
961/962	D923L			++	++
963/964	A62L	+	++	++	++
965/966	S916R			++	+
967/968	P914G			+	+
969/970	D523E	+		++	+
971/972	D923W			++	
973/974	A62F			+++	+
975/976	E188W			++	+
977/978	P71V			+	
979/980	P71W			++	+
981/982	L742V	+	+	++	+
983/984	N527V			+	+
985/986	P123V			+	+

Table 10-1. Activity of GAA Variants Relative to SEQ ID NO: 20 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC	Activity from C2C12 GAA-/- Lysate FIOPC
987/988	H193P	+	++	++	
989/990	P123L		++	++	+
991/992	P914Q			+	++
993/994	S124V			++	+
995/996	E858C			++	+
997/998	P57M			+	++
999/1000	S264M	+	+	++	+
1001/1002	T148K			+	
1003/1004	P57F	+	+	++	++
1005/1006	E614Q	+		++	++
1007/1008	A112H	++	+	++	+
1009/1010	R862Q	+	+	++	+
1011/1012	H48W			++	++
1013/1014	P914T			+	
1015/1016	S916G			++	+
1017/1018	K88R		+	++	+
1019/1020	P914K			++	++
1021/1022	G820A			++	+
1023/1024	V913W			++	+
1025/1026	D923V		++	++	++
1027/1028	P832A				+
1029/1030	P71L	+	++	++	+
1031/1032	A253M		++	+	++
1033/1034	L823F		+	++	+
1035/1036	L305F			++	+
1037/1038	P914S			+	+
1039/1040	S727W	+		++	++
1041/1042	L940Q	+	+	++	+
1043/1044	E188R			+	+
1045/1046	H48V	+		+	+
1047/1048	P832R		+	++	++
1049/1050	H48Q	+	+	+	
1051/1052	H193E	+	+	+	+
1053/1054	T148R	+		+++	++
1055/1056	G820V			++	
1057/1058	E381R			++	
1059/1060	S402N	+	+	++	++
1061/1062	L823V		+	++	++
1063/1064	R862M		+	++	+
1065/1066	K88L				+
1067/1068	A62W		++	++	++

Table 10-1. Activity of GAA Variants Relative to SEQ ID NO: 20 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC	Activity from C2C12 GAA-/- Lysate FIOPC
1069/1070	G108R	+	++	+	+
1071/1072	L748V		+	++	+
1073/1074	E42G		+	++	+
1075/1076	E858W		+	++	++
1077/1078	P57L	+	+	++	++
1079/1080	I333L			+	
1081/1082	V312A			+	
1083/1084	P197G	++	++	+	+
1085/1086	S916H	+	++	++	+
1087/1088	P914I			++	
1089/1090	N911R			++	
1091/1092	E381V			++	+
1093/1094	P71Y			++	+
1095/1096	R862I	+	++	++	++
1097/1098	S124M	+	+	++	+
1099/1100	P71G	+		+	+
1101/1102	P914R	+	+	++	++
1103/1104	S124G			++	+
1105/1106	N911G			+	+
1107/1108	E614R			++++	++
1109/1110	V913R	+	++	++	++
1111/1112	E937K		+	+++	++
1113/1114	E204A		+		
1115/1116	S402V/P781Q	+	++	++	+
1117/1118	V913G	+	+	++	+

¹All activities were determined relative to the reference polypeptide of SEQ ID NO: 20. Levels of increased activity are defined as follows: "+" = 0.9 to 1.1; "++" > 1.1; "+++" > 2; and "++++" > 3.5.

Table 10-2. Activity of GAA Variants Relative to SEQ ID NO: 20 ¹				
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC
19/20		+	+	+
1119/1120	K88G	++	++	+
1121/1122	L818V	+		++
1123/1124	T148E	+	+	+
1125/1126	A741D	+		+
1127/1128	N527P	+	+	++
1129/1130	L737M	++	++	++
1131/1132	G445L	+		

Table 10-2. Activity of GAA Variants Relative to SEQ ID NO: 20¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC
1133/1134	P197L	+	+	+
1135/1136	M260V	+	+	
1137/1138	A112W	+		++
1139/1140	H45M	+	+	+
1141/1142	G108S	+	+	++
1143/1144	L259N			+
1145/1146	P71Q	+	+	++
1147/1148	E42C			+
1149/1150	P57T	+	+	
1151/1152	P197I	+	+	+
1153/1154	L238Q	+		
1155/1156	I816L	+	+	++
1157/1158	E858V	+	+	+
1159/1160	S916A	+	+	+
1161/1162	P682A	+	+	+
1163/1164	M260F	+		+
1165/1166	H255Q	+		+
1167/1168	M432C	+	+	++
1169/1170	P71D	+		++
1171/1172	T325A	+		
1173/1174	E188Q/R377Q	+	+	
1175/1176	K106G	+	+	
1177/1178	K106N	++	+	+
1179/1180	L244I	+		+
1181/1182	A319F			+
1183/1184	I815M			+
1185/1186	F237L	+		+
1187/1188	H63A	++	+	+
1189/1190	P832E	+	+	+
1191/1192	A741T		+	+
1193/1194	S124F	++	+	++
1195/1196	T325V	+	++	++
1197/1198	P832V	++	+	+
1199/1200	P682K	++	++	+
1201/1202	H387R	+	+	+
1203/1204	R862G	+	+	++
1205/1206	H63K	+	+	+
1207/1208	S243R			+
1209/1210	H255R	+		+
1211/1212	E204G	+	+	++
1213/1214	V312F	+		+
1215/1216	S727T	++	++	++

Table 10-2. Activity of GAA Variants Relative to SEQ ID NO: 20¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC
1217/1218	L240I	++	++	++
1219/1220	L261T	+		
1221/1222	S264F	+	+	++
1223/1224	E614V	+		++
1225/1226	T91I			+
1227/1228	L240M	++	+	++
1229/1230	S727A	+	+	+
1231/1232	V913A	+		+
1233/1234	P832K		+	++
1235/1236	L940G	+		++
1237/1238	Q61V	+	+	+
1239/1240	R455T	+	+	+
1241/1242	H193S	++	+	++
1243/1244	M260R	+		
1245/1246	E204Q			+
1247/1248	S262L	+		+
1249/1250	T91K	+	+	++
1251/1252	S127G/P914F			+
1253/1254	R455V	++	+	
1255/1256	A62P	+	+	
1257/1258	E42Q	+	+	+
1259/1260	E614L	+	+	+
1261/1262	L305Y	+	+	++
1263/1264	S264L	+	+	+
1265/1266	P186G	+	+	
1267/1268	G820K	+	+	+
1269/1270	P832M	+	+	++
1271/1272	S264R	+	+	++
1273/1274	H63T	+	+	+
1275/1276	S262F			+
1277/1278	F237M	+		
1279/1280	K88V	+	+	+
1281/1282	S124R	+	+	+
1283/1284	Q61M	+	+	+
1285/1286	H193V	+	+	+
1287/1288	H63C	+		++
1289/1290	T148V	+	+	+
1291/1292	N911M			+
1293/1294	N527F	+		++
1295/1296	K88E	+	+	
1297/1298	L305G	+		
1299/1300	Q61G	+	+	++

Table 10-2. Activity of GAA Variants Relative to SEQ ID NO: 20¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC
1301/1302	S124D	+	+	++
1303/1304	F237V	+		
1305/1306	A253L	+	++	+
1307/1308	I815A			+
1309/1310	R862C	+	+	++
1311/1312	Q421R	+	+	++
1313/1314	H255T	+		
1315/1316	H63D	+	+	+
1317/1318	P832W	+	+	++
1319/1320	P57R	+	+	++
1321/1322	T148S		+	+
1323/1324	S262T	+	+	+
1325/1326	A62G			+
1327/1328	P914E	+	+	+
1329/1330	K88M	+	+	+
1331/1332	R862K	+	+	++
1333/1334	R455G	+		
1335/1336	S264C			+
1337/1338	K88I	+	+	
1339/1340	Y248A	+		
1341/1342	L305R	+	+	++
1343/1344	P279G	+		
1345/1346	A62S	+	+	++
1347/1348	R455H	+	+	+
1349/1350	L320M	+	++	++
1351/1352	L736V	+	+	+
1353/1354	A309C	+	+	+
1355/1356	E204D	+	+	+
1357/1358	S264A	++	++	++
1359/1360	H193R	+	+	++
1361/1362	E614G	+	+	++
1363/1364	K154R	+	+	+
1365/1366	K154E	+	+	+
1367/1368	T29S	+	+	++
1369/1370	Q421Y			+
1371/1372	P197V	+	+	
1373/1374	E188G	+	+	++
1375/1376	K88S		+	
1377/1378	Y248F	+	+	+
1379/1380	K88W		+	+
1381/1382	H45N	++	+	+
1383/1384	P914M	+	+	++

Table 10-2. Activity of GAA Variants Relative to SEQ ID NO: 20¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC
1385/1386	S727Q	+	+	+
1387/1388	P71A	+	+	+
1389/1390	P197T	+	+	+
1391/1392	T148A	+	+	
1393/1394	E858M	+		+
1395/1396	L823A	+	+	++
1397/1398	P57S	+	++	+
1399/1400	E204S	+		+
1401/1402	P186S	+		
1403/1404	K106V	+	+	+
1405/1406	L736W	+	+	
1407/1408	E614I	+		+
1409/1410	S916I	+	+	+
1411/1412	R862A	+	+	+
1413/1414	V913E	+	+	+
1415/1416	H63R	+	+	++
1417/1418	E937S	+	+	++
1419/1420	S262A	+	+	
1421/1422	P71R	+	+	++
1423/1424	E858R	+		
1425/1426	H63P	+	+	
1427/1428	L305V	++	++	
1429/1430	E42W	+	+	++
1431/1432	H48I	+	+	+
1433/1434	K154A	+	+	
1435/1436	Y248V	+		
1437/1438	A112F	+	+	+
1439/1440	L818T			+
1441/1442	A741C	+	+	+
1443/1444	P123Y	+	+	+
1445/1446	F556S	+	+	++
1447/1448	D523L	+		+
1449/1450	R862L	+		+
1451/1452	A62V	+	+	+
1453/1454	Q61P	+	+	+
1455/1456	H193A	+	+	
1457/1458	H255V	+		
1459/1460	L748I	+	+	+
1461/1462	M260W	++		++
1463/1464	R455L	+	+	+
1465/1466	I249L	+	+	+
1467/1468	G108N	+	+	++

Table 10-2. Activity of GAA Variants Relative to SEQ ID NO: 20¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC
1469/1470	Q421G	+	+	++
1471/1472	I816V	+	+	+
1473/1474	L818M	+	+	+
1475/1476	R195Y	+	+	
1477/1478	P832C			++
1479/1480	L748T	+		
1481/1482	N527W			++
1483/1484	T148R/V772I	+	+	++
1485/1486	K106T	+	+	
1487/1488	L240W/A374T		+	
1489/1490	F556Y		+	
1491/1492	P914F			+
1493/1494	P57Y	++	++	++
1495/1496	M432L	+	++	++
1497/1498	P123S	+	+	+
1499/1500	G108V	+	+	
1501/1502	P279R	+		
1503/1504	R195V	+	+	
1505/1506	E42D	+	+	
1507/1508	F205L	+	+	
1509/1510	H45S	+	+	++
1511/1512	Q421M			++
1513/1514	A112R	+	+	++
1515/1516	S916V	+	+	++
1517/1518	E381T	+		++
1519/1520	A253S	+		
1521/1522	F205V	+	+	+
1523/1524	E858S	+	++	+
1525/1526	E858G	+	++	++
1527/1528	E937Q	+	+	+
1529/1530	P914W			+
1531/1532	P197R	+	++	++
1533/1534	S264V	+	+	++
1535/1536	S727R			+
1537/1538	E858K		+	+
1539/1540	P832G	++	+	++
1541/1542	V913H			++
1543/1544	T148M	+		++
1545/1546	L252V	+		+
1547/1548	P123M	+	+	++
1549/1550	E937R			+
1551/1552	S243G			+

Table 10-2. Activity of GAA Variants Relative to SEQ ID NO: 20¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC
1553/1554	R455S			+
1555/1556	A741E			+
1557/1558	S243V			+
1559/1560	S402R	+	+	+
1561/1562	P832I	+	+	++
1563/1564	P57C	+	+	+
1565/1566	E614P			+
1567/1568	P279E	+		+
1569/1570	V913Q	+	+	++
1571/1572	P682W			++
1573/1574	E381G			++
1575/1576	N527S	+	+	+
1577/1578	E614Y	+	++	++
1579/1580	I816M			+
1581/1582	H45C	+	+	+
1583/1584	L259G	+		
1585/1586	S243E			+
1587/1588	E204V	+	+	+
1589/1590	P57N	+	+	+
1591/1592	H45Y			+
1593/1594	F237A			+
1595/1596	P71C	+		++
1597/1598	H48G	+	+	++
1599/1600	A253G	+		++
1601/1602	L736M	++	+	++
1603/1604	L940W	+		++
1605/1606	L823G			++
1607/1608	S262E	++	++	+
1609/1610	E188S	+	++	+
1611/1612	T325L	+	+	++
1613/1614	N527G			+
1615/1616	P197A	+	+	+
1617/1618	H193L	+	+	
1619/1620	S727G	+	+	++
1621/1622	F556R	+	+	++
1623/1624	L261E	+		+
1625/1626	A309G	++		++
1627/1628	E42Y			+
1629/1630	K154L	+		+
1631/1632	F556H	+	+	+
1633/1634	K106A	+	+	
1635/1636	S402G	++	++	++

Table 10-2. Activity of GAA Variants Relative to SEQ ID NO: 20¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC
1637/1638	H45T	++	++	+
1639/1640	E858P	+	+	++
1641/1642	S262P	+	++	+
1643/1644	P914H		+	++
1645/1646	P71E	++	++	+
1647/1648	P682L	+	+	+
1649/1650	E188T	+	++	+
1651/1652	L329F	+	++	++
1653/1654	S246G			+
1655/1656	H63M	+	++	+
1657/1658	H63L	+	+	+
1659/1660	R323A			+
1661/1662	T148G	++	++	++
1663/1664	S124C			++
1665/1666	R323L			+
1667/1668	P682V	+	++	++
1669/1670	G445Y	+		++
1671/1672	E858Y	+	++	++
1673/1674	P279L	+		++
1675/1676	R862N	++	++	+
1677/1678	P186A	+	+	
1679/1680	S262M	+	+	++
1681/1682	E937F			++
1683/1684	A741G	++	++	+
1685/1686	P123G	+	+	++
1687/1688	H45L		+	++
1689/1690	H63W	+	++	++
1691/1692	S246A	+		+
1693/1694	L940T	++	++	+
1695/1696	P57E	+	++	+
1697/1698	P123A	+	+	++
1699/1700	H63V	+	+	+
1701/1702	H255S	+		
1703/1704	H387L	+	++	+
1705/1706	K106W			+
1707/1708	P832S	++	++	++
1709/1710	S727C	++	+	+
1711/1712	S124L	++	++	++
1713/1714	P914V	++	++	++
1715/1716	S124N	+	+	+
1717/1718	L244V	+		++
1719/1720	R323M	+		++

Table 10-2. Activity of GAA Variants Relative to SEQ ID NO: 20¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC
1721/1722	S402D	+	+	+
1723/1724	S402M	+	+	++
1725/1726	L259M	+	+	++
1727/1728	D523W	+	+	+
1729/1730	R195G	++	++	+
1731/1732	Q421F			++
1733/1734	T148H	+	++	++
1735/1736	Y248R	+		
1737/1738	A62T	+	+	+
1739/1740	P279A	+		+
1741/1742	R455E		+	
1743/1744	P682G	++	++	+
1745/1746	E188M	+	++	++
1747/1748	P57Q	+	+	+
1749/1750	H193Y	+	+	+
1751/1752	Q421A	+	+	++
1753/1754	R195P	+	+	+
1755/1756	L320V	++	+	+
1757/1758	A253P	+	++	+
1759/1760	S298P/Q421R			++
1761/1762	H63G	++	++	++
1763/1764	A319M			++
1765/1766	P682F	+	+	++
1767/1768	H48R	+	+	++
1769/1770	I333V	+	+	++
1771/1772	A112P		+	+
1773/1774	Q61D	+		
1775/1776	R323K		+	+
1777/1778	P71F	+	+	+
1779/1780	V913L	+	+	+
1781/1782	E614S	+	++	+
1783/1784	M260L			++
1785/1786	L259S			+
1787/1788	L823R			++
1789/1790	L240Y			+
1791/1792	S402W	+		++
1793/1794	E937G			+
1795/1796	H63Y		+	++
1797/1798	I249V/S777N	+		++
1799/1800	E42M			+
1801/1802	E188V	+	+	+
1803/1804	S402L	+	+	+

Table 10-2. Activity of GAA Variants Relative to SEQ ID NO: 20¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Activity from Pompe Fibroblast Lysate FIOPC
1805/1806	A112V	++	++	++
1807/1808	G108H	+	++	++
1809/1810	P682R	++	+	+
1811/1812	M432I			+
1813/1814	Q61S	+	+	+
1815/1816	N527A	+	+	+
1817/1818	A741I			+
1819/1820	G251C			+
1821/1822	H63E	++	++	+
1823/1824	T91V			+
1825/1826	Q421P			+
1827/1828	E937L			+

¹All activities were determined relative to the reference polypeptide of SEQ ID NO: 20. Levels of increased activity are defined as follows: "+" = 0.9 to 1.1; and "++" > 1.1.

EXAMPLE 11**High-Throughput Growth of Suspension Mammalian Cells and GAA Assays Obtained Through Suspension Mammalian Expression****High-Throughput (HTP) Growth of GAA and GAA Variants in Suspension Mammalian Cells (Expi293F)**

[0214] EXPI293F™ cells (ThermoFisher Scientific) were transfected with pDH or PCR-amplified linear DNA (as described in Example 1) encoding a synthetic mouse IG signal peptide (SEQ ID NOS: 3381 and 3382) fusion to wild-type GAA or GAA variants using the lipofection method with EXPIFECTAMINE™ 293 Reagent (ThermoFisher Scientific) in EXPI293™ Expression Medium (ThermoFisher Scientific). EXPI293F™ cells (ThermoFisher Scientific) were cultured in EXPI293™ Expression Medium (ThermoFisher Scientific) and seeded into Axygen 1.1 mL deep well plate (Corning, P-DW-11-C-S), at densities of 1×10^6 cells/well/400 μ L. Cells were subjected to lipofection-mediated transfection and returned to a shaking incubator with 8% CO₂ and 70% humidity for 3-4 days to allow for expression and secretion of GAA variants into the conditioned media. Conditioned media was harvested by centrifugation of expression plates and transfer of conditioned media into a BioRad Hardshell PCR Plate (BioRad, HSP9601). Plates were centrifuged again and clarified conditioned media was transferred into new 96-well plates for activity, stability or uptake into cell analysis.

HTP-Analysis of Supernatants

[0215] GAA variant activity was determined by measuring the hydrolysis of 4-methylumbelliferyl α -D-glucopyranoside (4-MUGlu). For the unchallenged assay, 5-10 μ L of EXPI293F™ clarified conditioned media produced as described above was mixed with 50 μ L of 1.5 mM 4-MUGlu in McIlvaine Buffer (McIlvaine, J. Biol. Chem., 49:183-186 [1921]), pH 4.4, in a 96-well, black, opaque bottom plate. The reactions were incubated at 25 - 37 °C for 10 - 60 minutes with agitation at 400 rpm, prior to quenching with 100 μ L of 0.5 M sodium carbonate pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX® M2 microplate reader (Molecular Devices) or an ENVISION® microplate reader (Perkin Elmer) monitoring fluorescence (Ex. 355 nm, Em. 460 nm). Unchallenged activity FIOPC was calculated by dividing normalized GAA variant by the activity of the reference polypeptide with the indicated SEQ ID NO.

HTP-Analysis of Supernatants Challenged with Plasma

[0216] GAA variants were challenged with plasma to simulate the conditions that the variants encounter in the blood following their administration to a patient. First, 30 μ L of GAA variants in EXPI293F™ clarified conditioned media were combined with 30 μ L of plasma (Innovative Research, Innovative Grade US Origin Monkey Cynomolgus Plasma K2 EDTA) in a 96-well plate. The plates were sealed and incubated at 37 °C for 2-4 h. Next, 10 μ L of plasma-challenged sample were mixed with 50 μ L of 1.5 mM 4-MUGlu in McIlvaine buffer, pH 4.4. The reactions were incubated at 25-37 °C for 15-60 minutes with agitation at 400 rpm, prior to quenching with 100 μ L of 0.5 M sodium carbonate pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX® M2 microplate reader or an ENVISION® microplate reader (Perkin Elmer) monitoring fluorescence (Ex. 355 nm, Em. 460 nm). Plasma stability FIOPC was calculated by dividing normalized GAA variant activity following challenge by the activity of the reference polypeptide with the indicated SEQ ID NO following challenge.

HTP-Analysis of GAA Activity in Lysates of Pompe Fibroblasts and C2C12 GAA Knockout**Myoblasts**

[0217] GAA variants from HTP EXPI293F™ expression in clarified conditioned media were incubated with target cells and assayed for residual intracellular activity after 24-72 hours. For these experiments, mammalian cells lacking functional GAA activity were used, namely Pompe patient-derived fibroblasts (Coriell Institute for Medical Research # GM00248) and C2C12 myoblasts whose native GAA gene had been knocked out using Crispr-Cas9 editing. In these experiments, Pompe fibroblasts or C2C12 GAA knockout myoblasts were seeded into 96-well plates COSTAR® (3904, Corning) and allowed to grow to confluency in standard complete growth medium. Upon confluency, complete growth culture media was removed from the plates using an automated BIOMEK® i5 liquid handling robot. Clarified conditioned media from transient HPT transfections in EXPI293F™, were

transferred to Pompe patient-derived fibroblasts and C2C12 myoblasts, and allowed to incubate for 4-24 hours at 37°C, 5% CO₂. Medium was removed from the cultures using an automated BIOMEK® i5 liquid handling robot. The cells were briefly washed with 150 µL 1xDPBS /well, and DPBS was removed using an automated BIOMEK® i5 liquid handling robot. Then, 200 µL standard complete growth culture medium was added to each well, and the plates were returned to the incubator for 0-72 hours. At the conclusion of incubation, standard complete growth media was removed using an automated BIOMEK® i5 liquid handling robot. The cells were washed with 150 µL 1xDPBS /well, and the DPBS removed using an automated BIOMEK® i5 liquid handling. The cells were lysed via addition of 50 µL of McIlvaine buffer, pH 4.4, supplemented with 0.2-0.5% TRITON X-100™ non-ionic surfactant (Sigma #93443) and agitation at room temperature for 30 minutes. Activity was assessed by addition of 50 µL of 1.5 mM 4-MUGlu in McIlvaine buffer, pH 4.4. The plates were sealed, incubated at 37 °C for 300-360 minutes with agitation at 400 rpm, prior to quenching with 100 µL of 0.5 M sodium carbonate, pH 10.5. Hydrolysis was analyzed using a SPECTRAMAX® M2 microplate reader or an ENVISION® microplate reader (Perkin Elmer) monitoring fluorescence (Ex. 355 nm, Em. 460 nm). Cellular uptake FIOPC was calculated by dividing normalized GAA variant intracellular activity by the activity of the reference polypeptide with the indicated SEQ ID NO.

EXAMPLE 12

GAA Variants of SEQ ID NO: 946

[0218] In this Example, experiments for evolution and screening of GAA variants derived from SEQ ID NO: 946 for improved GAA activity after a series of challenges are described. Libraries of variant genes GAA encoded based off of by SEQ ID NO: 946 were constructed, plated, grown, and screened for GAA MU-Glu activity (“Unchallenged Activity FIOPC”), as well as after plasma challenge (“Plasma Stability and Activity FIOPC”), as described in Example 11. Variants were also tested for 4-MUGlu activity after lysis of Pompe fibroblasts treated for 24 hours (“Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC”) or GAA^{-/-} C2C12 cells (“Lysate Activity from Extended Treatment C2C12 GAA^{-/-} FIOPC”) or 6 hour treatments hours (“Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC”) or (“Lysate Activity from Acute Treatment C2C12 GAA^{-/-} FIOPC”), as described in Example 11. The results of these assays are presented in Table 12-1.

Table 12-1. Activity of GAA Variants Relative to SEQ ID NO: 946 ¹							
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 946)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
945/946		+	+	+	+	+	+
1829/1830	A62W/A89R/E188W/D489A/D500A/S727W/L823F			+++	++	+++	
1831/1832	S124V/D500A/S842G/R862Q	+		++	++	++	
1833/1834	W24R/D489A/D500A/S842G	+	+		++	+	++
1835/1836	W24R/Q39H/D489A/R862Q	++	+		++		
1837/1838	W24R/P57L/L823F/R862Q	+	+	+	+	++	
1839/1840	P57L	+	+	++	+	++	
1841/1842	W24R/A89R/D489A	+	+		+		
1843/1844	W24R/A89R/D489A/S727W/R862Q	+	+	++	++	+	
1845/1846	Q39H/P57L/A62W/E188W/D500A/S842G			+++	++	+++	
1847/1848	W24R/P57L/A62W/A89R/D489A/L823F/R862Q	+		++	++	++	
1849/1850	W24R/A62W/A89R/E188W/L823V/S842G/R862Q			+		++	

Table 12-1. Activity of GAA Variants Relative to SEQ ID NO: 946 ¹							
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 946)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
1851/1852	P57L/E188W/D489A/L823F/R862Q			++	++	+++	
1853/1854	Q39H/P57L/D500A/R862Q	+	+	++	++	++	
1855/1856	A62W/S124V/E188W/L823F/S842G/R862Q			++	++	+++	
1857/1858	P57F/A62L/L305F/D500A/V913R/S916G			+++	++	+++	
1859/1860	P57F/E614Q/S916G/S932A			++		++	
1861/1862	P57F/A62L/L305F/V913R			++	+	++	++
1863/1864	P57F/A62L/A437G/D500A/L761F/P914K/S916R		+	+++	++	+	
1865/1866	A62L/A437G/V913R/S916R		+	++	++		
1867/1868	A62L/D489A/E614Q/S916R			++	+	++	
1869/1870	P57F/A62L/V913R/S916R/S932A			+++	++	++	
1871/1872	A62L/N527R/S727W/S916G			+++			

Table 12-1. Activity of GAA Variants Relative to SEQ ID NO: 946 ¹							
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 946)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
1873/1874	P57F/A62W/L305F/A437G/D500A/S727W/V913R/S916R			+++	+++	++	
1875/1876	P57F/A437G/D500A/E614Q/S727W/P914R			+++	++		
1877/1878	D500A/S727W/V913R/S916R			+++	++	+	
1879/1880	E381V			++	+	+	
1881/1882	S124V/T148R/E381W/S727W/E858W/E937K			+++		++	
1883/1884	E381W/E858C/E937K			+++	++	++	
1885/1886	A62F/E614R			+++	++	+++	++
1887/1888	A89R/T148R/P149R/E381W	+		+++	++	++	++
1889/1890	A62F/P149R/E381V/E858W/E937K			+++	++	+++	
1891/1892	A62F/P149R/S727W			+++	++	+++	++
1893/1894	A89R/T148R/P149R	+	+	+++	++	+++	+++
1895/1896	A62F/A89R/S124V/T148R/E381W/E858W			++++	++	+++	++

Table 12-1. Activity of GAA Variants Relative to SEQ ID NO: 946 ¹							
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 946)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
1897/1898	A62F/A89R/T148R/E381V/E614R/E858W			+++	++	+++	++
1899/1900	A62W/S124V/P149R/E381V/P832R/E858C/E937K			+++	+	++	
1901/1902	S124V/P832R/E937K			+++	++	++	+
1903/1904	A62W/A89R/E381V	+	+	+++	++	++	+
1905/1906	A62W/P149R/E381W/E614R/E937K			+++	++	+++	+
1907/1908	S124V/E381W/E614R			+++	+	++	+
1909/1910	A62F/E96K/E614R			+++	+	++	
1911/1912	T148R/P832R/E858W/E937K			+++	++	+++	
1913/1914	A62F/A89R/E381V/D923W			+++	+	++	++
1915/1916	A62F/E381V			+++	+	+	
1917/1918	A62W/T148R/P149R/E381V/E858C/E937K			++++	++	++	+
1919/1920	A62W/T148R/E381W/E858C			+++	++	++	++
1921/1922	A62F/A89R/P149R/E381W/P832R			++++	++	++	+

Table 12-1. Activity of GAA Variants Relative to SEQ ID NO: 946 ¹							
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 946)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
1923/1924	A62W/A89R/P149R/E381W	+		+++	++	++	+
1925/1926	D923W			++			
1927/1928	A62W/A89R/S124V/E381W/E858C			+++	++	++	
1929/1930	E381V/E614R/P832R			+++	+	++	++
1931/1932	A62W/A89R/E381W/E858C			+++	+	++	++
1933/1934	A89R/E381W	+	+	++	++	++	
1935/1936	A62W/S124V/T148R/P149R/E381W/E614R			+++	+	++	+++
1937/1938	A62F/S124V/E381W/P832R			+++	+	++	
1939/1940	S19T/S124V/P149R/E381V/S727W			+++	+	++	
1941/1942	A62W/T148R/E381W/E614R/E937K			+++	++	++	+
1943/1944	A62F/A89R/T148R/D923W			+++	++	+++	++++
1945/1946	A62W/T148R/E381W/S727W			++++	++	++	+++
1947/1948	A62W/E381V			++	+		
1949/1950	A62F/A89R	+	+	+++	++	+	++
1951/1952	A62F/A89R/E858C			+++	++	++	++

Table 12-1. Activity of GAA Variants Relative to SEQ ID NO: 946 ¹							
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 946)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
1953/1954	A62W/A437G/D489A/E614Q/S727W/V913R			++	++	+	
1955/1956	A62L/A437G/D489A/N527R/S727W/V913R/S932A			+++	+++	++	+
1957/1958	P57F/A62W/L305F/D489A/Q907K/V913R/S916G			+++	++	+++	++
1959/1960	P57F/A62W/L305F/S916R			++	++	+++	++
1961/1962	P57F/A437G/D500A/N527R/S727W/S916R			+++	+++	++	++
1963/1964	A62L/L305F/S727W			++		+++	++
1965/1966	P57F/A62L/L305F/A437G/E614Q/A683S/V913R/S916R/S932A			+++	++	++	+
1967/1968	P57F/L305F		+	++	+	++	+
1969/1970	D489A/E614Q/S916G			++	+	++	
1971/1972	D489A/D500A	+	++	++	++	++	
1973/1974	P57F/A62W/A437G/V913R/S916G			++	++	++	++

Table 12-1. Activity of GAA Variants Relative to SEQ ID NO: 946 ¹							
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 946)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
1975/1976	P57F/A62W/L305F/D489A/V913R/S916G			+++	++	++	++
1977/1978	A62W/S727W			++		++	++
1979/1980	A62L/N527R			++	+	++	+
1981/1982	P57F/L305F/A437G/S916G		+	+++	++		
1983/1984	A62L/A437G/N527R/S727W	+	+	++++	+++		
1985/1986	P57F/A62L/A437G/N527R/S727W		+	++++	+++		
1987/1988	A62W/D489A/N527R/S916R/S932A			+++	+++		
1989/1990	A62L/D489A/D500A/S932A	+	++	++	++		
1991/1992	D500A/P914R/S916G		+	+++	++		
1993/1994	A62W/D500A		+	++	++		
1995/1996	A62L/L305F/A437G/D500A/S727W/V913R			+++	++	++	
1997/1998	A437G	+	+	+	+		
1999/2000	P57F/A62L/S916G	+	++	+++	++		

Table 12-1. Activity of GAA Variants Relative to SEQ ID NO: 946 ¹							
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 946)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
2001/2002	P57F/A62L/L305F/A437G/D500A/E614Q/S727W/S916R			++++	+++	++	+++
2003/2004	A437G/D489A/P914R/S916R	+	+	++	++		
2005/2006	P57F/A62W/F120I/N527R/V913R/S916R			++		++	++
2007/2008	A62L/A437G/N527R/S916G/S932A		+	+++	++		++
2009/2010	A437G/S727W/P914K		+	+++	++		
2011/2012	P57F/A437G/V913R/P914R	+	++	++	+++		++
2013/2014	A62W/S916G	+	++	++	+		
2015/2016	P57F/D489A/N527R/P914R/S916G			+++	++		
2017/2018	A437G/P914R/S916G		+	++	++		
¹ All activities were determined relative to the reference polypeptide of SEQ ID NO: 946. Levels of increased activity are defined as follows: "+" = 0.9 to 1.1; "++" > 1.1; "+++">> 2; and "++++">3.5.							

EXAMPLE 13

GAA Variants of SEQ ID NO: 1956

[0219] In this Example, experiments for evolution and screening of GAA variants derived from SEQ ID NO: 1956 for improved GAA activity after a series of challenges are described. Libraries of variant genes GAA encoded based off of by SEQ ID NO: 1956 were constructed, plated, grown, and screened for GAA MU-Glu activity ("Unchallenged Activity FIOPC"), as well as after plasma

challenge ("Plasma Stability and Activity FIOPC"), as described in Example 11. Variants were also tested for 4-MUGlu activity after lysis of Pompe fibroblasts treated for 24 hours ("Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC") or GAA^{-/-} C2C12 cells ("Lysate Activity from Extended Treatment C2C12 GAA^{-/-} FIOPC") or 6 hour treatments hours ("Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC") or ("Lysate Activity from Acute Treatment C2C12 GAA^{-/-} FIOPC"), as described in Example 11. The results of these assays are presented in Tables 13-1 and 13-2.

Table 13-1. Activity of GAA Variants Relative to SEQ ID NO: 1956¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA ^{-/-} FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA ^{-/-} FIOPC
1955/1956		+	+	+	+	+	+
2019/2020	Q795E	+		++	++	+	++
2021/2022	P78E/S372T/L390Q				++		
2023/2024	D87E/T266N/S372T/T483S		++	++	++		++
2025/2026	D87E/Q795E	+	+	+++	++	++	+++
2027/2028	T266N/S372T/T924N		++	+	++	++	++
2029/2030	P78E	+	++	++	+	+	
2031/2032	P78E/D87E/V536T	++	++	++	+++		
2033/2034	D87E/T266N/T483S	+	++	++	+++	+	++
2035/2036	D87E	+	+	++	++	+	++
2037/2038	P78E/D87E/K176T/T266N/V536T/Q615D	+	++	+	++	+	++
2039/2040	P78E/D87E/T266N/T483S/T924N		++	++	+++		++
2041/2042	D87E/S372T/S777G	+		+		++	+
2043/2044	T266N/S372T/V536T/Q615D/T763L/S777G						+
2045/2046	P78E/V536T/Q615D	+		++			

Table 13-1. Activity of GAA Variants Relative to SEQ ID NO: 1956¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/ FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/ FIOPC
2047/2048	P78E/T266N/T483S/V536T/Q615D	+	+		++		
2049/2050	T266N	+	++	++	+++	+	++
2051/2052	D87E/Q615D			++	+++	++	+
2053/2054	P78E/T266N/T763L	+	++	++	+++		
2055/2056	P78E/D87E/T266N/S372T/A386Y/S777G	+	+++	++	++		+
2057/2058	D87E/V536T/S777G	+	+	++		+	+
2059/2060	L390Q/Q615D	+		+		++	
2061/2062	P78E/T266N/T483S/Q795E	+	+++	++	+++		++
2063/2064	P78E/D87E/T483S/S777G	+	+				++
2065/2066	D87E/T266N	+	+++	+++	+++	++	+++
2067/2068	P78E/L390Q	+		++	+		
2069/2070	T266N/V536T/Q615D/Q795E	+	+	++	+++	+	+
2071/2072	D87E/T266N/T924N		++	++	++		+
2073/2074	P78E/D87E/T266N/S372T/V536T	+	+++	++	+++	+	++
2075/2076	S202K	+		++	+	++	++
2077/2078	T44G	+		+	++		++
2079/2080	S446T	+	+	++		++	++
2081/2082	Q344M	+	++	++	+	++	+++
2083/2084	L41I/S53M	+	+		++		++
2085/2086	L488M	+	+	+	++	+	++
2087/2088	Q615G			+	++	+	++
2089/2090	P779E	+		+		++	++

Table 13-1. Activity of GAA Variants Relative to SEQ ID NO: 1956¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
2091/2092	S777G	++	+	++		++	++
2093/2094	T141W	+				+	+
2095/2096	G65L	+			+	++	++
2097/2098	L857T	+				+	+
2099/2100	P779R	+	++	++	++	++	++
2101/2102	S202N	+		+		++	++
2103/2104	S678H	+	++			+	+
2105/2106	Q425R/S678I/G894C			++	+	++	+++
2107/2108	T543C	+		+	+	+	+
2109/2110	T543S	+	++	++	++	++	++
2111/2112	V859Y					+	+
2113/2114	P199V	+	++	++	++	++	++
2115/2116	G412Y	+				+	+
2117/2118	L909F	+	++	++	++	++	++
2119/2120	F679W	+	+				
2121/2122	G65V	+				+	+
2123/2124	P503S	+	+			++	++
2125/2126	G56L	+	++	+	++	++	+
2127/2128	G56A	+	+	+		++	+
2129/2130	L423V	+	+			+	
2131/2132	S777I	+		+	+	+	+
2133/2134	V797L				++		++
2135/2136	S678Y	++	++	+	+	+	++
2137/2138	N484L						++
2139/2140	Q768S	+		++	++	++	++
2141/2142	T145R	+	+	++		++	++
2143/2144	I156L					+	++
2145/2146	V797M				+	+	+
2147/2148	T44P/L347I					+	
2149/2150	E773V	++		++		++	++
2151/2152	A105T	+	++		+	+	+
2153/2154	L530V					+	++
2155/2156	T543Q	++		++		++	+
2157/2158	T543V	+		+	++	++	++
2159/2160	L488K	+	++	++	++	++	++
2161/2162	T826G						+
2163/2164	V859T	+		+	+	++	++
2165/2166	T141S	++		++	+	++	++

Table 13-1. Activity of GAA Variants Relative to SEQ ID NO: 1956¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/ FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/ FIOPC
2167/2168	S202L	+		+	++		++
2169/2170	S202R	+		+	+		+
2171/2172	A834H	+					+
2173/2174	L488G		++				
2175/2176	S777R	+		+	+		+
2177/2178	T44R	++	++	++	+	++	++
2179/2180	S136G					+	+
2181/2182	T44E	++	+	++	++	++	++
2183/2184	T145I	+		+	+	++	++
2185/2186	D726E						+
2187/2188	S202T	+		+	+	+	++
2189/2190	P779H					++	+
2191/2192	A496G	+		+	+		+
2193/2194	S678L	+	++			++	+++
2195/2196	G56R	+	++	+	++	++	+
2197/2198	D348G	+		++	++	++	++
2199/2200	Q344G		+				+
2201/2202	S678T	++	++	++	+++	++	++
2203/2204	T44L	++		++	+	++	++
2205/2206	L656V		+		++		+
2207/2208	L693F						+
2209/2210	V710N						+
2211/2212	V710L		+				
2213/2214	M663A						+
2215/2216	V710M	+					
2217/2218	L670I	+		++	+		+
2219/2220	L670F	+	++	+	++		+
2221/2222	L670R	+	+	++		+	++
2223/2224	V710S						+
2225/2226	F588L	+					
2227/2228	F705M		+				+
2229/2230	L670Q	+	+	+	+		
2231/2232	L670T	+	++	+	++		++
2233/2234	K154R/F588L	+					+
2235/2236	L693Y			+			
2237/2238	V710K		+	+			+
2239/2240	A708C			+	+		++
2241/2242	Y430F			+			+

Table 13-1. Activity of GAA Variants Relative to SEQ ID NO: 1956¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
2243/2244	I816L			++			
2245/2246	L656M			++			
2247/2248	L670W			+	+	++	++
2249/2250	L670V	+				+	
2251/2252	L693I			++	+		++
2253/2254	V589I/M663F	+	+		+	++	++
2255/2256	F588V			++			
2257/2258	M663F	+	+	++		+	+
2259/2260	M687L			++		+	++
2261/2262	L670E	+		+			+
2263/2264	L670S	++	++	++			+
2265/2266	L628M	+	+	+	++	+	+
2267/2268	V580I	++	++	++	++	++	++
2269/2270	V580L	+		+			
2271/2272	R583L	+		+		+	+
2273/2274	W3L/L569F	+	+			+	+
2275/2276	L629I	+				++	++
2277/2278	T692L	+	+	+		+	
2279/2280	A711C						+
2281/2282	R577T	+					
2283/2284	H499Y/A711F					+	
2285/2286	P503T	+	++				+
2287/2288	A690V			+			++
2289/2290	L628V						+
2291/2292	L569I	+	+				+
2293/2294	L691V		++				
2295/2296	T692V		++	++			
2297/2298	L569Y	+	++	++	+		++
2299/2300	I573Q	+	++				++
2301/2302	A711F		+	+	+		+
2303/2304	T692Y	+	+	+	++		++
2305/2306	L629C						+
2307/2308	T692I			+			
2309/2310	L157S			+			
2311/2312	S671A		+				+
2313/2314	R577M		+				+
2315/2316	L569S		++				+

Table 13-1. Activity of GAA Variants Relative to SEQ ID NO: 1956¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
2317/2318	I573M	+	++				+
2319/2320	R577V	+					
2321/2322	T692C	+	+				
2323/2324	I573H	+					
2325/2326	S671T	++	++				++
2327/2328	I573D	++					
2329/2330	S671M	+	++				
2331/2332	R577D	+	+				+
2333/2334	L629A		+			+	+
2335/2336	T692G	+	+	+		++	++
2337/2338	A711R	+	++	++	+		++
2339/2340	R577A	+	+				
2341/2342	L628I	+	++	++	+	+	++
2343/2344	T692F	+	+	+			++
2345/2346	R583G						+
2347/2348	R583K	+	+	+			+
2349/2350	V580Y						+
2351/2352	A572G						+
2353/2354	A711W			+			
2355/2356	L629G						+
2357/2358	I573C	+					+
2359/2360	L569V						+
2361/2362	L669R			++			
2363/2364	L579V						+
2365/2366	A572S	++	++	+			++
2367/2368	T692S	+	++	++			+
2369/2370	T692R	+	+	++			++
2371/2372	V580E	+	+	+			++
2373/2374	A574S		+				+
2375/2376	V631I		+	+			+
2377/2378	A711H	+	++			+	++
2379/2380	L691F	+		+	+		++
2381/2382	V580G	+					+
2383/2384	V580W	+					
2385/2386	R577E	++					+
2387/2388	R583C						+
2389/2390	A711L	+		+			
2391/2392	S671G			+			

Table 13-1. Activity of GAA Variants Relative to SEQ ID NO: 1956¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
2393/2394	A711G				+		+
2395/2396	V631L			++			+
2397/2398	L569T	+	++	++	+		++
2399/2400	L569H	++				+	+
2401/2402	L569Q	+					
2403/2404	V631M			++			+
2405/2406	H709S	+		+			++
2407/2408	T227A			+			+
2409/2410	V878R	+	+	++	+		
2411/2412	A578S	+					
2413/2414	T585L	+		++	+	+	+
2415/2416	L871E	+	+	+	+		
2417/2418	V222C	+		++	+		+
2419/2420	K581Y						+
2421/2422	V878G						++
2423/2424	T585V		+				
2425/2426	K581H						++
2427/2428	I877L		+	+			+
2429/2430	V868L		+				+
2431/2432	V878A	+	+		++	+	+
2433/2434	R873Y		+	+	+		+
2435/2436	H706V						+
2437/2438	K581G						+
2439/2440	V222P		+				+
2441/2442	T585M			+	+		++
2443/2444	T585Q		+		+		+
2445/2446	L633V			+			
2447/2448	A229C			++			
2449/2450	L871R						+
2451/2452	V878L	+	++	+			+
2453/2454	V868I		+	++			+
2455/2456	K581V						+
2457/2458	K581F						+
2459/2460	T585F		+				+
2461/2462	V878K			++			+
2463/2464	R873A	+	+	+			
2465/2466	K581L	+	++				+
2467/2468	K581T	+	+				

Table 13-1. Activity of GAA Variants Relative to SEQ ID NO: 1956¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
2469/2470	I869L	+	+	++	+		+
2471/2472	K581S	+	++				+
2473/2474	V878S	++	++		++	++	++
2475/2476	H706M		++				+
2477/2478	V878Q	++	++	++			
2479/2480	R873F			+			
2481/2482	I877V	+	++	+			
2483/2484	V878F	++	++	+	+		+
2485/2486	I869S	+	++	+			
2487/2488	V878W	++	++				+
2489/2490	H706F	+	+	++			
2491/2492	N225D		+	++			
2493/2494	L871K	++	++		+	++	+
2495/2496	P78E/D87E/T266N	+	+	+	+	+	+

¹All activities were determined relative to the reference polypeptide of SEQ ID NO: 1956. Levels of increased activity are defined as follows: "+" = 0.9 to 1.1; "++" > 1.1; and "+++" > 2.

Table 13-2. Activity of GAA Variants Relative to SEQ ID NO: 1956¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC
2497/2498	A142G	+		+	+
2499/2500	P788N		+		+
2501/2502	D493L		++	++	++
2503/2504	P788L		++		
2505/2506	S612T	+	++	+	+
2507/2508	W265H				+
2509/2510	K129E				+
2511/2512	K752N	+	+++		
2513/2514	H499E	+			++
2515/2516	V185L		++		
2517/2518	L857E	+			
2519/2520	K752S	+	+		
2521/2522	G56S	++	+	++	++

Table 13-2. Activity of GAA Variants Relative to SEQ ID NO: 1956 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC
2523/2524	T145L		++		+
2525/2526	T44F	+			+
2527/2528	T543K	+			++
2529/2530	P788T		+		+
2531/2532	K176R	+	++		
2533/2534	V859A	++	+		+
2535/2536	K129V	++	++		++
2537/2538	T44A	++		++	
2539/2540	P779S	+		++	
2541/2542	L857A	+			++
2543/2544	S202Q	+	+	+	
2545/2546	F354S	++	++		
2547/2548	H499I				+
2549/2550	S446K	+			
2551/2552	K752G	+	++		
2553/2554	G65F		+		
2555/2556	P503N	+	++		+
2557/2558	T44W	+		++	+
2559/2560	A834S	+	+	++	+
2561/2562	S446D	+			
2563/2564	Q615M	+	+	+	
2565/2566	L857V	++			
2567/2568	F152W	+	+	+	
2569/2570	F152S				+
2571/2572	S136K		++		
2573/2574	K129S	+	++		
2575/2576	T373S	+	++		
2577/2578	G822R	+	++		+
2579/2580	T826M	+			
2581/2582	G936S	+			+++
2583/2584	F679Y	+	++		
2585/2586	H63N	++	+		
2587/2588	V931L	+	+		
2589/2590	I156C	+	++		
2591/2592	H499V				++
2593/2594	P788A	+	+		
2595/2596	K752W				++
2597/2598	W730L			++	+++++
2599/2600	P788Q		+		
2601/2602	H499M	+		++	

Table 13-2. Activity of GAA Variants Relative to SEQ ID NO: 1956 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC
2603/2604	S202D	+	++		
2605/2606	T543L	++	++	++	
2607/2608	P199A	++	+	++	
2609/2610	S612G				+++
2611/2612	S856A	++		+	
2613/2614	K129W			++	++
2615/2616	I156R	+	+		
2617/2618	S136V	++		++	
2619/2620	F354L	++		++	
2621/2622	P186H	+			
2623/2624	Y677T	+			
2625/2626	P199W	++		+	
2627/2628	S136R	+	++	+	
2629/2630	K129T	++	++	+	
2631/2632	P337H				+++
2633/2634	I469M	++		++	
2635/2636	H499A	+			+++
2637/2638	Q344C			++	++++
2639/2640	D177Q			+	++
2641/2642	S202Y		++		
2643/2644	A105V	+	++		
2645/2646	E773P	++	++	++	++
2647/2648	W730K				+
2649/2650	L857R	+			+
2651/2652	A834W		++		
2653/2654	L488C	+	+++	+	+
2655/2656	L860S	+		++	++
2657/2658	L488E	+	+++		
2659/2660	L649M		+		
2661/2662	K267E				+
2663/2664	K752L	++	++		
2665/2666	S678R		+		
2667/2668	E855G	+			
2669/2670	N484A		++		
2671/2672	P788I	+	+		++
2673/2674	D348E	++	++	+	
2675/2676	L488S	++	+++		+
2677/2678	Q615S	+	+		
2679/2680	V859G	++	++	+	++
2681/2682	T141K			+	

Table 13-2. Activity of GAA Variants Relative to SEQ ID NO: 1956 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC
2683/2684	Q768I		++	++	+++
2685/2686	V926T	+	+	+	
2687/2688	P788H	+	++	+	+
2689/2690	S446C	+	++	+	
2691/2692	P779I	+	++	+	
2693/2694	V797F	+	++	++	
2695/2696	S372D	+			++
2697/2698	V350F				+
2699/2700	Y765W				++
2701/2702	T44V	++		++	++
2703/2704	T373A	+			
2705/2706	G936N	++			+
2707/2708	S446I	+			
2709/2710	S678Q	+		+	++
2711/2712	P526L			++	
2713/2714	P199G	+	++	+	
2715/2716	P199R	+		+	
2717/2718	L187I	+		+	
2719/2720	K267R	+		++	++
2721/2722	I469T	++			
2723/2724	T826I	++	++	++	+++
2725/2726	S856G	++	+	++	+++
2727/2728	P503C	+	++	++	
2729/2730	V797I	+	++		
2731/2732	G412W	+		+	++
2733/2734	S202A	++	++	++	+
2735/2736	T145A	+	+	+	
2737/2738	S777W	+		++	++
2739/2740	G412R	+		+	++
2741/2742	D348W				++
2743/2744	P199T	++	+	++	++
2745/2746	W265D				+++
2747/2748	K267G			+	++++
2749/2750	W265F	+	+		
2751/2752	H499Q	++	+		+
2753/2754	E855L	+			++
2755/2756	A496M	+	+	++	++
2757/2758	S202G	++		++	++
2759/2760	Q768V				++
2761/2762	T543H	++	++	++	++

Table 13-2. Activity of GAA Variants Relative to SEQ ID NO: 1956 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC
2763/2764	Y125W		++		
2765/2766	N484K	+	+		++
2767/2768	N484R				++
2769/2770	Q768K			+	++
2771/2772	S612L				+++
2773/2774	A834V	+			++
2775/2776	S678V	+	++	++	++
2777/2778	V926M	+	++	+	++
2779/2780	G139E				++
2781/2782	Y125H			++	++
2783/2784	P503H	+		+	++
2785/2786	V797R			+	+++
2787/2788	F152L	++		++	
2789/2790	W730R			++	+
2791/2792	T924A	+		++	+
2793/2794	V797W				++
2795/2796	V797E		++		+
2797/2798	T141R	+	+	+	++
2799/2800	S678W	++		++	++
2801/2802	G65Y	+		++	+
2803/2804	G936R	++			
2805/2806	T44Y	++	+	++	++
2807/2808	P788S	+		+	
2809/2810	P199V/L775I	++	+	++	++
2811/2812	S777M		+	++	++
2813/2814	I156K				++
2815/2816	P526V	+	+		
2817/2818	G56W	++	++	++	
2819/2820	P199I	+	+	+	
2821/2822	D401G	+			
2823/2824	G412S	++		++	
2825/2826	A834G	+		++	++
2827/2828	P788Y	+	++	+	+++
2829/2830	S612R	+			++
2831/2832	S202H	+	+		+
2833/2834	K129I	+	++		++
2835/2836	D401S	+			+
2837/2838	A496W		++		+
2839/2840	T543R	+		+	+++
2841/2842	S446G	+			+

Table 13-2. Activity of GAA Variants Relative to SEQ ID NO: 1956 ¹					
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC
2843/2844	I156S		+		++
2845/2846	G65R	+	++	++	++
2847/2848	G65A	+		+	++
2849/2850	P779M	++	+	++	++
2851/2852	K752F	++	++		
2853/2854	D162T		+		
2855/2856	I469V	++		+	++
2857/2858	S53I	+		+	++
2859/2860	A105W	++	+	++	++
2861/2862	L857S	+			
2863/2864	T543G	++			++
2865/2866	V350I	++		+	++
¹ All activities were determined relative to the reference polypeptide of SEQ ID NO: 1956. Levels of increased activity are defined as follows: "+" = 0.9 to 1.1; "++" > 1.1; "+++" > 2; "++++" > 3.5; and "+++++" > 5.					

EXAMPLE 14

GAA Variants of SEQ ID NO: 2496

[0220] In this Example, experiments for evolution and screening of GAA variants derived from SEQ ID NO: 2496 for improved GAA activity after a series of challenges are described. Libraries of variant genes GAA encoded based off of by SEQ ID NO: 2496 were constructed, plated, grown, and screened for GAA MU-Glu activity ("Unchallenged Activity FIOPC"), as well as after plasma challenge ("Plasma Stability and Activity FIOPC"), as described in Example 11. Variants were also tested for 4-MUGlu activity after lysis of Pompe fibroblasts treated for 24 hours ("Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC") or GAA^{-/-} C2C12 cells ("Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC") or 6 hour treatments hours ("Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC") or ("Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC"), as described in Example 11. The results of these assays are presented in Table 14-1 and Table 14-2.

Table 14-1. Activity of GAA Variants Relative to SEQ ID NO: 2496 ¹							
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/FIOPC
2495/2496		+	+	+	+	+	+
2867/2868	L569T/L628M/T692Y	++	+		++	++	+
2869/2870	L569Y/L670T/A711H	++		+	++	+	+
2871/2872	L569Y/L628M/L670T/T692Y/A711H/L871S	++	+	+	++	++	+
2873/2874	V589I/L670T/Q795E/L871S				+	+	+
2875/2876	L628M/L670F	+		++	++	++	++
2877/2878	L569H/L628M/L670T	++	+	+	++	+	++
2879/2880	L569T/L670T/A711H/L871E	++	+		++	++	++
2881/2882	L628M/L670T/T692G/A711H/Q795E	++		++	++	++	++
2883/2884	L569Y/L628M/L670T/A711H	++	+	+	++	++	++
2885/2886	L569Y/V589I/L628M/T692G/A711H/Q795E	+	+	+	++	++	++
2887/2888	L569H/T692G	+			+	+	+
2889/2890	L569Y/V589I/L670T/S678T/T692G/A711H/Q795E	++	++	++	++	++	++

Table 14-1. Activity of GAA Variants Relative to SEQ ID NO: 2496¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/FIOPC
2891/2892	L569T/L871K	++	++	+	++	++	++
2893/2894	L569Y/V589I/L628M/L670T/T692Y/A711H	++	++	+	+	++	+
2895/2896	V589I/L871E	+				+	
2897/2898	L670T/S678T/T692Y/L871S		+	+	+	+	++
2899/2900	L628M/A711H/Q795E	++		+	++	++	++
2901/2902	T692G/A711H	+		++	+	++	++
2903/2904	V589I/L670T/T692G/Q795E/L871K			+	+	+	++
2905/2906	L569Y/V589I/A711H/L871K	+	+		++	+	++
2907/2908	L670T/T692G/L871K	++		+	++	++	+
2909/2910	L569H/T692Y	+			+	+	+
2911/2912	L569H/L628M/L670T/T692Y/A711H/L871K	++	+	++	++	++	+
2913/2914	L670T/S678T	++		+	++	++	++
2915/2916	L569T/L670T/S678T/T692G/L871K	++	++	++	++	++	++
2917/2918	V589I						++

Table 14-1. Activity of GAA Variants Relative to SEQ ID NO: 2496 ¹							
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/FIOPC
2919/2920	L569H/V589I/L628M/L670T/S678T/T692G/A711H/Q795E/L871S	++	++	++	+	++	++
2921/2922	L569H/L670F/S678T/T692G/L871S	+	++	+	+	+	++
2923/2924	L569T/V589I/L628M/L670F/T692G/A711H	++	++	++	++	++	++
2925/2926	L569H/L670T/T692G	++	+				+
2927/2928	L569Y/L871E	++	+				+
2929/2930	L569H/L628M/L670F/S678T	++	+	+		+	++
2931/2932	L569Y/L670F/S678T	+	++				++
2933/2934	L569Y/V589I/L670F	++	++		+	+	++
2935/2936	L569Y/V589I/L670T/L871K	++	++	+	+	+	++
2937/2938	L569T/L871S	+	++			+	+
2939/2940	L871K	+		+		++	++
2941/2942	L569Y/L670T				++	+	++
2943/2944	L569Y/T692Y/A711H	++	++		+		+
2945/2946	L871S	+				+	++
2947/2948	L569Y/A711H	++	++				++

Table 14-1. Activity of GAA Variants Relative to SEQ ID NO: 2496¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
2949/2950	L628M/L871S	+			+	++	++
2951/2952	L569T/V589I/L871S	++	+		+	+	++
2953/2954	T692G/A711H/I869L/V878S	++		++	++	++	++
2955/2956	A572S/F588L/S678T/T692G/I869L/V878A	+			+	+	+
2957/2958	A711H	++		+	++	++	++
2959/2960	T692G/I869S				+	++	+
2961/2962	A572S/T692L/I869S/V878S	+	+	+	++	++	++
2963/2964	L629I/I869L/V878S			+		++	+
2965/2966	A572S/F588L/Q795E						+
2967/2968	A711H/I869S/V878S	+				+	+
2969/2970	V878S	++	+		++	+	+
2971/2972	S678T/T692G/A711H/I869S	++	+	+	++	++	++
2973/2974	A572S/T692G/I869S/V878S	+				+	
2975/2976	A572S/T692G/V878S	+			+	++	++
2977/2978	A711H/Q795E/I869S/V878S	+				+	++
2979/2980	Q795E/V878S	+	+			+	
2981/2982	T692G/A711H/Q795E/I869S	+		+		+	+

Table 14-1. Activity of GAA Variants Relative to SEQ ID NO: 2496¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/FIOPC
2983/2984	S678T/T692G/Q795E/I869S				+	++	++
2985/2986	T692G/A711H/I869S/V878A				+	++	
2987/2988	L569T/A711H	++			+	+	
2989/2990	L569T/L628M/S678T/A711H	+	+			+	+
2991/2992	L569T/V589I/T692G/Q795E/L871K/V878S	++	++		++	++	
2993/2994	L569T/V589I/S678T/L871K/V878S	++	++		++	++	+
2995/2996	L569H/A711H/I869S/V878S	+				++	
2997/2998	S678T/Q795E/L871K/V878A	+				+	++
2999/3000	L569H/S678T/T692G/Q795E	+			+	+	
3001/3002	F588L/V589I/L628M/S678T	++			+	+	+
3003/3004	T692Y	++	+	+	+	++	+
3005/3006	L569T	++	+		+	++	+
3007/3008	T692G/L871K	+		+	+	++	
3009/3010	L569T/F588L/V878S	++	++	+	++	++	+
3011/3012	L569H/A711H/L871K	++			++	+	

Table 14-1. Activity of GAA Variants Relative to SEQ ID NO: 2496 ¹							
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3013/3014	L569T/F588L/S678T/T692Y	++	++		+	+	+
3015/3016	L569T/F588L/A711H/I869L/L871K/V878S	+			+	++	
3017/3018	L569T/F588L	++	++			+	++
3019/3020	L569T/T692Y	+	++	+	++	++	+
3021/3022	L569T/S678T/I869S/V878S	++	+		++	+	++
3023/3024	L628M/L629I/T692Y/L871S/V878S	+			+	++	+
3025/3026	L569T/S678T/V878S	++	++	+	++	++	++
3027/3028	L569H/A711H/Q795E/L871S/V878S	++	++		+	++	++
3029/3030	A711H/L871K	++				+	+
3031/3032	T692G/A711H/Q795E/I869L/L871K/V878A	+			+	++	+
3033/3034	L569T/F588L/V589I/L628M/L629I/T692Y/A711H		+	+	+		++
3035/3036	S678T/T692G/I869S	++			+	++	+

Table 14-1. Activity of GAA Variants Relative to SEQ ID NO: 2496¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3037/3038	T692G/I869L/L871K/V878S/S916R	+			+	++	+
3039/3040	L569H/L628M	++	+			+	+
3041/3042	L628M/V878S	++			+	+	+
3043/3044	T692G/A711H/L871S/V878A	++		+	++	++	++
3045/3046	L569T/Q795E	++	++		+	+	++
3047/3048	L569T/T692Y/A711H/I869L/L871K/V878A	++	++	+	++	++	++
3049/3050	S246T	++			+	+	+
3051/3052	T585K	++				+	++
3053/3054	G584E	+				+	
3055/3056	A812E	++	+		+	+	+
3057/3058	V313L	+				+	
3059/3060	L304M	++			+	++	++
3061/3062	V313I	++			+	++	++
3063/3064	A582T	++			++	++	++

¹All activities were determined relative to the reference polypeptide of SEQ ID NO: 2496. Levels of increased activity are defined as follows: "+" = 0.9 to 1.1; and "++" > 1.1.

Table 14-2. Activity of GAA Variants Relative to SEQ ID NO: 2496¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC
2495/2496		+	+	+	+
3065/3066	L810V		++		
3067/3068	A60V/V589A				+
3069/3070	S307T				+

Table 14-2. Activity of GAA Variants Relative to SEQ ID NO: 2496¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC
3071/3072	V313T			+	
3073/3074	G584C				+

¹All activities were determined relative to the reference polypeptide of SEQ ID NO: 2496. Levels of increased activity are defined as follows: "+" = 0.9 to 1.1; and "++" > 1.1.

EXAMPLE 15**GAA Variants of SEQ ID NO: 2880**

[0221] In this Example, experiments for evolution and screening of GAA variants derived from SEQ ID NO: 2880 for improved GAA activity after a series of challenges are described. Libraries of variant genes GAA encoded based off of by SEQ ID NO: 2880 were constructed, plated, grown, and screened for GAA MU-Glu activity ("Unchallenged Activity FIOPC"), as well as after plasma challenge ("Plasma Stability and Activity FIOPC"), as described in Example 11. Variants were also tested for 4-MUGlu activity after lysis of GAA^{-/-} C2C12 cells ("Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC") or 6 hour treatments hours of Pompe fibroblasts ("Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC") or GAA^{-/-} C2C12 cells ("Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC"), as described in Example 11. The results of these assays are presented in Table 15-1.

Table 15-1. Activity of GAA Variants Relative to SEQ ID NO: 2880¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2880)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
2879/2880		+	+	+	+	+
3075/3076	R527N/L736M/A932S	++	++	+	++	++
3077/3078	L109D/S678T/A812E	++	++	++	+	++
3079/3080	A89R/S842G/V878S	++	+	+	+	+
3081/3082	R403H/R527N/S678T/T692G/L736M/A812E/S842G/L860F	++	++	++	++	++

Table 15-1. Activity of GAA Variants Relative to SEQ ID NO: 2880¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2880)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3083/3084	S678T/T692G/A812E	++	++	+	+	++
3085/3086	L109D/W727S/L860F/V878S	++	++	++	+	++
3087/3088	A89R/R527N/W727S/A812E/L860F	++	++	++	+	++
3089/3090	A89R/S678T/T692G/L736M/A932S	++	++	++	++	++
3091/3092	L109D/R527N/S678T/A812E	++	++	+	+	++
3093/3094	S678T/L860F/V878S	++	++	++	+	++
3095/3096	L109D/A812E	++	++	+		++
3097/3098	T692G/W727S/L736M/S842G/R913V	++	++	++	++	++
3099/3100	A89R/L109D/W727S/A932S	++	++	++	+	++
3101/3102	R527N/S678T/T692G/W727S/L736M/V878S	++	++	++	+	++
3103/3104	R527N/T692G/W727S/L736M/A812E	++	++	++	+	++
3105/3106	A89R/R527N	++	++	+		+
3107/3108	A89R/L109D/A932S	++	++	+		+
3109/3110	L109D/S842G	+		+	+	++
3111/3112	S678T/R913V	++	++	++	++	++
3113/3114	A89R/L109D/S678T/W727S/L736M/A812E/V878S	++	++	++	+	++
3115/3116	L109D/T692G/W727S/A812E/S842G/L860F	++	++	++	++	++
3117/3118	T692G/A812E	++	+	++	+	+
3119/3120	S678T/A932S	++	+	++	+	++
3121/3122	L109D/S678T/L736M/A812E/V878S	++	++	++	++	++

Table 15-1. Activity of GAA Variants Relative to SEQ ID NO: 2880¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2880)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3123/3124	L109D/A932S	++	++	++	+	++
3125/3126	R527N/T692G/ W727S/L736M/ S842G/L860F/V 878S	++	++	+	+	++
3127/3128	R527N/W727S/ L736M	++	++	+		++
3129/3130	L109D/S678T/S 842G/V878S	++	+	+	++	++
3131/3132	L109D/S678T/ W727S/L860F	++	++	++	+	++
3133/3134	R527N/S678T/T 692G/A812E/A 932S	++	++	++	++	+++
3135/3136	A89R/S678T/A 812E/V878S	++	+	++	++	++
3137/3138	R527N/A812E	++	++	+		++
3139/3140	S678T/T692G/ W727S/A812E/ S842G	++	+	+		++
3141/3142	L109D/L736M/ A932S	++	++	++	++	++
3143/3144	A89R/L109D/R 527N/S678T/W 727S/S842G	++	++	++		++
3145/3146	W727S	++	+	++	+	++
3147/3148	A89R/R527N/S 678T/A932S	++	++	++	++	+++
3149/3150	L109D/S678T/T 692G/S842G/L8 60F/V878S/A93 2S	++	++	++	++	+++
3151/3152	A89R/R527N/S 678T/T692G/L7 36M/S842G/V8 78S/A932S	++	+	++	+	++
3153/3154	R527N	++		+		+
3155/3156	L109D/T692G/ W727S/L736M/ A812E	++	++	+	+	++
3157/3158	W727S/A932S	++	++	++	+	++
3159/3160	V878S/A932S	++	++	++	+	++
3161/3162	S678T/A812E	++	+	++	++	+++

Table 15-1. Activity of GAA Variants Relative to SEQ ID NO: 2880 ¹						
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2880)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3163/3164	W24L/S28L/T29L/Q39P/V50Q/L62A/E78P/E87D/Q135S/S150T/N266T/K267R/V522E/R527N/A551V/T670L/W727S/P750A/K830Q/S842G/E871L/H883R/G894Q/A932S	++	++	+		++
3165/3166	S28L/V50Q/E78P/E87D/Q135S/N266T/K267R/G437A/E486T/R527N/A551V/T670L/W727S/P750A/K830Q/S842G/E871L/H883R/G894Q/R913V/A932S	++	++	+		+
3167/3168	W24L/S28L/Q39P/V50Q/L62A/E78P/E87D/Q135S/S150T/N266T/K267R/V522E/R527N/A551V/T569L/W727S/K830Q/S842G/E871L/H883R/G894Q/R913V	++	+			+
3169/3170	T29L/Q39P/V50Q/L62A/G65R/E78P/E87D/Q135S/S150T/G437A/A551V/T569L/T670L/W727S/P750A/K830Q/S842G/H883R/G894Q/A932S	++	+	++		

Table 15-1. Activity of GAA Variants Relative to SEQ ID NO: 2880¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2880)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3171/3172	W24L/V50Q/E78P/E87D/Q135S/S150T/K267R/E486T/V522E/R527N/A551V/T670L/W727S/P750A/K830Q/S842G/E871L/H883R/G894Q/R913V/A932S	++	++	++		++
3173/3174	W24L/T29L/Q39P/V50Q/L62A/E78P/E87D/Q135S/S150T/K267R/G437A/E486T/V522E/R527N/A551V/H711A/W727S/P750A/K830Q/S842G/E871L/H883R/G894Q/R913V/A932S	++	++	+		++
3175/3176	L62A/Q135S/V522E/H711A/W727S/P750A/S842G/E871L/G894Q	++	++	++	+	++
3177/3178	E87D/P750A	+	++	++		++
3179/3180	Q135S/H711A/P750A/A932S	+	+	++	+	++
3181/3182	V522E/K830Q	++	++	++		++
3183/3184	S28L/G437A/R527N/E871L	++	++	++	++	++
3185/3186	L62A/G437A	++	++	+		++
3187/3188	G437A/P750A/K830Q/A932S	++	++	+	++	++
3189/3190	T29L/E78P/Q135S/W727S/K830Q	++	++	++	+	++
3191/3192	L62A	++	++	++	++	++
3193/3194	V50Q/H711A	+	+	++	+	++
3195/3196	S150T/E871L/A932S	+	++	+	++	+
3197/3198	Q39P/V50Q	+	+	++	+	++

Table 15-1. Activity of GAA Variants Relative to SEQ ID NO: 2880¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2880)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3199/3200	E78P/E87D/E486T/R527N/T670L/W727S/P750A/K830Q/S842G/E871L/R913V/A932S	+	+	++		++
3201/3202	W24L/R527N/W727S/S842G/E871L/H883R/R913V/A932S	++	++	++		+++
3203/3204	W24L/T670L/W727S/P750A/S842G/E871L	++	++	++	++	++
3205/3206	V50Q/W727S/P750A/H883R/G894Q	++	++	++	++	++
3207/3208	T29L/Q135S/S150T/R527N/T670L/W727S/H883R	+	+	++	+	++
3209/3210	Q39P/W727S/P750A/A932S	++	++	++	+	++
3211/3212	S150T/H883R/A932S	+	+	+	+	
3213/3214	Q135S/T670L/W727S	+	+	+++	++	++
3215/3216	T29L/E78P/E87D/S150T/R527N/W727S	+	+	+++	++	++
3217/3218	W24L/S28L/V50Q/Q135S/S150T/G437A/V522E/R527N/E871L/H883R/G894Q/A932S	++	++	+++		++
3219/3220	W24L/L62A/E87D/E486T/W727S	+	+	+++	+	+
3221/3222	T29L/V522E/T670L/H711A/E871L	+	+	+++		+
3223/3224	S28L/V522E/R527N/T569L/H711A/K830Q/G894Q	+	+	++		

Table 15-1. Activity of GAA Variants Relative to SEQ ID NO: 2880¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2880)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3225/3226	S28L/W727S/E871L	+	+	+++	++	+++
3227/3228	T670L/H711A/E871L			+++	+	++
3229/3230	T29L/S150T/G437A/W727S	+		++		+
3231/3232	K267R/R527N/W727S	++	+	+++	++	++
3233/3234	S150T/R527N/S842G/E871L/R913V	+	+	+++	++	++
3235/3236	T29L/S150T/K267R/W727S/P750A/E871L/H883R/A932S	+	+	++	++	++
3237/3238	S28L/W727S	+		+++	++	++
3239/3240	T29L/E87D	+		+++	++	++
3241/3242	T29L/L62A/G437A/R527N	+		++	+	+
3243/3244	V50Q/Q135S/S150T/A932S	+	+	+++	++	++
3245/3246	V50Q/G437A/V522E/R527N	+	+	++		+
3247/3248	S28L/L62A	++	++	++	++	++
3249/3250	S28L/L62A/K267R/A932S	++	++	++	++	++
3251/3252	W24L/V50Q/E486T/R527N/H711A/W727S	++	++	+	+	++
3253/3254	L62A/E87D/S150T/W727S	++	+++	++	++	++
3255/3256	S28L	++	++	++	+++	++
3257/3258	W24L/L62A/W727S/K830Q/A932S	+++	+++	++	++	++
3259/3260	V522E/R527N/T569L/W727S	++	++	+	+	++
3261/3262	T29L/T670L/A932S	++	++	++	++	++
3263/3264	E871L	++	+++	++	+++	++
3265/3266	G437A/V522E/R527N/T670L/E871L	++	++	+		+

Table 15-1. Activity of GAA Variants Relative to SEQ ID NO: 2880¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2880)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3267/3268	V522E	++	++	++	+	++
3269/3270	L62A/E87D/S150T	++	++	++	++	++
3271/3272	W24L/S28L/L62A/V522E/T569L/A932S	++	++	++	++	+
3273/3274	W24L/E87D/Q135S/V522E/T670L/H711A/K830Q/S842G/R913V	++	+++	++	++	++
3275/3276	W24L/S150T/V522E/R527N/W727S/H883R/G894Q	++	+++	++	++	++
3277/3278	T670L/K830Q/E871L	++	+++	++	+	++
3279/3280	L62A/G437A/W727S	+++	+++	++	++	++
3281/3282	S28L/V50Q/V522E/R527N/H711A/W727S/E871L	+++	+++	++	++	++
3283/3284	W24L/S28L/G437A/E486T/R527N	++	++	++	++	++

¹All activities were determined relative to the reference polypeptide of SEQ ID NO: 2880. Levels of increased activity are defined as follows: "+" = 0.9 to 1.1; "++" > 1.1; "+++" > 2; and "++++" > 3.5.

EXAMPLE 16

GAA Variants of SEQ ID NO: 3104

[0222] In this Example, experiments for evolution and screening of GAA variants derived from SEQ ID NO: 3104 for improved GAA activity after a series of challenges are described. Libraries of variant genes GAA encoded based off of by SEQ ID NO: 3104 were constructed, plated, grown, and screened for GAA MU-Glu activity ("Unchallenged Activity FIOPC"), as well as after plasma challenge ("Plasma Stability and Activity FIOPC"), as described in Example 11. Variants were also tested for 4-MUGlu activity after lysis of Pompe fibroblasts treated for 20 hours ("Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC") or GAA^{-/-} C2C12 cells ("Lysate Activity from Extended Treatment C2C12 GAA^{-/-} FIOPC") or 4 hour treatments hours ("Lysate Activity from

Acute Treatment of Pompe Fibroblast FIOPC”) or (“Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC”), as described in Example 11. The results of these assays are presented in Table 16-1.

Table 16-1. Activity of GAA Variants Relative to SEQ ID NO: 3104 ¹							
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 3104)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3285/3286	L62A/L860F/A932S	+	+	+	+	+	++
3287/3288	L62A/S678T/K830Q/L860F/V878S/A932S	+	+	+	+	+	+++
3289/3290	L62A/K830Q/L860F/V878S/A932S	+	+	+	+	+	+
3291/3292	S678T/A932S	+	+	+	+	+	+
3293/3294	L62A/M833I/L860F/A932S	+	+	+	+	+	+
3295/3296	L62A/L860F/E871L/V878S	+	+	+	+	+	+
3297/3298	L62A/S678T/P785Q	+	+	+	+	+	+
3299/3300	L62A/V878S/A932S	+	+	+	+	+	++
3301/3302	L62A/K830Q/L860F/E871L/R873H	+	+	+	+	+	+
3303/3304	L62A/Y248H/S678T/K830Q/V878S/A932S	+	+	+	+	+	+
3305/3306	L62A/L860F/E871L	+	+	+	+	+	+
3307/3308	S678T	+	+	+	+	+	+
3309/3310	L62A/S678T/K830Q/E871L/A932S	+	+	+	+	+	+

Table 16-1. Activity of GAA Variants Relative to SEQ ID NO: 3104¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 3104)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3311/3312	L62A/S678T/L860F	+	+	+	+	+	++
3313/3314	L62A/A89D/K830Q	+	+	+	+	+	+
3315/3316	L62A/V878S	+	+	+	+	+	+
3317/3318	L62A/K830Q/A932S	+	+	+	+	+	+++
3319/3320	L62A/S678T/K830Q/L860F/A932S	+	+	+	+	+	++
3321/3322	L62A/A932S	+	+	+	+	+	++
3323/3324	L62A/S678T/L860F/V878S	+	+	+	+	+	+
3325/3326	L62A/S678T/E871L	+	+	+	+	+	+++
3327/3328	L62A	+	+	+	+	+	++
3329/3330	L62A/S678T/K830Q/L860F	+	+	+	+	+	+++
3331/3332	L62A/K830Q/V878S	+	+	+	+	+	+
3333/3334	L62A/E871L/A932S	+	+	+	+	+	+++
3335/3336	L62A/K830Q/E871L/A932S	+	+	+	+	+	+
3337/3338	L62A/S678T/K830Q/E871L	+	+	+	+	+	++
3339/3340	L860F/A932S	+	+	+	+	+	++
3341/3342	L62A/S678T/V878S/A932S	+	+	+	+	+	+
3343/3344	L62A/S678T/K830Q	+	+	+	+	+	+

Table 16-1. Activity of GAA Variants Relative to SEQ ID NO: 3104¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 3104)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3345/3346	L62A/L860F	+	+	+	+	+	+++
3347/3348	L62A/S678T/V878S/A932T	+	+	+	+	+	+++
3349/3350	L62A/S678T/K830Q/L860F/V878S	+	+	+	+	+	+
3351/3352	L62A/L860F/V878S/A932S	+	+	+	+	+	+
3353/3354	L62A/S678T/L860F/A932S	+	+	+	+	+	+
3355/3356	L62A/S678T	+	+	+	+	+	++
3357/3358	L62A/S678T/A932S	+	+	+	+	+	+
3359/3360	L62A/S678T/E871L/A932S	+	+	+	+	+	++
3361/3362	L62A/K830Q/L860F/A932S	+	+	+	+	+	++
3363/3364	L62A/S678T/K830Q/V878S/A932S	+	+	+	+	+	+
3365/3366	L62A/E871L/V878S/A932S	+	+	+	+	+	+++
3367/3368	L62A/K830Q/L860F	+	+	+	+	+	++
3369/3370	L62A/S678T/K830Q/A932S	+	+	+	+	+	++
3371/3372	L62A/L860F/E871L/A932S	+	+	+	+	+	+

Table 16-1. Activity of GAA Variants Relative to SEQ ID NO: 3104¹

SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 3104)	Unchallenged Activity FIOPC	Plasma Stability and Activity FIOPC	Lysate Activity from Extended Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Acute Treatment of Pompe Fibroblast FIOPC	Lysate Activity from Extended Treatment C2C12 GAA-/- FIOPC	Lysate Activity from Acute Treatment C2C12 GAA-/- FIOPC
3373/3374	L62A/S678T/K830Q/L860F/E871L/V878S/A932S	+	+	+	+	+	+++
3375/3376	L62A/K830Q	+	+	+	+	+	++
3377/3378	S678T/K830Q/A932S	+	+	+	+	+	+

¹All activities were determined relative to the reference polypeptide of SEQ ID NO: 3104. Levels of increased activity are defined as follows: "+" = 0.9 to 1.1; "++" > 1.1; "+++" > 2; and "++++" > 3.5.

EXAMPLE 17**Identification of active GAA variants with reduced immunogenicity**

[0223] Putative T-cell epitopes in a WT GAA (SEQ ID NO:6) were identified using the Immune Epitope Database (IEDB; Immune Epitope Database and Analysis Resource website) tools, as known in the art and proprietary statistical analysis tools (See e.g., iedb.org and Vita et al., Nucl. Acids Res., 38(Database issue): D854-62 [2010]. Epub 2009 Nov 11). The WT GAA was parsed into all possible 15-mer analysis frames, with each frame overlapping the last by 14 amino acids. The 15-mer analysis frames were evaluated for immunogenic potential by scoring their 9-mer core regions for predicted binding to eight common class II HLA-DR alleles (DRB1*0101, DRB1*0301, DRB1*0401, DRB1*0701, DRB1*0801, DRB1*1101, DRB1*1301, and DRB1*1501) that collectively cover nearly 95% of the human population (See e.g., Southwood et al., J. Immunol., 160:3363-3373 [1998]), using methods recommended on the IEDB website. Potential T-cell epitope clusters contained within the enzyme (i.e., sub-regions contained within GAA which have an unusually high potential for immunogenicity) were identified using statistical analysis tools, as known in the art.

[0224] GAA variants that were identified in Examples 3, 4, 6, 10, and 12-16 to be active in assays described in Examples 5 and 11 were analyzed for their levels of predicted immunogenicity by evaluating their binding to the eight common Class II HLA-DR alleles. The total immunogenicity score and immunogenic hit count were calculated for each variant. The total immunogenicity score (TIS) reflects the overall predicted immunogenicity of the variant (i.e., a higher score indicates a

higher level of predicted immunogenicity). The immunogenic "hit count" (IHC) indicates the number of 15-mer analysis frames with an unusually high potential for immunogenicity (i.e., a higher score indicates a higher potential for immunogenicity). Mutations resulting in a reduced total immunogenicity score and/or an immunogenic hit count compared to the reference sequence were considered to be potential "deimmunizing mutations" and are shown in Tables 17-1, 17-2, 17-3, 17-4, 17-5, 17-6, 17-7, 17-8, and 17-9. A collection of the most deimmunizing mutations were recombined to generate a number of variants that were active and predicted to be significantly less immunogenic than WT GAA.

Table 17-1. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 2¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	TIS Reduction (Relative to SEQ ID NO: 2)	IHC Reduction (Relative to SEQ ID NO: 2)
7/8	T150S/T486E/A750P/R883H/Q894G	+++	++++
101/102	L871E	+++	+++
105/106	M138A	+	
109/110	Q107G	++	
113/114	Q110G	++	
115/116	Q110L	+	
123/124	R403W	+	
125/126	R414P	+	
137/138	S135Q	+	
145/146	T148G	+	
147/148	T148Y	+	
149/150	T150G	+	
155/156	T692Q	++	+
159/160	V30G	+++	+++
161/162	V30K	++	+
163/164	V30T	+++	+++
183/184	L275M/A281V/S402A/M431V/M507L/I518V/W610R/S668D	++	
185/186	L275M/M507L/A547G/S668D/L669H/S671N	+++	+
187/188	L275M/M431V/V638I	++	
189/190	A281V/S402A/I518V/A547G/S668D	++	
193/194	L275V/M431V/M507L/I518V/A547G/S668D/L669H/S671N	+++	+
195/196	S402A/M431V/I518V/W610R	+	
197/198	K106P/H191R/G280D/S402A/R414G/A444P/S727P	++	
199/200	M431V/M507L/I518V/L669H/S671N	++	+
201/202	L275V/R377K/S402A/M507L/I518V/L669H/S671N/V715G	++	+
203/204	H191R/R414G/E522V/G842S/C944S	+	
207/208	L275M/A281V/W610R/V638I/S668D/L669H	++	

Table 17-1. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 2¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	TIS Reduction (Relative to SEQ ID NO: 2)	IHC Reduction (Relative to SEQ ID NO: 2)
209/210	A196V/S402A/M431V/A547G/W610R/V638I	++	
211/212	H191R/G280D/S402A/R414G/A444P/A489D/D500A/C944S	++	
213/214	L275V/S402A/V638I/L669H/S671N	+	+
215/216	L29Q/L240I/A596P/S668D/I869L	++++	++++
217/218	K106P/G280D/S402A/R414G/A444P/A489D/S727P/C944S	+++	
219/220	L29Q/L240I/A596S/S668D/H700F/I744V/I869T	++++	+++
221/222	L218S/S668D/H700F/I869T	+++	+
223/224	M507L/A547G/W610R	+	
225/226	A281V/M431V/M507L/I518V/A547G/W610R/V638I/S668D	++	
227/228	H191R/G280D/R414G/A444P/A489D/E522V/S727P/C944S	++	
229/230	L275V/S402A/M431V/I518V/W610R/V638I/L669H/S671N/P922L	+	
23/24	A276Y	+	
231/232	L29V/L218S/L240I/H700F/I869T	++	+
233/234	A547G/V638I/S668D	+++	+
235/236	H191R/G280D/R414G/C944S	+	
237/238	L275V/M431V/M507L/I518V/W610R/L669H/S671N	+	+
239/240	L275M/S402A/M431V/M507L/A547G/S671N	+++	
241/242	S402A/M431V/A547G/V638I/S671N	+++	
243/244	A281V/S402A/M507L/A547G/V638I/L669H/S671N	+	
245/246	L275M/A281V/M507L/A547G/L669H/S671N	++	
247/248	L275M/M431V/I518V/A547G/V638I/S668D	+++	+
249/250	A547G/W610R/V638I/S671N	++	
25/26	A418E/H499R	+	
251/252	L275V/M431V/M507L/A547G/W610R/V638I/S671N	++	
253/254	L275M/S402A/M507L/A547G/W610R/S671N	+++	
255/256	L275M/A281V/S402A/I518V/A547G/W610R/V638I/S671N	++	
257/258	A281V/S402A/I518V/A547G/W610R/V638I/S668D/L669H	++	
259/260	L275M/A281V/S402A/A547G/W610R/V638I/L669H/S671N	++	
261/262	L275M/M431V/I518V/W610R/V638I/L669H/S671N	++	+

Table 17-1. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 2¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	TIS Reduction (Relative to SEQ ID NO: 2)	IHC Reduction (Relative to SEQ ID NO: 2)
263/264	L29Q/L218S/L240I/S668D/H700F/I744V/I869L	+++	+
265/266	A281V/S402A/M507L/I518V/A547G/W610R/V638I/L669H/S671N	+	
267/268	S402A/M431V/I518V/A547G/S671N	+++	
269/270	L275V/A281V/S402A/M431V/I518V/A547G/W610R/L669H/S671N	+	
271/272	L224F/S402A/M507L/I518V/A547G/V638I/S668D	+++	++
273/274	N180H/S402A/M507L/A547G/W610R/S671N	++	
275/276	L275M/A281V/S402A/M507L/I518V/A547G/V638I/L669H/S671N	++	
277/278	K106P/T150S/T486E/Q749E/E793K/R883H/Q894G	+++	+++
279/280	L275V/A281V/M431V/I518V/A547G/V638I/L669H/S671N	+	
281/282	L275V/S402A/A547G/W610R/V638I/L669H/S671N	++	+
283/284	L275M/S402A/A547G/V638I/L669H/S671N	+++	+
285/286	K106P/T150S/T486E/N527D/A750P/E793K	+	
287/288	S402A/M431V/I518V/A547G/W610R/S668D	+++	+
29/30	A437S	++	++
291/292	L275V/M507L/I518V/A547G/V638I/L669H/S671N	++	+
293/294	N180H/S402A/M431V/M507L/A547G/W610R/L669H/S671N/E793G	+++	
295/296	L275V/M507L/I518V/A547G/W610R/V638I/S668D/L669H	+++	+
297/298	L275V/S402A/M507L/A547G/W610R/V638I/S668D/L669H	+++	+
299/300	M507L/A547G/V638I/L669H/S671N	++	+
303/304	T150S/R414G/T486E/Q749E/A750P/E793K	++	
305/306	K106P/T150S/L218S/R414G/T486E/L642F/A750P/E793K/R883H	+++	
307/308	N180H/L275M/S402A/I518V/A547G/W610R/V638I/L669H/S671N	+++	+
309/310	T150S/L218S/R414G/Q749E/E793K	+	
311/312	L275V/M507L/A547G/W610R/V638I/L669H/S671N	+	+
313/314	T150S/R414G/T486E/N527D/A750P/Q894R	++	
315/316	K106P/T150S/L218S/N527D/E793K/Q894G	++	+

Table 17-1. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 2 ¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	TIS Reduction (Relative to SEQ ID NO: 2)	IHC Reduction (Relative to SEQ ID NO: 2)
321/322	K106P/T150S/R414G/Q749E/A750P/E793K/Q894R	++	
323/324	K106P/T150S/N169S/N527D/Q749E/E793K/R883H	++	
325/326	L275V/S402A/M507L/A547G/W610R/V638I/L669H/S671N	++	+
329/330	M431V/M507L/I518V/G541E/A547G/V638I/L669H/S671N	+++	+
331/332	T150S/L218S/R414G/T486A/A750P/E793K	+	
335/336	K106P/T150S/N169S/L218S/R414G/T486E/Q894R	+	
337/338	T150S/L218S/R414G/T486E/A750P/E793K/R883H	++	
339/340	K106P/T150S/N169S/L218S/R414G/Q749E/E793K	+	
341/342	T269N/L275M/M431V/I518V/A547G/V638I/S668D/L669H	+++	+
343/344	K106P/T150S/R414G/Q749E/E793K/Q894R	++	
347/348	K106P/T150S/N169S/Q749E/E793K/R883H/Q894R	++	
35/36	A547G	+	
353/354	K106P/T150S/L218S/R414G/T486E/A750P/E793K/Q894R	+	
355/356	K106P/T150S/N169S/L218S/R414G/T486E/Q749E/E793K/R883H/Q894R	++	
357/358	K106P/T150S/N169S/L218S/T486E/R883H	++	+
359/360	K106P/T150S/N169S/L218S/R414G/T486E/A750P/E793K/R883H/Q894R	++	
361/362	G36R/K106P/T150S/L218S/N527D/A750P/R883H/Q894R	+	
363/364	K106P/T150S/N169S/L218S/T486E/Q749E/R883H	++	+
365/366	K106P/N169S/V185G/L218S/R414G/Q749E/A750P/E793K	+	
367/368	K106P/T150S/P245S/E793K/R883H/Q894R	+	
369/370	K106P/T150S/N169S/L218S/R414G/Q749E/A750P/E793K/R883H	++	
37/38	A750P	+	
375/376	T150S/L218S/R414G/Q749E/A750P/E793K/Q894R	+	
377/378	K106P/T150S/N169S/R414G/T486E/Q749E/A750P/R883H	+++	+
379/380	K106P/T150S/N169S/L218S/R414G/T486E/E793K/R883H	++	

Table 17-1. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 2¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	TIS Reduction (Relative to SEQ ID NO: 2)	IHC Reduction (Relative to SEQ ID NO: 2)
381/382	K106P/T150S/L218S/T486E/N527D/Q749E/Q894R	+	
383/384	K106P/T150S/N169S/L218S/R414G/T486E/N527D/Q894R	+	
385/386	T150S/L218S/R414G/T486E/Q749E/A750P	++	
387/388	K106P/T150S/L218S/R414G/Q749E/E793K/R883H	++	
389/390	K106P/T150S/N169S/L218S/R414G/T486E/N527D/A750P/Q894R	++	
39/40	A753T	+	
391/392	N169S/T486E/A750P/E793K/R883H	++	
393/394	K106P/T150S/L218S/R414G/N527D/Q749E/A750P/R883H	+++	+
395/396	K106P/T150S/L218S/R414G/Q749E/A750P/E793K/R883H/Q894R	++	
397/398	K106P/T150S/N169S/L218S/R414G/Q749E/E793K/R883H	++	
399/400	K106P/T150S/L218S/T486E/E793K/R883H	+	
401/402	K106P/T150S/L218S/V331A/R414G/T486E/N527D/D733E/Q749E/E793K	++	
403/404	L275M/A281V/S402A/I518V/A547G/W610R/S668D/L669H/E887D	+++	+
405/406	K106P/T150S/L218S/R414G/N527D/Q749E/E793K/R883H/Q894G	+++	++
407/408	K106P/T150S/Q749E/E793K/R883H	++	
409/410	K106P/A112S/T150S/L218S/R414G/N527D/E793K/R883H	++	
43/44	C944G	+	
47/48	D274G	+	
53/54	F27P	+	+
55/56	F27P/C944W	+	+
57/58	F27R	+	+
61/62	G820E	+	
63/64	I375E	+	
65/66	K471Q/A478S	+	
69/70	K581G	++	+
71/72	K581T	+	
81/82	L109G/G842E	+	
83/84	L109P	+	
85/86	L28P	++	++
89/90	L28S	++	++
91/92	L29T/A478T	+++	++
95/96	L642Q	+	
97/98	L642S	+	
99/100	L670N	+++	+

Table 17-1. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 2¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2)	TIS Reduction (Relative to SEQ ID NO: 2)	IHC Reduction (Relative to SEQ ID NO: 2)
¹ TIS reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 2, and defined as follows: "+" = 1-10; "++" > 10; "+++" > 20; and "++++" > 50. IHC reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 2, and defined as follows: "+" = 1-3; "++" > 3; "+++" > 5; and "++++" > 8.			

Table 17-2. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 8¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 8)	TIS Reduction (Relative to SEQ ID NO: 8)	IHC Reduction (Relative to SEQ ID NO: 8)
13/14	L28S/L29T/S135Q	++++	+++
413/414	A418E	+++	+
415/416	A418E/A489R	+++	+
431/432	K106P	+++	+
433/434	K106P/A489R/S671N	+++	+
445/446	K106P/V638I	+++	+
447/448	L157M	+++	+
469/470	L218S/V638I/S671N	+++	
509/510	Q113S/A418E/A478T/A489R/K581T	+++	
511/512	Q113S/A418E/R455V/A478T/K581T	+++	
519/520	S135P/C944Y	+++	+
521/522	S135Q	+++	+
523/524	S135Q/A478T/A489R/K581T	+++	
531/532	S671N/Q749E	+++	+
441/442	K106P/M138A/L218S/M431V/S671N/Q749E	+++	
513/514	Q113S/S135Q/A418E	+++	+
527/528	S135Q/T148G/S150G/A418E	+++	+
429/430	F27R	+++	+
425/426	F27P/A418E/A478T	+++	+
501/502	M507L/S668D	+++	+
495/496	M138A/L218S/S668D/S671N	+++	+
499/500	M431V/S668D/S671N	++++	++
471/472	L28S	+++	++
481/482	L29T/A478T	+++	+++
483/484	L29T/Q113S/M138A	+++	+++
485/486	L29T/Q113S/P126Q/S135Q/H193Q	+++	+++
487/488	L29T/Q113S/S135Q	+++	+++
489/490	L29T/Q113S/S135Q/R455V	+++	+++
491/492	L29T/T148G	+++	+++
473/474	L28S/L29T	+++	+++
475/476	L28S/L29T/A418E	++++	+++
477/478	L28S/L29T/Q113S/S135Q/A418E	++++	+++
479/480	L28S/L29T/Q113S/S135Q/M138A	++++	+++

Table 17-2. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 8¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 8)	TIS Reduction (Relative to SEQ ID NO: 8)	IHC Reduction (Relative to SEQ ID NO: 8)
¹ TIS reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 8, and defined as follows: "+" = 1-10; "++" > 10; "+++" > 20; and "++++" > 50. IHC reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 8, and defined as follows: "+" = 1-3; "++" > 5; and "+++" > 8.			

Table 17-3. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 18¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 18)	TIS Reduction (Relative to SEQ ID NO: 18)	IHC Reduction (Relative to SEQ ID NO: 18)
19/20	P39Q/R267K/A489D/E522V/S612D/Q830K/G842S	+	
371/372	P39D	+	
533/534	V70A/R267K/K725E/C944S	+	
535/536	R267K/A489D/D500A/K725E/Q830K/C930P	+	
537/538	L109P/E522V/Q830K/C944S	+	
539/540	V70A/R267K/C930P/C944S	+	
543/544	H734K	+	
553/554	P39Q/V70A/L109P/Q830K/G842S	+	
555/556	P39Q/V70A/K725E	+	
557/558	P39Q/R267K/A489D/Q830K/C944S	+	
559/560	C930P	+	
561/562	D500A/C930P/C944S	+	
567/568	P39Q/R267K	+	
575/576	R267K/E522V/K725E	+	
577/578	V70A/A489D/C930P	+	
585/586	P39Q	+	
601/602	S37F/N528S/I790V	+	
611/612	S37F/A62E	+	
625/626	S37F/A196T	+	
627/628	S37F/A62E/D523N	+	
633/634	S37F/A62E/N79S/A196T/A696S/R862Q	+	
655/656	L34D	+	
657/658	Y352K	+	
667/668	N875D	++	++
673/674	I22R	+	
675/676	L24W	++	
677/678	L778Q	+	
685/686	Y352V	+	
691/692	F27K	+	

Table 17-3. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 18¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 18)	TIS Reduction (Relative to SEQ ID NO: 18)	IHC Reduction (Relative to SEQ ID NO: 18)
695/696	S932A	+	
697/698	L24R	+	
699/700	L24E	++	
709/710	F27G	+	
713/714	L34M	+	
731/732	A774G	+	
735/736	F27W	+	
745/746	R385G	+	
747/748	F27G/M165I	+	
749/750	V30D	+	
753/754	V40W	+	
761/762	E33G	+	
765/766	L109D	+	
793/794	Q107G	++	
801/802	F27R	+	
803/804	L934F	+	
813/814	Q217D	+	
817/818	L34T	+	
821/822	P673N	+	
¹ TIS reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 18, and defined as follows: "+" = 1-10; "++" > 10; "+++" > 20; and "++++" > 50. IHC reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 18, and defined as follows: "+" = 1-2; "++" > 2; and "+++" > 5.			

Table 17-4. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 20¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	TIS Reduction (Relative to SEQ ID NO: 20)	IHC Reduction (Relative to SEQ ID NO: 20)
829/830	L24R/F27G/A89R/D500A/S842G	+	
831/832	L24R/F27G/D500A/S842G	+	
833/834	D500A/S932A	+	
835/836	L24R/Q39H/S842G/S932A	+	
839/840	D500A/S842G	+	
841/842	S842G/S932A	+	
843/844	A89R/A97G/Q107G	++	
847/848	F27G/Q39H/Q49G/A97G/D500A/S842G	+	
849/850	R68N/A89R/A97G/S932A	+	
851/852	A89R/S842G	+	

Table 17-4. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 20¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	TIS Reduction (Relative to SEQ ID NO: 20)	IHC Reduction (Relative to SEQ ID NO: 20)
853/854	L24R/Q39D/R68S/A89R/Q107G/D500A/S842G	++	
855/856	L24R/F27G/D500A/S842G/S932A	+	
857/858	L24R/Q39H/Q49G/A89R/A97G/S842G/S932A	+	
859/860	L24R/S842G/S932A	+	
861/862	L24R/A89R/D500A	+	
863/864	L24R/Q39H/A89R/A97D/S842G/S932A	+	
865/866	L24R	+	
867/868	F27G/Q49G/R68S/D500A/S842G	+	
869/870	Q107G/D500A/S842G	++	
871/872	L34T/Q39D/D500A/S932A	+	
875/876	L24W/L109D/D612S	+++	
877/878	L24W/V70A/K267R/A774S	+	
879/880	I22R/L24W/D612S/K725E/C944R	+++	
881/882	L24W	++	
883/884	I22R/L24W/F27A/D489A	+++	
887/888	I22R/L24W/F27A/Q50V/Q107G/L109D/D489A/D612S/K725E	+++	
889/890	Q50V/D612S/C944S	+	
891/892	D612S/K725E/C944S	+	
893/894	L109D/D612S/A774S/C944S	+	
895/896	I22R/L24W/F27A/D612S/C944R	+++	
899/900	Q50V/L109D/D489A/D612S	+	
901/902	I22R/L24W/Q50V/L109D/K267R/D489A/A774S/C944R	++	
903/904	I22R/L24W	++	
905/906	L24W/Q50V/D489A/C944S	++	
907/908	I22R/Q50V/L109D/K267R/D489A	+	
909/910	L24W/D489A/C944R	++	
911/912	L24W/K725E	++	
913/914	L24W/L109D/C944S	++	
915/916	D612S/P922E	+	
917/918	L24W/K267R/K725E/C944S	+	
919/920	Q107G/L109D	++	
923/924	I22R/L24W/Q50V/K267R/D612S/P922E/C944R	++	
925/926	I22R/L24W/F27A/D489A/D612S/A774S	+++	
927/928	L24W/F27A/Q107G/K267R/D612S/C944S	+++	
929/930	L24W/F27A/K267R/C944R	++	
931/932	I22R/L24W/D489A	++	
933/934	I22R/Q50V/Q107G/K267R/D489A/D612S/C944S	++	

Table 17-4. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 20¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	TIS Reduction (Relative to SEQ ID NO: 20)	IHC Reduction (Relative to SEQ ID NO: 20)
935/936	L24W/V70A/Q107G/L109D/D489A/D612S/K725E	+++	
937/938	I22R/L24W/Q107G/K267R/D489A/P922E	+++	
939/940	L24W/Q107G/L109D/K267R/D489A/D612S/K725E/A774S	+++	
941/942	L24W/F27A/C944R	+++	
943/944	D612S	+	
945/946	L24W/Q50V/D612S	++	
947/948	I22R/K267R/D489A/D612S	+	
949/950	L24W/F27A/Q50V/Q107G/K267R/A774S/C944S	+++	
951/952	L24W/C944S	++	
1121/1122	L818V	+	
1123/1124	T148E	+	
1125/1126	A741D	++	
1129/1130	L737M	+	
1141/1142	G108S	++	
1143/1144	L259N	+	
1153/1154	L238Q	+++	+++
1159/1160	S916A	++	
1167/1168	M432C	++	++
1173/1174	E188Q/R377Q	+	
1175/1176	K106G	+	
1177/1178	K106N	+	
1179/1180	L244I	+	
1181/1182	A319F	+	+
1183/1184	I815M	+	++
1189/1190	P832E	+	
1191/1192	A741T	+	
1207/1208	S243R	+	
1215/1216	S727T	+	
1217/1218	L240I	+	
1231/1232	V913A	++	
965/966	S916R	+	
1235/1236	L940G	+	
967/968	P914G	++	
1261/1262	L305Y	+	+
971/972	D923W	+	
1297/1298	L305G	+++	+++
981/982	L742V	+	
1307/1308	I815A	+	
991/992	P914Q	+	
1327/1328	P914E	+	

Table 17-4. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 20¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	TIS Reduction (Relative to SEQ ID NO: 20)	IHC Reduction (Relative to SEQ ID NO: 20)
1335/1336	S264C	+	
1339/1340	Y248A	+	
1341/1342	L305R	+	+
1001/1002	T148K	+	
1349/1350	L320M		+
1351/1352	L736V	+	
1353/1354	A309C	++	+
1385/1386	S727Q	+	
1007/1008	A112H	+	
1395/1396	L823A	+	
1405/1406	L736W	+	
1409/1410	S916I	+	
1413/1414	V913E	++	
1427/1428	L305V	++	+
1435/1436	Y248V	+	
1439/1440	L818T	++	
1441/1442	A741C	++	
1445/1446	F556S	++	
1013/1014	P914T	+	
1015/1016	S916G	++	
1459/1460	L748I	+	
1461/1462	M260W	+	
1467/1468	G108N	++	
1019/1020	P914K	+	
1471/1472	I816V	+	++
1479/1480	L748T	+	
1023/1024	V913W	++	
1483/1484	T148R/V772I	+	
1485/1486	K106T	+	
1487/1488	L240W/A374T	++	+
1489/1490	F556Y	+	
1499/1500	G108V	+	
1515/1516	S916V	+	
1033/1034	L823F	+	
1527/1528	E937Q	+	
1035/1036	L305F	+	+
1539/1540	P832G	+	
1541/1542	V913H	++	
1545/1546	L252V	+	
1551/1552	S243G	++	+
1555/1556	A741E	++	
1557/1558	S243V	+	

Table 17-4. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 20¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	TIS Reduction (Relative to SEQ ID NO: 20)	IHC Reduction (Relative to SEQ ID NO: 20)
1037/1038	P914S	+	
1039/1040	S727W	+	
1041/1042	L940Q	+	
1567/1568	P279E	+	
1569/1570	V913Q	++	
1583/1584	L259G	+	
1585/1586	S243E	+++	++
1059/1060	S402N	+	
1599/1600	A253G	+	
1601/1602	L736M	++	
1603/1604	L940W	+	
1069/1070	G108R	+	
1605/1606	L823G	+	
1071/1072	L748V	+	
1619/1620	S727G	+	
1621/1622	F556R	++	
1623/1624	L261E	+	
1625/1626	A309G	++	
1631/1632	F556H	++	
1633/1634	K106A	+	
1635/1636	S402G	+	
1641/1642	S262P	+	
1643/1644	P914H	+	
1079/1080	I333L	+	
1651/1652	L329F	+	
1653/1654	S246G	++	+
1081/1082	V312A	+++	+++
1661/1662	T148G	+	
1085/1086	S916H	++	
1683/1684	A741G	+	
1691/1692	S246A	+	
1693/1694	L940T	+	
1703/1704	H387L	+	
1717/1718	L244V	++	+
1733/1734	T148H	+	
1735/1736	Y248R	+	
1101/1102	P914R	+	
1757/1758	A253P	+	
1105/1106	N911G	++	
1769/1770	I333V	+	
1771/1772	A112P	+	
1779/1780	V913L	++	

Table 17-4. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 20¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 20)	TIS Reduction (Relative to SEQ ID NO: 20)	IHC Reduction (Relative to SEQ ID NO: 20)
1785/1786	L259S	+	
1787/1788	L823R	+	
1789/1790	L240Y	+	
1797/1798	I249V/S777N	+	
1109/1110	V913R	+	
1807/1808	G108H	+	
1825/1826	Q421P	+	
1117/1118	V913G	++	
¹ TIS reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 20, and defined as follows: "+" = 1-10; "++" > 10; and "+++" > 20. IHC reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 20, and defined as follows: "+" = 1-3; "++" > 3; "+++" > 5; and "++++" > 8.			

Table 17-5. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 946¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 946)	TIS Reduction (Relative to SEQ ID NO: 946)	IHC Reduction (Relative to SEQ ID NO: 946)
1829/1830	A62W/A89R/E188W/D489A/D500A/S727W/L823F	+	
1831/1832	S124V/D500A/S842G/R862Q	+	
1851/1852	P57L/E188W/D489A/L823F/R862Q	+	
1855/1856	A62W/S124V/E188W/L823F/S842G/R862Q	+	
1859/1860	P57F/E614Q/S916G/S932A	+	
1869/1870	P57F/A62L/V913R/S916R/S932A	+	
1871/1872	A62L/N527R/S727W/S916G	++	
1877/1878	D500A/S727W/V913R/S916R	+	
1925/1926	D923W	+	
1969/1970	D489A/E614Q/S916G	+	
1977/1978	A62W/S727W	+	
1987/1988	A62W/D489A/N527R/S916R/S932A	+	
1989/1990	A62L/D489A/D500A/S932A	+	
1991/1992	D500A/P914R/S916G	+	
1999/2000	P57F/A62L/S916G	++	
2005/2006	P57F/A62W/F120I/N527R/V913R/S916R	+	
2013/2014	A62W/S916G	++	
2015/2016	P57F/D489A/N527R/P914R/S916G	+	
1857/1858	P57F/A62L/L305F/D500A/V913R/S916G	+++	+
1861/1862	P57F/A62L/L305F/V913R	++	+
1863/1864	P57F/A62L/A437G/D500A/L761F/P914K/S916R	+	++

Table 17-5. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 946¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 946)	TIS Reduction (Relative to SEQ ID NO: 946)	IHC Reduction (Relative to SEQ ID NO: 946)
1865/1866	A62L/A437G/V913R/S916R	+++	++
1873/1874	P57F/A62W/L305F/A437G/D500A/S727W/V913R/S916R	+++	++
1875/1876	P57F/A437G/D500A/E614Q/S727W/P914R	++	++
1953/1954	A62W/A437G/D489A/E614Q/S727W/V913R	++	++
1955/1956	A62L/A437G/D489A/N527R/S727W/V913R/S932A	+++	++
1957/1958	P57F/A62W/L305F/D489A/Q907K/V913R/S916G	++	+
1959/1960	P57F/A62W/L305F/S916R	++	+
1961/1962	P57F/A437G/D500A/N527R/S727W/S916R	+++	++
1963/1964	A62L/L305F/S727W	++	+
1965/1966	P57F/A62L/L305F/A437G/E614Q/A683S/V913R/S916R/S932A	+++	++
1967/1968	P57F/L305F	+	+
1973/1974	P57F/A62W/A437G/V913R/S916G	+++	++
1975/1976	P57F/A62W/L305F/D489A/V913R/S916G	+++	+
1981/1982	P57F/L305F/A437G/S916G	+++	++
1983/1984	A62L/A437G/N527R/S727W	++	++
1985/1986	P57F/A62L/A437G/N527R/S727W	++	++
1995/1996	A62L/L305F/A437G/D500A/S727W/V913R	+++	++
1997/1998	A437G	++	++
2001/2002	P57F/A62L/L305F/A437G/D500A/E614Q/S727W/S916R	+++	++
2003/2004	A437G/D489A/P914R/S916R	++	++
2007/2008	A62L/A437G/N527R/S916G/S932A	+++	++
2009/2010	A437G/S727W/P914K	+++	++
2011/2012	P57F/A437G/V913R/P914R	++	++
2017/2018	A437G/P914R/S916G	+++	++
¹ TIS reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 946, and defined as follows: "+" = 1-10; "++" > 10; "+++" > 20; and "++++" > 50. IHC reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 946, and defined as follows: "+" = 1-3; "++" > 3; "+++" > 5; and "++++" > 8.			

Table 17-6. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 1956¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	TIS Reduction (Relative to SEQ ID NO: 1956)	IHC Reduction (Relative to SEQ ID NO: 1956)
2021/2022	P78E/S372T/L390Q	+	
2023/2024	D87E/T266N/S372T/T483S	+	
2027/2028	T266N/S372T/T924N	+	

Table 17-6. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 1956¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	TIS Reduction (Relative to SEQ ID NO: 1956)	IHC Reduction (Relative to SEQ ID NO: 1956)
2037/2038	P78E/D87E/K176T/T266N/V536T/Q615D	+	
2041/2042	D87E/S372T/S777G	+	
2043/2044	T266N/S372T/V536T/Q615D/T763L/S777G	+	
2053/2054	P78E/T266N/T763L	+	
2055/2056	P78E/D87E/T266N/S372T/A386Y/S777G	+	
2073/2074	P78E/D87E/T266N/S372T/V536T	+	
2497/2498	A142G	+	
2501/2502	D493L	+	
2509/2510	K129E	+	
2513/2514	H499E	+	
2517/2518	L857E	+	
2089/2090	P779E	+	
2541/2542	L857A	+	
2547/2548	H499I	+	
2559/2560	A834S	+	
2565/2566	L857V	+	
2567/2568	F152W	+	
2097/2098	L857T	+	
2569/2570	F152S	+	
2571/2572	S136K	+	
2573/2574	K129S	+	
2589/2590	I156C	+	
2601/2602	H499M	+	
2617/2618	S136V	+	
2117/2118	L909F	+	
2133/2134	V797L	+	
2143/2144	I156L	+	
2145/2146	V797M	+	
2651/2652	A834W	+	
2661/2662	K267E	+	
2667/2668	E855G	+	
2151/2152	A105T	++	
2683/2684	Q768I	+	
2685/2686	V926T	++	
2693/2694	V797F	+	
2695/2696	S372D	+	
2699/2700	Y765W	+	
2161/2162	T826G	+	
2725/2726	S856G	+	
2729/2730	V797I	++	
2165/2166	T141S	+	
2751/2752	H499Q	+	

Table 17-6. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 1956¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	TIS Reduction (Relative to SEQ ID NO: 1956)	IHC Reduction (Relative to SEQ ID NO: 1956)
2171/2172	A834H	+	
2763/2764	Y125W	+	
2777/2778	V926M	+	
2779/2780	G139E	+	
2791/2792	T924A	+	
2179/2180	S136G	+	
2793/2794	V797W	++	
2795/2796	V797E	++	
2191/2192	A496G	+	
2821/2822	D401G	+	
2825/2826	A834G	+	
2859/2860	A105W	+	
2861/2862	L857S	+	
2207/2208	L693F	+	+
2209/2210	V710N	++	++
2211/2212	V710L	+	
2213/2214	M663A	++	+
2215/2216	V710M	+++	++
2217/2218	L670I	+	
2219/2220	L670F	+	
2221/2222	L670R	+	+
2223/2224	V710S	+	+
2225/2226	F588L	++	+
2229/2230	L670Q	+++	+
2231/2232	L670T	+	+
2233/2234	K154R/F588L	+	+
2235/2236	L693Y	+	+
2237/2238	V710K	+	++
2239/2240	A708C	+++	++
2247/2248	L670W	++	+
2249/2250	L670V	+	
2251/2252	L693I	++	+
2253/2254	V589I/M663F	+	
2255/2256	F588V	++	+
2257/2258	M663F	+	
2261/2262	L670E	+++	+
2263/2264	L670S	+	+
2275/2276	L629I	++	+
2277/2278	T692L	+	+
2279/2280	A711C	+	++
2281/2282	R577T	+	+
2289/2290	L628V	+	

Table 17-6. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 1956¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	TIS Reduction (Relative to SEQ ID NO: 1956)	IHC Reduction (Relative to SEQ ID NO: 1956)
2293/2294	L691V	+	
2297/2298	L569Y	+	
2299/2300	I573Q	++	+
2303/2304	T692Y	++	+
2305/2306	L629C	+++	++
2307/2308	T692I	+	+
2311/2312	S671A	+	
2315/2316	L569S	+	+
2317/2318	I573M	+	+
2321/2322	T692C	+++	+++
2323/2324	I573H	++	+
2325/2326	S671T	+	
2327/2328	I573D	+++	+
2331/2332	R577D	+++	++
2333/2334	L629A	++	+
2335/2336	T692G	++	+
2339/2340	R577A	+	+
2343/2344	T692F	++	++
2345/2346	R583G	+++	+
2351/2352	A572G	+++	+
2353/2354	A711W		++
2355/2356	L629G	+++	++
2357/2358	I573C	+++	+
2359/2360	L569V		+
2361/2362	L669R	+	
2363/2364	L579V	++	+
2365/2366	A572S	+	
2371/2372	V580E	+++	+
2373/2374	A574S	+	+
2375/2376	V631I	+	+
2377/2378	A711H	+	++
2381/2382	V580G	+++	+
2383/2384	V580W	++	+
2385/2386	R577E	+++	++
2387/2388	R583C	++	+
2391/2392	S671G	+	
2393/2394	A711G	+	++
2395/2396	V631L	++	+
2397/2398	L569T	++	+
2399/2400	L569H	++	+
2401/2402	L569Q	++	+
2403/2404	V631M	+	+

Table 17-6. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 1956¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 1956)	TIS Reduction (Relative to SEQ ID NO: 1956)	IHC Reduction (Relative to SEQ ID NO: 1956)
2407/2408	T227A	+	+
2415/2416	L871E	+++	+++
2417/2418	V222C	++	++
2421/2422	V878G	+	+
2425/2426	K581H	+	+
2431/2432	V878A	+	
2433/2434	R873Y	+	
2437/2438	K581G	++	+
2439/2440	V222P	+++	++
2443/2444	T585Q	++	+
2447/2448	A229C	+	+
2461/2462	V878K		+
2463/2464	R873A	+	
2467/2468	K581T	+	
2469/2470	I869L	+	
2473/2474	V878S	+	+
2477/2478	V878Q	+	+
2479/2480	R873F	+	
2485/2486	I869S	+++	++
2487/2488	V878W	+	+
2489/2490	H706F	+	
2491/2492	N225D	+	++
2493/2494	L871K	+	+
¹ TIS reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 1956, and defined as follows: "+" = 1-10; "++" > 10; "+++" > 20; and "++++" > 50. IHC reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 1956, and defined as follows: "+" = 1-3; "++" > 3; "+++" > 5; and "++++" > 8.			

Table 17-7. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 2496¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	TIS Reduction (Relative to SEQ ID NO: 2496)	IHC Reduction (Relative to SEQ ID NO: 2496)
2867/2868	L569T/L628M/T692Y	+++	+++
2869/2870	L569Y/L670T/A711H	++	+++
2871/2872	L569Y/L628M/L670T/T692Y/A711H/L871S	+++	++++
2873/2874	V589I/L670T/Q795E/L871S	+++	++
2875/2876	L628M/L670F	+	
2877/2878	L569H/L628M/L670T	+++	++
2879/2880	L569T/L670T/A711H/L871E	++++	++++
2881/2882	L628M/L670T/T692G/A711H/Q795E	+++	++++

Table 17-7. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 2496¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	TIS Reduction (Relative to SEQ ID NO: 2496)	IHC Reduction (Relative to SEQ ID NO: 2496)
2883/2884	L569Y/L628M/L670T/A711H	++	+++
2885/2886	L569Y/V589I/L628M/T692G/A711H/Q795E	+++	+++
2887/2888	L569H/T692G	+++	+++
2889/2890	L569Y/V589I/L670T/S678T/T692G/A711H/Q795E	+++	++++
2891/2892	L569T/L871K	++	++
2893/2894	L569Y/V589I/L628M/L670T/T692Y/A711H	+++	++++
2895/2896	V589I/L871E	+++	+++
2897/2898	L670T/S678T/T692Y/L871S	+++	+++
2899/2900	L628M/A711H/Q795E	+	++
2901/2902	T692G/A711H	+++	+++
2903/2904	V589I/L670T/T692G/Q795E/L871K	+++	+++
2905/2906	L569Y/V589I/A711H/L871K	++	++
2907/2908	L670T/T692G/L871K	+++	+++
2909/2910	L569H/T692Y	+++	+++
2911/2912	L569H/L628M/L670T/T692Y/A711H/L871K	+++	++++
2913/2914	L670T/S678T	+	+
2915/2916	L569T/L670T/S678T/T692G/L871K	+++	++++
2917/2918	V589I	+	
2919/2920	L569H/V589I/L628M/L670T/S678T/T692G/A711H/Q795E/L871S	++++	++++
2921/2922	L569H/L670F/S678T/T692G/L871S	++++	+++
2923/2924	L569T/V589I/L628M/L670F/T692G/A711H	++++	++++
2925/2926	L569H/L670T/T692G	+++	+++
2927/2928	L569Y/L871E	+++	+++
2929/2930	L569H/L628M/L670F/S678T	++	+
2931/2932	L569Y/L670F/S678T	+	
2933/2934	L569Y/V589I/L670F	++	
2935/2936	L569Y/V589I/L670T/L871K	++	+
2937/2938	L569T/L871S	+++	++
2939/2940	L871K	+	+
2941/2942	L569Y/L670T	+	+
2943/2944	L569Y/T692Y/A711H	++	+++
2945/2946	L871S	++	+
2947/2948	L569Y/A711H	+	++
2949/2950	L628M/L871S	++	+
2951/2952	L569T/V589I/L871S	+++	++
2953/2954	T692G/A711H/I869L/V878S	+++	+++
2955/2956	A572S/F588L/S678T/T692G/I869L/V878A	+++	++
2957/2958	A711H	+	++
2959/2960	T692G/I869S	+++	+++
2961/2962	A572S/T692L/I869S/V878S	+++	+++
2963/2964	L629I/I869L/V878S	++	+

Table 17-7. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 2496¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	TIS Reduction (Relative to SEQ ID NO: 2496)	IHC Reduction (Relative to SEQ ID NO: 2496)
2965/2966	A572S/F588L/Q795E	+++	+
2967/2968	A711H/I869S/V878S	+++	++++
2969/2970	V878S	+	+
2971/2972	S678T/T692G/A711H/I869S	+++	++++
2973/2974	A572S/T692G/I869S/V878S	++++	++++
2975/2976	A572S/T692G/V878S	+++	++
2977/2978	A711H/Q795E/I869S/V878S	+++	++++
2979/2980	Q795E/V878S	+	+
2981/2982	T692G/A711H/Q795E/I869S	+++	++++
2983/2984	S678T/T692G/Q795E/I869S	+++	+++
2985/2986	T692G/A711H/I869S/V878A	++++	++++
2987/2988	L569T/A711H	++	+++
2989/2990	L569T/L628M/S678T/A711H	++	+++
2991/2992	L569T/V589I/T692G/Q795E/L871K/V878S	+++	+++
2993/2994	L569T/V589I/S678T/L871K/V878S	+++	++
2995/2996	L569H/A711H/I869S/V878S	++++	++++
2997/2998	S678T/Q795E/L871K/V878A	+	+
2999/3000	L569H/S678T/T692G/Q795E	+++	+++
3001/3002	F588L/V589I/L628M/S678T	+	
3003/3004	T692Y	++	+
3005/3006	L569T	++	+
3007/3008	T692G/L871K	+++	++
3009/3010	L569T/F588L/V878S	+++	++
3011/3012	L569H/A711H/L871K	+++	+++
3013/3014	L569T/F588L/S678T/T692Y	+++	+++
3015/3016	L569T/F588L/A711H/I869L/L871K/V878S	+++	++++
3017/3018	L569T/F588L	+++	++
3019/3020	L569T/T692Y	+++	+++
3021/3022	L569T/S678T/I869S/V878S	+++	++++
3023/3024	L628M/L629I/T692Y/L871S/V878S	+++	+++
3025/3026	L569T/S678T/V878S	++	++
3027/3028	L569H/A711H/Q795E/L871S/V878S	+++	++++
3029/3030	A711H/L871K	+	++
3031/3032	T692G/A711H/Q795E/I869L/L871K/V878A	+++	+++
3033/3034	L569T/F588L/V589I/L628M/L629I/T692Y/A711H	+++	++++
3035/3036	S678T/T692G/I869S	+++	+++
3037/3038	T692G/I869L/L871K/V878S/S916R	+++	++
3039/3040	L569H/L628M	++	+
3041/3042	L628M/V878S	+	+
3043/3044	T692G/A711H/L871S/V878A	+++	++++
3045/3046	L569T/Q795E	++	+

Table 17-7. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 2496¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2496)	TIS Reduction (Relative to SEQ ID NO: 2496)	IHC Reduction (Relative to SEQ ID NO: 2496)
3047/3048	L569T/T692Y/A711H/I869L/L871K/V878A	+++	++++
3065/3066	L810V	+	++
3049/3050	S246T	++	+
3067/3068	A60V/V589A	+++	+
3051/3052	T585K	+	+
3069/3070	S307T		+
3053/3054	G584E	+	+
3055/3056	A812E	++	++
3057/3058	V313L	+	+++
3059/3060	L304M		+
3071/3072	V313T	++	+++
3061/3062	V313I	+	++
3073/3074	G584C	+	+
3063/3064	A582T	+	+
¹ TIS reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 2496, and defined as follows: "+" = 1-10; "++" > 10; "+++" > 20; and "++++" > 50. IHC reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 2496, and defined as follows: "+" = 1-3; "++" > 3; "+++" > 5; and "++++" > 8.			

Table 17-8. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 2880¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2880)	TIS Reduction (Relative to SEQ ID NO: 2880)	IHC Reduction (Relative to SEQ ID NO: 2880)
3075/3076	R527N/L736M/A932S	+	
3107/3108	A89R/L109D/A932S	+	
3109/3110	L109D/S842G	+	
3123/3124	L109D/A932S	+	
3127/3128	R527N/W727S/L736M	+	
3141/3142	L109D/L736M/A932S	++	
3143/3144	A89R/L109D/R527N/S678T/W727S/S842G	+	
3181/3182	V522E/K830Q	+	
3269/3270	L62A/E87D/S150T	+	
3077/3078	L109D/S678T/A812E	++	++
3081/3082	R403H/R527N/S678T/T692G/L736M/A812E/S842G/L860F	+++	+++
3083/3084	S678T/T692G/A812E	+++	+++
3087/3088	A89R/R527N/W727S/A812E/L860F	++	++
3089/3090	A89R/S678T/T692G/L736M/A932S	+++	+
3091/3092	L109D/R527N/S678T/A812E	++	++
3095/3096	L109D/A812E	++	++

Table 17-8. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 2880¹			
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 2880)	TIS Reduction (Relative to SEQ ID NO: 2880)	IHC Reduction (Relative to SEQ ID NO: 2880)
3097/3098	T692G/W727S/L736M/S842G/R913V	+++	+
3101/3102	R527N/S678T/T692G/W727S/L736M/V878S	+++	+
3103/3104	R527N/T692G/W727S/L736M/A812E	+++	+++
3113/3114	A89R/L109D/S678T/W727S/L736M/A812E/V878S	++	++
3115/3116	L109D/T692G/W727S/A812E/S842G/L860F	+++	+++
3117/3118	T692G/A812E	+++	+++
3121/3122	L109D/S678T/L736M/A812E/V878S	+++	++
3125/3126	R527N/T692G/W727S/L736M/S842G/L860F/V878S	+++	+
3133/3134	R527N/S678T/T692G/A812E/A932S	+++	+++
3135/3136	A89R/S678T/A812E/V878S	+	++
3137/3138	R527N/A812E	++	++
3139/3140	S678T/T692G/W727S/A812E/S842G	+++	+++
3149/3150	L109D/S678T/T692G/S842G/L860F/V878S/A932S	++	+
3151/3152	A89R/R527N/S678T/T692G/L736M/S842G/V878S/A932S	+++	+
3155/3156	L109D/T692G/W727S/L736M/A812E	+++	+++
3161/3162	S678T/A812E	++	++
¹ TIS reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 2880, and defined as follows: "+" = 1-10; "++" > 10; "+++" > 20; and "++++" > 50. IHC reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 2880, and defined as follows: "+" = 1-3; "++" > 3; "+++" > 5; and "++++" > 8.			

Table 17-9. Reduction of Total Immunogenicity Score (TIS) and Immunogenic Hit Count (IHC) for GAA Variants Relative to SEQ ID NO: 3104¹		
SEQ ID NO: (nt/aa)	Amino Acid Differences (Relative to SEQ ID NO: 3104)	TIS Reduction (Relative to SEQ ID NO: 3104)
3313/3314	L62A/A89D/K830Q	+
3317/3318	L62A/K830Q/A932S	+
3361/3362	L62A/K830Q/L860F/A932S	+
3367/3368	L62A/K830Q/L860F	+
3375/3376	L62A/K830Q	+
¹ TIS reduction was measured as a reduction in the number of counts compared to the reference polypeptide of SEQ ID NO: 3104, and defined as follows: "+" = 1-10.		

[0225] While the invention has been described with reference to the specific embodiments, various changes can be made and equivalents can be substituted to adapt to a particular situation, material,

composition of matter, process, process step or steps, thereby achieving benefits of the invention without departing from the scope of what is claimed.

[0226] For all purposes in the United States of America, each and every publication and patent document cited in this disclosure is incorporated herein by reference as if each such publication or document was specifically and individually indicated to be incorporated herein by reference. Citation of publications and patent documents is not intended as an indication that any such document is pertinent prior art, nor does it constitute an admission as to its contents or date.

CLAIMS

WHAT IS CLAIMED IS:

1. A recombinant acid alpha glucosidase and/or biologically active recombinant acid alpha glucosidase fragment comprising an amino acid sequence comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

2. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 2, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 27, 27/944, 28, 29/478, 30, 88, 107, 109, 109/842, 110, 113, 135, 137, 138, 148, 150, 247, 274, 276, 278, 375, 403, 414, 418, 418/499, 421, 426, 437, 444, 455, 463, 471, 471/478, 476, 489, 527, 547, 581, 610, 642, 668, 670, 692, 725/732, 750, 753, 786, 820, 862, 871, 895, 897, 930, 934, and 944, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 2.

3. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 2, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 29/218/240/668/700/744/869, 29/218/240/700/869, 29/240/596/668/700/744/869, 29/240/596/668/869, 36/106/150/218/527/750/883/894, 106/112/150/218/414/527/793/883, 106/150/169/218/414/486/527/750/894, 106/150/169/218/414/486/527/894, 106/150/169/218/414/486/749/793/883/894, 106/150/169/218/414/486/750/793/883/894, 106/150/169/218/414/486/793/883, 106/150/169/218/414/486/894, 106/150/169/218/414/749/750/793/883, 106/150/169/218/414/749/793, 106/150/169/218/414/749/793/883, 106/150/169/218/486/527/749/793/894, 106/150/169/218/486/749/883, 106/150/169/218/486/883, 106/150/169/218/749/800, 106/150/169/414/486/749/750/883, 106/150/169/527/749/793/883, 106/150/169/749/793/883/894, 106/150/218/331/414/486/527/733/749/793, 106/150/218/414/486/642/750/793/883, 106/150/218/414/486/750/793/894, 106/150/218/414/527/749/750/883, 106/150/218/414/527/749/793/883/894, 106/150/218/414/749/750/793/883/894, 106/150/218/414/749/793/883, 106/150/218/486/527/749/894, 106/150/218/486/793/883,

106/150/218/527/749/750/793, 106/150/218/527/793/894, 106/150/218/749/750/793,
 106/150/218/793, 106/150/218/793/894, 106/150/245/793/883/894, 106/150/414/749/750/793/894,
 106/150/414/749/793/894, 106/150/486/527/750/793, 106/150/486/749/793/883/894,
 106/150/749/793/883, 106/169/185/218/414/749/750/793, 106/191/280/402/414/444/727,
 106/191/414/444/522/928/944, 106/191/414/489/928/944, 106/280/402/414/444/489/727/944,
 150/169/218/414/527/793, 150/218/414/486/749/750, 150/218/414/486/750/793,
 150/218/414/486/750/793/883, 150/218/414/749/750/793/894, 150/218/414/749/793,
 150/218/527/749/793, 150/218/749/750/793, 150/218/749/793, 150/414/486/527/750/894,
 150/414/486/749/750/793, 150/486/750/883/894, 169/486/750/793/883,
 180/275/402/518/547/610/638/669/671, 180/402/431/507/547/610/669/671/793,
 180/402/507/547/610/671, 191/280/402/414/444/465/842/928, 191/280/402/414/444/489/500/944,
 191/280/414/444/489/500/522/842/928/944, 191/280/414/444/489/522/727/944,
 191/280/414/489/842/928/944, 191/280/414/944, 191/414/522/842/944, 196/402/431/547/610/638,
 218/668/700/869, 224/402/507/518/547/638/668, 269/275/431/518/547/638/668/669,
 275/281/402/431/507/518/610/668, 275/281/402/431/518/547/610/669/671,
 275/281/402/507/518/547/638/669/671, 275/281/402/518/547/610/638/671,
 275/281/402/518/547/610/668/669/887, 275/281/402/547/610/638/669/671,
 275/281/431/518/547/638/669/671, 275/281/507/547/669/671, 275/281/610/638/668/669,
 275/281/671, 275/377/402/507/518/669/671/715, 275/402/431/507/547/671,
 275/402/431/518/610/638/669/671/922, 275/402/507/547/610/638/668/669,
 275/402/507/547/610/638/669/671, 275/402/507/547/610/671, 275/402/547/610/638/669/671,
 275/402/547/638/669/671, 275/402/638/669/671, 275/431/507/518/547/668/669/671,
 275/431/507/518/610/669/671, 275/431/507/547/610/638/671, 275/431/518/547/638/668,
 275/431/518/610/638/669/671, 275/431/638, 275/507/518/547/610/638/668/669,
 275/507/518/547/638/669/671, 275/507/547/610/638/669/671, 275/507/547/668/669/671,
 275/518/671, 280/402/536/928, 281/402/507/518/547/610/638/669/671,
 281/402/507/547/638/669/671, 281/402/518/547/610/638/668/669, 281/402/518/547/668,
 281/431/507/518/547/610/638/668, 402/431/518/547/610/668, 402/431/518/547/671,
 402/431/518/610, 402/431/547/638/671, 431/507/518/541/547/638/669/671, 431/507/518/669/671,
 507/547/610, 507/547/638/669/671, 547/610/638/671, and 547/638/668, wherein the amino acid
 positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 2.

4. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 2, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 27, 27/944, 28,

29/218/240/668/700/744/869, 29/218/240/700/869, 29/240/596/668/700/744/869,
29/240/596/668/869, 29/478, 30, 36/106/150/218/527/750/883/894,
106/112/150/218/414/527/793/883, 106/150/169/218/414/486/527/750/894,
106/150/169/218/414/486/527/894, 106/150/169/218/414/486/749/793/883/894,
106/150/169/218/414/486/750/793/883/894, 106/150/169/218/414/486/793/883,
106/150/169/218/414/486/894, 106/150/169/218/414/749/750/793/883,
106/150/169/218/414/749/793, 106/150/169/218/414/749/793/883, 106/150/169/218/486/749/883,
106/150/169/218/486/883, 106/150/169/414/486/749/750/883, 106/150/169/527/749/793/883,
106/150/169/749/793/883/894, 106/150/218/331/414/486/527/733/749/793,
106/150/218/414/486/642/750/793/883, 106/150/218/414/486/750/793/894,
106/150/218/414/527/749/750/883, 106/150/218/414/527/749/793/883/894,
106/150/218/414/749/750/793/883/894, 106/150/218/414/749/793/883,
106/150/218/486/527/749/894, 106/150/218/486/793/883, 106/150/218/527/793/894,
106/150/245/793/883/894, 106/150/414/749/750/793/894, 106/150/414/749/793/894,
106/150/486/527/750/793, 106/150/486/749/793/883/894, 106/150/749/793/883,
106/169/185/218/414/749/750/793, 106/191/280/402/414/444/727,
106/280/402/414/444/489/727/944, 107, 109, 109/842, 110, 135, 138, 148, 150,
150/218/414/486/749/750, 150/218/414/486/750/793, 150/218/414/486/750/793/883,
150/218/414/749/750/793/894, 150/218/414/749/793, 150/414/486/527/750/894,
150/414/486/749/750/793, 150/486/750/883/894, 169/486/750/793/883,
180/275/402/518/547/610/638/669/671, 180/402/431/507/547/610/669/671/793,
180/402/507/547/610/671, 191/280/402/414/444/489/500/944, 191/280/414/444/489/522/727/944,
191/280/414/944, 191/414/522/842/944, 196/402/431/547/610/638, 218/668/700/869,
224/402/507/518/547/638/668, 269/275/431/518/547/638/668/669, 274,
275/281/402/431/507/518/610/668, 275/281/402/431/518/547/610/669/671,
275/281/402/507/518/547/638/669/671, 275/281/402/518/547/610/638/671,
275/281/402/518/547/610/668/669/887, 275/281/402/547/610/638/669/671,
275/281/431/518/547/638/669/671, 275/281/507/547/669/671, 275/281/610/638/668/669,
275/377/402/507/518/669/671/715, 275/402/431/507/547/671,
275/402/431/518/610/638/669/671/922, 275/402/507/547/610/638/668/669,
275/402/507/547/610/638/669/671, 275/402/507/547/610/671, 275/402/547/610/638/669/671,
275/402/547/638/669/671, 275/402/638/669/671, 275/431/507/518/547/668/669/671,
275/431/507/518/610/669/671, 275/431/507/547/610/638/671, 275/431/518/547/638/668,
275/431/518/610/638/669/671, 275/431/638, 275/507/518/547/610/638/668/669,
275/507/518/547/638/669/671, 275/507/547/610/638/669/671, 275/507/547/668/669/671, 276,
281/402/507/518/547/610/638/669/671, 281/402/507/547/638/669/671,
281/402/518/547/610/638/668/669, 281/402/518/547/668, 281/431/507/518/547/610/638/668, 375,

402/431/518/547/610/668, 402/431/518/547/671, 402/431/518/610, 402/431/547/638/671, 403, 414, 418/499, 431/507/518/541/547/638/669/671, 431/507/518/669/671, 437, 471/478, 507/547/610, 507/547/638/669/671, 547, 547/610/638/671, 547/638/668, 581, 642, 670, 692, 750, 753, 820, 871, and 944, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 2.

5. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 6, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 27, 27/944, 28, 29/218/240/668/700/744/869, 29/218/240/700/869, 29/240/596/668/700/744/869, 29/240/596/668/869, 29/478, 30, 36/106/150/218/527/750/883/894, 88, 106/112/150/218/414/527/793/883, 106/150/169/218/414/486/527/750/894, 106/150/169/218/414/486/527/894, 106/150/169/218/414/486/749/793/883/894, 106/150/169/218/414/486/750/793/883/894, 106/150/169/218/414/486/793/883, 106/150/169/218/414/486/894, 106/150/169/218/414/749/750/793/883, 106/150/169/218/414/749/793, 106/150/169/218/414/749/793/883, 106/150/169/218/486/527/749/793/894, 106/150/169/218/486/749/883, 106/150/169/218/486/883, 106/150/169/218/749/800, 106/150/169/414/486/749/750/883, 106/150/169/527/749/793/883, 106/150/169/749/793/883/894, 106/150/218/331/414/486/527/733/749/793, 106/150/218/414/486/642/750/793/883, 106/150/218/414/486/750/793/894, 106/150/218/414/527/749/750/883, 106/150/218/414/527/749/793/883/894, 106/150/218/414/749/750/793/883/894, 106/150/218/414/749/793/883, 106/150/218/486/527/749/894, 106/150/218/486/793/883, 106/150/218/527/749/750/793, 106/150/218/527/793/894, 106/150/218/749/750/793, 106/150/218/793, 106/150/218/793/894, 106/150/245/793/883/894, 106/150/414/749/750/793/894, 106/150/414/749/793/894, 106/150/486/527/750/793, 106/150/486/749/793/883/894, 106/150/749/793/883, 106/169/185/218/414/749/750/793, 106/191/280/402/414/444/727, 106/191/414/444/522/928/944, 106/191/414/489/928/944, 106/280/402/414/444/489/727/944, 107, 109, 109/842, 110, 113, 135, 137, 138, 148, 150, 150/169/218/414/527/793, 150/218/414/486/749/750, 150/218/414/486/750/793, 150/218/414/486/750/793/883, 150/218/414/749/750/793/894, 150/218/414/749/793, 150/218/527/749/793, 150/218/749/750/793, 150/218/749/793, 150/414/486/527/750/894, 150/414/486/749/750/793, 150/486/750/883/894, 169/486/750/793/883, 180/275/402/518/547/610/638/669/671, 180/402/431/507/547/610/669/671/793, 180/402/507/547/610/671, 191/280/402/414/444/465/842/928, 191/280/402/414/444/489/500/944, 191/280/414/444/489/500/522/842/928/944, 191/280/414/444/489/522/727/944,

191/280/414/489/842/928/944, 191/280/414/944, 191/414/522/842/944, 196/402/431/547/610/638, 218/668/700/869, 224/402/507/518/547/638/668, 247, 269/275/431/518/547/638/668/669, 274, 275/281/402/431/507/518/610/668, 275/281/402/431/518/547/610/669/671, 275/281/402/507/518/547/638/669/671, 275/281/402/518/547/610/638/671, 275/281/402/518/547/610/668/669/887, 275/281/402/547/610/638/669/671, 275/281/431/518/547/638/669/671, 275/281/507/547/669/671, 275/281/610/638/668/669, 275/281/671, 275/377/402/507/518/669/671/715, 275/402/431/507/547/671, 275/402/431/518/610/638/669/671/922, 275/402/507/547/610/638/668/669, 275/402/507/547/610/638/669/671, 275/402/507/547/610/671, 275/402/547/610/638/669/671, 275/402/547/638/669/671, 275/402/638/669/671, 275/431/507/518/547/668/669/671, 275/431/507/518/610/669/671, 275/431/507/547/610/638/671, 275/431/518/547/638/668, 275/431/518/610/638/669/671, 275/431/638, 275/507/518/547/610/638/668/669, 275/507/518/547/638/669/671, 275/507/547/610/638/669/671, 275/507/547/668/669/671, 275/518/671, 276, 278, 280/402/536/928, 281/402/507/518/547/610/638/669/671, 281/402/507/547/638/669/671, 281/402/518/547/610/638/668/669, 281/402/518/547/668, 281/431/507/518/547/610/638/668, 375, 402/431/518/547/610/668, 402/431/518/547/671, 402/431/518/610, 402/431/547/638/671, 403, 414, 418, 418/499, 421, 426, 431/507/518/541/547/638/669/671, 431/507/518/669/671, 437, 444, 455, 463, 471, 471/478, 476, 489, 507/547/610, 507/547/638/669/671, 527, 547, 547/610/638/671, 547/638/668, 581, 610, 642, 668, 670, 692, 725/732, 750, 753, 786, 820, 862, 871, 895, 897, 934, and 944, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 6.

6. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 8, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 4, 27, 27/28/489, 27/418/478, 28, 28/29, 28/29/113/135/138, 28/29/113/135/418, 28/29/135, 28/29/418, 29/113/126/135/193, 29/113/135, 29/113/135/455, 29/113/138, 29/148, 29/478, 106, 106/138/218/431/671/749, 106/218/281, 106/218/455, 106/218/455/507/749, 106/489/671, 106/638, 106/671/934, 113, 113/135/418, 113/418/455/478/581, 113/418/478/489/581, 135, 135/148/150/418, 135/478/489/581, 135/489, 135/944, 138/218/668/671, 138/218/749/934, 138/671/749/934, 157, 218, 218/281, 218/281/431, 218/281/671, 218/431, 218/431/489/507/749/934, 218/455, 218/507/749, 218/507/934, 218/638/671, 218/749, 281/431/489/668, 345/934, 418, 418/489, 431/668/671, 489/638/934, 489/671/934, 489/749, 489/934, 507/668, 507/671/934, 671/749, 671/934, and 749/784, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 8.

7. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 8, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 27, 27/418/478, 28, 28/29, 28/29/113/135/138, 28/29/113/135/418, 28/29/135, 28/29/418, 29/113/126/135/193, 29/113/135, 29/113/135/455, 29/113/138, 29/148, 29/478, 106, 106/138/218/431/671/749, 106/489/671, 106/638, 113/135/418, 113/418/455/478/581, 113/418/478/489/581, 135, 135/148/150/418, 135/478/489/581, 135/944, 138/218/668/671, 157, 218/638/671, 418, 418/489, 431/668/671, 507/668, and 671/749, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 8.

8. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 12, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 4, 27, 27/28/489, 27/418/478, 28, 28/29, 28/29/113/135/138, 28/29/113/135/418, 28/29/135, 28/29/418, 29/113/126/135/193, 29/113/135, 29/113/135/455, 29/113/138, 29/148, 29/478, 106, 106/138/218/431/671/749, 106/218/281, 106/218/455, 106/218/455/507/749, 106/489/671, 106/638, 106/671/934, 113, 113/135/418, 113/418/455/478/581, 113/418/478/489/581, 135, 135/148/150/418, 135/478/489/581, 135/489, 135/944, 138/218/668/671, 138/218/749/934, 138/671/749/934, 157, 218, 218/281/431, 218/281/671, 218/431, 218/455, 218/507/749, 218/638/671, 218/749, 281/431/489/668, 345/934, 418, 418/489, 431/668/671, 489/638/934, 489/671/934, 489/749, 489/934, 507/668, 507/671/934, 671/749, 671/934, and 749/784, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 12.

9. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 14, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 22, 24, 27, 27/165, 30, 33, 34, 37/62, 37/62/79/196/696/862, 37/62/523, 37/62/523/793, 37/64/66/79/154/523/681/793/862, 37/79/154/793, 37/196, 37/528/696/793, 37/528/790, 37/528/790/793/862, 37/790/793, 39, 39/58/489/725/830/842/930/944, 39/70/109/830/842, 39/70/489/612, 39/70/725, 39/267, 39/267/489/522/612/830/842, 39/267/489/830/944, 39/489/500/612, 39/500/612, 40, 44/157, 47, 49,

50, 55, 60/500/612, 62/79/154/862, 62/79/196/681/862, 62/79/523/528/790, 62/79/790/793, 62/79/862, 62/92, 62/92/790/793, 62/106/523/528/696/793/862, 62/154/696/793/862, 62/793/862, 68, 70, 70/267/725/944, 70/267/930/944, 70/489/930, 70/725/830/860/930/944, 77, 79/154/681, 79/154/793/862, 79/862, 89, 97, 106/154, 107, 109, 109/522/612/725, 109/522/830/944, 109/612, 118, 149, 157, 158, 178, 179, 196/528/681/790/793, 207, 208, 217, 267/489/500/725/830/930, 267/522/725, 352, 385, 424, 448, 463, 489/830/944, 500, 500/612/830/860, 500/860/930, 500/930/944, 522/725, 523, 523/790/793, 528/681, 528/793, 528/862, 672, 673, 725, 734, 740, 753, 774, 778, 793, 830, 844, 862, 875, 880, 892, 902, 922, 925, 930, 932, 934, 938, and 944, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 14.

10. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 18, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 22, 24, 27, 27/165, 30, 33, 34, 37/62, 37/62/79/196/696/862, 37/62/523, 37/196, 37/528/790, 39, 39/70/109/830/842, 39/70/725, 39/267, 39/267/489/522/612/830/842, 39/267/489/830/944, 40, 70/267/725/944, 70/267/930/944, 70/489/930, 107, 109, 109/522/830/944, 217, 267/489/500/725/830/930, 267/522/725, 352, 385, 500/930/944, 673, 734, 774, 778, 875, 930, 932, and 934, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 18.

11. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 20, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 22/24, 22/24/27/50/107/109/489/612/725, 22/24/27/489, 22/24/27/489/612/774, 22/24/27/612/944, 22/24/50/109/267/489/774/944, 22/24/50/267/612/922/944, 22/24/107/267/489/922, 22/24/489, 22/24/612/725/944, 22/50/107/267/489/612/944, 22/50/109/267/489, 22/267/489/612, 24, 24/27/50/107/267/774/944, 24/27/89/500/842, 24/27/107/267/612/944, 24/27/267/944, 24/27/500/842, 24/27/500/842/932, 24/27/944, 24/39/49/89/97/842/932, 24/39/68/89/107/500/842, 24/39/89/97/842/932, 24/39/842/932, 24/50/489/944, 24/50/612, 24/70/107/109/489/612/725, 24/70/267/774, 24/89/500, 24/107/109/267/489/612/725/774, 24/109/612, 24/109/944, 24/267/725/944, 24/489/944, 24/725, 24/842/932, 24/944, 27/39/49/97/500/842, 27/49/68/500/842, 34/39/500/932, 39/89/97/500, 42, 48, 50/109/489/612, 50/489/774, 50/612/944, 57, 62, 68, 68/89/97/932, 71, 88, 89/97/107, 89/97/500, 89/842, 107/109, 107/500/842, 108, 109/612/774/944, 112, 123, 124, 148, 188, 193, 197, 204, 253, 264, 305, 312, 333, 381, 402, 402/781, 489, 489/944,

500/842, 500/932, 523, 527, 612, 612/725/944, 612/922, 614, 727, 742, 748, 820, 823, 832, 842/932, 858, 862, 911, 913, 914, 916, 923, 937, and 940, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 20.

12. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 20, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 22/24, 22/24/27/50/107/109/489/612/725, 22/24/27/489, 22/24/27/489/612/774, 22/24/27/612/944, 22/24/50/109/267/489/774/944, 22/24/50/267/612/922/944, 22/24/107/267/489/922, 22/24/489, 22/24/612/725/944, 22/50/107/267/489/612/944, 22/50/109/267/489, 22/267/489/612, 24, 24/27/50/107/267/774/944, 24/27/89/500/842, 24/27/107/267/612/944, 24/27/267/944, 24/27/500/842, 24/27/500/842/932, 24/27/944, 24/39/49/89/97/842/932, 24/39/68/89/107/500/842, 24/39/89/97/842/932, 24/39/842/932, 24/50/489/944, 24/50/612, 24/70/107/109/489/612/725, 24/70/267/774, 24/89/500, 24/107/109/267/489/612/725/774, 24/109/612, 24/109/944, 24/267/725/944, 24/489/944, 24/725, 24/842/932, 24/944, 27/39/49/97/500/842, 27/49/68/500/842, 34/39/500/932, 50/109/489/612, 50/612/944, 68/89/97/932, 89/97/107, 89/842, 106, 107/109, 107/500/842, 108, 109/612/774/944, 112, 148, 148/772, 188/377, 238, 240, 240/374, 243, 244, 246, 248, 249/777, 252, 253, 259, 260, 261, 262, 264, 279, 305, 309, 312, 319, 320, 329, 333, 387, 402, 421, 432, 500/842, 500/932, 556, 612, 612/725/944, 612/922, 727, 736, 737, 741, 742, 748, 815, 816, 818, 823, 832, 842/932, 911, 913, 914, 916, 923, 937, and 940, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 20.

13. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 20, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 29, 42, 45, 48, 57, 61, 62, 63, 71, 88, 91, 106, 108, 112, 123, 124, 127/914, 148, 148/772, 154, 186, 188, 188/377, 193, 195, 197, 204, 205, 237, 238, 240, 240/374, 243, 244, 246, 248, 249, 249/777, 251, 252, 253, 255, 259, 260, 261, 262, 264, 279, 298/421, 305, 309, 312, 319, 320, 323, 325, 329, 333, 381, 387, 402, 421, 432, 445, 455, 523, 527, 556, 614, 682, 727, 736, 737, 741, 748, 815, 816, 818, 820, 823, 832, 858, 862, 911, 913, 914, 916, 937, and 940, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 20.

14. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 946, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 19/124/149/381/727, 24/39/489/862, 24/57/62/89/489/823/862, 24/57/823/862, 24/62/89/188/823/842/862, 24/89/489, 24/89/489/727/862, 24/489/500/842, 39/57/62/188/500/842, 39/57/500/862, 57, 57/62/120/527/913/916, 57/62/305/437/500/614/727/916, 57/62/305/437/500/727/913/916, 57/62/305/437/614/683/913/916/932, 57/62/305/489/907/913/916, 57/62/305/489/913/916, 57/62/305/500/913/916, 57/62/305/913, 57/62/305/916, 57/62/437/500/761/914/916, 57/62/437/527/727, 57/62/437/913/916, 57/62/913/916/932, 57/62/916, 57/188/489/823/862, 57/305, 57/305/437/916, 57/437/500/527/727/916, 57/437/500/614/727/914, 57/437/913/914, 57/489/527/914/916, 57/614/916/932, 62/89, 62/89/124/148/381/858, 62/89/124/381/858, 62/89/148/381/614/858, 62/89/148/923, 62/89/149/381, 62/89/149/381/832, 62/89/188/489/500/727/823, 62/89/381, 62/89/381/858, 62/89/381/923, 62/89/858, 62/96/614, 62/124/148/149/381/614, 62/124/149/381/832/858/937, 62/124/188/823/842/862, 62/124/381/832, 62/148/149/381/858/937, 62/148/381/614/937, 62/148/381/727, 62/148/381/858, 62/149/381/614/937, 62/149/381/858/937, 62/149/727, 62/305/437/500/727/913, 62/305/727, 62/381, 62/437/489/527/727/913/932, 62/437/489/614/727/913, 62/437/527/727, 62/437/527/916/932, 62/437/913/916, 62/489/500/932, 62/489/527/916/932, 62/489/614/916, 62/500, 62/527, 62/527/727/916, 62/614, 62/727, 62/916, 89/148/149, 89/148/149/381, 89/381, 124/148/381/727/858/937, 124/381/614, 124/500/842/862, 124/832/937, 148/832/858/937, 381, 381/614/832, 381/858/937, 437, 437/489/914/916, 437/727/914, 437/914/916, 489/500, 489/614/916, 500/727/913/916, 500/914/916, and 923, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 946.

15. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 946, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 57/62/120/527/913/916, 57/62/305/437/500/614/727/916, 57/62/305/437/500/727/913/916, 57/62/305/437/614/683/913/916/932, 57/62/305/489/907/913/916, 57/62/305/489/913/916, 57/62/305/500/913/916, 57/62/305/913, 57/62/305/916, 57/62/437/500/761/914/916, 57/62/437/527/727, 57/62/437/913/916, 57/62/913/916/932, 57/62/916, 57/188/489/823/862, 57/305, 57/305/437/916, 57/437/500/527/727/916, 57/437/500/614/727/914, 57/437/913/914, 57/489/527/914/916, 57/614/916/932, 62/89/188/489/500/727/823, 62/124/188/823/842/862,

62/305/437/500/727/913, 62/305/727, 62/437/489/527/727/913/932, 62/437/489/614/727/913, 62/437/527/727, 62/437/527/916/932, 62/437/913/916, 62/489/500/932, 62/489/527/916/932, 62/527/727/916, 62/727, 62/916, 124/500/842/862, 437, 437/489/914/916, 437/727/914, 437/914/916, 489/614/916, 500/727/913/916, 500/914/916, and 923, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 946.

16. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 1956, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 3/569, 41/53, 44, 44/347, 56, 65, 78, 78/87/176/266/536/615, 78/87/266, 78/87/266/372/386/777, 78/87/266/372/536, 78/87/266/483/924, 78/87/483/777, 78/87/536, 78/266/483/536/615, 78/266/483/795, 78/266/763, 78/372/390, 78/390, 78/536/615, 87, 87/266, 87/266/372/483, 87/266/483, 87/266/924, 87/372/777, 87/536/777, 87/615, 87/795, 105, 136, 141, 145, 154/588, 156, 157, 199, 202, 222, 225, 227, 229, 266, 266/372/536/615/763/777, 266/372/924, 266/536/615/795, 344, 348, 390/615, 412, 423, 425/678/894, 430, 446, 484, 488, 496, 499/711, 503, 530, 543, 569, 572, 573, 574, 577, 578, 579, 580, 581, 583, 585, 588, 589/663, 615, 628, 629, 631, 633, 656, 663, 669, 670, 671, 678, 679, 687, 690, 691, 692, 693, 705, 706, 708, 709, 710, 711, 726, 768, 773, 777, 779, 795, 797, 816, 826, 834, 857, 859, 868, 869, 871, 873, 877, 878, and 909, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 1956.

17. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 1956, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 44, 53, 56, 63, 65, 105, 125, 129, 136, 139, 141, 142, 145, 152, 156, 162, 176, 177, 185, 186, 187, 199, 199/775, 202, 265, 267, 337, 344, 348, 350, 354, 372, 373, 401, 412, 446, 469, 484, 488, 493, 496, 499, 503, 526, 543, 612, 615, 649, 677, 678, 679, 730, 752, 765, 768, 773, 777, 779, 788, 797, 822, 826, 834, 855, 856, 857, 859, 860, 924, 926, 931, and 936, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 1956.

18. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 1956, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at

least one substitution or substitution set at one or more positions selected from 78/87/176/266/536/615, 78/87/266/372/386/777, 78/87/266/372/536, 78/266/763, 78/372/390, 87/266/372/483, 87/372/777, 105, 125, 129, 136, 139, 141, 142, 152, 154/588, 156, 222, 225, 227, 229, 266/372/536/615/763/777, 266/372/924, 267, 372, 401, 493, 496, 499, 569, 572, 573, 574, 577, 579, 580, 581, 583, 585, 588, 589/663, 628, 629, 631, 663, 669, 670, 671, 691, 692, 693, 706, 708, 710, 711, 765, 768, 779, 797, 826, 834, 855, 856, 857, 869, 871, 873, 878, 909, 924, and 926, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 1956.

19. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 2496, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 246, 304, 313, 569, 569/588, 569/588/589/628/629/692/711, 569/588/678/692, 569/588/711/869/871/878, 569/588/878, 569/589/628/670/678/692/711/795/871, 569/589/628/670/692/711, 569/589/628/692/711/795, 569/589/670, 569/589/670/678/692/711/795, 569/589/670/871, 569/589/678/871/878, 569/589/692/795/871/878, 569/589/711/871, 569/589/871, 569/628, 569/628/670, 569/628/670/678, 569/628/670/692/711/871, 569/628/670/711, 569/628/678/711, 569/628/692, 569/670, 569/670/678, 569/670/678/692/871, 569/670/692, 569/670/711, 569/670/711/871, 569/678/692/795, 569/678/869/878, 569/678/878, 569/692, 569/692/711, 569/692/711/869/871/878, 569/711, 569/711/795/871/878, 569/711/869/878, 569/711/871, 569/795, 569/871, 572/588/678/692/869/878, 572/588/795, 572/692/869/878, 572/692/878, 582, 584, 585, 588/589/628/678, 589, 589/670/692/795/871, 589/670/795/871, 589/871, 628/629/692/871/878, 628/670, 628/670/692/711/795, 628/711/795, 628/871, 628/878, 629/869/878, 670/678, 670/678/692/871, 670/692/871, 678/692/711/869, 678/692/795/869, 678/692/869, 678/795/871/878, 692, 692/711, 692/711/795/869, 692/711/795/869/871/878, 692/711/869/878, 692/711/871/878, 692/869, 692/869/871/878/916, 692/871, 711, 711/795/869/878, 711/869/878, 711/871, 795/878, 812, 871, and 878, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 2496.

20. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 2496, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 60/589, 307, 313, 584,

and 810, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 2496.

21. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 2496, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 60/589, 246, 304, 307, 313, 569, 569/588, 569/588/589/628/629/692/711, 569/588/678/692, 569/588/711/869/871/878, 569/588/878, 569/589/628/670/678/692/711/795/871, 569/589/628/670/692/711, 569/589/628/692/711/795, 569/589/670, 569/589/670/678/692/711/795, 569/589/670/871, 569/589/678/871/878, 569/589/692/795/871/878, 569/589/711/871, 569/589/871, 569/628, 569/628/670, 569/628/670/678, 569/628/670/692/711/871, 569/628/670/711, 569/628/678/711, 569/628/692, 569/670, 569/670/678, 569/670/678/692/871, 569/670/692, 569/670/711, 569/670/711/871, 569/678/692/795, 569/678/869/878, 569/678/878, 569/692, 569/692/711, 569/692/711/869/871/878, 569/711, 569/711/795/871/878, 569/711/869/878, 569/711/871, 569/795, 569/871, 572/588/678/692/869/878, 572/588/795, 572/692/869/878, 572/692/878, 582, 584, 585, 588/589/628/678, 589, 589/670/692/795/871, 589/670/795/871, 589/871, 628/629/692/871/878, 628/670, 628/670/692/711/795, 628/711/795, 628/871, 628/878, 629/869/878, 670/678, 670/678/692/871, 670/692/871, 678/692/711/869, 678/692/795/869, 678/692/869, 678/795/871/878, 692, 692/711, 692/711/795/869, 692/711/795/869/871/878, 692/711/869/878, 692/711/871/878, 692/869, 692/869/871/878/916, 692/871, 711, 711/795/869/878, 711/869/878, 711/871, 795/878, 810, 812, 871, and 878, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 2496.

22. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 2880, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 24/28/29/39/50/62/78/87/135/150/266/267/522/527/551/670/727/750/830/842/871/883/894/932, 24/28/39/50/62/78/87/135/150/266/267/522/527/551/569/727/830/842/871/883/894/913, 24/28/50/135/150/437/522/527/871/883/894/932, 24/28/62/522/569/932, 24/28/437/486/527, 24/29/39/50/62/78/87/135/150/267/437/486/522/527/551/711/727/750/830/842/871/883/894/913/932, 24/50/78/87/135/150/267/486/522/527/551/670/727/750/830/842/871/883/894/913/932, 24/50/486/527/711/727, 24/62/87/486/727, 24/62/727/830/932, 24/87/135/522/670/711/830/842/913, 24/150/522/527/727/883/894, 24/527/727/842/871/883/913/932, 24/670/727/750/842/871, 28,

28/50/78/87/135/266/267/437/486/527/551/670/727/750/830/842/871/883/894/913/932,
 28/50/522/527/711/727/871, 28/62, 28/62/267/932, 28/437/527/871, 28/522/527/569/711/830/894,
 28/727, 28/727/871, 29/39/50/62/65/78/87/135/150/437/551/569/670/727/750/830/842/883/894/932,
 29/62/437/527, 29/78/87/150/527/727, 29/78/135/727/830, 29/87, 29/135/150/527/670/727/883,
 29/150/267/727/750/871/883/932, 29/150/437/727, 29/522/670/711/871, 29/670/932, 39/50,
 39/727/750/932, 50/135/150/932, 50/437/522/527, 50/711, 50/727/750/883/894, 62, 62/87/150,
 62/87/150/727, 62/135/522/711/727/750/842/871/894, 62/437, 62/437/727,
 78/87/486/527/670/727/750/830/842/871/913/932, 87/750, 89/109/527/678/727/842,
 89/109/678/727/736/812/878, 89/109/727/932, 89/109/932, 89/527, 89/527/678/692/736/842/878/932,
 89/527/678/932, 89/527/727/812/860, 89/678/692/736/932, 89/678/812/878, 89/842/878,
 109/527/678/812, 109/678/692/842/860/878/932, 109/678/727/860, 109/678/736/812/878,
 109/678/812, 109/678/842/878, 109/692/727/736/812, 109/692/727/812/842/860, 109/727/860/878,
 109/736/932, 109/812, 109/842, 109/932, 135/670/727, 135/711/750/932, 150/527/842/871/913,
 150/871/932, 150/883/932, 267/527/727, 403/527/678/692/736/812/842/860, 437/522/527/670/871,
 437/750/830/932, 522, 522/527/569/727, 522/830, 527, 527/678/692/727/736/878,
 527/678/692/812/932, 527/692/727/736/812, 527/692/727/736/842/860/878, 527/727/736,
 527/736/932, 527/812, 670/711/871, 670/830/871, 678/692/727/812/842, 678/692/812, 678/812,
 678/860/878, 678/913, 678/932, 692/727/736/842/913, 692/812, 727, 727/932, 871, and 878/932,
 wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ
 ID NO: 2880.

23. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid
 alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%,
 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 2880,
 or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at
 least one substitution or substitution set at one or more positions selected from 62/87/150,
 89/109/527/678/727/842, 89/109/678/727/736/812/878, 89/109/932,
 89/527/678/692/736/842/878/932, 89/527/727/812/860, 89/678/692/736/932, 89/678/812/878,
 109/527/678/812, 109/678/692/842/860/878/932, 109/678/736/812/878, 109/678/812,
 109/692/727/736/812, 109/692/727/812/842/860, 109/736/932, 109/812, 109/842, 109/932,
 403/527/678/692/736/812/842/860, 522/830, 527/678/692/727/736/878, 527/678/692/812/932,
 527/692/727/736/812, 527/692/727/736/842/860/878, 527/727/736, 527/736/932, 527/812,
 678/692/727/812/842, 678/692/812, 678/812, 692/727/736/842/913, and 692/812, wherein the amino
 acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 2880.

24. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid
 alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%,

91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 3104, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 62, 62/89/830, 62/248/678/830/878/932, 62/678, 62/678/785, 62/678/830, 62/678/830/860, 62/678/830/860/871/878/932, 62/678/830/860/878, 62/678/830/860/878/932, 62/678/830/860/932, 62/678/830/871, 62/678/830/871/932, 62/678/830/878/932, 62/678/830/932, 62/678/860, 62/678/860/878, 62/678/860/932, 62/678/871, 62/678/871/932, 62/678/878/932, 62/678/932, 62/830, 62/830/860, 62/830/860/871/873, 62/830/860/878/932, 62/830/860/932, 62/830/871/932, 62/830/878, 62/830/932, 62/833/860/932, 62/860, 62/860/871, 62/860/871/878, 62/860/871/932, 62/860/878/932, 62/860/932, 62/871/878/932, 62/871/932, 62/878, 62/878/932, 62/932, 678, 678/830/932, 678/932, and 860/932, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 3104.

25. The recombinant acid alpha glucosidase of Claim 1, wherein said recombinant acid alpha glucosidase comprises a polypeptide sequence having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO: 3104, or a functional fragment thereof, and wherein said recombinant acid alpha glucosidase comprises at least one substitution or substitution set at one or more positions selected from 62/89/830, 62/830, 62/830/860, 62/830/860/932, and 62/830/932, wherein the amino acid positions of said polypeptide sequence are numbered with reference to SEQ ID NO: 3104.

26. The recombinant acid alpha glucosidase of Claim 1, wherein said acid alpha glucosidase comprises at least one mutation in at least one position as provided in Tables 3-1, 3-2, 4-1, 6-1, 10-1, 10-2, 12-1, 13-1, 13-2, 14-1, 14-2, 15-1, 16-1, 17-1, 17-2, 17-3, 17-4, 17-5, 17-6, 17-7, 17-8, and/or 17-9.

27. The recombinant acid alpha glucosidase of any of Claims 1-26, wherein said recombinant acid alpha glucosidase is derived from a human acid alpha glucosidase .

28. A recombinant acid alpha glucosidase comprising the polypeptide sequence of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

29. The recombinant acid alpha glucosidase of any of Claims 1-28, wherein said recombinant acid alpha glucosidase is more thermostable than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

30. The recombinant acid alpha glucosidase of any of Claims 1-29, wherein said recombinant acid alpha glucosidase is more stable at pH 7 than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

31. The recombinant acid alpha glucosidase of any of Claims 1-30, wherein said recombinant acid alpha glucosidase is more stable at pH 4 than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

32. The recombinant acid alpha glucosidase of any of Claims 1-31, wherein said recombinant acid alpha glucosidase exhibits increased expression than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

33. The recombinant acid alpha glucosidase of any of Claims 1-32, wherein said recombinant acid alpha glucosidase is more lysosomally stable than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

34. The recombinant acid alpha glucosidase of any of Claims 1-33, wherein said recombinant acid alpha glucosidase is more readily taken up by cells than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

35. The recombinant acid alpha glucosidase of any of Claims 1-34, wherein said recombinant acid alpha glucosidase exhibits greater enzymatic activity in cell lysates than the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

36. The recombinant acid alpha glucosidase of any of Claims 1-35, wherein said recombinant acid alpha glucosidase exhibits reduced immunogenicity, as compared to the acid alpha glucosidase of SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.

37. The recombinant acid alpha glucosidase of any of Claims 1-36, wherein said recombinant acid alpha glucosidase is purified.

38. The recombinant acid alpha glucosidase of any of Claims 1-37, wherein said recombinant acid alpha glucosidase exhibits at least one improved property selected from: i) enhanced catalytic activity; ii) increased tolerance to pH 7; iii) increased tolerance to pH 4; iv) increased expression; v) increased uptake into cells; vi) increased enzymatic activity in cell lysates; vii) reduced immunogenicity; or a combination of any of i), ii), iii), iv), v), vi), and/or vii), as compared to a reference sequence.

39. The recombinant acid alpha glucosidase of Claim 38, wherein said reference sequence is selected from SEQ ID NO: 2, 6, 8, 12, 14, 16, 18, 20, 946, 1956, 2496, 2880, and/or 3104.
40. A composition comprising at least one recombinant acid alpha glucosidase of any of Claims 1-39.
41. A recombinant polynucleotide sequence encoding at least one recombinant acid alpha glucosidase set forth in any of Claims 1-40.
42. The recombinant polynucleotide sequence of Claim 41, wherein said polynucleotide sequence is selected from DNA, RNA, and mRNA.
43. The recombinant polynucleotide sequence of Claim 41 and/or 42, wherein said polynucleotide sequence is codon-optimized.
44. An expression vector comprising the recombinant polynucleotide sequence of Claim 41, 42, and/or 43.
45. The expression vector of Claim 44, wherein said recombinant polynucleotide sequence is operably linked to a control sequence.
46. The expression vector of Claim 45, wherein said control sequence is a promoter.
47. The expression vector of Claim 46, wherein said promoter is a heterologous promoter.
48. The expression vector pDH.
49. The expression vector pDH of Claim 48, wherein said expression vector comprises the nucleotide sequence of SEQ ID NO: 3379 or SEQ ID NO: 3380.
50. The expression vector of Claim 48, wherein said expression vector comprises a polynucleotide sequence of any of the odd-numbered sequences of SEQ ID NO: 7-3377.
51. A host cell comprising the expression vector of any of Claims 44-50.

52. The host cell of Claim 51, wherein said host cell is selected from eukaryotes and prokaryotes.
53. The host cell of Claim 51 and/or 52, wherein said host cell is a mammalian cell.
54. A method of producing a recombinant acid alpha glucosidase variant, comprising culturing said host cell of any of Claims 52-53, under conditions that said acid alpha glucosidase encoded by said recombinant polynucleotide is produced.
55. The method of Claim 54, further comprising the step of recovering said acid alpha glucosidase.
56. The method of Claim 55, further comprising the step of purifying said acid alpha glucosidase.
57. A recombinant acid alpha glucosidase variant produced according to the method of any of Claims 54-56.
58. A composition comprising the recombinant acid alpha glucosidase of Claim 57.
59. A composition comprising at least one polynucleotide of any of Claims 41-43.
60. A pharmaceutical composition for the treatment of Pompe disease, comprising the composition of Claim 40, 58, and/or 59.
61. A pharmaceutical composition for the treatment of Pompe disease, comprising the composition of Claim 40 and/or 58.
62. A pharmaceutical composition comprising the recombinant polynucleotide of any of Claims 41-43.
63. The pharmaceutical composition of any of Claims 60-62, further comprising a pharmaceutically acceptable carrier and/or excipient.
64. The pharmaceutical composition of any of Claims 60-63, wherein said composition is suitable for parenteral injection or infusion to a human.

65. A method for treating and/or preventing the symptoms of Pompe disease in a subject, comprising providing a subject having Pompe disease, and providing the pharmaceutical composition and/or composition of any of Claims 40, 58, 59, 60, 61, 62, 63, and/or 64, to said subject.

66. The method of Claim 65, wherein said symptoms of Pompe disease are ameliorated.

67. The method of any of Claims 65-66, wherein said subject is an infant or child.

68. The method of any of Claims 65-67, wherein said subject is an adult or young adult.

69. Use of the compositions provided in any of Claims 40, and 58-64.