PHARMACEUTICAL COMPOSITIONS FOR THE TREATMENT OF ALCOHOL ADDICTION

The present invention relates to pharmaceutical compositions for the treatment of alcohol addiction, characterized in that they contain - as the active principle - a purified lipophilic extract of Salvia miltiorrhyza Bunge. The invention also relates to a process for the preparation of the purified lipophilic extract of Salvia miltiorrhyza Bunge, and the extract thus obtained. Furthermore, the invention relates to the use of the pure active principles contained in this extract (Tanshinone IIA and Miltirone) in the treatment of alcohol addiction and to pharmaceutical compositions for said treatment which contain as the active principles Tanshinone IIA and/or Miltirone.

Alcohol abuse and alcohol addiction, phenomena which can be collectively referred to by the term alcoholism, represent a serious problem for the whole of modern society (Gessa G.L., Bisogno compulsivo di bere e "principio del piacere" [The compulsive need to drink and the pleasure principle] in Medicina delle tossicodipendenze [Drug Addiction Medicine] II, 5 (1994)). In Italy, for example, more than 9% of the population (about 5 million people) are heavy drinkers and more than 1 million people are alcohol-addict (Calamo-Specchia F.P. - Epidemiologia dell'alcolismo in Italia [Epidemiology of alcoholism in Italy] in Atti del VII Congresso Nazionale della S.I.A., [Records of the 7th National Congress of the S.I.A.] Mediserve, Rome, 295-301, (1991)). These figures become much higher if we

CONFIRMATION COPY

consider countries such as the United States of America where there are more than 13 million alcohol-addicts. Alcohol abuse and actual alcohol addiction result in an enormous outlay of public money (recently, since 1991, in the United States about 200 thousand million dollars a year have been spent) and are the cause of enormous social and psychological damage for the individuals involved.

The existing approaches for the treatment of alcoholism, in addition to those of a psychological nature (group therapy, etc.), consist in the use of drugs such as disulfiram and calcium carbamide which act on the metabolism of alcohol, inhibiting hepatic aldehyde-dehydrogenase and therefore raising the hematic levels of acetaldehyde, with all the undesiderable phenomena which occur each time ethanol is taken.

According to the present state of the art, the sole plant whose derivatives have been used for the treatment of alcoholism is the Pueraria lobata (Radix puerarie), which is widely used in traditional Chinese medicine and forms the subject of Patent Application WO 93/00896.

It has now been surprisingly found that the purified lipophilic extract of the roots of Salvia miltiorrhyza Bunge can be used with success for reducing the voluntary consumption of alcohol and other drugs which induce addiction. Furthermore, individual pure products isolated from the extract, such as Tanshinone IIA and Miltirone, have also proved to be useful. The dried roots of Salvia miltiorrhyza Bunge (Danshen, Radix Salviae miltiorrhyzae) are officially listed in the Chinese Pharmacopeia; they are widely used for the

treatment of menstrual disturbances, cardiovascular diseases such as, for example, angina pectoris and abnormalities of platelet function, as well as insomnia (Tang, W. Eisenbrand, G., Salvia miltiorrhyza Bge., in Chinese drugs of plant origin; Chemistry, Pharmacology, and Use in Traditional Chinese Medicine, pp. 891-902 - Springer-Verlag, Berlin (1992); Kee Chang Huang, Antianginal Herbs in The Pharmacology of Chinese Herbs, pp.81-84 - CRC Press, Boca Raton (1993)).

The use of hydro-alcoholic extracts of Salvia miltiorrhyza for the topical treatment of excess fat deposits is also known (Italian Patent 1,239,281).

The extract according to the invention, unlike that which is known, is of the lipophilic type and is obtained by extraction of the roots of Salvia miltiorrhyza with acetone, at temperatures of between about 20 and 50°C, using a drug/solvent ratio (w/v) of between 1:2 and 1:10, preferably about 1:5. Following concentration down to about 1/20th to 1/50th of the initial volume, water and a solvent which is immiscible in water are added. The acetone is removed by means of repeated washings with water while the organic phase containing the active principles is concentrated, following dilution, where necessary, with 95° ethanol.

The lipophilic extract according to the present invention is characterized by having a content of from 5 to 30% w/w, preferably from 15 to 25%, of Tanshinone IIA and of from 0.5 to 3% w/w, preferably of from 0.8 to 2%, of Miltirone.

Determination of the inhibiting action on alcohol consumption was performed using alcohol-consuming rats

of the strain called "Sardinian alcohol-preferring" (Sp) (Fadda F., Mosca E., Colombo G., Gessa G.L., Alcohol-preferring rats: Genetic sensitivity to alcohol-induced stimulation of dopamine metabolism, in Physiol. Behav. 47, 727 (1990)).

These animals, which given a free choice between alcohol and water consume daily 6 to 7 g of alcohol per kg of body weight (with a water-to-alcohol ratio higher than 2:1), during the last few years have been used with success to determine the effect of various substances on the voluntary consumption of alcohol; see, for example, Balakleevsky A., Colombo G., Fadda F., Gessa G.L., Ro 19-4603, a benzodiazepine receptor inverse agonist, attenuates voluntary ethanol consumption in rats selectively bred for high ethanol preference, in Alcohol Alcohol. 25, 449-452 (1990); Fadda F., Garau B., Colombo G., Gessa G.L., Isradipine and other calcium channel antagonists attenuate ethanol consumption in ethanol-preferring rats, in Alcoholism: Clinical and Experimental Research 16(3), 449-452 (1992).

The animals, which were kept under normal housing conditions, were given a free choice between water (which was always present) and alcohol (a 10% solution v/v) which was offered for a period of 4 hours a day (i.e. the first 4 hours of darkness during the day/night cycle). The amounts of water and alcohol consumed were recorded every day at the same time. Food was offered ad libitum. Once a stable consumption of alcohol and water was reached, different doses of purified lipophilic extract of Salvia miltiorrhyza (prepared as above) dissolved in dimethylsulfoxide, were administered

orally in a volume amount of 2 ml/kg once a day for 5 consecutive days. An equal volume of the carrier was used as the control. At the end of treatment, the alcohol and water consumption was recorded until the values recorded prior to the treatment were re-established. Statistical significant difference from the mean of the values obtained in the group treated with the carrier alone (* $p<0.05$, ** $p<0.01$) was evaluated by the Dunnet test for multiple comparisons.

Figure 1 shows the effect of repeated oral administration of increasing doses of the extract on alcohol consumption; Figure 2 shows the effect of increasing doses on water consumption. Figure 3, on the other hand, shows the effect on alcohol consumption resulting from repeated doses both of the extract and of the principles contained therein (Tanshinone IIA and Miltirone); Figure 4, finally, illustrates the effect on water consumption of the extract and the said principles when administered in repeated doses.

From an examination of Figures 1 and 2 it can be concluded that the extract reduces alcohol consumption significantly and in a dose-related manner, this result being achieved in some cases (i.e. with the highest dose) already after the first day of administration. The reduction in alcohol consumption remains constant until the fifth day and then regresses following suspension of the treatment. Moreover, it is surprising to note that this trend is accompanied by the tendency for a gradual increase in the water consumption, as if the animal were substituting it for the alcohol. This latter observation is of particular importance since it

shows that the treatment carried out with the product being tested is perfectly well-tolerated and the animal returns without any difficulty to a more physiological life cycle, using water instead of alcohol.

The administration in constant doses, repeated for 5 days, of both the extract and Tanshinone IIA and Miltirone (Figures 3 and 4) confirms the results indicated above, but demonstrates that the pure components - taking account of the doses - are the main factors responsible for the effect of the extract.

The invention therefore provides pharmaceutical compositions which can be administered orally and which contain as the active principle either the lipophilic extract of Salvia miltiorrhyza or the purified active principles Tanshinone IIA and/or Miltirone, the latter being obtainable from commercial sources or by means of conventional purification of the lipophilic extract. The compositions of the invention, in addition to conventional excipients or carriers, will contain from about 10 to about 500 mg of the extract or the equivalent doses of Tanshinone IIA and Miltirone, taking into account their extract content.

EXAMPLE 1

Preparation of the lipophilic extract

1 kg of finely ground roots of Salvia miltiorrhyza are extracted 4 times using 5 l of acetone at 50°C. The recombined extracts are concentrated under a vacuum to 500 ml; the concentrate is diluted with 1 l of methylene chloride and 500 ml of H_2O. The phases are separated and the organic phase containing the active principles is washed with water until the acetone and the unwanted

RECTIFIED SHEET (RULE 91)
ISA/EP

polar substances are eliminated.

The chloromethylene phase is concentrated to a small volume; the residue is diluted with 300 ml of 95° ethanol. The solution is concentrated under a vacuum to dryness at a temperature of 60°C. After drying overnight, 11.3 g of extract are obtained, containing 15.2% of Tanshinone IIA and 1.1% of Miltirone.

EXAMPLE 2

Formulation in capsule form

Each 130 mg capsule contains:

Extract of Example 1	25 mg
Pregelatinized starch	25 mg
Microcrystalline cellulose	49 mg
Lactose	15 mg
Colloidal silica	6 mg
Cross-linked sodium carboxymethylcellulose	6.5 mg
Polyvinylpyrrolidone	2.5 mg
Magnesium steareate	1 mg

EXAMPLE 3

Formulation in tablet form

Each 400 mg tablet contains:

Extract of Example 1	100 mg
Pregelatinized starch	100 mg
Microcrystalline cellulose	96 mg
Lactose	45 mg
Colloidal silica	25 mg
Cross-linked sodium carboxymethylcellulose	20 mg
Polyvinylpyrrolidone	10 mg
Magnesium steareate	4 mg

RECTIFIED SHEET (RULE 91)
ISA/EP

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

CLAIMS

1. Lipophilic extract of the roots of Salvia miltiorrhyza Bunge characterized in that it contains from 5 to 30% w/w of Tanshinone IIA and from 0.5 to 3% w/w of Miltirone.

2. An extract according to claim 1, containing from 15 to 25% w/w of Tanshinone IIA.

3. An extract according to claim 1 or 2, containing from 0.8 to 2% of Miltirone.

4. A process for the preparation of the extract of claims 1-3, comprising the extraction of the roots of Salvia miltiorrhyza with acetone, subsequent concentration, treatment of the concentrate with water and a water-immiscible solvent, removal of the acetone by washing with water and concentration of the organic phase.

5. Lipophilic extract of Salvia miltiorrhyza Bunge obtainable by extraction of the roots of Salvia miltiorrhyza with acetone, subsequent concentration, treatment of the concentrate with water and a water-immiscible solvent, removal of the acetone by washing with water and concentration of the organic phase.

6. Oral pharmaceutical compositions containing the extract according to Claims 1-3 or 5 in admixture with suitable carriers.

7. Composition according to Claim 6 in the form of capsules or tablets.

8. Use of the extract according to Claims 1-3 or 5 for the preparation of medicaments for the treatment of alcohol addiction.

* C L M S *

THIS SHEET HAS BEEN REPLACED

9. Use of Tanshinone IIA for the preparation of medicaments for the treatment of alcohol addiction.

10. Use of Miltirone for the preparation of medicaments for the treatment of alcohol addiction.

11. Oral pharmaceutical compositions containing as the active principle Tanshinone IIA and/or Miltirone in admixture with suitable carriers.

This Sheet Has Been Replaced





* C L M S *

FIGURE 1


Controls
25 mg/kg
50 mg/kg
100 mg/kg
Alcohol consumption (g/kg)
3.0
2.5
2.0
1.5
1.0
0.5
0.0
1
2
3
4
5
6
7
8
9
10
days
*
**
***


Controls
25 mg/kg
50 mg/kg
100 mg/kg
Water consumption (ml/kg)
25,0
22,5
20,0
17,5
15,0
12,5
10,0
7,5
5,0
2,5
0,0
1
2
3
4
5
6
7
8
9
10
days

FIGURE 2

FIGURE 3


Carrier
Salvia miltiorrhyza
Tanshinone IIA
Miltirone
Alcohol consumption (g/kg)
3.0
2.5
2.0
1.5
1.0
0.5
0.0
1
2
3
4
5
6
7
8
9
10 days

FIGURE 4


Carrier
Salvia miltiorrhyza
Tanshinone IIA
Miltirone
days
Water consumption (ml/g)
10.0
9.0
8.0
7.0
6.0
5.0
4.0
3.0
2.0
1.0
0.0
1
2
3
4
5
6
7
8
9
10

9705042-1
IPOS

PCT WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: A61K 35/78, 31/34, 31/12 — A1

(11) International Publication Number: WO 96/35441

(43) International Publication Date: 14 November 1996 (14.11.96)

(21) International Application Number: PCT/EP96/01916

(22) International Filing Date: 8 May 1996 (08.05.96)

(30) Priority Data:
MI95A000958 12 May 1995 (12.05.95) IT

(71) Applicant *(for all designated States except US)*: INDENA S.P.A. [IT/IT]; Viale Orties, 12, I-20139 Milano (IT).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: MORAZZONI, Paolo [IT/IT]; Viale Orties, 12, I-20139 Milano (IT). BOMBARDELLI, Ezio [IT/IT]; Viale Orties, 12, I-20139 Milano (IT).

(74) Agent: BIANCHETTI, Giuseppe; Studio Consulenza Brevettuale, Via Rossini, 8, I-20122 Milano (IT).

(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published
With international search report.
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: PHARMACEUTICAL COMPOSITIONS FOR THE TREATMENT OF ALCOHOL ADDICTION



(57) Abstract

The present invention relates to pharmaceutical compositions for the treatment of alcohol addiction, characterized in that they contain - as the active principle - a purified lipophilic extract of Salvia miltiorrhyza Bunge.

CLAIMS

1. Use of lypophilic extracts of the roots of Salvia miltiorrhyza Bunge containing from 5 to 30% w/w of Tanshinone IIA and from 0,5 to 3% w/w of Miltirone, for the preparation of medicaments for the treatment of alcohol addition.

2. Use of Tanshinone IIA for the preparation of medicaments for the treatment of alcohol addiction.

3. Use of Miltirone for the preparation of medicaments for the treatment of alcohol addiction.

* C L M S *

AMENDED SHEET