

1/17




Figure 1

2/17




Figure 2

3/17




Figure 3




Figure 4

5/17

COUNTS




Figure 5

6/17




Figure 6

7/17




Figure 7

8/17




Figure 8

9/17




Figure 9

10/17





Figure 10

11/17

EFFECTS OF Cys481Ser ON THE ACTIVITY OF BTK INHIBITORS^a

IC ₅₀ (nM), [ATP] = 50 μ M	WT BTK	Cys481Ser BTK	FOLD CHANGE
COMPOUND 2	4.6	1.1	0.2
IBRUTINIB	0.1	6.6	66
ACALABRUTINIB	4.2	707	168

^aDIRECT MEASUREMENT OF KINASE ACTIVITY OF HUMAN RECOMBINANT WT BTK OR Cys481Ser MUTANT BTK WAS PERFORMED IN A FRET KINASE ASSAY

Figure 11

12/17





Figure 12


13/17

PERCENT BTK INHIBITION VS. TIME^a

%BTK INHIBITION (MEAN, SD) DETERMINED BY ASSESSMENT OF pBTK AS DESCRIBED
IN METHODS BTK, BRUTON'S TYROSINE KINASE; pBTK, PHOSPHORYLATED BTK.

Figure 13

14/17

PERCENT BTK INHIBITION VS. COMPOUND 1 PLASMA CONCENTRATION^a

%BTK INHIBITION DETERMINED BY ASSESSMENT OF pBTK AS DESCRIBED
IN METHODS; COMPOUND 1 FREE BASE CONCENTRATION DETERMINED BY LC-MS/MS
BTK, BRUTON'S TYROSINE KINASE; pBTK, PHOSPHORYLATED BTK.

Figure 14

15/17




Figure 15

16/17




Figure 16

17/17

COMPOUND	WT C481 BTK IC ₅₀ nM	C481S BTK IC ₅₀ nM	FOLD CHANGE ^a
COMPOUND 1	570	800	1.4
IBRUTINIB	16	1700	>100

^aEXPRESSED AS IC₅₀ OF COMPOUND AGAINST C481S-BTK DIVIDED BY IC₅₀
OF COMPOUND AGAINST WT BTK.

Figure 17