

DESCRIPTION

NITROGEN-CONTAINING HETEROCYCLIC COMPOUND

Technical Field

[0001]

5 The present invention relates to a nitrogen-containing heterocyclic compound that is useful as a chemokine CXCL10 inhibitor.

Background Art

[0002]

10 CXCL10 (C-X-C motif chemokine 10) is a CXC chemokine also known as IP-10 (interferon gamma inducible protein 10). Upon partial damage of living tissues by infection or the like, chemotactic factors such as CXCL10 and MCP-1 (monocyte chemotactic protein-1) are produced. As a result, monocytes, lymphocytes and neutrophils are infiltrated into inflammatory tissues.

15 It has been suggested that immune diseases such as inflammatory bowel disease, arthritis, psoriasis, systemic sclerosis, systemic lupus erythematosus and autoimmune neuroinflammatory disease develop due to the increased production of CXCL10. Therefore, a compound having an excellent CXCL10 inhibitory activity is considered to be useful as an agent for treatment such as prophylaxis and/or therapy of a disease involving the overproduction of
20 CXCL10 (Patent Literature 1).

Prior Art Documents

Patent Document

[0003]

25 Patent Document 1: International Publication No. WO 2009/041026

Summary of Invention

Technical Problem

[0004]

30 There has been a demand for a compound that has an excellent CXCL10 inhibitory activity and is useful as an agent for treatment such as prophylaxis and/or therapy of a disease involving the overproduction of CXCL10.

An object of the present invention is to provide a compound that has an excellent CXCL10 inhibitory activity and is useful as an agent for treatment such as prophylaxis and/or therapy of a disease involving the overproduction of CXCL10.

5 Solution to Problem

[0005]


Under these circumstances, the present inventors have conducted diligent studies and consequently completed the present invention by finding that a nitrogen-containing heterocyclic compound shown below has an excellent CXCL10 inhibitory activity and is useful
10 as an agent for treatment such as prophylaxis and/or therapy of a disease involving the overproduction of CXCL10.

[0006]

Thus, the present invention provides the following:

<1>

15 A compound represented by the general formula [1] or a salt thereof:



[1]

wherein

R¹ represents an optionally substituted C₁₋₆ alkyl group;

R² represents a hydrogen atom, a halogen atom or an optionally substituted C₁₋₆ alkyl group;

20 R³ represents a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₃₋₈ cycloalkyl group, an optionally substituted C₄₋₈ cycloalkenyl group, an optionally substituted aryl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted C₁₋₆ alkylamino group, an optionally substituted di(C₁₋₆ alkyl)amino group or an optionally substituted heterocyclic group;

25 Z¹, Z² and Z³ are the same or different and each represent a nitrogen atom or a group represented by the general formula CR⁵ wherein R⁵ represents a hydrogen atom, a halogen atom or an optionally substituted C₁₋₆ alkyl group;

X¹ represents

(1) a group represented by the general formula C(=O)N(R⁶) wherein the carbon atom is bonded

to ring A, and R^6 represents a hydrogen atom, an amino-protective group or an optionally substituted C_{1-6} alkyl group,

(2) a group represented by the general formula $N(R^7)C(=O)$ wherein the nitrogen atom is bonded to ring A, and R^7 represents a hydrogen atom, an amino-protective group or an optionally

5 substituted C_{1-6} alkyl group; or R^7 represents, together with one substituent R^4 on ring A, an optionally substituted C_{2-4} alkylene group, a group represented by the general formula $O-Y^1$ wherein the oxygen atom is bonded to ring A, and Y^1 represents an optionally substituted C_{1-3} alkylene group, a group represented by the general formula $S(O)_n-Y^2$ wherein the sulfur atom is bonded to ring A, Y^2 represents an optionally substituted C_{1-3} alkylene group, and n represents an
10 integer of 0 to 2, or a group represented by the general formula $N(R^8)-Y^3$ wherein the nitrogen atom is bonded to ring A, Y^3 represents an optionally substituted C_{1-3} alkylene group, and R^8 represents a hydrogen atom, an amino-protective group, an optionally substituted C_{1-6} alkyl group or an optionally substituted aryl group,

(3) an optionally substituted divalent cyclic hydrocarbon group resulting from the removal of one
15 hydrogen atom from each of two adjacent atoms or

(4) an optionally substituted divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms;

ring A represents a cyclic hydrocarbon group or a heterocyclic group;

m number of R^4 are the same or different and each represent a halogen atom, a cyano group, a

20 nitro group, an amino-protective group, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{3-8} cycloalkyl group, an optionally substituted C_{4-8} cycloalkenyl group, an optionally substituted aryl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted aryloxy group, an optionally substituted C_{1-6} alkylamino group, an optionally substituted di(C_{1-6}
25 alkyl)amino group, an optionally substituted arylamino group, an optionally substituted carbamoyl group, an optionally substituted sulfamoyl group, an optionally substituted C_{1-6} alkylthio group, an optionally substituted arylthio group, an optionally substituted C_{1-6} alkylsulfonyl group, an optionally substituted arylsulfonyl group, an optionally substituted heterocyclic group, an optionally protected amino group, an optionally protected hydroxyl group,
30 an optionally protected carboxyl group,

an optionally substituted C_{2-5} alkylene group formed together by two adjacent R^4 ,

an optionally substituted C_{2-4} alkylene group formed together by one R^4 and R^7 ,

a group represented by the general formula $O-Y^1$ formed together by one R^4 and R^7 wherein the oxygen atom is bonded to ring A, and Y^1 represents an optionally substituted C_{1-3} alkylene

group,

a group represented by the general formula $S(O)_n-Y^2$ formed together by one R^4 and R^7 wherein the sulfur atom is bonded to ring A, Y^2 represents an optionally substituted C_{1-3} alkylene group, and n represents an integer of 0 to 2, or

- 5 a group represented by the general formula $N(R^8)-Y^3$ formed together by one R^4 and R^7 wherein the nitrogen atom is bonded to ring A, Y^3 represents an optionally substituted C_{1-3} alkylene group, and R^8 represents a hydrogen atom, an amino-protective group, an optionally substituted C_{1-6} alkyl group or an optionally substituted aryl group; and m represents an integer of 0 to 5.

10 [0007]

<2>

The compound according to <1> or a salt thereof, wherein

R^2 is a hydrogen atom or an optionally substituted C_{1-6} alkyl group; and each of Z^1 , Z^2 and Z^3 is CH.

15 <3>

The compound according to <1> or <2> or a salt thereof, wherein

R^1 is a C_{1-3} alkyl group, and

R^3 is an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{3-8} cycloalkyl group, an optionally substituted aryl group or an optionally substituted heterocyclic group.

20 <4>

The compound according to any one of <1> to <3> or a salt thereof, wherein ring A is a cyclic hydrocarbon group.

[0008]

<5>

- 25 The compound according to any one of <1> to <4> or a salt thereof, wherein X^1 is a group represented by the general formula $C(=O)N(R^6)$ wherein the carbon atom is bonded to ring A, and R^6 represents a hydrogen atom, an amino-protective group or an optionally substituted C_{1-6} alkyl group.

<6>

- 30 The compound according to any one of <1> to <4> or a salt thereof, wherein X^1 is a group represented by the general formula $C(=O)N(R^{6a})$ wherein the carbon atom is bonded to ring A, and R^{6a} represents a hydrogen atom or an optionally substituted C_{1-3} alkyl group;

m number of R^4 are the same or different and each are a halogen atom, an optionally substituted

C₁₋₃ alkyl group, an optionally substituted C₁₋₃ alkylamino group or an optionally protected amino group; and

m is an integer of 0 to 2.

[0009]

5 <7>

The compound according to any one of <1> to <4> or a salt thereof, wherein

R³ is an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₃₋₈ cycloalkyl group or an optionally substituted heterocyclic group;

X¹ is a group represented by the general formula C(=O)N(R^{6b}) wherein the carbon atom is

10 bonded to ring A, and R^{6b} represents an optionally substituted C₁₋₃ alkyl group;

m number of R⁴ are the same or different and each are a halogen atom or an optionally substituted C₁₋₃ alkyl group; and

m is an integer of 0 to 2.

<8>

15 The compound according to <7> or a salt thereof, wherein

m number of R⁴ are the same or different and each are a halogen atom; and

m is an integer of 0 to 2.

[0010]

<9>

20 The compound according to any one of <1> to <4> or a salt thereof, wherein

X¹ is a group represented by the general formula N(R⁷)C(=O) wherein the nitrogen atom is

bonded to ring A, and R⁷ represents a hydrogen atom, an amino-protective group or an optionally substituted C₁₋₆ alkyl group; or R⁷ represents, together with one substituent R⁴ on ring A, an optionally substituted C₂₋₄ alkylene group, a group represented by the general formula O-Y¹

25 wherein the oxygen atom is bonded to ring A, and Y¹ represents an optionally substituted C₁₋₃

alkylene group, a group represented by the general formula S(O)_n-Y² wherein the sulfur atom is bonded to ring A, Y² represents an optionally substituted C₁₋₃ alkylene group, and n represents an

integer of 0 to 2, or a group represented by the general formula N(R⁸)-Y³ wherein the nitrogen atom is bonded to ring A, Y³ represents an optionally substituted C₁₋₃ alkylene group, and R⁸

30 represents a hydrogen atom, an amino-protective group, an optionally substituted C₁₋₆ alkyl group or an optionally substituted aryl group.

<10>

The compound according to any one of <1> to <4> or a salt thereof, wherein

R³ is an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₃₋₈ cycloalkyl group or

an optionally substituted heterocyclic group;

X^1 is a group represented by the general formula $N(R^{7a})C(=O)$ wherein the nitrogen atom is bonded to ring A, and R^{7a} represents an optionally substituted C_{1-3} alkyl group, or R^{7a} represents, together with one substituent R^4 on ring A, an optionally substituted C_{2-3} alkylene group or a group represented by the general formula $O-Y^{1a}$ wherein the oxygen atom is bonded to ring A, and Y^{1a} represents an ethylene group;

m number of R^4 are the same or different and each are a halogen atom, an optionally substituted C_{1-3} alkyl group, an optionally substituted C_{1-3} alkylsulfonyl group,

an optionally substituted C_{2-3} alkylene group formed together by two adjacent R^4 ,

an optionally substituted C_{2-3} alkylene group formed together by one R^4 and R^{7a} , or a group represented by the general formula $O-Y^{1a}$ formed together by one R^4 and R^{7a} wherein the oxygen atom is bonded to ring A, and Y^{1a} represents an ethylene group; and m is an integer of 0 to 2.

[0011]

<11>

The compound according to any one of <1> to <4> or a salt thereof, wherein

R^3 is an optionally substituted heterocyclic group;

X^1 is a group represented by the general formula $N(R^{7b})C(=O)$ wherein the nitrogen atom is bonded to ring A, and R^{7b} represents an optionally substituted C_{1-3} alkyl group, or R^{7b} represents, together with one substituent R^4 on ring A, an optionally substituted C_{2-3} alkylene group;

m number of R^4 are the same or different and each are an optionally substituted C_{1-3} alkyl group, an optionally substituted C_{2-3} alkylene group formed together by two adjacent R^4 , or an optionally substituted C_{2-3} alkylene group formed together by one R^4 and R^{7b} ; and m is an integer of 0 to 2.

<12>

The compound according to <11> or a salt thereof, wherein

X^1 is a group represented by the general formula $N(R^{7b})C(=O)$ wherein the nitrogen atom is bonded to ring A, and R^{7b} represents an optionally substituted C_{1-3} alkyl group, or R^{7b} represents, together with one substituent R^4 on ring A, an optionally substituted C_{2-3} alkylene group;

m number of R^4 are the same or different and each are an optionally substituted C_{1-3} alkyl group, or

an optionally substituted C_{2-3} alkylene group formed together by one R^4 and R^{7b} ; and

m is an integer of 0 to 2.

[0012]

<13>

The compound according to any one of <1> to <4> or a salt thereof, wherein X^1 is an optionally substituted divalent cyclic hydrocarbon group resulting from the removal of one hydrogen atom from each of two adjacent atoms or an optionally substituted divalent

- 5 heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms.

<14>

The compound according to any one of <1> to <4> or a salt thereof, wherein R^3 is an optionally substituted C_{1-6} alkyl group or an optionally substituted heterocyclic group;
 10 X^1 is an optionally substituted divalent cyclic hydrocarbon group resulting from the removal of one hydrogen atom from each of two adjacent atoms or an optionally substituted divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms;

m number of R^4 are the same or different and each are a halogen atom; and

- 15 m is an integer of 0 to 2.

[0013]

<15>

The compound according to <14> or a salt thereof, wherein

R^3 is an optionally substituted heterocyclic group; and

- 20 X^1 is an optionally substituted divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms.

<16>

A pharmaceutical composition comprising a compound according to any one of <1> to <15> or a salt thereof.

- 25 <17>

An agent for treatment of an immune disease, comprising a compound according to any one of <1> to <15> or a salt thereof.

Advantageous Effects of Invention

- 30 [0014]

The compound of the present invention has an excellent CXCL10 inhibitory activity and is useful as an agent for treatment such as prophylaxis and/or therapy of a disease involving the overproduction of CXCL10.

Description of Embodiments

[0015]

The present invention will be described below in detail.

In the present invention, % means % by mass, unless otherwise specified.

5 In the present invention, each term has the following meaning, unless otherwise specified.

The halogen atom means a fluorine atom, a chlorine atom, a bromine atom or an iodine atom.

10 The C₁₋₆ alkyl group means a linear or branched C₁₋₆ alkyl group such as a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a pentyl group, an isopentyl group and a hexyl group.

The C₁₋₃ alkyl group means a methyl group, an ethyl group, a propyl group or an isopropyl group.

15 The C₂₋₆ alkenyl group means a linear or branched C₂₋₆ alkenyl group such as a vinyl group, an allyl group, a propenyl group, an isopropenyl group, a butenyl group, an isobutenyl group, a 1,3-butadienyl group, a pentenyl group and a hexenyl group.

The C₂₋₆ alkynyl group means a linear or branched C₂₋₆ alkynyl group such as an ethynyl group, a propynyl group, a butynyl group, a pentynyl group and a hexynyl group.

20 The C₃₋₈ cycloalkyl group means a C₃₋₈ cycloalkyl group such as a cyclopropyl group, a cyclobutyl group, a cyclopentyl group and a cyclohexyl group.

The C₄₋₈ cycloalkenyl group means a C₄₋₈ cycloalkenyl group such as a cyclobutenyl group, a cyclopentenyl group, a cyclohexenyl group and a cyclohexadienyl group.

The aryl group means a phenyl group or a naphthyl group.

25 The ar-C₁₋₆ alkyl group means an ar-C₁₋₆ alkyl group such as a benzyl group, a diphenylmethyl group, a trityl group, a phenethyl group and a naphthylmethyl group.

[0016]

The C₂₋₅ alkylene group means a linear or branched C₂₋₅ alkylene group such as an ethylidene group, an ethylene group, a trimethylene group, a propylene group, a tetramethylene group and a pentamethylene group.

30 The C₂₋₄ alkylene group means a linear or branched C₂₋₄ alkylene group such as an ethylidene group, an ethylene group, a trimethylene group, a propylene group and a tetramethylene group.

The C₂₋₃ alkylene group means an ethylidene group, an ethylene group, a trimethylene group or a propylene group.

The C₁₋₃ alkylene group means a methylene group, an ethylidene group, an ethylene group, a trimethylene group or a propylene group.

[0017]

The C₁₋₆ alkoxy group means a linear or branched C₁₋₆ alkyloxy group such as a methoxy group, an ethoxy group, a propoxy group, an isopropoxy group, a butoxy group, an isobutoxy group, a sec-butoxy group, a tert-butoxy group, a pentyloxy group and a hexyloxy group.

The aryloxy group means a phenoxy group or a naphthyloxy group.

The C₁₋₆ alkoxy-C₁₋₆ alkyl group means a C₁₋₆ alkyloxy-C₁₋₆ alkyl group such as a methoxymethyl group and a 1-ethoxyethyl group.

The ar-C₁₋₆ alkoxy-C₁₋₆ alkyl group means an ar-C₁₋₆ alkyloxy-C₁₋₆ alkyl group such as a benzyloxymethyl group and a phenethyloxymethyl group.

[0018]

The C₂₋₆ alkanoyl group means a linear or branched C₂₋₆ alkanoyl group such as an acetyl group, a propionyl group, a valeryl group, an isovaleryl group and a pivaloyl group.

The aroyl group means a benzoyl group or a naphthoyl group.

The heterocyclic carbonyl group means a furoyl group, a thenoyl group, a pyrrolidinylcarbonyl group, a piperidinylcarbonyl group, a piperazinylcarbonyl group, a morpholinylcarbonyl group or a pyridinylcarbonyl group.

The acyl group means a formyl group, a C₂₋₆ alkanoyl group, an aroyl group or a heterocyclic carbonyl group.

The C₁₋₆ alkoxy carbonyl group means a linear or branched C₁₋₆ alkyloxy carbonyl group such as a methoxycarbonyl group, an ethoxycarbonyl group, a propoxycarbonyl group, an isopropoxycarbonyl group, a tert-butoxycarbonyl group and a 1,1-dimethylpropoxycarbonyl group.

The aryloxy carbonyl group means a phenyloxy carbonyl group or a naphthyloxy carbonyl group.

The ar-C₁₋₆ alkoxy carbonyl group means an ar-C₁₋₆ alkyloxy carbonyl group such as a benzyloxy carbonyl group and a phenethyloxy carbonyl group.

[0019]

The C₁₋₆ alkylamino group means a linear or branched C₁₋₆ alkylamino group such as a methylamino group, an ethylamino group, a propylamino group, an isopropylamino group, a butylamino group, a sec-butylamino group, a tert-butylamino group, a pentylamino group and a hexylamino group.

The C₁₋₃ alkylamino group means a linear or branched C₁₋₃ alkylamino group such as a methylamino group, an ethylamino group, a propylamino group and an isopropylamino group.

5 The di(C₁₋₆ alkyl)amino group means a linear or branched di(C₁₋₆ alkyl)amino group such as a dimethylamino group, a diethylamino group, a dipropylamino group, a diisopropylamino group, a dibutylamino group, a di(tert-butyl)amino group, a dipentylamino group, a dihexylamino group, an (ethyl)(methyl)amino group and a (methyl)(propyl)amino group.

10 The arylamino group means a phenylamino group or a naphthylamino group.
[0020]

The C₁₋₆ alkylthio group means a C₁₋₆ alkylthio group such as a methylthio group, an ethylthio group and a propylthio group.

The arylthio group means a phenylthio group or a naphthylthio group.

15 The C₁₋₆ alkylsulfonyl group means a C₁₋₆ alkylsulfonyl group such as a methylsulfonyl group, an ethylsulfonyl group and a propylsulfonyl group.

The C₁₋₃ alkylsulfonyl group means a C₁₋₃ alkylsulfonyl group such as a methylsulfonyl group, an ethylsulfonyl group and a propylsulfonyl group.

The arylsulfonyl group means a benzenesulfonyl group, a p-toluenesulfonyl or a naphthalenesulfonyl group.

20 The C₁₋₆ alkylsulfonyloxy group means a C₁₋₆ alkylsulfonyloxy group such as a methylsulfonyloxy group, an ethylsulfonyloxy group and a propylsulfonyloxy group.

The arylsulfonyloxy group means a benzenesulfonyloxy group, a p-toluenesulfonyloxy group or a naphthalenesulfonyloxy group.

[0021]

25 The silyl group means a trimethylsilyl group, a triethylsilyl group or a tributylsilyl group.

[0022]

30 The cyclic amino group means, for example, a cyclic amino group which contains one or more nitrogen atoms as heteroatoms constituting the ring and may further contain one or more atoms selected from an oxygen atom and a sulfur atom, such as aziridinyl, azetidiny, pyrrolidinyl, piperidinyl, homopiperidinyl, pyrrolyl, dihydropyrrolyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, thiazolinyl, thiazolidinyl, dihydrothiadiazolyl, piperazinyl, homopiperazinyl, morpholinyl, homomorpholinyl, thiomorpholinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, decahydroisoquinolinyl,

benzomorpholinyl, dihydropyridoxazinyl and quinuclidinyl.

[0023]

The cyclic hydrocarbon group means a C₃₋₈ cycloalkyl group, a C₄₋₈ cycloalkenyl group or an aryl group.

5 [0024]

The monocyclic nitrogen-containing heterocyclic group means a monocyclic nitrogen-containing heterocyclic group which contains only a nitrogen atom as a heteroatom constituting the ring and is optionally substituted by an oxo group, such as an azetidiny group, a pyrrolidinyl group, an oxopyrrolidinyl group, a pyrrolinyl group, a pyrrolyl group, a piperidyl group, an oxopiperidyl group, a tetrahydropyridyl group, a dihydropyridyl group, a pyridyl group, a homopiperidinyl group, an octahydroazocinyl group, an imidazolidinyl group, an oxoimidazolidinyl group, an imidazoliny group, a dihydrooxoimidazolyl group, an imidazolyl group, a pyrazolidinyl group, a pyrazoliny group, a pyrazolyl group, a piperazinyl group, a homopiperazinyl group, a pyrazinyl group, a pyridazinyl group, a pyrimidinyl group, a triazolidinyl group, a dioxotriazolidinyl group, a triazolyl group and a tetrazolyl group.

The monocyclic oxygen-containing heterocyclic group means an oxetanyl group, a tetrahydrofuranyl group, an oxotetrahydrofuranyl group, a furanyl group, a tetrahydropyranyl group, an oxotetrahydropyranyl group, a dihydropyranyl group or a pyranyl group.

The monocyclic sulfur-containing heterocyclic group means a tetrahydrothienyl group, an oxotetrahydrothienyl group or a thienyl group.

The monocyclic nitrogen- and oxygen-containing heterocyclic group means a monocyclic nitrogen- and oxygen-containing heterocyclic group which contains only a nitrogen atom and an oxygen atom as heteroatoms constituting the ring and is optionally substituted by an oxo group, such as an oxazolyl group, an isoxazolyl group, an oxoisoxazolyl group, an oxadiazolyl group and a morpholinyl group.

The monocyclic nitrogen- and sulfur-containing heterocyclic group means a monocyclic nitrogen- and sulfur-containing heterocyclic group which contains only a nitrogen atom and a sulfur atom as heteroatoms constituting the ring and is optionally substituted by an oxo group, such as a thiazolyl group, an isothiazolyl group, a thiadiazolyl group, a thiomorpholinyl group, a 1-oxidethiomorpholinyl group and a 1,1-dioxidethiomorpholinyl group.

The monocyclic heterocyclic group means a monocyclic nitrogen-containing heterocyclic group, a monocyclic oxygen-containing heterocyclic group, a monocyclic sulfur-containing heterocyclic group, a monocyclic nitrogen- and oxygen-containing heterocyclic group

or a monocyclic nitrogen- and sulfur-containing heterocyclic group.

[0025]

The bicyclic nitrogen-containing heterocyclic group means a bicyclic nitrogen-containing heterocyclic group which contains only a nitrogen atom as a heteroatom constituting the rings and is optionally substituted by an oxo group, such as an indolinyl group, an oxoindolinyl group, an indolyl group, an isoindolinyl group, an oxoisoindolinyl group, an isoindolyl group, a pyrrolopyridinyl group, an indazolyl group, a benzimidazolyl group, a benzotriazolyl group, a tetrahydroquinolinyl group, an oxotetrahydroquinolinyl group, a dihydroquinolinyl group, an oxodihydroquinolinyl group, a quinolinyl group, a dihydroisoquinolinyl group, an octahydroisoquinolinyl group, a oxooctahydroisoquinolinyl group, a tetrahydroisoquinolinyl group, a decahydroisoquinolinyl group, an isoquinolinyl group, a dihydroquinazolinyl group, a cinnolinyl group, a phthalazinyl group, a quinazolinyl group, a tetrahydroquinoxalinyl group, an oxotetrahydroquinoxalinyl group, a hydroquinoxalinyl group, a quinoxalinyl group, a naphthyridinyl group, a purinyl group, a pteridinyl group and a quinuclidinyl group.

The bicyclic oxygen-containing heterocyclic group means a bicyclic oxygen-containing heterocyclic group which contains only an oxygen atom as a heteroatom constituting the rings and is optionally substituted by an oxo group, such as a 2,3-dihydrobenzofuranyl group, an oxo-2,3-dihydrobenzofuranyl group, a benzofuranyl group, an isobenzofuranyl group, a chromanyl group, an oxochromanyl group, a chromenyl group, an isochromanyl group, an oxoisochromanyl group, a 1,3-benzodioxolyl group, a 1,3-benzodioxanyl and a 1,4-benzodioxanyl group.

The bicyclic sulfur-containing heterocyclic group means a bicyclic sulfur-containing heterocyclic group which contains only a sulfur atom as a heteroatom constituting the rings and is optionally substituted by an oxo group, such as a 2,3-dihydrobenzothienyl group, an oxo-2,3-dihydrobenzothienyl group and a benzothienyl group.

The bicyclic nitrogen- and oxygen-containing heterocyclic group means a bicyclic nitrogen- and oxygen-containing heterocyclic group which contains only a nitrogen atom and an oxygen atom as heteroatoms constituting the rings and is optionally substituted by an oxo group, such as a dihydrobenzoxazolyl group, an oxodihydrobenzoxazolyl group, a benzoxazolyl group, a benzisoxazolyl group, a benzoxadiazolyl group, a benzomorpholinyl group, an oxobenzomorpholinyl group, a dihydropyranopyridyl group, an oxodihydropyranopyridyl group, a dihydrodioxinopyridyl group, an oxodihydrodioxinopyridyl group and a dihydropyridoxazinyl group.

The bicyclic nitrogen- and sulfur-containing heterocyclic group means a bicyclic nitrogen- and sulfur-containing heterocyclic group which contains a nitrogen atom and a sulfur atom as heteroatoms constituting the rings and is optionally substituted by an oxo group, such as a dihydrobenzothiazolyl group, an oxodihydrobenzothiazolyl group, a benzothiazolyl group, a
 5 benzisothiazolyl group and a benzothiadiazolyl group.

The bicyclic heterocyclic group means a bicyclic nitrogen-containing heterocyclic group, a bicyclic oxygen-containing heterocyclic group, a bicyclic sulfur-containing heterocyclic group, a bicyclic nitrogen- and oxygen-containing heterocyclic group or a bicyclic nitrogen- and sulfur-containing heterocyclic group.

10 [0026]

The heterocyclic group means a monocyclic heterocyclic group or a bicyclic heterocyclic group.

[0027]

The C₃₋₈ cycloalkane means cyclopropane, cyclobutane, cyclopentane,
 15 cyclohexane, cycloheptane or cyclooctane.

The C₄₋₈ cycloalkene means C₄₋₈ cycloalkene such as cyclobutene, cyclopentene, cyclopentadiene, cyclohexene, cyclohexadiene and cycloheptene.

The cyclic hydrocarbon means C₃₋₈ cycloalkane, C₄₋₈ cycloalkene, benzene or naphthalene.

20 [0028]

The nitrogen-containing heterocyclic ring means a nitrogen-containing heterocyclic ring which contains only a nitrogen atom as a heteroatom constituting the ring and is optionally substituted by an oxo group, such as azetidine, pyrrolidine, oxopyrrolidine, pyrroline, pyrrole, piperidine, oxopiperidine, tetrahydropyridine, dihydropyridine,
 25 oxodihydropyridine, pyridine, homopiperidine, octahydroazocine, imidazolidine, oxoimidazolidine, imidazoline, dihydrooxoimidazole, imidazole, pyrazolidine, dioxypyrazolidine, pyrazoline, oxopyrazoline, pyrazole, piperazine, homopiperazine, pyrazine, pyridazine, pyrimidine, triazolidine, dioxotriazolidine, triazole, tetrazole, 1H-benzimidazole and quinoxaline.

30 The oxygen-containing heterocyclic ring means oxetane, tetrahydrofuran, oxotetrahydrofuran, furan, tetrahydropyran, oxotetrahydropyran, dihydropyran or pyran.

The sulfur-containing heterocyclic ring means tetrahydrothiophene, oxotetrahydrothiophene or thiophene.

The nitrogen- and oxygen-containing heterocyclic ring means a monocyclic

nitrogen- and oxygen-containing heterocyclic ring which contains only a nitrogen atom and an oxygen atom as heteroatoms constituting the ring and is optionally substituted by an oxo group, such as oxazole, isoxazole, oxoisoxazole, oxadiazole and morpholine.

The nitrogen- and sulfur-containing heterocyclic ring means a monocyclic
 5 nitrogen- and sulfur-containing heterocyclic ring which contains only a nitrogen atom and a sulfur atom as heteroatoms constituting the ring and is optionally substituted by an oxo group, such as thiazole, isothiazole, thiadiazole, thiomorpholine, 1-oxidithiomorpholine and 1,1-dioxidithiomorpholine.

The heterocyclic ring means a nitrogen-containing heterocyclic ring, an oxygen-
 10 containing heterocyclic ring, a sulfur-containing heterocyclic ring, a nitrogen- and oxygen-containing heterocyclic ring or a nitrogen- and sulfur-containing heterocyclic ring.

[0029]

The divalent cyclic hydrocarbon group resulting from the removal of one
 hydrogen atom from each of two adjacent atoms means a group which is formed by the removal
 15 of two hydrogen atoms bonded to two adjacent atoms from cyclic hydrocarbon, such as cyclopropane-1,2-diyl, cyclobutane-1,2-diyl, cyclobutene-1,2-diyl, cyclopentane-1,2-diyl, cyclopentene-1,2-diyl, cyclopentadiene-1,2-diyl, cyclohexane-1,2-diyl, cyclohexene-1,2-diyl, cyclohexadiene-1,2-diyl, cycloheptane-1,2-diyl, cycloheptene-1,2-diyl, cyclooctane-1,2-diyl, benzene-1,2-diyl, naphthalene-1,2-diyl and naphthalene-2,3-diyl.

20 [0030]

The divalent heterocyclic group resulting from the removal of one hydrogen atom
 from each of two adjacent atoms means a group which is formed by the removal of two
 hydrogen atoms bonded to two adjacent atoms from a heterocyclic ring and optionally
 substituted by an oxo group, such as azetidine-1,2-diyl, pyrrolidine-1,2-diyl, oxopyrrolidine-1,2-
 25 diyl, oxopyrrolidine-3,4-diyl, pyrroline-3,4-diyl, pyrrole-3,4-diyl, piperidine-2,3-diyl, piperidine-3,4-diyl, oxopiperidine-2,3-diyl, tetrahydropyridine-1,2-diyl, tetrahydropyridine-2,3-diyl, tetrahydropyridine-3,4-diyl, dihydropyridine-2,3-diyl, dihydropyridine-3,4-diyl, dihydropyridine-1,2-diyl, oxodihydropyridine-1,2-diyl, pyridine-2,3-diyl, pyridine-3,4-diyl, homopiperidine-2,3-diyl, homopiperidine-3,4-diyl, octahydroazocine-2,3-diyl, imidazolidine-
 30 1,5-diyl, oxoimidazolidine-1,5-diyl, 1,2-dihydroimidazole-3,4-diyl, 4,5-dihydroimidazole-1,2-diyl, dihydroxoimidazole-1,5-diyl, 2H-imidazole-4,5-diyl, imidazole-1,2-diyl, imidazole-1,5-diyl, imidazole-4,5-diyl, pyrazolidine-1,2-diyl, dioxypyrazolidine-1,2-diyl, pyrazoline-1,2-diyl, oxopyrazoline-1,2-diyl, pyrazoline-1,5-diyl, 1H-pyrazole-3,4-diyl, 1H-pyrazole-4,5-diyl, 1H-pyrazole-1,5-diyl, piperazine-1,2-diyl, piperazine-2,3-diyl, homopiperazine-1,2-diyl,

homopiperazine-2,3-diyl, pyrazine-2,3-diyl, pyridazine-3,4-diyl, pyrimidine-4,5-diyl, 1,2,4-triazolidine-1,2-diyl, dioxotriazolidine-1,2-diyl, 1,2,3-triazole-1,5-diyl, 1,2,4-triazole-1,5-diyl, 1,2,4-triazole-3,4-diyl, tetrazole-1,5-diyl, 1H-benzimidazole-1,2-diyl, quinoxaline-2,3-diyl, oxetane-2,3-diyl, tetrahydrofuran-2,3-diyl, oxotetrahydrofuran-2,3-diyl, furan-2,3-diyl, 5 tetrahydropyran-2,3-diyl, oxotetrahydropyran-2,3-diyl, dihydropyran-2,3-diyl, pyran-2,3-diyl, tetrahydrothiophene-2,3-diyl, oxotetrahydrothiophene-2,3-diyl, thiophene-3,4-diyl, oxazole-4,5-diyl, isoxazole-3,4-diyl, dihydrooxoisoxazole-3,4-diyl, 1,2,3-oxadiazole-4,5-diyl, morpholine-2,3-diyl, morpholine-3,4-diyl, thiazole-4,5-diyl, isothiazole-4,5-diyl, 1,2,3-thiadiazole-4,5-diyl, thiomorpholine-2,3-diyl, 1-oxidithiomorpholine-2,3-diyl and 1,1-dioxidithiomorpholine-2,3-
10 diyl.
[0031]

The leaving group means a halogen atom, a C₁₋₆ alkylsulfonyloxy group or an arylsulfonyloxy group. The C₁₋₆ alkylsulfonyloxy group and the arylsulfonyloxy group are each optionally substituted by one or more groups selected from substituent group A.

15 [0032]

Substituent group A: a halogen atom, a carbamoyl group optionally substituted by one or more groups selected from substituent group B, a sulfamoyl group optionally substituted by one or more groups selected from substituent group B, an acyl group optionally substituted by one or more groups selected from substituent group B, a C₁₋₆ alkyl group optionally substituted
20 by one or more groups selected from substituent group B, a C₂₋₆ alkenyl group optionally substituted by one or more groups selected from substituent group B, a C₃₋₈ cycloalkyl group optionally substituted by one or more groups selected from substituent group B, a C₄₋₈ cycloalkenyl group optionally substituted by one or more groups selected from substituent group B, a C₁₋₆ alkoxy group optionally substituted by one or more groups selected from substituent
25 group B, a C₁₋₆ alkylsulfonyl group optionally substituted by one or more groups selected from substituent group B, an arylsulfonyl group optionally substituted by one or more groups selected from substituent group B, a C₁₋₆ alkylamino group optionally substituted by one or more groups selected from substituent group B, a di(C₁₋₆ alkyl)amino group optionally substituted by one or more groups selected from substituent group B, an aryl group optionally substituted by one or
30 more groups selected from substituent group B, a heterocyclic group optionally substituted by one or more groups selected from substituent group B, an arylamino group optionally substituted by one or more groups selected from substituent group B, a cyano group, an optionally protected amino group, an optionally protected hydroxyl group, an optionally protected carboxyl group and an oxo group.

[0033]

Substituent group B: a halogen atom, an acyl group, a C₁₋₆ alkyl group optionally substituted by one or more groups selected from substituent group C, a C₃₋₈ cycloalkyl group optionally substituted by one or more groups selected from substituent group C, a C₁₋₆ alkoxy group optionally substituted by one or more groups selected from substituent group C, a C₁₋₆ alkylamino group, a di(C₁₋₆ alkyl)amino group, an aryl group optionally substituted by one or more groups selected from substituent group C, a heterocyclic group optionally substituted by one or more groups selected from substituent group C, a cyano group, an optionally protected amino group, an optionally protected hydroxyl group, an optionally protected carboxyl group and an oxo group.

[0034]

Substituent group C: a halogen atom, an optionally protected amino group, an optionally protected hydroxyl group and an optionally protected carboxyl group.

[0035]

The hydroxyl-protective group includes every group that can be used as a usual protective group for a hydroxyl group. Examples thereof include a group described in Greene's Protective Groups in Organic Synthesis, 5th ed., p. 17-471, 2014, John Wiley & Sons, INC. Specific examples thereof include a C₁₋₆ alkyl group, an ar-C₁₋₆ alkyl group, a C₁₋₆ alkoxy-C₁₋₆ alkyl group, an acyl group, a C₁₋₆ alkoxycarbonyl group, an ar-C₁₋₆ alkoxycarbonyl group, a C₁₋₆ alkylsulfonyl group, an arylsulfonyl group, a silyl group, a tetrahydrofuranyl group and a tetrahydropyranyl group. These groups are each optionally substituted by one or more groups selected from substituent group A.

[0036]

The carboxyl-protective group includes every group that can be used as a usual protective group for an amino group. Examples thereof include a group described in Greene's Protective Groups in Organic Synthesis, 5th ed., p. 686-836, 2014, John Wiley & Sons, INC. Specific examples thereof include a C₁₋₆ alkyl group, an ar-C₁₋₆ alkyl group, a C₁₋₆ alkoxy-C₁₋₆ alkyl group, an ar-C₁₋₆ alkoxy-C₁₋₆ alkyl group and a silyl group. These groups are each optionally substituted by one or more groups selected from substituent group A.

[0037]

The amino-protective group includes every group that can be used as a usual protective group for an amino group. Examples thereof include a group described in Greene's Protective Groups in Organic Synthesis, 5th ed., p. 895-1193, 2014, John Wiley & Sons, INC. Specific examples thereof include an ar-C₁₋₆ alkyl group, a C₁₋₆ alkoxy-C₁₋₆ alkyl group, an acyl

group, a C₁₋₆ alkoxy carbonyl group, an ar-C₁₋₆ alkoxy carbonyl group, an aryloxy carbonyl group, a C₁₋₆ alkylsulfonyl group, an arylsulfonyl group and a silyl group. These groups are each optionally substituted by one or more groups selected from substituent group A.

[0038]

5 The aliphatic hydrocarbons mean pentane, hexane, heptane, cyclohexane, methylcyclohexane or ethylcyclohexane.

 The halogenated hydrocarbons mean dichloromethane, chloroform or dichloroethane.

 The ethers mean diethyl ether, diisopropyl ether, tetrahydrofuran, 2-
10 methyltetrahydrofuran, 1,4-dioxane, anisole, ethylene glycol dimethyl ether, diethylene glycol dimethyl ether or diethylene glycol diethyl ether.

 The alcohols mean methanol, ethanol, propanol, 2-propanol, butanol, 2-methyl-2-propanol, ethylene glycol, propylene glycol or diethylene glycol.

 The ketones mean acetone, 2-butanone or 4-methyl-2-pentanone.

15 The esters mean methyl acetate, ethyl acetate, propyl acetate, isopropyl acetate or butyl acetate.

 The amides mean N,N-dimethylformamide, N,N-dimethylacetamide or N-methylpyrrolidone.

 The nitriles mean acetonitrile or propionitrile.

20 The sulfoxides mean dimethyl sulfoxide or sulfolane.

 The aromatic hydrocarbons mean benzene, toluene or xylene.

[0039]

 The inorganic base means sodium hydroxide, potassium hydroxide, sodium bicarbonate, sodium carbonate, sodium hydride, potassium carbonate, tripotassium phosphate,
25 potassium acetate, cesium fluoride or cesium carbonate.

 The organic base means sodium methoxide, sodium ethoxide, sodium tert-butoxide, potassium tert-butoxide, triethylamine, N,N-diisopropylethylamine, 1,8-diazabicyclo(5.4.0)undec-7-ene (DBU), pyridine, N,N-dimethyl-4-aminopyridine or 4-methylmorpholine.

30 [0040]

 The C₁₋₆ alkyl group represented by R¹; the C₁₋₆ alkyl group, the C₂₋₆ alkenyl group, the C₃₋₈ cycloalkyl group, the C₄₋₈ cycloalkenyl group, the aryl group, the C₁₋₆ alkoxy group, the C₁₋₆ alkylamino group, the di(C₁₋₆ alkyl)amino group and the heterocyclic group represented by R³; the C₁₋₆ alkyl group, the C₁₋₃ alkyl group, the C₂₋₆ alkenyl group, the C₂₋₆

alkynyl group, the C₃₋₈ cycloalkyl group, the C₄₋₈ cycloalkenyl group, the aryl group, the C₁₋₆ alkoxy group, the aryloxy group, the C₁₋₆ alkylamino group, the C₁₋₃ alkylamino group, the di(C₁₋₆ alkyl)amino group, the arylamino group, the carbamoyl group, the sulfamoyl group, the C₁₋₆ alkylthio group, the arylthio group, the C₁₋₆ alkylsulfonyl group, the C₁₋₃ alkylsulfonyl group, the arylsulfonyl group and the heterocyclic group represented by R⁴; the C₂₋₃ alkylene group and the C₂₋₅ alkylene group formed together by two adjacent R⁴; the C₁₋₆ alkyl group represented by R⁵; the C₁₋₆ alkyl group represented by R⁶; the C₁₋₃ alkyl group represented by R^{6a}; the C₁₋₃ alkyl group represented by R^{6b}; the C₁₋₆ alkyl group represented by R⁷; the C₁₋₃ alkylene group represented by Y¹; the C₁₋₃ alkylene group represented by Y²; the C₁₋₃ alkylene group represented by Y³; the C₁₋₃ alkyl group represented by R^{7a}; the C₁₋₃ alkyl group represented by R^{7b}; the C₂₋₄ alkylene group formed together by R⁴ and R⁷; the C₂₋₃ alkylene group formed together by R⁴ and R^{7a}; the C₂₋₃ alkylene group formed together by R⁴ and R^{7b}; the C₁₋₆ alkyl group and the aryl group represented by R⁸; the divalent cyclic hydrocarbon group resulting from the removal of one hydrogen atom from each of two adjacent atoms and the divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms, represented by X¹; the divalent cyclic hydrocarbon group resulting from the removal of one hydrogen atom from each of two adjacent atoms and the divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms, represented by X^{1A}; the C₁₋₆ alkyl group represented by R^a; the C₁₋₆ alkyl group represented by R^b; the C₁₋₃ alkylene group formed together by R^a and R^b; the C₁₋₆ alkyl group represented by R^c; the C₁₋₆ alkyl group represented by R^d; and the C₁₋₃ alkylene group formed together by R^c and R^d are each optionally substituted by one or more groups selected from substituent group A.

The C₁₋₆ alkyl group represented by R² is optionally substituted by one or more groups selected from substituent group B.

[0041]

In the present invention, examples of preferred compounds include the following compounds or salts thereof.

R¹ is an optionally substituted C₁₋₆ alkyl group.

A compound in which R¹ is a C₁₋₃ alkyl group is preferred. A compound in which R¹ is an ethyl group is more preferred.

[0042]

R² is a hydrogen atom, a halogen atom or an optionally substituted C₁₋₆ alkyl group.

A compound in which R² is a hydrogen atom or an optionally substituted C₁₋₆

alkyl group is preferred. A compound in which R^2 is a hydrogen atom is more preferred.
[0043]

R^3 is a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{3-8} cycloalkyl group, an optionally substituted C_{4-8} cycloalkenyl group, an optionally substituted aryl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted C_{1-6} alkylamino group, an optionally substituted di(C_{1-6} alkyl)amino group or an optionally substituted heterocyclic group.

A compound in which R^3 is a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{3-8} cycloalkyl group, an optionally substituted C_{4-8} cycloalkenyl group, an optionally substituted aryl group, an optionally substituted di(C_{1-6} alkyl)amino group or an optionally substituted heterocyclic group is preferred.

A compound in which R^3 is an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{3-8} cycloalkyl group, an optionally substituted aryl group or an optionally substituted heterocyclic group is more preferred.

A compound in which R^3 is an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{3-8} cycloalkyl group or an optionally substituted heterocyclic group is further preferred.

A compound in which R^3 is an optionally substituted C_{1-6} alkyl group or an optionally substituted heterocyclic group is still further preferred.

A compound in which R^3 is an optionally substituted heterocyclic group is particularly preferred.

[0044]

Z^1 , Z^2 and Z^3 are the same or different and each are a nitrogen atom or a group represented by the general formula CR^5 wherein R^5 is as defined above.

A compound in which each of Z^1 , Z^2 and Z^3 is a group represented by the general formula CR^5 wherein R^5 is as defined above is preferred. A compound in which each of Z^1 , Z^2 and Z^3 is CH is more preferred.

[0045]

X^1 is (1) a group represented by the general formula $C(=O)N(R^6)$ wherein R^6 is as defined above, (2) a group represented by the general formula $N(R^7)C(=O)$ wherein R^7 is as defined above, (3) an optionally substituted divalent cyclic hydrocarbon group resulting from the removal of one hydrogen atom from each of two adjacent atoms or (4) an optionally substituted divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two

adjacent atoms.

[0046]

A compound in which X^1 is a group represented by the general formula $C(=O)N(R^6)$ wherein R^6 is as defined above is preferred. A compound in which X^1 is a group represented by the general formula $C(=O)N(R^{6a})$ wherein R^{6a} is as defined above is more preferred. A compound in which X^1 is a group represented by the general formula $C(=O)N(R^{6b})$ wherein R^{6b} is as defined above is further preferred.

[0047]

In an alternative embodiment, a compound in which X^1 is a group represented by the general formula $N(R^7)C(=O)$ wherein R^7 is as defined above is preferred. A compound in which X^1 is a group represented by the general formula $N(R^{7a})C(=O)$ wherein R^{7a} is as defined above is more preferred. A compound in which X^1 is a group represented by the general formula $N(R^{7b})C(=O)$ wherein R^{7b} is as defined above is further preferred.

[0048]

In a further alternative embodiment, a compound in which X^1 is an optionally substituted divalent cyclic hydrocarbon group resulting from the removal of one hydrogen atom from each of two adjacent atoms or an optionally substituted divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms is preferred.

A compound in which X^1 is an optionally substituted divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms is more preferred.

[0049]

Ring A is a cyclic hydrocarbon group or a heterocyclic group.

A compound in which ring A is a cyclic hydrocarbon group is preferred.

A compound in which ring A is a phenyl group is more preferred.

[0050]

“m” number of R^4 are the same or different and each are a halogen atom, a cyano group, a nitro group, an amino-protective group, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{3-8} cycloalkyl group, an optionally substituted C_{4-8} cycloalkenyl group, an optionally substituted aryl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted aryloxy group, an optionally substituted C_{1-6} alkylamino group, an optionally substituted di(C_{1-6} alkyl)amino group, an optionally substituted arylamino group, an optionally substituted carbamoyl group, an optionally substituted sulfamoyl group, an optionally substituted

C₁₋₆ alkylthio group, an optionally substituted arylthio group, an optionally substituted C₁₋₆ alkylsulfonyl group, an optionally substituted arylsulfonyl group, an optionally substituted heterocyclic group, an optionally protected amino group, an optionally protected hydroxyl group or an optionally protected carboxyl group. In this context, two adjacent R⁴ may together form an optionally substituted C₂₋₅ alkylene group, and one R⁴ may form, together with R⁷, an optionally substituted C₂₋₄ alkylene group, a group represented by the general formula O-Y¹ wherein Y¹ is as defined above, a group represented by the general formula S(O)_n-Y² wherein Y² and n are as defined above or a group represented by the general formula N(R⁸)-Y³ wherein Y³ and R⁸ are as defined above.

10 [0051]

A compound in which m number of R⁴ are the same or different and each are a halogen atom, a cyano group, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₃₋₈ cycloalkyl group, an optionally substituted C₄₋₈ cycloalkenyl group, an optionally substituted aryl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted C₁₋₆ alkylamino group, an optionally substituted di(C₁₋₆ alkyl)amino group, an optionally substituted arylamino group, an optionally substituted carbamoyl group, an optionally substituted sulfamoyl group, an optionally substituted C₁₋₆ alkylsulfonyl group, an optionally substituted heterocyclic group, an optionally protected amino group, an optionally protected hydroxyl group or an optionally protected carboxyl group is preferred.

A compound in which m number of R⁴ are the same or different and each are a halogen atom, an optionally substituted C₁₋₃ alkyl group, an optionally substituted C₁₋₃ alkylamino group, an optionally substituted C₁₋₃ alkylsulfonyl group or an optionally protected amino group is more preferred.

25 A compound in which m number of R⁴ are the same or different and each are a halogen atom, an optionally substituted C₁₋₃ alkyl group, an optionally substituted C₁₋₃ alkylamino group or an optionally protected amino group is further preferred.

A compound in which m number of R⁴ are the same or different and each are a halogen atom or an optionally substituted C₁₋₃ alkyl group is still further preferred. A compound in which each of m number of R⁴ is a halogen atom is particularly preferred.

30 [0052]

In an alternative embodiment, a compound in which m number of R⁴ are the same or different and each are a halogen atom, a cyano group, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally

substituted carbamoyl group, an optionally substituted C₁₋₆ alkylsulfonyl group, an optionally substituted heterocyclic group, an optionally protected amino group, an optionally protected carboxyl group,

- an optionally substituted C₂₋₅ alkylene group formed together by two adjacent R⁴, an optionally substituted C₂₋₄ alkylene group formed together by one R⁴ and R⁷,
 5 a group represented by the general formula O-Y¹ formed together by one R⁴ and R⁷ wherein Y¹ is as defined above,
 a group represented by the general formula S(O)_n-Y² formed together by one R⁴ and R⁷ wherein Y² is as defined above, or
 10 a group represented by the general formula N(R⁸)-Y³ formed together by one R⁴ and R⁷ wherein R⁸ and Y³ are as defined above
 is preferred.

[0053]

A compound in which

- 15 m number of R⁴ are the same or different and each are a halogen atom, an optionally substituted C₁₋₃ alkyl group, an optionally substituted C₁₋₃ alkylamino group, an optionally substituted C₁₋₃ alkylsulfonyl group, an optionally protected amino group,
 an optionally substituted C₂₋₅ alkylene group formed together by two adjacent R⁴, an optionally substituted C₂₋₄ alkylene group formed together by one R⁴ and R⁷,
 20 a group represented by the general formula O-Y¹ formed together by one R⁴ and R⁷ wherein Y¹ is as defined above,
 a group represented by the general formula S(O)_n-Y² formed together by one R⁴ and R⁷ wherein Y² is as defined above, or
 a group represented by the general formula N(R⁸)-Y³ formed together by one R⁴ and R⁷ wherein
 25 R⁸ and Y³ are as defined above
 is more preferred.

[0054]

A compound in which

- m number of R⁴ are the same or different and each are a halogen atom, an optionally substituted
 30 C₁₋₃ alkyl group, an optionally substituted C₁₋₃ alkylsulfonyl group,
 an optionally substituted C₂₋₃ alkylene group formed together by two adjacent R⁴, an optionally substituted C₂₋₄ alkylene group formed together by one R⁴ and R⁷,
 a group represented by the general formula O-Y¹ formed together by one R⁴ and R⁷ wherein Y¹ is as defined above,

a group represented by the general formula $S(O)_n-Y^2$ formed together by one R^4 and R^7 wherein Y^2 is as defined above, or

a group represented by the general formula $N(R^8)-Y^3$ formed together by one R^4 and R^7 wherein R^8 and Y^3 are as defined above

5 is further preferred.

[0055]

A compound in which

m number of R^4 are the same or different and each are a halogen atom, an optionally substituted C_{1-3} alkyl group, an optionally substituted C_{1-3} alkylsulfonyl group,

10 an optionally substituted C_{2-3} alkylene group formed together by two adjacent R^4 , an optionally substituted C_{2-3} alkylene group formed together by one R^4 and R^{7a} , or

a group represented by the general formula $O-Y^{1a}$ formed together by one R^4 and R^{7a} wherein Y^{1a} is as defined above

is still further preferred.

15 [0056]

A compound in which

m number of R^4 are the same or different and each are an optionally substituted C_{1-3} alkyl group, an optionally substituted C_{2-3} alkylene group formed together by two adjacent R^4 , or an optionally substituted C_{2-3} alkylene group formed together by one R^4 and R^{7b}

20 is particularly preferred.

A compound in which m number of R^4 are the same or different and each are an optionally substituted C_{1-3} alkyl group or an optionally substituted C_{2-3} alkylene group formed together by one R^4 and R^{7b} is most preferred.

[0057]

25 In a further alternative embodiment, a compound in which m number of R^4 are the same or different and each are a halogen atom is preferred.

[0058]

“m” is an integer of 0 to 5.

A compound in which m is an integer of 0 to 2 is preferred.

30 [0059]

R^5 is a hydrogen atom, a halogen atom or an optionally substituted C_{1-6} alkyl group.

A compound in which R^5 is a hydrogen atom is preferred.

[0060]

R^6 is a hydrogen atom, an amino-protective group or an optionally substituted C_{1-6} alkyl group.

A compound in which R^6 is a hydrogen atom or an optionally substituted C_{1-3} alkyl group is preferred. A compound in which R^6 is an optionally substituted C_{1-3} alkyl group is more preferred.
[0061]

R^7 is a hydrogen atom, an amino-protective group or an optionally substituted C_{1-6} alkyl group. Alternatively, R^7 is, together with one substituent R^4 on ring A, an optionally substituted C_{2-4} alkylene group, a group represented by the general formula $O-Y^1$ wherein Y^1 is as defined above, a group represented by the general formula $S(O)_n-Y^2$ wherein Y^2 and n are as defined above, or a group represented by the general formula $N(R^8)-Y^3$ wherein Y^3 and R^8 are as defined above.

A compound in which R^7 is an optionally substituted C_{1-3} alkyl group, or a compound in which R^7 is, together with one substituent R^4 on ring A, an optionally substituted C_{2-3} alkylene group, a group represented by the general formula $O-Y^{1a}$ wherein Y^{1a} is as defined above, or a group represented by the general formula $N(R^{8a})-Y^3$ wherein R^{8a} represents an optionally substituted C_{1-3} alkyl group; and Y^3 is as defined above is preferred.

A compound in which R^7 is an optionally substituted C_{1-3} alkyl group, or a compound in which R^7 is, together with one substituent R^4 on ring A, an optionally substituted C_{2-3} alkylene group or a group represented by the general formula $O-Y^{1a}$ wherein Y^{1a} is as defined above is more preferred.

A compound in which R^7 is an optionally substituted C_{1-3} alkyl group, or a compound in which R^7 is, together with one substituent R^4 on ring A, an optionally substituted C_{2-3} alkylene group is further preferred.
[0062]

R^8 is a hydrogen atom, an amino-protective group, an optionally substituted C_{1-6} alkyl group or an optionally substituted aryl group.

A compound in which R^8 is a hydrogen atom or an optionally substituted C_{1-6} alkyl group is preferred. A compound in which R^8 is a hydrogen atom is more preferred.
[0063]

Y^1 is an optionally substituted C_{1-3} alkylene group.

A compound in which Y^1 is an ethylene group is preferred.

Y^2 is an optionally substituted C_{1-3} alkylene group.

A compound in which Y^2 is a C_{1-3} alkylene group is preferred.

Y^3 is an optionally substituted C_{1-3} alkylene group.

A compound in which Y^3 is a C_{1-3} alkylene group is preferred.

[0064]

In an alternative embodiment, a compound in which

- 5 R^2 is a hydrogen atom or an optionally substituted C_{1-6} alkyl group; and
each of Z^1 , Z^2 and Z^3 is CH
is preferred.

[0065]

In an alternative embodiment, a compound in which

- 10 R^1 is a C_{1-3} alkyl group, and
 R^3 is an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{3-8} cycloalkyl group,
an optionally substituted aryl group or an optionally substituted heterocyclic group
is preferred.

[0066]

- 15 In an alternative embodiment, a compound in which

X^1 is a group represented by the general formula $C(=O)N(R^{6a})$ wherein R^{6a} is as defined above;
m number of R^4 are the same or different and each are a halogen atom, an optionally substituted
 C_{1-3} alkyl group, an optionally substituted C_{1-3} alkylamino group or an optionally protected
amino group; and

- 20 m is an integer of 0 to 2
is preferred.

[0067]

In an alternative embodiment, a compound in which

- R^3 is an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{3-8} cycloalkyl group or
25 an optionally substituted heterocyclic group;
 X^1 is a group represented by the general formula $C(=O)N(R^{6b})$ wherein R^{6b} is as defined above;
m number of R^4 are the same or different and each are a halogen atom or an optionally
substituted C_{1-3} alkyl group; and
m is an integer of 0 to 2

- 30 is preferred.

[0068]

In an alternative embodiment, a compound in which

R^3 is an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{3-8} cycloalkyl group or
an optionally substituted heterocyclic group;

X^1 is a group represented by the general formula $N(R^{7a})C(=O)$ wherein R^{7a} is as defined above;
 m number of R^4 are the same or different and each are a halogen atom, an optionally substituted
 C_{1-3} alkyl group, an optionally substituted C_{1-3} alkylsulfonyl group,
 an optionally substituted C_{2-3} alkylene group formed together by two adjacent R^4 ,
 5 an optionally substituted C_{2-3} alkylene group formed together by one R^4 and R^{7a} , or
 a group represented by the general formula $O-Y^{1a}$ formed together by one R^4 and R^{7a} wherein
 Y^{1a} is as defined above; and
 m is an integer of 0 to 2
 is preferred.

10 [0069]

A compound in which

R^3 is an optionally substituted heterocyclic group;

X^1 is a group represented by the general formula $N(R^{7b})C(=O)$ wherein R^{7b} is as defined above;
 m number of R^4 are the same or different and each are an optionally substituted C_{1-3} alkyl group,
 15 an optionally substituted C_{2-3} alkylene group formed together by two adjacent R^4 , or
 an optionally substituted C_{2-3} alkylene group formed together by one R^4 and R^{7b} ; and
 m is an integer of 0 to 2
 is more preferred.

[0070]

20

A compound in which

R^3 is an optionally substituted heterocyclic group;

X^1 is a group represented by the general formula $N(R^{7b})C(=O)$ wherein R^{7b} is as defined above;
 m number of R^4 are the same or different and each are an optionally substituted C_{1-3} alkyl group,
 or

25 an optionally substituted C_{2-3} alkylene group formed together by one R^4 and R^{7b} ; and
 m is an integer of 0 to 2
 is further preferred.

[0071]

In an alternative embodiment, a compound in which

30 R^3 is an optionally substituted C_{1-6} alkyl group or an optionally substituted heterocyclic group;
 X^1 is an optionally substituted divalent cyclic hydrocarbon group resulting from the removal of
 one hydrogen atom from each of two adjacent atoms or an optionally substituted divalent
 heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent
 atoms;

m number of R⁴ are the same or different and each are a halogen atom;
 m is an integer of 0 to 2
 is preferred.

A compound in which

- 5 R³ is an optionally substituted heterocyclic group;
 X¹ is an optionally substituted divalent heterocyclic group resulting from the removal of one
 hydrogen atom from each of two adjacent atoms;
 m number of R⁴ are the same or different and each are a halogen atom; and
 m is an integer of 0 to 2
 10 is more preferred.
 [0072]

Examples of preferred compounds according to the present invention include the
 following compounds.

[0073]

- 15 6-(3,4-dihydroquinolin-1(2H)-ylcarbonyl)-1-ethyl-4-(1-methylpiperidin-4-
 yl)quinolin-2(1H)-one, 1-ethyl-N-methyl-N-(4-methylphenyl)-4-(1-methylpiperidin-4-yl)-2-oxo-
 1,2-dihydroquinoline-6-carboxamide, N-(2,3-dihydro-1H-inden-5-yl)-1-ethyl-N-methyl-4-(1-
 methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide, 6-(5-(4-chlorophenyl)-2-oxo-
 2,3-dihydro-1H-imidazol-1-yl)-1-ethyl-4-(morpholin-4-yl)quinolin-2(1H)-one, 6-(5-(4-
 20 chlorophenyl)-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-1-yl)-1-ethyl-4-(morpholin-4-
 yl)quinolin-2(1H)-one, 1-ethyl-4-(morpholin-4-yl)-6-(2-oxo-5-phenyl-3-(propan-2-yl)-2,3-
 dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one, 1-ethyl-N-(3-fluoro-4-methylphenyl)-N-methyl-
 4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide, N-(3-chloro-4-
 methylphenyl)-1-ethyl-N-methyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-
 25 carboxamide and N-(3,4-dimethylphenyl)-1-ethyl-N-methyl-4-(1-methylpiperidin-4-yl)-2-oxo-
 1,2-dihydroquinoline-6-carboxamide.

[0074]

- Examples of the salt of the compound represented by the general formula [1] can
 include a usually known salt of a basic group such as an amino group or an acidic group such as
 30 a hydroxyl group or a carboxyl group.

[0075]

Examples of the salt of a basic group include: a salt with a mineral acid such as
 hydrochloric acid, hydrobromic acid, nitric acid and sulfuric acid; a salt with an organic
 carboxylic acid such as formic acid, acetic acid, citric acid, oxalic acid, fumaric acid, maleic

acid, succinic acid, malic acid, tartaric acid, aspartic acid, trichloroacetic acid and trifluoroacetic acid; and a salt with a sulfonic acid such as methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, mesitylenesulfonic acid and naphthalenesulfonic acid.

[0076]

5 Examples of the salt of an acidic group include: a salt with an alkali metal such as sodium and potassium; a salt with an alkaline earth metal such as calcium and magnesium; an ammonium salt; and a salt with a nitrogen-containing organic base such as trimethylamine, triethylamine, tributylamine, pyridine, N,N-dimethylaniline, N-methylpiperidine, 4-methylmorpholine, diethylamine, dicyclohexylamine, procaine, dibenzylamine, N-benzyl-β-phenethylamine, 1-phenamine and N,N'-dibenzylethylenediamine.

[0077]

 Among the salts described above, examples of preferred salts include pharmacologically acceptable salts.

 When the compound represented by the general formula [1] has isomers (e.g.,
15 optical isomers, geometric isomers and tautomers), the present invention encompasses these isomers and also encompasses solvates, hydrates and various forms of crystals.


[0078]

 Next, methods for producing the compound of the present invention will be described.

20 The compound of the present invention is produced by a combination of methods known per se in the art and can be produced according to, for example, the following production methods:

[0079]

[Production method 1]



25

wherein L^1 represents a hydroxyl group or a leaving group; and R^1 , R^2 , R^3 , R^4 , R^6 , Z^1 , Z^2 , Z^3 , A and m are as defined above.

[0080]

(1-1) In the case where L^1 is a hydroxyl group

For example, p-chlorobenzoic acid is known as the compound represented by the general formula [3].

The compound represented by the general formula [1a] can be produced by reacting the compound represented by the general formula [2] with the compound represented by the general formula [3] in the presence of a condensing agent or an acid halide and in the presence of a base.

The solvent for use in this reaction is not particularly limited as long as the solvent does not influence the reaction. Examples thereof include halogenated hydrocarbons, ethers, esters, amides, nitriles, sulfoxides and aromatic hydrocarbons. These solvents may be used as a mixture.

Preferred examples of the solvent include halogenated hydrocarbons, ethers, esters and amides. Halogenated hydrocarbons and amides are more preferred.

The amount of the solvent used is not particularly limited and can be 1 to 500 times (v/w) the amount of the compound represented by the general formula [2].

Examples of the base for use in this reaction include an inorganic base and an organic base.

Preferred examples of the base include an organic base. Triethylamine, N,N-diisopropylethylamine and 4-methylmorpholine are more preferred, and N,N-diisopropylethylamine and 4-methylmorpholine are further preferred.

The amount of the base used can be 1 to 50 times, preferably 1 to 10 times the mol of the compound represented by the general formula [2].

[0081]

Examples of the condensing agent for use in this reaction include: carbodiimides such as N,N'-diisopropylcarbodiimide (DIC), N,N'-di-(tert-butyl)carbodiimide, N,N'-dicyclohexylcarbodiimide (DCC), N-(tert-butyl)-N'-ethylcarbodiimide (BEC), N-cyclohexyl-N'-(2-morpholinoethyl)carbodiimide (CMC) and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC); imidazoliums such as 1,1'-carbonyldiimidazole (CDI) and 1,1'-carbonyldi(1,2,4-triazole) (CDT); acid azides such as diphenylphosphorylazide; acid cyanides such as diethylphosphorylcyanide; 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline; and uroniums such as O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), O-(benzotriazol-1-yl)-N,N,N',N'-bis(tetramethylene)uronium hexafluorophosphate (HBPYU), O-(benzotriazol-1-yl)-N,N,N',N'-bis(pentamethylene)uronium hexafluorophosphate (HBPipU), O-(6-chlorobenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HCTU), O-

(3,4-dihydro-4-oxo-1,2,3-benzotriazin-3-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HDBTU), O-(2-oxo-1(2H)pyridyl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (TPTU), O-((ethoxycarbonyl)cyanomethyleneamino)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HOTU), O-((ethoxycarbonyl)cyanomethyleneamino)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TOTU), N,N,N',N'-tetramethyl-O-(N-succinimidyl)uronium hexafluorophosphate (HSTU), N,N,N',N'-tetramethyl-O-(N-succinimidyl)uronium tetrafluoroborate (TSTU) dipyrrolidino(N-succinimidyl)oxy)carbenium hexafluorophosphate (HSPyU) and S-(1-oxido-2-pyridyl)-N,N,N',N'-tetramethylthiuronium tetrafluoroborate (TOTT).

Preferred examples of the condensing agent include carbodiimides. EDC is more preferred.

The amount of the condensing agent used can be 1 to 50 times, preferably 1 to 5 times the mol of the compound represented by the general formula [2].

[0082]

In the case of using a carbodiimide as the condensing agent, it is preferred to add an additive.

Examples of the additive include 1-hydroxybenzotriazole (HOBT), 1-hydroxy-7-azabenzotriazole (HOAT) and ethyl (hydroxyimino)cyanoacetate. HOBT and ethyl (hydroxyimino)cyanoacetate are preferred.

The amount of the additive used can be 0.01 to 10 times, preferably 0.1 to 1 times the mol of the compound represented by the general formula [2].

[0083]

Examples of the acid halide for use in this reaction include: oxalyl chloride; carboxylic acid halides such as acetyl chloride and trifluoroacetyl chloride; sulfonic acid halides such as methanesulfonyl chloride and tosyl chloride; and chloroformic acid esters such as ethyl chloroformate and isobutyl chloroformate.

The amount of the compound represented by the general formula [3] used is not particularly limited and can be 1 to 10 times the mol of the compound represented by the general formula [2].

This reaction can be carried out at -30 to 150°C, preferably 0 to 100°C, for 30 minutes to 48 hours.

[0084]

(1-2) In the case where L¹ is a leaving group

For example, p-chlorobenzoyl chloride is known as the compound represented by

the general formula [3].

The compound represented by the general formula [1a] can be produced by reacting the compound represented by the general formula [2] with the compound represented by the general formula [3] in the presence of a base.

5 The solvent for use in this reaction is not particularly limited as long as the solvent does not influence the reaction. Examples thereof include halogenated hydrocarbons, ethers, esters, amides, nitriles, sulfoxides and aromatic hydrocarbons. These solvents may be used as a mixture.

 Preferred examples of the solvent include halogenated hydrocarbons, ethers and
10 esters. Halogenated hydrocarbons are more preferred.

The amount of the solvent used is not particularly limited and can be 1 to 500 times (v/w) the amount of the compound represented by the general formula [2].

[0085]

 Examples of the base for use in this reaction include an inorganic base and an
15 organic base.

 Preferred examples of the base include an organic base. Triethylamine, N,N-diisopropylethylamine, 4-methylmorpholine and pyridine are more preferred, and N,N-diisopropylethylamine, 4-methylmorpholine and pyridine are further preferred.


 The amount of the base used can be 1 to 50 times, preferably 1 to 10 times the
20 mol of the compound represented by the general formula [2].

 The amount of the compound represented by the general formula [3] used is not particularly limited and can be 1 to 10 times the mol of the compound represented by the general formula [2].

 This reaction can be carried out at -30 to 150°C, preferably 0 to 100°C, for 30
25 minutes to 48 hours.

[0086]

[Production method 2]



wherein R^{6A} represents an optionally substituted C_{1-6} alkyl group; L^2 represents a leaving group; and R^1 , R^2 , R^3 , R^4 , L^1 , Z^1 , Z^2 , Z^3 , A and m are as defined above.

[0087]

5 <Step 1>

The compound represented by the general formula [1b] can be produced by reacting the compound represented by the general formula [4] with the compound represented by the general formula [3].

This reaction can be carried out according to Production method 1.

10 [0088]

<Step 2>

For example, methyl iodide is known as the compound represented by the general formula [5].

The compound represented by the general formula [1c] can be produced by reacting the compound represented by the general formula [1b] with the compound represented by the general formula [5] in the presence of a base.

The solvent for use in this reaction is not particularly limited as long as the solvent does not influence the reaction. Examples thereof include halogenated hydrocarbons, ethers, ketones, esters, amides, nitriles, sulfoxides and aromatic hydrocarbons. These solvents may be used as a mixture.

Preferred examples of the solvent include ethers and amides. Amides are more preferred.

The amount of the solvent used is not particularly limited and can be 1 to 500 times (v/w) the amount of the compound represented by the general formula [1b].

25 [0089]

Examples of the base for use in this reaction include: an organic base such as sodium methoxide, sodium ethoxide, potassium tert-butoxide, pyridine, dimethylaminopyridine and triethylamine; and an inorganic base such as sodium hydride, sodium hydroxide, potassium hydroxide, sodium bicarbonate, potassium carbonate and sodium carbonate. Preferred

5 examples of the base include sodium hydride and potassium carbonate.


The amount of the base used is not particularly limited and can be 1 to 20 times, preferably 1 to 5 times the mol of the compound represented by the general formula [1b].

The amount of the compound represented by the general formula [5] used is not particularly limited and can be 1 to 10 times the mol of the compound represented by the general
10 formula [1b].

This reaction can be carried out at -30 to 150°C, preferably 0 to 100°C, for 30 minutes to 48 hours.

[0090]

[Production method 3]



wherein L^3 represents a hydroxyl group or a leaving group; and $\text{R}^1, \text{R}^2, \text{R}^3, \text{R}^4, \text{R}^7, \text{Z}^1, \text{Z}^2, \text{Z}^3, \text{A}$ and m are as defined above.

[0091]


For example, N-methylaniline is known as the compound represented by the
20 general formula [7].

The compound represented by the general formula [1d] can be produced by reacting the compound represented by the general formula [6] with the compound represented by the general formula [7].

This reaction can be carried out according to Production method 1.

25 [0092]

[Production method 4]



wherein R^{7A} represents an optionally substituted C_{1-6} alkyl group; L^4 represents a leaving group; and R^1 , R^2 , R^3 , R^4 , L^3 , Z^1 , Z^2 , Z^3 , A and m are as defined above.

[0093]

5 <Step 1>

The compound represented by the general formula [1e] can be produced by reacting the compound represented by the general formula [6] with the compound represented by the general formula [8].

This reaction can be carried out according to Production method 1.

10 [0094]

<Step 2>


For example, methyl iodide is known as the compound represented by the general formula [9].

The compound represented by the general formula [1f] can be produced by
 15 reacting the compound represented by the general formula [1e] with the compound represented by the general formula [9].

This reaction can be carried out according to <Step 2> of Production method 2.

[0095]

[Production method 5]



wherein R^a represents a hydrogen atom or an optionally substituted C₁₋₆ alkyl group; R^b represents a hydrogen atom or an optionally substituted C₁₋₆ alkyl group; or R^a and R^b together represent an optionally substituted C₁₋₃ alkylene group; L⁵ represents a leaving group; X^{1A} represents an optionally substituted divalent cyclic hydrocarbon group resulting from the removal of one hydrogen atom from each of two adjacent atoms or an optionally substituted divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms; and R¹, R², R³, R⁴, Z¹, Z², Z³, A and m are as defined above.

[0096]

For example, 2-biphenylboric acid is known as the compound represented by the general formula [11].

The compound represented by the general formula [1g] can be produced by reacting the compound represented by the general formula [10] with the compound represented by the general formula [11] in the presence or absence of a base, in the presence of a palladium catalyst and in the presence or absence of a ligand.

The solvent for use in this reaction is not particularly limited as long as the solvent does not influence the reaction. Examples thereof include water, alcohols, halogenated hydrocarbons, ethers, ketones, esters, amides, nitriles, sulfoxides and aromatic hydrocarbons. These solvents may be used as a mixture.

Preferred examples of the solvent include mixed solvents of aromatic hydrocarbons and water and mixed solvents of ethers and water.

The amount of the solvent used is not particularly limited and can be 1 to 500 times (v/w) the amount of the compound represented by the general formula [10].

[0097]

Examples of the palladium catalyst for use in this reaction include: metal palladium such as palladium-carbon and palladium black; an inorganic palladium salt such as palladium chloride; an organic palladium salt such as palladium acetate; chloro(2-(dicyclohexylphosphino)-3,6-dimethoxy-2',4',6'-triisopropyl-1,1'-biphenyl)(2-(2-aminoethyl)phenyl)palladium(II); an organic palladium complex such as tetrakis(triphenylphosphine)palladium(0), bis(triphenylphosphine)palladium(II) dichloride, bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 1,1'-bis(diphenylphosphino)ferrocene palladium(II) dichloride, (E)-di(μ-acetato)bis(o-(di-*o*-tolylphosphino)benzyl)dipalladium(II) and tris(dibenzylideneacetone)dipalladium(0); and a polymer-supported organic palladium complex such as polymer-supported bis(acetato)triphenylphosphine palladium(II) and polymer-supported

di(acetato)dicyclohexylphenylphosphine palladium(II). An organic palladium complex is preferred.

The amount of the palladium catalyst used can be 0.00001 to 1 times, preferably, 0.01 to 0.2 times the mol of the compound represented by the general formula [10].

5 [0098]

Examples of the ligand that is used in this reaction, if desired, include:

trialkylphosphines such as trimethylphosphine and tri-tert-butylphosphine;
 tricycloalkylphosphines such as tricyclohexylphosphine; triarylphosphines such as
 triphenylphosphine and tritolylphosphine; trialkyl phosphites such as trimethyl phosphite,
 10 triethyl phosphite and tributyl phosphite; tricycloalkyl phosphites such as tricyclohexyl
 phosphite; triaryl phosphites such as triphenyl phosphite; imidazolium salts such as 1,3-
 bis(2,4,6-trimethylphenyl)imidazolium chloride; diketones such as acetylacetone and
 octafluoroacetylacetone; amines such as trimethylamine, triethylamine, tripropylamine,
 triisopropylamine and tributylamine; 1,1'-bis(diphenylphosphino)ferrocene; 2,2'-
 15 bis(diphenylphosphino)-1,1'-binaphthyl; 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl; 2-
 dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl; 2-(di-tert-butylphosphino)-2',4',6'-
 triisopropylbiphenyl; 2-dicyclohexylphosphino-2'-(N,N-dimethylamino)biphenyl; 4,5-
 bis(diphenylphosphino)-9,9-dimethylxanthene; and 2-(di-tert-butylphosphino)biphenyl.

The amount of the ligand used can be 0.00001 to 1 times, preferably 0.02 to 0.5
 20 times the mol of the compound represented by the general formula [10].

[0099]

Examples of the base that is used in this reaction, if desired, include an inorganic
 base and an organic base. An inorganic base such as sodium bicarbonate, sodium carbonate,
 potassium carbonate, cesium carbonate and tripotassium phosphate is preferred.


25 The amount of the base used can be 1 to 50 times, preferably 2 to 10 times the
 mol of the compound represented by the general formula [10].

The amount of the compound represented by the general formula [11] used can be
 1 to 50 times, preferably 1 to 2 times the mol of the compound represented by the general
 formula [10].

30 This reaction can usually be carried out at 0 to 160°C, preferably 20 to 120°C, for
 1 minute to 96 hours in an inert gas (e.g. nitrogen and/or argon) atmosphere.

[0100]

[Production method 6]



wherein L^6 represents a leaving group; and R^1 , R^2 , R^3 , R^4 , R^a , R^b , X^{1A} , Z^1 , Z^2 , Z^3 , A and m are as defined above.


[0101]

- 5 The compound represented by the general formula [1h] can be produced by reacting the compound represented by the general formula [12] with the compound represented by the general formula [13].

This reaction can be carried out according to Production method 5.

[0102]

- 10 [Production method 7]



wherein R^{3A} represents an optionally substituted C_{1-6} alkylamino group, an optionally substituted di(C_{1-6} alkyl)amino group or an optionally substituted cyclic amino group; L^7 represents a leaving group; and R^1 , R^2 , R^4 , X^1 , Z^1 , Z^2 , Z^3 , A and m are as defined above.

- 15 [0103]

For example, morpholine is known as the compound represented by the general formula [15].

- The compound represented by the general formula [1i] can be produced by reacting the compound represented by the general formula [14] with the compound represented by the general formula [15] in the presence or absence of a base.

The solvent for use in this reaction is not particularly limited as long as the solvent does not influence the reaction. Examples thereof include halogenated hydrocarbons, ethers, esters, amides, nitriles, sulfoxides and aromatic hydrocarbons. These solvents may be used as a mixture.

- 25 Preferred examples of the solvent include halogenated hydrocarbons, ethers,

esters and amides. Amides are more preferred.

The amount of the solvent used is not particularly limited and can be 1 to 500 times (v/w) the amount of the compound represented by the general formula [14].

5 Examples of the base for use in this reaction include an inorganic base and an organic base.

Preferred examples of the base include an organic base. Triethylamine, N,N-diisopropylethylamine and 4-methylmorpholine are more preferred, and N,N-diisopropylethylamine and 4-methylmorpholine are further preferred.

10 The amount of the base used can be 1 to 50 times, preferably 1 to 10 times the mol of the compound represented by the general formula [14].

The amount of the compound represented by the general formula [15] used is not particularly limited and can be 1 to 10 times the mol of the compound represented by the general formula [14].


15 This reaction can be carried out at -30 to 150°C, preferably 0 to 150°C, for 30 minutes to 48 hours.

[0104]

Next, methods for producing starting materials for the production of the compound of the present invention will be described.

[0105]

20 [Production method A]



wherein $\text{R}^{3\text{B}}$ represents an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{3-8} cycloalkyl group, an optionally substituted aryl group or an optionally substituted heterocyclic group; R^c represents a hydrogen atom or an optionally substituted C_{1-6} alkyl group; R^d represents
 25 a hydrogen atom or an optionally substituted C_{1-6} alkyl group; or R^c and R^d together represent an optionally substituted C_{1-3} alkylene group; L^8 represents a leaving group; L^9 represents a leaving

group; and R^1 , R^2 , Z^1 , Z^2 and Z^3 are as defined above.

[0106]

<Step 1>

For example, 4-chloro-6-nitroquinolin-2(1H)-one is known as the compound
5 represented by the general formula [S1].

For example, ethyl iodide is known as the compound represented by the general
formula [S2].

The compound represented by the general formula [S3] can be produced by
reacting the compound represented by the general formula [S1] with the compound represented
10 by the general formula [S2] in the presence of a base.

The solvent for use in this reaction is not particularly limited as long as the
solvent does not influence the reaction. Examples thereof include halogenated hydrocarbons,
ethers, esters, amides, nitriles, sulfoxides and aromatic hydrocarbons. These solvents may be
used as a mixture.

15 Preferred examples of the solvent include amides and sulfoxides. Amides are
more preferred.

The amount of the solvent used is not particularly limited and can be 1 to 500
times (v/w) the amount of the compound represented by the general formula [S1].

Examples of the base for use in this reaction include an inorganic base and an
20 organic base.

Preferred examples of the base include an inorganic base. Cesium carbonate is
more preferred.

The amount of the base used can be 1 to 50 times, preferably 1 to 10 times the
mol of the compound represented by the general formula [S1].

25 The amount of the compound represented by the general formula [S2] used is not
particularly limited and can be 1 to 10 times the mol of the compound represented by the general
formula [S1].

This reaction can be carried out at -30 to 150°C, preferably 0 to 100°C, for 30
minutes to 48 hours.

30 [0107]

<Step 2>

For example, cyclopropylboric acid is known as the compound represented by the
general formula [S4].

The compound represented by the general formula [S5] can be produced by

reacting the compound represented by the general formula [S3] with the compound represented by the general formula [S4] in the presence of a base, in the presence of a palladium catalyst and in the presence or absence of a ligand.

This reaction can be carried out according to Production method 5.

5 [0108]

<Step 3>

The compound represented by the general formula [4] can be produced by subjecting the compound represented by the general formula [S5] to reduction reaction. This reaction can be carried out according to a method described in Richard C. Larock et al.,

10 Comprehensive Organic Transformations, 2nd edition, p. 823-827, 1999, John Wiley & Sons, INC. or a method equivalent thereto. Specific examples thereof include catalytic hydrogenation reaction using a metal catalyst, and reduction reaction using a metal such as iron or zinc.

[0109]

In the case of subjecting the compound represented by the general formula [S5] to
15 the catalytic hydrogenation reaction, the solvent used is not particularly limited as long as the solvent has no adverse effect on the reaction. Examples thereof include water, alcohols, amides, halogenated hydrocarbons, aromatic hydrocarbons, ethers, acetonitrile, ketones, esters, acetic acid and pyridine. These solvents may be used as a mixture.

Examples of the metal catalyst for use in this reaction include: metal palladium
20 such as palladium-carbon and palladium black; a palladium salt such as palladium oxide and palladium hydroxide; nickel metal such as Raney nickel; and a platinum salt such as platinum oxide.

The amount of the metal catalyst used is 0.001 to 5 times (w/w), preferably 0.01 to 1 times (w/w) the amount of the compound represented by the general formula [S5].

25 Examples of the hydrogen source include: hydrogen; formic acid; formate such as sodium formate, ammonium formate and triethylammonium formate; cyclohexene; and cyclohexadiene.

The amount of the hydrogen source used can be 2 to 100 times, preferably, 2 to 10 times the mol of the compound represented by the general formula [S5].

30 This reaction can be carried out at 0 to 200°C, preferably 0 to 100°C, for 1 minute to 24 hours.

[0110]

In the case of subjecting the compound represented by the general formula [S5] to the reduction reaction using a metal, the solvent used is not particularly limited as long as the

solvent has no adverse effect on the reaction. Examples thereof include water, alcohols, amides, halogenated hydrocarbons, aromatic hydrocarbons, ethers, acetonitrile, ketones and esters. These solvents may be used as a mixture.

Examples of the metal for use in this reaction include iron, zinc, tin and tin(II)

5 chloride.

The amount of the metal used is 1 to 50 times, preferably 1 to 10 times the mol of the compound represented by the general formula [S5].


Examples of the acid that is used in this reaction, if desired, include hydrogen chloride, hydrogen bromide, acetic acid and ammonium chloride.

10 The amount of the acid used can be 0.001 to 100 times (v/w), preferably 0.01 to 20 times (v/w) the amount of the compound represented by the general formula [S5].

This reaction can be carried out at 0 to 200°C, preferably 0 to 100°C, for 1 minute to 24 hours.

[0111]

15 [Production method B]



wherein R^e represents a carboxyl-protective group; and R^1 , R^2 , R^{3B} , R^c , R^d , L^5 , L^8 , Z^1 , Z^2 and Z^3 are as defined above.

[0112]

20 <Step 1>

For example, N-ethyl-4-iodoaniline is known as the compound represented by the general formula [S6].

For example, 3-tert-butoxy-3-oxopropionic acid is known as the compound represented by the general formula [S7].

The compound represented by the general formula [S8] can be produced by reacting the compound represented by the general formula [S6] with the compound represented by the general formula [S7] in the presence of a condensing agent or an acid halide and in the presence of a base.

5 This reaction can be carried out according to Production method 1.

[0113]

<Step 2>

The compound represented by the general formula [S9] can be produced by reacting the compound represented by the general formula [S8] with a dehydrating agent.

10 The solvent for use in this reaction is not particularly limited as long as the solvent does not influence the reaction. Examples thereof include halogenated hydrocarbons, ethers, esters, amides, nitriles, sulfoxides and aromatic hydrocarbons. These solvents may be used as a mixture.

15 Preferred examples of the solvent include amides and sulfoxides. Amides are more preferred.

The amount of the solvent used is not particularly limited and can be 1 to 500 times (v/w) the amount of the compound represented by the general formula [S8].

Examples of the dehydrating agent for use in this reaction include diphosphorus pentoxide, phosphorus pentachloride, phosphoryl chloride and thionyl chloride.

20 The amount of the dehydrating agent used can be 1 to 50 times, preferably 1 to 10 times the mol of the compound represented by the general formula [S8].

This reaction can be carried out at -30 to 150°C, preferably 0 to 100°C, for 30 minutes to 48 hours.

[0114]

25 <Step 3>

When L⁸ is, for example, a halogen atom, the compound represented by the general formula [S10] can be produced by reacting the compound represented by the general formula [S9] with a halogenating agent.

30 The solvent for use in this reaction is not particularly limited as long as the solvent does not influence the reaction. Examples thereof include aliphatic hydrocarbons, halogenated hydrocarbons, ethers, ketones, esters, amides, nitriles, sulfoxides and aromatic hydrocarbons. These solvents may be used as a mixture.

Preferred examples of the solvent include aromatic hydrocarbons.

The amount of the solvent used is not particularly limited and can be 1 to 500

times (v/w) the amount of the compound represented by the general formula [S9].

Examples of the halogenating agent for use in this reaction include phosphorus oxychloride, phosphoryl chloride and thionyl chloride.

The amount of the halogenating agent used can be 1 to 50 times, preferably 1 to 10 times the mol of the compound represented by the general formula [S9]. Also, the halogenating agent may be used as a solvent.

This reaction can be carried out at -30 to 150°C, preferably 0 to 100°C, for 30 minutes to 48 hours.

[0115]

10 <Step 4>

The compound represented by the general formula [S11] can be produced by reacting the compound represented by the general formula [S10] with sodium formate in the presence of a base and in the presence of a palladium catalyst.

The solvent for use in this reaction is not particularly limited as long as the solvent does not influence the reaction. Examples thereof include halogenated hydrocarbons, ethers, esters, amides, nitriles, sulfoxides and aromatic hydrocarbons. These solvents may be used as a mixture.

Preferred examples of the solvent include amides and sulfoxides. Amides are more preferred.

The amount of the solvent used is not particularly limited and can be 1 to 500 times (v/w) the amount of the compound represented by the general formula [S10].

Examples of the base for use in this reaction include an organic base.

The amount of the base used can be 1 to 50 times, preferably 1 to 10 times the mol of the compound represented by the general formula [S10].

Examples of the palladium catalyst for use in this reaction include: metal palladium such as palladium-carbon and palladium black; an inorganic palladium salt such as palladium chloride; an organic palladium salt such as palladium acetate; chloro(2-(dicyclohexylphosphino)-3,6-dimethoxy-2',4',6'-triisopropyl-1,1'-biphenyl)(2-(2-aminoethyl)phenyl)palladium(II); an organic palladium complex such as tetrakis(triphenylphosphine)palladium(0), bis(triphenylphosphine)palladium(II) dichloride, bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 1,1'-bis(diphenylphosphino)ferrocene palladium(II) dichloride, (E)-di(μ -acetato)bis(o-(di-*o*-tolylphosphino)benzyl)dipalladium(II) and tris(dibenzylideneacetone)dipalladium(0); and a polymer-supported organic palladium complex such as polymer-supported

bis(acetato)triphenylphosphine palladium(II) and polymer-supported di(acetato)dicyclohexylphenylphosphine palladium(II). An organic palladium complex is preferred.

The amount of the palladium catalyst used can be 0.00001 to 1 times, preferably 0.01 to 0.2 times the mol of the compound represented by the general formula [S10].

The amount of sodium formate used can be 1 to 50 times, preferably 1 to 10 times the mol of the compound represented by the general formula [S10].

This reaction can be carried out at -30 to 150°C, preferably 0 to 100°C, for 30 minutes to 48 hours.

[0116]

<Step 5>

For example, tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydropyridine-1(2H)-carboxylate is known as the compound represented by the general formula [S4].

The compound represented by the general formula [6a] can be produced by reacting the compound represented by the general formula [S11] with the compound represented by the general formula [S4] in the presence of a base, in the presence of a palladium catalyst and in the presence or absence of a ligand.

This reaction can be carried out according to Production method 5.

[0117]

When each compound used in these production methods has solvates, hydrates and various forms of crystals, these solvates, hydrates and various forms of crystals can also be used.

Among the compounds used in these production methods, a compound having, for example, an amino group, a hydroxyl group or a carboxyl group can be subjected to the protection of the group with a usual protective group in advance, and after the reaction, the protective group can be eliminated by a method known per se in the art.

Each compound obtained by these production methods can be converted to a different compound, for example, by subjecting the compound to a reaction known per se in the art such as condensation, addition, oxidation, reduction, dislocation, substitution, halogenation, dehydration or hydrolysis or by appropriately combining these reactions.

[0118]

In the case of using the compound represented by the general formula [1] as a medicament, the compound represented by the general formula [1] may be appropriately mixed

with a pharmaceutical aid usually used in formulation, such as an excipient, a carrier and a diluent. Such a preparation can be administered orally or parenterally in a form such as a tablet, a capsule, a powder, a syrup, granules, a pill, a suspension, an emulsion, a solution, a dust, a suppository, eye drops, nasal drops, ear drops, a patch, an ointment or an injection according to a routine method. The administration method, the dose and the number of doses can be appropriately selected according to the age, body weight and symptoms of a patient. Usually, the compound represented by the general formula [1] can be administered orally or parenterally (e.g. administered through injection, administered through intravenous drip, and administered to a rectal site) to an adult at a daily dose of 0.01 to 1000 mg/kg, which is administered in one portion or several portions.

[0119]

Examples of the disease involving the overproduction of CXCL10 include an immune disease such as inflammatory bowel disease, arthritis, psoriasis, systemic sclerosis, systemic lupus erythematosus and autoimmune neuroinflammatory disease.

The pharmaceutical composition means a composition containing the compound of the present invention or the salt thereof as an active ingredient appropriately mixed with a pharmaceutical aid usually used in formulation, such as an excipient, a carrier and a diluent.

The agent for treatment means a pharmaceutical composition aimed at treatment.

The treatment includes prophylaxis or therapy. The prophylaxis includes the inhibition of development, reduction in the risk of development and the delay of development. The therapy includes the amelioration of a target disease or condition and the suppression (sustentation or delay) of progression of the disease or the condition. The recipient for the treatment includes a human or a nonhuman animal in need of the treatment.

[0120]

Next, the present invention will be described with reference to Reference Examples, Examples and Test Examples. However, the present invention is not intended to be limited by them.

[0121]

Purification by column chromatography employed an automatic purification apparatus ISOLERA (Biotage Japan Ltd.) or a medium-pressure liquid chromatograph YFLC-W Prep 2XY.N (Yamazen Corp.), unless otherwise specified.

The carrier used in silica gel column chromatography was SNAP KP-Sil Cartridge (Biotage Japan Ltd.) or Hi-Flash Column W001, W002, W003, W004 or W005 (Yamazen Corp.), and the carrier used in basic silica gel column chromatography was SNAP KP-

NH Cartridge (Biotage Japan Ltd.), unless otherwise specified.

The mixing ratio of an eluent is a volume ratio. For example, "hexane:ethyl acetate gradient elution = 100:0 - 50:50" means that an eluent composed of 100% hexane and 0% ethyl acetate was finally changed to an eluent composed of 50% hexane and 50% ethyl acetate.

5 [0122]

The flow hydrogenation reaction apparatus used was H-Cube (ThalesNano Nanotechnology Inc.).

The microwave apparatus used was Initiator+ or Initiator Sixty (both from Biotage Japan Ltd.).

10 MS spectra were measured using ACQUITY SQD LC/MS System (Waters Corp., ionization method: ESI (electrospray ionization), model M-8000 (Hitachi, Ltd., ionization: ESI), LCMS-2010EV (Shimadzu Corp., ionization: simultaneous ionization method of ESI and APCI (atmospheric pressure chemical ionization)) or JMS-T100LP (DART) (JEOL Ltd., ionization: DART (direct analysis in real time)).

15 NMR spectra were measured using tetramethylsilane as internal standards and Bruker AV300 (Bruker Corp.) or model JNM-AL400 (JEOL Ltd.), and total δ values were indicated by ppm.

[0123]

Abbreviations in NMR measurement have the following meanings:

20

s: Singlet

brs: Broad singlet

d: Doublet

dd: Double doublet

t: Triplet

25

q: Quartet

quint: Quintet

m: Multiplet

DMSO-D₆: deuterated dimethyl sulfoxide

[0124]


30 Abbreviations in Reference Examples and Examples have the following meanings:

Bn: Benzyl

Me: Methyl

[0125]

Reference Example 1




To a suspension of 18.5 g of 4-chloro-6-nitroquinolin-2(1H)-one in 150 mL of N,N-dimethylacetamide was added 53.8 g of cesium carbonate at 40-50°C. To the reaction mixture under ice cooling was added 7.91 mL of ethyl iodide, and then the mixture was stirred at room temperature for 3 hours. To the reaction mixture were added ethyl acetate and water, and the pH of the mixture was adjusted to 2.0 with 2 mol/L hydrochloric acid. The solid matter was filtered and washed with water to obtain a slightly brown solid. To the obtained solid were added ethyl acetate and diisopropyl ether, and then the solid matter was filtered and washed with ethyl acetate to obtain 8.88 g of 4-chloro-1-ethyl-6-nitroquinolin-2(1H)-one as a slightly brown solid.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.39 (3H, t, $J = 7.3$ Hz), 4.39 (2H, q, $J = 7.3$ Hz), 7.01 (1H, s), 7.51 (1H, d, $J = 9.9$ Hz), 8.48 (1H, dd, $J = 9.2, 2.6$ Hz), 8.94 (1H, d, $J = 2.6$ Hz).

[0126]

15 Reference Example 2



A mixture of 5.17 g of 4-chloro-1-ethyl-6-nitroquinolin-2(1H)-one, 4.26 g of cyclopropylboronic acid monohydrate, 10.9 g of sodium carbonate, 0.44 g of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 50 mL of ethylene glycol dimethyl ether, and 5.0 mL of water was heated at reflux under nitrogen atmosphere for 3 hours. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, washed successively with water and saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. To the obtained residue were added diisopropyl ether, ethyl acetate, and hexane, and the solid matter was filtered and washed with diisopropyl ether to obtain 4.81 g of 4-cyclopropyl-1-ethyl-6-nitroquinolin-2(1H)-one as a slightly brown solid.


$^1\text{H-NMR}$ (CDCl_3) δ : 0.79-0.87 (2H, m), 1.15-1.24 (2H, m), 1.37 (3H, t, $J = 7.3$ Hz), 2.11-2.23

(1H, m), 4.37 (2H, q, J = 7.1 Hz), 6.54 (1H, s), 7.47 (1H, d, J = 9.2 Hz), 8.42 (1H, dd, J = 9.2, 2.6 Hz), 9.00 (1H, d, J = 2.6 Hz).

[0127]

Reference Example 3

5



A mixture of 4.8 g of 4-cyclopropyl-1-ethyl-6-nitroquinolin-2(1H)-one, 0.68 g of ammonium chloride, 3.91 g of iron powder, 48 mL of ethanol, and 9.6 mL of water was heated at reflux for 1 hour. The reaction mixture was cooled to room temperature, and then water and ethyl acetate were added, and the insoluble matter was filtered off. The filter residue was washed with ethyl acetate and water. The filtrate was combined with the wash solution, and the organic layer was fractionated, washed with water and saturated sodium chloride aqueous solution, then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. To the obtained residue were added diisopropyl ether and ethyl acetate, and the solid matter was filtered to obtain 3.92 g of 6-amino-4-cyclopropyl-1-ethylquinolin-2(1H)-one as a light yellow solid.


10

15

¹H-NMR (CDCl₃) δ: 0.71-0.79 (2H, m), 0.99-1.08 (2H, m), 1.33 (3H, t, J = 7.3 Hz), 1.96-2.08 (1H, m), 3.74 (2H, brs), 4.31 (2H, q, J = 7.1 Hz), 6.42 (1H, s), 6.99 (1H, dd, J = 9.2, 2.6 Hz), 7.23 (1H, d, J = 9.2 Hz), 7.38 (1H, d, J = 2.6 Hz).

[0128]

20 Reference Example 4



A mixture of 2.0 g of 4-chloro-1-ethyl-6-nitroquinolin-2(1H)-one, 10 mL of N,N-dimethylacetamide and 3.46 mL of morpholine was stirred at the ambient temperature of 130-140°C for 2 hours. The reaction mixture was cooled to room temperature and the solvent was distilled off under reduced pressure. To the obtained residue was added water, and the solid matter was filtered and washed with diisopropyl ether to obtain 2.29 g of 1-ethyl-4-(morpholin-


25

4-yl)-6-nitroquinolin-2(1H)-one as a slightly brown solid.

¹H-NMR (CDCl₃) δ: 1.37 (3H, t, J = 7.3 Hz), 3.10-3.18 (4H, m), 3.95-4.03 (4H, m), 4.36 (2H, q, J = 7.0 Hz), 6.27 (1H, s), 7.46 (1H, d, J = 9.2 Hz), 8.38 (1H, dd, J = 9.2, 2.6 Hz), 8.71 (1H, d, J = 2.6 Hz).

5 [0129]

Reference Example 5




A mixture of 2.27 g of 1-ethyl-4-(morpholin-4-yl)-6-nitroquinolin-2(1H)-one, 0.26 g of ammonium chloride, 1.46 g of iron powder, 20 mL of ethanol and 4.0 mL of water was heated at reflux for 3 hours and 20 minutes. To the reaction mixture were added 20 mL of dioxane and 20 mL of ethyl acetate, and the mixture was heated at reflux for 3 hours. The reaction mixture was cooled to room temperature, allowed to stand overnight, and then heated at reflux for 1 hour. To the reaction mixture was added 100 mL of chloroform, and the mixture was heated at reflux for 1 hour. To the reaction mixture were added 0.26 g of ammonium chloride and 0.73 g of iron powder, and the mixture was heated at reflux for 5 hours. The reaction mixture was cooled to room temperature and the insoluble matter was filtered off. The filter residue was washed with chloroform and water. The filtrate was combined with the wash solution, and then the organic layer was fractionated, washed with saturated sodium chloride aqueous solution, then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. To the obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 1.76 g of 6-amino-1-ethyl-4-(morpholin-4-yl)quinolin-2(1H)-one as slightly brown solid.

¹H-NMR (CDCl₃) δ: 1.32 (3H, t, J = 7.3 Hz), 3.04-3.12 (4H, m), 3.71 (2H, brs), 3.89-3.96 (4H, m), 4.29 (2H, q, J = 7.0 Hz), 6.18 (1H, s), 6.96 (1H, dd, J = 8.6, 2.6 Hz), 7.09 (1H, d, J = 3.3 Hz), 7.24 (1H, d, J = 9.2 Hz).

25 [0130]

Reference Example 6




To a solution of 0.7 g of 6-amino-1-ethyl-4-(morpholin-4-yl)quinolin-2(1H)-one in 14 mL of dichloromethane was added 0.46 g of 1,1'-carbonyldiimidazole, and the mixture was stirred at room temperature for 4 hours. To the reaction mixture was added 0.54 mL of triethylamine. To the reaction mixture under ice cooling was added 0.48 g of 2-amino-1-phenylethanone hydrochloride, and the mixture was stirred at room temperature for 3 hours. To the reaction mixture were added chloroform and water, and the pH of the mixture was adjusted to 2.0 with 2 mol/L hydrochloric acid. The organic layer was fractionated, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. To the obtained residue were added diisopropyl ether and ethyl acetate, and the solid matter was filtered to obtain 0.86 g of 1-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-3-(2-oxo-2-phenylethyl)urea as a slightly brown solid.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.34 (3H, t, $J = 6.9$ Hz), 3.10-3.17 (4H, m), 3.89-3.97 (4H, m), 4.32 (2H, q, $J = 7.0$ Hz), 4.86 (2H, d, $J = 4.0$ Hz), 5.93-6.01 (1H, m), 6.22 (1H, s), 6.98 (1H, s), 7.34 (1H, d, $J = 9.2$ Hz), 7.42-7.56 (3H, m), 7.59-7.68 (1H, m), 7.96-8.03 (3H, m).

[0131]

Reference Example 7




To a solution of 5 mL of N-ethylaniline in 50 mL of dichloromethane and 25 mL of methanol was added 6.7 g of sodium bicarbonate at room temperature, followed by the addition of 13.8 g of benzyltrimethylammonium dichloroiodate. After stirring the reaction mixture at room temperature for 30 minutes, ethyl acetate and water were added to the mixture. The organic layer was fractionated, washed successively with water and saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure to obtain 9.8 g of N-ethyl-4-iodoaniline as a slightly brown solid.

¹H-NMR (CDCl₃) δ: 1.24 (3H, t, J = 6.9 Hz), 3.12 (2H, q, J = 7.0 Hz), 3.60 (1H, brs), 6.34-6.42 (2H, m), 7.37-7.45 (2H, m).

[0132]

Reference Example 8




5

To a mixture of 9.8 g of N-ethyl-4-iodoaniline, 7.0 g of 3-tert-butoxy-3-oxopropionic acid and 50 mL of dichloromethane was added 8.38 g of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride under ice cooling, and the obtained mixture was stirred at room temperature for 2 hours. To the reaction mixture were added 6.1 mL of triethylamine and 5.34 g of N,N-dimethyl-4-aminopyridine, and the mixture was stirred at room temperature for 1 hour. To the reaction mixture were added 1.75 g of 3-tert-butoxy-3-oxopropionic acid and 4.2 g of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, and the mixture was stirred at room temperature for 1 hour. To the reaction mixture were added water and ethyl acetate. The organic layer was fractionated, washed twice with saturated sodium hydrogen carbonate aqueous solution and 1 mol/L hydrochloric acid, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. To the obtained residue was added diisopropyl ether and ethyl acetate, and the solid matter was filtered and washed with diisopropyl ether to obtain 8.1 g of tert-butyl 3-(ethyl(4-iodophenyl)amino)-3-oxopropanoate as a white solid.

¹H-NMR (CDCl₃) δ: 1.12 (3H, t, J = 7.3 Hz), 1.42 (9H, s), 3.07 (2H, s), 3.75 (2H, q, J = 7.0 Hz), 6.94-7.01 (2H, m), 7.71-7.79 (2H, m).

[0133]

Reference Example 9




25

To 40 mL of methanesulfonic acid was added 8.1 g of tert-butyl 3-(ethyl (4-iodophenyl)amino)-3-oxopropanoate at room temperature. To the obtained mixture was added

5.91 g of diphosphorus pentaoxide, and then the mixture was stirred at the ambient temperature of 50-60°C for 30 minutes. The reaction mixture was warmed and stirred at the ambient temperature of 100-110°C for 30 minutes. The reaction mixture was cooled to room temperature, and then the mixture was put into iced water. The solid matter was filtered and washed with water to obtain 6.55 g of 1-ethyl-4-hydroxy-6-iodoquinolin-2(1H)-one as a slightly brown solid.

¹H-NMR (DMSO-D₆) δ: 1.14 (3H, t, J = 7.3 Hz), 4.17 (2H, q, J = 7.0 Hz), 5.86 (1H, s), 7.36 (1H, d, J = 9.2 Hz), 7.88 (1H, dd, J = 8.6, 2.0 Hz), 8.14 (1H, d, J = 2.0 Hz), 11.56 (1H, brs).
[0134]

10 Reference Example 10




To 25 mL of phosphorous oxychloride was added 6.5 g of 1-ethyl-4-hydroxy-6-iodoquinolin-2(1H)-one at room temperature, and the mixture was stirred at the ambient temperature of 90-100°C for 30 minutes. The reaction mixture was cooled to room

15 temperature, and then the mixture was put into water. To the obtained mixture was added ethyl acetate, the organic layer was fractionated, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [ethyl acetate]. To the obtained residue were added diisopropyl ether, ethyl
20 acetate, and hexane, and the solid matter was filtered to obtain 4.87 g of 4-chloro-1-ethyl-6-iodoquinolin-2(1H)-one as a slightly brown solid.

¹H-NMR (DMSO-D₆) δ: 1.19 (3H, t, J = 7.3 Hz), 4.25 (2H, q, J = 7.0 Hz), 6.98 (1H, s), 7.52 (1H, d, J = 9.2 Hz), 8.01 (1H, dd, J = 9.2, 2.0 Hz), 8.19 (1H, d, J = 2.0 Hz).
[0135]

25 Reference Example 11




A mixture of 1.48 g of 4-chloro-1-ethyl-6-iodoquinolin-2(1H)-one, 0.9 g of sodium formate, 0.56 g of lithium chloride, 1.51 mL of N,N-diisopropylethylamine, 0.86 mL of

acetic anhydride, 0.12 g of tris(dibenzylideneacetone)dipalladium (0), and 10 mL of N,N-dimethylacetamide was stirred under nitrogen atmosphere at the ambient temperature of 90°C for 4 hours. The reaction mixture was cooled to room temperature and the insoluble matter was filtered off. The filter residue was washed with 5 mol/L sodium hydroxide aqueous solution and water. After combining the filtrate with the wash solution and adjusting the pH to 2.0 with 6 mol/L hydrochloric acid, diisopropyl ether and ethyl acetate were added to the mixture. The solid matter was filtered and washed with diisopropyl ether to obtain 1.06 g of 4-chloro-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylic acid as a slightly brown solid.

¹H-NMR (DMSO-D₆) δ: 1.23 (3H, t, J = 7.3 Hz), 4.31 (2H, q, J = 7.0 Hz), 7.06 (1H, s), 7.80 (1H, d, J = 9.2 Hz), 8.22 (1H, dd, J = 8.6, 2.0 Hz), 8.51 (1H, d, J = 2.0 Hz), 13.27 (1H, brs). [0136]

Reference Example 12




A mixture of 10.1 g of 4-chloro-1-ethyl-6-nitroquinolin-2(1H)-one, 8.07 g of isopropenylboronic acid pinacol ester, 17.38 g of tripotassium phosphate, 0.28 g of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 225 mL of dioxane and 90 mL of water was heated at reflux under nitrogen atmosphere for 3 hours and 10 minutes. To the reaction mixture was added 0.28 g of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), and the mixture was heated at reflux for 30 minutes. To the reaction mixture was added 0.28 g of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), and the mixture was heated at reflux for 1 hour and 30 minutes. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, and the aqueous layer was extracted twice with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. To the obtained residue were added diisopropyl ether and ethyl acetate, and the solid matter was filtered to obtain 10.19 g of 1-ethyl-6-nitro-4-(prop-1-en-2-yl)quinolin-2(1H)-one as a yellow solid.

¹H-NMR (CDCl₃) δ: 1.40 (3H, t, J = 7.2 Hz), 2.14-2.19 (3H, m), 4.40 (2H, q, J = 7.2 Hz), 5.13-5.18 (1H, m), 5.49-5.53 (1H, m), 6.66 (1H, s), 7.48 (1H, d, J = 9.3 Hz), 8.40 (1H, dd, J = 9.3, 2.7 Hz), 8.64 (1H, d, J = 2.7 Hz).

[0137]

Reference Example 13




A solution of 5.0 g of 1-ethyl-6-nitro-4-(prop-1-en-2-yl)quinolin-2(1H)-one in
 5 450 mL of methanol, and 50 mL of dioxane was added 2.5 g of 10% palladium-carbon, and the
 mixture was stirred at room temperature under hydrogen atmosphere for 2 hours. The insoluble
 matter was filtered off, and the solvent was distilled off under reduced pressure, to obtain 3.86 g
 of 6-amino-1-ethyl-4-(propan-2-yl)quinolin-2(1H)-one as a yellow foam.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.31 (6H, d, $J = 6.8$ Hz), 1.27-1.38 (3H, m), 3.23-3.35 (1H, m), 4.32 (2H, q,
 10 $J = 7.2$ Hz), 6.63 (1H, s), 6.98 (1H, dd, $J = 9.0, 2.4$ Hz), 7.10 (1H, d, $J = 2.4$ Hz), 7.23-7.30 (1H,
 m).

[0138]

Reference Example 14




To a suspension of 1.01 g of 4-chloro-1-ethyl-2-oxo-1,2-dihydroquinoline-6-
 15 carboxylic acid in 5 mL of N,N -dimethylformamide were added 0.83 g of potassium carbonate
 and 0.51 mL of benzyl chloride, and the mixture was stirred at the ambient temperature of 80°C
 for 1 hour. The reaction mixture was cooled to room temperature, and then ethyl acetate and
 water were added, and the pH of the mixture was adjusted to 2.0 with 6 mol/L hydrochloric acid.
 20 The organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The
 organic layer was combined with the extract, washed successively with water and saturated
 sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the
 solvent was distilled off under reduced pressure. The obtained residue was purified by silica
 gel column chromatography [hexane:ethyl acetate gradient elution = 80:20 to 70:30]. To the
 25 obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 1.09 g
 of benzyl 4-chloro-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylate as a white solid.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.36 (3H, t, $J = 7.2$ Hz), 4.36 (2H, q, $J = 7.2$ Hz), 5.43 (2H, s), 6.93 (1H, s),
 7.34-7.52 (6H, m), 8.30 (1H, dd, $J = 9.0, 2.0$ Hz), 8.75 (1H, d, $J = 2.0$ Hz).

[0139]

Reference Example 15




A mixture of 1.09 g of benzyl 4-chloro-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylate, 1.18 g of tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydropyridine-1(2H)-carboxylate, 1.36 g of tripotassium phosphate, 68 mg of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 23 mL of dioxane and 9 mL of water was heated at reflux under nitrogen atmosphere for 30 minutes. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, and the aqueous layer was extracted twice with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [hexane:ethyl acetate gradient elution = 70:30 to 60:40] to obtain benzyl 4-(1-(tert-butoxycarbonyl)-1,2,3,6-tetrahydropyridin-4-yl)-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylate as a yellow oil.

To a solution of the obtained benzyl 4-(1-(tert-butoxycarbonyl)-1,2,3,6-tetrahydropyridin-4-yl)-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylate in 4 mL of formic acid was added 2.5 mL of 37% formaldehyde aqueous solution, and the mixture was stirred at the ambient temperature of 80°C for 1 hour and 30 minutes. To sodium hydrogen carbonate aqueous solution were added the reaction mixture and ethyl acetate. The organic layer was fractionated, and the aqueous layer was extracted twice with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [chloroform:methanol gradient elution = 100:0 to 70:30] to obtain 1.17 g of benzyl 1-ethyl-4-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxylate as a light brown solid.

¹H-NMR (CDCl₃) δ: 1.37 (3H, t, J = 7.2 Hz), 2.45 (3H, s), 2.42-2.53 (2H, m), 2.71 (2H, t, J = 5.6 Hz), 3.13-3.21 (2H, m), 4.37 (2H, q, J = 7.2 Hz), 5.39 (2H, s), 5.80-5.87 (1H, m), 6.58 (1H, s), 7.33-7.50 (6H, m), 8.21 (1H, dd, J = 9.0, 2.0 Hz), 8.45 (1H, d, J = 2.0 Hz).

[0140]

5 Reference Example 16




To a solution of 118 mg of benzyl 1-ethyl-4-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxylate in 20 mL of methanol was added 118 mg of 10% palladium-carbon, and the mixture was stirred under hydrogen atmosphere at room temperature for 3 hours and 15 minutes. The insoluble matter was filtered off, and the solvent was distilled off under reduced pressure. To the obtained residue was added ethyl acetate, and the solid matter was filtered to obtain 62 mg of 1-ethyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxylic acid.

¹H-NMR (D₂O) δ: 1.12 (3H, t, J = 7.3 Hz), 1.70-1.85 (2H, m), 2.01-2.12 (2H, m), 2.81 (3H, s), 3.13-3.23 (2H, m), 3.24-3.34 (1H, m), 3.48-3.57 (2H, m), 4.13 (2H, q, J = 7.3 Hz), 6.41 (1H, s), 7.51 (1H, d, J = 9.0 Hz), 7.98 (1H, d, J = 9.0 Hz), 8.25 (1H, s).

[0141]

Reference Example 17




A mixture of 0.39 g of 4-chloro-1-ethyl-6-iodoquinolin-2(1H)-one, 2.0 mL of N,N-dimethylacetamide and 0.51 mL of morpholine was stirred in a sealed tube at the ambient temperature of 130-140°C for 1 hour and 30 minutes. The reaction mixture was cooled to room temperature and then the solvent was distilled off under reduced pressure. To the obtained residue were added ethyl acetate and water, and the pH of the mixture was adjusted to 2.0 with 2 mol/L hydrochloric acid. The organic layer was fractionated, washed with saturated sodium

chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. To the obtained residue were added diisopropyl ether and hexane, and the solid matter was filtered to obtain 0.33 g of 1-ethyl-6-iodo-4-(morpholin-4-yl)quinolin-2(1H)-one as slightly brown solid.

- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.32 (3H, t, $J = 7.3$ Hz), 3.04-3.13 (4H, m), 3.90-3.99 (4H, m), 4.28 (2H, q, $J = 7.3$ Hz), 6.18 (1H, s), 7.14 (1H, d, $J = 9.2$ Hz), 7.79 (1H, dd, $J = 8.9, 2.3$ Hz), 8.10 (1H, d, $J = 8.9$ Hz).

[0142]

Reference Example 18




10

- A mixture of 0.25 g of 1-ethyl-6-iodo-4-(morpholin-4-yl)quinolin-2(1H)-one, 53 mg of 1,1'-bis(diphenyl phosphino)ferrocene palladium(II) dichloride, 0.19 g of potassium acetate, 0.17 g of bis(pinacolato)diboron, and 3.0 mL of dioxane was heated at reflux under nitrogen atmosphere for 4 hours. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [hexane:ethyl acetate gradient elution = 80:20 to 0:100] to obtain 66 mg of 1-ethyl-4-(morpholin-4-yl)-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinolin-2(1H)-one as a slightly brown oil.
- 15
- 20

$^1\text{H-NMR}$ (CDCl_3) δ : 1.20-1.45 (15H, m), 3.05-3.22 (4H, m), 3.90-4.05 (4H, m), 4.25-4.41 (2H, m), 6.14-6.22 (1H, m), 7.33-7.43 (1H, m,), 7.89-8.00 (1H, m), 8.23-8.33 (1H, m).

[0143]

Reference Example 19




25

To a solution of 1.0 g of 6-amino-1-ethyl-4-(morpholin-4-yl)quinolin-2(1H)-one in 10 mL of dichloromethane was added 0.65 g of 1,1'-carbonyldiimidazole, and the mixture was stirred at the ambient temperature of 40-50°C for 2 hours. To the reaction mixture under ice cooling were added 0.76 mL of triethylamine and 0.84 g of 2-amino-1-(4-chlorophenyl)ethanone hydrochloride, and the mixture was stirred at room temperature for 2 hours. To the reaction mixture were added chloroform and water, and the pH of the mixture was adjusted to 2.0 with 2 mol/L hydrochloric acid. The organic layer was fractionated, washed with water and saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [chloroform:methanol gradient elution = 100:0 to 90:10]. To the obtained residue were added diisopropyl ether and ethyl acetate, and the solid matter was filtered to obtain 0.74 g of 1-(2-(4-chlorophenyl)-2-oxoethyl)-3-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)urea as a slightly brown solid.

¹H-NMR (DMSO-D₆) δ: 1.17 (3H, t, J = 7.0 Hz), 3.01-3.08 (4H, m), 3.79-3.86 (4H, m), 4.20 (2H, q, J = 7.1 Hz), 4.70 (2H, d, J = 5.1 Hz), 5.99 (1H, s), 6.50 (1H, t, J = 5.2 Hz), 7.47 (1H, d, J = 9.3 Hz), 7.56 (1H, dd, J = 9.1, 2.6 Hz), 7.61-7.67 (2H, m), 8.01-8.07 (3H, m), 9.07 (1H, s).

[0144]


Reference Example 20



A mixture of 11.14 g of 4-chloro-1-ethyl-6-iodoquinolin-2(1H)-one, 7.93 g of tert-butyl 1-phenylhydrazinecarboxylate, 581 mg of tri-tert-butyl phosphonium tetrafluoroborate, 375 mg of palladium acetate, 16.32 g of cesium carbonate, and 160 mL of toluene was heated at reflux under nitrogen atmosphere for 1 hour and 30 minutes. The reaction mixture was cooled to room temperature, and then water was added. The organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [hexane:ethyl acetate gradient elution = 80:20 to 60:40] to obtain 6.01 g of tert-butyl 2-(4-chloro-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-1-phenylhydrazinecarboxylate as a yellow solid.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.33 (3H, t, $J = 7.2$ Hz), 1.43 (9H, s), 4.31 (2H, q, $J = 7.1$ Hz), 6.58 (1H, s), 6.87 (1H, s), 7.13-7.20 (2H, m), 7.29-7.39 (3H, m), 7.45 (1H, d, $J = 2.7$ Hz), 7.54-7.60 (2H, m).
[0145]


Reference Example 21



A mixture of 690 mg of tert-butyl 2-(4-chloro-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-1-phenylhydrazinecarboxylate, 172 mg of cyclopropylboric acid, 1.06 g of tripotassium phosphate, 35 mg of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 12 mL of dioxane and 4.0 mL of water was stirred at 140°C for 5 minutes using microwave
10 equipment. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica
15 gel column chromatography [hexane:ethyl acetate gradient elution = 60:40 to 40:60] to obtain 527 mg of tert-butyl 2-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-1-phenylhydrazinecarboxylate as a light yellow solid.

$^1\text{H-NMR}$ (CDCl_3) δ : 0.67-0.73 (2H, m), 0.94-1.01 (2H, m), 1.32 (3H, t, $J = 7.1$ Hz), 1.41 (9H, s), 1.90-2.00 (1H, m), 4.31 (2H, q, $J = 7.2$ Hz), 6.44 (1H, d, $J = 0.98$ Hz), 6.56 (1H, s), 7.08-7.19
20 (2H, m), 7.25-7.39 (3H, m), 7.52 (1H, d, $J = 2.4$ Hz), 7.57-7.63 (2H, m).
[0146]

Reference Example 22




A mixture of 200 mg of tert-butyl 2-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-1-phenylhydrazinecarboxylate and 2 mL of 20% sodium ethoxide-ethanol
25 solution was stirred at 140°C for 2 minutes using microwave equipment. The reaction mixture

was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, washed successively with water and saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. To the obtained residue was added diisopropyl ether, and the solid matter was filtered and washed with diisopropyl ether to obtain 115 mg of 4-cyclopropyl-1-ethyl-6-(2-phenylhydrazinyl)quinolin-2(1H)-one as a light yellow solid.

$^1\text{H-NMR}$ (DMSO-D_6) δ : 0.62-0.69 (2H, m), 0.87-0.95 (2H, m), 1.15 (3H, t, $J = 7.0$ Hz), 1.90-2.00 (1H, m), 4.18 (2H, q, $J = 7.0$ Hz), 6.20 (1H, s), 6.65 (1H, t, $J = 7.2$ Hz), 6.74-6.81 (2H, m), 7.08-7.15 (3H, m), 7.41 (1H, d, $J = 9.3$ Hz), 7.45 (1H, d, $J = 2.4$ Hz), 7.69-7.77 (2H, m).

[0147]

Reference Example 23




To a suspension of 4.0 g of 4-chloro-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylic acid in 25 mL of tetrahydrofuran at room temperature were added 2.6 mL of oxalyl chloride and 10 μL of N,N-dimethylformamide, and the mixture was stirred for 3 hours. To a 28% ammonium aqueous solution under ice cooling were added the reaction mixture and 100 mL of water. The solid matter was filtered and washed with water to obtain 3.70 g of 4-chloro-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxamide as a slightly brown solid.

$^1\text{H-NMR}$ (DMSO-D_6) δ : 1.22 (3H, t, $J = 7.1$ Hz), 4.31 (2H, q, $J = 7.2$ Hz), 7.02 (1H, s), 7.52

(1H, brs), 7.76 (1H, d, $J = 9.0$ Hz), 8.22 (1H, dd, $J = 8.9, 2.1$ Hz), 8.25 (1H, brs), 8.50 (1H, d, $J = 2.2$ Hz).

[0148]

Reference Example 24




A mixture of 4-chloro-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxamide and 10 mL of 1,1-dimethoxy-N,N-dimethylethaneamine was heated at reflux for 4 hours. After cooling the reaction mixture to room temperature, diisopropyl ether was added to the reaction mixture, the solid matter was filtered and washed with diisopropyl ether to obtain 2.13 g of 4-

chloro-N-((1E)-1-(dimethylamino)ethylidene)-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxamide as a slightly brown solid.

¹H-NMR (DMSO-D₆) δ: 1.22 (3H, t, J = 7.1 Hz), 2.31 (3H, s), 3.16 (3H, s), 3.19 (3H, s), 4.30 (2H, q, J = 7.1 Hz), 7.00 (1H, s), 7.72 (1H, d, J = 8.8 Hz), 8.33 (1H, dd, J = 8.8, 2.0 Hz), 8.64 (1H, d, J = 2.0 Hz).

[0149]

Reference Example 25




To a solution of 0.50 g of 6-amino-4-cyclopropyl-1-ethylquinolin-2(1H)-one in 9 mL of acetonitrile was added 1.25 g of p-toluenesulfonic acid monohydrate followed by the addition of 1.3 mL of aqueous solution of 0.91 g of potassium iodide and 0.30 g of sodium nitrite under ice cooling. The mixture was stirred for 10 minutes, and then stirred at room temperature for 1 hour. To the reaction mixture were added water, saturated sodium carbonate aqueous solution, 1% sodium thiosulfate aqueous solution, and ethyl acetate. The organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [hexane:ethyl acetate gradient elution = 70:30 to 40:60]. To the obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 0.31 g of 4-cyclopropyl-1-ethyl-6-iodoquinolin-2(1H)-one as a light yellow solid.

¹H-NMR (CDCl₃) δ: 0.75-0.79 (2H, m), 1.07-1.12 (2H, m), 1.32 (3H, t, J = 7.2 Hz), 2.01-2.08 (1H, m), 4.30 (2H, q, J = 7.2 Hz), 6.43 (1H, d, J = 1.2 Hz), 7.14 (1H, d, J = 8.8 Hz), 7.81 (1H, dd, J = 8.8, 2.0 Hz), 8.38 (1H, d, J = 2.0 Hz).

[0150]


Reference Example 26



By the same method as Reference Example 18, from 4-chloro-1-ethyl-6-iodoquinolin-2(1H)-one was obtained 4-chloro-1-ethyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinolin-2(1H)-one.

¹H-NMR (CDCl₃) δ: 1.35 (3H, t, J = 7.2 Hz), 1.38 (12H, s), 4.36 (2H, q, J = 7.2 Hz), 6.88 (1H, s), 7.39 (1H, d, J = 8.5 Hz), 8.03 (1H, dd, J = 8.5, 1.5 Hz), 8.46 (1H, d, J = 1.5 Hz).
[0151]


Reference Example 27



A mixture of 334 mg of 4-chloro-1-ethyl-6-iodoquinolin-2(1H)-one, 112 mg of ethynylbenzene, 2 mg of copper iodide(I), 5 mg of triphenylphosphine, 2 mg of dichloropalladium, 0.42 mL of triethylamine, and 5 mL of tetrahydrofuran was stirred under nitrogen atmosphere at room temperature for 2 hours. To the reaction mixture were added ethyl acetate, water and 1 mol/L hydrochloric acid. The organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. To the obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 160 mg of 4-chloro-1-ethyl-6-(phenylethynyl)quinolin-2(1H)-one as a brown solid.

¹H-NMR (CDCl₃) δ: 1.37 (3H, t, J = 7.2 Hz), 4.36 (2H, q, J = 7.2 Hz), 6.92 (1H, s), 7.35-7.42 (4H, m), 7.55-7.60 (2H, m), 7.76 (1H, dd, J = 8.8, 2.0 Hz), 8.20 (1H, d, J = 2.0 Hz).
[0152]

Reference Example 28




A mixture of 154 mg of 4-chloro-1-ethyl-6-(phenylethynyl)quinolin-2(1H)-one, 52 mg of cyclopropylboric acid, 318 mg of tripotassium phosphate, 11 mg of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 3 mL of dioxane and 1 mL of water was

stirred at 140°C for 15 minutes using microwave equipment. The reaction mixture was cooled to room temperature, and then ethyl acetate, water and 2 mol/L hydrochloric acid were added. The organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel chromatography [hexane:ethyl acetate gradient elution = 80:20 to 60:40]. To the obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 100 mg of 4-cyclopropyl-1-ethyl-6-(phenylethynyl)quinolin-2(1H)-one as a white solid.

¹H-NMR (CDCl₃) δ: 0.77-0.83 (2H, m), 1.08-1.16 (2H, m), 1.36 (3H, t, J = 7.1 Hz), 2.08-2.19 (1H, m), 4.35 (2H, q, J = 7.2 Hz), 6.45-6.49 (1H, m), 7.34-7.42 (4H, m), 7.55-7.61 (2H, m), 7.71 (1H, dd, J = 8.9, 1.8 Hz), 8.27 (1H, d, J = 2.0 Hz).

[0153]

Reference Example 29



15

A mixture of 31 mg of 4-cyclopropyl-1-ethyl-6-(phenylethynyl)quinolin-2(1H)-one, 24 mg of magnesium sulfate, 19 mg of potassium permanganate, 5 mg of sodium hydrogen carbonate, 3 mL of acetone, and 1.7 mL of water was stirred at room temperature for 20 minutes. To the reaction mixture was added 43 mg of potassium permanganate, and the mixture was stirred at room temperature for 1 hour. To the reaction mixture were added 30 mg of sodium nitrite and 10% sulfuric acid aqueous solution, and then the insoluble matter was filtered off. The filter residue was washed with ethyl acetate and water. The filtrate was combined with the wash solution, and the organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel chromatography [hexane:ethyl acetate gradient elution = 60:40 to 30:70] to obtain 20 mg of 1-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-2-phenylethane-1,2-dione.


¹H-NMR (CDCl₃) δ: 0.74-0.80 (2H, m), 1.05-1.12 (2H, m), 1.35 (3H, t, J = 7.1 Hz), 2.07-2.16 (1H, m), 4.36 (2H, q, J = 7.2 Hz), 6.46-6.49 (1H, m), 7.46 (1H, d, J = 9.0 Hz), 7.54 (2H, t, J = 7.9

30

Hz), 7.69 (1H, t, J = 7.4 Hz), 8.02 (2H, dd, J = 8.3, 1.2 Hz), 8.14 (1H, dd, J = 8.9, 2.1 Hz), 8.77 (1H, d, J = 2.0 Hz).

[0154]

Reference Example 30




5

- A mixture of 503 mg of 4-chloro-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylic acid, 639 mg of 8-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-(1,4-dioxaspiro[4.5]dec-7-ene, 849 mg of tripotassium phosphate, 42 mg of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 12 mL of dioxane and 4 mL of water
- 10 was stirred at 140°C for 5 minutes using microwave equipment. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added to the reaction mixture. The aqueous layer was fractionated and washed twice with ethyl acetate. To the aqueous layer was added 6 mol/L hydrochloric acid to adjust the pH to 2.0, and then added tetrahydrofuran. The organic layer was fractionated, and the aqueous layer was extracted twice with ethyl acetate.
- 15 The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. To the obtained residue was added ethyl acetate, and the solid matter was filtered to obtain 590 mg of 4-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylic acid as a light brown solid.
- 20 ¹H-NMR (DMSO-D₆) δ: 1.21 (3H, t, J = 7.0 Hz), 1.85 (2H, t, J = 6.2 Hz), 2.39-2.56 (4H, m), 3.94-4.00 (4H, m), 4.29 (2H, q, J = 7.1 Hz), 5.71-5.76 (1H, m), 6.41 (1H, s), 7.69 (1H, d, J = 9.0 Hz), 8.11 (1H, dd, J = 8.9, 2.1 Hz), 8.26 (1H, d, J = 2.2 Hz), 13.02 (1H, brs).

[0155]

Reference Example 31




To a mixture of 583 mg of 4-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylic acid, 20 mL of dioxane and 100 mL of methanol was added 291 mg of 20% hydroxide palladium-carbon, and the mixture was stirred under hydrogen

- 5 atmosphere at room temperature for 30 minutes. The insoluble matter was filtered off, and the solvent was distilled off under reduced pressure. To the obtained residue was added diisopropyl ether, the solid matter was filtered to obtain 502 mg of 4-(1,4-dioxaspiro[4.5]dec-8-yl)-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylic acid as a white solid.

- ¹H-NMR (DMSO-D₆) δ: 1.22 (3H, t, J = 7.0 Hz), 1.60-1.93 (8H, m), 3.10-3.25 (1H, m), 3.87-3.96 (4H, m), 4.29 (2H, q, J = 6.9 Hz), 6.46-6.51 (1H, m), 7.66-7.73 (1H, m), 8.13 (1H, dd, J = 8.7 Hz, 1.6 Hz), 8.45 (1H, d, J = 1.5 Hz), 13.06 (1H, brs).

[0156]

Reference Example 32




- 15 To a suspension of 250 mg of 4-(1,4-dioxaspiro[4.5]dec-8-yl)-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylic acid in 10 mL of tetrahydrofuran at room temperature was added 1.0 mL of concentrated hydrochloric acid, and the mixture was stirred for 4 hours. The solid matter was filtered and washed with ethyl acetate to obtain 182 mg of 1-ethyl-2-oxo-4-(4-oxocyclohexyl)-1,2-dihydroquinoline-6-carboxylic acid as a white solid.

- 20 ¹H-NMR (DMSO-D₆) δ: 1.22 (3H, t, J = 7.1 Hz), 1.83-2.00 (2H, m), 2.09-2.19 (2H, m), 2.26-2.37 (2H, m), 2.71-2.83 (2H, m), 3.67-3.77 (1H, m), 4.30 (2H, q, J = 7.1 Hz), 6.58-6.62 (1H, m), 7.73 (1H, d, J = 9.0 Hz), 8.16 (1H, dd, J = 8.9 Hz, 1.8 Hz), 8.57 (1H, d, J = 2.0 Hz).

[0157]

Example 1




To a suspension of 1.2 g of 6-amino-4-cyclopropyl-1-ethylquinolin-2(1H)-one in 6 mL of pyridine was added 0.73 mL of benzoyl chloride, and the mixture was stirred at room temperature for 1 hour. To the reaction mixture were added ethyl acetate and water, and the pH of the mixture was adjusted to 2.0 with 2 mol/L hydrochloric acid. The organic layer was fractionated, washed with water and saturated sodium chloride aqueous solution, then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. To the obtained residue were added diisopropyl ether and ethyl acetate, and the solid matter was filtered to obtain 1.75 g of N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)benzamide as a slightly brown solid.

¹H-NMR (DMSO-D₆) δ: 0.75-0.84 (2H, m), 1.02-1.11 (2H, m), 1.20 (3H, t, J = 6.9 Hz), 2.09-2.21 (1H, m), 4.26 (2H, q, J = 7.1 Hz), 6.32 (1H, s), 7.44-7.66 (4H, m), 7.91-8.09 (3H, m), 8.65 (1H, d, J = 2.0 Hz), 10.45 (1H, s).

[0158]

Example 2




To a solution of 1.75 g of N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)benzamide in 15 mL of N,N-dimethylacetamide, 0.25 g of 60% sodium hydride was added under ice cooling and the mixture was stirred for 10 minutes. To the reaction mixture under ice cooling was added 0.43 mL of methyl iodide, the mixture was stirred at room temperature for 1 hour. To the reaction mixture were added ethyl acetate and water, and the pH of the mixture was adjusted to 2.0 with 6 mol/L hydrochloric acid, and then the solid matter was filtered. The organic layer of the filtrate was fractionated, washed with water and saturated sodium chloride aqueous solution, then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. To the obtained residue and filtered solid matter were added diisopropyl ether, ethyl acetate and water, and then the solid matter was filtered, washed with

water and diisopropyl ether to obtain 1.45 g of N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide as a slightly brown solid.

¹H-NMR (DMSO-D₆) δ: 0.48-0.57 (2H, m), 0.86-0.95 (2H, m), 1.13 (3H, t, J = 6.9 Hz), 1.93-2.06 (1H, m), 3.44 (3H, s), 4.19 (2H, q, J = 6.8 Hz), 6.24 (1H, s), 7.16-7.33 (5H, m), 7.46-7.58 (2H, m), 7.76-7.81 (1H, m).

[0159]

Example 3




To a suspension of 0.86 g of 1-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-3-(2-oxo-2-phenylethyl)urea in 5.0 mL of dioxane at room temperature was added 5.0 mL of concentrated hydrochloric acid and the mixture was stirred for 2 hours. To the reaction mixture was added water, and the solid matter was filtered and washed with water and diisopropyl ether to obtain 0.72 g of 1-ethyl-4-(morpholin-4-yl)-6-(2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one as a slightly brown solid.

¹H-NMR (DMSO-D₆) δ: 1.19 (3H, t, J = 7.3 Hz), 2.54-2.65 (4H, m), 3.44-3.53 (4H, m), 4.17-4.29 (2H, m), 5.98 (1H, s), 6.87 (1H, d, J = 2.6 Hz), 7.05-7.12 (2H, m), 7.14-7.31 (4H, m), 7.59-7.68 (2H, m), 10.57-10.63 (1H, m).

[0160]

Example 4



20

To a suspension of 3.0 g of 4-chloro-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylic acid in 30 mL of tetrahydrofuran at room temperature were added 1.25 mL of oxalyl chloride and 10 μL of N,N-dimethylformamide, and the mixture was stirred for 2 hours. To the reaction mixture under ice cooling were added 1.55 mL of N-methylaniline and 1.81 mL of triethylamine, and then the mixture was stirred at room temperature for 1 hour. To the reaction mixture were added ethyl acetate and water. The organic layer was fractionated, washed with


25

water and saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [hexane:ethyl acetate gradient elution = 50:50 to 20:80]. To the obtained residue were added diisopropyl ether and ethyl acetate, and then the solid matter was filtered to obtain 2.01 g of 4-chloro-1-ethyl-N-methyl-2-oxo-N-phenyl-1,2-dihydroquinoline-6-carboxamide as a slightly brown solid.

¹H-NMR (CDCl₃) δ: 1.28 (3H, t, J = 6.9 Hz), 3.55 (3H, s), 4.26 (2H, q, J = 7.3 Hz), 6.81 (1H, s), 7.07-7.13 (2H, m), 7.15-7.32 (4H, m), 7.65 (1H, dd, J = 9.2, 2.0 Hz), 7.94 (1H, d, J = 2.0 Hz).

[0161]

10 Example 5




A mixture of 2.0 g of 4-chloro-1-ethyl-N-methyl-2-oxo-N-phenyl-1,2-dihydroquinoline-6-carboxamide, 2.18 g of tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydropyridine-1(2H)-carboxylate, 1.24 g of sodium carbonate, 0.21 g of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 20 mL of ethylene glycol dimethyl ether, and 4.0 mL of water was heated at reflux under nitrogen atmosphere for 3 hours. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, washed successively with water and saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel chromatography [hexane:ethyl acetate gradient elution = 50:50 to 0:100] to obtain 2.9 g of tert-butyl 4-(1-ethyl-6-(methyl(phenyl)carbamoyl)-2-oxo-1,2-dihydroquinolin-4-yl)-3,6-dihydropyridine-1(2H)-carboxylate as a slightly brown foam.

¹H-NMR (CDCl₃) δ: 1.32 (3H, t, J = 6.9 Hz), 1.56 (9H, s), 1.94-2.14 (2H, m), 3.53 (3H, s), 3.57 (2H, t, J = 5.6 Hz), 3.99-4.06 (2H, m), 4.30 (2H, q, J = 7.0 Hz), 5.42 (1H, brs), 6.41 (1H, s), 6.99-7.07 (2H, m), 7.10-7.19 (1H, m), 7.22-7.33 (3H, m), 7.35-7.41 (1H, m), 7.77 (1H, dd, J = 9.2, 2.0 Hz).

[0162]

Example 6




A mixture of 1.47 g of tert-butyl 4-(1-ethyl-6-(methyl(phenyl)carbamoyl)-2-oxo-1,2-dihydroquinolin-4-yl)-3,6-dihydropyridine-1(2H)-carboxylate, 0.44 g of 5% palladium-carbon, 0.29 g of ammonium formate, and 15 mL of methanol was heated at reflux under nitrogen atmosphere for 1 hour and 30 minutes. The reaction mixture was cooled to room temperature and the insoluble matter was filtered off. The filter residue was washed with ethyl acetate. The filtrate was combined with the wash solution, and the solvent was distilled off under reduced pressure. To the obtained residue were added 0.44 g of 5% palladium-carbon, 0.29 g of ammonium formate, and 15 mL of methanol, and the mixture was heated at reflux under nitrogen atmosphere for 2 hours. The reaction mixture was cooled to room temperature and the insoluble matter was filtered off. The filter residue was washed with ethyl acetate. The filtrate was combined with the wash solution, and the solvent was distilled off under reduced pressure. To the obtained residue was added diisopropyl ether, and the solid matter was filtered and washed with diisopropyl ether to obtain 1.26 g of tert-butyl 4-(1-ethyl-6-(methyl(phenyl)carbamoyl)-2-oxo-1,2-dihydroquinolin-4-yl)piperidine-1-carboxylate as a white solid.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.31 (3H, t, $J = 7.3$ Hz), 1.40-1.58 (4H, m), 1.50 (9H, s), 2.62-2.85 (3H, m), 3.56 (3H, s), 4.15-4.34 (4H, m), 6.49 (1H, s), 7.04-7.11 (2H, m), 7.12-7.20 (1H, m), 7.23-7.33 (3H, m), 7.61 (1H, d, $J = 2.0$ Hz), 7.73 (1H, dd, $J = 8.6, 2.0$ Hz).

[0163]


Example 7



A mixture of 1.26 g of tert-butyl 4-(1-ethyl-6-(methyl(phenyl)carbamoyl)-2-oxo-1,2-dihydroquinolin-4-yl)-piperidine-1-carboxylate in 5 mL of dichloromethane and 5 mL of trifluoroacetic acid was stirred at room temperature for 30 minutes, and then the solvent was
 5 distilled off under reduced pressure. To the obtained residue were added ethyl acetate and saturated sodium hydrogen carbonate aqueous solution, and the organic layer was fractionated. The aqueous layer was extracted with ethyl acetate, and further extracted twice with chloroform. The organic layer was combined with the extract, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure to obtain 1.0 g of 1-ethyl-N-
 10 methyl-2-oxo-N-phenyl-4-(piperidin-4-yl)-1,2-dihydroquinoline-6-carboxamide as a white solid.
¹H-NMR (CDCl₃) δ: 1.31 (3H, t, J = 7.3 Hz), 1.55-1.67 (4H, m), 2.67-2.90 (3H, m), 3.22-3.34 (2H, m), 3.56 (3H, s), 4.29 (2H, q, J = 7.3 Hz), 6.55 (1H, s), 7.05-7.20 (3H, m), 7.22-7.32 (3H, m), 7.62 (1H, d, J = 2.0 Hz), 7.71 (1H, dd, J = 8.6, 2.0 Hz).

[0164]

15 Example 8



To a suspension of 1.0 g of 1-ethyl-N-methyl-2-oxo-N-phenyl-4-(piperidin-4-yl)-1,2-dihydroquinoline-6-carboxamide in 15 mL of acetone and 8 mL of tetrahydrofuran were added 0.71 g of potassium carbonate and 0.25 mL of ethyl iodide, and the mixture was stirred at
 20 room temperature for 30 minutes, and then stirred at the ambient temperature of 40°C for 2 hours. After adding 0.24 g of potassium carbonate and 82 μL of ethyl iodide to the reaction mixture and stirring at the ambient temperature of 40°C for 1 hour, the solvent was distilled off


under reduced pressure. To the obtained residue were added ethyl acetate and water, the organic layer was fractionated, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by basic silica gel column chromatography

5 [hexane:ethyl acetate gradient elution = 50:50 to 0:100]. To the obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 0.76 g of 1-ethyl-4-(1-ethylpiperidin-4-yl)-N-methyl-2-oxo-N-phenyl-1,2-dihydroquinoline-6-carboxamide as a white solid.

¹H-NMR (DMSO-D₆) δ: 1.05 (3H, t, J = 7.3 Hz), 1.15 (3H, t, J = 7.3 Hz), 1.31-1.52 (4H, m),
 10 1.92-2.06 (2H, m), 2.40 (2H, q, J = 7.0 Hz), 2.47-2.63 (1H, m), 2.86-2.98 (2H, m), 3.43 (3H, s),
 4.20 (2H, q, J = 7.0 Hz), 6.36 (1H, s), 7.10-7.32 (5H, m), 7.50-7.58 (2H, m), 7.73 (1H, dd, J =
 8.9, 1.7 Hz).

[0165]

Example 9




By the same method as Example 4, from 4-chloro-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylic acid was obtained 4-chloro-6-(3,4-dihydroquinolin-1(2H)-ylcarbonyl)-1-ethylquinolin-2(1H)-one.

¹H-NMR (CDCl₃) δ: 1.31 (3H, t, J = 7.2 Hz), 2.10 (2H, quint, J = 6.6 Hz), 2.88 (2H, t, J = 6.6
 20 Hz), 3.96 (2H, t, J = 6.7 Hz), 4.30 (2H, q, J = 7.2 Hz), 6.65 (1H, d, J = 7.8 Hz), 6.83-6.90 (2H,
 m), 7.00-7.06 (1H, m), 7.18-7.28 (2H, m), 7.56 (1H, dd, J = 8.8, 2.0 Hz), 8.09 (1H, d, J = 2.2
 Hz).

[0166]

Example 10




A mixture of 64 mg of 4-chloro-6-(3,4-dihydroquinolin-1(2H)-ylcarbonyl)-1-

ethylquinolin-2(1H)-one, 47 mg of 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2,3,6-tetrahydropyridine, 55 mg of sodium carbonate, 12.3 mg of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 2 mL of ethylene glycol dimethyl ether, and 0.2 mL of water was heated at reflux under nitrogen atmosphere for 2 hours. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, washed successively with water and saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by basic silica gel chromatography [hexane:ethyl acetate gradient elution = 50:50 to 0:100] to obtain 62 mg of 6-(3,4-dihydroquinolin-1(2H)-ylcarbonyl)-1-ethyl-4-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)quinolin-2(1H)-one as a slightly brown foam.

¹H-NMR (CDCl₃) δ: 1.35 (3H, t, J = 6.9 Hz), 2.05-2.18 (4H, m), 2.41 (3H, s), 2.45-2.55 (2H, m), 2.87 (2H, t, J = 6.6 Hz), 2.94-3.02 (2H, m), 3.95 (2H, t, J = 6.6 Hz), 4.33 (2H, q, J = 7.1 Hz), 5.29-5.36 (1H, m), 6.47 (1H, s), 6.54 (1H, d, J = 7.9 Hz), 6.80-6.89 (1H, m), 6.94-7.02 (1H, m), 7.18 (1H, d, J = 7.3 Hz), 7.35 (1H, d, J = 9.2 Hz), 7.49 (1H, d, J = 2.0 Hz), 7.77 (1H, dd, J = 8.6, 2.0 Hz).

[0167]

Example 11




A mixture of 60 mg of 6-(3,4-dihydroquinolin-1(2H)-ylcarbonyl)-1-ethyl-4-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)quinolin-2(1H)-one in 15 mL of methanol and 15 mL of ethyl acetate was subjected to hydrogenation reaction (25°C, 1 bar, flow rate 2 mL/min, 10% palladium-carbon) using the flow type hydrogenation reaction apparatus. The solvent was distilled off under reduced pressure, and then the obtained residue was purified by basic silica gel column chromatography [hexane:ethyl acetate gradient elution = 50:50 to 0:100]. To the obtained residue were added diisopropyl ether and ethyl acetate, and the solid matter was filtered to obtain 10 mg of 6-(3,4-dihydroquinolin-1(2H)-ylcarbonyl)-1-ethyl-4-(1-methylpiperidin-4-yl)quinolin-2(1H)-one as a white solid.

¹H-NMR (DMSO-D₆) δ: 1.18 (3H, t, J = 6.9 Hz), 1.24-1.52 (4H, m), 1.80-1.94 (2H, m), 1.95-

2.10 (2H, m), 2.19 (3H, s), 2.50-2.65 (1H, m), 2.70-2.81 (2H, m), 2.87 (2H, t, $J = 6.3$ Hz), 3.81 (2H, t, $J = 6.6$ Hz), 4.24 (2H, q, $J = 6.8$ Hz), 6.38 (1H, s), 6.58 (1H, d, $J = 7.9$ Hz), 6.84 (1H, t, $J = 7.3$ Hz), 6.96 (1H, t, $J = 7.3$ Hz), 7.22 (1H, d, $J = 7.3$ Hz), 7.50-7.55 (1H, m), 7.62 (1H, d, $J = 9.2$ Hz), 7.73-7.81 (1H, m).

5 [0168]

Example 12




By the same method as Example 4, from 4-chloro-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxylic acid was obtained 4-chloro-1-ethyl-N-methyl-N-(4-methylphenyl)-2-oxo-1,2-dihydroquinoline-6-carboxamide.

10

$^1\text{H-NMR}$ (CDCl_3) δ : 1.29 (3H, t, $J = 7.2$ Hz), 2.29 (3H, s), 3.51 (3H, s), 4.26 (2H, q, $J = 7.1$ Hz), 6.81 (1H, s), 6.98 (2H, d, $J = 8.3$ Hz), 7.07 (2H, d, $J = 8.3$ Hz), 7.20 (1H, d, $J = 8.8$ Hz), 7.65 (1H, dd, $J = 8.8, 2.0$ Hz), 7.94 (1H, d, $J = 2.0$ Hz).

[0169]

15 Example 13




By the same method as Example 10, from 4-chloro-1-ethyl-N-methyl-N-(4-methylphenyl)-2-oxo-1,2-dihydroquinoline-6-carboxamide was obtained 1-ethyl-N-methyl-N-(4-methylphenyl)-4-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide.

20

$^1\text{H-NMR}$ (CDCl_3) δ : 1.31 (3H, t, $J = 7.2$ Hz), 2.14-2.21 (2H, m), 2.26 (3H, s), 2.45 (3H, s), 2.61 (2H, t, $J = 5.6$ Hz), 3.04-3.11 (2H, m), 3.50 (3H, s), 4.29 (2H, q, $J = 7.1$ Hz), 5.36-5.42 (1H, m), 6.45 (1H, s), 6.92 (2H, d, $J = 8.3$ Hz), 7.03 (2H, d, $J = 8.5$ Hz), 7.27 (1H, d, $J = 8.8$ Hz), 7.51 (1H, d, $J = 2.0$ Hz), 7.69 (1H, dd, $J = 8.9, 2.1$ Hz).

25 [0170]

Example 14




By the same method as Reference Example 13, from 1-ethyl-N-methyl-N-(4-methylphenyl)-4-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide was obtained 1-ethyl-N-methyl-N-(4-methylphenyl)-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide.

¹H-NMR (CDCl₃) δ: 1.30 (3H, t, J = 7.1 Hz), 1.50-1.71 (4H, m), 1.99-2.10 (2H, m), 2.26 (3H, s), 2.36 (3H, s), 2.48-2.59 (1H, m), 2.91-3.00 (2H, m), 3.53 (3H, s), 4.28 (2H, q, J = 7.2 Hz), 6.55 (1H, s), 6.96 (2H, d, J = 8.3 Hz), 7.06 (2H, d, J = 8.0 Hz), 7.25-7.30 (1H, m), 7.60 (1H, d, J = 2.0 Hz), 7.71 (1H, dd, J = 8.8, 2.0 Hz).

[0171]

Example 15




By the same method as Example 1, from 6-amino-1-ethyl-4-(propan-2-yl)quinolin-2(1H)-one and 4-chlorobenzoyl chloride was obtained 4-chloro-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)benzamide.

¹H-NMR (CDCl₃) δ: 1.34 (6H, d, J = 6.6 Hz), 1.31-1.40 (3H, m), 3.34-3.47 (1H, m), 4.37 (2H, q, J = 7.1 Hz), 6.67 (1H, s), 7.42 (1H, d, J = 9.2 Hz), 7.48 (2H, d, J = 8.3 Hz), 7.77 (1H, dd, J = 9.2, 2.3 Hz), 7.91 (2H, d, J = 8.3 Hz), 8.23 (1H, s), 8.33 (1H, d, J = 2.3 Hz).

[0172]

Example 16




By the same method as Example 2, from 4-chloro-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)benzamide was obtained 4-chloro-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide.

- 5 ¹H-NMR (CDCl₃) δ: 1.12 (6H, d, J = 6.8 Hz), 1.34 (3H, t, J = 7.2 Hz), 3.01-3.14 (1H, m), 3.53 (3H, s), 4.31 (2H, q, J = 7.2 Hz), 6.61 (1H, s), 7.12-7.19 (2H, m), 7.24-7.29 (2H, m), 7.31-7.37 (3H, m).

[0173]

Example 17



10


- A mixture of 80 mg of 1-ethyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxylic acid, 80 μL of N-methyl-2,3-dihydro-1H-inden-5-amine, 3 mL of dichloromethane, 90 mg of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, 63 mg of 1-hydroxy-7-azabenzotriazole and 0.3 mL of triethylamine was stirred in a sealed tube at the ambient temperature of 70-80°C for 3 hours. The reaction mixture was cooled to room temperature and then the solvent was distilled off under reduced pressure. To the obtained residue were added 1 mL of 2 mol/L sodium hydroxide aqueous solution, 4 mL of tetrahydrofuran, and 2 mL of methanol, and the mixture was stirred at the ambient temperature of 40-50°C for 20 minutes, and the solvent was distilled off under reduced pressure. To the reaction mixture were added ethyl acetate and saturated sodium chloride aqueous solution, the organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [chloroform:methanol gradient elution = 100:0 to 80:20] to obtain 8 mg of N-(2,3-dihydro-1H-inden-5-yl)-1-ethyl-N-methyl-4-(1-methylpiperidin-4-yl)-2-
- 15
- 20
- 25

oxo-1,2-dihydroquinoline-6-carboxamide as a white solid.

¹H-NMR (DMSO-D₆) δ: 1.15 (3H, t, J = 7.0 Hz), 1.36-1.57 (4H, m), 1.89-2.06 (4H, m), 2.24 (3H, s), 2.69-2.89 (6H, m), 3.29-3.37 (1H, m), 3.39 (3H, s), 4.21 (2H, q, J = 7.1 Hz), 6.38 (1H, s), 6.84-6.92 (1H, m), 7.07 (1H, d, J = 8.0 Hz), 7.15 (1H, s), 7.52-7.61 (2H, m), 7.75 (1H, dd, J = 8.9, 1.6 Hz).

[0174]

Example 18



By the same method as Reference Example 2, from 1-ethyl-4-(morpholin-4-yl)-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinolin-2(1H)-one and 4-bromo-1-methyl-5-phenyl-1H-pyrazole was obtained 1-ethyl-6-(1-methyl-5-phenyl-1H-pyrazol-4-yl)-4-(morpholin-4-yl)quinolin-2(1H)-one.

¹H-NMR (DMSO-D₆) δ: 1.17 (3H, t, J = 6.9 Hz), 2.65-2.76 (4H, m), 3.42-3.53 (4H, m), 3.74 (3H, s), 4.21 (2H, q, J = 6.8 Hz), 5.95 (1H, s), 7.35 (1H, d, J = 2.0 Hz), 7.37-7.64 (7H, m), 7.81 (1H, s).

[0175]

Examples 19-244

Confirming to the procedure described in the specification, the compounds indicated in Tables 1-25 were produced by publicly known reactions, such as condensation, addition, oxidization, reduction, rearrangement, substitution, halogenation, dehydration, hydrolysis or the like, or the appropriate combination of these reactions.

[0176]

[Table 1]

Example No.	Structural Formula	Compound Name	MS
19		N-(1-ethyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	404(M+H)
20		N-(1-ethyl-4-(octahydroisoquinolin-2(1H)-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	444(M+H)
21		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-2-hydroxy-N-methylbenzamide	408(M+H)
22		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-2-(trifluoromethoxy)benzamide	476(M+H)
23		2-cyano-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	417(M+H)
24		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-propylbenzamide	434(M+H)
25		3-ethoxy-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	436(M+H)
26		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-2-(trifluoromethyl)benzamide	460(M+H)
27		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-2-fluoro-N-methylbenzamide	410(M+H)

[0177]

[Table 2]

Example No.	Structural Formula	Compound Name	MS
28		N-(4-((E)-2-cyclopropylvinyl)-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	373(M+H)
29		N-(1-ethyl-4-((1E)-4-hydroxybut-1-en-1-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	377(M+H)
30		N-(4-butyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	363(M+H)
31		4-cyclopropyl-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	432(M+H)
32		4-((E)-2-cyclopropylvinyl)-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	458(M+H)
33		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)benzamide	487(M+H)
34		4-(cyclohex-1-en-1-yl)-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	472(M+H)
35		N-(4-(2-cyclopropylethyl)-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	375(M+H)
36		N-(1,4-diethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	335(M+H)

[0178]

[Table 3]

Example No.	Structural Formula	Compound Name	MS
37		N-(1-ethyl-4-(4-hydroxybutyl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	379(M+H)
38		4-(2-cyclopropylethyl)-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	460(M+H)
39		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-(1-methylpiperidin-4-yl)benzamide	489(M+H)
40		4-cyclohexyl-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	474(M+H)
41		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-4-((1E)-4-hydroxybut-1-en-1-yl)-N-methylbenzamide	462(M+H)
42		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-sulfamoylbenzamide	471(M+H)
43		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-(methylsulfonyl)benzamide	470(M+H)
44		4-(dimethylamino)-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	435(M+H)
45		4-ethyl-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	420(M+H)

[0179]

[Table 4]

Example No.	Structural Formula	Compound Name	MS
46		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-(propan-2-yl)benzamide	434(M+H)
47		2-amino-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-(trifluoromethyl)benzamide	430(M+H)
48		2-amino-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-4-fluoro-N-methylbenzamide	380(M+H)
49		2-amino-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	362(M+H)
50		2-amino-4-cyclopropyl-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	402(M+H)
51		2-amino-4-(2-cyclopropylethyl)-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	430(M+H)
52		2-amino-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-(propan-2-yl)benzamide	404(M+H)
53		2-amino-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-4-ethyl-N-methylbenzamide	390(M+H)
54		2-amino-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-propylbenzamide	404(M+H)
55		N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	349(M+H)

[0180]

[Table 5]

Example No.	Structural Formula	Compound Name	MS
56		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-propylbenzamide	389(M+H)
57		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-(propan-2-yl)benzamide	389(M+H)
58		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-4-ethyl-N-methylbenzamide	375(M+H)
59		4-(2-cyclopropylethyl)-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	415(M+H)
60		4-cyclopropyl-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	387(M+H)
61		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-4-fluoro-N-methylbenzamide	365(M+H)
62		2-amino-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-4-fluoro-N-methylbenzamide	382(M+H)
63		2-amino-4-cyclopropyl-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	404(M+H)
64		2-amino-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	364(M+H)
65		2-amino-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methyl-4-(trifluoromethyl)benzamide	432(M+H)

[0181]

[Table 6]

Example No.	Structural Formula	Compound Name	MS
66		2-amino-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methyl-4-(propan-2-yl)benzamide	406(M+H)
67		2-amino-4-ethyl-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	392(M+H)
68		2-amino-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methyl-4-propylbenzamide	406(M+H)
69		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-4-(dimethylamino)-N-methylbenzamide	390(M+H)
70		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-4-(ethylamino)-N-methylbenzamide	390(M+H)
71		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-(phenylamino)benzamide	438(M+H)
72		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-(methyl(propyl)amino)benzamide	418(M+H)
73		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-(piperidin-1-yl)benzamide	430(M+H)
74		2-amino-N-(1-ethyl-4-methyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	354(M+H)
75		N-(4-(butan-2-yl)-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	363(M+H)

[0182]

[Table 7]

Example No.	Structural Formula	Compound Name	MS
76		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N,2-dimethylbenzamide	361(M+H)
77		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N,3-dimethylbenzamide	361(M+H)
78		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N,4-dimethylbenzamide	361(M+H)
79		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-2-methoxy-N-methylbenzamide	377(M+H)
80		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-3-methoxy-N-methylbenzamide	377(M+H)
81		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-4-(diethylamino)-N-methylbenzamide	418(M+H)
82		4-tert-butyl-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	403(M+H)
83		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-2-hydroxy-N-methylbenzamide	363(M+H)
84		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-3-hydroxy-N-methylbenzamide	363(M+H)
85		3-amino-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	362(M+H)

[0183]

[Table 8]

Example No.	Structural Formula	Compound Name	MS
86		4-amino-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	362(M+H)
87		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-3-(dimethylamino)-N-methylbenzamide	390(M+H)
88		4-(acetylamino)-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	404(M+H)
89		3-(acetylamino)-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	404(M+H)
90		2-(acetylamino)-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	404(M+H)
91		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-((methylsulfonyl)amino)benzamide	440(M+H)
92		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-3-((methylsulfonyl)amino)benzamide	440(M+H)
93		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-2-((methylsulfonyl)amino)benzamide	440(M+H)
94		2-amino-N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N,4-methylbenzamide	376(M+H)
95		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-2-(methylamino)benzamide	376(M+H)
96		N-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-2-(dimethylamino)-N-methylbenzamide	390(M+H)

[0184]

[Table 9]

Example No.	Structural Formula	Compound Name	MS
97		N-(1-ethyl-4-(1-(2-hydroxyethyl)piperidin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	434(M+H)
98		N-(1-ethyl-4-(1-(2-methoxyethyl)piperidin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	448(M+H)
99		3-chloro-N-(1-ethyl-2-oxo-4-(piperidin-4-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	425(M+H)
100		N-(1-(ethyl-2-oxo-4-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)-1,2-dihydroquinolin-6-yl)-N,4-dimethylbenzamide	486(M+H)
101		3-chloro-N-(1-ethyl-2-oxo-4-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	507(M+H)
102		3-chloro-N-(4-(1-(cyanomethyl)piperidin-4-yl)-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	464(M+H)
103		N-(1-ethyl-2-oxo-4-(4-oxocyclohexyl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	401(M-H)

[0185]

[Table 10]

Example No.	Structural Formula	Compound Name	MS
104		N-(4-(1-(cyanomethyl)piperidin-4-yl)-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	429(M+H)
105		N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-2-hydroxy-N,4-dimethylbenzamide	377(M-H)
106		3-chloro-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	384(M+H)
107		N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-2-hydroxy-N,5-dimethylbenzamide	377(M-H)
108		2-(acetilamino)-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N,4-dimethylbenzamide	418 (M-H)
109		2-(acetilamino)-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-4-fluoro-N-methylbenzamide	422(M-H)
110		4-chloro-N-(1-ethyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	439(M+H)
111		2-amino-4-chloro-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	399(M+H)
112		3-amino-4-chloro-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	399(M+H)
113		4-amino-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-2-hydroxy-N-methylbenzamide	380(M+H)

[0186]

[Table 11]

Example No.	Structural Formula	Compound Name	MS
114		4-chloro-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-2-methoxy-N-methylbenzamide	414(M+H)
115		4-chloro-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-3-methoxy-N-methylbenzamide	414(M+H)
116		4-chloro-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-3-(3-hydroxypropoxy)-N-methylbenzamide	458(M+H)
117		4-chloro-3-(3-(dimethylamino)propoxy)-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	484(M+H)
118		4-chloro-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-2-(3-hydroxypropoxy)-N-methylbenzamide	458(M+H)
119		4-chloro-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-2-((2-hydroxyethyl)amino)-N-methylbenzamide	443(M+H)
120		4-chloro-2-((2-(dimethylamino)ethyl)amino)-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	469(M+H)
121		N-(1-ethyl-4-methyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylthiophene-2-carboxamide	327(M+H)
122		N-(1-ethyl-4-methyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylthiophene-3-carboxamide	327(M+H)
123		N-(1-ethyl-4-methyl-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-1H-pyrrole-2-carboxamide	310(M+H)

[0187]

[Table 12]

Example No.	Structural Formula	Compound Name	MS
124		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylnaphthalene-2-carboxamide	442(M+H)
125		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylnaphthalene-1-carboxamide	442(M+H)
126		4-tert-butyl-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	448(M+H)
127		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-4-methoxy-N-methylbenzamide	422(M+H)
128		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbiphenyl-4-carboxamide	468(M+H)
129		4-cyano-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	417(M+H)
130		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-3-fluoro-N-methylbenzamide	410(M+H)
131		3-cyano-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	417(M+H)
132		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-3-(trifluoromethyl)benzamide	460(M+H)

[0188]

[Table 13]

Example No.	Structural Formula	Compound Name	MS
133		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-3-methoxy-N-methylbenzamide	422(M+H)
134		3-(dimethylamino)-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	435(M+H)
135		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-4-fluoro-N-methylbenzamide	410(M+H)
136		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methyl-4-(trifluoromethyl)benzamide	460(M+H)
137		N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-3-hydroxy-N-methylbenzamide	408(M+H)
138		1-ethyl-N-methyl-4-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)-2-oxo-N-phenyl-1,2-dihydroquinoline-6-carboxamide	402(M+H)
139		1-ethyl-N-methyl-4-(1-methylpiperidin-4-yl)-2-oxo-N-phenyl-1,2-dihydroquinoline-6-carboxamide	404(M+H)
140		4-cyclohexyl-1-ethyl-N-methyl-2-oxo-N-phenyl-1,2-dihydroquinoline-6-carboxamide	389(M+H)
141		4-(cyclohex-1-en-1-yl)-1-ethyl-N-methyl-2-oxo-N-phenyl-1,2-dihydroquinoline-6-carboxamide	387(M+H)

[0189]

[Table 14]

Example No.	Structural Formula	Compound Name	MS
142		4-cyclopropyl-1-ethyl-N-methyl-2-oxo-N-phenyl-1,2-dihydroquinoline-6-carboxamide	347(M+H)
143		6-(2,3-dihydro-1H-indol-1-ylcarbonyl)-1-ethyl-4-(1-methylpiperidin-4-yl)quinolin-2(1H)-one	416(M+H)
144		4-(((1-ethyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)carbonyl)-3,4-dihydroquinoxalin-2(1H)-one	445(M+H)
145		N-(2-cyanophenyl)-4-cyclopropyl-1-ethyl-N-methyl-2-oxo-1,2-dihydroquinoline-6-carboxamide	372(M+H)
146		4-cyclopropyl-1-ethyl-N-(4-fluorophenyl)-N-methyl-2-oxo-1,2-dihydroquinoline-6-carboxamide	365(M+H)
147		4-cyclopropyl-1-ethyl-N-methyl-2-oxo-N-(4-(trifluoromethyl)phenyl)-1,2-dihydroquinoline-6-carboxamide	415(M+H)
148		4-cyclopropyl-1-ethyl-N-(2-fluorophenyl)-N-methyl-2-oxo-1,2-dihydroquinoline-6-carboxamide	365(M+H)
149		4-cyclopropyl-1-ethyl-N-methyl-2-oxo-N-(2-(trifluoromethyl)phenyl)-1,2-dihydroquinoline-6-carboxamide	415(M+H)
150		4-cyclopropyl-1-ethyl-N-methyl-N-(2-methylphenyl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	361(M+H)
151		4-cyclopropyl-1-ethyl-N-methyl-N-(3-methylphenyl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	361(M+H)

[0190]

[Table 15]

Example No.	Structural Formula	Compound Name	MS
152		4-cyclopropyl-1-ethyl-N-methyl-N-(4-methylphenyl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	361(M+H)
153		4-cyclopropyl-1-ethyl-N-(3-fluorophenyl)-N-methyl-2-oxo-1,2-dihydroquinoline-6-carboxamide	365(M+H)
154		N-(3-cyanophenyl)-4-cyclopropyl-1-ethyl-N-methyl-2-oxo-1,2-dihydroquinoline-6-carboxamide	372(M+H)
155		N-(4-cyanophenyl)-4-cyclopropyl-1-ethyl-N-methyl-2-oxo-1,2-dihydroquinoline-6-carboxamide	372(M+H)
156		4-cyclopropyl-1-ethyl-N-methyl-2-oxo-N-(3-(trifluoromethyl)phenyl)-1,2-dihydroquinoline-6-carboxamide	415(M+H)
157		4-cyclopropyl-1-ethyl-N-methyl-2-oxo-N-(pyridin-2-yl)-1,2-dihydroquinoline-6-carboxamide	348(M+H)
158		4-cyclopropyl-N-(2,4-difluorophenyl)-1-ethyl-N-methyl-2-oxo-1,2-dihydroquinoline-6-carboxamide	383(M+H)
159		4-cyclopropyl-1-ethyl-N-methyl-2-oxo-N-(2,4,6-trifluorophenyl)-1,2-dihydroquinoline-6-carboxamide	401(M+H)
160		4-cyclopropyl-1-ethyl-N-(3-fluoro-4-methylphenyl)-N-methyl-2-oxo-1,2-dihydroquinoline-6-carboxamide	379(M+H)
161		4-cyclopropyl-N-(3,4-difluorophenyl)-1-ethyl-N-methyl-2-oxo-1,2-dihydroquinoline-6-carboxamide	383(M+H)
162		4-cyclopropyl-1-ethyl-N-(4-fluoro-3-methylphenyl)-N-methyl-2-oxo-1,2-dihydroquinoline-6-carboxamide	379(M+H)

[Table 16]

Example No.	Structural Formula	Compound Name	MS
163		4-cyclopropyl-1-ethyl-N-methyl-2-oxo-N-(pyridin-4-yl)-1,2-dihydroquinoline-6-carboxamide	348(M+H)
164		4-cyclopropyl-1-ethyl-N-methyl-2-oxo-N-(pyridin-3-yl)-1,2-dihydroquinoline-6-carboxamide	348(M+H)
165		1-ethyl-N,4-dimethyl-N-(3-methylphenyl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	335(M+H)
166		1-ethyl-N,4-dimethyl-N-(4-methylphenyl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	335(M+H)
167		1-ethyl-4-(1-(2-hydroxyethyl)piperidin-4-yl)-N-methyl-2-oxo-N-phenyl-1,2-dihydroquinoline-6-carboxamide	434(M+H)
168		1-ethyl-4-(1-(2-methoxyethyl)piperidin-4-yl)-N-methyl-2-oxo-N-phenyl-1,2-dihydroquinoline-6-carboxamide	448(M+H)
169		4-cyclopropyl-1-ethyl-N-methyl-N-(6-methylpyridin-2-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	362(M+H)
170		4-cyclopropyl-N-(4,6-dimethylpyridin-2-yl)-1-ethyl-N-methyl-2-oxo-1,2-dihydroquinoline-6-carboxamide	376(M+H)
171		4-cyclopropyl-1-ethyl-N-methyl-N-(5-methylpyridin-2-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	362(M+H)
172		4-cyclopropyl-1-ethyl-N-methyl-N-(6-methylpyridin-3-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	362(M+H)

[0192]

[Table 17]

Example No.	Structural Formula	Compound Name	MS
173		4-cyclopropyl-1-ethyl-N-methyl-N-(2-methylpyridin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	362(M+H)
174		1-ethyl-N-methyl-2-oxo-N-phenyl-4-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)-1,2-dihydroquinoline-6-carboxamide	472(M+H)
175		methyl 2-(((4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)carbonyl)(methyl)amino)benzoate	405(M+H)
176		3-(((4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)carbonyl)(methyl)amino)benzoic acid	391(M+H)
177		methyl 4-(((4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)carbonyl)(methyl)amino)benzoate	405(M+H)
178		2-(((4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)carbonyl)(methyl)amino)benzoic acid	391(M+H)
179		1-ethyl-4-(1-(2-hydroxyethyl)piperidin-4-yl)-N-methyl-2-oxo-N-(5,6,7,8-tetrahydronaphthalen-2-yl)-1,2-dihydroquinoline-6-carboxamide	488(M+H)
180		4-(1-benzoylpiperidin-4-yl)-1-ethyl-N-methyl-N-(4-methylphenyl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	508(M+H)

[0193]

[Table 18]

Example No.	Structural Formula	Compound Name	MS
181		1-ethyl-N-methyl-N-(4-methylphenyl)-4-(1-(methylsulfonyl)piperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	480(M-H)
182		1-ethyl-N-(2-(hydroxymethyl)-4-methylphenyl)-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	393(M+H)
183		1-ethyl-N-methyl-N-(4-methylphenyl)-4-(1-(2-methylpropanoyl)piperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	474(M+H)
184		N-(4-chlorophenyl)-1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	384(M+H)
185		1-ethyl-N-methyl-N-(4-methyl-3-(phenylcarbamoyl)phenyl)-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	480(M-H)
186		1-ethyl-N-methyl-N-(4-methyl-3-(propan-2-ylcarbamoyl)phenyl)-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	446(M-H)
187		N-(2-(dimethylcarbamoyl)-4-methylphenyl)-1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	434(M+H)
188		1-ethyl-N-methyl-N-(4-methyl-2-(phenylcarbamoyl)phenyl)-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	480 (M-H)
189		1-ethyl-N-methyl-N-(4-methyl-2-(propan-2-ylcarbamoyl)phenyl)-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	446(M-H)

[0194]

[Table 19]

Example No.	Structural Formula	Compound Name	MS
190		1-ethyl-N-methyl-4-(1-methylpiperidin-4-yl)-2-oxo-N-(5,6,7,8-tetrahydronaphthalen-2-yl)-1,2-dihydroquinoline-6-carboxamide	458(M+H)
191		1-ethyl-4-(1-(2-hydroxyethyl)piperidin-4-yl)-N-methyl-2-oxo-N-(6,7,8,9-tetrahydro-5H-benzo(7)annulen-2-yl)-1,2-dihydroquinoline-6-carboxamide	502(M+H)
192		N-(4-chlorophenyl)-1-ethyl-N-methyl-4-(1-(2-methylpropanoyl)piperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	494, 496(M+H)
193		1-ethyl-N-methyl-4-(1-(2-methylpropanoyl)piperidin-4-yl)-2-oxo-N-(4-(trifluoromethyl)phenyl)-1,2-dihydroquinoline-6-carboxamide	528(M+H)
194		4-cyano-N-(1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)-N-methylbenzamide	374(M+H)
195		1-ethyl-N-(2-methoxy-1-methyl-1H-indol-6-yl)-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	430(M+H)
196		tert-butyl 6-(((1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)carbonyl)(methyl)amino)-2,3-dihydro-1H-indole-1-carboxylate	490(M+H)
197		N-(4-chloro-3-(1H-tetrazol-5-yl)phenyl)-1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	451(M+H)

[0195]

[Table 20]

Example No.	Structural Formula	Compound Name	MS
198		(2-chloro-5-(((1-ethyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinolin-6-yl)carbonyl)(methyl)amino)phenyl)acetic acid	442(M+H)
199		N-(4-chloro-3-(2-hydroxyethyl)phenyl)-1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	428(M+H)
200		1-ethyl-N-(3-((2-hydroxyethyl)carbonyl)-4-methylphenyl)-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	450(M+H)
201		N-(4-chloro-3-methoxyphenyl)-1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	414(M+H)
202		N-(3-amino-4-chlorophenyl)-1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	399(M+H)
203		1-ethyl-N-methyl-N-(4-(methylsulfonyl)phenyl)-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	427(M+H)
204		N-(3-(butylcarbonyl)-4-methylphenyl)-1-ethyl-N-methyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	517(M+H)
205		1-ethyl-N-(3-((2-hydroxyethyl)carbonyl)-4-methylphenyl)-N-methyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	505(M+H)
206		1-ethyl-N-methyl-N-(4-methyl-2-(methylcarbonyl)phenyl)-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	475(M+H)

[0196]

[Table 21]

Example No.	Structural Formula	Compound Name	MS
207		1-ethyl-N-methyl-4-(1-methylpiperidin-4-yl)-N-(4-methyl-2-(propan-2-ylcarbamoyl)phenyl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	503(M+H)
208		1-ethyl-N-(2-((2-hydroxyethyl)carbamoyl)-4-methylphenyl)-N-methyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	505(M+H)
209		1-ethyl-N-(3-(2-methoxyethyl)-4-methylphenyl)-N-methyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	476(M+H)
210		1-ethyl-N-(3-(2-hydroxyethyl)-4-methylphenyl)-N-methyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	462(M+H)
211		1-ethyl-N-methyl-2-oxo-N-phenyl-4-(trifluoromethyl)-1,2-dihydroquinoline-6-carboxamide	375(M+H)
212		1-ethyl-N-methyl-N-(4-methylphenyl)-2-oxo-4-(trifluoromethyl)-1,2-dihydroquinoline-6-carboxamide	389(M+H)
213		1-ethyl-N-(3-fluoro-4-(trifluoromethyl)phenyl)-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	435(M+H)
214		1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-N-(3-(trifluoromethyl)phenyl)-1,2-dihydroquinoline-6-carboxamide	417(M+H)
215		N-(2-cyano-4-methylphenyl)-1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	388(M+H)

[0197]

[Table 22]

Example No.	Structural Formula	Compound Name	MS
216		1-ethyl-N-methyl-N-(6-methylpyridin-3-yl)-2-oxo-4-(tetrahydro-2H-pyran-4-yl)-1,2-dihydroquinoline-6-carboxamide	406(M+H)
217		1-ethyl-N-methyl-N-(3-methylphenyl)-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	363(M+H)
218		1-ethyl-N-(isoquinolin-6-yl)-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	400(M+H)
219		N-(4-chloro-2-cyanophenyl)-1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	408(M+H)
220		1-ethyl-N-methyl-N-(4-methyl-3-(trifluoromethyl)phenyl)-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	431(M+H)
221		1-ethyl-N-methyl-N-(4-methyl-3-(methylsulfonyl)phenyl)-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	441(M+H)
222		1-ethyl-N-methyl-N-(3-methyl-4-(methylsulfonyl)phenyl)-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	441(M+H)
223		1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-N-(quinoxalin-6-yl)-1,2-dihydroquinoline-6-carboxamide	401(M+H)
224		1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-N-(quinolin-7-yl)-1,2-dihydroquinoline-6-carboxamide	400(M+H)
225		1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-N-(2-(trifluoromethyl)phenyl)-1,2-dihydroquinoline-6-carboxamide	417(M+H)

[0198]

[Table 23]

Example No.	Structural Formula	Compound Name	MS
226		1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-N-(4-(trifluoromethyl)phenyl)-1,2-dihydroquinoline-6-carboxamide	417(M+H)
227		N-(3,4-difluorophenyl)-1-ethyl-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	385(M+H)
228		1-ethyl-N-(4-fluoro-3-(trifluoromethyl)phenyl)-N-methyl-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	435(M+H)
229		1-ethyl-N-methyl-2-oxo-N-(3-oxo-2,3-dihydro-1H-isoindol-5-yl)-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	404(M+H)
230		1-ethyl-4-(4-hydroxypiperidin-1-yl)-N-methyl-2-oxo-N-(3-(trifluoromethyl)phenyl)-1,2-dihydroquinoline-6-carboxamide	474(M+H)
231		1-ethyl-N-methyl-N-(2-methyl-3-oxo-2,3-dihydro-1H-isoindol-5-yl)-2-oxo-4-(propan-2-yl)-1,2-dihydroquinoline-6-carboxamide	418(M+H)
232		1-ethyl-4-(3-hydroxyazetidin-1-yl)-N-methyl-2-oxo-N-(3-(trifluoromethyl)phenyl)-1,2-dihydroquinoline-6-carboxamide	446(M+H)
233		1-ethyl-4-(4-(2-hydroxyethyl)piperidin-1-yl)-N-methyl-2-oxo-N-(3-(trifluoromethyl)phenyl)-1,2-dihydroquinoline-6-carboxamide	502(M+H)
234		4-(dimethylamino)-1-ethyl-N-methyl-2-oxo-N-(3-(trifluoromethyl)phenyl)-1,2-dihydroquinoline-6-carboxamide	418(M+H)

[0199]

[Table 24]

Example No.	Structural Formula	Compound Name	MS
235		N,1-dimethyl-N-(4-methylphenyl)-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	404(M+H)
236		1-ethyl-4-methyl-6-(2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one	346(M+H)
237		1-ethyl-6-(2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)-4-(piperidin-1-yl)quinolin-2(1H)-one	415(M+H)
238		1-ethyl-4-methyl-6-(3-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one	360(M+H)
239		6-(3-benzyl-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)-1-ethyl-4-methylquinolin-2(1H)-one	436(M+H)
240		1-ethyl-6-(3-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)-4-(piperidin-1-yl)quinolin-2(1H)-one	429(M+H)
241		6-(3-benzyl-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)-1-ethyl-4-(piperidin-1-yl)quinolin-2(1H)-one	505(M+H)
242		1-ethyl-4-(1-methylpiperidin-4-yl)-6-(2-oxo-5-phenylimidazolidin-1-yl)quinolin-2(1H)-one	431(M+H)


[0200]

[Table 25]

Example No.	Structural Formula	Compound Name	MS
243		1-ethyl-4-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)-6-(2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one	427(M+H)
244		1-ethyl-4-(1-methylpiperidin-4-yl)-6-(2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one	429(M+H)

[0201]

Example 245



5


To a suspension of 0.70 g of 1-(2-(4-chlorophenyl)-2-oxoethyl)-3-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)urea in 4.0 mL of dioxane was added 4.0 mL of concentrated hydrochloric acid at room temperature and the mixture was stirred for 2 hours. Water was added to the reaction mixture, and the solid matter was filtered and washed with water and diisopropyl ether. The obtained solid matter was purified by silica gel column chromatography [chloroform:methanol gradient elution = 100:0 to 90:10]. To the obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 0.59 g of 6-(5-(4-chlorophenyl)-2-oxo-2,3-dihydro-1H-imidazol-1-yl)-1-ethyl-4-(morpholin-4-yl)quinolin-2(1H)-one as a white solid.

¹H-NMR (DMSO-D₆) δ: 1.19 (3H, t, J = 6.8 Hz), 2.64-2.73 (4H, m), 3.50-3.57 (4H, m), 4.23 (2H, q, J = 6.8 Hz), 6.00 (1H, s), 6.95 (1H, s), 7.07-7.13 (2H, m), 7.24 (1H, d, J = 2.4 Hz), 7.32-7.38 (2H, m), 7.57 (1H, dd, J = 9.1, 2.3 Hz), 7.65 (1H, d, J = 9.0 Hz), 10.67 (1H, s).

MS (ESI, m/z): 451(M+H), 449(M-H)

[0202]

20 Example 246




To a solution of 300 mg of 6-(5-(4-chlorophenyl)-2-oxo-2,3-dihydro-1H-imidazol-1-yl)-1-ethyl-4-(morpholin-4-yl)quinolin-2(1H)-one in 3 mL of N,N-dimethylformamide was added under ice cooling 30 mg of 60% sodium hydride and the mixture was stirred for 5 minutes. To the reaction mixture, under ice cooling was added 50 μ L of methyl iodide, the mixture was stirred at room temperature for 2 hours. To the reaction mixture were added under ice cooling ethyl acetate and water, and the pH of the mixture was adjusted to 2.0 with 1 mol/L hydrochloric acid, the solid matter was filtered, and washed with water and diisopropyl ether to obtain 203 mg of 6-(5-(4-chlorophenyl)-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-1-yl)-1-ethyl-4-(morpholin-4-yl)quinolin-2(1H)-one as a white solid.

$^1\text{H-NMR}$ (DMSO- D_6) δ : 1.19 (3H, t, $J = 7.0$ Hz), 2.64-2.72 (4H, m), 3.28 (3H, s), 3.50-3.57 (4H, m), 4.23 (2H, q, $J = 6.8$ Hz), 6.01 (1H, s), 7.05-7.12 (2H, m), 7.07 (1H, s), 7.26 (1H, d, $J = 2.4$ Hz), 7.34-7.39 (2H, m), 7.57 (1H, dd, $J = 9.0, 2.4$ Hz), 7.66 (1H, d, $J = 9.0$ Hz).

MS (ESI, m/z): 465(M+H)

[0203]

Example 247




To a suspension of 330 mg of 1-ethyl-4-(morpholin-4-yl)-6-(2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one in 2 mL of N,N-dimethylacetamide under ice cooling was added 35 mg of 60% sodium hydride and the mixture was stirred for 5 minutes. To the reaction mixture under ice cooling was added 119 μ L of 2-iodopropane, and the mixture was stirred at room temperature for 1 hour. To the reaction mixture was added 60 μ L of 2-iodopropane, and the mixture was stirred at room temperature for 1 hour. To the reaction mixture under ice cooling were added ethyl acetate and water, and the pH of the mixture was

adjusted to 2.0 with 1 mol/L hydrochloric acid. The organic layer was fractionated, washed with water and saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [ethyl acetate to chloroform:methanol gradient elution = 95:5 to 90:10]. To the obtained residue were added diisopropyl ether and ethyl acetate, and the solid matter was filtered. The obtained solid matter was purified by silica gel column chromatography [ethyl acetate]. To the obtained residue were added diisopropyl ether and ethyl acetate, and the solid matter was filtered to obtain 68 mg of 1-ethyl-4-(morpholin-4-yl)-6-(2-oxo-5-phenyl-3-(propan-2-yl)-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one as a light brown solid.

¹H-NMR (CDCl₃) δ: 1.36 (3H, t, J = 7.1 Hz), 1.43 (6H, d, J = 6.8 Hz), 2.63-2.70 (4H, m), 3.55-3.62 (4H, m), 4.32 (2H, q, J = 7.2 Hz), 4.55 (1H, quint, J = 6.8 Hz), 6.09 (1H, s), 6.51 (1H, s), 7.07-7.12 (2H, m), 7.15-7.28 (3H, m), 7.33 (1H, d, J = 2.4 Hz), 7.43 (1H, d, J = 9.0 Hz), 7.72 (1H, dd, J = 9.0, 2.4 Hz).

[0204]

Example 248




A mixture of 100 mg of 4-cyclopropyl-1-ethyl-6-(2-phenylhydrazinyl)quinolin-2(1H)-one, 46 mg of ethyl carbamoyl carbamate and 3 mL of toluene was stirred at 140°C for 1 hour and 30 minutes using microwave equipment. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, washed successively with water and saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [hexane:ethyl acetate gradient elution = 50:50 to 0:100]. To the obtained residue was added chloroform, and the solid matter was filtered to obtain 15 mg of 1-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-2-phenyl-1,2,4-triazolidine-3,5-dione as a light brown solid.

¹H-NMR (DMSO-D₆) δ: 0.57-0.65 (2H, m), 0.92-1.01 (2H, m), 1.14 (3H, t, J = 7.0 Hz), 2.03-2.14 (1H, m), 4.20 (2H, q, J = 7.1 Hz), 6.29 (1H, s), 7.20-7.27 (1H, m), 7.34-7.47 (4H, m), 7.60 (1H, d, J = 9.3 Hz), 7.70 (1H, dd, J = 9.1, 2.6 Hz), 8.03 (1H, d, J = 2.4 Hz), 12.09 (1H, s).

[0205]

Example 249




To a suspension of 2.0 g of 6-amino-1-ethyl-4-(morpholin-4-yl)quinolin-2(1H)-
 5 one in 10 mL of pyridine under ice cooling was added 1.12 mL of 4-chlorobenzoyl chloride and the mixture was stirred at room temperature for 1 hour. To the reaction mixture under ice cooling were added ethyl acetate and water, and the pH of the mixture was adjusted to 2.0 with 2 mol/L hydrochloric acid. The solid matter was filtered and washed with water and diisopropyl ether to obtain 3.01 g of 4-chloro-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)benzamide as a white solid.
 10

¹H-NMR (DMSO-D₆) δ: 1.19 (3H, t, J = 7.0 Hz), 3.03-3.13 (4H, m), 3.81-3.90 (4H, m), 4.24 (2H, q, J = 7.1 Hz), 6.04 (1H, s), 7.59 (1H, d, J = 9.3 Hz), 7.64 (2H, d, J = 8.5 Hz), 7.98 (1H, dd, J = 9.3, 2.4 Hz), 8.01 (2H, d, J = 8.5 Hz), 8.35 (1H, d, J = 2.2 Hz), 10.48 (1H, s).

[0206]

15 Example 250




To a suspension of 3.0 g of 4-chloro-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)benzamide in 24 mL of N,N-dimethylacetamide, 0.35 g of 60% sodium hydride was added under ice cooling and the mixture was stirred for 10 minutes. To the
 20 reaction mixture under ice cooling was added 0.68 mL of methyl iodide, the mixture was stirred at room temperature for 1 hour. To the reaction mixture were added ethyl acetate and iced water, and the pH of the mixture was adjusted to 2.0 with 2 mol/L hydrochloric acid. The organic layer was fractionated, washed with water and saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off

under reduced pressure. To the obtained residue were added diisopropyl ether and ethyl acetate, the solid matter was filtered, washed with diisopropyl ether to obtain 2.55 g of 4-chloro-N-(1-ethyl-4-(morpholin-4-yl)-2-oxo-1,2-dihydroquinolin-6-yl)-N-methylbenzamide as a white solid.

¹H-NMR (DMSO-D₆) δ: 1.16 (3H, t, J = 7.1 Hz), 2.45-2.60 (4H, m), 3.42 (3H, s), 3.60-3.71 (4H, m), 4.19 (2H, q, J = 6.9 Hz), 5.96 (1H, s), 7.21 (1H, s), 7.28 (2H, d, J = 8.3 Hz), 7.33 (2H, d, J = 8.3 Hz), 7.60 (1H, d, J = 9.0 Hz), 7.68 (1H, dd, J = 9.0, 2.2 Hz).

[0207]

Example 251




A mixture of 1.5 g of 4-chloro-N-((1E)-1-(dimethylamino)ethylidene)-1-ethyl-2-oxo-1,2-dihydroquinoline-6-carboxamide, 0.51 ml of phenylhydrazine and 12 mL of acetic acid was stirred at the ambient temperature of 120°C for 2 hours. The reaction mixture was cooled to room temperature and then the solvent was distilled off under reduced pressure. To the obtained residue were added diisopropyl ether and ethyl acetate, and the solid matter was filtered and washed with diisopropyl ether to obtain 1.37 g of 4-chloro-1-ethyl-6-(3-methyl-1-phenyl-1H-1,2,4-triazol-5-yl)quinolin-2(1H)-one as a light brown solid.

¹H-NMR (DMSO-D₆) δ: 1.18 (3H, t, J = 7.1 Hz), 2.41 (3H, s), 4.25 (2H, q, J = 6.9 Hz), 6.97 (1H, s), 7.44-7.58 (5H, m), 7.72 (1H, d, J = 9.0 Hz), 7.77 (1H, dd, J = 9.0, 2.0 Hz), 7.99 (1H, d, J = 2.0 Hz).

[0208]

Example 252




A mixture of 1.37 g of 4-chloro-1-ethyl-6-(3-methyl-1-phenyl-1H-1,2,4-triazol-5-yl)quinolin-2(1H)-one, 0.48 g of cyclopropylboric acid, 1.6 g of tripotassium phosphate, 0.26 g of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 15 mL of dioxane and 3.0 mL of water was stirred under nitrogen atmosphere at the ambient temperature of 100-

110°C for 3 hours. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, washed successively with water and saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [hexane:ethyl acetate gradient elution = 50:50 to 0:100]. To the obtained residue were added diisopropyl ether and cyclohexane, and the solid matter was filtered to obtain 1.10 g of 4-cyclopropyl-1-ethyl-6-(3-methyl-1-phenyl-1H-1,2,4-(triazol-5-yl)quinolin-2(1H)-one as a slightly brown solid.

¹H-NMR (DMSO-D₆) δ: 0.54-0.61 (2H, m), 0.70-0.79 (2H, m), 1.17 (3H, t, J = 7.1 Hz), 1.70-1.79 (1H, m), 2.40 (3H, s), 4.23 (2H, q, J = 7.1 Hz), 6.31 (1H, s), 7.42-7.57 (5H, m), 7.64 (1H, d, J = 9.0 Hz), 7.81 (1H, dd, J = 8.9, 2.1 Hz), 8.03 (1H, d, J = 2.0 Hz).

[0209]

Example 253




A mixture of 41 mg of 1-ethyl-6-iodo-4-(morpholin-4-yl)quinolin-2(1H)-one, 17 mg of (5S)-5-phenylpyrrolidin-2-one, 1 mg of copper iodide(I), 4 mg of 4,7-dimethoxy-1,10-phenanthroline, 48 mg of cesium carbonate, and 0.5 mL of N-methyl-2-pyrrolidone was stirred at the ambient temperature of 130-135°C for 30 minutes. The reaction mixture was cooled to room temperature, and then ethyl acetate was added to the reaction mixture, and the insoluble matter was filtered off. To the filtrate were added water and saturated sodium chloride aqueous solution, and the organic layer was fractionated, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by basic silica gel chromatography [hexane:ethyl acetate gradient elution = 70:30 to 40:60]. To the obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 23 mg of 1-ethyl-4-(morpholin-4-yl)-6-((5S)-2-oxo-5-phenylpyrrolidin-1-yl)quinolin-2(1H)-one as a light yellow solid.

¹H-NMR (CDCl₃) δ: 1.29 (3H, t, J = 7.2 Hz), 2.02-2.08 (1H, m), 2.60-2.97 (7H, m), 3.73-3.80 (4H, m), 4.20-4.33 (2H, m), 5.29-5.32 (1H, m), 6.11 (1H, s), 7.23-7.35 (6H, m), 7.75 (1H, d, J = 2.4 Hz), 7.81 (1H, dd, J = 8.8, 2.4 Hz).

[0210]

Example 254




A mixture of 38 mg of 4-cyclopropyl-1-ethyl-6-iodoquinolin-2(1H)-one, 18 mg of (5S)-5-phenylpyrrolidin-2-one, 1 mg of copper iodide(I), 4 mg of 4,7-dimethoxy-1,10-

5 phenanthroline, 51 mg of cesium carbonate, and 0.5 mL of N-methyl-2-pyrrolidone was stirred at the ambient temperature of 130-135°C for 1 hour and 15 minutes. The reaction mixture was cooled to room temperature, and then ethyl acetate was added to the reaction mixture, and the insoluble matter was filtered off. To the filtrate were added water and saturated sodium chloride aqueous solution, the organic layer was fractionated, washed with saturated sodium
10 chloride aqueous solution, and then dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by basic silica gel chromatography [hexane:ethyl acetate gradient elution = 80:20 to 60:40] to obtain 22 mg of 4-cyclopropyl-1-ethyl-6-((5S)-2-oxo-5-phenylpyrrolidin-1-yl)quinolin-2(1H)-one as a light yellow solid.

15 ¹H-NMR (CDCl₃) δ: 0.50-0.57 (1H, m), 0.64-0.71 (1H, m), 0.82-0.99 (2H, m), 1.28 (3H, t, J = 7.2 Hz), 1.79-1.86 (1H, m), 2.05-2.12 (1H, m), 2.63-2.85 (3H, m), 4.26 (2H, q, J = 7.5 Hz), 5.27-5.30 (1H, m), 6.39 (1H, d, J = 1.2 Hz), 7.25-7.40 (6H, m), 7.79 (1H, dd, J = 9.2, 2.4 Hz), 7.99 (1H, d, J = 2.4 Hz).

[0211]

20 Example 255




A mixture of 60 mg of 1-ethyl-6-iodo-4-(morpholin-4-yl)quinolin-2(1H)-one, 25 mg of 2-phenyl-1H-imidazole, 1.1 mg of copper oxide(I), 5.6 mg of 4,7-dimethoxy-1,10-phenanthroline, 29 mg of polyethylene glycol, 71 mg of cesium carbonate, and 4 mL of
25 butyronitrile was stirred under nitrogen atmosphere at 170°C for 40 minutes using microwave equipment.. The reaction mixture was cooled to room temperature, and then ethyl acetate and

water were added. The organic layer was fractionated, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by thin layer chromatography for fractionation. To the obtained residue was added hexane, and the solid matter was filtered to obtain 10 mg of 1-ethyl-4-(morpholin-4-yl)-6-(2-phenyl-1H-imidazol-1-yl)-quinolin-2(1H)-one as a white solid.

¹H-NMR (DMSO-D₆) δ: 1.20 (3H, t, J = 7.3 Hz), 2.57-2.66 (4H, m), 3.42-3.52 (4H, m), 4.26 (2H, q, J = 6.8 Hz), 6.03 (1H, s), 7.22 (1H, d, J = 1.3 Hz), 7.28-7.38 (6H, m), 7.59 (1H, s), 7.71-7.77 (2H, m).

MS (ESI, m/z):401(M+H)
[0212]

Example 256



A mixture of 50 mg of 1-ethyl-6-iodo-4-(morpholin-4-yl)quinolin-2(1H)-one, 23 mg of 4-methyl-2-phenyl-1H-imidazole, 1.9 mg of copper oxide(I), 6.2 mg of 4,7-dimethoxy-1,10-phenanthroline, 25 mg of polyethylene glycol, 64 mg of cesium carbonate, and 1.5 mL of butyronitrile was stirred under nitrogen atmosphere at 210°C for 45 minutes using microwave equipment. To the reaction mixture was added 1.9 mg of copper oxide(I), and the mixture was stirred at 225°C for 45 minutes using microwave equipment. To the reaction mixture were added 6.0 mg of copper oxide(I) and 6.2 mg of 4,7-dimethoxy-1,10-phenanthroline, and the mixture was stirred at 230°C for 1 hour and 30 minutes using microwave equipment. The reaction mixture was cooled to room temperature, and then the insoluble matter was filtered off, and ethyl acetate and water were added. The organic layer was fractionated, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by thin layer chromatography for fractionation. To the obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 5 mg of 1-ethyl-6-(4-methyl-2-phenyl-1H-imidazol-1-yl)-4-(morpholin-4-yl)quinolin-2(1H)-one as a white solid.


¹H-NMR (DMSO-D₆) δ: 1.20 (3H, t, J = 6.9 Hz), 2.23 (3H, s), 2.56-2.66 (4H, m), 3.41-3.51 (4H,

m), 4.20-4.30 (2H, m), 6.02 (1H, s), 7.22-7.36 (7H, m), 7.67-7.76 (2H, m).

MS (ESI, m/z):415(M+H)

[0213]

Example 257



5

A mixture of 50 mg of 1-ethyl-6-iodo-4-(morpholin-4-yl)quinolin-2(1H)-one, 28 mg of 1-phenyl-1H-imidazole, 50 mg of copper iodide(I), 1.5 mg of palladium acetate, and 2 mL of N,N-dimethylacetamide was stirred under nitrogen atmosphere at 185°C for 30 minutes using microwave equipment. The mixture was further stirred at 205°C for 20 minutes. The reaction mixture was cooled to room temperature and then N,N-dimethylacetamide was distilled off under reduced pressure. To the obtained residue was added methanol, and the solid matter was filtered. To the obtained solid matter was added chloroform, and the mixture was heated at reflux for 10 minutes. The reaction mixture was cooled to room temperature, and then the insoluble matter was filtered off. The filtrate was purified by silica gel column chromatography [chloroform:methanol]. To the obtained residue were added hexane and ethyl acetate, and the solid matter was filtered to obtain 10 mg of 1-ethyl-4-(morpholin-4-yl)-6-(1-phenyl-1H-imidazol-2-yl)quinolin-2(1H)-one as a light brown solid.

10

15


¹H-NMR (DMSO-D₆) δ: 1.16 (3H, t, J = 6.9 Hz), 2.64-2.73 (4H, m), 3.47-3.57 (4H, m), 4.19 (2H, q, J = 7.0 Hz), 5.97 (1H, s), 7.23-7.38 (3H, m), 7.39-7.63 (6H, m), 7.74-7.85 (1H, m).

20

MS (ESI, m/z):401(M+H)

[0214]

Example 258



25

A mixture of 50 mg of 1-ethyl-6-iodo-4-(morpholin-4-yl)quinolin-2(1H)-one, 28 mg of biphenyl-2-ylboronic acid, 6 mg of tris(dibenzylideneacetone)dipalladium(0), 85 mg of

cesium carbonate, and 2 mL of dioxane was stirred under nitrogen atmosphere at 150°C for 30 minutes using microwave equipment. The mixture was further stirred at 160°C for 30 minutes. The reaction mixture was cooled to room temperature, and then the insoluble matter was filtered off, and the solvent was distilled off under reduced pressure. The obtained residue was purified

5 by silica gel column chromatography [hexane:ethyl acetate]. To the obtained residue were added hexane and ethyl acetate, and the solid matter was filtered to obtain 6 mg of 6-(biphenyl-2-yl)-1-ethyl-4-(morpholin-4-yl)quinolin-2(1H)-one as a white solid.

¹H-NMR (DMSO-D₆) δ: 1.19 (3H, t, J = 6.9 Hz), 2.46-2.55 (4H, m), 3.50-3.58 (4H, m), 4.23 (2H, q, J = 7.0 Hz), 5.93 (1H, s), 7.13-7.22 (3H, m), 7.23-7.31 (3H, m), 7.42-7.53 (4H, m), 7.59-

10 7.70 (2H, m).

MS (ESI, m/z):411(M+H)

[0215]

Examples 259-276

Confirming to the procedure described in the specification, the compounds
15 indicated in Tables 26-28 were produced by publicly known reactions, such as condensation, addition, oxidization, reduction, rearrangement, substitution, halogenation, dehydration, hydrolysis or the like, or the appropriate combination of these reactions.

[0216]

[Table 26]

Example No.	Structural Formula	Compound Name	MS
259		1-ethyl-4-(morpholin-4-yl)-6-(2-oxo-5-phenyl-3-(2-(piperidin-1-yl)ethyl)-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one	528(M+H)
260		1-ethyl-4-(morpholin-4-yl)-6-(3-(2-(morpholin-4-yl)ethyl)-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one	530(M+H)
261		6-(3-(2-(dimethylamino)ethyl)-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)-1-ethyl-4-(morpholin-4-yl)quinolin-2(1H)-one	488(M+H)
262		1-ethyl-6-(3-(2-hydroxyethyl)-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)-4-(morpholin-4-yl)quinolin-2(1H)-one	461(M+H)
263		1-ethyl-6-(3-(2-methoxyethyl)-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)-4-(morpholin-4-yl)quinolin-2(1H)-one	475(M+H)
264		6-(3-butyl-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)-1-ethyl-4-(morpholin-4-yl)quinolin-2(1H)-one	473(M+H)
265		1-ethyl-4-(morpholin-4-yl)-6-(2-oxo-5-phenyl-3-propyl-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one	459(M+H)
266		1-ethyl-6-(3-ethyl-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-1-yl)-4-(morpholin-4-yl)quinolin-2(1H)-one	445(M+H)

[0217]

[Table 27]

Example No.	Structural Formula	Compound Name	MS
267		4-cyclopropyl-1-ethyl-6-(3-methyl-2-oxo-5-(4-(trifluoromethyl)phenyl)-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one	454(M+H)
268		4-cyclopropyl-1-ethyl-6-(2-oxo-5-(4-(trifluoromethyl)phenyl)-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one	440(M+H)
269		6-(5-(4-chlorophenyl)-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-1-yl)-4-cyclopropyl-1-ethylquinolin-2(1H)-one	420(M+H)
270		1-ethyl-6-(3-methyl-2-oxo-5-(4-(trifluoromethyl)phenyl)-2,3-dihydro-1H-imidazol-1-yl)-4-(morpholin-4-yl)quinolin-2(1H)-one	499(M+H)
271		6-(5-(4-chlorophenyl)-2-oxo-2,3-dihydro-1H-imidazol-1-yl)-4-cyclopropyl-1-ethylquinolin-2(1H)-one	406(M+H)
272		1-ethyl-6-(2-oxo-5-(4-(trifluoromethyl)phenyl)-2,3-dihydro-1H-imidazol-1-yl)-4-(morpholin-4-yl)quinolin-2(1H)-one	485(M+H)
273		1-ethyl-6-(3-methyl-5-(4-methylphenyl)-2-oxo-2,3-dihydro-1H-imidazol-1-yl)-4-(morpholin-4-yl)quinolin-2(1H)-one	445(M+H)


[0218]

[Table 28]

Example No.	Structural Formula	Compound Name	MS
274		1-ethyl-6-(5-(4-methylphenyl)-2-oxo-2,3-dihydro-1H-imidazol-1-yl)-4-(morpholin-4-yl)quinolin-2(1H)-one	431(M+H)
275		4-cyclopropyl-1-ethyl-6-(3-methyl-5-(4-methylphenyl)-2-oxo-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one	400(M+H)
276		4-cyclopropyl-1-ethyl-6-(5-(4-methylphenyl)-2-oxo-2,3-dihydro-1H-imidazol-1-yl)quinolin-2(1H)-one	386(M+H)

[0219]

Example 277



5


A mixture of 0.17 g of 4-chloro-1-ethyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinolin-2(1H)-one, 0.11 g of 3-bromo-2-phenylpyridine, 0.20 g of tripotassium phosphate, 33 mg of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 6 mL of dioxane and 2 mL of water was stirred under nitrogen atmosphere at the ambient temperature of 100-110°C for 1 hour. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel chromatography [hexane:ethyl acetate gradient elution = 100:0 to 60:40] to obtain 134 mg of 4-chloro-1-ethyl-6-(2-phenylpyridin-3-yl)quinolin-2(1H)-one as a white foam.

¹H-NMR (CDCl₃) δ: 1.33 (3H, t, J = 7.1 Hz), 4.31 (2H, q, J = 7.1 Hz), 6.89 (1H, s), 7.24-7.29 (3H, m), 7.32-7.42 (5H, m), 7.81 (1H, dd, J = 7.6, 1.7 Hz), 7.94 (1H, d, J = 2.2 Hz), 8.74 (1H,

dd, $J = 4.8, 1.7$ Hz).

[0220]

Example 278




5 A mixture of 36 mg of 4-chloro-1-ethyl-6-(2-phenylpyridin-3-yl)quinolin-2(1H)-one, 17 mg of cyclopropylboric acid, 43 mg of tripotassium phosphate, 8 mg of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 0.6 mL of dioxane and 0.2 mL of water was stirred at 120°C for 15 minutes using microwave equipment. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel chromatography [hexane:ethyl acetate gradient elution = 100:0 to 50:50]. To the obtained residue was added hexane, and the solid matter was filtered to obtain 3 mg of 4-cyclopropyl-1-ethyl-6-(2-phenylpyridin-3-yl)quinolin-2(1H)-one as a white solid.

$^1\text{H-NMR}$ (CDCl_3) δ : 0.53-0.59 (2H, m), 0.81-0.92 (2H, m), 1.34 (3H, t, $J = 7.1$ Hz), 1.71-1.80 (1H, m), 4.33 (2H, q, $J = 7.3$ Hz), 6.39-6.44 (1H, m), 7.20-7.49 (8H, m), 7.82 (1H, dd, $J = 7.6, 1.5$ Hz), 7.87 (1H, d, $J = 2.2$ Hz), 8.73 (1H, dd, $J = 4.6, 1.5$ Hz).

[0221]

20 Example 279




A mixture of 30 mg of 4-chloro-1-ethyl-6-(2-phenylpyridin-3-yl)quinolin-2(1H)-one, 0.7 mL of N,N-dimethylformamide, 35 mg of potassium carbonate and 21.5 μL of morpholine was stirred at 150°C for 5 minutes using microwave equipment. The mixture was further stirred at 200°C for 40 minutes. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, and the

aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel chromatography [hexane:ethyl acetate gradient elution = 100:0 to 0:100 to chloroform:methanol = 95:5]. To the obtained residue were added hexane and ethyl acetate, and the solid matter was filtered to obtain 8 mg of 1-ethyl-4-(morpholin-4-yl)-6-(2-phenylpyridin-3-yl)quinolin-2(1H)-one as a brown solid.

¹H-NMR (CDCl₃) δ: 1.38 (3H, t, J = 7.1 Hz), 2.57-2.65 (4H, m), 3.57-3.66 (4H, m), 4.34 (2H, q, J = 7.2 Hz), 6.09 (1H, s), 7.20-7.29 (2H, m), 7.35-7.49 (6H, m), 7.61 (1H, dd, J = 8.7, 2.1 Hz), 7.74 (1H, dd, J = 7.6, 1.7 Hz), 8.73 (1H, dd, J = 4.9, 1.7 Hz).

[0222]

Example 280




A mixture of 20 mg of tert-butyl 2-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-1-phenylhydrazinecarboxylate, 0.5 mL of diethyl methylmalonate and 1 mL of 20% sodium ethoxide-ethanol solution was stirred at 180°C for 3 minutes using microwave equipment. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added, and the pH of the mixture was adjusted to 1.0 with 2 mol/L hydrochloric acid. The organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [chloroform:methanol gradient elution = 100:0 to 95:5]. To the obtained residue was added ethyl acetate, and the solid matter was filtered to obtain 6 mg of 1-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-4-methyl-2-phenylpyrazolidine-3,5-dione as a white solid.

¹H-NMR (CDCl₃) δ: 0.58-0.65 (2H, m), 0.95-1.03 (2H, m), 1.30 (3H, t, J = 7.1 Hz), 1.64 (3H, d, J = 7.8 Hz), 1.84-1.94 (1H, m), 3.47 (1H, q, J = 7.7 Hz), 4.27 (2H, q, J = 7.2 Hz), 6.44 (1H, d, J = 1.2 Hz), 7.18-7.24 (1H, m), 7.31-7.39 (5H, m), 7.61 (1H, dd, J = 9.1 Hz, 2.6 Hz), 7.94 (1H, d, J = 2.4 Hz).

[0223]

Example 281




A mixture of 20 mg of tert-butyl 2-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-1-phenylhydrazinecarboxylate, 0.5 mL of diethyl diethylmalonate and 1 mL of 20% sodium ethoxide-ethanol solution was stirred at 150°C for 3 minutes using microwave equipment. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added, and the pH of the mixture was adjusted to 1.0 with 2 mol/L hydrochloric acid. The organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [chloroform:methanol gradient elution = 100:0 to 95:5]. To the obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 8 mg of 1-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-4-ethyl-2-phenylpyrazolidine-3,5-dione as a light yellow solid.

¹H-NMR (CDCl₃) δ: 0.58-0.65 (2H, m), 0.95-1.02 (2H, m), 1.10-1.18 (3H, m), 1.30 (3H, t, J = 7.2 Hz), 1.85-1.94 (1H, m), 2.15-2.24 (2H, m), 3.42 (1H, t, J = 5.5 Hz), 4.27 (2H, q, J = 7.2 Hz), 6.42-6.45 (1H, m), 7.18-7.24 (1H, m), 7.31-7.40 (5H, m), 7.61 (1H, dd, J = 9.1 Hz, 2.6 Hz), 7.95 (1H, d, J = 2.4 Hz).

[0224]

Examples 282a, 282b



A mixture of 56 mg of 4-cyclopropyl-1-ethyl-6-(2-phenylhydrazinyl)quinolin-2(1H)-one, 14 μL of 4-methylideneoxetan-2-one, 27 μL of triethylamine, and 1 mL of chloroform was heated at reflux for 2 hours. The reaction mixture was cooled to room temperature and then the solvent was distilled off under reduced pressure. The obtained residue

was purified by silica gel column chromatography [ethyl acetate:methanol gradient elution = 100:0 to 95:5] to obtain 7 mg of a) 4-cyclopropyl-1-ethyl-6-(3-methyl-5-oxo-2-phenyl-2,5-dihydro-1H-pyrazol-1-yl)quinolin-2(1H)-one as a white solid, and 4 mg of b) 4-cyclopropyl-1-ethyl-6-(5-methyl-3-oxo-2-phenyl-2,3-dihydro-1H-pyrazol-1-yl)quinolin-2(1H)-one as a white solid.

a) 4-cyclopropyl-1-ethyl-6-(3-methyl-5-oxo-2-phenyl-2,5-dihydro-1H-pyrazol-1-yl)quinolin-2(1H)-one


¹H-NMR (CDCl₃) δ: 0.63-0.69 (2H, m), 0.99-1.05 (2H, m), 1.28 (3H, t, J = 7.2 Hz), 1.91-2.00 (1H, m), 2.12 (3H, d, J = 1.0 Hz), 4.26 (2H, q, J = 7.0 Hz), 5.61 (1H, d, J = 0.7 Hz), 6.40 (1H, d, J = 1.0 Hz), 7.21-7.39 (6H, m), 7.71 (1H, dd, J = 9.3 Hz, 2.4 Hz), 7.95 (1H, d, J = 2.4 Hz).

b) 4-cyclopropyl-1-ethyl-6-(5-methyl-3-oxo-2-phenyl-2,3-dihydro-1H-pyrazol-1-yl)quinolin-2(1H)-one

¹H-NMR (CDCl₃) δ: 0.64-0.70 (2H, m), 1.02-1.09 (2H, m), 1.30 (3H, t, J = 7.1 Hz), 1.90-1.99 (1H, m), 2.14 (3H, s), 4.27 (2H, q, J = 7.0 Hz), 5.61-5.65 (1H, m), 6.44-6.48 (1H, m), 7.13 (1H, t, J = 7.3 Hz), 7.22-7.42 (6H, m), 7.87 (1H, d, J = 2.2 Hz).

[0225]

Example 283




A mixture of 40 mg of 1-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-2-phenylethane-1,2-dione, 1 mL of acetic acid, 89 mg of ammonium acetate, and 32 μL of acetaldehyde was stirred at 120°C for 5 minutes using microwave equipment. The reaction mixture was further stirred at 140°C for 5 minutes. The reaction mixture was cooled to room temperature, and then ethyl acetate and saturated sodium hydrogen carbonate aqueous solution were added. The organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel chromatography [ethyl acetate:methanol gradient elution = 100:0 to 90:10]. To the obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 4.7 mg of 4-cyclopropyl-1-ethyl-6-(2-methyl-5-phenyl-1H-imidazol-4-yl)quinolin-2(1H)-one as a white

solid.

$^1\text{H-NMR}$ (CDCl_3) δ : 0.59-0.67 (2H, m), 0.75-0.85 (2H, m), 1.34 (3H, t, $J = 7.2$ Hz), 1.79-1.90 (1H, m), 2.55 (3H, s), 4.33 (2H, q, $J = 7.2$ Hz), 6.36-6.42 (1H, m), 6.72 (1H, s), 7.20-7.41 (4H, m), 7.42-7.55 (1H, m), 7.75-7.79 (1H, m), 8.19-8.26 (1H, m).

5 [0226]

Example 284




A mixture of 27 mg of 1-(4-cyclopropyl-1-ethyl-2-oxo-1,2-dihydroquinolin-6-yl)-2-phenylethane-1,2-dione and 1 mL of polyethylene glycol 400 was stirred at 100°C for 5 minutes using microwave equipment. To the reaction mixture was added 5.2 μL of ethan-1,2-diamine, and the mixture was stirred at 150°C for 7 minutes using microwave equipment. The reaction mixture was further stirred at 160°C for 10 minutes. The reaction mixture was cooled to room temperature, and then ethyl acetate and water were added. The organic layer was fractionated, washed with water and saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel chromatography [hexane:ethyl acetate gradient elution = 50:50 to 0:100]. To the obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 2 mg of 4-cyclopropyl-1-ethyl-6-(3-phenylpyridin-2-yl)quinolin-2(1H)-one as a white solid.

20 $^1\text{H-NMR}$ (CDCl_3) δ : 0.52-0.58 (2H, m), 0.77-0.85 (2H, m), 1.34 (3H, t, $J = 7.2$ Hz), 1.65-1.75 (1H, m), 4.33 (2H, q, $J = 7.2$ Hz), 6.37-6.41 (1H, m), 7.31-7.41 (4H, m), 7.47-7.54 (2H, m), 7.86 (1H, dd, $J = 8.9$ Hz, 2.1 Hz), 8.13 (1H, d, $J = 2.2$ Hz), 8.63 (2H, dd, $J = 6.8$ Hz, 2.4 Hz).

[0227]

Example 285




To a suspension of 134 mg of 1-ethyl-2-oxo-4-(4-oxocyclohexyl)-1,2-dihydroquinoline-6-carboxylic acid in 5 mL of dichloromethane were added 72 mg of 4-chloro-N-methylaniline, 0.27 mL of triethylamine and 216 mg of bis(2-oxo-3-oxazolidinyl)phosphinic acid chloride, and the mixture was stirred at room temperature for 9 hours. To the reaction mixture were added 72 mg of N-methyl-4-chloroaniline, 0.27 mL of triethylamine and 216 mg of bis(2-oxo-3-oxazolidinyl)phosphinic acid chloride, and the mixture was stirred at room temperature for 15 hours. The obtained reaction mixture was subjected to solvent elimination by distillation under reduced pressure. The obtained residue was purified by silica gel column chromatography [ethyl acetate]. To the obtained residue were added diisopropyl ether and ethyl acetate, the solid matter was filtered to obtain 73 mg of N-(4-chlorophenyl)-1-ethyl-N-methyl-2-oxo-4-(4-oxocyclohexyl)-1,2-dihydroquinoline-6-carboxamide as a white solid.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.32 (3H, t, $J = 7.2$ Hz), 1.74-1.88 (2H, m), 1.95-2.04 (2H, m), 2.42-2.61 (4H, m), 3.07-3.17 (1H, m), 3.55 (3H, s), 4.30 (2H, q, $J = 7.1$ Hz), 6.55-6.58 (1H, m), 7.03 (2H, d, $J = 8.8$ Hz), 7.22-7.33 (3H, m), 7.66 (1H, dd, $J = 8.9$ Hz, 2.1 Hz), 7.71 (1H, d, $J = 1.7$ Hz).

[0228]

Example 286




A mixture of 51 mg of N-(4-chlorophenyl)-1-ethyl-N-methyl-2-oxo-4-(4-oxocyclohexyl)-1,2-dihydroquinoline-6-carboxamide, 8.8 mg of sodium borohydride, and 1.0 mL of methanol was stirred at room temperature for 1 hour. To the reaction mixture were added ethyl acetate and 1 mol/L hydrochloric acid. The organic layer was fractionated, and the aqueous layer was extracted with ethyl acetate. The organic layer was combined with the extract, washed with saturated sodium chloride aqueous solution, and then dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography [ethyl acetate:methanol gradient elution = 100:0 to 90:10]. To the obtained residue was added diisopropyl ether, and the solid matter was filtered to obtain 41 mg of N-(4-chlorophenyl)-1-ethyl-4-(4-hydroxycyclohexyl)-N-methyl-2-oxo-1,2-dihydroquinoline-6-carboxamide as a white solid.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.32 (3H, t, $J = 7.2$ Hz), 1.36-1.70 (6H, m), 2.09-2.20 (2H, m), 2.50-2.61 (1H, m), 3.54 (3H, s), 3.60-3.75 (1H, m), 4.29 (2H, q, $J = 7.0$ Hz), 6.51-6.55 (1H, m), 7.02 (2H, d, $J = 8.8$ Hz), 7.20-7.35 (3H, m), 7.55 (1H, d, $J = 2.0$ Hz), 7.72 (1H, dd, $J = 8.8$ Hz, 2.0 Hz).

[0229]

5 Example 287




By the same method as Example 5, from 4-chloro-1-ethyl-N-methyl-N-(4-methylphenyl)-2-oxo-1,2-dihydroquinoline-6-carboxamide was obtained tert-butyl 4-(1-ethyl-6-(methyl(4-methylphenyl)carbamoyl)-2-oxo-1,2-dihydroquinolin-4-yl)-3,6-dihydropyridine-1(2H)-carboxylate.

MS (ESI, m/z): 502($M+H$)

[0230]

Example 288




By the same method as Example 6, from tert-butyl 4-(1-ethyl-6-(methyl(4-methylphenyl)carbamoyl)-2-oxo-1,2-dihydroquinolin-4-yl)-3,6-dihydropyridine-1(2H)-carboxylate was obtained tert-butyl 4-(1-ethyl-6-(methyl(4-methylphenyl)carbamoyl)-2-oxo-1,2-dihydroquinolin-4-yl)piperidine-1-carboxylate.

MS (ESI, m/z): 504($M+H$)

20 [0231]

Example 289




By the same method as Example 7, from tert-butyl 4-(1-ethyl-6-(methyl(4-methylphenyl)carbamoyl)-2-oxo-1,2-dihydroquinolin-4-yl)piperidine-1-carboxylate was

- 5 obtained 1-ethyl-N-methyl-N-(4-methylphenyl)-2-oxo-4-piperidin-4-yl)-1,2-dihydroquinoline-6-carboxamide.

MS (ESI, m/z):404(M+H)

[0232]

Example 290



10


A mixture of 50 mg of 1-ethyl-N-methyl-N-(4-methylphenyl)-2-oxo-4-(piperidin-4-yl)-1,2-dihydroquinoline-6-carboxamide, 20 μ L of acetyl chloride, 0.1 mL of triethylamine, and 1 mL of dichloromethane was stirred at room temperature for 20 minutes. The reaction mixture was purified by silica gel column chromatography [chloroform:methanol gradient

15 elution = 100:0 to 80:20]. To the obtained residue were added hexane and ethyl acetate, and the solid matter was filtered to obtain 13 mg of 4-(1-acetylpiperidin-4-yl)-1-ethyl-N-methyl-N-(4-methylphenyl)-2-oxo-1,2-dihydroquinoline-6-carboxamide as a yellow solid.

- $^1\text{H-NMR}$ (DMSO- D_6) δ : 1.12-1.55 (7H, m), 2.05 (3H, s), 2.20 (3H, s), 2.57-2.69 (1H, m), 2.86-2.99 (1H, m), 3.10-3.23 (1H, m), 3.40 (3H, s), 3.87 (1H, d, $J = 13.9$ Hz), 4.20 (2H, q, $J = 6.9$ Hz),
- 20 4.49 (1H, d, $J = 12.9$ Hz), 6.37 (1H, s), 7.06-7.13 (4H, m), 7.52-7.62 (2H, m), 7.74 (1H, d, $J = 8.8$ Hz).

[0233]

Example 291



A mixture of 30 mg of 4-chloro-1-ethyl-6-(2-phenylpyridin-3-yl)quinolin-2(1H)-one, 10 mg of methylboric acid, 35 mg of tripotassium phosphate, 6 mg of bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II), 0.5 mL of dioxane and 0.2 mL of water was stirred at 120°C for 10 minutes using microwave equipment. The reaction mixture was cooled to room temperature. The reaction mixture was purified by silica gel chromatography [hexane:ethyl acetate gradient elution = 100:0 to 30:70]. To the obtained residue was added hexane, and the solid matter was filtered to obtain 5 mg of 1-ethyl-4-methyl-6-(2-phenylpyridin-3-yl)quinolin-2(1H)-one as a white solid.

¹H-NMR (CDCl₃) δ: 1.34 (3H, t, J = 7.3 Hz), 2.21 (3H, d, J = 1.0 Hz), 4.33 (2H, q, J = 7.3 Hz), 6.55 (1H, d, J = 0.96 Hz), 7.23-7.32 (4H, m), 7.34-7.42 (4H, m), 7.48 (1H, d, J = 1.9 Hz), 7.81 (1H, dd, J = 7.8, 1.7 Hz), 8.73 (1H, dd, J = 4.9, 1.7 Hz).

MS (ESI, m/z):341(M+H)




[0234]

Examples 292-294

Confirming to the procedure described in the specification, the compounds indicated in Table 29 were produced by publicly known reactions, such as condensation, addition, oxidization, reduction, rearrangement, substitution, halogenation, dehydration, hydrolysis or the like, or the appropriate combination of these reactions.

[0235]

[Table 29]

Example No.	Structural Formula	Compound Name	MS
292		1-ethyl-N-(3-fluoro-4-methylphenyl)-N-methyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	436(M+H)
293		N-(3-chloro-4-methylphenyl)-1-ethyl-N-methyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	452(M+H)
294		N-(3,4-dimethylphenyl)-1-ethyl-N-methyl-4-(1-methylpiperidin-4-yl)-2-oxo-1,2-dihydroquinoline-6-carboxamide	432(M+H)

[0236]

Test Example 1

5 (CXCL10 production inhibition test)

0.5 × 10⁴ or 1 × 10⁴ human umbilical vein-derived endothelial cells were suspended in 100 μL of an EGM-2 BulletKit medium (without GA) or a Medium-200 medium supplemented with an EGM-2 SingleQuots kit (without GA), inoculated to a 96-well plate and cultured overnight. 100 μL of a solution of each test compound diluted with the same medium as that used in the inoculation was added to the cultured cells (final concentration of the test compound: 0.1 μmol/L). 1 hour thereafter, TNFα was added thereto at a final concentration of 10 ng/mL. 72 hours after the stimulation, the supernatant was recovered, and the amount of CXCL10 produced in the culture supernatant was determined using an ELISA kit (Peprotech, 900-K39).

15 The rate of inhibition was calculated according to the following expression:

[0237]

Rate of inhibition (%) = 100 - (Amount of CXCL10 produced in the presence of the test compound / Amount of CXCL10 produced in the absence of the test compound) × 100

20 The compounds of Example 2, Example 3, Example 8, Example 11, Example 14, Example 16, Example 17, Examples 245 to 248, Example 250, Examples 252 to 276, Examples 278 to 281, Example 282a, Example 282b, Example 283, Example 284, Example 286 and

Examples 290 to 294 inhibited 50% or more of CXCL10 production at 0.1 $\mu\text{mol/L}$.

The compound of the present invention exhibited an excellent CXCL10 production inhibitory effect.

5 Industrial Applicability


[0238]

The compound represented by the general formula [1] or the salt thereof has an excellent CXCL10 inhibitory activity and is useful as an agent for treatment such as prophylaxis and/or therapy of a disease involving the overproduction of CXCL10.

CLAIMS

[Claim 1]

A compound represented by the general formula [1] or a salt thereof:



wherein

R^1 represents an optionally substituted C_{1-6} alkyl group;

R^2 represents a hydrogen atom, a halogen atom or an optionally substituted C_{1-6} alkyl group;

R^3 represents a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{3-8} cycloalkyl group, an optionally substituted C_{4-8} cycloalkenyl group, an optionally substituted aryl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted C_{1-6} alkylamino group, an optionally substituted $di(C_{1-6} \text{ alkyl})$ amino group or an optionally substituted heterocyclic group;

Z^1 , Z^2 and Z^3 are the same or different and each represent a nitrogen atom or a group represented by the general formula CR^5 wherein R^5 represents a hydrogen atom, a halogen atom or an optionally substituted C_{1-6} alkyl group;

X^1 represents

(1) a group represented by the general formula $C(=O)N(R^6)$ wherein the carbon atom is bonded to ring A, and R^6 represents a hydrogen atom, an amino-protective group or an optionally substituted C_{1-6} alkyl group,

(2) a group represented by the general formula $N(R^7)C(=O)$ wherein the nitrogen atom is bonded to ring A, and R^7 represents a hydrogen atom, an amino-protective group or an optionally substituted C_{1-6} alkyl group; or R^7 represents, together with one substituent R^4 on ring A, an optionally substituted C_{2-4} alkylene group, a group represented by the general formula $O-Y^1$ wherein the oxygen atom is bonded to ring A, and Y^1 represents an optionally substituted C_{1-3} alkylene group, a group represented by the general formula $S(O)_n-Y^2$ wherein the sulfur atom is bonded to ring A, Y^2 represents an optionally substituted C_{1-3} alkylene group, and n represents an integer of 0 to 2, or a group represented by the general formula $N(R^8)-Y^3$ wherein the nitrogen atom is bonded to ring A, Y^3 represents an optionally substituted C_{1-3} alkylene group, and R^8 represents a hydrogen atom, an amino-protective group, an optionally substituted C_{1-6} alkyl

group or an optionally substituted aryl group,

(3) an optionally substituted divalent cyclic hydrocarbon group resulting from the removal of one hydrogen atom from each of two adjacent atoms or

(4) an optionally substituted divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms;

ring A represents a cyclic hydrocarbon group or a heterocyclic group;

m number of R^4 are the same or different and each represent a halogen atom, a cyano group, a nitro group, an amino-protective group, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{3-8} cycloalkyl group, an optionally substituted C_{4-8} cycloalkenyl group, an optionally substituted aryl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted aryloxy group, an optionally substituted C_{1-6} alkylamino group, an optionally substituted di(C_{1-6} alkyl)amino group, an optionally substituted arylamino group, an optionally substituted carbamoyl group, an optionally substituted sulfamoyl group, an optionally substituted C_{1-6} alkylthio group, an optionally substituted arylthio group, an optionally substituted C_{1-6} alkylsulfonyl group, an optionally substituted arylsulfonyl group, an optionally substituted heterocyclic group, an optionally protected amino group, an optionally protected hydroxyl group, an optionally protected carboxyl group,

an optionally substituted C_{2-5} alkylene group formed together by two adjacent R^4 ,

an optionally substituted C_{2-4} alkylene group formed together by one R^4 and R^7 ,

a group represented by the general formula $O-Y^1$ formed together by one R^4 and R^7 wherein the oxygen atom is bonded to ring A, and Y^1 represents an optionally substituted C_{1-3} alkylene group,

a group represented by the general formula $S(O)_n-Y^2$ formed together by one R^4 and R^7 wherein the sulfur atom is bonded to ring A, Y^2 represents an optionally substituted C_{1-3} alkylene group, and n represents an integer of 0 to 2, or

a group represented by the general formula $N(R^8)-Y^3$ formed together by one R^4 and R^7 wherein the nitrogen atom is bonded to ring A, Y^3 represents an optionally substituted C_{1-3} alkylene group, and R^8 represents a hydrogen atom, an amino-protective group, an optionally substituted C_{1-6} alkyl group or an optionally substituted aryl group; and

m represents an integer of 0 to 5.

[Claim 2]

The compound according to claim 1 or a salt thereof, wherein

R^2 is a hydrogen atom or an optionally substituted C_{1-6} alkyl group; and

each of Z^1 , Z^2 and Z^3 is CH.

[Claim 3]

The compound according to claim 1 or 2 or a salt thereof, wherein

R^1 is a C_{1-3} alkyl group, and

R^3 is an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{3-8} cycloalkyl group, an optionally substituted aryl group or an optionally substituted heterocyclic group.

[Claim 4]

The compound according to any one of claims 1 to 3 or a salt thereof, wherein

ring A is a cyclic hydrocarbon group.

[Claim 5]

The compound according to any one of claims 1 to 4 or a salt thereof, wherein X^1

is a group represented by the general formula $C(=O)N(R^6)$ wherein the carbon atom is bonded to ring A, and R^6 represents a hydrogen atom, an amino-protective group or an optionally substituted C_{1-6} alkyl group.

[Claim 6]

The compound according to any one of claims 1 to 4 or a salt thereof, wherein

X^1 is a group represented by the general formula $C(=O)N(R^{6a})$ wherein the carbon atom is bonded to ring A, and R^{6a} represents a hydrogen atom or an optionally substituted C_{1-3} alkyl group;

m number of R^4 are the same or different and each are a halogen atom, an optionally substituted C_{1-3} alkyl group, an optionally substituted C_{1-3} alkylamino group or an optionally protected amino group; and

m is an integer of 0 to 2.

[Claim 7]

The compound according to any one of claims 1 to 4 or a salt thereof, wherein

R^3 is an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{3-8} cycloalkyl group or an optionally substituted heterocyclic group;

X^1 is a group represented by the general formula $C(=O)N(R^{6b})$ wherein the carbon atom is bonded to ring A, and R^{6b} represents an optionally substituted C_{1-3} alkyl group;

m number of R^4 are the same or different and each are a halogen atom or an optionally substituted C_{1-3} alkyl group; and

m is an integer of 0 to 2.

[Claim 8]

The compound according to claim 7 or a salt thereof, wherein

m number of R^4 are the same or different and each are a halogen atom; and
m is an integer of 0 to 2.

[Claim 9]

The compound according to any one of claims 1 to 4 or a salt thereof, wherein X^1 is a group represented by the general formula $N(R^7)C(=O)$ wherein the nitrogen atom is bonded to ring A, and R^7 represents a hydrogen atom, an amino-protective group or an optionally substituted C_{1-6} alkyl group; or R^7 represents, together with one substituent R^4 on ring A, an optionally substituted C_{2-4} alkylene group, a group represented by the general formula $O-Y^1$ wherein the oxygen atom is bonded to ring A, and Y^1 represents an optionally substituted C_{1-3} alkylene group, a group represented by the general formula $S(O)_n-Y^2$ wherein the sulfur atom is bonded to ring A, Y^2 represents an optionally substituted C_{1-3} alkylene group, and n represents an integer of 0 to 2, or a group represented by the general formula $N(R^8)-Y^3$ wherein the nitrogen atom is bonded to ring A, Y^3 represents an optionally substituted C_{1-3} alkylene group, and R^8 represents a hydrogen atom, an amino-protective group, an optionally substituted C_{1-6} alkyl group or an optionally substituted aryl group.

[Claim 10]

The compound according to any one of claims 1 to 4 or a salt thereof, wherein R^3 is an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{3-8} cycloalkyl group or an optionally substituted heterocyclic group;
 X^1 is a group represented by the general formula $N(R^{7a})C(=O)$ wherein the nitrogen atom is bonded to ring A, and R^{7a} represents an optionally substituted C_{1-3} alkyl group, or R^{7a} represents, together with one substituent R^4 on ring A, an optionally substituted C_{2-3} alkylene group or a group represented by the general formula $O-Y^{1a}$ wherein the oxygen atom is bonded to ring A, and Y^{1a} represents an ethylene group;
m number of R^4 are the same or different and each are a halogen atom, an optionally substituted C_{1-3} alkyl group, an optionally substituted C_{1-3} alkylsulfonyl group,
an optionally substituted C_{2-3} alkylene group formed together by two adjacent R^4 ,
an optionally substituted C_{2-3} alkylene group formed together by one R^4 and R^{7a} , or
a group represented by the general formula $O-Y^{1a}$ formed together by one R^4 and R^{7a} wherein the oxygen atom is bonded to ring A, and Y^{1a} represents an ethylene group; and
m is an integer of 0 to 2.

[Claim 11]

The compound according to any one of claims 1 to 4 or a salt thereof, wherein R^3 is an optionally substituted heterocyclic group;

X^1 is a group represented by the general formula $N(R^{7b})C(=O)$ wherein the nitrogen atom is bonded to ring A, and R^{7b} represents an optionally substituted C_{1-3} alkyl group, or R^{7b} represents, together with one substituent R^4 on ring A, an optionally substituted C_{2-3} alkylene group; m number of R^4 are the same or different and each are an optionally substituted C_{1-3} alkyl group, an optionally substituted C_{2-3} alkylene group formed together by two adjacent R^4 , or an optionally substituted C_{2-3} alkylene group formed together by one R^4 and R^{7b} ; and m is an integer of 0 to 2.

[Claim 12]

The compound according to claim 11 or a salt thereof, wherein

X^1 is a group represented by the general formula $N(R^{7b})C(=O)$ wherein the nitrogen atom is bonded to ring A, and R^{7b} represents an optionally substituted C_{1-3} alkyl group, or R^{7b} represents, together with one substituent R^4 on ring A, an optionally substituted C_{2-3} alkylene group; m number of R^4 are the same or different and each are an optionally substituted C_{1-3} alkyl group, or an optionally substituted C_{2-3} alkylene group formed together by one R^4 and R^{7b} ; and m is an integer of 0 to 2.

[Claim 13]

The compound according to any one of claims 1 to 4 or a salt thereof, wherein X^1 is an optionally substituted divalent cyclic hydrocarbon group resulting from the removal of one hydrogen atom from each of two adjacent atoms or an optionally substituted divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms.

[Claim 14]

The compound according to any one of claims 1 to 4 or a salt thereof, wherein R^3 is an optionally substituted C_{1-6} alkyl group or an optionally substituted heterocyclic group; X^1 is an optionally substituted divalent cyclic hydrocarbon group resulting from the removal of one hydrogen atom from each of two adjacent atoms or an optionally substituted divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms; m number of R^4 are the same or different and each are a halogen atom; and m is an integer of 0 to 2.

[Claim 15]

The compound according to claim 14 or a salt thereof, wherein

R^3 is an optionally substituted heterocyclic group; and

X¹ is an optionally substituted divalent heterocyclic group resulting from the removal of one hydrogen atom from each of two adjacent atoms.

[Claim 16]

A pharmaceutical composition comprising a compound according to any one of claims 1 to 15 or a salt thereof.

[Claim 17]

An agent for treatment of an immune disease, comprising a compound according to any one of claims 1 to 15 or a salt thereof.