

Fluorinated Ligands for Targeting Peripheral Benzodiazepine Receptors

Technical Field

The present invention relates to radiolabelled and non-radiolabelled heterocyclic compounds which bind to peripheral benzodiazepine receptors and are useful for imaging such receptors and providing therapeutic treatment of conditions associated with PBR receptors.

Background of the Invention

The peripheral benzodiazepine receptors, which are also commonly referred to as the peripheral benzodiazepine binding sites, mitochondrial benzodiazepine receptors or ω -3-receptors, are distinct from the central benzodiazepine receptors in their pharmacology, subcellular location and structural requirements. Peripheral benzodiazepine receptors are predominantly found in the peripheral organs such as kidney, heart, adrenal cortex, testis, ovaries, plasma (platelets) as well as in the glial cells and olfactory bulbs in the brain. It has also been reported that peripheral benzodiazepine receptor density is higher in tumours, such as breast, prostate, glioma, ovarian and colon carcinoma, than in corresponding normal tissue. Recently high concentrations of peripheral benzodiazepine receptors has been reported in Dunning rat prostate tumours compared to the normal ventral or dorsolateral prostate.

The peripheral benzodiazepine receptors appear to be associated (but not exclusively) with the outer mitochondrial membrane in many tissues where they are modulated by hormones and drugs and reflect the effects of emotional stress, and hypertension. Peripheral benzodiazepine receptors in the brain have been investigated for use as markers of neurodegeneration and inflammation including Huntington's disease, Alzheimer's disease, multiple sclerosis, anxiety, stress, emotional disturbances, cognitive impairment, stroke, and cerebral ischemia. The Peripheral Benzodiazepine Receptor has also been implicated in apoptotic processes.

PBR ligands themselves modulate various cellular processes including heme transport, cholesterol and steroid synthesis, apoptosis and calcium transport among others. They are also involved in cellular proliferation and differentiation.

Several classes of ligands have been shown to exhibit high affinity binding to the peripheral benzodiazepine receptor, the most widely investigated being the benzodiazepine Ro 5-4864 (7-chloro-5-(4-chlorophenyl)-1,3-dihydro-1-methyl-2H-1,4-benzodiazepin-2-one) and the isoquinoline PK-11195 (1-(2-chlorophenyl)-N-methyl-N-(1-methylpropyl)-3-isoquinolinecarboxamide). Labelled with ^{11}C , ^{14}C and ^3H , these ligands have been used to map peripheral benzodiazepine receptors in the human brain and heart by *in vivo* and *in vitro* means. Enhanced uptake of [^3H]PK-11195 has been reported in a variety of tumour cells including breast, ovarian, prostate, adrenal, brain and colon. Radiolabelling with suitable levels of radioactive isotopes such as carbon-11, fluorine-18

and iodine-123/124 may be used firstly to diagnose these tumours and subsequently to treat them with therapeutic doses (for instance using ^{123}I , ^{125}I or ^{131}I). Furthermore, recent work has also revealed the existence of several binding domains which differ in affinity for isoquinoline and benzodiazepine ligands in different organs and species.


Various 2-aryl substituted imidazo[1,2-a]pyridines imidazopyridazines and aryl-pyrazolo pyrimidines having anxiolytic, hypnotic, anticonvulsant, analgesic and other properties have been reported (Almirante L. *et al.*, *J. Med. Chem.* **12** 122-126 (1969); Langer S.Z. *et al.*, *Pharmacol. Biochem. Behav.* **29** 763-766 (1988); Langer S.Z. *et al.*, *Pharmacopsychiatry* **23** 103-107 (1990); Bourguignon, J.-J., "Endogenous and Synthetic Ligands of Mitochondrial Benzodiazepine Receptors: Structure Affinity Relationships" in Giesen-Grouse, E. ed. *Peripheral Benzodiazepine Receptors*, Academic Press, London (1993)). For example ^{123}I labelled 6-methyl-2-(4'-iodophenyl)imidazo[1,2-a]pyridine-3-(N,N-dimethyl)-acetamide has been reported as a potential tracer for the study of the peripheral benzodiazepine receptor using SPECT (Katsifis A. *et al.*, *J. Lab. Comp. Radiopharm.* **40** 620-622 (1997)).

In published international application WO 99/51594 it was discovered that certain 2-(iodophenyl)-imidazo[1,2-a]pyridines having an electronegative substituent, especially halogen, in the pyridine nucleus exhibit strong binding to peripheral benzodiazepine receptors and much weaker binding to central benzodiazepine receptors.

The present invention relates to a class of fluorinated compounds for use as PBR pharmacological agents in the treatment of various disorders such as neuroinflammation, neurodegeneration, stroke, apoptosis and other disorders. The invention also relates to compounds labelled with the PET radionuclide ^{18}F that are useful for PET imaging using positron emission tomography. Appropriately radiolabelled compounds of the present invention are clinically useful in PET scanning, for example to detect those cancers which express a high density of the peripheral benzodiazepine receptors and/or to detect, or non-invasively diagnose, neurodegenerative disorders. The compounds of the present invention are also useful for the treatment of disorders characterised by an abnormal density of peripheral benzodiazepine receptors, such as neurodegenerative disorders and tumours. The present invention also relates to intermediates of the fluorinated compounds.

Summary of the Invention

In a first aspect the present invention refers to a fluorinated compound of formula (I):



wherein,

D, G, and L are independently selected from the group consisting of: CH, C and N, and J and M are independently selected from the group consisting of C and N provided that one of J and M is C and the other is N, wherein at least two of D, G, M, J and L are N;

X is CH₂;

Y is absent;

Z is NR₁R₂;

R₁ and R₂ are independently selected from the group consisting of: hydrogen, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, aryl and heteroaryl, each being optionally substituted with one or more of the following substituents: halogen, an C₁-C₆ alkyl;

or R₁ and R₂, together with the nitrogen to which they are attached, form a heterocyclic ring having between 3 and 7 ring members, optionally substituted with one or more of the following substituents: halogen and C₁-C₆ alkyl;

R₃ is selected from the group consisting of: halogen, C₁-C₁₀ alkyl and O-(C₁-C₁₀ alkyl), wherein the C₁-C₁₀ alkyl group is optionally substituted;

E is an aryl group or a heteroaryl group, wherein each is optionally substituted with one or more fluoro substituents, or with one or more of the following substituents: C₁-C₆ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, QC₁-C₁₀ alkyl, QC₂-C₁₀ alkenyl, QC₂-C₁₀ alkynyl, Q(CH₂)_p-Q-(CH₂)_qCH₃ or Q(CH₂)_p-Q-(CH₂)_q-Q-(CH₂)_rCH₃, each of which is optionally substituted with one or more fluoro substituents, and wherein p, q and r are, independently, integers between 1 and 3, and wherein Q is selected from the group consisting of: NH, O and S,

m is a number between 0 and 3;

with the proviso that:


R₃ is a fluoro substituent, or

the group E comprises a fluoro substituent, or

the group Z comprises a fluoro substituent,

with the further proviso that E is not 4-fluorophenyl,

or a compound of formula (1a):



wherein,

D, G and L are independently selected from the group consisting of: CH, C and N, and J and M are independently selected from the group consisting of C and N provided that one of J and M is C and the other is N, wherein at least two of D, G, M, J and L are N;

X is CH₂;

Y is absent;

Z is NR₁R₂;

R₁ and R₂ are independently selected from the group consisting of: hydrogen, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, (CH₂)_naryl, aryl and heteroaryl, each of which is optionally substituted with one or more of the following substituents: chloro, bromo, iodo, C₁-C₆ alkyl and hydroxy;

or R₁ and R₂, together with the nitrogen to which they are attached, form a heterocyclic ring having between 3 and 7 ring members, which is optionally substituted with one or more of the following substituents: chloro, bromo, iodo and C₁-C₆ alkyl;

R₃ is selected from the group consisting of: chloro, bromo, iodo, C₁-C₁₀ alkyl and O-(C₁-C₁₀ alkyl), wherein the C₁-C₁₀ alkyl group is optionally substituted;

E is an aryl group or a heteroaryl group, wherein each is optionally substituted with a chloro, bromo or iodo substituent, and/or with one or more of the following substituents: C₁-C₆ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, QC₁-C₁₀ alkyl, QC₁-C₁₀ alkenyl, QC₂-C₁₀ alkynyl, Q(CH₂)_p-Q-(CH₂)_qCH₃ or Q(CH₂)_p-Q-(CH₂)_q-Q-(CH₂)_rCH₃, each of which is optionally substituted with one or more chloro, bromo, iodo or hydroxy substituents, and wherein p, q and r are, independently, integers between 1 and 3, and wherein Q is selected from the group consisting of: NH, O and S, wherein when E is phenyl, E does not have an iodo substituent attached directly to it in the 4 position;

m is a number between 0 and 3;


wherein at least one of Z, E or R₃ comprises iodine,

with the proviso that E is not 4-iodophenyl,

and with the proviso that said compound is not a compound of formula (1a) as defined in the following table:

D	G	J	L	M	R ₃	E	Z
CH	CH	C	N	N	6-Cl	4-chlorophenyl	NHCH ₂ CF ₃
CH	CH	C	N	N	6-CH ₃	4-chlorophenyl	NHCH ₂ CF ₃

In another embodiment the disclosure provides a fluorinated compound of



wherein,

D, G, and L are independently selected from the group consisting of: CH, C and N, and J and M are independently selected from the group consisting of C and N provided that at least one of J and M is C, wherein at least two of D, G, M, J and L are N;

X is selected from the group consisting of: O, NH, $(CH_2)_n$ and S;

Y is absent, or is selected from the group consisting of: O, NH, $(CH_2)_n$, and S;

Z is selected from the group consisting of: NR_1R_2 and aryl;

R_1 and R_2 are independently selected from the group consisting of: hydrogen, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, aryl and heteroaryl, each of which may optionally be substituted with one or more of the following substituents: halogen and C_1 - C_6 alkyl;

or R_1 and R_2 , together with the nitrogen to which they are attached, form a heterocyclic ring having between 3 and 7 ring members, which may optionally be substituted with one or more of the following substituents: halogen and C_1 - C_6 alkyl;

R_3 is selected from the group consisting of: halogen, C_1 - C_{10} alkyl and O- $(C_1$ - C_{10} alkyl), wherein the C_1 - C_{10} alkyl group is optionally substituted;

E is an aryl group or a heteroaryl group, wherein each group is optionally substituted with one or more fluoro substituents, or with one or more of the following substituents: C_1 - C_6 alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, QC_1 - C_{10} alkyl, QC_2 - C_{10} alkenyl, QC_2 - C_{10} alkynyl, $Q(CH_2)_p-Q-(CH_2)_qCH_3$ or $Q(CH_2)_p-Q-(CH_2)_q-Q-(CH_2)_rCH_3$, each of which may be optionally substituted with one or more fluoro substituents, and wherein p, q and r are, independently, integers between 1 and 3, and wherein Q is selected from the group consisting of: NH, O and S;

m is a number between 0 and 3 (e.g. 0, 1, 2 or 3) wherein if m is 2 or 3, the R_3 groups may be the same or may be different;

n is a number between 1 and 4, e.g. 1, 2, 3 or 4 (wherein n in X may be the same as or different to n in Y);

with the proviso that R_3 is a fluoro substituent, or

the group E comprises a fluoro substituent, or

the group Z comprises a fluoro substituent, with the further proviso that E is not 4-fluorophenyl.

The further proviso may be that E is not -PhF.

The disclosure provides a fluorinated compound of formula (I) wherein:

D, G, and L are independently selected from the group consisting of: CH, C and N, and J and M are independently selected from the group consisting of C and N provided that at least one of J and M is C, wherein at least two of D, G, M, J and L are N;

X is independently selected from the group consisting of: O, NH, $(CH_2)_n$ and S;

5 Y is absent, or is independently selected from the group consisting of: O, NH and $(CH_2)_n$, and S;

Z is selected from the group consisting of: NR_1R_2 and aryl;

R_1 and R_2 are independently selected from the group consisting of: hydrogen, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, aryl and heteroaryl, each of which may optionally be substituted with
10 one or more of the following substituents: halogen and C_1 - C_6 alkyl;

or R_1 and R_2 , together with the nitrogen to which they are attached, form a heterocyclic ring having between 3 and 7 ring members, which may optionally be substituted with one or more of the following substituents: halogen and C_1 - C_6 alkyl;

R_3 is selected from the group consisting of: halogen and C_1 - C_{10} alkyl;

15 E is an aryl group or a heteroaryl group, which may be optionally substituted with one or more fluoro substituents, or one or more of the following substituents: C_1 - C_6 alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, QC_1 - C_{10} alkyl, QC_2 - C_{10} alkenyl, QC_2 - C_{10} alkynyl, $Q(CH_2)_p-Q(CH_2)_qCH_3$ or $Q(CH_2)_p-Q(CH_2)_r-Q(CH_2)_rCH_3$, each of which may be optionally substituted with one or more fluoro substituents, and wherein p, q and r are integers between 1 and 3, and wherein Q is independently
20 selected from the group consisting of: NH, O and S;

m is a number between 0 and 3;

n is a number between 1 and 4;

with the proviso that R_3 is a fluoro substituent, or

the group E comprises a fluoro substituent, or

25 the group Z comprises a fluoro substituent, with the further proviso that E may not be -PhF.

The term fluoro substituent(s) may refer to ^{19}F (also referred to as F) or may refer to ^{18}F , or may refer to a mixture of the two.

E may be an aryl group or a heteroaryl group, which may be optionally substituted with one or more fluoro substituents selected from ^{19}F , ^{18}F and a combination thereof, and/or one or more of
30 the following substituents: C_1 - C_6 alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, QC_1 - C_{10} alkyl, QC_2 - C_{10} alkenyl, QC_2 - C_{10} alkynyl, $Q(CH_2)_p-Q(CH_2)_qCH_3$ or $Q(CH_2)_p-Q(CH_2)_r-Q(CH_2)_rCH_3$, each of which may be optionally substituted with one or more fluoro substituents selected from ^{19}F , ^{18}F and a combination thereof, and wherein p, q and r are, independently, integers between 1 and 3, and wherein Q is selected from the group consisting of: NH, O and S.

E may be an aryl group or a heteroaryl group, which may be optionally substituted with ^{18}F , or one or more of the following substituents: $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_{10}$ alkenyl, $\text{C}_2\text{-C}_{10}$ alkynyl, $\text{QC}_1\text{-C}_{10}$ alkyl, $\text{QC}_2\text{-C}_{10}$ alkenyl, $\text{QC}_2\text{-C}_{10}$ alkynyl, $\text{Q}(\text{CH}_2)_p\text{-Q}(\text{CH}_2)_q\text{CH}_3$ or $\text{Q}(\text{CH}_2)_p\text{-Q}(\text{CH}_2)_q\text{-Q}(\text{CH}_2)_r\text{CH}_3$, each of which may be optionally substituted with ^{18}F , and wherein p, q and r are, independently, integers between 1 and 3, and wherein Q is selected from the group consisting of: NH, O and S.

E may be an aryl group or a heteroaryl group, which is substituted with ^{18}F .

E may be an aryl group or a heteroaryl group, which is substituted with ^{19}F .

One embodiment may be with the proviso that R_3 is a fluoro substituent which is ^{19}F or ^{18}F , or the group E comprises a fluoro substituent which is ^{19}F or ^{18}F or a combination thereof, or the group Z comprises a fluoro substituent which is ^{19}F or ^{18}F or a combination thereof, with the further proviso that E may not be $-\text{PhF}$ (or may not be 4-fluorophenyl).

R_3 may be selected from the group consisting of: ^{19}F and ^{18}F . R_3 may be ^{18}F . R_3 may be ^{19}F .

^{18}F is a radioactive fluoro substituent. ^{19}F , which is alternatively referred to as F throughout the specification, is a non-radioactive fluoro substituent.

In an embodiment of the invention Q may be oxygen.

In another embodiment of the invention, m is 1 or 2, and n is 1 or 2, optionally 1.

In a further embodiment of the invention, p may be 1 or 2, q and r may be 1, and Q may be oxygen.

In some embodiments, L is N. In some embodiments one of M and J is N.

In one embodiment, M and L are N, D and G are CH and J is C. In another embodiment, D, M and L are N, J is C and G is CH. In an alternative embodiment, D, J and L are N, M is C and G is CH.

R_3 may be attached to a carbon atom.

R_3 may be optionally substituted with one or more halogens, such as fluorine, chlorine, bromine and/or iodine.

R_3 may be selected from the group consisting of: halogen and $\text{C}_1\text{-C}_6$ alkyl, optionally substituted (e.g. with one or more halogens such as fluorine, chlorine, bromine and/or iodine). R_3 may be methyl. It may be methoxy. It may be trifluoromethyl. It may be trifluoromethoxy.

In another embodiment, R_3 may be selected from the group consisting of chloro and $\text{C}_1\text{-C}_3$ alkyl. In a further embodiment, R_3 may be chloro or methyl and m may be 1 or 2. Alternatively, R_3 may be chloro and m may be 1 or 2, or R_3 may be fluoro and m may be 1 or 2. R_3 may be chloro and m may be 1 or 2, or R_3 may be ^{18}F fluoro and m may be 1 or 2. If $m > 1$, the R_3 groups may be the same or they may be different. For example, if $m=2$, the R_3 groups may be both methyl, or one may be methyl and the other trifluoromethyl, or one may be F and the other trifluoromethyl.

R_3 may be chloro and m may be 1 or 2, or R_3 may be ^{19}F fluoro and m may be 1 or 2.

In one embodiment, Y is absent and X is $(\text{CH}_2)_n$, wherein n is 1 or 2.

In another embodiment, X and Y may be selected from the group consisting of: NH and $(\text{CH}_2)_n$, wherein n is 1.

5 Y may be $(\text{CH}_2)_n$ and X may be NH , wherein n is 1 or 2.

In one embodiment, $-\text{X}-\text{Y}-\text{C}(\text{O})-\text{Z}$ may have a structure selected from the group consisting of: $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{Z}$, wherein n is 1 or 2, or $-(\text{CH}_2)_n-\text{NH}-\text{C}(\text{O})-\text{Z}$, wherein n is 1 or 2.

Z may be NR_1R_2 or phenyl.

In one embodiment Z is NR_1R_2 .

10 R_1 and R_2 may be independently selected from the group consisting of: hydrogen $\text{C}_1\text{-C}_6$ alkyl, phenyl, pyridyl, pyrimidinyl, pyridazinyl and where R_1 and R_2 , together with the nitrogen to which they are attached, form a heterocyclic ring having between 4 and 6 ring members, wherein the $\text{C}_1\text{-C}_6$ alkyl group, the phenyl group, the pyridyl group, the pyrimidinyl group, the pyridazinyl group or the heterocyclic ring may optionally be substituted with between one and three fluoro substituents.

15 The heterocyclic ring may be a pyrrolidine ring.

R_1 and R_2 may be independently selected from the group consisting of: hydrogen $\text{C}_1\text{-C}_6$ alkyl, phenyl, pyridyl, pyrimidinyl, pyridazinyl and where R_1 and R_2 , together with the nitrogen to which they are attached, form a heterocyclic ring having between 4 and 6 ring members, wherein the $\text{C}_1\text{-C}_6$ alkyl group, the phenyl group, the pyridyl group, the pyrimidinyl group, the pyridazinyl group or the heterocyclic ring may optionally be substituted with between one or more fluoro substituents selected from the group consisting of 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent. The heterocyclic ring may be a pyrrolidine ring.

20 R_1 and R_2 may be independently selected from the group consisting of: hydrogen $\text{C}_1\text{-C}_6$ alkyl, phenyl, pyridyl, and where R_1 and R_2 , together with the nitrogen to which they are attached, form a heterocyclic ring having between 4 and 6 ring members, wherein the heterocyclic ring, the $\text{C}_1\text{-C}_6$ alkyl group, the phenyl group or the pyridyl group may optionally be substituted with between 1 and 3 fluoro substituents (e.g. 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent).

30 In another embodiment, R_1 and R_2 may be independently selected from the group consisting of: hydrogen, phenyl, pyridyl, methyl, ethyl, propyl, isopropyl, butyl, *t*-butyl, and pentyl, wherein each of these substituents may optionally be substituted with between 1 and 3 fluoro substituents (e.g. 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent).

In an alternative embodiment, R₁ and R₂ may be selected from the group consisting of: hydrogen, phenyl, pyridyl, methyl, ethyl, isopropyl and wherein R₁ and R₂ together with the nitrogen to which they are attached, form a pyrrolidine ring, and wherein the phenyl, pyridyl, methyl, ethyl, isopropyl or pyrrolidine ring may optionally be substituted with between 1 and 3 fluoro substituents (e.g. 1, 2, or 3 ¹⁹F substituents or an ¹⁸F substituent or a combination of a ¹⁹F substituent and an ¹⁸F substituent or a combination of two ¹⁹F substituents and an ¹⁸F substituent).

When E is phenyl, E may optionally not have a halogen attached directly to it in the 4 position. In particular E may not have a chloro, bromo, fluoro or iodo substituent attached directly to it in the 4 position. E may be phenyl or substituted phenyl which is not substituted by a fluoro or iodo substituent in the 4 position. If E is phenyl, it may in some embodiments have no halogen attached directly to it.

E may be a pyridyl group, a pyrimidinyl group or a pyridazinyl group, each of which may optionally be substituted with one or more fluoro substituents (e.g. 1, 2, or 3 ¹⁹F substituents or an ¹⁸F substituent or a combination of a ¹⁹F substituent and an ¹⁸F substituent or a combination of two ¹⁹F substituents and an ¹⁸F substituent), and/or substituted with one or more of the following substituents: C₁-C₆ alkyl or C₁-C₁₀ alkoxy, each of which may optionally be substituted with one or more fluoro substituents (e.g. 1, 2, or 3 ¹⁹F substituents or an ¹⁸F substituent or a combination of a ¹⁹F substituent and an ¹⁸F substituent or a combination of two ¹⁹F substituents and an ¹⁸F substituent).

E may be a pyridyl group, a pyrimidinyl group or a pyridazinyl group, each of which may optionally be substituted with between one and three fluoro substituents (e.g. 1, 2, or 3 ¹⁹F substituents or an ¹⁸F substituent or a combination of a ¹⁹F substituent and an ¹⁸F substituent or a combination of two ¹⁹F substituents and an ¹⁸F substituent), or substituted with one of the following substituents: C₁-C₆ alkyl or C₁-C₆ alkoxy, either of which may optionally be substituted with between one and three fluoro substituents (e.g. 1, 2, or 3 ¹⁹F substituents or an ¹⁸F substituent or a combination of a ¹⁹F substituent and an ¹⁸F substituent or a combination of two ¹⁹F substituents and an ¹⁸F substituent).

E may be a pyridyl group, a pyrimidinyl group or a pyridazinyl group, each of which may optionally be substituted with between one and three fluoro substituents (e.g. 1, 2, or 3 ¹⁹F substituents or an ¹⁸F substituent or a combination of a ¹⁹F substituent and an ¹⁸F substituent or a combination of two ¹⁹F substituents and an ¹⁸F substituent).

E may be a phenyl group, a pyridyl group, a pyrimidinyl group or a pyridazinyl group, each of which may optionally be substituted with one or more of the following substituents: C₁-C₆ alkyl or C₁-C₁₀ alkoxy, each of which may optionally be substituted with one or more fluoro substituents

(e.g. 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent).

E may be a phenyl group, a pyridyl group, a pyrimidinyl group or a pyridazinyl group, each of which may optionally be substituted with one of the following substituents: $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_1\text{-C}_{10}$ alkoxy, either of which may optionally be substituted with between one and three fluoro substituents (e.g. 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent).

E may be a phenyl group, a pyridyl group, a pyrimidinyl group or a pyridazinyl group, each of which may optionally be substituted with one of the following substituents: $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_1\text{-C}_{10}$ alkoxy, either of which may optionally be substituted with between one and three fluoro substituents (e.g. 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent), or E may be a pyridyl group, a pyrimidinyl group or a pyridazinyl group, each of which may optionally be substituted with between one and three fluoro substituents (e.g. 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent).

E may be a phenyl group which is substituted with one or more $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_1\text{-C}_{10}$ alkoxy groups, each of which may optionally be substituted with one or more fluoro substituents (e.g. 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent).

E may be a phenyl group which is substituted with one or more $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_1\text{-C}_{10}$ alkoxy groups, each of which may optionally be substituted with between one to three fluoro substituents (e.g. 1, 2, or 3 fluoro substituents or 1, 2 or 3 ^{18}F substituents or any combination of 2 or 3 F and ^{18}F substituents).

E may be a phenyl group which is substituted with one or more $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_1\text{-C}_6$ alkoxy groups, each of which may optionally be substituted with one or more fluoro substituents (e.g. 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent).

E may be a phenyl group which is substituted with one or more $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_1\text{-C}_6$ alkoxy groups, each of which may optionally be substituted with between one and three fluoro substituents (e.g. 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent).

E may be a phenyl group which is substituted with a C₁-C₆ alkyl group or a C₁-C₁₀ alkoxy group, each of which may optionally be substituted with a fluoro substituent (e.g. an ¹⁹F substituent or an ¹⁸F substituent).

5 E may be a phenyl group which is substituted with a C₁-C₆ alkyl group or a C₁-C₆ alkoxy group, each of which may optionally be substituted with a fluoro substituent (e.g. an ¹⁹F substituent or an ¹⁸F substituent).

E may be a phenyl group which is substituted with a C₁-C₆ alkoxy group, which may optionally be substituted with a fluoro substituent (e.g. an ¹⁹F substituent or an ¹⁸F substituent).

10 In an embodiment, R₃ is chloro, m is 1 or 2, E is fluoropyridyl, fluoropyrimidinyl or fluoropyridazinyl, Z is NR₁R₂, wherein R₁ and R₂ are selected from the group consisting of: C₁-C₆ alkyl and phenyl, Y is absent, and X is (CH₂)_n, wherein n is 1 or 2.

In another embodiment, R₃ is chloro, m is 1 or 2, E is ¹⁸fluoropyridyl, ¹⁸fluoropyrimidinyl or ¹⁸fluoropyridazinyl, Z is NR₁R₂, wherein R₁ and R₂ are selected from the group consisting of: C₁-C₆ alkyl and phenyl, Y is absent, and X is (CH₂)_n, wherein n is 1 or 2.

15 In another embodiment, R₃ is chloro, m is 1 or 2, E is ¹⁹fluoropyridyl, ¹⁹fluoropyrimidinyl or ¹⁹fluoropyridazinyl, Z is NR₁R₂, wherein R₁ and R₂ are selected from the group consisting of: C₁-C₆ alkyl and phenyl, Y is absent, and X is (CH₂)_n, wherein n is 1 or 2.

In a further embodiment, R₃ is chloro, m is 1 or 2, E is phenyl substituted with a C₁-C₆ alkoxy group, Z is NR₁R₂, wherein R₁ is C₁-C₆ alkyl and R₂ is fluoropyridyl or fluorophenyl.

20 In a further embodiment, R₃ is chloro, m is 1 or 2, E is phenyl substituted with a C₁-C₆ alkoxy group, Z is NR₁R₂, wherein R₁ is C₁-C₆ alkyl and R₂ is ¹⁸fluoropyridyl or ¹⁸fluorophenyl.

In a further embodiment, R₃ is chloro, m is 1 or 2, E is phenyl substituted with a C₁-C₆ alkoxy group, Z is NR₁R₂, wherein R₁ is C₁-C₆ alkyl and R₂ is ¹⁹fluoropyridyl or ¹⁹fluorophenyl.

25 In one embodiment, R₃ is F, m is 1 and E is phenyl substituted with a C₁-C₆ alkyl group or a C₁-C₆ alkoxy group.

In one embodiment, R₃ is ¹⁸F, m is 1 and E is phenyl substituted with a C₁-C₆ alkyl group or a C₁-C₆ alkoxy group.

In another embodiment, R₃ is F, m is 1 and E is phenyl substituted with a C₁-C₃ alkoxy group or a C₁-C₆ alkyl group.

30 In another embodiment, R₃ is ¹⁸F, m is 1 and E is phenyl substituted with a C₁-C₃ alkoxy group or a C₁-C₆ alkyl group.

In a further embodiment, R₃ is C₁-C₆ alkyl or chloro, m is 1 or 2, Y is absent, X is (CH₂)_n, wherein n is 1 or 2, and E is a phenyl group to which is attached one or more C₁-C₆ alkoxy groups, said C₁-C₆ alkoxy groups comprising one or more fluoro substituents (e.g. 1, 2, or 3 ¹⁹F substituents).

or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent).

In a further embodiment, R_3 is $\text{C}_1\text{-C}_6$ alkyl or chloro, m is 1 or 2 and E is a phenyl group to which is attached one or more $\text{C}_1\text{-C}_6$ alkoxy groups, said $\text{C}_1\text{-C}_6$ alkoxy groups comprising between one and three fluoro substituents (e.g. 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent).

In another embodiment, R_3 is methyl or chloro, m is 1 or 2 and E is a phenyl group to which is attached one or more $\text{C}_1\text{-C}_6$ alkoxy groups, said $\text{C}_1\text{-C}_6$ alkoxy groups comprising one or more fluoro substituents (e.g. 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent).

In another embodiment, R_3 is methyl or chloro, m is 1 or 2 and E is a phenyl group to which is attached one or more $\text{C}_1\text{-C}_6$ alkoxy groups, said $\text{C}_1\text{-C}_6$ alkoxy groups comprising between one and three fluoro substituents (e.g. 1, 2, or 3 ^{19}F substituents or an ^{18}F substituent or a combination of a ^{19}F substituent and an ^{18}F substituent or a combination of two ^{19}F substituents and an ^{18}F substituent).

In another embodiment, R_3 is methyl or chloro, m is 1 or 2 and E is a phenyl group to which is attached a $\text{C}_1\text{-C}_6$ alkoxy group, said $\text{C}_1\text{-C}_6$ alkoxy group comprising a single fluoro substituent (^{19}F or ^{18}F).

In another embodiment, R_3 is methyl or chloro, m is 1 or 2 and E is a phenyl group to which is attached an alkoxy group of the following formula: $-\text{O}(\text{CH}_2)_n\text{F}$, wherein n is a number between 1 and 4.

In another embodiment, R_3 is methyl or chloro, m is 1 or 2 and E is a phenyl group to which is attached an alkoxy group of the following formula: $-\text{O}(\text{CH}_2)_n^{18}\text{F}$, wherein n is a number between 1 and 4.

In one embodiment, the alkoxy group attached to the phenyl group is in the para position.

In a particular embodiment of the disclosure, R_3 may be chloro or $\text{C}_1\text{-C}_6$ alkyl, m may be 1 or 2, G may be CH , D may be CH , M may be N , J may be C , L may be N , Y may be absent, X may be CH_2 or CH_2CH_2 , Z may be NR_1R_2 , wherein R_1 and R_2 are independently selected from hydrogen, pyridyl, phenyl, $\text{C}_1\text{-C}_4$ alkyl, or R_1 and R_2 together with the nitrogen to which they are attached form a pyrrolidine ring, and E may be phenyl group substituted with a single substituent of the formula $-\text{O}(\text{CH}_2)_n\text{F}$, wherein n is a number between 1 and 4.

In a particular embodiment of the disclosure, R_3 may be chloro or $\text{C}_1\text{-C}_6$ alkyl, m may be 1 or 2, G may be CH , D may be CH , M may be N , J may be C , L may be N , Y may be absent, X may be

CH₂ or CH₂CH₂, Z may be NR₁R₂, wherein R₁ and R₂ are independently selected from hydrogen, pyridyl, phenyl, C₁-C₄ alkyl, or R₁ and R₂ together with the nitrogen to which they are attached form a pyrrolidine ring, and E may be phenyl group substituted with a single substituent of the formula -O(CH₂)_n¹⁸F, wherein n is a number between 1 and 4.

5 In another embodiment of the disclosure, R₃ may be methyl, m may be 1 or 2, G may be CH, D may be N, M may be C, J may be N, L may be N, Y may be absent, X may be CH₂ or CH₂CH₂, Z may be NR₁R₂, wherein R₁ and R₂ are independently C₁-C₆ alkyl, and E may be a phenyl group substituted with a single substituent of the formula -O(CH₂)_nF, wherein n is a number between 1 and 4.

10 In a further particular embodiment of the invention, R₃ may be fluoro, m may be 1, G may be CH, D may be N, M may be N, J may be C, L may be N, Y may be absent, X may be CH₂, Z may be NR₁R₂, wherein R₁ and R₂ are independently C₁-C₄ alkyl, and E may be a phenyl group substituted with a C₁-C₃ alkoxy group or a C₁-C₆ alkyl group.

In a further particular embodiment of the disclosure, R₃ may be fluoro, m may be 1, G may be
15 CH, D may be N, M may be N, J may be C, L may be N, Y may be NH, X may be CH₂ or CH₂CH₂, Z may be phenyl, and E may be a phenyl group substituted with one or two C₁-C₆ alkyl groups.

In another particular embodiment of the disclosure, R₃ may be chloro, m may be 1 or 2, G may be CH, D may be CH, M may be N, J may be C, L may be N, Y may be absent, X may be CH₂ or CH₂CH₂, Z may be NR₁R₂, wherein R₁ and R₂ are independently selected from hydrogen, phenyl and C₁-C₄ alkyl, and E may be fluoropyridine, fluoropyridazine or fluoropyrimidine.
20

In another particular embodiment of the disclosure, R₃ may be chloro, m may be 1 or 2, G may be CH, D may be CH, M may be N, J may be C, L may be N, Y may be absent, X may be CH₂ or CH₂CH₂, Z may be NR₁R₂, wherein R₁ and R₂ are independently selected from hydrogen, phenyl and C₁-C₄ alkyl, and E may be ¹⁹F fluoropyridine, ¹⁹F fluoropyridazine or ¹⁹F fluoropyrimidine.

25 In another particular embodiment of the disclosure, R₃ may be chloro, m may be 1 or 2, G may be CH, D may be CH, M may be N, J may be C, L may be N, Y may be absent, X may be CH₂ or CH₂CH₂, Z may be NR₁R₂, wherein R₁ and R₂ are independently selected from hydrogen, phenyl and C₁-C₄ alkyl, and E may be ¹⁸F fluoropyridine, ¹⁸F fluoropyridazine or ¹⁸F fluoropyrimidine.

In another particular embodiment of the disclosure, R₃ may be chloro, m may be 1 or 2, G may be CH, D may be CH, M may be N, J may be C, L may be N, Y may be absent, X may be CH₂ or CH₂CH₂, Z may be NR₁R₂, wherein R₁ and R₂ are independently selected from hydrogen, C₁-C₄ alkyl and fluoropyridyl, and E may be a phenyl group substituted with a C₁-C₆ alkoxy group.
30

In another particular embodiment of the disclosure, R₃ may be chloro, m may be 1 or 2, G may be CH, D may be CH, M may be N, J may be C, L may be N, Y may be absent, X may be CH₂

or CH_2CH_2 , Z may be NR_1R_2 , wherein R_1 and R_2 are independently selected from hydrogen, $\text{C}_1\text{-C}_4$ alkyl and ^{18}F fluoropyridyl, and E may be a phenyl group substituted with a $\text{C}_1\text{-C}_6$ alkoxy group.

In another particular embodiment of the disclosure, R_3 may be chloro, m may be 1 or 2, G may be CH, D may be CH, M may be N, J may be C, L may be N, Y may be absent, X may be CH_2 or CH_2CH_2 , Z may be NR_1R_2 , wherein R_1 and R_2 are independently selected from hydrogen, $\text{C}_1\text{-C}_4$ alkyl and ^{18}F fluoropyridyl, and E may be a phenyl group substituted with a $\text{C}_1\text{-C}_6$ alkoxy group.

In an embodiment of the disclosure:

G is C,

D, M, J and L are, respectively, N, C, N and N; C, N, C and N; or N, N, C and N,

10 R_3 is Cl, F or Me

m is 1 or 2


E is 4-(2-fluoroethoxy)phenyl, 4-(3-fluoropropoxy)phenyl, 4-*t*-butylphenyl, 4-methoxyphenyl, 2-fluoropyridin-5-yl or 2-fluoropyridazin-5-yl

X is CH_2 (i.e. n is 1)

15 Y is absent or is NH, and

Z is NMe_2 , NEt_2 , NPr_2 , NMeH , NHPr , NMeEt , N-pyrrolidinyl, Ph, NMePh or $\text{NMe(5-(2-fluoro)pyridinyl)}$.

In still another embodiment the disclosure provides a compound of formula (Ia):



20

wherein,

D, G, and L are independently selected from the group consisting of: CH, C and N, and J and M are independently selected from the group consisting of C and N provided that at least one of J and M is C, wherein at least two of D, G, M, J and L are N;

25 X is selected from the group consisting of: O, NH, $(\text{CH}_2)_n$ and S;

Y is absent, or is selected from the group consisting of: O, NH, $(\text{CH}_2)_n$, and S;

Z is selected from the group consisting of: NR_1R_2 and aryl;

R_1 and R_2 are independently selected from the group consisting of: hydrogen, $\text{C}_1\text{-C}_{10}$ alkyl, $\text{C}_2\text{-C}_{10}$ alkenyl, $\text{C}_2\text{-C}_{10}$ alkynyl, $(\text{CH}_2)_n$ aryl, aryl and heteroaryl, each of which may optionally be substituted with one or more of the following substituents: chloro, bromo, iodo, $\text{C}_1\text{-C}_6$ alkyl and hydroxy;

30

or R_1 and R_2 , together with the nitrogen to which they are attached, form a heterocyclic ring having between 3 and 7 ring members, which may optionally be substituted with one or more of the following substituents: chloro, bromo, iodo and C_1-C_6 alkyl;

R_3 is selected from the group consisting of: chloro, bromo, iodo, C_1-C_{10} alkyl and O-(C_1-C_{10} alkyl), wherein the C_1-C_{10} alkyl group is optionally substituted;

E is an aryl group or a heteroaryl group, wherein each group is optionally substituted with a chloro, bromo or iodo substituent, and/or with one or more of the following substituents: C_1-C_6 alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, QC_1-C_{10} alkyl, QC_1-C_{10} alkenyl, QC_2-C_{10} alkynyl, $Q(CH_2)_p-Q-(CH_2)_qCH_3$ or $Q(CH_2)_p-Q-(CH_2)_q-Q-(CH_2)_rCH_3$, each of which may be substituted with one or more chloro, bromo, iodo or hydroxy substituents, and wherein p, q and r are, independently, integers between 1 and 3, and wherein Q is selected from the group consisting of: NH, O and S;

m is a number between 0 and 3 (e.g. 0, 1, 2 or 3) wherein if m is 2 or 3, the R_3 groups may be the same or may be different;

n is a number between 1 and 4, e.g. 1, 2, 3 or 4 (wherein n in X may be the same as or different to n in Y);
with the proviso that E is not 4-iodophenyl.

The disclosure provides a compound of formula (Ia) wherein:

D, G, and L are independently selected from the group consisting of: CH, C and N, and J and M are independently selected from the group consisting of C and N provided that at least one of J and M is C, wherein at least two of D, G, M, J and L are N;

X is independently selected from the group consisting of: O, NH, $(CH_2)_n$ and S;

Y is absent, or is independently selected from the group consisting of: O, NH and $(CH_2)_n$, and S;

Z is selected from the group consisting of: NR_1R_2 and aryl;

R_1 and R_2 are independently selected from the group consisting of: hydrogen, C_1-C_{10} alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, $(CH_2)_n$ aryl, aryl and heteroaryl, each of which may optionally be substituted with one or more of the following substituents: chloro, bromo, iodo, C_1-C_6 alkyl and hydroxy;

or R_1 and R_2 , together with the nitrogen to which they are attached, form a heterocyclic ring having between 3 and 7 ring members, which may optionally be substituted with one or more of the following substituents: chloro, bromo, iodo and C_1-C_6 alkyl;

R_3 is selected from the group consisting of: chloro, bromo, iodo and C_1-C_{10} alkyl;

E may be an aryl group or a heteroaryl group, which is substituted with a chloro, bromo or iodo substituent, and/or one or more of the following substituents: C_1-C_6 alkyl, C_2-C_{10} alkenyl, C_2-

C₁₀ alkynyl, QC₁-C₁₀ alkyl, QC₁-C₁₀ alkenyl, QC₂-C₁₀ alkynyl, Q(CH₂)_p-Q-(CH₂)_qCH₃ or Q(CH₂)_p-Q-(CH₂)_q-Q-(CH₂)_rCH₃, each of which may be substituted with one or more chloro, bromo, iodo or hydroxy substituents, and wherein p, q and r are integers between 1 and 3, and wherein Q is independently selected from the group consisting of: NH, O and S;

5 m is a number between 0 and 3;

n is a number between 1 and 4;

with the proviso that E is not -PhI.

In some embodiments E is not -PhI

In an embodiment of the invention Q may be oxygen.

10 In another embodiment of the invention, m is 1 or 2, and n is 1.

In a further embodiment of the invention, p may be 1 or 2, q and r may be 1, and Q may be oxygen.

In some embodiments, L is N. In some embodiments one of M and J is N.

15 In one embodiment, M and L are N, D and G are CH and J is C. In another embodiment, D, M and L are N, J is C and G is CH. In an alternative embodiment, D, J and L are N, M is C and G is CH.

R₃ may be attached to a carbon atom.

20 R₃ may be selected from the group consisting of: chloro, bromo and iodo and C₁-C₆ alkyl. In another embodiment, R₃ may be selected from the group consisting of chloro and C₁-C₃ alkyl. In a further embodiment, R₃ may be chloro or methyl and m may be 1 or 2. Alternatively, R₃ may be chloro and m may be 1 or 2, or R₃ may be fluoro (¹⁹F or ¹⁸F) and m may be 1 or 2.

In one embodiment, Y is absent and X is (CH₂)_n, wherein n is 1 or 2.

In another embodiment, X and Y may be independently selected from the group consisting of: NH and (CH₂)_n, wherein n is 1.

25 Y may be (CH₂)_n and X may be NH, wherein n is a number between 1 and 3.

In one embodiment, -X-Y-C(O)-Z may have a structure selected from the group consisting of: -(CH₂)_n-C(O)-Z, wherein n is 1 or 2, or -(CH₂)_n-NH-C(O)-Z, wherein n is 1 or 2, and -NH-(CH₂)_n-C(O)-Z, wherein n is 1 or 2.

Z may be NR₁R₂ or phenyl.

30 R₁ and R₂ may be independently selected from the group consisting of: hydrogen C₁-C₆ alkyl, benzyl, phenyl, pyridyl, pyrimidinyl or pyridazinyl, and where R₁ and R₂, together with the nitrogen to which they are attached, form a heterocyclic ring having between 4 and 6 ring members. The heterocyclic ring may be a pyrrolidine ring. The C₁-C₆ alkyl group, the heterocyclic group, the

phenyl group, the pyridyl group, the pyrimidinyl group, the pyridazinyl group or the benzyl group may optionally be substituted with between 1 and 3 chloro, bromo, iodo or hydroxy substituents.

R₁ and R₂ may be independently selected from the group consisting of: hydrogen C₁-C₆ alkyl, phenyl, pyridyl, pyrimidinyl or pyridazinyl, and where R₁ and R₂, together with the nitrogen to which they are attached, form a heterocyclic ring having between 4 and 6 ring members. The heterocyclic ring may be a pyrrolidine ring. The C₁-C₆ alkyl group, the heterocyclic group, the phenyl group, the pyridyl group, the pyrimidinyl group, or the pyridazinyl group may optionally be substituted with between 1 and 3 chloro, bromo, iodo or hydroxy substituents.

In another embodiment, R₁ and R₂ may be independently selected from the group consisting of: hydrogen, phenyl, pyridyl, pyrimidinyl or pyridazinyl, methyl, ethyl, propyl, isopropyl, butyl, *t*-butyl and pentyl, each of which may optionally be substituted with between 1 and 3 chloro, bromo, iodo or hydroxy substituents.

In an alternative embodiment, R₁ and R₂ may be selected from the group consisting of: hydrogen, phenyl, pyridyl, pyrimidinyl, pyridazinyl, methyl, ethyl, isopropyl, propyl, butyl and *t*-butyl, and wherein R₁ and R₂ together with the nitrogen to which they are attached, form a pyrrolidine ring, each of which may optionally be substituted with between 1 and 3 chloro, bromo, iodo or hydroxy substituents.

R₃ may be selected from the group consisting of: halogen and C₁-C₆ alkyl, optionally substituted (e.g. with one or more halogens such as fluorine, chlorine, bromine and/or iodine). R₃ may be methyl. It may be methoxy. It may be trifluoromethyl. It may be trifluoromethoxy.

When E is phenyl, E may not have an iodo substituent attached directly to it in the 4 position. E may be substituted phenyl which does not have an iodo substituent attached directly to the 4 position. It may be substituted phenyl which does not have an iodo substituent attached directly to the phenyl ring.

E may be a phenyl group, a pyridyl group, a pyrimidinyl group or a pyridazinyl group, each of which may optionally be substituted with one or more chloro, bromo, iodo or hydroxy substituents, and/or substituted with one or more of the following substituents: C₁-C₆ alkyl, C₂-C₆ alkenyl or C₁-C₁₀ alkoxy, each of which may optionally be substituted with one or more of the following substituents: chloro, bromo, iodo and hydroxy.

E may be a phenyl group, a pyridyl group, a pyrimidinyl group or a pyridazinyl group, each of which may optionally be substituted with between one and three chloro, bromo, iodo or hydroxy substituents, or substituted with one of the following substituents: C₁-C₆ alkyl, C₂-C₆ alkenyl or C₁-C₆ alkoxy, either of which may optionally be substituted with between one and three chloro, bromo, iodo or hydroxy substituents.

E may be a phenyl group, a pyridyl group, a pyrimidinyl group or a pyridazinyl group, each of which may optionally be substituted with between one and three chloro, bromo, iodo or hydroxyl substituents.

In another aspect, the present invention provides a compound that is radiolabelled with ^{18}F .

In another aspect, the present invention provides a pharmaceutical composition comprising a compound of the present invention, said compound being not radiolabelled with ^{18}F , together with at least one pharmaceutically acceptable carrier, diluent, excipient or adjuvant.

In another embodiment, the present disclosure provides a pharmaceutical composition comprising a compound of one embodiment described in the present disclosure, together with at least one pharmaceutically acceptable carrier, diluent, excipient or adjuvant.

In another embodiment, the present disclosure provides a method for diagnosis of a disorder in a mammal characterised by an abnormal density of peripheral benzodiazepine receptors, the method including the steps of:

- (i) administering to the mammal a compound of the third aspect in an amount sufficient to allow a detectable image of the location of the radiolabel in the body of the mammal to be recorded.
- (ii) diagnosing the presence or absence of the disorder from the image.

The method of diagnosis may comprise recording an image of the distribution of the radiolabel in at least part of the body of the mammal.

The method for diagnosis may be PET. Thus step (ii) may comprise conducting positron emission tomography of the body or part thereof the mammal and determining from a resulting image the presence or absence of the disorder.

In another aspect, the present invention provides the use of a compound of the present invention in the manufacture of a medicament for the treatment of a disorder characterised by an abnormal density of peripheral benzodiazepine receptors in a mammal in need of said treatment, the method comprising administering to the mammal a therapeutically effective amount of a compound of the present invention, said compound being not radiolabelled with ^{18}F .

In another aspect, the invention provides the use of a compound of the present invention in the manufacture of a diagnostic composition in the diagnosis of a disorder in a mammal characterised by an abnormal density of peripheral benzodiazepine receptors.

The compound of the present invention may be selected from the group consisting of: PBRs 099*, 102*, 103*, 104*, 105*, 106*, 107*, 110*, 111*, 130*, 120*, 132*, 133*, 136*, 139*, 143*, 146*, 147*, 149*, 080*, 081*, 082*, 083* (as defined herein), or any combination thereof. These compounds are all radiolabelled with ^{18}F .

Another compound of the present disclosure may be selected from the group consisting of: PBRs 099, 102, 103, 104, 105, 106, 107, 110, 111, 130, 120, 132, 133, 136, 139, 143, 146, 147, 149, 080, 081, 082, 083, 115, 138, 145, 155, 142, 153, 154, 156, 131, 134, 135, 141, 125, 126, 137, 98, 139, 132, 133, 117, 118, 121, 123, 124, 128, 129, 116, 119, 140, 144, 101, 113, and 114, or any combination thereof. None of these compounds are radiolabelled with ^{18}F .

In another aspect, the invention provides the use of a compound according to the present invention in the manufacture of a medicament for the treatment of a condition selected from the group consisting of apoptosis, neurodegenerative disorders, including Huntington's disease, Alzheimer's disease, anxiety, stress, emotional disturbances, cognitive impairment, stroke, and cerebral ischemia; tumours, such as glioma, and carcinoma of the ovary, colon, breast, prostate, brain and adrenals; neurotoxic injury, including that associated with anoxia or ischemia which results from stroke or cardiac arrest; and cognitive disorders, or for cognitive enhancement, said compound being not radiolabelled with ^{18}F .

In another embodiment, the disclosure provides a process for making a compound according to the first, second or third aspect, said process being substantially as described herein with reference to Schemes 1 to 8 (*vide infra*).

Brief Description of the Drawings

Figure 1 shows a plot of the biodistribution of compound PBR111 in rats ($n=4$) over 4 hours. The plot depicts percent of injected dose per gram of tissue v. time.

Figure 2 indicates reduced uptake of [^{18}F]PBR111 radiotracer distribution in organs expressing the PBR receptor when pre-treated with the selective PBR ligands PK 11195, Ro 5-4864 and the non-radioactive analogue of PBR111 all at 1 mg/kg. The central benzodiazepine ligand flumazenil does not compete and hence not reduce the [^{18}F]PBR 111 activity in these tissues.

Figure 3 indicates the percentage ID of [^{18}F]PBR111/gram of tissue v. time. The plot indicates that the tracers follow normal distribution of PBR expression over time.

Figure 4 shows biodistribution of [^{18}F]PBR102 in rodents over a 4 hour time period. The plot shows percentage uptake of [^{18}F]PBR102.

Figure 5 shows the biodistribution data of a selection of tissues from Figure 4 that are relevant to PBR expression.

Figure 6 shows the blocking effect on the uptake of [^{18}F]PBR102 by the selective PBR ligands PK 11195, Ro 5-4864, and the non-radioactive analogue PBR102 all at 1mg/kg. The central benzodiazepine ligand flumazenil does not compete and hence does not reduce the [^{18}F]PBR102 uptake in these tissues.

Figure 7* shows the blocking effect on the uptake of [^{18}F]PBR111 by the selective PBR ligands PK 11195, Ro 5-4864, and the non-radioactive analogue PBR111 all at 1mg/kg. The central benzodiazepine ligand flumazenil does not compete and hence does not reduce the [^{18}F]PBR111 uptake in these tissues.

Figure 8* shows the blocking effect on the uptake of [^{18}F]PBR102 by the selective PBR ligands PK 11195, Ro 5-4864 and the non-radioactive analogue PBR102 all at 1mg/kg. The central benzodiazepine ligand flumazenil does not compete and hence does not reduce the [^{18}F]PBR102 uptake in these tissues.

Figure 9* shows the blocking effect on the uptake of [^{18}F]PBR111 by the selective PBR ligands PK 11195, Ro 5-4864 and the non-radioactive analogue PBR111 all at 1mg/kg. The central benzodiazepine ligand flumazenil does not compete and hence does not reduce the [^{18}F]PBR111 uptake in these tissues.

Figure 10 shows the blocking effect on the uptake of [^{18}F]PBR132 by the selective PBR ligands PK 11195, Ro 5-4864 and the non-radioactive analogue PBR132 all at 1mg/kg. The central benzodiazepine ligand flumazenil does not compete and hence does not reduce the [^{18}F]PBR132 uptake in these tissues.

Figure 11 shows the blocking effect on the uptake of [^{18}F]PBR147 by the selective PBR ligands PK 11195, Ro 5-4864 and the non-radioactive analogue PBR147 all at 1mg/kg. The central benzodiazepine ligand flumazenil does not compete and hence does not reduce the [^{18}F]PBR147 uptake in these tissues.

Figure 12 shows the blocking effect on the uptake of [^{18}F]PBR099 by the selective PBR ligands PK 11195, Ro 5-4864 and the non-radioactive analogue PBR099 all at 1mg/kg. The central benzodiazepine ligand flumazenil does not compete and hence does not reduce the [^{18}F]PBR099 uptake in these tissues.

Figure 13 shows the blocking effect on the uptake of [^{18}F]PBR146 by the selective PBR ligands PK 11195, Ro 5-4864 and the non-radioactive analogue PBR146 all at 1mg/kg. The central benzodiazepine ligand flumazenil does not compete and hence does not reduce the [^{18}F]PBR146 uptake in these tissues.

* The results in the graphs showing "stars" relate to statistical analysis of the results, i.e. Results from the competition studies were analysed by one-way analysis of variance (ANOVA) for comparing the tissue radioactivity concentrations in the treated animals ($n = 3-4$) and in the controls ($n = 4$). The criteria for significance were $**p < 0.01$ (highly significant) and $*p < 0.05$ (significant).

5 No stars implies not statistical difference from normals or controls.

Detailed Description of the Invention

Definitions

The following are some definitions that may be helpful in understanding the description of the present invention. These are intended as general definitions and should in no way limit the scope of the present invention to those terms alone, but are put forth for a better understanding of the following description.

Unless the context requires otherwise or it is specifically stated to the contrary, integers, steps, or elements of the invention recited herein as singular integers, steps or elements clearly encompass both singular and plural forms of the recited integers, steps or elements.

15 Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated step or element or integer or group of steps or elements or integers, but not the exclusion of any other step or element or integer or group of elements or integers. Thus, in the context of this specification, the term "comprising" means "including principally, but not necessarily solely".

20 Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations or any two or more of said steps or features.

25 As used herein, the term "therapeutically effective amount(s)" includes within its meaning a sufficient but non-toxic amount of a compound or composition of the invention to provide the desired therapeutic effect. The exact amount required will vary from subject to subject depending on factors such as the species being treated, the age and general condition of the subject, the severity of the condition being treated, the particular compound being administered, the mode of administration and so forth. Thus, it is not possible to specify an exact "therapeutically effective amount", however for any given case an appropriate "therapeutically effective amount" may be determined by one of ordinary skill in the art using only routine trial and experimentation.

As used herein, the term "treatment" refers to any and all uses which remedy a disease state or symptoms, prevent the establishment of a disease, or otherwise prevent, hinder, retard or reverse the progression of disease or other undesirable symptoms in any way whatsoever.

As used herein, the term "alkyl" includes within its meaning monovalent straight chain or
5 branched chain saturated aliphatic groups having from 1 to 10 carbon atoms, e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms. The alkyl group may have from 1 to 4, 1 to 6 or 1 to 8 carbon atoms. Examples of such alkyl groups include but are not limited to: methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, amyl, isoamyl, sec-amyl, 1,2-dimethylpropyl, 1,1-dimethyl-propyl, hexyl, 4-methylpentyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 1,1-dimethylbutyl, 2,2-
10 dimethylbutyl, 3,3-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 1,2,2-trimethylpropyl and 1,1,2-trimethylpropyl, octyl, nonyl, decyl and the like.

As used herein, the term "alkoxy" refers to a group of the formula alkyl-O-, wherein the alkyl group is as defined above. Examples include methoxy, ethoxy, n-propoxy, isopropoxy, tert-butoxy, and the like.

As used herein, the term "alkenyl" includes within its meaning monovalent straight chain or
15 branched chain unsaturated aliphatic hydrocarbon groups having from 2 to 10 carbon atoms, e.g. 2, 3, 4, 5, 6, 7, 8, 9, or 10 carbon atoms. The alkenyl group may have from 2 to 4, 2 to 6 or 2 to 8 carbon atoms. Examples of such alkenyl groups include but are not limited to vinyl, allyl, 1-methylvinyl, butenyl, iso-butenyl, 3-methyl-2-butenyl, 1-pentenyl, cyclopentenyl, 1-methyl-
20 cyclopentenyl, 1-hexenyl, 3-hexenyl, cyclohexenyl, 1,3-butadienyl, 1,4-pentadienyl, 1,3-cyclopentadienyl, 1,3-hexadienyl, 1,4-hexadienyl, 1,3-cyclohexadienyl and 1,4-cyclohexadienyl.

As used herein, the term "alkynyl" includes within its meaning monovalent and straight chain
or branched chain unsaturated aliphatic hydrocarbon groups having from 2 to 10 carbon atoms (e.g. 2, 3, 4, 5, 6, 7, 8, 9, or 10 carbon atoms) and having at least one triple bond. The alkynyl
25 group may have from 2 to 4, 2 to 6 or 2 to 8 carbon atoms. Examples of such alkynyl groups include but are not limited to: ethynyl, 1-propynyl, 1-butyne, 2-butyne, 1-methyl-2-butyne, 3-methyl-1-butyne, 1-pentyne, 1-hexyne, methylpentynyl, 1-heptyne, 2-heptyne, 1-octyne, 2-octyne, 1-nonyne, 1-decynyl, and the like.

As used herein, the term "aryl" refers to monovalent single, polynuclear, conjugated and
30 fused residues of aromatic hydrocarbons having from 6 and 20 carbon atoms. The aryl group may have from 6 to 10, 6 to 12, 6 to 14, 6 to 16 or 6 to 18 carbon atoms. Examples of such aryl groups include but are not limited to phenyl, biphenyl, naphthyl, tetrahydronaphthyl, indenyl, azulenyl, phenantryl, pyrenyl and the like. Any available position of the aromatic residue can be used for attachment to the remainder of the molecule of formula (I).


As used herein, the term "heteroaryl" refers to single, polynuclear, conjugated and fused aromatic radicals having between 6 and 20 ring atoms, wherein 1 to 6, or 1 to 5, or 1 to 4, or 1 to 3, or 1 or 2 of these rings atoms are heteroatoms independently selected from the group consisting of: N, NH, O and S. The heteroaryl group may have from 6 to 10, 6 to 12, 6 to 14, 6 to 16 or 6 to 18 carbon atoms. The heteroaryl group may have 1 to 2, 1 to 3, 1 to 4, 1 to 5 or 1 to 6 heteroatoms. The hetero atoms may be independently selected from the group consisting of: N and NH, N and O, NH and O, N and S, NH and S and S and O. Examples of such heteroaryl groups include but are not limited to pyridyl, pyridyl, pyridyl, thienyl, furyl, pyrrol, indolyl, pyridazinyl, pyrazolyl, pyrazinyl, thiazolyl, pyrimidinyl, quinolinyl, isoquinolinyl, benzofuranyl, benzothienyl, purinyl, quinazolinyl, phenazinyl, acridinyl, benzoxazolyl, benzothiazolyl and the like. Any available position of the heteroaromatic residue can be used for attachment to the remainder of the molecule of formula (I). Nitrogen-containing heteroaryl groups may be substituted at nitrogen with an oxygen atom to form an N-oxide. Sulfur-containing heteroaryl groups may be substituted at sulfur with one or two oxygen atoms to form a sulfoxide or a sulfone respectively.

Synthesis of compounds

Compounds in accordance with the invention may be synthesised employing combinations of reactions which are generally known in the art. Starting materials for synthesis of compounds of the invention are either commercially available, or are known compounds, or are compounds that can be easily prepared by a person of ordinary skill in the art. Those skilled in the art will also appreciate that various protecting groups may be used throughout a synthesis where a functional group on a molecule may be incompatible with a desired synthetic step. Suitable protecting groups are known to those skilled in the art and have been described, for example, in Greene *et al.*, "Protective Groups in Organic Synthesis"; John Wiley & Sons, 2nd Edition, New York (1991).


Scheme 1 shows one possible synthesis of functionalised imidazo[1,2-a]pyridines starting with an appropriately functionalised acetophenone derivative 1 and an appropriately functionalised aminopyridine 2. In schemes 1 to 6, R_3 may be selected from the group consisting of: halogen, C_1 - C_{10} alkyl and O-(C_1 - C_{10} alkyl), wherein the C_1 - C_{10} alkyl group is optionally substituted. R_3 may be $(R_3)_m$, where m is a number between 0 and 3. If m is 2 or 3, the R_3 groups may be the same or may be different or, in the case where $m=3$, two may be the same and one may be different. The reaction proceeds in a solvent such as ethanol, or similar solvent, with heat to provide the imidazo[1,2-a]pyridines 3.

22




Scheme 1

In an alternative synthesis shown in Scheme 2, the side chain at the 3-position of the imidazo[1,2-a]pyridine can be incorporated into the starting acetophenone derivative (compound 1a). Reaction of 1a with aminopyridine 2 in ethanol, or a similar solvent, gives compound 3a which includes an amide side chain.



Scheme 2

An alternative method for introducing an appropriate side chain, for example $-\text{CH}_2\text{C}(\text{O})\text{NR}_1\text{R}_2$, is shown in Scheme 3. This procedure involves treating an imidazo[1,2-a]pyridine 3, with formaldehyde and dimethylamine in an aqueous medium to give the methanamine derivative 4, which is easily converted to the trimethylammonium salt 5 using methyl iodide. Substitution of the trimethylammonium group of 5 with cyanide ion provides the cyanomethyl compound 6, which may be hydrolysed to the corresponding acid 7 by known methods. The acid 7 may then be converted to a range of amide derivatives 8.





Scheme 3

As an alternative, compounds of formula 4 may be demethylated by known methods and acylated to form compounds having a $-\text{CH}_2\text{NHC(O)R}$ side chain.


5 Scheme 4 shows another method for preparing an imidazo[1,2-a]pyridine derivative from an appropriately functionalised aryl aldehyde derivative **1b**, and an appropriately functionalised aminopyridine **2**, wherein the resulting product **9** may possess a $-\text{NHCOOR}$ side chain. The condensation reaction of compounds **1b** and **2** is carried out in the presence of an isocyanoacetate derivative and scandium(III) triflate. The procedure depicted in Scheme 4 provides compounds of

10 formula **9**, which possess an $-\text{NHCOOEt}$ side chain.




Scheme 5 shows a further alternative procedure for introducing a $-\text{CH}_2\text{NHC(O)-R}$ group into an imidazo[1,2-a]pyridine compound of formula 3. In this procedure, the imidazo[1,2-a]pyridine compound is treated with N-hydroxybenzamide in the presence of sulfuric acid.

Compounds of the type represented by formula 11 may be synthesised according to a literature method (Selleri, S. *et al.* Bioorganic and Medicinal Chemistry **9**, 2661-2671 (2001). One example involves condensation of pentane-2,4-dione with 4-(2-(diethylamino) ethyl)-3-(4-methoxyphenyl)-1H-pyrazol-5-amine.




25

Compounds of the general formula 12 may be prepared by the methods described above in Schemes 1 to 5, except the compound of formula 2 is an appropriately functionalised amino pyridazine as opposed to an amino pyridine.



12

5




Scheme 6

The substituent B depicted in schemes 1 to 5 can be manipulated using simple functional group interconversions known to those skilled in the art, so as to produce a wide range of derivatives once the initial condensation reaction has been performed. Scheme 6 shows one such possibility for preparing a fluorinated ether derivative. The methoxy compound 13 is cleaved by a standard procedure using BCl_3 and tetrabutylammonium iodide to give hydroxy compound 14. Reaction of compound 14 with an appropriate alkyl fluoride in the presence of a base gives the fluorinated compound 15.


Schemes 7 to 10 set out synthetic routes to selected compounds of the invention. Examples of carrying out the various methods A to J referred to in schemes 7 to 10 can be found in Examples 3 to 17.

It should be noted that the synthesis of the compounds of the present invention is simpler than the synthesis disclosed in WO 99/51594 in that no iodo substituent is necessary on the 2-phenyl ring.


20



Scheme 7




Scheme 8



Scheme 9

** Salleri, S. *et al.* Bioorganic and Medicinal Chemistry 9, 2661-2671 (2001).


27




Scheme 10

*** Katsifis *et al.* Radiochim Acta 92, 305-309 (2004).

Compounds in accordance with the present invention include the following:





Intermediate compounds of the present invention also include the following: PBRs 115, 138, 145, 155, 142, 153, 154, 156, 131, 134, 135, 141, 125, 126, 137, 98, 139, 132, 133, 117, 118, 121, 123, 124, 128, 129, 116, 119, 140, 144, 101, 113, and 114 (see Schemes 7 to 10)

Radiolabelling of compounds

Diagram 1 illustrates the general structure of the compounds of the invention, where A and C are unsaturated ring systems, B indicates the different configuration of the amide groups, and R₁ and R₂ designate various substituents.





Diagram 1


Radiolabelling is possible on rings B or C, or at positions R1 or R2 to form radiolabelled products. Three different paths to form ^{18}F labelled products in the form of alkyl fluorides, fluoro pyridines and fluoro pyridazines are described below.

Alkyl fluorides can be introduced as substituents on rings A or C, or as part of the functional groups in R1 or R2, while fluoropyridines and fluoropyridazines can be introduced as part of rings A or C, or as a part of the group in R1 or R2.


Radiofluorination may be achieved by methods known to those skilled in the art. One such method involves the use of the $[\text{K}2.2.2][\text{K}_2\text{CO}_3][^{18}\text{F}]$ complex, wherein the complex is reacted with a terminal alkyl group possessing a suitable leaving group, for example a halogen or *O*-tosylate or *O*-mesylate. This method is depicted in Schemes 11, 12 and 13 below.



Scheme 11 – Formation of alkyl fluorides



Scheme 12 – Formation of Fluoropyridines




Scheme 13 – Formation of Fluoropyridazines



In a typical radiofluorination reaction, ^{18}F -fluoride, in the form of H^{18}F , is produced from an $^{18}\text{O}(\text{p},\text{n})^{18}\text{F}$ nuclear reaction by bombardment of an ^{18}O -enriched water target. The aqueous ^{18}F -fluoride solution is added to a 2.5 mL vial containing dry acetonitrile (1 mL), Kryptofix 2.2.2 and potassium carbonate. The solvent is evaporated under a stream of nitrogen at a temperature of about 80-120°C. This azeotropic drying is then repeated twice by further addition of acetonitrile (2 x 1 mL). The precursor compound is dissolved in an appropriate solvent, for example acetonitrile, DMF or DMSO (1 ml) and then added to the dried $\text{K}_{222}.\text{K}_2\text{CO}_3.\text{KF}$ complex. Stirring and heating is continued for about 5 to 15 mins before the reaction mixture is diluted with mobile phase and purified by reverse phase HPLC. The radioactive peak corresponding to the fluorinated product is collected, dried *in vacuo* and then formulated in saline (0.9%) for biological evaluation.

Radiolabelled compounds





Radiolabelled compounds in accordance with the present invention include the following:



31




5



10





Pharmaceutical compositions

Pharmaceutical compositions in accordance with the invention may be administered orally, topically, parenteral, or subcutaneous, e.g. by injection and by intra-arterial infusion, rectally or by inhalation spray.

For oral administration, the pharmaceutical composition may be in the form of tablets, lozenges, pills, troches, capsules, elixirs, powders, granules, suspensions, emulsions, syrups and tinctures. Slow-release or delayed-release forms may also be prepared, for example in the form of coated particles, multi-layer tablets or microgranules. Solid forms for oral administration may contain pharmaceutically acceptable binders, sweeteners, disintegrating agents, diluents, flavourings, coating agents, preservatives, lubricants and/or time delay agents. Suitable binders include gum acacia, gelatin, corn starch, gum tragacanth, sodium alginate, carboxymethylcellulose or polyethylene glycol. Suitable sweeteners include sucrose, lactose, glucose, aspartame or saccharine. Suitable disintegrating agents include corn starch, methylcellulose, polyvinylpyrrolidone, xanthan gum, bentonite, alginic acid or agar. Suitable diluents include lactose, sorbitol, mannitol, dextrose, kaolin, cellulose, calcium carbonate, calcium silicate or dicalcium phosphate. Suitable flavouring agents include peppermint oil, oil of wintergreen, cherry, orange or raspberry flavouring. Suitable coating agents include polymers or copolymers of acrylic acid and/or methacrylic acid and/or their esters, waxes, fatty alcohols, zein, shellac or gluten. Suitable preservatives include sodium benzoate, vitamin E, alpha-tocopherol, ascorbic acid, methyl paraben, propyl paraben or sodium bisulphite. Suitable lubricants include magnesium stearate, stearic acid, sodium oleate, sodium chloride or talc. Suitable time delay agents include glyceryl monostearate or glyceryl distearate.

Liquid forms for oral administration may contain, in addition to the above agents, a liquid carrier. Suitable liquid carriers include water, oils such as olive oil, peanut oil, sesame oil,

sunflower oil, safflower oil, arachis oil, coconut oil, liquid paraffin, ethylene glycol, propylene glycol, polyethylene glycol, ethanol, propanol, isopropanol, glycerol, fatty alcohols, triglycerides or mixtures thereof.

Suspensions for oral administration may further comprise dispersing agents and/or
5 suspending agents. Suitable suspending agents include sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, sodium alginate or cetyl alcohol. Suitable dispersing agents include lecithin, polyoxyethylene esters of fatty acids such as stearic acid, polyoxyethylene sorbitol mono- or di-oleate, -stearate or -laurate, polyoxyethylene sorbitan mono- or di-oleate, -stearate or -laurate and the like. The emulsions for oral administration
10 may further comprise one or more emulsifying agents. Suitable emulsifying agents include dispersing agents as exemplified above or natural gums such as gum acacia or gum tragacanth. For topical administration, the pharmaceutical composition may be in the form of a cream, ointment, gel, jelly, tincture, suspension or emulsion. The pharmaceutical composition may contain pharmaceutically acceptable binders, diluents, disintegrating agents, preservatives, lubricants,
15 dispersing agents, suspending agents and/or emulsifying agents as exemplified above.

For parenteral administration, the compound of formula (I) or (Ia) or a salt thereof, may be prepared in sterile aqueous or oleaginous solution or suspension. Suitable non-toxic parenterally acceptable diluents or solvents include water, Ringer's solution, isotonic salt solution, 1,3-butanediol, ethanol, propylene glycol or polyethylene glycols in mixtures with water. Aqueous
20 solutions or suspensions may further comprise one or more buffering agents. Suitable buffering agents include sodium acetate, sodium citrate, sodium borate or sodium tartrate, for example.

For rectal administration, the compound of formula (I) or (Ia) is suitably administered in the form of an enema or suppository. A suitable suppository may be prepared by mixing the active substance with a non-irritating excipient which is solid at ordinary temperatures but which will melt
25 in the rectum. Suitable such materials are cocoa butter and polyethylene glycols. Suitable enemas may comprise agents as exemplified above with reference to forms for topical administration.

Suitably, an inhalation spray comprising a compound of formula (I) or (Ia) will be in the form of a solution, suspension or emulsion as exemplified above. The inhalation spray composition may further comprise an inhalable propellant of low toxicity. Suitable propellants include carbon dioxide
30 or nitrous oxide.

Pharmaceutical compositions, diagnostic compositions and medicaments including compounds of the first, second or third aspects may be manufactured by methods which are generally known in the art. Compositions or medicaments may be prepared by grinding, crushing, blending, dispersing, dissolving, suspending, mixing, admixing, combining, emulsifying or

homogenising a compound of the first or second aspect with one or more suitable carriers, adjuvants, diluents or excipients. Combinations of two or more of the foregoing steps may also be employed.

The dosage form of the compound of formula (I) or (Ia) will include from 0.01% to 99% by weight of the compound of formula (I) or (Ia). Usually, dosage forms according to the invention will
5 comprise from 0.1% to about 10% by weight of the active substance.

In a diagnostic method in accordance with the sixth aspect of the invention, a dosage of typically from about 3 to about 10mCi of a compound of the third aspect is typically administered to a mammal, usually a human, in whom it is desired to diagnose the presence or absence of a
10 disorder characterised by an abnormal density of peripheral benzodiazepine receptors. After a period of from 0.5 to 4 hours following administration of a compound of the second embodiment, or alternatively from 1 to 40 hours, an image of the distribution of radioactivity in the body, or part of the body, of the mammal may be obtained. It will be appreciated that the lower doses are appropriate for administration to a child, and the higher dosages are more appropriate for
15 administration for diagnosis of a tumour where imaging is to be carried out 2 to 4 hours or more after administration of the substance. The presence of a concentration of radioactivity at a site where an abnormal density of peripheral benzodiazepine receptors occurs in a mammal suffering from the disorder indicates a positive diagnosis.

A method in accordance with the seventh aspect includes the administration to a mammal,
20 typically a human, of a compound of the first or second aspects, or a pharmaceutical composition of the fourth aspect. Single or multiple administrations of the pharmaceutical compositions of the fourth aspect may be carried out. One skilled in the art would be able, by routine experimentation, to determine a therapeutically effective amount (which is non-toxic) of the compound and/or compositions of the invention and an administration pattern which would be suitable for treating the
25 diseases to which the compounds and compositions are applicable.

Further, it will be apparent to one of ordinary skill in the art that the optimal course of treatment, such as the number of doses of therapeutically effective amounts of the compound or compositions of the invention given per day for a defined number of days, can be ascertained using conventional treatment determination tests. Generally, a therapeutically effective amount per 24
30 hours may be in the range of about 0.0001 mg to about 1000 mg per kg body weight; suitably, about 0.001 mg to about 750 mg per kg body weight; about 0.01 mg to about 500 mg per kg body weight; about 0.1 mg to about 500 mg per kg body weight; about 0.1 mg to about 250 mg per kg body weight; or about 1.0 mg to about 250 mg per kg body weight. More suitably, an effective dosage per 24 hours may be in the range of about 1.0 mg to about 200 mg per kg body weight;

about 1.0 mg to about 100 mg per kg body weight; about 1.0 mg to about 50 mg per kg body weight; about 1.0 mg to about 25 mg per kg body weight ; about 5.0 mg to about 50 mg per kg body weight; about 5.0 mg to about 20 mg per kg body weight; or about 5.0 mg to about 15 mg per kg body weight. The therapeutically effective amount per 24 hrs may be continued on a daily, or every
 5 second, third, fourth, fifth, sixth or seventh day, until treatment of the disorder in the subject is achieved.

A therapeutic dosage of a compound of the first or second aspects of the invention may be determined by the attending physician in any given circumstance, depending on factors such as the condition which is to be treated in the patient, and its severity.

10 Conditions for the treatment of which compounds in accordance with the invention are useful are conditions associated with abnormal density of peripheral benzodiazepine receptors, such as inflammation, neurodegenerative disorders, including Huntington's disease, Alzheimer's disease, multiple sclerosis, anxiety, stress, emotional disturbances, cognitive impairment, stroke, and cerebral ischemia; certain tumours, such as glioma, and carcinoma of the ovary, colon, breast,
 15 prostate, brain and adrenals; neurotoxic injury, including that associated with anoxia or ischemia which may result from stroke or cardiac arrest; cognitive disorders, in cognitive enhancement and in the modulation of apoptotic processes. A compound in accordance with the invention may be administered in a single dose, or in two doses, or in multiple doses, depending on the disorder, the stage of the disorder, its response to the treatment, and any undesirable effects which may become
 20 apparent.

In vivo biodistribution studies

In vivo biodistribution of the radiolabelled compounds of the invention indicated high uptake in the tissues which have known PBR sites. Tables 1, 1A, 2, 2A, 3, 4, 5 and 6 show the biodistribution of the compounds of PBR111*, PBR102*, PBR132*, PBR146*, PBR147*, and
 25 PBR099* in which the fluorine atom is radiolabelled, when administered to rats. Pretreatment of the rats with known standards such as PK 11195 without radiolabel (1 mg/kg) significantly reduced the uptake of activity in the tissues of interest. Pretreatment with Flumazenil and other central acting benzodiazepines did not reduce the uptake of activity of the radiolabelled compounds of the invention, thereby indicating that the binding of the compounds of the invention *in vivo* is specific
 30 and selective for the peripheral benzodiazepine receptors.

Table 1: Biodistribution of the PBR 111* in normal rodents (rats). The uptake of the tracer in the different organs is consistent with the normal known peripheral benzodiazepine receptor distribution†.

35

Time\Organ	15 min	30 min	1 hour	4 hours
------------	--------	--------	--------	---------

LIVER	1.54±0.20	0.94±0.28	0.82±0.22	0.44±0.07
SPLEEN	8.25±2.25	6.54±1.35	6.72±1.91	3.45±0.43
KIDNEY	5.18±0.67	4.77±1.05	5.08±1.23	2.57±0.48
MUSCLE	0.68±0.16	0.62±0.15	0.55±0.11	0.19±0.05
SKIN	0.39±0.06	0.41±0.06	0.54±0.07	0.71±0.07
BONE	0.99±0.18	1.35±0.27	2.41±0.24	4.86±0.86
LUNGS	16.45±2.81	9.27±1.60	7.29±2.12	3.37±0.40
HEART	7.75±1.08	7.30±0.90	8.41±1.91	4.92±0.66
BLOOD	0.48±0.11	0.33±0.11	0.24±0.08	0.12±0.02
BLADDER	0.67±0.36	0.76±0.16	1.24±0.29	1.33±0.17
STOMACH	0.95±0.23	1.18±0.34	0.91±0.06	0.51±0.06
GIT	1.21±0.11	1.38±0.36	1.89±0.37	2.01±0.27
BRAIN	0.31±0.06	0.19±0.04	0.17±0.04	0.10±0.02
OLF BULBS	0.74±0.14	0.60±0.14	0.65±0.14	0.47±0.06
THYROID	11.40±0.33	13.24±4.62	13.04±3.22	8.91±1.19
PANCREAS	2.33±0.33	1.49±0.29	1.24±0.36	0.61±0.10
THYMUS	1.07±0.23	1.23±0.47	1.39±0.29	1.88±0.40
ADRENALS	8.36±2.02	9.06±3.47	12.11±3.32	17.81±3.61
TESTES	0.35±0.05	0.38±0.07	0.51±0.09	0.67±0.08

Results are mean ± SD percent injected dose per gram (% ID/g), n = 4 rats.

† Results not corrected for the decay of ¹⁸F

Table 1A: Biodistribution of PBR111* in rats. Results are mean ± SD percent injected dose per gram (% ID/g), n = 4 rats.

5

	15 min	30 min	60 min	4 hours
Liver	1.37±0.20	0.86±0.17	0.65±0.09	0.39±0.06
Spleen	7.25±1.57	5.98±0.63	5.27±0.85	3.12±0.37
Kidney	4.56±0.34	4.34±0.52	4.11±0.66	2.32±0.37
Muscle	0.61±0.16	0.57±0.10	0.44±0.04	0.17±0.05
Skin	0.34±0.03	0.37±0.02	0.46±0.02	0.63±0.04
Bone	0.87±0.12	1.22±0.10	2.04±0.09	4.32±0.33
Lungs	14.39±2.19	8.41±0.77	5.68±1.03	3.02±0.39
Heart	6.76±0.82	6.63±0.35	6.72±0.80	4.38±0.42
Blood	0.41±0.08	0.30±0.06	0.18±0.04	0.11±0.01
Bladder	0.60±0.28	0.68±0.08	1.01±0.28	1.20±0.23
Stomach	0.82±0.20	1.06±0.23	0.78±0.12	0.45±0.06

37

GIT	1.05±0.08	1.22±0.20	1.54±0.19	1.77±0.12
Brain	0.27±0.04	0.17±0.02	0.13±0.01	0.08±0.01
Olfactory Bulbs	0.64±0.10	0.54±0.07	0.54±0.10	0.41±0.04
Thyroid	9.82±0.74	10.33±0.99	10.72±2.14	8.25±1.10
Pancreas	1.99±0.18	1.28±0.12	0.92±0.10	0.53±0.09
Thymus	0.91±0.17	1.10±0.35	1.12±0.21	1.65±0.41
Adrenals	6.26±0.94	6.49±0.93	9.23±1.25	16.46±0.97
Testes	0.29±0.03	0.33±0.03	0.40±0.01	0.57±0.03

† Results corrected for the decay of ^{18}F

Table 2: Biodistribution of PBR 102* in normal rats, expressed as a % of ID/g tissue over a 4 hour period†.

	15 min	30 min	60 min	4 hours
Liver	0.75±0.21	0.63±0.21	0.40±0.04	0.34±0.05
Spleen	5.35±0.89	4.68±0.90	3.89±0.54	2.29±0.50
Kidney	3.90±0.53	3.95±1.01	3.10±0.51	1.61±0.29
Muscle	0.49±0.03	0.46±0.08	0.39±0.07	0.27±0.03
Skin	0.29±0.05	0.37±0.05	0.43±0.07	0.59±0.06
Bone	0.56±0.07	0.64±0.12	0.73±0.13	1.22±0.22
Lungs	17.25±1.70	14.80±3.67	13.44±2.71	10.26±1.09
Heart	5.23±0.33	5.41±0.71	5.32±0.74	3.38±0.44
Blood	0.25±0.06	0.21±0.03	0.20±0.03	0.26±0.03
Bladder	0.59±0.14	0.67±0.16	0.60±0.11	1.08±0.14
Stomach	0.80±0.19	0.71±0.21	0.52±0.13	0.49±0.13
Gut	0.82±0.11	1.07±0.26	1.09±0.13	1.43±0.34
Brain	0.24±0.03	0.20±0.04	0.20±0.03	0.23±0.03
Olf Bulbs	0.68±0.12	0.44±0.05	0.52±0.05	0.50±0.13
Thyroid	7.74±1.90	10.70±2.08	11.55±4.24	8.29±3.22
Pancreas	1.74±0.22	1.42±0.45	0.88±0.18	0.49±0.12
Thymus	0.68±0.17	0.81±0.06	0.91±0.16	1.21±0.27
Adrenals	8.30±2.13	9.75±1.53	7.88±1.86	9.77±2.42
Testes	0.22±0.02	0.29±0.05	0.37±0.06	0.52±0.06

5 Results are mean ± SD percent injected dose per gram (% ID/g), n = 4 rats.

† Results not corrected for the decay of ¹⁸F.

Table 2A: Biodistribution of PBR102* in rats. Results are mean ± SD percent injected dose per gram (% ID/g), n = 4 rats#

	15 min	30 min	60 min	4 hours
Liver	0.63±0.15	0.52±0.14	0.35±0.01	0.29±0.02
Spleen	4.48±0.52	3.88±0.44	3.42±0.19	1.92±0.23
Kidney	3.26±0.27	3.25±0.50	2.72±0.23	1.35±0.11
Muscle	0.41±0.02	0.38±0.03	0.34±0.04	0.23±0.03
Skin	0.24±0.03	0.31±0.03	0.37±0.03	0.49±0.03
Bone	0.46±0.04	0.52±0.05	0.64±0.07	1.01±0.09
Lungs	14.30±1.45	12.10±1.98	11.63±1.52	8.09±1.14

39


Heart	4.35±0.33	4.44±0.26	4.62±0.27	2.82±0.15
Blood	0.20±0.04	0.17±0.02	0.18±0.01	0.21±0.01
Bladder	0.49±0.10	0.55±0.10	0.52±0.08	0.90±0.07
Stomach	0.66±0.15	0.57±0.12	0.45±0.08	0.41±0.12
GIT	0.67±0.06	0.86±0.13	0.94±0.05	1.17±0.17
Brain	0.20±0.01	0.16±0.02	0.17±0.01	0.19±0.01
Olfactory bulbs	0.55±0.07	0.35±0.01	0.44±0.02	0.40±0.07
Thyroid	6.28±1.36	8.77±2.27	8.24±0.99	5.75±0.89
Pancreas	1.41±0.09	1.13±0.26	0.75±0.11	0.40±0.06
Thymus	0.55±0.13	0.65±0.07	0.77±0.10	0.98±0.18
Adrenals	6.66±1.31	7.92±1.70	7.31±1.14	7.14±1.04
Testes	0.17±0.01	0.23±0.03	0.31±0.03	0.42±0.01

Results corrected for the decay of ^{18}F **Table 3: Biodistribution of PBR132* in SD rats. Results are mean ± SD percent injected dose per gram (% ID/g), n = 4 rats*.**

	15 min	30 min	60 min	4 hours
Liver	1.22±0.17	1.18±0.33	0.67±0.06	0.31±0.03
Spleen	5.85±0.93	6.89±0.80	5.56±0.67	2.64±0.36
Kidney	4.10±0.28	3.94±0.54	3.24±0.15	2.31±0.17
Muscle	0.89±0.28	0.91±0.10	0.50±0.03	0.24±0.06
Skin	0.60±0.09	0.65±0.08	0.64±0.07	0.67±0.17
Bone	1.04±0.14	1.25±0.18	2.06±0.10	3.28±0.65
Lungs	16.09±2.56	11.58±2.11	6.40±0.50	2.42±0.25
Heart	8.13±0.77	8.34±0.86	6.87±0.53	4.71±0.31
Blood	0.41±0.05	0.32±0.05	0.17±0.00	0.07±0.00
Bladder	0.73±0.33	0.87±0.36	0.84±0.08	0.99±0.30
Stomach	0.58±0.19	0.64±0.11	0.47±0.05	0.72±0.24
GIT	1.08±0.08	1.36±0.22	1.78±0.04	2.93±0.39
Brain	0.35±0.03	0.23±0.02	0.12±0.00	0.07±0.00
Olfactory Bulbs	0.63±0.09	0.46±0.11	0.36±0.05	0.25±0.03
Thyroid	8.87±1.06	9.15±0.87	18.30±1.00	7.41±0.68
Pancreas	2.01±0.26	1.97±0.32	1.20±0.19	0.57±0.10
Thymus	0.78±0.20	0.80±0.06	0.88±0.07	1.02±0.06
Adrenals	10.03±1.11	13.21±1.16	11.24±2.03	13.51±0.61
Testes	0.31±0.03	0.31±0.03	0.32±0.01	0.42±0.06

5 * Results corrected for the decay of ^{18}F

Table 4: Biodistribution of PBR146* in rat. Results are mean $\pm$ SD percent injected dose per gram (% ID/g), n = 4 rats*.



5

	15 min	30 min	60 min	4 hours
Liver	1.08 $\pm$ 0.05	0.98 $\pm$ 0.21	0.69 $\pm$ 0.26	0.38 $\pm$ 0.09
Spleen	5.49 $\pm$ 0.90	5.53 $\pm$ 0.36	5.14 $\pm$ 0.38	3.61 $\pm$ 0.19
Kidney	5.53 $\pm$ 1.20	4.91 $\pm$ 0.21	4.70 $\pm$ 0.24	3.23 $\pm$ 0.26
Muscle	1.10 $\pm$ 0.15	0.82 $\pm$ 0.13	0.67 $\pm$ 0.04	0.31 $\pm$ 0.06
Skin	0.53 $\pm$ 0.11	0.55 $\pm$ 0.05	0.67 $\pm$ 0.06	0.71 $\pm$ 0.07
Bone	1.06 $\pm$ 0.17	1.46 $\pm$ 0.27	2.07 $\pm$ 0.27	4.12 $\pm$ 0.23
Lungs	20.42 $\pm$ 4.32	12.36 $\pm$ 1.28	7.12 $\pm$ 0.95	3.25 $\pm$ 0.21
Heart	8.99 $\pm$ 0.69	9.05 $\pm$ 0.51	8.57 $\pm$ 1.01	5.76 $\pm$ 0.92
Blood	0.38 $\pm$ 0.08	0.26 $\pm$ 0.03	0.18 $\pm$ 0.02	0.10 $\pm$ 0.01
Bladder	0.63 $\pm$ 0.14	0.71 $\pm$ 0.22	0.89 $\pm$ 0.25	1.26 $\pm$ 0.16
Stomach	0.69 $\pm$ 0.12	1.03 $\pm$ 0.16	1.27 $\pm$ 0.45	0.34 $\pm$ 0.04
GIT	1.05 $\pm$ 0.13	1.25 $\pm$ 0.09	1.51 $\pm$ 0.21	1.54 $\pm$ 0.11
Brain	0.27 $\pm$ 0.03	0.15 $\pm$ 0.02	0.12 $\pm$ 0.01	0.08 $\pm$ 0.01
Olfactory Bulbs	0.41 $\pm$ 0.10	0.45 $\pm$ 0.07	0.25 $\pm$ 0.06	0.31 $\pm$ 0.08
Thyroid	7.71 $\pm$ 2.92	7.69 $\pm$ 2.21	13.37 $\pm$ 0.46	10.68 $\pm$ 1.44
Pancreas	2.20 $\pm$ 0.68	1.59 $\pm$ 0.10	1.41 $\pm$ 0.22	0.77 $\pm$ 0.16
Thymus	0.80 $\pm$ 0.07	0.84 $\pm$ 0.11	0.96 $\pm$ 0.17	1.35 $\pm$ 0.15
Adrenals	8.05 $\pm$ 0.66	7.94 $\pm$ 1.52	11.02 $\pm$ 1.88	12.21 $\pm$ 1.56
Testes	0.269 $\pm$ 0.022	0.285 $\pm$ 0.024	0.319 $\pm$ 0.028	0.426 $\pm$ 0.043

* Results corrected for the decay of ^{18}F

Table 5: Biodistribution of PBR147* in SD rats. Results are mean $\pm$ SD percent injected dose per gram (% ID/g), n = 4 rats*.

	15 min	30 min	60 min	4 hours
Liver	1.88 $\pm$ 0.16	1.37 $\pm$ 0.38	1.08 $\pm$ 0.14	0.57 $\pm$ 0.02
Spleen	8.07 $\pm$ 1.11	7.25 $\pm$ 1.13	6.98 $\pm$ 0.29	4.79 $\pm$ 0.31
Kidney	4.75 $\pm$ 0.75	3.82 $\pm$ 0.45	3.68 $\pm$ 0.28	3.26 $\pm$ 0.16
Muscle	0.78 $\pm$ 0.09	0.64 $\pm$ 0.19	0.55 $\pm$ 0.11	0.44 $\pm$ 0.04

41

Skin	0.45±0.03	0.51±0.06	0.49±0.06	0.57±0.04
Bone	0.94±0.05	1.18±0.08	1.49±0.16	2.56±0.03
Lungs	26.34±3.04	18.00±2.29	11.36±0.97	5.62±0.24
Heart	6.53±0.38	5.69±0.42	5.82±0.24	6.09±0.54
Blood	0.63±0.08	0.34±0.06	0.23±0.03	0.13±0.01
Bladder	0.61±0.15	0.58±0.05	0.58±0.12	0.80±0.05
Stomach	0.41±0.12	0.40±0.06	0.64±0.27	0.83±0.17
GIT	0.98±0.15	1.14±0.04	1.44±0.29	2.71±0.10
Brain	0.43±0.05	0.26±0.04	0.16±0.01	0.09±0.01
Olfactory Bulbs	0.58±0.09	0.48±0.09	0.27±0.03	0.24±0.09
Thyroid	7.23±1.35	7.43±1.00	8.17±1.71	6.44±0.45
Pancreas	2.12±0.16	1.88±0.24	1.66±0.13	0.95±0.14
Thymus	0.76±0.12	0.63±0.11	0.69±0.08	0.81±0.08
Adrenals	7.59±1.30	8.47±1.46	8.45±0.84	9.30±0.55
Testes	0.23±0.02	0.21±0.02	0.22±0.01	0.28±0.02

° Results corrected for the decay of ^{18}F

Table 6: Biodistribution of PBR099* in SD rats. Results are mean ± SD percent injected dose per gram (% ID/g).

	15 min		30 min		60 min		4 hours	
Liver	0.63	± 0.06	0.54	± 0.18	0.40	± 0.05	0.29	± 0.04
Spleen	4.60	± 0.29	4.71	± 0.61	4.51	± 0.70	2.88	± 0.56
Kidney	3.85	± 0.24	3.62	± 0.53	3.53	± 0.24	2.41	± 0.40
Muscle	0.65	± 0.14	0.55	± 0.03	0.43	± 0.02	0.30	± 0.05
Skin	0.35	± 0.04	0.49	± 0.02	0.48	± 0.04	0.65	± 0.05
Bone	0.64	± 0.03	0.70	± 0.02	0.79	± 0.03	1.07	± 0.05
Lungs	14.55	± 1.35	8.82	± 0.53	5.66	± 0.56	2.80	± 0.51
Heart	5.86	± 0.44	5.59	± 0.26	6.45	± 0.43	4.29	± 0.86
Blood	0.23	± 0.03	0.16	± 0.01	0.14	± 0.01	0.16	± 0.01
Bladder	0.46	± 0.18	0.42	± 0.08	0.72	± 0.17	0.80	± 0.11
Stomach	0.45	± 0.18	0.32	± 0.04	0.41	± 0.09	0.79	± 0.16
GIT	0.83	± 0.09	0.90	± 0.03	1.31	± 0.08	1.90	± 0.09
Brain	0.21	± 0.01	0.14	± 0.01	0.13	± 0.02	0.15	± 0.01
Olfactory Bulbs	0.35	± 0.03	0.28	± 0.03	0.29	± 0.03	0.31	± 0.05
Thyroid	6.94	± 1.04	7.99	± 1.83	9.90	± 3.67	7.97	± 1.37
Pancreas	1.70	± 0.13	1.33	± 0.16	1.08	± 0.19	0.56	± 0.09
Thymus	0.62	± 0.17	0.66	± 0.14	0.77	± 0.12	0.91	± 0.05

Adrenals	5.64	±	0.53	6.56	±	1.20	7.82	±	1.73	9.69	±	1.49
Testes	0.184	±	0.006	0.209	±	0.007	0.266	±	0.019	0.349	±	0.011

In vitro binding studies

The inhibition constant (IC_{50}) of selected compounds for the peripheral benzodiazepine binding site was determined by incubating, in triplicate, aliquots (0.1 ml) of diluted kidney membrane preparation (200 μ g), at 4°C for 1 h with 7 concentrations of the compound (10^{-5} to 10^{-10} M) and [3 H]-PK11195 (2 nM) in a final volume of 0.5 ml. Non-specific binding was determined with PK11195 (10 μ M).

Similarly, the IC_{50} of the compounds for central benzodiazepine receptors was determined by incubating aliquots (0.1 ml), in triplicate, of diluted brain cortex membrane preparation (200 μ g) at 25°C for 45 min with 7 concentrations of the compounds (10^{-5} to 10^{-10} M) and of [3 H]-Flumazenil (2 nM) in a final volume of 0.5 mL. Non-specific binding was determined with Flumazenil (20 μ M). In both cases, incubations were terminated by rapid filtration through Whatman GF/B glass fibre. Filters were immediately washed 3 times with 5 mL ice-cold 50 mM Tris/HCl at pH 7.4. Filters were counted in a β -scintillation counter (Packard) to measure the amount of bound radioactivity. The IC_{50} of the compounds for the peripheral benzodiazepine binding site and the central benzodiazepine receptor were calculated using Kell 6 software. These results are shown in Table 7.

Table 7: Binding Affinity for PBR and CBR of all compounds

Compound	PBR IC_{50} (nM)	CBR IC_{50} (nM)	LogP HPLC
PBR099	14.95	>5000	2.54±0.08
PBR102	13.2	1560	2.70±0.09
PBR103	51.3	1843.6	3.44±0.11
PBR104	575.2	1635.5	2.45±0.08
PBR105	674.9	1759.9	2.04±0.06
PBR106	169.6	547.7	2.05±0.07
PBR107	54.8	1393.6	2.34±0.07
PBR110	14.1	2735.7	2.74±0.09
PBR111	7.5	1588.1	3.20±0.10
PBR 120	35	941.6	3.53±0.11
PBR 130	20	50% @ 10^{-5} M >5000	3.43±0.10
PBR 132	38	343.2	3.60±0.11
PBR 133	32	18.8	2.35±0.07
PBR 136	17.6	588.1	2.99±0.09
PBR 139	800	2616.7	4.23±0.13
PBR 143	6.4	50% @ 10^{-5} M	3.85±0.12


43

PBR 146	10.5	8000 50% @ 10 ⁻⁵ M	3.16±0.10
PBR 147	7.4	4087.3	4.31±0.13
PBR 149	4.7	357.0	3.57±0.11

Examples

The invention will now be described in more detail, by way of illustration only, with respect to the following examples. The examples are intended to serve to illustrate this invention and should not be construed as limiting the generality of the disclosure of the description throughout this specification.

Comparative example



10

Compounds 1 to 4 were prepared in order to assess the effect on selectivity of replacing the 2-iodo substituent with a 2-fluoro substituent. As can be seen from the IC₅₀ values, where a 2-fluoro substituent is substituted for the 2-iodo substituent, the selectivity for the peripheral benzodiazepine receptor is lost. The above results surprisingly show that the simple replacement of the iodo substituent of the compounds disclosed in WO 99/51594 with a fluoro substituent results in compounds that are selective for the central benzodiazepine receptor as opposed to the peripheral benzodiazepine receptor. Unexpectedly, removing the iodo substituent from the phenyl ring and attaching a fluoro substituted alkoxy group results in fluorinated compounds that retain selectivity for the peripheral benzodiazepine receptor such that they are useful for PET imaging and also for therapeutic applications of disorders characterised by an abnormal density of peripheral benzodiazepine receptors.

20

Example 1 - Preparation of precursor bromoacetophenones

A solution of bromine (2.52 g, 15.75 mmoles) in carbon tetrachloride (10 ml) was added over 1 h to a stirred, ice-cooled solution of *N,N*-diethyl-4-(4-methoxyphenyl)-4-oxobutanamide (3.945 g, 15.0 mmoles) in carbon tetrachloride (50 ml) containing 1 drop of conc. HBr. Stirring was continued at this temperature for a further 2 h, and then at room temperature for 30 min. The pale yellow solution was evaporated and the residue stirred with ethyl acetate (100 ml) and water (50 ml). Solid sodium bicarbonate was added cautiously to neutralize hydrogen bromide, the organic phase separated, washed (brine, 3 x 10 ml), dried (Na₂SO₄), and evaporated to give the crude bromoketone, 3-bromo-*N,N*-diethyl-4-(4-methoxyphenyl)-4-oxobutanamide as a colourless oil (5.15 g).

Example 2 - Preparation of pyridine-based bromoacetophenones

In one reaction vessel, 2-chloro-5-acetylpyridine (1.55 g, 10 mmol) was dissolved in 45% HBr in acetic acid (30 ml) and left stirring in an ice-water bath for 20 min to form a dark brown solution. In a separate reaction vessel, bromine (1.61 g, 10 mmol) was added to HBr in acetic acid (45%) (6 ml) and the resulting solution was added to the first reaction vessel. The combined reaction mixture was left stirring in an ice-water bath for 1h. After this time, anhydrous diethyl ether was added to the reaction solution, which resulted in precipitation of the reaction. The diethyl ether was added until no further precipitate was formed (~ 60 mls). The solid was collected by filtration and washed with cold diethyl ether (dried over sodium) to yield a light yellow powder of 2-bromo-1-(6-chloropyridin-3-yl)ethanone as a HBr salt (2.44 g, 77%).

Example 3 - METHOD A1 – Condensation 1

A mixture of 2-bromo-4'-methoxyacetophenone (1.4 g, 6.13 mmol), 2-amino-3,5-dichloropyridine (1.0 g, 6.13 mmol) in absolute ethanol (15 mls) was heated to reflux. After 2 h, NaHCO₃ (309 mg, 3.68 mmol) was slowly added to the dark brown solution. The reaction mixture continued to reflux for another 15 h before another portion of NaHCO₃ was slowly added (206 mg, 2.45 mmol). After 15 mins, the reaction was cooled and kept in the fridge for 20 h. The resulting precipitate was collected by filtration, washed with cold ethanol followed by water. The solid was then boiled in hot ethanol for 10 mins before it was cooled to RT and then collected by filtration to give 6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridine as beige coloured crystals (1.34 g, 75%).

Example 4 - METHOD A2 – Condensation 2

A solution of 3-bromo-*N,N*-diethyl-4-(4-methoxyphenyl)-4-oxobutanamide (60%), 3-bromo-4-(3-bromo-4-methoxyphenyl)-*N,N*-diethyl-4-oxobutanamide (20%) and *N,N*-diethyl-4-(4-methoxyphenyl)-4-oxobutanamide (20%) (total weight 4 g) in dry DMF (10 ml) was added

portionwise over 5 h to a stirred solution of 3,5-dichloro-2-aminopyridine (1.63 g, 10.0 mmol) in DMF at 110°C; heating was continued at this temperature for 6 h, and then at 100°C for 15 h. The cooled mixture was diluted with ethyl acetate (75 ml), washed (water, 3 x 20 ml; 10% aqueous tartaric acid, 20 ml; 2% aqueous NaHCO₃, 10 ml; water, 2 x 10 ml; brine, 2 x 10 ml), dried (Na₂SO₄) and run through a short silica column (ca, 4 g) with elution being continued with ethyl acetate (30 ml). The solvent was evaporated to give a red-brown tar, which was purified by preparative HPLC on an Alltech Alltima C18, 10μ, 250 x 22mm column eluting with 80% acetonitrile and 20% ammonium bicarbonate (20mM) at 10mLs /min. The major product, eluting at 11.0 mins, was recrystallized from ethyl acetate / hexane to give white needles of 2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide (465 mg, 9.54% overall). The minor product, eluting at 13.8 mins gave white woolly needles of 2-(2-(3-bromo-4-methoxyphenyl)-6,8-dichloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide (209 mg, 3.59%) from the same solvents.

Example 5 - METHOD B – UGI reaction

A mixture of 2-amino-5-chloropyridine (200 mg, 1.56 mmol), 4-bromobenzaldehyde (346 mg, 1.87 mmol) and scandium (III) triflate (38mg, 0.078 mmol) was dissolved in methanol dichloromethane (1:3, 3 mls). After 30 mins of stirring at RT, ethyl isocyanoacetate (212 mg, 1.87 mmol) was added and the reaction continued to stir at RT for 72 hrs. After this time the reaction mixture was diluted with DCM (10 mls) and then washed with water. The water phase was extracted with DCM (2x 10 ml) and the combined organic extracts was then vigorously stirred with a solution of sodium bisulfite (2M, 25mls) for 10 mins before the organic extract was separated, dried (Na₂SO₄), filtered and the filtrate concentrated *in vacuo* to give a yellow oil. The yellow oil was crystallised from hot ethyl acetate with addition of hexane. The yellow crystals were collected by filtration to give ethyl 2-(2-(4-bromophenyl)-6-chloroimidazo[1,2-a]pyridin-3-ylamino)acetate (411 mg, 65%).

Example 6 - METHOD C1

Step 1. To a solution of 2-(3-bromophenyl)-6-chloroimidazo[1,2-a]pyridine (1.93 g, 6.3 mmol) in acetic acid (25 ml) was slowly added aqueous dimethylamine (40%, 11.8 ml, 1.2 equiv.) with stirring. The temperature rose and a precipitate formed. Aqueous formaldehyde (37%, 4.7 ml) was then added, and the mixture stirred at 55-60°C for 20h. After cooling down, the solvent was concentrated and the brown coloured residue was dissolved in a mixture of water (10 ml) and chloroform (20ml) and then 2.5 M aqueous NaOH was slowly added, with stirring, until the pH was 12. The chloroform layer was separated and the aqueous phase was further extracted with chloroform (10 ml + 2x 5 ml). The combined chloroform solutions was then extracted with 2N HCl

(3 × 10 ml). The extract was then basified with NaOH to pH 12. A yellow gummy solid formed. The solid was redissolved in EtOH and recrystallised from ethanol / water to give 2-(3-bromophenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-*N,N*-dimethylmethanamine as a beige crystalline solid (1.36 g, 60%).

5 **Step 2.** The product from Step 1 [(2-(3-bromophenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-*N,N*-dimethylmethanamine (1.3 g, 3.6 mmol)] was dissolved in benzene (15 ml) and iodomethane (0.8 ml (1.8 g), 13 mmol) was added. The reaction was left stirring in the absence of light for 18h. A white solid precipitated out, and collected by filtration and washed with benzene to give a trimethyl ammonium iodide salt (0.8 g).

10 **Step 3.** The trimethyl ammonium iodide salt from obtained from Step 2 (0.51 g, 1.1 mmol) was dissolved in EtOH/H₂O (1/1, 10 ml) and KCN (0.50 g, 7.8 mmol) was added to form a suspension. The mixture was heated to reflux for 24 h. Upon cooling, the suspension was blown with N₂ for 1h, and then evaporated to remove EtOH. Water (10 ml) was added and resulting brown solid was filtered and collected. The solid was dried in a desiccator to give the acetamide 2-(2-(3-bromophenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)acetamide as a light yellow solid (0.4 g).

Step 4. The acetamide from step 3 (0.4 g) was added to a solution of concentrated HCl (6 ml) and acetic acid (6 ml). The suspension was heated to reflux over night and an off-white precipitate formed upon cooling. The mixture was then evaporated to dryness and water (10 ml) and 10% NaOH (5 ml) was added. The resulting solution was heated for 40 min at 80 °C and then filtered.

20 The brown coloured filtrate was cooled and acidified with concentrated HCl to pH 4 to precipitate the hydrolysed product 2-(2-(3-bromophenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)acetic acid as a creamy solid (0.24 g, 60%).

Example 7 -METHOD C2

Steps 1, 2 and 3 are the same as described above in METHOD C1, however the hydrolysis in Step

25 4 was modified. Instead of using a solution of HCl / Acetic acid, a solution of HBr (48%, 200 ml) and acetic acid (100 ml) was added to the amide, for example 2-(6-chloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)acetamide (9.96g, 0.031 mol). The reaction mixture was heated to reflux for 48h where upon a white precipitate formed after cooling. The resulting crystalline solid was collected by filtration and washed with water to give the acid 2-(6-chloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)acetic acid as a hydrobromide salt (10.8g, HBr salt,

30 89%).

Example 8 - METHOD C3 – Reverse amide reaction

Concentrated sulfuric acid (18 µl) was added dropwise to a vigorously stirred mixture of (6-chloro-2-(6-chloropyridin-3-yl)imidazo[1,2-a]pyridine) (40 mg, 0.15 mmole), *N*-hydroxybenzamide (22.7 mg,

0.15 mmole) and acetic acid (1.0 ml). The cloudy solution was then heated in an oil bath at 110°C for 12 h, cooled, diluted with ethyl acetate (15 ml) and washed with excess sodium carbonate solution (2.5%). The organic layer was washed with water (2 x 3 ml) and saturated brine (2 ml), dried (Na₂SO₄), and evaporated to give an oily residue which was separated by preparative HPLC on an Alltech Alltima Column (250x22mm, 10μ), eluting at 10mls/min with 50% acetonitrile and 50% ammonium bicarbonate (20 mM). The product that eluted at 14.8 mins gave *N*-((6-chloro-2-(6-hydroxypyridin-3-yl)imidazo[1,2-*a*]pyridin-3-yl)methyl)benzamide as fine white needles (8.2 mg, 14.3% yield) after crystallisation from chloroform/ethanol. The product that eluted at 16.9 mins gave *N*-((6-chloro-2-(6-chloropyridin-3-yl)imidazo[1,2-*a*]pyridin-3-yl)methyl) benzamide (10.2 mg, 17.0%) as white needles after crystallisation from ethyl acetate / hexane.

Example 9 - METHOD D1 – Coupling (condensation) using EDCI

Diisopropylethylamine (620 μl, 3.6 mmoles) was added to a stirred mixture of (2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-*a*]pyridin-3-yl)acetic acid) (702 mg, 2.0 mmoles), *N*-methylaniline (260 μl, 2.4 mmoles) and HOBt (270 mg, 2.0 mmoles) in dry DMF (6 ml). After all solids were dissolved, EDCI (500 mg, 2.6 mmoles) was then added. Four days later, acetic acid (350 μl, 6 mmoles) and water (ca. 15 ml) were added slowly with stirring. The mixture was cooled to 5 °C for 3 h before the resulting precipitate was collected by filtration and recrystallised from chloroform / ethanol to give 2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-*a*]pyridin-3-yl)-*N*-methyl-*N*-phenylacetamide (842 mg, 95.6% yield) as white woolly needles.

Exception

Diisopropylethylamine (2.09 ml, 12.0 mmoles) was added to a stirred mixture of (2-(6-chloro-2-(4-hydroxyphenyl)imidazo[1,2-*a*]pyridin-3-yl)acetic acid) (hydrobromide, 1.1505 g, 3.0 mmoles), diethylamine (372 μl, 3.6 mmoles) and HOBt (405 mg, 3.0 mmoles) in dry DMF (8 ml). After all solids had dissolved, EDCI (748 mg, 3.9 mmoles) was added. After 18 h, acetic acid (500 μl) and water (50 ml) were added slowly with stirring. The solid product collected, after cooling the mixture to 5°C for 6 h, was washed well with water and dried. This crude product (901 mg), which contained dimers and trimers formed from condensation of carboxylic acid and phenol moieties, was stirred with dry DMF (3 ml) and diethylamine (0.5 ml) at ca. 100 °C in a sealed container for 2 h. It was heated further, with the lid removed, until all solids dissolved and most of the excess diethylamine evaporated. Ethyl acetate (10 ml) was added rapidly, and the product was allowed to crystallise. The crystals were collected after 4 h at 5 °C, washed with a small volume of cold ethyl acetate and dried to give 2-(6-chloro-2-(4-hydroxyphenyl)imidazo[1,2-*a*]pyridin-3-yl)-*N,N*-diethylacetamide (816 mg, 76.1% yield) as white woolly needles.

Example 10 - METHOD D2 – Coupling (condensation) using CDI

A mixture of 2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)acetic acid (176 mg, 0.50 mmole), carbonyl diimidazole (122 mg, 0.75 mmole) and dry DMF (1.5 ml) was stirred at room temperature for 6 h. After this time, 3-Amino-2-chloropyridine (71 mg, 0.55 mmole) was added and the reaction mixture heated at 85°C (oil bath temperature) for 16 h. Water (ca. 10 ml) was added slowly to the cooled mixture with vigorous stirring to precipitate the product as a purple-red solid, which was collected 2 h later, washed with water and recrystallized from ethanol to give 2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)-N-(6-chloropyridazin-3-yl)-N-methylacetamide (145.4 mg, 63.0%) as pale pink woolly needles.

Example 11 - METHOD E1 – Hydrolysis 1

A solution of sodium hydroxide (0.40 g, 10 mmoles) in water (10 ml) was added to a stirred mixture of methyl 2-(2-(4-(2-fluoroethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)acetate (1.45 g, 4.0 mmoles) in methanol (40 ml). The mixture was boiled under reflux for 1 h. Most of the methanol was distilled off (at atmospheric pressure) to a final volume of ca. 20 ml. The solution was cooled slightly, acetic acid (1 ml) added rapidly, and the mixture left undisturbed to crystallize the product. This was collected after being refrigerated overnight, washed with cold 1:1 methanol / water and dried to give 2-(2-(4-(2-fluoroethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)acetic acid (1.371 g, 98.4%) as white flakes.

Example 12 - METHOD E2 – Hydrolysis 2

To a solution of 2-(2-(4-(4-acetoxymethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide (180 mg, 0.38 mmol) in MeOH (3 ml) was slowly added a solution of cesium carbonate (276 mg, 0.76 mmol) in H₂O (1 ml). After 2 days of stirring at ambient temperature, a precipitate was observed and was collected by filtration. The solid was collected by filtration and washed with water. The solid was then recrystallised from hot ethyl acetate and hexanes to give 2-(2-(4-(4-hydroxymethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide as colourless crystals (146 mg, 90%).

Example 13 - METHOD F – Esterification

To stirring methanol (50 ml) was added concentrated sulphuric acid (1.5 ml) followed by 2-(6-chloro-2-(4-hydroxyphenyl)imidazo[1,2-a]pyridin-3-yl)acetic acid (hydrobromide, 1.918 g, 5.0 mmoles). The mixture was heated at 50°C for 1.25 h, cooled to room temperature and trimethyl orthoformate (530 mg, 5.0 mmoles) was added. The following day, the mixture was poured slowly into a vigorously stirred mixture of ethyl acetate (400 ml), water (100 ml) and sodium bicarbonate (6.0 g). The separated organic layer was washed (water, 2 x 50 ml; 5% aqueous sodium carbonate, 10 ml; water, 2 x 50 ml; brine, 25 ml), dried (Na₂SO₄), and concentrated (Rotavapor) until crystallization was well advanced (ca. 20 ml remaining). Hexane (20 ml) was then added

slowly. The resulting product was collected 2 h later, washed with ethyl acetate / hexane (1:1, 20 ml) and dried to give methyl 2-(6-chloro-2-(4-hydroxyphenyl)imidazo[1,2-a]pyridin-3-yl)acetate (1.4673 g, 92.7%) as white needles.

Example 14 - METHOD G – Alkylation of a phenol

5 A mixture of 2-(6-chloro-2-(4-hydroxyphenyl)imidazo[1,2-a]pyridin-3-yl)-*N,N*-diethylacetamide (71.5 mg, 0.2 mmole), 2-bromoethyl methyl ether (55.6mg, 38 μ l, 0.4 mmole), potassium carbonate (83 mg, 0.6 mmole), potassium iodide (5 mg), tetrabutylammonium iodide (1.4 mg) and dry DMF (1 ml) was stirred vigorously for 2 days. The mixture was diluted with ethyl acetate (15 ml), washed (water, 4 x 3ml; brine, 3 ml), dried (Na_2SO_4) and evaporated. The resulting residue was
10 recrystallized from ethyl acetate / hexane to give 2-(2-(4-(2-methoxyethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-*N,N*-diethylacetamide (76 mg, 91.5% yield) as pale yellow flakes.

Example 15 - METHOD H – Fluorination of an alkyl alcohol

Diisopropylethylamine (210 μ l, 1.2 mmoles), perfluoro-1-butanefluoride (69 μ l, 0.40 mmole) and triethylamine trihydrofluoride (67 μ l, 0.40 mmole) were added to a stirred mixture of 2-(2-(4-(3-
15 hydroxypropoxy)phenyl)-5,7-dimethylpyrazolo[1,5-a]pyrimidin-3-yl)-*N,N*-diethylacetamide (82 mg, 0.20 mmole) and dry acetonitrile (0.80 ml). The mixture was warmed to 40°C for 30 min and then left overnight at room temperature. Volatiles were removed, and a solution of the residual oil in ethyl acetate (10 ml) was washed (water, 3 x 3ml; 10% aqueous tartaric acid, 3 ml; water, 2 x 3 ml; brine, 3 ml), dried (Na_2SO_4) and evaporated. The residue was purified by preparative HPLC on an
20 Alltech Alltima C18 250x22mm column eluting with 70% Acetonitrile 30% water + 0.1% TFA at 10mls/min. The peak at 15.4 mins was collected, concentrated and recrystallization from ethyl acetate / hexane gave 2-(2-(4-(3-fluoropropoxy)phenyl)-5,7-dimethylpyrazolo[1,5-a]pyrimidin-3-yl)-*N,N*-diethylacetamide (48.6 mg, 59.0%) as chunky needles.

Example 16 - METHOD I – Demethylation


25 A stirred solution of (*N,N*-diethyl-2-(2-(4-methoxyphenyl)-5,7-dimethylpyrazolo[1,5-a]pyrimidin-3-yl)acetamide) (732 mg, 2.0 mmoles) and tetrabutylammonium iodide (812 mg, 2.2 mmoles) in dry dichloromethane (10 ml) was cooled in an ethanol / liquid nitrogen bath (ca. -70°C) while being flushed with N_2 . A solution of boron trichloride in hexane (1.0 M, 9.0 ml, 9.0 mmoles) was then added drop-wise *via* syringe over 5 min, with vigorous stirring. Five mins later, the ethanol / N_2 bath
30 was replaced with ice, and stirring continued for 2.5 h. The reaction was quenched by careful addition of ice and water (ca. 30 ml total) and the mixture stirred until all solid had dispersed (ca. 1 h). The semi-solid mass was shaken vigorously with water (50 ml), chloroform (50 ml) and ethanol (5 ml); solid sodium bicarbonate was added in small portions to neutralize the acids from hydrolysis of boron compounds, which was complete when the mixture was no longer purple in the presence

of excess bicarbonate. The organic layer was separated, and the aqueous phase extracted with more chloroform (2 x 20 ml). The combined organic solution was washed with dilute brine (50 ml), dried (Na_2SO_4) and evaporated to give a crystalline semi-solid, which was slurried with acetonitrile (4 ml), the crystals filtered off, washed with minimal cold acetonitrile and dried to give *N,N*-diethyl-2-(2-(4-hydroxyphenyl)-5,7-dimethylpyrazolo[1,5-*a*]pyrimidin-3-yl)acetamide (620 mg, 88.1% yield) as white needles.

Example 17 - METHOD J – Fluorination of halopyridazines

A mixture of 2-(2-(4-*tert*-butylphenyl)-6-bromoimidazo[1,2-*b*]pyridazin-3-yl)-*N,N*-diethylacetamide (66.5 mg, 0.15 mmole), potassium fluoride (87 mg, 1.50 mmoles) dimethyl sulfoxide (0.60 ml) and toluene (1.5 ml) in a 4-ml container was stirred and heated slowly to 170°C (oil bath temperature) to evaporate the toluene and remove water. The container was flushed with N_2 , sealed, and kept at 165-170°C for 7 h. The cooled mixture was diluted with ethyl acetate (3 ml), washed (water, 3 x 1 ml; brine, 1 ml), dried (Na_2SO_4) and evaporated. The semi-crystalline residue was purified by preparative HPLC on an Alltech Alltima C18 250x22mm column eluting with 80% Acetonitrile 20% water + 0.1% TFA at 20mls/min. The peak at 7.4 mins was collected, concentrated and recrystallization of the resulting solid from ethyl acetate / hexane gave 38.5 mg of 2-(2-(4-*tert*-butylphenyl)-6-fluoroimidazo[1,2-*b*]pyridazin-3-yl)-*N,N*-diethylacetamide (67.2%) as colourless plates.

Example 18 - Radiosynthesis to form alkyl fluorides.



^{18}F Fluoride, in the form of H^{18}F , was produced by the $^{18}\text{O}(\text{p},\text{n})^{18}\text{F}$ nuclear reaction by bombardment of ^{18}O -enriched water target. The aqueous ^{18}F fluoride solution was added to a 2.5- mL wheaton vial containing dry acetonitrile (1 mL), Kryptofix 2.2.2 (5 mg, 13.3 μmol) and potassium carbonate (2 mg, 14.5 μmol). The solvent was evaporated under a stream of nitrogen at 90°C with a reducing vacuum. This azeotropic drying was repeated twice by further addition of acetonitrile (2 x 1 mL). The tosylate precursor 2-(2-(4-(2-tosylethoxy)phenyl)-6-chloroimidazo[1,2-*a*]pyridin-3-yl)-*N,N*-diethylacetamide (2 mg, 3.6 μmol) was dissolved in acetonitrile (1 ml) and added to the dried $\text{K}_{222}\cdot\text{K}_2\text{CO}_3\cdot\text{KF}$ complex. Stirring and heating at 90°C continued for 10 to 15 mins before the reaction mixture was diluted with mobile phase and purified by reverse phase HPLC on an Alltech Alltima, C18, 10 μ , 250 x 22 mm column eluting at 10 mLs/min with 45% acetonitrile, 55% water with

0.1% TFA. The radioactive peak between 13 to 14 mins was collected, dried *in vacuo* and then formulated in saline for biological evaluation, typically 25 $\mu\text{Ci}/100\mu\text{l}$ saline solution per rat for biodistribution studies.

Example 19 - Characterisation data for various compounds

- 5 By the general procedures depicted in Schemes 1 to 4, and by utilising methods A - J, the following compounds were synthesised and characterised.

PBR098

N,N-diethyl-2-(2-(4-hydroxyphenyl)-5,7-dimethylpyrazolo[1,5-a]pyrimidin-3-yl)acetamide

m.p.: 234-235°C.

- 10 ^1H NMR (400 MHz, DMSO) δ 1.02 (t, $J = 7.0$ Hz, 3H, CH_3), 1.17 (t, $J = 7.0$ Hz, 3H, CH_3), 2.48 (s, 3H, CH_3), 2.69 (s, 3H, CH_3), 3.28 (q, $J = 7.0$ Hz, 1H, NCH_2), 3.52 (q, $J = 7.0$ Hz, 1H, NCH_2), 3.82 (s, 2H, CH_2), 6.82 (d, $J = 8.8$ Hz, 2H, ArCH), 6.83 (s, 1H, ArCH), 7.58 (d, $J = 8.8$ Hz, 2H, ArCH), 9.63 (br s, 1H, OH).

MS: ES(+ve) m/z 353 ($M+1$).

- 15 Elemental Analysis: Calculated for $\text{C}_{20}\text{H}_{24}\text{N}_4\text{O}_2$, C = 68.16; H = 6.86; N = 15.90. Found C = 67.97, H = 6.95, N = 15.85.

PBR099

2-(2-(4-(2-fluoroethoxy)phenyl)-5,7-dimethylpyrazolo[1,5-a]pyrimidin-3-yl)-*N,N*-diethylacetamide

m.p.: 148°C.

- 20 ^1H NMR (400 MHz, DMSO) δ 1.02 (t, $J = 7.0$ Hz, 3H, CH_3), 1.18 (t, $J = 7.0$ Hz, 3H, CH_3), 2.48 (s, 3H, CH_3), 2.69 (s, 3H, CH_3), 3.28 (q, $J = 7.0$ Hz, 1H, NCH_2), 3.52 (q, $J = 7.0$ Hz, 1H, NCH_2), 3.83 (s, 2H, CH_2), 4.29 (dm $J = 30.1$ Hz, 2H, OCH_2), 4.76 (dm, $J = 47.6$ Hz, 2H, CH_2F), 6.84 (s, 1H, ArCH), 7.06 (d, $J = 8.8$ Hz, 2H, ArCH), 7.72 (d, $J = 8.8$ Hz, 2H, ArCH).

- ^{13}C NMR (100 MHz, DMSO) δ 13.1 (CH_3), 14.2 (CH_3), 16.3 (CH_3), 24.2 (CH_3), 27.5 (CH_2), 39.81 (NCH_2), 41.6 (NCH_2), 67.1 (d, $J = 18.9$ Hz, OCH_2), 82.1 (d, $J = 66.0$ Hz, CH_2F), 100.6 (C), 108.3 (ArCH), 114.5 (ArCH), 126.3 (C), 129.3 (ArCH), 144.5 (C), 147.2 (C), 153.4 (C), 157.3 (C), 158.3 (C), 169.0 (CO).

Elemental Analysis: Calculated for $\text{C}_{22}\text{H}_{27}\text{FN}_4\text{O}_2$, C = 66.31; H = 6.83; N = 14.06. Found C = 66.47, H = 6.89, N = 14.17.

- 30 PBR101

2-(6-chloro-2-(4-hydroxyphenyl)-imidazo[1,2-a]pyridin-3-yl)-*N,N*-diethylacetamide

^1H NMR (400 MHz, DMSO), δ 1.06 (t, $J = 7.1$ Hz, 3H, CH_3), 1.16 (t, $J = 7.1$ Hz, 3H, CH_3), 3.32 (q, $J = 7.1$ Hz, 2H, NCH_2), 3.44 (q, $J = 7.1$ Hz, 2H, NCH_2), 4.18 (s, 2H, CH_2), 6.83 (d, $J = 8.8$ Hz, 2H,

ArCH), 7.26 (dd, $J = 9.4, 1.8$ Hz, 1H, ArCH), 7.41 (d, $J = 8.8$ Hz, 2H, ArCH), 7.57 (d, $J = 9.4$ Hz, 1H, ArCH), 8.49 (d, $J = 1.8$ Hz, 1H, ArCH), 9.61 (br s, 1H, OH).

^{13}C NMR (100 MHz, DMSO) δ 13.0 (CH₃), 14.11 (CH₃), 28.7 (CH₂), 39.9 (NCH₂), 41.6 (NCH₂), 115.4 (ArCH), 116.1 (C), 117.0 (ArCH), 118.3 (C), 122.9 (ArCH), 124.5 (ArCH), 125.0 (C), 129.1 (ArCH), 142.1 (C), 143.9 (C), 157.2 (C), 167.1 (CO).

MS: ES(+ve) m/z 358 (M+1).

PBR102

2-(2-(4-(2-fluoroethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide

m.p.: 142°C.

^1H NMR (400 MHz, DMSO), δ 1.06 (t, $J = 7.0$ Hz, 3H, CH₃), 1.18 (t, $J = 7.0$ Hz, 3H, CH₃), 3.34 (q, $J = 7.0$ Hz, 2H, NCH₂), 3.46 (q, $J = 7.0$ Hz, 2H, NCH₂), 4.22 (s, 2H, CH₂), 4.30 (dm, $J = 30.1$ Hz, 2H, OCH₂), 4.78 (dm, $J = 47.6$ Hz, 2H, CH₂F), 7.08 (d, $J = 9.1$ Hz, 2H, ArCH), 7.29 (dd, $J = 9.4, 2.0$ Hz, 1H, ArCH), 7.55 (d, $J = 9.1$ Hz, 2H, ArCH), 7.62 (d, $J = 9.4$ Hz, 1H, ArCH), 8.52 (d, $J = 2.0$ Hz, 1H, ArCH).

^{13}C NMR (100 MHz, DMSO) δ 13.0 (CH₃), 14.1 (CH₃), 28.7 (CH₂), 39.9 (NCH₂), 41.7 (NCH₂), 67.1 (d, $J = 18.8$ Hz, OCH₂), 82.3 (d, $J = 166.8$ Hz, CH₂F), 114.7 (ArCH), 116.5 (C), 117.1 (ArCH), 118.4 (C), 123.0 (ArCH), 124.7 (ArCH), 127.0 (C), 129.0 (ArCH), 142.2 (C), 143.3 (C), 157.9 (C), 167.0 (CO).

MS: ES(+ve) m/z 404 (M+1), 442 (M+K).

Elemental Analysis: Calculated for C₂₁H₂₃ClFNO₂, C = 62.45; H = 5.74; N = 10.40. Found C = 62.51, H = 5.79, N = 10.43.

PBR103

2-(2-(4-(2-fluoroethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-dipropylacetamide

m.p.: 158-160°C.

^1H NMR (400 MHz, DMSO) δ 0.81 (t, $J = 7.3$ Hz, 3H, CH₃), 0.85 (t, $J = 7.3$ Hz, 3H, CH₃), 1.5 (m, 2H, CH₂), 1.6 (m, 2H, CH₂), 3.25 (m, 2H, NCH₂), 3.32 (m, 2H, NCH₂), 4.23 (s, 2H, CH₂), 4.29 (dm, $J = 30.1$ Hz, 2H, OCH₂), 4.76 (dm, $J = 47.9$ Hz, 2H, CH₂F), 7.06 (d, $J = 9.0$ Hz, 2H, ArCH), 7.29 (dd, $J = 9.6, 2.0$ Hz, 1H, ArCH), 7.56 (d, $J = 9.0$ Hz, 2H, ArCH), 7.61 (d, $J = 9.6$ Hz, 1H, ArCH), 8.50 (d, $J = 2.0$ Hz, 1H, ArCH).

^{13}C NMR (100 MHz, DMSO), δ 11.0 (CH₃), 11.2 (CH₃), 20.5 (CH₂), 21.6 (CH₂), 28.8 (CH₂), 47.1 (NCH₂), 49.0 (NCH₂), 67.1 (d, $J = 19$ Hz, OCH₂), 82.3 (d, $J = 165.7$ Hz, CH₂F), 114.6 (ArCH₂), 116.4 (C), 117.1 (ArCH), 118.4 (C), 123.0 (ArCH), 124.7 (ArCH), 127.0 (C), 129.1 (ArCH), 142.2 (C), 143.3 (C), 157.9 (C), 167.5 (CO).

MS: ES(+ve) m/z 432 (M+1).

Elemental Analysis: Calculated for $C_{23}H_{27}ClFN_3O_2$; C = 63.96, H = 6.30, N = 9.73. Found, C = 64.15, H = 6.36, N = 9.78.

PBR104

2-(2-(4-(2-fluoroethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-1-(pyrrolidin-1-yl)ethanone

5 m.p.: 150-152°C.

1H NMR (400 MHz, DMSO) δ 1.84 (m, 2H, CH_2), 1.95 (m, 2H, CH_2), 3.36 (t, J = 6.7 Hz, 2H, NCH_2), 3.58 (t, J = 6.7 Hz, 2H, NCH_2), 4.15 (s, 2H, CH_2), 4.29 (dm, J = 30.1 Hz, 2H, OCH_2), 4.77 (dm, J = 47.9 Hz, 2H, CH_2F), 7.07 (d, J = 9.0 Hz, 2H, ArCH), 7.29 (dd, J = 9.4, 2.0 Hz, 1H, ArCH), 7.59 (d, J = 9.0 Hz, 2H, ArCH), 7.61 (d, J = 9.0 Hz, 1H, ArCH), 8.55 (d, J = 2.0 Hz, 1H, ArCH).

10 ^{13}C NMR (100 MHz, DMSO) δ 24.0 (CH_2), 25.7 (CH_2), 29.9 (CH_2), 45.7 (NCH_2), 46.1 (NCH_2), 67.1 (d, J = 19.0 Hz, OCH_2), 82.3 (d, J = 165.7 Hz, CH_2F), 114.7 (ArCH), 116.1 (C), 117.1 (ArCH), 118.5 (C), 123.1 (ArCH), 124.8 (ArCH), 127.0 (C), 129.1 (ArCH), 142.2 (C), 143.5 (C), 157.8 (C), 166.1 (CO).

MS: ES(+ve) m/z 402 ($M+1$).

15 Elemental Analysis: Calculated for $C_{21}H_{21}ClFN_3O_2$; C = 62.76; H = 5.27; N = 10.46. Found C = 63.02, H = 5.38, N = 10.41.

PBR105

2-(2-(4-(2-fluoroethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-dimethylacetamide

m.p.: 194-196°C.

20 1H NMR (400 MHz, DMSO), δ 2.90 (s, 3H, CH_3), 3.13 (s, 3H, CH_3), 4.21 (s, 2H, CH_2), 4.29 (dm, J = 30.1 Hz, 2H, OCH_2), 4.76 (dm, J = 47.9 Hz, 2H, CH_2F), 7.08 (d, J = 8.8 Hz, 2H, ArCH), 7.28 (dd, J = 9.6, 2.0 Hz, 1H, ArCH), 7.57 (d, J = 8.8 Hz, 2H, ArCH), 7.61 (d, J = 9.6 Hz, 1H, ArCH), 8.54 (d, J = 2.0 Hz, 1H, ArCH).

25 ^{13}C NMR (100 MHz, DMSO), δ 28.9 (CH_2), 35.3 (NCH_3), 37.0 (NCH_3), 66.1 (d, J = 19.0 Hz, OCH_2), 82.3 (d, J = 165.7 Hz, CH_2F), 114.7 (ArCH), 116.3 (C), 117.1 (ArCH), 118.5 (C), 123.0 (ArCH), 124.7 (ArCH), 127.0 (C), 129.0 (ArCH), 142.2 (C), 143.5 (C), 157.8 (C), 167.9 (CO).

MS: ES(+ve) m/z 376 ($M+1$).

Elemental Analysis: Calculated for $C_{19}H_{19}ClFN_3O_2$; C = 60.72; H = 5.10; N = 11.18. Found C = 60.60, H = 5.09, N = 11.15

30 **PBR106**

2-(2-(4-(2-fluoroethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N-methylacetamide

m.p.: 255°C.

1H NMR (400 MHz, DMSO) δ 2.65 (d, J = 4.7 Hz, 3H, NCH_3), 3.99 (s, 2H, CH_2), 4.30 (dm, J = 30.1 Hz, 2H, OCH_2), 4.77 (dm, J = 47.6 Hz, 2H, CH_2F), 7.08 (d, J = 9.0 Hz, 2H, ArCH), 7.29 (dd, J = 9.6,

2.0 Hz, 1H, ArCH), 7.62 (d, $J = 9.6$ Hz, 1H, ArCH), 7.73 (d, $J = 9.0$ Hz, 1H, ArCH), 8.21 (q, $J = 4.7$ Hz, 1H, NH), 8.61 (d, $J = 2.0$ Hz, 1H, ArCH).

^{13}C NMR (100 MHz, DMSO) δ 25.8 (CH_3), 30.7 (CH_2), 67.1 (d, $J = 19.2$ Hz, OCH_2), 82.3 (d, $J = 165.8$ Hz, CH_2F), 114.6 (ArCH), 115.8 (C), 117.2 (ArCH), 118.6 (C), 122.9 (ArCH), 124.9 (ArCH),
5 126.9 (C), 129.1 (ArCH), 142.2 (C), 143.4 (C), 157.9 (C), 168.6 (CO).

MS: ES(+ve) m/z 362 ($M+1$).

Elemental Analysis: Calculated for $\text{C}_{18}\text{H}_{17}\text{ClFN}_3\text{O}_2$; C = 59.76; H = 4.74; N = 11.61. Found C = 59.85, H = 4.77, N = 11.67.

PBR107

10 2-(2-(4-(2-fluoroethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N-ethyl-N-methylacetamide
m.p.: 117 – 118 °C.

Two rotamers observed in NMR.

^1H NMR (400 MHz, DMSO), δ 1.06 (t, $J = 7.0$ Hz, 1.5 H, CH_3), 1.17 (t, $J = 7.0$ Hz, 1.5 H, CH_3), 2.87 (s, 1.5H, NCH_3), 3.10 (s, 1.5H, NCH_3), 3.37 (q, $J = 7.0$ Hz, 1H, NCH_2), 3.47 (q, $J = 7.0$ Hz, 1H, NCH_2), 4.19 (s, 1H, CH_2), 4.23 (s, 1H, CH_2), 4.29 (dm, $J = 29.8$ Hz, 2H, OCH_2), 4.76 (dm, $J = 47.9$ Hz, 2H, CH_2F), 7.08 (dm, $J = 8.8$ Hz, 2H, ArCH), 7.28 (dd, $J = 9.4$, 2.0 Hz, 0.5H, CH), 7.29 (dd, $J = 9.4$, 2.0 Hz, 0.5H, CH), 7.55 (d, $J = 8.8$ Hz, 1H, ArCH), 7.57 (d, $J = 8.8$ Hz, 1H, ArCH), 7.61 (d, $J = 9.4$ Hz, 0.5H, CH), 7.61 (d, $J = 9.4$ Hz, 0.5H, CH), 8.52 (d, $J = 2.0$ Hz, 0.5H, CH), 8.53 (d, $J = 2.0$ Hz, 0.5H, CH).

20 ^{13}C NMR (100 MHz, DMSO), δ 12.3, 13.3 (CH_3), 28.5, 29.0 (CH_2), 32.5, 34.5 (NCH_3), 42.0, 43.8 (NCH_2), 67.1 (d, $J = 19.0$ Hz, OCH_2), 81.3 (d, $J = 165.6$ Hz, CH_2F), 114.7 (ArCH), 116.4 (C), 117.1 (ArCH), 118.4 (C), 123.0 (ArCH), 124.7 (ArCH), 127.0 (C), 129.0 (ArCH), 142.2 (C), 143.4 (C), 157.8 (C), 167.4 (CO).

Elemental Analysis: Calculated for $\text{C}_{20}\text{H}_{21}\text{ClFN}_3\text{O}_2$; C = 61.62; H = 5.43; N = 10.78. Found C = 61.34, H = 5.48, N = 10.70.

PBR110

2-(2-(4-(2-fluoroethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N-isopropylacetamide
m.p.: 205-206 °C.

30 ^1H NMR (400 MHz, DMSO) δ 1.11 (d, $J = 6.7$ Hz, 6H, 2 x CH_3), 3.89 (m, 1H, NCH), 3.96 (s, 2H, CH_2), 4.29 (dm, $J = 30.1$ Hz, 2H, OCH_2), 4.76 (dm, $J = 47.6$ Hz, 2H, CH_2F), 7.07 (d, $J = 8.8$ Hz, 2H, ArCH), 7.29 (dd, $J = 9.6$, 2.0 Hz, 1H., CH), 7.62 (d, $J = 9.6$ Hz, 1H, CH), 7.74 (d, $J = 8.8$ Hz, 2H, ArCH), 8.29 (d, $J = 7.9$ Hz, 1H, CH), 8.61 (d, $J = 2.0$ Hz, 1H, CH).

¹³C NMR (100 MHz, DMSO) δ 22.3 (CH₃), 30.7 (CH₂), 40.7 (NCH), 67.1 (d, *J* = 19.0 Hz, OCH₂), 81.3 (d, *J* = 165.6 Hz, CH₂F), 114.6 (ArCH), 116.2 (C), 117.2 (CH), 118.6 (C), 122.8 (CH), 124.8 (CH), 126.9 (C), 129.2 (ArCH), 142.1 (C), 143.5 (C), 157.9 (C), 167.1 (CO).

MS: ES(+ve) *m/z* 390 (M+1).

- 5 Elemental Analysis: Calculated for C₂₀H₂₁ClFN₃O₂, C = 61.62; H = 5.43; N = 10.78. Found C = 61.75, H = 5.51, N = 10.89.

PBR111

2-(2-(4-(3-fluoropropoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide

m.p.: 110-114°C

- 10 ¹H NMR (400 Hz, CD₃CN) δ 1.13 (t, *J* = 7.1 Hz, 3H, CH₃); 1.18 (t, *J* = 7.1 Hz, 3H, CH₃); 2.20 (m, 2H, CH₂CH₂CH₂CH₂CH₂); 3.40 (q, *J* = 7.1 Hz, 2H, CH₂CH₃); 3.44 (q, *J* = 7.1 Hz, 2H, CH₂CH₃); 4.11 (s, 2H, CH₂C=O); 4.17 (t, *J* = 6.2 Hz, 2H, OCH₂); 4.59-4.75 (dt, *J* = 5.9, 47.3 Hz, 2H, FCH₂); 7.05 (d, *J* = 8.8 Hz, 2H, ArCH₂); 7.24 (dd, *J* = 2.0, 9.5 Hz, 1H, CH), 7.54 (dd, *J* = 0.8, 9.5 Hz, 1H, CH), 7.58 (d, *J* = 8.8 Hz, 2H, ArCH₂); 8.24 (dd, *J* = 0.8, 2.0 Hz, 1H, CH).

- 15 PBR113

2-(2-(4-(3-hydroxypropoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide

m.p.: 119-120°C.

- 20 ¹H NMR (400 Hz, DMSO) δ 1.08 (t, *J* = 6.9 Hz, 3H, CH₃), 1.23 (t, *J* = 6.9 Hz, 3H, CH₃), 1.90 (m, 2H, CH₂), 3.36 (q, *J* = 6.9 Hz, 2H, CH₂CH₃, superimposed with H₂O peak), 3.45 (q, *J* = 6.9 Hz, 2H, CH₂CH₃), 3.58 (t, *J* = 6.1 Hz, 2H, HOCH₂CH₂), 4.10 (t, *J* = 6.2 Hz, 2H, OCH₂CH₂), 4.22 (s, 2H, COCH₂), 7.05 (d, *J* = 8.6 Hz, 2H, ArCH₂), 7.31 (dd, *J* = 1.7, 9.5 Hz, 1H, CH), 7.55 (d, *J* = 8.6 Hz, 2H, ArCH₂), 7.62 (d, *J* = 9.5 Hz, 1H, CH), 8.52 (d, *J* = 1.7 Hz, 1H, CH).

PBR114

2-(2-(4-(2-hydroxyethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide

- 25 ¹H NMR (400 Hz, DMSO) δ 1.07 (t, *J* = 7.1 Hz, 3H, CH₃), 1.17 (t, *J* = 7.1 Hz, 3H, CH₃), 3.34 (q, *J* = 7.1 Hz, 2H, CH₂), 3.45 (q, *J* = 7.1 Hz, 2H, CH₂), 3.73 (dt, *J* = 4.9, 5.3 Hz, 2H, CH₂OH), 4.03 (t, *J* = 5.3 Hz, 2H, CH₂O), 4.20 (s, 2H, CH₂CO), 4.87 (t, *J* = 5.5 Hz, 1H, OH), 7.03 (d, *J* = 8.8 Hz, 2H, ArCH₂), 7.29 (dd, *J* = 9.5, 2.0 Hz, 1H, CH), 7.52 (d, *J* = 8.8 Hz, 2H, ArCH₂), 7.61 (dd, *J* = 0.7, 9.5 Hz, 1H, CH), 8.50 (dd, *J* = 0.7, 2.0 Hz, 1H, CH).

- 30 ¹³C NMR (400 MHz, DMSO) δ 13.06, 14.16 (CH₃), 28.72, 39.90, 41.68 (CH₂), 59.57 (OHCH₂), 69.60 (OCH₂), 114.62 (ArCH₂), 116.41 (C), 117.13 (CH), 118.43 (C), 123.02, 124.72 (CH), 126.51 (C), 129.03 (ArCH₂), 142.22, 143.47, 158.44 (C), 167.06 (C=O).

MS: ES(+ve) *m/z* 402 (M+1).

PBR115

2-(2-(3-bromophenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide

m.p.: 154-155°C.

¹H NMR (400 MHz, DMSO) δ 1.10 (t, *J* = 7.1 Hz, 3H, CH₃), 1.23 (t, *J* = 7.1 Hz, 3H, CH₃), 3.36 (q, *J* = 7.1 Hz, 2H, CH₂), 3.52 (q, *J* = 7.1 Hz, 2H, CH₂), 4.28 (s, 2H, COCH₂), 7.35 (dd, *J* = 1.9, 9.5 Hz, 1H, CH), 7.46 (m, 1H, ArCH), 7.60 (m, 1H, ArCH), 7.68 (d, *J* = 9.5 Hz, 1H, CH), 7.66 (m, 1H, ArCH, superimposed), 7.72 (m, 1H, ArCH), 8.65 (d, *J* = 1.9 Hz, CH).

¹³C NMR (400 MHz, DMSO) δ 13.03, 14.14 (CH₃), 28.49, 39.90 (CH₂), 41.68 (COCH₂), 119.19, 119.65, 120.66, 123.58, 124.91, 127.09, 128.26, 131.68, 132.04, 132.51, 138.24, 143.34, 144.04 (C), 168.60 (C=O).

MS: ES(+ve) *m/z* 420, 422 (*M*+1).

PBR117*2-(6-chloro-2-(4-methoxyphenyl)imidazo[1,2-b]pyridazin-3-yl)-N,N-diethylacetamide*

m.p.: 138-139°C.

¹H NMR (400 MHz, CD₃CN) δ 1.11 (t, *J* = 7.1 Hz, 3H, CH₃); 1.28 (t, *J* = 7.1 Hz, 3H, CH₃); 3.40 (q, *J* = 7.1 Hz, 2H, NCH₂); 3.55 (q, *J* = 7.1 Hz, 2H, NCH₂); 3.84 (s, 3H, OCH₃); 4.17 (s, 2H, CH₂C=O); 7.03 (d, *J* = 8.9 Hz, 2H, 2 × ArCH); 7.15 (d, *J* = 9.4 Hz, 1H, CH); 7.73 (d, *J* = 8.9 Hz, 2H, 2 × ArCH); 7.93 (d, *J* = 9.4 Hz, 1H, CH).

MS: ES(+ve) *m/z* 373 (*M*+1), 395 (*M*+Na).

PBR118*2-(2-(3-bromo-4-methoxyphenyl)-6-chloroimidazo[1,2-b]pyridazin-3-yl)-N,N-diethylacetamide*

m.p.: 188°C.

¹H NMR (400 MHz, DMSO) δ 1.07 (t, *J* = 7.1 Hz, 3H, CH₃); 1.26 (t, *J* = 7.1 Hz, 3H, CH₃); 3.33 (q, *J* = 7.1 Hz, 2H, NCH₂); 3.57 (q, *J* = 7.1 Hz, 2H, NCH₂); 3.90 (s, 3H, OCH₃); 4.21 (s, 2H, CH₂C=O); 7.24 (d, *J* = 8.7 Hz, 1H, ArCH), 7.38 (d, *J* = 9.4 Hz, 1H, CH); 7.78 (dd, *J* = 8.7, 2.1 Hz, 1H, ArCH); 7.90 (d, *J* = 2.1 Hz, 1H, ArCH); 8.22 (d, *J* = 9.4 Hz, 1H, CH).

PBR119*2-(2-(4-butoxyphenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide*

m.p.: 118-119°C.

¹H NMR (400 MHz, CD₃CN) δ 0.98 (t, *J* = 7.4 Hz, 3H, CH₃); 1.11 (t, *J* = 7.1 Hz, 3H, CH₃); 1.15 (t, *J* = 7.1 Hz, 3H, CH₃); 1.50 (m, *J* = 7.4 Hz, 2H, CH₂CH₂CH₃); 1.76 (m, *J* = 7.2 Hz, 2H, OCH₂CH₂CH₃); 3.38 (m, *J* = 7.2 Hz, 4H, 2 × 2H, NCH₂CH₃); 4.03 (t, *J* = 6.5 Hz, 2H, OCH₂); 4.09 (s, 2H, CH₂C=O); 7.0 (d, *J* = 8.8 Hz, 2H, 2 × ArCH); 7.2 (dd, *J* = 9.6, 2.0 Hz, 1H, CH); 7.5 (dd, *J* = 9.6, 0.7 Hz, 1H, NCH); 7.5 (d, *J* = 8.8 Hz, 2H, 2 × ArCH); 8.21 (dd, *J* = 2.0, 0.7 Hz, 1H, NCH).

PBR120

2-(2-(4-(4-fluorobutoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide

m.p.: 102-103°C.

¹H NMR (400 MHz, CD₃CN) δ 1.15 (t, *J* = 7.1 Hz, 3H, CH₃); 1.20 (t, 3H, *J* = 7.1 Hz, 3H, CH₃); 1.9 (m, 4H, 2×CH₂, OCH₂CH₂CH₂); 3.4 (q, *J* = 7.1 Hz, 2H, CH₂CH₃); 3.45 (q, *J* = 7.1 Hz, 2H, CH₂CH₃); 4.10 (t, *J* = 6.1 Hz, 2H, OCH₂); 4.14 (s, 2H, CH₂C=O); 4.48-4.64 (dm, *J* = 47.5 Hz, 2H, FCH₂); 7.00 (d, *J* = 8.8 Hz, 2H, ArCH×2); 7.20 (dd, *J* = 9.5, 2.1 Hz, 1H, CH); 7.54 (d, *J* = 9.5 Hz, 1H, NCCH); 7.60 (d, *J* = 8.8 Hz, 2H, ArCH×2); 8.25 (d, *J* = 2.1 Hz, 1H, NCH).

MS: ES(+ve) *m/z* 432 (M+1).PBR121*2-(2-(4-tert-butylphenyl)-6-chloroimidazo[1,2-b]pyridazin-3-yl)-N,N-diethylacetamide*

m.p.: 184-185°C.

¹H NMR (400 MHz, DMSO) δ 1.07 (t, *J* = 7.08 Hz, 3H, CH₂CH₃); 1.27 (t, *J* = 7.08 Hz, 3H, CH₂CH₃); 1.32 (s, 9H, 3×CH₃); 3.33 (q, *J* = 7.08 Hz, 2H, CH₂CH₃); 3.55 (q, *J* = 7.08 Hz, 2H, CH₂CH₃); 4.21 (s, 2H, CH₂C=O); 7.36 (d, *J* = 9.40 Hz, 1H, CH=CH-C=N); 7.50 (d, *J* = 8.5 Hz, 2H, 2×ArCH); 7.5 (d, *J* = 8.5 Hz, 2H, 2×ArCH); 8.2 (d, *J* = 9.40 Hz, 1H, CH=CH-C=N).

PBR123*2-(2-(4-tert-butylphenyl)-6-iodoimidazo[1,2-b]pyridazin-3-yl)-N,N-diethylacetamide*

m.p.: 175-176°C.

¹H NMR (400 MHz, DMSO) δ 1.07 (t, *J* = 7.08 Hz, 3H, CH₂CH₃); 1.29 (t, *J* = 7.08 Hz, 3H, CH₂CH₃); 1.32 (s, 9H, 3×CH₃); 3.33 (q, *J* = 7.08 Hz, 2H, CH₂CH₃); 3.55 (q, *J* = 7.08 Hz, 2H, CH₂CH₃); 4.19 (s, 2H, CH₂C=O); 7.50 (d, *J* = 8.5 Hz, 2H, 2×ArCH); 7.52 (d, *J* = 9.40 Hz, 1H, CH=CH-C=N); 7.7 (d, *J* = 8.5 Hz, 2H, 2×ArCH); 7.9 (d, *J* = 9.40 Hz, CH=CH-C=N).

PBR124*2-(2-(4-tert-butylphenyl)-6-bromoimidazo[1,2-b]pyridazin-3-yl)-N,N-diethylacetamide*

m.p.: 174-175°C.

¹H NMR (400 MHz, DMSO) δ 1.04 (t, *J* = 7.08 Hz, 3H, CH₂CH₃); 1.24 (t, *J* = 7.08 Hz, 3H, CH₂CH₃); 1.30 (s, 9H, 3×CH₃); 3.33 (q, *J* = 7.08 Hz, 2H, CH₂CH₃); 3.52 (q, *J* = 7.08 Hz, 2H, CH₂CH₃); 4.20 (s, 2H, CH₂C=O); 7.40 (d, *J* = 9.40 Hz, 1H, CH=CH-C=N); 7.48 (d, *J* = 8.5 Hz, 2H, 2×ArCH); 7.7 (d, *J* = 8.5 Hz, 2H, 2×ArCH); 8.10 (d, *J* = 9.40 Hz, 1H, CH=CH-C=N).

PBR125*2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide*

m.p.: 180-181°C.

¹H NMR (400 MHz, DMSO) δ 1.07 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 1.17 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 3.33 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 3.44 (q, *J* = 7.1 Hz, 2H, CH₂CH₃); 3.81 (s, 3H, OCH₃), 4.22 (s,

2H, CH₂C=O), 7.04 (d, *J* = 8.9 Hz, 2H, 2×ArCH), 7.55 (d, *J* = 8.9 Hz, 2H, 2×ArCH), 7.63 (d, *J* = 1.7 Hz, 1H, CH), 8.59 (d, *J* = 1.7 Hz, 1H, CH).

¹³C NMR (400 MHz, DMSO) δ 13.05, 14.16 (CH₃), 28.91 (CH₂), 39.91, 41.69 (CH₂CH₃), 55.23 (OCH₃), 114.14 (ArCH×2), 117.54, 118.60, 121.48 (C), 122.55, 123.63 (CH), 126.01 (C), 129.15 (ArCH×2), 139.52, 143.75 (C), 159.21 (ArC-O), 166.84 (C=O)

MS: ES(+ve) *m/z* 406 (M+1).

PBR126

2-(2-(3-bromo-4-methoxyphenyl)-6,8-dichloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide

m.p.: 190-191°C.

¹H NMR(400MHz, DMSO) δ 1.09 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 1.20 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 3.35 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 3.50 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 3.90 (s, 3H, CH₃), 4.25 (s, 2H, CH₂C=O), 7.24 (d, *J* = 8.7 Hz, 2H, ArCH), 7.63 (dd, *J* = 8.7, 2.2 Hz, 1H, ArCH), 7.66 (d, *J* = 1.7 Hz, CH), 7.72 (d, *J* = 2.2 Hz, 1H, ArCH), 8.67 (d, *J* = 1.7 Hz, CH).

MS: ES(+ve) *m/z* 484, 486 (M+1).

PBR128

2-(6-bromo-2-(3-bromo-4-methoxyphenyl)imidazo[1,2-b]pyridazin-3-yl)-N,N-diethylacetamide

m.p.: 180°C.

¹H NMR(400MHz, DMSO) δ 1.07 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 1.25 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 3.33 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 3.57 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 3.90 (s, 3H, OCH₃), 4.20 (s, 2H, CH₂C=O), 7.22 (d, *J* = 6.8 Hz, 1H, ArCH), 7.43 (d, *J* = 9.4 Hz, 2H, CH), 7.78 (dd, *J* = 6.8, 2.2 Hz, 2H, 2×ArCH), 7.90 (d, *J* = 2.2 Hz, 1H, ArCH), 8.12 (d, *J* = 9.4 Hz, 1H, CH).

MS: ES(+ve) *m/z* 495, 497 (M+1).

PBR129

2-(6-bromo-2-(4-methoxyphenyl)imidazo[1,2-b]pyridazin-3-yl)-N,N-diethylacetamide

m.p.: 124-125°C.

¹H NMR(400MHz, CD₃CN) δ 1.13 (t, *J* = 7.1 Hz, 3H, CH₂CH₃); 1.32 (t, *J* = 7.1 Hz, 3H, CH₂CH₃); 3.40 (q, *J* = 7.1 Hz, 2H, CH₂CH₃); 3.57 (q, *J* = 7.1 Hz, 2H, CH₂CH₃); 3.85 (s, 3H, OCH₃), 4.15 (s, 2H, CH₂C=O); 7.04 (d, *J* = 8.9 Hz, 2H, 2×ArCH); 7.24 (d, *J* = 9.40 Hz, 1H, CH); 7.74 (d, *J* = 8.9 Hz, 2H, 2×ArCH); 7.84 (d, *J* = 9.40 Hz, 1H, CH).

MS: ES(+ve) *m/z* 417 (M+1).

PBR130

2-(2-(4-(3-fluoropropoxy)phenyl)-6,8-dichloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide

m.p.: 132-134°C.

¹H NMR (400 MHz, CD₃CN) δ 1.07 (t, *J* = 7.05 Hz, 3H, CH₃); 1.18 (t, *J* = 7.05 Hz, 3H, CH₃); 2.1 (m, 2H, CH₂CH₂CH₂CH₂CH₂); 3.34 (q, *J* = 7.05 Hz, 2H, CH₂CH₃); 3.45 (q, *J* = 7.05 Hz, 2H, CH₂CH₃); 4.1 (t, *J* = 6.3 Hz, 2H, OCH₂); 4.22 (s, 2H, CH₂C=O); 4.55-4.70 (dt, *J* = 6.1, 47.2 Hz, 2H, FCH₂); 7.1 (d, *J* = 8.8 Hz, 2H, ArCH₂); 7.54 (d, *J* = 8.8 Hz, 2H, ArCH₂); 7.63 (d, *J* = 1.8 Hz, 1H, CH); 8.59 (d, *J* = 1.8 Hz, 1H, CH).

¹³C NMR (400 MHz, DMSO) δ 13.02 (CH₃), 14.14 (CH₃), 28.88 (CH₂CH₂CH₂CH₂CH₂), 29.65, 29.84 (CH₂CH₃), 41.66 (CH₂C=O), 63.64 (OCH₂), 80.01, 81.62 (FCH₂), 114.64 (ArCH × 2), 117.52, 118.59, 121.46 (C), 122.52, 123.60 (CH), 139.49, 143.68 (C), 158.33 (ArC-O), 166.81 (C=O).

PBR131

10 *N*-butyl-2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-*a*]pyridin-3-yl)-*N*-methylacetamide
m.p.: 118-120°C.

Rotamers were observed in the NMR, 58:42 based on ¹H NMR integration.

Rotamer 1 (major):

15 ¹H NMR (400 MHz, DMSO) δ 0.89 (t, *J* = 7.2 Hz, 3H, CH₂CH₃), 1.26 (m, 2H, CH₂CH₃), 1.50 (m, 2H, CH₂CH₂CH₃), 3.10 (s, 3H, NCH₃), 3.35 (m, 2H, NCH₂), 3.81 (s, 3H, OCH₃), 4.22 (s, 2H, CH₂), 4.45 (m, 1H, CHCH₃), 7.05 (d, *J* = 8.8 Hz, 2H, ArCH₂), 7.57 (d, *J* = 8.8 Hz, 2H, ArCH₂), 7.6 (d, *J* = 1.6 Hz, 1H, CH), 8.58 (d, *J* = 1.6 Hz, 1H, CH).

20 ¹³C NMR (100 MHz, DMSO) δ 15.37 (CH₂CH₃), 21.02 (CH₂CH₃), 30.49 (CH₂CH₂CH₃), 36.62 (NCH₃), 48.39 (CH₂), 55.88 (OCH₃), 115.78 (ArCH₂), 119.24 (C), 120.17 (CH), 123.16 (ArC), 124.11 (C-Cl), 125.05 (C-Cl), 127.5 (CH), 130.79 (ArCH₂), 145.55 (C-phenyl), 160.82 (ArC-OMe), 169.02 (C=O).

Rotamer 2 (minor):

25 ¹H NMR (400 MHz, DMSO) δ 0.91 (t, *J* = 7.2 Hz, 3H, CH₂CH₃); 1.26 (m, 2H, CH₂CH₂CH₃); 1.50 (m, 2H, CH₂CH₃); 2.86 (s, 3H, NCH₃), 3.35 (m, NCH₂), 3.81 (s, 3H, OCH₃), 4.23 (s, 2H, CH₂), 3.32 (dd, *J* = 2.0, 9.5 Hz, 1H, CH), 7.03 (d, *J* = 8.8 Hz, 2H, ArCH₂), 7.55 (d, *J* = 8.8 Hz, 2H, ArCH₂), 7.64 (d, *J* = 2.0 Hz, 1H, CH), 8.59 (d, *J* = 2.0 Hz, 1H, CH).

30 ¹³C NMR (100 MHz, DMSO) δ 15.50 (CH₂CH₃), 21.10 (CH₂CH₃), 31.64 (CH₂CH₂CH₃), 34.8 (NCH₃), 48.39 (CH₂), 55.88 (OCH₃), 118.80 (ArCH₂), 119.24 (C), 120.17 (CH), 123.16 (ArC), 124.11 (C-Cl), 125.05 (C-Cl), 127.5 (CH), 130.93 (ArCH₂), 145.55 (C-phenyl), 160.82 (ArC-OMe), 169.02 (C=O).

MS: ES(+ve) *m/z* 420 (M+1).

PBR132

2-(2-(4-*tert*-butylphenyl)-6-fluoroimidazo[1,2-*b*]pyridazin-3-yl)-*N,N*-diethylacetamide
m.p.: 160-162°C

¹H NMR(400MHz, CD₃CN) δ 1.12 (t, *J* = 7.1 Hz, 3H, CH₂CH₃); 1.31 (t, *J* = 7.1 Hz, 3H, CH₂CH₃); 1.37 (s, 9H, 3×CH₃); 3.4 (q, *J* = 7.1 Hz, 2H, CH₂CH₃); 3.56 (q, *J* = 7.1 Hz, 2H, CH₂CH₃); 4.17 (s, 2H, CH₂C=O); 7.02 (d, *J* = 9.6 Hz, 1H, CH=CH-C=N); 7.56 (d, *J* = 8.5 Hz, 2H, 2×ArCH); 7.72 (d, *J* = 8.5 Hz, 2H, 2×ArCH); 8.10 (d, *J* = 9.6 Hz, 1H, CH=CH-C=N).

5 PBR133

2-(6-chloro-2-(4-methoxyphenyl)imidazo[1,2-b]pyridazin-3-yl)-N,N-diethylacetamide

m.p.: 132-134°C.

¹H NMR (400 MHz, DMSO) δ 1.05 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 1.25 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 3.30 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 3.50 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 3.80 (s, 3H, OCH₃), 4.14 (s, 2H, CH₂C=O), 7.05 (d, *J* = 8.9 Hz, 2H, 2×ArCH), 7.24 (d, *J* = 9.6 Hz, 1H, CH), 7.69 (d, *J* = 8.9 Hz, 2H, 2×ArCH), 8.32 (dd, *J* = 9.6, 7.5 Hz, CH-CF).

¹³C NMR (400 MHz, DMSO) δ 13.06, 14.22 (CH₃), 28.46, 39.95, 41.69 (CH₂), 55.21(OCH₃), 106.93, 107.30 (CHCF), 114.18 (ArCH×2), 120.85 (C), 126.31 (C), 128.72 (ArCH×2), 129.26, 129.37 (CHCHCF), 136.82 (C), 144.00 (CCHCHCF), 158.72, 161.00 (CF), 166.57 (C=O).

15 PBR134

2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)-N-methyl-N-phenylacetamide

m.p.: 186-188°C.

¹H NMR (400 MHz, DMSO) δ 3.24 (s, 3H, NCH₃), 3.80 (s, 3H, OCH₃), 3.88 (s, 2H, CH₂), 7.03 (d, *J* = 8.7 Hz, 2H, ArCH×2), 7.43 (d, *J* = 8.7 Hz, 2H, ArCH×2), 7.3-7.57 (m, 5H, ArCH), 7.61 (d, *J* = 1.6 Hz, 1H, CH), 8.72(d, *J* = 1.6 Hz, 1H, CH)

MS: ES(+ve) *m/z* 440 (M+1).

PBR135

N-benzyl-2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)-N-methylacetamide

m.p.: 164-167°C.

25 Rotamers were observed in the NMR, 60:40 based on ¹H NMR integration.

Rotamer 1 (major):

¹H NMR (400 MHz, DMSO) δ 3.09 (s, 3H, NCH₃), 3.80 (s, 3H, OCH₃), 4.33 (s, 2H, CH₂), 4.56 (CH₂-phenyl), 7.01 (d, *J* = 8.8 Hz, 2H, ArCH×2), 7.2-7.45 (m, 5H, ArCH), 7.57 (d, *J* = 8.8 Hz, 2H, ArCH×2), 7.62 (d, *J* = 1.7 Hz, 1H, CH), 8.65(d, *J* = 1.7 Hz, 1H, CH).

30 Rotamer 2 (minor):

¹H NMR (400 MHz, DMSO) δ 2.94 (s, 3H, NCH₃), 3.78 (s, 3H, OCH₃), 4.23 (s, 2H, CH₂), 4.72 (CH₂-phenyl), 6.93 (d, *J* = 8.8 Hz, 2H, ArCH×2), 7.2-7.45 (m, 5H, ArCH), 7.42 (d, *J* = 8.8 Hz, 2H, ArCH×2), 7.62 (d, *J* = 1.7 Hz, 1H, CH), 8.57(d, *J* = 1.7 Hz, 1H, CH).

MS: ES(+ve) *m/z* 454 (M+1).

PBR136

2-(2-(4-(2-fluoroethoxy)phenyl)-6,8-dichloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide

m.p.: 144-146°C.

¹H NMR (400 MHz, CD₃CN) δ 1.07 (t, J = 7.1 Hz, 3H, CH₃), 1.18 (t, J = 7.1 Hz, 3H, CH₃), 3.34 (q, J = 7.1 Hz, 2H, CH₂CH₃), 3.45 (q, J = 7.1 Hz, 2H, CH₂CH₃), 4.23 (s, 2H, CH₂C=O); 4.30 (dm, J = 30.1 Hz, 2H, OCH₂), 4.82 (dm, J = 47.9 Hz, 1H, FCH₂), 7.08 (d, J = 8.9 Hz, 2H, ArCHx2), 7.56 (d, J = 8.9 Hz, 2H, ArCHx2), 7.63 (d, J = 1.7 Hz, 1H, CH), 8.59 (d, J = 1.7 Hz, 1H, CH).

PBR137

2-(2-(4-tert-butylphenyl)-6,8-dichloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide

m.p.: 160°C.

¹H NMR (400 MHz, DMSO) δ 1.06 (t, J = 7.1 Hz, 3H, CH₂CH₃), 1.17 (t, J = 7.1 Hz, 3H, CH₂CH₃), 1.32 (s, 9H, 3xCH₃), 3.30 (q, J = 7.1 Hz, 2H, CH₂CH₃), 3.45 (q, J = 7.1 Hz, 2H, CH₂CH₃), 4.25 (s, 2H, CH₂C=O), 7.50 (d, J = 8.6 Hz, 2H, 2xArCH), 7.55 (d, J = 8.6 Hz, 2H, 2xArCH); 7.64 (d, J = 1.7 Hz, 1H, CH); 8.60 (d, J = 1.7 Hz, CH).

¹³C NMR (400 MHz, DMSO) δ 13.06, 14.14 (CH₃), 28.90 (CH₂), 31.08 (CH₃x3), 34.39 (CH₂), 41.69 (CH₂), 117.63, 119.16, 121.62 (C), 122.62 (ArCHx2), 123.76 (ArCHx2), 130.81, 143.71, 150.57 (ArC-O), 166.81 (C=O).

PBR138

N-((6-chloro-2-(6-chloropyridin-3-yl)imidazo[1,2-a]pyridin-3-yl)methyl)benzamide

m.p.: 288-290°C.

¹H NMR (400 MHz, DMSO) δ 5.0 (d, J = 5.4 Hz, 2H, CH₂NH); 7.29 (dd, J = 2.0, 9.6 Hz, 1H, Cl-C-CH); 7.4-7.6 (m, 5H, ArCH), 7.8 (d, J = 9.6 Hz, 1H, Cl-C-CH=CH, superimposed with d, J = 8.3 Hz, 1H, pyridyl-CH), 8.27 (dd, J = 2.5, 8.3 Hz, 1H, pyridyl-CH), 8.64 (d, J = 2.0 Hz, 1H, Cl-C=CH), 8.9 (d, J = 2.5 Hz, 1H, pyridyl-CH-N).

MS: ES(+ve) m/z 397 (M+1).

PBR139

N-((2-(4-tert-butylphenyl)-6-fluoroimidazo[1,2-b]pyridazin-3-yl)methyl)benzamide

m.p.: 237°C.

¹H NMR (400 MHz, DMSO) δ 1.35 (s, 9H, CH₃x3), 5.06 (d, J = 5.0 Hz, 2H, CH₂NH); 7.06 (d, J = 9.6 Hz, 1H, CH); 7.4-7.5 (m, 3H, ArCH), 7.55 (d, J = 8.5 Hz, 2H, ArCHx2), 7.72 (m, 2H, ArCH), 7.9 (d, J = 8.5 Hz, 2H, ArCHx2), 8.1 (dd, J = 7.4, 9.6 Hz, F-C-CH)

MS: ES(+ve) m/z 303 (M+1).

PBR140

2-(2-(4-(allyloxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide

m.p.: 126°C.

¹H NMR (400 MHz, DMSO) δ 1.05 (t, *J* = 7.1 Hz, 3H, CH₃), 1.15 (t, *J* = 7.1 Hz, 3H, CH₃), 3.3 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 3.4 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 4.2 (s, 2H, CH₂), 4.6 (ddd, *J* = 1.46, 1.46, 5.3 Hz, 2H, OCH₂), 5.25 (ddt, *J* = 1.46, 1.75, 10.5 Hz, 1H, CH), 5.4 (ddt, *J* = 1.46, 17.2 Hz, 1H, CH), 6.05 (ddt, *J* = 10.5, 17.2, 5.2 Hz, 1H, CH).

PBR141

2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)-N-phenylacetamide

m.p.: 220-222°C.

¹H NMR (400 MHz, DMSO) δ 3.80 (s, 3H, OCH₃), 4.29 (s, 2H, CH₂), 7.05 (d, *J* = 8.8 Hz, 2H, ArCH₂), 7.06 (m, 1H, ArCH, superimposed), 7.3-7.60 (m, 4H, ArCH), 7.66 (d, *J* = 1.7 Hz, 1H, CH), 7.70 (d, *J* = 8.8 Hz, 2H, ArCH₂), 8.72 (d, *J* = 1.7 Hz, 1H, CH), 10.40 (s, 1H, NH).

¹³C NMR (100 MHz, DMSO) δ 31.79 (CH₂), 55.20 (OCH₃), 114.21 (ArCH₂), 117.65, 117.78 (C), 119.29 (ArCH₂), 121.55 (C), 122.58, 123.52, 123.91 (CH), 125.88 (C), 128.80, 129.18 (ArCH₂), 138.90, 139.63 (C), 143.99 (ArC-NH), 159.23 (ArC-O), 166.91 (C=O).

PBR142

N-((6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)methyl)benzamide

m.p.: 260-263°C.

¹H NMR (400 MHz, DMSO) δ 3.80 (s, 3H, OCH₃), 4.94 (s, 2H, CH₂), 7.05 (d, *J* = 8.8 Hz, 2H, ArCH₂), 7.4-7.6 (m, 3H, ArCH), 7.68 (d, *J* = 1.6 Hz, CH), 7.85 (m, 2H, ArCH), 7.88 (d, *J* = 8.8 Hz, 2H, ArCH₂), 8.82 (d, *J* = 1.7 Hz, 1H, CH).

MS: ES(+ve). *m/z* 426 (M+1).

PBR143

2-(2-(4-(2-fluoroethoxy)phenyl)-6,8-dichloroimidazo[1,2-a]pyridin-3-yl)-N-methyl-N-phenylacetamide

m.p.: 189-190°C

¹H NMR (400 MHz, DMSO) δ 3.25 (s, 3H, NCH₃), 3.90 (s, 2H, CH₂C=O), 4.32 (dm, *J* = 30.1 Hz, 2H, OCH₂), 4.70-4.85 (dm, *J* = 47.8 Hz, 2H, FCH₂), 7.07 (d, *J* = 8.7 Hz, 2H, ArCH₂), 7.42 (d, *J* = 8.7 Hz, 2H, ArCH₂), 7.32-7.58 (m, 5H, ArCH), 7.63 (d, *J* = 1.8 Hz, 1H, CH); 8.74 (d, *J* = 1.8 Hz, 1H, CH).

PBR144

2-(2-(4-(2-methoxyethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide

m.p.: 134-135°C.

¹H NMR (400 MHz, DMSO) δ 1.06 (t, *J* = 7.1 Hz, 3H, CH₃), 1.16 (t, *J* = 7.1 Hz, 3H, CH₃), 3.29 (s, 3H, OCH₃), 3.32 (q, *J* = 7.1 Hz, 2H, NCH₂), 3.47 (q, *J* = 7.1 Hz, 2H, NCH₂), 3.67 (m, 2H, OCH₂), 4.14 (m, 2H, OCH₂), 4.21 (s, 2H, CH₂), 7.03 (d, *J* = 8.9 Hz, 2H, ArCH₂), 7.30 (dd, *J* = 9.5, 2.0 Hz,

1H, CH), 7.50 (d, $J = 8.9$ Hz, 2H, ArCH), 7.60 (dd, $J = 9.4, 0.7$ Hz, 1H, CH), 8.46 (dd, $J = 2.0, 0.7$ Hz, 1H, CH).

PBR145

3-(2-(4-bromophenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N,N-diethylpropanamide

5 m.p.: 119-120°C.

¹H NMR (400 MHz, DMSO), δ 0.90 (t, $J = 7.1$ Hz, 3H, CH₃), 0.98 (t, $J = 7.1$ Hz, 3H, CH₃), 3.15 (q, $J = 7.1$ Hz, 2H, NCH₂), 3.25 (q, $J = 7.1$ Hz, 2H, NCH₂), 3.82 (d, $J = 5.6$ Hz, 2H, NHCH₂), 5.23 (t, $J = 5.6$ Hz, 1H, NHCH₂), 7.20 (dd, $J = 9.5, 2.0$ Hz, 1H, CH), 7.50 (dd, $J = 9.5, 0.8$ Hz, 1H, CH), 7.61 (d, $J = 8.6$ Hz, 2H, ArCH₂), 8.05 (d, $J = 8.6$ Hz, 2H, ArCH₂), 8.65 (dd, $J = 2.0, 0.8$ Hz, 1H, CH).

10 PBR146

2-(2-(4-(3-fluoropropoxy)phenyl)-5,7-dimethylpyrazolo[1,5-a]pyrimidin-3-yl)-N,N-diethylacetamide

m.p.: 106-108°C.

¹H NMR (400 MHz, DMSO) δ 1.02 (t, $J = 7.1$ Hz, 3H, CH₃), 1.18 (t, $J = 7.1$ Hz, 3H, CH₃), 2.14 (dm, $J = 25.7$ Hz, 2H, CH₂), 2.49 (s, 3H, CCH₃), 2.69 (s, 3H, CCH₃), 3.29 (q, $J = 7.1$ Hz, 2H, NCH₂), 3.52 (q, $J = 7.1$ Hz, 2H, NCH₂), 3.84 (s, 2H, CH₂C=O), 4.13 (t, $J = 6.3$ Hz, 2H, OCH₂), 4.55-4.70 (dt, $J = 47.3, 5.9$ Hz, 2H, FCH₂), 6.84 (s, 1H, CH), 7.05 (d, $J = 8.9$ Hz, 2H, ArCH₂), 7.70 (d, $J = 8.9$ Hz, 2H, ArCH₂).

¹³C NMR (400 MHz, DMSO) δ 13.08, 14.25 (CH₃CH₂), 16.27, 24.20 (CH₃C), 27.50 (CH₂C=O), 29.66, 29.86 (OC(CH₂)CF), 39.82, 41.57 (CH₃CH₂), 63.56, 63.61 (O CH₂), 80.04, 81.65 (FCH₂), 100.61 (C), 108.33 (CH), 114.45 (ArCH₂), 126.07 (C), 129.28 (ArCH₂), 144.55, 147.19, 153.49, 157.34, 158.57 (C), 169.05 (C=O).

20 PBR147

2-(2-(4-(3-fluoropropoxy)phenyl)-6,8-dichloroimidazo[1,2-a]pyridin-3-yl)-N-methyl-N-phenylacetamide

25 m.p.: 128°C.

¹H NMR (400 MHz, DMSO) δ 2.14 (dm, $J = 25.8$ Hz, 2H, CH₂), 3.30 (s, 3H, NCH₃), 3.90 (s, 2H, CH₂C=O), 4.14 (t, $J = 6.3$ Hz, 2H, OCH₂), 4.48-4.64 (dt, $J = 47.3, 5.9$ Hz, 2H, FCH₂), 7.07 (d, $J = 8.7$ Hz, 2H, ArCH₂), 7.45 (d, $J = 8.7$ Hz, 2H, ArCH₂), 7.34-7.60 (m, 5H, ArCH), 7.63 (d, $J = 1.7$ Hz, 1H, CH); 8.74 (d, $J = 1.7$ Hz, 1H, CH).

¹³C NMR (400 MHz, DMSO) δ 29.66, 29.86 (CH₂CF), 30.24 (CH₂), 37.12 (NCH₃), 63.61, 63.67 (OCH₂CCF), 80.06, 81.67 (FCH₂), 114.71 (ArCH), 117.51, 118.12, 121.40 (C), 122.84 (CH), 123.79 (CH), 125.78 (C), 127.34 (CH), 127.98 (CH), 128.93 (ArCH), 129.86 (ArCH), 139.58 (C), 143.26 (N-ArC), 158.30 (O-ArC), 167.59 (C=O).

PBR149

2-(2-(4-(2-fluoroethoxy)phenyl)-6-chloroimidazo[1,2-a]pyridin-3-yl)-N-methyl-N-phenylacetamide

m.p.: 198-200°C

¹H NMR (400 MHz, DMSO) δ 3.26 (s, 3H, NCH₃), 3.90 (s, 2H, CH₂), 4.3 (dm, *J* = 30 Hz, 2H, OCH₂), 4.8 (dm, *J* = 48 Hz, 2H, FCH₂), 7.06 (d, *J* = 8.5 Hz, 2H, ArCH_{x2}), 7.28 (dd, *J* = 1.9, 9.5 Hz, 1H, CH), 7.3-7.55 (m, 5H, ArCH), 7.57 (d, *J* = 8.5 Hz, 2H, ArCH_{x2}), 7.57 (d, *J* = 8.5 Hz, 1H, CH, superimposed), 8.68 (d, *J* = 1.9 Hz, 1H, CH).

¹³C NMR (400 MHz, DMSO) δ 29.95 (CH₃), 37.12 (CH₂), 67.03, 67.22 (OCH₂), 81.33, 82.99 (FC H₂), 114.68 (ArCH_{x2}), 116.06 (ArCH_{x2}), 117.06 (CH), 118.445 (C), 123.32 (CH), 124.94 (CH), 126.60 (C), 127.34 (CH), 127.95 (C), 128.83 (ArCH_{x2}), 129.86 (ArCH_{x2}), 142.28 (N-ArC), 142.90 (C), 143.34 (C), 157.81 (O-ArC), 167.85 (C=O).

PBR153*2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)-N-(6-chloropyridazin-3-yl)-N-methylacetamide*

m.p.: 178-179°C.

¹H NMR (400 MHz, DMSO) δ 3.57 (s, 3H, CH₃), 4.43 (s, 2H, CH₂), 7.04 (d, *J* = 8.9 Hz, 2H, ArCH_{x2}), 7.59 (d, *J* = 8.9 Hz, 2H, ArCH_{x2}), 7.65 (d, *J* = 1.7 Hz, 1H, CH), 7.93 (d, *J* = 9.2 Hz, 1H, CH), 8.03 (d, *J* = 9.2 Hz, 1H, CHCl), 8.79 (d, *J* = 1.7 Hz, 1H, CH).

PBR154*2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)-N-(2-chloropyridin-3-yl)acetamide*

m.p.: 110-112°C.

¹H NMR (400 MHz, DMSO) δ 3.82 (s, 3H, CH₃), 4.47 (s, 2H, CH₂), 7.06 (d, *J* = 8.9 Hz, 2H, ArCH_{x2}), 7.44 (dd, *J* = 4.7, 8.0 Hz, 1H, CH), 7.67 (d, *J* = 1.7 Hz, 1H, CH), 7.75 (d, *J* = 8.9 Hz, 2H, ArCH_{x2}), 8.18 (dd, *J* = 1.8, 8.0 Hz, 1H, CH), 8.23 (dd, *J* = 1.8, 4.7 Hz, 1H, CH), 8.81 (d, *J* = 1.7 Hz, 1H, CH), 10.20 (s, 1H, NH).

PBR155*2-(6-chloro-2-(6-chloropyridin-3-yl)imidazo[1,2-a]pyridin-3-yl)-N,N-diethylacetamide*

¹H NMR (400 MHz, DMSO) δ 1.05 (t, *J* = 7.1 Hz, 3H, CH₃), 1.16 (t, *J* = 7.1 Hz, 3H, CH₃), 3.36 (q, *J* = 7.1 Hz, 2H, CH₂), 3.52 (q, *J* = 7.1 Hz, 2H, CH₂), 4.33 (s, 2H, COCH₂), 7.37 (dd, *J* = 2.0, 9.6 Hz, 1H, CH), 7.66 (d, *J* = 8.3 Hz, 1H, CH), 7.69 (d, *J* = 9.6 Hz, 1H, CH), 8.10 (dd, *J* = 2.5, 8.3 Hz, 1H, CH), 8.62 (s, 1H, CH, superimposed), 8.62 (s, 1H, CH, super-imposed).

¹³C NMR (400 MHz, DMSO) δ 12.99, 14.14 (CH₃), 28.36, 39.88, 41.67 (CH₂), 117.55 (CH), 118.45, 119.12 (C), 123.29, 124.39, 125.68 (CH), 129.58 (C), 138.40 (CH), 139.25, 142.67 (C), 148.10 (CH), 149.21 (C), 166.69 (C=O)

MS: ES(+ve) *m/z* 377 (M+1).

PBR156

2-(6,8-dichloro-2-(4-methoxyphenyl)imidazo[1,2-a]pyridin-3-yl)-N-(6-chloropyridin-3-yl)acetamide

m.p.: 150-154°C.

¹H NMR (400 MHz, DMSO) δ 3.80 (s, 3H, OCH₃), 4.32 (s, 2H, CH₂), 7.06 (d, *J* = 8.9 Hz, 2H, ArCH_x2), 7.48 (d, *J* = 8.7 Hz, 1H, CH), 7.66 (d, *J* = 1.8 Hz, 1H, CH), 7.67 (d, *J* = 8.9 Hz, 2H, ArCH_x2), 8.09 (dd, *J* = 2.8, 8.7 Hz, 1H, CH), 8.61 (d, *J* = 2.8 Hz, 1H, CH), 8.81 (d, *J* = 1.8 Hz, 1H, CH).

¹³C NMR (400 MHz, DMSO) δ 31.76 (CH₂), 55.21 (OCH₃), 114.26 (ArCH_x2), 117.11, 117.88, 121.57 (C), 122.63, 124.04, 124.28 (CH), 125.80 (C), 129.20 (ArCH_x2), 130.02 (CH), 135.19, 139.71 (C), 140.50 (CH), 143.83, 144.15, 159.26 (C), 167.65 (C=O)

MS: ES(+ve) *m/z* 461, 463 (M+1).

PBR158

N-(6-(1H-imidazol-1-yl)pyridazin-3-yl)-2-(6-chloro-2-(4-iodophenyl)imidazo[1,2-a]pyridin-3-yl)-N-methylacetamide

m.p.: 218-219 °C.

¹H NMR (400 MHz, DMSO) δ 3.55 (s, 3H, NCH₃), 4.48 (s, 2H, CH₂), 7.21 (b, 1H, CH-imidazole), 7.32 (dd, *J* = 2.0, 9.6 Hz, 1H, CH-pyridine), 7.46 (d, *J* = 8.4 Hz, 2H, ArCH_x2), 7.62 (dd, *J* = 9.6 Hz, 1H, CH-pyridine), 7.81 (d, *J* = 8.4 Hz, 2H, ArCH_x2), 8.05 (dd, *J* = 1.2, 1.6 Hz, 1H, CH-imidazole, not fully resolved), 8.10 (d, *J* = 9.4 Hz, 1H, CH-pyridazine), 8.25 (d, *J* = 9.4 Hz, 1H, CH-pyridazine), 8.62 (b, 1H, CH-imidazole), 8.74 (d, *J* = 2.0 Hz, 1H, CH-pyridine, not fully resolved).

PBR159

2-(6-chloro-2-(4-iodophenyl)imidazo[1,2-a]pyridin-3-yl)-N-(2-chloropyridin-3-yl)-N-methylacetamide

m.p.: 258 °C.

Rotamers were observed in NMR.

Major:

¹H NMR (400 MHz, DMSO) δ 3.17 (s, 3H, NCH₃), 4.46 (ABq, *J* = 17.5, 2H, CH₂), 7.32 (dd, *J* = 2.0, 9.6 Hz, 1H, CH), 7.36 (d, *J* = 8.5 Hz, 2H, ArCH_x2), 7.55 (dd, *J* = 4.7, 7.9 Hz, 1H, Cl-Py-CH), 7.64 (d, *J* = 9.6 Hz, 1H, CH), 7.80 (d, *J* = 8.5 Hz, 2H, ArCH_x2), 8.31 (dd, *J* = 1.8, 7.9 Hz, 1H, Cl-Py-CH), 8.44 (dd, *J* = 1.8, 4.7 Hz, 1H, Cl-Py-CH), 8.60 (d, *J* = 2.0 Hz, 1H, CH).

¹³C NMR (400 MHz, DMSO) δ 30.39 (CH₂), 35.72 (NCH₃), 94.15 (ArC-I), 116.03 (C), 117.48 (CH), 118.93 (C), 123.22, 124.65, 125.56 (CH), 129.76 (ArCH_x2), 133.27, 136.63 (C), 137.35 (ArCH_x2), 140.00 (CH), 142.52, 142.65, 148.71 (C), 149.63 (CH), 167.38 (C=O).

minor:


¹H NMR (400 MHz, DMSO) δ 3.48 (s, 3H, NCH₃), 4.53 (m, 2H, CH₂), 7.34 (m, 1H, CH, superimposed), 7.51 (d, J = 8.5 Hz, 2H, ArCH_x2), 7.53 (m, 1H, Cl-Py-CH, superimposed), 7.65 (d, J = 9.6 Hz, 1H, CH), 7.85 (d, J = 8.5 Hz, 2H, ArCH_x2), 7.94 (dd, J = 1.8, 7.9, Hz, 1H, Cl-Py-CH), 8.38 (dd, J = 1.8, 4.7 Hz, 1H, Cl-Py-CH), 8.62 (m, 1H, CH).

¹³C NMR (400 MHz, DMSO) δ 29.24 (CH₂), 37.62 (NCH₃), 94.15 (ArC-I), 116.35 (C), 117.48 (CH), 118.93 (C), 123.16, 124.23, 125.44 (CH), 129.86 (ArCH_x2), 133.61(C), 137.44 (ArCH_x2), 137.95 (C), 139.05(CH), 142.49, 143.00 (C), 148.35 (CH), 148.53 (C), 168.49 (C=O).

Example 20

IN VIVO EVALUATION OF THE [¹⁸F] LABELLED IMIDAZOPYRIDAZINE PBR-132*,

Synthesis and evaluation of the imidazopyridazine: 2-(2-(4-*tert*-butylphenyl)-6-fluoroimidazo[1,2-*b*]pyridazin-3-yl)-*N,N*-diethylacetamide **PBR132***.



PBR132*


[¹⁸F]**PBR132*** has been prepared by nucleophilic substitution of the Br precursor with ¹⁸F-fluoride in the presence of K222 and K₂CO₃ in DMF at 150°C for 5 mins.

The biodistribution of **PBR132*** was undertaken in SD rats and analysis up to 4h p.i. in the brain and peripheral tissues was performed. The specificity and selectivity of the tracer were assessed by pre-treatment with PBR and Central benzodiazepine receptor (CBR) specific ligands (1 mg/kg) 5 min prior to injection of **PBR132***.

In vitro binding of **1** indicated an IC₅₀ of 29 nM for PBR and 340 nM for the CBR. [¹⁸F]**1** was synthesised in 40-50% radiochemical yield (unoptimised) >95% radiochemical purity and specific activity of 40-80 GBq/μmol. The in vivo biodistribution of **PBR132*** showed high uptake in tissues of known PBR. In the adrenals was found an uptake of 13 % ID/g at 30 min p.i and maintained over the 4 h. In kidney, heart and lung the activity peaked (4, 8 and 16 % ID/g) at 15 min p.i. and decreased over time to less than 2.3% ID/g at 4h. Bone uptake ranged from 1 (at 15 min) to 3.3% ID/g at 4h. The uptake in the olfactory bulbs ranged from an initial 0.63 % at 15 min to 0.25 % at 4h p.i. . The concentration in the blood was low throughout the time of measurement. Pre-treatment with PK 11195 and Ro 5-4864 decreased the uptake in the brain and peripheral organs except in the adrenals which showed an activity increase. Flumazenil had no effect in the uptake of **PBR132*** in the brain or peripheral organs.

Example 21**SYNTHESIS AND EVALUATION OF THE [¹⁸F] LABELLED IMIDAZOPYRIDINE PBR 147***

Synthesis and evaluation of the fluorine-18 imidazopyridine analogue 2-(2-(4-(3-fluoropropoxy)phenyl)-6,8-dichloroimidazo[1,2-a]pyridin-3-yl)-N-methyl-N-phenylacetamide 1.

**PBR 147***

PBR 147* has been prepared by nucleophilic substitution of the tosylate precursor with ¹⁸F-fluoride in the presence of K₂₂₂, K₂CO₃ in ACN at 100°C for 5 mins.

The biodistribution of **PBR 147*** was performed in SD rats and brain and peripheral tissues were analysed at 15, 30 min, 1, and 4 h p.i.. The specificity and selectivity of the tracer was assessed by pre-treatment of the animals with the PBR ligands PK11195 and Ro 5-4864 and with Flumazenil for CBR at 1 mg/kg 5 min prior to injection of **PBR 147***


In vitro binding of **PBR147** indicated an IC₅₀ of 7.4 nM for PBR and a low IC₅₀ of >4000 nM for the CBR. [¹⁸F]**147*** was synthesised in 40-55 % radiochemical yield non-decay corrected and with > 95 % radiochemical purity. The specific activity ranged from 37-80 GBq/μmol. The in vivo biodistribution showed uptake of **PBR 147*** in tissue rich in PBR. The adrenals showed high uptake, increasing from 9 % ID/g at 30 min p.i to 11 % at 4 h. In the kidneys the activity peaked at 15 min p.i.(5 % ID/g) and decreased over time to 4 and 3.6 % ID/g at 4h. In the heart the uptake was maintained at 7% throughout of the time of the experiment (15 min to 4 h). Bone uptake ranged from 1 (at 15 min) to 2.9 % ID/g at 4h. The uptake in the olfactory bulbs ranged from an initial 0.66% at 15 min to 0.26% at 4h p.i. while the concentration in the blood was significantly lower. Pre-treatment with PK 11195 and Ro 5-4864 decreased the uptake in the brain and peripheral organs except in the adrenals which showed an increase of **PBR 147*** uptake . Flumazenil had no effect in the uptake of **PBR 147*** in the brain or peripheral organs.

Example 22**SYNTHESIS AND EVALUATION OF [¹⁸F] LABELLED IMIDAZOPYRIDINES PBR102* and PBR111***

Synthesis and evaluation of two fluorine-18 imidazopyridine analogues **PBR102*** and **PBR111***.

Tracers **PBR102*** and **PBR111*** were prepared by nucleophilic substitution of the corresponding ethyl and propyl tosylate precursors with ^{18}F -fluoride in the presence of K_{222} , K_2CO_3 in ACN at 100°C for 5 mins.

The biodistribution of **PBR102*** and **PBR111*** were performed in SD rats and brain and peripheral tissues analysed at 15, 30 min, 1, and 4 h p.i. The specificity and selectivity of the tracers were assessed by pre-treatment with the PBR ligands PK11195 and Ro 5-4864 and with Flumazenil for CBR at 1 mg/kg 5 min prior to injection of the corresponding tracers.



PBR102* R = $\text{CH}_2\text{CH}_2^{18}\text{F}$

PBR111* R = $\text{CH}_2\text{CH}_2\text{CH}_2^{18}\text{F}$

The IC_{50} values of **PBR102** and **PBR111** are 13.2 and 7.5 for the PBR and >1500 nM for the CBR. **PBR102*** and **PBR111*** were synthesised in 40-55% radiochemical yield non-decay corrected and with > 95 % radiochemical purity. The specific activity ranged from 37-80 GBq/ μmol (un-optimised). The in vivo biodistribution of both tracers showed radioactivity uptake in tissue rich in PBR. The uptake of **PBR102***


in the olfactory bulbs ranged from 0.56 % at 15 min to 0.40 % ID/g at 4 h p.i whilst for **PBR111*** it ranged from 0.64 % to 0.41 % at 15 min and 4 h respectively. Whilst blood activity levels for **PBR102*** remained constant (0.20%) over the 4 h study period **PBR111*** showed a decrease from 0.4 at 15 min to 0.1 at 4 h. The adrenals showed high uptake of **PBR102*** increasing from 9 % ID/g at 30 min p.i to 11 % at 4 h whilst **PBR111*** increased from 6.5 at 30 min to 16 % at 4 h. In the kidneys the activity peaked at 15 min p.i. for both tracers (4.5% ID/g) for **PBR102*** and (3.2 % ID/g) for **PBR111*** decreasing to 1.3 and 2.3% at 4 h. Bone uptake for **PBR102*** ranged from 0.4 % at 15 min to 1.0 % at 4 h whereas for **PBR111*** was 0.87 to 4.3 % for the same period. Pre-treatment with PK 11195 and Ro 5-4864 decreased the uptake in peripheral organs except in the adrenals which showed an increase of **PBR102*** and **PBR111*** uptake. In the olfactory regions a non-significant decrease was observed with **PBR111*** Flumazenil had no effect in the uptake of **PBR102*** or **PBR111*** in the brain or peripheral organs.

Example 23

SYNTHESIS AND EVALUATION OF [^{18}F] LABELLED PYRAZOLOPYRIMIDINES **PBR099* and **PBR146*****

Tracers **PBR099*** and **PBR146*** were prepared by nucleophilic substitution of the corresponding ethyl and propyl tosylate precursors with ^{18}F -fluoride in the presence of K_{222} , K_2CO_3 in ACN at 100°C for 5 mins.

The biodistribution of **PBR099*** and **PBR146*** were performed in SD rats and brain and peripheral tissues analysed at 15, 30 min, 1, and 4 h p.i. The specificity and selectivity of the tracers were assessed by pre-treatment with the PBR ligands PK11195 and Ro 5-4864 and with Flumazenil for CBR at 1 mg/kg 5 min prior to injection of the corresponding tracers.




PBR099* R = $\text{CH}_2\text{CH}_2^{18}\text{F}$

PBR146* R = $\text{CH}_2\text{CH}_2\text{CH}_2^{18}\text{F}$

The IC_{50} values of **PBR099** and **PBR146** are 14.9 and 10.5 for the PBR and >5000 nM for the CBR. **PBR099*** and **PBR146*** were synthesised in 40-50% radiochemical yield non-decay corrected and with > 95 % radiochemical purity. The specific activity ranged from 40-80 GBq/ μmol (un-optimised). The in vivo biodistribution of both tracers showed radioactivity uptake in tissue rich in PBR. The uptake of **PBR099*** in the olfactory bulbs ranged from 0.35 % at 15 min to 0.31 % ID/g at 4 h p.i whilst for **PBR146*** it ranged from 0.41 % to 0.31 % at 15 min and 4 h respectively. Whilst blood activity levels for **PBR099*** remained constant (0.23%) over the 4 h study period **PBR146*** showed a decrease from 0.4 at 15 min to 0.1 at 4 h. The adrenals showed high uptake of **PBR099*** increasing from 6.6 % ID/g at 30 min p.i to 9.7 % at 4 h whilst **PBR146*** increased from 7.9 at 30 min to 12.2 % at 4 h. In the kidneys the activity peaked at 15 min p.i. for both tracers (3.9% ID/g) for **PBR099*** and (5.5 % ID/g) for **PBR146*** decreasing to 2.4 and 3.2% at 4 h. Bone uptake for **PBR099*** ranged from 0.4 % at 15 min to 1.0 % at 4 h whereas for **PBR146*** was 1.0 to 5.0 % for the same period. Pre-treatment with PK 11195 and Ro 5-4864 decreased the uptake in peripheral organs except in the adrenals which showed an increase of **PBR099*** and **PBR146*** uptake. In the olfactory regions a small decrease was observed with **PBR146*** Flumazenil had no effect in the uptake of **PBR099*** or **PBR146*** in the brain or peripheral organs.

Claims:

1. A fluorinated compound of formula (I):



wherein,

D, G, and L are independently selected from the group consisting of: CH, C and N, and J and M are independently selected from the group consisting of C and N provided that one of J and M is C and the other is N, wherein at least two of D, G, M, J and L are N;

X is CH₂;

Y is absent;

Z is NR₁R₂;

R₁ and R₂ are independently selected from the group consisting of: hydrogen, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, aryl and heteroaryl, each being optionally substituted with one or more of the following substituents: halogen, an C₁-C₆ alkyl;

or R₁ and R₂, together with the nitrogen to which they are attached, form a heterocyclic ring having between 3 and 7 ring members, optionally substituted with one or more of the following substituents: halogen and C₁-C₆ alkyl;

R₃ is selected from the group consisting of: halogen, C₁-C₁₀ alkyl and O-(C₁-C₁₀ alkyl), wherein the C₁-C₁₀ alkyl group is optionally substituted;

E is an aryl group or a heteroaryl group, wherein each is optionally substituted with one or more fluoro substituents, or with one or more of the following substituents: C₁-C₆ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, QC₁-C₁₀ alkyl, QC₂-C₁₀ alkenyl, QC₂-C₁₀ alkynyl, Q(CH₂)_p-Q-(CH₂)_qCH₃ or Q(CH₂)_p-Q-(CH₂)_q-Q-(CH₂)_rCH₃, each of which is optionally substituted with one or more fluoro substituents, and wherein p, q and r are, independently, integers between 1 and 3, and wherein Q is selected from the group consisting of: NH, O and S,

m is a number between 0 and 3;

with the proviso that:


R₃ is a fluoro substituent, or

the group E comprises a fluoro substituent, or

the group Z comprises a fluoro substituent,

with the further proviso that E is not 4-fluorophenyl,

or a compound of formula (1a):



wherein,

D, G and L are independently selected from the group consisting of: CH, C and N, and J and M are independently selected from the group consisting of C and N provided that one of J and M is C and the other is N, wherein at least two of D, G, M, J and L are N;

X is CH₂;

Y is absent;

Z is NR₁R₂;

R₁ and R₂ are independently selected from the group consisting of: hydrogen, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, (CH₂)_naryl, aryl and heteroaryl, each of which is optionally substituted with one or more of the following substituents: chloro, bromo, iodo, C₁-C₆ alkyl and hydroxy;

or R₁ and R₂, together with the nitrogen to which they are attached, form a heterocyclic ring having between 3 and 7 ring members, which is optionally substituted with one or more of the following substituents: chloro, bromo, iodo and C₁-C₆ alkyl;

R₃ is selected from the group consisting of: chloro, bromo, iodo, C₁-C₁₀ alkyl and O-(C₁-C₁₀ alkyl), wherein the C₁-C₁₀ alkyl group is optionally substituted;

E is an aryl group or a heteroaryl group, wherein each is optionally substituted with a chloro, bromo or iodo substituent, and/or with one or more of the following substituents: C₁-C₆ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, QC₁-C₁₀ alkyl, QC₁-C₁₀ alkenyl, QC₂-C₁₀ alkynyl, Q(CH₂)_p-Q-(CH₂)_qCH₃ or Q(CH₂)_p-Q-(CH₂)_q-Q-(CH₂)_rCH₃, each of which is optionally substituted with one or more chloro, bromo, iodo or hydroxy substituents, and wherein p, q and r are, independently, integers between 1 and 3, and wherein Q is selected from the group consisting of: NH, O and S, wherein when E is phenyl, E does not have an iodo substituent attached directly to it in the 4 position;

m is a number between 0 and 3;

wherein at least one of Z, E or R₃ comprises iodine,

with the proviso that E is not 4-iodophenyl,

and with the proviso that said compound is not a compound of formula (1a) as defined in the following table:

D	G	J	L	M	R ₃	E	Z
CH	CH	C	N	N	6-Cl	4-chlorophenyl	NHCH ₂ CF ₃
CH	CH	C	N	N	6-CH ₃	4-chlorophenyl	NHCH ₂ CF ₃

2. The compound of formula (1) of claim 1 wherein E is an aryl group or a heteroaryl group, which is substituted with ¹⁸F or with ¹⁹F.

3. The compound of formula (1) of claim 1 or claim 2 wherein:
R₃ is a fluoro substituent which is ¹⁹F or ¹⁸F, or
the group E comprises a fluoro substituent which is ¹⁹F or ¹⁸F or a combination thereof, or
the group Z comprises a fluoro substituent which is ¹⁹F or ¹⁸F or a combination thereof, with the proviso that E is not-PhF.

4. The compound of formula (1) of any one of claims 1 to 3 wherein Q is oxygen.

5. The compound of formula (1) of any one of claims 1 to 4 wherein m is 1 or 2.

6. The compound of formula (1) of any one of claims 1 to 5 wherein p is 1 or 2, q is 1 or 2, r is 1, and Q is oxygen.

7. The compound of formula (1) of any one of claims 1 to 6 wherein:
M and L are N, D and G are CH and J is C, or
D, M and L are N, J is C and G is CH; or
D, J and L are N, M is C and G is CH.

8. The compound of formula (1) of any one of claims 1 to 7 wherein R₃ is attached to a carbon atom.

9. The compound of formula (1) of any one of claims 1 to 8 wherein R₃ is chloro or methyl or fluoro and m is 1 or 2.

10. The compound of formula (1) of any one of claims 1 to 9 wherein R₁ and R₂ are independently selected from the group consisting of: hydrogen, C₁-C₆ alkyl, phenyl, pyridyl, pyrimidinyl, pyridazinyl and where R₁ and R₂, together with the nitrogen to which they are attached, form a heterocyclic ring having between 4 and 6 ring members, wherein the C₁-C₆ alkyl group, the phenyl group, the pyridyl group, the pyrimidinyl group, the pyridazinyl group or the heterocyclic ring are optionally substituted with between one and three fluoro substituents.

11. The compound of formula (1) of claim 1 wherein:
R₃ is chloro, m is 1 or 2, E is fluoropyridyl, fluoropyrimidinyl or fluoropyridazinyl, Z is NR₁R₂, R₁ and R₂ are selected from the group consisting of: C₁-C₆ alkyl and phenyl and Y is absent; or
R₃ is chloro, m is 1 or 2, E is ¹⁸F fluoropyridyl, ¹⁸F fluoropyrimidinyl or ¹⁸F fluoropyridazinyl, Z is NR₁R₂, wherein R₁ and R₂ are selected from the group consisting of: C₁-C₆ alkyl and phenyl and Y is absent, or

R_3 is chloro, m is 1 or 2, E is ^{19}F fluoropyridyl, ^{19}F fluoropyrimidinyl or ^{19}F fluoropyridazinyl, Z is NR_1R_2 , wherein R_1 and R_2 are selected from the group consisting of: $\text{C}_1\text{-C}_6$ alkyl and phenyl, Y is absent, or

R_3 is chloro, m is 1 or 2, E is phenyl substituted with a $\text{C}_1\text{-C}_6$ alkoxy group, Z is NR_1R_2 , wherein R_1 is $\text{C}_1\text{-C}_6$ alkyl and R_2 is fluoropyridyl or fluorophenyl or fluoropyridazinyl, or

R_3 is chloro, m is 1 or 2, E is phenyl substituted with a $\text{C}_1\text{-C}_6$ alkoxy group, Z is NR_1R_2 , wherein R_1 is $\text{C}_1\text{-C}_6$ alkyl and R_2 is ^{18}F fluoropyridyl or ^{18}F fluorophenyl or ^{18}F fluoropyridazinyl, or

R_3 is chloro, m is 1 or 2, E is phenyl substituted with a $\text{C}_1\text{-C}_6$ alkoxy group, Z is NR_1R_2 , wherein R_1 is $\text{C}_1\text{-C}_6$ alkyl and R_2 is ^{19}F fluoropyridyl or ^{19}F fluorophenyl or ^{19}F fluoropyridazinyl, or

R_3 is F , m is 1 and E is phenyl substituted with a $\text{C}_1\text{-C}_6$ alkyl group or a $\text{C}_1\text{-C}_6$ alkoxy group, or R_3 is ^{18}F , m is 1 and E is phenyl substituted with a $\text{C}_1\text{-C}_6$ alkyl group or a $\text{C}_1\text{-C}_6$ alkoxy group, or R_3 is F , m is 1 and E is phenyl substituted with a $\text{C}_1\text{-C}_3$ alkoxy group or a $\text{C}_1\text{-C}_6$ alkyl group, or alkoxy group or a $\text{C}_1\text{-C}_6$ alkyl group; or

R_3 is $\text{C}_1\text{-C}_6$ alkyl or chloro, m is 1 or 2, Y is absent, and E is a phenyl group to which is attached one or more $\text{C}_1\text{-C}_6$ alkoxy groups, said $\text{C}_1\text{-C}_6$ alkoxy groups comprising one or more fluoro substituents, or

R_3 is $\text{C}_1\text{-C}_6$ alkyl or chloro, m is 1 or 2 and E is a phenyl group to which is attached one or more $\text{C}_1\text{-C}_6$ alkoxy groups, said $\text{C}_1\text{-C}_6$ alkoxy groups comprising between one and three fluoro substituents, or

R_3 is methyl or chloro, m is 1 or 2 and E is a phenyl group to which is attached one or more $\text{C}_1\text{-C}_6$ alkoxy groups, said $\text{C}_1\text{-C}_6$ alkoxy groups comprising one or more fluoro substituents, or

R_3 is methyl or chloro, m is 1 or 2 and E is a phenyl group to which is attached one or more $\text{C}_1\text{-C}_6$ alkoxy groups, said $\text{C}_1\text{-C}_6$ alkoxy groups comprising between one and three fluoro substituents, or

R_3 is methyl or chloro, m is 1 or 2 and E is a phenyl group to which is attached a $\text{C}_1\text{-C}_6$ alkoxy group, said $\text{C}_1\text{-C}_6$ alkoxy group comprising a single fluoro substituent (^{19}F or ^{18}F), or

R_3 is methyl or chloro, m is 1 or 2 and E is a phenyl group to which is attached an alkoxy group of the following formula: $-\text{O}(\text{CH}_2)_n\text{F}$, wherein n is a number between 1 and 4.

or R_3 is methyl or chloro, m is 1 or 2 and E is a phenyl group to which is attached an alkoxy group of the following formula: $-\text{O}(\text{CH}_2)_n^{18}\text{F}$, wherein n is a number between 1 and 4, or

R_3 is chloro or $\text{C}_1\text{-C}_6$ alkyl, m is 1 or 2, G is CH , D is CH , M is N , J is C , L is N , Y is absent, Z is NR_1R_2 , wherein R_1 and R_2 are independently selected from hydrogen, pyridyl, phenyl, $\text{C}_1\text{-C}_4$ alkyl, or R_1 and R_2 together with the nitrogen to which they are attached form a pyrrolidine ring,

and E is phenyl group substituted with a single substituent of the formula $-O(CH_2)_nF$, wherein n is a number between 1 and 4, or

R_3 is chloro or C_1-C_6 alkyl, m is 1 or 2, G is CH, D is CH, M is N, J is C, L is N, Y is absent, Z is NR_1R_2 , wherein R_1 and R_2 are independently selected from hydrogen, pyridyl, phenyl, C_1-C_4 alkyl, or R_1 and R_2 together with the nitrogen to which they are attached form a pyrrolidine ring, and E is phenyl group substituted with a single substituent of the formula $-O(CH_2)_n^{18}F$, wherein n is a number between 1 and 4, or

R_3 is methyl, m is 1 or 2, G is CH, D is N, M is C, J is N, L is N, Y is absent, Z is NR_1R_2 , wherein R_1 and R_2 are independently C_1-C_6 alkyl, and E is a phenyl group substituted with a single substituent of the formula $-O(CH_2)_nF$, wherein n is a number between 1 and 4, or

R_3 is fluoro, m is 1, G is CH, D is N, M is N, J is C, L is N, Y is absent, Z is NR_1R_2 , wherein R_1 and R_2 are independently C_1-C_4 alkyl, and E is a phenyl group substituted with a C_1-C_3 alkoxy group or a C_1-C_6 alkyl group.

R_3 is fluoro, m is 1, G is CH, D is N, M is N, J is C, L is N, Y is NH, Z is phenyl, and E is a phenyl group substituted with one or two C_1-C_6 alkyl groups, or

R_3 is chloro, m is 1 or 2, G is CH, D is CH, M is N, J is C, L is N, Y is absent, Z is NR_1R_2 , wherein R_1 and R_2 are independently selected from hydrogen, phenyl and C_1-C_4 alkyl, and E is fluoropyridine, fluoropyridazine or fluoropyrimidine, or

R_3 is chloro, m is 1 or 2, G is CH, D is CH, M is N, J is C, L is N, Y is absent, Z is NR_1R_2 , wherein R_1 and R_2 are independently selected from hydrogen, phenyl and C_1-C_4 alkyl, and E is ^{18}F fluoropyridine, ^{18}F fluoropyridazine or ^{18}F fluoropyrimidine, or

R_3 is chloro, m is 1 or 2, G is CH, D is CH, M is N, J is C, L is N, Y is absent, Z is NR_1R_2 , wherein R_1 and R_2 are independently selected from hydrogen, phenyl and C_1-C_4 alkyl, and E is ^{18}F fluoropyridine, ^{18}F fluoropyridazine or ^{18}F fluoropyrimidine, or

R_3 is chloro, m is 1 or 2, G is CH, D is CH, M is N, J is C, L is N, Y is absent, Z is NR_1R_2 , wherein R_1 and R_2 are independently selected from hydrogen, C_1-C_4 alkyl, fluoropyridyl and fluoropyridazinyl, and E is a phenyl group substituted with a C_1-C_6 alkoxy group, or

R_3 is chloro, m is 1 or 2, G is CH, D is CH, M is N, J is C, L is N, Y is absent, Z is NR_1R_2 , wherein R_1 and R_2 are independently selected from hydrogen, C_1-C_4 alkyl, ^{18}F fluoropyridyl and ^{18}F fluoropyridazinyl, and E is a phenyl group substituted with a C_1-C_6 alkoxy group, or

R_3 is chloro, m is 1 or 2, G is CH, D is CH, M is N, J is C, L is N, Y is absent, Z is NR_1R_2 , wherein R_1 and R_2 are independently selected from hydrogen, C_1-C_4 alkyl, ^{18}F fluoropyridyl and ^{18}F fluoropyridazinyl, and E is a phenyl group substituted with a C_1-C_6 alkoxy group.

12. The compound of formula (1a) of claim 1 wherein Q is oxygen.

13. The compound of formula (1a) of claim 1 or claim 12 wherein m is 1 or 2.
14. The compound of formula (1a) of any one of claims 1, 12 or 13 wherein p is 1 or 2, q is 1 or 2, r is 1, and Q is oxygen.
15. The compound of formula (1a) of any one of claims 1 or 12 to 14 wherein:
M and L are N, D and G are CH and J is C, or
D, M and L are N, J is C and G is CH, or
D, J and L are N, M is C and G is CH.
16. The compound of formula (1a) of any one of claims 1 or 12 to 15 wherein R₃ is attached to a carbon atom.
17. The compound of any one of formula (1a) of claims 1 or 12 to 16 wherein:
R₃ is chloro or methyl and m is 1 or 2, or
R₃ is chloro and m is 1 or 2.
18. The compound of any one of formula (1a) of claims 1 or 12 to 17 wherein E is a phenyl group, a pyridyl group, a pyrimidinyl group or a pyridazinyl group, each of which is optionally substituted with one or more chloro, bromo, iodo or hydroxy substituents, and/or substituted with one or more of the following substituents: C₁-C₆ alkyl, C₂-C₆ alkenyl or C₁-C₁₀ alkoxy, each of which is optionally substituted with one or more of the following substituents: chloro, bromo, iodo and hydroxy.
19. A compound according to any one of claims 1 to 18 that is radiolabelled with ¹⁸F.
20. A pharmaceutical composition comprising a compound according to any one of claims 1 to 18, together with at least one pharmaceutically acceptable carrier, diluent, excipient or adjuvant, said compound being not radiolabelled with ¹⁸F.
21. A pharmaceutical composition comprising a compound according to claim 19, together with at least one pharmaceutically acceptable carrier, diluent, excipient or adjuvant.

22. Use of a compound according to claim 19 in the manufacture of a diagnostic composition in the diagnosis of a disorder in a mammal characterised by an abnormal density of peripheral benzodiazepine receptors.

23. Use of a compound according to any one of claims 1 to 18 in the manufacture of a medicament for the treatment of a disorder characterised by an abnormal density of peripheral benzodiazepine receptors in a mammal in need of said treatment, said compound being not radiolabelled with ^{18}F .

24. A compound according to claim 19, said compound being selected from the group consisting of: PBRs 099*, 102*, 103*, 104*, 105*, 106*, 107*, 110*, 111*, 130*, 120*, 132*, 133*, 136*, 139*, 143*, 146*, 147*, 149*, 080*, 081*, 082* and 083* (as defined herein), or any combination thereof.

25. A compound according to claim 1, said compound being selected from the group consisting of: PBRs 099, 102, 103, 104, 105, 106, 107, 110, 111, 130, 120, 132, 133, 138, 139, 143, 146, 147, 149, 080, 081, 082, 083, 115, 138, 145, 155, 142, 153, 154, 156, 131, 134, 135, 141, 125, 126, 137, 98, 139, 132, 133, 117, 118, 121, 123, 124, 128, 129, 116, 119, 140, 144, 101, 113, and 114 (as defined herein), or any combination thereof.