

DESCRIPTION

ANTIOXIDANT INFLAMMATION MODULATORS: OLEANOLIC ACID

DERIVATIVES WITH SATURATION IN THE C-RING

5

BACKGROUND OF THE INVENTION

The present application claims the benefit of priority to U.S. Provisional Application Nos. 61/046,332, filed April 18, 2008, and 61/111,333, filed November 4, 2008, the entire contents of both applications are incorporated herein by reference in their entireties.

I. Field of the Invention

The present invention relates generally to the fields of biology and medicine. More particularly, it concerns compounds and methods for the treatment and prevention of diseases such as those associated with oxidative stress and inflammation.

II. Description of Related Art

15 Many serious and intractable human diseases are associated with dysregulation of inflammatory processes, including diseases such as cancer, atherosclerosis, and diabetes, which were not traditionally viewed as inflammatory conditions. Similarly, autoimmune diseases such as rheumatoid arthritis, lupus, psoriasis, and multiple sclerosis involve inappropriate and chronic activation of inflammatory processes in affected tissues, arising
20 from dysfunction of self vs. non-self recognition and response mechanisms in the immune system. In neurodegenerative diseases such as Alzheimer's and Parkinson's diseases, neural damage is correlated with activation of microglia and elevated levels of pro-inflammatory proteins such as inducible nitric oxide synthase (iNOS).

One aspect of inflammation is the production of inflammatory prostaglandins such as
25 prostaglandin E, whose precursors are produced by the enzyme cyclo-oxygenase (COX-2). High levels of COX-2 are found in inflamed tissues. Consequently, inhibition of COX-2 is known to reduce many symptoms of inflammation and a number of important anti-inflammatory drugs (*e.g.*, ibuprofen and celecoxib) act by inhibiting COX-2 activity. Recent research, however, has demonstrated that a class of cyclopentenone prostaglandins (*e.g.*, 15-deoxy prostaglandin J₂, a.k.a. PGJ₂) plays a role in stimulating the orchestrated resolution of
30 inflammation. COX-2 is also associated with the production of cyclopentenone


prostaglandins. Consequently, inhibition of COX-2 may interfere with the full resolution of inflammation, potentially promoting the persistence of activated immune cells in tissues and leading to chronic, “smoldering” inflammation. This effect may be responsible for the increased incidence of cardiovascular disease in patients using selective COX-2 inhibitors for long periods of time. Corticosteroids, another important class of anti-inflammatory drugs, have many undesirable side effects and frequently are not suitable for chronic use. Newer protein-based drugs, such as anti-TNF monoclonal antibodies, have proven to be effective for the treatment of certain autoimmune diseases such as rheumatoid arthritis. However, these compounds must be administered by injection, are not effective in all patients, and may have severe side effects. In many severe forms of inflammation (*e.g.*, sepsis, acute pancreatitis), existing drugs are ineffective. In addition, currently available drugs typically do not have significant antioxidant properties, and are not effective in reducing oxidative stress associated with excessive production of reactive oxygen species and related molecules such as peroxynitrite. Accordingly, there is a pressing need for improved therapeutics with antioxidant and anti-inflammatory properties.

A series of synthetic triterpenoid analogs of oleanolic acid have been shown to be inhibitors of cellular inflammatory processes, such as the induction by IFN- γ of inducible nitric oxide synthase (iNOS) and of COX-2 in mouse macrophages. See Honda *et al.* (2000a); Honda *et al.* (2000b), and Honda *et al.* (2002), which are all incorporated herein by reference. For example, one of these, 2-cyano-3,12-dioxooleane-1,9(11)-dien-28-oic acid methyl ester (CDDO-Me), is currently in clinical trials for a variety of disorders related to inflammation, including cancer and diabetic nephropathy. The pharmacology of these molecules is complex, as they have been shown to affect the function of multiple protein targets and thereby modulate the function of several important cellular signaling pathways related to oxidative stress, cell cycle control, and inflammation (*e.g.*, Dinkova-Kostova *et al.*, Ahmad *et al.*, 2006; Ahmad *et al.*, 2008; Liby *et al.*). Given that the biological activity profiles of the known oleanolic acid derivatives vary, and in view of the wide variety of diseases that may be treated with compounds having potent antioxidant and anti-inflammatory effects, it is desirable to synthesize new candidates for the treatment or prevention of disease.

SUMMARY OF THE INVENTION

The present disclosure provides new compounds with antioxidant and anti-inflammatory properties, methods for their manufacture, and methods for their use. Compounds covered by the generic or specific formulas below or specifically named are referred to as “compounds of the invention,” “compounds of the present disclosure,” “the present oleanolic acid derivatives” in the present application.

In some aspects, the disclosure provides compounds of the formula:



(I),

wherein:

Y is cyano, heteroaryl_(C≤12), substituted heteroaryl_(C≤12), or -C(O)R_a, further wherein R_a is:


hydrogen, hydroxy, halo, amino, hydroxyamino, azido, silyl or mercapto;

alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12), heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyloxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12), heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12), alkylamino_(C≤12), dialkylamino_(C≤12), alkoxyamino_(C≤12), alkenylamino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12), aralkylamino_(C≤12), heteroarylamino_(C≤12), heteroaralkylamino_(C≤12), alkylsulfonylamino_(C≤12), amido_(C≤12), alkyl-

thio_(C≤12), alkenylthio_(C≤12), alkynylthio_(C≤12), arylthio_(C≤12),
 aralkylthio_(C≤12), heteroarylthio_(C≤12), heteroaralkylthio_(C≤12),
 acylthio_(C≤12), alkylammonium_(C≤12), alkylsulfonium_(C≤12),
 alkylsilyl_(C≤12), or a substituted version of any of these groups;

or

R_a comprises a nitrogen atom that is also attached to carbon atom 13
 and R_d to form:



wherein R_d is alkyl_(C≤8), alkenyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8),
 heteroaryl_(C≤8), heteroaralkyl_(C≤8), or a substituted
 version of any of these groups;

Z is a single or double bond, -O- or -NR_e-, wherein R_e is hydrogen, hydroxy,
 alkyl_(C≤8) or alkoxy_(C≤8);

X is OR_b, NR_bR_c, or SR_b, wherein R_b and R_c are each independently:

hydrogen or hydroxy;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8),
 alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of
 any of these groups; or

a substituent convertible *in vivo* to hydrogen;

provided that R_b is absent when the atom to which it is bound is part of a
 double bond, further provided that when R_b is absent the atom to which
 it is bound is part of a double bond;

R₁ is:

hydrogen, cyano, hydroxy, halo or amino; or

alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8),
 heteroaralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8),

alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups;

R₂ is:

cyano, hydroxy, halo or amino; or

5 fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8), acyl_(C≤8),
alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8),
amido_(C≤8), or a substituted version of any of these groups;

R₃ is:

absent or hydrogen;

10 alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any of
these groups; or

a substituent convertible *in vivo* to hydrogen;

provided that R₃ is absent when the oxygen atom to which it is bound is part of
a double bond, further provided that when R₃ is absent the oxygen
15 atom to which it is bound is part of a double bond;

R₄ and R₅ are each independently alkyl_(C≤8) or substituted alkyl_(C≤8);

R₆ is hydrogen, hydroxy or oxo;


R₇ is hydrogen or hydroxy; and

R₈, R₉, R₁₀ and R₁₁ are each independently hydrogen, hydroxy, alkyl_(C≤8), substituted

20 alkyl_(C≤8), alkoxy_(C≤8) or substituted alkoxy_(C≤8);

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

In some embodiments, the compound is further defined as:



wherein:

Y is cyano, or $-C(O)R_a$, further wherein:

5 R_a is:

hydrogen, hydroxy, halo, amino, hydroxyamino, azido or mercapto; or
 alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12),
 heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyl-
 oxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12),
 10 heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12),
 alkylamino_(C≤12), dialkylamino_(C≤12), alkoxyamino_(C≤12),
 alkenylamino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12),
 aralkylamino_(C≤12), heteroarylamino_(C≤12), heteroaralkyl-
 amino_(C≤12), alkylsulfonylamino_(C≤12), amido_(C≤12), alkyl-
 15 thio_(C≤12), alkenylthio_(C≤12), alkynylthio_(C≤12), arylthio_(C≤12),
 aralkylthio_(C≤12), heteroarylthio_(C≤12), heteroaralkylthio_(C≤12),
 acylthio_(C≤12), alkylammonium_(C≤12), alkylsulfonium_(C≤12),
 alkylsilyl_(C≤12), or a substituted version of any of these groups;

X is OR_b , NR_bR_c , or SR_b , wherein R_b and R_c are each independently:

20 hydrogen or hydroxy;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8),
acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a
substituted version of any of these groups; or
a substituent convertible *in vivo* to hydrogen;
5 provided that R_b is absent when the atom to which it is bound is part of
a double bond, further provided that when R_b is absent the atom
to which it is bound is part of a double bond;

R₁ is:

hydrogen, cyano, hydroxy, halo or amino; or
10 alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8),
heteroaralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8),
alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of
any of these groups;

R₂ is:

15 cyano, hydroxy, halo or amino; or
fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8), acyl_(C≤8),
alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8),
amido_(C≤8), or a substituted version of any of these groups;

R₃ is:


20 absent or hydrogen;
alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any of
these groups; or
a substituent convertible *in vivo* to hydrogen;
provided that R₃ is absent when the oxygen atom to which it is bound is part of
25 a double bond, further provided that when R₃ is absent the oxygen
atom to which it is bound is part of a double bond;

R₄ and R₅ are each independently alkyl_(C≤8) or substituted alkyl_(C≤8); and

R₆ and R₇ are each independently hydrogen or hydroxy;

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical
30 isomers thereof.

In some embodiments, the compound is further defined as:



wherein:

Y is cyano or $-\text{C}(\text{O})\text{R}_a$, further wherein:

5

R_a is:

hydrogen, hydroxy, halo, amino, azido, mercapto or silyl; or

alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12),

heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyl-

oxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12),

10

heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12),

alkylamino_(C≤12), alkoxyamino_(C≤12), dialkylamino_(C≤12),

alkenylamino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12),

aralkylamino_(C≤12), heteroarylamino_(C≤12), heteroaralkyl-

amino_(C≤12), amido_(C≤12), or a substituted version of any of these

15

groups;

X is OR_b , NR_bR_c , or SR_b , wherein R_b and R_c are each independently:

hydrogen or hydroxy;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8),

acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a

20

substituted version of any of these groups; or

a substituent convertible *in vivo* to hydrogen;

provided that R_b is absent when the atom to which it is bound is part of a double bond, further provided that when R_b is absent the atom to which it is bound is part of a double bond;

R_1 is:

5 hydrogen, cyano, hydroxy, halo or amino; or
alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8), heteroaralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups;


10 R_2 is:

cyano, hydroxy, halo or amino; or
fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups; and

15 R_4 is alkyl_(C≤8) or substituted alkyl_(C≤8);

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

In some embodiments, the compound is further defined as:



20 wherein:

R_a is:

hydrogen, hydroxy, halo, or amino; or

alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8), heteroaralkyl_(C≤8), alkoxy_(C≤8), alkenyloxy_(C≤8), alkynyloxy_(C≤8), aryloxy_(C≤8), aralkoxy_(C≤8), heteroaryloxy_(C≤8), heteroaralkoxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), alkoxyamino_(C≤8), dialkylamino_(C≤8), alkenylamino_(C≤8), alkynylamino_(C≤8), arylamino_(C≤8), aralkylamino_(C≤8), heteroarylamino_(C≤8), heteroaralkylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups;

X is OR_b or NR_bR_c, wherein R_b and R_c are each independently:

hydrogen or hydroxy;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups; or

a substituent convertible *in vivo* to hydrogen;

provided that R_b is absent when the atom to which it is bound is part of a double bond, further provided that when R_b is absent the atom to which it is bound is part of a double bond; and


R₂ is:

cyano, hydroxy, halo or amino; or

fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups;

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

In some embodiments, the compound is further defined as:



wherein:

R_a is:


- 5 hydrogen, hydroxy, halo, or amino; or
- alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8),
 heteroaralkyl_(C≤8), alkoxy_(C≤8), alkenyloxy_(C≤8), alkynyloxy_(C≤8),
 aryloxy_(C≤8), aralkoxy_(C≤8), heteroaryloxy_(C≤8), heteroaralkoxy_(C≤8),
 acyloxy_(C≤8), alkylamino_(C≤8), alkoxyamino_(C≤8), dialkylamino_(C≤8),
 10 alkenylamino_(C≤8), alkynylamino_(C≤8), arylamino_(C≤8), aralkylamino_(C≤8),
 heteroarylamino_(C≤8), heteroaralkylamino_(C≤8), amido_(C≤8), or a
 substituted version of any of these groups; and

R_2 is:

- cyano or fluoro; or
- 15 fluoroalkyl_(C≤5), alkenyl_(C≤5), alkynyl_(C≤5), heteroaryl_(C≤5), acyl_(C≤5), acyloxy_(C≤5),
 amido_(C≤5), or a substituted version of any of these groups;

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

In some embodiments, the compound is further defined as:




wherein R_a is:

hydrogen, hydroxy, halo or amino; or

- 5 alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8),
 heteroaralkyl_(C≤8), alkoxy_(C≤8), alkenyloxy_(C≤8), alkynyloxy_(C≤8), aryloxy_(C≤8),
 aralkoxy_(C≤8), heteroaryloxy_(C≤8), heteroaralkoxy_(C≤8), acyloxy_(C≤8),
 alkylamino_(C≤8), alkoxyamino_(C≤8), dialkylamino_(C≤8), alkenylamino_(C≤8),
 alkynylamino_(C≤8), arylamino_(C≤8), aralkylamino_(C≤8), heteroarylamino_(C≤8),
 10 heteroaralkylamino_(C≤8), amido_(C≤8), or a substituted version of any of these
 groups;


or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

In some embodiments, the compound is further defined as:




- 15 wherein R_a is alkoxy_(C1-4), alkylamino_(C1-4), alkoxyamino_(C1-4), dialkylamino_(C2-4), or a
 substituted version of any of these groups; or pharmaceutically acceptable salts, esters,
 hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

In some embodiments, the compound is further defined as:




wherein R_a is alkyl_(C1-4) or aralkoxy_(C7-8) or a substituted version of either of these groups; or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

In some embodiments, the compound is further defined as:




wherein R_a is hydrogen, hydroxy, amino, dimethylamino, methyl, methoxy, methoxyamino, benzyloxy, or 2,2,2-trifluoroethylamino; or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

In some embodiments, the compound is further defined as:



wherein R_a is hydrogen, hydroxy, amino, methoxy, or 2,2,2-trifluoroethylamino; or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.


In some embodiments, the compound is further defined as:



(XI),

wherein R_d is $\text{alkyl}_{(C \leq 8)}$, $\text{alkenyl}_{(C \leq 8)}$, $\text{aryl}_{(C \leq 8)}$, $\text{aralkyl}_{(C \leq 8)}$, $\text{heteroaryl}_{(C \leq 8)}$, $\text{heteroaralkyl}_{(C \leq 8)}$, or a substituted version of any of these groups; or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.


In some embodiments, the compound is further defined as:



(XII),

wherein Y is $\text{heteroaryl}_{(C \leq 8)}$ or a substituted $\text{heteroaryl}_{(C \leq 8)}$; or a pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

In some embodiments, the compound is further defined as:



(XIII),

wherein:

wherein Y is cyano or $-\text{C}(\text{O})\text{R}_a$, further wherein:

R_a is:

hydrogen, hydroxy, halo, amino, azido, mercapto or silyl; or

alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12),
 heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyl-
 oxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12),
 heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12),
 alkylamino_(C≤12), alkoxyamino_(C≤12), dialkylamino_(C≤12),
 alkenylamino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12),
 aralkylamino_(C≤12), heteroarylamino_(C≤12), heteroaralkyl-
 amino_(C≤12), amido_(C≤12), or a substituted version of any of these
 groups;

R₂ is:

cyano, hydroxy, halo or amino; or

fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8), acyl_(C≤8),
 alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8),
 amido_(C≤8), or a substituted version of any of these groups;

R₃ is:

absent or hydrogen;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any of
 these groups; or


a substituent convertible *in vivo* to hydrogen;

provided that R₃ is absent when the oxygen atom to which it is bound is part of
 a double bond, further provided that when R₃ is absent the oxygen
 atom to which it is bound is part of a double bond; and

R₄ is alkyl_(C≤8) or substituted alkyl_(C≤8);

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical
 isomers thereof.

In some embodiments, the compound is further defined as:



wherein:

Y is cyano or $-\text{C}(\text{O})\text{R}_a$, wherein

5 R_a is:
hydrogen, hydroxy, halo, amino, hydroxyamino, azido or mercapto; or
 $\text{alkyl}_{(\text{C}\leq 12)}$, $\text{alkenyl}_{(\text{C}\leq 12)}$, $\text{alkynyl}_{(\text{C}\leq 12)}$, $\text{aryl}_{(\text{C}\leq 12)}$, $\text{aralkyl}_{(\text{C}\leq 12)}$,
 $\text{heteroaryl}_{(\text{C}\leq 12)}$, $\text{heteroaralkyl}_{(\text{C}\leq 12)}$, $\text{alkoxy}_{(\text{C}\leq 12)}$, alkenyl-
 $\text{oxy}_{(\text{C}\leq 12)}$, $\text{alkynyloxy}_{(\text{C}\leq 12)}$, $\text{aryloxy}_{(\text{C}\leq 12)}$, $\text{aralkoxy}_{(\text{C}\leq 12)}$,
10 $\text{heteroaryloxy}_{(\text{C}\leq 12)}$, $\text{heteroaralkoxy}_{(\text{C}\leq 12)}$, $\text{acyloxy}_{(\text{C}\leq 12)}$,
 $\text{alkylamino}_{(\text{C}\leq 12)}$, $\text{alkoxyamino}_{(\text{C}\leq 12)}$, $\text{dialkylamino}_{(\text{C}\leq 12)}$,
 $\text{alkenylamino}_{(\text{C}\leq 12)}$, $\text{alkynylamino}_{(\text{C}\leq 12)}$, $\text{arylamino}_{(\text{C}\leq 12)}$,
 $\text{aralkylamino}_{(\text{C}\leq 12)}$, $\text{heteroarylamino}_{(\text{C}\leq 12)}$, heteroaralkyl-
 $\text{amino}_{(\text{C}\leq 12)}$, $\text{alkylsulfonylamino}_{(\text{C}\leq 12)}$, $\text{amido}_{(\text{C}\leq 12)}$, or a
15 substituted version of any of these groups;

R_2 is:


cyano, hydroxy, halo or amino; or
 $\text{fluoroalkyl}_{(\text{C}\leq 8)}$, $\text{alkenyl}_{(\text{C}\leq 8)}$, $\text{alkynyl}_{(\text{C}\leq 8)}$, $\text{aryl}_{(\text{C}\leq 8)}$, $\text{heteroaryl}_{(\text{C}\leq 8)}$, $\text{acyl}_{(\text{C}\leq 8)}$,
 $\text{alkoxy}_{(\text{C}\leq 8)}$, $\text{aryloxy}_{(\text{C}\leq 8)}$, $\text{acyloxy}_{(\text{C}\leq 8)}$, $\text{alkylamino}_{(\text{C}\leq 8)}$, $\text{arylamino}_{(\text{C}\leq 8)}$,
20 $\text{amido}_{(\text{C}\leq 8)}$, or a substituted version of any of these groups; and

R_3 is:

hydrogen;
 $\text{alkyl}_{(\text{C}\leq 8)}$, $\text{aryl}_{(\text{C}\leq 8)}$, $\text{aralkyl}_{(\text{C}\leq 8)}$, $\text{acyl}_{(\text{C}\leq 8)}$, or a substituted version of any of
these groups; or
25 a substituent convertible *in vivo* to hydrogen;

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

In some embodiments, the compound is further defined as:



wherein:

X is OR_b , NR_bR_c , or SR_b , wherein R_b and R_c are each independently:

- 5 hydrogen;
 $alkyl_{(C\leq 8)}$, $aryl_{(C\leq 8)}$, $aralkyl_{(C\leq 8)}$, $acyl_{(C\leq 8)}$, or a substituted version of any
 of these groups; or
 a substituent convertible *in vivo* to hydrogen;
 provided that R_b is absent when the atom to which it is bound is part of
 10 a double bond, further provided that when R_b is absent the atom
 to which it is bound is part of a double bond;

R_1 is:


- hydrogen, cyano, hydroxy, halo or amino; or
 $alkyl_{(C\leq 8)}$, $alkenyl_{(C\leq 8)}$, $alkynyl_{(C\leq 8)}$, $aryl_{(C\leq 8)}$, $aralkyl_{(C\leq 8)}$, $heteroaryl_{(C\leq 8)}$,
 15 $heteroaralkyl_{(C\leq 8)}$, $acyl_{(C\leq 8)}$, $alkoxy_{(C\leq 8)}$, $aryloxy_{(C\leq 8)}$, $acyloxy_{(C\leq 8)}$,
 $alkylamino_{(C\leq 8)}$, $arylamino_{(C\leq 8)}$, $amido_{(C\leq 8)}$, or a substituted version of
 any of these groups;

R_2 is:

- cyano, hydroxy, halo or amino; or
 20 $fluoroalkyl_{(C\leq 8)}$, $alkenyl_{(C\leq 8)}$, $alkynyl_{(C\leq 8)}$, $aryl_{(C\leq 8)}$, $heteroaryl_{(C\leq 8)}$, $acyl_{(C\leq 8)}$,
 $alkoxy_{(C\leq 8)}$, $aryloxy_{(C\leq 8)}$, $acyloxy_{(C\leq 8)}$, $alkylamino_{(C\leq 8)}$, $arylamino_{(C\leq 8)}$,
 $amido_{(C\leq 8)}$, or a substituted version of any of these groups; and

R_4 is alkyl_(C≤8) or substituted alkyl_(C≤8);
or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

In some embodiments, the compound is further defined as:



wherein:

X is OR_b or NR_bR_c, wherein R_b and R_c are each independently:

hydrogen;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any
of these groups; or

a substituent convertible *in vivo* to hydrogen;

provided that R_b is absent when the atom to which it is bound is part of
a double bond, further provided that when R_b is absent the atom
to which it is bound is part of a double bond; and


R₂ is:

cyano, hydroxy, halo or amino; or

fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8), acyl_(C≤8),
alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8),
amido_(C≤8), or a substituted version of any of these groups;

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

In some embodiments, the compound is further defined as:



wherein:

R_a is:

- 5 hydrogen, hydroxy, halo, amino, azido, mercapto or silyl; or
 alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12), heteroaryl_(C≤12),
 heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyloxy_(C≤12), alkynyloxy_(C≤12),
 aryloxy_(C≤12), aralkoxy_(C≤12), heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12),
 acyloxy_(C≤12), alkylamino_(C≤12), alkoxyamino_(C≤12), dialkylamino_(C≤12),
 10 alkenylamino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12),
 aralkylamino_(C≤12), heteroarylamino_(C≤12), heteroaralkylamino_(C≤12),
 amido_(C≤12), or a substituted version of any of these groups;

R₁ is:


- hydrogen, cyano, hydroxy, halo or amino; or
 15 alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8),
 heteroaralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8),
 alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of
 any of these groups;

R₂ is:

- 20 cyano, hydroxy, halo or amino; or

fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8), acyl_(C≤8),
 alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8),
 amido_(C≤8), or a substituted version of any of these groups; and
 or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical
 5 isomers thereof.

In some embodiments, the compound is further defined as:




wherein R_a is:

hydrogen, hydroxy, halo or amino; or

10 alkyl_(C≤6), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8), alkoxy_(C≤6), aryloxy_(C≤8), aralkoxy_(C≤8),
 alkylamino_(C≤6), alkoxyamino_(C≤6), alkoxyamino_(C≤6), dialkylamino_(C≤6),
 arylamino_(C≤8), aralkylamino_(C≤8), heteroarylamino_(C≤6), heteroarylamino_(C≤8),
 amido_(C≤6), or a substituted version of any of these groups;
 or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical
 15 isomers thereof.

In some aspects, the disclosure provides compounds of the formula:



wherein:

Y is cyano or $-C(O)R_a$, wherein

R_a is:

hydrogen, hydroxy, halo, amino, hydroxyamino, azido or mercapto; or

5 alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12),
 heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyl-
 oxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12),
 heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12),
 alkylamino_(C≤12), dialkylamino_(C≤12), alkoxyamino_(C≤12), alkenyl-
 10 amino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12),
 aralkylamino_(C≤12), heteroarylamino_(C≤12), heteroaralkyl-
 amino_(C≤12), alkylsulfonylamino_(C≤12), amido_(C≤12), alkyl-
 thio_(C≤12), alkenylthio_(C≤12), alkynylthio_(C≤12), arylthio_(C≤12),
 aralkylthio_(C≤12), heteroarylthio_(C≤12), heteroaralkylthio_(C≤12),
 15 acylthio_(C≤12), alkylammonium_(C≤12), alkylsulfonium_(C≤12),
 alkylsilyl_(C≤12), or a substituted version of any of these groups;

X is OR_b , NR_bR_c , or SR_b , wherein R_b and R_c are each independently:

hydrogen;

20 alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any
 of these groups; or


provided that R_b is absent when the atom to which it is bound is part of
 a double bond, further provided that when R_b is absent the atom
 to which it is bound is part of a double bond; and

R_1 is:

25 hydrogen, cyano, hydroxy, halo or amino; or
 alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8),
 heteroaralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8),
 alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of
 any of these groups;

30 or salts, esters, hydrates, solvates, tautomers, or optical isomers thereof.

In some aspects, the disclosure provides compounds of the formula:



wherein:

Y is cyano or $-C(O)R_a$, wherein

5 R_a is:

hydrogen, hydroxy, halo, amino, hydroxyamino, azido or mercapto; or
 alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12),
 heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyl-
 oxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12),
 10 heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12),
 alkylamino_(C≤12), dialkylamino_(C≤12), alkoxyamino_(C≤12),
 alkenylamino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12),
 aralkylamino_(C≤12), heteroarylamino_(C≤12), heteroaralkyl-
 amino_(C≤12), alkylsulfonylamino_(C≤12), amido_(C≤12), alkyl-
 15 thio_(C≤12), alkenylthio_(C≤12), alkynylthio_(C≤12), arylthio_(C≤12),
 aralkylthio_(C≤12), heteroarylthio_(C≤12), heteroaralkylthio_(C≤12),
 acylthio_(C≤12), alkylammonium_(C≤12), alkylsulfonium_(C≤12),
 alkylsilyl_(C≤12), or a substituted version of any of these groups;

X is OR_b , NR_bR_c , or SR_b , wherein R_b and R_c are each independently:

20 hydrogen;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any
 of these groups; or

provided that R_b is absent when the atom to which it is bound is part of a double bond, further provided that when R_b is absent the atom to which it is bound is part of a double bond; and

R_1 is:

5 hydrogen, cyano, hydroxy, halo or amino; or
alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8), heteroaralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups;

10 R' is hydroxy, alkoxy_(C≤12), substituted alkoxy_(C≤12), aryloxy_(C≤12), substituted aryloxy_(C≤12), aralkoxy_(C≤12), substituted aralkoxy_(C≤12), acyloxy_(C≤12), or substituted acyloxy_(C≤12);

or salts, esters, hydrates, solvates, tautomers, or optical isomers thereof.

In a variation of each of the above embodiments containing a Z group, Z can be a
15 single bond, $-O-$, or $-NH-$. In a variation of each of the above embodiments containing an X group, X can be OR_b . In some variations, R_b is absent. In other variations, R_b is hydrogen. In other variations, X can be NR_b . In some variations, R_b can be hydroxy. In a variation of each of the above embodiments containing a Y group, Y can be cyano or $-C(O)R_a$. In some variations, R_a can be hydroxy. In some variations, R_a can be alkoxy_(C≤6), aryloxy_(C≤8),
20 aralkyloxy_(C≤8), or a substituted version of any of these groups. In some of these variations, R_a can be alkoxy_(C2-6). In some of these variations, R_a can be alkoxy_(C1-5) or substituted alkoxy_(C1-5). In some of these variations, R_a can be alkoxy_(C2-4) or substituted alkoxy_(C2-4). In some of these variations, R_a can be alkoxy_(C1-4) or substituted alkoxy_(C1-4). In some of these variations, R_a can be alkoxy_(C1-2) or substituted alkoxy_(C1-2). For example, R_a can be methoxy.
25 In some variations, R_a can be amino. In some variations, R_a can be alkylamino_(C1-6), alkoxyamino_(C1-6), arylamino_(C1-8), aralkylamino_(C1-8), dialkylamino_(C2-8), or a substituted version of any of these groups. In some of these variations, R_a can be alkylamino_(C2-6) or substituted alkylamino_(C2-6). In some of these variations, R_a can be alkylamino_(C3-6). In some variations, R_a can be alkylamino_(C1-5), dialkylamino_(C2-6), or substituted version of either of
30 these groups. In some variations, R_a can be alkylamino_(C2-4), dialkylamino_(C2-5), or substituted version of either of these groups. In some variations, R_a can be alkylamino_(C1-4) or substituted alkylamino_(C1-4). In some variations, R_a can be alkylamino_(C1-3). In some of these variations,









R_a can be methylamino or ethylamino. In some of these variations, R_a can be substituted alkylamino_(C1-3). For example, R_a can be 2,2,2-trifluoroethylamino. In some of these variations, R_a can be alkyl_(C1-5), aryl_(C≤8), aralkyl_(C≤8), heteroaralkyl_(C≤8), or a substituted version of any of these groups. In some of these variations, R_a can be heteroaryl_(C1-8) or substituted heteroaryl_(C1-8). For example, wherein R_a can be imidazolyl. In some of these variations, R_a can be -H.


In a variation of each of the above embodiments containing an R₁ group, R₁ can be -H, -OH or -F. For example, R₁ can be -H. In a variation of each of the above embodiments containing an R₂ group, R₂ can be -CN. In some variations, R₂ can be a substituted acyl_(C1-3), such as -C(=O)NHS(=O)₂CH₃. In some variations, R₂ is fluoroalkyl_(C≤8). For example, R₂ can be -CF₃. In other variations, R₂ is not fluoroalkyl_(C≤8).


In a variation of each of the above embodiments containing an R₃ group, R₃ can be hydrogen or acetyl. In another variation, R₃ can be absent. In a variation of each of the above embodiments containing an R₄ group, R₄ can be methyl or hydroxymethyl. In a variation of each of the above embodiments containing an R₆, R₇, R₈, or R₉ group, R₆, R₇, R₈, or R₉ can independently be hydrogen. In a variation of each of the above embodiments containing an R₁₀ or R₁₁ group, R₁₀ or R₁₁ can independently be methyl. In a variation of each of the above embodiments containing an R' group, R' can be acetyloxy or hydroxy.

In a variation of each of the above embodiments containing an R_d group, R_d can be alkyl_(C1-5), aryl_(C≤8), aralkyl_(C≤8), heteroaralkyl_(C≤8), or a substituted version of any of these groups. In some variations, R_d can be alkyl_(C1-4) or a substituted version thereof. In some variations, R_d can be alkyl_(C1-3) or a substituted version thereof. In some variations, R_d can be alkyl_(C1-2) or a substituted version thereof.

Non-limiting examples of compounds provided by this invention include the compounds according to the formulas shown below, as well as or pharmaceutically acceptable salts thereof. In certain embodiments, these compounds are substantially free from other optical isomers thereof.

 63101 (402-02) (dh402)	 63250 (402-46)
 63196 (402-51) (dh401)	 63252 (402-57)
 63205 (402-59)	 63206 (402-64)
 63207 (402-66)	 63219 (402-78)





Examples of specific compounds provided by the present disclosure include:

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,14a*R*,14b*S*)-methyl 11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydricene-4a-carboxylate,

5 (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,14a*R*,14b*S*)-methyl 11-cyano-10-hydroxy-2,2,6a,6b,9,9,12a-heptamethyl-14-oxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,12,12a,12b,13,14,14a,14b-icosahydricene-4a-carboxylate,

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,14a*R*,14b*S*)-11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxylic acid,

5 (4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14b*R*)-4,4,6a,6b,11,11,14b-heptamethyl-3,13-dioxo-3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydronicene-2,8a-dicarbonitrile,

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,14a*R*,14b*S*)-11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxamide,

10 (4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14b*R*)-8a-formyl-4,4,6a,6b,11,11,14b-heptamethyl-3,13-dioxo-3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydronicene-2-carbonitrile,

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,14*R*,14a*R*,14b*S*)-methyl 11-cyano-14-hydroxy-2,2,6a,6b,9,9,12a-heptamethyl-10-oxo-

15 1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxylate,

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,14a*R*,14b*S*)-11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-N-(2,2,2-trifluoroethyl)-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxamide,

20 (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14*R*,14a*R*,14b*S*)-2,2,2-trifluoroethyl 11-cyano-14-hydroxy-2,2,6a,6b,9,9,12a-heptamethyl-10-oxo-

1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxylate,

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-N-(2,2,2-trifluoroethyl)-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxamide,

25 (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-N'-acetyl-11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carbohydrazide,

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-N-(2,2,2-trifluoroethyl)docosahydronicene-4a-carboxamide,

30 (4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14a*R*,14b*R*)-4,4,6a,6b,11,11,14b-heptamethyl-3,13-dioxo-8a-(2H-tetrazol-5-yl)-3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydronicene-2-carbonitrile,

(4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14a*R*,14b*R*)-4,4,6a,6b,11,11,14b-heptamethyl-8a-(2-methyl-2H-tetrazol-5-yl)-3,13-dioxo-

3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydronicene-2-carbonitrile,

(4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14a*R*,14b*R*)-4,4,6a,6b,11,11,14b-heptamethyl-8a-(5-methyl-1,3,4-oxadiazol-2-yl)-3,13-dioxo-

3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydronicene-2-carbonitrile, and

(4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14a*R*,14b*R*)-4,4,6a,6b,11,11,14b-heptamethyl-8a-(5-methyl-1,3,4-thiadiazol-2-yl)-3,13-dioxo-

3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydronicene-2-carbonitrile,

(4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14a*R*,14b*R*)-8a-acetyl-4,4,6a,6b,11,11,14b-heptamethyl-3,13-dioxo-3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydronicene-2-carbonitrile,

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-benzyl 11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxylate,

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-11-cyano-*N,N*,2,2,6a,6b,9,9,12a-nonamethyl-10,14-dioxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxamide,

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-11-cyano-*N*-methoxy-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxamide

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-methyl 11-cyano-14-(hydroxyimino)-2,2,6a,6b,9,9,12a-heptamethyl-10-oxo-

1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxylate,

(4a*R*,6a*R*,6b*S*,8a*S*,12a*R*,15a*R*,15b*R*)-methyl 2-cyano-14-hydroxy-4,4,6a,6b,11,11,15b-heptamethyl-3-oxo-3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,14,15,15a,15b-icosahydrodinaphtho[1,2-*b*:2',1'-*d*]oxepine-8a-carboxylate, and

(4a*R*,6a*R*,6b*S*,8a*S*,12a*R*,12b*R*,15a*R*,15b*R*)-methyl 2-cyano-4,4,6a,6b,11,11,15b-heptamethyl-3,14-dioxo-4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,15,15a,15b-icosahydro-3*H*-dinaphtho[1,2-*b*:2',1'-*d*]azepine-8a-carboxylate.

In some embodiments, compounds of the present disclosure are in the form of pharmaceutically acceptable salts. In other embodiments, compounds of the present

disclosure are not in the form of a pharmaceutically acceptable salts. In some embodiments, compounds of the present disclosure are in the form of a hydrate. In other embodiments, compounds of the present disclosure are not in the form of a hydrate. In some embodiments, compounds of the present disclosure are in the form of a solvate. In other embodiments, compounds of the present disclosure are not in the form of a solvate.

In some embodiments, compounds of the present disclosure can be esters of the above formulas. The ester may, for example, result from a condensation reaction between a hydroxy group of the formula and the carboxylic acid group of biotin. In other embodiments, compounds of the present disclosure are not an ester.

In some embodiments, the compounds of the present disclosure can be present as a mixture of stereoisomers. In other embodiments, the compounds of the present disclosure are present as single stereoisomers.

In some embodiments, compounds of the present disclosure may be inhibitors of IFN- γ -induced nitrous oxide (NO) production in macrophages, for example, having an IC₅₀ value of less than 0.2 μ M.

Other general aspects of the present disclosure contemplate a pharmaceutical composition comprising as an active ingredient a compound of the present disclosure and a pharmaceutically acceptable carrier. The composition may, for example, be adapted for administration by a route selected from the group consisting of orally, intraadiposally, intraarterially, intraarticularly, intracranially, intradermally, intralesionally, intramuscularly, intranasally, intraocularly, intrapericardially, intraperitoneally, intrapleurally, intraprostatically, intrarectally, intrathecally, intratracheally, intratumorally, intraumbilically, intravaginally, intravenously, intravesicularly, intravitreally, liposomally, locally, mucosally, orally, parenterally, rectally, subconjunctival, subcutaneously, sublingually, topically, transbuccally, transdermally, vaginally, in crèmes, in lipid compositions, via a catheter, via a lavage, via continuous infusion, via infusion, via inhalation, via injection, via local delivery, via localized perfusion, bathing target cells directly, or any combination thereof. In particular embodiments, the composition may be formulated for oral delivery. In particular embodiments, the composition is formulated as a hard or soft capsule, a tablet, a syrup, a suspension, a wafer, or an elixir. In certain embodiments, the soft capsule is a gelatin capsule. Certain compositions may comprise a protective coating, such as those compositions formulated for oral delivery. Certain compositions further comprise an agent that delays absorption, such as those compositions formulated for oral delivery. Certain compositions

may further comprise an agent that enhances solubility or dispersibility, such as those compositions formulated for oral delivery. Certain compositions may comprise a compound of the present disclosure, wherein the compound is dispersed in a liposome, an oil in water emulsion or a water in oil emulsion.

5 Yet another general aspect of the present disclosure contemplates a therapeutic method comprising administering a pharmaceutically effective compound of the present disclosure to a subject. The subject may, for example, be a human. These or any other methods of the present disclosure may further comprise identifying a subject in need of treatment.

10 Another method of the present disclosure contemplates a method of treating cancer in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound of the present disclosure. The cancer may be any type of cancer, such as a carcinoma, sarcoma, lymphoma, leukemia, melanoma, mesothelioma, multiple myeloma, or seminoma. Other types of cancers include cancer of the bladder, blood, bone, brain, breast,
15 central nervous system, colon, endometrium, esophagus, genitourinary tract, head, larynx, liver, lung, neck, ovary, pancreas, prostate, spleen, small intestine, large intestine, stomach, or testicle. In these or any other methods, the subject may be a primate. In these or any other methods, the subject may be a human. This or any other method may further comprise identifying a subject in need of treatment. The subject may have a family or patient history of
20 cancer. In certain embodiments, the subject has symptoms of cancer. The compounds of the invention may be administered *via* any method described herein, such as locally. In certain embodiments, the compound is administered by direct intratumoral injection or by injection into tumor vasculature. In certain embodiments, the compounds may be administered systemically. The compounds may be administered intravenously, intra-arterially,
25 intramuscularly, intraperitoneally, subcutaneously or orally, in certain embodiments.

In certain embodiments regarding methods of treating cancer in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound of the present disclosure, the pharmaceutically effective amount is 0.1 – 1000 mg/kg. In certain
30 embodiments, the pharmaceutically effective amount is administered in a single dose per day. In certain embodiments, the pharmaceutically effective amount is administered in two or more doses per day. The compound may be administered by contacting a tumor cell during *ex vivo* purging, for example. The method of treatment may comprise any one or more of the following: a) inducing cytotoxicity in a tumor cell; b) killing a tumor cell; c) inducing

apoptosis in a tumor cell; d) inducing differentiation in a tumor cell; or e) inhibiting growth in a tumor cell. The tumor cell may be any type of tumor cell, such as a leukemia cell. Other types of cells include, for example, a bladder cancer cell, a breast cancer cell, a lung cancer cell, a colon cancer cell, a prostate cancer cell, a liver cancer cell, a pancreatic cancer cell, a stomach cancer cell, a testicular cancer cell, a brain cancer cell, an ovarian cancer cell, a lymphatic cancer cell, a skin cancer cell, a brain cancer cell, a bone cancer cell, or a soft tissue cancer cell.

Combination treatment therapy is also contemplated by the present disclosure. For example, regarding methods of treating cancer in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound of the present disclosure, the method may further comprise a treatment selected from the group consisting of administering a pharmaceutically effective amount of a second drug, radiotherapy, gene therapy, and surgery. Such methods may further comprise (1) contacting a tumor cell with the compound prior to contacting the tumor cell with the second drug, (2) contacting a tumor cell with the second drug prior to contacting the tumor cell with the compound, or (3) contacting a tumor cell with the compound and the second drug at the same time. The second drug may, in certain embodiments, be an antibiotic, anti-inflammatory, anti-neoplastic, anti-proliferative, anti-viral, immunomodulatory, or immunosuppressive. The second drug may be an alkylating agent, androgen receptor modulator, cytoskeletal disruptor, estrogen receptor modulator, histone-deacetylase inhibitor, HMG-CoA reductase inhibitor, prenyl-protein transferase inhibitor, retinoid receptor modulator, topoisomerase inhibitor, or tyrosine kinase inhibitor. In certain embodiments, the second drug is 5-azacitidine, 5-fluorouracil, 9-cis-retinoic acid, actinomycin D, alitretinoin, all-trans-retinoic acid, annamycin, axitinib, belinostat, bevacizumab, bexarotene, bosutinib, busulfan, capecitabine, carboplatin, carmustine, CD437, cediranib, cetuximab, chlorambucil, cisplatin, cyclophosphamide, cytarabine, dacarbazine, dasatinib, daunorubicin, decitabine, docetaxel, dolastatin-10, doxifluridine, doxorubicin, doxorubicin, epirubicin, erlotinib, etoposide, etoposide, gefitinib, gemcitabine, gemtuzumab, ozogamicin, hexamethylmelamine, idarubicin, ifosfamide, imatinib, irinotecan, isotretinoin, ixabepilone, lapatinib, LBH589, lomustine, mechlorethamine, melphalan, mercaptopurine, methotrexate, mitomycin, mitoxantrone, MS-275, neratinib, nilotinib, nitrosourea, oxaliplatin, paclitaxel, plicamycin, procarbazine, semaxanib, semustine, sodium butyrate, sodium phenylacetate, streptozotocin, suberoylanilide hydroxamic acid, sunitinib, tamoxifen, teniposide, thiopeta, tioguanine,

topotecan, TRAIL, trastuzumab, tretinoin, trichostatin A, valproic acid, valrubicin, vandetanib, vinblastine, vincristine, vindesine, or vinorelbine.

Methods of treating or preventing a disease with an inflammatory component in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound of the present disclosure are also contemplated. The disease may be, for example, lupus or rheumatoid arthritis. The disease may be an inflammatory bowel disease, such as Crohn's disease or ulcerative colitis. The disease with an inflammatory component may be a cardiovascular disease. The disease with an inflammatory component may be diabetes, such as type 1 or type 2 diabetes. Compounds of the present disclosure may also be used to treat complications associated with diabetes. Such complications are well-known in the art and include, for example, obesity, hypertension, atherosclerosis, coronary heart disease, stroke, peripheral vascular disease, hypertension, nephropathy, neuropathy, myonecrosis, retinopathy and metabolic syndrome (syndrome X). The disease with an inflammatory component may be a skin disease, such as psoriasis, acne, or atopic dermatitis. Administration of a compound of the present disclosure in treatment methods of such skin diseases may be, for example, topical or oral.

The disease with an inflammatory component may be metabolic syndrome (syndrome X). A patient having this syndrome is characterized as having three or more symptoms selected from the following group of five symptoms: (1) abdominal obesity; (2) hypertriglyceridemia; (3) low high-density lipoprotein cholesterol (HDL); (4) high blood pressure; and (5) elevated fasting glucose, which may be in the range characteristic of Type 2 diabetes if the patient is also diabetic. Each of these symptoms is defined in the Third Report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III, or ATP III), National Institutes of Health, 2001, NIH Publication No. 01-3670, incorporated herein by reference. Patients with metabolic syndrome, whether or not they have or develop overt diabetes mellitus, have an increased risk of developing the macrovascular and microvascular complications that are listed above that occur with type 2 diabetes, such as atherosclerosis and coronary heart disease.

Another general method of the present disclosure entails a method of treating or preventing a cardiovascular disease in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound of the present disclosure. The cardiovascular disease may be, for example, atherosclerosis, cardiomyopathy, congenital

heart disease, congestive heart failure, myocarditis, rheumatic heart disease, valve disease, coronary artery disease, endocarditis, or myocardial infarction. Combination therapy is also contemplated for such methods. For example, such methods may further comprise administering a pharmaceutically effective amount of a second drug. The second drug may be, for example, a cholesterol lowering drug, an anti-hyperlipidemic, a calcium channel blocker, an anti-hypertensive, or an HMG-CoA reductase inhibitor. Non-limiting examples of second drugs include amlodipine, aspirin, ezetimibe, felodipine, lacidipine, lercanidipine, nicardipine, nifedipine, nimodipine, nisoldipine or nitrendipine. Other non-limiting examples of second drugs include atenolol, bucindolol, carvedilol, clonidine, doxazosin, indoramin, labetalol, methyldopa, metoprolol, nadolol, oxprenolol, phenoxybenzamine, phentolamine, pindolol, prazosin, propranolol, terazosin, timolol or tolazoline. The second drug may be, for example, a statin, such as atorvastatin, cerivastatin, fluvastatin, lovastatin, mevastatin, pitavastatin, pravastatin, rosuvastatin or simvastatin.

Methods of treating or preventing a neurodegenerative disease in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound of the present disclosure are also contemplated. The neurodegenerative disease may, for example, be selected from the group consisting of Parkinson's disease, Alzheimer's disease, multiple sclerosis (MS), Huntington's disease and amyotrophic lateral sclerosis. In particular embodiments, the neurodegenerative disease is Alzheimer's disease. In particular embodiments, the neurodegenerative disease is MS, such as primary progressive, relapsing-remitting secondary progressive or progressive relapsing MS. The subject may be, for example, a primate. The subject may be a human.

In particular embodiments of methods of treating or preventing a neurodegenerative disease in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound of the present disclosure, the treatment suppresses the demyelination of neurons in the subject's brain or spinal cord. In certain embodiments, the treatment suppresses inflammatory demyelination. In certain embodiments, the treatment suppresses the transection of neuron axons in the subject's brain or spinal cord. In certain embodiments, the treatment suppresses the transection of neurites in the subject's brain or spinal cord. In certain embodiments, the treatment suppresses neuronal apoptosis in the subject's brain or spinal cord. In certain embodiments, the treatment stimulates the remyelination of neuron axons in the subject's brain or spinal cord. In certain embodiments, the treatment restores lost function after an MS attack. In certain embodiments, the treatment prevents a new MS

attack. In certain embodiments, the treatment prevents a disability resulting from an MS attack.

One general aspect of the present disclosure contemplates a method of treating or preventing a disorder characterized by overexpression of iNOS genes in a subject, comprising
5 administering to the subject a pharmaceutically effective amount of a compound of the present disclosure.

Another general aspect of the present disclosure contemplates a method of inhibiting IFN- γ -induced nitric oxide production in cells of a subject, comprising administering to said subject a pharmaceutically effective amount of a compound of the present disclosure.

10 Yet another general method of the present disclosure contemplates a method of treating or preventing a disorder characterized by overexpression of COX-2 genes in a subject, comprising administering to the subject a pharmaceutically effective amount of compound of the present disclosure.

Methods of treating renal/kidney disease (RKD) in a subject, comprising
15 administering to the subject a pharmaceutically effective amount of a compound of the present disclosure are also contemplated. See U.S. Patent Application 12/352,473, which is incorporated by reference herein in its entirety. The RKD may result from, for example, a toxic insult. The toxic insult may result from, for example, an imaging agent or a drug. The drug may be a chemotherapeutic, for example. The RKD may result from
20 ischemia/reperfusion injury, in certain embodiments. In certain embodiments, the RKD results from diabetes or hypertension. The RKD may result from an autoimmune disease. The RKD may be further defined as chronic RKD, or acute RKD.

In certain methods of treating renal/kidney disease (RKD) in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound of the
25 present disclosure, the subject has undergone or is undergoing dialysis. In certain embodiments, the subject has undergone or is a candidate to undergo kidney transplant. The subject may be a primate. The primate may be a human. The subject in this or any other method may be, for example, a cow, horse, dog, cat, pig, mouse, rat or guinea pig.


Also contemplated by the present disclosure is a method for improving glomerular
30 filtration rate or creatinine clearance in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound of the present disclosure.

In some embodiments, the invention provides compounds useful for preventing and/or treating diseases or disorders whose pathology involves oxidative stress, inflammation, and/or


dysregulation of inflammatory signaling pathways. In some variations, the diseases or disorders can be characterized by overexpression of inducible nitric oxide synthase (iNOS) and/or inducible cyclooxygenase (COX-2) in affected tissues. In some variations, the diseases or disorders can be characterized by overproduction of reactive oxygen species (ROS) or reactive nitrogen species (RNS) such as superoxide, hydrogen peroxide, nitric oxide or peroxynitrite in affected tissues. In some variations, the disease or disorder is characterized by excessive production of inflammatory cytokines or other inflammation-related proteins such as TNF α , IL-6, IL-1, IL-8, ICAM-1, VCAM-1, and VEGF. Such diseases or disorders may, in some embodiments, involve undesirable proliferation of certain cells, as in the case of cancer (*e.g.*, solid tumors, leukemias, myelomas, lymphomas, and other cancers), fibrosis associated with organ failure, or excessive scarring. Non limiting examples of the disease or disorder include: lupus, rheumatoid arthritis, juvenile-onset diabetes, multiple sclerosis, psoriasis, and Crohn's disease. Further non-limiting examples include cardiovascular diseases, such as atherosclerosis, heart failure, myocardial infarction, acute coronary syndrome, restenosis following vascular surgery, hypertension, and vasculitis; neurodegenerative or neuromuscular diseases such as Alzheimer's disease, Parkinson's disease, Huntington's disease, ALS, and muscular dystrophy; neurological disorders such as epilepsy and dystonia; neuropsychiatric conditions such as major depression, bipolar disorder, post-traumatic stress disorder, schizophrenia, anorexia nervosa, ADHD, and autism-spectrum disorders; retinal diseases such as macular degeneration, diabetic retinopathy, glaucoma, and retinitis; chronic and acute pain syndromes, including inflammatory and neuropathic pain; hearing loss and tinnitus; diabetes and complications of diabetes, including metabolic syndrome, diabetic nephropathy, diabetic neuropathy and diabetic ulcers; respiratory diseases such as asthma, chronic obstructive pulmonary disease, acute respiratory distress syndrome, and cystic fibrosis; inflammatory bowel diseases; osteoporosis, osteoarthritis, and other degenerative conditions of bone and cartilage; acute or chronic organ failure, including renal failure, liver failure (including cirrhosis and hepatitis), and pancreatitis; ischemia-reperfusion injury associated with thrombotic or hemorrhagic stroke, subarachnoid hemorrhage, cerebral vasospasm, myocardial infarction, shock, or trauma; complications of organ or tissue transplantation including acute or chronic transplant failure or rejection and graft-versus-host disease; skin diseases including atopic dermatitis and acne; sepsis and septic shock; excessive inflammation associated with infection, including respiratory inflammation associated with

influenza and upper respiratory infections; mucositis associated with cancer therapy, including radiation therapy or chemotherapy; and severe burns.

Methods of synthesizing compounds of the present disclosure are also contemplated. In particular embodiments, such methods can comprise a method of making a target compound defined of the formula:



wherein R_a is alkoxy_(C1-4), comprising reacting a compound of the formula:



with an oxidizing agent under a set of conditions to form the target compound.

Kits are also contemplated by the present disclosure, such as a kit comprising: a compound of the present disclosure; and instructions which comprise one or more forms of information selected from the group consisting of indicating a disease state for which the compound is to be administered, storage information for the compound, dosing information and instructions regarding how to administer the compound. The kit may comprise a compound of the present disclosure in a multiple dose form.

Other objects, features and advantages of the present disclosure will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description. Note that simply because a particular compound is ascribed to one particular generic formula doesn't mean that it cannot also belong to another generic formula.

BRIEF DESCRIPTION OF THE DRAWINGS

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present disclosure. The invention may be better understood by reference to one of these drawings in combination with the detailed description of specific embodiments presented herein.

FIGS. 1-8 and 32-34. Inhibition of NO Production. RAW264.7 macrophages were pre-treated with DMSO or drugs at various concentrations (nM) for 2 hours, then treated with 20 ng/ml IFN γ for 24 hours. NO concentration in media was determined using a Griess reagent system; cell viability was determined using WST-1 reagent.

FIG. 9. Suppression of COX-2 Induction. RAW264.7 cells were pre-treated for 2 hours with indicated compounds and subsequently stimulated with 10 ng/ml IFN γ for an additional 24 hours. COX-2 protein levels were assayed by immunoblotting. Actin was used as a loading control. **RTA 402** and **RTA 404** refer to comparison compounds **402** and **404** (see Example 1).

FIGS. 10-12. Inhibition of IL-6 Induced STAT3 Phosphorylation. HeLa cells were treated with the indicated compounds and concentrations for 6 hours and subsequently stimulated with 20 ng/ml IL-6 for 15 minutes. Phosphorylated STAT3 and total STAT3 levels were assayed by immunoblotting. Compounds **402-52** and **402-53** are comparison compounds (see Example 1).

FIG. 13. Suppression of IL-6 Induced STAT3 Phosphorylation. HeLa cells were treated with DMSO or the indicated compounds at 2 μ M for 6 hours and subsequently stimulated with 20 ng/ml IL-6 for 15 minutes. Phosphorylated STAT3 and total STAT3 levels were assayed by immunoblotting. Compounds **402-54**, **402-55** and **402-56** are comparison compounds (see Example 1).

FIG. 14. Inhibition of TNF α -induced I κ B α degradation. HeLa cells were treated with indicated compounds and concentrations for 6 hours and subsequently stimulated with 20 ng/ml TNF α for 15 minutes. Lysates were analyzed with antibodies against I κ B α and actin.

FIGS. 15 and 16. Inhibition of NF κ B activation. HeLa cells were transfected with pNF- κ B-Luc (inducible) and pRL-TK (constitutive) reporter plasmids. Twenty-four hours later cells were pre-treated with the indicated compounds for 2 hours. DMSO served as a

vehicle control. Following pre-treatment, cells were stimulated with 20 ng/ml TNF α for 3 hours. Reporter activity was measured using DualGlo luciferase reporter assay and pNF- κ B luciferase activity was normalized against pRL-TK luciferase activity. Fold-induction of mean luciferase activity relative to unstimulated (-TNF α) samples is shown. Error bars represent the SD of the mean of 6 samples.

FIGS. 17- 20. Induction of HO-1. MDA-MB-435 human melanoma cells were treated with vehicle (DMSO) or the indicated compounds and concentrations for 16 hours. HO-1 mRNA levels were quantified using qPCR and were normalized relative to a DMSO-treated sample run in parallel. Values are averages of duplicate wells.

FIG. 21. Induction of HO-1, TrxR1 and γ -GCS. MDA-MB-435 human melanoma cells were treated with vehicle (DMSO) or the indicated compounds and concentrations for 16 hours. HO-1, thioredoxin reductase-1 (TrxR1), and γ -glutamylcysteine synthetase (γ -GCS) mRNA levels were quantified using qPCR and were normalized relative to a DMSO-treated sample run in parallel. Values are averages of duplicate wells.

FIG. 22. Induction of TrxR1. MDA-MB-435 human melanoma cells were treated with vehicle (DMSO) or the indicated compounds and concentrations for 16 hours. Thioredoxin reductase-1 (TrxR1) mRNA levels were quantified using qPCR and were normalized relative to a DMSO-treated sample run in parallel. Values are averages of duplicate wells. Compounds **401**, **402-19** and **402-53** are comparison compounds (see Example 1). Comparison with the results of FIG. 25 demonstrates that higher concentrations of **402-02** and **404-02** are required to approach effects seen with the unsaturated counterpart compounds **402** and **404**.

FIG. 23. Induction of γ -GCS. MDA-MB-435 human melanoma cells were treated with vehicle (DMSO) or the indicated compounds and concentrations for 16 hours. γ -glutamylcysteine synthetase (γ -GCS) mRNA levels were quantified using qPCR and were normalized relative to a DMSO-treated sample run in parallel. Values are averages of duplicate wells. Comparison with the results of FIG. 26 demonstrates that higher concentrations of **402-02** and **404-02** are required to approach effects seen with the unsaturated counterpart compounds **402** and **404**.

FIG. 24. Induction of Ferritin Heavy Chain. MDA-MB-435 human melanoma cells were treated with vehicle (DMSO) or the indicated compounds and concentrations for

16 hours. Ferritin heavy chain mRNA levels were quantified using qPCR and were normalized relative to a DMSO-treated sample run in parallel. Values are averages of duplicate wells. Comparison with the results of FIG. 27 demonstrates that higher concentrations of **402-02** and **404-02** are required to approach effects seen with the unsaturated counterpart compounds **402** and **404**.

FIG. 25. Induction of TrxR1. MDA-MB-435 human melanoma cells were treated with vehicle (DMSO) or the indicated compounds and concentrations for 16 hours. Thioredoxin reductase-1 (TrxR1) mRNA levels were quantified using qPCR and were normalized relative to a DMSO-treated sample run in parallel. Values are averages of duplicate wells. Comparison with the results of FIG. 22 demonstrates that higher concentrations of **402-02** and **404-02** are required to approach effects seen with the unsaturated counterpart compounds **402** and **404**.

FIG. 26. Induction of γ -GCS. MDA-MB-435 human melanoma cells were treated with vehicle (DMSO) or the indicated compounds and concentrations for 16 hours. γ -glutamylcysteine synthetase (γ -GCS) mRNA levels were quantified using qPCR and were normalized relative to a DMSO-treated sample run in parallel. Values are averages of duplicate wells. Comparison with the results of FIG. 23 demonstrates that higher concentrations of **402-02** and **404-02** are required to approach effects seen with the unsaturated counterpart compounds **402** and **404**.

FIG. 27. Induction of Ferritin Heavy Chain. MDA-MB-435 human melanoma cells were treated with vehicle (DMSO) or the indicated compounds and concentrations for 16 hours. Ferritin heavy chain mRNA levels were quantified using qPCR and were normalized relative to a DMSO-treated sample run in parallel. Values are averages of duplicate wells. Comparison with the results of FIG. 24 demonstrates that higher concentrations of **402-02** and **404-02** are required to approach effects seen with the unsaturated counterpart compounds **402** and **404**.

FIG. 28 – CDDO-TFEA (TP-500) Is Detected at Higher Levels in Mouse Brain than CDDO-EA (TP-319). CD-1 mice were fed either 200 or 400 mg/kg diet of either TP-319 or TP-500 for 3.5 days, and TP levels in the brains of the mice were analyzed by LC/MS. The structures of TP-319 and TP-500 are shown below.

FIGS. 29 A & B – Weight Change Data from a Head to Head Tox Study of Compounds 401 (FIG 29A) versus 401-02 (FIG. 29B). Compounds were assessed for toxicity in mice in a 14-day study. Each compound was formulated in sesame oil and administered daily by oral gavage at doses of 10, 50, 100, or 250 mg/kg (n = 4 per group).

5 **FIGS. 30 A & B – Weight Change Data from a Head to Head Tox Study of Compounds 402 (FIG 30A) versus 402-02 (FIG. 30B).** Compounds were assessed for toxicity in mice in a 14-day study. Each compound was formulated in sesame oil and administered daily by oral gavage at doses of 10, 50, 100, or 250 mg/kg (n = 4 per group).

10 **FIGS. 31 A & B – Weight Change Data from a Head to Head Tox Study of Compounds 404 (FIG 31A) versus 404-02 (FIG. 31B).** Compounds were assessed for toxicity in mice in a 14-day study. Each compound was formulated in sesame oil and administered daily by oral gavage at doses of 10, 50, 100, or 250 mg/kg (n = 4 per group).

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

15 Disclosed herein are, for example, new compounds with antioxidant and anti-inflammatory properties, methods for their manufacture, and methods for their use, including for the treatment and/or prevention of disease.

I. Definitions

As used herein, “hydrogen” means –H; “hydroxy” means –OH; “oxo” means =O; 20 “halo” means independently –F, –Cl, –Br or –I; “amino” means –NH₂ (see below for definitions of groups containing the term amino, *e.g.*, alkylamino); “hydroxyamino” means –NHOH; “nitro” means –NO₂; imino means =NH (see below for definitions of groups containing the term imino, *e.g.*, alkylamino); “cyano” means –CN; “azido” means –N₃; “mercapto” means –SH; “thio” means =S; “sulfonamido” means –NHS(O)₂– (see below for 25 definitions of groups containing the term sulfonamido, *e.g.*, alkylsulfonamido); “sulfonyl” means –S(O)₂– (see below for definitions of groups containing the term sulfonyl, *e.g.*, alkylsulfonyl); and “silyl” means –SiH₃ (see below for definitions of group(s) containing the term silyl, *e.g.*, alkylsilyl).


For the groups below, the following parenthetical subscripts further define the groups 30 as follows: “(Cn)” defines the exact number (n) of carbon atoms in the group. “(C≤n)” defines the maximum number (n) of carbon atoms that can be in the group, with the minimum number of carbon atoms in such at least one, but otherwise as small as possible for the group

in question. *E.g.*, it is understood that the minimum number of carbon atoms in the group “alkenyl_(C≤8)” is 2. For example, “alkoxy_(C≤10)” designates those alkoxy groups having from 1 to 10 carbon atoms (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, or any range derivable therein (*e.g.*, 3-10 carbon atoms)). (C_n-n') defines both the minimum (n) and maximum number (n') of carbon atoms in the group. Similarly, “alkyl_(C2-10)” designates those alkyl groups having from 2 to 10 carbon atoms (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or 10, or any range derivable therein (*e.g.*, 3-10 carbon atoms)).

The term “alkyl” when used without the “substituted” modifier refers to a non-aromatic monovalent group with a saturated carbon atom as the point of attachment, a linear or branched, cyclo, cyclic or acyclic structure, no carbon-carbon double or triple bonds, and no atoms other than carbon and hydrogen. The groups, -CH₃ (Me), -CH₂CH₃ (Et), -CH₂CH₂CH₃ (*n*-Pr), -CH(CH₃)₂ (*iso*-Pr), -CH(CH₂)₂ (cyclopropyl), -CH₂CH₂CH₂CH₃ (*n*-Bu), -CH(CH₃)CH₂CH₃ (*sec*-butyl), -CH₂CH(CH₃)₂ (*iso*-butyl), -C(CH₃)₃ (*tert*-butyl), -CH₂C(CH₃)₃ (*neo*-pentyl), cyclobutyl, cyclopentyl, cyclohexyl, and cyclohexylmethyl are non-limiting examples of alkyl groups. The term “substituted alkyl” refers to a non-aromatic monovalent group with a saturated carbon atom as the point of attachment, a linear or branched, cyclo, cyclic or acyclic structure, no carbon-carbon double or triple bonds, and at least one atom independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. The following groups are non-limiting examples of substituted alkyl groups:

-CH₂OH, -CH₂Cl, -CH₂Br, -CH₂SH, -CF₃, -CH₂CN, -CH₂C(O)H, -CH₂C(O)OH, -CH₂C(O)OCH₃, -CH₂C(O)NH₂, -CH₂C(O)NHCH₃, -CH₂C(O)CH₃, -CH₂OCH₃, -CH₂OCH₂CF₃, -CH₂OC(O)CH₃, -CH₂NH₂, -CH₂NHCH₃, -CH₂N(CH₃)₂, -CH₂CH₂Cl, -CH₂CH₂OH, -CH₂CF₃, -CH₂CH₂OC(O)CH₃, -CH₂CH₂NHCO₂C(CH₃)₃, and -CH₂Si(CH₃)₃.


The term “alkanediyl” when used without the “substituted” modifier refers to a non-aromatic divalent group, wherein the alkanediyl group is attached with two σ-bonds, with one or two saturated carbon atom(s) as the point(s) of attachment, a linear or branched, cyclo, cyclic or acyclic structure, no carbon-carbon double or triple bonds, and no atoms other than carbon and hydrogen. The groups, -CH₂- (methylene), -CH₂CH₂-, -CH₂C(CH₃)₂CH₂-,

-CH₂CH₂CH₂-, and , are non-limiting examples of alkanediyl groups. The term “substituted alkanediyl” refers to a non-aromatic monovalent group, wherein the alkynediyl group is attached with two σ-bonds, with one or two saturated carbon atom(s) as

the point(s) of attachment, a linear or branched, cyclo, cyclic or acyclic structure, no carbon-carbon double or triple bonds, and at least one atom independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. The following groups are non-limiting examples of substituted alkanediyl groups: $-\text{CH}(\text{F})-$, $-\text{CF}_2-$, $-\text{CH}(\text{Cl})-$, $-\text{CH}(\text{OH})-$, $-\text{CH}(\text{OCH}_3)-$,
 5 and $-\text{CH}_2\text{CH}(\text{Cl})-$.

The term “alkenyl” when used without the “substituted” modifier refers to a monovalent group with a nonaromatic carbon atom as the point of attachment, a linear or branched, cyclo, cyclic or acyclic structure, at least one nonaromatic carbon-carbon double bond, no carbon-carbon triple bonds, and no atoms other than carbon and hydrogen. Non-
 10 limiting examples of alkenyl groups include: $-\text{CH}=\text{CH}_2$ (vinyl), $-\text{CH}=\text{CHCH}_3$, $-\text{CH}=\text{CHCH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}=\text{CH}_2$ (allyl), $-\text{CH}_2\text{CH}=\text{CHCH}_3$, and $-\text{CH}=\text{CH}-\text{C}_6\text{H}_5$. The term “substituted alkenyl” refers to a monovalent group with a nonaromatic carbon atom as the point of attachment, at least one nonaromatic carbon-carbon double bond, no carbon-carbon triple bonds, a linear or branched, cyclo, cyclic or acyclic structure, and at least one atom
 15 independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. The groups, $-\text{CH}=\text{CHF}$, $-\text{CH}=\text{CHCl}$ and $-\text{CH}=\text{CHBr}$, are non-limiting examples of substituted alkenyl groups.

The term “alkenediyl” when used without the “substituted” modifier refers to a non-aromatic divalent group, wherein the alkenediyl group is attached with two σ -bonds, with two
 20 carbon atoms as points of attachment, a linear or branched, cyclo, cyclic or acyclic structure, at least one nonaromatic carbon-carbon double bond, no carbon-carbon triple bonds, and no atoms other than carbon and hydrogen. The groups, $-\text{CH}=\text{CH}-$, $-\text{CH}=\text{C}(\text{CH}_3)\text{CH}_2-$,

$-\text{CH}=\text{CHCH}_2-$, and , are non-limiting examples of alkenediyl groups. The
 term “substituted alkenediyl” refers to a non-aromatic divalent group, wherein the alkenediyl
 25 group is attached with two σ -bonds, with two carbon atoms as points of attachment, a linear or branched, cyclo, cyclic or acyclic structure, at least one nonaromatic carbon-carbon double bond, no carbon-carbon triple bonds, and at least one atom independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. The following groups are non-limiting examples of substituted alkenediyl groups: $-\text{CF}=\text{CH}-$, $-\text{C}(\text{OH})=\text{CH}-$, and $-\text{CH}_2\text{CH}=\text{C}(\text{Cl})-$.

The term “alkynyl” when used without the “substituted” modifier refers to a monovalent group with a nonaromatic carbon atom as the point of attachment, a linear or branched, cyclo, cyclic or acyclic structure, at least one carbon-carbon triple bond, and no


atoms other than carbon and hydrogen. The groups, $-\text{C}\equiv\text{CH}$, $-\text{C}\equiv\text{CCH}_3$, $-\text{C}\equiv\text{CC}_6\text{H}_5$ and $-\text{CH}_2\text{C}\equiv\text{CCH}_3$, are non-limiting examples of alkynyl groups. The term “substituted alkynyl” refers to a monovalent group with a nonaromatic carbon atom as the point of attachment and at least one carbon-carbon triple bond, a linear or branched, cyclo, cyclic or acyclic structure, and at least one atom independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. The group, $-\text{C}\equiv\text{CSi}(\text{CH}_3)_3$, is a non-limiting example of a substituted alkynyl group.

The term “alkynediyl” when used without the “substituted” modifier refers to a non-aromatic divalent group, wherein the alkynediyl group is attached with two σ -bonds, with two carbon atoms as points of attachment, a linear or branched, cyclo, cyclic or acyclic structure, at least one carbon-carbon triple bond, and no atoms other than carbon and hydrogen. The groups, $-\text{C}\equiv\text{C}-$, $-\text{C}\equiv\text{CCH}_2-$, and $-\text{C}\equiv\text{CCH}(\text{CH}_3)-$ are non-limiting examples of alkynediyl groups. The term “substituted alkynediyl” refers to a non-aromatic divalent group, wherein the alkynediyl group is attached with two σ -bonds, with two carbon atoms as points of attachment, a linear or branched, cyclo, cyclic or acyclic structure, at least one carbon-carbon triple bond, and at least one atom independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. The groups $-\text{C}\equiv\text{CCFH}-$ and $-\text{C}\equiv\text{CHCH}(\text{Cl})-$ are non-limiting examples of substituted alkynediyl groups.

The term “aryl” when used without the “substituted” modifier refers to a monovalent group with an aromatic carbon atom as the point of attachment, said carbon atom forming part of a six-membered aromatic ring structure wherein the ring atoms are all carbon, and wherein the monovalent group consists of no atoms other than carbon and hydrogen. Non-limiting examples of aryl groups include phenyl (Ph), methylphenyl, (dimethyl)phenyl, $-\text{C}_6\text{H}_4\text{CH}_2\text{CH}_3$ (ethylphenyl), $-\text{C}_6\text{H}_4\text{CH}_2\text{CH}_2\text{CH}_3$ (propylphenyl), $-\text{C}_6\text{H}_4\text{CH}(\text{CH}_3)_2$, $-\text{C}_6\text{H}_4\text{CH}(\text{CH}_2)_2$, $-\text{C}_6\text{H}_3(\text{CH}_3)\text{CH}_2\text{CH}_3$ (methylethylphenyl), $-\text{C}_6\text{H}_4\text{CH}=\text{CH}_2$ (vinylphenyl), $-\text{C}_6\text{H}_4\text{CH}=\text{CHCH}_3$, $-\text{C}_6\text{H}_4\text{C}\equiv\text{CH}$, $-\text{C}_6\text{H}_4\text{C}\equiv\text{CCH}_3$, naphthyl, and the monovalent group derived from biphenyl. The term “substituted aryl” refers to a monovalent group with an aromatic carbon atom as the point of attachment, said carbon atom forming part of a six-membered aromatic ring structure wherein the ring atoms are all carbon, and wherein the monovalent group further has at least one atom independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. Non-limiting examples of substituted aryl groups include the groups: $-\text{C}_6\text{H}_4\text{F}$, $-\text{C}_6\text{H}_4\text{Cl}$, $-\text{C}_6\text{H}_4\text{Br}$, $-\text{C}_6\text{H}_4\text{I}$, $-\text{C}_6\text{H}_4\text{OH}$, $-\text{C}_6\text{H}_4\text{OCH}_3$, $-\text{C}_6\text{H}_4\text{OCH}_2\text{CH}_3$, $-\text{C}_6\text{H}_4\text{OC}(\text{O})\text{CH}_3$, $-\text{C}_6\text{H}_4\text{NH}_2$, $-\text{C}_6\text{H}_4\text{NHCH}_3$, $-\text{C}_6\text{H}_4\text{N}(\text{CH}_3)_2$,

$-\text{C}_6\text{H}_4\text{CH}_2\text{OH}$, $-\text{C}_6\text{H}_4\text{CH}_2\text{OC}(\text{O})\text{CH}_3$, $-\text{C}_6\text{H}_4\text{CH}_2\text{NH}_2$, $-\text{C}_6\text{H}_4\text{CF}_3$, $-\text{C}_6\text{H}_4\text{CN}$, $-\text{C}_6\text{H}_4\text{CHO}$,
 $-\text{C}_6\text{H}_4\text{CHO}$, $-\text{C}_6\text{H}_4\text{C}(\text{O})\text{CH}_3$, $-\text{C}_6\text{H}_4\text{C}(\text{O})\text{C}_6\text{H}_5$, $-\text{C}_6\text{H}_4\text{CO}_2\text{H}$, $-\text{C}_6\text{H}_4\text{CO}_2\text{CH}_3$, $-\text{C}_6\text{H}_4\text{CONH}_2$,
 $-\text{C}_6\text{H}_4\text{CONHCH}_3$, and $-\text{C}_6\text{H}_4\text{CON}(\text{CH}_3)_2$.

The term “arenediyl” when used without the “substituted” modifier refers to a divalent
 5 group, wherein the arenediyl group is attached with two σ -bonds, with two aromatic carbon
 atoms as points of attachment, said carbon atoms forming part of one or more six-membered
 aromatic ring structure(s) wherein the ring atoms are all carbon, and wherein the monovalent
 group consists of no atoms other than carbon and hydrogen. Non-limiting examples of
 arenediyl groups include:



The term “substituted arenediyl” refers to a divalent group, wherein the arenediyl group is
 attached with two σ -bonds, with two aromatic carbon atoms as points of attachment, said
 carbon atoms forming part of one or more six-membered aromatic rings structure(s), wherein
 the ring atoms are all carbon, and wherein the divalent group further has at least one atom
 15 independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S.


The term “aralkyl” when used without the “substituted” modifier refers to the
 monovalent group $-\text{alkanediyl-aryl}$, in which the terms alkanediyl and aryl are each used in a
 manner consistent with the definitions provided above. Non-limiting examples of aralkyls
 are: phenylmethyl (benzyl, Bn), 1-phenyl-ethyl, 2-phenyl-ethyl, indenyl and 2,3-dihydro-
 20 indenyl, provided that indenyl and 2,3-dihydro-indenyl are only examples of aralkyl in so far
 as the point of attachment in each case is one of the saturated carbon atoms. When the term
 “aralkyl” is used with the “substituted” modifier, either one or both the alkanediyl and the
 aryl is substituted. Non-limiting examples of substituted aralkyls are: (3-chlorophenyl)-
 methyl, 2-oxo-2-phenyl-ethyl (phenylcarbonylmethyl), 2-chloro-2-phenyl-ethyl, chromanyl
 25 where the point of attachment is one of the saturated carbon atoms, and tetrahydroquinolinyl
 where the point of attachment is one of the saturated atoms.

The term “heteroaryl” when used without the “substituted” modifier refers to a
 monovalent group with an aromatic carbon atom or nitrogen atom as the point of attachment,
 said carbon atom or nitrogen atom forming part of an aromatic ring structure wherein at least
 30 one of the ring atoms is nitrogen, oxygen or sulfur, and wherein the monovalent group
 consists of no atoms other than carbon, hydrogen, aromatic nitrogen, aromatic oxygen and

aromatic sulfur. Non-limiting examples of aryl groups include acridinyl, furanyl, imidazoimidazolyl, imidazopyrazolyl, imidazopyridinyl, imidazopyrimidinyl, indolyl, indazolyl, methylpyridyl, oxazolyl, phenylimidazolyl, pyridyl, pyrrolyl, pyrimidyl, pyrazinyl, quinolyl, quinazolyl, quinoxalinyl, tetrahydroquinolyl, thienyl, triazinyl, pyrrolopyridinyl, pyrrolopyrimidinyl, pyrrolopyrazinyl, pyrrolotriazinyl, pyrroloimidazolyl, chromenyl (where the point of attachment is one of the aromatic atoms), and chromanyl (where the point of attachment is one of the aromatic atoms). The term “substituted heteroaryl” refers to a monovalent group with an aromatic carbon atom or nitrogen atom as the point of attachment, said carbon atom or nitrogen atom forming part of an aromatic ring structure wherein at least one of the ring atoms is nitrogen, oxygen or sulfur, and wherein the monovalent group further has at least one atom independently selected from the group consisting of non-aromatic nitrogen, non-aromatic oxygen, non aromatic sulfur F, Cl, Br, I, Si, and P.

The term “heteroarenediyl” when used without the “substituted” modifier refers to a divalent group, wherein the heteroarenediyl group is attached with two σ -bonds, with an aromatic carbon atom or nitrogen atom as the point of attachment, said carbon atom or nitrogen atom two aromatic atoms as points of attachment, said carbon atoms forming part of one or more six-membered aromatic ring structure(s) wherein the ring atoms are all carbon, and wherein the monovalent group consists of no atoms other than carbon and hydrogen.

Non-limiting examples of heteroarenediyl groups include:



The term “substituted heteroarenediyl” refers to a divalent group, wherein the heteroarenediyl group is attached with two σ -bonds, with two aromatic carbon atoms as points of attachment, said carbon atoms forming part of one or more six-membered aromatic rings structure(s), wherein the ring atoms are all carbon, and wherein the divalent group further has at least one atom independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S.

The term “heteroaralkyl” when used without the “substituted” modifier refers to the monovalent group –alkanediyl–heteroaryl, in which the terms alkanediyl and heteroaryl are each used in a manner consistent with the definitions provided above. Non-limiting examples of aralkyls are: pyridylmethyl, and thienylmethyl. When the term “heteroaralkyl” is used

with the “substituted” modifier, either one or both the alkanediyl and the heteroaryl is substituted.

The term “acyl” when used without the “substituted” modifier refers to a monovalent group with a carbon atom of a carbonyl group as the point of attachment, further having a linear or branched, cyclo, cyclic or acyclic structure, further having no additional atoms that are not carbon or hydrogen, beyond the oxygen atom of the carbonyl group. The groups, $-\text{CHO}$, $-\text{C}(\text{O})\text{CH}_3$ (acetyl, Ac), $-\text{C}(\text{O})\text{CH}_2\text{CH}_3$, $-\text{C}(\text{O})\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{C}(\text{O})\text{CH}(\text{CH}_3)_2$, $-\text{C}(\text{O})\text{CH}(\text{CH}_2)_2$, $-\text{C}(\text{O})\text{C}_6\text{H}_5$, $-\text{C}(\text{O})\text{C}_6\text{H}_4\text{CH}_3$, $-\text{C}(\text{O})\text{C}_6\text{H}_4\text{CH}_2\text{CH}_3$, $-\text{COC}_6\text{H}_3(\text{CH}_3)_2$, and $-\text{C}(\text{O})\text{CH}_2\text{C}_6\text{H}_5$, are non-limiting examples of acyl groups. The term “acyl” therefore encompasses, but is not limited to groups sometimes referred to as “alkyl carbonyl” and “aryl carbonyl” groups. The term “substituted acyl” refers to a monovalent group with a carbon atom of a carbonyl group as the point of attachment, further having a linear or branched, cyclo, cyclic or acyclic structure, further having at least one atom, in addition to the oxygen of the carbonyl group, independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. The groups, $-\text{C}(\text{O})\text{CH}_2\text{CF}_3$, $-\text{CO}_2\text{H}$ (carboxyl), $-\text{CO}_2\text{CH}_3$ (methylcarboxyl), $-\text{CO}_2\text{CH}_2\text{CH}_3$, $-\text{CO}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CO}_2\text{C}_6\text{H}_5$, $-\text{CO}_2\text{CH}(\text{CH}_3)_2$, $-\text{CO}_2\text{CH}(\text{CH}_2)_2$, $-\text{C}(\text{O})\text{NH}_2$ (carbamoyl), $-\text{C}(\text{O})\text{NHCH}_3$, $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_3$, $-\text{CONHCH}(\text{CH}_3)_2$, $-\text{CONHCH}(\text{CH}_2)_2$, $-\text{CON}(\text{CH}_3)_2$, $-\text{CONHCH}_2\text{CF}_3$, $-\text{CO}$ -pyridyl, $-\text{CO}$ -imidazolyl, and $-\text{C}(\text{O})\text{N}_3$, are non-limiting examples of substituted acyl groups. The term “substituted acyl” encompasses, but is not limited to, “heteroaryl carbonyl” groups.

The term “alkylidene” when used without the “substituted” modifier refers to the divalent group $=\text{CRR}'$, wherein the alkylidene group is attached with one σ -bond and one π -bond, in which R and R' are independently hydrogen, alkyl, or R and R' are taken together to represent alkanediyl. Non-limiting examples of alkylidene groups include: $=\text{CH}_2$, $=\text{CH}(\text{CH}_2\text{CH}_3)$, and $=\text{C}(\text{CH}_3)_2$. The term “substituted alkylidene” refers to the group $=\text{CRR}'$, wherein the alkylidene group is attached with one σ -bond and one π -bond, in which R and R' are independently hydrogen, alkyl, substituted alkyl, or R and R' are taken together to represent a substituted alkanediyl, provided that either one of R and R' is a substituted alkyl or R and R' are taken together to represent a substituted alkanediyl.

The term “alkoxy” when used without the “substituted” modifier refers to the group $-\text{OR}$, in which R is an alkyl, as that term is defined above. Non-limiting examples of alkoxy groups include: $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}_3$, $-\text{OCH}_2\text{CH}_2\text{CH}_3$, $-\text{OCH}(\text{CH}_3)_2$, $-\text{OCH}(\text{CH}_2)_2$, $-\text{O}$ -cyclopentyl, and $-\text{O}$ -cyclohexyl. The term “substituted alkoxy” refers to the group

–OR, in which R is a substituted alkyl, as that term is defined above. For example, –OCH₂CF₃ is a substituted alkoxy group.

Similarly, the terms “alkenyloxy”, “alkynyloxy”, “aryloxy”, “aralkoxy”, “heteroaryloxy”, “heteroaralkoxy” and “acyloxy”, when used without the “substituted” modifier, refers to groups, defined as –OR, in which R is alkenyl, alkynyl, aryl, aralkyl, heteroaryl, heteroaralkyl and acyl, respectively, as those terms are defined above. When any of the terms alkenyloxy, alkynyloxy, aryloxy, aralkyloxy and acyloxy is modified by “substituted,” it refers to the group –OR, in which R is substituted alkenyl, alkynyl, aryl, aralkyl, heteroaryl, heteroaralkyl and acyl, respectively.

The term “alkylamino” when used without the “substituted” modifier refers to the group –NHR, in which R is an alkyl, as that term is defined above. Non-limiting examples of alkylamino groups include: –NHCH₃, –NHCH₂CH₃, –NHCH₂CH₂CH₃, –NHCH(CH₃)₂, –NHCH(CH₂)₂, –NHCH₂CH₂CH₂CH₃, –NHCH(CH₃)CH₂CH₃, –NHCH₂CH(CH₃)₂, –NHC(CH₃)₃, –NH–cyclopentyl, and –NH–cyclohexyl. The term “substituted alkylamino” refers to the group –NHR, in which R is a substituted alkyl, as that term is defined above. For example, –NHCH₂CF₃ is a substituted alkylamino group.

The term “dialkylamino” when used without the “substituted” modifier refers to the group –NRR', in which R and R' can be the same or different alkyl groups, or R and R' can be taken together to represent an alkanediyl having two or more saturated carbon atoms, at least two of which are attached to the nitrogen atom. Non-limiting examples of dialkylamino groups include: –NHC(CH₃)₃, –N(CH₃)CH₂CH₃, –N(CH₂CH₃)₂, *N*-pyrrolidinyl, and *N*-piperidinyl. The term “substituted dialkylamino” refers to the group –NRR', in which R and R' can be the same or different substituted alkyl groups, one of R or R' is an alkyl and the other is a substituted alkyl, or R and R' can be taken together to represent a substituted alkanediyl with two or more saturated carbon atoms, at least two of which are attached to the nitrogen atom.

The terms “alkoxyamino”, “alkenylamino”, “alkynylamino”, “arylamino”, “aralkylamino”, “heteroarylamino”, “heteroaralkylamino”, and “alkylsulfonylamino” when used without the “substituted” modifier, refers to groups, defined as –NHR, in which R is alkoxy, alkenyl, alkynyl, aryl, aralkyl, heteroaryl, heteroaralkyl and alkylsulfonyl, respectively, as those terms are defined above. A non-limiting example of an arylamino group is –NHC₆H₅. When any of the terms alkoxyamino, alkenylamino, alkynylamino, arylamino, aralkylamino, heteroarylamino, heteroaralkylamino and alkylsulfonylamino is

modified by “substituted,” it refers to the group -NHR , in which R is substituted alkenyl, alkenyl, alkynyl, aryl, aralkyl, heteroaryl, heteroaralkyl and alkylsulfonyl, respectively.

The term “amido” (acylamino), when used without the “substituted” modifier, refers to the group -NHR , in which R is acyl, as that term is defined above. A non-limiting example of an acylamino group is -NHC(O)CH_3 . When the term amido is used with the “substituted” modifier, it refers to groups, defined as -NHR , in which R is substituted acyl, as that term is defined above. The groups -NHC(O)OCH_3 and -NHC(O)NHCH_3 are non-limiting examples of substituted amido groups.

The term “alkylimino” when used without the “substituted” modifier refers to the group =NR , wherein the alkylimino group is attached with one σ -bond and one π -bond, in which R is an alkyl, as that term is defined above. Non-limiting examples of alkylimino groups include: =NCH_3 , $\text{=NCH}_2\text{CH}_3$ and =N-cyclohexyl . The term “substituted alkylimino” refers to the group =NR , wherein the alkylimino group is attached with one σ -bond and one π -bond, in which R is a substituted alkyl, as that term is defined above. For example, $\text{=NCH}_2\text{CF}_3$ is a substituted alkylimino group.

Similarly, the terms “alkenylimino”, “alkynylimino”, “arylimino”, “aralkylimino”, “heteroarylimino”, “heteroaralkylimino” and “acylimino”, when used without the “substituted” modifier, refers to groups, defined as =NR , wherein the alkylimino group is attached with one σ -bond and one π -bond, in which R is alkenyl, alkynyl, aryl, aralkyl, heteroaryl, heteroaralkyl and acyl, respectively, as those terms are defined above. When any of the terms alkenylimino, alkynylimino, arylimino, aralkylimino and acylimino is modified by “substituted,” it refers to the group =NR , wherein the alkylimino group is attached with one σ -bond and one π -bond, in which R is substituted alkenyl, alkynyl, aryl, aralkyl, heteroaryl, heteroaralkyl and acyl, respectively.

The term “fluoroalkyl” when used without the “substituted” modifier refers to an alkyl, as that term is defined above, in which one or more fluorines have been substituted for hydrogens. The groups, $\text{-CH}_2\text{F}$, -CF_3 , and $\text{-CH}_2\text{CF}_3$ are non-limiting examples of fluoroalkyl groups. The term “substituted fluoroalkyl” refers to a non-aromatic monovalent group with a saturated carbon atom as the point of attachment, a linear or branched, cyclic, cyclic or acyclic structure, at least one fluorine atom, no carbon-carbon double or triple bonds, and at least one atom independently selected from the group consisting of N, O, Cl, Br, I, Si, P, and S. The following group is a non-limiting example of a substituted fluoroalkyl: -CFHOH .

The term “alkylthio” when used without the “substituted” modifier refers to the group –SR, in which R is an alkyl, as that term is defined above. Non-limiting examples of alkylthio groups include: –SCH₃, –SCH₂CH₃, –SCH₂CH₂CH₃, –SCH(CH₃)₂, –SCH(CH₂)₂, –S–cyclopentyl, and –S–cyclohexyl. The term “substituted alkylthio” refers to the group
 5 –SR, in which R is a substituted alkyl, as that term is defined above. For example, –SCH₂CF₃ is a substituted alkylthio group.

Similarly, the terms “alkenylthio”, “alkynylthio”, “arylthio”, “aralkylthio”, “heteroarylthio”, “heteroaralkylthio”, and “acylthio”, when used without the “substituted” modifier, refers to groups, defined as –SR, in which R is alkenyl, alkynyl, aryl, aralkyl,
 10 heteroaryl, heteroaralkyl and acyl, respectively, as those terms are defined above. When any of the terms alkenylthio, alkynylthio, arylthio, aralkylthio, heteroarylthio, heteroaralkylthio, and acylthio is modified by “substituted,” it refers to the group –SR, in which R is substituted alkenyl, alkynyl, aryl, aralkyl, heteroaryl, heteroaralkyl and acyl, respectively.

The term “thioacyl” when used without the “substituted” modifier refers to a
 15 monovalent group with a carbon atom of a thiocarbonyl group as the point of attachment, further having a linear or branched, cyclo, cyclic or acyclic structure, further having no additional atoms that are not carbon or hydrogen, beyond the sulfur atom of the carbonyl group. The groups, –CHS, –C(S)CH₃, –C(S)CH₂CH₃, –C(S)CH₂CH₂CH₃, –C(S)CH(CH₃)₂, –C(S)CH(CH₂)₂, –C(S)C₆H₅, –C(S)C₆H₄CH₃, –C(S)C₆H₄CH₂CH₃, –C(S)C₆H₃(CH₃)₂, and
 20 –C(S)CH₂C₆H₅, are non-limiting examples of thioacyl groups. The term “thioacyl” therefore encompasses, but is not limited to, groups sometimes referred to as “alkyl thiocarbonyl” and “aryl thiocarbonyl” groups. The term “substituted thioacyl” refers to a radical with a carbon atom as the point of attachment, the carbon atom being part of a thiocarbonyl group, further having a linear or branched, cyclo, cyclic or acyclic structure, further having at least one
 25 atom, in addition to the sulfur atom of the carbonyl group, independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. The groups, –C(S)CH₂CF₃, –C(S)O₂H, –C(S)OCH₃, –C(S)OCH₂CH₃, –C(S)OCH₂CH₂CH₃, –C(S)OC₆H₅, –C(S)OCH(CH₃)₂, –C(S)OCH(CH₂)₂, –C(S)NH₂, and –C(S)NHCH₃, are non-limiting examples of substituted thioacyl groups. The term “substituted thioacyl” encompasses, but is not limited to,
 30 “heteroaryl thiocarbonyl” groups.

The term “alkylsulfonyl” when used without the “substituted” modifier refers to the group –S(O)₂R, in which R is an alkyl, as that term is defined above. Non-limiting examples of alkylsulfonyl groups include: –S(O)₂CH₃, –S(O)₂CH₂CH₃, –S(O)₂CH₂CH₂CH₃,

$-\text{S}(\text{O})_2\text{CH}(\text{CH}_3)_2$, $-\text{S}(\text{O})_2\text{CH}(\text{CH}_2)_2$, $-\text{S}(\text{O})_2$ -cyclopentyl, and $-\text{S}(\text{O})_2$ -cyclohexyl. The term “substituted alkylsulfonyl” refers to the group $-\text{S}(\text{O})_2\text{R}$, in which R is a substituted alkyl, as that term is defined above. For example, $-\text{S}(\text{O})_2\text{CH}_2\text{CF}_3$ is a substituted alkylsulfonyl group.

Similarly, the terms “alkenylsulfonyl”, “alkynylsulfonyl”, “arylsulfonyl”,
 5 “aralkylsulfonyl”, “heteroarylsulfonyl”, and “heteroaralkylsulfonyl” when used without the “substituted” modifier, refers to groups, defined as $-\text{S}(\text{O})_2\text{R}$, in which R is alkenyl, alkynyl, aryl, aralkyl, heteroaryl, and heteroaralkyl, respectively, as those terms are defined above. When any of the terms alkenylsulfonyl, alkynylsulfonyl, arylsulfonyl, aralkylsulfonyl, heteroarylsulfonyl, and heteroaralkylsulfonyl is modified by “substituted,” it refers to the
 10 group $-\text{S}(\text{O})_2\text{R}$, in which R is substituted alkenyl, alkynyl, aryl, aralkyl, heteroaryl and heteroaralkyl, respectively.

The term “alkylammonium” when used without the “substituted” modifier refers to a group, defined as $-\text{NH}_2\text{R}^+$, $-\text{NHR}'^+$, or $-\text{NRR}'\text{R}''^+$, in which R, R' and R'' are the same or different alkyl groups, or any combination of two of R, R' and R'' can be taken together to
 15 represent an alkanediyl. Non-limiting examples of alkylammonium cation groups include: $-\text{NH}_2(\text{CH}_3)^+$, $-\text{NH}_2(\text{CH}_2\text{CH}_3)^+$, $-\text{NH}_2(\text{CH}_2\text{CH}_2\text{CH}_3)^+$, $-\text{NH}(\text{CH}_3)_2^+$, $-\text{NH}(\text{CH}_2\text{CH}_3)_2^+$, $-\text{NH}(\text{CH}_2\text{CH}_2\text{CH}_3)_2^+$, $-\text{N}(\text{CH}_3)_3^+$, $-\text{N}(\text{CH}_3)(\text{CH}_2\text{CH}_3)_2^+$, $-\text{N}(\text{CH}_3)_2(\text{CH}_2\text{CH}_3)^+$, $-\text{NH}_2\text{C}(\text{CH}_3)_3^+$, $-\text{NH}(\text{cyclopentyl})_2^+$, and $-\text{NH}_2(\text{cyclohexyl})^+$. The term “substituted alkylammonium” refers $-\text{NH}_2\text{R}^+$, $-\text{NHR}'^+$, or $-\text{NRR}'\text{R}''^+$, in which at least one of R, R' and
 20 R'' is a substituted alkyl or two of R, R' and R'' can be taken together to represent a substituted alkanediyl. When more than one of R, R' and R'' is a substituted alkyl, they can be the same or different. Any of R, R' and R'' that are not either substituted alkyl or substituted alkanediyl, can be either alkyl, either the same or different, or can be taken together to represent a alkanediyl with two or more carbon atoms, at least two of which are
 25 attached to the nitrogen atom shown in the formula.

The term “alkylsulfonium” when used without the “substituted” modifier refers to the group $-\text{SRR}'^+$, in which R and R' can be the same or different alkyl groups, or R and R' can be taken together to represent an alkanediyl. Non-limiting examples of alkylsulfonium groups include: $-\text{SH}(\text{CH}_3)^+$, $-\text{SH}(\text{CH}_2\text{CH}_3)^+$, $-\text{SH}(\text{CH}_2\text{CH}_2\text{CH}_3)^+$, $-\text{S}(\text{CH}_3)_2^+$,
 30 $-\text{S}(\text{CH}_2\text{CH}_3)_2^+$, $-\text{S}(\text{CH}_2\text{CH}_2\text{CH}_3)_2^+$, $-\text{SH}(\text{cyclopentyl})^+$, and $-\text{SH}(\text{cyclohexyl})^+$. The term “substituted alkylsulfonium” refers to the group $-\text{SRR}'^+$, in which R and R' can be the same or different substituted alkyl groups, one of R or R' is an alkyl and the other is a substituted

alkyl, or R and R' can be taken together to represent a substituted alkanediyl. For example, $-\text{SH}(\text{CH}_2\text{CF}_3)^+$ is a substituted alkylsulfonium group.

The term "alkylsilyl" when used without the "substituted" modifier refers to a monovalent group, defined as $-\text{SiH}_2\text{R}$, $-\text{SiHRR}'$, or $-\text{SiRR}'\text{R}''$, in which R, R' and R'' can be the same or different alkyl groups, or any combination of two of R, R' and R'' can be taken together to represent an alkanediyl. The groups, $-\text{SiH}_2\text{CH}_3$, $-\text{SiH}(\text{CH}_3)_2$, $-\text{Si}(\text{CH}_3)_3$ and $-\text{Si}(\text{CH}_3)_2\text{C}(\text{CH}_3)_3$, are non-limiting examples of unsubstituted alkylsilyl groups. The term "substituted alkylsilyl" refers $-\text{SiH}_2\text{R}$, $-\text{SiHRR}'$, or $-\text{SiRR}'\text{R}''$, in which at least one of R, R' and R'' is a substituted alkyl or two of R, R' and R'' can be taken together to represent a substituted alkanediyl. When more than one of R, R' and R'' is a substituted alkyl, they can be the same or different. Any of R, R' and R'' that are not either substituted alkyl or substituted alkanediyl, can be either alkyl, either the same or different, or can be taken together to represent a alkanediyl with two or more saturated carbon atoms, at least two of which are attached to the silicon atom.

In addition, atoms making up the compounds of the present disclosure are intended to include all isotopic forms of such atoms. Isotopes, as used herein, include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium, and isotopes of carbon include ^{13}C and ^{14}C . Similarly, it is contemplated that one or more carbon atom(s) of a compound of the present disclosure may be replaced by a silicon atom(s). Furthermore, it is contemplated that one or more oxygen atom(s) of a compound of the present disclosure may be replaced by a sulfur or selenium atom(s).

A compound having a formula that is represented with a dashed bond is intended to include the formulae optionally having zero, one or more double bonds. Thus, for example,


the structure  includes the structures , , ,  and .

As will be understood by a person of skill in the art, no one such ring atom forms part of more than one double bond.


Any undefined valency on an atom of a structure shown in this application implicitly represents a hydrogen atom bonded to the atom.

A ring structure shown with an unconnected "R" group, indicates that any implicit hydrogen atom on that ring can be replaced with that R group. In the case of a divalent R

group (*e.g.*, oxo, imino, thio, alkylidene, etc.), any pair of implicit hydrogen atoms attached to one atom of that ring can be replaced by that R group. This concept is as exemplified below:



represents



As used herein, a “chiral auxiliary” refers to a removable chiral group that is capable of influencing the stereoselectivity of a reaction. Persons of skill in the art are familiar with such compounds, and many are commercially available.

The use of the word “a” or “an,” when used in conjunction with the term “comprising” in the claims and/or the specification may mean “one,” but it is also consistent with the meaning of “one or more,” “at least one,” and “one or more than one.”

Throughout this application, the term “about” is used to indicate that a value includes the inherent variation of error for the device, the method being employed to determine the value, or the variation that exists among the study subjects.

The terms “comprise,” “have” and “include” are open-ended linking verbs. Any forms or tenses of one or more of these verbs, such as “comprises,” “comprising,” “has,” “having,” “includes” and “including,” are also open-ended. For example, any method that “comprises,” “has” or “includes” one or more steps is not limited to possessing only those one or more steps and also covers other unlisted steps.

The term “effective,” as that term is used in the specification and/or claims, means adequate to accomplish a desired, expected, or intended result.

The term “hydrate” when used as a modifier to a compound means that the compound has less than one (*e.g.*, hemihydrate), one (*e.g.*, monohydrate), or more than one (*e.g.*, dihydrate) water molecules associated with each compound molecule, such as in solid forms of the compound.

As used herein, the term “IC₅₀” refers to an inhibitory dose which is 50% of the maximum response obtained.

An "isomer" of a first compound is a separate compound in which each molecule contains the same constituent atoms as the first compound, but where the configuration of those atoms in three dimensions differs.

As used herein, the term "patient" or "subject" refers to a living mammalian organism, such as a human, monkey, cow, sheep, goat, dog, cat, mouse, rat, guinea pig, or transgenic species thereof. In certain embodiments, the patient or subject is a primate. Non-limiting examples of human subjects are adults, juveniles, infants and fetuses.

"Pharmaceutically acceptable" means that which is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable and includes that which is acceptable for veterinary use as well as human pharmaceutical use.

"Pharmaceutically acceptable salts" means salts of compounds of the present disclosure which are pharmaceutically acceptable, as defined above, and which possess the desired pharmacological activity. Such salts include acid addition salts formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or with organic acids such as 1,2-ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, 2-naphthalenesulfonic acid, 3-phenylpropionic acid, 4,4'-methylenebis(3-hydroxy-2-ene-1-carboxylic acid), 4-methylbicyclo[2.2.2]oct-2-ene-1-carboxylic acid, acetic acid, aliphatic mono- and dicarboxylic acids, aliphatic sulfuric acids, aromatic sulfuric acids, benzenesulfonic acid, benzoic acid, camphorsulfonic acid, carbonic acid, cinnamic acid, citric acid, cyclopentanepropionic acid, ethanesulfonic acid, fumaric acid, glucoheptonic acid, gluconic acid, glutamic acid, glycolic acid, heptanoic acid, hexanoic acid, hydroxynaphthoic acid, lactic acid, laurylsulfuric acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, muconic acid, *o*-(4-hydroxybenzoyl)benzoic acid, oxalic acid, *p*-chlorobenzenesulfonic acid, phenyl-substituted alkanolic acids, propionic acid, *p*-toluenesulfonic acid, pyruvic acid, salicylic acid, stearic acid, succinic acid, tartaric acid, tertiarybutylacetic acid, trimethylacetic acid, and the like. Pharmaceutically acceptable salts also include base addition salts which may be formed when acidic protons present are capable of reacting with inorganic or organic bases. Acceptable inorganic bases include sodium hydroxide, sodium carbonate, potassium hydroxide, aluminum hydroxide and calcium hydroxide. Acceptable organic bases include ethanolamine, diethanolamine, triethanolamine, tromethamine, *N*-methylglucamine and the like. It should be recognized that the particular anion or cation forming a part of any salt of this invention is not critical, so long as the salt, as

a whole, is pharmacologically acceptable. Additional examples of pharmaceutically acceptable salts and their methods of preparation and use are presented in *Handbook of Pharmaceutical Salts: Properties, and Use* (2002).

As used herein, “predominantly one enantiomer” means that a compound contains at least about 85% of one enantiomer, or more preferably at least about 90% of one enantiomer, or even more preferably at least about 95% of one enantiomer, or most preferably at least about 99% of one enantiomer. Similarly, the phrase “substantially free from other optical isomers” means that the composition contains at most about 15% of another enantiomer or diastereomer, more preferably at most about 10% of another enantiomer or diastereomer, even more preferably at most about 5% of another enantiomer or diastereomer, and most preferably at most about 1% of another enantiomer or diastereomer.

“Prevention” or “preventing” includes: (1) inhibiting the onset of a disease in a subject or patient which may be at risk and/or predisposed to the disease but does not yet experience or display any or all of the pathology or symptomatology of the disease, and/or (2) slowing the onset of the pathology or symptomatology of a disease in a subject or patient which may be at risk and/or predisposed to the disease but does not yet experience or display any or all of the pathology or symptomatology of the disease.

“Prodrug” means a compound that is convertible *in vivo* metabolically into an inhibitor according to the present disclosure. The prodrug itself may or may not also have activity with respect to a given target protein. For example, a compound comprising a hydroxy group may be administered as an ester that is converted by hydrolysis *in vivo* to the hydroxy compound. Suitable esters that may be converted *in vivo* into hydroxy compounds include acetates, citrates, lactates, phosphates, tartrates, malonates, oxalates, salicylates, propionates, succinates, fumarates, maleates, methylene-bis- β -hydroxynaphthoate, gentisates, isethionates, di-*p*-toluoyltartrates, methanesulfonates, ethanesulfonates, benzenesulfonates, *p*-toluenesulfonates, cyclohexylsulfamates, quinate, esters of amino acids, and the like. Similarly, a compound comprising an amine group may be administered as an amide that is converted by hydrolysis *in vivo* to the amine compound.

The term “saturated” when referring to an atom means that the atom is connected to other atoms only by means of single bonds.

A “stereoisomer” or “optical isomer” is an isomer of a given compound in which the same atoms are bonded to the same other atoms, but where the configuration of those atoms in three dimensions differs. “Enantiomers” are stereoisomers of a given compound that are

mirror images of each other, like left and right hands. "Diastereomers" are stereoisomers of a given compound that are not enantiomers.

The invention contemplates that for any stereocenter or axis of chirality for which stereochemistry has not been defined, that stereocenter or axis of chirality can be present in its *R* form, *S* form, or as a mixture of the *R* and *S* forms, including racemic and non-racemic mixtures.

"Substituent convertible to hydrogen *in vivo*" means any group that is convertible to a hydrogen atom by enzymological or chemical means including, but not limited to, hydrolysis and hydrogenolysis. Examples include acyl groups, groups having an oxycarbonyl group, amino acid residues, peptide residues, *o*-nitrophenylsulfenyl, trimethylsilyl, tetrahydropyranyl, diphenylphosphinyl, hydroxy or alkoxy substituents on imino groups, and the like. Examples of acyl groups include formyl, acetyl, trifluoroacetyl, and the like. Examples of groups having an oxycarbonyl group include ethoxycarbonyl, *tert*-butoxycarbonyl ($-\text{C}(\text{O})\text{OC}(\text{CH}_3)_3$), benzyloxycarbonyl, *p*-methoxybenzyloxycarbonyl, vinyloxycarbonyl, β -(*p*-toluenesulfonyl)ethoxycarbonyl, and the like. Suitable amino acid residues include, but are not limited to, residues of Gly (glycine), Ala (alanine), Arg (arginine), Asn (asparagine), Asp (aspartic acid), Cys (cysteine), Glu (glutamic acid), His (histidine), Ile (isoleucine), Leu (leucine), Lys (lysine), Met (methionine), Phe (phenylalanine), Pro (proline), Ser (serine), Thr (threonine), Trp (tryptophan), Tyr (tyrosine), Val (valine), Nva (norvaline), Hse (homoserine), 4-Hyp (4-hydroxyproline), 5-Hyl (5-hydroxylysine), Orn (ornithine) and β -Ala. Examples of suitable amino acid residues also include amino acid residues that are protected with a protecting group. Examples of suitable protecting groups include those typically employed in peptide synthesis, including acyl groups (such as formyl and acetyl), arylmethyloxycarbonyl groups (such as benzyloxycarbonyl and *p*-nitrobenzyloxycarbonyl), *tert*-butoxycarbonyl groups ($-\text{C}(\text{O})\text{OC}(\text{CH}_3)_3$), and the like. Suitable peptide residues include peptide residues comprising two to five, and optionally amino acid residues. The residues of these amino acids or peptides can be present in stereochemical configurations of the D-form, the L-form or mixtures thereof. In addition, the amino acid or peptide residue may have an asymmetric carbon atom. Examples of suitable amino acid residues having an asymmetric carbon atom include residues of Ala, Leu, Phe, Trp, Nva, Val, Met, Ser, Lys, Thr and Tyr. Peptide residues having an asymmetric carbon atom include peptide residues having one or more constituent amino acid residues having an asymmetric carbon atom. Examples of suitable amino acid protecting groups include those typically employed in peptide

synthesis, including acyl groups (such as formyl and acetyl), arylmethyloxycarbonyl groups (such as benzyloxycarbonyl and *p*-nitrobenzyloxycarbonyl), *tert*-butoxycarbonyl groups ($-\text{C}(\text{O})\text{OC}(\text{CH}_3)_3$), and the like. Other examples of substituents “convertible to hydrogen *in vivo*” include reductively eliminable hydrogenolyzable groups. Examples of suitable reductively eliminable hydrogenolyzable groups include, but are not limited to, arylsulfonyl groups (such as *o*-toluenesulfonyl); methyl groups substituted with phenyl or benzyloxy (such as benzyl, trityl and benzyloxymethyl); arylmethoxycarbonyl groups (such as benzyloxycarbonyl and *o*-methoxy-benzyloxycarbonyl); and haloethoxycarbonyl groups (such as β,β,β -trichloroethoxycarbonyl and β -iodoethoxycarbonyl).

“Therapeutically effective amount” or “pharmaceutically effective amount” means that amount which, when administered to a subject or patient for treating a disease, is sufficient to effect such treatment for the disease.

“Treatment” or “treating” includes (1) inhibiting a disease in a subject or patient experiencing or displaying the pathology or symptomatology of the disease (*e.g.*, arresting further development of the pathology and/or symptomatology), (2) ameliorating a disease in a subject or patient that is experiencing or displaying the pathology or symptomatology of the disease (*e.g.*, reversing the pathology and/or symptomatology), and/or (3) effecting any measurable decrease in a disease in a subject or patient that is experiencing or displaying the pathology or symptomatology of the disease.

As used herein, the term “water soluble” means that the compound dissolves in water at least to the extent of 0.010 mole/liter or is classified as soluble according to literature precedence.

Other abbreviations used herein are as follows: DMSO, dimethyl sulfoxide; NO, nitric oxide; iNOS, inducible nitric oxide synthase; COX-2, cyclooxygenase-2; NGF, nerve growth factor; IBMX, isobutylmethylxanthine; FBS, fetal bovine serum; GPDH, glycerol 3-phosphate dehydrogenase; RXR, retinoid X receptor; TGF- β , transforming growth factor- β ; IFN γ or IFN- γ , interferon- γ ; LPS, bacterial endotoxic lipopolysaccharide; TNF α or TNF- α , tumor necrosis factor- α ; IL-1 β , interleukin-1 β ; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; MTT, 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide; TCA, trichloroacetic acid; HO-1, inducible heme oxygenase.

The above definitions supersede any conflicting definition in any of the reference that is incorporated by reference herein. The fact that certain terms are defined, however, should not be considered as indicative that any term that is undefined is indefinite. Rather, all terms

used are believed to describe the invention in terms such that one of ordinary skill can appreciate the scope and practice the present disclosure.

II. Synthetic Methods

Compounds of the present disclosure may be made using the methods outlined in the Examples section (Example 2 and 3). These methods can be further modified and optimized using the principles and techniques of organic chemistry as applied by a person skilled in the art. Such principles and techniques are taught, for example, in March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure (2007), which is incorporated by reference herein.

III. Biological Activity of Oleanolic Acid Derivatives

Biological activity results, both *in vivo* and *in vitro* are provided throughout the present disclosure. These include: inhibition of NO Production, suppression of COX-2 induction, inhibition of IL-6 induced STAT3 phosphorylation, suppression of IL-6 induced STAT3 phosphorylation, inhibition of TNF α -induced I κ B α degradation, inhibition of NF κ B activation, induction of HO-1, Nrf2 induction of HO-1, TrxR1 and γ -GCS, induction of TrxR1, induction of γ -GCS, induction of ferritin heavy chain, induction of TrxR1, induction of γ -GCS, induction of ferritin heavy chain, and various *in vivo* toxicity studies. See figures and figure descriptions. Suppression of NO production and induction of Nrf2 induction results can be respectively summarized as shown Tables 1a and 1b, below. Further results, including toxicity studies, are provided in the Examples section.

Table 1a. Suppression of IFN γ -Induced NO Production.

Compound ID(s)	MW	RAW264.7 (20 ng/ml IFN γ)		
		NO IC ₅₀	WST-1 IC ₅₀	iNOS suppr. WB
63101 / 402-02 / dh402	507.70	~12 nM	200 nM	>90%
63102 / 404-02 / dh404	574.72	~45 nM	> 200 nM	
63250 / 402-46	509.70	> 200 nM	> 200 nM	
63197 / 402-48	512.72	> 200 nM	> 200 nM	
63195 / 402-49	509.72	> 200 nM	> 200 nM	
63196 / 402-51 / dh401	493.68	~75 nM	> 200 nM	

Compound ID(s)	MW	RAW264.7 (20 ng/ml IFN γ)		
		NO IC ₅₀	WST-1 IC ₅₀	iNOS suppr. WB
63252 / 402-57	474.68	~5 nM	> 200 nM	
63205 / 402-59	492.69	~25 nM	> 200 nM	
63206 / 402-64	477.68	~10 nM	> 200 nM	
63207 / 402-66	509.72	~50 nM	> 200 nM	
63219 / 402-78	509.72	~150 nM	> 200 nM	
63229	517.71	> 200nM	> 200nM	
63230	531.74	~50 nM	> 200 nM	
63227	576.73	> 200nM	> 200nM	
63219	509.72	~150 nM	> 200 nM	
63223	576.73	> 200nM	> 200nM	
63237	572.70	~80 nM	> 200 nM	
63268	576.75	> 200 nM	> 200 nM	
63274	547.81	~ 70 nM	> 200 nM	
63289	525.73	> 200 nM	> 200 nM	
63295	522.73	~ 20 nM	> 200 nM	
63296	522.73	~ 200 nM	> 200 nM	
63308	491.70	~ 25 nM	> 200 nM	
63323	583.80	~ 40 nM	See FIG. 33	
63325	520.75	~ 50 nM	See FIG. 32	
63326	552.79	~ 50 nM	See FIG. 34	

Table 1b. Induction of HO-1, TrxR1 and γ -GCS in Human Melanoma Cells.

Compound Code	Nrf2 target gene induction in MDA-MB-435 cells					
	400 nM*			250 nM**		
	HO-1	TrxR1	γ -GCS	HO-1	NQO1	γ -GCS
63101 (dh402)	4	47	53	3	2	5
63102 (dh404)				3	2	4.5
63196 (dh401)	1	48	26			
63252				5	3	
63205				1.7	1.7	3.5
63206				2.5	1.8	4.5
63207				1.2	1.6	3.1
63237				3	2.3	6.4

Blank entry: Not determined.

* Data expressed as a percent of induction observed for **402** (see below for structure).

** Data expressed as fold induction above DMSO control.


5 In certain embodiments, the compounds of the present disclosure are capable of crossing the blood brain barrier and achieving therapeutically effective concentrations in the brain. They may therefore be used to treat neurodegenerative diseases, brain cancer and other inflammatory conditions affecting the central nervous system. For example, 404-02 has been shown to cross the blood-brain barrier and achieve high concentrations in the central nervous system tissue following oral dosing. Like the other compounds of the present disclosure, it promotes the resolution of innate and adaptive immune-mediated inflammation by restoring redox homeostasis in inflamed tissues. It is a potent inducer of the antioxidant transcription factor Nrf2 and inhibitor of the pro-oxidant/pro-inflammatory transcription factors NF- κ B and the STATs. These biological pathways are implicated in a wide variety of diseases, including autoimmune conditions and several neurodegenerative diseases.

10

15

IV. Improved Rodent Toxicology

In certain embodiments, the invention provides compounds possessing low toxicity in rodents. In some cases, toxicity in rodents has been observed in preclinical studies with some analogues containing a carbon-carbon double bond in the C-ring, including 402 and 401. Compounds having a saturated C-ring, in contrast, have consistently shown low toxicity in rodents. Predictably low rodent toxicity provides an advantage since high rodent toxicity can be a significant complication in conducting preclinical studies required for development and registration of therapeutic compounds for use in humans or non-human animals. Illustrations of this effect are provided below.



For example, an initial study (Example 6) was performed in Sprague Dawley rats using both 402 and 402-02 and showed that 402-02 was less toxic. In a further study (Example 7), six compounds (401, 402, 404, 401-2, 402-2, and 404-2) were assessed for toxicity in mice in a 14-day study. At higher doses (above 10 mg/kg/day) both 401 and 402 caused at least 50% mortality, while 404 was non-toxic. In contrast, no mortality was observed in the 402-2 and 404-2 groups and only the highest dose of 401-02 caused any lethality (Table 5). Body weight measurements (FIGS. 29-31) were consistent with the mortality observations. Notably, the two highest doses of 401 and 402 were lethal within 4 days, in contrast to the effects of 401-2 and 402-2.

V. Diseases Associated with Inflammation and/or Oxidative Stress

Inflammation is a biological process that provides resistance to infectious or parasitic organisms and the repair of damaged tissue. Inflammation is commonly characterized by localized vasodilation, redness, swelling, and pain, the recruitment of leukocytes to the site of infection or injury, production of inflammatory cytokines such as TNF- α and IL-1, and production of reactive oxygen or nitrogen species such as hydrogen peroxide, superoxide and peroxynitrite. In later stages of inflammation, tissue remodeling, angiogenesis, and scar

formation (fibrosis) may occur as part of the wound healing process. Under normal circumstances, the inflammatory response is regulated and temporary and is resolved in an orchestrated fashion once the infection or injury has been dealt with adequately. However, acute inflammation can become excessive and life-threatening if regulatory mechanisms fail.

5 Alternatively, inflammation can become chronic and cause cumulative tissue damage or systemic complications.

Many serious and intractable human diseases involve dysregulation of inflammatory processes, including diseases such as cancer, atherosclerosis, and diabetes, which were not traditionally viewed as inflammatory conditions. In the case of cancer, the inflammatory
10 processes are associated with tumor formation, progression, metastasis, and resistance to therapy. Atherosclerosis, long viewed as a disorder of lipid metabolism, is now understood to be primarily an inflammatory condition, with activated macrophages playing an important role in the formation and eventual rupture of atherosclerotic plaques. Activation of inflammatory signaling pathways has also been shown to play a role in the development of
15 insulin resistance, as well as in the peripheral tissue damage associated with diabetic hyperglycemia. Excessive production of reactive oxygen species and reactive nitrogen species such as superoxide, hydrogen peroxide, nitric oxide, and peroxynitrite is a hallmark of inflammatory conditions. Evidence of dysregulated peroxynitrite production has been reported in a wide variety of diseases (Szabo *et al.*, 2007; Schulz *et al.*, 2008; Forstermann,
20 2006; Pall, 2007).

Autoimmune diseases such as rheumatoid arthritis, lupus, psoriasis, and multiple sclerosis involve inappropriate and chronic activation of inflammatory processes in affected tissues, arising from dysfunction of self vs. non-self recognition and response mechanisms in the immune system. In neurodegenerative diseases such as Alzheimer's and Parkinson's
25 diseases, neural damage is correlated with activation of microglia and elevated levels of pro-inflammatory proteins such as inducible nitric oxide synthase (iNOS). Chronic organ failure such as renal failure, heart failure, and chronic obstructive pulmonary disease is closely associated with the presence of chronic oxidative stress and inflammation, leading to the development of fibrosis and eventual loss of organ function.

30 Many other disorders involve oxidative stress and inflammation in affected tissues, including inflammatory bowel disease; inflammatory skin diseases; mucositis related to radiation therapy and chemotherapy; eye diseases such as uveitis, glaucoma, macular degeneration, and various forms of retinopathy; transplant failure and rejection; ischemia-

reperfusion injury; chronic pain; degenerative conditions of the bones and joints including osteoarthritis and osteoporosis; asthma and cystic fibrosis; seizure disorders; and neuropsychiatric conditions including schizophrenia, depression, bipolar disorder, post-traumatic stress disorder, attention deficit disorders, autism-spectrum disorders, and eating disorders such as anorexia nervosa. Dysregulation of inflammatory signaling pathways is believed to be a major factor in the pathology of muscle wasting diseases including muscular dystrophy and various forms of cachexia.

A variety of life-threatening acute disorders also involve dysregulated inflammatory signaling, including acute organ failure involving the pancreas, kidneys, liver, or lungs, myocardial infarction or acute coronary syndrome, stroke, septic shock, trauma, severe burns, and anaphylaxis.

Many complications of infectious diseases also involve dysregulation of inflammatory responses. Although an inflammatory response can kill invading pathogens, an excessive inflammatory response can also be quite destructive and in some cases can be a primary source of damage in infected tissues. Furthermore, an excessive inflammatory response can also lead to systemic complications due to overproduction of inflammatory cytokines such as TNF- α and IL-1. This is believed to be a factor in mortality arising from severe influenza, severe acute respiratory syndrome, and sepsis.

The aberrant or excessive expression of either iNOS or cyclooxygenase-2 (COX-2) has been implicated in the pathogenesis of many disease processes. For example, it is clear that NO is a potent mutagen (Tamir and Tannebaum, 1996), and that nitric oxide can also activate COX-2 (Salvemini *et al.*, 1994). Furthermore, there is a marked increase in iNOS in rat colon tumors induced by the carcinogen, azoxymethane (Takahashi *et al.*, 1997). A series of synthetic triterpenoid analogs of oleanolic acid have been shown to be powerful inhibitors of cellular inflammatory processes, such as the induction by IFN- γ of inducible nitric oxide synthase (iNOS) and of COX-2 in mouse macrophages. See Honda *et al.* (2000a); Honda *et al.* (2000b), and Honda *et al.* (2002), which are all incorporated herein by reference.

In one aspect, compounds of the invention are characterized by their ability to inhibit the production of nitric oxide in macrophage-derived RAW 264.7 cells induced by exposure to γ -interferon. They are further characterized by their ability to induce the expression of antioxidant proteins such as NQO1 and reduce the expression of pro-inflammatory proteins such as COX-2 and inducible nitric oxide synthase (iNOS). These properties are relevant to the treatment of a wide array of diseases involving oxidative stress and dysregulation of

inflammatory processes including cancer, mucositis resulting from radiation therapy or chemotherapy, autoimmune diseases, cardiovascular diseases including atherosclerosis, ischemia-reperfusion injury, acute and chronic organ failure including renal failure and heart failure, respiratory diseases, diabetes and complications of diabetes, severe allergies, transplant rejection, graft-versus-host disease, neurodegenerative diseases, diseases of the eye and retina, acute and chronic pain, degenerative bone diseases including osteoarthritis and osteoporosis, inflammatory bowel diseases, dermatitis and other skin diseases, sepsis, burns, seizure disorders, and neuropsychiatric disorders.

Without being bound by theory, the activation of the antioxidant/anti-inflammatory Keap1/Nrf2/ARE pathway is believed to be implicated in both the anti-inflammatory and anti-carcinogenic properties of the present oleanolic acid derivatives.

In another aspect, compounds of the invention may be used for treating a subject having a condition caused by elevated levels of oxidative stress in one or more tissues. Oxidative stress results from abnormally high or prolonged levels of reactive oxygen species such as superoxide, hydrogen peroxide, nitric oxide, and peroxynitrite (formed by the reaction of nitric oxide and superoxide). The oxidative stress may be accompanied by either acute or chronic inflammation. The oxidative stress may be caused by mitochondrial dysfunction, by activation of immune cells such as macrophages and neutrophils, by acute exposure to an external agent such as ionizing radiation or a cytotoxic chemotherapy agent (*e.g.*, doxorubicin), by trauma or other acute tissue injury, by ischemia/reperfusion, by poor circulation or anemia, by localized or systemic hypoxia or hyperoxia, by elevated levels of inflammatory cytokines and other inflammation-related proteins, and/or by other abnormal physiological states such as hyperglycemia or hypoglycemia.

In animal models of many such conditions, stimulating expression of inducible heme oxygenase (HO-1), a target gene of the Nrf2 pathway, has been shown to have a significant therapeutic effect including models of myocardial infarction, renal failure, transplant failure and rejection, stroke, cardiovascular disease, and autoimmune disease (*e.g.*, Sacerdoti *et al.*, 2005; Abraham & Kappas, 2005; Bach, 2006; Araujo *et al.*, 2003; Liu *et al.*, 2006; Ishikawa *et al.*, 2001; Kruger *et al.*, 2006; Satoh *et al.*, 2006; Zhou *et al.*, 2005; Morse and Choi, 2005; Morse and Choi, 2002). This enzyme breaks free heme down into iron, carbon monoxide (CO), and biliverdin (which is subsequently converted to the potent antioxidant molecule, bilirubin).

In another aspect, compounds of this invention may be used in preventing or treating tissue damage or organ failure, acute and chronic, resulting from oxidative stress exacerbated by inflammation. Examples of diseases that fall in this category include: heart failure, liver failure, transplant failure and rejection, renal failure, pancreatitis, fibrotic lung diseases (cystic fibrosis and COPD, among others), diabetes (including complications), atherosclerosis, ischemia-reperfusion injury, glaucoma, stroke, autoimmune disease, autism, macular degeneration, and muscular dystrophy. For example, in the case of autism, studies suggest that increased oxidative stress in the central nervous system may contribute to the development of the disease (Chauhan and Chauhan, 2006).

Evidence also links oxidative stress and inflammation to the development and pathology of many other disorders of the central nervous system, including psychiatric disorders such as psychosis, major depression, and bipolar disorder; seizure disorders such as epilepsy; pain and sensory syndromes such as migraine, neuropathic pain or tinnitus; and behavioral syndromes such as the attention deficit disorders. See, *e.g.*, Dickerson *et al.*, 2007; Hanson *et al.*, 2005; Kendall-Tackett, 2007; Lencz *et al.*, 2007; Dudhgaonkar *et al.*, 2006; Lee *et al.*, 2007; Morris *et al.*, 2002; Ruster *et al.*, 2005; McIver *et al.*, 2005; Sarchielli *et al.*, 2006; Kawakami *et al.*, 2006; Ross *et al.*, 2003, which are all incorporated by reference herein. For example, elevated levels of inflammatory cytokines, including TNF, interferon- γ , and IL-6, are associated with major mental illness (Dickerson *et al.*, 2007). Microglial activation has also been linked to major mental illness. Therefore, downregulating inflammatory cytokines and inhibiting excessive activation of microglia could be beneficial in patients with schizophrenia, major depression, bipolar disorder, autism-spectrum disorders, and other neuropsychiatric disorders.

Accordingly, in pathologies involving oxidative stress alone or oxidative stress exacerbated by inflammation, treatment may comprise administering to a subject a therapeutically effective amount of a compound of this invention, such as those described above or throughout this specification. Treatment may be administered preventively, in advance of a predictable state of oxidative stress (*e.g.*, organ transplantation or the administration of radiation therapy to a cancer patient), or it may be administered therapeutically in settings involving established oxidative stress and inflammation.

The compounds of the invention may be generally applied to the treatment of inflammatory conditions, such as sepsis, dermatitis, autoimmune disease and osteoarthritis.

In one aspect, the compounds of this invention may be used to treat inflammatory pain and/or neuropathic pain, for example, by inducing Nrf2 and/or inhibiting NF- κ B.

In one aspect, the compounds of the invention may be used to function as antioxidant inflammation modulators (AIMs) having potent anti-inflammatory properties that mimic the biological activity of cyclopentenone prostaglandins (cyPGs). In one embodiment, the compounds of the invention may be used to control the production of pro-inflammatory cytokines by selectively targeting regulatory cysteine residues (RCRs) on proteins that regulate the transcriptional activity of redox-sensitive transcription factors. Activation of RCRs by cyPGs or AIMs has been shown to initiate a pro-resolution program in which the activity of the antioxidant and cytoprotective transcription factor Nrf2 is potently induced, and the activities of the pro-oxidant and pro-inflammatory transcription factors NF- κ B and the STATs are suppressed. This increases the production of antioxidant and reductive molecules (*e.g.*, NQO1, HO-1, SOD1, and/or γ -GCS) and/or decreases oxidative stress and the production of pro-oxidant and pro-inflammatory molecules (*e.g.*, iNOS, COX-2, and/or TNF- α).

In some embodiments, the compounds of the invention may be used in the treatment and prevention of diseases such as cancer, inflammation, Alzheimer's disease, Parkinson's disease, multiple sclerosis, autism, amyotrophic lateral sclerosis, autoimmune diseases such as rheumatoid arthritis, lupus, and MS, inflammatory bowel disease, all other diseases whose pathogenesis is believed to involve excessive production of either nitric oxide or prostaglandins, and pathologies involving oxidative stress alone or oxidative stress exacerbated by inflammation.

Another aspect of inflammation is the production of inflammatory prostaglandins such as prostaglandin E. These molecules promote vasodilation, plasma extravasation, localized pain, elevated temperature, and other symptoms of inflammation. The inducible form of the enzyme COX-2 is associated with their production, and high levels of COX-2 are found in inflamed tissues. Consequently, inhibition of COX-2 may relieve many symptoms of inflammation and a number of important anti-inflammatory drugs (*e.g.*, ibuprofen and celecoxib) act by inhibiting COX-2 activity. Recent research, however, has demonstrated that a class of cyclopentenone prostaglandins (cyPGs) (*e.g.*, 15-deoxy prostaglandin J2, a.k.a. PGJ2) plays a role in stimulating the orchestrated resolution of inflammation (*e.g.*, Rajakariar *et al.*, 2007). COX-2 is also associated with the production of cyclopentenone prostaglandins. Consequently, inhibition of COX-2 may interfere with the full resolution of inflammation,

potentially promoting the persistence of activated immune cells in tissues and leading to chronic, “smoldering” inflammation. This effect may be responsible for the increased incidence of cardiovascular disease in patients using selective COX-2 inhibitors for long periods of time.

5 In one aspect, the compounds of the invention may be used to control the production of pro-inflammatory cytokines within the cell by selectively activating regulatory cysteine residues (RCRs) on proteins that regulate the activity of redox-sensitive transcription factors. Activation of RCRs by cyPGs has been shown to initiate a pro-resolution program in which the activity of the antioxidant and cytoprotective transcription factor Nrf2 is potently induced and
10 the activities of the pro-oxidant and pro-inflammatory transcription factors NF- κ B and the STATs are suppressed. In some embodiments, this increases the production of antioxidant and reductive molecules (NQO1, HO-1, SOD1, γ -GCS) and decreases oxidative stress and the production of pro-oxidant and pro-inflammatory molecules (iNOS, COX-2, TNF- α). In some
15 embodiments, the compounds of this invention may cause the cells that host the inflammatory event to revert to a non-inflammatory state by promoting the resolution of inflammation and limiting excessive tissue damage to the host.

A. Cancer

 Further, the compounds of the present disclosure may be used to induce apoptosis in tumor cells, to induce cell differentiation, to inhibit cancer cell proliferation, to inhibit an
20 inflammatory response, and/or to function in a chemopreventative capacity. For example, the invention provides new compounds that have one or more of the following properties: (1) an ability to induce apoptosis and differentiate both malignant and non-malignant cells, (2) an activity at sub-micromolar or nanomolar levels as an inhibitor of proliferation of many malignant or premalignant cells, (3) an ability to suppress the *de novo* synthesis of the
25 inflammatory enzyme inducible nitric oxide synthase (iNOS), (4) an ability to inhibit NF- κ B activation, and (5) an ability to induce the expression of heme oxygenase-1 (HO-1).

 The levels of iNOS and COX-2 are elevated in certain cancers and have been implicated in carcinogenesis and COX-2 inhibitors have been shown to reduce the incidence of primary colonic adenomas in humans (Rostom *et al.*, 2007; Brown and DuBois, 2005;
30 Crowel *et al.*, 2003). iNOS is expressed in myeloid-derived suppressor cells (MDSCs) (Angulo *et al.*, 2000) and COX-2 activity in cancer cells has been shown to result in the production of prostaglandin E₂ (PGE₂), which has been shown to induce the expression of

arginase in MDSCs (Sinha *et al.*, 2007). Arginase and iNOS are enzymes that utilize L-arginine as a substrate and produce L-ornithine and urea, and L-citrulline and NO, respectively. The depletion of arginine from the tumor microenvironment by MDSCs, combined with the production of NO and peroxynitrite has been shown to inhibit proliferation and induce apoptosis of T cells (Bronte *et al.*, 2003). Inhibition of COX-2 and iNOS has been shown to reduce the accumulation of MDSCs, restore cytotoxic activity of tumor-associated T cells, and delay tumor growth (Sinha *et al.*, 2007; Mazzoni *et al.*, 2002; Zhou *et al.*, 2007).

Inhibition of the NF- κ B and JAK/STAT signaling pathways has been implicated as a strategy to inhibit proliferation of cancer epithelial cells and induce their apoptosis. Activation of STAT3 and NF- κ B has been shown to result in suppression of apoptosis in cancer cells, and promotion of proliferation, invasion, and metastasis. Many of the target genes involved in these processes have been shown to be transcriptionally regulated by both NF- κ B and STAT3 (Yu *et al.*, 2007).

In addition to their direct roles in cancer epithelial cells, NF- κ B and STAT3 also have important roles in other cells found within the tumor microenvironment. Experiments in animal models have demonstrated that NF- κ B is required in both cancer cells and hematopoietic cells to propagate the effects of inflammation on cancer initiation and progression (Greten *et al.*, 2004). NF- κ B inhibition in cancer and myeloid cells reduces the number and size, respectively, of the resultant tumors. Activation of STAT3 in cancer cells results in the production of several cytokines (IL-6, IL-10) which suppress the maturation of tumor-associated dendritic cells (DC). Furthermore, STAT3 is activated by these cytokines in the dendritic cells themselves. Inhibition of STAT3 in mouse models of cancer restores DC maturation, promotes antitumor immunity, and inhibits tumor growth (Kortylewski *et al.*, 2005).

B. Treatment of Multiple Sclerosis and Other Neurodegenerative Conditions

The compounds and methods of this invention may be used for treating patients for multiple sclerosis (MS). MS is known to be an inflammatory condition of the central nervous system (Williams *et al.*, 1994; Merrill and Benvenist, 1996; Genain and Nauser, 1997). Based on several investigations, there is evidence suggesting that inflammatory, oxidative, and/or immune mechanisms are involved in the pathogenesis of Alzheimer's disease (AD), Parkinson's disease (PD), amyotrophic lateral sclerosis (ALS), and MS (Bagasra *et al.*, 1995;

McGeer and McGeer, 1995; Simonian and Coyle, 1996; Kaltschmidt *et al.*, 1997). Both reactive astrocytes and activated microglia have been implicated in causation of neurodegenerative disease (NDD) and neuroinflammatory disease (NID); there has been a particular emphasis on microglia as cells that synthesize both NO and prostaglandins as products of the respective enzymes, iNOS and COX-2. *De novo* formation of these enzymes may be driven by inflammatory cytokines such as interferon- γ or interleukin-1. In turn, excessive production of NO may lead to inflammatory cascades and/or oxidative damage in cells and tissues of many organs, including neurons and oligodendrocytes of the nervous system, with consequent manifestations in AD and MS, and possible PD and ALS (Coyle and Puttfarcken, 1993; Beal, 1996; Merrill and Benvenist, 1996; Simonian and Coyle, 1996; Vodovotz *et al.*, 1996). Epidemiologic data indicate that chronic use of NSAID's which block synthesis of prostaglandins from arachidonate, markedly lower the risk for development of AD (McGeer *et al.*, 1996; Stewart *et al.*, 1997). Thus, agents that block formation of NO and prostaglandins, may be used in approaches to prevention and treatment of NDD. Successful therapeutic candidates for treating such a disease typically require an ability to penetrate the blood-brain barrier. See, for example, U.S. Patent Publication 2009/0060873, which is incorporated by reference herein in its entirety. See also, for example, the results presented for compound 404-02 in Examples 4 and 5, below.

C. Neuroinflammation

The compounds and methods of this invention may be used for treating patients with neuroinflammation. Neuroinflammation encapsulates the idea that microglial and astrocytic responses and actions in the central nervous system have a fundamentally inflammation-like character, and that these responses are central to the pathogenesis and progression of a wide variety of neurological disorders. This idea originated in the field of Alzheimer's disease (Griffin *et al.*, 1989; Rogers *et al.*, 1988), where it has revolutionized our understanding of this disease (Akiyama *et al.*, 2000). These ideas have been extended to other neurodegenerative diseases (Eikelenboom *et al.*, 2002; Ishizawa and Dickson, 2001), to ischemic/toxic diseases (Gehrmann *et al.*, 1995; Touzani *et al.*, 1999), to tumor biology (Graeber *et al.*, 2002) and even to normal brain development.

Neuroinflammation incorporates a wide spectrum of complex cellular responses that include activation of microglia and astrocytes and induction of cytokines, chemokines, complement proteins, acute phase proteins, oxidative injury, and related molecular processes.

These events may have detrimental effects on neuronal function, leading to neuronal injury, further glial activation, and ultimately neurodegeneration.

D. Treatment of Renal Failure

The compounds and methods of this invention may be used for treating patients with renal failure. See U.S. Patent Application 12/352,473, which is incorporated by reference herein in its entirety. Another aspect of the present disclosure concerns new methods and compounds for the treatment and prevention of renal disease. Renal failure, resulting in inadequate clearance of metabolic waste products from the blood and abnormal concentrations of electrolytes in the blood, is a significant medical problem throughout the world, especially in developed countries. Diabetes and hypertension are among the most important causes of chronic renal failure, also known as chronic kidney disease (CKD), but it is also associated with other conditions such as lupus. Acute renal failure may arise from exposure to certain drugs (*e.g.*, acetaminophen) or toxic chemicals, or from ischemia-reperfusion injury associated with shock or surgical procedures such as transplantation, and may result in chronic renal failure. In many patients, renal failure advances to a stage in which the patient requires regular dialysis or kidney transplantation to continue living. Both of these procedures are highly invasive and associated with significant side effects and quality of life issues. Although there are effective treatments for some complications of renal failure, such as hyperparathyroidism and hyperphosphatemia, no available treatment has been shown to halt or reverse the underlying progression of renal failure. Thus, agents that can improve compromised renal function would represent a significant advance in the treatment of renal failure.

Inflammation contributes significantly to the pathology of CKD. There is also a strong mechanistic link between oxidative stress and renal dysfunction. The NF- κ B signaling pathway plays an important role in the progression of CKD as NF- κ B regulates the transcription of MCP-1, a chemokine that is responsible for the recruitment of monocytes/macrophages resulting in an inflammatory response that ultimately injures the kidney (Wardle, 2001). The Keap1/Nrf2/ARE pathway controls the transcription of several genes encoding antioxidant enzymes, including heme oxygenase-1 (HO-1). Ablation of the Nrf2 gene in female mice results in the development of lupus-like glomerular nephritis (Yoh *et al.*, 2001). Furthermore, several studies have demonstrated that HO-1 expression is

induced in response to renal damage and inflammation and that this enzyme and its products – bilirubin and carbon monoxide – play a protective role in the kidney (Nath *et al.*, 2006).

The glomerulus and the surrounding Bowman's capsule constitute the basic functional unit of the kidney. Glomerular filtration rate (GFR) is the standard measure of renal function.

5 Creatinine clearance is commonly used to measure GFR. However, the level of serum creatinine is commonly used as a surrogate measure of creatinine clearance. For instance, excessive levels of serum creatinine are generally accepted to indicate inadequate renal function and reductions in serum creatinine over time are accepted as an indication of improved renal function. Normal levels of creatinine in the blood are approximately 0.6 to
10 1.2 milligrams (mg) per deciliter (dl) in adult males and 0.5 to 1.1 milligrams per deciliter in adult females.

Acute kidney injury (AKI) can occur following ischemia-reperfusion, treatment with certain pharmacological agents such as cisplatin and rapamycin, and intravenous injection of radiocontrast media used in medical imaging. As in CKD, inflammation and oxidative stress
15 contribute to the pathology of AKI. The molecular mechanisms underlying radiocontrast-induced nephropathy (RCN) are not well understood; however, it is likely that a combination of events including prolonged vasoconstriction, impaired kidney autoregulation, and direct toxicity of the contrast media all contribute to renal failure (Tumlin *et al.*, 2006). Vasoconstriction results in decreased renal blood flow and causes ischemia-reperfusion and
20 the production of reactive oxygen species. HO-1 is strongly induced under these conditions and has been demonstrated to prevent ischemia-reperfusion injury in several different organs, including the kidney (Nath *et al.*, 2006). Specifically, induction of HO-1 has been shown to be protective in a rat model of RCN (Goodman *et al.*, 2007). Reperfusion also induces an inflammatory response, in part through activation of NF- κ B signaling (Nichols, 2004).
25 Targeting NF- κ B has been proposed as a therapeutic strategy to prevent organ damage (Zingarelli *et al.*, 2003).

E. Cardiovascular Disease

The compounds and methods of this invention may be used for treating patients with cardiovascular disease. See U.S. Patent Application 12/352,473, which is incorporated by
30 reference herein in its entirety. Cardiovascular (CV) disease is among the most important causes of mortality worldwide, and is the leading cause of death in many developed nations. The etiology of CV disease is complex, but the majority of causes are related to inadequate or

completely disrupted supply of blood to a critical organ or tissue. Frequently such a condition arises from the rupture of one or more atherosclerotic plaques, which leads to the formation of a thrombus that blocks blood flow in a critical vessel. Such thrombosis is the principal cause of heart attacks, in which one or more of the coronary arteries is blocked and blood flow to the heart itself is disrupted. The resulting ischemia is highly damaging to cardiac tissue, both from lack of oxygen during the ischemic event and from excessive formation of free radicals after blood flow is restored (a phenomenon known as ischemia-reperfusion injury). Similar damage occurs in the brain during a thrombotic stroke, when a cerebral artery or other major vessel is blocked by thrombosis. Hemorrhagic strokes, in contrast, involve rupture of a blood vessel and bleeding into the surrounding brain tissue. This creates oxidative stress in the immediate area of the hemorrhage, due to the presence of large amounts of free heme and other reactive species, and ischemia in other parts of the brain due to compromised blood flow. Subarachnoid hemorrhage, which is frequently accompanied by cerebral vasospasm, also causes ischemia/reperfusion injury in the brain.

Alternatively, atherosclerosis may be so extensive in critical blood vessels that stenosis (narrowing of the arteries) develops and blood flow to critical organs (including the heart) is chronically insufficient. Such chronic ischemia can lead to end-organ damage of many kinds, including the cardiac hypertrophy associated with congestive heart failure.

Atherosclerosis, the underlying defect leading to many forms of cardiovascular disease, occurs when a physical defect or injury to the lining (endothelium) of an artery triggers an inflammatory response involving the proliferation of vascular smooth muscle cells and the infiltration of leukocytes into the affected area. Ultimately, a complicated lesion known as an atherosclerotic plaque may form, composed of the above-mentioned cells combined with deposits of cholesterol-bearing lipoproteins and other materials (*e.g.*, Hansson *et al.*, 2006).

Pharmaceutical treatments for cardiovascular disease include preventive treatments, such as the use of drugs intended to lower blood pressure or circulating levels of cholesterol and lipoproteins, as well as treatments designed to reduce the adherent tendencies of platelets and other blood cells (thereby reducing the rate of plaque progression and the risk of thrombus formation). More recently, drugs such as streptokinase and tissue plasminogen activator have been introduced and are used to dissolve the thrombus and restore blood flow. Surgical treatments include coronary artery bypass grafting to create an alternative blood supply, balloon angioplasty to compress plaque tissue and increase the diameter of the arterial

lumen, and carotid endarterectomy to remove plaque tissue in the carotid artery. Such treatments, especially balloon angioplasty, may be accompanied by the use of stents, expandable mesh tubes designed to support the artery walls in the affected area and keep the vessel open. Recently, the use of drug-eluting stents has become common in order to prevent post-surgical restenosis (renarrowing of the artery) in the affected area. These devices are wire stents coated with a biocompatible polymer matrix containing a drug that inhibits cell proliferation (*e.g.*, paclitaxel or rapamycin). The polymer allows a slow, localized release of the drug in the affected area with minimal exposure of non-target tissues. Despite the significant benefits offered by such treatments, mortality from cardiovascular disease remains high and significant unmet needs in the treatment of cardiovascular disease remain.

As noted above, induction of HO-1 has been shown to be beneficial in a variety of models of cardiovascular disease, and low levels of HO-1 expression have been clinically correlated with elevated risk of CV disease. Compounds of the invention, therefore, may be used in treating or preventing a variety of cardiovascular disorders including but not limited to atherosclerosis, hypertension, myocardial infarction, chronic heart failure, stroke, subarachnoid hemorrhage, and restenosis.

F. Diabetes

The compounds and methods of this invention may be used for treating patients with diabetes. See U.S. Patent Application 12/352,473, which is incorporated by reference herein in its entirety. Diabetes is a complex disease characterized by the body's failure to regulate circulating levels of glucose. This failure may result from a lack of insulin, a peptide hormone that regulates both the production and absorption of glucose in various tissues. Deficient insulin compromises the ability of muscle, fat, and other tissues to absorb glucose properly, leading to hyperglycemia (abnormally high levels of glucose in the blood). Most commonly, such insulin deficiency results from inadequate production in the islet cells of the pancreas. In the majority of cases this arises from autoimmune destruction of these cells, a condition known as type 1 or juvenile-onset diabetes, but may also be due to physical trauma or some other cause.

Diabetes may also arise when muscle and fat cells become less responsive to insulin and do not absorb glucose properly, resulting in hyperglycemia. This phenomenon is known as insulin resistance, and the resulting condition is known as Type 2 diabetes. Type 2 diabetes, the most common type, is highly associated with obesity and hypertension. Obesity

is associated with an inflammatory state of adipose tissue that is thought to play a major role in the development of insulin resistance (*e.g.*, Hotamisligil, 2006; Guilherme *et al.*, 2008).

Diabetes is associated with damage to many tissues, largely because hyperglycemia (and hypoglycemia, which can result from excessive or poorly timed doses of insulin) is a significant source of oxidative stress. Chronic kidney failure, retinopathy, peripheral neuropathy, peripheral vasculitis, and the development of dermal ulcers that heal slowly or not at all are among the common complications of diabetes. Because of their ability to protect against oxidative stress, particularly by the induction of HO-1 expression, compounds of the invention may be used in treatments for many complications of diabetes. As noted above (Cai *et al.*, 2005), chronic inflammation and oxidative stress in the liver are suspected to be primary contributing factors in the development of Type 2 diabetes. Furthermore, PPAR γ agonists such as thiazolidinediones are capable of reducing insulin resistance and are known to be effective treatments for Type 2 diabetes.

The effect of treatment of diabetes may be evaluated as follows. Both the biological efficacy of the treatment modality as well as the clinical efficacy are evaluated, if possible. For example, because the disease manifests itself by increased blood sugar, the biological efficacy of the treatment therefore can be evaluated, for example, by observation of return of the evaluated blood glucose towards normal. Measurement of glycosylated hemoglobin, also called A1c or HbA1c, is another commonly used parameter of blood glucose control. Measuring a clinical endpoint which can give an indication of b-cell regeneration after, for example, a six-month period of time, can give an indication of the clinical efficacy of the treatment regimen.

G. Rheumatoid Arthritis

The compounds and methods of this invention may be used for treating patients with RA. Typically the first signs of rheumatoid arthritis (RA) appear in the synovial lining layer, with proliferation of synovial fibroblasts and their attachment to the articular surface at the joint margin (Lipsky, 1998). Subsequently, macrophages, T cells and other inflammatory cells are recruited into the joint, where they produce a number of mediators, including the cytokines interleukin-1 (IL-1), which contributes to the chronic sequelae leading to bone and cartilage destruction, and tumour necrosis factor (TNF- α), which plays a role in inflammation (Dinarello, 1998; Arend and Dayer, 1995; van den Berg, 2001). The concentration of IL-1 in plasma is significantly higher in patients with RA than in healthy individuals and, notably,

plasma IL-1 levels correlate with RA disease activity (Eastgate *et al.*, 1988). Moreover, synovial fluid levels of IL-1 are correlated with various radiographic and histologic features of RA (Kahle *et al.*, 1992; Rooney *et al.*, 1990).

5 In normal joints, the effects of these and other proinflammatory cytokines are balanced by a variety of anti-inflammatory cytokines and regulatory factors (Burger and Dayer, 1995). The significance of this cytokine balance is illustrated in juvenile RA patients, who have cyclical increases in fever throughout the day (Prieur *et al.*, 1987). After each peak in fever, a factor that blocks the effects of IL-1 is found in serum and urine. This factor has been isolated, cloned and identified as IL-1 receptor antagonist (IL-1ra), a member of the IL-1
10 gene family (Hannum *et al.*, 1990). IL-1ra, as its name indicates, is a natural receptor antagonist that competes with IL-1 for binding to type I IL-1 receptors and, as a result, blocks the effects of IL-1 (Arend *et al.*, 1998). A 10- to 100-fold excess of IL-1ra may be needed to block IL-1 effectively; however, synovial cells isolated from patients with RA do not appear to produce enough IL-1ra to counteract the effects of IL-1 (Firestein *et al.*, 1994; Fujikawa *et al.*, 1995).
15

H. Psoriatic Arthritis

The compounds and methods of this invention may be used for treating patients with psoriatic arthritis. Psoriasis is an inflammatory and proliferative skin disorder with a prevalence of 1.5-3%. Approximately 20% of patients with psoriasis develop a characteristic
20 form of arthritis that has several patterns (Gladman, 1992; Jones *et al.*, 1994; Gladman *et al.*, 1995). Some individuals present with joint symptoms first but in the majority, skin psoriasis presents first. About one-third of patients have simultaneous exacerbations of their skin and joint disease (Gladman *et al.*, 1987) and there is a topographic relationship between nail and distal interphalangeal joint disease (Jones *et al.*, 1994; Wright, 1956). Although the
25 inflammatory processes which link skin, nail and joint disease remain elusive, an immune-mediated pathology is implicated.

Psoriatic arthritis (PsA) is a chronic inflammatory arthropathy characterized by the association of arthritis and psoriasis and was recognized as a clinical entity distinct from rheumatoid arthritis (RA) in 1964 (Blumberg *et al.*, 1964). Subsequent studies have revealed
30 that PsA shares a number of genetic, pathogenic and clinical features with other spondyloarthropathies (SpAs), a group of diseases that comprise ankylosing spondylitis, reactive arthritis and enteropathic arthritis (Wright, 1979). The notion that PsA belongs to the

SpA group has recently gained further support from imaging studies demonstrating widespread enthesitis in the, including PsA but not RA (McGonagle *et al.*, 1999; McGonagle *et al.*, 1998). More specifically, enthesitis has been postulated to be one of the earliest events occurring in the SpAs, leading to bone remodeling and ankylosis in the spine, as well as to articular synovitis when the inflamed entheses are close to peripheral joints. However, the link between enthesitis and the clinical manifestations in PsA remains largely unclear, as PsA can present with fairly heterogeneous patterns of joint involvement with variable degrees of severity (Marsal *et al.*, 1999; Salvarani *et al.*, 1998). Thus, other factors must be posited to account for the multifarious features of PsA, only a few of which (such as the expression of the HLA-B27 molecule, which is strongly associated with axial disease) have been identified. As a consequence, it remains difficult to map the disease manifestations to specific pathogenic mechanisms, which means that the treatment of this condition remains largely empirical.

Family studies have suggested a genetic contribution to the development of PsA (Moll and Wright, 1973). Other chronic inflammatory forms of arthritis, such as ankylosing spondylitis and rheumatoid arthritis, are thought to have a complex genetic basis. However, the genetic component of PsA has been difficult to assess for several reasons. There is strong evidence for a genetic predisposition to psoriasis alone that may mask the genetic factors that are important for the development of PsA. Although most would accept PsA as a distinct disease entity, at times there is a phenotypic overlap with rheumatoid arthritis and ankylosing spondylitis. Also, PsA itself is not a homogeneous condition and various subgroups have been proposed.

Increased amounts of TNF- α have been reported in both psoriatic skin (Ettehadi *et al.*, 1994) and synovial fluid (Partsch *et al.*, 1997). Recent trials have shown a positive benefit of anti-TNF treatment in both PsA (Mease *et al.*, 2000) and ankylosing spondylitis (Brandt *et al.*, 2000).

I. Reactive Arthritis

The compounds and methods of this invention may be used for treating patients with reactive arthritis. In reactive arthritis (ReA) the mechanism of joint damage is unclear, but it is likely that cytokines play critical roles. A more prevalent Th1 profile high levels of interferon gamma (IFN- γ) and low levels of interleukin 4 (IL-4) has been reported (Lahesmaa *et al.*, 1992; Schlaak *et al.*, 1992; Simon *et al.*, 1993; Schlaak *et al.*, 1996; Kotake *et al.*,

1999; Ribbens *et al.*, 2000), but several studies have shown relative predominance of IL-4 and IL-10 and relative lack of IFN- γ and tumour necrosis factor alpha (TNF- α) in the synovial membrane (Simon *et al.*, 1994; Yin *et al.*, 1999) and fluid (SF) (Yin *et al.*, 1999; Yin *et al.*, 1997) of reactive arthritis patients compared with rheumatoid arthritis (RA) patients. A lower
5 level of TNF- α secretion in reactive arthritis than in RA patients has also been reported after *ex vivo* stimulation of peripheral blood mononuclear cells (PBMC) (Braun *et al.*, 1999).

It has been argued that clearance of reactive arthritis-associated bacteria requires the production of appropriate levels of IFN- γ and TNF- α , while IL-10 acts by suppressing these responses (Autenrieth *et al.*, 1994; Sieper and Braun, 1995). IL-10 is a regulatory cytokine
10 that inhibits the synthesis of IL-12 and TNF- γ by activated macrophages (de Waal *et al.*, 1991; Hart *et al.*, 1995; Chomarat *et al.*, 1995) and of IFN- γ by T cells (Macatonia *et al.*, 1993).

J. Enteropathic Arthritis

The compounds and methods of this invention may be used for treating patients with
15 enteropathic arthritis. Typically enteropathic arthritis (EA) occurs in combination with inflammatory bowel diseases (IBD) such as Crohn's disease or ulcerative colitis. It also can affect the spine and sacroiliac joints. Enteropathic arthritis involves the peripheral joints, usually in the lower extremities such as the knees or ankles. It commonly involves only a few or a limited number of joints and may closely follow the bowel condition. This occurs in
20 approximately 11% of patients with ulcerative colitis and 21% of those with Crohn's disease. The synovitis is generally self-limited and non-deforming.

Enteropathic arthropathies comprise a collection of rheumatologic conditions that share a link to GI pathology. These conditions include reactive (*i.e.*, infection-related) arthritis due to bacteria (*e.g.*, *Shigella*, *Salmonella*, *Campylobacter*, *Yersinia* species,
25 *Clostridium difficile*), parasites (*e.g.*, *Strongyloides stercoralis*, *Taenia saginata*, *Giardia lamblia*, *Ascaris lumbricoides*, *Cryptosporidium* species), and spondyloarthropathies associated with inflammatory bowel disease (IBD). Other conditions and disorders include intestinal bypass (jejunoileal), arthritis, celiac disease, Whipple disease, and collagenous colitis.

K. Juvenile Rheumatoid Arthritis

The compounds and methods of this invention may be used for treating patients with JRA. Juvenile rheumatoid arthritis (JRA), a term for the most prevalent form of arthritis in children, is applied to a family of illnesses characterized by chronic inflammation and hypertrophy of the synovial membranes. The term overlaps, but is not completely synonymous, with the family of illnesses referred to as juvenile chronic arthritis and/or juvenile idiopathic arthritis in Europe.

Both innate and adaptive immune systems use multiple cell types, a vast array of cell surface and secreted proteins, and interconnected networks of positive and negative feedback (Lo *et al.*, 1999). Furthermore, while separable in thought, the innate and adaptive wings of the immune system are functionally intersected (Fearon and Locksley, 1996), and pathologic events occurring at these intersecting points are likely to be highly relevant to our understanding of pathogenesis of adult and childhood forms of chronic arthritis (Warrington, *et al.*, 2001).

Polyarticular JRA is a distinct clinical subtype characterized by inflammation and synovial proliferation in multiple joints (four or more), including the small joints of the hands (Jarvis, 2002). This subtype of JRA may be severe, because of both its multiple joint involvement and its capacity to progress rapidly over time. Although clinically distinct, polyarticular JRA is not homogeneous, and patients vary in disease manifestations, age of onset, prognosis, and therapeutic response. These differences very likely reflect a spectrum of variation in the nature of the immune and inflammatory attack that can occur in this disease (Jarvis, 1998).

L. Early Inflammatory Arthritis

The compounds and methods of this invention may be used for treating patients with early inflammatory arthritis. The clinical presentation of different inflammatory arthropathies is similar early in the course of disease. As a result, it is often difficult to distinguish patients who are at risk of developing the severe and persistent synovitis that leads to erosive joint damage from those whose arthritis is more self-limited. Such distinction is critical in order to target therapy appropriately, treating aggressively those with erosive disease and avoiding unnecessary toxicity in patients with more self-limited disease. Current clinical criteria for diagnosing erosive arthropathies such as rheumatoid arthritis (RA) are less effective in early disease and traditional markers of disease activity such as joint counts and acute phase

response do not adequately identify patients likely to have poor outcomes (Harrison *et al.*, 1998). Parameters reflective of the pathologic events occurring in the synovium are most likely to be of significant prognostic value.

Recent efforts to identify predictors of poor outcome in early inflammatory arthritis have identified the presence of RA specific autoantibodies, in particular antibodies towards citrullinated peptides, to be associated with erosive and persistent disease in early inflammatory arthritis cohorts. On the basis of this, a cyclical citrullinated peptide (CCP) has been developed to assist in the identification of anti-CCP antibodies in patient sera. Using this approach, the presence of anti-CCP antibodies has been shown to be specific and sensitive for RA, can distinguish RA from other arthropathies, and can potentially predict persistent, erosive synovitis before these outcomes become clinically manifest. Importantly, anti-CCP antibodies are often detectable in sera many years prior to clinical symptoms suggesting that they may be reflective of subclinical immune events (Nielen *et al.*, 2004; Rantapaa-Dahlqvist *et al.*, 2003).

M. Ankylosing Spondylitis

The compounds and methods of this invention may be used for treating patients with ankylosing spondylitis. AS is a disease subset within a broader disease classification of spondyloarthropathy. Patients affected with the various subsets of spondyloarthropathy have disease etiologies that are often very different, ranging from bacterial infections to inheritance. Yet, in all subgroups, the end result of the disease process is axial arthritis. Despite the early clinically differences seen in the various patient populations, many of them end up nearly identical after a disease course of ten-to-twenty years. Recent studies suggest the mean time to clinical diagnosis of ankylosing spondylitis from disease onset of disease is 7.5 years (Khan, 1998). These same studies suggest that the spondyloarthropathies may have prevalence close to that of rheumatoid arthritis (Feldtkeller *et al.*, 2003; Doran *et al.*, 2003).

AS is a chronic systemic inflammatory rheumatic disorder of the axial skeleton with or without extraskkeletal manifestations. Sacroiliac joints and the spine are primarily affected, but hip and shoulder joints, and less commonly peripheral joints or certain extra-articular structures such as the eye, vasculature, nervous system, and gastrointestinal system may also be involved. Its etiology is not yet fully understood (Wordsworth, 1995; Calin and Taurog, 1998). It is strongly associated with the major histocompatibility class I (MHC I) HLA-B27 allele (Calin and Taurog, 1998). AS affects individuals in the prime of their life and is feared

because of its potential to cause chronic pain and irreversible damage of tendons, ligaments, joints, and bones (Brewerton *et al.*, 1973a; Brewerton *et al.*, 1973b; Schlosstein *et al.*, 1973). AS may occur alone or in association with another form of spondyloarthropathy such as reactive arthritis, psoriasis, psoriatic arthritis, enthesitis, ulcerative colitis, irritable bowel disease, or Crohn's disease, in which case it is classified as secondary AS.

Typically, the affected sites include the discovertebral, apophyseal, costovertebral, and costotransverse joints of the spine, and the paravertebral ligamentous structures. Inflammation of the entheses, which are sites of musculotendinous and ligamentous attachment to bones, is also prominent in this disease (Calin and Taurog, 1998). The site of enthesitis is known to be infiltrated by plasma cells, lymphocytes, and polymorphonuclear cells. The inflammatory process frequently results in gradual fibrous and bony ankylosis, (Ball, 1971; Khan, 1990).

Delayed diagnosis is common because symptoms are often attributed to more common back problems. A dramatic loss of flexibility in the lumbar spine is an early sign of AS. Other common symptoms include chronic pain and stiffness in the lower back which usually starts where the lower spine is joined to the pelvis, or hip. Although most symptoms begin in the lumbar and sacroiliac areas, they may involve the neck and upper back as well. Arthritis may also occur in the shoulder, hips and feet. Some patients have eye inflammation, and more severe cases must be observed for heart valve involvement.

The most frequent presentation is back pain, but disease can begin atypically in peripheral joints, especially in children and women, and rarely with acute iritis (anterior uveitis). Additional early symptoms and signs are diminished chest expansion from diffuse costovertebral involvement, low-grade fever, fatigue, anorexia, weight loss, and anemia. Recurrent back pain - often nocturnal and of varying intensity - is an eventual complaint, as is morning stiffness typically relieved by activity. A flexed or bent-over posture eases back pain and paraspinal muscle spasm; thus, some degree of kyphosis is common in untreated patients.

Systemic manifestations occur in 1/3 of patients. Recurrent, usually self-limited, acute iritis (anterior uveitis) rarely is protracted and severe enough to impair vision. Neurologic signs can occasionally result from compression radiculitis or sciatica, vertebral fracture or subluxation, and cauda equina syndrome (which consists of impotence, nocturnal urinary incontinence, diminished bladder and rectal sensation, and absence of ankle jerks). Cardiovascular manifestations can include aortic insufficiency, angina, pericarditis, and ECG

conduction abnormalities. A rare pulmonary finding is upper lobe fibrosis, occasionally with cavitation that may be mistaken for TB and can be complicated by infection with *Aspergillus*.

AS is characterized by mild or moderate flares of active spondylitis alternating with periods of almost or totally inactive inflammation. Proper treatment in most patients results in minimal or no disability and in full, productive lives despite back stiffness. Occasionally, the course is severe and progressive, resulting in pronounced incapacitating deformities. The prognosis is bleak for patients with refractory iritis and for the rare patient with secondary amyloidosis.

N. Ulcerative Colitis

The compounds and methods of this invention may be used for treating patients with ulcerative colitis. Ulcerative colitis is a disease that causes inflammation and sores, called ulcers, in the lining of the large intestine. The inflammation usually occurs in the rectum and lower part of the colon, but it may affect the entire colon. Ulcerative colitis rarely affects the small intestine except for the end section, called the terminal ileum. Ulcerative colitis may also be called colitis or proctitis. The inflammation makes the colon empty frequently, causing diarrhea. Ulcers form in places where the inflammation has killed the cells lining the colon; the ulcers bleed and produce pus.

Ulcerative colitis is an inflammatory bowel disease (IBD), the general name for diseases that cause inflammation in the small intestine and colon. Ulcerative colitis can be difficult to diagnose because its symptoms are similar to other intestinal disorders and to another type of IBD, Crohn's disease. Crohn's disease differs from ulcerative colitis because it causes inflammation deeper within the intestinal wall. Also, Crohn's disease usually occurs in the small intestine, although it can also occur in the mouth, esophagus, stomach, duodenum, large intestine, appendix, and anus.

Ulcerative colitis may occur in people of any age, but most often it starts between ages 15 and 30, or less frequently between ages 50 and 70. Children and adolescents sometimes develop the disease. Ulcerative colitis affects men and women equally and appears to run in some families. Theories about what causes ulcerative colitis abound, but none have been proven. The most popular theory is that the body's immune system reacts to a virus or a bacterium by causing ongoing inflammation in the intestinal wall. People with ulcerative colitis have abnormalities of the immune system, but doctors do not know whether these abnormalities are a cause or a result of the disease. Ulcerative colitis is not caused by

emotional distress or sensitivity to certain foods or food products, but these factors may trigger symptoms in some people.

The most common symptoms of ulcerative colitis are abdominal pain and bloody diarrhea. Patients also may experience fatigue, weight loss, loss of appetite, rectal bleeding, and loss of body fluids and nutrients. About half of patients have mild symptoms. Others suffer frequent fever, bloody diarrhea, nausea, and severe abdominal cramps. Ulcerative colitis may also cause problems such as arthritis, inflammation of the eye, liver disease (hepatitis, cirrhosis, and primary sclerosing cholangitis), osteoporosis, skin rashes, and anemia. No one knows for sure why problems occur outside the colon. Scientists think these complications may occur when the immune system triggers inflammation in other parts of the body. Some of these problems go away when the colitis is treated.

A thorough physical exam and a series of tests may be required to diagnose ulcerative colitis. Blood tests may be done to check for anemia, which could indicate bleeding in the colon or rectum. Blood tests may also uncover a high white blood cell count, which is a sign of inflammation somewhere in the body. By testing a stool sample, the doctor can detect bleeding or infection in the colon or rectum. The doctor may do a colonoscopy or sigmoidoscopy. For either test, the doctor inserts an endoscope - a long, flexible, lighted tube connected to a computer and TV monitor - into the anus to see the inside of the colon and rectum. The doctor will be able to see any inflammation, bleeding, or ulcers on the colon wall. During the exam, the doctor may do a biopsy, which involves taking a sample of tissue from the lining of the colon to view with a microscope. A barium enema x ray of the colon may also be required. This procedure involves filling the colon with barium, a chalky white solution. The barium shows up white on x-ray film, allowing the doctor a clear view of the colon, including any ulcers or other abnormalities that might be there.

Treatment for ulcerative colitis depends on the seriousness of the disease. Most people are treated with medication. In severe cases, a patient may need surgery to remove the diseased colon. Surgery is the only cure for ulcerative colitis. Some people whose symptoms are triggered by certain foods are able to control the symptoms by avoiding foods that upset their intestines, like highly seasoned foods, raw fruits and vegetables, or milk sugar (lactose). Each person may experience ulcerative colitis differently, so treatment is adjusted for each individual. Emotional and psychological support is important. Some people have remissions - periods when the symptoms go away - that last for months or even years. However, most patients' symptoms eventually return. This changing pattern of the disease means one cannot

always tell when a treatment has helped. Some people with ulcerative colitis may need medical care for some time, with regular doctor visits to monitor the condition.

O. Crohn's Disease

The compounds and methods of this invention may be used for treating patients with Crohn's disease. Another disorder for which immunosuppression has been tried is Crohn's disease. Crohn's disease symptoms include intestinal inflammation and the development of intestinal stenosis and fistulas; neuropathy often accompanies these symptoms. Anti-inflammatory drugs, such as 5-aminosalicylates (*e.g.*, mesalamine) or corticosteroids, are typically prescribed, but are not always effective (reviewed in Botoman *et al.*, 1998). Immunosuppression with cyclosporine is sometimes beneficial for patients resistant to or intolerant of corticosteroids (Brynskov *et al.*, 1989).

Efforts to develop diagnostic and treatment tools against Crohn's disease have focused on the central role of cytokines (Schreiber, 1998; van Hogezaand and Verspaget, 1998). Cytokines are small secreted proteins or factors (5 to 20 kD) that have specific effects on cell-to-cell interactions, intercellular communication, or the behavior of other cells. Cytokines are produced by lymphocytes, especially T_H1 and T_H2 lymphocytes, monocytes, intestinal macrophages, granulocytes, epithelial cells, and fibroblasts (reviewed in Rogler and Andus, 1998; Galley and Webster, 1996). Some cytokines are pro-inflammatory (*e.g.*, TNF- α , IL-1(α and β), IL-6, IL-8, IL-12, or leukemia inhibitory factor [LIF]); others are anti-inflammatory (*e.g.*, IL-1 receptor antagonist, IL-4, IL-10, IL-11, and TGF- β). However, there may be overlap and functional redundancy in their effects under certain inflammatory conditions.

In active cases of Crohn's disease, elevated concentrations of TNF- α and IL-6 are secreted into the blood circulation, and TNF- α , IL-1, IL-6, and IL-8 are produced in excess locally by mucosal cells (*id.*; Funakoshi *et al.*, 1998). These cytokines can have far-ranging effects on physiological systems including bone development, hematopoiesis, and liver, thyroid, and neuropsychiatric function. Also, an imbalance of the IL-1 β /IL-1ra ratio, in favor of pro-inflammatory IL-1 β , has been observed in patients with Crohn's disease (Rogler and Andus, 1998; Saiki *et al.*, 1998; Dionne *et al.*, 1998; but see Kuboyama, 1998). One study suggested that cytokine profiles in stool samples could be a useful diagnostic tool for Crohn's disease (Saiki *et al.*, 1998).

Treatments that have been proposed for Crohn's disease include the use of various cytokine antagonists (*e.g.*, IL-1ra), inhibitors (*e.g.*, of IL-1 β converting enzyme and

antioxidants) and anti-cytokine antibodies (Rogler and Andus, 1998; van Hogezaand and Verspaget, 1998; Reimund *et al.*, 1998; Luger *et al.*, 1998; McAlindon *et al.*, 1998). In particular, monoclonal antibodies against TNF- α have been tried with some success in the treatment of Crohn's disease (Targan *et al.*, 1997; Stack *et al.*, 1997; van Dullemen *et al.*, 1995). These compounds may be used in combination therapy with compounds of the present disclosure.

Another approach to the treatment of Crohn's disease has focused on at least partially eradicating the bacterial community that may be triggering the inflammatory response and replacing it with a non-pathogenic community. For example, U.S. Patent 5,599,795 discloses a method for the prevention and treatment of Crohn's disease in human patients. Their method was directed to sterilizing the intestinal tract with at least one antibiotic and at least one anti-fungal agent to kill off the existing flora and replacing them with different, select, well-characterized bacteria taken from normal humans. Borody taught a method of treating Crohn's disease by at least partial removal of the existing intestinal microflora by lavage and replacement with a new bacterial community introduced by fecal inoculum from a disease-screened human donor or by a composition comprising *Bacteroides* and *Escherichia coli* species. (U.S. Patent 5,443,826).

P. Systemic Lupus Erythematosus

The compounds and methods of this invention may be used for treating patients with SLE. There has also been no known cause for autoimmune diseases such as systemic lupus erythematosus. Systemic lupus erythematosus (SLE) is an autoimmune rheumatic disease characterized by deposition in tissues of autoantibodies and immune complexes leading to tissue injury (Kotzin, 1996). In contrast to autoimmune diseases such as MS and type 1 diabetes mellitus, SLE potentially involves multiple organ systems directly, and its clinical manifestations are diverse and variable (reviewed by Kotzin and O'Dell, 1995). For example, some patients may demonstrate primarily skin rash and joint pain, show spontaneous remissions, and require little medication. At the other end of the spectrum are patients who demonstrate severe and progressive kidney involvement that requires therapy with high doses of steroids and cytotoxic drugs such as cyclophosphamide (Kotzin, 1996).

The serological hallmark of SLE, and the primary diagnostic test available, is elevated serum levels of IgG antibodies to constituents of the cell nucleus, such as double-stranded DNA (dsDNA), single-stranded DNA (ss-DNA), and chromatin. Among these

autoantibodies, IgG anti-dsDNA antibodies play a major role in the development of lupus glomerulonephritis (G N) (Hahn and Tsao, 1993; Ohnishi *et al.*, 1994). Glomerulonephritis is a serious condition in which the capillary walls of the kidney's blood purifying glomeruli become thickened by accretions on the epithelial side of glomerular basement membranes.

5 The disease is often chronic and progressive and may lead to eventual renal failure.

Q. Irritable Bowel Syndrome

The compounds and methods of this invention may be used for treating patients with Irritable bowel syndrome (IBS). IBS is a functional disorder characterized by abdominal pain and altered bowel habits. This syndrome may begin in young adulthood and can be
10 associated with significant disability. This syndrome is not a homogeneous disorder. Rather, subtypes of IBS have been described on the basis of the predominant symptom--diarrhea, constipation, or pain. In the absence of "alarm" symptoms, such as fever, weight loss, and gastrointestinal bleeding, a limited workup is needed. Once a diagnosis of IBS is made, an integrated treatment approach can effectively reduce the severity of symptoms. IBS is a
15 common disorder, although its prevalence rates have varied. In general, IBS affects about 15% of US adults and occurs about three times more often in women than in men (Jailwala *et al.*, 2000).

IBS accounts for between 2.4 million and 3.5 million visits to physicians each year. It not only is the most common condition seen by gastroenterologists but also is one of the most
20 common gastrointestinal conditions seen by primary care physicians (Everhart *et al.*, 1991; Sandler, 1990).

IBS is also a costly disorder. Compared with persons who do not have bowel symptoms, persons with IBS miss three times as many workdays and are more likely to report being too sick to work (Drossman *et al.*, 1993; Drossman *et al.*, 1997). Moreover, those with
25 IBS incur hundreds of dollars more in medical charges than persons without bowel disorders (Talley *et al.*, 1995).

No specific abnormality accounts for the exacerbations and remissions of abdominal pain and altered bowel habits experienced by patients with IBS. The evolving theory of IBS suggests dysregulation at multiple levels of the brain-gut axis. Dysmotility, visceral
30 hypersensitivity, abnormal modulation of the central nervous system (CNS), and infection have all been implicated. In addition, psychosocial factors play an important modifying role. Abnormal intestinal motility has long been considered a factor in the pathogenesis of IBS.

Transit time through the small intestine after a meal has been shown to be shorter in patients with diarrhea-predominant IBS than in patients who have the constipation-predominant or pain-predominant subtype (Cann *et al.*, 1983).

In studies of the small intestine during fasting, the presence of both discrete, clustered
5 contractions and prolonged, propagated contractions has been reported in patients with IBS (Kellow and Phillips, 1987). They also experience pain with irregular contractions more often than healthy persons (Kellow and Phillips, 1987; Horwitz and Fisher, 2001)

These motility findings do not account for the entire symptom complex in patients with IBS; in fact, most of these patients do not have demonstrable abnormalities (Rothstein,
10 2000). Patients with IBS have increased sensitivity to visceral pain. Studies involving balloon distention of the rectosigmoid colon have shown that patients with IBS experience pain and bloating at pressures and volumes much lower than control subjects (Whitehead *et al.*, 1990). These patients maintain normal perception of somatic stimuli.

Multiple theories have been proposed to explain this phenomenon. For example,
15 receptors in the viscera may have increased sensitivity in response to distention or intraluminal contents. Neurons in the dorsal horn of the spinal cord may have increased excitability. In addition, alteration in CNS processing of sensations may be involved (Drossman *et al.*, 1997). Functional magnetic resonance imaging studies have recently shown that compared with control subjects, patients with IBS have increased activation of the
20 anterior cingulate cortex, an important pain center, in response to a painful rectal stimulus (Mertz *et al.*, 2000).

Increasingly, evidence suggests a relationship between infectious enteritis and subsequent development of IBS. Inflammatory cytokines may play a role. In a survey of patients with a history of confirmed bacterial gastroenteritis (Neal *et al.*, 1997), 25% reported
25 persistent alteration of bowel habits. Persistence of symptoms may be due to psychological stress at the time of acute infection (Gwee *et al.*, 1999).

Recent data suggest that bacterial overgrowth in the small intestine may have a role in IBS symptoms. In one study (Pimentel *et al.*, 2000), 157 (78%) of 202 IBS patients referred for hydrogen breath testing had test findings that were positive for bacterial overgrowth. Of
30 the 47 subjects who had follow-up testing, 25 (53%) reported improvement in symptoms (*i.e.*, abdominal pain and diarrhea) with antibiotic treatment.

IBS may present with a range of symptoms. However, abdominal pain and altered bowel habits remain the primary features. Abdominal discomfort is often described as crampy

in nature and located in the left lower quadrant, although the severity and location can differ greatly. Patients may report diarrhea, constipation, or alternating episodes of diarrhea and constipation. Diarrheal symptoms are typically described as small-volume, loose stools, and stool is sometimes accompanied by mucus discharge. Patients also may report bloating, fecal
5 urgency, incomplete evacuation, and abdominal distention. Upper gastrointestinal symptoms, such as gastroesophageal reflux, dyspepsia, or nausea, may also be present (Lynn and Friedman, 1993).

Persistence of symptoms is not an indication for further testing; it is a characteristic of IBS and is itself an expected symptom of the syndrome. More extensive diagnostic evaluation
10 is indicated in patients whose symptoms are worsening or changing. Indications for further testing also include presence of alarm symptoms, onset of symptoms after age 50, and a family history of colon cancer. Tests may include colonoscopy, computed tomography of the abdomen and pelvis, and barium studies of the small or large intestine.

R. Sjögren's Syndrome

The compounds and methods of this invention may be used for treating patients with SS. Primary Sjögren's syndrome (SS) is a chronic, slowly progressive, systemic autoimmune disease, which affects predominantly middle-aged women (female-to-male ratio 9:1), although it can be seen in all ages including childhood (Jonsson *et al.*, 2002). It is characterized by lymphocytic infiltration and destruction of the exocrine glands, which are
15 infiltrated by mononuclear cells including CD4+, CD8+ lymphocytes and B-cells (Jonsson *et al.*, 2002). In addition, extraglandular (systemic) manifestations are seen in one-third of patients (Jonsson *et al.*, 2001).

The glandular lymphocytic infiltration is a progressive feature (Jonsson *et al.*, 1993), which, when extensive, may replace large portions of the organs. Interestingly, the glandular
25 infiltrates in some patients closely resemble ectopic lymphoid microstructures in the salivary glands (denoted as ectopic germinal centers) (Salomonsson *et al.*, 2002; Xanthou *et al.*, 2001). In SS, ectopic GCs are defined as T and B cell aggregates of proliferating cells with a network of follicular dendritic cells and activated endothelial cells. These GC-like structures formed within the target tissue also portray functional properties with production of
30 autoantibodies (anti-Ro/SSA and anti-La/SSB) (Salomonsson and Jonsson, 2003).

In other systemic autoimmune diseases, such as RA, factors critical for ectopic GCs have been identified. Rheumatoid synovial tissues with GCs were shown to produce

chemokines CXCL13, CCL21 and lymphotoxin (LT)- β (detected on follicular center and mantle zone B cells). Multivariate regression analysis of these analytes identified CXCL13 and LT- β as the solitary cytokines predicting GCs in rheumatoid synovitis (Weyand and Goronzy, 2003). Recently CXCL13 and CXCR5 in salivary glands has been shown to play an essential role in the inflammatory process by recruiting B and T cells, therefore contributing to lymphoid neogenesis and ectopic GC formation in SS (Salomonsson *et al.*, 2002).

S. Psoriasis

The compounds and methods of this invention may be used for treating patients with psoriasis. Psoriasis is a chronic skin disease of scaling and inflammation that affects 2 to 2.6 percent of the United States population, or between 5.8 and 7.5 million people. Although the disease occurs in all age groups, it primarily affects adults. It appears about equally in males and females. Psoriasis occurs when skin cells quickly rise from their origin below the surface of the skin and pile up on the surface before they have a chance to mature. Usually this movement (also called turnover) takes about a month, but in psoriasis it may occur in only a few days. In its typical form, psoriasis results in patches of thick, red (inflamed) skin covered with silvery scales. These patches, which are sometimes referred to as plaques, usually itch or feel sore. They most often occur on the elbows, knees, other parts of the legs, scalp, lower back, face, palms, and soles of the feet, but they can occur on skin anywhere on the body. The disease may also affect the fingernails, the toenails, and the soft tissues of the genitals and inside the mouth. While it is not unusual for the skin around affected joints to crack, approximately 1 million people with psoriasis experience joint inflammation that produces symptoms of arthritis. This condition is called psoriatic arthritis.

Psoriasis is a skin disorder driven by the immune system, especially involving a type of white blood cell called a T cell. Normally, T cells help protect the body against infection and disease. In the case of psoriasis, T cells are put into action by mistake and become so active that they trigger other immune responses, which lead to inflammation and to rapid turnover of skin cells. In about one-third of the cases, there is a family history of psoriasis. Researchers have studied a large number of families affected by psoriasis and identified genes linked to the disease. People with psoriasis may notice that there are times when their skin worsens, then improves. Conditions that may cause flareups include infections, stress, and changes in climate that dry the skin. Also, certain medicines, including lithium and beta

blockers, which are prescribed for high blood pressure, may trigger an outbreak or worsen the disease.

T. Infectious diseases

Compounds of the present disclosure may be useful in the treatment of infectious diseases, including viral and bacterial infections. As noted above, such infections may be associated with severe localized or systemic inflammatory responses. For example, influenza may cause severe inflammation of the lung and bacterial infection can cause the systemic hyperinflammatory response, including the excessive production of multiple inflammatory cytokines, that is the hallmark of sepsis. In addition, compounds of the invention may be useful in directly inhibiting the replication of viral pathogens. Previous studies have demonstrated that related compounds such as CDDO can inhibit the replication of HIV in macrophages (Vazquez *et al.*, 2005). Other studies have indicated that inhibition of NF-kappa B signaling may inhibit influenza virus replication, and that cyclopentenone prostaglandins may inhibit viral replication (*e.g.*, Mazur *et al.*, 2007; Pica *et al.*, 2000).

VI. Pharmaceutical Formulations and Routes of Administration

The compounds of the present disclosure may be administered by a variety of methods, *e.g.*, orally or by injection (*e.g.* subcutaneous, intravenous, intraperitoneal, *etc.*). Depending on the route of administration, the active compounds may be coated in a material to protect the compound from the action of acids and other natural conditions which may inactivate the compound. They may also be administered by continuous perfusion/infusion of a disease or wound site.

To administer the therapeutic compound by other than parenteral administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation. For example, the therapeutic compound may be administered to a patient in an appropriate carrier, for example, liposomes, or a diluent. Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Liposomes include water-in-oil-in-water CGF emulsions as well as conventional liposomes (Strejan *et al.*, 1984).

The therapeutic compound may also be administered parenterally, intraperitoneally, intraspinally, or intracerebrally. Dispersions can be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. In all cases, the composition must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (such as, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, sodium chloride, or polyalcohols such as mannitol and sorbitol, in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate or gelatin.

Sterile injectable solutions can be prepared by incorporating the therapeutic compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the therapeutic compound into a sterile carrier which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient (*i.e.*, the therapeutic compound) plus any additional desired ingredient from a previously sterile-filtered solution thereof.

The therapeutic compound can be orally administered, for example, with an inert diluent or an assimilable edible carrier. The therapeutic compound and other ingredients may also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. For oral therapeutic administration, the therapeutic compound may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. The percentage of the therapeutic compound in the compositions and preparations may, of

course, be varied. The amount of the therapeutic compound in such therapeutically useful compositions is such that a suitable dosage will be obtained.

It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit containing a predetermined quantity of therapeutic compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the therapeutic compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such a therapeutic compound for the treatment of a selected condition in a patient.

The therapeutic compound may also be administered topically to the skin, eye, or mucosa. Alternatively, if local delivery to the lungs is desired the therapeutic compound may be administered by inhalation in a dry-powder or aerosol formulation.

Active compounds are administered at a therapeutically effective dosage sufficient to treat a condition associated with a condition in a patient. A “therapeutically effective amount” preferably reduces the amount of symptoms of the condition in the infected patient by at least about 20%, more preferably by at least about 40%, even more preferably by at least about 60%, and still more preferably by at least about 80% relative to untreated subjects. For example, the efficacy of a compound can be evaluated in an animal model system that may be predictive of efficacy in treating the disease in humans, such as the model systems shown in the examples and drawings.

The actual dosage amount of a compound of the present disclosure or composition comprising a compound of the present disclosure administered to a subject may be determined by physical and physiological factors such as age, sex, body weight, severity of condition, the type of disease being treated, previous or concurrent therapeutic interventions, idiopathy of the subject and on the route of administration. These factors may be determined by a skilled artisan. The practitioner responsible for administration will typically determine the concentration of active ingredient(s) in a composition and appropriate dose(s) for the individual subject. The dosage may be adjusted by the individual physician in the event of any complication.

An effective amount typically will vary from about 0.001 mg/kg to about 1,000 mg/kg, from about 0.01 mg/kg to about 750 mg/kg, from about 100 mg/kg to about 500

mg/kg, from about 1.0 mg/kg to about 250 mg/kg, from about 10.0 mg/kg to about 150 mg/kg in one or more dose administrations daily, for one or several days (depending, of course, of the mode of administration and the factors discussed above). Other suitable dose ranges include 1 mg to 10,000 mg per day, 100 mg to 10,000 mg per day, 500 mg to 10,000 mg per day, and 500 mg to 1,000 mg per day. In some particular embodiments, the amount is less than 10,000 mg per day with a range, for example, of 750 mg to 9,000 mg per day.

The effective amount may be less than 1 mg/kg/day, less than 500 mg/kg/day, less than 250 mg/kg/day, less than 100 mg/kg/day, less than 50 mg/kg/day, less than 25 mg/kg/day or less than 10 mg/kg/day. It may alternatively be in the range of 1 mg/kg/day to 200 mg/kg/day. For example, regarding treatment of diabetic patients, the unit dosage may be an amount that reduces blood glucose by at least 40% as compared to an untreated subject. In another embodiment, the unit dosage is an amount that reduces blood glucose to a level that is $\pm 10\%$ of the blood glucose level of a non-diabetic subject.

In other non-limiting examples, a dose may also comprise from about 1 microgram/kg/body weight, about 5 microgram/kg/body weight, about 10 microgram/kg/body weight, about 50 microgram/kg/body weight, about 100 microgram/kg/body weight, about 200 microgram/kg/body weight, about 350 microgram/kg/body weight, about 500 microgram/kg/body weight, about 1 milligram/kg/body weight, about 5 milligram/kg/body weight, about 10 milligram/kg/body weight, about 50 milligram/kg/body weight, about 100 milligram/kg/body weight, about 200 milligram/kg/body weight, about 350 milligram/kg/body weight, about 500 milligram/kg/body weight, to about 1,000 mg/kg/body weight or more per administration, and any range derivable therein. In non-limiting examples of a derivable range from the numbers listed herein, a range of about 5 mg/kg/body weight to about 100 mg/kg/body weight, about 5 microgram/kg/body weight to about 500 milligram/kg/body weight, *etc.*, can be administered, based on the numbers described above.

In certain embodiments, a pharmaceutical composition of the present disclosure may comprise, for example, at least about 0.1% of a compound of the present disclosure. In other embodiments, the compound of the present disclosure may comprise between about 2% to about 75% of the weight of the unit, or between about 25% to about 60%, for example, and any range derivable therein.

Single or multiple doses of the agents are contemplated. Desired time intervals for delivery of multiple doses can be determined by one of ordinary skill in the art employing no more than routine experimentation. As an example, subjects may be administered two doses

daily at approximately 12 hour intervals. In some embodiments, the agent is administered once a day.

The agent(s) may be administered on a routine schedule. As used herein a routine schedule refers to a predetermined designated period of time. The routine schedule may encompass periods of time which are identical or which differ in length, as long as the schedule is predetermined. For instance, the routine schedule may involve administration twice a day, every day, every two days, every three days, every four days, every five days, every six days, a weekly basis, a monthly basis or any set number of days or weeks therebetween. Alternatively, the predetermined routine schedule may involve administration on a twice daily basis for the first week, followed by a daily basis for several months, *etc.* In other embodiments, the invention provides that the agent(s) may be taken orally and that the timing of which is or is not dependent upon food intake. Thus, for example, the agent can be taken every morning and/or every evening, regardless of when the subject has eaten or will eat.

VII. Combination Therapy

In addition to being used as a monotherapy, the compounds of the present disclosure may also find use in combination therapies. Effective combination therapy may be achieved with a single composition or pharmacological formulation that includes both agents, or with two distinct compositions or formulations, at the same time, wherein one composition includes a compound of this invention, and the other includes the second agent(s). Alternatively, the therapy may precede or follow the other agent treatment by intervals ranging from minutes to months.

Various combinations may be employed, such as when a compound of the present disclosure is "A" and "B" represents a secondary agent, non-limiting examples of which are described below:

A/B/A	B/A/B	B/B/A	A/A/B	A/B/B	B/A/A	A/B/B/B	B/A/B/B
B/B/B/A	B/B/A/B	A/A/B/B	A/B/A/B	A/B/B/A	B/B/A/A		
B/A/B/A	B/A/A/B	A/A/A/B	B/A/A/A	A/B/A/A	A/A/B/A		

Administration of the compounds of the present disclosure to a patient will follow general protocols for the administration of pharmaceuticals, taking into account the toxicity, if any, of the drug. It is expected that the treatment cycles would be repeated as necessary.

Beta interferons may be suitable secondary agents. These are medications derived from human cytokines which help regulate the immune system. They include interferon β -1b and interferon β -1a. Betaseron has been approved by the FDA for relapsing forms of secondary progressive MS. Furthermore, the FDA has approved the use of several β -interferons as treatments for people who have experienced a single attack that suggests multiple sclerosis, and who may be at risk of future attacks and developing definite MS. For example, risk of MS may be suggested when an MRI scan of the brain shows lesions that predict a high risk of conversion to definite MS.

Glatiramer acetate is a further example of a secondary agent that may be used in a combination treatment. Glatiramer is presently used to treat relapsing remitting MS. It is made of four amino acids that are found in myelin. This drug is reported to stimulate T cells in the body's immune system to change from harmful, pro-inflammatory agents to beneficial, anti-inflammatory agents that work to reduce inflammation at lesion sites.

Another potential secondary agent is mitoxantrone, a chemotherapy drug used for many cancers. This drug is also FDA-approved for treatment of aggressive forms of relapsing remitting MS, as well as certain forms of progressive MS. It is given intravenously, typically every three months. This medication is effective, but is limited by cardiac toxicity. Novantrone has been approved by the FDA for secondary progressive, progressive-relapsing, and worsening relapsing-remitting MS.

Another potential secondary agent is natalizumab. In general, natalizumab works by blocking the attachment of immune cells to brain blood vessels, which is a necessary step for immune cells to cross into the brain, thus reducing the immune cells' inflammatory action on brain neurons. Natalizumab has been shown to significantly reduce the frequency of attacks in people with relapsing MS.

In the case of relapsing remitting MS, patients may be given intravenous corticosteroids, such as methylprednisolone, as a secondary agent, to end the attack sooner and leave fewer lasting deficits.

Other common drugs for MS that may be used in combination with the present oleanolic acid derivatives include immunosuppressive drugs such as azathioprine, cladribine and cyclophosphamide.

It is contemplated that other anti-inflammatory agents may be used in conjunction with the treatments of the current invention. Other COX inhibitors may be used, including arylcarboxylic acids (salicylic acid, acetylsalicylic acid, diflunisal, choline magnesium

trisalicylate, salicylate, benorylate, flufenamic acid, mefenamic acid, meclofenamic acid and triflumic acid), arylalkanoic acids (diclofenac, fenclofenac, alclofenac, fentiazac, ibuprofen, flurbiprofen, ketoprofen, naproxen, fenoprofen, fenbufen, suprofen, indoprofen, tiaprofenic acid, benoxaprofen, piroprofen, tolmetin, zomepirac, clopinac, indomethacin and sulindac) and enolic acids (phenylbutazone, oxyphenbutazone, azapropazone, feprazone, piroxicam, and isoxicam. See also U.S. Pat. No. 6,025,395, which is incorporated herein by reference.

Histamine H₂ receptor blocking agents may also be used in conjunction with the compounds of the current invention, including cimetidine, ranitidine, famotidine and nizatidine.

Treatment with acetylcholinesterase inhibitors such as tacrine, donepezil, metrifonate and rivastigmine for the treatment of Alzheimer's and other disease in conjunction with the compounds of the present disclosure is contemplated. Other acetylcholinesterase inhibitors may be developed which may be used once approved include rivastigmine and metrifonate. Acetylcholinesterase inhibitors increase the amount of neurotransmitter acetylcholine at the nerve terminal by decreasing its breakdown by the enzyme cholinesterase.

MAO-B inhibitors such as selegilene may be used in conjunction with the compounds of the current invention. Selegilene is used for Parkinson's disease and irreversibly inhibits monoamine oxidase type B (MAO-B). Monoamine oxidase is an enzyme that inactivates the monoamine neurotransmitters norepinephrine, serotonin and dopamine.

Dietary and nutritional supplements with reported benefits for treatment or prevention of Parkinson's, Alzheimer's, multiple sclerosis, amyotrophic lateral sclerosis, rheumatoid arthritis, inflammatory bowel disease, and all other diseases whose pathogenesis is believed to involve excessive production of either nitric oxide (NO) or prostaglandins, such as acetyl-L-carnitine, octacosanol, evening primrose oil, vitamin B₆, tyrosine, phenylalanine, vitamin C, L-dopa, or a combination of several antioxidants may be used in conjunction with the compounds of the current invention.

For the treatment or prevention of cancer, compounds of the invention may be combined with one or more of the following: radiation, chemotherapy agents (*e.g.*, cytotoxic agents such as anthracyclines, vincristine, vinblastin, microtubule-targeting agents such as paclitaxel and docetaxel, 5-FU and related agents, cisplatin and other platinum-containing compounds, irinotecan and topotecan, gemcitabine, temozolomide, etc.), targeted therapies (*e.g.*, imatinib, bortezomib, bevacizumab, rituximab), or vaccine therapies designed to promote an enhanced immune response targeting cancer cells.

For the treatment or prevention of autoimmune disease, compounds of the invention may be combined with one or more of the following: corticosteroids, methotrexate, anti-TNF antibodies, other TNF-targeting protein therapies, and NSAIDs. For the treatment or prevention of cardiovascular diseases, compounds of the invention may be combined with antithrombotic therapies, anticholesterol therapies such as statins (*e.g.*, atorvastatin), and surgical interventions such as stenting or coronary artery bypass grafting. For the treatment of osteoporosis, compounds of the invention may be combined with antiresorptive agents such as bisphosphonates or anabolic therapies such as teriparatide or parathyroid hormone. For the treatment of neuropsychiatric conditions, compounds of the invention may be combined with antidepressants (*e.g.*, imipramine or SSRIs such as fluoxetine), antipsychotic agents (*e.g.*, olanzapine, sertindole, risperidone), mood stabilizers (*e.g.*, lithium, valproate semisodium), or other standard agents such as anxiolytic agents. For the treatment of neurological disorders, compounds of the invention may be combined with anticonvulsant agents (*e.g.*, valproate semisodium, gabapentin, phenytoin, carbamazepine, and topiramate), antithrombotic agents (*e.g.*, tissue plasminogen activator), or analgesics (*e.g.*, opioids, sodium channel blockers, and other antinociceptive agents).

For the treatment of disorders involving oxidative stress, compounds of the present disclosure may be combined with tetrahydrobiopterin (BH4) or related compounds. BH4 is a cofactor for constitutive forms of nitric oxide synthase, and may be depleted by reactions with peroxynitrite. Peroxynitrite is formed by the reaction of nitric oxide and superoxide. Thus, under conditions of oxidative stress excessive levels of superoxide can deplete normal, beneficial levels of nitric oxide by converting NO to peroxynitrite. The resulting depletion of BH4 by reaction with peroxynitrite results in the “uncoupling” of nitric oxide synthases so that they form superoxide rather than NO. This adds to the oversupply of superoxide and prolongs the depletion of NO. Addition of exogenous BH4 can reverse this uncoupling phenomenon, restoring the production of NO and reducing the level of oxidative stress in tissues. This mechanism is expected to complement the actions of compounds of the invention, which reduce oxidative stress by other means, as discussed above and throughout this invention.

VIII. Examples

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in

the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1 – Methods and Materials

Nitric Oxide production and cell viability. RAW264.7 macrophages were pre-treated with DMSO or drugs for 2 hours, then treated with recombinant mouse IFN γ (Sigma) for 24 hours. NO concentration in media was determined using the Griess reagent system (Promega). Cell viability was determined using WST-1 reagent (Roche).

STAT3 phosphorylation. HeLa cells were treated with the indicated compounds and concentrations for 6 hours and subsequently stimulated with 20 ng/ml recombinant human IL-6 (R&D Systems) for 15 minutes. Lysates were immunoblotted with antibodies against phosphorylated or total STAT3 (Cell Signaling).

NF- κ B activation. HeLa cells were transfected with pNF- κ B-Luc (inducible, Stratagene) and pRL-TK (constitutive, Promega) reporter plasmids. Twenty-four hours later cells were pre-treated with the indicated compounds for 2 hours. DMSO served as a vehicle control. Following pre-treatment, cells were stimulated with 20 ng/ml recombinant human TNF α (BD Biosciences) for 3 hours. Reporter activity was measured using DualGlo luciferase reporter system (Promega) and pNF- κ B luciferase activity was normalized against pRL-TK luciferase activity. Fold-induction of mean luciferase activity relative to unstimulated (-TNF α) samples is shown. Error bars represent the SD of the mean of 6 samples.

I κ B α degradation. HeLa cells were treated with indicated compounds and concentrations for 6 hours and subsequently stimulated with 20 ng/ml TNF α for 15 minutes. Lysates were blotted with antibodies against I κ B α (Santa Cruz) and actin (Chemicon).

COX-2 induction Western blot. RAW264.7 cells were pre-treated for 2 hours with indicated compounds and subsequently stimulated with 10 ng/ml IFN γ for an additional 24 hours. COX-2 protein levels were assayed by immunoblotting using an antibody from Santa Cruz. Actin was used as a loading control.

Nrf2 target gene induction. MDA-MB-435 human melanoma cells were treated with vehicle (DMSO) or the indicated compounds and concentrations for 16 hours. HO-1, thioredoxin reductase-1 (TrxR1), γ -glutamylcysteine synthetase (γ -GCS), and ferritin heavy chain mRNA levels were quantified using qPCR and were normalized relative to a DMSO-treated sample run in parallel. Values are averages of duplicate wells. Primer sequences are as follows.

HO-1 FW: TCCGATGGGTCCTTACACTC (SEQ ID NO:1),

HO-1 REV: TAGGCTCCTTCCTCCTTTCC (SEQ ID NO:2),

TrxR1 FW: GCAGCACTGAGTGGTCAAAA (SEQ ID NO:3),

TrxR1 REV: GGTCAACTGCCTCAATTGCT (SEQ ID NO:4),

γ -GCS FW: GCTGTGGCTACTGCGGTATT (SEQ ID NO:5),

γ -GCS REV ATCTGCCTCAATGACACCAT (SEQ ID NO:6),





Ferritin HC FW: ATGAGCAGGTGAAAGCCATC (SEQ ID NO:7),









Ferritin HC REV: TAAAGGAAACCCCAACATGC (SEQ ID NO:8),

S9 FW: GATTACATCCTGGGCCTGAA (SEQ ID NO:9),

S9 REV: GAGCGCAGAGAGAAGTCGAT (SEQ ID NO:10).

Comparison Compounds. Some of the experimental results presented below and throughout this application present data for not only the compounds discussed above, but also for one or more of the triterpenoid derivatives shown in the table below.

 401 (RTA 401)	 402 (RTA 402)
 402-19	 402-52

 402-53	 402-54
 402-55	 402-56
 404 (RTA 404)	 63112
 63324	 63166

Several of the above compounds, including **401**, **402**, **402-56** and **404** can be prepared according to the methods taught by Honda *et al.* (1998), Honda *et al.* (2000b), Honda *et al.* (2002), Yates *et al.* (2007), U.S. Patent 6,974,801, and U.S. Provisional Applications 61/046,342, 61/046,352, 61/046,366, 61/111,269, and 61/111,294, which are all incorporated

5 herein by reference. The synthesis of the other compounds may be prepared according to the methods disclosed in one or more of the following applications filed concurrently herewith, each of which is incorporated herein by reference in their entireties: U.S. Patent Application

by Eric Anderson, Xin Jiang and Melean Visnick, entitled "Antioxidant Inflammation Modulators: Oleanolic Acid Derivatives with Amino and Other Modifications At C-17," filed April 20, 2009; U.S. Patent Application by Xin Jiang, Jack Greiner, Lester Maravetz, Stephen S. Szucs, Melean Visnick, entitled "Antioxidant Inflammation Modulators: Novel Derivatives of Oleanolic Acid," filed April 20, 2009; U.S. Patent Application by Xin Jiang, Xiaofeng Liu, Jack Greiner, Stephen S. Szucs, Melean Visnick entitled, "Antioxidant Inflammation Modulators: C-17 Homologated Oleanolic Acid Derivatives," filed April 20, 2009.

Aqueous Solubility Determination. The following procedure was used to obtain the aqueous solubility results summarized in Example 8. Step 1. Determination of optimal UV/vis wavelengths and generation of standard curves for a compound of interest:

- (1) For eight standard calibration curves (one plate), prepare 34 mL of 50:50 (v:v) universal buffer:acetonitrile in a 50 mL tube.
- (2) Using a multichannel pipet, dispense (in μL) the buffer:acetonitrile in a deep well plate as follows:

	1	2	3	4	5	6	7	8	9	10	11	12
A	285	285	380	380	285	285	285	285	285	285	285	285
B												
C												
D												
E												
F												
G												
H												

- (3) Using a multichannel pipet, dispense DMSO into the same plate as follows:


	1	2	3	4	5	6	7	8	9	10	11	12
A			12 μL	12 μL	15 μL	15 μL	15 μL	15 μL	15 μL	15 μL	15 μL	15 μL
B												
C												
D												
E												
F												
G												
H												

(4) Add 10 mM compound in DMSO into the plates as follows:


	1	2	3	4	5	6	7	8	9	10	11	12
A	15 μ L cmpd1	15 μ L cmpd1	8 μ L cmpd1	8 μ L cmpd1								
B	15 μ L cmpd2	15 μ L cmpd2	8 μ L cmpd2	8 μ L cmpd2								
C	15 μ L cmpd3	15 μ L cmpd3	8 μ L cmpd3	8 μ L cmpd3								
D	15 μ L cmpd4	15 μ L cmpd4	8 μ L cmpd4	8 μ L cmpd4								
E	15 μ L cmpd5	15 μ L cmpd5	8 μ L cmpd5	8 μ L cmpd5								
F	15 μ L cmpd6	15 μ L cmpd6	8 μ L cmpd6	8 μ L cmpd6								
G	15 μ L cmpd7	15 μ L cmpd7	8 μ L cmpd7	8 μ L cmpd7								
H	15 μ L cmpd8	15 μ L cmpd8	8 μ L cmpd8	8 μ L cmpd8								

(5) Mix columns 1 and 2 by pipetting each up and down 10 times. Mix columns 3 and 4 by pipetting up and down 10 times. Serially dilute as follows (pipet up and down 10 times after each transfer):

5



	1	2	3	4	5	6	7	8	9	10	11	12
A												
B												
C												
D												
E												
F												
G												
H												



Note columns 11 and 12 contain DMSO only and so compound should not be transferred to these wells.

10 (6) Cover plate with lid and shake (200-300 rpm) at room temperature for 20 minutes.

(7) Mix all wells by pipetting up and down 10 times.

- (8) Transfer 120 μ L from each well to a UV transparent plate. Cover and shake for 3-5 minutes. Remove any bubbles in the wells using a pipet.
- (9) Read from 220 nm to 500 nm at 10 nm increments on a spectrophotometer (*e.g.*, SpectraMax®).

5 Step 2. Compound Solubility Testing Procedures using the Millipore™ Multiscreen® Solubility Filter Plate.

Consumables: Millipore™ Multiscreen® Solubility Filter Plate #MSSLBPC10

Greiner® 96 well disposable UV-Star analysis plate, VWR#655801

Greiner® 96 well polypropylene V-bottom collection plate, VWR#651201

10 Universal Aqueous Buffer:

- (a) To prepare 500 mL of universal buffer, add the following: 250 mL Nanopure water; 1.36 mL (45 mM) ethanolamine; 3.08 g (45 mM) potassium dihydrogen phosphate; 2.21 g (45 mM) potassium acetate; thoroughly mix.
- (b) Adjust pH to 7.4 with HCl and q.s. to 500 mL with 0.15 M KCl.
- (c) Filter to remove particulates and reduce bacterial growth.
- (d) Store at 4°C in the dark.

Solubility Protocol:

- (a) Add 285 μ L of Universal Aqueous Buffer to desired wells of the Millipore™ Multiscreen® Solubility filter plate.
- (b) Add 15 μ L of 10 mM compound in DMSO to the appropriate wells. Add 15 μ L of 100% DMSO only to 6 wells of the filter plate for blanks.
- (c) Using a multichannel pipet, mix wells by pipetting up and down 10 times. Be careful not to touch the filters in the plate with the tips.
- (d) Cover and gently shake (200-300 rpm) filter plate for 90 minutes at room temperature.
- (e) Vacuum filter the aqueous solution from the Multiscreen® solubility filter plate into a polypropylene V-bottom plate.
- (f) Transfer 60 μ L of filtrate to a UV transparent plate (Greiner® UV-Star Analysis Plate).

- (g) Add 60 μ L of acetonitrile to each well and mix by pipetting up and down 10 times.
- (h) Cover and gently shake for 3-5 minutes. Remove any bubbles with a pipet.
- 5 (i) Measure the absorbance of each well in the plate on the spectrophotometer (UV/vis) at the desired wavelength. For compounds in a plate with different absorbance peaks, set the spectrophotometer to read a spectrum (*e.g.*, from 220 nm to 460 nm).
- 10 (j) Identify concentration using measured absorbance for each compound and the predetermined standard curve (see Step 1).


Example 2 – Synthesis of Oleanolic Acid Derivatives

The synthesis of compounds **402-02** and **402-51** started from compound **1** (Scheme 1). Compound **1** was oxidized with bleach to give ketone **2** in 80% yield. Formylation of **2** with ethyl formate using sodium methoxide as the base afforded compound **402-48** (70% yield),

15 which was then treated with hydroxylamine hydrochloride in aqueous EtOH at 55 °C to give isoxazole **402-49** in 93% yield. Cleavage of the isoxazole under basic conditions gave α -cyanoketone **402-46** in quantitative yield as a mixture of ketone and enol forms. Compound **402-46** was treated with 1,3-dibromo-5,5-dimethylhydantoin, followed by elimination of HBr using pyridine as the base, to give compound **402-02** in 81% yield (from **402-49**), which was

20 demethylated with LiI in refluxing DMF to give acid **402-51** in 95% yield.


Scheme 1:



Reagents and conditions applicable to Scheme 1 are: (a) AcOH, bleach, room temperature, 1 h, 80%; (b) HCO₂Et, NaOMe, 55 °C, 24 h, 70%; (c) NH₂OH·HCl, 55 °C, 16 h, 93%; (d) NaOMe, 55 °C, 2 h, 100%; (e) (i) 1,3-dibromo-5,5-dimethylhydantoin, room temperature, 2 h; (ii) pyridine, 55 °C, 15 h, 81%; (f) LiI, 160 °C, 8 h, 95%.

Compound **402-63** was oxidized with Dess-Martin periodinane to give aldehyde **402-64** in 47% yield (Scheme 2).


Scheme 2:



Reagents and conditions applicable to Scheme 2 are: (a) Dess-Martin periodinane, NaHCO₃, room temperature, 1 h, 47%.

The synthesis of compounds **402-59** and **402-57** began from **402-51**. Compound **402-51** was treated with oxalyl chloride and catalytic DMF to give acid chloride **3**. Compound **3** was treated with ammonia in methanol to give **402-59** (99% from **402-51**). Dehydration of **402-59** utilizing TFAA and Et₃N gave dicyano compound **402-57** (45% yield).


Scheme 3:



Reagents and conditions applicable to Scheme 3 are: (a) $(\text{COCl})_2$, DMF (cat.), 0 °C to room temperature, 3 h; (b) NH_3 (MeOH), 0 °C to room temperature, 5 h, 99%; (c) TFAA, Et_3N , 0 °C, 3 h, 45%.


404-02 was synthesized from compound **3** as summarized in Scheme 4. Compound **3** was reacted with 2,2,2-trifluoroethylamine-HCl in toluene and water at 70 °C with NaHCO_3 as the base, giving **404-02** in 69% yield.

10 Scheme 4:



Reagents and conditions applicable to Scheme 4 are: (a) 2,2,2-trifluoroethylamine-HCl, NaHCO_3 , toluene, H_2O , 70 °C, 69%.


Scheme 5:



Reagents and conditions applicable to Scheme 5 are: (a) LiAlH_4 , THF, 0 °C, 1.5 h, 38% for **4**; 34% for **5**; (b) NaOMe, 55 °C, 7 h, 94% for **6**; 89% for **7**; (c) (i) 1,3-dibromo-5,5-dimethylhydantoin, r.t., 40 min; (ii) pyridine, 55 °C, 25 h, 35% for **402-66**; 32% for **63219**.


5 The synthesis of **402-66** began from isoxazole compound **402-49**. Reduction of the ketone in **402-49** was achieved by treatment with LiAlH_4 in THF at 0 °C, giving compounds **4** and **5** (as a 1:1 mixture of diastereotopic alcohols). Compound **4** was treated with NaOMe in MeOH at 55 °C to give **6**, which exists as a 3:2 mixture of keto and enol tautomeric forms. Bromination and subsequent dehydrobromination of **6** by treatment with 1,3-dibromo-5,5-dimethylhydantoin and then pyridine gave **402-66** in 33% yield (from **4**).
10 Using the same synthetic sequence, compound **5** was converted into **63219** in 28% overall yield.

Scheme 6:




Reagent and conditions applicable to Scheme 6 are: (a) oxalyl chloride, r.t., 2 h; (b) $\text{NH}_2\text{NH}_2\text{-H}_2\text{O}$, 0 °C, 30 min, 97%; (c) AcCl , Et_3N , r.t., 1 h, 77%; (d) NaOMe , r.t., 10 min, 72%; (e) TsOH , 110 °C, 1 h, 33%.

Scheme 7:




Reagent and conditions applicable to Scheme 7 are: (a) Et₃N, TFAA, r.t., 1.5 h, 85%; (b) Bu₃SnN₃, 150 °C, 67%; (c) (i) HCO₂Et, NaOMe, 0 °C to room temperature, 1 h; (ii) NH₂OH·HCl, 60 °C, 3 h, 54%; (d) NaOMe, 55 °C, 2 h; (e) 1,3-dibromo-5,5-dimethylhydantoin, r.t., 2 h; (f) pyridine, 55 °C, 16 h, 60% for **63229**; 5 69% for **63230**; (g) TMSCHN₂, 0 °C, 10 min, 77%.

Scheme 8:




Reagents and conditions applicable to Scheme 8 are: (a) i) NaOMe, HCO₂Et, 0 °C to room temperature; concentrated HCl ii) NH₂OH-HCl, EtOH, H₂O, 60 °C, 14 h, 47%; (b) NaOMe, 55 °C, 16 h, quantitative; (c) (i) 1,3-dibromo-5,5-dimethylhydantoin, room temperature, 3 h; (ii) pyridine, 55 °C, 16 h, 12%.

5 Scheme 9:




Reagents and conditions applicable to Scheme 9 are: (a) LiAlH₄, THF, 0 °C, 2 h, 19%; (b) NaOMe, 55 °C, 7 h, 92%; (c) (i) 1,3-dibromo-5,5-dimethylhydantoin, r.t., 50 min; (ii) pyridine, 55 °C, 7 h, 56%.

Scheme 10:




Reagents and conditions pertaining to Scheme 10: (a) (i) HCO₂Et, NaOMe, 0 °C, 1.5 h; (ii) NH₂OH-HCl, 65 °C, 3.5 h, 78%; (b) (i) oxalyl chloride, 0 °C to rt, 2 h; (ii) AcNHNH₂, Et₃N, 0 °C to rt, 30 min, 99%; (c) Lawesson's reagent, 110 °C, 30 min, 10% (for compound **21**) and 29% (for compound **22**); (d) NaOMe, 55 °C, 2 h, 73%; (e) (i) DBDMH, 0 °C, 1 h; (ii) 5 pyridine, 55 °C, 3 h, 79%. Compound **18** was reported by Honda *et al.* (2000a), which is incorporated herein by reference in its entirety.

Scheme 11:




(continued on next page)


(continued from previous page):




Reagents and conditions applicable to Scheme 11: (a) LAH, rt to 65 °C, 1.5 h, 27%; (b) TEMPO, IPh(OAc)₂, rt, 72 h, 77%; (c) (Ph₃PCH₂Cl)Cl / n-BuLi, THF, HMPA, 0 °C to rt, 87%; (d) MeLi, THF, 0 °C to rt, 91%; (e) PCC, NaOAc, CH₂Cl₂, rt, 78%; (f) HCO₂Et, NaOMe, 0 °C to rt; (g) NH₂OH·HCl, EtOH-H₂O, 60 °C, 89% from **28**; (h) NaOMe, 55 °C, 3 h; (i) DDQ, Benzene, 85 °C, 39% from **30**.

Scheme 12:


Reagents and conditions applicable to Scheme 12 are: (a) HgSO₄, H₂SO₄, acetone/H₂O, 55 °C, 20 h, 91%; (b) (i) 1,3-dibromo-5,5-dimethylhydantoin, 0 °C, 3 h; (ii) pyridine, 55 °C, 19 h, 73%.

Scheme 13:


Reagents and conditions pertaining to Scheme 13: (a) benzyl bromide, DBU, 100 °C, 6 h, 65%.

5 Scheme 14:


Reagents and conditions pertaining to Scheme 14: (a) MeONH₂-HCl, Et₃N, 40 °C, 4 h, 37%.

Scheme 15:


Reagents and conditions pertaining to Scheme 15: (a) Me₂NH, 40 °C, 71 h, 61%.

Scheme 16:


Reagents and conditions applicable to Scheme 16 are: (a) $\text{NH}_2\text{OH}\cdot\text{HCl}$, NaOAc , EtOH , H_2O , $80\text{ }^\circ\text{C}$, 27 h, 72%; (b) NaOMe , MeOH , $55\text{ }^\circ\text{C}$, 1 h; (c) (i) 1,3-dibromo-5,5-dimethylhydantoin, $0\text{ }^\circ\text{C}$, 40 min; (ii) pyridine, $55\text{ }^\circ\text{C}$, 7 h, 27% from **33**.

Scheme 17:

Reagents and conditions applicable to Scheme 17 are: (a) POCl_3 , pyridine, rt, 5 h, 75%; (b) NaOMe , MeOH , $55\text{ }^\circ\text{C}$, 4 h, 93% (i) 1,3-dibromo-5,5-dimethylhydantoin, $0\text{ }^\circ\text{C}$, 1 h; (ii) pyridine, $55\text{ }^\circ\text{C}$, 23 h, 93%.

Scheme 18:

Reagents and conditions applicable to Scheme 18 are: (a) *m*-CPBA, rt, 48 h, 22%; (b) NaOMe, 55 °C, 2 h, 66%; (c) (i) DBDMH, 0 °C, 1 h; (ii) pyridine, 55 °C, 3 h, 72%.

5 **Scheme 19:**

Reagents and conditions pertaining applicable to Scheme 19 are: (a) LiAlH₄, 0 °C, 40 min, 62%; (b) NaOMe, 55 °C, 2 h, 83%; (c) (i) DBDMH, 0 °C, 1 h; (ii) pyridine, 55 °C, 4 h, 40%.

10 **Example 3 – Synthesis and Characterization of Oleanolic Acid Derivatives**

Compound 2: Bleach (5.25 wt% of NaClO (aq), 129 mL, 91 mmol) was added to a stirred solution of compound 1 (34.67 g, 71 mmol) in AcOH (471 mL) at room temperature.

After stirring for 40 min, the reaction mixture was poured into ice-water (1.5 L) and stirred for 5 min. The white precipitate was collected by filtration and washed thoroughly with water. The filtered solid was then dissolved in EtOAc and washed with NaHCO₃ (aq) solution, dried with MgSO₄ and concentrated. The residue obtained was purified by column chromatography (silica gel, 10% to 25% EtOAc in hexanes) to give product **2** (27.8 g, 80%) as a white foam solid: ¹H NMR (400 MHz, CDCl₃) δ 3.69 (s, 3H), 2.80 (m, 1H), 2.65 (d, 1H, *J* = 4.0 Hz), 2.53 (ddd, 1H, *J* = 7.2, 10.8, 16.0 Hz), 2.38 (ddd, 1H, *J* = 3.6, 6.8, 16.0 Hz), 2.16-2.30 (m, 2 H), 1.95 (m, 1H), 1.89 (m, 1H), 1.80 (m, 2H), 1.62-1.73 (m, 3H), 1.57 (m, 2H), 1.47 (m, 2H), 1.15-1.40 (m, 7H), 1.09 (s, 3H), 1.05 (s, 3H), 1.01 (s, 3H), 0.99 (s, 3H), 0.98 (s, 3H), 0.95 (s, 3H), 0.90 (s, 3H); *m/z* 485.3 (M+1).

Compound 402-48: NaOMe solution (25% w/w in MeOH, 132.3 mL, 570 mmol) was added to a solution of compound **2** (27.6 g, 57 mmol) in MeOH (250 mL) under nitrogen. The reaction mixture was heated to 55 °C in an oil bath, and HCO₂Et (93 mL, 1.15 mmol, 20 eq) was added dropwise via an addition funnel. The reaction mixture was stirred at 55 °C for 24 h and then at room temperature for another 40 h. After removal of MeOH (150 mL) by evaporation, *t*-BuOMe (200 mL) was added, and the mixture was cooled to 0 °C. 12 N HCl (aq) (50 mL, 600 mmol, 10.5 eq) was then added over 10 min, and the mixture was extracted with EtOAc. The combined extracts were washed with water, dried with MgSO₄, and concentrated. The brown oil obtained was purified by column chromatography (silica gel, 5% to 10% EtOAc in hexanes) to give product **402-48** (20.5 g, 70%) as a white foam solid: ¹H NMR (400 MHz, CDCl₃) δ 14.89 (d, 1H, *J* = 3.2 Hz), 8.61 (d, 1H, *J* = 3.2 Hz), 3.69 (s, 3H), 2.80 (m, 1H), 2.67 (d, 1H, *J* = 4.0 Hz), 2.20-2.34 (m, 3H), 1.98 (m, 1H), 1.62-1.92 (m, 6H), 1.10-1.56 (m, 10H), 1.20 (s, 3H), 1.12 (s, 3H), 1.02 (s, 3H), 0.99 (s, 3H), 0.96 (s, 3H), 0.91 (s, 3H), 0.85 (s, 3H); *m/z* 513.3 (M+1).

Compound 402-49: A mixture of compound **402-48** (20.3 g, 40 mmol) and NH₂OH·HCl (4.12 g, 59 mmol) in EtOH (300 mL) and water (60 mL) was heated at 55 °C for 14 h. After cooling to room temperature, EtOH was removed by evaporation, and the white slurry obtained was extracted with EtOAc. The combined extracts were washed with water, dried with MgSO₄, and concentrated. The residue obtained was purified by column chromatography (silica gel, 10% to 20% EtOAc in hexanes) to give product **402-49** (18.8 g, 93%) as a white solid: ¹H NMR (400 MHz, CDCl₃) δ 7.99 (s, 1H), 3.70 (s, 3H), 2.81 (m, 1H), 2.68 (d, 1H, *J* = 4.4 Hz), 2.37 (d, 1H, *J* = 15.2 Hz), 2.23-2.33 (m, 2H), 1.76-1.98 (m, 5H),

1.68 (m, 3H), 1.11-1.62 (m, 9H), 1.32 (s, 3H), 1.23 (s, 3H), 1.02 (s, 3H), 0.99 (s, 3H), 0.97 (s, 3H), 0.91 (s, 3H), 0.84 (s, 3H); m/z 510.3 (M+1).

Compound 402-46: NaOMe (25% w/w in MeOH, 8.75 mL, 38 mmol) was added dropwise to a suspension of **402-49** (16.16 g, 31.7 mmol) in MeOH (55 mL) at 0 °C under N₂.

5 The reaction mixture was heated at 55 °C for 2 h and then cooled to 0 °C. *t*-BuOMe (150 mL) and 1 N HCl (aq) (50 mL) were added successively, and the mixture was extracted with EtOAc. The combined extracts were washed with water, dried with MgSO₄, and concentrated to give compound **402-46** (17.80 g, 100%) as a white foam solid. **402-46** is a mixture of two equilibrium forms, the enol form (as shown in Scheme 1) and the ketone form,
10 in the ratio of 2:3. ¹H NMR of the mixture: (400 MHz, CDCl₃) δ 5.69 (s, 0.4H), 3.87 (m, 0.6H), 2.80 (m, 1H), 2.65 (m, 1H), 0.82-2.30 (m, 44H); m/z 510.3 (M+1).

Compound 402-02: 1,3-Dibromo-5,5-dimethylhydantoin (5.98 g, 20.9 mmol) was added to a solution of compound **402-46** (17.76 g, 35 mmol) in DMF (75 mL) at 10 °C. After stirring at room temperature for 2 h, pyridine (8.5 mL, 105 mmol) was added, and the
15 reaction mixture was heated at 55 °C for 15 h. After cooling to room temperature, the mixture was poured into water (700 mL) and stirred for 5 min. The pale brown precipitate was collected by filtration and washed with water. The solid was dissolved in CH₂Cl₂, and the solution was washed with 1 N HCl (aq) and water, then dried with MgSO₄ and concentrated. The residue obtained was purified by column chromatography (silica gel, 0%
20 to 70% EtOAc in hexanes) to give product **402-02** (14.3 g, 81%) as a white solid: ¹H NMR (400 MHz, CDCl₃) δ 7.65 (s, 1H), 3.69 (s, 3H), 2.82 (m, 1H), 2.68 (d, 1H, *J* = 4.4 Hz), 2.44 (dd, 1H, *J* = 4.8, 16.0 Hz), 2.35 (dd, 1H, *J* = 12.8, 16.0 Hz), 1.86-2.00 (m, 3H), 1.81 (m, 1H), 1.60-1.71 (m, 4H), 1.42-1.55 (m, 3H), 1.24 (m, 1H), 1.10-1.24 (m, 4H), 1.22 (s, 3H), 1.16 (s, 3H), 1.15 (s, 3H), 1.07 (s, 3H), 0.99 (s, 3H), 0.97 (s, 3H), 0.92 (s, 3H); m/z 508.2 (M+1).

25 **Compound 402-51:** A stream of nitrogen was bubbled through a stirring solution of compound **402-02** (6.31 g, 12.4 mmol) and LiI (33.35 g, 248 mmol) in DMF (87 mL) at 160 °C for 8 h. After cooling to 50 °C, the reaction mixture was diluted with EtOAc (100 mL). 1 N HCl (aq) solution (30 mL) was then added at room temperature and stirred for 5 min. The mixture was extracted with EtOAc, and the combined extracts were washed with water, 10%
30 Na₂S₂O₃ (aq) and water, then dried with MgSO₄ and concentrated. The residue obtained was purified by column chromatography (silica gel, 5% to 50% EtOAc in CH₂Cl₂) to give acid **402-51** (6.02 g, 95%) as a white solid: ¹H NMR (400 MHz, CDCl₃) δ 10.41 (bs, 1H), 7.65 (s, 1H), 2.80 (m, 1H), 2.74 (d, 1H, *J* = 4.4 Hz), 2.46 (dd, 1H, *J* = 4.8, 16.0 Hz), 2.37 (dd, 1H, *J* =

12.8, 16.0 Hz), 1.86-2.02 (m, 4H), 1.44-1.79 (m, 8H), 1.35 (m, 1H), 1.12-1.29 (m, 3H), 1.22 (s, 3H), 1.16 (s, 3H), 1.14 (s, 3H), 1.11 (s, 3H), 0.99 (s, 3H), 0.98 (s, 3H), 0.93 (s, 3H); m/z 494.3 (M+1).

Compound 402-64: NaHCO₃ (78 mg, 0.93 mmol) and Dess-Martin periodinane (99 mg, 0.23 mmol) were added successively to a solution of **402-63** (45 mg, 94 μ mol) in CH₂Cl₂ (5 mL) at room temperature. After stirring for 1 h, 5% Na₂S₂O₃ (aq) solution was added. The reaction mixture was extracted with *t*-BuOMe, and the combined extracts were washed with NaHCO₃ (aq) solution, dried with MgSO₄, and concentrated. The crude product obtained was purified by column chromatography (silica gel, 0% to 35% EtOAc in hexanes) to give **402-64** (22 mg, 49%) as a white foam solid: ¹H NMR (300 MHz, CDCl₃) δ 9.33 (s, 1H), 7.62 (s, 1H), 2.61 (m, 1H), 2.50 (d, 1H, J = 4.4 Hz), 2.45 (dd, 1H, J = 4.8, 16.4 Hz), 2.34 (dd, 1H, J = 13.2, 16.4 Hz), 1.92-2.00 (m, 2H), 1.88 (m, 1H), 1.42-1.74 (m, 9 H), 1.28-1.35 (m, 2H), 1.21 (s, 3H), 1.20 (m, 1H), 1.15 (s, 3H), 1.14 (s, 3H), 1.12 (m, 1H), 1.06 (s, 3H), 0.97 (s, 6H), 0.93 (s, 3H); m/z 478.2 (M+1).

Compound 402-59: To a solution of **402-51** (2.08 g, 4.21 mmol) in CH₂Cl₂ (28 mL) were successively added oxalyl chloride (1.07 mL, 12.64 mmol) and DMF (5 drops, cat.) at 0 °C. The reaction was allowed to warm to room temperature and was stirred for 3 h. The reaction mixture was concentrated and dried *in vacuo* 30 min, to give acid chloride **3** as a yellow solid, which was used directly in the next step. To a solution of **3** (2.16 g, 4.21 mmol) in THF (28 mL) at 0 °C was added ammonia (2.0 M solution in MeOH, 11 mL, 22.00 mmol). The reaction was allowed to warm to room temperature and was stirred for 5 h. The solvents were then evaporated, and the residue was extracted with EtOAc. The extracts were washed with water, 1 N HCl (aq), and water, then dried over MgSO₄, filtered, and concentrated to give **402-59** (2.06 g, 99%) as a pale yellow solid. A small amount (53 mg) was purified by column chromatography (silica gel, 0% to 25% EtOAc in CH₂Cl₂) to give higher purity **402-59** (14 mg, white solid) for biological assay: ¹H NMR (400 MHz, CDCl₃) δ 7.65 (s, 1H), 5.63 (br s, 1H), 5.36 (br s, 1H), 2.90 (br d, 1H, J = 5.2 Hz), 2.71 (br d, 1H, J = 12 Hz), 2.42 (m, 2H), 1.96-2.10 (m, 4H), 1.78-1.90 (m, 2H), 1.45-1.69 (m, 6H), 1.23-1.40 (m, 4H), 1.22 (s, 3H), 1.16 (s, 3H), 1.15 (s, 3H), 1.13 (s, 3H), 0.99 (s, 3H), 0.98 (s, 3H), 0.93 (s, 3H); m/z 493.3 (M+1)

Compound 402-57: A solution of **402-59** (2.01 g, 4.08 mmol) in CH₂Cl₂ (28 mL) was prepared and cooled to 0 °C. To this solution were added TFAA (0.91 mL, 6.55 mmol) and Et₃N (1.48 mL, 10.62 mmol). The reaction was stirred at 0 °C for 3 h, after which it was

quenched by the addition of saturated NaHCO_3 (aq) solution (40 mL). After stirring for 10 min, the reaction mixture was extracted with CH_2Cl_2 and washed with saturated NaHCO_3 (aq), water, 1 N HCl (aq), and water. The extracts were dried over MgSO_4 , filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 5% to 35% EtOAc in hexanes). The purified product was triturated with EtOH, then filtered and dried on the filter to give **402-57** (0.87 g, 45%) as a powdery white solid: ^1H NMR (400 MHz, CDCl_3) δ 7.63 (s, 1H), 3.04 (d, 1H, $J = 4.4$ Hz), 2.38-2.57 (m, 3H), 1.91-2.19 (m, 5H), 1.61-1.78 (m, 4H), 1.44-1.54 (m, 3H), 1.32 (s, 3H), 1.26-1.30 (m, 4H), 1.22 (s, 3H), 1.19 (s, 3H), 1.16 (s, 3H), 0.99 (s, 3H), 0.95 (s, 3H), 0.92 (s, 3H); m/z 475.2 (M+1).

Compound 404-02: To a solution of **3** (3.03 g, 5.92 mmol) in toluene (84 mL) was added NaHCO_3 (1.98 g). A solution of trifluoroethylamine hydrochloride (5.64 g, 41.62 mmol) in water (14 mL) was prepared, then added to the reaction. The reaction was heated to 70 °C and stirred for 2 h. After cooling to room temperature, the reaction mixture was extracted with EtOAc and washed with brine. The combined extracts were dried over MgSO_4 , filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 40% EtOAc in CH_2Cl_2) to give **404-02** (2.35 g, 69%) as a white solid: ^1H NMR (400 MHz, CDCl_3) δ 7.65 (s, 1H), 5.94 (br t, 1H, $J = 8$ Hz), 4.10 (m, 1H), 3.69-3.88 (m, 1H), 2.84 (d, 1H, $J = 8$ Hz), 2.78 (br d, 1H, $J = 16$ Hz), 2.38 (m, 2H), 2.12 (m, 1H), 2.06 (m, 2H), 1.61-1.83 (m, 5H), 1.24-1.52 (m, 8H), 1.22 (s, 3H), 1.16 (s, 3H), 1.14 (s, 3H), 1.07 (s, 3H), 0.99 (s, 6H), 0.93 (s, 3H); m/z 575.3 (M+1).

Compounds 4, 5: To a solution of **402-49** (395 mg, 0.775 mmol) in THF (7.8 mL) was added LiAlH_4 (1.0 M solution in THF, 0.78 mL, 0.780 mmol) at 0 °C. The reaction was stirred at 0 °C for 40 min, after which it was quenched by the addition of water (5 mL) and stirred 5 min. The reaction mixture was extracted with EtOAc and washed with water. Solid NaCl was added to break up emulsions. The combined extracts were dried over Na_2SO_4 , filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 10% to 70% EtOAc in hexanes) to give both **4** (151 mg, 38%) as a white solid and to give **5** (134 mg, 34%) as a white solid:

Compound 4: ^1H NMR (400 MHz, CDCl_3) δ 7.99 (s, 1H), 3.79 (m, 1H), 3.72 (s, 3H), 2.75 (m, 1H), 2.52 (d, 1H, $J = 14.4$ Hz), 1.95-2.11 (m, 2H), 1.57-1.88 (m, yyH), 1.24-1.54 (m, yyH), 1.30 (s, 3H), 1.20 (s, 3H), 1.00 (s, 3H), 0.93 (s, 6H), 0.92 (s, 3H), 0.82 (s, 3H); m/z 512.3 (M+1).

Compound 5: m/z 494.3 (M-17), 434.3 (M-17-60).

Compound 6: To a solution of **4** (371 mg, 66 μ mol) in MeOH (7.3 mL) was added NaOMe (25 wt % solution in MeOH, 0.42 mL, 1.837 mmol) at room temperature. The reaction was heated to 55 °C and stirred 7 h. After cooling the reaction to room temperature, the reaction mixture was diluted with MTBE (10 mL), then quenched with 1 N HCl (aq) (10 mL). The reaction mixture was extracted with EtOAc and washed with 1 N HCl (aq) and brine. The combined extracts were dried over Na₂SO₄, filtered, and concentrated to give **6** (361 mg, 94%) as a white solid. Compound **6** is a mixture of two equilibrium forms, the enol form (as shown in Scheme 5) and the ketone form, in the ratio of 2:3. ¹H NMR of the mixture: (400 MHz, CDCl₃) δ 5.66 (d, 0.4H, *J* = 4.8 Hz), 4.09 (br, 1H), 3.90 (m, 0.6H), 3.71 (s, 1.2H), 3.68 (s, 1.8H), 2.73 (m, 1H), 2.48 (m, 1H), 2.14-2.26 (m, 2H), 0.80-2.02 (m, 39H); m/z 494.3 (M-17), 434.3 (M-77).

Compound 402-66: A solution of **6** (361 mg, 0.705 mmol) in DMF (7.1 mL) was prepared. 1,3-Dibromo-5,5-dimethylhydantoin (120 mg, 0.420 mmol) was added, and the reaction was stirred at room temperature for 1 h. Pyridine (0.23 mL, 2.858 mmol) was added, and the reaction was heated to 55 °C and stirred 10 h. After cooling the reaction to room temperature, the reaction mixture was extracted with EtOAc and washed with 5% Na₂S₂O₃ (aq), water, 1 N HCl (aq), and water. The EtOAc extracts were dried over Na₂SO₄, filtered, and evaporated. The crude product was purified by column chromatography (silica gel, 5% to 40% EtOAc in hexanes) to give **402-66** (127 mg, 35% yield) as a white solid; ¹H NMR (400 MHz, CDCl₃) δ 7.81 (s, 1H), 3.81 (ddd, 1H, *J* = 4.8, 10.8, 15.6 Hz), 3.72 (s, 3H), 2.75 (m, 1H), 2.01 (m, 2H), 1.74-1.88 (m, 3H), 1.24-1.72 (m, 15 H), 1.19 (s, 3H), 1.13 (s, 3H), 1.12 (s, 3H), 0.98 (s, 3H), 0.96 (s, 3H), 0.95 (s, 3H), 0.94 (s, 3H); m/z 492.3 (M-17), 432.3 (M-77).

Compound 7: Using the procedure described for the synthesis of compound **6** from compound **4**, compound **7** (96 mg, 89% yield) was produced from compound **5** (108 mg, 0.211 mmol): m/z 494.3 (M-17).

Compound 63219: Using the procedure described for the synthesis of compound **402-66** from compound **6**, compound **63219** (30 mg, 32% yield) was produced from compound **7** (95 mg, 0.186 mmol): ¹H NMR (400 MHz, CDCl₃) δ 7.81 (1H, s), 4.16 (1H, bs), 3.68 (3H, s), 2.44-2.54 (2H, m), 1.98-2.10 (2H, m), 1.78-1.94 (4H, m), 1.42-1.76 (7H, m), 1.00-1.42 (6H, m), 1.32 (3H, s), 1.23 (3H, s), 1.13 (3H, s), 1.11 (3H, s), 0.94 (3H, s), 0.93 (3H, s), 0.92 (3H, s); m/z 492.3 (M-17).

Compound 8: Oxalyl chloride (0.11 mL, 1.30 mmol) and catalytic amount of DMF were added sequentially to a solution of compound **402-51** (200 mg, 0.41 mmol) in CH₂Cl₂ (4 mL) at 0 °C. The reaction mixture was warmed to room temperature and stirred for 2 h. After removing the solvent by evaporation, the crude acid chloride was obtained as a light yellow foam solid. Hydrazine hydrate (64% of hydrazine, 0.50 mL) was added to a solution of acid chloride in Et₂O (8 mL) at 0 °C. After stirring for 30 min, CH₂Cl₂ was added. The mixture was washed with water, dried over MgSO₄, filtered and evaporated to give compound **8** (200 mg, 97% yield) as white solid, which was used in the next step without further purification: m/z 508.3 (M+1).

Compound 9: Et₃N (0.12 mL, 0.86 mmol) and acetyl chloride (37 µL, 0.52 mmol) were added sequentially to a solution of compound **8** (200 mg, 0.39 mmol) in CH₂Cl₂ (4 mL) at r.t. After stirring for 30 min, Et₃N (0.36 mL, 2.59 mmol) and acetyl chloride (110 µL, 1.55 mmol) were added again. After stirring for another 30 min, NaHCO₃ (aq.) solution was added to quench the reaction. The reaction mixture was transferred to a separatory funnel, and extracted with EtOAc. The combined extracts were washed with water, dried over MgSO₄, filtered and evaporated. The residue was purified by silica gel chromatography (0% to 75% EtOAc in hexanes) to give compound **9** (180 mg, 77% yield) as a white foam solid: m/z 592.3 (M+1).

Compound 63264: NaOMe (25% w/w in MeOH, 0.14 mL, 0.61 mmol) was added to a solution of compound **9** (180 mg, 0.30 mmol) in MeOH (3 mL) at 0 °C. After stirring at r.t. for 10 min, the reaction mixture was treated with t-BuOMe (10 mL) and 1N HCl (aq.) (1 mL), which was then transferred to a separatory funnel and extracted with EtOAc. The combined extracts were washed with NaHCO₃ (aq.) solution, dried over MgSO₄, filtered and evaporated. The residue was purified by silica gel chromatography (0% to 100% EtOAc in hexanes) to give compound **63264** (121 mg, 72% yield) as a white solid: ¹H NMR (400 MHz, CDCl₃) δ 7.98 (d, 1H, *J* = 4.4 Hz), 7.77 (d, 1H, *J* = 4.4 Hz), 7.65 (s, 1H), 2.89 (d, 1H, *J* = 4.4 Hz), 2.82 (m, 1H), 2.34-2.45 (m, 2H), 2.10 (m, 1H), 2.08 (s, 3H), 1.82-2.02 (m, 4H), 1.60-1.69 (m, 3H), 1.44-1.53 (m, 4H), 1.16-1.40 (m, 4H), 1.22 (s, 3H), 1.16 (s, 3H), 1.15 (s, 3H), 1.13 (s, 3H), 0.99 (s, 3H), 0.98 (s, 3H), 0.94 (s, 3H); m/z 550.3 (M+1).

Compound 63267: A solution of compound **63264** (74 mg, 0.13 mmol), TsOH (13 mg, 0.068 mmol) in toluene (5 mL) was heated at reflux with dean-stark trap for 1 h. After cooling to r.t., the reaction mixture was transferred to a separatory funnel, washed with NaHCO₃ (aq.) solution, dried over MgSO₄, filtered and evaporated. The residue was purified

by silica gel chromatography (0% to 100% EtOAc in hexanes) to give compound **63267** (24 mg, 33% yield) as a white foam solid: ^1H NMR (400 MHz, CDCl_3) δ 7.64 (s, 1H), 2.94 (m, 1H), 2.79 (d, 1H, $J = 4.4$ Hz), 2.54 (s, 3H), 2.46 (m, 1H), 2.34 (m, 1H), 2.21 (m, 1H), 1.84-2.06 (m, 5H), 1.56-1.70 (m, 4H), 1.24-1.47 (m, 5H), 1.23 (s, 3H), 1.18 (m, 1H), 1.15 (s, 3H), 1.13 (s, 3H), 1.05 (s, 3H), 1.02 (s, 3H), 0.98 (s, 3H), 0.93 (s, 3H); m/z 532.3 ($M+1$).

Compound 11: Et_3N (1.46 mL, 10.49 mmol) and TFAA (0.88 mL, 6.33 mmol) were added sequentially to a solution of compound **10** (1.97 g, 4.20 mmol) in CH_2Cl_2 (42 mL) at 0 °C. After stirring for 1.5 h, NaHCO_3 (aq.) solution was added to the reaction mixture, which was then transferred to a separatory funnel and extracted with CH_2Cl_2 . The combined extracts were dried over MgSO_4 , filtered and evaporated. The residue was purified by silica gel chromatography (0% to 35% EtOAc in hexanes) to give compound **11** (1.62 g, 85% yield) as a white solid: m/z 452.3.

Compound 12: A solution of Bu_3SnN_3 (1.00 mL, 3.62 mmol) and compound **11** (1.36 g, 3.02 mmol) in xylene (5.0 mL) was heated at reflux for 48 h. After cooling to r.t., the reaction mixture was purified by silica gel chromatography (0% to 30% EtOAc in CH_2Cl_2) to give compound **12** (994 mg, 67% yield) as a light yellow foam solid: m/z 493.3 ($M+1$).

Compound 13: NaOMe solution (25% w/w in MeOH, 1.16 mL, 5.07 mmol) was added dropwise to a mixture of compound **12** (168 mg, 0.34 mmol) and HCO_2Et (0.82 mL, 10.19 mmol) at 0 °C under N_2 . After stirring at room temperature for 1 h, $t\text{-BuOMe}$ (10 mL) was added. The mixture was cooled to 0 °C, and 12 N HCl (aq) (0.42 mL, 5.04 mmol) was added slowly. The mixture was transferred to a separatory funnel and extracted with EtOAc. The combined extracts were washed with water, dried over MgSO_4 and concentrated to give crude 2-formyl ketone, which was then mixed with $\text{NH}_2\text{OH}\cdot\text{HCl}$ (36 mg, 0.51 mmol), EtOH (4 mL), and water (0.4 mL), and heated at 60 °C for 3 h. After removing EtOH by evaporation, the white slurry obtained was transferred to a separatory funnel and extracted with CH_2Cl_2 . The combined extracts were washed with water, dried over MgSO_4 , and concentrated. The residue was purified by column chromatography (silica gel, 0% to 30% EtOAc in CH_2Cl_2) to give compound **13** (95 mg, 54% yield) as a white foam solid: m/z 520.3 ($M+1$).

Compound 63229: Using the procedure described for the synthesis of compound **402-66** from compound **4**, **63229** (12 mg, 60% yield) was produced from compound **13** (20 mg, 0.038 mmol): ^1H NMR (400 MHz, CDCl_3) δ 7.70 (s, 1H), 3.03 (m, 1H), 2.69 (d, 1H, $J = 4.0$ Hz), 2.52 (dd, 1H, $J = 4.4, 16.8$ Hz), 2.29-2.36 (m, 2H), 1.96-2.03 (m, 3H), 1.56-1.82 (m,

6H), 1.25-1.57 (m, 6H), 1.22 (s, 3H), 1.18 (m, 1H), 1.13 (s, 3H), 1.11 (s, 3H), 1.04 (s, 3H), 1.03 (s, 3H), 0.98 (s, 3H), 0.75 (s, 3H); m/z 518.3 (M+1).

Compound 14: TMSCHN₂ (2.0 M in Et₂O, 89 μ L, 0.18 mmol) was added to a solution of compound **13** (84 mg, 0.16 mmol) in THF (1.25 mL) and MeOH (0.31 mL) at 0 °C. After stirring at room temperature for 10 min, acetic acid was added to quench the reaction. The reaction mixture was diluted with EtOAc, transferred to a separatory funnel, washed with NaHCO₃ (aq.) solution, dried over MgSO₄ and evaporated. The residue was purified by column chromatography (silica gel, 0% to 60% EtOAc in hexanes) to give compound **14** (67 mg, 77% yield) as a white solid: m/z 534.3 (M+1).

Compound 63230: Using the procedure described for the synthesis of compound **402-66** from compound **4**, **63230** (45 mg, 69% yield) was produced from compound **14** (65 mg, 0.12 mmol): ¹H NMR (400 MHz, CDCl₃) δ 7.62 (s, 1H), 4.32 (s, 3H), 3.11 (m, 1H), 2.68 (d, 1H, *J* = 4.4 Hz), 2.42 (dd, 1H, *J* = 4.8, 16.4 Hz), 2.27 (dd, 1H, *J* = 13.2, 16.4 Hz), 2.22 (dd, 1H, *J* = 4.4, 14.8 Hz), 1.94-2.04 (m, 3H), 1.79 (m, 1H), 1.54-1.63 (m, 5H), 1.36-1.50 (m, 4H), 1.26 (m, 1H), 1.20 (s, 3H), 1.13 (m, 1H), 1.12 (s, 3H), 1.09 (s, 3H), 1.05 (s, 3H), 1.01 (s, 3H), 0.97 (s, 3H), 0.70 (s, 3H); m/z 532.3 (M+1).

Compound 63223: Using the procedure described for the synthesis of compound **13** from compound **12**, compound **63223** (1.95 g, 47% yield) was produced from compound **15** (3.93 g, 7.12 mmol) as a pale yellow solid: ¹H NMR (400 MHz, CDCl₃) δ 7.98 (1H, s), 5.91 (1H, t, *J* = 6.0 Hz), 4.00-4.15 (1H, m), 3.55-3.90 (1H, m), 2.72-2.82 (2H, m), 2.20-2.40 (3H, m), 1.88-2.16 (4H, m), 1.10-1.84 (13H, m), 1.31 (3H, s), 1.21 (3H, s), 1.00 (3H, s), 0.98 (6H, s), 0.91 (3H, s), 0.82 (3H, s); m/z 577.3 (M+H).

Compound 63227: Using the procedure described for the synthesis of compound **6** from compound **4**, compound **63227** (1.64 g, quantitative yield) was produced from compound **63223** (1.61 g, 2.79 mmol): ¹H NMR (400 MHz, CDCl₃) for enol form: δ 5.91 (1H, t, *J* = 6.0 Hz), 5.78 (1H, bs, enol), 4.00-4.16 (1H, m), 3.75-3.94 (1H, m), 2.70-2.85 (2H, m), 1.90-2.30 (5H, m), 0.80-1.88 (36H, m); m/z 577.3 (M+H) (for both enol and ketone isomers).

Compound 63237: A solution of **63227** (1.61 g, 2.79 mmol) in DMF (9.3 mL) was prepared. 1,3-Dibromo-5,5-dimethylhydantoin (456 mg, 1.59 mmol) was added, and the reaction was stirred at room temperature for 3 h. Pyridine (0.67 mL, 8.33 mmol) was added, and the reaction was heated to 55 °C and stirred 16 h. After cooling the reaction to room temperature, the reaction mixture was extracted with EtOAc and washed with 5% Na₂S₂O₃

(aq), 1 N HCl (aq), and water. The EtOAc extracts were dried over Na₂SO₄, filtered, and evaporated. Compound **63237** was a minor component (18%) by crude LC-MS analysis. The crude product was purified by column chromatography (silica gel, 5% to 35% EtOAc in hexanes) to give **63237** (188 mg, 12%) as a yellow foam solid: ¹H NMR (400 MHz, CDCl₃) δ 7.67 (s, 1H), 3.82 (m, 2H), 2.91 (m, 1H), 2.54 (m, 1H), 2.41 (m, 1H), 2.08 (m, 1H), 1.83-1.94 (m, 2H), 1.52-1.74 (m, 6H), 1.39 (s, 3H), 1.22 (s, 6H), 1.20-1.49 (m, 7H), 1.16 (s, 3H), 0.98 (s, 3H), 0.96 (s, 3H), 0.95 (s, 3H); m/z 573.3 (M+1).

Compound 16: Using the procedure described for the synthesis of compound **4** from compound **402-49**, compound **16** (100 mg, 19% yield) was produced from compound **63223** (531 mg, 0.921 mmol): m/z 579.3 (M+1)

Compound 17: Using the procedure described for the synthesis of compound **6** from compound **4**, compound **17** (90 mg, 92% yield) was produced from compound **16** (98 mg, 0.169 mmol): m/z 561.3 (M-17).

Compound 63268: Using the procedure described for the synthesis of compound **402-66**, compound **63268** (50 mg, 56% yield) was produced from compound **17** (90 mg, 0.156 mmol): ¹H NMR (400 MHz, CDCl₃) δ 7.78 (s, 1H), 6.09 (t, 1H, *J* = 6.4 Hz), 4.06 (m, 1H), 3.89 (m, 1H), 3.78 (m, 1H), 2.70 (m, 1H), 1.98-2.05 (m, 2H), 1.84 (dd, 1H, *J* = 4.4, 10.8 Hz), 1.75 (ddd, 1H, *J* = 4.4, 13.6, 13.6 Hz), 1.20-1.62 (m, 15H), 1.18 (s, 3H), 1.11 (s, 3H), 1.10 (s, 3H), 1.06 (m, 1H), 0.98 (s, 3H), 0.95 (s, 3H), 0.94 (s, 3H), 0.91 (s, 3H); m/z 559.3 (M-17).

Compound 19: NaOMe (25 w/w% solution in MeOH, 7.29 mL, 31.88 mmol) was added to a solution of compound **18** (1.00 g, 2.12 mmol) in ethyl formate (5.13 mL, 63.78 mmol) at 0 °C. After stirring for 1.5 h, *t*-BuOMe (10 mL) and 12 N (aq.) HCl (2.66 mL, 31.92 mmol) were added sequentially. After stirring for another 5 min, the reaction mixture was transferred to a separatory funnel, which was extracted with EtOAc. The combined extracts were washed with water. The organic layer was separated, which was dried over MgSO₄, filtered, and concentrated. The crude product was mixed with NH₂OH-HCl (0.22 g, 3.17 mmol), water (2 mL) and EtOH (35 mL). The reaction mixture was heated at 65 °C for 3.5 h, after which EtOH was removed by evaporation. The residue was partitioned between EtOAc and water. The organic extract was separated, which was dried over MgSO₄, filtered, and evaporated. The residue was purified by silica gel chromatography (0% to 60% EtOAc in hexanes) to give compound **19** (820 mg, 78% yield) as a white solid: m/z 496.3 (M+1).

Compound 20: Oxalyl chloride (110 μ L, 1.30 mmol) was added to a solution of compound **19** (195 mg, 0.39 mmol) in CH_2Cl_2 (4 mL) at 0 $^\circ\text{C}$, followed by the addition of catalytic amount of DMF. The reaction was stirred at room temperature for 2 h, after which CH_2Cl_2 was evaporated under vacuum to give acid chloride as a light yellow foam solid.

5 Et_3N (113 μ L, 0.81 mmol) and a solution of acethydrazide (50 mg, 0.67 mmol) in CH_2Cl_2 (2 mL) were added sequentially to a suspension of the acid chloride in ether (4 mL) at 0 $^\circ\text{C}$. The reaction was warmed to room temperature and stirred for 30 min. EtOAc was then added, and the crude mixture was transferred to a separatory funnel, which was washed with water, 1N (aq.) HCl, water. The organic layer was separated, which was dried over MgSO_4 , filtered,
10 and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 100% EtOAc in hexanes) to give product **20** (215 mg, 99% yield) as a white foam solid: ^1H NMR (400 MHz, CDCl_3) δ 8.14 (d, 1H, $J = 5.2$ Hz), 8.05 (d, 1H, $J = 5.2$ Hz), 8.00 (s, 1H), 2.86 (m, 2H), 2.34 (m, 3H), 2.09 (s, 3H), 1.80-2.18 (m, 8H), 1.34-1.74 (m, 7H), 1.33 (s, 3H), 1.23 (s, 3H), 1.16-1.26 (m, 2H), 1.08 (s, 3H), 1.00 (s, 6H), 0.93 (s, 3H), 0.84 (s, 3H).

15 **Compound 21 and Compound 22:** A suspension of compound **20** (215 mg, 0.39 mmol) and Lawesson's reagent (190 mg, 0.47 mmol) in toluene was heated at reflux for 30 min. After cooling to room temperature, the reaction mixture was purified by column chromatography (silica gel, 0% to 65% EtOAc in hexanes) to give product **21** (21 mg, 10% yield) as a light yellow foam solid: m/z 550.3 ($M+1$). From the column, compound **22** (60
20 mg, 29% yield) was also obtained as a white foam solid: m/z 534.3 ($M+1$).

Compound 23: NaOMe (25 w/w% solution in MeOH, 17 μ L, 0.074 mmol) was added to a solution of compound **21** (33 mg, 0.060 mmol) in MeOH (0.6 mL) at room temperature. The reaction was then heated to 55 $^\circ\text{C}$, and stirred for 1 h. After cooling to 0 $^\circ\text{C}$, $t\text{-BuOMe}$ and 1 N (aq.) HCl were added, and stirred for 5 min. The reaction mixture was transferred to
25 a separatory funnel, which was extracted with EtOAc. The combined EtOAc extracts were washed with water, dried over MgSO_4 , filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 45% EtOAc in hexanes) to give product **23** (24 mg, 73% yield) as a white foam solid: m/z 550.3 ($M+1$). Compound **23** is an isomeric mixture of ketone and enol forms.

30 **Compound 63274:** To a solution of compound **23** (23 mg, 0.041 mmol) in DMF (0.3 mL) was added 1,3-dibromo-5,5-dimethylhydantion (6.1 mg, 0.021 mmol) at 0 $^\circ\text{C}$, and the reaction was stirred at 0 $^\circ\text{C}$ for 1 h. Pyridine (14 μ L, 0.17 mmol) was then added, and the mixture was heated at 55 $^\circ\text{C}$ for 3 h. After cooling to room temperature, the reaction was diluted with EtOAc, and transferred to a separatory funnel, which was then washed with

Na₂SO₃ (aq.) solution and water. The organic layer was separated, which was dried over MgSO₄, filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 45% EtOAc in hexanes) to give product **63274** (18 mg, 79% yield) as a white foam solid: ¹H NMR (400 MHz, CDCl₃) δ 7.61 (s, 1H), 2.97 (d, 1H, *J* = 4.4 Hz), 2.89 (m, 1H), 2.74 (s, 3H), 2.42 (dd, 1H, *J* = 4.8, 16.4 Hz), 2.29 (dd, 1H, *J* = 13.6, 16.4 Hz), 2.29 (m, 1H), 2.02 (m, 1H), 1.94 (dd, 1H, *J* = 4.8, 13.2 Hz), 1.77-1.91 (m, 3H), 1.52-1.66 (m, 4H), 1.36-1.50 (m, 4H), 1.26 (m, 1H), 1.19 (s, 3H), 1.13 (m, 1H), 1.11 (s, 3H), 1.09 (s, 3H), 1.02 (s, 3H), 1.00 (s, 3H), 0.96 (s, 3H), 0.80 (s, 3H); *m/z* 548.3 (M+1).

Compound 24: LiAlH₄ solution (1.0 M in THF, 42 mL, 42 mmol) was added to a solution of compound **1** (5.0 g, 10.3 mmol) in THF (100 mL) at room temperature under N₂. After stirring for 20 min at room temperature, LiAlH₄ solution (1.0 M in THF, 21 mL, 21 mmol) was added again and the reaction mixture was refluxed for 1 h. After cooling to 0 °C, water (10 mL) was added dropwise, followed by the addition of 1N HCl (aq) (300 mL). The mixture was extracted with EtOAc. The combined extracts were washed with water, dried with MgSO₄, and concentrated. The residue obtained was mixed with CH₂Cl₂ (200 mL). The white solid that precipitated was collected by filtration and washed with CH₂Cl₂ (2×100mL) to give compound **24** (500 mg, 10%) as a white solid. The combined filtrate was loaded on a silica gel column and eluted with 0% to 100% EtOAc in hexanes to give additional compound **24** (800 mg, 17%) as a white solid: ¹H NMR (400 MHz, CDCl₃) δ 3.79 (m, 1H), 3.54 (m, 2H), 3.20 (dd, 1H, *J* = 4.8, 10.8 Hz), 1.98 (m, 1H), 1.12-1.88 (m, 23H), 1.03 (s, 3H), 0.98 (s, 6H), 0.91 (s, 3H), 0.86 (s, 3H), 0.85 (s, 3H), 0.77 (s, 3H), 0.65-1.10 (m, 3H); *m/z* 443.3 (M-H₂O+1), 425.3 (100%, M-2×H₂O+1).

Compound 25: TEMPO (27 mg × 4, 0.17 mmol × 4) and IPh(OAc)₂ (563 mg × 4, 1.74 mmol × 4) were added to a white slurry of compound **24** (725 mg, 1.59 mmol) in CH₂Cl₂ (200 mL) and water (0.1 mL) at 0 h, 2 h, 24 h and 48 h at room temperature. After stirring at room temperature for 72 h (overall reaction time), the reaction mixture turned into a clear pink solution, which was then transferred to a separatory funnel and washed with Na₂SO₃ (aq) solution. The organic phase was separated, dried over MgSO₄, filtered, and evaporated. The residue was purified by silica gel chromatography (0% to 75% EtOAc in hexanes) to give compound **25** (560 mg, 77%) as a white solid: ¹H NMR (400 MHz, CDCl₃) δ 9.37 (d, 1H, *J* = 1.2 Hz), 3.77 (m, 1H), 3.18 (dd, 1H, *J* = 4.8, 11.2 Hz), 2.51 (m, 1H), 0.98-1.87 (m, 23H), 0.97 (s, 3H), 0.96 (s, 3H), 0.94 (s, 3H), 0.92 (m, 1H), 0.90 (s, 3H), 0.86 (s, 3H), 0.82 (s, 3H), 0.75 (s, 3H), 0.65 (m, 1H); *m/z* 441.3 (M-H₂O+1), 423.3 (M-2×H₂O+1).

Compound 26: To a stirred suspension of $(\text{Ph}_3\text{PCH}_2\text{Cl})\text{Cl}$ (4.224 g, 12.1 mmol) in THF (13 mL) was added a solution of *n*-BuLi (4.8 mL, 11.64 mmol, 2.5 M in Hexanes) dropwise within 5 minutes at 0 °C, followed by the addition of HMPA (2.4 mL). The reaction was stirred at r.t. for 20 minutes and then Compound **25** (1.332 g, 2.90 mmol) in THF (13.0 mL) was added within 1 minute. The reaction mixture was stirred at room temperature for 2 h, then quenched with HCl (1N, 20 mL) and extracted with EtOAc (100 mL). The organic phase was washed by HCl (1N, 10 mL), NaCl (Sat., 20 mL), dried over Na_2SO_4 , filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 10% to 30% EtOAc in hexanes) to give compound **26** (1.2508 g, 87.8%, a mixture of E/Z isomers) as a white solid:

Compound 27: To a stirred solution of **26** (1.2508 g, 2.55 mmol) in THF (17 mL) was added a solution of MeLi (5.16 mL, 15.44 mmol, 3 M in $\text{CH}_2(\text{OEt})_2$) dropwise within 1 minute at 0 °C. The mixture was then stirred at room temperature for 28 h and quenched with HCl (1N, 15 mL). The aqueous solution was extracted with EtOAc (2×100 mL). The combined organic phase was washed with water, NaCl (sat.), dried over Na_2SO_4 , filtered, and concentrated to give compound **27** (1.0630 g, 91.7%) as a white solid: m/z 437.3 (M-OH).

Compound 28: To a stirred mixture of **27** (881.7 mg, 1.94 mmol), NaOAc (628.6 mg, 4 eq.) in CH_2Cl_2 (40 mL) was added PCC (1.257 g, 3 eq.) in one-portion at room temperature. The mixture was then stirred at room temperature for 5 h and diluted with a solvent mixture of EtOAc/Hexanes (1:1, 50 mL). The mixture was directly loaded on a silica gel pad, which was then eluted throughout with a solvent mixture of EtOAc/Hexanes (1:1). The eluate was collected and concentrated to give a colorless crystalline product. This crude product was purified by column chromatography (silica gel, 0% to 10% to 25% EtOAc in hexanes) to give compound **28** (685 mg, 78.8%) as a white solid: m/z 451.3 (M+1).

Compound 29: To a stirred suspension of **28** (22.0 mg, 0.0488 mmol) in HCO_2Et (0.118 mL, 1.46 mmol) was added a solution of MeONa (0.167 mL, 0.732 mmol, 25%w/w in MeOH) at 0 °C. The mixture was then stirred at room temperature for 25 h, diluted with TBME (1.4 mL) and quenched with HCl (0.126 mL, conc.) followed by water (3 mL). The aqueous solution was extracted with EtOAc (10 mL). The combined organic phase was washed with brine (5 mL), dried over Na_2SO_4 , filtered, and concentrated to give compound **29** as a light yellow foam, which was used directly in the next step.

Compound 30: Compound **29** was dissolved in EtOH (2.1 mL). To this solution were added $\text{NH}_2\text{OH}\cdot\text{HCl}$ (5.1 mg, 0.0732 mmol) and H_2O (0.27 mL) at room temperature. The mixture was heated at 60 °C for 18 h and then cooled to room temperature. The organic

volatiles were removed *in vacuo*. The remaining mixture was extracted with EtOAc (10 mL). The organic phase was washed with water, brine, dried over Na₂SO₄, filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 10% to 25% EtOAc in hexanes) to give **30** (20.8 mg, 89.6% from **6**) as a colorless crystalline solid: m/z 476.3 (M+1).

Compound 31: To a stirred suspension of **30** (20.8 mg, 0.0437 mmol) in a solvent mixture of MeOH (0.66 mL) and THF (0.11 mL) was added a solution of MeONa (23.8 µL, 0.105 mmol, 25%w/w in MeOH) at 55 °C. The mixture was then stirred at 55 °C for 3 h, cooled to room temperature and quenched with 1N HCl (aq) (5 mL). The mixture was extracted by EtOAc (15 mL). The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated to give compound **31** as a light yellow foam: m/z 476.3 (M-17).

Compound 63303: Compound **31** was dissolved in benzene (2 mL). To this solution was added a solution of DDQ (10.4 mg, 0.0458 mmol) in benzene (1 mL) at 85 °C. The mixture was stirred at 85 °C for 1.5 h, cooled to room temperature, and quenched with sat. NaHCO₃ (aq) (5 mL). The mixture was extracted with EtOAc (30 mL). The organic phase was washed with sat. NaHCO₃ (aq) and brine, then was dried over Na₂SO₄, filtered, and concentrated to give a solid residue (a mixture of starting material and desired product), which was then dissolved in pyridine (0.5 mL). To this solution were added Ac₂O (50 µL) and DMAP (cat.) at room temperature. The mixture was stirred at room temperature for 30 min and then quenched with NaHCO₃ (sat.). The mixture was extracted with EtOAc (20 mL). The organic phase was washed with NaHCO₃ (sat.), HCl (1N), brine, dried over Na₂SO₄, filtered, and concentrated to give a crude mixture, which was purified by column chromatography (silica gel, 0% to 10% to 25% EtOAc in hexanes) to give **63303** (8.1 mg, 39.1% from **30**) as a colorless solid: ¹H NMR (400 MHz, CDCl₃) δ 7.66 (s, 1H), 3.25 (d, 1H, *J* = 4.0 Hz), 2.25-2.52 (m, 3H), 2.22 (s, 1H), 1.78-2.15 (m, 5H), 1.44-1.76 (m, 9H), 1.08-1.36 (m, 2H), 1.29 (s, 3H), 1.23 (s, 3H), 1.19 (s, 3H), 1.17 (s, 3H), 0.89 (s, 3H), 0.94 (s, 3H), 0.91 (s, 3H); m/z 474.3 (M+1).

Compound 32: Compound **31** (95 mg, 0.2 mmol) was dissolved in a solvent mixture of acetone (3.5 mL) and water (1.5 mL). To this solution were added HgSO₄ (5.9 mg, 0.02 mmol) and H₂SO₄ (2 drops, conc.) at room temperature. The mixture was stirred at 55 °C for 20 h, cooled to room temperature, and quenched with water (20 mL) and 1 N HCl(aq) (10 mL). The mixture was extracted with EtOAc (30 mL). The organic phase was washed with 1 N HCl(aq), water, sat. NaHCO₃ (aq), brine, dried over Na₂SO₄, filtered, and concentrated to give a white solid, which was purified by column chromatography (silica gel, 0% to 10% to

25% EtOAc in hexanes) to give compound **32** (90.3 mg, 91.5%) as a white foam: m/z 494.3 (M+1).

Compound 63308: The procedure described for the synthesis of product **63274** from compound **23** was then employed to convert compound **32** into product **TX63308** (36.1 mg, 73.4%) as a white foam: ¹H NMR (400 MHz, CDCl₃) δ 7.65 (s, 1H), 2.75-2.85 (m, 1H), 2.67 (d, 1H, *J* = 4.4 Hz), 2.44 (dt, 1H, *J* = 16.4, 4.8 Hz), 2.35 (dt, 1H, *J* = 16.0, 13.2 Hz), 2.16 (s, 3H), 1.92-2.06 (m, 3H), 1.30-1.76 (m, 12H), 1.18-1.29 (m, 1H), 1.22 (s, 3H), 1.16 (s, 3H), 1.15 (s, 3H), 1.04 (s, 3H), 0.97 (s, 3H), 0.96 (s, 3H), 0.93 (s, 3H); m/z 492.3 (M+1).

Compound 63323: 1,8-Diazabicyclo[5,4,0]undec-7-ene (0.18 mL, 1.204 mmol) was added to a suspension of compound **402-51** (402 mg, 0.814 mmol) in toluene (5.4 mL) at room temperature. After stirring for 2 min, benzyl bromide (0.12 mL, 1.009 mmol) was added. The reaction mixture was heated at 100 °C for 6 h, after which it was cooled to room temperature. The reaction was then diluted with EtOAc, and was transferred to a separatory funnel, which was washed with 1 N HCl(aq) and brine. The organic extracts were separated, which was dried over Na₂SO₄, filtered and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 20% EtOAc in hexanes) to give product **63323** (308 mg, 65% yield) as a pale yellow foam solid: ¹H NMR (400 MHz, CDCl₃) δ 7.56 (s, 1H), 7.30-7.37 (m, 5H), 5.21 (d, 1H, *J* = 12.4 Hz), 5.07 (d, 1H, *J* = 12.4 Hz), 2.84 (m, 1H), 2.47 (d, 1H, *J* = 4.0 Hz), 2.36 (dd, 1H, *J* = 4.4, 16.0 Hz), 2.17 (dd, 1H, *J* = 13.6, 16.0 Hz), 1.81-1.92 (m, 4H), 1.20-1.72 (m, 12H), 1.19 (s, 3H), 1.12 (s, 3H), 1.07 (s, 3H), 0.99 (s, 3H), 0.90 (s, 6H), 0.65 (s, 3H); m/z 584.4 (M+1).

Compound 63325: MeONH₂-HCl (109 mg, 1.305 mmol), water (0.4 mL) and Et₃N (0.24 mL, 1.722 mmol) were added sequentially to a solution of compound **3** (439 mg, 0.857 mmol) in THF (4.2 mL) at room temperature. The reaction was then heated at 40 °C for 4 h. After cooling to room temperature, the reaction was diluted with EtOAc, and was transferred to a separatory funnel, which was washed with 1 N HCl(aq) and brine. The organic extracts were separated, which was dried over Na₂SO₄, filtered and concentrated. The crude product was purified by column chromatography (silica gel, 10% to 75% EtOAc in hexanes) to give product **63325** (165 mg, 37% yield) as a white solid: ¹H NMR (400 MHz, CDCl₃) δ 8.37 (s, 1H), 7.63 (s, 1H), 3.76 (s, 3H), 2.86 (d, 1H, *J* = 4.0 Hz), 2.71 (m, 1H), 2.44 (dd, 1H, *J* = 4.8, 16.4 Hz), 2.34 (dd, 1H, *J* = 13.2, 16.4 Hz), 1.91-2.08 (m, 3H), 1.74-1.90 (m, 2H), 1.59-1.68 (m, 3H), 1.40-1.50 (m, 4H), 1.21 (s, 3H), 1.18-1.38 (m, 4H), 1.15 (s, 3H), 1.14 (s, 3H), 1.12 (s, 3H), 0.97 (s, 3H), 0.96 (s, 3H), 0.91 (s, 3H); m/z 523.4 (M+1).

Compound 63326: Me₂NH (2.0 M solution in THF, 1.23 mL, 2.460 mmol) was added to a solution of compound **3** (408 mg, 0.797 mmol) in THF (4.1 mL) at room temperature. The reaction was then heated at 40 °C for 71 h. After cooling to room temperature, the reaction was diluted with EtOAc, and was transferred to a separatory funnel, which was washed with 1 N HCl(aq) and brine. The organic extracts were separated, which was dried over Na₂SO₄, filtered and concentrated. The crude product was purified by column chromatography (silica gel, 10% to 70% EtOAc in hexanes) to give product **63326** (254 mg, 61% yield) as a white solid: ¹H NMR (400 MHz, CDCl₃) δ 7.64 (s, 1H), 3.06 (s, 6H), 2.97 (m, 1H), 2.29-2.42 (m, 2H), 1.94-2.06 (m, 3H), 1.74-1.85 (m, 2H), 1.61-1.67 (m, 5H), 1.24-1.54 (m, 6H), 1.20 (s, 3H), 1.14 (s, 3H), 1.13 (s, 3H), 1.10 (m, 1H), 1.05 (s, 3H), 0.99 (s, 3H), 0.96 (s, 3H), 0.91 (s, 3H); m/z 521.4 (M+1).

Compound 33: NH₂OH-HCl (705 mg, 10.145 mmol), NaOAc (1.169 mg, 14.251 mmol) and water (3.3 mL) were added to a suspension of compound **402-49** (520 mg, 1.020 mmol) in EtOH (9.8 mL) at room temperature. The reaction mixture was heated at 80 °C for 27 h, after which it was cooled to room temperature. The reaction mixture was transferred to a separatory funnel, which was extracted with EtOAc. The combined organic extracts were washed with water and brine, then dried over Na₂SO₄, filtered and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 20% EtOAc in hexanes) to give product **33** (387 mg, 72% yield) as a white solid: m/z 525.3 (M+1).

Compound 34: NaOMe (25 w/w% solution in MeOH, 0.12 mL, 0.525 mmol) was added to a solution of compound **33** (128 mg, 0.244 mmol) in MeOH (1.2 mL) at room temperature. The reaction was then heated to 55 °C and stirred for 1 h. After cooling to room temperature, the reaction was diluted with *t*-BuOMe (3 mL) and was cooled to 0 °C. 1 N HCl (aq) (5 mL) was added. After stirring for another 5 min, the reaction mixture was transferred to a separatory funnel, which was extracted with EtOAc. The combined EtOAc extracts were washed with water, dried over Na₂SO₄, filtered, and concentrated to give product **34** (131 mg) as a white solid: m/z 525.3 (M+1). Compound **34** is an isomeric mixture of C3 ketone and enol forms.

Compound 63295: 1,3-Dibromo-5,5-dimethylhydantoin (40 mg, 0.140 mmol) in DMF (0.5 mL) was added to a solution of compound **34** (126 mg, 0.240 mmol) in DMF (1.6 mL) at 0 °C. After stirring at 0 °C for 40 min, the reaction was treated with pyridine (40 µL, 0.495 mmol), and was heated at 55 °C for 7 h. After cooling to room temperature, brine was added, and the reaction mixture was transferred to a separatory funnel, which was extracted with EtOAc. The combined organic extracts were washed with brine, 10% Na₂SO₃(aq)

solution, 1 N HCl(aq), and water. The organic layer was separated, which was dried over Na₂SO₄, filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 20% EtOAc in hexanes) to give product **63295** (34 mg, 27% yield from **33**) as a white solid: ¹H NMR (400 MHz, CDCl₃) δ 7.83 (s, 1H), 3.68 (s, 3H), 3.36 (dd, 1H, *J* = 16.8, 4.8 Hz), 2.82-2.91 (m, 1H), 2.54 (d, 1H, *J* = 3.6 Hz), 1.76-2.06 (m, 4H), 1.52-1.74 (m, 6H), 1.04-1.50 (m, 8H), 1.20 (s, 3H), 1.15 (s, 3H), 1.13 (s, 3H), 0.93 (s, 3H), 0.92 (s, 6H), 0.90 (s, 3H); *m/z* 523.3 (M+1);

Compound 35: POCl₃ (0.14 mL, 1.502 mmol) was added to a solution of compound **33** (200 mg, 0.381 mmol) in pyridine (1.9 mL) at room temperature. After stirring for 5 h, the reaction mixture was diluted with EtOAc (5 mL), and was quenched with 1 N HCl(aq) (5 mL). The reaction mixture was transferred to a separatory funnel, which was extracted with EtOAc. The combined organic extracts were washed with 1 N HCl(aq) and brine, then dried over Na₂SO₄, filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 50% EtOAc in hexanes) to give product **35** (185 mg, 75% yield) as a colorless glassy solid: *m/z* 525.4 (M+1);

Compound 36: NaOMe (25 w/w% solution in MeOH, 0.17 mL, 0.743 mmol) was added to a solution of compound **35** (177 mg, 0.337 mmol) in MeOH (1.7 mL) at room temperature. The reaction was then heated to 55 °C and stirred for 4 h. After cooling to 0 °C, *t*-BuOMe and 1 N HCl(aq) were added, and the reaction mixture was stirred for 5 min. The reaction mixture was then transferred to a separatory funnel, which was extracted with EtOAc. The combined EtOAc extracts were washed with 1 N HCl(aq) and brine, then dried over Na₂SO₄, filtered, and concentrated to give product **36** (164 mg, 93% yield) as a white solid: *m/z* 525.4 (M+1). Compound **36** is an isomeric mixture of C3 ketone and enol forms.

Compound 63296: 1,3-Dibromo-5,5-dimethylhydantion (53 mg, 0.185 mmol) in DMF (0.8 mL) was added to a solution of compound **36** (163 mg, 0.311 mmol) in DMF (2.1 mL) at 0 °C. After stirring at 0 °C for 1 h, the reaction was treated with pyridine (50 μL, 0.618 mmol) and was heated at 55 °C for 23 h. After cooling to room temperature, brine was added, and the reaction mixture was transferred to a separatory funnel, which was extracted with EtOAc. The combined organic extracts were washed with brine, 10% Na₂SO₃(aq) solution, 1 N HCl(aq), and water, then dried over Na₂SO₄, filtered, and concentrated to give product **63296** (150 mg, 93% yield) as a white solid: ¹H NMR (400 MHz, CDCl₃) δ 7.88 (s, 1H), 5.57 (d, 1H, *J* = 4.8 Hz), 4.06 (t, 1H, *J* = 6.0 Hz), 3.73 (s, 3H), 2.68 (dd, 1H, *J* = 14.4, 10.4 Hz), 2.48-2.60 (m, 1H), 2.13 (d, 1H, *J* = 14.4 Hz), 1.65-1.87 (m, 3H), 1.19-1.64 (m,

13H), 1.17 (s, 3H), 1.13 (s, 3H), 1.11 (s, 3H), 1.06 (s, 3H), 1.00 (s, 3H), 0.97 (s, 3H), 0.88 (s, 3H); m/z 523.3 (M+1).

Compound 37: *m*-CPBA (77%, 7.04 g, 31.52 mmol) was added to a solution of compound **402-49** (1.60 g, 3.15 mmol) in CH₂Cl₂ (28 mL) at room temperature. After stirring for 8 h, additional *m*-CPBA (77%, 3.52 g, 15.71 mmol) was added, and the reaction was stirred for another 40 h. Na₂SO₃ (aq.) solution was then added. After another 10 min, the reaction mixture was transferred to a separatory funnel, which was extracted with EtOAc. The combined EtOAc extracts were washed with NaHCO₃ (aq.) solution, dried over MgSO₄, filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 45% EtOAc in hexanes) to give product **37** (358 mg, 22% yield) as a white foam solid: m/z 526.3 (M+1).

Compound 38: NaOMe (25 w/w% solution in MeOH, 20 µL, 0.087 mmol) was added to a solution of compound **37** (38 mg, 0.072 mmol) in MeOH (0.7 mL) at room temperature. The reaction was then heated to 55 °C, and stirred for 2 h. After cooling to 0 °C, *t*-BuOMe and 1 N (aq.) HCl were added. The reaction mixture was then transferred to a separatory funnel, which was extracted with EtOAc. The combined EtOAc extracts were washed with water, dried over MgSO₄, filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 50% EtOAc in hexanes) to give product **38** (25 mg, 66% yield) as a white foam solid: m/z 526.4 (M+1). Compound **38** is an isomeric mixture of C3 ketone and enol forms.

Compound 63263: A solution of 1,3-dibromo-5,5-dimethylhydantion (6.9 mg, 0.024 mmol) in DMF (0.2 mL) was added to a solution of compound **38** (25 mg, 0.048 mmol) in DMF (0.8 mL) at 0 °C. After stirring at 0 °C for 1 h, the reaction was treated with pyridine (12 µL, 0.15 mmol), and was heated at 55 °C for 3 h. After cooling to room temperature, the reaction was diluted with EtOAc, and transferred to a separatory funnel, which was then washed with Na₂SO₃ (aq.) solution, 1N (aq.) HCl and water. The organic extract was separated, dried over MgSO₄, filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 45% EtOAc in hexanes) to give product **63263** (18 mg) as a white foam solid: ¹H NMR (400 MHz, CDCl₃) δ 7.83 (s, 1H), 4.95 (d, 1H, *J* = 7.2 Hz), 3.73 (s, 3H), 2.79 (m, 2H), 2.37 (d, 1H, *J* = 14.4 Hz), 1.92 (m, 2H), 1.77 (d, 1H, *J* = 10.4 Hz), 1.44-1.74 (m, 8H), 1.20-1.41 (m, 5H), 1.19 (s, 3H), 1.15 (s, 3H), 1.13 (s, 3H), 1.08 (s, 3H), 1.07 (s, 3H), 0.96 (s, 3H), 0.89 (s, 3H); m/z 524.3 (M+1).

Compound 39: LiAlH₄ (2.0 M in THF, 48 µL, 0.096 mmol) was added to a solution of compound **37** (50 mg, 0.095 mmol) in THF (0.95 mL) at 0 °C. After stirring for 40 min,

the reaction was quenched by adding water (1 mL) carefully. After stirring at room temperature for 10 min, the reaction mixture was transferred to a separatory funnel, which was extracted with EtOAc. The combined organic extracts were washed with 1N (aq.) HCl, and water, dried over MgSO₄, filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 40% EtOAc in hexanes) to give product **39** (31 mg, 62% yield) as a white foam solid: m/z 528.3 (M+1). The stereochemical configuration of C12 was not assigned.

Compound 40: NaOMe (25 w/w% solution in MeOH, 16 µL, 0.070 mmol) was added to a solution of compound **39** (30 mg, 0.057 mmol) in MeOH (0.6 mL) at room temperature. The reaction was then heated to 55 °C, and stirred for 2 h. After cooling to 0 °C, t-BuOMe and 1 N (aq.) HCl were added. The reaction mixture was then transferred to a separatory funnel, which was extracted with EtOAc. The combined EtOAc extracts were washed with water, dried over MgSO₄, filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 40% EtOAc in hexanes) to give product **40** (25 mg, 83% yield) as a white foam solid: m/z 510.3 (M-18+1). Compound **40** is an isomeric mixture of C3 ketone and enol forms. The stereochemical configuration of C12 was not assigned.

Compound 63289: 1,3-Dibromo-5,5-dimethylhydantion (6.8 mg, 0.024 mmol) was added to a solution of compound **40** (25 mg, 0.047 mmol) in DMF (0.47 mL) at 0 °C. After stirring at 0 °C for 1 h, the reaction was treated with pyridine (12 µL, 0.15 mmol), and was heated at 55 °C for 3 h. After cooling to room temperature, the reaction was diluted with EtOAc, and transferred to a separatory funnel, which was then washed with Na₂SO₃ (aq.) solution, 1N (aq.) HCl and water. The organic extract was separated, dried over MgSO₄, filtered, and concentrated. The crude product was purified by column chromatography (silica gel, 0% to 40% EtOAc in hexanes) to give partially purified product **63289** (16 mg), which was purified again by preparative TLC plate (silica gel, eluted with 8% EtOAc in hexanes) to give product **63289** (10 mg, 40% yield) as a white foam solid: ¹H NMR (400 MHz, CDCl₃) δ 7.64 (s, 1H), 5.12 (m, 1H), 4.33 (d, 1H, *J* = 6.8 Hz), 3.71 (s, 3H), 2.54 (m, 1H), 2.43 (d, 1H, *J* = 2.8 Hz), 1.21-1.96 (m, 18H), 1.19 (s, 6H), 1.13 (s, 3H), 1.05 (s, 3H), 1.02 (s, 3H), 0.95 (s, 3H), 0.88 (s, 3H); m/z 508.3 (M-18+1). The stereochemical configuration of C12 was not assigned.

Example 4 –Uptake of 404-02 into CNS and Lungs of Monkeys

Plasma concentrations following oral dosing: Compound 404-02 shows high uptake in the CNS and lung in monkeys following oral dosing: 2 male and 2 female cynomolgus

monkeys were administered 404-02 at 0.5, 5, 25 or 75 mg/kg/day doses via oral gavage. Doses were prepared in sesame oil and individualized to body weight on the day of dosing. Blood was drawn prior to dosing and at 1, 2, 4, 8 and 24 hours post-dose on days 1 and 12. Blood samples were collected from the femoral artery/vein for determination of 404-02 plasma concentrations. Blood was placed in tubes containing K3EDTA and stored on ice until centrifugation at room temperature. The isolated plasma was transferred to cryovials and stored at -80°C until sample processing and LC-MS/MS analysis. Extracted plasma standard curves were prepared from fresh stock solutions and analyzed prior to study samples. Summary results are shown in Tables 2a and 2b.

Population mean pharmacokinetic parameter estimates were obtained by performing non-compartmental analysis of the 404-02 plasma concentration-versus-time data using WinNonlinTM software version 5.2. Across the investigated dose range, 404-02 demonstrated dose-dependent kinetics with increased oral clearance (Cl/F), reciprocal shortening of elimination half-life ($T_{1/2}$) and an increasing apparent volume of distribution (V_z/F) with increasing dose. A 1.6-fold increase in the area under the concentration versus time curve over 24 hours (AUC_{0-24hr}) was observed at the 75 mg/kg dose level after 12 days of dosing compared to the corresponding AUC on day 1. Accumulation was not observed at any of the other dose levels. The observed mean maximum plasma concentration (C_{max}) for the 0.5, 5, 25 and 75 mg/kg/day dose groups on day 12 were 4.6, 12.7, 17.5 and 48.6 nM 404-02, respectively.

Table 2a shows the population mean plasma pharmacokinetics of 404-02 in cynomolgus monkey on day 1 of study (n=4). Pharmacokinetic parameters obtained using non-compartmental analysis, WinNonlinTM version 5.2.

Table 2a: Day 1 Plasma Pharmacokinetics of 404-02:

Dose (mg/kg/d)	Mean Maximum 404-02 Plasma Concentration (ng/mL) $\pm$ SEM	Mean Maximum 404-02 Plasma Concentration (nM) $\pm$ SEM
0.5	1.95 $\pm$ 0.35	3.4 $\pm$ 0.61
5	10.1 $\pm$ 3.8	17.6 $\pm$ 6.6
25	16.8 $\pm$ 1.3	29.3 $\pm$ 2.3
75	19.4 $\pm$ 2.7	33.8 $\pm$ 4.7

Table 2b shows the population mean plasma pharmacokinetics of 404-02 in cynomolgus monkey on day 12 of study (n=4). Pharmacokinetic parameters were obtained using non-compartmental analysis using WinNonlin™ version 5.2 software.

Table 2b: Day 12 Plasma Pharmacokinetics of 404-02:

Dose (mg/kg/d)	Mean Maximum 404-02 Plasma Concentration (ng/mL) $\pm$ SEM	Mean Maximum 404-02 Plasma Concentration (nM) $\pm$ SEM
0.5	2.65 $\pm$ 0.6	4.6 $\pm$ 1.0
5	7.3 $\pm$ 1.7	12.7 $\pm$ 2.9
25	10.0 $\pm$ 3.7	17.5 $\pm$ 6.4
75	28.0 $\pm$ 12.2	48.6 $\pm$ 12.2

5 **CNS and lung concentrations following oral dosing:** 2 male and 2 female cynomolgus monkeys were administered 404-02 at 0.5, 5, 25 or 75 mg/kg/day doses via oral gavage, in addition to a control (non-treatment) group with 2 animals per sex. Animals were sacrificed approximately 3 hours after dosing on day 15 following harvest of brain and lung tissues. Each sample collected was rinsed in 1 $\times$ isotonic phosphate buffered saline and
10 blotted dry before weighing. Harvested tissue slices were transferred to cryovials and stored at -80°C until processing and LC-MS/MS analysis. Standard curves were derived for 404-02 in homogenates of these tissues and were used to quantify the day 15 samples.

 The concentration of 404-02 necessary for 50% suppression of nitric oxide (NO) production in macrophages stimulated with interferon-gamma is approximately 45 nM. As
15 evidenced by the data, 0.5 mg/kg administered orally as the lowest dose in this study for 15 days resulted in a mean 404-02 CNS concentration of 2,162 nM, which markedly exceeds the IC₅₀ value for NO production *in vitro*. The CNS penetration provides a large therapeutic margin at all doses tested (Table 2c). For instance, the mean 404-02 CNS concentrations following administration of 0.5, 5, 25 and 75 mg/kg/day 404-02 indicates a 48-, 44-, 37- and
20 75-fold excess compared to the dose necessary to suppress inflammation *in vitro*. For comparison, the mean 404-02 lung tissue exposure at 75 mg/kg/day 404-02 showed a 133-fold increase (Table 2d). Non-linear disposition of 404-02 in CNS and lung tissue was observed suggesting 404-02 is taken up by a saturable mechanism(s) for transport across membranes and/or intracellular binding. In addition, concentrations of 404-02 in monkey
25 CNS and lung tissue exceed plasma levels. Table 2c shows the population mean 404-02

cynomolgus monkey CNS tissue exposure on day 15. Table 2d shows the population mean 404-02 cynomolgus monkey lung tissue content on day 15.

Table 2c: Day 15 CNS Tissue Content/Concentration of 404-02:

Dose (mg/kg/day)	Mean 404-02 CNS Tissue Content (ng/g) $\pm$ SEM	Mean 404-02 CNS Tissue Concentration (nM)* $\pm$ SEM
0.5	1198 $\pm$ 712	2087 $\pm$ 1239
5	1090 $\pm$ 564	1900 $\pm$ 982
25	930 $\pm$ 352	1618 $\pm$ 612
75	1898 $\pm$ 496	3305 $\pm$ 864

* Conversion based on the assumption the density of tissue is equal to water, 1 g/mL.

5 Table 2d: Day 15 Lung Tissue Content/Concentration of 404-02:

Dose Level (mg/kg/day)	Mean 404-02 Lung Tissue Content (ng/g) $\pm$ SEM	Mean 404-02 Lung Tissue Concentration (nM)* $\pm$ SEM
0.5	705 $\pm$ 292	1227 $\pm$ 508
5	24 $\pm$ 15	55 $\pm$ 27
25	141 $\pm$ 50	246 $\pm$ 87
75	3405 $\pm$ 824	5929 $\pm$ 1435

Example 5 –Uptake of 404-02 into CNS and Lungs of Rats

Compound 404-02 reaches high concentrations in the lung and CNS of rats following oral dosing: To assess basic pharmacokinetic parameters following oral dosing, 9 male and 9 female Sprague Dawley (SD) rats were administered 404-02 at 1, 10 or 50 mg/kg doses via oral gavage. Doses were prepared in sesame oil and individualized to body weight on the day of dosing. Blood was drawn at 0, 1, 2, 4, 8 and 24 hours post-dose on days 1 and 15. Blood was collected from the orbital sinus after carbon dioxide/oxygen inhalation for determination of 404-02 plasma concentrations. Plasma was transferred to cryovials and stored at -80°C until processing and LC-MS/MS analysis. Summary results for Day 1 and Day 15 are shown in Tables 2a and 2b, respectively. A standard curve was derived for 404-02 in rat plasma, and quantification of experimental results was based on this standard curve.

Table 3a shows population mean plasma pharmacokinetics of 404-02 in SD rats on day 1 of study (n=9/sex/dose level). Pharmacokinetic parameters were obtained using non-compartmental analysis, WinNonlin™ version 5.2.

Table 3a: Day 1 Pharmacokinetic Parameters

Dose (mg/kg/day)	Mean Maximum 404-02 Plasma Concentration (ng/mL) $\pm$ SEM	Mean Maximum 404-02 Plasma Concentration (nM) $\pm$ SEM
1	3.3 $\pm$ 0.65	5.7 $\pm$ 1.1
10	66 $\pm$ 16.5	116 $\pm$ 28.7
50	713 $\pm$ 98.2	1241 $\pm$ 171

5 Table 3b shows the population mean plasma pharmacokinetics of 404-02 in SD rats on day 15 of study (n=9/sex/dose level). Pharmacokinetic parameters were obtained using non-compartmental analysis, WinNonlin™ version 5.2.

Table 3b: Day 15 Pharmacokinetic Parameters:

Dose (mg/kg/day)	Mean Maximum 404-02 Plasma Concentration (ng/mL) $\pm$ SEM	Mean Maximum 404-02 Plasma Concentration (nM) $\pm$ SEM
1	6.3 $\pm$ 0.9	11.0 $\pm$ 1.6
10	144 $\pm$ 24.3	210 $\pm$ 42.3
50	1129 $\pm$ 114	1966 $\pm$ 199

To examine tissue concentrations following oral dosing, 5 male and 5 female Sprague
 10 Dawley (SD) rats were administered 404-02 at 1, 10, 50 or 150 mg/kg/day via oral gavage. Doses were prepared in sesame oil and individualized to body weight on the day of dosing. Animals were sacrificed 3 hours post-dose on day 15 of study following harvest of brain and lung specimens. Each sample collected was rinsed in 1x isotonic phosphate buffered saline and blotted dry before weighing. Tissue slices were transferred to cryovials and stored at -
 15 80°C until processing and LC-MS/MS analysis. Summary results for CNS and lung samples are shown in Tables 3c and 3d, respectively. Standard curves were derived for 404-02 in these tissues.

Table 3c: Population Mean 404-02 CNS Tissue Content in SD Rats on Day 15.

Dose (mg/kg/day)	Mean 404-02 CNS Tissue Content (ng/g = ng/mL)* $\pm$ SEM	Mean 404-02 CNS Tissue Concentration (nM) $\pm$ SEM
1	29 $\pm$ 6.8	51 $\pm$ 11.8
10	421 $\pm$ 99	733 $\pm$ 173
50	750 $\pm$ 108	1306 $\pm$ 188
150	640 $\pm$ 82	1114 $\pm$ 143

* Conversion based on the assumption the density of tissue is equal to water, 1 g/mL.

Table 3d: Population Mean 404-02 Lung Tissue Content in SD Rats on Day 15.

Dose (mg/kg/day)	Mean dh404 Lung Tissue Content (ng/g = ng/mL)* $\pm$ SEM	Mean dh404 Lung Tissue Concentration (nM) $\pm$ SEM
1	558 $\pm$ 123	972 $\pm$ 215
10	4032 $\pm$ 928	7020 $\pm$ 1616
50	7165 $\pm$ 1221	12474 $\pm$ 2126
150	9719 $\pm$ 1643	16921 $\pm$ 2860

* Conversion based on the assumption the density of tissue is equal to water, 1 g/mL.

5 Example 6 – Rodent Toxicity Comparison between 402 and 402-02

A study was performed in Sprague Dawley rats using both **402** and **402-02**. Animals were dosed orally once daily for 7 days. The low dose **402** group had elevated total bilirubin and GGT levels as well as suppressed weight gain. The high dose animals that were treated with **402** were all sacrificed *in extremis* on Day 6 before study completion. GGT and total bilirubin levels were elevated in these animals as well. However, no toxicity as assessed by clinical observations, weight gain, GGT, and total bilirubin was observed in any animal treated with **402-02** (Table 4). In a second study involving oral administration to Sprague-Dawley rats for 14 days, **402-02**, achieved comparable blood levels to that of **402**. However, no significant toxicity was observed as assessed by weight loss, clinical observations, and GGT and total bilirubin elevations relative to controls at doses up to 1,500 mg/m²/day for 14 days, which is 50-fold higher than the MTD of RTA 402 in this species (Table 2).

Table 4: Comparing Compounds 402-02 and 402 for Rodent Toxicity

Timepoint/ Dose Level	Survival	Weight vs Control	GGT (U/L)	Total Bilirubin (mg/dL)
7 Day Study – Crystalline Forms of 402 and 402-02				
Vehicle Control	4/4 (100%)	100%	< 5	0.13
402 - 60 mg/m ² /day	4/4 (100%)	35%	8.7	0.25
402 - 180 mg/m ² /day	0/4 (0%)	NA	5.0	0.78
402-02 - 60 mg/m ² /day	4/4 (100%)	106%	< 5	0.20
402-02 - 180 mg/m ² /day	4/4 (100%)	132%	< 5	0.20
14 Day Study – Crystalline Form of 402-02				
Vehicle Control	10/10 (100%)	100%	< 5	0.20
402-02 - 60 mg/m ² /day	10/10 (100%)	95%	< 5	0.24
402-02 - 180 mg/m ² /day	5/5 (100%)	102%	< 5	0.22
402-02 - 600 mg/m ² /day	5/5 (100%)	96%	< 5	0.20
402-02 - 1,500 mg/m ² /day	5/5 (100%)	105%	< 5	0.20

Example 7 –Toxicity Comparison in Mice

In this study, six compounds (401, 402, 404, 401-2, 402-2, and 404-2) were assessed for toxicity in mice in a 14-day study. Each compound was formulated in sesame oil and administered daily by oral gavage at doses of 10, 50, 100, or 250 mg/kg (n = 4 per group). At higher doses (above 10 mg/kg/day) both 401 and 402 caused at least 50% mortality; 404 was non-toxic. In contrast, no mortality was observed in the 402-2 and 404-2 groups and only the highest dose of 401-02 caused any lethality (Table 5). Body weight measurements (FIGS. 29-31) were consistent with the mortality observations. The two highest doses of 401 and 402 were lethal within 4 days, in contrast to the effects of 401-2 and 402-2.

Table 5: Mortality Observations in 14-Day Toxicity Study

Group	Compound	Dose (mg/kg)	Schedule	N	Number of Deaths	Comments
1	vehicle		QD×14, D1-14	4	0	
2	401	10	QD×14, D1-14	4	0	
3	401	50	QD×14, D1-14	4	2	
4	401	100	QD×14, D1-14	4	4	
5	401	250	QD×14, D1-14	4	4	
6	401-02	10	QD×14, D1-14	4	0	
7	401-02	50	QD×14, D1-14	4	1*	*Due to gavage injury
8	401-02	100	QD×14, D1-14	4	0	
9	401-02	250	QD×14, D1-14	4	1	Sacrificed due to weightless on Day 11
10	402	10	QD×14, D1-14	4	0	
11	402	50	QD×14, D1-14	4	4	
12	402	100	QD×14, D1-14	4	4	
13	402	250	QD×14, D1-14	4	4	
14	402-02	10	QD×14, D1-14	4	0	
15	402-02	50	QD×14, ID1-14	4	0	
15	402-02	100	QD×14, D1-14	4	0	
17	402-02	250	QD×14, D1-14	4	0	
1a	404	10	QD×14, D1-14	4	0	
19	404	50	QD×14, D1-14	4	0	
20	404	100	QD×14, D1-14	4	0	
21	404	250	QD×14, D1-14	4	0	
22	404-02	10	QD×14, D1-14	4	0	

Group	Compound	Dose (mg/kg)	Schedule	N	Number of Deaths	Comments
23	404-02	50	QD×14, D1-14	4	0	
22	404-02	100	QD×14, D1-14	4	0	
23	404-02	250	QD×14, D1-14	4	0	

In a second experiment, six additional compound differing only in the saturation or non-saturation of the C ring were tested for toxicity in mice by daily oral administration for 9 days, using sesame oil as the vehicle. In this study, no significant toxicity was observed. The deaths of two animals were attributed to gavage errors during the administration of the test article. No significant differences in weight were observed in any group compared to the vehicle-treated controls. Results are summarized in Table 6 below. As with compounds 402-2, 401-2 and 404-2 above, compounds with saturation in the C ring consistently show low toxicity in rodents. Compounds lacking saturation in the C ring show significant rodent toxicity in some cases (*e.g.*, 401 and 402). Predictably low rodent toxicity provides an advantage since high rodent toxicity can be a significant complication in conducting preclinical studies required for development and registration of therapeutic compounds for use in humans or non-human animals.



Table 6: Further Mouse Toxicity Results.






Compound	Dose (per day, p.o.)	Mortality
63112	3 mg/kg	0/5
	10 mg/kg	0/5
	30 mg/kg	1/5
63323	3 mg/kg	0/5
	10 mg/kg	0/5
	30 mg/kg	0/5
63324	3 mg/kg	0/5
	10 mg/kg	0/5
	30 mg/kg	0/5






Compound	Dose (per day, p.o.)	Mortality
63325	3 mg/kg	0/5
	10 mg/kg	0/5
	30 mg/kg	0/5
63166	3 mg/kg	0/5
	10 mg/kg	0/5
	30 mg/kg	0/5
63326	3 mg/kg	0/5
	10 mg/kg	1/5
	30 mg/kg	0/5

Example 8 – Aqueous Solubility of Oleanolic Acid Derivatives

The aqueous solubility of the compounds shown here was determined using the procedures outlined in Example 1.

Compound ID(s)	Structure	Aqueous Solubility (μM)
63097 (402)		1.46
63102 (dh404)		0.06

Compound ID(s)	Structure	Aqueous Solubility (μM)
63198		163.6
63202		1.89
63208		9.49
63214		112.2
63219		13.58

Compound ID(s)	Structure	Aqueous Solubility (μM)
63221		8.78
63226		0.71
63231		1.23
63232		0.75
63237		5.16

All of the methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

REFERENCES

The following references, and those listed in the Appendix, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

- 5
- U.S. Patent 5,443,826
- U.S. Patent 5,599,795
- U.S. Patent 6,025,395
- U.S. Patent 6,974,801
- 10 U.S. Provisional Application No. 61/046,342
- U.S. Provisional Application No. 61/046,352
- U.S. Provisional Application No. 61/046,363
- U.S. Provisional Application No. 61/046,366
- U.S. Provisional Application No. 61/111,269
- 15 U.S. Provisional Application No. 61/111,294
- U.S. Patent Publication 2009/0060873
- U.S. Serial No. 12/352,473
- U.S. Serial No. 12/151,425
- 20 U.S. Patent Application by Eric Anderson, Gary L. Bolton, Deborah Ferguson, Xin Jiang, Robert M. Kral, Jr., Patrick M. O'Brian and Melean Visnick, entitled "Natural Products Including an Anti-Inflammatory Pharmacore and Methods of Use," filed April 20, 2009.
- U.S. Patent Application by Eric Anderson, Xin Jiang and Melean Visnick, entitled
- 25 "Antioxidant Inflammation Modulators: Oleanolic Acid Derivatives with Amino and Other Modifications At C-17," filed April 20, 2009.
- U.S. Patent Application by Xin Jiang, Jack Greiner, Lester L. Maravetz, Stephen S. Szucs, Melean Visnick, entitled "Antioxidant Inflammation Modulators: Novel Derivatives of Oleanolic Acid," filed April 20, 2009.
- 30 U.S. Patent Application by Xin Jiang, Xioafeng Liu, Jack Greiner, Stephen S. Szucs, Melean Visnick entitled, "Antioxidant Inflammation Modulators: C-17 Homologated Oleanolic Acid Derivatives," filed April 20, 2009.

- Abraham and Kappas, *Free Radic. Biol. Med.*, 39(1):1-25, 2005.
- Adult Treatment Panel III, or ATP III
- Ahmad *et al.*, *Cancer Res.*, 68(8):2920-2926, 2008.
- Ahmad *et al.*, *J Biol Chem.*, 281(47):35764-35769, 2006.
- 5 Akiyama *et al.*, *Alzheimer Dis. Assoc. Disord.*, 14(1):S47-53, 2000.
- Angulo *et al.*, *Eur. J. Immunol.*, 30:1263-1271, 2000.
- Araujo *et al.*, *J. Immunol.*, 171(3):1572-1580, 2003.
- Arend and Dayer, *Arthritis Rheum.*, 38:151-160, 1995.
- Arend *et al.*, *Annu. Rev. Immunol.*, 16:27-55, 1998.
- 10 Autenrieth *et al.*, *Infect. Immun.*, 62:2590-2599, 1994.
- Bach, *Hum. Immunol.*, 67(6):430-432, 2006
- Bagasra *et al.*, *Proc. Natl. Acad. Sci. USA*, 92:12041-12045, 1995.
- Ball, *Ann. Rheum. Dis.*, 30:213-223, 1971.
- Beal, *Curr. Opin. Neurobiol.*, 6:661-666, 1996.
- 15 Bendzen *et al.*, *Scand. J. Rheumatol.*, 28:599-606, 1988.
- Blumberg *et al.*, *Arthritis Rheum.*, 7:93-97, 1964.
- Botoman *et al.*, *Am. Fam. Physician*, 57(1):57-68, 1998.
- Brandt *et al.*, *Arthritis Rheum.*, 43:1346-1352, 2000.
- Braun *et al.*, *Arthritis Rheum.*, 42:2039-2044, 1999.
- 20 Brewerton *et al.*, *Lancet.*, 1:904-907, 1973a.
- Brewerton *et al.*, *Lancet.*, 1:956-957, 1973b.
- Bronte *et al.*, *Trends Immunol.*, 24:302-306, 2003.
- Brown and DuBois, *J. Clin. Oncol.*, 23:2840-2855, 2005.
- Brynskov *et al.*, *N. Engl. J. Med.*, 321(13):845-850, 1989.
- 25 Burger and Dayer, *Neurology*, 45(6S-6):S39-43, 1995.
- Cai *et al.*, *Nat. Med.*, 11(2):183-190, 2005.
- Calin and Taurog, In: *The Spondylarthritides*, Calin *et al.* (Eds.), Oxford, UK. Oxford University Press, 179, 1998.
- Cann *et al.*, *Gut.*, 24(12):1135-1140, 1983.
- 30 Chauhan and Chauhan, *Pathophysiology*, 13(3):171-181 2006.
- Chomarar *et al.*, *Arthritis Rheum.*, 38:1046-1054, 1995.
- Coyle and Puttfarcken, *Science*, 262:689-695, 1993.
- Crowell *et al.*, *Mol. Cancer Ther.*, 2:815-823, 2003.
- Culver *et al.*, *Science*, 256:1550-1552, 1992.

- de Waal *et al.*, *J. Exp. Med.*, 174:1209–1220, 1991.
- Dickerson *et al.*, *Prog Neuropsychopharmacol Biol. Psychiatry*, March 6, 2007.
- Dinareello, *Int. Rev. Immunol.*, 16:457–499, 1998.
- Dinkova-Kostova *et al.*, *Proc Natl Acad Sci USA*, 102(12):4584–4589, 2005.
- 5 Dionne *et al.*, *Clin. Exp. Immunol.*, 112(3):435–442, 1998.
- Doran *et al.*, *J. Rheumatol.*, 30(2):316–320, 2003.
- Drossman *et al.*, *Dig. Dis. Sci.*, 38(9):1569–1580, 1993.
- Drossman *et al.*, *Gastroenterol.*, 112(6):2120–2137, 1997.
- Dudhgaonkar *et al.*, *Eur. J. Pain*, 10(7):573–9, 2006.
- 10 Eikelenboom *et al.*, *Glia*, 40(2):232–239, 2002.
- Ettehad *et al.*, *Clin. Exp. Immunol.*, 96(1):146–151, 1994.
- Everhart *et al.*, *Gastroenterol.*, 100(4):998–1005, 1991.
- Fearon and Locksley, *Science*, 272(5258):50–53, 1996.
- Feldtkeller *et al.*, *Rheumatol. Int.*, 23(2):61–66, 2003.
- 15 Firestein *et al.*, *Arthritis Rheum.*, 37:644–652, 1994.
- Forstermann, *Biol. Chem.*, 387:1521, 2006.
- Fujikawa *et al.*, *Ann. Rheum. Dis.*, 54:318–320, 1995.
- Funakoshi *et al.*, *Digestion*, 59(1):73–78, 1998.
- Galley and Webster, *Br. J. Anaesth.*, 77:11–16, 1996.
- 20 Gehrmann *et al.*, *Glia*, 15(2):141–151, 1995.
- Genain and Nauser, *J. Mol. Med.*, 75:187–197, 1997.
- Gladman *et al.*, *Br. J. Rheumatol.*, 22:675–679, 1995.
- Gladman *et al.*, *J. Med.*, 62:127–141, 1987.
- Gladman, *Rheum. Dis. Clin. North Am.*, 18:247–256, 1992.
- 25 Goodman *et al.*, *Kidney Int.*, 72(8):945–953, 2007.
- Graeber *et al.*, *Glia*, 40(2):252–259, 2002.
- Greten *et al.*, *Cell*, 118:285–296, 2004.
- Griffin *et al.*, *Proc. Natl. Acad. Sci. USA*, 86(19):7611–7615, 1989.
- Guilherme *et al.*, *Nat. Rev. Mol. Cell Biol.*, 9(5):367–77, 2008.
- 30 Gwee *et al.*, *Gut*, 44(3):400–406, 1999.
- Hahn and Tsao, In: *Dubois' Lupus Erythematosus*, 4th Ed, Wallace and Hahn (Eds.), Lea and Febiger, Philadelphia, 195–201, 1993.
- Handbook of Pharmaceutical Salts: Properties, and Use* (Stahl & Wermuth eds., Verlag Helvetica Chimica Acta, 2002.

- Handbook of Pharmaceutical Salts: Properties, Selection and Use* (Stahl & Wermuth, Eds.), Verlag Helvetica Chimica Acta, 2002.
- Hannum *et al.*, *Nature*, 343:336–340, 1990.
- Hanson *et al.*, *BMC Medical Genetics*, 6(7), 2005.
- 5 Hansson *et al.*, *Annu. Rev. Pathol. Mech. Dis.*, 1:297–329, 2006.
- Harrison and Symmons *et al.*, *Ann. Rheum. Dis.*, 57(6):375–377, 1998.
- Harrison *et al.*, *J. Rheumatol.*, 25(12):2324–2330, 1998.
- Hart *et al.*, *Immunology*, 84:536–542, 1995.
- Hohler *et al.*, *Arthritis Rheum.*, 41:1489–1492, 1998.
- 10 Hohler *et al.*, *J. Invest. Dermatol.*, 109:562–565, 1997.
- Honda *et al.*, *Bioorg. Med. Chem. Lett.*, 12:1027–1030, 2002.
- Honda *et al.*, *Bioorg. Med. Chem. Lett.*, 19:2711–2714, 1998.
- Honda *et al.*, *Bioorg. Med. Chem. Lett.*, 9:3429–3434, 1999.
- Honda *et al.*, *J. Med. Chem.*, 43:1866–1877, 2000a.
- 15 Honda *et al.*, *J. Med. Chem.*, 43:4233–4246, 2000b.
- Horwitz and Fisher, *N. Engl. J. Med.*, 344(24):1846–1850, 2001.
- Hotamisligil, *Nature*, 444(7121):860–7, 2006.
- Ishikawa *et al.*, *Circulation*, 104(15):1831–1836, 2001.
- Ishizawa and Dickson, *J. Neuropathol. Exp. Neurol.*, 60(6):647–657, 2001.
- 20 Jacob *et al.*, *Proc. Natl. Acad. Sci. USA*, 87:1233–1237, 1990.
- Jailwala *et al.*, *Ann. Intern. Med.*, 133(2):136–147, 2000.
- Jarvis, *Curr. Opin. Rheumatol.*, 10(5):459–467, 1998.
- Jarvis, *Pediatr. Ann.*, 31(7):437–446, 2002.
- Jones *et al.*, *Br. J. Rheumatol.*, 33(9):834–839, 1994.
- 25 Jonsson *et al.*, *Br. J. Rheumatol.*, 32(7):578–581 1993.
- Jonsson *et al.*, *Oral Dis.*, 8(3):130–140, 2002.
- Jonsson *et al.*, *Trends Immunol.*, 22(12):653–654, 2001.
- Kahle *et al.*, *Ann. Rheum. Dis.*, 51:731–734, 1992.
- Kaltschmidt *et al.*, *Proc. Natl. Acad. Sci. USA*, 94:2642–2647, 1997.
- 30 Kawakami *et al.*, *Brain Dev.*, 28(4):243–246, 2006.
- Kellow and Phillips, *Gastroenterol.*, 92(6):1885–1893, 1987.
- Kendall-Tackett, *Trauma Violence Abuse*, 8(2):117–126, 2007.
- Khan *et al.*, *J. Neurochem.*, 71:78–87, 1998.
- Khan *et al.*, *Toxicol. Applied Pharmacol.*, 103:482–490, 1990.

- Kortylewski *et al.*, *Nat. Med.*, 11:1314-1321, 2005.
- Kotake *et al.*, *Infect. Immun.*, 67:2682–2686, 1999.
- Kotzin and O'Dell, In: *Samler's Immunologic Diseases*, 5th Ed., Frank *et al.* (Eds.), Little Brown & Co., Boston, 667-697, 1995.
- 5 Kotzin, *Cell*, 85:303-306, 1996.
- Kruger *et al.*, *J. Pharmacol. Exp. Ther.*, 319(3):1144-1152, 2006.
- Kuboyama, *Kurume Med. J.*, 45(1):33-37, 1998.
- Lahesmaa *et al.*, *J. Immunol.*, 148:3079–3085, 1992.
- Lee *et al.*, *Glia.*, 55(7):712-22, 2007.
- 10 Lencz *et al.*, *Mol. Psychiatry*, 12(6):572-80, 2007.
- Liby *et al.*, *Nat Rev. Cancer*, 7(5):357-369, 2007.
- Lipsky, In: *Harrison's principles of internal medicine*, Fauci *et al.* (Eds.), 14th Ed., NY, McGraw-Hill, 1880–1888, 1998.
- Liu *et al.*, *FASEB J.*, 20(2):207-216, 2006.
- 15 Lo *et al.*, *Curr. Dir. Autoimmun.*, 1:226-246, 1999.
- Lugering *et al.*, *Ital. J. Gastroenterol. Hepatol.*, 30(3):338-344, 1998.
- Lynn and Friedman, *N. Engl. J. Med.*, 329(26):1940-1945, 1993.
- Macatonia *et al.*, *J. Immunol.*, 150:3755–3765, 1993.
- March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure, 2007.
- 20 March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure (March's Advanced Organic Chemistry), Smith and March (Eds.), 2007.
- Marsal *et al.*, *Rheumatology*, 38:332-337, 1999.
- Mazur *et al.*, *Cell Microbiol.*, 9(7):1683-94, 2007.
- Mazzoni *et al.*, *J. Immunol.*, 168:689-695, 2002.
- 25 McAlindon *et al.*, *Gut*, 42(2):214-219, 1998.
- McGeer and McGeer, *Brain Res. Brain Res. Rev.*, 21:195–218, 1995.
- McGeer *et al.*, *Neurology*, 19:331–338, 1996.
- McGonagle *et al.*, *Arthritis Rheum.*, 41:694-700, 1998.
- McGonagle *et al.*, *Curr. Opin. Rheumatol.*, 11:244-250, 1999.
- 30 McIver *et al.*, *Pain*, 120(1-2):161-9, 2005.
- Mease *et al.*, *Lancet*, 356:385–390, 2000.
- Merrill and Benvenist, *Trends Neurosci.*, 19:331–338, 1996.
- Mertz *et al.*, *Gastroenterol.*, 118(5):842-848, 2000.
- Moll and Wright, *Ann. Rheum. Dis.*, 32:181–201, 1973.


- Moll and Wright, *Semin. Arthritis Rheum.*, 3:55-78, 1973.
- Morris *et al.*, *J. Mol. Med.*, 80(2):96-104, 2002.
- Morse and Choi, *Am. J. Respir. Crit. Care Med.*, 172(6):660-670, 2005.
- Morse and Choi, *Am. J. Respir. Crit. Care Med.*, 27(1):8-16, 2002.
- 5 Nath *et al.*, *Neurology*, 66(1):149-150, 2006.
- National Institutes of Health, NIH Publication No. 01-3670, 2001.
- Neal *et al.*, *BMJ.*, 314(7083):779-782, 1997.
- Nichols, *Drug News Perspect.*, 17(2):99-104, 2004.
- Nielen *et al.*, *Arthritis Rheum.*, 50(2):380-386, 2004.
- 10 Ohnishi *et al.*, *Int. Immunol.*, 6:817-830, 1994.
- Pall, *Med. Hypoth.*, 69:821-825, 2007.
- Partschi *et al.*, *Br. J. Rheumatol.*, 24:518-523, 1997.
- Pica *et al.*, *Antimicrob Agents Chemother.*, 44(1):200-4, 2000.
- Pimentel *et al.*, *Am. J. Gastroenterol.*, 95(12):3503-3506, 2000.
- 15 Pociot *et al.*, *Scand. J. Immunol.*, 42(4):501-504, 1995.
- Prieur *et al.*, *Lancet.*, 2:1240-1242, 1987.
- Rajakariar *et al.*, *Proc. Natl. Acad. Sci. USA*, 104(52):20979-84, 2007.
- Rantapaa-Dahlqvist *et al.*, *Arthritis Rheum.*, 48(10):2741-2749, 2003.
- Reimund *et al.*, *Eur. J. Clin. Invest.*, 28(2):145-150, 1998.
- 20 Ribbens *et al.*, *Eur. Cytokine Netw.*, 11:669-676, 2000.
- Rogers *et al.*, *Neurobiol Aging*, 9(4):339-349, 1988.
- Rogler and Andus, *World J. Surg.*, 22(4):382-389, 1998.
- Rooney *et al.*, *Rheumatol. Int.*, 10:217-219, 1990.
- Ross *et al.*, *Nutr. Neurosci.*, 6(5):277-81, 2003.
- 25 Rostom *et al.*, *Ann. Intern. Med.*, 146, 376-389, 2007.
- Rothstein, *Med. Clin. North Am.*, 84(5):1247-1257, 2000.
- Ruster *et al.*, *Scand. J. Rheumatol.*, 34(6):460-3, 2005.
- Sacerdoti *et al.*, *Curr Neurovasc Res.* 2(2):103-111, 2005.
- Saiki *et al.*, *Scand. J. Gastroenterol.*, 33(6):616-622, 1998.
- 30 Salomonsson and Jonsson, *Arthritis Rheum.*, 48(11):3187-3201, 2003.
- Salomonsson *et al.*, *Scand. J. Immunol.*, 55(4):336-342, 2002.
- Salvarani *et al.*, *Curr. Opin. Rheumatol.* 1998; 10:299-305, 1998.
- Salvemini *et al.*, *J. Clin. Invest.*, 93:1940-1947, 1994.
- Sandler, *Gastroenterol.*, 99(2):409-415, 1990.

- Sarchielli *et al.*, *Cephalalgia*, 26(9):1071-1079, 2006.
- Satoh *et al.*, *Proc. Natl. Acad. Sci. USA*, 103(3):768-773, 2006.
- Schellekens *et al.*, *Arthritis Rheum.*, 43(1):155-163, 2000.
- Schlaak *et al.*, *Clin. Exp. Rheumatol.*, 14:155-162, 1996.
- 5 Schlaak *et al.*, *Eur. J. Immunol.*, 22:2771-2776, 1992.
- Schlosstein *et al.*, *NE J. Medicine*, 288:704-706, 1973.
- Schreiber, *Neth. J. Med.*, 53(6):S24-31, 1998.
- Schulz *et al.*, *Antioxid. Redox. Sig.*, 10:115, 2008.
- Sieper and Braun, *Arthritis Rheum.*, 38:1547-1554, 1995.
- 10 Simon *et al.*, *Clin. Exp. Immunol.*, 94:122-126, 1993.
- Simon *et al.*, *Proc. Natl. Acad. Sci. USA*, 91:8562-8566, 1994.
- Simonian and Coyle, *Annu. Rev. Pharmacol. Toxicol.*, 36:83-106, 1996.
- Sinha *et al.*, *Cancer Res.*, 67:4507-4513, 2007.
- Stack *et al.*, *Lancet*, 349(9051):521-524, 1997.
- 15 Stewart *et al.*, *Neurology*, 48:626-632, 1997.
- Strejan *et al.*, *J. Neuroimmunol.*, 7:27, 1984.
- Szabo *et al.*, *Nature Rev. Drug Disc.*, 6:662-680, 2007.
- Takahashi *et al.*, *Cancer Res.*, 57:1233-1237, 1997.
- Talley *et al.*, *Gastroenterol.*, 109(6):1736-1741, 1995.
- 20 Tamir and Tannenbaum, *Biochim. Biophys. Acta.*, 1288:F31-F36, 1996.
- Targan *et al.*, *N. Engl. J. Med.*, 337(15):1029-1035, 1997.
- Third Report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III, or ATP III), National Institutes of Health, 2001, NIH Publication No. 01-
- 25 3670.
- Touzani *et al.*, *J. Neuroimmunol.*, 100(1-2):203-215, 1999.
- Tumlin *et al.*, *Am. J. Cardiol.*, 98(6A):14K-20K, 2006.
- van den Berg, *Semin. Arthritis Rheum.*, 30(5S-2):7-16, 2001.
- van Dullemen *et al.*, *Gastroenterol.*, 109(1):129-135, 1995.
- 30 van Hogezaand and Verspaget, *Drugs*, 56(3):299-305, 1998.
- Vazquez *et al.*, *J. Virol.*, 79(7):4479-91, 2005.
- Vodovotz *et al.*, In; *Handbook of Experimental Immunology*, Volumes I-IV, 1996.
- Wardle, *Nephrol. Dial. Transplant.*, 16(9):1764-8, 2001.
- Warrington *et al.*, *Arthritis and Rheumatism*, 44:13-20, 2001.

- Weyand and Goronzy, *Ann. NY Acad. Sci.*, 987:140-149, 2003.
- Whitehead *et al.*, *Gastroenterol.*, 98(5 Pt 1):1187-1192, 1990.
- Williams *et al.*, *Clin. Neurosci.*, 2(3-4):229-245, 1994.
- Wordsworth, In: *Genes and Arthritis*, Brit. Medical Bulletin, 51:249-266, 1995.
- 5 Wright, *Ann. Rheum. Dis.*, 15:348-356, 1956.
- Wright, *Clin. Orthop. Related Res.*, 143:8-14, 1979.
- Xanthou *et al.*, *Arthritis Rheum.*, 44(2):408-418, 2001.
- Yates *et al.*, *Cancer Res.*, 66(4): 2488-2494, 2006.
- Yin *et al.*, *Arthritis Rheum.*, 40:1788-1797, 1997.
- 10 Yin *et al.*, *Rheumatology*, 38:1058-1067, 1999.
- Yoh *et al.*, *Kidney Int.*, 60(4):1343-1353, 2001.
- Yu *et al.*, *Nat. Rev. Immunol.*, 7:41-51, 2007.
- Zhou *et al.*, *Am. J. Pathol.*, 166(1):27-37, 2005.
- Zhou *et al.*, *Cancer Sci.*, 98:882-889, 2007.
- 15 Zingarelli *et al.*, *J. Immunol.*, 171(12):6827-6837, 2003.

CLAIMS

1. A compound of the formula:



wherein:

5 Y is cyano, heteroaryl_(C≤12), substituted heteroaryl_(C≤12), or $-C(O)R_a$, further wherein R_a is:


hydrogen, hydroxy, halo, amino, hydroxyamino, azido, silyl or mercapto;

10 alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12), heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyloxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12), heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12), alkylamino_(C≤12), dialkylamino_(C≤12), alkoxyamino_(C≤12), alkenylamino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12), aralkylamino_(C≤12), heteroarylamino_(C≤12), heteroaralkylamino_(C≤12), alkylsulfonylamino_(C≤12), amido_(C≤12), alkylthio_(C≤12), alkenylthio_(C≤12), alkynylthio_(C≤12), arylthio_(C≤12), aralkylthio_(C≤12), heteroarylthio_(C≤12), heteroaralkylthio_(C≤12), acylthio_(C≤12), alkylammonium_(C≤12), alkylsulfonium_(C≤12), alkylsilyl_(C≤12), or a substituted version of any of these groups; or

15

20

R_a comprises a nitrogen atom that is also attached to carbon atom 13 and R_d to form:



wherein R_d is $\text{alkyl}_{(C \leq 8)}$, $\text{alkenyl}_{(C \leq 8)}$, $\text{aryl}_{(C \leq 8)}$, $\text{aralkyl}_{(C \leq 8)}$, $\text{heteroaryl}_{(C \leq 8)}$, $\text{heteroaralkyl}_{(C \leq 8)}$, or a substituted version of any of these groups;

Z is a single or double bond, $-\text{O}-$ or $-\text{NR}_e-$, wherein R_e is hydrogen, hydroxy, $\text{alkyl}_{(C \leq 8)}$ or $\text{alkoxy}_{(C \leq 8)}$;

X is OR_b , NR_bR_c , or SR_b , wherein R_b and R_c are each independently:

hydrogen or hydroxy;

$\text{alkyl}_{(C \leq 8)}$, $\text{aryl}_{(C \leq 8)}$, $\text{aralkyl}_{(C \leq 8)}$, $\text{acyl}_{(C \leq 8)}$, $\text{alkoxy}_{(C \leq 8)}$, $\text{aryloxy}_{(C \leq 8)}$, $\text{acyloxy}_{(C \leq 8)}$, $\text{alkylamino}_{(C \leq 8)}$, $\text{arylamino}_{(C \leq 8)}$, $\text{amido}_{(C \leq 8)}$, or a substituted version of any of these groups; or

a substituent convertible *in vivo* to hydrogen;

provided that R_b is absent when the atom to which it is bound is part of a double bond, further provided that when R_b is absent the atom to which it is bound is part of a double bond;

R_1 is:

hydrogen, cyano, hydroxy, halo or amino; or

$\text{alkyl}_{(C \leq 8)}$, $\text{alkenyl}_{(C \leq 8)}$, $\text{alkynyl}_{(C \leq 8)}$, $\text{aryl}_{(C \leq 8)}$, $\text{aralkyl}_{(C \leq 8)}$, $\text{heteroaryl}_{(C \leq 8)}$, $\text{heteroaralkyl}_{(C \leq 8)}$, $\text{acyl}_{(C \leq 8)}$, $\text{alkoxy}_{(C \leq 8)}$, $\text{aryloxy}_{(C \leq 8)}$, $\text{acyloxy}_{(C \leq 8)}$, $\text{alkylamino}_{(C \leq 8)}$, $\text{arylamino}_{(C \leq 8)}$, $\text{amido}_{(C \leq 8)}$, or a substituted version of any of these groups;

R_2 is:

cyano, hydroxy, halo or amino; or

$\text{fluoroalkyl}_{(C \leq 8)}$, $\text{alkenyl}_{(C \leq 8)}$, $\text{alkynyl}_{(C \leq 8)}$, $\text{aryl}_{(C \leq 8)}$, $\text{heteroaryl}_{(C \leq 8)}$, $\text{acyl}_{(C \leq 8)}$, $\text{alkoxy}_{(C \leq 8)}$, $\text{aryloxy}_{(C \leq 8)}$, $\text{acyloxy}_{(C \leq 8)}$, $\text{alkylamino}_{(C \leq 8)}$,

arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups;

R₃ is:

absent or hydrogen;

5 alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any of these groups; or

a substituent convertible *in vivo* to hydrogen;

10 provided that R₃ is absent when the oxygen atom to which it is bound is part of a double bond, further provided that when R₃ is absent the oxygen atom to which it is bound is part of a double bond;

R₄ and R₅ are each independently alkyl_(C≤8) or substituted alkyl_(C≤8);


R₆ is hydrogen, hydroxy or oxo;

R₇ is hydrogen or hydroxy; and

15 R₈, R₉, R₁₀ and R₁₁ are each independently hydrogen, hydroxy, alkyl_(C≤8), substituted alkyl_(C≤8), alkoxy_(C≤8) or substituted alkoxy_(C≤8);

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

2. The compound of claim 1, further defined as:



20

wherein:

Y is cyano or $-C(O)R_a$, further wherein:

R_a is:

hydrogen, hydroxy, halo, amino, hydroxyamino, azido or
mercapto; or

alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12),
heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyl-
oxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12),
heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12),
alkylamino_(C≤12), dialkylamino_(C≤12), alkoxyamino_(C≤12),
alkenylamino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12),
aralkylamino_(C≤12), heteroarylamino_(C≤12), heteroaralkyl-
amino_(C≤12), alkylsulfonylamino_(C≤12), amido_(C≤12),
alkylthio_(C≤12), alkenylthio_(C≤12), alkynylthio_(C≤12), aryl-
thio_(C≤12), aralkylthio_(C≤12), heteroarylthio_(C≤12),
heteroaralkylthio_(C≤12), acylthio_(C≤12),
alkylammonium_(C≤12), alkylsulfonium_(C≤12),
alkylsilyl_(C≤12), or a substituted version of any of these
groups;

X is OR_b , NR_bR_c , or SR_b , wherein R_b and R_c are each independently:

hydrogen or hydroxy;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8),
aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8),
arylamino_(C≤8), amido_(C≤8), or a substituted version of
any of these groups; or

a substituent convertible *in vivo* to hydrogen;

provided that R_b is absent when the atom to which it is bound is
part of a double bond, further provided that when R_b is
absent the atom to which it is bound is part of a double
bond;

R_1 is:

hydrogen, cyano, hydroxy, halo or amino; or

alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8), heteroaralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups;

5 R₂ is:

cyano, hydroxy, halo or amino; or

fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of
10 these groups;

R₃ is:

absent or hydrogen;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any
of these groups; or

15 a substituent convertible *in vivo* to hydrogen;


provided that R₃ is absent when the oxygen atom to which it is bound
is part of a double bond, further provided that when R₃ is
absent the oxygen atom to which it is bound is part of a double
bond;

20 R₄ and R₅ are each independently alkyl_(C≤8) or substituted alkyl_(C≤8); and

R₆ and R₇ are each independently hydrogen or hydroxy;

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs,
or optical isomers thereof.

3. The compound of claim 2, further defined as:



wherein:

Y is cyano or $-\text{C}(\text{O})\text{R}_a$, further wherein:

5

R_a is:

hydrogen, hydroxy, halo, amino, azido, mercapto or silyl; or
 alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12),
 heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyl-
 oxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12),
 heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12),
 10 alkylamino_(C≤12), alkoxyamino_(C≤12), dialkylamino_(C≤12),
 alkenylamino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12),
 aralkylamino_(C≤12), heteroarylamino_(C≤12), heteroaralkyl-
 amino_(C≤12), amido_(C≤12), or a substituted version of any
 15 of these groups;

X is OR_b , NR_bR_c , or SR_b , wherein R_b and R_c are each independently:

hydrogen or hydroxy;
 alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8),
 aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8),
 20 arylamino_(C≤8), amido_(C≤8), or a substituted version of
 any of these groups; or
 a substituent convertible *in vivo* to hydrogen;

provided that R_b is absent when the atom to which it is bound is part of a double bond, further provided that when R_b is absent the atom to which it is bound is part of a double bond;

5 R_1 is:

hydrogen, cyano, hydroxy, halo or amino; or

alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8), heteroaralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups;

10 R_2 is:


cyano, hydroxy, halo or amino; or

fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups; and

15 R_4 is alkyl_(C≤8) or substituted alkyl_(C≤8);

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

20 4. The compound of claim 3, further defined as:



wherein:

R_a is:

hydrogen, hydroxy, halo, or amino; or

alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8),
heteroaralkyl_(C≤8), alkoxy_(C≤8), alkenyloxy_(C≤8), alkynyloxy_(C≤8),
aryloxy_(C≤8), aralkoxy_(C≤8), heteroaryloxy_(C≤8),
heteroaralkoxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8),
alkoxyamino_(C≤8), dialkylamino_(C≤8), alkenylamino_(C≤8),
alkynylamino_(C≤8), arylamino_(C≤8), aralkylamino_(C≤8),
heteroarylamino_(C≤8), heteroaralkylamino_(C≤8), amido_(C≤8), or a
substituted version of any of these groups;

X is OR_b or NR_bR_c, wherein R_b and R_c are each independently:

hydrogen or hydroxy;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8),
aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8),
arylamino_(C≤8), amido_(C≤8), or a substituted version of
any of these groups; or

a substituent convertible *in vivo* to hydrogen;

provided that R_b is absent when the atom to which it is bound is
part of a double bond, further provided that when R_b is
absent the atom to which it is bound is part of a double
bond; and


R₂ is:

cyano, hydroxy, halo or amino; or

fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8),
acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8),
arylamino_(C≤8), amido_(C≤8), or a substituted version of any of
these groups;

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs,
or optical isomers thereof.

5. The compound of claim 4, further defined as:



wherein:

R_a is:

hydrogen, hydroxy, halo, or amino; or

alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8), heteroaralkyl_(C≤8), alkoxy_(C≤8), alkenyloxy_(C≤8), alkynyloxy_(C≤8), aryloxy_(C≤8), aralkoxy_(C≤8), heteroaryloxy_(C≤8), heteroaralkoxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), alkoxyamino_(C≤8), dialkylamino_(C≤8), alkenylamino_(C≤8), alkynylamino_(C≤8), arylamino_(C≤8), aralkylamino_(C≤8), heteroarylamino_(C≤8), heteroaralkylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups; and


R_2 is:

cyano or fluoro; or

fluoroalkyl_(C≤5), alkenyl_(C≤5), alkynyl_(C≤5), heteroaryl_(C≤5), acyl_(C≤5), acyloxy_(C≤5), amido_(C≤5), or a substituted version of any of these groups;

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

6. The compound of claim 5, further defined as:




wherein R_a is:

hydrogen, hydroxy, halo or amino; or

- 5 alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8),
 heteroaralkyl_(C≤8), alkoxy_(C≤8), alkenyloxy_(C≤8), alkynyloxy_(C≤8),
 aryloxy_(C≤8), aralkoxy_(C≤8), heteroaryloxy_(C≤8), heteroaralkoxy_(C≤8),
 acyloxy_(C≤8), alkylamino_(C≤8), alkoxyamino_(C≤8), dialkylamino_(C≤8),
 alkenylamino_(C≤8), alkynylamino_(C≤8), arylamino_(C≤8), aralkylamino_(C≤8),
 10 heteroarylamino_(C≤8), heteroaralkylamino_(C≤8), amido_(C≤8), or a
 substituted version of any of these groups;


or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs,
 or optical isomers thereof.

7. The compound of claim 6, further defined as:




- 15 wherein R_a is alkoxy_(C1-4), alkylamino_(C1-4), alkoxyamino_(C1-4), dialkylamino_(C2-4), or a
 substituted version of any of these groups; or pharmaceutically acceptable salts,
 esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

8. The compound of claim 6, further defined as:




wherein R_a is alkyl_(C1-4) or aralkoxy_(C7-8) or a substituted version of either of these groups; or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

9. The compound of claim 6, further defined as:




wherein R_a is hydrogen, hydroxy, amino, dimethylamino, methyl, methoxy, methoxyamino, benzyloxy, or 2,2,2-trifluoroethylamino; or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

10. The compound of claim 6, further defined as:




wherein R_a is hydrogen, hydroxy, amino, methoxy, or 2,2,2-trifluoroethylamino; or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

11. The compound of claim 1, further defined as:




wherein R_d is $\text{alkyl}_{(C \leq 8)}$, $\text{alkenyl}_{(C \leq 8)}$, $\text{aryl}_{(C \leq 8)}$, $\text{aralkyl}_{(C \leq 8)}$, $\text{heteroaryl}_{(C \leq 8)}$, $\text{heteroaralkyl}_{(C \leq 8)}$, or a substituted version of any of these groups; or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

12. The compound of claim 1, further defined as:



wherein Y is $\text{heteroaryl}_{(C \leq 8)}$ or a substituted $\text{heteroaryl}_{(C \leq 8)}$; or a pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

13. The compound of claim 2, further defined as:



wherein:

wherein Y is cyano or $-\text{C}(\text{O})\text{R}_a$, further wherein:

R_a is:

hydrogen, hydroxy, halo, amino, azido, mercapto or silyl; or
 alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12),
 heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyl-
 oxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12),
 heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12),
 alkylamino_(C≤12), alkoxyamino_(C≤12), dialkylamino_(C≤12),
 alkenylamino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12),
 aralkylamino_(C≤12), heteroarylamino_(C≤12), heteroaralkyl-
 amino_(C≤12), amido_(C≤12), or a substituted version of any
 of these groups;

R₂ is:

cyano, hydroxy, halo or amino; or
 fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8),
 acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8),
 arylamino_(C≤8), amido_(C≤8), or a substituted version of any of
 these groups;


R₃ is:

absent or hydrogen;
 alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any
 of these groups; or
 a substituent convertible *in vivo* to hydrogen;
 provided that R₃ is absent when the oxygen atom to which it is bound
 is part of a double bond, further provided that when R₃ is
 absent the oxygen atom to which it is bound is part of a double
 bond; and

R₄ is alkyl_(C≤8) or substituted alkyl_(C≤8);

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs,
 or optical isomers thereof.

14. The compound of claim 2 further defined as:



wherein:

Y is cyano or $-C(O)R_a$, wherein

5

R_a is:

hydrogen, hydroxy, halo, amino, hydroxyamino, azido or
mercapto; or

alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12),
heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12),
10 alkenyloxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralk-
oxy_(C≤12), heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyl-
oxy_(C≤12), alkylamino_(C≤12), alkoxyamino_(C≤12), dialkyl-
amino_(C≤12), alkenylamino_(C≤12), alkynylamino_(C≤12), aryl-
amino_(C≤12), aralkylamino_(C≤12), heteroarylamino_(C≤12),
15 heteroaralkylamino_(C≤12), alkylsulfonylamino_(C≤12),
amido_(C≤12), or a substituted version of any of these
groups;

R_2 is:

20

cyano, hydroxy, halo or amino; or
fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8),
acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8),
arylamino_(C≤8), amido_(C≤8), or a substituted version of any of
these groups; and


R_3 is:

25

hydrogen;
alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any
of these groups; or
a substituent convertible *in vivo* to hydrogen;

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

15. The compound of claim 3, further defined as:



,

5 wherein:

X is OR_b, NR_bR_c, or SR_b, wherein R_b and R_c are each independently:

hydrogen;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any of these groups; or

10 a substituent convertible *in vivo* to hydrogen;

provided that R_b is absent when the atom to which it is bound is part of a double bond, further provided that when R_b is absent the atom to which it is bound is part of a double bond;

15 R₁ is:

hydrogen, cyano, hydroxy, halo or amino; or

alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8), heteroaralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups;

20


R₂ is:

cyano, hydroxy, halo or amino; or

fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups; and

5 R₄ is alkyl_(C≤8) or substituted alkyl_(C≤8);
or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

16. The compound of claim 15, further defined as:



10 wherein:

X is OR_b or NR_bR_c, wherein R_b and R_c are each independently:

hydrogen;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any of these groups; or

15 a substituent convertible *in vivo* to hydrogen;

provided that R_b is absent when the atom to which it is bound is part of a double bond, further provided that when R_b is absent the atom to which it is bound is part of a double bond; and

20 R₂ is:


cyano, hydroxy, halo or amino; or

fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8),

arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups;

or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

5 17. The compound of claim 3, further defined as:



wherein:

R_a is:

hydrogen, hydroxy, halo, amino, azido, mercapto or silyl; or

10 alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12),
heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12),
alkenyloxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12),
heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12), alkyl-
amino_(C≤12), alkoxyamino_(C≤12), dialkylamino_(C≤12), alkenyl-
15 amino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12),
aralkylamino_(C≤12), heteroarylamino_(C≤12),
heteroaralkylamino_(C≤12), amido_(C≤12), or a substituted version of
any of these groups;

R₁ is:

20 hydrogen, cyano, hydroxy, halo or amino; or
alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8),
heteroaralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8),

acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups;


R₂ is:

cyano, hydroxy, halo or amino; or

5 fluoroalkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), heteroaryl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups; and

10 or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

18. The compound according to claim 17, further defined as:




wherein R_a is:

hydrogen, hydroxy, halo or amino; or

15 alkyl_(C≤6), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8), alkoxy_(C≤6), aryloxy_(C≤8), aralkoxy_(C≤8), alkylamino_(C≤6), alkoxyamino_(C≤6), alkoxyamino_(C≤6), dialkylamino_(C≤6), arylamino_(C≤8), aralkylamino_(C≤8), heteroarylamino_(C≤6), heteroarylamino_(C≤8), amido_(C≤6), or a substituted version of any of these groups;

20 or pharmaceutically acceptable salts, esters, hydrates, solvates, tautomers, prodrugs, or optical isomers thereof.

19. A compound of the formula:



wherein:

Y is cyano or $-\text{C}(\text{O})\text{R}_a$, wherein

R_a is:

hydrogen, hydroxy, halo, amino, hydroxyamino, azido or mercapto; or

alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12), heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyloxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12), heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12), alkylamino_(C≤12), dialkylamino_(C≤12), alkoxyamino_(C≤12), alkenylamino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12), aralkylamino_(C≤12), heteroarylamino_(C≤12), heteroaralkylamino_(C≤12), alkylsulfonylamino_(C≤12), amido_(C≤12), alkylthio_(C≤12), alkenylthio_(C≤12), alkynylthio_(C≤12), arylthio_(C≤12), aralkylthio_(C≤12), heteroarylthio_(C≤12), heteroaralkylthio_(C≤12), acylthio_(C≤12), alkylammonium_(C≤12), alkylsulfonium_(C≤12), alkylsilyl_(C≤12), or a substituted version of any of these groups;

X is OR_b , NR_bR_c , or SR_b , wherein R_b and R_c are each independently:

hydrogen;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted version of any of these groups; or

provided that R_b is absent when the atom to which it is bound is part of a double bond, further provided that when R_b is absent the atom to which it is bound is part of a double bond; and


R₁ is:

hydrogen, cyano, hydroxy, halo or amino; or

alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8), heteroaralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8), acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a substituted version of any of these groups;

or salts, esters, hydrates, solvates, tautomers, or optical isomers thereof.

20. A compound of the formula:



wherein:

Y is cyano or $-C(O)R_a$, wherein

R_a is:

hydrogen, hydroxy, halo, amino, hydroxyamino, azido or mercapto; or

alkyl_(C≤12), alkenyl_(C≤12), alkynyl_(C≤12), aryl_(C≤12), aralkyl_(C≤12), heteroaryl_(C≤12), heteroaralkyl_(C≤12), alkoxy_(C≤12), alkenyloxy_(C≤12), alkynyloxy_(C≤12), aryloxy_(C≤12), aralkoxy_(C≤12),

heteroaryloxy_(C≤12), heteroaralkoxy_(C≤12), acyloxy_(C≤12),
 alkylamino_(C≤12), dialkylamino_(C≤12), alkoxyamino_(C≤12),
 alkenylamino_(C≤12), alkynylamino_(C≤12), arylamino_(C≤12),
 aralkylamino_(C≤12), heteroarylmino_(C≤12), heteroaralkyl-
 amino_(C≤12), alkylsulfonylamino_(C≤12), amido_(C≤12),
 alkylthio_(C≤12), alkenylthio_(C≤12), alkynylthio_(C≤12),
 arylthio_(C≤12), aralkylthio_(C≤12), heteroarylthio_(C≤12),
 heteroaralkylthio_(C≤12), acylthio_(C≤12),
 alkylammonium_(C≤12), alkylsulfonium_(C≤12),
 alkylsilyl_(C≤12), or a substituted version of any of these
 groups;

X is OR_b, NR_bR_c, or SR_b, wherein R_b and R_c are each independently:

hydrogen;

alkyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), acyl_(C≤8), or a substituted
 version of any of these groups; or

provided that R_b is absent when the atom to which it is bound is
 part of a double bond, further provided that when R_b is
 absent the atom to which it is bound is part of a double
 bond; and

R₁ is:

hydrogen, cyano, hydroxy, halo or amino; or

alkyl_(C≤8), alkenyl_(C≤8), alkynyl_(C≤8), aryl_(C≤8), aralkyl_(C≤8), heteroaryl_(C≤8),
 heteroaralkyl_(C≤8), acyl_(C≤8), alkoxy_(C≤8), aryloxy_(C≤8),
 acyloxy_(C≤8), alkylamino_(C≤8), arylamino_(C≤8), amido_(C≤8), or a
 substituted version of any of these groups;

R' is hydroxy, alkoxy_(C≤12), substituted alkoxy_(C≤12), aryloxy_(C≤12), substituted
 aryloxy_(C≤12), aralkoxy_(C≤12), substituted aralkoxy_(C≤12), acyloxy_(C≤12), or
 substituted acyloxy_(C≤12);

or salts, esters, hydrates, solvates, tautomers, or optical isomers thereof.

21. The compound according to claim 1 wherein Z is a single bond.

22. The compound according to claim 1 wherein Z is —O—.

23. The compound according to claim 1 wherein Z is -NH- .
24. The compound according to any one of claims 1-3, 13, 14, and 19-23, wherein Y is cyano.
25. The compound according to any one of claims 1-3, 13, 14, and 19-23, wherein Y is -C(O)R_a .
26. The compound according to any one of claims 1-3, 13, 14, and 19-25, wherein X is OR_b .
27. The compound of claim 26, wherein R_b is absent.
28. The compound of claim 26, wherein R_b is hydrogen.
29. The compound according to any one of claims 1-3, 13, 14, and 19-25, wherein X is NR_b .
30. The compound of claim 29, wherein R_b is hydroxy.
31. The compound of claim 25, wherein R_a is hydroxy.
32. The compound of claim 25, wherein R_a is $\text{alkoxy}_{(\text{C}\leq 6)}$, $\text{aryloxy}_{(\text{C}\leq 8)}$, $\text{aralkyloxy}_{(\text{C}\leq 8)}$, or a substituted version of any of these groups.
33. The compound of claim 32, wherein R_a is $\text{alkoxy}_{(\text{C}2-6)}$.
34. The compound of claim 32, wherein R_a is $\text{alkoxy}_{(\text{C}1-5)}$ or substituted $\text{alkoxy}_{(\text{C}1-5)}$.
35. The compound of claim 34, wherein R_a is $\text{alkoxy}_{(\text{C}2-4)}$ or substituted $\text{alkoxy}_{(\text{C}2-4)}$.
36. The compound of claim 35, wherein R_a is $\text{alkoxy}_{(\text{C}1-4)}$ or substituted $\text{alkoxy}_{(\text{C}1-4)}$.
37. The compound of claim 36, wherein R_a is $\text{alkoxy}_{(\text{C}1-2)}$ or substituted $\text{alkoxy}_{(\text{C}1-2)}$.
38. The compound of claim 37, wherein R_a is methoxy.
39. The compound of claim 25, wherein R_a is amino.


40. The compound of claim 25, wherein R_a is alkylamino_(C1-6), alkoxyamino_(C1-6), arylamino_(C1-8), aralkylamino_(C1-8), dialkylamino_(C2-8), or a substituted version of any of these groups.
41. The compound of claim 40 wherein R_a is alkylamino_(C2-6) or substituted
5 alkylamino_(C2-6).
42. The compound of claim 41, wherein R_a is alkylamino_(C3-6).
43. The compound of claim 40, wherein R_a is alkylamino_(C1-5), dialkylamino_(C2-6), or substituted version of either of these groups.
44. The compound of claim 43, wherein R_a is alkylamino_(C2-4), dialkylamino_(C2-5), or
10 substituted version of either of these groups.
45. The compound of claim 43, wherein R_a is alkylamino_(C1-4) or substituted alkylamino_(C1-4).
46. The compound of claim 45, wherein R_a is alkylamino_(C1-3).
47. The compound of claim 46, wherein R_a is methylamino or ethylamino.
- 15 48. The compound of claim 43, wherein R_a is substituted alkylamino_(C1-3).
49. The compound of claim 48, wherein R_a is 2,2,2-trifluoroethylamino.
50. The compound of claim 25, wherein R_a is alkyl_(C1-5), aryl_(C≤8), aralkyl_(C≤8), heteroaralkyl_(C≤8), or a substituted version of any of these groups.
51. The compound of claim 25, wherein R_a is heteroaryl_(C1-8) or substituted heteroaryl<sub>(C1-
20 8)</sub>.
52. The compound of claim 51, wherein R_a is imidazolyl.
53. The compound of claim 25, wherein R_a is -H.
54. The compound according to any one of claims 1-3, 15, 17, 19 and 20, wherein R₁ is -H, -OH or -F.
- 25 55. The compound of claim 54, wherein R₁ is -H.

56. The compound according to any one of claims 1-5 and 13-17, wherein R_2 is $-\text{CN}$.
57. The compound according to any one of claims 1-5 and 13-17, wherein R_2 is fluoroalkyl_(C $\leq$ 8).
58. The compound of claim 57, wherein R_2 is $-\text{CF}_3$.
- 5 59. The compound according to any one of claims 1-5 and 13-17, wherein R_2 is not fluoroalkyl_(C $\leq$ 8).
60. The compound according to any one of claims 1-5 and 13-17, wherein R_2 is a substituted acyl_(C1-3).
61. The compound of claim 60, wherein R_2 is $-\text{C}(=\text{O})\text{NHS}(=\text{O})_2\text{CH}_3$.
- 10 62. The compound according to any one of claims 1, 2, 13 and 14, wherein R_3 is hydrogen or acetyl.
63. The compound according to any one of claims 1, 2, 13 and 14, wherein R_3 is absent.
64. The compound according to any one of claims 1-3, 13 and 15, wherein R_4 is methyl.
65. The compound according to any one of claims 1-3, 13 and 15, wherein R_4 is hydroxy-
15 methyl.
66. The compound according to any one of claims 1 and 2, wherein R_6 and R_7 are each hydrogen.
67. The compound of claim 1, wherein R_8 and R_9 are each hydrogen.
68. The compound of claim 1, wherein R_{10} and R_{11} are each methyl.
- 20 69. The compound of claim 20, wherein R' is acetyloxy.
70. The compound of claim 20, wherein R' is hydroxy.
71. The compound according to any one of claims 1 and 11, wherein R_d is alkyl_(C1-5), aryl_(C $\leq$ 8), aralkyl_(C $\leq$ 8), heteroaralkyl_(C $\leq$ 8), or a substituted version of any of these groups
72. The compound of claim 71, wherein R_d is alkyl_(C1-4) or a substituted version thereof.

73. The compound of claim 72, wherein R_d is alkyl_(C1-3) or a substituted version thereof.


74. The compound of claim 73, wherein R_d is alkyl_(C1-2) or a substituted version thereof.

75. The compound of claim 6, further defined as:




5 or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

76. The compound of claim 75, further defined as:




10 or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

77. The compound of claim 6, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

78. The compound of claim 77, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

5 79. The compound of claim 7, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

80. The compound of claim 79, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

81. The compound of claim 7, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

5 82. The compound of claim 81, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

83. The compound of claim 6, wherein the compound is:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

84. The compound of claim 83, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

5 85. The compound of claim 8, wherein the compound is:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

86. The compound of claim 85, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

87. The compound of claim 7, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

5 88. The compound of claim 87, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

89. The compound of claim 6, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

90. The compound of claim 89, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

5 91. The compound of claim 6, wherein the compound is:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

92. The compound of claim 91, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

93. The compound of claim 6, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

5 94. The compound of claim 29, further defined as:



or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.


95. The compound of claim 94, further defined as:



10


or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

96. The compound of claim 18, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

5 97. The compound of claim 96, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

98. The compound of claim 96, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

99. The compound of claim 96, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

5 100. The compound of claim 16, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

101. The compound of claim 100, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

102. The compound of claim 14, wherein the compound is:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

5 103. The compound of claim 102, further defined as:




or pharmaceutically acceptable salts thereof, and substantially free from other optical isomers thereof.

104. The compound of claim 11, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

105. The compound of claim 2, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

5 106. The compound of claim 12, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

107. The compound of claim 12, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

108. The compound of claim 12, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

5 109. The compound of claim 12, further defined as:




or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

110. The compound of claim 22, further defined as:



or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

111. The compound of claim 23, further defined as:



or pharmaceutically acceptable salts, hydrates, solvates, tautomers, or optical isomers thereof.

5 112. A compound selected from the group consisting of:

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,14a*R*,14b*S*)-methyl 11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydropicene-4a-carboxylate,

10 (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,14a*R*,14b*S*)-methyl 11-cyano-10-hydroxy-2,2,6a,6b,9,9,12a-heptamethyl-14-oxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,12,12a,12b,13,14,14a,14b-icosahydropicene-4a-carboxylate,

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,14a*R*,14b*S*)-11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydropicene-4a-carboxylic acid,

15 (4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14b*R*)-4,4,6a,6b,11,11,14b-heptamethyl-3,13-dioxo-3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydropicene-2,8a-dicarbonitrile,

(4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,14a*R*,14b*S*)-11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydropicene-4a-carboxamide,


20 (4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14b*R*)-8a-formyl-4,4,6a,6b,11,11,14b-heptamethyl-3,13-dioxo-3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydropicene-2-carbonitrile,

25 (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,14a*R*,14b*S*)-methyl 11-cyano-14-hydroxy-2,2,6a,6b,9,9,12a-heptamethyl-10-oxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydropicene-4a-carboxylate,

- (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,14a*R*,14b*S*)-11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-N-(2,2,2-trifluoroethyl)-
1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxamide,
- 5 (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14*R*,14a*R*,14b*S*)-2,2,2-trifluoroethyl 11-cyano-14-hydroxy-2,2,6a,6b,9,9,12a-heptamethyl-10-oxo-
1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxylate,
- 10 (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-N-(2,2,2-trifluoroethyl)-
1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxamide,
- (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-N'-acetyl-11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-
15 1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carbohydrazide,
- (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-N-(2,2,2-trifluoroethyl)docosahydronicene-4a-carboxamide,
- (4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14a*R*,14b*R*)-4,4,6a,6b,11,11,14b-heptamethyl-3,13-dioxo-8a-(2H-tetrazol-5-yl)-
20 3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydronicene-2-carbonitrile,
- (4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14a*R*,14b*R*)-4,4,6a,6b,11,11,14b-heptamethyl-8a-(2-methyl-2H-tetrazol-5-yl)-3,13-dioxo-
25 3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydronicene-2-carbonitrile,
- (4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14a*R*,14b*R*)-4,4,6a,6b,11,11,14b-heptamethyl-8a-(5-methyl-1,3,4-oxadiazol-2-yl)-3,13-dioxo-
30 3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydronicene-2-carbonitrile,
- (4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14a*R*,14b*R*)-4,4,6a,6b,11,11,14b-heptamethyl-8a-(5-methyl-1,3,4-thiadiazol-2-yl)-3,13-dioxo-
3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydronicene-2-carbonitrile,


- (4a*R*,6a*R*,6b*R*,8a*S*,12a*S*,12b*R*,14a*R*,14b*R*)-8a-acetyl-4,4,6a,6b,11,11,14b-heptamethyl-3,13-dioxo-3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,14a,14b-icosahydronicene-2-carbonitrile,
- 5 (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-benzyl 11-cyano-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxylate,
- 10 (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-11-cyano-*N,N*,2,2,6a,6b,9,9,12a-nonamethyl-10,14-dioxo-1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxamide,
- (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-11-cyano-*N*-methoxy-2,2,6a,6b,9,9,12a-heptamethyl-10,14-dioxo-
- 15 1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxamide
- (4a*S*,6a*R*,6b*R*,8a*R*,12a*R*,12b*R*,14a*R*,14b*S*)-methyl 11-cyano-14-(hydroxyimino)-2,2,6a,6b,9,9,12a-heptamethyl-10-oxo-
- 20 1,2,3,4,4a,5,6,6a,6b,7,8,8a,9,10,12a,12b,13,14,14a,14b-icosahydronicene-4a-carboxylate,
- (4a*R*,6a*R*,6b*S*,8a*S*,12a*R*,15a*R*,15b*R*)-methyl 2-cyano-14-hydroxy-4,4,6a,6b,11,11,15b-heptamethyl-3-oxo-
- 3,4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,14,15,15a,15b-icosahydrodinaphtho[1,2-*b*:2',1'-*d*]oxepine-8a-carboxylate, and
- 25 (4a*R*,6a*R*,6b*S*,8a*S*,12a*R*,12b*R*,15a*R*,15b*R*)-methyl 2-cyano-4,4,6a,6b,11,11,15b-heptamethyl-3,14-dioxo-
- 4,4a,5,6,6a,6b,7,8,8a,9,10,11,12,12a,12b,13,14,15,15a,15b-icosahydro-3*H*-dinaphtho[1,2-*b*:2',1'-*d*]azepine-8a-carboxylate.

113. The compound of claim 19, further defined as:




or salts, hydrates, solvates, tautomers, or optical isomers thereof.

114. The compound of claim 113, further defined as:




substantially free from other optical isomers thereof.

115. The compound of claim 19, further defined as:




or salts, hydrates, solvates, tautomers, or optical isomers thereof.

10 116. The compound of claim 20, further defined as:



or salts, hydrates, solvates, tautomers, or optical isomers thereof.

117. The compound of claim 116, further defined as:



substantially free from other optical isomers thereof.

118. The compound according to any one of claims 1-74, 75, 77, 83, 87, 91, 96, 100, 102, 104, 105, 106, 107 and 109, wherein the compound is in the form of a pharmaceutically acceptable salt.

119. The compound according to any one of claims 1-74, 75, 77, 83, 87, 91, 96, 100, 102, 104, 105, 106, 107, 109, 113, 115 and 116, wherein the compound is not a salt.

120. The compound according to any one of claims 1-74, 75, 77, 83, 87, 91, 96, 100, 102, 104, 105, 106, 107, 109, 113, 115 and 116, wherein the compound is a hydrate.

121. The compound according to any one of claims 1-74, 75, 77, 83, 87, 91, 96, 100, 102, 104, 105, 106, 107, 109, 113, 115 and 116, wherein the compound is not a hydrate.

122. The compound according to any one of claims 1-74, 75, 77, 83, 87, 91, 96, 100, 102, 104, 105, 106, 107, 109, 113, 115 and 116, wherein the compound is a solvate.

123. The compound according to any one of claims 1-74, 75, 77, 83, 87, 91, 96, 100, 102, 104, 105, 106, 107, 109, 113, 115 and 116, wherein the compound is not a solvate.

124. The compound according to any one of claims 1-7 and 13-20 wherein the compound is an ester thereof.

125. The compound of claim 124, wherein the ester results from a condensation reaction between a hydroxy group of the formula and the carboxylic acid group of biotin.

126. The compound according to any one of claims 1-7 and 13-20 wherein the compound is not an ester thereof.

127. The compound according to any one of claims 1-74, 75, 77, 83, 87, 91, 96, 100, 102, 104, 105, 106, 107, 109, 113, 115 and 116, wherein the compound is present as a mixture of stereoisomers.
- 5 128. The compound according to any one of claims 1-74, 75, 77, 83, 87, 91, 96, 100, 102, 104, 105, 106, 107, 109, 113, 115 and 116, wherein the compound is predominantly present as a single stereoisomer.
129. The compound of any of claims 1-112, wherein the compound is effective for inhibiting IFN- γ -induced NO production in macrophages, further wherein the compound has an IC₅₀ value of less than 0.2 μ M.
- 10 130. A pharmaceutical composition comprising as an active ingredient a compound according to any one of claims 1-112 and a pharmaceutically acceptable carrier.
131. The pharmaceutical composition of claim 130, wherein the composition is adapted for administration by a route selected from the group consisting of orally, intraadiposally, intraarterially, intraarticularly, intracranially, intradermally, intralesionally, 15 intramuscularly, intranasally, intraocularly, intrapericardially, intraperitoneally, intrapleurally, intraprostatically, intrarectally, intrathecally, intratracheally, intratumorally, intraumbilically, intravaginally, intravenously, intravesicularly, intravitreally, liposomally, locally, mucosally, orally, parenterally, rectally, subconjunctival, subcutaneously, sublingually, topically, transbuccally, transdermally, 20 vaginally, in crèmes, in lipid compositions, via a catheter, via a lavage, via continuous infusion, via infusion, via inhalation, via injection, via local delivery, via localized perfusion, bathing target cells directly, or any combination thereof.
132. The composition of claim 131, wherein the composition is formulated for oral delivery.
- 25 133. The composition of claim 132, wherein the composition is formulated as a hard or soft capsule, a tablet, a syrup, a suspension, a wafer, or an elixir.
134. The composition of claim 132, wherein the composition is prepared as a solid dispersion.
135. The composition of claim 133, wherein the soft capsule is a gelatin capsule.

136. The composition of claim 132, further comprising a protective coating.
137. The composition of claim 132, further comprising an agent that delays absorption.
138. The composition of claim 132, further comprising an agent that enhances solubility or dispersibility.
- 5 139. The composition of claim 130, wherein the compound is dispersed in a liposome, an oil in water emulsion or a water in oil emulsion.
140. A therapeutic method comprising administering a pharmaceutically effective amount of a compound of any one of claims 1-112 to a subject.
141. The method of claim 140, wherein the subject is a human.
- 10 142. The method of claim 140, further comprising identifying a subject in need of treatment.
143. A method of treating cancer in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound according to any of claims 1-112.
144. The method of claim 143, wherein the cancer is a carcinoma, sarcoma, lymphoma, leukemia, melanoma, mesothelioma, multiple myeloma, or seminoma.
- 15 145. The method of claim 143, wherein the cancer is of the bladder, blood, bone, brain, breast, central nervous system, colon, endometrium, esophagus, genitourinary tract, head, larynx, liver, lung, neck, ovary, pancreas, prostate, spleen, small intestine, large intestine, stomach, or testicle.
- 20 146. The method of claim 143, wherein the subject is a primate.
147. The method of claim 143, wherein the subject is a human.
148. The method of claim 143, further comprising identifying a subject in need of treatment.
- 25 149. The method of claim 148, wherein the subject has a family or patient history of cancer.

150. The method of claim 143, wherein the subject has symptoms of cancer.
151. The method of claim 143, wherein the compound is administered locally.
152. The method of claim 151, wherein the compound is administered by direct intratumoral injection or by injection into tumor vasculature.
- 5 153. The method of claim 143, wherein the compound is administered systemically.
154. The method of claim 153, wherein the compound is administered intravenously, intra-arterially, intramuscularly, intraperitoneally, subcutaneously or orally.
155. The method of claim 143, wherein the compound is administered by contacting a tumor cell during *ex vivo* purging.
- 10 156. The method of claim 143, wherein the method comprises:
- a) inducing cytotoxicity in a tumor cell;
 - b) killing a tumor cell;
 - c) inducing apoptosis in a tumor cell;
 - d) inducing differentiation in a tumor cell; or
 - 15 e) inhibiting growth in a tumor cell.
157. The method of claim 156, wherein the tumor cell is a leukemia cell.
158. The method of claim 156, wherein the tumor cell is a bladder cancer cell, a breast cancer cell, a lung cancer cell, a colon cancer cell, a prostate cancer cell, a liver cancer cell, a pancreatic cancer cell, a stomach cancer cell, a testicular cancer cell, a brain cancer cell, an ovarian cancer cell, a lymphatic cancer cell, a skin cancer cell, a brain cancer cell, a bone cancer cell, or a soft tissue cancer cell.
- 20 159. The method of claim 143, further comprising a treatment selected from the group consisting of administering a pharmaceutically effective amount of a second drug, radiotherapy, gene therapy, and surgery.
- 25 160. The method of claim 159, further comprising (1) contacting a tumor cell with the compound prior to contacting the tumor cell with the second drug, (2) contacting a tumor cell with the second drug prior to contacting the tumor cell with the compound,

or (3) contacting a tumor cell with the compound and the second drug at the same time.

161. The method of claim 159, wherein the second drug is an antibiotic, anti-inflammatory, anti-neoplastic, anti-proliferative, anti-viral, immunomodulatory, or immunosuppressive.

162. The method of claim 159, wherein the second drug is an alkylating agent, androgen receptor modulator, cytoskeletal disruptor, estrogen receptor modulator, histone-deacetylase inhibitor, HMG-CoA reductase inhibitor, prenyl-protein transferase inhibitor, retinoid receptor modulator, topoisomerase inhibitor, or tyrosine kinase inhibitor.

163. The method of claim 159, wherein the second drug is 5-azacitidine, 5-fluorouracil, 9-cis-retinoic acid, actinomycin D, alitretinoin, all-trans-retinoic acid, annamycin, axitinib, belinostat, bevacizumab, bexarotene, bosutinib, busulfan, capecitabine, carboplatin, carmustine, CD437, cediranib, cetuximab, chlorambucil, cisplatin, cyclophosphamide, cytarabine, dacarbazine, dasatinib, daunorubicin, decitabine, docetaxel, dolastatin-10, doxifluridine, doxorubicin, doxorubicin, epirubicin, erlotinib, etoposide, etoposide, gefitinib, gemcitabine, gemtuzumab ozogamicin, hexamethylmelamine, idarubicin, ifosfamide, imatinib, irinotecan, isotretinoin, ixabepilone, lapatinib, LBH589, lomustine, mechlorethamine, melphalan, mercaptopurine, methotrexate, mitomycin, mitoxantrone, MS-275, neratinib, nilotinib, nitrosourea, oxaliplatin, paclitaxel, plicamycin, procarbazine, semaxanib, semustine, sodium butyrate, sodium phenylacetate, streptozotocin, suberoylanilide hydroxamic acid, sunitinib, tamoxifen, teniposide, thiopeta, tioguanine, topotecan, TRAIL, trastuzumab, tretinoin, trichostatin A, valproic acid, valrubicin, vandetanib, vinblastine, vincristine, vindesine, or vinorelbine.

164. A method of treating or preventing a disease with an inflammatory component in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound of any of claims 1-112.

165. The method of claim 164, wherein the disease is lupus or rheumatoid arthritis.

166. The method of claim 164, wherein the disease is an inflammatory bowel disease.

167. The method of claim 166, wherein the inflammatory bowel disease is Crohn's disease or ulcerative colitis.
168. The method of claim 164, wherein the disease with an inflammatory component is a cardiovascular disease.
- 5 169. The method of claim 164, wherein the disease with an inflammatory component is diabetes.
170. The method of claim 169, wherein the diabetes is type 1 diabetes.
171. The method of claim 169, wherein the diabetes is type 2 diabetes.
172. The method of claim 169, wherein the method also effectively treats one or more
10 complications associated with diabetes.
173. The method of claim 172, wherein the complications are selected from the group consisting of obesity, hypertension, atherosclerosis, coronary heart disease, stroke, peripheral vascular disease, hypertension, nephropathy, neuropathy, myonecrosis, retinopathy and metabolic syndrome (syndrome X).
- 15 174. The method of claim 164, wherein the disease with an inflammatory component is metabolic syndrome (syndrome X).
175. The method of claim 164, wherein the disease with an inflammatory component is a skin disease.
176. The method of claim 175, wherein the administration is topical or oral.
- 20 177. The method of claim 175, wherein the skin disease is psoriasis, acne, or atopic dermatitis.
178. A method of treating or preventing a cardiovascular disease in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound of any of claims 1-112.
- 25 179. The method of claim 178, wherein the cardiovascular disease is atherosclerosis, cardiomyopathy, congenital heart disease, congestive heart failure, myocarditis,

rheumatic heart disease, valve disease, coronary artery disease, endocarditis, or myocardial infarction.

180. The method of claim 178, further comprising administering a pharmaceutically effective amount of a second drug.

5 181. The method of claim 180, wherein the second drug is a cholesterol lowering drug, an anti-hyperlipidemic, a calcium channel blocker, an anti-hypertensive, or an HMG-CoA reductase inhibitor.

182. The method of claim 181, wherein the second drug is amlodipine, aspirin, ezetimibe, felodipine, lacidipine, lercanidipine, nicardipine, nifedipine, nimodipine, nisoldipine
10 or nitrendipine.

183. The method of claim 181, wherein the second drug is atenolol, bucindolol, carvedilol, clonidine, doxazosin, indoramin, labetalol, methyldopa, metoprolol, nadolol, oxprenolol, phenoxybenzamine, phentolamine, pindolol, prazosin, propranolol, terazosin, timolol or tolazoline.

15 184. The method of claim 180, wherein the second drug is a statin.

185. The method of claim 184, wherein the statin is atorvastatin, cerivastatin, fluvastatin, lovastatin, mevastatin, pitavastatin, pravastatin, rosuvastatin or simvastatin.

186. A method of treating or preventing a neurodegenerative disease in a subject, comprising administering to the subject a pharmaceutically effective amount of a
20 compound of any of claims 1-112.

187. The method of claim 186, wherein said neurodegenerative disease is selected from the group consisting of Parkinson's disease, Alzheimer's disease, multiple sclerosis (MS), Huntington's disease and amyotrophic lateral sclerosis.

188. The method of claim 187, wherein said neurodegenerative disease is Alzheimer's
25 disease.

189. The method of claim 187, wherein said neurodegenerative disease is MS.


190. The method of claim 189, wherein the MS is primary progressive, relapsing-remitting secondary progressive or progressive relapsing.
191. The method of claim 189, wherein the treatment suppresses the demyelination of neurons in the subject's brain or spinal cord.
- 5 192. The method of claim 191, wherein the treatment suppresses inflammatory demyelination.
193. The method of claim 189, wherein the treatment suppresses the transection of neuron axons in the subject's brain or spinal cord.
194. The method of claim 189, wherein the treatment suppresses the transection of neurites
10 in the subject's brain or spinal cord.
195. The method of claim 189, wherein the treatment suppresses neuronal apoptosis in the subject's brain or spinal cord.
196. The method of claim 189, wherein the treatment stimulates the remyelination of neuron axons in the subject's brain or spinal cord.
- 15 197. The method of claim 189, wherein the treatment restores lost function after an MS attack.
198. The method of claim 189, wherein the treatment prevents a new MS attack.
199. The method of claim 189, wherein the treatment prevents a disability resulting from an MS attack.
- 20 200. The method of claim 189, wherein the subject is a primate.
201. The method of claim 200, wherein the primate is a human.
202. A method of treating or preventing a disorder characterized by overexpression of iNOS genes in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound according to any of claims 1-112.

203. A method of inhibiting IFN- γ -induced nitric oxide production in cells of a subject, comprising administering to said subject a pharmaceutically effective amount of a compound of any of claims 1-112.
- 5 204. A method of treating or preventing a disorder characterized by overexpression of COX-2 genes in a subject, comprising administering to the subject a pharmaceutically effective amount of compound of any of claims 1-112.
- 10 205. A method for treating renal/kidney disease (RKD), insulin resistance, diabetes, endothelial dysfunction, fatty liver disease, or cardiovascular disease (CVD) in a subject comprising, administering to said subject a pharmaceutically effective amount of a compound of any of claims 1-112.
206. The method of claim 205, wherein the method treats RKD.
207. The method of claim 206, wherein the RKD is diabetic nephropathy (DN).
208. The method of claim 206, wherein the RKD results from a toxic insult.
- 15 209. The method of claim 208, wherein the toxic insult results from an imaging agent or a drug.
210. The method of claim 209, wherein the drug is a chemotherapeutic.
211. The method of claim 206, wherein the RKD results from ischemia/reperfusion injury.
212. The method of claim 206, wherein the RKD results from diabetes or hypertension.
213. The method of claim 206, wherein the RKD results from an autoimmune disease.
- 20 214. The method of claim 206, wherein the RKD is chronic RKD.
215. The method of claim 206, wherein the RKD is acute RKD.
216. The method of claim 205, wherein the subject has undergone or is undergoing dialysis.
- 25 217. The method of claim 205, wherein the subject has undergone or is a candidate to undergo kidney transplant.


218. The method of claim 205, wherein the method treats RKD and insulin resistance.
219. The method of claim 218, wherein the method treats RKD, insulin resistance and endothelial dysfunction.
220. The method of claim 205, wherein the method treats RKD and diabetes.
- 5 221. The method of claim 205, wherein the method treats insulin resistance.
222. The method of claim 205, wherein the method treats diabetes.
223. The method of claim 222, wherein the method also effectively treats one or more complications associated with diabetes.
- 10 224. The method of claim 223, wherein the complications are selected from the group consisting of obesity, hypertension, atherosclerosis, coronary heart disease, stroke, peripheral vascular disease, hypertension, nephropathy, neuropathy, myonecrosis, retinopathy and metabolic syndrome (syndrome X).
225. The method of claim 224, wherein the complication is metabolic syndrome (syndrome X).
- 15 226. The method of claim 222, wherein the diabetes results from insulin resistance.
227. The method of claim 205, wherein the method treats RKD and endothelial dysfunction.
228. The method of claim 205, wherein the method treats RKD and cardiovascular disease.
229. The method of claim 205, wherein the method treats CVD.
- 20 230. The method of claim 229, wherein the CVD results from endothelial dysfunction.
231. The method of claim 205, wherein the method treats endothelial dysfunction.
232. The method of claim 231, wherein the method treats endothelial dysfunction and insulin resistance.
233. The method of claim 205, wherein the method treats fatty liver disease.

234. The method of claim 233, wherein the fatty liver disease is non alcoholic fatty liver disease.
235. The method of claim 233, wherein the fatty liver disease is alcoholic fatty liver disease.
- 5 236. The method of claim 233, wherein the method treats fatty liver disease and one or more of the following disorders: renal/kidney disease (RKD), insulin resistance, diabetes, endothelial dysfunction, and cardiovascular disease (CVD).
- 10 237. The method according to any of claims 205-236, further comprising identifying a subject in need of treatment of any of the listed diseases, dysfunctions, resistances or disorders.
238. The method according to any of claims 205-236, wherein the subject has a family or patient history of any of the listed diseases, dysfunctions, resistances or disorders.
239. The method according to any of claims 205-236, wherein the subject has symptoms of any of the listed diseases, dysfunctions, resistances or disorders.
- 15 240. A method for improving glomerular filtration rate or creatinine clearance in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound of any of claims 1-112.
241. The method according to any of claims 140-240, wherein the pharmaceutically effective amount is a daily dose of 25 mg to 500 mg of the compound.
- 20 242. The method according to any of claims 140-240, wherein the pharmaceutically effective amount is a daily dose is 0.1 – 1000 mg of compound per kg of body weight.
243. The method of claim 242, wherein the daily dose is 0.15 – 20 mg of compound per kg of body weight.
244. The method of claim 243, wherein the daily dose is 0.20 – 10 mg of compound per kg
25 of body weight.
245. The method of claim 244, wherein the daily dose is 0.40 – 3 mg of compound per kg of body weight.

246. The method according to any of claims 140-240, wherein the pharmaceutically effective amount is administered in a single dose per day.
247. The method according to any of claims 140-240, wherein the pharmaceutically effective amount is administered in two or more doses per day.
- 5 248. A kit comprising:
a compound of any one of claims 1-112; and
instructions which comprise one or more forms of information selected from the
group consisting of indicating a disease state for which the compound is to be
administered, storage information for the compound, dosing information and
10 instructions regarding how to administer the compound.
249. The kit according to claim 248, wherein the kit comprises the compound in a multiple dose form.
250. An article of manufacture comprising:
a compound of any one of claims 1-112; and
15 packaging materials.
251. The article of manufacture according to claim 250, wherein the packaging materials comprise a container for housing the compound.
252. The article of manufacture according to claim 251, wherein the container comprises a label indicating one or more members of the group consisting of a disease state for
20 which the compound is to be administered, storage information, dosing information and/or instructions regarding how to administer the compound.
253. The article of manufacture according to claim 250, wherein the article of manufacture comprises the compound in a multiple dose form.
254. A method of making a target compound defined of the formula:



wherein R_a is alkoxy_(C1-4), comprising reacting a compound of the formula:



with an oxidizing agent under a set of conditions to form the target compound.