

FRACTION I-IV-1 PRECIPITATION OF IMMUNOGLOBINS FROM PLASMA

CROSS REFERENCES TO APPLICATIONS

[0001] This application claims priority to U.S. Provisional Patent Application Serial No. 61/602,488 filed February 23, 2012.

BACKGROUND OF THE INVENTION

[0002] Immune globulin products from human plasma were first used in 1952 to treat immune deficiency. Initially, intramuscular or subcutaneous administration of immunoglobulin isotype G (IgG) isolated from plasma were the methods of choice. However, these methods did not allow for the administration of larger amounts of IgG necessary for effective treatment of various diseases. Accordingly, IgG products that could be administered intravenously were developed. Usually, intravenous immunoglobulin (IVIG) contains the pooled immunoglobulin G (IgG) immunoglobulins from the plasma of more than a thousand blood donors. The preparations usually contain more than 95% unmodified IgG, which has intact Fc-dependent effector functions and only trace amounts of immunoglobulin A (IgA) and immunoglobulin M (IgM). Typically, IVIGs are sterile filtered and the manufacturing process contains steps to inactivate and/or remove viruses. These purified IgG products are primarily used in treating three main categories of medical conditions: (1) immune deficiencies: X-linked agammaglobulinemia, hypogammaglobulinemia (primary immune deficiencies), and acquired compromised immunity conditions (secondary immune deficiencies), featuring low antibody levels; (2) inflammatory and autoimmune diseases; and (3) acute infections.

[0003] Specifically, many people with primary immunodeficiency disorders lack antibodies needed to resist infection. In certain cases these deficiencies can be supplemented by the infusion of purified IgG, commonly through intravenous administration (*i.e.*, IVIG therapy). Several primary immunodeficiency disorders are commonly treated in the fashion, including X-linked Agammaglobulinemia (XLA), Common Variable Immunodeficiency (CVID), Hyper-IgM Syndrome (HIM), Severe Combined Immunodeficiency (SCID), and some IgG subclass

deficiencies (Blaese and Winkelstein, *J. Patient & Family Handbook for Primary Immunodeficiency Diseases*. Towson, MD: Immune Deficiency Foundation; 2007).

[0004] While IgG treatment can be very effective for managing primary immunodeficiency disorders, this therapy is only a temporary replacement for antibodies that are not being produced in the body, rather than a cure for the disease. Accordingly, patients depend upon repeated doses of IgG therapy, typically about once a month for life. This therapy places a great demand on the continued production of IgG compositions. However, unlike other biologics that are produced via *in vitro* expression of recombinant DNA vectors, IgG is fractionated from human blood and plasma donations. Thus, the level of commercially available IgG is limited by the available supply of blood and plasma donations.

[0005] Several factors drive the demand for IgG, including the acceptance of IgG treatments, the identification of additional indications for which IgG therapy is effective, and increasing patient diagnosis and IgG prescription. Notably, the global demand for IgG more than quadrupled between 1990 and 2009, and continues to increase at an annual rate between about 7% and 10% (Robert P., *Pharmaceutical Policy and Law*, 11 (2009) 359-367). For example, the Australian National Blood Authority reported that the demand for IgG in Australia grew by 10.6% for the 2008-2009 fiscal year (National Blood Authority Australia Annual Report 2008-2009).

[0006] Due in part to the increasing global demand and fluctuations in the available supply of immunoglobulin products, several countries, including Australia and England, have implemented demand management programs to protect supplies of these products for the highest demand patients during times of product shortages.

[0007] It has been reported that in 2007, 26.5 million liters of plasma were fractionated, generating 75.2 metric tons of IgG, with an average production yield of 2.8 grams per liter (Robert P., *supra*). This same report estimated that global IgG yields are expected to increase to about 3.43 grams per liter by 2012. However, due to the continued growth in global demand for IgG, projected at between about 7% and 13% annually between now and 2015, further improvement of the overall IgG yield will be needed to meet global demand.

[0008] A number of IgG preparation methods are used by commercial suppliers of IgG products. A common problem with the current IgG production methods is the substantial loss of IgG

during the purification process, estimated to be at least 30% to 35% of the total IgG content of the starting material. One challenge is to maintain the quality of viral inactivation and removal of impurities which can cause adverse reactions, while improving the process efficiency to increase the yield of IgG. At the current production levels of IgG, what may be considered small increases in the yield are in fact highly significant. For example at 2007 production levels, a 2% increase in efficiency, equal to an additional 56 milligrams per liter, would generate 1.5 additional metric tons of IgG.

[0009] In the fourth installment of a series of seminal papers published on the preparation and properties of serum and plasma proteins, Cohn *et al.* (J. Am. Chem. Soc., 1946, 68(3): 459-475) first described a method for the alcohol fractionation of plasma proteins (method 6), which allows for the isolation of a fraction enriched in IgG from human plasma. Several years later, Oncley *et al.* (J. Am. Chem. Soc., 1949, 71(2): 541-550) expanded upon the Cohn methods by publishing a method (method 9) that resulted in the isolation of a purer IgG preparation.

[0010] These methods, while laying the foundation for an entire industry of plasma derived blood proteins, were unable to provide IgG preparations having sufficiently high purity for the treatment of several immune-related diseases, including Kawasaki syndrome, immune thrombocytopenic purpura, and primary immune deficiencies. As such, additional methodologies employing various techniques, such as ion exchange chromatography, were developed to provide higher purity IgG formulations. Hoppe *et al.* (Munch Med Wochenschr 1967 (34): 1749-1752), Falksveden (Swedish Patent No. 348942), and Falksveden and Lundblad (Methods of Plasma Protein Fractionation 1980) were among the first to employ ion exchange chromatography for this purpose.

[0011] Various modern methods for the purification of immunoglobulins from plasma employ a precipitation step, such as caprylate precipitation (Lebing *et al.*, Vox Sang 2003 (84):193-201) and Cohn Fraction (I+II+III) ethanol precipitation (Tanaka *et al.*, Braz J Med Biol Res 2000 (33)37-30) coupled to column chromatography. Most recently, Teschner *et al.* (Vox Sang, 2007 (92):42-55) have described a method for production of a 10% IgG product in which cryo-precipitate is first removed from pooled plasma and then a modified Cohn-Oncley cold ethanol fractionation is performed, followed by S/D treatment of the intermediate, ion exchange chromatography, nanofiltration, and optionally ultrafiltration/diafiltration.

[0012] Despite the purity, safety, and yield afforded by these IgG isolation methods, the yield of IgG recovered from plasma can still be improved. For example, Teschner *et al.* report that their method results in an increased IgG yield of 65% (Teschner *et al.*, supra). As reported during various plasma product meetings, the average yields for large-scale preparation of IgG, such as from Baxter, CSL Behring, Upfront Technology, Cangene, Prometric BioTherapeutics, and the Finnish Red Cross, range from about 61% to about 65% in the final container. Although better than methods previously employed, this amount of IgG recovery still represents a loss of at least about a third of the IgG present in the pooled plasma fraction during the isolation process.

[0013] Due to the limited supply of plasma available for the manufacture of plasma-derived products, the isolation of several blood proteins from a common starting plasma pool can be achieved by integrating the purifications into a single framework. For example, IgG is commonly enriched through the formation of a Cohn Fraction II+III precipitate or Kistler-Nitschmann precipitate A, the corresponding supernatants of which are then used for the manufacture of alpha-1-antitrypsin (A1PI) and albumin. Similarly, several methods have been described for the manufacture of Factor H from by-products formed during the manufacture of IgG immunoglobulins, including WO 2008/113589 and WO 2011/011753.

[0014] As such, a need exists for improved and more efficient methods for manufacturing therapeutic IgG products. Furthermore, these methods should also allow for the manufacture of additional plasma-derived products from a single plasma source. The present invention satisfies these and other needs by providing IgG isolation methods that produce yields that are approximately 20 to 25% higher than currently achievable, as well as IgG compositions provided there from. Advantageously, the methods provided herein allow for the co-isolation of other therapeutically important plasma-derived proteins, including alpha-1-antitrypsin (A1PI), Factor H, inter-alpha-inhibitor proteins (IaIp), Prothrombin complexes, Factor VII (FVII), Factor VIII (FVIII), antithrombin III (ATIII), fibrinogen, butyrylcholinesterase, and others.

BRIEF SUMMARY OF INVENTION

[0015] Current methods for the production of IVIG and alpha-1-antitrypsin (A1PI) rely on multiple protein precipitation steps to separate immunoglobulin IgG and A1PI from other components found in human plasma. For example, many manufactures use a variation of the

Cohn-Oncley method 6, in which three initial alcohol precipitation steps are employed. The first precipitation step, referred to as Fraction I precipitation, is performed at high pH (7.2) and low ethanol concentration (8-10% v/v) to precipitate protein such as fibrinogen and Factor XIII away from IgG and A1PI, which remain in the supernatant. IgG is then precipitated from the Fraction I supernatant in a second precipitation reaction, referred to as a Fraction II+III precipitation, performed at moderate pH (6.8) and high ethanol concentration (20%-25%). The bulk of the A1PI remains in the supernatant of the Fraction II+III precipitation reaction, as is subsequently separated from albumin in a third initial precipitation reaction, referred to as a Fraction I-IV-1 precipitation, performed at low pH (5.2) and moderate ethanol concentration (18%).

[0016] Unfortunately, because the Cohn-Oncley method described above, as well as the comparable Kistler-Nitschmann process that employs four initial precipitations to fractionate IgG and A1PI, separate individual components in an intricate series of precipitation reactions, the fractionation is inefficient. A significant loss in the yield of IgG and A1PI, in these initial precipitation steps, can be attributed to the partial precipitation into non-targeted fractions as well as incomplete precipitation in the targeted fractions. For example, IgG co-precipitates to some extent with fibrinogen and Factor XIII in the Fraction I precipitate and some IgG is not precipitated by Fraction II+III precipitation. After clarification of a dissolved Fraction II+III precipitate, the IgG yield is typically between 75% and 85% of the IgG present in the starting Cohn pool. Accordingly, 15% to 25% of the total IgG content of the starting material is lost after these two fractionation steps.

[0017] The present disclosure improves upon the recovery of IgG and A1PI from pooled plasma by removing the need for multiple initial precipitation steps. Rather, the methods described herein rely on a single initial precipitation step that captures all of the proteins normally precipitated in the Fraction I, Fraction II+III, and Fraction IV-1 precipitates combined. This single precipitation step is referred herein as a "Fraction I+II+III+IV-1 precipitation," a "Fraction I-IV-1 precipitation," or an "initial low pH, high alcohol precipitation." Advantageously, it was found that IgG and A1PI could be efficiently extracted from the Fraction I-IV-1 precipitate without the use of subsequent protein precipitations. Furthermore, it was found that the Fraction I-IV-1 precipitate contained nearly all of the IgG and A1PI content of the source plasma, while

albumin remains in the supernatant. Taken together, these advantages result in a significant increase in the overall recovery of these important commercial products.

[0018] As shown in the Examples, the use of a low pH, high alcohol precipitation step (Fraction I-IV-1 precipitation) as an initial step in the purification of IgG from cryo-poor plasma, allows for the production of pharmaceutical grade IgG compositions with unprecedented yields of 4.3-4.7 g IgG/L source plasma. These yields represent an approximate 20% to 25% increase in yield as compared to state-of-the-art manufacturing processes, such as those used to manufacture GAMMAGARD LIQUID[®] (Baxter International; Deerfield, IL) from the same plasma type.

[0018a] In one aspect, the present invention relates to a method for manufacturing an enriched immunoglobulin composition from a Cohn pool, the method comprising the steps of: (a) co-precipitating immunoglobulins and alpha-1-antitrypsin (A1PI) from a Cohn pool, in a first precipitation step by the addition of ethanol to the Cohn pool at a final concentration of from 20% to 30% (v/v) at a pH of from 5.0 to 6.0, to form a first precipitate and a first supernatant; (b) solubilizing immunoglobulins in the first precipitate, to form a first suspension having a soluble portion comprising immunoglobulins and an insoluble portion comprising A1PI; (c) separating the soluble portion of the first suspension from the insoluble portion of the first suspension; and (d) recovering the soluble fraction of the first suspension, thereby forming an enriched immunoglobulin composition, wherein no alcohol precipitation is performed prior to step (a).

[0018b] In another embodiment, there is provided an enriched immunoglobulin composition prepared by a method disclosed herein.

[0018c] In another embodiment, there is provided the use of an enriched immunoglobulin composition as disclosed herein in the preparation of a medicament for treating an immune deficiency, an inflammatory disease, an autoimmune disease, or an acute infection in an individual in need thereof, the medicament being suitable for administration to the subject.

[0019] Accordingly, among other aspects, the present invention provides a new plasma fractionation process which separates plasma or cryo-poor plasma in an initial step into a Fraction I-IV-1 precipitate and a Fraction I-IV-1 supernatant. The Fraction I-IV-1 precipitate contains nearly all immunoglobulins (*e.g.*, IgG, IgA, and IgM) and alpha 1 elastase inhibitor (A1PI), while the supernatant contains mainly albumin.

[0020] In one embodiment, the present disclosure provides a method for preparing an enriched immunoglobulin composition from a Cohn pool, the method comprising the steps of: (a) co-precipitating immunoglobulins and alpha-1-antitrypsin (A1PI) from cryo-poor plasma, in a first precipitation step, to form a first precipitate and a first supernatant; (b) solubilizing immunoglobulins in the first precipitate, to form a first suspension having a soluble portion comprising immunoglobulins and an insoluble portion comprising A1PI; (c) separating the soluble portion of the first suspension from the insoluble portion of the first suspension; and (d) recovering the soluble fraction of the first suspension, thereby forming an enriched immunoglobulin composition.

[0021] In one embodiment of the methods described above, the first precipitation step is an alcohol precipitation step.

[0022] In one embodiment of the methods described above, the alcohol precipitation step comprises adding ethanol to the cryo-poor plasma to achieve a final concentration of from 20% to 30% ethanol (v/v) at a pH of from 5 to 6.

[0023] In one embodiment of the methods described above, the final concentration of ethanol in the first precipitation step is $25\pm 4\%$. In one embodiment of the methods described above, the final concentration of ethanol is $25\pm 3\%$. In one embodiment of the methods described above, the final concentration of ethanol is $25\pm 2\%$. In one embodiment of the methods described above, the final concentration of ethanol is $25\pm 1\%$. In one embodiment of the methods described above, the final concentration of ethanol is 25%.

[0024] In one embodiment of the methods described above, the pH of the first precipitation step is 5.5 ± 0.4 . In one embodiment of the methods described above, the pH is 5.5 ± 0.3 . In one embodiment of the methods described above, the pH is 5.5 ± 0.2 . In one embodiment of the methods described above, the pH is 5.5 ± 0.1 . In one embodiment of the methods described above, the pH is 5.5.

[0025] In one embodiment of the methods described above, the pH is maintained for the duration of the first precipitation step.

[0026] In one embodiment of the methods described above, the first alcohol precipitation step comprises addition of the alcohol by spraying.

[0027] In one embodiment of the methods described above, the first alcohol precipitation step comprises addition of the alcohol at a site adjacent to an impeller. In another embodiment, alcohol is introduced into the solution at more than one site. In one embodiment, alcohol is introduced into the solution at a plurality of small ports. In a specific embodiment, the multiple sites of alcohol addition are located at or near an impeller or other dispersive element. In another embodiment, alcohol is introduced into the solution through a diffuser port comprising a plurality of openings. In a specific embodiment, one or more of the respective openings in the plurality of openings in the diffuser port is located at or proximal to an impeller or other dispersive element.

[0028] In one embodiment of the methods described above, the first precipitation step is performed at a temperature of from -3°C to -10°C . In one embodiment of the methods described above, the first precipitation step is performed at a temperature of from -5°C to -9°C .

[0029] In one embodiment of the methods described above, the first precipitate is suspended with from 4 L to 60 L of buffer per kg precipitate. In one embodiment of the methods described above, the first precipitate is suspended with from 8 L to 15 L of buffer per kg precipitate.

[0030] In one embodiment of the methods described above, the first suspension has a pH of from 4.0 to 5.4. In one embodiment of the methods described above, the first suspension has a pH of from 4.7 to 5.1.

[0031] In one embodiment of the methods described above, the first suspension has a conductivity of from 0 mS/cm to 4 mS/cm. In one embodiment of the methods described above, the first suspension has a conductivity of from 0.5 mS/cm to 2 mS/cm.

[0032] In one embodiment of the methods described above, the first precipitate is suspended in a buffer comprising acetate and/or phosphate.

[0033] In one embodiment of the methods described above, the soluble portion of the first suspension is separated from the insoluble portion of the first suspension by centrifugation or filtration.

[0034] In one embodiment of the methods described above, the step of separating the soluble portion of the first suspension from the insoluble portion of the first suspension comprises: (i) mixing finely divided silicon dioxide (SiO_2) with the first suspension; and (ii) separating the SiO_2 from the suspension.

[0035] In one embodiment of the methods described above, the finely divided silicon dioxide (SiO_2) has an average surface area of from $350 \text{ m}^2/\text{g}$ to $410 \text{ m}^2/\text{g}$.

[0036] In one embodiment of the methods described above, the finely divided silicon dioxide (SiO_2) is added to the first suspension at a final concentration of from 15 g/kg first precipitate to 80 g/kg first precipitate.

[0037] In one embodiment of the methods described above, the method comprises the steps of: (a) precipitating immunoglobulins from cryo-poor plasma, in a first precipitation step, with between 24% and 26% alcohol at a pH of between 5.3 and 5.7 and temperature of between -6°C and -8°C to form a first precipitate and a first supernatant; (b) suspending the first precipitate to

form a first suspension; (c) treating the first suspension with finely divided silicon dioxide (SiO_2); (d) separating the soluble fraction of the first suspension from the insoluble fraction of the first suspension; and (e) recovering the soluble fraction of the first suspension, thereby forming an enriched immunoglobulin composition.

[0038] In one embodiment of the methods described above, the enriched immunoglobulin composition contains at least 90% of the immunoglobulin content of IgG present in the Cohn pool used in step (a).

[0039] In one embodiment of the methods described above, the enriched immunoglobulin composition contains at least 95% of the immunoglobulin content of IgG present in the Cohn pool used in step (a).

[0040] In one embodiment of the methods described above, the method further comprises the steps of (f) precipitating immunoglobulins from the soluble fraction of the first suspension, in a second precipitation step, to obtain a second precipitate and a second supernatant; (g) suspending the second precipitate to form a second suspension; and (h) recovering the soluble fraction of the second suspension, thereby forming a further enriched immunoglobulin composition.

[0041] In one embodiment of the methods described above, the second precipitation step is an alcohol precipitation step.

[0042] In one embodiment of the methods described above, the second alcohol precipitation step comprises adding ethanol to the soluble fraction of the first suspension to achieve a final concentration of between 22% and 28% ethanol at a pH between 6.5 and 7.5.

[0043] In one embodiment of the methods described above, the second alcohol precipitation step comprises addition of the alcohol by spraying.

[0044] In one embodiment of the methods described above, the second alcohol precipitation step comprises addition of the alcohol at a site adjacent to an impeller. In another embodiment, alcohol is introduced into the solution at more than one site. In one embodiment, alcohol is introduced into the solution at a plurality of small ports. In a specific embodiment, the multiple sites of alcohol addition are located at or near an impeller or other dispersive element. In another embodiment, alcohol is introduced into the solution through a diffuser port comprising a plurality

of openings. In a specific embodiment, one or more of the respective openings in the plurality of openings in the diffuser port is located at or proximal to an impeller or other dispersive element.

[0045] In one embodiment of the methods described above, the second precipitation step is performed at a temperature between -3°C and -10°C.

[0046] In one embodiment of the methods described above, the enriched immunoglobulin composition contains at least 90% of the IgG content present in the cryo-poor plasma used in step (a).

[0047] In one embodiment of the methods described above, the enriched immunoglobulin composition contains at least 95% of the IgG content present in the cryo-poor plasma used in step (a).

[0048] In one embodiment of the methods described above, the method further comprises an anion exchange chromatography enrichment step.

[0049] In one embodiment of the methods described above, the method further comprises a cation exchange chromatography enrichment step.

[0050] In one embodiment of the methods described above, the method further comprises a viral inactivation or removal step.

[0051] In one embodiment of the methods described above, the method comprises a solvent/detergent (S/D) viral inactivation step.

[0052] In one embodiment of the methods described above, the method comprises a nanofiltration step.

[0053] In one embodiment of the methods described above, the method comprises a step of incubating the composition at a pH from 4.0 to 5.0 and temperature from 20°C to 40°C for at least one week.

[0054] In one embodiment of the methods described above, the final enriched IgG composition comprises at least 98% IgG.

[0055] In one embodiment of the methods described above, the final enriched IgG composition comprises at least 99% IgG.

[0056] In one embodiment of the methods described above, the method yields at least 4 g of IgG per L of cryo-poor plasma used in step (a).

[0057] In one embodiment of the methods described above, the method yields at least 4.25 g of IgG per L of cryo-poor plasma used in step (a).

[0058] In one embodiment of the methods described above, the method yields at least 4.5 g of IgG per L of cryo-poor plasma used in step (a).

[0059] In one embodiment of the methods described above, alpha-1-antitrypsin (A1PI) is further purified from an insoluble fraction of the first suspension.

[0060] In one embodiment of the methods described above, fibrinogen is further purified from an insoluble fraction of the first suspension

[0061] In one embodiment of the methods described above, Factor H is further purified from an insoluble fraction of the first suspension.

[0062] In one embodiment of the methods described above, an Inter-alpha-Trypsin Inhibitor protein (I α Ip) is further purified from an insoluble fraction of the first suspension.

[0063] In one embodiment of the methods described above, the insoluble fraction of the first suspension is treated with finely divided silicon dioxide (SiO₂).

[0064] In one embodiment of the methods described above, albumin is further purified from the first supernatant.

[0065] In one embodiment, the present disclosure provides an enriched immunoglobulin composition prepared by a method according to any one of claims 1 to 63.

[0066] In one embodiment, the present disclosure provides a method for treating an immune deficiency, an inflammatory disease, an autoimmune disease, or an acute infection in an individual in need thereof, the method comprising administering an enriched immunoglobulin composition of claim 64 to the subject.

BRIEF DESCRIPTION OF DRAWINGS

[0067] **Figure 1.** Diagram of an plasma fractionation scheme according to one embodiment described herein.

DETAILED DESCRIPTION OF INVENTION

I. Introduction

[0068] Among other aspects, the present invention provides a more efficient method for the isolation of therapeutic proteins from pooled plasma. The methods provided herein for the fractionation of pooled plasma provide higher yields of multiple therapeutically important plasma-derived blood proteins, including immunoglobulins and alpha-1-antitrypsin (A1PI). In a particularly important aspect, the present invention provides methods that significantly increase the yield of immunoglobulin G (IgG) isolated from plasma, as compared to state of the art methods used for the manufacture of therapeutic IgG compositions. In one embodiment, these improved yields are achieved by precipitating immunoglobulins and A1PI from a starting plasma sample (herein referred to as a “Cohn pool”) under low pH, high alcohol conditions. The low pH, high alcohol precipitation step results in a mass capture of the immunoglobulin composition of the starting Cohn pool. As compared to state-of-the art immunoglobulin manufacturing processes, the methods provided herein significantly reduce the amount of immunoglobulin lost during the upstream fractionation of plasma.

[0069] The inventive methods reduce the number of protein precipitation steps required for the isolation of proteins from human plasma at sufficiently high purities for therapeutic administration, while simultaneously increasing the recovery yield of therapeutically important blood proteins such as immunoglobulin G (IgG). Specifically, the methods eliminate the need for an initial high pH, low alcohol precipitation step, commonly referred to a Fraction I precipitation, used in manufacturing processes derived from Cohn-Onoley, Kistler-Nitschmann, and Deutsch fractionation schemes. Instead, the methods provided herein employ an initial low pH, high alcohol precipitation step (herein referred to as Fraction I-IV-1 precipitation) that partitions the bulk of the IgG, IgA, IgM, and alpha-1-antitrypsin (A1PI) content of the plasma in the precipitate and the bulk of the albumin in the supernatant. IgG is then separated from IgA,

IgM, and A1PI through various means, resulting in high yield IgG compositions. As compared to methods that employ a Fraction I precipitation step, the mass capture of IgG in the initial Fraction I-IV-1 precipitation step increases the final yield of the manufacturing process by at least 10 to 25%.

[0070] E. J. Cohn first established a method for the fractionation of plasma using ethanol, rather than salts, in 1946. Since this time, ethanol precipitation has become the method of choice for the large-scale manufacture of plasma-derived products, such as IgG and A1PI. Typically, these manufacturing processes employ a series of ethanol fractionation steps, providing crude fractions of various plasma-derived blood proteins, which are further enriched by various different downstream procedures/techniques.

[0071] The Cohn procedures were initially developed to obtain albumin at relatively high (95%) purity by fractional precipitation with alcohol. However, it was quickly realized by Oncley *et al.*, Deutsch *et al.*, and Kistler and Nitschmann, that a particular protein precipitate (Fraction II+III) from Cohn method number 6 could be used as a starting material for the purification of highly enriched immunoglobulin compositions. In order to achieve the higher purity and safety required for the intravenous administration of IgG compositions, several purification and polishing steps (*e.g.* adsorption in general or *all* different chromatographic techniques, Cross-flow-filtration, *etc.*) have been added to IgG manufacturing processes after the alcohol fractionation steps.

[0072] Typically, IgG manufactures rely on either a Cohn *method* 6 Fraction II+III precipitate or a Kistler-Nitschmann precipitate A as the starting material for downstream processing. Both fractions are formed by a two step process in which: i.) proteins such as fibrinogen and Factor XIII are removed by an initial precipitation step (Fraction I precipitation) performed at high pH (7.2) with low ethanol concentration (8-10% v/v); and ii.) IgG is precipitated from the Fraction I supernatant at pH 6.8 with 20-25% (v/v) ethanol (Fraction II+III) or at pH 5.85 with 19% ethanol (v/v) ethanol (precipitate A), while albumin and a significant portion of A1PI remain in the supernatant. However, the use of a Fraction II+III precipitate or precipitate A as the starting material for the manufacture of IgG compositions results in the loss IgG at several steps in the process, as described above.

[0073] To overcome these issues, the inventors have developed an initial purification step that co-precipitates immunoglobulins and A1PI to provide a starting material containing nearly all of the IgG and A1PI content of the source plasma, while the albumin remains in the supernatant. This fractionation step essentially collapses fractional precipitation steps I, II+III, and IV-1 as described by Oncley *et al. (supra)* into a single precipitation reaction, referred to herein as Fraction I+II+III+IV-1 (or Fraction I-IV-1) precipitation.

[0074] In one aspect, the present invention provides a process for manufacturing a high yield immunoglobulin composition for intravenous, subcutaneous, or intramuscular administration using a Fraction I-IV-1 precipitate as a starting material. In various embodiments, the method further comprises the separation of alpha-1-antitrypsin (A1PI) into an insoluble fraction of a suspension formed during the extraction of immunoglobulin from the Fraction I-IV-1 precipitate. The separated insoluble fraction can then be used as a starting material for the manufacture of a plasma-derived A1PI composition.

[0075] In one embodiment, the present invention provides methods for manufacturing a therapeutic composition of a protein isolated from pooled plasma. In some embodiments, these manufacturing methods include an initial low pH, high alcohol (e.g., ethanol) precipitation step that increases the recovery of various blood proteins, as compared to traditional methods. In some embodiments, the methods are useful for manufacturing one or more therapeutic compositions containing an immunoglobulin (e.g., IgG), albumin, alpha-1-antitrypsin (A1PI), butyrylcholinesterase, Factor H, or an inter-alpha-trypsin inhibitor (IaI) protein isolated from human blood, plasma, or a derivative thereof.

II. Definitions

[0076] As used herein, the term "IgG treatment" refers generally to a therapeutic method of intravenously, subcutaneously, or intramuscularly administering a composition of IgG immunoglobulins to a patient for treating a number of conditions such as immune deficiencies, inflammatory diseases, and autoimmune diseases. The IgG immunoglobulins are typically pooled and prepared from plasma. Whole antibodies or fragments can be used. IgG immunoglobulins can be formulated in higher concentrations (e.g., greater than 10%) for subcutaneous administration, or formulated for intramuscular administration. This is particularly

common for specialty IgG preparations which are prepared with higher than average titres for specific antigens (e.g., Rho D factor, pertussis toxin, tetanus toxin, botulism toxin, rabies, etc.).

[0077] As used herein, “cryo-poor plasma” refers to the supernatant created after the removal of cryo-precipitate formed by thawing plasma or pooled plasma at temperatures near freezing, e.g., at temperatures below about 10°C, preferably at a temperature no higher than about 6°C. In the context of the present invention, plasma may refer interchangeably to recovered plasma (i.e., plasma that has been separated from whole blood *ex vivo*) or source plasma (i.e., plasma collected via plasmapheresis). Cryo-precipitation is commonly performed, for example, by thawing previously frozen pooled plasma, which has already been assayed for safety and quality considerations, although fresh plasma may also be used. After complete thawing of the frozen plasma at low temperature, separation of the solid cryo-precipitates from the liquid supernatant is performed in the cold (e.g., $\leq 6^{\circ}\text{C}$) by centrifugation or filtration.

[0078] As used herein, a “Cohn pool” refers to the starting material used for the fractionation of a plasma sample or pool of plasma samples. Cohn pools may include one or more of whole plasma, cryo-poor plasma, and cryo-poor plasma that has been subjected to a pre-processing step. In certain embodiments, a Cohn pool is a cryo-poor plasma sample from which one or more blood factor have been removed in a pre-processing step, for example, by adsorption onto a solid phase (e.g., aluminum hydroxide, finely divided silicon dioxide, etc.), or chromatographic step (e.g., ion exchange or heparin affinity chromatography). Various blood proteins, including but not limited to Factor Eight Inhibitor Bypass Activity (FEIBA), Factor IX-complex, Factor VII, Prothrombin complexes, and/or antithrombin III, may be isolated from the cryo-poor plasma sample prior to fractionation.

[0079] As used herein, the term “enriched composition” refers to a protein composition isolated from a plasma sample, in which the purity of the protein is higher than the purity of the protein in the starting plasma sample. In one embodiment, a protein in an enriched composition is at least 25% more pure than in the starting plasma sample. In other embodiments, an enriched composition is at least 50%, 75%, 100%, 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, or more pure than the starting plasma sample. For example, an enriched IgG composition in which 70% of the total protein is IgG is 7-fold enriched as compared to a starting plasma sample in which 10% of the total protein is IgG.

[0080] As used herein, the term "diffuse addition" refers to a means of adding a substance to a liquid system in a delocalized fashion. Diffuse addition may be achieved, for example, by spraying or misting a liquid (*e.g.*, alcohol, pH modifying agent, solvent, detergent, or other liquid) into an liquid system (*e.g.*, plasma fraction), introducing a liquid into a system at multiple sites, introducing a liquid into a system using a diffuser port, introducing a liquid at or proximal to an impeller or other dispersive element, or distributing a chemical present in a solid-state over an expanded area of a liquid system.

[0081] As used herein, the term "spraying" refers to a means of delivering a liquid substance into a system, *e.g.*, during an alcohol precipitation step, such as a Fraction I-IV-1 precipitation step, in the form of fine droplets or mist of the liquid substance. Spraying may be achieved by any pressurized device, such as a container (*e.g.*, a spray bottle), that has a spray head or a nozzle and is operated manually or automatically to generate a fine mist from a liquid. Typically, spraying is performed while the system receiving the liquid substance is continuously stirred or otherwise mixed to ensure rapid and equal distribution of the liquid within the system.

[0082] As used herein, the term "solvent" encompasses any liquid substance capable of dissolving or dispersing one or more other substances. A solvent may be inorganic in nature, such as water, or it may be an organic liquid, such as ethanol, acetone, methyl acetate, ethyl acetate, hexane, petrol ether, etc. As used in the term "solvent detergent treatment," solvent denotes an organic solvent (*e.g.*, tri-N-butyl phosphate), which is part of the solvent detergent mixture used to inactivate lipid-enveloped viruses in solution.

[0083] As used herein, the term "detergent" is used in this application interchangeably with the term "surfactant" or "surface acting agent." Surfactants are typically organic compounds that are amphiphilic, *i.e.*, containing both hydrophobic groups ("tails") and hydrophilic groups ("heads"), which render surfactants soluble in both organic solvents and water. A surfactant can be classified by the presence of formally charged groups in its head. A non-ionic surfactant has no charge groups in its head, whereas an ionic surfactant carries a net charge in its head. A zwitterionic surfactant contains a head with two oppositely charged groups. Some examples of common surfactants include: Anionic (based on sulfate, sulfonate or carboxylate anions): perfluorooctanoate (PFOA or PFO), perfluorooctanesulfonate (PFOS), sodium dodecyl sulfate (SDS), ammonium lauryl sulfate, and other alkyl sulfate salts, sodium laureth sulfate (also

known as sodium lauryl ether sulfate, or SLES), alkyl benzene sulfonate; cationic (based on quaternary ammonium cations): cetyl trimethylammonium bromide (CTAB) a.k.a. hexadecyl trimethyl ammonium bromide, and other alkyltrimethylammonium salts, cetylpyridinium chloride (CPC), polyethoxylated tallow amine (POEA), benzalkonium chloride (BAC), benzethonium chloride (BZT); Long chain fatty acids and their salts: including caprylate, caprylic acid, heptanoat, hexanoic acid, heptanoic acid, nanoic acid, decanoic acid, and the like; Zwitterionic (amphoteric): dodecyl betaine; cocamidopropyl betaine; coco ampho glycinate; nonionic: alkyl poly(ethylene oxide), alkylphenol poly(ethylene oxide), copolymers of poly(ethylene oxide) and poly(propylene oxide) (commercially known as Poloxamers or Poloxamines), alkyl polyglucosides, including octyl glucoside, decyl maltoside, fatty alcohols (e.g., cetyl alcohol and oleyl alcohol), cocamide MEA, cocamide DEA, polysorbates (Tween 20, Tween 80, etc.), Triton detergents, and dodecyl dimethylamine oxide.

[0084] By "therapeutically effective amount or dose" or "sufficient/effective amount or dose," it is meant a dose that produces effects for which it is administered. The exact dose will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques (see, e.g., Lieberman, *Pharmaceutical Dosage Forms* (vols. 1-3, 1992); Lloyd, *The Art, Science and Technology of Pharmaceutical Compounding* (1999); Pickar, *Dosage Calculations* (1999); and Remington: *The Science and Practice of Pharmacy*, 20th Edition, 2003, Gennaro, Ed., Lippincott, Williams & Wilkins).

[0085] As used herein, the terms "initial low pH, high alcohol precipitation" and "Fraction I-IV-1 precipitation" interchangeably refer to the precipitation of proteins from a Cohn pool at a final alcohol concentration (typically ethanol) of from 20% to 30% (v/v) and pH from 5.0 to 6.0. Generally, the resulting Fraction I-IV-1 precipitate contains at least 90% of the immunoglobulin content of the starting Cohn pool, preferably at least 95%, more preferably at least 98%, most preferably at least 99% of the immunoglobulin content of the starting Cohn pool. In a preferred embodiment, the immunoglobulins comprise IgG immunoglobulins. In another preferred embodiment, alpha-1-antitrypsin (A1PI) is co-precipitated with the immunoglobulins in the Fraction I-IV-1 precipitation.

III. Fractionation of Plasma

[0086] In one aspect, the present invention provides methods for fractionating therapeutic proteins from pooled plasma employing an initial precipitation step in which the majority of the immunoglobulin and alpha-1-antitrypsin (A1PI) content of the starting plasma is precipitated and the majority of the albumin content of the starting plasma is retained in the supernatant. Starting from this initial precipitation step, a number of therapeutically important blood proteins can be manufactured at high recovery yields.

[0087] In one embodiment, the present invention provides a method for fractionating blood proteins in a plasma sample, the method comprising precipitating immunoglobulins and A1PI from the starting plasma under low pH, high alcohol conditions. This low pH, high alcohol precipitation step results in the formation of a precipitate (referred to herein as a Fraction I-IV-1 precipitate) and a supernatant (referred to herein as a Fraction I-IV-1 supernatant).

[0088] The Fraction I-IV-1 supernatant will contain at least 70%, preferably at least 80%, most preferably at least 90% of the albumin content of the starting plasma. Accordingly, this supernatant can be used as a starting material for the manufacture of a pharmaceutical albumin composition.

[0089] The Fraction I-IV-1 precipitate will contain at least 90%, preferably at least 95%, more preferably at least 98%, most preferably at least 99% of the immunoglobulin content of the starting plasma. In a specific embodiment, the Fraction I-IV-1 precipitate contains at least 98%, preferably 99% of the IgG content of the starting plasma. Accordingly, this precipitate can be used as a starting material for the manufacture of pharmaceutical immunoglobulin compositions. In one embodiment, the Fraction I-IV-1 precipitate is used as a starting material for the manufacture of a pharmaceutical IgG composition. In yet another embodiment, the Fraction I-IV-1 precipitate is used as a starting material for the manufacture of a pharmaceutical immunoglobulin composition containing more than one immunoglobulin sub-type.

[0090] Likewise, the Fraction I-IV-1 precipitate will contain at least 90%, preferably at least 95%, more preferably at least 97%, most preferably at least 98% of the A1PI content of the starting plasma. Accordingly, this precipitate can be used as a starting material for the manufacture of a pharmaceutical A1PI composition.

[0091] The Fraction I-IV-1 precipitate will also contain at least 70%, preferably at least 80%, more preferably at least 90%, most preferably at least 95% of the Factor H content of the starting plasma. Accordingly, this precipitate can be used as a starting material for the manufacture of a pharmaceutical Factor H composition.

[0092] The Fraction I-IV-1 precipitate will also contain at least 70%, preferably at least 80%, more preferably at least 90%, most preferably at least 95% of the inter-alpha-inhibitor (IaIp) content of the starting plasma. Accordingly, this precipitate can be used as a starting material for the manufacture of a pharmaceutical IaIp composition.

[0093] In one aspect, the present invention provides a method for separating immunoglobulins from A1PI found in the Fraction I-IV-1 precipitate. In one embodiment, the method comprises solubilizing immunoglobulins in a suspension of the Fraction I-IV-1 precipitate, while retaining A1PI in the insoluble portion of the suspension and then separating the soluble and insoluble portions, for example by filtration or centrifugation. In one embodiment, the separation is aided by treating the Fraction I-IV-1 suspension with finely divided silicon dioxide prior to separating the soluble and insoluble portions. Without being bound by theory, the silicon dioxide may bind A1PI that is co-solubilized with the immunoglobulins, enhancing the amount of A1PI partitioned into the insoluble portion of the suspension.

[0094] Furthermore, it is known that Factor H and IaIp bind particulate silicon dioxide under certain conditions (*see*, WO/2011/011753, WO/2012/012773, and WO/2011/150284).

Accordingly, in one embodiment, the present invention provides a method for separating immunoglobulins from A1PI, Factor H, and IaIp found in a Fraction I-IV-1 precipitate, the method comprising suspending a Fraction I-IV-1 precipitate in water or a low conductivity buffer sufficient to solubilize immunoglobulins; treating the suspension with silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion, wherein soluble portion contains immunoglobulins and the insoluble portion contains A1PI, Factor H, and IaIp.

[0095] In one embodiment of the methods provided above, the separated soluble portion of the Fraction I-IV-1 suspension contains at least 80% of the IgG content of the starting plasma

sample. In a preferred embodiment, the separated soluble portion of the Fraction I-IV-1 suspension contains at least 90% of the IgG content of the starting plasma sample. In a more preferred embodiment, the separated soluble portion of the Fraction I-IV-1 suspension contains at least 95% of the IgG content of the starting plasma sample. In yet other embodiments, the soluble portion of the Fraction I-IV-1 suspension contains at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more of the IgG content of the starting plasma sample.

[0096] In one embodiment of the methods provided above, the separated insoluble portion of the Fraction I-IV-1 suspension contains at least 70% of the A1PI content of the starting plasma sample. In a preferred embodiment, the separated insoluble portion of the Fraction I-IV-1 suspension contains at least 80% of the A1PI content of the starting plasma sample. In a more preferred embodiment, the separated insoluble portion of the Fraction I-IV-1 suspension contains at least 90% of the A1PI content of the starting plasma sample. In a more preferred embodiment, the separated insoluble portion of the Fraction I-IV-1 suspension contains at least 95% of the A1PI content of the starting plasma sample. In yet other embodiments, the separated insoluble portion of the Fraction I-IV-1 suspension contains at least 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more of the A1PI content of the starting plasma sample.

[0097] Blood proteins fractionated using the initial mass precipitation step (*e.g.*, Fraction I-IV-1 precipitation) may be further enriched by suitable procedures, for example, precipitation (*e.g.*, alcohol fractionation or polyethylene glycol fractionation), chromatographic methods (ion exchange chromatography, affinity chromatography, immunoaffinity chromatography, etc.), filtration (ultrafiltration/diafiltration, nanofiltration), ultracentrifugation, electrophoretic preparation, and the like.

A. Preparation of Cryo-poor Plasma

[0098] The starting material used for the preparation of concentrated IgG compositions generally consists of either recovered plasma (*i.e.*, plasma that has been separated from whole blood ex

vivo) or source plasma (*i.e.*, plasma collected via plasmapheresis). The purification process typically starts with thawing previously frozen pooled plasma, which has already been assayed for safety and quality considerations. Thawing is typically carried out at a temperature no higher than 6°C. After complete thawing of the frozen plasma at low temperature, centrifugation is performed in the cold (*e.g.*, $\leq 6^{\circ}\text{C}$) to separate solid cryo-precipitates from the liquid supernatant. Alternatively, the separation step can be performed by filtration rather than centrifugation. The liquid supernatant (also referred to as “cryo-poor plasma,” after cold-insoluble proteins removed by centrifugation from fresh thawed plasma) is then processed in the next step. Various additional steps can be taken at this juncture for the isolation of factor eight inhibitor bypass activity (FEIBA), Factor IX-complex, Factor VII, antithrombin III, Prothrombin complexes, *etc.*

B. First Precipitation Event – Fraction I-IV-1 Precipitation

[0099] In one embodiment, the present invention provides a method for fractionating blood proteins in a plasma sample, the method comprising precipitating immunoglobulins and A1PI from the starting plasma in a first precipitation step under low pH, high alcohol conditions. This low pH, high alcohol precipitation step results in the formation of a precipitate (Fraction I-IV-1 precipitate) and a supernatant (Fraction I-IV-1 supernatant).

[00100] In one embodiment, the first precipitation step is performed by admixing ethanol with a starting plasma pool (the Cohn pool) to a final concentration of from 20% to 30% (v/v) at a pH between 5.0 and 6.0. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 2\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 1\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of 25% (v/v). Likewise, in one embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.5 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.4 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.3 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.2 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.1 . In another embodiment, the first precipitation step is performed at a pH of 5.5. In yet other embodiments, the final ethanol concentration and pH of the first precipitation step is

selected from variations 1 to 2166, as listed in **Table 1**, **Table 2**, **Table 3**, **Table 4**, **Table 5**, **Table 6**, and **Table 7**.

Table 1. Combinations of pH and final ethanol concentration useful for low pH, high alcohol precipitation of a Cohn pool.

pH	Final Ethanol Concentration [% (v/v)]							
	20-30	21-30	22-30	23-30	24-30	25-30	20-29	21-29
5.0-6.0	Var. 1	Var. 39	Var. 77	Var. 115	Var. 153	Var. 191	Var. 229	Var. 267
5.1-5.9	Var. 2	Var. 40	Var. 78	Var. 116	Var. 154	Var. 192	Var. 230	Var. 268
5.2-5.8	Var. 3	Var. 41	Var. 79	Var. 117	Var. 155	Var. 193	Var. 231	Var. 269
5.3-5.7	Var. 4	Var. 42	Var. 80	Var. 118	Var. 156	Var. 194	Var. 232	Var. 270
5.4-5.6	Var. 5	Var. 43	Var. 81	Var. 119	Var. 157	Var. 195	Var. 233	Var. 271
5.0±0.2	Var. 6	Var. 44	Var. 82	Var. 120	Var. 158	Var. 196	Var. 234	Var. 272
5.1±0.2	Var. 7	Var. 45	Var. 83	Var. 121	Var. 159	Var. 197	Var. 235	Var. 273
5.2±0.2	Var. 8	Var. 46	Var. 84	Var. 122	Var. 160	Var. 198	Var. 236	Var. 274
5.3±0.2	Var. 9	Var. 47	Var. 85	Var. 123	Var. 161	Var. 199	Var. 237	Var. 275
5.4±0.2	Var. 10	Var. 48	Var. 86	Var. 124	Var. 162	Var. 200	Var. 238	Var. 276
5.5±0.2	Var. 11	Var. 49	Var. 87	Var. 125	Var. 163	Var. 201	Var. 239	Var. 277
5.6±0.2	Var. 12	Var. 50	Var. 88	Var. 126	Var. 164	Var. 202	Var. 240	Var. 278
5.7±0.2	Var. 13	Var. 51	Var. 89	Var. 127	Var. 165	Var. 203	Var. 241	Var. 279
5.8±0.2	Var. 14	Var. 52	Var. 90	Var. 128	Var. 166	Var. 204	Var. 242	Var. 280
5.9±0.2	Var. 15	Var. 53	Var. 91	Var. 129	Var. 167	Var. 205	Var. 243	Var. 281
6.0±0.2	Var. 16	Var. 54	Var. 92	Var. 130	Var. 168	Var. 206	Var. 244	Var. 282
5.0±0.1	Var. 17	Var. 55	Var. 93	Var. 131	Var. 169	Var. 207	Var. 245	Var. 283
5.1±0.1	Var. 18	Var. 56	Var. 94	Var. 132	Var. 170	Var. 208	Var. 246	Var. 284
5.2±0.1	Var. 19	Var. 57	Var. 95	Var. 133	Var. 171	Var. 209	Var. 247	Var. 285
5.3±0.1	Var. 20	Var. 58	Var. 96	Var. 134	Var. 172	Var. 210	Var. 248	Var. 286
5.4±0.1	Var. 21	Var. 59	Var. 97	Var. 135	Var. 173	Var. 211	Var. 249	Var. 287
5.5±0.1	Var. 22	Var. 60	Var. 98	Var. 136	Var. 174	Var. 212	Var. 250	Var. 288
5.6±0.1	Var. 23	Var. 61	Var. 99	Var. 137	Var. 175	Var. 213	Var. 251	Var. 289
5.7±0.1	Var. 24	Var. 62	Var. 100	Var. 138	Var. 176	Var. 214	Var. 252	Var. 290
5.8±0.1	Var. 25	Var. 63	Var. 101	Var. 139	Var. 177	Var. 215	Var. 253	Var. 291
5.9±0.1	Var. 26	Var. 64	Var. 102	Var. 140	Var. 178	Var. 216	Var. 254	Var. 292
6.0±0.1	Var. 27	Var. 65	Var. 103	Var. 141	Var. 179	Var. 217	Var. 255	Var. 293
5	Var. 28	Var. 66	Var. 104	Var. 142	Var. 180	Var. 218	Var. 256	Var. 294
5.1	Var. 29	Var. 67	Var. 105	Var. 143	Var. 181	Var. 219	Var. 257	Var. 295
5.2	Var. 30	Var. 68	Var. 106	Var. 144	Var. 182	Var. 220	Var. 258	Var. 296
5.3	Var. 31	Var. 69	Var. 107	Var. 145	Var. 183	Var. 221	Var. 259	Var. 297
5.4	Var. 32	Var. 70	Var. 108	Var. 146	Var. 184	Var. 222	Var. 260	Var. 298
5.5	Var. 33	Var. 71	Var. 109	Var. 147	Var. 185	Var. 223	Var. 261	Var. 299
5.6	Var. 34	Var. 72	Var. 110	Var. 148	Var. 186	Var. 224	Var. 262	Var. 300
5.7	Var. 35	Var. 73	Var. 111	Var. 149	Var. 187	Var. 225	Var. 263	Var. 301
5.8	Var. 36	Var. 74	Var. 112	Var. 150	Var. 188	Var. 226	Var. 264	Var. 302
5.9	Var. 37	Var. 75	Var. 113	Var. 151	Var. 189	Var. 227	Var. 265	Var. 303

6	Var. 38	Var. 76	Var. 114	Var. 152	Var. 190	Var. 228	Var. 266	Var. 304
---	---------	---------	----------	----------	----------	----------	----------	----------

Table 2. Combinations of pH and final ethanol concentration useful for low pH, high alcohol precipitation of a Cohn pool.

pH	Final Ethanol Concentration [% (v/v)]						
	22-29	23-29	24-29	25-29	20-28	21-28	22-28
5.0-6.0	Var. 305	Var. 343	Var. 381	Var. 419	Var. 457	Var. 495	Var. 533
5.1-5.9	Var. 306	Var. 344	Var. 382	Var. 420	Var. 458	Var. 496	Var. 534
5.2-5.8	Var. 307	Var. 345	Var. 383	Var. 421	Var. 459	Var. 497	Var. 535
5.3-5.7	Var. 308	Var. 346	Var. 384	Var. 422	Var. 460	Var. 498	Var. 536
5.4-5.6	Var. 309	Var. 347	Var. 385	Var. 423	Var. 461	Var. 499	Var. 537
5.0±0.2	Var. 310	Var. 348	Var. 386	Var. 424	Var. 462	Var. 500	Var. 538
5.1±0.2	Var. 311	Var. 349	Var. 387	Var. 425	Var. 463	Var. 501	Var. 539
5.2±0.2	Var. 312	Var. 350	Var. 388	Var. 426	Var. 464	Var. 502	Var. 540
5.3±0.2	Var. 313	Var. 351	Var. 389	Var. 427	Var. 465	Var. 503	Var. 541
5.4±0.2	Var. 314	Var. 352	Var. 390	Var. 428	Var. 466	Var. 504	Var. 542
5.5±0.2	Var. 315	Var. 353	Var. 391	Var. 429	Var. 467	Var. 505	Var. 543
5.6±0.2	Var. 316	Var. 354	Var. 392	Var. 430	Var. 468	Var. 506	Var. 544
5.7±0.2	Var. 317	Var. 355	Var. 393	Var. 431	Var. 469	Var. 507	Var. 545
5.8±0.2	Var. 318	Var. 356	Var. 394	Var. 432	Var. 470	Var. 508	Var. 546
5.9±0.2	Var. 319	Var. 357	Var. 395	Var. 433	Var. 471	Var. 509	Var. 547
6.0±0.2	Var. 320	Var. 358	Var. 396	Var. 434	Var. 472	Var. 510	Var. 548
5.0±0.1	Var. 321	Var. 359	Var. 397	Var. 435	Var. 473	Var. 511	Var. 549
5.1±0.1	Var. 322	Var. 360	Var. 398	Var. 436	Var. 474	Var. 512	Var. 550
5.2±0.1	Var. 323	Var. 361	Var. 399	Var. 437	Var. 475	Var. 513	Var. 551
5.3±0.1	Var. 324	Var. 362	Var. 400	Var. 438	Var. 476	Var. 514	Var. 552
5.4±0.1	Var. 325	Var. 363	Var. 401	Var. 439	Var. 477	Var. 515	Var. 553
5.5±0.1	Var. 326	Var. 364	Var. 402	Var. 440	Var. 478	Var. 516	Var. 554
5.6±0.1	Var. 327	Var. 365	Var. 403	Var. 441	Var. 479	Var. 517	Var. 555
5.7±0.1	Var. 328	Var. 366	Var. 404	Var. 442	Var. 480	Var. 518	Var. 556
5.8±0.1	Var. 329	Var. 367	Var. 405	Var. 443	Var. 481	Var. 519	Var. 557
5.9±0.1	Var. 330	Var. 368	Var. 406	Var. 444	Var. 482	Var. 520	Var. 558
6.0±0.1	Var. 331	Var. 369	Var. 407	Var. 445	Var. 483	Var. 521	Var. 559
5	Var. 332	Var. 370	Var. 408	Var. 446	Var. 484	Var. 522	Var. 560
5.1	Var. 333	Var. 371	Var. 409	Var. 447	Var. 485	Var. 523	Var. 561
5.2	Var. 334	Var. 372	Var. 410	Var. 448	Var. 486	Var. 524	Var. 562
5.3	Var. 335	Var. 373	Var. 411	Var. 449	Var. 487	Var. 525	Var. 563
5.4	Var. 336	Var. 374	Var. 412	Var. 450	Var. 488	Var. 526	Var. 564
5.5	Var. 337	Var. 375	Var. 413	Var. 451	Var. 489	Var. 527	Var. 565
5.6	Var. 338	Var. 376	Var. 414	Var. 452	Var. 490	Var. 528	Var. 566
5.7	Var. 339	Var. 377	Var. 415	Var. 453	Var. 491	Var. 529	Var. 567
5.8	Var. 340	Var. 378	Var. 416	Var. 454	Var. 492	Var. 530	Var. 568
5.9	Var. 341	Var. 379	Var. 417	Var. 455	Var. 493	Var. 531	Var. 569
6	Var. 342	Var. 380	Var. 418	Var. 456	Var. 494	Var. 532	Var. 570

Table 3. Combinations of pH and final ethanol concentration useful for low pH, high alcohol precipitation of a Cohn pool.

pH	Final Ethanol Concentration [% (v/v)]						
	23-28	24-28	25-28	20-27	21-27	22-27	23-27
5.0-6.0	Var. 571	Var. 609	Var. 647	Var. 685	Var. 723	Var. 761	Var. 799
5.1-5.9	Var. 572	Var. 610	Var. 648	Var. 686	Var. 724	Var. 762	Var. 800
5.2-5.8	Var. 573	Var. 611	Var. 649	Var. 687	Var. 725	Var. 763	Var. 801
5.3-5.7	Var. 574	Var. 612	Var. 650	Var. 688	Var. 726	Var. 764	Var. 802
5.4-5.6	Var. 575	Var. 613	Var. 651	Var. 689	Var. 727	Var. 765	Var. 803
5.0±0.2	Var. 576	Var. 614	Var. 652	Var. 690	Var. 728	Var. 766	Var. 804
5.1±0.2	Var. 577	Var. 615	Var. 653	Var. 691	Var. 729	Var. 767	Var. 805
5.2±0.2	Var. 578	Var. 616	Var. 654	Var. 692	Var. 730	Var. 768	Var. 806
5.3±0.2	Var. 579	Var. 617	Var. 655	Var. 693	Var. 731	Var. 769	Var. 807
5.4±0.2	Var. 580	Var. 618	Var. 656	Var. 694	Var. 732	Var. 770	Var. 808
5.5±0.2	Var. 581	Var. 619	Var. 657	Var. 695	Var. 733	Var. 771	Var. 809
5.6±0.2	Var. 582	Var. 620	Var. 658	Var. 696	Var. 734	Var. 772	Var. 810
5.7±0.2	Var. 583	Var. 621	Var. 659	Var. 697	Var. 735	Var. 773	Var. 811
5.8±0.2	Var. 584	Var. 622	Var. 660	Var. 698	Var. 736	Var. 774	Var. 812
5.9±0.2	Var. 585	Var. 623	Var. 661	Var. 699	Var. 737	Var. 775	Var. 813
6.0±0.2	Var. 586	Var. 624	Var. 662	Var. 700	Var. 738	Var. 776	Var. 814
5.0±0.1	Var. 587	Var. 625	Var. 663	Var. 701	Var. 739	Var. 777	Var. 815
5.1±0.1	Var. 588	Var. 626	Var. 664	Var. 702	Var. 740	Var. 778	Var. 816
5.2±0.1	Var. 589	Var. 627	Var. 665	Var. 703	Var. 741	Var. 779	Var. 817
5.3±0.1	Var. 590	Var. 628	Var. 666	Var. 704	Var. 742	Var. 780	Var. 818
5.4±0.1	Var. 591	Var. 629	Var. 667	Var. 705	Var. 743	Var. 781	Var. 819
5.5±0.1	Var. 592	Var. 630	Var. 668	Var. 706	Var. 744	Var. 782	Var. 820
5.6±0.1	Var. 593	Var. 631	Var. 669	Var. 707	Var. 745	Var. 783	Var. 821
5.7±0.1	Var. 594	Var. 632	Var. 670	Var. 708	Var. 746	Var. 784	Var. 822
5.8±0.1	Var. 595	Var. 633	Var. 671	Var. 709	Var. 747	Var. 785	Var. 823
5.9±0.1	Var. 596	Var. 634	Var. 672	Var. 710	Var. 748	Var. 786	Var. 824
6.0±0.1	Var. 597	Var. 635	Var. 673	Var. 711	Var. 749	Var. 787	Var. 825
5	Var. 598	Var. 636	Var. 674	Var. 712	Var. 750	Var. 788	Var. 826
5.1	Var. 599	Var. 637	Var. 675	Var. 713	Var. 751	Var. 789	Var. 827
5.2	Var. 600	Var. 638	Var. 676	Var. 714	Var. 752	Var. 790	Var. 828
5.3	Var. 601	Var. 639	Var. 677	Var. 715	Var. 753	Var. 791	Var. 829
5.4	Var. 602	Var. 640	Var. 678	Var. 716	Var. 754	Var. 792	Var. 830
5.5	Var. 603	Var. 641	Var. 679	Var. 717	Var. 755	Var. 793	Var. 831
5.6	Var. 604	Var. 642	Var. 680	Var. 718	Var. 756	Var. 794	Var. 832
5.7	Var. 605	Var. 643	Var. 681	Var. 719	Var. 757	Var. 795	Var. 833
5.8	Var. 606	Var. 644	Var. 682	Var. 720	Var. 758	Var. 796	Var. 834
5.9	Var. 607	Var. 645	Var. 683	Var. 721	Var. 759	Var. 797	Var. 835
6	Var. 608	Var. 646	Var. 684	Var. 722	Var. 760	Var. 798	Var. 836

Table 4. Combinations of pH and final ethanol concentration useful for low pH, high alcohol precipitation of a Cohn pool.

pH	Final Ethanol Concentration [% (v/v)]							
	24-27	25-27	20-26	21-26	22-26	23-26	24-26	25-26
5.0-6.0	Var. 837	Var. 875	Var. 913	Var. 951	Var. 989	Var. 1027	Var. 1065	Var. 1103
5.1-5.9	Var. 838	Var. 876	Var. 914	Var. 952	Var. 990	Var. 1028	Var. 1066	Var. 1104
5.2-5.8	Var. 839	Var. 877	Var. 915	Var. 953	Var. 991	Var. 1029	Var. 1067	Var. 1105
5.3-5.7	Var. 840	Var. 878	Var. 916	Var. 954	Var. 992	Var. 1030	Var. 1068	Var. 1106
5.4-5.6	Var. 841	Var. 879	Var. 917	Var. 955	Var. 993	Var. 1031	Var. 1069	Var. 1107
5.0±0.2	Var. 842	Var. 880	Var. 918	Var. 956	Var. 994	Var. 1032	Var. 1070	Var. 1108
5.1±0.2	Var. 843	Var. 881	Var. 919	Var. 957	Var. 995	Var. 1033	Var. 1071	Var. 1109
5.2±0.2	Var. 844	Var. 882	Var. 920	Var. 958	Var. 996	Var. 1034	Var. 1072	Var. 1110
5.3±0.2	Var. 845	Var. 883	Var. 921	Var. 959	Var. 997	Var. 1035	Var. 1073	Var. 1111
5.4±0.2	Var. 846	Var. 884	Var. 922	Var. 960	Var. 998	Var. 1036	Var. 1074	Var. 1112
5.5±0.2	Var. 847	Var. 885	Var. 923	Var. 961	Var. 999	Var. 1037	Var. 1075	Var. 1113
5.6±0.2	Var. 848	Var. 886	Var. 924	Var. 962	Var. 1000	Var. 1038	Var. 1076	Var. 1114
5.7±0.2	Var. 849	Var. 887	Var. 925	Var. 963	Var. 1001	Var. 1039	Var. 1077	Var. 1115
5.8±0.2	Var. 850	Var. 888	Var. 926	Var. 964	Var. 1002	Var. 1040	Var. 1078	Var. 1116
5.9±0.2	Var. 851	Var. 889	Var. 927	Var. 965	Var. 1003	Var. 1041	Var. 1079	Var. 1117
6.0±0.2	Var. 852	Var. 890	Var. 928	Var. 966	Var. 1004	Var. 1042	Var. 1080	Var. 1118
5.0±0.1	Var. 853	Var. 891	Var. 929	Var. 967	Var. 1005	Var. 1043	Var. 1081	Var. 1119
5.1±0.1	Var. 854	Var. 892	Var. 930	Var. 968	Var. 1006	Var. 1044	Var. 1082	Var. 1120
5.2±0.1	Var. 855	Var. 893	Var. 931	Var. 969	Var. 1007	Var. 1045	Var. 1083	Var. 1121
5.3±0.1	Var. 856	Var. 894	Var. 932	Var. 970	Var. 1008	Var. 1046	Var. 1084	Var. 1122
5.4±0.1	Var. 857	Var. 895	Var. 933	Var. 971	Var. 1009	Var. 1047	Var. 1085	Var. 1123
5.5±0.1	Var. 858	Var. 896	Var. 934	Var. 972	Var. 1010	Var. 1048	Var. 1086	Var. 1124
5.6±0.1	Var. 859	Var. 897	Var. 935	Var. 973	Var. 1011	Var. 1049	Var. 1087	Var. 1125
5.7±0.1	Var. 860	Var. 898	Var. 936	Var. 974	Var. 1012	Var. 1050	Var. 1088	Var. 1126
5.8±0.1	Var. 861	Var. 899	Var. 937	Var. 975	Var. 1013	Var. 1051	Var. 1089	Var. 1127
5.9±0.1	Var. 862	Var. 900	Var. 938	Var. 976	Var. 1014	Var. 1052	Var. 1090	Var. 1128
6.0±0.1	Var. 863	Var. 901	Var. 939	Var. 977	Var. 1015	Var. 1053	Var. 1091	Var. 1129
5	Var. 864	Var. 902	Var. 940	Var. 978	Var. 1016	Var. 1054	Var. 1092	Var. 1130
5.1	Var. 865	Var. 903	Var. 941	Var. 979	Var. 1017	Var. 1055	Var. 1093	Var. 1131
5.2	Var. 866	Var. 904	Var. 942	Var. 980	Var. 1018	Var. 1056	Var. 1094	Var. 1132
5.3	Var. 867	Var. 905	Var. 943	Var. 981	Var. 1019	Var. 1057	Var. 1095	Var. 1133
5.4	Var. 868	Var. 906	Var. 944	Var. 982	Var. 1020	Var. 1058	Var. 1096	Var. 1134
5.5	Var. 869	Var. 907	Var. 945	Var. 983	Var. 1021	Var. 1059	Var. 1097	Var. 1135
5.6	Var. 870	Var. 908	Var. 946	Var. 984	Var. 1022	Var. 1060	Var. 1098	Var. 1136
5.7	Var. 871	Var. 909	Var. 947	Var. 985	Var. 1023	Var. 1061	Var. 1099	Var. 1137
5.8	Var. 872	Var. 910	Var. 948	Var. 986	Var. 1024	Var. 1062	Var. 1100	Var. 1138
5.9	Var. 873	Var. 911	Var. 949	Var. 987	Var. 1025	Var. 1063	Var. 1101	Var. 1139
6	Var. 874	Var. 912	Var. 950	Var. 988	Var. 1026	Var. 1064	Var. 1102	Var. 1140

Table 5. Combinations of pH and final ethanol concentration useful for low pH, high alcohol precipitation of a Cohn pool.

pH	Final Ethanol Concentration [% (v/v)]							
	20-25	21-25	22-25	23-25	24-25	20±1	21±1	22±1
5.0-6.0	Var. 1141	Var. 1179	Var. 1217	Var. 1255	Var. 1293	Var. 1331	Var. 1369	Var. 1407
5.1-5.9	Var. 1142	Var. 1180	Var. 1218	Var. 1256	Var. 1294	Var. 1332	Var. 1370	Var. 1408
5.2-5.8	Var. 1143	Var. 1181	Var. 1219	Var. 1257	Var. 1295	Var. 1333	Var. 1371	Var. 1409
5.3-5.7	Var. 1144	Var. 1182	Var. 1220	Var. 1258	Var. 1296	Var. 1334	Var. 1372	Var. 1410
5.4-5.6	Var. 1145	Var. 1183	Var. 1221	Var. 1259	Var. 1297	Var. 1335	Var. 1373	Var. 1411
5.0±0.2	Var. 1146	Var. 1184	Var. 1222	Var. 1260	Var. 1298	Var. 1336	Var. 1374	Var. 1412
5.1±0.2	Var. 1147	Var. 1185	Var. 1223	Var. 1261	Var. 1299	Var. 1337	Var. 1375	Var. 1413
5.2±0.2	Var. 1148	Var. 1186	Var. 1224	Var. 1262	Var. 1300	Var. 1338	Var. 1376	Var. 1414
5.3±0.2	Var. 1149	Var. 1187	Var. 1225	Var. 1263	Var. 1301	Var. 1339	Var. 1377	Var. 1415
5.4±0.2	Var. 1150	Var. 1188	Var. 1226	Var. 1264	Var. 1302	Var. 1340	Var. 1378	Var. 1416
5.5±0.2	Var. 1151	Var. 1189	Var. 1227	Var. 1265	Var. 1303	Var. 1341	Var. 1379	Var. 1417
5.6±0.2	Var. 1152	Var. 1190	Var. 1228	Var. 1266	Var. 1304	Var. 1342	Var. 1380	Var. 1418
5.7±0.2	Var. 1153	Var. 1191	Var. 1229	Var. 1267	Var. 1305	Var. 1343	Var. 1381	Var. 1419
5.8±0.2	Var. 1154	Var. 1192	Var. 1230	Var. 1268	Var. 1306	Var. 1344	Var. 1382	Var. 1420
5.9±0.2	Var. 1155	Var. 1193	Var. 1231	Var. 1269	Var. 1307	Var. 1345	Var. 1383	Var. 1421
6.0±0.2	Var. 1156	Var. 1194	Var. 1232	Var. 1270	Var. 1308	Var. 1346	Var. 1384	Var. 1422
5.0±0.1	Var. 1157	Var. 1195	Var. 1233	Var. 1271	Var. 1309	Var. 1347	Var. 1385	Var. 1423
5.1±0.1	Var. 1158	Var. 1196	Var. 1234	Var. 1272	Var. 1310	Var. 1348	Var. 1386	Var. 1424
5.2±0.1	Var. 1159	Var. 1197	Var. 1235	Var. 1273	Var. 1311	Var. 1349	Var. 1387	Var. 1425
5.3±0.1	Var. 1160	Var. 1198	Var. 1236	Var. 1274	Var. 1312	Var. 1350	Var. 1388	Var. 1426
5.4±0.1	Var. 1161	Var. 1199	Var. 1237	Var. 1275	Var. 1313	Var. 1351	Var. 1389	Var. 1427
5.5±0.1	Var. 1162	Var. 1200	Var. 1238	Var. 1276	Var. 1314	Var. 1352	Var. 1390	Var. 1428
5.6±0.1	Var. 1163	Var. 1201	Var. 1239	Var. 1277	Var. 1315	Var. 1353	Var. 1391	Var. 1429
5.7±0.1	Var. 1164	Var. 1202	Var. 1240	Var. 1278	Var. 1316	Var. 1354	Var. 1392	Var. 1430
5.8±0.1	Var. 1165	Var. 1203	Var. 1241	Var. 1279	Var. 1317	Var. 1355	Var. 1393	Var. 1431
5.9±0.1	Var. 1166	Var. 1204	Var. 1242	Var. 1280	Var. 1318	Var. 1356	Var. 1394	Var. 1432
6.0±0.1	Var. 1167	Var. 1205	Var. 1243	Var. 1281	Var. 1319	Var. 1357	Var. 1395	Var. 1433
5	Var. 1168	Var. 1206	Var. 1244	Var. 1282	Var. 1320	Var. 1358	Var. 1396	Var. 1434
5.1	Var. 1169	Var. 1207	Var. 1245	Var. 1283	Var. 1321	Var. 1359	Var. 1397	Var. 1435
5.2	Var. 1170	Var. 1208	Var. 1246	Var. 1284	Var. 1322	Var. 1360	Var. 1398	Var. 1436
5.3	Var. 1171	Var. 1209	Var. 1247	Var. 1285	Var. 1323	Var. 1361	Var. 1399	Var. 1437
5.4	Var. 1172	Var. 1210	Var. 1248	Var. 1286	Var. 1324	Var. 1362	Var. 1400	Var. 1438
5.5	Var. 1173	Var. 1211	Var. 1249	Var. 1287	Var. 1325	Var. 1363	Var. 1401	Var. 1439
5.6	Var. 1174	Var. 1212	Var. 1250	Var. 1288	Var. 1326	Var. 1364	Var. 1402	Var. 1440
5.7	Var. 1175	Var. 1213	Var. 1251	Var. 1289	Var. 1327	Var. 1365	Var. 1403	Var. 1441
5.8	Var. 1176	Var. 1214	Var. 1252	Var. 1290	Var. 1328	Var. 1366	Var. 1404	Var. 1442
5.9	Var. 1177	Var. 1215	Var. 1253	Var. 1291	Var. 1329	Var. 1367	Var. 1405	Var. 1443
6	Var. 1178	Var. 1216	Var. 1254	Var. 1292	Var. 1330	Var. 1368	Var. 1406	Var. 1444

Table 6. Combinations of pH and final ethanol concentration useful for low pH, high alcohol precipitation of a Cohn pool.

pH	Final Ethanol Concentration [% (v/v)]							
	23±1	24±1	25±1	26±1	27±1	28±1	29±1	30±1
5.0-6.0	Var. 1445	Var. 1483	Var. 1521	Var. 1559	Var. 1597	Var. 1635	Var. 1673	Var. 1711
5.1-5.9	Var. 1446	Var. 1484	Var. 1522	Var. 1560	Var. 1598	Var. 1636	Var. 1674	Var. 1712
5.2-5.8	Var. 1447	Var. 1485	Var. 1523	Var. 1561	Var. 1599	Var. 1637	Var. 1675	Var. 1713
5.3-5.7	Var. 1448	Var. 1486	Var. 1524	Var. 1562	Var. 1600	Var. 1638	Var. 1676	Var. 1714
5.4-5.6	Var. 1449	Var. 1487	Var. 1525	Var. 1563	Var. 1601	Var. 1639	Var. 1677	Var. 1715
5.0±0.2	Var. 1450	Var. 1488	Var. 1526	Var. 1564	Var. 1602	Var. 1640	Var. 1678	Var. 1716
5.1±0.2	Var. 1451	Var. 1489	Var. 1527	Var. 1565	Var. 1603	Var. 1641	Var. 1679	Var. 1717
5.2±0.2	Var. 1452	Var. 1490	Var. 1528	Var. 1566	Var. 1604	Var. 1642	Var. 1680	Var. 1718
5.3±0.2	Var. 1453	Var. 1491	Var. 1529	Var. 1567	Var. 1605	Var. 1643	Var. 1681	Var. 1719
5.4±0.2	Var. 1454	Var. 1492	Var. 1530	Var. 1568	Var. 1606	Var. 1644	Var. 1682	Var. 1720
5.5±0.2	Var. 1455	Var. 1493	Var. 1531	Var. 1569	Var. 1607	Var. 1645	Var. 1683	Var. 1721
5.6±0.2	Var. 1456	Var. 1494	Var. 1532	Var. 1570	Var. 1608	Var. 1646	Var. 1684	Var. 1722
5.7±0.2	Var. 1457	Var. 1495	Var. 1533	Var. 1571	Var. 1609	Var. 1647	Var. 1685	Var. 1723
5.8±0.2	Var. 1458	Var. 1496	Var. 1534	Var. 1572	Var. 1610	Var. 1648	Var. 1686	Var. 1724
5.9±0.2	Var. 1459	Var. 1497	Var. 1535	Var. 1573	Var. 1611	Var. 1649	Var. 1687	Var. 1725
6.0±0.2	Var. 1460	Var. 1498	Var. 1536	Var. 1574	Var. 1612	Var. 1650	Var. 1688	Var. 1726
5.0±0.1	Var. 1461	Var. 1499	Var. 1537	Var. 1575	Var. 1613	Var. 1651	Var. 1689	Var. 1727
5.1±0.1	Var. 1462	Var. 1500	Var. 1538	Var. 1576	Var. 1614	Var. 1652	Var. 1690	Var. 1728
5.2±0.1	Var. 1463	Var. 1501	Var. 1539	Var. 1577	Var. 1615	Var. 1653	Var. 1691	Var. 1729
5.3±0.1	Var. 1464	Var. 1502	Var. 1540	Var. 1578	Var. 1616	Var. 1654	Var. 1692	Var. 1730
5.4±0.1	Var. 1465	Var. 1503	Var. 1541	Var. 1579	Var. 1617	Var. 1655	Var. 1693	Var. 1731
5.5±0.1	Var. 1466	Var. 1504	Var. 1542	Var. 1580	Var. 1618	Var. 1656	Var. 1694	Var. 1732
5.6±0.1	Var. 1467	Var. 1505	Var. 1543	Var. 1581	Var. 1619	Var. 1657	Var. 1695	Var. 1733
5.7±0.1	Var. 1468	Var. 1506	Var. 1544	Var. 1582	Var. 1620	Var. 1658	Var. 1696	Var. 1734
5.8±0.1	Var. 1469	Var. 1507	Var. 1545	Var. 1583	Var. 1621	Var. 1659	Var. 1697	Var. 1735
5.9±0.1	Var. 1470	Var. 1508	Var. 1546	Var. 1584	Var. 1622	Var. 1660	Var. 1698	Var. 1736
6.0±0.1	Var. 1471	Var. 1509	Var. 1547	Var. 1585	Var. 1623	Var. 1661	Var. 1699	Var. 1737
5	Var. 1472	Var. 1510	Var. 1548	Var. 1586	Var. 1624	Var. 1662	Var. 1700	Var. 1738
5.1	Var. 1473	Var. 1511	Var. 1549	Var. 1587	Var. 1625	Var. 1663	Var. 1701	Var. 1739
5.2	Var. 1474	Var. 1512	Var. 1550	Var. 1588	Var. 1626	Var. 1664	Var. 1702	Var. 1740
5.3	Var. 1475	Var. 1513	Var. 1551	Var. 1589	Var. 1627	Var. 1665	Var. 1703	Var. 1741
5.4	Var. 1476	Var. 1514	Var. 1552	Var. 1590	Var. 1628	Var. 1666	Var. 1704	Var. 1742
5.5	Var. 1477	Var. 1515	Var. 1553	Var. 1591	Var. 1629	Var. 1667	Var. 1705	Var. 1743
5.6	Var. 1478	Var. 1516	Var. 1554	Var. 1592	Var. 1630	Var. 1668	Var. 1706	Var. 1744
5.7	Var. 1479	Var. 1517	Var. 1555	Var. 1593	Var. 1631	Var. 1669	Var. 1707	Var. 1745
5.8	Var. 1480	Var. 1518	Var. 1556	Var. 1594	Var. 1632	Var. 1670	Var. 1708	Var. 1746
5.9	Var. 1481	Var. 1519	Var. 1557	Var. 1595	Var. 1633	Var. 1671	Var. 1709	Var. 1747
6	Var. 1482	Var. 1520	Var. 1558	Var. 1596	Var. 1634	Var. 1672	Var. 1710	Var. 1748

Table 7. Combinations of pH and final ethanol concentration useful for low pH, high alcohol precipitation of a Cohn pool.

pH	Final Ethanol Concentration [% (v/v)]							
	20	21	22	23	24	25	26	27
5.0-6.0	Var. 1749	Var. 1787	Var. 1825	Var. 1863	Var. 1901	Var. 1939	Var. 1977	Var. 2015
5.1-5.9	Var. 1750	Var. 1788	Var. 1826	Var. 1864	Var. 1902	Var. 1940	Var. 1978	Var. 2016
5.2-5.8	Var. 1751	Var. 1789	Var. 1827	Var. 1865	Var. 1903	Var. 1941	Var. 1979	Var. 2017
5.3-5.7	Var. 1752	Var. 1790	Var. 1828	Var. 1866	Var. 1904	Var. 1942	Var. 1980	Var. 2018
5.4-5.6	Var. 1753	Var. 1791	Var. 1829	Var. 1867	Var. 1905	Var. 1943	Var. 1981	Var. 2019
5.0±0.2	Var. 1754	Var. 1792	Var. 1830	Var. 1868	Var. 1906	Var. 1944	Var. 1982	Var. 2020
5.1±0.2	Var. 1755	Var. 1793	Var. 1831	Var. 1869	Var. 1907	Var. 1945	Var. 1983	Var. 2021
5.2±0.2	Var. 1756	Var. 1794	Var. 1832	Var. 1870	Var. 1908	Var. 1946	Var. 1984	Var. 2022
5.3±0.2	Var. 1757	Var. 1795	Var. 1833	Var. 1871	Var. 1909	Var. 1947	Var. 1985	Var. 2023
5.4±0.2	Var. 1758	Var. 1796	Var. 1834	Var. 1872	Var. 1910	Var. 1948	Var. 1986	Var. 2024
5.5±0.2	Var. 1759	Var. 1797	Var. 1835	Var. 1873	Var. 1911	Var. 1949	Var. 1987	Var. 2025
5.6±0.2	Var. 1760	Var. 1798	Var. 1836	Var. 1874	Var. 1912	Var. 1950	Var. 1988	Var. 2026
5.7±0.2	Var. 1761	Var. 1799	Var. 1837	Var. 1875	Var. 1913	Var. 1951	Var. 1989	Var. 2027
5.8±0.2	Var. 1762	Var. 1800	Var. 1838	Var. 1876	Var. 1914	Var. 1952	Var. 1990	Var. 2028
5.9±0.2	Var. 1763	Var. 1801	Var. 1839	Var. 1877	Var. 1915	Var. 1953	Var. 1991	Var. 2029
6.0±0.2	Var. 1764	Var. 1802	Var. 1840	Var. 1878	Var. 1916	Var. 1954	Var. 1992	Var. 2030
5.0±0.1	Var. 1765	Var. 1803	Var. 1841	Var. 1879	Var. 1917	Var. 1955	Var. 1993	Var. 2031
5.1±0.1	Var. 1766	Var. 1804	Var. 1842	Var. 1880	Var. 1918	Var. 1956	Var. 1994	Var. 2032
5.2±0.1	Var. 1767	Var. 1805	Var. 1843	Var. 1881	Var. 1919	Var. 1957	Var. 1995	Var. 2033
5.3±0.1	Var. 1768	Var. 1806	Var. 1844	Var. 1882	Var. 1920	Var. 1958	Var. 1996	Var. 2034
5.4±0.1	Var. 1769	Var. 1807	Var. 1845	Var. 1883	Var. 1921	Var. 1959	Var. 1997	Var. 2035
5.5±0.1	Var. 1770	Var. 1808	Var. 1846	Var. 1884	Var. 1922	Var. 1960	Var. 1998	Var. 2036
5.6±0.1	Var. 1771	Var. 1809	Var. 1847	Var. 1885	Var. 1923	Var. 1961	Var. 1999	Var. 2037
5.7±0.1	Var. 1772	Var. 1810	Var. 1848	Var. 1886	Var. 1924	Var. 1962	Var. 2000	Var. 2038
5.8±0.1	Var. 1773	Var. 1811	Var. 1849	Var. 1887	Var. 1925	Var. 1963	Var. 2001	Var. 2039
5.9±0.1	Var. 1774	Var. 1812	Var. 1850	Var. 1888	Var. 1926	Var. 1964	Var. 2002	Var. 2040
6.0±0.1	Var. 1775	Var. 1813	Var. 1851	Var. 1889	Var. 1927	Var. 1965	Var. 2003	Var. 2041
5	Var. 1776	Var. 1814	Var. 1852	Var. 1890	Var. 1928	Var. 1966	Var. 2004	Var. 2042
5.1	Var. 1777	Var. 1815	Var. 1853	Var. 1891	Var. 1929	Var. 1967	Var. 2005	Var. 2043
5.2	Var. 1778	Var. 1816	Var. 1854	Var. 1892	Var. 1930	Var. 1968	Var. 2006	Var. 2044
5.3	Var. 1779	Var. 1817	Var. 1855	Var. 1893	Var. 1931	Var. 1969	Var. 2007	Var. 2045
5.4	Var. 1780	Var. 1818	Var. 1856	Var. 1894	Var. 1932	Var. 1970	Var. 2008	Var. 2046
5.5	Var. 1781	Var. 1819	Var. 1857	Var. 1895	Var. 1933	Var. 1971	Var. 2009	Var. 2047
5.6	Var. 1782	Var. 1820	Var. 1858	Var. 1896	Var. 1934	Var. 1972	Var. 2010	Var. 2048
5.7	Var. 1783	Var. 1821	Var. 1859	Var. 1897	Var. 1935	Var. 1973	Var. 2011	Var. 2049
5.8	Var. 1784	Var. 1822	Var. 1860	Var. 1898	Var. 1936	Var. 1974	Var. 2012	Var. 2050
5.9	Var. 1785	Var. 1823	Var. 1861	Var. 1899	Var. 1937	Var. 1975	Var. 2013	Var. 2051
6	Var. 1786	Var. 1824	Var. 1862	Var. 1900	Var. 1938	Var. 1976	Var. 2014	Var. 2052

Table 8. Combinations of pH and final ethanol concentration useful for low pH, high alcohol precipitation of a Cohn pool.

Final Ethanol Concentration [% (v/v)]			
pH	28	29	30
5.0-6.0	Var. 2053	Var. 2091	Var. 2129
5.1-5.9	Var. 2054	Var. 2092	Var. 2130
5.2-5.8	Var. 2055	Var. 2093	Var. 2131
5.3-5.7	Var. 2056	Var. 2094	Var. 2132
5.4-5.6	Var. 2057	Var. 2095	Var. 2133
5.0±0.2	Var. 2058	Var. 2096	Var. 2134
5.1±0.2	Var. 2059	Var. 2097	Var. 2135
5.2±0.2	Var. 2060	Var. 2098	Var. 2136
5.3±0.2	Var. 2061	Var. 2099	Var. 2137
5.4±0.2	Var. 2062	Var. 2100	Var. 2138
5.5±0.2	Var. 2063	Var. 2101	Var. 2139
5.6±0.2	Var. 2064	Var. 2102	Var. 2140
5.7±0.2	Var. 2065	Var. 2103	Var. 2141
5.8±0.2	Var. 2066	Var. 2104	Var. 2142
5.9±0.2	Var. 2067	Var. 2105	Var. 2143
6.0±0.2	Var. 2068	Var. 2106	Var. 2144
5.0±0.1	Var. 2069	Var. 2107	Var. 2145
5.1±0.1	Var. 2070	Var. 2108	Var. 2146
5.2±0.1	Var. 2071	Var. 2109	Var. 2147
5.3±0.1	Var. 2072	Var. 2110	Var. 2148
5.4±0.1	Var. 2073	Var. 2111	Var. 2149
5.5±0.1	Var. 2074	Var. 2112	Var. 2150
5.6±0.1	Var. 2075	Var. 2113	Var. 2151
5.7±0.1	Var. 2076	Var. 2114	Var. 2152
5.8±0.1	Var. 2077	Var. 2115	Var. 2153
5.9±0.1	Var. 2078	Var. 2116	Var. 2154
6.0±0.1	Var. 2079	Var. 2117	Var. 2155
5	Var. 2080	Var. 2118	Var. 2156
5.1	Var. 2081	Var. 2119	Var. 2157
5.2	Var. 2082	Var. 2120	Var. 2158
5.3	Var. 2083	Var. 2121	Var. 2159
5.4	Var. 2084	Var. 2122	Var. 2160
5.5	Var. 2085	Var. 2123	Var. 2161
5.6	Var. 2086	Var. 2124	Var. 2162
5.7	Var. 2087	Var. 2125	Var. 2163
5.8	Var. 2088	Var. 2126	Var. 2164
5.9	Var. 2089	Var. 2127	Var. 2165
6	Var. 2090	Var. 2128	Var. 2166

[00101] Accordingly, in one embodiment, the present invention provides a method for fractionating blood proteins in a plasma sample, the method comprising the steps of: precipitating immunoglobulins and A1PI in a first precipitation step by adding ethanol to a Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; separating the first precipitate from the first supernatant; wherein the first supernatant contains at least 75% of the albumin content of the Cohn pool. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 3\%$ (v/v).

[00102] In one embodiment of the methods provided herein, at least 90% of the immunoglobulin content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a preferred embodiment, at least 95% of the immunoglobulin content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a more preferred embodiment, at least 99% of the immunoglobulin content of the starting Cohn pool is precipitated in the initial precipitation reaction. In certain embodiments, at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more of the immunoglobulin content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a more specific embodiment, the immunoglobulin content of the starting Cohn pool refers to the IgG, IgA, and IgM content of the starting Cohn pool. In one specific embodiment, the immunoglobulin content of the starting Cohn pool refers to the IgG content of the starting Cohn pool.

[00103] In one embodiment of the methods provided herein, at least 80% of the alpha-1-antitrypsin (A1PI) content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a preferred embodiment, at least 90% of the A1PI content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a more preferred embodiment, at least 95% of the A1PI content of the starting Cohn pool is precipitated in the initial precipitation reaction. In certain embodiments, at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more of the A1PI content of the starting Cohn pool is precipitated in the initial precipitation reaction.

[00104] In one embodiment of the methods provided herein, at least 70% of the Factor H content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a preferred embodiment, at least 80% of the Factor H content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a preferred embodiment, at least 90% of the

Factor H content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a more preferred embodiment, at least 95% of the Factor H content of the starting Cohn pool is precipitated in the initial precipitation reaction. In certain embodiments, at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more of the Factor H content of the starting Cohn pool is precipitated in the initial precipitation reaction.

[00105] In one embodiment of the methods provided herein, at least 70% of the inter-alpha-inhibitor (IaIp) content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a preferred embodiment, at least 80% of the IaIp content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a preferred embodiment, at least 90% of the IaIp content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a more preferred embodiment, at least 95% of the IaIp content of the starting Cohn pool is precipitated in the initial precipitation reaction. In certain embodiments, at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more of the IaIp content of the starting Cohn pool is precipitated in the initial precipitation reaction.

[00106] Accordingly, in one embodiment, the present invention provides a method for fractionating blood proteins in a Cohn pool, the method comprising the steps of: precipitating immunoglobulins, A1PI, Factor H, and IaIp in a first precipitation step by adding ethanol to a Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; separating the first precipitate from the first supernatant; wherein the first precipitate contains i.) at least 95%, preferably at least 97%, more preferably at least 99% of the IgG content of the starting Cohn pool, ii.) at least 90%, preferably at least 95%, most preferably at least 97% of the A1PI content of the starting Cohn pool, iii.) at least 80%, preferably at least 90%, more preferably at least 95% of the Factor H content of the starting Cohn pool, and iv.) at least 80%, preferably at least 90%, more preferably at least 95% of the IaIp content of the starting Cohn pool, and further wherein the first supernatant contains at least 70%, preferably at least 80%, more preferably at least 90% of the albumin content of the Cohn pool. In one embodiment, the first precipitation step is performed by admixing ethanol with a starting plasma pool (the Cohn pool) to a final concentration of from 22% to 28% (v/v) at a pH

between 5.0 and 6.0. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 2\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 1\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of 25% (v/v). Likewise, in one embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.5 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.4 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.3 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.2 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.1 . In another embodiment, the first precipitation step is performed at a pH of 5.5. In yet other embodiments, the final ethanol concentration and pH of the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7.

IV. Preparation of Immunoglobulin Compositions

[00107] Generally, immunoglobulin preparations according to the present invention can be prepared from any suitable starting plasma material, for example, recovered plasma or source plasma. In a typical example, blood or plasma is collected from healthy donors. Usually, the blood is collected from the same species of animal as the subject to which the immunoglobulin preparation will be administered (typically referred to as “homologous” immunoglobulins). The recovered or source plasma may or may not be cryo-precipitated to provide a cryo-poor plasma. Furthermore, the plasma or cryo-poor plasma may also be treated to remove one or more blood factor by adsorption, ion exchange chromatography, or other chromatographic method. Immunoglobulins are then precipitated from the plasma or cryo-poor plasma starting material, referred to as a “Cohn pool,” under low pH, high alcohol conditions to form a Fraction I-IV-1 precipitate.

[00108] The immunoglobulins may be further enriched from the Fraction I-IV-1 precipitate by suitable procedures, for example, precipitation (alcohol fractionation or polyethylene glycol fractionation), chromatographic methods (*e.g.*, ion exchange chromatography, affinity chromatography, immunoaffinity chromatography, etc.), filtration (Ultrafiltration/diafiltration,

nanofiltration), ultracentrifugation, electrophoretic preparation, and the like. (*See, e.g., Cohn et al., J. Am. Chem. Soc.* 68:459-75 (1946); *Oncley et al., J. Am. Chem. Soc.* 71:541-50 (1949); *Barundern et al., Vox Sang.* 7:157-74 (1962); *Koblet et al., Vox Sang.* 13:93-102 (1967); U.S. Patent Nos. 5,122,373 and 5,177,194; WO/2011/149472; and WO/2011/150284).

A. Upstream Processing

[00109] Among other aspects, the present invention provides methods for fractionating proteins present in pooled plasma by performing an initial precipitation step that precipitates a majority of the immunoglobulin and alpha-1-antitrypsin content of the starting plasma. To further increase the purity of individual plasma proteins (*e.g., IgG, A1PI, Factor H, IaIp, etc.*) found in the precipitate and supernatant of the initial precipitation step, these compositions can be further enriched by fractionation (*e.g., by alcohol precipitation, PEG precipitation, salting out, etc.*), chromatography, filtration, or other method. Although use of these various techniques for the enrichment of a plasma-derived protein may be performed in any order, in a particular embodiment, the first precipitate and/or supernatant is first fractionated (*e.g., by ethanol precipitation*) and then enriched using chromatographic and/or filtration techniques. In the context of the present invention, the initial precipitation reaction and subsequent fractionation are aggregately referred to as “upstream processing” steps, while the chromatographic and filtration steps are aggregately referred to as “downstream processing” steps.

1. Mass Immunoglobulin Precipitation

[00110] As described herein, the inventors have discovered an improved method for fractionating human plasma in order to purify therapeutically beneficial blood proteins such as immunoglobulins, alpha-1-antitrypsin (A1PI), factor H, Inter-alpha-Inhibitor proteins (IaIp), albumin, fibrinogen, *etc.* This method incorporates an initial purification step in which the majority of the immunoglobulin, A1PI, factor H, IaIp, and fibrinogen content of the starting Cohn pool (*i.e., blood, plasma, and/or pre-treated plasma*) is precipitated and the majority of the albumin content of the starting Cohn pool remains in the supernatant. Advantageously, the inventors have developed a method of separating immunoglobulins from A1PI, factor H, and

IaIp by extracting the immunoglobulins, but not the other proteins, from the initial precipitate. In particular, the inventors have found that treating a suspension of the initial precipitate with finely divided silicon dioxide (SiO_2) improves the retention of A1PI, factor H, and IaIp in the insoluble fraction. Without being bound by theory, the inventors believe that under certain conditions, A1PI, factor H, and IaIp, which may be initially extracted from the precipitate, bind to the finely divided silicon dioxide, which is subsequently removed with the insoluble portion of the suspension. Separation of the resulting supernatant (*i.e.*, the soluble fraction of the suspension) provides an enriched solution containing the majority of the immunoglobulin content of the starting Cohn Fraction.

[00111] In one aspect, the present invention provides a method for preparing an enriched immunoglobulin composition from a Cohn pool, the method comprising the steps of: co-precipitating immunoglobulins and alpha-1-antitrypsin (A1PI) from a Cohn pool, in a first precipitation step, to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing the A1PI from the suspension; and recovering the soluble fraction of the first suspension, thereby forming an enriched immunoglobulin composition. Generally, any chemical means for the precipitation of immunoglobulins and A1PI may be used, including but not limited to, alcohol precipitation (*e.g.*, using ethanol or methanol), precipitation using water soluble polymers (*e.g.*, PEG or dextran), and salting out (*e.g.*, using ammonium phosphate, ammonium sulfate, sodium citrate, *etc.*). In a preferred embodiment, the precipitation is alcohol precipitation, preferably ethanol precipitation. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00112] In one embodiment, the first precipitation step is performed by admixing ethanol with the Cohn pool to a final concentration of from 20% to 30% (v/v) at a pH between 5.0 and 6.0. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 3\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 2\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 1\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of 25% (v/v). Likewise, in one embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.5 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.4 . In another embodiment, the first precipitation step is performed at a pH of

5.5±0.3. In another embodiment, the first precipitation step is performed at a pH of 5.5±0.2. In another embodiment, the first precipitation step is performed at a pH of 5.5±0.1. In another embodiment, the first precipitation step is performed at a pH of 5.5. In yet other embodiments, the final ethanol concentration and pH of the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00113] The methods provided herein provide significantly higher yields of immunoglobulin recovery in the final enriched product due to the precipitation of a majority of the immunoglobulin content of the starting Cohn pool in the initial precipitation reaction, as compared to state of the art purification procedures that rely on initial low alcohol precipitation steps (*i.e.*, Fraction I precipitation).

[00114] Accordingly, in one embodiment of the methods provided herein, at least 90% of the immunoglobulin content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a preferred embodiment, at least 95% of the immunoglobulin content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a more preferred embodiment, at least 99% of the immunoglobulin content of the starting Cohn pool is precipitated in the initial precipitation reaction. In certain embodiments, at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more of the immunoglobulin content of the starting Cohn pool is precipitated in the initial precipitation reaction. In one specific embodiment, the immunoglobulin content of the starting Cohn pool refers to the IgG content of the starting Cohn pool.

[00115] As shown in the examples provided herein, the use of an initial low pH, high alcohol precipitation step (*i.e.*, Fraction I-IV-1 precipitation) results in the precipitation of at least 99% of the IgG content of the starting Cohn pool. Accordingly, in one embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating from 99% to 100% of IgG content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; separating the first precipitate from the first supernatant; and recovering IgG from the first precipitate, thereby forming an enriched immunoglobulin composition. In one embodiment, IgG recovered from the first precipitate is further enriched. In certain embodiments, the final ethanol

concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In one embodiment, the enriched immunoglobulin composition is an IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v).

[00116] As evidenced by the data presented in Table 15 and Table 31, greater than 97% of the alpha-1-antitrypsin (A1PI) content of a Cohn pool is also precipitated by an initial low pH, high alcohol precipitation step. Accordingly, in one embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing the A1PI from the suspension; and recovering the soluble fraction of the first suspension, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the first precipitate is further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00117] Furthermore, it was found that greater than 90% of the albumin content of a Cohn pool is not precipitated by an initial low pH, high alcohol precipitation step (Table 33), and can thus be recovered from the supernatant. Accordingly, in one embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing the A1PI from the suspension; and recovering the soluble fraction of the first suspension, wherein from 80% to

100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the first precipitate is further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In yet another embodiment, albumin found in the first supernatant is further enriched. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

2. Extraction and Separation of Immunoglobulins

[00118] As compared to a Cohn Fraction II+III precipitate or Kistler-Nitschmann Precipitate A, the initial precipitate formed by the methods provided herein contains substantially higher levels of non-immunoglobulin proteins, including A1PI and fibrinogen. For example, as shown in Table 30, nearly 100% of the fibrinogen content of the starting Cohn pool is co-precipitated with immunoglobulins in an initial low pH, high alcohol precipitation reaction. This is in contrast to the Cohn-Oncley and Kistler-Nitschmann purification schemes, in which the bulk of the fibrinogen is removed in an initial low alcohol precipitation step (Fraction I precipitation). Likewise, as shown in Table 15 and Table 31, greater than 97% of the alpha-1-antitrypsin (A1PI) content of the starting Cohn pool is co-precipitated with immunoglobulins in an initial low pH, high alcohol precipitation reaction. In contrast, the bulk of A1PI is not co-precipitated with immunoglobulins in the Cohn-Oncley and Kistler-Nitschmann purification schemes. Rather, A1PI is found in the Fraction II+III supernatant or Supernatant A.

[00119] Accordingly, in order to provide pharmaceutical grade immunoglobulin compositions, contaminants such as A1PI and fibrinogen present in the initial precipitate need to be removed from the immunoglobulin composition. This may be achieved, for example, by further fractionation of the first precipitate (*e.g.*, by differential precipitation using alcohol, non-ionic hydrophilic polymers, or salting-out), chromatographic methodologies, or filtration methodologies.

[00120] In one embodiment, the first precipitate is suspended in Water for Injection (WFI) or a low ionic strength buffer suitable to extract immunoglobulins from the precipitate. In certain embodiments, the suspension is then treated with finely divided silicon dioxide (SiO_2), and the soluble portion of the suspension containing the immunoglobulins is separated from the insoluble portion of the suspension containing the bulk of the A1PI and fibrinogen.

[00121] Suitable solutions for the extraction of the first precipitate will generally have a pH between 4.0 and 5.5. In certain embodiments, the solution will have a pH between 4.5 and 5.0, in other embodiments, the extraction solution will have a pH of about 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5.0, 5.1, 5.2, 5.3, 5.4, or 5.5. In a preferred embodiment, the pH of the extraction buffer is 4.7 ± 0.1 . In another preferred embodiment, the pH of the extraction buffer is 4.8 ± 0.1 . In another preferred embodiment, the pH of the extraction buffer is 4.9 ± 0.1 . Generally, these pH requirements can be met using a buffering agent selected from, for example, acetate, citrate, monobasic phosphate, dibasic phosphate, mixtures thereof, and the like. Suitable buffer concentrations typically range from 5 to 100 mM, or from 10 to 50 mM, or 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 mM buffering agent.

[00122] The extraction buffer will preferably have a conductivity of from 0.5 mS/cm to 2.0 mS/cm. For example, in certain embodiments, the conductivity of the extraction buffer will be 0.5 ± 0.1 mS/cm, or 0.6 ± 0.1 , 0.7 ± 0.1 , 0.8 ± 0.1 , 0.9 ± 0.1 , 1.0 ± 0.1 , 1.1 ± 0.1 , 1.2 ± 0.1 , 1.3 ± 0.1 , 1.4 ± 0.1 , 1.5 ± 0.1 , 1.6 ± 0.1 , 1.7 ± 0.1 , 1.8 ± 0.1 , 1.9 ± 0.1 , or 2.0 ± 0.1 mS/cm. One of ordinary skill in the art will know how to generate extraction buffers having an appropriate conductivity. In one particular embodiment, the extraction buffer contains 5 mM monobasic sodium phosphate and 5 mM acetate at a pH of 4.8 ± 0.2 and conductivity of from 0.7 to 0.9 mS/cm.

[00123] In one embodiment, finely divided silicon dioxide is admixed with the Fraction I-IV-1 suspension prior to filtration. In one embodiment, this pretreatment step comprises addition of finely divided silica dioxide particles (*e.g.*, fumed silica; Aerosil®) followed by a 40 to 80 minute incubation period during which the suspension is constantly mixed. In certain embodiments, the incubation period will be between about 50 minutes and about 70 minutes, or about 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, or more minutes. Generally, the treatment will be performed at between 0°C and 10°C , or between 2°C and 8°C . In certain embodiments, the treatment may be performed at about 0°C , 1°C , 2°C , 3°C , 4°C , 5°C , 6°C , 7°C , 8°C , 9°C , or

10°C. In a particular embodiment, the treatment is performed at between 2°C and 10°C. In a preferred embodiment, the treatment is performed at between 5°C and 10°C.

[00124] In one embodiment, fumed silica is added at a concentration of between 20 g/kg Fraction I-IV-1 precipitate and 100 g/kg Fraction I-IV-1 precipitate. In another embodiment, fumed silica is added at a concentration of between 30 g/kg Fraction I-IV-1 precipitate and 80 g/kg Fraction I-IV-1 precipitate. In certain embodiments, the fumed silica may be added at a concentration of about 20 g/kg Fraction I-IV-1 precipitate, or about 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 g/kg Fraction I-IV-1 precipitate. In one specific embodiment, fumed silica (*e.g.*, Aerosil 380 or equivalent) is added to the Fraction I-IV-1 suspension to a final concentration of 40±20 g/kg Fraction I-IV-1 precipitate. In another specific embodiment, fumed silica is added to the Fraction I-IV-1 suspension to a final concentration of 40±10 g/kg Fraction I-IV-1 precipitate. Mixing takes place at 2°C to 8°C for at least 50 to 70 minutes.

[00125] In certain embodiments, SiO₂ is added to an immunoglobulin composition at a concentration of from 0.01 g/g total protein to 10 g/g total protein. In another embodiment, SiO₂ is added to an immunoglobulin composition at a concentration of from 0.01 g/g total protein to 5 g/g total protein. In another embodiment, SiO₂ is added to an immunoglobulin composition at a concentration of from 0.02 g/g total protein to 4 g/g total protein. In one embodiment, SiO₂ is added at a final concentration of at least 0.1 g per gram total protein. In another specific embodiment, fumed silica is added at a concentration of at least 0.2 g per gram total protein. In another specific embodiment, fumed silica is added at a concentration of at least 0.25 g per gram total protein. In other specific embodiments, fumed silica is added at a concentration of at least 1 g per gram total protein. In another specific embodiment, fumed silica is added at a concentration of at least 2 g per gram total protein. In another specific embodiment, fumed silica is added at a concentration of at least 2.5 g per gram total protein. In yet other specific embodiments, finely divided silicon dioxide is added at a concentration of at least 0.01 g/g total protein or at least 0.02 g, 0.03 g, 0.04 g, 0.05 g, 0.06 g, 0.07 g, 0.08 g, 0.09 g, 0.1 g, 0.2 g, 0.3 g, 0.4 g, 0.5 g, 0.6 g, 0.7 g, 0.8 g, 0.9 g, 1.0 g, 1.5 g, 2.0 g, 2.5 g, 3.0 g, 3.5 g, 4.0 g, 4.5 g, 5.0 g, 5.5 g, 6.0 g, 6.5 g, 7.0 g, 7.5 g, 8.0 g, 8.5 g, 9.0 g, 9.5 g, 10.0 g, or more per gram total protein.

[00126] Accordingly, in one embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating

from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains immunoglobulins and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the first precipitate is further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00127] Furthermore, it was found that greater than 90% of the albumin content of a Cohn pool is not precipitated by an initial low pH, high alcohol precipitation step (Table 33), and can thus be recovered from the supernatant. Accordingly, in one embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains immunoglobulins and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp, and further wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched immunoglobulin composition.. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG present in the soluble portion of the suspension is further enriched. In

another embodiment, A1PI present in the insoluble portion of the suspension is further enriched. In another embodiment, fibrinogen present in the insoluble portion of the suspension is further enriched. In another embodiment, Factor H present in the insoluble portion of the suspension is further enriched. In another embodiment, IaIp present in the insoluble portion of the suspension is further enriched. In yet another embodiment, albumin found in the first supernatant is further enriched. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

3. Fractionation

[00128] In one embodiment, immunoglobulins recovered from the soluble portion of the first suspension (I-IV-1 paste suspension) are further enriched by fractionation. Generally, any method of fractionation (*e.g.*, alcohol or polymer precipitation, salting out, *etc.*) may be used. In a preferred embodiment, immunoglobulins are further enriched by fractionating the soluble portion of the first suspension by ethanol precipitation. In one embodiment, immunoglobulins are enriched by the addition of ethanol to the soluble portion of the first suspension at a final concentration and pH suitable to precipitate the immunoglobulins, while at least one contaminant is not precipitated. In another embodiment, immunoglobulins are enriched by the addition ethanol to the soluble portion of the first suspension at a final concentration and pH suitable to precipitate at least one contaminant, while the immunoglobulins are not precipitated.

[00129] Additional fractionation of the material recovered from the initial low pH, high alcohol precipitation step is optional. In certain embodiments, the immunoglobulin composition recovered from the initial low pH, high alcohol precipitation step may be further enriched using one or more of the various downstream processing steps described herein. In other embodiments, the composition recovered from the initial low pH, high alcohol precipitation step may be further processed through the use of one or more additional precipitation steps. In

certain embodiments, an additional immunoglobulin precipitation step may be used to concentrate an immunoglobulin composition, prepare an immunoglobulin for storage, and/or prepare an immunoglobulin composition for transport.

[00130] In one embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition from a Cohn pool, the method comprising the steps of: co-precipitating immunoglobulins and alpha-1-antitrypsin (A1PI) from a Cohn pool, in a first precipitation step, to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing A1PI from the suspension; precipitating immunoglobulins from the first suspension in a second precipitation step, to form a second precipitate comprising immunoglobulins and a second supernatant comprising at least one contaminant; and recovering immunoglobulins from the second precipitate, thereby forming an enriched immunoglobulin composition. In one embodiment, IgG recovered from the second precipitate is further enriched. In another embodiment, A1PI removed from the first suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00131] In an exemplary embodiment, the second precipitation step is performed by admixing ethanol to the soluble portion of the second suspension (*e.g.*, a filtrate or supernatant of the suspension formed after filtration or centrifugation) at a final concentration of from 22% to 28% (v/v) at a pH between 6.5 and 7.5. In one embodiment, ethanol is admixed to a final concentration of $25 \pm 3\%$ (v/v). In another embodiment, ethanol is admixed to a final concentration of $25 \pm 2\%$ (v/v). In another embodiment, ethanol is admixed to a final concentration of $25 \pm 1\%$ (v/v). In another embodiment, ethanol is admixed to a final concentration of 25% (v/v). Likewise, in one embodiment, the second precipitation step is performed at a pH of 7.0 ± 0.5 . In another embodiment, the second precipitation step is performed at a pH of 7.0 ± 0.4 . In another embodiment, the second precipitation step is performed at a pH of 7.0 ± 0.3 . In another embodiment, the second precipitation step is performed at a pH of 7.0 ± 0.2 . In another embodiment, the second precipitation step is performed at a pH of 7.0 ± 0.1 . In another embodiment, the second precipitation step is

performed at a pH of 7.0. In a particular embodiment, the second precipitation step is performed using a final ethanol concentration of $25\pm 3\%$ (v/v) at a pH of 7.0 ± 0.3 .

[00132] In a preferred embodiment, the first suspension or soluble fraction thereof, is treated with a detergent prior to performing the second precipitation step. In one embodiment, the first suspension or soluble fraction thereof is further treated with citrate prior to performing the second precipitation step. In a particular embodiment, polysorbate-80 is added to the first suspension or soluble fraction thereof and the composition is incubated for at least 30 minutes. In another particular embodiment, Sodium citrate dihydrate is further added to the first suspension or soluble fraction thereof and the composition is incubated for at least an additional 30 minutes. In one embodiment, polysorbate-80 is added at a final concentration of about 0.2% (w/v). In one embodiment, sodium citrate dihydrate is then mixed into the solution at a final concentration of about 8 g/L. In a particular embodiment, the incubations are performed at a temperature between about 2 to 8°C with continuous of stirring.

[00133] In a particular embodiment, the method comprises the steps of: precipitating from 99% to 100% of IgG content from a Cohn pool plasma in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; separating the first precipitate from the first supernatant; suspending the first precipitate to form a first suspension; precipitating immunoglobulins from the first suspension in a second precipitation step, to form a second precipitate comprising immunoglobulins and a second supernatant comprising at least one contaminant; and recovering immunoglobulins from the second precipitate, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00134] In a specific embodiment, the method comprises the steps of: precipitating from 99% to 100% of IgG content from a Cohn pool plasma in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a

first precipitate and a first supernatant; separating the first precipitate from the first supernatant; suspending the first precipitate to form a first suspension; precipitating immunoglobulins from the first suspension in a second precipitation step by admixing ethanol to the first suspension at a final concentration of from $25\pm 3\%$ (v/v) at a pH of 7.0 ± 0.3 , to form a second precipitate comprising immunoglobulins and a second supernatant comprising at least one contaminant; and recovering immunoglobulins from the second precipitate, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00135] In one embodiment, the method comprises the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing A1PI from the suspension; recovering the soluble fraction of the first suspension; precipitating immunoglobulins from the first suspension in a second precipitation step, to form a second precipitate comprising immunoglobulins and a second supernatant comprising at least one contaminant; and recovering immunoglobulins from the second precipitate, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In one embodiment, A1PI separated from the first suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, between 99% and 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00136] In a specific embodiment, the method comprises the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to

the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing A1PI from the suspension; recovering the soluble fraction of the first suspension; precipitating immunoglobulins from the first suspension in a second precipitation step by admixing ethanol to the first suspension at a final concentration of from $25\pm 3\%$ (v/v) at a pH of 7.0 ± 0.3 , to form a second precipitate comprising immunoglobulins and a second supernatant comprising at least one contaminant; and recovering immunoglobulins from the second precipitate, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In one embodiment, A1PI separated from the first suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, between 99% and 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00137] In another embodiment, the method comprises the steps of: precipitating from 99% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing the A1PI from the suspension; recovering the soluble portion of the first suspension; precipitating immunoglobulins from the soluble portion of the first suspension in a second precipitation step, to form a second precipitate comprising immunoglobulins and a second supernatant comprising at least one contaminant; and recovering immunoglobulins from the second precipitate, wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In one embodiment, A1PI separated from the first suspension is further enriched. In another embodiment, albumin present in the first supernatant is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is

selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, between 99% and 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In a preferred embodiment, between 90% and 100% of the albumin content of the Cohn pool is present in the first supernatant. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00138] In another embodiment, the method comprises the steps of: precipitating from 99% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing the A1PI from the suspension; recovering the soluble portion of the first suspension; precipitating immunoglobulins from the soluble portion of the first suspension in a second precipitation step by admixing ethanol to the soluble portion of the first suspension at a final concentration of from $25\pm 3\%$ (v/v) at a pH of 7.0 ± 0.3 , to form a second precipitate comprising immunoglobulins and a second supernatant comprising at least one contaminant; and recovering immunoglobulins from the second precipitate, wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In one embodiment, A1PI separated from the first suspension is further enriched. In another embodiment, albumin present in the first supernatant is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, between 99% and 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In a preferred embodiment, between 90% and 100% of the albumin content of the Cohn pool is present in the first supernatant. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00139] In one embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to

the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains immunoglobulins and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp, recovering the soluble portion of the first suspension; precipitating immunoglobulins from the soluble portion of the first suspension in a second precipitation step, to form a second precipitate comprising immunoglobulins and a second supernatant comprising at least one contaminant; and recovering immunoglobulins from the second precipitate, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the first precipitate is further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00140] In a specific embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains immunoglobulins and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp, recovering the soluble portion of the first suspension; precipitating immunoglobulins from the soluble portion of the first suspension in a second precipitation step by admixing ethanol to the soluble portion of the first suspension at a final concentration of from $25\pm 3\%$ (v/v) at a pH of 7.0 ± 0.3 , to form a second precipitate

comprising immunoglobulins and a second supernatant comprising at least one contaminant; and recovering immunoglobulins from the second precipitate, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the first precipitate is further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00141] In one embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains immunoglobulins and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp; precipitating immunoglobulins from the soluble portion of the first suspension in a second precipitation step, to form a second precipitate comprising immunoglobulins and a second supernatant comprising at least one contaminant; and recovering immunoglobulins from the second precipitate, wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG present in the soluble portion of the suspension is further enriched. In another embodiment, A1PI present in the insoluble portion of the suspension is further enriched. In another embodiment, fibrinogen present in the insoluble portion of the suspension is further enriched. In another embodiment, Factor H present in the insoluble portion of the suspension is further enriched. In another embodiment, IaIp present in the insoluble portion of the suspension is further enriched. In yet another embodiment, albumin

found in the first supernatant is further enriched. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00142] In one embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains immunoglobulins and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp; precipitating immunoglobulins from the soluble portion of the first suspension in a second precipitation step by admixing ethanol to the soluble portion of the first suspension at a final concentration of from 25±3% (v/v) at a pH of 7.0±0.3, to form a second precipitate comprising immunoglobulins and a second supernatant comprising at least one contaminant; and recovering immunoglobulins from the second precipitate, wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of 25±3% (v/v). In one embodiment, IgG present in the soluble portion of the suspension is further enriched. In another embodiment, A1PI present in the insoluble portion of the suspension is further enriched. In another embodiment, fibrinogen present in the insoluble portion of the suspension is further enriched. In another embodiment, Factor H present in the insoluble portion of the suspension is further enriched. In another embodiment, IaIp present in the insoluble portion of the suspension is further enriched. In yet another embodiment, albumin found in the first supernatant is further enriched. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In certain embodiments, the final ethanol

concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00143] In one embodiment, immunoglobulins are recovered from the second precipitate by suspending the precipitate with a cold extraction buffer. For example, the second precipitate is suspended at a ratio of 1 part precipitate to 2-15 parts of Water for Injection (WFI) or low conductivity buffer. In a preferred embodiment, the second precipitate suspended at a ratio of 1 part precipitate to 4-5 parts, preferably 3.5 parts, WFI. In one embodiment, the suspension step is performed at a temperature between 0°C and 8°C. In one embodiment, the final pH of the solution is adjusted to from 4.5 to 5.6, preferably to 5.2 ± 0.2 . In one embodiment, this pH adjustment is performed with acetic acid. In one embodiment, the conductivity of the suspension is increased to between 2.5 and 6.0 mS/cm to increase the solubility of immunoglobulins. In one embodiment, the conductivity is increased by the addition of sodium chloride.

[00144] Suitable solutions for the extraction of the second precipitate include WFI and low conductivity buffers. In one embodiment, a low conductivity buffer has a conductivity of less than about 10 mS/cm. In other embodiments, the low conductivity buffer has a conductivity of less than about 9, 8, 7, 6, 5, 4, 3, 2, or 1 mS/cm. In a preferred embodiment, the low conductivity buffer has a conductivity of less than about 6 mS/cm. In another preferred embodiment, the low conductivity buffer has a conductivity of less than about 4 mS/cm. In another preferred embodiment, the low conductivity buffer has a conductivity of less than about 2 mS/cm.

[00145] In one embodiment, the soluble portion of the suspension, containing immunoglobulins, is separated from the insoluble portion. In one embodiment, this is done by filtering the suspension with a depth filter having a nominal pore size of from 0.1 μm and 0.4 μm . In one embodiment, the nominal pore size of the depth filter is 0.2 μm (*e.g.*, Cuno VR06 filter or equivalent). In one embodiment, the filter is washed with WFI or a suitable buffer after filtration to recover additional immunoglobulin and the post-wash added to the filtrate. In a preferred embodiment, the post-wash of the filter is performed using a sodium chloride solution with a

conductivity of between about 2.5 and about 6.0 mS/cm. In another embodiment, the second suspension is centrifuged to recover the soluble portion.

B. Downstream Processing

[00146] Immunoglobulin fractions obtained after fractionation using an initial precipitation step that precipitates immunoglobulins and alpha-1-antitrypsin (A1PI), but not albumin, can be further enriched according to well known methods in the art, including without limitation: chromatography (*e.g.*, anion exchange chromatography, cation exchange chromatography, hydrophobic interaction chromatography (HIC), hydroxyapatite chromatography (HAP), Protein A affinity chromatography, immuno-affinity chromatography, size exclusion chromatography, *etc.*); filtration (*e.g.*, ultrafiltration and/or diafiltration); and one or more viral reduction steps (*e.g.*, nanofiltration, solvent and detergent treatment, UV irradiation, heat treatment, low pH incubation, *etc.*).

[00147] In one embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition from a Cohn pool comprising an initial low pH, high alcohol precipitation step and at least one downstream processing step (*e.g.*, chromatography). In certain embodiments, the method further comprises a second precipitation step prior to the at least one downstream processing step (*e.g.*, a PptG precipitation step). The second precipitation is optional, as a suspension of the initial low pH, high alcohol precipitate may be used directly for downstream purification.

[00148] In one embodiment, immunoglobulins present in a plasma-derived composition prepared using an initial low pH, high alcohol precipitation step are further enriched by performing at least one chromatographic step. In one embodiment, the chromatographic step is selected from anion exchange chromatography, cation exchange chromatography, hydrophobic interaction chromatography (HIC), hydroxyapatite chromatography (HAP), Protein A affinity chromatography, immuno-affinity chromatography, and size exclusion chromatography. In a particular embodiment, immunoglobulins are enriched by performing anion exchange chromatography. In another embodiment, immunoglobulins are further enriched by performing cation exchange chromatography. In a specific embodiment, immunoglobulins are further enriched by performing both cation and anion exchange chromatography.

[00149] In a particular embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains immunoglobulins and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp, binding the immunoglobulins to a cation exchange resin; eluting the immunoglobulins from the cation exchange to form a cation exchange eluate; contacting the cation exchange eluate with an anion exchange resin; and recovering immunoglobulins that do not bind to the resin, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 3\%$ (v/v). In one embodiment, IgG recovered from the anion exchange chromatographic step is further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In another embodiment, between 80% and 100%, preferably between 90% and 100%, of the albumin content of the Cohn pool is present in the first supernatant. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, the method further comprises at least one viral reduction/inactivation step. In a preferred embodiment, the method further comprises at least two viral reduction/inactivation steps. In a more preferred embodiment, the method further comprises at least three viral reduction/inactivation steps. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00150] In a particular embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to

6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains immunoglobulins and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp, recovering the soluble portion of the first suspension; precipitating immunoglobulins from the soluble portion of the first suspension in a second precipitation step by admixing ethanol to the soluble portion of the first suspension at a final concentration of from $25\pm 3\%$ (v/v) at a pH of 7.0 ± 0.3 , to form a second precipitate comprising immunoglobulins and a second supernatant comprising at least one contaminant; recovering immunoglobulins from the second precipitate; binding the immunoglobulins to a cation exchange resin; eluting the immunoglobulins from the cation exchange to form a cation exchange eluate; contacting the cation exchange eluate with an anion exchange resin; and recovering immunoglobulins that do not bind to the resin, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the anion exchange chromatographic step is further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In another embodiment, between 80% and 100%, preferably between 90% and 100%, of the albumin content of the Cohn pool is present in the first supernatant. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, the method further comprises at least one viral reduction/inactivation step. In a preferred embodiment, the method further comprises at least two viral reduction/inactivation steps. In a more preferred embodiment, the method further comprises at least three viral reduction/inactivation steps. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00151] In one particular embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating immunoglobulins and A1PI from a Cohn pool in a low pH, high alcohol precipitation step;

separating the precipitated immunoglobulins and A1PI; enriching the immunoglobulin composition by performing cation exchange chromatography; and further enriching the immunoglobulin composition by performing anion exchange chromatography. In one embodiment, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In one embodiment, the method further comprises at least one viral reduction/inactivation step. In a preferred embodiment, the method further comprises at least two viral reduction/inactivation steps. In a more preferred embodiment, the method further comprises at least three viral reduction/inactivation steps. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00152] In another particular embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating immunoglobulins and A1PI from a Cohn pool in a low pH, high alcohol precipitation step; separating the precipitated immunoglobulins and A1PI; precipitating the separated immunoglobulins in a second precipitation step; enriching the immunoglobulin composition by performing cation exchange chromatography; and further enriching the immunoglobulin composition by performing anion exchange chromatography. In one embodiment, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In one embodiment, the method further comprises at least one viral reduction/inactivation step. In a preferred embodiment, the method further comprises at least two viral reduction/inactivation steps. In a more preferred embodiment, the method further comprises at least three viral reduction/inactivation steps. In one embodiment, the enriched immunoglobulin composition is an IgG composition.

[00153] In another particular embodiment, the present invention provides a method for preparing an enriched immunoglobulin composition, the method comprising the steps of: precipitating immunoglobulins and A1PI from a Cohn pool in a low pH, high alcohol precipitation step; separating the precipitated immunoglobulins and A1PI; precipitating the separated immunoglobulins in a second precipitation step; treating the immunoglobulin composition with a solvent and detergent to inactivate viruses; enriching the immunoglobulin

composition by performing cation exchange chromatography; further enriching the immunoglobulin composition by performing anion exchange chromatography; and nanofiltering the immunoglobulin composition to remove viruses. In one embodiment, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In one embodiment, the method further comprises concentrating the immunoglobulin composition by ultrafiltration/diafiltration to a final protein concentration of from 5 g/L to 25 g/L. In a particular embodiment, the final concentration of the enriched immunoglobulin composition is $5\pm 1\%$ (w/v). In another particular embodiment, the final concentration of the enriched immunoglobulin composition is $10\pm 1\%$ (w/v). In another particular embodiment, the final concentration of the enriched immunoglobulin composition is $15\pm 1\%$ (w/v). In another particular embodiment, the final concentration of the enriched immunoglobulin composition is $20\pm 2\%$ (w/v). In another particular embodiment, the final concentration of the enriched immunoglobulin composition is $25\pm 2\%$ (w/v). In one embodiment, the enriched immunoglobulin composition is an IgG composition.

C. Exemplary Enrichment of IgG from Fraction I-IV-1 Precipitate

1. Extraction of Fraction I-IV-1 Precipitate

[00154] In order to solubilize the IgG content of the Fraction I-IV-1 precipitate, a cold extraction buffer is used to re-suspend the Fractionation II+III precipitate at a typical ratio of 1 part precipitate to 15 parts of extraction buffer. Other suitable re-suspension ratios may be used, for example from 1:8 to 1:30, or from 1:10 to 1:20, or from 1:12 to 1:18, or from 1:13 to 1:17, or from 1:14 to 1:16. In certain embodiments, the re-suspension ratio may be 1:8, 1:9, 1:10, 1:11, 1:12, 1:13, 1:14, 1:15, 1:16, 1:17, 1:18, 1:19, 1:20, 1:21, 1:22, 1:23, 1:24, 1:25, 1:26, 1:27, 1:28, 1:29, 1:30, or higher.

[00155] Suitable solutions for the extraction of the first precipitate will generally have a pH between 4.0 and 5.5. In certain embodiments, the solution will have a pH between 4.5 and 5.0, in other embodiments, the extraction solution will have a pH of about 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5.0, 5.1, 5.2, 5.3, 5.4, or 5.5. In a preferred embodiment, the pH of the

extraction buffer is 4.7 ± 0.1 . In another preferred embodiment, the pH of the extraction buffer is 4.8 ± 0.1 . In another preferred embodiment, the pH of the extraction buffer is 4.9 ± 0.1 . Generally, these pH requirements can be met using a buffering agent selected from, for example, acetate, citrate, monobasic phosphate, dibasic phosphate, mixtures thereof, and the like. Suitable buffer concentrations typically range from 5 to 100 mM, or from 10 to 50 mM, or 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 mM buffering agent.

[00156] The extraction buffer will preferably have a conductivity of from 0.5 mS/cm to 2.0 mS/cm. For example, in certain embodiments, the conductivity of the extraction buffer will be 0.5 ± 0.1 mS/cm, or about 0.6 ± 0.1 , 0.7 ± 0.1 , 0.8 ± 0.1 , 0.9 ± 0.1 , 1.0 ± 0.1 , 1.1 ± 0.1 , 1.2 ± 0.1 , 1.3 ± 0.1 , 1.4 ± 0.1 , 1.5 ± 0.1 , 1.6 ± 0.1 , 1.7 ± 0.1 , 1.8 ± 0.1 , 1.9 ± 0.1 , or 2.0 ± 0.1 mS/cm. One of ordinary skill in the art will know how to generate extraction buffers having an appropriate conductivity.

[00157] In an exemplary embodiment, the Fraction I-IV-1 precipitate is extracted with a paste to buffer ratio of 1:15 using an extraction buffer containing 5 mM monobasic sodium phosphate and 5 mM acetate, the pH of which is adjusted to 4.8 ± 0.2 with acetic acid. In one embodiment, the pH of the solution is maintained at a pH of 4.8 ± 0.3 for the duration of the extraction process. In a specific embodiment, the pH of the solution is maintained at a pH of 4.8 ± 0.2 for the duration of the extraction process.

[00158] Generally, the extraction is performed at a temperature between 0°C and 10°C , or between 2°C and 8°C . In certain embodiments, the extraction may be performed at $0 \pm 1^{\circ}\text{C}$, $1 \pm 1^{\circ}\text{C}$, $2 \pm 1^{\circ}\text{C}$, $3 \pm 1^{\circ}\text{C}$, $4 \pm 1^{\circ}\text{C}$, $5 \pm 1^{\circ}\text{C}$, $6 \pm 1^{\circ}\text{C}$, $7 \pm 1^{\circ}\text{C}$, $8 \pm 1^{\circ}\text{C}$, $9 \pm 1^{\circ}\text{C}$, or $10 \pm 1^{\circ}\text{C}$. In a particular embodiment, the extraction is performed at between 2°C and 10°C . Typically, the extraction process will proceed for between 60 and 300 minutes, or for between 120 and 240 minutes, or for between 150 and 210 minutes, while the suspension is continuously stirred. In certain embodiments, the extraction process will proceed for about 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300 minutes, or more. In a preferred embodiment, the extraction process will proceed for at least 160 minutes with continuous stirring.

2. Silicon Dioxide (SiO₂) Treatment

[00159] As compared to Cohn-Onoley and Kistler-Nitschmann purifications, the initial immunoglobulin-containing precipitates formed by the methods provided herein contain substantially higher levels of non-immunoglobulin proteins, including A1PI, fibrinogen, Factor H, and IaIp. It is known that Factor H and IaIp can be bound by and subsequently eluted off of finely divided silicon dioxide (*see*, WO 2011/011753 and WO 2012/012773). Advantageously, it was found that inclusion of a SiO₂ treatment step after extraction of the Fraction I-IV-1 precipitate and prior to filtration of the Fraction I-IV-1 suspension aids in the separation of A1PI, fibrinogen, Factor H, and IaIp into the insoluble filter cake formed during filtration of the Fraction I-IV-1 suspension.

[00160] Accordingly, in one embodiment, finely divided silicon dioxide is admixed with the Fraction I-IV-1 suspension prior to filtration. In one embodiment, this pretreatment step comprises addition of finely divided silica dioxide particles (*e.g.*, fumed silica; Aerosil®) followed by a 40 to 80 minute incubation period during which the suspension is constantly mixed. In certain embodiments, the incubation period will be between about 50 minutes and about 70 minutes, or about 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, or more minutes. Generally, the treatment will be performed at between 0°C and 10°C, or between 2°C and 8°C. In certain embodiments, the treatment may be performed at about 0°C, 1°C, 2°C, 3°C, 4°C, 5°C, 6°C, 7°C, 8°C, 9°C, or 10°C. In a particular embodiment, the treatment is performed at between 2°C and 10°C. In a preferred embodiment, the treatment is performed at between 5°C and 10°C.

[00161] In certain embodiments, fumed silica is added at a concentration of between 20 g/kg Fraction I-IV-1 precipitate and 100 g/kg Fraction I-IV-1 precipitate. In certain embodiments, the fumed silica may be added at a concentration of about 20 g/kg Fraction I-IV-1 precipitate, or about 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 g/kg Fraction I-IV-1 precipitate. In one specific embodiment, fumed silica (*e.g.*, Aerosil 380 or equivalent) is added to the Fraction I-IV-1 suspension to a final concentration of 40±10 g/kg Fraction I-IV-1 precipitate. Mixing takes place at 2°C to 8°C for at least 50 to 70 minutes.

[00162] In certain embodiments, SiO₂ is added to an IgG composition at a concentration of from 0.01 g/g total protein to 10 g/g total protein. In another embodiment, SiO₂ is added to an IgG composition at a concentration of from 0.01 g/g total protein to 5 g/g total protein. In another embodiment, SiO₂ is added to an IgG composition at a concentration of from 0.02 g/g total protein to 4 g/g total protein. In one embodiment, SiO₂ is added at a final concentration of at least 0.1 g per gram total protein. In another specific embodiment, fumed silica is added at a concentration of at least 0.2 g per gram total protein. In another specific embodiment, fumed silica is added at a concentration of at least 0.25 g per gram total protein. In other specific embodiments, fumed silica is added at a concentration of at least 1 g per gram total protein. In another specific embodiment, fumed silica is added at a concentration of at least 2 g per gram total protein. In another specific embodiment, fumed silica is added at a concentration of at least 2.5 g per gram total protein. In yet other specific embodiments, finely divided silicon dioxide is added at a concentration of at least 0.01 g/g total protein or at least 0.02 g, 0.03 g, 0.04 g, 0.05 g, 0.06 g, 0.07 g, 0.08 g, 0.09 g, 0.1 g, 0.2 g, 0.3 g, 0.4 g, 0.5 g, 0.6 g, 0.7 g, 0.8 g, 0.9 g, 1.0 g, 1.5 g, 2.0 g, 2.5 g, 3.0 g, 3.5 g, 4.0 g, 4.5 g, 5.0 g, 5.5 g, 6.0 g, 6.5 g, 7.0 g, 7.5 g, 8.0 g, 8.5 g, 9.0 g, 9.5 g, 10.0 g, or more per gram total protein.

3. Filtration of the Fraction I-IV-1 Suspension

[00163] In order to separate the soluble portion of the Fraction I-IV-1 suspension containing immunoglobulins from the insoluble portion containing fibrinogen, A1PI, Factor H, and IaIp, the suspension is filtered, typically using depth filtration. Depth filters that may be employed in the methods provided herein include, metallic, glass, ceramic, organic (such as diatomaceous earth) depth filters, and the like. Examples of suitable filters include, without limitation, Cuno 50SA, Cuno 90SA, and Cuno VR06 filters (3M). Alternatively, the separation step can be performed by centrifugation rather than filtration.

[00164] In certain embodiments, filter aid, for example Celpure C300 (Advanced Minerals) or Hyflo-Super-Cel (World Minerals), is added to the suspension after the silica dioxide treatment to facilitate depth filtration. Filter aid is added at a final concentration of from 0.1 kg/kg Fraction I-IV-1 precipitate to 1.0 kg/kg Fraction I-IV-1 precipitate, or from 0.2 kg/kg Fraction I-IV-1 precipitate to 0.8 kg/kg Fraction I-IV-1 precipitate, or from 0.3 kg/kg Fraction I-IV-1

precipitate to 0.7 kg/kg Fraction I-IV-1 precipitate. In other embodiments, filter aid can be added at a final concentration of from 0.1 kg/kg Fraction I-IV-1 precipitate to 0.7 kg/kg Fraction I-IV-1 precipitate, or from 0.2 kg/kg Fraction I-IV-1 precipitate to 0.6 kg/kg Fraction I-IV-1 precipitate, or from about 0.3 kg/kg Fraction I-IV-1 precipitate to about 0.05 kg/kg Fraction I-IV-1 precipitate. In certain embodiments, the filter aid will be added at a final concentration of about 0.01 kg/kg Fraction I-IV-1 precipitate, or about 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, or 1.0 kg/kg Fraction I-IV-1 precipitate.

[00165] In order to minimize the loss of immunoglobulins during filtration, the filter cake formed should be washed with at least one dead volume, preferably at least two dead volumes, more preferably at least three dead volumes, of suspension buffer or a similar buffer thereto, which is not sufficient to solubilize non-immunoglobulin proteins present in the filter cake. In one embodiment, the filter and filter cake are post washed with at least 3.0 dead volumes of buffer. In another embodiment, the filter and filter cake are post washed with at least 3.6 dead volumes of buffer. In another embodiment, the filter and filter cake are post washed with at least 50% of the suspension volume filtered, using a suitable buffer. In another embodiment, the filter and filter cake are post washed with at least 75% of the suspension volume filtered, using a suitable buffer. In another embodiment, the filter and filter cake are post washed with at least 100% of the suspension volume filtered, using a suitable buffer. Typically, no more than 200% of the suspension volume filtered should be used to wash the filter and filter cake.

[00166] In a specific embodiment, the wash buffer is the dissolution buffer containing from 90 to 150 mL of glacial acetic acid per 1000 L. In one embodiment, the pH of the post-wash extraction buffer is between about 4.6 and about 5.3. In a preferred embodiment, the pH of the post-wash buffer is between about 4.7 and about 5.2. In another preferred embodiment, the pH of the post-wash buffer is between about 4.8 and about 5.1. In yet another preferred embodiment, the pH of the post-wash buffer is between about 4.9 and about 5.0.

4. Detergent Treatment

[00167] To remove additional contaminants from the Fraction I-IV-1 filtrate, the sample is next subjected to a detergent treatment. Methods for the detergent treatment of plasma derived fractions are well known in the art. Generally, any standard non-ionic detergent treatment may

be used in conjunction with the methods provided herein. For example, an exemplary protocol for a detergent treatment is provided below.

[00168] Polysorbate-80 is added to the Fraction I-IV-1 filtrate at a final concentration of about 0.2% (w/v) with stirring and the sample is incubated for at least 30 minutes at a temperature between about 2 to 8°C. Sodium citrate dihydrate is then mixed into the solution at a final concentration of about 8 g/L and the sample is incubated for an additional 30 minutes, with continuous of stirring at a temperature between about 2 to 8°C.

[00169] In certain embodiments, any suitable non-ionic detergent can be used. Examples of suitable non-ionic detergents include, without limitation, Octylglucoside, Digitonin, C12E8, Lubrol, Triton X-100, Nonidet P-40, Tween-20 (i.e., polysorbate-20), Tween-80 (i.e., polysorbate-80), an alkyl poly(ethylene oxide), a Brij detergent, an alkylphenol poly(ethylene oxide), a poloxamer, octyl glucoside, decyl maltoside, and the like.

[00170] In one embodiment, a process improvement is realized by adding the detergent reagents (e.g., polysorbate-80 and sodium citrate dehydrate) by diffuse addition. The diffuse addition of these reagents minimizes local fluctuations of alcohol concentration and pH as compared to addition at a single entry point. In one embodiment, the diffuse addition comprises spraying, rather than fluent addition. In another embodiment, the diffuse addition comprises addition of a reagent from multiple ports. In one embodiment, the diffuse addition comprises addition of a reagent from a diffuser port. In certain embodiments, at least one port used to introduce a reagent into the system is located at or near an impeller or other dispersive element. In other embodiments, the detergent reagents may be added as solids to the Modified Fraction II+III filtrate while the sample is being mixed to ensure rapid distribution of the additives. In certain embodiments, it is preferable to add solid reagents by sprinkling the solids over a delocalized surface area of the filtrate such that local overconcentration does not occur, such as in fluent addition.

5. Second Precipitation Event – Precipitation G

[00171] In order to remove residual impurities, a second precipitation is performed at pH 6.5 to 7.5 using 25% alcohol. Briefly, the Fraction I-IV-1 extraction is adjusted to a pH from 6.5 to

7.5, preferably from 6.8 to 7.2, more preferably from 6.9 to 7.1, most preferably 7.0 with a suitable pH modifying solution (*e.g.*, sodium hydroxide or acetic acid). Cold alcohol is then added to the solution to a final concentration of about 25% (v/v) and the mixture is incubated while stirring at between about -6°C to about -10°C for at least 1 hour to form a second precipitate (*i.e.*, precipitate G). In one embodiment, the mixture is incubated for at least 2 hours, or at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or more hours. In a preferred embodiment, the mixture is incubated for at least 2 hours. In a more preferred embodiment, the mixture is incubated for at least 4 hours. In an even more preferred embodiment, the mixture is incubated for at least 8 hours.

[00172] In certain embodiments, a higher percentage of IgG may be precipitated from the I-IV-1 extraction by adjusting the pH of the solution during and/or after the addition of alcohol to the precipitation reaction. Likewise, due to fluctuations in the pH of the solution during the precipitation process, the pH of the reaction mixture may be monitored and/or adjusted throughout the entirety of the incubation. In another embodiment, improvement in the yield of IgG is achieved by adding the precipitating alcohol and/or the solution used to adjust the pH of the mixture by diffuse addition of the alcohol or pH modifying agent. The diffuse addition of these reagents minimizes local fluctuations of alcohol concentration and pH as compared to addition at a single entry point. In one embodiment, the diffuse addition comprises spraying, rather than fluent addition. In another embodiment, the diffuse addition comprises addition of a reagent from multiple ports. In one embodiment, the diffuse addition comprises addition of a reagent from a diffuser port. In certain embodiments, at least one port used to introduce a reagent into the system is located at or near an impeller or other dispersive element.

6. Suspension and Filtration of Precipitate G (Ppt G)

[00173] In order to solubilize the IgG content of the precipitate G, a cold extraction buffer is used to re-suspend the PptG. Briefly, the precipitate G is dissolved at a ratio of 1 part precipitate to 2-5 parts Water for Injection (WFI) or low conductivity buffer. In a preferred embodiment, the precipitate G is dissolved at a ratio of 1 part precipitate to 1-1.5 parts, preferably 3.5 parts, WFI. The suspension step is typically done at a temperature between 0°C and 8°C to achieve an AU₂₈₀₋₃₂₀ value of from 40 to 95. The final pH of the solution, which is stirred for at least 2

hours, is then adjusted to from 4.5 to 5.6, preferably to 5.2 ± 0.2 . In one embodiment, this pH adjustment is performed with acetic acid. To increase the solubility of IgG, the conductivity of the suspension is increased to between 2.5 and 6.0 mS/cm. In one embodiment, the conductivity is increased by the addition of sodium chloride.

[00174] Suitable solutions for the extraction of precipitate G include WFI and low conductivity buffers. In one embodiment, a low conductivity buffer has a conductivity of less than about 10 mS/cm. In other embodiments, the low conductivity buffer has a conductivity of less than about 9, 8, 7, 6, 5, 4, 3, 2, or 1 mS/cm. In a preferred embodiment, the low conductivity buffer has a conductivity of less than about 6 mS/cm. In another preferred embodiment, the low conductivity buffer has a conductivity of less than about 4 mS/cm. In another preferred embodiment, the low conductivity buffer has a conductivity of less than about 2 mS/cm.

[00175] The suspended PptG solution is then filtered with a suitable depth filter having a nominal pore size of between 0.1 μm and 0.4 μm in order to remove any undissolved particles. In one embodiment, the nominal pore size of the depth filter is about 0.2 μm (e.g., Cuno VR06 filter or equivalent). In another embodiment, the suspended PptG solution is centrifuged to recover a clarified supernatant. The filter is washed with WFI or a suitable buffer after filtration to recover additional IgG and the post-wash added to the filtrate. In a preferred embodiment, the post-wash of the filter is performed using a sodium chloride solution with a conductivity of between about 2.5 and about 6.0 mS/cm.

7. Solvent and Detergent (S/D) Treatment

[00176] In order to inactivate various viral contaminants present in plasma-derived products, the clarified PptG filtrate is next subjected to a solvent detergent (S/D) treatment. Methods for the detergent treatment of plasma derived fractions are well known in the art (for review *see*, Pelletier JP *et al.*, Best Pract Res Clin Haematol. 2006;19(1):205-42). Generally, any standard S/D treatment may be used in conjunction with the methods provided herein. For example, an exemplary protocol for an S/D treatment is provided below.

[00177] Triton X-100, Tween-20, and tri(n-butyl)phosphate (TNBP) are added to the clarified PptG filtrate at final concentrations of about 1.0%, 0.3%, and 0.3%, respectively. The mixture is then stirred at a temperature between 18°C and 25°C for at least an hour.

[00178] In one embodiment, the S/D reagents (e.g., Triton X-100, Tween-20, and TNBP) are added by diffuse addition. In a specific embodiment, the S/D reagents are added by spraying rather than by fluent addition. In another embodiment, the detergent reagents are added as solids to the clarified PptG filtrate, which is being mixed to ensure rapid distribution of the S/D components. In certain embodiments, it is preferable to add solid reagents by sprinkling the solids over a delocalized surface area of the filtrate such that local overconcentration does not occur, such as in fluent addition.

8. Ion Exchange Chromatography

[00179] In order to further purify and concentrate IgG, cation exchange and/or anion exchange chromatography can be employed. Methods for purifying and concentrating IgG using ion exchange chromatography are well known in the art. For example, U.S. Patent No. 5,886,154 describes a method in which a Fraction II+III precipitate is extracted at low pH (between about 3.8 and 4.5), followed by precipitation of IgG using caprylic acid, and finally implementation of two anion exchange chromatography steps. U.S. Patent No. 6,069,236 describes a chromatographic IgG purification scheme that does not rely on alcohol precipitation at all. PCT Publication No. WO 2005/073252 describes an IgG purification method involving the extraction of a Fraction II+III precipitate, caprylic acid treatment, PEG treatment, and a single anion exchange chromatography step. U.S. Patent No. 7,186,410 describes an IgG purification method involving the extraction of a Fraction I+II+III or Fraction II precipitate followed by a single anion exchange step performed at an alkaline pH. U.S. Patent No. 7,553,938 describes a method involving the extraction of a Fraction I+II+III or Fraction II+III precipitate, caprylate treatment, and either one or two anion exchange chromatography steps. U.S. Patent No. 6,093,324 describes a purification method comprising the use of a macroporous anion exchange resin operated at a pH between about 6.0 and about 6.6. U.S. Patent No. 6,835,379 describes a purification method that relies on cation exchange chromatography in the absence of alcohol

fractionation.

[00180] In one embodiment, the S/D treated PptG filtrate may be subjected to both cation exchange chromatography and anion exchange chromatography. For example, in one embodiment, the S/D treated PptG filtrate is contacted by a cation exchange resin under suitable conditions to binds the IgG to the resin. The S/D reagents can then be washed away from the adsorbed IgG, which is subsequently eluted off of the resin with a suitable elution buffer. In this fashion, the cation exchange chromatography step can be used to remove the S/D reagents from the preparation, concentrate the IgG containing solution, and/or remove impurities from the composition.

[00181] In one embodiment, IgG is eluted from the cation exchange resin using an elution buffer having a pH between about 8.0 and 9.0. In certain embodiments, the pH elution buffer may have a pH between about 8.2 and about 8.8, or between about 8.4 and about 8.6, or a pH of about 8.0, 8.1, 8.2, 8.3, 8.4, 8.5, 8.6, 8.7, 8.8, 8.9, or 9.0. In a preferred embodiment, the pH of the elution buffer is about 8.5 ± 0.1 .

[00182] Likewise, anion exchange chromatography can be used to reduce impurities in the IgG composition. In one embodiment, anion exchange chromatography is performed under solution conditions in which one or more impurities bind to the anion exchange resin but IgG does not. In a specific embodiment, anion exchange chromatography is performed at slightly acidic conditions, *i.e.*, at a pH between 5.0 and 7.0, preferably between 5.5 and 6.5, at a low ionic strength suitable for binding of contaminants but not IgG to the anion exchange resin.

[00183] In a specific embodiment, the eluate from the cation exchange column may be adjusted to a lower pH, for example between about 5.5 and about 6.5, and diluted with an appropriate buffer such that the conductivity of the solution is reduced. In certain embodiments, the pH of the cation exchange eluate may be adjusted to a pH between about 5.7 and about 6.7, or between about 5.9 and about 6.5, or a pH of about 5.5, 5.6, 5.7, 5.8, 5.9, 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, or 6.7. In a preferred embodiment, the pH of the eluate is adjusted to a pH of about 6.4 ± 0.1 . The eluate is then loaded onto an anion exchange column, which binds several contaminants found in the preparation. The column flow through, containing the IgG fraction, is collected during

column loading and washing. In certain embodiments, the ion exchange chromatographic steps of the present invention can be performed in column mode, batch mode, or in a combination of the two. In one embodiment, the solution used to adjust the pH of the IgG composition prior to anion exchange chromatography is added by diffuse addition. In a specific embodiment, the solution is added by spraying, rather than by fluent addition.

9. Nanofiltration

[00184] In order to reduce the viral load of an IgG composition provided herein, the composition may be nanofiltered using a suitable nanofiltration device. In certain embodiments, the nanofiltration device will have a mean pore size of between about 15 nm and about 200 nm. Examples of nanofilters suitable for this use include, without limitation, DVD, DV 50, DV 20 (Pall), Viresolve NFP, Viresolve NFR (Millipore), Planova 15N, 20N, 35N, and 75N (Planova). In a specific embodiment, the nanofilter may have a mean pore size of between about 15 nm and about 72 nm, or between about 19 nm and about 35 nm, or of about 15 nm, 19 nm, 35 nm, or 72 nm. In a preferred embodiment, the nanofilter will have a mean pore size of about 35 nm, such as an Asahi PLANOVA 35N filter or equivalent thereof. In a particular embodiment, the IgG composition recovered from the anion exchange step is nanofiltered using a nanofilter having a pore size between 30 nm and 40 nm, preferably 35 ± 2 nm. In another preferred embodiment, the nanofilter will have a mean pore size of about 19 or 20 nm, such as an Asahi PLANOVA 20N filter (19 ± 2 nm) or equivalent thereof. In a particular embodiment, the IgG composition recovered from the anion exchange step is nanofiltered using a nanofilter having a pore size between 15 nm and 25 nm, preferably 19 ± 2 nm.

10. Ultra-/Diafiltration (UF/DF)

[00185] Ultrafiltration/diafiltration may be performed to concentrate the IgG composition at any step during the purification process. In one embodiment, the nanofiltrate is concentrated by UF/DF. In one embodiment, the nanofiltrate may be concentrated by ultrafiltration to a protein concentration of between about 2% and about 10% (w/v). In certain embodiments, the ultrafiltration is carried out in a cassette with an open channel screen and the ultrafiltration membrane has a nominal molecular weight cut off (NMWCO) of less than about 100 kDa or less

than about 90, 80, 70, 60, 50, 40, 30, or fewer kDa. In a preferred embodiment, the ultrafiltration membrane has a NMWCO of no more than 50 kDa.

[00186] In one embodiment, an open channel membrane is used with a specifically designed post-wash and formulation near the end the production process to render the resulting IgG compositions about twice as high in protein concentration (200 mg/mL) compared to state of the art IVIGs (*e.g.*, GAMMAGARD[®] LIQUID, 10% IgG) without affecting yield and storage stability. With most of the commercial available ultrafiltration membranes a concentration of 200 mg/mL IgG cannot be reached without major protein losses. These membranes are blocked early and therefore adequate post-wash is difficult to achieve. Therefore open channel membrane configurations have to be used. Even with open channel membranes, a specifically designed post-wash procedure has to be used to obtain the required concentration without significant protein loss (less than 2% loss). Even more surprising is the fact that the higher protein concentration of 200 mg/mL does not reduce the virus inactivation capacity of a low pH storage step.

[00187] Upon completion of the ultrafiltration step, the concentrate may further be concentrated via diafiltration against a solution suitable for intravenous or intramuscular administration. In certain embodiments, the diafiltration solution may comprise a stabilizing and/or buffering agent. In a preferred embodiment, the stabilizing and buffering agent is glycine at an appropriate concentration, for example between about 0.20 M and about 0.30 M, or between about 0.22 M and about 0.28 M, or between about 0.24 M and about 0.26 M, or at a concentration of 0.20±0.01 M, 0.21±0.01 M, 0.22±0.01 M, 0.23±0.01 M, 0.24±0.01 M, 0.25±0.01 M, 0.26±0.01 M, 0.27±0.01 M, 0.28±0.01 M, 0.29±0.01 M, or 0.3±0.01 M. In a preferred embodiment, the diafiltration buffer contains 0.25±0.01 M glycine.

[00188] Typically, the minimum exchange volume for diafiltration is at least about 3 times the original concentrate volume or at least about 4, 5, 6, 7, 8, 9, or more times the original concentrate volume. The IgG solution may be concentrated to a final protein concentration of between about 5% and about 25% (w/v), or between about 6% and about 18% (w/v), or between about 7% and about 16% (w/v), or between about 8% and about 14% (w/v), or between about 9% and about 12%, or to a final concentration of about 5%, or 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25% or higher. In

one embodiment, a final protein concentration of at least about 23% is achieved without adding the post-wash fraction to the concentrated solution. In another embodiment, a final protein concentration of at least about 24% is achieved without adding the post-wash fraction to the concentrated solution. In another embodiment, a final protein concentration of at least about 24% is achieved without adding the post-wash fraction to the concentrated solution. In one embodiment, the pH of the solution will be from 4.6 to 5.1 after diafiltration.

[00189] In an exemplary embodiment, the pH of the IgG composition is adjusted to about 4.5 prior to ultrafiltration. The solution is concentrated to a protein concentration of $5 \pm 2\%$ w/v through ultrafiltration. The UF membrane has a nominal molecular weight cut off (NMWCO) of 50,000 Daltons or less (*e.g.*, Millipore Pellicon Polyether sulfon membrane). The concentrate is then diafiltered against ten volumes of 0.25 M glycine solution, pH 4.5 ± 0.2 . Throughout the ultra-diafiltration operation the solution is maintained at a temperature of between about 2°C to about 8°C. After diafiltration, the solution is concentrated to a protein concentration of at least 11 % (w/v).

11. Formulation

[00190] Upon completion of the diafiltration step, the protein concentration of the solution is adjusted to with the diafiltration buffer to a final concentration of between about 5% and about 20% (w/v), or between about 6% and about 18% (w/v), or between about 7% and about 16% (w/v), or between about 8% and about 14% (w/v), or between about 9% and about 12%, or to a final concentration of about 5%, or 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, or 25%. In one embodiment, the final protein concentration of the solution is between about 9% and about 11%, more preferably about 10%.

[00191] The formulated bulk solution is further sterilized by filtering through a membrane filter with an absolute pore size of no more than about 0.22 micron, for example about 0.2 micron. Then the solution is aseptically dispensed into final containers for proper sealing, with samples taken for testing.

[00192] In one embodiment, the IgG composition is further adjusted to a concentration of $10.2 \pm 0.2\%$ (w/v) with diafiltration buffer. In another embodiment, the IgG composition is adjusted

to a concentration of $15\pm 1\%$ (w/v). In another embodiment, the IgG composition is adjusted to a concentration of $20\pm 1\%$ (w/v). In another embodiment, the IgG composition is adjusted to a concentration of $25\pm 1\%$ (w/v). The pH is adjusted to about 4.4 to about 4.9 if necessary. Finally, the solution is sterile filtered and incubated for three weeks at or about 30°C .

D. Immunoglobulin G (IgG)

[00193] In one aspect, the present invention provides a method for preparing an enriched immunoglobulin G (IgG) composition from a Cohn pool, the method comprising the steps of: co-precipitating IgG and alpha-1-antitrypsin (A1PI) from a Cohn pool, in a first precipitation step, to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing the A1PI from the suspension; and recovering the soluble fraction of the first suspension containing IgG, thereby forming an enriched IgG composition. Generally, any chemical means for the precipitation of immunoglobulins and A1PI may be used, including but not limited to, alcohol precipitation (*e.g.*, using ethanol or methanol), precipitation using water soluble polymers (*e.g.*, PEG or dextran), and salting out (*e.g.*, using ammonium phosphate, ammonium sulfate, sodium citrate, *etc.*). In a preferred embodiment, the precipitation is alcohol precipitation, preferably ethanol precipitation.

[00194] In one embodiment, the present invention provides a method for preparing an enriched immunoglobulin G (IgG) composition by precipitating IgG from cryo-poor plasma in a low pH, high alcohol precipitation step. In a specific embodiment, the method comprises the steps of: precipitating IgG from cryo-poor plasma in a first precipitation step by adding ethanol to the cryo-poor plasma to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; separating the first precipitate from the first supernatant; and recovering IgG from the first precipitate, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the first precipitate is further enriched.

[00195] In one embodiment, the method comprises the steps of: precipitating IgG from cryo-poor plasma in a first precipitation step by adding ethanol to the cryo-poor plasma to a final concentration and pH selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7 to form a first precipitate and a first supernatant;

separating the first precipitate from the first supernatant; and recovering IgG from the first precipitate, thereby forming an enriched IgG composition. In one embodiment, IgG recovered from the first precipitate is further enriched.

[00196] In one embodiment, the first precipitation step is performed by admixing ethanol with the Cohn pool to a final concentration of from 20% to 30% (v/v) at a pH between 5.0 and 6.0. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 3\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 2\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 1\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of 25% (v/v). Likewise, in one embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.5 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.4 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.3 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.2 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.1 . In another embodiment, the first precipitation step is performed at a pH of 5.5. In yet other embodiments, the final ethanol concentration and pH of the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7.

[00197] The methods provided herein provide significantly higher yields of IgG recovery in the final enriched product due to the precipitation of a majority of the immunoglobulin content of the starting Cohn pool in the initial precipitation reaction, as compared to state of the art purification procedures that rely on initial low alcohol precipitation steps (*i.e.*, Fraction I precipitation).

[00198] Accordingly, in one embodiment of the methods provided herein, at least 90% of the IgG content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a preferred embodiment, at least 95% of the IgG content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a more preferred embodiment, at least 99% of the IgG content of the starting Cohn pool is precipitated in the initial precipitation reaction. In certain embodiments, at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more of the IgG content of the starting Cohn pool is precipitated in the initial precipitation reaction.

[00199] As shown in the examples provided herein, the use of an initial low pH, high alcohol precipitation step (*i.e.*, Fraction I-IV-1 precipitation) results in the precipitation of at least 99% of the IgG content of the starting Cohn pool. Accordingly, in one embodiment, the present invention provides a method for preparing an enriched IgG composition, the method comprising the steps of: precipitating from 99% to 100% of the IgG content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; separating the first precipitate from the first supernatant; and recovering IgG from the first precipitate, thereby forming an enriched immunoglobulin composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the first precipitate is further enriched. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7.

[00200] In one embodiment, the present invention provides a method for preparing an enriched IgG composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing the A1PI from the suspension; and recovering the soluble fraction of the first suspension, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the first precipitate is further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step.

[00201] In one embodiment, the present invention provides a method for preparing an enriched IgG composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn

pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing the A1PI from the suspension; and recovering the soluble fraction of the first suspension, wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the first precipitate is further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In yet another embodiment, albumin found in the first supernatant is further enriched. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7.

[00202] In order to provide pharmaceutical grade IgG compositions, contaminants such as A1PI and fibrinogen present in the initial precipitate need to be removed from the immunoglobulin composition. This may be achieved, for example, by further fractionation of the first precipitate (*e.g.*, by differential precipitation using alcohol, non-ionic hydrophilic polymers, or salting-out), chromatographic methodologies, or filtration methodologies.

[00203] In one embodiment, the first precipitate is suspended in Water for Injection (WFI) or a low ionic strength buffer suitable to extract IgG from the precipitate, the suspension is then treated with finely divided silicon dioxide (SiO_2), and the soluble portion of the suspension containing the IgG is separated from the insoluble portion of the suspension containing the bulk of the A1PI and fibrinogen.

[00204] Suitable solutions for the extraction of the first precipitate will generally have a pH between 4.0 and 5.5. In certain embodiments, the solution will have a pH between 4.5 and 5.0, in other embodiments, the extraction solution will have a pH of about 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5.0, 5.1, 5.2, 5.3, 5.4, or 5.5. In a preferred embodiment, the pH of the extraction buffer is 4.7 ± 0.1 . In another preferred embodiment, the pH of the extraction buffer is 4.8 ± 0.1 . In another preferred embodiment, the pH of the extraction buffer is 4.9 ± 0.1 . Generally, these pH requirements can be met using a buffering agent selected from, for example, acetate,

citrate, monobasic phosphate, dibasic phosphate, mixtures thereof, and the like. Suitable buffer concentrations typically range from 5 to 100 mM, or from 10 to 50 mM, or 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 mM buffering agent.

[00205] The extraction buffer will preferably have a conductivity of from 0.5 mS/cm to 2.0 mS/cm. For example, in certain embodiments, the conductivity of the extraction buffer will be 0.5 ± 0.1 mS/cm, or 0.6 ± 0.1 , 0.7 ± 0.1 , 0.8 ± 0.1 , 0.9 ± 0.1 , 1.0 ± 0.1 , 1.1 ± 0.1 , 1.2 ± 0.1 , 1.3 ± 0.1 , 1.4 ± 0.1 , 1.5 ± 0.1 , 1.6 ± 0.1 , 1.7 ± 0.1 , 1.8 ± 0.1 , 1.9 ± 0.1 , or 2.0 ± 0.1 mS/cm. One of ordinary skill in the art will know how to generate extraction buffers having an appropriate conductivity. In one particular embodiment, the extraction buffer contains 5 mM monobasic sodium phosphate and 5 mM acetate at a pH of 4.8 ± 0.2 and conductivity of from 0.7 to 0.9 mS/cm.

[00206] In one embodiment, fumed silica is added prior to separation (*e.g.*, filtration or centrifugation) at a concentration of between 20 g/kg Fraction I-IV-1 precipitate and 100 g/kg Fraction I-IV-1 precipitate. In another embodiment, fumed silica is added at a concentration of between 30 g/kg Fraction I-IV-1 precipitate and 80 g/kg Fraction I-IV-1 precipitate. In certain embodiments, the fumed silica may be added at a concentration of about 20 g/kg Fraction I-IV-1 precipitate, or about 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 g/kg Fraction I-IV-1 precipitate. In one specific embodiment, fumed silica (*e.g.*, Aerosil 380 or equivalent) is added to the Fraction I-IV-1 suspension to a final concentration of 40 ± 20 g/kg Fraction I-IV-1 precipitate. In another specific embodiment, fumed silica is added to the Fraction I-IV-1 suspension to a final concentration of 40 ± 10 g/kg Fraction I-IV-1 precipitate. Mixing takes place at 2°C to 8°C for at least 50 to 70 minutes.

[00207] Accordingly, in one embodiment, the present invention provides a method for preparing an enriched IgG composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains IgG and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp, thereby forming an enriched IgG composition. In one

embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the first precipitate is further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, SiO_2 is added to a final concentration of 40 ± 10 g/kg Fraction I-IV-1 precipitate.

[00208] In one embodiment, the present invention provides a method for preparing an enriched IgG composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains IgG and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp, and further wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG present in the soluble portion of the suspension is further enriched. In another embodiment, A1PI present in the insoluble portion of the suspension is further enriched. In another embodiment, fibrinogen present in the insoluble portion of the suspension is further enriched. In another embodiment, Factor H present in the insoluble portion of the suspension is further enriched. In another embodiment, IaIp present in the insoluble portion of the suspension is further enriched. In yet another embodiment, albumin found in the first supernatant is further enriched. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a specific embodiment, the soluble and insoluble portions of the first

suspension are separated by filtration. In one embodiment, SiO₂ is added to a final concentration of 40±10 g/kg Fraction I-IV-1 precipitate.

[00209] In one embodiment, IgG recovered from the soluble portion of the first suspension is further enriched by fractionation. Generally, any method of fractionation (*e.g.*, alcohol or polymer precipitation, salting out, *etc.*) may be used. In a preferred embodiment, IgG is further enriched by fractionating the soluble portion of the first suspension by ethanol precipitation. In one embodiment, IgG is enriched by the addition of ethanol to the soluble portion of the first suspension at a final concentration and pH suitable to precipitate the IgG, while at least one contaminant is not precipitated. In another embodiment, IgG is enriched by the addition of ethanol to the soluble portion of the first suspension at a final concentration and pH suitable to precipitate at least one contaminant, while IgG is not precipitated.

[00210] In one embodiment, the present invention provides a method for preparing an enriched IgG composition from a Cohn pool, the method comprising the steps of: co-precipitating IgG and alpha-1-antitrypsin (A1PI) from a Cohn pool, in a first precipitation step, to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing A1PI from the suspension; precipitating IgG from the first suspension in a second precipitation step, to form a second precipitate comprising IgG and a second supernatant comprising at least one contaminant; and recovering IgG from the second precipitate, thereby forming an enriched IgG composition. In one embodiment, IgG recovered from the second precipitate is further enriched. In another embodiment, A1PI removed from the first suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7.

[00211] In an exemplary embodiment, the second precipitation step is performed by admixing ethanol to the soluble portion of the second suspension (*e.g.*, a filtrate or supernatant of the suspension formed after filtration or centrifugation) at a final concentration of from 22% to 28% (v/v) at a pH between 6.5 and 7.5. In one embodiment, ethanol is admixed to a final concentration of 25±3% (v/v). In another embodiment, ethanol is admixed to a final concentration of 25±2% (v/v). In another embodiment, ethanol is admixed to a final concentration of 25±1% (v/v). In another embodiment, ethanol is admixed to a final

concentration of 25% (v/v). Likewise, in one embodiment, the second precipitation step is performed at a pH of 7.0 ± 0.5 . In another embodiment, the second precipitation step is performed at a pH of 7.0 ± 0.4 . In another embodiment, the second precipitation step is performed at a pH of 7.0 ± 0.3 . In another embodiment, the second precipitation step is performed at a pH of 7.0 ± 0.2 . In another embodiment, the second precipitation step is performed at a pH of 7.0 ± 0.1 . In another embodiment, the second precipitation step is performed at a pH of 7.0. In a particular embodiment, the second precipitation step is performed using a final ethanol concentration of $25 \pm 3\%$ (v/v) at a pH of 7.0 ± 0.3 .

[00212] In a particular embodiment, the method comprises the steps of: precipitating from 99% to 100% of IgG content from a Cohn pool plasma in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; separating the first precipitate from the first supernatant; suspending the first precipitate to form a first suspension; precipitating IgG from the first suspension in a second precipitation step, to form a second precipitate comprising IgG and a second supernatant comprising at least one contaminant; and recovering IgG from the second precipitate, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7.

[00213] In a specific embodiment, the method comprises the steps of: precipitating from 99% to 100% of IgG content from a Cohn pool plasma in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; separating the first precipitate from the first supernatant; suspending the first precipitate to form a first suspension; precipitating IgG from the first suspension in a second precipitation step by admixing ethanol to the first suspension at a final concentration of from $25 \pm 3\%$ (v/v) at a pH of 7.0 ± 0.3 , to form a second precipitate comprising IgG and a second supernatant comprising at least one contaminant; and recovering IgG from the second precipitate, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 3\%$ (v/v). In one embodiment, IgG

recovered from the second precipitate is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7.

[00214] In one embodiment, the method comprises the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing A1PI from the suspension; recovering the soluble fraction of the first suspension; precipitating IgG from the first suspension in a second precipitation step, to form a second precipitate comprising IgG and a second supernatant comprising at least one contaminant; and recovering IgG from the second precipitate, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In one embodiment, A1PI separated from the first suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, between 99% and 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step.

[00215] In a specific embodiment, the method comprises the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing A1PI from the suspension; recovering the soluble fraction of the first suspension; precipitating IgG from the first suspension in a second precipitation step by admixing ethanol to the first suspension at a final concentration of from $25\pm 3\%$ (v/v) at a pH of 7.0 ± 0.3 , to form a second precipitate comprising IgG and a second supernatant comprising at least one contaminant; and recovering IgG from the second precipitate, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In one embodiment, A1PI separated from the first suspension is further enriched. In

certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, between 99% and 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step.

[00216] In another embodiment, the method comprises the steps of: precipitating from 99% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing the A1PI from the suspension; recovering the soluble portion of the first suspension; precipitating IgG from the soluble portion of the first suspension in a second precipitation step, to form a second precipitate comprising IgG and a second supernatant comprising at least one contaminant; and recovering IgG from the second precipitate, wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In one embodiment, A1PI separated from the first suspension is further enriched. In another embodiment, albumin present in the first supernatant is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, between 99% and 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In a preferred embodiment, between 90% and 100% of the albumin content of the Cohn pool is present in the first supernatant.

[00217] In another embodiment, the method comprises the steps of: precipitating from 99% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; removing the A1PI from the suspension; recovering the soluble portion of the first suspension; precipitating IgG from the soluble portion of the first suspension in a second precipitation step by admixing ethanol to the soluble portion of the first suspension at a final concentration of from $25\pm 3\%$ (v/v) at a pH of 7.0 ± 0.3 , to form a second precipitate comprising IgG and a second

supernatant comprising at least one contaminant; and recovering IgG from the second precipitate, wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In one embodiment, A1PI separated from the first suspension is further enriched. In another embodiment, albumin present in the first supernatant is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, between 99% and 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In a preferred embodiment, between 90% and 100% of the albumin content of the Cohn pool is present in the first supernatant.

[00218] In one embodiment, the present invention provides a method for preparing an enriched IgG composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains IgG and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp, recovering the soluble portion of the first suspension; precipitating IgG from the soluble portion of the first suspension in a second precipitation step, to form a second precipitate comprising IgG and a second supernatant comprising at least one contaminant; and recovering IgG from the second precipitate, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the first precipitate is further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In a specific embodiment, the soluble and

insoluble portions of the first suspension are separated by filtration. In one embodiment, SiO_2 is added to a final concentration of 40 ± 10 g/kg Fraction I-IV-1 precipitate.

[00219] In a specific embodiment, the present invention provides a method for preparing an enriched IgG composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains IgG and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp, recovering the soluble portion of the first suspension; precipitating IgG from the soluble portion of the first suspension in a second precipitation step by admixing ethanol to the soluble portion of the first suspension at a final concentration of from $25 \pm 3\%$ (v/v) at a pH of 7.0 ± 0.3 , to form a second precipitate comprising IgG and a second supernatant comprising at least one contaminant; and recovering IgG from the second precipitate, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 3\%$ (v/v). In one embodiment, IgG recovered from the first precipitate is further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, SiO_2 is added to a final concentration of 40 ± 10 g/kg Fraction I-IV-1 precipitate.

[00220] In one embodiment, the present invention provides a method for preparing an enriched IgG composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension;

treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains IgG and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp; precipitating IgG from the soluble portion of the first suspension in a second precipitation step, to form a second precipitate comprising IgG and a second supernatant comprising at least one contaminant; and recovering IgG from the second precipitate, wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In another embodiment, A1PI present in the insoluble portion of the suspension is further enriched. In another embodiment, fibrinogen present in the insoluble portion of the suspension is further enriched. In another embodiment, Factor H present in the insoluble portion of the suspension is further enriched. In another embodiment, IaIp present in the insoluble portion of the suspension is further enriched. In yet another embodiment, albumin found in the first supernatant is further enriched. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, SiO_2 is added to a final concentration of 40 ± 10 g/kg Fraction I-IV-1 precipitate.

[00221] In one embodiment, the present invention provides a method for preparing an enriched IgG composition, the method comprising the steps of: precipitating from 95% to 100% of IgG and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains IgG and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp; precipitating IgG from the soluble portion of the first suspension in a second precipitation step by admixing ethanol to the soluble portion of the first

suspension at a final concentration of from $25\pm 3\%$ (v/v) at a pH of 7.0 ± 0.3 , to form a second precipitate comprising IgG and a second supernatant comprising at least one contaminant; and recovering IgG from the second precipitate, wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched IgG composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, IgG recovered from the second precipitate is further enriched. In another embodiment, A1PI present in the insoluble portion of the suspension is further enriched. In another embodiment, fibrinogen present in the insoluble portion of the suspension is further enriched. In another embodiment, Factor H present in the insoluble portion of the suspension is further enriched. In another embodiment, IaIp present in the insoluble portion of the suspension is further enriched. In yet another embodiment, albumin found in the first supernatant is further enriched. In a preferred embodiment, from 99% to 100% of the IgG content of the Cohn pool is precipitated in the first precipitation step. In certain embodiments, the final ethanol concentration and pH used in the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, SiO_2 is added to a final concentration of 40 ± 10 g/kg Fraction I-IV-1 precipitate.

[00222] In one embodiment, IgG are recovered from the second precipitate by suspending the precipitate with a cold extraction buffer. For example, the second precipitate is suspended at a ratio of 1 part precipitate to 2-5 parts Water for Injection (WFI) or low conductivity buffer. In a preferred embodiment, the second precipitate suspended at a ratio of 1 part precipitate to 4-5 parts, preferably 3.5 parts, WFI. In one embodiment, the suspension step is performed at a temperature between 0°C and 8°C . In one embodiment, the final pH of the solution is adjusted to from 4.8 to 5.6, preferably to 5.2 ± 0.2 . In one embodiment, this pH adjustment is performed with acetic acid. In one embodiment, the conductivity of the suspension is increased to between 2.5 and 6.0 mS/cm to increase the solubility of IgG. In one embodiment, the conductivity is increased by the addition of sodium chloride.

[00223] Suitable solutions for the extraction of the second precipitate include WFI and low conductivity buffers. In one embodiment, a low conductivity buffer has a conductivity of less

than about 10 mS/cm. In other embodiments, the low conductivity buffer has a conductivity of less than about 9, 8, 7, 6, 5, 4, 3, 2, or 1 mS/cm. In a preferred embodiment, the low conductivity buffer has a conductivity of less than about 6 mS/cm. In another preferred embodiment, the low conductivity buffer has a conductivity of less than about 4 mS/cm. In another preferred embodiment, the low conductivity buffer has a conductivity of less than about 2 mS/cm.

[00224] In one embodiment, the soluble portion of the first suspension, containing IgG, is separated from the insoluble portion. In one embodiment, this is done by filtering the suspension with a depth filter having a nominal pore size of from 0.1 μm and 0.4 μm . In one embodiment, the nominal pore size of the depth filter is 0.2 μm (*e.g.*, Cuno VR06 filter or equivalent). In one embodiment, the filter is washed with WFI or a suitable buffer after filtration to recover additional IgG and the post-wash added to the filtrate. In a preferred embodiment, the post-wash of the filter is performed using a sodium chloride solution with a conductivity of between about 2.5 and about 6.0 mS/cm. In another embodiment, the second suspension is centrifuged to recover the soluble portion.

[00225] In one aspect of the present invention, the methods provided for preparing an enriched IgG composition further comprise purifying at least one additional blood factor from the same Cohn pool. In one embodiment, the methods further comprise purifying at least two additional blood proteins from the same Cohn pool. In another embodiment, the methods further comprise purifying at least three additional blood proteins from the same Cohn pool. In yet other embodiments, the methods further comprise purifying at least four, five, six, or more additional blood proteins from the same Cohn pool. Exemplary blood proteins that may be co-purified from the same Cohn pool as IgG include, without limitation, albumin, alpha-1-antitrypsin (A1PI), Factor H, inter-alpha-inhibitor proteins (IaIp), Prothrombin complexes, Factor VII (FVII), Factor VIII (FVIII), antithrombin III (ATIII), fibrinogen, butyrylcholinesterase, and others.

[00226] In one embodiment, alpha-1-antitrypsin (A1PI) is further purified from the same Cohn pool as IgG. In a specific embodiment, A1PI is purified from the insoluble fraction of the first suspension.

[00227] In another embodiment, fibrinogen is further purified from the same Cohn pool as IgG. In a specific embodiment, fibrinogen is purified from the insoluble fraction of the first suspension. In another specific embodiment, both A1PI and fibrinogen are purified from the insoluble fraction of the first suspension.

[00228] In another embodiment, Factor H is further purified from the same Cohn pool as IgG. In a specific embodiment, Factor H is purified from the insoluble fraction of the first suspension. In another specific embodiment, both A1PI and Factor H are purified from the insoluble fraction of the first suspension

[00229] In another embodiment, inter-alpha-inhibitor proteins (IaIp) are further purified from the same Cohn pool as IgG. In a specific embodiment, IaIp is purified from the insoluble fraction of the first suspension. In another specific embodiment, both A1PI and IaIp are purified from the insoluble fraction of the first suspension.

[00230] In another embodiment, albumin is further purified from the same Cohn pool as IgG. In a specific embodiment, albumin is purified from the first supernatant formed. In a specific embodiment, albumin is purified from the first supernatant formed, while at least one additional blood factor is purified from the insoluble portion of the first suspension formed.

[00231] In one embodiment, immunoglobulins recovered from an initial low pH, high alcohol precipitation step are further enriched according to a downstream Kistler-Nitschmann or Deutsch *et al.* fractionation. For example, a recovered immunoglobulin composition may be subjected to a step similar to a Kistler-Nitschmann B precipitation or Deutsch *et al.* β -globulin precipitation. The conditions used in these steps result in the precipitation of α - and β -globulin, while immunoglobulin G remains in the supernatant.

[00232] Accordingly, in one aspect the present invention provides a method for preparing an enriched immunoglobulin G (IgG) composition comprising the steps of (i) co-precipitating immunoglobulins and alpha-1-antitrypsin (A1PI) from a Cohn pool in an initial low pH, high alcohol precipitation; (ii) recovering immunoglobulins from the precipitate; (iii) precipitating at least one non-gamma globulin protein from the immunoglobulin composition recovered from the first precipitate; and (iv) recovering the supernatant from the second precipitation step. In certain embodiments, these methods may further comprise additional precipitation steps (*e.g.*, a

Precipitate G precipitation), an anion and/or cation exchange step, one or more ultrafiltration/diafiltration steps, and one or more viral reduction or inactivation steps (*e.g.*, S/D treatment, nanofiltration, incubation at low pH, *etc.*).

V. Preparation of Alpha-1-Antitrypsin (A1PI) Compositions

[00233] Modern methods for manufacturing blood proteins from donated plasma are derived from early work done in the 1940's and 1950's. Since this work was focused primarily on the purification of albumin and gamma immunoglobulins (IgG) the fractionation schemes developed were not optimized for the recovery of additional blood proteins, which are becoming increasingly more therapeutically important. For example, the use of an initial high pH, low alcohol precipitation of plasma (*e.g.*, Fraction I precipitation) results in the partial loss of Factor H, IaIp, and to some extent immunoglobulins in the Fraction I precipitate. Furthermore, the separation of immunoglobulins and A1PI achieved by subsequent precipitation steps (*e.g.*, Fraction II+III precipitation or Precipitation A) is incomplete, with immunoglobulins lost in the supernatant due to incomplete precipitation and A1PI lost in the precipitate due to partial precipitation. Accordingly, updated fractionation methods that provide higher yields of multiple blood proteins are needed.

[00234] The present invention provides methods that satisfy these requirements through the use of an initial precipitation step that fractionates immunoglobulins, A1PI, and other blood proteins in an initial precipitate and albumin in an initial supernatant. In a preferred embodiment, this initial precipitation is performed under low pH, high alcohol conditions.

[00235] In one aspect, the present invention provides a method for the manufacture of alpha-1-antitrypsin (A1PI) from pooled plasma using an initial step that precipitates the bulk of the immunoglobulin and A1PI content of a Cohn pool. In a particular embodiment, this initial step is a low pH, high alcohol precipitation step. As shown in the examples provided herein, greater than 95% of the A1PI content of the starting Cohn pool is precipitated under these conditions. Furthermore, the present invention provides methods for efficiently separating A1PI and immunoglobulins in this initial precipitation.

[00236] In one aspect, the present invention provides a method for preparing an enriched alpha-1-antitrypsin (A1PI) composition from a Cohn pool, the method comprising the steps of: co-precipitating immunoglobulins and A1PI from a Cohn pool, in a first precipitation step, to form a first precipitate and a first supernatant; solubilizing immunoglobulins present in the first precipitate, to form a first suspension having a soluble portion comprising immunoglobulins and an insoluble portion comprising A1PI; separating the soluble and insoluble portions of the first suspension; and recovering A1PI from the insoluble portion of the first suspension, thereby forming an enriched A1PI composition. Generally, any chemical means for the precipitation of immunoglobulins and A1PI may be used, including but not limited to, alcohol precipitation (*e.g.*, using ethanol or methanol), precipitation using water soluble polymers (*e.g.*, PEG or dextran), and salting out (*e.g.*, using ammonium phosphate, ammonium sulfate, sodium citrate, *etc.*). In a preferred embodiment, the precipitation is alcohol precipitation, preferably ethanol precipitation.

[00237] In one embodiment, the present invention provides a method for preparing an enriched A1PI composition by precipitating A1PI from cryo-poor plasma in a low pH, high alcohol precipitation step. In a specific embodiment, the method comprises the steps of: precipitating A1PI from cryo-poor plasma in a first precipitation step by adding ethanol to the cryo-poor plasma to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; separating the first precipitate from the first supernatant; and recovering A1PI from the first precipitate, thereby forming an enriched A1PI composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 3\%$ (v/v). In one embodiment, A1PI recovered from the first precipitate is further enriched.

[00238] In one embodiment, the method comprises the steps of: precipitating A1PI from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration and pH selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7 to form a first precipitate and a first supernatant; separating the first precipitate from the first supernatant; and recovering A1PI from the first precipitate, thereby forming an enriched A1PI composition. In one embodiment, A1PI recovered from the first precipitate is further enriched.

[00239] In one embodiment, the first precipitation step is performed by admixing ethanol with the Cohn pool to a final concentration of from 22% to 28% (v/v) at a pH between 5.0 and 6.0. In

one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 2\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 1\%$ (v/v). In another embodiment, ethanol is admixed with the Cohn pool to a final concentration of 25% (v/v). Likewise, in one embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.5 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.4 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.3 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.2 . In another embodiment, the first precipitation step is performed at a pH of 5.5 ± 0.1 . In another embodiment, the first precipitation step is performed at a pH of 5.5. In yet other embodiments, the final ethanol concentration and pH of the first precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7.

[00240] The methods provided herein provide significantly higher yields of A1PI recovery in the final enriched product due to the precipitation of a majority of the A1PI content of the starting Cohn pool in the initial precipitation reaction, as compared to state of the art purification procedures that rely on initial low alcohol precipitation steps (*i.e.*, Fraction I precipitation).

[00241] Accordingly, in one embodiment of the methods provided herein, at least 80% of the alpha-1-antitrypsin (A1PI) content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a preferred embodiment, at least 90% of the A1PI content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a more preferred embodiment, at least 95% of the A1PI content of the starting Cohn pool is precipitated in the initial precipitation reaction. In a most preferred embodiment, at least 97% of the A1PI content of the starting Cohn pool is precipitated in the initial precipitation reaction. In certain embodiments, at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more of the A1PI content of the starting Cohn pool is precipitated in the initial precipitation reaction.

[00242] As shown in the examples provided herein, the use of an initial low pH, high alcohol precipitation step (*i.e.*, Fraction I-IV-1 precipitation) results in the precipitation of at least 95% of the A1PI content of the starting Cohn pool. Accordingly, in one embodiment, the present invention provides a method for preparing an enriched A1PI composition, the method

comprising the steps of: precipitating from 95% to 100% of the A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; separating the first precipitate from the first supernatant; and recovering A1PI from the first precipitate, thereby forming an enriched A1PI composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, A1PI recovered from the first precipitate is further enriched. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7.

[00243] In one embodiment, the present invention provides a method for preparing an enriched A1PI composition, the method comprising the steps of: precipitating from 95% to 100% of immunoglobulin and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; solubilizing immunoglobulins present in the first precipitate, to form a first suspension having a soluble portion comprising immunoglobulins and an insoluble portion comprising A1PI; separating the soluble and insoluble portions of the first suspension; and recovering A1PI from the insoluble portion of the first suspension, thereby forming an enriched A1PI composition. In one embodiment, immunoglobulins recovered from the soluble portion of the first suspension are further enriched. In another embodiment, A1PI removed from the insoluble fraction of the first suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, from 99% to 100% of the immunoglobulin content of the Cohn pool is precipitated in the first precipitation step.

[00244] In one embodiment, the present invention provides a method for preparing an enriched A1PI composition, the method comprising the steps of: precipitating from 95% to 100% of immunoglobulin and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; solubilizing immunoglobulins present in the first

precipitate, to form a first suspension having a soluble portion comprising immunoglobulins and an insoluble portion comprising A1PI; separating the soluble and insoluble portions of the first suspension; and recovering A1PI from the insoluble portion of the first suspension, wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant, thereby forming an enriched A1PI composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25\pm 3\%$ (v/v). In one embodiment, immunoglobulins recovered from the soluble portion of the first suspension are further enriched. In another embodiment, A1PI removed from the insoluble portion of the first suspension is further enriched. In yet another embodiment, albumin found in the first supernatant is further enriched. In a preferred embodiment, from 99% to 100% of the immunoglobulin content of the Cohn pool is precipitated in the first precipitation step. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7.

[00245] In one embodiment, the first precipitate is suspended in Water for Injection (WFI) or a low ionic strength buffer suitable to extract immunoglobulins from the precipitate, the suspension is then treated with finely divided silicon dioxide (SiO_2), and the soluble portion of the suspension containing the immunoglobulins is separated from the insoluble portion of the suspension containing the bulk of the A1PI content.

[00246] Suitable solutions for the extraction of immunoglobulins from the first precipitate will generally have a pH between 4.0 and 5.5. In certain embodiments, the solution will have a pH between 4.5 and 5.0, in other embodiments, the extraction solution will have a pH of about 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5.0, 5.1, 5.2, 5.3, 5.4, or 5.5. In a preferred embodiment, the pH of the extraction buffer is 4.7 ± 0.1 . In another preferred embodiment, the pH of the extraction buffer is 4.8 ± 0.1 . In another preferred embodiment, the pH of the extraction buffer is 4.9 ± 0.1 . Generally, these pH requirements can be met using a buffering agent selected from, for example, acetate, citrate, monobasic phosphate, dibasic phosphate, mixtures thereof, and the like. Suitable buffer concentrations typically range from 5 to 100 mM, or from 10 to 50 mM, or 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 mM buffering agent.

[00247] The extraction buffer will preferably have a conductivity of from 0.5 mS/cm to 2.0 mS/cm. For example, in certain embodiments, the conductivity of the extraction buffer will be

0.5±0.1 mS/cm, or 0.6±0.1, 0.7±0.1, 0.8±0.1, 0.9±0.1, 1.0±0.1, 1.1±0.1, 1.2±0.1, 1.3±0.1, 1.4±0.1, 1.5±0.1, 1.6±0.1, 1.7±0.1, 1.8±0.1, 1.9±0.1, or 2.0±0.1 mS/cm. One of ordinary skill in the art will know how to generate extraction buffers having an appropriate conductivity. In one particular embodiment, the extraction buffer contains 5 mM monobasic sodium phosphate and 5 mM acetate at a pH of 4.8 ± 0.2 and conductivity of from 0.7 to 0.9 mS/cm.

[00248] In one embodiment, fumed silica is added prior to separation (*e.g.*, filtration or centrifugation) at a concentration of between 20 g/kg Fraction I-IV-1 precipitate and 100 g/kg Fraction I-IV-1 precipitate. In another embodiment, fumed silica is added at a concentration of between 30 g/kg Fraction I-IV-1 precipitate and 80 g/kg Fraction I-IV-1 precipitate. In certain embodiments, the fumed silica may be added at a concentration of about 20 g/kg Fraction I-IV-1 precipitate, or about 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 g/kg Fraction I-IV-1 precipitate. In one specific embodiment, fumed silica (*e.g.*, Aerosil 380 or equivalent) is added to the Fraction I-IV-1 suspension to a final concentration of 40±20 g/kg Fraction I-IV-1 precipitate. In another specific embodiment, fumed silica is added to the Fraction I-IV-1 suspension to a final concentration of 40±10 g/kg Fraction I-IV-1 precipitate. Mixing takes place at 2°C to 8°C for at least 50 to 70 minutes.

[00249] Accordingly, in one embodiment, the present invention provides a method for preparing an enriched A1PI composition, the method comprising the steps of: precipitating from 95% to 100% of immunoglobulin and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of between 20% and 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains immunoglobulins and the insoluble portion of the suspension contains A1PI; and recovering A1PI from the insoluble portion of the first suspension, thereby forming an enriched A1PI composition. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of 25±3% (v/v). In one embodiment, immunoglobulins recovered from the first precipitate are further enriched. In another embodiment, A1PI removed from the suspension is further enriched. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is

selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a preferred embodiment, from 99% to 100% of the immunoglobulin content of the Cohn pool is precipitated in the first precipitation step. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, SiO_2 is added to a final concentration of 40 ± 10 g/kg Fraction I-IV-1 precipitate.

[00250] In one embodiment, the present invention provides a method for preparing an enriched A1PI composition, the method comprising the steps of: precipitating from 95% to 100% of immunoglobulin and A1PI content from a Cohn pool in a first precipitation step by adding ethanol to the Cohn pool to a final concentration of from 20% to 30% at a pH from 5.0 to 6.0 to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; treating the first suspension with finely divided silicon dioxide; and separating the soluble portion of the suspension from the insoluble portion of the suspension, wherein the soluble portion of the suspension contains immunoglobulins and the insoluble portion of the suspension contains A1PI, fibrinogen, Factor H, and IaIp, and further wherein from 80% to 100% of the albumin content of the Cohn pool is present in the first supernatant. In one embodiment, ethanol is admixed with the Cohn pool to a final concentration of $25 \pm 3\%$ (v/v). In one embodiment, immunoglobulins present in the soluble portion of the suspension are further enriched. In another embodiment, A1PI present in the insoluble portion of the suspension is further enriched. In another embodiment, fibrinogen present in the insoluble portion of the suspension is further enriched. In another embodiment, Factor H present in the insoluble portion of the suspension is further enriched. In another embodiment, IaIp present in the insoluble portion of the suspension is further enriched. In yet another embodiment, albumin found in the first supernatant is further enriched. In a preferred embodiment, from 99% to 100% of the immunoglobulin content of the Cohn pool is precipitated in the first precipitation step. In certain embodiments, the final ethanol concentration and pH used in the low pH, high alcohol precipitation step is selected from variations 1 to 2166, as listed in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7. In a specific embodiment, the soluble and insoluble portions of the first suspension are separated by filtration. In one embodiment, SiO_2 is added to a final concentration of 40 ± 10 g/kg Fraction I-IV-1 precipitate.

[00251] In one embodiment, the soluble portion of the first suspension, containing IgG, is separated from the insoluble portion, containing A1PI by filtering the suspension with a depth filter having a nominal pore size of from 0.1 μm and 0.4 μm . In one embodiment, the nominal pore size of the depth filter is 0.2 μm (*e.g.*, Cuno VR06 filter or equivalent). In one embodiment, the filter is washed with WFI or a suitable buffer after filtration to recover additional IgG and the post-wash added to the filtrate. In another embodiment, the first suspension is centrifuged to separate the soluble and insoluble portions.

[00252] In one embodiment, A1PI is recovered from the separated insoluble portion by suspending the insoluble portion in an A1PI extraction buffer. For example, in one embodiment, the separated insoluble portion is suspended at a ratio of 1 part insoluble material to 2-15 parts Water for Injection (WFI) or low conductivity buffer, to form a second suspension. In another embodiment, the separated insoluble portion is suspended at a ratio of 1 part insoluble material to 2-10 parts Water for Injection (WFI) or low conductivity buffer. In another embodiment, the separated insoluble portion is suspended at a ratio of 1 part insoluble material to 3-7 parts Water for Injection (WFI) or low conductivity buffer. In yet other embodiments, the separated insoluble portion is suspended at a ratio of 1 part insoluble material to 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more parts Water for Injection (WFI) or low conductivity buffer. In one embodiment, the extraction is stirred for at least 2 hours. In another embodiment, the extraction is stirred for at least 4 hours. In another embodiment, the extraction is stirred for at least 8 hours. In another embodiment, the extraction is stirred for 6 to 12 hours. In one embodiment, the extraction is performed at from 10°C to 25°C. In another embodiment, the extraction is performed at 17 $\pm$ 5°C. In another embodiment, the extraction is performed at 17 $\pm$ 4°C. In another embodiment, the extraction is performed at 17 $\pm$ 3°C. In another embodiment, the extraction is performed at 17 $\pm$ 2°C. In another embodiment, the extraction is performed at 17 $\pm$ 1°C. In one embodiment, the soluble portion containing A1PI is separated from the insoluble portion by filtration, for example, depth filtration. In another embodiment, the soluble portion containing A1PI is separated from the insoluble portion by centrifugation.

[00253] Suitable solutions for the extraction of A1PI include WFI and low conductivity buffers. In one embodiment, a low conductivity buffer has a conductivity of less than about 10 mS/cm. In other embodiments, the low conductivity buffer has a conductivity of less than about 9, 8, 7, 6,

5, 4, 3, 2, or 1 mS/cm. In a preferred embodiment, the low conductivity buffer has a conductivity of less than about 6 mS/cm. In another preferred embodiment, the low conductivity buffer has a conductivity of less than about 4 mS/cm. In another preferred embodiment, the low conductivity buffer has a conductivity of less than about 2 mS/cm. In one embodiment, the WFI or low conductivity buffer has a pH of between 8.0 and 9.5. In a particular embodiment, the pH of the A1PI extraction buffer is 8.8 ± 0.5 . In a specific embodiment, the pH of the A1PI extraction buffer is 8.8 ± 0.4 . In a specific embodiment, the pH of the A1PI extraction buffer is 8.8 ± 0.3 . In a specific embodiment, the pH of the A1PI extraction buffer is 8.8 ± 0.2 . In a specific embodiment, the pH of the A1PI extraction buffer is 8.8 ± 0.1 . In yet other embodiments, the pH of the A1PI extraction buffer is 8.0 ± 0.2 , 8.1 ± 0.2 , 8.2 ± 0.2 , 8.3 ± 0.2 , 8.4 ± 0.2 , 8.5 ± 0.2 , 8.6 ± 0.2 , 8.7 ± 0.2 , 8.8 ± 0.2 , 8.9 ± 0.2 , 9.0 ± 0.2 , 9.1 ± 0.2 , 9.2 ± 0.2 , 9.3 ± 0.2 , 9.4 ± 0.2 , or 9.5 ± 0.2 .

[00254] In one embodiment, the separated insoluble portion of the first suspension is suspended in WFI or a low ionic strength buffer which is not suitable for extracting A1PI to form an intermediate suspension, prior to extraction using an A1PI buffer. In this embodiment, the method comprises suspending the insoluble portion in WFI or a low ionic strength buffer at a pH of no more than 7.0; separating the soluble and insoluble portions of the intermediate suspension; and recovering A1PI in the insoluble portion of the intermediate suspension. In one embodiment, the intermediate suspension is formed by suspending the insoluble material at a ratio of 1 part insoluble material to 2-15 parts WFI or buffer. In one embodiment, the WFI or low ionic strength buffer has a pH of from 4.5 to 6.5. In another embodiment, the low ionic strength buffer has a pH of 5.5 ± 0.4 . In another embodiment, the low ionic strength buffer has a pH of 5.5 ± 0.3 . In another embodiment, the low ionic strength buffer has a pH of 5.5 ± 0.2 . In another embodiment, the low ionic strength buffer has a pH of 5.5 ± 0.1 . In yet other embodiments, the low ionic strength buffer has a pH of 4.5 ± 0.2 , 4.6 ± 0.2 , 4.7 ± 0.2 , 4.8 ± 0.2 , 4.9 ± 0.2 , 5.0 ± 0.2 , 5.1 ± 0.2 , 5.2 ± 0.2 , 5.3 ± 0.2 , 5.4 ± 0.2 , 5.5 ± 0.2 , 5.6 ± 0.2 , 5.7 ± 0.2 , 5.8 ± 0.2 , 5.9 ± 0.2 , 6.0 ± 0.2 , 6.1 ± 0.2 , 6.2 ± 0.2 , 6.3 ± 0.2 , 6.4 ± 0.2 , or 6.5 ± 0.2 . In one embodiment, the soluble portion is separated from the insoluble portion containing A1PI by filtration, for example, depth filtration. In another embodiment, the soluble portion is separated from the insoluble portion containing A1PI by centrifugation.

[00255] In one embodiment, A1PI extracted from the insoluble portion of the first suspension is further enriched according to well known methods in the art, including without limitation: chromatography (*e.g.*, anion exchange chromatography, cation exchange chromatography, hydrophobic interaction chromatography (HIC), hydroxyapatite chromatography (HAP), Protein A affinity chromatography, immuno-affinity chromatography, size exclusion chromatography, *etc.*); filtration (*e.g.*, ultrafiltration and/or diafiltration); and one or more viral reduction steps (*e.g.*, nanofiltration, solvent and detergent treatment, UV irradiation, heat treatment, low pH incubation, *etc.*).

[00256] In one embodiment, A1PI extracted from the insoluble portion of the first suspension is further enriched by fractionation. Generally, any method of fractionation (*e.g.*, alcohol or polymer precipitation, salting out, *etc.*) may be used. In a preferred embodiment, the composition is further enriched by precipitating A1PI from the clarified extracted solution using a divalent cation, preferably zinc. In one embodiment, zinc chloride is added to the solution to a final concentration of from 1 mM to 15 mM. In a specific embodiment, zinc chloride is added to the solution to a final concentration of from 1 mM to 4 mM. In a more specific embodiment, zinc chloride is added to the solution to a final concentration of from 2.0 mM to 3.0 mM.

[00257] In certain embodiments, A1PI is further enriched according to purification methods known in the art, for example, those described in WO 1995/35306, WO 1998/00154, and U.S. Patent Nos. 5,981,715; 7,807,435; and 7,879,800.

VI. Pharmaceutical Compositions

[00258] In one aspect, the present invention provides compositions of plasma-derived proteins which are prepared according to any of the methods described herein. In certain embodiments, these compositions will be formulated for pharmaceutical administration (*i.e.*, pharmaceutical compositions).

[00259] In one embodiment, the present invention provides a plasma-derived protein composition prepared by a method comprising the steps of: co-precipitating immunoglobulins and alpha-1-antitrypsin (A1PI) from a Cohn pool, in a first precipitation step, to form a first

precipitate and a first supernatant; suspending the first precipitate to form a first suspension; solubilizing immunoglobulins from the first suspension to form a soluble portion of the suspension comprising immunoglobulins and an insoluble portion of the suspension comprising A1PI, fibrinogen, Factor H, and IaIp; and separating the soluble portion of the suspension from the insoluble portion of the suspension. In certain embodiments, the composition comprises a plasma-derived protein is selected from an immunoglobulin (Ig), albumin, alpha-1-antitrypsin (A1PI), butyrylcholinesterase, Factor H, a protein of the complement system, and an inter-alpha-trypsin inhibitor (IaI) protein.

[00260] Generally, any chemical means for the precipitation of immunoglobulins and A1PI may be used, including but not limited to, alcohol precipitation (*e.g.*, using ethanol or methanol), precipitation using water soluble polymers (*e.g.*, PEG or dextran), and salting out (*e.g.*, using ammonium phosphate, ammonium sulfate, sodium citrate, *etc.*). In a preferred embodiment, the precipitation is alcohol precipitation, preferably ethanol precipitation.

[00261] In one embodiment, the composition is formulated for pharmaceutical administration. In a specific embodiment, the composition is formulated for intravenous administration. In another specific embodiment, the composition is formulated for intramuscular administration. In another embodiment, the composition is formulated for subcutaneous administration. In yet another embodiment, the composition is formulated for intraocular administration.

[00262] In one embodiment, the pharmaceutical compositions provided herein are prepared by formulating a plasma-derived protein composition isolated using a method provided herein. Generally, the formulated composition will have been subjected to at least one, preferably at least two, most preferably at least three, viral inactivation or removal steps. Non-limiting examples of viral inactivation or removal steps that may be employed with the methods provided herein include, solvent detergent treatment (Horowitz et al., Blood Coagul fibrinogenolysis 1994 (5 Suppl 3):S21-S28 and Kreil et al., Transfusion 2003 (43):1023-1028), nanofiltration (Hamamoto et al., Vox Sang 1989 (56)230-236 and Yuasa et al., J Gen Virol. 1991 (72 (pt 8)):2021-2024), and low pH incubation at high temperatures (Kempf et al., Transfusion 1991 (31)423-427 and Louie et al., Biologicals 1994 (22):13-19). In certain embodiments, the composition comprises a

plasma-derived protein is selected from an immunoglobulin (Ig), albumin, alpha-1-antitrypsin (A1PI), butyrylcholinesterase, Factor H, a protein of the complement system, and an inter-alpha-trypsin inhibitor (I α I) protein.

[00263] In one embodiment, the pharmaceutical compositions provided herein will comprise one or more buffering agents or pH stabilizing agents suitable for intravenous, subcutaneous, intramuscular, and/or intraocular administration. Non-limiting examples of buffering agents suitable for formulating a plasma-derived protein composition provided herein include glycine, citrate, phosphate, acetate, glutamate, tartrate, benzoate, lactate, histidine or other amino acids, gluconate, malate, succinate, formate, propionate, carbonate, or any combination thereof adjusted to an appropriate pH. Generally, the buffering agent will be sufficient to maintain a suitable pH in the formulation for an extended period of time. In a preferred embodiment, the buffering agent is glycine. In certain embodiments, the composition comprises a plasma-derived protein is selected from an immunoglobulin (Ig), albumin, alpha-1-antitrypsin (A1PI), butyrylcholinesterase, Factor H, a protein of the complement system, and an inter-alpha-trypsin inhibitor (I α I) protein.

[00264] In some embodiments, the pharmaceutical compositions provided herein may optionally further comprise an agent for adjusting the osmolarity of the composition. Non-limiting examples of osmolarity agents include mannitol, sorbitol, glycerol, sucrose, glucose, dextrose, levulose, fructose, lactose, polyethylene glycols, phosphates, sodium chloride, potassium chloride, calcium chloride, calcium gluconoglucoheptonate, dimethyl sulfone, and the like.

[00265] Typically, the formulations provided herein will have osmolarities that are comparable to physiologic osmolarity, about 285 to 295 mOsmol/kg (Lacy et al., Drug Information Handbook – Lexi-Comp 1999:1254. In certain embodiments, the osmolarity of the formulation will be between about 200 mOsmol/kg and about 350 mOsmol/kg, preferably between about 240 and about 300 mOsmol/kg. In particular embodiments, the osmolarity of the formulation will be about 200 mOsmol/kg, or 210 mOsmol/kg, 220 mOsmol/kg, 230 mOsmol/kg, 240 mOsmol/kg, 245 mOsmol/kg, 250 mOsmol/kg, 255 mOsmol/kg, 260 mOsmol/kg, 265 mOsmol/kg, 270 mOsmol/kg, 275 mOsmol/kg, 280 mOsmol/kg, 285 mOsmol/kg, 290 mOsmol/kg, 295

mOsmol/kg, 300 mOsmol/kg, 310 mOsmol/kg, 320 mOsmol/kg, 330 mOsmol/kg, 340 mOsmol/kg, 340 mOsmol/kg, or 350 mOsmol/kg.

[00266] The plasma-derived formulations provided herein are generally stable in liquid form for an extended period of time. In certain embodiments, the formulations are stable for at least about 3 months at room temperature, or at least about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 months at room temperature. The formulation will also generally be stable 6 or at least about 18 months under refrigerated conditions (typically between about 2°C and about 8°C), or for at least about 21, 24, 27, 30, 33, 36, 39, 42, or 45 months under refrigerated conditions.

VII. Method of Treatment

[00267] In one aspect, the present invention provides methods for treating a disease or disorder associated with a blood protein deficiency or dysfunction in a subject in need thereof by administering a therapeutically effective dose of a plasma-derived protein prepared according to a method provided herein. In certain embodiments, the composition comprises a plasma-derived protein is selected from an immunoglobulin (Ig), albumin, alpha-1-antitrypsin (A1PI), butyrylcholinesterase, Factor H, a protein of the complement system, and an inter-alpha-trypsin inhibitor (I α I) protein.

[00268] In one aspect, the present invention provides the use of a plasma-derived protein composition prepared according to a method provided herein for the manufacture of a medicament for use in the treatment of a condition associated with a blood protein deficiency or dysfunction. In certain embodiments, the plasma-derived protein is selected from an immunoglobulin (Ig), albumin, alpha-1-antitrypsin (A1PI), butyrylcholinesterase, Factor H, a protein of the complement system, and an inter-alpha-trypsin inhibitor (I α I) protein.

[00269] In one embodiment, the plasma-derived protein composition is prepared by a method comprising the steps of: co-precipitating immunoglobulins and alpha-1-antitrypsin (A1PI) from a Cohn pool, in a first precipitation step, to form a first precipitate and a first supernatant; suspending the first precipitate to form a first suspension; solubilizing immunoglobulins from the first suspension to form a soluble portion of the suspension comprising immunoglobulins and an

insoluble portion of the suspension comprising A1PI, fibrinogen, Factor H, and IaI_p; and separating the soluble portion of the suspension from the insoluble portion of the suspension. In certain embodiments, the composition comprises a plasma-derived protein is selected from an immunoglobulin (Ig), albumin, alpha-1-antitrypsin (A1PI), butyrylcholinesterase, Factor H, a protein of the complement system, and an inter-alpha-trypsin inhibitor (IaI) protein.

VIII. Examples

Example 1 – NG2 C91

[00270] Immunoglobulin G (IgG) compositions are commercially manufactured by fractionating plasma through a series of precipitation reactions. In a common methodology, impurities are precipitated out of an immunoglobulin-containing cryo-poor plasma in a first alcohol precipitation step. IgG immunoglobulins are then precipitated from the resulting supernatant and subsequently further enriched. In an attempt to increase the yield of IgG recovered from human plasma during alcohol fractionation, pH values from 5.5 to 7.0 and ethanol concentrations from 20% to 25% were tested in IgG precipitation reactions.

[00271] Briefly, Fraction I supernatant, an intermediate used in the commercial manufacturing of plasma-derived Immunoglobulin G (IgG), alpha-1-antitrypsin (A1PI), and albumin products, was formed by admixing ethanol at a final concentration of 8% (v/v) to cryo-poor plasma at pH 7.2±0.2, incubating the mixture at 0°C, and separating the precipitate formed (Fraction I) from the supernatant (Fraction I supernatant). The resulting Fraction I supernatant was then aliquoted into 600 g samples. The pH of the individual samples were adjusted to from 5.5 to 7.0 and ethanol was admixed to a final concentration of from 20% to 25%, as shown in Table 9.

[00272] As reported in Table 9, conditions similar to those used to prepare a traditional Cohn Fraction II+III precipitate (25% ethanol; pH 6.8) and the modified Fraction II+III precipitate used in the manufacture of GAMMAGARD LIQUID[®] (Baxter International; Deerfield, IL) result in a loss of 2.9% and 6.7% of total IgG in the supernatant, due to incomplete precipitation. However, it was found that by reducing the pH of the reaction to 6.0 or lower and using a final ethanol concentration of at least 22.5%, IgG loss is significantly reduced. For example, the use of 22.5% alcohol at pH 5.5 resulted in the loss of only 0.4% of total IgG in the supernatant.

Similarly, the use of 25% alcohol at pH 6.0 or 5.5 resulted in the loss of only 0.5% and 0.1% of total IgG in the supernatant. As reported in Table 10, this equates to an increase of more than 300 mg IgG/L plasma and 125 mg IgG/L plasma in the precipitate as compared to the traditional and modified Fraction II+III precipitates, respectively.

Table 9. Yield of proteins in precipitation supernatant, expressed as a percent of the protein content of the starting material (Fraction I supernatant).

EtOH (%)	pH	protein biuret	IgG ELISA	IgA ELISA	IgM ELISA	fibrinogen ELISA	transferrin ELISA	a1-antitrypsin ELISA	albumin neph
20	5.5	60.8	1.1	3.5	0.2	0.2	73.9	6.1	93.1
20	6	68.0	2.9	12.8	0.5	0.2	94.3	38.9	92.4
20	6.5	70.0	4.1	20.6	1.0	0.7	100.2	63.2	97.4
20	7	73.0	6.7	45.8	1.9	4.6	98.4	76.9	96.5
22.5	5.5	56.7	0.4	1.2	0.1	0.2	47.5	1.1	90.6
22.5	6	64.6	1.1	4.8	0.4	0.4	78.2	21.7	98.0
22.5	6.5	68.0	1.8	9.0	0.6	0.4	85.9	27.3	96.6
22.5	7	70.3	3.0	19.4	1.2	0.9	97.3	27.1	96.5
25	5.5	45.6	0.1	0.6	0.0	0.2	8.6	0.8	76.8
25	6	60.2	0.5	1.4	0.3	0.3	59.2	7.6	92.9
25	6.5	66.9	2.2	5.7	1.6	1.7	86.0	18.3	95.6
25	7	67.8	2.9	8.5	2.3	1.7	95.1	18.9	96.5

Table 10. Yield of proteins in precipitation supernatant, expressed as g/L plasma.

EtOH (%)	pH	protein biuret	IgG ELISA	IgA ELISA	IgM ELISA	fibrinogen ELISA	transferrin ELISA	a1-antitrypsin ELISA	albumin neph
20	5.5	26.457	0.061	0.045	0.001	0.000	1.327	0.071	29.402
20	6	29.603	0.157	0.162	0.002	0.001	1.692	0.459	29.174
20	6.5	30.445	0.223	0.261	0.004	0.002	1.799	0.745	30.737
20	7	31.792	0.368	0.579	0.008	0.012	1.765	0.907	30.460
22.5	5.5	24.673	0.021	0.015	0.000	0.000	0.852	0.013	28.591
22.5	6	28.095	0.062	0.061	0.002	0.001	1.403	0.255	30.946
22.5	6.5	29.577	0.097	0.114	0.002	0.001	1.542	0.322	30.488
22.5	7	30.609	0.165	0.246	0.005	0.001	1.747	0.320	30.455
25	5.5	19.833	0.004	0.007	0.000	0.000	0.155	0.009	24.263
25	6	26.212	0.025	0.018	0.001	0.001	1.062	0.089	29.325
25	6.5	29.097	0.119	0.072	0.006	0.004	1.795	0.215	30.198
25	7	29.512	0.157	0.107	0.009	0.004	1.706	0.223	30.458

Example 2 – NG2 C96

[00273] An experimental comparison of IgG purification schemes performed with and without an initial 8% ethanol precipitation was undertaken to determine if further increases in the recovery of IgG could be obtained using a low pH (5.5), high ethanol (25%) precipitation, as outlined in Example 1.

[00274] Briefly, 100 kg of cryo-poor plasma was processed with (NG2C96/1) or without (NG2C96/2) an initial precipitation step (Fraction I) performed using 8% (v/v) final concentration ethanol at pH 7.4 and 0°C. The pH of the cryo-poor plasma (NG2C96/1) or Fraction I supernatant (NG2C96/2) was adjusted to pH 5.5±0.1 and ethanol was admixed to a final concentration of 25% (v/v). The mixtures were then incubated at -5±2°C for 5 hours to allow for protein precipitation. The resulting precipitates (precipitate (I)+II+III) and supernatants (supernatant (I)+II+III) were separated by centrifugation. Due to the size of the precipitation reactions, two centrifugations were required for each reaction. The resulting precipitates from each centrifugation were processed separately.

[00275] The precipitates were suspended in buffer containing 5 mM monosodium phosphate, 5 mM sodium acetate, and 600 mg/L glacial acetic acid at a ratio of 1 part precipitate per 15 parts buffer, the pH was adjusted to 4.8±0.2, and stirred to extract proteins for 2 hours at between 4°C and 8°C. After the extraction incubation, 40 g of fumed silica (SiO₂)/kg precipitate (I)+II+III was admixed into the reactions and stirred for 50 minutes at between 4°C and 8°C. 400 g Cellpure filter aid/kg precipitate (I)+II+III was then admixed and the reactions were filtered through Cuno 50SA membranes to remove insoluble material, forming a (I)+II+III filtrate and a filtercake. The filtercake was washed with buffer containing 5 mM monosodium phosphate, 5 mM sodium acetate, and 150 mg/L glacial acetic acid, the filtrate of which was added to the (I)+II+III filtrate.

[00276] Polysorbate 80 was added to the filtrates to a final concentration of 0.2% (w/w) and the solutions were incubated for 30 minutes. Sodium citrate dihydrate was then added to the solutions to a final concentration of 0.8% (w/w) and the solution was incubated for an additional 30 minutes. After the second incubation, the pH of the solution was adjusted to 7.0 using sodium hydroxide and ethanol was admixed to a final concentration of 25%. The temperature was lowered to between 0°C and -10°C to allow for precipitation. The precipitate (PptG) and supernatant (PptG supernatant) were separated by centrifugation.

[00277] The IgG, IgA, and IgM immunoglobulin contents of each sub-step of the purification reactions were determined. As shown in Table 11, removal of the initial Fraction I precipitation step resulted in a significantly greater yield of IgG in the final composition (compare PptG dissolved NG2C96/2 (without Fraction I precipitation) to NG2C96/1 (with Fraction I

precipitation). Likewise, the IgA (Table 12) and IgM (Table 13) content of the dissolved PptG fraction formed without the initial Fraction I precipitation step.

[00278] Notably, the Fraction I precipitation step is used in part to remove fibrinogen from immunoglobulin-containing compositions. Although the final fibrinogen content of the dissolved PptG fraction formed without the initial Fraction I precipitation step is roughly 6 times greater than the content of the PptG fraction formed with the initial Fraction I precipitation step, the fractionation scheme removes greater than 99% of the fibrinogen content present in the cryo-poor plasma starting material (Table 14).

Table 11. IgG content, expressed as g/L starting plasma, of the various fractions formed during fractionation of plasma using a low pH, high alcohol precipitation reaction with and without an initial 8% ethanol Fraction I precipitation step.

	IgG			
	g/L Plasma			
	NG2C96/1	NG2C96/1 B	NG2C96/2	NG2C96/2 B
	precipitation II+III		precipitation I+II+III	
plasma	5.36		5.36	
supernatant I	5.32		-	
Ppt I dissolved supernatant	0.11		-	
supernatant (I)+II+III	0.004		0.007	
(I)+II+III suspension	4.37	4.72	5.07	4.40
(I)+II+III filtrate	4.55	5.33	5.36	4.77
filtercake dissolved supernatant	0.10	0.18	0.08	0.10
PptG supernatant	0.08	0.06	0.09	0.05
PptG dissolved	4.03	-	5.03	-

Table 12. IgA content, expressed as g/L starting plasma, of the various fractions formed during fractionation of plasma using a low pH, high alcohol precipitation reaction with and without an initial 8% ethanol Fraction I precipitation step.

	IgA			
	g/L Plasma			
	NG2C96/1	NG2C96/1 B	NG2C96/2	NG2C96/2 B
	precipitation II+III		precipitation I+II+III	
plasma	1.20		1.20	
supernatant I	1.14		-	
Ppt I dissolved supernatant	-		-	
supernatant (I)+II+III	0.001		0.002	
(I)+II+III suspension	0.92	1.37	1.08	1.21
(I)+II+III filtrate	0.80	1.02	1.01	0.96
filtercake dissolved supernatant	-	-	-	-
PptG supernatant	-	-	-	-
PptG dissolved	0.66	-	0.79	-

Table 13. IgM content, expressed as g/L starting plasma, of the various fractions formed during fractionation of plasma using a low pH, high alcohol precipitation reaction with and without an initial 8% ethanol Fraction I precipitation step.

	IgM			
	g/L Plasma			
	NG2C96/1	NG2C96/1 B	NG2C96/2	NG2C96/2 B
	precipitation II+III		precipitation I+II+III	
plasma	0.41		0.41	
supernatant I	0.39		-	
Ppt I dissolved supernatant	-		-	
supernatant (I)+II+III	0.0001		0.0004	
(I)+II+III suspension	0.38	0.45	0.40	0.41
(I)+II+III filtrate	0.22	0.24	0.29	0.28
filtercake dissolved supernatant	-	-	-	-
PptG supernatant	-	-	-	-
PptG dissolved	0.19	-	0.29	-

Table 14. fibrinogen content, expressed as g/L starting plasma, of the various fractions formed during fractionation of plasma using a low pH, high alcohol precipitation reaction with and without an initial 8% ethanol Fraction I precipitation step.

	Fibrinogen			
	g/L Plasma			
	NG2C96/1	NG2C96/1 B	NG2C96/2	NG2C96/2 B
	precipitation II+III		precipitation I+II+III	
plasma	1.45		1.45	
supernatant I	0.28		-	
Ppt I dissolved supernatant	-		-	
supernatant (I)+II+III	0.0001		0.0004	
(I)+II+III suspension	0.25	0.34	1.27	1.34
(I)+II+III filtrate	0.002	0.005	0.017	0.018
filtercake dissolved supernatant	-	-	-	-
PptG supernatant	-	-	-	-
PptG dissolved	0.0021	-	0.013	-

Example 3 – NG2 C99

[00279] To further explore the use of a low pH, high alcohol precipitation in place of traditional upstream Cohn-Oncley and Kistler-Nitschmann alcohol fraction steps (*i.e.*, Cohn Fraction I and II; Cohn Fraction I and II+III; or Kistler Nitschmann 8-10% ethanol, precipitate A, and precipitate B), the NG2C96/2 and NG2C96/2B PptG precipitates were combined and further enriched for IgG immunoglobulins.

[00280] Briefly, the combined NG2C96/2 and NG2C96/2B PptG precipitates were suspended in water for injection (WFI) adjusted to pH 4.75±0.75 with acetic acid and a conductivity of 4.5±1.5 mS/cm with sodium chloride. Filter aid was added and the suspension was filtered

through Cuno VR06 membranes to form a clarified PptG filtrate (VR06). The clarified filtrate was then treated with solvents and detergents (S/D treatment) according to standard procedures.

[00281] The S/D treated filtrate was loaded onto a CM sepharose column equilibrated with buffer containing 10 mM sodium acetate (pH 5.0) and the flow through (CM D/N) was collected. The column was then washed with buffer containing 10 mM sodium acetate (pH 5.5) and the wash (CM W) was collected. The immunoglobulins were eluted off of the column using an elution buffer containing 55 mM monosodium phosphate and 10 mM TRIS (pH 8.5). The eluate was collected in three fractions: Fraction 1 (CM E-VL) including the first portion of the eluate having an OD₂₈₀ of less than 100 mAU; Fraction 2 (CM E) including the peak of the eluate having an OD₂₈₀ of greater than 100 mAU on the leading edge of the elution peak and an OD₂₈₀ of greater than 400 mAU on the lagging edge of the elution peak; and Fraction 3 (CM E-NL) including the portion of the eluate on the lagging side of the peak having an OD₂₈₀ of less than 400 mAU.

[00282] The pH and conductivity of the Fraction 2 eluate was adjusted to 6.4 and 2.3 mS/cm using glacial acetic acid and water for injection (WFI), respectively. The Fraction 2 eluate was then loaded onto an ANX Sepharose fast flow anion exchange resin equilibrated with buffer containing 15 mM monosodium phosphate (pH 6.4) at a load concentration of 100 mg protein/mL resin. The anion exchange flow through, containing the majority of the IgG was collected (ANX D/N). Protein binding to the anion exchange resin was eluted off the column using a solution containing 2 M sodium chloride (ANX 2M NaCl).

[00283] Glycine was added to the anion exchange flow through fraction and the sample was ultra- and diafiltered (UF/DF) using a Pellicon[®] 2 Mini (Millipore) against solution containing 0.25 M glycine (pH 5.2) until a protein concentration of 5% was achieved. The immunoglobulin solution was further concentrated to a final protein concentration of 10% and the pH was adjusted to 4.5 (dia-concentrate). The 10% immunoglobulin solution was then sterile filtered using a Millipak 20 (Millipore) to form the final immunoglobulin composition (final container).

[00284] The biochemical profile for each sub-step of the purification reaction was determined (Table 15 and Table 16). As shown in the data, the final composition contained only low levels of non-IgG proteins, within acceptable parameters for pharmaceutical administration.

Table 15. Biochemical characterization of the various fractions formed during fractionation of plasma using a low pH, high alcohol precipitation reaction without an initial 8% ethanol Fraction I precipitation step.

	Protein		IgG	IgA	IgM	α 1 Antitrypsin
	g/L Cohn Pool		g/L Cohn Pool	g/L Cohn Pool	g/L Cohn Pool	g/L Cohn Pool
	Biuret	UV				
Cohn pool	47.38		5.36	1.20	0.41	1.01
supernatant I+II+III	19.79		< 0.0073	0.002	0.0004	0.004
I+II+III suspension	25.83		4.62	1.17	0.41	0.33
I+II+III filtrate	18.24		4.97	0.98	0.28	
filtercake dissolved supernatant	4.52		0.09			0.61
PptG supernatant	11.00		0.06			
PptG dissolved	7.46		5.46	1.01	0.32	0.01
VR 06	7.47		5.20	0.95	0.31	
VR 06 diluted	7.23		5.20	0.99	0.31	
CM D/N	0.66		<0.04 / 0.026			
CM W	0.11	0.09	<0.13 / 0.0066			
CM E-VL	0.001	0.001	<0.0031 / <0.00001			
CM E	6.16	5.38	4.85	0.60	0.042	
CM E-NL	0.08	0.04	0.011 / 0.014			
CM 2M NaCl	0.19	0.11	0.05 / 0.04	0.014	0.02	
CM E diluted		5.33	4.61			
ANX D/N		4.45	4.52	0.0014	0.00007	
ANX 2M NaCl	1.34		0.40 / 0.28	0.59	0.044	
dia-concentrate		4.16	2.55			
post wash 1		0.59				
post wash 2		0.05				
post wash 3		0.01				
dia-concentrate diluted		3.81				
final container		3.86	3.94	0.0012	0.00004	

Table 16. Further biochemical characterization of the various fractions formed during fractionation of plasma using a low pH, high alcohol precipitation reaction without an initial 8% ethanol Fraction I precipitation step.

	α 2 Macroglobulin	Fibrinogen	Transferrin	Albumin	Complement C3
	g/L Cohn Pool	g/L Cohn Pool	g/L Cohn Pool	g/L Cohn Pool	g/L Cohn Pool
Cohn pool		1.45	2.17		0.80
supernatant I+II+III		0.0004	2.44		< 0.001
I+II+III suspension		1.32	1.99	6.40	0.55
I+II+III filtrate		0.02	1.95	6.82	0.05
filtercake dissolved supernatant					
PptG supernatant					
PptG dissolved	0.46	0.02	0.02		0.04
VR 06		0.01	0.02		0.03
VR 06 diluted					
CM D/N	0.17				
CM W					
CM E-VL					
CM E	0.22	0.011	0.024		0.020
CM E-NL					
CM 2M NaCl					
CM E diluted					
ANX D/N	< 0.00002	< 0.00006	0.003		< 0.00004
ANX 2M NaCl					
dia-concentrate					
post wash 1					

post wash 2					
post wash 3					
dia-concentrate diluted					
final container	0.000001	0.00001	0.0028		0.00003

Example 4 – NG2 C100

[00285] To validate the use of a low pH, high alcohol precipitation without an initial low alcohol precipitation step to remove fibrinogen, 200 kg of cryo-poor plasma was fractionated and enriched as described for sample NG2C96/2 in Examples 2 and 3, with a couple of alterations to the process. First, the low pH, high alcohol precipitation step (I-IV1 precipitation) was performed using 25% ethanol at pH 5.1, rather than 5.5. Secondly, the suspended I-IV1 precipitate was treated with 33 g of fumed silica (SiO₂)/kg precipitate I-IV1, rather than 40 g SiO₂/kg precipitate. Thirdly, the CM cation exchange eluate was loaded onto the ANX anion exchange resin at a concentration of 150 mg protein/mL ANX resin, rather than 100 mg/mL ANX resin. Finally, the Cuno VR06 filtered anion exchange flow through was passed through a PLANOVA® 35N (Asahi Kasei Medical Co., Ltd.; Tokyo, Japan) to perform a nanofiltration step prior to ultra-/diafiltration.

[00286] The IgG content of each step of the purification was determined and the results are provided in Table 17. As compared to the average IgG yield for the GAMMAGARD LIQUID® manufacturing process (3.7 g IgG/L source plasma), the purification method used in this example results in a significantly higher yield of 4.4 g IgG/L source plasma. This represents almost a 20% increase in the IgG yield. Biochemical characterization of the final IgG composition is reported in Table 10. As seen in the table, the final IgG composition has greater than 99% purity and the IgG monomer/dimer content is greater than 99.8%. The final container contains only traces of impurities, the levels of which are within regulatory standards.

Table 17. IgG content of the fractions formed during enrichment from cryo-poor plasma using an initial low pH, high alcohol precipitation step.

		IgG				
		mg/ml	g	%	purity (%)	g/l plasma
I-IV-1 precipitation	Cohn pool	5.52	1113.94	100.00	11.43	5.52
	I-IV1 prior EtOH addition					
	I-IV1 after EtOH addition					
	I-IV1 supernatant	< 0.00121	< 0.029	< 0.026	< 0.013	< 0.0014
	I-IV1 supernatant centrifuge 1	< 0.00121	< 0.029	< 0.026	< 0.013	< 0.0014
	I-IV1 supernatant centrifuge 2	< 0.00121	< 0.029	< 0.026	< 0.013	< 0.0014
	I-IV1 precipitate					
I-IV1 extraction a.	I-IV1 precipitate used					
	I-IV1 suspension	2.69	975.30	87.55	13.43	4.83

filtration	CUNO filtrate	1.69	1186.24	106.49	21.41	5.88
	REZ 1	0.09	18.23	1.64	1.78	0.090
filter cake	REZ 2	0.02	2.31	0.21	1.04	0.011
PptG-precipitation	PptG precipitation					
	PptG supernatant	0.02	14.02	1.26	0.35	0.069
	PptG used					
PptG extraction, clarification.	PptG suspension	47.4	1152.39	103.45	69.55	5.71
	VR06 (PptG)	22.8	1074.82	96.49	66.31	5.33
	VR06 diluted	13.80	1126.98	101.17	80.00	5.58
SD and CM sepharose ff	SD					
	Load CM					
	D/N	0.140	18.38	1.65	7.93	0.091
	D/N	0.228	29.84	2.68	12.88	0.15
	W	0.005	1.96	0.18	16.45	0.010
	W	< 0.068	< 27.91	< 2.51	< 234.48	< 0.14
	E-VL	< 0.0712	< 0.59	< 0.05	< 334.63	< 0.00
	E	23.2	1026.43	92.14	86.02	5.09
	E-NL	0.661	4.55	0.41	89.26	0.023
	2M NaCl	0.114	5.82	0.52	43.92	0.029
	2M NaCl	0.139	7.08	0.64	53.46	0.035
	E diluted	9.005	1113.29	99.94	92.64	5.52
ANX SEPH 4FF low sub	Load ANX					
	D/N	6.35	903.11	81.07	94.49	4.48
	2M NaCl	3	78.99	7.09	26.31	0.39
VR06	VR06 (ANX)	6.185	972.42	87.30	101.23	4.82
Asahi	ASAHI filtrate	5.27	907.55	81.47	94.95	4.50
UF/DF	Diaconcentrate	102.5	910.38	81.73	100.89	4.51
	post wash 1	8.255	51.70	4.64	102.78	0.26
	post wash 2					
	post wash 3					
	Diaconcentrate (final)	96.1	871.30	78.22	96.56	4.32
FC	final container (calculated)	100	888.18	79.73		4.40

Table 18. Biochemical characterization of the final IgG composition enriched from 200 L of cryo-poor plasma using an initial low pH, high alcohol precipitation step.

Test	Unit	Result
Total Protein - Kjeldahl	mg/ml	109.8
Total Protein - Kjeldahl	mg/ml	108.5213
Purity (CAE)	Albumin %	0
	α/β globulin %	0
	γ -globulin %	100
	denat. Protein %	0
Molecular size distribution	Monomer [%]	93.2085
	Dimer [%]	6.6806
	Polymer [%]	0.0828
	Fragments [%]	0.0281
IgG neph	mg/ml	100
IgA (ELISA)	mg/ml	0.03
IgM (ELISA)	mg/dl	0
IgE (ELISA)	IU/ml	66.2
Albumin (ELISA)	mg/ml	0.00184
Fibrinogen (ELISA)	μ g/ml	<0.17
Plasminogen (ELISA)	μ g/ml	0.26
C3 complement	mg/dl	<19.4
Test	Subclass group	Result (mg/ml)
IgG-subclass distribution	IgG1	54.7158
	IgG2	29.22
	IgG3	7.2029
	IgG4	2.3634
Test	Unit	Result
IgG Fc Function (FACS)		108 $\pm$ 15
IgG Fc Function Hemaggl.	%	103
Protein identity		positive

Amidolytic activity (PL-1)	nmol/ml	87.3
Anticomple. Activity	%	42
PKA	% of ref lot 3	1.59
PKA	IU/ml	1.2
NAPPT	mg	0.773
Factor Xia	mU/ml	24.9
	pM	771.6
Appearance		satisfactory
ph		5.643
Silicon	µg/l	664
Aluminum	µg/l	<25
Glycine	M	0.074
Osmolality	mosmol/kg	90
Pyrogen	EU/US	sat./sat.
Density	g/cm ³	1.0301
SD Reagents		
Octoxynol 9 (Triton X100)	ppm	0.50
Polysorbate 80	ppm	<26
Tri(n-butyl)phosphate	ppm	<0.2
Test	Unit	Result
Antibodies		
Diphtheria	IU/mL	8.0
Diphtheria	IU/mL	9
Measles	IU/mL	39.63
HBs antigen (ELISA)		negative
HBs (ELISA)	mIU/mL	5775
Poliomyelitis	Quotient gg 176	0.90
HAV (ELISA)	IU/mL	13.8
Parvo B19 (ELISA)	IU/mL	426
Haemophilus Influenzae	Titer	1:6400
Anti D Antibodies		satisfactory
Anti A&AntiB Hemagglutinins	Anti A	16
	Anti B	8

Example 5 – P02910NG2

[00287] To compare the effect of passing the CM cation exchange eluate over an ANX anion exchange column at loading concentrations of 100 g protein/mL ANX resin and 150 g protein/mL ANX resin, 200 L of cryo-poor plasma was fractionated via an initial low pH (5.4), high alcohol (25%) precipitation step and processed to form a PptG precipitate as described for sample NG2C96/2 in Examples 2 and 3. The IgG, IgA, and IgM contents of the upstream fractions formed were determined and are reported in Table 19, Table 20, and Table 21, respectively.

[00288] The resulting PptG precipitate was divided into two 1.8 kg samples, which were enriched to the final container as described for sample NG2C96/2 in Examples 2 and 3, except that one sample was passed over the ANX anion exchange resin at a loading concentration of 100 mg protein/mL ANX resin (P03010NG2) and the other was passed over the ANX anion exchange resin at a loading concentration of 150 mg protein/mL ANX resin (P03110NG2).

[00289] The IgG content of each downstream enrichment step of the purifications was determined and the results are provided in Table 22 and Table 23. As compared to the average IgG yield for the GAMMAGARD LIQUID[®] manufacturing process (3.7 g IgG/L source plasma), the purification methods used in this example results in significantly higher yields of 4.3 g IgG/L source plasma. This represents a 16% increase in the IgG yield. Biochemical characterizations of the final IgG compositions are reported in Table 16 and Table 17. As seen in the tables, the final IgG compositions have greater than 99% purity and the IgG monomer/dimer contents are greater than 99.8%. The final container contains only traces of impurities, the levels of which are within regulatory standards.

Table 19. IgG content of the upstream fractions formed during fractionation of plasma using an initial low pH (5.4), high alcohol (25% ethanol) precipitation reaction.

		weight kg	weight calc. kg	IgG nephelometry IPC / ELISA				
				mg/ml	g	%	purity (%)	g/l plasma
	Cohn Pool	204.60	204.60	5.26	1076.20	100.00	10.72	5.26
I-IV-1 precipitation	I-IV1 prior EtOH addition	206.46	206.81					
	I-IV1 after EtOH addition	266.56	267.14					
	I-IV1 supernatant	247.80	248.64	0.00517	1.29	0.12	0.03	0.01
	I-IV1 precipitate	17.10	17.16					
I-IV1 extraction a. filtration	I-IV1 precipitate used	11.21	17.16					
	I-IV1 suspension	180.21	275.80	4.09	1128.02	104.82	23.15	5.51
	CUNO filtrate	340.70	522.00	2.11	1101.42	102.34	34.72	5.38
PptG- precipitation	PptG precipitation	2.59	3.96					
	PptG supernatant	442.20	677.32	0.0235	15.92	1.48	0.93	0.08

ELISA

Table 20. IgA content of the upstream fractions formed during fractionation of plasma using an initial low pH (5.4), high alcohol (25% ethanol) precipitation reaction.

		weight kg	weight calc. kg	IgA ELISA				
				mg/ml	g	%	purity (%)	g/l plasma
	Cohn Pool	204.60	204.60	1.18	241.43	100.00	2.41	1.18
I-IV1- precipitation	I-IV1 prior EtOH addition	206.46	206.81					
	I-IV1 after EtOH addition	266.56	267.14					
	I-IV1 supernatant	247.80	248.64	0.00251	0.62	0.26	0.01	0.0031
	I-IV1 precipitate	17.10	17.16					
I-IV1 extraction a. filtration	I-IV1 precipitate used	11.21	17.16					
	I-IV1 suspension	180.21	275.80	0.861	237.46	98.36	4.87	1.16
	CUNO filtrate	340.70	522.00	0.365	190.53	78.92	6.01	0.93
PptG- precipitation	PptG precipitation	2.59	3.96					
	PptG supernatant	442.20	677.32					

Table 21. IgM content of the upstream fractions formed during fractionation of plasma using an initial low pH (5.4), high alcohol (25% ethanol) precipitation reaction.

		weight kg	weight calc. kg	IgM ELISA				
				mg/ml	g	%	purity (%)	g/l plasma
	Cohn Pool	204.60	204.60	0.451	92.27	100.00	0.92	0.45

I-IV1-precipitation	I-IV1 prior EtOH addition	206.46	206.81					
	I-IV1 after EtOH addition	266.56	267.14					
	I-IV1 supernatant	247.80	248.64	0.00021	0.052	0.0566	< 0.0011	< 0.00026
	I-IV1 precipitate	17.10	17.16					
I-IV1 extraction a. filtration	I-IV1 precipitate used	11.21	17.16					
	I-IV1 suspension	180.21	275.80	0.353	97.36	105.51	2.00	0.48
	CUNO filtrate	340.70	522.00	0.110	57.42	62.23	1.81	0.28
PptG-precipitation	PptG precipitation	2.59	3.96					
	PptG supernatant	442.20	677.32					

Table 22. IgG content of the downstream fractions formed during fractionation of plasma using an initial low pH (5.4), high alcohol (25% ethanol) precipitation reaction, and an ANX anion exchange loading concentration of 100 mg/mL ANX resin (P03010NG2).

Sample		Weight (kg)	Corr. Weight (kg) corr.	Protein (g)	IgG					Step Yield %
					mg/ml	g	Yield (%)	Purity(%)	g/l Plasma	
PptG Suspension and Filtration										
PptG Susp.	10/2	8.36	8.36	644.59	55.7	465.47	100	72.2	4.99	
PptG Filtrat.	10/6	15.91	16.24	669.20	30	487.34	104.70	72.8	5.22	
PptG Filtrat verd.	10/7	32.50	33.43	658.56	13.7	458.05	98.41	69.6	4.91	100.00
CM Sepharose										
D	12/12 A	38.00	53.88	48.9	0.0937	5.048	1.085	10.325	0.054	1.10
W	12/12 B	127.60	180.91	0.54	< 0.0712					0.00
VL	12/12 C	2.41	3.41	0.04	< 0.0712					0.00
Euate	12/12 D	13.40	19.00	580.49	23.8	452.17	97.14	77.9	4.85	98.72
NL	12/12 E	1.18	1.67	2.18	0.642	1.073	0.23	49.1	0.012	0.23
2M NaCl	12/12 F	15.01	21.28	9.69	0.163	3.47	0.75	35.8	0.037	0.76
Sum							99.20		4.95	100.81
Diluted Euate	12/13	4.3377	37.49	599.85	10.7	401.16	86.18	66.9	4.30	100.00
ANX-Sepharose										
D/N	13/6 A	7.83	53.94	419.63	8.27	446.05	95.83	106.3	4.78	111.19
2M NaCl	13/6 B	2.34	16.12	144.67	2.38	38.37	8.24	26.5	0.41	9.56
Sum							10.407		5.19	120.76
UF / DF										
UF/DF Concentrate	15/14A	0.42	2.91	412.48	134	390.39	83.87	94.6	4.18	87.52
Post-Wash 1	15/14B	0.20	1.39	13.63	10.4	14.41	3.10	105.7	0.15	3.23
Sum										100% UF/DF
Bulk	16/1	0.53	3.97	415.03	101	400.71	86.09	96.55	4.30	102.64
EB	19/1	0.53	3.97	415.03	102	404.68	86.94	97.51	4.34	103.66

Table 23. IgG content of the downstream fractions formed during fractionation of plasma using an initial low pH (5.4), high alcohol (25% ethanol) precipitation reaction, and an ANX anion exchange loading concentration of 150 mg/mL ANX resin (P03110NG2).

Sample		Weight (kg)	Corr. Weight (kg) corr.	Protein (g)	IgG					Step Yield %
					mg/ml	g	Yield (%)	Purity(%)	g/l Plasma	
PptG Suspension and Filtration										
PptG Susp.	10/2	8.36	8.36	644.59	56.7	465.47	100	72.2	4.99	
PptG Filtrat.	10/6	15.91	16.24	669.20	30	487.34	104.70	72.8	5.22	
PptG Filtrat verd.	10/7	32.50	33.43	658.56	13.7	458.05	98.41	69.6	4.91	100.00
CM Sepharose										
D	12/12 A	38.00	53.88	48.9	0.0037	5.05	1.08	10.3	0.054	1.10
W	12/12 B	127.60	180.91	0.54	< 0.0712					0.00

VL	12/12 C	2.41	3.41	0.04	< 0.0712					0.00
Eluate	12/12 D	13.40	19.00	580.49	23.5	452.17	97.14	77.9	4.88	95.72
NL	12/12 E	1.18	1.67	2.18	0.642	1.07	0.23	49.1	0.01	0.23
2M NaCl	12/12 F	15.01	21.28	9.69	0.163	3.47	0.75	35.8	0.04	0.76
Sum							99.20		4.95	100.81
Diluted Eluate	12/13	4.3377	37.49	599.85	10.9	408.65	87.79	68.1	4.38	100.00
ANX-Sepharose										
D/N	13/6 A	11.18	51.03	416.92	8.79	448.56	96.37	107.6	4.81	109.76
2M NaCl	13/6 B	2.37	10.83	144.25	3.11	33.68	7.24	23.3	0.36	8.24
Sum							103.60		5.17	118.01
UF / DF										
UF/DF Concentrate	15/14A	0.57	2.61	425.26	150	391.90	84.19	92.2	4.20	87.37
Post-Wash 1	15/14B	0.55	2.51	13.46	5.87	14.76	3.17	109.7	0.16	3.29
Sum										100% UF/DF
Bulk	16/1	0.84	4.07	445.39	97	394.91	84.84	88.67	4.23	100.77
EB	19/1	0.84	4.07	445.39	99.1	403.45	85.68	90.59	4.32	102.95

Table 24. Biochemical characterization of the final IgG composition enriched from cryo-poor plasma using an initial low pH (5.4), high alcohol (25% ethanol) precipitation reaction, and an ANX anion exchange loading concentration of 100 mg/mL ANX resin (P03010NG2).

Test	Unit	Result
Total Protein - Kjeldahl	mg/ml	109.9996
Purity (CAE)	Albumin %	0
	α/β globulin %	0
	γ -globulin %	100
	cenat. Protein %	0
Molecular size distribution	Monomer [%]	94.8683
	Dimer [%]	4.9579
	Polymer [%]	0.0698
	Fragments [%]	0.104
IgG neph	g/l	91.3
IgA (ELISA)	mg/ml	0.05
IgM (ELISA)	mg/dl	0
Fibrinogen (ELISA)	μ g/ml	<0.17
Plasminogen (ELISA)	μ g/ml	0.26
C3 complement	mg/dl	<19.4
Test	Subclass group	Result [mg/mL]
IgG-subclass distribution	IgG1 (mg/ml)	47.897
	IgG2 (mg/ml)	25.8157
	IgG3 (mg/ml)	6.2317
	IgG4 (mg/ml)	2.0201
Test	Unit	Result
IgG1 Fc Function tion Hemaggl.	%	109
Amidolytic activity (PL-1)	nmol/ml min	< 10
Anticomple. Activity	%	38
PKA	% ref lot 3	1.399
	IU/ml	1
		1.486
		0.957
NAPTT	mg	< 4
Appearance		> 10
pH		Satisfactory
Factor Xla	mIU/mL	4.84
SN13a	mIU/ml	< 0.04
Osmolality	mosmol/kg	< 0.375
Density	g/cm ³	266
Thrombin generation assay		1.0346
Complement factor I	μ g/ml	122.66
Amidolytic activity profil	S-2288	69.1
		< 5

	S-2266	< 5
	S-2222	< 5
	S-2251	< 5
	S-2302	< 5
SD reagents		
Octoxynol 9 (Triton X100)	ppm	<0.1
Polysorbate 80	ppm	<26
Tri(n-butyl)phosphate	ppm	0.1178
Test	Unit	Result
Antibodies		
Anti D Antibodies		Satisfactory
Anti A&AntiB Hemagglutinins	Anti A	16
	Anti B	4

Table 25. Biochemical characterization of the final IgG composition enriched from cryo-poor plasma using an initial low pH (5.4), high alcohol (25% ethanol) precipitation reaction, and an ANX anion exchange loading concentration of 150 mg/mL ANX resin (P03110NG2).

Test	Unit	Result
Total Protein - Kjeldahl	mg/ml	108.3393
Purity (CAE)	Albumin %	0
	α/β globulin %	0
	γ -globulin %	100
	denat. Protein %	0
Molecular size distribution	Monomer [%]	94.9047
	Dimer [%]	4.9122
	Polymer [%]	0.0762
	Fragments [%]	0.1068
IgG neph	g/l	95.7
IgA (ELISA)	mg/ml	0.08
IgM (ELISA)	mg/dl	0
Fibrinogen (ELISA)	μ g/ml	0.33
Plasminogen (ELISA)	μ g/ml	0.27
C3 complement	mg/dl	<19.4
Test	Subclassgroup	Result [mg/ML]
IgG-subclass distribution	IgG1 (mg/ml)	54.0113
	IgG2 (mg/ml)	28.6462
	IgG3 (mg/ml)	6.0688
	IgG4 (mg/ml)	2.3599
Test	Unit	Result
IgG Fc Function Hemaggl.	%	91
Amidolytic activity (PL-1)	nmol/ml min	< 10
Anticomple. Activity	%	36
PKA	% ref lot 3	2.261
	IU/ml	1.8
		1.864
		0.763
		< 4
NAPTT	mg	> 10
Appearance		Satisfactory
pH		4.853
Factor XIa	mU/ml	< 0.04
SN13a	mU/ml	31.3
Osmolality	mosmol/kg	266
Thrombin generation assay		109.81
Density	g/cm ³	1.0344
Complement factor I	μ g/ml	147
Amidolytic activity profil	S-2288	< 5
	S-2266	< 5
	S-2222	< 5
	S-2251	< 5

	S-2302	< 5
SD reagents		
Octoxynol 9 (Triton X100)	ppm	<0.1
Polysorbate 80	ppm	<26
Tri(n-butyl)phosphate	ppm	0.1414
Test	Unit	Result
Antibodies		
Anti D Antibodies		Satisfactory
Anti A&AntiB Hemagglutinins	Anti A	16
	Anti B	4

Example 6 – NG2C107B

[00290] The feasibility of extracting alpha-1-antitrypsin (A1PI) from the insoluble portion of a low pH, high alcohol precipitate used in the manufacture of IgG immunoglobulins was investigated. Briefly, two 500 kg samples (#1 and #3) of the insoluble filtercake formed during filtration of the extracted Fraction I-IV-1 precipitate were first suspended in 2.8 parts water (w/w) at $7\pm 2^{\circ}\text{C}$ and the pH of the samples was adjusted to 5.5 ± 0.1 . The suspensions were filtered through a CUNO 10CP membrane, forming filtrates (1.Filtrat) and second filtercakes. The second filtercakes were then suspended in 5 parts water (w/w), the temperature of the suspensions was adjusted to $17\pm 2^{\circ}\text{C}$, the pH of the samples adjusted to 8.8 ± 0.2 , and the samples were then stirred for 8 hours while maintaining the temperature and pH to extract A1PI from the insoluble material.

[00291] After the incubation, 10 mmol Tris and 150 mmol sodium chloride per kg suspension were added to the samples and the pH was adjusted to 8.0 ± 0.1 . 176 g of PEG 3350 per kg suspension was added to the samples, which were incubated at $17\pm 2^{\circ}\text{C}$ for 1 hour. The suspensions were then filtered to remove the insoluble material from the supernatant (2.Filtrat) containing the extracted A1PI.

[00292] Crude A1PI was then precipitated from the second filtrate by the addition of either 0.35 g zinc chloride/kg filtrate (2.5 mM ZnCl₂; sample #1) or 0.54 g/kg zinc chloride/kg filtrate (4.0 mM ZnCl₂; sample #3), adjusting the pH to 7.5 ± 0.2 , and incubating the samples at $7\pm 2^{\circ}\text{C}$ for 3 hours. The zinc chloride precipitate was recovered by centrifugation (ZnCl₂ cent). A1PI was extracted from the zinc chloride precipitate with 50 mM sodium EDTA (Zn-EDTA-conz).

[00293] The A1PI (a1A) content and activity was determined for each step in the enrichment process and is presented in Table 26. About 400 mg of active A1PI/L source plasma was

recovered in the EDTA extraction step. Notably, when A1PI was extracted at $38 \pm 2^\circ\text{C}$ and a pH of about 9.2, approximately 600 mg of active A1PI/L source plasma was recovered.

[00294] **Table 26.** alpha-1-antitrypsin content of the downstream fractions formed during fractionation of plasma using an initial low pH (5.4), high alcohol (25% ethanol) precipitation reaction, and zinc precipitation.

		weight (g)	weight corr (g)	a1A ELISA				a1A-Activity OC			
				(mg/mL)	mg	mg/L plasma	purity (% of Biuret)	(mg/mL)	mg	mg/L plasma	% activity vs ELISA
1	Filtercake	500.50									
	Cake suspension	1900.40	1900.40								
	Cake susp. cent	32.71	1209.85	0.0964	117	32.32	7.69	0.02	24	6.71	20.7
	1.Filtrate	3291.90	3380.85	<0.0175	<59	<16.40	<5.32	0.01	34	9.37	57.1
	Extr. n.pH	3956.90	4063.82								
	Extr. n.pH cent	42.61	3385.99	0.636	2153	596.84	10.09	0.43	1456	403.52	67.6
	2.Filtrate	7357.80	7653.33	0.295	2258	625.73	15.96	0.2	1531	424.22	67.8
	ZnCl ₂ cent	7487.40	7841.42								
	ZnCl ₂ cent excl. conc	7305.73	7651.16	0.00646	49	13.70	1.13				
	Zn-EDTA-conc	1150.58	1204.98	1.83	2205	611.15	23.41	1.13	1362	377.37	61.7
3	Filtercake	506.5									
	Cake suspension	1926.30	1926.30								
	Cake susp. cent	32.50	1224.18	0.102	125	34.20	7.92	0.02	24	6.71	19.6
	1.Filtrate	3448.10	3539.99	<0.0175	<62	<16.97	<5.40	0.01	35	9.69	57.1
	Extr. n.pH	4175.90	4287.18								
	Extr. n.pH cent	43.03	3611.54	0.589	2127	582.57	9.33	0.42	1517	415.41	71.3
	2.Filtrate	7860.20	8167.45	0.314	2565	702.35	17.97	0.15	1225	335.52	47.8
	ZnCl ₂ cent	8112.00	8483.06								
	ZnCl ₂ cent excl. conc	7913.50	8275.48	0.00319	26	7.23	0.59				
	Zn-EDTA-conz	1485.86	1553.83	1.38	2144	587.25	22.87	0.86	1336	365.96	62.3

Example 7 – P03410NG2A and B

[00295] The effect of pH on the recovery of IgG from a low pH, high alcohol precipitation of cryo poor plasma was investigated. Briefly, 100 L of cryo-poor plasma that had been treated to remove Factor IX, Factor VII, and antithrombin III (ATIII) by adsorption, was fractionated by precipitation using 25% ethanol at a pH of either 5.4 (Sample A; P03410NG2A) or 5.6 (Sample B; P03410NG2B). The samples were then processed to form a Precipitate G precipitate (PptG precipitate) and supernatant (PptG supernatant) as described in Example 2, except that 50 g SiO₂/kg precipitate I-IV-1 was admixed with the precipitate I-IV-1 suspension, rather than 40 g/kg precipitate.

[00296] The IgG, IgA, and IgM contents of the fractions formed were determined and are reported in Table 27, Table 28, and Table 29, respectively. Notably, about 90% of the IgG content of the source plasma is recovered in the Fraction I-IV-1 CUNO filtrate when the initial precipitation is performed at pH 5.4 (93.8%) or pH 5.6 (88.8%). Similarly, nearly all of the IgA and IgM content of the source plasma is recovered in the Fraction I-IV-1 suspension.

Table 27. IgG content of the upstream fractions formed during fractionation of plasma using a low pH (A = 5.4; B = 5.6), high alcohol (25%) initial precipitation step.

			weight kg	weight calc. kg	IgG nephelometry IPC / ELISA				
					mg/ml	g	%	purity (%)	g/l plasma
A		Cohn Pool	101.20	101.20	6.01	608.21	100.00	12.65	6.01
		Cohn Pool	101.82	101.20	5.96	603.15	100.00	12.55	5.96
	I-IV-1 precipitation	I-IV1 prior EtOH addition	101.82	102.28					
		I-IV1 after EtOH addition	131.62	132.21					
		I-IV1 supernatant	123.40	124.23	0.00427	0.53	0.09	0.02	0.01
		I-IV1 precipitate	7.11	7.16					
	I-IV-1 extraction a. filtration	I-IV1 precipitate used	7.07	7.16					
		I-IV1 suspension	113.87	115.33	4.47	515.52	84.76	24.20	5.09
		CUNO filtrate	222.10	225.45	2.53	570.39	93.78	40.77	5.64
	PptG-precipitation	PptG precipitation	2.40	2.44					
		PptG supernatant	288.20	292.81	0.0276	8.08	1.33	1.29	0.08
B		Cohn Pool	100.80	100.80	6.01	605.81	100.00	12.65	6.01
	I-IV-1 precipitation	I-IV1 prior EtOH addition	101.43	101.53					
		I-IV1 after EtOH addition	131.43	131.56					
		I-IV1 supernatant	123.40	123.81	0.0113	1.40	0.23	0.05	0.01
		I-IV1 precipitate	6.52	6.55					
	I-IV-1 extraction a. filtration	I-IV1 precipitate used	6.49	6.55					
		I-IV1 suspension	104.49	105.46	3.80	400.74	66.15	19.67	3.98
		CUNO filtrate	201.60	203.86	2.64	538.19	88.84	46.74	5.34
	PptG-precipitation	PptG precipitation	2.19	2.21					
		PptG supernatant	261.40	264.60	0.0200	5.29	0.87	1.25	0.05

ELISA
IgG neph PPD/PS

Table 28. IgA content of the upstream fractions formed during fractionation of plasma using a low pH (A = 5.4; B = 5.6), high alcohol (25%) initial precipitation step.

			weight kg	weight calc. kg	IgA ELISA				
					mg/ml	g	%	purity (%)	g/l plasma
A		Cohn Pool	101.20	101.20	1.1187	113.21	100.00	2.36	1.12
	I-IV-1 precipitation	I-IV1 prior EtOH addition	101.91	102.37					
		I-IV1 after EtOH addition	131.71	132.30					
		I-IV1 supernatant	123.40	124.23	0.0027	0.34	0.06	0.01	0.003
		I-IV1 precipitate	7.11	7.16					
	I-IV-1 extraction a.	I-IV1 precipitate	7.07	7.16					

	filtration	used							
		I-IV1 suspension	113.87	115.33	1.03	118.79	19.53	5.58	1.17
		CUNO filtrate	222.10	225.45	0.41	93.09	15.31	6.65	0.92
	PptG-precipitation	PptG precipitation	2.40	2.44					
		PptG supernatant	288.20	292.81					

B		Cohn Pool	100.80	100.80	1.1187	112.76	100.00	2.36	1.12
	I-IV-1 precipitation	I-IV1 prior EtOH addition	101.43	101.53					
		I-IV1 after EtOH addition	131.43	131.56					
		I-IV1 supernatant	123.40	123.81	0.0073	0.90	0.15	0.03	0.009
		I-IV1 precipitate	6.52	6.55					
	I-IV-1 extraction a. filtration	I-IV1 precipitate used	6.49	6.55					
		I-IV1 suspension	104.49	105.46	1.03	108.62	17.93	5.33	1.08
		CUNO filtrate	201.60	203.86	0.47	95.00	15.68	8.25	0.94
	PptG-precipitation	PptG precipitation	2.19	2.21					
		PptG supernatant	261.40	264.60					

Table 29. IgM content of the upstream fractions formed during fractionation of plasma using a low pH (A = 5.4; B = 5.6), high alcohol (25%) initial precipitation step.

			weight kg	weight calc. kg	IgM ELISA				
					mg/ml	g	%	purity (%)	g/l plasma
A		Cohn Pool	101.20	101.20	0.4381	44.34	100.00	0.92	0.44
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.91	102.37					
		I-IV1 after EtOH addition	131.71	132.30					
		I-IV1 supernatant	123.40	124.23	0.00014	0.02	0.00	0.00	0.0002
		I-IV1 precipitate	7.11	7.16					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	7.07	7.16					
		I-IV1 suspension	113.87	115.33	0.4042	46.62	7.66	2.19	0.46
		CUNO filtrate	222.10	225.45	0.1506	33.95	5.58	2.43	0.34
	PptG-precipitation	PptG precipitation	2.40	2.44					
		PptG supernatant	288.20	292.81					

B		Cohn Pool	100.80	100.80	0.4381	44.16	100.00	0.92	0.44
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.43	101.53					
		I-IV1 after EtOH addition	131.43	131.56					
		I-IV1 supernatant	123.40	123.81	0.00029	0.04	0.01	0.00	0.0004
		I-IV1 precipitate	6.52	6.55					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	6.49	6.55					
		I-IV1 suspension	104.49	105.46	0.4323	45.59	7.53	2.24	0.45
		CUNO filtrate	201.60	203.86	0.1360	27.73	4.58	2.41	0.28
	PptG-precipitation	PptG precipitation	2.19	2.21					
		PptG supernatant	261.40	264.60					

[00297] To further characterize the fractionation scheme, the contents of fibrinogen (Table 30), A1PI (Table 31), Alpha-2-Macroglobulin (Table 32), albumin (Table 33), Transferrin (Table 34), and C3 (Table 35) were determined for various upstream fractions.

[00298] **Table 30.** fibrinogen content of the upstream fractions formed during fractionation of plasma using a low pH (A = 5.4; B = 5.6), high alcohol (25%) initial precipitation step.

			weight kg	weight calc. kg	Fibrinogen ELISA				
					mg/ml	g	%	purity (%)	g/l plasma
A		Cohn Pool	101.20	101.20	1.698	171.84	100.00	3.57	1.70
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.91	102.37					
		I-IV1 after EtOH addition	131.71	132.30					
		I-IV1 supernatant	123.40	124.23	<0.0001	0.01	0.00	0.00	0.00
		I-IV1 precipitate	7.11	7.16					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	7.07	7.16					
		I-IV1 suspension	113.87	115.33	1.14	131.01	21.54	6.15	1.29
		CUNO filtrate	222.10	225.45	0.03	6.52	1.07	0.47	0.06
	PptG-precipitation	PptG precipitation	2.40	2.44					
		PptG supernatant	288.20	292.81					
B		Cohn Pool	100.80	100.80	1.698	171.16	100.00	3.57	1.70
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.43	101.53					
		I-IV1 after EtOH addition	131.43	131.56					
		I-IV1 supernatant	123.40	123.81	0.00087	0.11	0.02	0.00	0.00
		I-IV1 precipitate	6.52	6.55					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	6.49	6.55					
		I-IV1 suspension	104.49	105.46	1.58	166.52	27.49	8.17	1.65
		CUNO filtrate	201.60	203.86	0.02	4.65	0.77	0.40	0.05
	PptG-precipitation	PptG precipitation	2.19	2.21					
		PptG supernatant	261.40	264.60					

[00299] **Table 31.** alpha-1-antitrypsin content of the upstream fractions formed during fractionation of plasma using a low pH (A = 5.4; B = 5.6), high alcohol (25%) initial precipitation step.

			weight kg	weight calc. kg	alpha 1 antitrypsin ELISA				
					mg/ml	g	%	purity (%)	g/l plasma
A		Cohn Pool	101.20	101.20	1.15	116.38	100.00	2.42	1.15
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.91	102.37					
		I-IV1 after EtOH addition	131.71	132.30					
		I-IV1 supernatant	123.40	124.23	0.0097	1.21	0.20	0.05	0.01
		I-IV1 precipitate	7.11	7.16					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	7.07	7.16					
		I-IV1 suspension	113.87	115.33	0.257	29.64	4.87	1.39	0.29
		CUNO filtrate	222.10	225.45	0.0502	11.32	1.86	0.81	0.11
	PptG-precipitation	PptG precipitation	2.40	2.44					
		PptG supernatant	288.20	292.81	0.0267	7.82	1.29	1.25	0.08
B		Cohn Pool	100.80	100.80	1.15	115.92	100.00	2.42	1.15
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.43	101.53					
		I-IV1 after EtOH	131.43	131.56					

	I-IV1 extraction a. filtration	addition							
		I-IV1 supernatant	123.40	123.81	0.0221	2.74	0.45	0.10	0.03
		I-IV1 precipitate	6.52	6.55					
		I-IV1 precipitate used	6.49	6.55					
		I-IV1 suspension	104.49	105.46	0.365	38.49	6.35	1.89	0.38
		CUNO filtrate	201.60	203.86	0.0597	12.17	2.01	1.06	0.12
	PptG-precipitation	PptG precipitation	2.19	2.21					
		PptG supernatant	261.40	264.60	0.0300	7.94	1.31	1.88	0.08

[00300] **Table 32.** Alpha-2-Macroglobulin content of the upstream fractions formed during fractionation of plasma using a low pH (A = 5.4; B = 5.6), high alcohol (25%) initial precipitation step.

			weight kg	weight calc. kg	alpha 2 macroglobulin ELISA				
					mg/ml	g	%	purity (%)	g/l plasma
A		Cohn Pool	101.20	101.20	1.08	109.30	100.00	2.27	1.08
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.91	102.37					
		I-IV1 after EtOH addition	131.71	132.30					
		I-IV1 supernatant	123.40	124.23	0.00041	0.05	0.01	0.002	0.001
		I-IV1 precipitate	7.11	7.16					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	7.07	7.16					
		I-IV1 suspension	113.87	115.33	0.945	108.99	17.92	5.12	1.08
		CUNO filtrate	222.10	225.45	0.533	120.17	19.76	8.59	1.19
	PptG-precipitation	PptG precipitation	2.40	2.44					
		PptG supernatant	288.20	292.81	0.158	46.26	7.61	7.38	0.46
B		Cohn Pool	100.80	100.80	1.08	108.86	100.00	2.27	1.08
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.43	101.53					
		I-IV1 after EtOH addition	131.43	131.56					
		I-IV1 supernatant	123.40	123.81	0.00105	0.13	0.02	0.005	0.001
		I-IV1 precipitate	6.52	6.55					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	6.49	6.55					
		I-IV1 suspension	104.49	105.46	1.06	111.78	18.45	5.49	1.11
		CUNO filtrate	201.60	203.86	0.559	113.96	18.81	9.90	1.13
	PptG-precipitation	PptG precipitation	2.19	2.21					
		PptG supernatant	261.40	264.60	0.142	37.57	6.20	8.88	0.37

[00301] **Table 33.** albumin content of the upstream fractions formed during fractionation of plasma using a low pH (A = 5.4; B = 5.6), high alcohol (25%) initial precipitation step.

			weight kg	weight calc. kg	Albumin ELISA				
					mg/ml	g	%	purity (%)	g/l plasma
		Cohn Pool	101.20	101.20	25.1	2540.12	100.00	52.84	25.10
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.91	102.37					

A		I-IV1 after EtOH addition	131.71	132.30					
		I-IV1 supernatant	123.40	124.23	18.5	2298.33	377.88	92.24	22.71
		I-IV1 precipitate	7.11	7.16					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	7.07	7.16					
		I-IV1 suspension	113.87	115.33	2.33	268.72	44.18	12.61	2.66
		CUNO filtrate	222.10	225.45	1.19	268.29	44.11	19.18	2.65
	PptG-precipitation	PptG precipitation	2.40	2.44					
		PptG supernatant	288.20	292.81	0.866	253.57	41.69	40.43	2.51
B	I-IV1-precipitation	Cohn Pool	100.80	100.80	25.1	2530.08	100.00	52.84	25.10
		I-IV1 prior EtOH addition	101.43	101.53					
		I-IV1 after EtOH addition	131.43	131.56					
		I-IV1 supernatant	123.40	123.81	19.9	2463.72	406.68	91.43	24.44
		I-IV1 precipitate	6.52	6.55					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	6.49	6.55					
		I-IV1 suspension	104.49	105.46	1.57	165.57	27.33	8.13	1.64
		CUNO filtrate	201.60	203.86	0.693	141.28	23.32	12.27	1.40
	PptG-precipitation	PptG precipitation	2.19	2.21					
		PptG supernatant	261.40	264.60	0.543	143.68	23.72	33.96	1.43

[00302] **Table 34.** Transferrin content of the upstream fractions formed during fractionation of plasma using a low pH (A = 5.4; B = 5.6), high alcohol (25%) initial precipitation step.

			weight kg	weight calc. kg	Transferrin ELISA				
					mg/ml	g	%	purity (%)	g/l plasma
A		Cohn Pool	101.20	101.20	1.95	197.34	100.00	4.11	1.95
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.91	102.37					
		I-IV1 after EtOH addition	131.71	132.30					
		I-IV1 supernatant	123.40	124.23	0.343	42.61	7.01	1.71	0.42
		I-IV1 precipitate	7.11	7.16					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	7.07	7.16					
		I-IV1 suspension	113.87	115.33	1.38	159.16	26.17	7.47	1.57
		CUNO filtrate	222.10	225.45	0.655	147.67	24.28	10.56	1.46
	PptG-precipitation	PptG precipitation	2.40	2.44					
B		PptG supernatant	288.20	292.81	0.541	158.41	26.05	25.26	1.57
		Cohn Pool	100.80	100.80	1.95	196.56	100.00	4.11	1.95
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.43	101.53					
		I-IV1 after EtOH addition	131.43	131.56					
		I-IV1 supernatant	123.40	123.81	0.825	102.14	16.86	3.79	1.01
		I-IV1 precipitate	6.52	6.55					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	6.49	6.55					
		I-IV1 suspension	104.49	105.46	1.13	119.17	19.67	5.85	1.18
		CUNO filtrate	201.60	203.86	0.527	107.44	17.73	9.33	1.07
	PptG-precipitation	PptG precipitation	2.19	2.21					
		PptG supernatant	261.40	264.60	0.383	101.34	16.73	23.95	1.01

[00303] **Table 35.** Complement Component 3 (C3) content of the upstream fractions formed during fractionation of plasma using a low pH (A = 5.4; B = 5.6), high alcohol (25%) initial precipitation step.

			weight kg	weight calc. kg	C3 ELISA				
					mg/ml	g	%	purity (%)	g/l plasma
A		Cohn Pool	101.20	101.20	0.0648	6.56	100.00	0.14	0.06
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.91	102.37					
		I-IV1 after EtOH addition	131.71	132.30					
		I-IV1 supernatant	123.40	124.23	<0.00006	<0.007	<0.001	<0.0003	<0.0001
		I-IV1 precipitate	7.11	7.16					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	7.07	7.16					
		I-IV1 suspension	113.87	115.33	0.673	77.62	12.76	3.64	0.77
		CUNO filtrate	222.10	225.45	0.0476	10.73	1.76	0.77	0.11
	PptG-precipitation	PptG precipitation	2.40	2.44					
		PptG supernatant	288.20	292.81					
B		Cohn Pool	100.80	100.80	0.0648	6.53	100.00	0.14	0.06
	I-IV1-precipitation	I-IV1 prior EtOH addition	101.43	101.53					
		I-IV1 after EtOH addition	131.43	131.56					
		I-IV1 supernatant	123.40	123.81	<0.00006	<0.007	<0.001	<0.0003	<0.0001
		I-IV1 precipitate	6.52	6.55					
	I-IV1 extraction a. filtration	I-IV1 precipitate used	6.49	6.55					
		I-IV1 suspension	104.49	105.46	0.570	60.11	9.92	2.95	0.60
		CUNO filtrate	201.60	203.86	0.0487	9.93	1.64	0.86	0.10
	PptG-precipitation	PptG precipitation	2.19	2.21					
		PptG supernatant	261.40	264.60					

Example 8 – P03510NG2 and P03610NG2

[00304] The effect of passing the CM cation exchange eluate over an ANX anion exchange column at loading concentrations of 100 g protein/mL ANX resin and 150 g protein/mL ANX resin during the downstream processing of a PptG precipitate formed using an initial low pH, high alcohol precipitation step was further investigated. Briefly, 2 kg of P03410NG2A PptG precipitate was processed as described in Example 4, with a first sample (P03510NG2) loaded onto the ANX anion exchange resin at a concentration of 157 mg protein/mL ANX resin and a second sample (P03610NG2) loaded onto the ANX anion exchange resin at a concentration of 106 mg protein/mL ANX resin.

[00305] The IgG contents of each downstream step of the purifications were determined and the results are provided in Table 22 and Table 23. As compared to the average IgG yield for the GAMMAGARD LIQUID[®] manufacturing process (3.7 g IgG/L source plasma), the purification

methods used in this example results in a significantly higher yields of 4.72 g IgG and 4.67 g IgG per L source plasma. This represents greater than a 25% increase in the IgG yield. Biochemical characterization of the final IgG compositions are reported in Table 38 and Table 39. As seen in the tables, the final IgG compositions have greater than 99% purity and the IgG monomer/dimer content is greater than 99.8%. The final container contains only traces of impurities, the levels of which are within regulatory standards.

Table 36. IgG content of the upstream fractions formed during fractionation of plasma using an initial low pH (5.4), high alcohol (25%) initial precipitation step and an anion exchange loading concentration of 157 mg/mL ANX resin (P03510NG2).

Sample		Weight (kg)	Corr. Weight (kg) corr.	Protein (g)	mg/ml	g	IgG			Step Yield %
							Yield (%)	Purity(%)	g/l Plasma	
PptG Suspension and Filtration										
PptG Suspension	10/2	9.63	9.63	683.69	46.4	446.62	100	65.3	5.37	
PptG Filtrate	10/6	18.80	19.15	646.26	25	482.54	108.04	74.7	5.81	
PptG Filtrate verd.	10/7	31.46	32.24	643.35	14.8	477.17	106.84	74.2	5.74	100.00
CM Sepharose										
D	12/12 A	36.40	47.57	73.2	0.188	8.942	2.002	12.212	0.108	1.87
W	12/12 B	109.60	143.22	3.72	< 0.0712					0.00
VL	12/12 C	2.41	3.15	0.00	< 0.0712					0.00
Eluate	12/12 D	13.40	17.51	532.56	23.1	404.49	90.57	76.0	4.87	84.77
NL	12/12 E	1.41	1.84	2.36	0.544	0.999	0.22	42.3	0.012	0.21
2M NaCl	12/12 F	14.81	19.35	10.44	0.196	3.79	0.85	36.3	0.046	0.79
Sum							93.64		5.03	87.65
Diluted Eluate	12/13	4.3377	34.55	535.67	11.6	400.83	89.75	74.8	4.82	100.00
ANX-Sepharose										
D/N	13/6 A	15.60	41.90	373.70	9.73	407.64	91.27	109.1	4.91	101.70
2M NaCl	13/6 B	3.76	10.10	120.56	3.32	33.54	7.51	27.8	0.40	8.37
Sum							98.78		5.31	110.07
VR06 Filtrate	14/2	17.15	46.30	373.66						
Asahi 35 nm Filtrate	14/5	18.82	50.84	371.10	8.19	416.35	93.22	112.2	5.01	103.87
UF / DF										
UF/DF Concentrate	15/14 A	1.02	2.74	391.05	133	364.39	81.59	93.2	4.39	89.39
Post-Wash 1	15/14 B	0.74	2.00	6.94	3.53	7.08	1.58	102.0	0.09	1.74
Sum										100% UF/DF
Bulk	16/1	1.38	3.85	375.89	96.5	371.96	83.28	98.95	4.48	102.08
EB	19/1	1.21	3.81	378.80	103	392.60	87.90	103.64	4.72	107.74

Table 37. IgG content of the upstream fractions formed during fractionation of plasma using an initial low pH (5.4), high alcohol (25%) initial precipitation step and an anion exchange loading concentration of 106 mg/mL ANX resin (P03610NG2).

Sample		Weight (kg)	Corr. Weight (kg) corr.	Protein (g)	mg/ml	g	IgG			Step Yield %
							Yield (%)	Purity(%)	g/l Plasma	
PptG Suspension and Filtration										
PptG Suspension	10/2	9.63	9.63	683.69	46.4	446.62	100	65.3	5.37	
PptG Filtrate	10/6	18.80	19.15	646.26	25	482.54	108.04	74.7	5.81	
PptG Filtrate verd.	10/7	31.46	32.24	643.35	14.8	477.17	106.84	74.2	5.74	100.00
CM Sepharose										
D	12/12 A	36.40	47.57	73.2	0.188	8.942	2.002	12.212	0.108	1.87
W	12/12 B	109.60	143.22	3.72	< 0.0712					0.00
VL	12/12 C	2.41	3.15	0.00	< 0.0712					0.00
Euate	12/12 D	13.40	17.51	532.56	23.1	404.49	90.57	76.0	4.87	84.77
NL	12/12 E	1.41	1.84	2.36	0.544	0.999	0.22	42.3	0.012	0.21
2M NaCl	12/12 F	14.81	19.35	10.44	0.196	3.79	0.85	36.3	0.046	0.79
Sum							93.64		5.03	87.65
Diluted Euate	12/13	4.3377	34.55	535.67	11.6	400.83	89.75	74.8	4.82	100.00
ANX-Sepharose										
D/N	13/6 A	11.80	46.90	384.12	8.46	396.78	88.84	103.3	4.77	98.99
2M NaCl	13/6 B	3.72	14.79	123.43	2.23	32.98	7.38	26.7	0.40	8.23
Sum							96.23		5.17	107.22
VR06 Filtrate	14/2	12.88	51.53	381.83						
Asahi 35 nm Filtrate	14/5	14.10	56.47	378.34	6.93	391.33	87.62	103.4	4.71	97.63
UF/DF										
UF/DF Concentrate	15/14A	0.69	2.77	385.58	125	346.49	77.58	89.9	4.17	87.33
Post-Wash 1	15/14B	0.78	3.12	9.96	3.4	10.62	2.38	106.6	0.13	2.68
Sum										100% UF/DF
Bulk	16/1	0.89	3.73	376.90	93.8	350.10	78.39	92.89	4.21	101.04
EB	19/1	0.89	3.73	376.90	104	388.17	86.91	102.99	4.67	112.03

Table 38. Biochemical characterization of the final IgG composition enriched from cryo-poor plasma using an initial low pH (5.4), high alcohol (25%) initial precipitation step and an anion exchange loading concentration of 157 mg/mL ANX resin (P03510NG2).

Test	Unit	Result
Total Protein - Kjeldahl	mg/ml	100.5061
Purity (CAE)	Albumin %	0
	α/β globulin %	0
	γ -globulin %	100
	denat. Protein %	0
Molecular size distribution	Monomer [%]	93.1042
	Dimer [%]	6.7664
	Polymer [%]	0.0499
	Fragments [%]	0.0795
IgG neph	g/l	92.4
IgA (ELISA)	mg/ml	0.07
IgM (ELISA)	mg/dl	< 0.95
Fibrinogen (ELISA)	μ g/ml	< 0.17
Plasminogen (ELISA)	μ g/ml	0.32
C3 complement	mg/dl	< 19.4
Test	Subclassgroup	Result [mg/mL]
IgG-subclass distribution	IgG1 (mg/ml)	54.4779
	IgG2 (mg/ml)	33.6428
	IgG3 (mg/ml)	6.3377
	IgG4 (mg/ml)	1.9745
Test	Unit	Result
IgG Fc Function Hemaggl.	%	84
Amidolytic activity (PL-1)	nmol/ml min	14.8

		1.37
Anticomp. Activity	%	35
	% ref lot 3	0.741
PKA	IU/ml	0.6
		1.700
		2.036
		< 4
NAPTT	mg	> 10
Appearance		Satisfactory
pH		4.7
Factor Xla	mIU/mL	0.15
SN13a	mU/ml	21.9
Thrombin generation assay		142.23
Osmolality	mosmol/kg	259
Density	g/cm ³	1.0326
Complement factor I	μg/ml	107
Amidolytic activity profil	S-2288	26.9
	S-2266	18.8
	S-2222	< 5
	S-2251	< 5
	S-2302	55.2
SD reagents		
Octoxynol 9 (Triton X100)	ppm	< 0.1
Polysorbate 80	ppm	< 26
Tri(n-butyl)phosphate	ppm	< 0.2
Test	Unit	Result
Antibodies		
Anti D Antibodies		Satisfactory
Anti A&AntiB Hemagglutinins	Anti A	8
	Anti B	8

Table 39. Biochemical characterization of the final IgG composition enriched from cryo-poor plasma using an initial low pH (5.4), high alcohol (25%) initial precipitation step and an anion exchange loading concentration of 106 mg/mL ANX resin (P03610NG2).

Test	Unit	Result
Total Protein - Kjeldahl	mg/ml	103.5231
Purity (CAE)	Albumin %	0
	α/β globulin %	0
	γ-globulin %	100
	denat. Protein %	0
Molecular size distribution	Monomer [%]	92.9679
	Dimer [%]	6.9018
	Polymer [%]	0.0526
	Fragments [%]	0.0778
IgG neph	g/l	95
IgA (ELISA)	mg/ml	0.07
IgM (ELISA)	mg/dl	< 0.95
Fibrinogen (ELISA)	μg/ml	0.20
Plasminogen (ELISA)	μg/ml	0.29
C3 complement	mg/dl	< 19.4
Test	Subclassgroup	Result [mg/mL]
IgG-subclass distribution	IgG1 (mg/ml)	59.3788
	IgG2 (mg/ml)	34.8724
	IgG3 (mg/ml)	6.7758
	IgG4 (mg/ml)	2.1446
Test	Unit	Result
IgG Fc Function Hemaggl.	%	101
Amidolytic activity (PL-1)	nmol/ml min	< 10
		0.951
Anticomp. Activity	%	39
PKA	% ref lot 3	< 1

	IU/ml	0
		< 4
NAPTT	mg	> 10
Appearance		Satisfactory
pH		4.7
Factor Xla	mlU/mL	0.06
SN13a	mU/ml	14.9
Thrombin generation assay		123.96
Osmolality	mosmol/kg	260
Density	g/cm ³	1.0333
Complement factor I	µg/ml	80.0
Amidolytic activity profil	S-2288	20.3
	S-2266	17.2
	S-2222	< 5
	S-2251	< 5
	S-2302	38.9
SD reagents		
Octoxynol 9 (Triton X100)	ppm	< 0.1
Polysorbate 80	ppm	< 26
Tri(n-butyl)phosphate	ppm	< 0.2
Test	Unit	Result
Antibodies		
Anti D Antibodies		Satisfactory
Anti A&AntiB Hemagglutinins	Anti A	16
	Anti B	8

Example 9 – Additional Characterization of Fractionation Products

[00306] The immunoglobulin G yield and purity achieved using purification methods that include an initial low pH, high alcohol precipitation were compared to those achieved using traditional Cohn precipitation I and precipitation II+III intermediates.

[00307] The yield and purity of three large scale IgG purifications employing an initial low pH, high alcohol precipitation - lot 3010, as described in Example 5 (“P03010NG2”); lot 3610 as described in Example 8 (“P03610NG2”); and a third lot 1111, prepared similarly to lots 3010 and 3610, was compared to the average yield for GAMMAGARD LIQUID (10% Immune Globulin Infusion (Human), Baxter International) prepared at a single manufacturing plant in 2010. The GAMMAGARD LIQUID purification process was substantially similar to the processes used to prepare lots 3010, 3610, and 1111, except that the GAMMAGARD LIQUID process proceeded through Cohn Fraction I and Cohn Fraction II+III intermediates, rather than an initial low pH, high alcohol precipitation. Characterization of the precipitate G products formed during the preparation of the experimental lots and averaged for GAMMAGARD LIQUID is shown in Table 40.

Table 40. Biochemical characterization of the precipitate G (Ppt G) intermediate formed during preparation of experimental IgG lots 3010, 3610, and 1111, proceeding through an initial low

pH, high alcohol precipitation step, and averaged for GAMMAGARD LIQUID preparation as a single manufacturing plant in the year 2010, proceeding through Cohn Fraction I and II+III precipitation steps.

Lot Number		3010	3610	1111	GAMMAGARD LIQUID
IgG in Cohn Pool	g/L plasma (PPD/PS-test)	5.26	6.01	6.91	5.82 (Avg)
Characterization of Precipitate G					
IgG	g/L plasma (PPD/PS-test)	4.99	5.81	6.14	82.9 (Avg)
IgG	% of Cohn Pool	94.9	96.6	88.9	82.9
C3	% of protein	0.4	1.8	1.5	0.1 - 0.6 (mean = 0.2)
Fibrinogen	% of protein	0.12	0.68	1.02	0.01 - 1.0 (mean = 0.13)

[00308] As seen in Table 40, use of an initial low pH, high alcohol precipitation step resulted in a significant increase in the IgG recovery at the precipitate G intermediate step as compared to the GAMMAGARD LIQUID purification method proceeding through Cohn Fraction I and II+III precipitation steps (88.9-96.6% IgG recovery for initial high alcohol, low pH precipitation step vs. 82.9% for Cohn Fraction I and II+III precipitation steps).

[00309] The IgG precipitate G intermediates formed in experimental lots 3010, 3610, and 1111 have slightly lower purity than does the average GAMMAGARD LIQUID precipitate G intermediate, as evidenced by the higher levels of complement component 3 (C3) and fibrinogen, although the fibrinogen content of these intermediates is still within the range of values seen for the GAMMAGARD LIQUID manufacturing. However, when the experimental lots were further processed as described above, the final IgG product met all manufacturing specifications tested, as shown in Table 41.

Table 41. Biochemical characterization of final pooled human IgG compositions for experimental lots 3010, 3610, and 1111, proceeding through an initial low pH, high alcohol precipitation step.

Lot number	Lot	3010	3610	1111
Protein	g/L plasma	4.45	4.54	4.82
IgA	%	0.05	0.07	0.04
IgM	mg/dL	<1.6	<1.6	<1.6
Fibrinogen	μg/mL	<0.17	0.2	<0.17
C3	mg/mL	<0.194	<0.194	<0.194
Fc Function	%	109	101	78

HP-SEC	% monomers and dimers	99.8	99.4	99.8
CAE	% g-globulin	100	100	100
ACA	%	38	39	46
PKA	IU/mL	1	0	1
PL-1	nmol/ml min (QC)	<10	<10	<10
NAPIT	mg	>10	>10	>10
TGA	% of normal plasma control @5%	123	124	112
Specific F-XIa	mU/mL	<0.04	0.06	<0.2

Example 10 – Characterization of Albumin Compositions

[00310] To further validate that the new immunoglobulin G manufacturing processes described herein (e.g., those proceeding through an initial low pH, high alcohol precipitation step) could support the manufacture of plasma protein by-products, albumin was purified from a low pH, high alcohol supernatant.

[00311] Methods for purifying albumin from Cohn Fraction IV-1 and IV-4 supernatants are well known in the art. Typically, the Cohn Fraction IV-1 and IV-4 supernatants are formed in a process that proceeds through intermediate alcohol precipitations steps, such as Cohn Fraction I and Cohn Fraction II+III precipitations. In the present Example, albumin was prepared according to standard methods, except that the supernatant from an initial low pH, high alcohol precipitation was used in place of a Fraction IV-1 supernatant. As shown in **Table 42**, the purification from a low pH, high alcohol supernatant resulted in a high yield of albumin (18.92 g/L plasma), which met all quality control requirements tested.

Table 42. Biochemical characterization of an albumin composition prepared from a large-scale low pH, high alcohol precipitation supernatant, as described above.

Test	Unit	AUF00111NG2 Final container
Protein yield	g/L plasma	18.92
Albumin (CAE)	(%)	97.3
	(%) after 2 weeks at 30°C	97.8

HPLC	(% monomers)	98.1
Citrate	$\mu\text{mol/mL}$	<0.05
PKA	IE/mL	<4
Visual inspection	Vials rejected (e.g., due to particles)	None

WHAT IS CLAIMED IS:

1. A method for manufacturing an enriched immunoglobulin composition from a Cohn pool, the method comprising the steps of:

(a) co-precipitating immunoglobulins and alpha-1-antitrypsin (A1PI) from a Cohn pool, in a first precipitation step by the addition of ethanol to the Cohn pool at a final concentration of from 20% to 30% (v/v) at a pH of from 5.0 to 6.0, to form a first precipitate and a first supernatant;

(b) solubilizing immunoglobulins in the first precipitate, to form a first suspension having a soluble portion comprising immunoglobulins and an insoluble portion comprising A1PI;

(c) separating the soluble portion of the first suspension from the insoluble portion of the first suspension; and

(d) recovering the soluble fraction of the first suspension, thereby forming an enriched immunoglobulin composition, wherein no alcohol precipitation is performed prior to step (a).

2. The method of claim 1, wherein the final concentration of ethanol is $25 \pm 2\%$.
3. The method of claim 1, wherein the final concentration of ethanol is 25%.
4. The method according to any one of claims 1 to 3, wherein the pH is 5.5 ± 0.2 .
5. The method according to any one of claims 1 to 3, wherein the pH is 5.5.

6. The method according to any one of claims 1 to 5, wherein the first precipitation step is performed at a temperature of from -5°C to -9°C .

7. The method according to any one of claims 1 to 6, wherein the first suspension has a pH of from 4.0 to 5.4.

8. The method according to any one of claims 1 to 6, wherein the first suspension has a pH of from 4.7 to 5.1.

9. The method according to any one of claims 1 to 8, wherein the first suspension has a conductivity of from 0 mS/cm to 4 mS/cm.

10. The method according to any one of claims 1 to 9, wherein the first suspension has a conductivity of from 0.5 mS/cm to 2 mS/cm.

11. The method according to any one of claims 1 to 10, wherein the step of separating the soluble portion of the first suspension from the insoluble portion of the first suspension comprises:

- (i) mixing finely divided silicon dioxide (SiO_2) with the first suspension; and
- (ii) separating the SiO_2 from the suspension

, wherein the finely divided silicon dioxide (SiO_2) has an average surface area of from $350 \text{ m}^2/\text{g}$ to $410 \text{ m}^2/\text{g}$.

12. The method according to any one of claims 1 to 11, wherein the finely divided silicon dioxide (SiO_2) is added to the first suspension at a final concentration of from 15 g/kg first precipitate to 80 g/kg first precipitate.

13. The method according to any one of claims 1 to 12, wherein the method comprises the steps of:

(a) precipitating immunoglobulins from a Cohn pool, in a first precipitation step, with between 24% and 26% alcohol at a pH of between 5.3 and 5.7 and temperature of between -6°C and -8°C to form a first precipitate and a first supernatant;

(b) suspending the first precipitate to form a first suspension;

(c) treating the first suspension with finely divided silicon dioxide (SiO_2);

(d) separating the soluble fraction of the first suspension from the insoluble fraction of the first suspension; and

(e) recovering the soluble fraction of the first suspension, thereby forming an enriched immunoglobulin composition.

14. The method according to any one of claims 1 to 13, wherein the method further comprises the steps of

(f) precipitating immunoglobulins from the soluble fraction of the first suspension, in a second precipitation step, to obtain a second precipitate and a second supernatant;

(g) suspending the second precipitate to form a second suspension; and

(h) recovering the soluble fraction of the second suspension, thereby forming a further enriched immunoglobulin composition

, wherein the second precipitation step is an alcohol precipitation step, comprising adding ethanol to the soluble fraction of the first suspension to achieve a final concentration of between 22% and 28% ethanol at a pH between 6.5 and 7.5.

15. The method according to any one of claims 1 to 14, wherein the method further comprises a cation exchange chromatography enrichment step and/or an anion exchange chromatography enrichment step.

16. The method according to any one of claims 1 to 15, wherein the method comprises at least one of: (i) a solvent/detergent (S/D) viral inactivation step;

(ii) a nanofiltration step; and

(iii) a step of incubating the composition at a pH from 4.0 to 5.0 and temperature from 20°C to 40°C for at least one week.

17. The method according to any one of claims 1 to 16, wherein the method yields at least 4 g of IgG per L of Cohn pool used in step (a).

18. The method of claim 17, wherein the method yields at least 4.5 g of IgG per L of Cohn pool used in step (a).

19. The method according to any one of claims 1 to 18, wherein alpha-1-antitrypsin (A1PI) is further purified from an insoluble fraction of the first suspension.

20. The method according to any one of claims 1 to 18, wherein Factor H is further purified from an insoluble fraction of the first suspension.

21. The method according to any one of claims 1 to 18, wherein an Inter-alpha-Trypsin Inhibitor protein (IaIp) is further purified from an insoluble fraction of the first suspension.

22. The method according to any one of claims 1 to 21, wherein the Cohn pool is cryo-poor plasma.