

HETEROCYCLIC MODULATORS OF LIPID SYNTHESIS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit under 35 U.S.C. §119(e) of U.S. Provisional Patent Application No. 61/667,894, filed July 3, 2012, U.S. Provisional Patent Application No. 61/698,511, filed September 7, 2012, U.S. Provisional Patent Application No. 61/699,819, filed September 11, 2012, and U.S. Provisional Patent Application No. 61/785,933, filed March 14, 2103. The foregoing applications are incorporated herein by reference in their entireties.

FIELD

The present disclosure relates generally to heterocyclic modulators of lipid synthesis and methods of use thereof. The present heterocyclic modulators of lipid synthesis can be used for the treatment of disorders characterized by dysregulation in the fatty acid synthase function in a subject by modulating the fatty acid synthase pathway and/or the fatty acid synthase function.

BACKGROUND

Viral disease is a significant health concern that threatens large segments of human populations. Some of the features related to viral infection which are of concern to health care professionals include its highly contagious nature (e.g., HIV, SARS, etc.) and high mutability. Some viruses are also oncogenic (such as HPV, EBV and HBV). While viruses are structurally amongst the simplest of organisms, they are regarded to be among the most difficult to control and present a formidable challenge for antiviral drug R&D.

Thus far, there have been a few antiviral drugs widely used in patients, such as Amantadine and Oseltamivir for influenza, Acyclovir for HSV-related infections, Ganciclovir for CMV infection, and multiple agents including co-formulated drugs (Efavirenz, emtricitabine, and tenofovir disoproxil fumarate) for AIDS treatments. These drugs possess a variety of undesirable neurological, metabolic and immunological side-effects. Therefore, development of new antiviral therapy has become a major focus of medical and pharmaceutical research and development.

Infection by hepatitis C virus (HCV) is a serious health issue. It is estimated that 170 million people worldwide are chronically infected with HCV. HCV infection can lead to

chronic hepatitis, cirrhosis, liver failure and hepatocellular carcinoma. Chronic HCV infection is thus a major worldwide cause of liver-related premature mortality.


The present standard of care treatment regimen for HCV infection involves combination therapy with interferon-alpha and ribavirin, often with the addition of a direct-acting protease inhibitor (Telaprevir or Boceprevir). The treatment is cumbersome and sometimes has debilitating and severe side effects. For this reason, many patients are not treated in early stages of the disease. Additionally, some patient populations do not durably respond to treatment. New and effective methods of treating HCV infection are urgently needed.

The dominant therapeutic approaches that are currently employed to treat cancer include surgical removal of primary tumors, tumor irradiation, and parenteral application of anti-mitotic cytotoxic agents. Unfortunately, only a relatively small cross-section of cancer patients have tumors that are "addicted" to a specific pathway, and can therefore be treated with newer targeted agents. The continued dominance of these long established therapies is mirrored by the lack of improvement in survival rates for most cancers. In addition to limited clinical success, devastating side effects accompany classic therapies. Both radiation- and cytotoxic-based therapies result in the destruction of rapidly dividing hematopoietic and intestinal epithelial cells leading to compromised immune function, anemia, and impaired nutrient absorption. Surgical intervention often results in a release of tumor cells into the circulation or lymph systems from which metastatic tumors can subsequently be established. Improved methods for the treatment of cancer are needed.

SUMMARY

The present disclosure addresses the deficiencies for antiviral and anticancer treatments by providing novel heterocyclic modulators of lipid synthesis having improved antiviral and anticancer activities.

In various aspects, the present disclosure provides for compounds of Structure I:






I, or pharmaceutically acceptable salts thereof, wherein:

L^3 is $-CH_2-$, $-CHR^{50}-$, $-O-$, $-NR^{50}-$, $-NC(O)R^{50}-$ or $-NC(O)OR^{50}-$, wherein R^{50} is C_1-C_6 alkyl, C_3-C_5 cycloalkyl, or 4- to 6-membered heterocycle;

n is 1, 2, or 3;

m is 1 or 2 with the proviso that $n+m \geq 3$;



Het is a 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered heterocycle) or -O-(C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is

10 optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;


R^3 is H or F;

R^{11} is H or -CH₃;


R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or a 4- to 6-membered heterocycle; and


15 R^{22} is H, halogen, or C_1 - C_2 alkyl.

In various aspects, the present disclosure provides for compounds of Structure II:



II, or pharmaceutically acceptable salts thereof, wherein:







Het is a 5- to 6-membered heteroaryl;


- 5 R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered heterocycle) or -O-(C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;
- each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;
- R^3 is H or F;
- 10 R^{11} is H or -CH₃;
- R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle;
- R^{22} is H, halogen, or C_1 - C_2 alkyl; and
- R^{24} is H, C_1 - C_4 alkyl, -(C_1 - C_4 alkyl)-OH, -(C_1 - C_4 alkyl)_t-N(R^{241})₂, -(C_1 - C_4 alkyl)_t-O-(C_3 - C_5 cycloalkyl), -(C_1 - C_4 alkyl)_t-O-(4- to 6-membered heterocycle) or -(C_1 - C_4 alkyl)_t-O-(C_1 - C_4 alkyl), wherein:
- 15 each t is independently 0 or 1; and
- each R^{241} is independently H or C_1 - C_2 alkyl.

In various aspects, the present disclosure provides for compounds of Structure III:



III, or pharmaceutically acceptable salts thereof, wherein:





Het is a 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered

5 heterocycle) or -O-(C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;

R^3 is H or F;

R^{11} is H or - CH_3 ;

10 R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle;

R^{22} is H, halogen, or C_1 - C_2 alkyl;

R^{24} is H, -CN, -(C_1 - C_4 alkyl)-CN, C_1 - C_4 alkyl, -(C_1 - C_4 alkyl)-OH, -(C_1 - C_4 alkyl)- $N(R^{241})_2$, -
(C_1 - C_4 alkyl) $_t$ -O $_u$ -(C_3 - C_6 cycloalkyl), -(C_1 - C_4 alkyl) $_t$ -O $_u$ -(4- to 6-membered
heterocycle) or -(C_1 - C_4 alkyl)-O-(C_1 - C_4 alkyl), wherein:

15 t is 0 or 1;


u is 0 or 1;

with the proviso that when u is 1, t is 1; and


each R^{241} is independently H or C_1 - C_2 alkyl; and


R^{25} is halogen, -CN, -(C_1 - C_4 alkyl)-CN, C_1 - C_2 alkyl or cyclopropyl.

20 In various aspects, the present disclosure provides for compounds of Structure IIIb:



IIIb, or pharmaceutically acceptable salts thereof, wherein:





Het is a 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered

5 heterocycle) or -O-(C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;

R^3 is H or F;

R^{11} is H or -CH₃;

10 R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle;

R^{22} is H, halogen or C_1 - C_2 alkyl; and

each R^{24} and R^{25} is independently H, halogen, -CN, -(C_1 - C_4 alkyl)-CN, C_1 - C_4 alkyl, -(C_1 - C_4 alkyl)-OH, -(C_1 - C_4 alkyl)-N(R^{241})₂, -(C_1 - C_4 alkyl)_t-O_u-(C_3 - C_5 cycloalkyl), -(C_1 - C_4 alkyl)_t-O_u-(4- to 6-membered heterocycle) or -(C_1 - C_4 alkyl)_t-O-(C_1 - C_4 alkyl),


15 wherein:

each t is independently 0 or 1;

each u is independently 0 or 1; and


each R^{241} is independently H or C_1 - C_2 alkyl,

wherein the compound is not:




20

In various aspects, the present disclosure provides for compounds of Structure IIIc:





IIIc, or pharmaceutically acceptable salts thereof, wherein:




Het is a 5- to 6-membered heteroaryl;

- 5 R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered heterocycle) or -O-(C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;
- each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;
- R^3 is H or F;
- 10 R^{11} is H or - CH_3 ;
- R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle;
- R^{22} is H, halogen or C_1 - C_2 alkyl; and
- each of R^{24} and R^{25} is independently H, - C_1 - C_4 alkyl, or halogen.

In various aspects, the present disclosure provides for compounds of Structure IV:





Het is a 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl),

-O-(4- to 6-membered heterocycle) or -O-(C_1 - C_4 alkyl), wherein when R^1 is not H,

5 -CN or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;

R^3 is H or F;

R^{11} is H or -CH₃;

R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle;

10 R^{22} is H, halogen or C_1 - C_2 alkyl; and

R^{24} is H, C_1 - C_4 alkyl, -(C_1 - C_4 alkyl)-OH, -(C_1 - C_4 alkyl)-N(R^{241})₂, -(C_1 - C_4 alkyl)_t-O_u-(C_3 - C_5 cycloalkyl), -(C_1 - C_4 alkyl)_t-O_u-(4- to 6-membered heterocycle) or -(C_1 - C_4 alkyl)-O-(C_1 - C_4 alkyl), wherein:


t is 0 or 1;

15 u is 0 or 1;


with the proviso that when u is 1, t is 1; and


R^{241} is H or C_1 - C_2 alkyl.

In various aspects, the present disclosure provides for compounds of Structure V:



V, or pharmaceutically acceptable salts thereof, wherein:





L² is -NHR³⁵ or -C(O)NHR³⁵¹, wherein R³⁵¹ is C₁-C₆ alkyl, C₃-C₅ cycloalkyl,

4- to 6- membered heterocycle, aryl or heteroaryl;

Het is a 5- to 6-membered heteroaryl;

- 5 R¹ is H, -CN, halogen, C₁-C₄ alkyl, -O-(C₃-C₅ cycloalkyl),
-O-(4- to 6-membered heterocycle), -O-(C₁-C₄ alkyl) wherein when R¹ is not H, -CN
or halogen, R¹ is optionally substituted with one or more halogens;

each R² is independently hydrogen, halogen or C₁-C₄ alkyl;

R³ is H or F;


- 10 R¹¹ is H or -CH₃;

R²¹ is H, halogen, C₁-C₄ alkyl, C₃-C₅ cycloalkyl or 4- to 6-membered heterocycle;

R²² is H, halogen, or C₁-C₂ alkyl; and

R³⁵ is -C(O)R³⁵¹, -C(O)NHR³⁵¹, C(O)OR³⁵¹ or S(O)₂R³⁵¹ wherein R³⁵¹ is C₁-C₆ alkyl, C₃-
C₅ cycloalkyl, 4- to 6- membered heterocycle, aryl or heteroaryl.

- 15 In various aspects, the present disclosure provides for compounds of Structure VI:




VI, or pharmaceutically acceptable salts thereof,

wherein:

each W, X, Y and Z is independently -N- or -CR²⁶- with the proviso that not more than 2
of W, X, Y and Z are -N-;

- 20 each R²⁶ is independently H, C₁-C₄ alkyl, -O-(C₁-C₄ alkyl), -N(R²⁷)₂, -S(O)₂-(C₁-C₄
alkyl), or -C(O)-(C₁-C₄ alkyl);

each R²⁷ is independently H or C₁-C₄ alkyl or both R²⁷ are C₁-C₄ alkyl and join to form a
3- to 6-membered ring together with the N to which they are attached and wherein the
ring optionally includes one oxygen atom as one of the members of the ring;



Het is a 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered heterocycle), -O-(C_1 - C_4 alkyl) wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;

R^3 is H or F;

R^{11} is H or - CH_3 ;

R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or a 4- to 6-membered heterocycle; and

R^{22} is H, halogen or C_1 - C_2 alkyl.

In various aspects, the present disclosure provides pharmaceutical compositions comprising any one of the compounds of Structures I, II, III, IIIb, IIIc, IV, V and VI and a pharmaceutically acceptable carrier, excipient, or diluent.

In various aspects, the present disclosure provides methods of treating a condition characterized by dysregulation of a fatty acid synthase pathway in subject, the method comprising administering to a subject in need of such treatment an effective amount of a compound of any one of the Structures I, II, III, IIIb, IIIc, IV, V and VI. In various aspects, the condition characterized by dysregulation of a fatty acid synthase pathway is a viral infection or cancer. In various aspects, the viral infection is treated using a compound of any one of the Structures I, II, III, IIIb, IIIc, IV, V and VI in combination with one or more additional antiviral treatments. In various aspects, the cancer is treated using a compound of any one of the Structures I, II, III, IIIb, IIIc, IV, V and VI in combination with one or more additional cancer treatments. In various aspects, the viral infection is hepatitis C. In various aspects, the cancer is breast cancer. In various aspects, the cancer is pancreatic cancer. In various aspects, the cancer is colon cancer.

DETAILED DESCRIPTION

The present disclosure addresses the deficiencies in treating conditions characterized by dysregulation of the FASN function in a subject, such as viral infection, cancer and metabolic disorders, by providing novel heterocyclic modulators of lipid synthesis.

5 In certain aspects, the present disclosure provides compositions and methods for treatment of viral infections. In general, the compositions and methods for treatment of viral infections are directed toward modulation of the fatty acid synthesis pathway. The fatty acid synthesis pathway is involved in the replication of viruses in the host cells. The present invention embodies methods for the treatment of viral infections that interact with the fatty
10 acid synthesis pathway, such as hepatitis C.

In certain aspects, the present disclosure provides compositions and methods for the treatment of cancer. Fatty acid synthase is responsible for conversion of malonyl-CoA into long-chain fatty acids, which is an early reaction in fatty acid biosynthesis. Fatty acid synthase is overexpressed in many cancer cells. Without being bound by any particular
15 theory, it is hypothesized that inhibition of fatty acid synthase expression or fatty acid synthase activity selectively suppresses proliferation and induces cell death of cancer cells, with little toxicity towards normal cells.

Further, the present disclosure provides compounds and methods for modulating host cell targets that are targeted by viruses. Such modulation of host cell targets can include
20 either activation or inhibition of the host cell targets. Accordingly, compounds that modulate components of the fatty acid synthesis pathway, such as the activity of a non-viral protein, e.g., a host cell protein, can be used as antiviral pharmaceutical agents.

Definitions

Chemical moieties referred to as univalent chemical moieties (e.g., alkyl, aryl, etc.)
25 also encompass structurally permissible multivalent moieties, as understood by those skilled in the art. For example, while an “alkyl” moiety generally refers to a monovalent radical (e.g., $\text{CH}_3\text{CH}_2\cdot$), in appropriate circumstances an “alkyl” moiety can also refer to a divalent radical (e.g., $\cdot\text{CH}_2\text{CH}_2\cdot$, which is equivalent to an “alkylene” group). Similarly, under circumstances where a divalent moiety is required, those skilled in the art will understand that the term
30 “aryl” refers to the corresponding divalent arylene group.

All atoms are understood to have their normal number of valences for bond formation (e.g., 4 for carbon, 3 for N, 2 for O, and 2, 4, or 6 for S, depending on the atom's oxidation state). On occasion a moiety can be defined, for example, as (A)_aB, wherein a is 0 or 1. In such instances, when a is 0 the moiety is B and when a is 1 the moiety is AB.

Where a substituent can vary in the number of atoms or groups of the same kind (e.g., alkyl groups can be C₁, C₂, C₃, etc.), the number of repeated atoms or groups can be represented by a range (e.g., C₁-C₆ alkyl) which includes each and every number in the range and any and all sub ranges. For example, C₁-C₃ alkyl includes C₁, C₂, C₃, C₁₋₂, C₁₋₃, and C₂₋₃ alkyl.

"Alkanoyl" refers to a carbonyl group with a lower alkyl group as a substituent.

"Alkylamino" refers to an amino group substituted by an alkyl group.

"Alkoxy" refers to an O-atom substituted by an alkyl group as defined herein, for example, methoxy [–OCH₃, a C₁alkoxy]. The term "C₁₋₆ alkoxy" encompasses C₁ alkoxy, C₂ alkoxy, C₃ alkoxy, C₄ alkoxy, C₅ alkoxy, C₆ alkoxy, and any sub-range thereof.

"Alkoxy carbonyl" refers to a carbonyl group with an alkoxy group as a substituent.

"Alkyl carbonyloxy" refers to the group –O-(C=O)-alkyl.


"Alkyl," "alkenyl," and "alkynyl," refer to optionally substituted, straight and branched chain aliphatic groups having from 1 to 30 carbon atoms, or preferably from 1 to 15 carbon atoms, or more preferably from 1 to 6 carbon atoms. Examples of alkyl groups include, without limitation, methyl, ethyl, propyl, isopropyl, butyl, *tert*-butyl, isobutyl, pentyl, hexyl, vinyl, allyl, isobutenyl, ethynyl, and propynyl. The term "heteroalkyl" as used herein contemplates an alkyl with one or more heteroatoms. The term "haloalkyl" as used herein contemplates an alkyl having one to three halogen substituents.

"Alkylene" refers to an optionally substituted divalent radical which is a branched or unbranched hydrocarbon fragment containing the specified number of carbon atoms, and having two points of attachment. An example is propylene [–CH₂CH₂CH₂–, a C₃alkylene].

"Amino" refers to the group –NH₂.


"Aryl" refers to optionally substituted aromatic groups which have at least one ring having a conjugated pi electron system and includes carbocyclic aryl, and biaryl groups, all of which can be optionally substituted. Phenyl and naphthyl groups are preferred carbocyclic aryl groups.

“Aralkyl” or “arylalkyl” refer to alkyl-substituted aryl groups. Examples of aralkyl groups include butylphenyl, propylphenyl, ethylphenyl, methylphenyl, 3,5-dimethylphenyl, tert-butylphenyl.



“Carbamoyl” as used herein contemplates a group of the structure

5 where in R^N is selected from the group consisting of hydrogen, -OH, C_1 to C_{12} alkyl, C_1 to C_{12} heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycle, aryl, heteroaryl, aralkyl, alkoxy, alkoxycarbonyl, alkanoyl, carbamoyl, sulfonyl, sulfonate and sulfonamide.



“Carbonyl” refers to a group of the structure

10 “Cycloalkyl” refers to an optionally substituted ring, which can be saturated or unsaturated and monocyclic, bicyclic, or tricyclic formed entirely from carbon atoms. An example of a cycloalkyl group is the cyclopentenyl group (C_5H_7-), which is a five carbon (C_5) unsaturated cycloalkyl group.

15 “Heterocycle” refers to an optionally substituted 5- to 7-membered cycloalkyl ring system containing 1, 2 or 3 heteroatoms, which can be the same or different, selected from N, O or S, and optionally containing one double bond. “Heterocycle” also refers to an optionally substituted 4- to 8-membered cycloalkyl ring system containing 1, 2 or 3 heteroatoms, which can be the same or different, selected from N, O or S, and optionally containing one double bond.

20 “Halogen” refers to a chloro, bromo, fluoro or iodo atom radical. The term “halogen” also contemplates terms “halo” or “halide.”

“Heteroatom” refers to a non-carbon atom, where boron, nitrogen, oxygen, sulfur and phosphorus are preferred heteroatoms, with nitrogen, oxygen and sulfur being particularly preferred heteroatoms in the compounds of the present disclosure.

25 “Heteroaryl” refers to optionally substituted aryl groups having from 1 to 9 carbon atoms and the remainder of the atoms are heteroatoms, and includes those heterocyclic systems described in “Handbook of Chemistry and Physics,” 49th edition, 1968, R. C. Weast, editor; The Chemical Rubber Co., Cleveland, Ohio. See particularly Section C, Rules for Naming Organic Compounds, B. Fundamental Heterocyclic Systems. Suitable heteroaryls


include thienyl, pyrrolyl, furyl, pyridyl, pyrimidyl, pyrazinyl, pyrazolyl, oxazolyl, isoxazolyl, imidazolyl, thiazolyl, pyranal, tetrazolyl, pyrrolyl, pyrrolinyl, pyridazinyl, triazolyl, indolyl, isoindolyl, indolizynyl, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl, oxadiazolyl, benzoxazolyl, benzoxadiazolyl, thiadiazolyl,

5 benzothiazolyl, benzothiadiaazolyl, and the like.

An “optionally substituted” moiety can be substituted with from one to four, or preferably from one to three, or more preferably one or two non-hydrogen substituents. Unless otherwise specified, when the substituent is on a carbon, it is selected from the group consisting of -OH, -CN, -NO₂, halogen, C₁-C₁₂ alkyl, C₁-C₁₂ heteroalkyl, cycloalkyl, heterocycle, aryl, heteroaryl, aralkyl, alkoxy, alkoxycarbonyl, alkanoyl, carbamoyl, sulfonyl, sulfonate, sulfonamide and amino, none of which are further substituted. Unless otherwise specified, when the substituent is on a carbon, it may also be selected from the group consisting of oxo. Unless otherwise specified, when the substituent is on a carbon, it may also be selected from the group consisting of alkylcarbonyloxy, which is not further substituted. Unless otherwise specified, when the substituent is on a carbon, it may also be selected from the group consisting of alkylamino, which is not further substituted. Unless otherwise specified, when the substituent is on a carbon, it may also be selected from the group consisting of C₁-C₁₂ alkenyl and C₁-C₁₂ alkynyl, neither of which are further substituted. Unless otherwise specified, when the substituent is on a nitrogen, it is selected from the group consisting of C₁-C₁₂ alkyl, C₁-C₁₂ heteroalkyl, cycloalkyl, heterocycle, aryl, heteroaryl, aralkyl, alkoxy, alkoxycarbonyl, alkanoyl, carbamoyl, sulfonyl, sulfonate and sulfonamide, none of which are further substituted. Unless otherwise specified, when the substituent is on a nitrogen, it may also be selected from the group consisting of C₁-C₁₂ alkenyl and C₁-C₁₂ alkynyl, neither of which are further substituted.


25 “Oxo” refers to the =O substituent.

The term “sulfonamide” as used herein contemplates a group having the structure



wherein R^N is selected from the group consisting of hydrogen, -OH, C₁-C₁₂ alkyl, C₁-C₁₂ heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycle, aryl, heteroaryl, aralkyl, alkoxy, alkoxycarbonyl, alkanoyl, carbamoyl, sulfonyl, sulfonate and sulfonamide.


The term “sulfonate” as used herein contemplates a group having the structure



wherein R^{S} is selected from the group consisting of hydrogen, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_1 - C_{10} alkanoyl, or C_1 - C_{10} alkoxy carbonyl.

“Sulfonyl” as used herein alone or as part of another group, refers to an SO_2 group.

- 5 The SO_2 moiety is optionally substituted. In particular, “sulfonyl” as used herein



contemplates a group having the structure , wherein R^{M} is selected from the group consisting of hydrogen, C_1 - C_{12} alkyl, C_1 - C_{12} heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycle, aryl, heteroaryl, aralkyl and alkoxy.

- Compounds of the present disclosure can exist as stereoisomers, wherein asymmetric or chiral centers are present. Stereoisomers are designated (R) or (S) depending on the configuration of substituents around the chiral carbon atom. The terms (R) and (S) used herein are configurations as defined in IUPAC 1974 Recommendations for Section E, Fundamental Stereochemistry, Pure Appl. Chem., (1976), 45: 13-30, hereby incorporated by reference. The present disclosure contemplates various stereoisomers and mixtures thereof and are specifically included within the scope of the present disclosure. Stereoisomers include enantiomers, diastereomers, and mixtures of enantiomers or diastereomers. Individual stereoisomers of compounds of the present disclosure can be prepared synthetically from commercially available starting materials which contain asymmetric or chiral centers or by preparation of racemic mixtures followed by resolution well-known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography and liberation of the optically pure product from the auxiliary or (2) direct separation of the mixture of optical enantiomers on chiral chromatographic columns.
- Also, moieties disclosed herein which exist in multiple tautomeric forms include all such forms encompassed by a given tautomeric structure.

Individual atoms in the disclosed compounds may be any isotope of that element. For example hydrogen may be in the form of deuterium.

“Pharmaceutically acceptable” means approved or approvable by a regulatory agency of the Federal or state government or listed in the U.S. Pharmacopoeia or other generally recognized pharmacopoeia for use in animals, and more particularly in humans. It can be material which is not biologically or otherwise undesirable, i.e., the material can be administered to an individual without causing any undesirable biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained.

The term “pharmaceutically acceptable salt” of a compound means a salt that is pharmaceutically acceptable and that possesses the desired pharmacological activity of the parent compound. Such salts include, for example, acid addition salts and base addition salts.

“Acid addition salts” according to the present disclosure, are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 2-naphthalenesulfonic acid, 4-methylbicyclo-[2.2.2]oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis-(3-hydroxy-2-ene-1-carboxylic acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like.

“Base addition salts” according to the present disclosure are formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base.

Acceptable organic bases include ethanolamine, diethanolamine, triethanolamine, tromethamine, N-methylglucamine, and the like. Acceptable inorganic bases include aluminum hydroxide, calcium hydroxide, potassium hydroxide, sodium carbonate, sodium hydroxide, and the like.

It should be understood that a reference to a pharmaceutically acceptable salt includes the solvent addition forms or crystal forms thereof, particularly solvates or polymorphs.

Solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and are often formed during the process of crystallization. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. Polymorphs include the different crystal packing arrangements of the same elemental composition of a compound. Polymorphs usually have different X-ray diffraction patterns, infrared spectra, melting points, density, hardness, crystal shape, optical and electrical properties, stability, and solubility. Various factors such as the recrystallization solvent, rate of crystallization, and storage temperature can cause a single crystal form to dominate.

The term “treating” includes the administration of the compounds or agents of the present invention to a subject to prevent or delay, to alleviate, or to arrest or inhibit development of the symptoms or conditions associated with fatty acid synthase-associated disorders.

A “therapeutically effective amount” or “pharmaceutically effective amount” means the amount that, when administered to a subject, produces effects for which it is administered. For example, a “therapeutically effective amount,” when administered to a subject to inhibit fatty acid synthase activity, is sufficient to inhibit fatty acid synthase activity. A “therapeutically effective amount,” when administered to a subject for treating a disease, is sufficient to effect treatment for that disease.

Except when noted, the terms “subject” or “patient” are used interchangeably and refer to mammals such as human patients and non-human primates, as well as experimental animals such as rabbits, rats, and mice, and other animals. Accordingly, the term “subject” or “patient” as used herein means any mammalian patient or subject to which the compounds of the invention can be administered. In an exemplary aspect of the present invention, to identify subject patients for treatment according to the methods of the invention, accepted screening methods are employed to determine risk factors associated with a targeted or suspected disease or condition or to determine the status of an existing disease or condition in a subject. These screening methods include, for example, conventional work-ups to determine risk factors that are associated with the targeted or suspected disease or condition. These and

other routine methods allow the clinician to select patients in need of therapy using the methods and formulations of the present invention.


Chemical names for the compounds of the present disclosure were generated using ChemDraw Ultra version 12.0 (CambridgeSoft Corp., Cambridge Mass).

5 FASN Pathway Modulators

One aspect of the present disclosure includes a method of inhibiting viral infection or treating cancer by contacting a cell with an agent that modulates the fatty acid synthesis pathway. This method of inhibiting viral infection or treating cancer can be performed *in vitro* by contacting virally infected/cancerous cells with an agent that modulates the fatty acid
10 synthesis pathway, or *in vivo* by administering an agent that modulates the fatty acid synthesis pathway to a subject infected with a virus/having cancer. In one aspect, an agent can be an inhibitor of the fatty acid synthesis pathway.

Examples of inhibitors of the fatty acid synthesis pathway that can be used in the methods and compositions of the present disclosure are described below.

15 In various aspects, the present disclosure provides for compounds of Structure I:




I, or pharmaceutically acceptable salts thereof, wherein:


L^3 is $-\text{CH}_2-$, $-\text{CHR}^{50}-$, $-\text{O}-$, $-\text{NR}^{50}-$, $-\text{NC}(\text{O})\text{R}^{50}-$ or $-\text{NC}(\text{O})\text{OR}^{50}-$, wherein R^{50} is C_1 - C_6 alkyl,

C_3 - C_5 cycloalkyl, or 4- to 6-membered heterocycle;

n is 1, 2, or 3;

20 m is 1 or 2 with the proviso that $n+m \geq 3$;





Het is a 5- to 6-membered heteroaryl;

- 5 R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered heterocycle) or -O-(C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;

R^3 is H or F;

- 10 R^{11} is H or -CH₃;

R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or a 4- to 6-membered heterocycle; and


R^{22} is H, halogen, or C_1 - C_2 alkyl.


- As noted above, each of the C_1 - C_2 alkyl, C_1 - C_4 alkyl, C_1 - C_6 alkyl, C_3 - C_5 cycloalkyl, 4- to 6-membered heterocycle and 5- to 6-membered heteroaryl moieties may be optionally substituted. Accordingly, the present disclosure provides for compounds of Structure I wherein:

L^3 is -CH₂-, CHR⁵⁰-, -O-, -NR⁵⁰-, -NC(O)R⁵⁰- or -NC(O)OR⁵⁰-, wherein R⁵⁰ is optionally substituted C_1 - C_6 alkyl, optionally substituted C_3 - C_5 cycloalkyl or optionally substituted 4- to 6-membered heterocycle;

- 20 n is 1, 2 or 3;

m is 1 or 2 with the proviso that $n+m \geq 3$;





Het is a optionally substituted 5- to 6-membered heteroaryl;

- 5 R^1 is H, -CN, halogen, optionally substituted C_1 - C_4 alkyl, -O-(optionally substituted C_3 - C_5 cycloalkyl), -O-(optionally substituted 4- to 6-membered heterocycle) or -O-(optionally substituted C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or optionally substituted C_1 - C_4 alkyl;


- 10 R^3 is H or F;

R^{11} is H or -CH₃;


R^{21} is H, halogen, optionally substituted C_1 - C_4 alkyl, optionally substituted C_3 - C_5 cycloalkyl or an optionally substituted 4- to 6-membered heterocycle; and


R^{22} is H, halogen or optionally substituted C_1 - C_2 alkyl.

- 15 In some embodiments, the present disclosure provides for compounds of Structure I



In some embodiments, the present disclosure provides for compounds of Structure I





In some embodiments, the present disclosure provides for compounds of Structure I wherein R^1 is H, -CN, -C₁-C₄ alkyl, -O-(C₃-C₅ cycloalkyl), -O-(4- to 6-membered heterocycle) or -O-(C₁-C₄ alkyl) wherein when R^1 is not H or -CN, R^1 is optionally substituted with one or more halogens.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^1 is halogen, -CN or C₁-C₂ haloalkyl.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^1 is -CN or C₁-C₂ haloalkyl.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^1 is -CN.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^1 is -Cl.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^2 is H.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^{21} is halogen, C₁-C₄ alkyl, C₃-C₅ cycloalkyl or 4- to 6-membered heterocycle.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^{21} is C₁-C₂ alkyl or C₃-C₅ cycloalkyl.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^{21} is C₃-C₅ cycloalkyl.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^{22} is H or C₁-C₂ alkyl.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^{22} is H.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^{22} is C₁-C₂ alkyl.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^{22} is -CH₃.

In some embodiments, the present disclosure provides for compounds of Structure I wherein L^3 is $-N(CH_3)-$.

In some embodiments, the present disclosure provides for compounds of Structure I wherein n is 2 and m is 2.

5 In some embodiments, the present disclosure provides for compounds of Structure I wherein n is 1 or 2.

In some embodiments, the present disclosure provides for compounds of Structure I wherein n is 1 and m is 2.


10 In some embodiments, the present disclosure provides for compounds of Structure I wherein R^{21} is C_1 - C_2 alkyl or C_3 - C_5 cycloalkyl and R^{22} is C_1 - C_2 alkyl.

In some embodiments, the present disclosure provides for compounds of Structure I wherein R^{21} is C_1 - C_2 alkyl or C_3 - C_5 cycloalkyl and R^{22} is H or C_1 - C_2 alkyl.





In some embodiments, the present disclosure provides for compounds of Structure I wherein R^{21} is C_3 - C_5 cycloalkyl and R^{22} is H or C_1 - C_2 alkyl.

15 In some embodiments, the present disclosure provides for compounds of Structure I wherein R^{21} is C_3 - C_5 cycloalkyl and R^{22} is H or $-CH_3$.

In various aspects, the present disclosure provides for compounds of Structure II:



II, or pharmaceutically acceptable salts thereof, wherein:



Het is a 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered heterocycle) or -O-(C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

5 each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;

R^3 is H or F;

R^{11} is H or -CH₃;

R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle;


R^{22} is H, halogen, or C_1 - C_2 alkyl; and

10 R^{24} is H, C_1 - C_4 alkyl, -(C_1 - C_4 alkyl)-OH, -(C_1 - C_4 alkyl)_t-N(R^{241})₂, -(C_1 - C_4 alkyl)_t-O-(C_3 - C_5 cycloalkyl), -(C_1 - C_4 alkyl)_t-O-(4- to 6-membered heterocycle) or -(C_1 - C_4 alkyl)_t-O-(C_1 - C_4 alkyl), wherein:

each t is independently 0 or 1; and

each R^{241} is independently H or C_1 - C_2 alkyl.

15 As noted above, each of the C_1 - C_2 alkyl, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl, 4- to 6-membered heterocycle and 5- to 6-membered heteroaryl moieties may be optionally substituted. Accordingly, the present disclosure provides for compounds of Structure II wherein:



Het is an optionally substituted 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, optionally substituted C_1 - C_4 alkyl, -O-(optionally substituted C_3 - C_5 cycloalkyl), -O-(optionally substituted 4- to 6-membered heterocycle) or -O-(optionally substituted C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

5 each R^2 is independently hydrogen, halogen or optionally substituted C_1 - C_4 alkyl;

R^3 is H or F;

R^{11} is H or - CH_3 ;

R^{21} is H, halogen, optionally substituted C_1 - C_4 alkyl, optionally substituted C_3 - C_5 cycloalkyl or optionally substituted 4- to 6-membered heterocycle;


10 R^{22} is H, halogen or optionally substituted C_1 - C_2 alkyl; and

R^{24} is H, optionally substituted C_1 - C_4 alkyl, -(optionally substituted C_1 - C_4 alkyl)-OH, -(optionally substituted C_1 - C_4 alkyl) $_t$ -N(R^{241}) $_2$, -(optionally substituted C_1 - C_4 alkyl) $_t$ -O-(optionally substituted C_3 - C_5 cycloalkyl), -(optionally substituted C_1 - C_4 alkyl) $_t$ -O-(optionally substituted 4- to 6-membered heterocycle) or -(optionally substituted C_1 - C_4 alkyl) $_t$ -O-(optionally substituted C_1 - C_4 alkyl), wherein:


t is 0 or 1; and

R^{241} is H or optionally substituted C_1 - C_2 alkyl.


In some embodiments, the present disclosure provides for compounds of Structure II



In some embodiments, the present disclosure provides for compounds of Structure II



In some embodiments, the present disclosure provides for compounds of Structure II



wherein Ar is

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^1 is halogen, -CN or C_1 - C_2 haloalkyl.

5 In some embodiments, the present disclosure provides for compounds of Structure II wherein R^1 is -CN.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^2 is H.

10 In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is H, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is C_1 - C_2 alkyl or C_3 - C_5 cycloalkyl.

15 In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is C_1 - C_2 alkyl.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is C_3 - C_5 cycloalkyl.

20 In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{22} is H or C_1 - C_2 alkyl.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{22} is H.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{22} is C_1 - C_2 alkyl.

25 In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{22} is -CH₃.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{24} is C_1 - C_4 alkyl or -(C_1 - C_4 alkyl)_t-O-(C_1 - C_4 alkyl).

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{24} is $-(C_1-C_2 \text{ alkyl})_t-O-(C_1-C_2 \text{ alkyl})$.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{24} is C_1-C_4 alkyl or $-(C_1-C_4 \text{ alkyl})_t-O-(C_1-C_4 \text{ alkyl})$ wherein t is 0 or 1.

5 In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is C_1-C_2 alkyl or C_3-C_5 cycloalkyl and R^{22} is H or C_1-C_2 alkyl.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is C_1-C_2 alkyl or C_3-C_5 cycloalkyl and R^{22} is C_1-C_2 alkyl.

10 In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is C_1-C_2 alkyl or C_3-C_5 cycloalkyl and R^{22} is $-CH_3$.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is C_1-C_2 alkyl or C_3-C_5 cycloalkyl and R^{22} is H.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is C_3-C_5 cycloalkyl and R^{22} is H or C_1-C_2 alkyl.

15 In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is C_3-C_5 cycloalkyl and R^{22} is H or $-CH_3$.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is C_3-C_5 cycloalkyl and R^{22} is C_1-C_2 alkyl.


20 In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is C_3-C_5 cycloalkyl and R^{22} is $-CH_3$.

In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{21} is C_3-C_5 cycloalkyl and R^{22} is H.


In some embodiments, the present disclosure provides for compounds of Structure II wherein R^{24} is $-(C_1-C_2 \text{ alkyl})_t-O-(C_1-C_2 \text{ alkyl})$ and wherein t is 0 or 1.

25 In some embodiments, the present disclosure provides for compounds of Structure II wherein R^1 is $-CN$ and R^2 is H.

In various aspects, the present disclosure provides for compounds of Structure III:




III, or pharmaceutically acceptable salts thereof, wherein:




Het is a 5- to 6-membered heteroaryl;

- 5 R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered heterocycle) or -O-(C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;
- each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;
- R^3 is H or F;
- 10 R^{11} is H or -CH₃;
- R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle;
- R^{22} is H, halogen, or C_1 - C_2 alkyl;
- R^{24} is H, -CN, -(C_1 - C_4 alkyl)-CN, C_1 - C_4 alkyl, -(C_1 - C_4 alkyl)-OH, -(C_1 - C_4 alkyl)-N(R^{241})₂, -(C_1 - C_4 alkyl)_t-O_u-(C_3 - C_6 cycloalkyl), -(C_1 - C_4 alkyl)_t-O_u-(4- to 6-membered
- 15 heterocycle) or -(C_1 - C_4 alkyl)-O-(C_1 - C_4 alkyl), wherein:
- t is 0 or 1;
- u is 0 or 1;
- with the proviso that when u is 1, t is 1; and
- each R^{241} is independently H or C_1 - C_2 alkyl; and
- 20 R^{25} is halogen, -CN, -(C_1 - C_4 alkyl)-CN, C_1 - C_2 alkyl or cyclopropyl.

As noted above, each of the C_1 - C_2 alkyl (i.e., methyl and ethyl), cyclopropyl, C_1 - C_2 alkyl, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl, C_3 - C_6 cycloalkyl, 4- to 6-membered heterocycle and 5- to 6-membered heteroaryl moieties may be optionally substituted. Accordingly, the present disclosure provides for compounds of Structure III wherein:





Het is an optionally substituted 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, optionally substituted C_1 - C_4 alkyl, -O-(optionally substituted C_3 - C_5 cycloalkyl), -O-(optionally substituted 4- to 6-membered heterocycle) or -O-(optionally substituted C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or optionally substituted C_1 - C_4 alkyl;

R^3 is H or F;

R^{11} is H or - CH_3 ;

R^{21} is H, halogen, optionally substituted C_1 - C_4 alkyl, optionally substituted C_3 - C_5 cycloalkyl or optionally substituted 4- to 6-membered heterocycle;

R^{22} is H, halogen or optionally substituted C_1 - C_2 alkyl;

R^{24} is H, -CN, -(optionally substituted C_1 - C_4 alkyl)-CN, optionally substituted C_1 - C_4 alkyl, -(optionally substituted C_1 - C_4 alkyl)-OH, -(optionally substituted C_1 - C_4 alkyl)- $N(R^{241})_2$, -(optionally substituted C_1 - C_4 alkyl) $_t$ -O $_u$ -(optionally substituted C_3 - C_6 cycloalkyl), -(optionally substituted C_1 - C_4 alkyl) $_t$ -O $_u$ -(optionally substituted 4- to 6-membered heterocycle) or -(optionally substituted C_1 - C_4 alkyl)-O-(optionally substituted C_1 - C_4 alkyl), wherein:

t is 0 or 1;


u is 0 or 1;

with the proviso that when u is 1, t is 1; and


R^{241} is H or optionally substituted C_1 - C_2 alkyl; and

R^{25} is halogen, -CN, -(optionally substituted C_1 - C_4 alkyl)-CN, optionally substituted methyl, optionally substituted ethyl or optionally substituted cyclopropyl.


In some embodiments, the present disclosure provides for compounds of Structure III



In some embodiments, the present disclosure provides for compounds of Structure III



In some embodiments, the present disclosure provides for compounds of Structure III



In some embodiments, the present disclosure provides for compounds of Structure III



In some embodiments, the present disclosure provides for compounds of Structure III wherein R¹ is halogen, -CN or C₁-C₂ haloalkyl.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R¹ is -CN.

15 In some embodiments, the present disclosure provides for compounds of Structure III wherein R² is H.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R²¹ is halogen, C₁-C₄ alkyl or C₃-C₅ cycloalkyl.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{21} is C_1 - C_4 alkyl or C_3 - C_5 cycloalkyl.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{21} is C_1 - C_2 alkyl or C_3 - C_5 cycloalkyl.

5 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{21} is C_1 - C_2 alkyl.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{21} is $-CH_3$.

10 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{22} is H or C_1 - C_2 alkyl.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{22} is H or $-CH_3$.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{22} is $-CH_3$.

15 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is H, $-CN$, $-(C_1-C_4 \text{ alkyl})-CN$, C_1-C_4 alkyl, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_6 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4- \text{ to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$.

20 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is H, C_1-C_4 alkyl, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_6 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4- \text{ to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is C_1 - C_4 alkyl or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$.

25 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is $-(C_1-C_2 \text{ alkyl})-O-(C_1-C_2 \text{ alkyl})$.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is $-CH_2-O-CH_3$.

30 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is C_1 - C_2 alkyl.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is $-CH_3$.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is C_3 - C_6 cycloalkyl.

5 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is C_3 - C_5 cycloalkyl.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is $-CN$ or $-(C_1$ - C_2 alkyl)- CN .

10 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is $-CN$.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is $-(C_1$ - C_2 alkyl)- CN .

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is H , $-CH_3$, $-CH_2OH$, $-CH_2OCH_3$, $-(CH_2)_2OH$, $-(CH_2)_2OCH_3$ or $-(CH_2)_2N(CH_3)_2$.

15 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is methyl, isopropyl, cyclopropyl, $-CN$, or $-(C_1$ - C_2 alkyl)- CN .

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is substituted with one or more substituents selected from C_1 - C_2 alkyl, oxo, $-CN$, halogen, alkanoyl, alkoxy, $-OH$ and C_1 - C_2 alkoxy.

20 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is substituted with one or more substituents selected from methyl, $-F$, methoxy, $-C(=O)CH_3$ and $-C(=O)-OCH_3$.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is substituted with two substituents that are the same or different.

25 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is substituted with three substituents that are the same or different.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{25} is halogen, $-CN$, C_1 - C_2 alkyl or cyclopropyl.

30 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{25} is halogen, C_1 - C_2 alkyl or cyclopropyl.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{25} is -CN, -Cl or -CH₃.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{25} is -Cl.

5 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{25} is -CH₃.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{25} is substituted with one or more substituents selected from -OH, halogen, C₁-C₂ alkyl and alkylcarbonyloxy.

10 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{25} is substituted with one or more substituents selected from -F, methyl and -O-C(=O)-CH₃.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{25} is substituted with two substituents that are the same or different.

15 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{25} is substituted with three substituents that are the same or different.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{24} is C₁-C₄ alkyl, -(C₁-C₄ alkyl)-CN or -(C₃-C₆ cycloalkyl).

In some embodiments, the present disclosure provides for compounds of Structure III
20 wherein R^{24} is -CN, -(C₁-C₂ alkyl)-CN, -(C₃-C₆ cycloalkyl) or methyl.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{25} is is halogen, methyl, ethyl or cyclopropyl.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{25} is halogen, -CN, methyl, ethyl or cyclopropyl.

25 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{21} is C₁-C₂ alkyl or C₃-C₆ cycloalkyl and R^{22} is H or -CH₃.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{21} is C₁-C₂ alkyl or C₃-C₆ cycloalkyl, R^{22} is H or -CH₃, R^{24} is -CH₂-O-CH₃ and R^{25} is -CH₃.

30 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{21} is -CH₃ and R^{22} is H.


In some embodiments, the present disclosure provides for compounds of Structure III wherein R^1 is $-\text{CN}$ and R^2 is H

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{21} is $\text{C}_1\text{-C}_2$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl and R^{22} is H or $\text{C}_1\text{-C}_2$ alkyl.

5 In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{21} is $\text{C}_1\text{-C}_2$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl, R^{22} is H or $\text{C}_1\text{-C}_2$ alkyl, R^{24} is $-\text{CH}_2\text{-O-CH}_3$ and R^{25} is $-\text{CH}_3$.

In some embodiments, the present disclosure provides for compounds of Structure III wherein R^{21} is $\text{C}_1\text{-C}_2$ alkyl and R^{22} is H .

10 In various aspects, the present disclosure provides for compounds of Structure IIIb:



15 Het is a 5- to 6-membered heteroaryl;

R^1 is H , $-\text{CN}$, halogen, $\text{C}_1\text{-C}_4$ alkyl, $-\text{O-}(\text{C}_3\text{-C}_5 \text{ cycloalkyl})$, $-\text{O-}(4\text{- to } 6\text{-membered heterocycle})$ or $-\text{O-}(\text{C}_1\text{-C}_4 \text{ alkyl})$, wherein when R^1 is not H , $-\text{CN}$ or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or $\text{C}_1\text{-C}_4$ alkyl;

20 R^3 is H or F ;

R^{11} is H or $-\text{CH}_3$;


R^{21} is H , halogen, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_3\text{-C}_5$ cycloalkyl or 4- to 6-membered heterocycle;

R^{22} is H , halogen or $\text{C}_1\text{-C}_2$ alkyl; and


each R^{24} and R^{25} is independently H, halogen, -CN, -(C₁-C₄ alkyl)-CN, C₁-C₄ alkyl, -(C₁-C₄ alkyl)-OH, -(C₁-C₄ alkyl)-N(R^{241})₂, -(C₁-C₄ alkyl)_t-O_u-(C₃-C₅ cycloalkyl), -(C₁-C₄ alkyl)_t-O_u-(4- to 6-membered heterocycle) or -(C₁-C₄ alkyl)_t-O-(C₁-C₄ alkyl),
wherein:

- 5 each t is independently 0 or 1;
each u is independently 0 or 1; and
each R^{241} is independently H or C₁-C₂ alkyl,

wherein the compound is not:



- 10 As noted above, each of the C₁-C₂ alkyl, C₁-C₄ alkyl, C₃-C₅ cycloalkyl, 4- to 6-membered heterocycle and 5- to 6-membered heteroaryl moieties may be optionally substituted. Accordingly, the present disclosure provides for compounds of Structure IIIb wherein:



R^{11} is H or $-CH_3$;

R^{21} is H, halogen, optionally substituted C_1 - C_4 alkyl, optionally substituted C_3 - C_5 cycloalkyl or optionally substituted 4- to 6-membered heterocycle;

R^{22} is H, halogen or optionally substituted C_1 - C_2 alkyl; and


5 each R^{24} and R^{25} is independently H, halogen, $-CN$, $-(\text{optionally substituted } C_1\text{-}C_4 \text{ alkyl})-CN$, optionally substituted C_1 - C_4 alkyl, $-(\text{optionally substituted } C_1\text{-}C_4 \text{ alkyl})-OH$, $-(\text{optionally substituted } C_1\text{-}C_4 \text{ alkyl})-N(R^{241})_2$, $-(\text{optionally substituted } C_1\text{-}C_4 \text{ alkyl})_t-O_u-(\text{optionally substituted } C_3\text{-}C_5 \text{ cycloalkyl})$, $-(\text{optionally substituted } C_1\text{-}C_4 \text{ alkyl})_t-O_u-(\text{optionally substituted 4- to 6-membered heterocycle})$ or $-(\text{optionally substituted } C_1$ -
10 $C_4 \text{ alkyl})_t-O-(\text{optionally substituted } C_1\text{-}C_4 \text{ alkyl})$, wherein:

t is 0 or 1;

u is 0 or 1; and


R^{241} is H or optionally substituted C_1 - C_2 alkyl,

wherein the compound is not:




15

In some embodiments, the present disclosure provides for compounds of Structure




In some embodiments, the present disclosure provides for compounds of Structure



20

In some embodiments, the present disclosure provides for compounds of Structure



In some embodiments, the present disclosure provides for compounds of Structure



In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R¹ is halogen, -CN or C₁-C₂ haloalkyl.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R¹ is -CN.

10 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R² is H.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R²¹ is halogen, C₁-C₄ alkyl, C₃-C₅ cycloalkyl or 4- to 6-membered heterocycle.

15 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R²¹ is C₁-C₄ alkyl, C₃-C₅ cycloalkyl or 4- to 6-membered heterocycle.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R²¹ is C₁-C₂ alkyl or C₃-C₅ cycloalkyl.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R²¹ is C₁-C₂ alkyl.

20 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R²¹ is -CH₃.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R²² is H or C₁-C₂ alkyl.

25 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R²² is H or -CH₃.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{22} is $-\text{CH}_3$.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein each R^{24} and R^{25} is independently H, $-\text{CN}$, $\text{C}_1\text{-C}_4$ alkyl, $-(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-OH}$, $-(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-N}(\text{R}^{241})_2$, $-(\text{C}_1\text{-C}_4 \text{ alkyl})_t\text{-O}_u\text{-(C}_3\text{-C}_5 \text{ cycloalkyl)}$, $-(\text{C}_1\text{-C}_4 \text{ alkyl})_t\text{-O}_u\text{-(4- to 6-membered heterocycle)}$ or $-(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-O-(C}_1\text{-C}_4 \text{ alkyl)}$.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein each R^{24} and R^{25} is independently H, $\text{C}_1\text{-C}_4$ alkyl, $-(\text{C}_1\text{-C}_4 \text{ alkyl})_t\text{-O}_u\text{-(4- to 6-membered heterocycle)}$ or $-(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-O-(C}_1\text{-C}_4 \text{ alkyl)}$.

10 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{24} is H, $\text{C}_1\text{-C}_4$ alkyl, $-(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-OH}$, $-(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-N}(\text{R}^{241})_2$, $-(\text{C}_1\text{-C}_4 \text{ alkyl})_t\text{-O}_u\text{-(C}_3\text{-C}_5 \text{ cycloalkyl)}$, $-(\text{C}_1\text{-C}_4 \text{ alkyl})_t\text{-O}_u\text{-(4- to 6-membered heterocycle)}$ or $-(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-O-(C}_1\text{-C}_4 \text{ alkyl)}$.

15 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{24} is $-\text{CN}$, $-\text{Cl}$, $\text{C}_1\text{-C}_4$ alkyl or $-(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-O-(C}_1\text{-C}_4 \text{ alkyl)}$.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{24} is $\text{C}_1\text{-C}_4$ alkyl or $-(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-O-(C}_1\text{-C}_4 \text{ alkyl)}$.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{24} is $-(\text{C}_1\text{-C}_2 \text{ alkyl})\text{-O-(C}_1\text{-C}_2 \text{ alkyl)}$.

20 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{24} is $\text{C}_1\text{-C}_4$ alkyl.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{24} is $-\text{CH}_3$.

25 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{24} is hydrogen.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{24} is substituted with one or more substituents selected from halogen, $\text{C}_3\text{-C}_5$ cycloalkyl and $\text{C}_1\text{-C}_2$ alkoxy.

30 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{24} is substituted with one or more substituents selected from $-\text{F}$, cyclopropyl and $-\text{OCH}_3$.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{24} is substituted with two substituents that are the same or different.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{24} is substituted with three substituents that are the same or different.

5 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{25} is halogen, methyl, ethyl or cyclopropyl.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{25} is -CN, -Cl, C_1 - C_4 alkyl, $-(C_1-C_4 \text{ alkyl})_t-O-(C_3-C_5 \text{ cycloalkyl})$ or $-(C_1-C_4 \text{ alkyl})_t-O-(C_1-C_4 \text{ alkyl})$.

10 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{25} is -CN, -Cl, -CH₃, $-O-(C_3-C_5 \text{ cycloalkyl})$ or $-O-(C_1-C_2 \text{ alkyl})$.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{25} is -CN, -Cl or C_1 - C_4 alkyl.

15 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{25} is -CH₃.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{25} is -Cl.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{25} is substituted with one or more halogen.

20 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{25} is substituted with one or more -F.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{25} is substituted by two substituents.

25 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{25} is substituted by three substituents.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{21} is -CH₃ and R^{22} is H or methyl.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{21} is C_1 - C_2 alkyl or C_3 - C_5 cycloalkyl and R^{22} is H or -CH₃.

30 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{21} is -CH₃ and R^{22} is H.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{24} is H or $-\text{CH}_3$ and R^{25} is $-\text{Cl}$.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^1 is $-\text{CN}$ and R^2 is H.


5 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{21} is C_1 - C_2 alkyl or C_3 - C_5 cycloalkyl and R^{22} is C_1 - C_2 alkyl.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{21} is C_1 - C_2 alkyl or C_3 - C_5 cycloalkyl and R^{22} is H or C_1 - C_2 alkyl.


10 In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{21} is C_1 - C_2 alkyl and R^{22} is H or $-\text{CH}_3$.

In some embodiments, the present disclosure provides for compounds of Structure IIIb wherein R^{21} is C_1 - C_2 alkyl and R^{22} is H.

In various aspects, the present disclosure provides for compounds of Structure IIIc:



IIIc, or pharmaceutically acceptable salts thereof, wherein:



Het is a 5- to 6-membered heteroaryl;

R^1 is H, $-\text{CN}$, halogen, C_1 - C_4 alkyl, $-\text{O}-(\text{C}_3$ - C_5 cycloalkyl), $-\text{O}-(4$ - to 6-membered

20 heterocycle) or $-\text{O}-(\text{C}_1$ - C_4 alkyl), wherein when R^1 is not H, $-\text{CN}$ or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;

R^3 is H or F;


R^{11} is H or $-\text{CH}_3$;

R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle;

R^{22} is H, halogen or C_1 - C_2 alkyl; and

each of R^{24} and R^{25} is independently H, $-C_1$ - C_4 alkyl, or halogen.

As noted above, each of the C_1 - C_2 alkyl, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl, 4- to 6-membered heterocycle and 5- to 6-membered heteroaryl moieties may be optionally substituted. Accordingly, the present disclosure provides for compounds of Structure IIIc wherein:



Het is an optionally substituted 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, optionally substituted C_1 - C_4 alkyl, -O-(optionally substituted C_3 - C_5 cycloalkyl), -O-(optionally substituted 4- to 6-membered heterocycle) or -O-(optionally substituted C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or optionally substituted C_1 - C_4 alkyl;

R^3 is H or F;


R^{11} is H or $-CH_3$;

R^{21} is H, halogen, optionally substituted C_1 - C_4 alkyl, optionally substituted C_3 - C_5 cycloalkyl or optionally substituted 4- to 6-membered heterocycle;


R^{22} is H, halogen or optionally substituted C_1 - C_2 alkyl; and

each of R^{24} and R^{25} is independently H, optionally substituted C_1 - C_4 alkyl, or halogen.


In some embodiments, the present disclosure provides for compounds of Structure



In some embodiments, the present disclosure provides for compounds of Structure



In some embodiments, the present disclosure provides for compounds of Structure




In some embodiments, the present disclosure provides for compounds of Structure IIIc wherein R¹ is halogen, -CN or C₁-C₂ haloalkyl.



10 In some embodiments, the present disclosure provides for compounds of Structure IIIc wherein R²¹ is halogen, C₁-C₄ alkyl, C₃-C₅ cycloalkyl or 4- to 6-membered heterocycle.


In some embodiments, the present disclosure provides for compounds of Structure IIIc wherein R²¹ is -CH₃.

15 In some embodiments, the present disclosure provides for compounds of Structure IIIc wherein R²² is H.

In some embodiments, the present disclosure provides for compounds of Structure IIIc wherein R²¹ is methyl, R²² is H and L-Ar is .

In various aspects, the present disclosure provides for compounds of Structure IV:







Het is a 5- to 6-membered heteroaryl;

- 5 R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered heterocycle) or -O-(C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;
- each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;
- R^3 is H or F;
- 10 R^{11} is H or -CH₃;
- R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle;
- R^{22} is H, halogen or C_1 - C_2 alkyl; and
- R^{24} is H, C_1 - C_4 alkyl, -(C_1 - C_4 alkyl)-OH, -(C_1 - C_4 alkyl)-N(R^{241})₂, -(C_1 - C_4 alkyl)_t-O_u-(C_3 - C_5 cycloalkyl), -(C_1 - C_4 alkyl)_t-O_u-(4- to 6-membered heterocycle) or -(C_1 - C_4 alkyl)-O-(C_1 - C_4 alkyl), wherein:
- 15 t is 0 or 1;
- u is 0 or 1;
- with the proviso that when u is 1, t is 1; and
- R^{241} is H or C_1 - C_2 alkyl.

- 20 As noted above, each of the C_1 - C_2 alkyl, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl, 4- to 6-membered heterocycle and 5- to 6-membered heteroaryl moieties may be optionally substituted. Accordingly, the present disclosure provides for compounds of Structure IV wherein:






Het is an optionally substituted 5- to 6-membered heteroaryl;

- 5 R^1 is H, -CN, halogen, optionally substituted C_1 - C_4 alkyl, -O-(optionally substituted C_3 - C_5 cycloalkyl), -O-(optionally substituted 4- to 6-membered heterocycle) or -O-(optionally substituted C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;
- each R^2 is independently hydrogen, halogen or optionally substituted C_1 - C_4 alkyl;
- 10 R^3 is H or F;
- R^{11} is H or - CH_3 ;
- R^{21} is H, halogen, optionally substituted C_1 - C_4 alkyl, optionally substituted C_3 - C_5 cycloalkyl or optionally substituted 4- to 6-membered heterocycle;
- R^{22} is H, halogen or optionally substituted C_1 - C_2 alkyl;
- 15 R^{24} is H, optionally substituted C_1 - C_4 alkyl, -(optionally substituted C_1 - C_4 alkyl) -OH, -(optionally substituted C_1 - C_4 alkyl)- $N(R^{241})_2$, -(optionally substituted C_1 - C_4 alkyl) $_t$ - O_u - (optionally substituted C_3 - C_5 cycloalkyl), -(optionally substituted C_1 - C_4 alkyl) $_t$ - O_u - (optionally substituted 4- to 6-membered heterocycle) or -(optionally substituted C_1 - C_4 alkyl) -O-(optionally substituted C_1 - C_4 alkyl), wherein:
- 20 t is 0 or 1;
- u is 0 or 1;
- with the proviso that when u is 1, t is 1; and
- R^{241} is H or optionally substituted C_1 - C_2 alkyl.

In some embodiments, the present disclosure provides for compounds of Structure IV



In some embodiments, the present disclosure provides for compounds of Structure IV
5 wherein R¹ is halogen, -CN or C₁-C₂ haloalkyl.

In some embodiments, the present disclosure provides for compounds of Structure IV
wherein R¹ is -CN.

In some embodiments, the present disclosure provides for compounds of Structure IV
wherein R² is H.

10 In some embodiments, the present disclosure provides for compounds of Structure IV
wherein R²¹ is halogen, C₁-C₄ alkyl, C₃-C₅ cycloalkyl or 4- to 6-membered heterocycle.

In some embodiments, the present disclosure provides for compounds of Structure IV
wherein R²¹ is C₁-C₂ alkyl or C₃-C₅ cycloalkyl.

15 In some embodiments, the present disclosure provides for compounds of Structure IV
wherein R²¹ is C₁-C₂ alkyl.

In some embodiments, the present disclosure provides for compounds of Structure IV
wherein R²¹ is C₃-C₅ cycloalkyl.

In some embodiments, the present disclosure provides for compounds of Structure IV
wherein R²² is H or C₁-C₂ alkyl.

20 In some embodiments, the present disclosure provides for compounds of Structure IV
wherein R²² is H.

In some embodiments, the present disclosure provides for compounds of Structure IV
wherein R²² is C₁-C₂ alkyl.

25 In some embodiments, the present disclosure provides for compounds of Structure IV
wherein R²² is -CH₃.

In some embodiments, the present disclosure provides for compounds of Structure IV
wherein R²⁴ is C₁-C₄ alkyl or -(C₁-C₄ alkyl)-O-(C₁-C₄ alkyl).

In some embodiments, the present disclosure provides for compounds of Structure IV wherein R^{24} is $-(C_1-C_2 \text{ alkyl})-O-(C_1-C_2 \text{ alkyl})$.

In some embodiments, the present disclosure provides for compounds of Structure IV wherein R^{21} is C_1-C_2 alkyl or C_3-C_5 cycloalkyl and R^{22} is C_1-C_2 alkyl.


5 In some embodiments, the present disclosure provides for compounds of Structure IV wherein R^{21} is C_3-C_5 cycloalkyl and R^{22} is H or C_1-C_2 alkyl.

In some embodiments, the present disclosure provides for compounds of Structure IV wherein R^{21} is C_3-C_5 cycloalkyl and R^{22} is H or $-CH_3$.


10 In some embodiments, the present disclosure provides for compounds of Structure IV wherein R^{21} is C_3-C_5 cycloalkyl and R^{22} is H or $-CH_3$.

In some embodiments, the present disclosure provides for compounds of Structure IV wherein R^1 is $-CN$ and R^2 is H.

In various aspects, the present disclosure provides for compounds of Structure V:



V, or pharmaceutically acceptable salts thereof, wherein:




L^2 is $-NHR^{35}$ or $-C(O)NHR^{351}$, wherein R^{351} is C_1-C_6 alkyl, C_3-C_5 cycloalkyl,

20 4- to 6- membered heterocycle, aryl or heteroaryl;

Het is a 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl),
 -O-(4- to 6-membered heterocycle), -O-(C_1 - C_4 alkyl) wherein when R^1 is not H, -CN
 or halogen, R^1 is optionally substituted with one or more halogens;
 each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;
 R^3 is H or F;
 R^{11} is H or -CH₃;
 R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle;
 R^{22} is H, halogen, or C_1 - C_2 alkyl; and
 R^{35} is -C(O) R^{351} , -C(O)NHR³⁵¹, C(O)OR³⁵¹ or S(O)₂R³⁵¹ wherein R^{351} is C_1 - C_6 alkyl, C_3 -
 C_5 cycloalkyl, 4- to 6- membered heterocycle, aryl or heteroaryl.

As noted above, each of the C_1 - C_2 alkyl, C_1 - C_4 alkyl, C_1 - C_6 alkyl, C_3 - C_5
 cycloalkyl, 4- to 6-membered heterocycle, 5- to 6-membered heteroaryl, aryl and heteroaryl
 moieties may be optionally substituted. Accordingly, the present disclosure provides for
 compounds of Structure V wherein:



L^2 is -NHR³⁵ or -C(O)NHR³⁵¹, wherein R^{351} is optionally substituted C_1 - C_6 alkyl,
 optionally substituted C_3 - C_5 cycloalkyl, optionally substituted 4- to 6-membered
 heterocycle, optionally substituted aryl or optionally substituted heteroaryl;
 Het is an optionally substituted 5- to 6-membered heteroaryl;
 R^1 is H, -CN, halogen, optionally substituted C_1 - C_4 alkyl, -O-(optionally substituted C_3 -
 C_5 cycloalkyl), -O-(optionally substituted 4- to 6-membered heterocycle) or -O-

(optionally substituted C₁-C₄ alkyl), wherein when R¹ is not H, -CN or halogen, R¹ is optionally substituted with one or more halogens;

each R² is independently hydrogen, halogen or optionally substituted C₁-C₄ alkyl;

R³ is H or F;


5 R¹¹ is H or -CH₃;

R²¹ is H, halogen, optionally substituted C₁-C₄ alkyl, optionally substituted C₃-C₅ cycloalkyl or optionally substituted 4- to 6-membered heterocycle;

R²² is H, halogen, or optionally substituted C₁-C₂ alkyl; and

10 R³⁵ is -C(O)R³⁵¹, -C(O)NHR³⁵¹, -C(O)OR³⁵¹ or -S(O)₂R³⁵¹, wherein R³⁵¹ is optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₅ cycloalkyl, optionally substituted 4- to 6-membered heterocycle, optionally substituted aryl or optionally substituted heteroaryl.


In some embodiments, the present disclosure provides for compounds of Structure V



15 In some embodiments, the present disclosure provides for compounds of Structure V wherein L² is -NHR³⁵.

In some embodiments, the present disclosure provides for compounds of Structure V wherein L² is -C(O)NHR³⁵¹.

In various aspects, the present disclosure provides for compounds of Structure VI:




wherein:

each W, X, Y and Z is independently -N- or -CR²⁶- with the proviso that not more than 2 of W, X, Y and Z are -N-;

25 each R²⁶ is independently H, C₁-C₄ alkyl, -O-(C₁-C₄ alkyl), -N(R²⁷)₂, -S(O)₂-(C₁-C₄ alkyl), or -C(O)-(C₁-C₄ alkyl);

each R^{27} is independently H or C_1 - C_4 alkyl or both R^{27} are C_1 - C_4 alkyl and join to form a 3- to 6-membered ring together with the N to which they are attached and wherein the ring optionally includes one oxygen atom as one of the members of the ring;



Het is a 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered heterocycle), -O-(C_1 - C_4 alkyl) wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;

R^3 is H or F;

R^{11} is H or -CH₃;

R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or a 4- to 6-membered heterocycle; and


R^{22} is H, halogen or C_1 - C_2 alkyl.

As noted above, each of the C_1 - C_2 alkyl, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl, 4- to 6-membered heterocycle and 5- to 6-membered heteroaryl moieties may be optionally substituted. Accordingly, the present disclosure provides for compounds of Structure VI wherein:

each W, X, Y and Z is independently -N- or -CR²⁶- with the proviso that not more than 2 of W, X, Y and Z are -N-;

R^{26} is H, optionally substituted C_1 - C_4 alkyl, -O-(optionally substituted C_1 - C_4 alkyl), -N(R^{27})₂, -S(O)₂-(optionally substituted C_1 - C_4 alkyl) or -C(O)-(optionally substituted C_1 - C_4 alkyl);

each R^{27} is independently H or optionally substituted C_1 - C_4 alkyl or both R^{27} are optionally substituted C_1 - C_4 alkyl and join to form an optionally substituted 3- to 6-membered ring together with the N to which they are attached and wherein the ring optionally includes one oxygen atom as one of the members of the ring;



Het is an optionally substituted 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, optionally substituted C_1 - C_4 alkyl, -O-(optionally substituted C_3 - C_5 cycloalkyl), -O-(optionally substituted 4- to 6-membered heterocycle) or -O-(optionally substituted C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or optionally substituted C_1 - C_4 alkyl;


R^3 is H or F;

R^{11} is H or - CH_3 ;

R^{21} is H, halogen, optionally substituted C_1 - C_4 alkyl, optionally substituted C_3 - C_5 cycloalkyl or an optionally substituted 4- to 6-membered heterocycle; and

R^{22} is H, halogen or optionally substituted C_1 - C_2 alkyl.

In some embodiments, the present disclosure provides for compounds of Structure VI



wherein Ar is

In some embodiments, the present disclosure provides for compounds of Structure VI wherein Y is - CR^{26} - wherein R^{26} is - $N(R^{27})_2$.



In some embodiments, the present disclosure provides for compounds of Structure VI wherein X is -N-.

Synthesis of Compounds


Also described herein are methods of synthesizing the compounds of the present disclosure. Compounds of the present disclosure can be synthesized according to the synthetic schemes provided below.

Scheme 1 provides methods useful for synthesizing the L moiety of Structures I-IV.


In Scheme 1, each of m' and n' are independently 1 or 2. Ar is as defined in Structures I-V.

Scheme 1


Scheme 2 provides methods useful for preparing the R^{21} - R^{22} moiety in each of Structures I-IV.

5 *Scheme 2*


Schemes 3-5 provide methods useful in the synthesis of compounds of Structure III.

Scheme 3

Scheme 4




Scheme 5




5 Schemes 6-7 provide methods useful in the synthesis of compounds of Structure II.

Scheme 6




Scheme 7




Scheme 8 provides methods useful in the synthesis of compounds of Structure IIIb.

Scheme 8



Scheme 9 provides methods useful in the synthesis of compounds of Structure I.

Scheme 9



Additional methods for producing particular compounds according to the present disclosure are provided in the Examples. One skilled in the art will recognize that other compounds of structures can be made by modifications to the specifically disclosed schemes employing methods known to those of skill in the art. Additional examples can be found in **TABLE 26**.

Many such techniques are well known in the art. However, many of the known techniques are elaborated in *Compendium of Organic Synthetic Methods* (Vol. 1, 1971; Vol. 2, 1974; Vol. 3, 1977; Vol. 4, 1980; Vol. 5, 1984; and Vol. 6 as well as *March in Advanced Organic Chemistry* (1985); *Comprehensive Organic Synthesis. Selectivity, Strategy & Efficiency in Modern Organic Chemistry*. In 9 Volumes (1993); *Advanced Organic Chemistry Part B: Reactions and Synthesis*, Second Edition (1983); *Advanced Organic Chemistry, Reactions, Mechanisms, and Structure*, Second Edition (1977); *Protecting Groups in Organic Synthesis*, Second Edition; and *Comprehensive Organic Transformations* (1999).

Antiviral Methods of Treatment

In various aspects, the present disclosure provides methods for treating viral infection in a subject, the method comprising administering to a subject in need of such treatment an effective amount of a compound of Structures I, II, III, IIIb, IIIc, IV, V and VI or as provided in **TABLE 26**.

In various aspects, the disclosure provides methods for treating a viral infection, the method comprising administering the compounds of the present disclosure to a subject in need thereof the agent.

In various aspects, the present disclosure provides methods for treating hepatitis C infection by administering to the subject one or more compounds disclosed herein.

In certain aspects the compounds of the present disclosure may be used in combination with other antiviral treatments in the treating of viral infection.

5 In various aspects the compounds of the present disclosure can be used for the treatment of infection of an animal subject, such as a human.

Anticancer Activity

In various aspects, the present disclosure provides methods for treating cancer in subject, the method comprising administering to a subject in need of such treatment an effective amount of a compound of Structures I, II, III, IIIb, IIIc, IV, V and VI or as provided
10 in **TABLE 26**. In further aspects, compounds having Structures I, II, III, IIIb, IIIc, IV, V and VI or as provided in **TABLE 26** can be used for the manufacture of a medicament for treating cancer.

In certain aspects, the present disclosure provides a method for inhibiting tumor cell
15 growth in a subject, the method comprising administering to a subject in need of such treatment an effective amount of a compound of Structure I, II, III, IIIb, IIIc, IV, V and VI or as provided in **TABLE 26**. In further aspects, the tumor can be derived from breast, lung, thyroid, lymph node, kidney, ureter, bladder, ovary, teste, prostate, bone, skeletal muscle, bone marrow, stomach, esophagus, small bowel, colon, rectum, pancreas, liver, smooth
20 muscle, brain, spinal cord, nerves, ear, eye, nasopharynx, oropharynx, salivary gland, or heart tissue. In certain aspects, the present compounds can be administered concurrently with one or more additional anti-cancer treatments.

In certain aspects, the present disclosure provides a method for treating pancreatic cancer in a subject, the method comprising administering to a subject in need of such
25 treatment an effective amount of a compound of Structure I, II, III, IIIb, IIIc, IV, V and VI or as provided in **TABLE 26**.

In certain aspects, the present disclosure provides for a method of treating colon cancer in a subject, the method comprising administering to a subject in need of such treatment an effective amount of a compound of Structure I, II, III, IIIb, IIIc, IV, V and VI or
30 as provided in **TABLE 26**.

Rapidly proliferating cancer cells activate the fatty acid synthesis pathway to supply the high levels of lipids needed for membrane assembly and oxidative metabolism. (Flavin, R. *et al.* (2010) *Future Oncology*. 6(4):551-562) Inhibitors of fatty acid synthesis have demonstrated *in vivo* activity in preclinical cancer models. (Orita, H. *et al.* (2007) *Clinical Cancer Research*. 13(23):7139-7145 and Puig, T. *et al.* (2011) *Breast Cancer Research*, 13(6):R131) Additionally, fatty acid synthesis supports new blood vessel formation and inhibitors of this pathway have activity in *in vitro* models of angiogenesis. (Browne, C.D., *et al.* (2006) *The FASEB Journal*, 20(12):2027-2035).

Utility in Metabolic Disorders

In various aspects, the compounds of the present disclosure have utility in the treating of metabolic diseases. FASN has been demonstrated to be involved in regulation of glucose, lipids and cholesterol metabolism. Mice with a liver-specific inactivation of FASN have normal physiology unless fed a zero-fat diet, in which case they develop hypoglycemia and fatty liver, both of which are reversed with dietary fat. (Chakravarthy, M. V., *et al.* (2005) *Cell Metabolism* 1:309-322). *Db/+* mice fed a high fructose diet exhibit reduced liver triglyceride levels and improved insulin sensitivity when treated for 28 days with platensimycin, a covalent inhibitor of FASN. (Wu, M. *et al.* (2011) *PNAS* 108(13):5378-5383). Ambient glucose levels are also reduced in *db/db* mice following treatment with platensimycin. These results provide evidence that inhibiting FASN can yield therapeutically relevant benefits in animal models of diabetes and related metabolic disorders. Thus the disclosed FASN inhibitors are useful in the treatment of disorders characterized by dysregulation in these systems. Without limitation, examples include steatosis and diabetes.

Pharmaceutical Compositions, Formulations, Routes of Administration, and Effective Doses

Also provided herein are pharmaceutical compositions comprising the compounds of the present disclosure.

In various aspects, the present disclosure provides pharmaceutical compositions comprising any one of the compounds of Structures I, II, III, IIIb, IIIc, IV, V and VI and a pharmaceutically acceptable carrier, excipient, or diluent.

In certain aspects, the present disclosure provides pharmaceutical compositions comprising any one of the compounds of **TABLE 26** and a pharmaceutically acceptable carrier, excipient, or diluent.

Yet another aspect of the present invention relates to formulations, routes of
5 administration and effective doses for pharmaceutical compositions comprising an agent or combination of agents of the instant invention. Such pharmaceutical compositions can be used to treat viral infections as described above.

Compounds of the invention can be administered as pharmaceutical formulations including those suitable for oral (including buccal and sub-lingual), rectal, nasal, topical,
10 transdermal patch, pulmonary, vaginal, suppository, or parenteral (including intramuscular, intraarterial, intrathecal, intradermal, intraperitoneal, subcutaneous and intravenous) administration or in a form suitable for administration by aerosolization, inhalation or insufflation. General information on drug delivery systems can be found in Ansel et al., *Pharmaceutical Dosage Forms and Drug Delivery Systems* (Lippencott Williams & Wilkins,
15 Baltimore Md. (1999).

In various aspects, the pharmaceutical composition includes carriers and excipients (including but not limited to buffers, carbohydrates, mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, bacteriostats, chelating agents, suspending agents, thickening agents and/or preservatives), water, oils including those of petroleum, animal,
20 vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like, saline solutions, aqueous dextrose and glycerol solutions, flavoring agents, coloring agents, detackifiers and other acceptable additives, adjuvants, or binders, other pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH buffering agents, tonicity adjusting agents, emulsifying agents,
25 wetting agents and the like. Examples of excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. In another aspect, the pharmaceutical preparation is substantially free of preservatives. In another aspect, the pharmaceutical preparation can contain at least one preservative. General
30 methodology on pharmaceutical dosage forms is found in Ansel et al., *Pharmaceutical Dosage Forms and Drug Delivery Systems* (Lippencott Williams & Wilkins, Baltimore Md.

(1999)). It will be recognized that, while any suitable carrier known to those of ordinary skill in the art can be employed to administer the compositions of this invention, the type of carrier will vary depending on the mode of administration.

Compounds can also be encapsulated within liposomes using well-known technology.

5 Biodegradable microspheres can also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Pat. Nos. 4,897,268; 5,075,109; 5,928,647; 5,811,128; 5,820,883; 5,853,763; 5,814,344 and 5,942,252.

The compound can be administered in liposomes or microspheres (or microparticles).

10 Methods for preparing liposomes and microspheres for administration to a patient are well known to those of skill in the art. U.S. Pat. No. 4,789,734, the contents of which are hereby incorporated by reference, describes methods for encapsulating biological materials in liposomes. Essentially, the material is dissolved in an aqueous solution, the appropriate phospholipids and lipids added, along with surfactants if required, and the material dialyzed
15 or sonicated, as necessary. A review of known methods is provided by G. Gregoriadis, Chapter 14, "Liposomes," *Drug Carriers in Biology and Medicine*, pp. 2.sup.87-341 (Academic Press, 1979).

Microspheres formed of polymers or proteins are well known to those skilled in the art, and can be tailored for passage through the gastrointestinal tract directly into the blood
20 stream. Alternatively, the compound can be incorporated and the microspheres, or composite of microspheres, implanted for slow release over a period of time ranging from days to months. See, for example, U.S. Pat. Nos. 4,906,474, 4,925,673 and 3,625,214, and Jain, TIPS 19:155-157 (1998), the contents of which are hereby incorporated by reference.

The concentration of drug can be adjusted, the pH of the solution buffered and the
25 isotonicity adjusted to be compatible with intravenous injection, as is well known in the art.

The compounds of the invention can be formulated as a sterile solution or suspension, in suitable vehicles, well known in the art. The pharmaceutical compositions can be sterilized by conventional, well-known sterilization techniques, or can be sterile filtered. The resulting aqueous solutions can be packaged for use as is, or lyophilized, the lyophilized preparation
30 being combined with a sterile solution prior to administration. Suitable formulations and additional carriers are described in Remington *The Science and Practice of Pharmacy* (20th

Ed., Lippincott Williams & Wilkins, Baltimore MD), the teachings of which are incorporated by reference in their entirety herein.

Typical salts are those of the inorganic ions, such as, for example, sodium, potassium, calcium, magnesium ions, and the like. Such salts include salts with inorganic or organic acids, such as hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulfuric acid, methanesulfonic acid, p-toluenesulfonic acid, acetic acid, fumaric acid, succinic acid, lactic acid, mandelic acid, malic acid, citric acid, tartaric acid or maleic acid. In addition, if the agent(s) contain a carboxy group or other acidic group, it can be converted into a pharmaceutically acceptable addition salt with inorganic or organic bases. Examples of suitable bases include sodium hydroxide, potassium hydroxide, ammonia, cyclohexylamine, dicyclohexyl-amine, ethanolamine, diethanolamine, triethanolamine, and the like.

The agent(s) (or pharmaceutically acceptable salts can be administered per se or in the form of a pharmaceutical composition wherein the active agent(s) is in an admixture or mixture with one or more pharmaceutically acceptable carriers.


EXAMPLES

EXAMPLE 1

Synthesis of Compounds of the Present Disclosure


General: All reactions and manipulations described were carried out in well ventilated fume-hoods. Operations and reactions carried out at elevated or reduced pressure were carried out behind blast shields. Abbreviations: ACN, acetonitrile; AcOH, acetic acid; AIBN, azobisisobutyronitrile; BF₃-Et₂O, boron trifluoride diethyl etherate; (Boc)₂O, di-*tert*-butyl dicarbonate; BuLi, butyl lithium; CDI, 1,1'-Carbonyldiimidazole; DBU, 1,8-Diazabicyclo[5.4.0]undec-7-ene; DCE, 1,2-dichloroethane; DCM, dichloromethane or methylene chloride; DIEA, N,N-Diisopropylethylamine; DMA, N,N-dimethylacetamide; DMAP, 4-dimethylaminopyridine; DME, 1,2-dimethoxyethane; DMEDA - *N,N'*-dimethylethylenediamine; DMF, N,N-dimethylformamide; DMSO, dimethylsulfoxide; DPPP, 1,3-bis(diphenylphosphino)propane; EDC, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide; EDCI, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride; EtOAc, ethyl acetate; EtOH, Ethanol; HATU, 2-(1H-7-Azabenzotriazol-1-

yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate; HBTU, O-Benzotriazole-N,N,N',N'-tetramethyl-uronium-hexafluoro-phosphate or 2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethylaminium hexafluorophosphate; HMPA, hexamethylphosphoramide; HOAc, acetic acid; HOBT, 1-Hydroxybenzotriazole; LDA, lithium diisopropylamine; m-CPBA, 3-chloroperbenzoic acid; MeOH, methanol; MsCl, methanesulfonyl chloride; MsOH, methanesulfonic acid; NaHMDS, sodium hexamethyldisilazane, NBS, N-bromosuccinimide; NCS, N-chlorosuccinimide; NIS, N-iodosuccinimide; Pd(dppf)Cl₂, [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II); PE, petroleum ether; PPA, polyphosphoric acid; PTAT, phenyltrimethylammonium tribromide; PTSA, p-toluenesulfonic acid; Py, pyridine; Pyr, pyridine; TBAF, tetrabutylammonium fluoride; TEA, triethylamine; TFA, trifluoroacetic acid; TFAA, trifluoroacetic anhydride; THF, tetrahydrofuran; TMSCl, chlorotrimethylsilane; TMSCN, trimethylsilyl cyanide; TsOH, p-toluenesulfonic acid.




Compound 1.1. *tert*-Butyl 4-(4-cyanophenyl)piperidine-1-carboxylate. Into a 500-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a suspension of Zn (21.6 g, 330 mmol) in DMA (53 mL). A 7:5 v/v mixture of TMSCl/1,2-dibromoethane (5.8 mL) was added to the mixture drop-wise at a rate to maintain the temperature below 65 °C. The mixture was stirred for an additional 10 min, then a solution of *tert*-butyl 4-iodopiperidine-1-carboxylate (68.7 g, 220 mmol) in DMA (122 mL) was added drop-wise at 40-45 °C and the mixture was stirred at the same temperature for 30 min. The mixture was cooled to room temperature and stirring was ceased to allow for the zinc powder to settle. Into another 500-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a mixture of 4-bromobenzonitrile (20 g, 110 mmol), CuI (2.1 g, 11 mmol), Pd(dppf)Cl₂ (4.51 g, 5.5 mmol) and DMA (100 mL). The freshly prepared zinc reagent solution was decanted into an addition funnel and added drop-wise to the mixture at room temperature. The resulting mixture was stirred at 85 °C for 4 h, then cooled to 20 °C and diluted with methyl *tert*-butyl ether (500 mL) and carefully quenched with 1 M ammonium chloride (500 mL). The mixture


was stirred at room temperature for 30 min and then filtered to remove solids. The organic layer was washed with saturated aqueous ammonium chloride (100 mL), followed by brine (3 x 100 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:10) as the eluent to yield the title compound as a brown oil (20g, crude) and used in the next step without additional purification.



Compound 1.2. 4-(Piperidin-4-yl)benzonitrile hydrochloride. Into a 500-mL three neck round-bottom flask, was placed a solution of tert-butyl 4-(4-cyanophenyl)piperidine-1-carboxylate (compound **1.1**, 20 g, crude) in ethyl acetate (200 mL). Hydrogen chloride (gas) was introduced to the solution and the resulting mixture was stirred for 30 min at room temperature. The solids were collected by filtration, then washed with ethyl acetate (100 mL) and ether (100 mL) to yield the title compound as a white solid (14g, 57% over two steps).




Compound 1.3. 5-Iodo-2,4-dimethylbenzoic acid. A solution of 2,4-dimethylbenzoic acid (20.0 g, 133 mmol), sodium periodate (14.27 g, 66.72 mmol), iodine (37.25 g, 146.8 mmol), and sulfuric acid (1.96 g, 20.0 mmol) in acetic acid (150 mL) was stirred at 110 °C for 6 h. The mixture was allowed to cool to ambient temperature then carefully diluted into water (1.2 L). To this mixture was carefully added saturated aqueous Na₂S₂O₃ (800 mL). The resulting solids were collected by filtration, then dissolved in ethyl acetate (1.2 L) and washed with saturated aqueous Na₂S₂O₃ (300 mL) followed by brine (400 mL). The organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude residue was re-crystallized from ethanol:H₂O (2:1) to yield the title compound as a white solid (30 g, 82%).




Compound 1.4. 2,4-Dimethyl-5-propionylbenzoic acid. Into a 100-mL autoclave (30 atm), was placed a solution of 5-iodo-2,4-dimethylbenzoic acid (compound **1.3**, 2.00 g,


7.24 mmol) in ethylene glycol dimethyl ether (20 mL). Triphenylphosphine (190 mg, 0.73 mmol), Pd(PPh₃)₂Cl₂ (500 mg, 0.71 mmol) and diethylaluminum chloride (2M, 10.8 mL, 21.6 mmol) were added to the reaction mixture. The resulting mixture was stirred under pressure with carbon monoxide (gas, 30 atm) at 80 °C for 15 h. (**CAUTION:** Highly toxic gas at high pressure. All necessary safety precautions were performed). After cooling to room temperature, the mixture was carefully purged, then quenched with 20 mL of water. Aqueous HCl (2M) was added carefully to adjust the pH to 5-6 and the aqueous layer was extracted with ethyl acetate (100 mL). The combined organic layers were washed with brine (2 x 100 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:40-1:20) as the eluent to yield the title compound as a light yellow solid (1.2 g, 80%).



Compound 1.5. Methyl 2,4-dimethyl-5-propionylbenzoate. To a 100-mL round-bottom flask, was carefully added 2,4-dimethyl-5-propionylbenzoic acid (compound **1.4**, 1.2 g, 5.8 mmol), sulfuric acid (1.0 mL, 19 mmol) and methanol (30 mL). The resulting solution was stirred at 70 °C for 5 h, then concentrated under reduced pressure. The residue was carefully diluted with H₂O (50 mL) and extracted ethyl acetate (100 mL). The organic layer was washed with H₂O (2 x 100 mL) and brine (100 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:100) as the eluent to yield the title compound as a light yellow solid (0.90 g, 70%).




Compound 1.6. Methyl 5-(2-bromopropanoyl)-2,4-dimethylbenzoate. Into a 100-mL round-bottom flask, was placed a solution of methyl 2,4-dimethyl-5-propionylbenzoate (compound **1.5**, 600 mg, 2.72 mmol) in chloroform (20 mL). Bromine (154 µL, 3.00 mmol) was added and the resulting solution was stirred at 20 °C for 2 h. The mixture was concentrated under reduced pressure to yield the title compound as a yellow oil (1.0 g, crude), which was used in the next step without further purification.



Compound 1.7. Methyl 5-(2,4-dimethyl-1*H*-imidazol-5-yl)-2,4-dimethylbenzoate.


Into a round-bottom flask, was placed a solution of methyl 5-(2-bromopropanoyl)-2,4-dimethylbenzoate (compound **1.6**, 400 mg, 1.34 mmol) in acetonitrile (30 mL).

- 5 Acetimidamide hydrochloride (260 mg, 2.75 mmol) and potassium carbonate (550 mg, 3.99 mmol) were added and the resulting mixture was stirred at 80 °C for 15 h. After cooling to room temperature, the mixture was concentrated under reduced pressure and the residue was diluted with ethyl acetate (50 mL) and washed with brine (2 x 25 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel
- 10 column chromatography with ethyl acetate/petroleum ether (1:1) to ethyl acetate as the eluent to yield the title compound as a yellow oil (0.20 g, 58%).




Compound 1.8. 5-(2,4-Dimethyl-1*H*-imidazol-5-yl)-2,4-dimethylbenzoic acid.

- Into a 50-mL round-bottom flask, were placed methyl 5-(2,4-dimethyl-1*H*-imidazol-5-yl)-2,4-dimethylbenzoate (compound **1.7**, 300 mg, 1.16 mmol) and sodium hydroxide (465 mg, 11.6 mmol) in methanol (20 mL) and H₂O (5 mL). The resulting solution was stirred at 55 °C
- 15 for 4 h, then after cooling to room temperature, aqueous HCl (2 M) was added to adjust the pH to 5. The resulting mixture was concentrated under reduced pressure and the residue was dissolved in methanol (5 mL). The solids were removed by filtration and the filtrate was
- 20 concentrated under reduced pressure to yield the title compound as a light yellow solid (280 mg, crude), which was used in the next step without further purification.




Compound 1. 4-(1-(5-(2,4-Dimethyl-1*H*-imidazol-5-yl)-2,4-dimethylbenzoyl)piperidin-4-yl)benzonitrile. A mixture of 5-(2,4-dimethyl-1*H*-imidazol-5-

yl)-2,4-dimethylbenzoic acid (compound **1.8**, 280 mg, 1.15 mmol), EDC•HCl (330 mg, 1.72 mmol), 4-dimethylaminopyridine (420 mg, 3.44 mmol), and HOBT (180 mg, 1.33 mmol) in *N,N*-dimethylformamide (6 mL) was stirred at room temperature. After 5 min, 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**, 230 mg, 1.03 mmol) was added and the resulting solution was stirred at room temperature for 15 h, then quenched with ice water (20 mL). The aqueous was extracted with ethyl acetate (50 mL) and the combined organics was washed with brine (2 x 50 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-001(SHIMADZU)): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150 mm; mobile phase, water with 0.05% TFA and CH₃CN (22% CH₃CN up to 37% in 7 min, up to 100% in 2 min, down to 22% in 1 min); Detector, Waters 2489, 254 & 220nm. The fractions containing pure compound were combined and lyophilized to yield the title compound as a white solid (214 mg, 50%). *m/z* (ES⁺) 413 (M+H)⁺. ¹H NMR (300 MHz, CD₃OD): δ 7.68 (d, *J* = 7.8 Hz, 2H), 7.47 (d, *J* = 8.4 Hz, 2H), 7.36 (br s, 1H), 7.27 & 7.16 (2 singlets, rotamers, Ar-H, 1H), ~4.9-4.82 (m, 1H partially obscured by water peak), 3.72-3.55 (m, 1H), ~3.35-3.20 (m, 1H partially overlapped with methanol solvent peak), 3.08-2.92 (m, 2H), 2.65(s, 3H), 2.44 & 2.34 (2 singlets, rotamers, Ar-CH₃, 3H), 2.29 (s, 3H), 2.22 (s, 3H), 2.10-1.96 (m, 1H), 1.93-1.53 (m, 3H).




Compounds 2.1 and 2.2. Methyl 5-(2-(methoxymethyl)-4-methyl-1*H*-imidazol-5-yl)-2,4-dimethylbenzoate and methyl 5-(2-(methoxymethyl)-4-methyloxazol-5-yl)-2,4-dimethylbenzoate. Into a 100-mL round-bottom flask, was placed a mixture of methyl 5-(2-bromopropanoyl)-2,4-dimethylbenzoate (compound **1.6**, 600 mg, 2.01 mmol), 2-methoxyethanimidamide hydrochloride (510 mg, 4.09 mmol), potassium carbonate (840 mg, 6.08 mmol) and acetonitrile (30 mL). The resulting mixture was stirred at 80 °C overnight, then cooled to room temperature and concentrated under reduced pressure. The residue was diluted with H₂O (50 mL) and extracted with ethyl acetate (100 mL). The organics was washed brine (2 x 50 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:2) as the eluent to yield methyl 5-(2-(methoxymethyl)-4-methyl-1*H*-imidazol-5-yl)-2,4-

dimethylbenzoate (compound **2.1**) (0.11 g, 19%) and methyl 5-(2-(methoxymethyl)-4-methyloxazol-5-yl)-2,4-dimethylbenzoate (compound **2.2**) (0.30 g, 52%), both as a yellow oils.




Compound 2.3. 5-(2-(Methoxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid. Into a 50-mL round-bottom flask, was placed a solution of methyl 5-(2-(methoxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **2.1**, 150 mg, 0.52 mmol) in methanol (15 mL) and H₂O (5 mL). Sodium hydroxide (280 mg, 7.00 mmol) was added and the resulting solution was stirred at 40 °C for 15 h. After cooled to room temperature, the solution was adjusted to pH 5 with aqueous HCl (2 M) and the resulting mixture was concentrated under reduced pressure. The residue was dissolved in methanol (5 mL) and the solids were filtered off. The filtrate was concentrated under reduced pressure to yield the title compound as a light yellow solid (140 mg, crude), which was used in the next step without further purification.




Compound 2. 4-(1-(5-(2-(Methoxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoyl)piperidin-4-yl)benzonitrile. A mixture of 5-(2-(methoxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **2.3**, 140 mg, 0.51 mmol), EDC•HCl (150 mg, 0.78 mmol), 4-dimethylaminopyridine (190 mg, 1.56 mmol), and HOBt (80 mg, 0.59 mmol) in *N,N*-dimethylformamide (6 mL). The mixture was stirred at room temperature, then after 15 min, 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**, 100 mg, 0.45 mmol) was added. The resulting solution was stirred for 15 h at room temperature, then quenched with 20 mL of H₂O. The aqueous was extracted with ethyl acetate (40 mL) and the combined organics was washed with brine (2 x 30 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-001(SHIMADZU)): Column,


SunFire Prep C18 OBD Column, 5 μ m, 19*150 mm; mobile phase, water with 0.05% TFA and CH₃CN (23% CH₃CN up to 38% in 7 min, up to 100% in 2 min, down to 23% in 1 min); Detector, Waters 2489, 254 & 220 nm. The fractions containing pure compound were combined and lyophilized to yield the title compound as a white solid (70.4 mg, 35%).




Compound 3. 4-(1-(5-(2-(Hydroxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoyl)piperidin-4-yl)benzonitrile. Into a 50-mL 3-necked round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a mixture of 4-(1-(5-(2-(methoxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoyl)piperidin-4-yl)benzonitrile (compound **2**, 30 mg, 0.068 mmol), NaI (20 mg, 0.13 mmol), 15-crown-5 (30 mg, 0.14 mmol) and dichloromethane (10 mL). The mixture was cooled to -30 °C and boron tribromide (70 mg, 0.28 mmol) was added. The resulting mixture was allowed to warm to room temperature and stirred for 15 h. The reaction mixture was carefully quenched by the addition of saturated aqueous sodium bicarbonate (10 mL) and the resulting mixture was extracted dichloromethane (2 x 20 mL). The combined organics was washed with saturated aqueous Na₂S₂O₃ (2 x 20 mL), then dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude product was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-001(SHIMADZU)): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150mm; mobile phase, water with 0.05% TFA and CH₃CN (21% CH₃CN up to 35% in 8 min, up to 100% in 2 min, down to 21% in 1 min); Detector, Waters 2489, 254&220nm. The fractions containing pure compound were combined and lyophilized to yield the title compound as a white solid (5.0 mg, 17%). *m/z* (ES⁺) 429 (M+H)⁺. ¹H NMR (400 MHz, CD₃OD): δ 7.74-7.65 (m, 2), 7.48 (d, J = 8.0 Hz, 2H), 7.38 (d, J = 5.6 Hz, 1H), 7.29 & 7.16 (2 singlets, rotamers, Ar-H, 1H), ~4.9 (1H obscured by water peak), 3.64 (app t, J = 15.0 Hz, 1H), ~3.35-3.21 (m, 1H partially overlapped with methanol solvent peak), 3.09-2.93 (m, 1H), 2.45 & 2.34 (2 singlets, rotamers, Ar-CH₃, 3H), 2.29 (s, 3H), 2.24 (s, 3H), 2.09-1.97 (m, 1H), 1.92-1.71 (m, 2H), 1.70-1.55 (m, 1H).



Compound 4.1. 3-Formyl-4-methylbenzoic acid. To a stirred solution of 3-bromo-4-methylbenzoic acid (2.14 g, 10.0 mmol) in tetrahydrofuran (30 mL) under nitrogen at -78 °C was added n-BuLi (10 mL, 2.5 M in THF, 25 mmol) drop-wise. The mixture was stirred for 1 h below -70 °C, then DMF (5 mL) was slowly added. The resulting solution was slowly warmed to room temperature and stirred for 1 h, then carefully quenched by slow addition of water (50 mL). The pH was adjusted to ~3-4 using aqueous HCl (6 M) and the resulting mixture was extracted with ethyl acetate (2 x 50 mL). The combined organics was dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield the title compound as a yellow solid (1.6 g, 98%).




Compound 4.2. 3-(1-Hydroxybut-3-en-1-yl)-4-methylbenzoic acid. Into a 100-mL 3-necked round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 3-formyl-4-methylbenzoic acid (compound 4.1, 2.00 g, 12.2 mmol) in tetrahydrofuran (50 mL). The mixture was cooled to -10 to 0 °C then allylmagnesium bromide (1M in Et₂O, 24.4 mL, 24.4 mmol) was added drop-wise. The resulting mixture was stirred for 1 hr at -10-0 °C, then carefully quenched with saturated aqueous NH₄Cl (50 mL) and diluted with ethyl acetate (200 mL). The organic layer was washed with saturated aqueous NH₄Cl (80 mL) and brine (2 x 80 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield the title compound as a light red solid (2.4 g, crude), which was used in the next step without further purification.




Compound 4.3. Methyl 3-(1-hydroxybut-3-en-1-yl)-4-methylbenzoate. Into a 100-mL round-bottom flask, was added a solution of 3-(1-hydroxybut-3-en-1-yl)-4-methylbenzoic acid (compound 4.2, 1.4 g, 6.79 mmol) in *N,N*-dimethylformamide (20 mL). Sodium bicarbonate (1.14 g, 13.6 mmol) and methyl iodide (0.847 mL, 13.6 mmol) were added and the resulting mixture was stirred overnight at 25 °C. The reaction was quenched with

saturated aqueous $\text{Na}_2\text{S}_2\text{O}_3$ (50 mL) and diluted with EtOAc (150 mL). The organic layer was washed with brine (3 x 50 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:3) as the eluent to yield the title compound as a light yellow oil (800 mg, 53%).




Compound 4.4. Methyl 3-(2-cyclopropyl-1-hydroxyethyl)-4-methylbenzoate. Into

a 100-mL 3-necked round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of methyl 3-(1-hydroxybut-3-en-1-yl)-4-methylbenzoate (compound **4.3**, 50 mg, 0.23 mmol). Diethylzinc (1 M in toluene) (3.45 mL, 3.45 mmol) in toluene (10 mL) was added and the mixture was cooled to 0-5 °C, then diiodomethane (924 mg, 3.45 mmol) was added drop-wise. The resulting mixture was stirred for 2 h at room temperature, then carefully quenched with 1 M aqueous HCl (50 mL) and diluted with MTBE (50 mL). The aqueous phase was extracted with MTBE (3 x 20 mL) and the combined organics was washed with saturated sodium bicarbonate (2 x 20 mL), brine (2 x 20 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:3) as the eluent to yield the title compound as a yellow oil (40 mg, 74%).




Compound 4.5. Methyl 3-(2-cyclopropylacetyl)-4-methylbenzoate. Into a 100-mL


round-bottom flask, was placed a solution of methyl 3-(2-cyclopropyl-1-hydroxyethyl)-4-methylbenzoate (compound **4.4**, 200 mg, 0.85 mmol) in dichloromethane (30 mL). This was followed by the addition of Dess-Martin periodinane (721 mg, 1.70 mmol) in portions at room temperature. The resulting solution was stirred for 1 h at room temperature, then quenched with saturated aqueous $\text{Na}_2\text{S}_2\text{O}_3$ (30 mL). The resulting mixture was extracted with DCM (3 x 20 mL) and the combined organics was washed with brine (2 x 20 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:5) as the eluent to yield the title compound as a yellow oil (150 mg, 75%).



Compound 4.6. Methyl 3-(2-bromo-2-cyclopropylacetyl)-4-methylbenzoate. Into a 100-mL round-bottom flask, was placed a solution of methyl 3-(2-cyclopropylacetyl)-4-methylbenzoate (compound **4.5**, 150 mg, 0.65 mmol) in chloroform (15 mL). Bromine (40 μ L, 0.78 mmol) in chloroform (2 mL) was added drop-wise to the reaction mixture. The resulting solution was stirred for 2 h at room temperature, then concentrated under reduced pressure to yield the title compound as a yellow oil (200 mg, crude), which was used in the next step without further purification.




Compound 4.7. 2-Methoxyacetimidamide hydrochloride. Into a 250-mL round-bottom flask, was placed a solution of 2-methoxyacetonitrile (6.00 g, 84.4 mmol) in methanol (60 mL). Sodium methoxide (860 mg, 15.9 mmol) was added and the mixture was stirred at room temperature for 40 h. Ammonium chloride (4.52 g, 84.5 mmol) was then added and the mixture was stirred at 40 °C for 12 h then concentrated under reduced pressure. The residue was diluted with H₂O (20 mL) and washed with ethyl acetate (2 x 20 mL). The aqueous was concentrated under reduced pressure to yield the title compound as a yellow solid (5 g, crude), which was used in the next step without further purification.




Compound 4.8 and compound 4.9. Methyl 3-(4-cyclopropyl-2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoate and methyl 3-(4-cyclopropyl-2-(methoxymethyl)oxazol-5-yl)-4-methylbenzoate. Into a 100-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed methyl 3-(2-bromo-2-cyclopropylacetyl)-4-methylbenzoate (compound **4.6**, 150 mg, 0.48 mmol), 2-methoxyacetimidamide hydrochloride (compound **4.7**, 90 mg, 0.72 mmol), potassium carbonate (200 mg, 1.44 mmol), and *N,N*-dimethylformamide (15 mL). The resulting mixture was stirred at 80 °C for 3 h, then diluted with ethyl acetate (100 mL). The mixture was washed with brine (3 x 30 mL) and water (3 x 30 mL), dried (Na₂SO₄), filtered, and

concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:2) as the eluent to yield methyl 3-(4-cyclopropyl-2-(methoxymethyl)-1*H*-imidazol-5-yl)-4-methylbenzoate (compound **4.8**) (30 mg, 21%) and methyl 3-(4-cyclopropyl-2-(methoxymethyl)oxazol-5-yl)-4-methylbenzoate (compound **4.9**) (60 mg, 41%), both as a yellow oils.



Compound 4.10. 3-(4-Cyclopropyl-2-(methoxymethyl)-1*H*-imidazol-5-yl)-4-methylbenzoic acid. Into a round-bottom flask, was placed a solution of methyl 3-(4-


cyclopropyl-2-(methoxymethyl)-1*H*-imidazol-5-yl)-4-methylbenzoate (compound **4.8**, 35 mg, 0.12 mmol) in methanol (5 mL), a solution of sodium hydroxide (9.6 mg, 0.24 mmol) in water (0.3 mL) was added to the reaction mixture. The resulting solution was stirred for 2 h at room temperature, then diluted with water (5 mL). Aqueous HCl (6 M) was carefully added to adjust the pH to 1-2 and the mixture was extracted with dichloromethane (4 x 10 mL). The combined organics were dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield the title compound as a yellow oil (30 mg, crude), which was used in the next step without further purification.




Compound 4. 4-(1-(3-(4-Cyclopropyl-2-(methoxymethyl)-1*H*-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile. Into a round-bottom flask, was placed a

solution of 3-(4-cyclopropyl-2-(methoxymethyl)-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **4.10**, 30 mg, 0.10 mmol) in dichloromethane (5 mL). HBTU (76 mg, 0.20 mmol), 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**, 35 mg, 0.15 mmol, 1.50 equiv) and triethylamine (28 μ L, 0.20 mmol) were added and the mixture was stirred overnight at room temperature. The mixture was diluted with DCM (30 mL) and washed with water (3 x 15 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-001(SHIMADZU)):

Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150 mm; mobile phase, water with 0.05% TFA and CH₃CN (25.0% CH₃CN up to 38.0% in 7 min, up to 100.0% in 3 min, down to 25.0% in 1 min); Detector, Waters 2489, 254 & 220 nm. The fractions containing pure compound were combined and lyophilized to yield the title compound as a white solid (5.9 mg, 12%). m/z (ES+) 455 (M+H)⁺. ¹H NMR (400 MHz, CDCl₃): δ 7.69 (d, J = 8.0 Hz, 2H), 7.59-7.53 (m, 2H), 7.53-7.47 (m, 3H), ~4.9-4.75 (m, 1H partially obscured by water peak), 4.72 (s, 2H), 3.99-3.85 (m, 1H), 3.52 (s, 3H), ~3.3 (m, 1H overlapped with methanol solvent peak), 3.08-2.93 (m, 2H), 2.37 (s, 3H), 2.08-1.93 (m, 1H), 1.93-1.65 (m, 4H), 1.02-0.94 (m, 2H), 0.74-0.68 (m, 2H).




Compound 5.1. *tert*-Butyl 3-(4-cyanophenyl)azetidine-1-carboxylate. A modified procedure to that described in *Org. Lett.* **2008**, *10*, 3259 was performed as follows. To a 20-mL vial was added (4-cyanophenyl)boronic acid (1.01 g, 6.87 mmol), trans-2-aminocyclohexanol hydrochloride (32 mg, 0.21 mmol), nickel (II) iodide (66 mg, 0.21 mmol) and sodium hexamethyldisilazane (1.29 g, 7.06 mmol). The system was purged with nitrogen and charged with isopropyl alcohol (7 mL). The mixture was stirred at room temperature for 10 minutes then sonicated for 1 min. While stirring, *tert*-butyl 3-iodoazetidine-1-carboxylate (1.00 g, 3.53 mmol) was added and the syringe rinsed with isopropyl alcohol (2 x 500 μ L). The suspension was stirred at 80 °C for 1 hour then concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexanes to 25% ethyl acetate) to yield the title compound as a pale yellow oil which solidified upon standing (0.832 g, 46%). m/z (ES+) 203 (M-C₄H₈+H)⁺. ¹H NMR (400 MHz, CDCl₃): δ 7.65 (d with fine str., J = 8.4 Hz, 2H), 7.43 (d with fine str., J = 8.4 Hz, 2H), 4.37 (app t, J = 8.8 Hz, 2H), 3.98-3.91 (m, 2H), 3.81-7.73 (m, 1H).




Compound 5.2. 4-(Azetidin-3-yl)benzonitrile hydrochloride. *tert*-Butyl 3-(4-cyanophenyl)azetidine-1-carboxylate (compound **5.1**, 100 mg, 0.387 mmol) was added to a

4-mL vial. HCl in dioxane (4 M, 500 μ L, 2 mmol) was added and the unsealed mixture was stirred at room temperature for 1.5 hours. The mixture was concentrated under reduced pressure and the residue was dissolved in DCM and concentrated under reduced pressure.

This was repeated with DCM twice to chase off any excess HCl to yield the title compound as a white powder (80 mg, over theory). m/z (ES⁺) 159 (M+H)⁺.




Compound 5.3. Methyl 3-iodo-4-methylbenzoate. To a solution of 3-iodo-4-methylbenzoic acid (28.0 g, 0.107 mol) in MeOH (300 mL) at 0 °C was carefully added concentrated H₂SO₄ (30 mL). The mixture was heated at 60 °C overnight, then cooled and the solvent removed under reduced pressure. The residue was carefully poured onto ice-water (200 mL) and the mixture was extracted with EtOAc (500 mL). The organics was washed with water (100 mL), saturated NaHCO₃ (100 mL), brine (100 mL), dried (MgSO₄), filtered, and concentrated to yield the title compound as a brown oil (29.0 g, 98%). ¹H NMR (400 MHz, CDCl₃) δ 8.47 (d, J = 1.7 Hz, 1H), 7.90 (dd, J = 7.9 Hz, 1.7 Hz, 1H), 7.29 (d, J = 7.9 Hz, 1H), 3.90 (s, 2H), 2.48 (s, 3H).




Compound 5.4. Methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate. A mixture of methyl 3-iodo-4-methylbenzoate (compound 5.3, 5.00 g, 18.1 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (5.20 g, 20.5 mmol), KOAc (5.33 g, 54.3 mmol) and PdCl₂(dppf)•CH₂Cl₂ (0.74 g, 0.91 mmol) in DMSO (50 mL) was degassed with argon. The mixture was then heated at 80 °C under argon overnight. The mixture was allowed to cool then partitioned between EtOAc (400 mL) and water (80 mL). The organic phase was washed with water (80 mL), saturated aqueous NaHCO₃ (80 mL), brine (80 mL), dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was purified with silica gel chromatography (hexanes:EtOAc 20:1) to yield the title compound as a white crystalline solid (3.56 g, 71%). ¹H NMR (400 MHz, CDCl₃) δ 8.41 (d,

$J = 1.9$ Hz, 1H), 7.97 (dd, $J = 8.0$ Hz, 2.0 Hz, 1H), 7.23 (d, $J = 8.0$ Hz, 1H), 3.90 (s, 3H), 2.58 (s, 3H), 1.35 (s, 12H).



Compound 5.5. 5-Iodo-2,4-dimethyl-1H-imidazole. NIS (14.0g, 62.4mmol) was


added portion-wise to a solution of 2,4-dimethyl-1H-imidazole (5.00 g, 52.0 mmol) in acetonitrile (100 mL). The mixture was heated at 80 °C for 16 hours, then cooled to room temperature. The solvent was removed under reduced pressure and the residue was partitioned between EtOAc (300 mL) and water (80 mL). The organic layer was washed with saturated sodium thiosulfate (50 mL), brine (50 mL), dried (MgSO₄) and concentrated under reduced pressure. The residue was purified with by silica gel column chromatography (hexanes:EtOAc 1:1 to 10% MeOH in EtOAc) to yield the title compound as a light yellow solid (8.56 g, 74%). m/z (ES+) 223(M+H)⁺. ¹H NMR (400 MHz, CDCl₃) δ 9.69 (br s, 1H), 2.38 (s, 3H), 2.19 (s, 3H).



Compound 5.6. Methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate.


Methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**, 3.56 g, 12.9 mmol), 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**, 3.43 g, 15.4 mmol), K₂CO₃ (5.33 g, 38.6 mmol) and PdCl₂(dppf)·CH₂Cl₂ (1.05 g, 1.29 mmol) were added to a round bottom flask. The flask was purged with argon, then dioxane (70 mL) and water (20 mL) were added and the mixture was heated at 90 °C for 16 hours. The mixture was cooled then additional K₂CO₃ (1M, 25 mL, 25.0 mmol) and catalyst PdCl₂(dppf)·CH₂Cl₂ (1.0g, 1.2 mmol) were added. The mixture was heated at 90 °C for an additional 10 hours, then cooled to room temperature and filtered through Celite®. The solvent was removed under reduced pressure and the residue was cooled to 0 °C and acidified to pH 3-4 with aqueous HCl (2 M). The acidic mixture was washed with EtOAc (150 mL) and then the aqueous material was adjusted to pH 10-11 with aqueous sodium hydroxide (2 M) and extracted with EtOAc (5 x 200 mL). The combined organic phases were dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was purified by column chromatography (DCM to 5% MeOH

in DCM) to yield the title compound as a thick brown oil (2.42 g, 77%). m/z (ES+) 245(M+H)⁺. ¹H NMR (400 MHz, CDCl₃) δ 7.89-7.87 (m, 2H), 7.31 (d with fine str, J = 8.6 Hz, 1H), 3.89 (s, 3H), 2.38 (s, 3H), 2.31 (s, 3H), 2.11 (s, 3H).



Compound 5.7. 3-(2,4-Dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid


hydrochloride. To a solution of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **5.6**, 0.66 g, 2.7 mmol) in MeOH (5 mL) was added aqueous NaOH (2 M, 4.8 mL, 9.6 mmol). The mixture was stirred at room temperature for 3 hours then the organic solvent was removed under reduced pressure. The aqueous residue was cooled to 0 °C and acidified to pH 3-4 with aqueous HCl (1 M). The mixture was concentrated to dryness and 5% methanol in DCM (20 mL) was added to the residue. The mixture was stirred at room temperature for 5 minutes and the solids (inorganic salts) were filtered from solution. The filtrate was concentrated to yield the title compound as a brown foam (0.484 g, 67%). m/z (ES+) 231(M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆) δ 14.22 (br s, 1H), 14.16 (br s, 1H), 13.11 (br s, 1H), 7.97 (dd, J = 8.0 Hz, 1.8 Hz, 1H), 7.85 (d, J = 1.8 Hz, 1H), 7.54 (d, J = 8.0 Hz, 1H), 2.58 (s, 3H), 2.31 (s, 3H), 2.15 (s, 3H).




Compound 5. 4-(1-(3-(2,4-Dimethyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile.

To a mixture of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**, 0.484 g, 2.10 mmol), 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**, 0.41 g, 2.10 mmol), HOBt (0.085 g, 7.40 mmol) and EDCI (0.603 g, 3.15 mmol) in DMF (8 mL) was added DIEA (1.09 mL, 6.3 mmol). The mixture was stirred at room temperature for 16 hours, then partitioned between EtOAc (300 mL) and water (30 mL). The organic layer was washed with brine (3 x 30 mL) and the combined aqueous phases were back extracted with EtOAc (2 x 50 mL). All organic extracts were combined, dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (EtOAc to 5% MeOH in EtOAc) to yield the title compound as a


white solid (0.35g, 45%). m/z (ES+) 371 (M+H)⁺. ¹H NMR (400 MHz, CDCl₃) δ 7.66 (d with fine str., J = 8.4 Hz, 2H), 7.54-7.48 (m, 2H), 7.44 (d with fine str., J = 8.2 Hz, 2H), 7.30 (d, J = 7.9 Hz, 1H), 4.77-4.56 (m, 2H), 4.35-4.18 (m, 2H), 3.97-3.87 (m, 1H), 2.38 (s, 3H), 2.30 (s, 3H), 2.10 (s, 3H).



Compound 6.1. Methyl 4-bromo-2-methylbenzoate. To a solution of 4-bromo-2-methylbenzoic acid (5.11 g, 23.8 mmol, 1.0 equiv) in methanol (25 mL) was added dropwise sulfuric acid (2.0 mL) over about 3 minutes (mildly exothermic). The resulting mixture was refluxed for 4 hours. After cooling to room temperature, the reaction mixture was carefully quenched into saturated aqueous NaHCO₃ (100 mL) (note – significant gas evolution) and extracted with dichloromethane (200 mL x 1 then 50 mL x 1). The combined organic phases were washed with a mixture of brine/saturated NaHCO₃ (9:1)(50 mL), dried (Na₂SO₄), and concentrated under reduced pressure to yield the title compound as a colorless oil (5.28 g, 97%). ¹H NMR (400 MHz, CDCl₃): δ 7.78 (d, J = 8.0 Hz, 1 H), 7.42 (d, J = 1.6 Hz, 1H), 7.38 (dd, J = 1.6 Hz, 1H), 3.89 (s, 3H), 2.58 (s, 3H).




Compound 6.2. Methyl 4-cyclobutyl-2-methylbenzoate. Cyclobutylzinc(II) bromide (50 mL, 0.5 M in THF, 25.0 mmol) was added to a mixture of methyl 4-bromo-2-methylbenzoate (compound **6.1**, 5.2 g, 22.7 mmol) and PdCl₂(dppf)•CH₂Cl₂ (1.85 g, 2.27 mmol). The mixture was degassed and the flask was filled with argon through a balloon. The mixture was heated at 65 °C under argon for 24 hours, then cooled to 0 °C and carefully quenched with water (10 mL). The mixture was diluted with EtOAc (200 mL) and washed with water then brine, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified using silica gel column chromatography (hexanes : EtOAc 30:1 to 20:1) to yield the title compound as a clear oil (4.1 g, 89%). ¹H NMR (400 MHz, CDCl₃) δ 7.86 (d, 1H), 7.12–7.02 (m, 2H), 3.88 (s, 3H), 3.59 – 3.48 (m, 1H), 2.59 (s, 3H), 2.35 (m, 2H), 2.22–1.96 (m, 3H), 1.86-1.84 (m, 1H).



Compound 6.3. Methyl 4-cyclobutyl-5-iodo-2-methylbenzoate. N-

Iodosuccinimide (3.52 g, 15.6 mmol) was added portion-wise to a solution of methyl 4-cyclobutyl-2-methylbenzoate (compound **6.2**, 3.2 g, 15.6 mmol) in concentrated sulfuric acid (25 mL) at 0 °C. The mixture was stirred at 0 °C for 30 min and at RT for 2 hours, upon where the mixture turned very thick. The mixture was again cooled to 0 °C and MeOH (30 mL) was slowly and carefully added. The mixture was heated at 60 °C for 2 hours. After cooling to room temperature, the solvent was removed under reduced pressure and the residue was poured onto ice water (100 mL). The mixture was extracted with EtOAc (2x).

The combined organic layers were washed with brine, then aqueous 1M NaHCO₃ (note-significant gas evolution), dried (Na₂SO₄), filtered, and concentrated. The residue was purified by silica gel column chromatography (hexanes : EtOAc 30:1 to 20:1) to yield the title compound as a light yellow oil (4.17 g, 81%). ¹H NMR (400 MHz, CDCl₃) δ 8.33 (s, 1H), 7.14 (s, 1H), 3.87 (s, 3H), 3.67 – 3.54 (m, 1H), 2.57 (s, 3H), 2.51 – 2.40 (m, 2H), 2.14 – 1.97 (m, 3H), 1.82-1.79 (m, 1H).




Compound 6.4. Methyl 5-cyano-4-cyclobutyl-2-methylbenzoate. A mixture of

methyl 4-cyclobutyl-5-iodo-2-methylbenzoate (compound **6.3**, 4.17 g, 12.6 mmol), Zn(CN)₂ (2.96 g, 25.2 mmol) and Pd(PPh₃)₄ (0.73 g, 0.63 mmol) in DMF (30 mL) was degassed and the flask was charged with argon. The mixture was heated at 100 °C under argon overnight.


After cooling to ambient temperature, the mixture was quenched with saturated aq. FeSO₄ (20 mL) and diluted with EtOAc (200 mL). The greenish solid was removed by filtration through Celite® and the filtrate was partitioned between water and EtOAc. The organics was washed with brine, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified using silica gel column chromatography (hexanes:EtOAc 30 : 1 to 20 : 1) to yield the title compound as a white solid (2.55 g, 88%). ¹H NMR (400 MHz, CDCl₃) δ

8.16 (s, 1H), 7.28 (s, 1H), 3.90 (s, 3H), 3.86-3.82 (m, 1H), 2.68 (s, 3H), 2.55 – 2.45 (m, 2H), 2.27 – 2.04 (m, 3H), 1.89-1.87 (m, 1H).



Compound 6.5. 5-Carbamoyl-4-cyclobutyl-2-methylbenzoic acid. To a 1-L round


5 bottom flask was added methyl 5-cyano-4-cyclobutyl-2-methylbenzoate (compound **6.4**, 12.00 g, 52.3 mmol) and dissolved in DMSO (100 mL). With stirring, aqueous sodium hydroxide (1 M, 260 mL, 260 mmol) was added carefully and the mixture was purged with nitrogen. The mixture was stirred at 95 °C for 13 hours and then cooled to room temperature. The solution was washed with diethyl ether (100 mL) and the basic aqueous was acidified to
 10 pH~2 by slow addition of aqueous HCl (1M) followed by aqueous H₃PO₄ (1M). The precipitated solids were filtered, and washed with water (2 x 100 mL), then dried to constant mass to yield the title compound as a white powder (11.68 g, 96%). *m/z* (ES-) 232 (M-H)⁻. ¹H
 NMR (400 MHz, DMSO-d₆): δ 12.75 (br s, 1H), 7.75 (s, 1H), 7.74 (br s, 1H), 7.34 (br s, 1H),
 7.28 (s, 1H), 3.90 (app p, *J* = 9.0 Hz, 1H), 2.56 (s, 3H), 2.32-2.20 (m, 2H), 2.16-2.02 (m, 2H),
 15 2.00-1.87 (m, 1H), 1.82-1.71 (m, 1H).




Compound 6. 5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-cyclobutyl-4-

methylbenzamide. To a 4-mL vial was added 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**, 27 mg, ~90% pure, 0.13 mmol), HOBT (20 wt % H₂O)(22 mg, 0.13 mmol),
 20 EDC (27 mg, 0.14 mmol). A solution of 5-carbamoyl-4-cyclobutyl-2-methylbenzoic acid (compound **6.5**, 28 mg, 0.12 mmol) in DMF (500 μL) was added followed by DIEA (83 μL, 0.48 mmol). The mixture was sealed and stirred at room temperature for 18 hours. Additional amine (3 mg, ~0.015 mmol) and EDC (5 mg, ~0.026 mmol) were added and the mixture was stirred at room temperature for an additional 27 hours. The mixture was diluted with ethyl
 25 acetate (5 mL) and washed with brine (8 mL). The aqueous was extracted with ethyl acetate (2 mL) and the combined organics was washed with brine, 1 M NaH₂PO₄, saturated NaHCO₃ and brine (7 mL each). The product began precipitating out of solution after the final brine

wash, so the organics was diluted with DCM (3 mL) and MeOH (~200 μ L). The organics was dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The crude product was triturated with Et_2O (1.5 mL), filtered, and washed with Et_2O (0.5 mL) to yield the title compound as an off white powder (41.4 mg, 92%). m/z (ES+) 374 ($\text{M}+\text{H}$)⁺. ^1H NMR (400 MHz, CDCl_3): δ 7.67 (d with fine str., J = 8.4 Hz, 2H), 7.40 (d with fine str., J = 8.4 Hz, 2H), 7.32 (s, 1H), 7.25 (s, 1H), 5.79 (br s, 1H), 5.70 (br s, 1H), 4.67-4.57 (m, 1H), 4.39-4.30 (m, 1H), 4.28-4.18 (m, 1H), 4.00-3.86 (m, 3H), 2.46 (s, 3H), 2.42-2.31 (m, 2H), 2.19-1.96 (m, 3H), 1.88-1.77 (m, 1H).




Compound 7.1. Methyl 4-methyl-3-(2-methyl-1H-imidazol-5-yl)benzoate. To a solution of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**, 600 mg, 2.17 mmol) in dioxane (20 mL) was added 2-methyl-4-bromoimidazole (419 mg, 2.6 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2\cdot\text{CH}_2\text{Cl}_2$ (180 mg, 0.22 mmol). The mixture was degassed argon and stirred for 10 minutes then aqueous potassium carbonate (1M, 10 mL, 10 mmol) was added and the mixture was stirred at 90 °C for 18 h. After cooling to ambient temperature, the reaction mixture was diluted with EtOAc and filtered through Celite®. The organic phase was washed by brine, dried (MgSO_4), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (0-5 % MeOH in EtOAc) to yield the title compound as a foam (324 mg, 65%). m/z (ES+) 231 ($\text{M}+\text{H}$)⁺. ^1H NMR (400 MHz, CDCl_3) δ 8.30 (br s, 1H), 7.86 (dd, J = 7.9, 1.9 Hz, 1H), 7.32 (d, J = 7.9 Hz, 1H), 7.08 (s, 1H), 3.92 (s, 3H), 2.53 (s, 3H), 2.51 (s, 3H).




Compound 7.2. Methyl 3-(5-chloro-2-methyl-1H-imidazol-4-yl)-4-methylbenzoate. To a solution of methyl 4-methyl-3-(2-methyl-1H-imidazol-5-yl) benzoate (compound **7.1**, 317 mg, 1.38 mmol) in carbon tetrachloride (30 mL) was added NCS (184 mg, 1.38 mmol). The mixture was stirred at 50 °C for 16 hours, then cooled to room temperature and washed with brine, dried (MgSO_4), filtered, and concentrated under reduced

pressure. The residue was purified by silica gel column chromatography (0-50 % EtOAc in hexanes) to yield the title compound as a solid (300 mg, 82%). m/z (ES+) 265 (M+H)⁺. ¹H NMR (400 MHz, CDCl₃) δ 9.55 (br s, 1H), 8.01 – 7.87 (m, 2H), 7.37 (d, J = 7.8 Hz, 1H), 3.91 (s, 3H), 2.42 (s, 3H), 2.40 (s, 3H).




Compound 7.3. 3-(5-Chloro-2-methyl-1H-imidazol-4-yl)-4-methylbenzoic acid. A

mixture of methyl 3-(5-chloro-2-methyl-1H-imidazol-4-yl)-4-methylbenzoate (compound 7.2, 10 mg, 0.038 mmol) in methanol (2 mL) and aqueous NaOH (2M, 0.2 mL, 0.4 mmol) was stirred at 50 °C for 16 hrs. The organic solvent was removed under reduced pressure and aqueous HCl (2 M) was added to the residue until a pH ~3-4 was attained. The solvents were removed under reduced pressure to produce a white solid which was a mixture of the title compound and salts and used in the next step without further purification. m/z (ES-) 249 (M-H)⁻.




Compound 7. 4-(1-(3-(4-Chloro-2-methyl-1H-imidazol-5-yl)-4-


methylbenzoyl)azetidin-3-yl)benzonitrile. A mixture of 3-(5-chloro-2-methyl-1H-imidazol-4-yl)-4-methylbenzoic acid (compound 7.3, ~0.038 mmol), 4-(azetidin-3-yl)benzonitrile hydrochloride (compound 5.2, 8.6 mg, 0.044 mmol), EDCI (12 mg, 0.063 mmol), HOBt (8 mg, 0.044 mmol) and DIEA (28 μ l, 0.16 mmol) in DMF (1 mL) was stirred at room temperature for 16 hours. The reaction was diluted with water and extracted with EtOAc. The organic phase was washed with brine (how much), dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel preparative TLC (8 % MeOH in DCM) to yield the title compound as a foam (2.8 mg, 19% over 2 steps). m/z (ES+) 391 (M+H)⁺. ¹H NMR (400 MHz, CDCl₃) δ 10.62 (s, 1H), 7.69 (d, J = 8.3 Hz, 2H), 7.54 – 7.40 (m, 3H), 7.37 (d, J = 1.9 Hz, 1H), 7.23 (d, J = 8.0 Hz, 1H), 4.80-4.58 (m, 2H), 4.41 – 4.19 (m, 2H), 4.03-3.92 (m, 1H), 2.49 (s, 3H), 2.32 (s, 3H).




Compound 8. 4-(1-(4-Cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)azetidin-3-yl)benzonitrile. To a 4-mL vial was added 5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-cyclobutyl-4-methylbenzamide (compound **6**, 39 mg, 0.104 mmol), dioxane (200 μ L) and 1,1-dimethoxy-*N,N*-dimethylethanamine (76 μ L, 0.52 mmol). The mixture was heated at 90 $^{\circ}$ C for 3 hours then cooled to room temperature. Acetic acid (42 μ L, 0.73 mmol) and hydrazine hydrate (30 μ L, 0.62 mmol) were added and the mixture was stirred at 90 $^{\circ}$ C for 1 hour. The mixture was diluted with DCM (5 mL) and washed with aqueous NaH_2PO_4 (1M, 5 mL) then saturated aqueous NaHCO_3 (5 mL). The organics was dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was purified by silica gel preparative thin layer chromatography (DCM/8% MeOH) to yield the title compound as a white powder (18.7 mg, 43%). m/z (ES+) 412 ($\text{M}+\text{H}$) $^{+}$. ^1H NMR (400 MHz, CD_3OD): δ 7.65 (d with fine str., $J = 8.4$ Hz, 2H), 7.57 (s, 1H), 7.39 (d with fine str., $J = 8.4$ Hz, 2H), 7.29 (s, 1H), 4.61 (app t, $J = 9.6$ Hz, 1H), 4.39 (app t, $J = 8.8$ Hz, 1H), 4.27-4.18 (m, 1H), 4.17-4.07 (m, 1H), 4.02-3.94 (m, 1H), 3.94-3.84 (m, 1H), 2.50 (s, 3H), 2.48 (s, 3H), 2.25-2.11 (m, 2H), 2.11-1.88 (m, 3H), 1.82-1.71 (m, 1H).




Compound 9.1. Methyl 4-cyclobutylbenzoate. To a stirred mixture of ZnBr_2 (83.0 g, 369 mmol) in THF (500 mL) under nitrogen at 0 $^{\circ}$ C was added a solution of bromo(cyclobutyl)magnesium (242 mL, 363 mmol, 1.5 M in THF) dropwise over 20 min. The resulting mixture was cooled to -40 $^{\circ}$ C and $\text{Pd}(\text{dppf})\text{Cl}_2$ (2.00 g, 2.73 mmol) and methyl 4-bromobenzoate (20.0 g, 93.0 mmol) were added. The resulting mixture was stirred at -40 $^{\circ}$ C for 1 h under nitrogen, and then carefully quenched with saturated aqueous NH_4Cl (500 mL). The mixture was extracted with ethyl acetate (3 x 500 mL) and the combined organic layers were washed with brine (3 x 500 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure to yield the title compound as a light yellow oil (18.0 g, crude).



Compound 9.2. Methyl 4-cyclobutyl-3-iodobenzoate. To a solution of methyl 4-cyclobutylbenzoate (compound **9.1**, 2.00 g, 10.5 mmol) in acetic acid (30 mL) was carefully added sodium periodate (1.00 g, 4.68 mmol), iodine (3.00 g, 11.8 mmol) and sulfuric acid (0.15 g). The resulting mixture was stirred overnight at 100 °C. After cooling to room temperature, the reaction was then carefully quenched with saturated aqueous Na₂S₂O₃ (30 mL) and the resulting mixture was extracted with ethyl acetate (3 x 20 mL). The combined organic layers were washed with brine (3 x 20 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield of the title compound as a yellow oil (1.50 g, 45%).




Compound 9.3. 4-Cyclobutyl-3-iodobenzoic acid. A solution of methyl 4-cyclobutyl-3-iodobenzoate (compound **9.2**, 11.0 g, 34.8 mmol) and sodium hydroxide (4.00 g, 100 mmol) in methanol (100 mL) and water (50 mL) was stirred at 50 °C overnight. After cooling to ambient temperature, the volatile solvent was removed under reduced pressure. The residual aqueous material was washed with ethyl acetate (20 mL). The pH of the aqueous was then adjusted to 3-4 with aqueous hydrogen chloride (6 M). The resulting precipitate was collected by filtration and dried to yield the title compound as a white solid (8.60 g, 82%).




Compound 9. 4-(1-(4-Cyclobutyl-3-(2-(methoxymethyl)-4-methyl-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1**, except 4-cyclobutyl-3-iodobenzoic acid (compound **9.3**) was used in place of 5-iodo-2,4-

dimethylbenzoic acid (compound **1.3**) and 2-methoxyacetimidamide hydrochloride (compound **4.7**) was used in place of acetimidamide hydrochloride. m/z (ES⁺) 469 (M+H)⁺.




Compound 10.1. Methyl 4-cyclopropylbenzoate. Into a 1-L three neck round-

5 bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution mixture of ZnBr₂ (41.5 g, 184 mmol) and tetrahydrofuran (500 mL). The mixture was cooled to 0 °C, then cyclopropylmagnesium bromide (2 M in THF)(92 mL, 184 mmol) was added drop-wise with stirring over 30 min. The mixture was then cooled to -40 °C and Pd(dppf)Cl₂ (3.00 g, 4.1 mmol) was added portion-wise over 1 min. Methyl 4-
10 bromobenzoate (10.0 g, 46.50 mmol) in THF (50 mL) was added drop-wise over 30 min at -40 °C. The resulting mixture was allowed to warm to room temperature and stirred overnight. The reaction was then carefully quenched by the addition of aqueous NH₄Cl (sat., 500 mL). The mixture was extracted with ethyl acetate (3 x 300 mL) and the combined organic layers were washed with brine (3 x 300 mL), dried (Na₂SO₄), filtered, and concentrated under
15 reduced pressure. The residue was purified by silica gel column chromatography with petroleum ether as the eluent to yield 8.87 g (crude) of the title compound as a yellow oil.




Compound 10.2. Methyl 3-bromo-4-cyclopropylbenzoate. Into a 50-mL round-


bottom flask, was placed a mixture of methyl 4-cyclopropylbenzoate (compound **10.1**, 500
20 mg, 2.84 mmol), *N*-bromosuccinimide (500 mg, 2.81 mmol) and trifluoroacetic acid (20 mL). The mixture was stirred overnight at 50 °C, then cooled and carefully quenched with water (10 mL). The mixture was extracted with ethyl acetate (3 x 30 mL) and the combined organic layers were washed with brine (3 x 30 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with
25 petroleum ether as the eluent to yield 0.5 g (69%) of the title compound as a yellow oil.



Compound 10.3. 3-Bromo-4-cyclopropylbenzoic acid. Into a 50-mL round-bottom flask, was placed a solution of methyl 3-bromo-4-cyclopropylbenzoate (compound **10.2**, 500 mg, 1.96 mmol) in methanol (10 mL) and a solution of sodium hydroxide (500 mg, 12.5 mmol) in water (5 mL). The resulting solution was stirred overnight at 50 °C, then the volatiles were removed under reduced pressure. The pH of the residual solution was adjusted to 6-7 with aqueous HCl (6 M). The resulting solids were collected by filtration to yield 0.3 g (63%) of the title compound as a white solid.




Compound 10.4. 4-Cyclopropyl-3-formylbenzoic acid. Into a 100-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 3-bromo-4-cyclopropylbenzoic acid (compound **10.3**, 500 mg, 2.07 mmol) in tetrahydrofuran/Et₂O (1:1, 20 mL). The solution was cooled to -78 °C then n-BuLi (2.5 M in THF) (1.8 mL, 4.5 mmol) was added drop-wise with stirring. The resulting solution was stirred for an additional 30 min at -78 °C, then *N,N*-dimethylformamide (0.49 mL, 6.3 mmol) was added drop-wise over 10 min. The mixture was stirred for 2 h at -78 °C, then quenched with water (100 mL). The pH of the solution was adjusted to 3-4 with aqueous HCl (6 M), then extracted with ethyl acetate (3 x 30 mL). The combined organic layers were washed with brine (3 x 30 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:5) as the eluent to yield 0.3 g (76%) of the title compound as a white solid.




Compound 10.5. 4-Cyclopropyl-3-(1-hydroxypropyl)benzoic acid. Into a 250-mL 3-necked round-bottom flask, purged and maintained with an inert atmosphere of nitrogen

and containing THF (60 mL), was placed a solution of ethylmagnesium bromide (3 M in diethyl ether) (24.6 mL, 73.8 mmol). A solution of 4-cyclopropyl-3-formylbenzoic acid (compound **10.4**, 3.5 g, 18.4 mmol) in THF (40 mL) was added drop-wise at room temperature over 45 min. The resulting mixture was stirred for 1.5 h at room temperature, then carefully quenched with water/ice (50 mL). The mixture was diluted with EtOAc (100 mL) and the organic layer was washed with aqueous NH_4Cl (sat., 2 x 100 mL) and the aqueous layers combined. The pH of the aqueous phase was adjusted to 3-4 with aqueous hydrogen chloride (2 M) and extracted with ethyl acetate (2 x 150 mL). The combined organic extracts were washed with brine (30 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate:PE (1:10-2:1) as the eluent to yield 3.4 g (84%) of the title compound as a white solid.




Compound 10.6. 4-Cyclopropyl-3-propionylbenzoic acid. Into a 500-mL round-bottom flask, was placed a mixture of 4-cyclopropyl-3-(1-hydroxypropyl)benzoic acid (compound **10.5**, 3.4 g, 15.44 mmol) in 1,2-dichloroethane (300 mL) and Dess-Martin periodinane (7.2 g, 16.9 mmol). The mixture was stirred for 1.5 h at room temperature and then the solids were removed with filtration. The filtrate was concentrated under reduced pressure and the residue was purified by silica gel column chromatography with ethyl acetate:PE (1:9) to methanol/ethyl acetate (20:1) as the eluent to yield 2.3 g (68%) of the title compound as a light yellow solid.




Compound 10.7. Methyl 4-cyclopropyl-3-propionylbenzoate. Into a 50-mL round-bottom flask, was placed a solution of 4-cyclopropyl-3-propanoylbenzoic acid (compound **10.6**, 500 mg, 2.29 mmol) in *N,N*-dimethylformamide (20 mL) and potassium carbonate (633 mg, 4.55 mmol). The mixture was cooled to 0-5 °C then iodomethane (157 μL , 2.52 mmol) was added drop-wise. The resulting mixture was stirred for 1 h at 5-10 °C, then quenched


with water/ice (80 mL). The aqueous phase was extracted with ethyl acetate (2 x 150 mL) and the combined organic layers were washed with brine (20 mL), sodium carbonate (sat., 20 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. Obtained 400 mg (75%) of the title compound as a light brown oil.



Compound 10. 4-(1-(4-Cyclopropyl-3-(2-(methoxymethyl)-4-methyl-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1**, except methyl 4-cyclopropyl-3-propionylbenzoate (compound **10.7**) was used in place of methyl 2,4-dimethyl-5-propionylbenzoate (compound **1.5**) and 2-methoxyacetimidamide hydrochloride (compound **4.7**) was used in place of acetimidamide hydrochloride. *m/z* (ES⁺) 455 (M+H)⁺.




Compound 11.1. 4-Methyl-3-propionylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **10.6**, except 3-bromo-4-methylbenzoic acid was used in place of 3-bromo-4-cyclopropylbenzoic acid (compound **10.3**).




Compound 11. 4-(1-(3-(2-(Methoxymethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1**, except 4-methyl-3-propionylbenzoic acid (compound **11.1**) was used in place of 2,4-dimethyl-5-propionylbenzoic acid (compound **1.4**) and 2-methoxyacetimidamide


hydrochloride (compound **4.7**) was used in place of acetimidamide hydrochloride. m/z (ES+) 429 (M+H)⁺.



Compound 12. 4-(1-(3-(2-(Hydroxymethyl)-4-methyl-1*H*-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **3**, except 4-(1-(3-(2-(methoxymethyl)-4-methyl-1*H*-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile (compound **11**) was used in place of 4-(1-(5-(2-(methoxymethyl)-4-methyl-1*H*-imidazol-5-yl)-2,4-dimethylbenzoyl)piperidin-4-yl)benzonitrile (compound **2**). m/z (ES+) 415 (M+H)⁺.




Compound 13.1. *tert*-Butyl 4-(4-bromophenyl)-4-hydroxypiperidine-1-carboxylate. To a stirred solution of 1-bromo-4-iodobenzene (93.7 g, 331 mmol) in tetrahydrofuran (800 mL) under nitrogen at -78 °C was added drop-wise a solution of butyllithium (150 mL, 2.43 M in THF) over 30 min. The resulting solution was stirred for 2 h at -78 °C. A solution of *tert*-butyl 4-oxopiperidine-1-carboxylate (60 g, 301 mmol) in tetrahydrofuran (800 mL) was then added drop-wise with stirring at -78 °C over 30 min. The mixture was stirred for 1 h at -78 °C, then the reaction was carefully quenched with water (350 mL). The resulting mixture was extracted with ethyl acetate (2 x 400 mL) and the combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified using silica gel column chromatography with ethyl acetate/petroleum ether (1:200-1:10) as the eluent to yield 91 g (85%) of the title compound as a yellow oil.




Compound 13.2. *tert*-Butyl 4-(4-cyanophenyl)-4-hydroxypiperidine-1-carboxylate.

A mixture of *tert*-butyl 4-(4-bromophenyl)-4-hydroxypiperidine-1-carboxylate (compound **13.1**, 36 g, 101 mmol), $\text{Pd(PPh}_3)_4$ (11.7 g, 10.1 mmol), and Zn(CN)_2 (17.9 g, 152.4 mmol) in DMF (400 mL) under nitrogen was stirred overnight at 80 °C. The mixture was cooled to ambient temperature, then the reaction was quenched by the addition of 600 mL of FeSO_4 (aq., sat.) and diluted with ethyl acetate. The resulting mixture was stirred vigorously then filtered through Celite® and washed with 1 M FeSO_4 , water, and ethyl acetate. The layers were separated and the aqueous phase was extracted with ethyl acetate (2 x 300 mL). The combined organic layers were washed with potassium carbonate (aq., sat., 200 mL), followed by brine (200 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was purified using silica gel column chromatography with ethyl acetate/petroleum ether (1:200-1:5) as the eluent to yield 23 g (75%) of the title compound as a white solid.




Compound 13.3. *tert*-Butyl 4-(4-cyanophenyl)-4-fluoropiperidine-1-carboxylate.


To a stirred solution of compound **13.2** (5.00 g, 16.5 mmol) in dichloromethane (250 mL) at -78 °C under nitrogen was added drop-wise Deoxo-Fluor® (4.4 g, 19.9 mmol). The resulting mixture was stirred for 1 h at -78 °C. The reaction mixture was then *carefully* quenched by the addition of sodium bicarbonate (aq., sat., 50 mL) and extracted with dichloromethane (3 x 100 mL). The combined organic layers were washed with brine (150 mL), dried over anhydrous magnesium sulfate, filtered, and concentrated under reduced pressure. The residue was purified using silica gel column chromatography with ethyl acetate/petroleum ether (1:30) as the eluent to yield 2.5 g (35%) of the title compound as a white solid.



Compound 13.4. 4-(4-Fluoropiperidin-4-yl)benzonitrile hydrochloride. The title compound was prepared using standard chemical manipulations and a procedure similar to that used for the preparation of compound **1.2**, except using compound **13.3** in place of compound **1.1**. m/z (ES⁺) 205 (M+H)⁺. ¹H NMR (300 MHz, CD₃OD): δ 7.83 (d, J = 6.3 Hz, 2H), 7.68 (d, J = 6.3 Hz, 2H), 3.55-3.32 (m, 4H), 2.58-2.40 (m, 2H), 2.28-2.22 (m, 2H).




Compound 14.1. 4-Methyl-2-(trifluoromethyl)-1H-imidazole. Into a 250-mL round-bottom flask, purged and maintained with an inert atmosphere of nitrogen, was placed 2-oxopropanal (5 g, 27.8 mmol), acetic acid (150 mL), 1-ethoxy-2,2,2-trifluoroethanol (13.3 g, 83.2 mmol), and ammonium acetate (17.1 g, 222 mmol). The mixture was stirred for 18 h at 110 °C, then cooled to room temperature and concentrated under reduced pressure. The residue was quenched with water/ice (30 mL) and the aqueous phase was extracted with ethyl acetate (2 x 100 mL). The combined organic layers were washed with brine (30 mL) and concentrated under reduced pressure. The residue was dissolved in aqueous hydrogen chloride (5 M, 20 mL) and washed with ethyl acetate (2 x 50 mL). The pH of the aqueous layer was adjusted to 8-9 with sodium carbonate, then extracted with ethyl acetate (2 x 100 mL). The combined organic layers were washed with brine (30 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude product was purified by silica gel column chromatography with ethyl acetate:PE (1:5) as the eluent to yield 1.2 g (29%) of the title compound as a brown oil.




Compound 14.2. 4-Methyl-1H-imidazole-2-carbonitrile. Into a 250-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed 4-methyl-2-(trifluoromethyl)-1H-imidazole (compound **14.1**, 800 mg, 5.33 mmol) and 5% aqueous ammonium hydroxide (50 mL). The resulting solution was stirred for 40 h at 25-30 °C, then the pH of the solution was adjusted to 5-6 with acetic acid. The aqueous phase


was extracted with ethyl acetate (3 x 150 mL) and the combined organic extracts were washed with brine (2 x 50 mL) and aqueous Na₂CO₃ (sat., 2 x 20 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate:PE (1:20-1:10) as the eluent to yield 350 mg (61%) of the title compound as a white solid.




Compound 14.3. 5-Iodo-4-methyl-1H-imidazole-2-carbonitrile. Into a 25-mL round bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a mixture of 4-methyl-1H-imidazole-2-carbonitrile (compound **14.2**, 350 mg, 3.27 mmol) and aqueous sodium hydroxide (2 M, 5 mL) and stirred for 15 min at room temperature. This was followed by the drop-wise addition of a solution of iodine (1.25 g, 4.92 mmol) in dichloromethane (5 mL). The resulting mixture was stirred for 24 h at room temperature, then diluted with water/ice (10 mL). The aqueous layer was washed with DCM (10 mL), and then the aqueous layer was acidified to pH 5-6 with acetic acid. The aqueous phase was extracted with EtOAc (5 x 50 mL) and the combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield 500 mg (66%) of the title compound as a brown solid.




Compound 14. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-4-methyl-1H-imidazole-2-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 5-Iodo-4-methyl-1H-imidazole-2-carbonitrile (compound **14.3**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**). *m/z* (ES⁺) 382 (M+H)⁺.




Compound 15. 5-(5-(4-(4-Cyanophenyl)piperidine-1-carbonyl)-2-methylphenyl)-4-methyl-1*H*-imidazole-2-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 5, except 5-Iodo-4-methyl-1*H*-imidazole-2-carbonitrile (compound **14.3**) was used in place of 5-iodo-2,4-dimethyl-1*H*-imidazole (compound **5.5**) and 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 410 (M+H)⁺.




Compound 16.1. 3,3,3-Trifluoro-2-oxopropanal. Into a 100-mL round-bottom flask, was placed a mixture of 3,3-dibromo-1,1,1-trifluoropropan-2-one (10.37 g, 38.43 mmol) and sodium acetate (12.61 g, 153.7 mmol) in water (50 mL). The resulting solution was stirred overnight at 100 °C, then cooled to room temperature and extracted with ethyl acetate (3 x 20 mL). The combined organic extracts were dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield 1.52 g (31%) of the title compound as light yellow oil.




Compound 16.2. 2-Methyl-4-(trifluoromethyl)-1*H*-imidazole. Into a 10-mL sealed tube, was placed a solution of acetaldehyde (296 μL, 5.29 mmol), 3,3,3-trifluoro-2-oxopropanal (compound **16.1**, 1.0 g, 7.9 mmol), and 25% ammonium hydroxide (0.8 mL) in methanol (5 mL). The solution was stirred for 3 h at 0 °C, then stirred at room temperature overnight. The resulting mixture was concentrated under reduced pressure and then the residue was diluted with water (50 mL). The aqueous phase was extracted with ethyl acetate (3 x 20 mL) and the combined organic extracts were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude product was purified by chromatography to yield 430 mg (54%) of the title compound as a light yellow solid.




Compound 16.3. 2-Methyl-1*H*-imidazole-4-carbonitrile. Into a 100-mL round-bottom flask, was placed a solution of 2-methyl-4-(trifluoromethyl)-1*H*-imidazole (compound **16.2**, 300 mg, 2.00 mmol) in 5% ammonium hydroxide (35 mL). The resulting solution was stirred for 4 days at room temperature, then the pH of the solution was adjusted to 7 with acetic acid. The aqueous phase was extracted with of ethyl acetate (3 x 10 mL) and the combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield 170 mg (79%) of the title compound as a light yellow solid.




Compound 16.4. 5-Iodo-2-methyl-1*H*-imidazole-4-carbonitrile. Into a 25-mL round-bottom flask, was placed a mixture of 2-methyl-1*H*-imidazole-4-carbonitrile (compound **16.3**, 50 mg, 0.47 mmol), *N*-iodosuccinimide (116 mg, 0.52 mmol) in ACN (5 mL). The resulting solution was heated at reflux overnight, then cooled to room temperature and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1/2) to yield 60 mg (55%) of the title compound as a light yellow solid.




Compound 16. 5-(5-(4-(4-Cyanophenyl)piperidine-1-carbonyl)-2-methylphenyl)-2-methyl-1*H*-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 5-iodo-2-methyl-1*H*-imidazole-4-carbonitrile (compound **16.4**) was used in place of 5-iodo-2,4-dimethyl-1*H*-imidazole (compound **5.5**) and 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). *m/z* (ES⁺) 410 (M+H)⁺.




Compound 17.1. 3-Butyryl-4-methylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **10.6**, except 3-bromo-4-methylbenzoic acid was used in place of 3-bromo-4-cyclopropylbenzoic acid (compound **10.3**) and propylmagnesium bromide was used in place of ethylmagnesium bromide.



Compound 17. 4-(1-(3-(4-ethyl-2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1**, except 3-butyryl-4-methylbenzoic acid (compound **17.1**) was used in place of 2,4-dimethyl-5-propionylbenzoic acid (compound **1.4**) and 2-methoxyacetimidamide hydrochloride (compound **4.7**) was used in place of acetimidamide hydrochloride. m/z (ES+) 443 (M+H)⁺.




Compound 18.1. 2-Methyl-5-propionylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **10.6**, except 5-bromo-2-methylbenzoic acid was used in place of 3-bromo-4-cyclopropylbenzoic acid (compound **10.3**).




Compound 18. 4-(1-(5-(2,4-Dimethyl-1H-imidazol-5-yl)-2-methylbenzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of


compound **1**, except 2-methyl-5-propionylbenzoic acid (compound **18.1**) was used in place of 2,4-dimethyl-5-propionylbenzoic acid (compound **1.4**). m/z (ES⁺) 399 (M+H)⁺.



Compound 19. 4-(1-(4-Ethyl-3-(2-(methoxymethyl)-4-methyl-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1**, except 4-ethylbenzoic acid was used in place of 2,4-dimethylbenzoic acid and 2-methoxyacetimidamide hydrochloride (compound **4.7**) was used in place of acetimidamide hydrochloride. m/z (ES⁺) 443 (M+H)⁺.




Compound 20.1. Methyl 4-(bromomethyl)-3-iodobenzoate. Into a 100-mL round-bottom flask, was placed a mixture of methyl 3-iodo-4-methylbenzoate (compound **5.3**, 3.00 g, 10.9 mmol) in CCl₄ (50 mL), NBS (2.9 g, 16.3 mmol), azobisisobutyronitrile (360 mg, 2.19 mmol) and potassium carbonate (1.65 g, 11.9 mmol). The resulting mixture was stirred overnight at 70 °C, then cooled to room temperature and concentrated under reduced pressure. The residue was diluted with EtOAc (100 mL) and the mixture was washed with brine (2 x 50 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:20) as the eluent to yield 3.0 g (78%) of the title compound as a yellow solid.




Compound 20.2. Methyl 3-iodo-4-(methoxymethyl)benzoate. Into a 250-mL round-bottom flask, was placed a mixture of methyl 4-(bromomethyl)-3-iodobenzoate (compound **20.1**, 3.0 g, 8.5 mmol) and sodium methoxide (1.8 g, 33 mmol) in methanol (100 mL). The resulting solution was stirred for 2 h at 50 °C, then cooled to room temperature and concentrated under reduced pressure. The residue was diluted with ethyl acetate (50 mL) and

the solids were filtered off. The filtrate was concentrated under reduced pressure to yield 2.0 g (77%) of the title compound as a yellow solid.




Compound 20. 4-(1-(3-(2,4-dimethyl-1H-imidazol-5-yl)-4-


- 5 **(methoxymethyl)benzoyl)piperidin-4-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except methyl 3-iodo-4-(methoxymethyl)benzoate (compound **20.2**) was used in place of methyl 3-iodo-4-methylbenzoate (compound **5.3**) and 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 429 (M+H)⁺.




- Compound 21.1. 2,5-Diiodo-4-methyl-1H-imidazole.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.5**, except 4-methyl-1H-imidazole (2.0 g, 24.4 mmol) was used in place of 2,4-dimethyl-1H-imidazole and 2 equivalents of NIS was used to yield 4.0 g (49%) of the title compound as a white solid.




- Compound 21.2. 5-Iodo-4-methyl-1H-imidazole.** Into a 250-mL round-bottom flask, was placed 2,5-diiodo-4-methyl-1H-imidazole (compound **21.1**, 1.0 g, 3.0 mmol), Na₂SO₃ (3.0 g, 25.4 mmol) and ethanol/water (20/40 mL). The resulting mixture was stirred overnight at reflux, then cooled and concentrated under reduced pressure. The residue was dissolved in water (100 mL) and extracted with ethyl acetate (2 x 100 mL). The combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield 0.60 g (96%) of the title compound as a white solid.




Compound 21. 4-(1-(4-methyl-3-(4-methyl-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 5-iodo-4-methyl-1H-imidazole (compound **21.2**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**) and 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 385 (M+H)⁺.




Compound 22. 4-(1-(3-(2,4-Dimethyl-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1**, except 3-iodobenzoic acid was used in place of 5-iodo-2,4-dimethylbenzoic acid (compound **1.3**). m/z (ES+) 385 (M+H)⁺.




Compound 23.1. Methyl 3-hydroxypropanimidate hydrochloride. Into a 50-mL 3-necked round-bottom flask, was placed a solution of 3-hydroxypropanenitrile (2.00 g, 28.1 mmol) in ether (10 mL) and methanol (5 mL). HCl gas was introduced by bubbling through the solution. The resulting solution was stirred for 2 h at room temperature, then the resulting mixture was carefully concentrated under reduced pressure. The residue was washed with ether (2 x 20 mL) and the solids were collected by filtration to yield 1.3 g (33%) of the title compound as a white solid.




Compound 23.2. 3-Hydroxypropanimidamide hydrochloride. Into a 50-mL 3-necked round-bottom flask, was placed a solution of methyl 3-hydroxypropanecarboximidate hydrochloride (compound **23.1**, 2.00 g, 14.3 mmol) in methanol (5 mL). The solution was cooled to 0 °C and ammonia (gas) was introduced over 20 min. The resulting solution was stirred for 2 h at room temperature, then concentrated under reduced pressure. The residue was washed with petroleum ether (2 x 20 mL) and EtOAc (2 x 20 mL) and the solids were collected by filtration to yield 1.6 g (crude) of the title compound as a white solid.



Compound 23. 4-(1-(3-(2-(2-Hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1**, except 4-methyl-3-propionylbenzoic acid (compound **11.1**) was used in place of 2,4-dimethyl-5-propionylbenzoic acid (compound **1.4**) and 3-hydroxypropanimidamide hydrochloride (compound **23.2**) was used in place of acetimidamide hydrochloride. *m/z* (ES⁺) 429 (M+H)⁺.




Compound 24.1. 4-(1-(3-(2-(2-Chloroethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile. Into a 50-mL round-bottom flask, was placed a solution of 4-(1-(3-(2-(2-hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile (compound **23**, 75 mg, 0.18 mmol) in dichloromethane (5 mL). Thionyl chloride (25 µL, 0.35 mmol) was added and the resulting solution was stirred for 3 h at room temperature. The solvents were carefully removed under reduced pressure to yield 50 mg (crude) of the title compound as a yellow solid.




Compound 24. 4-(1-(3-(2-(2-(Dimethylamino)ethyl)-4-methyl-1*H*-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile. Into a vessel with condensor, was placed a solution of 4-(1-(3-(2-(2-chloroethyl)-4-methyl-1*H*-imidazol-5-yl)-4-

5 methylbenzoyl)piperidin-4-yl)benzonitrile (compound **24.1**, 26 mg, 0.06 mmol, 1.00 equiv) in EtOH (10 mL). Dimethylamine (1 M in THF, 0.3 mL, 0.3 mmol) was added and the mixture was sealed under a nitrogen balloon. The resulting solution was stirred overnight with a 100 °C oil bath. After cooling to room temperature, the resulting mixture was concentrated under reduced pressure and the crude product was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-001(SHIMADZU)): Column, XBridge Shield RP18
10 OBD Column, 5 μ m, 19*150 mm; mobile phase, WATER WITH 0.03% $\text{NH}_3\text{H}_2\text{O}$ and CH_3CN (31% CH_3CN up to 43% in 7 min, up to 100% in 0.5 min, down to 31% in 3.5 min); Detector, Waters 2489, 254 & 220 nm. The fractions containing pure compound were combined and lyophilized to yield 6.3 mg (24%) of the title compound as a white solid. m/z
15 (ES+) 456 (M+H)⁺.




Compound 25.1. 4-(1-(3-(2,4-Dimethyl-1*H*-imidazol-5-yl)-4-fluorobenzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound
20 **1**, except 4-fluorobenzoic acid was used in place of 2,4-dimethylbenzoic acid.




Compound 25. 4-(1-(4-(Azetidin-1-yl)-3-(2,4-dimethyl-1*H*-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. Into a 10-mL sealed tube, was placed a solution of

4-(1-(3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-fluorobenzoyl)piperidin-4-yl)benzonitrile (compound **25.1**, 200 mg, 0.50 mmol) in 1,4-dioxane (4 mL). Azetidine hydrochloride (50 mg, 0.53 mmol) and cesium carbonate (500 mg, 1.53 mmol) were added and the mixture was stirred for 48 h at 120 °C behind a blast shield. The mixture was cooled and the solids were removed by filtration. The filtrate was concentrated under reduced pressure and the crude product was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-001(SHIMADZU)): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150 mm; mobile phase, Water with 50 mmol NH_4HCO_3 and CH_3CN (33.0% CH_3CN up to 46.0% in 10 min, up to 100.0% in 3 min, down to 33.0% in 1 min); Detector, Waters 2489, 254 & 220 nm. The fractions containing pure compound were combined and lyophilized to yield 59.6 mg (27%) of the title compound as a white solid. m/z (ES⁺) 440 (M+H)⁺.




Compound 26.1. 4-Methyl-3-(2-methyl-1*H*-imidazol-4-yl)benzoic acid. Methyl 4-methyl-3-(2-methyl-1*H*-imidazol-5-yl) benzoate (compound **7.1**, 219 mg, 0.95 mmol) was dissolved in a mixture of methanol (20 mL) and aqueous sodium hydroxide (5 mL, 2M). The resulting solution was heated at 50 °C for 16 hrs, then the reaction was cooled to room temperature and the organic solvent was removed under reduced pressure. The resulting aqueous residue was acidified to pH 3-4 with aqueous HCl (2 M). The resulting solids were collected and dried to yield 128 mg (62%) of the title compound as a white solid. m/z (ES⁻) 215 (M-H)⁻.




Compound 26. 4-(1-(4-Methyl-3-(2-methyl-1*H*-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 4-methyl-3-(2-methyl-1*H*-imidazol-4-yl)benzoic acid (compound **26.1**, 65 mg, 0.30 mmol) was used in place of 3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**, 74 mg, 0.33 mmol) was used in

place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). 79 mg (68%) of the title compound was obtained as a white solid. m/z (ES+) 385 (M+H)⁺.




Compound 27.1. Methyl 3-acetyl-4-methylbenzoate. Into a 250-mL round-bottom

flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a mixture of methyl 3-iodo-4-methylbenzoate (compound **5.3**, 4.50 g, 16.3 mmol), 1-(vinylloxy)butane (4.21 mL, 32.6 mmol), TEA (4.53 mL, 32.5 mmol), 1,3-bis(diphenylphosphino)propane (672 mg, 1.63 mmol) and Pd(OAc)₂ (349 mg, 1.55 mmol) in DMSO (50 mL). The mixture was stirred for 12 hours at 120 °C, then cooled to room temperature. The pH was adjusted to 1-2 with aqueous hydrogen chloride (2 M) and stirred for 1 hour. The aqueous phase was extracted with ethyl acetate (3 x 200 mL) and the combined organic layers were washed with water (100 mL), then brine (3 x 100 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:50) as the eluent to yield 1.45 g (46%) of the title compound as a yellow solid.



Compound 27.2. Methyl 3-(2-bromoacetyl)-4-methylbenzoate. Into a 50-mL


round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of methyl 3-acetyl-4-methylbenzoate (compound **27.1**, 200 mg, 1.04 mmol) in chloroform (4 mL). Bromine (53 μL, 1.04 mmol) was added drop-wise and the solution was stirred for 2 h at room temperature, then concentrated under reduced pressure. Obtained 300 mg (crude) of the title compound as a brown solid.




Compound 27.3. Methyl 3-(2-(methoxymethyl)-1H-imidazol-5-yl)-4-

methylbenzoate. Into a 50-mL round-bottom flask, which was purged and maintained with

an inert atmosphere of nitrogen, was placed a solution of methyl 3-(2-bromoacetyl)-4-methylbenzoate (compound **27.2**, 281 mg, 1.04 mmol) in ACN (5 mL). 2-Methoxyacetimidamide (compound **4.7**, 194 mg, 1.56 mmol) and potassium carbonate (434 mg, 3.14 mmol) were added and the resulting mixture was stirred for 12 hours at 80 °C, then concentrated under reduced pressure. The residue was diluted with water (50 mL) and extracted with ethyl acetate (3 x 30 mL). The combined organic layers were washed with brine (3 x 50 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:1) as the eluent to yield 50 mg (19%) of the title compound as a brown solid.




Compound 27.4. Methyl 3-(4-chloro-2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoate. Into a 50-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of methyl 3-(2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoate (compound **27.3**, 40 mg, 0.15 mmol) and NCS (24.6 mg, 0.18 mmol) in CCl₄ (20 mL). The resulting mixture was stirred for 2 h at 50 °C, then cooled to room temperature and quenched with water (100 mL). The aqueous phase was extracted with ethyl acetate (100 mL) and the combined organic layers were washed with brine (3 x 30 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield 30 mg (crude) of the title compound as a yellow solid.




Compound 27.5. 3-(4-Chloro-2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoic acid. Into a 50-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was added methyl 3-(4-chloro-2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoate (compound **27.4**, 50 mg, 0.19 mmol) and sodium hydroxide (31 mg, 0.76 mmol) in methanol/water (3 mL/3 mL). The resulting solution was stirred for 12 hours at room temperature. The pH of the solution was adjusted to 1-2 with hydrogen

chloride (6 M), then concentrated under reduced pressure to yield 150 mg (crude) of the title compound as the HCl salt as white solid.



Compound 27. 4-(1-(3-(4-Chloro-2-(methoxymethyl)-1H-imidazol-5-yl)-4-


methylbenzoyl)piperidin-4-yl)benzonitrile. Into a 50-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 3-(4-chloro-2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **27.5**, 20 mg, 0.07 mmol) in *N,N*-dimethylformamide (3 mL). 4-Dimethylaminopyridine (17.4 mg, 0.14 mmol), EDC•HCl (27 mg, 0.14 mmol), 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**, 15.5 mg, 0.07 mmol, 1.00 equiv) were added and the resulting solution was stirred for 12 h at room temperature. The reaction was quenched with water (10 mL) and extracted with ethyl acetate (3 x 10 mL). The combined organic layers were washed with brine (10 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-001(SHIMADZU)): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150 mm; mobile phase, water with 0.05% TFA and CH₃CN (35% CH₃CN up to 50% in 7 min, up to 100% in 3 min, down to 35% in 1 min); Detector, Waters 2489, 254 & 220 nm. The fractions containing pure compound were combined and lyophilized to yield 15 mg (47%) of the title compound as a white solid. *m/z* (ES⁺) 449 (M+H)⁺.




Compound 28.1. 3-(4-Chloro-2-(hydroxymethyl)-1H-imidazol-5-yl)-4-

methylbenzoic acid. Into a 50-mL round-bottom flask, was placed a solution of methyl 3-(4-chloro-2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoate (compound **27.4**, 380 mg, 1.29 mmol, 1.00 equiv) in HBr (40% in AcOH)(10 mL). The solution was stirred overnight at 80 °C, then cooled and concentrated under reduced pressure. The crude residue was purified by prep-HPLC (WATER WITH 0.05% TFA and CH₃CN (0% CH₃CN in 3 min, then up to


100% for 5 min, down to 0% in 1 min); Detector, 254 & 220 nm. The fractions containing clean product were combined and lyophilized to yield 171 mg (50%) of the title compound as yellow crude oil.



- 5 **Compound 28. 4-(1-(3-(4-Chloro-2-(hydroxymethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 27, except 3-(4-chloro-2-(hydroxymethyl)-1H-imidazol-5-yl)-4-methylbenzoic acid (compound 28.1) was used in place of 3-(4-chloro-2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoic acid (compound 27.5) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound 5.2) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound 1.2). m/z (ES+) 407 (M+H)⁺.




- 15 **Compound 29. 4-(1-(3-(4-Chloro-2-(hydroxymethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 27, except 3-(4-chloro-2-(hydroxymethyl)-1H-imidazol-5-yl)-4-methylbenzoic acid (compound 28.1) was used in place of 3-(4-chloro-2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoic acid (compound 27.5). m/z (ES+) 435 (M+H)⁺.




- 20 **Compound 30. 4-(1-(3-(4-Chloro-2-(hydroxymethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile.** The title compound was prepared using


standard chemical manipulations and procedures similar to those used for the preparation of compound **27**, except 3-(4-chloro-2-(hydroxymethyl)-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **28.1**) was used in place of 3-(4-chloro-2-(methoxymethyl)-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **27.5**) and 4-(3-Fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**). m/z (ES+) 425 (M+H)⁺.




Compound 31.1. Methyl 5-iodo-2,4-dimethylbenzoate. A solution of 5-iodo-2,4-dimethylbenzoic acid (compound **1.3**, 10.0 g, 32.6 mmol, 90%) and sulfuric acid (10 mL) in methanol (100 mL) was stirred overnight at 80 °C. After cooling to room temperature, the mixture was concentrated under reduced pressure and the residue was diluted with of ethyl acetate (200 mL). The resulting mixture was washed with water (3 x 50 mL), sodium bicarbonate (aq. sat., 2 x 50 mL, caution: gas evolution), followed by brine (2 x 50 mL). The organic phase was dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure to yield 9.2 g (88%) of the title compound as a yellow oil.




Compound 31. 4-(1-(5-(4-Chloro-2-methyl-1*H*-imidazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for preparation of compound **27**, except methyl 5-iodo-2,4-dimethylbenzoate (compound **31.1**) was used in place of methyl 3-iodo-4-methylbenzoate (compound **5.3**), acetimidamide hydrochloride was used in place of 2-methoxyacetimidamide (compound **4.7**) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**). m/z (ES+) 405 (M+H)⁺.




Compound 32. 4-(1-(5-(4-Chloro-2-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for preparation of compound **27**, except methyl 5-iodo-2,4-dimethylbenzoate (compound **31.1**) was used in place of methyl 3-iodo-4-methylbenzoate (compound **5.3**) and acetimidamide hydrochloride was used in place of 2-methoxyacetimidamide (compound **4.7**). m/z (ES⁺) 433 (M+H)⁺.




Compound 33. 4-(1-(3-(4-Chloro-2-methyl-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for preparation of compound **27**, except methyl 3-iodobenzoate was used in place of methyl 3-iodo-4-methylbenzoate (compound **5.3**) and acetimidamide hydrochloride was used in place of 2-methoxyacetimidamide (compound **4.7**). m/z (ES⁺) 405 (M+H)⁺.




Compound 34. 4-(1-(3-(4-Chloro-2-methyl-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for preparation of compound **27**, except methyl 3-iodobenzoate was used in place of methyl 3-iodo-4-methylbenzoate (compound **5.3**), acetimidamide hydrochloride was used in place of 2-methoxyacetimidamide (compound **4.7**) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**). m/z (ES⁺) 377 (M+H)⁺.




Compound 35. (3-(4-chlorophenyl)pyrrolidin-1-yl)(3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylphenyl)methanone. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 5, except 3-(4-chlorophenyl)pyrrolidine was used in place 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 394 (M+H)⁺.




Compound 36.1. *tert*-Butyl 4-([1,2,4]triazolo[4,3-*a*]pyridin-6-yl)piperidine-1-carboxylate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1.1**, except 6-bromo-[1,2,4]triazolo[4,3-*a*]pyridine was used in place of 4-bromobenzonitrile. m/z (ES⁺) 303 (M+H)⁺.




Compound 36.2. 6-(Piperidin-4-yl)-[1,2,4]triazolo[4,3-*a*]pyridine. To a solution of *tert*-Butyl 4-([1,2,4]triazolo[4,3-*a*]pyridin-6-yl)piperidine-1-carboxylate (compound **36.1**, 0.05g, 0.165 mmol) in dichloromethane (2 mL) was added TFA (0.2 mL). The mixture was stirred at room temperature for 3 hours. The solvents were removed under reduced pressure. The residue was purified by prep-TLC (10% MeOH in dichloromethane + ~0.5% NH₄OH) to give 0.25 g (76%) of the title compound as a light brown oil. m/z (ES⁺) 203 (M+H)⁺.



Compound 36. (4-([1,2,4]Triazolo[4,3-*a*]pyridin-6-yl)piperidin-1-yl)(3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-methylphenyl)methanone. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 6-(piperidin-4-yl)-[1,2,4]triazolo[4,3-*a*]pyridine (compound **36.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 415 (M+H)⁺. ¹H NMR (400 MHz, Methanol-*d*₄) δ 9.12 (d, J = 0.8 Hz, 1H), 8.41 (d, J = 1.4 Hz, 1H), 7.73 (dt, J = 9.6, 1.0, 1.0 Hz, 1H), 7.53 (dd, J = 9.5, 1.6 Hz, 1H), 7.47 – 7.37 (m, 2H), 7.33 (d, J = 1.7 Hz, 1H), 4.84 (m, 1H), 4.00 (m, 1H), 3.00 (m, 2H), 2.42 (s, 3H), 2.32 (s, 3H), 2.14 (s, 3H), 2.10 (m, 1H), 1.96 (m, 1H), 1.77 (m, 3H).




Compound 37. 4-(1-(3-(2,4-Dimethyl-1*H*-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 399 (M+H)⁺. ¹H NMR (400 MHz, Methanol-*d*₄) δ 7.65 (d, J =8.4Hz, 2H), 7.48 (d, J =8.0Hz, 2H), 7.38 (d, J =8.0Hz, 1H), 7.34 (dd, J =7.20, 2.0Hz, 1H), 7.28 (d, J =2.0Hz, 1H) 4.84-4.71 (m, 1H), 4.02-3.89 (m, 1H), 3.03-2.91 (m, 2H), 2.36 (s, 3H), 2.29 (s, 3H), 2.09 (s, 3H), 2.00-1.64 (m, 5H).




Compound 38. (3-(2,4-Dimethyl-1*H*-imidazol-5-yl)-4-methylphenyl)(4-(imidazo[1,5-*a*]pyridin-7-yl)piperidin-1-yl)methanone. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 7-(piperidin-4-yl)imidazo[1,5-*a*]pyridine hydrochloride (compound **39.5**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 399 (M+H)⁺. ¹H NMR (400 MHz, Methanol-*d*₄) δ 8.25 (s, 1H), 8.18 (d,


$J=7.6\text{Hz}$, 1H), 7.43-7.34 (m, 3H), 7.31-7.27 (m, 2H), 6.68 (dd, $J=7.6$, 1.6Hz, 1H), 4.85-4.76 (m, 1H), 4.04-3.93 (m, 1H), 3.05-2.80 (m, 2H), 2.37 (s, 3H), 2.31 (s, 3H), 2.11 (s, 3H), 1.95-1.58 (m, 5H).



- 5 **Compound 39.1. (4-Bromopyridin-2-yl)methanamine.** To a 1-L round-bottom flask, was placed a solution of 4-bromopyridine-2-carbonitrile (10 g, 95%, 51.9 mmol; patent US 2009/0239876 A1, example 2) in tetrahydrofuran (220 mL), then BH_3 -THF complex (1 M) (330 mL) was added drop-wise with stirring at room temperature. The resulting solution was stirred overnight at room temperature, then carefully quenched with formic acid (100 mL).
- 10 The mixture was concentrated under reduced pressure to yield the title compound as the formate salt which was a light yellow solid and was used in the next step without further purification (8 g, crude).




- 15 **Compound 39.2. *N*-((4-Bromopyridin-2-yl)methyl)formamide.** To a 500-mL round-bottom flask, was placed (4-bromopyridin-2-yl)methanamine (compound **39.1**, 8.0 g, crude) and formic acid (200 mL). The solution was stirred for 2 h at 100 °C, then cooled and the pH was adjusted to 7 by careful and slow addition of aqueous sodium carbonate (sat.). The aqueous phase was extracted with ethyl acetate (3 x 200 mL) and the combined organic layers were washed with brine (3 x 30 mL), dried (Na_2SO_4), filtered, and concentrated under
- 20 reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate as the eluent to yield the title compound as a yellow oil (8.0 g, 90% pure, 65% yield over 2 steps).




- 25 **Compound 39.3. 7-bromoimidazo[1,5-*a*]pyridine.** To a 100-mL round-bottom flask, was placed a solution of *N*-((4-bromopyridin-2-yl)methyl)formamide (compound **39.2**, 3.0 g, 90%, 12.6 mmol) in dichloromethane (20 mL). The solution was cooled to 0-5 °C then

trifluoroacetic anhydride (1.93 mL, 13.9 mmol) was added drop-wise. The resulting solution was stirred for 1 h at room temperature, then the pH was adjusted to 7 by careful and slow addition of aqueous sodium carbonate (sat.). The aqueous phase was extracted with dichloromethane (3 x 20 mL) and the combined organic layers were washed with brine (3 x 20 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1/1) as the eluent to yield the title compound as a brown solid (1.0 g, 40%).




Compound 39.4. *tert*-Butyl 4-(imidazo[1,5-*a*]pyridin-7-yl)piperidine-1-

carboxylate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1.1**, except 7-bromoimidazo[1,5-*a*]pyridine (compound **39.3**) was used in place of 4-bromobenzonitrile.




Compound 39.5. 7-(Piperidin-4-yl)imidazo[1,5-*a*]pyridine hydrochloride.

The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1.2**, except *tert*-butyl 4-(imidazo[1,5-*a*]pyridin-7-yl)piperidine-1-carboxylate (compound **39.4**) was used in place of *tert*-butyl 4-(4-cyanophenyl)piperidine-1-carboxylate (compound **1.1**).




Compound 39. 2-Cyclobutyl-5-(4-(imidazo[1,5-*a*]pyridin-7-yl)piperidine-1-carbonyl)-*N*,4-dimethylbenzamide. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **62**, except 7-(piperidin-4-yl)imidazo[1,5-*a*]pyridine hydrochloride (compound **39.5**) was used

in place of 5-(piperidin-4-yl)-1*H*-indazole (compound **62.5**) to yield the title compound as a white solid. m/z (ES+) 431 (M+H)⁺.




Compound 40.1. (E)-N'-((4-Bromopyridin-2-yl)methylene)-4-

5 methylbenzenesulfonohydrazide. 4-Bromopyridinaldehyde (2.0 g, 10.8 mmol) and 4-methylbenzenesulfonohydrazide (2.0 g, 10.8 mmol) were mixed in MeOH (20 mL) and dichloromethane (20 mL). The mixture was stirred at room temperature for 1 hour. The solvents were removed under reduced pressure to give 3.80 g (theoretical) of the title compound as a yellow solid. m/z ES+ 354, 356 (M+H)⁺.




Compound 40.2. 5-Bromo-[1,2,3]triazolo[1,5-*a*]pyridine. A solution of (E)-N'-((4-bromopyridin-2-yl)methylene)-4-methylbenzenesulfonohydrazide (compound **40.1**, 3.8 g, 10.7 mmol) in morpholine (12 mL) was heated at 130 °C for 3 hours. The reaction mixture was cooled to room temperature, then diluted with EtOAc (150 mL) and washed with water (2 x 30 mL). The organic layer was dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexanes:EtOAc 4:1) to yield 2.10 g (99%) of the title compound as a light yellow solid. m/z (ES+) 198, 200 (M+H)⁺.




Compound 40.3. *tert*-Butyl 4-([1,2,3]triazolo[1,5-*a*]pyridin-5-yl)piperidine-1-carboxylate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1.1**, except *tert*-butyl 4-

([1,2,3]triazolo[1,5-*a*]pyridin-5-yl)piperidine-1-carboxylate (compound **40.2**) was used in place of 4-bromobenzonitrile. m/z (ES⁺) 303(M+H)⁺.




Compound 40.4. 5-(Piperidin-4-yl)-[1,2,3]triazolo[1,5-*a*]pyridine. The title

compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **36.2**, except *tert*-butyl 4-([1,2,3]triazolo[1,5-*a*]pyridin-5-yl)piperidine-1-carboxylate (compound **40.3**) was used in place of *tert*-butyl 4-([1,2,4]triazolo[4,3-*a*]pyridin-6-yl)piperidine-1-carboxylate (compound **36.1**). m/z (ES⁺) 203(M+H)⁺.




Compound 40. (4-([1,2,3]Triazolo[1,5-*a*]pyridin-5-yl)piperidin-1-yl)(3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylphenyl)methanone. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 5-(piperidin-4-yl)-[1,2,3]triazolo[1,5-*a*]pyridine (compound **40.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 415 (M+H)⁺. ¹H NMR (400 MHz, Methanol-*d*₄) δ 8.84 (d, J =7.2Hz, 1H), 8.07 (d, J =0.8Hz, 1H), 7.77 (t, J =0.8, 0.8Hz, 1H), 7.44-7.34 (m, 2H), 7.32 (d, J =1.6Hz, 1H), 7.18 (dd, J =7.6, 2.0Hz, 1H), 4.87-4.76 (m, 1H), 4.08-3.92 (m, 1H), 3.11-2.95 (m, 2H), 2.37 (s, 3H), 2.31 (s, 3H), 2.11 (s, 3H), 2.00-1.67 (m, 5H).




Compound 41.1. 4-Bromo-2-hydrazinylpyridine. To a solution of 4-bromo-2-fluoropyridine (2.0 g, 11.4 mmol) in pyridine (10 mL) was added hydrazine (5 mL, 159 mmol). The mixture was heated at 70 °C for 2 hours, then cooled to room temperature. The volatile organics were removed under reduced pressure, then water (60 mL) was added to the

residue and an off-white solid precipitated. The solid was filtered, washed with water, and dried under reduced pressure at 50 °C to give 1.77 g (84%) of the title compound as an off-white solid. m/z (ES+) 188, 190 (M+H)⁺.




Compound 41.2. 7-Bromo-[1,2,4]triazolo[4,3-*a*]pyridine. 4-Bromo-2-

hydrazinylpyridine (compound **41.1**) was suspended in formic acid (3 mL). The mixture was heated at 100 °C for one hour, then upon complete reaction, the mixture cooled to room temperature. The volatile organics were removed under reduced pressure, then water (50 mL) was added to the residue. The solids that formed were filtered, washed with water and dried under reduced pressure at 50 °C to give 1.68 g (90%) of the title compound as an off-white solid. m/z (ES+) 198, 200 (M+H)⁺.



Compound 41.3. *tert*-Butyl 4-([1,2,4]triazolo[4,3-*a*]pyridin-7-yl)piperidine-1-


carboxylate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1.1**, except 7-bromo-[1,2,4]triazolo[4,3-*a*]pyridine (compound **41.2**) was used in place of 4-bromobenzonitrile. m/z (ES+) 303(M+H)⁺.



Compound 41.4. 7-(Piperidin-4-yl)-[1,2,4]triazolo[4,3-*a*]pyridine. The title


compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **36.2**, except *tert*-butyl 4-([1,2,4]triazolo[4,3-*a*]pyridin-7-yl)piperidine-1-carboxylate (compound **41.3**) was used in place of *tert*-butyl 4-

([1,2,4]triazolo[4,3-*a*]pyridin-6-yl)piperidine-1-carboxylate (compound **36.1**) m/z (ES+) 203(M+H)⁺.



Compound 41. (4-([1,2,4]Triazolo[4,3-*a*]pyridin-7-yl)piperidin-1-yl)(3-(2,4-


5 dimethyl-1*H*-imidazol-5-yl)-4-methylphenyl)methanone. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 7-(piperidin-4-yl)-[1,2,4]triazolo[4,3-*a*]pyridine (compound **41.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 415 (M+H)⁺. ¹H NMR (400 MHz, Methanol-*d*₄) δ 9.13 (s, 1H), 8.45 (d, $J=8.0$ Hz, 1H), 7.59 (s, 1H), 7.44-7.36 (m, 2H), 7.33 (d, $J=1.6$ Hz, 1H), 7.07 (dd, $J=7.2, 1.6$ Hz, 1H), 4.87-4.78 (m, 1H), 4.08-3.96 (m, 1H), 3.11-2.95 (m, 2H), 2.40 (s, 3H), 2.32 (s, 3H), 2.13 (s, 3H), 2.10-1.73 (m, 5H).



Compound 42.1. *tert*-Butyl 4-(4-chlorophenyl)-4-hydroxypiperidine-1-


15 carboxylate. Into a 1-L 3-necked round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 1-chloro-4-iodobenzene (10.0 g, 41.9 mmol) in tetrahydrofuran (150 mL). The solution was cooled to -78 °C, then *n*-BuLi (2.4 M) (16.6 mL, 39.8 mmol) was added dropwise and the resulting mixture was stirred for 0.5 h at -78 °C. A solution of *tert*-butyl 4-oxopiperidine-1-carboxylate (7.60 g, 38.1 mmol) in tetrahydrofuran (50 mL) was added dropwise and the resulting mixture was stirred for 1 h at -78 °C. The reaction was then warmed to 0 °C, and carefully quenched by the slow addition of water (150 mL). The layers were separated and the aqueous phase was extracted with ethyl acetate (150 mL). The combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column

chromatography with ethyl acetate/petroleum ether (1:50 to 1:3) to yield the title compound as a light yellow solid (9.3 g, 78%).




Compound 42.2. *tert*-Butyl 4-(4-chlorophenyl)-5,6-dihydropyridine-1(2*H*)-


carboxylate. Into a 250-mL round-bottom flask, was placed a solution of *tert*-butyl 4-(4-chlorophenyl)-4-hydroxypiperidine-1-carboxylate (compound **42.1**, 9.00 g, 28.9 mmol) in pyridine (50 mL). With stirring, phosphoroyl trichloride (7.93 mL, 84.8 mmol) was added drop-wise. The resulting mixture was stirred overnight at room temperature and then concentrated under reduced pressure. The residue was diluted with ethyl acetate (20 mL) and carefully quenched by slow addition of aqueous sodium bicarbonate. The layers were separated and the organic layer was washed with aqueous sodium bicarbonate (2 x 30 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:100 to 1:10) to yield the title compound as a colorless oil (6.1 g, 72%).



Compound 42.3. *tert*-Butyl 4-(4-chlorophenyl)piperidine-1-carboxylate. Into a 250-mL round-bottom flask, was placed a mixture of Pt₂O (200 mg) and methanol (50 mL). The mixture was purged with nitrogen, then hydrogen was introduced and the mixture was stirred for 15 min. A solution of *tert*-butyl 4-(4-chlorophenyl)-5,6-dihydropyridine-1(2*H*)-carboxylate (compound **42.2**, 6.00 g, 20.4 mmol) in methanol (50 mL) was added and the resulting mixture was stirred under hydrogen overnight at room temperature. After purging with nitrogen, the solids were filtered off and the resulting solution was concentrated under vacuum to yield the title compound as a light green oil (5.30 g, 88%).




Compound 42.4. 4-(4-Chlorophenyl)piperidine. Into a 250-mL round-bottom flask, was placed a solution of *tert*-butyl 4-(4-chlorophenyl)piperidine-1-carboxylate (compound 42.3, 5.00 g, 16.9 mmol) in dichloromethane (100 mL), trifluoroacetic acid (9.6 g, 84 mmol). The resulting solution was stirred overnight at room temperature and then concentrated under reduced pressure. The residue was carefully diluted with ethyl acetate (100 mL) and aqueous sodium bicarbonate was added until a pH of 8 was attained. The resulting mixture was washed with brine (100 mL) and the organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield the title compound as a light yellow solid (2.30 g, 70%).



Compound 43.1. *tert*-Butyl 3-(4-bromophenyl)-3-hydroxyazetidine-1-carboxylate.


Into a 1-L 3-necked round-bottom flask, purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 1-bromo-4-iodobenzene (25.0 g, 88.4 mmol) in tetrahydrofuran/diethyl ether (400/200 mL). The solution was cooled to -78 °C then *n*-BuLi (2.5 M, 37.1 mL, 92.8 mmol) was added drop-wise over 10 min. To the resulting mixture was added *tert*-butyl 3-oxoazetidine-1-carboxylate (16.6 g, 97.0 mmol) in THF (100 mL) drop-wise at -78 °C. The resulting mixture was stirred for 1.5 h at -78 °C, then carefully quenched with water (300 mL). The aqueous phase was extracted with ethyl acetate (200 mL) and the combined organic layers were washed with brine (2 x 100 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:20-1:5) as the eluent followed by re-crystallized from ethyl acetate:PE in the ratio of 1:100 to yield 8.0 g (28%) of the title compound as a white solid.



Compound 43.2. *tert*-Butyl 3-(4-cyanophenyl)-3-hydroxyazetidine-1-carboxylate.


Into a 500-mL round-bottom flask, purged and maintained with an inert atmosphere of nitrogen, was placed a mixture of *tert*-butyl 3-(4-bromophenyl)-3-hydroxyazetidine-1-

carboxylate (compound **43.1**, 16.3 g, 49.7 mmol) in *N,N*-dimethylformamide (250 mL), zinc cyanide (8.7 g, 75 mmol) and Pd(PPh₃)₄ (5.77 g, 5.00 mmol). The resulting mixture was stirred for 15 h at 100 °C, then cooled to room temperature. The reaction was quenched with saturated aqueous FeSO₄ (500 mL) and stirred vigorously. The mixture was filtered through
 5 Celite® and the layers from the filtrate were separated. The aqueous phase was extracted with ethyl acetate (300 mL) and the combined organic layers were washed with brine (2 x 100 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1/100-1/3) as the eluent to yield 14 g (crude) of the title compound as a white solid.




Compound 43.3. *tert*-Butyl 3-(4-cyanophenyl)-3-fluoroazetidine-1-carboxylate.

Into a 250-mL 3-necked round-bottom flask, purged and maintained with an inert atmosphere of nitrogen, was placed a solution of *tert*-butyl 3-(4-cyanophenyl)-3-hydroxyazetidine-1-carboxylate (compound **43.2**, 5.00 g, 18.2 mmol) in dichloromethane (120 mL). The solution
 15 was cooled to -78 °C and Deoxo-Fluor® (bis(2-methoxyethyl)aminosulfur trifluoride) (5.99 g, 27.1 mmol) was added drop-wise. The resulting mixture was stirred for 1.5 h at -78 °C, then carefully quenched with sodium bicarbonate (50 mL, 1 M). The organic layer was additionally washed with brine (2 x 50 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl
 20 acetate/petroleum ether (1/50-1/30) as the eluent to yield 3.2 g (64%) of the title compound as colorless oil.




Compound 43.4. 4-(3-Fluoroazetidin-3-yl)benzonitrile hydrochloride. Into a 100-mL round-bottom flask, was placed a solution of *tert*-butyl 3-(4-cyanophenyl)-3-fluoroazetidine-1-carboxylate (compound **43.3**, 1.00 g, 3.62 mmol) in dioxane (10 mL) and
 25 HCl in dioxane (4 M in dioxane, 10 mL, 40 mmol). The resulting mixture was stirred for 1 h at 60 °C, then cooled and concentrated under reduced pressure. The residue was washed with


EtOAc (20 mL) and the product solids were collected by filtration to yield 522 mg (68%) of the title compound as a white solid. m/z (ES⁺) 177 (M+H)⁺. ¹H NMR (300 MHz, CD₃OD): δ 7.91 (d, J = 7.8 Hz, 2H), 7.82 (d, J = 8.1 Hz, 2H), 4.85-4.52 (m, 4H).




5 **Compound 44.1. 2-Cyclopropyl-1H-imidazole.** Into a 25-mL round-bottom flask, was placed a solution of cyclopropanecarbaldehyde (500 mg, 7.13 mmol), oxaldehyde (455 mg, 7.84 mmol) in methanol (5 mL). The solution was cooled to 0 °C, then 25% ammonium hydroxide (1 mL) was added drop-wise. The resulting solution was stirred for 3 h at 0 °C, then stirred at room temperature overnight. The resulting mixture was concentrated under
10 reduced pressure and the residue was dissolved in brine (50 mL). The aqueous phase was extracted with ethyl acetate (3 x 10 mL) and the combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield 600 mg (78%) of the title compound as a light brown solid.



15 **Compound 44.2. 2-Cyclopropyl-4,5-diiodo-1H-imidazole.** Into a 100-mL round-bottom flask, was placed a solution of 2-cyclopropyl-1H-imidazole (compound **44.1**, 1.8 g, 16.6 mmol) in sodium hydroxide (2 M, 40 mL). A solution of iodine (8.5 g, 33.5 mmol) in dichloromethane (40 mL) was added drop-wise and the resulting mixture was stirred overnight at room temperature. The aqueous layer was separated and neutralized with acetic
20 acid and quenched by the addition of Na₂S₂O₃ (sat. aq.). The solids were collected by filtration to yield 3.8 g (63%) of the title compound as a brown solid.



Compound 44.3. 2-Cyclopropyl-4-iodo-1H-imidazole. Into a 100-mL round-bottom flask, was placed a solution of sodium sulfite (11.3 g, 89.7 mmol) in H₂O/EtOH (30/15 mL).
25 2-Cyclopropyl-4,5-diiodo-1H-imidazole (compound **44.2**, 3.8 g, 10.6 mmol) was added and the resulting solution was heated at reflux overnight. The reaction was cooled and the volatiles were removed under reduced pressure. The solids were collected by filtration to yield 1.8 g (73%) of the title compound as a light brown solid.










Compound 45.1. 5-Iodo-2-isopropyl-1H-imidazole. The title compound was prepared using standard chemical manipulations and a procedure similar to that used for the preparation of compound **44.3**, except 2-isopropyl-1H-imidazole was used in place of 2-cyclopropyl-1H-imidazole (compound **44.1**).


The compounds in **TABLE 1** were prepared using standard chemical manipulations and procedures with readily available starting materials or building blocks described in this manuscript. The utilized procedures were similar to that used for the preparation of compound **7** using the respective imidazoles (2-methyl-4-bromo imidazole, compound **44.3**, or compound **45.1**) and the respective amines (compound **1.2**, compound **13.4**, compound **42.4**, compound **5.2**, or compound **43.4**).

TABLE 1

Cpd	Name	Structure	<i>m/z</i> (ES+) (M+H) ⁺
46	4-(1-(3-(4-chloro-2-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile		419
13	4-(1-(3-(4-chloro-2-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)-4-fluoropiperidin-4-yl)benzonitrile		437

Cpd	Name	Structure	m/z (ES ⁺) (M+H) ⁺
42	(3-(4-chloro-2-methyl-1 <i>H</i> -imidazol-5-yl)-4-methylphenyl)(4-(4-chlorophenyl)piperidin-1-yl)methanone		408
44	4-(1-(3-(4-chloro-2-cyclopropyl-1 <i>H</i> -imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		417
47	4-(1-(3-(4-chloro-2-cyclopropyl-1 <i>H</i> -imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile		445
43	4-(1-(3-(4-chloro-2-cyclopropyl-1 <i>H</i> -imidazol-5-yl)-4-methylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile		435


Cpd	Name	Structure	<i>m/z</i> (ES+) (M+H) ⁺
48	4-(1-(3-(4-chloro-2-cyclopropyl-1 <i>H</i> -imidazol-5-yl)-4-methylbenzoyl)-4-fluoropiperidin-4-yl)benzonitrile		463
45	4-(1-(3-(4-chloro-2-isopropyl-1 <i>H</i> -imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		419
49	4-(1-(3-(4-chloro-2-isopropyl-1 <i>H</i> -imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile		447



Compound 50.1. Methyl 4-chloro-3-iodo-benzoate. 4-Chloro-3-iodo-benzoic acid


(5.31 g, 18.8 mmol) was dissolved in methanol (50 mL) and concentrated sulfuric acid (3 mL) was carefully added. The solution was stirred at 80 °C for 4 hours, then cooled to room temperature and the volatile organics were removed under reduced pressure. The residue was partitioned between EtOAc (50 mL) and water (50 mL) and the organic layer was washed with brine, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by flash chromatography (SiO₂; 0-10 % EtOAc in hexanes) to yield 5.32 g (95

%) of the title compound as an oil. ^1H NMR (400 MHz, Chloroform- d) δ 8.54 (d, J = 2.0 Hz, 1H), 7.96 (dd, J = 8.4, 2.0 Hz, 1H), 7.53 (d, J = 8.4 Hz, 1H), 3.94 (s, 3H).




Compound 50.2. Methyl 4-chloro-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate.

5 **yl)benzoate.** A mixture of methyl 4-chloro-3-iodobenzoate (compound **50.1**, 3.85 g, 13.0 mmol), bis(pinacolato)diboron (3.96 g, 15.6 mmol), Pd(dppf)Cl₂·DCM (531 mg, 0.65 mmol) and potassium acetate (3.83 g, 39.0 mmol) in DMSO (40 mL) was degassed with argon and then heated to 80 °C for 18 hours. The reaction mixture was cooled then diluted with ethyl acetate (200 mL) and sequentially washed with water, aqueous HCl (1 M), saturated aqueous
10 NaHCO₃, and brine, then dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was purified by flash chromatography (SiO₂; 0-10 % EtOAc in hexanes) to yield 1.92 g (49 %) of the title compound as a white solid. ^1H NMR (400 MHz, Chloroform- d) δ 8.36 (d, J = 2.3 Hz, 1H), 8.01 (dd, J = 8.4, 2.3 Hz, 1H), 7.44 (d, J = 8.4 Hz, 1H), 3.94 (s, 3H), 1.40 (s, 12H).




15 **Compound 50.3. Methyl 4-chloro-3-(2,4-dimethyl-1H-imidazol-5-yl)benzoate.** To methyl 4-chloro-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **50.2**, 600 mg, 2.02 mmol) in dioxane (20 mL) was added 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**, 538 mg, 2.42 mmol) and Pd(dppf)Cl₂·DCM (165 mg, 0.20 mmol). The
20 mixture was degassed with argon and stirred for 10 minutes at room temperature, then an aqueous potassium carbonate solution (1M, 10 mL) was added and the mixture was stirred at 90 °C for 18 h. The mixture was cooled and diluted with EtOAc, then filtered through Celite®. The filtrate was washed with brine, dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was purified by flash chromatography (SiO₂; 0-100 % EtOAc
25 in hexanes) to yield 270 mg (50 %) of the title compound as a foam. m/z (ES⁺) 265 (M+H)⁺. ^1H NMR (400 MHz, Chloroform- d) δ 8.07 (d, J = 2.2 Hz, 1H), 7.90 (dd, J = 8.4, 2.2 Hz, 1H), 7.50 (d, J = 8.4 Hz, 1H), 3.91 (s, 3H), 2.40 (s, 3H), 2.19 (s, 3H).




Compound 50.4. 4-Chloro-3-(2,4-dimethyl-1H-imidazol-5-yl)benzoic acid.

Methyl 4-chloro-3-(2,4-dimethyl-1H-imidazol-5-yl)benzoate (compound **50.3**, 270 mg, 1.02 mmol) was dissolved in methanol (20 mL) and aqueous NaOH (2 M, 6 mL) then heated to 50 °C for 16 hrs. The volatile solvents were removed under reduced pressure and the resulting aqueous phase was acidified to pH 5-6 with aqueous HCl (2M). The precipitated solids were filtered, and dried to yield 230 mg (94%) of the title compound as a white solid. m/z (ES-) 249 (M-H)⁻.



Compound 50. 4-(1-(4-Chloro-3-(2,4-dimethyl-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile.


The title compound was obtained as a white solid (77 mg, 54%) using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 4-chloro-3-(2,4-dimethyl-1H-imidazol-5-yl)benzoic acid (compound **50.4**, 85 mg, 0.34 mmol) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 419 (M+H)⁺.




Compound 51.1. Methyl 4-methyl-3-(1H-pyrazol-5-yl)benzoate.

To methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**, 800 mg, 2.9 mmol) in dioxane (30 mL) was added 5-iodo-1H-pyrazole (674 mg, 3.5 mmol), and Pd(dppf)Cl₂·DCM (237 mg, 0.29 mmol). The mixture was degassed with argon and stirred for 10 minutes then an aqueous potassium carbonate solution (2 M, 8 mL) was added. The mixture was heated at 90 °C for 18 h, then cooled and diluted with EtOAc and filtered


through Celite®. The filtrate was washed with brine, dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was purified by flash chromatography (SiO₂; 0-30 % EtOAc in hexanes) to yield 258 mg (41 %) of the title compound as a solid. m/z (ES⁺) 217 (M+H)⁺.



Compound 51.2. Methyl 3-(4-chloro-1H-pyrazol-5-yl)-4-methylbenzoate. Methyl 4-methyl-3-(1H-pyrazol-5-yl)benzoate (compound **51.1**, 385 mg, 1.78 mmol) was dissolved in 1,2-dichloroethane (60 mL), then *N*-chlorosuccinimide (250 mg, 1.87 mmol) was added. The mixture was stirred at room temperature for 16 hours then washed with brine, dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was purified by flash chromatography (SiO₂; 0-20 % EtOAc in hexanes) to yield 152 mg (34 %) of the title compound as an oil. m/z (ES⁺) 251 (M+H)⁺. ¹H NMR (400 MHz, Chloroform-d) δ 8.07 – 7.94 (m, 2H), 7.63 (d, J = 0.7 Hz, 1H), 7.47 – 7.34 (m, 1H), 3.93 (s, 3H), 2.36 (s, 3H).




Compound 51.3. 3-(4-Chloro-1H-pyrazol-5-yl)-4-methylbenzoic acid. Methyl 3-(4-chloro-1H-pyrazol-5-yl)-4-methylbenzoate (compound **51.2**, 133 mg, 0.53 mmol) was dissolved in a mixture of aqueous NaOH (2 M, 3 mL) and methanol (10 mL). The solution was heated at 50 °C for 16 hrs then the volatiles were removed under reduced pressure. Aqueous HCl (2 M) was added to adjust the pH to 4-5 then concentrated to yield 150 mg of a white solid which was used in the next step without further purification. m/z (ES⁻) 235 (M-H)⁻.




Compound 51. 4-(1-(3-(4-Chloro-1H-pyrazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was obtained as a white solid (69 mg, 34% over 2 steps)


using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 3-(4-chloro-1*H*-pyrazol-5-yl)-4-methylbenzoic acid (compound **51.3**, ~0.53 mmol) was used in place of 3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). m/z (ES+) 377 (M+H)⁺.




Compound 52. 4-(1-(3-(4-Chloro-1*H*-pyrazol-5-yl)-4-methylbenzoyl)-4-fluoropiperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for preparation of compound **5**, except 3-(4-chloro-1*H*-pyrazol-5-yl)-4-methylbenzoic acid (compound **51.3**) was used in place of 3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(4-fluoropiperidin-4-yl)benzonitrile hydrochloride (compound **13.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 423 (M+H)⁺.



Compound 53. 4-(1-(3-(4-Chloro-1*H*-pyrazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for preparation of compound **5**, except 3-(4-chloro-1*H*-pyrazol-5-yl)-4-methylbenzoic acid (compound **51.3**) was used in place of 3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 405 (M+H)⁺.



Compound 54. 4-(1-(3-(4-Chloro-3-methyl-1*H*-pyrazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for preparation of compound **51** and compound **5**, except 3-bromo-5-methyl-1*H*-pyrazole was used in place of 5-iodo-1*H*-pyrazole. m/z (ES+) 391 (M+H)⁺. ¹H NMR (300 MHz, Methanol-d₄): δ 7.77-7.69 (m, 3H), 7.64-7.59 (m, 3H), 7.46 (d, J = 8.1 Hz, 1H), 4.84 (m, 1H), 4.64 (m, 1H), 4.48 (m, 1H), 4.23 (m, 1H), 4.09 (m, 1H), 2.34 (s, 3H), 2.33 (s, 3H).







Compound 55. 4-(1-(3-(4-Chloro-3-methyl-1*H*-pyrazol-5-yl)-4-methylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for preparation of compound **51** and compound **5**, except 3-bromo-5-methyl-1*H*-pyrazole was used in place of 5-iodo-1*H*-pyrazole and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 409 (M+H)⁺.

The compounds in **TABLE 2** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **51**, **52**, **53**, **54** and **55**.

TABLE 2

Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
94	4-(1-(5-(4-chloro-1 <i>H</i> -pyrazol-5-yl)-2,4-dimethylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile		409


Cpd	Name	Structure	<i>m/z</i> (ES+) (M+H) ⁺
136	4-(1-(5-(4-chloro-3-methyl-1 <i>H</i> -pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile		405
140	4-(1-(3-(4-chloro-3-methyl-1 <i>H</i> -pyrazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile		419
141	4-(1-(3-(4-chloro-3-methyl-1 <i>H</i> -pyrazol-5-yl)-4-methylbenzoyl)-4-fluoropiperidin-4-yl)benzonitrile		437
142	4-(1-(5-(4-chloro-1 <i>H</i> -pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile		391




Compound 56.1. 5-Iodo-4-methyl-1*H*-pyrazole. To a solution of 4-methyl-1*H*-

pyrazole (2.15 g, 26.1 mmol) dissolved in DMF (20 mL) was added *N*-iodosuccinimide (6.19 g, 26.1 mmol). The mixture was stirred at RT for 16 hours and then diluted with water and filtered. The filtrate was extracted with EtOAc (2 x 50 mL), then the combined organic extracts were washed with brine, dried (Na₂SO₄), filtered, and concentrated under reduced


pressure. The residue was purified by flash chromatography (SiO₂; 0-50 % EtOAc in hexanes) to yield 2.19 g (41%) of the title compound as a white solid. m/z (ES+) 209 (M+H)⁺.



Compound 56. 4-(1-(4-Methyl-3-(4-methyl-1H-pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for preparation of compound **5**, except 5-iodo-4-methyl-1H-pyrazole (compound **56.1**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**) to yield the title compound as white solid. m/z (ES+) 357 (M+H)⁺.







Compound 57. 4-(1-(4-Methyl-3-(4-methyl-1H-pyrazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for preparation of compound **5**, except 5-iodo-4-methyl-1H-pyrazole (compound **56.1**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**) and 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 385 (M+H)⁺.




Compound 58. 4-(3-Fluoro-1-(4-methyl-3-(4-methyl-1H-pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for preparation of compound **5**, except 5-iodo-4-methyl-1H-pyrazole (compound **56.1**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 375 (M+H)⁺.


The compounds in **TABLE 3** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **51**, **56**, **57** and **58**.

5 **TABLE 3**


Cpd	Name	Structure	<i>m/z</i> (ES+) (M+H) ⁺
145	4-(1-(4-methyl-3-(1 <i>H</i> -pyrazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile		371
146	4-(1-(2,4-dimethyl-5-(4-methyl-1 <i>H</i> -pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		371
147	4-(1-(2,4-dimethyl-5-(4-methyl-1 <i>H</i> -pyrazol-5-yl)benzoyl)-3-fluoroazetidin-3-yl)benzonitrile		389
148	4-(1-(2,4-dimethyl-5-(4-methyl-1 <i>H</i> -pyrazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile		399




Compound 59.1. 5-Iodo-3,4-dimethyl-1H-pyrazole. NIS (5.62g, 24.9mmol) was added portion-wise to a solution of 3,4-dimethyl-1H-pyrazole (2.0 g, 20.8 mmol) in CH₃CN (50 mL). The mixture was heated at 80 °C for 16 hours, upon where thick off-white solids formed. The mixture was cooled to room temperature and the solid were filtered, washed with cold CH₃CN and dried under reduced pressure to yield 4.32g (94%) of the title compound as an off-white solid. *m/z* (ES⁺) 223 (M+H)⁺.



Compound 59.2. Methyl 3-(3,4-dimethyl-1H-pyrazol-5-yl)-4-methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 5.6, except 5-iodo-3,4-dimethyl-1H-pyrazole (compound 59.1) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound 5.5). *m/z* (ES⁺) 245 (M+H)⁺.




Compound 59.3. 3-(3,4-Dimethyl-1H-pyrazol-5-yl)-4-methylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 5.7, except methyl 3-(3,4-dimethyl-1H-pyrazol-5-yl)-4-methylbenzoate (compound 59.2) was used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound 5.6). *m/z* (ES⁺) 231 (M+H)⁺.




Compound 59. 4-(1-(3-(3,4-Dimethyl-1H-pyrazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations

and procedures similar to those used for the preparation of compound **5**, except 3-(3,4-dimethyl-1*H*-pyrazol-5-yl)-4-methylbenzoic acid (compound **59.3**) was used in place of 3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). m/z (ES⁺) 371 (M+H)⁺. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.43-12.17 (br, 1H), 7.83 (d, J =8.4Hz, 2H), 7.63 (d, J =8.4Hz, 2H), 7.60 (dd, J =8.4Hz, 2.0Hz, 1H), 7.47 (d, J =1.6Hz, 1H), 7.38 (d, J =8.0Hz, 1H), 4.71 (m, 1H), 4.54-4.35 (m, 2H), 4.09-4.00 (m, 2H), 2.24 (s, 3H), 2.17 (s, 3H), 1.83 (s, 3H).




Compound 60. 4-(1-(3-(3,4-Dimethyl-1*H*-pyrazol-5-yl)-4-methylbenzoyl)-3-


fluoroazetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 3-(3,4-dimethyl-1*H*-pyrazol-5-yl)-4-methylbenzoic acid (compound **59.3**) and 4-(3-fluoroazetidin-3-yl)benzonitrile (compound **43.4**) were used in place of 3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**), respectively. m/z (ES⁺) 389 (M+H)⁺. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.45-12.26 (br, 1H), 7.95 (d, J =8.0Hz, 2H), 7.79 (d, J =8.0Hz, 2H), 7.65 (d, J =7.2Hz, 1H), 7.52 (d, J =2.0Hz, 1H), 7.40 (d, J =8.0Hz, 1H), 4.98-4.42 (m, 4H), 2.25 (s, 3H), 2.17 (s, 3H), 1.83 (s, 3H).




Compound 61. 4-(1-(4-Methyl-3-(5-methyl-1*H*-pyrazol-4-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for preparation of compound **5**, except 4-bromo-5-methyl-1*H*-pyrazole was used in place of 5-iodo-2,4-dimethyl-1*H*-imidazole (compound **5.5**) and 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 385 (M+H)⁺.




Compound 62.1. 4-Cyclobutyl-5-iodo-2-methylbenzoic acid. To a solution of methyl 4-cyclobutyl-5-iodo-2-methylbenzoate (compound **6.3**, 35.0 g, 106 mmol) in methanol (200 mL) at 0-5 °C was added aqueous sodium hydroxide (12.7 g, 318 mmol in 100 mL water) drop-wise. The resulting mixture was stirred for 3 h at 60 °C, then cooled to ambient temperature and the volatile organics were removed under reduced pressure. The pH of the remaining aqueous material was adjusted to ~4 with hydrogen chloride (aqueous, 2 M). The resulting solids were collected by filtration and dried in an oven under reduced pressure to yield the title compound as a white solid (31.0 g, 93%).




Compound 62.2. 4-Cyclobutyl-5-(methoxycarbonyl)-2-methylbenzoic acid. Into a 50-mL high pressure autoclave reactor, was placed a solution of 4-cyclobutyl-5-iodo-2-methylbenzoic acid (compound **62.1**, 1.50 g, 4.74 mmol) in methanol (20 mL). Pd(dppf)Cl₂ (320 mg, 0.44 mmol) and triethylamine (1.27 mL, 9.09 mmol) were added and carbon monoxide (gas, 40 atm) was introduced. (**CAUTION:** Highly toxic gas at high pressure. All necessary safety precautions were performed). The resulting mixture was stirred overnight at 90 °C, then cooled to room temperature. The reaction was vented carefully using the necessary precautions, then the resulting mixture was concentrated under reduced pressure. The residue was dissolved in water (20 mL) and the pH of the solution was adjusted to 3-4 with aqueous HCl (1 M) and the resulting mixture was extracted with ethyl acetate (3 x 20 mL). The combined organic extracts were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1/2-1/1) as the eluent to yield the title compound as a white solid (1.0 g, 85%).




Compound 62.3. 4-Cyclobutyl-2-methyl-5-(methoxycarbonyl)benzoic acid. Into a 20-mL sealed tube, was placed 4-cyclobutyl-5-(methoxycarbonyl)-2-methylbenzoic acid (compound **62.2**, 1.0 g, 4.0 mmol) and methylamine (30% in ethanol) (8 mL). The resulting solution was stirred overnight at 120 °C behind a blast shield, then cooled and diluted with H₂O (20 mL). The pH of the solution was adjusted to 4-5 with hydrogen chloride (1 M) and the solids were collected by filtration. The crude product was re-crystallized from ethyl acetate/petroleum ether in the ratio of 1:10 to yield the title compound as an off-white solid (500 mg, 50%).




Compound 62.4. *tert*-Butyl 4-(1H-indazol-5-yl)piperidine-1-carboxylate. Into a 50-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a suspension of Zn (990 mg, 15.1 mmol) in DMA (1 mL). A 7:5 v/v mixture of TMSCl/1,2-dibromoethane (0.12 mL) was added drop-wise at a rate to maintain the temperature below 65 °C, then the mixture was stirred for an additional 10 min. A solution of *tert*-butyl 4-iodopiperidine-1-carboxylate (3.17 g, 10.2 mmol) in DMA (2 mL) was added drop-wise and stirred at 40-45 °C for 30 min. The resulting mixture was added to a mixture of 5-bromo-1H-indazole (1.00 g, 5.08 mmol), CuI (80 mg, 0.42 mmol) and Pd(dppf)Cl₂ (260 mg, 0.36 mmol) in DMA (1 mL) in a 50-mL round-bottom flask under a nitrogen atmosphere. The resulting mixture was stirred overnight at 85 °C, then cooled and diluted with ethyl acetate (50 mL) and washed with brine (3 x 20 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1/5) as the eluent to yield the title compound as a yellow solid (200 mg, 12%).




Compound 62.5. 5-(Piperidin-4-yl)-1*H*-indazole. Into a 50-mL round-bottom flask, was placed a solution of *tert*-butyl 4-(1*H*-indazol-5-yl)piperidine-1-carboxylate (compound **62.4**, 200 mg, 0.60 mmol) in dichloromethane (3 mL) and trifluoroacetic acid (1 mL). The resulting solution was stirred for 1 h at room temperature, then the pH of the solution was carefully adjusted to 8-9 with aqueous sodium bicarbonate (aq. sat.). The mixture was extracted with dichloromethane (3 x 30 mL) and the combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield the title compounds as a yellow oil (100 mg, 75%).




Compound 62. 5-(4-(1*H*-Indazol-5-yl)piperidine-1-carbonyl)-2-cyclobutyl-*N,N*-dimethylbenzamide. Into a 50-mL round-bottom flask, was placed a solution of 5-(piperidin-4-yl)-1*H*-indazole (compound **62.5**, 100 mg, 0.50 mmol) in *N,N*-dimethylformamide (3 mL). EDC•HCl (192 mg, 1.00 mmol), 4-dimethylaminopyridine (122 mg, 1.00 mmol) and 4-cyclobutyl-2-methyl-5-(methylcarbamoyl)benzoic acid (compound **62.3**, 122 mg, 0.49 mmol) were added and the resulting solution was stirred for 4 h at room temperature. The reaction was diluted with water (20 mL) and extracted with of ethyl acetate (2 x 25 mL). The combined organic layers were washed with brine (2 x 15 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate as the eluent, followed by additional purification by Prep-HPLC with the following conditions (1#-Pre-HPLC-001(SHIMADZU)): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150 mm; mobile phase, water with 0.05% TFA and CH₃CN (30% CH₃CN up to 47% in 7 min, up to 100% in 3 min, down to 30% in 1 min); Detector, Waters 2489, 254 & 220 nm. The fractions containing pure compound were combined and lyophilized to yield the title compound as a white solid (71.2 mg, 33%). *m/z* (ES⁺) 431 (M+H)⁺.




Compound 63.1. 1-Bromo-3-(2,2-diethoxyethoxy)benzene. Into a 3-L three neck round-bottom flask, was placed a solution of 3-bromophenol (50.00 g, 289.0 mmol) in *N,N*-dimethylformamide (1 L). The system was purged with nitrogen and the solution was cooled to 0 °C, then sodium hydride (60%, 12.8 g, 320 mmol) was added portion-wise. To the resulting mixture was added 2-bromo-1,1-diethoxyethane (53.1 mL, 345 mmol) drop-wise at 0 °C. The resulting mixture was stirred overnight at 120 °C, behind a blast shield (**CAUTION:** NaH and DMF can become a runaway reaction. All necessary safety precautions were performed). The mixture was cooled to room temperature, then diluted with ethyl acetate (3 L). The mixture was washed with brine (4 x 500 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1/50-1/30) as the eluent to yield the title compound as a light yellow oil (78.9 g, 94%).



Compound 63.2 and compound 63.3. 6-Bromo-3a,7a-dihydrobenzofuran and 4-bromo-3a,7a-dihydrobenzofuran. Into a 1-L round-bottom flask, was carefully placed a mixture of 1-bromo-3-(2,2-diethoxyethoxy)benzene (compound **63.1**, 62.9 g, 218 mmol) and polyphosphoric acid (157 g) in chlorobenzene (320 mL). The mixture was stirred overnight at 90 °C, then the mixture was cooled to room temperature and concentrated under reduced pressure. The residue was carefully and slowly diluted with water (200 mL) and extracted with EtOAc (3 x 200 mL). The organic extracts were dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether(1:50-1:30) as the eluent to yield a mixture of the title compounds as a brown oil (20 g, crude).




Compound 63.4 and compound 63.5. *tert*-Butyl 4-(benzofuran-6-yl)piperidine-1-carboxylate and *tert*-butyl 4-(benzofuran-4-yl)piperidine-1-carboxylate. Into a 50-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a suspension of Zn (1.13 g, 17.3 mmol) in DMA (5 mL). A 7:5 v/v mixture of TMSCl/1,2-dibromoethane (0.5 mL) was added to the reaction flask drop-wise to maintain the temperature below 65 °C, then the mixture was stirred for an additional 10 min. To this mixture was added a solution of *tert*-butyl 4-iodopiperidine-1-carboxylate (5.40 g, 17.4 mmol) in DMA (40 mL) drop-wise with stirring and the resulting mixture was stirred at room temperature for 1 hour. The above mixture was filtered, and added to a mixture of 6-bromo-3a,7a-dihydrobenzofuran (compound **63.2**) and 4-bromo-3a,7a-dihydrobenzofuran (compound **63.3**) (2.83 g, 14.2 mmol), and CuI (274 mg, 1.44 mmol), Pd(dppf)Cl₂ (1.18 g, 1.6 mmol) in DMA (30 mL). The resulting mixture was stirred overnight at 85 °C, then cooled to room temperature. The solids were removed by filtration and the filtrate was diluted with ethyl acetate (200 mL) and washed with brine (3 x 80 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1/30~1/20) as the eluent to yield *tert*-butyl 4-(1-benzofuran-6-yl)piperidine-1-carboxylate (compound **63.4**) as a colorless oil (267 mg, 6%) and *tert*-butyl 4-(1-benzofuran-4-yl)piperidine-1-carboxylate (compound **63.5**) as an off-white solid (320 mg, 7%).




Compound 63.6. 4-(Benzofuran-6-yl)piperidine. Into a 50-mL round-bottom flask, was placed a solution of *tert*-butyl 4-(1-benzofuran-6-yl)piperidine-1-carboxylate (compound **63.4**, 200 mg, 0.66 mmol) in dichloromethane (3 mL) and trifluoroacetic acid (1 mL). The resulting solution was stirred at 15 °C for 1 hour, then the pH was carefully adjusted to ~9 with aqueous sodium hydroxide (2 M). The mixture was diluted with H₂O (20 mL) and the layers were separated. The aqueous phase was extracted with dichloromethane (4 x 20 mL)

and the combined organic layers were washed with brine (30 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield the title compound as a brown oil (150 mg, crude).




5 **Compound 63. 5-(4-(Benzofuran-6-yl)piperidine-1-carbonyl)-2-cyclobutyl-N,4-dimethylbenzamide.** Into a 25-mL round-bottom flask, was placed a solution of 4-(benzofuran-6-yl)piperidine (compound **63.6**) (110 mg, 0.55 mmol), 4-cyclobutyl-2-methyl-5-(methylcarbamoyl)benzoic acid (compound **62.3**, 135 mg, 0.55 mmol), EDC•HCl (210 mg, 1.10 mmol) and 4-dimethylaminopyridine (133.5 mg, 1.09 mmol) in *N,N*-dimethylformamide (3 mL). The resulting solution was stirred for 2 h at 15 °C then diluted EtOAc (60 mL). The resulting mixture was washed with aqueous saturated NH₄Cl (2 x 20 mL) and brine (20 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude product (110 mg) was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-001(SHIMADZU)): Column, SunFire Prep C18 OBD Column, 5 µm, 19*150 mm; mobile phase, water with 0.05% TFA and CH₃CN (51.0% CH₃CN up to 60.0% in 9 min, up to 100.0% in 5 min, down to 51.0% in 1 min); Detector, Waters 2489, 254 & 220 nm. The fractions containing pure compound were combined and lyophilized to yield the title compound as a white solid (46.0 mg, 20%). *m/z* (ES⁺) 431 (M+H)⁺.




20 **Compound 64.1. (Aminoxy)diphenylphosphine oxide.** Into a 500-mL three neck round-bottom flask, was placed a solution of hydroxylamine hydrochloride (30.0 g, 432 mmol) in H₂O/dioxane (90/45 mL). The solution was cooled to 0-5 °C, then sodium bicarbonate (36.5 g, 434 mmol) was added portion-wise over 10 min and the mixture was stirred at 0-5 °C for 30 min. A solution of diphenylphosphinoyl chloride (41.0 g, 173 mmol) in dioxane (45 mL) was added drop-wise at 0-5 °C over 30 min, then the resulting mixture was stirred for an additional 2 h at ambient temperature. The resulting solids were collected

by filtration and washed with water (200 mL), NaOH (0.25 M, 200 mL) and PE (200 mL). The product was dried to yield the title compound as a white solid (20 g, crude).




Compound 64.2. 1-Amino-4-bromopyridin-1-ium iodide. Into a 250-mL round-

bottom flask, was placed a solution of 4-bromopyridine hydrochloride (13.8 g, 71.0 mmol) in water (50 mL). The solution was cooled to 0-5 °C then sodium bicarbonate (12.0 g, 141mmol) was added portion-wise over 10 min and the mixture was stirred at 0-5°C for 30 min. The mixture was extracted with DCM (4 x 50 mL) and the combined organic layers were washed with brine (50 mL), dried (Na₂SO₄), and filtered. The filtrate was placed into a 500-mL round-bottom flask, then purged and maintained with an inert atmosphere of nitrogen. (Aminooxy)diphenylphosphine oxide (compound **64.1**, 20 g, ~70% purity, 60 mmol) was added and the mixture was stirred overnight at room temperature. The reaction was carefully quenched with aqueous HI (8 mL, 45%) and stirred for 30 min. The solids were collected by filtration, and washed with DCM (200 mL) and hexanes (200 mL) to yield the title compound as a brown solid (15 g, crude).




Compound 64.3. Ethyl 5-bromopyrazolo[1,5-a]pyridine-3-carboxylate. Into a


250-mL three neck round-bottom flask, was placed a solution of 1-amino-4-bromopyridin-1-ium iodide (compound **64.2**, 15 g, ~50% purity, 24.9 mmol) in *N,N*-dimethylformamide (80 mL). Potassium carbonate (10.6 g, 76.7 mmol) was added portion-wise followed by the addition of ethyl propiolate (11.7 mL, 115 mmol) drop-wise over 10 min. The resulting mixture was stirred overnight at room temperature, then diluted with EtOAc (300 mL) and water (100 mL). The solids were removed by filtration and the organic layer was washed with brine (3 x 50 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:10) as the eluent to yield the title compound as a brown solid (300 mg, 6%).



Compound 64.4. 5-Bromopyrazolo[1,5-*a*]pyridine. Into a 50-mL round-bottom flask, was placed ethyl 5-bromopyrazolo[1,5-*a*]pyridine-3-carboxylate (compound **64.3**, 100 mg, 0.37 mmol). Sulfuric acid (50%, 4 mL) was added carefully in portions at room temperature, then the resulting solution was stirred overnight at 80 °C. After cooling to room temperature, the pH of the solution was carefully adjusted to 8-9 with aqueous sodium hydroxide (5 M) and then extracted with ethyl acetate (2 x 20 mL). The combined organic extracts were washed with brine (20 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:10) as the eluent to yield the title compound as a brown solid (40 mg, 55%).




Compound 64. 2-Cyclobutyl-N,4-dimethyl-5-(4-(pyrazolo[1,5-*a*]pyridin-5-yl)piperidine-1-carbonyl)benzamide. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **62**, except 5-bromopyrazolo[1,5-*a*]pyridine (compound **64.4**) was used in place of 5-bromo-1*H*-indazole to yield the title compound as a white solid. *m/z* (ES⁺) 431 (M+H)⁺.




Compound 65. (4-Cyclobutyl-3-(2,4-dimethyl-1*H*-imidazol-5-yl)phenyl)(4-(imidazo[1,5-*a*]pyridin-7-yl)piperidin-1-yl)methanone. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1**, except 4-cyclobutyl-3-iodobenzoic acid (compound **9.3**) was used in place of 5-iodo-2,4-dimethylbenzoic acid (compound **1.3**) and 7-(piperidin-4-yl)imidazo[1,5-*a*]pyridine hydrochloride (compound **39.5**) was used in place of 4-(piperidin-


4-yl)benzonitrile hydrochloride (compound **1.2**) to yield the title compound. m/z (ES+) 454



Compound 66.1. 1-(4-Bromopyridin-2-yl)ethanamine. Into a 500-mL 3-necked round-bottom flask, purged and maintained with an inert atmosphere of nitrogen, was placed a solution of methylmagnesium bromide (3 M in THF) (63.4 mL, 190 mmol) in THF (100 mL). A solution of 4-bromopyridine-2-carbonitrile (11.6 g, 63.4 mmol; patent US 2009/0239876 A1, example 2) in THF (40 mL) was added drop-wise at room temperature over 40 min. Methanol (40 mL) was then added drop-wise followed by portion-wise addition of sodium borohydride (11.8 g, 312 mmol) in several batches. The resulting mixture was stirred at room temperature for 2 h, then diluted with ethyl acetate (200 mL). The pH of the mixture was adjusted to 9 with aqueous sodium hydroxide (1 M) and the solids were removed by filtration. The filtrate was extracted with ethyl acetate (2 x 100 mL) and the combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure to yield 10.3 g (crude) of the title compound as a yellow oil.




Compound 66.2. 1-Methyl-7-(piperidin-4-yl)imidazo[1,5-a]pyridine hydrochloride. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **39.5**, except 1-(4-bromopyridin-2-yl)ethanamine (compound **66.1**) was used in place of (4-bromopyridin-2-yl)methanamine (compound **39.1**).




Compound 67.1. N-((4-Bromopyridin-2-yl)methyl)acetamide. Into a 500-mL round-bottom flask, purged and maintained with an inert atmosphere of nitrogen, was placed a solution of (4-bromopyridin-2-yl)methanamine (compound **39.1**, 4.5 g, 24.1 mmol) in dichloromethane (160 mL). Triethylamine (6.72 mL, 48.2 mmol) and acetic anhydride (2.29

mL, 24.2 mmol) were carefully added and the resulting solution was stirred at room temperature for 12 h. The volatiles were removed under reduced pressure and the residue was slowly quenched with water (200 mL). The pH of the solution was slowly adjusted to 9-12 with aqueous sodium carbonate (3 M) and the aqueous phase was extracted with ethyl acetate (3 x 60 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with dichloromethane/methanol (100:1-10:1) as the eluent to yield 4.0 g (73%) of the title compound as yellow oil.




Compound 67.2. 3-Methyl-7-(piperidin-4-yl)imidazo[1,5-a]pyridine

hydrochloride. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **39.5**, except *N*-((4-bromopyridin-2-yl)methyl)acetamide (compound **67.1**) was used in place of *N*-((4-bromopyridin-2-yl)methyl)formamide (compound **39.2**).




Compound 68.1 and compound 68.2. *tert*-Butyl 4-(1-chloroimidazo[1,5-a]pyridin-7-yl)piperidine-1-carboxylate and *tert*-butyl 4-(3-chloroimidazo[1,5-a]pyridin-7-yl)piperidine-1-carboxylate. Into a 25-mL round-bottom flask, was placed a solution of *tert*-butyl 4-(imidazo[1,5-a]pyridin-7-yl)piperidine-1-carboxylate (compound **39.4**, 170 mg, 0.56 mmol) in CCl₄ (3 mL). *N*-Chlorosuccinimide (75 mg, 0.56 mmol) was added and the resulting mixture was stirred for 5 h at room temperature. The mixture was diluted with EtOAc (30 mL) and washed with brine (3 x 20 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by preparative TLC with ethyl acetate/petroleum ether (1/1) to yield 45 mg (24%) of *tert*-butyl 4-(1-chloroimidazo[1,5-a]pyridin-7-yl)piperidine-1-carboxylate (compound **68.1**) as a yellow solid and 64 mg (34%)

of *tert*-butyl 4-(3-chloroimidazo[1,5-*a*]pyridin-7-yl)piperidine-1-carboxylate (compound **68.2**) as a yellow solid.



Compound 68.3. 1-Chloro-7-(piperidin-4-yl)imidazo[1,5-*a*]pyridine

- 5 **hydrochloride.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1.2**, except *tert*-butyl 4-(1-chloroimidazo[1,5-*a*]pyridin-7-yl)piperidine-1-carboxylate (compound **68.1**) was used in place of *tert*-butyl 4-(4-cyanophenyl)piperidine-1-carboxylate (compound **1.1**) to yield the title compound as a yellow solid.



Compound 71.1. 3-Chloro-7-(piperidin-4-yl)imidazo[1,5-*a*]pyridine




- 10 **hydrochloride.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1.2**, except *tert*-butyl 4-(3-chloroimidazo[1,5-*a*]pyridin-7-yl)piperidine-1-carboxylate (compound **68.2**) was used in place of *tert*-butyl 4-(4-cyanophenyl)piperidine-1-carboxylate (compound **1.1**) to yield the title compound as a yellow solid.




The compounds in **TABLE 4** were prepared using standard chemical manipulations and procedures with readily available starting materials or building blocks described in this manuscript. The utilized procedures were similar to that used for the preparation of


20 compound **1**, using 5-(2,4-dimethyl-1*H*-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **1.8**) and the respective amines (compound **5.2**, compound **39.5**, compound **66.2**, compound **67.2**, compound **68.3**, or compound **71.1**).

25


TABLE 4

Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
69	4-(1-(5-(2,4-dimethyl-1 <i>H</i> -imidazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile		385
70	(5-(2,4-dimethyl-1 <i>H</i> -imidazol-5-yl)-2,4-dimethylphenyl)(4-(imidazo[1,5- <i>a</i>]pyridin-7-yl)piperidin-1-yl)methanone		428
66	(5-(2,4-dimethyl-1 <i>H</i> -imidazol-5-yl)-2,4-dimethylphenyl)(4-(1-methylimidazo[1,5- <i>a</i>]pyridin-7-yl)piperidin-1-yl)methanone		442


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
67	(5-(2,4-dimethyl-1 <i>H</i> -imidazol-5-yl)-2,4-dimethylphenyl)(4-(3-methylimidazo[1,5- <i>a</i>]pyridin-7-yl)piperidin-1-yl)methanone		442
68	(4-(1-chloroimidazo[1,5- <i>a</i>]pyridin-7-yl)piperidin-1-yl)(5-(2,4-dimethyl-1 <i>H</i> -imidazol-5-yl)-2,4-dimethylphenyl)methanone		462
71	(4-(3-chloroimidazo[1,5- <i>a</i>]pyridin-7-yl)piperidin-1-yl)(5-(2,4-dimethyl-1 <i>H</i> -imidazol-5-yl)-2,4-dimethylphenyl)methanone		462



Compound 72. (4-Cyclopropyl-3-(2,4-dimethyl-1H-imidazol-5-yl)phenyl)(4-(1-methylimidazo[1,5-a]pyridin-7-yl)piperidin-1-yl)methanone. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1**, except methyl 4-cyclopropyl-3-propionylbenzoate (compound **10.7**) was used in place of methyl 2,4-dimethyl-5-propionylbenzoate (compound **1.5**) and 1-methyl-7-(piperidin-4-yl)imidazo[1,5-a]pyridine hydrochloride (compound **66.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**). m/z (ES+) 454 (M+H)⁺.




Compound 73. (4-Cyclopropyl-3-(2,4-dimethyl-1H-imidazol-5-yl)phenyl)(4-(3-methylimidazo[1,5-a]pyridin-7-yl)piperidin-1-yl)methanone. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1**, except methyl 4-cyclopropyl-3-propionylbenzoate (compound **10.7**) was used in place of methyl 2,4-dimethyl-5-propionylbenzoate (compound **1.5**) and 3-methyl-7-(piperidin-4-yl)imidazo[1,5-a]pyridine hydrochloride (compound **67.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**). m/z (ES+) 454 (M+H)⁺.




Compound 74.1. 1-(4-Bromophenyl)-1H-pyrazole. Into a 100-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a mixture of 1-bromo-4-iodobenzene (2.82 g, 9.97 mmol), 1H-pyrazole (680 mg, 9.99 mmol), CuI (380 mg, 2.00 mmol), DMEDA (430 μ L, 4.00 mmol, 0.40 equiv), Cs₂CO₃ (6.52 g, 20.00 mmol) and CH₃CN (40 mL). The resulting mixture was stirred overnight at 82 °C. After cooling to ambient temperature, the solids were removed by filtration and the filtrate was


concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:12) as the eluent to yield the title compound as a white solid (2.1 g, 94%).




Compound 74. 5-(4-(4-(1H-pyrazol-1-yl)phenyl)piperidine-1-carbonyl)-2-cyclobutyl-N,4-dimethylbenzamide. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **62**, except 1-(4-bromophenyl)-1H-pyrazole (compound **74.1**) was used in place of 5-bromo-1H-indazole to yield the title compound as a white solid. m/z (ES⁺) 457 (M+H)⁺.




Compound 75.1. 4-(4-Bromophenyl)-4H-1,2,4-triazole. Into a 100-mL round-bottom flask, was placed a solution of 4-bromoaniline (1.71 g, 9.94 mmol), *N'*-formylformohydrazide (2.64 g, 30.0 mmol), and triethylamine (9.74 mL, 69.9 mmol, 7.00 equiv) in pyridine (40 mL). Chlorotrimethylsilane (19.2 mL, 151 mmol) was added drop-wise and the resulting solution was stirred for 18 h at 100 °C, then cooled to room temperature. The resulting mixture was concentrated under reduced pressure and the residue was diluted with brine (50 mL) and extracted with ethyl acetate (6 x 50 mL). The combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was washed with ether (30 mL) and the solids were collected by filtration to yield the title compound as a pink solid (1.6 g, 72%).




Compound 75. 5-(4-(4-(4*H*-1,2,4-Triazol-4-yl)phenyl)piperidine-1-carbonyl)-2-cyclobutyl-*N*,4-dimethylbenzamide. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 62, except 4-(4-bromophenyl)-4*H*-1,2,4-triazole (compound 75.1) was used in place of 5-bromo-1*H*-indazole to yield the title compound as a yellow solid. *m/z* (ES⁺) 458 (M+H)⁺.




Compound 76.1. ((4-Bromophenyl)ethynyl)trimethylsilane. Into a 100-mL three neck round-bottom flask, which was maintained with an inert atmosphere of nitrogen, was placed a solution of 1-bromo-4-iodobenzene (1.00 g, 3.53 mmol) in tetrahydrofuran/TEA(9:1) (30 mL). PdCl₂(PPh₃)₂ (50 mg, 0.07 mmol), CuI (13.4 mg, 0.07 mmol), and ethynyltrimethylsilane (748 μL, 5.29 mmol) were added and the mixture was stirred for 18 h at room temperature, then concentrated under reduced pressure. The residue was diluted with water (50 mL) and extracted with ethyl acetate (3 x 20 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with petroleum ether as the eluent to yield the title compound as a light yellow solid (0.83 g, 93%).




Compound 76.2. *tert*-Butyl 4-(4-((trimethylsilyl)ethynyl)phenyl)piperidine-1-carboxylate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 62.4, except ((4-bromophenyl)ethynyl)trimethylsilane (compound 76.1, 850 mg, 3.36 mmol) was used in place of 5-bromo-1*H*-indazole to yield the title compound as a yellow oil (0.80 g, 67%).




Compound 76.3. *tert*-Butyl 4-(4-ethynylphenyl)piperidine-1-carboxylate. Into a 50-mL round-bottom flask, was placed a solution of *tert*-butyl 4-(4-((trimethylsilyl)ethynyl)phenyl)piperidine-1-carboxylate (compound **76.2**, 1.34 g, 3.75 mmol) in tetrahydrofuran (20 mL). Tetrabutylammonium fluoride (1.95 g, 7.47 mmol) was added and the resulting solution was stirred for 10 min at room temperature. The mixture was diluted with water (30 mL) and extracted with EtOAc (3 x 20 mL). The organic layers were combined, dried (Na₂SO₄), filtered and concentrated under reduced pressure to yield the title compound as a brown oil (1.0 g, crude).



Compound 76.4. 4-(4-Ethynylphenyl)piperidine hydrochloride. Into a 100-mL round-bottom flask, was placed a solution of *tert*-butyl 4-(4-ethynylphenyl)piperidine-1-carboxylate (compound **76.3**, 1.0 g, 3.5 mmol) in ethyl acetate (20 mL). Hydrogen chloride (g) was introduced by bubbling through the solution and the resulting solution was stirred for 1 hour at room temperature. The solids that formed were collected by filtration and washed with hexanes (3 x 10 mL) to yield the title compound as a brown solid (630 mg, 81%).




Compound 76. 2-Cyclobutyl-5-(4-(4-ethynylphenyl)piperidine-1-carbonyl)-*N*,4-dimethylbenzamide. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **62**, except 4-(4-ethynylphenyl)piperidine hydrochloride (compound **76.4**, 178 mg, 0.81 mmol) was used in place of 5-(piperidin-4-yl)-1*H*-indazole (compound **62.5**) to yield the title compound as a white solid (20.1 mg, 12%). *m/z* (ES⁺) 415 (M+H)⁺.




Compound 77.1. 1-Bromo-4-(prop-1-yn-1-yl)benzene. Into a 250-mL three neck round-bottom flask, which was maintained with an inert atmosphere of nitrogen, was placed

1-bromo-4-iodobenzene (2.00 g, 7.07 mmol), $\text{PdCl}_2(\text{PPh}_3)_2$ (99.2 mg, 0.14 mmol), CuI (26.8 mg, 0.14 mmol), trimethyl(prop-1-yn-1-yl)silane (2.08 mL, 14.1 mmol) and tetrahydrofuran/TEA(9:1) (100 mL). Stirring was initiated and tetrabutylammonium fluoride (3.69 g, 14.1 mmol) was added rapidly to the mixture. The resulting mixture was stirred for 18 h at room temperature, then concentrated under reduced pressure. The residue diluted with water (100 mL) and extracted with ethyl acetate (3 x 50 mL). The organic extracts were dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with petroleum ether as the eluent to yield the title compound as a light yellow oil (1.0 g, 73%).




Compound 77. 2-Cyclobutyl-N,4-dimethyl-5-(4-(4-(prop-1-yn-1-yl)phenyl)piperidine-1-carbonyl)benzamide. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **62**, except 1-bromo-4-(prop-1-yn-1-yl)benzene (compound **77.1**) was used in place of 5-bromo-1*H*-indazole to yield the title compound as a white solid. m/z (ES+) 429 (M+H)+.




Compound 78.1. Methyl 5-carbamothioyl-4-cyclobutyl-2-methylbenzoate. To a round-bottom flask was added methyl 5-cyano-4-cyclobutyl-2-methylbenzoate (compound **6.4**, 3.63 g, 0.015 mol), *O,O'*-diethyl dithiophosphate (10 mL) and water (1 mL). The reaction mixture was heated to 80 °C for 3 hours (**CAUTION**: significant gas evolution occurs - this and all other reactions described herein should be carried out in well ventilated fume hoods). After cooling to room temperature, the reaction mixture was partitioned between ethyl acetate (50 mL) and water (50 mL). The organic layer was washed successively with saturated aqueous NaHCO_3 (50 mL) and brine (50 mL), dried (Na_2SO_4), filtered, and concentrated

under reduced pressure. The crude product was purified by silica gel flash chromatography (hexanes/ethyl acetate = 80/20 to 50/50) to yield the title compound as a yellow solid (3.06 g, 78% yield). m/z (ES+) 264 (M+H)⁺. ¹H NMR (400 MHz, CDCl₃): δ 7.93 (s, 1H), 7.82 (s, 1H), 7.26 (s, 1H), 6.92 (s, 1H), 4.19 (m, 1H), 3.89 (s, 3H), 2.64 (s, 3H), 2.40 (m, 2H), 2.29 – 2.15 (m, 2H), 2.12 – 2.00 (m, 1H), 1.95 – 1.84 (m, 1H).



Compound 78.2. Methyl 4-cyclobutyl-5-(imino(methylthio)methyl)-2-


methylbenzoate. To a round-bottom flask was added methyl 5-carbamothioyl-4-cyclobutyl-2-methylbenzoate (compound **78.1**, 861 mg, 3.27 mmol) in THF (10 mL). Iodomethane (400 μ L, 6.42 mmol) was added drop-wise and the reaction mixture was stirred at room temperature for 7 hours. The reaction mixture was concentrated under reduced pressure and the residue was purified by silica gel flash chromatography (ethyl acetate to ethyl acetate/methanol = 95/5) to yield the title compound as a yellowish oil (807 mg, 89% yield). m/z (ES+) 278 (M+H)⁺. ¹H NMR (400 MHz, DMSO-*d*₆): δ 7.67 (s, 1H), 7.40 (s, 1H), 3.88 – 3.71 (m, 4H), 2.57 (s, 3H), 2.44 (s, 3H), 2.22–2.19 (m, 2H), 2.12 (m, 2H), 1.98 – 1.86 (m, 1H), 1.82 – 1.70 (m, 1H).



Compound 78.3. Methyl 4-cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-


yl)benzoate. To a round-bottom flask was added methyl 4-cyclobutyl-5-(imino(methylthio)methyl)-2-methylbenzoate (compound **78.2**, 556 mg, 2.00 mmol) and aceto-hydrazide (223 mg, 3.00 mol) in acetic acid (6 mL). The mixture was heated at 90 °C for 3 hours then cooled to room temperature. The reaction mixture was partitioned between water (50 mL) and ethyl acetate (50 mL) and the organic layer was washed with brine (2 x 50 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel flash chromatography (hexanes/ethyl acetate = 50/50 to 30/70) to yield the compound as a white solid (243 mg, 43% yield). m/z (ES+) 286 (M+H)⁺. ¹H NMR (400 MHz, CDCl₃): δ 8.23 (s, 1H), 7.32 (s, 1H), 4.24 – 4.05 (m, 1H), 3.89 (s, 3H), 2.69 (s, 3H),

2.54 (s, 3H), 2.23-2.20 (m, 2H), 2.16 – 2.05 (m, 2H), 2.05 – 1.88 (m, 1H), 1.88 – 1.71 (m, 1H).




Compound 78.4. 4-Cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-

yl)benzoic acid. To a solution of methyl 4-cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoate (compound **78.3**, 240 mg, 0.842 mmol) in methanol (5 mL) was added aqueous NaOH (6 mL, 1 M). The resulting mixture was heated to 50 °C for 6 hours then cooled to ambient temperature and acidified to pH 2 with aqueous 1 M HCl and extracted with ethyl acetate (3 x 50 mL). The combined organic extracts were washed with brine (50 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield the title compound as a white solid (260 mg, quantitative). *m/z* (ES⁺) 272 (M+H)⁺.



Compound 78.5. 6-(Piperidin-4-yl)imidazo[1,2-*a*]pyridine hydrochloride. The


title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1.2** except 6-bromoimidazo[1,2-*a*]pyridine (500 mg) was used in place of 4-bromobenzonitrile to yield the title compound as a brown solid (400 mg, 66% over 2 steps).



Compound 78. (4-Cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-


yl)phenyl)(4-(imidazo[1,2-*a*]pyridin-6-yl)piperidin-1-yl)methanone. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **62**, except 4-cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoic acid (compound **78.4**, 100 mg) was used in place of 4-cyclobutyl-2-methyl-5-(methylcarbamoyl)benzoic acid (compound **62.3**) and 6-(piperidin-4-yl)imidazo[1,2-*a*]pyridine hydrochloride (compound **78.5**, 88 mg) was used in place of 5-(piperidin-4-yl)-

1*H*-indazole (compound **62.5**). The title compound was obtained as a white solid (35.6 mg, 21%). m/z (ES+) 455 (M+H)⁺.




Compound 79. 4-(1-(4-Cyclobutyl-2-methyl-5-(5-methyl-4*H*-1,2,4-triazol-3-


yl)benzoyl)-3-fluoroazetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **62**, except 4-cyclobutyl-2-methyl-5-(5-methyl-4*H*-1,2,4-triazol-3-yl)benzoic acid (compound **78.4**) was used in place of 4-cyclobutyl-2-methyl-5-(methylcarbamoyl)benzoic acid (compound **62.3**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 5-(piperidin-4-yl)-1*H*-indazole (compound **62.5**). m/z (ES+) 430 (M+H)⁺.




Compound 80.1. 3-Methoxypropanehydrazide. Into a 500-mL round-bottom flask, was placed a solution of methyl 3-methoxypropanoate (30.0 g, 254 mmol) in ethanol (100 mL) and hydrazine hydrate (24.7 mL, 507 mmol). The resulting solution was stirred overnight at 80 °C, then cooled and concentrated under reduced pressure to yield 26.3 g (88%, crude) of the title compound as colorless oil.




Compound 80. 4-(1-(4-Cyclobutyl-5-(5-(2-methoxyethyl)-4*H*-1,2,4-triazol-3-yl)-2-methylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **78**, except 3-methoxypropanehydrazide (compound **80.1**) was used in place of acetohydrazide and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 6-(piperidin-4-yl)imidazo[1,2-*a*]pyridine hydrochloride (compound **78.5**). m/z (ES+) 474 (M+H)⁺.




Compound 81. 4-(1-(4-Cyclobutyl-5-(5-(2-methoxyethyl)-4H-1,2,4-triazol-3-yl)-2-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 78, except 3-methoxypropanehydrazide (compound **80.1**) was used in place of acetohydrazide and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 6-(piperidin-4-yl)imidazo[1,2-*a*]pyridine hydrochloride (compound **78.5**). *m/z* (ES⁺) 456 (M+H)⁺.




Compound 82.1. 1-*tert*-Butyl 4-ethyl 5-oxoazepane-1,4-dicarboxylate. To a solution of *tert*-butyl 4-oxopiperidine-1-carboxylate (20.0 g, 100 mmol) in diethylether (60 mL) under nitrogen at -30 °C, was added drop-wise a solution of BF₃·Et₂O (16.0 mL, 130 mmol) in ether (20 mL). After stirring for 30 min at -30 °C, a solution of ethyl 2-diazoacetate (16.0 g, 140 mmol) in ether (20 mL) was added drop-wise to the reaction at -30 °C. The resulting solution was stirred for 1 h at -30 °C, then at room temperature for 2 h. The reaction was carefully quenched with 30% aqueous potassium carbonate (100 mL) and the resulting mixture was extracted with ethyl acetate (2 x 250 mL). The combined organic extracts were washed with brine (2 x 50 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1/10) as the eluent to furnish 19 g (66%) of the title compound as a light yellow oil.




Compound 82.2. *tert*-Butyl 4-oxoazepane-1-carboxylate. To a solution 1-*tert*-butyl 4-ethyl 5-oxoazepane-1,4-dicarboxylate (compound **82.1**, 19.0 g, 66.6 mmol) in 1,4-dioxane (190 mL) was added drop-wise a solution of sodium hydroxide (4.00 g, 100 mmol) in water (100 mL). The resulting mixture was stirred at room temperature overnight. The pH was then adjusted to 4-5 with hydrogen chloride (aq. 3 M) and the resulting solution was extracted with ethyl acetate (2 x 50 mL). The combined organic extracts were washed with brine (2 x 10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure. The crude residue was purified by silica gel column chromatography using ethyl acetate/petroleum ether (1:3) as the eluent to furnish 11 g (77%) of the title compound as a yellow oil.




Compound 82.3. *tert*-Butyl 4-bromo-5-oxoazepane-1-carboxylate. To a solution *tert*-butyl 4-oxoazepane-1-carboxylate (compound **82.2**, 11.0 g, 51.6 mmol) in chloroform (220 mL) at 0 °C was added drop-wise a solution of bromine (3.98 mL, 77.6 mmol) in chloroform (110 mL). The resulting mixture was stirred at room temperature overnight, then the solids that formed were collected by filtration and dissolved in dichloromethane (200 mL). Triethylamine (16.8 mL, 121 mmol) and (Boc)₂O (8.70 g, 40.3 mmol) were added to the mixture at 0 °C and the resulting solution was stirred for 3 h at room temperature, and then concentrated under reduced pressure. The crude residue was purified by silica gel chromatography using ethyl acetate/petroleum ether (1:10) as the eluent to give 4.0 g (27%) of the title compound as a yellow oil.




Compound 82.4. 4-Cyclobutyl-5-formyl-2-methylbenzoic acid. Into a three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 4-cyclobutyl-5-iodo-2-methylbenzoic acid (compound **62.1**, 5.00 g, 80%, 12.7 mmol) in a solvent mixture of tetrahydrofuran and Et₂O (50mL/50 mL). The solution was cooled to -78 °C then n-butyllithium (15 mL, 2.5 M in hexanes) was added drop-wise with stirring. *N,N*-Dimethylformamide (2.64 mL, 34.2 mmol) was added and the resulting mixture was stirred for 1 h at -78 °C, then carefully quenched by slow addition of aqueous NH₄Cl (sat., 50 mL). The pH was adjusted to 1-2 with aqueous hydrogen chloride (6 M), then diluted with ethyl acetate (100 mL) and washed with brine (4 x 50 mL). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified using silica gel column chromatography with ethyl acetate/petroleum ether (1:1) as the eluent to furnish 1.62 g (41%) of the title compound as a white solid.




Compound 82.5. Methyl 4-cyclobutyl-5-formyl-2-methylbenzoate. Into a 100-mL round-bottom flask, was placed a mixture of 4-cyclobutyl-5-formyl-2-methylbenzoic acid (compound **82.4**, 500 mg, 2.29 mmol) in *N,N*-dimethylformamide (10 mL) and sodium bicarbonate (390 mg, 4.64 mmol). With stirring, methyl iodide (430 µL, 6.90 mmol) was added drop-wise and the resulting mixture was stirred for 5 h at room temperature. The reaction was then diluted with EtOAc (50 mL) and the mixture was washed with brine (4 x 10 mL). The organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield 0.40 g (crude) of the title compound as a brown oil



Compound 82.6. *tert*-Butyl 2-(2-cyclobutyl-5-(methoxycarbonyl)-4-methylphenyl)-4,5,7,8-tetrahydroimidazo[4,5-*d*]azepine-6(1*H*)-carboxylate. Into a 100-mL round-bottom flask, was placed a mixture of methyl 4-cyclobutyl-5-formyl-2-methylbenzoate (compound **82.5**, 300 mg, 1.29 mmol), ammonium acetate (449 mg, 5.83 mmol), *tert*-butyl 4-bromo-5-oxoazepane-1-carboxylate (compound **82.3**, 564 mg, 1.93 mmol) and ammonium hydroxide (25%) (597 μ L, 3.87 mmol) in *N,N*-dimethylformamide (8 mL). The resulting mixture was stirred for 4 h at 130 °C, then cooled and quenched with water/ice (10 mL). The aqueous phase was extracted with ethyl acetate (2 x 30 mL) and the combined organic layers were washed with brine (3 x 10 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1/1) as the eluent to yield 0.10 g (18%) of the title compound as a white solid.




Compound 82.7. Methyl 4-cyclobutyl-5-(1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2-methylbenzoate hydrochloride. Into a 50-mL 3-necked round-bottom flask, was placed a solution of *tert*-butyl 2-(2-cyclobutyl-5-(methoxycarbonyl)-4-methylphenyl)-4,5,7,8-tetrahydroimidazo[4,5-*d*]azepine-6(1*H*)-carboxylate (compound **82.6**, 200 mg, 0.46 mmol) in EtOAc (10 mL). Hydrogen chloride (gas) was introduced into the solution by bubbling and the solution was stirred for 30 min at room temperature. The resulting mixture was concentrated under reduced pressure to yield 136 mg (crude) of the title compound as a yellow solid.




Compound 82.8. Methyl 4-cyclobutyl-2-methyl-5-(6-methyl-1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)benzoate. Into a 100-mL round-bottom flask, was placed a mixture of methyl 4-cyclobutyl-5-(1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-

2-methylbenzoate hydrochloride (compound **82.7**, 40 mg, 0.11 mmol), NaBH(OAc)₃ (75 mg, 0.35 mmol), and formaldehyde (37 wt %) (26 μ L, 0.33 mmol) in tetrahydrofuran (4 mL). The resulting mixture was stirred for 2 h at 40 °C, then cooled and the pH of the solution was adjusted to 8-9 with sodium bicarbonate (sat.). The aqueous phase was extracted with ethyl acetate (2 x 20 mL) and the combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield 20 mg (crude) of the title compound as a yellow solid.



Compound 82.9. 4-Cyclobutyl-2-methyl-5-(6-methyl-1,4,5,6,7,8-


hexahydroimidazo[4,5-d]azepin-2-yl)benzoic acid. Into a 50-mL round-bottom flask, was placed a mixture of methyl 4-cyclobutyl-2-methyl-5-(6-methyl-1,4,5,6,7,8-hexahydroimidazo[4,5-d]azepin-2-yl)benzoate (compound **82.8**, 40 mg, 0.11 mmol) and NaOH (18 mg, 0.44 mmol) in methanol (4 mL), water (2 mL). The resulting solution was stirred for 2 h at 60 °C. After cooling to room temperature, the volatiles were removed under reduced pressure. The pH of the residual solution was adjusted to about 1 with hydrogen chloride (3 M) and concentrated under reduced pressure to yield 0.10 g (crude) of the title product as the HCl salt as a yellow solid.




Compound 82. 4-(1-(4-Cyclobutyl-2-methyl-5-(6-methyl-1,4,5,6,7,8-

hexahydroimidazo[4,5-d]azepin-2-yl)benzoyl)azetidin-3-yl)benzonitrile. Into a 100-mL round-bottom flask, was placed a mixture of 4-cyclobutyl-2-methyl-5-(6-methyl-1,4,5,6,7,8-hexahydroimidazo[4,5-d]azepin-2-yl)benzoic acid (compound **82.9**, 40 mg, 0.12 mmol), EDC•HCl (45.4 mg, 0.24 mmol, 2.00 equiv), 4-dimethylaminopyridine (29 mg, 0.24 mmol) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**, 23 mg, 0.12 mmol) in *N,N*-dimethylformamide (5 mL). The resulting solution was stirred for 4 h at room temperature

then quenched by the addition of water/ice (10 mL). The resulting mixture was extracted with ethyl acetate (2 x 20 mL) and the combined organic layers were washed with brine (2 x 10 mL). The organic layer was dried (Na_2SO_4), filtered, and concentrated under reduced pressure to yield the crude product which was purified by Prep-HPLC using the following conditions (1#-Pre-HPLC-001(SHIMADZU)): Column, XBridge Shield RP18 OBD Column, 5 μm , 19*150 mm; mobile phase, WATER WITH 0.03% NH_4OH and CH_3CN (30% CH_3CN up to 43% in 8 min, up to 100% in 4 min, down to 30% in 2 min); Detector, Waters 2489, 254 & 220 nm. The fractions containing clean product were combined to yield 2.2 mg (4%) of the title compound as a white solid. m/z (ES^+) 480 ($\text{M}+\text{H}$)⁺.




Compound 83.1. 5-Formyl-2,4-dimethylbenzoic acid. To a stirred solution of 5-iodo-2,4-dimethylbenzoic acid (compound **1.3**, 5.00 g, 18.1 mmol) in tetrahydrofuran (150 mL) under nitrogen at -78 °C was added n-BuLi (2.5 M in THF, 18 mL, 45 mmol) drop-wise. The mixture was stirred at -78 °C for 1 h and then DMF (5.3 mL, 68 mmol) was added drop-wise. The resulting mixture was stirred at -78 °C for 0.5 h and then carefully quenched by slow addition of water (50 mL). The pH of the mixture was adjusted to ~3-4 with aqueous HCl (6 M) and then extracted with ethyl acetate (3 x 200 mL). The combined organic layers were dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was purified using silica gel column chromatography with ethyl acetate/petroleum ether (1:10-1:5) as the eluent to yield the title compound as a white solid (2.4 g, 74%).




Compound 83.2. Methyl 5-formyl-2,4-dimethylbenzoate. Into a 250-mL round-bottom flask, was placed 5-formyl-2,4-dimethylbenzoic acid (compound **83.1**, 2.00 g, 11.2 mmol) and methanol (50 mL). Concentrated sulfuric acid (2 mL) was carefully added drop-wise and the resulting solution was stirred for 2 h at 80 °C, then cooled and the volatiles were removed under reduced pressure. The pH of the residue was adjusted to 9 with sodium bicarbonate (sat.), then the aqueous phase was extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with brine (2 x 20 mL), dried (Na_2SO_4), filtered, and

concentrated under reduced pressure to yield 2.0 g (crude) of the title compound as yellow oil. The crude product was used in next step without further purification.




Compound 83.3. *tert*-Butyl 2-(5-(methoxycarbonyl)-2,4-dimethylphenyl)-4,5,7,8-tetrahydroimidazo[4,5-*d*]azepine-6(1*H*)-carboxylate. Into a 10-mL sealed tube, was placed methyl 5-formyl-2,4-dimethylbenzoate (compound **83.2**, 500 mg, 2.60 mmol), *tert*-butyl 4-bromo-5-oxoazepane-1-carboxylate (compound **82.3**, 1.1 g, 3.8 mmol), ammonium hydroxide (25%) (1.2 mL, 7.8 mmol), ammonium acetate (900 mg, 11.7 mmol) in *N,N*-dimethylformamide (6 mL). The resulting mixture was stirred for 3 h at 130 °C behind a blast shield, then cooled to room temperature and diluted with ethyl acetate (150 mL). The mixture was washed with brine (5 x 20 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with dichloromethane/methanol (10:1) as the eluent to yield 0.80 g (46%) of the title compound as yellow oil.




Compound 83.4. Methyl 5-(1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2,4-dimethylbenzoate. Into a 100-mL round-bottom flask, was placed a solution of *tert*-butyl 2-(5-(methoxycarbonyl)-2,4-dimethylphenyl)-4,5,7,8-tetrahydroimidazo[4,5-*d*]azepine-6(1*H*)-carboxylate (compound **83.3**, 800 mg, 2.00 mmol) in dichloromethane (16 mL).


Trifluoroacetic acid (4 mL) was added drop-wise and the resulting solution was stirred for 5 h at room temperature. The pH of the solution was carefully adjusted to 8-9 with NaHCO₃ (sat.) and the aqueous phase was extracted with DCM (3 x 30 mL), and the combined organics were dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield 0.60 mg (crude) of the title compound as brown oil.



Compound 83.5. Methyl 5-(6-isopropyl-1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2,4-dimethylbenzoate. Into a 100-mL round-bottom flask, was placed methyl 5-(1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2,4-dimethylbenzoate (compound **83.4**, 347 mg, 0.84 mmol), 2-bromopropane (790 μ L, 8.40 mmol), *N,N*-diisopropylethylamine (1.46 mL, 8.4 mmol) in *N,N*-dimethylformamide (5 mL). The solution was stirred for 4 h at 80 $^{\circ}$ C, then cooled to room temperature. The resulting solution was diluted with ethyl acetate (50 mL) and washed with brine (4 x 20 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure to yield 0.22 g (crude) of the title compound as brown oil.




Compound 83. 4-(1-(5-(6-Isopropyl-1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **82**, except methyl 5-(6-isopropyl-1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2,4-dimethylbenzoate (compound **83.5**) was used in place of 4-cyclobutyl-2-methyl-5-(6-methyl-1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)benzoate (compound **82.8**). *m/z* (ES⁺) 468 (M+H)⁺.




Compound 84. 4-(1-(2,4-Dimethyl-5-(6-methyl-1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **82**, except methyl 5-(1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2,4-


dimethylbenzoate (compound **83.4**) was used in place of methyl 4-cyclobutyl-5-(1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2-methylbenzoate hydrochloride (compound **82.7**). m/z (ES+) 440 (M+H)⁺.



- 5 **Compound 85.1. 3-(4-Chlorophenyl)azetidine hydrochloride.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.2**, except (4-chlorophenyl)boronic acid was used in place of (4-cyanophenyl)boronic acid. The title compound was obtained in 20% yield over two steps.




- 10 **Compound 85. (3-(4-Chlorophenyl)azetidin-1-yl)(2,4-dimethyl-5-(6-methyl-1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)phenyl)methanone.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **82**, except methyl 5-(1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2,4-dimethylbenzoate (compound **83.4**) was used in place of methyl 4-cyclobutyl-5-(1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2-methylbenzoate hydrochloride (compound **82.7**) and 3-(4-chlorophenyl)azetidine hydrochloride (compound **85.1**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 449 (M+H)⁺.




- 20 **Compound 86.1. Methyl 4-cyclopropyl-2-methylbenzoate.** To a solution of methyl 4-bromo-2-methylbenzoate (compound **6.1**, 5.00 g, 20.7 mmol, 95%) in a mixture of toluene and H₂O (20 mL/1 mL) were added potassium carbonate (6.10 g, 44.1 mmol), cyclopropylboronic acid (2.30 g, 26.8 mmol), Pd(dppf)Cl₂ (900 mg, 1.23 mmol), and Pd(OAc)₂ (250 mg, 1.12 mmol). The reaction mixture was purged with nitrogen and stirred at


80 °C overnight. After cooling to room temperature, the mixture was then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:50) as the eluent to yield 2.68 g (61%) of the title compound as a colorless oil.




Compound 86.2. Methyl 4-cyclopropyl-5-iodo-2-methylbenzoate. To a solution of methyl 4-cyclopropyl-2-methylbenzoate (compound **86.1**, 2.68 g, 13.4 mmol, 95%) in AcOH (50 mL) were added NaIO₄ (1.51 g, 7.08 mmol), iodine (3.58 g, 14.1 mmol), and sulfuric acid (106 µL, 2.0 mmol, 0.15 equiv). The reaction mixture was stirred overnight at 110 °C. After cooling to ambient temperature, water (100 mL) was slowly added and the mixture was extracted with ethyl acetate (100 mL). The organic layer was washed with Na₂S₂O₃ (aq., sat., 3 x 30 mL) and brine (30 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1/50) as the eluent to yield 2.0 g (45%) of the title compound as a colorless oil.




Compound 86.3. 4-Cyclopropyl-5-iodo-2-methylbenzoic acid. Into a 500-mL round-bottom flask, was placed a solution of methyl 4-cyclopropyl-5-iodo-2-methylbenzoate (compound **86.2**, 15.0 g, 47.5 mmol) in methanol (150 mL). A solution of sodium hydroxide (5.70 g, 143 mmol) in water (75 mL) was added and the resulting solution was stirred for 4 h at 60 °C, then cooled to room temperature. The volatiles were removed under reduced pressure and the remaining solution was adjusted to pH 3 with aqueous hydrogen chloride (12 M). The mixture was extracted with ethyl acetate (2 x 200 mL) and the combined organic extracts were washed with NH₄Cl (aq.) (2 x 400 mL) and brine (400 mL). The organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield 13.0 g (91%) of the title compound as a light yellow solid.




Compound 86.4. Methyl 4-cyclopropyl-5-(1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2-methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **83.4**, except 4-cyclopropyl-5-iodo-2-methylbenzoic acid (compound **86.3**) was used in place of 5-iodo-2,4-dimethylbenzoic acid (compound **1.3**).




Compound 86. 4-(1-(4-Cyclopropyl-2-methyl-5-(6-methyl-1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)benzoyl)azetid-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **82**, except methyl 4-cyclopropyl-5-(1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2-methylbenzoate (compound **86.4**) was used in place of methyl 4-cyclobutyl-5-(1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)-2-methylbenzoate hydrochloride (compound **82.7**). m/z (ES⁺) 466 (M+H)⁺.




Compound 87.1. Methyl 4-cyclopropyl-5-formyl-2-methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **83.2**, except 4-cyclopropyl-5-iodo-2-methylbenzoic acid (compound **86.3**) was used in place of 5-iodo-2,4-dimethylbenzoic acid (compound **1.3**).



Compound 87.2. Methyl 4-cyclopropyl-5-(3*H*-imidazo[4,5-*c*]pyridin-2-yl)-2-methylbenzoate. Into a 100-mL round-bottom flask was placed a mixture of methyl 4-cyclopropyl-5-formyl-2-methylbenzoate (compound **87.1**, 500 mg, 2.29 mmol), pyridine-3,4-diamine (500 mg, 4.58 mmol), NH₄OAc (1.42 g, 18.4 mmol) and ethanol (50 mL). The resulting mixture was stirred open to the air for 3 days at 70 °C, then cooled to room temperature and diluted with aqueous sodium bicarbonate (sat., 50 mL). The aqueous phase was extracted with ethyl acetate (2 x 50 mL) and the combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with ethyl acetate as the eluent to yield 268 mg (38%) of the title compound as a yellow semi-solid.




Compound 87.3. 2-(2-Cyclopropyl-5-(methoxycarbonyl)-4-methylphenyl)-5-methyl-3*H*-imidazo[4,5-*c*]pyridin-5-ium iodide. Into a 50-mL round-bottom flask, was placed a solution of methyl 4-cyclopropyl-5-(3*H*-imidazo[4,5-*c*]pyridin-2-yl)-2-methylbenzoate (compound **87.2**, 500 mg, 1.63 mmol) in dichloromethane (15 mL). Iodomethane (203 µL, 3.26 mmol) was added drop-wise and the resulting solution was stirred overnight at room temperature. The mixture was concentrated under reduced pressure to yield 0.30 g (41%) of the title compound as a yellow solid.




Compound 87.4. Methyl 4-cyclopropyl-2-methyl-5-(5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridin-2-yl)benzoate. Into a 100-mL round-bottom flask, was placed a mixture of 2-(2-cyclopropyl-5-(methoxycarbonyl)-4-methylphenyl)-5-methyl-3*H*-


imidazo[4,5-*c*]pyridin-5-ium iodide (compound **87.3**, 300 mg, 0.67 mmol) and NaBH₄ (1.42 g, 37.5 mmol) in methanol (30 mL). The resulting mixture was stirred for 4 h at room temperature, then concentrated under reduced pressure. The residue was diluted with EtOAc (120 mL) and the mixture was washed with brine (2 x 40 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with EtOAc/MeOH (20/1) as the eluent to yield 170 mg (78%) the title compound as a light yellow oil.



Compound 87. 4-(1-(4-Cyclopropyl-2-methyl-5-(5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridin-2-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **82**, except methyl 4-cyclopropyl-2-methyl-5-(5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridin-2-yl)benzoate (compound **87.4**) was used in place of 4-cyclobutyl-2-methyl-5-(6-methyl-1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)benzoate (compound **82.8**). *m/z* (ES⁺) 452 (M+H)⁺.




Compound 88.1. Methyl 4-cyclobutyl-2-methyl-5-(5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridin-2-yl)benzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **87.4**, except methyl 4-cyclobutyl-5-formyl-2-methylbenzoate (compound **82.5**) was used in place of methyl 4-cyclopropyl-5-formyl-2-methylbenzoate (compound **87.1**).




Compound 88. 4-(1-(4-cyclobutyl-2-methyl-5-(5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridin-2-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was


prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **82**, except methyl 4-cyclobutyl-2-methyl-5-(5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridin-2-yl)benzoate (compound **88.1**) was used in place of 4-cyclobutyl-2-methyl-5-(6-methyl-1,4,5,6,7,8-hexahydroimidazo[4,5-*d*]azepin-2-yl)benzoate (compound **82.8**). m/z (ES⁺) 466 (M+H)⁺.




Compound 89.1. Methyl 2-bromo-4-methylbenzoate. A solution of 2-bromo-4-methylbenzoic acid (10.0 g, 46.5 mmol) in MeOH (50 mL) was cooled to 0 °C, then concentrated sulfuric acid (10 mL) was carefully added. The mixture was heated at 70 °C for 2 hours. After cooling to room temperature, the volatile organics were removed under reduced pressure, and the residue was poured onto ice-water (100 mL). The mixture was extracted with EtOAc (x2) and the combined organic extracts were washed with aq. NaHCO₃, brine, dried (MgSO₄), filtered, and concentrated to yield 10.5 g (99%) of the title compound as a clear oil. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.73 (d, 1H), 7.50 (d, 1H), 7.19 – 7.11 (m, 1H), 3.92 (s, 3H), 2.36 (s, 3H).



Compound 89.2. Methyl 2-cyclobutyl-4-methylbenzoate. Cyclobutylzinc(II) bromide (50 mL, 0.5 M in THF, 25.0 mmol) was added to a mixture of methyl 2-bromo-4-methylbenzoate (compound **89.1**, 5.0 g, 21.8 mmol) and PdCl₂(dppf)CH₂Cl₂ (1.78 g, 2.20 mmol). The mixture was degassed with argon, then heated at 65°C under argon for 24 hours. The mixture was cooled to 0 °C, then carefully quenched with water (10 mL). The mixture was diluted with EtOAc (200 mL) and washed with water then brine. The organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexanes:EtOAc 30:1 to 20:1) to yield 3.6 g (81%) of the title compound. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.68 (d, 1H), 7.23 – 7.17 (s, 1H), 7.03 (d, 1H), 4.16 (m, 1H), 3.86 (s, 3H), 2.39 (s, 3H), 2.34 (m, 2H), 2.16 – 1.96 (m, 3H), 1.80 (m, 1H).




Compound 89.3. Methyl 2-cyclobutyl-5-iodo-4-methylbenzoate. To a solution methyl 2-cyclobutyl-4-methylbenzoate (compound **89.2**, 4.77 g, 23.3 mmol) in concentrated sulfuric acid (100 mL) at 0 °C, was added *N*-iodosuccinimide (5.25 g, 23.3 mmol) portion-
 5 wise. The mixture was stirred at 0 °C for 30 min and then at RT for 2 hours. The thick, dark mixture was cooled back to 0 °C, then MeOH (100 mL) was added slowly and carefully. The mixture was heated at 60 °C for 2 hours. After cooling to room temperature, the volatile solvents were removed under reduced pressure and the residue was carefully poured onto ice water (200 mL). The mixture was extracted with EtOAc (2x) and the combined organic
 10 extracts were washed with brine, aqueous NaHCO₃ (1 M), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexanes:EtOAc 30:1 to 20:1) to yield 5.0 g (65%) of the title compound as a clear oil. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.19 (s, 1H), 7.24 (s, 1H), 4.17 – 4.04 (m,
 1H), 3.86 (s, 3H), 2.48 – 2.44 (s, 3H), 2.40 – 2.28 (m, 2H), 2.13 – 1.92 (m, 3H), 1.85 – 1.75
 15 (m, 1H).




Compound 89.4. Methyl 5-cyano-2-cyclobutyl-4-methylbenzoate. A mixture of methyl 2-cyclobutyl-5-iodo-4-methylbenzoate (compound **89.3**, 3.0 g, 9.1 mmol), Zn(CN)₂ (2.3 g, 19.6 mmol) and Pd(PPh₃)₄ (0.55 g, 0.47 mmol) in DMF (50 mL) was degassed and the
 20 system was charged with argon. The mixture was heated at 100 °C overnight, then cooled to room temperature. The mixture was quenched with saturated aqueous FeSO₄ (20 mL), then diluted with EtOAc (200 mL). The solids were removed by filtration through Celite® and the filtrate was partitioned between water and EtOAc. The organic layer was washed with brine, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified
 25 by silica gel column chromatography (hexanes:EtOAc 30:1 to 20:1) to yield 2.0 g (96%) of the title compound. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.03 (s, 1H), 7.34 (s, 1H), 4.26 –

4.13 (m, 1H), 3.89 (s, 3H), 2.59 (s, 3H), 2.46 – 2.32 (m, 2H), 2.16 – 1.98 (m, 3H), 1.90 – 1.78 (m, 1H).




Compound 89.5. 5-Cyano-2-cyclobutyl-4-methylbenzohydrazide. To a solution of

5 methyl 5-cyano-2-cyclobutyl-4-methylbenzoate (compound **89.4**, 2.0g, 8.73mmol) in EtOH (10 mL) was added anhydrous hydrazine (2mL, excess) at room temperature. The mixture was heated at 90 °C overnight, then the mixture was cooled to room temperature and partitioned between water (60 mL) and EtOAc (200 mL). The organic layer was washed with water (x2), brine, dried (Na₂SO₄), filtered, and concentrated to yield 1.9 g (95%) of the title compound as a white solid. *m/z* (ES⁺) 230 (M+H)⁺. ¹H NMR (400 MHz, Chloroform-d) δ 7.52 (s, 1H), 7.32 (s, 1H), 6.91 (br, 1H), 4.08 (br, 2H), 3.89 (m, 1H), 2.61 – 2.52 (m, 3H), 2.42 – 2.28 (m, 2H), 2.18 – 1.98 (m, 3H), 1.91 – 1.78 (m, 1H).



Compound 89.6. 5-(5-Amino-1,3,4-oxadiazol-2-yl)-4-cyclobutyl-2-


15 **methylbenzonitrile.** To a solution of 5-cyano-2-cyclobutyl-4-methylbenzohydrazide (compound **89.5**, 0.5 g, 2.18 mmol) in H₂O (10 mL) and dioxane (15 mL) was added NaHCO₃ (0.55 g, 6.55 mmol). After the mixture was stirred at room temperature for 5 minutes, BrCN (1.3 mL, 5 M in CH₃CN, 6.55 mmol) was added drop-wise. The mixture was stirred at room temperature for 30 minutes, where upon white solids formed. The mixture was
20 diluted with EtOAc and washed with water, then brine. The organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexanes:EtOAc 1:1 to EtOAc) to yield 0.55 g (theoretical) of the title compound as a white solid. *m/z* (ES⁺) 255 (M+H)⁺. ¹H NMR (400 MHz, Chloroform-d) δ 7.93 (s, 1H), 7.45 (s, 1H), 5.10 (br, 2H), 4.38 (m, 1H), 2.61 (s, 3H), 2.48 –
25 2.34 (m, 2H), 2.17 – 1.98 (m, 3H), 1.91 – 1.79 (m, 1H).



Compound 89.7. 4-Cyclobutyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-


methylbenzonitrile. To a solution of 5-(5-amino-1,3,4-oxadiazol-2-yl)-4-cyclobutyl-2-methylbenzonitrile (compound **89.6**, 0.5 g, 2.0 mmol) in MeOH (40 mL) was added KOH (1.11 g, 20.0 mmol). The mixture was heated at 85 °C overnight, then cooled to 0 °C and neutralized to pH 7 with aqueous 1 M HCl. The mixture was extracted with EtOAc (x2), and the combined organic extracts were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexanes:EtOAc 1:1 to EtOAc) to yield 0.2 g (34%) of the title compound as a white solid. *m/z* (ES⁺) 269

(M+H)⁺. ¹H NMR (400 MHz, Chloroform-d) δ 10.96 (br, 1H), 7.82 (s, 1H), 7.35 (s, 1H), 4.11 (s, 3H), 4.15-4.05 (m, 1H), 2.59 (s, 3H), 2.31 – 2.16 (m, 2H), 2.14 – 1.89 (m, 3H), 1.87 – 1.71 (m, 1H).



Compound 89.8. 4-Cyclobutyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-


methylbenzamide. To a solution of 4-cyclobutyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-methylbenzonitrile (compound **89.7**, 0.15 g, 0.53 mmol) in EtOH (10 mL) was added NH₄OH (0.18 mL, 2.66 mmol, 14.8 M in H₂O), followed by H₂O₂ (1.8 mL, 26.6 mmol, 50% in H₂O). The mixture was stirred at room temperature overnight, then cooled to 0 °C and carefully quenched with 1 M Na₂S₂O₃ solution (26 mL). The mixture was extracted with EtOAc (x2) and the combined organic extracts were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by prep-TLC (5% MeOH in CH₂Cl₂) to yield 0.1 g (63%) of the title compound as a white solid. *m/z* (ES⁺) 287(M+H)⁺. ¹H NMR (400 MHz, Methanol-d₄) δ 7.50 (s, 1H), 7.33 (s, 1H), 4.03 (s, 3H), 3.95 – 4.05 (m, 1H), 2.51 (s, 3H), 2.23 – 2.11 (m, 2H), 2.11 – 1.88 (m, 3H), 1.83 – 1.71 (m, 1H).



Compound 89.9 4-Cyclobutyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-


methylbenzoic acid To a solution of 4-cyclobutyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-

methylbenzamide (compound **89.8**, 0.1 g, 0.33 mmol) in TFA (5 mL) at 0 °C, was added
 5 NaNO₂ (46 mg, 0.66 mmol). The mixture was stirred at 0 °C for 1 hour, then at room
 temperature for 2 hours. The mixture was concentrated under reduced pressure and the
 residue was partitioned between EtOAc and brine. The aqueous layer was extracted with
 EtOAc and the combined organic layers were dried (Na₂SO₄), filtered and concentrated under
 reduced pressure to yield 0.1 g (theoretical) of the title compound as a clear oil. *m/z* (ES⁺) 288
 10 (M+H)⁺.




Compound 89. 4-(1-(4-Cyclobutyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-


methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard
 chemical manipulations and procedures similar to those used for the preparation of compound
 15 **5**, except 4-cyclobutyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-methylbenzoic acid (compound
89.9) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid
 (compound **5.7**). *m/z* (ES⁺) 428 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆) δ 13.63-13.17
 (br, 1H), 7.83 (d, *J*=8.4Hz, 2H), 7.58 (d, *J*=8.4Hz, 2H), 7.49 (s, 1H), 7.39 (s, 1H), 4.48 (m,
 1H), 4.32 (m, 1H), 4.15 (m, 1H), 4.07-3.95 (m, 3H), 3.93 (s, 3H), 2.41 (s, 3H), 2.21-2.11 (m,
 20 2H), 2.06-1.97 (m, 2H), 1.94-1.84 (m, 1H), 1.78-1.68 (m, 1H).



Compound 90. 4-(1-(4-Cyclobutyl-5-(5-methoxy-4*H*-1,2,4-triazol-3-yl)-2-methylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 4-cyclobutyl-5-(5-methoxy-4*H*-1,2,4-triazol-3-yl)-2-methylbenzoic acid (compound **89.9**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) were used in place of 3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**), respectively. *m/z* (ES+) 446 (M+H)⁺. ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.64-13.05 (br, 1H), 7.96 (d, *J*=8.0 Hz, 2H), 7.76 (d, *J*=8.0Hz, 2H), 7.56 (s, 1H), 7.41 (s, 1H), 4.61-4.38(m, 4H), 4.19 (m, 1H), 3.92 (s, 3H), 2.42 (s, 3H), 2.16 (m, 1H), 2.02 (m, 1H), 1.89 (m, 1H), 1.75 (m, 1H).




Compound 91.1. Methyl 4-cyclopropyl-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate. A mixture of methyl 4-cyclopropyl-5-iodo-2-methylbenzoate (compound **86.2**, 4.0 g, 12.7 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (3.86 g, 15.2 mmol), PdCl₂(dppf)CH₂Cl₂ (0.52 g, 0.64 mmol) and potassium acetate (3.73g, 38.10mmol) in DMSO (50 mL) was degassed with argon. The mixture was heated at 80 °C for 18 hours under argon, then cooled to room temperature and diluted with ethyl acetate (300 mL). The mixture was washed with water, aqueous HCl (1 M), saturated aqueous NaHCO₃, and brine, then dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexanes:EtOAc 50:1 to 30:1) to yield 2.63 g (65.6%) of the title compound as a white solid. *m/z* (ES+) 317 (M+H)⁺.




Compound 91.2. 4-Benzyl-3,5-dibromo-4*H*-1,2,4-triazole. To a solution of 3,5-dibromo-4*H*-1,2,4-triazole (3.0 g, 13.3 mmol) in DMF (10 mL) at 0 °C was added NaH (60% in mineral oil, 0.58g, 14.5 mmol)(CAUTION: NaH and DMF can become a runaway

reaction. All necessary safety precautions were performed). After the mixture was stirred at 0 °C for 30 minutes, benzyl bromide (1.57 mL, 13.2 mmol) was added. The mixture was stirred at 0 °C for 2 hours, then the mixture was partitioned between EtOAc (150 mL) and water (30 mL). The organic layer was washed with brine (2 x 30 mL), dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexanes:EtOAc 50:1 to 10:1) to yield 3.71 g (89%) of the title compound as a white solid. *m/z* (ES+) 316, 318, 320 (M+H)⁺.




Compound 91.3. 4-Benzyl-3-bromo-5-methoxy-4H-1,2,4-triazole. To a solution of 4-benzyl-3,5-dibromo-4H-1,2,4-triazole (compound **91.2**, 3.71 g, 11.7 mmol) in MeOH 15 mL) was added NaOMe (1.26 g, 23.4 mmol) and the mixture was refluxed for 18 hours. The mixture was cooled somewhat and additional NaOMe was added (1.26 g, 23.4 mmol). The mixture was refluxed for an additional 5 hours, then cooled to room temperature, and the solvent was removed under reduced pressure. The residue was dissolved in EtOAc (200 mL) and washed with water (30 mL). The organic layer was dried (MgSO₄), filtered and concentrated under reduced pressure to yield 3.13 g (theoretical) of the title compound as a clear oil. *m/z* (ES+) 269 (M+H)⁺.




Compound 91.4. Methyl 5-(4-benzyl-5-methoxy-4H-1,2,4-triazol-3-yl)-4-cyclopropyl-2-methylbenzoate. A mixture of 4-benzyl-3-bromo-5-methoxy-4H-1,2,4-triazole (compound **91.3**, 1.2 g, 4.48 mmol), methyl 4-cyclopropyl-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **91.1**, 1.56 g, 4.93 mmol), PdCl₂(dppf)CH₂Cl₂ (0.37g, 0.45 mmol) and potassium carbonate (3.10 g, 22.5 mmol) in dioxane (50 mL) and water (20 mL) was degassed with argon. The mixture was heated at 90 °C for 18 hours under argon, then cooled to room temperature. The mixture was diluted with ethyl acetate (300 mL) and washed with water then brine, dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel column

chromatography (hexanes:EtOAc 10:1 to 4:1) to yield 1.51 g (89%) of the title compound as thick oil. m/z (ES⁺) 378 (M+H)⁺.




Compound 91.5. Methyl 4-cyclopropyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-

methylbenzoate. A flask containing methyl 5-(4-benzyl-5-methoxy-4H-1,2,4-triazol-3-yl)-4-cyclopropyl-2-methylbenzoate (compound **91.4**, 0.55 g, 1.46 mmol) and Pd/C (10%, 0.25 g) was purged with nitrogen, then MeOH (10 mL) and HCl (4 M in dioxane, 37 μ L, 0.15 mmol) were carefully added. The system was then charged with hydrogen and the mixture was stirred at room temperature for 3 hours. Upon completion, the system was purged with nitrogen, then the mixture was neutralized with a few drops of NH₄OH and filtered through Celite®. The filtrate was concentrated under reduced pressure and the residue was purified using prep-TLC (hexanes: EtOAc 1:1) to yield 0.3 g (71%) of the title compound as a clear oil. m/z (ES⁺) 288 (M+H)⁺.




Compound 91.6. 4-Cyclopropyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-

methylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except methyl 4-cyclopropyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-methylbenzoate (compound **91.5**) was used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **5.6**). m/z (ES⁺) 274 (M+H)⁺.



Compound 91. 4-(1-(4-Cyclopropyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound




5, except 4-cyclopropyl-5-(5-methoxy-4*H*-1,2,4-triazol-3-yl)-2-methylbenzoic acid (compound **91.6**) was used in place of 3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). m/z (ES+) 414 (M+H)⁺. ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.84-12.92 (br, 1H), 7.86-7.80 (m, 2H), 7.61-7.53 (m, 3H), 6.85 (s, 1H), 4.47 (m, 1H), 4.32 (m, 1H), 4.08-3.94 (m, 3H), 2.79 (m, 1H), 0.94 (m, 2H), 0.70 (m, 2H).



Compound 92. 4-(1-(4-Cyclopropyl-5-(5-methoxy-4*H*-1,2,4-triazol-3-yl)-2-methylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 4-cyclopropyl-5-(5-methoxy-4*H*-1,2,4-triazol-3-yl)-2-methylbenzoic acid (compound **91.6**) and 4-(3-fluoroazetidin-3-yl)benzonitrile (compound **43.4**) were used in place of 3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**), respectively. m/z (ES+) 432 (M+H)⁺. ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.66-13.01 (br, 1H), 7.96 (d, $J=8.0$ Hz, 2H), 7.76 (d, $J=8.0$ Hz, 2H), 7.61 (s, 1H), 6.86 (s, 1H), 4.60-4.38 (m, 4H), 3.91 (s, 3H), 2.82 (m, 1H), 2.34 (s, 3H), 0.94 (m, 2H), 0.71 (m, 2H).

The compounds in **TABLE 5** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **78** and **79**.

TABLE 5



Cpd	Name	Structure	<i>m/z</i> (ES+) (M+H) ⁺
96	4-(1-(4-((1S,3S)-3-methoxycyclobutyl)-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile		470
97	4-(1-(4-((1R,3R)-3-methoxycyclobutyl)-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile		470
98	4-(1-(4-((1S,3S)-3-hydroxycyclobutyl)-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile		456

Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
104	4-(1-(4-((1R,3R)-3-hydroxycyclobutyl)-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile		456
107	4-(1-(4-((1S,3S)-3-fluorocyclobutyl)-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile		458
108	4-(1-(4-((1R,3R)-3-fluorocyclobutyl)-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile		458
112	4-(1-(2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)-4-(tetrahydrofuran-2-yl)benzoyl)piperidin-4-yl)benzonitrile		456

The compounds in **TABLE 6** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **89**, **90**, **91**, and **92**.


TABLE 6

Cpd	Name	Structure	<i>m/z</i> (ES+) (M+H) ⁺
99	4-(1-(4-cyclobutyl-5-(5-(2-methoxyethoxy)-4H-1,2,4-triazol-3-yl)-2-methylbenzoyl)piperidin-4-yl)benzonitrile		500
100	4-(1-(4-cyclobutyl-5-(5-(2-hydroxyethoxy)-4H-1,2,4-triazol-3-yl)-2-methylbenzoyl)piperidin-4-yl)benzonitrile		486
133	4-(1-(4-cyclopropyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-methylbenzoyl)-4-fluoropiperidin-4-yl)benzonitrile		460
134	4-(1-(4-cyclopropyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)-2-methylbenzoyl)piperidin-4-yl)benzonitrile		442

Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
137	4-(1-(4-cyclopropyl-2-ethyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)benzoyl)azetidin-3-yl)benzonitrile		428
138	4-(1-(4-cyclopropyl-2-ethyl-5-(5-methoxy-4H-1,2,4-triazol-3-yl)benzoyl)-3-fluoroazetidin-3-yl)benzonitrile		446

The compounds in **TABLE 7** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **62**, **63** and **64**.

5 **TABLE 7**





Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
103	5-(4-(4-cyanophenyl)piperidine-1-carbonyl)-2-cyclobutyl-4-methylbenzamide		402



Cpd	Name	Structure	<i>m/z</i> (ES+) (M+H) ⁺
106	2-cyclobutyl-5-(4-(imidazo[1,2-a]pyridin-7-yl)piperidine-1-carbonyl)-N,4-dimethylbenzamide		431
110	2-cyclobutyl-5-(4-(imidazo[1,2-a]pyridin-6-yl)piperidine-1-carbonyl)-N,4-dimethylbenzamide		431
116	5-(4-(benzofuran-7-yl)piperidine-1-carbonyl)-2-cyclobutyl-N,4-dimethylbenzamide		431

The compounds in **TABLE 8** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **78**, **79**, **80** and **81**.

TABLE 8


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
111	4-(1-(4-cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)-2-methylbenzonitrile		454
113	4-(1-(4-cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)-2-fluorobenzonitrile		458
114	4-(1-(4-cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)-3-fluorobenzonitrile		458
115	2-chloro-4-(1-(4-cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile		474




Cpd	Name	Structure	<i>m/z</i> (ES+) (M+H) ⁺
117	4-(1-(4-cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)-3-methylbenzonitrile		454
118	3-chloro-4-(1-(4-cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile		474
119	4-(1-(4-cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)-3-methoxybenzonitrile		470
120	4-(1-(4-cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)-2-methoxybenzonitrile		470


Cpd	Name	Structure	<i>m/z</i> (ES+) (M+H) ⁺
135	(4-(4-bromophenyl)piperidin-1-yl)(4-cyclobutyl-2-methyl-5-(5-methyl-4H-1,2,4-triazol-3-yl)phenyl)methanone		493
150	4-(1-(4-cyclobutyl-3-(5-methyl-4H-1,2,4-triazol-3-yl)benzoyl)azetidin-3-yl)benzonitrile		398

The compounds in **TABLE 9** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **78**, **79**, **80** and **81**.


5 **TABLE 9**

Cpd	Name	Structure	<i>m/z</i> (ES+) (M+H) ⁺
121	4-(1-(4-cyclobutyl-5-(5-(hydroxymethyl)-4H-1,2,4-triazol-3-yl)-2-methylbenzoyl)-4-fluoropiperidin-4-yl)benzonitrile		474


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
123	5-(5-(4-(4-cyanophenyl)piperidine-1-carbonyl)-2-cyclobutyl-4-methylphenyl)-4H-1,2,4-triazole-3-carbonitrile		451
130	methyl 2-(5-(5-(4-(4-cyanophenyl)piperidine-1-carbonyl)-2-cyclobutyl-4-methylphenyl)-4H-1,2,4-triazol-3-yl)acetate		498
131	2-(5-(5-(4-(4-cyanophenyl)piperidine-1-carbonyl)-2-cyclobutyl-4-methylphenyl)-4H-1,2,4-triazol-3-yl)acetic acid		484




Compound 95. 4-(1-(3-(3-Chloro-4-methyl-1*H*-pyrazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 250, except methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound 5.4) was used in place of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound 160.1). m/z (ES⁺) 391 (M+H)⁺.




Compound 101.1. 4-Fluoro-5-iodo-2-methylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 1.3, except 4-fluoro-2-methylbenzoic acid was used in place of 2,4-dimethylbenzoic acid.




Compound 101.2. Methyl 4-fluoro-5-iodo-2-methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 6.1, except 4-fluoro-5-iodo-2-methylbenzoic acid (compound 101.1) was used in place of 4-bromo-2-methylbenzoic acid.



Compound 101.3. Methyl 5-cyano-4-fluoro-2-methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 6.4, except methyl 4-fluoro-5-iodo-2-methylbenzoate (compound 101.2) was used in place of methyl 4-cyclobutyl-5-iodo-2-methylbenzoate (compound 6.3).



Compound 101.4. 4-(1-(4-Fluoro-2-methyl-5-(5-methyl-4*H*-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 78, except methyl 5-cyano-4-fluoro-2-methylbenzoate (compound **101.3**) was used in place of methyl 5-cyano-4-cyclobutyl-2-methylbenzoate (compound **6.4**) and 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 6-(piperidin-4-yl)imidazo[1,2-*a*]pyridine hydrochloride (compound **78.5**).








Compound 101. 4-(1-(2-Methyl-5-(5-methyl-4*H*-1,2,4-triazol-3-yl)-4-(piperidin-1-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **25**, except 4-(1-(4-fluoro-2-methyl-5-(5-methyl-4*H*-1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile (compound **101.4**) was used in place of 4-(1-(3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-fluorobenzoyl)piperidin-4-yl)benzonitrile (compound **25.1**) and piperidine was used in place of azetidine hydrochloride. m/z (ES⁺) 469 (M+H)⁺.


The compounds in **TABLE 10** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **101**.

TABLE 10

Cpd	Name	Structure	m/z (ES ⁺) (M+H) ⁺
102	4-(1-(2-methyl-5-(5-methyl-4 <i>H</i> -1,2,4-triazol-3-yl)-4-morpholinobenzoyl)piperidin-4-yl)benzonitrile		471


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
105	4-(1-(2-methyl-5-(5-methyl-4 <i>H</i> -1,2,4-triazol-3-yl)-4-(3-methylpyrrolidin-1-yl)benzoyl)piperidin-4-yl)benzonitrile		469
109	4-(1-(4-(dimethylamino)-2-methyl-5-(5-methyl-4 <i>H</i> -1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile		429
122	4-(1-(2-methyl-5-(5-methyl-4 <i>H</i> -1,2,4-triazol-3-yl)-4-(4-methylpiperazin-1-yl)benzoyl)piperidin-4-yl)benzonitrile		484
124	(<i>S</i>)-4-(1-(2-methyl-5-(5-methyl-4 <i>H</i> -1,2,4-triazol-3-yl)-4-(2-methylpyrrolidin-1-yl)benzoyl)piperidin-4-yl)benzonitrile		469
125	(<i>S</i>)-4-(1-(4-(3-methoxypyrrolidin-1-yl)-2-methyl-5-(5-methyl-4 <i>H</i> -1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile		485

Cpd	Name	Structure	<i>m/z</i> (ES+) (M+H) ⁺
126	(<i>R</i>)-4-(1-(4-(3-methoxypyrrolidin-1-yl)-2-methyl-5-(5-methyl-4 <i>H</i> -1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile		485
127	(<i>R</i>)-4-(1-(2-methyl-5-(5-methyl-4 <i>H</i> -1,2,4-triazol-3-yl)-4-(2-methylpyrrolidin-1-yl)benzoyl)piperidin-4-yl)benzonitrile		469
128	4-(1-(2-methyl-5-(5-methyl-4 <i>H</i> -1,2,4-triazol-3-yl)-4-(3-methylazetidin-1-yl)benzoyl)piperidin-4-yl)benzonitrile		455
129	4-(1-(4-(diethylamino)-2-methyl-5-(5-methyl-4 <i>H</i> -1,2,4-triazol-3-yl)benzoyl)piperidin-4-yl)benzonitrile		457




Compound 132.1. Methyl 4-fluoro-3-iodobenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the

preparation of compound **6.1**, except 4-fluoro-3-iodobenzoic acid was used in place of 4-bromo-2-methylbenzoic acid.




Compound 132.2. Methyl 3-(2-bromoacetyl)-4-fluorobenzoate. The title

- 5 compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **27.3**, except methyl 4-fluoro-3-iodobenzoate (compound **132.1**) was used in place of methyl 3-(2-bromoacetyl)-4-methylbenzoate (compound **27.2**).



10 **Compound 132.3. 4-(1-(4-Fluoro-3-(2-methyl-1H-imidazol-5-**


yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1**, except methyl 3-(2-bromoacetyl)-4-fluorobenzoate (compound **132.1**) was used in place of methyl 5-(2-bromopropanoyl)-2,4-dimethylbenzoate (compound **1.6**).




15 **Compound 132. 4-(1-(4-(Azetidin-1-yl)-3-(2-methyl-1H-imidazol-5-**

yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **25**, except 4-(1-(4-fluoro-3-(2-methyl-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile (compound **132.3**) was used in place of 4-(1-(3-(2,4-dimethyl-1H-imidazol-5-yl)-4-fluorobenzoyl)piperidin-4-yl)benzonitrile (compound **25.1**). m/z (ES⁺) 426 (M+H)⁺.


20




Compound 139. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-2-methyl-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 16, except 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used instead of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**). m/z (ES⁺) 382 (M+H)⁺.




Compound 151.1. 2-(Morpholin-4-yl)-5-nitropyridin-4-amine. Into a 50-mL sealed tube, were placed a solution of 2-chloro-5-nitropyridin-4-amine (500 mg, 2.88 mmol) in *N,N*-dimethylformamide (20 mL) and morpholine (503 μ L, 5.77 mmol). The reaction mixture was stirred overnight at 55 $^{\circ}$ C. The reaction was then quenched by the addition of 100 mL of water. The reaction mixture was extracted with 3 x 150 mL of ethyl acetate. The organic layers were combined, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. This resulted in 733 mg (crude) of the title compound as a yellow solid.



Compound 151.2. 6-Morpholinopyridine-3,4-diamine. Into a 50-mL round-bottom flask were placed 2-(morpholin-4-yl)-5-nitropyridin-4-amine (compound **151.1**, 350 mg, 1.56 mmol), methanol (20 mL), THF (10 mL), and Pd/C (35 mg). The above solution was purged with N₂ and then H₂. The reaction mixture was stirred for 4h at room temperature. The solids were filtered out. The resulting mixture was concentrated under reduced pressure. This resulted in 280 mg (92%) of the title compound as a pink solid.




Compound 151.3. 5-Formyl-2,4-dimethylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **4.1**, except 5-iodo-2,4-dimethylbenzoic acid (compound **1.3**) was used in place of 3-bromo-4-methylbenzoic acid.



Compound 151.4. 4-(1-(5-Formyl-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile.


Into a 50-mL round-bottom flask, was placed a solution of 5-formyl-2,4-dimethylbenzoic acid (compound **151.3**, 500 mg, 2.81 mmol) in DCM (10 mL). 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**, 545 mg, 2.80 mmol), DIEA (1.4 mL, 8.41 mmol) and HBTU (1.60 g, 4.22 mmol) were added. The reaction mixture was stirred for 3h at room temperature. The reaction mixture was diluted with 150 mL of EtOAc, then washed with 2 x 50 mL of NH₄Cl (sat.) and 1 x 50 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/1) as eluent to furnish 480 mg (54%) of the title compound as a white solid.




Compound 151. 4-(1-(2,4-Dimethyl-5-(6-morpholino-3H-imidazo[4,5-c]pyridin-2-yl)benzoyl)azetidin-3-yl)benzonitrile. Into a 50-mL round-bottom flask, was placed a solution of 4-(1-(5-formyl-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile (compound **151.4**, 159 mg, 0.50 mmol) in ethanol (16 mL). 6-Morpholinopyridine-3,4-diamine (compound **151.2**, 194 mg, 1.00 mmol) and NH₄OAc (308 mg, 4.00 mmol) were added to the reaction. The reaction mixture was stirred for 3 days at 70 °C under air. The pH of the solution was

adjusted to 8-9 with sodium bicarbonate (sat.). The reaction mixture was extracted with 1 x 100 mL of ethyl acetate. The organic layer was washed with 3 x 50 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The crude product (200 mg) was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-010(Waters)):


5 Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150 mm; mobile phase, WATER WITH 0.05% TFA and CH₃CN (20.0% CH₃CN up to 36.0% in 10 min, up to 100.0% in 1 min, down to 20.0% in 2 min); Detector, UV 254 and 220 nm. This resulted in 94.1 mg (38%) of the title compound as a brown solid. m/z (ES⁺) 493 (M+H)⁺.




10 **Compound 152. 4-(1-(4-Methyl-3-(6-morpholino-3H-imidazo[4,5-c]pyridin-2-yl)benzoyl)azetidin-3-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **151**, except 3-formyl-4-methylbenzoic acid (compound **4.1**) was used in place of 5-formyl-2,4-dimethylbenzoic acid (compound **151.3**). m/z (ES⁺) 479 (M+H)⁺.




15 **Compound 153.1. 6-(Pyrrolidin-1-yl)pyridine-3,4-diamine.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **151.1** and **151.2**, except pyrrolidine was used in place of morpholine.




Compound 153. 4-(1-(2,4-Dimethyl-5-(6-(pyrrolidin-1-yl)-3H-imidazo[4,5-c]pyridin-2-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **151**, except 6-(pyrrolidin-1-yl)pyridine-3,4-diamine (compound **153.1**) was used in place of 6-morpholinopyridine-3,4-diamine (compound **151.2**). m/z (ES⁺) 477 (M+H)⁺.




Compound 154. 4-(1-(4-Methyl-3-(6-(pyrrolidin-1-yl)-3H-imidazo[4,5-c]pyridin-2-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **151**, except 3-formyl-4-methylbenzoic acid (compound **4.1**) was used in place of 5-formyl-2,4-dimethylbenzoic acid (compound **151.3**) and 6-(pyrrolidin-1-yl)pyridine-3,4-diamine (compound **153.1**) was used in place of 6-morpholinopyridine-3,4-diamine (compound **151.2**). m/z (ES⁺) 463 (M+H)⁺.




Compound 155.1. 6-(Azetidin-1-yl)pyridine-3,4-diamine. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **151.1** and **151.2**, except azetidine was used in place of morpholine.




Compound 155. 4-(1-(3-(6-(Azetidin-1-yl)-3H-imidazo[4,5-c]pyridin-2-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 151, except 3-formyl-4-methylbenzoic acid (compound 4.1) was used in place of 5-formyl-2,4-dimethylbenzoic acid (compound 151.3) and 6-(azetidin-1-yl)pyridine-3,4-diamine (compound 155.1) was used in place of 6-morpholinopyridine-3,4-diamine (compound 151.2). m/z (ES⁺) 449 (M+H)⁺.



Compound 156.1. 3-(2-(Methoxymethyl)-4-methyl-1H-imidazol-5-yl)benzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 2.3, except 3-iodobenzoic acid was used in place of 5-iodo-2,4-dimethylbenzoic acid (compound 1.3).




Compound 156. 4-(1-(3-(2-(Methoxymethyl)-4-methyl-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 2, except 3-(2-(methoxymethyl)-4-methyl-1H-imidazol-5-yl)benzoic acid (compound 156.1) was used in place of 5-(2-(methoxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound 2.3). m/z (ES⁺) 415 (M+H)⁺.




Compound 157.1. Methyl 3-(2-(2-hydroxyethyl)-1H-imidazol-5-yl)-4-

methylbenzoate. Into a 250-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of methyl 3-(2-bromoacetyl)-4-methylbenzoate (compound **27.2**, 10 g, 36.89 mmol) in *N,N*-dimethylformamide (150 mL). K_2CO_3 (30 g, 215.5 mmol) and 3-hydroxypropanimidamide hydrochloride (compound **23.1**, 15 g, 120.4 mmol) were added to the reaction. The reaction mixture was stirred for 12h at 80 °C, then concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (2:1) as eluent to furnish 2 g (21%) of the title compound as a light yellow oil.




Compound 157.2. Methyl 3-(2-(2-(azetidin-1-yl)ethyl)-1H-imidazol-5-yl)-4-

methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **24.1** and **24**, except methyl 3-(2-(2-hydroxyethyl)-1H-imidazol-5-yl)-4-methylbenzoate (compound **157.1**) was used in place of 4-(1-(3-(2-(2-hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile (compound **23**) and azetidine was used in place of dimethylamine.




Compound 157.3. 3-(2-(2-(Azetidin-1-yl)ethyl)-4-chloro-1H-imidazol-5-yl)-4-


methylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **27.4** and **27.5**, except methyl 3-(2-(2-(azetidin-1-yl)ethyl)-1H-imidazol-5-yl)-4-methylbenzoate (compound **157.2**) was used in place of methyl 3-(2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoate (compound **27.3**).



Compound 157. 4-(1-(3-(2-(2-(Azetidin-1-yl)ethyl)-4-chloro-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 27, except 3-(2-(2-(azetidin-1-yl)ethyl)-4-chloro-1H-imidazol-5-yl)-4-methylbenzoic acid (compound 157.3) was used in place of 3-(4-chloro-2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoic acid (compound 27.5) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound 5.2) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound 1.2). m/z (ES⁺) 460 (M+H)⁺.




Compound 158. 4-(1-(3-(2-(2-(Azetidin-1-yl)ethyl)-4-chloro-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 27, except 3-(2-(2-(azetidin-1-yl)ethyl)-4-chloro-1H-imidazol-5-yl)-4-methylbenzoic acid (compound 157.3) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound 1.2). m/z (ES⁺) 488 (M+H)⁺.




Compound 159.1. 2-Isopropyl-4-methyl-1H-imidazole. A solution of isobutyraldehyde (4.5 mL, 50 mmol) in ethanol (25 mL) was treated with ammonium hydroxide (28% w/w, 25 mL) at 55 °C. Methylglyoxal (40% in H₂O, 28 mL, 63 mmol) was added dropwise. The resulting mixture was stirred at 60 °C for 16 hours and the solvent was removed under reduced pressure. The residue was partitioned between EtOAc (50 mL) and water (30 mL). The organic phase was washed by brine, dried over MgSO₄ and concentrated


under reduced pressure. The residue was purified by flash chromatography (SiO₂; 0-4 % MeOH in dichloromethane) to give 3.96 g (64%) of the title compound as a yellow solid. m/z (ES+) 125 (M+H)⁺.




- 5 **Compound 159.2. 5-Iodo-2-isopropyl-4-methyl-1H-imidazole.** 2-Isopropyl-4-methyl-1H-imidazole (3.96 g, 32 mmol) and *N*-iodosuccinimide (8.33 g, 35 mmol) were dissolved in acetonitrile (100 mL) and heated to reflux for 16 hours. The reaction was concentrated under reduced pressure and the residue was partitioned between EtOAc (50 mL) and water (50 mL). The organic phase was washed with brine, dried over MgSO₄ and
- 10 concentrated under reduced pressure. The residue was purified by flash chromatography (SiO₂; 0-40 % EtOAc in hexane) to give 4.87 g (61%) of the title compound as a yellow solid. m/z (ES+) 251 (M+H)⁺.



- 15 **Compound 160.1. Methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.4**, except methyl 5-iodo-2,4-dimethylbenzoate (compound **31.1**) was used in place of methyl 3-iodo-4-methylbenzoate (compound **5.3**). m/z (ES+) 291(M+H)⁺.




- 20 **Compound 160.2. 2-Cyclopropyl-5-iodo-4-methyl-1H-imidazole.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159.2**, except cyclopropanecarbaldehyde was used in place of isobutyraldehyde.




Compound 160.3. Methyl 5-(2-cyclopropyl-4-methyl-1H-imidazol-5-yl)-2,4-

dimethylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.6**, except methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) and 2-cyclopropyl-5-iodo-4-methyl-1H-imidazole (compound **160.2**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**). m/z (ES⁺) 285 (M+H)⁺.



Compound 160.4. 5-(2-Cyclopropyl-4-methyl-1H-imidazol-5-yl)-2,4-


dimethylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except methyl 5-(2-cyclopropyl-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **160.3**) was used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **5.6**). m/z (ES⁺) 271 (M+H)⁺.



Compound 160. 4-(1-(5-(2-Cyclopropyl-4-methyl-1H-imidazol-5-yl)-2,4-


dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 5-(2-cyclopropyl-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **160.4**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). m/z (ES⁺) 411 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆): δ 11.51 (br, 1H), 7.85-7.80 (m, 2H), 7.62-7.56 (m, 2H), 7.20 and 7.13 (2 singlets, Ar-H, 1H), 7.10 and 7.07 (2 singlets, Ar-H, 1H), 4.49-4.40 (m, 1H), 4.36-4.28 (m, 1H), 4.06-3.88 (m, 3H), 2.33

and 2.30 (2 singlets, CH₃, 3H), 2.23 and 2.18 (2 singlets, CH₃, 3H), 2.07 and 1.93 (2 singlets, amide rotamers, CH₃, 3H), 1.89-1.82 (m, 1H), 0.88-0.74 (m, 4H).




Compound 161. 4-(1-(5-(2-Cyclopropyl-4-methyl-1H-imidazol-5-yl)-2,4-

dimethylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 5-(2-cyclopropyl-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **160.4**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). *m/z* (ES⁺) 429 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆): δ 11.52 (br, 1H), 7.95 (d, *J*=8.4Hz, 2H), 7.76 (dd, *J*=8.4Hz, 3.2Hz, 2H), 7.21 (d, *J*=6.8Hz, 1H), 7.13(s, 1H), 4.56-4.34(m, 4H), 2.35 and 2.32 (2 singlets, CH₃, 3H), 2.43 and 2.18 (2 singlets, CH₃, 3H), 2.07 and 1.93 (2 singlets, CH₃, 3H), 1.90-1.80 (m, 1H), 0.88-0.74(m, 4H).




Compound 159.3. 5-(2-Isopropyl-4-methyl-1H-imidazol-5-yl)-2,4-


dimethylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except 5-iodo-2-isopropyl-4-methyl-1H-imidazole (compound **159.2**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**).



Compound 159. 4-(1-(5-(2-Isopropyl-5-methyl-1H-imidazol-4-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The mixture of 5-(2-isopropyl-5-methyl-1H-imidazol-4-yl)-2, 4-dimethylbenzoic acid (compound **159.3**, 250 mg, 0.92 mmol), 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**, 195 mg, 1.0 mmol), EDCI (264 mg, 1.4 mmol), HOBT (170 mg, 1.0 mmol) and DIEA (640 μ L, 3.7 mmol) in DMF (10 mL) was stirred at room temperature for 16 hours. The reaction was diluted with saturated NaHCO_3 and extracted with EtOAc (60 mL). The organic phase was washed with brine (3 X 20 mL), dried over MgSO_4 and concentrated under reduced pressure. The residue was purified by preparative TLC with 8% methanol in dichloromethane and lyophilized to give 85 mg (22%) of the title compound as a white solid. m/z (ES+) 413 (M+H)⁺.




Compound 162. 4-(1-(3-(2-Isopropyl-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159**, except methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used in place of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**). m/z (ES+) 399 (M+H)⁺.




Compound 163. 4-(1-(3-(2-Isopropyl-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159**, except 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) and methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used in place of

methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**). m/z (ES⁺) 427 (M+H)⁺.




Compound 164.1. 5-Iodo-4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazole.

- 5 The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159.2**, except tetrahydro-2H-pyran-4-carbaldehyde was used in place of isobutyraldehyde.



Compound 164.2. 2,4-Dimethyl-5-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-


- 10 **imidazol-5-yl)benzoic acid.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except 5-iodo-4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazole (compound **164.1**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in
15 place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**).




Compound 164. 4-(1-(2,4-Dimethyl-5-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile.

- 20 The mixture of 2,4-dimethyl-5-(5-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-4-yl)benzoic acid (compound **164.2**, 157 mg, 0.5 mmol), 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**, 126 mg, 0.65 mmol), EDCI (143 mg, 0.75 mmol), HOBt (93 mg, 0.55 mmol) and DIEA (345 μ L, 2.00 mmol) in DMF (4 mL) was stirred at room temperature for 16 hours. The reaction was diluted with water and extracted with EtOAc (30 mL). The organic phase was washed with
25 saturated NaHCO₃ (10 mL), brine (3 X 20 mL), dried over MgSO₄ and concentrated under


reduced pressure. The residue was purified by preparative TLC with 5% methanol in dichloromethane and lyophilized to give 95 mg (42%) of the title compound as a white solid. m/z (ES+) 455 (M+H)⁺.



5 **Compound 165. 4-(1-(2,4-Dimethyl-5-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **164**, except 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound
10 **5.2**). m/z (ES+) 483 (M+H)⁺.




Compound 166. 4-(4-Fluoro-1-(4-methyl-3-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **164**, except 4-(4-fluoropiperidin-4-yl)benzonitrile hydrochloride (compound **13.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound
15 **5.2**). m/z (ES+) 501 (M+H)⁺.




20 **Compound 167. 4-(1-(2,4-Dimethyl-5-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoyl)-3-fluoroazetidin-3-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **164**, except 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride


(compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 459 (M+H)⁺.



Compound 168.1. 4-Methyl-3-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except 5-iodo-4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazole (compound **164.1**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**).




Compound 168. 4-(1-(4-Methyl-3-(5-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-4-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **164**, except 4-methyl-3-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoic acid (compound **168.1**) was used in place of 2,4-dimethyl-5-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoic acid (compound **164.2**). m/z (ES+) 441 (M+H)⁺.




Compound 169. 4-(1-(4-Methyl-3-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **164**, except 4-methyl-3-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoic acid (compound **168.1**) was used in place of 2,4-dimethyl-5-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoic acid (compound **164.2**) and 4-


(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) m/z (ES+) 469 (M+H)⁺.




Compound 170. 4-(3-Fluoro-1-(4-methyl-3-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **164**, except 4-methyl-3-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoic acid (compound **168.1**) was used in place of 2,4-dimethyl-5-(4-methyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoic acid (compound **164.2**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 459 (M+H)⁺.




Compound 171.1. 5-Iodo-4-methyl-2-(tetrahydrofuran-3-yl)-1H-imidazole. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159.2**, except tetrahydrofuran-3-carbaldehyde was used in place of isobutyraldehyde.




Compound 171.2. 2,4-Dimethyl-5-(4-methyl-2-(tetrahydrofuran-3-yl)-1H-imidazol-5-yl)benzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except 5-iodo-4-methyl-2-(tetrahydrofuran-3-yl)-1H-imidazole (compound **171.1**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**).



Compound 171. 4-(1-(2,4-Dimethyl-5-(5-methyl-2-(tetrahydrofuran-3-yl)-1H-imidazol-4-yl)benzoyl)azetidin-3-yl)benzonitrile. The mixture of 2,4-dimethyl-5-(5-methyl-2-(tetrahydrofuran-3-yl)-1H-imidazol-4-yl)benzoic acid (compound **171.2**, 100 mg, 0.30 mmol), 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**, 71 mg, 0.37 mmol), EDCI (95 mg, 0.50 mmol), HOBt (20 mg, 0.10 mmol) and DIEA (207 μ L, 1.20 mmol) in DMF (5 mL) was stirred at room temperature for 16 hours. The reaction was concentrated under reduced pressure, diluted with saturated NaHCO_3 (10 mL) and extracted with EtOAc (30 mL). The organic phase was washed with brine (3 X 10 mL), dried over MgSO_4 and concentrated under reduced pressure. The residue was purified by preparative TLC with 6 % methanol in dichloromethane and lyophilized to give 16 mg (12%) of the title compound as a white solid. m/z (ES⁺) 441 (M+H)⁺.




Compound 172.1. 4-Methyl-3-(4-methyl-2-(tetrahydrofuran-3-yl)-1H-imidazol-5-yl)benzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except 5-iodo-4-methyl-2-(tetrahydrofuran-3-yl)-1H-imidazole (compound **171.1**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**).




Compound 172. 4-(1-(4-Methyl-3-(4-methyl-2-(tetrahydrofuran-3-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **171**, except 4-methyl-3-(4-methyl-2-(tetrahydrofuran-3-yl)-1H-imidazol-5-


yl)benzoic acid (compound **172.1**) was used in place of 2,4-dimethyl-5-(4-methyl-2-(tetrahydrofuran-3-yl)-1H-imidazol-5-yl)benzoic acid (compound **171.2**). m/z (ES+) 427 (M+H)⁺.




- 5 **Compound 173.1. 5-Iodo-4-methyl-2-(tetrahydrofuran-2-yl)-1H-imidazole.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159.2**, except tetrahydrofuran-2-carbaldehyde was used in place of isobutyraldehyde. m/z (ES+) 279 (M+H)⁺.




- 10 **Compound 173.2. 2,4-Dimethyl-5-(4-methyl-2-(tetrahydrofuran-2-yl)-1H-imidazol-5-yl)benzoic acid.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except 5-iodo-4-methyl-2-(tetrahydrofuran-2-yl)-1H-imidazole (compound **173.1**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**).




- 20 **Compound 173. 4-(1-(2,4-Dimethyl-5-(5-methyl-2-(tetrahydrofuran-2-yl)-1H-imidazol-4-yl)benzoyl)azetidin-3-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **171**, except 2,4-dimethyl-5-(4-methyl-2-(tetrahydrofuran-2-yl)-1H-imidazol-5-yl)benzoic acid (compound **173.2**) was used in place of 2,4-dimethyl-5-(4-methyl-2-(tetrahydrofuran-3-yl)-1H-imidazol-5-yl)benzoic acid (compound **171.2**). m/z (ES+) 441 (M+H)⁺.




Compound 174.1. 4-Methyl-3-(4-methyl-2-(tetrahydrofuran-2-yl)-1H-imidazol-5-yl)benzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except 5-iodo-4-methyl-2-(tetrahydrofuran-2-yl)-1H-imidazole (compound **173.1**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**).



Compound 174. 4-(1-(4-Methyl-3-(4-methyl-2-(tetrahydrofuran-2-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **171**, except 4-methyl-3-(4-methyl-2-(tetrahydrofuran-2-yl)-1H-imidazol-5-yl)benzoic acid (compound **174.1**) was used in place of 2,4-dimethyl-5-(4-methyl-2-(tetrahydrofuran-3-yl)-1H-imidazol-5-yl)benzoic acid (compound **171.2**). m/z (ES⁺) 427 (M+H)⁺.




Compound 175.1. 4-Methyl-2-(3-methyloxetan-3-yl)-1H-imidazole. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16.2**, except 3-methyloxetane-3-carbaldehyde was used in place of acetaldehyde and 2-oxopropanal was used in place of 3,3,3-trifluoro-2-oxopropanal (compound **16.1**). m/z (ES⁺) 153 (M+H)⁺.




Compound 175.2. 5-Iodo-4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazole. NIS (2.61g, 11.58mmol) was added portion-wise to a solution of 4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazole (compound **175.1**, 1.76 g, 11.58 mmol) in acetonitrile (60mL). The mixture


was stirred at room temperature for 1 hour, then was partitioned between EtOAc (300 mL) and water (80 mL). The organic layer was washed with saturated sodium thiosulfate (50 mL), brine (50 mL), dried (MgSO₄) and concentrated under reduced pressure to give the title compound as a light yellow solid (3.0 g, 93%). *m/z* (ES⁺) 279 (M+H)⁺.



Compound 175.3. Methyl 2,4-dimethyl-5-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 5.6, except methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound 160.1) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound 5.4) and 5-iodo-4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazole (compound 175.2) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound 5.5). *m/z* (ES⁺) 315 (M+H)⁺.




Compound 175.4. 2,4-Dimethyl-5-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 5.7, except methyl 2,4-dimethyl-5-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoate (compound 175.3) was used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound 5.6). *m/z* (ES⁺) 301 (M+H)⁺.




Compound 175. 4-(1-(2,4-Dimethyl-5-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using

standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 2,4-dimethyl-5-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoic acid (compound **175.4**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). m/z (ES⁺) 441 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆):

5 δ 11.72 (br, 1H), 7.83 (d, $J=8.4$ Hz, 2H), 7.59 (d, $J=8.0$ Hz, 2H), 7.21 and 7.18 (2 singlets, Ar-H, 1H), 7.13 (s, 1H), 4.90 (d, $J=4.8$ Hz, 2H), 4.5-4.43 (m, 1H), 4.12 (d, $J=5.6$ Hz, 2H), 4.36-4.28 (m, 1H), 4.07-3.90 (m, 3H), 2.32 (s, 3H), 2.28 and 2.18 (2 singlets, CH₃, 3H), 2.13 and 2.00 (2 singlets, CH₃, 3H), 1.67 (s, 3H).




10 **Compound 176. 4-(1-(2,4-Dimethyl-5-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoyl)-3-fluoroazetidin-3-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 2,4-dimethyl-5-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoic acid (compound **175.4**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 459 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆): δ 11.73 (br, 1H), 7.95 (d, $J=8.0$ Hz, 2H), 7.76 (d, $J=8.4$ Hz, 2H), 7.25 and 7.24 (2 singlets, Ar-H, 1H), 7.19 and 7.16 (2 singlets, Ar-H, 1H), 4.94-4.86 (m, 2H), 4.57-4.35 (m, 6H), 2.36 and 2.34 (2 singlets, CH₃, 3H), 2.29 and 2.19 (2 singlets, CH₃, 3H), 2.14 and 2.00 (2 singlets, CH₃, 3H), 1.67 (s, 3H).



15 **Compound 177.1. Methyl 4-methyl-3-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoate.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.6**,


25

except 5-iodo-4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazole (compound **175.2**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**). m/z (ES+) 301 (M+H)⁺.




Compound 177.2. 4-Methyl-3-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-

yl)benzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except methyl 4-methyl-3-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoate (compound **177.1**) was used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **5.6**). m/z (ES+) 287 (M+H)⁺.



Compound 177. 4-(1-(4-Methyl-3-(4-methyl-2-(3-methyloxetan-3-yl)-1H-


imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 4-methyl-3-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoic acid (compound **177.2**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). m/z (ES+) 427 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆): δ 11.79 (br, 1H), 7.84 (d, J =8.4Hz, 2H), 7.63 (d, J =8.4Hz, 2H), 7.60-7.46 (m, 2H), 7.42-7.28 (m, 1H), 4.91 (d, J =5.2Hz, 2H), 4.76-4.67 (m, 1H), 4.52-4.36 (m, 4H), 4.09-3.97 (m, 2H), 2.34 and 2.24 (2 singlets, CH₃, 3H), 2.16 and 2.02 (2 singlets, CH₃, 3H), 1.68 (s, 3H).




Compound 178. 4-(3-Fluoro-1-(4-methyl-3-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the

preparation of compound **5**, except 4-methyl-3-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoic acid (compound **177.2**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-

yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 445 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆): δ 11.80 (br, 1H), 7.96 (dd, $J=8.0$ Hz, 2.4Hz, 2H), 7.80 (d, $J=8.4$ Hz, 2H), 7.67-7.52 (m, 2H), 7.44-7.33 (m, 1H), 4.95-4.89 (m, 2H), 4.89-4.46 (m, 4H), 4.46-4.40 (m, 2H), 3.36 and 2.26 (2 singlets, CH₃, 3H), 2.17 and 2.03 (2 singlets, CH₃, 3H), 1.69 and 1.68 (2 singlets, CH₃, 3H).




Compound 179. 4-(1-(4-Methyl-3-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 4-methyl-3-(4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)benzoic acid (compound **177.2**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 455 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆): δ 11.78 (br, 1H), 7.78 (d, $J=8.0$ Hz, 2H), 7.53 (d, $J=8.0$ Hz, 2H), 7.40-7.23 (m, 3H), 4.92 (t, $J=6.0$ Hz, 5.2Hz, 2H), 4.78-4.52 (m, 1H), 4.43 (t, $J=5.2$ Hz, 4.8Hz, 2H), 3.91-3.71 (m, 1H), 3.24-3.08 (m, 1H), 2.98-2.88 (m, 2H), 2.34 and 2.25 (2 singlets, CH₃, 3H), 2.16 and 2.03 (2 singlets, amide rotamers, CH₃, 3H), 1.91-1.59 (m, 4H), 1.69 and 1.68 (2 singlets, CH₃, 3H).




Compound 180.1. 4-Methyl-2-(oxetan-3-yl)-1H-imidazole. Dess-Martin reagent (4.33g, 10.21mmol) was added to a solution of oxetan-3-ylmethanol (0.9g, 10.21 mmol) in CH₂Cl₂ (10 mL) at 0 °C. The mixture was stirred at 0 °C for 30 minutes and at room

temperature for 2 hours. CH_2Cl_2 was then removed under reduced pressure. The residue was dissolved in EtOH (10 mL). NH_4OH (5 mL) was added, followed by 2-oxopropanal (40% in water, 2.76 mL, 15.32 mmol). The mixture was stirred at room temperature for 4 hours.


Water (10 mL) was added to the mixture and the mixture was lyophilized. The dried residue was purified with column chromatography (2.5% MeOH to 5% MeOH in CH_2Cl_2) to give the title product as a brown oil (0.61 g, 43.3% for two steps). m/z (ES+) 139 (M+H)⁺.



Compound 180.2. 5-Iodo-4-methyl-2-(oxetan-3-yl)-1H-imidazole. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **175.2**, except 4-methyl-2-(oxetan-3-yl)-1H-imidazole (compound **180.1**) was used in place of 4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazole (compound **175.1**). m/z (ES+) 265 (M+H)⁺.




Compound 180.3. Methyl 2,4-dimethyl-5-(4-methyl-2-(oxetan-3-yl)-1H-imidazol-5-yl)benzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.6**, except methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) and 5-iodo-4-methyl-2-(oxetan-3-yl)-1H-imidazole (compound **180.2**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**). m/z (ES+) 301 (M+H)⁺.




Compound 180.4. 2,4-Dimethyl-5-(4-methyl-2-(oxetan-3-yl)-1H-imidazol-5-yl)benzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except methyl 2,4-dimethyl-5-(4-methyl-2-(oxetan-3-yl)-1H-imidazol-5-yl)benzoate (compound **180.3**) was

used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **5.6**). m/z (ES⁺) 287 (M+H)⁺.




Compound 180. 4-(1-(2,4-Dimethyl-5-(4-methyl-2-(oxetan-3-yl)-1H-imidazol-5-

yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 2,4-dimethyl-5-(4-methyl-2-(oxetan-3-yl)-1H-imidazol-5-yl)benzoic acid (compound **180.4**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). m/z (ES⁺) 427 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆): δ 11.81 (br, 1H), 7.84 (d, J =8.4Hz, 2H), 7.60 (d, J =8.4Hz, 2H), 7.26-7.10 (m, 2H), 4.87-4.75 (m, 4H), 4.51-4.42 (m, 1H), 4.37-4.21 (m, 2H), 4.07-3.91 (m, 3H), 2.33 (s, 3H), 2.27 and 2.19 (2 singlets, CH₃, 3H), 2.12 and 2.02 (2 singlets, CH₃, 3H).




Compound 181. 4-(1-(2,4-Dimethyl-5-(4-methyl-2-(oxetan-3-yl)-1H-imidazol-5-


yl)benzoyl)-3-fluoroazetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 2,4-dimethyl-5-(4-methyl-2-(oxetan-3-yl)-1H-imidazol-5-yl)benzoic acid (compound **180.4**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 445 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆): δ 11.80 (br, 1H), 7.96 (d, J =8.4Hz, 2H), 7.77 (d, J =8.0Hz, 2H), 7.24 (s, 1H), 7.18 (d, J =7.2Hz, 2H), 4.87-4.75 (m, 4H), 4.60-4.36 (m, 4H), 4.31-4.21 (m, 1H), 2.35 (s, CH₃, 3H), 2.90 and 2.20 (2 singlets, CH₃, 3H), 2.13 and 2.02 (2 singlets, CH₃, 3H).



Compound 182.1. tert-Butyl 4-(5-iodo-4-methyl-1H-imidazol-2-yl)piperidine-1-carboxylate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159.2**, except tert-butyl 4-formylpiperidine-1-carboxylate was used in place of isobutyraldehyde. m/z (ES⁺) 392 (M+H)⁺.




Compound 182.2. 3-(2-(1-(tert-butoxycarbonyl)piperidin-4-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except tert-butyl 4-(5-iodo-4-methyl-1H-imidazol-2-yl)piperidine-1-carboxylate (compound **182.1**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**).




Compound 182.3. tert-Butyl 4-(4-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-5-methyl-1H-imidazol-2-yl)piperidine-1-carboxylate. A mixture of 3-(2-(1-(tert-butoxycarbonyl)piperidin-4-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **182.2**, 506 mg), 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**, 296 mg, 1.52 mmol), EDCI (315 mg, 1.65 mmol), HOBt (107 mg, 0.64 mmol) and DIEA (877 μ L, 5.08 mmol) in DMF (25 mL) was stirred at room temperature for 16 hours. The reaction was concentrated under reduced pressure, diluted with saturated NaHCO₃ (20 mL) and extracted with EtOAc (50 mL). The organic phase was washed with brine (3 X 10 mL), dried over MgSO₄ and concentrated under reduced pressure. The residue was purified by preparative TLC with 8% methanol in dichloromethane to give 476 mg (69%) of the title

compound as a foam. m/z (ES+) 540 (M+H)⁺.




Compound 182. 4-(1-(4-Methyl-3-(4-methyl-2-(1-methylpiperidin-4-yl)-1H-imidazol-5-yl) benzoyl) azetidin-3-yl)benzonitrile. To a solution of tert-butyl 4-(4-(5-(3-(4-cyanophenyl) azetidine-1-carbonyl)-2-methylphenyl)-5-methyl-1H-imidazol-2-yl)piperidine-1-carboxylate (compound **182.3**, 475 mg, 0.88 mmol) in 20 mL CH₂Cl₂, was added trifluoroacetic acid (4 mL). The reaction stirred for 1.5 hours and concentrated under reduced pressure, neutralized with 10 mL Na₂CO₃ (1M) and lyophilized to afford 4-(1-(4-methyl-3-(4-methyl-2-(piperidin-4-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The mixture of 4-(1-(4-methyl-3-(4-methyl-2-(piperidin-4-yl)-1H-imidazol-5-yl) benzoyl)azetidin-3-yl)benzonitrile (135 mg, 0.31 mmol), formic acid (300 μ L, 6.76 mmol), formaldehyde (37% in water, 300 μ L, 3.41 mmol) and water (1.5 mL) was heated to reflux for 10 hours. The reaction was diluted with saturated NaHCO₃ until pH 9 and extracted with CH₂Cl₂ (3 X 20 mL). The residue was purified by preparative TLC with 10 % methanol and 1% ammonium hydroxide in dichloromethane and lyophilized to give 43 mg (30% two steps) of the title compound as a white solid. m/z (ES+) 540 (M+H)⁺.




Compound 183. 4-(1-(3-(2-(1-Acetyl-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl) azetidin-3-yl)benzonitrile. To a solution of tert-butyl 4-(4-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-5-methyl-1H-imidazol-2-yl)piperidine-1-carboxylate (compound **182.3**, 475 mg, 0.88 mmol) in 20 mL DCM, was added trifluoroacetic acid (4 mL). The reaction was stirred for 1.5 hours, concentrated under reduced pressure and neutralized with 10 mL Na₂CO₃ (1M) to obtain a crude product after lyophilization. A mixture of 4-(1-(4-methyl-3-(4-methyl-2-(piperidin-4-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile (252 mg, 0.57 mmol), acetic anhydride (71 μ L, 0.75 mmol) and triethylamine (120 μ L, 0.86 mmol) was stirred at room temperature for 16 hours.


The reaction was concentrated under reduced pressure, then diluted with saturated NaH_2PO_4 (20 mL) and extracted with EtOAc (3 X 20 mL). The organic phase was dried over MgSO_4 and concentrated under reduced pressure. The residue was purified by preparative TLC with 10 % methanol in dichloromethane and lyophilized to give 235 mg (80%-2 steps) of the title compound as a white solid. m/z (ES+) 482 (M+H)⁺.




Compound 184. Methyl 4-(5-(5-(3-(4-cyanophenyl) azetidine-1-carbonyl)-2-methylphenyl)-4-methyl-1H-imidazol-2-yl) piperidine-1-carboxylate. To a solution of tert-butyl 4-(4-(5-(3-(4-cyanophenyl) azetidine-1-carbonyl)-2-methylphenyl)-5-methyl-1H-imidazol-2-yl) piperidine-1-carboxylate (compound **182.3**, 227 mg, 0.42 mmol) in CH_2Cl_2 (10 mL), was added trifluoroacetic acid (2 mL). The reaction was stirred for 1.5 hours and concentrated under reduced pressure. The reaction mixture was neutralized with 2 mL Na_2CO_3 (1M) to afford the crude intermediate after lyophilization. A mixture of 4-(1-(4-methyl-3-(4-methyl-2-(piperidin-4-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile (185 mg, 0.42 mmol) and diisopropyl ethylamine (363 μL , 2.1 mmol) were dissolved in DMF (6 mL). Methyl chloroformate (36 μL , 0.46 mmol) was added and the mixture was stirred at room temperature for 1 hour. The reaction was concentrated, diluted with saturated NaH_2PO_4 (20 mL) and extracted with EtOAc (3 X 20 mL). The organic phase was washed by brine, dried over MgSO_4 and concentrated under reduced pressure. The residue was purified by preparative TLC with 10 % methanol in dichloromethane and lyophilized to give 141 mg (67%, over 2 steps) of the title compound as a white solid. m/z (ES+) 498 (M+H)⁺.




Compound 185.1. tert-Butyl 4-(5-iodo-4-methyl-1H-imidazol-2-yl)-4-methylpiperidine-1-carboxylate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159.2**, except tert-butyl 4-formyl-4-methylpiperidine-1-carboxylate was used in place of isobutyraldehyde. m/z (ES+) 406 (M+H)⁺.



Compound 185.2. 3-(2-(1-(tert-butoxycarbonyl)-4-methylpiperidin-4-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except tert-butyl 4-(5-iodo-4-methyl-1H-imidazol-2-yl)-4-methylpiperidine-1-carboxylate (compound **185.1**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**).




Compound 185.3. tert-Butyl 4-(5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-4-methyl-1H-imidazol-2-yl)-4-methylpiperidine-1-carboxylate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **182.3**, except 3-(2-(1-(tert-butoxycarbonyl)-4-methylpiperidin-4-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **185.2**) was used in place of 3-(2-(1-(tert-butoxycarbonyl)piperidin-4-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **182.2**).




Compound 185. Methyl 4-(5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-4-methyl-1H-imidazol-2-yl)-4-methylpiperidine-1-carboxylate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **184**, except tert-butyl 4-(5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-4-methyl-1H-imidazol-2-yl)-4-methylpiperidine-1-carboxylate (compound **185.3**) was used in place of tert-butyl 4-(4-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-5-methyl-1H-imidazol-2-yl)

piperidine-1-carboxylate (compound **182.3**). m/z (ES+) 512 $M+H^+$.




Compound 186. Methyl 4-(5-(5-(3-(4-cyanophenyl)-3-fluoroazetidine-1-carbonyl)-2-methylphenyl)-4-methyl-1H-imidazol-2-yl)-4-methylpiperidine-1-

carboxylate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **184**, except tert-butyl 4-(5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-4-methyl-1H-imidazol-2-yl)-4-methylpiperidine-1-carboxylate (compound **185.3**) was used in place of tert-butyl 4-(4-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-5-methyl-1H-imidazol-2-yl)piperidine-1-carboxylate (compound **182.3**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 530 $(M+H)^+$.




Compound 187. 4-(1-(4-Methyl-3-(4-methyl-2-((tetrahydro-2H-pyran-4-

yl)methyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159**, except 2-(tetrahydro-2H-pyran-4-yl)acetaldehyde was used in place of isobutyraldehyde and methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used in place of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**). m/z (ES+) 455 $(M+H)^+$.



Compound 188.1. Ethyl 4-methyltetrahydro-2H-pyran-4-carboxylate. Into a 500-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of ethyl tetrahydro-2H-pyran-4-carboxylate (8 g, 50.6 mmol) in tetrahydrofuran (100 mL). This was followed by the addition of LDA (50 mL, 101.1 mmol, 2M in THF) dropwise at -78 °C and stirred for 3h. To this was added a solution of CH₃I (9.5 mL, 151.9 mmol) in tetrahydrofuran (50 mL) dropwise at -78 °C. The reaction mixture was stirred for 3 h at -78 °C, then carefully quenched with 400 mL of NH₄Cl (sat.). The aqueous phase was extracted with 300 mL of ethyl acetate and the combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 8 g (92%) of the title compound as a yellow oil.




Compound 188.2. (4-Methyltetrahydro-2H-pyran-4-yl)methanol. Into a 100-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of ethyl 4-methyltetrahydro-2H-pyran-4-carboxylate (compound **188.1**, 500 mg, 2.90 mmol) in tetrahydrofuran (12 mL). This was followed by the addition of lithium aluminum hydride (221 mg, 5.82 mmol) in portions at 0 °C. The reaction mixture was stirred for 1 h at room temperature, then carefully quenched with 1.2 mL of H₂O, 1.2 mL of NaOH (15%), 3.5 mL of H₂O. The solids were removed by filtration and the filtrate was concentrated under reduced pressure. This resulted in 300 mg (crude) of the title compound as a yellow oil.




Compound 188.3. 4-Methyltetrahydro-2H-pyran-4-carbaldehyde. Into a 100-mL 3-neck round-bottom flask, was placed a solution of (4-methyltetrahydro-2H-pyran-4-yl)methanol (compound **188.2**, 300 mg, 2.30 mmol) in dichloromethane (15 mL). Dess-Martin reagent (1.17 g, 2.76 mmol) was added to the reaction. The reaction mixture was stirred for 2 h at room temperature, then quenched with 15 mL of water. The aqueous phase was extracted with 20 mL of dichloromethane. The combined organic layers were dried over


anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 200 mg (68%) of the title compound as a yellow oil.



Compound 188. 4-(1-(4-Methyl-3-(4-methyl-2-(4-methyltetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159**, except 4-methyltetrahydro-2H-pyran-4-carbaldehyde (compound **188.3**) was used in place of isobutyraldehyde and methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used in place of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**). m/z (ES⁺) 455 (M+H)⁺.




Compound 189.1. tert-Butyl 3-(5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-4-methyl-1H-imidazol-2-yl)azetidine-1-carboxylate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **182.3**, except tert-butyl 3-formylazetidine-1-carboxylate was used in place of tert-butyl 4-formylpiperidine-1-carboxylate.




Compound 189.2. 4-(1-(3-(2-(Azetidin-3-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. Into a 250-mL round-bottom flask, was placed a solution of tert-butyl 3-(5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-4-methyl-1H-imidazol-2-yl)azetidine-1-carboxylate (compound **189.1**, 2 g, 3.91 mmol) in 1,4-dioxane (20 mL). Hydrogen chloride (4 M) (10 mL) was added to the reaction. The reaction

mixture was stirred for 2h at 30 °C, then concentrated under reduced pressure. The pH of the solution was adjusted to 7-8 with sodium bicarbonate (sat.). The resulting mixture was concentrated under reduced pressure to give 800 mg (crude) of the title compound as a white solid.




Compound 189. 4-(1-(3-(2-(1-Acetylazetidin-3-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. Into a 25-mL round-bottom flask, was placed a solution of 4-(1-(3-(2-(azetidin-3-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile (compound **189.2**, 80 mg, 0.19 mmol) in *N,N*-dimethylformamide (1 mL).

Triethylamine (27 μ L, 0.20 mmol) and acetic anhydride (19 μ L, 0.20 mmol) were added to the reaction. The reaction mixture was stirred for 2h at room temperature, then quenched with 10 mL of water. The aqueous phase was extracted with 3 x 3 mL of ethyl acetate and the combined organic extracts were concentrated under reduced pressure. The crude product (100 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-020): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150 mm,; mobile phase, Water with 50 mmol NH_4HCO_3 and MeCN (28% MeCN up to 40% in 7 min, up to 100% in 2 min, down to 28% in 1 min); Detector, Waters 2489 255 and 220 nm. This resulted in 32.8 mg (37%) of the title compound as a white solid. m/z (ES⁺) 454 (M+H)⁺.




Compound 190. Methyl 3-(5-(5-(3-(4-cyanophenyl)azetidine-1-carboxylate)-2-methylphenyl)-4-methyl-1H-imidazol-2-yl)azetidine-1-carboxylate. Into a 25-mL round-bottom flask, was placed a solution of 4-(1-(3-(2-(azetidin-3-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile (compound **189.2**, 100 mg, 0.24 mmol) in *N,N*-dimethylformamide (2 mL). Methyl chloroformate (33 μ L, 0.42 mmol) and triethylamine (34 μ L, 0.25 mmol) were added to the reaction. The reaction mixture was stirred for 2h at room temperature, then was quenched with 10 mL of water. The aqueous phase was extracted with

3 x 5 mL of ethyl acetate. The combined organic layers were washed with 3 x 3 mL of brine and concentrated under reduced pressure. The residue was purified by silica gel chromatography with dichloromethane/methanol (10:1). The crude product (100 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-020): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150 mm; mobile phase, Water with 50 mmol NH_4HCO_3 and MeCN (33.0% MeCN up to 45.0% in 7 min, up to 100.0% in 2 min, down to 33.0% in 1 min); Detector, Waters 2489, 254 and 220 nm. This resulted in 21.5 mg (19%) of the title compound as a white solid. m/z (ES+) 470 (M+H)⁺.




Compound 191. 4-(1-(4-Methyl-3-(4-methyl-2-(1-methylazetidin-3-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. Into a 50-mL round-bottom flask, was placed a solution of 4-(1-(3-(2-(azetidin-3-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile (compound **189.2**, 50 mg, 0.12 mmol) in tetrahydrofuran (1 mL). Formaldehyde (65 μ L, 37% wt, 0.57 mmol) and NaBH_3CN (30 mg, 0.48 mmol) were added to the reaction. The reaction mixture was stirred for 1h at room temperature. The reaction mixture was extracted with 3 x 3 mL of ethyl acetate and the combined organic extracts were washed with 3 x 1 mL of brine and concentrated under reduced pressure. The crude product (20 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-020): Column, XBridge Prep C18 OBD Column, 5 μ m, 19*150 mm; mobile phase, WATER WITH 0.03% $\text{NH}_3\text{H}_2\text{O}$ and MeCN (26.0% MeCN up to 40.0% in 7 min, up to 100.0% in 2 min, down to 26.0% in 1 min); Detector, Waters 2489, 254 and 220 nm. This resulted in 1.8 mg (3%) of the title compound as a white solid. m/z (ES+) 426 (M+H)⁺.




Compound 192.1. 4-Methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazole. Into a 1-L three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 4-methyl-1H-imidazole (10 g, 121.8 mmol)

in tetrahydrofuran (200 mL). This was followed by the addition of sodium hydride (7.32 g, 182.7 mmol, 60%) in several batches at 0 °C and stirred for 1h at room temperature. To this was added SEM-Cl (30.5 g, 199.7 mmol) at 0 °C. The mixture was stirred for 1h at room temperature, then carefully quenched with 50 mL of brine. The aqueous phase was extracted with 1 x 800 mL of ethyl acetate. The organic layer was washed with 1 x 300 mL of brine, 2 x 300 mL of sodium bicarbonate (sat.), dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:1) ethyl acetate as eluent to furnish 9 g (35%) of the title compound as a yellow oil.




Compound 192.2. 3-(4-Methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazol-2-yl)oxetan-3-ol. Into a 1000-mL three-neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazole (compound **192.1**, 9 g, 42.38 mmol) in tetrahydrofuran (200 mL). This was followed by the addition of n-BuLi (34 mL, 2.5 M in THF) dropwise at -78 °C and stirred for 1h. To this was added a solution of oxetan-3-one (6.11 g, 84.79 mmol) in tetrahydrofuran (30 mL) dropwise at -78 °C. The reaction mixture was stirred for 1h at -78 °C, then quenched with 50 mL of NH₄Cl (sat.). The aqueous phase was extracted with 1 x 600 mL of ethyl acetate. The organic layer was washed with 3 x 300 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 13 g (crude) of the title compound as a light yellow solid.




Compound 192.3. 3-(5-Iodo-4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazol-2-yl)oxetan-3-ol. Into a 100-mL round-bottom flask, was placed a solution of 3-(4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazol-2-yl)oxetan-3-ol (compound **192.2**, 200 mg, 0.70 mmol) in ACN (10 mL). NIS (237.7 mg, 1.06 mmol) was added to the reaction. The reaction mixture was stirred overnight at room temperature, then concentrated under reduced pressure. The reaction mixture was diluted with 40 mL of EtOAc. The organic layer

was washed with 3 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (2:3) as eluent to furnish 100 mg (35%) of the title compound as an orange oil.




Compound 192.4. Methyl 3-(2-(3-hydroxyoxetan-3-yl)-4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazol-5-yl)-4-methylbenzoate. Into a 100-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of methyl 4-methyl-3-(tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**, 282.7 mg, 1.02 mmol) in dioxane (12 mL). A solution of K_3PO_4 (904.9 mg, 4.26 mmol) in water (1.2 mL), 3-(5-iodo-4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazol-2-yl)oxetan-3-ol (compound **192.3**, 350 mg, 0.85 mmol) and $Pd(dppf)_2Cl_2$ (62.4 mg, 0.09 mmol) were added to the reaction. The mixture was stirred for 7h. The reaction mixture was cooled, then diluted with 60 mL of EtOAc. The solids were removed by filtration, the filtrate was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (2:3) eluent to furnish 220 mg (60%) of the title compound as an orange oil.




Compound 192.5. Methyl 3-(2-(3-hydroxyoxetan-3-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoate. Into a 100-mL round-bottom flask, was placed methyl 3-(2-(3-hydroxyoxetan-3-yl)-4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazol-5-yl)-4-methylbenzoate (compound **192.4**, 180 mg, 0.42 mmol), Et_3SiH (1 mL) and trifluoroacetic acid (2 mL). The reaction mixture was stirred for 4h at room temperature. The pH of the solution was adjusted to 8 with sodium hydroxide (1 M). The reaction mixture was diluted with 100 mL of brine. The aqueous phase was extracted with 2 x 30 mL of ethyl acetate, the

combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 150 mg (crude) of the title compound as an orange oil.




Compound 192. 4-(1-(3-(2-(3-Hydroxyoxetan-3-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except methyl 3-(2-(3-hydroxyoxetan-3-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **192.5**) was used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **5.6**). m/z (ES⁺) 429 (M+H)⁺.




Compound 193. 4-(1-(3-(2-(3-Fluorooxetan-3-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. Into a 50-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 4-(1-(3-(2-(3-hydroxyoxetan-3-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile (compound **192**, 200 mg, 0.47 mmol) in dichloromethane (5 mL). This was followed by the addition of a solution of DAST (76.2 μ L, 0.58 mmol) in dichloromethane (1 mL) dropwise at -78 °C. The resulting solution was stirred for 1h at room temperature, then quenched by the addition of 2 mL of sodium bicarbonate (sat.). The resulting solution was diluted with 60 mL of EtOAc and additional water. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate as eluent. The crude product (50 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-020): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150 mm; mobile phase, WATER WITH 0.05% TFA and MeCN (18.0% MeCN up to 28.0% in 9 min, up to 100.0% in 2 min, down to 18.0% in 1 min);

Detector, Waters 2489, 254 and 220 nm. This resulted in 15.1 mg (8%) of the title compound as a white solid. m/z (ES+) 431 (M+H)⁺.




Compound 194.1. 5-Iodo-2-(3-methoxyoxetan-3-yl)-4-methyl-1-((2-


5 **(trimethylsilyl)ethoxy)methyl)-1H-imidazole.** Into a 100-mL round-bottom flask, was placed a solution of 3-(5-iodo-4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazol-2-yl)oxetan-3-ol (compound **192.3**, 800 mg, 1.95 mmol) in tetrahydrofuran (30 mL). This was followed by the addition of sodium hydride (156.2 mg, 3.9 mmol, 60%) at 0 °C and stirred for 20 min at room temperature. To this was added CH₃I (554.1 mg, 3.90 mmol) at 0 °C. The
10 resulting solution was stirred for 5h at room temperature, then quenched with 30 mL of Na₂S₂O₃ (sat.). The aqueous phase was extracted with 3 x 40 mL of ethyl acetate, the combined organic layers were washed with 3 x 40 mL of brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:2) eluent to furnish 700 mg (85%) of
15 the title compound as colorless oil.




Compound 194. 4-(1-(3-(2-(3-Methoxyoxetan-3-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound
20 **192**, except 5-iodo-2-(3-methoxyoxetan-3-yl)-4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazole (compound **194.1**) was used in place of 3-(5-iodo-4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazol-2-yl)oxetan-3-ol (compound **192.3**). m/z (ES+) 443 (M+H)⁺.




Compound 195. 4-(1-(3-(2-(4-Hydroxytetrahydro-2H-pyran-4-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **192**, except dihydro-2H-pyran-4(3H)-one was used in place of oxetan-3-one. m/z (ES+) 457 (M+H)⁺.



Compound 196. 4-(1-(3-(2-(4-Methoxytetrahydro-2H-pyran-4-yl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **194**, except dihydro-2H-pyran-4(3H)-one was used in place of oxetan-3-one. m/z (ES+) 471 (M+H)⁺.




Compound 197.1. Pyrrolidine-1-carboximidamide hydrochloride. Into a 100-mL round-bottom flask, was placed a solution of pyrrolidine (5.8 mL, 70.30 mmol) in CH₃CN (30 mL). Triethylamine (9.8 mL, 70.17 mmol) and 1H-pyrazole-1-carboximidamide hydrochloride (10.2 g, 69.59 mmol) was added to the reaction. The reaction mixture was stirred overnight at 60 °C. The product was collected by filtration to yield 7.5 g (71%) of the title compound as a white solid.




Compound 197.2. Methyl 3-(2-bromopropanoyl)-4-methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to

those used for the preparation of compound **1.6**, except 4-methylbenzoic acid was used in place of 2,4-dimethylbenzoic acid.




Compound 197.3. Methyl 4-methyl-3-(4-methyl-2-(pyrrolidin-1-yl)-1H-imidazol-

5-yl)benzoate. Into a 50-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of methyl 3-(2-bromopropanoyl)-4-methylbenzoate (compound **197.2**, 500.0 mg, 1.75 mmol) in *N,N*-dimethylformamide (15 mL). Pyrrolidine-1-carboximidamide hydrochloride (compound **197.1**, 260.8 mg, 1.75 mmol) was treated with K_2CO_3 , then added to the reaction. The reaction mixture was stirred for 1 h at 50 °C. The reaction mixture was diluted with 30 mL of H_2O . The aqueous phase was extracted with 3 x 30 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with PE/EtOAc (1:1)~EtOAc/methanol (15:1) as eluent to furnish 230.0 mg (44%) of the title compound as a dark blue solid.




Compound 197. 4-(1-(4-Methyl-3-(4-methyl-2-(pyrrolidin-1-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **1**, except methyl 4-methyl-3-(4-methyl-2-(pyrrolidin-1-yl)-1H-imidazol-5-yl)benzoate (compound **197.3**) was used in place of methyl 5-(2,4-dimethyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **1.7**) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**). m/z (ES⁺) 426 (M+H)⁺.



Compound 198.1. 1-(4-Methyl-1H-imidazol-2-yl)ethanone. Into a 500-mL round-bottom flask, was placed an aqueous solution of 2-oxopropanal (25.2 mL, 222.0 mmol, 50%). Ammonium acetate (85 g, 1.10 mol) and acetic acid (200 mL) were added to the reaction.


- 5 The reaction mixture was stirred overnight at 100 °C, then concentrated under reduced pressure. The residue was diluted with 300 mL of EtOAc. The organic layer was washed with 4 x 30 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/hexane (1:1) as eluent to furnish 1.5 g (11%) of the title compound as a yellow solid.



10

Compound 198.2. Methyl 5-(2-acetyl-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.6**, except methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) and 1-(4-ethyl-1H-imidazol-2-yl)ethanone (compound **198.1**) was used in place of 2,4-dimethyl-1H-imidazole.


15




Compound 198.3. Methyl 5-(2-(1-hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate. Into a 100-mL round-bottom flask, was placed a solution of methyl 5-(2-acetyl-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **198.2**, 300 mg, 1.10 mmol) in methanol (30 mL). This was followed by the addition of NaBH₄ (84 mg, 2.22 mmol) in portions at 0 °C. The resulting solution was stirred for 1h at room temperature, then carefully quenched with 1 mL of aqueous hydrogen chloride (2 M). The resulting mixture was concentrated under reduced pressure. This resulted in 330 mg (crude) of the title compound as a white solid.

20


25




Compound 198. 4-(1-(5-(2-(1-Hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except methyl 5-(2-(1-hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **198.3**) was used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **5.6**). m/z (ES⁺) 415 (M+H)⁺.




Compound 199. 4-(3-Fluoro-1-(5-(2-(1-hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **198**, except 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 433 (M+H)⁺.




Compound 200.1. Methyl 3-(2-(1-hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **198.3**, except methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used in place of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**).




Compound 200. 4-(1-(3-(2-(1-Hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 198, except methyl 3-(2-(1-hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **200.1**) was used in place of methyl 5-(2-(1-hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **198.3**). m/z (ES⁺) 401 (M+H)⁺.



Compound 201. 4-(3-Fluoro-1-(3-(2-(1-hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **200**, except 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 419 (M+H)⁺.




Compound 202.1. 2-(3-(trifluoromethyl)-1H-imidazol-5-yl)-2-methyl-1,3-dioxolane. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16.2**, except 3-methyloxetane-3-carbaldehyde was used in place of acetaldehyde. m/z (ES⁺) 207(M+H)⁺.




Compound 202.2. 5-Iodo-2-(3-methyloxetan-3-yl)-4-(trifluoromethyl)-1H-imidazole. The title compound was prepared using standard chemical manipulations and

procedures similar to those used for the preparation of compound **175.2**, except 2-(3-methyloxetan-3-yl)-4-(trifluoromethyl)-1H-imidazole (compound **202.1**) was used in place of 4-methyl-2-(3-methyloxetan-3-yl)-1H-imidazole (compound **175.1**). m/z (ES+) 333 (M+H)⁺.




Compound 202.3. 5-Iodo-2-(3-methyloxetan-3-yl)-1H-imidazole-4-carbonitrile.


The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16.3**, except 5-iodo-2-(3-methyloxetan-3-yl)-4-(trifluoromethyl)-1H-imidazole (compound **202.2**) was used in place of 2-methyl-4-(trifluoromethyl)-1H-imidazole (compound **16.2**). m/z (ES+) 290 (M+H)⁺.




Compound 202.4. Methyl 3-(4-cyano-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)-4-methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.6**, except 5-iodo-2-(3-methyloxetan-3-yl)-1H-imidazole-4-carbonitrile (compound **202.3**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**). m/z (ES+) 312 (M+H)⁺.




Compound 202.5. 3-(4-Cyano-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)-4-methylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except methyl 3-(4-cyano-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)-4-methylbenzoate (compound **202.4**) was used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **5.6**). m/z (ES+) 298 (M+H)⁺.



Compound 202. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-2-(3-methyloxetan-3-yl)-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 3-(4-cyano-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **202.5**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). m/z (ES⁺) 438 (M+H)⁺. ¹H NMR (400 MHz, DMSO-*d*₆): δ 13.13 (br, 1H), 7.85 (d, J = 8.4Hz, 2H), 7.73-7.66 (m, 2H), 7.63 (d, J = 8.4Hz, 2H), 7.47 (d, J = 8.0Hz, 1H), 4.95 (d, J = 5.6Hz, 2H), 4.78-4.70 (m, 1H), 4.57-4.45 (m, 2H), 4.46 (d, J = 5.6Hz, 2H), 4.11-4.01 (m, 2H), 2.39 (s, 3H), 1.71 (s, 3H).




Compound 203.1. Methyl 5-(4-cyano-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)-2,4-dimethylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.6**, except methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) and 5-iodo-2-(3-methyloxetan-3-yl)-1H-imidazole-4-carbonitrile (compound **202.3**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**). m/z (ES⁺) 326 (M+H)⁺.



Compound 203.2. 5-(4-Cyano-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.7**, except methyl 5-(4-cyano-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)-2,4-dimethylbenzoate


(compound **203.1**) was used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **5.6**). m/z (ES⁺) 312 (M+H)⁺.



Compound 203. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2,4-


dimethylphenyl)-2-(3-methyloxetan-3-yl)-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 5-(4-cyano-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **203.2**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). m/z (ES⁺) 452 (M+H)⁺.

¹H NMR (400 MHz, DMSO-d₆): δ 13.01 (br, 1H), 7.84 (d, J = 8.4 Hz, 2H), 7.60 (d, J = 8.4 Hz, 2H), 7.38 (s, 1H), 7.24 (s, 1H), 4.94 (d, J = 5.2 Hz, 2H), 4.53-4.32 (m, 4H), 4.09-3.96 (m, 3H), 2.36 (s, 3H), 2.35 (s, 3H), 1.68 (s, 3H).




Compound 204. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-


2-cyclopropyl-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except cyclopropanecarbaldehyde was used in place of acetaldehyde and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**). m/z (ES⁺) 408 (M+H)⁺.



Compound 205. 5-(5-(4-(4-Cyanophenyl)piperidine-1-carbonyl)-2-methylphenyl)-2-cyclopropyl-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except cyclopropanecarbaldehyde was used in place of acetaldehyde. m/z (ES⁺) 436 (M+H)⁺.




Compound 206. 5-(5-(3-(4-Cyanophenyl)-3-fluoroazetidine-1-carbonyl)-2,4-dimethylphenyl)-2-cyclopropyl-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except cyclopropanecarbaldehyde was used in place of acetaldehyde and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES⁺) 440 (M+H)⁺.




Compound 207. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2,4-dimethylphenyl)-2-cyclopropyl-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except cyclopropanecarbaldehyde was used in place of acetaldehyde, 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place

of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**).
 m/z (ES+) 422 (M+H)⁺.




Compound 208. 5-(5-(3-(4-Cyanophenyl)-3-fluoroazetidine-1-carbonyl)-2,4-

5 dimethylphenyl)-2-methyl-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES+) 414 (M+H)⁺.



Compound 209. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2,4-


dimethylphenyl)-2-methyl-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES+) 396 (M+H)⁺.



Compound 210. 5-(5-(3-(4-Cyanophenyl)-3-fluoroazetidine-1-carbonyl)-2,4-dimethylphenyl)-2-isopropyl-1H-imidazole-4-carbonitrile. The title compound was


prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**), methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate, and isobutyraldehyde was used in place of acetaldehyde.

m/z (ES⁺) 442 (M+H)⁺.




Compound 211. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazole-4-carbonitrile. The title compound was


prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) and tetrahydro-2H-pyran-4-carbaldehyde was used in place of acetaldehyde. m/z (ES⁺) 452 (M+H)⁺.




Compound 212. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2,4-dimethylphenyl)-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**), tetrahydro-2H-pyran-4-carbaldehyde was used in place of acetaldehyde and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES+) 466 (M+H)⁺.




Compound 213. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-2-(tetrahydrofuran-2-yl)-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) and tetrahydrofuran-2-carbaldehyde was used in place of acetaldehyde. m/z (ES+) 438 (M+H)⁺.




Compound 214. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-2-(tetrahydrofuran-3-yl)-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) and tetrahydrofuran-3-carbaldehyde was used in place of acetaldehyde. m/z (ES+) 438 (M+H)⁺.



Compound 215. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-2-(4-methyltetrahydro-2H-pyran-4-yl)-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) and 4-methyltetrahydro-2H-pyran-4-carbaldehyde (compound **188.3**) was used in place of acetaldehyde. m/z (ES⁺) 466 (M+H)⁺.




Compound 216.1. 4-(Trifluoromethyl)-1H-imidazole. Into a 1000-mL round-bottom flask, was placed a solution of 3,3,3-trifluoro-2-oxopropanal (143 mL, 111.1 mmol) in a solvent mixture of methanol and water (200/200 mL). An aqueous solution of formaldehyde (350 mL, 116.67 mmol, 35%) and ammonium hydroxide (30 mL, 25%) were added to the reaction. The resulting solution was stirred for 2 h at room temperature, then concentrated under reduced pressure. The solids were collected by filtration to give 2 g (13%) of the title compound as a white solid.



Compound 216.2. 4-(Trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazole. Into a 250-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 4-(trifluoromethyl)-1H-imidazole (compound **216.1**, 5 g, 36.74 mmol) in tetrahydrofuran (100 mL). This was followed by the addition of sodium hydride (1.6 g, 40.00 mmol, 60%) in portions at 0 °C and stirred for 1h at 0 °C. To this was added SEMCl (7.1 mL, 40.36 mmol) dropwise at 0 °C. The

resulting solution was stirred for 4h at 0 °C, then carefully quenched with 100 mL of brine. The pH of the solution was adjusted to 7-8 with hydrogen chloride (1 M). The aqueous phase was extracted with 2 x 100 mL of ethyl acetate and the combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was


5 purified by silica gel chromatography with ethyl acetate/petroleum ether (1:5) as eluent to furnish 6 g (61%) of the title compound as a light yellow oil.



Compound 216.3. 4-(4-(Trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazol-2-yl)tetrahydro-2H-pyran-4-ol. Into a 100-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 4-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazole (compound **216.2**, 5 g, 18.77 mmol) in tetrahydrofuran (50 mL). This was followed by the addition of n-butyllithium (9 mL, 22.5 mmol, 2.5N in hexane) dropwise at -78 °C and stirred for 1h at -60 °C. To this was added dihydro-2H-pyran-4(3H)-one (6 g, 59.93 mmol). The resulting solution

15 was stirred for 4h at 0 °C in an ice/salt bath, then quenched with 10 mL of water. The resulting solution was diluted with 100 mL of NH₄Cl (sat.). The aqueous phase was extracted with 2 x 100 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:4) as eluent to furnish 6 g (87%) of


20 the title compound as a light yellow oil.




Compound 216.4. 4-(4-(Trifluoromethyl)-1H-imidazol-2-yl)tetrahydro-2H-pyran-4-ol. Into a 25-mL round-bottom flask, was placed 4-(4-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxymethyl)-1H-imidazol-2-yl)tetrahydro-2H-pyran-4-ol (compound **216.3**, 2 g, 5.46 mmol), Et₃SiH (2 mL) and trifluoroacetic acid (4 mL). The resulting solution was

25 stirred for 2h at room temperature. The pH of the solution was adjusted to 8 with sodium hydroxide (1 M). The resulting solution was diluted with 100 mL of brine. The aqueous


phase was extracted with 2 x 100 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 1.2 g (93%) of the title compound as a white solid.



Compound 216. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-2-(4-hydroxytetrahydro-2H-pyran-4-yl)-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except 4-(4-(trifluoromethyl)-1H-imidazol-2-yl)tetrahydro-2H-pyran-4-ol (compound **216.4**) was used in place of 2-methyl-4-(trifluoromethyl)-1H-imidazole (compound **16.2**) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**). m/z (ES⁺) 468 (M+H)⁺.




Compound 217.1. 2-(4-Methoxytetrahydro-2H-pyran-4-yl)-4-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazole. Into a 250-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 4-(4-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazol-2-yl)tetrahydro-2H-pyran-4-ol (compound **216.3**, 2 g, 5.46 mmol) in tetrahydrofuran (100 mL). This was followed by the addition of sodium hydride (262 mg, 6.55 mmol, 60%) at -70 °C and stirred for 30 min. To this was added MeI (930 mg, 6.55 mmol). The resulting solution was stirred for 1h at room temperature, then carefully quenched with 10 mL of water. The resulting mixture was diluted with 50 mL of brine. The aqueous phase was extracted with 2 x 50 mL of ethyl acetate. The combined organic layers were washed with 1 x 50 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:3) as eluent to furnish 1.6 g (77%) of the title compound as light brown oil.




Compound 217.2. 2-(4-Methoxytetrahydro-2H-pyran-4-yl)-4-(trifluoromethyl)-1H-imidazole. Into a 100-mL round-bottom flask, was placed 2-(4-methoxytetrahydro-2H-pyran-4-yl)-4-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazole

(compound **217.1**, 1.3 g, 3.42 mmol), trifluoroacetic acid (4 mL), Et₃SiH (2 mL). The resulting solution was stirred overnight at 20 °C, then quenched by the addition of 20 mL of water. The pH of the solution was adjusted to 8 with sodium hydroxide (1 M). The aqueous phase was extracted with 2 x 100 mL of ethyl acetate. The combined organic layers were washed with 2 x 50 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 1 g (crude) of the title compound as a light yellow oil.




Compound 217. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-2-(4-methoxytetrahydro-2H-pyran-4-yl)-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **16**, except 2-(4-methoxytetrahydro-2H-pyran-4-yl)-4-(trifluoromethyl)-1H-imidazole (compound **217.2**) was used in place of 2-methyl-4-(trifluoromethyl)-1H-imidazole (compound **16.2**) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**). *m/z* (ES⁺) 482 (M+H)⁺.




Compound 218.1. 5-Iodo-2-methyl-4-(trifluoromethyl)-1H-imidazole. Into a 50-mL round-bottom flask, was placed a solution of 2-methyl-4-(trifluoromethyl)-1H-imidazole (compound **16.2**, 1.72 g, 11.46 mmol) in CH₃CN (25 mL). NIS (3.87 g, 17.20 mmol) was added to the reaction. The reaction mixture was stirred overnight at 85 °C. The reaction


mixture diluted with 50 mL of H₂O and extracted with 3 x 30 mL of ethyl acetate. The combined organic layers were washed with 2 x 20 mL of Na₂S₂O₃(sat.) and 2 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 4.32 g (crude) of the title compound as a brown oil.



Compound 218. 4-(1-(4-Methyl-3-(2-methyl-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 16, except 5-iodo-2-methyl-4-(trifluoromethyl)-1H-imidazole (compound 218.1) was used in place of 2-methyl-1H-imidazole-4-carbonitrile (Compound 16.3). *m/z* (ES⁺) 425 (M+H)⁺.




Compound 219.1. 2-((Benzyloxy)methyl)-4-(trifluoromethyl)-1H-imidazole. Into a 100-mL round-bottom flask, was placed a solution of 3,3,3-trifluoro-2-oxopropanal (2 g, 15.87 mmol) in methanol (30 mL). 2-(benzyloxy)acetaldehyde (2.8 g, 18.64 mmol) and ammonium hydroxide (25%) (36 mL, 63.48 mmol) were added to the reaction. The reaction mixture was stirred for 15h at 20 °C, then concentrated under reduced pressure. The residue was diluted with 20 mL of H₂O. The aqueous phase was extracted with 2 x 30 mL of ethyl acetate and the combined organic layers were washed with 2 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 4.2 g (crude) of the title compound as a yellow crude oil.




Compound 219.2. 2-((Benzyloxy)methyl)-5-iodo-4-(trifluoromethyl)-1H-imidazole. Into a 50-mL round-bottom flask, was placed a solution of 2-((benzyloxy)methyl)-4-(trifluoromethyl)-1H-imidazole (compound 219.1, 1.5 g, 5.85 mmol)

in CH₃CN (18 mL). NIS (1.6 g, 7.02 mmol) was added to the reaction. The reaction mixture was stirred for 15 h at 85 °C. The reaction mixture was diluted with 30 mL of H₂O. The aqueous phase was extracted with 2 x 30 mL of ethyl acetate. The combined organic layers were washed with 2 x 20 mL of Na₂S₂O₃(sat.) and 2 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:100-1:2) as eluent to furnish 0.7 g (31%) of the title compound as a yellow oil.



Compound 219.3. Methyl 5-(2-((benzyloxy)methyl)-4-(trifluoromethyl)-1H-


imidazol-5-yl)-2,4-dimethylbenzoate. Into a 50-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 2-((benzyloxy)methyl)-5-iodo-4-(trifluoromethyl)-1H-imidazole (compound **219.2**, 500 mg, 1.31 mmol) in dioxane (8 mL). Methyl 2,4-dimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**, 450 mg, 1.55 mmol), Pd(dppf)Cl₂ (0.1 g) and an aqueous solution of potassium carbonate (2 M) (3.25 mL) were added to the reaction. The reaction mixture was stirred for 1 h at 80 °C, then diluted with 20 mL of H₂O. The aqueous phase was extracted with 2 x 30 mL of ethyl acetate. The combined organic layers were washed with 2 x 30 mL of brine and dried over anhydrous sodium sulfate. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/20-1/4) as eluent to yield 0.2 g (37%) of the title compound as a light yellow solid.




Compound 219.4. Methyl 5-(2-(hydroxymethyl)-4-(trifluoromethyl)-1H-

imidazol-5-yl)-2,4-dimethylbenzoate. Into a 50-mL round-bottom flask, under a nitrogen atmosphere, was placed a solution of methyl 5-(2-((benzyloxy)methyl)-4-(trifluoromethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **219.3**, 150 mg, 0.36 mmol) in methanol (8 mL). Palladium on carbon (150 mg, 1.00 equiv) and HCl (4 M, 2 mL) were added to the


reaction under a nitrogen atmosphere. To the above hydrogen (1 atm) was introduced. The reaction mixture was stirred for 2h at room temperature. The system was purged with nitrogen, then the solids were removed by filtration and the filtrate was concentrated under reduced pressure to give 100 mg (85%) of the title compound as a light yellow oil.




Compound 219.5. Methyl 5-(4-cyano-2-(hydroxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoate. Into a 50-mL round-bottom flask, was placed a solution of methyl 5-(2-((benzyloxy)methyl)-4-(trifluoromethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **219.4**, 100 mg, 0.30 mmol) in ammonium hydroxide (5%) (30 mL). The reaction mixture was stirred for 2h at 60 °C. The reaction mixture was extracted with 2 x 50 mL of dichloromethane, the organic layers combined, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 80 mg (92%) of the title compound as a light yellow oil.




Compound 219.6. 5-(4-Cyano-2-(hydroxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid. Into a 25-mL round-bottom flask, was placed a solution of methyl 5-(4-cyano-2-(hydroxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **219.6**, 100 mg, 0.35 mmol) in methanol (10 mL) and a solution of NaOH (0.14 g, 3.5 mmol) in water (5 mL). The reaction mixture was stirred for 15h at room temperature, then concentrated under reduced pressure. The pH of the solution was adjusted to 1-2 with hydrogen chloride (4 M). The resulting mixture was concentrated under reduced pressure. Methanol (5 mL) was added to the residue. The salt was filtered off, and the filtrate was concentrated under reduced pressure to give 200 mg (crude) of the title compound as a light yellow solid.




Compound 219. 5-(5-(3-(4-Cyanophenyl)-3-fluoroazetidine-1-carbonyl)-2,4-dimethylphenyl)-2-(hydroxymethyl)-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 5-(4-cyano-2-(hydroxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **219.6**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 430 (M+H)⁺.




Compound 220. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2,4-dimethylphenyl)-2-(hydroxymethyl)-1H-imidazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 5-(4-cyano-2-(hydroxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **219.6**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). m/z (ES⁺) 412 (M+H)⁺.




Compound 221.1. 3-(4-Cyano-2-(hydroxymethyl)-1H-imidazol-5-yl)-4-methylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **219.6**, except methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used in place of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**).




Compound 221. 4-(1-(4-Methyl-3-(4-methyl-2-(pyrrolidin-1-yl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 5, except 3-(4-cyano-2-(hydroxymethyl)-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **221.1**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). m/z (ES⁺) 398 (M+H)⁺.



Compound 222.1. 2-(Methoxymethyl)-4-methyl-4,5-dihydro-1H-imidazole. Into a 500-mL round-bottom flask, was placed a solution of 2-methoxyacetic acid (20 g, 222.0 mmol) in toluene (200 mL). Propane-1,2-diamine (50 g, 674.5 mmol) was added to the reaction. The reaction mixture was stirred overnight at 130 °C, then concentrated under reduced pressure. This resulted in 25 g (crude) of the title compound as a yellow oil.




Compound 222.2. 2-(Methoxymethyl)-4-methyl-1H-imidazole. Into a 500-mL round-bottom flask, was placed a solution of 2-(methoxymethyl)-4-methyl-4,5-dihydro-1H-imidazole (compound **222.1**, 19 g, 148.4 mmol) in acetonitrile (200 mL). Al₂O₃ (19 g, 182.7 mmol) was added to the reaction. This was followed by the addition of potassium permanganate (58 g, 367.1 mmol) in several batches at 0 °C. The reaction mixture was stirred for 2h at 0 °C, then warmed to room temperature overnight. The reaction was quenched with 20 mL of sodium sulfite (sat.). The solids were removed by filtration and the filtrate was concentrated under reduced pressure. This resulted in 19 g (crude) of the title compound as yellow oil.




Compound 222.3. 5-Iodo-2-(methoxymethyl)-4-methyl-1H-imidazole. Into a 100-mL round-bottom flask, was placed a solution of 2-(methoxymethyl)-4-methyl-1H-imidazole (compound **222.2**, 1.9 g, 15.06 mmol) in sodium hydroxide aqueous solution (30 mL, 2M).

5 This was followed by the addition of a solution of iodine (7.7 g, 30.34 mmol) in dichloromethane (30 mL). The reaction mixture was stirred for 1h at room temperature. The aqueous phase was collected and the pH was adjusted to 4 with hydrogen chloride (2M). The reaction mixture was extracted with 4 x 30 mL of dichloromethane and the organic layers combined. The resulting mixture was washed with 2 x 20 mL of sodium sulfite (sat.) and 3 x
10 30 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 1.2 g (32%) of the title compound as a yellow solid.




Compound 222.4. Methyl 5-(2-(methoxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate. Into a 50-mL three neck round-bottom flask, which was purged and

15 maintained with an inert atmosphere of nitrogen, was placed a solution of 5-iodo-2-(methoxymethyl)-4-methyl-1H-imidazole (compound **222.3**, 400 mg, 1.59 mmol) in 1,4-dioxane (15 mL). Methyl 2,4-dimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**, 510 mg, 1.76 mmol), a solution of potassium carbonate (662 mg, 4.79 mmol) in water (1 mL) and Pd(dppf)Cl₂ (234 mg, 0.32 mmol) were added to the reaction.
20 The reaction mixture was stirred overnight at 90 °C . The solids were removed by filtration and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography with dichloromethane/ethyl acetate (1:1) as eluent to furnish 220 mg (48%) of the title compound as a yellow solid.




Compound 222.5. 5-(2-(Hydroxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-

dimethylbenzoic acid. Into a 100-mL round-bottom flask, was placed methyl 5-(2-(methoxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **222.4**, 220 mg, 0.76 mmol) and HBr (20 mL, 40% in HOAc). The reaction mixture was stirred overnight at 80 °C. The resulting mixture was concentrated under reduced pressure. This resulted in 190 mg (crude) of the title compound as a brown solid.

**Compound 222.6. 5-(2-(Cyanomethyl)-4-methyl-1H-imidazol-5-yl)-2,4-**


dimethylbenzoic acid. Into a 100-mL round-bottom flask, was placed a solution of trimethylsilanecarbonitrile (376 mg, 3.79 mmol) in acetonitrile (20 mL). Tetrabutylammonium fluoride (3.8 mL, 1 M in THF) was added to the reaction. This was followed by the addition of 5-(2-(hydroxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **222.5**, 190 mg, 0.73 mmol), in portions. The reaction mixture was stirred for 1h at room temperature, then concentrated under reduced pressure.

The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:1) as eluent to furnish 190 mg (97%) of the title compound as a yellow solid.


**Compound 222. 4-(1-(5-(2-(Cyanomethyl)-4-methyl-1H-imidazol-5-yl)-2,4-**

dimethylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile. Into a 100-mL round-bottom flask, was placed a solution of 5-(2-(cyanomethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **222.6**, 190 mg, 0.71 mmol) and HBTU (538 mg, 1.42 mmol) in *N,N*-dimethylformamide (3 mL). To the above were added a solution of 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**, 160 mg, 0.75 mmol) and triethylamine (197 µL, 1.42 mmol) in *N,N*-dimethylformamide (3 mL) dropwise. The reaction mixture was stirred for 1h at room temperature. The crude product (150 mg) was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-001(SHIMADZU)): Column, XBridge Prep C18 OBD Column, 5 µm, 19*150 mm; mobile phase, Water with 50


mmol NH_4HCO_3 and acetonitrile (30% acetonitrile up to 44% in 7 min, hold 44% in 1 min, up to 100% in 1 min, down to 30% in 1 min); Detector, Waters 2489, 254 and 220 nm. This resulted in 45.9 mg (15%) of the title compound as a white solid. m/z (ES⁺) 428 (M+H)⁺.




5 **Compound 223. 4-(1-(3-(2-(Cyanomethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 222, except methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound 5.4) was used in place of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound 160.1) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound 5.2) was used in place of 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound 43.4). m/z (ES⁺) 396 (M+H)⁺.




15 **Compound 224. 4-(1-(5-(2-(Hydroxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 222, except 5-(2-(hydroxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound 222.5) was used in place of 5-(2-(cyanomethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound 222.6) and 4-(azetidin-3-yl)benzonitrile hydrochloride (compound 5.2) was used in place of 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound 43.4). m/z (ES⁺) 401 (M+H)⁺.




Compound 225. 4-(3-Fluoro-1-(5-(2-(hydroxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **222**, except 5-(2-(hydroxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **222.5**) was used in place of 5-(2-(cyanomethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **222.6**). m/z (ES+) 419 (M+H)⁺.



Compound 226.1. 3-(2-(Hydroxymethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of 5-(2-(hydroxymethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **222.5**), except methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used in place of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**).




Compound 226. 4-(1-(3-(2-(Hydroxymethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **222**, except 3-(2-(hydroxymethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **226.1**) was used in place of 5-(2-(cyanomethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **222.6**). m/z (ES+) 387 (M+H)⁺.




Compound 227.1. 3-(Benzyloxy)propanal. Into a 100-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed DMSO (5.1 mL, 71.79 mmol) in dichloromethane (40 mL). This was followed by the addition of oxalyl chloride (3.1 mL, 54.16 mmol) dropwise at -78 °C and stirred for 30 min at


-78 °C. To this was added a solution of 3-(benzyloxy)propan-1-ol (4.8 mL, 30.08 mmol) in dichloromethane (10 mL) dropwise at -78 °C. The resulting solution was stirred for 1 h at -78 °C, then triethylamine (16.5 mL, 118.59 mmol) was added to the reaction. The resulting solution was stirred for 1 h at -78 to -20 °C, then quenched with 50 mL of NH₄Cl (sat.). The aqueous phase was extracted with 2 x 50 mL of dichloromethane. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with EtOAc:PE (1:5) as eluent to furnish 2.0 g (40%) of the title compound as light yellow oil.



Compound 227.2. 2-(2-(Benzyloxy)ethyl)-5-iodo-4-methyl-1H-imidazole. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **160.2**, except 3-(benzyloxy)propanal (compound **227.1**) was used in place of cyclopropanecarbaldehyde.




Compound 227.3. Methyl 3-(2-(2-(benzyloxy)ethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.6**, except 2-(2-(benzyloxy)ethyl)-5-iodo-4-methyl-1H-imidazole (compound **227.2**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**).




Compound 227.4. Methyl 3-(2-(2-hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoate. Into a 100-mL round-bottom flask, was placed a solution of methyl 3-(2-(2-(benzyloxy)ethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **227.3**, 100 mg, 0.27 mmol) in methanol (10 mL). Palladium on carbon (100 mg), hydrogen chloride (4M) (2.5 mL) were added to the reaction under N₂. To the above hydrogen was introduced. The reaction mixture was stirred for 3h at room temperature. The solids were filtered off and the


filtrate was concentrated under reduced pressure. The pH of the solution was adjusted to 7 with sodium bicarbonate (sat.). The resulting solution was extracted with 4 x 20 mL of ethyl acetate. The organic layers were combined, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 60 mg (80%) of the title compound as a yellow oil.




Compound 227. 4-(1-(3-(2-(2-Hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except methyl 3-(2-(2-hydroxyethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **227.4**) was used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **5.6**). m/z (ES⁺) 401 (M+H)⁺.




Compound 228. 5-(5-(3-(4-Cyanophenyl)-3-fluoroazetidine-1-carbonyl)-2,4-dimethylphenyl)-4-methyl-1H-imidazole-2-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **14**, except 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES⁺) 414 (M+H)⁺.




Compound 229. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2,4-dimethylphenyl)-4-methyl-1H-imidazole-2-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **14**, except methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES⁺) 396 (M+H)⁺.



Compound 230.1. 2-(Methoxymethyl)-4,5-dihydro-1H-imidazole. Into a 250-mL round-bottom flask, was placed a solution of 2-methoxyacetic acid (20 g, 222.03 mmol) in toluene (60 mL). Ethane-1,2-diamine (133g, 2.22 mol) was added to the reaction. The reaction mixture was stirred for 2 days at 130 °C. The resulting mixture was concentrated under reduced pressure. This resulted in 20 g (79%) of the title compound as yellow crude oil.




Compound 230.2. 2-(Methoxymethyl)-1H-imidazole. Into a 500-mL 3-neck round-bottom flask, was placed a solution of 2-(methoxymethyl)-4,5-dihydro-1H-imidazole (compound **230.1**, 20 g, 175.21 mmol) in ACN (150 mL). Al₂O₃ (9 g, 87.6 mmol), KMnO₄ (27.7 g, 175.21 mmol) were added to the reaction. The reaction mixture was stirred for 6h at room temperature, then quenched with 30 mL of Na₂SO₃(sat). The solids were removed by filtration and the filtrate was concentrated under reduced pressure. This resulted in 15 g (crude) of the title compound as a yellow solid.




Compound 230.3. 4,5-Diiodo-2-(methoxymethyl)-1H-imidazole. Into a 500-mL round-bottom flask, was placed a solution of 2-(methoxymethyl)-1H-imidazole (compound **230.2**, 15 g, 133.77 mmol) in sodium hydroxide aqueous solution (100 mL, 2M). A solution of I₂ (60 g, 236.22 mmol) in dichloromethane (100 mL) was added to the reaction. The reaction mixture was stirred for 3h at 25°C. The aqueous phase was collected and diluted with 50 mL of Na₂SO₃(aq). The pH of the solution was adjusted to 6-7 with hydrogen chloride


(2M). The reaction mixture was extracted with 3 x 50 mL of dichloromethane. The aqueous layers were combined and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/hexane (1:2) as eluent to furnish 4 g (8%) of the title compound as a white solid.




Compound 230.4. 5-Iodo-2-(methoxymethyl)-1H-imidazole. Into a 250-mL round-bottom flask, was placed a solution of 4,5-diiodo-2-(methoxymethyl)-1H-imidazole (compound **230.3**, 2 g, 5.50 mmol) in ethanol (40 mL). A solution of Na_2SO_3 (5.9 g, 46.83 mmol) in water (80 mL) was added to the reaction. The reaction mixture was stirred overnight at 90 °C. The resulting mixture was concentrated under reduced pressure. The residue was extracted with 3 x 20 mL of dichloromethane and the organic layers combined and concentrated under reduced pressure. This resulted in 1 g (76%) of the title compounds as yellow oil.




Compound 230.5. Methyl 5-(2-(methoxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoate. Into a 100-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 5-iodo-2-(methoxymethyl)-1H-imidazole (compound **230.4**, 70 mg, 0.29 mmol) in dioxane (10 mL). Pd(dppf)Cl_2 (22 mg, 0.029 mmol), a solution of potassium carbonate (160 mg, 1.16 mmol) in water (2 mL), methyl 2,4-dimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**, 168 mg, 0.58 mmol) were added to the reaction. The reaction mixture was stirred overnight at 90 °C, then quenched with 10 mL of H_2O . The aqueous phase was extracted with 3 x 10 mL of ethyl acetate and the combined organic layers were washed with 1 x 10 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:1) as eluent to furnish 70 mg (87%) of the title compound as a white solid.



Compound 230.6. Methyl 5-(4-chloro-2-(methoxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoate. Into a 100-mL round-bottom flask, was placed a solution of methyl 5-(2-(methoxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **230.5**, 70 mg, 0.26 mmol) in chloroform (6 mL). NCS (35 mg, 0.26 mmol) was added to the reaction. The reaction mixture was stirred overnight at room temperature, then quenched with 1 mL of water. The aqueous phase was extracted with 2 x 5 mL of dichloromethane and the combined organic layers were concentrated under reduced pressure. The residue was applied onto a silica gel chromatography with ethyl acetate/petroleum ether (1:10) as eluent to yield 30 mg (38%) of the title compound as a white crude solid.




Compound 230.7. 5-(4-Chloro-2-(hydroxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid. Into a 50-mL round-bottom flask, was placed a solution of methyl 5-(4-chloro-2-(methoxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **230.5**, 20 mg, 0.06 mmol) in HBr (5 mL, 40% in AcOH). The reaction mixture was stirred overnight at 90°C. The resulting mixture was concentrated under reduced pressure. This resulted in 10 mg (55%) of the title compound as a yellow crude solid.




Compound 230. 4-(1-(5-(4-Chloro-2-(hydroxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile. Into a 50-mL round-bottom flask, was placed a solution of 5-(4-chloro-2-(hydroxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoic acid (compound **230.6**, 180 mg, 0.64 mmol) in *N,N*-dimethylformamide (15 mL). 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**, 165 mg, 0.78 mmol), EDC·HCl (245 mg, 1.28 mmol) and 4-dimethylaminopyridine (310 mg, 2.54 mmol)


were added to the reaction. The reaction mixture was stirred overnight at 25°C. The reaction was then quenched with 5 mL of water. The reaction mixture was extracted with 3 x 10 mL of ethyl acetate and the organic layers combined. The resulting mixture was washed with 5 x 10 mL of brine. The resulting mixture was concentrated under reduced pressure. The crude product (200 mg) was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-001(SHIMADZU)): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150 mm; mobile phase, WATER WITH 0.05% TFA and ACN (26.0% ACN up to 39.0% in 10 min, up to 100.0% in 2 min, down to 26.0% in 1 min); Detector, Waters 2489, 254 and 220 nm. This resulted in 84 mg (30%) of the title compound as a white solid. m/z (ES+) 439 (M+H)⁺.




Compound 231. 4-(1-(5-(4-Chloro-2-(hydroxymethyl)-1H-imidazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **230**, except 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**). m/z (ES+) 421 (M+H)⁺.




Compound 232.1. Ethyl 4-oxobutanoate. Into a 500-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of ethyl 4-chloro-4-oxobutanoate (12 g, 72.91 mmol) in tetrahydrofuran (200 mL). The system was purged with nitrogen and palladium on carbon (2 g) was added to the reaction. This was followed by the addition of triethylsilane (16.5 mL, 103.20 mmol) dropwise with stirring at room temperature. The reaction mixture was stirred for 2h at room temperature. The solids were filtered off and the filtrate was concentrated under reduced pressure to give 18 g (crude) of the title compound as a light yellow solid.




Compound 232.2. Methyl 3-(4-methyl-1H-imidazol-2-yl)propanoate. Into a 500-mL round-bottom flask, was placed a solution of ethyl 4-oxobutanoate (compound **232.1**, 5.4 g, 41.49 mmol) in methanol (125 mL). An aqueous solution of 2-oxopropanal (35%) (28.5 mL, 49.96 mmol) and ammonium hydroxide (25%) (100 mL, 165.9 mmol) were added to the reaction. The reaction mixture was stirred for 2 h at 50 °C. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate as eluent to furnish 600 mg (8%) of the title compound as light yellow oil.



Compound 232.3. 3-(4-Methyl-1H-imidazol-2-yl)propanamide. Into a 100-mL round-bottom flask, was placed a solution of ethyl 3-(4-methyl-1H-imidazol-2-yl)propanoate (compound **232.2**, 500 mg, 2.97 mmol) in methanol (10 mL). To the above $\text{NH}_3(\text{g})$ was bubbled through the solution. The reaction mixture was stirred for overnight at 50 °C. The resulting mixture was concentrated under reduced pressure. This resulted in 346 mg (82%) of the title compound as a light yellow solid.



Compound 232.4. 3-(4-Methyl-1H-imidazol-2-yl)propanenitrile. Into a 250-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, were placed a solution of 3-(4-methyl-1H-imidazol-2-yl)propanamide (compound **232.3**, 2 g, 13.06 mmol) in toluene (100 mL) and P_2O_5 (2 g, 14.08 mmol). The reaction mixture was stirred for 12h at 120 °C, then quenched with 100 mL of water. The aqueous phase was extracted with 2 x 200 mL of ethyl acetate and the combined organic layers were washed with 2 x 100 mL of sodium carbonate (sat.), dried over sodium sulfate and concentrated under reduced pressure. This resulted in 1 g (57%) of 3-(4-methyl-1H-imidazol-2-yl)propanenitrile as light yellow oil







Compound 232. 4-(1-(5-(2-(2-Cyanoethyl)-4-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 3-(4-methyl-1H-imidazol-2-yl)propanenitrile (compound **232.4**) was used in place of 2,4-dimethyl-1H-imidazole, methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**), and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 442 (M+H)⁺.

The compounds in **TABLE 11** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **222**, **223**, **231**, and **232**.


TABLE 11

Cpd	Name	Structure	m/z (ES ⁺) (M+H) ⁺
93	4-(1-(3-(4-chloro-2-(cyanomethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile		444


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
143	4-(1-(3-(4-chloro-2-(2-cyanoethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		430
144	4-(1-(3-(4-chloro-2-(2-cyanoethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile		458
149	4-(1-(3-(4-chloro-2-(cyanomethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		416




Compound 233. 4-(1-(3-(2-(2-Cyanoethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 5, except 3-(4-methyl-1H-imidazol-2-yl)propanenitrile (compound **232.4**) was used in place of 2,4-dimethyl-1H-imidazole. m/z (ES+) 410 (M+H)⁺.



Compound 234.1. 1H-Imidazole-2-carbonitrile. Into a 1-L round-bottom flask, was placed a solution of 1H-imidazole-2-carbaldehyde (5 g, 52.04 mmol) in *N,N*-dimethylformamide (200 mL). Triethylamine (10.8 mL, 77.97 mmol), hydroxylamine hydrochloride (3.95 g, 56.84 mmol, 1.10 equiv), 1-propanephosphonic anhydride solution and 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphorinane-2,4,6-trioxide solution (T₃P) (36.4 g, 114.40 mmol) were added to the reaction. The reaction mixture was stirred for 4h at 100 °C, cooled and then quenched with 500 mL of water/ice. The aqueous phase was extracted with 3 x 1L of ethyl acetate, then the combined organic layers was washed with 2 x 1L of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/2) as eluent to furnish 2 g (41%) of the title compound as an off-white solid.




Compound 234.2. 5-Iodo-1H-imidazole-2-carbonitrile. Into a 100-mL round-bottom flask, was placed a solution 1H-imidazole-2-carbonitrile (compound **234.1**, 3.39 g, 36.42 mmol) in sodium hydroxide (54.7 mL, 2 M). Iodine (9.26 g, 36.46 mmol) was added to the reaction. The reaction mixture was stirred for 5h at room temperature. The pH of the solution was adjusted to 5 with HCl (2 M). The aqueous phase was extracted with 3 x 80 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/1.5) as eluent to furnish 1.02 g (13%) of the title compound as a white solid.




Compound 234.3. Methyl 3-(2-cyano-1H-imidazol-5-yl)-4-methylbenzoate. Into a 100-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 5-iodo-1H-imidazole-2-carbonitrile (compound **234.2**, 700 mg, 3.20 mmol) in a solvent mixture of DME and H₂O (30/3 mL). Methyl 4-methyl-3-(tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**, 1.06 g,

3.84 mmol), K_3PO_4 (2.71 g, 12.8 mmol), Xphos (152 mg, 0.32 mmol), $Pd(PPh_3)_4$ (369 mg, 0.32 mmol) were added to the reaction. The reaction mixture was stirred overnight at 90 °C, then concentrated under reduced pressure. The residue was diluted with 50 mL of ethyl acetate, then was washed with 3 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/petroleum ether (1/2) as eluent to furnish 160 mg (21%) of the title compound as a white solid.



Compound 234.4. Methyl 3-(4-chloro-2-cyano-1H-imidazol-5-yl)-4-


methylbenzoate. Into a 50-mL round-bottom flask, was placed a solution of methyl 3-(2-cyano-1H-imidazol-5-yl)-4-methylbenzoate (compound **234.3**, 290 mg, 1.20 mmol) in a solvent mixture of DMF and CHCl₃ (4/20 mL). NCS (161 mg, 1.21 mmol, 1.00 equiv) was added to the above reaction mixture. The reaction mixture was stirred overnight at room temperature, then was concentrated under reduced pressure. The residue was diluted with 30 mL of EtOAc, then was washed with 3 x 15 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/10-1/7) as eluent to furnish 160 mg (48%) of the title compound as light yellow oil.



Compound 234.5. 3-(4-Chloro-2-cyano-1H-imidazol-5-yl)-4-methylbenzoic acid.


Into a 25-mL round-bottom flask, was placed a solution of methyl 3-(4-chloro-2-cyano-1H-imidazol-5-yl)-4-methylbenzoate (compound **234.4**, 130 mg, 0.47 mmol) in methanol (4 mL). Then a solution of LiOH (45.4 mg, 1.90 mmol) in water (2 mL) was added to the reaction. The reaction mixture was stirred overnight at 30 °C, then was concentrated under reduced pressure. The pH of the solution was adjusted to 3 with hydrogen chloride (1M) and was concentrated under reduced pressure. Methanol (5 mL) was added to the residue. The salt was

filtered off, and the filtrate was concentrated under reduced pressure to give 150 mg (crude) of the title compound as a white solid.



Compound 234. 4-Chloro-5-(5-(4-(4-cyanophenyl)piperidine-1-carbonyl)-2-

methylphenyl)-1H-imidazole-2-carbonitrile. Into a 25-mL round-bottom flask, was placed a solution of 3-(4-chloro-2-cyano-1H-imidazol-5-yl)-4-methylbenzoic acid (compound 234.5, 60 mg, 0.23 mmol) in *N,N*-dimethylformamide (2 mL). 4-(Piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**, 61 mg, 0.27 mmol), HBTU (131 mg, 0.35 mmol), DIEA (123 μ L, 0.69 mmol) were added to the reaction. The reaction mixture was stirred for 30 min at 15 $^{\circ}$ C. The reaction mixture was diluted with 40 mL of EtOAc, then was washed with 2 x 30 mL of brine and 2 x 30 mL of NH_4Cl (sat.), dried over anhydrous sodium sulfate and concentrated under reduced pressure. The crude product (160 mg) was purified by Prep-HPLC with the following conditions (2#-AnalyseHPLC-SHIMADZU(HPLC-10)): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150 mm.; mobile phase, WATER WITH 0.05% TFA and MeCN (49.0% MeCN up to 62.0% in 7 min, up to 100.0% in 3 min, down to 49.0% in 1 min); Detector, Waters 2489, 254 and 220 nm. This resulted in 25.1 mg (25%) of the title compound as a white solid. m/z (ES+) 430 (M+H) $^{+}$.




Compound 235. 4-Chloro-5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-


methylphenyl)-1H-imidazole-2-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **234**, except 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**). m/z (ES+) 402 (M+H) $^{+}$.




Compound 236.1. 5-Iodo-2-methyl-1H-imidazole-4-carbaldehyde. Into a 100-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 2-methyl-1H-imidazole-4-carbaldehyde (1.5 g, 13.6 mmol) in CH₃CN (50 mL). NIS (3.7 g, 16.4 mmol) was added to the reaction. The reaction mixture was stirred for 4 h at 80 °C, then was concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (3:1) as eluent to furnish 2.11 g (66%) of the title compound as a yellow solid.




Compound 236.2. Methyl 5-(4-formyl-2-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate. Into a 100-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 5-iodo-2-methyl-1H-imidazole-4-carbaldehyde (compound **236.1**, 600 mg, 2.54 mmol) in dioxane (30 mL). Methyl 2,4-dimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**, 800 mg, 2.76 mmol), a solution of potassium carbonate (1.8 g, 13.0 mmol) in water (10 mL), and Pd(dppf)Cl₂ (400 mg, 0.55 mmol, 0.20 equiv) were added to the reaction. The reaction mixture was stirred for overnight at 80 °C, then was quenched with 20 mL of water. The aqueous phase was extracted with 2 x 100 mL of ethyl acetate and the combined organic layers were washed with 2 x 50 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (3:1) as eluent to furnish 400 mg (58%) of the title compound as a light yellow solid.




Compound 236.3. Methyl 5-(4-(hydroxymethyl)-2-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate. Into a 50-mL round-bottom flask, was placed a solution of methyl 5-(4-formyl-2-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **236.2**, 400 mg, 1.47 mmol) in methanol (20 mL). This was followed by the addition of NaBH₄ (100 mg, 2.64 mmol) in several batches at 0 °C. The reaction mixture was stirred for 1 h at room temperature, then was used for next step directly.




Compound 236. 4-(1-(5-(4-(Hydroxymethyl)-2-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except methyl 5-(4-(hydroxymethyl)-2-methyl-1H-imidazol-5-yl)-2,4-dimethylbenzoate (compound **236.3**) was used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **5.6**). *m/z* (ES⁺) 401 (M+H)⁺.




Compound 237.1. Methyl 3-(4-(hydroxymethyl)-2-methyl-1H-imidazol-5-yl)-4-methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **236.3**, except methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used in place of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**).




Compound 237. 4-(1-(3-(4-(Hydroxymethyl)-2-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 5, except methyl 3-(4-(hydroxymethyl)-2-methyl-1H-imidazol-5-yl)-4-methylbenzoate (compound 237.1) was used in place of methyl 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoate (compound 5.6). m/z (ES+) 387 (M+H)⁺.




Compound 238.1. 2-Cyclopropyl-1H-imidazole. Into a 500-mL round-bottom flask, was placed a solution of cyclopropanecarbaldehyde (10.7 mL, 142.7 mmol) in methanol (60 mL). An aqueous solution of oxalaldehyde (18.0 mL, 157.1 mmol, 40%) was added to the reaction. This was followed by the addition of NH₄OH (89 mL, 571.4 mmol) dropwise at 0 °C in 30 min. The reaction mixture was stirred overnight at room temperature, then concentrated under reduced pressure. The resulting solution was extracted with 3 x 100 mL of ethyl acetate. The combined organic layers were washed with 2 x 30 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 12 g (crude) of the title compound as a brown solid.




Compound 238.2. 2-Cyclopropyl-N,N-dimethyl-1H-imidazole-1-sulfonamide. Into a 500-mL round-bottom flask, was placed a solution of 2-cyclopropyl-1H-imidazole (compound 238.1, 12 g, 110.9 mmol) in dichloromethane (60 mL). Triethylamine (31.4 mL, 225.3 mmol), *N,N*-dimethylsulfamoyl chloride (13.2 mL, 121.87 mmol, 1.10 equiv) were added to the reaction. The reaction mixture was stirred for 4h at room temperature, then quenched with 20 mL of water. The aqueous phase was extracted with 3 x 100 mL of dichloromethane. The combined organic layers were washed with 2 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:4) as eluent to furnish 16 g (67%) of the title compound as a white solid.




Compound 238.3. 2-Cyclopropyl-1H-imidazole-4-carbaldehyde. Into a 500-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 2-cyclopropyl-*N,N*-dimethyl-1H-imidazole-1-sulfonamide (compound **238.2**, 5 g, 23.23 mmol) in tetrahydrofuran (100 mL). This was followed by the addition of *n*-butyllithium (11.2 mL, 2.5M in THF) dropwise at -78 °C over a period of 30 min. To this was added DMF (11.8 mL, 152.55 mmol). The resulting solution was stirred for 30 min at -50 °C, then hydrogen chloride (50 mL, 1M) was added. The reaction mixture was stirred for 2h at room temperature. The pH of the solution was adjusted to 7-8 with sodium bicarbonate (sat). The resulting solution was extracted with 3 x 30 mL of ethyl acetate and the organic layers combined. The resulting mixture was washed with 2 x 10 mL of brine. The mixture was dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 3 g (97%) of the title compound as a white solid.




Compound 238. 4-(1-(3-(2-Cyclopropyl-4-(hydroxymethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **236**, except 2-cyclopropyl-1H-imidazole-4-carbaldehyde (compound **238.3**) was used in place of 2-methyl-1H-imidazole-4-carbaldehyde and methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used in place of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**). *m/z* (ES⁺) 413 (M+H)⁺.




Compound 239.1. Ethyl 4-methyl-1H-imidazole-2-carboxylate. Into a 50-mL round-bottom flask, was placed a solution of ethyl 2-oxoacetate (400 mg, 3.92 mmol) in methanol (4 mL). A aqueous solution of 2-oxopropanal (565 mg, 3.92 mmol, 50%), ammonia (25%) (8.8 mL, 15.68 mmol) were added to the reaction. The reaction mixture was stirred for overnight at room temperature, then was concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/1) as eluent to furnish 200 mg (33%) of the title compound as a yellow solid.




Compound 239.2. Ethyl 5-iodo-4-methyl-1H-imidazole-2-carboxylate. Into a 100-mL round-bottom flask, was placed a solution of ethyl 4-methyl-1H-imidazole-2-carboxylate (compound **239.1**, 100 mg, 0.65 mmol) in CH₃CN (10 mL). NIS (218 mg, 0.97 mmol) was added to the reaction. The reaction mixture was stirred for overnight at room temperature, then concentrated under reduced pressure. The residue was dissolved with 30 mL of EtOAc, then was washed with 2 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 80 mg (44%) of the title compound as a yellow solid.




Compound 239.3. 5-Iodo-4-methyl-1H-imidazole-2-carboxylic acid. Into a 100-mL round-bottom flask, was placed a solution of ethyl 5-iodo-4-methyl-1H-imidazole-2-carboxylate (compound **239.2**, 200 mg, 0.71 mmol) in methanol (4 mL). A solution of sodium hydroxide (114 mg, 2.85 mmol) in water (2 mL) was added to the reaction. The reaction mixture was stirred for 2 h at 60 °C. The pH of the solution was adjusted to 1-2 with hydrogen chloride (1 M). The resulting mixture was concentrated under reduced pressure. Methanol (5 mL) was added to the residue. The salt was filtered off and the filtrate was concentrated under reduced pressure to give 290 mg (crude) of the title compound as a white solid.




Compound 239.4. 5-Iodo-*N,N*,4-trimethyl-1H-imidazole-2-carboxamide. Into a 100-mL round-bottom flask, was placed a solution of 5-iodo-4-methyl-1H-imidazole-2-carboxylic acid (compound **239.3**, 200 mg, 0.79 mmol) in *N,N*-dimethylformamide (4 mL). HBTU (602 mg, 1.59 mmol), dimethylamine hydrochloride (96 mg, 1.18 mmol) and triethylamine (552 μ L, 3.96 mmol) were added to the reaction. The reaction mixture was stirred for 3 h at room temperature, then diluted with 50 mL of EtOAc. The organic layer was washed with 2 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 150 mg (68%) of the title compound as yellow oil.




Compound 239. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2,4-dimethylphenyl)-*N,N*,4-trimethyl-1H-imidazole-2-carboxamide. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 5-iodo-*N,N*,4-trimethyl-1H-imidazole-2-carboxamide (compound **239.4**) was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES⁺) 442 (M+H)⁺.




Compound 240.1. 5-Iodo-4-methyl-2-(trifluoromethyl)-1H-imidazole. Into a 250-mL round-bottom flask, was placed a solution of 4-methyl-2-(trifluoromethyl)-1H-imidazole (compound **14.1**, 4.41 g, 29.38 mmol) in CH₃CN (100 mL). NIS (6.6 g, 29.34 mmol) was added to the reaction. The reaction mixture was stirred overnight at room temperature, then concentrated under reduced pressure. The reaction mixture was diluted with 300 mL of EtOAc. The organic layer was washed with 3 x 150 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/30-1/20) as eluent to furnish 2.125 g (26%) of the title compound as a light yellow solid.




Compound 240. 4-(1-(4-Methyl-3-(4-methyl-2-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 16, except 5-iodo-4-methyl-2-(trifluoromethyl)-1H-imidazole (compound **240.1**) was used in place of 5-iodo-2-methyl-1H-imidazole-4-carbonitrile (compound **16.4**). m/z (ES⁺) 425 (M+H)⁺.




Compound 241.1. 5-Iodo-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazole. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **44.3**, except tetrahydro-2H-pyran-4-carbaldehyde was used in place of cyclopropanecarbaldehyde.




Compound 241.2. Methyl 4-methyl-3-(2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoate. Into a 50-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 5-iodo-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazole (compound **241.1**, 400 mg, 1.44 mmol) in dioxane (15 mL). Methyl 4-methyl-3-(tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**, 397 mg, 1.44 mmol), potassium carbonate (993 mg, 7.18 mmol), water (1.5 mL) and Pd(dppf)Cl₂ (105 mg, 0.14 mmol, 0.10 equiv) were added to the reaction. The reaction mixture was stirred overnight at 90 °C, then diluted with 120 mL of EtOAc. The organic layer was washed with 3 x 40 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate as eluent to furnish 260 mg (60%) of the title compound as red oil.



Compound 241.3. Methyl 3-(4-chloro-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)-4-methylbenzoate. Into a 25-mL round-bottom flask, was placed a solution of methyl 4-methyl-3-(2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)benzoate (compound 241.2, 260 mg, 0.87 mmol) in chloroform (3 mL). NCS (116 mg, 0.87 mmol) was added to the reaction. The reaction mixture was stirred for 4 h at 25°C, then concentrated under reduced pressure. The residue was dissolved in 50 mL of EtOAc, then was washed with 3 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 313 mg (crude) of the title compound as brown oil.




Compound 241.4. 3-(4-Chloro-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)-4-methylbenzoic acid. Into a 50-mL round-bottom flask, was placed a solution of methyl 3-(4-chloro-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)-4-methylbenzoate (compound 241.3, 290 mg, 0.87 mmol) in methanol (6 mL). Then a solution of sodium hydroxide (139 mg, 3.48 mmol) in water (3 mL) was added to the reaction. The reaction mixture was stirred for 1 h at 60°C, then concentrated under reduced pressure. The pH of the solution was adjusted to 1 with hydrogen chloride (2 M). The resulting mixture was concentrated under reduced pressure. Methanol (10 mL) was added to the residue. The salt was filtered off, and the filtrate was concentrated under reduced pressure to give 300 mg of the title compound as a brown solid.




Compound 241. 4-(1-(3-(4-Chloro-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. Into a 25-mL round-bottom flask, was placed a solution of 3-(4-chloro-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)-4-methylbenzoic acid (compound 241.4, 150 mg, 0.47 mmol) in *N,N*-dimethylformamide (3


mL). 4-(Azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**, 91 mg, 0.47 mmol), EDC·HCl (180 mg, 0.94 mmol), 4-dimethylaminopyridine (114 mg, 0.93 mmol) were added to the reaction. The reaction mixture was stirred for 3 h at 25°C, then diluted with 80 mL of EtOAc. The organic layer was washed with 3 x 50 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The crude product (300 mg) was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-001(SHIMADZU)): Column, SunFire Prep C18 OBD Column, 5 µm, 19*150 mm; mobile phase, WATER WITH 0.05% TFA and ACN (25.0% ACN up to 39.0% in 8 min, hold 39.0% in 1 min, up to 100.0% in 3 min, down to 25.0% in 1 min); Detector, Waters 2489, 254 and 220 nm. This resulted in 133.3 mg (62%) of the title compound as a light yellow solid. m/z (ES+) 461 (M+H)⁺.




Compound 242.1. Methyl 5-(3,4-dimethyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **59.2**, except methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES+) 259 (M+H)⁺.




Compound 242.2. 5-(3,4-Dimethyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoic acid. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **59.3**, except methyl 5-(3,4-dimethyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoate (compound **242.1**) was used in place of methyl 3-(3,4-dimethyl-1H-pyrazol-5-yl)-4-methylbenzoate (compound **59.2**). m/z (ES+) 245 (M+H)⁺.



Compound 242. 4-(1-(5-(3,4-Dimethyl-1*H*-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **59**, except 5-(3,4-dimethyl-1*H*-pyrazol-5-yl)-2,4-dimethylbenzoic acid (compound **242.2**) was used in place of 3-(3,4-dimethyl-1*H*-pyrazol-5-yl)-4-methylbenzoic acid (compound **59.3**). m/z (ES⁺) 385 (M+H)⁺.




Compound 243. 4-(1-(5-(3,4-Dimethyl-1*H*-pyrazol-5-yl)-2,4-dimethylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **60**, except 5-(3,4-dimethyl-1*H*-pyrazol-5-yl)-2,4-dimethylbenzoic acid (compound **242.2**) was used in place of 3-(3,4-dimethyl-1*H*-pyrazol-5-yl)-4-methylbenzoic acid (compound **59.3**). m/z (ES⁺) 403 (M+H)⁺.



Compound 244.1. 1-Benzyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole. Into a 100-mL three neck round-bottom flask, was placed a solution of 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole (3.60 g, 18.6 mmol) in THF (50 mL). Sodium hydride (742 mg, 18.6 mmol, 60% dispersion in mineral oil) was carefully added in portions at 0 °C. The resulting mixture was stirred for 30 min at 0 °C, then benzyl bromide (2.21 mL, 18.6 mmol) was added. The resulting mixture was stirred overnight at room temperature, then carefully quenched with water (10 mL). The pH of the mixture was adjusted to 9-10 with aqueous HCl (2 M) and the aqueous phase was extracted with EtOAc (300 mL). The combined organic layers were washed with brine (2 x 150 mL), dried (Na₂SO₄), filtered and concentrated under vacuum. The residue was purified by silica gel

chromatography with ethyl acetate/petroleum ether (1/2) as eluent to yield the title compound as a yellow oil (3.47 g, 66%).




Compound 244.2. 1-Benzyl-1H-pyrazol-4-ol. Into a 100-mL three neck round-

bottom flask, was placed a solution of 1-benzyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (compound **244.1**, 3.47 g, 12.2 mmol) in tetrahydrofuran (35 mL). Sodium hydroxide (980 mg, 24.5 mmol) was added and then the mixture was cooled to 0 °C.

Hydrogen peroxide (2.51 mL, 24.4 mmol) was carefully added drop-wise and the resulting mixture was stirred for 2 h at room temperature. The reaction was carefully quenched with

aqueous Na₂S₂O₃(sat.) (20 mL). The aqueous phase was extracted with EtOAc (300 mL) and the combined organic layers were washed with brine (2 x 150 mL), dried (Na₂SO₄), filtered,


and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/1) as eluent to furnish the title compound as a white solid (1.64 g, 77%).




Compound 244.3. 1-Benzyl-4-methoxy-1H-pyrazole. Into a 100-mL round-bottom

flask, was placed a solution of 1-benzyl-1H-pyrazol-4-ol (compound **244.2**, 1.40 g, 8.04 mmol) in *N,N*-dimethylformamide (20 mL). Iodomethane (703 µL, 11.3 mmol) and Cs₂CO₃ (3.68 g, 11.3 mmol,) were added and the resulting mixture was stirred for 3 h at room


temperature. The mixture was concentrated under reduced pressure, then diluted with EtOAc (200 mL). The mixture was washed with brine (3 x 80 mL), and the organic layer was dried (Na₂SO₄), filtered and concentrated under reduced pressure to obtain the title compound as a yellow oil (1.30 g, 86%).



Compound 244.4. 4-Methoxy-1*H*-pyrazole. Into a 100-mL round-bottom flask, was placed a solution of 1-benzyl-4-methoxy-1*H*-pyrazole (compound **244.3**, 666 mg, 3.54 mmol) in methanol (10 mL). Hydrogen chloride (1 M, 1 mL) and Pd(OH)₂/C (20 wt %, 670 mg) were added and the mixture was purged with nitrogen then charged with hydrogen. The mixture was stirred under hydrogen overnight at room temperature, then purged with nitrogen. The solids were removed by filtration and the filtrate was concentrated under reduced pressure to obtain the title compound as a brown oil (0.36 g, crude).




Compound 244.5. 3-Iodo-4-methoxy-1*H*-pyrazole. Into a 50-mL round-bottom flask, was placed 4-methoxy-1*H*-pyrazole (compound **244.4**, 360 mg, 3.67 mmol) in *N,N*-dimethylformamide (4 mL). NIS (830 mg, 3.69 mmol) was added portion-wise and the resulting mixture was stirred overnight at room temperature. The mixture was diluted with ethyl acetate (40 mL), then washed with brine (3 x 10 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:10) as the eluent to obtain the title compound as a white solid (0.54 g, 66%).




Compound 244.6. Methyl 3-(4-methoxy-1*H*-pyrazol-5-yl)-4-methylbenzoate. Into a 50-mL three neck round-bottom flask, purged and maintained with nitrogen, was placed a solution of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (414 mg, 1.50 mmol) in dioxane (6 mL). Potassium carbonate (2 M, 2.5 mL, 5.00 mmol), 3-iodo-4-methoxy-1*H*-pyrazole (compound **244.5**, 224 mg, 1.00 mmol), Pd(dppf)Cl₂ (73 mg, 0.10 mmol) were added to the reaction and the resulting mixture was stirred overnight at 90 °C. The resulting mixture was cooled, then diluted with EtOAc (80 mL) and washed with brine (2 x 30 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was

purified by a silica gel column chromatography with ethyl acetate/petroleum ether (1/1) as eluent to obtain the title compound as a yellow oil (180 mg, 73%).



Compound 244.7. 3-(4-Methoxy-1H-pyrazol-5-yl)-4-methylbenzoic acid. Into a


5 50-mL round-bottom flask, was placed a solution of methyl 3-(4-methoxy-1H-pyrazol-5-yl)-4-methylbenzoate (compound **244.6**, 180 mg, 0.73 mmol) in methanol (4 mL). Sodium hydroxide (117 mg, 2.92 mmol) in water (2 mL) was added and the resulting solution was stirred for 3 h at 60 °C, then concentrated under reduced pressure. The pH of the residue was adjusted to 1 with aqueous hydrogen chloride (1 M), then concentrated under reduced
10 pressure. Methanol (5 mL) was added to the residue and the solids were filtered from the solution. The filtrate was concentrated under reduced pressure to obtain the title compound as a light yellow solid (353 mg, crude).




Compound 244. 4-(1-(3-(4-Methoxy-1H-pyrazol-5-yl)-4-methylbenzoyl)azetidin-

15 **3-yl)benzonitrile.** Into a 25-mL round-bottom flask, was placed a solution of 3-(4-methoxy-1H-pyrazol-5-yl)-4-methylbenzoic acid (compound **244.7**, 60 mg, 0.26 mmol) in *N,N*-dimethylformamide (2 mL). 4-(Azetidin-3-yl)benzonitrile hydrochloride (75 mg, 0.39 mmol), EDC·HCl (99 mg, 0.52 mmol), and 4-dimethylaminopyridine (63 mg, 0.52 mmol) were added and the solution was stirred for 2 h at room temperature. The reaction was diluted with
20 EtOAc (30 mL) and washed with brine (3 x 15 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The crude product (150 mg) was purified by Prep-HPLC with the following conditions (1#-Pre-HPLC-010(Waters)): Column, SunFire Prep C18 OBD Column, 5 μm, 19*150 mm; mobile phase, water with 0.05% TFA and CH₃CN (35.0% CH₃CN up to 50.0% in 10 min, up to 100% in 1 min, down to 35.0% in 2 min);
25 Detector, UV 254 & 220nm. The fractions contained pure product were combined and

lyophilized to obtain the title compound as an off-white solid (44.3 mg, 46%). m/z (ES+) 373 ($M+H^+$).




5 **Compound 245. 4-(1-(3-(4-Methoxy-1H-pyrazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 401 ($M+H^+$).



10


Compound 246. 4-(3-Fluoro-1-(3-(4-methoxy-1H-pyrazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 391 ($M+H^+$).

15




20


Compound 247. 4-(1-(5-(4-Methoxy-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES+) 387 ($M+H^+$).




Compound 248.1. 1-Benzyl-4-bromo-3-methyl-1H-pyrazole. Into a 500-mL 3-neck round-bottom flask, was placed a solution of 4-bromo-3-methyl-1H-pyrazole (20.0 g, 124 mmol) in *N,N*-dimethylformamide (100 mL). The system was purged with nitrogen, then the solution was cooled to 0 °C. Sodium hydride (60% dispersion in mineral oil, 10.0 g, 250 mmol) was carefully added portion-wise and then the resulting mixture was stirred for 30 min at room temperature. The mixture was cooled to 0 °C, then benzyl bromide (22.1 mL, 186 mmol) was added drop-wise and the resulting mixture was stirred overnight at room temperature. The mixture was carefully quenched by drop-wise addition of water (40 mL), then the mixture was extracted with ethyl acetate (3 x 200 mL). The combined organic extracts were washed with brine (4 x 100 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure to obtain the title compound as a yellow oil (24.9 g, 80%).




Compound 248.2. 1-Benzyl-3-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole. Into a 250-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 1-benzyl-4-bromo-3-methyl-1H-pyrazole (compound **248.1**, 24.5 g, 97.6 mmol) in DMSO (160 mL). 4,4,4',4',5,5,5',5'-Octamethyl-2,2'-bi(1,3,2-dioxaborolane) (29.9 g, 118 mmol), Pd(dppf)Cl₂ (7.16 g, 9.79 mmol) and KOAc (28.8 g, 293 mmol) were added. The resulting mixture was stirred overnight at 90 °C, then cooled and quenched with water (100 mL). The mixture was extracted with ethyl acetate (3 x 100 mL) and the combined organic extracts were washed with brine (4 x 100 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by a silica gel column with ethyl acetate/petroleum ether (1:20) as eluent to obtain the title compound as a yellow oil (10.64 g, 37%).



Compound 248. 4-(1-(3-(4-Methoxy-3-methyl-1H-pyrazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 244, except 1-benzyl-3-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (compound **248.2**) was used in place of 1-benzyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (compound **244.1**). m/z (ES⁺) 387 (M+H⁺).




Compound 249.1. Ethyl 2-methyl-3-oxopropanoate. Into a 500-mL 3-neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of ethyl propionate (11.3 mL, 97.9 mmol) in tetrahydrofuran (160 mL). The solution was cooled to 0 °C, then sodium hydride (4.1 g, 103mmol, 60% dispersion in mineral oil) was added in portions. The mixture was stirred at 0 °C for 10 min, then ethyl formate (7.92 mL, 98.5 mmol) was added and the resulting mixture was stirred overnight at room temperature. The reaction was carefully quenched with water (100 mL) by drop-wise addition and the mixture was extracted with ethyl acetate (2 x 100 mL). The combined organic extracts were dried (Na₂SO₄), filtered and concentrated under reduced pressure. The crude product was purified by distillation under reduced pressure (50 mm Hg) and the fraction between 70-80 °C was collected to obtain the title compound as a colorless liquid (3.0 g, 24%).




Compound 249.2. 1-Benzyl-4-methyl-1H-pyrazol-5-ol. Into a 100-mL round-bottom flask, was placed a solution of ethyl 2-methyl-3-oxopropanoate (compound **249.1**, 500 mg, 3.84 mmol) in ethanol (15 mL). Benzylhydrazine hydrochloride (730 mg, 4.66 mmol) was added and the reaction was stirred for 5 h at 80 °C, then concentrated under reduced


pressure. The residue was purified by a silica gel chromatography with dichloromethane/methanol (30:1) as eluent to obtain the title compound as a yellow oil (430 mg, 59%).




Compound 249.3. 1-Benzyl-5-methoxy-4-methyl-1H-pyrazole. Into a 100-mL round-bottom flask, was placed a solution of 1-benzyl-4-methyl-1H-pyrazol-5-ol (compound **249.2**, 2.0 g, 10.6 mmol) in *N,N*-dimethylformamide (15 mL). The solution was cooled to 0 °C, then sodium hydride (600 mg, 15.00 mmol, 60% dispersion in mineral oil) was added in portions and the resulting mixture was stirred for 10 min at 0 °C. Iodomethane (878 µL, 14.1 mmol) was added and the resulting mixture was stirred for 3 h at room temperature, then carefully quenched by drop-wise addition of water (10 mL). The mixture was extracted ethyl acetate (3 x 40 mL) and the combined organic extracts were washed with brine (3 x 20 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/petroleum ether (1:8) as eluent to obtain the title compound as a yellow oil (500 mg, 23%).




Compound 249.4. 5-Methoxy-4-methyl-1H-pyrazole. Into a 100-mL round-bottom flask, was placed a solution of 1-benzyl-5-methoxy-4-methyl-1H-pyrazole (compound **249.3**, 300 mg, 1.48 mmol) in DMSO (10 mL). Potassium *tert*-butoxide (11 mL, 11 mmol, 1 M in THF) was added slowly and the resulting mixture was stirred overnight at room temperature under dry air conditions. The pH of the solution was adjusted to 7-8 with aqueous HCl (2M), diluted with additional water and the mixture was extracted with ethyl acetate (3 x 30 mL). The combined organic extracts were washed with brine (3 x 20 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:8) as eluent to obtain the title compound as a yellow oil (150 mg, 90%).



Compound 249. 4-(1-(5-(3-Methoxy-4-methyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 5-methoxy-4-methyl-1H-pyrazole (compound **249.4**) was used in place 4-methoxy-1H-pyrazole (compound **244.4**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) m/z (ES⁺) 401 (M+H⁺).




Compound 250.1. 5-Iodo-4-methyl-1H-pyrazole. NIS (5.62g, 24.98mmol) was added portion-wise to a solution of 4-methyl-1H-pyrazole (Aldrich, 2.05g, 24.97mmol) in acetonitrile (50 mL). The mixture was heated at 60 °C for 30 minutes, then cooled to room temperature. The mixture was partitioned between EtOAc (300 mL) and water (80 mL). The organic layer was washed with saturated sodium thiosulfate (50 mL), brine (50 mL), dried (MgSO₄) and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexanes:EtOAc, 5:1) to yield the title compound as a white solid (2.3g, 45%). m/z (ES⁺) 209 (M+H)⁺.




Compound 250.2. 5-Iodo-4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole. NaH (0.48 g, 12.2 mmol, 60%) was added to a solution of 5-iodo-4-methyl-1H-pyrazole (compound **250.1**, 2.3 g, 11.1 mmol) in THF (20 mL) at 0 °C. The mixture was stirred at 0 °C for 30 minutes, then 2-(trimethylsilyl)ethoxymethyl chloride (2.15 mL, 12.2 mmol) was added. The resulting mixture was stirred at room temperature for 3 hours, and then carefully quenched with a small amount of water and partitioned between EtOAc (200

mL) and water (50 mL). The organic layer was washed with brine (50 mL), dried (MgSO₄) and concentrated under reduced pressure. The residue was purified with by silica gel column chromatography (hexanes:EtOAc, 10:1) to yield the title compound as a light yellow oil (3.15 g, 84%). *m/z* (ES⁺) 339 (M+H)⁺.




Compound 250.3. Methyl 2,4-dimethyl-5-(4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-5-yl)benzoate. Methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-

dioxaborolan-2-yl)benzoate (compound **160.1**, 2.7 g, 9.32 mmol), 5-iodo-4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole (compound **250.2**, 3.15 g, 9.32 mmol), K₂CO₃ (6.42 g, 46.6 mmol) and PdCl₂(dppf)·CH₂Cl₂ (0.76 g, 0.93 mmol) were added to a round bottom flask. The flask was purged with argon, then dioxane (50mL) and water (20 mL) were added and the mixture was heated at 90 °C for 16 hours. The mixture was cooled to room temperature and then partitioned between EtOAc (300 mL) and water (50 mL). The organic layer was washed with brine (50 mL), dried (MgSO₄) and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexanes:EtOAc, 10:1) to yield the title compound as a light brown oil (3.0 g, 86%). *m/z* (ES⁺) 375(M+H)⁺.



Compound 250.4. Methyl 5-(3-chloro-4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-5-yl)-2,4-dimethylbenzoate. To a solution of methyl 2,4-dimethyl-5-(4-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-5-yl)benzoate (compound **250.3**, 3.0 g, 8.02 mmol) in acetonitrile (20 mL) was added NCS (1.07 g, 8.02 mmol). The mixture was stirred at room temperature overnight, then partitioned between EtOAc (200 mL) and water (50 mL). The organic layer was washed with saturated sodium thiosulfate (50 mL), brine (50 mL), dried (MgSO₄) and concentrated under reduced pressure. The residue was purified with by silica gel column chromatography (hexanes:EtOAc, 20:1) to yield the title compound as a colorless oil (2.0 g, 61%). *m/z* (ES⁺) 409 (M+H)⁺.




Compound 250.5. Methyl 5-(3-chloro-4-methyl-1H-pyrazol-5-yl)-2,4-

dimethylbenzoate. TFA (10mL) was carefully added to a solution of 5-(3-chloro-4-methyl-1-((2-(trimethylsilyl)ethoxy) methyl)-1H-pyrazol-5-yl)-2,4-dimethylbenzoate (compound


250.4, 2.0 g, 4.90 mmol) in CH₂Cl₂ (10mL). The mixture was stirred at room temperature for 2 hours, then the solvents were removed under reduced pressure. The residue was dissolved in EtOAc (200 mL) and washed with sat. NaHCO₃ (50 mL), brine (50 mL), dried (MgSO₄) and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexanes:EtOAc, 4:1) to yield the title compound as white foam (1.13 g,

83%). *m/z* (ES⁺) 279(M+H)⁺.



Compound 250.6. 5-(3-Chloro-4-methyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoic acid.


NaOH (2 M in water, 8 mL, 16 mmol) was added to a solution of methyl 5-(3-chloro-4-methyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoate (compound **250.5**, 1.10 g, 3.96 mmol) in MeOH (10 mL). The mixture was stirred at room temperature overnight and the solvents were removed under the reduced pressure. The residue was cooled to 0 °C and acidified to pH ~3-4 with 1M HCl in water. The resulting white solids were filtered, washed with water and dried to give the title compound as a white solid (1.07g, 90%). *m/z* (ES⁺) 265 (M+H)⁺.




Compound 250. 4-(1-(5-(3-Chloro-4-methyl-1H-pyrazol-5-yl)-2,4-

dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of


compound **5**, except 5-(3-chloro-4-methyl-1*H*-pyrazol-5-yl)-2,4-dimethylbenzoic acid (compound **250.6**) was used in place of 3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). m/z (ES⁺) 405 (M+H)⁺.




Compound 251. 4-(1-(5-(3-Chloro-4-methyl-1*H*-pyrazol-5-yl)-2,4-dimethylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 5-(3-chloro-4-methyl-1*H*-pyrazol-5-yl)-2,4-dimethylbenzoic acid (compound **250.6**) was used in place of 3-(2,4-dimethyl-1*H*-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 423 (M+H)⁺.




Compound 252. 4-(1-(3-(3-Chloro-4-methyl-1*H*-pyrazol-5-yl)-4-methylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **250**, except methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used in place of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 409 (M+H)⁺.



Compound 253. 4-(1-(3-(3-Chloro-4-methyl-1H-pyrazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **250**, except methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used in place of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) and 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES⁺) 419 (M+H⁺).




Compound 136. 4-(1-(5-(4-Chloro-3-methyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **51**, except 5-bromo-3-methyl-1H-pyrazole was used in place of 5-iodo-1H-pyrazole. m/z (ES⁺) 405 (M+H⁺).



Compound 255.1. 5-(Methoxycarbonyl)-2,4-dimethylbenzoic acid. A solution of 5-iodo-2,4-dimethylbenzoic acid (5.0 g, 18.1 mmol) in methanol (45 mL) was added to a 100-mL pressure tank reactor (40 atm). Triethylamine (7.57 mL, 54.3 mmol) and Pd(dppf)Cl₂ (1.3 g, 1.8 mmol) were added and the system was purged with nitrogen. CO (g) was very carefully introduced. **Caution:** carbon monoxide is a very toxic gas. The reaction was heated at 90 °C overnight, then cooled to room temperature and very carefully evacuated. The solvent was removed under reduced pressure. The pH of the residue was adjusted to 1-2 with


aq HCl (6 M), diluted with additional water, then extracted with EtOAc (2 x 100 mL). The combined organic extracts were washed brine (3 x 10 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by a silica gel column with ethyl acetate/hexane (1:1) as eluent to obtain the title compound as a red solid (2.3 g, 61%).



Compound 255.2. Methyl 5-(chlorocarbonyl)-2,4-dimethylbenzoate. Into a 100-

mL round-bottom flask, was placed a solution of 5-(methoxycarbonyl)-2,4-dimethylbenzoic acid (compound **255.1**, 100 mg, 0.48 mmol) in dichloromethane (10 mL). Oxalyl chloride (80 μL, 0.95 mmol) was added and the resulting solution was stirred for 1 h at 40 °C, then

concentrated under reduced pressure to obtain the title compound as a white solid (80 mg, 74%).




Compound 255.3. Methyl 5-(2-(difluoromethylene)butanoyl)-2,4-

dimethylbenzoate. Into a 100-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of argon, was placed a solution of 2,2,2-trifluoroethyl 4-methylbenzenesulfonate (1.00 g, 3.93 mmol) in THF (30 mL). The solution was cooled to -78 °C, then n-BuLi (5.0 mL, 1.6 M in hexane, 8.0 mmol) was added drop-wise and the mixture was stirred at -78 °C for 40 min. Triethylborane (TEB) (5.0 mL, 1 M in THF, 5.0 mmol) was slowly added and the mixture was stirred at -78 °C for 1 h, then at room temperature for 3 h.


The solution was cooled to 0 °C, then hexamethylphosphoramide (HMPA) (6 mL) and CuI (1.5 g, 7.9 mmol) were added and the mixture was stirred at room temperature for 30 min. A solution of methyl 5-(chlorocarbonyl)-2,4-dimethylbenzoate (compound **255.2**, 530 mg, 2.34 mmol) in THF (30 mL) was carefully added and stirred at room temperature for 30 min. The pH of the mixture was carefully adjusted to 7 with aqueous H₃PO₄ and the mixture was partitioned. The aqueous phase was extracted with ethyl acetate (3 x 20 mL) and the combined organic layers were concentrated under reduced pressure. The residue was purified

by silica gel column chromatography with ethyl acetate/petroleum ether (1:40) as eluent to obtain the title compound as a yellow oil (250 mg, 38%).




Compound 255.4. Methyl 5-(4-ethyl-3-fluoro-1H-pyrazol-5-yl)-2,4-

- 5 **dimethylbenzoate.** Into a 250-mL round-bottom flask, was placed a solution of methyl 5-(2-(difluoromethylene)butanoyl)-2,4-dimethylbenzoate (compound **255.3**, 175 mg, 0.62 mmol) in THF (100 mL). At room temperature, hydrazine hydrate (120 μ L, 2.48 mmol) was added followed by trifluoroacetic acid (144 μ L, 1.88 mmol), then the solution was stirred at 50 $^{\circ}$ C for 1.5 h. The reaction was cooled to room temperature, carefully diluted with water and the
- 10 mixture was extracted with ethyl acetate (3 x 50 mL). The combined organic extracts were dried (Na_2SO_4), filtered and concentrated under reduced pressure. The residue was purified by a silica gel column with ethyl acetate/hexane (1:10) as eluent to obtain the title compound as a yellow oil (90 mg, 53%).




- 15 **Compound 255. 4-(1-(5-(4-Ethyl-3-fluoro-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except methyl 5-(4-ethyl-3-fluoro-1H-pyrazol-5-yl)-2,4-dimethylbenzoate (compound **255.4**) was used in place methyl 3-(4-methoxy-1H-pyrazol-5-yl)-4-
- 20 methylbenzoate (compound **244.6**). m/z (ES $^{+}$) 403 ($\text{M}+\text{H}^{+}$).




Compound 256.1. 3-Iodo-1H-pyrazole-4-carbonitrile. Into a 100-mL three neck round-bottom flask, was placed a solution of 3-amino-1H-pyrazole-4-carbonitrile (1.00 g, 9.25 mmol) in hydrogen chloride (12 M, 8 mL). The reaction solution was cooled to 0 $^{\circ}$ C,

then a solution of sodium nitrite (800 mg, 11.6 mmol) in water (5.8 mL) was carefully added dropwise and the mixture was stirred for 1 h at 0 °C. KI (1.91 g, 11.6 mmol) was added and the resulting mixture was stirred for 3 h at room temperature. The pH of the solution was carefully adjusted to 10 with aqueous sodium hydroxide (3 M) and then extracted with ethyl acetate (3 x 100 mL). The combined organic layers were dried (Na₂SO₄), filtered and concentrated under reduced pressure to obtain the title compound as a white solid (700 mg, 35%).



Compound 256.2. 3-Iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole-4-


carbonitrile. Into a 100-mL round-bottom flask, was placed a solution of 3-iodo-1H-pyrazole-4-carbonitrile (compound **256.1**, 5.00 g, 22.8 mmol) in tetrahydrofuran (10 mL). The solution was cooled to 0 °C under nitrogen, then sodium hydride (1.1 g, 27.4 mmol, 60% dispersion in mineral oil) was added in portions. The resulting mixture was stirred at 0 °C for 30 min, then (2-(chloromethoxy)ethyl)trimethylsilane (6.05 mL, 34.2 mmol) was added to the reaction. The resulting mixture was stirred for overnight at room temperature, then carefully quenched with water/ice (30 mL). The aqueous phase was extracted with ethyl acetate (2 x 100 mL) and the combined organic layers were dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/ petroleum ether (20:1) as eluent to obtain the title compound as a colorless oil (7.2 g, 90%).



Compound 256.3. Methyl 3-(4-cyano-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-5-yl)-4-methylbenzoate. Into a 250-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole-4-carbonitrile (compound **256.2**, 2.73 g, 7.82 mmol, 1.00 equiv) in dioxane (30 mL). Methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-

dioxaborolan-2-yl)benzoate (2.59 g, 9.38 mmol), Pd(dppf)Cl₂ (570 mg, 0.78 mmol), and a solution of potassium carbonate (3.24 g, 23.4 mmol) in water (3 mL) were added to the reaction. The resulting mixture was stirred overnight at 90 °C, then cooled and diluted with EtOAc (500 mL). The resulting mixture was washed with brine (2 x 100 mL), dried


5 (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/petroleum ether (40:1) as eluent to obtained the title compound as a yellow oil (1.9 g, 65%).



Compound 256.4. Methyl 3-(4-cyano-1H-pyrazol-5-yl)-4-methylbenzoate. Into a

50-mL round-bottom flask, was placed a solution of methyl 3-(4-cyano-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-5-yl)-4-methylbenzoate (compound **256.3**, 1.9 g, 5.11 mmol) in triethylsilane (4 mL) and trifluoroacetic acid (8 mL). The resulting solution was stirred for 4 h at room temperature, then concentrated under reduced pressure. The pH of the residue was carefully adjusted to 7-8 with aqueous sodium bicarbonate (sat.), then the


15 aqueous phase was extracted with ethyl acetate (2 x 100 mL). The combined organic layers were dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (20/1) as eluent to obtain the title compound as a white solid (600 mg, 49%).




Compound 256.5. 3-(4-Cyano-1H-pyrazol-5-yl)-4-methylbenzoic acid. The title

compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244.7**, except methyl 3-(4-cyano-1H-pyrazol-5-yl)-4-methylbenzoate (compound **256.4**) was used in place of methyl 3-(4-methoxy-1H-


pyrazol-5-yl)-4-methylbenzoate (compound **244.6**).




Compound 256. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-1H-pyrazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 3-(4-cyano-1H-pyrazol-5-yl)-4-methylbenzoic acid (compound **256.5**) was used in place of 3-(4-methoxy-1H-pyrazol-5-yl)-4-methylbenzoic acid (compound **244.7**). m/z (ES+) 368 (M+H⁺).




Compound 257.1. 3-Oxobutanenitrile. Into a 250-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was carefully placed (Z)-3-aminobut-2-enenitrile (20.0 g, 244 mmol) and aqueous hydrogen chloride (6 M) (80 mL). The mixture was stirred for 3 h at 80 °C, then the solids were removed by filtration. The filtrate was extracted with ethyl acetate (3 x 100 mL) and the combined organic extracts were dried (Na₂SO₄), filtered and concentrated under reduced pressure to obtain the title compound as a yellow solid (10 g, 49%).




Compound 257.2. (Z)-2-(Ethoxymethylene)-3-oxobutanenitrile. Into a round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 3-oxobutanenitrile (compound **257.1**, 16.0 g, 193 mmol) in acetic anhydride (48 mL). Triethyl orthoformate (96 mL, 577 mmol) was added and the reaction was stirred for 2 h at 130 °C, then concentrated under reduced pressure to obtain the title compound as a red solid (10 g, 37%).



Compound 257.3. 3-Methyl-1H-pyrazole-4-carbonitrile. Into a 50-mL round-bottom flask, was placed a solution of (Z)-2-(ethoxymethylene)-3-oxobutanenitrile (compound **257.2**, 5.00 g, 35.9 mmol) in ethanol (10 mL). The solution was cooled to 0 °C, then hydrazine monohydrate (2.09 mL, 43.1 mmol) was added and the reaction was stirred for 1 h at 10 °C. The reaction was diluted with water (10 mL) and then extracted with ethyl acetate (3 x 50 mL). The combined organic extracts were dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by a silica gel column with ethyl acetate/petroleum ether (1:10) as eluent to obtain the title compound as a yellow solid (2.0 g, 52%).




Compound 257.4. 5-Iodo-3-methyl-1H-pyrazole-4-carbonitrile. Into a 10-mL sealed tube, was placed a solution of 3-methyl-1H-pyrazole-4-carbonitrile (compound **257.3**, 2.0 g, 18.7 mmol) in DCE (6 mL). NIS (4.20 g, 18.7 mmol) was added to the reaction and the mixture was irradiated with microwave radiation for 1 h at 150 °C. The reaction was cooled, then carefully quenched with aqueous Na₂S₂O₃ (sat., 20 mL). The mixture was partitioned and the aqueous layer was extracted with ethyl acetate (100 mL). The combined organic layers were washed with brine (5 x 30 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:10) as eluent to obtain the title compound as a yellow solid (900 mg, 21%).



Compound 257.5. 3-(4-Cyano-3-methyl-1H-pyrazol-5-yl)-4-methylbenzoic acid.


The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **256.5**, except 5-iodo-3-methyl-1H-

pyrazole-4-carbonitrile (compound **257.4**) was used in place 3-iodo-1*H*-pyrazole-4-carbonitrile (compound **256.1**).




Compound 257. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-

3-methyl-1*H*-pyrazole-4-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 3-(4-cyano-3-methyl-1*H*-pyrazol-5-yl)-4-methylbenzoic acid (compound **257.5**) was used in place of 3-(4-methoxy-1*H*-pyrazol-5-yl)-4-methylbenzoic acid (compound **244.7**). m/z (ES⁺) 382 (M+H⁺).



Compound 258.1. Ethyl 5-iodo-4-methyl-1*H*-pyrazole-3-carboxylate. Into a 250-


mL round-bottom flask, was placed a solution of ethyl 4-methyl-1*H*-pyrazole-3-carboxylate (600 mg, 3.89 mmol) in *N,N*-dimethylformamide (20 mL). NIS (2.2 g, 9.78 mmol) was added and the resulting mixture was stirred for 6 h at 50 °C, then quenched with water (10 mL). The mixture was extracted with ethyl acetate (3 x 40 mL) and the combined organic extracts were washed with aqueous Na₂S₂O₃ (sat., 2 x 30 mL) and brine (2 x 20 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/petroleum ether (1:10) as eluent to obtain the title compound as a white solid (800 mg, 73%).




Compound 258.2. 5-Iodo-4-methyl-1*H*-pyrazole-3-carboxamide. Into a 100-mL

round-bottom flask, was placed a solution of ethyl 5-iodo-4-methyl-1*H*-pyrazole-3-carboxylate (compound **258.1**, 500 mg, 1.79 mmol) in methanol (5 mL). Then ammonium


hydroxide (15 mL, 25% aq.) was added and the resulting solution was stirred overnight at 60 °C. The mixture was concentrated under reduced pressure to obtain the title compound as a white solid (400 mg, 89%).




Compound 258.3. 5-Iodo-4-methyl-1H-pyrazole-3-carbonitrile. Into a 100-mL round-bottom flask, was placed a solution of 5-iodo-4-methyl-1H-pyrazole-3-carboxamide (compound **258.2**, 800 mg, 3.19 mmol) in dichloromethane (10 mL). The solution was cooled to 0 °C, then trifluoroacetic anhydride (1.32 mL, 9.5 mmol) and triethylamine (3.16 mL, 22.7 mmol) were each added carefully drop-wise. The mixture was stirred for 2 h at 0 °C, then the pH was carefully adjusted to 7-8 with aqueous sodium bicarbonate (10%). The aqueous phase was extracted with ethyl acetate (2 x 40 mL) and the combined organic layers were washed with 2x10 mL of brine (2 x 10 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/petroleum ether (1:10) as eluent to obtain the title compound as a white solid (280 mg, 38%).




Compound 258. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-4-methyl-1H-pyrazole-3-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 5-iodo-4-methyl-1H-pyrazole-3-carbonitrile (compound **258.3**) was used in place of 3-iodo-4-methoxy-1H-pyrazole (compound **244.5**). *m/z* (ES⁺) 382 (M+H⁺).




Compound 259. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2,4-dimethylphenyl)-4-methyl-1H-pyrazole-3-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **258**, except methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES+) 396 (M+H⁺).




Compound 260.1. Methyl 5-nitro-1H-pyrazole-3-carboxylate. Into a 100-mL round-bottom flask, was placed a solution of 5-nitro-1H-pyrazole-3-carboxylic acid (3.0 g, 19.1 mmol) in methanol (50 mL). Sulfuric acid (3 mL) was carefully added and the resulting solution was stirred overnight at 80 °C, then concentrated under reduced pressure. The pH of the solution was carefully adjusted to 7 with aqueous sodium bicarbonate (sat.), then extracted with ethyl acetate (3 x 50 mL). The combined organic extracts were washed with brine (3 x 50 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure to obtain the title compound as a yellow solid (2.3 g, 70%).



Compound 260.2. Methyl 5-amino-1H-pyrazole-3-carboxylate. Into a 100-mL round-bottom flask, was placed a solution of methyl 5-nitro-1H-pyrazole-3-carboxylate (compound **260.1**, 2.5 g, 14.6 mmol) in methanol (10 mL). The system was purged with nitrogen, then palladium/carbon (10 wt % Pd, 0.8 g) was added. The system was charged with hydrogen (1 atm) and the mixture was stirred overnight at room temperature. The system was purged with nitrogen, then the solids were removed by filtration. The filtrate was concentrated under reduced pressure to obtain the title compound as an off-white solid (1.5 g, 73%).



Compound 260.3. Methyl 5-iodo-1H-pyrazole-3-carboxylate. Into a 100-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of methyl-5-amino-1H-pyrazole-3-carboxylate (compound **260.2**, 1.0 g, 7.1 mmol) in a carefully prepared solution of sulfuric acid (15 mL) into water (30 mL) (**Caution:** addition of sulfuric acid to water is exothermic). The solution was cooled to 0 °C, then a solution of NaNO₂ (0.53 g, 7.7 mmol) in water (5 mL) was added drop-wise. The resulting mixture was stirred for 2 h at 0 °C, then a solution of KI (1.4 g, 8.4 mmol) in water (5 mL) was added drop-wise at 0 °C. The resulting mixture was stirred overnight at room temperature, then the pH of the mixture was carefully adjusted to 7 with aqueous sodium bicarbonate (sat.). The mixture was extracted with ethyl acetate (3 x 50 mL) and the combined organic extracts were washed with brine (3 x 50 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/petroleum ether (1:5) as eluent to obtain the title compound as a yellow solid (0.6 g, 30%).




Compound 260.4. 5-Iodo-1H-pyrazole-3-carboxamide. Into a 100-mL round-bottom flask, was placed a solution of methyl 5-iodo-1H-pyrazole-3-carboxylate (compound **260.3**, 900 mg, 3.57 mmol) in methanol (5 mL). Ammonia (25% aq.) (15 mL) was added and the resulting solution was stirred overnight at 60 °C, then cooled and concentrated under reduced pressure to obtain the title compound as a white solid (0.8 g, 95%).




Compound 260.5. 5-Iodo-1H-pyrazole-3-carbonitrile. Into a 100-mL round-bottom flask, was placed a solution of 5-iodo-1H-pyrazole-3-carboxamide (compound **260.4**, 1.00 g, 4.22 mmol) in dichloromethane (10 mL). The solution was cooled to 0 °C, then trifluoroacetic


anhydride (1.8 mL, 12.9 mmol) and triethylamine (4.11 mL, 29.5 mmol) were added carefully drop-wise. The mixture was stirred for 2 h at 0 °C, then the pH was carefully adjusted to 7-8 with aqueous sodium bicarbonate (10%). The aqueous phase was extracted with ethyl acetate (3 x 30 mL) and the combined organic layers were washed with brine (2 x 20 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:10) as eluent to obtain the title compound as a light yellow solid (180 mg, 19%).



Compound 260.6. 4-Chloro-5-iodo-1H-pyrazole-3-carbonitrile. Into a 50-mL round-bottom flask, was placed a solution of 5-iodo-1H-pyrazole-3-carbonitrile (compound **260.5**, 100 mg, 0.46 mmol) in *N,N*-dimethylformamide (10 mL). NCS (61.6 mg, 0.46 mmol) was added and the resulting mixture was stirred for 2 h at 50 °C, then quenched with water (10 mL). The mixture was extracted with ethyl acetate (3 x 10 mL) and the combined organic extracts were washed with brine (3 x 10 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by silica gel column with ethyl acetate/petroleum ether (1:2) as eluent to obtain the title compound as a yellow oil (100 mg, 86%).




Compound 260. 4-Chloro-5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-1H-pyrazole-3-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 4-chloro-5-iodo-1H-pyrazole-3-carbonitrile (compound **260.6**) was used in place of 3-iodo-4-methoxy-1H-pyrazole (compound **244.5**). *m/z* (ES⁺) 402 (M+H⁺).




Compound 261.1. (E)-4-Ethoxy-1,1,1-trifluoro-3-methylbut-3-en-2-one. Into a 50-mL 3-neck round-bottom flask, which was purged and maintained with an inert atmosphere


of nitrogen, was placed a solution trifluoroacetic anhydride (3.31 mL, 23.8 mmol) and diethyl ether (20 mL). The solution was cooled to -10 °C, then a solution mix of (*E*)-1-ethoxyprop-1-ene (2.82 mL, 25.5 mmol) and pyridine (1.94 mL, 24.0 mmol) was added drop-wise at -10°C. The resulting solution was stirred overnight at room temperature, then diluted with diethyl ether (50 mL). The solids were removed by filtration and the filtrate was concentrated under reduced pressure. The residue was diluted with water (100 mL) and extracted with DCM (3 x 50 mL). The combined organic extracts were washed with aq HCl (0.1 M, 3 x 100 mL) and water (100 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure to obtain the title compound as a colorless oil (2.6 g, 60%).




Compound 261.2. 4-Methyl-3-(trifluoromethyl)-1H-pyrazole. Into a 100-mL round-bottom flask, was placed (*E*)-4-ethoxy-1,1,1-trifluoro-3-methylbut-3-en-2-one (compound **261.1**, 2.6 g, 14.3 mmol) in EtOH (20 mL). Hydrazine hydrate (1.8 mL, 38 mmol) was added and the resulting solution was stirred for 2 h at 80 °C. The reaction was concentrated under reduced pressure to obtain the title compound as a yellow oil (2 g, crude).




Compound 261. 4-(1-(2,4-Dimethyl-5-(4-methyl-3-(trifluoromethyl)-1H-pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 4-methyl-3-(trifluoromethyl)-1H-pyrazole (compound **261.2**) was used in place of 4-methoxy-1H-pyrazole (compound **244.4**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). *m/z* (ES⁺) 439 (M+H⁺).




Compound 262.1. (Z)-4-(Dimethylamino)-3-(trifluoromethyl)but-3-en-2-one. Into a 100-mL round-bottom flask, was placed 4,4,4-trifluorobutan-2-one (1.7 mL, 15.86 mmol, 1.00 equiv) in toluene (45 mL). 1,1-Dimethoxy-*N,N*-dimethylmethanamine (4.21 mL, 31.7 mmol) was added and the resulting solution was stirred for 4 h at 100 °C, then concentrated under reduced pressure to obtain the title compound as yellow oil (2.0 g, crude).




Compound 262.2. 3-Methyl-4-(trifluoromethyl)-1H-pyrazole. Into a 50-mL round-bottom flask, was placed a solution of (Z)-4-(dimethylamino)-3-(trifluoromethyl)but-3-en-2-one (compound **262.1**, 2.0 g, 11.0 mmol) in ethanol (30 mL). Hydrazine hydrate (3.7 mL, 77 mmol) was added and the resulting solution was stirred overnight at 75 °C, then cooled and concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/petroleum ether (1:2) as eluent to obtain the title compound as a white solid (1.0 g, 61%).



Compound 262. 4-(1-(2,4-Dimethyl-5-(3-methyl-4-(trifluoromethyl)-1H-pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 3-methyl-4-(trifluoromethyl)-1H-pyrazole (compound **262.2**) was used in place of 4-methoxy-1H-pyrazole (compound **244.4**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). *m/z* (ES⁺) 439 (M+H⁺).



Compound 263.1. *N*-Methoxy-*N*-methyltetrahydrofuran-3-carboxamide. Into a 250-mL round-bottom flask, was placed a solution of tetrahydrofuran-3-carboxylic acid (10 g, 86.1 mmol) in dichloromethane (200 mL). EDC·HCl (18.2 g, 94.9 mmol), *N*,*O*-dimethylhydroxylamine hydrochloride (10 g, 103 mmol) and triethylamine (24.0 mL, 172 mmol) were added and the resulting mixture was stirred overnight at room temperature. The mixture was diluted with DCM (50 mL) and washed with aqueous HCl (1 M, 3 x 10 mL), sodium bicarbonate (sat., 2 x 30 mL) and brine (3 x 20 mL). The organic layer was dried (Na₂SO₄), filtered and concentrated under reduced pressure to obtain the title compound as light yellow oil (6.71 g, 49%).




Compound 263.2. 1-(Tetrahydrofuran-3-yl)propan-1-one. Into a 250-mL 3-neck round-bottom flask, was placed a solution of *N*-methoxy-*N*-methyltetrahydrofuran-3-carboxamide (compound **263.1**, 6.71 g, 42.2 mmol) in tetrahydrofuran (100 mL). The system was purged with nitrogen, then cooled to 0 °C. Ethylmagnesium bromide (3M in diethyl ether, 17 mL, 51 mmol) was added drop-wise at 0 °C over 30 min. The resulting solution was stirred overnight at 40 °C, then cooled and carefully quenched with aqueous HCl (1M, 10 mL). The resulting mixture was concentrated under reduced pressure and the residue was diluted with water (10 mL). The mixture was extracted with ethyl acetate (3 x 30 mL) and the combined organic extracts were washed with brine (2 x 10 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. Obtained the title compound as a light yellow oil (4.5 g, 83%).




Compound 263.3. 2-Methyl-3-oxo-3-(tetrahydrofuran-3-yl)propanal. Into a 100-mL round-bottom flask, was placed 1-(tetrahydrofuran-3-yl)propan-1-one (compound **263.2**, 2.5 g, 19.5 mmol), sodium methoxide (1.4 g, 25.9 mmol) and ethyl formate (20 mL). The mixture was stirred overnight at room temperature, then concentrated under reduced pressure.

The residue was purified by a silica gel column chromatography with ethyl acetate/petroleum ether (1:20 to 1:5) as eluent to obtain the title compound as a yellow solid (2.0 g, 66%).




Compound 263.4. 4-Methyl-5-(tetrahydrofuran-3-yl)-1H-pyrazole. Into a 100-mL

5 round-bottom flask, was placed a solution of 2-methyl-3-oxo-3-(tetrahydrofuran-3-yl)propanal (compound **263.3**, 3.00 g, 19.2 mmol) in ethanol (25 mL). Hydrazine hydrate (1.16 mL, 24 mmol) was added and the resulting solution was stirred for 3 h at 80 °C. The resulting mixture was concentrated under reduced pressure to obtain the title compound as a colorless oil (3.5 g, crude).




10 **Compound 263. 4-(1-(4-Methyl-3-(4-methyl-3-(tetrahydrofuran-3-yl)-1H-pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 4-methyl-5-(tetrahydrofuran-3-yl)-1H-pyrazole (compound **263.4**) was used in place of 4-methoxy-1H-pyrazole (compound **244.4**). m/z (ES⁺) 427 (M+H⁺).




20 **Compound 264.1. 1-Cyclopropylpropan-1-one.** Into a 250-mL 3-neck round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed cyclopropanecarbonitrile (20.0 g, 22.0 mL, 298 mmol) in diethyl ether (100 mL). The solution was cooled to 10 °C, then ethylmagnesium bromide (3 M in diethyl ether, 109 mL, 327 mmol) was added drop-wise over 10 min. The resulting solution was stirred overnight at 40 °C, then cooled and carefully quenched with aqueous NH₄Cl (sat., 100 mL). The layers were partitioned and the aqueous phase was extracted with ethyl acetate (3 x 200 mL). The combined organic layers were washed with brine (3 x 200 mL), dried (Na₂SO₄), filtered and


concentrated under reduced pressure. The crude product was purified by distillation under reduced pressure (50 mm Hg) and the fraction collected at 50 °C was the title compound (11 g, 38%).



5 **Compound 264. 4-(1-(3-(3-Cyclopropyl-4-methyl-1H-pyrazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **263**, except 1-cyclopropylpropan-1-one (compound **264.1**) was used in place of 1-(tetrahydrofuran-3-yl)propan-1-one (compound **263.2**). m/z (ES⁺) 397 (M+H)⁺.




10 **Compound 265. 4-(1-(5-(3-Cyclopropyl-4-methyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile.** The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **263**, except 1-cyclopropylpropan-1-one (compound **264.1**) was used in place of 1-(tetrahydrofuran-3-yl)propan-1-one (compound **263.2**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES⁺) 411 (M+H)⁺.




20 **Compound 266.1. 2-Cyclopropyl-3-oxobutanenitrile.** Into a 100-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of argon, was placed a solution of 2-cyclopropylacetonitrile (3.70 mL, 40.1 mmol) in THF (20 mL).

The solution was cooled to -78 °C, then LDA (2 M in THF, 30 mL, 60 mmol) was added drop-wise and the mixture was stirred for 1 h at -78 °C. A solution of acetic anhydride (4.54 mL, 48.0 mmol) in tetrahydrofuran (10 mL) was added drop-wise at -78 °C. The mixture was stirred for 1 h at 15 °C, then carefully quenched with citric acid (10% aq, 200 mL). The mixture was extracted with diethyl ether (2 x 70 mL) and the combined organic extracts were washed with brine (50 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure to obtain the title compound, which was used in the next step without purification.




Compound 266.2. 4-Cyclopropyl-3-methyl-1H-pyrazol-5-amine. Into a 250-mL round-bottom flask, was placed a solution of 2-cyclopropyl-3-oxobutanenitrile (compound **266.1**, 4.0 g, 32.5 mmol) in ethanol (18 mL). AcOH (3 mL) and hydrazine hydrate (3.15 mL, 65 mmol) were added and the reaction was stirred overnight at 80 °C. The reaction was cooled, then the pH was carefully adjusted to 9 with NaHCO₃ (aq. sat.). The mixture was diluted with water, then extracted with ethyl acetate (6 x 60 mL) and the combined organic extracts were washed with brine (80 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure to obtain the title compound as a brown oil (4.17 g, crude), which was used in the next step without purification.




Compound 266.3. 4-Cyclopropyl-5-iodo-3-methyl-1H-pyrazole. Into a 250-mL 3-neck round-bottom flask, was placed a solution of 4-cyclopropyl-3-methyl-1H-pyrazol-5-amine (compound **266.2**, 3.8 g, 27.7 mmol) in ACN (40 mL). The solution was cooled to 0 °C, then AcOH (4.8 mL, 83 mmol) was carefully added followed by a drop-wise addition of a solution of KI (11.5 g, 69.3 mmol) in water (40 mL). The mixture was stirred for 10 min at 0 °C, then tert-butyl nitrite (8.24 mL, 69.3 mmol) in ACN (40 mL) was added drop-wise. The resulting mixture was stirred at 0 °C for 10 min, then at room temperature for 2 h. The pH of the reaction was carefully adjusted to 8 with sodium bicarbonate (aq. sat.) and extracted with ethyl acetate (4 x 50 mL). The combined organic extracts were washed with Na₂S₂O₃ (aq.


sat., 80 mL) and brine (2 x 80 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/petroleum ether (1:2) as eluent to obtain the title compound as a yellow solid (1.71 g, 25%).




Compound 266. 4-(1-(5-(4-Cyclopropyl-3-methyl-1*H*-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 4-cyclopropyl-5-iodo-3-methyl-1*H*-pyrazole (compound **266.3**.) was used in place of 3-iodo-4-methoxy-1*H*-pyrazole (compound **244.5**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). *m/z* (ES⁺) 411 (M+H)⁺.




Compound 267.1. Methyl 4-cyclopropyl-2-methylbenzoate. To a solution of methyl 4-bromo-2-methylbenzoate (5.00 g, 20.7 mmol, 1.00 equiv, 95%) in a mixture of toluene and H₂O (20 mL/1 mL) were added potassium carbonate (6.10 g, 44.1 mmol, 2.00 equiv), cyclopropylboronic acid (2.30 g, 26.8 mmol, 1.20 equiv), Pd(dppf)Cl₂ (900 mg, 1.23 mmol, 0.05 equiv), and Pd(OAc)₂ (250 mg, 1.12 mmol, 0.05 equiv). The reaction mixture was purged with nitrogen and stirred at 80 °C overnight. After cooling to room temperature, the mixture was then concentrated under reduced pressure. The resulting residue was purified via silica gel column chromatography with ethyl acetate/petroleum ether (1:50) as eluent to yield 2.68 g (61%) of methyl 4-cyclopropyl-2-methylbenzoate as a colorless oil.



Compound 267.2. Methyl 4-cyclopropyl-5-iodo-2-methylbenzoate. To a solution of methyl 4-cyclopropyl-2-methylbenzoate (compound **267.1**, 2.68 g, 13.4 mmol, 1.00 equiv, 95%) in AcOH (50 mL) were added NaIO₄ (1.51 g, 7.08 mmol, 0.50 equiv), I₂ (3.58 g, 14.1 mmol, 1.00 equiv), and sulfuric acid (201 mg, 2.01 mmol, 0.15 equiv, 98%). The reaction mixture was stirred at 110 °C overnight. After cooling to ambient temperature, 100 mL of water was added. The resulting mixture was diluted with 100 mL of ethyl acetate, then washed with 3 x 30 mL of Na₂S₂O₃ (aq., sat.) and 1 x 30 mL of brine. The organic phase was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified via silica gel column chromatography with ethyl acetate/petroleum ether (1/50) as eluent to yield 2.00 g (45%) of methyl 4-cyclopropyl-5-iodo-2-methylbenzoate as a colorless oil.




Compound 267.3. Methyl 4-cyclopropyl-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5.4**, except methyl 4-cyclopropyl-5-iodo-2-methylbenzoate (compound **267.2**) was used in place of methyl 3-iodo-4-methylbenzoate (compound **5.3**).




Compound 267. 4-(1-(4-Cyclopropyl-5-(3,4-dimethyl-1H-pyrazol-5-yl)-2-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **59**, except methyl 4-cyclopropyl-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-

yl)benzoate (compound **267.3**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES+) 411 (M+H)⁺.




Compound 268.1. Ethyl 1-benzyl-4-methyl-1H-pyrazole-3-carboxylate. Into a

5 100-mL 3-neck round-bottom flask, was placed a solution of ethyl 4-methyl-1H-pyrazole-3-carboxylate (1.0 g, 6.5 mmol) in tetrahydrofuran (20 mL). The solution was cooled to 0 °C, then sodium hydride (360 mg, 9.00 mmol, 60% dispersion in mineral oil) was added in portions and the mixture was stirred for 20 min. Benzyl bromide (904 µL, 7.60 mmol) was added at 0 °C, then the resulting mixture was stirred for 1.5 h at room temperature. The
10 reaction was carefully quenched with by drop-wise addition of water (35 mL) and the mixture was extracted with ethyl acetate (3 x 100 mL). The combined organic extracts were washed with of brine (3 x 100 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/petroleum ether (1:20-1:2) as eluent to obtain the title compound as a yellow solid (1.0 g, 63%).




15 **Compound 268.2. (1-Benzyl-4-methyl-1H-pyrazol-3-yl)methanol.** Into a 100-mL 3-neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of ethyl 1-benzyl-4-methyl-1H-pyrazole-3-carboxylate (compound **268.1**, 2.3 g, 9.42 mmol) in THF (30 mL). The solution was cooled to 0 °C, then
20 LiAlH₄ (360 mg, 9.49 mmol, 1.00 equiv) was carefully added in portions. The resulting mixture was stirred for 1 h at room temperature, then carefully quenched by drop-wise addition of water (400 µL), then aqueous NaOH (10%, 400 µL), followed by additional water (1.2 mL). The solids were removed by filtration and the filtrate was dried (Na₂SO₄), filtered

and concentrated under reduced pressure to obtain the title compound as a yellow oil (1.7 g, 89%).




Compound 268.3. 1-Benzyl-3-(methoxymethyl)-4-methyl-1H-pyrazole. Into a 250-

mL 3-neck round-bottom flask, was placed a solution of (1-benzyl-4-methyl-1H-pyrazol-3-yl)methanol (compound **268.2**, 1.7 g, 8.4 mmol) in THF (50 mL). The solution was cooled to 0 °C, then sodium hydride (470 mg, 11.8 mmol, 60% dispersion in mineral oil) was added in portions at 0 °C, and then the mixture was stirred for 20 min. Iodomethane (750 µL, 12 mmol) was added at 0 °C, then the resulting mixture was stirred for 2 h at room temperature. The reaction was carefully quenched by drop-wise addition of water (30 mL) and the mixture was extracted with ethyl acetate (3 x 100 mL). The combined organic extracts were washed with brine (3 x 100 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/petroleum ether (1:10) as eluent to obtain the title compound as a yellow oil (1.5 g, 83%).




Compound 268.4. 3-(Methoxymethyl)-4-methyl-1H-pyrazole. Into a 100-mL

round-bottom flask, was placed a solution of 1-benzyl-3-(methoxymethyl)-4-methyl-1H-pyrazole (compound **268.3**, 700 mg, 3.24 mmol) in methanol (10 mL). The system was purged with nitrogen, then Pd(OH)₂/C (70 mg, 20 wt %) and aqueous HCl (2 M, 0.3 mL) were carefully added. The system was then charged with hydrogen (1 atm) and the resulting mixture was stirred for 2 days at room temperature. The system was purged with nitrogen, then the solids were removed by filtration and the filtrate was concentrated under reduced pressure. The pH of the residue was adjusted to 7 with aqueous sodium bicarbonate (sat.) and extracted with ethyl acetate (3 x 50 mL). The combined organic extracts were washed with brine (2 x 30 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure to obtain the title compound as a yellow oil (500 mg, crude).






Compound 268. 4-(1-(5-(3-(Methoxymethyl)-4-methyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 3-(methoxymethyl)-4-methyl-1H-pyrazole (compound **268.4**) was used in place 4-methoxy-1H-pyrazole (compound **244.4**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) m/z (ES+) 415 (M+H)⁺.



Compound 269. 4-(1-(3-(3,5-Dimethyl-1H-pyrazol-4-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 3-iodobenzoic acid was used in place of 3-iodo-4-methylbenzoic acid and 4-bromo-3,5-dimethyl-1H-pyrazole was used in place of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**). m/z (ES+) 357 (M+H)⁺.

The compounds in **TABLE 12** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **263**.



TABLE 12

Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
330	4-(1-(2,4-dimethyl-5-(4-methyl-3-(tetrahydrofuran-3-yl)-1H-pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		441
331	4-(1-(2,4-dimethyl-5-(4-methyl-3-(tetrahydro-2H-pyran-4-yl)-1H-pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		455
274	4-(1-(2,4-dimethyl-5-(4-methyl-3-(tetrahydrofuran-2-yl)-1H-pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		441

The compounds in **TABLE 13** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **268**.


TABLE 13


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
275	4-(1-(5-(3-((cyclopropylmethoxy)methyl)-4-methyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile		455
286	4-(1-(5-(3-(hydroxymethyl)-4-methyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile		401
309	4-(1-(2,4-dimethyl-5-(4-methyl-3-((oxetan-3-yloxy)methyl)-1H-pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		457
356	(5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2,4-dimethylphenyl)-4-methyl-1H-pyrazol-3-yl)methyl acetate		443


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
350	4-(1-(5-(3-(isopropoxymethyl)-4-methyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile		443
312	4-(1-(2,4-dimethyl-5-(4-methyl-3-((2,2,2-trifluoroethoxy)methyl)-1H-pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		483

The compounds in **TABLE 14** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **249**.


5 **TABLE 14**

Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
277	4-(1-(5-(3-(cyclopropylmethoxy)-4-methyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile		441


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
276	4-(1-(5-(3-(2-methoxyethoxy)-4-methyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile		445




Compound 279. 4-(1-(5-(3-Chloro-4-methoxy-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **244** and **250**. *m/z* (ES⁺) 421 (M+H)⁺.




Compound 281.1. 3-(Methoxymethyl)-1H-pyrazole. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **268.4**, except methyl 1H-pyrazole-3-carboxylate was used instead of methyl 4-methyl-1H-pyrazole-3-carboxylate.




Compound 281.2. 4-Chloro-3-(methoxymethyl)-1H-pyrazole. To a solution of 3-(methoxymethyl)-1H-pyrazole (compound **281.1**, 200 mg, 1.78 mmol) in chloroform (20 mL) was added NCS (237 mg, 1.77 mmol). The reaction mixture was stirred overnight at room temperature, then quenched with 20 mL of H₂O. The aqueous phase was extracted with 3 x 50 mL of ethyl acetate. The combined organic layers were washed with 3 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with EtOAc:PE (1:2) as the eluent to furnish 50 mg (19%) of the title compound as a white solid.



Compound 281.3. Methyl 5-(4-chloro-3-(methoxymethyl)-1H-pyrazol-5-yl)-2,4-dimethylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244.6**, except 4-chloro-3-(methoxymethyl)-1H-pyrazole (compound **281.2**) was used instead of 4-methoxy-1H-pyrazole (compound **244.4**).




Compound 281. 4-(1-(5-(4-Chloro-3-(methoxymethyl)-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except methyl 5-(4-chloro-3-(methoxymethyl)-1H-pyrazol-5-yl)-2,4-dimethylbenzoate (compound **281.3**) was used instead of methyl 3-(4-methoxy-1H-pyrazol-5-yl)-4-methylbenzoate (compound **244.6**). m/z (ES⁺) 435 (M+H)⁺.




Compound 304.1. 5-(3-(Bromomethyl)-4-chloro-1H-pyrazol-5-yl)-2,4-

dimethylbenzoic acid. A mixture of methyl 5-(4-chloro-3-(methoxymethyl)-1H-pyrazol-5-yl)-2,4-dimethylbenzoate (compound **281.3**, 20 mg, 0.06 mmol) and HBr-HOAc (5 mL) was stirred overnight at 80 °C, then concentrated under reduced pressure to yield 20 mg (90%) of the title compound as a yellow oil.




Compound 304.2. 5-(4-Chloro-3-(hydroxymethyl)-1H-pyrazol-5-yl)-2,4-

dimethylbenzoic acid. A mixture of 5-(3-(bromomethyl)-4-chloro-1H-pyrazol-5-yl)-2,4-dimethylbenzoic acid (compound **304.1**, 200 mg, 0.58 mmol) and NaHCO₃ (0.3 M, 10 mL) was stirred for 30 min at room temperature, then concentrated under reduced pressure. This resulted in 200 mg (crude) of the title compound as a white solid.




Compound 304. 4-(1-(5-(4-Chloro-3-(hydroxymethyl)-1H-pyrazol-5-yl)-2,4-


dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 5-(4-chloro-3-(hydroxymethyl)-1H-pyrazol-5-yl)-2,4-dimethylbenzoic acid (compound **304.2**) was used instead of 3-(4-methoxy-1H-pyrazol-5-yl)-4-methylbenzoic acid (compound **244.7**). *m/z* (ES⁺) 421 (M+H)⁺.



Compound 280.1. 4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole. Into a 250-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 4-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (5.82 g, 30.0 mmol) in tetrahydrofuran (80 mL). This was followed by the addition of NaH (70%) (2.05 g, 85.4 mmol) in portions at 0 °C. To this was added SEMCl (6.4 mL, 36.1 mmol) dropwise. The reaction mixture was stirred overnight at room temperature, then quenched with 50 mL of NH₄Cl (sat). The aqueous phase was extracted with 2 x 100 mL of ethyl acetate and the combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 7 g (72%) of the title compound as colorless oil.




Compound 280.2. 1-((2-(Trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-ol. To a solution of 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole (compound **280.1**, 9.7 g, 30.0 mmol) in THF (100 mL) was added 2M sodium hydroxide (aq.) (30 mL) dropwise. To this was added H₂O₂ (30%) (22.7 mL) dropwise at 0 °C. The reaction mixture was stirred for 2h at room temperature, then quenched with 50 mL of water. The aqueous phase was extracted with 2 x 150 mL of ethyl acetate and the combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 4.3 g (67%) of the title compound as a white solid.




Compound 280.3. 4-Methoxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole.

To a solution of 1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-ol (compound **280.2**, 2.14 g, 10.0 mmol) in tetrahydrofuran (50 mL) was added Cs₂CO₃ (6.5 g, 20.0 mmol) in portions. To this was added CH₃I (1.25 mL, 20.0 mmol) dropwise. The reaction mixture was stirred for 3h at room temperature, then concentrated under reduced pressure. The resulting residue was dissolved in 100 mL of EtOAc. The solids were filtered off and the filtrate was concentrated under reduced pressure. This resulted in 2.1 g (92%) of the title compound as a colorless oil.


**Compound 280.4. 4-Methoxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole-**

3-carboxylic acid. Into a 250-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 4-methoxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole (compound **280.3**, 4.6 g, 20.0 mmol) in tetrahydrofuran (50 mL). This was followed by the addition of n-BuLi (2.5 M in hexane) (10 mL) dropwise at -78 °C. The reaction mixture was stirred for 1h at -60 °C. To this was added CO₂(solid) (5 g) with stirring at -78 °C. The reaction mixture was warmed slowly to room temperature and was stirred for 1h, then quenched with 10 mL of NH₄Cl (sat.). The pH value of the solution was adjusted to 5-6 with HCl (1 M). The aqueous phase was extracted with 2 x 100 mL of ethyl acetate and the combined organic layers were dried over sodium sulfate and concentrated under reduced pressure. This resulted in 5 g (92%) of the title compound as a white solid.

**Compound 280.5. 4-Methoxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole-**


3-carboxamide. To a solution of 4-methoxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole-3-carboxylic acid (compound **280.4**, 5 g, 18.4 mmol) in *N,N*-dimethylformamide (50 mL) were added EDC·HCl (7 g, 36.5 mmol), NH₄Cl (1.97 g, 36.8 mmol) and 4-dimethylaminopyridine (8.95 g, 73.3 mmol). The mixture was stirred for 2h at room

temperature, then warmed to 50 °C for 3h. The reaction mixture was quenched with 30 mL of NH₄Cl (sat). The aqueous phase was extracted with 2 x 100 mL of ethyl acetate. The combined organic layers were washed with 3 x 100 mL of NH₄Cl (sat), dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:5) as the eluent to furnish 1.3 g (26%) of the title compound.




Compound 280.6. 4-Methoxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole-3-carbonitrile. To a solution of 4-methoxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-


pyrazole-3-carboxamide (compound **280.5**, 1.08 g, 4.0 mmol) in toluene (25 mL) was added POCl₃ (1.86 mL, 20.0 mmol) dropwise at 0 °C. The mixture was stirred for 2h at 50 °C, then quenched with 30 mL of sodium bicarbonate (sat.). The aqueous phase was extracted with 2 x 100 mL of ethyl acetate and the combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:6) as the eluent to yield 500 mg (49%) of the title compound as a white solid.




Compound 280.7. 4-Methoxy-1H-pyrazole-3-carbonitrile. A mixture of 4-methoxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole-3-carbonitrile (compound **280.6**, 500 mg, 1.97 mmol) and trifluoroacetic acid (10 mL) in TES (5 mL) was stirred for 2h at room temperature, then concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:2) as the eluent to furnish 210 mg (86%) of the title compound as a white solid.




Compound 280.8. 5-Iodo-4-methoxy-1H-pyrazole-3-carbonitrile. To a solution of 4-methoxy-1H-pyrazole-3-carbonitrile (compound **280.7**, 200 mg, 1.62 mmol) in *N,N*-dimethylformamide (15 mL) was added NIS (549 mg, 2.44 mmol). The reaction mixture was stirred for 2h at 70 °C, then quenched with 20 mL of Na₂S₂O₃ (sat.). The aqueous phase was extracted with 2 x 50 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:6) as the eluent to furnish 330 mg (82%) of the title compound as a white solid.




Compound 280. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2,4-dimethylphenyl)-4-methoxy-1H-pyrazole-3-carbonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 5-iodo-4-methoxy-1H-pyrazole-3-carbonitrile (compound **280.8**) was used instead of 4-chloro-5-iodo-1H-pyrazole-3-carbonitrile (compound **260.6**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). *m/z* (ES⁺) 412 (M+H⁺).




Compound 294.1. (E)-3-(Dimethylamino)-1-(tetrahydro-2H-pyran-4-yl)prop-2-en-1-one. To a sealed tube were added 1-(tetrahydro-2H-pyran-4-yl)ethanone (3.00 g, 23.4 mmol) and DMF-DMA (6.2 mL, 46.8 mmol). The reaction mixture was heated at 100 °C overnight. The resulting mixture was concentrated under reduced pressure to afford 4.00 g (crude) of the title compound as light yellow oil.



Compound 294.2. 3-(Tetrahydro-2H-pyran-4-yl)-1H-pyrazole. To a solution of (E)-3-(dimethylamino)-1-(tetrahydro-2H-pyran-4-yl)prop-2-en-1-one (compound **294.1**, 4.3 g, 23.4 mmol) in ethanol (20 mL) was added $\text{NH}_2\text{NH}_2 \cdot \text{H}_2\text{O}$ (1.41 g, 28.2 mmol). The reaction mixture was stirred for 3 h at 80 °C. The resulting mixture was concentrated under reduced pressure to yield 4.45 g (crude) of the title compound as a light yellow solid.




Compound 294.3. 4-Chloro-3-(tetrahydro-2H-pyran-4-yl)-1H-pyrazole. To a solution of 3-(tetrahydro-2H-pyran-4-yl)-1H-pyrazole (compound **294.2**, 3.5 g, 23.0 mmol) in tetrahydrofuran (20 mL) was added NCS (3.4 g, 25.4 mmol). The reaction mixture was stirred overnight at 40 °C. The resulting mixture was concentrated under reduced pressure. This resulted in 4.75 g (crude) of the title compound as a light yellow solid.




Compound 294.4. 4-Chloro-5-iodo-3-(tetrahydro-2H-pyran-4-yl)-1H-pyrazole.


To a solution of 4-chloro-3-(tetrahydro-2H-pyran-4-yl)-1H-pyrazole (compound **294.3**, 3.00 g, 16.1 mmol) in *N,N*-dimethylformamide (20 mL) was added NIS (7.26 g, 32.3 mmol). The reaction mixture was stirred overnight at 80 °C. The reaction mixture was diluted with 50 mL $\text{Na}_2\text{S}_2\text{O}_3$ (sat.). The aqueous phase was extracted with 3 x 100 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/8-1/5) as the eluent to furnish 4.05 g (81%) of the title compound as a white solid.



Compound 294. 4-(1-(5-(4-Chloro-3-(tetrahydro-2H-pyran-4-yl)-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 4-chloro-5-iodo-3-(tetrahydro-2H-pyran-4-yl)-1H-pyrazole (compound **294.4**) was used instead of was used in place 4-methoxy-1H-pyrazole (compound **244.4**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES+) 475 (M+H)⁺.




Compound 293.1. 4-Chloro-3-cyclopropyl-5-iodo-1H-pyrazole. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **294.4**, except 1-cyclopropylethanone was used instead of 1-(tetrahydro-2H-pyran-4-yl)ethanone.



Compound 293. 4-(1-(5-(4-Chloro-3-cyclopropyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **294**, except 4-chloro-3-cyclopropyl-5-iodo-1H-pyrazole (compound **293.1**) was


used instead of 4-chloro-5-iodo-3-(tetrahydro-2H-pyran-4-yl)-1H-pyrazole (compound **294.4**). m/z (ES⁺) 431(M+H)⁺.



Compound 305.1. *N,N*,4-Trimethyl-1H-pyrazole-1-sulfonamide. Into a 500-mL

5 round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 4-methyl-1H-pyrazole (10.4 g, 127.2 mmol) in CH₂Cl₂ (200 mL). *N,N*-dimethylsulfamoyl chloride (17.14 mL, 167.1 mmol) and triethylamine (35 mL, 251.1 mmol) were added to the reaction. The reaction mixture was stirred for 30 min at 0 °C in a water/ice bath and then stirred overnight at room temperature. The reaction was quenched with

10 NaHCO₃ (sat.) (40 mL) and the aqueous phase was extracted with 3 x 200 mL of EtOAc. The combined organic phase was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/hexanes (1:6) as the eluent to afford the title compound (22.2 g, 92%) as a light yellow oil.




Compound 305.2. 3-(2-Hydroxyethyl)-*N,N*,4-trimethyl-1H-pyrazole-1-

sulfonamide. Into a 50-mL three-neck round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed a solution of *N,N*,4-trimethyl-1H-pyrazole-1-sulfonamide (compound **305.1**, 1.19 g, 6.26 mmol) in THF (24 mL). *n*-BuLi (3 mL, 7.5


20 mmol, 2.5 M in hexanes) was added to the reaction dropwise at -78 °C. The resulting mixture was stirred for 30 min at -78 °C. This was followed by the addition of a solution of oxirane (5.6 mL, 113.51 mmol) in THF (4 mL) at -78 °C. The reaction mixture was then brought to room temperature and stirred for 3h. The reaction was quenched with 10 mL of NH₄Cl (sat.) and the aqueous phase was extracted with 3 x 50 mL of EtOAc. The combined organic layers

were dried over Na_2SO_4 . The solvent was removed to give the crude product which was purified by silica gel chromatography with hexanes/EtOAc (3:1) as the eluent to furnish 636 mg (44%) of the title compound as light yellow oil.




Compound 305.3. 3-(2-Methoxyethyl)-N,N,4-trimethyl-1H-pyrazole-1-

sulfonamide. Into a 50-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 3-(2-hydroxyethyl)-N,N,4-trimethyl-1H-pyrazole-1-sulfonamide (compound **305.2**, 636 mg, 2.73 mmol) in THF (10 mL). Sodium hydride (131 mg, 60% in mineral oil, 3.27 mmol) was added at 0 °C and stirred for 30 min at 0 °C. This was followed by the addition of CH_3I (204 μL , 3.28 mmol). The reaction mixture was stirred for 3h at room temperature. The reaction was quenched with 10 mL of NH_4Cl (sat.) and the aqueous phase was extracted with 3 x 50 mL of EtOAc. The combined organic phases were dried over Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure to give the crude product which was purified by silica gel chromatography with hexanes/EtOAc (15:1-8:1) as the eluent to give 452 mg (67%) of the title compound as a yellow oil.




Compound 305.4. 3-(2-Methoxyethyl)-4-methyl-1H-pyrazole. Into a 50-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 3-(2-methoxyethyl)-N,N,4-trimethyl-1H-pyrazole-1-sulfonamide (compound **305.3**, 440 mg, 1.78 mmol) in dioxane (10 mL). HCl (10 mL, 6 M, 30 mmol) was added to the reaction mixture. The reaction mixture was stirred for 1h at 85 °C, then cooled to room temperature and carefully neutralized with NaHCO_3 aq (sat.) to pH~8. The aqueous phase


was extracted with 3 x 50 mL of EtOAc and the combined organic layers were dried over Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure to give the crude product which was used in the next step without further purification.




Compound 305.5. 5-Iodo-3-(2-methoxyethyl)-4-methyl-1H-pyrazole. Into a 50-mL round-bottom flask (1 atm) purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 3-(2-methoxyethyl)-4-methyl-1H-pyrazole (compound **305.4**, 249 mg, 1.78 mmol) in *N,N*-dimethylformamide (10 mL). NIS (482 mg, 2.14 mmol, 1.20 equiv) was added to the reaction. The reaction mixture was stirred for 2h at 60 °C, then cooled and quenched with 5 mL of NH₄Cl (sat.). The aqueous phase was extracted with 3 x 30 mL of EtOAc and the combined organic layers were dried over Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure to give the crude product which was purified by silica gel chromatography with hexanes/EtOAc (5:1) as the eluent to yield 328 mg (69% over two steps) of the title compound as a brown oil.




Compound 305. 4-(1-(5-(3-(2-Methoxyethyl)-4-methyl-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 5-iodo-3-(2-methoxyethyl)-4-methyl-1H-pyrazole (compound **305.5**) was used instead of 3-iodo-4-methoxy-1H-pyrazole (compound **244.5**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). *m/z* (ES⁺) 429 (M+H)⁺.




Compound 307.1. 3-(2-Methoxyethyl)-1H-pyrazole. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **305.4**, except 1H-pyrazole was used instead of 4-methyl-1H-pyrazole.



Compound 307.2. 4-Chloro-5-iodo-3-(2-methoxyethyl)-1H-pyrazole. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **294.4**, except 3-(2-methoxyethyl)-1H-pyrazole (compound **307.1**) was used instead of 3-(tetrahydro-2H-pyran-4-yl)-1H-pyrazole (compound **294.2**).




Compound 307. 4-(1-(5-(4-Chloro-3-(2-methoxyethyl)-1H-pyrazol-5-yl)-2,4-dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 4-chloro-5-iodo-3-(2-methoxyethyl)-1H-pyrazole (compound **307.2**) was used instead of 3-iodo-4-methoxy-1H-pyrazole (compound **244.5**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES⁺) 449 (M+H)⁺.




Compound 357.1. (Z)-Methyl 2-((dimethylamino)methylene)-3-oxobutanoate.

Into a 250-mL round-bottom flask, was placed methyl 3-oxobutanoate (10.8 g, 92.6 mmol) and (dimethoxymethyl)dimethylamine (12.7 mL, 95.3 mmol). The resulting solution was stirred for 2.5h at 80 °C, then concentrated under reduced pressure to yield 14.5 g (crude) of the title compound as a red solid.



Compound 357.2. Methyl 1-benzyl-3-methyl-1H-pyrazole-4-carboxylate.


To a solution of (Z)-methyl 2-((dimethylamino)methylene)-3-oxobutanoate (15.8 g, 92.5 mmol) in ethanol (50 mL) was added 1-benzylhydrazine dihydrochloride (18 g, 92.3 mmol). The resulting solution was stirred for 3.5h at 80 °C, then concentrated under reduced pressure. The residue was dissolved in 200 mL of ethyl acetate. The organic layer was washed with 1 x 40 mL of water and 2 x 40 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:4) as the eluent to furnish 12.4 g (58%) of the title compound as a yellow solid.




Compound 357. 4-(1-(5-(4-(Methoxymethyl)-3-methyl-1H-pyrazol-5-yl)-2,4-

dimethylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **268**, except methyl 1-benzyl-3-methyl-1H-pyrazole-4-carboxylate (compound


357.2) was used instead of ethyl 1-benzyl-4-methyl-1*H*-pyrazole-3-carboxylate (compound 268.1). m/z (ES⁺) 415 (M+H)⁺.



Compound 314.1. 1-Benzyl-3-methyl-1*H*-pyrazole. To a solution of 3-methyl-1*H*-pyrazole (5.00 g, 60.9 mmol) in tetrahydrofuran (30 mL) was added sodium hydride (60%) (4.88 g, 203 mmol) in portions at 0 °C and stirred for 30 min under nitrogen. To the above mixture was added benzyl bromide (8.7 mL, 73.1 mmol). The reaction mixture was stirred for 1h at room temperature, then carefully quenched with 150 mL of water. The aqueous phase was extracted with 3 x 30 mL of EtOAc. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford 5.66 g (54%) of the title compound as light yellow oil.




Compound 314.2. 1-Benzyl-3-methyl-1*H*-pyrazole-4-carbaldehyde. To a solution of 1-benzyl-3-methyl-1*H*-pyrazole (compound 314.1, 1.00 g, 5.81 mmol) in *N,N*-dimethylformamide (3 mL) was added POCl₃ (493 μL, 5.28 mmol) dropwise. The reaction mixture was stirred for 3 h at 90 °C under nitrogen. The reaction mixture was diluted with 50 mL of H₂O. The pH of the solution was adjusted to 7 with sodium hydroxide (2 M). The aqueous phase was extracted with 3 x 20 mL of ethyl acetate and the combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure to yield 1.42 g (crude) of the title compound as a brown oil.




Compound 314.3. 1-(1-Benzyl-3-methyl-1*H*-pyrazol-4-yl)-2,2,2-trifluoroethanol. To a solution of 1-benzyl-3-methyl-1*H*-pyrazole-4-carbaldehyde (compound 314.2, 1.20 g, 5.99 mmol) in tetrahydrofuran (15 mL) was added CF₃SiMe₃ (1.33 mL, 9.01 mmol) and TBAF (0.11 mL, 0.11 mmol, 1M solution in THF). The mixture was stirred for 10 min at

room temperature under nitrogen. This was followed by the addition of aqueous HCl (6 M) (1mL). The reaction mixture was stirred for another 10 min at room temperature, then diluted with 50 mL of H₂O. The aqueous phase was extracted with 3 x 20 mL of ethyl acetate and the combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/5-1/2) as the eluent to furnish 1.15 g (71%) of the title compound as a light yellow solid.




Compound 314.4. O-(1-(1-Benzyl-1H-pyrazol-4-yl)-2,2,2-trifluoroethyl) O-phenyl

carbonothioate. To a solution of 1-(1-benzyl-3-methyl-1H-pyrazol-4-yl)-2,2,2-trifluoroethanol (compound **314.3**, 1.15 g, 4.26 mmol) in toluene (15 mL) were added 4-dimethylaminopyridine (1.47 g, 12.0 mmol) and phenyl chloromethanethioate (1.04 g, 6.02 mmol). The reaction mixture was stirred for 1 h at 60 °C under nitrogen, then concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/15) as the eluent to yield 1.34 g (77%) of the title compound as a light yellow oil.




Compound 314.5. 1-Benzyl-3-methyl-4-(2,2,2-trifluoroethyl)-1H-pyrazole.


To a solution of O-(1-(1-benzyl-1H-pyrazol-4-yl)-2,2,2-trifluoroethyl) O-phenyl carbonothioate (compound **314.4**, 1.34 g, 3.30 mmol) in toluene (20 mL) was added Bn₃SnH (3.84 mL, 13.2 mmol), AIBN (320 mg, 1.95 mmol). The reaction mixture was stirred for 1 h at 80 °C under nitrogen, then concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:20) as the eluent to furnish 650 mg (78%) of the title compound as a colorless oil.



Compound 314.6. 3-Methyl-4-(2,2,2-trifluoroethyl)-1H-pyrazole. To a solution of 1-benzyl-3-methyl-4-(2,2,2-trifluoroethyl)-1H-pyrazole (compound **314.5**, 700 mg, 2.75 mmol) in methanol (10 mL) was added palladium on carbon (700 mg, 10 wt. %) and HCl (4 M) (3 mL). The reaction mixture was hydrogenated for 48h at room temperature under 1 atmosphere of hydrogen. The solids were removed via filtration. The filtrate was evaporated under reduced pressure and diluted with 50 mL of water. The pH of the solution was adjusted to 7 with NaOH (2 M). The aqueous phase was extracted with 5 x 20 mL of ethyl acetate and the combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford 230 mg (51%) of the title compound as a light yellow oil.




Compound 314. 4-(1-(2,4-Dimethyl-5-(3-methyl-4-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **244**, except 3-methyl-4-(2,2,2-trifluoroethyl)-1H-pyrazole (compound **314.6**) was used instead of 4-methoxy-1H-pyrazole (compound **244.4**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). m/z (ES⁺) 453 (M+H)⁺.




Compound 313.1. 1-Benzyl-4-methyl-1H-pyrazole-3-carbaldehyde. To a solution of (1-benzyl-4-methyl-1H-pyrazol-3-yl)methanol (compound **268.2**, 3 g, 14.8 mmol) in dichloromethane (100 mL) was added PDC (11.3 g, 30.1 mmol), molecular sieves (3 g) and


NaOAc (300 mg) were added to the reaction mixture. The reaction mixture was stirred for 4h at room temperature. The solids were removed by filtration and the filtrate was concentrated under reduced pressure. The residue was purified by a silica gel chromatography with ethyl acetate/petroleum ether (1/10) as the eluent to yield 1 g (34%) of the title compound as a colorless oil.




Compound 313. 4-(1-(2,4-Dimethyl-5-(4-methyl-3-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **314**, except 1-benzyl-4-methyl-1H-pyrazole-3-carbaldehyde (compound **313.1**) was used instead of 1-benzyl-3-methyl-1H-pyrazole-4-carbaldehyde (compound **314.2**). m/z (ES⁺) 453 (M+H)⁺.




Compound 374.1. N,N,4-Trimethyl-1H-pyrazole-3-carboxamide. To a mixture of 4-methyl-1H-pyrazole-3-carboxylic acid (1.00 g, 7.93 mmol), HOBT (1.07g, 7.93 mmol) and EDCI (2.27 g, 11.89 mmol) in DMF (10 mL) was added DIEA (2.74 mL, 15.86 mmol). After the mixture was stirred at room temperature for 16 hours, dimethylamine (2M in THF, 11.89 mL, 23.79 mmol) was added. The mixture was stirred at room temperature for 3 hours, then partitioned between EtOAc (300 mL) and water (30 mL). The organics was washed with brine (3 x 30 mL) and the combined aqueous material was back extracted with EtOAc (2 x 50 mL). All organic extracts were combined, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (5% MeOH in CH₂Cl₂) to obtain the title compound as a brown oil (1.1 g, 92%). m/z (ES⁺) 154 (M+H)⁺.



Compound 374.2. 5-Iodo-N,N,4-trimethyl-1H-pyrazole-3-carboxamide. To a solution of N,N,4-trimethyl-1H-pyrazole-3-carboxamide (compound **374.1**, 0.98 g, 6.40 mmol) in CH₃CN (10 mL) was added NIS (1.59g, 7.04 mmol). The mixture was stirred at room temperature for 16 hours then quenched with sat. Na₂SO₃ solution (5mL). The mixture was partitioned between EtOAc (200 mL) and water (30 mL). The EtOAc layer was washed with brine (30 mL), dried (MgSO₄) and concentrated. The residue was purified with column chromatography (50% EtOAc in hexanes to 100% EtOAc) to give the product as a brown solid (1.79 g, 100%). *m/z* (ES⁺) 280 (M+H)⁺.




Compound 374. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2,4-dimethylphenyl)-N,N,4-trimethyl-1H-pyrazole-3-carboxamide. The title compound was prepared using the standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 5-iodo-N,N,4-trimethyl-1H-pyrazole-3-carboxamide (compound **374.2**) was used instead of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**) and methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). *m/z* (ES⁺) 442 (M+H)⁺.




Compound 375. 5-(5-(3-Fluoro-3-(p-tolyl)azetidine-1-carbonyl)-2,4-dimethylphenyl)-N,N,4-trimethyl-1H-pyrazole-3-carboxamide. The title compound was prepared using the standard chemical manipulations and procedures similar to those used for


the preparation of compound **5**, except 5-iodo-*N,N*,4-trimethyl-1*H*-pyrazole-3-carboxamide (compound **374.2**) was used instead of 5-iodo-2,4-dimethyl-1*H*-imidazole (compound **5.5**), methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) and 4-(3-fluoroazetidin-3-yl)benzonitrile hydrochloride (compound **43.4**) was used in place of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 460 (M+H)⁺.




Compound 329. (3-(2,4-Dimethyl-1*H*-imidazol-5-yl)-4-methylphenyl)(4-(4-methoxyphenyl)piperidin-1-yl)methanone. The title compound was prepared using the standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 4-(4-methoxyphenyl)piperidine hydrochloric acid salt was used instead of 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**). m/z (ES+) 404 (M+H)⁺.



Compound 284.1. Methyl 3-(2-(hydroxymethyl)-4-(trifluoromethyl)-1*H*-imidazol-5-yl)-4-methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **219.4**, except methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used instead of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**).



Compound 284.2. 3-(2-(Hydroxymethyl)-4-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoic acid. Into a 100-mL round-bottom flask, was placed methyl 3-(2-(hydroxymethyl)-4-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoate (compound **284.1**,
 5 2 g, 6.36 mmol), HBr (40% in AcOH) (50 mL). The reaction mixture was stirred for 12h at 80 °C. The crude product (100 mL) was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): mobile phase, CH₃CN:H₂O=1:1; Detector, UV 254 nm to yield 1 g (52%) of the title compound as a light yellow solid.




Compound 284. 4-(1-(3-(2-(Hydroxymethyl)-4-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 3-(2-(hydroxymethyl)-4-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **284.2**) was used in place of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **5.7**). *m/z* (ES⁺) 441 (M+H)⁺.


The compounds in **TABLE 15** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **284**.

TABLE 15


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
285	4-(3-fluoro-1-(3-(2-(hydroxymethyl)-4-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		459
352	4-(1-(3-(2-(2-hydroxyethyl)-4-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		455
362	4-(1-(3-(2-(2-methoxyethyl)-4-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		468
360	4-(1-(3-(2-(methoxymethyl)-4-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		455




Compound 287.1. 5-Iodo-2-(tetrahydro-2H-pyran-4-yl)-4-(trifluoromethyl)-1H-imidazole. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **202.2**, except tetrahydro-2H-pyran-4-carbaldehyde was used instead of 3-methyloxetane-3-carbaldehyde.



Compound 287.2. Methyl 4-methyl-3-(2-(tetrahydro-2H-pyran-4-yl)-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **202.4**, except 5-iodo-2-(tetrahydro-2H-pyran-4-yl)-4-(trifluoromethyl)-1H-imidazole (compound **287.1**) was used instead of 5-iodo-2-(3-methyloxetan-3-yl)-1H-imidazole-4-carbonitrile (compound **202.3**).



Compound 287.3. 4-Methyl-3-(2-(tetrahydro-2H-pyran-4-yl)-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoic acid. A solution of methyl 4-methyl-3-(2-(tetrahydro-2H-pyran-4-yl)-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoate (compound **287.2**, 50 mg, 0.14 mmol) in HCl (6 M) (2 mL) was stirred for 5h at 80 °C, then concentrated under reduced pressure. This resulted 200 mg (crude) of the title compound as a brown solid which was used in the next step without further purification.





Compound 287. 4-(1-(4-Methyl-3-(2-(tetrahydro-2H-pyran-4-yl)-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to




those used for the preparation of compound **202**, except 4-methyl-3-(2-(tetrahydro-2H-pyran-4-yl)-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoic acid (compound **287.3**) was used instead of 3-(4-cyano-2-(3-methyloxetan-3-yl)-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **202.5**). m/z (ES⁺) 495 (M+H)⁺.

5

The compounds in **TABLE 16** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **287** and **202**.

TABLE 16


Cpd	Name	Structure	m/z (ES ⁺) (M+H) +
288	4-(1-(4-methyl-3-(2-(4-methyltetrahydro-2H-pyran-4-yl)-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		509
290	4-(1-(4-methyl-3-(2-(1-methylpiperidin-4-yl)-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		508

Cpd	Name	Structure	m/z (ES+) (M+H) +
291	methyl 4-(5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl)piperidine-1-carboxylate		552
292	4-(1-(4-methyl-3-(2-(1-methylpyrrolidin-3-yl)-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		494
289	4-(1-(4-methyl-3-(2-(3-methyloxetan-3-yl)-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		481

The compounds in **TABLE 17** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **230** and **231**.

TABLE 17


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
353	4-(1-(3-(4-chloro-2-(2-hydroxyethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		421
354	4-(1-(3-(4-chloro-2-(2-hydroxyethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile		439
359	4-(1-(3-(4-chloro-2-(methoxymethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		421
361	4-(1-(3-(4-chloro-2-(2-methoxyethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		435




Compound 283. (4-Chloro-5-(5-(3-(4-cyanophenyl)-3-fluoroazetidine-1-carbonyl)-2-methylphenyl)-1H-imidazol-2-yl)methyl acetate. Into a 50-mL round-bottom flask, was placed a solution of 4-(1-(3-(4-chloro-2-(hydroxymethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)piperidin-4-yl)benzonitrile (compound **29**, 40 mg, 0.09 mmol) in *N,N*-dimethylformamide (3 mL). Triethylamine (14 μ L, 0.10 mmol) was added to the reaction at 0 °C, followed by acetic anhydride (10 μ L, 0.10 mmol). The reaction mixture was stirred for 2h at room temperature, then diluted with 10 mL of ethyl acetate. The organic layer was washed with 3 x 5 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The crude product (35 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-020): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*100 mm; mobile phase, WATER WITH 0.05% TFA and MeCN (10% MeCN up to 70% in 6 min, up to 95% in 1 min, down to 10% in 1 min); Detector, Waters 2489 254 & 220 nm. This resulted in 15 mg (34%) of the title compound as a white solid. *m/z* (ES⁺) 467 (M+H)⁺.

The compounds in **TABLE 18** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **185**, **192** and **194**.


TABLE 18

Cpd	Name	Structure	<i>m/z</i> (ES ⁺) (M+H) +
344	methyl 4-(5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-4-methyl-1H-imidazol-2-yl)-4-hydroxypiperidine-1-carboxylate		514

Cpd	Name	Structure	m/z (ES+) (M+H) +
345	methyl 4-(5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-4-methyl-1H-imidazol-2-yl)-4-methoxypiperidine-1-carboxylate		528

The compounds in **TABLE 19** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **185** and **241**.

5 **TABLE 19**






Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
346	4-(1-(3-(4-chloro-2-(4-methyltetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		475

Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
347	4-(1-(3-(4-chloro-2-(4-methyltetrahydro-2H-pyran-4-yl)-1H-imidazol-5-yl)-4-methylbenzoyl)-3-fluoroazetidin-3-yl)benzonitrile		494
348	methyl 4-(4-chloro-5-(5-(3-(4-cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-1H-imidazol-2-yl)-4-methylpiperidine-1-carboxylate		532

The compounds in **TABLE 20** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **14**.

5 **TABLE 20**

Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
358	4-(3-fluoro-1-(4-methyl-3-(4-methyl-2-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		443


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
366	4-(1-(2,4-dimethyl-5-(4-methyl-2-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		439
367	4-(1-(2,4-dimethyl-5-(4-methyl-2-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)-3-fluoroazetidin-3-yl)benzonitrile		457
363	4-(1-(3-(4-ethyl-2-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		439
364	4-(1-(3-(4-(methoxymethyl)-2-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		455
365	4-(1-(3-(4-cyclopropyl-2-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile		451

The compounds in **TABLE 21** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **218**.

TABLE 21


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
368	4-(3-fluoro-1-(4-methyl-3-(2-methyl-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		443
369	4-(1-(2,4-dimethyl-5-(2-methyl-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		439
370	4-(1-(2,4-dimethyl-5-(2-methyl-4-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)-3-fluoroazetidin-3-yl)benzonitrile		457

5




Compound 282.1. *N,N*,2-Trimethyl-1H-imidazole-1-sulfonamide. To a solution of 2-methyl-1H-imidazole (10 g, 121.8 mmol) in dichloromethane (100 mL) was added *N,N*-dimethylsulfamoyl chloride (14.4 mL, 133.7 mmol) and triethylamine (34 mL, 244.1 mmol).


The reaction mixture was stirred overnight at room temperature, then quenched with 10 mL of water and extracted with 1 x 50 mL of DCM. The organic layer was washed with 2 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:2) as the eluent to furnish 20 g (87%) of the title compound as a yellow oil.




Compound 282.2. 4-(2-Hydroxyethyl)-*N,N*,2-trimethyl-1H-imidazole-1-sulfonamide. Into a 250-mL three neck round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of *N,N*,2-trimethyl-1H-imidazole-1-sulfonamide (compound **282.1**, 6.4 g, 33.8 mmol) in tetrahydrofuran (50 mL). This was followed by the addition of *n*-BuLi (2.5M in hexane) (16.3 mL, 40.6 mmol) dropwise with stirring at -78 °C. The reaction mixture was stirred for 1h at -78 °C. Oxirane (12.1 mL, 242.89 mmol) was added dropwise at -30 °C. The reaction mixture was stirred for 2h at room temperature, then quenched with 10 mL of water. The aqueous phase was extracted with 3 x 30 mL of ethyl acetate. The combined organic layers were washed with 3 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with EtOAc:MeOH (40:1) as the eluent to yield 3.4 g (43%) of the title compound as a white solid.




Compound 282.3. 2-(2-Methyl-1H-imidazol-4-yl)ethanol. To a solution of 4-(2-hydroxyethyl)-*N,N*,2-trimethyl-1H-imidazole-1-sulfonamide (compound **282.2**, 1 g, 4.29 mmol) in 1,4-dioxane (30 mL) was added HCl (6 M, 30 mL). The reaction mixture was stirred for 4h at 85 °C, then concentrated under reduced pressure. The pH of the solution was adjusted to 7-8 with NaHCO₃ (sat.). The resulting mixture was concentrated under reduced pressure to afford 3 g (crude) of the title compound as a white solid.



Compound 282. 4-(1-(3-(4-(2-Hydroxyethyl)-2-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 5, except 2-(2-methyl-1H-imidazol-4-yl)ethanol (compound **282.3**) was used instead of 2,4-dimethyl-1H-imidazole. m/z (ES+) 401 (M+H)⁺.




Compound 295.1. 4-(1-(3-(4-Formyl-2-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 5, except 5-iodo-2-methyl-1H-imidazole-4-carbaldehyde (compound **236.1**) was used instead of 5-iodo-2,4-dimethyl-1H-imidazole (compound **5.5**).



Compound 295. 4-(1-(3-(4-(Difluoromethyl)-2-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. To a solution of 4-(1-(3-(4-formyl-2-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile (compound **295.1**, 100 mg, 0.26 mmol) in dichloromethane (10 mL) was added DAST (103 μ L, 0.78 mmol) at -70 °C. The reaction mixture was stirred for 5h at room temperature. The pH of the solution was adjusted to 7 with sodium bicarbonate (sat.). The aqueous phase was extracted with 20 mL of


dichloromethane and the combined organic layers were washed with 3 x 10 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:20-1:2) as the eluent. The crude product (100 mg) was purified by Prep-HPLC with the following

5 conditions (Prep-HPLC-020): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*100mm; mobile phase, WATER WITH 0.05% TFA and ACN (16.0% up to 30.0% in 7 min, up to 95.0% in 1 min, down to 16.0% in 1 min); Detector, Waters 2489 254 & 220nm. This resulted in 48.7 mg (36%) of the title compound as a white solid. m/z (ES+) 407 (M+H)⁺.




10 **Compound 300.1. Ethyl 4-methoxycyclohexanecarboxylate.** Into a 100-mL round-bottom flask, was placed a solution of ethyl 4-hydroxycyclohexane-1-carboxylate (1 g, 5.81 mmol) in tetrahydrofuran (20 mL). Sodium hydride (349 mg, 8.72 mmol, 60%) was added to the reaction at 0 °C in portions, then stirred for 20 min. This was followed by the addition of CH₃I (865 μ L, 11.6 mmol) at 0 °C. The reaction mixture was stirred overnight at room

15 temperature, quenched with 6 mL of brine and extracted with 3 x 40 mL of ether. The combined organic layers were washed with 2 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford 800 mg (74%) of the title compound as a colorless oil.




20 **Compound 300.2. 4-Methoxycyclohexanecarbaldehyde.** Into a 100-mL 3-neck round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed a solution of ethyl 4-methoxycyclohexane-1-carboxylate (1.3 g, 6.98 mmol) in dichloromethane (30 mL). This was followed by the addition of DIBAL-H (1 M in hexanes) (9 mL, 9 mmol) dropwise at -78 °C. The reaction mixture was stirred for 3h at -78 °C, then


25 quenched with 10 mL of NH₄Cl (sat.). The aqueous phase was extracted with 3 x 50 mL of ethyl acetate. The combined organic layers were washed with 3 x 50 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 1 g (crude) of the title compound as a colorless oil.



Compound 300.2. methyl 3-(2-(4-methoxycyclohexyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **160.3**, except 4-methoxycyclohexanecarbaldehyde (compound **300.2**) was used instead of cyclopropanecarbaldehyde and methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used instead of except methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**).




Compound 300.3. 3-(2-(4-Methoxycyclohexyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoic acid. Into a 50-mL round-bottom flask, was placed a solution of methyl 3-(2-(4-methoxycyclohexyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **300.2**, 150 mg, 0.44 mmol) in methanol (4 mL). A solution of NaOH (70 mg, 1.75 mmol) in water (2 mL) was added to the reaction. The reaction mixture was stirred for 3 h at 50 °C, then concentrated under reduced pressure. The reaction mixture was diluted with 2 mL of water. The pH of the solution was adjusted to 1 with HCl (2 M) and concentrated under reduced pressure. This resulted in 240 mg (crude) of the title compound as a light yellow solid.




Compounds 300.A and 300.B. 4-(1-(3-(2-(4-Methoxycyclohexyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. Into a 50-mL round-bottom flask, was placed a solution of 3-(2-(4-methoxycyclohexyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **300.3**, 140 mg, 0.43 mmol) in *N,N*-dimethylformamide (3 mL). 4-(Azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**, 82.8 mg, 0.43 mmol), HBTU (242.6 mg, 0.64 mmol) and DIEA (228 μ L, 1.28 mmol) were added to the reaction. The reaction mixture was stirred for 1 overnight at room temperature, then diluted with 60

mL of EtOAc. The organic layer was washed with 3 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The crude product (150 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-020): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*100mm.; mobile phase, and (20.0% up to 27.0% in 10 min, up to 95.0% in 3 min, down to 20.0% in 1 min); Detector, Waters 2489 254 & 220nm. This resulted in 18.4 mg (9%) of compound **300.A** as a white solid and 59.2 mg (30%) of compound **300.B** as an off-white solid. m/z (ES⁺) 469 (M+H)⁺.




Compound 332.1. 3-(2-(4-Hydroxycyclohexyl)-4-methyl-1H-imidazol-5-yl)-4-

methylbenzoic acid. A solution of methyl 3-(2-(4-methoxycyclohexyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoate (compound **300.2**, 300 mg, 0.88 mmol) in HI (45% in water) (3 mL) was stirred for 2h at 60 °C, then concentrated under reduced pressure. This resulted in 500 mg (crude) of the title compound as brown oil which was used into the next step without further purification.




Compound 332. 4-(3-(3-(2-(4-Hydroxycyclohexyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)cyclobutyl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except 3-(2-(4-hydroxycyclohexyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoic acid (compound **332.1**) was used instead of 3-(2,4-dimethyl-1H-imidazol-5-yl)-4-methylbenzoic acid hydrochloride (compound **5.7**). m/z (ES⁺) 455 (M+H)⁺.



Compound 355.1. 4-(1-(3-(2-Acetyl-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard

chemical manipulations and procedures similar to those used for the preparation of compound 5, except 1-(4-methyl-1H-imidazol-2-yl)ethanone was used in place of 2,4-dimethyl-1H-imidazole.




5

Compound 355. 4-(1-(3-(2-(1,1-Difluoroethyl)-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. Into a 50-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed a solution of 4-(1-(3-(2-acetyl-4-methyl-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile (compound **355.1**, 85 mg, 0.21 mmol) in dichloromethane (15 mL). This was followed by the addition of DAST (82 μ L, 0.62 mmol). The reaction mixture was stirred for 4 days at room temperature, while adding fifteen additional equivalent of DAST over 4 days. The reaction was then quenched by the addition of 10 mL of water. The aqueous phase was extracted with 3 x 50 mL of dichloromethane and the combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/hexanes (6:1) as the eluent. The crude product (54 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-020): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*100mm; mobile phase, WATER WITH 0.05% TFA and MeCN (24.0% MeCN up to 38.0% in 5 min, up to 95.0% in 1 min, down to 34.0% in 1 min); Detector, Waters 2489 254 & 220 nm. This resulted in 42.8 mg (38%) of the title compound (trifluoroacetic acid salt) as a white solid. m/z (ES⁺) 421 (M+H)⁺.

15


20




Compound 371.1. N,2-Dimethyl-1H-imidazole-4-carboxamide. 2-Methyl-1H-imidazole-4-carboxylic acid (378 mg, 3 mmol), EDCI (745 mg, 3.9 mmol), HOBt (253 mg, 1.5 mmol), DIEA (1.6 mL, 9 mmol) and methyl amine (2M in THF) (3.6 mL, 7.2 mmol) were dissolved in DMF (10 mL) and stirred at room temperature for 16 hours. After removal

25


of DMF, the residue was dry loaded and purified by flash chromatography (SiO₂; 0-10 % methanol in DCM and 1% NH₄OH) to give 280 mg (67%) of the title compound. *m/z* (ES⁺) 140 (M+H)⁺.




Compound 371.1. 5-(5-(3-(4-Cyanophenyl)azetidine-1-carbonyl)-2-methylphenyl)-N,2-dimethyl-1H-imidazole-4-carboxamide. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except N,2-dimethyl-1H-imidazole-4-carboxamide (compound **371.1**) was used instead of 2,4-dimethyl-1H-imidazole. *m/z* (ES⁺) 414 (M+H)⁺.




Compound 349.1. 4,4,4-Trifluoro-2-oxobutanal. To a solution of SeO₂ (4.23 g, 38.1 mmol) in dioxane (25 mL) was added AcOH /H₂O (1mL/1 mL). 4,4,4-Trifluorobutan-2-one (3.81 mL, 31.7 mmol) was added dropwise to the above reaction mixture. The resulting solution was stirred overnight at 100 °C. The solution was used in the next step without any further work-up or purification.




Compound 349.2. 2-Methyl-4-(2,2,2-trifluoroethyl)-1H-imidazole. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159.1**, except 4,4,4-trifluoro-2-oxobutanal (compound **349.1**) was used instead of methylglyoxal.



Compound 349. 4-(1-(2,4-Dimethyl-5-(2-methyl-4-(2,2,2-trifluoroethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159**, except 2-methyl-4-(2,2,2-trifluoroethyl)-1H-imidazole (compound **349.2**) was used instead of 2-isopropyl-4-methyl-1H-imidazole (compound **159.1**). m/z (ES⁺) 453 (M+H)⁺.




Compound 296. 4-(1-(4-Methyl-3-(2-methyl-4-(2,2,2-trifluoroethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **159**, except 2-methyl-4-(2,2,2-trifluoroethyl)-1H-imidazole (compound **349.2**) was used instead of 2-isopropyl-4-methyl-1H-imidazole (compound **159.1**) and methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**) was used instead of methyl 2,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **160.1**). m/z (ES⁺) 439 (M+H)⁺.




Compound 377.1. Methyl 4-fluoro-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate. Into a 250-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed a solution of methyl 4-fluoro-5-iodo-2-

methylbenzoate (compound **101.2**, 5.04 g, 17.1 mmol) in DMSO (50 mL). 4,4,5,5-Tetramethyl-2-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (5.23 g, 20.6 mmol), KOAc (5.04 g, 51.4 mmol), Pd(dppf)Cl₂ (2.51 g, 3.43 mmol) were added to the reaction. The reaction mixture was stirred for 2h at 90 °C, then cooled and diluted with 300 mL of EtOAc. The solids were filtered off and the filtrate was washed with 3 x 30 mL of brine (sat.). The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/30) as the eluent to furnish 3.97 g (79%) of the title compound as an off-white solid.




Compound 377.2. 4-(1-(5-(3,4-Dimethyl-1H-pyrazol-5-yl)-4-fluoro-2-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **59**, except methyl 4-fluoro-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **377.1**) was used in place of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**). *m/z* (ES⁺) 389 (M+H)⁺.



Compound 377. 4-(1-(4-(Azetidin-1-yl)-5-(3,4-dimethyl-1H-pyrazol-5-yl)-2-methylbenzoyl)azetidin-3-yl)benzonitrile. Into a 5-mL sealed tube, was placed a solution of 4-(1-(5-(3,4-dimethyl-1H-pyrazol-5-yl)-4-fluoro-2-methylbenzoyl)azetidin-3-yl)benzonitrile (compound **377.2**, 120 mg, 0.31 mmol) in dioxane (3 mL). Azetidine (833 μL, 12.4 mmol) and Cs₂CO₃ (1.01 g, 3.09 mmol) were added to the reaction. The reaction mixture was stirred for 20h at 120 °C, then cooled with a water/ice bath. The resulting solution was diluted with 30 mL of EtOAc. The organic layer was washed with 3 x 20 mL of brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The crude product (200 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-020): Column,


SunFire Prep C18 OBD Column, 5 μ m, 19*100mm; mobile phase, Water with 10 mmol NH₄HCO₃ and MeCN (36.0% MeCN up to 50.0% in 6 min, up to 95.0% in 2 min, down to 36.0% in 1 min); Detector, waters 2489 254 & 220nm. This resulted in 56 mg (43%) of the title compound as a white solid. *m/z* (ES⁺) 426 (M+H)⁺.



5


Compound 376.1. Methyl 4-fluoro-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **377.1**, except methyl 4-fluoro-3-iodobenzoate (compound **132.1**) was used instead of methyl 4-fluoro-5-iodo-2-methylbenzoate (compound **101.2**).

10



Compound 376.2. 4-(1-(3-(2,4-Dimethyl-1H-imidazol-5-yl)-4-fluorobenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **5**, except methyl 4-fluoro-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **376.1**) was used instead of methyl 4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (compound **5.4**).

15



Compound 376. 4-(1-(4-(Azetidin-1-yl)-3-(2,4-dimethyl-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **377**, except 4-(1-(3-(2,4-dimethyl-1H-imidazol-5-yl)-4-fluorobenzoyl)azetidin-3-yl)benzonitrile (compound **376.2**) was used instead of 4-(1-(5-(3,4-dimethyl-1H-pyrazol-5-


20

yl)-4-fluoro-2-methylbenzoyl)azetidin-3-yl)benzonitrile (compound **377.2**). m/z (ES+) 412 (M+H)⁺.


The compounds in **TABLE 22** were prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **14**, **101**, **375** and **376**.

TABLE 22


Cpd	Name	Structure	m/z (ES+) (M+H) ⁺
382	4-(1-(4-(azetidin-1-yl)-3-(4-methyl-2-(trifluoromethyl)-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile		466
383	4-(1-(3-(2,4-dimethyl-1H-imidazol-5-yl)-4-(3-methylazetidin-1-yl)benzoyl)azetidin-3-yl)benzonitrile		426
384	4-(1-(3-(2,4-dimethyl-1H-imidazol-5-yl)-4-(pyrrolidin-1-yl)benzoyl)azetidin-3-yl)benzonitrile		426
385	4-(1-(3-(2,4-dimethyl-1H-imidazol-5-yl)-4-(3-methoxypyrrolidin-1-yl)benzoyl)azetidin-3-yl)benzonitrile		456




Compound 378. 4-(1-(4-Chloro-3-(2,4-dimethyl-1H-imidazol-5-yl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound 50, except 4-(azetidin-3-yl)benzonitrile hydrochloride (compound **5.2**) was used in place of 4-(piperidin-4-yl)benzonitrile hydrochloride (compound **1.2**). m/z (ES⁺) 391 (M+H)⁺.



Compound 379. 4-(1-(3-(2,4-dimethyl-1H-imidazol-5-yl)-4-(trifluoromethyl)benzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compound **50** and **378**. m/z (ES⁺) 425 (M+H)⁺.



Compound 380. 4-(1-(3-(2-((cis)-4-methoxycyclohexyl)-4-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **202**, **287**, **300A** and **300B**. m/z (ES⁺) 523 (M+H)⁺.



Compound 381. 4-(1-(3-(2-((trans)-4-methoxycyclohexyl)-4-(trifluoromethyl)-1H-imidazol-5-yl)-4-methylbenzoyl)azetidin-3-yl)benzonitrile. The title compound was prepared using standard chemical manipulations and procedures similar to those used for the preparation of compounds **202**, **287**, **300A** and **300B**. m/z (ES⁺) 523 (M+H)⁺.

The compounds of the present disclosure in **TABLE 26** below were, or may be, prepared using the foregoing procedures, as well as standard chemical manipulations and procedures similar to the foregoing procedures.

EXAMPLE 2

Activity of Compound of the Present Disclosure

Antiviral Activity

The antiviral activities of the compounds were assessed using the HCV1b replicon system. The replicon was constructed using the ET (luc-ubi-neo/ET) cell line, a Huh7 human hepatoma cell line harboring an HCV replicon with a stable luciferase (Luc) reporter and three cell culture-adaptive mutations (Pietschmann, *et al* (2002) *J. Virol.* 76:4008-4021). The HCV replicon antiviral evaluation assay examined the effects of compounds at ten three-fold dilutions. Sub-confluent cultures of the ET line were plated out into 96-well plates that were dedicated for the analysis of cell viability (cytotoxicity) or antiviral activity and the next day drugs were added to the appropriate wells. Cells were processed 72 hr later when the cells are still sub-confluent. EC₅₀ (concentrations inhibiting the HCV RNA replicon by 50%), CC₅₀ (concentration decreasing cell viability by 50%) and SI (selective index: CC₅₀/EC₅₀) values were determined. HCV RNA replicon levels were assessed using the Bright-GloTM Luciferase Assay System (Promega) to measure replicon-derived Luc activity. The CellTiter-GloTM Luminescent Cell Viability Assay (Promega) was used to estimate cell viability. Inhibition of HCV for selected compounds is disclosed in **TABLE 26**. None of the compounds in **TABLE 26** caused material cytotoxicity at the highest concentrations tested.

FASN Inhibition by Compounds of the Present Disclosure

Determination of FASN biochemical activity: The FASN enzyme was isolated from SKBr3 cells. SKBr3 is a human breast cancer cell-line with high levels of FASN expression.

5 It is estimated that FASN comprises about 25% of the cytosolic proteins in this cell line. SKBr3 cells were homogenized in a dounce homogenizer then centrifuged for 15 minutes at 4 °C to remove particulate matter. The supernatant was then analyzed for protein content, diluted to the appropriate concentration, and used to measure FASN activity. The presence of FASN was confirmed by western blot analysis. A similar method for isolation of FASN from
10 SKBr3 cells is described in Teresa, P. et al. (*Clin. Cancer Res.* 2009; 15(24), 7608-7615).

FASN activity of the SKBr3 cell extract was determined by measuring either NADPH oxidation or the amount of thiol-containing coenzyme A (CoA) released during the fatty acid synthase reaction. The dye CPM (7-diethylamino-3-(4'-maleimidyl-phenyl)-4-methylcoumarin) contains a thiol reactive group that increases its fluorescence emission on
15 reaction with the sulfhydryl group of CoA. The biochemical activities shown in **TABLE 26** were determined using the fluorescence measurement of CoA release via a procedure described in Chung C.C. et al. (*Assay and Drug Development Technologies*, 2008, 6(3), 361-374).

FASN Inhibitors Have Oncology Efficacy

Experimental Procedures

Cell viability assay: For drug treatment, PANC-1 cells were plated in 96-well plates at a concentration of 1.0×10^3 per well in medium (Advanced MEM containing 1% or 10% FBS). Serial 3-fold dilutions of compounds **160**, **161**, **242**, or **164** were prepared in DMSO
25 and further diluted 1:10 in Advanced MEM for addition to the assay solution. 24-hours after cell plating, advanced MEM-diluted compound was added to make a final reaction volume of 100 ul per well. Final drug concentrations in assay wells were 10,000, 3,300, 1,100, 370, 123, 41, 14, 5, and 0 nmol/L. Assays were done in triplicate at each drug concentration 10 days after compound addition. The number of viable cells was measured using the Cell Titer-Glo
30 assay (Promega) according to the manufactures instructions. Luminosity per well was determined and the signal intensity was analyzed versus drug concentration. For each FASN

inhibitor, the concentration of drug resulting in 50% inhibition of the maximum signal was determined, and this value was reported as the IC₅₀.

Xenograft tumor growth inhibition efficacy study: For xenograft tumor development and drug treatment, female BALB/c-nude mice, 8-9 weeks of age, were inoculated subcutaneously at the right flank with PANC-1 tumor cells (5×10^6) in 0.1 ml of PBS. The day of tumor cell inoculation is denoted as day 0. Drug treatment was initiated when the mean tumor size reached 149 mm³ (day 15). Drug administration to tumor-bearing mice (n=10 per group) was according to the following regimen: compound **161**: 30 or 60 mg/kg once daily (Qd); compound **242**: 30 or 60 mg/kg once daily (Qd); C75 (trans-4-methylene-2-octyl-5-oxotetrahydrofuran-3-carboxylic acid): 30 mg/kg once every 5 days (Q5d); gemcitabine 40 mg/kg once every 3 days (Q3d). The administered dose of C75 was lowered to 20 mg/kg once every 5 days following the first dose, due to drug-induced weight loss and toxicity. Mice were assigned into treatment groups using a randomized block design based upon their tumor volumes. This ensured that all the groups were comparable at the start of treatment. Tumor sizes were measured twice weekly in two dimensions using a caliper, and the volumes were expressed in mm³ using the formula: $V = 0.5 a \times b^2$ where a and b were the long and short diameters of the tumor, respectively. The study was terminated 6 hours post the 18th Qd dose on day 32 after tumor inoculation. Tumor growth inhibition (TGI) was calculated as the percentage of tumor growth, relative to tumor size at the start of treatment, in drug-treated groups compared to vehicle-treated groups. The Mann-Whitney U test was used to assess statistical significance of the mean tumor size between drug and vehicle-treated groups.

Results

FASN inhibition inhibits tumor cell growth and induces cell death: To test for effects of FASN inhibition on tumor cell survival, a pancreatic tumor cell line (PANC-1) was treated with multiple FASN inhibitors, and IC₅₀ values were determined from a 9-point dose response cell viability assay. Cell survival IC₅₀ values were determined 10 days following addition of FASN inhibitor compounds to the tumor cells (**TABLE 23**). FASN cell survival IC₅₀ values were in agreement with previously determined FASN biochemical and cell-based IC₅₀ values in palmitate synthesis and HCV replicon assays. Alignment of IC₅₀ values in

phenotypic (cell survival) and mechanistic (palmitate synthesis) assays demonstrates that tumor cell death induced by FASN inhibitor treatment is a direct result of FASN inhibition.

TABLE 23. FASN inhibitor in vitro cell assay IC₅₀ values

Treatment	<u>Survival</u>	<u>HCV</u>	<u>Palmitate</u>	<u>FASN</u>
	PANC-1	HeLa	HeLa	Biochem.
	(nM)	(nM)	(nM)	(nM)
Compound 160	16 ^a	7	24	17
Compound 161	112 ^a	34	51	32
Compound 242	42 ^b	77	74	42
Compound 164	1.2 ^b	2	27	19

5 a -Advanced MEM cell medium with 10% FBS

b -Advanced MEM cell medium with 1% FBS

FASN inhibition inhibits in vivo xenograft tumor growth: To test for the effects of FASN inhibition on in vivo tumor growth, PANC-1 xenograft tumors were inoculated subcutaneously in female BALB/c-nude mice. When the xenograft tumors grew to an average of 149 mm³ (day 15 following tumor cell inoculation), drug treatment was initiated with 3-V Biosciences reversible FASN inhibitors (compound **161** or **242**) or control compounds (C75 or gemcitabine) to determine tumor growth inhibition activity of the different compounds in the PANC-1 xenograft tumor model. Drug treatment groups consisted of 10-tumor-bearing mice per group. The FASN inhibitors, compounds **161** and **242** both demonstrated tumor growth inhibition activity in a dose-responsive manner. Compound **161** dosed at 60 mg/kg once daily for 18 days resulted in 52% tumor growth inhibition compared to vehicle treated mice, and compound **242** dosed at 100 mg/kg once daily for 18 days resulted in 57% tumor growth inhibition compared to vehicle treated mice (**TABLES 24** and **25**). Once daily 30 mg/kg doses of compounds **161** and **242** resulted in 31% and 19% tumor growth inhibition, respectively. C75, the irreversible FASN inhibitor control compound was dosed initially at 30 mg/kg once every 5 days; however, toxicity of this compound required a dose adjustment to 20 mg/kg once every 5 days. The C75 treatment group after 18 days (3 doses) showed 65% tumor growth inhibition. Six of the ten mice in the C75 treatment group died from C75-related toxicity during the 18-day study. Gemcitabine treated PANC-1 tumor-bearing mice

(40 mg/kg once every 3 days) showed 96 percent tumor growth inhibition. Tumor growth inhibition induced by compounds **161** and **242**, C75, and gemcitabine was statistically significant compared to vehicle treated mice. The results from this study demonstrate that reversible FASN inhibitors discovered at 3-V Biosciences are well tolerated in PANC-1 xenograft tumor-bearing mice, and that once-daily dosing of the inhibitors as a single agent cancer therapeutic is associated with statistically significant tumor growth inhibition compared to vehicle treated mice. Additional studies to investigate dosing levels, schedules, and in combination with existing standard of care chemotherapies are being pursued.

Tumor Volumes

The tumor sizes of each treatment group at different time points are shown in **TABLE 24**.

TABLE 24. Mean tumor sizes in the different treatment groups

Treatment	Tumor volume (mm ³) ^a					
	15	19	22	26	29	32
Compound 161 (30 mg/kg)	150±13	159±10	211±17	297±31	379±53	479±69
Compound 161 (60 mg/kg)	150±13	178±23	213±32	207±33	261±51	413±83
Compound 242 (30 mg/kg)	149±12	219±24	240±18	304±21	394±24	543±38
Compound 242 (100 mg/kg)	149±13	157±15	187±22	247±29	300±39	378±59
C75 (30/20 mg/kg)	150±15	164±22	156±26	213±43	367±77	403±83
Gemcitabine (40 mg/kg)	149±12	144±16	152±14	177±20	172±21	165±27
Vehicle 1 (0.5% CMC)	148±11	200±23	242±32	423±57	491±57	638±108
Vehicle 2 (20% DMSO, 80% PBS)	146±11	212±11	277±18	409±18	474±24	614±29

Note: a. Mean ± SEM.






Tumor Growth Inhibition






- 5 The tumor growth inhibition (TGI) and statistical analysis of each treatment group at the study conclusion (day 32) is summarized in **TABLE 25**.






TABLE 25. Antitumor activity of compounds 161 and 242 as a single agent in the treatment of subcutaneous PANC-1 human pancreatic cancer xenograft model






Treatment	TGI vs. Vehicle 1 (%)	<i>P</i> -value vs. Vehicle 1	TGI vs. Vehicle 2 (%)	<i>P</i> -value vs. Vehicle 2
Compound 161 (30 mg/kg)	31	0.310	29	0.052
Compound 161 (60 mg/kg)	52	0.188	51	0.014
Compound 242 (30 mg/kg)	19	0.950	16	0.244
Compound 242 (100 mg/kg)	57	0.043	56	0.002
C75 (30/20 mg/kg)	65	0.024	64	0.002
Gemcitabine (40 mg/kg)	96	<0.001	96	<0.001
Vehicle 1 (0.5% CMC)	--	--	--	1.000
Vehicle 2 (20% DMSO, 80% PBS)	--	1.000	--	--






TABLE 26. Compounds of the present disclosure.






Structure	Cpd	Hu FASN IC ₅₀ (μM)	HCV GT1b EC ₅₀ (μM)
	1	0.024	0.027
	2	0.041	0.037
	3	0.032	0.068
	4	0.053	0.081
	5	0.010	0.009






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	6	0.105	0.086
	7	0.006	0.003
	8	0.040	0.050
	9	0.025	0.030
	10	0.027	0.030






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	11	0.036	0.044
	12	0.036	0.160
	13	0.006	0.005
	14	0.027	0.031
	15	0.026	0.022






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	16	0.018	0.013
	17	0.045	0.081
	18	0.540	
	19	0.030	0.042
	20	0.165	0.130






Structure	Cpd	Hu FASN IC ₅₀ (μM)	HCV GT1b EC ₅₀ (μM)
	21	0.207	0.390
	22	1.690	2.220
	23	0.029	0.093
	24	0.041	0.097
	25		0.019






Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	26	0.245	
	27	0.009	0.005
	28	0.048	0.011
	29	0.024	0.009
	30	0.016	0.013






Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	31	0.011	0.003
	32	0.012	0.003
	33	0.080	
	34	0.048	
	35	2.870	2.780






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	36	9.190	
	37	0.032	0.032
	38	0.104	0.640
	39	0.420	0.980
	40	0.620	1.700






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	41	50.000	
	42	0.032	0.115
	43	0.012	0.004
	44	0.007	0.003
	45	0.009	0.002






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	46	0.005	0.004
	47	0.007	0.003
	48	0.009	0.004
	49	0.009	0.003
	50	0.145	0.250






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	51	0.019	0.051
	52	0.046	0.280
	53	0.033	0.100
	54	0.020	0.033
	55	0.086	0.320






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	56	0.021	0.032
	57	0.097	
	58	0.102	0.310
	59	0.033	0.063
	60	0.110	0.360






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	61	0.310	
	62	0.650	0.760
	63	1.180	6.170
	64	1.550	1.940
	65	0.050	0.096






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	66	2.370	15.000
	67	2.490	
	68	0.150	0.240
	69	0.018	0.016
	70	0.205	0.680

Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	71	0.300	0.710
	72	0.420	2.220
	73	1.000	2.570
	74	0.460	0.730
	75	0.073	0.111






Structure	Cpd	Hu FASN IC ₅₀ (μM)	HCV GT1b EC ₅₀ (μM)
	76	0.069	0.245
	77	0.125	0.170
	78	3.670	5.420
	79	0.059	0.150
	80	0.057	0.295






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	81	0.054	0.200
	82	0.020	0.036
	83	0.062	0.089
	84	0.112	0.540
	85	0.057	






Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	86	0.086	0.850
	87	0.040	0.029
	88	0.021	0.009
	89	0.049	0.087
	90	0.071	0.165






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	91	0.140	0.140
	92	0.140	0.230
	93	0.034	
	94	0.029	0.630
	95	0.030	0.008

Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	96	0.270	
	97	0.190	
	98	0.240	0.960
	99	0.045	
	100	0.055	0.070






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	101	0.530	
	102	0.890	
	103	0.135	
	104	0.070	0.270
	105	0.070	0.030






Structure	Cpd	Hu FASN IC ₅₀ (μM)	HCV GT1b EC ₅₀ (μM)
	106	50.000	20.000
	107	5.580	20.000
	108	2.250	15.000
	109	0.280	0.350
	110	11.020	20.000






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	111	0.230	1.980
	112	0.840	1.490
	113	0.145	0.270
	114	0.200	0.450
	115	0.320	1.210






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	116	50.000	20.000
	117	0.980	3.330
	118	0.680	3.180
	119	4.480	10.000
	120	1.300	10.000






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	121	0.030	0.240
	122	3.440	
	123	3.710	
	124	0.140	0.230
	125	0.110	0.110






Structure	Cpd	Hu FASN IC ₅₀ (μM)	HCV GT1b EC ₅₀ (μM)
	126	1.130	
	127	3.270	
	128	0.098	0.120
	129	1.210	2.150
	130	0.082	3.200






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	131	0.185	2.200
	132	0.210	0.340
	133	0.065	0.080
	134	0.099	0.230
	135		






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	136	0.026	0.052
	137	0.105	0.230
	138	0.120	0.250
	139	0.017	0.014
	140	0.080	0.240






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	141	0.099	0.280
	142	0.024	0.018
	143	0.018	0.002
	144	0.020	0.003
	145	2.170	20.000






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	146	0.062	0.017
	147	0.120	
	148	0.084	
	149	0.021	
	150	0.097	






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	151	0.057	0.026
	152	0.290	0.380
	153	0.036	0.032
	154	0.094	0.380
	155	0.590	






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	156	1.495	
	157	0.210	0.056
	158		0.028
	159	0.021	0.016
	160	0.017	0.007






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	161	0.026	0.034
	162	0.019	0.006
	163	0.037	0.027
	164	0.019	0.002
	165	0.038	0.018






Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	166	0.040	0.016
	167	0.061	0.025
	168	0.011	0.002
	169	0.035	0.010
	170	0.040	






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	171	0.021	0.005
	172	0.011	0.001
	173	0.035	0.018
	174	0.037	0.011
	175	0.020	0.006






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	176	0.024	0.016
	177	0.011	0.001
	178	0.027	0.006
	179	0.025	0.013
	180	0.034	0.007






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	181	0.068	0.021
	182	0.038	0.079
	183	0.029	
	184	0.009	0.002
	185	0.033	






Structure	Cpd	Hu FASN IC ₅₀ (μM)	HCV GT1b EC ₅₀ (μM)
	186	0.066	0.025
	187	0.063	
	188	0.052	0.007
	189	0.020	0.045
	190	0.008	






Structure	Cpd	Hu FASN IC ₅₀ (μM)	HCV GT1b EC ₅₀ (μM)
	191	0.065	0.150
	192	0.024	0.010
	193	0.065	0.370
	194	0.025	0.004
	195	0.013	0.004






Structure	Cpd	Hu FASN IC ₅₀ (μM)	HCV GT1b EC ₅₀ (μM)
	196	0.015	
	197	0.870	0.730
	198	0.054	0.018
	199	0.110	0.024
	200	0.027	0.019






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	201	0.130	0.054
	202	0.020	0.012
	203	0.035	0.021
	204	0.010	0.005
	205	0.016	0.009






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	206	0.024	0.026
	207	0.035	0.012
	208	0.036	0.025
	209	0.024	0.005
	210	0.020	0.017






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	211	0.011	0.006
	212	0.018	0.013
	213	0.065	0.051
	214	0.014	0.006
	215	0.048	






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	216	0.079	0.026
	217	0.052	
	218	0.015	0.003
	219	0.059	0.040
	220	0.050	0.079






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	221	0.043	0.120
	222	0.053	0.033
	223	0.023	0.015
	224	0.031	0.038
	225	0.059	0.066






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	226	0.088	0.070
	227	0.012	0.013
	228	0.027	0.022
	229	0.042	0.012
	230	0.017	0.007






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	231	0.019	0.002
	232	0.020	0.013
	233	0.011	0.002
	234	0.053	1.600
	235	0.039	1.100






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	236	0.019	0.019
	237	0.016	0.014
	238	0.010	0.001
	239	0.440	0.430
	240	0.053	0.032






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	241	0.009	0.000
	242	0.060	0.069
	243	0.170	0.450
	244	0.150	0.120
	245	0.109	





Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	246	0.340	0.740
	247	0.110	
	248	0.075	0.077
	249	0.043	0.026
	250	0.051	0.075





Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	251	0.178	0.340
	252	0.095	0.230
	253	0.080	0.220
	255	0.063	0.083
	256	0.210	





Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	257	0.440	0.660
	258	0.058	
	259	0.140	0.082
	260	0.067	0.250
	261	0.190	0.170





Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	262	0.140	0.190
	263	0.094	0.067
	264	0.056	0.088
	265	0.110	
	266	0.200	0.260





Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	267	0.100	0.150
	268	0.059	0.029
	269	50.000	
	270		
	271		





Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	272		
	273		
	274	0.051	0.016
	275	0.057	0.038





Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	276	50	20
	277	50	20
	278		
	279	0.11	0.039






Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	280	0.089	0.15
	281	0.072	0.021
	282	0.091	0.15
	283	0.064	0.082





Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	284	0.044	0.01
	285	0.048	0.027
	286	0.061	0.027
	287	0.029	0.003





Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	288	0.026	0.01
	289	0.018	0.01
	290	0.021	0.029
	291	0.02	0.003






Structure	Cpd	Hu FASN IC ₅₀ (μM)	HCV GT1b EC ₅₀ (μM)
	292	0.044	0.033
	293	0.064	0.12
	294	0.049	0.041
	295	0.031	0.037

Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	296	0.024	
	297		
	298		
	299		







Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	300A	0.026	0.007
	300B	0.024	0.003
	301		
	302		
	303		






Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	304	0.021	
	305	0.078	0.077
	306		
	307	0.056	






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	308		
	309	0.037	
	310		
	311		

Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	312	0.037	
	313	0.1	
	314	0.46	
	315		
	316		





Structure	Cpd	Hu FASN IC ₅₀ (μM)	HCV GT1b EC ₅₀ (μM)
	317		
	318		
	320		
	321		
	322		
	323		






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	324		
	325		
	326		
	327		
	328		
	329	0.089	






Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	330	0.1	0.028
	331	0.11	0.021
	332	0.01	
	333		
	334		






Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	335		
	336		
	337		
	338		
	339		






Structure	Cpd	Hu FASN IC ₅₀ (μM)	HCV GT1b EC ₅₀ (μM)
	340		
	341		
	342		
	343		
	344	0.024	0.013
	345	0.035	0.006







Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	346	0.018	0.0001
	347	0.016	0.005
	348	0.014	0.002
	349	0.019	







Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	350	0.055	
	351		
	352	0.011	
	353	0.009	
	354	0.018	



Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	355	0.082	
	356	0.851	
	357	0.087	
	358	0.110	
	359	0.014	

Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	360	0.012	
	361	0.013	
	362	0.011	
	363	0.021	
	364	0.040	

Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	365	0.023	
	366	0.022	
	367	0.040	
	368	0.012	
	369	0.014	

Structure	Cpd	Hu FASN IC50 (μM)	HCV GT1b EC50 (μM)
	370	0.012	
	371	0.82	
	374	0.024	
	375		
	376	0.029	
	377	0.029	

Structure	Cpd	Hu FASN IC ₅₀ (μM)	HCV GT1b EC ₅₀ (μM)
	378	0.072	
	379	0.059	
	380	0.013	
	381	0.009	
	382	0.018	
	383	0.024	


Structure	Cpd	Hu FASN IC50 (μ M)	HCV GT1b EC50 (μ M)
	384	0.012	
	385	0.04	

While preferred aspects of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such aspects are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the aspects of the invention described herein can be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.


All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

CLAIMS

1. A compound having Structural Formula III:



III, or pharmaceutically acceptable salts thereof, wherein:



Het is a 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered heterocycle) or -O-(C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;

R^3 is H or F;

R^{11} is H or -CH₃;

R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle;


R^{22} is H, halogen, or C_1 - C_2 alkyl;


R^{24} is H, -CN, -(C_1 - C_4 alkyl)-CN, C_1 - C_4 alkyl, -(C_1 - C_4 alkyl)-OH, -(C_1 - C_4 alkyl)-N(R^{241})₂, -(C_1 - C_4 alkyl)-O-(C_3 - C_6 cycloalkyl), -(C_1 - C_4 alkyl)-O-(4- to 6-membered heterocycle) or -(C_1 - C_4 alkyl)-O-(C_1 - C_4 alkyl), wherein:



t is 0 or 1;



u is 0 or 1;

with the proviso that when u is 1, t is 1; and
each R^{241} is independently H or C_1 - C_2 alkyl; and
 R^{25} is halogen, -CN, $-(C_1-C_4 \text{ alkyl})-CN$, C_1 - C_2 alkyl or cyclopropyl.


2. The compound of claim 1, wherein when L-Ar is , Ar is not



3. The compound of claim 1, wherein i) L-Ar is  or  and Ar



4. The compound of claim 3, wherein Ar is



5. The compound of any of claims 1-4, wherein i) R^1 is halogen, -CN or C_1 - C_2 haloalkyl; or ii) R^1 is -CN.

6. The compound of any of claims 1-5, wherein R^2 is H.
7. The compound of any of claims 1-6, wherein i) R^{21} is halogen, C_1 - C_4 alkyl or C_3 - C_5 cycloalkyl; or ii) R^{21} is C_1 - C_4 alkyl or C_3 - C_5 cycloalkyl; or iii) R^{21} is C_1 - C_2 alkyl or C_3 - C_5 cycloalkyl; or iv) R^{21} is C_1 - C_2 alkyl; or v) R^{21} is $-CH_3$.
8. The compound of any of claims 1-7, wherein i) R^{22} is H or C_1 - C_2 alkyl; or ii) R^{22} is H or $-CH_3$; or iii) R^{22} is $-CH_3$.
9. The compound of any of claims 1-8, wherein i) R^{24} is H, $-CN$, $-(C_1-C_4 \text{ alkyl})-CN$, C_1 - C_4 alkyl, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_6 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4\text{- to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or ii) R^{24} is H, C_1 - C_4 alkyl, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_6 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4\text{- to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or iii) R^{24} is C_1 - C_4 alkyl or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or iv) R^{24} is $-(C_1-C_2 \text{ alkyl})-O-(C_1-C_2 \text{ alkyl})$; or v) R^{24} is C_1 - C_2 alkyl; or vi) R^{24} is C_3 - C_6 cycloalkyl; or vii) R^{24} is $-CN$ or $-(C_1-C_2 \text{ alkyl})-CN$; or viii) R^{24} is $-(C_1-C_2 \text{ alkyl})-CN$; or ix) R^{24} is methyl, isopropyl, cyclopropyl, $-CN$, or $-(C_1-C_2 \text{ alkyl})-CN$.
10. The compound of claim 9, wherein i) R^{24} is $-CH_2-O-CH_3$; or ii) R^{24} is $-CH_3$; or iii) R^{24} is $-CN$; or iv) R^{24} is H, $-CH_3$, $-CH_2OH$, $-CH_2OCH_3$, $-(CH_2)_2OH$, $-(CH_2)_2OCH_3$ or $-(CH_2)_2N(CH_3)_2$.
11. The compound of claim 9, wherein R^{24} is substituted with one or more substituents selected from C_1 - C_2 alkyl, oxo, $-CN$, halogen, alkanoyl, alkoxycarbonyl, $-OH$ and C_1 - C_2 alkoxy when R^{24} is $-(C_1-C_4 \text{ alkyl})-CN$, C_1 - C_4 alkyl, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_6 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4\text{- to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or R^{24} is substituted with one or more substituents selected from methyl, $-F$, methoxy, $-C(=O)CH_3$ and $-C(=O)-OCH_3$ when R^{24} is $-(C_1-C_4 \text{ alkyl})-CN$, C_1 - C_4 alkyl, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_6 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4\text{- to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$.
12. The compound of claim 11, wherein i) R^{24} is substituted with two substituents that

are the same or different when R^{24} is $-(C_1-C_4 \text{ alkyl})-CN$, $C_1-C_4 \text{ alkyl}$, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_6 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4- \text{ to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or ii) R^{24} is substituted with three substituents that are the same or different when R^{24} is $-(C_1-C_4 \text{ alkyl})-CN$, $C_1-C_4 \text{ alkyl}$, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_6 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4- \text{ to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$.


13. The compound of any of claims 1-12, wherein i) R^{25} is halogen, $-CN$, $C_1-C_2 \text{ alkyl}$ or cyclopropyl; or ii) R^{25} is halogen, $C_1-C_2 \text{ alkyl}$ or cyclopropyl.

14. The compound of claim 13, wherein i) R^{25} is $-CN$, $-Cl$ or $-CH_3$; or ii) R^{25} is $-Cl$; or R^{25} is $-CH_3$.


15. The compound of any of claims 1-13, wherein when R^{25} includes alkyl or cyclopropyl, i) R^{25} is substituted with one or more substituents selected from $-OH$, halogen, $C_1-C_2 \text{ alkyl}$ and alkylcarbonyloxy; or ii) R^{25} is substituted with one or more substituents selected from $-F$, methyl and $-O-C(=O)-CH_3$.


16. The compound of claim 15, wherein when R^{25} includes alkyl or cyclopropyl, i) R^{25} is substituted with two substituents that are the same or different; or ii) R^{25} is substituted with three substituents that are the same or different.

17. A compound of Structural Formula IIIb:



IIIb, or pharmaceutically acceptable salts thereof, wherein:





Het is a 5- to 6-membered heteroaryl;

R^1 is H, -CN, halogen, C_1 - C_4 alkyl, -O-(C_3 - C_5 cycloalkyl), -O-(4- to 6-membered heterocycle) or -O-(C_1 - C_4 alkyl), wherein when R^1 is not H, -CN or halogen, R^1 is optionally substituted with one or more halogens;

each R^2 is independently hydrogen, halogen or C_1 - C_4 alkyl;

R^3 is H or F;

R^{11} is H or -CH₃;

R^{21} is H, halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle;

R^{22} is H, halogen or C_1 - C_2 alkyl; and


each R^{24} and R^{25} is independently H, halogen, -CN, -(C_1 - C_4 alkyl)-CN, C_1 - C_4 alkyl, -(C_1 - C_4 alkyl)-OH, -(C_1 - C_4 alkyl)-N(R^{241})₂, -(C_1 - C_4 alkyl)_t-O_u-(C_3 - C_5 cycloalkyl), -(C_1 - C_4 alkyl)_t-O_u-(4- to 6-membered heterocycle) or -(C_1 - C_4 alkyl)_t-O-(C_1 - C_4 alkyl), wherein:

each t is 1;


each u is independently 0 or 1; and



each R^{241} is independently H or C_1 - C_2 alkyl,




wherein the compound is not:




18. The compound of claim 17, wherein when L-Ar is , Ar is not



19. The compound of claim 17, wherein i) L-Ar is  or  Ar and



20. The compound of claim 19, wherein Ar is



21. The compound of any of claims 17-20, wherein i) R¹ is halogen, -CN or C₁-C₂ haloalkyl; or ii) R¹ is -CN.

22. The compound of any of claims 17-21, wherein R² is H.

23. The compound of any of claims 17-22, wherein i) R²¹ is halogen, C₁-C₄ alkyl, C₃-C₅ cycloalkyl or 4- to 6-membered heterocycle; or ii) R²¹ is C₁-C₄ alkyl, C₃-C₅ cycloalkyl or 4- to

6-membered heterocycle; or iii) R^{21} is C_1 - C_2 alkyl or C_3 - C_5 cycloalkyl; or iv) R^{21} is C_1 - C_2 alkyl; or v) R^{21} is $-CH_3$.

24. The compound of any of claims 17-23, wherein i) R^{22} is H or C_1 - C_2 alkyl; or ii) R^{22} is H or $-CH_3$; or iii) R^{22} is $-CH_3$.

25. The compound of any of claims 17-24, wherein i) each R^{24} and R^{25} is independently H, $-CN$, C_1 - C_4 alkyl, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_5 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4- \text{ to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or ii) each R^{24} and R^{25} is independently H, C_1 - C_4 alkyl, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4- \text{ to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or iii) R^{24} is H, C_1 - C_4 alkyl, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_5 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4- \text{ to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or iv) R^{24} is $-CN$, $-Cl$, C_1 - C_4 alkyl or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or v) R^{24} is C_1 - C_4 alkyl or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or vi) R^{24} is $-(C_1-C_2 \text{ alkyl})-O-(C_1-C_2 \text{ alkyl})$; or vii) R^{24} is C_1 - C_4 alkyl.

26. The compound of claim 17-25, wherein i) R^{24} is $-CH_3$; or ii) R^{24} is hydrogen.

27. The compound of any of claims 17-25, wherein i) R^{24} is substituted with one or more substituents selected from halogen, C_3 - C_5 cycloalkyl and C_1 - C_2 alkoxy when R^{24} is $-(C_1-C_4 \text{ alkyl})-CN$, C_1 - C_4 alkyl, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_6 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4- \text{ to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or ii) R^{24} is substituted with one or more substituents selected from $-F$, cyclopropyl and $-OCH_3$ when R^{24} is $-(C_1-C_4 \text{ alkyl})-CN$, C_1 - C_4 alkyl, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_6 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4- \text{ to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$.

28. The compound of claim 27, wherein i) R^{24} is substituted with two substituents that are the same or different when R^{24} is $-(C_1-C_4 \text{ alkyl})-CN$, C_1 - C_4 alkyl, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_6 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4- \text{ to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or ii) R^{24} is substituted with three substituents that


are the same or different when R^{24} is $-(C_1-C_4 \text{ alkyl})-CN$, $C_1-C_4 \text{ alkyl}$, $-(C_1-C_4 \text{ alkyl})-OH$, $-(C_1-C_4 \text{ alkyl})-N(R^{241})_2$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(C_3-C_6 \text{ cycloalkyl})$, $-(C_1-C_4 \text{ alkyl})_t-O_u-(4- \text{ to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$.

29. The compound of any of claims 17-28, wherein i) R^{25} is halogen, methyl, ethyl or cyclopropyl; or ii) R^{25} is $-CN$, $-Cl$, $C_1-C_4 \text{ alkyl}$, $-(C_1-C_4 \text{ alkyl})_t-O-(C_3-C_5 \text{ cycloalkyl})$ or $-(C_1-C_4 \text{ alkyl})_t-O-(C_1-C_4 \text{ alkyl})$; or iii) R^{25} is $-CN$, $-Cl$, $-CH_3$, $-O-(C_3-C_5 \text{ cycloalkyl})$ or $-O-(C_1-C_2 \text{ alkyl})$; or iv) R^{25} is $-CN$, $-Cl$ or $C_1-C_4 \text{ alkyl}$.

30. The compound of claim 17-29, wherein i) R^{25} is $-CH_3$; or ii) R^{25} is $-Cl$.

31. The compound of any of claims 17-29, wherein when R^{25} includes alkyl, cycloalkyl or heterocycle, i) R^{25} is substituted with one or more halogen; or ii) R^{25} is substituted with one or more $-F$; or iii) R^{25} is substituted by two substituents; or iv) R^{25} is substituted by three substituents.

32. A compound having Structural Formula I or II:





or pharmaceutically acceptable salts thereof, wherein:

L^3 is $-CH_2-$, $-CHR^{50}-$, $-O-$, $-NR^{50}-$, $-NC(O)R^{50}-$ or $-NC(O)OR^{50}-$, wherein R^{50} is $C_1-C_6 \text{ alkyl}$, $C_3-C_5 \text{ cycloalkyl}$, or 4- to 6-membered heterocycle;

n is 1, 2, or 3;

m is 1 or 2 with the proviso that $n+m \geq 3$;





Het is a 5- to 6-membered heteroaryl;

R¹ is H, -CN, halogen, C₁-C₄ alkyl, -O-(C₃-C₅ cycloalkyl), -O-(4- to 6-membered heterocycle) or -O-(C₁-C₄ alkyl), wherein when R¹ is not H, -CN or halogen, R¹ is optionally substituted with one or more halogens;

each R² is independently hydrogen, halogen or C₁-C₄ alkyl;

R³ is H or F;

R¹¹ is H or -CH₃;


R²¹ is H, halogen, C₁-C₄ alkyl, C₃-C₅ cycloalkyl or 4- to 6-membered heterocycle;



R²² is H, halogen, or C₁-C₂ alkyl; and

R²⁴ is H, C₁-C₄ alkyl, -(C₁-C₄ alkyl)-OH, -(C₁-C₄ alkyl)_t-N(R²⁴¹)₂, -(C₁-C₄ alkyl)_t-O-(C₃-C₅ cycloalkyl), -(C₁-C₄ alkyl)_t-O-(4- to 6-membered heterocycle) or -(C₁-C₄ alkyl)_t-O-(C₁-C₄ alkyl), wherein:

each t is independently 0 or 1; and

each R²⁴¹ is independently H or C₁-C₂ alkyl.





35. The compound of any of claims 32-34, wherein i) R^1 is halogen, -CN or C_1 - C_2 haloalkyl; or ii) R^1 is -CN or C_1 - C_2 haloalkyl or iii) R^1 is -CN; or iv) R^1 is -Cl.

36. The compound of any of claims 32-35, wherein R^2 is H.

37. The compound of any of claims 32-36 wherein i) R^{21} is halogen, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle; or ii) R^{21} is H, C_1 - C_4 alkyl, C_3 - C_5 cycloalkyl or 4- to 6-membered heterocycle; or iii) R^{21} is C_1 - C_2 alkyl or C_3 - C_5 cycloalkyl; or iv) R^{21} is C_3 - C_5 cycloalkyl; or v) R^{21} is C_1 - C_2 alkyl.


38. The compound of any of claims 32-37, wherein i) R^{22} is H or C_1 - C_2 alkyl or ii) R^{22} is H; or iii) R^{22} is C_1 - C_2 alkyl; or iv) R^{22} is -CH₃.

39. The compound of any of claims 32-38, wherein i) R^{24} is C_1 - C_4 alkyl or -(C_1 - C_4 alkyl)_t-O-(C_1 - C_4 alkyl) or ii) R^{24} is -(C_1 - C_2 alkyl)_t-O-(C_1 - C_2 alkyl).





40. The compound of any of claims 32-39, wherein L^3 is -N(CH₃)-.

41. The compound of any of claims 32-40, wherein i) n is 2 and m is 2; or ii) n is 1 or 2; or iii) n is 1 and m is 2.

42. A compound having Structural Formula IV:



IV, or pharmaceutically acceptable salts thereof, wherein:



Het is a 5- to 6-membered heteroaryl;

R¹ is H, -CN, halogen, C₁-C₄ alkyl, -O-(C₃-C₅ cycloalkyl), -O-(4- to 6-membered heterocycle) or -O-(C₁-C₄ alkyl), wherein when R¹ is not H, -CN or halogen, R¹ is optionally substituted with one or more halogens;

each R² is independently hydrogen, halogen or C₁-C₄ alkyl;

R³ is H or F;

R¹¹ is H or -CH₃;

R²¹ is H, halogen, C₁-C₄ alkyl, C₃-C₅ cycloalkyl or 4- to 6-membered heterocycle;

R²² is H, halogen or C₁-C₂ alkyl; and

R²⁴ is H, C₁-C₄ alkyl, -(C₁-C₄ alkyl)-OH, -(C₁-C₄ alkyl)-N(R²⁴¹)₂, -(C₁-C₄ alkyl)_t-O-(C₃-C₅


cycloalkyl), $-(C_1-C_4 \text{ alkyl})_t-O_u-(4\text{- to } 6\text{-membered heterocycle})$ or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$, wherein:

t is 0 or 1;

u is 0 or 1;

with the proviso that when u is 1, t is 1; and

R^{241} is H or C_1-C_2 alkyl.



44. The compound of claim 42 or 43, wherein i) R^1 is halogen, $-CN$ or C_1-C_2 haloalkyl; or ii) R^1 is $-CN$.

45. The compound of any of claims 42-44, wherein R^2 is H.

46. The compound of any of claims 42-45, wherein i) R^{21} is halogen, C_1-C_4 alkyl, C_3-C_5 cycloalkyl or 4- to 6-membered heterocycle; or ii) R^{21} is C_1-C_2 alkyl or C_3-C_5 cycloalkyl; or iii) R^{21} is C_1-C_2 alkyl; or iv) R^{21} is C_3-C_5 cycloalkyl.

47. The compound of any of claims 42-46, wherein i) R^{22} is H or C_1-C_2 alkyl; or ii) R^{22} is H; or iii) R^{22} is C_1-C_2 alkyl; or iv) R^{22} is $-CH_3$.

48. The compound of any of claims 42-47, wherein i) R^{24} is C_1-C_4 alkyl or $-(C_1-C_4 \text{ alkyl})-O-(C_1-C_4 \text{ alkyl})$; or ii) R^{24} is $-(C_1-C_2 \text{ alkyl})-O-(C_1-C_2 \text{ alkyl})$.

49. A pharmaceutical composition comprising any of the compounds of claims 1-48 and a pharmaceutically acceptable carrier, excipient, or diluent.

50. Use of a therapeutically effective amount of any of the compounds of claims 1-48 or a pharmaceutical composition of claim 49 in the manufacture of a medicament for treating a viral infection in a subject.

51. The use of claim 50, wherein the viral infection comprises hepatitis C infection.

52. Use of a therapeutically effective amount of any of the compounds of claims 1-48 or a pharmaceutical composition of claim 49 in the manufacture of a medicament for treating a metabolic disorder in a subject in need thereof.

53. Use of a therapeutically effective amount of any of the compounds of claims 1-48 or a pharmaceutical composition of claim 49 in the manufacture of a medicament for treating a cancer in a subject in need thereof.

54. The use of claim 53, wherein the cancer is breast cancer, pancreatic cancer, or colon cancer.

55. The use of claim 53, wherein the cancer is pancreatic cancer.