




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07D 217/02, 217/04, 217/06, 409/12, A61K 31/47	A1	(11) International Publication Number: WO 99/31068 (43) International Publication Date: 24 June 1999 (24.06.99)
(21) International Application Number: PCT/GB98/03785 (22) International Filing Date: 16 December 1998 (16.12.98) (30) Priority Data: 9726695.1 17 December 1997 (17.12.97) GB (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): COULTON, Steven [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). PORTER, Roderick, Alan [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). (74) Agent: WEST, Vivien; SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>

(54) Title: SUBSTITUTED ISOQUINOLINE DERIVATIVES AND THEIR USE AS ANTICONVULSANTS

**(57) Abstract**

Compounds of formula (I) or pharmaceutically acceptable salts or solvates thereof, where Q is a monocyclic or bicyclic aryl or heteroaryl ring, R¹ is hydrogen, C₁₋₆alkyl (optionally substituted by hydroxy or C₁₋₄alkoxy), C₁₋₆alkenyl, C₁₋₆alkynyl, C₁₋₆alkylCO-formyl, CF₃CO- or C₁₋₆alkylSO₂-, R² is hydrogen or up to three substituents selected from halogen, NO₂, CN, N₃, CF₃O-, CF₃S-, CF₃CO-, trifluoromethyl-diaziriny, C₁₋₆alkyl, C₁₋₆alkenyl, C₁₋₆alkynyl, C₁₋₆perfluoroalkyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₄alkyl-, C₁₋₆alkylO-, C₁₋₆alkylCO-, C₃₋₆cycloalkylO-, C₃₋₆cycloalkylCO-, C₃₋₆cycloalkyl-C₁₋₄alkylO-, C₃₋₆cycloalkyl-C₁₋₄alkylCO-, phenyl, phenoxy, benzyloxy, benzoyl, phenyl-C₁₋₄alkyl-, C₁₋₆alkylS-, C₁₋₆alkylSO₂-, (C₁₋₄alkyl)₂NSO₂-, (C₁₋₄alkyl)NHSO₂-, (C₁₋₄alkyl)₂NCO-, (C₁₋₄alkyl)NHCO-, CONH₂, CF₃SO₂, C₁₋₆alkenyl, C₁₋₆alkynyl or C₁₋₆hydroxyalkyl; or -NR³R⁴ where R³ is hydrogen or C₁₋₄alkyl, and R⁴ is hydrogen, C₁₋₄alkyl, formyl, -CO₂C₁₋₄alkyl or -COC₁₋₄alkyl; or two R² groups together form a carbocyclic ring that is saturated or unsaturated and unsubstituted or substituted by -OH or =O; and X is halogen, C₁₋₆alkoxy, C₁₋₆alkyl, C₂₋₆alkenyl optionally substituted by phenyl; are useful in the treatment and/or prophylaxis of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntington's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia and narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesis in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, or amyotrophic lateral sclerosis (ALS).

SUBSTITUTED ISOQUINOLINE DERIVATIVES AND THEIR USE AS ANTICONVULSANTS


This invention relates to novel compounds, to processes for preparing them, and to their use as therapeutic agents.

5

US-A-4022900 (Marion) discloses benzamido-tetrahydroisoquinolines having anti-hypertensive and vasodilator properties.

WO97/48683 (SmithKline Beecham) discloses that benzamide compounds of formula (A) below possess anti-convulsant activity and are therefore believed to be useful in the treatment and/or prevention of anxiety, mania, and related depression disorders.

10



15

(A)

where n and p are independently integers from 1 to 4 and $(n+p)$ is from 2 to 5;

R^1 is C_{1-6} alkylo-;

R^2 is hydrogen, halogen, CN, N_3 , trifluoromethyldiazirinyl, CF_3 , CF_3O -,
 20 CF_3S -, CF_3CO -, C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl- C_{1-4} alkyl-,
 C_{1-6} alkylo-, C_{1-6} alkylCO-, C_{3-6} cycloalkylCO-,
 C_{3-6} cycloalkyl- C_{1-4} alkylCO-, phenyl, phenoxy, benzyloxy, benzoyl,
 phenyl- C_{1-4} alkyl-, C_{1-6} alkylS-, C_{1-6} alkylSO₂-, $(C_{1-4}$ alkyl)₂NSO₂- or
 $(C_{1-4}$ alkyl)NHSO₂-;

R^3 is hydrogen, halogen, NO₂, CN, N_3 , trifluoromethyldiazirinyl,
 C_{1-6} alkylo-, C_{1-6} alkylS-, C_{1-6} alkyl, C_{3-6} cycloalkyl,
 C_{3-6} cycloalkyl- C_{1-4} alkyl-, C_{1-6} alkenyl, C_{1-6} alkynyl, CF_3CO -,
 C_{1-6} alkylCO-, C_{3-6} cycloalkylCO-, C_{3-6} cycloalkyl- C_{1-4} alkylCO-,
 phenyl, phenoxy, benzyloxy, benzoyl, phenyl- C_{1-4} alkyl-,
 or -NR⁵R⁶ where R⁵ is hydrogen or C_{1-4} alkyl, and

30

R^6 is hydrogen, C_{1-4} alkyl, -CHO, -CO₂ C_{1-4} alkyl or -COC C_{1-4} alkyl;

R^4 is hydrogen, C_{1-6} alkyl, C_{1-6} alkenyl, or C_{1-6} alkynyl;

and also the compounds: N-(7-iodo-2-methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-5-benzoyl-2-methoxybenzamide, N-(7-iodo-1,2,3,4-tetrahydroisoquinolin-5-yl)-5-benzoyl-2-methoxybenzamide, N-(5-iodo-1,2,3,4-tetrahydroisoquinolin-7-yl)-5-benzoyl-2-methoxybenzamide, N-(5-iodo-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methoxy-4-

35

trifluoromethyldiazirinybenzamide, N-(5-iodo-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methoxy-5-trifluoromethyldiaziriny-benzamide, N-(7-iodo-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-methoxy-5-trifluoromethyldiaziriny-benzamide and N-(8-fluoro-2-methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-4-*t*-butyl-2-methoxybenzamide.


5

It has now been surprisingly found that tetrahydroisoquinoliny-carboxamide compounds of formula (I) below possess anti-convulsant activity and are believed to be useful in the for the treatment and/or prophylaxis of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, and amyotrophic lateral sclerosis (ALS).

20

Accordingly, the present invention provides a compound of formula (I) or pharmaceutically acceptable salt thereof:

25



where Q is a monocyclic or bicyclic aryl or heteroaryl ring,

R¹ is hydrogen, C₁₋₆ alkyl (optionally substituted by hydroxy or C₁₋₄alkoxy),
 30 C₁₋₆alkenyl, C₁₋₆alkynyl, C₁₋₆alkylCO-, formyl, CF₃CO- or C₁₋₆alkylSO₂-,
 R² is hydrogen or up to three substituents selected from halogen, NO₂, CN, N₃,
 CF₃O-, CF₃S-, CF₃CO-, trifluoromethyldiaziriny, C₁₋₆alkyl, C₁₋₆alkenyl,
 C₁₋₆alkynyl, C₁₋₆perfluoroalkyl, C₃₋₆cycloalkyl,
 35 C₃₋₆cycloalkyl-C₁₋₄alkyl-, C₁₋₆alkylO-, C₁₋₆alkylCO-,
 C₃₋₆cycloalkylO-, C₃₋₆cycloalkylCO-, C₃₋₆cycloalkyl-C₁₋₄alkylO-,


C_{3-6} cycloalkyl- C_{1-4} alkylCO-, phenyl, phenoxy, benzyloxy, benzoyl,
 phenyl- C_{1-4} alkyl-, C_{1-6} alkylS-, C_{1-6} alkylSO₂-, (C_{1-4} alkyl)₂NSO₂-,
 (C_{1-4} alkyl)NHSO₂-, (C_{1-4} alkyl)₂NCO-, (C_{1-4} alkyl)NHCO, CONH₂,
 CF₃SO₂, C_{1-6} alkenyl, C_{1-6} alkynyl or C_{1-6} hydroxyalkyl;
 or -NR³R⁴ where R³ is hydrogen or C_{1-4} alkyl, and
 R⁴ is hydrogen, C_{1-4} alkyl, formyl, -CO₂ C_{1-4} alkyl or -COC C_{1-4} alkyl;
 or two R² groups together form a carbocyclic ring that is saturated or
 unsaturated and unsubstituted or substituted by -OH or =O; and
 X is halogen, C_{1-6} alkoxy, C_{1-6} alkyl, C_{2-6} alkenyl optionally substituted by
 phenyl.

When compounds of the present invention possess chiral centres and as such may exist in
 different enantiomeric forms, the present invention extends to each enantiomeric form and
 mixtures thereof including diastereoisomers and racemates. Alkenyl compounds exist as
 geometric isomers and the present invention extends to each isomeric form and mixtures
 thereof.


The ring system Q is typically optionally substituted phenyl or optionally substituted
 thiophenyl. When two R² groups form a carbocyclic ring, this is typically a 5-7 membered
 ring, so that Q may be a naphthalene or an indane or indanone ring system.

In the formula (I), alkyl groups, including alkyl groups that are part of other moieties, such
 as alkoxy or acyl, may be straight chain or branched. Phenyl groups, including phenyl
 groups that are part of other moieties, in R² may optionally be substituted with one or more
 independently selected halogen or C_{1-6} alkyl, C_{1-6} alkoxy or C_{1-6} alkylcarbonyl.
 Suitable C_{3-6} cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, and
 cyclohexyl. Suitable halo substituents include fluoro, chloro, iodo and bromo.

One suitable group of compounds of this invention are of formula (IA)



and another suitable group are of formula (IB)



A suitable group of compounds of formula (I) have

- R^1 as hydrogen, methyl, ethyl, propyl, hydroxyethyl, formyl, acetyl, trifluoroacetyl or methanesulfonyl,
- 5 R^2 as hydrogen or one or more of methyl, ethyl, *n*-butyl, *iso*-propyl, *t*-butyl, phenyl, methoxy, ethoxy, *iso*-propoxy, *n*-butoxy, cyclopropylmethoxy, phenoxy, benzyloxy, amino, acetylamino, nitro, azido, cyano, bromo, chloro, fluoro, iodo, acetyl, propionyl, pivaloyl, *iso*-butyryl, benzoyl, iodobenzoyl, trifluoromethyl, perfluoroethyl, trifluoromethoxy,
- 10 trifluoroacetyl, trifluoromethyldiaziriny, methanesulfonyl, *n*-propylsulfonyl, isopropylsulfonyl, dimethylsulfamoyl, CF_3SO_2 ;
- or two groups R^2 form a benzene, cyclopentane or cyclopentanone ring;
- X as methyl, ethyl, chloro, bromo, iodo, fluoro, phenylethenyl.

15 A preferred group of compounds of formula (I) have

- R^1 as hydrogen, methyl, hydroxyethyl, formyl or trifluoroacetyl.
- R^2 as hydrogen or one or more of methyl, ethyl, *iso*-propyl, methoxy, ethoxy, *iso*-propoxy, acetyl, propionyl, pivaloyl, cyano, bromo, chloro, fluoro, iodo, trifluoromethyl.
- 20 X as methyl, ethyl, chloro or bromo.

In a special class of compounds of formula (I), suitable for use as mechanistic probes, R^2 groups are photolabile groups, such as N_3 , benzoyl and trifluoromethyldiaziriny. Also radiolabels such as ^{125}I can be incorporated at R, and 3H can be located at suitable

25 positions.

Examples of compounds of formula (I) are:

- 30 N-(8-chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-ethoxybenzamide
- N-(8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-ethoxybenzamide
- N-(8-chloro-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-ethoxybenzamide
- N-(8-chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- 35 N-(8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

- N-(8-chloro-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- N-(8-bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- 5 N-(8-bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
N-(8-bromo-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
N-(2,8-Dimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- 10 N-(2-(2-Hydroxyethyl)-8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
N-(2-formyl-8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
N-(8-ethyl-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- 15 N-(8-ethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
N-(8-ethyl-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
N-(8-chloro-2-methanesulfonyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- 20 N-(8-Styryl-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
N-(8-Styryl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
N-(8-Styryl-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- 25 N-(8-chloro-2-methanesulfonyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide hydrochloride
N-(8-Bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-acetyl-4-ethoxybenzamide
N-(8-Bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-acetyl-4-ethoxybenzamide
N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-fluoro-4-methoxybenzamide
- 30 N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-chloro-4-ethoxybenzamide
N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-methoxybenzamide
N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-acetyl-4-ethoxybenzamide
N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-acetyl-4-*iso*-propoxybenzamide..
- 35 When synthesised, these compounds are often in salt form, such as the hydrochloride or trifluoroacetate, and such salts also form part of this invention. Such salts may be used in preparing pharmaceutically acceptable salts. The compounds and their salts may be obtained as solvates, such as hydrates, and these also form part of this invention.

The above compounds and pharmaceutically acceptable salts thereof, especially the hydrochloride, and pharmaceutically acceptable solvates, especially hydrates, form a preferred aspect of the present invention.

- 5 The administration of such compounds to a mammal may be by way of oral, parenteral, sub-lingual, nasal, rectal, topical or transdermal administration.

An amount effective to treat the disorders hereinbefore described depends on the usual factors such as the nature and severity of the disorders being treated and the weight of the
10 mammal. However, a unit dose will normally contain 1 to 1000 mg, suitably 1 to 500 mg, for example an amount in the range of from 2 to 400 mg such as 2, 5, 10, 20, 30, 40, 50, 100, 200, 300 and 400 mg of the active compound. Unit doses will normally be administered once or more than once per day, for example 1, 2, 3, 4, 5 or 6 times a day, more usually 1 to 4 times a day, such that the total daily dose is normally in the range, for a
15 70 kg adult of 1 to 1000 mg, for example 1 to 500 mg, that is in the range of approximately 0.01 to 15 mg/kg/day, more usually 0.1 to 6 mg/kg/day, for example 1 to 6 mg/kg/day.

It is greatly preferred that the compound of formula (I) is administered in the form of a unit-dose composition, such as a unit dose oral, including sub-lingual, rectal, topical or
20 parenteral (especially intravenous) composition.

Such compositions are prepared by admixture and are suitably adapted for oral or parenteral administration, and as such may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable and
25 infusible solutions or suspensions or suppositories. Orally administrable compositions are preferred, in particular shaped oral compositions, since they are more convenient for general use.

Tablets and capsules for oral administration are usually presented in a unit dose, and
30 contain conventional excipients such as binding agents, fillers, diluents, tableting agents, lubricants, disintegrants, colorants, flavourings, and wetting agents. The tablets may be coated according to well known methods in the art. Suitable fillers for use include cellulose, mannitol, lactose and other similar agents. Suitable disintegrants include starch, polyvinylpyrrolidone and starch derivatives such as sodium starch glycollate. Suitable
35 lubricants include, for example, magnesium stearate. Suitable pharmaceutically acceptable wetting agents include sodium lauryl sulphate.

These solid oral compositions may be prepared by conventional methods of blending, filling, tableting or the like. Repeated blending operations may be used to distribute the

active agent throughout those compositions employing large quantities of fillers. Such operations are, of course, conventional in the art.

5 Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethyl cellulose, aluminium stearate gel or hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan
10 monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example, almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl *p*-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents. Oral formulations also include conventional sustained release formulations, such as tablets or granules having
15 an enteric coating.

For parenteral administration, fluid unit dose forms are prepared containing the compound and a sterile vehicle. The compound, depending on the vehicle and the concentration, can be either suspended or dissolved. Parenteral solutions are normally prepared by dissolving
20 the compound in a vehicle and filter sterilising before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are also dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum.

25 Parenteral suspensions are prepared in substantially the same manner except that the compound is suspended in the vehicle instead of being dissolved and sterilised by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound of the invention.

30 As is common practice, the compositions will usually be accompanied by written or printed directions for use in the medical treatment concerned.

Accordingly, the present invention further provides a pharmaceutical composition for use
35 in the treatment and/or prophylaxis of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis,

migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, or amyotrophic lateral sclerosis (ALS) which comprises a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier.

The present invention also provides a method of treatment and/or prophylaxis of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, or amyotrophic lateral sclerosis (ALS) comprising administering to the sufferer in need thereof an effective or prophylactic amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof.

In a further aspect the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, for the manufacture of a medicament for the treatment and/or prophylaxis of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury,

tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, or amyotrophic lateral sclerosis (ALS).

5


In a further aspect the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate, thereof as a therapeutic agent, in particular for the treatment and/or prophylaxis of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, or amyotrophic lateral sclerosis (ALS).

10

15

20

Another aspect of the invention is a process for the preparation of compounds of formula (I), which comprises reacting a compound of formula (II)



25

(II)

where R^{1A} is R^1 as defined for formula (I) or a group convertible to R^1 and X^A is X as defined in claim 1 or a group convertible to X

30

with a compound of formula (III)



(III)

where Q is as defined in formula (I), Y is Cl or OH, and R^{2A} groups are independently R^2 as defined for formula (I) or groups convertible to R^2 ,
 and where required converting an X^A , R^{1A} or R^{2A} group to an X, R^1 or R^2 group,
 5 converting one X, R^1 or R^2 group to another X, R^1 or R^2 group,
 converting a salt product to the free base or another pharmaceutically acceptable salt, or
 converting a free base product to a pharmaceutically acceptable salt.


Conventional conditions for condensation of amines with carboxylic acids or active
 10 derivatives thereof, such as acid chlorides, may be used. For example the amides and acids
 may be reacted in the presence of a mixture of ethyl(dimethylaminopropyl)-
 carbodiimide/hydroxybenzotriazole in a suitable solvent such as dimethyl formamide, and
 amines and acid chlorides may be reacted together in a suitable solvent such as ethyl
 acetate or dichloromethane, optionally in the presence of a base such as triethylamine.

15 Reaction of a compound of formula (III) which is an acid chloride ($Y=Cl$) typically results
 in formation of the hydrochloride salt of the compound of formula (I). Hydrochloride salts
 can also be obtained by passing HCl gas into a solution of the free base, or adding a
 solution of HCl in ether.

20 Conversions of an R^{1A} or R^{2A} group to a R^1 or R^2 group typically arise when a protecting
 group is needed during the above coupling reaction or during the preparation of the
 reactants by the procedures described below. Interconversion of one R^1 or R^2 group to
 another typically arises when one compound of formula (I) is used as the immediate
 25 precursor of another compound of formula (I) or when it is easier to introduce a more
 complex or reactive substituent at the end of a synthetic sequence.


Compounds of formula (II) may be prepared from a nitro-tetrahydroisoquinoline of formula
 (IV).

30



by reaction with a compound $R^{1A}Z$ where Z is a leaving group such as halogen, especially
 iodo, or tosylate to obtain an intermediate of formula (V)

35




(V)

- 5 which can be hydrogenated, for example using either tin (II) chloride and HCL or hydrogen and a palladium/activated carbon catalyst, to obtain an amino-tetrahydroisoquinoline of formula (II).

- 10 When the intended R^{1A} group is methyl, the compound of formula (IV) may also be reacted with formic acid and formaldehyde to introduce the N-methyl group.

- 15 The nitro-tetrahydroisoquinoline of formula (IV) in which X^A is hydrogen may be prepared by hydrolysis of 2-trifluoroacetyl-7-nitro-tetrahydroisoquinoline obtained by reaction of an N-(nitrophenyl)ethyl-trifluoroacetamide and paraformaldehyde in acidic conditions using the procedure of Stokker, Tet.Lett.,1996, 37, 5453. N-(nitrophenyl)ethyl-trifluoroacetamides can be prepared from readily available materials by reaction of trifluoroacetic anhydride with lutidine and nitrophenethylamine hydrochloride, as illustrated in the Descriptions below.

- 20 Compounds of formula (II) may also be prepared from the corresponding amino-isoquinoline (or its nitro-analogue) of formula (VI)




(VI)

25

where R^N is NH₂ or NO₂

by reaction with a compound R^{1A}Z where Z is a leaving group such as halogen, especially iodo, or tosylate to obtain an intermediate of formula (VII)



30

(VII)

which can be reduced, for example using sodium borohydride, or hydrogenated, for example using hydrogen and a palladium/activated carbon catalyst, to obtain a tetrahydroisoquinoline of formula (II). When the compound of formula (VII) is replaced by a nitro-isoquinoline, the nitro group is converted to an amino group in the hydrogenation step.

When the intended R^1 is hydrogen, the N of the tetrahydroisoquinoline or isoquinoline is preferably protected conventionally, prior to the coupling step that forms the carboxamide of formula (I), for example by *tert.*-butoxycarbonyl or trifluoroacetyl. The compound can be deprotected under standard conditions, for example using trifluoroacetic acid/methylene chloride or potassium carbonate in aqueous methanol.

Amino/nitro-isoquinolines of formulae (VI) and the reagents used are commercially available, or can be prepared from commercially available materials using conventional procedures described in the literature.

The substituent X^A may be a group X already present on commercially available starting materials usable in the above described procedures. When the substituent X^A is a group convertible to X, then X may be introduced during any of the procedures above, for example by conventional substitution of the aromatic ring of compounds of formula (IV), (V) or (VII). Most suitably the substituent X as halogen is introduced to a compound of formula (II) in which X^A is hydrogen. For example, X as halogen may be incorporated by reaction with a halo-succinimide as illustrated in the Descriptions below. As a further example, X as alkyl may be introduced by reaction of a compound of formula (II) in which X is bromo with an alkyl stannane, as illustrated in the Descriptions and Examples below. The compound of formula (II) in which X is bromo may be obtained by halogenation of an amino-tetrahydroisoquinoline of formula (II) in which X^A is hydrogen with NBS, again using procedures illustrated in the Descriptions below.

Compounds of formula (III) may be prepared by further substitution of commercially available benzoic acid or thiophene carboxylic acid derivatives using conventional procedures, or by oxidation of corresponding substituted benzyl alcohols. Alternatively benzoic acids can be prepared from correspondingly substituted phenols, for example by formation of the acetate, conversion to an acetophenone and then to the desired acid.

Examples of these procedures are documented in WO 98/41507 and WO98/41508.

Where the above described intermediates are novel compounds, they also form part of this invention.

The preparation of compounds of this invention is further illustrated by the following Descriptions and Examples. The utility of compounds of this invention is shown by the Pharmacological Data that follow the Examples.

5

Description 1**N-2-(4-Nitrophenyl)ethyl-trifluoroacetamide**

A solution of trifluoroacetic anhydride (10.6ml) in dichloromethane (100ml) was added dropwise to a stirred solution of 2,6-lutidine (17.44ml) and 4-nitrophenethylamine hydrochloride (15.2g; 75 mmol) at 0°C. The mixture was stirred at 25°C overnight under argon and then washed with dilute citric acid (x2), brine and dried over Na₂SO₄. The material in the organic phase gave the title compound D1 as a pale yellow solid (19.04g).

Description 2**7-Nitro-1,2,3,4-tetrahydro-2-trifluoroacetylisoquinoline**

The compound D1 (2.26g; 9.15 mmol) and paraformaldehyde (0.45g; 14.4 mmol) in acetic acid (10ml) and conc. H₂SO₄ (15ml) were stirred at 25°C for 20h according to the procedure of G.E. Stokker., Tet. Lett., 1996, 37, 5453. Work up afforded the title compound D2 as a white solid (2.17g).

20

¹H NMR (CDCl₃) δ: 3.10 (2H, m), 3.92 (2H, m), 4.85 + 4.92 (2H, 2xs), 7.38 (1H, t), 8.10 (2H, m); m/z (EI): 274 (M⁺)

Description 3**7-Nitro-1,2,3,4-tetrahydroisoquinoline**

The compound D2 (17.22g; 63 mmol) was hydrolysed at room temperature using a solution of potassium carbonate (46.6g) in 10% aqueous methanol (660ml). Work-up with dichloromethane gave the title compound (11g).

30

Description 3**7-Amino-1,2,3,4-tetrahydro-2-trifluoroacetylisoquinoline**

The 7-nitro compound D2 (0.99g; 3.6 mmol) in ethanol (50ml) was hydrogenated over 10% palladium on carbon (450mg) at atmospheric pressure for 4h. The catalyst was removed by filtration through a pad of Celite and evaporation *in vacuo* gave the title compound as a white solid (840mg).

35

¹H NMR (250MHz, CDCl₃) δ: 2.84 (2H, t), 3.23 (2H, br s), 3.82 (2H, m), 4.66 (2H, d, restricted rotation around C-1), 6.47 (1H, m), 6.57 (1H, m), 6.96 (1H, m)

Description 4**7-Amino-8-chloro-1,2,3,4-tetrahydro-2-trifluoroacetylisoquinoline**

To a solution of amine D3 (1.00g) in acetonitrile (20ml) N-chlorosuccinimide (0.60g) was added and the solution stirred at room temperature for 6 days. The solution was diluted with ethyl acetate, washed with water and the organic phase dried (MgSO₄) and solvent removed at reduced pressure. The residue was column chromatographed (silica gel, dichloromethane then 2% methanol/dichloromethane) to give 7-amino-8-chloro-1,2,3,4-tetrahydro-2-trifluoroacetyl-isoquinoline (0.72g) as a pale yellow solid.

10

¹H NMR (250MHz, CDCl₃) δ: 2.85 (2H, m), 3.83 (2H, dt, restricted amide rotation), 4.76 (2H, s), 6.68 (1H, m) and 6.89 (1H, m).

Description 5**7-Amino-8-bromo-1,2,3,4-tetrahydro-2-trifluoroacetyl-isoquinoline**

The title compound (0.27g) was prepared from amine D3 (0.24g) and N-bromosuccinimide (0.20g) according to the method of Description 4.

¹H NMR (250MHz, CDCl₃) δ: 2.85 (2H, m), 3.76 - 3.87 (2H, m, restricted amide rotation), 4.72 (2H, d due to restricted amide rotation), 6.68 (1H, m) and 6.93 (1H, m).

20

Preparation 1**3-Bromo-4-ethoxybenzoic acid**

To a solution of 4-ethoxybenzoic acid (3.6g, 0.022mol) in chloroform (60ml) was added bromine (1.13ml, 0.022mol) in chloroform (20ml) dropwise. After stirring overnight at room temperature the precipitate was filtered off and dried to afford the title compound as a white solid.

¹H NMR (DMSO-D₆) δ: 1.45 (3H, t, J = 7 Hz), 4.26 (2H, q, J = 7 Hz), 7.26 (1H, d, J = 9 Hz), 7.98 (1H, dd, J = 2, 9 Hz), 8.12 (1H, d, J = 2 Hz)

30

Preparation 2**4-Methoxy-3-trifluoromethylbenzoic acid**

3-Bromo-4-ethoxybenzoic acid methyl ester (1.4g; 5.4 mmol) in DMF (25ml) and toluene (8ml) under argon was treated with potassium trifluoroacetate (1.53g; 10.1 mmol) and copper (I) iodide (2.1g, 10.9 mmol). The mixture was heated to 170°C with removal of water (Dean/Stark), and then at 155°C overnight. The mixture was allowed to cool, poured into ether and water and filtered through Kieselguhr. The organic layer was dried (Na₂SO₄)

35

and concentrated *in vacuo* to give a brown solid. Chromatography on Kieselgel 60 with 1:1 ether/petrol gave a solid which was hydrolysed in 1:1 methanolic: aqueous NaOH (50ml) at 50°C. Work-up gave the title compound as a white solid (1g).

- 5 ¹H NMR (DMSO-D₆) δ: 3.78 (3H, s), 7.18 (1H, d, J = 9 Hz), 7.90 (1H, d, J = 2 Hz), 8.00 (1H, dd, J = 2, 9 Hz), 12.70 - 13.10 (1H, br,exchangeable)

Preparation 3

4-Methoxy-3-trifluoromethylbenzoyl chloride

- 10 The title compound was prepared from 4-methoxy-3-trifluoromethylbenzoic acid with oxalyl chloride and DMF in chloroform at room temperature [D. Levin, Chem. Br., 1977, 20] followed by evaporation *in vacuo*.

Preparation 4

- 15 3-Chloro-4-ethoxybenzoic acid

¹H NMR (DMSO-D₆) δ: 1.39 (3H, t, J = 7 Hz), 4.20 (2H, q, J = 7 Hz), 7.22 (1H, d, J = 7 Hz), 7.87 (2H, m).

- 20 Preparation 5

Methyl 3-Bromo-4-*iso*-propoxybenzoate

Methyl 3-bromo-4-hydroxybenzoate (2.5g, 10.8mmol) in DMF (35ml) was treated with potassium carbonate (3.0g, 21.6mmol), 2-iodopropane (2.76, 21.6mmol) and then stirred at 25°C for 48h. Work-up with ethyl acetate gave the title compound (3.0g).

25

¹H NMR (250MHz, CDCl₃) δ: 1.41 (6H, d, J=7 Hz), 3.89 (3H, s), 4.66 (1H, m), 6.90 (1H, d, J = 8 Hz), 7.93 (1H, dd, J = 8, 2 Hz), 8.22 (1H, d, J = 2 Hz)

Preparation 6

- 30 Methyl 3-Acetyl-4-*iso*-propoxybenzoate

The bromo ester P5 (2.5g, 8.3mmol) in dry dioxan (30ml) was treated with (1-ethoxyvinyl)-tributyl tin (3.58g, 9.9mmol) followed by tetrakis triphenylphosphine palladium(o) (0.48g, 0.4mmol) and heated at 100° for 18h. After cooling, the mixture was acidified and aqueous work-up and extraction into ethyl acetate gave a coloured oil (5.6g).

- 35 Flash chromatography on Kieselgel 60 [hexane to 20% EtAc/hexane] gave the title compound as a yellow oil (2.3g).

Preparation 7

3-Acetyl-4-*iso*-propoxybenzoic acid

Saponification of the ester **P6** (2.3g) gave the title compound as a white solid (1.3g).

¹H NMR (250MHz, CDCl₃) δ: 1.48 (6H, d, J=7 Hz), 2.63 (3H, s), 4.80 (1H, m), 7.00 (1H, d, J = 8 Hz), 8.17 (1H, dd, J = 8, 2 Hz), 8.46 (1H, d, J = 2 Hz)

Preparation 8

3-Acetyl-4-ethoxybenzoic acid

Prepared in a similar manner to that described for Preparations 6 and 7.

¹H NMR (250MHz, CDCl₃) δ: 1.53 (3H, t, J=7 Hz), 2.65 (3H, s), 4.23 (2H, q, J=7 Hz), 7.01 (1H, d, J = 8 Hz), 8.19 (1H, dd, J = 8, 2 Hz), 8.48 (1H, d, J = 2 Hz).

Example 1

N-(8-Chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-ethoxybenzamide

To a solution of amine **D4** (0.28g) in tetrahydrofuran (5ml) containing triethylamine (0.21ml), 3-bromo-4-ethoxybenzoyl chloride (0.37g) in tetrahydrofuran (7ml) was added. The mixture was stirred overnight partitioned between ethyl acetate and water and the organic phase dried (MgSO₄) and solvent removed at reduced pressure. The residue was purified by column chromatography (silica gel, and an ethyl acetate/hexane gradient to give after combining of appropriate fractions the title compound (0.31g) as a colourless solid.

¹H NMR (250MHz, CDCl₃) δ: 1.26 (3H, t), 2.98 (2H, m), 3.85 (2H, dt, restricted amide rotation), 4.11 (2H, q), 4.80 (2H, d due to restricted amide rotation) 6.98 (1H, d), 7.18 (1H, t), 7.84 (1H, dd), 8.14 (1H, d), 8.28 (1H, m) and 8.39 (1H, d); ^{m/z} (API): 507 (MH⁺; 100%)

Example 2

N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-ethoxybenzamide

A solution of N-(8-chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-ethoxybenzamide (0.28g) in methanol/water (5ml 9:1) was treated with potassium carbonate (0.38g) and stirred 12h. The mixture was diluted with dichloromethane and washed with water. The organic phase was dried (MgSO₄) solvent removed at reduced

pressure. The residue was chromatographed (silica gel, dichloromethane/methanol/-ammonia upto 9:1:0.1 eluant) to give the title compound (0.18g) as a colourless solid.

^1H NMR (250MHz, CDCl_3) δ : 1.52 (3H, t), 2.79 (2H, t), 3.11 (2H, t), 4.04 (2H, s) 4.20 (2H, q) 6.96 (1H, d, $J = 6\text{Hz}$), 7.09 (1H, d, $J = 8.5\text{Hz}$), 7.84 (1H, dd, $J = 8.5, 2\text{Hz}$), 8.13 (1H, d, $J = 2\text{Hz}$), 8.26 - 8.29 (2H, m); m/z (APD): 409, 411 (MH^+ ; 100%)

Example 3

10 N-(8-chloro-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-ethoxybenzamide

A solution of N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-ethoxybenzamide (0.12g) in 37% aqueous formaldehyde (0.63ml) and formic acid (0.34ml) and stirred at 80°C for 3h. Solid sodium hydroxide was added to neutralise the solution and the aqueous phase extracted with dichloromethane. The combined organic extracts were dried (MgSO_4) and solvent removed *in vacuo* to give the title compound (0.11g).

^1H NMR (250MHz, CDCl_3) δ : 1.52 (3H, t), 2.52 (3H, s), 2.68 (2H, t), 2.93 (2H, t), 3.61 (2H, s), 4.18 (2H, q), 6.96 (1H, d, $J = 6\text{Hz}$), 7.10 (1H, d, $J = 8.5\text{Hz}$), 7.84 (1H, dd, $J = 8.5, 2\text{Hz}$), 8.13 (1H, d, $J = 2\text{Hz}$), 8.25 - 8.28 (2H, m); m/z (APD): 423, 425 (MH^+ ; 100%)

Example 4

25 N-(8-Chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

The title compound (0.32g) was prepared from amine D4 (0.28g) and 4-methoxy-3-trifluoromethylbenzoyl chloride (0.33g) according to the procedure of Example 1.

^1H NMR (250MHz, CDCl_3) δ : 2.99 (2H, m), 3.85 (2H, dt, restricted amide rotation), 4.81 (2H, d due to restricted amide rotation) 7.12 - 7.21 (2H, m), 8.08 (1H, d), 8.16 (1H, s), 8.28 - 8.40 (2H, m); m/z (APD): 481 (MH^+ ; 100%)

35 Example 5

N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

The title compound (0.05g) was prepared from N-(8-chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.29g) according to the procedure of Example 2.

5

^1H NMR (250MHz, CDCl_3) δ : 2.79 (2H, t), 3.11 (2H, t), 4.00 (3H, s), 4.05 (2H, s), 7.11(2H, t), 8.09 (1H, d), 8.16 (1H, s), 8.26 (1H, d) and 8.31 (1H, br. s);
 m/z (API): 385 (MH^+ ; 100%)

10

Example 6

N-(8-Chloro-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

15

The title compound (0.13g) was prepared from N-(8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.19g) according to the procedure of Example 3.

20 ^1H NMR (250MHz, CDCl_3) δ : 2.53 (3H, s), 2.68 (2H, t), 2.92 (2H, t), 3.61 (2H, s) 4.00 (3H, s), 7.12(2H, d), 8.09 (1H, d), 8.15 (1H, s), 8.26 (1H, d) and 8.29 (1H, br. s);
 m/z (API): 385 (MH^+ ; 100%)

25

Example 7

N-(8-Bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

30 The title compound (0.32g) was prepared from amine D5 (0.22g) and 4-methoxy-3-trifluoromethylbenzoyl chloride (0.22g) according to the procedure of Example 1.

^1H NMR (250MHz, CDCl_3) δ : 2.94 - 3.01 (2H, m), 3.82 - 3.93 (2H, m, restricted amide rotation), 4.01 (3H, s), 4.78 (2H, d due to restricted amide rotation) 6.95 (1H, d), 7.14 (1H, d), 8.04 (1H, d), 8.18 (1H, s), 8.36 (1H, d) and 8.39 (1H, s);
35 m/z (API): 524, 526 (MH^+ ; 100%).

Example 8

N-(8-Bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

- 5 The title compound (0.12g) was prepared from N-(8-bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.24g) according to the procedure of Example 2.

¹H NMR (250MHz, CDCl₃) δ: 2.79 (2H, t), 3.11 (2H, t), 4.00 (5H, s), 7.13 (2H, dd), 8.10 (1H, dd, J = 2.3 and 8.7Hz) 8.18 (1H, d, J = 2.1Hz), 8.25 (1H, d, J = 8.5Hz) and 8.38 (1H, s); m/z (API): 429, 431 (MH⁺; 100%)

Example 9

15

N-(8-Bromo-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

- 20 The title compound (0.08g) was prepared from N-(8-bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.09g) according to the procedure of Example 3.

¹H NMR (250MHz, CDCl₃) δ: 2.53 (3H, s), 2.67 (2H, t), 2.94 (2H, t), 3.59 (2H, s), 4.00 (3H, s), 7.15 (2H, t), 8.11 (1H, dd) 8.18 (1H, d), 8.25 (1H, d, J = 8.5Hz) and 8.37 (1H, s); m/z (API): 443, 445 (MH⁺; 100%)

25

Example 10

- 30 **N-(2,8-Dimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide**

a) N-(8-Bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.25g), was combined with lithium chloride (0.06g), tetramethyltin (0.08ml) and *bis*(triphenylphosphine)palladium(II)chloride (0.025g) in dimethylformamide (5ml) and the mixture warmed to 100°C for 24 h. Solvent was removed at reduced pressure, filtered (Celite pad) and the filtrate column chromatographed (silica gel, diethyl ether eluant) to give N-(8-methyl-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.16g).

35

¹H NMR (CDCl₃) δ: 2.31 (3H, m), 2.95 – 2.99 (2H, m), 3.81 – 3.90 (2H, m), 4.07 (3H, s), 4.70, 4.73 (2H, s, appears as 2 singlets due to restricted rotation), 7.04 – 7.12 (2H, m), 7.41 – 7.47 (1H, m), 7.63 (1H, br. s.), 8.08 – 8.11 (2H, m); m/z (API⁺): 461 (MH⁺)

- 5 b) N-(8-Methyl-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.15g) in methanol:2N sodium hydroxide(15ml 2:1) was stirred at room temperature for 1h. 2M hydrochloric acid (4.5ml) was added and solvent (10ml) removed at reduced pressure. The residual solvent was extracted with dichloromethane, the organic phase washed with brine, dried MgSO₄) and solvent removed at reduced
10 pressure to give N-(8-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.08g)

¹H NMR (CDCl₃) δ: 2.31 (3H, m), 2.07 (3H, s), 2.80 (2H, t), 3.09 (2H, t), 3.95 (2H, s), 3.98 (3H, s), 6.98 (1H, d), 7.09 (1H, d), 7.32 (1H, s), 7.69 (1H, br. s.), 8.07 – 8.10 (2H, m);
15 m/z (API⁺): 365 (MH⁺)

- c) The title compound (0.03g) was prepared from N-(8-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.08g), according to the method of Example 3.

20 ¹H NMR (CDCl₃) δ: 2.01 (3H, s), 2.48 (3H, s), 2.68 (2H, t), 2.95 (2H, t), 3.48 (2H, s), 3.99 (3H, s), 7.00 (1H, d), 7.09 (1H, d), 7.30 (1H, d), 7.92 (1H br. s.), 8.08 (1H, br. s.), 8.11 (1H, br. s.); m/z (API⁺): 379 (MH⁺)

25

Example 11

N-(2-(2-Hydroxyethyl)-8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

30

- a) A mixture of N-(8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.192g) and 2-bromoethoxy-*tert*-butyldimethylsilane (0.24g) were combined in dimethylformamide and warmed to 80°C for 18 h. Solvent was removed at reduced pressure, the residue dissolved in dichloromethane and washed with saturated
35 sodium hydrogen carbonate. The organic phase was dried (MgSO₄) and solvent removed at reduced pressure. The residue was column chromatographed (silica gel, ammonia/methanol/dichloromethane mixtures as eluant) to give N-(2-(2-*tert*-butyldimethylsilyloxyethyl)-8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.09g)

- b) N-(2-(2-*Tert*-butyldimethylsilyloxyethyl)-8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.09g) in tetrahydrofuran (10ml) was treated with tetrabutylammonium fluoride in tetrahydrofuran (1M, 0.16 ml) at room temperature overnight. The reaction mixture was diluted with ethyl acetate, washed with water, the organic phase dried (MgSO₄) and solvent removed at reduced pressure. The residue was column chromatographed (silica gel, ammonia/methanol/dichloromethane mixtures as eluant) to give the title compound (0.04g)
- ¹H NMR CDCl₃) δ: 2.80 (4H, m), 2.91 (2H, m), 3.73 (2H, s), 3.75 (2H, t), 3.99(3H, s), 7.11 (2H, dd), 8.07 (1H, dd), 8.16 (1H, d), 8.25 (1H, d) and 8.32 (1H, br. s.); m/z (API⁺): 429(MH⁺)

Example 12

N-(2-Formyl-8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

- Formic acetic anhydride (0.21g) was added to N-(8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.15g) and dimethylaminopyridine (0.05g) in dichloromethane. The mixture was stirred at room temperature overnight and quenched with saturated aqueous sodium hydrogen carbonate (20ml). The organic phase was separated and washed with brine, dried (MgSO₄) and solvent removed at reduced pressure. The residue was column chromatographed (silica gel, ammonia/methanol/dichloromethane mixtures as eluant) to give the title compound (0.10g)
- ¹H NMR (d⁶-DMSO, 353°K) δ: 3.79 and 2.90 (2H, t, rotamers), 3.68 (2H, m), 3.99 (3H, s), 4.55 and 4.62 (2H, s, rotamers), 7.23 (1H, d), 7.39 (1H, d), 7.44 (1H, d), 8.21, 8.27, 8.32 (3H, s), 10.20 (1H, s); m/z (API⁺): 413 (MH⁺)

Example 13

- #### **N-(8-Ethyl-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide**

N-(8-Bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.525g), lithium chloride (0.127g), tetraethyltin (0.47g) and

- bis*(triphenylphosphine)palladium(II)chloride (0.04g) were combined in dimethylformamide (10ml) and the mixture stirred under argon at 120°C for 20 h. Solvent was removed at reduced pressure, the residue dissolved in dichloromethane and filtered through Celite. Solvent was removed at reduced pressure and the residue column chromatographed (silica gel, ethyl acetate/hexane mixtures) to give the title compound (0.18g) as an oil.
- ¹H NMR (CDCl₃) δ: 1.24 (3H, t), 2.66 (2H, m), 2.99 (2H, m), 3.84 (2H, m), 4.00 (3H, s), 4.77 and 4.82 (2H, s, rotamers), 7.10 (2H, m), 7.62 (2H, m), 8.08 (2H, m);
- m/z (API⁺): 475 (MH⁺)

Example 14

- N-(8-Ethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide**

- N-(8-Ethyl-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.157g) was dissolved in methanol/water (5:1 6 ml), potassium carbonate (0.228g) added and the mixture stirred overnight. The mixture was partitioned between dichloromethane and water the organic phase separated and solvent removed at reduced pressure. The residue was column chromatographed (silica gel, ammonia/methanol/dichloromethane mixtures as eluant) to give the title compound (0.97g)
- ¹H NMR (CDCl₃) δ: 1.15 (3H, t), 2.57 (2H, q), 2.82 (2H, t), 3.10(2H, t), 3.99 (3H, s), 4.04 (2H, s), 7.00 (1H, d), 7.10 (1H, d), 7.49(1H, d), 7.62 (1H, br. s.), 8.05 – 8.10 (2H, m);
- m/z (API⁺): 379 (MH⁺)

Example 15

N-(8-Ethyl-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

- The title compound (0.08g) was prepared from N-(8-ethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.08g) according to the method of Example 3.

¹H NMR (CDCl₃) δ: 1.17 (3H, t), 2.56 (3H, s), 2.57 (2H, q), 2.68 (2H, t), 2.95 (2H, m), 3.61 (2H, s), 3.99 (3H, s), 7.04 (1H, d), 7.10 (1H, d), 7.51 (1H, d), 7.59 (1H, s), 8.04 – 8.09 (2H, m); m/z (API⁺): 393 (MH⁺)

5

Example 16

N-(8-Chloro-2-methanesulfonyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

10

N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.193g) in dichloromethane (10ml) containing triethylamine (0.055ml) was treated with methanesulfonyl chloride ((0.063g) and the mixture stirred for 48 h. Water was added and the organic phase separated and solvent removed at reduced pressure. The residue was trituated with diethyl ether to give the title compound (0.19g).

15

¹H NMR (CDCl₃) δ: 2.91 (3H, s), 3.00 (2H, t), 3.56 (2H, t), 4.01 (3H, s), 4.48 (2H, s), 7.13 (1H, d), 7.23 (1H, d), 8.08 (1H, dd), 8.16 (1H, s), 8.29 (1H, s), 8.35 (1H, d); m/z (API⁺): 463 (MH⁺)

20

Example 17

N-(8-Styryl-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

25

A mixture of N-(8-bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.525g), palladium diacetate (0.011g), tris(*o*-tolyl)phosphine (0.03g), triethylamine (0.15g) and styrene (0.16g) were combined in acetonitrile and warmed to reflux for 23 h. The mixture was cooled, partitioned between dichloromethane/water, the organic phase separated and solvent removed at reduced pressure. The residue was column chromatographed (silica gel, ethyl acetate/hexane mixtures as eluant) to provide the title compound (0.46g).

30

¹H NMR (CDCl₃) δ: 2.99 (2H, m), 3.85 (2H, m), 3.93, 4.01 (3H, s, rotamers), 4.75 (2H, m), 6.81 – 7.26 (4H, m), 7.39 – 7.53 (3H, m), 7.92 – 8.40 (4H, m); m/z (API⁺): 549 (MH⁺)

35

Example 18

N-(8-Styryl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

- 5 The title compound (0.08g) was prepared from N-(8-styryl-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.439g) according to the method of Example 2.

¹H NMR (CDCl₃) δ: 2.83 (2H, t), 3.12 (2H, t), 3.93 (3H, s), 3.99 (2H, s), 6.77 (1H, d), 6.94
10 (1H, d), 7.00 (1H, s), 7.11 (1H, d), 7.30 – 7.50 (5H, m), 7.93 – 7.99 (2H, m), 8.16 (1H, d), 8.26 (1H, s); m/z (API⁺): 453 (MH⁺)

Example 19

15

N-(8-Styryl-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide hydrochloride

- The title compound (0.033g) was prepared from N-(8-styryl-1,2,3,4-tetrahydroisoquinolin-
20 7-yl)-4-methoxy-3-trifluoromethylbenzamide (0.068g) according to the method of Example 3. The free base was converted into the hydrochloride salt by treatment of a methanol solution with excess ethereal HCl.

¹H NMR (CDCl₃) δ: 2.46 (3H, s), 2.66 (2H, t), 2.95 (2H, t), 3.52 (2H, s), 3.93 (3H, s), 6.79
25 (1H, d), 6.95 (1H, d), 7.00 (1H, s), 7.14 (1H, d), 7.35 – 7.53 (5H, m), 7.93 – 7.98 (2H, m), 8.20 (1H, d), 8.25 (1H, s); m/z (API⁺): 467(MH⁺)

The following Examples were prepared using the methods described for the Descriptions, Preparations and Examples above.

30

Example 20

N-(8-Bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-acetyl-4-ethoxybenzamide

35

¹H NMR (250MHz, CDCl₃) δ: 1.54 (3H, t, J = 7 Hz), 2.68 (3H, s), 2.98 (2H, m), 3.87 (2H, m, restricted amide rotation), 4.25 (2H, q, J = 7 Hz), 4.80 (2H, br) 7.05 (1H, d, J = 8 Hz), 7.19 (1H, m), 8.04 (1H, d), 8.10 (1H, dd, J = 8, 2 Hz), 8.22 - 8.36 (2H, m) and 8.32 (1H, m); m/z (API): 537, 535 (MNa⁺; 100%), 515, 513 (MH⁺; 10%).

Example 21**5 N-(8-Bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-acetyl-4-ethoxybenzamide**

¹H NMR (250MHz, d₆DMSO) δ: 1.46 (3H, t, J = 7 Hz), 2.60 (3H, s), 2.73 (2H, t), 2.93 (2H, t), 3.82 (2H, s), 4.26 (2H, q, J = 7 Hz), 7.12 (1H, d, J = 8 Hz), 7.28 (2H, m), 8.15 (1H, dd, J = 8, 2 Hz), 8.26 (2H, s), 9.95 (1H, s); m/z (API): 419, 417 (MH⁺; 100%)

10

Example 22**N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-fluoro-4-methoxybenzamide**

15 ¹H NMR (250MHz, CDCl₃) δ: 2.79 (2H, t), 3.11 (2H, t), 3.97 (3H, s), 4.05 (2H, s), 7.09 (2H, m), 7.68 (2H, m), 8.28 (2H, m); m/z (API): 335.1 (MH⁺; 100%)

Example 23**20 N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-chloro-4-ethoxybenzamide**

¹H NMR (250MHz, CDCl₃) δ: 1.51 (3H, t, J = 7 Hz), 2.78 (2H, t), 3.10 (2H, t), 4.03 (2H, s), 4.18 (2H, q, J = 7 Hz), 6.98 (1H, d, J = 8 Hz), 7.06 (1H, d), 7.77 (1H, dd, J = 8, 2 Hz), 7.94 (1H, d), 8.23 (1H, d), 8.31 (1H, s); m/z (API): 365.1 (MH⁺; 100%)

25

Example 24**N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-methoxybenzamide**

30 ¹H NMR (250MHz, d₆DMSO) δ: 2.70 (2H, t), 2.91 (2H, t), 3.83 (2H, s), 3.94 (3H, s), 7.10 (1H, d, J = 8 Hz), 7.22 - 7.32 (2H, m), 8.02 (1H, dd, J = 8, 2 Hz), 8.22 (1H, d, J = 2 Hz), 9.98 (1H, s); m/z (API): 397, 395 (MH⁺; 100%)

Example 25

35

N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-acetyl-4-ethoxybenzamide

¹H NMR (250MHz, CDCl₃) δ: 1.54 (3H, t, J = 7 Hz), 2.68 (3H, s), 2.80 (2H, t), 3.12 (2H, t), 4.05 (2H, s), 4.24 (2H, q, J = 7 Hz), 7.07 (2H, m), 8.11 (1H, dd, J = 8, 2 Hz), 8.25 (2H, m), 8.37 (1H, s); m/z (API): 373.2 (MH⁺; 100%)

5 Example 26

N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-acetyl-4-*iso*-propoxybenzamide

¹H NMR (250MHz, CDCl₃) δ: 1.46 (6H, d, J = 7 Hz), 2.66 (3H, s), 2.78 (2H, t), 3.10 (2H, t), 4.04 (2H, s), 4.80 (1H, sep, J = 7 Hz), 7.07 (2H, d, J = 8 Hz), 8.10 (1H, dd, J = 8, 2 Hz), 8.21 (1H, d, J = 8 Hz), 8.24 (1H, d, J = 2 Hz), 8.39 (1H, s); m/z (API): 409 (MNa⁺; 100%), 387 (MH⁺; 18%)

15

PHARMACOLOGICAL DATA

1. Binding Assay Method

- 20 WO 92/22293 (SmithKline Beecham) discloses compounds having anti-convulsant activity, including *inter alia* the compound *trans*-(+)-6-acetyl-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (hereinafter referred to as Compound A). It has been found that the compounds of WO 92/22293 bind to a novel receptor obtainable from rat forebrain tissue, as described in WO 96/18650 (SmithKline Beecham).
- 25 The affinity of test compounds to the novel receptor site is assessed as follows.

Method

- Whole forebrain tissue is obtained from rats. The tissue is first homogenised in buffer (usually 50mM Tris/HCl, pH 7.4). The homogenised tissue is washed by centrifugation and resuspension in the same buffer, then stored at -70°C until used.

- To carry out the radioligand binding assay, aliquots of tissue prepared as above (usually at a concentration of 1-2mg protein/ml) are mixed with aliquots of [3H]-Compound A dissolved in buffer. The final concentration of [3H]-Compound A in the mixture is usually 20nM. The mixture is incubated at room temperature for 1 hour. [3H]-Compound A bound to the tissue is then separated from unbound [3H]-Compound A by filtration through Whatman GF/B glass fibre filters. The filters are then washed rapidly with ice-cold buffer. The amount of radioactivity bound to the tissue trapped on the filters is measured by

addition of liquid scintillation cocktail to the filters followed by counting in a liquid scintillation counter.

5 In order to determine the amount of "specific" binding of [3H]-Compound A, parallel assays are carried out as above in which [3H]-Compound A and tissue are incubated together in the presence of unlabelled Compound A (usually 3 μ M). The amount of binding of [3H]-Compound A remaining in the presence of this unlabelled compound is defined as "non-specific" binding. This amount is subtracted from the total amount of [3H]-Compound A binding (i.e. that present in the absence of unlabelled compound) to
10 obtain the amount of "specific" binding of [3H]-Compound A to the novel site.

The affinity of the binding of test compounds to the novel site can be estimated by incubating together [3H]-Compound A and tissue in the presence of a range of concentrations of the compound to be tested. The decrease in the level of specific [3H]-
15 Compound A binding as a result of competition by increasing concentrations of the compound under test is plotted graphically, and non-linear regression analysis of the resultant curve is used to provide an estimate of compound affinity in terms of pKi value.

Results

20 Compounds of this invention were active in this test with pKi's greater than 6. For example, compounds of Examples 2, 3, 5, 6, 8-12, 14, 15, 18, 19 gave pKi values greater than 7.

25 2. MEST Test

The maximal electroshock seizure (MEST) threshold test in rodents is particularly sensitive for detecting potential anticonvulsant properties¹. In this model, anticonvulsant agents elevate the threshold to electrically-induced seizures whilst proconvulsants lower the
30 seizure threshold.

Method for mouse model

35 Mice (naive male, Charles River, U.K. CD-1 strain, 25 - 30g) are randomly assigned to groups of 10 - 20 and dosed orally or intraperitoneally at a dose volume of 10 ml/kg with various doses of compound (0.3 - 300 mg/kg) or vehicle. Mice are then subjected at 30 or 60 min post dose to a single electroshock (0.1 sec, 50Hz, sine wave form) administered via corneal electrodes. The mean current and standard error required to induce a tonic seizure in 50% (CC₅₀) of the mice in a particular treatment group is determined by the 'up and

down' method of Dixon and Mood (1948)². Statistical comparisons between vehicle- and drug-treated groups are made using the method of Litchfield and Wilcoxon (1949)³.

5 In control animals the CC₅₀ is usually 14 - 18 mA. Hence the first animal in the control group is subjected to a current of 16 mA. If a tonic seizure does not ensue, the current is increased for a subsequent mouse. If a tonic convulsion does occur, then the current is decreased, and so on until all the animals in the group have been tested.

10 Studies are carried out using a Hugo Sachs Elektronik Constant Current Shock Generator with totally variable control of shock level from 0 to 300 mA and steps of 2 mA are usually used.

Results

15 Compounds of this invention dosed at 10 mg/kg by the oral route as a suspension in methyl cellulose and tested one hour post dosing showed an increase in seizure threshold.

Method for rat model

20 The threshold for maximal (tonic hindlimb extension) electroshock seizures in male rats (Sprague Dawley, 80 - 150g, 6 weeks old) was determined by a Hugo Sachs Elektronik stimulator which delivered a constant current (0.3 sec duration; from 1-300mA in steps of 5-20mA). The procedure is similar to that outlined above for mouse and full details are as published by Upton et al.,⁴

25 The percentage increase or decrease in CC₅₀ for each group compared to the control is calculated.

Drugs are suspended in 1% methyl cellulose.

30

Results

At a dosage of 2 mg/kg p.o. at 2h, the compounds of Examples 2, 3, 5, 6, 8 and 9 show statistically significant increases of 120%, 160%, 320%, 260%, 400% and 340% respectively.

35

References


1. Loscher, W. and Schmidt, D. (1988). *Epilepsy Res.*, **2**, 145-181
2. Dixon, W.J. and Mood, A.M. (1948). *J. Amer. Stat. Assn.*, **43**, 109-126

3. Litchfield, J.T. and Wilcoxon, F.(1949). J. Pharmacol. exp. Ther., 96, 99-113
4. N.Upton, T.P.Blackburn, C.A.Campbell, D.Cooper, M.L.Evans, H.J.Herdon, P.D.King, A.M.Ray, T.O.Stean, W.N.Chan, J.M.Evans and M.Thompson. (1997). B. J. Pharmacol., 121, 1679-1686

CLAIMS

1. A compound of formula (I) or pharmaceutically acceptable salt or solvate thereof:


5




- where Q is a monocyclic or bicyclic aryl or heteroaryl ring,
- 10 R^1 is hydrogen, C_{1-6} alkyl (optionally substituted by hydroxy or C_{1-4} alkoxy), C_{1-6} alkenyl, C_{1-6} alkynyl, C_{1-6} alkylCO-, formyl, CF_3 CO- or C_{1-6} alkylSO₂-,
- R^2 is hydrogen or up to three substituents selected from halogen, NO₂, CN, N₃, CF_3 O-, CF_3 S-, CF_3 CO-, trifluoromethyldiaziriny, C_{1-6} alkyl, C_{1-6} alkenyl,
- 15 C_{1-6} alkynyl, C_{1-6} perfluoroalkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl- C_{1-4} alkyl-, C_{1-6} alkylO-, C_{1-6} alkylCO-, C_{3-6} cycloalkylO-, C_{3-6} cycloalkylCO-, C_{3-6} cycloalkyl- C_{1-4} alkylO-, C_{3-6} cycloalkyl- C_{1-4} alkylCO-, phenyl, phenoxy, benzyloxy, benzoyl, phenyl- C_{1-4} alkyl-, C_{1-6} alkylS-, C_{1-6} alkylSO₂-, $(C_{1-4}$ alkyl)₂NSO₂-,
- 20 $(C_{1-4}$ alkyl)NHCO, CONH₂, CF_3 SO₂, C_{1-6} alkenyl, C_{1-6} alkynyl or C_{1-6} hydroxyalkyl; or -NR³R⁴ where R³ is hydrogen or C_{1-4} alkyl, and R⁴ is hydrogen, C_{1-4} alkyl, formyl, -CO₂ C_{1-4} alkyl or -COC C_{1-4} alkyl; or two R² groups together form a carbocyclic ring that is saturated or
- 25 unsaturated and unsubstituted or substituted by -OH or =O; and
- X is halogen, C_{1-6} alkoxy, C_{1-6} alkyl, C_{2-6} alkenyl optionally substituted by phenyl.

2. A compound according to claim 1 of formula (IA)

30



3. A compound according to claim 1 of formula (IB)




(IB)

- 5 4. A compounds of claim 1, 2 or 3 in which
 R^1 is hydrogen, methyl, ethyl, propyl, hydroxyethyl, formyl, acetyl, trifluoroacetyl or methanesulfonyl,
 R^2 is hydrogen or one or more of methyl, ethyl, *n*-butyl, *iso*-propyl, *t*-butyl, phenyl, methoxy, ethoxy, *iso*-propoxy, *n*-butoxy, cyclopropylmethoxy,
 10 phenoxy, benzyloxy, amino, acetylamino, nitro, azido, cyano, bromo, chloro, fluoro, iodo, acetyl, propionyl, pivaloyl, *iso*-butyryl, benzoyl, iodobenzoyl, trifluoromethyl, perfluoroethyl, trifluoromethoxy, trifluoroacetyl, trifluoromethyldiaziriny, methanesulfonyl, *n*-propylsulfonyl, isopropylsulfonyl, dimethylsulfamoyl, CF_3SO_2 ;
 15 or two groups R^2 form a benzene, cyclopentane or cyclopentanone ring;
 X is methyl, ethyl, chloro, bromo, iodo, fluoro, phenylethenyl.
5. A compounds of claim 1, 2 or 3 in which
 R^1 is hydrogen, methyl, formyl, hydroxyethyl or trifluoroacetyl,
 20 R^2 is hydrogen or one or more of methyl, ethyl, *iso*-propyl, methoxy, ethoxy, *iso*-propoxy, acetyl, propionyl, pivaloyl, cyano, bromo, chloro, fluoro, iodo, trifluoromethyl,.
 X is methyl, ethyl, chloro or bromo.
- 25 6. A compound of formula (I) selected from:
 N-(8-chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-ethoxybenzamide
 N-(8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-ethoxybenzamide
 30 N-(8-chloro-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-ethoxybenzamide
 N-(8-chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
 N-(8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide

- N-(8-chloro-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- N-(8-bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- 5 N-(8-bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- N-(8-bromo-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- N-(2,8-Dimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- 10 N-(2-(2-Hydroxyethyl)-8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- N-(2-formyl-8-chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- N-(8-ethyl-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- 15 N-(8-ethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- N-(8-ethyl-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- N-(8-chloro-2-methanesulfonyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- 20 N-(8-Styryl-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- N-(8-Styryl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- N-(8-Styryl-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxy-3-trifluoromethylbenzamide
- 25 N-(8-Bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-acetyl-4-ethoxybenzamide
- N-(8-Bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-acetyl-4-ethoxybenzamide
- N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-fluoro-4-methoxybenzamide
- 30 N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-chloro-4-ethoxybenzamide
- N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-bromo-4-methoxybenzamide
- N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-acetyl-4-ethoxybenzamide
- N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-acetyl-4-*iso*-propoxybenzamide..
7. A process for the preparation of compounds of formula (I) according to any one of
- 35 claims 1 to 6, which comprises reacting a compound of formula (II)





where R^{1A} is R^1 as defined for formula (I) or a group convertible to R^1

- 5 and X^A is X as defined in claim 1 or a group convertible to X
with a compound of formula (III)



- 10 where Q is as defined in formula (I), Y is Cl or OH, and R^{2A} groups are independently R^2
as defined for formula (I) or groups convertible to R^2 ,
and where required converting an X^A , R^{1A} or R^{2A} group to an X, R^1 or R^2 group,
converting one X, R^1 or R^2 group to another X, R^1 or R^2 group,
converting a salt product to the free base or another pharmaceutically acceptable salt, or
15 converting a free base product to a pharmaceutically acceptable salt.

8. A pharmaceutical composition, in particular for use in the treatment and/or
prophylaxis of anxiety, mania, depression, panic disorders and/or aggression, disorders
associated with a subarachnoid haemorrhage or neural shock, the effects associated with
20 withdrawal from substances of abuse such as cocaine, nicotine, alcohol and
benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such
as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine,
cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as
Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological
25 deficits associated with AIDS, sleep disorders (including circadian rhythm disorders,
insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury,
tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer
pain, inappropriate neuronal activity resulting in neurodysthesias in diseases such as
diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity
30 (spasticity), temporomandibular joint dysfunction, or amyotrophic lateral sclerosis (ALS)
which comprises a compound of formula (I) according to any one of claims 1 to 6, or a



pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier.

9. A method of treatment and/or prophylaxis of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesis in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, or amyotrophic lateral sclerosis (ALS) comprising administering to the sufferer in need thereof an effective or prophylactic amount of a compound of formula (I) according to any one of claims 1 to 6, or a pharmaceutically acceptable salt or solvate thereof.

10. Use of a compound of formula (I) according to any one of claims 1 to 6, or a pharmaceutically acceptable salt or solvate thereof, for the manufacture of a medicament for the treatment and/or prophylaxis of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesis in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, or amyotrophic lateral sclerosis (ALS).

11. Use of a compound of formula (I) according to any one of claims 1 to 6, or a pharmaceutically acceptable salt or solvate, thereof as a therapeutic agent, in particular for the treatment and/or prophylaxis of anxiety, mania, depression, panic disorders and/or



- aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis,
- 5 migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer
- 10 pain, inappropriate neuronal activity resulting in neurodysthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, or amyotrophic lateral sclerosis (ALS).