CLAIMS

What is claimed is:

1. A liquid formulation comprising a polypeptide and a surfactant, wherein at least about 70% (wt%) of the surfactant are isosorbide polyoxyethylene (POE) fatty acid esters.
2. The liquid formulation of claim 1, wherein the isosorbide POE fatty acid esters comprise about 5-30 POE units, or


wherein the isosorbide POE fatty acid esters comprise about 20 POE units, or

wherein the isosorbide POE fatty acid esters comprise fatty acid chains selected from the group consisting of an optionally substituted C_{4-28} alkyl and an optionally substituted C_{4-28} alkenyl.

3. The liquid formulation of claim 1 or 2, wherein the isosorbide POE fatty acid esters are monoesters, or

wherein the isosorbide POE fatty acid esters are monoesters selected from the group consisting of isosorbide POE monolaurate, isorsobide POE monomyristate, isosorbide POE monopalmitate, isosorbide POE monostearate and isosorbide POE monooelate, or wherein the isosorbide POE fatty acid ester is isosorbide POE monolaurate.

4. The liquid formulation of claim 1 or 2, wherein the isosorbide POE fatty acid esters are monoesters, diesters, or a mixture of the foregoing.
5. The liquid formulation of claim 1, wherein the isosorbide POE fatty acid ester is a compound of Formula (I):



(I);

wherein:

a and b are independently integers from 2 to 28, provided that the sum of a and b is an integer from 5-30;

R^1 and R^2 are independently selected from the group consisting of hydrogen and

-C(O)R", wherein R" is an optionally substituted C_{3-27} alkyl or an optionally substituted C_{3-27} alkenyl; and

R^3 and R^4 are independently hydrogen.

6. The liquid formulation of claim 5, wherein the sum of a and b is 9,

or wherein R^1 is H and R^2 is -C(O)R", or

R^1 is -C(O)R" and R^2 is H, or

both R^1 and R^2 are -C(O)R".

7. The liquid formulation of claim 5 or 6, wherein R" is an unsubstituted C_{3-27} alkyl,
or wherein R" is an unsubstituted C_{11} alkyl.

8. The liquid formulation of claim 5 or 6, wherein R" is an unsubstituted C_{3-27} alkenyl,
or wherein R" is an unsubstituted C_{17} alkenyl.

9. The liquid formulation of any one of claims 1-8, wherein at least about 80% (wt%) of the surfactant are isosorbide POE fatty acid esters.

10. The liquid formulation of any one of claims 1-9, wherein the surfactant further comprises a POE fatty acid ester,

or wherein the surfactant comprises a greater amount of isosorbide POE fatty acid esters than POE fatty acid esters, or

wherein the POE fatty acid ester comprises a fatty acid chains selected from the group consisting of an optionally substituted C_{4-28} alkyl and an optionally substituted C_{4-28} alkenyl, or

wherein the POE fatty acid ester is selected from a group consisting of POE monolaurate, POE monomyristate, POE monopalmitate, POE monostearate, and POE monooelate, or

wherein less than about 20% (wt%) of the surfactant are POE fatty acid esters, or

wherein the surfactant is about 0.0005% to 0.2% (w:v) of the liquid formulation, or

wherein the surfactant further comprises a sorbitan POE fatty acid ester.

11. The liquid formulation of any one of claims 1-10, wherein the polypeptide is a protein, or

wherein the polypeptide is a protein that is an antibody, or

wherein the polypeptide is a protein that is an antibody selected from the group consisting of a polyclonal antibody, a monoclonal antibody, a humanized antibody, a human antibody, a chimeric antibody, and an antibody fragment,

or wherein the polypeptide is a protein that is an antibody fragment selected from the group consisting of Fab, Fab', $F(ab')_2$, and Fv fragments.

12. The liquid formulation of claim 11, wherein the antibody concentration is about 0.1 mg/mL to about 300 mg/mL, or

about 100 mg/mL to about 300 mg/mL.

13. The liquid formulation of any one of claims 1-12, wherein the liquid formulation is a reconstituted lyophilized formulation.

14. The liquid formulation of claim 12, wherein the liquid formulation is further diluted with an infusion solution to a polypeptide concentration of about 0.001 mg/mL to about 0.5 mg/mL.

15. The liquid formulation of any one of claims 1-14, wherein the liquid formulation is substantially free of aggregates, or

comprises less free fatty acid particle formation.

16. The liquid formulation of any one of claims 10-15, wherein at least about 80% (wt%) of the surfactant are isosorbide polyoxyethylene (POE) fatty acid esters and POE fatty acid esters, or

wherein at least about 90% (wt%) of the surfactant are isosorbide polyoxyethylene (POE) fatty acid esters and POE fatty acid esters.

17. A lyophilized formulation comprising a polypeptide and a surfactant, wherein at least about 70% (wt%) of the surfactant are isosorbide polyoxyethylene (POE) fatty acid esters and POE fatty acid esters; and wherein the lyophilized formulation is prepared by lyophilizing the liquid formulation according to any one of claims 10-16.

18. An article of manufacture comprising a container enclosing the liquid formulation of any one of claims 1-16.

19. The article of manufacture of claim 18, wherein the container is an IV bag, or

wherein the container is an IV bag that comprises an injection device, or

wherein the container is an IV bag that comprises an infusion solution.

20. The article of manufacture of claim 18, wherein the container is a vial or a pre-filled syringe.

21. An article of manufacture comprising a container enclosing the lyophilized formulation of claim 17.

22. The article of manufacture of claim 18 or 21, further comprising a second container enclosing a diluent.

23. A method of making a liquid formulation comprising adding a polypeptide and a surfactant to an aqueous solution, wherein at least 70% (wt%) of the surfactant are isosorbide POE fatty acid esters.

24. The method of claim 23, wherein the isosorbide POE fatty acid esters comprises about 5-30 POE units, or


wherein the isosorbide POE fatty acid esters comprises about 20 POE units.

25. The method of claim 23 or 24, wherein the isosorbide POE fatty acid esters comprise fatty acid chains selected from the group consisting of an optionally substituted C_{4-28} alkyl and an optionally substituted C_{4-28} alkenyl, or wherein the isosorbide POE fatty acid esters are monoesters, diesters, or a mixture of the foregoing.

26. The method of claim 23 or 24, wherein the isosorbide POE fatty acid esters are monoesters, or

wherein the isosorbide POE fatty acid esters are monoesters selected from the group consisting of isosorbide POE monolaurate, isorsobide POE monomyristate, isosorbide POE monopalmitate, isosorbide POE monostearate, and isosorbide POE monooelate, or wherein the isosorbide POE fatty acid ester is isosorbide POE monolaurate.

27. The method of claim 23, wherein the isosorbide POE fatty acid ester is a compound of Formula (I):



(I);

wherein:

a and b are independently integers from 2 to 28, provided that the sum of a and b is an integer from 5-30;

R^1 and R^2 are independently selected from the group consisting of hydrogen and

-C(O)R", wherein R" is an optionally substituted C_{3-27} alkyl or an optionally substituted C_{3-27} alkenyl; and

R^3 and R^4 are independently hydrogen.

28. The method of claim 27, wherein the sum of a and b is 9, or

wherein R^1 is H and R^2 is -C(O)R", or

R^2 is -C(O)R" and R^1 is H, or

both R^1 and R^2 are -C(O)R".

29. The method of claim 27 or 28, wherein R" is an unsubstituted C_{3-27} alkyl, or

wherein R" is an unsubstituted C_{11} alkyl.

30. The method of claim 27 or 28, wherein R" is an unsubstituted C_{3-27} alkenyl, or

wherein R" is an unsubstituted C_{17} alkenyl.

31. The method of any one of claims 23-30, wherein the surfactant further comprises a POE fatty acid ester, or

wherein at least about 80% (wt%) of the surfactant are isosorbide POE fatty acid esters and POE fatty acid esters, or

wherein at least about 90% (wt%) of the surfactant are isosorbide POE fatty acid esters and POE fatty acid esters, or

wherein the surfactant comprises a greater amount of isosorbide POE fatty acid esters than POE fatty acid esters, or

wherein the surfactant further comprises a POE fatty acid ester that comprises a fatty acid chain selected from the group consisting of an optionally substituted C_{4-28} alkyl and an optionally substituted C_{4-28} alkenyl, or

wherein the surfactant further comprises a POE fatty acid ester selected from the group consisting of POE monolaurate, POE monomyristate, POE monopalmitate, POE monostearate, and POE monooelate, or

wherein less than about 20% (wt%) of the surfactant are POE fatty acid esters, or

wherein the surfactant further comprises a sorbitan POE fatty acid ester, or

wherein the surfactant is about 0.0005% to 0.2% (w:v) of the liquid formulation.

32. The method of any one of claims 23-31, wherein the polypeptide is a protein, or

wherein the polypeptide is a protein that is an antibody, or

wherein the polypeptide is a protein that is an antibody selected from the group consisting of a polyclonal antibody, a monoclonal antibody, a humanized antibody, a human antibody, a chimeric antibody, and an antibody fragment, or

wherein the polypeptide is a protein that is an antibody fragment selected from the group consisting of Fab, Fab', $F(ab')_2$, and Fv fragments, or

wherein the polypeptide is a protein that is an antibody at a concentration of about 0.1 mg/mL to about 300 mg/mL, or

about 100 mg/mL to about 300 mg/mL, or

wherein the liquid formulation is further diluted with an infusion solution to a concentration of about 0.1 mg/mL to about 2 mg/mL.

33. The method of any one of claims 23-32, further comprising lyophilizing the liquid formulation to make a lyophilized formulation, or

wherein the liquid formulation is substantially free of aggregates, or comprises less free fatty acid particle formation.