

Precision Medicine for Schizophrenia and Psychotic Disorders: Objective Assessment, Risk Prediction, Pharmacogenomics, and Repurposed Drugs

STATEMENT OF GOVERNMENT SUPPORT

[0001] This invention was made with government support under OD007363 awarded by the National Institutes of Health and CX000139 merit award by the Veterans Administration. The government has certain rights in the invention.

CROSS-REFERENCE TO RELATED APPLICATION

[0002] This application claims priority to U.S. Patent Application No. 17/351,132, filed on June 17, 2021, the disclosure of which is hereby expressly incorporated by reference in its entirety.

BACKGROUND

[0003] Schizophrenia is a heterogeneous disorder, composed of positive and negative psychotic symptoms. Psychotic symptoms, more broadly, are also often present in other psychiatric disorders. They can be difficult to assess, as they are based on the patient's self-reporting and on the clinician's clinical impression. Continued improvements are needed to adequately diagnose and treat individuals suffering psychotic symptoms.

SUMMARY

[0004] Provided here are newly identified blood gene expression biomarkers for hallucinations, and for delusions. The biomarkers provide a means of assessing state severity, short-term risk, and long-term risk. The biomarkers can also be used for drug repurposing.

[0005] Some aspects of the invention include methods for treating an individual experiencing or at a heightened risk for developing symptoms such as delusions and/or hallucinations. These symptoms may be indicative of certain psychiatric disorders such as psychosis and/or schizophrenia. Treating involves administered at least one course of treatment, treatment may include psychiatric counseling, administering certain physical intervention, and/or prescribing and/or administering at least one therapeutic compound. Treating may include at least one of the following outcomes, curing, mitigating, managing or otherwise recuing the severity and/or the number of frequency delusions and or hallucinations. In some aspects of the invention treating

may include identifying individuals with or at an increased risk for developing delusions and/or hallucinations by measuring the level of certain RNA biomarkers as identified in, for example, Tables 2 and 3.

[0006] Some aspects of the invention include methods for diagnosing an individual experiencing or at a heightened risk for developing symptoms such as delusions and/or hallucinations. These symptoms may be indicative of certain psychiatric disorders such as psychosis and/or schizophrenia. Diagnosing does not require treatment, although it may lead to, or become, part of treating an individual who is manifesting or at an increased risk for manifesting symptom of certain types of mental illness such as psychosis and or schizophrenia. In some aspects of the invention diagnosing may include identifying individuals with or at an increased risk for developing delusions and/or hallucinations by measuring the level of certain RNA biomarkers as identified in, for example, Tables 2 and 3.

[0007] A first embodiment of the invention is a method for treating at least one psychiatric disorder, for example delusions or an increased risk for developing delusions in an individual, comprising the steps of: (a) obtaining a biological sample from an individual and quantifying the amounts of RNA biomarkers in the biological sample, to create a panel of RNA biomarkers, (b) quantifying the amounts of the RNA biomarkers in the panel in a clinically relevant population to generate a reference expression level for the RNA biomarkers in a panel of RNA biomarkers; (c) comparing the amounts of the biomarkers in the biological sample from the individual with the amounts of the RNA biomarkers present in the reference standard to generate a score for each biomarker; wherein the biomarkers in the first panel (a) comprise one or more of the following RNA biomarkers: Activator Of Transcription and Developmental Regulator 2 (AUTS2), Pyruvate Dehydrogenase Phosphatase Catalytic Subunit 1 (PDP1), Nuclear Receptor Subfamily 4 Group A Member 2 (NR4A2), GNAS Complex Locus (GNAS), Interleukin 6 Signal Transducer (IL6ST), Chromodomain Helicase DNA Binding Protein 9 (CHD9), X-Ray Repair Cross Complementing 6 (XRCC6), RAR Related Orphan Receptor A (RORA), Actinin Alpha 4 (ACTN4), and Acyl-CoA Synthetase Long Chain Family Member 4 (ACSL4), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting delusions or an increased risk for developing delusions; and biomarkers in a second panel (b) comprise one or more of the following RNA biomarkers: Zinc Finger And BTB Domain Containing 20 (ZBTB20), Forkhead Box P1 (FOXP1), Spondin 1 (SPON1), and

(NRP2), wherein the expression level of the RNA biomarker(s) in the sample is decreased relative to a reference expression level of the RNA biomarkers in the panel, denoting delusions or an increased risk for developing delusions; (d) generating a score for the panel of RNA biomarkers, based on the scores of the biomarker(s) in the panel; (e) determining a reference score for the panel in a clinically normal relevant population; (f) identifying a difference between the score of the panel of biomarker(s) in the sample and the reference score of the panel of biomarker(s); (g) identifying the individual as having delusions or of having an elevated risk for developing delusions, based on the difference between the biomarker panel score of the individual relative to the biomarker panel score of the reference; (h) treating the individual identified as having delusions or an elevated risk of delusions with at least one treatment selected from the group consisting of: a treatment based on clinical practice guidelines, administering a therapeutically effective amount of at least one therapeutic drug wherein the mode of treatment is on the specific biomarkers scores indicating that individual will benefit from a particular therapy, treating includes curing, mitigating, reducing or even eliminating symptoms of psychotic disorders such as schizophrenia. In some embodiments of the invention samples are taken from an individual two or more time in order to treat, diagnose, and/or monitor the presence of at least one psychotic disorder such as schizophrenia. In some embodiments, an individual may be treated for symptoms such as delusions or hallucinations without a formal diagnosis of a specific psychotic disorder. In some embodiments, the individual may be treated with drugs known to treat mental illness and/or drugs repurposed to treat mental illness. Samples from an individual may include any or all of the following, tissue samples, bodily fluids such as blood serum, plasma, saliva, cerebral fluid and the like. The samples may be further processed such as by extraction or purification before being analyzed for the presence of one or more biomarkers of interest.

[0008] In a third embodiment an individual exhibiting symptoms of a psychotic disorder such as those noted the first and second embodiments may be treated with at least drug selected from the group consisting of: adenosine phosphate, N-acetyl-L-leucine, eldeline, pempidine, verteporfin, C-75, oxprenolol, Prestwick-675, meglumine, guanethidine, pancuronium bromide, karakoline, 15(S)-15-methylprostaglandin E2, hexylcaine, dicoumarol, apramycin, mephentoin, estriol, 528116.cdx, Cyclopiazonic Acid, SB 218078, BRD-A36630025, Quinacrine hydrochloride, GF-109203X, BRD-A36630025, N9-isopropylolomoucine, BMS-536924, BRD-

K76951091, BRD-K26304855, trichostatin A, ALW-II-38-3, mitoxantrone, HG-6-64-0, alvocidib, SB-216763, and Caffeic acid phenethyl ester. In some embodiments the drug used to treat the individual may be a drug repurposed from another use, see for example the drugs in Table 4, some repurposed drugs may be identified as efficacious for this purpose because their use correlates in a beneficial change in at least one of the Biomarkers listed in Tables 1 and/or 2.

[0009] A fourth embodiment includes at least portion of the first through the third embodiments, wherein the individual is male, and the biomarkers in a first panel (a) comprise one or more biomarkers selected from the group consisting of: Activator Of Transcription And Developmental Regulator (AUTS2), Pyruvate Dehydrogenase Phosphatase Catalytic Subunit 1 (PDP1), Forkhead Box P1 (FOXP1), GNAS Complex Locus (GNAS), Serine Racemase (SRR), Nuclear Receptor Subfamily 4 Group A Member 2 (NR4A2), X-Ray Repair Cross Complementing 6 (XRCC6), RAR Related Orphan Receptor A (RORA), and Actinin Alpha 4 (ACTN4), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting increased delusions, or the biomarkers in a second panel (b) comprising one or more biomarkers selected from the group consisting of: Zinc Finger And BTB Domain Containing 20 (ZBTB20), Forkhead Box P1 (FOXP1), Spondin 1 (SPON1), NRP2, wherein the expression level of the biomarker(s) in the sample is decreased relative to a reference expression level, denoting increased delusions. The males identified in this embodiment may be treated with at least one therapeutic drug is selected from the group consisting of: flunisolide, apramycin, adenosine phosphate, guanethidine, 15(S)-15-methylprostaglandin E2, meteneprost, methyldopate, hydralazine, rotenone, phthalylsulfathiazole, N-acetyl-L-leucine, eldeline, tocainide, laudanosine, pempidine, 7-aminocephalosporanic acid, Sulfachlorpyridazine, finasteride, 528116.cdx, SB 218078, Quinacrine hydrochloride, N9-isopropylolomoucine, ALW-II-38-3, mitoxantrone, HG-6-64-01, Alvocidib, SB-216763, Syk Inhibitor, Cyclopiazonic Acid, GW 441756, LY 225910, AG 82, doxorubicin, mitomycin, and terfenadine.

[0010] A fifth embodiment includes at least portion of the first through the third embodiments, wherein the individual is a female and the biomarker is at least one biomarker selected from the group consisting of: Phosphodiesterase 4D Interacting Protein (PDE4DIP), Pyruvate Dehydrogenase Phosphatase Catalytic Subunit 1 (PDP1), Transcription Factor 4 (TCF4), Nuclear Receptor Subfamily 4 Group A Member 2 (NR4A2), Chromodomain Helicase DNA

Binding Protein 9 (CHD9), (CLCN3), Activator Of Transcription And Developmental Regulator (AUTS2), and (LDB2), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting increased delusions; and the biomarkers in a second panel (b) comprise one or more biomarkers selected from the group consisting of (FGFR1), (DISC1), (FGFR2), (SPTBN1), (INSR), (GRIK3), Zinc Finger, and BTB Domain Containing 20 (ZBTB20), wherein the expression level of the biomarker(s) in the sample is decreased relative to a reference expression level, denoting increased delusions. The females identified in this embodiments may be treated with at least one therapeutic drug is selected from the group consisting of: erastin, harpagoside, metacycline, amiodarone, furaladone, metformin, timolol, Repaglinide, sulfafurazole, PNU-0230031, Probenecid, furosemide, fluphenazine, myricetin, sulfacetamide, lomustine, BCB000039, Harmalol, I-BET151, Nylidrin hydrochloride, AMG 9810, Doxorubicin, Mitomycin C, Fludrocortisone acetate, Purvalanol A, Teniposide, Geldanamycin, Importazole, BRD-A36630025, YM-155, Auranofin, 7643453, G-221, BRD-A49680073, BRD-K08547377, and Cladribine.

[0011] A sixth embodiment includes a method of assessing and/or treating schizophrenia and other psychotic disorders in general, and delusions in particular, in an individual, comprising: calculating combined biomarkers and clinical information Up- based on the equation: (Biomarker Panel Score) + (Delusions Score) - (Grooming Score) = Up-Delusions Score; wherein the Biomarker Panel Score is obtained as per the method of claim 1; wherein the Delusions Score is calculated with a clinical rating or self-report scales; wherein the Grooming Score is calculated with a rating scale; assessing the level of delusions of the individual by comparing the individual's Up-Delusions Score to a reference Up-Delusions Score; administering a treatment for delusions to the individual when the individual's Up-Delusions Score is greater than a reference Up-Suicide Score; and monitoring the individual's response to a treatment for delusions by determining changes in the Up-Delusions Score after initiating a treatment.

[0012] A seventh embodiment is a method for assessing and/or treating schizophrenia and other psychotic disorders in general, in particular hallucinations and risk of developing hallucinations in an individual, comprising the steps of: (a) obtaining a biological sample from an individual and quantifying the amounts of one or more RNA biomarkers in the biological sample, to create at least one panel of RNA biomarkers, (b) quantifying the amounts of the RNA biomarkers in the at least one panel in a clinically relevant population to generate a reference

expression level for the RNA biomarkers in a panel of RNA biomarkers; (c) comparing the amounts of the biomarkers in the biological sample from the individual with the amounts of the RNA biomarkers present in the reference standard to generate a score for each biomarker a first panel and a second panel; wherein the biomarkers in the first panel comprise one or more of the following RNA biomarkers: (PRICKLE1), (NCAM1), (B3GALT5), (ARHGAP18), (PTP4A2), Acylphosphatase 2 (ACYP2), Reticulon 4 (RTN4), Cullin 4A (CUL4A), Zinc Finger E-Box Binding Homeobox 2 (ZEB2), Dystonin (DST), and Discs Large MAGUK Scaffold Protein 1 (DLG1), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting hallucinations or an increased risk for developing hallucinations; and wherein the biomarkers in the second panel comprise one or more of the following RNA biomarkers: (PRL), (SERPING1), Ectonucleotide Pyrophosphatase/Phosphodiesterase 2 (ENPP2), (LAMA4), (KCNV1), Catenin Delta 1 (CTNND1), and FAT Atypical Cadherin 4 (FAT4), wherein the expression level of the RNA biomarker(s) in the sample is decreased relative to a reference expression level of the RNA biomarkers in the panel, denoting hallucinations or an increased risk for developing hallucinations; (d) generating a score for the panel of RNA biomarkers, based on the scores of the biomarker(s) in the panel; (e) determining a reference score for the panel in a clinically normal relevant population; (f) identifying a difference between the score of the panel of biomarker(s) in the sample and the reference score of the panel of biomarker(s); (g) identifying the individual as manifesting hallucinations or of having an elevated risk for developing hallucinations, based on the difference between the biomarker panel score of the individual relative to the biomarker panel score of the reference; (h) treating the individual identified as having hallucinations or an elevated risk of hallucinations with one or more of the following: 1) a treatment based on clinical practice guidelines, 2) administering a therapeutically effective amount of a therapeutic drug (s), selected based on the specific biomarkers whose scores indicate that they are changed in the individual compared to a reference standard.

[0013] An eighth embodiment is any embodiment from the first through the seventh embodiments wherein the biomarkers are quantified in samples taken on two or more occasions from the individual. As noted earlier, samples can be taken from tissue or any bodily fluid harboring RNA biomarkers.

[0014] A ninth embodiment is a method of the seventh embodiment wherein each biomarker

is assigned a weighted coefficient based on the biomarkers importance in assessing and predicting hallucinations risk; and the biomarker panel score is based on the weighted coefficients of each of the biomarkers.

[0015] A tenth embodiment is any embodiment from the seventh through the ninth embodiments, wherein the one or more therapeutic is one or more compounds selected from the group consisting of: clioquinol, pirinixic acid, moxislyte, Prestwick-685, exemestane, azacytidine, C-75, estradiol, tetraethylenepentamine, sparteine, guanethidine, idoxuridine, gliclazide, nitrendipine, N-acetyl-L-aspartic acid, sulfanilamide, doxazosin, pimozide, Proscillaridin, oxetacaine, BRD-K71489689, trichostatin A, A443654, AG 825, Proscillaridin A, Ala-Ala-Phe-CMK, Fluocinolone acetonide, manumycin A, curcumin, BRD-K68548958, CHR 2797, Tyrphostin AG 1478, Wortmannin, HY-50878, 598226, S1003, BRD-A52530684, CGP-60474, Buparlisib, and AS-601245.

[0016] An eleventh embodiment is the method according to the seventh embodiment wherein the individual is male, and the biomarkers in a first panel (a) comprise one or more biomarkers selected from the group consisting of: (SH3PXD2A), Zinc Finger E-Box Binding Homeobox 2 (ZEB2), (PRICKLE1), (ARHGAP18), Acylphosphatase 2 (ACYP2), Reticulon 4 (RTN4), and Dystonin (DST), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting increased hallucinations; and biomarkers in a second panel (b) comprise one or more biomarkers selected from the group consisting of: (PRL), (SERPING1), Ectonucleotide Pyrophosphatase/Phosphodiesterase 2 (ENPP2), (KCNV1), Mab-21 Like 1 (MAB21L1), Catenin Delta 1 (CTNND1), and FAT Atypical Cadherin 4 (FAT4), wherein the expression level of the biomarker(s) in the sample is decreased relative to a reference expression level, denoting increased hallucinations. When the individual is an male the at least one therapeutic compound selected from the group consisting of: digoxigenin, doxazosin, meptazinol, promethazine, cefixime, velnacrine, cetirizine, eldeline, atropine oxide, clioquinol, nicotinic acid, clioquinol, galantamine, rolitetracycline, betahistine, sulconazole, monocrotaline, lanatoside C, Prestwick-1084, Naftidrofuryl, sulfachlorpyridazine, helveticoside, bezafibrate, mifepristone, trichostatin A, manumycin A, NCGC00189555-02, Buparlisib, linifanib, AZD-7762, Dinaciclib, Piretanide, KN-62, Fluticasone propionate, JAK3 Inhibitor VI, Sarmentogenin, Digoxin, Megestrol acetate, Oxymetazoline hydrochloride, U-0126, Tracazolate hydrochloride, Flufenamic acid, Fenofibrate, and U 99194 maleate, may be particularly effective in treating the

individual, although any therapeutically effective drug may be used to treat the individual.

[0017] A twelfth embodiment is the method according to the sixth embodiment wherein the individual is female, and the biomarkers in a first panel (a) comprise one or more biomarkers selected from the group consisting of: (CELSR2), (KALRN), (B3GALT5), Protein Phosphatase 3 Catalytic Subunit Beta (PPP3CB), (ZFR), (THNSL1), (TNIK), Nuclear Receptor Subfamily 4 Group A Member 2 (NR4A2), Zinc Finger E-Box Binding Homeobox 2 (ZEB2), and (TNIK), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting increased hallucinations; and biomarkers in a second panel (b) comprise one or more biomarkers selected from the group consisting of GNAS Complex Locus (GNAS), and Catenin Delta 1 (CTNND1), wherein the expression level of the biomarker(s) in the sample is decreased relative to a reference expression level, denoting increased hallucinations. When the individual is a female at least one therapeutic compound selected from the group consisting of: proglumide, quinethazone, esculin, MG-262, GW-8510, haloperidol, guanethidine, deferoxamine, citiolone, meteneprost, amylocaine, CP-944629, Clemizole, IC-86621, Nortriptyline, CP-944629, Tanesprimycin, Prestwick-674, 0317956-0000, and Pioglitazone, may be particularly effective in treating the individual, although any therapeutically effective drug may be used to treat the individual.

[0018] Still another embodiment is a method of treating and or accessing schizophrenia and other psychotic disorders in general, and hallucinations in particular, in an individual, comprising: calculating combined biomarkers and clinical information Up- based on the equation: (Biomarker Panel Score) + (Hallucinations Score) - (Grooming Score) = Up-Hallucinations Score; wherein the Biomarker Panel Score is obtained as per the seventh embodiment; wherein the Hallucinations Score is calculated with a clinical rating or self-report scales; wherein the Grooming Score is calculated with a rating scale; assessing the level of hallucinations of the individual by comparing the individual's Up-Hallucinations Score to a reference Up-Hallucinations Score; administering a treatment for hallucinations to the individual when the individual's Up-Hallucinations Score is greater than a reference Up-Suicide Score; and monitoring the individual's response to a treatment for hallucinations by determining changes in the Up-Hallucinations Score after initiating a treatment.

[0019] Some non-limiting aspects of the invention include the following aspects.

[0020] Aspect 1, A method for assessing and treating schizophrenia and other psychotic

disorders in general, in particular delusions and risk of developing delusions in an individual, comprising the steps of: (a) obtaining a biological sample from an individual and quantifying the amounts of RNA biomarkers in the biological sample, to create a panel of RNA biomarkers, (b) quantifying the amounts of the RNA biomarkers in the panel in a clinically relevant population to generate a reference expression level for the RNA biomarkers in a panel of RNA biomarkers; (c) comparing the amounts of the biomarkers in the biological sample from the individual with the amounts of the RNA biomarkers present in the reference standard to generate a score for each biomarker; wherein the biomarkers in the a first panel (a) comprise one or more of the following RNA biomarkers: Activator Of Transcription and Developmental Regulator 2 (AUTS2), Pyruvate Dehydrogenase Phosphatase Catalytic Subunit 1 (PDP1), Nuclear Receptor Subfamily 4 Group A Member 2 (NR4A2), GNAS Complex Locus (GNAS), Interleukin 6 Signal Transduce (IL6ST), Chromodomain Helicase DNA Binding Protein 9 (CHD9), X-Ray Repair Cross Complementing 6 (XRCC6), RAR Related Orphan Receptor A (RORA), Actinin Alpha 4 (ACTN4), and Acyl-CoA Synthetase Long Chain Family Member 4 (ACSL4), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting delusions or an increased risk for developing delusions; and biomarkers in a second panel (b) comprise one or more of the following RNA biomarkers: Zinc Finger And BTB Domain Containing 20 (ZBTB20), Forkhead Box P1 (FOXP1), Spondin 1 (SPON1), and (NRP2), wherein the expression level of the RNA biomarker(s) in the sample is decreased relative to a reference expression level of the RNA biomarkers in the panel, denoting delusions or an increased risk for developing delusions; (d) generating a score for the panel of RNA biomarkers, based on the scores of the biomarker(s) in the panel; (e) determining a reference score for the panel in a clinically normal relevant population; (f) identifying a difference between the score of the panel of biomarker(s) in the sample and the reference score of the panel of biomarker(s); (g) identifying the individual as having delusions or of having an elevated risk for developing delusions, based on the difference between the biomarker panel score of the individual relative to the biomarker panel score of the reference; and (h) treating the individual identified as having delusions or an elevated risk of delusions with at least one treatment selected from the group consisting of: a treatment based on clinical practice guidelines, administering a therapeutically effective amount of at least one therapeutic drug wherein the mode of treatment is on the specific biomarkers scores indicating that individual will benefit

from a particular therapy.

[0021] Aspect 2, the method of aspect 1, wherein the biomarkers are quantified in samples taken on two or more occasions from the individual.

[0022] Aspect 3, the method of aspect 1, wherein each biomarker is assigned a weighted coefficient based on each biomarkers importance in in assessing and predicting delusions risk; and the biomarker panel score is based on the weighted coefficients of each of the biomarkers.

[0023] Aspect 4, the method of aspect 1, wherein the biological sample is at least sample from the individual selected from the group consisting of: tissue, a fluid such as cerebrospinal fluid, whole blood, blood serum, plasma, saliva, or other bodily fluid, or an extract or purification therefrom, or a dilution thereof.

[0024] Aspect 5, the method of aspect 1, wherein the therapeutic is at least drug selected from the group consisting of: adenosine phosphate, N-acetyl-L-leucine, eldeline, pempidine, verteporfin, C-75, oxprenolol, Prestwick-675, meglumine, guanethidine, pancuronium bromide, karakoline, 15(S)-15-methylprostaglandin E2, hexylcaine, dicoumarol, apramycin, mephénytoin, estriol, 528116.cdx, Cyclopiazonic Acid, SB 218078, BRD-A36630025, Quinacrine hydrochloride, GF-109203X, BRD-A36630025, N9-isopropylolomoucine, BMS-536924, BRD-K76951091, BRD-K26304855, trichostatin A, ALW-II-38-3, mitoxantrone, HG-6-64-0, alvocidib, SB-216763, and caffeic acid phenethyl ester.

[0025] Aspect 6, the method of aspect 1, wherein when the individual is male, and the biomarkers in a first panel (a) comprise one or more biomarkers selected from the group consisting of: Activator Of Transcription And Developmental Regulator (AUTS2), Pyruvate Dehydrogenase Phosphatase Catalytic Subunit 1 (PDP1), Forkhead Box P1 (FOXP1), GNAS Complex Locus (GNAS), Serine Racemase (SRR), Nuclear Receptor Subfamily 4 Group A Member 2 (NR4A2), X-Ray Repair Cross Complementing 6 (XRCC6), RAR Related Orphan Receptor A (RORA), and Actinin Alpha 4 (ACTN4), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting increased delusions, or the biomarkers in a second panel (b) comprising one or more biomarkers selected from the group consisting of: Zinc Finger And BTB Domain Containing 20 (ZBTB20), Forkhead Box P1 (FOXP1), Spondin 1 (SPON1), NRP2, wherein the expression level of the biomarker(s) in the sample is decreased relative to a reference expression level, denoting increased delusions.

[0026] Aspect 7, the method of aspect 6, wherein the at least one therapeutic drug is one or more drugs selected from the group consisting of: flunisolide, apramycin, adenosine phosphate, guanethidine, 15(S)-15-methylprostaglandin E2, meteneprost, methyl dopate, hydralazine, rotenone, phthalylsulfathiazole, N-acetyl-L-leucine, eldeline, tocainide, laudanosine, pempidine, 7-aminocephalosporanic acid, Sulfachlorpyridazine, finasteride, 528116.cdx, SB 218078, Quinacrine hydrochloride, N9-isopropylolomoucine, ALW-II-38-3, mitoxantrone, HG-6-64-01, Alvocidib, SB-216763, Syk Inhibitor, Cyclopiazonic Acid, GW 441756, LY 225910, AG 82, doxorubicin, mitomycin, and terfenadine.

[0027] Aspect 8, the method of aspect 1, wherein when the individual is female, and the biomarkers in the panel comprise one or biomarkers in a first panel (a) comprise one or more of the biomarkers selected from the group consisting of: Phosphodiesterase 4D Interacting Protein (PDE4DIP), Pyruvate Dehydrogenase Phosphatase Catalytic Subunit 1 (PDP1), Transcription Factor 4 (TCF4), Nuclear Receptor Subfamily 4 Group A Member 2 (NR4A2), Chromodomain Helicase DNA Binding Protein 9 (CHD9), (CLCN3), Activator Of Transcription And Developmental Regulator (AUTS2), and (LDB2), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting increased delusions; and the biomarkers in a second panel (b) comprise one or more biomarkers selected from the group consisting of (FGFR1), (DISC1), (FGFR2), (SPTBN1), (INSR), (GRIK3), Zinc Finger, and BTB Domain Containing 20 (ZBTB20), wherein the expression level of the biomarker(s) in the sample is decreased relative to a reference expression level, denoting increased delusions.

[0028] Aspect 9, the method of aspect 8, wherein the at least one therapeutic drug is at least drug selected from the group consisting of: erastin, harpagoside, metacycline, amiodarone, furaltadone, metformin, timolol, Repaglinide, sulfafurazole, PNU-0230031, Probenecid, furosemide, fluphenazine, myricetin, sulfacetamide, lomustine, BCB000039, Harmalol, I-BET151, Nylidrin hydrochloride, AMG 9810, Doxorubicin, Mitomycin C, Fludrocortisone acetate, Purvalanol A, Teniposide, Geldanamycin, Importazole, BRD-A36630025, YM-155, Auranofin, 7643453, G-221, BRD-A49680073, BRD-K08547377, and Cladribine.

[0029] Aspect 10, a method of assessing and treating schizophrenia and other psychotic disorders in general, and delusions in particular, in an individual, comprising: calculating combined biomarkers and clinical information Up-based on the equation: (Biomarker Panel

Score) + (Delusions Score) - (Grooming Score) = Up-Delusions Score; wherein the Biomarker Panel Score is obtained as per the method of claim 1; wherein the Delusions Score is calculated with a clinical rating or self-report scales; wherein the Grooming Score is calculated with a rating scale; assessing the level of delusions of the individual by comparing the individual's Up-Delusions Score to a reference Up-Delusions Score; administering a treatment for delusions to the individual when the individual's Up-Delusions Score is greater than a reference Up-Suicide Score; and monitoring the individual's response to a treatment for delusions by determining changes in the Up-Delusions Score after initiating a treatment.

[0030] Aspect 11, a method for assessing and treating schizophrenia and other psychotic disorders in general, in particular hallucinations and risk of developing hallucinations in an individual, comprising the steps of: (a) obtaining a biological sample from an individual and quantifying the amounts of one or more RNA biomarkers in the biological sample, to create at least one panel of RNA biomarkers, (b) quantifying the amounts of the RNA biomarkers in the at least one panel in a clinically relevant population to generate a reference expression level for the RNA biomarkers in a panel of RNA biomarkers; (c) comparing the amounts of the biomarkers in the biological sample from the individual with the amounts of the RNA biomarkers present in the reference standard to generate a score for each biomarker a first panel and a second panel; wherein the biomarkers in the first panel comprise one or more of the following RNA biomarkers: (PRICKLE1), (NCAM1), (B3GALT5), (ARHGAP18), (PTP4A2), Acylphosphatase 2 (ACYP2), Reticulon 4 (RTN4), Cullin 4A (CUL4A), Zinc Finger E-Box Binding Homeobox 2 (ZEB2), Dystonin (DST), and Discs Large MAGUK Scaffold Protein 1 (DLG1), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting hallucinations or an increased risk for developing hallucinations; and wherein the biomarkers in the second panel comprise one or more of the following RNA biomarkers: (PRL), (SERPING1), Ectonucleotide Pyrophosphatase/Phosphodiesterase 2 (ENPP2), (LAMA4), (KCNV1), Catenin Delta 1 (CTNND1), and FAT Atypical Cadherin 4 (FAT4), wherein the expression level of the RNA biomarker(s) in the sample is decreased relative to a reference expression level of the RNA biomarkers in the panel, denoting hallucinations or an increased risk for developing hallucinations; (d) generating a score for the panel of RNA biomarkers, based on the scores of the biomarker(s) in the panel; (e) determining a reference score for the panel in a clinically normal relevant population; (f) identifying a difference between the score of the panel

of biomarker(s) in the sample and the reference score of the panel of biomarker(s); (g) identifying the individual as manifesting hallucinations or of having an elevated risk for developing hallucinations, based on the difference between the biomarker panel score of the individual relative to the biomarker panel score of the reference; and (h) treating the individual identified as having hallucinations or an elevated risk of hallucinations with one or more of the following: 1) a treatment based on clinical practice guidelines, 2) administering a therapeutically effective amount of a therapeutic drug (s), selected based on the specific biomarkers whose scores indicate that they are changed in the individual compared to a reference standard.

[0031] Aspect 12, the method of aspect 11, wherein the biomarkers are quantified in samples taken on two or more occasions from the individual.

[0032] Aspect 13, the method of aspect 11, wherein each biomarker is assigned a weighted coefficient based on the biomarkers importance in assessing and predicting hallucinations risk; and the biomarker panel score is based on the weighted coefficients of each of the biomarkers.

[0033] Aspect 14, the method of aspect 11, wherein the biological sample is a tissue sample or a fluid, such as cerebrospinal fluid, whole blood, blood serum, plasma, saliva, or other bodily fluid, or an extract or purification therefrom, or dilution thereof.

[0034] Aspect 15, the method of aspect 11, wherein the one or more therapeutic is one or more compounds selected from the group consisting of: clioquinol, pirinixic acid, moxislyte, Prestwick-685, exemestane, azacytidine, C-75, estradiol, tetraethylenepentamine, sparteine, guanethidine, idoxuridine, gliclazide, nitrendipine, N-acetyl-L-aspartic acid, sulfanilamide, doxazosin, pimozide, Proscillaridin, oxetacaine, BRD-K71489689, trichostatin A, A443654, AG 825, Proscillaridin A, Ala-Ala-Phe-CMK, Fluocinolone acetonide, manumycin A, curcumin, BRD-K68548958, CHR 2797, Tyrphostin AG 1478, Wortmannin, HY-50878, 598226, S1003, BRD-A52530684, CGP-60474, Buparlisib, and AS-601245.

[0035] Aspect 16, the method of aspect 11, wherein when the individual is male, and the biomarkers in a first panel (a) comprise one or more biomarkers selected from the group consisting of: (SH3PXD2A), Zinc Finger E-Box Binding Homeobox 2 (ZEB2), (PRICKLE1), (ARHGAP18), Acylphosphatase 2 (ACYP2), Reticulon 4 (RTN4), and Dystonin (DST), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting increased hallucinations; and biomarkers in a second panel (b) comprise one or more biomarkers selected from the group consisting of: (PRL), (SERPING1),

Ectonucleotide Pyrophosphatase/Phosphodiesterase 2 (ENPP2), (KCNV1), Mab-21 Like 1 (MAB21L1), Catenin Delta 1 (CTNND1), and FAT Atypical Cadherin 4 (FAT4), wherein the expression level of the biomarker(s) in the sample is decreased relative to a reference expression level, denoting increased hallucinations.

[0036] Aspect 17, the method of aspect 16, wherein the at least one therapeutic is at least one compound selected from the group consisting of: digoxigenin, doxazosin, meptazinol, promethazine, cefixime, velnacrine, cetirizine, eldeline, atropine oxide, clioquinol, nicotinic acid, clioquinol, galantamine, rolitetracycline, betahistine, sulconazole, monocrotaline, lanatoside C, Prestwick-1084, Naftidrofuryl, sulfachlorpyridazine, helveticoside, bezafibrate, mifepristone, trichostatin A, manumycin A, NCGC00189555-02, Buparlisib, linifanib, AZD-7762, Dinaciclib, Piretanide, KN-62, Fluticasone propionate, JAK3 Inhibitor VI, Sarmentogenin, Digoxin, Megestrol acetate, Oxymetazoline hydrochloride, U-0126, Tracazolate hydrochloride, Flufenamic acid, Fenofibrate, and U 99194 maleate.

[0037] Aspect 18, the method of aspect 11, wherein when the individual is female, and the biomarkers in a first panel (a) comprise one or more biomarkers selected from the group consisting of: (CELSR2), (KALRN), (B3GALT5), Protein Phosphatase 3 Catalytic Subunit Beta (PPP3CB), (ZFR), (THNSL1), (TNIK), Nuclear Receptor Subfamily 4 Group A Member 2 (NR4A2), Zinc Finger E-Box Binding Homeobox 2 (ZEB2), and (TNIK), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting increased hallucinations; and biomarkers in a second panel (b) comprise one or more biomarkers selected from the group consisting of GNAS Complex Locus (GNAS), and Catenin Delta 1 (CTNND1), wherein the expression level of the biomarker(s) in the sample is decreased relative to a reference expression level, denoting increased hallucinations.

[0038] Aspect 19, the method of aspect 18, wherein the at least one therapeutic is at least one compound selected from the group consisting of: proglumide, quinethazone, esculin, MG-262, GW-8510, haloperidol, guanethidine, deferoxamine, citiolone, meteneprost, amylocaine, CP-944629, Clemizole, IC-86621, Nortriptyline, CP-944629, Tanespimycin, Prestwick-674, 0317956-0000, and Pioglitazone.

[0039] Aspect 20, a method of assessing and treating schizophrenia and other psychotic disorders in general, and hallucinations in particular in an individual, comprising: calculating combined biomarkers and clinical information Up- based on the equation: (Biomarker Panel

Score) + (Hallucinations Score) - (Grooming Score) = Up-Hallucinations Score; wherein the Biomarker Panel Score is obtained as per the method of aspect 11; wherein the Hallucinations Score is calculated with a clinical rating or self-report scales; wherein the Grooming Score is calculated with a rating scale; assessing the level of hallucinations of the individual by comparing the individual's Up-Hallucinations Score to a reference Up-Hallucinations Score; administering a treatment for hallucinations to the individual when the individual's Up-Hallucinations Score is greater than a reference Up-Suicide Score; and monitoring the individual's response to a treatment for hallucinations by determining changes in the Up-Hallucinations Score after initiating a treatment.

BRIEF DESCRIPTION OF THE FIGURES

- [0040] FIG. 1 A. Cohorts used in study, depicting flow of discovery, prioritization, and validation of biomarkers from each step (N of 74).
- [0041] FIG. 1 B. Discovery cohort longitudinal within-subject analysis. Phchp### is study ID for each subject. V# denotes visit number.
- [0042] FIG. 1 C. The clinical phenotypic measure (item from PANSS) used for discovery.
- [0043] FIG. 1 D. Validation in independent cohort of psychiatric patients with clinically severe psychosis.
- [0044] FIG. 1 E. Validation in independent cohort of psychiatric patients with clinically severe psychosis.
- [0045] FIG. 1 F. The 4 step process of discovery, prioritization, validation, and testing.
- [0046] FIG. 1 G. Schematic illustrating the process of biomarker validation.
- [0047] FIG. 1 H. Cohorts used in study, depicting flow of discovery, prioritization, and validation of biomarkers from each step.
- [0048] FIG. 1 I. Discovery cohort longitudinal within-subject analysis. Phchp### is the study identifier (ID) for each subject. V# denotes visit number.
- [0049] FIG. 1 J. The clinical phenotypic measure (item from PANSS) used for discovery (N of 36).
- [0050] FIG. 1 K. Validation in independent cohort of psychiatric patients with clinically severe psychosis (N of 36).
- [0051] FIG. 1 L. The 4 step process of discovery, prioritization, validation, and testing.

- [0052] FIG. 1 M. Schematic illustrating the process of biomarker validation.
- [0053] FIG. 2A. Top Predictive Biomarkers for Different Demographic and Disease Groups. Delusions A-C, high delusions state predictions ($p \geq 4$).
- [0054] FIG. 2B. Top Predictive Biomarkers for Different Demographic and Disease Groups. Delusions A-C, delusions trait predictions first year
- [0055] FIG. 2C. Top Predictive Biomarkers for Different Demographic and Disease Groups. Delusions A-C, delusion trait all future years.
- [0056] FIG. 2D. Top Predictive Biomarkers for Different Demographic and Disease Groups. Hallucinations D-F, high hallucinations state predictions.
- [0057] FIG. 2E. Top Predictive Biomarkers for Different Demographic and Disease Groups. Hallucinations, hallucinations trait predictions first year.
- [0058] FIG. 2F. Top Predictive Biomarkers for Different Demographic and Disease Groups. Hallucinations, hallucinations trait predictions all future years.
- [0059] FIG. 3A. Network analysis of top candidate biomarkers, delusions.
- [0060] FIG. 4A. Example of a possible report to communicate to clinicians – delusion severity.
- [0061] FIG. 4B. Example of a possible report to communicate to clinicians – hallucinations severity.

DETAILED DESCRIPTION

[0062] We endeavored to find objective blood biomarkers for hallucinations and delusions, two core positive psychotic symptoms that are transdiagnostic, and sought to see whether these biomarkers can be used to track and predict clinical course. Our initial work over a decade ago (Kurian, Le-Niculescu et al. 2009)¹ indicated that may be possible. We now used a more advanced methodology, with a longitudinal within-subject design for discovery, followed by prioritization, validation, and testing in independent cohorts. This comprehensive four-step approach is similar to the one used in our more recent studies on mood disorders, memory, stress, pain, and delusions. Provided here are newly identified blood gene expression biomarkers for hallucinations, and for delusions. These biomarkers opened a window into disease biology and core gene networks involved. Second, they permit objective assessment of state severity, short-term risk, and long-term risk. Third, they were used for drug repurposing. Lastly, we provide an example of how a precision medicine report for a patient would look, with objective

scores for severity, and list of prioritized suggested medications. Such tools can be used for informing assessment, treatment choices, and monitoring response to treatment, and ultimately in prevention. Their integration in routine clinical practice and new drug development, including use and development of psychedelic drugs, can be transformative in an area that is in great need of progress.

[0063] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the disclosure belongs. Although any methods and materials similar to or equivalent to those described herein can be used in the practice or testing of the present disclosure, the preferred methods and materials are described below.

[0064] The present disclosure is generally directed at methods for assessing psychosis and/or schizophrenia and early identification of risk for future schizophrenia, as well as methods for matching patients and drugs for prevention and mitigation of schizophrenia and symptoms such as hallucinations and/or delusion, and for monitoring response to treatment. The methods may further include the generation of a report providing a risk score and/or personalized treatment options. Further, the present disclosure generally is directed to drugs for mitigating schizophrenia in subjects. Particular drugs have been found that can mitigate schizophrenia in subjects universally; that is, drugs that can be used for mitigating schizophrenia across psychiatric diagnoses and genders. Some drugs, however, have been found that can be used more effectively for mitigating schizophrenia dependent on gender, psychiatric diagnoses, and combinations thereof.

[0065] In additional embodiments, the present disclosure is directed to blood gene expression biomarkers that are more universal in nature; that is, blood biomarkers that can be used for predicting schizophrenia across psychiatric diagnoses and genders. Accordingly, a longitudinal within-participant design and large cohorts were used.

[0066] Additionally, subtypes of schizophrenia were identified based on mental state (hallucinations, delusions, psychosis) at the time that schizophrenia was manifest.

Materials and Methods

Cohorts

[0067] For each of the 2 core schizophrenia psychotic symptoms, delusions and

hallucinations, we used three independent cohorts:

1. discovery (a longitudinal psychiatric subjects cohort with diametric changes in state from at least two consecutive testing visits);
2. validation (an independent psychiatric subjects cohort with clinically severe psychosis); and
3. testing (an independent psychiatric subjects test cohort for predicting psychosis state, and for predicting future hospitalizations with psychotic symptoms) (Figure 1 and Table 1).

[0068] Similar to our previous studies, the live psychiatric subjects are part of a larger longitudinal cohort of adults that we are continuously collecting. Subjects are recruited primarily from the patient population at the Indianapolis VA Medical Center. All subjects understood and signed informed consent forms detailing the research goals, procedure, caveats and safeguards, per IRB approved protocol. Subjects completed diagnostic assessments by structured clinical interviews. They had an initial testing visit in the lab or on the inpatient psychiatric unit, followed by up to six testing visits, 3–6 months apart or whenever a new psychiatric hospitalization occurred. At each testing visit, they received a series of psychiatric rating scales, and their blood was drawn. The rating scales included the clinically used Positive and Negative Symptoms Scale (PANSS), containing the PANSS Positive sub-scale, that measures positive psychotic symptoms including delusions and hallucinations, as well as a new visual analog scale for assessing psychosis state, which provides a score that is the average of several items. This scale provides a score for psychosis state at a particular moment in time.

[0069] At each visit, we collected whole blood (5 ml) in two RNA-stabilizing PAXgene tubes, labeled with an anonymized study ID number, and stored at -80°C. in a locked freezer until the time of future processing. Whole-blood RNA was extracted for microarray gene expression studies from the PAXgene tubes, as detailed below.

[0070] For this study, for the delusions biomarker part, our within-subject discovery cohort, from which the biomarker data were derived, consisted of 31 subjects with psychiatric disorders and multiple testing visits, who each had at least one diametric change in PANSS item P1 Delusion score from no symptoms (score =1) to high symptoms (score $\geq$ 4), or vice versa, from one testing visit to another. There are a total of 95 blood samples for subsequent gene expression microarray studies (Figure 1, Table 1).

[0071] For the hallucinations biomarker part, our within-subject discovery cohort, from which the biomarker data were derived, consisted of 25 subjects with psychiatric disorders and multiple testing visits, who each had at least one diametric change in PANSS item P3 Hallucinations score from no symptoms (score =1) to high symptoms (score $\geq$ 4), or vice versa, from one testing visit to another. There are a total of 65 blood samples for subsequent gene expression microarray studies (Figure 1, Table 1).

[0072] Our independent validation cohort, in which the top biomarker findings were validated for being even more changed in expression, consisted of 43 subjects for delusions, and 36 subjects for hallucinations, with clinically severe psychosis (Table 1). Our independent test cohort for predicting consisted of 120 subjects for delusions, and 196 subjects for hallucinations (Figure 1 and Table 1).

[0073] *Medications.* The subjects in the discovery cohort were all diagnosed with various psychiatric disorders (Table 1) and had various medical co-morbidities. Their medications were listed in their electronic medical record and documented by us at the time of each testing visit. Medications can have a strong influence on gene expression. However, there was no consistent pattern of any particular type of medication, as our subjects were on a wide variety of different medications, psychiatric and non-psychiatric. Furthermore, the independent validation and testing cohort's gene expression data was Z-scored by gender and by diagnosis before being combined, to normalize for any such effects. Some subjects may be non-compliant with their treatment and may thus have changes in medications or drug of abuse not reflected in their medical records. That being said our goal is to find biomarkers that track psychosis, regardless if the reason for it is endogenous biology or it is driven by medications or drugs. Moreover, the prioritization step that occurs after discovery is based on a field-wide convergence with literature that includes genetic data and animal model data, that are unrelated to medication effects. Overall, the discovery, validation and replication by testing in independent cohorts of the biomarkers, with our design, occurs despite the subjects having different genders, diagnoses, being on various different medications, and other lifestyle variables.

Blood gene expression experiments

[0074] *RNA extraction.* Whole blood (2.5 ml) was collected into each PaxGene tube by routine venipuncture. PaxGene tubes contain proprietary reagents for the stabilization of RNA.

RNA was extracted and processed as previously described^{2,3,4}.

Microarrays. Microarray work was carried out using previously described methodology^{2,3,4,5}.

[0075] Of note, all genomic data was normalized (RMA for technical variability, then z-scoring for biological variability), by gender and psychiatric diagnosis, before being combined and analyzed.

Biomarkers

Step 1: Discovery

[0076] We have used the subject's score from a visual-analog scale PANSS scale P1 and P3 items, assessed at the time of blood collection (Figure 1). We analyzed gene expression differences between visits with low psychosis (defined as a score of 1) and visits with high psychosis (defined as a score of ≥ 4), using a powerful within-subject design, then an across-subjects summation (Figure 1).

[0077] We analyzed the data in two ways: an Absent-Present (AP) approach, and a differential expression (DE) approach, as in previous work by us on suicide biomarkers²⁻⁴. The AP approach may capture turning on and off of genes, and the DE approach may capture gradual changes in expression. Analyses were performed as previously described³⁻⁵. In brief, we imported all Affymetrix microarray data as CEL files into Partek Genomic Suites 6.6 software package (Partek Incorporated, St Louis, MI, USA). Using only the perfect match values, we ran a robust multi-array analysis (RMA) by gender and diagnosis, background corrected with quantile normalization and a median polish probeset summarization of all chips, to obtain the normalized expression levels of all probesets for each chip. Then, to establish a list of differentially expressed probesets we conducted a within-subject analysis, using a fold change in expression of at least 1.2 between consecutive high- and low mood visits within each subject. Probesets that have a 1.2-fold change are then assigned either a 1 (increased in high mood) or a -1 (decreased in high mood) in each comparison. Fold changes between 1.1 and 1.2 are given 0.5, and fold changes less than 1.1 are given 0. These values were then summed for each probeset across all the comparisons and subjects, yielding a range of raw scores. The probesets above the 33.3% of raw scores were carried forward in analyses (Figure 1), and received an internal score of 2 points; those above 50% 4 points, and those above 80% 6 points^{3 4 5}. We have developed in our labs R scripts to automate and conduct all these large dataset analyses in bulk, checked against

human manual scoring⁵.

[0078] Gene Symbol for the probesets were identified using NetAffyx (Affymetrix) for Affymetrix HG-U133 Plus 2.0 GeneChips, followed by GeneCards to confirm the primary gene symbol. In addition, for those probesets that were not assigned a gene symbol by NetAffyx, we used GeneAnnot (<https://genecards.weizmann.ac.il/geneannot/index.shtml>), or if need be UCSC (<https://genome.ucsc.edu>), to obtain gene symbol for these uncharacterized probesets, followed by GeneCard. Genes were then scored using our manually curated CFG databases as described below (Figure 1D).

Step 2: Prioritization using Convergent Functional Genomics (CFG)

[0079] *Databases.* We have established in our laboratory (Laboratory of Neurophenomics, www.neurophenomics.info) manually curated databases of the human gene expression/protein expression studies (postmortem brain, peripheral tissue/fluids: CSF, blood and cell cultures), human genetic studies (association, copy number variations and linkage), and animal model gene expression and genetic studies, published to date on psychiatric disorders. Only findings deemed statistically scientifically using particular experimental design and thresholds are included in the databases. Our databases include only primary literature data and do not include review papers or other secondary data integration analyses to avoid redundancy and circularity. We also favored unbiased discovery studies over candidate genes hypothesis-driven studies. These large and constantly updated databases have been used in our CFG (Convergent Functional Genomics) cross validation and prioritization platform (Figure 1).

Step 3: Validation analyses

[0080] We examined which of the top candidate genes (score of 6 or above after the first two steps), were stepwise changed in expression from the clinically depressed validation group to the low mood discovery group to the high mood discovery group to the clinically manic validation group. A total score of 6 or above after the first two steps permits the inclusion of potentially novel genes with maximal internal score of 6 from Discovery but no external evidence CFG score from Prioritization.

[0081] The AP derived and DE derived lists of genes were combined, and the gene expression data corresponding to them was used for the validation analysis. We transferred the log transformed expression data to an Excel sheet, and non-log transformed the data by taking 2

to the power of the transformed expression value. We then Z-scored the values by gender and diagnosis. We then imported the Excel sheets with the Z-scored by gender and diagnosis expression data into Partek, and statistical analyses were performed using a one-way ANOVA for the stepwise changed probesets, and also did a stringent Bonferroni correction for all the probesets tested in ANOVA (Figure 1).

Top Candidate Biomarkers (after the first 3 steps)

[0082] Adding the scores from the first three steps into an overall convergent functional evidence (CFE) score (Figure 1), we ended up with a list of 70 top candidate biomarkers for delusions, and 213 for hallucinations (Table 2 A and 2B). These top candidate biomarkers were carried forward into additional testing for clinical utility.

Networks

[0083] For network analyses we performed STRING Interaction network (<https://string-db.org>) by inputting the genes into the search window and performed Multiple Proteins Homo sapiens analysis. (Figure 3).

Testing for Clinical Utility in Independent Cohorts

[0084] We tested in independent cohorts of psychiatric patients the ability of each of the top candidate biomarkers to assess state severity, and predict trait risk (future hospitalizations with psychosis). We conducted our analyses across all patients, as well as personalized by gender and diagnosis.

[0085] The test cohort for predicting psychosis (delusions, hallucinations) were assembled out of data that was RMA normalized by gender and diagnosis. The cohort was completely independent from the discovery and validation cohorts, there was no subject overlap with them. Individual markers used for predictions were Z scored by gender and diagnosis, to be able to combine different biomarkers into panels and to avoid potential artefacts due to different ranges of expression in different gender and diagnoses. For panels, biomarkers were combined by simple summation of the increased risk biomarkers minus the decreased risk biomarkers. Predictions were performed using R-studio (opensource). For cross-sectional analyses, we used biomarker expression levels, z-scored by gender and diagnosis. For longitudinal analyses, we combined four measures: biomarker expression levels, slope (defined as ratio of levels at current testing visit vs. previous visit, divided by time between visits), maximum levels (at any of the

current or past visits), and maximum slope (between any adjacent current or past visits). For decreased biomarkers, we used the minimum rather than the maximum for level calculations. All four measures were Z-scored, then combined in an additive fashion into a single measure. The longitudinal analysis was carried out in a sub-cohort of the testing cohort consisting of subjects that had at least two subject visits (timepoints).

[0086] *Predicting State Severity.* Receiver-operating characteristic (ROC) analyses between marker levels and psychosis state were performed.

[0087] *Predicting Trait- Future Psychiatric Hospitalization with Psychosis as a Symptom/Reason for Admission.* We conducted analyses for predicting future psychiatric hospitalizations with psychosis as a symptom/reason for admission in the first year following each testing visit, in subjects that had at least one year of follow-up in the VA system, in which we have access to complete electronic medical records. ROC analyses between biomarkers measures (cross-sectional, longitudinal) at a specific testing visit and future hospitalizations within the first year were performed. A Cox regression was performed for all future hospitalizations. The odds ratio was calculated such that a value greater than 1 always indicates increased risk for hospitalization, regardless if the biomarker is increased or decreased in expression.

Therapeutics

[0088] *Pharmacogenomics.* We analyzed which of the top biomarkers for psychosis after Steps 1- 4 are known to be changed in expression by existing drugs in a direction opposite to the one in disease, using our CFG databases, and using Ingenuity Drugs analyses (Table 3).

[0089] *New drug discovery/repurposing.* We also analyzed which drugs and natural compounds are an opposite match for the gene expression signatures of our top biomarkers, using the Connectivity Map (<https://portals.broadinstitute.org>, Broad Institute, MIT) (Table 4). Of note, not all the probesets from the HG-U133 Plus 2.0 array we used were present in the HGU-133A array used for the Connectivity Map. We stayed with exact probeset level matches, not gene level imputation. We also used the NIH LINCS database to conduct similar analyses, at a gene level.

Report generation

[0090] We present an example of how a report to doctors might look, using the described

methods and identified biomarkers. Out of a dataset of 794 subject visits, we chose, as case studies, a visit from a patient with self-reported high delusions, and one from a patient with self-reported high hallucinations (Figure 4).

[0091] For each biomarker in the panel, we also have a list of existing psychiatric medications that modulate the expression of the biomarker in the direction of high mood. Each medication got a score of 1 or 0 whether it modulated a particular biomarker in the panel or not, and that score is multiplied with the risk score of the biomarker, i.e. 1 or 0.5 or 0. A medication can modulate more than one biomarker. We then calculated an average score for each medication based on its effects on all the biomarkers in the panel, and multiplied that by 100, resulting in a % score for each medication. Thus, psychiatric medications are matched to the patient and ranked in order of impact on the panel.

Table 1. Demographics of patient cohorts used. A. Delusions, B. Hallucinations

A. Delusions	Number of subjects (number of visits)	Gender	Diagnosis	Ethnicity	Age in years at time of lab visit Mean (SD) (Range)	T-test for age at time of lab visit
Discovery						
Discovery Cohort (Longitudinal Within-Subject Changes in Delusion) P1>=4 high Delusions P1<=1 No Delusions	31 (with 95 visits)	Male 27 (86) Female 4 (9)	BP = 10 (35) MDD= 2 (5) PSYCH= 1 (3) PTSD= 3 (12) SZ= 7 (21) SZA= 8 (19)	AA= 10 (28) EA= 21 (67)	49.45 (7.38) (30-63)	
Validation						
Independent Validation Cohort for Gene Expression (P1>=4 high Delusion and PANSS Positive Total >= 21)	43 (with 62 visits)	Male 36 (52) Female 7 (10)	BP=7(8) MOOD= 1(2) PSYCH= 1 (1) PTSD=3(6) SZ=15(23) SZA=16(22)	AA= 16(25) EA= 27(37)	46.11 (11.55) (22-62)	T-test for age Validation (Clinical) vs Stepwise Discovery (High and Low) =.11
Stepwise Discovery Cohorts Used for Validation P1>=4 high Delusions & PANSS Positive <21 P1<=1 No Delusions	Stepwise Discovery High					
	23 (with 25 visits)	Male 21 (23) Female 2 (2)	BP=6(6) MDD= 1(1) PTSD= 3 (3) SZ=6(8) SZA=7(7)	AA= 7(8) EA= 16(17)	48.52 (6.87) (33-61)	
	Stepwise Discovery Low					
	31 (with 56 visits)	Male 27 (52) Female 4(4)	BP=10(23) MDD= 2(3) PSYCH= 1 (2) PTSD= 3 (9) SZ=7(9) SZA=8(10)	AA= 10(15) EA= 21(41)	49 (7.1) (31-63)	
Testing						
Independent Testing Cohort For Predicting State (High Delusions P1 ≥4, P1 <= 3 (others) at Time of Assessment)	120 (with 315 visits)	Male 109 (285) Female 11 (30)	BP = 51 (147) MDD= 10 (28) SZ= 27 (64) SZA= 32 (76)	AA= 25 (63) EA= 94 (249) Hispanic= 1 (3)	All 51 (9.1) (23-74) High Delusions (N= 36) 51.3 Others (N= 279) 51	T-test for age High Delusions (N=36) vs. Others (N= 279) 0.8
Independent Testing Cohort For Predicting Trait (All Future Hospitalizations with Delusions following Assessment)	113(with 292 visits)	Male 99 (253) Female 14 (39)	BP = 44 (122) MDD= 9 (25) PSYCH= 3(7) PTSD= 1(3) SZ= 25 (60) SZA= 31(75)	AA= 35 (87) EA= 77 (202) Hispanic= 1 (3)	All = 49.6 (8.21) (23-63) No Hosp for Delusions (N= 212) 49.8 Hosp for Delusions (N= 80)	T-test for age No Hosp with Delusions (N= 212) vs. Hosp with Delusions (N= 80) 0.5177799

					49.1	
Independent Testing Cohort For Predicting Trait (Hospitalizations with Delusions in the First Year Following Assessment)	95 (with 254 visits)	Male 91 (243) Female 4 (11)	BP = 45 (135) PSYCH= 3(7) SZ=27(64) SZA=20(48)	AA= 18 (48) EA= 77 (206)	All = 50.9 (9.21) (23-74) No Hospitalizations visit for Delusions (N= 239) 50.9 Hospitalizations for Delusions (N= 15) 50.5	T-test for age No Hosp with Delusions (N= 239) vs. Hosp with Delusions within the first Year (N= 15) 0.815349

B. Hallucinations	Number of subjects (number of visits)	Gender	Diagnosis	Ethnicity	Age in years at time of lab visit Mean (SD) (Range)	T-test for age at time of lab visit
Discovery						
Discovery Cohort (Longitudinal Within-Subject Changes in Hallucination) P3>=4 high Hallucinations P3<=1 No Hallucinations	25 (with 65 visits)	Male 20(53) Female 5 (12)	BP =5(14) MDD=3(6) MOOD=1(2) PSYCH=2(5) SZ=6(18) SZA=8(20)	AA=12(31) EA=13(34)	49.57 (10) (24-62)	
Validation						
Independent Validation Cohort for Gene Expression (P3>=4 high Hallucination and PANSS Positive Total >= 21)	36 (with 52 visits)	Male 27(39) Female 9 (13)	BP=10 (13) PTSD=4 (6) SZ=12 (21) SZA=10 (12)	AA= 12(17) EA= 24(35)	46.92 (10) (22-63)	T-test for age Validation (Clinical) vs Stepwise Discovery (High and Low) =.542434
Stepwise Discovery Cohorts Used for Validation P3>=4 high Hallucination & PANSS Positive <21 P3<=1 No Hallucination	Stepwise Discovery High					
	16 (with 21 visits)	Male 13(18) Female 3 (3)	BP=4(5) MDD=2(2) PSYCH=1(2) SZ=4(7) SZA=5(5)	AA= 7(11) EA= 9(10)	50.33 (8.5) (24-61)	
	Stepwise Discovery Low					
	19 (with 24 visits)	Male 15(19) Female 4(5)	BP=5(8) MDD=2(2) MOOD=1(1) PSYCH= 1(1) SZ=3(3) SZA=7(9)	AA= 9(11) EA= 10(13)	46.17 (9) (24-56)	
Testing						
Independent Testing Cohort For Predicting State	196 (with 513 visits)	Male 162 (426)	BP = 74 (207) MDD= 22 (55)	AA= 55 (135) EA= 139 (372)	All 50.1	T-test for age High P3 (N=62)

(High Hallucination P3 ≥4, P3 ≤3 (others) at Time of Assessment)		Female 34 (87)	MOOD= 6 (17) PTSD= 25 (64) SZ= 30 (74) SZA= 39 (96)	Hispanic= 2 (6)	(9.2) (20-74) High P3 (N= 62) 50.7 Others (N= 451) 50.0	vs. Others (N= 451) 0.54722
Independent Testing Cohort for Predicting Trait (Hospitalizations with Hallucinations in the First Year Following Assessment)	163 (with 437 visits)	Male 146 (389) Female 17 (48)	BP = 55 (165) MDD= 30 (77) PTSD= 14 (39) SZ= 29 (72) SZA= 35 (84)	AA= 33 (84) EA= 130 (353)	All = 50.3 (8.3) (23-65) No Hosp for P3 (N= 415) 50.5 Hosp for P3 (N= 22) 49.3	T-test for age No Hosp with Hallucination (N= 415) vs. Hosp with Hallucination within the first Year (N= 22) 0.509856
Independent Testing Cohort for Predicting Trait (Hospitalizations with Hallucinations in All Future Years Following Assessment)	170 (with 442 visits)	Male 153 (392) Female 17 (50)	BP = 54 (152) MDD= 40 (101) PTSD= 14 (39) SZ= 27 (66) SZA= 35 (84)	AA= 51 (123) EA= 118 (316) Hispanic= 1 (3)	All = 49.4 (7.9) (23-66) No Hosp for P3 (N= 334) 49.5 Hosp for P3 (N= 109) 48.5	T-test for age Future Hosp with no Hallucination (N= 334) vs. Hallucination with Hosp (N= 109) 0.50054434

[0092] Referring now to Figure 1. Steps 1-3: Discovery, Prioritization and Validation of Biomarkers. See for examples Figures 1A-G Delusions; Figures 1. H-M Hallucinations (A.) Cohorts used in study, depicting flow of discovery, prioritization, and validation of biomarkers from each step. (B.) Discovery cohort longitudinal within-subject analysis. Phchp### is study ID for each subject. V# denotes visit number. (C.) The clinical phenotypic measure (item from PANSS) used for discovery. (D.) Prioritization with CFG for prior evidence of involvement in mood disorders. In the prioritization step probesets are converted to their associated genes using Affymetrix annotation and GeneCards. Genes are prioritized and scored using CFG for schizophrenia/psychosis evidence with a maximum of 12 points. Genes scoring at least 6 points out of a maximum possible of 18 total discovery and prioritization scores points are carried to the validation step. (E.) Validation in independent cohort of psychiatric patients with clinically severe psychosis

Table 2. Top Candidate biomarkers after Steps 1-3. A. Delusions. B. Hallucinations.

[0093] In Table 2, “D” indicates a decrease in RNA biomarker expression measured in a population identified with delusions or hallucinations related to a population that do not manifest either delusion or hallucinations; “I” indicates an increase in RNA biomarker expression measured in a population identified with either delusion or hallucinations related to a population that do not manifest either delusion or hallucinations.

A. For Delusions (n=70)	
Gene	Direction of Change
DAG1biom212128_s_at	D
LPAR1biom204037_at	I
CTNND1biom1557944_s_at	D
POLBbiom234352_x_at	D
ERBB4biom241581_at	D
INSRbiom207851_s_at	D
IL6biom205207_at	D
KMObiom205307_s_at	I
NR4A2biom204621_s_at	I
PDE4DIPbiom205872_x_at	I
PTPRZ1biom204469_at	I
RORAbiom240951_at	I
SNAP25biom1556629_a_at	D
TNIBbiom213107_at	D
TCF4biom212382_at	I
ZFRbiom33148_at	I
CNPbiom1557943_at	D
ACTN4biom241788_x_at	I
AUTS2biom242721_at	I
ACSL4biom224091_at	I
AKAP10biom213396_s_at	D
ANKRD11biom1566001_at	D
CREB1biom234289_x_at	D
COMTbiom236804_at	I
CDC42biom214230_at	D
CLCN3biom240237_at	I
CHD9biom220586_at	I
DPYDbiom1554534_at	D
DISC1biom206090_s_at	D
DISC1biom207759_s_at	D
EGR1biom227404_s_at	I

A. For Delusions (n=70)	
Gene	Direction of Change
EDIL3biom225275_at	I
ELMO1biom1562894_at	D
FGFR1biom226705_at	D
FGFR2biom230842_at	D
FOXP1biom223937_at	I
FOXP1biom240666_at	D
FOXP1biom223936_s_at	I
GPHNbiom215578_at	I
GRIK3biom207454_at	D
GRM7biom217008_s_at	I
GLRA1biom207972_at	I
GNASbiom242975_s_at	I
HEATR5Bbiom1569503_at	I
IL6STbiom204863_s_at	I
LDB2biom206481_s_at	I
LIX1biom230865_at	I
KAT14biom228544_s_at	I
MACROD2biom242468_at	I
MEF2Cbiom239966_at	D
NRXN1biom216096_s_at	D
NRP2biom211844_s_at	D
NRP2biom228699_at	D
NR3C2biom1564236_at	I
PDE4Dbiom236610_at	D
KCNB1biom240869_at	I
KCNB1biom211006_s_at	I
PRKCAbiom215195_at	I
PRKCBbiom228795_at	I
PDP1biom218273_s_at	I
RRBP1biom213973_at	D
RBMS3biom238447_at	I
SRRbiom235677_at	I
SPTBN1biom200672_x_at	D
SPON1biom209436_at	D
SOX6biom1563454_at	D
TIA1biom1554889_at	D
XRCC6biom215308_at	I
ZBTB20biom240216_at	D
ZBTB20biom239955_at	D

B. For Hallucinations (n=213)	
Gene	Direction of Change
DLG1	I
PPP3CB	I
ERBB4	D
RTN4	I
CALM1	I
GNAO1	D
GNAS	D
MAN1A1	I
NDEL1	I
SF3B1	I
NFATC2IP	I
ARHGAP18	I
ESD	I
KCNV1	D
LSM3	I
NCAM1	I
NEK1	I
PDE4B	I
TLK1	I
ZNF24	I
MBP	I
PRKCH	D
THBS1	I
ANXA1	I
ASAP1	D
ATP5C1	I
ATP5C1	I
ATP5C1	I
ATP6V1G1	I
B3GALT5	I
B3GALT5	I
CD9	I
COX7C	I
CRYZL1	I
DDX56	I
DISC1	I
DLG4	D

B. For Hallucinations (n=213)	
Gene	Direction of Change
DST	I
ELMO1	D
ENPP2	D
EVI2A	I
FAT4	D
FGF14	D
FGFR2	D
FN1	D
FN1	D
FNBP1	I
GLS	I
GLS	I
GNAS	D
GNAS	D
GNAS	D
GNAS	D
HDAC8	I
HMGB2	I
HTR2A	D
INPP5D	I
ITPR3	D
KALRN	I
CSRP2BP	I
LAMA4	D
LHPP	D
LPAR1	I
MAB21L1	D
MACROD2	D
MAPK8	I
MLLT10	I
MKL1	I
MYH10	I
NCOA7	I
NOS1	I
ORMDL1	I
PDE7B	I
PPM1B	I
PPP3CB	I
PPP3R1	I

B. For Hallucinations (n=213)	
Gene	Direction of Change
PRICKLE1	I
PRKCB	D
PRKCB	D
PRKCB	I
PRRC2C	D
PSMA4	I
RAPH1	I
SAMHD1	I
SEC31A	I
SERPING1	D
SF3A1	D
SF3B1	I
SLC2A1	D
SPTBN1	D
TBCD	I
THBD	D
THNSL1	I
UNC13C	I
ZBTB20	I
ZEB2	I
ZEB2	I
ZIC1	I
AKAP13	I
AKAP13	I
AKAP13	I
DLGAP1	I
DNAJC1	I
HINT1	I
HINT1	I
HRH1	I
NAT1	I
POLR3GL	I
SEPTIN5	D
UBE2L3	I
ACTR2	I
ACYP2	I
ANGPT1	D
AP1S2	I
ARHGAP18	I

B. For Hallucinations (n=213)	
Gene	Direction of Change
ATXN2	D
CAPZA1	I
CELSR2	I
CTNND1	D
CUL4A	I
DGKZ	D
DHX36	I
DUSP3	D
EPB42	D
ESF1	I
FABP7	I
FBXO7	D
FOXO3	D
FRAS1	I
FRMD4A	I
FRMD4A	D
FYN	D
GRIN2B	I
HACD4	I
ITGAV	I
KCNH2	D
KPNA3	D
LPL	I
LUZP1	I
MAGI1	I
ME1	D
FAM63B	I
NEK1	I
NFKB1	D
NPEPL1	D
NR4A2	I
NR4A2	I
NRG1	D
PAPSS2	I
PDE4DIP	D
PRL	D
PRUNE	D
PSMA6	I
PTP4A2	I

B. For Hallucinations (n=213)	
Gene	Direction of Change
PPP2R4	D
PPP2R4	D
RAPGEF6	D
RORA	I
RORA	D
RTN4	I
SCPEP1	I
SEC14L1	D
SH3PXD2A	I
SLC25A37	D
SLC25A37	D
SLC9A3R1	D
SMPD3	D
SOS1	I
SPTLC1	I
STAT6	D
TATDN3	D
TCF4	I
TECR	I
TGIF1	I
TLE3	D
TMF1	I
TNIK	I
TRPS1	D
UBE3A	D
UQCRR	I
WIZ	D
YTHDC1	I
ZMYND11	I
ZNF24	I
AGO2	D
AGO2	D
ARL16	I
C9orf16	I
CLTA	D
CLTA	D
CPEB4	D
GEM	I
HSPA1B	I

B. For Hallucinations (n=213)	
Gene	Direction of Change
IGF2R	D
IGHM	D
KCNJ4	D
MBP	D
MOBP	I
MOBP	D
MYH9	D
MYH9	I
MYL6	I
PRKCH	I
RAB18	I
RASGRP2	D
RASGRP2	I
RASGRP2	D
S100A8	I
STX7	I
TFRC	D
THBS1	I
TPI1	D
ZFR	I

Table 3 Convergent Functional Evidence (CFE) for Top Biomarkers after Steps 1-4.

A. Delusions B. Hallucinations.

A. Top Biomarkers for Delusions.										
Gene Symbol/ Gene Name	Prob eset	Step 1 Discovery in Blood (Direct ion of Chang e in High Delusions) Method/ Score/ % 6pts	Step 2 External Convergent Functional Genomics (CFG) Evidence For Involvement in Delusion Disorders Score 12pts	Step 3 Validation in Blood AN OVA p-value/ Score 6 pts	Step 4 Best Significant Prediction of High Delusional State ROC AUC/ p-value 4 pts All 2pts Gender 1pts Gender /Dx	Step 4 Significant Prediction of First Year Hosp for Delusions All Gender Best in Individualiz ed Gender /Dx ROC AUC/ p-value 4 pts All 2pts Gender 1pts Gender /Dx	Step 4 Best Significant Predictions of All Future Hosp for Delusions OR/OR p- value Updated 1- tailed p-value and added new data for genes that has now signif p-value 4 pts All 2pts Gender 1pts Gender /Dx	Step 5 Other Psychiatric and Related Disorders Evidence	Step 6 Drugs that Modulate the Biomarker in Same Direction as High Delusions	CFE Polyvi dence Score for Involvement in Delusions (Based on Steps 1- 4)
AUTS 2 Activator Of Transcription And Developmental Regulator AUTS 2	2427 21_a t	(l) DE/4 52.78 %	10	0.49 6/0 Not Step wise	All C:(36/3 15) 0.62/1.1 4E-02 Gender Male C:(32/2 85) 0.62/1.2 8E-02 Gender/ Dx M- PSYCHO SIS C:(29/1 38) 0.67/2.7	All L:(7/158) 0.74/1.58E- 02 Gender/Dx M-BP L:(1/90) 0.99/4.70E- 02	Gender Female C:(12/39) 1.43/1.64E- 02 Gender/Dx F-MDD C:(2/25) 1.74/2.61E- 02 F-SZA C:(6/9) 4.45/2.21E- 02 M-SZ L:(9/34) 2.7/2.69E-02	Alzheimer's Disease Aging Alcohol ASD BP PTSD Cannabis	Lithium Citalopram	24

					4E-03 M-SZA C:(16/76) 0.71/4.85E-03					
NR4 A2 Nucl ear Rece ptor Subf amily 4 Grou p A Mem ber 2	2046 21_s _at	(I) DE/2 33.33 %	11	0.95 5/2 Step wise		All C:(15/254) 0.69/5.87E-03 L:(7/158) 0.79/5.29E-03 Gender Female C:(4/11) 0.96/7.01E-03 L:(2/6) 1/3.20E-02 Male L:(5/152) 0.72/4.66E-02 Gender/Dx M-BP C:(2/135) 0.97/1.09E-02 L:(1/90) 1/4.34E-02 F- PSYCHOSIS C:(4/11) 0.96/7.01E-03 L:(2/6) 1/3.20E-02 F-SZA C:(3/9) 0.94/1.94E-02 L:(2/6) 1/3.20E-02	All C:(80/292) 1.19/2.88E-02 Gender Female C:(12/39) 1.96/8.64E-03 Gender/Dx F-PSYCHOSIS C:(7/11) 3.61/1.24E-02 L:(4/6) 7.9/4.92E-02 F-SZA C:(6/9) 3.42/2.36E-02 L:(4/6) 7.9/4.92E-02	Alcoh ol MDD Alzhei mer's Disea se Aging PTSD		23
CHD 9 Chro	2205 86_a t	(I) AP/6	8	0.01 21/ 0		All L:(7/158)	All C:(80/292)			22

mod omai n Helic ase DNA Bindi ng Prote in 9		98.67 %		Not Step wise		0.69/4.51E- 02 Gender Female L:(2/6) 1/3.20E-02 Gender/Dx M-BP C:(2/135) 0.92/1.98E- 02 L:(1/90) 0.99/4.70E- 02 F- PSYCHOSIS L:(2/6) 1/3.20E-02 F-SZA C:(3/9) 0.89/3.54E- 02 L:(2/6) 1/3.20E-02	1.2/2.40E-02 L:(43/176) 1.48/3.96E- 02 Gender Male C:(68/253) 1.19/3.72E- 02 Gender/Dx F-MDD C:(2/25) 2.19/2.58E- 02 M- PSYCHOSIS C:(39/124) 1.3/2.76E-02 M-SZA C:(22/66) 1.4/2.19E-02 L:(11/38) 2.99/5.69E- 04			
IL6ST Interl eukin 6 Signa l Trans duce r	2048 63_s _at	(I) AP/6 84%	6	0.96 8/2 Step wise		All C:(15/254) 0.63/4.98E- 02 Gender/Dx M-BP C:(2/135) 0.89/3.03E- 02	All C:(80/292) 1.19/4.53E- 02 Gender Male C:(68/253) 1.24/2.47E- 02 Gender/Dx M-BP C:(24/122) 1.62/2.67E- 03 L:(15/78) 1.94/2.05E- 02 M-SZA C:(22/66) 1.33/4.85E- 02	Male MDD BP Later- Life Depr essio n (LLD) Stress	Fluox etine	22

MACROD2 Mononucleoside diphosphate-ribosyltransferase 2	242468_at	(l) AP/6 100%	8	0.108/0 Not Stepwise	Gender Female C:(4/30) 0.77/4.38E-02	Gender/Dx M-BP C:(2/135) 0.86/3.86E-02	All C:(80/292) 1.22/1.66E-02 Gender Male C:(68/253) 1.23/1.90E-02 Gender/Dx F-MDD C:(2/25) 2.2/3.64E-02 M-SZA C:(22/66) 1.32/4.97E-02			21
PDP1 Pyruvate Dehydrogenase Phosphatase Catalytic Subunit 1	218273_subat	(l) DE/4 55.56%	8	0.972/2 Stepwise	All L:(20/194) 0.67/5.28E-03 Gender Female L:(2/18) 0.97/1.75E-02 Male L:(18/176) 0.64/2.72E-02 Gender/Dx F-MDD L:(2/18) 0.97/1.75E-02 M-PSYCHOSIS L:(17/80) 0.68/1.22E-02 M-SZ	Gender Male C:(11/243) 0.68/2.22E-02 Gender/Dx M-BP C:(2/135) 1/7.71E-03	M-SZA C:(22/66) 1.49/2.02E-02	Omega-3 fatty acids Fluoxetine Lithium CPI-613	21	

					L:(8/36) 0.73/2.6 2E-02					
ROR A RAR Relat ed Orph an Rece ptor A	2409 51_a t	(I) DE/4 52.78 %	11	0.10 8/0 Not Step wise	Gender/ Dx M-SZ C:(13/6 2) 0.65/4.9 3E-02	Gender/Dx M-BP C:(2/135) 0.9/2.57E- 02 L:(1/90) 1/4.34E-02	All C:(80/292) 1.3/4.60E-03 Gender Male C:(68/253) 1.35/1.99E- 03 L:(38/154) 1.44/2.73E- 02 M-BP C:(24/122) 2.08/1.21E- 05 L:(15/78) 2.72/8.99E- 04 M-SZA L:(11/38) 2.06/1.28E- 02	Suicid e Aging ASD	Risper idone Fluox etine Lithiu m	21
FOXP 1 Fork head Box P1	2239 36_s _at	(I) AP/4 53.33 %	10	0.5/ 0 Not Step wise	Gender Male C:(11/243) 0.69/1.88E- 02 Gender/Dx M-BP C:(2/135) 0.86/4.02E- 02 L:(1/90) 0.99/4.70E- 02 M- PSYCHOSIS C:(6/101) 0.71/4.37E- 02	Gender Male C:(11/243) 0.69/1.88E- 02 Gender/Dx M-BP C:(2/135) 0.86/4.02E- 02 L:(1/90) 0.99/4.70E- 02 M- PSYCHOSIS C:(6/101) 0.71/4.37E- 02	All C:(80/292) 1.19/4.73E- 02 Gender/Dx M-BP C:(24/122) 1.49/2.38E- 02 L:(15/78) 1.85/2.57E- 02	Alcoh ol	Keta mine Omeg a-3 fatty acids	20
FOXP 1 Fork head	2406 66_a t	(D) DE/4 51.35 %	10	0.19 2/0 Not	All C:(36/3 15)	Gender/Dx M-SZ L:(2/36)	Gender/Dx F-PSYCHOSIS C:(7/11)	MDD Aging	Omeg a-3 fatty acids	20

Box P1				Step wise	0.6/2.25 E-02 L:(20/19 4) 0.62/3.4 6E-02 Gender Female C:(4/30) 0.8/2.93 E-02 Gender/ Dx M- PSYCHO SIS C:(29/1 38) 0.63/1.8 6E-02 M-SZ C:(13/6 2) 0.7/1.57 E-02 L:(8/36) 0.71/3.3 9E-02	0.87/4.21E- 02	2.53/2.52E- 02	Circa dian abnor maliti es Cann abis BP Stress	Cloza pine	
GNA S GNA S Com plex Locu s	2429 75_s _at	(I) DE/4 58.33 %	10	0.21 4/0 Not Step wise		All C:(15/254) 0.66/2.17E- 02 Gender Male C:(11/243) 0.67/2.90E- 02 Gender/Dx M-BP C:(2/135) 0.97/1.14E- 02 L:(1/90) 1/4.34E-02	Gender Male C:(68/253) 1.2/4.69E-02 L:(38/154) 1.32/4.45E- 02 Gender/Dx M-BP C:(24/122) 1.35/4.86E- 02 L:(15/78) 1.77/1.00E- 02 M-SZA C:(22/66) 1.49/1.32E- 02		Risper idone Valpr oic acid Cloza pine	20

XRCC 6 X- Ray Repa ir Cross Com plem entin g 6	2153 08_a t	(I) AP/4 66.67 %	8	0.52 3/2 Step wise	Gender/ Dx M-BP C:(3/14 7) 0.8/3.86 E-02	Gender/Dx M-BP C:(2/135) 1/8.11E-03 L:(1/90) 1/4.34E-02	All C:(80/292) 1.4/3.02E-04 Gender Male C:(68/253) 1.45/9.00E-05 Gender/Dx M-BP C:(24/122) 1.72/1.41E-04 L:(15/78) 2.34/1.40E-02 M- PSYCHOSIS C:(39/124) 1.38/9.06E-03 M-SZA C:(22/66) 1.5/6.88E-03	Social Isolati on Aging Suicid e Comp leters Depr essio n BP		20
ZBTB 20 Zinc Finge r And BTB Dom ain Cont ainin g 20	2399 55_a t	(D) AP/2 35.29 %	10	0.62 2/2 Step wise	All C:(36/3 15) 0.63/6.2 3E-03 L:(20/19 4) 0.65/1.5 7E-02 Gender Female C:(4/30) 0.81/2.5 5E-02 Male C:(32/2 85) 0.61/2.5 5E-02 L:(18/17 6)	Gender/Dx F-SZA C:(6/9) 5.31/4.76E-02		Social Isolati on Aging Suicid e Comp leters Depr essio n BP	Risper idone Fluox etine	20

					0.65/1.7 9E-02 Gender/ Dx F-MDD C:(3/28) 0.8/4.73 E-02 M- PSYCHO SIS C:(29/1 38) 0.64/1.1 7E-02 L:(17/80) 0.65/3.3 7E-02 M-SZ C:(13/6 2) 0.74/3.5 8E-03 L:(8/36) 0.79/7.4 4E-03					
ACSL 4 Acyl- CoA Synt hetas e Long Chai n Famili y Mem ber 4	2240 91_a t	(I) AP/4 50.67 %	10	0.74 4/0 Not Step wise	Gender/ Dx M-SZA C:(16/7 6) 0.64/3.8 3E-02 L:(9/44) 0.71/2.6 5E-02		All C:(80/292) 1.25/1.44E- 02 Gender Male C:(68/253) 1.23/3.30E- 02 Gender/Dx M- PSYCHOSIS C:(39/124) 1.35/1.88E- 02 L:(20/72) 1.93/1.34E- 02 M-SZ	MDD Suicid e Comp leters BP Hunti ngton 's Disea se Parki nson Stress ASD	Lithiu m Omeg a-3 fatty acids	19

							C:(17/58) 1.51/1.27E-02 L:(9/34) 4.14/1.59E-03			
ACTN4 Actin in Alpha 4	241788_x_at	(l) DE/4 50%	6	0.025/4 Step wise		Gender/Dx M-BP L:(1/90) 0.99/4.70E-02	All C:(80/292) 1.2/1.37E-02 Gender Male C:(68/253) 1.21/1.16E-02 Gender/Dx M-BP C:(24/122) 1.26/2.15E-02			19
FOXP1 Fork head Box P1	223937_at	(l) DE/4 54.17 %	10	0.281/0 Not Step wise		Gender/Dx M-BP C:(2/135) 0.98/1.04E-02 L:(1/90) 1/4.34E-02	All C:(80/292) 1.17/4.07E-02 Gender Male C:(68/253) 1.17/4.29E-02 M-BP C:(24/122) 1.39/2.14E-03 L:(15/78) 2.15/2.85E-02 M-SZA L:(11/38) 1.84/2.54E-02	Alcohol	Ketamine Omega-3 fatty acids	19
GLRA1 Glycine Receptor	207972_at	(l) AP/4 56%	10	0.354/0 Not Step wise	Gender/Dx M-PSYCHOSIS C:(29/138)		All C:(80/292) 1.2/3.44E-02	Aging MDD Suicide BP	Clozapine Haloperidol	19

Alpha 1					0.6/4.45E-02 M-SZA C: (16/76) 0.64/4.16E-02 L: (9/44) 0.7/3.45E-02			Anxiety PTSD		
PDE4D Phosphodiesterase 4D	236610_at	(D) AP/4 60.29%	10	0.418/0 Not Step wise	All L: (20/194) 0.62/4.08E-02	M-PSYCH C: (3/7) 1/1.69E-02 M-PSYCHOSIS L: (2/58) 0.95/1.66E-02 M-SZ L: (2/36) 0.96/1.61E-02		MDD Autism Alzheimer's Disease Social Isolation Chronic Stress Longevity	TCA Olanzapine Risperidone Clozapine	19
PDE4DIP Phosphodiesterase 4D Interacting Protein	205872_xat	(I) DE/2 43.06%	9	0.0129/4 Step wise	All C: (36/315) 0.6/2.87E-02 Gender Female C: (4/30) 0.83/1.90E-02 Gender/Dx F-MDD C: (3/28) 0.81/4.04E-02 M-				Clozapine	19

					PSYCHO SIS C: (29/138) 0.61/3.47E-02 M-SZ C: (13/62) 0.7/1.57E-02					
SPO N1 Spon din 1	2094 36_a t	(D) DE/4 54.05 %	10	0.24 8/0 Not Step wise	All L: (20/194) 0.65/1.60E-02 Gender Male L: (18/176) 0.65/2.19E-02 Gender/ Dx M- PSYCHO SIS C: (29/138) 0.61/3.47E-02 L: (17/80) 0.67/1.69E-02 M-SZA L: (9/44) 0.72/2.16E-02		Gender/Dx F-PSYCHOSIS C: (7/11) 2.34/4.39E-02 M-SZ C: (17/58) 1.63/3.84E-02	MDD Suicid e	Cloza pine	19
SRR Serin e Race mase	2356 77_a t	(I) DE/4 56.94 %	10	0.71 2/0 Not Step wise	Gender Female C: (4/30) 0.77/4.38E-02 Gender/ Dx F-MDD	Gender Male L: (5/152) 0.77/1.92E-02 Gender/Dx M-BP C: (2/135)	Gender/Dx F-MDD C: (2/25) 19.59/4.90E-02			19

					C:(3/28) 0.93/7.8 9E-03	0.98/9.40E- 03 M- PSYCHOSIS L:(2/58) 0.85/4.83E- 02 M-SZ L:(2/36) 0.87/4.21E- 02				
TCF4 Trans cripti on Facto r 4	2123 82_a t	(I) DE/2 33.33 %	11	0.27 5/2 Step wise	Gender Female L:(2/18) 0.97/1.7 5E-02 Gender/ Dx F-MDD L:(2/18) 0.97/1.7 5E-02	Gender/Dx M-BP C:(2/135) 0.91/2.26E- 02	Gender/Dx M-BP C:(24/122) 1.41/1.46E- 02	Alzhei mer's Disea se BP Males Suicid e Comp leters Alcoh ol	Dexa meth asone Valpr oate CBD	19
ZBTB 20 Zinc Finge r And BTB Dom ain Cont ainin g 20	2402 16_a t	(D) AP/4 69.12 %	10	0.85 /0 Not Step wise	Gender/ Dx M-SZ L:(8/36) 0.71/3.3 9E-02	Gender Female C:(4/11) 0.82/4.45E- 02 Gender/Dx F- PSYCHOSIS C:(4/11) 0.82/4.45E- 02 M-PSYCH C:(3/7) 1/1.69E-02 M-SZ L:(2/36) 0.9/3.11E- 02	Gender Female C:(12/39) 1.84/2.69E- 02 Gender/Dx F-PSYCHOSIS C:(7/11) 5.31/1.27E- 02 F-SZA C:(6/9) 4.69/2.76E- 02	Social Isolati on Aging Suicid e Comp leters Depr essio n BP	Risper idone Fluox etine	19

B. Top Biomarkers for Hallucinations										
Gene Symbol/ Gene Name	Probesets	Step 1 Discovery in Blood (Direction of Change in High Hallucinations) Method/ Score/ % 6pts	Step 2 External Convergent Functional Genomics (CFG) Evidence For Involvement in Hallucination Disorders Score 12pts	Step 3 Validation in Blood ANOVA p-value / Score 6 pts	Step 4 Best Significant Prediction of High Hallucinations State ROC AUC/ p-value 4 pts All Gender 1pts Gender /Dx	Step 4 Significant Prediction of First Year Hosp for Hallucinations All Gender Best in Individualized Gender /Dx ROC AUC/ p-value 4 pts All 2pts Gender	Step 4 Best Significant Predictions of All Future Hosp for Hallucinations OR/OR p-value Updated 1-tailed p-value and added new data for genes that has now signif p-value 4 pts All 2pts Gender 1pts Gender /Dx	Step 5 Other Psychiatric and Related Disorders Evidence	Step 6 Drugs that Modulate the Biomarker in Opposite Direction as High Hallucinations	CFE Polyevidence Score for Involvement in Hallucinations (Based on Steps 1-4)
DLG1 Discs Large MAG UK Scaffold Protein 1	233869_xapt	(l) AP/4 55.1%	11	3.79E-02/4 Step wise	All L:(29/317) 0.61/2.43E-02 Gender Male L:(29/264) 0.61/2.67E-02 Gender/ Dx Male-SZA C:(21/96) 0.64/2.64E-02 M-BP L:(5/107)	All C:(22/437) 0.61/4.18E-02 Gender Female C:(4/48) 0.84/1.39E-02 Male L:(10/243) 0.68/3.01E-02 Gender/Dx M-PTSD C:(4/39) 0.76/4.34E-02 M-MDD L:(2/34) 0.94/2.02E-02	All C:(108/442) 1.16/1.59E-02 L:(60/270) 1.43/8.39E-03 Gender Male C:(99/392) 1.14/4.07E-02 L:(57/239) 1.4/1.40E-02 Gender/Dx M-MDD C:(9/78) 1.75/1.16E-02 M-PTSD C:(10/39) 2.07/6.48E-	Suicide Stress Pain Anxiety Addiction	Clozapine	31

					0.8/1.11 E-02 M-PSYCHOSIS L:(19/101) 0.64/3.34E-02 M-SZA L:(11/57) 0.68/3.44E-02		04 M-PTSD L:(7/25) 3.93/2.31E-03			
PPP3CB Protein Phosphatase 3 Catalytic Subunit Beta	215586_at	(l) AP/4 53.06%	10	3.51E-02/4 Stepwise	All L:(29/317) 0.6/4.51E-02 Gender Male L:(29/264) 0.6/4.68E-02 M-PSYCHOSIS C:(35/170) 0.62/1.30E-02 M-SZA C:(21/96) 0.69/4.41E-03 M-BP L:(5/107) 0.8/1.24E-02 M-PSYCHOSIS L:(19/101) 0.64/2.5	All C:(22/437) 0.64/1.47E-02 Gender Female C:(4/48) 0.88/6.22E-03 F-BP C:(2/24) 0.89/3.79E-02 M-MDD C:(3/56) 0.82/3.31E-02	All C:(108/442) 1.2/7.59E-03 Gender Female C:(9/50) 2.31/1.10E-02 Male C:(99/392) 1.15/4.26E-02 Gender/Dx M-MDD C:(9/78) 1.59/4.17E-02 M-PTSD C:(10/39) 1.84/1.01E-02	Addictions Aging Alzheimer's Disease Suicide Depression Bipolar Disorder PTSD Pain ASD	Clozapine	30

					8E-02 M-SZA L:(11/57) 0.67/4.1 0E-02					
ENPP 2 Ecto nucle otide Pyrop hosph atase/ Phosp hodie steras e 2	210 839 _s_a _t	(D) AP/6 94%	8	4.13E - 01/0 Not Step wise	All C: (62/5 13) 0.57/3.9 8E-02 L: (29/31 7) 0.63/9.0 0E-03 Gender Male L: (29/26 4) 0.63/1.1 0E-02 Gender/ Dx M- PSYCHO SIS C: (35/1 70) 0.61/2.6 4E-02 M-SZA C: (21/9 6) 0.67/1.0 0E-02 M- PSYCHO SIS L: (19/10 1) 0.68/6.4 7E-03	All C: (22/437) 0.65/8.18E- 03 Gender Female C: (18/389) 0.66/9.16E- 03 Male L: (10/243) 0.68/2.60E- 02 Gender/Dx M- PSYCHOSIS C: (8/153) 0.72/1.76E- 02 M-SZ C: (4/72) 0.76/3.84E- 02 L: (3/43) 0.83/2.83E- 02	All L: (60/270) 1.54/1.51E- 02 Gender Male L: (57/239) 1.61/1.00E- 02 Gender/Dx M-BP L: (13/82) 2.09/3.92E- 02 M- PSYCHOSIS C: (54/147) 1.5/2.80E-02 L: (31/86) 1.67/3.07E- 02 M-SZ C: (37/81) 1.98/2.69E- 02 L: (10/39) 3.89/2.49E- 02	MDD Addict ions Alzhei mer's Diseas e Aging Stress Pain	Fluox etine Valpr oic acid	26
ZEB2 Zinc Finger E-Box Bindin g	239 296 _at	(I) AP/2 40.82 %	10	1.13E - 01/2 Step wise	All C: (62/5 13) 0.58/1.7 7E-02 Gender	All C: (22/437) 0.64/1.28E- 02 Gender Female	All C: (108/442) 1.24/1.50E- 03 Female-BP C: (2/24)	Deme ntia Depre ssion Aging Stress	Cloza pine, Celas trolo mega -3	26

Homeobox 2					Male C: (58/426) 0.58/2.30E-02 Gender/Dx M-PTSD C: (4/39) 0.81/2.08E-02 L: (3/25) 0.88/1.83E-02 C: (35/170) 0.64/6.71E-03 M-SZA C: (21/96) 0.68/6.30E-03	C: (4/48) 0.83/1.52E-02 Gender/Dx M-PTSD C: (4/39) 0.81/2.08E-02 L: (3/25) 0.88/1.83E-02	2.05/4.00E-02 Gender Female C: (9/50) 1.99/6.11E-03 Male C: (99/392) 1.19/1.14E-02 Gender/Dx M-PSYCHOSIS C: (54/147) 1.21/4.76E-02 M-PTSD C: (10/39) 2.51/9.48E-04 L: (7/25) 5.65/3.94E-03	Suicide	fatty acids	
FNBP1 Formin Binding Protein 1	244286_at	(l) DE/450%	8	2.84E-01/2 Step wise	All L: (29/317) 0.59/4.93E-02 Gender Male L: (29/264) 0.6/4.31E-02 M-BP L: (5/107) 0.83/6.85E-03 M-PSYCHOSIS C: (35/170) 0.6/3.96E-02 L: (19/10	Gender Female C: (4/48) 0.81/2.20E-02 Gender/Dx M-PTSD C: (4/39) 0.76/4.78E-02 M-SZ L: (3/43) 0.79/4.76E-02	All C: (108/442) 1.14/3.82E-02 M-PTSD C: (10/39) 3.42/8.70E-05 L: (7/25) 4.4/4.44E-03	Depression Alzheimer's Disease, Aging ASD Pain Suicide Stress Addictions	Olanzapine	24

					1) 0.63/4.0 4E-02 M-SZ L:(8/44) 0.69/4.4 2E-02					
RTN4 Reticu lon 4	243 031 _at	(I) DE/4 64.81 %	11	3.79E - 01/2 Step wise	All L:(29/31 7) 0.64/6.9 6E-03 Gender Male L:(29/26 4) 0.64/6.8 4E-03 Gender/ Dx M-PTSD C:(5/45) 0.76/3.0 2E-02 M-BP L:(5/107) 0.88/1.9 1E-03 M- PSYCHO SIS L:(19/10 1) 0.66/1.6 6E-02 M-SZ L:(8/44) 0.79/5.7 7E-03	Gender Male L:(10/243) 0.68/2.57E- 02 M-PTSD C:(4/39) 0.89/6.21E- 03 L:(3/25) 0.82/3.95E- 02	M-PTSD C:(10/39) 2.82/1.74E- 04 L:(7/25) 3.68/8.69E- 03	Depre ssion Bipola r Disord er Stress Suicid e Deme ntia, Pain Addict ions	Valpr oate Ome ga-3 fatty acids	24
ZNF24 / Zinc Finger Protei n 24	203 247 _s_a _t	(I) DE/4 50%	9	4.73E -01/2 Step wise	All L:(29/31 7) 0.63/1.2 9E-02 Gender Males L:(29/26		All C:(108/442) 1.19/1.63E- 02 Gender Female C:(9/50) 1.69/3.75E-	BP Alzhei mer's Diseas e Aggre ssion Stress		23

					4) 0.63/1.2 0E-02 Gender/ Dx M-SZA C:(21/9 6) 0.64/2.3 3E-02 M-BP L:(5/107) 0.83/6.5 7E-03 M-SZ L:(8/44) 0.78/6.8 6E-03 M- PSYCHO SIS L:(19/10 1) 0.67/9.2 6E-03		02 L:(3/31) 3.11/4.68E- 02 Male C:(99/392) 1.16/4.76E- 02 Gender/Dx M-PTSD L:(7/25) 3.75/1.99E- 02 C:(10/39) 3.09/3.18E- 04			
DST Dysto nin	215 016 _x_a t	(I) AP/4 51.02 %	10	6.44E - 02/0 Not Step wise	All C:(62/5 13) 0.57/3.9 3E-02 L:(29/31 7) 0.65/4.6 1E-03 Gender Male C:(58/4 26) 0.57/4.8 0E-02 L:(29/26 4) 0.65/3.9 8E-03 Gender/ Dx	Gender/Dx M-MDD L:(2/34) 0.94/2.02E- 02	All C:(108/442) 1.2/1.84E-02 L:(60/270) 1.4/5.42E-03 Gender Male C:(99/392) 1.22/1.22E- 02 L:(57/239) 1.41/4.57E- 03 Gender/Dx M-MDD C:(9/78) 2.39/2.28E- 05 L:(6/46) 6.86/2.17E- 04	Depre ssion Alzhei mer's Diseas e ASD Pain Suicid e	Rispe ridon e	23

					M-PSYCHOSIS C: (35/170) 0.61/1.99E-02 L: (19/101) 0.78/6.13E-05 M-SZ L: (8/44) 0.84/1.43E-03 M-SZA C: (21/96) 0.63/3.10E-02 L: (11/57) 0.75/5.42E-03					
FAT4 FAT Atypical Cadherin 4	219 427 _at	(D) AP/6 80%	8	8.09E - 01/0 Not Step wise	Gender/Dx M-SZ L: (8/44) 0.69/4.71E-02	All C: (22/437) 0.61/3.54E-02	All C: (108/442) 1.21/3.11E-02 L: (60/270) 1.91/2.18E-04 Gender Male C: (99/392) 1.2/4.38E-02 L: (57/239) 1.77/1.62E-03 M-PTSD C: (10/39) 2.02/3.31E-02 L: (7/25) 3.23/3.69E-02	Aging Suicide Stress	Citalopram	23
ITGAV Integrin	236 251 _at	(I) DE/4	7	3.94E - 01/2	All L: (29/317)	Gender Female C: (4/48)	All C: (108/442)	ASD Alzheimer's	Valproate	23

Subunit Alpha V		51.85 %		Step wise	0.6/3.33 E-02 Gender Male L:(29/264) 0.6/3.31 E-02 Gender/Dx M-PTSD C:(5/45) 0.74/4.15 E-02 M-BP L:(5/107) 0.79/1.50 E-02 M-PSYCHOSIS L:(19/101) 0.65/1.81 E-02 M-SZ L:(8/44) 0.7/4.14 E-02	0.79/2.86 E-02 Gender/Dx M-PTSD C:(4/39) 0.86/1.03 E-02	1.14/4.95 E-02 L:(60/270) 1.28/4.90 E-02 Gender/Dx M-PTSD C:(10/39) 3.08/4.74 E-04 M-MDD L:(6/46) 1.49/4.76 E-02	Disease Depression PTSD Addictions Aging		
ORMDL1 ORMDL Sphingolipid Biosynthesis Regulator 1	228801 -at	(l) DE/4 55.56 %	8	5.70 E-01/2 Step wise	All L:(29/317) 0.63/1.02 E-02 Gender Male L:(29/264) 0.63/1.07 E-02 Gender/Dx M-BP L:(5/107) 0.81/9.47 E-03	All C:(22/437) 0.61/4.16 E-02 Gender Female C:(4/48) 0.82/1.67 E-02	Gender/Dx M-PTSD C:(10/39) 2.17/1.59 E-03 L:(7/25) 2.72/2.96 E-02	Stress Depression Aging PTSD Pain		23

					M-PSYCHOSIS C: (35/170) 0.6/3.15E-02 L: (19/101) 0.71/1.96E-03 M-SZ L: (8/44) 0.84/1.30E-03				
ACYP2 / Acyl phosphatas e 2	217536_x_ant	(I) DE/681.48%	5	8.46E-02/2 Step wise	Gender/Dx M-PTSD C: (5/45) 0.77/2.56E-02 M-PSYCHOSIS C: (35/170) 0.61/2.59E-02 M-SZA C: (21/96) 0.63/3.04E-02 M-BP L: (5/107) 0.84/5.80E-03	All C: (22/437) 0.65/9.03E-03 Gender Females C: (4/48) 0.81/2.20E-02 Males L: (10/243) 0.65/4.86E-02 Gender/Dx M-PTSD C: (4/39) 0.88/7.08E-03 M-SZ L: (3/43) 0.89/1.25E-02	All C: (108/442) 1.14/3.10E-02 All L: (60/270) 1.33/2.08E-02 Gender Male L: (57/239) 1.29/3.81E-02 Gender/Dx M-PTSD C: (10/39) 1.77/5.50E-03 L: (7/25) 2.68/6.06E-03 M-PSYCHOSIS L: (31/86) 1.49/2.26E-02 M-SZA L: (21/47) 1.49/3.69E-02	Alzheimer's Disease Progression Aging Stress	22

CALM 1/ Calmo dulin 1	244 869 _at	(I) DE/4 77.78 %	8	3.40E -02/4 Step wise	Gender/ Dx M-PTSD C:(5/45) 0.83/8.5 7E-03 M-SZA C:(21/9 6) 0.63/3.2 3E-02 M-SZ L:(8/44) 0.76/1.2 2E-02 M- PSYCHO SIS L:(19/10 1) 0.66/1.7 0E-02	All C:(22/437) 0.63/2.18E- 02 Gender Females C:(4/48) 0.78/3.38E- 02 F-MDD C:(1/21) 1/4.93E-02 Males L:(10/243) 0.67/3.62E- 02 Gender/Dx M-PTSD C:(4/39) 0.89/5.45E- 03 M-MDD L:(2/34) 0.86/4.61E- 02	All C:(108/442) 1.16/1.66E- 02 L:(60/270) 1.27/3.46E- 02 Gender Male C:(99/392) 1.15/2.71E- 02 Gender/Dx F-MDD C:(4/23) 2.19/4.72E- 02 M-MDD C:(9/78) 1.62/6.95E- 03 M-PTSD C:(10/39) 1.58/1.14E- 02 M- PSYCHOSIS L:(31/86) 1.34/4.05E- 02 All	Suicid e Compl eters Male MDD Alzhei mer's Diseas e Childh ood Traum a ASD Parkin son Aging Alcohol		22
CUL4 A/ Cullin 4A	240 971 _x_a t	(I) DE/2 44.44 %	9	5.46E -02/2 Step wise	Gender/ Dx M- PSYCHO SIS L:(19/10 1) 0.68/8.6 3E-03 M-SZA L:(11/57) 0.67/3.9 3E-02		All C:(108/442) 1.14/3.76E- 02 L:(60/270) 1.37/9.52E- 03 Gender Male L:(57/239) 1.36/1.39E- 02 M-MDD C:(9/78) 1.54/1.92E- 02 M-PTSD	Alzhei mer's Diseas e Aging Stress	Cloza pine Ome ga-3 fatty acids	22

							C:(10/39) 2.44/4.88E-04 L:(7/25) 5.03/9.29E-04			
DHX3 6/ DEAH- Box Helica se 36	155 903 9_at	(I) AP/4 57.14 %	5	3.55E -02/4 Step wise	All L:(29/31 7) 0.62/1.4 4E-02 Gender Males L:(29/26 4) 0.63/8.9 5E-03 Gender/ Dx M- PSYCHO SIS C:(35/1 70) 0.6/3.89 E-02 L:(19/10 1) 0.67/1.2 2E-02 M-PTSD C:(5/45) 0.73/4.8 3E-02 M-SZA C:(21/9 6) 0.64/2.9 2E-02 L:(11/57) 0.68/3.6 0E-02	All C:(22/437) 0.61/4.30E-02	Gender/Dx M-PTSD C:(10/39) 2.66/1.46E-03 L:(7/25) 2.45/2.61E-02	MDD	Rispe ridon e	22
GNAS / GNAS Compl	211 858 _x_a t	(D) DE/4 53.85 %	10	2.52E -01/0 Not	All L:(29/31 7) 0.6/4.31	Gender Females C:(4/48)	Gender Female C:(9/50)	BP Alzhei mer's Diseas	Valpr oic acid Antip	22

ex Locus				Step wise	E-02 Gender Males L:(29/26 4) 0.6/4.55 E-02	0.82/1.84E- 02 Gender/Dx M-MDD L:(2/34) 0.86/4.61E- 02	2.54/6.84E- 03 Gender/Dx F-MDD C:(4/23) 3.93/1.36E- 02 M-MDD C:(9/78) 2.24/1.98E- 02 L:(6/46) 4.23/5.03E- 03	e Suicid e Childh ood Traum a Aging Cocai ne Anxiet y Alcoh ol	sych otics Cloza pine	
HDAC 8/ Histon e Deace tylase 8	223 908 _at	(l) DE/6 81.48 %	6	1.11E -01/2 Step wise	All L:(29/31 7) 0.63/1.1 9E-02 Gender Females C:(4/87) 0.79/2.7 1E-02 Males L:(29/26 4) 0.63/1.2 1E-02 Gender/ Dx M-SZ L:(8/44) 0.89/2.9 2E-04 M- PSYCHO SIS L:(19/10 1) 0.72/1.2 5E-03		All C:(108/442) 1.16/4.46E- 02 Gender Male C:(99/392) 1.17/3.36E- 02 Gender/Dx M-MDD C:(9/78) 2.38/1.17E- 05 L:(6/46) 3.58/3.55E- 04 M-SZ C:(17/66) 1.42/1.65E- 02	Panic Disord er MDD Stress	Lithiu m	22

MAB2 1L1/ Mab- 21 Like 1	206 163 _at	(D) DE/2 36.92 %	10	3.06E -01/2 Step wise	All L:(29/31 7) 0.62/1.6 2E-02 Gender Males C:(58/4 26) 0.57/4.8 0E-02 L:(29/26 4) 0.62/1.6 4E-02 M- PSYCHO SIS C:(35/1 70) 0.64/5.9 6E-03 L:(19/10 1) 0.68/8.2 3E-03 M-SZ C:(14/7 4) 0.68/1.7 4E-02 L:(8/44) 0.77/8.1 1E-03		All C:(108/442) 1.24/2.32E- 02 Gender Male C:(99/392) 1.3/1.08E-02 M-MDD C:(9/78) 2.02/3.15E- 02 M-BP C:(26/128) 1.6/1.72E-02	MDD Stress	Escit alopr am (SSRI) Cloza pine	22
ZBTB2 0/ Zinc Finger And BTB Domai n Contai ning 20	235 308 _at	(I) DE/4 51.85 %	6	3.90E -02/4 Step wise	All L:(29/31 7) 0.61/2.3 7E-02 Gender Males L:(29/26 4) 0.61/2.3 7E-02 Gender/Dx M-		All C:(108/442) 1.17/3.08E- 02 Gender Male C:(99/392) 1.16/4.50E- 02 Gender/Dx M-PTSD C:(10/39) 2.19/3.65E- 03	Alcoh ol Alzhei mer's Diseas e Suicid e Parkin son Hunti ngton' s	Rispe ridon e Fluox etine	22

					PSYCHO SIS C: (35/170) 0.61/2.05E-02 L: (19/101) 0.69/4.81E-03 M-SZ C: (21/96) 0.65/1.93E-02 L: (8/44) 0.75/1.42E-02		L: (7/25) 3.57/5.36E-03	Disease Depression		
ZNF24 / Zinc Finger Protein 24	242 210 _at	(l) DE/4 53.7%	9	9.54E-01/0 Not Step wise	All L: (29/317) 0.62/1.52E-02 Gender Males L: (29/264) 0.62/1.45E-02 Gender/Dx Dx M-BP L: (5/107) 0.87/2.76E-03 M- PSYCHO SIS L: (19/101) 0.63/3.53E-02 M-SZ L: (8/44) 0.69/4.71E-02	Gender/Dx M-PTSD C: (4/39) 0.8/2.59E-02	All L: (60/270) 1.47/8.69E-03 Gender Male L: (57/239) 1.42/1.87E-02 Gender/Dx M-MDD L: (6/46) 4.42/1.39E-03 M-PTSD C: (10/39) 2.36/7.01E-04 L: (7/25) 2.74/2.82E-02 M-SZA L: (21/47) 1.63/3.43E-02	BP Alzheimer's Disease Aggression Stress		22

CTNN D1 Catenin Delta 1	155 794 4_s_at	(D) DE/2 33.85	9	8.94E - 02/2 Step wise	Gender/ Dx M-PTSD C:(5/45) 0.8/1.66 E-02 M-BP L:(5/107) 0.73/4.2 1E-02	All C:(22/437) 0.61/4.76E- 02 Gender/Dx M-MDD C:(3/56) 0.81/3.58E- 02 M-PTSD C:(4/39) 0.8/2.59E- 02	All C:(108/442) 1.29/4.38E- 03 Gender Female L:(5/33) 5.89/7.40E- 03 Male C:(99/392) 1.28/6.59E- 03 Gender/Dx M-BP C:(26/128) 1.38/3.83E- 02 M-MDD C:(9/78) 2.48/8.37E- 03 M-PTSD C:(10/39) 2.35/1.04E- 02	Stress Addict ion Suicid e Alzhei mer	Cloza pine	22
-------------------------------------	----------------------	----------------------	---	------------------------------------	---	--	--	---	---------------	----

[0094] Referring now to Figure 2. Top Predictive Biomarkers for Different Demographic and Disease Groups. Delusions A-C. A. State Severity. B. Short-Term Risk. C. Long-term Risk. Hallucinations D-F. D. State Severity. E. Short-Term Risk. F. Long-term Risk. “*”- nominally significant. “***”- significant after Bonferroni correction for number of biomarkers tested. For Delusions, n=70 probesets, 64 genes. For Hallucinations, n=213 probesets, 178 genes.

[0095] The number of biomarkers with nominally significant AUCs or Odds Ratios are depicted in the tables underneath the graphs. Bar graph shows best predictive biomarkers in each group. * nominally significant p<0.05. ** survived Bonferroni correction for the number of candidate biomarkers tested. Table underneath the figures displays the actual number of biomarkers for each group whose ROC AUC p-values or Cox Odds Ratio p-values are at least nominally significant. Cross-sectional is based on levels at one visit. Longitudinal is based on

levels at multiple visits (integrates levels at most recent visit, maximum levels, slope into most recent visit, and maximum slope). Dividing lines represent the cutoffs for a test performing at chance levels (white), and at the same level as the best biomarkers for all subjects in cross-sectional (gray) and longitudinal (black) based predictions. All depicted biomarkers perform better than chance. Biomarkers performed better when personalized by gender and diagnosis, particularly in females.

[0096] Referring now to Figure 4. Example of Possible Report to Clinicians. Using a panel of the top predictive biomarkers after Step 4. A. Delusions, B. Hallucinations. Each panel contains 18 biomarkers: the top 3 best biomarkers for Current Severity (State) in All, Gender, Gender and Diagnosis; Short-Term Risk (1st Tear) in All, Gender, Gender and Diagnosis; Long-Term Risk (All future) in All, Gender, Gender and Diagnosis. For the participant and visit for which the report is generated, the raw expression values of the biomarkers were Z-scored by gender and diagnosis with the 793 other participants and visit datasets in our dataset.

[0097] The Z-scored expression value of each biomarker in our participant tested was compared to the average value for the biomarker in the 793 dataset, from the severely psychotic group (PANSS item ≥ 4 for state, or having future hospitalizations with psychosis for trait) and from the non-psychotic group (PANSS item = 1 for state, or not having future hospitalizations with psychosis for trait.). For increased in expression biomarkers, the comparison resulted in scores of 1 if above the first average, 0 if below the second average, and 0.5 if it was in between. The reverse was done for decreased biomarkers. The comparison groups in the 793 cohort were all, the same gender, and the same gender and diagnosis corresponding to the participant for which the report is generated.

[0098] The “digitized” biomarkers were then added into a polygenic risk score, and a percentile calculated. If above the 75%, the patient is deemed high severity/ risk (red), if between 75 and 50% is intermediate high, if between 50 and 25% intermediate low, and if below 25% low.

[0099] The stars depict each biomarker, and are filled corresponding to the score, and colored corresponding to the level of risk. The “digitized” biomarkers were also used for matching with existing psychiatric medications. Biomarkers were matched based on our CFG literature databases with existing psychiatric medications that had effects on gene expression

opposite to psychosis. Each medication matched to a biomarker got a score of one (1) that was then multiplied with the biomarker score of 1, 0.5 or 0. The scores for the medications were added, a percentile calculated, and medications prioritized by this percentile for consideration and use by the clinician.

A. Delusions.

[0100] The participant had a delusions severity score of 63% for current state, 44% for short-term risk, and 11% for long-term risk. This participant's clinical measures were fairly concordant with the blood test results (high delusions scores, poor grooming).

B. Hallucinations.

[0101] The participant had a delusions severity score of 11% for current state, 6% for short-term risk, and 33% for long-term risk. This participant's clinical measures were discordant among themselves and with the blood test results (self-reported high-hallucination scores, but above average grooming), pointing out to the need for objective measures.

Table 4. Repurposed Drugs. A-C Delusions. D-F Hallucinations.

A. Delusions - All Patients			
CMAP Biomarker Panel Used		NIH LINCS Biomarker Panel Used	
Decreased Biomarkers: SPON1 Increased Biomarkers: PDP1, NR4A2, IL6ST, XRCC6, CHD9		Decreased Biomarkers: ZBTB20, FOXP1, SPON1, NRP2 Increased Biomarkers: AUTS2, PDP1, NR4A2, GNAS, IL6ST, CHD9, XRCC6, RORA, ACTN4, ACSL4	
Drug	Score	Drug	Score
adenosine phosphate	-1	528116.cdx	0.3077
N-acetyl-L-leucine	-0.974	Cyclopiazonic Acid	0.2308
eldeline	-0.96	SB 218078	0.2308
pempidine	-0.958	BRD-A36630025	0.2308
verteporfin	-0.945	QUINACRINE HYDROCHLORIDE	0.2308
C-75	-0.91	GF-109203X	0.2308
oxprenolol	-0.909	BRD-A36630025	0.2308
Prestwick-675	-0.901	N9- isopropylolomoucine	0.2308
meglumine	-0.898	BMS-536924	0.2308
guanethidine	-0.891	BRD-K76951091	0.2308
pancuronium bromide	-0.889	BRD-K26304855	0.2308
karakoline	-0.886	trichostatin A	0.2308
15(S)-15-methylprostaglandin E2	-0.885	ALW-II-38-3	0.2308
hexylcaine	-0.878	Mitoxantrone	0.2308
dicoumarol	-0.878	HG-6-64-01	0.2308
apramycin	-0.878	Alvocidib	0.2308
mephenytoin	-0.877	SB-216763	0.2308
estriol	-0.876	Caffeic acid phenethyl ester	0.1538
sodium phenylbutyrate	-0.868	FR 139317	0.1538
dienestrol	-0.867	Syk Inhibitor	0.1538

B. Delusions – Males			
CMAP Biomarker Panel Used		NIH LINCS Biomarker Panel Used	
Decreased Biomarkers: ZBTB20, FOXP1, ZBTB20, SPON1, NRP2 Increased Biomarkers: AUTS2, PDP1, FOXP1, PDP1, GNAS, SRR, NR4A2, XRCC6, RORA, ACTN4		Decreased Biomarkers: ZBTB20, FOXP1, SPON1, NRP2 Increased Biomarkers: AUTS2, PDP1, FOXP1, GNAS, SRR, NR4A2, XRCC6, RORA, ACTN4	
Drug	Score	Drug	Score
flunisolide	-1	528116.cdx	0.3636
apramycin	-0.995	SB 218078	0.2727
adenosine phosphate	-0.974	QUINACRINE HYDROCHLORIDE	0.2727
guanethidine	-0.958	N9-isopropylolomoucine	0.2727
15(S)-15-methylprostaglandin E2	-0.952	ALW-II-38-3	0.2727
meteneprost	-0.948	Mitoxantrone	0.2727
methyldopate	-0.941	Mitoxantrone	0.2727
hydralazine	-0.939	HG-6-64-01	0.2727
rotenone	-0.938	Alvocidib	0.2727
phthalylsulfathiazole	-0.936	SB-216763	0.2727
N-acetyl-L-leucine	-0.933	Syk Inhibitor	0.1818
eldeline	-0.92	Cyclopiazonic Acid	0.1818
tocainide	-0.919	GW 441756	0.1818
laudanoline	-0.918	LY 225910	0.1818
pempidine	-0.918	AG 82	0.1818
7-aminocephalosporanic acid	-0.915	DOXORUBICIN	0.1818
sulfachlorpyridazine	-0.909	MITOMYCIN C	0.1818
finasteride	-0.907	TERFENADINE	0.1818
verteporfin	-0.905	DOXORUBICIN	0.1818
pempidine	-0.904	Syk Inhibitor	0.1538

C. Delusions – Females			
CMAP Biomarker Panel Used		NIH LINCS Biomarker Panel Used	
Decreased Biomarkers: FGFR1, DISC1, FGFR2, SPTBN1, INSR, GRIK3, ZBTB20 Increased Biomarkers: PDE4DIP, PDP1, TCF4, NR4A2, CHD9, CLCN3, AUTS2, LDB2, NR4A2		Decreased Biomarkers: FGFR1, DISC1, FGFR2, SPTBN1, INSR, GRIK3, ZBTB20 Increased Biomarkers: PDE4DIP, PDP1, TCF4, NR4A2, CHD9, CLCN3, AUTS2, LDB2	
Drug	Score	Drug	Score
erastin	-1	I-BET151	0.2667
harpagoside	-0.976	NYLIDRIN HYDROCHLORIDE	0.2
metacycline	-0.972	AMG 9810	0.2
amiodarone	-0.969	DOXORUBICIN	0.2
furaltadone	-0.958	MITOMYCIN C	0.2
metformin	-0.951	FLUDROCORTISONE ACETATE	0.2
timolol	-0.935	Purvalanol A	0.2
repaglinide	-0.928	TENIPOSIDE	0.2
sulfafurazole	-0.92	Geldanamycin	0.2
PNU-0230031	-0.908	Importazole	0.2
probenecid	-0.907	BRD-A36630025	0.2
furosemide	-0.904	YM-155	0.2
fluphenazine	-0.896	Auranofin	0.2
myricetin	-0.893	7643453	0.2
sulfacetamide	-0.892	G-221	0.2
lomustine	-0.891	BRD-A49680073	0.2
BCB000039	-0.889	BRD-K08547377	0.2
harmalol	-0.885	Cladribine	0.2
cimetidine	-0.88	NVP-AUY922	0.2
acenocoumarol	-0.877	TWS-119	0.2

D. Hallucinations - All Patients			
CMAP Biomarker Panel Used		NIH LINCS Biomarker Panel Used	
Decreased Biomarkers: PRL, SERPING1, ENPP2, KCNV1, FAT4 Increased Biomarkers: NCAM1, B3GALT5, PTP4A2, ACYP2, DST		Decreased Biomarkers: PRL, SERPING1, ENPP2, LAMA4, KCNV1, CTNND1, FAT4 Increased Biomarkers: PRICKLE1, NCAM1, B3GALT5, ARHGAP18, PTP4A2, ACYP2, RTN4, CUL4A, ZEB2, DST, DLG1	
Drug	Score	Drug	Score
clioquinol	-1	BRD-K71489689	0.25
pirinixic acid	-0.949	trichostatin A	0.25
moxisylyte	-0.931	A443654	0.25
Prestwick-685	-0.93	AG 825	0.1875
exemestane	-0.926	Proscillaridin A	0.1875
azacitidine	-0.914	Ala-Ala-Phe-CMK	0.1875
C-75	-0.913	FLUOCINOLONE ACETONIDE	0.1875
estradiol	-0.894	manumycin A	0.1875
tetraethylenepentamine	-0.893	curcumin	0.1875
sparteine	-0.887	BRD-K68548958	0.1875
guanethidine	-0.883	CHR 2797	0.1875
idoxuridine	-0.883	Tyrphostin AG 1478	0.1875
gliclazide	-0.878	wortmannin	0.1875
nitrendipine	-0.877	HY-50878	0.1875
N-acetyl-L-aspartic acid	-0.872	598226	0.1875
sulfanilamide	-0.871	S1003	0.1875
doxazosin	-0.87	BRD-A52530684	0.1875
pimozide	-0.865	CGP-60474	0.1875
proscillaridin	-0.864	buparlisib	0.1875
oxetacaine	-0.86	AS-601245	0.1875

E. Hallucinations – Males			
CMAP Biomarker Panel Used		NIH LINCS Biomarker Panel Used	
Decreased Biomarkers: PRL, SERPING1, ENPP2, KCNV1, MAB21L1, FAT4 Increased Biomarkers: ACYP2, DST		Decreased Biomarkers: PRL, SERPING1, ENPP2, KCNV1, MAB21L1, CTNND1, FAT4 Increased Biomarkers: SH3PXD2A, ZEB2, PRICKLE1, ARHGAP18, ACYP2, RTN4, DST	
Drug	Score	Drug	Score
digoxigenin	-1	trichostatin A	0.3333
doxazosin	-0.984	manumycin A	0.25
meptazinol	-0.972	NCGC00189555-02	0.25
promethazine	-0.963	buparlisib	0.25
cefixime	-0.942	linifanib	0.25
velnacrine	-0.929	AZD-7762	0.25
cetirizine	-0.927	dinaciclib	0.25
eldeline	-0.924	Piretanide	0.1667
atropine oxide	-0.922	KN-62	0.1667
clioquinol	-0.92	Fluticasone propionate	0.1667
nicotinic acid	-0.916	JAK3 Inhibitor VI	0.1667
clioquinol	-0.915	SARMENTOGENIN	0.1667
galantamine	-0.911	Digoxin	0.1667
rolitetracycline	-0.909	MEGESTROL ACETATE	0.1667
betahistine	-0.903	Oxymetazoline hydrochloride	0.1667
sulconazole	-0.9	U-0126	0.1667
monocrotaline	-0.899	Tracazolate hydrochloride	0.1667
lanatoside C	-0.895	FLUFENAMIC ACID	0.1667
Prestwick-1084	-0.89	FENOFIBRATE	0.1667
naftidrofuryl	-0.887	U 99194 maleate	0.1667

F. Hallucinations - Females	
CMAP Biomarker Panel Used	
Decreased Biomarkers: GNAS Increased Biomarkers: CELSR2, B3GALT5, PPP3CB, THNSL1, NR4A2	
Drug	Score
proglumide	-1
quinethazone	-0.951
esculin	-0.938
MG-262	-0.935
GW-8510	-0.932
haloperidol	-0.931
guanethidine	-0.93
deferoxamine	-0.925
citolone	-0.919
meteneprost	-0.913
amylocaine	-0.91
CP-944629	-0.907
clemizole	-0.899
IC-86621	-0.899
nortriptyline	-0.89
CP-944629	-0.888
tanespimycin	-0.885
Prestwick-674	-0.883
0317956-0000	-0.883
pioglitazone	-0.879

REFERENCES

1. Kurian SM, Le-Niculescu H, Patel SD, et al. Identification of blood biomarkers for psychosis using convergent functional genomics. *Mol Psychiatry* 2011;16:37-58.
2. Le-Niculescu H, Levey DF, Ayalew M, et al. Discovery and validation of blood biomarkers for suicidality. *Mol Psychiatry* 2013;18:1249-64.
3. Niculescu AB, Levey DF, Phalen PL, et al. Understanding and predicting suicidality using a combined genomic and clinical risk assessment approach. *Mol Psychiatry* 2015;20:1266-85.
4. Levey DF, Niculescu EM, Le-Niculescu H, et al. Towards understanding and predicting suicidality in women: biomarkers and clinical risk assessment. *Mol Psychiatry* 2016;21:768-85.
5. Niculescu AB, Le-Niculescu H, Levey DF, et al. Precision medicine for suicidality: from universality to subtypes and personalization. *Mol Psychiatry* 2017;22:1250-73.

CLAIMS

1. A method for assessing and treating schizophrenia and other psychotic disorders in general, in particular delusions and risk of developing delusions in an individual, comprising the steps of:

(a) obtaining a biological sample from an individual and quantifying the amounts of RNA biomarkers in the biological sample, to create a panel of RNA biomarkers,

(b) quantifying the amounts of the RNA biomarkers in the panel in a clinically relevant population to generate a reference expression level for the RNA biomarkers in a panel of RNA biomarkers;

(c) comparing the amounts of the biomarkers in the biological sample from the individual with the amounts of the RNA biomarkers present in the reference standard to generate a score for each biomarker;

wherein the biomarkers in the a first panel (a) comprise one or more of the following RNA biomarkers: Activator Of Transcription and Developmental Regulator 2 (AUTS2), Pyruvate Dehydrogenase Phosphatase Catalytic Subunit 1 (PDP1), Nuclear Receptor Subfamily 4 Group A Member 2 (NR4A2), GNAS Complex Locus (GNAS), Interleukin 6 Signal Transduce (IL6ST), Chromodomain Helicase DNA Binding Protein 9 (CHD9), X-Ray Repair Cross Complementing 6 (XRCC6), RAR Related Orphan Receptor A (RORA), Actinin Alpha 4 (ACTN4), and Acyl-CoA Synthetase Long Chain Family Member 4 (ACSL4), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting delusions or an increased risk for developing delusions; and

biomarkers in a second panel (b) comprise one or more of the following RNA biomarkers: Zinc Finger And BTB Domain Containing 20 (ZBTB20), Forkhead Box P1 (FOXP1), Spondin 1 (SPON1), and (NRP2), wherein the expression level of the RNA biomarker(s) in the sample is decreased relative to a reference expression level of the RNA biomarkers in the panel, denoting delusions or an increased risk for developing delusions;

- (d) generating a score for the panel of RNA biomarkers, based on the scores of the biomarker(s) in the panel;
- (e) determining a reference score for the panel in a clinically normal relevant population;
- (f) identifying a difference between the score of the panel of biomarker(s) in the sample and the reference score of the panel of biomarker(s);
- (g) identifying the individual as having delusions or of having an elevated risk for developing delusions, based on the difference between the biomarker panel score of the individual relative to the biomarker panel score of the reference;
- (h) treating the individual identified as having delusions or an elevated risk of delusions with at least one treatment selected from the group consisting of: a treatment based on clinical practice guidelines, administering a therapeutically effective amount of at least one therapeutic drug wherein the mode of treatment is on the specific biomarkers scores indicating that individual will benefit from a particular therapy.
2. The method of claim 1, wherein the biomarkers are quantified in samples taken on two or more occasions from the individual.
3. The method of claim 1, wherein each biomarker is assigned a weighted coefficient based on each biomarkers importance in in assessing and predicting delusions risk; and the biomarker panel score is based on the weighted coefficients of each of the biomarkers.
4. The method of claim 1, wherein the biological sample is at least sample from the individual selected from the group consisting of: tissue, a fluid such as cerebrospinal fluid, whole blood, blood serum, plasma, saliva, or other bodily fluid, or an extract or purification therefrom, or a dilution thereof.
5. The method of claim 1, wherein the therapeutic is at least drug selected from the group consisting of: adenosine phosphate, N-acetyl-L-leucine, eldeline, pempidine, verteporfin, C-75, oxprenolol, Prestwick-675, meglumine, guanethidine, pancuronium bromide, karakoline, 15(S)-15-methylprostaglandin E2, hexylcaine, dicoumarol, apramycin, mephentyoin, estriol, 528116.cdx, Cyclopiazonic Acid, SB 218078, BRD-A36630025, Quinacrine hydrochloride, GF-109203X, BRD-A36630025, N9-isopropylolomoucine, BMS-536924, BRD-K76951091, BRD-

K26304855, trichostatin A, ALW-II-38-3, mitoxantrone, HG-6-64-0, alvocidib, SB-216763, and caffeic acid phenethyl ester.

6. The method of claim 1, wherein when the individual is male, and the biomarkers in a first panel (a) comprise one or more biomarkers selected from the group consisting of: Activator Of Transcription And Developmental Regulator (AUTS2), Pyruvate Dehydrogenase Phosphatase Catalytic Subunit 1 (PDP1), Forkhead Box P1 (FOXP1), GNAS Complex Locus (GNAS), Serine Racemase (SRR), Nuclear Receptor Subfamily 4 Group A Member 2 (NR4A2), X-Ray Repair Cross Complementing 6 (XRCC6), RAR Related Orphan Receptor A (RORA), and Actinin Alpha 4 (ACTN4), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting increased delusions, or

the biomarkers in a second panel (b) comprising one or more biomarkers selected from the group consisting of: Zinc Finger And BTB Domain Containing 20 (ZBTB20), Forkhead Box P1 (FOXP1), Spondin 1 (SPON1), NRP2, wherein the expression level of the biomarker(s) in the sample is decreased relative to a reference expression level, denoting increased delusions.

7. The method of claim 6, wherein the at least one therapeutic drug is one or more drugs selected from the group consisting of: flunisolide, apramycin, adenosine phosphate, guanethidine, 15(S)-15-methylprostaglandin E2, meteneprost, methyl dopate, hydralazine, rotenone, phthalylsulfathiazole, N-acetyl-L-leucine, eldeline, tocainide, laudanosine, pempidine, 7-aminocephalosporanic acid, Sulfachlorpyridazine, finasteride, 528116.cdx, SB 218078, Quinacrine hydrochloride, N9-isopropylolomoucine, ALW-II-38-3, mitoxantrone, HG-6-64-01, Alvocidib, SB-216763, Syk Inhibitor, Cyclopiazonic Acid, GW 441756, LY 225910, AG 82, doxorubicin, mitomycin, and terfenadine.

8. The method of claim 1, wherein when the individual is female, and the biomarkers in the panel comprise one or biomarkers in a first panel (a) comprise one or more of the biomarkers selected from the group consisting of: Phosphodiesterase 4D Interacting Protein (PDE4DIP), Pyruvate Dehydrogenase Phosphatase Catalytic Subunit 1 (PDP1), Transcription Factor 4 (TCF4), Nuclear Receptor Subfamily 4 Group A Member 2 (NR4A2), Chromodomain Helicase DNA Binding Protein 9 (CHD9), (CLCN3), Activator Of Transcription And Developmental Regulator (AUTS2), and (LDB2), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting increased delusions; and

the biomarkers in a second panel (b) comprise one or more biomarkers selected from the group consisting of (FGFR1), (DISC1), (FGFR2), (SPTBN1), (INSR), (GRIK3), Zinc Finger, and BTB Domain Containing 20 (ZBTB20), wherein the expression level of the biomarker(s) in the sample is decreased relative to a reference expression level, denoting increased delusions.

9. The method of claim 8, wherein the at least one therapeutic drug is at least drug selected from the group consisting of: erastin, harpagoside, metacycline, amiodarone, furaltadone, metformin, timolol, Repaglinide, sulfafurazole, PNU-0230031, Probenecid, furosemide, fluphenazine, myricetin, sulfacetamide, lomustine, BCB000039, Harmalol, I-BET151, Nylidrin hydrochloride, AMG 9810, Doxorubicin, Mitomycin C, Fludrocortisone acetate, Purvalanol A, Teniposide, Geldanamycin, Importazole, BRD-A36630025, YM-155, Auranofin, 7643453, G-221, BRD-A49680073, BRD-K08547377, and Cladribine.

10. A method of assessing and treating schizophrenia and other psychotic disorders in general, and delusions in particular, in an individual, comprising:

calculating combined biomarkers and clinical information Up- based on the equation:

$(\text{Biomarker Panel Score}) + (\text{Delusions Score}) - (\text{Grooming Score}) = \text{Up-Delusions Score};$

wherein the Biomarker Panel Score is obtained as per the method of claim 1;

wherein the Delusions Score is calculated with a clinical rating or self-report scales;

wherein the Grooming Score is calculated with a rating scale;

assessing the level of delusions of the individual by comparing the individual's Up-Delusions Score to a reference Up-Delusions Score;

administering a treatment for delusions to the individual when the individual's Up-Delusions Score is greater than a reference Up-Suicide Score; and

monitoring the individual's response to a treatment for delusions by determining changes in the Up-Delusions Score after initiating a treatment.

11. A method for assessing and treating schizophrenia and other psychotic disorders in general, in particular hallucinations and risk of developing hallucinations in an individual, comprising the steps of:

(a) obtaining a biological sample from an individual and quantifying the amounts of one or more RNA biomarkers in the biological sample, to create at least one panel of RNA biomarkers,

(b) quantifying the amounts of the RNA biomarkers in the at least one panel in a clinically relevant population to generate a reference expression level for the RNA biomarkers in a panel of RNA biomarkers;

(c) comparing the amounts of the biomarkers in the biological sample from the individual with the amounts of the RNA biomarkers present in the reference standard to generate a score for each biomarker a first panel and a second panel; wherein the biomarkers in the first panel comprise one or more of the following RNA biomarkers: (PRICKLE1), (NCAM1), (B3GALT5), (ARHGAP18), (PTP4A2), Acylphosphatase 2 (ACYP2), Reticulon 4 (RTN4), Cullin 4A (CUL4A), Zinc Finger E-Box Binding Homeobox 2 (ZEB2), Dystonin (DST), and Discs Large MAGUK Scaffold Protein 1 (DLG1), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting hallucinations or an increased risk for developing hallucinations; and

wherein the biomarkers in the second panel comprise one or more of the following RNA biomarkers: (PRL), (SERPING1), Ectonucleotide Pyrophosphatase/Phosphodiesterase 2 (ENPP2), (LAMA4), (KCNV1), Catenin Delta 1 (CTNND1), and FAT Atypical Cadherin 4 (FAT4), wherein the expression level of the RNA biomarker(s) in the sample is decreased relative to a reference expression level of the RNA biomarkers in the panel, denoting hallucinations or an increased risk for developing hallucinations;

(d) generating a score for the panel of RNA biomarkers, based on the scores of the biomarker(s) in the panel;

(e) determining a reference score for the panel in a clinically normal relevant population;

(f) identifying a difference between the score of the panel of biomarker(s) in the sample and the reference score of the panel of biomarker(s);

(g) identifying the individual as manifesting hallucinations or of having an elevated risk for developing hallucinations, based on the difference between the biomarker panel score of the individual relative to the biomarker panel score of the reference;

(h) treating the individual identified as having hallucinations or an elevated risk of hallucinations with one or more of the following: 1) a treatment based on clinical practice guidelines, 2) administering a therapeutically effective amount of a therapeutic drug (s), selected based on the specific biomarkers whose scores indicate that they are changed in the individual compared to a reference standard.

12. The method of claim 11, wherein the biomarkers are quantified in samples taken on two or more occasions from the individual.

13. The method of claim 11, wherein each biomarker is assigned a weighted coefficient based on the biomarkers importance in assessing and predicting hallucinations risk; and the biomarker panel score is based on the weighted coefficients of each of the biomarkers.

14. The method of claim 11, wherein the biological sample is a tissue sample or a fluid, such as cerebrospinal fluid, whole blood, blood serum, plasma, saliva, or other bodily fluid, or an extract or purification therefrom, or dilution thereof.

15. The method of claim 11, wherein the one or more therapeutic is one or more compounds selected from the group consisting of: clioquinol, pirinixic acid, moxislyte, Prestwick-685, exemestane, azacytidine, C-75, estradiol, tetraethylenepentamine, sparteine, guanethidine, idoxuridine, gliclazide, nitrendipine, N-acetyl-L-aspartic acid, sulfanilamide, doxazosin, pimozide, Proscillaridin, oxetacaine, BRD-K71489689, trichostatin A, A443654, AG 825, Proscillaridin A, Ala-Ala-Phe-CMK, Fluocinolone acetonide, manumycin A, curcumin, BRD-K68548958, CHR 2797, Tyrphostin AG 1478, Wortmannin, HY-50878, 598226, S1003, BRD-A52530684, CGP-60474, Buparlisib, and AS-601245.

16. The method of claim 11, wherein when the individual is male, and the biomarkers in a first panel (a) comprise one or more biomarkers selected from the group consisting of: (SH3PXD2A), Zinc Finger E-Box Binding Homeobox 2 (ZEB2), (PRICKLE1), (ARHGAP18), Acylphosphatase 2 (ACYP2), Reticulon 4 (RTN4), and Dystonin (DST), wherein the expression

level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting increased hallucinations; and

biomarkers in a second panel (b) comprise one or more biomarkers selected from the group consisting of: (PRL), (SERPING1), Ectonucleotide Pyrophosphatase/Phosphodiesterase 2 (ENPP2), (KCNV1), Mab-21 Like 1 (MAB21L1), Catenin Delta 1 (CTNND1), and FAT Atypical Cadherin 4 (FAT4), wherein the expression level of the biomarker(s) in the sample is decreased relative to a reference expression level, denoting increased hallucinations.

17. The method of claim 16, wherein the at least one therapeutic is at least one compound selected from the group consisting of: digoxigenin, doxazosin, meptazinol, promethazine, cefixime, velnacrine, cetirizine, eldeline, atropine oxide, clioquinol, nicotinic acid, clioquinol, galantamine, rolitetracycline, betahistine, sulconazole, monocrotaline, lanatoside C, Prestwick-1084, Naftidrofuryl, sulfachlorpyridazine, helveticoside, bezafibrate, mifepristone, trichostatin A, manumycin A, NCGC00189555-02, Buparlisib, linifanib, AZD-7762, Dinaciclib, Piretanide, KN-62, Fluticasone propionate, JAK3 Inhibitor VI, Sarmentogenin, Digoxin, Megestrol acetate, Oxymetazoline hydrochloride, U-0126, Tracazolate hydrochloride, Flufenamic acid, Fenofibrate, and U 99194 maleate.

18. The method of claim 11, wherein when the individual is female, and the biomarkers in a first panel (a) comprise one or more biomarkers selected from the group consisting of: (CELSR2), (KALRN), (B3GALT5), Protein Phosphatase 3 Catalytic Subunit Beta (PPP3CB), (ZFR), (THNSL1), (TNIK), Nuclear Receptor Subfamily 4 Group A Member 2 (NR4A2), Zinc Finger E-Box Binding Homeobox 2 (ZEB2), and (TNIK), wherein the expression level of the biomarker(s) in the sample is increased relative to a reference expression level, denoting increased hallucinations; and

biomarkers in a second panel (b) comprise one or more biomarkers selected from the group consisting of GNAS Complex Locus (GNAS), and Catenin Delta 1 (CTNND1), wherein the expression level of the biomarker(s) in the sample is decreased relative to a reference expression level, denoting increased hallucinations.

19. The method of claim 18, wherein the at least one therapeutic is at least one compound selected from the group consisting of: proglumide, quinethazone, esculin, MG-262, GW-8510,

haloperidol, guanethidine, deferoxamine, citiolone, meteneprost, amylocaine, CP-944629, Clemizole, IC-86621, Nortriptyline, CP-944629, Tanespimycin, Prestwick-674, 0317956-0000, and Pioglitazone.

20. A method of assessing and treating schizophrenia and other psychotic disorders in general, and hallucinations in particular in an individual, comprising:

calculating combined biomarkers and clinical information Up- based on the equation:

$$(\text{Biomarker Panel Score}) + (\text{Hallucinations Score}) - (\text{Grooming Score}) = \text{Up-Hallucinations Score};$$

wherein the Biomarker Panel Score is obtained as per the method of claim 11;

wherein the Hallucinations Score is calculated with a clinical rating or self-report scales;

wherein the Grooming Score is calculated with a rating scale;

assessing the level of hallucinations of the individual by comparing the individual's Up-Hallucinations Score to a reference Up-Hallucinations Score;

administering a treatment for hallucinations to the individual when the individual's Up-Hallucinations Score is greater than a reference Up-Suicide Score; and

monitoring the individual's response to a treatment for hallucinations by determining changes in the Up-Hallucinations Score after initiating a treatment.