

[Figure 1]

Study No. 370264




Fig. 1. Kaplan-Meier plot.

a): Start of administration in the control group and the EF-2001 group at 1.6 mg/kg/day for 17 days.
 b): Start of administration in the EF-2001 group at 1.6 mg/kg/day for 10 days and 4.8 mg/kg/day for 10 days.
 Significantly different from the control group (#: $p < 0.05$ by Generalized wilcoxon test).

[Figure 2]

Study No. 370264




Fig. 2. Rectal temperature.

a): Start of administration in the control group and the EF-2001 group at 1.6 mg/kg/day for 17 days.
b): Start of administration in the EF-2001 group at 1.6 mg/kg/day for 10 days and 4.8 mg/kg/day for 10 days.
Significantly different from the control group (*; p<0.05).
Significantly different from the EF-2001 group at 1.6 mg/kg/day for 17 days (#; p<0.05).
Significantly different from the EF-2001 group at 1.6 mg/kg/day for 10 days (††; p<0.01).

[Figure 3]

Study No. 370264




Fig. 3. Body weights.

- a) Start of administration in the control group and the EF-2001 group at 1.6 mg/kg/day for 17 days.
 b) Start of administration in the EF-2001 group at 1.6 mg/kg/day for 10 days and 4.8 mg/kg/day for 10 days.

[Figure 4]

Study No. 370264




Fig. 4. Food consumption.

Significantly different from the control group (*; $p < 0.05$).

Administration began on 7 days before infection in the control group, the EF-2001 group at 1.6 mg/kg/day for 17 days.

a): Start of administration in the EF-2001 group at 1.6 mg/kg/day for 10 days and 4.8 mg/kg/day for 10 days.

[Figure 5]

Study No. 370264




Fig. 5. Water consumption.

Significantly different from the control group (*: $p < 0.05$, **: $p < 0.01$).Significantly different from the EF-2001 group at 1.6 mg/kg/day for 17 days (#: $p < 0.05$, ##: $p < 0.01$).Significantly different from the EF-2001 group at 1.6 mg/kg/day for 10 days (†: $p < 0.05$, ††: $p < 0.01$).

Administration began on 7 days before infection in the control group, the EF-2001 group at 1.6 mg/kg/day for 17 days.

a): Start of administration in the EF-2001 group at 1.6 mg/kg/day for 10 days and 4.8 mg/kg/day for 10 days.

[Figure 6]

