

PSMA BINDING PROTEINS AND USES THEREOF**CROSS-REFERENCE TO RELATED APPLICATIONS**

This application claims the benefit of U.S. Serial No. 63/142,921 filed January 28, 2021; and U.S. Serial No. 63/165,448 filed March 24, 2021, the disclosure of each of which is incorporated by reference herein in its entirety.

REFERENCE TO SEQUENCE LISTING SUBMITTED ELECTRONICALLY

This application contains a sequence listing, which is submitted electronically via EFS-Web as an ASCII formatted sequence listing with a file "14620-625-228_SL.txt" and a creation date of January 7, 2022 and having a size of 561,620 bytes. The sequence listing submitted via EFS-Web is part of the specification and is herein incorporated by reference in its entirety.

FIELD

[0001] Provided herein, in certain aspects, are antibodies that bind to prostate-specific membrane antigen (PSMA), as well as recombinant cells containing the vectors, and compositions comprising the antibodies. Methods of making and using the antibodies are also provided. Also provided herein are, in certain aspects, are multispecific antibodies that bind PSMA, as well as recombinant cells containing the vectors, and compositions comprising the antibodies. In some embodiments, the multispecific antibodies bind to PSMA and cluster of differentiation 3 (CD3).

SUMMARY

[0002] In one aspect, provided herein is an isolated antibody that binds PSMA. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:32. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3,

respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:66. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and

a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:371; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:372. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a

VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:405; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:406. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:473; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:507; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:541; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:542. In one

embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:575; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:576. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:643; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:677; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:678. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:711; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3,

respectively, of a VL having an amino acid sequence of SEQ ID NO:474. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:745; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:746. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:779; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:780. In one embodiment, the PSMA antibody comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:813; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:814.

[0003] In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Kabat numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Chothia numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the AbM numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Contact numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the IMGT numbering system.

In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to a combination of the numbering systems provided herein.

[0004] In some embodiments, the isolated antibody comprises, consists of and/or consists essentially of a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:31, 99, 167, 235, 303, 371, 405, 439, 473, 507, 541, 575, 643, 677, 711, 779 or 813 and a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:32, 66, 100, 134, 236, 270, 372, 406, 474, 508, 542, 576, 678, 746, 780 or 814.

[0005] In some embodiments, the isolated antibody comprises, consists of and/or consists essentially of a heavy chain (HC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:33, 101, 169, 237, 305, 373, 407 or 441 and a light chain (LC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:34, 68, 102, 136, 238, 272, 374, or 408.

[0006] In some embodiments, provided herein is an isolated antibody that binds PSMA comprising, consisting of and/or consisting essentially of (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In some embodiments, provided herein is an isolated antibody that binds PSMA comprising, consisting of and/or consisting essentially of (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:205, 206, and 411, respectively, and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:242, 209, and 244, respectively. In some embodiments, provided herein is an isolated antibody that binds PSMA comprising, consisting of and/or consisting essentially of (i) a VH having an amino acid sequence of SEQ ID NO:SEQ ID NO:439; and (ii) a VL having an amino acid sequence of SEQ ID NO:270. In some embodiments, provided herein is an isolated antibody that binds PSMA comprising, consisting of and/or consisting essentially of (i) a HC having

an amino acid sequence of SEQ ID NO:441; and (ii) a LC having an amino acid sequence of SEQ ID NO:272.

[0007] In some embodiments, the isolated antibody binds a PSMA antigen. In some embodiments, isolated antibody binds a PSMA epitope. In some embodiments, the isolated antibody specifically binds to PSMA. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 form a binding site for an antigen of the PSMA. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 form a binding site for an epitope of the PSMA. In some embodiments, the PSMA is present on the surface of a cell. In some embodiments, the PSMA is present on the surface of a prostate cell. In some embodiments, the PSMA is present on the surface of a prostate cancer cell. In some embodiments, the PSMA is present on the surface of a renal cell. In some embodiments, the PSMA is present on the surface of a renal cancer cell.

[0008] In some embodiments of the PSMA antibody provided herein the binding domain that binds to PSMA is a scFv, an scFv dimer, a Fv, a Fab, a Fab, a F(ab')₂, a dsFv, a sdAb, a VHH or a single chain antibody.

[0009] In some embodiments, the PSMA antibody is a humanized antibody. In some embodiments, the PSMA antibody is a human antibody. In some embodiments, the isolated antibody is an IgG antibody. In some embodiments, the PSMA antibody is an IgG1 antibody. In some embodiments, the PSMA antibody is an IgG2 antibody. In some embodiments, the PSMA antibody is an IgG3 antibody. In some embodiments, the PSMA antibody is an IgG4 antibody. In some embodiments, the PSMA antibody comprises, consists of and/or consists essentially of a kappa light chain. In some embodiments, the PSMA antibody comprises, consists of and/or consists essentially of a lambda light chain. In some embodiments, the PSMA antibody is a monoclonal antibody. In some embodiments, the PSMA antibody is multivalent. In some embodiments, the PSMA antibody is capable of binding at least three antigens. In some embodiments, the PSMA antibody is capable of binding at least four antigens. In some embodiments, the PSMA antibody is capable of binding at least five antigens. In some embodiments, the PSMA antibody is a multispecific antibody. In some embodiments, the PSMA antibody is a bispecific antibody. In some embodiments, the PSMA antibody is a trispecific antibody. In some embodiments, the PSMA antibody is a quadraspecific antibody.

[0010] In another aspect, provided is a nucleic acid encoding a PSMA antibody provided herein. Also provided is a vector comprising, consisting of and/or consisting essentially of a nucleic acid encoding a PSMA antibody provided herein. Also provided is a host cell comprising, consisting of and/or consisting essentially of a vector comprising, consisting of and/or consisting essentially of a nucleic acid encoding a PSMA antibody provided herein. Also provided is a kit comprising, consisting of and/or consisting essentially of vector comprising, consisting of and/or consisting essentially of a nucleic acid encoding a PSMA antibody provided herein, and packaging for the same.

[0011] In another aspect, provided is a kit comprising, consisting of and/or consisting essentially of a PSMA antibody provided herein, and packaging for the same.

[0012] In another aspect, provided is a pharmaceutical composition comprising, consisting of and/or consisting essentially of a PSMA antibody provided herein, and a pharmaceutically acceptable carrier.

[0013] In another aspect, provided is a method of producing a pharmaceutical composition comprising, consisting of and/or consisting essentially of a PSMA antibody provided herein, comprising, consisting of and/or consisting essentially of combining the PSMA antibody with a pharmaceutically acceptable carrier to obtain the pharmaceutical composition.

[0014] In another aspect, provided is an isolated multispecific PSMAxCD3 antibody, comprising, consisting of and/or consisting essentially of: (a) a first binding domain that binds to PSMA, and (b) a second binding domain that binds CD3. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:32. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ

ID NO:31; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:66. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH

comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:371; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL

CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:372. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:405; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:406. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:473; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:507; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3,

respectively, of a VH having an amino acid sequence of SEQ ID NO:541; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:542. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:575; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:576. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:643; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:677; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:678. In some embodiments, the first binding domain that binds to PSMA comprises,

consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:711; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:745; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:746. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:779; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:780. In some embodiments, the first binding domain that binds to PSMA comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:813; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:814. In some embodiments, the second binding domain that binds to CD3 comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:1505; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3

having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:1464. In some embodiments, the second binding domain that binds to CD3 comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:847; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:848. In some embodiments, the second binding domain that binds to CD3 comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:915; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:916. In some embodiments, the second binding domain that binds to CD3 comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:983; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:984. In some embodiments, the second binding domain that binds to CD3 comprises, consists of and/or consists essentially of: (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:1463; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:1464. In some embodiments, the second binding domain that binds to CD3 comprises, consists of and/or consists essentially of a scFv comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, a VL CDR3 having an amino acid sequence

of a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, a VL CDR3 respectively, of a scFv having an amino acid sequence of SEQ ID NO:1524.

[0015] In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the multispecific PSMAxCD3 antibody are according to the Kabat numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the multispecific PSMAxCD3 antibody are according to the Chothia numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the multispecific PSMAxCD3 antibody are according to the AbM numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the multispecific PSMAxCD3 antibody are according to the Contact numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the multispecific PSMAxCD3 antibody are according to the IMGT numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the multispecific PSMAxCD3 antibody are according to a combination of the numbering systems provided herein.

[0016] In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises, consists of and/or consists essentially of a heavy chain (HC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:33, 101, 169, 237, 305, 373, 407, 441, 1242, 1244, 1248, 1250, 1252, 1254, 1256, 1258, 1260, 1262, 1264, 1266, 1268 or 1270 and/or a light chain (LC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:34, 68, 102, 136, 238, 272, 374 or 408.

[0017] In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises, consists of and/or consists essentially of a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of any one of SEQ ID NOs:1485-1500 or SEQ ID NOs:1526-1531.

[0018] In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises, consists of and/or consists essentially of (i) a HC having an amino acid sequence of SEQ ID NO:SEQ ID NO:441; and (ii) a LC having an amino acid sequence of SEQ ID NO:SEQ ID NO:272.

[0019] In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the second binding domain that binds CD3 comprises, consists of and/or consists essentially of a heavy chain (HC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:849, 883, 917, 951, 985, 1019, 1504, 1455, 1192, 1194, 1167, 1218 or 1238 and/or a light chain (LC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:850, 918, 986, 1193, 1195 or 1219.

[0020] In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the second binding domain that binds CD3 comprises, consists of and/or consists essentially of a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:1186, 1187, 1523 or 1524.

[0021] Provided herein is an isolated bispecific antibody comprising, consisting of and/or consisting essentially of a first binding domain that binds PSMA and a second binding domain that binds CD3, wherein the first binding domain that binds PSMA comprises, consists of and/or consists essentially of (i) a VH comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising, consisting of and/or consisting essentially of a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270, and the second binding domain that binds CD3 comprises, consists of and/or consists essentially of a scFv comprising, consisting of and/or consisting essentially of a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, and a VL CDR3 of a scFv having an amino acid sequence of SEQ ID NO:1524. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises, consists of and/or consists essentially of a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:205, 206, 411, 242, 209 and 244, respectively, wherein the amino acid sequences are according to the Kabat numbering system; and the second binding domain that binds CD3 comprises, consists of and/or consists essentially of a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:1467, 1468, 1506, 1470, 1471 and 1472, respectively; wherein the amino acid sequences are according to the Kabat numbering system. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises, consists of and/or consists essentially of (i) a VH having an amino

acid sequence of SEQ ID NO:439; and (ii) a VL having an amino acid sequence of SEQ ID NO:270, and the second binding domain that binds CD3 comprises, consists of and/or consists essentially of a scFv of SEQ ID NO:1524, respectively. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises, consists of and/or consists essentially of (i) a HC2 having an amino acid sequence of SEQ ID NO:441; and (ii) a LC2 having an amino acid sequence of SEQ ID NO:272, respectively; and the second binding domain that binds CD3 comprises, consists of and/or consists essentially of a HC1 of SEQ ID NO:1455.

[0022] In some embodiments of the multispecific antibody provided herein, the first binding domain, the second binding domain and/or the first and second is a scFv, an scFv dimer, a Fv, a Fab, a Fab, a F(ab')₂, a dsFv, a sdAb, a VHH or a single chain antibody.

[0023] In some embodiments, the multispecific PSMAxCD3 antibody is a humanized antibody. In some embodiments, the multispecific PSMAxCD3 antibody is a human antibody. In some embodiments, the multispecific PSMAxCD3 antibody is an IgG antibody. In some embodiments, the multispecific PSMAxCD3 antibody is an IgG1 antibody. In some embodiments, the multispecific PSMAxCD3 antibody is an IgG2 antibody. In some embodiments, the multispecific PSMAxCD3 antibody is an IgG3 antibody. In some embodiments, the multispecific PSMAxCD3 antibody is an IgG4 antibody. In some embodiments, the multispecific PSMAxCD3 antibody comprises, consists of and/or consists essentially of a kappa light chain. In some embodiments, the multispecific PSMAxCD3 antibody comprises, consists of and/or consists essentially of a lambda light chain. In some embodiments, the multispecific PSMAxCD3 antibody is a monoclonal antibody.

[0024] In some embodiments of the multispecific PSMAxCD3 antibody, the first binding domain binds a PSMA antigen. In some embodiments of the multispecific PSMAxCD3 antibody, the first binding domain binds a PSMA epitope. In some embodiments of the multispecific PSMAxCD3 antibody, the first binding domain specifically binds to PSMA. In some embodiments of the multispecific PSMAxCD3 antibody, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the first binding domain form a binding site for an antigen of the PSMA. In some embodiments of the multispecific PSMAxCD3 antibody, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the first binding domain form a binding site for an epitope of the PSMA. In some embodiments of the multispecific PSMAxCD3 antibody, the second binding domain binds a CD3 antigen. In some embodiments of the multispecific PSMAxCD3 antibody, the second binding

domain binds a CD3 epitope. In some embodiments of the multispecific PSMAxCD3 antibody, the second binding domain specifically binds to CD3. In some embodiments of the multispecific PSMAxCD3 antibody, the second binding domain form a binding site for an antigen of the CD3. In some embodiments of the multispecific PSMAxCD3 antibody, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the multispecific PSMAxCD3 antibody binding domain form a binding site for an epitope of the CD3.

[0025] In some embodiments, the PSMA is present on the surface of a cell. In some embodiments, the cell is a prostate cell. In some embodiments, the cell is a prostate cancer cell. In some embodiments, the cell is a renal cell. In some embodiments, the cell is a renal cancer cell.

[0026] In some embodiments of the multispecific PSMAxCD3 antibody, the antibody is a bispecific antibody. In some embodiments of the multispecific PSMAxCD3 antibody, is a trispecific antibody. In some embodiments of the multispecific PSMAxCD3 antibody, is a quadraspecific antibody.

[0027] In another aspect, provided is a nucleic acid encoding a multispecific PSMAxCD3 antibody provided herein. Also provided is a vector comprising, consisting of and/or consisting essentially of a nucleic acid encoding a multispecific PSMAxCD3 antibody provided herein. Also provided is a host cell comprising, consisting of and/or consisting essentially of a vector comprising, consisting of and/or consisting essentially of a nucleic acid encoding a multispecific PSMAxCD3 antibody provided herein. Also provided is a kit comprising, consisting of and/or consisting essentially of vector comprising, consisting of and/or consisting essentially of a nucleic acid encoding a multispecific PSMAxCD3 antibody provided herein, and packaging for the same.

[0028] In another aspect, provided is a kit comprising, consisting of and/or consisting essentially of a multispecific PSMAxCD3 antibody provided herein, and packaging for the same.

[0029] In another aspect, provided is a pharmaceutical composition comprising, consisting of and/or consisting essentially of a multispecific PSMAxCD3 antibody provided herein, and a pharmaceutically acceptable carrier.

[0030] In another aspect, provided is a method of producing a pharmaceutical composition comprising, consisting of and/or consisting essentially of a multispecific PSMAxCD3 antibody provided herein, comprising, consisting of and/or consisting

essentially of combining the multispecific PSMAxCD3 antibody with a pharmaceutically acceptable carrier to obtain the pharmaceutical composition.

[0031] In another aspect, provided is a method of directing a CD3-expressing T cell to a PSMA-expressing target cell, comprising, consisting of and/or consisting essentially of contacting the T cell with the multispecific PSMAxCD3 antibody provided herein. In some embodiments, the contacting directs the T cell to the target cell. In another aspect, provided is a method of inhibiting the growth or proliferation of a PSMA-expressing target cell, comprising, consisting of and/or consisting essentially of contacting the target cell with the multispecific PSMAxCD3 antibody provided herein. In some embodiments, the contacting is in the presence of CD3-expressing T cells. In some embodiments, the target cell expresses PSMA on the cell surface. In some embodiments, the target cell is a prostate cell. In some embodiments, the target cell is a prostate cancer cell. In some embodiments, the target cell is a renal cell. In some embodiments, the target cell is a renal cancer cell.

[0032] In another aspect, provided is a method of eliminating a PSMA-expressing target cell in a subject, comprising, consisting of and/or consisting essentially of administering to the subject an effective amount of the multispecific PSMAxCD3 antibody provided herein. In another aspect, provided is a method of treating a disease or disorder in a subject, comprising, consisting of and/or consisting essentially of administering to the subject an effective amount of the multispecific PSMAxCD3 antibody provided herein. In some embodiments, the disease or disorder is a disease or disorder of the prostate. In some embodiments, the disease or disorder of the prostate is prostate inflammation. In some embodiments, the disease or disorder of the prostate is Benign prostatic hyperplasia. In some embodiments, the disease or disorder of the prostate is prostate cancer. In some embodiments, the disease or disorder of the prostate is metastatic castration-resistant prostate cancer (mCRPC). In some embodiments, the disease or disorder is a renal disease or disorder. In some embodiments, the renal disease or disorder is renal cancer. In some embodiments, the renal disease or disorder is a renal cell carcinoma. In some embodiments, the renal cell carcinoma is a metastatic renal cell carcinoma (mRCC). In some embodiments, the subject is a subject in need thereof. In some embodiments, the subject is a human.

BRIEF DESCRIPTION OF THE DRAWINGS

[0033] The foregoing summary, as well as the following detailed description of specific embodiments of the present application, will be better understood when read in conjunction with the appended drawings. It should be understood, however, that the application is not limited to the precise embodiments shown in the drawings.

[0034] FIG. 1 shows HDX-MS epitope mapping of PSMA against PS3B1352 (top) and PS3B1353 (bottom). G is glycosylation site. Black box is epitope and gray is probable epitope. White box indicates no/little change in deuteration level in the presence of the antibody. The residues without box indicate the HDX behaviors were not monitored, because there is no peptide to cover the residues or the residues are the first two residues of a peptide. The epitopes of PS3B1352 and PS3B1353 are identical. FIG. 1 discloses SEQ ID NO:1483.

[0035] FIG. 2 shows HDX-MS identified epitopes of PSMA overlaid on X-ray crystal structure. The circled alpha-helix represents the HDX-MS identified epitope. The black: epitope; gray: potential epitope.

[0036] FIG. 3 shows PAN-T cell binding assay. Human PAN-T cells were treated with various concentrations of PSMA/CD3 bispecific antibodies and incubated at 37°C for 30 minutes followed by CD3 cell surface expression analysis by flow cytometry.

[0037] FIG. 4 shows the non-linear regression fit of four-parameter function of PSMA ligand binding of C4-2B human prostate tumor cells.

[0038] FIG. 5 shows a target cell binding assay. C4-2B human prostate tumor cells were treated with various concentrations of PSMA/CD3 bispecific antibodies and incubated at 37°C for 30 minutes followed by PSMA cell surface expression analysis by flow cytometry.

[0039] FIG. 6 shows internalization of PSMA. Human C4-2B prostate tumor cells were incubated with PSMA/CD3 bispecific antibodies conjugated to IncuCyte® Human Fab-fluor-pH Red Antibody Labeling Dye for 24 hours.

[0040] FIGS. 7A-7H show bispecific anti-PSMA/ anti-T cell redirection antibodies evaluated in an IncuCyte®-based cytotoxicity assay. Isolated PAN-T cells were co-incubated with PSMA+ C4-2B cells in the presence of bispecific PSMA/ T cell redirection

antibodies for 120 hours. Shown are data for (A) PS3B1352, (B) PS3B1356, (C) PS3B1353, (D) PS3B1357, (E) PS3B1354, (F) PS3B937, (G) PS3B1355, and (H) PS3B1358.

[0041] FIG. 8 shows T cell redirected killing assay. Normal human PBMCs were combined with C4-2B human prostate tumor cells transduced with IncuCyte® NucLight red nuclear dye and treated with PSMA/CD3 bispecific antibodies for 5 days.

[0042] FIG. 9 shows cytokine induction by bispecific anti-PSMA/ anti-T cell redirection antibodies. Isolated PAN-T cells were co-incubated with PSMA+ C4-2B cells in the presence of bispecific anti-PSMA/ anti-T cell redirection antibodies for the indicated time points. IFN-gamma concentration was measured from supernatants collected at the indicated time points.

DETAILED DESCRIPTION

[0043] Various publications, articles and patents are cited or described in the background and throughout the specification; each of these references is herein incorporated by reference in its entirety as though fully set forth. Discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is for the purpose of providing context for the invention. Such discussion is not an admission that any or all of these matters form part of the prior art with respect to any inventions disclosed or claimed.

[0044] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention pertains. Otherwise, certain terms used herein have the meanings as set forth in the specification. Although any methods and materials similar or equivalent to those described herein can be used in the practice for testing embodiments of the present disclosure, exemplary materials and methods are described herein.

[0045] When a list is presented, unless stated otherwise, it is to be understood that each individual element of that list, and every combination of that list, is a separate embodiment. For example, a list of embodiments presented as “A, B, or C” is to be interpreted as including the embodiments, “A,” “B,” “C,” “A or B,” “A or C,” “B or C,” or “A, B, or C.”

[0046] It must be noted that as used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to “a cell” includes a combination of two or more cells, and the like.

[0047] Unless otherwise stated, any numerical values, such as a concentration or a concentration range described herein, are to be understood as being modified in all instances by the term “about.” Thus, a numerical value typically includes $\pm 10\%$ of the recited value. For example, a concentration of 1 mg/mL includes 0.9 mg/mL to 1.1 mg/mL. Likewise, a concentration range of 1% to 10% (w/v) includes 0.9% (w/v) to 11% (w/v). “About” means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, i.e., the limitations of the measurement system. Unless explicitly stated otherwise within the Examples or elsewhere in the Specification in the context of a particular assay, result or embodiment, “about” means within one standard deviation per the practice in the art, or a range of up to 5%, whichever is larger.

[0048] As used herein, the use of a numerical range expressly includes all possible subranges, all individual numerical values within that range, including integers within such ranges and fractions of the values unless the context clearly indicates otherwise.

[0049] Unless otherwise indicated, the term “at least” preceding a series of elements is to be understood to refer to every element in the series.

[0050] Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments described herein. Such equivalents are intended to be encompassed by the invention.

[0051] As used herein, the terms “comprises,” “comprising,” “includes,” “including,” “has,” “having,” “contains” or “containing,” or any other variation thereof, will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers and are intended to be non-exclusive or open-ended. For example, a composition, a mixture, a process, a method, an article, or an apparatus that comprises a list of elements is not necessarily limited to only those elements but can include other elements not expressly listed or inherent to such composition, mixture, process, method, article, or apparatus. Further, unless expressly stated to the contrary, “or” refers to an inclusive or and not to an exclusive or. For example, a condition A or B is satisfied by any one of the following: A is true (or present) and B is false (or not present), A is false (or not present) and B is true (or present), and both A and B are true (or present).

[0052] As used herein, the conjunctive term “and/or” between multiple recited elements is understood as encompassing both individual and combined options. For instance, where

two elements are conjoined by “and/or,” a first option refers to the applicability of the first element without the second. A second option refers to the applicability of the second element without the first. A third option refers to the applicability of the first and second elements together. Any one of these options is understood to fall within the meaning, and therefore satisfy the requirement of the term “and/or” as used herein. Concurrent applicability of more than one of the options is also understood to fall within the meaning, and therefore satisfy the requirement of the term “and/or.”

[0053] As used herein, the term “consists of,” or variations such as “consist of” or “consisting of,” as used throughout the specification and claims, indicate the inclusion of any recited integer or group of integers, but that no additional integer or group of integers can be added to the specified method, structure, or composition.

[0054] As used herein, the term “consists essentially of,” or variations such as “consist essentially of” or “consisting essentially of,” as used throughout the specification and claims, indicate the inclusion of any recited integer or group of integers, and the optional inclusion of any recited integer or group of integers that do not materially change the basic or novel properties of the specified method, structure or composition. See M.P.E.P. § 2111.03.

[0055] As used herein, “subject” means any animal, such as a mammal or a human. The term “mammal” as used herein, encompasses any mammal. Examples of mammals include, but are not limited to, cows, horses, sheep, pigs, cats, dogs, mice, rats, rabbits, guinea pigs, monkeys, humans, *etc.* In specific embodiments, the subject is a human.

[0056] It should also be understood that the terms “about,” “approximately,” “generally,” “substantially,” and like terms, used herein when referring to a dimension or characteristic of a component of embodiments provided herein, indicate that the described dimension/characteristic is not a strict boundary or parameter and does not exclude minor variations therefrom that are functionally the same or similar, as would be understood by one having ordinary skill in the art. At a minimum, such references that include a numerical parameter would include variations that, using mathematical and industrial principles accepted in the art (*e.g.*, rounding, measurement or other systematic errors, manufacturing tolerances, *etc.*), would not vary the least significant digit.

[0057] The terms “identical” or percent “identity,” in the context of two or more nucleic acids or polypeptide sequences (*e.g.*, PSMA antibodies and polynucleotides that encode

them, CD3 antibodies and polynucleotides that encode them), refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same, when compared and aligned for maximum correspondence, as measured using one of the following sequence comparison algorithms or by visual inspection.

[0058] For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

[0059] Optimal alignment of sequences for comparison can be conducted, *e.g.*, by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (see generally, *Current Protocols in Molecular Biology*, F.M. Ausubel *et al.*, eds., *Current Protocols*, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (1995 Supplement) (Ausubel)).

[0060] Examples of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul *et al.* (1990) *J. Mol. Biol.* 215: 403-410 and Altschul *et al.* (1997) *Nucleic Acids Res.* 25: 3389-3402, respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length *W* in the query sequence, which either match or satisfy some positive-valued threshold score *T* when aligned with a word of the same length in a database sequence. *T* is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased.

[0061] Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a word length (W) of 11, an expectation (E) of 10, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a word length (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, Proc. Natl. Acad. Sci. USA 89:10915 (1989)).

[0062] In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, *e.g.*, Karlin & Altschul, Proc. Nat'l. Acad. Sci. USA 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P (N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more such as less than about 0.01, and or less than about 0.001.

[0063] A further indication that two nucleic acid sequences or polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules hybridize to each other under stringent conditions.

[0064] As used herein, the term "polynucleotide," synonymously referred to as "nucleic acid molecule," "nucleotides" or "nucleic acids," refers to any polyribonucleotide or polydeoxyribonucleotide, which can be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA,

DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that can be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, “polynucleotide” refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. “Modified” bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, “polynucleotide” embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. “Polynucleotide” also embraces relatively short nucleic acid chains, often referred to as oligonucleotides.

[0065] As used herein, “variant” refers to a polypeptide or a polynucleotide that differs from a reference polypeptide or a reference polynucleotide by one or more modifications, for example one or more substitutions, insertions or deletions.

[0066] As used herein, the term “vector” is a replicon in which another nucleic acid segment can be operably inserted so as to bring about the replication or expression of the segment. Vector polynucleotides typically contain elements, such as origins of replication, polyadenylation signal or selection markers, that function to facilitate the duplication or maintenance of these polynucleotides in a biological system, such as a cell, virus, animal, plant, and reconstituted biological systems utilizing biological components capable of duplicating a vector. The vector polynucleotide can be DNA or RNA molecules or a hybrid of these, single stranded or double stranded.

[0067] As used herein, “expression vector” refers to a vector that can be utilized in a biological system or in a reconstituted biological system to direct the translation of a polypeptide encoded by a polynucleotide sequence present in the expression vector. As used herein, the term “host cell” refers to a cell comprising a nucleic acid molecule provided herein.

[0068] As used herein, “host cell” can be any type of cell, *e.g.*, a primary cell, a cell in culture, or a cell from a cell line. In one embodiment, a “host cell” is a cell transfected with a nucleic acid molecule provided herein. In another embodiment, a “host cell” is a progeny

or potential progeny of such a transfected cell. A progeny of a cell may or may not be identical to the parent cell, *e.g.*, due to mutations or environmental influences that can occur in succeeding generations or integration of the nucleic acid molecule into the host cell genome.

[0069] The term “expression” as used herein, refers to the biosynthesis of a gene product. The term encompasses the transcription of a gene into RNA. The term also encompasses translation of RNA into one or more polypeptides, and further encompasses all naturally occurring post-transcriptional and post-translational modifications. The expressed antibody can be within the cytoplasm of a host cell, into the extracellular milieu such as the growth medium of a cell culture or anchored to the cell membrane.

[0070] As used herein, “flow cytometry” is a technology that is used to analyze the physical and chemical characteristics of particles in a fluid as it passes through at least one laser. Cell components are fluorescently labelled and then excited by the laser to emit light at varying wavelengths (Adan *et al.*, Crit. Rev. Biotech. (2016) 1549-7801).

[0071] As used herein, “overexpress”, “overexpressed” and “overexpressing” interchangeably refers to a sample such as a cancer cell, malignant cell or cancer tissue that has measurably higher levels of PSMA when compared to a reference sample. The overexpression can be caused by gene amplification or by increased transcription or translation. Expression and overexpression of protein in the sample can be measured using well known assays using, for example ELISA, immunofluorescence, flow cytometry or radioimmunoassay on live or lysed cells. Expression and overexpression of a polynucleotide in the sample can be measured, for example, using fluorescent in situ hybridization, Southern blotting, or PCR techniques. A protein or a polynucleotide is overexpressed when the level of the protein or the polynucleotide in the sample is at least 1.5-fold higher when compared to the reference sample. Selection of the reference sample is well known.

[0072] As used herein, “sample” refers to a collection of similar fluids, cells, or tissues isolated from a subject, as well as fluids, cells, or tissues present within a subject. Exemplary samples are of biological fluids such as blood, serum and serosal fluids, plasma, lymph, urine, saliva, cystic fluid, tear drops, feces, sputum, mucosal secretions of the secretory tissues and organs, vaginal secretions, ascites fluids such as those associated with non-solid tumors, fluids of the pleural, pericardial, peritoneal, abdominal and other body

cavities, fluids collected by bronchial lavage, liquid solutions contacted with a subject or biological source, for example, cell and organ culture medium including cell or organ conditioned medium, lavage fluids and the like, tissue biopsies, fine needle aspirations or surgically resected tumor tissue.

[0073] A “cancer cell” or a “tumor cell” as used herein refers to a cancerous, pre-cancerous or transformed cell, either *in vivo*, *ex vivo*, or in tissue culture, that has spontaneous or induced phenotypic changes. These changes do not necessarily involve the uptake of new genetic material. Although transformation can arise from infection with a transforming virus and incorporation of new genomic nucleic acid or uptake of exogenous nucleic acid, it can also arise spontaneously or following exposure to a carcinogen, thereby mutating an endogenous gene. Transformation/cancer is exemplified by morphological changes, immortalization of cells, aberrant growth control, foci formation, proliferation, malignancy, modulation of tumor specific marker levels, invasiveness, tumor growth in suitable animal hosts such as nude mice, and the like, *in vitro*, *in vivo*, and *ex vivo* (Freshney, *Culture of Animal Cells: A Manual of Basic Technique* (3rd ed. 1994)). Unless otherwise stated, any numerical values, such as a concentration or a concentration range described herein, are to be understood as being modified in all instances by the term “about.” Thus, a numerical value typically includes $\pm 10\%$ of the recited value. For example, a concentration of 1 mg/mL includes 0.9 mg/mL to 1.1 mg/mL. Likewise, a concentration range of 1% to 10% (w/v) includes 0.9% (w/v) to 11% (w/v). As used herein, the use of a numerical range expressly includes all possible subranges, all individual numerical values within that range, including integers within such ranges and fractions of the values unless the context clearly indicates otherwise.

[0074] As used herein, the terms “peptide,” “polypeptide,” or “protein” can refer to a molecule comprised of amino acids and can be recognized as a protein by those of skill in the art. The conventional one-letter or three-letter code for amino acid residues is used herein. The terms “peptide,” “polypeptide,” and “protein” can be used interchangeably herein to refer to polymers of amino acids of any length. The polymer can be linear or branched, it can comprise modified amino acids, and it can be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified naturally or by intervention; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation or modification, such as conjugation with a labeling component. Also included within the definition are, for example, polypeptides

containing one or more analogs of an amino acid (including, for example, unnatural amino acids, *etc.*), as well as other modifications known in the art.

[0075] The peptide sequences described herein are written according to the usual convention whereby the N-terminal region of the peptide is on the left and the C-terminal region is on the right. Although isomeric forms of the amino acids are known, it is the L-form of the amino acid that is represented unless otherwise expressly indicated.

[0076] Conventional one and three-letter amino acid codes are used herein as shown in Table 1.

Table 1. Conventional Amino Acid Codes

Amino acid	Three-letter code	One-letter code
Alanine	Ala	A
Arginine	Arg	R
Asparagine	Asn	N
Aspartate	Asp	D
Cysteine	Cys	C
Glutamate	Gln	E
Glutamine	Glu	Q
Glycine	Gly	G
Histidine	His	H
Isoleucine	Ile	I
Lysine	Lys	K
Methionine	Met	M
Phenylalanine	Phe	F
Proline	Pro	P
Serine	Ser	S
Threonine	Thr	T
Tryptophan	Trp	W
Tyrosine	Tyr	Y
Valine	Val	V

[0077] As used herein, “effector antigens” are antigens from cells of the immune system, which can stimulate or trigger cytotoxicity, phagocytosis, antigen presentation, cytokine release. Such effector antigens are from, for example but not limited to, T cells and natural killer (NK) cells. Examples of suitable specificities for effector antigens include but are not limited to CD3 or CD3 subunits such as CD3ε for T cells and CD16 for NK cells. Such cell surface molecules of effector cells are suitable for mediating cell killing. Effector cells are

cells of the immune system, which can stimulate or trigger cytotoxicity, phagocytosis, antigen presentation, cytokine release. Such effector cells are, for example but not limited to, T cells, natural killer (NK) cells, granulocytes, monocytes, macrophages, dendritic cells, and antigen-presenting cells. Examples of suitable specificities for effector cells include but are not limited to CD2, CD3 and CD3 subunits such as CD3e, CD5, CD28 and other components of the T cell receptor (TCR) for T cells; CD16, CD16A, CD25, CD38, CD44, CD56, CD69, CD94, CD335 (NKp46), CD336, (NKp44), CD337 (NKp30), NKp80, NKG2C and NKG2D, DNAM, NCRs for NK cells; CD18, CD64 and CD89 for granulocytes; CD18, CD32, CD64, CD89 and mannose receptor for monocytes and macrophages; CD64 and mannose receptor for dendritic cells; as well as CD35. In certain embodiments, those specificities, i. e. cell surface molecules, of effector cells are suitable for mediating cell killing upon binding of a bispecific or multispecific molecules to such cell surface molecule and, thereby, inducing cytolysis or apoptosis.

[0078] As used herein, “multispecific PSMAxCD3 antibody”, “PSMA/CD3 antibody”, “multispecific anti-PSMAxCD3 antibody” or “anti-PSMA/CD3 antibody” refers to a molecule comprising at least one binding domain specifically binding PSMA and at least one binding domain specifically binding CD3. The domains specifically binding PSMA and CD3 are typically VH/VL pairs. The bispecific anti-PSMAxCD3 antibody can be monovalent in terms of its binding to either PSMA or CD3.

[0079] “Valent” refers to the presence of a specified number of binding sites specific for an antigen in a molecule. As such, the terms “monovalent”, “bivalent”, “tetravalent”, and “hexavalent” refer to the presence of one, two, four and six binding sites, respectively, specific for an antigen in a molecule. “Multivalent” refers to the presence of two or more binding sites specific for an antigen in a molecule.

Antibodies

[0080] Provided herein are PSMA antibodies or antigen-binding fragments thereof, nucleic acids and expression vectors encoding the antibodies, recombinant cells containing the vectors, and compositions comprising the antibodies. Methods of making the antibodies, and methods of using the antibodies to treat diseases are also provided. The antibodies disclosed herein possess one or more desirable functional properties, including but not limited to high-affinity binding to PSMA or high specificity to PSMA. In certain

embodiments, the antibodies disclosed herein possess the ability to treat or prevent a disease or disorder when administered to a subject alone or in combination with other therapies.

[0081] Also provided herein are PSMA bispecific antibodies or antigen-binding fragments thereof, nucleic acids and expression vectors encoding the bispecific antibodies, recombinant cells containing the vectors, and compositions comprising the bispecific antibodies. Methods of making the antibodies, and methods of using the bispecific antibodies to treat diseases, including cancer, are also provided. The antibodies disclosed herein possess one or more desirable functional properties. In some embodiments, the bispecific antibodies provided herein have high-affinity binding to PSMA. In some embodiments, the bispecific antibodies provided herein have high-affinity binding to a second target antigen. In some embodiments, the bispecific antibodies provided herein have high specificity to PSMA. In some embodiments, the bispecific antibodies provided herein have high specificity to a second target antigen. In some, embodiments, the bispecific antibodies provided herein have high specificity to CD3. In some embodiments, the bispecific antibodies provided herein have the ability to treat or prevent a disease or disorder when administered alone. In some embodiments, the bispecific antibodies provided herein have the ability to treat or prevent a disease or disorder when administered in combination with other therapies.

[0082] As used herein, the term “antibody” is used in a broad sense and includes immunoglobulin or antibody molecules including human, humanized, composite and chimeric antibodies and antibody fragments that are monoclonal or polyclonal. In general, antibodies are proteins or peptide chains that exhibit binding specificity to a specific antigen. Antibody structures are well known. Immunoglobulins can be assigned to five major classes (*i.e.*, IgA, IgD, IgE, IgG and IgM), depending on the heavy chain constant domain amino acid sequence. IgA and IgG are further sub-classified as the isotypes IgA1, IgA2, IgG1, IgG2, IgG3 and IgG4. Accordingly, the antibodies provided herein can be of any of the five major classes or corresponding sub-classes. In specific embodiments, the antibodies provided herein are IgG1, IgG2, IgG3 or IgG4. Antibody light chains of vertebrate species can be assigned to one of two clearly distinct types, namely kappa and lambda, based on the amino acid sequences of their constant domains. Accordingly, the antibodies provided herein can, in certain embodiments, contain a kappa light chain constant domain. The antibodies provided herein can, in certain embodiments, also contain a lambda light chain constant domain. According to particular embodiments, the antibodies provided herein

include heavy and/or light chain constant regions from rat or human antibodies. In specific embodiments, the constant region is a human constant region.

[0083] In addition to the heavy and light constant domains, antibodies contain an antigen-binding region that is made up of a light chain variable region (VL) and a heavy chain variable region (VH), each of which contains three domains (*i.e.*, complementarity determining regions 1 (CDR1), CDR2 and CDR3). A “CDR” refers to one of three hypervariable regions (HCDR1, HCDR2 or HCDR3) within the non-framework region of the immunoglobulin (Ig or antibody) VH β -sheet framework, or one of three hypervariable regions (LCDR1, LCDR2 or LCDR3) within the non-framework region of the antibody VL β -sheet framework. Accordingly, CDRs are variable region sequences interspersed within the framework region sequences. CDR regions are well known to those skilled in the art and have been defined by, for example, Kabat as the regions of most hypervariability within the antibody variable (V) domains (Kabat *et al.*, *J. Biol. Chem.* 252:6609-6616 (1977); Kabat, *Adv. Prot. Chem.* 32:1-75 (1978); Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)). CDR region sequences also have been defined structurally by Chothia as those residues that are not part of the conserved β -sheet framework, and thus are able to adapt different conformations (Chothia and Lesk, *J. Mol. Biol.* 196:901-917 (1987)). Both terminologies are well recognized in the art. CDR region sequences have also been defined by AbM, Contact and IMGT. Exemplary CDR region sequences are illustrated herein, for example, in the tables provided in the Examples below. The positions of CDRs within a canonical antibody variable region have been determined by comparison of numerous structures (Al-Lazikani *et al.*, *J. Mol. Biol.* 273:927-948 (1997); Morea *et al.*, *Methods* 20:267-279 (2000)). Because the number of residues within a hypervariable region varies in different antibodies, additional residues relative to the canonical positions are conventionally numbered with a, b, c and so forth next to the residue number in the canonical variable region numbering scheme (Al-Lazikani *et al.*, *supra* (1997)). Such nomenclature is similarly well known to those skilled in the art.

[0084] The light chain variable region CDR1 domain is interchangeably referred to herein as LCDR1 or VL CDR1. The light chain variable region CDR2 domain is interchangeably referred to herein as LCDR2 or VL CDR2. The light chain variable region CDR3 domain is interchangeably referred to herein as LCDR3 or VL CDR3. The heavy chain variable region CDR1 domain is interchangeably referred to herein as HCDR1 or VH

CDR1. The heavy chain variable region CDR2 domain is interchangeably referred to herein as HCDR2 or VH CDR2. The heavy chain variable region CDR1 domain is interchangeably referred to herein as HCDR3 or VH CDR3.

[0085] The term “hypervariable region”, such as a VH or VL, when used herein refers to the regions of an antibody variable region that are hypervariable in sequence and/or form structurally defined loops. Generally, antibodies comprise six hypervariable regions; three in the VH (HCDR1, HCDR2, HCDR3), and three in the VL (LCDR1, LCDR2, LCDR3). A number of hypervariable region delineations are in use and are encompassed herein. The “Kabat” CDRs are based on sequence variability and are the most commonly used (see, *e.g.*, Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)). “Chothia” refers instead to the location of the structural loops (see, *e.g.*, Chothia and Lesk, *J. Mol. Biol.* 196:901-917 (1987)). The end of the Chothia CDR-HCDR1 loop when numbered using the Kabat numbering convention varies between H32 and H34 depending on the length of the loop (this is because the Kabat numbering scheme places the insertions at H35A and H35B; if neither 35A nor 35B is present, the loop ends at 32; if only 35A is present, the loop ends at 33; if both 35A and 35B are present, the loop ends at 34). The “AbM” hypervariable regions represent a compromise between the Kabat CDRs and Chothia structural loops, and are used by Oxford Molecular’s AbM antibody modeling software (see, *e.g.*, Martin, in Antibody Engineering, Vol. 2, Chapter 3, Springer Verlag). “Contact” hypervariable regions are based on an analysis of the available complex crystal structures.

[0086] Recently, a universal numbering system has been developed and widely adopted, ImMunoGeneTics (IMGT) Information System® (Lafranc *et al.*, *Dev. Comp. Immunol.* 27(1):55-77 (2003)). IMGT is an integrated information system specializing in immunoglobulins (IG), T cell receptors (TR) and major histocompatibility complex (MHC) of human and other vertebrates. Herein, the CDRs are referred to in terms of both the amino acid sequence and the location within the light or heavy chain. As the “location” of the CDRs within the structure of the immunoglobulin variable domain is conserved between species and present in structures called loops, by using numbering systems that align variable domain sequences according to structural features, CDR and framework residues are readily identified. This information can be used in grafting and replacement of CDR residues from immunoglobulins of one species into an acceptor framework from, typically, a human antibody. An additional numbering system (AHon) has been developed by Honnegger

and Plückthun, *J. Mol. Biol.* 309: 657-670 (2001). Correspondence between the numbering system, including, for example, the Kabat numbering and the IMGT unique numbering system, is well known to one skilled in the art (see, *e.g.*, Kabat, *supra*; Chothia and Lesk, *supra*; Martin, *supra*; Lefranc *et al.*, *supra*). An Exemplary system, shown herein, combines Kabat and Chothia.

	Exemplary	IMGT	Kabat	AbM	Chothia	Contact
V _H CDR1	26-35	27-38	31-35	26-35	26-32	30-35
V _H CDR2	50-65	56-65	50-65	50-58	53-55	47-58
V _H CDR3	95-102	105-117	95-102	95-102	96-101	93-101
V _L CDR1	24-34	27-38	24-34	24-34	26-32	30-36
V _L CDR2	50-56	56-65	50-56	50-56	50-52	46-55
V _L CDR3	89-97	105-117	89-97	89-97	91-96	89-96

[0087] Hypervariable regions can comprise “extended hypervariable regions” as follows: 24-36 or 24-34 (LCDR1), 46-56 or 50-56 (LCDR2) and 89-97 or 89-96 (LCDR3) in the VL and 26-35 or 26-35A (HCDR1), 50-65 or 49-65 (HCDR2) and 93-102, 94-102, or 95-102 (HCDR3) in the VH. CDR sequences, reflecting each of the above numbering schemes, are provided herein, including in the tables in the Examples below, including Tables 4-12 and 15-20.

[0088] The term “constant region” or “constant domain” refers to a carboxy terminal portion of the light and heavy chain which is not directly involved in binding of the antibody to antigen but exhibits various effector function, such as interaction with the Fc receptor. The terms refer to the portion of an immunoglobulin molecule having a more conserved amino acid sequence relative to the other portion of the immunoglobulin, the variable region, which contains the antigen binding site. The constant region can contain the CH1, CH2 and CH3 regions of the heavy chain and the CL region of the light chain.

[0089] The term “framework” or “FR” residues are those variable region residues flanking the CDRs. FR residues are present, for example, in chimeric, humanized, human, domain antibodies, diabodies, linear antibodies, and bispecific antibodies. FR residues are those variable domain residues other than the hypervariable region residues or CDR residues.

[0090] As used herein, the term an “isolated antibody” refers to an antibody, which is substantially free of other antibodies having different antigenic specificities (*e.g.*, an isolated

antibody that specifically binds to PSMA is substantially free of antibodies that do not bind to PSMA). In addition, an isolated antibody is substantially free of other cellular material and/or chemicals. In the case of bispecific PSMA $\times$ CD3 antibodies, the bispecific antibody specifically binds both PSMA and CD3, and is substantially free of antibodies that specifically bind antigens other than PSMA and CD3. "Isolated antibody" encompasses antibodies that are isolated to a higher purity, such as antibodies that are 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% pure.

[0091] As used herein, the term "monoclonal antibody" refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that can be present in minor amounts. Monoclonal antibodies provided herein can be made by the hybridoma method, phage display technology, single lymphocyte gene cloning technology, or by recombinant DNA methods. For example, the monoclonal antibodies can be produced by a hybridoma, which includes a B cell obtained from a transgenic nonhuman animal, such as a transgenic mouse or rat, having a genome comprising a human heavy chain transgene and a light chain transgene.

[0092] As used herein, the term "antigen-binding fragment" refers to an antibody fragment such as, for example, a diabody, a Fab, a Fab', a F(ab')₂, an Fv fragment, a disulfide stabilized Fv fragment (dsFv), a (dsFv)₂, a bispecific dsFv (dsFv-dsFv'), a disulfide stabilized diabody (ds diabody), a single-chain antibody molecule (scFv), a single domain antibody (sdAb) an scFv dimer (bivalent diabody), a multispecific antibody formed from a portion of an antibody comprising one or more CDRs, a camelized single domain antibody, a nanobody, a domain antibody, a bivalent domain antibody, or any other antibody fragment that binds to an antigen but does not comprise a complete antibody structure. An antigen-binding fragment is capable of binding to the same antigen to which the parent antibody or a parent antibody fragment binds. According to particular embodiments, the antigen-binding fragment comprises a light chain variable region, a light chain constant region, and an Fd segment of the heavy chain. According to other particular embodiments, the antigen-binding fragment comprises Fab and F(ab').

[0093] As used herein, the term "single-chain antibody" refers to a conventional single-chain antibody in the field, which comprises a heavy chain variable region and a light chain variable region connected by a short peptide of about 15 to about 20 amino acids. As used

herein, the term “single domain antibody” refers to a conventional single domain antibody in the field, which comprises a heavy chain variable region and a heavy chain constant region or which comprises only a heavy chain variable region.

[0094] As used herein, the term “human antibody” refers to an antibody produced by a human or an antibody having an amino acid sequence corresponding to an antibody produced by a human made using any technique known in the art. This definition of a human antibody includes intact or full-length antibodies, fragments thereof, and/or antibodies comprising at least one human heavy and/or light chain polypeptide. If the antibody contains a constant region or a portion of the constant region, the constant region also is derived from sequences of human origin.

[0095] Human antibody comprises heavy or light chain variable regions that are “derived from” sequences of human origin if the variable regions of the antibody are obtained from a system that uses human germline immunoglobulin or rearranged immunoglobulin genes. Such exemplary systems are human immunoglobulin gene libraries displayed on phage, and transgenic non-human animals such as mice or rats carrying human immunoglobulin loci as described herein. “Human antibody” can contain amino acid differences when compared to the human germline immunoglobulin or rearranged immunoglobulin genes due to for example naturally occurring somatic mutations or intentional introduction of substitutions into the framework or antigen binding site, or both. Typically, “human antibody” is at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% identical in amino acid sequence to an amino acid sequence encoded by human germline immunoglobulin or rearranged immunoglobulin genes. In some cases, “human antibody” can contain consensus framework sequences derived from human framework sequence analyses, for example as described in Knappik *et al.*, (2000) J Mol Biol 296:57-86, or synthetic HCDR3 incorporated into human immunoglobulin gene libraries displayed on phage, for example as described in Shi *et al.*, (2010) J Mol Biol 397:385-96, and in Int. Patent Publ. No. WO2009/085462. Human antibodies derived from human immunoglobulin sequences can be generated using systems such as phage display incorporating synthetic CDRs and/or synthetic frameworks, or can be subjected to *in vitro* mutagenesis to improve antibody properties, resulting in antibodies that are not expressed by the human antibody germline repertoire *in vivo*. Antibodies in which antigen binding sites are derived from a non-human species are not included in the definition of “human antibody”.

[0096] As used herein, the term “humanized antibody” refers to a non-human antibody that is modified to increase the sequence homology to that of a human antibody, such that the antigen-binding properties of the antibody are retained, but its antigenicity in the human body is reduced. Humanized antibody includes an antibody in which the antigen binding sites are derived from non-human species and the variable region frameworks are derived from human immunoglobulin sequences. Humanized antibody can include substitutions in the framework so that the framework may not be an exact copy of expressed human immunoglobulin or human immunoglobulin germline gene sequences.

[0097] As used herein, the term “chimeric antibody” refers to an antibody wherein the amino acid sequence of the immunoglobulin molecule is derived from two or more species. The variable region of both the light and heavy chains often corresponds to the variable region of an antibody derived from one species of mammal (*e.g.*, mouse, rat, rabbit, *etc.*) having the desired specificity, affinity, and capability, while the constant regions correspond to the sequences of an antibody derived from another species of mammal (*e.g.*, human) to avoid eliciting an immune response in that species.

[0098] “Recombinant” refers to DNA, antibodies and other proteins that are prepared, expressed, created or isolated by recombinant means when segments from different sources are joined to produce recombinant DNA, antibodies or proteins.

[0099] “Epitope” refers to a portion of an antigen to which an antibody specifically binds. Epitopes typically consist of chemically active (such as polar, non-polar or hydrophobic) surface groupings of moieties such as amino acids or polysaccharide side chains and can have specific three-dimensional structural characteristics, as well as specific charge characteristics. An epitope can be composed of contiguous and/or discontinuous amino acids that form a conformational spatial unit. For a discontinuous epitope, amino acids from differing portions of the linear sequence of the antigen come in close proximity in 3-dimensional space through the folding of the protein molecule. Antibody “epitope” depends on the methodology used to identify the epitope.

[00100] “Paratope” refers to a portion of an antibody to which an antigen specifically binds. A paratope can be linear in nature or can be discontinuous, formed by a spatial relationship between non-contiguous amino acids of an antibody rather than a linear series of amino acids. A “light chain paratope” and a “heavy chain paratope” or “light chain paratope amino acid residues” and “heavy chain paratope amino acid residues” refer to

antibody light chain and heavy chain residues in contact with an antigen, respectively, or in general, “antibody paratope residues” refer to those antibody amino acids that are in contact with antigen.

[00101] “Anti-idiotypic (anti-Id) antibody” is an antibody, which recognizes the antigenic determinants (*e.g.* the paratope or CDRs) of the antibody. It is generally known in the art the process of producing or preparing an anti-idiotypic antibody. (Lathey, J. et al Immunology 1986 57(1):29-35). The anti-Id antibody can be antigen-blocking or non-blocking. The antigen-blocking anti-Id antibody can be used to detect the free antibody in a sample (*e.g.* anti-PSMA, anti-CD3 or the bispecific PSMAxCD3 antibody provided herein). The non-blocking anti-Id antibody can be used to detect the total antibody (free, partially bound to antigen, or fully bound to antigen) in a sample. An anti-Id antibody can be prepared by immunizing an animal with the antibody to which an anti-Id antibody is being prepared. In some embodiments described herein, the anti-idiotypic antibody is used for detecting the level of the therapeutic antibodies (*e.g.* anti-PSMA, anti-CD3 or the bispecific PSMAxCD3 antibody provided herein) in a sample. An anti-Id antibody can also be used as an immunogen to induce an immune response in yet another animal, producing a so-called anti-anti-Id antibody. An anti-anti-Id can be epitopically identical to the original mAb, which induced the anti-Id antibody. Thus, by using antibodies to the idiotypic determinants of a mAb, it is possible to identify other clones expressing antibodies of identical specificity. Anti-Id antibodies can be varied (thereby producing anti-Id antibody variants) and/or derivatized by any suitable technique, such as those described elsewhere herein with respect to the antibodies specifically binding PSMA or CD3, or the bispecific PSMAxCD3 antibodies.

[00102] As used herein, the term “multispecific antibody” refers to an antibody that comprises a plurality of immunoglobulin variable domain sequences, wherein a first immunoglobulin variable domain sequence of the plurality has binding specificity for a first epitope and a second immunoglobulin variable domain sequence of the plurality has binding specificity for a second epitope. In an embodiment, the first and second epitopes do not overlap or do not substantially overlap. In an embodiment, the first and second epitopes are on different antigens, *e.g.*, the different proteins (or different subunits of a multimeric protein). In an embodiment, a multispecific antibody comprises a third, fourth or fifth immunoglobulin variable domain. In an embodiment, a multispecific antibody is a

bispecific antibody molecule, a trispecific antibody molecule, or a tetraspecific antibody molecule.

[00103] As used herein, the term “bispecific antibody” refers to a multispecific antibody that binds no more than two epitopes or two antigens. A bispecific antibody is characterized by a first immunoglobulin variable domain sequence which has binding specificity for a first epitope (*e.g.*, an epitope on a PSMA antigen) and a second immunoglobulin variable domain sequence that has binding specificity for a second epitope. In an embodiment, the first and second epitopes are on different antigens, *e.g.*, the different proteins (or different subunits of a multimeric protein). In an embodiment, a bispecific antibody comprises a heavy chain variable domain sequence and a light chain variable domain sequence which have binding specificity for a first epitope and a heavy chain variable domain sequence and a light chain variable domain sequence which have binding specificity for a second epitope. In an embodiment, a bispecific antibody comprises a half antibody, or fragment thereof, having binding specificity for a first epitope and a half antibody, or fragment thereof, having binding specificity for a second epitope. In an embodiment, a bispecific antibody comprises a scFv, or fragment thereof, having binding specificity for a first epitope, and a scFv, or fragment thereof, having binding specificity for a second epitope. In an embodiment, the first epitope is located on PSMA and the second epitope is located on CD3.

[00104] As used herein, the term “prostate-specific membrane antigen” or “PSMA” refers to a type II membrane protein expressed on certain cells. The amino acid sequence of the *Pan troglodytes* (also referred to as chimpanzee or chimp) PSMA is shown in SEQ ID NO:1416 (H2Q3K5_PANTR). The extracellular domain spans residues 44 - 750, the transmembrane domain spans residues 20 - 43 and the cytoplasmic domain spans residues 1 - 19 of SEQ ID NO:1416.

MWNLHETDSAVATARRPRWLCAGALVLAGGFLLGFLFGWFIKSSNEATNITPKHNMKAFLDELKA
ENIKKFLYNFTQIPHLAGTEQNFQIAKQIQSQWKEFGLDSVELAHYDVLLSYPNKTHPNYISIIINED
GNEIFNTSLFEPPPPGYENVLDIVPPFSAFSPQGMPEGLVYVNYARTEDFFKLERDMKINCSCGKIV
IARYGKVFRGNKVKNAQLAGAKGVILYSDPADYFAPGVKSYPDGWNLPGGGVQRGNI LNLNGAGDPL
TPGYPANNEYAYRHGIAEAVGLPSIPVHPIGYYDAQKLLKMGGSAPPDSSWRGSLKVPYNVGPFGFTG
NFSTQKVKMHIHSTNEVTIRIYNVIGTLRGAVEPDYVILGGHRDSWVFGGIDPQSGAAVVHEIVRSF
GTLKKEGWRPRRTILFASWDAEEFGLLGSTEWAEENSRLQERGVAYINADSSIEGNYTLRVDCTPL
MYSLVYNLTKEKLSPDEGFEGKSLYESWTKKSPSPFSGMPRISKLKSGNDFEVFFQRLGIASGRAR
YTKNWETNKFSGYPLYHSVYETVELVEKFYDPMFKYHLTVQVRGGMVFELANSIVLPFDCRDYAVV
LRKYADKIYNI SMKHPQEMKTYSVSFDLSFAVKNFTEIASKFTERLQDFDKSNPILLRMMNDQLMF
LERAFIDPLGLPDRPFYRHVIYAPSSHKNKYAGESFPGIYDALFDIESKVDPSKAWGDVKRQISVAAF
TVQAAAETLSEVA (SEQ ID NO:1416)

[00105] The amino acid sequence of the *Macaca fascicularis* (also referred to as cynomolgus monkey, macaque or cyno) PSMA is shown in SEQ ID NO:1417 (EHH56646.1). The extracellular domain spans residues 44 - 750, the transmembrane domain spans residues 20 - 43 and the cytoplasmic domain spans residues 1 - 19 of SEQ ID NO:1417.

```
MWNLLHETDSAVATARRPRWLCAGALVLAGGFFLLGFLFGWFIKSSSEATNITPKHNMKAFLDELKA
ENIKKFLHNFTQIPHLAGTEQNFQIAKQIQSQWKEFGLDSVELTHYDVLLSYPNKTHPNYISINED
GNEIFNTSLFEPPPPAGYENVSDIVPPFSAFSPQGMPEGDLVYVNYARTEDFFKLERDMKINC SGKIV
IARYGKVFGRGNKVNAQLAGATGVILYSDPDYFAPGVKSYPDGWNLPGGGVQQRGNI LNLNGAGDPL
TPGYPANNEYAYRRGMAEAVGLPSIPVHPIGYYDAQKLEKMGGSASPDSSWRGSLKVPYNVGPGFTG
NFSTQKVKMHIHSTSEVTRIYNVIGTLRGAVEPDYVILGGHRDSWVFGGIDPQSGAAVVEIVRSF
GMLKKEGWRRPRTILFASWDAEEFGLLGSTEWAEENSRLQERGVAYINADSSIEGNYTLRVDTPL
MYSLVYNLTKELESPDEGFEGKSLYESWTKKSPSEFSGMPRISKLGSNDFEVFFQRLGIASGRAR
YTKNWETNKFSSYPLYHSVYETVELVEKFYDPMFKYHLTVAQVRGGMVFELANSVVLFPDCRDYAVV
LRKYADKIYNI SMKHPQEMKTYSVSFDLSFAVKNFTEIASKFSERLRDFDKSNPILLRMMNDQLMF
LERAFIDPLGLPDRPFYRHVIYAPSSH NKYAGESFPGIYDALFDIESKVDPSQAWGEVKRQISIATF
TVQAAAETLSEVA (SEQ ID NO:1417)
```

[00106] The amino acid sequence of the human PSMA is shown in SEQ ID NO:1418. The extracellular domain spans residues 44 - 750, the transmembrane domain spans residues 20 - 43 and the cytoplasmic domain spans residues 1 - 19 of SEQ ID NO:1418.

```
MWNLLHETDSAVATARRPRWLCAGALVLAGGFFLLGFLFGWFIKSSNEATNITPKHNMKAFLDELKA
ENIKKFLYNFTQIPHLAGTEQNFQIAKQIQSQWKEFGLDSVELAHYDVLLSYPNKTHPNYISINED
GNEIFNTSLFEPPPPAGYENVSDIVPPFSAFSPQGMPEGDLVYVNYARTEDFFKLERDMKINC SGKIV
IARYGKVFGRGNKVNAQLAGAKGVILYSDPADYFAPGVKSYPDGWNLPGGGVQQRGNI LNLNGAGDPL
TPGYPANNEYAYRRGIAEAVGLPSIPVHPIGYYDAQKLEKMGGSAPPDSSWRGSLKVPYNVGPGFTG
NFSTQKVKMHIHSTNEVTRIYNVIGTLRGAVEPDYVILGGHRDSWVFGGIDPQSGAAVVEIVRSF
GTLKKEGWRRPRTILFASWDAEEFGLLGSTEWAEENSRLQERGVAYINADSSIEGNYTLRVDTPL
MYSLVHNLTKELKSPDEGFEGKSLYESWTKKSPSEFSGMPRISKLGSNDFEVFFQRLGIASGRAR
YTKNWETNKFSGYPLYHSVYETVELVEKFYDPMFKYHLTVAQVRGGMVFELANSIVLFPDCRDYAVV
LRKYADKIYNI SMKHPQEMKTYSVSFDLSFAVKNFTEIASKFSERLQDFDKSNPIVLRMMNDQLMF
LERAFIDPLGLPDRPFYRHVIYAPSSH NKYAGESFPGIYDALFDIESKVDPSKAWGEVKRQIYVAAF
TVQAAAETLSEVA (SEQ ID NO:1418)
```

[00107] The term “PSMA” includes any PSMA variant, isoform, and species homolog, which is naturally expressed by cells (including prostate cells) or can be expressed on cells transfected with genes or cDNA encoding the polypeptide. In specific embodiments, the PSMA is a human PSMA.

[00108] As used herein, the term “CD3” refers to an antigen that is expressed on T cells as part of the multimeric T cell receptor (TCR) complex and which consists of a homodimer or heterodimer formed from the association of two or four receptor chains: CD3 epsilon, CD3 delta, CD3 zeta and CD3 gamma. In some embodiments, CD3 antibodies provided herein bind to the CD3-epsilon polypeptide, which together with CD3-gamma, -delta and -zeta, and the T cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3

complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The CD3 complex mediates signal transduction, resulting in T cell activation and proliferation. CD3 is required for the immune response. The term “CD3” includes any CD3 variant, isoform, and species homolog, which is naturally expressed by cells (including T cells) or can be expressed on cells transfected with genes or cDNA encoding the polypeptide. In specific embodiments, the CD3 is a human CD3. All references to proteins, polypeptides and protein fragments herein are intended to refer to the human version of the respective protein, polypeptide or protein fragment unless explicitly specified as being from a non-human species. Thus, “CD3” means human CD3 unless specified as being from a non-human species, e.g., “mouse CD3” “monkey CD3,” *etc.*

[00109] An exemplary human CD3 epsilon comprises the amino acid sequence of SEQ ID NO:1021.

MQSGTHWRVLGLCLLSVGWVGQDGNEEMGGITQTPYKVSISGTTVILTCPQYPGSEILWQHNDKNIG
GDEDDKNIGSDEDHLSLKEFSELEQSGYYVCYPRGSKPEDANFYLYLRARVCENCMEMDMVSVATIV
IVDICITGGLLLLVYWSKNRKAKAPVTRGAGAGGRQGRQNKERPPVPNPDIPIRKGRDLYSG
LNQRRI (SEQ ID NO:1021)

[00110] An exemplary extracellular domain of a human CD3 epsilon comprises the amino acid sequence of SEQ ID NO:1022.

DGNEEMGGITQTPYKVSISGTTVILTCPQYPGSEILWQHNDKNIGGDEDDKNIGSDEDHLSLKEFSE
LEQSGYYVCYPRGSKPEDANFYLYLRARVCENCMEMD (SEQ ID NO:1022)

[00111] “Specific binding” or “specifically binds” or “binds” refers to an antibody binding to an antigen or an epitope within the antigen with greater affinity than for other antigens. Typically, the antibody binds to the antigen or the epitope within the antigen with an equilibrium dissociation constant (K_D) of about 1×10^{-7} M or less, about 5×10^{-8} M or less, about 1×10^{-8} M or less, or about 5×10^{-8} M or less, for example about 1×10^{-9} M or less, about 1×10^{-10} M or less, about 1×10^{-11} M or less, or about 1×10^{-12} M or less, typically with the K_D that is at least one hundred fold less than its K_D for binding to a non-specific antigen (e.g., BSA, casein). The dissociation constant can be measured using standard procedures. Antibodies that specifically bind to the antigen or the epitope within the antigen can, however, have cross-reactivity to other related antigens, for example to the same antigen from other species (homologs), such as human or monkey, for example *Macaca fascicularis* (cynomolgus, cyno) or *Pan troglodytes* (chimpanzee, chimp). While a monospecific

antibody specifically binds one antigen or one epitope, a bispecific antibody specifically binds two distinct antigens or two distinct epitopes.

[00112] Throughout the specification, “CD3-specific” or “specifically binds CD3” or “anti-CD3 antibody” refers to antibodies that bind specifically to the CD3-epsilon polypeptide (SEQ ID NO:1021), including antibodies that bind specifically to the CD3-epsilon extracellular domain (ECD) (SEQ ID NO:1022). CD3-epsilon, together with CD3-gamma, -delta and -zeta, and the T cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The CD3 complex mediates signal transduction, resulting in T cell activation and proliferation. CD3 is required for the immune response.

[00113] The term “KD” refers to the dissociation constant, which is obtained from the ratio of K_d to K_a (*i.e.*, K_d/K_a) and is expressed as a molar concentration (M). KD values for antibodies can be determined using methods in the art in view of the present disclosure. For example, the KD of an antibody can be determined by using surface plasmon resonance, such as by using a biosensor system, *e.g.*, a Biacore® system, or by using bio-layer interferometry technology, such as an Octet RED96 system. The smaller the value of the KD of an antibody, the higher affinity that the antibody binds to a target antigen.

[00114] As used herein, “Tagg” refers to the temperature at which the protein starts to aggregate either through dimerization or oligomerization. The aggregation temperature detects the onset of aggregation, the temperature at which a protein will show a tendency to aggregate. Tagg can be determined by differential scanning calorimetry (DSC), Differential Scanning Fluorimetry (DSF) or by circular dichroism (CD). These techniques can detect small changes in the conformation of the protein and therefore detect the starting point of aggregation. Tagg values can be lower or higher than T_m . In cases where Tagg is lower than T_m , the protein either dimerizes and/or oligomerizes first and then starts unfolding later at higher temperatures than the Tagg. In cases where Tagg is higher than T_m , the protein starts to unfold first and then aggregates at a higher temperature than the T_m . Both events are commonly observed and depend on amino acid composition and protein conformation.

[00115] As used herein, “ T_m ” or “mid-point temperature” is the temperature midpoint of a thermal unfolding curve. It refers to the temperature where 50% of the amino acid sequence is in its native conformation and the other 50% is denatured. A thermal unfolding

curve is typically plotted as a function of temperature. T_m is used to measure protein stability. In general, a higher T_m is an indication of a more stable protein. The T_m can be readily determined using methods well known to those skilled in the art such as Circular Dichroism Spectroscopy, Differential Scanning Calorimetry, Differential Scanning Fluorimetry (both intrinsic and extrinsic dye based), UV spectroscopy, FT-IR and Isothermal Calorimetry (ITC).

[00116] In one aspect, provided herein is an antibody that binds to PSMA. In some embodiments, the antibody comprises a heavy chain variable region and a light chain variable region. In some embodiments, the PSMA antibody is a humanized antibody. In some embodiments, the PSMA antibody is a human antibody.

[00117] In certain embodiments, provided herein is a PSMA antibody comprising a VH region, VL region, VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and/or VL CDR3 of any one of the antibodies described herein. In some embodiments, provided herein is a PSMA antibody comprising a VH region of any one of the antibodies described herein. In some embodiments, provided herein is a PSMA antibody comprising a VL region of any one of the antibodies described herein. In some embodiments, provided herein is a PSMA antibody comprising a VH region of any one of the antibodies described herein, and a VL region of any one of the antibodies described herein. In some embodiments, provided herein is a PSMA antibody comprising a VH CDR1, VH CDR2, and VH CDR3 of any one of the antibodies described herein. In some embodiments, provided herein is a PSMA antibody comprising a VL CDR1, VL CDR2, and VL CDR3 of any one of the antibodies described herein. In some embodiments, provided herein is a PSMA antibody comprising a VH CDR1, VH CDR2, and VH CDR3 of any one of the antibodies described herein; and a VL CDR1, VL CDR2, and VL CDR3 of any one of the antibodies described herein. Representative VH and VL amino acid sequences, including VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 amino acid sequences, of PSMA antibodies provided herein are provided in Tables 4-12.

[00118] In certain embodiments, provided herein is a PSMA bispecific antibody comprising a binding domain that binds to PSMA having a VH region, VL region, VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and/or VL CDR3 of any one of the antibodies described herein. In some embodiments, provided herein is a PSMA bispecific antibody comprising a binding domain that binds to PSMA having a VH region of any one of the antibodies described herein. In some embodiments, provided herein is a PSMA

bispecific antibody comprising a binding domain that binds to PSMA having a VL region of any one of the antibodies described herein. In some embodiments, provided herein is a PSMA bispecific antibody comprising a binding domain that binds to PSMA having a VH region of any one of the antibodies described herein, and a VL region of any one of the antibodies described herein. In some embodiments, provided herein is a PSMA bispecific antibody comprising a binding domain that binds to PSMA having a VH CDR1, VH CDR2, and VH CDR3 of any one of the antibodies described. In some embodiments, provided herein is a PSMA bispecific antibody comprising a binding domain that binds to PSMA having a VL CDR1, VL CDR2, and VL CDR3 of any one of the antibodies described herein. In some embodiments, provided herein is a PSMA bispecific antibody comprising a binding domain that binds to PSMA having a VH CDR1, VH CDR2, and VH CDR3 of any one of the antibodies described herein; and a VL CDR1, VL CDR2, and VL CDR3 of any one of the antibodies described herein. In certain embodiments, the PSMA antibody is a bispecific antibody. In some embodiments, the PSMA bispecific antibody further comprises a second binding domain that binds to CD3 having a VH region, VL region, VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and/or VL CDR3 of a CD3 antibody provided herein. In some embodiments, the PSMA bispecific antibody further comprises a second binding domain that binds to CD3 having a VH region of a CD3 antibody provided herein. In some embodiments, the PSMA bispecific antibody further comprises a second binding domain that binds to CD3 having a VL region of a CD3 antibody provided herein. In some embodiments, the PSMA bispecific antibody further comprises a second binding domain that binds to CD3 having a VH region of a CD3 antibody provided herein, and a VL region of a CD3 antibody provided herein. In some embodiments, the PSMA bispecific antibody further comprises a second binding domain that binds to CD3 having a VH CDR1, VH CDR2, and VH CDR3 of a CD3 antibody provided herein. In some embodiments, the PSMA bispecific antibody further comprises a second binding domain that binds to CD3 having a VL CDR1, VL CDR2, and VL CDR3 of a CD3 antibody provided herein. In some embodiments, the PSMA bispecific antibody further comprises a second binding domain that binds to CD3 having a VH CDR1, VH CDR2, and VH CDR3 of a CD3 antibody provided herein, and a VL CDR1, VL CDR2, and VL CDR3 of a CD3 antibody provided herein.

[00119] In some embodiments, the antibody specifically binds PSMA. In some embodiments, the antibody specifically binds to *Pan troglodytes* (chimpanzee, chimp)

PSMA. In other embodiments, the antibody specifically binds to *Macaca fascicularis* (cynomolgus monkey, macaque, cyno) PSMA. In yet other embodiments, the antibody specifically binds to and/or human PSMA. In some embodiments, the antibody specifically binds to *Pan troglodytes*, *Macaca fascicularis*, and human PSMA. In specific embodiments, the antibody specifically binds to both cyno and human PSMA.

[00120] In certain embodiments, the PSMA antibodies bind to the chimpanzee target antigen. In one embodiment, the antibodies bind to the human and macaque PSMA target antigens with affinities within 5-fold of each other. In other words, the difference in antibody binding is less than a multiple of 5. In this case, the identical antibody molecule can be used both for preclinical evaluation of safety, activity and/or pharmacokinetic profile of PSMA in primates and as a drug in humans. Put in other words, the same PSMA-specific molecule can be used in preclinical animal studies as well as in clinical studies in humans. This leads to highly comparable results and a much-increased predictive power of the animal studies compared to species-specific surrogate molecules. Since the PSMA domain is cross-species specific, *i.e.* reactive with the human and macaque antigens, the antibody or fragments thereof can be used both for preclinical evaluation of safety, activity and/or pharmacokinetic profile of these binding domains in primates and--in the identical form--as drug in humans.

[00121] In other embodiments, the PSMA is present on the surface of a cell.

[00122] In some embodiments, the antibody is a humanized antibody. In some embodiments, the antibody is a human antibody.

[00123] In certain embodiments, the antibody is an IgG antibody. In other embodiments, the IgG antibody is an IgG1, IgG2, IgG3, or IgG4 antibody.

[00124] In some embodiments, a multispecific PSMAxCD3 antibody provided herein is an IgG1, an IgG2, an IgG3 or an IgG4 isotype. In some embodiments, a multispecific PSMAxCD3 antibody provided herein is an IgG1 isotype. In some embodiments, a multispecific PSMAxCD3 antibody provided herein is an IgG2 isotype. In some embodiments, a multispecific PSMAxCD3 antibody provided herein is an IgG3 isotype. In some embodiments, a multispecific PSMAxCD3 antibody provided herein is an IgG4 isotype.

[00125] Immunogenicity of therapeutic antibodies can associated with increased risk of infusion reactions and decreased duration of therapeutic response (Baert *et al.*, (2003) N

Engl J Med 348:602-08). The extent to which therapeutic antibodies induce an immune response in the host can be determined in part by the allotype of the antibody (Stickler *et al.*, (2011) Genes and Immunity 12:213-21). Antibody allotype is related to amino acid sequence variations at specific locations in the constant region sequences of the antibody. The table below shows select IgG1, IgG2 and IgG4 allotypes of some embodiments.

Allotype	Amino acid residue at position of diversity (residue numbering: EU Index)							
	IgG2		IgG4		IgG1			
	189	282	309	422	214	356	358	431
G2m(n)	T	M						
G2m(n-)	P	V						
G2m(n)(n-)	T	V						
GnG4m(a)			L	R				
F1m(17)					K	E	M	A
G1m(17,1)					K	D	L	A

[00126] In some embodiments, a PSMA antibody provided herein is an G2m(n) allotype. In some embodiments, a PSMA antibody provided herein is an G2m(n-) allotype. In some embodiments, a PSMA antibody provided herein is an G2m(n)/(n-) allotype. In some embodiments, a PSMA antibody provided herein is an nG4m(a) allotype. In some embodiments, a PSMA antibody provided herein is an G1m(17) allotype. In some embodiments, a PSMA antibody provided herein is an G1m(17,1) allotype.

[00127] In some embodiments, a multispecific PSMAxCD3 antibody provided herein is an G2m(n) allotype. In some embodiments, a multispecific PSMAxCD3 antibody provided herein is an G2m(n-) allotype. In some embodiments, a multispecific PSMAxCD3 antibody provided herein is an G2m(n)/(n-) allotype. In some embodiments, a multispecific PSMAxCD3 antibody provided herein is an nG4m(a) allotype. In some embodiments, a multispecific PSMAxCD3 antibody provided herein is an G1m(17) allotype. In some embodiments, a multispecific PSMAxCD3 antibody provided herein is an G1m(17,1) allotype.

[00128] In some embodiments, the antibody is a bispecific antibody. In certain embodiments, the antibody is multivalent. In other embodiments, the antibody is capable of binding at least three antigens. In some embodiments, the antibody is capable of binding at least five antigens.

[00129] In certain embodiments, provided is a PSMA antibody that is an intact antibody. In other embodiments, provided is a PSMA antibody is an antigen binding fragment of the

PSMA antibody. In some embodiments, the antigen binding fragment of the PSMA antibody is a functional fragment.

[00130] In some embodiments, the antigen binding fragment is a diabody. In some embodiments, the antigen binding fragment is a Fab. In some embodiments, the antigen binding fragment is a Fab'. In some embodiments, the antigen binding fragment is a F(ab')₂. In some embodiments, the antigen binding fragment is a Fv fragment. In some embodiments, the antigen binding fragment is a disulfide stabilized Fv fragment (dsFv). In some embodiments, the antigen binding fragment is a (dsFv)₂. In some embodiments, the antigen binding fragment is a bispecific dsFv (dsFv-dsFv'). In some embodiments, the antigen binding fragment is a disulfide stabilized diabody (ds diabody). In some embodiments, the antigen binding fragment is a single-chain antibody molecule (scFv). In some embodiments, the antigen binding fragment is a single domain antibody (sdAb). In some embodiments, the antigen binding fragment is an scFv dimer (bivalent diabody). In some embodiments, the antigen binding fragment is a multispecific antibody formed from a portion of an antibody comprising one or more CDRs. In some embodiments, the antigen binding fragment is a camelized single domain antibody. In some embodiments, the antigen binding fragment is a nanobody. In some embodiments, the antigen binding fragment is a domain antibody. In some embodiments, the antigen binding fragment is a bivalent domain antibody. In some embodiments, the antigen binding fragment is an antibody fragment that binds to an antigen but does not comprise a complete antibody structure.

[00131] In specific embodiments, the PSMA antibody comprises a VH region and a VL region. In some embodiments, the PSMA antibody is a single chain antibody. In some embodiments, the PSMA antibody is a single domain antibody. In some embodiments, the PSMA antibody is a nanobody. In certain embodiments, the PSMA antibody is a VHH antibody. In certain embodiments, the PSMA antibody is a llama antibody. In some embodiments, the PSMA antibody is a multispecific antibody. In other embodiments, the PSMA is a bispecific antibody. In certain embodiments, the multispecific antibody comprises an antigen binding fragment of a PSMA antibody provided herein. In other embodiments, the bispecific antibody comprises an antigen binding fragment of a PSMA antibody provided herein. In some embodiments, the PSMA antibody is an agonistic antibody. In other embodiments, the PSMA antibody is an antagonistic antibody.

[00132] In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 sequences are according to the Kabat numbering system. In some

embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 sequences are according to the Chothia numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 sequences are according to the Exemplary numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 sequences are according to the Contact numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 sequences are according to the IMGT numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 sequences are according to a combination of the numbering systems provided herein. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 sequences are according to the AbM numbering system.

Exemplary sets of 6 CDRs (VH CDR1-3 and VL CDR1-3) of certain antibody embodiments are provided herein, including in the Examples and Tables 4-12 (PSMA antibodies), Tables 16-22 (CD3 antibodies), and Tables 23-28 (PSMAxCD3 antibodies) herein. Other sets of CDRs are contemplated and within the scope of the antibody embodiments provided herein.

[00133] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs: 1, 2, and 3, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs: 16, 5, and 6, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs: 7, 4, and 9, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs: 10, 11, and 12, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs: 13, 14, and 3, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs: 16, 5, and 6, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs: 19, 20, and 21, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs: 22, 23, and 24, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of

SEQ ID NOs:25, 26, and 27, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:28, 11, and 6, respectively.

In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:31; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:32. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:31. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:32. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:31, and a VL having an amino acid sequence of SEQ ID NO:32. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:33. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:34. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:33, and a light chain having an amino acid sequence of SEQ ID NO:34. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:31. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:32. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:31, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:32. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:33. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:34. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:33, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:34.

[00134] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1, 2, and 3, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:38, 39, and 40, respectively.

In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:7, 42, and 9, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:44, 45, and 46, respectively.

In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:13, 14, and 3, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:38, 39, and 40, respectively.

In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:19, 20, and 21, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:56, 57, and 58, respectively.

In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:25, 26, and 27, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:62, 45, and 40, respectively.

In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:31; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:66. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:31. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:66. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:31, and a VL having an amino acid sequence of SEQ ID NO:66. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:33. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:68. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an

amino acid sequence of SEQ ID NO:33, and a light chain having an amino acid sequence of SEQ ID NO:68. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:31. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:66. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:31, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:66. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:33. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:68. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:33, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:68.

[00135] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:69, 70, and 71, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:72, 73, and 74, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:75, 76, and 77, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:78, 79, and 80, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:81, 82, and 71, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:72, 73, and 74, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:87, 88, and 89, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:90,

91, and 92, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:93, 94, and 95, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:96, 79, and 74, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:99; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:99. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:99, and a VL having an amino acid sequence of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:101. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:102. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:101, and a light chain having an amino acid sequence of SEQ ID NO:102. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:99. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:99, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:101. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:102. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:101, and a

light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:102.

[00136] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:69, 70, and 71, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:106, 107, and 108, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:75, 76, and 77, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:112, 113, and 114, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:81, 82, and 71, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:106, 107, and 108, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:87, 88, and 89, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:124, 125, and 126, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:93, 94, and 95, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:130, 113, and 108, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:99; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:134. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:99. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:134. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:99, and a VL having an amino acid sequence of SEQ ID NO:134. In one aspect, provided herein is an antibody that binds PSMA,

comprising a heavy chain having an amino acid sequence of SEQ ID NO:101. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:136. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:101, and a light chain having an amino acid sequence of SEQ ID NO:136. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:99. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:134. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:99, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:134. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:101. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:136. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:101, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:136.

[00137] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:137, 138, and 139, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:72, 73, and 74, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:143, 144, and 145, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:78, 79, and 148, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:149, 150, and 139, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ

ID NOs:72, 73, and 74, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:155, 156, and 157, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:90, 91, and 92, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:161, 162, and 163, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:79, 165, and 74, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:167; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:167. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:167, and a VL having an amino acid sequence of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:169. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:102. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:169, and a light chain having an amino acid sequence of SEQ ID NO:102. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:167. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:167, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:169. In one

aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:102. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:169, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:102.

[00138] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:137, 138, and 139, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:106, 107, and 108, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:143, 144, and 145, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:112, 113, and 114, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:149, 150, and 139, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:106, 107, and 108, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:155, 156, and 157, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:124, 125, and 126, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:161, 162, and 163, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:130, 113, and 108, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:167; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:134. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:167. In one aspect,

provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:134. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:167, and a VL having an amino acid sequence of SEQ ID NO:134. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:169. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:136. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:169, and a light chain having an amino acid sequence of SEQ ID NO:136. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:167. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:134. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:167, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:134. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:169. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:136. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:169, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:136.

[00139] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:205, 206, and 207, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:208, 209, and 210, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:143, 212, and 213, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:214, 215,

and 216, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:217, 218, and 207, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:208, 209, and 210, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:223, 224, and 225, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:226, 227, and 228, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:229, 230, and 231, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:232, 215, and 210, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:235; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:236. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:235. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:236. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:235, and a VL having an amino acid sequence of SEQ ID NO:236. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:237. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:238. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:237, and a light chain having an amino acid sequence of SEQ ID NO:238. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:235. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:236. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an

amino acid sequence of SEQ ID NO:235, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:236. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:237. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:238. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:237, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:238.

[00140] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:205, 206, and 207, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:242, 209, and 244, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:143, 212, and 213, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:248, 215, and 250, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:217, 218, and 207, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:242, 209, and 244, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:223, 224, and 225, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:260, 227, and 262, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:229, 230, and 231, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:266, 215, and 244, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3,

respectively, of SEQ ID NO:235; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:235. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:235, and a VL having an amino acid sequence of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:237. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:272. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:237, and a light chain having an amino acid sequence of SEQ ID NO:272. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:235. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:235, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:237. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:272. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:237, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:272.

[00141] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:205, 274, and 207, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:208, 209, and 210,

respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:143, 280, and 213, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:214, 215, and 216, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:217, 286, and 207, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:208, 209, and 210, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:223, 292, and 225, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:226, 227, and 228, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:229, 298, and 231, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:232, 215, and 210, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:303; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:236. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:303. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:236. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:303, and a VL having an amino acid sequence of SEQ ID NO:236. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:305. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:238. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:305, and a light chain having an amino acid sequence of SEQ ID NO:238. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence

of SEQ ID NO:303. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:236. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:303, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:236. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:305. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:238. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:305, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:238.

[00142] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:205, 274, and 207, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:242, 209, and 244, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:143, 280, and 213, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:248, 215, and 250, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:217, 286, and 207, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:242, 209, and 244, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:223, 292, and 225, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:260, 227, and 262, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:229, 298, and 231, respectively, and

(ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:266, 215, and 244, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:303; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:303. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:303, and a VL having an amino acid sequence of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:305. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:272. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:305, and a light chain having an amino acid sequence of SEQ ID NO:272. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:303. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:303, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:305. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:272. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:305, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:272.

[00143] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:205, 342, and 343, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:208, 209, and 210, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:347, 348, and 213, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:214, 215, and 216, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:353, 354, and 343, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:208, 209, and 210, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:223, 360, and 225, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:226, 363, and 228, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:365, 366, and 367, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:232, 215, and 210, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:371; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:372. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:371. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:372. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:371, and a VL having an amino acid sequence of SEQ ID NO:372. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:373. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:374. In one aspect,

provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:373, and a light chain having an amino acid sequence of SEQ ID NO:374. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:371. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:372. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:371, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:372. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:373. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:374. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:373, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:374.

[00144] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:375, 376, and 377, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:378, 379, and 380, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:381, 382, and 383, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:384, 385, and 386, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:387, 388, and 377, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:378, 379, and 380, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:393, 394, and 395, respectively, and (ii) a

VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:396, 397, and 398, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:399, 400, and 401, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:402, 385, and 380, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:405; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:406. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:405. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:406. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:405, and a VL having an amino acid sequence of SEQ ID NO:406. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:407. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:408. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:407, and a light chain having an amino acid sequence of SEQ ID NO:408. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:405. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:406. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:405, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:406. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:407. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:408. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy

chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:407, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:408.

[00145] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:205, 206, and 411, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:242, 209, and 244, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:143, 212, and 417, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:248, 215, and 250, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:217, 218, and 411, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:242, 209, and 244, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:223, 224, and 429, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:260, 227, and 262, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:229, 230, and 435, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:266, 215, and 244, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:439; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:439. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:439, and a VL

having an amino acid sequence of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:441. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:272. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:441, and a light chain having an amino acid sequence of SEQ ID NO:272. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:439. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:439, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:270. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:441. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:272. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:441, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:272.

[00146] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:443, 444, and 445, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:446, 447, and 448, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:450, 451, and 452, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:453, 454, and 455, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:456, 457, and 458, respectively, and (ii) a VL

comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:446, 447, and 448, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:461, 462, and 463, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:464, 465, and 466, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:467, 468, and 469, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:470, 453, and 448, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:473; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:474. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:473. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:474. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:473, and a VL having an amino acid sequence of SEQ ID NO:474. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:475. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:476. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:475, and a light chain having an amino acid sequence of SEQ ID NO:476. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:473. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:474. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:473, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:474. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid

sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:475. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:476. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:475, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:476.

[00147] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:477, 478, and 479, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:480, 481, and 482, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:483, 484, and 485, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:486, 487, and 488, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:489, 490, and 479, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:480, 481, and 482, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:495, 496, and 497, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:498, 499, and 500, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:501, 502, and 503, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:504, 487, and 482, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:507; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:508. In one aspect, provided herein is an antibody that binds

PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:507. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:508. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:507, and a VL having an amino acid sequence of SEQ ID NO:508. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:509. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:510. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:509, and a light chain having an amino acid sequence of SEQ ID NO:510. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:507. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:508. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:507, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:508. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:509. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:510. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:509, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:510.

[00148] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:69, 512, and 513, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:514, 515, and 516, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:517, 518, and 519, respectively, and (ii) a VL comprising a VL

CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:520, 385, and 522, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:523, 524, and 513, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:514, 515, and 516, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:529, 530, and 531, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:532, 533, and 534, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:535, 536, and 537, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:538, 385, and 516, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:541; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:542. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:541. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:542. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:541, and a VL having an amino acid sequence of SEQ ID NO:542. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:543. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:544. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:543, and a light chain having an amino acid sequence of SEQ ID NO:544. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:541. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:542. In one aspect, provided herein is an antibody that binds

PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:541, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:542. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:543. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:544. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:543, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:544.

[00149] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:545, 546, and 547, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:548,549, and550, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:143, 518, and 553, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:554,555, and 556, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:557, 558, and 547, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:548, 549, and 550, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:563, 564, and 565, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:566, 567, and 568, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:161, 570, and 571, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:572, 555, and 550, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and

a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:575; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:576. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:575. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:576. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:575, and a VL having an amino acid sequence of SEQ ID NO:576. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:577. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:578. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:577, and a light chain having an amino acid sequence of SEQ ID NO:578. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:575. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:576. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:575, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:576. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:577. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:578. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:577, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:578.

[00150] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:69, 70, and 71, respectively, and (ii) a VL comprising a VL CDR1, VL

CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:72, 73, and 74, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:75, 76, and 587, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:78, 79, and 80, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:81, 82, and 71, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:72, 73, and 74, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:87, 88, and 89, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:90, 91, and 92, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:93, 94, and 95, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:96, 79, and 74, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:99; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:99. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:99, and a VL having an amino acid sequence of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:611. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:612. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:611, and a light chain having an amino acid sequence of SEQ ID NO:612. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence

having at least 95% identity to an amino acid sequence of SEQ ID NO:99. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:99, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:100. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:611. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:612. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:611, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:612.

[00151] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:477, 478, and 615, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:480, 481, and 482, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:483, 484, and 621, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:486, 523, and 488, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:489, 490, and 615, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:480, 481, and 482, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:495, 496, and 633, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:498, 499, and 500, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH

CDR3 having an amino acid sequence of SEQ ID NOs:501, 502, and 639, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:504, 487, and 482, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:643; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:508. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:643. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:508. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:643, and a VL having an amino acid sequence of SEQ ID NO:508. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:645. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:646. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:645, and a light chain having an amino acid sequence of SEQ ID NO:646. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:643. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:508. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:643, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:508. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:645. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:646. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:645, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:646.

[00152] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:647, 648, and 649, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:650, 549, and 652, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:653, 518, and 655, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:656, 555, and 658, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:659, 660, and 649, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:650, 549, and 652, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:665, 666, and 667, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:566, 567, and 670, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:671, 672, and 673, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:674, 555, and 652, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:677; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:678. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:677. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:678. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:677, and a VL having an amino acid sequence of SEQ ID NO:678. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:679. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:680. In one aspect,

provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:679, and a light chain having an amino acid sequence of SEQ ID NO:680. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:677. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:678. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:677, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:678. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:679. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:680. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:679, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:680.

[00153] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:443, 682, and 445, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:446, 447, and 448, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:143, 688, and 451, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:452, 453, and 454, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:455, 694, and 445, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:446, 447, and 448, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:461, 700, and 463, respectively, and (ii) a

VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:464, 465, and 466, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:467, 706, and 469, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:470, 453, and 448, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:711; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:474. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:711. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:474. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:711, and a VL having an amino acid sequence of SEQ ID NO:474. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:713. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:714. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:713, and a light chain having an amino acid sequence of SEQ ID NO:714. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:711. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:474. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:711, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:474. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:713. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:714. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy

chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:713, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:714.

[00154] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:715, 716, and 717, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:72, 719, and 720, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:721, 722, and 723, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:78, 725, and 726, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:727, 728, and 717, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:72, 719, and 720, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:733, 734, and 735, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:736, 737, and 738, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:739, 740, and 741, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:96, 725, and 720, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:745; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:746. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:745. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:746. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:745, and a VL

having an amino acid sequence of SEQ ID NO:746. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:747. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:748. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:747, and a light chain having an amino acid sequence of SEQ ID NO:748. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:745. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:746. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:745, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:746. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:747. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:748. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:747, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:748.

[00155] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:647, 750, and 751, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:752, 481, and 754, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:653, 518, and 757, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:758, 487, and 760, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:659, 762, and 751, respectively, and (ii) a VL

comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:752, 481, and 754, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:665, 768, and 769, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:498, 499, and 772, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:671, 536, and 775, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:504, 487, and 754, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:779; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:780. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:779. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:780. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:779, and a VL having an amino acid sequence of SEQ ID NO:780. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:781. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:782. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:781, and a light chain having an amino acid sequence of SEQ ID NO:782. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:779. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:780. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:779, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:780. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid

sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:781. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:782. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:781, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:782.

[00156] In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:783, 784, and 785, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:548, 549, and 788, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:653, 518, and 791, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:554, 555, and 794, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:795, 796, and 785, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:548, 549, and 788, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:801, 802, and 803, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:566, 567, and 806, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:671, 536, and 809, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:572, 555, and 788, respectively. In one aspect, provided herein is an antibody that binds PSMA, comprising: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:813; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:814. In one aspect, provided herein is an antibody that binds

PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:813. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence of SEQ ID NO:814. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence of SEQ ID NO:813, and a VL having an amino acid sequence of SEQ ID NO:814. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:815. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence of SEQ ID NO:816. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence of SEQ ID NO:815, and a light chain having an amino acid sequence of SEQ ID NO:816. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:813. In one aspect, provided herein is an antibody that binds PSMA, comprising a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:814. In one aspect, provided herein is an antibody that binds PSMA, comprising a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:813, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:814. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:815. In one aspect, provided herein is an antibody that binds PSMA, comprising a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:816. In one aspect, provided herein is an antibody that binds PSMA, comprising a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:815, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:816.

[00157] In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:31. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:99. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:167. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid

sequence of SEQ ID NO:235. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:303. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:371. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:405. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:439. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:473. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:507. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:541. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:575. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:643. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:677. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:711. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:779. In some embodiments, the isolated antibody comprises a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:813.

[00158] In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:32. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:66. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:100. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:134. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID

NO:236. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:270. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:372. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:406. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:474. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:508. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:542. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:576. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:678. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:746. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:780. In some embodiments, the isolated antibody comprises a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:814.

[00159] In some embodiments, the isolated antibody comprises a heavy chain (HC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:33. In some embodiments, the isolated antibody comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:101. In some embodiments, the isolated antibody comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:169. In some embodiments, the isolated antibody comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:237. In some embodiments, the isolated antibody comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:305. In some embodiments, the isolated antibody comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:373. In some embodiments, the isolated antibody comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:407. In some

embodiments, the isolated antibody comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:441.

[00160] In some embodiments, the isolated antibody comprises a light chain (LC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:34. In some embodiments, the isolated antibody comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:68. In some embodiments, the isolated antibody comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:102. In some embodiments, the isolated antibody comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:136. In some embodiments, the isolated antibody comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:238. In some embodiments, the isolated antibody comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:272. In some embodiments, the isolated antibody comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:374. In some embodiments, the isolated antibody comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:408.

[00161] In some embodiments, provided herein is an isolated antibody that binds PSMA comprising (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In some embodiments, provided herein is an isolated antibody that binds PSMA comprising (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:SEQ ID NOs:205, 206, and 411, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:SEQ ID NOs:242, 209, and 244, respectively. In some embodiments, provided herein is an isolated antibody that binds PSMA comprising (i) a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL having an amino acid sequence of SEQ ID NO:270. In some embodiments, provided herein is an isolated antibody that binds PSMA comprising (i) a HC having an amino acid sequence of SEQ ID NO:441; and (ii) a LC having an amino acid sequence of SEQ ID NO:272.

[00162] In another aspect, provided herein is an antibody that competes for binding to PSMA with any of the PSMA antibodies described herein. In another aspect, provided herein is an antibody that binds to the same epitope as any of the PSMA antibodies described herein. In another aspect, provided is a PSMA antibody that binds an epitope on PSMA that overlaps with the epitope on PSMA bound by a PSMA antibody described herein.

[00163] In one aspect, provided is an antibody that competes for binding to PSMA with a PSMA reference antibody. In another aspect, provided is a PSMA antibody that binds to the same PSMA epitope as a PSMA reference antibody. In another aspect, provided is a PSMA antibody that binds an epitope on PSMA that overlaps with the epitope on PSMA bound by a PSMA reference antibody.

[00164] In some embodiments, the epitope is as provided in FIG. 1. In some embodiments, the antibody binds to one or more of amino acid residues 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, or 612 of the amino acid sequences of PSMA as shown in FIG. 1.

[00165] In some embodiments, the antibody binds to amino acid residue 597 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 598 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 599 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 600 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 601 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 602 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 603 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 604 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 605 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 606 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 607 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 608 of the amino acid sequence of PSMA as provided in FIG. 1. In some

embodiments, the antibody binds to amino acid residue 609 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 610 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 611 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to amino acid residue 612 of the amino acid sequence of PSMA as provided in FIG. 1. In some embodiments, the antibody binds to one or more residues selected from 597-598 (CR) of PSMA as shown in FIG. 1. In some embodiments, the antibody binds to one or more residues selected from 593-594 (LP), 595 (F), 596 (D), 600 (Y), 601 (A), 604 (L), 606-607 (KY), 607-609 (YAD), and 610-612 (KIY) of PSMA as shown in FIG. 1.

[00166] In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:32. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:66. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134. In one embodiment, the PSMA reference antibody

comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH

CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:371; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:372. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:405; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:406. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:473; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:507; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:541; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:542. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH

CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:575; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:576. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:643; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:677; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:678. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:711; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:745; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:746. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid

sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:779; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:780. In one embodiment, the PSMA reference antibody comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:813; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:814.

[00167] In certain embodiments, provided are multispecific antibodies that specifically bind to PSMA. According to certain embodiments, a bispecific antibody can be used to engage two different therapeutic targets or perform two distinct functions. Such antibodies can be used for example to recruit an immune effector cell, *e.g.*, T- or NK-cell, towards a particular target cell. Various antibody-fragment based molecules are known and under investigation, for example for cancer therapy. A multispecific PSMA antibody provided herein can be a trispecific antibody for dual targeting of tumor cells - these are trifunctional structures that can be designed to target two different targets/epitopes on the tumor cell and with the third functionality bind with high affinity to either T cells or NK-cells. Trispecific antibodies targeting two distinct tumor epitopes and engaging T- or NK-cells lyse the tumor cells that express both targets. Such molecules can be generated by antibody formats known in the art and are fully described. (WO20151842071, WO2015158636, WO2010136172, WO2013174873). In a trispecific antibody embodiment, the multispecific antibody is specific for PSMA and a second distinct antigen on the same or another tumor cell and additionally specific for an effector cell, in particular a T cell or an NK cell.

[00168] Also provided are a PSMA X “effector antigen” bispecific antibody. In one embodiment, the effector antigen of the PSMA X “effector antigen” bispecific antibody is CD3. Certain multispecific PSMAxCD3 antibodies provided can be used in preclinical animal testing, as well as clinical studies and even in therapy in human. This is due to the identification of the PSMAxCD3 bispecific antibody, which, in addition to binding to human PSMA and human CD3, respectively, also binds to the homologs of antigens of chimpanzee and macaques. For example, PSMAxCD3 multispecific antibodies provided herein can be used as a therapeutic agent against various diseases, including, but not limited, to cancer. In

view of the above, the need to construct a surrogate target PSMAxCD3 multispecific antibody for testing in a phylogenetically distant (from humans) species disappears. As a result, the identical molecule can be used in animal preclinical testing as is intended to be administered to humans in clinical testing as well as following market approval and therapeutic drug administration. The ability to use the same molecule for preclinical animal testing as in later administration to humans virtually eliminates, or at least greatly reduces, the danger that the data obtained in preclinical animal testing have limited applicability to the human case. In short, obtaining preclinical safety data in animals using the same molecule as will actually be administered to humans does much to ensure the applicability of the data to a human-relevant scenario. In contrast, in conventional approaches using surrogate molecules, said surrogate molecules have to be molecularly adapted to the animal test system used for preclinical safety assessment. Thus, the molecule to be used in human therapy in fact differs in sequence and also likely in structure from the surrogate molecule used in preclinical testing in pharmacokinetic parameters and/or biological activity, with the consequence that data obtained in preclinical animal testing have limited applicability/transferability to the human case. The use of surrogate molecules requires the construction, production, purification and characterization of a completely new construct. This leads to additional development costs and time necessary to obtain that molecule. In sum, surrogates have to be developed separately in addition to the actual drug to be used in human therapy, so that two lines of development for two molecules have to be carried out. Therefore, a major advantage of the PSMAxCD3 multispecific antibodies provided herein exhibiting cross-species specificity described herein is that the identical molecule can be used for therapeutic agents in humans and in preclinical animal testing.

[00169] Another major advantage of the antibodies and multispecific antibodies provided herein is the applicability for preclinical testing in various primates. The behavior of a drug candidate in animals should ideally be indicative of the expected behavior of this drug candidate upon administration to humans. As a result, the data obtained from such preclinical testing should therefore generally have a highly predictive power for the human case. However, as learned from the recent Phase I clinical trial on TGN1412 (a CD28 monoclonal antibody), a drug candidate can act differently in a primate species than in humans: Whereas in preclinical testing of the antibody, no or only limited adverse effects have been observed in animal studies performed with cynomolgus monkeys, six human patients developed multiple organ failure upon administration of the antibody (Lancet 368

(2006), 2206-7). The results of these dramatic negative events suggest that it may not be sufficient to limit preclinical testing to only one (non-chimpanzee primate) species. The fact that certain antibodies and multispecific antibodies provided herein specifically bind PSMA of chimpanzee and cynomolgus monkey can help to overcome the problems faced in the case mentioned above. Accordingly, provided are means and methods for minimizing species differences in effects when drugs for human therapy are being developed and tested.

[00170] With the antibodies and multispecific antibodies provided herein it is also no longer necessary to adapt the test animal to the drug candidate intended for administration to humans, such as *e.g.*, the creation of transgenic animals. The cross-species specificity of the PSMA antibody or multispecific antibody provided herein allows the antibody to be directly used for preclinical testing in non-chimpanzee primates without any genetic manipulation of the animals. As well known to those skilled in the art, approaches in which the test animal is adapted to the drug candidate always bear the risk that the results obtained in the preclinical safety testing are less representative and predictive for humans due to the modification of the animal. For example, in transgenic animals, the proteins encoded by the transgenes are often highly over-expressed. Thus, data obtained for the biological activity of an antibody against this protein antigen can be limited in their predictive value for humans in which the protein is expressed at much lower, more physiological levels.

[00171] A further advantage of the uses of certain antibodies provided herein exhibiting cross-species specificity is the fact that the use of chimpanzees, an endangered species, can be avoided for animal testing. Chimpanzees are the closest relatives to humans and were recently grouped into the family of hominids based on the genome sequencing data (Wildman *et al.*, PNAS 100 (2003), 7181). Therefore, data obtained with chimpanzee is generally considered to be highly predictive for humans. However, due to their status as endangered species, the number of chimpanzees, which can be used for medical experiments, is highly restricted. As stated above, maintenance of chimpanzees for animal testing is therefore both costly and ethically problematic. The uses of the antibodies provided herein avoid both ethical objections and financial burden during preclinical testing without prejudicing the quality, *i.e.*, applicability, of the animal testing data obtained. In light of this, the uses of the antibody or multispecific antibody specifically binding PSMA provide for a reasonable alternative for studies in chimpanzees.

[00172] A still further advantage of certain antibodies or multispecific antibodies provided herein that bind to PSMA is the ability of extracting multiple blood samples when using it as

part of animal preclinical testing, for example in the course of pharmacokinetic animal studies. Multiple blood extractions can be much more readily obtained with a non-chimpanzee primate than with lower animals, *e.g.*, a mouse. The extraction of multiple blood samples allows continuous testing of blood parameters for the determination of the biological effects induced by the antibody or multispecific antibody specifically binding PSMA provided herein. Furthermore, the extraction of multiple blood samples enables the researcher to evaluate the pharmacokinetic profile of the antibody or multispecific antibody specifically binding PSMA provided herein as defined herein. In addition, potential side effects, which can be induced by said antibody or multispecific antibody specifically binding PSMA provided herein reflected in blood parameters can be measured in different blood samples extracted during the course of the administration of said antibody. This can allow the determination of the potential toxicity profile of the antibodies or multispecific antibodies provided herein.

[00173] In some embodiments, the PSMA antibody is PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A, PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P70_F02, PSMA_P72_G02, or PSMA_P72_A11. In some embodiments, the PSMA antibody is PSMB946 (PSMB895 with a C-terminal Lys (K) amino acid residue). In some embodiments, the PSMA antibody is PSMB947 (PSMB896 with a C-terminal Lys (K) amino acid residue). In some embodiments, the PSMA antibody is PSMB948 (PSMB897 with a C-terminal Lys (K) amino acid residue). In some embodiments, the PSMA antibody is PSMB949 (PSMB898 with a C-terminal Lys (K) amino acid residue). In certain embodiments, the PSMA antibody is PSMB889. In other embodiments, the PSMA antibody is PSMB890. In certain embodiments, the PSMA antibody is PSMB891. In certain embodiments, the PSMA antibody is PSMB892. In other embodiments, the PSMA antibody is PSMB893. In certain embodiments, the PSMA antibody is PSMB894. In other embodiments, the PSMA antibody is PSMB895. In certain embodiments, the PSMA antibody is PSMB896. In certain embodiments, the PSMA antibody is PSMB897. In other embodiments, the PSMA antibody is PSMB898. In certain embodiments, the PSMA antibody is PSMB899. In certain embodiments, the PSMA antibody is PSMHB49SC1133_011A11_1. In other embodiments, the PSMA antibody is PSMB896-

G100A. In certain embodiments, the PSMA antibody is PSMA_P72_A10-HC-G54E. In certain embodiments, the PSMA antibody is PSMA_P72_D01-HC-D95E. In other embodiments, the PSMA antibody is PSMA_P72_F01. In other embodiments, the PSMA antibody is PSMA_P75_F01. In certain embodiments, the PSMA antibody is PSMA_P72_F07. In certain embodiments, the PSMA antibody is PSMA_P72_E07. In other embodiments, the PSMA antibody is PSMA_P72_D01. In certain embodiments, the PSMA antibody is PSMA_P72_C01. In other embodiments, the PSMA antibody is PSMA_P72_A10. In certain embodiments, the PSMA antibody is PSMA_P72_F02. In certain embodiments, the PSMA antibody is PSMA_P70_F02. In other embodiments, the PSMA antibody is PSMA_P72_G02. In other embodiments, the PSMA antibody is PSMA_P72_A11. Each of these antibodies is further described in the Examples below, including Tables 4-12.

[00174] In some embodiments, provided is a PSMA antibody comprising a VH of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In some embodiments, provided is a PSMA antibody comprising a VL of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In some embodiments, provided is a PSMA antibody comprising a VH and VL of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In other embodiments, provided is a PSMA antibody comprising a VH CDR1, VH CDR2, and VH CDR3 of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In other embodiments, provided is a PSMA antibody comprising a VL CDR1, VL CDR2, and VL CDR3 of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In other embodiments, provided is a PSMA antibody comprising a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Kabat numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Chothia numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the AbM numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Contact numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences

of the PSMA antibody are according to the IMGT numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 sequences of the PSMA antibody are according to a combination of the numbering systems provided herein.

[00175] In some embodiments, the PSMA antibody comprises the VH of PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A, PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P70_F02, PSMA_P72_G02, or PSMA_P72_A11. In some embodiments, the PSMA antibody comprises the VL of PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A, PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P70_F02, PSMA_P72_G02, or PSMA_P72_A11. In some embodiments, the PSMA antibody comprises the both the VH and the VL of PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A, PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P70_F02, PSMA_P72_G02, or PSMA_P72_A11. In a specific embodiment, the VH and VL are of a single PSMA antibody clone.

[00176] In some embodiments, the PSMA antibody is a multispecific PSMA antibody, wherein the first binding domain that binds PSMA comprises the VH of PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A, PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P70_F02, PSMA_P72_G02, or PSMA_P72_A11. In some embodiments, the PSMA antibody is a multispecific PSMA antibody, wherein the first binding domain that binds PSMA comprises the VL of

PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A, PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P70_F02, PSMA_P72_G02, or PSMA_P72_A11. In some embodiments, the PSMA antibody is a multispecific PSMA antibody, wherein the first binding domain that binds PSMA comprises both the VH and VL of PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A, PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P70_F02, PSMA_P72_G02, or PSMA_P72_A11. In a specific embodiment, the VH and VL are of a single PSMA antibody clone. In certain embodiments, the multispecific PSMA antibody is a multispecific PSMAxCD3 antibody. In specific embodiments, the multispecific PSMAxCD3 antibody is a bispecific PSMAxCD3 antibody. In some embodiments, the second binding domain that binds CD3 comprises the VH of CD3B376. In some embodiments, the second binding domain that binds CD3 comprises the VL of CD3B376. In some embodiments, the second binding domain that binds CD3 comprises both the VH and VL of CD3B376. In some embodiments, the second binding domain that binds CD3 comprises the VH of CD3B450. In some embodiments, the second binding domain that binds CD3 comprises the VL of CD3B450. In some embodiments, the second binding domain that binds CD3 comprises both the VH and VL of CD3B450. In some embodiments, the second binding domain that binds CD3 comprises the VH of CD3W245. In some embodiments, the second binding domain that binds CD3 comprises the VL of CD3W245. In some embodiments, the second binding domain that binds CD3 comprises both the VH and VL of CD3W245. In some embodiments, the second binding domain that binds CD3 comprises the VH of CD3B2030. In some embodiments, the second binding domain that binds CD3 comprises the VL of CD3B2030. In some embodiments, the second binding domain that binds CD3 comprises both the VH and VL of CD3B2030.

[00177] In some embodiments, the PSMA antibody comprises the VH CDR1-3 of PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A,

PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P70_F02, PSMA_P72_G02, or PSMA_P72_A11. In some embodiments, the PSMA antibody comprises the VL CDR1-3 of PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A, PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P70_F02, PSMA_P72_G02, or PSMA_P72_A11. In some embodiments, the PSMA antibody comprises both the VH CDR1-3 and the VL CDR1-3 of PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A, PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P70_F02, PSMA_P72_G02, or PSMA_P72_A11. In a specific embodiment, the VH CDR1-3 and VL CDR1-3 are of a single PSMA antibody clone.

[00178] In some embodiments, the PSMA antibody is a multispecific PSMA antibody, wherein the first binding domain that binds PSMA comprises the VH CDR1-3 of PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A, PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P70_F02, PSMA_P72_G02, or PSMA_P72_A11. In some embodiments, the PSMA antibody is a multispecific PSMA antibody, wherein the first binding domain that binds PSMA comprises the VL CDR1-3 of PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A, PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P70_F02, PSMA_P72_G02, or PSMA_P72_A11. In some embodiments, the PSMA antibody is a multispecific PSMA antibody, wherein the first binding domain that binds PSMA comprises both the VH CDR1-

3 and VL CDR1-3 of PSMB889, PSMB890, PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898, PSMB899, PSMHB49SC1133_011A11_1, PSMB896-G100A, PSMA_P72_A10-HC-G54E, PSMA_P72_D01-HC-D95E, PSMA_P72_F01, PSMA_P75_F01, PSMA_P72_F07, PSMA_P70_F02, PSMA_P72_E07, PSMA_P72_D01, PSMA_P72_C01, PSMA_P72_A10, PSMA_P72_F02, PSMA_P72_G02, or PSMA_P72_A11. In a specific embodiment, the VH CDR1-3 and VL CDR1-3 are of a single PSMA antibody clone. In certain embodiments, the multispecific PSMA antibody is a multispecific PSMAxCD3 antibody. In specific embodiments, the multispecific PSMAxCD3 antibody is a bispecific PSMAxCD3 antibody. In some embodiments, the second binding domain that binds CD3 comprises the VH CDR1-3 of CD3B376. In some embodiments, the second binding domain that binds CD3 comprises the VL CDR1-3 of CD3B376. In some embodiments, the second binding domain that binds CD3 comprises both the VH CDR1-3 and VL CDR1-3 of CD3B376. In some embodiments, the second binding domain that binds CD3 comprises the VH CDR1-3 of CD3B450. In some embodiments, the second binding domain that binds CD3 comprises the VL CDR1-3 of CD3B450. In some embodiments, the second binding domain that binds CD3 comprises the both the VH CDR1-3 and VL CDR1-3 of CD3B450. In some embodiments, the second binding domain that binds CD3 comprises the VH CDR1-3 of CD3W245. In some embodiments, the second binding domain that binds CD3 comprises the VL CDR1-3 of CD3W245. In some embodiments, the second binding domain that binds CD3 comprises both the VH CDR1-3 and VL CDR1-3 of CD3W245. In some embodiments, the second binding domain that binds CD3 comprises the VH CDR1-3 of CD3B2030. In some embodiments, the second binding domain that binds CD3 comprises the VL CDR1-3 of CD3B2030. In some embodiments, the second binding domain that binds CD3 comprises both the VH CDR1-3 and VL CDR1-3 of CD3B2030.

[00179] In another aspect, provided herein is a multispecific antibody that binds PSMA. In some embodiments, the multispecific antibody is a bispecific antibody. In some embodiments, the multispecific antibody is a trispecific antibody. In some embodiments, the multispecific antibody is a quadraspecific antibody. In one embodiment, the multispecific PSMA antibody comprises: (a) a first binding domain that binds PSMA, and (b) a second binding domain that binds to a second target. In one embodiment, the multispecific PSMA antibody comprises: (a) a first binding domain that binds PSMA, and (b) a second binding domain that binds to a second target, and (c) a third binding domain that binds to a third target. In one embodiment, the multispecific PSMA antibody comprises: (a) a first binding

domain that binds PSMA, and (b) a second binding domain that binds to a second target, (c) a third binding domain that binds to a third target, and (d) a fourth binding domain that binds to a fourth target.

[00180] In another aspect, provided herein is a multispecific antibody comprising: (a) a first binding domain that binds to PSMA, and (b) a second binding domain that binds to a second target that is not PSMA. In another aspect, provided herein is a multispecific antibody comprising: (a) a first binding domain that binds to PSMA, and (b) a second binding domain that binds to CD3. In certain embodiments, the multispecific antibody is a bispecific antibody that comprises: (a) a first binding domain that binds to PSMA, and (b) a second binding domain that binds to CD3. Exemplary first binding domains that bind to PSMA are provided herein. Exemplary second binding domains that bind to CD3 are also provided herein. Also contemplated is any combination of (a) a first binding domain that binds PSMA provided herein, and (b) a second binding domain that binds CD3 provided herein.

[00181] In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:32. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:66. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a

VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a

VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:371; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:372. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:405; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:406. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:473; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:507; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH

comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:541; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:542. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:575; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:576. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:643; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:677; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:678. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:711; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474. In one embodiment, the first binding

domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:745; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:746. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:779; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:780. In one embodiment, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:813; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:814.

[00182] In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the first binding domain that binds PSMA are according to the Kabat numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the first binding domain that binds PSMA are according to the Chothia numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the first binding domain that binds PSMA are according to the AbM numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the first binding domain that binds PSMA are according to the Contact numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the first binding domain that binds PSMA are according to the IMGT numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the first binding domain that binds PSMA are according to a combination of the numbering systems provided herein. Exemplary sets of 6 CDRs (VH CDR1-3 and VL CDR1-3) of certain antibody embodiments are provided herein, including

in the Examples and Tables 4-12 (PSMA antibodies), Tables 16-22 (CD3 antibodies), and Tables 23-28 (PSMA $\times$ CD3 antibodies) herein. Other sets of CDRs are contemplated and within the scope of the antibody embodiments provided herein.

[00183] In some embodiments of the multispecific PSMA antibodies provided herein, the first binding domain binds a PSMA antigen. In some embodiments, the first binding domain binds a PSMA epitope. In some embodiments, the first binding domain specifically binds to PSMA. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the first binding domain form a binding site for an antigen of the PSMA. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the first binding domain form a binding site for an epitope of the PSMA. In some embodiments, the PSMA is present on the surface of a T cell.

[00184] In some embodiments of the multispecific PSMA antibodies provided herein, the second target is not a PSMA antigen. In some embodiments of the multispecific PSMA antibodies provided herein, the third target is not a PSMA antigen. In some embodiments of the multispecific PSMA antibodies provided herein, the fourth target is not a PSMA antigen. In some embodiments of the multispecific PSMA antibodies provided herein, the second target is not a PSMA antigen, and the third target is not a PSMA antigen. In some embodiments of the multispecific PSMA antibodies provided herein, the second target is not a PSMA antigen, and the fourth target is not a PSMA antigen. In some embodiments of the multispecific PSMA antibodies provided herein, the third target is not a PSMA antigen, and the fourth target is not a PSMA antigen. In some embodiments of the multispecific PSMA antibodies provided herein, the second target is not a PSMA antigen, the third target is not a PSMA antigen, and the fourth target is not a PSMA antigen. In some embodiments of the multispecific PSMA antibodies provided herein, the second target is not a PSMA epitope. In some embodiments of the multispecific PSMA antibodies provided herein, the third target is not a PSMA epitope. In some embodiments of the multispecific PSMA antibodies provided herein, the fourth target is not a PSMA epitope. In some embodiments of the multispecific PSMA antibodies provided herein, the second target is not a PSMA epitope, and the third target is not a PSMA epitope. In some embodiments of the multispecific PSMA antibodies provided herein, the second target is not a PSMA epitope, and the fourth target is not a PSMA epitope. In some embodiments of the multispecific PSMA antibodies provided herein, the third target is not a PSMA epitope, and the fourth target is not a PSMA epitope. In some embodiments of the multispecific PSMA antibodies provided herein, the second

target is not a PSMA epitope, the third target is not a PSMA epitope, and the fourth target is not a PSMA epitope.

[00185] In some embodiments of the multispecific PSMA antibodies provided herein, the second target is CD3 and the multispecific PSMA antibody comprises a second binding domain that binds to CD3.

[00186] In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:817, 818, and 819, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:820, 821, and 822, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:823, 824, and 825, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:826, 487, and 828, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:829, 830, and 819, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:820, 821, and 822, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:835, 836, and 837, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:838, 839, and 840, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:841, 842, and 843, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:844, 487, and 822, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:847; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a

VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:848. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:847. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence of SEQ ID NO:848. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:847, and a VL having an amino acid sequence of SEQ ID NO:848. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence of SEQ ID NO:849. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a light chain having an amino acid sequence of SEQ ID NO:850. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence of SEQ ID NO:849, and a light chain having an amino acid sequence of SEQ ID NO:850. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:847. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:848. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:847, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:848. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:849. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:850. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence

of SEQ ID NO:849, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:850.

[00187] In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:817, 818, and 819, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:820, 821, and 822, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:823, 824, and 825, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:826, 487, and 828, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:829, 830, and 819, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:820, 821, and 822, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:835, 836, and 837, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:838, 839, and 840, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:841, 842, and 843, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:844, 487, and 822, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:847; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:848. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:847.

In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence of SEQ ID NO:848. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:847, and a VL having an amino acid sequence of SEQ ID NO:848. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence of SEQ ID NO:883. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a light chain having an amino acid sequence of SEQ ID NO:850. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence of SEQ ID NO:883, and a light chain having an amino acid sequence of SEQ ID NO:850. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:847. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:848. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:847, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:848. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:883. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:850. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:883, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:850.

[00188] In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:817, 818, and 819, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:820, 821, and 822, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:823, 824, and 825, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:826, 487, and 828, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:829, 830, and 819, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:820, 821, and 822, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:835, 836, and 837, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:838, 907, and 840, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:841, 842, and 843, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:844, 487, and 822, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:915; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:916. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:915. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence of SEQ ID NO:916. In one embodiment of the multispecific PSMA antibodies provided herein, the

second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:915, and a VL having an amino acid sequence of SEQ ID NO:916. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence of SEQ ID NO:917. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a light chain having an amino acid sequence of SEQ ID NO:918. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence of SEQ ID NO:917, and a light chain having an amino acid sequence of SEQ ID NO:918. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:915. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:916. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:915, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:916. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:917. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:918. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:917, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:918.

[00189] In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:817, 818, and 819, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an

amino acid sequence of SEQ ID NOs:820, 821, and 822, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:823, 824, and 825, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:826, 487, and 828, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:829, 830, and 819, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:820, 821, and 822, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:835, 836, and 837, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:838, 907, and 840, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:841, 842, and 843, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:844, 487, and 822, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:915; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:916. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:915. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence of SEQ ID NO:916. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:915, and a VL having an amino acid sequence of SEQ ID NO:916. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence of SEQ ID

NO:951. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a light chain having an amino acid sequence of SEQ ID NO:918. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence of SEQ ID NO:951, and a light chain having an amino acid sequence of SEQ ID NO:918. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:915. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:916. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:915, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:916. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:951. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:918. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:951, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:918.

[00190] In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1, 954, and 955, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:956, 957, and 958, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:7, 960, and 961, respectively, and (ii) a VL comprising

a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:962, 963, and 964, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:13, 966, and 955, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:956, 957, and 958, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:19, 972, and 973, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:974, 975, and 976, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:25, 978, and 979, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:980, 963, and 958, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:983; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:984. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:983. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence of SEQ ID NO:984. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:983, and a VL having an amino acid sequence of SEQ ID NO:984. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence of SEQ ID NO:985. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a light chain having an amino acid sequence of SEQ ID NO:986. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having

an amino acid sequence of SEQ ID NO:985, and a light chain having an amino acid sequence of SEQ ID NO:986. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:983. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:984. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:983, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:984. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:985. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:986. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:985, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:986.

[00191] In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1, 954, and 955, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:956, 957, and 958, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:7, 960, and 961, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:962, 963, and 964, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID

NOs:13, 966, and 955, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:956, 957, and 958, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:19, 972, and 973, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:974, 975, and 976, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:25, 978, and 979, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:980, 963, and 958, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:983; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:984. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:983. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence of SEQ ID NO:984. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:983, and a VL having an amino acid sequence of SEQ ID NO:984. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence of SEQ ID NO:1019. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a light chain having an amino acid sequence of SEQ ID NO:986. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence of SEQ ID NO:1019, and a light chain having an amino acid sequence of SEQ ID NO:986. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID

NO:983. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:984. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:983, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:984. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:1019. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:986. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a heavy chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:1019, and a light chain having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:986.

[00192] In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1467, 1468, and 1469, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:1470, 1471, and 1472, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1473, 1474, and 1475, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:1476, 1477, and 1478, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1479, 1480, and 1481, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:1482, 1477, and 1472, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding

domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1508, 1509, and 1469, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:1470, 1471, and 1472, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1510, 1511, and 1512, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:1513, 1514, and 1515, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:1463; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:1464. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:1463. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence of SEQ ID NO:1464. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:1463, and a VL having an amino acid sequence of SEQ ID NO:1464. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:1463. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:1464. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:1463, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:1464.

[00193] In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH

CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1467, 1468, and 1506, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:1470, 1471, and 1472, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1473, 1474, and 1507, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:1476, 1477, and 1478, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1479, 1480, and 1518, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:1482, 1477, and 1472, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1508, 1509, and 1506, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:1470, 1471, and 1472, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:1516, 1511, and 1517, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:1513, 1514, and 1515, respectively. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3, comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:1505; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:1464. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence of SEQ ID NO:1505. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence of SEQ ID NO:1464. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3

comprises a VH having an amino acid sequence of SEQ ID NO:1505, and a VL having an amino acid sequence of SEQ ID NO:1464. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:1505. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:1464. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises a VH having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:1505, and a VL having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:1464.

[00194] In some embodiments, the second binding domain that binds to CD3 comprises a scFv comprising a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, a VL CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, a VL CDR3 respectively, of a scFv having an amino acid sequence of SEQ ID NO:1524. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises an scFv having an amino acid sequence of SEQ ID NO:1524. In one embodiment of the multispecific PSMA antibodies provided herein, the second binding domain that binds CD3 comprises an scFv having an amino acid sequence having at least 95% identity to an amino acid sequence of SEQ ID NO:1524.

[00195] In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises a heavy chain (HC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:33. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:101. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:169. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:237. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises

a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:305. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:373. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:407. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:441. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1242. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1244. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1248. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1250. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1252. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1254. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1256. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1258. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1260. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least

80% identical to the amino acid sequence of SEQ ID NO:1262. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1264. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1266. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1268. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1270.

[00196] In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises a light chain (LC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:34. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:68. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:102. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:136. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:238. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:272. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:374. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:408.

[00197] In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1485. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1486. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1487. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1488. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1489. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1490. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1491. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1492. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1493. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1494. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1495. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1496. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1497. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1498. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1499. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO: 1500. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence

of SEQ ID NO:1526. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO:1527. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO:1528. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO:1529. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO:1530. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence of SEQ ID NO:1531.

[00198] In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1485. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1486. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1487. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1488. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1489. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1490. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1491. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1492. In some embodiments of the multispecific antibody provided herein, the first binding domain that

binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1493. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1494. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1495. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1496. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1497. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1498. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1499. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1500. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1526. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1527. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1528. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1529. In some embodiments of the multispecific antibody provided herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1530. In some embodiments of the multispecific antibody provided

herein, the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1531.

[00199] In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises (i) a HC having an amino acid sequence of SEQ ID NO:441; and (ii) a LC having an amino acid sequence of SEQ ID NO:272.

[00200] In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the second binding domain that binds CD3 comprises a heavy chain (HC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:849. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:883. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:917. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:951. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:985. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1019. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1504. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1455. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1192. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1194. In some embodiments of the multispecific antibody provided

herein, the second binding domain that binds CD3 comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1167. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1218. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a HC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1238.

[00201] In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:850. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:918. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:986. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1193. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1195. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a LC having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1219.

[00202] In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the second binding domain that binds CD3 comprises a scFv having an amino acid sequence of SEQ ID NO:1186. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a scFv having an amino acid sequence of SEQ ID NO:1187. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a scFv having an amino acid sequence of SEQ ID NO:1523. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a scFv having an amino acid sequence of SEQ ID NO:1524. In some embodiments of the multispecific antibody provided herein,

the second binding domain that binds CD3 comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1186. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1187. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1523. In some embodiments of the multispecific antibody provided herein, the second binding domain that binds CD3 comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1524.

[00203] Provided herein is an isolated bispecific antibody comprising a first binding domain that binds PSMA and a second binding domain that binds CD3, wherein the first binding domain that binds PSMA comprises (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270, and the second binding domain that binds CD3 comprises a scFv comprising a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, and a VL CDR3 of a scFv having an amino acid sequence of SEQ ID NO:1524. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:205, 206, 411, 242, 209 and 244, respectively, wherein the amino acid sequences are according to the Kabat numbering system; and the second binding domain that binds CD3 comprises a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:1467, 1468, 1506, 1470, 1471 and 1472, respectively; wherein the amino acid sequences are according to the Kabat numbering system. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises (i) a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL having an amino acid sequence of SEQ ID NO:270, and the second binding domain that binds CD3 comprises a scFv of SEQ ID NO:1524, respectively. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises (i) a HC2 having an amino acid sequence of

SEQ ID NO:441; and (ii) a LC2 having an amino acid sequence of SEQ ID NO:272, respectively; and the second binding domain that binds CD3 comprises a HC1 of SEQ ID NO:1455.

[00204] Provided herein is an isolated bispecific antibody comprising a first binding domain that binds PSMA and a second binding domain that binds CD3, wherein the first binding domain that binds PSMA comprises (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270, and the second binding domain that binds CD3 comprises a scFv comprising a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, and a VL CDR3 of a scFv having an amino acid sequence of SEQ ID NO:1286. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:205, 206, 411, 242, 209 and 244, respectively, wherein the amino acid sequences are according to the Kabat numbering system; and the second binding domain that binds CD3 comprises a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:1, 954, 955, 956, 957 and 958, respectively; wherein the amino acid sequences are according to the Kabat numbering system. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises (i) a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL having an amino acid sequence of SEQ ID NO:270, and the second binding domain that binds CD3 comprises a scFv of SEQ ID NO:1186, respectively. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises (i) a HC2 having an amino acid sequence of SEQ ID NO:441; and (ii) a LC2 having an amino acid sequence of SEQ ID NO:272, respectively; and the second binding domain that binds CD3 comprises a HC1 of SEQ ID NO:1192.

[00205] Provided herein is an isolated bispecific antibody comprising a first binding domain that binds PSMA and a second binding domain that binds CD3, wherein the first binding domain that binds PSMA comprises (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:405; and (ii) a

VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:406, and the second binding domain that binds CD3 comprises a scFv comprising a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, and a VL CDR3 of a scFv having an amino acid sequence of SEQ ID NO:1523. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:375, 376, 377, 378, 379 and 380, respectively, wherein the amino acid sequences are according to the Kabat numbering system; and the second binding domain that binds CD3 comprises a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:1467, 1468, 1469, 1470, 1471 and 1472, respectively; wherein the amino acid sequences are according to the Kabat numbering system. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises (i) a VH having an amino acid sequence of SEQ ID NO:405; and (ii) a VL having an amino acid sequence of SEQ ID NO:406, and the second binding domain that binds CD3 comprises a scFv of SEQ ID NO:1523, respectively. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises (i) a HC2 having an amino acid sequence of SEQ ID NO:407; and (ii) a LC2 having an amino acid sequence of SEQ ID NO:408, respectively; and the second binding domain that binds CD3 comprises a HC1 of SEQ ID NO:1504.

[00206] Provided herein is an isolated bispecific antibody comprising a first binding domain that binds PSMA and a second binding domain that binds CD3, wherein the first binding domain that binds PSMA comprises (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270, and the second binding domain that binds CD3 comprises (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:847; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3,

respectively, of a VL having an amino acid sequence of SEQ ID NO:848. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:205, 206, 411, 242, 209 and 244, respectively, wherein the amino acid sequences are according to the Kabat numbering system; and the second binding domain that binds CD3 comprises a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:817, 818, 819, 820, 821 and 822, respectively; wherein the amino acid sequences are according to the Kabat numbering system. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises (i) a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL having an amino acid sequence of SEQ ID NO:270, and the second binding domain that binds CD3 comprises (i) a VH having an amino acid sequence of SEQ ID NO:847; and (ii) a VL having an amino acid sequence of SEQ ID NO:848, respectively. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises (i) a HC2 having an amino acid sequence of SEQ ID NO:1244; and (ii) a LC2 having an amino acid sequence of SEQ ID NO:272, respectively; and the second binding domain that binds CD3 comprises a (i) HC1 having an amino acid sequence of SEQ ID NO:849; and (ii) LC1 having an amino acid sequence of SEQ ID NO:850. In some embodiments of the multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds PSMA comprises (i) a HC2 having an amino acid sequence of SEQ ID NO:1244; and (ii) a LC2 having an amino acid sequence of SEQ ID NO:272, respectively; and the second binding domain that binds CD3 comprises a (i) HC1 having an amino acid sequence of SEQ ID NO:883; and (ii) LC1 having an amino acid sequence of SEQ ID NO:850.

[00207] Exemplary PSMA antibody embodiments are provided herein, including in the Examples and Tables 4-12. Exemplary CD3 antibody embodiments are provided herein, including in the Examples and Tables 16-22. Exemplary PSMAxCD3 antibody embodiments are provided herein, including in the Examples and Tables 23-28.

[00208] In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMB889. In other embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMB890. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMB891. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMB892. In other

embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMB893. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMB894. In other embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMB895. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMB896. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMB897. In other embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMB898. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMB899. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMHB49SC1133_011A11_1. In other embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMB896-G100A. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P72_A10-HC-G54E. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P72_D01-HC-D95E. In other embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P72_F01. In other embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P75_F01. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P72_F07. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P72_E07. In other embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P72_D01. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P72_C01. In other embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P72_A10. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P72_F02. In certain embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P70_F02. In other embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P72_G02. In other embodiments, the multispecific PSMA antibody comprises a first binding domain of PSMA_P72_A11. Each of these antibodies is further described in the Examples below. In a specific embodiment, the multispecific PSMA antibody is a multispecific PSMAxCD3 antibody.

[00209] In some embodiments, provided is a PSMA multispecific antibody comprising a first binding domain that comprises a VH of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In some embodiments, provided is a PSMA multispecific antibody

comprising a first binding domain that comprises a VL of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In some embodiments, provided is a PSMA multispecific antibody comprising a first binding domain that comprises a VH and VL of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In other embodiments, provided is a PSMA multispecific antibody comprising a first binding domain that comprises a VH CDR1, VH CDR3 and VH CDR3 of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In other embodiments, provided is a PSMA multispecific antibody comprising a first binding domain that comprises a VL CDR1, VL CDR3 and VL CDR3 of any of the PSMA antibodies provided Tables 4-12 or 23-28. In other embodiments, provided is a PSMA multispecific antibody comprising a first binding domain that comprises a VH CDR1, VH CDR3, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Kabat numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Chothia numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the AbM numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Contact numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the IMGT numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences pf the PSMA antibody are according to a combination of the numbering systems provided herein.

[00210] In some embodiments, provided is a multispecific PSMAxCD3 antibody comprising a first binding domain that comprises a VH of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In some embodiments, provided is a multispecific PSMAxCD3 antibody comprising a first binding domain that comprises a VL of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In some embodiments, provided is a multispecific PSMAxCD3 antibody comprising a first binding domain that comprises a VH and VL of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In other embodiments, provided is a multispecific PSMAxCD3 antibody comprising a first binding

domain that comprises a VH CDR1, VH CDR3 and VH CDR3 of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In other embodiments, provided is a multispecific PSMAxCD3 antibody comprising a first binding domain that comprises a VL CDR1, VL CDR3 and VL CDR3 of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In other embodiments, provided is a multispecific PSMAxCD3 antibody comprising a first binding domain that comprises a VH CDR1, VH CDR3, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 of any of the PSMA antibodies provided in Tables 4-12 or 23-28.

[00211] In some embodiments, provided is a multispecific PSMAxCD3 antibody comprising a second binding domain that comprises a VH of any of the CD3 antibodies provided in Tables 16-22 or 23-28. In some embodiments, provided is a multispecific PSMAxCD3 antibody comprising a second binding domain that comprises a VL of any of the CD3 antibodies provided in Tables 16-22 or 23-28. In some embodiments, provided is a multispecific PSMAxCD3 antibody comprising a second binding domain that comprises a VH and VL of any of the CD3 antibodies provided in Tables 16-22 or 23-28. In other embodiments, provided is a multispecific PSMAxCD3 antibody comprising a second binding domain that comprises a VH CDR1, VH CDR3 and VH CDR3 of any of the CD3 antibodies provided in Tables 16-22 or 23-28. In other embodiments, provided is a multispecific PSMAxCD3 antibody comprising a second binding domain that comprises a VL CDR1, VL CDR3 and VL CDR3 of any of the CD3 antibodies provided in Tables 16-22 or 23-28. In other embodiments, provided is a multispecific PSMAxCD3 antibody comprising a second binding domain that comprises a VH CDR1, VH CDR3, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 of any of the CD3 antibodies provided in Tables 16-22 or 23-28. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences are according to the Kabat numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences are according to the Chothia numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences are according to the AbM numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences are according to the Contact numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences are according to the IMGT numbering system. In some embodiments, the VH CDR1, VH

CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences are according to a combination of the numbering systems provided herein.

[00212] In some embodiments, the multispecific PSMAxCD3 antibody is PS3B1353, PS3B1505, PS3B1508, PS3B1917, PS3B1918, PS3B1919, PS3B1920, PS3B1921, PS3B1922, PS3B1923, PS3B1924, PS3B1925, PS3B1926, PS3B1927, PS3B1928, PSMB2908, PSMB2909, PS3B917, PS3B918, PS3B913, PS3B915, PS3B914, PS3B916, PS3B919, PS3B921, PS3B920, PS3B922, PS3B912, PS3B930, PS3B931, PS3B926, PS3B928, PS3B927, PS3B929, PS3B932, PS3B934, PS3B933, PS3B935, PS3B925, PS3B1352, PS3B1353, PS3B1354, PS3B1355, PS3B1356, PS3B1357, PS3B1358, PSMB937, PS3B1391, or PS3B1396. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1353. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1505. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1508. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1917. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1918. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1919. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1920. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1921. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1922. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1923. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1924. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1925. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1926. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1927. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1928. In one embodiment, the multispecific PSMAxCD3 antibody is PSMB2908. In one embodiment, the multispecific PSMAxCD3 antibody is PSMB2909. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B917. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B918. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B913. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B915. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B914. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B916. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B919. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B921. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B920. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B922. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B912. In one

embodiment, the multispecific PSMAxCD3 antibody is PS3B930. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B931. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B926. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B928. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B927. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B929. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B932. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B934. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B933. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B935. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B925. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1352. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1353. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1354. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1355. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1356. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1357. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1358. In one embodiment, the multispecific PSMAxCD3 antibody is PSMB937. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1391. In one embodiment, the multispecific PSMAxCD3 antibody is PS3B1396.

[00213] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1353 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1353 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1353 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1353 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B1353 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B1353 that binds to PSMA.

[00214] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1505 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1505 that binds to CD3. In some embodiments, the multispecific PSMAxCD3

antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1505 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1505 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B1505 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B1505 that binds to PSMA.

[00215] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1508 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1508 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B1508 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B1508 that binds to PSMA.

[00216] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1917 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1917 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1917 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1917 that binds to PSMA.

[00217] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1918 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1918 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1918 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1918 that binds to PSMA.

[00218] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1919 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1919 that binds to CD3. In some embodiments, the multispecific PSMAxCD3

antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1919 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1919 that binds to PSMA.

[00219] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1920 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1920 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1920 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1920 that binds to PSMA.

[00220] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1921 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1921 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1921 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1921 that binds to PSMA.

[00221] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1922 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1922 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1922 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1922 that binds to PSMA.

[00222] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1923 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1923 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1923 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1923 that binds to PSMA.

[00223] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1924 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1924 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1924 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1924 that binds to PSMA.

[00224] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1925 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1925 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1925 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1925 that binds to PSMA.

[00225] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1926 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1926 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1926 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1926 that binds to PSMA.

[00226] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1927 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1927 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1927 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1927 that binds to PSMA.

[00227] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1928 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1928 that binds to CD3. In some embodiments, the multispecific PSMAxCD3

antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1928 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1928 that binds to PSMA.

[00228] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B917 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B917 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B917 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B917 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B917 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B917 that binds to PSMA.

[00229] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B918 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B918 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B918 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B918 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B918 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B918 that binds to PSMA.

[00230] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B913 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B913 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B913 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B913 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of

PS3B913 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B913 that binds to PSMA.

[00231] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B915 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B915 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B915 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B915 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B915 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B915 that binds to PSMA.

[00232] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B914 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B914 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B914 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B914 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B914 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B914 that binds to PSMA.

[00233] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B916 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B916 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B916 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B916 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of

PS3B916 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B916 that binds to PSMA.

[00234] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B919 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B919 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B919 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B919 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B919 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B919 that binds to PSMA.

[00235] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B921 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B921 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B921 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B921 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B921 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B921 that binds to PSMA.

[00236] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B920 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B920 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B920 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B920 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of

PS3B920 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B920 that binds to PSMA.

[00237] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B922 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B922 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B922 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B922 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B922 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B922 that binds to PSMA.

[00238] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B912 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B912 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B912 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B912 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B912 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B912 that binds to PSMA.

[00239] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B930 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B930 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B930 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B930 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of

PS3B930 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B930 that binds to PSMA.

[00240] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B931 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B931 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B931 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B931 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B931 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B931 that binds to PSMA.

[00241] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B926 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B926 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B926 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B926 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B926 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B926 that binds to PSMA.

[00242] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B928 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B928 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B928 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B928 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of

PS3B928 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B928 that binds to PSMA.

[00243] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B927 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B927 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B927 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B927 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B927 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B927 that binds to PSMA.

[00244] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B929 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B929 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B929 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B929 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B929 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B929 that binds to PSMA.

[00245] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B932 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B932 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B932 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B932 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of

PS3B932 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B932 that binds to PSMA.

[00246] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B934 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B934 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B934 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B934 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B934 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B934 that binds to PSMA.

[00247] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B933 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B933 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B933 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B933 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B933 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B933 that binds to PSMA.

[00248] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B935 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B935 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B935 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B935 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of

PS3B935 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B935 that binds to PSMA.

[00249] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B925 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B925 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B925 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B925 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B925 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B925 that binds to PSMA.

[00250] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1352 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1352 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1352 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1352 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B1352 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B1352 that binds to PSMA.

[00251] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1353 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1353 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1353 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1353 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of

PS3B1353 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B1353 that binds to PSMA.

[00252] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1354 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1354 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1354 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1354 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B1354 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B1354 that binds to PSMA.

[00253] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1355 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1355 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1355 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1355 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B1355 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B1355 that binds to PSMA.

[00254] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1356 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1356 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1356 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1356 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of

PS3B1356 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B1356 that binds to PSMA.

[00255] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1357 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1357 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1357 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1357 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B1357 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B1357 that binds to PSMA.

[00256] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1358 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1358 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1358 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1358 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B1358 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B1358 that binds to PSMA.

[00257] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B937 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B937 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B937 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B937 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of

PS3B937 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B937 that binds to PSMA.

[00258] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1391 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1391 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1391 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1391 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B1391 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B1391 that binds to PSMA.

[00259] In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 of PS3B1396 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC1 of PS3B1396 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC1 and LC1 of PS3B1396 that binds to CD3. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 of PS3B1396 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the LC2 of PS3B1396 that binds to PSMA. In some embodiments, the multispecific PSMAxCD3 antibody comprises the amino acid sequence of the HC2 and LC2 of PS3B1396 that binds to PSMA.

[00260] In specific embodiments, provided is a multispecific antibody comprising a PSMA antibody provided herein in a knob-in-hole format. In specific embodiments, provided is a bispecific antibody comprising a PSMA antibody provided herein in a knob-in-hole format. In specific embodiments, provided is a trispecific antibody comprising a PSMA antibody provided herein in a knob-in-hole format. In specific embodiments, provided is a quadraspecific antibody comprising a PSMA antibody provided herein in a knob-in-hole format. Other specificities can be added to an antibody in knob-in-hole format using methods well known in the art (*e.g.*, adding an scFv to the N-terminus or C-terminus). In

addition, other formats and methods of making multispecific antibodies are also known in the art and contemplated. In some embodiments, a PSMA antibody provided herein is comprised in a bispecific antibody. In some embodiments, a PSMA antibody provided herein is comprised in a trispecific antibody. In some embodiments, a PSMA antibody provided herein is comprised in a quadraspecific antibody. In some embodiments, a PSMA bispecific antibody provided herein is comprised in a multispecific antibody.

[00261] In certain embodiments, a multispecific antibody provided herein comprises a first binding domain comprising a PSMA antibody provided herein that binds to a first PSMA epitope, and a second binding domain that binds to a second epitope, wherein the first PSMA epitope and the second epitope are not the same. In certain embodiments, a bispecific antibody provided herein comprises a first binding domain comprising a PSMA antibody provided herein that binds to a first PSMA epitope, and a second binding domain that binds to a second epitope, wherein the first PSMA epitope and the second epitope are not the same. In certain embodiments, a trispecific antibody provided herein comprises a first binding domain comprising a PSMA antibody provided herein that binds to a first PSMA epitope, a second binding domain that binds to a second epitope, and a third binding domain that binds to a third epitope, wherein the first PSMA epitope, the second epitope, and the third epitope are not the same. In certain embodiments, a quadraspecific antibody provided herein comprises a first binding domain comprising a PSMA antibody provided herein that binds to a first PSMA epitope, a second binding domain that binds to a second epitope, a third binding domain that binds to a third epitope, and a fourth binding domain that binds to a fourth epitope, wherein the first PSMA epitope, the second epitope, the third epitope, and the fourth epitope are not the same. In certain embodiments, a multispecific antibody provided herein comprises a first binding domain comprising a PSMA antibody provided herein that binds to a first PSMA antigen, and a second binding domain that binds to a second antigen, wherein the first PSMA antigen and the second antigen are not the same. In certain embodiments, a bispecific antibody provided herein comprises a first binding domain comprising a PSMA antibody provided herein that binds to a first PSMA antigen, and a second binding domain that binds to a second antigen, wherein the first PSMA antigen and the second antigen are not the same. In certain embodiments, a trispecific antibody provided herein comprises a first binding domain comprising a PSMA antibody provided herein that binds to a first PSMA antigen, a second binding domain that binds to a second antigen, and a third binding domain that binds to a third antigen, wherein the first PSMA antigen, the

second antigen, and the third antigen are not the same. In certain embodiments, a quadraspecific antibody provided herein comprises a first binding domain comprising a PSMA antibody provided herein that binds to a first PSMA antigen, a second binding domain that binds to a second antigen, a third binding domain that binds to a third antigen, and a fourth binding domain that binds to a fourth antigen, wherein the first PSMA antigen, the second antigen, the third antigen, and the fourth antigen are not the same. In a specific embodiment, a PSMA antibody, or antigen binding fragment thereof, provided herein specifically binds to PSMA.

[00262] In some embodiments, the multispecific antibody comprises heavy chain variable regions and light chain variable region. In some embodiments, the first binding domain comprises a heavy chain variable region and a light chain variable region. In some embodiments, the second binding domain comprises a heavy chain variable region and a light chain variable region. In some embodiments, the first binding domain comprises a heavy chain variable region and a light chain variable region, and the second binding domain comprises a heavy chain variable region and a light chain variable region. In some embodiments, the antibody is not a single domain antibody or nanobody. In some embodiments, the third binding domain comprises a heavy chain variable region and a light chain variable region. In some embodiments, the fourth binding domain comprises a heavy chain variable region and a light chain variable region.

[00263] In certain embodiments, the PSMA multispecific antibodies or antigen binding fragments thereof bind to a first epitope located on PSMA and a second epitope of a second target antigen. In some embodiments, provided herein is a multispecific antibody comprising: (a) a first binding domain that binds to a PSMA antigen, and (b) a second binding domain that binds to a second target antigen. In some embodiments, provided herein is a multispecific antibody comprising: (a) a first binding domain that specifically binds to a PSMA antigen, and (b) a second binding domain that specifically binds to a second target antigen. In some embodiments, provided herein is a multispecific antibody comprising: (a) a first binding domain that binds to a first epitope on a PSMA antigen, and (b) a second binding domain that binds to a second epitope on a second target antigen. In some embodiments, provided herein is a multispecific antibody comprising: (a) a first binding domain that specifically binds to a first epitope on a PSMA antigen, and (b) a second binding domain that specifically binds to a second epitope on a second target antigen.

[00264] In specific embodiments, the PSMA antigen is on the surface of a T cell. In certain embodiments, the second target antigen is not PSMA. The binding of the PSMA multispecific antibody to PSMA present on the surface of the T cell, and the binding of the second target antigen present on the surface of the second target cell can, for example, result in the killing of the second target cell. In other embodiment, the binding of the PSMA multispecific antibody to PSMA present on the surface of the T cell, and the binding of a second target antigen can, for example, result in the activation of the T cell.

[00265] In another aspect, provided herein is a multispecific antibody that comprises a first binding domain that binds to PSMA and a second binding domain that binds to CD3 ("multispecific PSMAxCD3 antibody"). In some embodiments, the multispecific PSMAxCD3 antibody is a bispecific antibody. In some embodiments, the multispecific PSMAxCD3 antibody is a trispecific antibody. In some embodiments, the multispecific PSMAxCD3 antibody is a quadraspecific antibody.

[00266] In one embodiment, the multispecific PSMAxCD3 antibody comprises: (a) a first binding domain that binds PSMA, and (b) a second binding domain that binds to CD3. In one embodiment, the multispecific PSMAxCD3 antibody comprises: (a) a first binding domain that binds PSMA, and (b) a second binding domain that binds to CD3, and (c) a third binding domain that binds to a third target. In one embodiment, the multispecific PSMAxCD3 antibody comprises: (a) a first binding domain that binds PSMA, and (b) a second binding domain that binds to CD3, (c) a third binding domain that binds to a third target, and (d) a fourth binding domain that binds to a fourth target.

[00267] In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:32. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2,

and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:66. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236. In one embodiment of the multispecific PSMAxCD3

antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:371; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:372. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:405; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:406. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a

VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:473; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:507; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:541; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:542. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:575; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:576. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid

sequence of SEQ ID NO:99; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:643; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:677; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:678. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:711; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:745; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:746. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:779; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a

VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:780. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain that binds PSMA comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:813; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:814.

[00268] In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the first binding domain that binds PSMA are according to the Kabat numbering system. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the first binding domain that binds PSMA are according to the Chothia numbering system. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the first binding domain that binds PSMA are according to the AbM numbering system. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the first binding domain that binds PSMA are according to the Contact numbering system. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the first binding domain that binds PSMA are according to the IMGT numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the first binding domain that binds PSMA are according to a combination of the numbering systems provided herein.

[00269] In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain binds a PSMA antigen. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain binds a PSMA epitope. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the first binding domain specifically binds to PSMA. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH

CDR3, VL CDR1, VL CDR2 and VL CDR3 of the first binding domain form a binding site for an antigen of the PSMA. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the first binding domain form a binding site for an epitope of the PSMA. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the PSMA is present on the surface of a T cell.

[00270] In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the second binding domain that binds CD3 comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:847; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:848. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the second binding domain that binds CD3 comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:847; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:848. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the second binding domain that binds CD3 comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:915; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:916. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the second binding domain that binds CD3 comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:915; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:916. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the second binding domain that binds CD3 comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:983; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2,

and a VL CDR3, respectively, of SEQ ID NO:984. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the second binding domain that binds CD3 comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:983; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:984. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the second binding domain that binds CD3 comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:1463; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:1464. In one embodiment of the multispecific PSMAxCD3 antibodies provided herein, the second binding domain that binds CD3 comprises: (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of SEQ ID NO:1505; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of SEQ ID NO:1464.

[00271] In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the second binding domain that binds CD3 are according to the Kabat numbering system. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the second binding domain that binds CD3 are according to the Chothia numbering system. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the second binding domain that binds CD3 are according to the AbM numbering system. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the second binding domain that binds CD3 are according to the Contact numbering system. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the second binding

domain that binds CD3 are according to the IMGT numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the second binding domain that binds CD3 are according to a combination of the numbering systems provided herein.

[00272] In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the second binding domain binds a CD3 antigen. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the second binding domain binds a CD3 epitope. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the second binding domain specifically binds to CD3. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the second binding domain form a binding site for an antigen of the CD3. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the second binding domain form a binding site for an epitope of the CD3. In some embodiments, the CD3 is present on the surface of a T cell.

[00273] In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the third target is not a PSMA antigen. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the fourth target is not a PSMA antigen. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the third target is not a PSMA antigen, and the fourth target is not a PSMA antigen. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the third target is not a CD3 antigen. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the fourth target is not a CD3 antigen. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the third target is not a CD3 antigen, and the fourth target is not a CD3 antigen. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the third target is not a PSMA epitope. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the fourth target is not a PSMA epitope. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the third target is not a PSMA epitope, and the fourth target is not a PSMA epitope. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the third target is not a CD3 epitope. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the fourth target is not a CD3 epitope. In some embodiments of the multispecific PSMAxCD3 antibodies provided herein, the third target is not a CD3 epitope, and the fourth target is not a CD3 epitope.

[00274] In a specific embodiment, the target is from a mammal. In a specific embodiment, the target is from a rat. In a specific embodiment, the target is from a mouse. In a specific embodiment, the target is from a primate. In a specific embodiment, the target is from a human.

[00275] In specific embodiments, provided is a multispecific PSMAxCD3 antibody in a knob-in-hole format. In specific embodiments, provided is a bispecific PSMAxCD3 antibody in a knob-in-hole format. In specific embodiments, provided is a trispecific antibody in a knob-in-hole format. In specific embodiments, provided is a quadraspecific antibody in a knob-in-hole format. Other specificities can be added to an antibody in knob-in-hole format using methods well known in the art (*e.g.*, adding an scFv to the N-terminus or C-terminus). In addition, other formats and methods of making multispecific antibodies are also known in the art and contemplated. In some embodiments, a PSMAxCD3 antibody provided herein is comprised in a bispecific antibody. In some embodiments, a PSMAxCD3 antibody provided herein is comprised in a trispecific antibody. In some embodiments, a PSMAxCD3 antibody provided herein is comprised in a quadraspecific antibody. In some embodiments, a PSMAxCD3 bispecific antibody provided herein is comprised in a multispecific antibody.

[00276] In certain embodiments, a trispecific PSMAxCD3 antibody provided herein comprises a first binding domain comprising a PSMA antibody provided herein that binds to a PSMA epitope, a second binding domain comprising a CD3 antibody provided herein that binds to a CD3 epitope, and a third binding domain that binds to a third epitope, wherein the PSMA epitope, the CD3 epitope, and the third epitope are not the same. In certain embodiments, a quadraspecific antibody provided herein comprises a first binding domain comprising a PSMA antibody provided herein that binds to a PSMA epitope, a second binding domain comprising a CD3 antibody provided herein that binds to a CD3 epitope, a third binding domain that binds to a third epitope, and a fourth binding domain that binds to a fourth epitope, wherein the PSMA epitope, the CD3 epitope, the third epitope, and the fourth epitope are not the same. In certain embodiments, a trispecific antibody provided herein comprises a first binding domain comprising a PSMA antibody provided herein that binds to a PSMA antigen, a second binding domain comprising a CD3 antibody provided herein that binds to a CD3 antigen, and a third binding domain that binds to a third antigen, wherein the PSMA antigen, the CD3 antigen, and the third antigen are not the same. In certain embodiments, a quadraspecific antibody provided herein that

binds to a PSMA antigen, a second binding domain comprising a CD3 antibody provided herein that binds to a CD3 antigen, a third binding domain that binds to a third antigen, and a fourth binding domain that binds to a fourth antigen, wherein the PSMA antigen, the CD3 antigen, the third antigen, and the fourth antigen are not the same. In certain embodiments of a multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds to PSMA specifically binds to the PSMA. In other embodiments of a multispecific PSMAxCD3 antibody provided herein, the second binding domain that binds to CD3 specifically binds to the CD3. In yet other embodiments of a multispecific PSMAxCD3 antibody provided herein, the first binding domain that binds to PSMA specifically binds to the PSMA, and the second binding domain that binds to CD3 specifically binds to the CD3.

[00277] In some embodiments, the multispecific PSMAxCD3 antibody comprises heavy chain variable regions and light chain variable region. In some embodiments, the first binding domain comprises a heavy chain variable region and a light chain variable region. In some embodiments, the second binding domain comprises a heavy chain variable region and a light chain variable region. In some embodiments, the first binding domain comprises a heavy chain variable region and a light chain variable region, and the second binding domain comprises a heavy chain variable region and a light chain variable region. In some embodiments, the PSMA antibody is not a single domain antibody or nanobody. In some embodiments, the third binding domain comprises a heavy chain variable region and a light chain variable region. In some embodiments, the fourth binding domain comprises a heavy chain variable region and a light chain variable region.

[00278] In certain embodiments, the PSMAxCD3 multispecific antibodies or antigen binding fragments thereof bind to a first epitope located on PSMA and a second epitope of located on CD3. In some embodiments, provided herein is a multispecific PSMAxCD3 antibody comprising: (a) a first binding domain that binds to a PSMA antigen, and (b) a second binding domain that binds to a CD3 antigen. In some embodiments, provided herein is a multispecific PSMAxCD3 antibody comprising: (a) a first binding domain that specifically binds to a PSMA antigen, and (b) a second binding domain that specifically binds to a CD3 antigen. In some embodiments, provided herein is a multispecific PSMAxCD3 antibody comprising: (a) a first binding domain that binds to a first epitope on a PSMA antigen, and (b) a second binding domain that binds to a second epitope on a CD3 antigen. In some embodiments, provided herein is a multispecific antibody comprising: (a) a

first binding domain that specifically binds to a first epitope on a PSMA antigen, and (b) a second binding domain that specifically binds to a second epitope on a CD3 antigen.

[00279] In specific embodiments, the PSMA antigen is on the surface of a T cell. In specific embodiments, the CD3 antigen is on the surface of a T cell. The binding of the PSMAxCD3 multispecific antibody to PSMA and CD3 present on the surface of T cells can, for example, result in the killing of the cell. In other embodiments, the binding of the PSMAxCD3 multispecific antibody to PSMA and CD3 present on the surface of T cells can, for example, result in the activation of the T cell.

[00280] In some embodiments, the PSMA antibody comprises a single chain antibody. In some embodiments, the PSMA antibody comprises a single domain antibody. In certain embodiments, the PSMA antibody comprises a nanobody. In certain embodiments, the PSMA antibody comprises a VHH antibody. In certain embodiments, the PSMA antibody comprises a llama antibody.

[00281] In some embodiments, the PSMA multispecific antibody comprises a single chain antibody. In some embodiments, the PSMA multispecific antibody comprises a single domain antibody. In certain embodiments, the PSMA multispecific antibody comprises a nanobody. In certain embodiments, the PSMA multispecific antibody comprises a VHH antibody. In certain embodiments, the PSMA multispecific antibody comprises a llama antibody. In some embodiments, the PSMA multispecific antibody does not comprise a single chain antibody. In some embodiments, the PSMA multispecific antibody does not comprise a single domain antibody. In certain embodiments, the PSMA multispecific antibody does not comprise a nanobody. In certain embodiments, the PSMA multispecific antibody does not comprise a VHH antibody. In certain embodiments, the PSMA multispecific antibody does not comprise a llama antibody.

[00282] According to another particular aspect, provided herein is a PSMA antibody or antigen-binding fragment thereof that induces antibody-dependent cell-mediated cytotoxicity (ADCC). The antibody or antigen-binding fragment thereof can, for example, induce ADCC *in vitro*.

[00283] In certain embodiments, the antibody or antigen-binding fragment thereof induces T cell dependent cytotoxicity of a second cell *in vitro* with an EC₅₀ of less than about 160 pM, when assessed *in vitro* at an effector to target cell ratio of 1:1.

[00284] In some embodiments, CD3 is present on the surface of a T cell. In some embodiments, the CD3 is present on the surface of a T cell, and the second target antigen is PSMA on the surface of a second cell. In some embodiments, the second cell is killed when the multispecific antibody binds to the CD3 on the surface of the T cell and the PSMA target antigen on the surface of the second cell. In some embodiments, the second cell is a prostate cell. In some embodiments, the second cell is a prostate cancer cell. In some embodiments, the second cell is a renal cell. In some embodiments, the second cell is a renal cancer cell.

[00285] In some embodiments, the multispecific antibody induces T cell dependent cytotoxicity of the second cell *in vitro* with an EC₅₀ of less than about 500 pM. In some embodiments, the multispecific antibody induces T cell dependent cytotoxicity of the second cell *in vitro* with an EC₅₀ of less than about 300 pM. In some embodiments, the multispecific antibody induces $\gamma\delta$ T cell dependent cytotoxicity of the second cell *in vitro* with an EC₅₀ of less than about 160 pM. In some embodiments, the EC₅₀ is assessed with a mixture of $\gamma\delta$ T effector cells and target cells expressing the second target antigen. In some embodiments, the effector cell to target cell ratio is about 0.01 to 1 to about 5 to 1. In some embodiments, the effector cell to target cell ratio is about 0.1 to 1 to about 2 to 1. In some embodiments, the effector cell to target cell ratio is about 1:1.

[00286] In certain embodiments, the EC₅₀ is less than about 1000 pM, less than about 900 pM, less than about 800 pM, less than about 700 pM, less than about 600 pM, less than about 500 pM, less than about 400 pM, less than about 300 pM, less than about 200 pM, less than about 190 pM, less than about 180 pM, less than about 170 pM, less than about 160 pM, less than about 150 pM, less than about 140 pM, less than about 130 pM, less than about 120 pM, less than about 110 pM, less than about 100 pM, less than about 90 pM, less than about 80 pM, less than about 70 pM, less than about 60 pM, less than about 50 pM, less than about 40 pM, less than about 30 pM, less than about 20 pM, or less than about 10 pM.

[00287] In certain embodiments, the effector to target cell ratio can, for example, be 0.01:1, 0.02:1, 0.03:1, 0.04:1, 0.05:1, 0.06:1, 0.07:1, 0.08:1, 0.09:1, 1:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, or 10:1. In certain embodiments, the concentration of the multispecific antibody or antigen-binding fragment thereof is about 0.000005 ng/mL, about 0.00005 ng/mL, about 0.0005, about 0.005 ng/mL, about 0.01 ng/mL, about 0.02 ng/mL, about 0.03 ng/mL, about 0.04 ng/mL, about 0.05 ng/mL, about 0.06 ng/mL, about 0.07 ng/mL, about 0.08 ng/mL, about 0.09 ng/mL, about 0.1 ng/mL, about 0.5 ng/mL, about 1.0 ng/mL, about 10 ng/mL, about 20 ng/mL, about 30 ng/mL, about 40 ng/mL, about 50 ng/mL, about

60 ng/mL, about 70 ng/mL, about 80 ng/mL, about 90 ng/mL, about 100 ng/mL, or about 1000 ng/mL.

[00288] In another aspect, provided herein is an antibody that competes for binding to PSMA with any of the PSMA antibodies described herein. In another aspect, provided herein is an antibody that binds to the same epitope as any of the PSMA antibodies described herein. In another aspect, provided is a PSMA antibody that binds an epitope on PSMA that overlaps with the epitope on PSMA bound by a PSMA antibody described herein. In some embodiments, the PSMA antibody comprises a VH CDR1, VH CDR2, and VH CDR3 of a PSMA antibody provided herein. In some embodiments, the PSMA antibody comprises a VL CDR1, VL CDR2, and VL CDR3 of a PSMA antibody provided herein. In some embodiments, the PSMA antibody comprises a VH CDR1, VH CDR2, VH CDR3, a VL CDR1, VL CDR2, and VL CDR3 of a PSMA antibody provided herein. In some embodiments, the PSMA antibody comprises a VH of a PSMA antibody provided herein. In some embodiments, the PSMA antibody comprises a VL of a PSMA antibody provided herein. In some embodiments, the PSMA antibody comprises a VH and a VL of a PSMA antibody provided herein. In some embodiments, the PSMA antibody comprises a VH CDR1, VH CDR2, VH CDR3, a VL CDR1, VL CDR2, and VL CDR3 of a PSMA antibody provided herein. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Kabat numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Chothia numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the AbM numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Contact numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the IMGT numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to a combination of the numbering systems provided herein. In certain embodiments, the PSMA antibody is a multispecific antibody. In some embodiments, the PSMA antibody is a bispecific antibody.

[00289] In another aspect, provided is an antibody that competes for binding to PSMA with a PSMA reference antibody. In another aspect, provided is a PSMA antibody that binds to the same PSMA epitope as a PSMA reference antibody. In another aspect, provided is a PSMA antibody that binds an epitope on PSMA that overlaps with the epitope on PSMA bound by a PSMA reference antibody. In some embodiments, the PSMA reference antibody comprises a VH CDR1, VH CDR2, and VH CDR3 of a PSMA reference antibody provided herein. In some embodiments, the PSMA reference antibody comprises a VL CDR1, VL CDR2, and VL CDR3 of a PSMA reference antibody provided herein. In some embodiments, the PSMA reference antibody comprises a VH CDR1, VH CDR2, VH CDR3, a VL CDR1, VL CDR2, and VL CDR3 of a PSMA reference antibody provided herein. In some embodiments, the PSMA reference antibody comprises a VH of a PSMA reference antibody provided herein. In some embodiments, the PSMA reference antibody comprises a VL of a PSMA reference antibody provided herein. In some embodiments, the PSMA reference antibody comprises a VH and a VL of a PSMA reference antibody provided herein. In some embodiments, the PSMA reference antibody comprises a VH CDR1, VH CDR2, VH CDR3, a VL CDR1, VL CDR2, and VL CDR3 of a PSMA reference antibody provided herein. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA reference antibody are according to the Kabat numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA reference antibody are according to the Chothia numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA reference antibody are according to the AbM numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA reference antibody are according to the Contact numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA reference antibody are according to the IMGT numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA reference antibody are according to a combination of the numbering systems provided herein. In certain embodiments, the antibody is a multispecific antibody. In some embodiments, the antibody is a bispecific antibody. In certain embodiments, the PSMA reference antibody is a multispecific antibody. In some embodiments, the PSMA reference antibody is a bispecific antibody.

[00290] The disclosure also provides an isolated multispecific antibody, comprising: a first half molecule and a second half molecule, wherein the first half molecule comprises a first antigen binding domain and a second antigen binding domain and the second half molecule comprises a third antigen binding domain, wherein the first antigen binding domain specifically binds PSMA, the second antigen binding domain specifically binds a second target, and the third antigen binding domain specifically binds a third target. In certain embodiments, the second target is CD3.

[00291] In some embodiments, the multispecific PSMAxCD3 antibody activates CD3+ T cells. In some embodiments, the multispecific PSMAxCD3 antibody binds to PSMA on prostate cells and binds to CD3 on T cells. In certain embodiments, the multispecific PSMAxCD3 antibody recruits CD3+ T cells to PSMA-expressing cells. In certain embodiments, the multispecific PSMAxCD3 antibody induces T cell proliferation. In certain embodiments, the multispecific PSMAxCD3 antibody induces T cell-mediated killing of the PSMA-expressing cells. In some embodiments, the multispecific PSMAxCD3 antibody specifically binds PSMA and the CD3 with an affinity that results in activation or recruitment of CD3+ T cells only upon co-engagement of the CD3 and PSMA. In specific embodiments, the PSMA-expressing cells are prostate cells. In some embodiments, the PSMA-expressing cells are prostate cancer cells. In some embodiments, the PSMA-expressing cells are renal cells. In some embodiments, the PSMA-expressing cells are renal cancer cells.

[00292] In some embodiments described herein, immune effector properties of the antibodies provided herein can be enhanced or silenced through Fc modifications by techniques known to those skilled in the art. For example, Fc effector functions such as C1q binding, complement dependent cytotoxicity (CDC), antibody-dependent cell-mediated cytotoxicity (ADCC), antibody-dependent cell-mediated phagocytosis (ADCP), down regulation of cell surface receptors (*e.g.*, B cell receptor; BCR), *etc.* can be provided and/or controlled by modifying residues in the Fc responsible for these activities.

[00293] “Antibody-dependent cell-mediated cytotoxicity” or “ADCC” refers to a cell-mediated reaction in which non-specific cytotoxic cells that express Fc receptors (FcRs) (*e.g.* Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell.

[00294] The ability of antibodies to induce ADCC can be enhanced by engineering their oligosaccharide component. Human IgG1 or IgG3 are N-glycosylated at Asn297 with the majority of the glycans in the well-known biantennary G0, G0F, G1, G1F, G2 or G2F forms. Antibodies produced by non-engineered CHO cells typically have a glycan fucose content of about at least 85%. The removal of the core fucose from the biantennary complex-type oligosaccharides attached to the Fc regions enhances the ADCC of antibodies via improved FcγRIIIa binding without altering antigen binding or CDC activity. Such Abs can be achieved using different methods reported to lead to the successful expression of relatively high defucosylated antibodies bearing the biantennary complex-type of Fc oligosaccharides such as control of culture osmolality (Konno *et al.*, *Cytotechnology* 64:249-65, 2012), application of a variant CHO line Lec13 as the host cell line (Shields *et al.*, *J Biol Chem* 277:26733-26740, 2002), application of a variant CHO line EB66 as the host cell line (Olivier *et al.*, *MAbs*; 2(4), 2010; Epub ahead of print; PMID:20562582), application of a rat hybridoma cell line YB2/0 as the host cell line (Shinkawa *et al.*, *J Biol Chem* 278:3466-3473, 2003), introduction of small interfering RNA specifically against the α -1,6-fucosyltransferase (FUT8) gene (Mori *et al.*, *Biotechnol Bioeng* 88:901-908, 2004), or coexpression of β -1,4-N-acetylglucosaminyltransferase III and golgi α -mannosidase II or a potent alpha-mannosidase I inhibitor, kifunensine (Ferrara *et al.*, *J Biol Chem* 281:5032-5036, 2006, Ferrara *et al.*, *Biotechnol Bioeng* 93:851-861, 2006; Xhou *et al.*, *Biotechnol Bioeng* 99:652-65, 2008).

[00295] In some embodiments described herein, ADCC elicited by the antibodies provided herein can also be enhanced by certain substitutions in the antibody Fc. Exemplary substitutions are for example substitutions at amino acid positions 256, 290, 298, 312, 356, 330, 333, 334, 360, 378 or 430 (residue numbering according to the EU index) as described in U.S. Pat. No. 6,737,056.

[00296] The IgG class is divided in four isotypes: IgG1, IgG2, IgG3 and IgG4 in humans. They share more than 95% homology in the amino acid sequences of the Fc regions but show major differences in the amino acid composition and structure of the hinge region. The Fc region mediates effector functions, such as antibody-dependent cellular cytotoxicity (ADCC) and complement-dependent cytotoxicity (CDC). In ADCC, the Fc region of an antibody binds to Fc receptors (FcγRs) on the surface of immune effector cells such as natural killers and macrophages, leading to the phagocytosis or lysis of the targeted cells. In CDC, the antibodies kill the targeted cells by triggering the complement cascade at the cell

surface. The antibodies described herein include antibodies with the described features of the variable domains in combination with any of the IgG isotypes, including modified versions in which the Fc sequence has been modified to effect different effector functions.

[00297] For many applications of therapeutic antibodies, Fc-mediated effector functions are not part of the mechanism of action. These Fc-mediated effector functions can be detrimental and potentially pose a safety risk by causing off-mechanism toxicity. Modifying effector functions can be achieved by engineering the Fc regions to reduce their binding to FcγRs or the complement factors. The binding of IgG to the activating (FcγRI, FcγRIIa, FcγRIIIa and FcγRIIIb) and inhibitory (FcγRIIb) FcγRs or the first component of complement (C1q) depends on residues located in the hinge region and the CH2 domain. Mutations have been introduced in IgG1, IgG2 and IgG4 to reduce or silence Fc functionalities. The antibodies described herein can include these modifications.

[00298] In one embodiment, the antibody comprises an Fc region with one or more of the following properties: (a) reduced effector function when compared to the parent Fc; (b) reduced affinity to Fcγ RI, Fcγ RIIa, Fcγ RIIf, Fcγ RIIfb and/or Fcγ RIIfa, (c) reduced affinity to FcγRI (d) reduced affinity to FcγRIIa (e) reduced affinity to FcγRIIf, (f) reduced affinity to Fcγ RIIfb or (g) reduced affinity to FcγRIIfa.

[00299] In some embodiments, the antibodies or antigen-binding fragments are IgG, or derivatives thereof, *e.g.*, IgG1, IgG2, IgG3, and IgG4 isotypes. In some embodiments wherein the antibody has an IgG4 isotype, the antibody contains S228P, L234A, and L235A substitutions in its Fc region. In some embodiments wherein the antibody has an IgG1 isotype, the antibody contains S228P, L234A, and L235A substitutions in its Fc region. The antibodies described herein can include these modifications. In some embodiments, the antibody has an IgG1 isotype.

[00300] In some embodiments, the antibody is of IgG4 isotype, optionally comprising a heavy chain substitution S228P when compared to the wild type IgG4.

[00301] In some embodiments, the antibody is of IgG1 isotype, optionally comprising heavy chain substitutions L234A, G237A, P238S, H268A, V309L, A330S and P331S when compared to the wild type IgG1.

[00302] In some embodiments, a PSMA antibody provided herein is chimeric. In some embodiments, a PSMA antibody provided herein is human. In some embodiments, a PSMA antibody provided herein is humanized. In certain embodiments, a PSMA antibody provided

herein is an isolated PSMA antibody. In some embodiments, a PSMA antigen binding fragment provided herein is chimeric. In some embodiments, a PSMA antigen binding fragment provided herein is human. In some embodiments, a PSMA antigen binding fragment provided herein is humanized. In certain embodiments, a PSMA antigen binding fragment provided herein is an isolated PSMA antigen binding fragment. In some embodiments, a PSMA antibody provided herein is an IgG antibody. In some embodiments, the IgG antibody is an IgG1 antibody. In some embodiments, the IgG antibody is an IgG2 antibody. In some embodiments, the IgG antibody is an IgG3 antibody. In some embodiments, the IgG antibody is an IgG4 antibody. In some embodiments, a PSMA antibody provided herein is multivalent. In some embodiments, the PSMA antibody is capable of binding at least three antigens. In some embodiments, the PSMA antibody is capable of binding at least four antigens. In some embodiments, the PSMA antibody is capable of binding at least five antigens.

[00303] In some embodiments, a PSMA multispecific antibody provided herein is chimeric. In some embodiments, a PSMA multispecific antibody provided herein is human. In some embodiments, a PSMA multispecific antibody provided herein is humanized. In certain embodiments, a PSMA multispecific antibody provided herein is an isolated PSMA multispecific antibody. In some embodiments, a PSMA multispecific antibody comprising a PSMA antigen binding fragment provided herein is chimeric. In some embodiments, a PSMA multispecific antibody comprising a PSMA antigen binding fragment provided herein is human. In some embodiments, a PSMA multispecific antibody comprising a PSMA antigen binding fragment provided herein is humanized. In certain embodiments, a PSMA multispecific antibody comprising a PSMA antigen binding fragment provided herein is an isolated PSMA multispecific antibody. In certain embodiments, the PSMA multispecific antibody is a multispecific PSMAxCD3 antibody.

[00304] In some embodiments of the PSMA multispecific antibodies provided herein, the first binding domain is human. In some embodiments, the second binding domain is human. In some embodiments of the PSMA multispecific antibodies provided herein, both the first binding domain and the second binding domain are human. In some embodiments of the PSMA multispecific antibodies provided herein, the first binding domain is humanized. In some embodiments of the PSMA multispecific antibodies provided herein, the second binding domain is humanized. In some embodiments of the PSMA multispecific antibodies provided herein, both the first binding domain and the second binding domain are

humanized. In some embodiments of the PSMA multispecific antibodies provided herein, both the first binding domain is human and the second binding domain is humanized. In some embodiments of the PSMA multispecific antibodies provided herein, both the first binding domain is humanized and the second binding domain is human. In certain embodiments, the PSMA multispecific antibody is a multispecific PSMAxCD3 antibody.

[00305] In some embodiments, a PSMA multispecific antibody provided herein is multivalent. In some embodiments, the multispecific antibody is capable of binding at least three antigens. In some embodiments, the multispecific antibody is capable of binding at least five antigens. In certain embodiments, the multispecific antibody is a multispecific antibody. In some embodiments, a PSMA multispecific antibody provided herein is an IgG antibody. In some embodiments, the IgG antibody is an IgG1 antibody. In some embodiments, the IgG antibody is an IgG2 antibody. In some embodiments, the IgG antibody is an IgG3 antibody. In some embodiments, the IgG antibody is an IgG4 antibody. In certain embodiments, the PSMA multispecific antibody is a multispecific PSMAxCD3 antibody.

[00306] In certain embodiments, the antibodies provided herein are part of a multispecific antibody. In some embodiments, the multispecific antibody comprises a first binding domain that binds to a PSMA antigen. In some embodiments, the multispecific antibody comprises a first binding domain that binds to a PSMA antigen and comprises a second binding domain that binds to a second target antigen, as provided herein. In certain embodiments, the multispecific antibody binds to a PSMA antigen, a second target antigen, and one or more additional antigens. In some embodiments of the various antibodies provided herein, the antibody binds to an epitope of a given antigen. In certain embodiments, the multispecific PSMA antibody is a multispecific PSMAxCD3 antibody, wherein the second target is CD3.

[00307] Variants of the antibodies specifically binding PSMA described herein are also contemplated as embodiments. In some embodiments, antibodies provided herein have altered amino acid sequences when compared to the parental antibodies can be generated using standard cloning and expression technologies. For example, site-directed mutagenesis or PCR-mediated mutagenesis can be performed to introduce the mutation(s) and the effect on antibody binding or other property of interest can be evaluated using well known methods and the methods described herein and in the Examples. For example, variants can comprise one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen or fifteen amino acid substitutions in the VH and/or the VL as long as the

homologous antibodies retain or have improved functional properties when compared to the parental antibodies. In one embodiment, a variant of a PSMA antibody provided herein comprises a VH with one amino acid substitution. In one embodiment, a variant of a PSMA antibody provided herein comprises a VH with two amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VH with three amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VH with four amino acid substitutions. In one embodiment, a variant of a PSMA antibody provided herein comprises a VH with five amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VH with six amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VH with seven amino acid substitutions. In one embodiment, a variant of a PSMA antibody provided herein comprises a VH with eight amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VH with nine amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VH with ten amino acid substitutions. In one embodiment, a variant of a PSMA antibody provided herein comprises a VH with eleven amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VH with twelve amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VH with thirteen amino acid substitutions. In one embodiment, a variant of a PSMA antibody provided herein comprises a VH with fourteen amino acid substitutions. In one embodiment, a variant of a PSMA antibody provided herein comprises a VH with fifteen amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VL with one amino acid substitution. In one embodiment, a variant of a PSMA antibody provided herein comprises a VL with two amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VL with three amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VL with four amino acid substitutions. In one embodiment, a variant of a PSMA antibody provided herein comprises a VL with five amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VL with six amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VL with seven amino acid substitutions. In one embodiment, a variant of a PSMA antibody provided herein comprises a VL with eight amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VL with nine amino acid substitutions. In

another embodiment, a variant of a PSMA antibody provided herein comprises a VL with ten amino acid substitutions. In one embodiment, a variant of a PSMA antibody provided herein comprises a VL with eleven amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VL with twelve amino acid substitutions. In another embodiment, a variant of a PSMA antibody provided herein comprises a VL with thirteen amino acid substitutions. In one embodiment, a variant of a PSMA antibody provided herein comprises a VL with fourteen amino acid substitutions. In one embodiment, a variant of a PSMA antibody provided herein comprises a VL with fifteen amino acid substitutions. In specific embodiments, the variation of the variant compared to the parental antibody is not within the CDRs of the variant. In certain embodiments, the amino acid substitution is a conservative modification.

[00308] In some embodiments, the sequence identity can be about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% to a VH or the VL amino acid sequences provided herein. In some embodiments, a variant of a PSMA antibody provided herein comprises a VH having about 90% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VH having about 91% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VH having about 92% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VH having about 93% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VH having about 94% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VH having about 95% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VH having about 96% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VH having about 97% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VH having about 98% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VH having about 99% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VL having about 90% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VL having about 91% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VL having about 92% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VL having about 93% sequence identity. In some embodiments, a variant

of a PSMA antibody provided herein comprises a VL having about 94% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VL having about 95% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VL having about 96% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VL having about 97% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VL having about 98% sequence identity. In some embodiments, a variant of a PSMA antibody provided herein comprises a VL having about 99% sequence identity. In specific embodiments, the variation of the variant compared to the parental antibody is not within the CDRs of the variant.

[00309] In some embodiments, a multispecific antibody provided herein is a diabody, a cross-body, or a multispecific antibody obtained via a controlled Fab arm exchange as those described herein.

Generation of monospecific antibodies

[00310] In some embodiments the PSMA antibodies are human. In some embodiments the PSMA antibodies are humanized.

[00311] Monospecific antibodies described herein (*e.g.*, antibodies specifically binding PSMA) can be generated using various technologies. For example, the hybridoma method of Kohler and Milstein, *Nature* 256:495, 1975 can be used to generate monoclonal antibodies. In the hybridoma method, a mouse or other host animal, such as a hamster, rat or monkey, is immunized with human chimpanzee or macaque PSMA or CD3 or fragments of PSMA or CD3, such as the extracellular domain of PSMA or CD3, followed by fusion of spleen cells from immunized animals with myeloma cells using standard methods to form hybridoma cells (Goding, *Monoclonal Antibodies: Principles and Practice*, pp. 59-103 (Academic Press, 1986)). Colonies arising from single immortalized hybridoma cells are screened for production of antibodies with desired properties, such as specificity of binding, cross-reactivity or lack thereof, and affinity for the antigen.

[00312] Various host animals can be used to produce the PSMA antibodies described herein. For example, Balb/c mice can be used to generate mouse anti-human PSMA antibodies. The antibodies made in Balb/c mice and other non-human animals can be humanized using various technologies to generate more human-like sequences.

[00313] Exemplary humanization techniques including selection of human acceptor frameworks are known and include CDR grafting (U.S. Patent No. 5,225,539), SDR grafting (U.S. Patent No. 6,818,749), Resurfacing (Padlan, (1991) Mol Immunol 28:489-499), Specificity Determining Residues Resurfacing (U.S. Patent Publ. No. 2010/0261620), human framework adaptation (U.S. Patent No. 8,748,356) or superhumanization (U.S. Patent No. 7,709, 226). In these methods, CDRs of parental antibodies are transferred onto human frameworks that can be selected based on their overall homology to the parental frameworks, based on similarity in CDR length, or canonical structure identity, or a combination thereof.

[00314] Humanized antibodies can be further optimized to improve their selectivity or affinity to a desired antigen by incorporating altered framework support residues to preserve binding affinity (back mutations) by techniques such as those described in Int. Patent Publ. Nos. WO1090/007861 and WO1992/22653, or by introducing variation at any of the CDRs for example to improve affinity of the antibody.

[00315] The framework sequences of the parental and engineered antibodies can further be modified, for example by back mutations to restore and/or improve binding of the generated antibodies to the antigen as described for example in U.S. Patent No. 6,180,370. The framework sequences of the parental or engineered antibodies can further be modified by mutating one or more residues within the framework region (or alternatively within one or more CDR regions) to remove T-cell epitopes to thereby reduce the potential immunogenicity of the antibody. This approach is also referred to as “deimmunization” and described in further detail in U.S. Patent Publ. No. US20070014796.

[00316] The CDR residues of the antibodies provided herein can be mutated to modulate affinity of the antibodies to PSMA and/or CD3.

[00317] The CDR residues of the antibodies provided herein can be mutated to minimize risk of post-translational modifications. Amino acid residues of putative motifs for deamination (NS), acid-catalyzed hydrolysis (DP), isomerization (DS), or oxidation (W) can be substituted with any of the naturally occurring amino acids to mutagenize the motifs, and the resulting antibodies can be tested for their functionality and stability using methods described herein.

[00318] Antibodies provided herein modified to improve stability, selectivity, cross-reactivity, affinity, immunogenicity or other desirable biological or biophysical property are

contemplated. Stability of an antibody is influenced by a number of factors, including (1) core packing of individual domains that affects their intrinsic stability, (2) protein/protein interface interactions that have impact upon the HC and LC pairing, (3) burial of polar and charged residues, (4) H-bonding network for polar and charged residues; and (5) surface charge and polar residue distribution among other intra- and inter-molecular forces (Worn *et al.*, (2001) *J Mol Biol* 305:989-1010). Potential structure destabilizing residues can be identified based upon the crystal structure of the antibody or by molecular modeling in certain cases, and the effect of the residues on antibody stability can be tested by generating and evaluating variants harboring mutations in the identified residues. One of the ways to increase antibody stability is to raise the thermal transition midpoint (T_m) as measured by differential scanning calorimetry (DSC). In general, the protein T_m is correlated with its stability and inversely correlated with its susceptibility to unfolding and denaturation in solution and the degradation processes that depend on the tendency of the protein to unfold (Remmele *et al.*, (2000) *Biopharm* 13:36-46). A number of studies have found correlation between the ranking of the physical stability of formulations measured as thermal stability by DSC and physical stability measured by other methods (Gupta *et al.*, (2003) *AAPS PharmSci* 5E8; Zhang *et al.*, (2004) *J Pharm Sci* 93:3076-89; Maa *et al.*, (1996) *Int J Pharm* 140:155-68; Bedu-Addo *et al.*, (2004) *Pharm Res* 21:1353-61; Remmele *et al.*, (1997) *Pharm Res* 15:200-8). Formulation studies suggest that a Fab T_m has implication for long-term physical stability of a corresponding mAb.

[00319] C-terminal lysine (CTL) can be removed from injected antibodies by endogenous circulating carboxypeptidases in the blood stream (Cai *et al.*, (2011) *Biotechnol Bioeng* 108:404-412). During manufacturing, CTL removal can be controlled to less than the maximum level by control of concentration of extracellular Zn^{2+} , EDTA or EDTA – Fe^{3+} as described in U.S. Patent Publ. No. US20140273092. CTL content in antibodies can be measured using known methods.

[00320] Fc substitutions can be made to the antibodies provided herein to modulate antibody effector functions and/or pharmacokinetic properties. In traditional immune function, the interaction of antibody-antigen complexes with cells of the immune system results in a wide array of responses, ranging from effector functions such as antibody-dependent cytotoxicity, mast cell degranulation, and phagocytosis to immunomodulatory signals such as regulating lymphocyte proliferation and antibody secretion. All these interactions are initiated through the binding of the Fc domain of antibodies or immune

complexes to specialized cell surface receptors on cells. The diversity of cellular responses triggered by antibodies and immune complexes results from the heterogeneity of the Fc receptors: FcγRI (CD64), FcγRIIa (CD32A), and FcγRIII (CD16) are activating Fcγ receptors (*i.e.*, immune system enhancing) whereas FcγRIIb (CD32B) is an inhibitory Fcγ receptor (*i.e.*, immune system dampening). Binding to the FcRn receptor modulates antibody half-life.

[00321] In some embodiments, the anti-PSMA antibodies or the bispecific anti-PSMA/anti-CD3 antibodies provided herein comprise at least one substitution in an Fc region.

[00322] In some embodiments, the anti-PSMA antibodies or the bispecific anti-PSMA/anti-CD3 antibodies provided herein comprise one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen or fifteen substitutions in the Fc region. As described elsewhere herein, Fc positions that can be substituted to modulate antibody half-life. Exemplary singular or combination substitutions that can be made to increase the half-life of the antibody are substitutions M428L/N434S, M252Y/S254T/T256E, T250Q/M428L, N434A and T307A/E380A/N434A. Exemplary singular or combination substitutions that can be made to reduce the half-life of the antibody are substitutions H435A, P257I/N434H, D376V/N434H, M252Y/S254T/T256E/H433K/N434F, T308P/N434A and H435R.

[00323] In some embodiments, the anti-PSMA antibodies or the bispecific anti-PSMA/anti-CD3 antibodies provided herein comprise at least one substitution in the Fc region selected from the group consisting of M428L/N434S, M252Y/S254T/T256E, T250Q/M428L, N434A, T307A/E380A/N434A, H435A, P257I/N434H, D376V/N434H, M252Y/S254T/T256E/H433K/N434F, T308P/N434A and H435R.

[00324] In some embodiments, the anti-PSMA antibodies or the bispecific anti-PSMA/anti-CD3 antibodies provided herein comprise at least one substitution in the Fc region that reduces binding of the antibody to an activating Fc-gamma receptor (FcγR) and/or reduces Fc effector functions such as C1q binding, complement dependent cytotoxicity (CDC), antibody-dependent cell-mediated cytotoxicity (ADCC) or phagocytosis (ADCP).

[00325] Fc positions that can be substituted to reduce binding of the antibody to the activating FcGR and subsequently to reduce effector function are substitutions L234A/L235A on IgG1, V234A/G237A/P238S/H268A/V309L/A330S/P331S on IgG2,

F234A/L235A on IgG4, S228P/F234A/ L235A on IgG4, N297A on all Ig isotypes, V234A/G237A on IgG2, K214T/E233P/ L234V/L235A/G236-deleted/A327G/P331A/D365E/L358M on IgG1, H268Q/V309L/ A330S/P331S on IgG2, S267E/L328F on IgG1, L234F/L235E/D265A on IgG1, L234A/L235A/G237A/P238S/H268A/A330S/P331S on IgG1, S228P/F234A/L235A/G237A/P238S on IgG4, and S228P/F234A/L235A/G236-deleted/G237A/P238S on IgG4.

[00326] An S228P substitution can also be made in IgG4 antibodies to enhance IgG4 stability. In some embodiments, the anti-PSMA antibodies or the bispecific anti-PSMA/anti-CD3 antibodies provided herein comprises a S228P substitution, wherein residue numbering is according to the EU Index. In some embodiments, the anti-PSMA antibodies or the bispecific anti-PSMA/anti-CD3 antibodies provided herein comprise a F234A, a L235A or a F234A/L235A substitution, wherein residue numbering is according to the EU Index. In some embodiments, the anti-PSMA antibodies or the bispecific anti-PSMA/anti-CD3 antibodies provided herein comprise a S228P, a F234A and a L235A substitution, wherein residue numbering is according to the EU Index.

[00327] In other embodiments, transgenic animals, such as mice or rat carrying human immunoglobulin (Ig) loci in their genome can be used to generate human antibodies against a target protein, and are described in for example U.S. Patent No. 6,150,584, Int. Patent Publ. No. WO99/45962, Int. Patent Publ. Nos. WO2002/066630, WO2002/43478, WO2002/043478 and WO1990/04036, Lonberg et al (1994) Nature 368:856-9; Green et al (1994) Nature Genet. 7:13-21; Green & Jakobovits (1998) Exp. Med. 188:483-95; Lonberg and Huszar (1995) Int Rev Immunol 13:65-93; Bruggemann *et al.*, (1991) Eur J Immunol 21:1323- 1326; Fishwild *et al.*, (1996) Nat Biotechnol 14:845-851; Mendez *et al.*, (1997) Nat Genet 15:146-156; Green (1999) J Immunol Methods 231:11-23; Yang *et al.*, (1999) Cancer Res 59:1236-1243; Bruggemann and Taussig (1997) Curr Opin Biotechnol 8:455-458. The endogenous immunoglobulin loci in such animal can be disrupted or deleted, and at least one complete or partial human immunoglobulin locus can be inserted into the genome of the animal using homologous or non-homologous recombination, using transchromosomes, or using minigenes. Companies such as Regeneron (http://_www_regeneron_com), Harbour Antibodies (http://_www_harbourantibodies_com), Open Monoclonal Technology, Inc. (OMT) (http://_www_omtinc_net), KyMab (http://_www_kymab_com), Trianni (http://_www.trianni_com) and Ablexis

(http://_www_ablexis_com) can be engaged to provide human antibodies directed against a selected antigen using technologies as described above.

[00328] Human antibodies can be selected from a phage display library, where the phage is engineered to express human immunoglobulins or portions thereof such as Fabs, single chain antibodies (scFv), or unpaired or paired antibody variable regions (Knappik *et al.*, (2000) J Mol Biol 296:57-86; Krebs *et al.*, (2001) J Immunol Meth 254:67-84; Vaughan *et al.*, (1996) Nature Biotechnology 14:309-314; Sheets *et al.*, (1998) PITAS (USA) 95:6157-6162; Hoogenboom and Winter (1991) J Mol Biol 227:381; Marks *et al.*, (1991) J Mol Biol 222:581). The antibodies provided herein can be isolated for example from phage display library expressing antibody heavy and light chain variable regions as fusion proteins with bacteriophage pIX coat protein as described in Shi *et al.*, (2010) J Mol Biol 397:385-96, and Int. Patent Publ. No. WO09/085462). The libraries can be screened for phage binding to human and/or cyno PSMA or CD3 and the obtained positive clones can be further characterized, the Fabs isolated from the clone lysates, and expressed as full length IgGs. Such phage display methods for isolating human antibodies are described in for example: U.S. Patent Nos. 5,223,409, 5,403,484, 5,571,698, 5,427,908, 5,580,717, 5,969,108, 6,172,197, 5,885,793; 6,521,404; 6,544,731; 6,555,313; 6,582,915 and 6,593,081.

[00329] Preparation of immunogenic antigens and monoclonal antibody production can be performed using any suitable technique, such as recombinant protein production. The immunogenic antigens can be administered to an animal in the form of purified protein, or protein mixtures including whole cells or cell or tissue extracts, or the antigen can be formed de novo in the animal's body from nucleic acids encoding said antigen or a portion thereof.

Generation of multispecific PSMAxCD3 antibodies

[00330] Multispecific PSMA antibodies, such as PSMAxCD3 bispecific antibodies provided herein can be generated by combining PSMA binding VH/VL domains with CD3 binding VH/VL domains isolated and characterized herein. Alternatively, the bispecific PSMAxCD3 antibodies can be engineered using VH/VL domains from publicly available monospecific anti-PSMA and anti-CD3 antibodies, and/or by mix-matching the PSMA or CD3 binding VH/VL domains identified herein with publicly available PSMA or CD3 binding VH/VL domains.

[00331] Exemplary anti-PSMA antibodies that can be used to engineer bispecific PSMAxCD3 molecules are for example those herein and in Tables 4-12 or 23-28. For

example, the VH/VL domains of the PSMA antibodies provided herein can be incorporated into bispecific antibodies comprising CD3 binding VH/VL domains described herein and in Tables 16-22 or 23-28. For example, the VH/VL domains of the CD3 antibodies CD3B376, CD3B450, CD3B2030 and CD3W245 described herein (see, *e.g.*, Tables 16-22) can be used to generate bispecific PSMAxCD3 antibodies.

[00332] Similarly, exemplary anti-CD3 antibodies that can be used to engineer bispecific PSMAxCD3 molecules are for example those described in Int. Patent Publ. Nos. WO2005/048935, WO2004/106380 and WO2015095392. These CD3 VH/VL domains can be incorporated into bispecific antibodies comprising PSMA binding VH/VL domains described herein and in Table 4-12 and 23-28. For example, the VH/VL domains of a PSMA antibody provided herein can be used to generate a PSMAxCD3 bispecific antibody. In certain embodiments, the PSMA antibody is PSMB889. In other embodiments, the PSMA antibody is PSMB890. In certain embodiments, the PSMA antibody is PSMB891. In certain embodiments, the PSMA antibody is PSMB892. In other embodiments, the PSMA antibody is PSMB893. In certain embodiments, the PSMA antibody is PSMB894. In other embodiments, the PSMA antibody is PSMB895. In certain embodiments, the PSMA antibody is PSMB896. In certain embodiments, the PSMA antibody is PSMB897. In other embodiments, the PSMA antibody is PSMB898. In certain embodiments, the PSMA antibody is PSMB899. In certain embodiments, the PSMA antibody is PSMHB49SC1133_011A11_1. In other embodiments, the PSMA antibody is PSMB896-G100A. In certain embodiments, the PSMA antibody is PSMA_P72_A10-HC-G54E. In certain embodiments, the PSMA antibody is PSMA_P72_D01-HC-D95E. In other embodiments, the PSMA antibody is PSMA_P72_F01. In other embodiments, the PSMA antibody is PSMA_P75_F01. In certain embodiments, the PSMA antibody is PSMA_P72_F07. In certain embodiments, the PSMA antibody is PSMA_P72_E07. In other embodiments, the PSMA antibody is PSMA_P72_D01. In certain embodiments, the PSMA antibody is PSMA_P72_C01. In other embodiments, the PSMA antibody is PSMA_P72_A10. In certain embodiments, the PSMA antibody is PSMA_P72_F02. In certain embodiments, the PSMA antibody is PSMA_P70_F02. In other embodiments, the PSMA antibody is PSMA_P72_G02. In other embodiments, the PSMA antibody is PSMA_P72_A11. In some embodiments, the PSMA antibody is PSMB946 (PSMB895 with a C-terminal Lys (K) amino acid residue). In some embodiments, the PSMA antibody is PSMB947 (PSMB896 with a C-terminal Lys (K) amino acid residue). In some

embodiments, the PSMA antibody is PSMB948 (PSMB897 with a C-terminal Lys (K) amino acid residue). In some embodiments, the PSMA antibody is PSMB949 (PSMB898 with a C-terminal Lys (K) amino acid residue).

[00333] The generated bispecific PSMAxCD3 antibodies can be tested for their binding to PSMA and CD3, and for their desired functional characteristics, such as T cell mediated killing of PSMA-expressing cells (*e.g.*, LNCaP).

[00334] Bispecific antibodies provided herein can comprise antibodies having a full length antibody structure. "Full length antibody" refers to an antibody having two full length antibody heavy chains and two full length antibody light chains. A full length antibody heavy chain (HC) consists of well-known heavy chain variable and constant domains VH, CH1, hinge, CH2, and CH3. A full length antibody light chain (LC) consists of well-known light chain variable and constant domains VL and CL. The full length antibody can be lacking the C-terminal lysine (K) in either one or both heavy chains. "Fab-arm" or "half molecule" refers to one heavy chain-light chain pair that specifically binds an antigen.

[00335] Full length bispecific antibodies can be generated for example using Fab arm exchange (or half molecule exchange) between two monospecific bivalent antibodies by introducing substitutions at the heavy chain CH3 interface in each half molecule to favor heterodimer formation of two antibody half molecules having distinct specificity either *in vitro* in cell-free environment or using co-expression. The Fab arm exchange reaction is the result of a disulfide-bond isomerization reaction and dissociation-association of CH3 domains. The heavy chain disulfide bonds in the hinge regions of the parental monospecific antibodies are reduced. The resulting free cysteines of one of the parental monospecific antibodies form an inter heavy-chain disulfide bond with cysteine residues of a second parental monospecific antibody molecule and simultaneously CH3 domains of the parental antibodies release and reform by dissociation-association. The CH3 domains of the Fab arms can be engineered to favor heterodimerization over homodimerization. The resulting product is a bispecific antibody having two Fab arms or half molecules which each bind a distinct epitope, *i.e.* an epitope on PSMA and an epitope on CD3. Other methods of making multispecific antibodies are known and contemplated.

[00336] "Homodimerization" refers to an interaction of two heavy chains having identical CH3 amino acid sequences. "Homodimer" refers to an antibody having two heavy chains with identical CH3 amino acid sequences. "Heterodimerization" refers to an interaction of

two heavy chains having non-identical CH3 amino acid sequences. "Heterodimer" refers to an antibody having two heavy chains with non-identical CH3 amino acid sequences.

[00337] As mentioned elsewhere, in some embodiments, the bispecific antibodies include designs such as the Triomab/Quadroma (Trion Pharma/Fresenius Biotech), Knob-in-Hole (Genentech), CrossMAbs (Roche) and the electrostatically-matched (Chugai, Amgen, NovoNordisk, Oncomed), the LUZ-Y (Genentech), the Strand Exchange Engineered Domain body (SEEDbody) (EMD Serono), the Biclonic (Merus) and the DuoBody (Genmab A/S).

[00338] In some embodiments, a PSMA multispecific antibody provided herein is in the knob-and-hole format. In some embodiments, a PSMA multispecific antibody provided herein is in a DuoBody format.

[00339] The Triomab quadroma technology can be used to generate full length bispecific antibodies provided herein. Triomab technology promotes Fab arm exchange between two parental chimeric antibodies, one parental mAb having IgG2a and the second parental mAb having rat IgG2b constant regions, yielding chimeric bispecific antibodies.

[00340] The "knob-in-hole" strategy (see, *e.g.*, PCT Publ. No. WO2006/028936) can be used to generate full length bispecific antibodies. Briefly, selected amino acids forming the interface of the CH3 domains in human IgG can be mutated at positions affecting CH3 domain interactions to promote heterodimer formation. An amino acid with a small side chain (hole) is introduced into a heavy chain of an antibody specifically binding a first antigen and an amino acid with a large side chain (knob) is introduced into a heavy chain of an antibody specifically binding a second antigen. After co-expression of the two antibodies, a heterodimer is formed as a result of the preferential interaction of the heavy chain with a "hole" with the heavy chain with a "knob." Exemplary CH3 substitution pairs forming a knob and a hole are (expressed as modified position in the first CH3 domain of the first heavy chain/modified position in the second CH3 domain of the second heavy chain): T366Y/F405A, T366W/F405W, F405W/Y407A, T394W/Y407T, T394S/Y407A, T366W/T394S, F405W/T394S and T366W/T366S_L368A_Y407V.

[00341] The CrossMAb technology can be used to generate full length bispecific antibodies provided herein. CrossMAbs, in addition to utilizing the "knob-in-hole" strategy to promote Fab arm exchange, have in one of the half arms the CH1 and the CL domains

exchanged to ensure correct light chain pairing of the resulting bispecific antibody (see *e.g.* U.S. Patent No. 8,242,247).

[00342] Other cross-over strategies can be used to generate full length bispecific antibodies provided herein by exchanging variable or constant, or both domains between the heavy chain and the light chain or within the heavy chain in the bispecific antibodies, either in one or both arms. These exchanges include for example VH-CH1 with VL-CL, VH with VL, CH3 with CL and CH3 with CH1 as described in Int. Patent Publ. Nos. WO2009/080254, WO2009/080251, WO2009/018386 and WO2009/080252.

[00343] Other strategies such as promoting heavy chain heterodimerization using electrostatic interactions by substituting positively charged residues at one CH3 surface and negatively charged residues at a second CH3 surface can be used, as described in US Pat. Publ. No. US2010/0015133; US Pat. Publ. No. US2009/0182127; US Pat. Publ. No. US2010/028637; or US Pat. Publ. No. US2011/0123532. In other strategies, heterodimerization can be promoted by the following substitutions (expressed as modified position in the first CH3 domain of the first heavy chain/modified position in the second CH3 domain of the second heavy chain): L351Y_F405AY407V/T394W, T366I_K392M_T394W/F405A_Y407V, T366L_K392M_T394W/F405A_Y407V, L351Y_Y407A/T366A_K409F, L351Y_Y407A/T366V_K409F Y407A/T366A_K409F, or T350V_L351Y_F405A_Y407V/T350V_T366L_K392L_T394W as described in U.S. Pat. Publ. No. US2012/0149876 or U.S. Pat. Publ. No. US2013/0195849.

[00344] LUZ-Y technology can be utilized to generate bispecific antibodies provided herein. In this technology, a leucine zipper is added into the C terminus of the CH3 domains to drive the heterodimer assembly from parental mAbs that is removed post-purification as described in Wranik *et al.*, (2012) J Biol Chem 287(52): 42221-9.

[00345] SEEDbody technology can be utilized to generate bispecific antibodies provided herein. SEEDbodies have, in their constant domains, select IgG residues substituted with IgA residues to promote heterodimerization as described in U.S. Patent No. US20070287170.

[00346] In some embodiments, also provided is a multispecific, multifunctional antibody that specifically binds to PSMA. Such a multispecific, multifunctional antibody that specifically binds to PSMA can be a trispecific antibody for dual targeting of tumor cells (*e.g.*, trifunctional structures that can be designed to target two different targets/epitopes on

the tumor cell and with the third functionality bind with high affinity to either T cells or NK-cells). Trispecific antibodies targeting two distinct tumor epitopes and engaging T- or NK-cells lyse the tumor cells that express both targets. Such molecules can be generated by antibody formats known in the art and are fully described. (WO2015/1842071, WO2015/158636, WO2010/136172, WO2013/174873). In a trispecific embodiment provided herein the antigen-binding polypeptide is bispecific for PSMA and a second distinct antigen on a tumor cell and additionally specific for an effector cell, in particular a T cell or a NK cell.

[00347] In addition to methods described above, bispecific antibodies provided herein can be generated *in vitro* in a cell-free environment by introducing asymmetrical mutations in the CH3 regions of two mono specific homodimeric antibodies and forming the bispecific heterodimeric antibody from two parent monospecific homodimeric antibodies in reducing conditions to allow disulfide bond isomerization according to methods described in PCT Pat. Publ. No. WO2011/131746. In the methods, the first monospecific bivalent antibody and the second monospecific bivalent antibody are engineered to have certain substitutions at the CH3 domain that promotes heterodimer stability; the antibodies are incubated together under reducing conditions sufficient to allow the cysteines in the hinge region to undergo disulfide bond isomerization; thereby generating the bispecific antibody by Fab arm exchange. The incubation conditions can optionally be restored to non-reducing conditions. Exemplary reducing agents that can be used are 2-mercaptoethylamine (2-MEA), dithiothreitol (DTT), dithioerythritol (DTE), glutathione, tris (2-carboxyethyl) phosphine (TCEP), L-cysteine and beta-mercaptoethanol, such as a reducing agent selected from the group consisting of: 2-mercaptoethylamine, dithiothreitol and tris (2-carboxyethyl) phosphine. For example, incubation for at least 90 min at a temperature of at least 20°C in the presence of at least 25 mM 2-MEA or in the presence of at least 0.5 mM dithiothreitol at a pH from 5-8, for example at pH of 7.0 or at pH of 7.4 can be used.

[00348] In some embodiments described herein, the bispecific antibody comprising a first domain specifically binding PSMA and a second domain specifically binding CD3 comprises at least one substitution in an antibody CH3 constant domain. In some embodiments, the at least one substitution in the antibody CH3 constant domain is 409R, F405L or F405L and R409K substitution, wherein residue numbering is according to the EU Index. Antibody domains and numbering are well known. "Asymmetrical" refers to non-identical substitutions in the two CH3 domains in two separate heavy chains in an antibody.

An IgG1 CH3 region typically consists of residues 341-446 on IgG1 (residue numbering according to the EU index). In some embodiments described herein, the bispecific PSMAxCD3 antibody comprises a F405L substitution in an antibody first heavy chain (HC1) and a 409R substitution in an antibody second heavy chain (HC2). In some embodiments, the bispecific PSMAxCD3 antibody comprises a S228P substitution in the HC1 and S228P, F405L and R409K substitutions in the HC2, wherein the antibody is of IgG4 isotype. In some embodiments, the HC1 contains the first domain specifically binding PSMA and the HC2 contains the second domain specifically binding CD3. In some embodiments, the bispecific antibody comprises at least one, two, three, four, five, six, seven or eight asymmetrical substitutions in the HC1 and the HC2 at residue positions 350, 366, 368, 370, 399, 405, 407 or 409, when residue numbering is according to the EU index. In some embodiments, the bispecific antibody comprises at least one, two, three or four asymmetrical substitutions in the HC1 and the HC2 at residue positions 350, 370, 405 or 409, when residue numbering is according to the EU index. In some embodiments, the bispecific antibody comprises at least one asymmetrical substitution in the HC1 and the HC2 at residue positions 405 or 409, when residue numbering is according to the EU index. In some embodiments, the bispecific antibody comprises a 409R or a F405L substitution in the HC1 and a 409R or a F405L substitution in the HC2, wherein residue numbering is according to the EU index. In some embodiments, the bispecific antibody comprises the F405L substitution in the HC1 and the 409R substitution in the HC2. In some embodiments, the bispecific antibody comprises at least one asymmetrical substitution in the HC1 and the HC2 at residue positions 366, 368, 370, 399, 405, 407 or 409, wherein residue numbering is according to the EU index. In some embodiments described herein, the HC1 position 409 has an amino acid substitution other than Lys, Leu or Met and the HC2 position 405 has an amino acid substitution other than Phe. In some embodiments described herein, the HC1 position 405 has an amino acid substitution other than Phe and the HC2 position 409 has an amino acid substitution other than Lys, Leu or Met. In some embodiments described herein, the HC1 position 409 has an amino acid substitution other than Lys, Leu or Met and the HC2 position 405 has an amino acid substitution other than Phe, Arg or Gly. In some embodiments described herein, the HC1 position 405 has an amino acid substitution other than Phe, Arg or Gly and the HC2 CH3 position 409 has an amino acid substitution other than Lys, Leu or Met. In some embodiments described herein, the HC1 CH3 has Phe at position 405 and an amino acid other than Lys, Leu or Met at position 409 and the HC2 has an amino acid other than Phe at position 405 and a Lys at position 409. In some

embodiments described herein, the HC1 has an amino acid other than Phe at position 405 and Lys at position 409 and the HC2 has Phe at position 405 and an amino acid other than Lys, Leu or Met at position 409. In some embodiments described herein, the HC1 has Phe at position 405 and an amino acid other than Lys, Leu or Met at position 409 and the HC2 has a substitution other than Phe, Arg or Gly at position 405 and Lys at position 409. In some embodiments described herein, the HC1 has a substitution other than Phe, Arg or Gly at position 405 and Lys at position 409 and the HC2 has Phe at position 405 and an amino acid other than Lys, Leu or Met at position 409. In some embodiments described herein, the HC1 has Phe at position 405 and an amino acid other than Lys, Leu or Met at position 409 and the HC2 has Leu at position 405 and Lys at position 409. In some embodiments described herein, the HC1 has Leu at position 405 and Lys at position 409 and the HC2 has Phe at position 405 and an amino acid other than Lys, Leu or Met at position 409. In some embodiments described herein, the HC1 has Phe at position 405 and Arg at position 409 and the HC2 has an amino acid other than Phe, Arg or Gly at position 405 and Lys at position 409. In some embodiments described herein, the HC1 has an amino acid other than Phe, Arg or Gly at position 405 and Lys at position 409 and the HC2 has Phe at position 405 and Arg at position 409. In some embodiments described herein, the HC1 has Phe at position 405 and Arg at position 409 and the HC2 has Leu at position 405 and Lys at position 409. In some embodiments described herein, the HC1 has Leu at position 405 and Lys at position 409 and the HC2 has Phe at position 405 and Arg at position 409. In some embodiments described herein, the HC1 has Phe at position 405 and Lys at position 409 and the HC2 has Leu at position 405 and Arg at position 409. In some embodiments described herein, the HC1 has Leu at position 405 and Arg at position 409 and the HC2 has Phe at position 405 and Lys at position 409. In some embodiments described herein, the HC1 has an amino acid other than Lys, Leu or Met at position 409 and the HC2 has Lys at position 409, Thr at position 370 and Leu at position 405. In some embodiments described herein, the HC1 has Lys at position 409, Thr at position 370 and Leu at position 405 and the HC2 has an amino acid other than Lys, Leu or Met at position 409. In some embodiments described herein, the HC1 has Arg at position 409 and the HC2 has Lys at position 409, Thr at position 370 and Leu at position 405. In some embodiments described herein, the HC1 has Lys at position 409, Thr at position 370 and Leu at position 405 and the HC2 has Arg at position 409. In some embodiments described herein, the HC1 has Lys at position 370, Phe at position 405 and Arg at position 409 and the HC2 has Lys at position 409, Thr at position 370 and Leu at position 405. In some embodiments described herein, the HC1 has Lys at position 409, Thr

at position 370 and Leu at position 405 and the HC2 has Lys at position 370, Phe at position 405 and Arg at position 409. In some embodiments described herein, the HC1 has an amino acid other than Lys, Leu or Met at position 409 and the HC2 has an amino acid other than Tyr, Asp, Glu, Phe, Lys, Gln, Arg, Ser or Thr at position 407. In some embodiments described herein, the HC1 has an amino acid other than Tyr, Asp, Glu, Phe, Lys, Gln, Arg, Ser or Thr at position 407 and the HC2 has an amino acid other than Lys, Leu or Met at position 409. In some embodiments described herein, the HC1 has an amino acid other than Lys, Leu or Met at position 409 and the HC2 has Ala, Gly, His, Ile, Leu, Met, Asn, Val or Trp at position 407. In some embodiments described herein, the HC1 has Ala, Gly, His, Ile, Leu, Met, Asn, Val or Trp at position 407 and the HC2 has an amino acid other than Lys, Leu or Met at position 409. In some embodiments described herein, the HC1 has an amino acid other than Lys, Leu or Met at position 409 and the HC2 has Gly, Leu, Met, Asn or Trp at position 407. In some embodiments described herein, the HC1 has Gly, Leu, Met, Asn or Trp at position 407 and the HC2 has an amino acid other than Lys, Leu or Met at position 409. In some embodiments described herein, the HC1 has Tyr at position 407 and an amino acid other than Lys, Leu or Met at position 409 and the HC2 has an amino acid other than Tyr, Asp, Glu, Phe, Lys, Gln, Arg, Ser or Thr at position 407 and Lys at position 409. In some embodiments described herein, the HC1 has an amino acid other than Tyr, Asp, Glu, Phe, Lys, Gln, Arg, Ser or Thr at position 407 and Lys at position 409 and the HC2 has Tyr at position 407 and an amino acid other than Lys, Leu or Met at position 409. In some embodiments described herein, the HC1 has Tyr at position 407 and an amino acid other than Lys, Leu or Met at position 409 and the HC2 has Ala, Gly, His, Ile, Leu, Met, Asn, Val or Trp at position 407 and Lys at position 409. In some embodiments described herein, the HC1 has Ala, Gly, His, Ile, Leu, Met, Asn, Val or Trp at position 407 and Lys at position 409 and the HC2 CH3 has Tyr at position 407 and an amino acid other than Lys, Leu or Met at position 409. In some embodiments described herein, the HC1 CH3 has Tyr at position 407 and an amino acid other than Lys, Leu or Met at position 409 and the HC2 has Gly, Leu, Met, Asn or Trp at position 407 and Lys at position 409. In some embodiments described herein, the HC1 has Gly, Leu, Met, Asn or Trp at position 407 and Lys at position 409 and the HC2 has Tyr at position 407 and an amino acid other than Lys, Leu or Met at position 409. In some embodiments described herein, the HC1 has Tyr at position 407 and Arg at position 409 and the HC2 has an amino acid other than Tyr, Asp, Glu, Phe, Lys, Gln, Arg, Ser or Thr at position 407 and Lys at position 409. In some embodiments described herein, the HC1 has an amino acid other than Tyr, Asp, Glu, Phe, Lys, Gln, Arg, Ser or Thr at

position 407 and Lys at position 409 and the HC2 has Tyr at position 407 and Arg at position 409. In some embodiments described herein, the HC1 has Tyr at position 407 and Arg at position 409 and the HC2 has Ala, Gly, His, Ile, Leu, Met, Asn, Val or Trp at position 407 and Lys at position 409. In some embodiments described herein, the HC1 has Ala, Gly, His, Ile, Leu, Met, Asn, Val or Trp at position 407 and Lys at position 409 and the HC2 has Tyr at position 407 and Arg at position 409. In some embodiments described herein, the HC1 CH3 has Tyr at position 407 and Arg at position 409 and the HC2 CH3 has Gly, Leu, Met, Asn or Trp at position 407 and Lys at position 409. In some embodiments described herein, the HC1 has Gly, Leu, Met, Asn or Trp at position 407 and Lys at position 409 and the HC2 has Tyr at position 407 and Arg at position 409. In some embodiments described herein, the HC1 has an amino acid other than Lys, Leu or Met at position 409, and the HC2 has (i) an amino acid other than Phe, Leu and Met at position 368, or (ii) a Trp at position 370, or (iii) an amino acid other than Asp, Cys, Pro, Glu or Gln at position 399. In some embodiments described herein, the HC1 has (i) an amino acid other than Phe, Leu and Met at position 368, or (ii) a Trp at position 370, or (iii) an amino acid other than Asp, Cys, Pro, Glu or Gln at position 399 and the HC2 has an amino acid other than Lys, Leu or Met at position 409. In some embodiments described herein, the HC1 has Arg, Ala, His or Gly at position 409, and the HC2 has (i) Lys, Gln, Ala, Asp, Glu, Gly, His, Ile, Asn, Arg, Ser, Thr, Val, or Trp at position 368, or (ii) Trp at position 370, or (iii) Ala, Gly, Ile, Leu, Met, Asn, Ser, Thr, Trp, Phe, His, Lys, Arg or Tyr at position 399. In some embodiments described herein, the HC1 has (i) Lys, Gln, Ala, Asp, Glu, Gly, His, Ile, Asn, Arg, Ser, Thr, Val, or Trp at position 368, or (ii) Trp at position 370, or (iii) Ala, Gly, Ile, Leu, Met, Asn, Ser, Thr, Trp, Phe, His, Lys, Arg or Tyr at position 399 and the HC2 has Arg, Ala, His or Gly at position 409. In some embodiments described herein, the HC1 has Arg at position 409, and the HC2 has (i) Asp, Glu, Gly, Asn, Arg, Ser, Thr, Val, or Trp at position 368, or (ii) Trp at position 370, or (iii) Phe, His, Lys, Arg or Tyr at position 399. In some embodiments described herein, the HC1 has (i) Asp, Glu, Gly, Asn, Arg, Ser, Thr, Val, or Trp at position 368, or (ii) Trp at position 370, or (iii) Phe, His, Lys, Arg or Tyr at position 399 and the HC2 has Arg at position 409. In some embodiments described herein, the HC1 comprises a 409R substitution or a F405L substitution and the HC2 comprises a 409R substitution or a F405L substitution, wherein residue numbering is according to the EU index. In some embodiments described herein, the HC1 comprises the F405L substitution and the HC2 comprises the 409R substitution.

[00349] Substitutions are typically made at the DNA level to a molecule such as the constant domain of the antibody using standard methods.

[00350] The antibodies provided herein can be engineered into various well-known antibody forms.

[00351] In some embodiments, the bispecific antibody is a diabody or a cross-body.

[00352] In some embodiments, the multispecific antibodies include IgG-like molecules with complementary CH3 domains that promote heterodimerization; recombinant IgG-like dual targeting molecules, wherein the two sides of the molecule each contain the Fab fragment or part of the Fab fragment of at least two different antibodies; IgG fusion molecules, wherein full length IgG antibodies are fused to an extra Fab fragment or parts of Fab fragment; Fc fusion molecules, wherein single chain Fv molecules or stabilized diabodies are fused to heavy-chain constant-domains, Fc-regions or parts thereof; Fab fusion molecules, wherein different Fab-fragments are fused together; ScFv- and diabody-based and heavy chain antibodies (*e.g.*, domain antibodies, nanobodies) wherein different single chain Fv molecules or different diabodies or different heavy-chain antibodies (*e.g.* domain antibodies, nanobodies) are fused to each other or to another protein or carrier molecule.

[00353] In some embodiments, recombinant IgG-like dual targeting molecules include Dual Targeting (DT)-Ig (GSK/Domantis), Two-in-one Antibody (Genentech), Cross-linked Mabs (Karmanos Cancer Center), mAb2 (F-Star) and CovX-body (CovX/Pfizer).

[00354] In some embodiments, IgG fusion molecules include Dual Variable Domain (DVD)-Ig (Abbott), IgG-like Bispecific (ImClone/Eli Lilly), Ts2Ab (MedImmune/AZ) and BsAb (Zymogenetics), HERCULES (Biogen Idec) and TvAb (Roche).

[00355] In some embodiments, Fc fusion molecules can include ScFv/Fc Fusions (Academic Institution), SCORPION (Emergent BioSolutions/Trubion, Zymogenetics/BMS), Dual Affinity Retargeting Technology (Fc-DART) (MacroGenics) and Dual(ScFv)₂-Fab (National Research Center for Antibody Medicine--China).

[00356] In some embodiments, Fab fusion bispecific antibodies include F(ab)₂ (Medarex/AMGEN), Dual-Action or Bis-Fab (Genentech), Dock-and-Lock (DNL) (ImmunoMedics), Bivalent Bispecific (Biotecol) and Fab-Fv (UCB-Celltech). ScFv-, diabody-based, and domain antibodies, include but are not limited to, Bispecific T Cell Engager (BiTE) (Micromet), Tandem Diabody (Tandab) (Affimed), Dual Affinity Retargeting Technology (DART) (MacroGenics), Single-chain Diabody (Academic), TCR-

like Antibodies (AIT, ReceptorLogics), Human Serum Albumin ScFv Fusion (Merrimack) and COMBODY (Epigen Biotech), dual targeting nanobodies (Ablynx), dual targeting heavy chain only domain antibodies. Various formats of bispecific antibodies have been described, for example in Chames and Baty (2009) *Curr Opin Drug Disc Dev* 12: 276 and in Nunez-Prado *et al.*, (2015) *Drug Discovery Today* 20(5):588-594.

[00357] Any of the VH and the VL domains identified herein (*e.g.*, those that bind PSMA) can be engineered into scFv format. In some embodiments, the scFv format is in the VH-linker-VL orientation. In other embodiments, the scFv format is in the VL-linker-VH orientation. Any of the VH and the VL domains identified herein can also be used to generate sc(Fv)₂ structures. In some embodiments, the sc(Fv)₂ structure is VH-linker-VL-linker-VL-linker-VH. In some embodiments, the sc(Fv)₂ structure is VH-linker-VL-linker-VH-linker-VL. In some embodiments, the sc(Fv)₂ structure is VH-linker-VH-linker-VL-linker-VL. In some embodiments, the sc(Fv)₂ structure is VL-linker-VH-linker-VH-linker-VL. In some embodiments, the sc(Fv)₂ structure is VL-linker-VH-linker-VL-linker-VH. In some embodiments, the sc(Fv)₂ structure is VL-linker-VL-linker-VH-linker-VH.

[00358] In specific embodiments, the linker is a peptide linker. In some embodiments, the linker comprises a naturally occurring amino acid. Exemplary amino acids that can be included into the linker are Gly, Ser Pro, Thr, Glu, Lys, Arg, Ile, Leu, His and The. In some embodiments, the linker has a length that is adequate to link the VH and the VL in such a way that they form the correct conformation relative to one another so that they retain the desired activity, such as binding to the target (*e.g.*, PSMA).

[00359] In certain embodiments, the linker is about 5-50 amino acids long. In some embodiments, the linker is about 10-40 amino acids long. In some embodiments, the linker is about 10-35 amino acids long. In some embodiments, the linker is about 10-30 amino acids long. In some embodiments, the linker is about 10-25 amino acids long. In some embodiments, the linker is about 10-20 amino acids long. In some embodiments, the linker is about 15-20 amino acids long. In some embodiments, the linker is 6 amino acids long. In some embodiments, the linker is 7 amino acids long. In some embodiments, the linker is 8 amino acids long. In some embodiments, the linker is 9 amino acids long. In some embodiments, the linker is 10 amino acids long. In some embodiments, the linker is 11 amino acids long. In some embodiments, the linker is 12 amino acids long. In some embodiments, the linker is 13 amino acids long. In some embodiments, the linker is 14 amino acids long. In some embodiments, the linker is 15 amino acids long. In some

embodiments, the linker is 16 amino acids long. In some embodiments, the linker is 17 amino acids long. In some embodiments, the linker is 18 amino acids long. In some embodiments, the linker is 19 amino acids long. In some embodiments, the linker is 20 amino acids long. In some embodiments, the linker is 21 amino acids long. In some embodiments, the linker is 22 amino acids long. In some embodiments, the linker is 23 amino acids long. In some embodiments, the linker is 24 amino acids long. In some embodiments, the linker is 25 amino acids long. In some embodiments, the linker is 26 amino acids long. In some embodiments, the linker is 27 amino acids long. In some embodiments, the linker is 28 amino acids long. In some embodiments, the linker is 29 amino acids long. In some embodiments, the linker is 30 amino acids long. In some embodiments, the linker is 31 amino acids long. In some embodiments, the linker is 32 amino acids long. In some embodiments, the linker is 33 amino acids long. In some embodiments, the linker is 34 amino acids long. In some embodiments, the linker is 35 amino acids long. In some embodiments, the linker is 36 amino acids long. In some embodiments, the linker is 37 amino acids long. In some embodiments, the linker is 38 amino acids long. In some embodiments, the linker is 39 amino acids long. In some embodiments, the linker is 40 amino acids long. Exemplary linkers that can be used are Gly rich linkers, Gly and Ser containing linkers, Gly and Ala containing linkers, Ala and Ser containing linkers, and other flexible linkers.

[00360] Exemplary linkers that can be used include any one of SEQ ID NOs.1419-1452. Exemplary linkers are shown in **Table 2**. Additional linkers are described for example in Int. Pat. Publ. No. WO2019/060695. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1419. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1420. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1421. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1422. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1423. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1424. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1425. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1426. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1427. In some embodiments, the linker comprises or consists of the amino acid

sequence of SEQ ID NO:1428. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1429. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1430. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1431. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1432. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1433. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1434. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1435. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1436. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1437. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1438. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1439. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1440. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1441. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1442. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1427. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1444. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1445. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1446. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1435. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1448. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1449. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1450. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1451. In some embodiments, the linker comprises or consists of the amino acid sequence of SEQ ID NO:1452.

Table 2: Exemplary Linkers

Linker name	Amino acid sequence	SEQ ID NO:
-------------	---------------------	------------

Linker 1	GGSEKSSSGSGSESKSTGGS	1419
Linker 2	GGSGGGS	1420
Linker 3	GGSGGGS	1421
Linker 4	GGSGGGS	1422
Linker 5	GGSGGGS	1423
Linker 6	GGSGGGS	1424
Linker 7	GGSGGGS	1425
Linker 8	GGSGGGS	1426
Linker 9	GSTSGSGKPGSGEGSTKG	1427
Linker 10	IRPRAIGGSKPRVA	1428
Linker 11	GKGGSGKGGSGKGG	1429
Linker 12	GKGGSGKGGSGKGG	1430
Linker 13	GKGGSGKGGSGKGG	1431
Linker 14	GKGGSGKGGSGKGG	1432
Linker 15	GKGGSGKGGSGKGG	1433
Linker 16	GKPGSGKPGSGKPGS	1434
Linker 17	GKPGSGKPGSGKPGS	1435
Linker 18	GKPGSGKPGSGKPGS	1436
Linker 19	STAGDTHLGGEDFD	1437
Linker 20	GEGSGEGSGEGSG	1438
Linker 21	GEGSGEGSGEGSG	1439
Linker 22	GEGSGEGSGEGSG	1440
Linker 23	GEGSGEGSGEGSG	1441
Linker 24	GEGSGEGSGEGSG	1442
Linker 25	GSTSGSGKPGSGEGSTKG	1427
Linker 26	PRGASKSGSASQTGSAPGS	1444
Linker 27	GTAAAGAGAAGGAAAGAAG	1445
Linker 28	GTSGSSGSGSGSGSGGGG	1446
Linker 29	GKPGSGKPGSGKPGSGKPGS	1435
Linker 30	GSGS	1448
Linker 31	APAPAPAPAP	1449
Linker 32	APAPAPAPAPAPAPAPAPAP	1450
Linker 33	AEAAAKEAAAKEAAAKEAAAKEAAAKAAA	1451
Linker 34	GTEGKSSSGSGSESKST	1452

[00361] In some embodiments, the scFv comprises, from the N- to C-terminus, a VH, a first linker (L1) and a VL (VH-L1-VL). In other embodiments, the scFv comprises, from the N- to C-terminus, the VL, the L1 and the VH (VL-L1-VH). In another embodiment, the L1 comprises the amino acid sequence of SEQ ID NO: 1419. In one embodiment, the L1 comprises the amino acid sequence of SEQ ID NO: 1420. In other embodiments, the L1 comprises the amino acid sequence of SEQ ID NO: 1421. In some embodiments, the L1

comprises the amino acid sequence of SEQ ID NO:1422. In one embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1423. In some embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1424. In other embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1425. In one embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1426. In some embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1427. In another embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1428. In one embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1429. In other embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1430. In some embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1431. In some embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1432. In one embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1433. In other embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1434. In one embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1435. In another embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1436. In some embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1437. In one embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1438. In other embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1439. In some embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1440. In another embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1441. In one embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1442. In other embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1427. In one embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1444. In some embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1445. In other embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1446. In one embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1435. In another embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1448. In one embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1449. In other embodiments, the L1 comprises the amino acid sequence of SEQ ID NO:1450. In another embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1451. In one embodiment, the L1 comprises the amino acid sequence of SEQ ID NO:1452.

[00362] In certain embodiments, antibodies, including provided herein comprise two linkers. In other embodiments antibodies provided herein comprise three linkers. In yet

other embodiments, antibodies provided herein comprise four or more linkers. In certain embodiments, the antibody is an antigen binding fragment thereof.

Polynucleotides, vectors and host cells

[00363] Also provided is a nucleic acid encoding an antibody provided herein. In another general aspect, provide is a vector comprising an isolated nucleic acid encoding an antibody provided herein. In another general aspect, provided is a vector comprising an isolated nucleic acid encoding an antibody provided herein. Also provided is a vector comprising a nucleic acid encoding an antibody provided herein. Also provided is a host cell comprising a vector comprising a nucleic acid encoding an antibody provided herein. Also provided is a kit comprising the vector comprising a nucleic acid encoding an antibody provided herein, and packaging for the same. In another general aspect, provided herein is an isolated nucleic acid encoding a monoclonal antibody or antigen-binding fragment thereof provided herein. In certain embodiments, the antibody is a PSMA antibody. In certain embodiments, the antibody is a multispecific PSMA antibody. In some embodiments, the antibody is a multispecific PSMAxCD3 antibody. Also provided is a nucleic acid encoding a bispecific antibody comprising: (a) a first binding domain that binds to PSMA, and (b) a second binding domain that binds to a second target that is not PSMA, as provided herein.

[00364] Also provided is a nucleic acid encoding a multispecific PSMAxCD3 antibody provided herein. In another general aspect, provide is a vector comprising an isolated nucleic acid encoding a multispecific PSMAxCD3 antibody provided herein. In another general aspect, provided is a vector comprising an isolated nucleic acid encoding a multispecific PSMAxCD3 antibody provided herein. Also provided is a vector comprising a nucleic acid encoding a multispecific PSMAxCD3 antibody provided herein. Also provided is a host cell comprising a vector comprising a nucleic acid encoding a multispecific PSMAxCD3 antibody provided herein. Also provided is a kit comprising the vector comprising a nucleic acid encoding a multispecific PSMAxCD3 antibody provided herein, and packaging for the same. Also provided is an isolated nucleic acid encoding a monoclonal antibody provided herein. Also provided is an isolated nucleic acid encoding an antigen binding fragment provided herein. Also provided is a nucleic acid encoding a multispecific antibody comprising: (a) a first binding domain that binds to PSMA, and (b) a second binding domain that binds to a CD3, as provided herein. In certain embodiments, the multispecific PSMAxCD3 antibody is a trispecific antibody. In certain embodiments, the multispecific PSMAxCD3 antibody is a quadraspecific antibody.

[00365] Also disclosed are isolated polynucleotides that encode the antibodies or antigen-binding fragments that immunospecifically bind to PSMA. The isolated polynucleotides capable of encoding the variable domain segments provided herein can be included on the same, or different, vectors to produce antibodies or antigen-binding fragments.

[00366] Polynucleotides encoding recombinant antigen-binding proteins also are within the scope of the disclosure. In some embodiments, the polynucleotides described (and the peptides they encode) include a leader sequence. Any leader sequence known in the art can be employed. The leader sequence can include, but is not limited to, a restriction site or a translation start site.

[00367] The PSMA-specific antibodies or antigen-binding fragments described herein include variants having single or multiple amino acid substitutions, deletions, or additions that retain the biological properties (*e.g.*, binding affinity or immune effector activity) of the described PSMA-specific antibodies or antigen-binding fragments. In certain embodiments, the following notations are, unless otherwise indicated, used to describe a mutation; i) substitution of an amino acid in a given position is written as *e.g.* K409R which means a substitution of a Lysine in position 409 with an Arginine; and ii) for specific variants the specific three or one letter codes are used, including the codes Xaa and X to indicate any amino acid residue. Thus, the substitution of Arginine for Lysine in position 409 is designated as: K409R, or the substitution of any amino acid residue for Lysine in position 409 is designated as K409X. In case of deletion of Lysine in position 409 it is indicated by K409*. Where a particular amino acid residue may vary among peptide isotypes or variants, and a substitution affects that residue in each isotype or variant or any of the isotypes or variants, the substitution is designated as, *e.g.*, 409R, meaning the amino acid corresponding to position 409 is substituted with Arginine. The skilled person can produce variants having single or multiple amino acid substitutions, deletions, or additions.

[00368] In some embodiments, these variants are: (a) variants in which one or more amino acid residues are substituted with conservative or nonconservative amino acids, (b) variants in which one or more amino acids are added to or deleted from the polypeptide, (c) variants in which one or more amino acids include a substituent group, and (d) variants in which the polypeptide is fused with another peptide or polypeptide such as a fusion partner, a protein tag or other chemical moiety, that can confer useful properties to the polypeptide, such as, for example, an epitope for an antibody, a polyhistidine sequence, a biotin moiety and the like. Antibodies or antigen-binding fragments described herein can include variants in

which amino acid residues from one species are substituted for the corresponding residue in another species, either at the conserved or nonconserved positions. In other embodiments, amino acid residues at nonconserved positions are substituted with conservative or nonconservative residues. The techniques for obtaining these variants, including genetic (deletions, mutations, etc.), chemical, and enzymatic techniques, are known to persons having ordinary skill in the art.

[00369] The PSMA-specific antibodies or antigen-binding fragments described herein can embody several antibody isotypes, such as IgM, IgD, IgG, IgA and IgE. In some embodiments the antibody isotype is IgG1, IgG2, IgG3, or IgG4 isotype, such as IgG1 or IgG4 isotype. Antibody or antigen-binding fragment thereof specificity is largely determined by the amino acid sequence, and arrangement, of the CDRs. Therefore, the CDRs of one isotype can be transferred to another isotype without altering antigen specificity. Alternatively, techniques have been established to cause hybridomas to switch from producing one antibody isotype to another (isotype switching) without altering antigen specificity. Accordingly, such antibody isotypes are within the scope of the described antibodies or antigen-binding fragments.

[00370] It will be appreciated by those skilled in the art that the coding sequence of a protein can be changed (*e.g.*, replaced, deleted, inserted, *etc.*) without changing the amino acid sequence of the protein. Accordingly, it will be understood by those skilled in the art that nucleic acid sequences encoding antibodies provided herein can be altered without changing the amino acid sequences of the proteins.

[00371] Any vector known to those skilled in the art in view of the present disclosure can be used, such as a plasmid, a cosmid, a phage vector or a viral vector. In some embodiments, the vector is a recombinant expression vector such as a plasmid. The vector can include any element to establish a conventional function of an expression vector, for example, a promoter, ribosome binding element, terminator, enhancer, selection marker, and origin of replication. The promoter can be a constitutive, inducible or repressible promoter. A number of expression vectors capable of delivering nucleic acids to a cell are known in the art and can be used herein for production of an antibody or antigen-binding fragment thereof in the cell. Conventional cloning techniques or artificial gene synthesis can be used to generate a recombinant expression vector according to certain embodiments. Such techniques are well known to those skilled in the art in view of the present disclosure.

[00372] Also provided, are vectors comprising the polynucleotides described herein. The vectors can be expression vectors. Recombinant expression vectors containing a sequence encoding a polypeptide of interest are thus contemplated as within the scope of this disclosure. The expression vector can contain one or more additional sequences such as but not limited to regulatory sequences (*e.g.*, promoter, enhancer), a selection marker, and a polyadenylation signal. Vectors for transforming a wide variety of host cells are well known and include, but are not limited to, plasmids, phagemids, cosmids, baculoviruses, bacmids, bacterial artificial chromosomes (BACs), yeast artificial chromosomes (YACs), as well as other bacterial, yeast and viral vectors.

[00373] Recombinant expression vectors within the scope of the description include synthetic, genomic, or cDNA-derived nucleic acid fragments that encode at least one recombinant protein which can be operably linked to suitable regulatory elements. Such regulatory elements can include a transcriptional promoter, sequences encoding suitable mRNA ribosomal binding sites, and sequences that control the termination of transcription and translation. Expression vectors, especially mammalian expression vectors, can also include one or more nontranscribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, other 5' or 3' flanking nontranscribed sequences, 5' or 3' nontranslated sequences (such as necessary ribosome binding sites), a polyadenylation site, splice donor and acceptor sites, or transcriptional termination sequences. An origin of replication that confers the ability to replicate in a host can also be incorporated.

[00374] The transcriptional and translational control sequences in expression vectors to be used in transforming vertebrate cells can be provided by viral sources. Exemplary vectors can be constructed as described by Okayama and Berg, 3 Mol. Cell. Biol. 280 (1983).

[00375] In some embodiments, the antibody- or antigen-binding fragment-coding sequence is placed under control of a powerful constitutive promoter, such as the promoters for the following genes: hypoxanthine phosphoribosyl transferase (HPRT), adenosine deaminase, pyruvate kinase, beta-actin, human myosin, human hemoglobin, human muscle creatine, and others. In addition, many viral promoters function constitutively in eukaryotic cells and are suitable for use with the described embodiments. Such viral promoters include without limitation, Cytomegalovirus (CMV) immediate early promoter, the early and late promoters of SV40, the Mouse Mammary Tumor Virus (MMTV) promoter, the long terminal repeats (LTRs) of Moloney leukemia virus, Human Immunodeficiency Virus

(HIV), Epstein Barr Virus (EBV), Rous Sarcoma Virus (RSV), and other retroviruses, and the thymidine kinase promoter of Herpes Simplex Virus. In one embodiment, the PSMA-specific antibody or antigen-binding fragment thereof coding sequence is placed under control of an inducible promoter such as the metallothionein promoter, tetracycline-inducible promoter, doxycycline-inducible promoter, promoters that contain one or more interferon-stimulated response elements (ISRE) such as protein kinase R 2',5'-oligoadenylate synthetases, Mx genes, ADAR1, and the like.

[00376] Vectors described herein can contain one or more Internal Ribosome Entry Site(s) (IRES). Inclusion of an IRES sequence into fusion vectors can be beneficial for enhancing expression of some proteins. In some embodiments the vector system will include one or more polyadenylation sites (*e.g.*, SV40), which can be upstream or downstream of any of the aforementioned nucleic acid sequences. Vector components can be contiguously linked, or arranged in a manner that provides optimal spacing for expressing the gene products (*i.e.*, by the introduction of "spacer" nucleotides between the ORFs), or positioned in another way. Regulatory elements, such as the IRES motif, can also be arranged to provide optimal spacing for expression.

[00377] The vectors can comprise selection markers, which are well known in the art. Selection markers include positive and negative selection markers, for example, antibiotic resistance genes (*e.g.*, neomycin resistance gene, a hygromycin resistance gene, a kanamycin resistance gene, a tetracycline resistance gene, a penicillin resistance gene), glutamate synthase genes, HSV-TK, HSV-TK derivatives for ganciclovir selection, or bacterial purine nucleoside phosphorylase gene for 6-methylpurine selection (Gadi *et al.*, 7 Gene Ther. 1738-1743 (2000)). A nucleic acid sequence encoding a selection marker or the cloning site can be upstream or downstream of a nucleic acid sequence encoding a polypeptide of interest or cloning site.

[00378] The vectors described herein can be used to transform various cells with the genes encoding the described antibodies or antigen-binding fragments. For example, the vectors can be used to generate PSMA-specific antibody or antigen-binding fragment-producing cells. Thus, another aspect features host cells transformed with vectors comprising a nucleic acid sequence encoding an antibody or antigen-binding fragment thereof that specifically binds PSMA, such as the antibodies or antigen-binding fragments described and exemplified herein.

[00379] Numerous techniques are known in the art for the introduction of foreign genes into cells and can be used to construct the recombinant cells for purposes of carrying out the described methods, in accordance with the various embodiments described and exemplified herein. The technique used should provide for the stable transfer of the heterologous gene sequence to the host cell, such that the heterologous gene sequence is heritable and expressible by the cell progeny, and so that the necessary development and physiological functions of the recipient cells are not disrupted. Techniques which can be used include but are not limited to chromosome transfer (*e.g.*, cell fusion, chromosome mediated gene transfer, micro cell mediated gene transfer), physical methods (*e.g.*, transfection, spheroplast fusion, microinjection, electroporation, liposome carrier), viral vector transfer (*e.g.*, recombinant DNA viruses, recombinant RNA viruses) and the like (described in Cline, 29 Pharmac. Ther. 69-92 (1985)). Calcium phosphate precipitation and polyethylene glycol (PEG)-induced fusion of bacterial protoplasts with mammalian cells can also be used to transform cells.

[00380] Also provided is a host cell comprising an isolated nucleic acid encoding an antibody provided herein. Also provided is a host cell comprising an isolated nucleic acid encoding an antigen binding fragment provided herein. Any host cell known to those skilled in the art in view of the present disclosure can be used for recombinant expression of antibodies or antigen-binding fragments thereof provided herein. In some embodiments, the host cells are *E. coli* TG1 or BL21 cells (for expression of, *e.g.*, an scFv or Fab antibody), CHO-DG44 or CHO-K1 cells or HEK293 cells (for expression of, *e.g.*, a full-length IgG antibody). Other cells suitable for use in the expression of the PSMA-specific antibodies or antigen-binding fragments described herein can be eukaryotic cells, such as cells of plant, rodent, or human origin, for example but not limited to NSO, CHO, CHOK1, perC.6, Tk-ts13, BHK, HEK293 cells, COS-7, T98G, CV-1/EBNA, L cells, C127, 3T3, HeLa, NS1, Sp2/0 myeloma cells, and BHK cell lines, among others. In addition, expression of antibodies can be accomplished using hybridoma cells. Methods for producing hybridomas are well established in the art.

[00381] According to particular embodiments, the recombinant expression vector is transformed into host cells by conventional methods such as chemical transfection, heat shock, or electroporation, where it is stably integrated into the host cell genome such that the recombinant nucleic acid is effectively expressed.

[00382] Cells transformed with expression vectors described herein can be selected or screened for recombinant expression of the antibodies or antigen-binding fragments described herein. Recombinant-positive cells are expanded and screened for subclones exhibiting a desired phenotype, such as high level expression, enhanced growth properties, or the ability to yield proteins with desired biochemical characteristics, for example, due to protein modification or altered post-translational modifications. These phenotypes can be due to inherent properties of a given subclone or to mutation. Mutations can be effected through the use of chemicals, UV-wavelength light, radiation, viruses, insertional mutagens, inhibition of DNA mismatch repair, or a combination of such methods.

[00383] Also provided are methods of producing an antibody disclosed herein. The methods comprise culturing a cell comprising a nucleic acid encoding the antibody under conditions to produce an antibody and recovering the antibody from the cell or cell culture (e.g., from the supernatant). Expressed antibodies can be harvested from the cells and purified according to conventional techniques known in the art and as described herein.

[00384] Also provided is a method of producing a multispecific antibody disclosed herein. The methods comprise culturing a cell comprising a nucleic acid encoding the multispecific antibody thereof under conditions to produce the multispecific antibody and recovering the multispecific antibody from the cell or cell culture (e.g., from the supernatant). Expressed multispecific antibodies thereof can be harvested from the cells and purified according to conventional techniques known in the art and as described herein. Other methods are described elsewhere herein.

Pharmaceutical Compositions

[00385] In another general aspect, provided is a pharmaceutical composition comprising a PSMA antibody provided herein and a pharmaceutically acceptable carrier. In certain embodiments, the antibody is isolated. Also provided is a method of producing the pharmaceutical composition, comprising combining the antibody with a pharmaceutically acceptable carrier to obtain the pharmaceutical composition.

[00386] In another general aspect, provided is a pharmaceutical composition comprising a PSMA multispecific antibody provided herein and a pharmaceutically acceptable carrier. In certain embodiments, the multispecific antibody is isolated. Also provided is a method of producing the pharmaceutical composition, comprising combining the multispecific antibody with a pharmaceutically acceptable carrier to obtain the pharmaceutical

composition. In another aspect, provided herein is a pharmaceutical composition comprising a comprising: (a) a first binding domain that binds to PSMA, and (b) a second binding domain that binds to a second target, and a pharmaceutically acceptable carrier. Any of the multispecific antibodies provided herein are contemplated in the pharmaceutical compositions. In certain embodiments, the second binding domain binds to CD3. Any of the antibodies provided herein are contemplated in the pharmaceutical compositions.

[00387] In another general aspect, provided is a pharmaceutical composition comprising a multispecific PSMAxCD3 antibody provided herein and a pharmaceutically acceptable carrier. In certain embodiments, the multispecific PSMAxCD3 antibody is isolated. Also provided is a method of producing the pharmaceutical composition, comprising combining the multispecific antibody with a pharmaceutically acceptable carrier to obtain the pharmaceutical composition. In another aspect, provided herein is a pharmaceutical composition comprising a comprising: (a) a first binding domain that binds to PSMA, and (b) a second binding domain that binds to CD3, and a pharmaceutically acceptable carrier. Any of the multispecific antibodies provided herein are contemplated in the pharmaceutical compositions.

[00388] The term “pharmaceutical composition” as used herein means a product comprising an antibody provided herein together with a pharmaceutically acceptable carrier. Antibodies of provided herein and compositions comprising them are also useful in the manufacture of a medicament for therapeutic applications.

[00389] As used herein, the term “carrier” refers to any excipient, diluent, filler, salt, buffer, stabilizer, solubilizer, oil, lipid, lipid containing vesicle, microsphere, liposomal encapsulation, or other material well known in the art for use in pharmaceutical formulations. It will be understood that the characteristics of the carrier, excipient or diluent will depend on the route of administration for a particular application. As used herein, the term “pharmaceutically acceptable carrier” refers to a non-toxic material that does not interfere with the effectiveness of a composition provided herein the biological activity of a composition provided herein. According to particular embodiments, in view of the present disclosure, any pharmaceutically acceptable carrier suitable for use in an antibody pharmaceutical composition can be used herein. Carrier can be liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. For example, 0.4% saline and 0.3% glycine can be used. These solutions are sterile and generally free of particulate matter. They can

be sterilized by conventional, well-known sterilization techniques (*e.g.*, filtration). The compositions can contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, stabilizing, thickening, lubricating and coloring agents, etc. The concentration of the antibodies provided herein in such pharmaceutical formulation can vary, from less than about 0.5%, usually to at least about 1% to as much as 15 or 20% by weight and can be selected primarily based on required dose, fluid volumes, viscosities, etc., according to the particular mode of administration selected. Suitable vehicles and formulations, inclusive of other human proteins, *e.g.*, human serum albumin. The formulation of pharmaceutically active ingredients with pharmaceutically acceptable carriers is known in the art, *e.g.*, Remington: The Science and Practice of Pharmacy (*e.g.* 21st edition (2005), and any later editions). Non-limiting examples of additional ingredients include: buffers, diluents, solvents, tonicity regulating agents, preservatives, stabilizers, and chelating agents. One or more pharmaceutically acceptable carriers can be used in formulating the pharmaceutical compositions provided herein.

[00390] In one embodiment, the pharmaceutical composition is a liquid formulation. An example of a liquid formulation is an aqueous formulation, *i.e.*, a formulation comprising water. The liquid formulation can comprise a solution, a suspension, an emulsion, a microemulsion, a gel, and the like. An aqueous formulation typically comprises at least 50% w/w water, or at least 60%, 70%, 75%, 80%, 85%, 90%, or at least 95% w/w of water.

[00391] In one embodiment, the pharmaceutical composition can be formulated as an injectable which can be injected, for example, via an injection device (*e.g.*, a syringe or an infusion pump). The injection can be delivered subcutaneously, intramuscularly, intraperitoneally, intravitreally, or intravenously, for example.

[00392] In another embodiment, the pharmaceutical composition is a solid formulation, *e.g.*, a freeze-dried or spray-dried composition, which can be used as is, or where to the physician or the patient adds solvents, and/or diluents prior to use. Solid dosage forms can include tablets, such as compressed tablets, and/or coated tablets, and capsules (*e.g.*, hard or soft gelatin capsules). The pharmaceutical composition can also be in the form of sachets, dragees, powders, granules, lozenges, or powders for reconstitution, for example.

[00393] The dosage forms can be immediate release, in which case they can comprise a water-soluble or dispersible carrier, or they can be delayed release, sustained release, or

modified release, in which case they can comprise water-insoluble polymers that regulate the rate of dissolution of the dosage form in the gastrointestinal tract or under the skin.

[00394] In other embodiments, the pharmaceutical composition can be delivered intranasally, intrabuccally, or sublingually.

[00395] The pH in an aqueous formulation can be between pH 3 and pH 10. In one embodiment, the pH of the formulation is from about 7.0 to about 9.5. In another embodiment, the pH of the formulation is from about 3.0 to about 7.0.

[00396] In another embodiment, the pharmaceutical composition comprises a buffer. Non-limiting examples of buffers include: arginine, aspartic acid, bicine, citrate, disodium hydrogen phosphate, fumaric acid, glycine, glycylglycine, histidine, lysine, maleic acid, malic acid, sodium acetate, sodium carbonate, sodium dihydrogen phosphate, sodium phosphate, succinate, tartaric acid, tricine, and tris(hydroxymethyl)-aminomethane, and mixtures thereof. The buffer can be present individually or in the aggregate, in a concentration from about 0.01 mgxml to about 50 mgxml, for example from about 0.1 mgxml to about 20 mgxml. Pharmaceutical compositions comprising each one of these specific buffers constitute alternative embodiments.

[00397] In another embodiment, the pharmaceutical composition comprises a preservative. Non-limiting examples of preservatives include: benzethonium chloride, benzoic acid, benzyl alcohol, bronopol, butyl 4-hydroxybenzoate, chlorobutanol, chlorocresol, chlorohexidine, chlorphenesin, o-cresol, m-cresol, p-cresol, ethyl 4-hydroxybenzoate, imidurea, methyl 4-hydroxybenzoate, phenol, 2-phenoxyethanol, 2-phenylethanol, propyl 4-hydroxybenzoate, sodium dehydroacetate, thiomerosal, and mixtures thereof. The preservative can be present individually or in the aggregate, in a concentration from about 0.01 mgxml to about 50 mgxml, for example from about 0.1 mgxml to about 20 mgxml. Pharmaceutical compositions comprising each one of these specific preservatives constitute alternative embodiments.

[00398] In another embodiment, the pharmaceutical composition comprises an isotonic agent. Non-limiting examples of isotonic agents include a salt (such as sodium chloride), an amino acid (such as glycine, histidine, arginine, lysine, isoleucine, aspartic acid, tryptophan, and threonine), an alditol (such as glycerol, 1,2-propanediol propyleneglycol), 1,3-propanediol, and 1,3-butanediol), polyethyleneglycol (*e.g.* PEG400), and mixtures thereof. Another example of an isotonic agent includes a sugar. Non-limiting examples of sugars can

include mono-, di-, or polysaccharides, or water-soluble glucans, including for example fructose, glucose, mannose, sorbose, xylose, maltose, lactose, sucrose, trehalose, dextran, pullulan, dextrin, cyclodextrin, alpha and beta- HPCD, soluble starch, hydroxyethyl starch, and sodium carboxymethyl-cellulose. Another example of an isotonic agent is a sugar alcohol, wherein the term “sugar alcohol” is defined as a C(4-8) hydrocarbon having at least one -OH group. Non-limiting examples of sugar alcohols include mannitol, sorbitol, inositol, galactitol, dulcitol, xylitol, and arabitol. The isotonic agent can be present individually or in the aggregate, in a concentration from about 0.01 mgxml to about 50 mgxml, for example from about 0.1 mgxml to about 20 mgxml. Pharmaceutical compositions comprising each one of these specific isotonic agents constitute alternative embodiments.

[00399] In another embodiment, the pharmaceutical composition comprises a chelating agent. Non-limiting examples of chelating agents include citric acid, aspartic acid, salts of ethylenediaminetetraacetic acid (EDTA), and mixtures thereof. The chelating agent can be present individually or in the aggregate, in a concentration from about 0.01 mgxml to about 50 mgxml, for example from about 0.1 mgxml to about 20 mgxml. Pharmaceutical compositions comprising each one of these specific chelating agents constitute alternative embodiments.

[00400] In another embodiment, the pharmaceutical composition comprises a stabilizer. Non-limiting examples of stabilizers include one or more aggregation inhibitors, one or more oxidation inhibitors, one or more surfactants, and/or one or more protease inhibitors.

[00401] In another embodiment, the pharmaceutical composition comprises a stabilizer, wherein said stabilizer is carboxy-xhydroxycellulose and derivatives thereof (such as HPC, HPC-SL, HPC-L and HPMC), cyclodextrins, 2-methylthioethanol, polyethylene glycol (such as PEG 3350), polyvinyl alcohol (PVA), polyvinyl pyrrolidone, salts (such as sodium chloride), sulphur-containing substances such as monothioglycerol), or thioglycolic acid. The stabilizer can be present individually or in the aggregate, in a concentration from about 0.01 mgxml to about 50 mgxml, for example from about 0.1 mgxml to about 20 mgxml. Pharmaceutical compositions comprising each one of these specific stabilizers constitute alternative embodiments.

[00402] In further embodiments, the pharmaceutical composition comprises one or more surfactants, such as a surfactant, at least one surfactant, or two different surfactants. The term “surfactant” refers to any molecules or ions that are comprised of a water-soluble

(hydrophilic) part, and a fat-soluble (lipophilic) part. The surfactant can, for example, be selected from the group consisting of anionic surfactants, cationic surfactants, nonionic surfactants, and/or zwitterionic surfactants. The surfactant can be present individually or in the aggregate, in a concentration from about 0.1 mg/ml to about 20 mg/ml. Pharmaceutical compositions comprising each one of these specific surfactants constitute alternative embodiments.

[00403] In a further embodiment, the pharmaceutical composition comprises one or more protease inhibitors, such as, *e.g.*, EDTA, and/or benzamidine hydrochloric acid (HCl). The protease inhibitor can be present individually or in the aggregate, in a concentration from about 0.1 mg/ml to about 20 mg/ml. Pharmaceutical compositions comprising each one of these specific protease inhibitors constitute alternative embodiments.

[00404] In another general aspect, provided herein is a method of producing a pharmaceutical composition comprising an antibody or antigen-binding fragment thereof provided herein, comprising combining an antibody or antigen-binding fragment thereof with a pharmaceutically acceptable carrier to obtain the pharmaceutical composition.

Methods of Administration

[00405] The mode of administration for therapeutic use of the antibodies provided herein can be any suitable route that delivers the antibody to the host, such as parenteral administration, *e.g.*, intradermal, intramuscular, intraperitoneal, intravenous or subcutaneous, pulmonary, transmucosal (oral, intranasal, intravaginal, rectal), using a formulation in a tablet, capsule, solution, powder, gel, particle; and contained in a syringe, an implanted device, osmotic pump, cartridge, micropump; or other means appreciated by the skilled artisan, as well known in the art. Site specific administration can be achieved by for example intratumoral, intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitary, intracelial, intracerebellar, intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intracardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravascular, intravesical, intralesional, vaginal, rectal, buccal, sublingual, intranasal, or transdermal delivery.

[00406] The antibodies provided herein can be administered to a subject by any suitable route, for example parentally by intravenous (i.v.) infusion or bolus injection,

intramuscularly or subcutaneously or intraperitoneally. i.v. infusion can be given over for example 15, 30, 60, 90, 120, 180, or 240 minutes, or from 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 hours.

[00407] The dose given to a subject is sufficient to alleviate or at least partially arrest the disease being treated ("therapeutically effective amount") and can be sometimes 0.005 mg to about 100 mg/kg, *e.g.* about 0.05 mg to about 30 mg/kg or about 5 mg to about 25 mg/kg, or about 4 mg/kg, about 8 mg/kg, about 16 mg/kg or about 24 mg/kg, or for example about 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 mg/kg, but can even higher, for example about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 40, 50, 60, 70, 80, 90 or 100 mg/kg.

[00408] A fixed unit dose can also be given, for example, 50, 100, 200, 500 or 1000 mg, or the dose can be based on the patient's surface area, *e.g.*, 500, 400, 300, 250, 200, or 100 mg/m². Usually between 1 and 8 doses, (*e.g.*, 1, 2, 3, 4, 5, 6, 7 or 8) can be administered to treat the patient, but 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more doses can be given.

[00409] The administration of the antibodies of provided herein, and in some embodiments of each and every one of the numbered embodiments listed below, can be repeated after one day, two days, three days, four days, five days, six days, one week, two weeks, three weeks, one month, five weeks, six weeks, seven weeks, two months, three months, four months, five months, six months or longer. Repeated courses of treatment are also possible, as is chronic administration. The repeated administration can be at the same dose or at a different dose. For example, the antibodies described herein can be administered at 8 mg/kg or at 16 mg/kg at weekly interval for 8 weeks, followed by administration at 8 mg/kg or at 16 mg/kg every two weeks for an additional 16 weeks, followed by administration at 8 mg/kg or at 16 mg/kg every four weeks by intravenous infusion.

[00410] For example, the antibodies in the methods described herein, and in some embodiments of each and every one of the numbered embodiments listed below, can be provided as a daily dosage in an amount of about 0.1-100 mg/kg, such as 0.5, 0.9, 1.0, 1.1, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, 45, 50, 60, 70, 80, 90 or 100 mg/kg, per day, on at least one of day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40, or alternatively, at least one of week 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 after initiation of treatment, or any

combination thereof, using single or divided doses of every 24, 12, 8, 6, 4, or 2 hours, or any combination thereof.

[00411] The antibodies in the methods described herein, and in some embodiments of each and every one of the numbered embodiments listed below, can also be administered prophylactically in order to reduce the risk of developing cancer, delay the onset of the occurrence of an event in cancer progression, and/or reduce the risk of recurrence when a cancer is in remission.

[00412] The antibodies provided herein can be lyophilized for storage and reconstituted in a suitable carrier prior to use. This technique has been shown to be effective with conventional protein preparations and well known lyophilization and reconstitution techniques can be employed.

Methods of use

[00413] The functional activity of antibodies provided herein can be characterized by methods known in the art and as described herein. Methods for characterizing antibodies and antigen-binding fragments thereof include, but are not limited to, affinity and specificity assays including Biacore, ELISA, and OCTETRED analysis; binding assays to detect the binding of antibodies to target cells by FACS; binding assays to detect the binding of antibodies to the target antigen on cells. According to particular embodiments, the methods for characterizing antibodies and antigen-binding fragments thereof include those described below. In certain embodiments, the antibody is a PSMA antibody. In some embodiments, the PSMA antibody is a PSMA multispecific antibody provided herein.

[00414] Also provided is a method of activating a T cell, comprising contacting the T cell with a multispecific PSMA antibody provided herein, wherein the second binding domain binds the T cell. In another general aspect, provided is a method of inactivating a T cell, comprising contacting the T cell with a multispecific PSMA antibody provided herein, wherein the second binding domain binds the T cell. In another general aspect, provided is a method of blocking activation of a T cell, comprising contacting the T cell with a multispecific PSMA antibody provided herein, wherein the second binding domain binds the T cell. In another general aspect, provided is a method of modulating the activation of a T cell, comprising contacting the T cell with a multispecific antibody provided herein, wherein the second binding domain binds the T cell. In some embodiments the T cell expresses CD3.

In some embodiments, the second binding domain binds CD3. In some embodiments, the multispecific PSMA antibody is a multispecific PSMA $\times$ CD3 antibody.

[00415] In another aspect, provided herein is a method of directing a T cell to a target cell, the method comprising contacting the T cell with a multispecific PSMA antibody provided herein. In another aspect, provided herein is a method of directing a T cell to a target cell, the method comprising contacting the T cell with a pharmaceutical composition comprising a multispecific PSMA antibody provided herein. In some embodiments, the contacting directs the T cell to the target cell. In some embodiments the target cell expresses PSMA. In some embodiments, the target cell is a prostate cell. In some embodiments, the target cell is a prostate cancer cell. In some embodiments, the target cell is a renal cell. In some embodiments, the target cell is a renal cancer cell. In some embodiments the T cell expresses CD3. In some embodiments, the multispecific PSMA antibody is a multispecific PSMA $\times$ CD3 antibody.

[00416] Also provided is a method of targeting an antigen on the surface of a target cell, the method comprising exposing the target cell to a PSMA antibody provided herein. Also provided is a method of targeting an antigen on the surface of a target cell, the method comprising exposing the target cell to a pharmaceutical composition comprising a PSMA antibody provided herein. In some embodiments the target cell expresses PSMA. In some embodiments, the target cell is a prostate cell. In some embodiments, the target cell is a prostate cancer cell. In some embodiments, the target cell is a renal cell. In some embodiments, the target cell is a renal cancer cell.

[00417] In another general aspect, provided is a method of targeting an antigen on the surface of a target cell, the method comprising exposing the target cell to a PSMA multispecific antibody provided herein. In another general aspect, provided is a method of targeting an antigen on the surface of a target cell, the method comprising exposing the target cell to a pharmaceutical composition comprising a PSMA multispecific antibody provided herein. In some embodiments the target cell expresses CD3. In some embodiments, the multispecific PSMA antibody is a multispecific PSMA $\times$ CD3 antibody.

[00418] PSMA is highly expressed in prostatic intraepithelial neoplasia (PIN), a condition in which some prostate cells have begun to look and behave abnormally, and in primary and metastatic prostate cancers (Bostwick *et al.* Cancer 1998;82 (11):2256–2261.); and the neovasculature of other solid tumors (*e.g.*, breast, lung, bladder, kidney). Expression of

PSMA in cancer tissues correlates with the stage of disease and Gleason score (Kawakami *et al.* Cancer Res 1997;57(12):2321–2324.). PSMA expression is also higher in prostate cancer cells from hormone-refractory patients (Wright *et al.* Urology 1996;48(2):326–334). Increased PSMA expression has been shown to be an independent marker of disease recurrence (Mitsiades *et al.* Clin Exp Metastasis 2004;21(6):495–505.). High-level PSMA expression is correlated with early prostate-specific antigen (PSA) recurrence in surgically treated prostate cancer. PSMA expression levels correlate with the aggressiveness of the disease. PSMA expression is increased in metastatic disease, hormone refractory cases, and higher-grade lesions, and it is further upregulated in androgen-insensitive tumors. Thus, this target is useful for prostate cancer characterization and subsequent therapy.

[00419] Blockade of PSMA can inhibit or decrease the growth of PSMA-expressing cancerous cells and tumors in a subject. It can also have antiangiogenic activity owing to expression of PSMA in tumor neovasculature (Milowsky, *et al.* 2007). PSMA is highly expressed in prostatic intraepithelial neoplasia, the most established precursor of prostatic carcinoma, and therefore blockade of PSMA can modulate progression of PIN to prostate cancer

[00420] In one aspect, provided is a method of inhibiting growth of tumor cells in a subject, comprising administering to the subject a therapeutically effective amount of a PSMA antibody provided herein. In some embodiments, the antibody is a multispecific PSMA antibody. In some embodiments, the multispecific PSMA antibody is a bispecific antibody. In certain embodiments, the multispecific PSMA antibody is a PSMAxCD3 antibody provided herein.

[00421] In one aspect, provided is a method of inhibiting the formation or growth of neovasculature of a tumor in a subject, comprising administering to the subject a therapeutically effective amount of a PSMA antibody provided herein. In some embodiments, the antibody is a multispecific PSMA antibody. In some embodiments, the multispecific PSMA antibody is a bispecific antibody. In certain embodiments, the multispecific PSMA antibody is a PSMAxCD3 antibody provided herein.

[00422] In one aspect, provided is a method of inhibiting progression of a precancerous state in a subject, comprising administering to the subject a therapeutically effective amount of a PSMA antibody provided herein. In some embodiments, the antibody is a multispecific PSMA antibody. In some embodiments, the multispecific PSMA antibody is a bispecific

antibody. In certain embodiments, the multispecific PSMA antibody is a PSMAxCD3 antibody provided herein.

[00423] In one aspect, provided is a method of treating a cancer by administering to the subject a PSMA antibody provided herein. In some embodiments, the antibody is a multispecific PSMA antibody. In some embodiments, the multispecific PSMA antibody is a bispecific antibody. In certain embodiments, the multispecific PSMA antibody is a PSMAxCD3 antibody provided herein.

[00424] Exemplary antibodies that can be used in the methods provided herein are antibodies specifically binding PSMA and bispecific PSMAxCD3 antibodies as described herein. Exemplary PSMA antibodies can be monospecific. Other exemplary PSMA antibodies can be part of a CD3 bispecific antibody, as provided herein. In certain embodiments, the PSMA antibody is PSMB889. In other embodiments, the PSMA antibody is PSMB890. In certain embodiments, the PSMA antibody is PSMB891. In certain embodiments, the PSMA antibody is PSMB892. In other embodiments, the PSMA antibody is PSMB893. In certain embodiments, the PSMA antibody is PSMB894. In other embodiments, the PSMA antibody is PSMB895. In certain embodiments, the PSMA antibody is PSMB896. In certain embodiments, the PSMA antibody is PSMB897. In other embodiments, the PSMA antibody is PSMB898. In certain embodiments, the PSMA antibody is PSMB899. In certain embodiments, the PSMA antibody is PSMHB49SC1133_011A11_1. In other embodiments, the PSMA antibody is PSMB896-G100A. In certain embodiments, the PSMA antibody is PSMA_P72_A10-HC-G54E. In certain embodiments, the PSMA antibody is PSMA_P72_D01-HC-D95E. In other embodiments, the PSMA antibody is PSMA_P72_F01. In other embodiments, the PSMA antibody is PSMA_P75_F01. In certain embodiments, the PSMA antibody is PSMA_P72_F07. In certain embodiments, the PSMA antibody is PSMA_P72_E07. In other embodiments, the PSMA antibody is PSMA_P72_D01. In certain embodiments, the PSMA antibody is PSMA_P72_C01. In other embodiments, the PSMA antibody is PSMA_P72_A10. In certain embodiments, the PSMA antibody is PSMA_P72_F02. In certain embodiments, the PSMA antibody is PSMA_P70_F02. In other embodiments, the PSMA antibody is PSMA_P72_G02. In other embodiments, the PSMA antibody is PSMA_P72_A11. Each of these antibodies is further described and characterized in the Examples below, including Tables 4-12 or 23-28. In some embodiments, the antibody is a PSMA antibody comprising a VH of any of the PSMA antibodies provided in Tables 4-12 or

23-28. In some embodiments, the antibody is a PSMA antibody comprising a VL of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In some embodiments, the antibody is a PSMA antibody comprising a VH and VL of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In other embodiments, the antibody is a PSMA antibody comprising a VH CDR1, VH CDR3 and VH CDR3 of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In other embodiments, the antibody is a PSMA antibody comprising a VL CDR1, VL CDR3 and VL CDR3 of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In other embodiments, the antibody is a PSMA antibody comprising a VH CDR1, VH CDR3, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 of any of the PSMA antibodies provided in Tables 4-12 or 23-28. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Kabat numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Chothia numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the AbM numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the Contact numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to the IMGT numbering system. In some embodiments, the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences of the PSMA antibody are according to a combination of the numbering systems provided herein.

[00425] Also provided herein is a method of treating a disease or disorder in a subject, comprising administering to the subject a PSMA antibody provided herein. Also provided herein is a method of treating a disease or disorder in a subject, comprising administering to the subject a PSMA antigen binding fragment provided herein. Also provided herein is a method of treating a disease or disorder in a subject, comprising administering to the subject a pharmaceutical composition comprising a PSMA antibody provided herein. Also provided herein is a method of treating a disease or disorder in a subject, comprising administering to the subject a pharmaceutical composition comprising a PSMA antigen binding fragment provided herein.

[00426] In some embodiments, the disease or disorder is a disease or disorder of the prostate. In some embodiments, the disease or disorder is a prostate-related disease or disorder. In some embodiments, the disease or disorder is prostate cancer. In some embodiments, the prostate cancer is a primary prostate cancer. In some embodiments, the prostate cancer is a metastatic prostate cancer. In another embodiment, the prostate cancer is a castration-resistant cancer. In one embodiment, the disease or disorder of the prostate is a prostate intraepithelial neoplasia. In some embodiments, the disease or disorder of the prostate is a prostate tumor. In some embodiments, the prostate tumor is a solid tumor.

[00427] In certain embodiments, the disease or disorder is a colorectal cancer. In certain embodiments, the disease or disorder is a clear cell renal carcinoma. In other embodiments, In certain embodiments, the disease or disorder is a renal cell carcinoma (RCC). In some embodiments, the RCC is a kidney clear cell carcinoma. In some embodiments, the RCC is a kidney papillary cell carcinoma. In certain embodiments, the disease or disorder is a metastatic lesion of a RCC. In certain embodiments, the disease or disorder is a bladder cancer. In certain embodiments, the disease or disorder is a breast cancer. In certain embodiments, the disease or disorder is a kidney cancer. In certain embodiments, the disease or disorder is a neovascular disorder. In certain embodiments, the disease or disorder is a cancer characterized by solid tumor growth. In some embodiments, the neovascular disorder is a clear cell renal carcinoma. In certain embodiments, the disease or disorder is a CCRCC colorectal cancer. In certain embodiments, the disease or disorder is a breast cancer. In certain embodiments, the disease or disorder is a bladder cancer. In certain embodiments, the disease or disorder is a lung cancer. In certain embodiments, the disease or disorder is a pancreatic cancer. In certain embodiments, the disease or disorder is a non-prostate cancers. In certain embodiments, the disease or disorder is a renal cancer. In certain embodiments, the disease or disorder is a urothelial cancer. In certain embodiments, the disease or disorder is a lung cancer. In certain embodiments, the disease or disorder is a colon cancer. In certain embodiments, the disease or disorder is a breast cancer. In certain embodiments, the disease or disorder is a liver adenocarcinoma.

[00428] The cancer can be a hyperproliferative condition or disorder, a solid tumor, a neovasculature, a soft tissue tumor, or a metastatic lesion. "Cancer" is meant to include all types of cancerous growths or oncogenic processes, metastatic tissues or malignantly transformed cells, tissues, or organs, irrespective of histopathology type or stage of invasiveness. Examples of cancers include solid tumors, hematological malignancies, soft

tissue tumors, and metastatic lesions. Exemplary solid tumors include malignancies, *e.g.*, sarcomas, and carcinomas (including adenocarcinomas and squamous cell carcinomas) of the various organ systems, such as those affecting prostate, liver, lung, breast, lymphoid, gastrointestinal (*e.g.*, colon), genitourinary tract (*e.g.*, renal, urothelial cells), prostate and pharynx. Adenocarcinomas include malignancies such as most colon cancers, a rectal cancer, a renal-cell carcinoma, a liver cancer, a non-small cell carcinoma of the lung, a cancer of the small intestine and a cancer of the esophagus. Squamous cell carcinomas include malignancies, *e.g.*, in the lung, esophagus, skin, head and neck region, oral cavity, anus, and cervix.

[00429] In one specific embodiment, the cancer is a prostate cancer.

[00430] Metastatic lesions of the aforementioned cancers can also be treated or prevented using the methods and antibodies described herein.

[00431] Exemplary cancers whose growth can be inhibited or reduced using the antibodies described herein include cancers that can overexpress PSMA. Exemplary such cancers include a prostate cancer or a prostatic intraepithelial neoplasia, a colorectal cancer, a gastric cancer, a clear cell renal carcinoma, a bladder cancer, a lung cancer, a squamous cell carcinoma, a glioma, a breast cancer, a kidney cancer, a neovascular disorder, a clear cell renal carcinoma (CCRCC), and a pancreatic cancer and various other non-prostate cancers, including but not limited to a renal cancer, a urothelial cancer and an adenocarcinoma to the liver. Refractory or recurrent malignancies can be treated using the antibodies described herein.

[00432] Exemplary other cancers that can be treated with the antibodies described herein are anal cancer, a basal cell carcinoma, a biliary tract cancer, a bladder cancer, a bone cancer, brain and CNS cancers, a carcinoma of the fallopian tubes, carcinoma of the vagina, a carcinoma of the vulva, a cutaneous or intraocular malignant melanoma, a astro-esophageal cancer, a testicular cancer, an ovarian cancer, a pancreatic cancer, a rectal cancer, an uterine cancer, a primary CNS lymphoma; a neoplasm of the central nervous system (CNS), a cervical cancer, a choriocarcinoma, a rectum cancer, a connective tissue cancer, a cancer of the digestive system, an endometrial cancer, an eye cancer; an intra-epithelial neoplasm, a kidney cancer, a larynx cancer, a liver cancer; a small cell lung cancer, a neuroblastoma, an oral cavity cancer (*e.g.*, lip, tongue, mouth, and pharynx), a nasopharyngeal cancer, a retinoblastoma, a rhabdomyosarcoma, a cancer of the respiratory

system, a sarcoma, a thyroid cancer, a cancer of the urinary system, a hepatocarcinoma, a cancer of the anal region, a carcinoma of the fallopian tubes, a carcinoma of the vagina, a carcinoma of the vulva, a cancer of the small intestine, a cancer of the endocrine system, a cancer of the parathyroid gland, a cancer of the adrenal gland, a sarcoma of soft tissue, a cancer of the urethra, a cancer of the penis, solid tumors of childhood, a tumor angiogenesis, a spinal axis tumor, a brain stem glioma, a pituitary adenoma, Kaposi's sarcoma, Merkel cell cancer, an epidermoid cancer, a squamous cell cancer, an environmentally induced cancers including those induced by asbestos, as well as other carcinomas and sarcomas, and combinations of said cancers.

[00433] In one embodiment, the cancer is a renal cancer. Renal cancer is one of the 10 most common cancers, affecting about 1 in every 63 people over a lifetime, mostly adults aged between 50 and 80 years. Worldwide, North America has the highest rate of renal cancer, but in developing countries, the incidence has been steadily increasing over the last three decades. Metastatic renal cell carcinoma (mRCC) is a disease that portends poor prognosis despite an increasing number of novel systemic treatment options including new targeted therapies and immunotherapy. PSMA is a transmembrane glycoprotein comprised of 750 amino acids and 3 protein domains; a small intracellular domain, a single-pass transmembrane domain, and a large extracellular domain. PSMA has been reported to be expressed within the neovasculature of other solid tumors including lung, bladder, and renal cancer (Chang SS, *et al.* Cancer Res. 1999;59(13):3192-3198). In a recent study examining PSMA expression in renal cell carcinomas (RCC), immunohistochemistry results revealed that endothelial PSMA protein was detected in 80% of clear cell renal carcinomas, 14% of papillary carcinomas, and 72% of chromophobe carcinomas (Spatz S, Tolkach *et al.* J Urol. 2018;199(2):370-377). Further analysis from the same study demonstrated that in both clear cell and papillary renal carcinomas, PSMA expression was significantly associated with lower overall survival rates in patients. In another clinical study, a PSMA-based radiotracer using ⁶⁸Ga was able to detect PSMA in metastatic lesions found in patients with clear cell carcinoma (Sawicki LM, *et al.* Eur J Nucl Med Mol Imaging. 2017;44(1):102-107). Thus, in addition to prostate cancer, PSMAxCD3 approaches can also have therapeutic benefit in patients with histologies such as clear cell renal cell carcinoma.

[00434] PSMA is highly expressed in prostate cancer and has also been reported to be expressed within the neovasculature of other solid tumors including lung, bladder, and renal cancer. In a recent study examining PSMA expression in renal cell carcinomas (RCC),

immunohistochemistry results revealed that endothelial PSMA protein was detected in 80% of clear cell renal carcinomas, 14% of papillary carcinomas, and 72% of chromophobe carcinomas. Further analysis from the same study demonstrated that in both clear cell and papillary renal carcinomas, PSMA expression was significantly associated with lower overall survival rates in patients. In another clinical study, a PSMA-based radiotracer using ^{68}Ga was able to detect PSMA in metastatic lesions found in patients with clear cell carcinoma. Thus, PSMA antibodies provided herein can have therapeutic benefit in patients with histologies, such as clear cell renal cell carcinoma.

[00435] In one specific embodiment, the cancer is a renal cancer. In another specific embodiment, the cancer is a renal cell carcinoma. In yet another specific embodiment, the cancer is a metastatic renal cell carcinoma. In other specific embodiments, the cancer is a clear cell renal cell carcinoma. In other specific embodiments, the cancer comprises PSMA-expressing cells.

[00436] Patients having cancer including metastatic cancer that express PSMA can be treated with the antibodies described herein. The cancer can be a prostate cancer or a prostatic intraepithelial neoplasia, a colorectal cancer, a gastric cancer, a clear cell renal carcinoma, a bladder cancer, a lung cancer, a squamous cell carcinoma, a glioma, a breast cancer, a kidney cancer, a neovascular disorder, a clear cell renal carcinoma (CCRCC), and a pancreatic cancer and various other non-prostate cancers, including but not limited to a renal cancer, a urothelial cancer and an adenocarcinoma to the liver.

[00437] In some embodiments, the subject has a solid tumor. In some embodiments, the subject has a prostate tumor. In some embodiments, the solid tumor is a colorectal cancer. In some embodiments, the solid tumor is a gastric cancer. In some embodiments, the solid tumor is a lung cancer. In some embodiments, the solid tumor is a bladder cancer. In some embodiments, the solid tumor is a squamous cell carcinoma. In some embodiments, the cancer is CCRCC. In some embodiments, the solid tumor is a breast cancer. In some embodiments, the solid tumor is glioma. In some embodiments, the solid tumor is a prostate cancer or castration-resistant prostate cancer. In some embodiments, the solid tumor is a kidney cancer. In some embodiments, the solid tumor is a pancreatic cancer. In some embodiments, the solid tumor is an adenocarcinoma to the liver. In some embodiments, the cancer is a neovascular. In some embodiments, the cancer is renal cancer. In some embodiments, the solid tumor is a urothelial cancer. In some embodiments, the solid tumor is a brain cancer.

[00438] In some embodiments, the subject has a tumor that expresses PSMA. In some embodiments, the subject has tumor-infiltrating T lymphocytes (TILs) in the tumor tissue.

[00439] “Increased number” refers to statistically significant increase in a subject when compared to a control. “Increased number” for example refers to statistically significant increase in the number of TILs in a subject (*e.g.* patient) pre- and post-treatment with a PSMA antibody or other therapeutic.

[00440] An exemplary PSMA positive cancer is a prostate cancer. Another exemplary PSMA positive cancer is a renal cancer. In some embodiments, prostate cancer is adenocarcinoma. In some embodiments, prostate cancer is a metastatic prostate cancer. In some embodiments, prostate cancer has metastasized to rectum, lymph node or bone, or any combination thereof. In some embodiments, prostate cancer is relapsed or refractory prostate cancer. In some embodiments, prostate cancer is castration resistant prostate cancer. In some embodiments, prostate cancer is sensitive to androgen deprivation therapy. In some embodiments, prostate cancer is insensitive to androgen deprivation therapy.

[00441] In some embodiments described herein, the subject has increased expression or activity of interferon-gamma (IFN- γ).

[00442] In some embodiments described herein the subject has been treated with an anti-PSMA antibody. In some embodiments described herein, the subject is refractory to treatment with the anti-PSMA antibody.

[00443] Any of the PSMA or bispecific PSMAxCD3 antibodies described herein can be used in the methods provided herein.

[00444] Also provided is a method of directing CD3-expressing T cells to a second target. The methods can comprise contacting the CD3-expressing T cell with a multispecific PSMAxCD3 antibody or antigen binding fragment thereof provided herein, wherein the multispecific PSMAxCD3 antibody or antigen binding fragment thereof directs the CD3-expressing T cell to the second target. Also provided is a method of directing a T cell expressing CD3 to a second target, the method comprising contacting the T cell with a multispecific PSMAxCD3 antibody provided herein, wherein the contacting directs the T cell to the second target. In certain embodiments, the antibody is a multispecific PSMAxCD3 antibody provided herein. In some embodiments, the second target is PSMA. In some embodiments, the second target is a cell expressing PSMA. In some embodiments, the second target is a prostate cell. In some embodiments, the second target is a prostate cell

expressing PSMA. In some embodiments, the second target is a prostate cancer cell. In some embodiments, the second target is a prostate cancer cell expressing PSMA. In some embodiments, the second target is a renal cell. In some embodiments, the second target is a renal cancer cell. In certain embodiments, the second target is a renal cell expressing PSMA.

[00445] Also provided is a method for inhibiting growth or proliferation of target cells. The methods can comprise contacting the CD3-expressing T cells with a multispecific PSMAxCD3 antibody or antigen binding fragment thereof provided herein, wherein contacting the target cells with the multispecific PSMAxCD3 antibody or antigen binding fragment thereof composition inhibits the growth or proliferation of the target cells. In some embodiments, the target cell is a cell expressing PSMA. In some embodiments, the target cell is a prostate cell. In some embodiments, the target cell is a prostate cell expressing PSMA. In some embodiments, the target cell is a prostate cancer cell. In some embodiments, the target cell is a prostate cancer cell expressing PSMA. In some embodiments, the target cell is a renal cell. In some embodiments, the target cell is a renal cancer cell. In certain embodiments, the target cell is a renal cell expressing PSMA.

[00446] Also provided is a method of inhibiting growth or proliferation of target cells expressing a second target antigen on the cell surface, the method comprising contacting the target cells with a multispecific PSMA antibody provided herein, wherein contacting the target cells with the pharmaceutical composition inhibits growth or proliferation of the target cells. In some embodiments, the target cells are in the presence of a T cell expressing CD3 while in contact with the multispecific antibody. In certain embodiments, the antibody is a multispecific PSMAxCD3 antibody provided herein. In some embodiments, the target cell is a cell expressing PSMA. In some embodiments, the target cell is a prostate cell. In some embodiments, the target cell is a prostate cell expressing PSMA. In some embodiments, the target cell is a prostate cancer cell. In some embodiments, the target cell is a prostate cancer cell expressing PSMA. In some embodiments, the target cell is a renal cell. In some embodiments, the target cell is a renal cancer cell. In certain embodiments, the target cell is a renal cell expressing PSMA.

[00447] Also provided herein is a method of treating a disease or disorder in a subject, comprising administering to the subject a PSMA multispecific antibody provided herein. Also provided herein is a method of treating a disease or disorder in a subject, comprising administering to the subject a multispecific antibody comprising a PSMA antigen binding fragment provided herein. Also provided herein is a method of treating a disease or disorder

in a subject, comprising administering to the subject a pharmaceutical composition comprising a PSMA multispecific antibody provided herein. Also provided herein is a method of treating a disease or disorder in a subject, comprising administering to the subject a pharmaceutical composition comprising a multispecific antibody comprising a PSMA antigen binding fragment provided herein. In certain embodiments, the antibody is a multispecific PSMAxCD3 antibody provided herein. In some embodiments, the disease or disorder is a disease or disorder of the prostate. In some embodiments, the disease or disorder is a prostate-related disease or disorder. In some embodiments, the disease or disorder is prostate cancer. In some embodiments, the disease or disorder is a renal disease or disorder. In some embodiments, the disease or disorder is a renal cancer. In some embodiments, the disease or disorder is a renal cell carcinoma. In some embodiments, the renal cell carcinoma is a metastatic renal cell carcinoma.

[00448] In certain embodiments of the methods provided herein, the disease or disorder is mediated by a PSMA-expressing cell.

[00449] In another general aspect, provided herein is a method of treating a disease or disorder in a subject, comprising administering to the subject an isolated multispecific antibody or antigen binding fragment thereof that specifically binds PSMA and a second target antigen presented on the surface of a T cell, or a pharmaceutical composition disclosed herein. In some embodiments, the second target antigen is CD3. In certain embodiments, the antibody is a multispecific PSMAxCD3 antibody provided herein. In some embodiments, the disease or disorder is a disease or disorder of the prostate. In some embodiments, the disease or disorder is a prostate-related disease or disorder. In some embodiments, the disease or disorder is prostate cancer. In some embodiments, provided is a method for eliminating target cells expressing PSMA or treating a disease caused all or in part by target cells expressing the PSMA in a subject, comprising administering an effective amount of a multispecific PSMA antibody provided herein to the subject. In certain embodiments, the antibody is a multispecific PSMAxCD3 antibody.

[00450] In some embodiments, the subject is a subject in need thereof. In some embodiments, the subject is a human. In specific embodiments, the subject is administered an effective amount. In some embodiments, the subject has a prostate cancer. In some embodiments, the subject has a primary prostate cancer. In one embodiment, the subject has a metastatic prostate cancer. In certain embodiments, the subject is hormone-refractory. PSMA can be used as a biomarker for disease, disease progression, or disease reoccurrence

in a subject. In some embodiments, the subject has detectable levels of PSMA in the blood. In some embodiments, the subject has detectable levels of PSMA in the bone marrow. Methods of detection of PSMA are known in the art. In some embodiments, the PSMA is detected using a PSMA antibody provided herein. In some embodiments, the subject has detectable levels of prostate-specific antigen (PSA) in the blood. In other embodiments, the subject has detectable levels of PSA in the bone marrow. Methods of detection of PSA are known in the art.

[00451] As used herein, the term “effective amount” refers to an amount of an active ingredient or component that elicits the desired biological or medicinal response in a subject.

[00452] According to particular embodiments, an effective amount refers to the amount of therapy which is sufficient to achieve one, two, three, four, or more of the following effects: (i) reduce or ameliorate the severity of the disease, disorder or condition to be treated or a symptom associated therewith; (ii) reduce the duration of the disease, disorder or condition to be treated, or a symptom associated therewith; (iii) prevent the progression of the disease, disorder or condition to be treated, or a symptom associated therewith; (iv) cause regression of the disease, disorder or condition to be treated, or a symptom associated therewith; (v) prevent the development or onset of the disease, disorder or condition to be treated, or a symptom associated therewith; (vi) prevent the recurrence of the disease, disorder or condition to be treated, or a symptom associated therewith; (vii) reduce hospitalization of a subject having the disease, disorder or condition to be treated, or a symptom associated therewith; (viii) reduce hospitalization length of a subject having the disease, disorder or condition to be treated, or a symptom associated therewith; (ix) increase the survival of a subject with the disease, disorder or condition to be treated, or a symptom associated therewith; (xi) inhibit or reduce the disease, disorder or condition to be treated, or a symptom associated therewith in a subject; and/or (xii) enhance or improve the prophylactic or therapeutic effect(s) of another therapy.

[00453] The effective amount or dosage can vary according to various factors, such as the disease, disorder or condition to be treated, the means of administration, the target site, the physiological state of the subject (including, *e.g.*, age, body weight, health), whether the subject is a human or an animal, other medications administered, and whether the treatment is prophylactic or therapeutic. Treatment dosages are optimally titrated to optimize safety and efficacy.

[00454] According to particular embodiments, the compositions described herein are formulated to be suitable for the intended route of administration to a subject. For example, the compositions described herein can be formulated to be suitable for intravenous, subcutaneous, or intramuscular administration.

[00455] As used herein, the terms “treat,” “treating,” and “treatment” are all intended to refer to an amelioration or reversal of at least one measurable physical parameter related to a disease or disorder (*e.g.*, a cancer), which is not necessarily discernible in the subject, but can be discernible in the subject. The terms “treat,” “treating,” and “treatment,” can also refer to causing regression, preventing the progression, or at least slowing down the progression of the disease, disorder, or condition. In a particular embodiment, “treat,” “treating,” and “treatment” refer to an alleviation, prevention of the development or onset, or reduction in the duration of one or more symptoms associated with the disease, disorder, or condition, such as a tumor or a cancer. In a particular embodiment, “treat,” “treating,” and “treatment” refer to prevention of the recurrence of the disease, disorder, or condition. In a particular embodiment, “treat,” “treating,” and “treatment” refer to an increase in the survival of a subject having the disease, disorder, or condition. In a particular embodiment, “treat,” “treating,” and “treatment” refer to elimination of the disease, disorder, or condition in the subject.

[00456] In some embodiments, a PSMA antibody provided herein is used in combination with a supplemental therapy. In some embodiments, a multispecific PSMAxCD3 antibody provided herein is used in combination with a supplemental therapy. In some embodiments, the supplemental therapy is surgery. In some embodiments, the supplemental therapy is radiation. In some embodiments, the supplemental therapy is hormone therapy. In certain embodiments, the hormone therapy is abiraterone. In certain embodiments, the hormone therapy is enzalutamide. In some embodiments, the supplemental therapy is chemotherapy. In certain embodiments, the chemotherapy is docetaxel. In certain embodiments, the chemotherapy is cabazitaxel. In some embodiments, the supplemental therapy is a cancer vaccine. In some embodiments, the supplemental therapy is a radiopharmaceutical agent (*e.g.*, radium-223 chloride).

[00457] As used herein, the term “in combination,” in the context of the administration of two or more therapies to a subject, refers to the use of more than one therapy. The use of the term “in combination” does not restrict the order in which therapies are administered to a subject. “In combination with” can include two or more therapeutics are administered to a

subject together in a mixture, concurrently as single agents or sequentially as single agents in any order. For example, a first therapy (*e.g.*, a composition described herein) can be administered prior to (*e.g.*, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 16 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (*e.g.*, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 16 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second therapy to a subject.

[00458] The antibodies described herein can also be administered in combination therapy, *i.e.*, combined with other therapeutic agents relevant for the disease or condition to be treated. Accordingly, in one embodiment, the antibody-containing medicament is for combination with one or more further therapeutic agent, such as a chemotherapeutic agent. In some embodiments, the other therapeutic agent is a radiopharmaceutical agent (such as radium-223 chloride), secondary hormone therapies (such as abiraterone or enzalutamide), and/or chemotherapies (docetaxel and cabazitaxel). Such combined administration can be simultaneous, separate or sequential, in any order. For simultaneous administration the agents can be administered as one composition or as separate compositions, as appropriate. a chemotherapeutic agent (*e.g.*, docetaxel, carboplatin, fludarabine), abiraterone, hormonal therapy (*e.g.*, flutamide, bicalutamide, nilutamide, cyproterone acetate, ketoconazole, aminoglutethimide, abarelix, degarelix, leuprolide, goserelin, triptorelin, buserelin, ARN-509), serine or tyrosine kinase inhibitor (*e.g.*, PI3 kinase inhibitor SF1126,) (*e.g.*, lapatanib), multikinase inhibitor (*e.g.*, sorafenib, sunitinib), VEGF inhibitor (*e.g.*, bevacizumab), TAK-700, cancer vaccine (*e.g.*, BPX-101, PEP223), Listeria-based vaccine, lenalidomide, TOK-001, IGF-1 receptor inhibitor (*e.g.*, cixutumumab), TRC105, Aurora A kinase inhibitor (*e.g.*, MLN8237), proteasome inhibitor (*e.g.*, bortezomib), OGX-011, radioimmunotherapy (*e.g.*, HuJ591-GS), HDAC inhibitor (*e.g.*, valproic acid, SB939, LBH589), hydroxychloroquine, mTOR inhibitor (*e.g.*, everolimus), dovitinib lactate, diindolylmethane, efavirenz, OGX-427, genistein, IMC-3G3, bafetinib, CP-675,206, radiation therapy, surgery, or a combination thereof.

[00459] In one embodiment, a method for treating a disorder involving cells expressing PSMA in a subject, which method comprises administration of a therapeutically effective amount of a bispecific antibody or fragment, such as a PSMA x CD3 bispecific antibody

described herein, and radiotherapy to a subject in need thereof is provided. In one embodiment is provided a method for treating or preventing cancer, which method comprises administration of a therapeutically effective amount of a bispecific antibody or fragment, such as a PSMA x CD3 antibody described herein, and radiotherapy to a subject in need thereof. Radiotherapy can comprise radiation or associated administration of radiopharmaceuticals to a patient is provided. The source of radiation can be either external or internal to the patient being treated (radiation treatment can, for example, be in the form of external beam radiation therapy (EBRT) or brachytherapy (BT)). Radioactive elements that can be used in practicing such methods include, *e.g.*, radium, cesium-137, iridium-192, americium-241, gold-198, cobalt-57, copper-67, technetium-99, iodide-123, iodide-131, and indium-111.

[00460] Antibodies described herein can be administered in combination with a vaccine. Exemplary vaccines are immunogenic agents, such as cancerous cells, purified tumor antigens (including recombinant proteins, antigen epitopes, peptides and carbohydrate molecules), tumor antigens delivered to a patient via gene therapy, cells, and cells transfected with genes encoding immune stimulating cytokines. Exemplary vaccines that can be used include peptides of melanoma antigens, such as peptides of gp100, MAGE antigens, Trp-2, MART1 and/or tyrosinase, or tumor cells transfected to express the cytokine GM-CSF, DNA-based vaccines, RNA-based vaccines, Listeria-based vaccines and viral transduction-based vaccines, peptides or prostate antigens (*e.g.*, PSMA, STEAP1, PSCA), the cancer vaccine sipuleucel-T or peptides of lung cancer antigens. The cancer vaccine can be prophylactic or therapeutic.

[00461] Many experimental strategies for vaccination against tumors have been devised (see Rosenberg, S., 2000, Development of Cancer Vaccines, ASCO Educational Book Spring: 60-62; Logothetis, C., 2000, ASCO Educational Book Spring: 300-302; Khayat, D. 2000, ASCO Educational Book Spring: 414-428; Foon, K. 2000, ASCO Educational Book Spring: 730-738; see also Restifo, N. and Sznol, M., Cancer Vaccines, Ch. 61, pp. 3023-3043 in DeVita, V. *et al.* (eds.), 1997, Cancer: Principles and Practice of Oncology. Fifth Edition). In one of these strategies, a vaccine is prepared using autologous or allogeneic tumor cells. These cellular vaccines have been shown to be most effective when the tumor cells are transduced to express GM-CSF. GM-CSF has been shown to be a potent activator of antigen presentation for tumor vaccination (Dranoff *et al.*, (1993) Proc Natl Acad Sci U.S.A. 90: 3539-43).

[00462] The antibodies described herein can be administered in combination with one or a collection of recombinant proteins and/or peptides expressed in or on a tumor in order to generate an immune response to these proteins. These proteins are normally viewed by the immune system as self-antigens and are therefore tolerant to them. The tumor antigen can also include the protein telomerase, which is required for the synthesis of telomeres of chromosomes and which is expressed in more than 85% of human cancers and in only a limited number of somatic tissues (Kim *et al.*, (1994) Science 266: 2011-2013). Tumor antigens can also be “neo-antigens” expressed in or on cancer cells as a result of somatic mutations that alter protein sequence or create fusion proteins between two unrelated sequences (*e.g.*, bcr-abl in the Philadelphia chromosome), or idiotype from B cell tumors. The tumor antigens can be antigen epitopes of prostate specific antigen (PSA), mesothelin, prostate-specific membrane antigen (PSMA), synovial sarcoma X2 (SSX2), NKX3.1, prostatic acidic phosphatase (PAP), or epidermal growth factor receptors, or peptides specific for variants of EGFR such as the well-known EGFRvIII overexpressed on tumor cells.

[00463] Other tumor vaccines can include the proteins from viruses implicated in human cancers such as Human Papilloma Viruses (HPV), Hepatitis Viruses (HBV and HCV) and Kaposi's Herpes Sarcoma Virus (KHSV), and Epstein-Barr virus (EBV). Another form of tumor specific antigens which can be used in combination with the antibodies described herein is purified heat shock proteins (HSP) isolated from the tumor tissue itself. HSP contain fragments of proteins from the tumor cells and are highly efficient at delivery to antigen presenting cells for eliciting tumor immunity (Suot and Srivastava (1995) Science 269:1585-1588; Tamura *et al.*, (1997) Science 278:117-120).

[00464] Dendritic cells (DC) are potent antigen presenting cells that can be used to prime antigen-specific responses. DC's can be produced *ex vivo* and loaded with various protein and peptide antigens as well as tumor cell extracts (Nestle *et al.*, (1998) Nature Medicine 4: 328-332). DCs can also be transduced by genetic means to express these tumor antigens. DCs have also been fused directly to tumor cells for the purposes of immunization (Kugler *et al.*, (2000) Nature Medicine 6:332-336). As a method of vaccination, DC immunization can be effectively combined with the antibodies described herein to activate more potent anti-tumor responses.

[00465] In some embodiments described herein, the antibodies specifically binding PSMA provided herein or the bispecific PSMAxCD3 antibodies provided herein are administered in

combination with a tumor vaccine comprising a peptide fragment of a prostate specific antigen, or a vector encoding the peptide fragment of a prostate specific antigen.

[00466] The antibodies described herein can be administered in combination with a standard of care cancer treatment.

[00467] The antibodies described herein can be administered in combination with a standard of care cancer chemotherapeutic regimes. In these instances, it can be possible to reduce the dose of chemotherapeutic reagent administered (Mokyr *et al.*, (1998) Cancer Research 58: 5301-5304).

[00468] In some embodiments described herein, the antibodies provided herein can be administered in combination with one or more of other antibody molecules, chemotherapy, other anti-cancer therapy (*e.g.*, targeted anti-cancer therapies, or oncolytic drugs), cytotoxic agents, cytokines, surgical and/or radiation procedures.

[00469] Exemplary cytotoxic agents that can be administered in combination with the antibodies described herein include hormone inhibitors, antimicrotubule agents, topoisomerase inhibitors, anti-metabolites, mitotic inhibitors, alkylating agents, anthracyclines, vinca alkaloids, intercalating agents, agents capable of interfering with a signal transduction pathway, agents that promote apoptosis, proteasome inhibitors, and radiation (*e.g.*, local or whole body irradiation).

[00470] Standard of care therapeutics include anastrozole (Arimidex®), bicalutamide (Casodex®), bleomycin sulfate (Blenoxane®), busulfan (Myleran®), busulfan injection (Busulfex®), capecitabine (Xeloda®), N4-pentoxycarbonyl-5-deoxy-5-fluorocytidine, carboplatin (Paraplatin®), carmustine (BiCNU®), chlorambucil (Leukeran®), cisplatin (Platinol®), cladribine (Leustatin®), cyclophosphamide (Cytosan® or Neosar®), cytarabine, cytosine arabinoside (Cytosar-U®), cytarabine liposome injection (DepoCyt®), dacarbazine (DTIC-Dome®), dactinomycin (Actinomycin D, Cosmegan), daunorubicin hydrochloride (Cerubidine®), daunorubicin citrate liposome injection (DaunoXome®), dexamethasone, docetaxel (Taxotere®), doxorubicin hydrochloride (Adriamycin®, Rubex®), etoposide (Vepesid®), fludarabine phosphate (Fludara®), 5-fluorouracil (Adrucil®, Efudex®), flutamide (Eulexin®), tezacitibine, Gemcitabine (difluorodeoxycytidine), hydroxyurea (Hydrea®), Idarubicin (Idamycin®), ifosfamide (IFEX®), irinotecan (Camptosar®), L-asparaginase (ELSPAR®), leucovorin calcium, melphalan (Alkeran®), 6-mercaptopurine (Purinethol®), methotrexate (Folex®),

mitoxantrone (Novantrone®), mylotarg, paclitaxel (Taxol®), phoenix (Yttrium90/MX-DTPA), pentostatin, polifeprosan 20 with carmustine implant (Gliadel®), tamoxifen citrate (Nolvadex®), teniposide (Vumon®), 6-thioguanine, thiotepa, tirapazamine (Tirazone®), topotecan hydrochloride for injection (Hycamptin®), vinblastine (Velban®), vincristine (Oncovin®), vinorelbine (Navelbine®), Ibrutinib, idelalisib, and brentuximab vedotin.

[00471] Exemplary alkylating agents include, nitrogen mustards, ethylenimine derivatives, alkyl sulfonates, nitrosoureas and triazenes: uracil mustard (Aminouracil Mustard®, Chlorethaminacil®, Demethyldopan®, Desmethyldopan®, Haemanthamine®, Nordopan®, Uracil Nitrogen Mustard®, Uracillost®, Uracilmostaza®, Uramustin®, Uramustine®), chlormethine (Mustargen®), cyclophosphamide (Cytosan®, Neosar®, Clafen®, Endoxan®, Procytox®, Revimmune™), ifosfamide (Mitoxana®), melphalan (Alkeran®), Chlorambucil (Leukeran®), pipobroman (Amedel®, Vercyte®), triethylenemelamine (Hemel®, Hexalen®, Hexastat®), triethylenethiophosphoramine, Temozolomide (Temodar®), thiotepa (Thioplex®), busulfan (Busilvex®, Myleran®), carmustine (BiCNU®), lomustine (CeeNU®), streptozocin (Zanosar®), and Dacarbazine (DTIC-Dome®). Additional exemplary alkylating agents include, Oxaliplatin (Eloxatin®); Temozolomide (Temodar® and Temodal®); Dactinomycin (also known as actinomycin-D, Cosmegen®); Melphalan (also known as L-PAM, L-sarcosylsin, and phenylalanine mustard, Alkeran®); Altretamine (also known as hexamethylmelamine (HMM), Hexalen®); Carmustine (BiCNU®); Bendamustine (Treanda®); Busulfan (Busulfex® and Myleran®); Carboplatin (Paraplatin®); Lomustine (also known as CCNU, CeeNU®); Cisplatin (also known as CDDP, Platinol® and Platinol®-AQ); Chlorambucil (Leukeran®); Cyclophosphamide (Cytosan® and Neosar®); Dacarbazine (also known as DTIC, DIC and imidazole carboxamide, DTIC-Dome®); Altretamine (also known as hexamethylmelamine (HMM), Hexalen®); Ifosfamide (Ifex®); Prednimustine; Procarbazine (Matulane®); Mechlorethamine (also known as nitrogen mustard, mustine and mechloroethamine hydrochloride, Mustargen®); Streptozocin (Zanosar®); Thiotepa (also known as thiophosphoamide, TESP and TSPA, Thioplex®); Cyclophosphamide (Endoxan®, Cytosan®, Neosar®, Procytox®, Revimmune®); and Bendamustine HCl (Treanda®).

[00472] Exemplary anthracyclines include, e.g., doxorubicin (Adriamycin® and Rubex®); bleomycin (Lenoxane®); daunorubicin (daunorubicin hydrochloride, daunomycin, and rubidomycin hydrochloride, Cerubidine®); daunorubicin liposomal (daunorubicin citrate liposome, DaunoXome®); mitoxantrone (DHAD, Novantrone®); epirubicin (Ellence™);

idarubicin (Idamycin®, Idamycin PFS®); mitomycin C (Mutamycin®); geldanamycin; herbimycin; ravidomycin; and desacetylavidomycin.

[00473] Exemplary vinca alkaloids that can be used in combination with the antibodies provided herein include vinorelbine tartrate (Navelbine®), Vincristine (Oncovin®), and Vindesine (Eldisine®); vinblastine (also known as vinblastine sulfate, vincalurekoblamine and VLB, Alkaban-AQ® and Velban®); and vinorelbine (Navelbine®).

[00474] Exemplary proteasome inhibitors that can be used in combination with the antibodies provided herein, alone or in combination with another immunomodulator are bortezomib (Velcade®); carfilzomib (PX-171-007, (S)-4-Methyl-N—((S)-1-(((S)-4-methyl-1-((R)-2-methyloxiran-2-yl)-1-oxopentan-2-yl)amino)-1-oxo-3-phenylpropan-2-yl)-2-((S)-2-(2-morpholinoacetamido)-4-phenylbutanamido)-pentanamide); marizomib (NPI-0052); ixazomib citrate (MLN-9708); delanzomib (CEP-18770); and O-Methyl-N-[(2-methyl-5-thiazolyl)carbonyl]-L-seryl-O-methyl-N-[(1S)-2-[(2R)-2-methyl-2-oxiranyl]-2-oxo-1-(phenylmethyl)ethyl]-L-serinamide (ONX-0912).

[00475] In some embodiments described herein, the antibodies provided herein are administered in combination with a serine or tyrosine kinase inhibitor (*e.g.*, a receptor tyrosine kinase (RTK) inhibitor). Exemplary tyrosine kinase inhibitor include an epidermal growth factor (EGF) pathway inhibitor (*e.g.*, an epidermal growth factor receptor (EGFR) inhibitor), a vascular endothelial growth factor (VEGF) pathway inhibitor (*e.g.*, a vascular endothelial growth factor receptor (VEGFR) inhibitor (*e.g.*, a VEGFR-1 inhibitor, a VEGFR-2 inhibitor, a VEGFR-3 inhibitor), a platelet derived growth factor (PDGF) pathway inhibitor (*e.g.*, a platelet derived growth factor receptor (PDGFR) inhibitor (*e.g.*, a PDGFR-β inhibitor), a RAF-1 inhibitor, a KIT inhibitor and a RET inhibitor. In some embodiments, the second therapeutic is axitinib (AG013736), bosutinib (SKI-606), cediranib (RECENTIN™, AZD2171), dasatinib (SPRYCEL®, BMS-354825), erlotinib (TARCEVA®), gefitinib (IRESSA®), imatinib (Gleevec®, CGP57148B, STI-571), lapatinib (TYKERB®, TYVERB®), lestaurtinib (CEP-701), neratinib (HKI-272), nilotinib (TASIGNA®), semaxanib (semaxinib, SU5416), sunitinib (SUTENT®, SU11248), toceranib (PALLADIA®), vandetanib (ZACTIMA®, ZD6474), vatalanib (PTK787, PTK/ZK), trastuzumab (HERCEPTIN®), bevacizumab (AVASTIN®), rituximab (RITUXAN®), cetuximab (ERBITUX®), panitumumab (VECTIBIX®), ranibizumab (Lucentis®), nilotinib (TASIGNA®), sorafenib (NEXAVAR®), alemtuzumab (CAMPATH®), gemtuzumab ozogamicin (MYLOTARG®), ENMD-2076, PCI-32765,

AC220, dovitinib lactate (TKI258, CHIR-258), BIBW 2992 (TOVOK™), SGX523, PF-04217903, PF-02341066, PF-299804, BMS-777607, ABT-869, MP470, BIBF 1120 (VARGATEF®), AP24534, JNJ-26483327, MGCD265, DCC-2036, BMS-690154, CEP-11981, tivozanib (AV-951), OSI-930, MM-121, XL-184, XL-647, XL228, AEE788, AG-490, AST-6, BMS-599626, CUDC-101, PD153035, pelitinib (EKB-569), vandetanib (zactima), WZ3146, WZ4002, WZ8040, ABT-869 (linifanib), AEE788, AP24534 (ponatinib), AV-951 (tivozanib), axitinib, BAY 73-4506 (regorafenib), brivanib alaninate (BMS-582664), brivanib (BMS-540215), cediranib (AZD2171), CHIR-258 (dovitinib), CP 673451, CYC116, E7080, Ki8751, masitinib (AB1010), MGCD-265, motesanib diphosphate (AMG-706), MP-470, OSI-930, Pazopanib Hydrochloride, PD173074, Sorafenib Tosylate (Bay 43-9006), SU 5402, TSU-68 (SU6668), vatalanib, XL880 (GSK1363089, EXEL-2880). Selected tyrosine kinase inhibitors are chosen from sunitinib, erlotinib, gefitinib, or sorafenib. In some embodiments, the EGFR inhibitor is a bispecific EGFRc-Met antibody (EM-1 mAb) comprising the heavy and the light chains of SEQ ID NOs 249, 250, 217 and 218 (US2014/0141000).

[00476] In some embodiments, the antibodies provided herein are administered in combination with Vascular Endothelial Growth Factor (VEGF) receptor inhibitors, including Bevacizumab (Avastin®), axitinib (Inlyta®); Brivanib alaninate (BMS-582664, (S)—((R)-1-(4-(4-Fluoro-2-methyl-1H-indol-5-yloxy)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yloxy)propan-2-yl)2-aminopropanoate); Sorafenib (Nexavar®); Pazopanib (Votrient®); Sunitinib malate (Sutent®); Cediranib (AZD2171, CAS 288383-20-1); Vargatef (BIBF1120, CAS 928326-83-4); Foretinib (GSK1363089); Telatinib (BAY57-9352, CAS 332012-40-5); Apatinib (YN968D1, CAS 811803-05-1); Imatinib (Gleevec®); Ponatinib (AP24534, CAS 943319-70-8); Tivozanib (AV951, CAS 475108-18-0); Regorafenib (BAY73-4506, CAS 755037-03-7); Vatalanib dihydrochloride (PTK787, CAS 212141-51-0); Brivanib (BMS-540215, CAS 649735-46-6); Vandetanib (Caprelsa® or AZD6474); Motesanib diphosphate (AMG706, CAS 857876-30-3, N-(2,3-dihydro-3,3-dimethyl-1H-indol-6-yl)-2-[(4-pyridinylmethyl)amino]-3-pyridinecarboxamide, described in PCT Publication No. WO 02/066470); Dovitinib dilactic acid (TKI258, CAS 852433-84-2); Linfanib (ABT869, CAS 796967-16-3); Cabozantinib (XL184, CAS 849217-68-1); Lestaurtinib (CAS 111358-88-4); N-[5-[[[5-(1,1-Dimethylethyl)-2-oxazolyl]methyl]thio]-2-thiazolyl]-4-piperidinecarboxamide (BMS38703, CAS 345627-80-7); (3R,4R)-4-Amino-1-((4-((3-methoxyphenyl)amino)pyrrolo[2,1-f][1,2,4]triazin-5-yl)methyl)piperidin-3-ol

(BMS690514); N-(3,4-Dichloro-2-fluorophenyl)-6-methoxy-7-[[[(3 α ,5 β ,6 α)-octahydro-2-methylcyclopenta[c]pyrrol-5-yl]methoxy]-4-quinazolinamine (XL647, CAS 781613-23-8); 4-Methyl-3-[[[1-methyl-6-(3-pyridinyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]amino]-N-[3-(trifluoromethyl)phenyl]-benzamide (BHG712, CAS 940310-85-0); and Aflibercept (Eylea®).

[00477] Exemplary VEGF inhibitors include a monoclonal antibody that binds to the same epitope as the monoclonal anti-VEGF antibody A4.6.1 produced by hybridoma ATCC HB 10709; a recombinant humanized anti-VEGF monoclonal antibody generated according to Presta *et al.*, (1997) Cancer Res 57:4593-4599. In one embodiment, the anti-VEGF antibody is Bevacizumab (BV), also known as rhuMAb VEGF or AVASTIN®. It comprises mutated human IgG1 framework regions and antigen-binding complementarity-determining regions from the murine anti-hVEGF monoclonal antibody A.4.6.1 that blocks binding of human VEGF to its receptors. Bevacizumab and other humanized anti-VEGF antibodies are further described in U.S. Patent No. 6,884,879. Additional anti-VEGF antibodies include the G6 or B20 series antibodies (*e.g.*, G6-31, B20-4.1), as described in Int. Patent Publication Nos. WO2005/012359 and WO2005/044853. For additional antibodies see U.S. Patent Nos. 7,060,269, 6,582,959, 6,703,020, 6,054,297, WO98/45332, WO 96/30046, WO94/10202, EP 0666868B1, U.S. Patent Application Publication Nos. US2006009360, US20050186208, US20030206899, US20030190317, US20030203409, and US20050112126; and Popkov *et al.*, (2004) Journal of Immunological Methods 288: 149-164. Other antibodies include those that bind to a functional epitope on human VEGF comprising of residues F17, M18, D19, Y21, Y25, Q89, 191, K101, E103, and C104 or, alternatively, comprising residues F17, Y21, Q22, Y25, D63, 183 and Q89.

[00478] In some embodiments described herein, the antibodies provided herein are administered in combination with a PI3K inhibitor. In one embodiment, the PI3K inhibitor is an inhibitor of delta and gamma isoforms of PI3K. In another embodiment, the PI3K inhibitor is an inhibitor of beta isoforms of PI3K. Exemplary PI3K inhibitors that can be used are described in, *e.g.*, WO 2010/036380, WO 2010/006086, WO 09/114870, WO 05/113556, GSK 2126458, GDC-0980, GDC-0941, Sanofi XL147, XL756, XL147, PF-46915032, BKM 120, CAL-101, CAL 263, SF1126, PX-886, and a dual PI3K inhibitor (*e.g.*, Novartis BEZ235).

[00479] In some embodiments described herein, the antibodies provided herein are administered in combination with a mTOR inhibitor, *e.g.*, one or more mTOR inhibitors

chosen from one or more of rapamycin, temsirolimus (TORISEL®), AZD8055, BEZ235, BGT226, XL765, PF-4691502, GDC0980, SF1126, OSI-027, GSK1059615, KU-0063794, WYE-354, Palomid 529 (P529), PF-04691502, or PKI-587. ridaforolimus (formally known as deferolimus, (1R,2R,4S)-4-[(2R)-2-[(1R,9S,12S,15R,16E,18R,19R,21R,23S,24E,26E,28Z,30S,32S,35R)-1,18-dihydroxy-19,30-dimethoxy-15,17,21,23,29,35-hexamethyl-2,3,10,14,20-pentaoxo-11,36-dioxo-4-azatricyclo[30.3.1.0^{4,9}] hexatriaconta-16,24,26,28-tetraen-12-yl]propyl]-2-methoxycyclohexyl dimethylphosphinate, also known as AP23573 and MK8669, and described in PCT Publication No. WO 03/064383); everolimus (Afinitor® or RAD001); rapamycin (AY22989, Sirolimus®); simapimod (CAS 164301-51-3); emsirolimus, (5-{2,4-Bis[(3S)-3-methylmorpholin-4-yl]pyrido[2,3-d]pyrimidin-7-yl}-2-methoxyphenyl)methanol (AZD8055); 2-Amino-8-[trans-4-(2-hydroxyethoxy)cyclohexyl]-6-(6-methoxy-3-pyridinyl)-4-methyl-pyrido[2,3-d]pyrimidin-7(8H)-one (PF04691502, CAS 1013101-36-4); and N2-[1,4-dioxo-4-[[4-(4-oxo-8-phenyl-4H-1-benzopyran-2-yl)morpholinium-4-yl]methoxy]butyl]-L-arginylglycyl-L- α -aspartyl-L-serine- (SEQ ID NO:237), inner salt (SF1126, CAS 936487-67-1), and XL765.

[00480] In some embodiments described herein, the antibodies provided herein are administered in combination with a BRAF inhibitor, *e.g.*, GSK2118436, RG7204, PLX4032, GDC-0879, PLX4720, and sorafenib tosylate (Bay 43-9006).

[00481] In some embodiments described herein, the antibodies provided herein are administered in combination with an immunomodulatory agent. Targeting immune checkpoints such as programmed cell death protein 1 (PD1), programmed cell death 1 ligand 1 (PDL1) and cytotoxic T lymphocyte antigen 4 (CTLA4) has achieved noteworthy benefit in multiple cancers by blocking immunoinhibitory signals and enabling patients to produce an effective antitumour response. In some embodiments, the antibodies provided herein are administered in combination with an anti-PD1 (*eg.* nivolumab), anti-PDL (*eg.* MDX-1105) or anti-CTLA4 (*eg.* Ipilimumab). The ability of agonistic CD40 antibodies (referred to as α CD40) or CD40 ligand to stimulate immune responses and target tumors suggests such reagents have promise as cancer immunotherapeutics. In some embodiments, the antibodies provided herein are administered in combination with an anti-CD40 (*eg.* SGN-40, CP-870,893) or anti-CD40L (*eg.* BG9588).

[00482] In some embodiments described herein, the antibodies provided herein are administered in combination with a MEK inhibitor.

[00483] In some embodiments described herein, the antibodies provided herein administered in combination with the MEK inhibitor are used in the treatment of a prostate cancer, a melanoma, a colorectal cancer, a non-small cell lung cancer, an ovarian cancer, a breast cancer, a prostate cancer, a pancreatic cancer, a hematological malignancy or a renal cell carcinoma. In certain embodiments, the tumor tissue or cancer cell has a BRAF mutation (*e.g.*, a BRAF V600E mutation), a BRAF wildtype, a KRAS wildtype or an activating KRAS mutation. The cancer can be at an early, intermediate or late stage. Any MEK inhibitor can be used in combination including, ARRY-142886, G02442104 (also known as GSK1120212), RDEA436, RDEA119/BAY 869766, AS703026, G00039805 (also known as AZD-6244 or selumetinib), BIX 02188, BIX 02189, CI-1040 (PD-184352), PD0325901, PD98059, U0126, GDC-0973 (Methanone, [3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]phenyl][[3-hydroxy-3-(25)-2-piperidinyl-1-azetidiny]-), G-38963, G02443714 (also known as AS703206), or a pharmaceutically acceptable salt or solvate thereof. Additional examples of MEK inhibitors are disclosed in WO 2013/019906, WO 03/077914, WO 2005/121142, WO 2007/04415, WO 2008/024725 and WO 2009/085983.

[00484] In some embodiments described herein, the antibodies provided herein are administered in combination with a JAK2 inhibitor, *e.g.*, CEP-701, INCB18424, CP-690550 (tasocitinib).

[00485] In some embodiments described herein, the antibodies provided herein are administered in combination with paclitaxel or a paclitaxel agent, *e.g.*, TAXOL®, protein-bound paclitaxel (*e.g.*, ABRAXANE®). Exemplary paclitaxel agents include nanoparticle albumin-bound paclitaxel (ABRAXANE, marketed by Abraxis Bioscience), docosahexaenoic acid bound-paclitaxel (DHA-paclitaxel, Taxoprexin, marketed by Protarga), polyglutamate bound-paclitaxel (PG-paclitaxel, paclitaxel poliglumex, CT-2103, XYOTAX, marketed by Cell Therapeutic), the tumor-activated prodrug (TAP), ANG105 (Angiopep-2 bound to three molecules of paclitaxel, marketed by ImmunoGen), paclitaxel-EC-1 (paclitaxel bound to the erbB2-recognizing peptide EC-1; see Li *et al.*, Biopolymers (2007) 87:225-230), and glucose-conjugated paclitaxel (*e.g.*, 2'-paclitaxel methyl 2-glucopyranosyl succinate, see Liu *et al.*, (2007) Bioorganic & Medicinal Chemistry Letters 17:617-620).

[00486] In some embodiments described herein, the antibodies provided herein are administered in combination with a cellular immunotherapy (*e.g.*, Provenge (*e.g.*, Sipuleucel)), and optionally in combination with cyclophosphamide.

[00487] Exemplary therapeutic agents that can be used in combination with the antibodies provided herein for treatment of a cancer include a chemotherapeutic agent (*e.g.*, docetaxel, carboplatin, fludarabine), abiraterone, hormonal therapy (*e.g.*, flutamide, bicalutamide, nilutamide, cyproterone acetate, ketoconazole, aminoglutethimide, abarelix, degarelix, leuprolide, goserelin, triptorelin, buserelin), tyrosine kinase inhibitor (*e.g.*, dual kinase inhibitor (*e.g.*, lapatanib), multikinase inhibitor (*e.g.*, sorafenib, sunitinib), VEGF inhibitor (*e.g.*, bevacizumab), TAK-700, cancer vaccine (*e.g.*, BPX-101, PEP223), lenalidomide, TOK-001, IGF-1 receptor inhibitor (*e.g.*, cixutumumab), TRC105, Aurora A kinase inhibitor (*e.g.*, MLN8237), proteasome inhibitor (*e.g.*, bortezomib), OGX-011, radioimmunotherapy (*e.g.*, Hu1591-GS), HDAC inhibitor (*e.g.*, valproic acid, SB939, LBH589), hydroxychloroquine, mTOR inhibitor (*e.g.*, everolimus), dovitinib lactate, diindolylmethane, efavirenz, OGX-427, genistein, IMC-3G3, bafetinib, CP-675,206, ARN-509, radiation therapy, surgery, or a combination thereof.

[00488] Exemplary therapeutic agents that can be used in combination with the antibodies provided herein for treatment of a cancer include a chemotherapeutic agent, *e.g.*, paclitaxel or a paclitaxel agent (*e.g.*, a paclitaxel formulation such as TAXOL, an albumin-stabilized nanoparticle paclitaxel formulation (*e.g.*, ABRAXANE) or a liposomal paclitaxel formulation); gemcitabine (*e.g.*, gemcitabine alone or in combination with AXP107-11); other chemotherapeutic agents such as oxaliplatin, 5-fluorouracil, capecitabine, rubitecan, epirubicin hydrochloride, NC-6004, cisplatin, docetaxel (*e.g.*, TAXOTERE), mitomycin C, ifosfamide; interferon; tyrosine kinase inhibitor (*e.g.*, EGFR inhibitor (*e.g.*, erlotinib, panitumumab, cetuximab, nimotuzumab); HER2/neu receptor inhibitor (*e.g.*, trastuzumab); dual kinase inhibitor (*e.g.*, bosutinib, saracatinib, lapatinib, vandetanib); multikinase inhibitor (*e.g.*, sorafenib, sunitinib, XL184, pazopanib); VEGF inhibitor (*e.g.*, bevacizumab, AV-951, brivanib); radioimmunotherapy (*e.g.*, XR303); cancer vaccine (*e.g.*, GVAX, survivin peptide); COX-2 inhibitor (*e.g.*, celecoxib); IGF-1 receptor inhibitor (*e.g.*, AMG 479, MK-0646); mTOR inhibitor (*e.g.*, everolimus, temsirolimus); IL-6 inhibitor (*e.g.*, CNTO 328); cyclin-dependent kinase inhibitor (*e.g.*, P276-00, UCN-01); Altered Energy Metabolism-Directed (AEMD) compound (*e.g.*, CPI-613); HDAC inhibitor (*e.g.*, vorinostat); TRAIL receptor 2 (TR-2) agonist (*e.g.*, conatumumab); MEK inhibitor (*e.g.*, AS703026, selumetinib, GSK1120212); Raf/MEK dual kinase inhibitor (*e.g.*, RO5126766); Notch signaling inhibitor (*e.g.*, MK0752); monoclonal antibody-antibody fusion protein (*e.g.*, L19IL2); curcumin; HSP90 inhibitor (*e.g.*, tanespimycin, STA-9090); rIL-2;

denileukin diftitox; topoisomerase I inhibitor (*e.g.*, irinotecan, PEP02); statin (*e.g.*, simvastatin); Factor VIIa inhibitor (*e.g.*, PCI-27483); AKT inhibitor (*e.g.*, RX-0201); hypoxia-activated prodrug (*e.g.*, TH-302); metformin hydrochloride, gamma-secretase inhibitor (*e.g.*, R04929097); ribonucleotide reductase inhibitor (*e.g.*, 3-AP); immunotoxin (*e.g.*, HuC242-DM4); PARP inhibitor (*e.g.*, KU-0059436, veliparib); CTLA-4 inhibitor (*e.g.*, CP-675,206, ipilimumab); AdV-tk therapy; proteasome inhibitor (*e.g.*, bortezomib (Velcade), NPI-0052); thiazolidinedione (*e.g.*, pioglitazone); NPC-1C; Aurora kinase inhibitor (*e.g.*, R763/AS703569), CTGF inhibitor (*e.g.*, FG-3019); siG12D LODER; and radiation therapy (*e.g.*, tomotherapy, stereotactic radiation, proton therapy), surgery, and a combination thereof. In certain embodiments, a combination of paclitaxel or a paclitaxel agent, and gemcitabine can be used with the antibodies provided herein.

[00489] Exemplary therapeutic agents that can be used in combination with the antibodies provided herein for treatment of small cell lung cancer include a chemotherapeutic agent, *e.g.*, etoposide, carboplatin, cisplatin, oxaliplatin, irinotecan, topotecan, gemcitabine, liposomal SN-38, bendamustine, temozolomide, belotecan, NK012, FR901228, flavopiridol); tyrosine kinase inhibitor (*e.g.*, EGFR inhibitor (*e.g.*, erlotinib, gefitinib, cetuximab, panitumumab); multikinase inhibitor (*e.g.*, sorafenib, sunitinib); VEGF inhibitor (*e.g.*, bevacizumab, vandetanib); cancer vaccine (*e.g.*, GVAX); Bcl-2 inhibitor (*e.g.*, oblimersen sodium, ABT-263); proteasome inhibitor (*e.g.*, bortezomib (Velcade), NPI-0052), paclitaxel or a paclitaxel agent; docetaxel; IGF-1 receptor inhibitor (*e.g.*, AMG 479); HGF/SF inhibitor (*e.g.*, AMG 102, MK-0646); chloroquine; Aurora kinase inhibitor (*e.g.*, MLN8237); radioimmunotherapy (*e.g.*, TF2); HSP90 inhibitor (*e.g.*, tanespimycin, STA-9090); mTOR inhibitor (*e.g.*, everolimus); Ep-CAM/CD3-bispecific antibody (*e.g.*, MT110); CK-2 inhibitor (*e.g.*, CX-4945); HDAC inhibitor (*e.g.*, belinostat); SMO antagonist (*e.g.*, BMS 833923); peptide cancer vaccine, and radiation therapy (*e.g.*, intensity-modulated radiation therapy (IMRT), hypofractionated radiotherapy, hypoxia-guided radiotherapy), surgery, and combinations thereof.

[00490] Exemplary therapeutic agents that can be used in combination with the antibodies provided herein for treatment of non-small cell lung cancer include a chemotherapeutic agent, *e.g.*, vinorelbine, cisplatin, docetaxel, pemetrexed disodium, etoposide, gemcitabine, carboplatin, liposomal SN-38, TLK286, temozolomide, topotecan, pemetrexed disodium, azacitidine, irinotecan, tegafur-gimeracil-oteracil potassium, sapacitabine); tyrosine kinase inhibitor (*e.g.*, EGFR inhibitor (*e.g.*, erlotinib, gefitinib, cetuximab, panitumumab,

necitumumab, PF-00299804, nimotuzumab, RO5083945), MET inhibitor (*e.g.*, PF-02341066, ARQ 197), PI3K kinase inhibitor (*e.g.*, XL147, GDC-0941), Raf/MEK dual kinase inhibitor (*e.g.*, RO5126766), PI3K/mTOR dual kinase inhibitor (*e.g.*, XL765), SRC inhibitor (*e.g.*, dasatinib), dual inhibitor (*e.g.*, BIBW 2992, GSK1363089, ZD6474, AZD0530, AG-013736, lapatinib, MEHD7945A, linifanib), multikinase inhibitor (*e.g.*, sorafenib, sunitinib, pazopanib, AMG 706, XL184, MGCD265, BMS-690514, R935788), VEGF inhibitor (*e.g.*, endostar, endostatin, bevacizumab, cediranib, BIBF 1120, axitinib, tivozanib, AZD2171), cancer vaccine (*e.g.*, BLP25 liposome vaccine, GVAX, recombinant DNA and adenovirus expressing L523S protein), Bel-2 inhibitor (*e.g.*, oblimersen sodium), proteasome inhibitor (*e.g.*, bortezomib, carfilzomib, NPI-0052, MLN9708), paclitaxel or a paclitaxel agent, docetaxel, IGF-1 receptor inhibitor (*e.g.*, cixutumumab, MK-0646, OSI 906, CP-751,871, BIIB022), hydroxychloroquine, HSP90 inhibitor (*e.g.*, tanespimycin, STA-9090, AU922, XL888), mTOR inhibitor (*e.g.*, everolimus, temsirolimus, ridaforolimus), Ep-CAM/CD3-bispecific antibody (*e.g.*, MT110), CK-2 inhibitor (*e.g.*, CX-4945), HDAC inhibitor (*e.g.*, MS 275, LBH589, vorinostat, valproic acid, FR901228), DHFR inhibitor (*e.g.*, pralatrexate), retinoid (*e.g.*, bexarotene, tretinoin), antibody-drug conjugate (*e.g.*, SGN-15), bisphosphonate (*e.g.*, zoledronic acid), cancer vaccine (*e.g.*, belagenpumatucel-L), low molecular weight heparin (LMWH) (*e.g.*, tinzaparin, enoxaparin), GSK1572932A, melatonin, talactoferrin, dimesna, topoisomerase inhibitor (*e.g.*, amrubicin, etoposide, karenitecin), nelfinavir, cilengitide, ErbB3 inhibitor (*e.g.*, MM-121, U3-1287), survivin inhibitor (*e.g.*, YM155, LY2181308), eribulin mesylate, COX-2 inhibitor (*e.g.*, celecoxib), pegfilgrastim, Polo-like kinase 1 inhibitor (*e.g.*, BI 6727), TRAIL receptor 2 (TR-2) agonist (*e.g.*, CS-1008), CNGRC peptide (SEQ ID NO:225)-TNF alpha conjugate, dichloroacetate (DCA), HGF inhibitor (*e.g.*, SCH 900105), SAR240550, PPAR-gamma agonist (*e.g.*, CS-7017), gamma-secretase inhibitor (*e.g.*, RO4929097), epigenetic therapy (*e.g.*, 5-azacitidine), nitroglycerin, MEK inhibitor (*e.g.*, AZD6244), cyclin-dependent kinase inhibitor (*e.g.*, UCN-01), cholesterol-Fus1, antitubulin agent (*e.g.*, E7389), farnesyl-OH-transferase inhibitor (*e.g.*, lonafamib), immunotoxin (*e.g.*, BB-10901, SS1 (dsFv) PE38), fondaparinux, vascular-disrupting agent (*e.g.*, AVE8062), PD-L1 inhibitor (*e.g.*, MDX-1105, MDX-1106), beta-glucan, NGR-hTNF, EMD 521873, MEK inhibitor (*e.g.*, GSK1120212), epothilone analog (*e.g.*, ixabepilone), kinesin-spindle inhibitor (*e.g.*, 4SC-205), telomere targeting agent (*e.g.*, KML-001), P70 pathway inhibitor (*e.g.*, LY2584702), AKT inhibitor (*e.g.*, MK-2206), angiogenesis inhibitor (*e.g.*, lenalidomide), Notch signaling

inhibitor (*e.g.*, OMP-21M18), EGFR/c-Met bispecific antibody EM-1 as described in US2014/0141000A1, radiation therapy, surgery, and combinations thereof.

[00491] Exemplary therapeutic agents that can be used in combination with the antibodies provided herein for treatment of ovarian cancer include a chemotherapeutic agent (*e.g.*, paclitaxel or a paclitaxel agent; docetaxel; carboplatin; gemcitabine; doxorubicin; topotecan; cisplatin; irinotecan, TLK286, ifosfamide, olaparib, oxaliplatin, melphalan, pemetrexed disodium, SJG-136, cyclophosphamide, etoposide, decitabine); ghrelin antagonist (*e.g.*, AEZS-130), immunotherapy (*e.g.*, APC8024, oregovomab, OPT-821), tyrosine kinase inhibitor (*e.g.*, EGFR inhibitor (*e.g.*, erlotinib), dual inhibitor (*e.g.*, E7080), multikinase inhibitor (*e.g.*, AZD0530, JI-101, sorafenib, sunitinib, pazopanib), ON 01910.Na), VEGF inhibitor (*e.g.*, bevacizumab, BIBF 1120, cediranib, AZD2171), PDGFR inhibitor (*e.g.*, IMC-3G3), paclitaxel, topoisomerase inhibitor (*e.g.*, karenitecin, Irinotecan), HDAC inhibitor (*e.g.*, valproate, vorinostat), folate receptor inhibitor (*e.g.*, farletuzumab), angiopoietin inhibitor (*e.g.*, AMG 386), epothilone analog (*e.g.*, ixabepilone), proteasome inhibitor (*e.g.*, carfilzomib), IGF-1 receptor inhibitor (*e.g.*, OSI 906, AMG 479), PARP inhibitor (*e.g.*, veliparib, AG014699, iniparib, MK-4827), Aurora kinase inhibitor (*e.g.*, MLN8237, ENMD-2076), angiogenesis inhibitor (*e.g.*, lenalidomide), DHFR inhibitor (*e.g.*, pralatrexate), radioimmunotherapeutic agent (*e.g.*, Hu3S193), statin (*e.g.*, lovastatin), topoisomerase 1 inhibitor (*e.g.*, NKTR-102), cancer vaccine (*e.g.*, p53 synthetic long peptides vaccine, autologous OC-DC vaccine), mTOR inhibitor (*e.g.*, temsirolimus, everolimus), BCR/ABL inhibitor (*e.g.*, imatinib), ET-A receptor antagonist (*e.g.*, ZD4054), TRAIL receptor 2 (TR-2) agonist (*e.g.*, CS-1008), HGF/SF inhibitor (*e.g.*, AMG 102), EGEN-001, Polo-like kinase 1 inhibitor (*e.g.*, BI 6727), gamma-secretase inhibitor (*e.g.*, RO4929097), Wee-1 inhibitor (*e.g.*, MK-1775), antitubulin agent (*e.g.*, vinorelbine, E7389), immunotoxin (*e.g.*, denileukin diftitox), SB-485232, vascular-disrupting agent (*e.g.*, AVE8062), integrin inhibitor (*e.g.*, EMD 525797), kinesin-spindle inhibitor (*e.g.*, 4SC-205), revlimid, HER2 inhibitor (*e.g.*, MGAH22), ErrB3 inhibitor (*e.g.*, MM-121), radiation therapy, and combinations thereof.

[00492] Exemplary therapeutics agents that can be used in combination with the antibodies provided herein for treatment of a renal cancer, *e.g.*, a renal cell carcinoma (RCC) or metastatic RCC include an immune-based strategy (*e.g.*, interleukin-2 or interferon- α), a targeted agent (*e.g.*, a VEGF inhibitor such as a monoclonal antibody to VEGF, *e.g.*, bevacizumab (Rini *et al.*, (2010) J Clin Oncol 28(13):2137-2143)); a VEGF tyrosine kinase

inhibitor such as sunitinib, sorafenib, axitinib and pazopanib (reviewed in Pal *et al.*, (2014) Clin Advances in Hematology & Oncology 12(2):90-99); an RNAi inhibitor, or an inhibitor of a downstream mediator of VEGF signaling, *e.g.*, an inhibitor of the mammalian target of rapamycin (mTOR), *e.g.*, everolimus and temsirolimus (Hudes *et al.*, (2007) N Engl J Med 356(22):2271-2281, Motzer *et al.*, (2008) Lancet 372: 449-456).

[00493] An “immunoconjugate” refers to the antibody provided herein conjugated to one or more heterologous molecule(s). In some embodiments, a PSMA antibody provided herein is an immunoconjugate. In some embodiments, a multispecific PSMA antibody provided herein is an immunoconjugate. In certain embodiments, the multispecific PSMA antibody is a multispecific PSMAxCD3 antibody provided herein.

[00494] In some embodiments, the antibody is conjugated to one or more cytotoxic agents. Exemplary such cytotoxic agents include chemotherapeutic agents or drugs, growth inhibitory agents, toxins (*e.g.*, protein toxins, enzymatically active toxins of bacterial, fungal, plant, or animal origin, or fragments thereof), and radioactive isotopes.

[00495] In some embodiments, an immunoconjugate is an antibody-drug conjugate (ADC) in which the antibody provided herein is conjugated to one or more drugs. In other embodiments, a PSMA antibody provided herein is an ADC. In other embodiments, a multispecific PSMA antibody provided herein is an ADC. In certain embodiments, the multispecific PSMA antibody is a multispecific PSMAxCD3 antibody provided herein. Exemplary drugs include a maytansinoid (see, *e.g.*, U.S. Patent No. 5,208,020, 5,416,066); an auristatin such as monomethylauristatin drug moieties DE and DF (MMAE and MMAF) (see, *e.g.*, U.S. Patent Nos. 5,635,483 and 5,780,588, and 7,498,298), a dolastatin, a calicheamicin or derivative thereof (see, *e.g.*, U.S. Patent Nos. 5,712,374, 5,714,586, 5,739, 116, 5,767,285, 5,770,701, 5,770,710, 5,773,001, and 5,877,296; Hinman *et al.*, (1993) Cancer Res 53:3336-3342; and Lode *et al.*, (1998) Cancer Res 58:2925-2928); an anthracycline such as daunomycin or doxorubicin (see, *e.g.*, Kratz *et al.*, (2006) Current Med. Chem 13:477-523; Jeffrey *et al.*, (2006) Bioorganic & Med Chem Letters 16:358-362; Torgov *et al.*, (2005) Bioconj Chem 16:717-721; Nagy *et al.*, (2000) Proc Natl Acad Sci USA 97:829-834; Dubowchik *et al.*, Bioorg. & Med. Chem. Letters 12: 1529-1532 (2002); King *et al.*, (2002) J Med Chem 45:4336-4343; and U.S. Patent No. 6,630,579), methotrexate, vindesine, a taxane such as docetaxel, paclitaxel, larotaxel, tesetaxel, and ortataxel.

[00496] In some embodiments, the immunoconjugate comprises the antibody described herein conjugated to an enzymatically active toxin or fragment thereof, such as diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolacca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes.

[00497] In some embodiments described herein, the antibody is conjugated to a radioactive atom to form a radioconjugate. A variety of radioactive isotopes are available for the production of radioconjugates. Examples include At211, 113I, 1125, Y90, Re186, Re188, Sm153, Bi212, P32, Pb212 and radioactive isotopes of Lu. When the radioconjugate is used for detection, it can comprise a radioactive atom for scintigraphic studies, for example tc99m or 1123, or a spin label for nuclear magnetic resonance (NMR) imaging (also known as magnetic resonance imaging, mri), such as iodine- 123 again, iodine- 131, indium-111, fluorine-19, carbon-13, nitrogen-15, oxygen-17, gadolinium, manganese or iron.

[00498] Conjugates of the antibodies described herein and the cytotoxic agent can be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP), succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HQ), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al.*, (1987) Science 238: 1098. Carbon- 14-labeled l-isothiocyanatobenzyl-3-methyldiethylene triaminopentaacetic acid (MX- DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See, *e.g.*, W094/11026. The linker can be a "cleavable linker" facilitating release of a cytotoxic drug in the cell. For example, an acid-labile linker, peptidase-sensitive linker, photolabile linker, dimethyl linker or disulfide-containing linker (Chari *et al.*, (1992) Cancer Res 52: 127-131; U.S. Patent No. 5,208,020) can be used.

[00499] The immunoconjugates or ADCs can be prepared with cross-linker reagents such as BMPS, EMCS, GMBS, HBVS, LC-SMCC, MBS, MPBH, SBAP, SIA, SIAB, SMCC,

SMPB, SMPH, sulfo-EMCS, sulfo-GMBS, sulfo-KMUS, sulfo-MBS, sulfo-SIAB, sulfo-SMCC, and sulfo-SMPB, and SVSB (succinimidyl-(4-vinylsulfone)benzoate) which are commercially available (*e.g.*, from Pierce Biotechnology, Inc., Rockford, IL., U.S.A).

[00500] In one embodiment, provided is an immunoconjugate comprising a PSMA antibody provided herein linked to a therapeutic agent or an imaging agent. In another embodiment, provided is an immunoconjugate comprising a CD3 antibody provided herein to a therapeutic agent or an imaging agent. In one embodiment, provided is an immunoconjugate comprising a bispecific PSMA/CD3 antibody provided herein linked to a therapeutic agent or an imaging agent.

[00501] PSMA antibodies provided herein can also be used as agents to detect PSMA-expressing cells. Thus, in another methods, provided is a method of detecting a cell expressing PSMA, comprising contacting a cell with a PSMA antibody provided herein. In some embodiments, the antibody is a multispecific PSMA antibody. In some embodiments, the PSMA antibody is a multispecific PSMA_xCD3 antibody. In certain embodiments, the detecting is by ELISA. In some embodiments, the detecting is by FACS analysis. Also provided are kits comprising a PSMA antibody provided herein, and instructions for use.

Enrichment and detection methods

[00502] In one aspect, the PSMA antibodies provided herein are used as agents to detect PSMA-expressing cells. Thus, in other methods, provided is a method of detecting a cell expressing PSMA, comprising contacting a cell with a PSMA antibody provided herein. In certain embodiments, the detecting is by ELISA. In some embodiments, the detecting is by FACS analysis.

[00503] In some embodiments, PSMA antibodies provided herein are used in a method to diagnose PSMA-expressing cancer progression. In some embodiments, PSMA antibodies provided herein are used in a method to diagnose PSMA-expressing cancer progression. In some embodiments, PSMA antibodies provided herein are used in a method to diagnose PSMA-expressing cancer regression. In some embodiments, PSMA antibodies provided herein are used in a method to diagnose PSMA-expressing cancer stability.

[00504] In some embodiments, PSMA antibodies provided herein are used in a method to monitor PSMA-expressing cancer progression. In some embodiments, PSMA antibodies provided herein are used in a method to monitor PSMA-expressing cancer progression. In some embodiments, PSMA antibodies provided herein are used in a method to monitor

PSMA-expressing cancer regression. In some embodiments, PSMA antibodies provided herein are used in a method to monitor PSMA-expressing cancer stability.

[00505] In other embodiments, PSMA antibodies provided herein are used in a method to determine whether a patient should be treated for cancer. In some embodiments, PSMA antibodies provided herein are used in a method to determine whether a subject is afflicted with PSMA-expressing cancer. In some embodiments, PSMA antibodies provided herein are used in a method to determine whether a subject is amenable to treatment with a PSMA-specific anti-cancer therapeutic. In certain embodiments, the PSMA-specific anti-cancer therapeutic is a PSMA antibody provided herein. In other embodiments, the PSMA-specific anti-cancer therapeutic is a multispecific PSMA antibody provided herein. In specific embodiments, the PSMA-specific anti-cancer therapeutic is a multispecific PSMAxCD3 antibody provided herein.

[00506] In one aspect, provided is a method of detecting PSMA in a sample, comprising obtaining the sample, contacting the sample with a PSMA antibody provided herein, and detecting the antibody bound to PSMA in the sample.

[00507] In another aspect, provided is a method of detecting PSMA and CD3 in a sample, comprising obtaining the sample, contacting the sample with the bispecific antibody comprising a first domain specifically binding PSMA and a second domain specifically binding CD3 provided herein, and detecting the antibody bound to PSMA and CD3 in the sample.

[00508] In some embodiments, the sample can be derived from urine, blood, serum, plasma, saliva, ascites, circulating cells, circulating tumor cells, cells that are not tissue associated (*i.e.*, free cells), tissues (*e.g.*, surgically resected tumor tissue, biopsies, including fine needle aspiration), histological preparations, and the like.

[00509] The antibodies described herein bound to PSMA or PSMA and CD3 can be detected using known methods. Exemplary methods include direct labeling of the antibodies using fluorescent or chemiluminescent labels, or radiolabels, or attaching to the antibodies provided herein a moiety, which is readily detectable, such as biotin, enzymes or epitope tags. Exemplary labels and moieties are ruthenium, ¹¹¹In-DOTA, ¹¹¹In-diethylenetriaminepentaacetic acid (DTPA), horseradish peroxidase, alkaline phosphatase and beta-galactosidase, poly-histidine (HIS tag), acridine dyes, cyanine dyes, fluorone dyes, oxazin dyes, phenanthridine dyes, rhodamine dyes and Alexafluor® dyes.

[00510] The antibodies provided herein can be used in a variety of assays to detect PSMA or PSMA and CD3 in the sample. Exemplary assays are western blot analysis, radioimmunoassay, surface plasmon resonance, immunoprecipitation, equilibrium dialysis, immunodiffusion, electrochemiluminescence (ECL) immunoassay, immunohistochemistry, fluorescence-activated cell sorting (FACS) or ELISA assay.

[00511] The PSMA antibodies or multispecific PSMA antibodies provided herein can be used to selectively enrich, isolate, separate, purify, sort, select, capture or detect PSMA-expressing cells. The PSMA antibodies or multispecific PSMA antibodies provided herein can be utilized in a bispecific format, *e.g.* containing a first antigen binding domain that specifically binds PSMA and a second antigen binding domain that specifically binds a second target. In some embodiments, the second target is CD3. In other embodiments, the multispecific PSMA antibodies provided herein can be utilized in a format that further incorporates a third antigen binding domain that specifically binds a third antigen (*e.g.*, at a trispecific antibody). In other embodiments, the multispecific PSMA antibodies provided herein can be utilized in a format that further incorporates a fourth antigen binding domain that specifically binds a fourth antigen. (*e.g.*, as a quadraspecific antibody).

[00512] In one aspect, provided herein is a method of enriching a PSMA-expressing cell comprising: providing a sample comprising the PSMA-expressing cell; contacting the sample with a PSMA antibody provided herein; and enriching the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of isolating a PSMA-expressing cell comprising: providing a sample comprising the PSMA-expressing cell; contacting the sample with a PSMA antibody provided herein; and isolating the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of separating a PSMA-expressing cell comprising: providing a sample comprising the PSMA-expressing cell; contacting the sample with a PSMA antibody provided herein; and separating the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of purifying a PSMA-expressing cell comprising: providing a sample comprising the PSMA-expressing cell; contacting the sample with a PSMA antibody provided herein; and purifying the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of sorting a PSMA-expressing cell comprising: providing a sample comprising the PSMA-expressing cell; contacting the sample with a PSMA antibody provided herein; and sorting the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of selecting a PSMA-expressing cell

comprising: providing a sample comprising the PSMA-expressing cell; contacting the sample with a PSMA antibody provided herein; and selecting the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of capturing a PSMA-expressing cell comprising: providing a sample comprising the PSMA-expressing cell; contacting the sample with a PSMA antibody provided herein; and capturing the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of detecting a PSMA-expressing cell comprising: providing a sample comprising the PSMA-expressing cell; contacting the sample with a PSMA antibody provided herein; and detecting the PSMA-expressing cell bound to the PSMA antibody.

[00513] In one aspect, provided herein is a method of enriching a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and enriching the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of isolating a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and isolating the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of separating a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and separating the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of purifying a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and purifying the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of sorting a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and sorting the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of selecting a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and selecting the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of capturing a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and capturing the PSMA-expressing cell bound to the PSMA antibody. In one aspect, provided herein is a method of detecting a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and detecting the PSMA-expressing cell bound to the PSMA antibody.

[00514] In one aspect, provided herein is a method of enriching a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and

enriching the PSMA-expressing cell based on binding of the PSMA-expressing cell to the PSMA antibody. In one aspect, provided herein is a method of isolating a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and isolating the PSMA-expressing cell based on binding of the PSMA-expressing cell to the PSMA antibody. In one aspect, provided herein is a method of separating a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and separating the PSMA-expressing cell based on binding of the PSMA-expressing cell to the PSMA antibody. In one aspect, provided herein is a method of purifying a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and purifying the PSMA-expressing cell based on binding of the PSMA-expressing cell to the PSMA antibody. In one aspect, provided herein is a method of sorting a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and sorting the PSMA-expressing cell based on binding of the PSMA-expressing cell to the PSMA antibody. In one aspect, provided herein is a method of selecting a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and selecting the PSMA-expressing cell based on binding of the PSMA-expressing cell to the PSMA antibody. In one aspect, provided herein is a method of capturing a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and capturing the PSMA-expressing cell based on binding of the PSMA-expressing cell to the PSMA antibody. In one aspect, provided herein is a method of detecting a PSMA-expressing cell comprising: contacting a PSMA-expressing cell with a PSMA antibody provided herein; and detecting the PSMA-expressing cell based on binding of the PSMA-expressing cell to the PSMA antibody.

[00515] In certain embodiments of the methods, the PSMA-expressing cell is a prostate cell. In some embodiments of the methods, the PSMA-expressing cell is in a population of cells. In some embodiments of the methods, the PSMA-expressing cell is in a population of prostate cells. In some embodiments of the methods, the PSMA-expressing cell is in a population of cancer cells. In some embodiments of the methods, the PSMA-expressing cell is provided as a population of cells. In some embodiments of the methods, the PSMA-expressing cell is provided as a population of prostate cells. In some embodiments of the methods, the PSMA-expressing cell is provided as a population of cancer cells. In some embodiments of the methods, the PSMA-expressing cell is provided as a sample comprising a population of cells. In some embodiments of the methods, the PSMA-expressing cell is

provided as a sample comprising a population of prostate cells. In some embodiments of the methods, the PSMA-expressing cell is provided as a sample comprising a population of cancer cells. In some embodiments of the methods, the sample is a blood sample. In some embodiments of the methods, the sample is a tissue sample. In some embodiments of the methods, the sample is a tissue culture sample.

[00516] Enrichment, isolation, separation, purification, sorting, selecting, capturing or detecting, or any combination thereof can be done using known technologies such as bead, microfluidics, solid support, columns, and the like. For example, PSMA cells can be separated or visualized using known methods when bound to the PSMA antibodies provided herein.

[00517] In some embodiments of the methods, the PSMA antibody is a multispecific PSMA antibody provided herein. In some embodiments of the methods, the PSMA antibody is a bispecific PSMA antibody provided herein. In some embodiments of the methods, the PSMA antibody is a trispecific PSMA antibody provided herein. In some embodiments of the methods, the PSMA antibody is a quadraspecific PSMA antibody provided herein. In certain embodiments, the PSMA antibody specifically binds to PSMA. In one embodiment, the multispecific PSMA antibody comprises: (a) a first binding domain that binds PSMA, and (b) a second binding domain that binds to a second target. In one embodiment, the multispecific PSMA antibody comprises: (a) a first binding domain that binds PSMA, and (b) a second binding domain that binds to a second target, and (c) a third binding domain that binds to a third target. In one embodiment, the multispecific PSMA antibody comprises: (a) a first binding domain that binds PSMA, and (b) a second binding domain that binds to a second target, (c) a third binding domain that binds to a third target, and (d) a fourth binding domain that binds to a fourth target. In one embodiment, the multispecific PSMA antibody comprises: (a) a first binding domain that specifically binds PSMA, and (b) a second binding domain that specifically binds to a second target. In one embodiment, the multispecific PSMA antibody comprises: (a) a first binding domain that specifically binds PSMA, and (b) a second binding domain that specifically binds to a second target, and (c) a third binding domain that specifically binds to a third target. In one embodiment, the multispecific PSMA antibody comprises: (a) a first binding domain that specifically binds PSMA, and (b) a second binding domain that specifically binds to a second target, (c) a third binding domain that specifically binds to a third target, and (d) a fourth binding domain that specifically binds to a fourth target.

[00518] In some embodiments of the methods, the PSMA antibody is a multispecific PSMA antibody, wherein the second target is CD3.

[00519] In some embodiments, the first binding domain of the multispecific PSMA antibody specifically binds PSMA. In some embodiments, the first binding domain of the multispecific PSMA antibody specifically binds to *Pan troglodytes* (chimpanzee, chimp) PSMA. In other embodiments, the first binding domain of the multispecific PSMA antibody specifically binds to *Macaca fascicularis* (cynomolgus monkey, macaque, cyno) PSMA. In yet other embodiments, the first binding domain of the multispecific PSMA antibody specifically binds to and/or human PSMA. In some embodiments, the first binding domain of the multispecific PSMA antibody specifically binds to *Pan troglodytes*, *Macaca fascicularis*, and human PSMA. In specific embodiments, the first binding domain of the multispecific PSMA antibody specifically binds to both cyno and human PSMA.

[00520] In specific embodiments of the methods provided herein, the method uses multi-marker detection. In some embodiments, the multi-marker detection uses a multispecific PSMA antibody provided herein. In some embodiments, the multi-marker detection uses a bispecific PSMA antibody provided herein. In some embodiments, the multi-marker detection uses a trispecific PSMA antibody provided herein. In some embodiments, the multi-marker detection uses a quadraspecific PSMA antibody provided herein.

[00521] In certain embodiments of the methods provided herein, the methods are included as steps in a diagnostic method. In certain embodiments of the methods provided herein, the methods are included as steps in a method to quantify the PSMA-expressing T cells.

[00522] In certain embodiments of the methods provided herein, the method further comprises expanding the enriched, isolated, separated, purified, sorted, selected, captured or detected PSMA-expressing cells. In certain embodiments, the expanding is *in vitro*. In certain embodiments, the expanding is *in vivo*. In certain embodiments of the methods provided herein, the method further comprises growing the enriched, isolated, separated, purified, sorted, selected, captured or detected PSMA-expressing cells. In certain embodiments, the growing is *in vitro*. In certain embodiments, the growing is *in vivo*. In certain embodiments of the methods provided herein, the method further comprises quantifying the enriched, isolated, separated, purified, sorted, selected, captured or detected PSMA-expressing cells.

Kits

[00523] Also provided are kits comprising a PSMA antibody provided herein, and instructions for use.

[00524] The described kits can be used to carry out the methods of using the PSMA-specific antibodies or antigen-binding fragments provided herein, or other methods known to those skilled in the art. In some embodiments, the described kits can include the antibodies or antigen-binding fragments described herein and reagents for use in detecting the presence of PSMA in a biological sample. Accordingly, the described kits can include one or more of the antibodies, or an antigen-binding fragment(s) thereof, described herein and a vessel for containing the antibody or fragment when not in use, instructions for use of the antibody or fragment, the antibody or fragment affixed to a solid support, and/or detectably labeled forms of the antibody or fragment, as described herein.

[00525] In one embodiment, provides is a kit comprising a PSMA antibody provided herein.

[00526] In another embodiment, provided is a kit comprising the bispecific PSMAxCD3 antibody comprising a first domain specifically binding PSMA and a second domain specifically binding CD3 provided herein.

[00527] The kit can be used for therapeutic uses and as diagnostic kits. The kit can also be used to detect the presence of PSMA, CD3 or PSMA and CD3 in a biological sample.

[00528] In some embodiments, the kit comprises an antibody described herein and reagents for detecting the antibody. The kit can further include one or more other elements including: instructions for use; other reagents, *e.g.*, a label, a therapeutic agent, or an agent useful for chelating, or otherwise coupling, an antibody to a label or therapeutic agent, or a radioprotective composition; devices or other materials for preparing the antibody for administration; pharmaceutically acceptable carriers; and devices or other materials for administration to a subject.

[00529] In some embodiments, the kit comprises the antibody provided herein in a container and instructions for use of the kit.

[00530] In some embodiments, the antibody in the kit is labeled.

EMBODIMENTS

[00531] This invention provides the following non-limiting embodiments:

1. An isolated antibody that binds to PSMA comprising:

- (1) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:32;
- (2) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:66;
- (3) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100;
- (4) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134;
- (5) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100;
- (6) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3,

- respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134;
- (7) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236;
- (8) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
- (9) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236;
- (10) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
- (11) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:371; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an

amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3,
respectively, of a VL having an amino acid sequence of SEQ ID NO:372;

- (12) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:405; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:406;
- (13) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
- (14) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:473; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474;
- (15) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:507; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508;
- (16) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:541; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:542;

- (17) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:575; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:576;
- (18) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100;
- (19) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:643; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508;
- (20) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:677; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:678;
- (21) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:711; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474;
- (22) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3,

- respectively, of a VH having an amino acid sequence of SEQ ID NO:745; and
- (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:746;
- (23) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:779; and
- (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:780; or
- (24) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:813; and
- (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:814.
2. The isolated antibody of embodiment 1, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences are according to the
- (i) Kabat numbering system;
- (ii) Chothia numbering system;
- (iii) AbM numbering system;
- (iv) Contact numbering system;
- (v) IMGT numbering system or a combination thereof.
3. The isolated antibody of embodiment 1 or embodiment 2, further comprising a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:31, 99, 167, 235, 303, 371, 405, 439, 473, 507, 541, 575, 643, 677, 711, 779 or 813 and a VL having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:32, 66, 100, 134, 236, 270, 372, 406, 474, 508, 542, 576, 678, 746, 780 or 814.
4. The isolated antibody of any one of embodiments 1 to 3, further comprising a heavy chain (HC) having an amino acid sequence at least 80% identical to the amino acid

sequence of SEQ ID NOs:33, 101, 169, 237, 305, 373, 407 or 441 and a light chain (LC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:34, 68, 102, 136, 238, 272, 374 or 408.

5. An isolated antibody that binds PSMA comprising
 - a. (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
 - b. (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:205, 206, and 411, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:242, 209, and 244, respectively;
 - c. (i) a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL having an amino acid sequence of SEQ ID NO:270; or
 - d. (i) a HC having an amino acid sequence of SEQ ID NO:441; and (ii) a LC having an amino acid sequence of SEQ ID NO:272.
6. The isolated antibody of any one of embodiments 1 to 5, wherein the antibody is a humanized antibody.
7. The isolated antibody of any one of embodiments 1 to 6, wherein the antibody is a human antibody.
8. The isolated antibody of any one of embodiments 1 to 7, wherein the binding domain that binds to PSMA is a scFv, an scFv dimer, a Fv, a Fab, a Fab, a F(ab')₂, a dsFv, a sdAb, a VHH or a single chain antibody.
9. The isolated antibody of any one of embodiments 1 to 8, wherein the antibody is an IgG antibody.
10. The isolated antibody of embodiment 9, wherein the IgG antibody is an IgG1, IgG2, IgG3, or IgG4 antibody.

11. The isolated antibody of any one of embodiments 1 to 10, wherein the antibody comprises a kappa light chain.
12. The isolated antibody of any one of embodiments 1 to 10, wherein the antibody comprises a lambda light chain.
13. The isolated antibody of any one of embodiments 1 to 12, wherein the antibody is a monoclonal antibody.
14. The isolated antibody of any one of embodiments 1 to 13, wherein the antibody binds a PSMA antigen.
15. The isolated antibody of any one of embodiments 1 to 14, wherein the antibody binds a PSMA epitope.
16. The isolated antibody of any one of embodiments 1 to 15, wherein the antibody specifically binds to PSMA.
17. The isolated antibody of any one of embodiments 1 to 16, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 form a binding site for an antigen of the PSMA.
18. The isolated antibody of any one of embodiments 1 to 16, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 form a binding site for an epitope of the PSMA.
19. The isolated antibody of any one of embodiments 1 to 18, wherein the PSMA is present on the surface of a cell.
20. The isolated antibody of embodiment 19, wherein the cell is a prostate cell.
21. The isolated antibody of embodiment 19, wherein the cell is a prostate cancer cell.
22. The isolated antibody of embodiment 19, wherein the cell is a renal cell.
23. The isolated antibody of embodiment 19, wherein the cell is a renal cancer cell.
24. The isolated antibody of embodiment 23, wherein the renal cancer is a metastatic renal cell carcinoma.
25. The isolated antibody of any one of embodiments 1 to 24, wherein the antibody is multivalent.

26. The isolated antibody of embodiment 25, wherein the antibody is capable of binding at least three antigens.
27. The isolated antibody of embodiment 25, wherein the antibody is capable of binding at least four antigens.
28. The isolated antibody of embodiment 25, wherein the antibody is capable of binding at least five antigens.
29. The isolated antibody of any one of embodiments 1 to 28, wherein the antibody is a multispecific antibody.
30. The isolated antibody of embodiment 29, wherein the antibody is a bispecific antibody.
31. The isolated antibody of embodiment 29, wherein the antibody is a trispecific antibody.
32. The isolated antibody of embodiment 29, wherein the antibody is a quadraspecific antibody.
33. A nucleic acid encoding the antibody of any one of embodiments 1 to 32.
34. A vector comprising the nucleic acid of embodiment 33.
35. A host cell comprising the vector of embodiment 34.
36. A kit comprising the vector of embodiment 34 and packaging for the same.
37. A kit comprising the antibody of any one of embodiments 1 to 32 and packaging for the same.
38. A pharmaceutical composition comprising the antibody of any one of embodiments 1 to 32, and a pharmaceutically acceptable carrier.
39. A method of producing the pharmaceutical composition of embodiment 38, comprising combining the antibody with a pharmaceutically acceptable carrier to obtain the pharmaceutical composition.
40. An isolated multispecific antibody, comprising:
 - (a) a first binding domain that binds to PSMA, wherein the first binding domain comprises:

- (1) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:32;
- (2) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:66;
- (3) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100;
- (4) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134;
- (5) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100;
- (6) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3,

- respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134;
- (7) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236;
- (8) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
- (9) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236;
- (10) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
- (11) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:371; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an

amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3,
respectively, of a VL having an amino acid sequence of SEQ ID NO:372;

- (12) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:405; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:406;
- (13) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
- (14) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:473; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474;
- (15) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:507; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508;
- (16) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:541; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:542;

- (17) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:575; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:576;
- (18) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100;
- (19) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:643; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508;
- (20) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:677; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:678;
- (21) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:711; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474;
- (22) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3,

- respectively, of a VH having an amino acid sequence of SEQ ID NO:745; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:746;
- (23) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:779; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:780; or
- (24) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:813; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:814; and
- (b) a second binding domain that binds to CD3, wherein the second binding domain comprises:
- (1) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:1505; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:1464;
- (2) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:915; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:916;
- (3) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3,

- respectively, of a VH having an amino acid sequence of SEQ ID NO:983; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:984;
- (4) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:1463; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:1464;
- (5) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:847; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:848; or
- (6) a scFv comprising a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, a VL CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, a VL CDR3 respectively, of a scFv having an amino acid sequence of SEQ ID NO:1524.
41. The isolated multispecific antibody of embodiment 40, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences are according to the
- (i) Kabat numbering system;
 - (ii) Chothia numbering system;
 - (iii) AbM numbering system;
 - (iv) Contact numbering system;
 - (v) IMGT numbering system or a combination thereof.
42. The isolated multispecific antibody of embodiment 40 or embodiment 41, wherein the first binding domain that binds PSMA comprises a heavy chain (HC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:33, 101, 169, 237, 305, 373, 407, 441, 1242, 1244, 1248, 1250, 1252, 1254, 1256, 1258,

- 1260, 1262, 1264, 1266, 1268 or 1270 and/or a light chain (LC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:34, 68, 102, 136, 238, 272, 374 or 408.
43. The isolated multispecific antibody of any one of embodiments 40 to 42, wherein the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of any one of SEQ ID NOs:1485-1500 or SEQ ID NOs:1526-1531.
44. The isolated multispecific antibody of any one of embodiments 40 to 43, wherein the first binding domain that binds PSMA comprises (i) a HC having an amino acid sequence of SEQ ID NO:441; and (ii) a LC having an amino acid sequence of SEQ ID NO:272.
45. The isolated multispecific antibody of any one of embodiments 40 to 44, wherein the second binding domain that binds CD3 comprises a heavy chain (HC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:849, 883, 917, 951, 985, 1019, 1504, 1455, 1192, 1194, 1167, 1218 or 1238 and/or a light chain (LC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:850, 918, 986, 1193, 1195 or 1219.
46. The isolated multispecific antibody of any one of embodiments 40 to 45, wherein the second binding domain that binds CD3 comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:1186, 1187, 1523 or 1524.
47. An isolated bispecific antibody comprising a first binding domain that binds PSMA and a second binding domain that binds CD3, wherein
- the first binding domain that binds PSMA comprises (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270, and the second binding domain that binds CD3 comprises a scFv comprising a VH CDR1, a VH CDR2, a VH CDR3, a

- VL CDR1, a VL CDR2, and a VL CDR3 of a scFv having an amino acid sequence of SEQ ID NO:1524;
- b. the first binding domain that binds PSMA comprises a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:205, 206, 411, 242, 209 and 244, respectively, wherein the amino acid sequences are according to the Kabat numbering system; and the second binding domain that binds CD3 comprises a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:1467, 1468, 1506, 1470, 1471 and 1472, respectively; wherein the amino acid sequences are according to the Kabat numbering system;
 - c. the first binding domain that binds PSMA comprises (i) a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL having an amino acid sequence of SEQ ID NO:270, and the second binding domain that binds CD3 comprises a scFv of SEQ ID NO:1524, respectively; and/or
 - d. the first binding domain that binds PSMA comprises (i) a HC2 having an amino acid sequence of SEQ ID NO:441; and (ii) a LC2 having an amino acid sequence of SEQ ID NO:272, respectively; and the second binding domain that binds CD3 comprises a HC1 of SEQ ID NO:1455.
48. The isolated multispecific antibody of any one of embodiments 40 to 47, wherein the antibody is a humanized antibody.
49. The isolated multispecific antibody of any one of embodiments 40 to 48, wherein the antibody is a human antibody.
50. The isolated multispecific antibody of any one of embodiments 40 to 49, wherein the first binding domain, the second binding domain and/or the first and second is a scFv, an scFv dimer, a Fv, a Fab, a Fab, a F(ab')₂, a dsFv, a sdAb, a VHH or a single chain antibody.
51. The isolated multispecific antibody of any one of embodiments 40 to 50, wherein the antibody is an IgG antibody.
52. The isolated multispecific antibody of embodiment 51, wherein the IgG antibody is an IgG1, IgG2, IgG3, or IgG4 antibody.

53. The isolated multispecific antibody of any one of embodiments 40 to 52, wherein the antibody comprises a kappa light chain.
54. The isolated multispecific antibody of any one of embodiments 40 to 52, wherein the antibody comprises a lambda light chain.
55. The isolated multispecific antibody of any one of embodiments 40 to 54, wherein the antibody is a monoclonal antibody.
56. The isolated multispecific antibody of any one of embodiments 40 to 55, wherein the first binding domain binds a PSMA antigen.
57. The isolated multispecific antibody of any one of embodiments 40 to 55, wherein first binding domain binds a PSMA epitope.
58. The isolated multispecific antibody of any one of embodiments 40 to 57, wherein the first binding domain specifically binds to PSMA.
59. The isolated multispecific antibody of any one of embodiments 40 to 58, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the first binding domain form a binding site for an antigen of the PSMA.
60. The isolated multispecific antibody of any one of embodiments 40 to 58, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the first binding domain form a binding site for an epitope of the PSMA.
61. The isolated multispecific antibody of any one of embodiments 40 to 60, wherein the second binding domain binds a CD3 antigen.
62. The isolated multispecific antibody of any one of embodiments 40 to 61, wherein the second binding domain binds a CD3 epitope.
63. The isolated multispecific antibody of any one of embodiments 40 to 62, wherein the second binding domain specifically binds to CD3.
64. The isolated multispecific antibody of any one of embodiments 40 to 63, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the second binding domain form a binding site for an antigen of the CD3.
65. The isolated multispecific antibody of any one of embodiments 40 to 63, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the second binding domain form a binding site for an epitope of the CD3.

66. The isolated multispecific antibody of any one of embodiments 40 to 65, wherein the PSMA is present on the surface of a cell.
67. The isolated multispecific antibody of embodiment 66, wherein the cell is a prostate cell.
68. The isolated multispecific antibody of embodiment 66, wherein the cell is a prostate cancer cell.
69. The isolated multispecific antibody of embodiment 66, wherein the cell is a renal cell.
70. The isolated multispecific antibody of embodiment 66, wherein the cell is a renal cancer cell.
71. The isolated multispecific antibody of embodiment 66, wherein the renal cancer is a metastatic renal cell carcinoma.
72. The isolated multispecific antibody of any one of embodiments 40 to 71, wherein the antibody is a bispecific antibody.
73. The isolated multispecific antibody of any one of embodiments 40 to 71, wherein the antibody is a trispecific antibody.
74. The isolated multispecific antibody of any one of embodiments 40 to 71, wherein the antibody is a quadraspecific antibody.
75. A nucleic acid encoding the multispecific antibody of any one of embodiments 40 to 74.
76. A vector comprising the nucleic acid of embodiment 75.
77. A host cell comprising the vector of embodiment 76.
78. A kit comprising the vector of embodiment 76 and packaging for the same.
79. A kit comprising the multispecific antibody of any one of embodiments 40 to 74 and packaging for the same.
80. A pharmaceutical composition comprising the multispecific antibody of any one of embodiments 40 to 74, and a pharmaceutically acceptable carrier.
81. A method of producing the pharmaceutical composition of embodiment 80, comprising combining the multispecific antibody with a pharmaceutically acceptable carrier to obtain the pharmaceutical composition.

82. A method of directing a CD3-expressing T cell to a PSMA-expressing target cell, comprising contacting the T cell with the multispecific antibody of any one of embodiments 40 to 74.
83. The method of embodiment 82, wherein the contacting directs the T cell to the target cell.
84. A method of inhibiting the growth or proliferation of a PSMA-expressing target cell, comprising contacting the target cell with the multispecific antibody of any one of embodiments 40 to 74.
85. The method of embodiment 84, wherein the contacting is in the presence of CD3-expressing T cells.
86. A method of eliminating a PSMA-expressing target cell in a subject, comprising administering to the subject an effective amount of the multispecific antibody of any one of embodiments 40 to 74.
87. The method of embodiment 86, wherein the subject has a disease or disorder of the prostate.
88. The method of embodiment 86, wherein the subject has prostate inflammation.
89. The method of embodiment 86, wherein the subject has Benign prostatic hyperplasia.
90. The method of embodiment 86, wherein the subject has a prostate cancer.
91. The method of embodiment 90, wherein the subject has metastatic castration-resistant prostate cancer (mCRPC).
92. The method of any one of embodiments 86 to 91, wherein the target cell expresses PSMA on the cell surface.
93. The method of any one of embodiments 86 to 91, wherein the target cell is a prostate cell.
94. The method of any one of embodiments 86 to 91, wherein the target cell is a prostate cancer cell.
95. The method of embodiment 86, wherein the subject has a renal disease or disorder.
96. The method of embodiment 86, wherein the subject has a renal cancer.

97. The method of embodiment 86, wherein the subject has a metastatic renal cell carcinoma.
98. The method of any one of embodiments 86 or 95 to 97, wherein the target cell expresses PSMA on the cell surface.
99. The method of any one of embodiments 86 or 95 to 97, wherein the target cell is a renal cell.
100. The method of any one of embodiments 86 or 95 to 97, wherein the target cell is a renal cancer cell.
101. The method of any one of embodiments 86 or 95 to 97, wherein the target cell is a metastatic renal cell carcinoma cell.
102. The method of any one of embodiments 86 to 101, wherein the subject is a subject in need thereof.
103. A method of treating a disease or disorder in a subject, comprising administering to the subject an effective amount of the multispecific antibody of any one of embodiments 40 to 74.
104. The method of embodiment 103, wherein the disease or disorder is a disease or disorder of the prostate.
105. The method of embodiment 103, wherein the disease or disorder of the prostate is prostate inflammation.
106. The method of embodiment 103, wherein the disease or disorder of the prostate is Benign prostatic hyperplasia.
107. The method of embodiment 103, wherein the disease or disorder of the prostate is a prostate cancer.
108. The method of embodiment 107, wherein the disease or disorder is metastatic castration-resistant prostate cancer (mCRPC).
109. The method of embodiment 103, wherein the disease or disorder is a renal disease or disorder.
110. The method of embodiment 103, wherein the renal disease or disorder is a renal cancer.

111. The method of embodiment 103, wherein the renal disease or disorder is a metastatic renal cell carcinoma.
112. The method of any one of embodiments 103 to 111, wherein the subject is a subject in need thereof.

[00532] Provided in the Examples herein are exemplary antibodies that bind to PSMA. Also provided in the Examples herein are exemplary bispecific antibodies that bind to PSMA and CD3.

[00533] In some embodiments, provided herein is a bispecific antibody comprising: (a) a first binding domain that binds to a PSMA antigen, and (b) a second binding domain that binds to a second target antigen. In some embodiments, provided herein is a bispecific antibody comprising: (a) a first binding domain that specifically binds to a PSMA antigen, and (b) a second binding domain that specifically binds to a second target antigen. In some embodiments, provided herein is a bispecific antibody comprising: (a) a first binding domain that binds to a first epitope on a PSMA antigen, and (b) a second binding domain that binds to a second epitope on a second target antigen. In some embodiments, provided herein is a bispecific antibody comprising: (a) a first binding domain that specifically binds to a first epitope on a PSMA antigen, and (b) a second binding domain that specifically binds to a second epitope on a second target antigen. In certain embodiments, the second target antigen is CD3.

[00534] Exemplary binding agents that bind to PSMA are provided herein, for example in the Examples, as well as Tables 4-12.

[00535] Exemplary binding agents (*e.g.*, multispecific PSMA antibodies) can comprise a second binding domain that binds to CD3, such as those provided in Tables 16-22.

[00536] In addition, exemplary binding agents that bind to PSMA and CD3 are provided elsewhere herein, for example in the Examples, as well as Tables 23-28.

[00537] Particular embodiments of this invention are described herein. Upon reading the foregoing description, variations of the disclosed embodiments can become apparent to individuals working in the art, and it is expected that those skilled artisans can employ such variations as appropriate. Accordingly, it is intended that the invention be practiced otherwise than as specifically described herein, and that the invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in

all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context. A number of embodiments of the invention have been described. Nevertheless, it will be understood that various modifications can be made without departing from the spirit and scope of the invention. Accordingly, the descriptions in the Examples section are intended to illustrate but not limit the scope of invention described in the claims.

EXAMPLES

[00538] The following examples are based on the premise that bispecific anti-PSMA/anti-CD3 antibodies, or antigen-binding fragments thereof, have potent anti-tumor functions. Prostate tumor cells express prostate specific membrane antigen (PSMA), a type II membrane protein that is highly expressed in prostatic intraepithelial neoplasia (PIN), and aggressive prostate cancers. CD3 is expressed on T cells and is involved in T cell activation and differentiation into an effector cell. This ability of CD3 to activate T cells is harnessed using bispecific antibodies constructed such that one arm binds to the PSMA, and the other arm binds to CD3 on the T cells. Thus, the bispecific antibody bridges together effector and target cells, resulting in tumor killing.

EXAMPLE 1: ANTIGEN GENERATION

[00539] Human PSMA extracellular domain (ECD) was produced based on Uniprot Accession #Q04609 sequence. The ECD construct was designed with a 6X His-tag sequence (SEQ ID NO:1484) at the N-terminus (construct PSMW39; SEQ ID NO:1023). The PSMA ECD from *Macaca fascicularis* was produced based on NCBI Accession# EHH56646.1, appending the sequence at the N-terminus with an Avi-tag fused to a 6x polyhistidine tag (SEQ ID NO:1484) (construct PSMW1; SEQ ID NO:1024). The extracellular domain (ECD) of PSMA from *Mus musculus* was produced based on Uniprot Accession# O35409, appending the ECD sequence at the N-terminus with an Avi-tag fused to a 6x polyhistidine tag (SEQ ID NO:1484) for purification (construct PSMW29; SEQ ID NO:1025). The cynomolgus and murine ECD expression constructs were used to transiently transfect HEK293-6E. Cells were incubated for six days at 37 °C with 5% CO₂ prior to harvest. The human ECD expression construct was used to stably transfect Expi-CHO using EXPIFECTAMINE-CHO. Stable transfectants were selected with G418 (Thermo Fisher Scientific; Cat# 10131027). Monoclones expressing the highest levels of human PSMA ECD protein were identified by ELISA. High PSMW39-expressing monoclones were

cultured in DYNAMIS media supplemented with 5% Cell Boost 5 and incubated at 37 °C, 125 rpm and 5% CO₂ until the cell viability dropped below 80%. On days 2, 4, 7, 9 and 11, 6 mL of 20% D-Glucose (2 g/L final conc.) and 6 mL of 200 mM L-Glutamine (2mM final conc.) was added as feed to each of the flasks. For the purification of all recombinant PSMA proteins, the expressed cells were removed by centrifugation and the soluble PSMA proteins with his-tags were purified from the media using immobilized metal affinity chromatography using His 60 Ni SUPERFLOW Resin, Clonotech, Cat # 635662 followed by SUPERDEX 200 preparative size exclusion chromatography (SEC) (GE Healthcare) and formulation into 1X DPBS, pH 7.2 containing 1 mM CaCl₂, 0.5 mM MgCl₂, and 0.5 mM ZnCl₂. The isolation of the homodimeric species was confirmed via analytical size exclusion chromatography. The amino acid sequences of the recombinant antigens are shown in Table 3.

Table 3: The amino acid sequences of the recombinant PSMA antigens.

Protein AA ID	Description	Amino Acid Sequence
PSMW39	Human PSMA ECD (Lys44-Ala750), N-6xHis	HHHHHHKSSNEATNITPKHNMKAFLDELKAENIKKFLYNFTQIPHLAGTEQNFQLAKQIQSQWKEFGLDSVELAHYDVLLSYPNKTHPNYISIIINEDGNEIFNTSLFEPPPPGYENVSDIVPPFSAFSPQGMPEGDLVYVNYARTEDFFKLERDMKINC SGKIVARIYKGVFRGNKVKNQAQLAGAKGVILYSDPADYFAPGVKSYPDGWNLPGGGVQQRGNILNLNGAGDPLTPGYPANAYAYRRGIAEAVGLPSIPVHPIGYYDAQKLEKMGGSAPPDSSWRGSLKVPYNVGPFGFTGNFSTQKVKMHIHSTNEVTRIYNVIGTLRGAVEPDRYVILGGHRDSWVFGGIDPQSGAAVVHEIVRSFGTLKKEGWRPRRTILFASWDAEEFGLLGSTEWAEENSRLLQERGVAYINADSSIEGNYTLRVDCTPLMYSLVNLTKEKSPDEGFEGKSLYESWTKKSPSPFSGMPRISKLGSNDFEVFFQRLGIASGRARYTKNWETNKFSGYPLYHSVYETVELVEKFYDPMFKYHLTVQVRGGMVFELANSIVLPFDCRDYAVVLRKYADKIYISMKHPQEMKTYSVSFDLSFSAVKNFTEIASKFSERLQDFDKSNPIVLRMMNDQLMFLEAFIDPLGLPDRPFYRHVIYAPSSHNKYAGESFFPGIYDALFDIESKVDPSKAWGEVQRQIYVAAFVQAAAETLSEVA (SEQ ID NO:1023)
PSMW1	Avi-6xHis-GS-cyno PSMA ECD	GLNDIFEAQKIEWHEHHHHHSGSKSSSEATNITPKHNMKAFLDELKAENIKKFLHNFTQIPHLAGTEQNFQLAKQIQSQWKEFGLDSVELTHYDVLLSYPNKTHPNYISIIINEDGNEIFNTSLFEPPPPAGYENVSDIVPPFSAFSPQGMPEGDLVYVNYARTEDFFKLERDMKINC SGKIVARIYKGVFRGNKVKNQAQLAGATGVILYSDPDYFAPGVKSYPDGWNLPGGGVQQRGNILNLNGAGDPLTPGYPANAYAYRRGMAEAVGLPSIPVHPIGYYDAQKLEKMGGSASPDSSWRGSLKVPYNVGPFGFTGNFSTQKVKMHIHSTSEVTRIYNVIGTLRGAVEPDRYVILGGHRDSWVFGGIDPQSGAAVVHEIVRSFGMLKKEGWRPRRTILFASWDAEEFGLLGSTEWAEENSRLLQERGVAYINADSSIEGNYTLRVDCTPLMYSLVNLTKELESPEDEGFEGKSLYESWTKKSPSPFSGMPRISKLGSNDFEVFFQRLGIASGRARYTKNWETNKFSSYPLYHSVYETVELVEKFYDPMFKYHLTVQVRGGMVFELANSVLPFDCRDYAVVLRKYADKIYINISMKHPQEMKTYSVSFDLSFSAVKNFTEIASKFSERLRDFDKSNPIVLRMMNDQLMFLERAFIDPLGLPDRPFYRHVIYAPSSHNKYAGESFFPGIYDALFDIESKVDPSQAWGEVQRQISIAFTTVQAAAETLSEVA (SEQ ID NO:1024)
PSMW29	Avi-6xHis-GS-mouse	GLNDIFEAQKIEWHEHHHHHSGSKPSNEATGNVSHSGMKKEFLHDLKAENIKKFLYNFTRTPHLAGTQNNFELAKQIHDQWKEFGLDLVELSHYDVLLSYPNKTHPNYISIIINEDGNEIFKTSLSEQPPPGYENISDVVPPYSAFSPQGTPEGDLVYVNYARTEDFFKLEREMKISCSGKIVARIYKGVFRGNMVKNAQL

	PSMA (45--752)	AGAKGMILYSDEADYFVPAVKSYPDGWNLPGGGVQRGNVLNLNGAGDPLT PGYPANEHAYRHELTNAVGLPSIPVHPIGYDDAQKLEHMGGFAPPDSSW KGGLKVPYNVGPGEAGNFSTQKVKMHIHSYTKVTRIYNVIGTLKGALEPD RYVILGGHRDAWVFGGIDPQSGAAVVHEIVRSFGTLKKKGRRPRRTILFA SWDAEEFGLLGSTEWAEHSRLLQERGVAYINADSSIENYTLRVDCTPL MYSLVYNLTKEQSPDEGFEGKSLYDSWKEKSPSPFEFIGMFRISKLGSGN DFEVFFQRLGIASGRARYTKNWKTNKVSSYPLYHSVYETVELVVKFYDPT FKYHLTVAQVRGAMVFELANSIVLFFDCQSYAVALKKYADTIYNISMKHP QEMKAYMISFDSLFSAVNNFTDVASKFNQRLQELDKSNPILLRIMNDQLM YLERAFIDPLGLPGRPFYRHIIYAPSSHKNKYAGESFPGIYDALFDISSKV NASKAWNEVKRQISIATFTVQAAAETLREVA (SEQ ID NO:1025)
--	----------------	--

EXAMPLE 2 – GENERATION OF ANTI-PSMA ANTIBODIES**EXAMPLE 2.1: ANTIBODY GENERATION USING TRANSGENIC ABLEXIS® MICE**

[00540] Ablexis® kappa, lambda, and kappa/lambda hybrid mice were immunized with recombinant human PSMA antigen (PSMW39.002) and cynomolgus PSMA antigen (PSMW1.009) in combination with CL413 adjuvant (InvivoGen, VAC-C413-5).

[00541] Ablexis® mice generate antibodies having human variable domains linked to human CH1 and CL domains, chimeric human/mouse hinge region, and mouse Fc regions. Ablexis® Kappa Mouse and Lambda Mouse strains are distinguished by which of their heavy chains are human or mouse as noted below. Antibodies produced by the Kappa Mouse lack sequence derived from mouse VH, DH and JH exons and mouse V κ , J κ and C κ exons. The endogenous mouse Ig λ is active in the Kappa Mouse. The human Ig κ chains comprise approximately 90-95% of the naïve repertoire and mouse Ig λ chains comprise approximately 5-10% of the naïve repertoire in this strain. Antibodies produced by the Lambda Mouse lack sequence derived from mouse VH, DH and JH exons and mouse V λ , J λ and C λ exons. The endogenous mouse Ig κ is active in the Lambda Mouse. The human Ig λ chains comprise approximately 40% of the naïve repertoire and mouse Ig κ chains comprise approximately 60% of the naïve repertoire. The preparation and use of Ablexis®, and the genomic modifications carried by such mice, is described in WO11/123708.

[00542] Mice were boosted at Days 0, 7, 14, 21, and 28 before being bled on Day 35 for serological analysis. Serology was performed on human PSMA (+) cells C4-2B (AG000002300) and a human PSMA knockout cell line (AG000002521). In total, 8 mice were selected for hybridoma fusion and final boosted on Day 56 with PSMW39.002 or an equimolar mixture of PSMW39.002 and PSMW1.009. This preparation also included recombinant anti-mouse CD40 mAb (R&D Systems, MAB440) to stimulate B cell expansion. On Day 60, spleen and draining lymph nodes were harvested from these mice, pooled and homogenized into a single cell suspension. Stable hybridomas were generated by PEG-mediated fusion of mouse myeloma cell line FO with the pooled mouse homogenate, followed by HAT selection.

[00543] Supernatants from these hybridomas were screened against C4-2B cells by MSD. From this primary screen, 440 positive samples were identified and re-arrayed for

confirmatory screening. Confirmatory screening was performed by ELISA as well as Fluorescence-activated cell sorting (FACS) to validate binding to PSMA protein and PSMA (+) C4-2B cells, respectively. To ensure specificity, samples were also screened against an irrelevant negative control sample TrRW2. Based on the screening results, 214 samples were advanced past confirmatory screening from the 440 samples identified and isotyped for kappa or lambda light chain expression.

EXAMPLE 2.2: V REGION CLONING

[00544] RNA from hybridoma were purified using RNEASY Plus Mini Kit (Qiagen) and used for cDNA synthesis using the SMARTER cDNA synthesis kit (Clontech, Mount View, CA). To facilitate cDNA synthesis, oligodT was used to prime reverse transcription of all messenger RNAs followed by “5’ capping” with a Smarter IIA oligonucleotide. Subsequent amplification of the VH and VL fragments was performed using a 2-step PCR amplification using 5’ primers targeting the Smarter IIA cap and 3’ primers targeting consensus regions in CH1. Briefly, each 50 µl PCR reaction consists of 20 µM of forward and reverse primer mixes, 25 µl of PRIMESTAR Max DNA polymerase premix (Clontech), 2 µl of unpurified cDNA, and 21 µl of double-distilled H₂O. The cycling program starts at 94°C for 3 min, followed by 35 cycles (94°C for 30 Sec, 55°C for 1 min, 68°C for 1 min), and ends at 72°C for 7 min. The second round PCR was performed with VL and VH 2nd round primers containing 15bp complementary extensions that “overlap” respective regions in their respective Lonza mother vector (VH and VL). Second round PCR was performed with the following program: 94°C for 3 min; 35 cycles (94°C for 30 Sec, 55°C for 1 min, 68°C for 1 min), and ends at 72°C for 7 min. In-Fusion® HD Cloning Kit (Clontech, U.S.A.) was used for directional cloning of VL gene into Lonza huIgK or Lambda vector and VH gene into Lonza huIgG1 vector. To facilitate In-Fusion® HD Cloning, PCR products were treated with Cloning Enhancer before In-Fusion HD Cloning. The resulting PCR fragments were sequenced. The amino-acid sequences of the recovered-regions were codon optimized and cloned into an expression vector carrying the IgG1 constant region with L234A, L235A and D265S mutations for Fc silencing (IgG1 AAS isotype), and the K409R mutation for heterodimerization or an expression vector carrying the IgG4 constant region with S228P, F234A and L235A mutations (IgG4PAA isotype). D365E mutation was also introduced. Cloning and transformation were performed according to manufacturer’s protocol

(Clontech, U.S.A.). Mini-prep DNAs were subjected to Sanger sequencing to confirm that complete V-gene fragments were obtained.

EXAMPLE 2.3: SCFV AND SCFV-FC FORMATTING

[00545] Following cloning of the variable regions, the VH and VL sequences were used to generate antibodies. Antibodies were expressed in a Fab format, a mAb format, a scFv format in the VH-linker-VL orientation or a scFv format in VL-linker-VH orientation and were further analyzed as described below. The linker sequence GGSEGGKSSGSGSESKSTGGS (SEQ ID NO:1419) was used to conjugate the VH/VL regions. A synthetic DNA fragment coding for the scFv moiety was designed using codons optimized for *Cricetulus griseus* (Chinese hamster) and was cloned between the HindIII and PstI sites within a mammalian expression vector containing the constant region.

EXAMPLE 2.4: EXPICHO TRANSFECTION AND PURIFICATION

[00546] PSMB890 identified from the immunization campaigns was cloned and expressed as IgG4PAA and purified. PSMB891, PSMB892, PSMB893, PSMB894, PSMB895, PSMB896, PSMB897, PSMB898 and PSMB899 were cloned and expressed as IgG1AAS and purified. ExpiCHO-S™ cells (ThermoFisher Scientific) were seeded at 1.25×10^5 – 2.25×10^5 viable cells/mL density in ExpiCHO™ Expression Medium and cultured in a 37°C, 7% CO₂ shaker incubator. For routine cell growth in 125 mL – 2L shake flasks; the shake speed was set to 130 rpm. Cells were sub-cultured when density reached log phase growth at 4×10^6 – 6×10^6 viable cells/mL with a 98-99% viability. Two days before transfection, ExpiCHO-S™ cells were seeded at 1.5×10^6 viable cells/mL density. On day of transfection, the viable cell density and percent viability is determined. Cells were transfected at a density of 6×10^6 viable cells/mL. For optimal transfection, sterile Heavy and Light Chain plasmid DNA at ≥ 1 mg/mL concentration in TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0) were used. ExpiCHO-S™ cells were transfected following manufacturer's Max Titer Transfection protocol. All amounts and volumes shown below are per mL of final transfected culture volume. Briefly, plasmid DNA were diluted with 0.04 mL cold OptiPRO™ medium at the following ratio: 0.125 µg Heavy Chain DNA: 0.375 µg Light Chain DNA: 0.5 µg pAdvantage. 6.4 µL of ExpiFectamine™ CHO Transfection Reagent was diluted and mixed gently with 0.04 mL cold OptiPRO™ medium and incubated for 1 min. The diluted ExpiFectamine™ CHO Reagent was added to the diluted

DNA, mixed gently and the ExpiFectamine™ CHO/plasmid DNA complexes were incubated at room temperature for 5 minutes. Post-incubation, the complexes were added to the ExpiCHO-S™ cells in a shaker flask and incubated overnight in a 37°C, 7% CO₂ shaker incubator. On Day 1 post-transfection, 6 µL ExpiFectamine™ CHO Enhancer and 160 µL ExpiCHO™ Feed were added and the flask was transferred to a 32°C, 7% CO₂ shaker incubator. Culture was harvested on Day 7 post-transfection, centrifuged at 850 x g for 15 mins and clarified through a 0.2 µm Acropak 1500 filter capsule (Pall).

EXAMPLE 2.5: PURIFICATION OF ANTI-PSMA ANTIBODIES

[00547] Antibodies were purified from the clarified supernatants using MABSELECT SURE Protein A columns equilibrated with 1X D-PBS, pH 7.2 prior. Unbound proteins were removed by washing extensively with 1X D-PBS, pH 7.2. Bound protein was eluted with 0.1 M Na-acetate, pH 3.5. Peak fractions were neutralized with 2.5 M Tris pH 7.2 and pooled. The neutralized fraction pools were either dialyzed into 1X dPBS for assays and biophysical characterization or utilized for bispecific DuoBody assembly. The protein concentration for each elution pool was determined by measuring absorbance at OD280 nm and calculated using absorbance extinction coefficient based on the amino acid sequence.

[00548] Sequences of representative PSMA antibodies are provided in Tables 4-12. Table 4 provides VH amino acid sequences, VL sequences, heavy chain sequences, and light chain sequences of various PSMA antibodies. VH CDR and VL CDR sequences are provided in Table 5 (Kabat), Table 6 (Chothia), Table 7 (AbM), Table 8 (Contact), and Table 9 (IMGT). Numbers beneath each sequence in the Tables denote corresponding SEQ ID NOs.

[00549] C-terminal Lysine 477 (CTL) can be added to the antibody heavy chain for optimized expression. The anti-PSMA antibodies PSMB895, PSMB896, PSMB897, and PSMB898 are identical to PSMB946, PSMB947, PSMB948, and PSMB949, respectively with the exception of the addition of a C-terminal lysine (CTL) in the antibody heavy chain (K477).

[00550] That is, PSMB895 and PSMB946 have the same VH amino acid sequences, the same VL amino acid sequences, the same VH CDR1-3 amino acid sequences, and the same VL CDR1-3 amino acid sequences. PSMB895 and PSMB946 also have the same light chain amino acid sequences, but their heavy chain amino acid sequences differ in that the heavy chain amino acid sequence of PSMB946 further comprises an additional Lys (K) amino acid

residue at the C-terminus of the PSMB895 heavy chain amino acid sequence. PSMB896 and PSMB947 have the same VH amino acid sequences, the same VL amino acid sequences, the same VH CDR1-3 amino acid sequences, and the same VL CDR1-3 amino acid sequences. PSMB896 and PSMB947 also have the same light chain amino acid sequences, but their heavy chain amino acid sequences differ in that the heavy chain amino acid sequence of PSMB947 further comprises an additional Lys (K) amino acid residue at the C-terminus of the PSMB896 heavy chain amino acid sequence. PSMB897 and PSMB948 have the same VH amino acid sequences, the same VL amino acid sequences, the same VH CDR1-3 amino acid sequences, and the same VL CDR1-3 amino acid sequences. PSMB897 and PSMB948 also have the same light chain amino acid sequences, but their heavy chain amino acid sequences differ in that the heavy chain amino acid sequence of PSMB948 further comprises an additional Lys (K) amino acid residue at the C-terminus of the PSMB897 heavy chain amino acid sequence. PSMB898 and PSMB949 have the same VH amino acid sequences, the same VL amino acid sequences, the same VH CDR1-3 amino acid sequences, and the same VL CDR1-3 amino acid sequences. PSMB898 and PSMB949 also have the same light chain amino acid sequences, but their heavy chain amino acid sequences differ in that the heavy chain amino acid sequence of PSMB949 further comprises an additional Lys (K) amino acid residue at the C-terminus of the PSMB898 heavy chain amino acid sequence. The presence of the C-terminal Lysine can improve expression and manufacturing. C-terminal lysine (CTL) can be removed from the Ig constant region by endogenous circulating carboxypeptidases in the blood stream (Cai *et al.*, (2011) *Biotechnol Bioeng* 108:404-412). During manufacturing, CTL removal can be controlled to less than the maximum level by control of concentration of extracellular Zn^{2+} , EDTA or EDTA – Fe^{3+} as described in U.S. Patent Publ. No. US20140273092.

Table 4. PSMA Antibody VH, VL, HC and LC Amino Acid Sequences

#	Protein Name (Target)	VH Name	VL Name	VH AA sequence	VL AA sequence	Heavy Chain AA sequence	Light Chain AA sequence
1	PSMB889 (PSMA)	PBD0001013 12	PBD0001013 13	EVQLVESGGGL VKPGSRLRSC AAGFTFSRYN MNVVRQAPGKG LEWVSSINSNS RYIYADSVKG RFTISRDSAKN SLYLQMSLRA EDTAVYYCAKT MGDYIYYGMD VMGQGTITVTS S	QSVLTQPPSV SGAPGQRTI SCTGSSFNIG AGYDVHWYQQ VPGTVPKLLI YDMSNRPSGV PDRFSGSKG TSASLAIITGL QAEDETYYIC QSYDSSLSGV VFGGGTKLTV L	EVQLVESGGGLVKPGGSLRSLSCAASGFTFS RYNNMVRQAPGKGLEWSSINSNSRYIY ADSVKGRFTISRDAKNSLYLQMSLRAED TAVYCAKTMGDYIYYGMDVMGQGTITTV SSASTKGPSVFPLAPSSKSTSGGTAALGCL VKDYFEPEPTVSNVNSGALTSGVHFFPAVLQ SSGLYSLSSVWTVPSSSLGTQTYICNVNHK PSNTKVDKVKVEPKCDKTHCTCPCPAPEAA GGPSVFLFPPKPKDTLMI SRTPEVTCVVVS VSHEDPEVKENWYVDGVEVHNAKTKPREEQ YNSTYRVSVLTILHQQDWLNGKEYCKVSN KALPAPIEKTI SKAGQGPPEPQVYTLPPSR EEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTTPPVLDSDGSEFLYSLRTVDKS RWQOGNVFSCSVMEALHNHYTQKSLSLSP G	QSVLTQPPSVSVGAPGQR VTISCTGSSFNIGAGYD VHWYQQVPGTVPKLLIY DMSNRPSGVDPDRFSGSK SGTSASLAIITGLQAEDE TVYYCQSYDSSLSGVF GGGKLTIVLGQPKAAPS VTLEFPSSSEELQANKAT LVCLI SDFYPGAVTVAM KADSSPVKAGVEITTPS KQSNKYAASSYLSLTP EQWKSHRYSYCQVTHEG STVEKTVAPTECS
2	PSMB890 (PSMA)	PBD0001013 12	PBD0001013 14	EVQLVESGGGL VKPGSRLRSC AAGFTFSRYN MNVVRQAPGKG LEWVSSINSNS RYIYADSVKG RFTISRDSAKN SLYLQMSLRA EDTAVYYCAKT MGDYIYYGMD VMGQGTITVTS S	SSELTQPPSV SGAPGQRTI SCAGSLSNIG AGYDVHWYQQ LPGTAPKLLI YGNINRLSGV PERFSGRSK TSASLAIITGL QAEEDGADYYC QSYDSSLSGY VFGTGTKVTI L	EVQLVESGGGLVKPGGSLRSLSCAASGFTFS RYNNMVRQAPGKGLEWSSINSNSRYIY ADSVKGRFTISRDAKNSLYLQMSLRAED TAVYCAKTMGDYIYYGMDVMGQGTITTV SSASTKGPSVFPLAPSSKSTSGGTAALGCL VKDYFEPEPTVSNVNSGALTSGVHFFPAVLQ SSGLYSLSSVWTVPSSSLGTQTYICNVNHK PSNTKVDKVKVEPKCDKTHCTCPCPAPEAA GGPSVFLFPPKPKDTLMI SRTPEVTCVVVS VSHEDPEVKENWYVDGVEVHNAKTKPREEQ YNSTYRVSVLTILHQQDWLNGKEYCKVSN KALPAPIEKTI SKAGQGPPEPQVYTLPPSR EEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTTPPVLDSDGSEFLYSLRTVDKS RWQOGNVFSCSVMEALHNHYTQKSLSLSP G	SSELTQPPSVSVGAPGQR VTISCAAGSLSNIGAGYD VHWYQQVPGTAPKLLIY GNINRLSGVPERFSGSK SGTSASLAIITGLQAEDE ADYICQSYDSSLSGVF GTGTKVTIVLGQPKAAPS VTLEFPSSSEELQANKAT LVCLI SDFYPGAVTVAM KADSSPVKAGVEITTPS KQSNKYAASSYLSLTP EQWKSHRYSYCQVTHEG STVEKTVAPTECS

3	PSMB891 (PSMA)	PBD0001013 16	PBD0001013 15	31	66	33	68
				EVQLVESGGGV VQGRSLRLSC AASGFTFTYG MHWVRQAPGKG LEWVAVSFEDE SNKYADSVK RFTISRDNKSN TLYLQMSLRA EDTAVYYCARA LRDGNNDYFN GMDVWGQGTIV TVSS	QSVLTQPPSA SGTFPGQVTI SCSGSSSNIG SNTVNWFOQL PGTAPKLLIY SDNQRPSGVP DRFSGSKSGT SASLAI SGLQ SEDEADYYCA AWDDSLNGV FGTGKVTVL	EVQLVESGGGVQGRSLRLSCAASGFTFI TYGHWVRQAPGKGLEWAVVSFDESNNKY ADSVKGRFTISRDNKNTLYLQMSLRAED TAVYCARALRDGNNDYFNQMDVWGQGT VTVSSASTKGPSVFLAPSSKSTSGGTAAL GCLVKDYFPEPTVYSWNSGALTSGVHTFPA VLQSSGLYSLSSVTVPSSSLGTQTYICNV NHKPSNTKVDKKVPPKSCDKTHTCPPCPAP EAAGPSFELPPPKDITLMISRTPEVTCV VVSVSHEDPEVKENWYVDGVEVHNATKPR EEQYNSTYRVVSVLTVLTQDMLNGKEYCK VSNKALPAPIEKTISKAKGQPREPOVYITLP PSREMTKNQVSLTCLVKGFYPSDIAVEWE SNGQPENNYKTTTPVLDSDGSFFLYSRLTV DKSRWQQGNVFSCSVNHEALHNHYTQKSLS LSPG	QSVLTQPPSASGTFGQ VTISCSGSSSNIGSNTV NWFQQLPGTAPKLLIYS DNQRPSSGVDRFSGSKS GTSSASLAISGLQSEDEA DYCAAWDDSLNGVYFVG TGKVTVLQGPKAAPSV TLFPPSSSEELQANKATL VCLISDFYPGAVTVAMK ADSSPVKAGVETTTFSK QSNKYAASSYLSLTPE QWKSHRYSYSCQVTHEGS TVEKTVAPTECS
4	PSMB892 (PSMA)	PBD0001013 16	PBD0001013 17	99	100	101	102
				EVQLVESGGGV VQGRSLRLSC AASGFTFTYG MHWVRQAPGKG LEWVAVSFEDE SNKYADSVK RFTISRDNKSN TLYLQMSLRA EDTAVYYCARA LRDGNNDYFN GMDVWGQGTIV TVSS	QSVLTQPPSV SGAPGQRTI SCTGSSSNIG ADYDVHWYQH LPGTAPKLLI YGNSSNRPSGV PDRFSGSKSG TSASLAITGL QAEDETYYC QSYDSSLSGW VFGGGTKLTV L	EVQLVESGGGVQGRSLRLSCAASGFTFI TYGHWVRQAPGKGLEWAVVSFDESNNKY ADSVKGRFTISRDNKNTLYLQMSLRAED TAVYCARALRDGNNDYFNQMDVWGQGT VTVSSASTKGPSVFLAPSSKSTSGGTAAL GCLVKDYFPEPTVYSWNSGALTSGVHTFPA VLQSSGLYSLSSVTVPSSSLGTQTYICNV NHKPSNTKVDKKVPPKSCDKTHTCPPCPAP EAAGPSFELPPPKDITLMISRTPEVTCV VVSVSHEDPEVKENWYVDGVEVHNATKPR EEQYNSTYRVVSVLTVLTQDMLNGKEYCK VSNKALPAPIEKTISKAKGQPREPOVYITLP PSREMTKNQVSLTCLVKGFYPSDIAVEWE SNGQPENNYKTTTPVLDSDGSFFLYSRLTV DKSRWQQGNVFSCSVNHEALHNHYTQKSLS LSPG	QSVLTQPPSVSGAPGQR VTISCTGSSSNIGADYD VHWYQHLPGTAPKLLIY GNSNRPSGVDRFSGSK SGTSASLAITGLQAEDE TDYYCQSYDSSLSGWVE GGGTKLTVLQGPKAAPS VTLPFPSSSEELQANKAT LVCLISDFYPGAVTVAM KADSSPVKAGVETTTFS KQSNKYAASSYLSLTPE EQWKSHRYSYSCQVTHEG STVEKTVAPTECS
5	PSMB893 (PSMA)	PBD0001013 18	PBD0001013 15	99	134	101	136
				EVQLVESGGGV VQGRSLRLSC VASGFTFSSYG IHWVRQAPGKG	QSVLTQPPSA SGTFPGQVTI SCSGSSSNIG SNTVNWFOQL	QVQLVESGGGVQGRSLRLSCVAGSTFIS SYGHWVRQAPGKGLEWAVIWDGNNKY ADSVKGRFTISRDNKNTLYLQMSLRAED TAVYSVRGVGPTSYIYNYGMDVWGQGTIV	QSVLTQPPSASGTFGQ VTISCSGSSSNIGSNTV NWFQQLPGTAPKLLIYS DNQRPSSGVDRFSGSKS

				LEWVAVIWDG SNKYADSVKG RFTISRDN SKN TLYLQMSLRA EDTAVIYSVRG VGPTSYYNIG MDVWGQGTTVT VSS			PGTAPKLLIY SDNQRPSPV DRFSGSKGT SASLAISGLQ SEDEADYYCA AWDDSLNGV FGTGKTVTL		TVSSASTKGPSVFEPLAPSSKSTSGGTAALG CLVKDYFPEPVTVSWNSGALTSVHTFPV LQSSGLYSLSSVTVVPSSSLGTQYICNVN HKPSNTKVDKKEPKSCDKHTTCPCPAPE AAGGPSVFLFPKPKDTLMI SRTPEVTCVV VSVSHEDPEVKFNWYDGVVEHNATKPRE EQYNSTYRVVSVLTVLHQDWLNGKEYKCKV SNKALPAPIEKTISKAKGQPREPVYTLPP SREEMTKNQVSLTCLVKGFYPSDIAVEWES NGQPENNYKTTPPVLDSDGSFFLYSRLTVD KSRWQQGNVFSCSVMEALHNHYTQKSLSL SPG	167	100	169	102
6	PSMB894 (PSMA)	PBD0001013 18	PBD0001013 17	QVQLVESGGGV VQPGSRLRSC VASGFTFSYG IHWVRQAPGKG LEWVAVIWDG SNKYADSVKG RFTISRDN SKN TLYLQMSLRA EDTAVIYSVRG VGPTSYYNIG MDVWGQGTTVT VSS		QSVLTQPPSV SGAPGQRTI SCTGSSSNIG ADYDVHWYQH LPGTAPKLLI YGNRNPSGV PDRFSGSKG TSASLAITGL QAEDETYYC QSYDSSLSGW VFGGGTKLTV L	QSVLTQPPSVSGAPGQR VTISCTGSSSNI GADYD VHWYQHLPGTAPKLLIY GNSNRPSGVPEDRFSGSK SGTSASLAITGLQAEDE TDYICQSDYDSSLSGWVF GGGTKLTVLGQPKAAP VTLEPPSSEELQANKAT LVCLISDFYFGAVTVAW KADSPVKAGVETITPS KQSNKYAASSYLSLTP EQMKSHRSYSCQVTHEG STVEKTVAPTECS	169	134	136			
7	PSMB895 (PSMA)	PBD0001013 20	PBD0001013 19	EVQLVESGGGL VQPGGSLRLSC AASGFTFSYA MSWVRQAPGKG LEWVSAISGGI GSTYYADSVKG RFTISRDN SKN TLYLQMSLRA EDTAVIYCAKD		QSVLTQPPSV SAAPGQRTI SCSGSSSNIG NNYVSWYQQL PGTAPKLLIY DNNKRP SGIP DRFSGSKGT SATLGITGLQ TGDEADYYCG	EVQLVESGGGLVQPGGSLRLSCAASGFTFS SYAMSWVRQAPGKGLEWVSAISGGIGSTYY ADSVKGRFTISRDN SKNTLWLQMSLPAED TAVYCAKDGVGATPYFDYWGQGLITVTS SASTKGPSVFEPLAPSSKSTSGGTAALGCLV KDYFPEPVTVSWNSGALTSVHTFPFVAVLQS SGLYSLSSVTVVPSSSLGTQYIICNVNHNKP SNTKVDKKEPKSCDKHTTCPCPAPEAAG GPSVLEFLPPKPKDTLMI SRTPEVTCVVSV	169	134	136			

				GVGATPYFYDY WGQGTLTVTSS	TWDSLSAYV FGTGKTVTL	SHEDPEVKFNWYVDGVEVHNAKTKPREEQY NSTYRVVSVLTVLHQDWLNGKEYKCKVSNK ALPAPIEKTIISAKAGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTPEVLDSDGSEFFLYSRLTVDKSR WQGNVFSCSVMHREALHNHYTOKLSLSLSPG	ADSSPVKAGVETTPSK QSNNKYAASSYLSLTPE QWKSHRYSYSCQVTHEGS TVEKTVAPTECS
8	PSMB896 (PSMA)	PBD0001013 20	PBD0001013 21	EVQLVESGGGL VQPGGSLRLSC AASGTFSSYA MSWVRQAPGKG LEWWSAISGGI GSTYYADSVKG RFTISRDN SKN TLWLQMSLRA EDTAVVYCAKD GVGATPYFYDY WGQGTLTVTSS	QSVLTQPPSV SAAPGQKVTI SCSGSSNIG INYSWYQQL PGTAPKLLIY DNNKRPSCGIP DRFSGSKSGT SATLIGITGLQ TWDSLSAVV FGGGTKLTVL	EVQLVESGGGLVQPGGSLRLSCAASGTFPS SYAMSWVRQAPGKGLEWWSAISGGIGSTYY ADSVKGRFTISRDN SKNTLWLQMSLRAED TAVYCAKDGVGATPYFYFDYWGQGTLTVTS SASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYFPEPVTVSWNSGALTSGVHTFPAVLQSL SGLYSLSVATVPS SLSGTQTYICNVAHKPE SNTKVDKKVEPKSKDKTHTCPPCPAPEAAAG GFSVFLRPPKPKDITMLSRTPETVTCVVVSV SHEDPEVKFNWYVDGVEVHNAKTKPREEQY NSTYRVVSVLTVLHQDWLNGKEYKCKVSNK ALPAPIEKTIISAKAGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTPEVLDSDGSEFFLYSRLTVDKSR WQGNVFSCSVMHREALHNHYTOKLSLSLSPG	QSVLTQPPSVSAAPGQK VTISCSGSSNIGINYY SWYQQLPGTAPKLLIYD NNKRPSCGIPDRFSGSKS GT SATLIGITGLQTGDEA DYCGTWDSSLSAVVFG TGT KVTVLGQPKRAAPS TLFPPSSEELQANKATL VCLISDFYPGAVTVAWK ADSSPVKAGVETTPSK QSNNKYAASSYLSLTPE QWKSHRYSYSCQVTHEGS TVEKTVAPTECS
9	PSMB897 (PSMA)	PBD0001013 22	PBD0001013 19	EVQLVESGGGL VQPGGSLRLSC AASGTFSSYA MSWVRQAPGKG LEWWSAISGGI GSTYYADSVKG RFTISRDN SKN TLYLQMSLRA EDSAVYCAKD GVGATPYFYDY WGQGTLTVTSS	QSVLTQPPSV SAAPGQKVTI SCSGSSNIG NNYSWYQQL PGTAPKLLIY DNNKRPSCGIP DRFSGSKSGT SATLIGITGLQ TWDSLSAYV FGTGKTVTL	EVQLVESGGGLVQPGGSLRLSCAASGTFPS SYAMSWVRQAPGKGLEWWSAISGSGSGSTYY ADSVKGRFTISRDN SKNTLYLQMSLRAED SAVYCAKDGVGATPYFYFDYWGQGTLTVTS SASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYFPEPVTVSWNSGALTSGVHTFPAVLQSL SGLYSLSVATVPS SLSGTQTYICNVAHKPE SNTKVDKKVEPKSKDKTHTCPPCPAPEAAAG GFSVFLRPPKPKDITMLSRTPETVTCVVVSV SHEDPEVKFNWYVDGVEVHNAKTKPREEQY NSTYRVVSVLTVLHQDWLNGKEYKCKVSNK ALPAPIEKTIISAKAGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTPEVLDSDGSEFFLYSRLTVDKSR WQGNVFSCSVMHREALHNHYTOKLSLSLSPG	QSVLTQPPSVSAAPGQK VTISCSGSSNIGNNYY SWYQQLPGTAPKLLIYD NNKRPSCGIPDRFSGSKS GT SATLIGITGLQTGDEA DYCGTWDSSLSAVVFG TGT KVTVLGQPKRAAPS TLFPPSSEELQANKATL VCLISDFYPGAVTVAWK ADSSPVKAGVETTPSK QSNNKYAASSYLSLTPE QWKSHRYSYSCQVTHEGS TVEKTVAPTECS

[illegible]

				SLHLHMNSLKA EDTAMVFCARS YSGSYDAFDW GQGTMTVSS	DFTLTISRLE PEDFAVYQC QYGVSPWTFG QGTKEIK	NTKVDKKVEPKSCDKTHTEPCPEAAGG PSVFLPPPKKDTLMISRTPEVTCVVSVS HEDPEKFNWYDGVVHNKTRPEEQYN STYRVSVLTVLHQDLNKEKYCKVSKA LPAIEKTIKAKGQPREQVYITLPPSREE MTKNQVLSLCAVKGFPYSDIAVWESNGQP ENNYKTTTPVLDSDGSFFLVSKLTVDKSRW QQGNVFSCSVMHREALHNRTQKSLSPGK	PPDEQLKSGTASVVCL LNNFYPREAKVQMKVDN ALQSGNSQESVTEQDSK DSTYSLSTLTLKADY EKHKVIACEVTHQGLSS PVTKSFNRGEC
13	PSME896 -G100A (PSMA)	NA	NA	405 EVQLVESGGGL VQPGGSLRLSC AAGGTFSSYA MSWVRQAPGKG LEWVAISGGI GSTYADSVKG RFTISRDNKSN TLWLQMSLRA EDTAVYYCAKD AVGATPYIFDY WGQGTLLVTVSS	406 QSVLTQPPSV SAAPGQKVTI SCSGSSNIG INYSWYQQL PGTAPKLLIY DNNKRPSCGP DRFSGSKSGT SATLIGITGLQ TGDEADYYCG TWDSSLSAVV FGGGTKLTVL	407 EVQLVESGGGLVQPGGSLRLSCAASGFTFS SYAMSWVRQAPGKGLEWVSALSGGIGSTYY ADSVKGRFTISRDNKNTLWLQMSLRAED TAVYYCAKDAVGATPYFDYWGQGTLLTVTS SASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYEEPEPTVSWNSGALTSGVHTFPAVLQV SGLYSLSSVTVTPSSSLGTQTYICNVNHKPF SNTKVDKKVPEPKSCDKTHTEPCPEAAG GPSVLEFPKPKDTLMISRTPEVTCVVSV SHEDPEVKFNWYDGVVHNKTRPEEQY NSTYRWVSIVTLVLDWLNKGEYCKVSKN ALPAIEKTIKAKGQPREQVYITLPPSRE EMTKNQVLSLCAVKGFPYSDIAVWESNGQ PENNYKTTTPVLDSDGSFFLVSKLTVDKSR WQQGNVFSCSVMHREALHNRTQKSLSPG K	408 QSVLTQPPSVSAAPGQK VTISCSGSSNIGINYV SWYQQLPFGTAPKLLIYD NNKRPSCGIDRFSGSKS GTSATLIGITGLQGTDEA DYCGTWDSSLSAVVFG GGTKLTVLQGPKAAPSV TLFPPSSEELQANKATL VCLISDFYFGAVTVAWK ADSPVKAGVETTTPSK QSNNKYAASYSLSLTPE QWKSHRYSYSCQVTHEGS TVEKTVAPTECS
14	PSMA_P7 2_A10- HC-G54E (PSMA)	NA	NA	439 QVQLVESGGGV VQPGSRLRSC AAGGTFSSYN MNWVRQAPGKG LEWVAIIYYDE SNKYADSVKG RFTISRDISKN TLYLQMSLRA EDTAVYYCARE RGRDYYGMDVW GQGTMTVTVSS	270 SYELMQPPSV SVSPGQTARI TCSGDALPKQ YATWYQQKPG QAPFLVIYKD SERPSGIPVR FSGSSSGTIV TLTITGVQAE DEADYYCOSA DSSGTYVEFT GKVTIVL	441 NA	272 NA
				473 GQGTMTVTVSS	474	475	476

15	PSMA_P7 2_D01- HC-D95E (PSMA)	NA	NA	EVQLVESGGDL VQPGSRLRSC AASGTFNNYN MHWVRQAPGKG LEWVSHISTSS SNKYADSVKG RFTISRDIKRN SNYLOMNSLRD EDTAVYYCARE GVGADYGDYYY YGMDFWGQGT VTVSS	QSVLTQPASV SGSPGQSITI SCTGTSSDVG GNYVSWYQQ HFGKAPKLMI YEVSNRPSGV SNRFSGRSKG NTASLTISGL QAEDEADYYC SSYTSSYTV FGTGKLTVL	NA	NA	510
16	PSMA_P7 2_F01 and PSMA_P7 5_F01 (PSMA)	NA	NA	QVQLQESGGGV VQPGSRLRSC AASGTFSTYG MHWVRQAPGKG LEWVAFISYDG SNKYADSVKG RFTISRDNKSH TLYLOMNSLRA EDTAVYYCAGR DNLRFLEWEMD VWGQGTITVTS S	EIVLTQSEPT LSVSPGERAT LSCRASQSVR SNLAWYQQKP GOAFRLIYG ASTRATGIPA RFSGSGSGTE FTLTISLSQS EDFAVYYCHQ YNDWPPYTFG QGTEKLEIK	NA	NA	509
17	PSMA_P7 2_F07 (PSMA)	NA	NA	EVQLVESGGGV VQPGSRLRSC AASGTFSSYG MHWVRQAPGKG LEWVAVTSYDG SNKYADSVKG RFTISRDISKN TLYLOMSSLRA EDTAVYYCARD PYSSSWNGAFD IWGPGMTVTS S	SYELTQPPSV SVAPGQTARI TCGNNIGSK SVHWYQQKEG QAEVLAVYDD SDRPSGIPER FSGNSGNTA TLTISRVEAG DEADYYCQVM DSSTDHVVFG GGTKLTVL	NA	NA	543
				541	542			544

18	PSMA_P7 2_E07 (PSMA)	NA	NA	EVQLVESGGGV VQPGSLRLSC AAGGTFITFG MHWVRQAPGKG LEWVAVVSEDE SNKYADSVKGG RFTISRDN SKN TLYLQMNSLRA EDTAVYYCARA LRDGNNDYFN GMDVWGQGT TVSS	99	QSVLTQPPSA SGHPGGQVTI SCSGSSNIG SNTVNWFOQL PGTAPKLLIY SDNQRPSCGP DRFSGSKSGT SASLAISGLQ SEDEADYYCA AWDDSLNGIV FGTGKTVTL	NA	NA	611	612
19	PSMA_P7 2_D01 (PSMA)	NA	NA	EVQLVESGGDL VQPGSLRLSC AAGGTFNNYN MHWVRQAPGKG LEWVSHISTSS SNKYADSVKGG RFTISRDIANK SMYLQMNSLRD EDTAVYYCARD GVGADYGDYY YGMVWGQGT TVSS	100	QSVLTQPPASV SGSPGQSITI SCTGTSSDVG GYNYVSWYQQ HPGKAPKIMI YEVSNRPSGV SNRPSGSKSG NTASLTISGL QAEDEADYYC SSYTSSTYV FGTGKTVTL	NA	NA	611	612
20	PSMA_P7 2_C01 (PSMA)	NA	NA	QVQLVESGGGE VQPGSLRLSC AAGGFSFGYG MHWVRQAPGKG LEWVAVMSYDG SNRFYVDSVRG RFTISRDN SKN TLYLQMNSLRP EDTAVYYCARD TVWGSHPDAPD IWGQGTIVTVS S	643	QSVLTQPPSV SVAPGQTARI TCGNNNSGSK SVHWYQQKPG QAPLVAVYDD SDRPSGIPER FSGNSGNTA TLTISRVEAG DEADYYCQVM DSSSDHGVFG GGTKLTVL	NA	NA	645	646
					677				679	680

21	PSMA_P7 2_A10 (PSMA)	NA	NA	QVQLVESGGGV VQPGSLRLSC AASGTFSSYN MHWVRQAPGKG LEWVAIIYYDG SNKYADSVKGG RFTLSRDISKN TLYLQMNLSRA EDTAVYYCARE RGRDYIGMDVW GGGTVTIVSS	711	SVELMQPESV SVSPGQTARI TCSGDALPKQ YAYWYQQKPG QAPFLVIYRD SERPSGIEVR FSGSSSGTIV TLTIIGVQAE DEADYYCOSA DSGGTYVFGT GTRKTVL	474	NA	NA	714
22	PSMA_P7 2_F02 and PSMA_P7 0_F02 (PSMA)	NA	NA	EVQLLESQGL VKPSETLSLT TVSGGSIISYY WSWIRQAPGKG LEWLGRIYSSG STNYPNLSKSR VTMSVDTSKNQ FSLKLSSTAA DTAVYYCAKVG VMPGAFDIWQG GTMVTIVSS	711	QSVLTQPPSA SGTFPGQRTI SCSGSSSNIG SNTVNWYQQL PGTAPKLLIY SSNQRPSPGV DRFSGSKSGT SASLAIISGLQ SEDEADYYCA AWDDSLNGW FSGGSKLTVL	474	NA	NA	714
23	PSMA_P7 2_G02 (PSMA)	NA	NA	EVQLVESGGGV VQPGSLRLSC AASGTFSSYG MHWVRQAPGKG LEWVAIIYYDG SNKYADSVKGG RFTLSRDNKN TLYLQMNLSRV EDTAVYYCARD RINGSRGYYG MDVWGQGTIVT VSS	745	QSAITQPASV SGSPGQSITI SCTGASSDVG GYNVSWYQQ HPGKAPKLM YEVSNRPSPGV SNRFSGSKSG NTASLTISGL QAEDEADYYC SSYTITSTLV FSGGSKLTVL	746	NA	NA	748
24	PSMA_P7 2_A11	NA	NA	QVQLQESGGDV VQPGSLRLSC	779	SVELTQPPSV SVAPGQTARI	780	NA	NA	782

	(PSMA)			AAGGFSFSGYG LHWVRQAPGRG LEWVTLISYDG SNKYADSVKVG RFTLSRDNSKN TLYLQMSLRRA EDTAVYYCART TVSDPYYGMD VWGQGTITVTS S	TCGNNIGSK SVHWYQQREG QAEVLVYDD SDRPSGIPER FSGTNSGNTA TLTISRAEAG DEADYYCQW DSSSDHVVFG GGTKLTVL			
				813	814	815	816	

Table 5. PSMA Antibody Kabat CDR Amino Acid Sequences

#	Protein Name	HC Kabat CDR1	HC Kabat CDR2	HC Kabat CDR3	LC Kabat CDR1	LC Kabat CDR2	LC Kabat CDR3
1	PSMB889	RYNMN	SINNSRYIYYADSVKG	TWGDYVYYYGMDV	TGSENLGAGYDVH	DNSNRPS	QSYDSSLGSAV
2	PSMB890	RYNMN	SINNSRYIYYADSVKG	TMGDYVYYYGMDV	AGLSNIGAGYDVH	GNIHPLS	QSYDSSLSSVY
3	PSMB891	TYGMH	VVSFDESNKYIYADSVKG	ALRDGNWDYFNGMDV	SGSSSNIGSNTVN	SDNQRP	AAMDDSLNGYV
4	PSMB892	TYGMH	VVSFDESNKYIYADSVKG	ALRDGNWDYFNGMDV	TGSSSNIGAGYDVH	GNSNRPS	QSYDSSLGSAV
5	PSMB893	SYGIH	VIWYDGSNKYIYADSVKG	GVGPTSYIYYNYGMDV	SGSSSNIGSNTVN	SDNQRP	AAMDDSLNGYV
6	PSMB894	SYGIH	VIWYDGSNKYIYADSVKG	GVGPTSYIYYNYGMDV	TGSSSNIGAGYDVH	GNSNRPS	QSYDSSLGSAV
7	PSMB895	SYAMS	AISGGIGSTYYADSVKG	DGVGATPYFFDY	SGSSSNIGNNYVS	DNNKRPS	GTWDSLSAVV
8	PSMB896	SYAMS	AISGGIGSTYYADSVKG	DGVGATPYFFDY	SGSSSNIGNNYVS	DNNKRPS	GTWDSLSAVV
9	PSMB897	SYAMS	AISGGIGSTYYADSVKG	DGVGATPYFFDY	SGSSSNIGNNYVS	DNNKRPS	GTWDSLSAVV
10	PSMB898	SYAMS	AISGGIGSTYYADSVKG	DGVGATPYFFDY	SGSSSNIGNNYVS	DNNKRPS	GTWDSLSAVV
11	PSMB899	SYAMS	AISGGYGAPYYADSVKG	DGVGATPYFFDD	SGSSSNIGNNYVS	DNNKRPS	GTWDSLSAVV
12	PSMHB49SC1133 011A11.1	FYSMN	SISSSGNVIYYADSVKG	SYSGSYDAEDF	RAQSVSSSFLA	GASSPAT	QQYGVSPWT
13	PSMB896-G100A	SYAMS	AISGGIGSTYYADSVKG	DAVGATPYFFDY	SGSSSNIGNNYVS	DNNKRPS	GTWDSLSAVV
14	PSMA_P72_A10- HC-G54E	SYNMN	IYYDESNKYIYADSVKG	ERGRDYVGMDV	SGDALPKQYAY	KDSRPS	QSADSSGTYV
15	PSMA_P72_D01- HC-D95E	NYNMN	HISTSSNKYIYADSVKG	EGVGADYGDYIYYGMD V	TGTSSDVGGYNYVS	EVSNRPS	SSYTSSTYTV
		477	478	479	480	481	482

16	PSMA_P72_F01 and PSMA_P75_F01	TYGMH	FTSYDGSNKYYADSVKG	RDLRFLEWFMV	RASQSVRSNLA	GASTRAT	HQYNDWPPYT
		69	512	513	514	515	516
17	PSMA_P72_F07	SYGMN	VTSYDGSNKYYADSVKG	DPYSSSWNGAFDI	GGNNIGSKSVH	DDSDRPS	QVWDSSTDHW
		545	546	547	548	549	550
18	PSMA_P72_E07	TYGMH	VVSFEDSNKYYADSVKG	ALRDGNNWDYFNGMDV	SGSSSNIGSNTVN	SDNQRPS	AAWDDSLNGV
		69	70	71	72	73	74
19	PSMA_P72_D01	NNNMN	HISTSSSNKYYADSVKG	DGVGADYGDYYYGMD V	TGTSSDVGGYNYVS	EVSNRPS	SSYTSSYTYV
		477	478	615	480	481	482
20	PSMA_P72_C01	GYGMH	VMSYDGSNRFYVDSVRG	DTVMGSHPDAFDI	GGNSGGSKSVH	DDSDRPS	QVWDSSSDHV
		647	648	649	650	549	652
21	PSMA_P72_A10	SYNMN	IIFYDGSNKYYADSVKG	ERGRDYGMV	SGDALPKQYAY	KDSERPS	QSADSSGTYY
		443	682	445	446	447	448
22	PSMA_P72_F02 and PSMA_P70_F02	SYYWS	RIYSSGSTNYNPSLKS	VGVWPGAFDI	SGSSSNIGSNTVN	SSNQRPS	AAWDDSLNGV
		715	716	717	72	719	720
23	PSMA_P72_G02	GYGMH	VTSYDGSNKYYADSVKG	DRWGRGYYYGMDV	TGASSDVGGYNYVS	EVSNRPS	SSYTITSTIV
		647	750	751	752	481	754
24	PSMA_P72_A11	GYGLH	LISYDGSNKYYADSVKG	TTVSDPFYYGMDV	GGNNIGSKSVH	DDSDRPS	QVWDSSSDHV
		783	784	785	548	549	788

Table 6. PSMA Antibody Chothia CDR Amino Acid Sequences

#	Protein Name	HC Chothia CDR1	HC Chothia CDR2	HC Chothia CDR3	LC Chothia CDR1	LC Chothia CDR2	LC Chothia CDR3
1	PSMB889	GTFESRY	NSNSRY	MGDYVYYVGM	SSENLGAGYD	DNS	YDSSLGGV
2	PSMB890	GTFESRY	NSNSRY	MGDYVYYVGM	SLSNIGAGYD	GNI	YDSSLSSY
3	PSMB891	GTFEITY	SFDES	LRDGNNDYFNGMD	SSSNIGSNT	SDN	WDDSLNGY
4	PSMB892	GTFEITY	SFDES	LRDGNNDYFNGMD	SSSNIGADYD	GNS	YDSSLGGV
5	PSMB893	GTFESSY	WYDGSN	VGPTSYVYVGM	SSSNIGSNT	SDN	WDDSLNGY
6	PSMB894	GTFESSY	WYDGSN	VGPTSYVYVGM	SSSNIGADYD	GNS	YDSSLGGV
7	PSMB895	GTFESSY	SGGIGS	GVGATPYVFD	SSSNIGNNY	DNN	WDDSLSAY
8	PSMB896	GTFESSY	SGGIGS	GVGATPYVFD	SSSNIGNNY	DNN	WDDSLSAY
9	PSMB897	GTFESSY	SGGIGS	GVGATPYVFD	SSSNIGNNY	DNN	WDDSLSAY
10	PSMB898	GTFESSY	SGGIGS	GVGATPYVFD	SSSNIGNNY	DNN	WDDSLSAY
11	PSMB899	GTFESSY	SGGYGA	GVGATPYVFD	SSSNIGNNY	DNN	WDDSLSAY
12	PSMB495C1133_011A11.1	GTFESFY	SSSGNY	SYSGSYDAFD	SQSVSSSF	GAS	YGVSPW
13	PSMB896-G100A	GTFESSY	SGGIGS	DAVGATPYVFD	SSSNIGNNY	DNN	WDDSLSAY
14	PSMA_P72_A10-HC-G54E	GTFESSY	YDESN	ERGRDYVGM	DALPKQY	KDS	ADSSGY
15	PSMA_P72_D01-HC-D95E	GTFENNY	STSSN	EGVGADYGYVGM	TSSDVGGNY	EVS	YTSSY
16	PSMA_P72_F01 and PSMA_P75_F01	GTFESTY	SYDGSN	RDLRFLEWFM	SQSVRSN	GAS	YNDWPPY

17	PSMA_P72_F07	GTFESSY	517	518	DPYSSSWNGAFD	519	NNIGSKS	520	DDS	385	WDSSTDHV	522
			143	518		553		554		555		556
18	PSMA_P72_E07	GTFITY		SFDES	ALRDGNNDYFNGMD		SSSNIQSNT		SDN		WDDSLNGY	
			75	76		587		78		79		80
19	PSMA_P72_D01	GTFENNY		STSSN	DGVGADYGDYYYGMD		TSSDVGGYNY		EVS		YTSSYTY	
			483	484		621		486		487		488
20	PSMA_P72_C01	GFSESGY		SYDGSN	DTVMGSHPDADF		NNSGSKS		DDS		WDSSSDHG	
			653	518		655		656		555		658
21	PSMA_P72_A10	GTFESSY		YDGSN	ERGRDYGMD		DALPKQY		KDS		ADSSGTY	
			143	688		451		452		453		454
22	PSMA_P72_F02 and PSMA_P70_F02	GGSIISY		YSSGS	VGVWPGAFD		SSSNIQSNT		SSN		WDDSLNGV	
			721	722		723		78		725		726
23	PSMA_P72_G02	GFSESGY		SYDGSN	DRWGSRGYYYGMD		ASSDVGGYNY		EVS		YTITSTL	
			653	518		757		758		487		760
24	PSMA_P72_A11	GFSESGY		SYDGSN	TTVSDPYYYGMD		NNIGSKS		DDS		WDSSTDHV	
			653	518		791		554		555		794

Table 7. PSMA Antibody AbM CDR Amino Acid Sequences

#	Protein Name	HC AbM CDR1	HC AbM CDR2	HC AbM CDR3	LC AbM CDR1	LC AbM CDR2	LC AbM CDR3
1	PSMB889	GFTFSRYNMN 13	SINNSRYTY 14	TMGDYYYYYGMVDV 3	TGSSFNIGAGYDVH 16	DNSNRPS 5	QSYDSSLGGVV 6
2	PSMB890	GFTFSRYNMN 13	SINNSRYTY 14	TMGDYYYYYGMVDV 3	AGSLNIGAGYDVH 38	GNINRLS 39	QSYDSSLSSYV 40
3	PSMB891	GFTFTITYGMH 81	VSPFDESNEY 82	ALRDGNNDYENGMDV 71	SGSSSNIGSNTVN 72	SDNQRP 73	AAWDDSLNGYV 74
4	PSMB892	GFTFTITYGMH 81	VSPFDESNEY 82	ALRDGNNDYENGMDV 71	TGSSSNIGAGYDVH 106	GNSNRPS 107	QSYDSSLGGVV 108
5	PSMB893	GFTFSSYGIH 149	VIWYDGSNKY 150	GVGFTSYYYNYGMVDV 139	SGSSSNIGSNTVN 72	SDNQRP 73	AAWDDSLNGYV 74
6	PSMB894	GFTFSSYGIH 149	VIWYDGSNKY 150	GVGFTSYYYNYGMVDV 139	TGSSSNIGAGYDVH 106	GNSNRPS 107	QSYDSSLGGVV 108
7	PSMB895	GFTFSSYAMS 217	AISGGIGSTY 218	DGVGATPYFYFDY 207	SGSSSNIGNNNYVS 208	DNNKRPS 209	GTWDSLSAYV 210
8	PSMB896	GFTFSSYAMS 217	AISGGIGSTY 218	DGVGATPYFYFDY 207	SGSSSNIGNNNYVS 242	DNNKRPS 209	GTWDSLSAYV 244
9	PSMB897	GFTFSSYAMS 217	AISGGIGSTY 218	DGVGATPYFYFDY 207	SGSSSNIGNNNYVS 208	DNNKRPS 209	GTWDSLSAYV 210
10	PSMB898	GFTFSSYAMS 217	AISGGIGSTY 218	DGVGATPYFYFDY 207	SGSSSNIGNNNYVS 242	DNNKRPS 209	GTWDSLSAYV 244
11	PSMB899	GFTFSSYAMS 353	AISGGYGA 354	DGVGATPYFYFDY 343	SGSSSNIGNNNYVS 208	DNNKRPS 209	GTWDSLSAYV 210
12	PSMB498C113 3 011A11 1	GFTFSPYSMN 387	SISSSGNYTY 388	SYSGSYDAFDF 377	RASQSVSSSEFLA 378	GASSRAT 379	QQYGVSPFWT 380
13	PSMB896- G100A	GFTFSSYAMS 217	AISGGIGSTY 218	DAVGATPYFYFDY 411	SGSSSNIGNNNYVS 242	DNNKRPS 209	GTWDSLSAYV 244
14	PSMA_P72_A10 -HC-G54E	GFTFSSYNMN 455	IYYDESNEY 456	ERGRDYGMVDV 445	SGDALPKQYAY 446	KDSERPS 447	QSADSSGTYV 448
15	PSMA_P72_D01 -HC-D95E	GFTFNNNMN 489	HISTSSSNKY 490	EGVGADYDYYGYGMVDV 479	TGTSSDVGGYNYVS 480	EVSNRPS 481	SSYTSSYTYV 482

16	PSMA_P72_F01 and PSMA_P75_F01	GFTFSTYGMH	FISYDGSNKY	RDLRFLEWEMDV	RASQSVPSNLA	GASTRAT	HQYNDWFEYFYT
		523	524	513	514	515	516
17	PSMA_P72_F07	GFTFSSYGMN	VTSYDGSNKY	DPYSSWNAGAFDI	GGNNIGSKSVH	DDSDRPS	QVWDSSTDHVV
		557	558	547	548	549	550
18	PSMA_P72_E07	GFTFTTYGMH	VVSFDES NKY	ALRDGNNWDYFENGMDV	SGSSSNIGSNTVN	SDNQ RPS	AAWDDSLNGYV
		81	82	71	72	73	74
19	PSMA_P72_D01	GFTFNNYNNM	HISTSSSNKY	DGVGADYGDYYYGYMDV	TGTSSDVGGYNYVS	EVSNRPS	SSYTSSYTYV
		489	490	615	480	481	482
20	PSMA_P72_C01	GFSFSGYGMH	VMSYDGSNRF	DTVWGSHPDAFDI	GGNNSGSKSVH	DDSDRPS	QVWDSSTDHGV
		659	660	649	650	549	652
21	PSMA_P72_A10	GFTFSSYNNM	IYYDGSNKY	ERGRDYGYMDV	SGDALFKQYAY	KDSERPS	QSADSSGTYV
		455	694	445	446	447	448
22	PSMA_P72_F02 and PSMA_P70_F02	GGSIIISYYWS	RIYSSSGSTN	VGVWPGAFDI	SGSSSNIGSNTVN	SSNQ RPS	AAWDDSLNGYV
		727	728	717	72	719	720
23	PSMA_P72_G02	GFSFSGYGMH	VISYDGSNKY	DRIWGSRGYYYGYMDV	TGASSDVGGYNYVS	EVSNRPS	SSYTITSTLV
		659	762	751	752	481	754
24	PSMA_P72_A11	GFSFSGYGLH	LISYDGSNKY	TTVSDPYYYGYMDV	GGNNIGSKSVH	DDSDRPS	QVWDSSTDHVV
		795	796	785	548	549	788

Table 8. PSMA Antibody Contact CDR Amino Acid Sequences

#	Protein Name	HC Contact CDR1	HC Contact CDR2	HC Contact CDR3	LC Contact CDR1	LC Contact CDR2	LC Contact CDR3
1	PSMB889	SRYNM 19	WVSINSNSRIY 20	AKTMGDYIIYYGMD 21	IGAGYDVHWY 22	LLIYDMSNRP 23	QSYDSSLSGV 24
2	PSMB890	SRYNM 19	WVSINSNSRIY 20	AKTMGDYIIYYGMD 21	IGAGYDVHWY 56	LLIYGNINRL 57	QSYDSSLSY 58
3	PSMB891	ITYGMH	WVAVVSFEDESNKY 88	APALRDGNNDYFNGMD 89	IGSNTVNWY 90	LLIYSDNQRP 91	AAWDDSLNGY 92
4	PSMB892	ITYGMH	WVAVVSFEDESNKY 88	APALRDGNNDYFNGMD 89	IGAGYDVHWY 124	LLIYGNINRL 125	QSYDSSLSGV 126
5	PSMB893	SSYGIH 155	WVAVIWDGSKNY 156	VRGVGPTSYIIYYGMD 157	IGSNTVNWY 90	LLIYSDNQRP 91	AAWDDSLNGY 92
6	PSMB894	SSYGIH 155	WVAVIWDGSKNY 156	VRGVGPTSYIIYYGMD 157	IGAGYDVHWY 124	LLIYGNINRL 125	QSYDSSLSGV 126
7	PSMB895	SSYAMS 223	WVSAISGGIGSTY 224	AKDGVGATPYIFD 225	IGNNYVSWY 226	LLIYDNNKRP 227	GTWDDSSLSAY 228
8	PSMB896	SSYAMS 223	WVSAISGGIGSTY 224	AKDGVGATPYIFD 225	IGNNYVSWY 260	LLIYDNNKRP 261	GTWDDSSLSAV 262
9	PSMB897	SSYAMS 223	WVSAISGGIGSTY 292	AKDGVGATPYIFD 225	IGNNYVSWY 226	LLIYDNNKRP 227	GTWDDSSLSAY 228
10	PSMB898	SSYAMS 223	WVSAISGGIGSTY 292	AKDGVGATPYIFD 225	IGNNYVSWY 260	LLIYDNNKRP 261	GTWDDSSLSAV 262
11	PSMB899	SSYAMS 223	WVSAISGGIGAPY 360	AKDGVGATPYIFD 225	IGNNYVSWY 226	LLIYDNNKRP 263	GTWDDSSLSAY 228
12	PSMB498C1133_011 A11_1	SFYSMN	WVSSISSSGNYIY 394	AKSYSGSYDAFD 395	SSSFLAWY 396	LLISGASSRA 397	QQYGVSPW 398
13	PSMB896-G100A	SSYAMS 223	WVSAISGGIGSTY 224	AKDGVGATPYIFD 429	IGNNYVSWY 260	LLIYDNNKRP 261	GTWDDSSLSAV 262
14	PSMA_P72_A10-HC- G54E	SSYNM 461	WVAILIYDESNKY 462	ABERGRDYIYGM 463	KQYAWY 464	LVYIKDSERP 465	QSADSSGTY 466
15	PSMA_P72_D01-HC- D95E	NNYNM 495	WVSHISTSSSNKY 496	AREGVGADYIYIYGM 497	VGGNYVSWY 498	LMYEVSNRP 499	SSYTSSITY 500

16	PSMA_P72_F01 and PSMA_P75_F01	STYGMH	WVAFISYDGSNKY	AGRDNLRFLEWFMD	RSNLAWY	LLIYGASTRA	HQYNDWPPY
		529	530	531	532	533	534
17	PSMA_P72_F07	SSYGMN	WVAVTSYDGSNKY	ARDPYSSSWNGAFD	SKSVHWY	LVVYDDSDRP	QVWDSSSDHV
		563	564	565	566	567	568
18	PSMA_P72_E07	ITYGMH	WVAVVSFDES NKY	ARALRDGNNDYFENGMD	IGSNTVNNWF	LLIYSDNQRP	AAWDDSLNGY
		87	88	89	90	91	92
19	PSMA_P72_D01	NNYNMN	WVSHISTSSNKY	ARDGVGADYGDYYYGMD	VGGYNYVSWY	LMIEVSNRP	SSYTSSYTY
		495	496	633	498	499	500
20	PSMA_P72_C01	SGYGMH	WVAVMSYDGSNRF	ARDTVWGSHPDAFD	SKSVHWY	LVVYDDSDRP	QVWDSSSDHG
		665	666	667	566	567	670
21	PSMA_P72_A10	SSYNMN	WVALLIYDGSNKY	APERGRDYGGMD	KQYAYWY	LVLYKDSERP	QADSSGTY
		461	700	463	464	465	466
22	PSMA_P72_F02 and PSMA_P70_F02	ISYYWS	WIGRIYSSGSTN	AKVGVWEGAFD	IGSNTVNNWY	LLIYSSNQRP	AAWDDSLNGV
		733	734	735	736	737	738
23	PSMA_P72_G02	SGYGMH	WVAVISYDGSNKY	ARDRIWGSRGYYYGMD	VGGYNYVSWY	LMIEVSNRP	SSYTITSTL
		665	768	769	498	499	772
24	PSMA_P72_A11	SGYGLH	WVTLLSYDGSNKY	AKTTVSDPYYYGMD	SKSVHWY	LVVYDDSDRP	QVWDSSSDHV
		801	802	803	566	567	806

Table 9. PSMA Antibody IMGT CDR Amino Acid Sequences

#	Protein Name	HC IMGT CDR1	HC IMGT CDR2	HC IMGT CDR3	LC IMGT CDR1	LC IMGT CDR2	LC IMGT CDR3
1	PSMB889	GFTFSRYN 25	INSNSRYI 26	AKTMGDYYYYYGMVDV 27	SENLGAGVD 28	DNS 11	QSYDSSLGGVV 6
2	PSMB890	GFTFSRYN 25	INSNSRYI 26	AKTMGDYYYYYGMVDV 27	LSNIGAGVD 62	GNI 45	QSYDSSLSSVV 40
3	PSMB891	GFTFITYG	VSFDES NK	ARALRDGNNWDYFNGMDV	SSNIGSNT	SDN	AAWDDSLNGYV
4	PSMB892	GFTFITYG	VSFDES NK	ARALRDGNNWDYFNGMDV	SSNIGADVD	GNS	QSYDSSLGGVV
5	PSMB893	GFTFSSYG 161	IWYDGS NK 162	VRGVGPTSYYYNYGMVDV 163	SSNIGSNT	SDN	AAWDDSLNGYV 74
6	PSMB894	GFTFSSYG 161	IWYDGS NK 162	VRGVGPTSYYYNYGMVDV 163	SSNIGADVD	GNS	QSYDSSLGGVV 108
7	PSMB895	GFTFSSYA 229	ISGGIGST 230	AKDGVGATPYFYFDY 231	SSNIGNNY	DNN	GTWDSLSAYV 210
8	PSMB896	GFTFSSYA 229	ISGGIGST 230	AKDGVGATPYFYFDY 231	SSNIGINY	DNN	GTWDSLSAVV 244
9	PSMB897	GFTFSSYA 229	ISGGSGST 298	AKDGVGATPYFYFDY 231	SSNIGNNY	DNN	GTWDSLSAYV 210
10	PSMB898	GFTFSSYA 229	ISGGSGST 298	AKDGVGATPYFYFDY 231	SSNIGINY	DNN	GTWDSLSAVV 244
11	PSMB899	GFTFSSYA 365	ISGGYGAP 366	AKDGVGATPYFYFDY 367	SSNIGNNY	DNN	GTWDSLSAYV 210
12	PSMHB49SC1133 011A11_1	GFTFSFYs	ISSSGNYI	ARSYSGSYDAFDF	QSVSSSF	GAS	QQYGVSPFWT
13	PSMB896-G100A	GFTFSSYA 229	ISGGIGST 230	AKDAVGATPYFYFDY 435	SSNIGINY	DNN	GTWDSLSAVV 244
14	PSMA_P72_A10- HC-G54E	GFTFSSYN	IYDES NK	ARERGRDYGMVDV	ALPKQY	KDS	QSADSSGTYV
15	PSMA_P72_D01- HC-D95E	GFTFNNNY	ISTSSNK	AREGVGADYDYYNYGMVDV	SSDVGGYNY	EVS	SSYTSSTYTV 448
		501	502	503	504	487	482

16	PSMA_P72_F01 and PSMA_P75_F01	GFTFSTYG	ISYDGSNK	AGRNLRLEWFMDV	QSVRSN	GAS	HQYNDWFFYT
		535	536	537	538	385	516
17	PSMA_P72_F07	GFTFSSYG	TSYDGSNK	ARDFYSSSWNGAFDI	NIGSKS	DDS	QVWDSSTDHVV
		161	570	571	572	555	550
18	PSMA_P72_E07	GFTFITYG	VSEDES NK	ARALRDGNNWDYFNGMDV	SSNIGSNT	SDN	AAWDDSLINGYV
		93	94	95	96	79	74
19	PSMA_P72_D01	GFTFNNYN	ISTSSNK	ARDGVGADYGDYYYGMDV	SSDVGGYNY	EVS	SSYTSSYTYV
		501	502	639	504	487	482
20	PSMA_P72_C01	GFSFSGYG	MSYDGSNR	ARDTVWGSHPDADFID	NSGSKS	DDS	QVWDSSTDHGV
		671	672	673	674	555	652
21	PSMA_P72_A10	GFTFSSYN	IYYDGSNK	ARERGRDYGMDV	ALPKQY	KDS	QSADSSGTYV
		467	706	469	470	453	448
22	PSMA_P72_F02 and PSMA_P70_F02	GGSLISYY	IYSSGST	AKVGWMPGAFDI	SSNIGSNT	SSN	AAWDDSLINGYV
		739	740	741	96	725	720
23	PSMA_P72_G02	GFSFSGYG	ISYDGSNK	ARDELWGRGYYYGMDV	SSDVGGYNY	EVS	SSYTITSTLV
		671	536	775	504	487	754
24	PSMA_P72_A11	GFSFSGYG	ISYDGSNK	AKTTVSDPYYYGMDV	NIGSKS	DDS	QVWDSSTDHVV
		671	536	809	572	555	788

[00551] Exemplary PSMA binders in scFv format LH (VL-Linker-VH) or HL (VH-Linker-VL) are provided below.

5 Table 10. PSMA binders in scFv format LH (VL-Linker-VH) or HL (VH-Linker-VL) using the Linker of SEQ ID NO:1419

	PSMA binders in scFv format LH (VL-Linker-VH) or HL (VH-Linker-VL)
PSMB895 scFv LH	QSVLTQPPSVSAAPGQKVTISCSGSSSNIGNNYVSWYQQLPGTAPKLLIYDNNKRPSGI PDRFSGSKSGTSATLGITGLQTGDEADYYCGTWDSLSAYVFGTGKTVLGGSEKSS GSGSESKSTGGSEVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEW VSAISGGIGSTYYADSVKGRFTISRDN SKNTLWLQMN SLRAEDTAVYYCAKDGVGATPY YFDYWGQGTTLTVSS (SEQ ID NO:1485)
PSMB896 scFv LH	QSVLTQPPSVSAAPGQKVTISCSGSSSNIGINIVSWYQQLPGTAPKLLIYDNNKRPSGI PDRFSGSKSGTSATLGITGLQTGDEADYYCGTWDSLSAVVFGGKTLTVLGGSEKSS GSGSESKSTGGSEVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEW VSAISGGIGSTYYADSVKGRFTISRDN SKNTLWLQMN SLRAEDTAVYYCAKDGVGATPY YFDYWGQGTTLTVSS (SEQ ID NO:1486)
PSMB897 scFv LH	QSVLTQPPSVSAAPGQKVTISCSGSSSNIGNNYVSWYQQLPGTAPKLLIYDNNKRPSGI PDRFSGSKSGTSATLGITGLQTGDEADYYCGTWDSLSAYVFGTGKTVLGGSEKSS GSGSESKSTGGSEVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEW VSAISGGIGSTYYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCAKDGVGATPY YFDYWGQGTTLTVSS (SEQ ID NO:1487)
PSMB898 scFb LH	QSVLTQPPSVSAAPGQKVTISCSGSSSNIGINIVSWYQQLPGTAPKLLIYDNNKRPSGI PDRFSGSKSGTSATLGITGLQTGDEADYYCGTWDSLSAVVFGGKTLTVLGGSEKSS GSGSESKSTGGSEVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEW VSAISGGIGSTYYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCAKDGVGATPY YFDYWGQGTTLTVSS (SEQ ID NO:1488)
PSMA_P75_F01 scFv LH	EIVLTQSPGTLSPGERATLSCRASQSVRSNLAWYQQKPGQAPRLLIYGASTRATGIP ARFSGSGSGTEFTLTITSLQSEDFAVYYCHQYNDWPPYTFGGGTKLEIKGGSEKSSGSG GSESKSTGGSQVLQESGGGVQPGRLRLSCAASGFTFSTYGMHWVRQAPGKGLEWVA FISYDGSNKYYADSVKGRFTISRDN SKHTLYLQMN SLRAEDTAVYYCAGRDNLRLEWF MDVWGQGTTLTVSS (SEQ ID NO:1489)
PSMA_P70_F02 scFv LH	QSVLTQPPSVASGTPGQKVTISCSGSSSNIGSNTVNWYQQLPGTAPKLLIYSSNQRPSPGV PDRFSGSKSGTSASLAI SGLQSEDEADYYCAWDDSLNGVVFGGGKTLTVLGGSEKSS GSGSESKSTGGSEVQLLESGLVLPKSETLSLTCTVSGGSIISYYWSWIRQAPGKGLEW IGRIYSSGSTNYNPSLKSRTVMSVDTSKNQFSLKLSVTAADTAVYYCAKGVWPGAFD IWGQGTMTTVSS (SEQ ID NO:1490)
PSMA_P72_A10-HC-G54E scFv LH	SYELMQPPSVSVSPGQTARITCSGDALPKQYAYWYQQKPGQAPVLVIYKDSERPSPGIPV RFGSSSGTTLTITIGVQAEDEADYYCQADSSGTYVFGTGKTVLGGSEKSSGSG SESKSTGGSQVLVESGGGVQPGRLRLSCAASGFTFSSYNMNVVRQAPGKGLEWVAI IYYDESNKYADSVKGRFTISRDISKNTLYLQMN SLRAEDTAVYYCAREGRDYYGMDV WGQGTTLTVSS (SEQ ID NO:1491)
PSMA_P72_D01-HC-D95E scFv LH	QSVLTQPPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMIEVSNRPSG VSNRFGSKSGNTASLTISGLQAEDEADYYCSTSSYTYVFGTGKTLTVLGGSEKSS GSGSESKSTGGSEVQLVESGGDLVQPGGSLRLSCAASGFTFNNYNMNVVRQAPGKGLEW VSHISTSSSNKYADSVKGRFISRDI AKNSMYLQMN SLRDEDTAVYYCAREGVGADYG DYYYGMDVWGQGTTLTVSS (SEQ ID NO:1492)
PSMA_P72_F07 scFv LH	SYELTQPPSVSVAPGQTARITCGGNNIGSKSVHWYQQKPGQAPVLVYDSDRPSGIPE RFGSNGSGNTATLTISRVEAGDEADYYCQVWDSSTDHVVFGGKTLTVLGGSEKSSGSG

	GSESKSTGGSEVQLVESGGGVVQPGRSRLRLSCAASGFTFSSYGMNWVRQAPGKGLEWVA VTSYDGSNKYYADSVKGRFTISRDISKNTLYLQMSLRAEDTAVYYCARDPYSSSSWNGA FDIWGPMTMTVSS (SEQ ID NO:1493)
PSMA_P72_E07 scFv LH	QSVLTQPPSASGTPGQGVITISCSGSSSNIGSNTVNWVQQLPGTAPKLLIYSDNQRPSGV PDRFSGSKSGTSASLAISGLQSEDEADYYCAAWDDSLNGYVFGTGKTVLGGSEGKSS GSGSESKSTGGSEVQLVESGGGVVQPGRSRLRLSCAASGFTFITYGHWVRQAPGKGLEW VAVVSFDESNKYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARALRDGNNW DYFNGMDVWGQGTMTVSS (SEQ ID NO:1494)
PSMA_P72_D01 scFv LH	QSVLTQPPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMIEVSNRPSG VSNRFGSKSGNTASLTISGLQAEDEADYYCSTSYTYVFGTGKTVLGGSEGKSS GSGSESKSTGGSEVQLVESGGDLVQPGGSLRLSCAASGFTFNINMNVVRQAPGKGLEW VSHISTSSSNKYADSVKGRFTISRDIKNSMYLQMNSLRDEDTAVYYCARDGVGADYG DYVYVMDVWGQGTMTVSS (SEQ ID NO:1495)
PSMA_P72_C01 scFv LH	QSVLTQPPSVSVAPGQTARITCGGNNSGSKSVHWYQQKPGQAPVLVYDSDRPSGIPV RFGSSSGTTLTITISRVQAEDEADYYCQVWDSSTDHGVFGGKTKTVLGGSEGKSSGS GSESKSTGGSQVQLVESGGGVVQPGRSRLRLSCAASGFSFSGYGMHWVRQAPGKGLEWVA VMSYDGSNRFYVDSVRGRFTISRDNKNTLYLQMNSLRPEDTAVYYCARDTVWGSHPDA FDIWGQGTMTVSS (SEQ ID NO:1496)
PSMA_P72_A10 scFv LH	SYELMQPPSVSVSPGQTARITCSGDALPKQYAYWYQQKPGQAPVLVIYKDSERPSGIPV RFGSSSGTTLTITISRVQAEDEADYYCQVWDSSTDHGVFGGKTKTVLGGSEGKSSGS SESKSTGGSQVQLVESGGGVVQPGRSRLRLSCAASGFTFSSYNMNVVRQAPGKGLEWVAI IYYDGSNKYYADSVKGRFTISRDISKNTLYLQMNSLRAEDTAVYYCARERGRDYVMDV WGQGTMTVSS (SEQ ID NO:1497)
PSMA_P70_F02 scFv LH	QSVLTQPPSASGTPGQRTVITISCSGSSSNIGSNTVNWVQQLPGTAPKLLIYSSNQRPSGV PDRFSGSKSGTSASLAISGLQSEDEADYYCAAWDDSLNGVVFVGKTKTVLGGSEGKSS GSGSESKSTGGSEVQLLESGLVKPSETLSLTCTVSGGSIISYVSWIRQAPGKGLEW IGRIYSSGSTNYNPSLKSRTVMSVDTSKNQFSLKLSSVTAADTAVYYCAKVGVPWPAFD IWGQGTMTVSS (SEQ ID NO:1498)
PSMA_P72_G02 scFv HL	EVQLVESGGGVVQPGRSRLRLSCAASGFSFSGYGMHWVRQAPGKGLEWVAISYDGSNKY YADSVKGRFTISRDNKNTLYLQMNSLRVEDTAVYYCARDRIWGSRGYYYGMDVWGQGT TVTSSGGSEGKSSGSGSESKSTGGSQSALTQPPASVSGSPGQSITISCTGASSDVGGYN YVSWYQQHPGKAPKLMIEVSNRPSGVSNRFGSKSGNTASLTISGLQAEDEADYYCST YTITSLVFGGKTKTVL (SEQ ID NO:1499)
PSMA_P72_C01 scFv HL	QVQLVESGGGVVQPGRSRLRLSCAASGFSFSGYGMHWVRQAPGKGLEWVAVMSYDGSNRF YVDSVRGRFTISRDNKNTLYLQMNSLRPEDTAVYYCARDTVWGSHPDAFDIWGQGTVV TVSSGGSEGKSSGSGSESKSTGGSQSVLTQPPSVSVAPGQTARITCGGNNSGSKSVHWY QQKPGQAPVLVYDSDRPSGIPERFSGNSGNTATLTISRVEAGDEADYYCQVWDSST DHGVFGGKTKTVL (SEQ ID NO:1500)
PSMA_P72_A11 scFv HL	QVQLQESGGDVVQPGRSRLRLSCAASGFSFSGYGLHWVRQAPGRGLEWVTLISYDGSNKY YADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAKTTVSDPYVYVMDVWGQGTTV TVSSGGSEGKSSGSGSESKSTGGSSYELTQPPSVSVAPGQTARITCGGNNIGSKSVHWY QQKPGQAPVLVYDSDRPSGIPERFSGTNSGNTATLTISRVEAGDEADYYCQVWDSST DHVVFVGKTKTVL (SEQ ID NO:1526)
PSMA_P72_A10 scFv LH	SYELMQPPSVSVSPGQTARITCSGDALPKQYAYWYQQKPGQAPVLVIYKDSERPSGIPV RFGSSSGTTLTITISRVQAEDEADYYCQVWDSSTDHGVFGGKTKTVLGGSEGKSSGS SESKSTGGSQVQLVESGGGVVQPGRSRLRLSCAASGFTFSSYNMNVVRQAPGKGLEWVAI

	IYYDGSNKYYADSVKGRFTISRDISKNTLYLQMNSLRAEDTAVYYCARERGRDYGMVDV WGQGTTVTVSS (SEQ ID NO:1527)
PSMA_P72_C01 scFv LH	QSVLTQPPSVSVAPGQTARITCGGNNSGSKSVHWYQQKPGQAPVLVYDSDRPSGIPE RFSGSNSGNTATLTISRVEAGDEADYYCQVWDSSSDHGVFGGKTLTVLGGSEGKSSGS GSESKSTGGSQVLVESGGGEVQPGRSRLRLSCAASGFSFSGYGMHWVRQAPGKGLEWVA VMSYDGSNRFXVDSVRGRFSISRDN SKNTLYLQMNSLRPEDTAVYYCARDTVWGSHPDA FDIWGQGTVVTVSS (SEQ ID NO:1528)
PSMA_P72_D01 scFv LH	QSVLTQPPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMIEVSNRPSG VSNRFSGSKSGNTASLTISGLQAEDEADYYCSSYTSSTYTVFGTGTCLTVLGGSEGKSS GSGSESKSTGGSEVQLVESGGDLVQPGGSLRLSCAASGFTFNNYMNWVRQAPGKGLEW VSHISTSSNKYYADSVKGRFSISRDIKNSMYLQMNSLRDEDTAVYYCARDGVGADYG DYYYGMDVWGQGTTVTVSS (SEQ ID NO:1529)
PSMA_P72_E07 scFv LH	QSVLTQPPSASGTPGQGVITISCGSSSNIGSNTVNWFOQLPGTAPKLLIYSDNRPSGV PDRFSGSKSGTSASLAISGLQSEDEADYYCAAWDDSLNGYVFGTGTCLTVLGGSEGKSS GSGSESKSTGGSEVQLVESGGGVVQPGRSRLRLSCAASGFTFITGMHWVRQAPGKGLEW VAVVSFDES NKYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARALRDGNW DYFNGMDVWGQGTTVTVSS (SEQ ID NO:1530)
PSMA_P72_A11 scFv HL	QVQLQESGGDVVQPGRSRLRLSCAASGFSFSGYGLHWVRQAPGRGLEWVTLISYDGSNKY YADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKTTVDFYYYGMDVWGQGTTV TVSSGGSEGKSSGSGSESKSTGGSSYELTQPPSVSVAPGQTARITCGGNNIGSKSVHWY QQKPGQAPVLVYDSDRPSGIPEFSGTNSGNTATLTISRVEAGDEADYYCQVWDSSS DHVVFGGGKTLTVL (SEQ ID NO:1531)

[00552] Exemplary PSMA cDNAs sequences are provided below.

[00553] SEQ ID NO: 1026 (PSMB889 and PSMB890 VH cDNA)

5 GAGGTGCAACTGGTCAATCTGGAGGGGGACTGGTTAAGCCTGGGGGTAGCTTGCCTCTTAGCTGTG
CCGCTAGCGGCTTCACCTTCTCACGTTACAATATGAATTGGGTTTCGACAGGCCCTGGCAAAGGTCT
TGAGTGGGTTTCATCCATTAAC TCAAATTCACGATATATCTACTACGCAGATTCTGTAAAGGGTCGA
TTCATATATCTCGAGACAGTGCCAAGAATTCCTTGTACTTGCAGATGAATAGCTTGAGGGCCGAAG
ATACTGCCGTGTACTACTGTGCAAAGACAATGGGAGATTATTATTACTATTACGGCATGGACGTGTG
GGGGCAGGGGACAACAGTCACCGTAAGCAGT

[00554] SEQ ID NO: 1028 (PSMB891 and PSMB892 VH cDNA)

10 GAGGTGCAATTGGTTGAGAGTGGCGGGGGGTCGTTCAACCAGGTGCGAGCTTGCAGCTCTCTTGTG
CCGCCAGCGGTTTCACATTCACTACGGGATGCACTGGGTGCGACAGGCACCCGAAAGGGACT
GGAGTGGGTTGCTGTGGTGTCTTTTCGATGAGTCTAATAAGTACTATGCAGACTCTGTAAAGCCGA
TTTACTATTAGCCGAGACAAC TCAAAAATACTCTGTACCTCCAGATGAATAGTTTTCGAGCAGAAG
ATACAGCAGTTTATTATTGTGCCAGGGCTCTTAGGGACGGTAATAATTGGGATTACTTTAACGGCAT
15 GGACGTATGGGGCAAGGCACTACCGTTACTGTATCATCC

[00555] SEQ ID NO: 1030 (PSMB893 and PSMB894 VH cDNA)

CAAGTGCACTTGGTTGAGTCAGGTGGAGGAGTGGTCCAGCCTGGTTCGAGCCTCAGATTGTCTGCG
TGGCAAGCGGATTACCTTTTCTTCTTACGGTATTCAGTGGGTACGGCAAGCACCAGGGAAAGGGTT
GGAGTGGGTCGCTGTGATTGGTACGACGGGTCAAATAAGTACTACGCCGATTCCGTGAAGGGTCGC
20 TTTACAATCTCTAGGATAACTCAAAAACACACTGTACCTGCAAATGAATAGTCTCCGTGCAGAAG
ACACAGCAGTATATTATTCTGTGCGCGGAGTAGGGCCCACTTCTTATTATTACAAC TACGGTATGGA
TGTATGGGGCCAGGGCACTACCGTTACTGTATCTTCC

[00556] SEQ ID NO: 1032 (PSMB895, PSMB946, PSMB896 and PSMB947 VH cDNA)

5 GAGGTACAACCTTGTGGAAAGTGGAGGCGGTCTTGTCCAACCTGGAGGATCTCTCCGATTGAGTTGCG
CAGCCAGCGGGTTTACTTTTTCTTCATACGCCATGTCTGGGTGCGGCAAGCACCAGGTAAAGGACT
TGAGTGGGTATCTGCTATTTTCAGGGGGGATAGGCTCAACATACTATGCTGATAGCGTGAAAGGTAGG
TTCACCATTTCCCGTGACAATAGTAAAAACACATTGTGGCTCCAAATGAACAGCCTTAGGGCTGAAG
ACACCGCTGTTTACTACTGCGCTAAAGACGGTGTAGGGGCAACTCCCTATTACTTCGATTATTGGGG
ACAAGGAACCTTGGTAACAGTTTCAAGC

[00557] SEQ ID NO: 1034 (PSMB897, PSMB948, PSMB898 and PSMB949 VH cDNA)

10 GAAGTCCAGTTGGTAGAATCTGGAGGCGGGCTGGTACAGCCTGGCGGTTCCCTGCGCCTCTCATGTG
CCGCAAGCGGGTTTACCTTCAGCTCTTACGCAATGTCATGGGTGCGTCAGGCCCTGGAAAAGGTCT
CGAGTGGGTGAGTCCATTTCTGGGGGCTCCGGCTCCACCTACTACGCAGATTGAGTTAAAGGGAGA
TTTACAATCTCAAGAGATAACAGTAAAAACACCTCTACCTCCAGATGAACCACTTCGAGCTGAGG
ATTGAGCAGTATATTACTGTGCTAAAGACGGTGTAGGTGCAACTCCCTACTATTTGAGTATTGGGG
CCAAGGGACTTTGGTGACAGTAAGTAGT

15 [00558] SEQ ID NO: 1036 (PSMB899 VH cDNA)

20 GAGGTGCAGCTTGTAGAATCAGGGGGAGGACTGGTGCAGCCAGGCGGATCATTGAGACTGAGTTGTA
CTGCCAGCGGATTTCATATTCTCCAGTTATGCCATGAGCTGGGTGCGGCAGGCACCTGGAAAAGGGCT
TGAGTGGGTCTCTGCCATAAGCGGTGGATACGGTGCTCCATACTATGCTGACACCGTAAAAGGGCGG
TTCACCATCTCCCGAGACAATAGTAAGAACACCTGTATCTCCAGATGAACCTCCCTCCGCGCTGAAG
ATACAGCCGTTTATTACTGCGCCAAAGATGGAGTCGGGGCCACTCCATACTACTTCGATGATTGGGG
ACAAGGGATCCTCGTCACTGTTAGCTCA

[00559] SEQ ID NO: 1532 (PSMB896-G100A VH cDNA)

25 GAGGTGCAGCTGGTTGAATCTGGCGGAGGATTGGTTCAGCCTGGCGGTTCTCTGAGACTGTCTGTG
CCGCTTCCGGCTTCACCTTCTCCAGCTACGCTATGTCTCTGGGTCCGACAGGCTCCTGGCAAAGGACT
GGAATGGGTGTCCGCTATCTCTGGCGGCATCGGCTCTACCTACTACGCCGACTCTGTGAAGGGCAGA
TTCACCATCAGCCGGGACAACCTCAAGAACACCTGTGGCTGCAGATGAACCTCCCTGAGAGCCGAGG
ATACCGCCGTGTACTACTGTGCCAAGGATGCCGTGGGCGCTACCCCTTACTACTTCGATTATTGGGG
CCAGGGCACCTGGTCACCGTTTCTTCT

[00560] SEQ ID NO: 1037 (PSMB889 VL cDNA)

30 CAATCCGTGCTTACACAACCACCATCAGTAAGCGGGGCTCCAGGCCAAAGAGTAACCATCTCATGCA
CTGGTTCAGCTTCAACCTTGGTGCCGGATATGATGTCCATTGGTATCAACAAGTCCCTGGAACCGT
CCCAAACTCCTGATCTACGATAATTCAAATCGACCTTCCGGAGTGCCAGATCGGTTGAGTGGATCT
AAAAGCGGCACCAAGTGCCTCTCTTGCCATTACTGGATTGCAAGCTGAAGATGAGACAGTTTATTACT
GTCAGTCTACGATTCTAGTTTGTCCGGAGTGTTTTCGGTGGTGGTACTAAGCTCACCGTGCTG

35 [00561] SEQ ID NO: 1038 (PSMB890 VL cDNA)

40 AGCAGCGAGCTGACACAACCTCCCTCCGTTTCTGGAGCCCCAGGTCAAAGAGTGACTATTAGTTGTG
CAGGCTCCCTTTCAAATATAGGAGCAGGTTATGATGTTTATTGGTATCAACAACCTCCAGGCACAGC
ACCAAACTCCTTATATACGGAACATAAATCGCTTGAGCGGTGTACCCGAACGCTTCAGTGGCAGC
AAGTCTGGCACCAGCGCTTCTTGGAATAACCGGACTCCAGGCTGAGGATGGCGCTGATTACTATT
GCCAGTCTATGATAGCTCCCTCTCTTCATATGTCTTCGGAACCTGGTACAAAAGTGACCGTGCTT

[00562] SEQ ID NO: 1039 (PSMB891 and PSMB893 VL cDNA)

CAGTCCGTGTTGACACAACCTCCAAGCGCCTCAGGGACCCAGGACAGGGTGTGACAATATCCTGTT
CCGGCAGTAGCAGTAACATTGGGTCTAATACCGTTAACTGGTTCAGCAACTGCCCGGCACAGCTCC
CAAACTTTTGATCTACAGTGACAACCAACGACCTCTGGCGTACCCGACCGTTTCTCAGGTTCCAAA

AGCGGGACCTCAGCATCTCTCGCTATCTCCGGTCTGCAAAGTGAGGACGAAGCCGATTACTACTGTG
CAGCCTGGGACGATTCAGTGAATGGGTATGTTTTCGGAACCTGGTACCAAAGTGACTGTGTTG

[00563] SEQ ID NO: 1040 (PSMB892 and PSMB894 VL cDNA)

5 CAGTCCGTTCTCACACAGCCACCATCAGTAAGTGGCGCTCCCGGTCAAAGAGTAACTATTTTCATGTA
CTGGTAGTTCAAGCAATATAGGTGCTGACTATGACGTCCATTGGTATCAGCACTTGCCCGGCACTGC
ACCTAACTTCTGATATACGGAAATAGTAACAGACCTAGTGGGGTTCCAGACCGTTTCTCCGGGAGT
AAAAGTGGAAACAAGCGCTTCACTTGCAATTACTGGCCTCCAGGCTGAAGATGAAACCGACTATTACT
GCCAAAGCTACGATTCTAGCCTGTCTGGGTGGGTTTTCGGCGGGGGAAGTAAATTGACCGTCTTG

[00564] SEQ ID NO: 1043 (PSMB895, PSMB946, PSMB897 and PSMB948 VL cDNA)

10 CAATCTGTCTGACTCAACCTCCCTCAGTCTCAGCCGACCCAGGACAGAAGGTGACAATTAGCTGTT
CAGGTTCTTCAAGTAACATCGGTAACAACACTACGTCTCATGGTATCAACAGCTTCCCGGAACAGCACC
CAAACCTGCTGATATATGATAACAACAAACGGCCATCTGGAATACCAGACCGGTTCTCAGGCTCCAAG
AGCGGTACTAGCGCAACTTTGGGAATCACCGGTTTGCACTGGGGATGAGGCAGACTATTACTGCG
GCACCTGGGATTCCAGTCTGTCTGCTTATGTTTTGGGACCGGGACAAAGGTGACTGTCCTT

15 [00565] SEQ ID NO: 1044 (PSMB896, PSMB947, PSMB898 and PSMB949 VL cDNA)

CAGAGTGTCTTACTCAGCCTCCTAGCGTTAGCGCCGCCCTGGACAGAAGGTTACTATCTCTCTGCT
CAGGGAGTTCAGTAATATTGGAATCAATTATGTGAGTTGGTATCAGCAGTTGCCCGGCACCGCTCC
TAAATFGCTTATCTATGACAACAATAAACGACCTAGTGGTATCCCTGATCGTTTTCTGGATCAAAA
20 TCTGGTACTAGCGCAACCCCTCGGTATCACCGGACTGCAAACAGGTGATGAAGCAGACTATTATTGCG
GCACCTGGGACTCATCACTCTCCGCCGTCGTTTTCGGGGGCGGAACCAACTTACAGTATTG

[00566] SEQ ID NO: 1047 (PSMB899 VL cDNA)

CAAAGCGTGCTGACACAGCCTCCATCAGTCAGCGCAGCCCCAGGACAGAAAGTCACTATTTCTTGCA
GCGGTAGCAGCTCTAATATCGGCAATAATTACGTCTCCTGGTATCAGCAGCTCCCCGGCACTGCACC
25 TAAGTTGCTCATCTTCGATAATAACAAACGTCTTCAGGCATTCCCGACCGATTCTCAGGTTCAAAA
TCAGGAACCAAGTGCTACTTTGGGAATAACAGGTCTGCAAACCTGGTGATGAAGCTGATTACTACTGCG
GTACCTGGGATAGCTCTCTCAGTGCTATGTATTGCGGACTGGCACAAAGGTTACTGTTCTG

[00567] SEQ ID NO: 1533 (PSMB896-G100A VL cDNA)

30 CAGTCTGTGCTGACCCAGCCTCCTTCTGTGTCTGCTGCTCCTGGCCAGAAAGTGACCATCTCCTGCT
CCGGCTCCTCCTCAACATCGGCATCAACTACGTGTCTGGTATCAGCAGCTGCCCGGCACAGCTCC
CAAACCTGCTGATCTACGACAACAACAAGCGGCCAGCGGCATCCCTGACAGATTCTCCGGATCTAAG
TCCGGCACCTCTGCTACCCTGGGCATCACAGGATTGCAGACAGGCGACGAGGCCGACTACTATTGCG
GCACCTGGGACTCTTCTCTGTCCGCCGTTGTTTTGGCGGTGGCACCAAGCTGACAGTGCTG

[00568] SEQ ID NO: 1048 (PSMB889 and PSMB890 HC cDNA)

35 GAGGTGCAACTGGTCAATCTGGAGGGGGACTGGTTAAGCCTGGGGGTAGCTTGCGTCTTAGCTGTG
CCGCTAGCGGCTTCACCTTCTCACGTTACAATATGAATTGGGTTCGACAGGCCCTTGGCAAAGGTCT
TGAGTGGGTTTCATCCATTAACCAAAATTCACGATATATCTACTACGCAGATTCTGTGAAGGGTCGA
TTCACTATATCTCGAGACAGTGCCAAGAATTCCCTGTACTTGCAAGATGAATAGCTTGAGGGCCGAAG
ATACTGCCGTGTACTACTGTGCAAAGACAATGGGAGATTATTATTACTATTACGGCATGGACGTGTG
40 GGGGCAGGGGACAACAGTCACCGTAAGCAGTGCTTCCACCAAGGGCCCATCCGTCTTCCCCCTGGCA
CCCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCG
AACCGGTGACGGTGTCTGGAACCTCAGGCGCCCTGACCAGCGCGTGCACACCTTCCCGGTGTCTCT
ACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAG
ACCTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCAAT
CTTGTGACAAAACACACATGTCCACCGTGCCAGCACCTGAAGCAGCAGGGGGACCGTCAGTCTT
45 CCTCTTCCCCCAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTG
GTGAGCGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATA

ATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCCT
CCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCC
CCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACACAGGTGTACACCCCTGCCCC
CATCCCGGGAGGAGATGACCAAGAACAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAG
5 CGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACTACAAGACCACGCCTCCCGTG
CTGGACTCCGACGGCTCCTTCTTCTCTACAGCCGGCTCACCCTGGACAAGTCTAGATGGCAGCAGG
GGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTC
CCTGTCTCCGGGT

[00569] SEQ ID NO: 1050 (PSMB891 and PSMB892 HC cDNA)

10 GAGGTGCAATTGGTTGAGAGTGGCGGGGGGGTCTTCAACCAGGTGCGAGCTTGCAGCTCTCTTGTG
CCGCCAGCGGTTTTCACATTCACTAATTACGGGATGCACTGGGTGCGACAGGCACCCGGAAGGGACT
GGAGTGGGTTGCTGTGGTGTCTTTGATGAGTCTAATAAGTACTATGCAGACTCTGTAAAGGCCGA
TTTACTATTAGCCGAGACAACCTCTAAAAATACTCTGTACCTCCAGATGAATAGTTTGCGAGCAGAAG
ATACAGCAGTTTATTATTGTGCCAGGGCTCTTAGGGACGGTAATAATTGGGATTACTTTAACGGCAT
15 GGACGTATGGGGGCAAGGCACCTACCGTTACTGTATCATCCGCTTCCACCAAGGGCCCATCCGTCTTC
CCCCGGCACCCTCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACT
ACTTCCCCGAACCGGTGACGGTGTCTGGAACCTCAGGCGCCCTGACCAGCGCGTGCACACCTTCCC
GGCTGTCTTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTG
GGCACCAGACCTACATCTGCAACGTGAATCACAAGCCAGCAACACCAAGGTGGACAAGAAAGTTG
20 AGCCCAAAATCTTGTGACAAAACCTCACACATGTCCACCGTCCCAGCACCTGAAGCAGCAGGGGGACC
GTCAGTCTTCTCTTCCCCCAAAACCAAGGACACCTCATGATCTCCCGGACCCCTGAGGTCAACA
TGCGTGGTGGTGAGCGTGAGCCACGAAGACCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGG
AGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGT
CCTCACCGTCTTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCC
25 CTCCCAGCCCCATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACACAGGTGTACA
CCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACAGGTGAGCCTGACCTGCCTGGTCAAAGGCTT
CTATCCAGCAGCATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACTACAAGACCAAG
CCTCCCGTGTCTGGACTCCGACGGCTCCTTCTTCTCTACAGCCGGCTCACCCTGGACAAGTCTAGAT
GGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAA
30 GAGCCTCTCCCTGTCTCCGGGT

[00570] SEQ ID NO: 1052 (PSMB893 and PSMB894 HC cDNA)

CAAGTGCAGTTGGTTGAGTCAGGTGGAGGAGTGGTCCAGCCTGGTCCGAGCCTCAGATTGTCTGCG
TGGCAAGCGGATTACCTTTTCTTCTTACGGTATTCAGTGGGTACGGCAAGCACCAGGGAAAGGGTT
GGAGTGGGTGCTGTGATTGGTACGACGGGTCAAATAAGTACTACGCCGATTCCGTGAAGGGTCGC
35 TTTACAATCTCTAGGATAACTCAAAAACACACTGTACCTGCAAATGAATAGTCTCCGTGCAGAAG
ACACAGCAGTATATTATTCTGTGCGCGGAGTAGGGCCCACTTCTTATTATTACAACTACGGTATGGA
TGTATGGGGCCAGGGCACTACCGTTACTGTATCTTCCGCTTCCACCAAGGGCCCATCCGTCTTCCCC
CTGGCACCTCTTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACT
TCCCCGAACCGGTGACGGTGTCTGGAACCTCAGGCGCCCTGACCAGCGGCGTGACACCTTCCCGGC
40 TGTCTTACAGTCTTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGC
ACCCAGACCTACATCTGCAACGTGAATCACAAGCCAGCAACACCAAGGTGGACAAGAAAGTTGAGC
CCAAATCTTGTGACAAAACCTCACACATGTCCACCGTGCCAGCACCTGAAGCAGCAGGGGGACCGTC
AGTCTTCTCTTCCCCCAAAACCAAGGACACCTCATGATCTCCCGGACCCCTGAGGTACATGC
GTGGTGGTGAGCGTGAGCCACGAAGACCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGG
45 TGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCT
CACCCTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTC
CCAGCCCCATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACACAGGTGTACACCC
TGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTCTA
TCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACTACAAGACCACGCCT
50 CCCGTGTGGACTCCGACGGCTCCTTCTTCTCTACAGCCGGCTCACCCTGGACAAGTCTAGATGGC
AGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAG
CCTCTCCCTGTCTCCGGGT

[00571] SEQ ID NO: 1054 (PSMB895 and PSMB896 HC cDNA)

GAGGTACAACTTGTGGAAAGTGGAGGCGGTCTTGTCCAACCTGGAGGATCTCTCCGATTGAGTTGCG
CAGCCAGCGGGTTTACTTTTTCTTCATACGCCATGTCTGGGTGCGGCAAGCACCAGGTAAAGGACT
TGAGTGGGTATCTGCTATTTTCAGGGGGGATAGGCTCAACATACTATGCTGATAGCGTGAAAGGTAGG
5 TTCACCATTTCCCGTGACAATAGTAAAAACACATTGTGGCTCCAAATGAACAGCCTTAGGGCTGAAG
ACACCGCTGTTTACTACTGCGCTAAAGACGGTGTAGGGGCAACTCCCTATTACTTCGATTATTGGGG
ACAAGGAACCTTGGTAACAGTTTCAAGCGCTTCCACCAAGGGCCCATCCGTCTTCCCCCTGGCACCC
TCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAAC
CGGTGACGGTGTCTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTCTACA
10 GTCTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACC
TACATCTGCAACGTGAATCACAAGCCCAGCAACCAAGGTGGACAAGAAAGTTGAGCCCCAAATCTT
GTGACAAAACCTACACATGTCCACCGTGCCAGCACCTGAAGCAGCAGGGGGACCGTCAGTCTTCTCT
CTTCCCCCAAAACCCAAGGACACCCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTG
AGCGTGAGCCACGAAGACCCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATG
15 CCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCTCTACCGTCTCT
GCACCAGGACTGGTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCC
ATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCCTGCCCCCAT
CCCCGGAGGAGATGACCAAGAACCAGGTGACCGTGACCTGCCCTGGTCAAAGGCTTCTATCCCAGCGA
CATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAGAACAACCTACAAGACCACGCCTCCCGTGCTG
20 GACTCCGACGGCTCTTCTTCTCTACAGCCGGCTCACCCTGGACAAGTCTAGATGGCAGCAGGGGA
ACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCT
GTCTCCGGGT

[00572] SEQ ID NO: 1501 (PSMB896-G100A HC cDNA)

GAGGTGCAGCTGGTTGAATCTGGCGGAGGATTGGTTGAGCCTGGCGGTCTCTGAGACTGTCTGTG
25 CCGCTTCCGGCTTCACCTTCTCCAGCTACGCTATGTCTGGGTCCGACAGGCTCCTGGCAAAGGACT
GGAATGGGTGTCCGCTATCTCTGGCGGCATCGGCTCTACCTACTACGCCGACTCTGTGAAGGGCAGA
TTCACCATCAGCCGGGACAACTCCAAGAACACCCCTGTGGCTGCAGATGAACCTCCCTGAGAGCCGAGG
ATACCGCCGTGTACTACTGTGCCAAGGATGCCGTGGGCGCTACCCCTTACTACTTCGATTATTGGGG
CCAGGGCACCCTGGTCACCGTTTCTTCTGCCCTCCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCC
30 TCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCGAAC
CGGTGACGGTGTCTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTCTACA
GTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACC
TACATCTGCAACGTGAATCACAAGCCCAGCAACCAAGGTGGACAAGAAAGTTGAGCCCCAAATCTT
GTGACAAAACCTACACATGCCACCGTGCCAGCACCTGAAGCCGCGGGGGACCGTCAGTCTTCTCT
35 CTTCCCCCAAAACCCAAGGACACCCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTG
AGCGTGAGCCACGAAGACCCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATG
CCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCTCTACCGTCTCT
GCACCAGGACTGGTGAATGGCAAGGAGTACAAGTGCAAGGTGTGAACAAAGCCCTCCCAGCCCCC
ATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCCTGCCCCAT
40 CCCGGGAGGAGATGACCAAGAACCAGGTGACCGTGTCTTGGCGCGTCAAAGGCTTCTATCCCAGCGA
CATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAGAACAACCTACAAGACCACGCCTCCCGTGCTG
GACTCCGACGGCTCTTCTTCTCTGAGCAAGCTCACCCTGGACAAGAGCAGATGGCAGCAGGGGA
ACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCGGTTACGCAGAAGTCTCTCTCCCT
GTCTCCGGGAAAA

[00573] SEQ ID NO: 1056 (PSMB897 and PSMB898 HC cDNA)

GAAGTCCAGTTGGTAGAATCTGGAGGCGGGCTGGTACAGCCTGGCGGTTCCTTGCGCCTCTCATGTG
CCGCAAGCGGGTTTACCTTCAGCTCTTACGCAATGTCTGCGGTGCGTCAGGCCCTGGAAAAGGTCT
CGAGTGGGTGAGTGCCATTTCTGGGGGCTCCGGCTCCACCTACTACGCAGATTCAGTTAAAGGGAGA
TTTACAATCTCAAGAGATAACAGTAAAAACACCCCTCTACCTCCAGATGAACCTCACTTCGAGCTGAGG
50 ATTACAGCAGTATATTACTGTGCTAAAGACGGTGTAGGTGCAACTCCCTACTATTTTCGACTATTGGGG
CCAAGGGACTTTGGTGACAGTAAGTAGTGCTCCACCAAGGGCCCATCCGTCTTCCCCCTGGCACCC

TCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAAC
CGGTGACGGTGTCTGGAACCTCAGGCGCCCTGACCAGCGGCGTGACACACCTTCCCGGCTGTCTACA
GTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACC
TACATCTGCAACGTGAATCACAAGCCCAGCAACCAAGGTGGACAAGAAAGTTGAGCCCCAATCTT
5 GTGACAAAACCTCACACATGTCCACCGTGCCAGCACCTGAAGCAGCAGGGGGACCGTCAGTCTTCCT
CTTCCCCCAAAACCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTG
AGCGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATG
CCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCT
GCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCC
10 ATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCAT
CCCGGGAGGAGATGACCAAGAACCAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGA
CATCGCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACTACAAGACCACGCCTCCCGTGCTG
GACTCCGACGGCTCTTCTTCTCTACAGCCGGCTCACCGTGGACAAGTCTAGATGGCAGCAGGGGA
ACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCT
15 GTCTCCGGGT

[00574] SEQ ID NO: 1058 (PSMB899 HC cDNA)

GAGGTGCAGCTTGTAGAATCAGGGGGAGGACTGGTGCAGCCAGGCGGATCATTGAGACTGAGTTGTA
CTGCCAGCGGATTCATATTCTCCAGTTATGCCATGAGCTGGGTGCGGCAGGCACCTGGAAAAGGGCT
TGAGTGGGTCTCTGCCATAAGCGGTGGATACGGTGCTCCATACTATGCTGACACCGTAAAAGGGCGG
20 TTCACCATCTCCCGAGACAATAGTAAGAACACCTGTATCTCCAGATGAATCCCTCCGCGCTGAAG
ATACAGCCGTTTATTACTGCGCCAAAGATGGAGTCGGGGCCACTCCATACTACTTCGATGATTGGGG
ACAAGGGATCCTCGTCACTGTTAGCTCAGCTTCCACCAAGGGCCCATCCGTCTTCCCCCTGGCACCC
TCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAAC
CGGTGACGGTGTCTGGAACCTCAGGCGCCCTGACCAGCGGCGTGACACACCTTCCCGGCTGTCTACA
25 GTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACC
TACATCTGCAACGTGAATCACAAGCCCAGCAACCAAGGTGGACAAGAAAGTTGAGCCCCAATCTT
GTGACAAAACCTCACACATGTCCACCGTGCCAGCACCTGAAGCAGCAGGGGGACCGTCAGTCTTCCT
CTTCCCCCAAAACCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTG
AGCGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATG
30 CCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCT
GCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCC
ATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCAT
CCCGGGAGGAGATGACCAAGAACCAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGA
CATCGCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACTACAAGACCACGCCTCCCGTGCTG
35 GACTCCGACGGCTCTTCTTCTCTACAGCCGGCTCACCGTGGACAAGTCTAGATGGCAGCAGGGGA
ACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCT
GTCTCCGGGT

[00575] SEQ ID NO: 1059 (PSMB946 and PSMB947 HC cDNA)

GAGGTGCAGCTGGTTGAATCTGGCGGAGGATTGGTTGAGCCTGGCGGTTCTCTGAGACTGTCTGTG
40 CCGCTTCCGGCTTCACCTTCTCCAGCTACGCTATGTCTGGGTCCGACAGGCTCCTGGCAAAGGACT
GGAATGGGTGTCCGCTATCTCTGGCGGCATCGGCTCTACCTACTACGCCGACTCTGTGAAGGGCAGA
TTCACCATCAGCCGGGACAACCTCAAGAACACCCCTGTGGCTGCAGATGAATCCCTGAGAGCCGAGG
ATACCGCCGTGTACTACTGTGCCAAAGATGGCGTGGGCGCTACCCCTTACTACTTCGATTATTGGGG
CCAGGGCACCCCTGGTCACCGTTTCTTCTGCCTCCACCAAGGGTCCATCGGTCTTCCCCCTGGCACCC
45 TCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAAC
CGGTGACGGTGTCTGGAACCTCAGGCGCCCTGACCAGCGGCGTGACACACCTTCCCGGCTGTCTACA
GTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACC
TACATCTGCAACGTGAATCACAAGCCCAGCAACCAAGGTGGACAAGAAAGTTGAGCCCCAATCTT
GTGACAAAACCTCACACTTGTCCACCGTGCCAGCACCTGAAGCAGCAGGGGGACCGTCAGTCTTCCT
50 CTTCCCCCAAAACCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTG
AGCGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATG
CCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCT

GCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTGTCCAACAAAGCCCTCCCAGCCCC
ATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCAT
CCCGGGAGGAGATGACCAAGAACCAGGTACGCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGA
CATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACCACGCCTCCCGTGCTG
5 GACTCCGACGGCTCCTTCTTCTCTACAGCCGGCTCACCCTGGACAAGTCCAGATGGCAGCAGGGGA
ACGCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGTCTCTCTCCCT
GTCTCCGGGAAAA

[00576] SEQ ID NO: 1061 (PSMB948 and PSMB949 HC cDNA)

GAGGTGCAGCTGGTTGAATCTGGCGGAGGATTGGTTCAGCCTGGCGGTTCTCTGAGACTGTCTTGTG
10 CCGCTTCCGGCTTACCTTCTCCAGCTACGCTATGTCTGGGTCCGACAGGCTCCTGGCAAAGGACT
GGAATGGGTGTCCGCTATCTCTGGCGGATCCGGCTCTACCTACTACGCCGACTCTGTGAAGGGCAGA
TTCACCATCAGCCGGGACAACCTCCAAGAACACCTGTACCTGCAGATGAACCTCCTGAGAGCCGAGG
ACTCCGCCGTGTACTACTGTGCTAAAGATGGCGTGGGCGCTACCCCTTACTACTTTCGATTATTGGGG
CCAGGGCACCCTGGTCACCGTTTCTTCTGCCTCCACCAAGGTTCCATCGGTCTTCCCCCTGGCACC
15 TCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCGAAC
CGGTGACGGTGTCTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTCTACA
GTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACC
TACATCTGCAACGTGAATCACAAGCCCAGCAACCAAGGTGGACAAGAAAGTTGAGCCCAAATCTT
GTGACAAAACCTCACACTTGTCCACCGTGCCAGCACCTGAAGCAGCAGGGGGACCGTCAGTCTTCCT
20 CTTCCCCCAAAACCAAGGACACCCTCATGATCTCCCGGACCCTGAGGTACATGCGTGGTGGTG
AGCGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCTGGAGGTGCATAATG
CCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCTCACCCTCCT
GCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTGTCCAACAAAGCCCTCCCAGCCCC
ATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCAT
25 CCCGGGAGGAGATGACCAAGAACCAGGTACGCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGA
CATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACCACGCCTCCCGTGCTG
GACTCCGACGGCTCCTTCTTCTCTACAGCCGGCTCACCCTGGACAAGTCCAGATGGCAGCAGGGGA
ACGCTCTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGTCTCTCTCCCT
GTCTCCGGGAAAA

[00577] SEQ ID NO: 1063 (PSMB889 LC cDNA)

CAATCCGTGCTTACACAACCACCATCAGTAAGCGGGGCTCCAGGCCAAAGAGTAACCATCTCATGCA
CTGGTTCAGCTTCAACCTTGGTGCCGGATATGATGTCCATTGGTATCAACAAGTCCCTGGAACCGT
CCCAAACTCCTGATCTACGATAATTCAAATCGACCTTCCGGAGTGCCAGATCGGTTCACTGGATCT
AAAAGCGGCACCAGTGCTCTCTTGCCATTACTGGATTGCAAGCTGAAGATGAGACAGTTTATTACT
35 GTCAGTCTACGATTCTAGTTTGTCCGGAGTGGTTTTCGGTGGTGGTACTAAGCTCACCCTGCTGGG
TCAGCCCAAGGCTGCACCCAGTGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAG
GCCACACTGGTGTGTCTATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCTGGAAGGCCGATA
GCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAAGCAACAACAAGTACGCGGC
CAGCAGCTATCTGAGCCTGACGCCGTGAGCAGTGGAAAGTCCACAGAAGCTACAGCTGCCAGGTCACG
40 CATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCTACAGAATGTTCA

[00578] SEQ ID NO: 1064 (PSMB890 LC cDNA)

AGCAGCGAGCTGACACAACCTCCCTCCGTTTCTGGAGCCCCAGGTCAAAGAGTGACTATTAGTTGTG
CAGGCTCCCTTTCAAATATAGGAGCAGGTTATGATGTTCAATGGTATCAACAACCTCCAGGCACAGC
ACCAAACTCCTTATATACGGAAACATAAATCGCTTGAGCGGTGTACCCGAACGCTTCACTGGCAGC
45 AAGTCTGGCACCAGCGCTTCTTGGCAATAACCGGACTCCAGGCTGAGGATGGCGCTGATTACTATT
GCCAGTCTATGATAGCTCCCTCTCTCATATGTCTTGGAACTGGTACAAAAGTGACCGTGCTTGG
TCAGCCCAAGGCTGCACCCAGTGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAG
GCCACACTGGTGTGTCTATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCTGGAAGGCCGATA
GCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAAGCAACAACAAGTACGCGGC
50 CAGCAGCTATCTGAGCCTGACGCCGTGAGCAGTGGAAAGTCCACAGAAGCTACAGCTGCCAGGTCACG
CATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCTACAGAATGTTCA

[00579] SEQ ID NO: 1065 (PSMB891 and PSMB893 LC cDNA)

CAGTCCGTGTTGACACAACCTCCAAGCGCCTCAGGGACCCCAGGACAGGGTGTGACAATATCCTGTT
CCGGCAGTAGCAGTAACATTGGGTCTAATACCGTTAACTGGTTCAGCAACTGCCCCGGCACAGCTCC
CAAACCTTTTGATCTACAGTGACAACCAACGACCTCTGGCGTACCCGACCGTTTCTCAGGTTCCAAA
5 AGCGGGACCTCAGCATCTCTCGCTATCTCCGGTCTGCAAAGTGAGGACGAAGCCGATTACTACTGTG
CAGCCTGGGACGATTCACTGAATGGGTATGTTTTCGGAAGTGGTACCAAAGTGACTGTGTTGGGTCA
GCCCCAAGGCTGCACCCAGTGCTACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCC
ACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGAAGGCCGATAGCA
GCCCCGTCAAGGGGGAGTGGAGACCACCACACCCTCCAAACAAAGCAACAACAAGTACGCGGCCAG
10 CAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAGCTGCCAGGTCACGCAT
GAAGGGAGCACCGTGAGAAGACAGTGGCCCCCTACAGAATGTTCA

[00580] SEQ ID NO: 1066 (PSMB892 and PSMB894 LC cDNA)

CAGTCCGTTCTCACACAGCCACCATCAGTAAGTGGCGCTCCCGGTCAAAGAGTAACTATTTTCATGTA
CTGGTAGTTCAAGCAATATAGGTGCTGACTATGACGTCCATTGGTATCAGCACTTGCCCCGGCACTGC
15 ACCTAACTTCTGATATACGGAAATAGTAACAGACCTAGTGGGGTTCAGACCGTTTCTCCGGGAGT
AAAAGTGAACAAGCGCTTCACTTGCAATTACTGGCCTCCAGGCTGAAGATGAAACCGACTATTACT
GCCAAAGCTACGATTCTAGCCTGTCTGGTGGGTTTTTCGGCGGGGAACTAAATTGACCGTCTTGGG
TCAGCCCAAGGCTGCACCCAGTGCTACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAG
GCCACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGAAGGCCGATA
20 GCAGCCCCGTCAAGGGGGAGTGGAGACCACCACACCCTCCAAACAAAGCAACAACAAGTACGCGGCC
CAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAGCTGCCAGGTCACG
CATGAAGGGAGCACCGTGAGAAGACAGTGGCCCCCTACAGAATGTTCA

[00581] SEQ ID NO: 1069 (PSMB895, PSMB846, PSMB897 and PSMB848 LC cDNA)

CAATCTGTCTGACTCAACCTCCCTCAGTCTCAGCCGCACCAGGACAGAAGGTGACAATTAGCTGTT
25 CAGGTTCTTCAAGTAACATCGGTAACAACCTACGTCTCATGGTATCAACAGCTTCCCGGAACAGCACC
CAAACCTGCTGATATATGATAACAACAACGGCCATCTGGAATACCAGACCGGTTCTCAGGCTCCAAG
AGCGGTACTAGCGCAACTTTGGGAATCACCGGTTTGACAGTGGGGATGAGGCAGACTATTACTGCG
GCACCTGGGATTCCAGTCTGTCTGCTTATGTTTTGGGACCGGGACAAAGGTGACTGTCTTGGTCA
GCCCCAAGGCTGCACCCAGTGCTACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCC
30 ACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGAAGGCCGATAGCA
GCCCCGTCAAGGGGGAGTGGAGACCACCACACCCTCCAAACAAAGCAACAACAAGTACGCGGCCAG
CAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAGCTGCCAGGTCACGCAT
GAAGGGAGCACCGTGAGAAGACAGTGGCCCCCTACAGAATGTTCA

[00582] SEQ ID NO: 1070 (PSMB896, PSMB847, PSMB898 and PSMB849 LC cDNA)

CAGAGTGTCTTACTCAGCCTCCTAGCGTTAGCGCCGCCCTGGACAGAAGGTTACTATCTCCTGCT
CAGGGAGTTCCAGTAATATTGGAATCAATTATGTGAGTTGGTATCAGCAGTTGCCCCGGCACCGCTCC
TAAATTGCTTATCTATGACAACAATAAACGACCTAGTGGTATCCCTGATCGTTTTTCTGGATCAAAA
TCTGGTACTAGCGCAACCCTCGGTATCACCGGACTGCAAACAGGTGATGAAGCAGACTATTATTGCG
GCACCTGGGACTCATCACTCTCCGCCGTGTTTTTCGGGGGCGGAACCAAACCTTACAGTATTGGGTCA
40 GCCCCAAGGCTGCACCCAGTGCTACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCC
ACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGAAGGCCGATAGCA
GCCCCGTCAAGGGGGAGTGGAGACCACCACACCCTCCAAACAAAGCAACAACAAGTACGCGGCCAG
CAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAGCTGCCAGGTCACGCAT
GAAGGGAGCACCGTGAGAAGACAGTGGCCCCCTACAGAATGTTCA

[00583] SEQ ID NO: 1073 (PSMB899 LC cDNA)

CAAAGCGTGCTGACACAGCCTCCATCAGTCAGCGCAGCCCCAGGACAGAAAGTCACTATTTCTTGCA
GCGGTAGCAGCTCTAATATCGGCAATAATTACGTCTCCTGGTATCAGCAGTCCCCGGCACTGCACC
TAAGTTGCTCATCTTCGATAATAACAACGTCCTTCAGGCATTCCCGACCGATTCTCAGGTTCAAAA

5 TCAGGAACCCAGTGCTACTTTGGGAATAACAGGTCTGCAAACCTGGTGATGAAGCTGATTACTACTGCG
GTACCTGGGATAGCTCTCTCAGTGCCTATGTATTCTGGGACTGGCACAAAGGTTACTGTTCTGGGTCA
GCCCCAAGGCTGCACCCAGTGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCC
ACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGAAGGCCGATAGCA
GCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAAGCAACAACAAGTACGCGGCCAG
CAGCTATCTGAGCCTGACGCCTGAGCAGTGAAGTCCCACAGAAGCTACAGCTGCCAGGTCACGCAT
GAAGGGAGCACCGTGGAGAAGACAGTGGCCCCCTACAGAATGTTCA

[00584] SEQ ID NO: 1534 (PSMB896-G100A LC cDNA)

10 CAGTCTGTGCTGACCCAGCCTCCTTCTGTGTCTGCTGCTCCTGGCCAGAAAGTGACCATCTCCTGCT
CCGGCTCCTCCTCCAACATCGGCATCAACTACGTGTCTGGTATCAGCAGCTGCCCCGGCACAGCTCC
CAAACCTGCTGATCTACGACAACAACAAGCGGGCCAGCGGCATCCCTGACAGATTCTCCGGATCTAAG
TCCGGCACCTCTGCTACCCCTGGGCATCACAGGATTGCAGACAGGCGACGAGGCCGACTACTATTGCG
GCACCTGGGACTCTTCTCTGTCCGCGTTGTTTTTGGCGGTGGCACCAGCTGACAGTGTGGGTCA
15 GCCCCAAGGCTGCACCCAGTGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCC
ACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGAAGGCCGATAGCA
GCCCCGTCAAGGCGGGAGTCGAAACCACCACACCCTCCAAACAAAGCAACAACAAGTACGCGGCCAG
CAGCTATCTGAGCCTGACGCCTGAGCAGTGAAGTCCCACAGAAGCTACAGCTGCCAGGTCACGCAT
GAAGGGAGCACCGTGGAGAAGACAGTGGCCCCCTACAGAATGTTCA

[00585] Exemplary amino acid and nucleic acid sequence of HC and LC of certain PSMA
20 antibodies in DuoBody format or in ScFv_LH-hinge CH2-CH3 format are provided below.

Table 11. PSMA Antibodies (DuoBody)

	HC AA	LC AA	HC DNA	LC DNA
PSMB889	SEQ ID NO:33	SEQ ID NO:34	SEQ ID NO:1048	SEQ ID NO:1063
PSMB890	SEQ ID NO:33	SEQ ID NO:68	SEQ ID NO:1048	SEQ ID NO:1064
PSMB891	SEQ ID NO:101	SEQ ID NO:102	SEQ ID NO:1050	SEQ ID NO:1065
PSMB892	SEQ ID NO:101	SEQ ID NO:136	SEQ ID NO:1050	SEQ ID NO:1066
PSMB893	SEQ ID NO:169	SEQ ID NO:102	SEQ ID NO:1052	SEQ ID NO:1065
PSMB894	SEQ ID NO:169	SEQ ID NO:136	SEQ ID NO:1052	SEQ ID NO:1066
PSMB895	SEQ ID NO:237	SEQ ID NO:238	SEQ ID NO:1054	SEQ ID NO:1069
PSMB896	SEQ ID NO:237	SEQ ID NO:272	SEQ ID NO:1054	SEQ ID NO:1070
PSMB896-G100A	SEQ ID NO:441	SEQ ID NO:272	SEQ ID NO:1501	SEQ ID NO:1534
PSMB897	SEQ ID NO:305	SEQ ID NO:238	SEQ ID NO:1056	SEQ ID NO:1069
PSMB898	SEQ ID NO:305	SEQ ID NO:272	SEQ ID NO:1056	SEQ ID NO:1070
PSMB899	SEQ ID NO:373	SEQ ID NO:374	SEQ ID NO:1058	SEQ ID NO:1073

[illegible]

	TCPPCPAPEAAGGSVFLFPKPKDIT LMI SRTPEVTCVVSVSHEDPEVFN WYDVEVHNNAKTRPREEQYNSTRV VSVLTVLHQDWLNGKEYKCKVSNKAL PAPHEKTI I SKAKGQPREPOFVIVLPPS REEMTKNQVSLTCLVKGFYPSDIAVE WESNGQPENNYLTWPPVLSDSDGSFEL YSKLTVDKSRWQQGNVFCFSVMHEAL HNHYTQKSLSLSPGK (SEQ ID NO: 1140)	GCACCCCTGTCACCGTTTCTTCTGAGCCCAAACTAGCGCAAAAACCTACACTTGTTCACACCGTGCACGCAACC TGAAGACGACAGGGGACCGTCACTTCTCTCTCCCCCAAAACCCAAAGGACACCCCTCATGATCTCCCGGACC CCTGAGGTCACATCGTGCTGGTGGAGCTGAGCCGACGAAGACCCCTGAGTCAAGTCTCAACTCGTACGTGAGG GCGTGGAGGTGCATAATCCCAAGCAAAAGCCGCGGGAGGAGCTACAACAGCAAGTACCGTGTGTCAGCGCT CTTCACGCTCTCCACACAGAGTGGCTGAATGCGAAGGATGCAAGTSCAAGTGTCCAAACAAGACCCCTCCCA GCCCCATCGAAGAAAACCATCTCCAAAGCCAAAGGGAGCCCGGAGAACACAGGTGTACGTGCTGCCCCCAT CCGGGAGGAGATGACCAAGAACACAGGTCAAGCTGCTGTGCTGCTCAAAAGGCTTCTATCCACGCAACATCCG TCTGGATGGGAGACGATGGCAGCGGAGAACATACATCTACCTGCGCCCTCCCTGCTGCACTCCGACGCGC TCCTCTTCTCTACAGCAAGCTCACCGTGGACAAGTCCAGATGGCAGGGGAAGCTCTTCTCATGCTGCTCCG TGATGTCAGGCTCTGCACACCACTACACGGAAGTCTCTCTCCCTGTCTCCGGGAAAA (SEQ ID NO: 1141)	
PSMA_P75_F01	EIVLTQSPGTLISVSPGERATLISCRAS QSVRSNLAWYQQKPGQAPRLLIYGAS TRATGIPARFSGSGSETEFTLIISL QSEDAFVYCHQYNDWPPYTFEGQTK LEIRGSEGRSGSGSESKSTGGSQV QLQESGGVQVQPSRLSLSCAASEPT FSTYGMHWVQAPRGKLEWVAFISVD GSNKYYADSVKGRFTI SRDLSKHTLY LQMSLRAEATVAYYCAGRDNSTRLLE WFMDVWGQGTITVVSSEPKSSDKDTH CPDPAPEAAGPSVFLFPKPKDITL MISRTPEVTCVVSVSHEDPEVKENW YVDGVEVHNNAKTRPREEQYNSTRVW SVLTVLHQDWLNGKEYKCKVSNKALP APIEKT I SKAKGQPREPOFVIVLPPSR EEMTKNQVSLTCLVKGFYPSDIAVEW ESNGQPENNYLTWPPVLSDSDGSFELY SKLTVDKSRWQQGNVFCFSVMHEALH HNHYTQKSLSLSPG (SEQ ID NO: 1252)	GAGATCTGCTGACCCACTCTCTCTGAACTCTGTCTGTCTCTGCGGAGAGAGTACCCCTGTCTTTGACAG CCCTCACTCCGTCGGGCTTAACCTGGCTGGTATCGACAGAAACCTGGACAGGCCCTAGACTGCTGATCTPA CGGCGCTTACAGAGGACACAGGATCCCTCGCAGAACTTTCTGGCTCTGGCTCCGGCACCGAGTTTCAACCTTG ACAATCAGTTCCTGCACTCCGAGGACTTCGCCGTGACTACTGCCACAGTACAACAGACTGSCCTCTCTTACA CCTTTGGCCAGGCAACAGCTGGAAATCAAAGCGGCTCCAGGGAAGAGCAGCGGACGCGCAGCAGAG CAAGAGCACCGGCGGACCAAGTTCAGTTCGAAGAATCTGTTGGCGGCTGGTGCAGCTGGAAGATCTGTG AGACTGCTGTGCGGCTCCGCTTCACTTCTCCACTTGGTGAATTCATTGGTCCGACAGGCTCCCGGCA AAGACTGGAATGGGTCCCTTTCATCTCTACAGCGGCTCCAAACAGTACTACGCCCACTCCGTGAAGGGCAG ATTCAACATCTCTCGGACAACCTCAAGCACACCTGTACTGTCAGATGAATCCCTGAGAGCCGAGGATACC CCACCGTCAGCTAGTTCTGAGCCCAAACTAGCGCAAAAACCTCAACATCATGATGTTGTGGGCGCAGGAA AGCAGAGGGGACCGTCAGTCTTCTCTTCCCCAAAACCAAGACACCCCTCATGATCTCCCGGACCCCT GAGTTCACATGCTGTGGTGGCTGAGCGCAGCAACCCGTGAGTCAAGTCAACTGATGTTGTCAGCGCGG TGGAGGTGCATAATGCCAAGACAAGCGCGGAGGAGCTATCAACAGCACTACCTGTGTGTCAGCGTCT CACCGTCTGCACAGGACTGGCTGAATGGCAGGATPACAACTGCAAGTCTCCAAACAAGCCCTCCAGCG CCCATCGAAGAAACCATCTCCAAAGCCAAAGGACGCCCCGAGAACACACAGGTGTAGTGTGTGCCCCCATCCC GGGAGGAGATGACCAAGAACAGGTCAAGCTGCTGTGCTGTGCTCAAGGCTTCTATCCAGGACATCCGCT GGAGTGGAGAGCAATGGCAGCGCGGAGAACAACTACTCACTGGCTCCCGTGTGTCATCCGAGCTCC TTCTTCTCTACAGCAACTCACCTGGACAACTCTAGATGGCAGCGGGGAACCTCTTCTCATGCTGCTCGTGA TGCAATGAGCTCTGCACAACCACTACACGCGAGAAGGCTCTCTCTGCTCTCCGGGT (SEQ ID NO: 1502)	
PSMA_P70_F02	QSVLTQPPSASGTGQQRVITISCSGSS SNI GSNVTNVWYQQLPGTAPKLLIYSS NQRPSGVPDRFSGSKGTSASLISG LQSEADAYYCAAMDLSLNGVWFGG TKLTVLGGSEKSGSGSGSKSTGGS FVOLLFSGPGLVKPSETLSITCTVSG	CAGTCCGCTGCTGACCCAGCCTCTCTTCTGCTTCTGGAACACCTGCGCAGAGAGTGAACATCTCTGCTCCGGCT CCCTCTCAACATCGGCTCTAACACCGTGAATCGGTATCAGCAGCTGCCCGCACAGCCCTTAAACTGCTGAT CTACTCTTCCAAACAGCGGCTTCTGGCTGCCGATAGATTTCTGGCTCAAAGTCTGGACCTCCGCTAGC CTGGCTATTTCTGGCTCTGAGTCTGAGCAGGAGCCGATCTACTCTGCTGCGCTGGATGATTTCTCTGAACG GCGTGTGTTTGGCGGAGCAACAAATTGACATCTTTTGGCGCTCCGAGGCAAGACAGCGGACGCGGCGAG CGCAGAGCAAGCCGCGGACGCAAGTCTCAGCTCTGGAACTTGCACTGCGCTGAGCTGGGCTCAAGCTTCCGAG FVOLLFSGPGLVKPSETLSITCTVSG	

	GSISYYWSWIRQFAGKGLEWIGRIY SSGSTNYNPSLKSRVTMSVDTSKNQF SLKLSVTAADTAVIYCAKVGVMFGA FDIWQQTMTVTVSSEPKSDKTHTCP PCFAPEAAGGSPSVLFFPKPKDTIMI SRPEVTCVVSVSHEDPEVKFNWIV DGEVHNARTKPREEQYNSTYRVVSV LTVLHQDWLNGKEYKCKVSNKALPAP IEKTISKARGQPREPPQVVLPSPREE MTKNQVSLCLVKGFPYPSDIAVEWES NGOPENNYLTWPPVLDSDGSFFLYSK LTVDKSRWQQGNVFSQSVNHEALNH YTQKSLSLSPG (SEQ ID NO: 1264)	ACACTGTCTCTGACCTGTACCGTGTCCGGCGGCTCCATCATCTCTACTACTGTCTCTGGATCAGACAGCCTG CCGGCAARAGGACTGGAATGGATCGGCAGAAATCTACTCCTCCGGCAGCACCACTACAAACCCACGCTGAAGTC CCGCGTGACCAATGTCTGTGGACACCTCCAAGAACCAAGTTCTCCCTGAAGCTGTCTCTGTGACCGCCGCTGAT ACCGCTGTGTACTACTGGCGCTAAAGTCGGAGTGTGGCGCTTGGATATCTGGGACAGGGCACAAATGG TCACCGTGTCTCT (SEQ ID NO: 1503)
--	---	--

EXAMPLE 3: CHARACTERIZATION OF ANTI-PSMA ANTIBODIES**EXAMPLE 3.1: ANALYTICAL CHARACTERIZATION OF PURIFIED ANTI-PSMA ANTIBODIES**

[00586] The protein concentration for each purified mAb was determined by measuring
5 the absorbance at 280 nm on a NANODROP1000 spectrophotometer or TRINEAN
DROPSense96 multichannel spectrophotometer and calculated using the extinction
coefficient based on the amino acid sequence. SE HPLC of the purified antibodies was
performed by running samples on a TOSOH TSKgel BioAssist G3SWxl column, in 0.2 M
Na Phosphate pH 6.8 at 1 mL/min on a Waters Alliance HPLC for 20 min. The column
10 effluent was monitored by absorbance at 280 nm. Bispecific antibodies (bsAbs) were
analyzed by Intact Mass Analysis to determine appropriate formation of heterodimers.

EXAMPLE 3.2: ANTI-PSMA ANTIBODIES BIND PSMA WITH HIGH AFFINITY

[00587] The binding affinity of selected anti-PSMA antibodies to the recombinant human,
15 cynomolgus or mouse PSMA ECD (PSMW39, PSMW1 and PSMW29, respectively) was
determined by surface plasmon resonance (SPR) using a BIACORE 8K instrument (ELN
PSMA-00702). The antibodies were captured on a goat anti-Fc antibody-modified C1 chip
and titrated with 3-fold serial dilutions of PSMA antigen spanning concentrations of 1 nM to
11.1 nM. The association and dissociation were monitored for 3 and 15 minutes,
20 respectively, using a flow rate of 50 μ L/min. Raw binding data was referenced by
subtracting the analyte binding signals from blanks and analyzed using a 1:1 Langmuir
binding model using the Biacore Insight evaluation software to obtain the kinetics, which
were used to calculate the binding affinity. The kinetic parameter of binding of selected
antibodies are shown in Table 13. The anti-PSMA were captured using an anti-human Fc
25 antibody and the antigens were injected in solution. The anti-PSMA antibodies were found
to bind both human PSMA and cyno PSMA with picomolar to nanomolar affinities.

Table 13: Affinities (K_D) for the interaction of anti-PSMA antibodies with Human, cyno or mouse PSMA as obtained by the BIAcore (SPR) method.

Antibody	Human, K _D (M)	Cyno, K _D (M)	Mouse, K _D (M)
PSMB889	≤2.27E-11	Poor fit	Low/no binding
PSMB890	poor fit	Low/No binding	Low/No binding
PSMB891	≤2.53E-10	≤2.10E-10	Low/No binding
PSMB892	3.22E-09	5.67E-09	4.04E-10
PSMB893	Low/No binding	2.60E-07	Low/No binding
PSMB894	No capture	No capture	No capture
PSMB895	7.04E-10	2.83E-09	Low/No binding
PSMB896	≤8.76E-11	6.58E-10	1.36E-09
PSMB897	1.28E-09	3.93E-09	Low/No binding
PSMB898	3.33E-10	1.19E-09	Low/No binding
PSMB899	1.32E-10	1.47E-09	Low/No binding

EXAMPLE 3.3: THERMAL STABILITY OF ANTI-PSMA ANTIBODIES

[00588] The thermal stability (conformational stability) of the anti-PSMA antibodies was determined by nanoDSF method using a Prometheus instrument. Measurements were made by loading sample into 24 well capillary from a 384 well sample plate. Duplicate runs were performed. The thermal scans span from 20°C to 95°C at a rate of 1.0°C/minute. The data was processed to obtain integrated data and first derivation analysis for 330 nm, 350 nm, Ratio 330/350, and scatter data from which thermal transitions, onset of unfolding, T_m and Tagg were obtained. Thermal unfolding parameters, T_m or midpoint temperature of thermal unfolding and Tagg or aggregation temperature of selected anti-PSMA antibodies are shown in Table 14. PSMB889, PSMB890, PSMB891, PSMB892, PSMB895, PSMB896, PSMB897 and PSMB898 showed similar temperatures of unfolding between 68 and 69.9°C. PSMB895, PSMB896, PSMB897 and PSMB898 showed the highest temperature of unfolding ranging from 80.3°C to 94.7°C

Table 14: Thermal stability data for anti-PSMA antibodies as obtained using a nanoDSF instrument.

Antibody	T _{agg}	T _{m1}	T _{m2}
PSMB889	68.6°C	69.1°C	
PSMB890	68.6°C	68.5°C	
PSMB891	75.2°C	69.2°C	76.0°C
PSMB892	69.0°C	68.0°C	
PSMB893	68.3°C	60.7°C	67.8°C
PSMB894	68.5°C	53.8°C	66.2°C
PSMB895	84.7°C	69.3°C	84.4°C

PSMB896	80.3°C	69.9°C	81.0°C
PSMB897	85.2°C	68.6°C	86.1°C
PSMB898	83.2°C	69.1°C	83.1°C
PSMB899	78.0°C	69.0°C	79.1°C

EXAMPLE 3.4: BINDING OF ANTI-PSMA ANTIBODIES ON PSMA+ CELLS

[00589] Selected anti-PSMA antibodies were evaluated for binding to commercial prostate cancer cell lines. C4-2B cells were originally developed at MD Anderson and are derived from LNCaP FGC grown *in vivo* and metastasize to bone marrow (Thalmann, et al 1994, Cancer Research 54, 2577-81). 22Rv1 cells are a human prostate carcinoma epithelial cell line derived from a xenograft that was serially propagated in mice after castration-induced regression and relapse of the parental, androgen-dependent CWR22 xenograft. 22RV1 and C4-2B cells were plated at 50,000 cells per well in 50 μ l of assay medium (RPMI, 10% HI FBS) in V bottom plates. Serial dilutions of antibodies were prepared in assay medium with 50 μ l of antibody dilutions added to the plates containing cells. The plates were incubated for 60 min at 37°C at which time 100 μ l of staining buffer (Becton Dickinson Cat#554657) was added to all wells of each plate. The plates were centrifuged at 300 X G for 5 minutes and the medium was removed from the wells. 200 μ l of staining buffer was added to all wells of each plate. The plates were centrifuged at 300 X G for 5 minutes and the medium was removed from the wells. 50 μ l of 2 μ g/ml AlexaFluor647-labeled goat anti-human Fc was added to all wells of the plates and the plates were incubated for 30 minutes at 4°C. 150 μ l of staining buffer was added to all wells of each plate. The plates were centrifuged at 300 X G for 5 minutes and the medium was removed from the wells. 200 μ l of running buffer (Staining buffer plus 1 mM EDTA, 0.1% pluronic acid) was added to all wells of the plates. The plates were centrifuged at 300 X G for 5 minutes and the medium was removed from the wells. 30 μ l running buffer was added to all wells with cells and the plates were analyzed on the IQUE PLUS instrument (Sartorius). Briefly, cells were gated on a FCS vs. SSC gate to eliminate cellular debris, then the cell populations were gated on singlet cells. Antibody binding was assessed in the red laser channel. Signal (Mab plus secondary antibody) to background (secondary antibody only) ratios were calculated for each plate and the resultant data was plotted vs. MAB concentration in GeneData Screener using 4 parameter curve fitting to generate EC50 values. Table 15 shows EC50 values of anti-PSMA antibodies binding to PSMA+ cell lines.

Table 15: EC50 values of anti-PSMA antibodies binding to PSMA-expressing cell lines 22RV1 and C4-2B measured by flow cytometry assays.

Antibody	22RV1 EC50 [M]	C4-2B EC50 [M]
PSMB889	9.80E-11	4.42E-10
PSMB890	1.71E-09	9.41E-10
PSMB891	5.54E-10	1.99E-09
PSMB892	6.00E-08	2.24E-08
PSMB893	6.00E-08	6.00E-08
PSMB894	6.75E-09	5.35E-08
PSMB895	1.71E-09	5.09E-09
PSMB896	1.24E-09	2.58E-08
PSMB897	2.26E-09	4.76E-09
PSMB898	2.07E-09	6.53E-09
PSMB899	1.02E-09	5.58E-09

EXAMPLE 4: GENERATION OF ANTI-CD3 ANTIBODIES

5 EXAMPLE 4.1: GENERATION OF CD3B376 and CD3B450

[00590] Ablexis AlivaMab or OMT transgenic mice can be used for the production of antibodies. The generation of anti-CD3 antibody CD3B376 and CD3B450 has been described in US Publ. No. 20200048349, which is incorporated by reference in its entirety. CD3B376 and CD3B450 have the same CDR sequences as shown in Tables 16-21 below.

10 EXAMPLE 4.2: GENERATION OF CD3W245

[00591] Anti-CD3 CD3W245 was generated using the Ablexis Transgenic mice technology. Ablexis mice were immunized with TRCW5 (SEQ ID NO:1142). TRCW5 is comprised of the extracellular region of CD3δ fused by a 26 amino acid linker to the extracellular region of CD3ε. This polypeptide has at its C-terminus a human IgG1 Fc domain with a C-terminal Avi-tag used for site-specific biotinylation).

[00592] SEQ ID NO:1142, TRCW5

FKIPIEELEDVRVFNCSITWVEGTVGTLTLLSDITRLDLGKRILDPGRIYRCNGTDIYKDKESTVQV
HYRMGSADDAKKDAKKDDAKKDDAKKDGSDGNEEMGGITQTPYKVSISGTTVILTCPOYPGSEILW
QHNDKNIGGEDDDKNIGSDEDHLSLKEFSELEQSGYYVCYPRGSKPEDANFYLYLRARVSPSPAPE
LLGGPSVFLFPPKPKDTLMISRTEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY
RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSRDELTKNQVSLTCL
LVKGFPYSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVVFSCVMHEALHN
HYTQKSLSLSPGKGGGLNDIFEAQKIEWHE

[00593] Mice were immunized twice weekly for the duration of 7 weeks. On day 42, mice were boosted for hybridoma fusion by administration of 50 µg TRCW5 and 50 µg CD40 mAb spread over 8 sites, including 6 subcutaneous and 2 intradermal injections. For a final boost, mice received 20 µL injections of Jurkat cells, a T cell line which endogenously expresses the T cell receptor complex, including CD3ε (Schneider et al (1977) Int. J. Cancer, 19 (5): 621-6), at 4.74×10^7 cells/mL.

[00594] Mice hybridoma were assayed by ELISA for binding to TRCW5 using TRCW5 either non-specifically immobilized on the plate (ELISA, Thermo cat. # 34022) or by streptavidin conjugation to biotinylated-TRCW5 (SPARCL ELISA, Lumigen). The two assay formats resulted in 426 hits (264 hits from ELISA, 194 from SPARCL ELISA, 70 hits were identified in both assays). Of these 426 initial hits, 49 ELISA and 32 SPARCL ELISA hits were confirmed. The hybridoma fusions corresponding to the positive binders were refed and tested for their abilities to bind Jurkat cells by flow cytometry. Three antibodies, clone 003_F12, clone 036_E10 and clone 065_D03, showed significant binding to Jurkat cells, endogenously expressing CD3. Next, these three clones were screened for their abilities to bind primary human and cyno T cells.

[00595] Clone 065_D03 was selected for further development. The variable region of the Clone 065_D03 was cloned into an IgG1 backbone, resulting in the antibody termed CD3B815.

[00596] SEQ ID NO: 1143, CD3B815 Heavy Chain

EVQLVESGGGLVKPGGSLRLSCAASGFTFSRYNMNVVRQAPGKGLEWVSSI STSSNYIYYADSVKGR
FTFSRDNAKNSLDLQMSGLRAEDTAIYYCTRGWGPFDYWGQGLTVTVSSASTKGPSVFPLAPSSKST
SGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNV
NHKPSNTKVDKKVEPKCDKHTCTPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDSHE
DPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI
SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGS
FFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

[00597] SEQ ID NO: 1144, CD3B815 Light Chain

DILLTQSPGILSVSPGERVSFSCRARQSIGTAIHVYQQRNGSPRLLIKYASESISGIPSRFSGSGS
GTDFTLTINSVESEDIADYYCQQSNSWPYTFGGGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVC
LLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTLSSTLTLSKADYEKHKVYACEVTHQGL
SSPVTKSFNRGEC

[00598] The light chain (LC) of the v-region of CD3B815 was humanized in scFv format. Briefly, the LC from CD3B815 was grafted onto the human IGKV1-39*01-IGKJ2*01

germline and two positions (Y49K and L78V) were identified for human to mouse back mutations. This resulted in variants, having either Y49K, L78V, or both Y49K and L78V. The LC also presented a risk of deamination because of a NS motif at positions 92-93. . These modifications resulted into an antibody termed CD3W245 scFv generated by grafting the CDRs of CD3B815 onto IGKVID-39*01 and introducing the mutations Y49K and N92G into the VL.

EXAMPLE 4.3: GENERATION OF CD3B2030 AND CD3B2030-N106A

[00599] The publicly available mouse Cris7 antibody, specific to human CD3ε (Alberola-Ila, J. et al., J Immunol 146, 1085-1092 (1991)) was used to generate additional exemplary CD3 binding domain provided herein. Murine Cris-7 was humanized in the single-chain fragment variable-domain (scFv) format. Two human heavy variable-domain (Hv) germline sequences and two human light variable-domain (Lv) germline sequences were selected for the antibody humanization: IGHV1-69*02-IGHJ1-01 and IGHV5-10-1*01-IIGHJ1-01 for Hv and IGKV3-11*02-IGKJ4-01 or the IGKV1-39*01-IGKJ4-01 germline for Lv. The CDR-grafted sequences were generated by grafting murine Cris7 CDRs onto these sequences while limiting the number of back mutations to enhance stability. The CDR-grafted v-regions were then expressed in E. coli in scFv format in both the heavy chain-linker-light chain (HL) and in the light chain-linker-heavy chain (LH) orientations. A matrix of Hv and Lv pairings was evaluated in scFv-format in both orientations of Lv followed by Hv or Hv followed by Lv and with a flexible linker between the variable domains. The IGHV1-69*02-IGHJ1-01 heavy chain germline with IGKV3-11*02-IGKJ4-01 light chain germline grafted construct in the heavy-light orientation exhibited the best expression, binding profile, and potential differentiation, and was chosen for human framework optimization.

[00600] Several sites in the VH, which may influence stability of the molecule were identified and thus selected for library-based mouse back-mutagenesis. In one VH library, 4 sites (M48I, A60N, V67A, and I69L - Kabat numbering) were mutated in binary libraries and R94 (Kabat numbering) was mutated to S, V, L, K, T, R, I, or Y generating a total of 128 variants. In a second library, 9 sites (K12A, V20M, R38K, M48I, A60N, R66, V67A, I69L, and R94S - Kabat numbering) were mutated in a binary library generating a total of 512 variants. Analogously, two libraries were generated for the VL sequence, by identifying sites which may influence stability of the molecule. In library 1, no changes were made to the LC. In library 2, 11 sites were selected for mouse back-mutagenesis in binary fashion

(L11M, L13A, A19V, L21M, Q42T, A43S, L46R, L47W, I58V, F71Y, and L78M) for a total of 2048 variants.

5 [00601] Clones exhibiting binding to CD38c greater than the murine parent were selected for sequencing and exposed to additional assay including titration, thermal stress and cell binding. Cell binding was performed against Pan T-cells and Jurkat-CD3-negative cells to look for increase in non-specific binding when thermally stressed. Molecules exhibiting an increase in binding to the negative cell line when thermally stressed were not chosen for additional characterization. Four humanized matrixed clones retained binding after 65° C heat shock and were selected. These clones include CD3B2030, whose CDR, VH and VL
10 sequences are included in the tables below.

[00602] CD3B2030 contains an NG motif within the Heavy Variable CDR3 (HCDR3). To mitigate this Post Translational Modification (PTM) risk, which could negatively impact potency of the T cell re-directing function, the N106A mutation was introduced using standard molecular biology techniques, generating the CD3B2030-N106A variant.

15 [00603] Sequences of representative CD3 antibodies are provided in Tables 16-21. Table 16 provides VH amino acid sequences, VL sequences, heavy chain sequences, and light chain sequences of various CD3 antibodies. VH CDR and VL CDR sequences are provided in Table 17 (Kabat), Table 18 (Chothia), Table 19 (AbM), Table 20 (Contact), and Table 21 (IMGT). Numbers beneath each sequence in the Tables denote corresponding SEQ ID NOs.

Table 16. CD3 Antibody VH, VL, HC and LC Amino Acid Sequences

#	Protein Name (Target)	VH AA sequence	VL AA sequence	Heavy Chain AA sequence	Light Chain AA sequence
25	CD3B376 +K477 (CD3) AKA CD3B2197	QVQLQSGGPRIVR PSQTLSTCAISG DSVFNNAAAWSWI RQSPSRGLEWIGR TYYSRKWLYDYAV SVKSRITVNEPDS RNQFTIQLNSVTP EDTALYYCARGYS SSFDYWGQGTLVTVSS	QSALTQPASVSGSPGQ SITISCTGTSNIGTY KEFVSWYQQHPDKAPKV LLYEVSKRPSGVSSRF SGSKSGNTASLTISGL QAEDQADYHCVSYAGS GTLFSGGGLKLTVL	QVQLQSGGPRIVRPSQTLSLTCAISGDSVFNNAAAWSWIRQ SPSRGLEWLGRTYYRSKWLYDYAVSVKSRITVNPDTSRNQF TLQLNSVTPEDTALYYCARGYSSSFDYWGQGLTVTVSSAST KGPSVPEPLAPSSKSTSGGTAALGCLVKDYFPEPTVYSWNSG ALTSGVHTFPAVLQSSGLYSLSSVATVPSSSLGTQYICNV NHKPSNTKVDKKVEPKSCDKTHCTCPCPAPAAAGGSVFLE PPKPKDTLMI SRTPEVTCVAVSVSHEDPEVKFNWYVDGVEV HNAKTRPREQYNSTYRVSVLTVHLQDNLNGKEYCKVSN KALPAPIEKTISKAKGQPREPQVYTLPPSRREMTKNQVSLT CLVKGFPYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFLLY SKLTVDKSRWQQGNVFPSCSVNHEALHNHYTQKSLSLSPGK	QSALTQPASVSGSPGQSITISC TGTSSNIGTYKEFVSWYQQHPDK APKVLLEYVSKRPSGVSSRFSG SKSGNTASLTISGLQAEDQADY HCVSYAGSGTLLFSGGGLKLTVL GQPKAAPSVTLFPPSSEELQAN KATLVCLISDFYPGAVTVAMKA DSSPVKAGVETTTTPSKQSNKKY AASSYLSLTPEQWKHRSYSQQ VTHEGSTVEKTVAPTECS
26	CD3B376 -K477 (CD3) AKA CD3B891	QVQLQSGGPRIVR PSQTLSTCAISG DSVFNNAAAWSWI RQSPSRGLEWIGR TYYSRKWLYDYAV SVKSRITVNEPDS RNQFTIQLNSVTP EDTALYYCARGYS SSFDYWGQGTLVTVSS	QSALTQPASVSGSPGQ SITISCTGTSNIGTY KEFVSWYQQHPDKAPKV LLYEVSKRPSGVSSRF SGSKSGNTASLTISGL QAEDQADYHCVSYAGS GTLFSGGGLKLTVL	QVQLQSGGPRIVRPSQTLSLTCAISGDSVFNNAAAWSWIRQ SPSRGLEWLGRTYYRSKWLYDYAVSVKSRITVNPDTSRNQF TLQLNSVTPEDTALYYCARGYSSSFDYWGQGLTVTVSSAST KGPSVPEPLAPSSKSTSGGTAALGCLVKDYFPEPTVYSWNSG ALTSGVHTFPAVLQSSGLYSLSSVATVPSSSLGTQYICNV NHKPSNTKVDKKVEPKSCDKTHCTCPCPAPAAAGGSVFLE PPKPKDTLMI SRTPEVTCVAVSVSHEDPEVKFNWYVDGVEV HNAKTRPREQYNSTYRVSVLTVHLQDNLNGKEYCKVSN KALPAPIEKTISKAKGQPREPQVYTLPPSRREMTKNQVSLT CLVKGFPYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFLLY SKLTVDKSRWQQGNVFPSCSVNHEALHNHYTQKSLSLSPG	(see above)
27	CD3B450 +K477 (CD3) AKA CD3B2200	QVQLQSGGPRIVR PSQTLSTCAISG DSVFNNAAAWSWI RQSPSRGLEWIGR TYYSRKWLYDYAV SVKSRITVNEPDS RNQFTIQLNSVTP EDTALYYCARGYS	QSALTQPASVSGSPGQ SITISCTGTSNIGTY KEFVSWYQQHPDKAPKV LLYEVSKRPSGVSSRF SGSKSGNTASLTISGL QAEDQADYHCVSYAGS GTLFSGGGLKLTVL	QVQLQSGGPRIVRPSQTLSLTCAISGDSVFNNAAAWSWIRQ SPSRGLEWLGRTYYRSKWLYDYAVSVKSRITVNPDTSRNQF TLQLNSVTPEDTALYYCARGYSSSFDYWGQGLTVTVSSAST KGPSVPEPLAPSSKSTSGGTAALGCLVKDYFPEPTVYSWNSG ALTSGVHTFPAVLQSSGLYSLSSVATVPSSSLGTQYICNV NHKPSNTKVDKKVEPKSCDKTHCTCPCPAPAAAGGSVFLE PPKPKDTLMI SRTPEVTCVAVSVSHEDPEVKFNWYVDGVEV HNAKTRPREQYNSTYRVSVLTVHLQDNLNGKEYCKVSN KALPAPIEKTISKAKGQPREPQVYTLPPSRREMTKNQVSLT CLVKGFPYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFLLY SKLTVDKSRWQQGNVFPSCSVNHEALHNHYTQKSLSLSPG	QSALTQPASVSGSPGQSITISC TGTSSNIGTYKEFVSWYQQHPDK APKVLLEYVSKRPSGVSSRFSG SKSGNTASLTISGLQAEDQADY HCVSYAGSGTLLFSGGGLKLTVL GQPKAAPSVTLFPPSSEELQAN KATLVCLISDFYPGAVTVAMKA DSSPVKAGVETTTTPSKQSNKKY

		SSFDYWGQGLVT VSS	915	916	917	918
28	CD3B450 -K477 (CD3) AKA CD3B2186	QVQLQSGPGGLVK PSQTLTLTCAISG DSVFENNAAWSWI RQSPSRGLEWIGR TYRRAKWLVDYAV SVKSRITINPDT KNQSLQLNSVTP EDTAVVYCARGYS SSFDYWGQGLVT VSS	QVQLQSGPGGLVK SITISCTGTSSNIGTY KEFVSQYQHPGKAPK MIYEVSKRPSGVSNRE SGSKSGNTASLTISGL QAEDADYVCVSYAGS GTLFEGGGLKTLVL	CLVKGYPSPDI SKLTVDKSRWQGNV QVQLQSGPGGLVK SPSRGLEWIGR SLQLNSVTP KGPSVFLP ALTSVHTFP NHKPSNTK PPKPKDTLMI HNAKTPREEQ KALPAPIEKT CLVKGYPSPDI SKLTVDKSRWQGNV	CLVKGYPSPDI SKLTVDKSRWQGNV QVQLQSGPGGLVK SPSRGLEWIGR SLQLNSVTP KGPSVFLP ALTSVHTFP NHKPSNTK PPKPKDTLMI HNAKTPREEQ KALPAPIEKT CLVKGYPSPDI SKLTVDKSRWQGNV	CLVKGYPSPDI SKLTVDKSRWQGNV QVQLQSGPGGLVK SPSRGLEWIGR SLQLNSVTP KGPSVFLP ALTSVHTFP NHKPSNTK PPKPKDTLMI HNAKTPREEQ KALPAPIEKT CLVKGYPSPDI SKLTVDKSRWQGNV
29	CD3W245 (-K477) (CD3) AKA CD3W245 or CD3B2183	EVQLVESGGGLVK PGSLRLSCAASG FTFSRYNMNVVRQ APGKLEWVSSIS TSSNVIYYADSVK GRFTFSRDNAKNS LDLQMSGLRAEDT AIYYCTRGWGPFDD YWGQGLTVTVSS	EVQLVESGGGLVK DRVTITCRARQSIGTA IHWYQOKPGKAPKLLI KYASEISGVPSRFSG SGSGTDFTLTISLQ EDFATYYCQSGSWPY TFGQGTGLEIK	EVQLVESGGGLVK GKGLEWVSSI MSGLEAEDTA VFPLAPSSK GVHTTTPAVLQ SNTKVDKKV KDTLMI TKPREEQ APIEKT GFYPSDI VDKSRWQGNV	EVQLVESGGGLVK GKGLEWVSSI MSGLEAEDTA VFPLAPSSK GVHTTTPAVLQ SNTKVDKKV KDTLMI TKPREEQ APIEKT GFYPSDI VDKSRWQGNV	EVQLVESGGGLVK GKGLEWVSSI MSGLEAEDTA VFPLAPSSK GVHTTTPAVLQ SNTKVDKKV KDTLMI TKPREEQ APIEKT GFYPSDI VDKSRWQGNV
30	CD3W245 +K477 (CD3) AKA N/A	EVQLVESGGGLVK PGSLRLSCAASG FTFSRYNMNVVRQ APGKLEWVSSIS TSSNVIYYADSVK GRFTFSRDNAKNS LDLQMSGLRAEDT AIYYCTRGWGPFDD YWGQGLTVTVSS	EVQLVESGGGLVK DRVTITCRARQSIGTA IHWYQOKPGKAPKLLI KYASEISGVPSRFSG SGSGTDFTLTISLQ EDFATYYCQSGSWPY TFGQGTGLEIK	EVQLVESGGGLVK GKGLEWVSSI MSGLEAEDTA VFPLAPSSK GVHTTTPAVLQ SNTKVDKKV KDTLMI TKPREEQ APIEKT GFYPSDI VDKSRWQGNV	EVQLVESGGGLVK GKGLEWVSSI MSGLEAEDTA VFPLAPSSK GVHTTTPAVLQ SNTKVDKKV KDTLMI TKPREEQ APIEKT GFYPSDI VDKSRWQGNV	EVQLVESGGGLVK GKGLEWVSSI MSGLEAEDTA VFPLAPSSK GVHTTTPAVLQ SNTKVDKKV KDTLMI TKPREEQ APIEKT GFYPSDI VDKSRWQGNV
			983	984	985	986

31	CD3E2030 (CD3)	QVQLVQSGAEVKK PGSSVKVSKRAG YTFRTSTMHWKQ APGQGLEWIGYIN PSSAYTNYNQKFK GRVTLTADKSTST AYMELSSLRSEDT AVYYCASPVHYD YNGFPYWGQGTIV TVSS	EIVLTQSPATLSASPG ERVTLSCSASSSVSYM NYYQQKPGQAPRRWIY DSKSLASGVPAREFSGS GSGRDYTLTISSLEPE DFAVYYCQQWSRNPFT FGGGRKVEIK	EIVLTQSPATLSASPGERVTLSCSASSSVSYMNYQQKPGQ APRRWIYDSSKLASGVPAERFSGSGSGRDYTLTISSLEPEDE AVYYCQQWSRNPFTFGGGRKVEIKGSGEGSKSGSGSEKST GGSQQLVQSGAEVKKPGSSVKVSKRAGYTFRTSTMHWK QAPGQGLEWIGYINPSSAYTNYNQKFGRVTLTADKSTSTA YMELSSLRSEDTAVYYCASPVHYDNGFPYWGQGTIVTVS SEPKSDKTHRCPPCPAPEAAGGPSVFLFPKPKDTIMI SR TPEVTCVVSVSHDEVEKFNWYVDGVEVHNAKTKPREEQY NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTIS KAKGQPREPQVYITLPPSRREMTKNQVSLWCLVKGFYPSDIA VEWESNGQPENNYKTTTPVLDSDGSEFLYSLKLTVDKSRWQQ GNVFCSSVMHEALHNHYTQKSLSLSPGK	NA	1466
32	CD3B2030 -N106A (CD3)	QVQLVQSGAEVKK PGSSVKVSKRAG YTFRTSTMHWKQ APGQGLEWIGYIN PSSAYTNYNQKFK GRVTLTADKSTST AYMELSSLRSEDT AVYYCASPVHYD YAGFPYWGQGTIV TVSS	EIVLTQSPATLSASPG (see above)	EIVLTQSPATLSASPGERVTLSCSASSSVSYMNYQQKPGQ APRRWIYDSSKLASGVPAERFSGSGSGRDYTLTISSLEPEDE AVYYCQQWSRNPFTFGGGRKVEIKGSGEGSKSGSGSEKST GGSQQLVQSGAEVKKPGSSVKVSKRAGYTFRTSTMHWK QAPGQGLEWIGYINPSSAYTNYNQKFGRVTLTADKSTSTA YMELSSLRSEDTAVYYCASPVHYDNGFPYWGQGTIVTVS SEPKSDKTHRCPPCPAPEAAGGPSVFLFPKPKDTIMI SR TPEVTCVVSVSHDEVEKFNWYVDGVEVHNAKTKPREEQY NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTIS KAKGQPREPQVYITLPPSRREMTKNQVSLWCLVKGFYPSDIA VEWESNGQPENNYKTTTPVLDSDGSEFLYSLKLTVDKSRWQQ GNVFCSSVMHEALHNHYTQKSLSLSPGK	NA	1464
						1505
						1455

Table 17. CD3 Antibody Kabat CDR Amino Acid Sequences

#	Protein Name	HC Kabat CDR1	HC Kabat CDR2	HC Kabat CDR3	LC Kabat CDR1	LC Kabat CDR2	LC Kabat CDR3
25	CD3B376+ K477	NNNAWS 817	RTYYRSKWLYDYAVSVK S 818	GYSSSEFY 819	TGTSSNIGTYKFVS 820	EVSKRPS 821	VSYAGSGTLL 822
26	CD3B376- K477	NNNAWS 817	RTYYRSKWLYDYAVSVK S 818	GYSSSEFY 819	TGTSSNIGTYKFVS 820	EVSKRPS 821	VSYAGSGTLL 822
27	CD3B450+ K477	NNNAWS 817	RTYYRSKWLYDYAVSVK S 818	GYSSSEFY 819	TGTSSNIGTYKFVS 820	EVSKRPS 821	VSYAGSGTLL 822
28	CD3B450- K477	NNNAWS 817	RTYYRSKWLYDYAVSVK S 818	GYSSSEFY 819	TGTSSNIGTYKFVS 820	EVSKRPS 821	VSYAGSGTLL 822
29	CD3W245+ K477	RNNMN 817	SISTSSNYIYADSVKG S 818	GWGPFY 819	PARQSIGTAIH 820	YASESIS 821	QQSGSWPYT 822
30	CD3W245- K477	1 RNNMN	954 SISTSSNYIYADSVKG	955 GWGPFY	956 PARQSIGTAIH	957 YASESIS	958 QQSGSWPYT
31	CD3B2030	1 RSTMH	954 YINPSSAYTNYNQRFQG	955 PQVHYDYGFFY	956 SASSSVSYMN	957 DSSKIAS	958 QQWSRNPPT
32	CD3B2030 -N106A	1467 RSTMH	1468 YINPSSAYTNYNQRFQG	1469 PQVHYDYGFFY	1470 SASSSVSYMN	1471 DSSKIAS	1472 QQWSRNPPT
		1467	1468	1506	1470	1471	1472

Table 18. CD3 Antibody Chothia CDR Amino Acid Sequences

#	Protein Name	HC Chothia CDR1	HC Chothia CDR2	HC Chothia CDR3	LC Chothia CDR1	LC Chothia CDR2	LC Chothia CDR3
25	CD3B376+ K477	GDSVFNN A	YRKS 824	YSSSF 825	TSSNIGTYKF 826	EVS 487	YAGSGTL 828
26	CD3B376-- K477	GDSVFNN A	YRKS 824	YSSSF 825	TSSNIGTYKF 826	EVS 487	YAGSGTL 828
27	CD3B450+ K477	GDSVFNN A	YRKS 824	YSSSF 825	TSSNIGTYKF 826	EVS 487	YAGSGTL 828
28	CD3B450-- K477	GDSVFNN A	YRKS 824	YSSSF 825	TSSNIGTYKF 826	EVS 487	YAGSGTL 828
29	CD3W245+ K477	GTFFSRY 7	STSSNY 960	GWGPF 961	RQSIGTA 962	YAS 963	SGSWPY 964
30	CD3W245-- K477	GTFFSRY 7	STSSNY 960	GWGPF 961	RQSIGTA 962	YAS 963	SGSWPY 964
31	CD3B2030	GYTFTRS 1473	NPSSAY 1474	PQVHYDYNGFP 1475	SSSVSY 1476	DSS 1477	WSPNPP 1478
32	CD3B2030 -N106A	GYTFTRS 1473	NPSSAY 1474	PQVHYDYNGFP 1507	SSSVSY 1476	DSS 1477	WSPNPP 1478

Table 19. CD3 Antibody AbM CDR Amino Acid Sequences

#	Protein Name	HC AbM CDR1	HC AbM CDR2	HC AbM CDR3	LC AbM CDR1	LC AbM CDR2	LC AbM CDR3
25	CD3B376+ K477	GDSVFNNNAAWS	RTYRSKWLYD	GYSSSEFY	TGTSSNIGTYKFVS	EVSKRPS	VSYAGSGTLL
26	CD3B376-- K477	829 GDSVFNNNAAWS	830 RTYRSKWLYD	819 GYSSSEFY	820 TGTSSNIGTYKFVS	821 EVSKRPS	822 VSYAGSGTLL
27	CD3B450+ K477	829 GDSVFNNNAAWS	830 RTYRSKWLYD	819 GYSSSEFY	820 TGTSSNIGTYKFVS	821 EVSKRPS	822 VSYAGSGTLL
28	CD3B450-- K477	829 GDSVFNNNAAWS	830 RTYRSKWLYD	819 GYSSSEFY	820 TGTSSNIGTYKFVS	821 EVSKRPS	822 VSYAGSGTLL
29	CD3W245+ K477	829 GFTFSRYNMN	830 SISTSSNYLY	819 GWGPFYD	820 RARQSIGTAIH	821 YASESIS	822 QQSGSWPYT
30	CD3W245-- K477	13 GFTFSRYNMN	966 SISTSSNYLY	955 GWGPFYD	956 RARQSIGTAIH	957 YASESIS	958 QQSGSWPYT
31	CD3B2030	13 GYTFTRSTMH	966 YINPSSAYTN	955 PQVHYDYGFPY	956 SASSSVSYMN	957 DSSKIAS	958 QQWSRNPPT
32	CD3B2030 -N106A	1508 GYTFTRSTMH	1509 YINPSSAYTN	1469 PQVHYDYGFPY	1470 SASSSVSYMN	1471 DSSKIAS	1472 QQWSRNPPT
		1508	1509	1506	1470	1471	1472

Table 20. CD3 Antibody Contact CDR Amino Acid Sequences

#	Protein Name	HC Contact CDR1	HC Contact CDR2	HC Contact CDR3	LC Contact CDR1	LC Contact CDR2	LC Contact CDR3
25	CD3B376+K477	FNNAAWS	WLGRTYYRSKWLVD	ARGYSSSF	IGTYKFVSWY	VLLYEVSRRP	VSYAGSGTL
		835	836	837	838	839	840
26	CD3B376-K477	FNNAAWS	WLGRTYYPSKWLVD	ARGYSSSF	IGTYKFVSWY	VLLYEVSRRP	VSYAGSGTL
27	CD3B450+K477	FNNAAWS	WLGRTYYRSKWLVD	ARGYSSSF	IGTYKFVSWY	VMIYEVSRRP	VSYAGSGTL
		835	836	837	838	907	840
28	CD3B450-K477	FNNAAWS	WLGRTYYRSKWLVD	ARGYSSSF	IGTYKFVSWY	VMIYEVSRRP	VSYAGSGTL
		835	836	837	838	907	840
29	CD3W245+K477	SRNMN	WVSSISTSSNYIY	TRGWGPF	GTAHWY	LLIKYASESI	QQSGSWPY
		19	972	973	974	975	976
30	CD3W245-K477	SRNMN	WVSSISTSSNYIY	TRGWGPF	GTAHWY	LLIKYASESI	QQSGSWPY
		19	972	973	974	975	976
31	CD3B2030	TRSTMH	WIGYINPSSAYTN	ASPOVHYDNGF	SYNNWY	RWLYDSSKLA	QQWSRNP
		1510	1511	1512	1513	1514	1515
32	CD3B2030-N106A	TRSTMH	WIGYINPSSAYTN	ASPOVHYDNGF	SYNNWY	RWLYDSSKLA	QQWSRNP
		1516	1517	1518	1519	1520	1521

Table 21. CD3 Antibody IMGT CDR Amino Acid Sequences

#	Protein Name	HC IMGT CDR1	HC IMGT CDR2	HC IMGT CDR3	LC IMGT CDR1	LC IMGT CDR2	LC IMGT CDR3
25	CD3B376+ K477	GDSVFNNAA	TYRSKWLY	ARGYSSSFY	SSNIGTYKF	EVS	VSYAGSGTLL
26	CD3B376- K477	GDSVFNNAA	TYRSKWLY	ARGYSSSFY	SSNIGTYKF	EVS	VSYAGSGTLL
27	CD3B450+ K477	GDSVFNNAA	TYRSKWLY	ARGYSSSFY	SSNIGTYKF	EVS	VSYAGSGTLL
28	CD3B450- K477	GDSVFNNAA	TYRSKWLY	ARGYSSSFY	SSNIGTYKF	EVS	VSYAGSGTLL
29	CD3W245+ K477	GFTFSRYN	ISTSSNYI	TRGWGPFY	QSIGTA	YAS	QQSGSWPYT
30	CD3W245- K477	GFTFSRYN	ISTSSNYI	TRGWGPFY	QSIGTA	YAS	QQSGSWPYT
31	CD3B2030	GYTFTRST	INPSSAYT	ASPQVHYDYGFPY	SSVS	DSS	QQWSRNPT
32	CD3B2030 -N106A	GYTFTRST	INPSSAYT	ASPQVHYDYGFPY	SSVS	DSS	QQWSRNPT
		1479	1480	1518	1482	1477	1472

[00604] Exemplary CD3 VH and VL nucleic acid sequences are provided below.

[00605] SEQ ID NO: 1453, VH nucleic acid sequence of CD3B2197 (CD3B376+K477)

CAGGTGCAGCTGCAGCAGTCTGGCCCTAGACTCGTGC GGCCCTCCAGACCCTGTCTCTGACCTGTG
CCATCTCCGGCGACTCCGTGTTCAACAACAACGCCGCCTGGTCCTGGATCCGGCAGTCTCCATCTCG
CGGTCTGGAGTGGCTCGGTGCGACCTACTACCGCTCTAAATGGCTGTACGACTACGCCGTGTCCGTG
AAGTCCCGGATCACCGTGAACCCCTGACACCTCCCGGAACCAGTTCACCCTGCAGCTGAACTCCGTGA
CCCCTGAGGACACCGCCCTGTACTACTGCGCCAGAGGCTACTCCTCCTCCTTCGACTATTGGGGCCA
AGGCACCCTCGTGACCGTGTCTCT

[00606] SEQ ID NO: 1170, VL nucleic acid sequence of CD3B2197 (CD3B376+K477)

CAGTCTGCTCTGACCCAGCCTGCCTCCGTGTCTGGCTCTCCCGGCCAGTCCATCACCATCAGCTGTA
CCGGCACCTCCTCCAACATCGGCACCTACAAGTTCGTGTCTGGTATCAGCAACACCCCGACAAGGC
CCCCAAAGTGCTGTGTACGAGGTGTCCAAGCGGCCCTCTGGCGTGTCTCCAGATTCTCCGGCTCC
AAGTCTGGCAACACCGCCTCCCTGACCATCAGCGGACTGCAGGCTGAGGACCAGGCCGACTACCACT
GTGTGTCTTACGCTGGCTCTGGCACCTGCTGTTTGGCGGAGGCACCAAGCTGACTGTCTCTG

[00607] SEQ ID NO: 1151, VH nucleic acid sequence of CD3B891 (CD3B376-K477)

CAGGTGCAGCTGCAGCAGTCTGGCCCTAGACTCGTGC GGCCCTCCAGACCCTGTCTCTGACCTGTG
CCATCTCCGGCGACTCCGTGTTCAACAACAACGCCGCCTGGTCCTGGATCCGGCAGAGCCCTTCTAG
AGGCCTGGGAATGGCTGGGCCGGACCTACTACCGGTCCAAGTGGCTGTACGACTACGCCGTGTCCGTG
AAGTCCCGGATCACCGTGAACCCCTGACACCTCCCGGAACCAGTTCACCCTGCAGCTGAACTCCGTGA
CCCCTGAGGACACCGCCCTGTACTACTGCGCCAGAGGCTACTCCTCCTCCTTCGACTATTGGGGCCA
GGGCACCCTCGTGACCGTGTCTCT

[00608] SEQ ID NO: 1152, VL nucleic acid sequence of CD3B891 (CD3B376-K477)

CAGTCTGCTCTGACCCAGCCTGCCTCCGTGTCTGGCTCTCCCGGCCAGTCCATCACCATCAGCTGTA
CCGGCACCTCCTCCAACATCGGCACCTACAAGTTCGTGTCTGGTATCAGCAGACCCCGACAAGGC
CCCCAAAGTGCTGTGTACGAGGTGTCCAAGCGGCCCTCTGGCGTGTCTCCAGATTCTCCGGCTCC
AAGTCTGGCAACACCGCCTCCCTGACCATCAGCGGACTGCAGGCTGAGGACCAGGCCGACTACCACT
GTGTGTCTTACGCTGGCTCTGGCACCTGCTGTTTGGCGGAGGCACCAAGCTGACCGTGTCTG

[00609] SEQ ID NO: 1153, VH nucleic acid sequence of CD3B2200 (CD3B450+K477)

CAGGTTCACTGCAGCAGTCTGGCCCTGGACTGGTCAAGCCCTCTCAGACCCTGTCTCTGACCTGTG
CCATCTCCGGCGACTCCGTGTTCAACAACAATGCCGCCTGGTCCTGGATTCGGCAGTCTCCTAGTAG
AGGCCTGGAATGGCTGGGCAGAACCTACTACCGGTCCAAGTGGCTGTACGACTACGCCGTGTCCGTG
AAGTCCCGGATCACCATCAATCCCGACACCTCCAAGAACCAGTTCCTCCTGCAGCTCAACAGCGTGA
CCCCTGAGGATACCGCGGTGTACTACTGTGCCAGAGGCTACTCCTCCTCCTTCGACTATTGGGGCCA
GGGCACACTGGTCACCGTTTCTTCT

[00610] SEQ ID NO: 1154, VL nucleic acid sequence of CD3B2200 (CD3B450+K477)

CAGTCTGCTCTGACCCAGCCTGCTTCTGTGTCTGGCTCTCCCGGCCAGTCCATCACCATCTCTTGTA
CCGGCACCTCCTCCAACATCGGCACCTACAAGTTCGTGTCTGGTATCAGCAGACCCCGCAAGGC
CCCTAAAGTGATGATCTACGAGGTGTCCAAGCGGCCCTCTGGCGTGTCCAACAGATTCTCCGGCTCC
AAGTCCGGCAATACCGCCTCTCTGACAATCAGCGGACTGCAGGCTGAGGACCAGGCCGACTACTACT
GTGTGTCTTACGCTGGCTCTGGCACCTGTTGTTTGGCGGCGGAACAAAGCTGACTGTGTCTG

[00611] SEQ ID NO: 1155, VH nucleic acid sequence of CD3B2186 (CD3B450-K477)

CAGGTGCAGCTGCAGCAGAGCGGCCCGGCCTGGTCAAGCCCAGCCAGACCCTGAGCCTGACCTGCG
CCATCAGCGGCGCAGCGTGTTCACAACAACGCGCCTGGTCCTGGATCCGCCAGAGCCCCAGCCG
CGGCCTGGAGTGGCTGGGCCGCACCTACTACCGCAGCAAGTGGCTGTACGACTACGCCGTGTCCGTG
AAGTCCCGCATCACCATCAACCCCGACACCAGCAAGAACCAGTTCTCCCTGCAGCTGAACAGCGTGA
CCCCGAGGACACCGCGTGTACTACTGCGCCCGCGGCTACAGCAGCAGCTTCGACTACTGGGGCCA
GGGCACCCTGGTCACCGTGTCCAGC

[00612] SEQ ID NO: 1156, VL nucleic acid sequence of CD3B2186 (CD3B450-K477)

CAGTCTGCTCTGACCCAGCCTGCCTCCGTGTCTGGCTCTCCCGGCCAGTCCATCACCATCAGCTGTA
CCGGCACCTCCTCCACATCGGCACCTACAAGTTCGTGTCTGGTATCAGCAGCACCCTCGCAAGGC
CCCCAAGTGATGATCTACGAGGTGTCCAAGCGGCCCTCCGGCGTGTCCAACAGATTCTCCGGCTCC
AAGTCCGGCAACACCGCCTCCCTGACAATCAGCGGACTGCAGGCCGAGGACGAGGCCGACTACTACT
GTGTCTCCTACGCCGCTCTGGCACCTGCTGTTTGGCGGCGGAACAAAGCTGACCGTGCTG

[00613] SEQ ID NO: 1157, VH nucleic acid sequence of CD3B2183 (CD3W245)

GAGGTGCAACTGGTGGAGTCTGGGGGAGGCCTGGTCAAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTCAGTAGATATAACATGAAGTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCATCCATTAGTACTAGTAGTAATTACATATACTACGCAGACTCAGTGAAGGGCCGA
TTCACCTTCTCCAGAGACAACGCCAAGAACTCACTGGATCTGCAAATGAGCGGCTGAGAGCCGAGG
ACACGGCTATTTATTACTGTACGAGAGGCTGGGGGCCTTTTGACTACTGGGGCCAGGGAACCTGGT
CACCGTCTCCTCA

[00614] SEQ ID NO: 1158, VL nucleic acid sequence of CD3B2183 (CD3W245)

GACATACAAATGACACAATCACCCTCTTCTCTTTCTGCAAGCGTTGGCGACCGTGTCACCTACCTT
GTCGAGCCCCGCGAGTCCATAGGTACTGCCATTCACTGGTATCAACAGAAGCCTGGCAAGGCTCCCAA
ACTCCTGATTAAAGTATGCCAGCGAGAGCATTTCCGGCGTACCTTCAAGATTTTCCGGCTCCGGTAGT
GGGACAGATTTCACTCTCACTATATCTAGCCTCCAACCAGAAGATTTGCGCACTTACTACTGTCAAC
AATCAGGTTTCATGGCCTTACACTTTCGGCCAGGGGACAAAATTGGAGATCAAG

[00615] SEQ ID NO: 1519, VH nucleic acid sequence of CD3B2030

CAGGTTCACTGGTTCAGTCTGGCGCCGAAGTGAAGAAACCTGGCTCCTCCGTCAAGGTGTCTTGCA
AGGCTTCCGGCTACACCTTTACCAGATCCACCATGCACTGGGTCAAACAGGCTCCAGGACAAGGCTT
GGAGTGGATCGGCTACATCAACCCAGCTCCGCCTACACCAACTACAACCAGAAATTCAGGGCAGA
GTCACCTCACCAGCCGACAAGTCTACCTCCACCGCTACATGGAACTGTCCAGCCTGAGATCTGAGG
ACACCGCGGTGTACTACTGCGCCAGCCCTCAGGTGCACTACGACTACAACGGCTTCCCTTATTGGGG
CCAGGGCACCCCTGGTTACCGTTTCTTCT

[00616] SEQ ID NO: 1520, VL nucleic acid sequence of CD3B2030

GAGATCGTGCTGACCCAGTCTCCTGCCACACTGTCAGCCTCTCCAGGCGAGAGAGTCAACCTGTCTT
GCTCCGCTTCTCCTCCGTGTCTACATGAAGTGGTATCAGCAGAAGCCCGGCCAGGCTCCTAGACG
GTGGATCTACGACTCCTCCAAGCTGGCCTCTGGCGTCCCTGCCCGCTTTTCCGGCTCTGGCTCTGGC
AGAGACTATACCTGACCATCTCCAGCCTGGAACCTGAGGACTTCGCCGTGTACTACTGCCAGCAGT
GGTCTAGAAACCTCCTACCTTTFGGCGGAGGCACCAAGGTGGAAATCAAG

[00617] SEQ ID NO: 1521, VH nucleic acid sequence of CD3B2030-N106A

CAGGTTCAACTGGTTCAGTCTGGCGCCGAAGTGAAGAAACCTGGCTCCTCCGTCAAGGTGTCTTGCA
AGGCTTCCGGCTACACCTTTACCAGATCCACCATGCACTGGGTCAAACAGGCCCCCTGGACAAGGCTT
GGAGTGGATCGGCTACATCAACCCAGCTCCGCCTACACCAACTACAACCAGAAATTCAGGGCAGA
GTGACCTTGACCGCCGACAAGTCTACCTCCACCGCTACATGGAACTGTCCAGCCTGAGATCTGAGG

ACACCGCCGTGTACTACTGCGCCTCTCCTCAGGTCCACTACGACTACGCCGGCTTTCCTTATTGGGG
CCAGGGCACACTGGTCACCGTTTCTTCT

[00618] SEQ ID NO: 1522, VL nucleic acid sequence of CD3B2030-N106A

GAGATCGTGCTGACCCAGTCTCCTGCCACACTGTCAGCCTCTCCAGGCGAGAGAGTCAACCCTGTCTT
GCTCCGCTTCTCCTCCGTGTCTACATGAAGTGGTATCAGCAGAAGCCCGGCCAGGCTCCTAGACG
GTGGATCTACGACTCTCCAAGCTGGCCTCTGGCGTCCCTGCCCGCTTTTCCGGCTCTGGCTCTGGC
AGAGACTATACCCTGACCATCTCCAGCCTGGAACCTGAGGACTTCGCCGTGTACTACTGCCAGCAGT
GGTCTAGAAACCTCCTACCTTTGGCGGAGGCACCAAGGTGGAAATCAAG

[00619] CD3 VH/VL regions were also engineered as scFvs in either VH-Linker-VL (also termed "HL") or VL-linker-VH (also termed "LH") orientations using the linker of SEQ ID NO: 1419. The VH-Linker-VL or VL-linker-VH scFv molecules binding CD3 were further engineered into a scFv-hinge-CH2-CH3 (also termed scFv-Fc) format comprising Fc silencing mutation (L234A/L235A/D265S) and mutations designed to promote selective heterodimerization.

[00620] Exemplary scFv amino acid sequences of CD3W245 are provided below (LH or HL orientation). In some embodiments, PSMAxCD3 multispecific antibodies provided herein comprise an amino acid sequence of CD3W245-LH. In some embodiments, PSMAxCD3 multispecific antibodies provided herein comprise an amino acid sequence of CD3W245-HL.

[00621] SEQ ID NO: 1186 CD3W245-LH

DIQMTQSPSSLSASVGDRVTITCRARQSIGTAIHWYQQKPGKAPKLLIKYASESISGVPSRFSGSGS
GTDFTLTITSSLPEDFATYYCQQSGSWPYTFGQGTKLEIKGGSEKSSGSGSESKSTGGSEVQLVES
GGGLVKPGGSLRLSCAASGFTFSRYNMNWRQAPGKGLEWVSSISTSSNYIYYADSVKGRFTFSRDN
AKNSLDLQMSGLRAEDTAIYYCTRGWGPFDYWGQGLTVTVSS

[00622] SEQ ID NO: 1187 CD3W245-HL

EVQLVESGGGLVKPGGSLRLSCAASGFTFSRYNMNWRQAPGKGLEWVSSISTSSNYIYYADSVKGR
FTFSRDNAKNSLDLQMSGLRAEDTAIYYCTRGWGPFDYWGQGLTVTVSSGGSEKSSGSGSESKSTG
GSDIQMTQSPSSLSASVGDRVTITCRARQSIGTAIHWYQQKPGKAPKLLIKYASESISGVPSRFSGSGS
GSGTDFTLTITSSLPEDFATYYCQQSGSWPYTFGQGTKLEIK

[00623] Exemplary scFv amino acid sequences of CD3B2030 and CD3B2030-N106A are provided below in the LH orientation. In some embodiments, PSMAxCD3 multispecific antibodies provided herein comprise an amino acid sequence of CD3B2030-LH. In some embodiments, PSMAxCD3 multispecific antibodies provided herein comprise an amino acid sequence of CD3B2030-N106A-LH.

[00624] SEQ ID NO: 1523, CD3B2030-LH scFv amino acid sequence

EIVLTQSPATLSASPGERVTLSCSASSSVSYMNWYQQKPGQAPRRWIYDSSKLASGVPARFSGSGSG
RDYTLTISSLEPEDFAVYYCQQWSRNPPTFGGGTKVEIKGGSEGKSSGSGSESKSTGGSQVQLVQSG
AEVKKPGSSSVKVSCKASGYTFTRSTMHWVKQAPGQGLEWIGYINPSSAYTNYNQKFQGRVTLTADKS
TSTAYMELSSLRSEDTAVYYCASPVHYDYNQFPYWGQGTLVTVSS

[00625] SEQ ID NO: 1524, CD3B2030-N106A-LH scFv amino acid sequence

EIVLTQSPATLSASPGERVTLSCSASSSVSYMNWYQQKPGQAPRRWIYDSSKLASGVPARFSGSGSG
RDYTLTISSLEPEDFAVYYCQQWSRNPPTFGGGTKVEIKGGSEGKSSGSGSESKSTGGSQVQLVQSG
AEVKKPGSSSVKVSCKASGYTFTRSTMHWVKQAPGQGLEWIGYINPSSAYTNYNQKFQGRVTLTADKS
TSTAYMELSSLRSEDTAVYYCASPVHYDYNQFPYWGQGTLVTVSS

[00626] SEQ ID NO: 1525, CD3B2030_N106A-LH scFv nucleotide sequence

GAAATTGTTTGTACCAATCACCTGCCACTCTCTCTGCCTCTCCTGGTGAGCGAGTTACTTTGTTCAT
GTAGCGCATCATCAAGTGTATCTTACATGAAGTGGTACCAACAAAAACCCGGACAGGCACCGTCTG
TTGGATTATGACAGTAGCAAGCTCGCCTCCGGGGTACCCGCAAGATTTCCGGGTGAGGCTCTGGC
AGGGACTATACCTGACAATCAGCAGTCTGGAACCTGAGGACTTTGCTGTGTATTACTGCCAACAGT
GGTCTCGCAACCCCTACTTTTCGGGGGAGGTACAAAGGTAGAAATTAAGGGCGGCTCCGAGGGCAA
GAGCAGCGGCAGCGGCAGCGAGAGCAAGAGCACCGCGCGGCAGCCAAGTGCAACTCGTGCAAGCGGG
GCTGAAGTGAAGAAGCCTGGATCAAGCGTGAAGGTGCAAGCCTCTGGATATACCTTCACTC
GATCAACCATGCATGGGTCAAGCAGGCCCCAGGGCAAGGGCTCGAATGGATAGGATATATTAACCC
AAGTTCTGCCTACCTAATAATCAGAAGTTTCAAGGCCGGGTAACACTTACAGCCGATAAGAGT
ACCTCAACAGCATACATGGAAGTCTTTGCGGAGCGAGGATACCGCTGTGTATTACTGCGCTT
CACCTCAGGTTCACTACGACTACGCTGGATTTCCCTATTGGGGTCAGGGTACACTGGTTACAGTTTC
CTCTG

[00627] In some embodiments, PSMAxCD3 multispecific antibodies provided herein
comprise an amino acid sequence of SEQ ID NO: 1186. In some embodiments, PSMAxCD3
multispecific antibodies provided herein comprise an amino acid sequence of SEQ ID
NO: 1187. In some embodiments, PSMAxCD3 multispecific antibodies provided herein
comprise an amino acid sequence of SEQ ID NO: 1523. In some embodiments, PSMAxCD3
multispecific antibodies provided herein comprise an amino acid sequence of SEQ ID
NO: 1524.

[00628] Other exemplary CD3 DuoBody HC and LC amino acid sequences are provided
below.

[00629] SEQ ID NO: 1167, HC amino acid sequence (DuoBody) of CD3B2183
(CD3W245)

EVQLVESGGGLVKGSSLRSLCAASGFTFSRYNMNWRQAPGKGLEWVSSI STSSNYIYYADSVKGR
FTFSRDNAKNSLDLQMSGLRAEDTAIYYCTRGWGFDFYWGQGLVTVSSASTKGPSVFPLAPSSKST
SGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNV
NHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVSVSHE
DPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI
SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGS
FLLYSKLTVDKSRWQQGNVFSQVMHEALHNHYTQKSLSLSP

[00630] Exemplary CD3 DuoBody HC and LC nucleic acid sequences are provided below.

[00631] SEQ ID NO: 1149, HC nucleic acid sequence (DuoBody) of CD3B2197 (CD3B376+K477)

```
CAGGTGCAGCTGCAGCAGTCTGGCCCTAGACTCGTGCGGCCCTCCAGACCCTGTCTCTGACCTGTG
CCATCTCCGGCGACTCCGTGTTCAACAACAACGCCGCTGGTCTGGATCCGGCAGTCTCCATCTCG
CGGTCTGGAGTGGCTCGGTGCGACCTACTACCGCTCTAAATGGCTGTACGACTACGCCGTGTCCGTG
AAGTCCCGGATCACCGTGAACCCGTGACACCTCCCGGAACAGTTACCCCTGCAGCTGAACCTCCGTGA
CCCCTGAGGACACCGCCCTGTACTACTGCGCCAGAGGCTACTCCTCCTTCGACTATTGGGGCCA
AGGCACCCTCGTGACCGTGTCTCTGCGCTCCACCAAGGGTCCATCGGTCTTCCCCCTGGCACCCCTCC
TCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCGTGGTCAAGGACTACTTCCCCGAACCGG
TGACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTCTTACAGTC
CTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCTCCAGCAGCTTGGGCACCCAGACCTAC
ATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCAAATCTTGTG
ACAAAACCTCACACTGTCCACCGTGCCCGACCTGAAGCAGCAGGGGGACCGTCAGTCTTCCCTCTT
CCCCCAAACCCCAAGGACACCCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGAGC
GTGAGCCACGAAGACCCCTGAGGTCAAGTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCA
AGACAAAGCCGCGGGAGGAGCAGTACAACAGCAGCTACCGTGTGGTCAAGCTCCTCACCCTCTGCA
CCAGGACTGGCTGAATGGCAAGGAGTACAAGTGAAGGTGTCCAACAAAGCCCTCCAGCCCCCATC
GAGAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCC
GGGAGGAGATGACCAAGAACCAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAGCGACAT
CGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCTCCCGTGTGGAC
TCCGACGGCTCCTTCTCTGCTCTACAGCAAGCTCACCGTGGACAAGTCCAGATGGCAGCAGGGGAACG
TCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGTCTCTCTCCCTGTC
TCCGGGAAAA
```

[00632] SEQ ID NO: 1150, LC nucleic acid sequence (DuoBody) of CD3B2197 (CD3B376+K477)

```
CAGTCTGCTCTGACCCAGCCTGCCTCCGTGTCTGGCTCTCCCGGCCAGTCCATCACCATCAGCTGTA
CCGGCACCTCCTCCAACATCGGCACCTACAAGTTCGTGTCTGGTATCAGCAACACCCCGACAAGGC
CCCCAAGTGCTGCTGTACGAGGTGTCCAAGCGGCCCTCTGGCGTGTCTCCAGATTCTCCGGCTCC
AAGTCTGGCAACACCGCCTCCCTGACCATCAGCGGACTGCAGGCTGAGGACCAGGCCGACTACCACT
GTGTGTCCTACGCTGGCTCTGGCACCTGCTGTTTGGCGGAGGCACCAAGCTGACTGTCTGGGTCA
GCCCCAAGGCTGCACCCAGTGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCC
ACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGAAGGCCGATAGCA
GCCCCGTCAAGGCGGGAGTCAAAACCACACCCCTCCAAACAAAGCAACAACAAGTACGCGGCCAG
CAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAAGTCCACAGAAGCTACAGCTGCCAGGTACGCAT
GAAGGGAGCACCGTGGAGAAGACAGTGGCCCCCTACAGAATGTTCA
```

[00633] SEQ ID NO: 1171, HC nucleic acid sequence (DuoBody) of CD3B891 (CD3B376-K477)

```
CAGGTGCAGCTGCAGCAGTCTGGCCCTAGACTCGTGCGGCCCTCCAGACCCTGTCTCTGACCTGTG
CCATCTCCGGCGACTCCGTGTTCAACAACAACGCCGCTGGTCTGGATCCGGCAGAGCCCTTCTAG
AGGCCCTGGAATGGCTGGGCGGACCTACTACCGTCCAAAGTGGCTGTACGACTACGCCGTGTCCGTG
AAGTCCCGGATCACCGTGAACCCGTGACACCTCCCGGAACAGTTACCCCTGCAGCTGAACCTCCGTGA
CCCCTGAGGACACCGCCCTGTACTACTGCGCCAGAGGCTACTCCTCCTTCGACTATTGGGGCCA
GGGCACCCTCGTGACCGTGTCTCTGCTTCCACCAAGGGCCCATCCGTCTTCCCCCTGGCACCCCTCC
TCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCGTGGTCAAGGACTACTTCCCCGAACCGG
TGACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTCTTACAGTC
```

CTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTAC
ATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCAAATCTTGTG
ACAAAATCACACATGTCCACCGTGCCAGCACCTGAAGCAGCAGGGGGACCGTCAGTCTTCTCTT
CCCCCAAACCCAAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGAGC
GTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCA
AGACAAAGCCCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCTTGCA
CCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCAGCCCCCATC
GAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCC
GGGAGGAGATGACCAAGAACCAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACAT
CGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTGTGGAC
TCCGACGGCTCCTTCTGCTCTACAGCAAGCTCACCGTGGACAAGTCTAGATGGCAGCAGGGGAACG
TCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTC
TCCGGGT

[00634] SEQ ID NO: 1172, LC nucleic acid sequence (DuoBody) of CD3B891
(CD3B376-K477)

CAGTCTGCTCTGACCCAGCCTGCCTCCGTGTCTGGCTCTCCCGGCCAGTCCATCACCATCAGCTGTA
CCGGCACCTCTCCAACATCGGCACCTACAAGTTCGTGTCTGGTATCAGCAGCACCCCGACAAGGC
CCCCAAGTGCTGCTGTACGAGGTGTCCAAGCGGCCCTCTGGCGTGTCTCCAGATTCTCCGGCTCC
AAGTCTGGCAACACCGCCTCCCTGACCATCAGCGGACTGCAGGCTGAGGACCAGGGCCGACTACCACT
GTGTGTCTACGCTGGCTCTGGCACCTGCTGTTGGCGGAGGCACCAAGCTGACCGTGTGGGTCA
GCCCCAAGGCTGCACCCAGTGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCC
ACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGAAGGCCGATAGCA
GCCCCGTCAAGGGGGAGTGGAGACCACCAACCCTCCAACAAAGCAACAACAAGTACGCGGCCAG
CAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAAGTCCACAGAAGCTACAGTGCAGGTACGCAT
GAAGGGAGCACCGTGGAGAAGACAGTGGCCCTACAGAATGTTCA

[00635] SEQ ID NO: 1173, HC nucleic acid sequence (DuoBody) of CD3B2200
(CD3B450+K477)

CAGGTTGAGCTGCAGCAGTCTGGCCCTGGACTGGTCAAGCCCTCTCAGACCCTGTCTCTGACCTGTG
CCATCTCCGGCGACTCCGTGTTCAACAACAATGCCGCCTGGTCTGGATTCCGCAGTCTCCTAGTAG
AGGCTTGAATGGCTGGGCAGAACCTACTACCGGTCCAAGTGGCTGTACGACTACGCCGTGTCCGTG
AAGTCCCGGATCACCATCAATCCCGACACCTCCAAGAACCAGTTCTCCCTGCAGCTCAACAGCGTGA
CCCCTGAGGATACCGCGTGTACTACTGTGCCAGAGGCTACTCCTCCTCCTTCGACTATTGGGGCCA
GGGCACACTGGTCACCGTTTCTTCTGCCTCCACCAAGGGTCCATCGGTCTTCCCCCTGGCACCCCTCC
TCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGG
TGACGGTGTCTGGAACTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTCTTACAGTC
CTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTAC
ATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCAAATCTTGTG
ACAAAATCACACTTGTCCACCGTGCCAGCACCTGAAGCAGCAGGGGGACCGTCAGTCTTCTCTT
CCCCCAAACCCAAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGAGC
GTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCA
AGACAAAGCCCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCTGCA
CCAGGACTGGCTGAATGGCAAGGAGTACAAGTGAAGGTGTCCAACAAAGCCCTCCAGCCCCCATC
GAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCC
GGGAGGAGATGACCAAGAACCAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACAT
CGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTGTGGAC
TCCGACGGCTCCTTCTGCTCTACAGCAAGCTCACCGTGGACAAGTCCAGATGGCAGCAGGGGAACG
TCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGTCTCTCTCCCTGTC
TCCGGGAAAA

[00636] SEQ ID NO: 1174, LC nucleic acid sequence (DuoBody) of CD3B2200
(CD3B450+K477)

CAGTCTGCTCTGACCCAGCCTGCTTCTGTGTCTGGCTCTCCCGGCCAGTCCATCACCATCTCTTGTA
CCGGCACCTCCTCCACATCGGCACCTACAAGTTCGTGTCTTGGTATCAGCAGCACCCCGGCAAGGC
CCCTAAAGTGATGATCTACGAGGTGTCCAAGCGGCCCTCTGGCGTGTCCAACAGATTCTCCGGCTCC
AAGTCCGGCAATACCGCCTCTCTGACAATCAGCGGACTGCAGGCTGAGGACGAGGCCGACTACTACT
GTGTGTCTTACGCTGGCTCTGGCACCTGTGTGTTGGCGGGCGGAACAAAGCTGACTGTGCTGGGTCA
GCCCCAAGGCTGCACCCAGTGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCC
ACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGAAGGCCGATAGCA
GCCCCGTCAAGGCGGGAGTCGAAACACACACCTTCCAAACAAGCAACAACAAGTACGCGGCCAG
CAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAAGTCCCACAGAAGCTACAGCTGCCAGGTACGCAT
GAAGGGAGCACCGTGGAGAAGACAGTGGCCCCCTACAGAATGTTCA

[00637] SEQ ID NO: 1175, HC nucleic acid sequence (DuoBody) of CD3B2186
(CD3B450-K477)

CAGGTGCAGCTGCAGCAGAGCGGCCCGGCCCTGGTCAAGCCCAGCCAGACCCTGAGCCTGACCTGCG
CCATCAGCGGCGACAGCGTGTTCACAACAACGCCCGCCTGGTCTGGATCCGCCAGAGCCCCAGCCG
CGGCCCTGGAGTGGCTGGGCCGCACCTACTACCGCAGCAAGTGGCTGTACGACTACGCCGTGTCCGTG
AAGTCCCGCATCACCATCAACCCCGACACAGCAAGAACCAGTTCTCCCTGCAGCTGAACAGCGTGA
CCCCCGAGGACACCGCGTGTACTACTGCGCCCGCGGCTACAGCAGCAGCTTCGACTACTGGGGCCA
GGGCACCCTGGTCAACGTGTCCAGCGCTTCCACCAAGGGCCCATCCGTCTTCCCCCTGGCACCCCTCC
TCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGG
TGACGGTGTCTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTCTACAGTC
CTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTAC
ATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCAAATCTTGTG
ACAAACTCACACATGTCCACCGTGCCAGCACCTGAAGCAGCAGGGGGACCGTCAGTCTTCTCTT
CCCCCAAAAACCCAAAGGACACCCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTGAGC
GTGAGCCACGAAGACCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCA
AGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAAGCGTCCCTACCGTCTCTGCA
CCAGGACTGGCTGAATGGCAAGGAGTACAAGTGAAGGTCTCCAACAAGCCCTCCCAGCCCCCATC
GAGAAAACCATCTCCAAGCCAAAGGGCAGCCCGAGAACCAAGGTGTACACCTGCCCCCATCCC
GGGAGGAGATGACCAAGAACCAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAGCGACAT
CGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACTACAAGACCACGCCTCCCGTGTGGAC
TCCGACGGCTCCTTCTGCTCTACAGCAAGCTCACCGTGGACAAGTCTAGATGGCAGCAGGGGAACG
TCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTC
TCCGGGT

[00638] SEQ ID NO: 1176, LC nucleic acid sequence (DuoBody) of CD3B2186
(CD3B450-K477)

CAGTCTGCTCTGACCCAGCCTGCCTCCGTGTCTGGCTCTCCCGGCCAGTCCATCACCATCAGCTGTA
CCGGCACCTCCTCCACATCGGCACCTACAAGTTCGTGTCTTGGTATCAGCAGCACCCCGGCAAGGC
CCCCAAAGTGATGATCTACGAGGTGTCCAAGCGGCCCTCCGGCGTGTCCAACAGATTCTCCGGCTCC
AAGTCCGGCAACACCGCCTCCCTGACAATCAGCGGACTGCAGGCGGAGGACGAGGCCGACTACTACT
GTGTGTCTTACGCCGCTCTGGCACCTGCTGTTTGGCGGGCGGAACAAAGCTGACCGTGTGGGTCA
GCCCCAAGGCTGCACCCAGTGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCC
ACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGAAGGCCGATAGCA
GCCCCGTCAAGGCGGGAGTGGAGACCACACCTTCCAAACAAGCAACAACAAGTACGCGGCCAG
CAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAAGTCCCACAGAAGCTACAGCTGCCAGGTACGCAT
GAAGGGAGCACCGTGGAGAAGACAGTGGCCCCCTACAGAATGTTCA

[00639] SEQ ID NO: 1177, HC nucleic acid sequence (DuoBody) of CD3B2183 (CD3W245)

GAGGTGCAACTGGTGGAGTCTGGGGGAGGCCTGGTCAAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTCAGTAGATATAACATGAACTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCATCCATTAGTACTAGTAGTAATTACATATACTACGCAGACTCAGTGAAGGGCCGA
TTCACCTTCTCCAGAGACAACGCCAAGAACTCACTGGATCTGCAAATGAGCGGCCTGAGAGCCGAGG
ACACGGCTATTTATTACTGTACGAGAGGCTGGGGGCCTTTTGACTACTGGGGCCAGGGAACCTGGT
CACCGTCTCCTCAGCTTCCACCAAGGGCCCATCCGTCTTCCCCCTGGCACCCTCCTCCAAGAGCACC
TCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGT
GGAACCTCAGGCGCCCTGACCAGCGGCGTGACACCTTCCCGGCTGTCTACAGTCTCAGGACTCTA
CTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTG
AATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCAAATCTTGTGACAAAACCTCACA
CATGTCCACCGTGCCAGCACCTGAAGCAGCAGGGGGACCGTCAGTCTTCTCTTCCCCCCTAAAACC
CAAGGACACCCCTCATGATCTCCCGGACCCCTGAGGTCAATGCGTGGTGGTGGTGGTGGTGGTGGTGGT
GACCCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGC
GGGACGAGCAGTACACAGCACGTACCGTGTGGTCAAGCGTCTCACCCTGCTGCACAGGACTGGCT
GAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCAGCCCCCATCGAGAAAACCATC
TCCAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCTGCCCCCATCCCGGGAGGAGATGA
CCAAGAACAGGTGACCTGACCTGCTTGGTCAAGGCTTCTATCCAGCGACATCGCCGTGGAGTG
GGAGAGCAATGGGCAGCCGAGAACAACTACAAGACCACGCTCCCGTGTGGTGGTGGTGGTGGTGGTGGT
TTCCTGCTCTACAGCAAGCTCACCGTGGACAAGTCTAGATGGCAGCAGGGGAACGTCTTCTCATGTCT
CCGTGATGCATGAGGCTCTGCACAACCACTACAGCAGAAGAGCCTCTCCCTGTCTCCGGGT

[00640] SEQ ID NO: 1178, LC nucleic acid sequence (DuoBody) of CD3B2183 (CD3W245)

GACATACAAATGACACAATCACCCCTCTTCTCTTCTGCAAGCGTTGGCGACCGTGTCACTATCACTT
GTCGAGCCCCCAGTCCATAGGTACTGCCATTCACTGGTATCAACAGAAGCCTGGCAAGGCTCCCAA
ACTCCTGATTAAAGTATGCCAGCGAGAGCATTTCCGGCGTACCTTCAAGATTTCCGGCTCCGGTAGT
GGGACAGATTTCACTCTCACTATATCTAGCCTCCAACCAGAAGATTTCCGCCACTTACTAGTGTCAAC
AATCAGGTTTCATGGCCTTACACTTTCGGCCAGGGGACAAAATTGGAGATCAAGCGGACAGTGGCCGC
TCCTTCCGTGTTTCATCTTCCACCTTCCGACGAGCAGCTGAAGTCCGGCACAGCTTCTGTCTGTGTG
CTGCTGAACAACCTTCTACCCTCGGGAAGCCAAGGTGCAGTGGAGGTGGACAATGCCCTGCAGTCCG
GCAACTCCCAAGAGTCTGTGACCGAGCAGGACTCCAAGGACAGCACCTACAGCCTGTCTTCCACACT
GACCTGTCCAAGGCCGACTACGAGAAGCACAAGGTGTACGCTTCCGAAGTGACCCATCAGGGCCCTG
TCTAGCCCTGTGACCAAGTCTTTCAACCGGGGCGAGTGT

EXAMPLE 5: GENERATION OF BISPECIFIC PSMA_xCD3 ANTIBODIES

EXAMPLE 5.1: FAB-ARM EXCHANGE OF ANTI-PSMA AND ANTI-CD3 ANTIBODIES

[00641] The formation of bispecific antibodies requires two parental monoclonal antibodies (mAbs), one specific for the targeting arm (*e.g.* PSMA) and one specific for the effector arm (*e.g.* CD3). Selected monospecific anti-PSMA and anti-CD3 antibodies were expressed as IgG1/ κ engineered to have L234A, L235A and D265S substitutions for cF silencing, (EU numbering). Selected monospecific anti-PSMA and anti-CD3 antibodies are also expressed as IgG4 antibodies. Mutations designed to promote selective

heterodimerization of the Fc domain were also engineered in the Fc domain. In some cases, the anti-CD3 and anti-PSMA binders were formatted as scFv or Fab to test their stability in both formats. In other cases, scFv-based binders were tested in a VH-Linker-VL (HL) orientation or in a VL-Linker-VH orientation (LH).

[00642] The monospecific antibodies were expressed in CHO cell lines under CMV promoters as described above). The parental PSMA and CD3 antibodies were purified using a protein A column with an elution buffer of 100 mM NaAc pH 3.5 and a neutralization puffer of 2 M Tris pH 7.5 and 150 mM NaCl. The anti-PSMA and anti-CD3 monoclonal antibodies were dialyzed into D-PBS, pH 7.2 buffer.

[00643] For DuoBody antibodies, post purification of parental monospecific antibodies, bispecific PSMAxCD3 antibodies were generated by mixing the parental PSMA antibodies with the desired parental CD3 antibody under reducing conditions in 75 mM cysteamine-HCl and incubated overnight at room temperature for *in vitro* Fab arm exchange as described in Int. Patent Publ. No. WO2011/131746. The recombination reactions were based on molar ratios, where a set amount of PSMA antibody (*e.g.*, 10 mg, or ~74.6 nanomoles) was combined with CD3 antibody (*e.g.*, ~70.87 nanomoles), where the PSMA antibody was added in a 5% excess of the CD3 antibody. The concentrations of the PSMA antibody stocks varied from 0.8 to 6 mg/mL, and the volumes the recombination reactions varied for each pairing. The recombinations were subsequently dialyzed overnight against PBS to remove the reductant. The PSMAxCD3 bispecific antibody reactions were performed with an excess of the PSMA antibody (ratio) to minimize the amount of unreacted CD3 parental antibody remaining after recombination.

[00644] Other bispecifics were generated via co-transfection of HC1:HC2:LC2, typically at a DNA ratio of 1:1:3. Purification was performed by protein A chromatography and CH1 affinity capture, followed by an ion exchange-based chromatography.

[00645] Exemplary PSMAxCD3 multispecific antibodies are provided in Tables 22-27.

Table 22. PSMA x CD3 Bispecific Antibodies: Clone Descriptions

Name	Bispecific Description
PS3B1353	HC1 (F405L): CD3B376 x HC2 (K409R); PSMB947
PS3B1505	HC1 CD3B376-Fab x HC2 PSMB896-G100A IgG1 DuoBody
PS3B1508	HC1 (Knob3): CD3W245-LH-scFv; HC2 (Hole3-RF) PSMB896-G100A-Fab-RF; IgG1 AAS
PS3B1917	HC1 (ZWA w/o K447_RF): CD3B376-Fab; HC2 (ZWB w/o K447): PSMA_P72_A10-HC-G54E-scFv LH
PS3B1918	HC1 (ZWA w/o K447_RF): CD3B376-Fab; HC2 (ZWB w/o K447): PSMA_P72_D01-HC-D95E-scFv LH
PS3B1919	HC1 (ZWA w/o K447_RF): CD3B376-Fab; HC2 (ZWB w/o K447): PSMA_P75_F01, LH
PS3B1920	HC1 (ZWA w/o K447_RF): CD3B376-Fab; HC2 (ZWB w/o K447): PSMA_P72_F07, LH
PS3B1921	HC1 (ZWA w/o K447_RF): CD3B376-Fab; HC2 (ZWB w/o K447): PSMA_P72_E07, LH
PS3B1922	HC1 (ZWA w/o K447_RF): CD3B376-Fab; HC2 (ZWB w/o K447): PSMA_P72_D01, LH
PS3B1923	HC1 (ZWA w/o K447_RF): CD3B376-Fab; HC2 (ZWB w/o K447): PSMA_P72_C01, LH
PS3B1924	HC1 (ZWA w/o K447_RF): CD3B376-Fab; HC2 (ZWB w/o K447): PSMA_P72_A10, LH
PS3B1925	HC1 (ZWA w/o K447_RF): CD3B376-Fab; HC2 (ZWB w/o K447): PSMA_P70_F02, LH
PS3B1926	HC1 (ZWA w/o K447_RF): CD3B376-Fab; HC2 (ZWB w/o K447): PSMA_P72_G02, HL
PS3B1927	HC1 (ZWA w/o K447_RF): CD3B376-Fab; HC2 (ZWB w/o K447): PSMA_P72_C01, HL
PS3B1928	HC1 (ZWA w/o K447_RF): CD3B376-Fab; HC2 (ZWB w/o K447): PSMA_P72_A11, HL
PSMB1041	HC1 (ZWA): B23B62-Fab; HC2 (ZWB): PSMA_P70_F02, LH
PSMB1045	HC1 (ZWA): B23B62-Fab; HC2 (ZWB): PSMA_P72_A10, LH
PSMB1047	HC1 (ZWA): B23B62-Fab; HC2 (ZWB): PSMA_P72_C01, LH
PSMB1049	HC1 (ZWA): B23B62-Fab; HC2 (ZWB): PSMA_P72_D01, LH
PSMB1051	HC1 (ZWA): B23B62-Fab; HC2 (ZWB): PSMA_P72_E07, LH

PSMB1052	HC1 (ZWA): B23B62-Fab, HC2 (ZWB): PSMA_P72_F07, LH
PSMB1060	HC1 (ZWA): B23B62-Fab, HC2 (ZWB): PSMA_P75_F01, LH
PSMB1068	HC1 (ZWA): B23B62-Fab, HC2 (ZWB): PSMA_P72_A11, HL
PSMB1069	HC1 (ZWA): B23B62-Fab, HC2 (ZWB): PSMA_P72_C01, HL
PSMB1075	HC1 (ZWA): B23B62-Fab, HC2 (ZWB): PSMA_P72_G02, HL
PSMB2908	HC1 (ZWA w/o K447): B23B62-Fab, HC2 (ZWB w/o K447): PSMA_P72_D01-HC-D93E-scFv LH
PSMB2909	HC1 (ZWA w/o K447): B23B62-Fab, HC2 (ZWB w/o K447): PSMA_P72_A10-HC-G54E-scFv LH
PS3B1391	HC1: N-term scFv LH CD3B2030 N106A LH-scFv (MA), HC2: N-term_Fab_PSMHB49SC1133_011A11_1
PS3B1396	HC1 (Knob3): CD3B2030-N106A-scFv LH, HC2 (Hole3): PSMB896-G100A-Fab

Table 23. PSMA x CD3 Bispecific Antibodies: CD3 Arm Descriptions

Name	HC1	SEQ ID NO.	LC1	SEQ ID NO.	CD3 Arm Description
PS3B1353	(see above)	849	(see above)	850	with CD3B376 arm (CD3B891 without K477 or CD3B2197 with K477 in HC1)
PS3B1505	(see above)	849	(see above)	850	with CD3B376 arm (CD3B891 without K477 or CD3B2197 with K477 in HC1)
PS3B1508	DIQMTQSPSSLSASVGDRVTITCFARQSIGTALHWYQKPGKAPK LLIKYASEISGVPSRFGSGSGTDFTLTITSSLOPEDEATYYCQQ SSGWPYTFGGTKLEIKRGSEKSGSGSEKSTGGSEVQLVESG GGLVKPGGSLRLSCAASGFTFPRYNMWRQAPGKGLIEWVSSIST SSNIYYADSVKGRFTFRDNARNSLDLQMSGGLPAEDTAIYYCTR GWGPFYWGQGLVTVSSEPKSSDKTHICPPCPAPEAAGGPSVFL FPPKPKDTLMI SRTPEVTCVVVSVSHEDPEVKFNWYVDGVEVHNA KTKPREQYNSTYRVSVLTVHLQDNLNGKEYCKKVSNAKALPAPI EKTISKAKGQPREPQVYTLPPSPEEMTKNQVSLVCLVKGFYPSDI AVEWESNGQPENNYKTPPPVLDSDGSGFFLYSKLTVDKSRWQQGNV FSCSVMEALHNHYTQKSLSLSPGK	1192	NA	1193	with CD3W245 arm (CD3B2183 without K477)
PS3B1917	QVQLQQSGFRLVRFSQTLSTLCAISGDSVFNNNAAWSWIRQSESR GLEWLGRTYYRSKWLYDYAVSVKSRITVNPDTSRNQFTLQLNSVT PEDTALYYCARGYSSTFDYWGQGLTVTVSSASTKGPVFPPLAPSS KSTSGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSS GLYSLSVWTVPSSSLGTQTYICNVNHKPSNTKVDKVEPKSCDK THTCPPCPAPEAAGGPSVFLFPPKPKDTLMI SRTPEVTCVVSVS HEDPEVKFNWYVDGVEVHNAKTKPREQYNSTYRVSVLTVHLQD NLNGKEYCKVSNKALPAPI EKTISKAKGQPREPQVYVPPSREE MTRNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPPVLDSDG SFALVSKLTVDKSRWQQGNVFCSSVMHEALHNRTQKLSLSLSPG	1194	QSALTQPAVSVSGSPGQSIT ISCTGTSSNIGTYKRVSWY QQHPDKAPKVLLEYVSKRP SGVSRFSGSKSGNTASLT ISGLQAEQADYHCVSYAG SGTLLFGGGTKLTVLRTVA APSVFIFPPSDEQLKSGTA SVVCLLNFFYPPEAKVQWK VDNALQSGNSQESVTEQDS KDTYSLSSTLTLSKADYE KHKVYACEVTHQGLSSPVT KSFNRGEC	1195	with CD3B376 arm (CD3B891 without K477 or CD3B2197 with K477 in HC1)
PS3B1918	(see above)	1194	(see above)	1195	with CD3B376 arm (CD3B891 without K477 or CD3B2197 with K477 in HC1)

						K477 or CD3B2197 with K477 in HCl)
PS3B1919	(see above)	1194	(see above)	1195		with CD3B376 am (CD3B891 without K477 or CD3B2197 with K477 in HCl)
PS3B1920	(see above)	1194	(see above)	1195		with CD3B376 am (CD3B891 without K477 or CD3B2197 with K477 in HCl)
PS3B1921	(see above)	1194	(see above)	1195		with CD3B376 am (CD3B891 without K477 or CD3B2197 with K477 in HCl)
PS3B1922	(see above)	1194	(see above)	1195		with CD3B376 am (CD3B891 without K477 or CD3B2197 with K477 in HCl)
PS3B1923	(see above)	1194	(see above)	1195		with CD3B376 am (CD3B891 without K477 or CD3B2197 with K477 in HCl)
PS3B1924	(see above)	1194	(see above)	1195		with CD3B376 am (CD3B891 without K477 or CD3B2197 with K477 in HCl)
PS3B1925	(see above)	1194	(see above)	1195		with CD3B376 am (CD3B891 without K477 or CD3B2197 with K477 in HCl)
PS3B1926	(see above)	1194	(see above)	1195		with CD3B376 am (CD3B891 without K477 or CD3B2197 with K477 in HCl)

PS3B1927	(see above)	1194	(see above)	1195	with CD3B376 arm (CD3B891 without K477 or CD3B2197 with K477 in HCl)
PS3B1928	(see above)	1194	(see above)	1195	with CD3B376 arm (CD3B891 without K477 or CD3B2197 with K477 in HCl)
PSMB1041	QITLKESGPTLVKPTQTTLTCTFSGFSLSTSGMGVSWIRQPPGK ALEWLAHIYWDGDKRYNPSLKSLTITKDTSKQVWLTMTNMDPV DTATYICARLYGFTYGFAYWGQTLTVSSASTKGPSVFPLAPSS KSTSGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVWTPSSSLGFTQYICNVNHNKPSNTKVDKVEPKSCDK THTCPPCPAPELGGPSVLEFPPKPKDTLMI SRTPETCVVVDVS HEDPEVFNWYVDGVEVHNKATKPREEQINSTRYVSVLTVLHQD WLNKKEYCKVSNKALPAPIEKTI SKAKGQPREPQVYVPPSREE MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPFVLDSDG SPALVSKLTVDKSPWQQGNVFCSCVMHEALHNRFTQKSLSLSPG	1218	DIVMTQSPDLSIAVSLGERA TINCPASQSYDNGISYMH WYQQKPGQPKLLIYAASN PESGVPDRFSGSGSGTDFT LTISLSQAEDVAVYYCQOI IEDPWTFGGGTKEIKRTV AAPSVFIFFPSDEQLKSGT ASVVCLLNNEFYPREAKVQW KVDNALQSGNSQESVTEQD SKDSTYSLSTLTLSKADY EKHKVYACEVTHQGLSPV TKSENRGEC	1219	Null CD3 arm
PSMB1045	(see above)	1218	(see above)	1219	Null CD3 arm
PSMB1047	(see above)	1218	(see above)	1219	Null CD3 arm
PSMB1049	(see above)	1218	(see above)	1219	Null CD3 arm
PSMB1051	(see above)	1218	(see above)	1219	Null CD3 arm
PSMB1052	(see above)	1218	(see above)	1219	Null CD3 arm
PSMB1060	(see above)	1218	(see above)	1219	Null CD3 arm
PSMB1068	(see above)	1218	(see above)	1219	Null CD3 arm
PSMB1069	(see above)	1218	(see above)	1219	Null CD3 arm
PSMB1075	(see above)	1218	(see above)	1219	Null CD3 arm
PSMB2908	QITLKESGPTLVKPTQTTLTCTFSGFSLSTSGMGVSWIRQPPGK ALEWLAHIYWDGDKRYNPSLKSLTITKDTSKQVWLTMTNMDPV DTATYICARLYGFTYGFAYWGQTLTVSSASTKGPSVFPLAPSS KSTSGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVWTPSSSLGFTQYICNVNHNKPSNTKVDKVEPKSCDK THTCPPCPAPEAAGGPSVLEFPPKPKDTLMI SRTPETCVVVDVS HEDPEVFNWYVDGVEVHNKATKPREEQINSTRYVSVLTVLHQD WLNKKEYCKVSNKALPAPIEKTI SKAKGQPREPQVYVPPSREE	1238	(see above)	1219	Null CD3 arm

	MTKNQVSLTCLVKGFYPSDIAVWESNQPENNYKTTFFVLDSDG SEALVSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPG	1238	(see above)		
PSMB2909	(see above)	1219			Null CD3 arm
PS3BI391	EIVLTQSEATLSASPGGERVTLSCASSSVSNMNYQQKPGQARR WIYDSSKLASGVPARFSGSGGFDYTLTISLPEDEFAVYQCQW SRNPTFGGKVEIKGSGEGKSSGSGESKSTGGQVQLVQSGA EVKKPGSSVKSCAKSGYTFTRSTMHWKQAPQGQLEWIGYINPS SAYTNYNQKFGQGRVTLTADKSTSTAYMELSSLRSEDTAVYICASP QVHYDYAGFPYWGQGLTVTSSEPKSSDKTHTCPPCPAPEAAGGP SVFLFPPKPKDTIMI SRTPEVTCVWSVSHEDPEVKFNWYVDGVE VHNAKTRPREEQYNSTYRVSVLTVLHQDWLNGKEYCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLWCLVKGFY PSDIAVWESNQPENNYKTTFFVLDSDGSEFFLYSKLTVDKSRWQ QGNVFCSCVMHEALHNHYTQKSLSLSPGK	1504	NA	1456	CD3B2030
PS3BI396	(see above)	1455	NA	1458	CD3B2030-N106A- scFv-LH-CH2

	YYGMDVWGQGTTVTVSSEPKSSDKTHTCCPPCPAPEAAGGSPVFLFP PKPKDTLMI SRTPEVTCVVSVSHEDPEVKFNWVVDGVEVHNKTK PREEQYNSTYRVSVLTLHQDLNGLREYKCKVSNKALPAPI EKT I SKAKGQPREPQVVVLPSPREEMTKNQVSLCLVKGFYPSDIAVEWE SNGQPENNYLTWPPVLDSDGSEFFLYSKLTVDKSRWQQGNVFCVM HEALHNHYTQKSLSLSPG	1250	NA	1251	PSMA_P72_ D01-HC- D95E
PS3B1918	QSVLTQPASVSGSPGQTTISCTGTSSDVGGVYVSWYQQHKGAP KLMIYEVSNRPSGVNRPFGSKSGNTASLTISGLQAEADYYCSSL YTSSYTVVFGTGKLTVLGGSEKSGSGSEKSTGGSEVQLVESG GDLVQPGGSLRLSCAASGFTFNNYNNWVRQAPGKGLEWVSHISTS SNKYADSVKGRFTISRDIANKSMYLMQNSLRDEDTAVYYCAREG VGADYGDYYYGMDVWGQGTTVTVSSEPKSSDKTHTCCPPCPAPEAA GGPSVFLFPPKPKDTLMI SRTPEVTCVVSVSHEDPEVKFNWVVDG VEVHNKTKPREEQYNSTYRVSVLTLHQDLNGLREYKCKVSNKA LPAPI EKTISKAKGQPREPQVVVLPSPREEMTKNQVSLCLVKGFY PSDIAVEWESNGQPENNYLTWPPVLDSDGSEFFLYSKLTVDKSRWQQ GNVFCVMHEALHNHYTQKSLSLSPG	1252	NA	1253	PSMA_P75_F 01
PS3B1919	EIVLTQSPGTTLSVSPGERATLSCRASQSVRSNLAWYQQKPGQAPRL LIYGASTRATGIPARFSGSGGTFTLTISLSQSEDEAVYYCHQYN DWPPYTFGGGTKLEIKGSGEGKSGSGSEKSTGGSOVQLQESGGG VVQPGRLRLSCAASGFTFSTYGMHWVRQAPGKGLEWVAFI SYDGS NKYYADSVKGRFTISRDNKHTLYLQNNLSRAEDTAVYYCAGRDNL RLEWFMDVWGQGTTVTVSSEPKSSDKTHTCCPPCPAPEAAGGSPVF LPPKPKDTLMI SRTPEVTCVVSVSHEDPEVKFNWVVDGVEVHNA KTKPREEQYNSTYRVSVLTLHQDLNGLREYKCKVSNKALPAPIE KTIISKAKGQPREPQVVVLPSPREEMTKNQVSLCLVKGFYPSDIAV EWESNGQPENNYLTWPPVLDSDGSEFFLYSKLTVDKSRWQQGNVFC SVMHEALHNHYTQKSLSLSPG	1254	NA	1255	PSMA_P72_F 07
PS3B1920	SYELTQPPPSVAPAGQTARITCGNNIGSKSVHWYQQKPGQAPVLV VYDDSDRPPSGIPERFSGSNSGNTATLTISRVEAGDEADYYCQWDS STIDHVFEGGKTLTLVGGSEKSGSGSEKSTGGSEVQLVESGGG VVQPGRLRLSCAASGFTFSSYGMHWVRQAPGKGLEWVAVTSYDGS NKYYADSVKGRFTISRDI SKNTLYLQMSLSRAEDTAVYYCARDPYS SSWNGAFDIWGPGTMVTVSSEPKSSDKTHTCCPPCPAPEAAGGSPVF LPPKPKDTLMI SRTPEVTCVVSVSHEDPEVKFNWVVDGVEVHNA KTKPREEQYNSTYRVSVLTLHQDLNGLREYKCKVSNKALPAPIE KTIISKAKGQPREPQVVVLPSPREEMTKNQVSLCLVKGFYPSDIAV EWESNGQPENNYLTWPPVLDSDGSEFFLYSKLTVDKSRWQQGNVFC SVMHEALHNHYTQKSLSLSPG				

PS3B1921	QSVLTQPPSASGTFPGQVTISCSGSSNIGSNTVNWFQQLPGTAPK LLIYSDNQRPSPGVDRFSGSKSGLAISGLQSEADYICAAW DDSLNGYVFGTGKTVTLGGSEKSGSGSEKSTGGSEVQLVESG GGVVQPGSLRLSCAASGFTTITYGMHWVRQAPGKGLEWVAVVSFD ESNKIYADSVKGRFTIISPDNSKNTLYLQMNLSLPAEDTAVIYCARAL RDGNWDYFNGMDVWGQTTVVSSEPKSSDKTHTCPCPAPAEAG GPSVFLFPKPKDTLMIISRTPEVTCVVSVSHEDPEVKFNWYVDGV EVHNAKTPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAL PAPIEKTISKAKGQPREPQVYVLPVPPSREEMTKNQVSLTCLVRGFY SDIAVESNGQPPENNYLTWPPVLDSDGSFFLYSKLTVDKSRWQOQ NVFSCSVMHENHHTQKSLSPG	1256	NA	1257	PSMA_P72_E 07
PS3B1922	QSVLTQPPASVSGSPGQSIITISCTGTSSDVGGYNYVSWYQQHPGKAP KLMIYEVSNNRPSGVNRFSGSKSGNTASLITISGLQAEADYICSS YTSSYTVFGTGKTLTVLGGSEKSGSGSEKSTGGSEVQLVESG GDLVQPGSLRLSCAASGFTFNYNMNVVRQAPGKGLEWVSHISTS SSNKYADSVKGRFSISRDIANKSMYLMQNSLRDEDTAVIYCARDG VGADYGDYIYGMVWGQTTVVSSEPKSSDKTHTCPCPAPAEAA GGPSVFLFPKPKDTLMIISRTPEVTCVVSVSHEDPEVKFNWYVDG VEVHNAKTPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKA LPAPIEKTISKAKGQPREPQVYVLPVPPSREEMTKNQVSLTCLVRGFY PSDIAVESNGQPPENNYLTWPPVLDSDGSFFLYSKLTVDKSRWQOQ GNVFCSSVMHEALHHTQKSLSPG	1258	NA	1259	PSMA_P72_ D01
PS3B1923	QSVLTQPPSVVAPGQTARITCGGNNSGSKSVHWYQQHPGQAPVLV VYDDSDRPSGIPERFSGNSGNTATLITSRVAGDEADYICQVWDS SSDHGVFGGKTLTVLGGSEKSGSGSEKSTGGSQVQLVESGGG EYQPGSLRLSCAASGFSFGYGMHWVRQAPGKGLEWVAVMSYDGS NRFYVDSVGRFISIIRDNSKNTLYLQMNLSLRDEDTAVIYCARDTVW GSHPDADIWGGQTTVVSSEPKSSDKTHTCPCPAPAEAGGVSF LFPKPKDTLMIISRTPEVTCVVSVSHEDPEVKFNWYVDGVEYHNA KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE KTIISKAKGQPREPQVYVLPVPPSREEMTKNQVSLTCLVRGFYPSDIAV EWSNGQPPENNYLTWPPVLDSDGSFFLYSKLTVDKSRWQOQNVFSC SYMHEALHHTQKSLSPG	1260	NA	1261	PSMA_P72_C 01
PS3B1924	SYELMQPPSVSVSPGQTARITCGDALPKQYAVWYQQHPGQAPVLV IYKDSERPSSIIVRFSGSSSGTTLITITGVQAEADYICQSDS SGTYVFGTGKTVTLGGSEKSGSGSEKSTGGSQVQLVESGGGV VQPGSLRLSCAASGFTFSSYNMNVVRQAPGKGLEWVAIIYYDGSN KYIADSVKGRFTISRDIANKNTLYLQMNLSLRDEDTAVIYCARERGRD YYGMDVWGQTTVVSSEPKSSDKTHTCPCPAPAEAGGSPVFLFP	1262	NA	1263	PSMA_P72_ A10

	PKPKDTLMI SRTPEVTCVVSVSHEDPEVKFNWYVDGVEVHNAKTK PREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYVLPFSREEMTKNQVSLTCLVKGFYPSDIAVEWE SNGQPENNYLTWPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVM HEALHNHYTQKSLSLSPG	1264	NA	1265	PSMA_P70_F 02
PS3B1925	QSVLTQFPASGTPGQRVTISCSGSSNIGSTVNWQQLPGTAPK LLIYSSNQRP SGVDFRFSGSKGTSASLAISGLQSEADYCAAW DDSLNGVFGGKTLTVLGGSEKSSGSGSEKSTGGSEVQLLESG PGLVKPSETLSLCTCTVSGGSIISYYWSWIRQAPAGKLEWIGRIYSS GSTNYPFSLKSRVTMSVDT SKNQFSLKLSSTAAADTA VYCAKGV WPGAFDIWGQGTMTVSSSEPKSSDKTHTCPPCPAPEAAGGPSVLEF PKPKDTLMI SRTPEVTCVVSVSHEDPEVKFNWYVDGVEVHNAKTK KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKT ISKAKGQPREPQVYVLPFSREEMTKNQVSLTCLVKGFYPSDIAVEW ESNGQPENNYLTWPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSV MHEALHNHYTQKSLSLSPG	1266	NA	1267	PSMA_P72_ G02
PS3B1926	EVQLVESGGGVQPGGRSLRLSCAASGFSFSGYGMHWVRQAPGKGLE WVAVI SYDGSNKYYADSVKGRFTISRDN SKNTLYLQMNSLRVEDTA VYYCARDRIWGSRGYYIGMDVWGQGTITVTVSSGSGSEKSSGSGSES KSTGGQSALTQPASVSGSPGQSITISCTGASDDVGGYVSWYQQ HFGKAPKLMIEVSNRPSGVNRFSGSKSGNTASLTISGLQADEA DYVCSSTITSTLVFGGKTLTVLEPKSSDKTHTCPPCPAPEAAGG PSVLFPPKPKDTLMI SRTPEVTCVVSVSHEDPEVKFNWYVDGVE VHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYVLPFSREEMTKNQVSLTCLVKGFYPS DIAVEWESNGQPENNYLTWPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVNHEALHNHYTQKSLSLSPG	1268	NA	1269	PSMA_P72_C 01
PS3B1927	QVQLVESGGGVQPGGRSLRLSCAASGFSFSGYGMHWVRQAPGKGLE WVAVMSYDGSNRFYVDVSVGRFISRDN SKNTLYLQMNSLRVEDTA VYYCARDTVWGSHPDADFIWGQGTITVTVSSGSGSEKSSGSGSEKSS TGGQSALTQPASVSGSPGQSITISCTGASDDVGGYVSWYQQEQA PVLVYDDSDRPSGIPERFSGNSGNTALTISRVEAGDEADYICQ VWDSSDHGVFGGKTLTVLEPKSSDKTHTCPPCPAPEAAGGPSVLEF LPPKPKDTLMI SRTPEVTCVVSVSHEDPEVKFNWYVDGVEVHNA KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE KTI SKAKGQPREPQVYVLPFSREEMTKNQVSLTCLVKGFYPSDIAV EWESNGQPENNYLTWPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSV SVNHEALHNHYTQKSLSLSPG				

PS3B1928	QVQLQESGGDVVQPGSRSLRLSCAASGFSFSGYGLHWVRQAPGRGLE WYTLISYDGSNKYYADSVKGRFTISRDNKNTILYLQNMNLSRAEDTA VYYCAKTTVSDPYIGMDVWGQGTITVSSGGSEKSSGSGSKS TGGSSYELTQPPSVSVAPGQTARITCGGNNIGSKSVHWYQQKPGQA FVLVYDDSDRPSGIPERFSGTNSGNTAIIITSPRAEAGDEADYYCQ VWDSSSDHWVFGGGTKLVLEPKSSDKTHTCPCPAPEAAGGPSVF LPPPKPKDTLMISRTPEVTCVVVSVSHEDPEVKFNWYVDGVEVHNA KTKPREQYNSTYRVVSVLTVHLQDNLNGKEYCKVSNKALPAPIE KTISKAKGQPRPPQVYVLPSPREEMTRNQVSLCLVKGFYPSDIAV EWESNGQPFENNYLTWPPVLDSDGSFFLYSKLTVDKSRWQQGNVFC SVMHEALHNHYTQKSLSLSPG	1270	NA	1271	PSMA_P72_ A11
PSMB1041	(see above)	1264	NA	1273	PSMA_P70_F 02
PSMB1045	(see above)	1262	NA	1275	PSMA_P72_ A10
PSMB1047	(see above)	1260	NA	1277	PSMA_P72_C 01
PSMB1049	(see above)	1258	NA	1279	PSMA_P72_ D01
PSMB1051	(see above)	1256	NA	1281	PSMA_P72_E 07
PSMB1052	(see above)	1254	NA	1283	PSMA_P72_F 07
PSMB1060	(see above)	1252	NA	1285	PSMA_P75_F 01
PSMB1068	(see above)	1270	NA	1287	PSMA_P72_ A11
PSMB1069	(see above)	1268	NA	1289	PSMA_P72_C 01
PSMB1075	(see above)	1266	NA	1291	PSMA_P72_ G02
PSMB2908	(see above)	1250	NA	1293	PSMA_P72_ D01-HC-D95E

PSMB2909	(see above)	1248	NA	1295	PSMA_P72- A10-HC- G54E
PS3B1391	EVQLVESGGGLVKPGGSLRLSCVASGFTFSFYSMNWVRQAPRGGLD WVSSISSSGNYIYYADSVKGRFTISRDAKNSLHLHMSLKAEDTA MYFCARSGSYDAFDEWGQGTMTVSSASTKGFSPFLAPSSKST SGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLYS LSSVWTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCP PCPAPEAAGGPSVFLFPPPKDTLMI SRTPEVTCVVSVSHEDPEV KENWYVDGVEVHNAKTRPREQYNSTYRVWVSLTLHQDWLNGKEY KCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREMTKNQVSL SCAVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSEFFLVSKLT VDKSRWQQGNVFSCSVMHEALHNRFQKSLSLSPGK (see above)	407	EIVMTQSPGTLSLSPGERATLS CRASQSVSSSFLAWYQQKPGQA PRLLISGASSRATGIPDRFSVS GSGTDFTLTISRLEPEDEFAVIY CQYGVSPWTFGGQTKVEIKRT VAAPSVFIFPPSDEQLKSGTAS VVCLLNNFYPPRAKVVQWKVDNA LQSGNSQESVTEQDSKDDSTYSL SSTLTLSKADYEKHKVYACEVT HQLSSPVTKSENRGEC (see above)	408	PSMHB49SC 1133_011A11 _1
PS3B1396	(see above)	441	(see above)	272	PSMB896- G100A-Fab

Table 25. PSMA x CD3 Bispecific Antibodies: Clone Descriptions

Name	Bispecific Description
PS3B917	FAB-A: CD3B891 (CD3B376-K477); FAB-B: PSMB889
PS3B918	FAB-A: CD3B891 (CD3B376-K477); FAB-B: PSMB890
PS3B913	FAB-A: CD3B891 (CD3B376-K477); FAB-B: PSMB891
PS3B915	FAB-A: CD3B891 (CD3B376-K477); FAB-B: PSMB892
PS3B914	FAB-A: CD3B891 (CD3B376-K477); FAB-B: PSMB893
PS3B916	FAB-A: CD3B891 (CD3B376-K477); FAB-B: PSMB894
PS3B919	FAB-A: CD3B891 (CD3B376-K477); FAB-B: PSMB895
PS3B921	FAB-A: CD3B891 (CD3B376-K477); FAB-B: PSMB896
PS3B920	FAB-A: CD3B891 (CD3B376-K477); FAB-B: PSMB897
PS3B922	FAB-A: CD3B891 (CD3B376-K477); FAB-B: PSMB898
PS3B912	FAB-A: CD3B891 (CD3B376-K477); FAB-B: PSMB899
PS3B930	FAB-A: CD3B2183 (CD3W245); FAB-B: PSMB889
PS3B931	FAB-A: CD3B2183 (CD3W245); FAB-B: PSMB890
PS3B926	FAB-A: CD3B2183 (CD3W245); FAB-B: PSMB891
PS3B928	FAB-A: CD3B2183 (CD3W245); FAB-B: PSMB892
PS3B927	FAB-A: CD3B2183 (CD3W245); FAB-B: PSMB893
PS3B929	FAB-A: CD3B2183 (CD3W245); FAB-B: PSMB894
PS3B932	FAB-A: CD3B2183 (CD3W245); FAB-B: PSMB895
PS3B934	FAB-A: CD3B2183 (CD3W245); FAB-B: PSMB896
PS3B933	FAB-A: CD3B2183 (CD3W245); FAB-B: PSMB897
PS3B935	FAB-A: CD3B2183 (CD3W245); FAB-B: PSMB898
PS3B925	FAB-A: CD3B2183 (CD3W245); FAB-B: PSMB899
PS3B1352	FAB-A: CD3B2197 (CD3B376+K477); FAB-B: PSMB946
PS3B1353	FAB-A: CD3B2197 (CD3B376+K477); FAB-B: PSMB947
PS3B1354	FAB-A: CD3B2197 (CD3B376+K477); FAB-B: PSMB948
PS3B1355	FAB-A: CD3B2197 (CD3B376+K477); FAB-B: PSMB949
PS3B1356	FAB-A: CD3B2200 (CD3B450+K477); FAB-B: PSMB946
PS3B1357	FAB-A: CD3B2200 (CD3B450+K477); FAB-B: PSMB947
PS3B1358	FAB-A: CD3B2200 (CD3B450+K477); FAB-B: PSMB949
PSMB937	FAB-A: CD3B2186 (CD3B450-K477); FAB-B: PSMB897 (PSMB948)

Table 26. PSMA x CD3 Bispecific Antibodies: CD3 Arm Descriptions

Name	HC1	LC1	CD3 Arm Description
PS3B917, PS3B918, PS3B913, PS3B915, PS3B914, PS3B916, PS3B919, PS3B921, PS3B920, PS3B922, PS3B912	SEQ ID NO: 883	SEQ ID NO: 850	with CD3B376 arm (CD3B891 without K477 or CD3B2197 with K477 in HC1) HC1 Construct ID: PBD000100300 LC1 Construct ID: PBD000044707
PS3B930, PS3B931, PS3B926, PS3B928, PS3B927, PS3B929, PS3B932, PS3B934, PS3B933, PS3B935, PS3B925	SEQ ID NO: 1167	SEQ ID NO: 986	with CD3W245 arm (CD3B2183 without K477) HC1 Construct ID: PBD000100302 LC1 Construct ID: PBD000084982
PS3B1352, PS3B1354, PS3B1355, PS3B1356, PS3B1357, PS3B1358	SEQ ID NO: 849	SEQ ID NO: 850	with CD3B376 arm (CD3B891 without K477 or CD3B2197 with K477 in HC1) HC1 Construct ID: PBD000108469 LC1 Construct ID: PBD000108469
PSMB937	SEQ ID NO: 951	SEQ ID NO: 918	with CD3B450 arm (CD3B2186 without K477 or CD3B2200 with K477 in HC1) HC1 Construct ID: PBD000100305 LC1 Construct ID: PBD000045576

Table 27. PSMA x CD3 Bispecific Antibodies: PSMA Arm Descriptions

Name	HC2	SEQ ID NO.	LC2	SEQ ID NO.	PSMA Arm Description
PS3B917	EVQLVESGGGVVQPGRLRLSCAASGFTFSRYNNMWVRQAPG KGLEWVSSINSRYIYADSVKGRFTISRDAKNSLYIQMN SLPAEDTAVYYCAKTMGDYYYGVMDVWGQGTFTVVSASTK GPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVSMNSGAL TSGVHTFPAVLQSSGLYSLSSVTVPPSSSLGTQTYICNVNHK PSNTKVDKKVEPKSCDKTHCTCPPEAAAGGSPVFLPPFKP KDTLMI SRTPEVTCVVVSVSHEDPEVKFNWYVDGVEVHNAKT KPREEQXNSTYRVVSVLTVHLQDWLNGKEYCKVSNKALPAP IEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFY PSDI AVEWESNGQPENNYKTTTPVLDSDGSEFFLYSRLTVDKS RWQGGNVFSCSVMEALHNHYTQKSLSLSPG (see above)	33	QSVLTQPPSVSGAPGQGVRTISC TGSSFNLGAGYDVHWYQQVPGT VPKLLIYDNSNRPSGVPDFRFG SKSGTSASLAITGLQAEDEVY YCQSYDSLSGVVFGGKLTIV LGQPKAAPSVTLFPPSSEELQA NKATLVCLISDFYPGAVTVAMK ADSSPVKAGVETTTTPSKQSNKK YAASSYLSLTPEQWKSHRSYSC QVTHEGSTVEKTVAPTECS	34	PSMB889 HC2 Construct ID: PBD000101312
PS3B918	(see above)	33	SSELTQPPSVSGAPGQGVRTISC AGSLNIGAGYDVHWYQQVPGT APKLLIYGNINRLSGVPERFSG SKSGTSASLAITGLQAEDEVY YCQSYDSLSGVVFGGKLTIV LGQPKAAPSVTLFPPSSEELQA NKATLVCLISDFYPGAVTVAMK ADSSPVKAGVETTTTPSKQSNKK YAASSYLSLTPEQWKSHRSYSC QVTHEGSTVEKTVAPTECS	68	PSMB890 HC2 Construct ID: PBD000101312
PS3B913	EVQLVESGGGVVQPGRLRLSCAASGFTFTTYGNHWRQAPG KGLEWVAVVSFDESNNKYADSVKGRFTISRDNKNTLYIQMN SLPAEDTAVYYCARALRDGNNDYFNGMDVWGQGTFTTVSSA STKGPSVPEPLAPSSKSTSGGTAALGCLVKDYFPEPTVSMNS GALTSGVHTFPAVLQSSGLYSLSSVTVFSSSLGTQTYICNV NHKPSNTKVDKKVEPKSCDKTHCTCPPEAAAGGSPVFLFP PAKPKDTLMI SRTPEVTCVVVSVSHEDPEVKFNWYVDGVEVHN AKTKPREEQXNSTYRVVSVLTVHLQDWLNGKEYCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVK GYPSDI AVEWESNGQPENNYKTTTPVLDSDGSEFFLYSRLTV DKSRWQGGNVFSCSVMEALHNHYTQKSLSLSPG (see above)	101	QSVLTQPPSVSGTPEGQVTISC SGSSNIGSNTVNWYQQVPGTA PKLLIYSDNQRPSSGVDFRFGS KSGTSASLAISGLQSEDEADYY CAAWDDSLNGYVFGTGTGKTVL GQPKAAPSVTLFPPSSEELQAN KATLVCLISDFYPGAVTVAMKA DSSPVKAGVETTTTPSKQSNKKY AASSYLSLTPEQWKSHRSYSCQ VTHEGSTVEKTVAPTECS	102	PSMB891 HC2 Construct ID: PBD000101316
PS3B915	(see above)	101	QSVLTQPPSVSGAPGQGVRTISC TGSSNIGAGYDVHWYQHLPGT APKLLIYGNNSNRPSGVPDFRFG	136	PSMB892 HC2 Construct ID: PBD000101316

				SKGTSASLAITGLQAEDETDY YCQSYDSSLSGWVFGGTKLTV LGQPKAAPSVTLFPPPSSEELQA NKATLVCLISDFYPGAVTAVWK ADSPVKAGVETTTPSKQSNK YAASSYLSTPEQWKSHRSYSC QVTHEGSTVEKTVAPTECS		
PS3B914	QVQLVESGGGVQPGGSLRLSCVASGFTFESSYGIHWVRQAPG KGLEWVAIVWYDGSNKKYADSVKGRFTISRDNKNTLYLQMN SLRAEDTAVYYSVRGVGPTSYNYNGMDVWGQTTVTVSSAS TKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPPTVSWNSG ALTSVHTFPAVLQSSGLYSLSSVTVFPSSSLGTQTYICNVN HKPSNTKVDKKVEPKSCDKTCTCPCCPAPEAAGGSPVFLFPP KPKDTLMI SRTPEVTCVVSVSHEDPEVKFNWYVDGVEVHNA KTKPREEQYNSTYRVSVLTVLHQDWLNGKEYCKVKSNKALP APIETISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKG FYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVVD KSRWQQGNVFSCSVNHEALHNHYTQKSLSLSPG (see above)	169	(see above)	102	PSMB893 HC2 Construct ID: PBD000101318	
PS3B916	(see above)	169	(see above)	136	PSMB894 HC2 Construct ID: PBD000101318	
PS3B919	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSVAMSWVRQAPG KGLEWVSAISGGIGSTYYADSVKGRFTISRDNKNTLWLQMN SLRAEDTAVYYCAKDGVGATPYFYDWGQGLTVTVSSASTKG PSVFFPLAPSSKSTSGGTAALGCLVKDYFPEPPTVSWNSGALT SGVHTFPAVLQSSGLYSLSSVTVFPSSSLGTQTYICNVNHPK SNTKVDKVEPKSCDKTHTCPCPAPEAAGGSPVFLFPPKPK DTLMI SRTPEVTCVVSVSHEDPEVKFNWYVDGVEVHNAKTK PREEQYNSTYRVSVLTVLHQDWLNGKEYCKVKSNKALPAPI EKTIISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFPY SDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVDKSR WQGNVFSCSVNHEALHNHYTQKSLSLSPG (see above)	237		238	PSMB895 HC2 Construct ID: PBD000101320	
PS3B921	(see above)	237		272	PSMB896 HC2 Construct ID: PBD000101320	

PS3B920	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPG KGLEWVSAISGGSTYYADSVKGRFTISRDNKNTLYLQMN SLRAEDTAVYCAKDGVGATPYFDYWGQGLTVTVSSASTKG PSVFPFLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALT SGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKP SNTKVDKKVEPKSCDKHTCCPCPAPEAAGGSPVFLFPKPK DTIMI SRTPEVTCVWVSVDHEDPEVKFNWYVDGVEVHNAKTK PREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPI EKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYP SDIAVEWESNGQPENNYKTTTPPVLDSDGSFFLYSRLLTVDKSR WQQGNVFSCSVMHEALHNHYTQKSLSLSPG (see above)	305	RATLVCLISDFYPGAVTVAWKA DSSPVKAGVETTPSKQSNKKY AASSYLSLTPEQWKSHPSYSCQ VTHEGSTVEKTVAPTECS (see above)	238	PSMB897 HC2 Construct ID: PBD000101322
PS3B922	(see above)	305	(see above)	272	PSMB898 HC2 Construct ID: PBD000101322
PS3B912	EVQLVESGGGLVQPGGSLRLSCTASGFTFSSYAMSWVRQAPG KGLEWVSAISGGYGAPYYADTVKGRFTISRDNKNTLYLQMN SLRAEDTAVYCAKDGVGATPYFDYWGQGLTVTVSSASTKG PSVFPFLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALT SGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKP SNTKVDKKVEPKSCDKHTCCPCPAPEAAGGSPVFLFPKPK DTIMI SRTPEVTCVWVSVDHEDPEVKFNWYVDGVEVHNAKTK PREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPI EKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYP SDIAVEWESNGQPENNYKTTTPPVLDSDGSFFLYSRLLTVDKSR WQQGNVFSCSVMHEALHNHYTQKSLSLSPG (see above)	373	QSVLTQPPSPVSAAPGQKVTISC SGSSSNIGNNVSWYQQLPGTA PKLLIFDNNKRPSSGIFDRFSGS KSGTSATLGITGLQGTDEADYY CGTWDSSLAYVFGTGKVTIVL GQPKAAPSVTLFPPPSSEELQAN RATLVCLISDFYPGAVTVAWKA DSSPVKAGVETTPSKQSNKKY AASSYLSLTPEQWKSHPSYSCQ VTHEGSTVEKTVAPTECS	374	PSMB899 HC2 Construct ID: PBD000101324
PS3B930	(see above)	33	(see above)	34	PSMB889 HC2 Construct ID: PBD000101312
PS3B931	(see above)	33	(see above)	68	PSMB890 HC2 Construct ID: PBD000101312
PS3B926	(see above)	101	(see above)	102	PSMB891 HC2 Construct ID:

PS3B928	(see above)	101	(see above)		136	PBD000101316 PSMB892 HC2 Construct ID: PBD000101316
PS3B927	(see above)	169	(see above)		102	PSMB893 HC2 Construct ID: PBD000101318
PS3B929	(see above)	169	(see above)		136	PSMB894 HC2 Construct ID: PBD000101318
PS3B932	(see above)	237	(see above)		238	PSMB895 HC2 Construct ID: PBD000101320
PS3B934	(see above)	237	(see above)		272	PSMB896 HC2 Construct ID: PBD000101320
PS3B933	(see above)	305	(see above)		238	PSMB897 HC2 Construct ID: PBD000101322
PS3B935	(see above)	305	(see above)		272	PSMB898 HC2 Construct ID: PBD000101322
PS3B925	(see above)	373	(see above)		374	PSMB899 HC2 Construct ID: PBD000101324
PS3B1352	EVQLVESGGGLYQPGGSLRLSCAASGFTFSSYAMSWVRQAPG KGLEWVSAISGIGSTYYADSVKGRFTISRDNKNTLWLQMN SLRAEDTAVYCAKDGVGATPYEDYWGQGLTVTVSSASTKG PSVFPFLAPSSKSTSGGTAALGCLVKDYFPEFVTVSWNSGALT SGVHTFPAVLQSSGLYSLSVTVTPSSSLGTQTYICNVNHPK SNTKVDKKEPKSCDKHTCCPCPAPEAAAGPSVFLFPKPK DTLMI SRTPEVTCVVSVSHEDPEVKFNWYVDGVEVHNAKTK PREEQNSTYRVVSVLVTHQDWLNGREYKCKVSNKALPAPI EKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYP SDIAVEWESNGQPENNYKTTTPFVLDSDGSFFFLYSRLTVDKSR WQQGVPFSCSVNHEALNHYTKQSLSLSPGK	1242	(see above)		238	PSMB946 HC2 Construct ID: PBD000108502

PS3B1353	(see above)	1242	(see above)	272	PSMB947 HC2 Construct ID: PBD000108503
PS3B1354	EVOLVESGGGLVQPGGSLRLSQAASGFTFSSYAMSWVRQAPG KLEWVSAISGGSGSTYYADSVKGRFTISRDNKNTLYLQMN SLPAEDGAVYCAKDGATPYFYFDYWGQGLTVTVSSASTKG PSVEPLAPSSKSTSGGTAALGCLVKDYFPEPPTVSWNSGALT SGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKP SNTKVDKKVEPKSCDKHTCTCPCPAPEAAGGSPVFLPPPKPK DTIMI SRTPETVTCVWVSVDHEDPEVKFNWYVDGVEVHNAKTK PREEQYNSTYRVVSVLTVLHQDWLNGREYKCKVSNKALPAPI ERTISKAAGQPRPEQVYVTLPPSREEMTKNQVSLTCLVKGFYP SDIAVEWESNGQPENNYKTTTPPVLDSDGSGFFLYSRLTVDKSR WQGGNVFSCSVNHEALHNHYTQKSLSLSPGK (see above)	1404	(see above)	238	PSMB848 HC2 Construct ID: PBD000108504
PS3B1355	(see above)	1404	(see above)	272	PSMB849 HC2 Construct ID: PBD000108505
PS3B1356	(see above)	1242	(see above)	238	PSMB946 HC2 Construct ID: PBD000108502
PS3B1357	(see above)	1242	(see above)	272	PSMB847 HC2 Construct ID: PBD000108503
PS3B1358	(see above)	1404	(see above)	272	PSMB949 HC2 Construct ID: PBD000108505
PSMB937	(see above)	305	(see above)	238	PSMB948 HC2 Construct ID: PBD000101322

EXAMPLE 5.2: ANALYTICAL CHARACTERIZATION OF BISPECIFIC ANTI-PSMA_xCD3 ANTIBODIES

[00646] The protein concentration for each purified bispecific Ab was determined by measuring the absorbance at 280 nm on a NANODROP1000 spectrophotometer or Trinean DROPSense96 multichannel spectrophotometer and calculated using the extinction coefficient based on the amino acid sequence. SE HPLC of the purified antibodies was performed by running samples on a TOSOH TSKgel BioAssist G3SWxl column, in 0.2 M Na Phosphate pH 6.8 at 1 mL/min on a Waters Alliance HPLC for 20 min. The column effluent was monitored by absorbance at 280 nm. Anti-PSMA-CD3 bispecific antibodies were further analyzed by Intact Mass Analysis to determine appropriate formation of heterodimers.

EXAMPLE 6: EPIOTOPE MAPPING OF ANTI-PSMA_xCD3 ANTIBODIES

EXAMPLE 6.1: HDX-MS EPIOTOPE MAPPING

[00647] The epitope of two anti-PSMA/CD3 bispecific antibodies PS3B1352 and PS3B1353 were determined by hydrogen-deuterium exchange mass spectrometry (HDX-MS). Human PSMA antigen was used for epitope mapping experiment.

[00648] *On-Exchange Experiment for HDX-MS.* Briefly, purified PSMA antigen was incubated with and without PSMA/CD3 bispecific antibody PS3B1352 or PS3B1353 in deuterium oxide labeled buffer. On-exchange reaction was initiated by mixing 10 μ L of 6.0 μ M human PSMA with or without 7.3 μ M antibody and 30 μ L of H₂O or a deuterated buffer (20 mM MES, pH 6.4, 150 mM NaCl in 95% D₂O or 20 mM Tris, pH 8.4, 150 mM NaCl in 95% D₂O). The reaction mixture was incubated for 15, 50, 150, 500, or 1,500 s at 23°C and quenched at the different time points described by the addition of 8 M urea, 1 M TCEP, pH 3.0. The quenched solutions were analyzed immediately.

[00649] *General Procedure for HDX-MS Data Acquisition.* HDX-MS sample preparation was performed using an automated HDx system (LEAP Technologies, Morrisville, NC). The samples were loaded through a protease type XIII (protease from *Aspergillus saitoi*, type XIII) /pepsin column (NovaBioAssays Inc) at 600 μ L/min in 1% acetonitrile, 0.1% formic acid. Peptide fragments were loaded on the trap column

ACQUITY UPLC BEH C18 VanGuard Pre-column (Waters, Milford, MA), with a linear gradient of 8% to 28% acetonitrile in 0.1% formic acid at 100 μ L/min over 20 min and analyzed by mass spectrometry. The analytical column used was a Accucore C18 (Thermo Fisher Scientific, Waltham, MA).

[00650] **MS Data Acquisition.** Mass spectrometric analyses were carried out using an LTQ™ Orbitrap Fusion Lumos mass spectrometer (Thermo Fisher Scientific) with the capillary temperature at 275°C, resolution 150,000, and mass range (m/z) 300 – 2,000.

[00651] **HDX-MS Data Extraction.** BioPharma Finder 2.0 (Thermo Fisher Scientific) was used for the peptide identification of non-deuterated samples prior to the HDX experiments. HDExaminer version 2.4 (Sierra Analytics, Modesto, CA) was used to extract centroid values from the MS raw data files for the HDX experiments.

[00652] Incubation of PS3B1352 and PS3B1353 antibodies with soluble PSMA protein resulted in identical patterns of hydrogen exchange and overall protection (FIG. 1). Residues 597-598 (CR) were identified as epitopes because they were significantly protected upon binding (average differences in deuteration levels >10%). Residues 599 (D), 602-603 (VV) and 605 (R) were also identified as epitopes because they were marginally protected upon binding (average differences in deuteration levels 5% – 10%). They may show bigger protections under longer time points. The binding epitopes could be larger than the four segments described herein. Residues located around the four segments describes above, such as 593-594 (LP), 595 (F), 596 (D), 600 (Y), 601 (A), 604 (L), 606-607 (KY), 607-609 (YAD), and 610-612, (KIY), did not exchange at all in the time window employed, but could be protected if longer time points were to be monitored. The protected segments were mapped onto the three-dimensional structure of PSMA to visualize the binding epitopes (FIG. 2).

EXAMPLE 7: CHARACTERIZATION OF BISPECIFIC ANTI-PSMA $\times$ CD3 ANTIBODIES

EXAMPLE 7.1: BINDING AFFINITY OF BISPECIFIC ANTI-PSMA $\times$ CD3 ANTIBODIES TO HUMAN PSMA

[00653] The binding affinity of anti-PSMA to the recombinant human, cynomolgus or mouse PSMA was determined by surface plasmon resonance (SPR) using a Biacore 8K instrument. The bispecific antibodies were captured on a goat anti-Fc antibody-modified

C1 chip and titrated with 3-fold serial dilutions of PSMA antigen spanning concentrations from 1 nM to 100 nM or 11.1 nM to 100 nM. The association and dissociation were monitored for 3 and 15 minutes, respectively, using a flow rate of 50 μ L/min. Raw binding data was referenced by subtracting the analyte binding signals from blanks and analyzed using a 1:1 Langmuir binding model using the Biacore Insight evaluation software to obtain the kinetics, which were used to calculate the binding affinity. Kd data are summarized in Table 28. The anti-PSMA were captured using an anti-human Fc antibody and the antigens were injected in solution.

Table 28: Affinities (KD) for the interaction of anti-PSMAxCD3 bispecific antibodies or anti-PSMA antibodies with human PSMA as obtained by the BIACORE (SPR) method.

Anti-PSMAxCD3		Anti-PSMA			
Name	KD (M)	Name	ka (1/Ms)	kd (1/s)	KD (M)
PS3B917	$\leq 2.33\text{E-}11$	PSMB1045	2.09E+05	6.98E-05	3.33E-10
PS3B918	poor fit	PSMB1049	1.07E+05	7.02E-05	6.57E-10
PS3B913	$\leq 2.40\text{E-}10$	PSMB1051	7.71E+04	7.75E-05	1.01E-09
PS3B915	1.89E-09	PSMB1041	2.07E+05	4.35E-04	2.11E-09
PS3B914	Low/No binding	PSMB1068	1.51E+05	1.68E-04	1.11E-09
PS3B916	1.75E-09	PSMB1052	1.96E+05	3.15E-04	1.61E-09
PS3B919	8.23E-10	PSMB1069	1.39E+05	1.97E-04	1.42E-09
PS3B921	$\leq 8.64\text{E-}11$	PSMB1047	6.94E+05	1.75E-03	2.52E-09
PS3B920	1.35E-09	PSMB1075	5.26E+05	4.92E-04	9.36E-10
PS3B922	4.10E-10	PSMB1060	1.17E+05	1.19E-04	1.02E-09
PS3B912	1.74E-10				
PS3B930	$\leq 2.26\text{E-}11$				
PS3B931	poor fit				
PS3B926	$\leq 2.53\text{E-}10$				
PS3B928	2.37E-09				
PS3B927	1.11E-09				
PS3B929	1.83E-09				
PS3B932	7.87E-10				
PS3B934	$\leq 8.73\text{E-}11$				
PS3B933	1.44E-09				
PS3B935	4.54E-10				
PS3B925	1.69E-10				

[00654] In addition, the binding affinity of selected anti-PSMA-CD3 bispecific antibodies to recombinant human PSMA was compared to the binding affinity to cynomolgus PSMA by SPR. The antibodies were captured on a goat anti-Fc antibody-modified C1 chip and titrated with 3-fold serial dilutions of PSMA antigen spanning concentrations of 100 nM to 3.7 nM (human PSMA) or 100 nM to 3.7 nM, or 22.2-600 nM for cyno PSMA. The

association and dissociation were monitored for 3 and 60 minutes, respectively, using a flow rate of 50 μ L/min. Raw binding data was referenced by subtracting the analyte binding signals from blanks and analyzed using a 1:1 Langmuir binding model by the Biacore Insight evaluation software to obtain the kinetics, which were used to calculate the binding affinity. The kinetic parameter of binding of selected antibodies are shown in **Table 29**. The anti-PSMA were captured using an anti-human Fc antibody and the antigens were injected in solution.

Table 29: Affinities (K_D)* for the interaction of anti-PSMA antibodies with Human, cyno or Mouse PSMA as obtained by the Biacore (SPR) method.

Protein name	Antigen	Avg. k_a (1/Ms)	95% CI	Avg. k_d (1/s)	95% CI	Avg. K_D (M)	95% CI
PS3B1391	cy PSMA	1.57E+05	5.77E+04	1.96E-05	6.59E-06	1.26E-10	3.72E-11
PS3B1391	hu PSMA	4.80E+05		4.80E-05		1.00E-10	
PS3B1508	cy PSMA	1.46E+06		3.23E-03		2.20E-09	
PS3B1508	hu PSMA	2.32E+05		8.89E-05		3.82E-10	
PS3B1396	cy PSMA	5.87E+05		1.90E-03		5.07E-09	
PS3B1396	hu PSMA	2.24E+05		8.82E-05		3.95E-10	
PS3B1505	cy PSMA	4.29E+05		1.24E-03		5.01E-09	
PS3B1505	hu PSMA	2.32E+05		9.10E-05		3.93E-10	

*Data in row 1 is an average of 3. Data in other rows an average of 2.

EXAMPLE 7.2: THERMAL STABILITY OF BISPECIFIC ANTI-PSMA $\times$ CD3 ANTIBODIES

[00655] The thermal stability (conformational stability information including, T_m and T_{agg}) of selected anti-PSMA $\times$ CD3 antibodies was determined by nanoDSF method using a Prometheus instrument. Briefly, measurements were made by loading sample into 24 well capillary from a 384 well sample plate. Duplicate runs were performed. The thermal scans span from 20°C to 95°C at a rate of 1.0°C/minute. The data was processed to obtain integrated data and first derivation analysis for 330 nm, 350 nm, Ratio 330/350, and scatter data from which thermal transitions, onset of unfolding, T_m and T_{agg} were obtained and summarized in **Table 30**.

Table 30: Thermal stability data for bispecific anti-PSMAxCD3 antibodies as obtained using a nanoDSF instrument.

Name	T _{agg}	T _{m1}	T _{m2}	T _{m3}
PS3B917	68.2°C	63.5°C	68.8°C	
PS3B918	67.8°C	63.4°C	68.2°C	
PS3B913	75.1°C	63.4°C	76.3°C	
PS3B915	69.2°C	63.5°C		
PS3B914	64.5°C	61.9°C	67.0°C	75.8°C
PS3B916	59.1°C	59.9°C		
PS3B919	81.0°C	63.5°C	68.3°C	85.7°C
PS3B921	78.9°C	63.5°C	80.8°C	
PS3B920	82.6°C	63.4°C	87.5°C	
PS3B922	80.5°C	63.3°C	83.4°C	
PS3B912	77.1°C	63.6°C	78.4°C	
PS3B930	72.2°C	70.1°C		
PS3B931	72.2°C	69.5°C		
PS3B926	75.6°C	70.2°C	75.6°C	
PS3B928	72.2°C	69.1°C		
PS3B927	69.1°C	66.5°C	69.4°C	
PS3B929	69.1°C	58.0°C	68.0°C	
PS3B932	82.7°C	70.4°C	85.8°C	
PS3B934	79.3°C	70.4°C	80.9°C	
PS3B933	83.7°C	70.0°C	87.8°C	
PS3B935	81.5°C	70.2°C	83.7°C	
PS3B925	77.4°C	70.3°C	77.9°C	

EXAMPLE 7.3: BINDING OF BISPECIFIC PSMAxCD3 ANTIBODIES ON PSMA+ CELLS

[00656] Selected bispecific PSMAxCD3 antibodies were assessed for their ability to bind prostate cancer cell lines expressing PSMA.

[00657] 22RV1 and C4-2B cells were plated at 50,000 cells per well in 50 µl of assay medium (RPMI, 10% HI FBS) in V bottom plates. Serial dilutions of antibodies were prepared in assay medium with 50 µl of antibody dilutions added to the plates containing cells. The plates were incubated for 60 min at 37°C at which time 100 µl of staining buffer (Becton Dickinson Cat#554657) was added to all wells of each plate. The plates were centrifuged at 300 X G for 5 minutes and the medium was removed from the wells. 200 µl of staining buffer was added to all wells of each plate. The plates were centrifuged at 300 X G for 5 minutes and the medium was removed from the wells. 50 µl of 2 µg/ml AlexaFluor647-labeled goat anti-human Fc was added to all wells of the plates and the plates were incubated for 30 minutes at 4°C. 150 µl of staining buffer was added to all wells

of each plate. The plates were centrifuged at 300 X G for 5 minutes and the medium was removed from the wells. Two hundred microliters of running buffer (Staining buffer plus 1 mM EDTA, 0.1% pluronic acid) was added to all wells of the plates. The plates were centrifuged at 300 X G for 5 minutes and the medium was removed from the wells. Thirty microliters running buffer was added to all wells with cells and the plates were analyzed on the IQUE Plus instrument (Sartorius). Briefly, cells were gated on a FCS vs. SSC gate to eliminate cellular debris, and then the cell populations were gated on singlet cells. Antibody binding was assessed in the red laser channel. Signal (Mab plus secondary antibody) to background (secondary antibody only) ratios were calculated for each plate and the resultant data was plotted vs. bispecific antibody concentration in GeneData Screener using 4 parameter curve fitting to generate EC50 values, summarized in Table 31.

Table 31: EC50 values of bispecific PSMA x CD3 antibodies binding to PSMA-expressing cell lines in flow cytometry assays.

Name	22RV1 binding EC50 [M]	C4-2B binding EC50 [M]
PS3B917	7.50E-10	1.90E-09
PS3B918	6.00E-08	6.00E-08
PS3B913	3.42E-09	4.56E-09
PS3B915	6.00E-08	6.00E-08
PS3B914	6.00E-08	6.00E-08
PS3B916	6.00E-08	6.00E-08
PS3B919	2.49E-08	2.33E-08
PS3B921	4.03E-09	6.44E-09
PS3B920	6.00E-08	6.00E-08
PS3B922	3.47E-08	4.41E-08
PS3B912	5.40E-08	3.68E-08
PS3B930	1.06E-09	3.01E-09
PS3B931	6.00E-08	6.00E-08
PS3B926	2.90E-09	4.55E-09
PS3B928	6.00E-08	6.00E-08
PS3B927	6.00E-08	6.00E-08
PS3B929	5.48E-08	6.00E-08
PS3B932	4.87E-08	6.00E-08
PS3B934	4.96E-09	1.10E-08
PS3B933	6.00E-08	6.00E-08
PS3B935	3.95E-08	8.22E-09
PS3B925	2.12E-08	1.17E-08

[00658] *Binding of anti-PSMA/CD3 bispecific on PAN-T cells via Flow.* Human PAN-T Cells (Biological Specialty Corporation, Colmar, PA) were thawed and transferred to a 15 mL conical with DPBS. The cells were centrifuged 1300 rpm for 5 minutes. DPBS was

aspirated and cells were re-suspended in DPBS. The cells were counted using the Vi-cell XR cell viability analyzer and were plated at 100K/well in 100uL DPBS. The plate was centrifuged 1200 rpm for 3 minutes and washed 2x with DPBS. Cells were stained with Violet Live/Dead stain (Thermo-Fisher) and incubated at RT in the dark for 25 min. The cells were centrifuged and washed 2x with FACS staining buffer (BD Pharmingen). Test antibodies were diluted to a final starting concentration of 1 μ M in FACS staining buffer and 3-fold serial dilutions were prepared from the starting concentration for a total of 10 dilution points. The serially diluted test antibodies (100 μ L/ well) were added to the cells and incubated for 30 min at 37°C. Cells were washed 2x with FACS staining buffer and AlexaFluor 647-conjugated Donkey anti-human secondary antibody (Jackson ImmunoResearch) was added and allowed to incubate with the cells for 30 min at 4°C. Cells were washed 2x with FACS staining buffer and re-suspended in 100 μ L FACS Buffer. Cells were run on BD Celesta using FACS Diva software and analyzed using FLOWJO. **FIG. 3** shows that the PSMA/CD3 bispecific antibodies display differential CD3 cell binding profiles detected by flow cytometry.

[00659] Binding curves demonstrated in **Table 32** below and **FIG. 4** were generated against the prostate cell line C4-2B at 37°C in RPMI media plus 10% fetal bovine serum after 1 hr incubation. Molecule concentrations ranged from 500 – 0 nM over 12-points at 3-fold dilution. Selective binding to PSMA was validated using an isotype control. Values reported in the **Table 32** were generated by fitting the data to a four-parameter function for ligand binding generating values for y-min, y-max, EC50, and Hill. EC90 were calculated using the equation $EC90 = (90 - (100 - 90))^{(1/Hill)} * EC50$. All curves exhibited a similar Y-min with an average of $7.1 \pm 1.3E+4$ for all curves. None of the Y-min values deviated significantly from the average value. The average fitted Y-max value was $1.7 \pm 0.6E+6$. Molecule PSMB1069 exhibited a 2-fold higher binding signal from the average. None of the other molecules exhibited a significant difference from the average. These molecules exhibited an average $EC50 = 17 \pm 12$ nM.

Table 32: EC50 values of a

anti-PSMA antibodies binding to PSMA-expressing cell line C4-2B measured by flow cytometry assays.

Molecule	Y-min	Y-max	EC50 (nM)	EC90 (nM)
PSMB1041	7.2E+04	1.1E+06	16	59
PSMB1045	5.3E+04	1.8E+06	7	248
PSMB1047	7.9E+04	1.6E+06	22	71
PSMB1049	8.4E+04	1.5E+06	5	40
PSMB1051	6.7E+04	1.5E+06	10	76
PSMB1052	6.8E+04	1.5E+06	13	100
PSMB1060	8.6E+04	1.6E+06	11	59
PSMB1068	4.8E+04	1.9E+06	8	247
PSMB1069	8.7E+04	3.3E+06	44	254
PSMB1075	6.6E+04	1.3E+06	30	156

[00660] *Binding of anti-PSMA variants/CD3 bispecific on T cells via flow cytometry.*

C4-2B human prostate tumor cells were washed with DPBS and 0.25% trypsin was added to allow cells to detach. Media was added to neutralize trypsin and the cells were transferred to a 15 mL conical with DPBS. The cells were centrifuged 1300 rpm for 5 minutes. DPBS was aspirated and cells were re-suspended in DPBS. The cells were counted using the Vi-cell XR cell viability analyzer and were plated at 100K/well in 100 μ L DPBS. The plate was centrifuged 1200 rpm for 3 minutes and washed 2x with DPBS. Cells were stained with Violet Live/Dead stain (Thermo-Fisher) and incubated at RT in the dark for 25 min. The cells were centrifuged and washed 2x with FACS staining buffer (BD Pharmingen). Test antibodies were diluted to a final starting concentration of 100 nM in FACS staining buffer and 3-fold serial dilutions were prepared from the starting concentration for a total of 10 dilution points. The serially diluted test antibodies (100 μ L/ well) were added to the cells and incubated for 30 min at 37°C. Cells were washed 2x with FACS staining buffer and AlexaFluor 647-conjugated Donkey anti-human secondary antibody (Jackson ImmunoResearch) was added and allowed to incubate with the cells for 30 min at 4°C. Cells were washed 2x with FACS staining buffer and re-suspended in 100 μ L FACS Buffer. Cells were run on BD CELESTA using FACS Diva software and analyzed using FLOWJO. FIG. 5 shows that PSMA/CD3 bispecific antibodies display similar PSMA cell binding profiles detected by flow cytometry.

EXAMPLE 7.4: INTERNALIZATION OF PSMA

[00661] C4-2B human prostate tumor cells were washed with DPBS and 0.25% trypsin was added to allow cells to detach. Media was added to neutralize trypsin and the cells were transferred to a 15 mL conical with DPBS. The cells were centrifuged 1300 rpm for 5 minutes. DPBS was aspirated and cells were re-suspended in DPBS. The cells were counted using the Vi-cell XR cell viability analyzer and were plated at 40K/well in 50 μ L Phenol Red-Free RPMI + 10% HI FBS. The PSMA/CD3 bispecific or control antibodies were incubated with IncuCyte® Human Fab-fluor-pH Red Antibody labeling dye for 15 minutes then 50 μ L of conjugated PSMA/CD3:Fab-fluor-pH Red complex was added to the wells containing C4-2B cells. The plates were placed in an IncuCyte S3® (Essen) at 37°C with 5% CO₂ for 24 hours. The Ab:Fab-fluor complex that is internalized by the target cells is processed by acidic lysosomes, which produces the red fluorogenic signal that is captured and analyzed by the IncuCyte®. FIG. 6 shows that PSMA/CD3 bispecific antibodies internalize but at a lesser rate than the transferrin receptor.

EXAMPLE 7.5: T-CELL MEDIATED KILLING OF BISPECIFIC PSMA $\times$ CD3 ANTIBODIES ON PSMA+ CELLS VIA FLOW CYTOMETRY

[00662] Selected bispecific PSMA $\times$ CD3 antibodies were assessed for their ability to mediate T cell mediated killing of prostate cancer cells.

[00663] T cell mediated killing of the PSMA $\times$ CD3 bispecific antibodies was measured using an assay that indirectly measures cell killing via flow cytometry. Target cell population are identified base on cell viability. Test samples and controls were prepared at 20 nM in assay medium (10% RPMI, 10% HI FCS). Half log serial dilutions for an 11-point titration of compounds in sterile polypropylene plates were prepared. Additional wells were used for controls without compounds, T cells or tumor cell containing wells only in assay medium. C4-2B cells were harvested from the cell culture flasks and cells were resuspended in PBS. Cells were stained with 20 pM CFSE for 10 minutes at room temperature. 25 mL of HI FBS was added to stop the staining reaction and the cells were centrifuged at 300 X G for 5 minutes. Cells were diluted to 1 $\times$ 10⁶/ml and then plated as tumor target cells in 50 μ L assay medium for 50,000 cells/well into a V-bottom tissue culture treated polystyrene assay plate. 50 μ L/well of assay media was added to the control wells that did not receive tumor cells. Human PAN-T cell vials were thawed in a water bath set at 37°C and washed twice by adding 10 ml assay medium and centrifuging at 400 X G

for 5 minutes. T cells were resuspended to $1 \times 10^6/\text{mL}$ in assay medium and 50 μL containing 50,000/well were added to the assay plates containing tumor target cells. 50 μL /well assay media was added to the control wells that did not receive tumor cells. 100 μL /well of serially diluted antibodies were added to the assay plates containing cell mixture of target and effector cells. Plates were incubated at 37°C , 5% CO_2 in a humidified cell culture incubator for 72 hours.

[00664] Following the incubation assay plates were centrifuged at 500 X G for 5 minutes and medium was removed from the wells. 150 μL DPBS was added to each well and the plates were centrifuged at 500 X G for 5 minutes and medium was removed from the wells. The cultures were assessed using flow cytometry on the INTELLICYT IQ Plus for viable tumor cells using near IR live/dead stain. T cell activation was assessed using a brilliant violet-labeled anti-CD25 MAB. Cells were gated in an FSC vs SSC gate to eliminate debris. Tumor cells were identified as CFSE positive cells. T cells were identified as CFSE negative cells. Tumor cell viability was calculated as number of live/dead stain positive tumor cells as a percentage of total CFSE cells. Activated T cells were calculated as the number of CD25 positive cells as a percentage of the total number of the live CFSE negative population. The data for percent dead tumor cells and activated T cells were plotted vs. antibody concentration in Gene Data Screener using 4 parameter curve fitting to generate EC50 values. Table 33 shows EC50 values for T cell activation and tumor cell killing.

Table 33: *In vitro* T cell mediated killing of tumor cells and T cell activation by bispecific PSMA x CD3 antibodies.

Name	Tumor killing EC50 [M]	T cell activation EC50 [M]
PS3B917	7.31E-12	4.48E-12
PS3B918	1.00E-08	1.00E-08
PS3B913	3.11E-10	1.55E-10
PS3B915	1.00E-08	1.00E-08
PS3B914	1.00E-08	1.00E-08
PS3B916	1.00E-08	1.00E-08
PS3B919	2.10E-10	1.87E-10
PS3B921	3.11E-11	2.11E-11
PS3B920	3.20E-09	1.77E-09
PS3B922	7.79E-11	5.85E-11
PS3B912	6.51E-11	4.73E-11
PS3B930	4.83E-12	2.40E-12
PS3B931	1.00E-08	1.00E-08
PS3B926	4.78E-11	3.45E-11

PS3B928	4.81E-10	1.84E-10
PS3B927	1.00E-08	1.00E-08
PS3B929	1.00E-08	1.00E-08
PS3B932	6.42E-12	1.20E-11
PS3B934	7.24E-12	4.81E-12
PS3B933	4.76E-11	6.01E-11
PS3B935	7.16E-12	6.72E-12
PS3B925	7.98E-12	4.59E-12

EXAMPLE 7.6: T-CELL MEDIATED KILLING OF BISPECIFIC PSMAxCD3 ANTIBODIES ON PSMA+ CELLS VIA INCUCYTE

[00665] Select bispecific PSMAxCD3 antibodies were assessed for their ability to mediate T cell mediated killing of prostate cancer cells via IncuCyte®-based cytotoxic assay.

[00666] *Healthy donor T cells.* PSMA+ C4-2B cells stably expressing red nuclear dye were generated to be used in the IncuCyte-based cytotoxicity assay. Frozen vials of healthy donor T cells (Biological Specialty Corporation, Colmar, PA) were thawed in a 37°C water bath, transferred to a 15 mL conical tube, and washed once with 5 mL phenol-red-free RPMI/ 10% HI FBS medium. The cells were counted using the VIACELL XR cell viability analyzer and the T cells were combined with target cells for a final effector T cell to target cell (E: T) ratio of 3:1. The cell mixture was combined in a 50 mL conical tube. The cell mixture (100 µL/ well) was added to a clear 96-well flat-bottom plate. Next, the test antibodies were diluted to a final starting concentration of 60 nM in phenol-red-free RPMI/ 10% HI FBS medium and 3-fold serial dilutions were prepared from the starting concentration for a total of 11 dilution points. The serially diluted test antibodies (100 µL/ well) were added to the combined cells. The plates were placed in either an IncuCyte® Zoom or an IncuCyte S3® (Essen) at 37°C with 5% CO₂ for 120 hours. The target cell lines stably express red nuclear dye, which is used to track the kinetics of target cell lysis. Percent cell growth inhibition (%) = (Initial viable target cell number- Current viable target cell number)/ Initial viable cell number * 100%. **Table 34** and **FIGS. 7A-7H** show cytotoxicity for C4-2B cells with increasing concentrations of anti-PSMA. Isolated PAN-T cells were co-incubated with PSMA+ C4-2B cells in the presence of bispecific PSMA/ T cell redirection antibodies for 120 hours.

Table 34: Bispecific anti-PSMA/ anti-T cell redirection antibodies evaluated in an IncuCyte®-based cytotoxicity assay.

Name	Cytotoxicity (C4-2B cells, 3:1 E:T ratio, 5 Day)				
	30 nM	10 nM	3.3 nM	1.1 nM	0.3 nM
PS3B1352	79%	85%	89%	88%	65%
PS3B1356	88%	83%	80%	55%	No lysis
PS3B1353	92%	95%	97%	98%	98%
PS3B1357	91%	96%	95%	96%	96%
PS3B1354	84%	72%	30%	No lysis	No lysis
PSMB937	41%	No lysis	No lysis	No lysis	No lysis
PS3B1355	94%	95%	96%	97%	92%
PS3B1358	88%	94%	93%	90%	68%

[00667] *Healthy PBMCs*. PSMA+ C4-2B human prostate tumor cells expressing red nuclear dye were generated to be used in the IncuCyte®-based cytotoxicity assay. Frozen vials of healthy PBMCs (Hemacare, Los Angeles, CA) were thawed in a 37°C water bath, transferred to a 15 mL conical tube, and washed once with 5 mL phenol-red-free RPMI/ 10% HI FBS medium. The cells were counted using the VIACELL XR cell viability analyzer and the PBMCs were combined with target cells for a final PBMC to target cell (E: T) ratio of 1:1. The cell mixture was combined in a 50 mL conical tube. The cell mixture (100 µL/ well) was added to a clear 96-well flat-bottom plate. Next, the test antibodies were diluted to a final starting concentration of 30 nM in phenol-red-free RPMI/ 10% HI FBS medium and 3-fold serial dilutions were prepared from the starting concentration for a total of 11 dilution points. The serially diluted test antibodies (100 µL/ well) were added to the combined cells. The plates were placed in either an IncuCyte® Zoom or an IncuCyte S3® (Essen) at 37°C with 5% CO₂ for 120 hours. The target cell lines stably express red nuclear dye, which is used to track the kinetics of target cell lysis. Percent cell growth inhibition (%) = (Initial viable target cell number- Current viable target cell number)/ Initial viable cell number * 100%. FIG. 8 shows that the PSMA/CD3 bispecific antibodies induce differential C4-2B cytotoxic effects.

EXAMPLE 7.7: EVALUATING CYTOKINE INDUCTION BY BISPECIFIC ANTI-PSMAxCD3 ANTIBODIES

[00668] Select bispecific PSMAxCD3 antibodies were assessed for their ability to induce cytokine release.

[00669] Supernatants collected from the in-vitro cytotoxicity experiment described above were analyzed using the Human Proinflammatory Panel 1 tissue culture kit (Meso Scale Discovery). Supernatants were thawed on wet ice, spun at 1,500 rpm for 5 minutes at 4°C, then placed on ice. The MULT-SPOT assay plates were pre-washed per the manufacturer's protocol. A standard curve was prepared by serial dilution of the provided calibrators in MSD Diluent 1. The standards and test antibody samples (25 µL/ well) were added to the pre-washed plates. Subsequent incubations and washes were all carried out per manufacturer's protocol. Assay plates were read on the SECTOR Imager 6000. IFN γ concentrations were quantified for each PSMAxCD3 bispecific antibody evaluated. FIG. 9 shows functional cytokine release by T cells activated by PSMAxCD3 antibodies.

EXAMPLE 7.8: T-CELL MEDIATED KILLING OF BISPECIFIC PSMAxCD3 ANTIBODIES ON PSMA+ CELLS VIA XCELLIGENCE

[00670] Select bispecific PSMAxCD3 antibodies were assessed for their ability to mediate T cell mediated killing of prostate cancer cells, C4-2B. C4-2B, a prostate cancer cell line expressing ~150,000 PSMA/cell was used at a 3:1 Effector to Target ratio (E:T), using three PAN-T donors. On day 0 of the experiment, xCelligence plates were blanked with 50 µl of growth media. Plates were then seeded with 20,000 C4-2B (50 µL out of 0.4×10^6 cells/ml) cells per well. Plates were then incubated on the xCelligence machine overnight. On day 1 of the experiment, three PAN-T donors were used to prepare the E:T ratio by adding 50 µL of 1.2×10^6 cells/mL (60,000 cells). Then 50 µL of the appropriate bispecific antibodies were added to the appropriate wells for each plate. CD3xnull was used as a control. Tumor/target only wells were assigned to be used for normalization in the percent cytotoxicity calculation. Final antibody concentrations were 50 nM, 10 nM, 2 nM, 0.4 nM, 80 pM and 0 nM. Plates were then placed in the XCELLIGENCE machine and impedance was recorded every 15 minutes for 120 hours. Percent cytotoxicity was calculated on the RTCA software using the equation $\% \text{ cytotoxicity} = [1 - (\text{NCI}) / (\text{AvgNCIR})] \times 100$, where NCI is the average cell

index of the well and AvgNCIR is the average cell index of the tumor only reference wells.

Table 35 summarizes cytotoxicity for each PSMA x CD3 bispecific molecule over time.

Table 35: Summary of % cytotoxicity at time point 120 hours for all four bispecific antibodies, for three PAN-T donors, at each dose concentration.

Donor ID	Name	50 nM	10 nM	2 nM	0.4 nM	80 pM
#1	PS3B1352	100%	100%	100%	100%	77%
	PS3B1353	100%	100%	100%	100%	100%
	PS3B1356	100%	100%	100%	100%	0%
	PS3B1357	100%	100%	100%	100%	100%
#2	PS3B1352	100%	100%	100%	100%	0%
	PS3B1353	100%	100%	100%	100%	100%
	PS3B1356	100%	100%	100%	24%	0%
	PS3B1357	100%	100%	100%	100%	100%
#3	PS3B1352	100%	100%	100%	100%	9%
	PS3B1353	100%	100%	100%	100%	100%
	PS3B1356	100%	100%	100%	100%	0%
	PS3B1357	100%	100%	100%	100%	100%

* * * * *

[00671] It will be appreciated by those skilled in the art that changes could be made to the embodiments described above without departing from the broad inventive concept thereof.

It is understood, therefore, that this invention is not limited to the particular embodiments disclosed, but it is intended to cover modifications within the spirit and scope of the present invention as defined by the present description.

What is claimed is:

1. An isolated antibody that binds to PSMA comprising:
 - (25) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:32;
 - (26) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:66;
 - (27) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100;
 - (28) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134;
 - (29) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an

amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3,
respectively, of a VL having an amino acid sequence of SEQ ID NO:100;

- (30) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134;
- (31) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236;
- (32) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
- (33) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236;
- (34) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;

- (35) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:371; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:372;
- (36) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:405; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:406;
- (37) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
- (38) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:473; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474;
- (39) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:507; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508;
- (40) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3,

- respectively, of a VH having an amino acid sequence of SEQ ID NO:541; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:542;
- (41) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:575; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:576;
- (42) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100;
- (43) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:643; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508;
- (44) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:677; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:678;
- (45) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:711; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an

- amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3,
respectively, of a VL having an amino acid sequence of SEQ ID NO:474;
- (46) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:745; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:746;
- (47) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:779; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:780; or
- (48) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:813; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:814.
2. The isolated antibody of claim 1, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences are according to the
(i) Kabat numbering system;
(ii) Chothia numbering system;
(iii) AbM numbering system;
(iv) Contact numbering system;
(v) IMGT numbering system or a combination thereof.
3. The isolated antibody of claim 1 or claim 2, further comprising a VH having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:31, 99, 167, 235, 303, 371, 405, 439, 473, 507, 541, 575, 643, 677, 711, 779 or 813 and a VL having an amino acid sequence at least 80% identical to the amino acid sequence

of SEQ ID NOs:32, 66, 100, 134, 236, 270, 372, 406, 474, 508, 542, 576, 678, 746, 780 or 814.

4. The isolated antibody of any one of claims 1 to 3, further comprising a heavy chain (HC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:33, 101, 169, 237, 305, 373, 407 or 441 and a light chain (LC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:34, 68, 102, 136, 238, 272, 374 or 408.
5. An isolated antibody that binds PSMA comprising
 - a. (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
 - b. (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of SEQ ID NOs:205, 206, and 411, respectively, and (ii) a VL comprising a VL CDR1, VL CDR2, and VL CDR3 having an amino acid sequence of SEQ ID NOs:242, 209, and 244, respectively;
 - c. (i) a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL having an amino acid sequence of SEQ ID NO:270; or
 - d. (i) a HC having an amino acid sequence of SEQ ID NO:441; and (ii) a LC having an amino acid sequence of SEQ ID NO:272.
6. The isolated antibody of any one of claims 1 to 5, wherein the antibody is a humanized antibody.
7. The isolated antibody of any one of claims 1 to 6, wherein the antibody is a human antibody.
8. The isolated antibody of any one of claims 1 to 7, wherein the binding domain that binds to PSMA is a scFv, an scFv dimer, a Fv, a Fab, a Fab, a F(ab')₂, a dsFv, a sdAb, a VHH or a single chain antibody.
9. The isolated antibody of any one of claims 1 to 8, wherein the antibody is an IgG antibody.

10. The isolated antibody of claim 9, wherein the IgG antibody is an IgG1, IgG2, IgG3, or IgG4 antibody.
11. The isolated antibody of any one of claims 1 to 10, wherein the antibody comprises a kappa light chain.
12. The isolated antibody of any one of claims 1 to 10, wherein the antibody comprises a lambda light chain.
13. The isolated antibody of any one of claims 1 to 12, wherein the antibody is a monoclonal antibody.
14. The isolated antibody of any one of claims 1 to 13, wherein the antibody binds a PSMA antigen.
15. The isolated antibody of any one of claims 1 to 14, wherein the antibody binds a PSMA epitope.
16. The isolated antibody of any one of claims 1 to 15, wherein the antibody specifically binds to PSMA.
17. The isolated antibody of any one of claims 1 to 16, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 form a binding site for an antigen of the PSMA.
18. The isolated antibody of any one of claims 1 to 16, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 form a binding site for an epitope of the PSMA.
19. The isolated antibody of any one of claims 1 to 18, wherein the PSMA is present on the surface of a cell.
20. The isolated antibody of claim 19, wherein the cell is a prostate cell.
21. The isolated antibody of claim 19, wherein the cell is a prostate cancer cell.
22. The isolated antibody of claim 19, wherein the cell is a renal cell.
23. The isolated antibody of claim 19, wherein the cell is a renal cancer cell.
24. The isolated antibody of claim 23, wherein the renal cancer is a metastatic renal cell carcinoma.

25. The isolated antibody of any one of claims 1 to 24, wherein the antibody is multivalent.
26. The isolated antibody of claim 25, wherein the antibody is capable of binding at least three antigens.
27. The isolated antibody of claim 25, wherein the antibody is capable of binding at least four antigens.
28. The isolated antibody of claim 25, wherein the antibody is capable of binding at least five antigens.
29. The isolated antibody of any one of claims 1 to 28, wherein the antibody is a multispecific antibody.
30. The isolated antibody of claim 29, wherein the antibody is a bispecific antibody.
31. The isolated antibody of claim 29, wherein the antibody is a trispecific antibody.
32. The isolated antibody of claim 29, wherein the antibody is a quadraspecific antibody.
33. A nucleic acid encoding the antibody of any one of claims 1 to 32.
34. A vector comprising the nucleic acid of claim 33.
35. A host cell comprising the vector of claim 34.
36. A kit comprising the vector of claim 34 and packaging for the same.
37. A kit comprising the antibody of any one of claims 1 to 32 and packaging for the same.
38. A pharmaceutical composition comprising the antibody of any one of claims 1 to 32, and a pharmaceutically acceptable carrier.
39. A method of producing the pharmaceutical composition of claim 38, comprising combining the antibody with a pharmaceutically acceptable carrier to obtain the pharmaceutical composition.
40. An isolated multispecific antibody, comprising:
 - (a) a first binding domain that binds to PSMA, wherein the first binding domain comprises:
 - (25) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3,

- respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:32;
- (26) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:31; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:66;
- (27) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100;
- (28) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:134;
- (29) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100;
- (30) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:167; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an

amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3,
respectively, of a VL having an amino acid sequence of SEQ ID NO:134;

- (31) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236;
- (32) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:235; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
- (33) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:236;
- (34) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:303; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
- (35) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:371; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:372;

- (36) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:405; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:406;
- (37) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270;
- (38) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:473; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474;
- (39) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:507; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508;
- (40) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:541; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:542;
- (41) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3,

- respectively, of a VH having an amino acid sequence of SEQ ID NO:575; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:576;
- (42) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:99; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:100;
- (43) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:643; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:508;
- (44) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:677; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:678;
- (45) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:711; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:474;
- (46) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:745; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an

amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3,
respectively, of a VL having an amino acid sequence of SEQ ID NO:746;

- (47) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:779; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:780; or
- (48) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:813; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:814; and
- (b) a second binding domain that binds to CD3, wherein the second binding domain comprises:
 - (7) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:1505; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:1464;
 - (8) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:915; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:916;
 - (9) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:983; and
(ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an

- amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3,
respectively, of a VL having an amino acid sequence of SEQ ID NO:984;
- (10) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:1463; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:1464;
- (11) (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:847; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:848; or
- (12) a scFv comprising a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, a VL CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, a VL CDR3 respectively, of a scFv having an amino acid sequence of SEQ ID NO:1524.
41. The isolated multispecific antibody of claim 40, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3 amino acid sequences are according to the
- (i) Kabat numbering system;
 - (ii) Chothia numbering system;
 - (iii) AbM numbering system;
 - (iv) Contact numbering system;
 - (v) IMGT numbering system or a combination thereof.
42. The isolated multispecific antibody of claim 40 or claim 41, wherein the first binding domain that binds PSMA comprises a heavy chain (HC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:33, 101, 169, 237, 305, 373, 407, 441, 1242, 1244, 1248, 1250, 1252, 1254, 1256, 1258, 1260, 1262, 1264, 1266, 1268 or 1270 and/or a light chain (LC) having an amino acid

sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:34, 68, 102, 136, 238, 272, 374 or 408.

43. The isolated multispecific antibody of any one of claims 40 to 42, wherein the first binding domain that binds PSMA comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of any one of SEQ ID NOs:1485-1500 or SEQ ID NOs:1526-1531.
44. The isolated multispecific antibody of any one of claims 40 to 43, wherein the first binding domain that binds PSMA comprises (i) a HC having an amino acid sequence of SEQ ID NO:441; and (ii) a LC having an amino acid sequence of SEQ ID NO:272.
45. The isolated multispecific antibody of any one of claims 40 to 44, wherein the second binding domain that binds CD3 comprises a heavy chain (HC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:849, 883, 917, 951, 985, 1019, 1504, 1455, 1192, 1194, 1167, 1218 or 1238 and/or a light chain (LC) having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:850, 918, 986, 1193, 1195 or 1219.
46. The isolated multispecific antibody of any one of claims 40 to 45, wherein the second binding domain that binds CD3 comprises a scFv having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NOs:1186, 1187, 1523 or 1524.
47. An isolated bispecific antibody comprising a first binding domain that binds PSMA and a second binding domain that binds CD3, wherein
 - a. the first binding domain that binds PSMA comprises (i) a VH comprising a VH CDR1, a VH CDR2, and a VH CDR3 having an amino acid sequence of a VH CDR1, a VH CDR2, and a VH CDR3, respectively, of a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL comprising a VL CDR1, a VL CDR2, and a VL CDR3 having an amino acid sequence of a VL CDR1, a VL CDR2, and a VL CDR3, respectively, of a VL having an amino acid sequence of SEQ ID NO:270, and the second binding domain that binds CD3 comprises a scFv comprising a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, and a VL CDR3 of a scFv having an amino acid sequence of SEQ ID NO:1524;

- b. the first binding domain that binds PSMA comprises a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:205, 206, 411, 242, 209 and 244, respectively, wherein the amino acid sequences are according to the Kabat numbering system; and the second binding domain that binds CD3 comprises a VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3 of SEQ ID NO:1467, 1468, 1506, 1470, 1471 and 1472, respectively; wherein the amino acid sequences are according to the Kabat numbering system;
 - c. the first binding domain that binds PSMA comprises (i) a VH having an amino acid sequence of SEQ ID NO:439; and (ii) a VL having an amino acid sequence of SEQ ID NO:270, and the second binding domain that binds CD3 comprises a scFv of SEQ ID NO:1524, respectively; and/or
 - d. the first binding domain that binds PSMA comprises (i) a HC2 having an amino acid sequence of SEQ ID NO:441; and (ii) a LC2 having an amino acid sequence of SEQ ID NO:272, respectively; and the second binding domain that binds CD3 comprises a HC1 of SEQ ID NO:1455.
48. The isolated multispecific antibody of any one of claims 40 to 47, wherein the antibody is a humanized antibody.
49. The isolated multispecific antibody of any one of claims 40 to 48, wherein the antibody is a human antibody.
50. The isolated multispecific antibody of any one of claims 40 to 49, wherein the first binding domain, the second binding domain and/or the first and second is a scFv, an scFv dimer, a Fv, a Fab, a Fab, a F(ab')₂, a dsFv, a sdAb, a VHH or a single chain antibody.
51. The isolated multispecific antibody of any one of claims 40 to 50, wherein the antibody is an IgG antibody.
52. The isolated multispecific antibody of claim 51, wherein the IgG antibody is an IgG1, IgG2, IgG3, or IgG4 antibody.
53. The isolated multispecific antibody of any one of claims 40 to 52, wherein the antibody comprises a kappa light chain.

54. The isolated multispecific antibody of any one of claims 40 to 52, wherein the antibody comprises a lambda light chain.
55. The isolated multispecific antibody of any one of claims 40 to 54, wherein the antibody is a monoclonal antibody.
56. The isolated multispecific antibody of any one of claims 40 to 55, wherein the first binding domain binds a PSMA antigen.
57. The isolated multispecific antibody of any one of claims 40 to 55, wherein first binding domain binds a PSMA epitope.
58. The isolated multispecific antibody of any one of claims 40 to 57, wherein the first binding domain specifically binds to PSMA.
59. The isolated multispecific antibody of any one of claims 40 to 58, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the first binding domain form a binding site for an antigen of the PSMA.
60. The isolated multispecific antibody of any one of claims 40 to 58, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the first binding domain form a binding site for an epitope of the PSMA.
61. The isolated multispecific antibody of any one of claims 40 to 60, wherein the second binding domain binds a CD3 antigen.
62. The isolated multispecific antibody of any one of claims 40 to 61, wherein the second binding domain binds a CD3 epitope.
63. The isolated multispecific antibody of any one of claims 40 to 62, wherein the second binding domain specifically binds to CD3.
64. The isolated multispecific antibody of any one of claims 40 to 63, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the second binding domain form a binding site for an antigen of the CD3.
65. The isolated multispecific antibody of any one of claims 40 to 63, wherein the VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of the second binding domain form a binding site for an epitope of the CD3.
66. The isolated multispecific antibody of any one of claims 40 to 65, wherein the PSMA is present on the surface of a cell.

67. The isolated multispecific antibody of claim 66, wherein the cell is a prostate cell.
68. The isolated multispecific antibody of claim 66, wherein the cell is a prostate cancer cell.
69. The isolated multispecific antibody of claim 66, wherein the cell is a renal cell.
70. The isolated multispecific antibody of claim 66, wherein the cell is a renal cancer cell.
71. The isolated multispecific antibody of claim 66, wherein the renal cancer is a metastatic renal cell carcinoma.
72. The isolated multispecific antibody of any one of claims 40 to 71, wherein the antibody is a bispecific antibody.
73. The isolated multispecific antibody of any one of claims 40 to 71, wherein the antibody is a trispecific antibody.
74. The isolated multispecific antibody of any one of claims 40 to 71, wherein the antibody is a quadraspecific antibody.
75. A nucleic acid encoding the multispecific antibody of any one of claims 40 to 74.
76. A vector comprising the nucleic acid of claim 75.
77. A host cell comprising the vector of claim 76.
78. A kit comprising the vector of claim 76 and packaging for the same.
79. A kit comprising the multispecific antibody of any one of claims 40 to 74 and packaging for the same.
80. A pharmaceutical composition comprising the multispecific antibody of any one of claims 40 to 74, and a pharmaceutically acceptable carrier.
81. A method of producing the pharmaceutical composition of claim 80, comprising combining the multispecific antibody with a pharmaceutically acceptable carrier to obtain the pharmaceutical composition.
82. A method of directing a CD3-expressing T cell to a PSMA-expressing target cell, comprising contacting the T cell with the multispecific antibody of any one of claims 40 to 74.
83. The method of claim 82, wherein the contacting directs the T cell to the target cell.

84. A method of inhibiting the growth or proliferation of a PSMA-expressing target cell, comprising contacting the target cell with the multispecific antibody of any one of claims 40 to 74.
85. The method of claim 84, wherein the contacting is in the presence of CD3-expressing T cells.
86. A method of eliminating a PSMA-expressing target cell in a subject, comprising administering to the subject an effective amount of the multispecific antibody of any one of claims 40 to 74.
87. The method of claim 86, wherein the subject has a disease or disorder of the prostate.
88. The method of claim 86, wherein the subject has prostate inflammation.
89. The method of claim 86, wherein the subject has Benign prostatic hyperplasia.
90. The method of claim 86, wherein the subject has a prostate cancer.
91. The method of claim 90, wherein the subject has metastatic castration-resistant prostate cancer (mCRPC).
92. The method of any one of claims 86 to 91, wherein the target cell expresses PSMA on the cell surface.
93. The method of any one of claims 86 to 91, wherein the target cell is a prostate cell.
94. The method of any one of claims 86 to 91, wherein the target cell is a prostate cancer cell.
95. The method of claim 86, wherein the subject has a renal disease or disorder.
96. The method of claim 86, wherein the subject has a renal cancer.
97. The method of claim 86, wherein the subject has a metastatic renal cell carcinoma.
98. The method of any one of claims 86 or 95 to 97, wherein the target cell expresses PSMA on the cell surface.
99. The method of any one of claims 86 or 95 to 97, wherein the target cell is a renal cell.
100. The method of any one of claims 86 or 95 to 97, wherein the target cell is a renal cancer cell.
101. The method of any one of claims 86 or 95 to 97, wherein the target cell is a metastatic renal cell carcinoma cell.

102. The method of any one of claims 86 to 101, wherein the subject is a subject in need thereof.
103. A method of treating a disease or disorder in a subject, comprising administering to the subject an effective amount of the multispecific antibody of any one of claims 40 to 74.
104. The method of claim 103, wherein the disease or disorder is a disease or disorder of the prostate.
105. The method of claim 103, wherein the disease or disorder of the prostate is prostate inflammation.
106. The method of claim 103, wherein the disease or disorder of the prostate is Benign prostatic hyperplasia.
107. The method of claim 103, wherein the disease or disorder of the prostate is a prostate cancer.
108. The method of claim 107, wherein the disease or disorder is metastatic castration-resistant prostate cancer (mCRPC).
109. The method of claim 103, wherein the disease or disorder is a renal disease or disorder.
110. The method of claim 103, wherein the renal disease or disorder is a renal cancer.
111. The method of claim 103, wherein the renal disease or disorder is a metastatic renal cell carcinoma.
112. The method of any one of claims 103 to 111, wherein the subject is a subject in need thereof.