

METHODS FOR PROCESSING NUCLEIC ACID MOLECULES**CROSS REFERENCE**

[0001] This application claims the benefit of U.S. Provisional Application No. 62/804,644, filed February 12, 2019, which application is entirely incorporated herein by reference.

BACKGROUND

[0002] A sample may be processed for various purposes, such as detection, identification, quantitation, and characterization of a type of moiety within the sample. The sample may be a biological sample. Biological samples may be processed, such as for detection of a disease (e.g., cancer) or identification of a particular species. There are various approaches for processing samples, such as polymerase chain reaction (PCR) and sequencing.

[0003] Biological samples may be processed within various reaction environments, such as partitions. Partitions may be wells or droplets. Droplets or wells may be employed to process biological samples in a manner that enables the biological samples to be partitioned and processed separately. For example, such droplets may be fluidically isolated from other droplets, enabling accurate control of respective environments in the droplets.

[0004] Biological samples in partitions may be subjected to various processes, such as chemical processes or physical processes. Samples in partitions may be subjected to heating or cooling, or chemical reactions, such as to yield species that may be qualitatively or quantitatively processed.

[0005] A biological sample may comprise one or more nucleic acid molecules, such as one or more deoxyribonucleic acid (DNA) molecules and/or one or more ribonucleic acid (RNA) molecules. Methods of processing such a biological sample may vary depending on the types of nucleic acid molecules included therein.

SUMMARY

[0006] The present disclosure provides methods, systems, and kits for processing multiple different types of nucleic acid molecules in tandem. The methods provided herein may allow for analysis of both deoxyribonucleic acid (DNA) molecules and ribonucleic acid (RNA) molecules originating from the same cells or cell beads or components thereof (e.g., nuclei). Analysis of different types of nucleic acid molecules may be performed simultaneously or near simultaneously. The methods provided herein may comprise use of a partitioning scheme in which materials (e.g., different types of target nucleic acid molecules, such as target nucleic acid molecules included within a cell, cell bead, or nucleus) are distributed between a plurality of partitions, such as a plurality of droplets or wells. Materials (e.g., target nucleic acid molecules)

may be co-partitioned with one or more reagents to facilitate processing of target nucleic acid molecules, such as one or more enzymes, beads (e.g., beads comprising nucleic acid barcode molecules), primers, oligos, lysing or permeabilizing agents, buffers, or other reagents. The methods provided herein may comprise generating a barcoded nucleic acid product (e.g., within a partition of a plurality of partitions) corresponding to each of various different target nucleic acid molecules (e.g., DNA and RNA molecules), prior to subjecting the barcoded nucleic acid products to one or more amplification processes (e.g., polymerase chain reaction (PCR), which may optionally be performed in bulk).

[0007] Target nucleic acid molecules (e.g., DNA and RNA molecules) may initially be included in a cell, cell bead, or cell nucleus. Accordingly, the methods provided herein provide sample preparation techniques that permit sequencing of nucleic acid molecules from single cells, cell beads, and nuclei of interest.

[0008] In eukaryotic genomes, chromosomal DNA winds itself around histone proteins (i.e., “nucleosomes”), thereby forming a complex known as chromatin. The tight or loose packaging of chromatin contributes to the control of gene expression. Tightly packed chromatin (“closed chromatin”) is usually not permissive for gene expression while more loosely packaged, accessible regions of chromatin (“open chromatin”) is associated with the active transcription of gene products. Methods for probing genome-wide DNA accessibility have proven extremely effective in identifying regulatory elements across a variety of cell types and quantifying changes that lead to both activation or repression of gene expression.

[0009] One such method is the Assay for Transposase Accessible Chromatin with high-throughput sequencing (ATAC-seq). The ATAC-seq method probes DNA accessibility with an artificial transposon, which inserts specific sequences into accessible regions of chromatin. Because the transposase can only insert sequences into accessible regions of chromatin not bound by transcription factors and/or nucleosomes, sequencing reads can be used to infer regions of increased chromatin accessibility.

[0010] Traditional approaches to the ATAC-seq methodology requires large pools of cells, processes cells in bulk, and result in data representative of an entire cell population, but lack information about cell-to-cell variation inherently present in a cell population (see, *e.g.*, Buenrostro, *et al.*, Curr. Protoc. Mol. Biol., 2015 Jan 5; 21.29.1–21.29.9). While single cell ATAC-seq (scATAC-seq) methods have been developed, they suffer from several limitations. For example, scATAC-seq methods that utilize sample pooling, cell indexing, and cell sorting (see, *e.g.*, Cusanovich, *et al.*, Science, 2015 May 22; 348(6237):910-14) result in high variability and a low number of reads associated with any single cell. Other scATAC-seq

methods that utilize a programmable microfluidic device to isolate single cells and perform scATAC-seq in nanoliter reaction chambers (see, *e.g.*, Buenrostro, *et al.*, Nature, 2015 Jul 23;523(7561):486-90) are limited by the throughput of the assay and may not generate personal epigenomic profiles on a timescale compatible with clinical decision-making.

[0011] In an aspect, the present disclosure provides a method for processing a nucleic acid sample, comprising: (a) providing a biological particle comprising a deoxyribonucleic acid (DNA) molecule and a ribonucleic acid (RNA) molecule within a partition among a plurality of partitions, wherein the partition comprises (i) a first nucleic acid barcode molecule, (ii) a second nucleic acid barcode molecule, and (iii) a splint sequence, wherein the first nucleic acid barcode molecule and the second nucleic acid barcode molecule comprise a common barcode sequence, wherein the first nucleic acid barcode molecule comprises an overhang sequence, and wherein the splint sequence comprises a sequence complementary to the DNA molecule and a sequence complementary to the overhang sequence; (b) within the partition, using the DNA molecule, the first nucleic acid barcode molecule, and the splint sequence to generate a first barcoded nucleic acid product corresponding to the DNA molecule and using the RNA molecule and the second nucleic acid barcode molecule to generate a second barcoded nucleic acid product corresponding to the RNA molecule, wherein the first barcoded nucleic acid product and the second barcoded nucleic acid product comprise the common barcode sequence, or a complement thereof; and (c) recovering the first barcoded nucleic acid product and the second barcoded nucleic acid product from the partition.

[0012] In some embodiments, the biological particle is a cell, cell bead, or cell nucleus. In some embodiments, the method further comprises lysing or permeabilizing the biological particle within the partition to provide access to the DNA molecule and the RNA molecule therein.

[0013] In some embodiments, the method further comprises, prior to (a), processing an open chromatin structure of the nucleic acid sample with a transposase to yield the DNA molecule. In some embodiments, the transposase is included in a transposase-nucleic acid complex that comprises (i) a first nucleic acid molecule comprising a first transposon end sequence and a first sequencing primer or portion thereof, or a complement thereof and (ii) a second nucleic acid molecule comprising a second transposon end sequence and a second sequencing primer or portion thereof, or a complement thereof. In some embodiments, the first transposon end sequence and the second transposon end sequence are the same, and wherein the first transposon end sequence and the second transposon end sequence are each hybridized to complementary sequences.

[0014] In some embodiments, generating the first barcoded nucleic acid product comprises (i) subjecting the DNA molecule and the splint sequence to conditions sufficient to hybridize to the sequence of the DNA molecule to the splint sequence and (ii) subjecting the first nucleic acid barcode molecule and the splint sequence to conditions sufficient to hybridize the overhang sequence of the first nucleic acid barcode molecule to the splint sequence. In some embodiments, the splint sequence is hybridized to the sequence of the DNA molecule outside of the partition. In some embodiments, the method further comprises ligating the first nucleic acid barcode molecule to the DNA molecule.

[0015] In some embodiments, the first nucleic acid barcode molecule comprises a flow cell sequence, or complement thereof; a barcode sequence; a sequencing primer, or portion thereof; a unique molecular identifier sequence; or a combination thereof.

[0016] In some embodiments, the DNA molecule comprises one or more gaps, and further comprising, prior to (c), subjecting the DNA molecule to an extension process to fill the one or more gaps.

[0017] In some embodiments, the second nucleic acid barcode molecule comprises a unique molecular identifier sequence, a barcode sequence, a sequencing primer or portion thereof, or a combination thereof.

[0018] In some embodiments, the method further comprises, prior to (b), within the partition, reverse transcribing the RNA molecule to provide a complementary DNA (cDNA) molecule. In some embodiments, prior to reverse transcribing the RNA molecule, the second nucleic acid barcode molecule is hybridized to a sequence of a precursor of the RNA molecule. In some embodiments, the precursor comprises a polyA sequence, and wherein the second nucleic acid barcode molecule comprises a polyT sequence configured to hybridize to the polyA sequence. In some embodiments, the method further comprises appending an additional sequence to the cDNA molecule, wherein the additional sequence is a poly(C) sequence. In some embodiments, the method further comprises contacting the cDNA molecule with a template-switching oligonucleotide under conditions sufficient to hybridize the template-switching oligonucleotide to the cDNA molecule, wherein the template-switching oligonucleotide hybridizes to the additional sequence of the cDNA molecule. In some embodiments, the method further comprises extending the cDNA molecule to generate an extended cDNA molecule, which extended cDNA molecule comprises sequences complementary to sequences of the template-switching oligonucleotide. In some embodiments, the template-switching oligonucleotide comprises a sequencing primer or portion thereof, or complement thereof; a unique molecular identifier sequence, or complement thereof; or a combination thereof. In some embodiments, the

template-switching oligonucleotide is the second nucleic acid barcode molecule, and wherein the extended cDNA molecule is the second barcoded nucleic acid product.

[0019] In some embodiments, the first nucleic acid barcode molecule and the second nucleic acid barcode molecule comprise a common nucleic acid sequence.

[0020] In some embodiments, the partition is a droplet among a plurality of droplets. In some embodiments, the partition is a droplet and wherein (c) comprises breaking or disrupting the droplet.

[0021] In some embodiments, the method further comprises, subsequent to (c), using (i) the first barcoded nucleic acid product to generate a first plurality of amplification products and (ii) the second barcoded nucleic acid product to generate a second plurality of amplification products. In some embodiments, generating the first plurality of amplification products and the second plurality of amplification products comprises performing one or more nucleic acid amplification reactions using the first barcoded nucleic acid product or the second barcoded nucleic acid product. In some embodiments, performing the one or more nucleic acid amplification reactions comprises attaching one or more flow cell sequences to the first barcoded nucleic acid product or the second barcoded nucleic acid product, or a derivative thereof.

[0022] In some embodiments, the first nucleic acid barcode molecule and the second nucleic acid barcode molecule are coupled to a bead. In some embodiments, the bead is a gel bead. In some embodiments, the first nucleic acid barcode molecule is coupled to the bead via a first labile moiety and the second nucleic acid barcode molecule is coupled to the bead via a second labile moiety. In some embodiments, the first nucleic acid barcode molecule and the second nucleic acid barcode molecule are releasably coupled to the bead. In some embodiments, the first nucleic acid barcode molecule is releasable from the bead upon application of a first stimulus and the second nucleic acid barcode molecule is releasable from the bead upon application of a second stimulus.

[0023] In another aspect, the present disclosure provides a method for processing a nucleic acid sample, comprising: (a) providing a biological particle comprising a deoxyribonucleic acid (DNA) molecule and a ribonucleic acid (RNA) molecule within a partition among a plurality of partitions, wherein the partition comprises a first nucleic acid barcode molecule and a second nucleic acid barcode molecule, wherein the first nucleic acid barcode molecule and the second nucleic acid barcode molecule comprise a common barcode sequence; (b) within the partition, generating (i) a first barcoded nucleic acid product from the DNA molecule and the first nucleic acid barcode molecule and (ii) a second barcoded nucleic acid product from the RNA molecule and the second nucleic acid barcode molecule, wherein the first barcoded nucleic acid product

and the second barcoded nucleic acid product comprise the common barcode sequence, or a complement thereof; (c) recovering the first barcoded nucleic acid product and the second barcoded nucleic acid product from the partition; (d) processing the first barcoded nucleic acid product to generate a processed nucleic acid product comprising a dA tail; and (e) using (i) the processed nucleic acid product to generate a first plurality of amplification products and (ii) the second barcoded nucleic acid product to generate a second plurality of amplification products.

[0024] In some embodiments, the biological particle is a cell, cell bead, or cell nucleus. In some embodiments, the method further comprises lysing or permeabilizing the biological particle within the partition to provide access to the DNA molecule and the RNA molecule therein.

[0025] In some embodiments, the method further comprises, prior to (a), processing an open chromatin structure of the nucleic acid sample with a transposase to yield the DNA molecule. In some embodiments, the transposase is included in a transposase-nucleic acid complex that comprises (i) a first nucleic acid molecule comprising a first transposon end sequence and a first sequencing primer or portion thereof, or a complement thereof and (ii) a second nucleic acid molecule comprising a second transposon end sequence and a second sequencing primer or portion thereof, or a complement thereof. In some embodiments, the first transposon end sequence and the second transposon end sequence are the same, and wherein the first transposon end sequence and the second transposon end sequence are each hybridized to complementary sequences.

[0026] In some embodiments, the first nucleic acid barcode molecule comprises a sequencing primer or portion thereof, or complement thereof. In some embodiments, (b) comprises subjecting the DNA molecule to conditions sufficient to hybridize the first nucleic acid barcode molecule to the DNA molecule. In some embodiments, the method further comprises (i) extending the first nucleic acid barcode molecule hybridized to the DNA molecule and (ii) ligating the first nucleic acid barcode molecule to the DNA molecule to generate the first barcoded nucleic acid product.

[0027] In some embodiments, the method further comprises ligating the first nucleic acid barcode molecule to the DNA molecule.

[0028] In some embodiments, the first nucleic acid barcode molecule comprises a flow cell sequence, or complement thereof; a barcode sequence; a unique molecular identifier sequence; or a combination thereof.

[0029] In some embodiments, the DNA molecule comprises one or more gaps, and further comprising, prior to (c), subjecting the DNA molecule to an extension process to fill the one or more gaps.

[0030] In some embodiments, the second nucleic acid barcode molecule comprises a unique molecular identifier sequence, a barcode sequence, a sequencing primer or portion thereof, or a combination thereof.

[0031] In some embodiments, the method further comprises, prior to (b), within the partition, reverse transcribing the RNA molecule to provide a complementary DNA (cDNA) molecule. In some embodiments, prior to reverse transcribing the RNA molecule, the second nucleic acid barcode molecule is hybridized to a sequence of a precursor of the RNA molecule. In some embodiments, the precursor comprises a polyA sequence, and wherein the second nucleic acid barcode molecule comprises a polyT sequence configured to hybridize to the polyA sequence. In some embodiments, the method further comprises appending an additional sequence to the cDNA molecule, wherein the additional sequence is a poly(C) sequence. In some embodiments, the method further comprises contacting the cDNA molecule with a template-switching oligonucleotide under conditions sufficient to hybridize the template-switching oligonucleotide to the cDNA molecule, wherein the template-switching oligonucleotide hybridizes to the additional sequence of the cDNA molecule. In some embodiments, the method further comprises extending the cDNA molecule to generate an extended cDNA molecule, which extended cDNA molecule comprises sequences complementary to sequences of the template-switching oligonucleotide. In some embodiments, the template-switching oligonucleotide comprises a sequencing primer or portion thereof, or complement thereof; a unique molecular identifier sequence, or complement thereof; or a combination thereof. In some embodiments, the template-switching oligonucleotide is the second nucleic acid barcode molecule, and wherein the extended cDNA molecule is the second barcoded nucleic acid product. In some embodiments, the first nucleic acid barcode molecule and the second nucleic acid barcode molecule comprise a common nucleic acid sequence.

[0032] In some embodiments, the partition is a droplet among a plurality of droplets. In some embodiments, the partition is a droplet and wherein (c) comprises breaking or disrupting the droplet.

[0033] In some embodiments, (e) comprises performing one or more nucleic acid amplification reactions using the first barcoded nucleic acid product or the second barcoded nucleic acid product. In some embodiments, (e) comprises attaching one or more flow cell

sequences to the first barcoded nucleic acid product or the second barcoded nucleic acid product, or a derivative thereof.

[0034] In some embodiments, the first nucleic acid barcode molecule and the second nucleic acid barcode molecule are coupled to a bead. In some embodiments, the bead is a gel bead. In some embodiments, the first nucleic acid barcode molecule is coupled to the bead via a first labile moiety and the second nucleic acid barcode molecule is coupled to the bead via a second labile moiety. In some embodiments, the first nucleic acid barcode molecule and the second nucleic acid barcode molecule are releasably coupled to the bead. In some embodiments, the first nucleic acid barcode molecule is releasable from the bead upon application of a first stimulus and the second nucleic acid barcode molecule is releasable from the bead upon application of a second stimulus.

[0035] In a further aspect, the present disclosure provides a method for processing a nucleic acid sample, comprising: (a) providing a biological particle comprising a deoxyribonucleic acid (DNA) molecule and a ribonucleic acid (RNA) molecule within a partition among a plurality of partitions, wherein the partition comprises a first nucleic acid barcode molecule and a second nucleic acid barcode molecule, wherein the first nucleic acid barcode molecule and the second nucleic acid barcode molecule comprise a common barcode sequence, and wherein the DNA molecule comprises a first hairpin moiety at a first end and a second hairpin moiety at a second end; (b) within the partition, generating (i) a first barcoded nucleic acid product from the DNA molecule and the first nucleic acid barcode molecule and (ii) a second barcoded nucleic acid product from the RNA molecule and the second nucleic acid barcode molecule, wherein the first barcoded nucleic acid product and the second barcoded nucleic acid product comprise the common barcode sequence, or a complement thereof; (c) recovering the first barcoded nucleic acid product and the second barcoded nucleic acid product from the partition; and (d) using (i) the first barcoded nucleic acid product to generate a first plurality of amplification products and (ii) the second barcoded nucleic acid product to generate a second plurality of amplification products.

[0036] In some embodiments, the biological particle is a cell, cell bead, or cell nucleus. In some embodiments, the method further comprises lysing or permeabilizing the biological particle within the partition to provide access to the DNA molecule and the RNA molecule therein.

[0037] In some embodiments, the method further comprises prior to (a), processing an open chromatin structure of the nucleic acid sample with a transposase to yield the DNA molecule. In some embodiments, the transposase is included in a transposase-nucleic acid complex that

comprises (i) a first nucleic acid molecule comprising a first transposon end sequence and a first sequencing primer or portion thereof, or a complement thereof and (ii) a second nucleic acid molecule comprising a second transposon end sequence and a second sequencing primer or portion thereof, or a complement thereof. In some embodiments, the first transposon end sequence and the second transposon end sequence are the same, and wherein the first transposon end sequence and the second transposon end sequence are each hybridized to complementary sequences.

[0038] In some embodiments, the first nucleic acid barcode molecule comprises a sequencing primer or portion thereof, or complement thereof. In some embodiments, (b) comprises subjecting the DNA molecule to conditions sufficient to hybridize the first nucleic acid barcode molecule to the DNA molecule. In some embodiments, the method further comprises (i) extending the first nucleic acid barcode molecule hybridized to the DNA molecule and (ii) ligating the first nucleic acid barcode molecule to the DNA molecule to generate the first barcoded nucleic acid product.

[0039] In some embodiments, the method further comprises ligating the first nucleic acid barcode molecule to the DNA molecule.

[0040] In some embodiments, the first nucleic acid barcode molecule comprises a flow cell sequence, or complement thereof; a barcode sequence; a unique molecular identifier sequence; or a combination thereof.

[0041] In some embodiments, the DNA molecule comprises one or more gaps, and further comprising, prior to (c), subjecting the DNA molecule to an extension process to fill the one or more gaps.

[0042] In some embodiments, the second nucleic acid barcode molecule comprises a unique molecular identifier sequence, a barcode sequence, a sequencing primer or portion thereof, or a combination thereof.

[0043] In some embodiments, the method further comprises, prior to (b), within the partition, reverse transcribing the RNA molecule to provide a complementary DNA (cDNA) molecule. In some embodiments, prior to reverse transcribing the RNA molecule, the second nucleic acid barcode molecule is hybridized to a sequence of a precursor of the RNA molecule. In some embodiments, the precursor comprises a polyA sequence, and wherein the second nucleic acid barcode molecule comprises a polyT sequence configured to hybridize to the polyA sequence. In some embodiments, the method further comprises appending an additional sequence to the cDNA molecule, wherein the additional sequence is a poly(C) sequence. In some embodiments, the method further comprises contacting the cDNA molecule with a template-switching

oligonucleotide under conditions sufficient to hybridize the template-switching oligonucleotide to the cDNA molecule, wherein the template-switching oligonucleotide hybridizes to the additional sequence of the cDNA molecule. In some embodiments, the method further comprises extending the cDNA molecule to generate an extended cDNA molecule, which extended cDNA molecule comprises sequences complementary to sequences of the template-switching oligonucleotide. In some embodiments, the template-switching oligonucleotide comprises a sequencing primer or portion thereof, or complement thereof; a unique molecular identifier sequence, or complement thereof; or a combination thereof. In some embodiments, the template-switching oligonucleotide is the second nucleic acid barcode molecule, and wherein the extended cDNA molecule is the second barcoded nucleic acid product.

[0044] In some embodiments, the first nucleic acid barcode molecule and the second nucleic acid barcode molecule comprise a common nucleic acid sequence.

[0045] In some embodiments, the partition is a droplet among a plurality of droplets. In some embodiments, the partition is a droplet and wherein (c) comprises breaking or disrupting the droplet.

[0046] In some embodiments, (e) comprises performing one or more nucleic acid amplification reactions using the first barcoded nucleic acid product or the second barcoded nucleic acid product. In some embodiments, (e) comprises attaching one or more flow cell sequences to the first barcoded nucleic acid product or the second barcoded nucleic acid product, or a derivative thereof.

[0047] In some embodiments, the first nucleic acid barcode molecule and the second nucleic acid barcode molecule are coupled to a bead. In some embodiments, the bead is a gel bead. In some embodiments, the first nucleic acid barcode molecule is coupled to the bead via a first labile moiety and the second nucleic acid barcode molecule is coupled to the bead via a second labile moiety. In some embodiments, the first nucleic acid barcode molecule and the second nucleic acid barcode molecule are releasably coupled to the bead. In some embodiments, the first nucleic acid barcode molecule is releasable from the bead upon application of a first stimulus and the second nucleic acid barcode molecule is releasable from the bead upon application of a second stimulus.

[0048] In an aspect, the present disclosure provides a method for processing a nucleic acid sample, comprising: (a) providing a partition among a plurality of partitions, wherein the partition comprises a deoxyribonucleic acid (DNA) molecule derived from the nucleic acid sample; (b) transcribing the DNA molecule to generate a ribonucleic acid (RNA) molecule; (c)

reverse transcribing the RNA molecule to generate a complementary DNA (cDNA) molecule; and (d) recovering the cDNA molecule or a derivative thereof from the partition.

[0049] In some embodiments, the partition is a well among a plurality of wells.

[0050] In some embodiments, the partition is a droplet among a plurality of droplets.

[0051] In some embodiments, the method further comprises, prior to (a), processing an open chromatin structure of the nucleic acid sample with a transposase to yield the DNA molecule. In some embodiments, the transposase is a Tn5 transposase. In some embodiments, the DNA molecule comprises one or more gaps. In some embodiments, the one or more gaps are filled using a reverse transcriptase. In some embodiments, the processing further comprises attachment of a nucleic acid molecule to the nucleic acid sample, wherein the nucleic acid molecule comprises a sequence having a first sub-sequence and a second sub-sequence, wherein the first sub-sequence is complementary to the second sub-sequence. In some embodiments, the nucleic acid molecule is a hairpin nucleic acid molecule. In some embodiments, the nucleic acid molecule comprises a promoter sequence. In some embodiments, the promoter sequence is a T7 promoter sequence. In some embodiments, the nucleic acid molecule is attached to a 3' end of the nucleic acid sample.

[0052] In some embodiments, (b) comprises transcribing the DNA molecule using a T7 polymerase.

[0053] In some embodiments, (c) comprises appending a poly(C) sequence to the RNA molecule.

[0054] In some embodiments, the method further comprises subsequent to (c), attaching a nucleic acid barcode molecule to a 5' end of the cDNA molecule. In some embodiments, the nucleic acid barcode molecule comprises a unique molecular identifier sequence. In some embodiments, the nucleic acid barcode molecule comprises a barcode sequence. In some embodiments, the nucleic acid barcode molecule comprises a primer sequence. In some embodiments, the nucleic acid barcode molecule comprises a capture sequence. In some embodiments, the capture sequence comprises a poly(G) sequence.

[0055] In some embodiments, the nucleic acid barcode molecule is coupled to a bead. In some embodiments, the bead is a gel bead. In some embodiments, the nucleic acid barcode molecule is releasably coupled to the bead. In some embodiments, the nucleic acid barcode molecule is releasable from the bead upon application of a stimulus. In some embodiments, the stimulus is a chemical stimulus. In some embodiments, the chemical stimulus is a reducing agent.

[0056] In some embodiments, the cDNA molecule is a single-stranded cDNA molecule. In some embodiments, the method further comprises using the single-stranded cDNA molecule to generate a double-stranded cDNA molecule. In some embodiments, the double-stranded cDNA molecule is generated using a 5' blocked primer and a Klenow fragment. In some embodiments, the Klenow fragment does not have exonuclease activity.

[0057] In some embodiments, the cDNA molecule comprises a DNA sequence coupled to an RNA sequence, and wherein subsequent to (d), the cDNA molecule is processed with an RNase enzyme to digest the RNA sequence or remove the RNA sequence from the cDNA molecule.

[0058] In some embodiments, the partition is a droplet, and wherein (d) comprises breaking or disrupting the droplet.

[0059] In some embodiments, the method further comprises subsequent to (d), attaching one or more flow cell sequences to the cDNA molecule or derivative thereof. In some embodiments, the one or more flow cell sequences are attached via one or more nucleic acid amplification reactions.

[0060] In some embodiments, the method further comprises subsequent to (d), attaching a primer to a 3' end of the cDNA molecule or derivative thereof. In some embodiments, the cDNA molecule includes an A tail and wherein the primer attaches to the A tail.

[0061] In some embodiments, the nucleic acid sample is from a cell or cell nucleus. In some embodiments, the cell or the cell nucleus is in the partition. In some embodiments, the cell or the cell nucleus comprises the DNA molecule. In some embodiments, the cell or the cell nucleus further comprises a second ribonucleic acid (RNA) molecule. In some embodiments, the method further comprises lysing or permeabilizing the cell or the cell nucleus to provide access to the DNA molecule and/or the second RNA molecule therein.

[0062] In some embodiments, the method further comprises prior to (a), processing a messenger RNA (mRNA) molecule to provide the second RNA molecule. In some embodiments, the method further comprises, within the partition, reverse transcribing the second RNA molecule to provide an additional cDNA molecule. In some embodiments, prior to reverse transcribing the second RNA molecule, a primer molecule is hybridized to a sequence of a precursor of the second RNA molecule. In some embodiments, the precursor comprises a polyA sequence, and wherein the primer molecule comprises a polyT sequence configured to hybridize to the polyA sequence. In some embodiments, the primer molecule comprises a unique molecular identifier sequence. In some embodiments, the primer molecule comprises a barcode sequence. In some embodiments, the primer molecule comprises a sequencing primer or portion

thereof. In some embodiments, the primer molecule is coupled to an additional bead. In some embodiments, the additional bead is a gel bead. In some embodiments, the primer molecule is coupled to the additional bead via a labile moiety. In some embodiments, the primer molecule is releasably coupled to the additional bead. In some embodiments, the primer molecule is releasable from the additional bead upon application of stimulus. In some embodiments, the stimulus is a chemical stimulus. In some embodiments, the chemical stimulus is a reducing agent.

[0063] In some embodiments, the method further comprises appending an additional sequence to the additional cDNA molecule. In some embodiments, the additional sequence comprises a poly(C) sequence. In some embodiments, the method further comprises contacting the additional cDNA molecule with a template-switching oligonucleotide under conditions sufficient to hybridize the template-switching oligonucleotide to the additional cDNA molecule. In some embodiments, the template-switching oligonucleotide hybridizes to the additional sequence of the additional cDNA molecule. In some embodiments, the method further comprises extending the additional cDNA molecule to generate an extended cDNA molecule, which extended cDNA molecule comprises sequences complementary to sequences of the template-switching oligonucleotide. In some embodiments, the template-switching oligonucleotide comprises a sequencing primer, or complement thereof. In some embodiments, the template-switching oligonucleotide comprises a barcode sequence, or complement thereof. In some embodiments, the template-switching oligonucleotide comprises a unique molecular identifier sequence, or complement thereof. In some embodiments, the template-switching oligonucleotide is coupled to a further bead. In some embodiments, the further bead is a gel bead. In some embodiments, the template-switching oligonucleotide is coupled to the further bead via a labile moiety. In some embodiments, the template-switching oligonucleotide is releasably coupled to the further bead. In some embodiments, the template-switching oligonucleotide is releasable from the further bead upon application of stimulus. In some embodiments, the stimulus is a chemical stimulus. In some embodiments, the chemical stimulus is a reducing agent.

[0064] In some embodiments, the template-switching oligonucleotide is a further nucleic acid barcode molecule, and wherein the extended cDNA molecule is a barcoded nucleic acid product. In some embodiments, the further nucleic acid barcode molecule comprises a flow cell sequence, or complement thereof. In some embodiments, the method further comprises contacting the extended cDNA molecule with an additional nucleic acid barcode molecule under conditions sufficient to hybridize the additional nucleic acid barcode molecule to a sequence of

the extended cDNA molecule. In some embodiments, the method further comprises subjecting the extended cDNA molecule hybridized to the additional nucleic acid barcode molecule to an extension reaction to generate a barcoded nucleic acid product, which barcoded nucleic acid product comprises sequences complementary to sequences of the extended cDNA molecule. In some embodiments, the additional nucleic acid barcode molecule comprises a sequencing primer or portion thereof, or complement thereof. In some embodiments, the additional nucleic acid barcode molecule comprises a flow cell sequence, or complement thereof. In some embodiments, the additional nucleic acid barcode molecule comprises a barcode sequence, or complement thereof. In some embodiments, the additional nucleic acid barcode molecule is coupled to another bead. In some embodiments, the other bead is a gel bead. In some embodiments, the additional nucleic acid barcode molecule is coupled to the other bead via a labile moiety. In some embodiments, the additional nucleic acid barcode molecule is releasably coupled to the other bead. In some embodiments, the additional nucleic acid barcode molecule is releasable from the other bead upon application of stimulus. In some embodiments, the stimulus is a chemical stimulus. In some embodiments, the chemical stimulus is a reducing agent.

[0065] In some embodiments, the method further comprises recovering the barcoded nucleic acid product or a derivative thereof from the partition. In some embodiments, the method further comprises attaching one or more flow cell sequences to the barcoded nucleic acid product or derivative thereof. In some embodiments, the one or more flow cell sequences are attached via one or more nucleic acid amplification reactions.

[0066] In another aspect, the present disclosure provides a method for processing a nucleic acid sample, comprising: (a) providing a biological particle comprising a deoxyribonucleic acid (DNA) molecule and a ribonucleic acid (RNA) molecule within a partition among a plurality of partitions, wherein the partition comprises a first plurality of nucleic acid barcode molecules and a second plurality of nucleic acid barcode molecules, wherein the first plurality of nucleic acid barcode molecules and the second plurality of nucleic acid barcode molecules comprise a common barcode sequence; (b) within the partition, generating a first barcoded nucleic acid product corresponding to the DNA molecule and a second barcoded nucleic acid product corresponding to the RNA molecule, wherein the first barcoded nucleic acid product and the second barcoded nucleic acid product comprise the common barcode sequence, or a complement thereof; (c) recovering the first barcoded nucleic acid product and the second barcoded nucleic acid product from the partition; and (d) using (i) the first barcoded nucleic acid product to generate a first plurality of amplification products and (ii) the second barcoded nucleic acid

product to generate a second plurality of amplification products, wherein in (b), the generating does not comprise performing an exponential amplification reaction.

[0067] In some embodiments, the partition is a droplet among a plurality of droplets.

[0068] In some embodiments, the partition is a well among a plurality of wells.

[0069] In some embodiments, the biological particle is a cell, cell bead, or cell nucleus.

[0070] In some embodiments, the method further comprises lysing or permeabilizing the biological particle within the partition to provide access to the DNA molecule and the RNA molecule therein.

[0071] In some embodiments, the method further comprises, prior to (a), processing an open chromatin structure of the nucleic acid sample with a transposase to yield the DNA molecule. In some embodiments, the transposase is a Tn5 transposase. In some embodiments, the transposase is included in a transposase-nucleic acid complex. In some embodiments, the transposase-nucleic acid complex comprises (i) a first nucleic acid molecule comprising a first transposon end sequence and a first sequencing primer or portion thereof, or a complement thereof and (ii) a second nucleic acid molecule comprising a second transposon end sequence and a second sequencing primer or portion thereof, or a complement thereof. In some embodiments, the first transposon end sequence and the second transposon end sequence are the same, and wherein the first transposon end sequence and the second transposon end sequence are each hybridized to complementary sequences. In some embodiments, the first sequencing primer or portion thereof, or complement thereof, is different than the second sequencing primer or portion thereof, or complement thereof.

[0072] In some embodiments, the first plurality of nucleic acid barcode molecules each comprise a sequencing primer or portion thereof, or complement thereof. In some embodiments, (b) comprises subjecting the DNA molecule to conditions sufficient to hybridize a first nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules to the DNA molecule. In some embodiments, the method further comprises extending the first nucleic acid barcode molecule hybridized to the DNA molecule to generate the first barcoded nucleic acid product. In some embodiments, the method further comprises ligating the first nucleic acid barcode molecule to the DNA molecule.

[0073] In some embodiments, the first plurality of nucleic acid barcode molecules each comprise an overhang sequence. In some embodiments, (b) comprises subjecting the DNA molecule to conditions sufficient to hybridize to a splint sequence, wherein the splint sequence comprises a sequence complementary to the DNA molecule and a sequence complementary to the overhang sequence. In some embodiments, the splint sequence comprises a 3' blocking

group. In some embodiments, (b) comprises subjecting the DNA molecule hybridized to the splint sequence to conditions sufficient to hybridize the overhang sequence of a first nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules to the sequence complementary to the overhang sequence. In some embodiments, the method further comprises ligating the first nucleic acid barcode molecule to the DNA molecule. In some embodiments, the split sequence is hybridized to the overhang sequence outside of the partition.

[0074] In some embodiments, the first plurality of nucleic acid barcode molecules each comprise a flow cell sequence, or complement thereof.

[0075] In some embodiments, the DNA molecule comprises one or more gaps. In some embodiments, the method further comprises, prior to (c), subjecting the DNA molecule to an extension process to fill the one or more gaps. In some embodiments, filling of the one or more gaps is blocked by the presence of a transposase. In some embodiments, the method further comprises subsequent to (c), subjecting the DNA molecule to an extension process to fill the one or more gaps.

[0076] In some embodiments, the method further comprises, prior to (a), processing a messenger RNA (mRNA) molecule to provide the RNA molecule. In some embodiments, the method further comprises, prior to (b), within the partition, reverse transcribing the RNA molecule to provide an additional cDNA molecule. In some embodiments, prior to reverse transcribing the RNA molecule, a primer molecule is hybridized to a sequence of a precursor of the second RNA molecule. In some embodiments, the precursor comprises a polyA sequence, and wherein the primer molecule comprises a polyT sequence configured to hybridize to the polyA sequence. In some embodiments, the primer molecule comprises a unique molecular identifier sequence. In some embodiments, the primer molecule comprises a barcode sequence. In some embodiments, the primer molecule comprises a sequencing primer or portion thereof. In some embodiments, the primer molecule is a second nucleic acid barcode molecule of the plurality of second nucleic acid barcode molecules. In some embodiments, the method further comprises appending an additional sequence to the additional cDNA molecule. In some embodiments, the additional sequence is a poly(C) sequence. In some embodiments, the method further comprises contacting the additional cDNA molecule with a template-switching oligonucleotide under conditions sufficient to hybridize the template-switching oligonucleotide to the additional cDNA molecule. In some embodiments, the template-switching oligonucleotide hybridizes to the additional sequence of the additional cDNA molecule. In some embodiments, the method further comprises extending the additional cDNA molecule to generate an extended cDNA molecule, which extended cDNA molecule comprises sequences complementary to

sequences of the template-switching oligonucleotide. In some embodiments, the template-switching oligonucleotide comprises a sequencing primer or portion thereof, or complement thereof. In some embodiments, the template-switching oligonucleotide comprises a unique molecular identifier sequence, or complement thereof. In some embodiments, the method further comprises contacting the extended cDNA molecule with a second nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules under conditions sufficient to hybridize the second nucleic acid barcode molecule to a sequence of the extended cDNA molecule. In some embodiments, the method further comprises subjecting the extended cDNA molecule hybridized to the second nucleic acid barcode molecule to an extension reaction to generate the second barcoded nucleic acid product, which second barcoded nucleic acid product comprises sequences complementary to sequences of the extended cDNA molecule. In some embodiments, the second nucleic acid barcode molecule comprises a sequencing primer or portion thereof, or complement thereof. In some embodiments, the second nucleic acid barcode molecule comprises a flow cell sequence, or complement thereof.

[0077] In some embodiments, the first plurality of nucleic acid barcode molecules and the second plurality of nucleic acid barcode molecules are the same.

[0078] In some embodiments, the template-switching oligonucleotide is a second nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules, and wherein the extended cDNA molecule is the second barcoded nucleic acid product. In some embodiments, the second nucleic acid barcode molecule comprises a flow cell sequence, or complement thereof.

[0079] In some embodiments, the partition is a droplet and wherein (c) comprises breaking or disrupting the droplet.

[0080] In some embodiments, (d) comprises performing one or more nucleic acid amplification reactions using the first barcoded nucleic acid product or the second barcoded nucleic acid product. In some embodiments, the one or more nucleic acid amplification reactions comprise one or more polymerase chain reactions (PCR). In some embodiments, (d) comprises attaching one or more flow cell sequences to the first barcoded nucleic acid product or the second barcoded nucleic acid product, or a derivative thereof.

[0081] In some embodiments, the first plurality of nucleic acid barcode molecules are coupled to a first bead and the second plurality of nucleic acid barcode molecules are coupled to a second bead. In some embodiments, the first bead and the second bead are gel beads. In some embodiments, the first bead is linked to the second bead. In some embodiments, the first plurality of nucleic acid barcode molecules are coupled to the first bead via a first labile moiety

and the second plurality of nucleic acid barcode molecules are coupled to the second bead via a second labile moiety. In some embodiments, the first labile moiety and the second labile moiety are the same. In some embodiments, the first plurality of nucleic acid barcode molecules are releasably coupled to the first bead and the second plurality of nucleic acid barcode molecules are releasably coupled to the second bead. In some embodiments, the first plurality of nucleic acid barcode molecules are releasable from the first bead upon application of a first stimulus and the second plurality of nucleic acid barcode molecules are releasable from the second bead upon application of a second stimulus. In some embodiments, the first stimulus and the second stimulus are the same. In some embodiments, the first stimulus and the second stimulus are chemical stimuli.

[0082] In some embodiments, the first plurality of nucleic acid barcode molecules and the second plurality of nucleic acid barcode molecules are coupled to a bead. In some embodiments, the bead is a gel bead. In some embodiments, the first plurality of nucleic acid barcode molecules are coupled to the bead via a first labile moiety and the second plurality of nucleic acid barcode molecules are coupled to the bead via a second labile moiety. In some embodiments, the first labile moiety and the second labile moiety are the same. In some embodiments, the first plurality of nucleic acid barcode molecules and the second plurality of nucleic acid barcode molecules are releasably coupled to the bead. In some embodiments, the first plurality of nucleic acid barcode molecules are releasable from the bead upon application of a first stimulus and the second plurality of nucleic acid barcode molecules are releasable from the bead upon application of a second stimulus. In some embodiments, the first stimulus and the second stimulus are the same. In some embodiments, the first stimulus and the second stimulus are chemical stimuli.

[0083] In some embodiments, the method further comprises, subsequent to (d), sequencing the first plurality of amplification products and the second plurality of amplification products to identify sequences of the DNA molecule and the RNA molecule, respectively.

[0084] Another aspect of the present disclosure provides a non-transitory computer readable medium comprising machine executable code that, upon execution by one or more computer processors, implements any of the methods above or elsewhere herein.

[0085] Another aspect of the present disclosure provides a system comprising one or more computer processors and computer memory coupled thereto. The computer memory comprises machine executable code that, upon execution by the one or more computer processors, implements any of the methods above or elsewhere herein.

[0086] Additional aspects and advantages of the present disclosure will become readily apparent to those skilled in this art from the following detailed description, wherein only illustrative embodiments of the present disclosure are shown and described. As will be realized, the present disclosure is capable of other and different embodiments, and its several details are capable of modifications in various obvious respects, all without departing from the disclosure. Accordingly, the drawings and description are to be regarded as illustrative in nature, and not as restrictive.

INCORPORATION BY REFERENCE

[0087] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference. To the extent publications and patents or patent applications incorporated by reference contradict the disclosure contained in the specification, the specification is intended to supersede and/or take precedence over any such contradictory material.

BRIEF DESCRIPTION OF THE DRAWINGS

[0088] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings (also “Figure” and “FIG.” herein), of which:

[0089] **FIG. 1** shows an example of a microfluidic channel structure for partitioning individual biological particles.

[0090] **FIG. 2** shows an example of a microfluidic channel structure for delivering barcode carrying beads to droplets.

[0091] **FIG. 3** shows an example of a microfluidic channel structure for co-partitioning biological particles and reagents.

[0092] **FIG. 4** shows an example of a microfluidic channel structure for the controlled partitioning of beads into discrete droplets.

[0093] **FIG. 5** shows an example of a microfluidic channel structure for increased droplet generation throughput.

[0094] **FIG. 6** shows another example of a microfluidic channel structure for increased droplet generation throughput.

[0095] FIG. 7A shows a cross-section view of another example of a microfluidic channel structure with a geometric feature for controlled partitioning. FIG. 7B shows a perspective view of the channel structure of FIG. 7A.

[0096] FIG. 8 illustrates an example of a barcode carrying bead.

[0097] FIG. 9 illustrates a transposase-nucleic acid complex comprising a transposase, a first double-stranded oligonucleotide comprising a transposon end sequence and a first primer sequence and a second double-stranded oligonucleotide comprising a transposon end sequence and a second primer sequence.

[0098] FIG. 10 illustrates a transposase-nucleic acid complex comprising a transposase, a first double-stranded oligonucleotide comprising a transposon end sequence and first and second primer sequences and a second double-stranded oligonucleotide comprising a transposon end sequence and third and fourth primer sequences.

[0099] FIG. 11 illustrates a transposase-nucleic acid complex comprising a transposase, a first hairpin molecule, and a second hairpin molecule.

[00100] FIG. 12 illustrates a scheme for tandem ATAC ligation and RNA template switching.

[00101] FIG. 13 illustrates an additional scheme for tandem ATAC ligation and RNA template switching.

[00102] FIG. 14 illustrates an exemplary scheme for tandem ATAC ligation and RNA template switching.

[00103] FIG. 15 illustrates an additional scheme for tandem ATAC ligation and RNA template switching.

[00104] FIG. 16 illustrates an additional scheme for tandem ATAC ligation and RNA template switching.

[00105] FIG. 17 illustrates an additional scheme for tandem ATAC ligation and RNA template switching.

[00106] FIG. 18 illustrates an additional scheme for tandem ATAC ligation and RNA template switching.

[00107] FIG. 19 illustrates an additional scheme for tandem ATAC ligation and RNA template switching.

[00108] FIG. 20 illustrates a scheme for T7 mediated linear amplification.

[00109] FIG. 21 shows a modified workflow T7 mediated linear amplification.

[00110] FIG. 22 illustrates a scheme for tandem ATAC and RNA processing.

[00111] FIG. 23 illustrates a scheme for tandem ATAC and RNA processing.

[00112] **FIGs. 24A and 24B** show beads for use according to the methods of the present disclosure.

[00113] **FIG. 25** shows a computer system that is programmed or otherwise configured to implement methods provided herein.

[00114] The instant application contained at least one drawing executed in color. Copies of this patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

DETAILED DESCRIPTION

[00115] While various embodiments of the invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions may occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed.

[00116] The terms “a,” “an,” and “the,” as used herein, generally refers to singular and plural references unless the context clearly dictates otherwise.

[00117] Whenever the term “at least,” “greater than,” or “greater than or equal to” precedes the first numerical value in a series of two or more numerical values, the term “at least,” “greater than” or “greater than or equal to” applies to each of the numerical values in that series of numerical values. For example, greater than or equal to 1, 2, or 3 is equivalent to greater than or equal to 1, greater than or equal to 2, or greater than or equal to 3.

[00118] Whenever the term “no more than,” “less than,” or “less than or equal to” precedes the first numerical value in a series of two or more numerical values, the term “no more than,” “less than,” or “less than or equal to” applies to each of the numerical values in that series of numerical values. For example, less than or equal to 3, 2, or 1 is equivalent to less than or equal to 3, less than or equal to 2, or less than or equal to 1.

[00119] Where values are described as ranges, it will be understood that such disclosure includes the disclosure of all possible sub-ranges within such ranges, as well as specific numerical values that fall within such ranges irrespective of whether a specific numerical value or specific sub-range is expressly stated.

[00120] The term “barcode,” as used herein, generally refers to a label, or identifier, that conveys or is capable of conveying information about an analyte. A barcode can be part of an analyte. A barcode can be independent of an analyte. A barcode can be a tag attached to an analyte (e.g., nucleic acid molecule) or a combination of the tag in addition to an endogenous characteristic of the analyte (e.g., size of the analyte or end sequence(s)). A barcode may be

unique. Barcodes can have a variety of different formats. For example, barcodes can include: polynucleotide barcodes; random nucleic acid and/or amino acid sequences; and synthetic nucleic acid and/or amino acid sequences. A barcode can be attached to an analyte in a reversible or irreversible manner. A barcode can be added to, for example, a fragment of a deoxyribonucleic acid (DNA) or ribonucleic acid (RNA) sample before, during, and/or after sequencing of the sample. Barcodes can allow for identification and/or quantification of individual sequencing-reads.

[00121] The term “real time,” as used herein, can refer to a response time of less than about 1 second, a tenth of a second, a hundredth of a second, a millisecond, or less. The response time may be greater than 1 second. In some instances, real time can refer to simultaneous or substantially simultaneous processing, detection or identification.

[00122] The term “subject,” as used herein, generally refers to an animal, such as a mammal (e.g., human) or avian (e.g., bird), or other organism, such as a plant. For example, the subject can be a vertebrate, a mammal, a rodent (e.g., a mouse), a primate, a simian or a human. Animals may include, but are not limited to, farm animals, sport animals, and pets. A subject can be a healthy or asymptomatic individual, an individual that has or is suspected of having a disease (e.g., cancer) or a pre-disposition to the disease, and/or an individual that is in need of therapy or suspected of needing therapy. A subject can be a patient. A subject can be a microorganism or microbe (e.g., bacteria, fungi, archaea, viruses).

[00123] The term “genome,” as used herein, generally refers to genomic information from a subject, which may be, for example, at least a portion or an entirety of a subject’s hereditary information. A genome can be encoded either in DNA or in RNA. A genome can comprise coding regions (e.g., that code for proteins) as well as non-coding regions. A genome can include the sequence of all chromosomes together in an organism. For example, the human genome ordinarily has a total of 46 chromosomes. The sequence of all of these together may constitute a human genome.

[00124] The terms “adaptor(s)”, “adapter(s)” and “tag(s)” may be used synonymously. An adaptor or tag can be coupled to a polynucleotide sequence to be “tagged” by any approach, including ligation, hybridization, or other approaches.

[00125] The term “sequencing,” as used herein, generally refers to methods and technologies for determining the sequence of nucleotide bases in one or more polynucleotides. The polynucleotides can be, for example, nucleic acid molecules such as deoxyribonucleic acid (DNA) or ribonucleic acid (RNA), including variants or derivatives thereof (e.g., single stranded DNA). Sequencing can be performed by various systems currently available, such as, without

limitation, a sequencing system by Illumina®, Pacific Biosciences (PacBio®), Oxford Nanopore®, or Life Technologies (Ion Torrent®). Alternatively or in addition, sequencing may be performed using nucleic acid amplification, polymerase chain reaction (PCR) (e.g., digital PCR, quantitative PCR, or real time PCR), or isothermal amplification. Such systems may provide a plurality of raw genetic data corresponding to the genetic information of a subject (e.g., human), as generated by the systems from a sample provided by the subject. In some examples, such systems provide sequencing reads (also “reads” herein). A read may include a string of nucleic acid bases corresponding to a sequence of a nucleic acid molecule that has been sequenced. In some situations, systems and methods provided herein may be used with proteomic information.

[00126] The term “bead,” as used herein, generally refers to a particle. The bead may be a solid or semi-solid particle. The bead may be a gel bead. The gel bead may include a polymer matrix (e.g., matrix formed by polymerization or cross-linking). The polymer matrix may include one or more polymers (e.g., polymers having different functional groups or repeat units). Polymers in the polymer matrix may be randomly arranged, such as in random copolymers, and/or have ordered structures, such as in block copolymers. Cross-linking can be via covalent, ionic, or inductive, interactions, or physical entanglement. The bead may be a macromolecule. The bead may be formed of nucleic acid molecules bound together. The bead may be formed via covalent or non-covalent assembly of molecules (e.g., macromolecules), such as monomers or polymers. Such polymers or monomers may be natural or synthetic. Such polymers or monomers may be or include, for example, nucleic acid molecules (e.g., DNA or RNA). The bead may be formed of a polymeric material. The bead may be magnetic or non-magnetic. The bead may be rigid. The bead may be flexible and/or compressible. The bead may be disruptable or dissolvable. The bead may be a solid particle (e.g., a metal-based particle including but not limited to iron oxide, gold or silver) covered with a coating comprising one or more polymers. Such coating may be disruptable or dissolvable.

[00127] The term “sample,” as used herein, generally refers to a biological sample of a subject. The biological sample may comprise any number of macromolecules, for example, cellular macromolecules. The sample may be a cell sample. The sample may be a cell line or cell culture sample. The sample can include one or more cells. The sample can include one or more microbes. The biological sample may be a nucleic acid sample or protein sample. The biological sample may also be a carbohydrate sample or a lipid sample. The biological sample may be derived from another sample. The sample may be a tissue sample, such as a biopsy, core biopsy, needle aspirate, or fine needle aspirate. The sample may be a fluid sample, such as a

blood sample, urine sample, or saliva sample. The sample may be a skin sample. The sample may be a cheek swab. The sample may be a plasma or serum sample. The sample may be a cell-free or cell free sample. A cell-free sample may include extracellular polynucleotides.

Extracellular polynucleotides may be isolated from a bodily sample that may be selected from the group consisting of blood, plasma, serum, urine, saliva, mucosal excretions, sputum, stool and tears.

[00128] The term “biological particle,” as used herein, generally refers to a discrete biological system derived from a biological sample. The biological particle may be a macromolecule. The biological particle may be a small molecule. The biological particle may be a virus. The biological particle may be a cell or derivative of a cell. The biological particle may be an organelle. The biological particle may be a rare cell from a population of cells. The biological particle may be any type of cell, including without limitation prokaryotic cells, eukaryotic cells, bacterial, fungal, plant, mammalian, or other animal cell type, mycoplasmas, normal tissue cells, tumor cells, or any other cell type, whether derived from single cell or multicellular organisms. The biological particle may be a constituent of a cell. The biological particle may be or may include DNA, RNA, organelles, proteins, or any combination thereof. The biological particle may be or may include a matrix (e.g., a gel or polymer matrix) comprising a cell or one or more constituents from a cell (e.g., cell bead), such as DNA, RNA, organelles, proteins, or any combination thereof, from the cell. The biological particle may be obtained from a tissue of a subject. The biological particle may be a hardened cell. Such hardened cell may or may not include a cell wall or cell membrane. The biological particle may include one or more constituents of a cell, but may not include other constituents of the cell. An example of such constituents is a nucleus or an organelle. A cell may be a live cell. The live cell may be capable of being cultured, for example, being cultured when enclosed in a gel or polymer matrix, or cultured when comprising a gel or polymer matrix. For a description of exemplary cell beads, polymer or cross-linked matrices, and cell bead generation methods, see, e.g., U.S. Pat. Pub. 20180216162 and U.S. Pat. Pub. 20190100632, each of which is herein incorporated by reference in their entireties.

[00129] The term “macromolecular constituent,” as used herein, generally refers to a macromolecule contained within or from a biological particle. The macromolecular constituent may comprise a nucleic acid. In some cases, the biological particle may be a macromolecule. The macromolecular constituent may comprise DNA. The macromolecular constituent may comprise RNA. The RNA may be coding or non-coding. The RNA may be messenger RNA (mRNA), ribosomal RNA (rRNA) or transfer RNA (tRNA), for example. The RNA may be a

transcript. The RNA may be small RNA that are less than 200 nucleic acid bases in length, or large RNA that are greater than 200 nucleic acid bases in length. Small RNAs may include 5.8S ribosomal RNA (rRNA), 5S rRNA, transfer RNA (tRNA), microRNA (miRNA), small interfering RNA (siRNA), small nucleolar RNA (snoRNAs), Piwi-interacting RNA (piRNA), tRNA-derived small RNA (tsRNA) and small rDNA-derived RNA (srRNA). The RNA may be double-stranded RNA or single-stranded RNA. The RNA may be circular RNA. The macromolecular constituent may comprise a protein. The macromolecular constituent may comprise a peptide. The macromolecular constituent may comprise a polypeptide.

[00130] The term “molecular tag,” as used herein, generally refers to a molecule capable of binding to a macromolecular constituent. The molecular tag may bind to the macromolecular constituent with high affinity. The molecular tag may bind to the macromolecular constituent with high specificity. The molecular tag may comprise a nucleotide sequence. The molecular tag may comprise a nucleic acid sequence. The nucleic acid sequence may be at least a portion or an entirety of the molecular tag. The molecular tag may be a nucleic acid molecule or may be part of a nucleic acid molecule. The molecular tag may be an oligonucleotide or a polypeptide. The molecular tag may comprise a DNA aptamer. The molecular tag may be or comprise a primer. The molecular tag may be, or comprise, a protein. The molecular tag may comprise a polypeptide. The molecular tag may be a barcode.

[00131] The term “partition,” as used herein, generally, refers to a space or volume that may be suitable to contain one or more species or conduct one or more reactions. A partition may be a physical compartment, such as a droplet or well. The partition may isolate space or volume from another space or volume. The droplet may be a first phase (e.g., aqueous phase) in a second phase (e.g., oil) immiscible with the first phase. The droplet may be a first phase in a second phase that does not phase separate from the first phase, such as, for example, a capsule or liposome in an aqueous phase. A partition may comprise one or more other (inner) partitions. In some cases, a partition may be a virtual compartment that can be defined and identified by an index (e.g., indexed libraries) across multiple and/or remote physical compartments. For example, a physical compartment may comprise a plurality of virtual compartments.

[00132] The present disclosure provides methods, systems, and kits for processing multiple types of nucleic acid molecules. The methods, systems, and kits provided herein may facilitate sample preparation for sequencing of nucleic acid molecules included in cells, cell beads, or cell nuclei of interest. For example, the present disclosure provides methods for processing both deoxyribonucleic acid (DNA) and ribonucleic acid (RNA) molecules included within a cell, cell bead, or cell nucleus. The methods may comprise performing Assay for Transposase Accessible

Chromatin with high-throughput sequencing (ATAC-seq) and RNA sequencing (RNA-seq) assays in tandem. Partitioning and barcoding schemes may be utilized to facilitate identification of resultant sequencing reads with the cell, cell bead, or cell nucleus from which they are derived.

[00133] The present disclosure also provides methods, systems, and kits for processing biological samples comprising nucleic acid molecules. The method may comprise providing one or more nucleic acid molecules (e.g., deoxyribonucleic acid (DNA) molecules and/or ribonucleic acid (RNA) molecules) from a nucleic acid sample (e.g., a sample comprising a cell, cell bead, or cell nucleus) in a partition of a plurality of partitions (e.g., a plurality of droplets or wells). The one or more nucleic acid molecules may be one or more DNA molecules. The one or more DNA molecules may be transcribed to generate one or more RNA molecules, where the one or more RNA molecules may be reverse transcribed to generate one or more complementary DNA (cDNA) molecules. The one or more cDNA molecules, or derivatives thereof, may then be recovered from the partition of the plurality of partitions (e.g., by pooling the contents of the plurality of partitions). The one or more cDNA molecules, or derivatives thereof, may comprise one or more nucleic acid barcode sequences, or complements thereof, where the one or more nucleic acid barcode sequences, or complements thereof, may be incorporated into nucleic acid molecules during any processing step (e.g., during transcription of a DNA molecule, reverse transcription of an RNA molecule, etc.). The one or more nucleic acid barcode sequences, or complements thereof, may be used to identify sequencing reads (e.g., sequencing reads obtained using a nucleic acid sequencing assay) corresponding to the one or more cDNA molecules from a nucleic acid molecule of the one or more nucleic acid molecules from the nucleic acid sample.

Tandem DNA and RNA Barcoding

[00134] In an aspect, the present disclosure provides a method for processing nucleic acid molecules from a cell, cell bead, or cell nucleus. The method may comprise contacting a cell, cell bead, or cell nucleus with a transposase-nucleic acid complex comprising a transposase molecule and one or more transposon end oligonucleotide molecules. The cell, cell bead or cell nucleus may be contacted with a transposase-nucleic acid complex in bulk solution, such that the cell, cell bead or cell nucleus undergoes “tagmentation” via a tagmentation reaction. Contacting the cell, cell bead, or cell nucleus with the transposase-nucleic acid complex may generate one or more template nucleic acid fragments (e.g., “tagmented fragments”). The one or more template nucleic acid fragments may correspond to one or more target nucleic acid molecules (e.g., deoxyribonucleic acid (DNA) molecules) within the cell, cell bead, or cell nucleus. In parallel,

the cell, cell bead, or cell nucleus may be contacted with a primer molecule (e.g., a primer molecule comprising a poly-T sequence) configured to interact with one or more additional target nucleic acid molecules (e.g., ribonucleic acid (RNA) molecules, such as messenger RNA (mRNA) molecules). The cell, cell bead, or cell nucleus may be contacted with a primer molecule in bulk solution. Alternatively or in addition to, the cell, cell bead, or cell nucleus may be contacted with a primer molecule within a partition. Interaction between these moieties may yield one or more additional template nucleic acid fragments (e.g., RNA fragments). For example, the primer molecule may have at least partial sequence complementarity to the one or more additional target nucleic acid molecules (e.g., mRNA molecules). The primer molecule may hybridize to a sequence of an additional target nucleic acid molecule of the one or more additional target nucleic acid molecules. The cell, cell bead, or cell nucleus may be partitioned (e.g., co-partitioned with one or more reagents) into a partition (e.g., of a plurality of partitions). The partition may be, for example, a droplet or a well. The partition may comprise one or more reagents, including, for example, one or more particles (e.g., beads) comprising one or more nucleic acid barcode molecules. The cell, cell bead, or cell nucleus may be lysed, permeabilized, fixed, cross-linked or otherwise manipulated to provide access to the one or more template nucleic acid fragments and the one or more additional template nucleic acid fragments therein. The one or more template nucleic acid fragments and the one or more additional template nucleic acid fragments therein may undergo one or more processing steps within the partition. For example, the one or more template nucleic acid fragments and/or the one or more additional template nucleic acid fragments may undergo a barcoding process, a ligation process, a reverse transcription process, a template switching process, a linear amplification process, and/or a gap filling process. The resultant one or more processed template nucleic acid fragments (e.g., tagged fragments) and/or the one or more processed additional template nucleic acid fragments (e.g., RNA fragments) may each include a barcode sequence (e.g., nucleic acid barcode sequence, as described herein). The one or more processed template nucleic acid fragments and/or the one or more processed additional template nucleic acid fragments may be released from the partition (e.g., pooled with contents of other partitions of a plurality of partitions) and may undergo one or more additional processing steps in bulk. For example, the one or more processed template nucleic acid fragments and/or the one or more processed additional template nucleic acid fragments may undergo a gap filling process, a dA tailing process, a terminal-transferase process, a phosphorylation process, a ligation process, a nucleic acid amplification process, or a combination thereof. For example, the one or more processed template nucleic acid fragments and/or the one or more processed additional template nucleic

acid fragments may be subjected to conditions sufficient to undergo one or more polymerase chain reactions (PCR, such as sequence independent PCR) to generate amplification products corresponding to the one or more processed template nucleic acid fragments (e.g., tagmented fragments) and/or the one or more processed additional template nucleic acid fragments (e.g., RNA fragments). Sequences of such amplification products can be detected using, for example, a nucleic acid sequencing assay and used to identify sequences of the one or more target nucleic acid molecules (e.g., DNA molecules) and the one or more additional target nucleic acid molecules (e.g., RNA molecules) of the cell, cell bead, or cell nucleus from which they derive.

[00135] A biological sample (e.g., a nucleic acid sample) may comprise one or more cells, cell beads, and/or cell nuclei. A biological sample may also comprise tissue, which tissue may comprise one or more cells, cell beads, and/or cell nuclei. In some cases, a biological sample may comprise a plurality of cells comprising a plurality of cell nuclei. In some cases, a biological sample may comprise a plurality of cell nuclei, which plurality of cell nuclei are not included within cells (e.g., other components of the cell have degraded, dissociated, dissolved, or otherwise been removed). A biological sample may comprise a plurality of cell-free nucleic acid molecules (e.g., nucleic acid molecules that are not included within cells). For example, a biological sample may comprise a plurality of cell-free fetal DNA (cffDNA) or circulating tumor DNA (ctDNA) or other cell-free nucleic acid molecules (e.g., deriving from degraded cells). Such a biological sample may be processed to separate such cell-free nucleic acid molecules from cells, cell beads, and/or cell nuclei, which cells, cell beads, and/or cell nuclei may be subjected to further processing (e.g., as described herein).

[00136] Nucleic acid molecules included within a biological sample may include, for example, DNA molecules and RNA molecules. For example, a biological sample may comprise genomic DNA comprising chromatin (e.g., within a cell, cell bead, or cell nucleus). A biological sample may comprise a plurality of RNA molecules, such as a plurality of pre-mRNA or mRNA molecules. mRNA molecules and other RNA molecules may comprise a polyA sequence. At least a subset of a plurality of RNA molecules included in a cell or cell bead may be present in a cell nucleus.

[00137] A nucleic acid molecule may undergo one or more processing steps within a cell, cell bead, or cell nucleus. For example, chromatin within a cell, cell bead, or cell nucleus may be contacted with a transposase. A transposase may be included within a transposase-nucleic acid complex, which transposase-nucleic acid complex may comprise a transposase molecule and one or more transposon end oligonucleotide molecules. A transposase may be a Tn transposase, such as a Tn3, Tn5, Tn7, Tn10, Tn552, Tn903 transposase. Alternatively, a transposase may be a

MuA transposase, a Vibhar transposase (e.g. from *Vibrio harveyi*), Ac-Ds, Ascot-1, Bs1, Cin4, Copia, En/Spm, F element, hobo, Hsmar1, Hsmar2, IN (HIV), IS1, IS2, IS3, IS4, IS5, IS6, IS10, IS21, IS30, IS50, IS51, IS150, IS256, IS407, IS427, IS630, IS903, IS911, IS982, IS1031, ISL2, L1, Mariner, P element, Tam3, Tc1, Tc3, Te1, THE-1, Tn/O, TnA, Tn3, Tn5, Tn7, Tn10, Tn552, Tn903, Tol1, Tol2, TnIO, Tyl, any prokaryotic transposase, or any transposase related to and/or derived from those listed above. For example, a transposase may be a Tn5 transposase or a mutated, hyperactive Tn5 transposase. A transposase related to and/or derived from a parent transposase may comprise a peptide fragment with at least about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% amino acid sequence homology to a corresponding peptide fragment of the parent transposase. The peptide fragment may be at least about 10, about 15, about 20, about 25, about 30, about 35, about 40, about 45, about 50, about 60, about 70, about 80, about 90, about 100, about 150, about 200, about 250, about 300, about 400, or about 500 amino acids in length. For example, a transposase derived from Tn5 may comprise a peptide fragment that is 50 amino acids in length and about 80% homologous to a corresponding fragment in a parent Tn5 transposase. Action of a transposase (e.g., insertion) may be facilitated and/or triggered by addition of one or more cations, such as one or more divalent cations (e.g., Ca^{2+} , Mg^{2+} , or Mn^{2+}).

[00138] A transposase-nucleic acid complex may comprise one or more nucleic acid molecules. For example, a transposase-nucleic acid complex may comprise one or more transposon end oligonucleotide molecules. A transposon end oligonucleotide molecule may comprise one or more adapter sequences (e.g., comprising one or more primer sequences) and/or one or more transposon end sequences. A transposon end sequence may be, for example, a Tn5 or modified Tn5 transposon end sequence or a Mu transposon end sequence. A transposon end sequence may have a sequence of, for example,

AGATGTGTATAAGAGACA (SEQ ID NO: 1).

A primer sequence of a transposon end oligonucleotide molecule may be a sequencing primer, such as an R1 or R2 sequencing primer, or a portion thereof. A sequencing primer may be, for example, a TrueSeq or Nextera sequencing primer. An R1 sequencing primer region may have a sequence of

TCTACACTCTTTCCCTACACGACGCTCTTCCGATCT (SEQ ID NO: 2),

or some portion thereof. An R1 sequencing primer region may have a sequence of

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG (SEQ ID NO: 3),

or some portion thereof. A transposon end oligonucleotide molecule may comprise a partial R1 sequence. A partial R1 sequence may be

ACTACACGACGCTCTTCCGATCT (SEQ ID NO: 4).

A transposon end oligonucleotide molecule may comprise an R2 sequencing priming region. An R2 sequencing primer region may have a sequence of

GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT (SEQ ID NO: 5),

or some portion thereof. An R2 sequencing primer region may have a sequence of

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG (SEQ ID NO: 6),

or some portion thereof. A transposon end oligonucleotide molecule may comprise a T7 promoter sequence. A T7 promoter sequence may be

TAATACGACTCACTATAG (SEQ ID NO: 7).

A transposon end oligonucleotide molecule may comprise a region at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to any one of SEQ ID NO: 1-7. A transposon end oligonucleotide molecule may comprise a P5 sequence and/or a P7 sequence. A transposon end oligonucleotide molecule may comprise a sample index sequence, such as a barcode sequence or unique molecular identifier sequence. One or more transposon end oligonucleotide molecules of a transposase-nucleic acid complex may be attached to a solid support (e.g., a solid or semi-solid particle such as a bead (e.g., gel bead)). A transposon end oligonucleotide molecule may be releasably coupled to a solid support (e.g., a bead). Examples of transposon end oligonucleotide molecules may be found in, for example, PCT Patent Publications Nos. WO2018/218226, WO2014/189957, US. Pat. Pub. 20180340171, and U.S. Pat. 10,059,989; each of which are herein incorporated by reference in their entireties.

[00139] FIG. 9 includes an example of a transposase-nucleic acid complex for use in the methods provided herein. Transposase-nucleic acid complex **900** (e.g., comprising a transposase dimer) comprises partially double-stranded oligonucleotide **901** and partially double-stranded oligonucleotide **905**. Partially double-stranded oligonucleotide **901** comprises transposon end sequence **903**, first primer sequence **902**, and a sequence **904** that is complementary to transposon end sequence **903**. Partially double-stranded oligonucleotide **905** comprises transposon end sequence **906**, first primer sequence **907**, and a sequence **908** that is complementary to transposon end sequence **906**. Primer sequences **902** and **907** may be the same or different. In some cases, primer sequence **902** may be designated “R1” and primer sequence **907** may be designated “R2”. Transposon end sequences **903** and **906** may be the same or different. Transposon end sequences **903** and **906** may alternately be referred to as “mosaic

end” or “ME” sequences, while their complementary sequences **904** and **908** may be referred to as “mosaic end reverse complement” or “MErc” sequences.

[00140] **FIG. 10** includes another example of a transposase-nucleic acid complex for use in the methods provided herein. Transposase-nucleic acid complex **1000** (e.g., comprising a transposase dimer) comprises forked adapters **1001** and **1006**, which forked adapters are partially double-stranded oligonucleotides. Partially double-stranded oligonucleotide **1001** comprises transposon end sequence **1003**, first primer sequence **1002**, second primer sequence **1005**, and a sequence **1004** that is complementary to transposon end sequence **1003**. Partially double-stranded oligonucleotide **1006** comprises transposon end sequence **1007**, first primer sequence **1008**, second primer sequence **1010**, and a sequence **1009** that is complementary to transposon end sequence **1007**. Primer sequences **1002**, **1005**, **1008**, and **1010** may be the same or different. In some cases, primer sequences **1002** and **1008** may be designated “R1” and primer sequences **1005** and **1010** may be designated “R2”. Alternatively, primer sequences **1002** and **1010** may be designated “R1” and primer sequences **1005** and **1008** may be designated “R2”. Alternatively, primer sequences **1002** and **1008** may be designated “R2” and primer sequences **1005** and **1010** may be designated “R1”. Alternatively, primer sequences **1002** and **1010** may be designated “R2” and primer sequences **1005** and **1008** may be designated “R1”. Transposon end sequences **1003** and **1007** may be the same or different. These sequences may alternately be referred to as “mosaic end” or “ME” sequences, while their complementary sequences **1004** and **1009** may be referred to as “mosaic end reverse complement” or “MErc” sequences.

[00141] **FIG. 11** shows transposase-nucleic acid complex **1100** (e.g., comprising a transposase dimer) comprising hairpin molecules **1101** and **1106**. Hairpin molecule **1101** comprises transposon end sequence **1103**, first hairpin sequence **1102**, second hairpin sequence **1105**, and a sequence **1104** that is complementary to transposon end sequence **1103**. Hairpin molecule **1106** comprises transposon end sequence **1107**, third hairpin sequence **1108**, fourth hairpin sequence **1110**, and a sequence **1109** that is complementary to transposon end sequence **1107**. Hairpin sequences **1102**, **1105**, **1108**, and **1110** may be the same or different. For example, hairpin sequence **1105** may be the same or different as hairpin sequence **1110**, and/or hairpin sequence **1102** may be the same or different as hairpin sequence **1108**. Hairpin sequences **1102** and **1108** may be spacer sequences or adapter sequences. Hairpin sequences **1105** and **1110** may be a promoter sequence such as T7 recognition or promoter sequences and/or UMI sequences. Transposon end sequences **1103** and **1107** may be the same or different. Transposon end sequences **1103** and **1107** may alternately be referred to as “mosaic end” or “ME” sequences, while their complementary sequences **1104** and **1109** may be referred to as “mosaic end reverse

complement” or “MErc” sequences. In some cases, sequence **1104** is a transposon end sequence and **1103** is a sequence complementary to sequence **1104**. In some cases, sequence **1109** is a transposon end sequence and **1107** is a sequence complementary to sequence **1109**.

[00142] Contacting a cell, cell bead, or cell nucleus comprising one or more target nucleic acid molecules (e.g., DNA molecules) with a transposase-nucleic acid complex may generate one or more template nucleic acid fragments (e.g., “tagmented fragments”). The one or more template nucleic acid fragments may each comprise a sequence of the one or more target nucleic acid molecules (e.g., a target sequence). The transposase-nucleic acid complex may be configured to target a specific region of the one or more target nucleic acid molecules to provide one or more template nucleic acid fragments comprising specific target sequences. The one or more template nucleic acid fragments may comprise target sequences corresponding to accessible chromatin. Generation of tagmented fragments may take place within a bulk solution. In other cases, generation of tagmented fragments may take place within a partition (e.g., a droplet or well). A template nucleic acid fragment (e.g., tagmented fragment) may comprise one or more gaps (e.g., between a transposon end sequence or complement thereof and a target sequence on one or both strands of a double-stranded fragment). Gaps may be filled via a gap filling process using, e.g., a polymerase (e.g., DNA polymerase), ligase, or reverse transcriptase. In some cases, a mixture of enzymes may be used to repair a partially double-stranded nucleic acid molecule and fill one or more gaps. Gap filling may not include strand displacement. Gaps may be filled within or outside of a partition.

[00143] Alternatively or in addition to, one or more additional nucleic acid molecules may be contacted with one or more capture nucleic acid molecules within a cell, cell bead, or cell nucleus to provide one or more additional template nucleic acid fragments. For example, an RNA molecule (e.g., an mRNA) molecule may be contacted with a primer molecule within a cell, cell bead, or cell nucleus. A primer molecule may comprise a primer sequence, which primer sequence may be a targeted primer sequence or a non-specific primer sequence (e.g., random N-mer). A targeted primer sequence may comprise, for example, a polyT sequence, which polyT sequence may interact with a polyA sequence of an RNA molecule. A primer nucleic acid molecule may also comprise one or more additional sequences, such as one or more sample index sequences, spacer or linker sequences, or one or more additional primer sequences. Generation of additional template nucleic acid fragments (e.g., RNA fragments) may take place within a bulk solution. In other cases, generation of additional template nucleic acid fragments may take place within a partition (e.g., a droplet or well).

[00144] Processing of nucleic acid molecules within a cell, cell bead, or cell nucleus (e.g., generation of template nucleic acid fragments using a transposase-nucleic acid complex and/or generation of additional template nucleic acid fragments using a capture nucleic acid molecule) may occur in a bulk solution comprising a plurality of cells, cell beads, and/or cell nuclei. In some cases, template nucleic acid fragments (e.g., tagmented fragments) may be generated in bulk solution and additional template nucleic acid fragments (e.g., RNA fragments) may be generated in a partition.

[00145] A plurality of cells, cell beads, and/or cell nuclei (e.g., a plurality of cells, cell beads, and/or cell nuclei that have undergone processing such as a tagmentation process) may be partitioned amongst a plurality of partitions. Partitions may be, for example, droplets or wells. Droplets (e.g., aqueous droplets) may be generated according to the methods provided herein. Partitioning may be performed according to the method provided herein. For example, partitioning a biological particle (e.g., cell, cell bead, or cell nucleus) and one or more reagents may comprise flowing a first phase comprising an aqueous fluid, the biological particle, and the one or more reagents and a second phase comprising a fluid that is immiscible with the aqueous fluid toward a junction. Upon interaction of the first and second phases, a discrete droplet of the first phase comprising the biological particle and the one or more reagents may be formed. The plurality of cells, cell beads, and/or cell nuclei may be partitioned amongst a plurality of partitions such that at least a subset of the plurality of partitions may comprise at most one cell, cell bead, or cell nucleus. Cells, cell beads, and/or cell nuclei may be co-partitioned with one or more reagents such that a partition of at least a subset of the plurality of partitions comprises a single cell, cell bead, or cell nucleus and one or more reagents. The one or more reagents may include, for example, enzymes (e.g., polymerases, reverse transcriptases, ligases, etc.), nucleic acid barcode molecules (e.g., nucleic acid barcode molecules comprising one or more barcode sequences, such as nucleic acid barcode molecules coupled to one or more beads), template switching oligonucleotides, deoxynucleotide triphosphates, buffers, lysis agents, primers, barcodes, detergents, reducing agents, chelating agents, oxidizing agents, nanoparticles, beads, antibodies, or any other useful reagents. Enzymes may include, for example, temperature-sensitive enzymes, pH-sensitive enzymes, light-sensitive enzymes, reverse transcriptases, proteases, ligases, polymerases, kinases, restriction enzymes, nucleases, protease inhibitors, exonucleases, and nuclease inhibitors.

[00146] A reagent of the one or more reagents may be useful for lysing or permeabilizing a cell, cell bead, or cell nucleus, or otherwise providing access to nucleic acid molecules and/or template nucleic acid fragments therein. A cell may be lysed using a lysis agent such as a

bioactive agent. A bioactive agent useful for lysing a cell may be, for example, an enzyme (e.g., as described herein). An enzyme used to lyse a cell may or may not be capable of carrying out additional actions such as degrading one or more RNA molecules. Alternatively, an ionic, zwitterionic, or non-ionic surfactant may be used to lyse a cell. Examples of surfactants include, but are not limited to, TritonX-100, Tween 20, sarcosyl, or sodium dodecyl sulfate. Cell lysis may also be achieved using a cellular disruption method such as an electroporation or a thermal, acoustic, or mechanical disruption method. Alternatively, a cell may be permeabilized to provide access to a plurality of nucleic acid molecules included therein. Permeabilization may involve partially or completely dissolving or disrupting a cell membrane or a portion thereof. Permeabilization may be achieved by, for example, contacting a cell membrane with an organic solvent or a detergent such as Triton X-100 or NP-40. By lysing or permeabilizing a cell, cell bead, or cell nucleus within a partition (e.g., droplet) to provide access to the plurality of nucleic acid molecules and/or template nucleic acid fragments therein, molecules originating from the same cell, cell bead, or cell nucleus may be isolated within the same partition.

[00147] A partition of a plurality of partitions (e.g., a partition comprising a cell, cell bead, and/or cell nucleus) may comprise one or more beads (e.g., gel beads). A bead may be a gel bead. A bead may comprise a plurality of nucleic acid barcode molecules (e.g., nucleic acid molecules each comprising one or more barcode sequences, as described herein). A bead may comprise at least 10,000 nucleic acid barcode molecules attached thereto. For example, the bead may comprise at least 100,000, 1,000,000, or 10,000,000 nucleic acid barcode molecules attached thereto. The plurality of nucleic acid barcode molecules may be releasably attached to the bead. The plurality of nucleic acid barcode molecules may be releasable from the bead upon application of a stimulus. Such a stimulus may be selected from the group consisting of a thermal stimulus, a photo stimulus, and a chemical stimulus. For example, the stimulus may be a reducing agent such as dithiothreitol. Application of a stimulus may result in one or more of (i) cleavage of a linkage between nucleic acid barcode molecules of the plurality of nucleic acid barcode molecules and the bead, and (ii) degradation or dissolution of the bead to release nucleic acid barcode molecules of the plurality of nucleic acid barcode molecules from the bead.

[00148] A plurality of nucleic acid barcode molecules attached (e.g., releasably attached) to a bead (e.g., gel bead) may be suitable for barcoding template nucleic acid fragments or additional template nucleic acid fragments deriving from DNA and/or RNA molecules of the plurality of cells, cell beads, and/or cell nuclei. For example, a nucleic acid barcode molecule of a plurality of nucleic acid barcode molecule may comprise a barcode sequence, unique molecular identifier (UMI) sequence, primer sequence, universal primer sequence, sequencing adapter or primer,

flow cell adapter sequence, or any other useful feature. In an example, a nucleic acid barcode molecule of a plurality of nucleic acid barcode molecules attached to a bead may comprise a flow cell adapter sequence (e.g., a P5 or P7 sequence), a barcode sequence, a capture sequence, and a sequencing primer sequence or portion thereof (e.g., an R1 or R2 sequence or portion thereof), or a complement of any of these sequences. These sequences may be arranged in any useful order and may be linked or may include one or more spacer sequences disposed between them. For instance, the flow cell adapter sequence, where present, may be disposed near (e.g., proximal to) an end of the nucleic acid barcode molecule that is closest to the bead, while the sequencing primer or portion thereof may be disposed at an end of the nucleic acid barcode molecule that is furthest from (e.g., distal to) the bead (e.g., most available to template nucleic acid fragments for interaction). In another example, a nucleic acid barcode molecule of a plurality of nucleic acid barcode molecules attached to a bead may comprise a flow cell adapter sequence (e.g., a P5 or P7 sequence), a barcode sequence, a sequencing primer sequence or portion thereof (e.g., an R1 or R2 sequence or portion thereof), and a UMI sequence, or a complement of any of these sequences. The nucleic acid barcode molecule may further comprise a capture sequence, which capture sequence may be a targeted capture sequence or comprise a template switch sequence (e.g., comprising a polyC or poly G sequence). These sequences may be arranged in any useful order and may be linked or may include one or more spacer sequences disposed between them. For instance, the flow cell adapter sequence may be disposed near (e.g., proximal to) an end of the nucleic acid barcode molecule that is closest to the bead, while the capture sequence or template switch sequence may be disposed at an end of the nucleic acid barcode molecule that is furthest from the bead (e.g., most available to template nucleic acid fragments for interaction).

[00149] All of the nucleic acid barcode molecules attached (e.g., releasably attached) to a bead (e.g., gel bead) of a plurality of beads may be the same. For example, all of the nucleic acid barcode molecules attached to the bead may have the same nucleic acid sequence. In such an instance, all of the nucleic acid barcode molecules attached to the bead may comprise the same flow cell adapter sequence, sequencing primer or portion thereof, and/or barcode sequence. The barcode sequence of a plurality of nucleic acid barcode molecules attached to a bead of a plurality of beads may be different from other barcode sequences of other nucleic acid barcode molecules attached to other beads of the plurality of beads. For example, a plurality of beads may comprise a plurality of barcode sequences, such that, for at least a subset of the plurality of beads, each bead comprises a different barcode sequence of the plurality of barcode sequences. This differentiation may permit template nucleic acid fragments (e.g., included within cells, cell

beads, and/or cell nuclei) co-partitioned with a plurality of beads between a plurality of partitions to be differentially barcoded within their respective partitions, such that the template nucleic acid fragments or molecules derived therefrom may be identified with the partition (and thus the cell, cell bead, and/or cell nucleus) to which they correspond (e.g., using a nucleic acid sequencing assay, as described herein). A barcode sequence may comprise between 4-20 nucleotides. A barcode sequence may comprise one or more segments, which segments may range in size from 2-20 nucleotides, such as from 4-20 nucleotides. Such segments may be combined to form barcode sequences using a combinatorial assembly method, such as a split-pool method. Details of such methods can be found, for example, in PCT/US2018/061391, filed November 15, 2018, and U.S. Pat. Pub. 20190249226, each of which are herein incorporated by reference in their entireties.

[00150] In some cases, nucleic acid barcode molecules attached to a bead may not be the same. For example, the plurality of nucleic acid barcode molecules attached to a bead may each comprise a UMI sequence, which UMI sequence varies across the plurality of nucleic acid barcode molecules. All other sequences of the plurality of nucleic acid barcode molecules attached to the bead may be the same.

[00151] In some cases, a bead may comprise multiple different nucleic acid barcode molecules attached thereto. For example, a bead may comprise a first plurality of nucleic acid barcode molecules and a second plurality of nucleic acid barcode molecules, which first plurality of nucleic acid barcode molecules is different than the second plurality of nucleic acid barcode molecules. The first plurality of nucleic acid barcode molecules and the second plurality of nucleic acid barcode molecules coupled to a bead may comprise one or more shared sequences. For example, each nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules and each nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules may comprise the same barcode sequence (e.g., as described herein). Such a barcode sequence may be prepared using a combinatorial assembly process (e.g., as described herein). For example, barcode sequences may comprise identical barcode sequence segments. Similarly, each nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules coupled to a bead may comprise the same flow cell adapter sequence and/or sequencing primer or portion thereof as each nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules coupled to the bead. In an example, each nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules coupled to a bead comprises a sequencing primer, and each nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules coupled to the bead comprises a portion of the same

sequencing primer. In some instances, each nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules coupled to a bead may comprise a first sequencing primer (e.g., a TruSeq R1 sequence), a barcode sequence, and a first functional sequence, and each nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules coupled to the bead may comprise a second sequencing primer (e.g., a Nextera R1 sequence, or a portion thereof), the barcode sequence, and a second functional sequence. Sequences shared between different sets of nucleic acid barcode molecules coupled to the same bead may be included in the same or different order and may be separated by the same or different sequences. Alternatively or in addition, the first plurality of nucleic acid barcode molecules and the second plurality of nucleic acid barcode molecules coupled to a bead may include one or more different sequences. For example, each nucleic acid barcode molecule of a first plurality of nucleic acid barcode molecules coupled to a bead of a plurality of beads may comprise one or more of a flow cell adapter sequence, a barcode sequence, UMI sequence, capture sequence, and a sequencing primer or portion thereof, while each nucleic acid barcode molecule of a second plurality of nucleic acid barcode molecules coupled to the bead may comprise one or more of a flow cell adapter sequence (e.g., the same flow cell adapter sequence), a barcode sequence (e.g., the same barcode sequence), UMI sequence, capture sequence, and a sequencing primer or portion thereof (e.g., the same sequencing primer or portion thereof). Nucleic acid barcode molecules of the first plurality of nucleic acid barcode molecules may not include a UMI sequence or capture sequence. A bead comprising multiple different populations of nucleic acid barcode molecules, such as a first plurality of nucleic acid molecules and a second plurality of nucleic acid molecules (e.g., as described above), may be referred to as a “multi-functional bead.”

[00152] A cell, cell bead, or cell nucleus comprising template nucleic acid fragments (e.g., template nucleic acid fragments and additional template nucleic acid fragments deriving from DNA or RNA molecules included within the cell, cell bead, or cell nucleus) may be co-partitioned with one or more beads (e.g., as described herein). For example, a cell, cell bead, or cell nucleus may be co-partitioned with a first bead (e.g., first gel bead) configured to interact with a first set of template nucleic acid fragments (e.g., template nucleic acid fragments deriving from DNA molecules, such as tagged fragments) and a second bead (e.g., second gel bead) configured to interact with a second set of template nucleic acid fragments (e.g., additional template nucleic acid fragments deriving from RNA molecules). The first bead may comprise a first nucleic acid molecule comprising a flow cell adapter sequence, a barcode sequence, and a sequencing primer or portion thereof, which sequencing primer or portion thereof may be configured to interact with (e.g., anneal or hybridize to) a complementary sequence included in

template nucleic acid fragments deriving from DNA molecules of the cell, cell bead, or cell nucleus, or derivatives thereof. The second bead may comprise a second nucleic acid molecule comprising the flow cell adapter sequence, the barcode sequence, the sequencing primer or a portion thereof, a UMI sequence, and a capture sequence, which capture sequence may be configured to interact with (e.g., anneal or hybridize to) a sequence of template nucleic acid fragments deriving from RNA molecules of the cell, cell bead, or cell nucleus, or derivatives thereof. In some cases, the capture sequence may be configured to interact with a sequence of a cDNA molecule generated upon reverse transcription of an RNA fragment. The first and second beads may be linked together (e.g., covalently or non-covalently). The first and second beads may each comprise a plurality of nucleic acid molecules. For example, the first bead may comprise a plurality of first nucleic acid molecules and the second bead may comprise a plurality of second nucleic acid molecules, where each first nucleic acid molecule of the plurality of first nucleic acid molecules comprises a first shared sequence and each second nucleic acid molecule of the plurality of second nucleic acid molecules comprises a second shared sequence. The first shared sequence and the second shared sequence may be the same or different. The first shared sequence and the second shared sequence may comprise one or more shared components, such as a shared barcode sequence or sequencing primer or portion thereof.

[00153] Alternatively, a cell, cell bead, or cell nucleus comprising template nucleic acid fragments (e.g., template nucleic acid fragments or additional template nucleic acid fragments deriving from DNA or RNA molecules included within the cell, cell bead, or cell nucleus) may be co-partitioned with a single bead (e.g., gel bead). For example, a cell, cell bead, or cell nucleus may be co-partitioned with a bead comprising (i) a first plurality of nucleic acid barcode molecules configured to interact with a first set of template nucleic acid fragments (e.g., template nucleic acid fragments deriving from DNA molecules, such as tagged fragments), or derivatives thereof, and (ii) a second plurality of nucleic acid barcode molecules configured to interact with a second set of template nucleic acid fragments (e.g., additional template nucleic acid fragments deriving from RNA molecules), or derivatives thereof (such as cDNA generated from an RNA fragment). A nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules may comprise a flow cell adapter sequence, a barcode sequence, and a sequencing primer or portion thereof, which sequencing primer or portion thereof may be configured to interact with (e.g., anneal or hybridize to) a complementary sequence included in template nucleic acid fragments deriving from DNA molecules of the cell, cell bead, or cell nucleus, or derivatives thereof. A nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules may comprise the flow cell adapter sequence, the barcode

sequence, the sequencing primer or a portion thereof, a UMI sequence, and a capture sequence, which capture sequence may be configured to interact with (e.g., anneal or hybridize to) a sequence of template nucleic acid fragments deriving from RNA molecules of the cell, cell bead, or cell nucleus, or derivatives thereof, such as cDNA generated from an RNA fragment. The first plurality of nucleic acid barcode molecules may comprise approximately the same number of nucleic acid barcode molecules as the second plurality of nucleic acid barcode molecules. Alternatively, the first plurality of nucleic acid barcode molecules may comprise a greater number of nucleic acid barcode molecules than the second plurality of nucleic acid barcode molecules, or vice versa. The distribution of nucleic acid barcode molecules on a bead may be controlled by, for example, sequence control, concentration control, and or blocking methods during assembly of the nucleic acid barcode molecules on the bead. Details of such processes are provided in, for example, PCT/US2018/061391, filed November 15, 2018, and U.S. Pat. Pub. 20190249226, each of which are incorporated by reference in their entireties.

[00154] **FIGs. 24A and 24B** show examples of beads for use according to the method provided herein. **FIG. 24A** shows a first bead **2401** and a second bead **2411** that may be co-partitioned with a cell, cell bead, or cell nucleus into a partition of a plurality of partitions (e.g., droplets or wells). First bead **2401** may comprise nucleic acid molecule **2402**. Nucleic acid molecule **2402** may comprise sequences **2403**, **2404**, and **2405**. Sequence **2403** may be, for example, a flow cell adapter sequence (e.g., a P5 or P7 sequence). Sequence **2404** may be, for example, a barcode sequence. Sequence **2405** may be, for example, a sequencing primer sequence or portion thereof (e.g., an R1 or R2 primer sequence, or portion thereof). Nucleic acid molecule **2402** may also include additional sequences, such as a UMI sequence. First bead **2401** may comprise a plurality of nucleic acid molecules **2402**. Second bead **2411** may comprise nucleic acid molecule **2412**. Nucleic acid molecule **2412** may comprise sequences **2413**, **2414**, and **2415**. Sequence **2413** may be, for example, a flow cell adapter sequence (e.g., a P5 or P7 sequence). Sequence **2414** may be, for example, a barcode sequence. Sequence **2415** may be, for example, a sequencing primer sequence or portion thereof (e.g., an R1 or R2 primer sequence, or portion thereof). Nucleic acid molecule **2412** may also include additional sequences, such as a UMI sequence and a capture sequence. Second bead **2401** may comprise a plurality of nucleic acid molecules **2412**.

[00155] **FIG. 24B** shows a bead **2421** (e.g., a multifunctional bead having two or more species of nucleic acid barcode molecules attached or coupled thereto) that may be co-partitioned with a cell, cell bead, or cell nucleus into a partition of a plurality of partitions (e.g., droplets or wells). Bead **2421** may comprise nucleic acid molecule **2422** and nucleic acid molecule **2426**.

Nucleic acid molecule **2422** may comprise sequences **2423**, **2424**, and **2425**. Sequence **2423** may be, for example, a flow cell adapter sequence (e.g., a P5 or P7 sequence). Sequence **2424** may be, for example, a barcode sequence. Sequence **2425** may be, for example, a sequencing primer or portion thereof (e.g., an R1 or R2 primer sequence, or portion thereof, such as a Nextera R1 sequence or portion thereof). In some instances, sequence **2425** may also be, for example, a sequence configured to hybridize to a splint oligonucleotide as described elsewhere herein. Nucleic acid molecule **2426** may comprise sequences **2427**, **2428**, and **2429**. Sequence **2427** may be, for example, a flow cell adapter sequence (e.g., a P5 or P7 sequence). Sequence **2428** may be, for example, a barcode sequence (e.g., the same barcode sequence as sequence **2424**). Sequence **2429** may be, for example, a sequencing primer or portion thereof (e.g., an R1 or R2 primer sequence, or portion thereof). Sequence **2427** may be, for example, a sequencing primer or portion thereof (e.g., an R1 or R2 primer sequence, or portion thereof, such as a TruSeq R1 sequence, or portion thereof). Sequence **2428** may be, for example, a barcode sequence (e.g., the same barcode sequence as **2424**). Sequence **2429** may be, for example, a capture sequence (e.g., a poly-T sequence), such as a capture sequence that is configured to hybridize with a target nucleic acid molecule (e.g., mRNA molecule). Sequence **2429** may be, for example, a template switching oligonucleotide (TSO) sequence configured to facilitate a template switching reaction with a target nucleic acid molecule (e.g., mRNA molecule). Sequence **2423** and sequence **2427** may be the same. Alternatively, sequence **2423** and sequence **2427** may be different. Sequence **2424** and sequence **2428** may be the same. Alternatively, sequence **2424** and sequence **2428** may be different. Sequence **2425** and sequence **2429** may be the same. Alternatively, sequence **2425** and sequence **2429** may be different. Nucleic acid molecules **2422** and **2426** may also include additional sequences, such as a UMI sequence and a capture sequence. Bead **2421** may comprise a plurality of nucleic acid molecules **2422** and a plurality of nucleic acid molecules **2426**.

[00156] Within a partition (e.g., as described herein), an RNA fragment (e.g., a molecule comprising a sequence of an RNA molecule of a cell, cell bead, or cell nucleus that is hybridized to a primer molecule) may be processed to provide a barcoded molecule. The RNA fragment may be reverse transcribed to generate a complementary cDNA strand, which cDNA strand may be barcoded. In some cases, template switching can be used to increase the length of a cDNA (e.g., via incorporation of one or more sequences, such as one or more barcode or unique molecular identifier sequences). In one example of template switching, cDNA can be generated from reverse transcription of a template (e.g., an mRNA molecule) where a reverse transcriptase with terminal transferase activity can add additional nucleotides, e.g., polyC, to the cDNA that

are not encoded by the **template**, such, as at an end of the cDNA. Template switch oligonucleotides (e.g., switch oligos) can include sequences complementary to the additional nucleotides, e.g. polyG (such as poly-riboG). The additional nucleotides (e.g., polyC) on the cDNA can hybridize to the sequences complementary to the additional nucleotides (e.g., polyG) on the template switch oligonucleotide, whereby the template switch oligonucleotide can be used by the reverse transcriptase as **template** to further extend the cDNA. Template switch oligonucleotides may comprise deoxyribonucleic acids, ribonucleic acids, modified nucleic acids including locked nucleic acids (LNA), or any combination thereof. A template switch oligonucleotide may comprise one or more sequences including, for example, one or more sequences selected from the group consisting of a sequencing primer, a barcode sequence, a unique molecular identifier sequence, and a homopolymer sequence (e.g., a polyG sequence), or a complement of any of the preceding sequence.

[00157] In some cases, the length of a template switch oligonucleotide may be 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250 nucleotides or longer.

[00158] In some cases, an adapter and/or barcode sequence may be added to an RNA molecule via a method other than template switching. For example, one or more sequences may be ligated to an end of an RNA molecule. Similarly, one or more sequences may be ligated to an end of a cDNA molecule generated via reverse transcription of an RNA molecule.

[00159] In an example, a cell, cell bead, or cell nucleus comprising chromatin and one or more RNA molecules is provided. The chromatin in the cell, cell bead, or cell nucleus may be processed to provide a first template nucleic acid fragment derived from the chromatin (e.g., a tagged fragment, as described herein). The chromatin may be processed in bulk solution. An RNA molecule may be processed to provide a second template nucleic acid fragment derived

from the RNA molecule (e.g., as described herein). The RNA molecule may be processed within a partition. The configuration of the first template nucleic acid fragment may be at least partially dependent on the structure of the transposase-nucleic acid complex used to generate the first template nucleic acid fragment. For example, a transposase-nucleic acid complex such as that shown in **FIG. 9** may be used to prepare the first template nucleic acid fragment. The first template nucleic acid fragment may be at least partially double-stranded. The first template nucleic acid fragment may comprise a double-stranded region comprising sequences of chromatin of the cell, cell bead, or cell nucleus. A first end of a first strand of the double-stranded region may be linked to a first transposon end sequence (e.g., mosaic end sequence), which first transposon end sequence may be linked to a first sequencing primer or portion thereof. A first end of the second strand of the double-stranded region, which end is opposite the first end of the first strand, may be linked to a second transposon end sequence (e.g., mosaic end sequence), which second transposon end sequence may be linked to a second sequencing primer or portion thereof. The second transposon end sequence may be the same as or different from the first transposon end sequence. The first sequencing primer or portion thereof may be the same as or different from the second sequencing primer or portion thereof. In some cases, the first sequencing primer or portion thereof may be an R1 sequence or portion thereof, and the second sequencing primer or portion thereof may be an R2 sequence or portion thereof. The first transposon end sequence may be hybridized to a first complementary sequence (e.g., mosaic end reverse complement sequence), which first complementary sequence may not be linked to a second end of the second strand of the double-stranded region of the first template nucleic acid fragment. Similarly, the second transposon end sequence may be hybridized to a second complementary sequence (e.g., mosaic end reverse complement sequence), which second complementary sequence may not be linked to a second end of the first strand of the double-stranded region of the first template nucleic acid fragment. In other words, the first template nucleic acid fragment may comprise one or more gaps. In some cases, the one or more gaps may be approximately 9 bp in length each. The second template nucleic acid fragment (e.g., an additional template nucleic acid fragment) may comprise a sequence of an RNA molecule of the cell, cell bead, or cell nucleus and a sequence hybridized to a primer molecule (e.g., a capture nucleic acid molecule). For example, the second template nucleic acid fragment may comprise a sequence of an RNA molecule of the cell, cell bead, or cell nucleus and a polyA sequence hybridized to a polyT sequence of a primer molecule. The primer molecule may also comprise an additional primer sequence.

[00160] The cell, cell bead, or cell nucleus comprising the first template nucleic acid fragment (e.g., tagged fragment) may be co-partitioned with one or more reagents into a partition of a plurality of partitions (e.g., as described herein). The partition may be, for example, a droplet or well. The partition may comprise one or more beads (e.g., as described herein). A bead of the one or more beads may comprise a first plurality of nucleic acid barcode molecules. A nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules may comprise one or more of a flow cell adapter sequence (e.g., P5 sequence), a barcode sequence, a sequencing primer or portion thereof (e.g., R1 sequence or portion thereof, or a complement thereof), and a sequence configured to hybridize to a splint oligonucleotide as described elsewhere herein. The sequencing primer or portion thereof may be complementary to a sequence of the first template nucleic acid fragment. A bead of the one or more beads may also comprise a second plurality of nucleic acid barcode molecules. A nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules may comprise one or more of a flow cell adapter sequence (e.g., P5 sequence), a barcode sequence, a sequencing primer or portion thereof (e.g., R1 sequence or portion thereof, or a complement thereof), and a sequence configured to hybridize to a splint oligonucleotide as described elsewhere herein. For example, a nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules may comprise one or more of a flow cell adapter sequence (e.g., P5 sequence), a barcode sequence, and a sequence configured to hybridize to a splint oligonucleotide as described elsewhere herein. A nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules may comprise a sequencing primer or portion thereof (e.g., R1 sequence or portion thereof, or a complement thereof), a barcode sequence, and a capture sequence (e.g., a poly T sequence) configured to hybridize to a nucleic acid molecule (e.g., RNA molecule). In some cases, the first plurality of nucleic acid barcode molecules and the second plurality of nucleic acid barcode molecules may be same.

[00161] Within the partition, the RNA molecule may be processed to provide the second template nucleic acid fragment (e.g., as described herein).

[00162] Within the partition, the cell, cell bead, or cell nucleus may be lysed or permeabilized to provide access to the first and/or second template nucleic acid fragments therein (e.g., as described herein). The second template nucleic acid fragment may be generated after the cell, cell bead, or cell nucleus is lysed or permeabilized.

[00163] The first and second template nucleic acid fragments may undergo processing within the partition. Within the partition, the gaps in the first template nucleic acid molecule may be filled via a gap filling extension process (e.g., using a DNA polymerase or reverse transcriptase). The resultant double-stranded nucleic acid molecule may be denatured to provide a single strand

comprising a chromatin sequence flanked by transposon end sequences and/or sequences complementary to transposon end sequences. Each transposon end sequence and/or sequence complementary to transposon end sequence may be linked to a sequencing primer or portion thereof, or a complement thereof (e.g., an R1 or R2 sequence or a portion thereof, or a complement thereof). A nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules may hybridize to a sequencing primer or portion thereof, or a complement thereof, of the single strand. A primer extension reaction may then be used to generate a complement of the single strand (e.g., using a DNA polymerase or reverse transcriptase). Such a process may amount to a linear amplification process. This process incorporates the barcode sequence of the nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules, or a complement thereof. The resultant double-stranded molecule may be denatured to provide a single strand comprising the flow cell adapter sequence, or complement thereof, of the nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules; barcode sequence, or complement thereof, of the nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules; sequencing primer or portion thereof, or complement thereof, of the nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules; transposon end sequences, and/or complements thereof; second sequencing primer or portion thereof, or complement thereof. An additional amplification process may or may not be performed within a partition. For example, exponential amplification may or may not be performed within a partition.

[00164] Within the partition, the second template nucleic acid fragment derived from the RNA molecule of the cell, cell bead, or cell nucleus may be reverse transcribed (e.g., using a reverse transcriptase) to provide a cDNA strand. The reverse transcription process may append a sequence to an end of a strand of the resultant double-stranded nucleic acid molecule comprising the RNA strand and the cDNA strand, such as a polyC sequence. A template switching oligonucleotide may comprise a sequence (e.g., a polyG sequence) that may hybridize to at least a portion of the double-stranded nucleic acid molecule (e.g., to the appended polyC sequence) and be used to further extend the strand of the double-stranded nucleic acid molecule to provide an extended double-stranded nucleic acid molecule. Such a sequence may comprise ribobases. The template switching oligonucleotide may comprise a UMI sequence, or complement thereof, and a sequencing primer or portion thereof, or complement thereof. The extended double-stranded nucleic acid molecule comprising the template switching oligonucleotide and a complement thereof, and the prior double-stranded nucleic acid molecule may be denatured to provide a single strand comprising a sequencing primer or portion thereof, or complement

thereof, of the nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules; the UMI sequence, or complement thereof; the poly(C) or poly(G) sequence; the sequence corresponding to the RNA molecule of the cell, cell bead, or cell nucleus, or complement thereof; and sequences of the capture nucleic acid molecule, or complements thereof. A nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules may hybridize to a sequencing primer or portion thereof, or a complement thereof, of the single strand. A primer extension reaction may then be used to generate a complement of the single strand (e.g., using a DNA polymerase). Such a process may amount to a linear amplification process. This process incorporates the barcode sequence of the nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules, or a complement thereof. The resultant double-stranded molecule may be denatured to provide a single strand comprising a flow cell adapter sequence, or complement thereof, of the nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules; a barcode sequence, or complement thereof, of the nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules; a sequencing primer or portion thereof, or complement thereof, of the nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules; the UMI sequence, or complement thereof; the poly(C) or poly(G) sequence; the sequence corresponding to the RNA molecule of the cell, cell bead, or cell nucleus, or complement thereof; and sequences of the capture nucleic acid molecule, or complements thereof. An additional amplification process may or may not be performed within a partition. For example, exponential amplification may or may not be performed within a partition.

[00165] The linear amplification products corresponding to the chromatin and the RNA molecule of the cell, cell bead, or cell nucleus included within the partition of the plurality of partitions may be recovered from the partition. For example, the contents of the plurality of partitions may be pooled to provide the linear amplification products in a bulk solution. The linear amplification product corresponding to the chromatin may then be subjected to conditions sufficient to undergo one or more nucleic acid amplification reactions (e.g., PCR) to generate one or more amplification products corresponding to the chromatin. A nucleic acid amplification process may incorporate one or more additional sequences, such as one or more additional flow cell adapter sequences. The linear amplification product corresponding to the RNA molecule may be subjected to fragmentation, end repair, and dA tailing processes. An additional primer sequence (e.g., a sequencing primer or portion thereof, such as an R2 sequence) may then be ligated to the resultant molecule. A nucleic acid amplification reaction (e.g., PCR) may then be performed to generate one or more amplification products

corresponding to the RNA molecule. A nucleic acid amplification process may incorporate one or more additional sequences, such as one or more additional flow cell adapter sequences (see, for example, **FIG. 12**).

[00166] In an RNA workflow, in-partition template switching may attach a sequencing primer (e.g., a TruSeq R1 or R2 sequence) to the 3' or 5' end of an RNA transcript. A bead (e.g., gel bead) carrying the sequencing primer, or portion thereof (e.g., partial TruSeq R1 or R2 sequence) may be also used for priming in a DNA (e.g., chromatin) workflow. This may allow for differential amplification of DNA (e.g., ATAC) and RNA libraries after removing materials from partitions (e.g., breaking emulsions) and sample splitting. Another advantage of this method is that the same enzyme (e.g. DNA polymerase or reverse transcriptase) may be used to barcode nucleic acid fragments derived from both DNA (e.g., chromatin) and RNA.

[00167] **FIG. 12** shows an example schematic corresponding to the preceding example. Panel **1200** shows a workflow corresponding to processing of chromatin from a cell, cell bead, or cell nucleus, and panel **1250** shows a workflow corresponding to processing of an mRNA molecule from the cell, cell bead, or cell nucleus. In the figure, two distinct beads (e.g., gel beads) are shown. However, the same bead (e.g., a single gel bead that may be a multifunctional bead) may be used in each workflow.

[00168] As shown in panel **1200**, in bulk solution, chromatin included within a cell, cell bead, or cell nucleus is processed (e.g., as described herein) to provide a template nucleic acid fragment (e.g., tagged fragment) **1204** comprising insert sequence **1208** (e.g., region of open chromatin) and a complement thereof, transposon end sequences **1206** and complements thereof, sequencing primer or portion thereof **1202** (e.g., an R1 sequence), sequencing primer or portion thereof **1210** (e.g., an R2 sequence), and gaps **1207**. Template nucleic acid fragment **1204** may then be partitioned within a partition (e.g., a droplet or well, as described herein). Within the partition, the cell, cell bead, or cell nucleus comprising template nucleic acid fragment **1204** may be lysed, permeabilized, or otherwise processed to provide access to template nucleic acid fragment **1204** (and one or more RNA molecules) therein. Gaps **1207** may be filled **1212** via a gap filling extension process (e.g., using a DNA polymerase). The partition may include a bead (e.g., gel bead) **1216a** coupled to a nucleic acid barcode molecule **1218a**. Nucleic acid barcode molecule **1218a** may comprise a flow cell adapter sequence **1220a** (e.g., a P5 sequence), a barcode sequence **1222a**, and a sequencing primer or portion thereof or complement thereof **1202'**. Sequence **1202'** may hybridize to sequence **1202** of template nucleic acid fragment **1204**, or its complement, and undergo primer extension **1214** to yield a strand comprising sequences **1220a**, **1222a**, **1202'**, **1210**, and insert sequence **1208** or a complement thereof. The contents of

the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the strand in bulk solution. This strand may undergo amplification (e.g., PCR) **1224** to provide a double-stranded amplification product **1226** that includes sequences of the nucleic acid barcode molecule **1218a**, the original chromatin molecule, and, optionally, an additional sequence **1228** that may be a flow cell adapter sequence (e.g., a P7 sequence).

[00169] In parallel to the chromatin workflow of panel **1200**, an RNA molecule deriving from the same cell, cell bead, or cell nucleus may be processed. As shown in panel **1250**, RNA molecule **1258** comprising RNA sequence **1260** and polyA sequence **1262** may be contacted **1264** with primer molecule **1252** comprising polyT sequence **1254** and additional primer sequence **1256**. RNA molecule **1258** may then be reverse transcribed **1266** off of polyT sequence **1254** using a reverse transcriptase with terminal transferase activity, which reverse transcriptase may append sequence **1268** to the resultant cDNA molecule comprising cDNA sequence **1270**. Sequence **1268** may be a polyC sequence. A template switch oligonucleotide **1272** comprising sequencing primer or portion thereof or complement thereof **1274**, unique molecule identifier sequence or complement thereof **1276**, and capture sequence (e.g., polyG sequence) **1278** may then hybridize **1280** to the RNA-cDNA molecule and template switching may take place. The partition may include a bead (e.g., gel bead) **1216b** coupled to a nucleic acid barcode molecule **1218b**. Nucleic acid barcode molecule **1218b** may comprise a flow cell adapter sequence **1220b** (e.g., a P5 sequence), a barcode sequence **1222b**, and a sequencing primer or portion thereof or complement thereof **1274'**. Bead **1216b** may be the same as bead **1216a** such that partition comprises a single bead (e.g., **1218a** and **1218b** are attached to a single bead). In such a case, nucleic acid barcode molecule **1218b** and nucleic acid barcode molecule **1218a** may have the same sequences. Sequence **1274'** may hybridize to sequence **1274** of the cDNA molecule, or its complement, and undergo primer extension **1282** to yield a strand comprising sequences **1220b**, **1222b**, **1274'**, **1276** or a complement thereof, **1268** or a complement thereof, and insert sequence **1270** or a complement thereof. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the strand in bulk solution. This strand may undergo amplification (e.g., PCR) **1284** to provide a double-stranded amplification product **1286** that includes sequences of the nucleic acid barcode molecule **1218b**, the original RNA molecule or cDNA corresponding thereto, and, optionally, an additional sequence **1288** that may comprise a sequencing primer or portion thereof (e.g., an R2 sequence) **1290**, a sample index sequence **1292**, and a flow cell adapter sequence (e.g., a P7 sequence) **1294**.

[00170] FIG. 13 shows another example schematic corresponding to the preceding example. Panel 1300 shows a workflow corresponding to processing of chromatin from a cell, cell bead, or cell nucleus, and panel 1350 shows a workflow corresponding to processing of an mRNA molecule from the cell, cell bead, or cell nucleus. In the figure, two distinct beads (e.g., gel beads) are shown. However, the same bead (e.g., gel bead) may be used in each workflow.

[00171] As shown in panel 1300, in bulk solution, chromatin included within a cell, cell bead, or cell nucleus is processed (e.g., as described herein) to provide a template nucleic acid fragment (e.g., tagged fragment) 1304 comprising insert sequence 1308 (e.g., region of open chromatin) and a complement thereof, transposon end sequences 1306 and complements thereof, sequencing primer or portion thereof 1302 (e.g., an R1 sequence), sequencing primer or portion thereof 1310 (e.g., an R2 sequence), and gaps 1307. Template nucleic acid fragment 1304 may then be partitioned within a partition (e.g., a droplet or well, as described herein). Within the partition, the cell, cell bead, or cell nucleus comprising template nucleic acid fragment 1304 may be lysed, permeabilized, or otherwise processed to provide access to template nucleic acid fragment 1304 (and one or more RNA molecules) therein. Gaps 1307 may be filled 1312 via a gap filling extension process (e.g., using a DNA polymerase). The partition may include a bead (e.g., gel bead) 1316a coupled to a nucleic acid barcode molecule 1318a. Nucleic acid barcode molecule 1318a may comprise a flow cell adapter sequence 1320a (e.g., a P5 sequence), a barcode sequence 1322a, and a sequencing primer or portion thereof or complement thereof 1302'. Sequence 1302' may hybridize to sequence 1302 of template nucleic acid fragment 1304, or its complement, and undergo primer extension 1314 to yield a strand comprising sequences 1320a, 1322a, 1302', 1310, and insert sequence 1308 or a complement thereof. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the strand in bulk solution. This strand may undergo amplification (e.g., PCR) 1324 to provide a double-stranded amplification product 1326 that includes sequences of the nucleic acid barcode molecule 1318a, the original chromatin molecule, and, optionally, an additional sequence 1328 that may be a flow cell adapter sequence (e.g., a P7 sequence).

[00172] In parallel to the chromatin workflow of panel 1300, an RNA molecule deriving from the same cell, cell bead, or cell nucleus may be processed. As shown in panel 1350, RNA molecule 1358 comprising RNA sequence 1360 and polyA sequence 1362 may be contacted with primer molecule 1352 comprising polyT sequence 1354, UMI sequence 1355, and sequencing primer or portion thereof (e.g., R1 sequence) 1356. RNA molecule 1358 may be reverse transcribed 1364 off of polyT sequence 1354 using a reverse transcriptase with terminal transferase activity, which reverse transcriptase may append sequence 1366 (e.g., a polyC

sequence) to the resultant cDNA molecule comprising cDNA sequence **1368**. A template switch oligonucleotide **1370** comprising additional primer sequence **1372** and a homopolymer sequence **1374** (e.g., a polyG) sequence that is complementary to sequence **1366** may then hybridize **1376** to the cDNA molecule and template switching may take place. The partition may include a bead (e.g., gel bead) **1316b** coupled to a nucleic acid barcode molecule **1318b**. Nucleic acid barcode molecule **1318b** may comprise a flow cell adapter sequence **1320b** (e.g., a P5 sequence), a barcode sequence **1322b**, and a sequencing primer or portion thereof or complement thereof **1356'**. Bead (e.g., gel bead) **1316b** may be the same as bead (e.g., gel bead) **1316a** such that partition comprises a single bead (i.e., **1318a** and **1318b** are attached to a single bead). In such a case, nucleic acid barcode molecule **1318b** and nucleic acid barcode molecule **1318a** may have the same sequences. Sequence **1356'** may hybridize to sequence **1356** of the cDNA molecule, or its complement, and undergo primer extension **1378** to yield a strand comprising sequences **1320b**, **1322b**, **1356'**, **1355** or a complement thereof, **1366** or a complement thereof, and insert sequence **1368** or a complement thereof. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the strand in bulk solution. This strand may undergo amplification (e.g., PCR) **1380** to provide a double-stranded amplification product **1382** that includes sequences of the nucleic acid barcode molecule **1318b**, the original RNA molecule or cDNA corresponding thereto, and, optionally, an additional sequence **1384** that may comprise a sequencing primer or portion thereof (e.g., an R2 sequence) **1390**, a sample index sequence **1388**, and a flow cell adapter sequence (e.g., a P7 sequence) **1386**.

[00173] In another example, a cell, cell bead, or cell nucleus comprising chromatin and one or more RNA molecules is provided. The chromatin in the cell, cell bead, or cell nucleus may be processed to provide a first template nucleic acid fragment derived from the chromatin (e.g., a tagged fragment, as described herein). The chromatin may be processed in bulk solution. An RNA molecule may be processed to provide a second template nucleic acid fragment derived from the RNA molecule (e.g., as described herein). The RNA molecule may be processed within a partition. The configuration of the first template nucleic acid fragment may be at least partially dependent on the structure of the transposase-nucleic acid complex used to generate the first template nucleic acid fragment. For example, a transposase-nucleic acid complex such as that shown in **FIG. 9** may be used to prepare the first template nucleic acid fragment. The first template nucleic acid fragment may be at least partially double-stranded. The first template nucleic acid fragment may comprise a double-stranded region comprising sequences of chromatin of the cell, cell bead, or cell nucleus. A first end of a first strand of the double-stranded region may be linked to a first transposon end sequence (e.g., mosaic end sequence),

which first transposon end sequence may be linked to a first sequencing primer or portion thereof. A first end of the second strand of the double-stranded region, which end is opposite the first end of the first strand, may be linked to a second transposon end sequence (e.g., mosaic end sequence), which second transposon end sequence may be linked to a second sequencing primer or portion thereof. The second transposon end sequence may be the same as or different from the first transposon end sequence. The first sequencing primer or portion thereof may be the same as or different from the second sequencing primer or portion thereof. In some cases, the first sequencing primer or portion thereof may be an R1 sequence or portion thereof, and the second sequencing primer or portion thereof may be an R2 sequence or portion thereof. The first transposon end sequence may be hybridized to a first complementary sequence (e.g., mosaic end reverse complement sequence), which first complementary sequence may not be linked to a second end of the second strand of the double-stranded region of the first template nucleic acid fragment. Similarly, the second transposon end sequence may be hybridized to a second complementary sequence (e.g., mosaic end reverse complement sequence), which second complementary sequence may not be linked to a second end of the first strand of the double-stranded region of the first template nucleic acid fragment. In other words, the first template nucleic acid fragment may comprise one or more gaps. In some cases, the one or more gaps may be approximately 9 bp in length each. For example, one or more gaps may be at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more bp in length. For example, one or more gaps may be at most about 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 bp in length. The second template nucleic acid fragment (e.g., an additional template nucleic acid fragment) may comprise a sequence of an RNA molecule of the cell, cell bead, or cell nucleus and a sequence hybridized to a primer molecule (e.g., a capture nucleic acid molecule). For example, the second template nucleic acid fragment may comprise a sequence of an RNA molecule of the cell, cell bead, or cell nucleus and a polyA sequence hybridized to a polyT sequence of a primer molecule. The primer molecule may also comprise an additional primer sequence.

[00174] The cell, cell bead, or cell nucleus comprising the first template nucleic acid fragment (e.g., tagmented fragment) may be co-partitioned with one or more reagents into a partition of a plurality of partitions (e.g., as described herein). The partition may be, for example, a droplet or well. The partition may comprise one or more beads (e.g., as described herein). A bead (e.g., gel bead) of the one or more beads may comprise a first plurality of nucleic acid barcode molecules. A nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules may comprise a flow cell adapter sequence (e.g., P5 sequence), a barcode sequence, and a sequencing primer or portion thereof (e.g., R1 sequence or portion thereof, or a

complement thereof). The sequencing primer or portion thereof may be complementary to a sequence of the first template nucleic acid fragment. The flow cell adapter sequence and/or barcode sequence may be hybridized to their complementary sequences. A bead (e.g., gel bead) of the one or more beads may also comprise a second plurality of nucleic acid barcode molecules. A nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules may comprise a flow cell adapter sequence (e.g., P5 sequence), a barcode sequence, a sequencing primer or portion thereof (e.g., R1 sequence or portion thereof, or a complement thereof), a UMI sequence, and a capture sequence (e.g., a polyG sequence, a polydT sequence or target specific sequence). In some cases, the first plurality of nucleic acid barcode molecules and the second plurality of nucleic acid barcode molecules may be coupled to the same bead, and the partition may comprise a single bead.

[00175] Within the partition, the RNA molecule may be processed to provide the second template nucleic acid fragment (e.g., as described herein).

[00176] Within the partition, the cell, cell bead, or cell nucleus may be lysed or permeabilized to provide access to the first and/or second template nucleic acid fragments therein (e.g., as described herein). The second template nucleic acid fragment may be generated after the cell, cell bead, or cell nucleus is lysed or permeabilized.

[00177] The first and second template nucleic acid fragments may undergo processing within the partition. Within the partition, a sequencing primer or portion thereof of the first template nucleic acid fragment corresponding to the chromatin of the cell, cell bead, or cell nucleus may hybridize to a sequencing primer or portion thereof of the nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules. The sequencing primer or portion thereof of the nucleic acid barcode molecule may then be ligated (e.g., using a ligase) to a transposon end sequence of the first template nucleic acid fragment, or a complement thereof to provide a partially double-stranded nucleic acid molecule corresponding to the chromatin of the cell, cell bead, or cell nucleus.

[00178] Within the partition, the second template nucleic acid fragment derived from the RNA molecule of the cell, cell bead, or cell nucleus may be reverse transcribed (e.g., using a reverse transcriptase) to provide a cDNA strand. The reverse transcription process may append a sequence to an end of a strand of the resultant double-stranded nucleic acid molecule comprising the RNA strand and the cDNA strand, such as a polyC sequence. The capture sequence of the nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules may hybridize to the appended sequence (e.g., polyC sequence) of the double-stranded nucleic acid molecule and a template switching process may take place to provide an extended double-

stranded nucleic acid molecule. Such a sequence may comprise ribobases. The sequence of the nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules may be considered a template switching oligonucleotide. Accordingly, barcoding and template switching may take place contemporaneously to provide a barcoded cDNA molecule. The cDNA strand of the barcoded cDNA molecule (e.g., a molecule comprising both a cDNA strand and an RNA strand) may comprise the polyC sequence, a sequence complementary to the sequence of the template switch oligonucleotide or a portion thereof (e.g., sequences complementary to the sequencing primer, barcode sequence, and UMI sequence of the template switch oligonucleotide), the cDNA sequence, the polyT sequence, and the additional primer sequence of the primer molecule. The RNA strand of the barcoded cDNA molecule may comprise the sequence of the template switch oligonucleotide, the mRNA sequence, and a sequence complementary to the additional primer sequence of the primer molecule.

[00179] The partially double-stranded molecule corresponding to the chromatin of the cell, cell bead, or cell nucleus and the barcoded cDNA molecule corresponding to the RNA molecule of the cell, cell bead, or cell nucleus included within the partition (e.g., droplet or well) of the plurality of partitions may be recovered from the partition. For example, the contents of the plurality of partitions may be pooled to provide these products in a bulk solution.

[00180] Outside of the partition, the gaps in the partially double-stranded nucleic acid molecule corresponding to the chromatin may be filled using via a gap filling extension process (e.g., using a DNA polymerase or reverse transcriptase). The gap filling extension process may not include strand displacement. The resultant gap-filled double-stranded nucleic acid molecule may be denatured to provide a single strand, which single strand may be subjected to conditions sufficient to perform one or more nucleic acid amplification reactions (e.g., PCR) to generate amplification products corresponding to the chromatin of the cell, cell bead, or cell nucleus. A nucleic acid amplification process may incorporate one or more additional sequences, such as one or more additional flow cell adapter sequences.

[00181] Outside of the partition, the barcoded cDNA molecule corresponding to the RNA molecule may be subjected to fragmentation, end repair, a dA tailing process, tagmentation, or a combination thereof. An additional primer sequence (e.g., a sequencing primer or portion thereof, such as an R2 sequence) may be ligated to the resultant molecule. Alternatively or in addition, a nucleic acid amplification reaction (e.g., PCR) may be performed to generate one or more amplification products corresponding to the RNA molecule or the cDNA molecule generated therefrom. A nucleic acid amplification process may incorporate one or more additional sequences, such as one or more additional flow cell adapter sequences.

[00182] FIG. 14 shows an example schematic corresponding to the preceding example. Panel 1400 shows a workflow corresponding to processing of chromatin from a cell, cell bead, or cell nucleus, and panel 1450 shows a workflow corresponding to processing of an mRNA molecule from the cell, cell bead, or cell nucleus.

[00183] As shown in panel 1400, in bulk solution, chromatin included within a cell, cell bead, or cell nucleus is processed (e.g., as described herein) to provide a template nucleic acid fragment (e.g., tagmented fragment) 1404 comprising insert sequence 1408 (e.g., region of open chromatin) and a complement thereof, transposon end sequences 1406 and complements thereof, sequencing primer or portion thereof 1402 (e.g., an R1 sequence), sequencing primer or portion thereof 1410 (e.g., an R2 sequence), and gaps 1407. Template nucleic acid fragment 1404 may then be partitioned within a partition (e.g., a droplet or well, as described herein). Within the partition, the cell, cell bead, or cell nucleus comprising template nucleic acid fragment 1404 may be lysed, permeabilized, or otherwise processed to provide access to template nucleic acid fragment 1404 (and one or more RNA molecules) therein. The partition may include a bead (e.g., gel bead) 1416 coupled to nucleic acid barcode molecules 1418a and 1418b. Nucleic acid barcode molecule 1418a may comprise a flow cell adapter sequence 1420a (e.g., a P5 sequence), a barcode sequence 1422a, and a sequencing primer or portion thereof or complement thereof 1402'. Sequences 1420a and 1422a may be hybridized to complementary sequences 1420' and 1422', respectively. Sequence 1402' may hybridize to sequence 1402 of template nucleic acid fragment 1404, or its complement, and sequence 1422' may be ligated 1412 to sequence 1402 of template nucleic acid fragment 1404. In some instances, template nucleic acid fragment 1404 may be phosphorylated using a suitable kinase enzyme (e.g., polynucleotide kinase (PNK), such as T4 PNK). In some instances, PNK and ATP may be added in bulk in the tagmentation (e.g., ATAC) reaction and/or prior to partitioning a cell, cell bead, or cell nucleus, or a plurality thereof. 15U of PNK with 1mM of ATP may be spiked into the tagmentation reaction. For example, less than 95U of PNK may be spiked into the tagmentation reaction. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the partially double-stranded nucleic acid molecule comprising nucleic acid barcode molecule 1418a attached to template nucleic acid fragment 1404 in bulk solution. In bulk solution, gaps 1407 may be filled 1424 via a gap filling extension process (e.g., using a DNA polymerase) to provide a double-stranded nucleic acid molecule. This molecule may undergo amplification (e.g., PCR) 1426 to provide a double-stranded amplification product 1428 that includes sequences of the nucleic acid barcode molecule 1418a, the original chromatin molecule, and,

optionally, an additional sequence **1430** that may be a flow cell adapter sequence (e.g., a P7 sequence). Gaps may be filled in the partition prior to bulk processing.

[00184] In parallel to the chromatin workflow of panel **1400**, an RNA molecule deriving from the same cell, cell bead, or cell nucleus may be processed. As shown in panel **1450**, RNA molecule **1458** comprising RNA sequence **1460** and polyA sequence **1462** may be contacted **1464** with primer molecule **1452** comprising polyT sequence **1454** and additional primer sequence **1456**. RNA molecule **1458** may then be reverse transcribed **1476** off of polyT sequence **1454** using a reverse transcriptase with terminal transferase activity, which reverse transcriptase may append sequence **1470** to the resultant cDNA molecule comprising cDNA sequence **1468**. Sequence **1470** may be a polyC sequence. Bead (e.g., gel bead) **1416** (e.g., the same bead described in panel **1400**) may be included within the partition and may be coupled to nucleic acid barcode molecule **1418b**. Nucleic acid barcode molecule **1418b** may comprise a flow cell adapter sequence **1420b** (e.g., a P5 sequence), a barcode sequence **1422b**, UMI sequence **1472**, and a sequence **1474** complementary to sequence **1470** (e.g., a polyG sequence). In some instances, nucleic acid barcode molecule **1418b** may comprise a sequencing primer sequence **1420b** (e.g., an R1 sequence or partial R1 sequence), a barcode sequence **1422b**, UMI sequence **1472**, and a template switching sequence **1474** complementary to sequence **1470** (e.g., a polyG sequence). Nucleic acid barcode molecule **1418b** may be used to perform template switching **1478**, which process may also result in the generation of a barcoded cDNA molecule. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the barcoded cDNA molecule in bulk solution. The barcoded cDNA molecule may undergo amplification (e.g., PCR) **1480** to provide a double-stranded amplification product **1484** that includes sequences of the nucleic acid barcode molecule **1418b**, the original RNA molecule or cDNA corresponding thereto, a flow cell adapter sequence **1486**, and an additional sequence **1488** that may comprise a sequencing primer or portion thereof (e.g., an R2 sequence) **1490**, a sample index sequence **1492**, and a flow cell adapter sequence (e.g., a P7 sequence) **1494**. The barcoded cDNA molecule may also or alternatively undergo fragmentation, end repair, dA tailing, ligation of one or more adapter sequences, and/or nucleic acid amplification.

[00185] **FIG. 15** shows another example schematic corresponding to the preceding example. Panel **1500** shows a workflow corresponding to processing of chromatin from a cell, cell bead, or cell nucleus, and panel **1550** shows a workflow corresponding to processing of an mRNA molecule from the cell, cell bead, or cell nucleus.

[00186] As shown in panel **1500**, in bulk solution, chromatin included within a cell, cell bead, or cell nucleus is processed (e.g., as described herein) to provide a template nucleic acid fragment (e.g., tagmented fragment) **1504** comprising insert sequence **1508** (e.g., region of open chromatin) and a complement thereof, transposon end sequences **1506** and complements thereof, sequencing primer or portion thereof **1502** (e.g., an R1 sequence), sequencing primer or portion thereof **1510** (e.g., an R2 sequence), and gaps **1507**. Template nucleic acid fragment **1504** may then be partitioned within a partition (e.g., a droplet or well, as described herein). Within the partition, the cell, cell bead, or cell nucleus comprising template nucleic acid fragment **1504** may be lysed, permeabilized, or otherwise processed to provide access to template nucleic acid fragment **1504** (and one or more RNA molecules) therein. The partition may include a bead (e.g., gel bead) **1516** coupled to nucleic acid barcode molecules **1518a** and **1518b**. Nucleic acid barcode molecule **1518a** may comprise a flow cell adapter sequence **1520a** (e.g., a P5 sequence), a barcode sequence **1522a**, and a sequencing primer or portion thereof or complement thereof **1502'**. Sequences **1520a** and **1522a** may be hybridized to complementary sequences **1520'** and **1522'**, respectively. Sequence **1502'** may hybridize to sequence **1502** of template nucleic acid fragment **1504**, or its complement, and sequence **1522'** may be ligated **1512** to sequence **1502** of template nucleic acid fragment **1504**. In some instances, template nucleic acid fragment **1504** may be phosphorylated using a suitable kinase enzyme (e.g., polynucleotide kinase (PNK), such as T4 PNK). PNK and ATP may be added in bulk in the tagmentation (e.g., ATAC) reaction and/or prior to partitioning a cell, cell bead, or cell nucleus, or plurality thereof. 15U of PNK with 1mM of ATP may be spiked into the tagmentation reaction. For example, less than 95U of PNK may be spiked into the tagmentation reaction. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the partially double-stranded nucleic acid molecule comprising nucleic acid barcode molecule **1518a** attached to template nucleic acid fragment **1504** in bulk solution. In bulk solution, gaps **1507** may be filled **1524** via a gap filling extension process (e.g., using a DNA polymerase) to provide a double-stranded nucleic acid molecule. This molecule may undergo amplification (e.g., PCR) **1526** to provide a double-stranded amplification product **1528** that includes sequences of the nucleic acid barcode molecule **1518a**, the original chromatin molecule, and, optionally, an additional sequence **1530** that may be a flow cell adapter sequence (e.g., a P7 sequence). Gaps may be filled in the partition prior to bulk processing.

[00187] In parallel to the chromatin workflow of panel **1500**, an RNA molecule deriving from the same cell, cell bead, or cell nucleus may be processed. As shown in panel **1550**, RNA molecule **1558** comprising RNA sequence **1560** and polyA sequence **1562** and bead (e.g., gel

bead) **1516** may be provided within a partition. Bead (e.g., gel bead) **1516** (e.g., the same bead described in panel **1500**) may be included within the partition and may be coupled to nucleic acid barcode molecule **1518b**. Nucleic acid barcode molecule **1518b** may comprise a flow cell adapter sequence **1568** (e.g., a P5 sequence), a barcode sequence **1522b** (e.g., the same barcode sequence as barcode sequence **1522a**), UMI sequence **1566**, and a polyT sequence **1564** complementary to polyA sequence **1562**. In some instances, nucleic acid barcode molecule **1518b** may comprise a sequencing primer sequence **1568** (e.g., an R1 sequence or partial R1 sequence), a barcode sequence **1522b** (e.g., the same barcode sequence as barcode sequence **1522a**), UMI sequence **1566**, and a polyT sequence **1564** complementary to polyA sequence **1562**. PolyT sequence **1564** may hybridize to polyA sequence **1562** of RNA molecule **1558**. RNA molecule **1558** may be reverse transcribed **1570** off of polyT sequence **1564** to provide an cDNA molecule comprising cDNA sequence **1572**. The reverse transcription process may use a reverse transcriptase with terminal transferase activity, which reverse transcriptase may append sequence **1574** to the resultant cDNA molecule comprising cDNA sequence **1572**. Sequence **1574** may be a polyC sequence. A template switch oligonucleotide **1578** comprising a primer sequence **1580** and a sequence complementary to sequence **1574** (e.g., a polyG sequence) may hybridize to the cDNA molecule and facilitate a template switching reaction onto template switch oligonucleotide **1578**. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the cDNA molecule in bulk solution. The cDNA molecule may undergo amplification (e.g., PCR) **1584**. Additional amplification (e.g., PCR) **1586** may be performed to provide a double-stranded amplification product **1588** that includes sequences of the nucleic acid barcode molecule **1518b**, the original RNA molecule or cDNA corresponding thereto, a flow cell adapter sequence **1598** (e.g., a P7 sequence), and an additional sequence **1590** that may comprise a sequencing primer or portion thereof (e.g., an R2 sequence) **1596**, a sample index sequence **1594**, and a flow cell adapter sequence (e.g., a P5 sequence) **1592**. The barcoded cDNA molecule may also or alternatively undergo fragmentation, end repair, dA tailing, ligation of one or more adapter sequences, and/or nucleic acid amplification.

[00188] In another example, a cell, cell bead, or cell nucleus comprising chromatin and one or more RNA molecules is provided. The chromatin in the cell, cell bead, or cell nucleus may be processed to provide a first template nucleic acid fragment derived from the chromatin (e.g., a tagged fragment, as described herein). The chromatin may be processed in bulk solution. An RNA molecule may be processed to provide a second template nucleic acid fragment derived from the RNA molecule (e.g., an additional nucleic acid fragment, as described herein). The

RNA molecule may be processed within a partition. The second template nucleic acid fragment derived from the RNA molecule may be processed according to the preceding examples. The configuration of the first template nucleic acid fragment may be at least partially dependent on the structure of the transposase-nucleic acid complex used to generate the first template nucleic acid fragment. For example, a transposase-nucleic acid complex such as that shown in **FIG. 9** may be used to prepare the first template nucleic acid fragment. Relative to the preceding examples, the polarities of the transposase-nucleic acid may be reversed such that sequencing primers (e.g., R1 and R2 sequencing primers) are not directly linked to the chromatin (see, e.g., **FIG. 17**). The first template nucleic acid fragment may be at least partially double-stranded. The first template nucleic acid fragment may comprise a double-stranded region comprising sequences of chromatin of the cell, cell bead, or cell nucleus. A first end of a first strand of the double-stranded region may be linked to a first transposon end sequence (e.g., mosaic end sequence). A first end of the second strand of the double-stranded region, which end is opposite the first end of the first strand, may be linked to a second transposon end sequence (e.g., mosaic end sequence). The second transposon end sequence may be the same as or different from the first transposon end sequence. The first transposon end sequence may be hybridized to a first complementary sequence (e.g., mosaic end reverse complement sequence), which first complementary sequence may not be linked to a second end of the second strand of the double-stranded region of the first template nucleic acid fragment. The first complementary sequence may be linked to a first sequencing primer or portion thereof. Similarly, the second transposon end sequence may be hybridized to a second complementary sequence (e.g., mosaic end reverse complement sequence), which second complementary sequence may not be linked to a second end of the first strand of the double-stranded region of the first template nucleic acid fragment. The second complementary sequence may be linked to a second sequencing primer or portion thereof. In other words, the first template nucleic acid fragment may comprise one or more gaps. In some cases, the one or more gaps may be approximately 9 bp in length each. For example, one or more gaps may be at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more bp in length. For example, one or more gaps may be at most about 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 bp in length. The first sequencing primer or portion thereof may be the same as or different from the second sequencing primer or portion thereof. In some cases, the first sequencing primer or portion thereof may be an R1 sequence or portion thereof, and the second sequencing primer or portion thereof may be an R2 sequence or portion thereof.

[00189] The cell, cell bead, or cell nucleus comprising the first template nucleic acid fragment (e.g., tagmented fragment) may be co-partitioned with one or more reagents into a partition of a

plurality of partitions (e.g., as described herein). The partition may be, for example, a droplet or well. The partition may comprise one or more beads (e.g., as described herein). A bead (e.g., gel bead) of the one or more beads may comprise a first plurality of nucleic acid barcode molecules. A nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules may comprise a flow cell adapter sequence (e.g., P5 sequence), a barcode sequence, and a sequencing primer or portion thereof (e.g., R1 sequence or portion thereof, or a complement thereof). The sequencing primer or portion thereof may be complementary to a sequence of the first template nucleic acid fragment. The flow cell adapter sequence and/or barcode sequence may be hybridized to their complementary sequences. The same bead or another bead may comprise a second plurality of nucleic acid barcode molecules. A nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules may comprise a sequencing primer or portion thereof (e.g., an R1 sequence or portion thereof, or complement thereof), a barcode sequence, a unique molecular identifier sequence, and a capture sequence.

[00190] Within the partition, the RNA molecule may be processed to provide the second template nucleic acid fragment (e.g., as described herein). For example, the RNA molecule (e.g., mRNA molecule) may be contacted with a primer molecule comprising a first primer sequence (e.g., a polyT sequence) and an additional primer sequence.

[00191] Within the partition, the cell, cell bead, or cell nucleus may be lysed or permeabilized to provide access to the first and/or second template nucleic acid fragments therein (e.g., as described herein). The second template nucleic acid fragment may be generated after the cell, cell bead, or cell nucleus is lysed or permeabilized.

[00192] The first and second template nucleic acid fragments may undergo processing within the partition. Within the partition, a sequencing primer or portion thereof of the first template nucleic acid fragment corresponding to the chromatin of the cell, cell bead, or cell nucleus may hybridize to a sequencing primer or portion thereof of the nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules. The sequencing primer or portion thereof of the nucleic acid barcode molecule may then be ligated (e.g., using a ligase) to a transposon end sequence of the first template nucleic acid fragment, or a complement thereof to provide a partially double-stranded nucleic acid molecule corresponding to the chromatin of the cell, cell bead, or cell nucleus. The second template nucleic acid fragment corresponding to the RNA molecule may be reverse transcribed using a reverse transcriptase with terminal transferase activity, which reverse transcriptase may append a sequence (e.g., a polyC sequence) to the cDNA strand of the resultant cDNA molecule. The cDNA molecule may then be contacted with a nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules that

may be a template switch oligonucleotide. The nucleic acid barcode molecule may comprise a sequencing primer or portion thereof (e.g., an R1 sequence or portion thereof, or complement thereof), a barcode sequence, a unique molecular identifier sequence, and a capture sequence. The capture sequence may be a sequence that is complementary to the sequence appended to the cDNA strand (e.g., a polyG sequence). Template switching and barcoding may then take place to provide a barcoded cDNA molecule.

[00193] The partially double-stranded molecule corresponding to the chromatin of the cell, cell bead, or cell nucleus and the barcoded cDNA molecule corresponding to the RNA molecule (e.g., prepared as described above) of the cell, cell bead, or cell nucleus included within the partition of the plurality of partitions may be recovered from the partition. For example, the contents of the plurality of partitions may be pooled to provide the linear amplification products in a bulk solution.

[00194] Outside of the partition, the gaps in the partially double-stranded nucleic acid molecule corresponding to the chromatin may be filled using via a gap filling extension process (e.g., using a DNA polymerase). Gaps may be filled in the partition prior to bulk processing. The resultant gap-filled double-stranded nucleic acid molecule may be denatured to provide a single strand, which single strand may be subjected to conditions sufficient to perform one or more nucleic acid amplification reactions (e.g., PCR) to generate amplification products corresponding to the chromatin of the cell, cell bead, or cell nucleus. A nucleic acid amplification process may incorporate one or more additional sequences, such as one or more additional flow cell adapter sequences. The barcoded cDNA molecule corresponding to the RNA molecule may also be processed and amplified according to the preceding examples.

[00195] **FIG. 16** shows an example schematic corresponding to the preceding example. Panel **1600** shows a workflow corresponding to processing of chromatin from a cell, cell bead, or cell nucleus, and panel **1650** shows a workflow corresponding to processing of an mRNA molecule from the cell, cell bead, or cell nucleus.

[00196] As shown in panel **1600**, in bulk solution, chromatin included within a cell, cell bead, or cell nucleus is processed (e.g., as described herein) to provide a template nucleic acid fragment (e.g., tagmented fragment) **1604** comprising insert sequence **1608** (e.g., region of open chromatin) and a complement thereof, transposon end sequences **1606** and complements thereof, sequencing primer or portion thereof **1602** (e.g., an R1 sequence), sequencing primer or portion thereof **1610** (e.g., an R2 sequence), and gaps **1607**. Template nucleic acid fragment **1604** may then be partitioned within a partition (e.g., a droplet or well, as described herein). Within the partition, the cell, cell bead, or cell nucleus comprising template nucleic acid fragment **1604** may

be lysed, permeabilized, or otherwise processed to provide access to template nucleic acid fragment **1604** (and one or more RNA molecules) therein. The partition may include a bead (e.g., gel bead) **1616** coupled to nucleic acid barcode molecules **1618a** and **1618b**. Nucleic acid barcode molecule **1618a** may comprise a flow cell adapter sequence **1620a** (e.g., a P5 sequence), a barcode sequence **1622a**, and a sequencing primer or portion thereof or complement thereof **1602'**. Sequence **1602'** may hybridize to sequence **1602** of template nucleic acid fragment **1604**, or its complement. Sequence **1602'** may then be ligated **1612** to a transposon end sequence **1606** of template nucleic acid fragment **1604**. In some instances, **1604** may be phosphorylated using a suitable kinase enzyme (e.g., polynucleotide kinase (PNK), such as T4 PNK). In some instances, PNK and ATP may be added in bulk in the tagmentation (e.g., ATAC) reaction and/or prior to partitioning a cell, cell bead, or cell nucleus, or plurality thereof. 15U of PNK with 1mM of ATP may be spiked into the tagmentation reaction. For example, less than 95U of PNK may be spiked into the tagmentation reaction. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the partially double-stranded nucleic acid molecule comprising nucleic acid barcode molecule **1618a** attached to template nucleic acid fragment **1604** in bulk solution. In bulk solution, gaps **1607** may be filled **1614** via a gap filling extension process (e.g., using a DNA polymerase) and the molecule extended from sequence **1602** to provide a double-stranded nucleic acid molecule. This molecule may undergo amplification (e.g., PCR) **1624** to provide a double-stranded amplification product **1626** that includes sequences of the nucleic acid barcode molecule **1618a**, the original chromatin molecule, and, optionally, an additional sequence **1628** that may be a flow cell adapter sequence (e.g., a P7 sequence). Gaps may be filled in the partition prior to bulk processing.

[00197] In parallel to the chromatin workflow of panel **1600**, an RNA molecule deriving from the same cell, cell bead, or cell nucleus may be processed. As shown in panel **1650**, RNA molecule **1658** comprising RNA sequence **1660** and polyA sequence **1662** may be contacted **1664** with primer molecule **1652** comprising polyT sequence **1654** and additional primer sequence **1656**. RNA molecule **1658** may then be reverse transcribed **1676** off of polyT sequence **1654** using a reverse transcriptase with terminal transferase activity, which reverse transcriptase may append sequence **1670** to the resultant cDNA molecule comprising cDNA sequence **1668**. Sequence **1670** may be a polyC sequence. Bead (e.g., gel bead) **1616** (e.g., the same bead described in panel **1600**) may be included within the partition and may be coupled to nucleic acid barcode molecule **1618b**. Nucleic acid barcode molecule **1618b** may comprise a flow cell adapter sequence **1620b** (e.g., a P5 sequence), a barcode sequence **1622b**, UMI sequence **1672**, and a sequence **1674** complementary to sequence **1670** (e.g., a polyG sequence).

In some instances, nucleic acid barcode molecule **1618b** may comprise a sequencing primer sequence **1620b** (e.g., an R1 sequence or partial R1 sequence), a barcode sequence **1622b**, UMI sequence **1672**, and a template switching sequence **1674** complementary to sequence **1670** (e.g., a polyG sequence). Nucleic acid barcode molecule **1618b** may be used to perform template switching **1678**, which process may also result in the generation of a barcoded cDNA molecule. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the barcoded cDNA molecule in bulk solution. The barcoded cDNA molecule may undergo amplification (e.g., PCR) **1680** to provide a double-stranded amplification product **1684** that includes sequences of the nucleic acid barcode molecule **1618b**, the original RNA molecule or cDNA corresponding thereto, a flow cell adapter sequence **1686**, and an additional sequence **1688** that may comprise a sequencing primer or portion thereof (e.g., an R2 sequence) **1690**, a sample index sequence **1692**, and a flow cell adapter sequence (e.g., a P7 sequence) **1694**. The barcoded cDNA molecule may also or alternatively undergo fragmentation, end repair, dA tailing, ligation of one or more adapter sequences, and/or nucleic acid amplification.

[00198] FIG. 17 shows another example schematic corresponding to the preceding example. Panel **1700** shows a workflow corresponding to processing of chromatin from a cell, cell bead, or cell nucleus, and panel **1750** shows a workflow corresponding to processing of an mRNA molecule from the cell, cell bead, or cell nucleus.

[00199] As shown in panel **1700**, in bulk solution, chromatin included within a cell, cell bead, or cell nucleus is processed (e.g., as described herein) to provide a template nucleic acid fragment (e.g., tagged fragment) **1704** comprising insert sequence **1708** (e.g., region of open chromatin) and a complement thereof, transposon end sequences **1706** and complements thereof, sequencing primer or portion thereof **1702** (e.g., an R1 sequence), sequencing primer or portion thereof **1710** (e.g., an R2 sequence), and gaps **1707**. Template nucleic acid fragment **1704** may then be partitioned within a partition (e.g., a droplet or well, as described herein). Within the partition, the cell, cell bead, or cell nucleus comprising template nucleic acid fragment **1704** may be lysed, permeabilized, or otherwise processed to provide access to template nucleic acid fragment **1704** (and one or more RNA molecules) therein. The partition may include a bead (e.g., gel bead) **1716** coupled to nucleic acid barcode molecules **1718a** and **1718b**. Nucleic acid barcode molecule **1718a** may comprise a flow cell adapter sequence **1720a** (e.g., a P5 sequence), a barcode sequence **1722a**, and a sequencing primer or portion thereof or complement thereof **1702'**. Sequence **1702'** may hybridize to sequence **1702** of template nucleic acid fragment **1704**, or its complement. Sequence **1702'** may then be ligated **1712** to a transposon end sequence **1706**

of template nucleic acid fragment **1704**. In some instances, **1704** may be phosphorylated using a suitable kinase enzyme (e.g., polynucleotide kinase (PNK), such as T4 PNK). In some instances, PNK and ATP may be added in bulk in the tagmentation reaction (e.g., ATAC) and/or prior to partitioning a cell, cell bead or cell nucleus, or plurality thereof. 15U of PNK with 1mM of ATP may be spiked into the tagmentation reaction. For example, less than 95U of PNK may be spiked into the tagmentation reaction. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the partially double-stranded nucleic acid molecule comprising nucleic acid barcode molecule **1718a** attached to template nucleic acid fragment **1704** in bulk solution. In bulk solution, gaps **1707** may be filled **1714** via a gap filling extension process (e.g., using a DNA polymerase) and the molecule extended from sequence **1702** to provide a double-stranded nucleic acid molecule. This molecule may undergo amplification (e.g., PCR) **1724** to provide a double-stranded amplification product **1726** that includes sequences of the nucleic acid barcode molecule **1718a**, the original chromatin molecule, and, optionally, an additional sequence **1728** that may be a flow cell adapter sequence (e.g., a P7 sequence). Gaps may be filled in the partition prior to bulk processing.

[00200] In parallel to the chromatin workflow of panel **1700**, an RNA molecule deriving from the same cell, cell bead, or cell nucleus may be processed. As shown in panel **1750**, RNA molecule **1758** comprising RNA sequence **1760** and polyA sequence **1762** and bead (e.g., gel bead) **1716** may be provided within a partition. Bead (e.g., gel bead) **1716** (e.g., the same bead described in panel **1700**) may be included within the partition and may be coupled to nucleic acid barcode molecule **1718b**. Nucleic acid barcode molecule **1718b** may comprise a flow cell adapter sequence **1768** (e.g., a P5 sequence), a barcode sequence **1722b** (e.g., the same barcode sequence as barcode sequence **1722a**), UMI sequence **1766**, and a polyT sequence **1764** complementary to polyA sequence **1762**. In some instances, nucleic acid barcode molecule **1718b** may comprise a sequencing primer sequence **1768** (e.g., an R1 sequence or partial R1 sequence), a barcode sequence **1722b** (e.g., the same barcode sequence as barcode sequence **1722a**), UMI sequence **1766**, and a polyT sequence **1764** complementary to polyA sequence **1762**. PolyT sequence **1764** may hybridize to polyA sequence **1762** of RNA molecule **1758**. RNA molecule **1758** may be reverse transcribed **1770** off of polyT sequence **1764** to provide an cDNA molecule comprising cDNA sequence **1772**. The reverse transcription process may use a reverse transcriptase with terminal transferase activity, which reverse transcriptase may append sequence **1774** to the resultant cDNA molecule comprising cDNA sequence **1772**. Sequence **1774** may be a polyC sequence. A template switch oligonucleotide **1778** comprising a primer sequence **1780** and a sequence complementary to sequence **1774** (e.g., a polyG sequence) may

hybridize to the cDNA molecule. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the cDNA molecule in bulk solution. The cDNA molecule may undergo amplification (e.g., PCR) **1784**. Additional amplification (e.g., PCR) **1786** may be performed to provide a double-stranded amplification product **1788** that includes sequences of the nucleic acid barcode molecule **1718b**, the original RNA molecule or cDNA corresponding thereto, a flow cell adapter sequence **1798** (e.g., a P7 sequence), and an additional sequence **1790** that may comprise a sequencing primer or portion thereof (e.g., an R2 sequence) **1796**, a sample index sequence **1794**, and a flow cell adapter sequence (e.g., a P5 sequence) **1792**. The barcoded cDNA molecule may also or alternatively undergo fragmentation, end repair, dA tailing, ligation of one or more adapter sequences, and/or nucleic acid amplification.

[00201] In another example, a cell, cell bead, or cell nucleus comprising chromatin and one or more RNA molecules is provided. The chromatin in the cell, cell bead, or cell nucleus may be processed to provide a first template nucleic acid fragment derived from the chromatin (e.g., a tagged fragment, as described herein). The chromatin may be processed in bulk solution. An RNA molecule may be processed to provide a second template nucleic acid fragment derived from an RNA molecule (e.g., as described herein). The RNA molecule may be processed within a partition. The configuration of the first template nucleic acid fragment may be at least partially dependent on the structure of the transposase-nucleic acid complex used to generate the first template nucleic acid fragment. For example, a transposase-nucleic acid complex such as that shown in **FIG. 9** may be used to prepare the first template nucleic acid fragment. The first template nucleic acid fragment may be at least partially double-stranded. The first template nucleic acid fragment may comprise a double-stranded region comprising sequences of chromatin of the cell, cell bead, or cell nucleus. A first end of a first strand of the double-stranded region may be linked to a first transposon end sequence (e.g., mosaic end sequence), which first transposon end sequence may be linked to a first sequencing primer or portion thereof. A first end of the second strand of the double-stranded region, which end is opposite the first end of the first strand, may be linked to a second transposon end sequence (e.g., mosaic end sequence), which second transposon end sequence may be linked to a second sequencing primer or portion thereof. The second transposon end sequence may be the same as or different from the first transposon end sequence. The first sequencing primer or portion thereof may be the same as or different from the second sequencing primer or portion thereof. In some cases, the first sequencing primer or portion thereof may be an R1 sequence or portion thereof, and the second sequencing primer or portion thereof may be an R2 sequence or portion thereof. The first

transposon end sequence may be hybridized to a first complementary sequence (e.g., mosaic end reverse complement sequence), which first complementary sequence may not be linked to a second end of the second strand of the double-stranded region of the first template nucleic acid fragment. Similarly, the second transposon end sequence may be hybridized to a second complementary sequence (e.g., mosaic end reverse complement sequence), which second complementary sequence may not be linked to a second end of the first strand of the double-stranded region of the first template nucleic acid fragment. In other words, the first template nucleic acid fragment may comprise one or more gaps. In some cases, the one or more gaps may be approximately 9 bp in length each. For example, one or more gaps may be at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more bp in length. For example, one or more gaps may be at most about 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 bp in length. The second template nucleic acid fragment (e.g., an additional template nucleic acid fragment) may comprise a sequence of an RNA molecule of the cell, cell bead, or cell nucleus and a sequence hybridized to a primer molecule (e.g., a capture nucleic acid molecule). For example, the second template nucleic acid fragment may comprise a sequence of an RNA molecule of the cell, cell bead, or cell nucleus and a polyA sequence hybridized to a polyT sequence of a primer molecule. The primer molecule may also comprise an additional primer sequence.

[00202] The cell, cell bead, or cell nucleus comprising the first template nucleic acid fragment (e.g., tagged fragment) may be co-partitioned with one or more reagents into a partition of a plurality of partitions (e.g., as described herein). The partition may be, for example, a droplet or well. The partition may comprise one or more beads (e.g., as described herein). A bead (e.g., gel bead) of the one or more beads may comprise a first plurality of nucleic acid barcode molecules. A nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules may comprise a flow cell adapter sequence (e.g., P5 sequence), a barcode sequence, and an overhang sequence. The partition may also comprise a splint sequence comprising a sequence complementary to the overhang sequence and a sequencing primer or portion thereof that may be complementary to a sequence of the first template nucleic acid fragment. A bead of the one or more beads may also comprise a second plurality of nucleic acid barcode molecules. A nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules may comprise a flow cell adapter sequence (e.g., P5 sequence), a barcode sequence, a sequencing primer or portion thereof (e.g., R1 sequence or portion thereof, or a complement thereof), a UMI sequence, and a capture sequence (e.g., a polyG sequence or a polydT sequence). In some cases, the first plurality of nucleic acid barcode molecules and the second plurality of nucleic acid

barcode molecules may be coupled to the same bead, and the partition may comprise a single bead.

[00203] Within the partition, the RNA molecule may be processed to provide the second template nucleic acid fragment (e.g., as described herein).

[00204] Within the partition, the cell, cell bead, or cell nucleus may be lysed or permeabilized to provide access to the first and/or second template nucleic acid fragments therein (e.g., as described herein). The second template nucleic acid fragment may be generated after the cell, cell bead, or cell nucleus is lysed or permeabilized.

[00205] The first and second template nucleic acid fragments may undergo processing within the partition. Within the partition, a sequencing primer or portion thereof of the first template nucleic acid fragment corresponding to the chromatin of the cell, cell bead, or cell nucleus may hybridize to a complementary sequence of the sequencing primer or portion thereof in the splint sequence. The splint sequence may also hybridize to the overhang sequence of the nucleic acid barcode molecule of the first plurality of nucleic acid barcode molecules. The overhang sequence of the nucleic acid barcode molecule may then be ligated (e.g., using a ligase) to a sequencing primer or portion thereof of the first template nucleic acid fragment. The resultant partially double-stranded nucleic acid molecule may comprise the barcode sequence as well as one or more gaps.

[00206] Within the partition, the second template nucleic acid fragment derived from the RNA molecule of the cell, cell bead, or cell nucleus may be reverse transcribed (e.g., using a reverse transcriptase) to provide a cDNA strand. The reverse transcription process may append a sequence to an end of a strand of the resultant double-stranded nucleic acid molecule comprising the RNA strand and the cDNA strand, such as a polyC sequence. The capture sequence of the nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules may hybridize to the appended sequence (e.g., polyC sequence) of the double-stranded nucleic acid molecule and a template switching process may take place to provide a second double-stranded nucleic acid molecule. The sequence of the nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules may be considered a template switching oligonucleotide. The template switch process may result in a barcoded cDNA molecule. The barcoded cDNA molecule may comprise the sequencing primer or portion thereof, or complement thereof, of the nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules; the barcode sequence, or complement thereof, of the nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules; the UMI sequence, or complement thereof, of the nucleic acid barcode molecule of the second plurality of nucleic acid

barcode molecules; the capture sequence, or complement thereof, of the nucleic acid barcode molecule of the second plurality of nucleic acid barcode molecules; the poly(C) or poly(G) sequence; the sequence corresponding to the RNA molecule of the cell, cell bead, or cell nucleus, or complement thereof; and sequences of the capture nucleic acid molecule, or complements thereof.

[00207] The partially double-stranded nucleic acid molecule corresponding to the chromatin of the cell, cell bead, or cell nucleus and the barcoded cDNA molecule corresponding to the RNA molecule of the cell, cell bead, or cell nucleus included within the partition of the plurality of partitions may be recovered from the partition. For example, the contents of the plurality of partitions may be pooled to provide the partially double-stranded nucleic acid molecule and the barcoded cDNA molecule in a bulk solution.

[00208] Outside of the partition, the gaps in the partially double-stranded nucleic acid molecule corresponding to the chromatin may be filled using via a gap filling extension process (e.g., using a DNA polymerase or reverse transcriptase). The DNA polymerase may lack strand displacement activity. The resultant gap-filled double-stranded nucleic acid molecule may be denatured to provide a single strand, which single strand may be subjected to conditions sufficient to perform one or more nucleic acid amplification reactions (e.g., PCR) to generate amplification products corresponding to the chromatin of the cell, cell bead, or cell nucleus. A nucleic acid amplification process may incorporate one or more additional sequences, such as one or more additional flow cell adapter sequences.

[00209] Outside of the partition, the barcoded cDNA molecule corresponding to the RNA molecule may be subjected to fragmentation, end repair, a dA tailing process, tagmentation, or a combination thereof. An additional primer sequence (e.g., a sequencing primer or portion thereof, such as an R2 sequence) may then be ligated to the resultant molecule. A nucleic acid amplification reaction (e.g., PCR) may then be performed to generate one or more amplification products corresponding to the RNA molecule. A nucleic acid amplification process may incorporate one or more additional sequences, such as one or more additional flow cell adapter sequences.

[00210] **FIG. 18** shows an example schematic corresponding to the preceding example. Panel **1800** shows a workflow corresponding to processing of chromatin from a cell, cell bead, or cell nucleus, and panel **1850** shows a workflow corresponding to processing of an mRNA molecule from the cell, cell bead, or cell nucleus.

[00211] As shown in panel **1800**, in bulk solution, chromatin included within a cell, cell bead, or cell nucleus is processed (e.g., as described herein) to provide a template nucleic acid

fragment (e.g., tagmented fragment) **1804** comprising insert sequence **1808** (e.g., region of open chromatin) and a complement thereof, transposon end sequences **1806** and complements thereof, sequencing primer or portion thereof **1802** (e.g., an R1 sequence), sequencing primer or portion thereof **1810** (e.g., an R2 sequence), and gaps **1807**. Template nucleic acid fragment **1804** may then be partitioned within a partition (e.g., a droplet or well, as described herein). Within the partition, the cell, cell bead, or cell nucleus comprising template nucleic acid fragment **1804** may be lysed, permeabilized, or otherwise processed to provide access to template nucleic acid fragment **1804** (and one or more RNA molecules) therein. The partition may comprise splint sequence **1812**, which splint sequence may comprise a first sequence **1802'** that is complementary to sequencing primer or portion thereof **1802** and a second sequence **1824**. Sequence **1824** may comprise a blocking group (e.g., a 3' blocking group), which blocking group may prevent extension by reverse transcription. The partition may also include a bead (e.g., gel bead) **1816** coupled to nucleic acid barcode molecules **1818a** and **1812b**. Nucleic acid barcode molecule **1818a** may comprise a flow cell adapter sequence **1820a** (e.g., a P5 sequence), a barcode sequence **1822a**, and an overhang sequence **1824'** that is complementary to sequence **1824** of the splint sequence. Sequence **1824** may hybridize to sequence **1824'** to provide a partially double-stranded nucleic acid molecule comprising the sequences of nucleic acid barcode molecule **1818a** and the template nucleic acid fragment **1804**. Sequence **1824'** of nucleic acid barcode molecule **1818a** may be ligated (e.g., using a ligase) **1826** to sequence **1802** of template nucleic acid fragment **1804**. In some instances, **1804** may be phosphorylated using a suitable kinase enzyme (e.g., polynucleotide kinase (PNK), such as T4 PNK). In some instances, PNK and ATP may be added in bulk in the tagmentation reaction (e.g., ATAC) and/or prior to partitioning a cell, cell bead, or cell nucleus, or plurality thereof. 15U of PNK with 1mM of ATP may be spiked into the tagmentation reaction. For example, less than 95U of PNK may be spiked into the tagmentation reaction. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the partially double-stranded nucleic acid molecule comprising nucleic acid barcode molecule **1818a** attached to template nucleic acid fragment **1804** in bulk solution. In bulk solution, gaps **1807** may be filled **1828** via a gap filling extension process (e.g., using a DNA polymerase) to provide a double-stranded nucleic acid molecule. This molecule may undergo amplification (e.g., PCR) **1830** to provide a double-stranded amplification product **1832** that includes sequences of the nucleic acid barcode molecule **1818a**, the original chromatin molecule, and, optionally, an additional sequence **1834** that may be a flow cell adapter sequence (e.g., a P7 sequence). Gaps may be filled in the partition prior to bulk processing.

[00212] In parallel to the chromatin workflow of panel **1800**, an RNA molecule deriving from the same cell, cell bead, or cell nucleus may be processed. As shown in panel **1850**, RNA molecule **1858** comprising RNA sequence **1860** and polyA sequence **1862** may be contacted **1864** with primer molecule **1852** comprising polyT sequence **1854** and additional primer sequence **1856**. RNA molecule **1858** may then be reverse transcribed **1876** off of polyT sequence **1854** using a reverse transcriptase with terminal transferase activity, which reverse transcriptase may append sequence **1870** to the resultant cDNA molecule comprising cDNA sequence **1868**. Sequence **1870** may be a polyC sequence. Bead (e.g., gel bead) **1816** (e.g., the same bead described in panel **1800**) may be included within the partition and may be coupled to nucleic acid barcode molecule **1818b**. Nucleic acid barcode molecule **1818b** may comprise a flow cell adapter sequence **1820b** (e.g., a P5 sequence), a barcode sequence **1822b**, UMI sequence **1872**, and a sequence **1874** complementary to sequence **1870** (e.g., a polyG sequence). In some instances, nucleic acid barcode molecule **1818b** may comprise a sequencing primer sequence **1820b** (e.g., an R1 sequence or partial R1 sequence), a barcode sequence **1822b**, UMI sequence **1872**, and a sequence **1874** complementary to sequence **1870** (e.g., a polyG sequence). Nucleic acid barcode molecule **1818b** may be used to perform template switching **1878**, which process may also result in the generation of a barcoded cDNA molecule. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the barcoded cDNA molecule in bulk solution. The barcoded cDNA molecule may undergo amplification (e.g., PCR) **1880** to provide a double-stranded amplification product **1884** that includes sequences of the nucleic acid barcode molecule **1818b**, the original RNA molecule or cDNA corresponding thereto, a flow cell adapter sequence **1886**, and an additional sequence **1888** that may comprise a sequencing primer or portion thereof (e.g., an R2 sequence) **1890**, a sample index sequence **1892**, and a flow cell adapter sequence (e.g., a P7 sequence) **1894**. The barcoded cDNA molecule may also or alternatively undergo fragmentation, end repair, dA tailing, ligation of one or more adapter sequences, and/or nucleic acid amplification.

[00213] **FIG. 19** shows an example schematic corresponding to the preceding example. Panel **1900** shows a workflow corresponding to processing of chromatin from a cell, cell bead, or cell nucleus, and panel **1950** shows a workflow corresponding to processing of an mRNA molecule from the cell, cell bead, or cell nucleus.

[00214] As shown in panel **1900**, in bulk solution, chromatin included within a cell, cell bead, or cell nucleus is processed (e.g., as described herein) to provide a template nucleic acid fragment (e.g., tagmented fragment) **1904** comprising insert sequence **1908** (e.g., region of open chromatin) and a complement thereof, transposon end sequences **1906** and complements thereof,

sequencing primer or portion thereof **1902** (e.g., an R1 sequence), sequencing primer or portion thereof **1910** (e.g., an R2 sequence), and gaps **1907**. Template nucleic acid fragment **1904** may then be partitioned within a partition (e.g., a droplet or well, as described herein). Within the partition, the cell, cell bead, or cell nucleus comprising template nucleic acid fragment **1904** may be lysed, permeabilized, or otherwise processed to provide access to template nucleic acid fragment **1904** (and one or more RNA molecules) therein. The partition may comprise splint sequence **1912**, which splint sequence may comprise a first sequence **1902'** that is complementary to sequencing primer or portion thereof **1902** and a second sequence **1924**. Sequence **1924** may comprise a blocking group (e.g., a 3' blocking group), which blocking group may prevent extension by reverse transcription. The partition may also include a bead (e.g., gel bead) **1916** coupled to nucleic acid barcode molecules **1918a** and **1918b**. Nucleic acid barcode molecule **1918a** may comprise a flow cell adapter sequence **1920a** (e.g., a P5 sequence), a barcode sequence **1922a**, and an overhang sequence **1924'** that is complementary to sequence **1924** of the splint sequence. Sequence **1924** may hybridize to sequence **1924'** to provide a partially double-stranded nucleic acid molecule comprising the sequences of nucleic acid barcode molecule **1918a** and the template nucleic acid fragment **1904**. Sequence **1924'** of nucleic acid barcode molecule **1918a** may be ligated (e.g., using a ligase) **1926** to sequence **1902** of template nucleic acid fragment **1904**. In some instances, **1904** may be phosphorylated using a suitable kinase enzyme (e.g., polynucleotide kinase (PNK), such as T4 PNK). In some instances, PNK and ATP may be added in bulk in the tagmentation reaction (e.g., ATAC) and/or prior to partitioning the cells, cell beads, or cell nuclei. 15U of PNK with 1mM of ATP may be spiked into the tagmentation reaction. For example, less than 95U of PNK may be spiked into the tagmentation reaction. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the partially double-stranded nucleic acid molecule comprising nucleic acid barcode molecule **1918a** attached to template nucleic acid fragment **1904** in bulk solution. In bulk solution, gaps **1907** may be filled **1928** via a gap filling extension process (e.g., using a DNA polymerase) to provide a double-stranded nucleic acid molecule. This molecule may undergo amplification (e.g., PCR) **1930** to provide a double-stranded amplification product **1932** that includes sequences of the nucleic acid barcode molecule **1918a**, the original chromatin molecule, and, optionally, an additional sequence **1934** that may be a flow cell adapter sequence (e.g., a P7 sequence). Gaps may be filled in the partition prior to bulk processing.

[00215] In parallel to the chromatin workflow of panel **1900**, an RNA molecule deriving from the same cell, cell bead, or cell nucleus may be processed. As shown in panel **1950**, RNA

molecule **1958** comprising RNA sequence **1960** and polyA sequence **1962** and bead (e.g., gel bead) **1916** may be provided within a partition. Bead (e.g., gel bead **1916** (e.g., the same bead described in panel **1900**)) may be included within the partition and may be coupled to nucleic acid barcode molecule **1918b**. Nucleic acid barcode molecule **1918b** may comprise a flow cell adapter sequence **1968** (e.g., a P5 sequence), a barcode sequence **1922b** (e.g., the same barcode sequence as barcode sequence **1922a**), UMI sequence **1966**, and a polyT sequence **1964** complementary to polyA sequence **1962**. In some instances, nucleic acid barcode molecule **1918b** may comprise a sequencing primer sequence **1968** (e.g., a R1 sequence or partial R1 sequence), a barcode sequence **1922b** (e.g., the same barcode sequence as barcode sequence **1922a**), UMI sequence **1966**, and a polyT sequence **1964** complementary to polyA sequence **1962**. PolyT sequence **1964** may hybridize to polyA sequence **1962** of RNA molecule **1958**. RNA molecule **1958** may be reverse transcribed **1970** off of polyT sequence **1964** to provide an cDNA molecule comprising cDNA sequence **1972**. The reverse transcription process may use a reverse transcriptase with terminal transferase activity, which reverse transcriptase may append sequence **1974** to the resultant cDNA molecule comprising cDNA sequence **1972**. Sequence **1974** may be a polyC sequence. A template switch oligonucleotide **1978** comprising a primer sequence **1980** and a sequence complementary to sequence **1974** (e.g., a polyG sequence) may hybridize to the cDNA molecule. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the cDNA molecule in bulk solution. The cDNA molecule may undergo amplification (e.g., PCR) **1984**. Additional amplification (e.g., PCR) **1986** may be performed to provide a double-stranded amplification product **1988** that includes sequences of the nucleic acid barcode molecule **1918b**, the original RNA molecule or cDNA corresponding thereto, a flow cell adapter sequence **1998** (e.g., a P7 sequence), and an additional sequence **1990** that may comprise a sequencing primer or portion thereof (e.g., an R2 sequence) **1996**, a sample index sequence **1994**, and a flow cell adapter sequence (e.g., a P5 sequence) **1992**. The barcoded cDNA molecule may also or alternatively undergo fragmentation, end repair, dA tailing, ligation of one or more adapter sequences, and/or nucleic acid amplification.

[00216] In another aspect, the present disclosure provides a method for processing a biological sample (e.g., a nucleic acid sample), which method may comprise performing sequential transcription and reverse transcription processes within a partition. The method may comprise providing a partition (e.g., droplet or well) of a plurality of partitions comprising a nucleic acid molecule (e.g., DNA molecule) derived from a nucleic acid sample. The nucleic acid molecule may be transcribed (e.g., using a transcriptase) to provide an RNA molecule. The

RNA molecule may then be reverse transcribed (e.g., using a reverse transcriptase) within the partition to generate a complementary DNA (cDNA) molecule. The cDNA molecule may undergo further processing within the partition to provide a derivative of the cDNA molecule. The cDNA molecule or derivative thereof may be recovered from the partition (e.g., by pooling the contents of the plurality of partitions). The partition may be a well among a plurality of wells. Alternatively, the partition may be a droplet among a plurality of droplets.

[00217] A nucleic acid molecule (e.g., DNA molecule) processed according to the method provided herein may derive from a cell, cell bead, or cell nucleus. In some cases, the nucleic acid molecule may be included within the cell, cell bead, or cell nucleus. The nucleic acid molecule may be chromatin. The cell, cell bead, or cell nucleus comprising the nucleic acid molecule may be included within the partition. For example, the cell, cell bead, or cell nucleus may be co-partitioned with one or more reagents (e.g., as described herein) into a partition (e.g., droplet or well). The cell, cell bead, or cell nucleus may be lysed or permeabilized (e.g., within a partition) to provide access to the nucleic acid molecule therein (e.g., as described herein).

[00218] A nucleic acid molecule processed according to the method provided herein may be a DNA molecule, such as chromatin. In some cases, the method may further comprise processing an open chromatin structure of the nucleic acid sample with a transposase (e.g., included within a transposase-nucleic acid complex) to provide the nucleic acid molecule. For example, a nucleic acid molecule (e.g., within a cell, cell bead, or cell nucleus) may be contacted with a transposase-nucleic acid complex (e.g., as described herein). A transposase used in such a process may be, for example, a Tn5 transposase. A transposase-nucleic acid complex may have a structure such as that of **FIG. 9** or **FIG. 10**. Alternatively, a transposase-nucleic acid complex may comprise one or more transposon end oligonucleotide molecules, which transposon end oligonucleotide molecules comprise hairpin molecules. An example of such a transposase-nucleic acid complex is shown in **FIG. 11**.

[00219] A nucleic acid molecule processed using a transposase-nucleic acid complex comprising one or more hairpin molecules may be a tagged fragment comprising a double-stranded region comprising sequences corresponding to the nucleic acid molecule (e.g., chromatin) of the cell, cell bead, or cell nucleus from which it originates or is derived, as well as one or more hairpin molecules appended to either end of the double-stranded region. For example, the double-stranded region may comprise a first hairpin molecule at one end and a second hairpin molecule at a second end. Generally, only one end of a hairpin molecule may be attached to the double-stranded region, such that the tagged fragment comprises a gap at either end. For example, a hairpin molecule may be attached to a 3' end of the double-stranded

region. The hairpin molecule may comprise a promoter sequence, such as a T7 promoter sequence, and/or a UMI sequence.

[00220] Within the partition, the nucleic acid molecule (e.g., tagmented fragment) may undergo a gap filling process with a reverse transcriptase. The reverse transcriptase enzyme may be a mutant reverse transcriptase enzyme such as, but not limited to, Moloney Murine Leukemia Virus (MMLV) reverse transcriptase. In one aspect, the reverse transcriptase is a mutant MMLV reverse transcriptase such as, but not limited to, enzyme “42B” (see, US Patent Publication No. 20180312822). Enzyme 42B may reduce inhibition of reverse transcription of mRNAs from a single cell due to one or more unknown components present in cell lysate of the single cell when prepared in reaction volumes of, e.g., less than 1 nanoliter (nL). Enzyme 42B as compared to a commercially available mutant MMLV RT enzyme (CA-MMLV) may show improved reverse transcriptase activity. Such a process may generate a double-stranded nucleic acid molecule comprising the double-stranded region corresponding to the nucleic acid molecule (e.g., chromatin) of the cell, cell bead, or cell nucleus from which it is derived, the sequences of the hairpin molecules at either end of the double-stranded region, and sequences complementary to the sequences of the hairpin molecules. The double-stranded nucleic acid molecule may then undergo transcription with a T7 polymerase, which process begins at an end of a T7 promoter sequences of a hairpin molecules. Both strands may be transcribed in this manner to provide two nucleic acid strands each comprising the T7 promoter sequence, and a complement thereof; one or more transposon end sequences, and one or more complements thereof; and a sequence of the original nucleic acid molecule of the cell, cell bead, or cell nucleus. The strands may also comprise one or more spacer, UMI, or other sequences (e.g., from the hairpin molecules). A strand may then undergo a self-priming process in which the transposon end sequence and complement thereof of a hairpin molecule hybridize to one another to regenerate a hairpin molecule at an end of the strand. The hairpin molecule may serve as the priming site for reverse transcription. A reverse transcriptase process may then be performed (e.g., using a reverse transcriptase). Before, during, or after this process, a sequence may be appended to the end of the molecule, which sequence may be a polyC sequence. A template switching oligonucleotide comprising a sequence complementary to the appended sequence (e.g., a polyG sequence) may hybridize to the appended sequence. The template switching oligonucleotide may comprise a UMI sequence (e.g., a second UMI sequence that may index transcripts that undergo template switching), a barcode sequence, and/or a priming sequence such as a sequencing primer sequence or portion thereof (e.g., an R1 or R2 sequence, or portion thereof). The template switching oligonucleotide may be attached to a bead (e.g., a gel bead) included within the

partition. For example, the template switching oligonucleotide may be a nucleic acid barcode molecule of a plurality of nucleic acid barcode molecules attached to the bead (e.g., as described herein). The resultant partially double-stranded nucleic acid molecule may comprise a hairpin moiety; sequences corresponding to the original nucleic acid molecule of the cell, cell bead, or cell nucleus; and the sequences of the template switching oligonucleotide, including a barcode sequence (see, e.g., **FIG. 20**).

[00221] The partially double-stranded nucleic acid molecule may be released from the partition (e.g., droplet or well). Releasing materials from the partition may comprise breaking or disrupting a droplet. The contents of multiple partitions of the plurality of partitions may be pooled together to provide a bulk solution for further processing. Nucleic acid molecules (e.g., partially double-stranded nucleic acid molecule) of the partitions of the plurality of partitions may each be differentially barcoded such that the nucleic acid molecule of each such partition comprises a different barcode sequence.

[00222] Outside of the partition, the partially double-stranded nucleic acid molecule may be partially denatured to provide a single-stranded molecule (e.g., a single-stranded cDNA molecule). An RNase treatment may be used to remove the hairpin molecule as well as the shorter strand (e.g., the RNA sequence) of the partially double-stranded nucleic acid molecule. The single-stranded molecule remaining may include the template switching oligonucleotide comprising the barcode sequence and, optionally, UMI sequence. A primer molecule comprising a priming sequence complementary to the priming sequence of the template switching oligonucleotide may be provided and may hybridize to the priming sequence of the template switching oligonucleotide. The priming sequence of the primer molecule may be a 5'-blocked priming sequence. A polymerase with dA tailing activity (e.g., a Klenow fragment having 5'→3' polymerase activity, such as an exo-Klenow fragment lacking exonuclease activity) may be used to generate a second nucleic acid strand. The resultant second strand may be dA tailed. The first strand may also be dA tailed. However, if a 5'-blocking priming sequence is used in the preceding processes, the dA tail appended to the first strand may not be available as a hybridization site for another moiety. Instead, a priming sequence comprising a sequencing primer (e.g., an R1 sequence or complement thereof) and a flow cell adapter sequence (e.g., a P5 sequence or complement thereof) may hybridize to a complementary sequence of the double-stranded nucleic acid molecule. At the opposite end of the double-stranded nucleic acid molecule, the dA moiety appended to the end of the second strand may serve as a site for hybridization of a priming sequence comprising a dT moiety at an end, a sequencing primer (e.g., an R2 sequence or complement thereof), and a flow cell adapter sequence (e.g., a P7

sequence or complement thereof). The double-stranded nucleic acid molecule may then be subjected to conditions sufficient to perform one or more nucleic acid amplification reactions (e.g., PCR) to provide amplification products corresponding to the original nucleic acid molecule of the cell, cell bead, or cell nucleus. The amplification products may comprise flow cell adapter sequences (e.g., P5 and P7 sequences) at either end to facilitate sequencing (e.g., as described herein).

[00223] The method provided herein may overcome certain challenges of performing reverse transcription within partitions. For example, reverse transcriptase may have a DNA-dependent DNA polymerase activity, and/or terminal transferase activities. The latter may result in generation of variable overhangs under certain reaction conditions. In the methods provided herein, every insertion site may be provided a T7 promoter, averting losses that may otherwise be encountered via R1-R1 and R2-R2 interactions. Moreover, both mRNA and chromatin-derived fragments may be barcoded using the same biochemistry (e.g., RT template switching). Performance of linear amplification of both such strands of a nucleic acid molecule may provide strand awareness and introduce a new dimension for, e.g., ATAC-seq processes. Further, this method may enable isothermal linear amplification of transposase derived nucleic acid fragments within partitions. Notably, this method may be combined with any of the RNA workflows described elsewhere herein.

[00224] **FIG. 20** shows a workflow **2000** corresponding to the preceding example. Workflow **2000** may be performed in parallel with an RNA workflow, such as an RNA workflow of any of **FIGs. 12-19**. Multiple beads, each comprising nucleic acid barcode molecules configured for analysis of DNA or RNA molecules, may be included within a partition. Alternatively, a single bead (e.g., gel bead) comprising nucleic acid barcode molecules configured for analysis of both DNA and RNA molecules (e.g., as described herein) may be included within a partition. The single bead (e.g., in a single partition) may comprise a plurality of identical nucleic acid barcode molecules for both RNA and DNA analysis. In some cases, a single bead (e.g., within a single partition) comprises a first plurality of nucleic acid barcode molecules for DNA analysis and a second plurality of nucleic acid barcode molecules for RNA molecules, where the first and second plurality of nucleic acid barcode molecules comprise a common barcode sequence.

[00225] Template nucleic acid fragment (e.g., tagmented fragment) **2002** may be prepared (e.g., using a transposase-nucleic acid complex such as that shown in **FIG. 11**) and provided in a partition (as described herein). Template nucleic acid fragment **2002** may comprise hairpin moieties **2003** and **2004** and target sequences **2005** and **2006**. Template nucleic acid fragment **2002** also comprises gaps **2007**. Gaps **2007** may be filled using a reverse transcriptase (e.g., a

42B enzyme), which process may result in the generation of a double-stranded nucleic acid molecule comprising the double-stranded region corresponding to the original nucleic acid molecule (e.g., chromatin) of the cell, cell bead, or cell nucleus comprising sequences **2005** and **2006** and sequences of the hairpin molecules **2003** and **2004**. The double-stranded nucleic acid molecule may comprise transposon end sequences **2008**, promoter (e.g., T7 promoter) sequences **2010**, and UMI sequences **2012**. The double-stranded nucleic acid molecule may then undergo transcription with a T7 polymerase, which process begins at an end of a T7 promoter sequences of a hairpin molecule. Both strands may be transcribed in this manner to provide two nucleic acid strands. **FIG. 20** shows one such strand comprising T7 promoter sequence **2010**, and a complement thereof; one or more transposon end sequences **2008**, and one or more complements thereof; UMI sequence **2012**, and a complement of a UMI sequence; and an RNA sequence **2006'** corresponding to sequence **2006** of the original nucleic acid molecule of the cell, cell bead, or cell nucleus. The strand may then undergo a self-priming process in which the transposon end sequence and complement thereof of hairpin molecule **2004** hybridize to one another to regenerate a hairpin molecule at an end of the strand. Regenerated hairpin molecule **2004** may serve as the priming site for reverse transcription. Reverse transcription and template switching may then be performed (e.g., using a reverse transcriptase). The reverse transcription process may append sequence **2014** (e.g., a polyC sequence) to the resultant cDNA molecule comprising cDNA sequence **2026** and sequences **2012'** and **2008'** that are complementary to sequences **2012** and **2008**, respectively. The template switching process may comprise the use of a template switch oligonucleotide coupled to bead (e.g., gel bead) **2016** included within the partition. Bead (e.g., gel bead) **2016** may be coupled to nucleic acid barcode molecule **2018** that is the template switch oligonucleotide that comprises sequencing primer or portion thereof **2020**, barcode sequence **2022**, UMI sequence **2024**, and a sequence **2014'** that is complementary to sequence **2014** (e.g., a polyG sequence). The resultant cDNA molecule may comprise a first strand comprising nucleic acid barcode molecule **2018** and RNA sequence **2006'** and a second strand comprising cDNA sequence **2026**, appended sequence **2014**, and sequences **2020'**, **2022'**, and **2024'** that are complementary to sequences **2020**, **2022**, and **2024**, respectively.

[00226] The cDNA molecule may be released from the partition (e.g., droplet or well). Releasing materials from the partition may comprise breaking or disrupting a droplet. The contents of multiple partitions of the plurality of partitions may be pooled together to provide a bulk solution for further processing. Outside of the partition, the cDNA molecule may be treated with RNase to remove the hairpin molecule as well as the shorter strand (e.g., the RNA sequence) of the partially double-stranded nucleic acid molecule. The single-stranded molecule

remaining may include sequences **2020'**, **2022'**, **2024'**, **2014**, **2012'**, **2008'**, and **2026**. Primer molecule **2028** may then hybridize to sequence **2020'**. Primer molecule **2028** may be a 5'-blocked priming sequence. A polymerase with dA tailing activity (e.g., a Klenow fragment having 5'→3' polymerase activity, such as an exo-Klenow fragment lacking exonuclease activity) may be used to generate a second nucleic acid strand comprising sequence **2026'** that is complementary to cDNA sequence **2026**. The resultant second strand may be dA tailed. The first strand may also be dA tailed at an end of sequence **2020'**. However, if a 5'-blocking priming sequence is used in the preceding processes, the dA tail appended to the first strand may not be available as a hybridization site for another moiety. A priming sequence **2030** comprising a dT moiety, a sequencing primer (e.g., an R2 sequence or complement thereof) **2032** and a flow cell adapter sequence (e.g., a P7 sequence or complement thereof) **2034** may hybridize to the dA moiety of the double-stranded nucleic acid molecule. A priming sequence **2036** comprising a sequencing primer (e.g., an R1 sequence or complement thereof) **2038** and a flow cell adapter sequence (e.g., a P5 sequence or complement thereof) **2040** may hybridize to sequence **2028** of the double-stranded nucleic acid molecule. The double-stranded nucleic acid molecule may then be amplified to provide amplified product **2042**, which amplification product may be subjected to further processing such as nucleic acid sequencing.

[00227] **FIG. 21** provides an overview of a workflow **2100** for processing a nucleic acid molecule (e.g., a nucleic acid molecule included within a cell, cell bead, or cell nucleus). The nucleic acid molecule (e.g., DNA molecule, such as chromatin) is tagmented (e.g., as described herein) to generate a tagmented fragment. The tagmented fragment then undergoes transcription, reverse transcription, and barcoding within a partition (e.g., as described herein). The resultant products are released from the partition and subjected to one of two processes, the first of which provides an ATAC library and the second of which provides a gene expression library. The first process may involve RNase treatment to remove RNA and provide cDNA, dA tailing and ligation of a sequencing primer, and PCR. The second process may involve cDNA amplification; fragmentation, dA tailing, and ligation of a sequencing primer; and PCR.

[00228] The present disclosure also provides a method of processing a nucleic acid molecule of a cell, cell bead, or cell nucleus using a reverse transcriptase fill-in process coupled with a barcoding process. The nucleic acid molecule (e.g., DNA molecule) may derive from a cell, cell bead, or cell nucleus. In some cases, the nucleic acid molecule may be included within the cell, cell bead, or cell nucleus. The nucleic acid molecule may be chromatin. The cell, cell bead, or cell nucleus comprising the nucleic acid molecule may be included within the partition. For example, the cell, cell bead, or cell nucleus may be co-partitioned with one or more reagents

(e.g., as described herein) into a partition (e.g., droplet or well). The cell, cell bead, or cell nucleus may be lysed or permeabilized (e.g., within a partition) to provide access to the nucleic acid molecule therein (e.g., as described herein).

[00229] A nucleic acid molecule processed according to the method provided herein may be a DNA molecule, such as chromatin. In some cases, the method may further comprise processing an open chromatin structure of the nucleic acid sample with a transposase (e.g., included within a transposase-nucleic acid complex) to provide the nucleic acid molecule. For example, a nucleic acid molecule (e.g., within a cell, cell bead, or cell nucleus) may be contacted with a transposase-nucleic acid complex (e.g., as described herein). A transposase used in such a process may be, for example, a Tn5 transposase. A transposase-nucleic acid complex may have a structure such as that of **FIG. 9, 10, or 11**. Subsequent to generation of a tagmented fragment (e.g., as described herein), the transposase of the transposase-nucleic acid complex may leave or be removed (e.g., displaced, for example, by an enzyme). Alternatively, the transposase may remain in place. The tagmented fragment may comprise sequences corresponding to the original nucleic acid molecule of the cell, cell bead, or cell nucleus; transposon end sequences and sequences complementary thereto; and one or more sequencing primers or portions thereof. A splint sequence comprising a sequence complementary to a sequencing primer or portion thereof of the tagmented fragment may hybridize to the sequencing primer or portion thereof. The splint sequence may be ligated to a transposon end sequence or complement thereof of the tagmented fragment (e.g., using a ligase). Prior to or after hybridization and/or ligation of the splint sequence, the tagmented fragment may be partitioned into a partition of a plurality of partitions (e.g., droplets or wells). The tagmented fragment may be co-partitioned with one or more reagents. The tagmented fragment may be included within a cell, cell bead, or cell nucleus, which cell, cell bead, or cell nucleus may be lysed or permeabilized to provide access to the tagmented fragment therein (e.g., as described herein). A sequence of the splint sequence may then hybridize to a nucleic acid barcode molecule (e.g., a nucleic acid barcode molecule coupled to a bead, as described herein). The bead (e.g., gel bead) may comprise a plurality of nucleic acid barcode molecules, where a nucleic acid barcode molecule of the plurality of nucleic acid barcode molecules may comprise, for example, a flow cell adapter sequence, a barcode sequence, and a UMI sequence. The nucleic acid barcode molecule may also comprise an overhang sequence having sequence complementarity to a sequence of the splint sequence. The overhang sequence may hybridize to the sequence of the splint sequence. A transposase reserved in the tagmented fragment may block gap filling during these processes. The splint sequence may then be extended within the partition (e.g., using a reverse transcriptase).

[00230] Subsequent to the barcoding/template switching and extension (e.g., reverse transcription) processes, the contents of the partition of the plurality of partitions may be released from the partition (e.g., as described herein). Prior or subsequent to release of the contents of the partition, the nucleic acid barcode molecule may be ligated to the sequencing primer of the processed tagmented fragment. Outside of the partition, the nucleic acid barcode molecule may hybridize to the sequencing primer or portion thereof of the template nucleic acid fragment. If a transposase is reserved in the tagmented fragment, the transposase may leave the processed tagmented fragment (e.g., via a strand displacing polymerase) and the remaining gaps may be filled to provide a double-stranded nucleic acid molecule. Alternatively, gaps may be filled as described elsewhere herein. The double-stranded nucleic acid molecule may then be subjected to a nucleic acid amplification process (e.g., PCR, as described herein). Amplification may comprise incorporation of one or more additional sequences, such as one or more flow cell adapter sequences (e.g., P7 sequences).

[00231] **FIG. 22** shows an example schematic corresponding to the preceding example. Panel **2200** shows a workflow corresponding to processing of chromatin from a cell, cell bead, or cell nucleus, and panel **2250** shows a workflow corresponding to processing of an mRNA molecule from the cell, cell bead, or cell nucleus. Multiple beads (e.g., gel beads), each comprising nucleic acid barcode molecules configured for analysis of DNA or RNA molecules, may be included within a partition. Alternatively, a single bead (e.g., gel bead) comprising nucleic acid barcode molecules configured for analysis of both DNA and RNA molecules (e.g., as described herein) may be included within a given partition.

[00232] As shown in panel **2200**, in bulk solution, chromatin included within a cell, cell bead, or cell nucleus is processed (e.g., as described herein) to provide a template nucleic acid fragment (e.g., tagmented fragment) **2204** comprising insert sequence **2208** (e.g., region of open chromatin) and a complement thereof, transposon end sequences **2206** and complements thereof, sequencing primer or portion thereof **2202** (e.g., an R1 sequence), sequencing primer or portion thereof **2210** (e.g., an R2 sequence), and gaps **2207**. The cell, cell bead, or cell nucleus comprising template nucleic acid fragment **2204** may be lysed, permeabilized, or otherwise processed to provide access to template nucleic acid fragment **2204** (and one or more RNA molecules) therein. Template nucleic acid fragment **2204** may be contacted with splint sequence **2212**, which splint sequence may comprise a first sequence **2202'** that is complementary to sequencing primer or portion thereof **2202** and a second sequence **2224**. Sequence **2224** may comprise a blocking group (e.g., a 3' blocking group), which blocking group may prevent extension by reverse transcription. Sequence **2202'** may hybridize **2214** to sequence **2202** of

template nucleic acid fragment **2204** to provide a partially double-stranded nucleic acid molecule comprising splint sequence **2212** and template nucleic acid fragment **2204**. Sequence **2202'** may be ligated **2226** to the complement of transposon end sequence **2206** of template nucleic acid fragment **2204** (e.g., using a ligase). Template nucleic acid fragment **2204** attached to splint sequence **2212** may then be partitioned within a partition (e.g., droplet or well) within a plurality of partitions (e.g., as described herein). The partition may also include a bead (e.g., gel bead) **2216** coupled to nucleic acid barcode molecules **2218a** and **2218b**. Nucleic acid barcode molecule **2218a** may comprise a flow cell adapter sequence **2220a** (e.g., a P5 sequence), a barcode sequence **2222a**, and an overhang sequence **2224'** that is complementary to sequence **2224** of the splint sequence **2212**. Sequence **2224** may hybridize **2228** to sequence **2224'**. Splint sequence **2212** may then be extended **2230** (e.g., using a reverse transcriptase or DNA polymerase) to provide sequences **2220a'** and **2222a'** that are complementary to sequences **2220a** and **2222a** of nucleic acid barcode molecule **2218a**. Alternatively, sequence **2224** may hybridize to sequence **2224'** to provide a partially double-stranded nucleic acid molecule and nucleic acid barcode molecule **2218a** may be ligated (e.g., using a ligase) to sequence **2202** of template nucleic acid fragment **2204**. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the partially double-stranded nucleic acid molecule comprising nucleic acid barcode molecule **2218a** attached to splint sequence **2212** and template nucleic acid fragment **2204** in bulk solution. Sequence **2224'** of nucleic acid barcode molecule **2218a** may be ligated (e.g., using a ligase) **2232** to sequence **2202** of template nucleic acid fragment **2204**. In bulk solution, gaps **2207** may be filled **2234** via a gap filling extension process (e.g., using a DNA polymerase) to provide a double-stranded nucleic acid molecule. This molecule may also undergo amplification (e.g., PCR) to provide a double-stranded amplification product **2236** that includes sequences of the nucleic acid barcode molecule **2218a**, the original chromatin molecule, and, optionally, an additional sequence **2238** that may be a flow cell adapter sequence (e.g., a P7 sequence). Gaps may be filled in the partition prior to bulk processing.

[00233] In parallel to the chromatin workflow of panel **2200**, an RNA molecule deriving from the same cell, cell bead, or cell nucleus may be processed. As shown in panel **2250**, RNA molecule **2258** comprising RNA sequence **2260** and polyA sequence **2262** may be contacted **2264** with primer molecule **2252** comprising polyT sequence **2254** and additional primer sequence **2256**. RNA molecule **2258** may then be reverse transcribed **2276** off of polyT sequence **2254** using a reverse transcriptase with terminal transferase activity, which reverse transcriptase may append sequence **2270** to the resultant cDNA molecule comprising cDNA

sequence **2268**. Sequence **2270** may be a polyC sequence. Bead (e.g., gel bead) **2216** (e.g., the same bead described in panel **2200**) may be included within the partition and may be coupled to nucleic acid barcode molecule **2218b**. Nucleic acid barcode molecule **2218b** may comprise a flow cell adapter sequence **2220b** (e.g., a P5 sequence), a barcode sequence **2222b**, UMI sequence **2272**, and a sequence **2274** complementary to sequence **2270** (e.g., a polyG sequence). In some instances, nucleic acid barcode molecule **2218b** may comprise a sequencing primer sequence **2220b** (e.g., an R1 sequence or partial R1 sequence), a barcode sequence **2222b**, UMI sequence **2272**, and a sequence **2274** complementary to sequence **2270** (e.g., a polyG sequence). Nucleic acid barcode molecule **2218b** may be used to perform template switching **2278**, which process may also result in the generation of a barcoded cDNA molecule. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the barcoded cDNA molecule in bulk solution. The barcoded cDNA molecule may undergo amplification (e.g., PCR) **2280** to provide a double-stranded amplification product **2284** that includes sequences of the nucleic acid barcode molecule **2218b**, the original RNA molecule or cDNA corresponding thereto, a flow cell adapter sequence **2286**, and an additional sequence **2288** that may comprise a sequencing primer or portion thereof (e.g., an R2 sequence) **2290**, a sample index sequence **2292**, and a flow cell adapter sequence (e.g., a P7 sequence) **2294**. The barcoded cDNA molecule may also or alternatively undergo fragmentation, end repair, dA tailing, ligation of one or more adapter sequences, and/or nucleic acid amplification.

[00234] **FIG. 23** shows another example schematic corresponding to the preceding example. Panel **2300** shows a workflow corresponding to processing of chromatin from a cell, cell bead, or cell nucleus, and panel **2350** shows a workflow corresponding to processing of an mRNA molecule from the cell, cell bead, or cell nucleus. Multiple beads (e.g., gel beads), each comprising nucleic acid barcode molecules configured for analysis of DNA or RNA molecules, may be included within a partition. Alternatively, a single bead (e.g., gel bead) comprising nucleic acid barcode molecules configured for analysis of both DNA and RNA molecules (e.g., as described herein) may be included within a given partition.

[00235] As shown in panel **2300**, in bulk solution, chromatin included within a cell, cell bead, or cell nucleus is processed (e.g., as described herein) to provide a template nucleic acid fragment (e.g., tagmented fragment) **2304** comprising insert sequence **2308** (e.g., region of open chromatin) and a complement thereof, transposon end sequences **2306** and complements thereof, sequencing primer or portion thereof **2302** (e.g., an R1 sequence), sequencing primer or portion thereof **2310** (e.g., an R2 sequence), and gaps **2307**. The cell, cell bead, or cell nucleus comprising template nucleic acid fragment **2304** may be lysed, permeabilized, or otherwise

processed to provide access to template nucleic acid fragment **2304** (and one or more RNA molecules) therein. Template nucleic acid fragment **2304** may be contacted with splint sequence **2312**, which splint sequence may comprise a first sequence **2302'** that is complementary to sequencing primer or portion thereof **2302** and a second sequence **2324**. Sequence **2324** may comprise a blocking group (e.g., a 3' blocking group), which blocking group may prevent extension by reverse transcription. Sequence **2302'** may hybridize **2314** to sequence **2302** of template nucleic acid fragment **2304** to provide a partially double-stranded nucleic acid molecule comprising splint sequence **2312** and template nucleic acid fragment **2304**. Sequence **2302'** may be ligated **2326** to the complement of transposon end sequence **2306** of template nucleic acid fragment **2304** (e.g., using a ligase). Template nucleic acid fragment **2304** attached to splint sequence **2312** may then be partitioned within a partition (e.g., droplet or well) within a plurality of partitions (e.g., as described herein). The partition may also include a bead (e.g., gel bead) **2316** coupled to nucleic acid barcode molecules **2318a** and **2318b**. Nucleic acid barcode molecule **2318a** may comprise a flow cell adapter sequence **2320a** (e.g., a P5 sequence), a barcode sequence **2322a**, and an overhang sequence **2324'** that is complementary to sequence **2324** of the splint sequence **2312**. Sequence **2324** may hybridize **2328** to sequence **2324'**. Splint sequence **2312** may then be extended **2330** (e.g., using a reverse transcriptase or DNA polymerase) to provide sequences **2320a'** and **2322a'** that are complementary to sequences **2320a** and **2322a** of nucleic acid barcode molecule **2318a**. Alternatively, sequence **2324** may hybridize to sequence **2324'** to provide a partially double-stranded nucleic acid molecule and nucleic acid barcode molecule **2318a** may be ligated (e.g., using a ligase) to sequence **2302** of template nucleic acid fragment **2304**. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the partially double-stranded nucleic acid molecule comprising nucleic acid barcode molecule **2318a** attached to splint sequence **2312** and template nucleic acid fragment **2304** in bulk solution. Sequence **2324'** of nucleic acid barcode molecule **2318a** may be ligated (e.g., using a ligase) **2332** to sequence **2302** of template nucleic acid fragment **2304**. In bulk solution, gaps **2307** may be filled **2334** via a gap filling extension process (e.g., using a DNA polymerase) to provide a double-stranded nucleic acid molecule. This molecule may also undergo amplification (e.g., PCR) to provide a double-stranded amplification product **2336** that includes sequences of the nucleic acid barcode molecule **2318a**, the original chromatin molecule, and, optionally, an additional sequence **2338** that may be a flow cell adapter sequence (e.g., a P7 sequence). Gaps may be filled in the partition prior to bulk processing.

[00236] In parallel to the chromatin workflow of panel **2300**, an RNA molecule deriving from the same cell, cell bead, or cell nucleus may be processed. As shown in panel **2350**, RNA molecule **2358** comprising RNA sequence **2360** and polyA sequence **2362** and gel bead **2316** may be provided within a partition. Bead (e.g., gel bead) **2316** (e.g., the same bead described in panel **2300**) may be included within the partition and may be coupled to nucleic acid barcode molecule **2318b**. Nucleic acid barcode molecule **2318b** may comprise a flow cell adapter sequence **2368** (e.g., a P5 sequence), a barcode sequence **2322b** (e.g., the same barcode sequence as barcode sequence **2322a**), UMI sequence **2366**, and a polyT sequence **2364** complementary to polyA sequence **2362**. In some instances, nucleic acid barcode molecule **2318b** may comprise a sequencing primer sequence **2368** (e.g., a R1 sequence or partial R1 sequence), a barcode sequence **2322b** (e.g., the same barcode sequence as barcode sequence **2322a**), UMI sequence **2366**, and a polyT sequence **2364** complementary to polyA sequence **2362**. PolyT sequence **2364** may hybridize to polyA sequence **2362** of RNA molecule **2358**. RNA molecule **2358** may be reverse transcribed **2370** off of polyT sequence **2364** to provide an cDNA molecule comprising cDNA sequence **2372**. The reverse transcription process may use a reverse transcriptase with terminal transferase activity, which reverse transcriptase may append sequence **2374** to the resultant cDNA molecule comprising cDNA sequence **2372**. Sequence **2374** may be a polyC sequence. A template switch oligonucleotide **2378** comprising a primer sequence **2380** and a sequence complementary to sequence **2374** (e.g., a polyG sequence) may hybridize to the cDNA molecule. The contents of the partition may then be recovered in bulk solution (e.g., a droplet may be broken) to provide the cDNA molecule in bulk solution. The cDNA molecule may undergo amplification (e.g., PCR) **2384**. Additional amplification (e.g., PCR) **2386** may be performed to provide a double-stranded amplification product **2388** that includes sequences of the nucleic acid barcode molecule **2318b**, the original RNA molecule or cDNA corresponding thereto, a flow cell adapter sequence **2398** (e.g., a P7 sequence), and an additional sequence **2390** that may comprise a sequencing primer or portion thereof (e.g., an R2 sequence) **2396**, a sample index sequence **2394**, and a flow cell adapter sequence (e.g., a P5 sequence) **2392**. The barcoded cDNA molecule may also or alternatively undergo fragmentation, end repair, dA tailing, ligation of one or more adapter sequences, and/or nucleic acid amplification.

Systems and methods for sample compartmentalization

[00237] In an aspect, the systems and methods described herein provide for the compartmentalization, depositing, or partitioning of one or more particles (e.g., biological particles, macromolecular constituents of biological particles, beads, reagents, etc.) into discrete

compartments or partitions (referred to interchangeably herein as partitions), where each partition maintains separation of its own contents from the contents of other partitions. The partition can be a droplet in an emulsion. A partition may comprise one or more other partitions.

[00238] A partition may include one or more particles. A partition may include one or more types of particles. For example, a partition of the present disclosure may comprise one or more biological particles and/or macromolecular constituents thereof. A partition may comprise one or more gel beads. A partition may comprise one or more cell beads. A partition may include a single gel bead, a single cell bead, or both a single cell bead and single gel bead. A partition may include one or more reagents. Alternatively, a partition may be unoccupied. For example, a partition may not comprise a bead. A cell bead can be a biological particle and/or one or more of its macromolecular constituents encased inside of a gel or polymer matrix, such as via polymerization of a droplet containing the biological particle and precursors capable of being polymerized or gelled. Unique identifiers, such as barcodes, may be injected into the droplets previous to, subsequent to, or concurrently with droplet generation, such as via a microcapsule (e.g., bead), as described elsewhere herein. Microfluidic channel networks (e.g., on a chip) can be utilized to generate partitions as described herein. Alternative mechanisms may also be employed in the partitioning of individual biological particles, including porous membranes through which aqueous mixtures of cells are extruded into non-aqueous fluids.

[00239] The partitions can be flowable within fluid streams. The partitions may comprise, for example, micro-vesicles that have an outer barrier surrounding an inner fluid center or core. In some cases, the partitions may comprise a porous matrix that is capable of entraining and/or retaining materials within its matrix. The partitions can be droplets of a first phase within a second phase, wherein the first and second phases are immiscible. For example, the partitions can be droplets of aqueous fluid within a non-aqueous continuous phase (e.g., oil phase). In another example, the partitions can be droplets of a non-aqueous fluid within an aqueous phase. In some examples, the partitions may be provided in a water-in-oil emulsion or oil-in-water emulsion. A variety of different vessels are described in, for example, U.S. Patent Application Publication No. 2014/0155295, which is entirely incorporated herein by reference for all purposes. Emulsion systems for creating stable droplets in non-aqueous or oil continuous phases are described in, for example, U.S. Patent Application Publication No. 2010/0105112, which is entirely incorporated herein by reference for all purposes.

[00240] In the case of droplets in an emulsion, allocating individual particles to discrete partitions may in one non-limiting example be accomplished by introducing a flowing stream of particles in an aqueous fluid into a flowing stream of a non-aqueous fluid, such that droplets are

generated at the junction of the two streams. Fluid properties (e.g., fluid flow rates, fluid viscosities, etc.), particle properties (e.g., volume fraction, particle size, particle concentration, etc.), microfluidic architectures (e.g., channel geometry, etc.), and other parameters may be adjusted to control the occupancy of the resulting partitions (e.g., number of biological particles per partition, number of beads per partition, etc.). For example, partition occupancy can be controlled by providing the aqueous stream at a certain concentration and/or flow rate of particles. To generate single biological particle partitions, the relative flow rates of the immiscible fluids can be selected such that, on average, the partitions may contain less than one biological particle per partition in order to ensure that those partitions that are occupied are primarily singly occupied. In some cases, partitions among a plurality of partitions may contain at most one biological particle (e.g., bead, DNA, cell or cellular material). The various parameters (e.g., fluid properties, particle properties, microfluidic architectures, etc.) may be selected or adjusted such that a majority of partitions are occupied, for example, allowing for only a small percentage of unoccupied partitions. The flows and channel architectures can be controlled as to ensure a given number of singly occupied partitions, less than a certain level of unoccupied partitions and/or less than a certain level of multiply occupied partitions.

[00241] FIG. 1 shows an example of a microfluidic channel structure **100** for partitioning individual biological particles. The channel structure **100** can include channel segments **102**, **104**, **106** and **108** communicating at a channel junction **110**. In operation, a first aqueous fluid **112** that includes suspended biological particles (or cells) **114** may be transported along channel segment **102** into junction **110**, while a second fluid **116** that is immiscible with the aqueous fluid **112** is delivered to the junction **110** from each of channel segments **104** and **106** to create discrete droplets **118**, **120** of the first aqueous fluid **112** flowing into channel segment **108**, and flowing away from junction **110**. The channel segment **108** may be fluidically coupled to an outlet reservoir where the discrete droplets can be stored and/or harvested. A discrete droplet generated may include an individual biological particle **114** (such as droplets **118**). A discrete droplet generated may include more than one individual biological particle **114** (not shown in FIG. 1). A discrete droplet may contain no biological particle **114** (such as droplet **120**). Each discrete partition may maintain separation of its own contents (e.g., individual biological particle **114**) from the contents of other partitions.

[00242] The second fluid **116** can comprise an oil, such as a fluorinated oil, that includes a fluorosurfactant for stabilizing the resulting droplets, for example, inhibiting subsequent coalescence of the resulting droplets **118**, **120**. Examples of particularly useful partitioning

fluids and fluorosurfactants are described, for example, in U.S. Patent Application Publication No. 2010/0105112, which is entirely incorporated herein by reference for all purposes.

[00243] As will be appreciated, the channel segments described herein may be coupled to any of a variety of different fluid sources or receiving components, including reservoirs, tubing, manifolds, or fluidic components of other systems. As will be appreciated, the microfluidic channel structure **100** may have other geometries. For example, a microfluidic channel structure can have more than one channel junction. For example, a microfluidic channel structure can have 2, 3, 4, or 5 channel segments each carrying particles (e.g., biological particles, cell beads, and/or gel beads) that meet at a channel junction. Fluid may be directed to flow along one or more channels or reservoirs via one or more fluid flow units. A fluid flow unit can comprise compressors (e.g., providing positive pressure), pumps (e.g., providing negative pressure), actuators, and the like to control flow of the fluid. Fluid may also or otherwise be controlled via applied pressure differentials, centrifugal force, electrokinetic pumping, vacuum, capillary or gravity flow, or the like.

[00244] The generated droplets may comprise two subsets of droplets: (1) occupied droplets **118**, containing one or more biological particles **114**, and (2) unoccupied droplets **120**, not containing any biological particles **114**. Occupied droplets **118** may comprise singly occupied droplets (having one biological particle) and multiply occupied droplets (having more than one biological particle). As described elsewhere herein, in some cases, the majority of occupied partitions can include no more than one biological particle per occupied partition and some of the generated partitions can be unoccupied (of any biological particle). In some cases, though, some of the occupied partitions may include more than one biological particle. In some cases, the partitioning process may be controlled such that fewer than about 25% of the occupied partitions contain more than one biological particle, and in many cases, fewer than about 20% of the occupied partitions have more than one biological particle, while in some cases, fewer than about 10% or even fewer than about 5% of the occupied partitions include more than one biological particle per partition.

[00245] In some cases, it may be desirable to minimize the creation of excessive numbers of empty partitions, such as to reduce costs and/or increase efficiency. While this minimization may be achieved by providing a sufficient number of biological particles (e.g., biological particles **114**) at the partitioning junction **110**, such as to ensure that at least one biological particle is encapsulated in a partition, the Poissonian distribution may expectedly increase the number of partitions that include multiple biological particles. As such, where singly occupied partitions are to be obtained, at most about 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%,

50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5% or less of the generated partitions can be unoccupied.

[00246] In some cases, the flow of one or more of the biological particles (e.g., in channel segment **102**), or other fluids directed into the partitioning junction (e.g., in channel segments **104**, **106**) can be controlled such that, in many cases, no more than about 50% of the generated partitions, no more than about 25% of the generated partitions, or no more than about 10% of the generated partitions are unoccupied. These flows can be controlled so as to present a non-Poissonian distribution of single-occupied partitions while providing lower levels of unoccupied partitions. The above noted ranges of unoccupied partitions can be achieved while still providing any of the single occupancy rates described above. For example, in many cases, the use of the systems and methods described herein can create resulting partitions that have multiple occupancy rates of less than about 25%, less than about 20%, less than about 15%, less than about 10%, and in many cases, less than about 5%, while having unoccupied partitions of less than about 50%, less than about 40%, less than about 30%, less than about 20%, less than about 10%, less than about 5%, or less.

[00247] As will be appreciated, the above-described occupancy rates are also applicable to partitions that include both biological particles and additional reagents, including, but not limited to, microcapsules or beads (e.g., gel beads) carrying barcoded nucleic acid molecules (e.g., oligonucleotides) (described in relation to **FIG. 2**). The occupied partitions (e.g., at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 99% of the occupied partitions) can include both a microcapsule (e.g., bead) comprising barcoded nucleic acid molecules and a biological particle.

[00248] In another aspect, in addition to or as an alternative to droplet based partitioning, biological particles may be encapsulated within a microcapsule that comprises an outer shell, layer or porous matrix in which is entrained one or more individual biological particles or small groups of biological particles. The microcapsule may include other reagents. Encapsulation of biological particles may be performed by a variety of processes. Such processes may combine an aqueous fluid containing the biological particles with a polymeric precursor material that may be capable of being formed into a gel or other solid or semi-solid matrix upon application of a particular stimulus to the polymer precursor. Such stimuli can include, for example, thermal stimuli (e.g., either heating or cooling), photo-stimuli (e.g., through photo-curing), chemical stimuli (e.g., through crosslinking, polymerization initiation of the precursor (e.g., through added initiators)), mechanical stimuli, or a combination thereof.

[00249] Preparation of microcapsules comprising biological particles may be performed by a variety of methods. For example, air knife droplet or aerosol generators may be used to dispense droplets of precursor fluids into gelling solutions in order to form microcapsules that include individual biological particles or small groups of biological particles. Likewise, membrane based encapsulation systems may be used to generate microcapsules comprising encapsulated biological particles as described herein. Microfluidic systems of the present disclosure, such as that shown in **FIG. 1**, may be readily used in encapsulating cells as described herein. In particular, and with reference to **FIG. 1**, the aqueous fluid **112** comprising (i) the biological particles **114** and (ii) the polymer precursor material (not shown) is flowed into channel junction **110**, where it is partitioned into droplets **118**, **120** through the flow of non-aqueous fluid **116**. In the case of encapsulation methods, non-aqueous fluid **116** may also include an initiator (not shown) to cause polymerization and/or crosslinking of the polymer precursor to form the microcapsule that includes the entrained biological particles. Examples of polymer precursor/initiator pairs include those described in U.S. Patent Application Publication No. 2014/0378345, which is entirely incorporated herein by reference for all purposes.

[00250] For example, in the case where the polymer precursor material comprises a linear polymer material, such as a linear polyacrylamide, PEG, or other linear polymeric material, the activation agent may comprise a cross-linking agent, or a chemical that activates a cross-linking agent within the formed droplets. Likewise, for polymer precursors that comprise polymerizable monomers, the activation agent may comprise a polymerization initiator. For example, in certain cases, where the polymer precursor comprises a mixture of acrylamide monomer with a N,N'-bis-(acryloyl)cystamine (BAC) comonomer, an agent such as tetraethylmethylenediamine (TEMED) may be provided within the second fluid streams **116** in channel segments **104** and **106**, which can initiate the copolymerization of the acrylamide and BAC into a cross-linked polymer network, or hydrogel.

[00251] Upon contact of the second fluid stream **116** with the first fluid stream **112** at junction **110**, during formation of droplets, the TEMED may diffuse from the second fluid **116** into the aqueous fluid **112** comprising the linear polyacrylamide, which will activate the crosslinking of the polyacrylamide within the droplets **118**, **120**, resulting in the formation of gel (e.g., hydrogel) microcapsules, as solid or semi-solid beads or particles entraining the cells **114**. Although described in terms of polyacrylamide encapsulation, other 'activatable' encapsulation compositions may also be employed in the context of the methods and compositions described herein. For example, formation of alginate droplets followed by exposure to divalent metal ions (e.g., Ca^{2+} ions), can be used as an encapsulation process using the described processes.

Likewise, agarose droplets may also be transformed into capsules through temperature based gelling (e.g., upon cooling, etc.).

[00252] In some cases, encapsulated biological particles can be selectively releasable from the microcapsule, such as through passage of time or upon application of a particular stimulus, that degrades the microcapsule sufficiently to allow the biological particles (e.g., cell), or its other contents to be released from the microcapsule, such as into a partition (e.g., droplet). For example, in the case of the polyacrylamide polymer described above, degradation of the microcapsule may be accomplished through the introduction of an appropriate reducing agent, such as DTT or the like, to cleave disulfide bonds that cross-link the polymer matrix. See, for example, U.S. Patent Application Publication No. 2014/0378345, which is entirely incorporated herein by reference for all purposes.

[00253] The biological particle can be subjected to other conditions sufficient to polymerize or gel the precursors. The conditions sufficient to polymerize or gel the precursors may comprise exposure to heating, cooling, electromagnetic radiation, and/or light. The conditions sufficient to polymerize or gel the precursors may comprise any conditions sufficient to polymerize or gel the precursors. Following polymerization or gelling, a polymer or gel may be formed around the biological particle. The polymer or gel may be diffusively permeable to chemical or biochemical reagents. The polymer or gel may be diffusively impermeable to macromolecular constituents of the biological particle. In this manner, the polymer or gel may act to allow the biological particle to be subjected to chemical or biochemical operations while spatially confining the macromolecular constituents to a region of the droplet defined by the polymer or gel. The polymer or gel may include one or more of disulfide cross-linked polyacrylamide, agarose, alginate, polyvinyl alcohol, polyethylene glycol (PEG)-diacrylate, PEG-acrylate, PEG-thiol, PEG-azide, PEG-alkyne, other acrylates, chitosan, hyaluronic acid, collagen, fibrin, gelatin, or elastin. The polymer or gel may comprise any other polymer or gel.

[00254] The polymer or gel may be functionalized to bind to targeted analytes, such as nucleic acids, proteins, carbohydrates, lipids or other analytes. The polymer or gel may be polymerized or gelled via a passive mechanism. The polymer or gel may be stable in alkaline conditions or at elevated temperature. The polymer or gel may have mechanical properties similar to the mechanical properties of the bead. For instance, the polymer or gel may be of a similar size to the bead. The polymer or gel may have a mechanical strength (e.g. tensile strength) similar to that of the bead. The polymer or gel may be of a lower density than an oil. The polymer or gel may be of a density that is roughly similar to that of a buffer. The polymer or gel may have a tunable pore size. The pore size may be chosen to, for instance, retain denatured nucleic acids.

The pore size may be chosen to maintain diffusive permeability to exogenous chemicals such as sodium hydroxide (NaOH) and/or endogenous chemicals such as inhibitors. The polymer or gel may be biocompatible. The polymer or gel may maintain or enhance cell viability. The polymer or gel may be biochemically compatible. The polymer or gel may be polymerized and/or depolymerized thermally, chemically, enzymatically, and/or optically.

[00255] The polymer may comprise poly(acrylamide-co-acrylic acid) crosslinked with disulfide linkages. The preparation of the polymer may comprise a two-step reaction. In the first activation step, poly(acrylamide-co-acrylic acid) may be exposed to an acylating agent to convert carboxylic acids to esters. For instance, the poly(acrylamide-co-acrylic acid) may be exposed to 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM). The polyacrylamide-co-acrylic acid may be exposed to other salts of 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium. In the second cross-linking step, the ester formed in the first step may be exposed to a disulfide crosslinking agent. For instance, the ester may be exposed to cystamine (2,2'-dithiobis(ethylamine)). Following the two steps, the biological particle may be surrounded by polyacrylamide strands linked together by disulfide bridges. In this manner, the biological particle may be encased inside of or comprise a gel or matrix (e.g., polymer matrix) to form a "cell bead." A cell bead can contain biological particles (e.g., a cell) or macromolecular constituents (e.g., RNA, DNA, proteins, etc.) of biological particles. A cell bead may include a single cell or multiple cells, or a derivative of the single cell or multiple cells. For example after lysing and washing the cells, inhibitory components from cell lysates can be washed away and the macromolecular constituents can be bound as cell beads. Systems and methods disclosed herein can be applicable to both cell beads (and/or droplets or other partitions) containing biological particles and cell beads (and/or droplets or other partitions) containing macromolecular constituents of biological particles.

[00256] Encapsulated biological particles can provide certain potential advantages of being more storable and more portable than droplet-based partitioned biological particles. Furthermore, in some cases, it may be desirable to allow biological particles to incubate for a select period of time before analysis, such as in order to characterize changes in such biological particles over time, either in the presence or absence of different stimuli. In such cases, encapsulation may allow for longer incubation than partitioning in emulsion droplets, although in some cases, droplet partitioned biological particles may also be incubated for different periods of time, e.g., at least 10 seconds, at least 30 seconds, at least 1 minute, at least 5 minutes, at least 10 minutes, at least 30 minutes, at least 1 hour, at least 2 hours, at least 5 hours, or at least 10 hours or more. The encapsulation of biological particles may constitute the partitioning of the

biological particles into which other reagents are co-partitioned. Alternatively or in addition, encapsulated biological particles may be readily deposited into other partitions (e.g., droplets) as described above.

Beads

[00257] A partition may comprise one or more unique identifiers, such as barcodes. Barcodes may be previously, subsequently or concurrently delivered to the partitions that hold the compartmentalized or partitioned biological particle. For example, barcodes may be injected into droplets previous to, subsequent to, or concurrently with droplet generation. The delivery of the barcodes to a particular partition allows for the later attribution of the characteristics of the individual biological particle to the particular partition. Barcodes may be delivered, for example on a nucleic acid molecule (e.g., an oligonucleotide), to a partition via any suitable mechanism. Barcoded nucleic acid molecules can be delivered to a partition via a microcapsule. A microcapsule, in some instances, can comprise a bead. Beads are described in further detail below.

[00258] In some cases, barcoded nucleic acid molecules can be initially associated with the microcapsule and then released from the microcapsule. Release of the barcoded nucleic acid molecules can be passive (e.g., by diffusion out of the microcapsule). In addition or alternatively, release from the microcapsule can be upon application of a stimulus which allows the barcoded nucleic acid nucleic acid molecules to dissociate or to be released from the microcapsule. Such stimulus may disrupt the microcapsule, an interaction that couples the barcoded nucleic acid molecules to or within the microcapsule, or both. Such stimulus can include, for example, a thermal stimulus, photo-stimulus, chemical stimulus (e.g., change in pH or use of a reducing agent(s)), a mechanical stimulus, a radiation stimulus; a biological stimulus (e.g., enzyme), or any combination thereof.

[00259] **FIG. 2** shows an example of a microfluidic channel structure **200** for delivering barcode carrying beads to droplets. The channel structure **200** can include channel segments **201**, **202**, **204**, **206** and **208** communicating at a channel junction **210**. In operation, the channel segment **201** may transport an aqueous fluid **212** that includes a plurality of beads **214** (e.g., with nucleic acid molecules, oligonucleotides, molecular tags) along the channel segment **201** into junction **210**. The plurality of beads **214** may be sourced from a suspension of beads. For example, the channel segment **201** may be connected to a reservoir comprising an aqueous suspension of beads **214**. The channel segment **202** may transport the aqueous fluid **212** that includes a plurality of biological particles **216** along the channel segment **202** into junction **210**. The plurality of biological particles **216** may be sourced from a suspension of biological

particles. For example, the channel segment **202** may be connected to a reservoir comprising an aqueous suspension of biological particles **216**. In some instances, the aqueous fluid **212** in either the first channel segment **201** or the second channel segment **202**, or in both segments, can include one or more reagents, as further described below. A second fluid **218** that is immiscible with the aqueous fluid **212** (e.g., oil) can be delivered to the junction **210** from each of channel segments **204** and **206**. Upon meeting of the aqueous fluid **212** from each of channel segments **201** and **202** and the second fluid **218** from each of channel segments **204** and **206** at the channel junction **210**, the aqueous fluid **212** can be partitioned as discrete droplets **220** in the second fluid **218** and flow away from the junction **210** along channel segment **208**. The channel segment **208** may deliver the discrete droplets to an outlet reservoir fluidly coupled to the channel segment **208**, where they may be harvested.

[00260] As an alternative, the channel segments **201** and **202** may meet at another junction upstream of the junction **210**. At such junction, beads and biological particles may form a mixture that is directed along another channel to the junction **210** to yield droplets **220**. The mixture may provide the beads and biological particles in an alternating fashion, such that, for example, a droplet comprises a single bead and a single biological particle.

[00261] Beads, biological particles and droplets may flow along channels at substantially regular flow profiles (e.g., at regular flow rates). Such regular flow profiles may permit a droplet to include a single bead and a single biological particle. Such regular flow profiles may permit the droplets to have an occupancy (e.g., droplets having beads and biological particles) greater than 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 95%. Such regular flow profiles and devices that may be used to provide such regular flow profiles are provided in, for example, U.S. Patent Publication No. 2015/0292988, which is entirely incorporated herein by reference.

[00262] The second fluid **218** can comprise an oil, such as a fluorinated oil, that includes a fluorosurfactant for stabilizing the resulting droplets, for example, inhibiting subsequent coalescence of the resulting droplets **220**.

[00263] A discrete droplet that is generated may include an individual biological particle **216**. A discrete droplet that is generated may include a barcode or other reagent carrying bead **214**. A discrete droplet generated may include both an individual biological particle and a barcode carrying bead, such as droplets **220**. In some instances, a discrete droplet may include more than one individual biological particle or no biological particle. In some instances, a discrete droplet may include more than one bead or no bead. A discrete droplet may be unoccupied (e.g., no beads, no biological particles).

[00264] Beneficially, a discrete droplet partitioning a biological particle and a barcode carrying bead may effectively allow the attribution of the barcode to macromolecular constituents of the biological particle within the partition. The contents of a partition may remain discrete from the contents of other partitions.

[00265] As will be appreciated, the channel segments described herein may be coupled to any of a variety of different fluid sources or receiving components, including reservoirs, tubing, manifolds, or fluidic components of other systems. As will be appreciated, the microfluidic channel structure **200** may have other geometries. For example, a microfluidic channel structure can have more than one channel junctions. For example, a microfluidic channel structure can have 2, 3, 4, or 5 channel segments each carrying beads that meet at a channel junction. Fluid may be directed flow along one or more channels or reservoirs via one or more fluid flow units. A fluid flow unit can comprise compressors (e.g., providing positive pressure), pumps (e.g., providing negative pressure), actuators, and the like to control flow of the fluid. Fluid may also or otherwise be controlled via applied pressure differentials, centrifugal force, electrokinetic pumping, vacuum, capillary or gravity flow, or the like.

[00266] A bead may be porous, non-porous, solid, semi-solid, semi-fluidic, fluidic, and/or a combination thereof. In some instances, a bead may be dissolvable, disruptable, and/or degradable. In some cases, a bead may not be degradable. In some cases, the bead may be a gel bead. A gel bead may be a hydrogel bead. A gel bead may be formed from molecular precursors, such as a polymeric or monomeric species. A semi-solid bead may be a liposomal bead. Solid beads may comprise metals including iron oxide, gold, and silver. In some cases, the bead may be a silica bead. In some cases, the bead can be rigid. In other cases, the bead may be flexible and/or compressible.

[00267] A bead may be of any suitable shape. Examples of bead shapes include, but are not limited to, spherical, non-spherical, oval, oblong, amorphous, circular, cylindrical, and variations thereof.

[00268] Beads may be of uniform size or heterogeneous size. In some cases, the diameter of a bead may be at least about 10 nanometers (nm), 100 nm, 500 nm, 1 micrometer (μm), 5 μm , 10 μm , 20 μm , 30 μm , 40 μm , 50 μm , 60 μm , 70 μm , 80 μm , 90 μm , 100 μm , 250 μm , 500 μm , 1mm, or greater. In some cases, a bead may have a diameter of less than about 10 nm, 100 nm, 500 nm, 1 μm , 5 μm , 10 μm , 20 μm , 30 μm , 40 μm , 50 μm , 60 μm , 70 μm , 80 μm , 90 μm , 100 μm , 250 μm , 500 μm , 1mm, or less. In some cases, a bead may have a diameter in the range of about 40-75 μm , 30-75 μm , 20-75 μm , 40-85 μm , 40-95 μm , 20-100 μm , 10-100 μm , 1-100 μm , 20-250 μm , or 20-500 μm .

[00269] In certain aspects, beads can be provided as a population or plurality of beads having a relatively monodisperse size distribution. Where it may be desirable to provide relatively consistent amounts of reagents within partitions, maintaining relatively consistent bead characteristics, such as size, can contribute to the overall consistency. In particular, the beads described herein may have size distributions that have a coefficient of variation in their cross-sectional dimensions of less than 50%, less than 40%, less than 30%, less than 20%, and in some cases less than 15%, less than 10%, less than 5%, or less.

[00270] A bead may comprise natural and/or synthetic materials. For example, a bead can comprise a natural polymer, a synthetic polymer or both natural and synthetic polymers. Examples of natural polymers include proteins and sugars such as deoxyribonucleic acid, rubber, cellulose, starch (e.g., amylose, amylopectin), proteins, enzymes, polysaccharides, silks, polyhydroxyalkanoates, chitosan, dextran, collagen, carrageenan, ispaghula, acacia, agar, gelatin, shellac, sterculia gum, xanthan gum, Corn sugar gum, guar gum, gum karaya, agarose, alginic acid, alginate, or natural polymers thereof. Examples of synthetic polymers include acrylics, nylons, silicones, spandex, viscose rayon, polycarboxylic acids, polyvinyl acetate, polyacrylamide, polyacrylate, polyethylene glycol, polyurethanes, polylactic acid, silica, polystyrene, polyacrylonitrile, polybutadiene, polycarbonate, polyethylene, polyethylene terephthalate, poly(chlorotrifluoroethylene), poly(ethylene oxide), poly(ethylene terephthalate), polyethylene, polyisobutylene, poly(methyl methacrylate), poly(oxymethylene), polyformaldehyde, polypropylene, polystyrene, poly(tetrafluoroethylene), poly(vinyl acetate), poly(vinyl alcohol), poly(vinyl chloride), poly(vinylidene dichloride), poly(vinylidene difluoride), poly(vinyl fluoride) and/or combinations (e.g., co-polymers) thereof. Beads may also be formed from materials other than polymers, including lipids, micelles, ceramics, glass-ceramics, material composites, metals, other inorganic materials, and others.

[00271] In some instances, the bead may contain molecular precursors (e.g., monomers or polymers), which may form a polymer network via polymerization of the molecular precursors. In some cases, a precursor may be an already polymerized species capable of undergoing further polymerization via, for example, a chemical cross-linkage. In some cases, a precursor can comprise one or more of an acrylamide or a methacrylamide monomer, oligomer, or polymer. In some cases, the bead may comprise prepolymers, which are oligomers capable of further polymerization. For example, polyurethane beads may be prepared using prepolymers. In some cases, the bead may contain individual polymers that may be further polymerized together. In some cases, beads may be generated via polymerization of different precursors, such that they comprise mixed polymers, co-polymers, and/or block co-polymers. In some cases, the bead may

comprise covalent or ionic bonds between polymeric precursors (e.g., monomers, oligomers, linear polymers), nucleic acid molecules (e.g., oligonucleotides), primers, and other entities. In some cases, the covalent bonds can be carbon-carbon bonds, thioether bonds, or carbon-heteroatom bonds.

[00272] Cross-linking may be permanent or reversible, depending upon the particular cross-linker used. Reversible cross-linking may allow for the polymer to linearize or dissociate under appropriate conditions. In some cases, reversible cross-linking may also allow for reversible attachment of a material bound to the surface of a bead. In some cases, a cross-linker may form disulfide linkages. In some cases, the chemical cross-linker forming disulfide linkages may be cystamine or a modified cystamine.

[00273] In some cases, disulfide linkages can be formed between molecular precursor units (e.g., monomers, oligomers, or linear polymers) or precursors incorporated into a bead and nucleic acid molecules (e.g., oligonucleotides). Cystamine (including modified cystamines), for example, is an organic agent comprising a disulfide bond that may be used as a crosslinker agent between individual monomeric or polymeric precursors of a bead. Polyacrylamide may be polymerized in the presence of cystamine or a species comprising cystamine (e.g., a modified cystamine) to generate polyacrylamide gel beads comprising disulfide linkages (e.g., chemically degradable beads comprising chemically-reducible cross-linkers). The disulfide linkages may permit the bead to be degraded (or dissolved) upon exposure of the bead to a reducing agent.

[00274] In some cases, chitosan, a linear polysaccharide polymer, may be crosslinked with glutaraldehyde via hydrophilic chains to form a bead. Crosslinking of chitosan polymers may be achieved by chemical reactions that are initiated by heat, pressure, change in pH, and/or radiation.

[00275] In some cases, a bead may comprise an acrydite moiety, which in certain aspects may be used to attach one or more nucleic acid molecules (e.g., barcode sequence, barcoded nucleic acid molecule, barcoded oligonucleotide, primer, or other oligonucleotide) to the bead. In some cases, an acrydite moiety can refer to an acrydite analogue generated from the reaction of acrydite with one or more species, such as, the reaction of acrydite with other monomers and cross-linkers during a polymerization reaction. Acrydite moieties may be modified to form chemical bonds with a species to be attached, such as a nucleic acid molecule (e.g., barcode sequence, barcoded nucleic acid molecule, barcoded oligonucleotide, primer, or other oligonucleotide). Acrydite moieties may be modified with thiol groups capable of forming a disulfide bond or may be modified with groups already comprising a disulfide bond. The thiol or disulfide (via disulfide exchange) may be used as an anchor point for a species to be attached or

another part of the acrydite moiety may be used for attachment. In some cases, attachment can be reversible, such that when the disulfide bond is broken (e.g., in the presence of a reducing agent), the attached species is released from the bead. In other cases, an acrydite moiety can comprise a reactive hydroxyl group that may be used for attachment.

[00276] Functionalization of beads for attachment of nucleic acid molecules (e.g., oligonucleotides) may be achieved through a wide range of different approaches, including activation of chemical groups within a polymer, incorporation of active or activatable functional groups in the polymer structure, or attachment at the pre-polymer or monomer stage in bead production.

[00277] For example, precursors (e.g., monomers, cross-linkers) that are polymerized to form a bead may comprise acrydite moieties, such that when a bead is generated, the bead also comprises acrydite moieties. The acrydite moieties can be attached to a nucleic acid molecule (e.g., oligonucleotide), which may include a priming sequence (e.g., a primer for amplifying target nucleic acids, random primer, primer sequence for messenger RNA) and/or one or more barcode sequences. The one more barcode sequences may include sequences that are the same for all nucleic acid molecules coupled to a bead and/or sequences that are different across all nucleic acid molecules coupled to the bead. The nucleic acid molecule may be incorporated into the bead.

[00278] In some cases, the nucleic acid molecule can comprise a functional sequence, for example, for attachment to a sequencing flow cell, such as, for example, a P5 sequence for Illumina® sequencing. In some cases, the nucleic acid molecule or derivative thereof (e.g., oligonucleotide or polynucleotide generated from the nucleic acid molecule) can comprise another functional sequence, such as, for example, a P7 sequence for attachment to a sequencing flow cell for Illumina sequencing. In some cases, the nucleic acid molecule can comprise a barcode sequence. In some cases, the primer can further comprise a unique molecular identifier (UMI). In some cases, the primer can comprise an R1 primer sequence for Illumina sequencing. In some cases, the primer can comprise an R2 primer sequence for Illumina sequencing. Examples of such nucleic acid molecules (e.g., oligonucleotides, polynucleotides, etc.) and uses thereof, as may be used with compositions, devices, methods and systems of the present disclosure, are provided in U.S. Patent Pub. Nos. 2014/0378345 and 2015/0376609, each of which is entirely incorporated herein by reference.

[00279] **FIG. 8** illustrates an example of a barcode carrying bead. A nucleic acid molecule **802**, such as an oligonucleotide, can be coupled to a bead **804** by a releasable linkage **806**, such as, for example, a disulfide linker. The same bead **804** may be coupled (e.g., via releasable

linkage) to one or more other nucleic acid molecules **818**, **820**. The nucleic acid molecule **802** may be or comprise a barcode. As noted elsewhere herein, the structure of the barcode may comprise a number of sequence elements. The nucleic acid molecule **802** may comprise a functional sequence **808** that may be used in subsequent processing. For example, the functional sequence **808** may include one or more of a sequencer specific flow cell attachment sequence (e.g., a P5 sequence for Illumina® sequencing systems) and a sequencing primer sequence (e.g., a R1 primer for Illumina® sequencing systems). The nucleic acid molecule **802** may comprise a barcode sequence **810** for use in barcoding the sample (e.g., DNA, RNA, protein, etc.). In some cases, the barcode sequence **810** can be bead-specific such that the barcode sequence **810** is common to all nucleic acid molecules (e.g., including nucleic acid molecule **802**) coupled to the same bead **804**. Alternatively or in addition, the barcode sequence **810** can be partition-specific such that the barcode sequence **810** is common to all nucleic acid molecules coupled to one or more beads that are partitioned into the same partition. The nucleic acid molecule **802** may comprise a specific priming sequence **812**, such as an mRNA specific priming sequence (e.g., poly-T sequence), a targeted priming sequence, and/or a random priming sequence. The nucleic acid molecule **802** may comprise an anchoring sequence **814** to ensure that the specific priming sequence **812** hybridizes at the sequence end (e.g., of the mRNA). For example, the anchoring sequence **814** can include a random short sequence of nucleotides, such as a 1-mer, 2-mer, 3-mer or longer sequence, which can ensure that a poly-T segment is more likely to hybridize at the sequence end of the poly-A tail of the mRNA.

[00280] The nucleic acid molecule **802** may comprise a unique molecular identifying sequence **816** (e.g., unique molecular identifier (UMI)). In some cases, the unique molecular identifying sequence **816** may comprise from about 5 to about 8 nucleotides. Alternatively, the unique molecular identifying sequence **816** may compress less than about 5 or more than about 8 nucleotides. The unique molecular identifying sequence **816** may be a unique sequence that varies across individual nucleic acid molecules (e.g., **802**, **818**, **820**, etc.) coupled to a single bead (e.g., bead **804**). In some cases, the unique molecular identifying sequence **816** may be a random sequence (e.g., such as a random N-mer sequence). For example, the UMI may provide a unique identifier of the starting mRNA molecule that was captured, in order to allow quantitation of the number of original expressed RNA. As will be appreciated, although **FIG. 8** shows three nucleic acid molecules **802**, **818**, **820** coupled to the surface of the bead **804**, an individual bead may be coupled to any number of individual nucleic acid molecules, for example, from one to tens to hundreds of thousands or even millions of individual nucleic acid molecules. The respective barcodes for the individual nucleic acid molecules can comprise both

common sequence segments or relatively common sequence segments (e.g., **808**, **810**, **812**, etc.) and variable or unique sequence segments (e.g., **816**) between different individual nucleic acid molecules coupled to the same bead.

[00281] In operation, a biological particle (e.g., cell, DNA, RNA, etc.) can be co-partitioned along with a barcode bearing bead **804**. The barcoded nucleic acid molecules **802**, **818**, **820** can be released from the bead **804** in the partition. By way of example, in the context of analyzing sample RNA, the poly-T segment (e.g., **812**) of one of the released nucleic acid molecules (e.g., **802**) can hybridize to the poly-A tail of a mRNA molecule. Reverse transcription may result in a cDNA transcript of the mRNA, but which transcript includes each of the sequence segments **808**, **810**, **816** of the nucleic acid molecule **802**. Because the nucleic acid molecule **802** comprises an anchoring sequence **814**, it will more likely hybridize to and prime reverse transcription at the sequence end of the poly-A tail of the mRNA. Within any given partition, all of the cDNA transcripts of the individual mRNA molecules may include a common barcode sequence segment **810**. However, the transcripts made from the different mRNA molecules within a partition may vary at the unique molecular identifying sequence **812** segment (e.g., UMI segment). Beneficially, even following any subsequent amplification of the contents of a partition, the number of different UMIs can be indicative of the quantity of mRNA originating from a given partition, and thus from the biological particle (e.g., cell). As noted above, the transcripts can be amplified, cleaned up and sequenced to identify the sequence of the cDNA transcript of the mRNA, as well as to sequence the barcode segment and the UMI segment. While a poly-T primer sequence is described, other targeted or random priming sequences may also be used in priming the reverse transcription reaction. Likewise, although described as releasing the barcoded oligonucleotides into the partition, in some cases, the nucleic acid molecules bound to the bead (e.g., gel bead) may be used to hybridize and capture the mRNA on the solid phase of the bead, for example, in order to facilitate the separation of the RNA from other cell contents.

[00282] In some cases, precursors comprising a functional group that is reactive or capable of being activated such that it becomes reactive can be polymerized with other precursors to generate gel beads comprising the activated or activatable functional group. The functional group may then be used to attach additional species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) to the gel beads. For example, some precursors comprising a carboxylic acid (COOH) group can co-polymerize with other precursors to form a gel bead that also comprises a COOH functional group. In some cases, acrylic acid (a species comprising free COOH groups), acrylamide, and bis(acryloyl)cystamine can be co-polymerized together to

generate a gel bead comprising free COOH groups. The COOH groups of the gel bead can be activated (e.g., via 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) and N-Hydroxysuccinimide (NHS) or 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM)) such that they are reactive (e.g., reactive to amine functional groups where EDC/NHS or DMTMM are used for activation). The activated COOH groups can then react with an appropriate species (e.g., a species comprising an amine functional group where the carboxylic acid groups are activated to be reactive with an amine functional group) comprising a moiety to be linked to the bead.

[00283] Beads comprising disulfide linkages in their polymeric network may be functionalized with additional species via reduction of some of the disulfide linkages to free thiols. The disulfide linkages may be reduced via, for example, the action of a reducing agent (e.g., DTT, TCEP, etc.) to generate free thiol groups, without dissolution of the bead. Free thiols of the beads can then react with free thiols of a species or a species comprising another disulfide bond (e.g., via thiol-disulfide exchange) such that the species can be linked to the beads (e.g., via a generated disulfide bond). In some cases, free thiols of the beads may react with any other suitable group. For example, free thiols of the beads may react with species comprising an acrydite moiety. The free thiol groups of the beads can react with the acrydite via Michael addition chemistry, such that the species comprising the acrydite is linked to the bead. In some cases, uncontrolled reactions can be prevented by inclusion of a thiol capping agent such as N-ethylmaleimide or iodoacetate.

[00284] Activation of disulfide linkages within a bead can be controlled such that only a small number of disulfide linkages are activated. Control may be exerted, for example, by controlling the concentration of a reducing agent used to generate free thiol groups and/or concentration of reagents used to form disulfide bonds in bead polymerization. In some cases, a low concentration (e.g., molecules of reducing agent:gel bead ratios of less than or equal to about 1:100,000,000,000, less than or equal to about 1:10,000,000,000, less than or equal to about 1:1,000,000,000, less than or equal to about 1:100,000,000, less than or equal to about 1:10,000,000, less than or equal to about 1:1,000,000, less than or equal to about 1:100,000, less than or equal to about 1:10,000) of reducing agent may be used for reduction. Controlling the number of disulfide linkages that are reduced to free thiols may be useful in ensuring bead structural integrity during functionalization. In some cases, optically-active agents, such as fluorescent dyes may be coupled to beads via free thiol groups of the beads and used to quantify the number of free thiols present in a bead and/or track a bead.

[00285] In some cases, addition of moieties to a gel bead after gel bead formation may be advantageous. For example, addition of an oligonucleotide (e.g., barcoded oligonucleotide) after gel bead formation may avoid loss of the species during chain transfer termination that can occur during polymerization. Moreover, smaller precursors (e.g., monomers or cross linkers that do not comprise side chain groups and linked moieties) may be used for polymerization and can be minimally hindered from growing chain ends due to viscous effects. In some cases, functionalization after gel bead synthesis can minimize exposure of species (e.g., oligonucleotides) to be loaded with potentially damaging agents (e.g., free radicals) and/or chemical environments. In some cases, the generated gel may possess an upper critical solution temperature (UCST) that can permit temperature driven swelling and collapse of a bead. Such functionality may aid in oligonucleotide (e.g., a primer) infiltration into the bead during subsequent functionalization of the bead with the oligonucleotide. Post-production functionalization may also be useful in controlling loading ratios of species in beads, such that, for example, the variability in loading ratio is minimized. Species loading may also be performed in a batch process such that a plurality of beads can be functionalized with the species in a single batch.

[00286] A bead injected or otherwise introduced into a partition may comprise releasably, cleavably, or reversibly attached barcodes. A bead injected or otherwise introduced into a partition may comprise activatable barcodes. A bead injected or otherwise introduced into a partition may be degradable, disruptable, or dissolvable beads.

[00287] Barcodes can be releasably, cleavably or reversibly attached to the beads such that barcodes can be released or be releasable through cleavage of a linkage between the barcode molecule and the bead, or released through degradation of the underlying bead itself, allowing the barcodes to be accessed or be accessible by other reagents, or both. In non-limiting examples, cleavage may be achieved through reduction of di-sulfide bonds, use of restriction enzymes, photo-activated cleavage, or cleavage via other types of stimuli (e.g., chemical, thermal, pH, enzymatic, etc.) and/or reactions, such as described elsewhere herein. Releasable barcodes may sometimes be referred to as being activatable, in that they are available for reaction once released. Thus, for example, an activatable barcode may be activated by releasing the barcode from a bead (or other suitable type of partition described herein). Other activatable configurations are also envisioned in the context of the described methods and systems.

[00288] In addition to, or as an alternative to the cleavable linkages between the beads and the associated molecules, such as barcode containing nucleic acid molecules (e.g., barcoded oligonucleotides), the beads may be degradable, disruptable, or dissolvable spontaneously or

upon exposure to one or more stimuli (e.g., temperature changes, pH changes, exposure to particular chemical species or phase, exposure to light, reducing agent, etc.). In some cases, a bead may be dissolvable, such that material components of the beads are solubilized when exposed to a particular chemical species or an environmental change, such as a change in temperature or a change in pH. In some cases, a gel bead can be degraded or dissolved at elevated temperature and/or in basic conditions. In some cases, a bead may be thermally degradable such that when the bead is exposed to an appropriate change in temperature (e.g., heat), the bead degrades. Degradation or dissolution of a bead bound to a species (e.g., a nucleic acid molecule, e.g., barcoded oligonucleotide) may result in release of the species from the bead.

[00289] As will be appreciated from the above disclosure, the degradation of a bead may refer to the disassociation of a bound or entrained species from a bead, both with and without structurally degrading the physical bead itself. For example, the degradation of the bead may involve cleavage of a cleavable linkage via one or more species and/or methods described elsewhere herein. In another example, entrained species may be released from beads through osmotic pressure differences due to, for example, changing chemical environments. By way of example, alteration of bead pore sizes due to osmotic pressure differences can generally occur without structural degradation of the bead itself. In some cases, an increase in pore size due to osmotic swelling of a bead can permit the release of entrained species within the bead. In other cases, osmotic shrinking of a bead may cause a bead to better retain an entrained species due to pore size contraction.

[00290] A degradable bead may be introduced into a partition, such as a droplet of an emulsion or a well, such that the bead degrades within the partition and any associated species (e.g., oligonucleotides) are released within the droplet when the appropriate stimulus is applied. The free species (e.g., oligonucleotides, nucleic acid molecules) may interact with other reagents contained in the partition. For example, a polyacrylamide bead comprising cystamine and linked, via a disulfide bond, to a barcode sequence, may be combined with a reducing agent within a droplet of a water-in-oil emulsion. Within the droplet, the reducing agent can break the various disulfide bonds, resulting in bead degradation and release of the barcode sequence into the aqueous, inner environment of the droplet. In another example, heating of a droplet comprising a bead-bound barcode sequence in basic solution may also result in bead degradation and release of the attached barcode sequence into the aqueous, inner environment of the droplet.

[00291] Any suitable number of molecular tag molecules (e.g., primer, barcoded oligonucleotide) can be associated with a bead such that, upon release from the bead, the molecular tag molecules (e.g., primer, e.g., barcoded oligonucleotide) are present in the partition

at a pre-defined concentration. Such pre-defined concentration may be selected to facilitate certain reactions for generating a sequencing library, e.g., amplification, within the partition. In some cases, the pre-defined concentration of the primer can be limited by the process of producing nucleic acid molecule (e.g., oligonucleotide) bearing beads.

[00292] In some cases, beads can be non-covalently loaded with one or more reagents. The beads can be non-covalently loaded by, for instance, subjecting the beads to conditions sufficient to swell the beads, allowing sufficient time for the reagents to diffuse into the interiors of the beads, and subjecting the beads to conditions sufficient to de-swell the beads. The swelling of the beads may be accomplished, for instance, by placing the beads in a thermodynamically favorable solvent, subjecting the beads to a higher or lower temperature, subjecting the beads to a higher or lower ion concentration, and/or subjecting the beads to an electric field. The swelling of the beads may be accomplished by various swelling methods. The de-swelling of the beads may be accomplished, for instance, by transferring the beads in a thermodynamically unfavorable solvent, subjecting the beads to lower or high temperatures, subjecting the beads to a lower or higher ion concentration, and/or removing an electric field. The de-swelling of the beads may be accomplished by various de-swelling methods. Transferring the beads may cause pores in the bead to shrink. The shrinking may then hinder reagents within the beads from diffusing out of the interiors of the beads. The hindrance may be due to steric interactions between the reagents and the interiors of the beads. The transfer may be accomplished microfluidically. For instance, the transfer may be achieved by moving the beads from one co-flowing solvent stream to a different co-flowing solvent stream. The swellability and/or pore size of the beads may be adjusted by changing the polymer composition of the bead.

[00293] In some cases, an acrydite moiety linked to a precursor, another species linked to a precursor, or a precursor itself can comprise a labile bond, such as chemically, thermally, or photo-sensitive bond e.g., disulfide bond, UV sensitive bond, or the like. Once acrydite moieties or other moieties comprising a labile bond are incorporated into a bead, the bead may also comprise the labile bond. The labile bond may be, for example, useful in reversibly linking (e.g., covalently linking) species (e.g., barcodes, primers, etc.) to a bead. In some cases, a thermally labile bond may include a nucleic acid hybridization based attachment, e.g., where an oligonucleotide is hybridized to a complementary sequence that is attached to the bead, such that thermal melting of the hybrid releases the oligonucleotide, e.g., a barcode containing sequence, from the bead or microcapsule.

[00294] The addition of multiple types of labile bonds to a gel bead may result in the generation of a bead capable of responding to varied stimuli. Each type of labile bond may be

sensitive to an associated stimulus (e.g., chemical stimulus, light, temperature, enzymatic, etc.) such that release of species attached to a bead via each labile bond may be controlled by the application of the appropriate stimulus. Such functionality may be useful in controlled release of species from a gel bead. In some cases, another species comprising a labile bond may be linked to a gel bead after gel bead formation via, for example, an activated functional group of the gel bead as described above. As will be appreciated, barcodes that are releasably, cleavably or reversibly attached to the beads described herein include barcodes that are released or releasable through cleavage of a linkage between the barcode molecule and the bead, or that are released through degradation of the underlying bead itself, allowing the barcodes to be accessed or accessible by other reagents, or both.

[00295] The barcodes that are releasable as described herein may sometimes be referred to as being activatable, in that they are available for reaction once released. Thus, for example, an activatable barcode may be activated by releasing the barcode from a bead (or other suitable type of partition described herein). Other activatable configurations are also envisioned in the context of the described methods and systems.

[00296] In addition to thermally cleavable bonds, disulfide bonds and UV sensitive bonds, other non-limiting examples of labile bonds that may be coupled to a precursor or bead include an ester linkage (e.g., cleavable with an acid, a base, or hydroxylamine), a vicinal diol linkage (e.g., cleavable via sodium periodate), a Diels-Alder linkage (e.g., cleavable via heat), a sulfone linkage (e.g., cleavable via a base), a silyl ether linkage (e.g., cleavable via an acid), a glycosidic linkage (e.g., cleavable via an amylase), a peptide linkage (e.g., cleavable via a protease), or a phosphodiester linkage (e.g., cleavable via a nuclease (e.g., DNAase)). A bond may be cleavable via other nucleic acid molecule targeting enzymes, such as restriction enzymes (e.g., restriction endonucleases), as described further below.

[00297] Species may be encapsulated in beads during bead generation (e.g., during polymerization of precursors). Such species may or may not participate in polymerization. Such species may be entered into polymerization reaction mixtures such that generated beads comprise the species upon bead formation. In some cases, such species may be added to the gel beads after formation. Such species may include, for example, nucleic acid molecules (e.g., oligonucleotides), reagents for a nucleic acid amplification reaction (e.g., primers, polymerases, dNTPs, co-factors (e.g., ionic co-factors), buffers) including those described herein, reagents for enzymatic reactions (e.g., enzymes, co-factors, substrates, buffers), reagents for nucleic acid modification reactions such as polymerization, ligation, or digestion, and/or reagents for template preparation (e.g., tagmentation) for one or more sequencing platforms (e.g., Nextera®).

for Illumina®). Such species may include one or more enzymes described herein, including without limitation, polymerase, reverse transcriptase, restriction enzymes (e.g., endonuclease), transposase, ligase, proteinase K, DNase, etc. Such species may include one or more reagents described elsewhere herein (e.g., lysis agents, inhibitors, inactivating agents, chelating agents, stimulus). Trapping of such species may be controlled by the polymer network density generated during polymerization of precursors, control of ionic charge within the gel bead (e.g., via ionic species linked to polymerized species), or by the release of other species. Encapsulated species may be released from a bead upon bead degradation and/or by application of a stimulus capable of releasing the species from the bead. Alternatively or in addition, species may be partitioned in a partition (e.g., droplet) during or subsequent to partition formation. Such species may include, without limitation, the abovementioned species that may also be encapsulated in a bead.

[00298] A degradable bead may comprise one or more species with a labile bond such that, when the bead/species is exposed to the appropriate stimuli, the bond is broken and the bead degrades. The labile bond may be a chemical bond (e.g., covalent bond, ionic bond) or may be another type of physical interaction (e.g., van der Waals interactions, dipole-dipole interactions, etc.). In some cases, a crosslinker used to generate a bead may comprise a labile bond. Upon exposure to the appropriate conditions, the labile bond can be broken and the bead degraded. For example, upon exposure of a polyacrylamide gel bead comprising cystamine crosslinkers to a reducing agent, the disulfide bonds of the cystamine can be broken and the bead degraded.

[00299] A degradable bead may be useful in more quickly releasing an attached species (e.g., a nucleic acid molecule, a barcode sequence, a primer, etc) from the bead when the appropriate stimulus is applied to the bead as compared to a bead that does not degrade. For example, for a species bound to an inner surface of a porous bead or in the case of an encapsulated species, the species may have greater mobility and accessibility to other species in solution upon degradation of the bead. In some cases, a species may also be attached to a degradable bead via a degradable linker (e.g., disulfide linker). The degradable linker may respond to the same stimuli as the degradable bead or the two degradable species may respond to different stimuli. For example, a barcode sequence may be attached, via a disulfide bond, to a polyacrylamide bead comprising cystamine. Upon exposure of the barcoded-bead to a reducing agent, the bead degrades and the barcode sequence is released upon breakage of both the disulfide linkage between the barcode sequence and the bead and the disulfide linkages of the cystamine in the bead.

[00300] As will be appreciated from the above disclosure, while referred to as degradation of a bead, in many instances as noted above, that degradation may refer to the disassociation of a

bound or entrained species from a bead, both with and without structurally degrading the physical bead itself. For example, entrained species may be released from beads through osmotic pressure differences due to, for example, changing chemical environments. By way of example, alteration of bead pore sizes due to osmotic pressure differences can generally occur without structural degradation of the bead itself. In some cases, an increase in pore size due to osmotic swelling of a bead can permit the release of entrained species within the bead. In other cases, osmotic shrinking of a bead may cause a bead to better retain an entrained species due to pore size contraction.

[00301] Where degradable beads are provided, it may be beneficial to avoid exposing such beads to the stimulus or stimuli that cause such degradation prior to a given time, in order to, for example, avoid premature bead degradation and issues that arise from such degradation, including for example poor flow characteristics and aggregation. By way of example, where beads comprise reducible cross-linking groups, such as disulfide groups, it will be desirable to avoid contacting such beads with reducing agents, e.g., DTT or other disulfide cleaving reagents. In such cases, treatment to the beads described herein will, in some cases be provided free of reducing agents, such as DTT. Because reducing agents are often provided in commercial enzyme preparations, it may be desirable to provide reducing agent free (or DTT free) enzyme preparations in treating the beads described herein. Examples of such enzymes include, e.g., polymerase enzyme preparations, reverse transcriptase enzyme preparations, ligase enzyme preparations, as well as many other enzyme preparations that may be used to treat the beads described herein. The terms “reducing agent free” or “DTT free” preparations can refer to a preparation having less than about 1/10th, less than about 1/50th, or even less than about 1/100th of the lower ranges for such materials used in degrading the beads. For example, for DTT, the reducing agent free preparation can have less than about 0.01 millimolar (mM), 0.005 mM, 0.001 mM DTT, 0.0005 mM DTT, or even less than about 0.0001 mM DTT. In many cases, the amount of DTT can be undetectable.

[00302] Numerous chemical triggers may be used to trigger the degradation of beads. Examples of these chemical changes may include, but are not limited to pH-mediated changes to the integrity of a component within the bead, degradation of a component of a bead via cleavage of cross-linked bonds, and depolymerization of a component of a bead.

[00303] A bead may be formed from materials that comprise degradable chemical crosslinkers, such as BAC or cystamine. Degradation of such degradable crosslinkers may be accomplished through a number of mechanisms. In some examples, a bead may be contacted with a chemical degrading agent that may induce oxidation, reduction or other chemical changes.

For example, a chemical degrading agent may be a reducing agent, such as dithiothreitol (DTT). Additional examples of reducing agents may include β -mercaptoethanol, (2S)-2-amino-1,4-dimercaptobutane (dithiobutylamine or DTBA), tris(2-carboxyethyl) phosphine (TCEP), or combinations thereof. A reducing agent may degrade the disulfide bonds formed between gel precursors forming the bead, and thus, degrade the bead. In other cases, a change in pH of a solution, such as an increase in pH, may trigger degradation of a bead. In other cases, exposure to an aqueous solution, such as water, may trigger hydrolytic degradation, and thus degradation of the bead. In some cases, any combination of stimuli may trigger degradation of a bead. For example, a change in pH may enable a chemical agent (e.g., DTT) to become an effective reducing agent.

[00304] Beads may also be induced to release their contents upon the application of a thermal stimulus. A change in temperature can cause a variety of changes to a bead. For example, heat can cause a solid bead to liquefy. A change in heat may cause melting of a bead such that a portion of the bead degrades. In other cases, heat may increase the internal pressure of the bead components such that the bead ruptures or explodes. Heat may also act upon heat-sensitive polymers used as materials to construct beads.

[00305] Any suitable agent may degrade beads. For example, changes in temperature or pH may be used to degrade thermo-sensitive or pH-sensitive bonds within beads. Alternatively or in addition to, chemical degrading agents may be used to degrade chemical bonds within beads by oxidation, reduction or other chemical changes. For example, a chemical degrading agent may be a reducing agent, such as DTT, wherein DTT may degrade the disulfide bonds formed between a crosslinker and gel precursors, thus degrading the bead. A reducing agent may be added to degrade the bead, which may or may not cause the bead to release its contents. Examples of reducing agents may include dithiothreitol (DTT), β -mercaptoethanol, (2S)-2-amino-1,4-dimercaptobutane (dithiobutylamine or DTBA), tris(2-carboxyethyl) phosphine (TCEP), or combinations thereof. The reducing agent may be present at a concentration of about 0.1mM, 0.5mM, 1mM, 5mM, 10mM. The reducing agent may be present at a concentration of at least about 0.1mM, 0.5mM, 1mM, 5mM, 10mM, or greater than 10 mM. The reducing agent may be present at concentration of at most about 10mM, 5mM, 1mM, 0.5mM, 0.1mM, or less.

[00306] Any suitable number of molecular tag molecules (e.g., primer, barcoded oligonucleotide) can be associated with a bead such that, upon release from the bead, the molecular tag molecules (e.g., primer, e.g., barcoded oligonucleotide) are present in the partition at a pre-defined concentration. Such pre-defined concentration may be selected to facilitate certain reactions for generating a sequencing library, e.g., amplification, within the partition. In

some cases, the pre-defined concentration of the primer can be limited by the process of producing oligonucleotide bearing beads.

[00307] Although **FIG. 1** and **FIG. 2** have been described in terms of providing substantially singly occupied partitions, above, in certain cases, it may be desirable to provide multiply occupied partitions, e.g., containing two, three, four or more cells and/or microcapsules (e.g., beads) comprising barcoded nucleic acid molecules (e.g., oligonucleotides) within a single partition. Accordingly, as noted above, the flow characteristics of the biological particle and/or bead containing fluids and partitioning fluids may be controlled to provide for such multiply occupied partitions. In particular, the flow parameters may be controlled to provide a occupancy rate at greater than about 50% of the partitions, greater than about 75%, and in some cases greater than about 80%, 90%, 95%, or higher.

[00308] In some cases, additional microcapsules can be used to deliver additional reagents to a partition. In such cases, it may be advantageous to introduce different beads into a common channel or droplet generation junction, from different bead sources (e.g., containing different associated reagents) through different channel inlets into such common channel or droplet generation junction (e.g., junction **210**). In such cases, the flow and frequency of the different beads into the channel or junction may be controlled to provide for a certain ratio of microcapsules from each source, while ensuring a pairing or combination of such beads into a partition with a given number of biological particles (e.g., one biological particle and one bead per partition).

[00309] The partitions described herein may comprise small volumes, for example, less than about 10 microliters (μL), 5 μL , 1 μL , 900 picoliters (pL), 800 pL, 700 pL, 600 pL, 500 pL, 400pL, 300 pL, 200 pL, 100pL, 50 pL, 20 pL, 10 pL, 1 pL, 500 nanoliters (nL), 100 nL, 50 nL, or less.

[00310] For example, in the case of droplet based partitions, the droplets may have overall volumes that are less than about 1000 pL, 900 pL, 800 pL, 700 pL, 600 pL, 500 pL, 400pL, 300 pL, 200 pL, 100pL, 50 pL, 20 pL, 10 pL, 1 pL, or less. Where co-partitioned with microcapsules, it will be appreciated that the sample fluid volume, e.g., including co-partitioned biological particles and/or beads, within the partitions may be less than about 90% of the above described volumes, less than about 80%, less than about 70%, less than about 60%, less than about 50%, less than about 40%, less than about 30%, less than about 20%, or less than about 10% of the above described volumes.

[00311] As is described elsewhere herein, partitioning species may generate a population or plurality of partitions. In such cases, any suitable number of partitions can be generated or

otherwise provided. For example, at least about 1,000 partitions, at least about 5,000 partitions, at least about 10,000 partitions, at least about 50,000 partitions, at least about 100,000 partitions, at least about 500,000 partitions, at least about 1,000,000 partitions, at least about 5,000,000 partitions at least about 10,000,000 partitions, at least about 50,000,000 partitions, at least about 100,000,000 partitions, at least about 500,000,000 partitions, at least about 1,000,000,000 partitions, or more partitions can be generated or otherwise provided. Moreover, the plurality of partitions may comprise both unoccupied partitions (e.g., empty partitions) and occupied partitions.

Reagents

[00312] In accordance with certain aspects, biological particles may be partitioned along with lysis reagents in order to release the contents of the biological particles within the partition. In such cases, the lysis agents can be contacted with the biological particle suspension concurrently with, or immediately prior to, the introduction of the biological particles into the partitioning junction/droplet generation zone (e.g., junction **210**), such as through an additional channel or channels upstream of the channel junction. In accordance with other aspects, additionally or alternatively, biological particles may be partitioned along with other reagents, as will be described further below.

[00313] **FIG. 3** shows an example of a microfluidic channel structure **300** for co-partitioning biological particles and reagents. The channel structure **300** can include channel segments **301**, **302**, **304**, **306** and **308**. Channel segments **301** and **302** communicate at a first channel junction **309**. Channel segments **302**, **304**, **306**, and **308** communicate at a second channel junction **310**.

[00314] In an example operation, the channel segment **301** may transport an aqueous fluid **312** that includes a plurality of biological particles **314** along the channel segment **301** into the second junction **310**. As an alternative or in addition to, channel segment **301** may transport beads (e.g., gel beads). The beads may comprise barcode molecules.

[00315] For example, the channel segment **301** may be connected to a reservoir comprising an aqueous suspension of biological particles **314**. Upstream of, and immediately prior to reaching, the second junction **310**, the channel segment **301** may meet the channel segment **302** at the first junction **309**. The channel segment **302** may transport a plurality of reagents **315** (e.g., lysis agents) suspended in the aqueous fluid **312** along the channel segment **302** into the first junction **309**. For example, the channel segment **302** may be connected to a reservoir comprising the reagents **315**. After the first junction **309**, the aqueous fluid **312** in the channel segment **301** can carry both the biological particles **314** and the reagents **315** towards the second junction **310**. In

some instances, the aqueous fluid **312** in the channel segment **301** can include one or more reagents, which can be the same or different reagents as the reagents **315**. A second fluid **316** that is immiscible with the aqueous fluid **312** (e.g., oil) can be delivered to the second junction **310** from each of channel segments **304** and **306**. Upon meeting of the aqueous fluid **312** from the channel segment **301** and the second fluid **316** from each of channel segments **304** and **306** at the second channel junction **310**, the aqueous fluid **312** can be partitioned as discrete droplets **318** in the second fluid **316** and flow away from the second junction **310** along channel segment **308**. The channel segment **308** may deliver the discrete droplets **318** to an outlet reservoir fluidly coupled to the channel segment **308**, where they may be harvested.

[00316] The second fluid **316** can comprise an oil, such as a fluorinated oil, that includes a fluorosurfactant for stabilizing the resulting droplets, for example, inhibiting subsequent coalescence of the resulting droplets **318**.

[00317] A discrete droplet generated may include an individual biological particle **314** and/or one or more reagents **315**. In some instances, a discrete droplet generated may include a barcode carrying bead (not shown), such as via other microfluidics structures described elsewhere herein. In some instances, a discrete droplet may be unoccupied (e.g., no reagents, no biological particles).

[00318] Beneficially, when lysis reagents and biological particles are co-partitioned, the lysis reagents can facilitate the release of the contents of the biological particles within the partition. The contents released in a partition may remain discrete from the contents of other partitions.

[00319] As will be appreciated, the channel segments described herein may be coupled to any of a variety of different fluid sources or receiving components, including reservoirs, tubing, manifolds, or fluidic components of other systems. As will be appreciated, the microfluidic channel structure **300** may have other geometries. For example, a microfluidic channel structure can have more than two channel junctions. For example, a microfluidic channel structure can have 2, 3, 4, 5 channel segments or more each carrying the same or different types of beads, reagents, and/or biological particles that meet at a channel junction. Fluid flow in each channel segment may be controlled to control the partitioning of the different elements into droplets. Fluid may be directed flow along one or more channels or reservoirs via one or more fluid flow units. A fluid flow unit can comprise compressors (e.g., providing positive pressure), pumps (e.g., providing negative pressure), actuators, and the like to control flow of the fluid. Fluid may also or otherwise be controlled via applied pressure differentials, centrifugal force, electrokinetic pumping, vacuum, capillary or gravity flow, or the like.

[00320] Examples of lysis agents include bioactive reagents, such as lysis enzymes that are used for lysis of different cell types, e.g., gram positive or negative bacteria, plants, yeast, mammalian, etc., such as lysozymes, achromopeptidase, lysostaphin, labiase, kitalase, lyticase, and a variety of other lysis enzymes available from, e.g., Sigma-Aldrich, Inc. (St Louis, MO), as well as other commercially available lysis enzymes. Other lysis agents may additionally or alternatively be co-partitioned with the biological particles to cause the release of the biological particle's contents into the partitions. For example, in some cases, surfactant-based lysis solutions may be used to lyse cells, although these may be less desirable for emulsion based systems where the surfactants can interfere with stable emulsions. In some cases, lysis solutions may include non-ionic surfactants such as, for example, TritonX-100 and Tween 20. In some cases, lysis solutions may include ionic surfactants such as, for example, sarcosyl and sodium dodecyl sulfate (SDS). Electroporation, thermal, acoustic or mechanical cellular disruption may also be used in certain cases, e.g., non-emulsion based partitioning such as encapsulation of biological particles that may be in addition to or in place of droplet partitioning, where any pore size of the encapsulate is sufficiently small to retain nucleic acid fragments of a given size, following cellular disruption.

[00321] Alternatively or in addition to the lysis agents co-partitioned with the biological particles described above, other reagents can also be co-partitioned with the biological particles, including, for example, DNase and RNase inactivating agents or inhibitors, such as proteinase K, chelating agents, such as EDTA, and other reagents employed in removing or otherwise reducing negative activity or impact of different cell lysate components on subsequent processing of nucleic acids. In addition, in the case of encapsulated biological particles, the biological particles may be exposed to an appropriate stimulus to release the biological particles or their contents from a co-partitioned microcapsule. For example, in some cases, a chemical stimulus may be co-partitioned along with an encapsulated biological particle to allow for the degradation of the microcapsule and release of the cell or its contents into the larger partition. In some cases, this stimulus may be the same as the stimulus described elsewhere herein for release of nucleic acid molecules (e.g., oligonucleotides) from their respective microcapsule (e.g., bead). In alternative aspects, this may be a different and non-overlapping stimulus, in order to allow an encapsulated biological particle to be released into a partition at a different time from the release of nucleic acid molecules into the same partition.

[00322] Additional reagents may also be co-partitioned with the biological particles, such as endonucleases to fragment a biological particle's DNA, DNA polymerase enzymes and dNTPs used to amplify the biological particle's nucleic acid fragments and to attach the barcode

molecular tags to the amplified fragments. Other enzymes may be co-partitioned, including without limitation, polymerase, transposase, ligase, proteinase K, DNase, etc. Additional reagents may also include reverse transcriptase enzymes, including enzymes with terminal transferase activity, primers and oligonucleotides, and switch oligonucleotides (also referred to herein as “switch oligos” or “template switching oligonucleotides”) which can be used for template switching. In some cases, template switching can be used to increase the length of a cDNA. In some cases, template switching oligonucleotides can be used to append a predefined nucleic acid sequence to the cDNA. In an example of template switching, cDNA can be generated from reverse transcription of a template, e.g., cellular mRNA, where a reverse transcriptase with terminal transferase activity can add additional nucleotides, e.g., polyC, to the cDNA in a template independent manner. Switch oligos can include sequences complementary to the additional nucleotides, e.g., polyG. The additional nucleotides (e.g., polyC) on the cDNA can hybridize to the additional nucleotides (e.g., polyG) on the switch oligo, whereby the switch oligo can be used by the reverse transcriptase as template to further extend the cDNA. Template switching oligonucleotides may comprise a hybridization region and a template region. The hybridization region can comprise any sequence capable of hybridizing to the target. In some cases, as previously described, the hybridization region comprises a series of G bases to complement the overhanging C bases at the 3' end of a cDNA molecule. The series of G bases may comprise 1 G base, 2 G bases, 3 G bases, 4 G bases, 5 G bases or more than 5 G bases. The template sequence can comprise any sequence to be incorporated into the cDNA. In some cases, the template region comprises at least 1 (e.g., at least 2, 3, 4, 5 or more) tag sequences and/or functional sequences. Switch oligos may comprise deoxyribonucleic acids; ribonucleic acids; modified nucleic acids including 2-Aminopurine, 2,6-Diaminopurine (2-Amino-dA), inverted dT, 5-Methyl dC, 2'-deoxyInosine, Super T (5-hydroxybutynl-2'-deoxyuridine), Super G (8-aza-7-deazaguanosine), locked nucleic acids (LNAs), unlocked nucleic acids (UNAs, e.g., UNA-A, UNA-U, UNA-C, UNA-G), Iso-dG, Iso-dC, 2' Fluoro bases (e.g., Fluoro C, Fluoro U, Fluoro A, and Fluoro G), or any combination.

[00323] In some cases, the length of a switch oligo may be at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147,

148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249 or 250 nucleotides or longer.

[00324] In some cases, the length of a switch oligo may be at most about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249 or 250 nucleotides.

[00325] Once the contents of the cells are released into their respective partitions, the macromolecular components (e.g., macromolecular constituents of biological particles, such as RNA, DNA, or proteins) contained therein may be further processed within the partitions. In accordance with the methods and systems described herein, the macromolecular component contents of individual biological particles can be provided with unique identifiers such that, upon characterization of those macromolecular components they may be attributed as having been derived from the same biological particle or particles. The ability to attribute characteristics to individual biological particles or groups of biological particles is provided by the assignment of unique identifiers specifically to an individual biological particle or groups of biological particles. Unique identifiers, e.g., in the form of nucleic acid barcodes can be assigned or associated with individual biological particles or populations of biological particles, in order to tag or label the biological particle's macromolecular components (and as a result, its characteristics) with the unique identifiers. These unique identifiers can then be used to attribute the biological particle's components and characteristics to an individual biological particle or group of biological particles.

[00326] In some aspects, this is performed by co-partitioning the individual biological particle or groups of biological particles with the unique identifiers, such as described above (with reference to **FIG. 2**). In some aspects, the unique identifiers are provided in the form of nucleic acid molecules (e.g., oligonucleotides) that comprise nucleic acid barcode sequences that may be attached to or otherwise associated with the nucleic acid contents of individual biological particle, or to other components of the biological particle, and particularly to fragments of those nucleic acids. The nucleic acid molecules are partitioned such that as between nucleic acid molecules in a partition, the nucleic acid barcode sequences contained therein are the same, but as between different partitions, the nucleic acid molecule can, and do have differing barcode sequences, or at least represent a large number of different barcode sequences across all of the partitions in a analysis. In some aspects, only one nucleic acid barcode sequence can be associated with a partition, although in some cases, two or more different barcode sequences may be present.

[00327] The nucleic acid barcode sequences can include from about 6 to about 20 or more nucleotides within the sequence of the nucleic acid molecules (e.g., oligonucleotides). The nucleic acid barcode sequences can include from about 6 to about 20, 30, 40, 50, 60, 70, 80, 90, 100 or more nucleotides. In some cases, the length of a barcode sequence may be about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or longer. In some cases, the length of a barcode sequence may be at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or longer. In some cases, the length of a barcode sequence may be at most about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or shorter. These nucleotides may be completely contiguous, i.e., in a single stretch of adjacent nucleotides, or they may be separated into two or more separate subsequences that are separated by 1 or more nucleotides. In some cases, separated barcode subsequences can be from about 4 to about 16 nucleotides in length. In some cases, the barcode subsequence may be about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or longer. In some cases, the barcode subsequence may be at least about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or longer. In some cases, the barcode subsequence may be at most about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or shorter.

[00328] The co-partitioned nucleic acid molecules can also comprise other functional sequences useful in the processing of the nucleic acids from the co-partitioned biological particles. These sequences include, e.g., targeted or random/universal amplification primer sequences for amplifying the genomic DNA from the individual biological particles within the partitions while attaching the associated barcode sequences, sequencing primers or primer recognition sites, hybridization or probing sequences, e.g., for identification of presence of the

sequences or for pulling down barcoded nucleic acids, or any of a number of other potential functional sequences. Other mechanisms of co-partitioning oligonucleotides may also be employed, including, e.g., coalescence of two or more droplets, where one droplet contains oligonucleotides, or microdispensing of oligonucleotides into partitions, e.g., droplets within microfluidic systems.

[00329] In an example, microcapsules, such as beads, are provided that each include large numbers of the above described barcoded nucleic acid molecules (e.g., barcoded oligonucleotides) releasably attached to the beads, where all of the nucleic acid molecules attached to a particular bead will include the same nucleic acid barcode sequence, but where a large number of diverse barcode sequences are represented across the population of beads used. Hydrogel beads, e.g., comprising polyacrylamide polymer matrices, may be used as a solid support and delivery vehicle for the nucleic acid molecules into the partitions, as they are capable of carrying large numbers of nucleic acid molecules, and may be configured to release those nucleic acid molecules upon exposure to a particular stimulus, as described elsewhere herein. In some cases, the population of beads provides a diverse barcode sequence library that includes at least about 1,000 different barcode sequences, at least about 5,000 different barcode sequences, at least about 10,000 different barcode sequences, at least about 50,000 different barcode sequences, at least about 100,000 different barcode sequences, at least about 1,000,000 different barcode sequences, at least about 5,000,000 different barcode sequences, or at least about 10,000,000 different barcode sequences, or more. Additionally, each bead can be provided with large numbers of nucleic acid (e.g., oligonucleotide) molecules attached. In particular, the number of molecules of nucleic acid molecules including the barcode sequence on an individual bead can be at least about 1,000 nucleic acid molecules, at least about 5,000 nucleic acid molecules, at least about 10,000 nucleic acid molecules, at least about 50,000 nucleic acid molecules, at least about 100,000 nucleic acid molecules, at least about 500,000 nucleic acids, at least about 1,000,000 nucleic acid molecules, at least about 5,000,000 nucleic acid molecules, at least about 10,000,000 nucleic acid molecules, at least about 50,000,000 nucleic acid molecules, at least about 100,000,000 nucleic acid molecules, at least about 250,000,000 nucleic acid molecules and in some cases at least about 1 billion nucleic acid molecules, or more. Nucleic acid molecules of a bead can include identical (or common) barcode sequences, different barcode sequences, or a combination of both. Nucleic acid molecules of a bead can include multiple sets of nucleic acid molecules. Nucleic acid molecules of a set can include identical barcode sequences. The identical barcode sequences can be different from barcode sequences of nucleic acid molecules of another set.

[00330] Moreover, when the population of beads is partitioned, the resulting population of partitions can also include a diverse barcode library that includes at least about 1,000 different barcode sequences, at least about 5,000 different barcode sequences, at least about 10,000 different barcode sequences, at least at least about 50,000 different barcode sequences, at least about 100,000 different barcode sequences, at least about 1,000,000 different barcode sequences, at least about 5,000,000 different barcode sequences, or at least about 10,000,000 different barcode sequences. Additionally, each partition of the population can include at least about 1,000 nucleic acid molecules, at least about 5,000 nucleic acid molecules, at least about 10,000 nucleic acid molecules, at least about 50,000 nucleic acid molecules, at least about 100,000 nucleic acid molecules, at least about 500,000 nucleic acids, at least about 1,000,000 nucleic acid molecules, at least about 5,000,000 nucleic acid molecules, at least about 10,000,000 nucleic acid molecules, at least about 50,000,000 nucleic acid molecules, at least about 100,000,000 nucleic acid molecules, at least about 250,000,000 nucleic acid molecules and in some cases at least about 1 billion nucleic acid molecules.

[00331] In some cases, it may be desirable to incorporate multiple different barcodes within a partition, either attached to a single or multiple beads within the partition. For example, in some cases, a mixed, but known set of barcode sequences may provide greater assurance of identification in the subsequent processing, e.g., by providing a stronger address or attribution of the barcodes to a partition, as a duplicate or independent confirmation of the output from a partition.

[00332] The nucleic acid molecules (e.g., oligonucleotides) are releasable from the beads upon the application of a particular stimulus to the beads. In some cases, the stimulus may be a photo-stimulus, e.g., through cleavage of a photo-labile linkage that releases the nucleic acid molecules. In other cases, a thermal stimulus may be used, where elevation of the temperature of the beads environment will result in cleavage of a linkage or other release of the nucleic acid molecules from the beads. In still other cases, a chemical stimulus can be used that cleaves a linkage of the nucleic acid molecules to the beads, or otherwise results in release of the nucleic acid molecules from the beads. In one case, such compositions include the polyacrylamide matrices described above for encapsulation of biological particles, and may be degraded for release of the attached nucleic acid molecules through exposure to a reducing agent, such as DTT.

[00333] In some aspects, provided are systems and methods for controlled partitioning. Droplet size may be controlled by adjusting certain geometric features in channel architecture

(e.g., microfluidics channel architecture). For example, an expansion angle, width, and/or length of a channel may be adjusted to control droplet size.

[00334] FIG. 4 shows an example of a microfluidic channel structure for the controlled partitioning of beads into discrete droplets. A channel structure **400** can include a channel segment **402** communicating at a channel junction **406** (or intersection) with a reservoir **404**. The reservoir **404** can be a chamber. Any reference to “reservoir,” as used herein, can also refer to a “chamber.” In operation, an aqueous fluid **408** that includes suspended beads **412** may be transported along the channel segment **402** into the junction **406** to meet a second fluid **410** that is immiscible with the aqueous fluid **408** in the reservoir **404** to create droplets **416**, **418** of the aqueous fluid **408** flowing into the reservoir **404**. At the junction **406** where the aqueous fluid **408** and the second fluid **410** meet, droplets can form based on factors such as the hydrodynamic forces at the junction **406**, flow rates of the two fluids **408**, **410**, fluid properties, and certain geometric parameters (e.g., w , h_0 , α , etc.) of the channel structure **400**. A plurality of droplets can be collected in the reservoir **404** by continuously injecting the aqueous fluid **408** from the channel segment **402** through the junction **406**.

[00335] A discrete droplet generated may include a bead (e.g., as in occupied droplets **416**). Alternatively, a discrete droplet generated may include more than one bead. Alternatively, a discrete droplet generated may not include any beads (e.g., as in unoccupied droplet **418**). In some instances, a discrete droplet generated may contain one or more biological particles, as described elsewhere herein. In some instances, a discrete droplet generated may comprise one or more reagents, as described elsewhere herein.

[00336] In some instances, the aqueous fluid **408** can have a substantially uniform concentration or frequency of beads **412**. The beads **412** can be introduced into the channel segment **402** from a separate channel (not shown in FIG. 4). The frequency of beads **412** in the channel segment **402** may be controlled by controlling the frequency in which the beads **412** are introduced into the channel segment **402** and/or the relative flow rates of the fluids in the channel segment **402** and the separate channel. In some instances, the beads can be introduced into the channel segment **402** from a plurality of different channels, and the frequency controlled accordingly.

[00337] In some instances, the aqueous fluid **408** in the channel segment **402** can comprise biological particles (e.g., described with reference to FIGS. 1 and 2). In some instances, the aqueous fluid **408** can have a substantially uniform concentration or frequency of biological particles. As with the beads, the biological particles can be introduced into the channel segment **402** from a separate channel. The frequency or concentration of the biological particles in the

aqueous fluid **408** in the channel segment **402** may be controlled by controlling the frequency in which the biological particles are introduced into the channel segment **402** and/or the relative flow rates of the fluids in the channel segment **402** and the separate channel. In some instances, the biological particles can be introduced into the channel segment **402** from a plurality of different channels, and the frequency controlled accordingly. In some instances, a first separate channel can introduce beads and a second separate channel can introduce biological particles into the channel segment **402**. The first separate channel introducing the beads may be upstream or downstream of the second separate channel introducing the biological particles.

[00338] The second fluid **410** can comprise an oil, such as a fluorinated oil, that includes a fluorosurfactant for stabilizing the resulting droplets, for example, inhibiting subsequent coalescence of the resulting droplets.

[00339] In some instances, the second fluid **410** may not be subjected to and/or directed to any flow in or out of the reservoir **404**. For example, the second fluid **410** may be substantially stationary in the reservoir **404**. In some instances, the second fluid **410** may be subjected to flow within the reservoir **404**, but not in or out of the reservoir **404**, such as via application of pressure to the reservoir **404** and/or as affected by the incoming flow of the aqueous fluid **408** at the junction **406**. Alternatively, the second fluid **410** may be subjected and/or directed to flow in or out of the reservoir **404**. For example, the reservoir **404** can be a channel directing the second fluid **410** from upstream to downstream, transporting the generated droplets.

[00340] The channel structure **400** at or near the junction **406** may have certain geometric features that at least partly determine the sizes of the droplets formed by the channel structure **400**. The channel segment **402** can have a height, h_0 and width, w , at or near the junction **406**. By way of example, the channel segment **402** can comprise a rectangular cross-section that leads to a reservoir **404** having a wider cross-section (such as in width or diameter). Alternatively, the cross-section of the channel segment **402** can be other shapes, such as a circular shape, trapezoidal shape, polygonal shape, or any other shapes. The top and bottom walls of the reservoir **404** at or near the junction **406** can be inclined at an expansion angle, α . The expansion angle, α , allows the tongue (portion of the aqueous fluid **408** leaving channel segment **402** at junction **406** and entering the reservoir **404** before droplet formation) to increase in depth and facilitate decrease in curvature of the intermediately formed droplet. Droplet size may decrease with increasing expansion angle. The resulting droplet radius, R_d , may be predicted by the following equation for the aforementioned geometric parameters of h_0 , w , and α :

$$R_d \approx 0.44 \left(1 + 2.2 \sqrt{\tan \alpha} \frac{w}{h_0} \right) \frac{h_0}{\sqrt{\tan \alpha}}$$

[00341] By way of example, for a channel structure with $w = 21 \mu\text{m}$, $h = 21 \mu\text{m}$, and $\alpha = 3^\circ$, the predicted droplet size is $121 \mu\text{m}$. In another example, for a channel structure with $w = 25 \mu\text{m}$, $h = 25 \mu\text{m}$, and $\alpha = 5^\circ$, the predicted droplet size is $123 \mu\text{m}$. In another example, for a channel structure with $w = 28 \mu\text{m}$, $h = 28 \mu\text{m}$, and $\alpha = 7^\circ$, the predicted droplet size is $124 \mu\text{m}$.

[00342] In some instances, the expansion angle, α , may be between a range of from about 0.5° to about 4° , from about 0.1° to about 10° , or from about 0° to about 90° . For example, the expansion angle can be at least about 0.01° , 0.1° , 0.2° , 0.3° , 0.4° , 0.5° , 0.6° , 0.7° , 0.8° , 0.9° , 1° , 2° , 3° , 4° , 5° , 6° , 7° , 8° , 9° , 10° , 15° , 20° , 25° , 30° , 35° , 40° , 45° , 50° , 55° , 60° , 65° , 70° , 75° , 80° , 85° , or higher. In some instances, the expansion angle can be at most about 89° , 88° , 87° , 86° , 85° , 84° , 83° , 82° , 81° , 80° , 75° , 70° , 65° , 60° , 55° , 50° , 45° , 40° , 35° , 30° , 25° , 20° , 15° , 10° , 9° , 8° , 7° , 6° , 5° , 4° , 3° , 2° , 1° , 0.1° , 0.01° , or less. In some instances, the width, w , can be between a range of from about 100 micrometers (μm) to about 500 μm . In some instances, the width, w , can be between a range of from about 10 μm to about 200 μm . Alternatively, the width can be less than about 10 μm . Alternatively, the width can be greater than about 500 μm . In some instances, the flow rate of the aqueous fluid **408** entering the junction **406** can be between about 0.04 microliters (μL)/minute (min) and about 40 $\mu\text{L}/\text{min}$. In some instances, the flow rate of the aqueous fluid **408** entering the junction **406** can be between about 0.01 microliters (μL)/minute (min) and about 100 $\mu\text{L}/\text{min}$. Alternatively, the flow rate of the aqueous fluid **408** entering the junction **406** can be less than about 0.01 $\mu\text{L}/\text{min}$. Alternatively, the flow rate of the aqueous fluid **408** entering the junction **406** can be greater than about 40 $\mu\text{L}/\text{min}$, such as 45 $\mu\text{L}/\text{min}$, 50 $\mu\text{L}/\text{min}$, 55 $\mu\text{L}/\text{min}$, 60 $\mu\text{L}/\text{min}$, 65 $\mu\text{L}/\text{min}$, 70 $\mu\text{L}/\text{min}$, 75 $\mu\text{L}/\text{min}$, 80 $\mu\text{L}/\text{min}$, 85 $\mu\text{L}/\text{min}$, 90 $\mu\text{L}/\text{min}$, 95 $\mu\text{L}/\text{min}$, 100 $\mu\text{L}/\text{min}$, 110 $\mu\text{L}/\text{min}$, 120 $\mu\text{L}/\text{min}$, 130 $\mu\text{L}/\text{min}$, 140 $\mu\text{L}/\text{min}$, 150 $\mu\text{L}/\text{min}$, or greater. At lower flow rates, such as flow rates of about less than or equal to 10 microliters/minute, the droplet radius may not be dependent on the flow rate of the aqueous fluid **408** entering the junction **406**.

[00343] In some instances, at least about 50% of the droplets generated can have uniform size. In some instances, at least about 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater of the droplets generated can have uniform size. Alternatively, less than about 50% of the droplets generated can have uniform size.

[00344] The throughput of droplet generation can be increased by increasing the points of generation, such as increasing the number of junctions (e.g., junction **406**) between aqueous fluid **408** channel segments (e.g., channel segment **402**) and the reservoir **404**. Alternatively or in addition, the throughput of droplet generation can be increased by increasing the flow rate of the aqueous fluid **408** in the channel segment **402**.

[00345] FIG. 5 shows an example of a microfluidic channel structure for increased droplet generation throughput. A microfluidic channel structure 500 can comprise a plurality of channel segments 502 and a reservoir 504. Each of the plurality of channel segments 502 may be in fluid communication with the reservoir 504. The channel structure 500 can comprise a plurality of channel junctions 506 between the plurality of channel segments 502 and the reservoir 504. Each channel junction can be a point of droplet generation. The channel segment 402 from the channel structure 400 in FIG. 4 and any description to the components thereof may correspond to a channel segment of the plurality of channel segments 502 in channel structure 500 and any description to the corresponding components thereof. The reservoir 404 from the channel structure 400 and any description to the components thereof may correspond to the reservoir 504 from the channel structure 500 and any description to the corresponding components thereof.

[00346] Each channel segment of the plurality of channel segments 502 may comprise an aqueous fluid 508 that includes suspended beads 512. The reservoir 504 may comprise a second fluid 510 that is immiscible with the aqueous fluid 508. In some instances, the second fluid 510 may not be subjected to and/or directed to any flow in or out of the reservoir 504. For example, the second fluid 510 may be substantially stationary in the reservoir 504. In some instances, the second fluid 510 may be subjected to flow within the reservoir 504, but not in or out of the reservoir 504, such as via application of pressure to the reservoir 504 and/or as affected by the incoming flow of the aqueous fluid 508 at the junctions. Alternatively, the second fluid 510 may be subjected and/or directed to flow in or out of the reservoir 504. For example, the reservoir 504 can be a channel directing the second fluid 510 from upstream to downstream, transporting the generated droplets.

[00347] In operation, the aqueous fluid 508 that includes suspended beads 512 may be transported along the plurality of channel segments 502 into the plurality of junctions 506 to meet the second fluid 510 in the reservoir 504 to create droplets 516, 518. A droplet may form from each channel segment at each corresponding junction with the reservoir 504. At the junction where the aqueous fluid 508 and the second fluid 510 meet, droplets can form based on factors such as the hydrodynamic forces at the junction, flow rates of the two fluids 508, 510, fluid properties, and certain geometric parameters (e.g., w , h_0 , α , etc.) of the channel structure 500, as described elsewhere herein. A plurality of droplets can be collected in the reservoir 504 by continuously injecting the aqueous fluid 508 from the plurality of channel segments 502 through the plurality of junctions 506. Throughput may significantly increase with the parallel channel configuration of channel structure 500. For example, a channel structure having five inlet channel segments comprising the aqueous fluid 508 may generate droplets five times as

frequently than a channel structure having one inlet channel segment, provided that the fluid flow rate in the channel segments are substantially the same. The fluid flow rate in the different inlet channel segments may or may not be substantially the same. A channel structure may have as many parallel channel segments as is practical and allowed for the size of the reservoir. For example, the channel structure may have at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 500, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1000, 1500, 5000 or more parallel or substantially parallel channel segments.

[00348] The geometric parameters, w , h_0 , and α , may or may not be uniform for each of the channel segments in the plurality of channel segments **502**. For example, each channel segment may have the same or different widths at or near its respective channel junction with the reservoir **504**. For example, each channel segment may have the same or different height at or near its respective channel junction with the reservoir **504**. In another example, the reservoir **504** may have the same or different expansion angle at the different channel junctions with the plurality of channel segments **502**. When the geometric parameters are uniform, beneficially, droplet size may also be controlled to be uniform even with the increased throughput. In some instances, when it is desirable to have a different distribution of droplet sizes, the geometric parameters for the plurality of channel segments **502** may be varied accordingly.

[00349] In some instances, at least about 50% of the droplets generated can have uniform size. In some instances, at least about 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater of the droplets generated can have uniform size. Alternatively, less than about 50% of the droplets generated can have uniform size.

[00350] **FIG. 6** shows another example of a microfluidic channel structure for increased droplet generation throughput. A microfluidic channel structure **600** can comprise a plurality of channel segments **602** arranged generally circularly around the perimeter of a reservoir **604**. Each of the plurality of channel segments **602** may be in fluid communication with the reservoir **604**. The channel structure **600** can comprise a plurality of channel junctions **606** between the plurality of channel segments **602** and the reservoir **604**. Each channel junction can be a point of droplet generation. The channel segment **402** from the channel structure **400** in **FIG. 4** and any description to the components thereof may correspond to a channel segment of the plurality of channel segments **602** in channel structure **600** and any description to the corresponding components thereof. The reservoir **404** from the channel structure **400** and any description to the components thereof may correspond to the reservoir **604** from the channel structure **600** and any description to the corresponding components thereof.

[00351] Each channel segment of the plurality of channel segments **602** may comprise an aqueous fluid **608** that includes suspended beads **612**. The reservoir **604** may comprise a second fluid **610** that is immiscible with the aqueous fluid **608**. In some instances, the second fluid **610** may not be subjected to and/or directed to any flow in or out of the reservoir **604**. For example, the second fluid **610** may be substantially stationary in the reservoir **604**. In some instances, the second fluid **610** may be subjected to flow within the reservoir **604**, but not in or out of the reservoir **604**, such as via application of pressure to the reservoir **604** and/or as affected by the incoming flow of the aqueous fluid **608** at the junctions. Alternatively, the second fluid **610** may be subjected and/or directed to flow in or out of the reservoir **604**. For example, the reservoir **604** can be a channel directing the second fluid **610** from upstream to downstream, transporting the generated droplets.

[00352] In operation, the aqueous fluid **608** that includes suspended beads **612** may be transported along the plurality of channel segments **602** into the plurality of junctions **606** to meet the second fluid **610** in the reservoir **604** to create a plurality of droplets **616**. A droplet may form from each channel segment at each corresponding junction with the reservoir **604**. At the junction where the aqueous fluid **608** and the second fluid **610** meet, droplets can form based on factors such as the hydrodynamic forces at the junction, flow rates of the two fluids **608**, **610**, fluid properties, and certain geometric parameters (e.g., widths and heights of the channel segments **602**, expansion angle of the reservoir **604**, etc.) of the channel structure **600**, as described elsewhere herein. A plurality of droplets can be collected in the reservoir **604** by continuously injecting the aqueous fluid **608** from the plurality of channel segments **602** through the plurality of junctions **606**. Throughput may significantly increase with the substantially parallel channel configuration of the channel structure **600**. A channel structure may have as many substantially parallel channel segments as is practical and allowed for by the size of the reservoir. For example, the channel structure may have at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1000, 1500, 5000 or more parallel or substantially parallel channel segments. The plurality of channel segments may be substantially evenly spaced apart, for example, around an edge or perimeter of the reservoir. Alternatively, the spacing of the plurality of channel segments may be uneven.

[00353] The reservoir **604** may have an expansion angle, α (not shown in **FIG. 6**) at or near each channel junction. Each channel segment of the plurality of channel segments **602** may have a width, w , and a height, h_0 , at or near the channel junction. The geometric parameters, w , h_0 , and α , may or may not be uniform for each of the channel segments in the plurality of channel segments **602**. For example, each channel segment may have the same or different widths at or

near its respective channel junction with the reservoir **604**. For example, each channel segment may have the same or different height at or near its respective channel junction with the reservoir **604**.

[00354] The reservoir **604** may have the same or different expansion angle at the different channel junctions with the plurality of channel segments **602**. For example, a circular reservoir (as shown in **FIG. 6**) may have a conical, dome-like, or hemispherical ceiling (e.g., top wall) to provide the same or substantially same expansion angle for each channel segments **602** at or near the plurality of channel junctions **606**. When the geometric parameters are uniform, beneficially, resulting droplet size may be controlled to be uniform even with the increased throughput. In some instances, when it is desirable to have a different distribution of droplet sizes, the geometric parameters for the plurality of channel segments **602** may be varied accordingly.

[00355] In some instances, at least about 50% of the droplets generated can have uniform size. In some instances, at least about 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater of the droplets generated can have uniform size. Alternatively, less than about 50% of the droplets generated can have uniform size. The beads and/or biological particle injected into the droplets may or may not have uniform size.

[00356] **FIG. 7A** shows a cross-section view of another example of a microfluidic channel structure with a geometric feature for controlled partitioning. A channel structure **700** can include a channel segment **702** communicating at a channel junction **706** (or intersection) with a reservoir **704**. In some instances, the channel structure **700** and one or more of its components can correspond to the channel structure **100** and one or more of its components. **FIG. 7B** shows a perspective view of the channel structure **700** of **FIG. 7A**.

[00357] An aqueous fluid **712** comprising a plurality of particles **716** may be transported along the channel segment **702** into the junction **706** to meet a second fluid **714** (e.g., oil, etc.) that is immiscible with the aqueous fluid **712** in the reservoir **704** to create droplets **720** of the aqueous fluid **712** flowing into the reservoir **704**. At the junction **706** where the aqueous fluid **712** and the second fluid **714** meet, droplets can form based on factors such as the hydrodynamic forces at the junction **706**, relative flow rates of the two fluids **712**, **714**, fluid properties, and certain geometric parameters (e.g., Δh , etc.) of the channel structure **700**. A plurality of droplets can be collected in the reservoir **704** by continuously injecting the aqueous fluid **712** from the channel segment **702** at the junction **706**.

[00358] A discrete droplet generated may comprise one or more particles of the plurality of particles **716**. As described elsewhere herein, a particle may be any particle, such as a bead, cell

bead, gel bead, biological particle, macromolecular constituents of biological particle, or other particles. Alternatively, a discrete droplet generated may not include any particles.

[00359] In some instances, the aqueous fluid **712** can have a substantially uniform concentration or frequency of particles **716**. As described elsewhere herein (e.g., with reference to **FIG. 4**), the particles **716** (e.g., beads) can be introduced into the channel segment **702** from a separate channel (not shown in **FIG. 7**). The frequency of particles **716** in the channel segment **702** may be controlled by controlling the frequency in which the particles **716** are introduced into the channel segment **702** and/or the relative flow rates of the fluids in the channel segment **702** and the separate channel. In some instances, the particles **716** can be introduced into the channel segment **702** from a plurality of different channels, and the frequency controlled accordingly. In some instances, different particles may be introduced via separate channels. For example, a first separate channel can introduce beads and a second separate channel can introduce biological particles into the channel segment **702**. The first separate channel introducing the beads may be upstream or downstream of the second separate channel introducing the biological particles.

[00360] In some instances, the second fluid **714** may not be subjected to and/or directed to any flow in or out of the reservoir **704**. For example, the second fluid **714** may be substantially stationary in the reservoir **704**. In some instances, the second fluid **714** may be subjected to flow within the reservoir **704**, but not in or out of the reservoir **704**, such as via application of pressure to the reservoir **704** and/or as affected by the incoming flow of the aqueous fluid **712** at the junction **706**. Alternatively, the second fluid **714** may be subjected and/or directed to flow in or out of the reservoir **704**. For example, the reservoir **704** can be a channel directing the second fluid **714** from upstream to downstream, transporting the generated droplets.

[00361] The channel structure **700** at or near the junction **706** may have certain geometric features that at least partly determine the sizes and/or shapes of the droplets formed by the channel structure **700**. The channel segment **702** can have a first cross-section height, h_1 , and the reservoir **704** can have a second cross-section height, h_2 . The first cross-section height, h_1 , and the second cross-section height, h_2 , may be different, such that at the junction **706**, there is a height difference of Δh . The second cross-section height, h_2 , may be greater than the first cross-section height, h_1 . In some instances, the reservoir may thereafter gradually increase in cross-section height, for example, the more distant it is from the junction **706**. In some instances, the cross-section height of the reservoir may increase in accordance with expansion angle, β , at or near the junction **706**. The height difference, Δh , and/or expansion angle, β , can allow the tongue (portion of the aqueous fluid **712** leaving channel segment **702** at junction **706** and

entering the reservoir **704** before droplet formation) to increase in depth and facilitate decrease in curvature of the intermediately formed droplet. For example, droplet size may decrease with increasing height difference and/or increasing expansion angle.

[00362] The height difference, Δh , can be at least about 1 μm . Alternatively, the height difference can be at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500 μm or more. Alternatively, the height difference can be at most about 500, 400, 300, 200, 100, 90, 80, 70, 60, 50, 45, 40, 35, 30, 25, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1 μm or less. In some instances, the expansion angle, β , may be between a range of from about 0.5° to about 4° , from about 0.1° to about 10° , or from about 0° to about 90° . For example, the expansion angle can be at least about 0.01° , 0.1° , 0.2° , 0.3° , 0.4° , 0.5° , 0.6° , 0.7° , 0.8° , 0.9° , 1° , 2° , 3° , 4° , 5° , 6° , 7° , 8° , 9° , 10° , 15° , 20° , 25° , 30° , 35° , 40° , 45° , 50° , 55° , 60° , 65° , 70° , 75° , 80° , 85° , or higher. In some instances, the expansion angle can be at most about 89° , 88° , 87° , 86° , 85° , 84° , 83° , 82° , 81° , 80° , 75° , 70° , 65° , 60° , 55° , 50° , 45° , 40° , 35° , 30° , 25° , 20° , 15° , 10° , 9° , 8° , 7° , 6° , 5° , 4° , 3° , 2° , 1° , 0.1° , 0.01° , or less.

[00363] In some instances, the flow rate of the aqueous fluid **712** entering the junction **706** can be between about 0.04 microliters (μL)/minute (min) and about 40 $\mu\text{L}/\text{min}$. In some instances, the flow rate of the aqueous fluid **712** entering the junction **706** can be between about 0.01 microliters (μL)/minute (min) and about 100 $\mu\text{L}/\text{min}$. Alternatively, the flow rate of the aqueous fluid **712** entering the junction **706** can be less than about 0.01 $\mu\text{L}/\text{min}$. Alternatively, the flow rate of the aqueous fluid **712** entering the junction **706** can be greater than about 40 $\mu\text{L}/\text{min}$, such as 45 $\mu\text{L}/\text{min}$, 50 $\mu\text{L}/\text{min}$, 55 $\mu\text{L}/\text{min}$, 60 $\mu\text{L}/\text{min}$, 65 $\mu\text{L}/\text{min}$, 70 $\mu\text{L}/\text{min}$, 75 $\mu\text{L}/\text{min}$, 80 $\mu\text{L}/\text{min}$, 85 $\mu\text{L}/\text{min}$, 90 $\mu\text{L}/\text{min}$, 95 $\mu\text{L}/\text{min}$, 100 $\mu\text{L}/\text{min}$, 110 $\mu\text{L}/\text{min}$, 120 $\mu\text{L}/\text{min}$, 130 $\mu\text{L}/\text{min}$, 140 $\mu\text{L}/\text{min}$, 150 $\mu\text{L}/\text{min}$, or greater. At lower flow rates, such as flow rates of about less than or equal to 10 microliters/minute, the droplet radius may not be dependent on the flow rate of the aqueous fluid **712** entering the junction **706**. The second fluid **714** may be stationary, or substantially stationary, in the reservoir **704**. Alternatively, the second fluid **714** may be flowing, such as at the above flow rates described for the aqueous fluid **712**.

[00364] In some instances, at least about 50% of the droplets generated can have uniform size. In some instances, at least about 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater of the droplets generated can have uniform size. Alternatively, less than about 50% of the droplets generated can have uniform size.

[00365] While **FIGs. 7A** and **7B** illustrate the height difference, Δh , being abrupt at the junction **706** (e.g., a step increase), the height difference may increase gradually (e.g., from about

0 μm to a maximum height difference). Alternatively, the height difference may decrease gradually (e.g., taper) from a maximum height difference. A gradual increase or decrease in height difference, as used herein, may refer to a continuous incremental increase or decrease in height difference, wherein an angle between any one differential segment of a height profile and an immediately adjacent differential segment of the height profile is greater than 90° . For example, at the junction **706**, a bottom wall of the channel and a bottom wall of the reservoir can meet at an angle greater than 90° . Alternatively or in addition, a top wall (e.g., ceiling) of the channel and a top wall (e.g., ceiling) of the reservoir can meet an angle greater than 90° . A gradual increase or decrease may be linear or non-linear (e.g., exponential, sinusoidal, etc.). Alternatively or in addition, the height difference may variably increase and/or decrease linearly or non-linearly. While **FIGs. 7A** and **7B** illustrate the expanding reservoir cross-section height as linear (e.g., constant expansion angle, β), the cross-section height may expand non-linearly. For example, the reservoir may be defined at least partially by a dome-like (e.g., hemispherical) shape having variable expansion angles. The cross-section height may expand in any shape.

[00366] The channel networks, e.g., as described above or elsewhere herein, can be fluidly coupled to appropriate fluidic components. For example, the inlet channel segments are fluidly coupled to appropriate sources of the materials they are to deliver to a channel junction. These sources may include any of a variety of different fluidic components, from simple reservoirs defined in or connected to a body structure of a microfluidic device, to fluid conduits that deliver fluids from off-device sources, manifolds, fluid flow units (e.g., actuators, pumps, compressors) or the like. Likewise, the outlet channel segment (e.g., channel segment **208**, reservoir **604**, etc.) may be fluidly coupled to a receiving vessel or conduit for the partitioned cells for subsequent processing. Again, this may be a reservoir defined in the body of a microfluidic device, or it may be a fluidic conduit for delivering the partitioned cells to a subsequent process operation, instrument or component.

[00367] The methods and systems described herein may be used to greatly increase the efficiency of single cell applications and/or other applications receiving droplet-based input. For example, following the sorting of occupied cells and/or appropriately-sized cells, subsequent operations that can be performed can include generation of amplification products, purification (e.g., via solid phase reversible immobilization (SPRI)), further processing (e.g., shearing, ligation of functional sequences, and subsequent amplification (e.g., via PCR)). These operations may occur in bulk (e.g., outside the partition). In the case where a partition is a droplet in an emulsion, the emulsion can be broken and the contents of the droplet pooled for additional operations. Additional reagents that may be co-partitioned along with the barcode

bearing bead may include oligonucleotides to block ribosomal RNA (rRNA) and nucleases to digest genomic DNA from cells. Alternatively, rRNA removal agents may be applied during additional processing operations. The configuration of the constructs generated by such a method can help minimize (or avoid) sequencing of the poly-T sequence during sequencing and/or sequence the 5' end of a polynucleotide sequence. The amplification products, for example, first amplification products and/or second amplification products, may be subject to sequencing for sequence analysis. In some cases, amplification may be performed using the Partial Hairpin Amplification for Sequencing (PHASE) method.

[00368] A variety of applications require the evaluation of the presence and quantification of different biological particle or organism types within a population of biological particles, including, for example, microbiome analysis and characterization, environmental testing, food safety testing, epidemiological analysis, e.g., in tracing contamination or the like.

Computer systems

[00369] The present disclosure provides computer systems that are programmed to implement methods of the disclosure. **FIG. 25** shows a computer system **2501** that is programmed or otherwise configured to implement one or more of the methods described herein. For example, computer system **2501** may be programmed or otherwise configured to control a microfluidics system (e.g., fluid flow), (ii) sort occupied droplets from unoccupied droplets, (iii) polymerize droplets, (iv) perform sequencing applications, and/or (v) generate and maintain sequencing libraries. The computer system **2501** can regulate various aspects of the present disclosure, such as, for example, regulating fluid flow rate in one or more channels in a microfluidic structure, regulating polymerization application units, etc. The computer system **2501** can be an electronic device of a user or a computer system that is remotely located with respect to the electronic device. The electronic device can be a mobile electronic device.

[00370] The computer system **2501** includes a central processing unit (CPU, also “processor” and “computer processor” herein) **2505**, which can be a single core or multi core processor, or a plurality of processors for parallel processing. The computer system **2501** also includes memory or memory location **2510** (e.g., random-access memory, read-only memory, flash memory), electronic storage unit **2515** (e.g., hard disk), communication interface **2520** (e.g., network adapter) for communicating with one or more other systems, and peripheral devices **2525**, such as cache, other memory, data storage and/or electronic display adapters. The memory **2510**, storage unit **2515**, interface **2520** and peripheral devices **2525** are in communication with the CPU **2505** through a communication bus (solid lines), such as a motherboard. The storage unit

2515 can be a data storage unit (or data repository) for storing data. The computer system **2501** can be operatively coupled to a computer network (“network”) **2530** with the aid of the communication interface **2520**. The network **2530** can be the Internet, an internet and/or extranet, or an intranet and/or extranet that is in communication with the Internet. The network **2530** in some cases is a telecommunication and/or data network. The network **2530** can include one or more computer servers, which can enable distributed computing, such as cloud computing. The network **2530**, in some cases with the aid of the computer system **2501**, can implement a peer-to-peer network, which may enable devices coupled to the computer system **2501** to behave as a client or a server.

[00371] The CPU **2505** can execute a sequence of machine-readable instructions, which can be embodied in a program or software. The instructions may be stored in a memory location, such as the memory **2510**. The instructions can be directed to the CPU **2505**, which can subsequently program or otherwise configure the CPU **2505** to implement methods of the present disclosure. Examples of operations performed by the CPU **2505** can include fetch, decode, execute, and writeback.

[00372] The CPU **2505** can be part of a circuit, such as an integrated circuit. One or more other components of the system **2501** can be included in the circuit. In some cases, the circuit is an application specific integrated circuit (ASIC).

[00373] The storage unit **2515** can store files, such as drivers, libraries and saved programs. The storage unit **2515** can store user data, e.g., user preferences and user programs. The computer system **2501** in some cases can include one or more additional data storage units that are external to the computer system **2501**, such as located on a remote server that is in communication with the computer system **2501** through an intranet or the Internet.

[00374] The computer system **2501** can communicate with one or more remote computer systems through the network **2530**. For instance, the computer system **2501** can communicate with a remote computer system of a user (e.g., operator). Examples of remote computer systems include personal computers (e.g., portable PC), slate or tablet PC's (e.g., Apple® iPad, Samsung® Galaxy Tab), telephones, Smart phones (e.g., Apple® iPhone, Android-enabled device, Blackberry®), or personal digital assistants. The user can access the computer system **2501** via the network **2530**.

[00375] Methods as described herein can be implemented by way of machine (e.g., computer processor) executable code stored on an electronic storage location of the computer system **2501**, such as, for example, on the memory **2510** or electronic storage unit **2515**. The machine executable or machine readable code can be provided in the form of software. During use, the

code can be executed by the processor **2505**. In some cases, the code can be retrieved from the storage unit **2515** and stored on the memory **2510** for ready access by the processor **2505**. In some situations, the electronic storage unit **2515** can be precluded, and machine-executable instructions are stored on memory **2510**.

[00376] The code can be pre-compiled and configured for use with a machine having a processor adapted to execute the code, or can be compiled during runtime. The code can be supplied in a programming language that can be selected to enable the code to execute in a pre-compiled or as-compiled fashion.

[00377] Aspects of the systems and methods provided herein, such as the computer system **2501**, can be embodied in programming. Various aspects of the technology may be thought of as “products” or “articles of manufacture” typically in the form of machine (or processor) executable code and/or associated data that is carried on or embodied in a type of machine readable medium. Machine-executable code can be stored on an electronic storage unit, such as memory (e.g., read-only memory, random-access memory, flash memory) or a hard disk. “Storage” type media can include any or all of the tangible memory of the computers, processors or the like, or associated modules thereof, such as various semiconductor memories, tape drives, disk drives and the like, which may provide non-transitory storage at any time for the software programming. All or portions of the software may at times be communicated through the Internet or various other telecommunication networks. Such communications, for example, may enable loading of the software from one computer or processor into another, for example, from a management server or host computer into the computer platform of an application server. Thus, another type of media that may bear the software elements includes optical, electrical and electromagnetic waves, such as used across physical interfaces between local devices, through wired and optical landline networks and over various air-links. The physical elements that carry such waves, such as wired or wireless links, optical links or the like, also may be considered as media bearing the software. As used herein, unless restricted to non-transitory, tangible “storage” media, terms such as computer or machine “readable medium” refer to any medium that participates in providing instructions to a processor for execution.

[00378] Hence, a machine readable medium, such as computer-executable code, may take many forms, including but not limited to, a tangible storage medium, a carrier wave medium or physical transmission medium. Non-volatile storage media include, for example, optical or magnetic disks, such as any of the storage devices in any computer(s) or the like, such as may be used to implement the databases, etc. shown in the drawings. Volatile storage media include dynamic memory, such as main memory of such a computer platform. Tangible transmission

media include coaxial cables; copper wire and fiber optics, including the wires that comprise a bus within a computer system. Carrier-wave transmission media may take the form of electric or electromagnetic signals, or acoustic or light waves such as those generated during radio frequency (RF) and infrared (IR) data communications. Common forms of computer-readable media therefore include for example: a floppy disk, a flexible disk, hard disk, magnetic tape, any other magnetic medium, a CD-ROM, DVD or DVD-ROM, any other optical medium, punch cards paper tape, any other physical storage medium with patterns of holes, a RAM, a ROM, a PROM and EPROM, a FLASH-EPROM, any other memory chip or cartridge, a carrier wave transporting data or instructions, cables or links transporting such a carrier wave, or any other medium from which a computer may read programming code and/or data. Many of these forms of computer readable media may be involved in carrying one or more sequences of one or more instructions to a processor for execution.

[00379] The computer system **2501** can include or be in communication with an electronic display **2535** that comprises a user interface (UI) **2540** for providing, for example, results of sequencing analysis, etc. Examples of UIs include, without limitation, a graphical user interface (GUI) and web-based user interface.

[00380] Methods and systems of the present disclosure can be implemented by way of one or more algorithms. An algorithm can be implemented by way of software upon execution by the central processing unit **2505**. The algorithm can, for example, perform a nucleic acid sequencing assay, etc.

[00381] Devices, systems, compositions and methods of the present disclosure may be used for various applications, such as, for example, processing a single analyte (e.g., RNA, DNA, or protein) or multiple analytes (e.g., DNA and RNA, DNA and protein, RNA and protein, or RNA, DNA and protein) from a single cell. For example, a biological particle (e.g., a cell or cell bead) is partitioned in a partition (e.g., droplet), and multiple analytes from the biological particle are processed for subsequent processing. The multiple analytes may be from the single cell. This may enable, for example, simultaneous proteomic, transcriptomic and genomic analysis of the cell.

[00382] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. It is not intended that the invention be limited by the specific examples provided within the specification. While the invention has been described with reference to the aforementioned specification, the descriptions and illustrations of the embodiments herein are not meant to be construed in a limiting sense. Numerous variations, changes, and substitutions

will now occur to those skilled in the art without departing from the invention. Furthermore, it shall be understood that all aspects of the invention are not limited to the specific depictions, configurations or relative proportions set forth herein which depend upon a variety of conditions and variables. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is therefore contemplated that the invention shall also cover any such alternatives, modifications, variations or equivalents. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

CLAIMS

WHAT IS CLAIMED IS:

1. A method for processing a nucleic acid sample, comprising:
 - (a) providing a biological particle comprising a deoxyribonucleic acid (DNA) molecule and a ribonucleic acid (RNA) molecule within a partition among a plurality of partitions, wherein said partition comprises (i) a first nucleic acid barcode molecule, (ii) a second nucleic acid barcode molecule, and (iii) a splint sequence, wherein said first nucleic acid barcode molecule and said second nucleic acid barcode molecule comprise a common barcode sequence, wherein said first nucleic acid barcode molecule comprises an overhang sequence, and wherein said splint sequence comprises a sequence complementary to said DNA molecule and a sequence complementary to said overhang sequence;
 - (b) within said partition, using said DNA molecule, said first nucleic acid barcode molecule, and said splint sequence to generate a first barcoded nucleic acid product corresponding to said DNA molecule and using said RNA molecule and said second nucleic acid barcode molecule to generate a second barcoded nucleic acid product corresponding to said RNA molecule, wherein said first barcoded nucleic acid product and said second barcoded nucleic acid product comprise said common barcode sequence, or a complement thereof; and
 - (c) recovering said first barcoded nucleic acid product and said second barcoded nucleic acid product from said partition.
2. The method of claim 1, wherein said biological particle is a cell, cell bead, or cell nucleus.
3. The method of claim 1 or 2, further comprising lysing or permeabilizing said biological particle within said partition to provide access to said DNA molecule and said RNA molecule therein.
4. The method of any one of claims 1-3, further comprising, prior to (a), processing an open chromatin structure of said nucleic acid sample with a transposase to yield said DNA molecule.
5. The method of claim 4, wherein said transposase is included in a transposase-nucleic acid complex that comprises (i) a first nucleic acid molecule comprising a first transposon end sequence and a first sequencing primer or portion thereof, or a complement thereof and (ii) a second nucleic acid molecule comprising a second transposon end sequence and a second sequencing primer or portion thereof, or a complement thereof.

6. The method of claim 5, wherein said first transposon end sequence and said second transposon end sequence are the same, and wherein said first transposon end sequence and said second transposon end sequence are each hybridized to complementary sequences.
7. The method of any one of claims 1-6, wherein generating said first barcoded nucleic acid product comprises (i) subjecting said DNA molecule and said splint sequence to conditions sufficient to hybridize to said sequence of said DNA molecule to said splint sequence and (ii) subjecting said first nucleic acid barcode molecule and said splint sequence to conditions sufficient to hybridize said overhang sequence of said first nucleic acid barcode molecule to said splint sequence.
8. The method of claim 7, wherein said splint sequence is hybridized to said sequence of said DNA molecule outside of said partition.
9. The method of any one of claims 1-8, further comprising ligating said first nucleic acid barcode molecule to said DNA molecule.
10. The method of any one of claims 1-9, wherein said first nucleic acid barcode molecule comprises a flow cell sequence, or complement thereof; a barcode sequence; a sequencing primer, or portion thereof; a unique molecular identifier sequence; or a combination thereof.
11. The method of any one of claims 1-10, wherein said DNA molecule comprises one or more gaps, and further comprising, prior to (c), subjecting said DNA molecule to an extension process to fill said one or more gaps.
12. The method of any one of claims 1-11, wherein said second nucleic acid barcode molecule comprises a unique molecular identifier sequence, a barcode sequence, a sequencing primer or portion thereof, or a combination thereof.
13. The method of any one of claims 1-12, further comprising, prior to (b), within said partition, reverse transcribing said RNA molecule to provide a complementary DNA (cDNA) molecule.
14. The method of claim 13, wherein, prior to reverse transcribing said RNA molecule, said second nucleic acid barcode molecule is hybridized to a sequence of a precursor of said RNA molecule.
15. The method of claim 14, wherein said precursor comprises a polyA sequence, and wherein said second nucleic acid barcode molecule comprises a polyT sequence configured to hybridize to said polyA sequence.

16. The method of claim 13, further comprising appending an additional sequence to said cDNA molecule, wherein said additional sequence is a poly(C) sequence.
17. The method of claim 16, further comprising contacting said cDNA molecule with a template-switching oligonucleotide under conditions sufficient to hybridize said template-switching oligonucleotide to said cDNA molecule, wherein said template-switching oligonucleotide hybridizes to said additional sequence of said cDNA molecule.
18. The method of claim 17, further comprising extending said cDNA molecule to generate an extended cDNA molecule, which extended cDNA molecule comprises sequences complementary to sequences of said template-switching oligonucleotide.
19. The method of claim 17 or 18, wherein said template-switching oligonucleotide comprises a sequencing primer or portion thereof, or complement thereof; a unique molecular identifier sequence, or complement thereof; or a combination thereof.
20. The method of claim 18 or 19, wherein said template-switching oligonucleotide is said second nucleic acid barcode molecule, and wherein said extended cDNA molecule is said second barcoded nucleic acid product.
21. The method of any one of claims 1-20, wherein said first nucleic acid barcode molecule and said second nucleic acid barcode molecule comprise a common nucleic acid sequence.
22. The method of any one of claims 1-21, wherein said partition is a droplet among a plurality of droplets.
23. The method of claim 22, wherein said partition is a droplet and wherein (c) comprises breaking or disrupting said droplet.
24. The method of any one of claims 1-23, further comprising, subsequent to (c), using (i) said first barcoded nucleic acid product to generate a first plurality of amplification products and (ii) said second barcoded nucleic acid product to generate a second plurality of amplification products.
25. The method of claim 24, wherein generating said first plurality of amplification products and said second plurality of amplification products comprises performing one or more nucleic acid amplification reactions using said first barcoded nucleic acid product or said second barcoded nucleic acid product.
26. The method of claim 25, wherein performing said one or more nucleic acid amplification reactions comprises attaching one or more flow cell sequences to said first barcoded nucleic acid product or said second barcoded nucleic acid product, or a derivative thereof.

27. The method of any one of claims 1-26, wherein said first nucleic acid barcode molecule and said second nucleic acid barcode molecule are coupled to a bead.
28. The method of claim 27, wherein said bead is a gel bead.
29. The method of claim 27 or 28, wherein said first nucleic acid barcode molecule is coupled to said bead via a first labile moiety and said second nucleic acid barcode molecule is coupled to said bead via a second labile moiety.
30. The method of any one of claims 27-29, wherein said first nucleic acid barcode molecule and said second nucleic acid barcode molecule are releasably coupled to said bead.
31. The method of claim 30, wherein said first nucleic acid barcode molecule is releasable from said bead upon application of a first stimulus and said second nucleic acid barcode molecule is releasable from said bead upon application of a second stimulus.
32. A method for processing a nucleic acid sample, comprising:
- a) providing a biological particle comprising a deoxyribonucleic acid (DNA) molecule and a ribonucleic acid (RNA) molecule within a partition among a plurality of partitions, wherein said partition comprises a first nucleic acid barcode molecule and a second nucleic acid barcode molecule, wherein said first nucleic acid barcode molecule and said second nucleic acid barcode molecule comprise a common barcode sequence;
 - b) within said partition, generating (i) a first barcoded nucleic acid product from said DNA molecule and said first nucleic acid barcode molecule and (ii) a second barcoded nucleic acid product from said RNA molecule and said second nucleic acid barcode molecule, wherein said first barcoded nucleic acid product and said second barcoded nucleic acid product comprise said common barcode sequence, or a complement thereof;
 - c) recovering said first barcoded nucleic acid product and said second barcoded nucleic acid product from said partition;
 - d) processing said first barcoded nucleic acid product to generate a processed nucleic acid product comprising a dA tail; and
 - e) using (i) said processed nucleic acid product to generate a first plurality of amplification products and (ii) said second barcoded nucleic acid product to generate a second plurality of amplification products.
33. The method of claim 32, wherein said biological particle is a cell, cell bead, or cell nucleus.

34. The method of claim 32 or 33, further comprising lysing or permeabilizing said biological particle within said partition to provide access to said DNA molecule and said RNA molecule therein.
35. The method of any one of claims 32-34, further comprising, prior to (a), processing an open chromatin structure of said nucleic acid sample with a transposase to yield said DNA molecule.
36. The method of claim 35, wherein said transposase is included in a transposase-nucleic acid complex that comprises (i) a first nucleic acid molecule comprising a first transposon end sequence and a first sequencing primer or portion thereof, or a complement thereof and (ii) a second nucleic acid molecule comprising a second transposon end sequence and a second sequencing primer or portion thereof, or a complement thereof.
37. The method of claim 36, wherein said first transposon end sequence and said second transposon end sequence are the same, and wherein said first transposon end sequence and said second transposon end sequence are each hybridized to complementary sequences.
38. The method of any one of claims 32-36, wherein said first nucleic acid barcode molecule comprises a sequencing primer or portion thereof, or complement thereof.
39. The method of claim 38, wherein (b) comprises subjecting said DNA molecule to conditions sufficient to hybridize said first nucleic acid barcode molecule to said DNA molecule.
40. The method of claim 39, further comprising (i) extending said first nucleic acid barcode molecule hybridized to said DNA molecule and (ii) ligating said first nucleic acid barcode molecule to said DNA molecule to generate said first barcoded nucleic acid product.
41. The method of any one of claims 32-40, further comprising ligating said first nucleic acid barcode molecule to said DNA molecule.
42. The method of any one of claims 32-41, wherein said first nucleic acid barcode molecule comprises a flow cell sequence, or complement thereof; a barcode sequence; a unique molecular identifier sequence; or a combination thereof.
43. The method of any one of claims 32-42, wherein said DNA molecule comprises one or more gaps, and further comprising, prior to (c), subjecting said DNA molecule to an extension process to fill said one or more gaps.

44. The method of any one of claims 32-43, wherein said second nucleic acid barcode molecule comprises a unique molecular identifier sequence, a barcode sequence, a sequencing primer or portion thereof, or a combination thereof.
45. The method of any one of claims 32-44, further comprising, prior to (b), within said partition, reverse transcribing said RNA molecule to provide a complementary DNA (cDNA) molecule.
46. The method of claim 45, wherein, prior to reverse transcribing said RNA molecule, said second nucleic acid barcode molecule is hybridized to a sequence of a precursor of said RNA molecule.
47. The method of claim 46, wherein said precursor comprises a polyA sequence, and wherein said second nucleic acid barcode molecule comprises a polyT sequence configured to hybridize to said polyA sequence.
48. The method of claim 46, further comprising appending an additional sequence to said cDNA molecule, wherein said additional sequence is a poly(C) sequence.
49. The method of claim 48, further comprising contacting said cDNA molecule with a template-switching oligonucleotide under conditions sufficient to hybridize said template-switching oligonucleotide to said cDNA molecule, wherein said template-switching oligonucleotide hybridizes to said additional sequence of said cDNA molecule.
50. The method of claim 49, further comprising extending said cDNA molecule to generate an extended cDNA molecule, which extended cDNA molecule comprises sequences complementary to sequences of said template-switching oligonucleotide.
51. The method of claim 49 or 50, wherein said template-switching oligonucleotide comprises a sequencing primer or portion thereof, or complement thereof; a unique molecular identifier sequence, or complement thereof; or a combination thereof.
52. The method of claim 50 or 51, wherein said template-switching oligonucleotide is said second nucleic acid barcode molecule, and wherein said extended cDNA molecule is said second barcoded nucleic acid product.
53. The method of any one of claims 32-52, wherein said first nucleic acid barcode molecule and said second nucleic acid barcode molecule comprise a common nucleic acid sequence.
54. The method of any one of claims 32-53, wherein said partition is a droplet among a plurality of droplets.
55. The method of claim 54, wherein said partition is a droplet and wherein (c) comprises breaking or disrupting said droplet.

56. The method of any one of claims 32-55, wherein (e) comprises performing one or more nucleic acid amplification reactions using said first barcoded nucleic acid product or said second barcoded nucleic acid product.
57. The method of claim 56, wherein (e) comprises attaching one or more flow cell sequences to said first barcoded nucleic acid product or said second barcoded nucleic acid product, or a derivative thereof.
58. The method of any one of claims 32-57, wherein said first nucleic acid barcode molecule and said second nucleic acid barcode molecule are coupled to a bead.
59. The method of claim 58, wherein said bead is a gel bead.
60. The method of claim 58 or 59, wherein said first nucleic acid barcode molecule is coupled to said bead via a first labile moiety and said second nucleic acid barcode molecule is coupled to said bead via a second labile moiety.
61. The method of any one of claims 58-60, wherein said first nucleic acid barcode molecule and said second nucleic acid barcode molecule are releasably coupled to said bead.
62. The method of claim 61, wherein said first nucleic acid barcode molecule is releasable from said bead upon application of a first stimulus and said second nucleic acid barcode molecule is releasable from said bead upon application of a second stimulus.
63. A method for processing a nucleic acid sample, comprising:
- a) providing a biological particle comprising a deoxyribonucleic acid (DNA) molecule and a ribonucleic acid (RNA) molecule within a partition among a plurality of partitions, wherein said partition comprises a first nucleic acid barcode molecule and a second nucleic acid barcode molecule, wherein said first nucleic acid barcode molecule and said second nucleic acid barcode molecule comprise a common barcode sequence, and wherein said DNA molecule comprises a first hairpin moiety at a first end and a second hairpin moiety at a second end;
 - b) within said partition, generating (i) a first barcoded nucleic acid product from said DNA molecule and said first nucleic acid barcode molecule and (ii) a second barcoded nucleic acid product from said RNA molecule and said second nucleic acid barcode molecule, wherein said first barcoded nucleic acid product and said second barcoded nucleic acid product comprise said common barcode sequence, or a complement thereof;
 - c) recovering said first barcoded nucleic acid product and said second barcoded nucleic acid product from said partition; and

- d) using (i) said first barcoded nucleic acid product to generate a first plurality of amplification products and (ii) said second barcoded nucleic acid product to generate a second plurality of amplification products.
64. The method of claim 63, wherein said biological particle is a cell, cell bead, or cell nucleus.
65. The method of claim 63 or 64, further comprising lysing or permeabilizing said biological particle within said partition to provide access to said DNA molecule and said RNA molecule therein.
66. The method of any one of claims 63-65, further comprising, prior to (a), processing an open chromatin structure of said nucleic acid sample with a transposase to yield said DNA molecule.
67. The method of claim 66, wherein said transposase is included in a transposase-nucleic acid complex that comprises (i) a first nucleic acid molecule comprising a first transposon end sequence and a first sequencing primer or portion thereof, or a complement thereof and (ii) a second nucleic acid molecule comprising a second transposon end sequence and a second sequencing primer or portion thereof, or a complement thereof.
68. The method of claim 67, wherein said first transposon end sequence and said second transposon end sequence are the same, and wherein said first transposon end sequence and said second transposon end sequence are each hybridized to complementary sequences.
69. The method of any one of claims 63-68, wherein said first nucleic acid barcode molecule comprises a sequencing primer or portion thereof, or complement thereof.
70. The method of claim 69, wherein (b) comprises subjecting said DNA molecule to conditions sufficient to hybridize said first nucleic acid barcode molecule to said DNA molecule.
71. The method of claim 70, further comprising (i) extending said first nucleic acid barcode molecule hybridized to said DNA molecule and (ii) ligating said first nucleic acid barcode molecule to said DNA molecule to generate said first barcoded nucleic acid product.
72. The method of any one of claims 63-71, further comprising ligating said first nucleic acid barcode molecule to said DNA molecule.

73. The method of any one of claims 63-72, wherein said first nucleic acid barcode molecule comprises a flow cell sequence, or complement thereof; a barcode sequence; a unique molecular identifier sequence; or a combination thereof.
74. The method of any one of claims 63-73, wherein said DNA molecule comprises one or more gaps, and further comprising, prior to (c), subjecting said DNA molecule to an extension process to fill said one or more gaps.
75. The method of any one of claims 63-74, wherein said second nucleic acid barcode molecule comprises a unique molecular identifier sequence, a barcode sequence, a sequencing primer or portion thereof, or a combination thereof.
76. The method of any one of claims 63-75, further comprising, prior to (b), within said partition, reverse transcribing said RNA molecule to provide a complementary DNA (cDNA) molecule.
77. The method of claim 76, wherein, prior to reverse transcribing said RNA molecule, said second nucleic acid barcode molecule is hybridized to a sequence of a precursor of said RNA molecule.
78. The method of claim 77, wherein said precursor comprises a polyA sequence, and wherein said second nucleic acid barcode molecule comprises a polyT sequence configured to hybridize to said polyA sequence.
79. The method of claim 77, further comprising appending an additional sequence to said cDNA molecule, wherein said additional sequence is a poly(C) sequence.
80. The method of claim 79, further comprising contacting said cDNA molecule with a template-switching oligonucleotide under conditions sufficient to hybridize said template-switching oligonucleotide to said cDNA molecule, wherein said template-switching oligonucleotide hybridizes to said additional sequence of said cDNA molecule.
81. The method of claim 80, further comprising extending said cDNA molecule to generate an extended cDNA molecule, which extended cDNA molecule comprises sequences complementary to sequences of said template-switching oligonucleotide.
82. The method of claim 80 or 81, wherein said template-switching oligonucleotide comprises a sequencing primer or portion thereof, or complement thereof; a unique molecular identifier sequence, or complement thereof; or a combination thereof.
83. The method of claim 81 or 82, wherein said template-switching oligonucleotide is said second nucleic acid barcode molecule, and wherein said extended cDNA molecule is said second barcoded nucleic acid product.

84. The method of any one of claims 63-83, wherein said first nucleic acid barcode molecule and said second nucleic acid barcode molecule comprise a common nucleic acid sequence.
85. The method of any one of claims 63-84, wherein said partition is a droplet among a plurality of droplets.
86. The method of claim 85, wherein said partition is a droplet and wherein (c) comprises breaking or disrupting said droplet.
87. The method of any one of claims 63-86, wherein (e) comprises performing one or more nucleic acid amplification reactions using said first barcoded nucleic acid product or said second barcoded nucleic acid product.
88. The method of claim 87, wherein (e) comprises attaching one or more flow cell sequences to said first barcoded nucleic acid product or said second barcoded nucleic acid product, or a derivative thereof.
89. The method of any one of claims 63-88, wherein said first nucleic acid barcode molecule and said second nucleic acid barcode molecule are coupled to a bead.
90. The method of claim 89, wherein said bead is a gel bead.
91. The method of claim 89 or 90, wherein said first nucleic acid barcode molecule is coupled to said bead via a first labile moiety and said second nucleic acid barcode molecule is coupled to said bead via a second labile moiety.
92. The method of any one of claims 89-91, wherein said first nucleic acid barcode molecule and said second nucleic acid barcode molecule are releasably coupled to said bead.
93. The method of claim 92, wherein said first nucleic acid barcode molecule is releasable from said bead upon application of a first stimulus and said second nucleic acid barcode molecule is releasable from said bead upon application of a second stimulus.