

METHODS FOR TREATING AND/OR PREVENTING GRAFT-VERSUS-HOST
DISEASE AND/OR DIFFUSE ALVEOLAR HEMORRHAGE AND/OR VENO-
OCCLUSIVE DISEASE ASSOCIATED WITH HEMATOPOIETIC STEM CELL
TRANSPLANT

5

STATEMENT REGARDING SEQUENCE LISTING

The sequence listing associated with this application is provided in text format in lieu of a paper copy and is hereby incorporated by reference into the specification. The
10 name of the text file containing the sequence listing is MP_1_0278_PCT_SequenceListingasFiled_20180807; the txt file is 116 KB; was created on August 7, 2018, and is being submitted via EFS-Web with the filing of the specification.

BACKGROUND

15 The complement system provides an early acting mechanism to initiate, amplify and orchestrate the immune response to microbial infection and other acute insults (M.K. Liszewski and J.P. Atkinson, 1993, in *Fundamental Immunology*, Third Edition, edited by W.E. Paul, Raven Press, Ltd., New York), in humans and other vertebrates. While complement activation provides a valuable first-line defense against potential
20 pathogens, the activities of complement that promote a protective immune response can also represent a potential threat to the host (K.R. Kalli, et al., *Springer Semin. Immunopathol.* 15:417-431, 1994; B.P. Morgan, *Eur. J. Clinical Investig.* 24:219-228, 1994). For example, C3 and C5 proteolytic products recruit and activate neutrophils. While indispensable for host defense, activated neutrophils are indiscriminate in their
25 release of destructive enzymes and may cause organ damage. In addition, complement activation may cause the deposition of lytic complement components on nearby host cells as well as on microbial targets, resulting in host cell lysis.

The complement system has also been implicated in the pathogenesis of numerous acute and chronic disease states, including: myocardial infarction, stroke, ARDS,
30 reperfusion injury, septic shock, capillary leakage following thermal burns, postcardiopulmonary bypass inflammation, transplant rejection, rheumatoid arthritis, multiple sclerosis, myasthenia gravis, and Alzheimer's disease. In almost all of these

conditions, complement is not the cause but is one of several factors involved in pathogenesis. Nevertheless, complement activation may be a major pathological mechanism and represents an effective point for clinical control in many of these disease states. The growing recognition of the importance of complement-mediated tissue injury
5 in a variety of disease states underscores the need for effective complement inhibitory drugs. To date, Eculizumab (Solaris®), an antibody against C5, is the only complement-targeting drug that has been approved for human use. Yet, C5 is one of several effector molecules located “downstream” in the complement system, and blockade of C5 does not inhibit activation of the complement system. Therefore, an inhibitor of the initiation
10 steps of complement activation would have significant advantages over a “downstream” complement inhibitor.

Currently, it is widely accepted that the complement system can be activated through three distinct pathways: the classical pathway, the lectin pathway, and the alternative pathway. The classical pathway is usually triggered by a complex composed
15 of host antibodies bound to a foreign particle (*i.e.*, an antigen) and thus requires prior exposure to an antigen for the generation of a specific antibody response. Since activation of the classical pathway depends on a prior adaptive immune response by the host, the classical pathway is part of the acquired immune system. In contrast, both the lectin and alternative pathways are independent of adaptive immunity and are part of the
20 innate immune system.

The activation of the complement system results in the sequential activation of serine protease zymogens. The first step in activation of the classical pathway is the binding of a specific recognition molecule, C1q, to antigen-bound IgG and IgM molecules. C1q is associated with the C1r and C1s serine protease proenzymes as a
25 complex called C1. Upon binding of C1q to an immune complex, autoproteolytic cleavage of the Arg-Ile site of C1r is followed by C1r-mediated cleavage and activation of C1s, which thereby acquires the ability to cleave C4 and C2. C4 is cleaved into two fragments, designated C4a and C4b, and, similarly, C2 is cleaved into C2a and C2b. C4b fragments are able to form covalent bonds with adjacent hydroxyl or amino groups and
30 generate the C3 convertase (C4b2a) through noncovalent interaction with the C2a fragment of activated C2. C3 convertase (C4b2a) activates C3 by proteolytic cleavage into C3a and C3b subcomponents leading to generation of the C5 convertase (C4b2a3b), which, by cleaving C5 leads to the formation of the membrane attack complex (C5b

combined with C6, C7, C8 and C-9, also referred to as “MAC”) that can disrupt cellular membranes leading to cell lysis. The activated forms of C3 and C4 (C3b and C4b) are covalently deposited on the foreign target surfaces, which are recognized by complement receptors on multiple phagocytes.

5 Independently, the first step in activation of the complement system through the lectin pathway is also the binding of specific recognition molecules, which is followed by the activation of associated serine protease proenzymes. However, rather than the binding of immune complexes by C1q, the recognition molecules in the lectin pathway comprise a group of carbohydrate-binding proteins (mannan-binding lectin (MBL),
10 H-ficolin, M-ficolin, L-ficolin and C-type lectin CL-11), collectively referred to as lectins. See J. Lu et al., *Biochim. Biophys. Acta* 1572:387-400, (2002); Holmskov et al., *Annu. Rev. Immunol.* 21:547-578 (2003); Teh et al., *Immunology* 101:225-232 (2000)). See also J. Luet et al., *Biochim Biophys Acta* 1572:387-400 (2002); Holmskov et al., *Annu Rev Immunol* 21:547-578 (2003); Teh et al., *Immunology* 101:225-232 (2000); Hansen et
15 al., *J. Immunol* 185(10):6096-6104 (2010).

Ikeda et al. first demonstrated that, like C1q, MBL could activate the complement system upon binding to yeast mannan-coated erythrocytes in a C4-dependent manner (Ikeda et al., *J. Biol. Chem.* 262:7451-7454, (1987)). MBL, a member of the collectin protein family, is a calcium-dependent lectin that binds carbohydrates with 3- and
20 4-hydroxy groups oriented in the equatorial plane of the pyranose ring. Prominent ligands for MBL are thus D-mannose and N-acetyl-D-glucosamine, while carbohydrates not fitting this steric requirement have undetectable affinity for MBL (Weis et al., *Nature* 360:127-134, (1992)). The interaction between MBL and monovalent sugars is extremely weak, with dissociation constants typically in the single-digit millimolar range.
25 MBL achieves tight, specific binding to glycan ligands by avidity, i.e., by interacting simultaneously with multiple monosaccharide residues located in close proximity to each other (Lee et al., *Archiv. Biochem. Biophys.* 299:129-136, (1992)). MBL recognizes the carbohydrate patterns that commonly decorate microorganisms such as bacteria, yeast, parasites and certain viruses. In contrast, MBL does not recognize D-galactose and sialic
30 acid, the penultimate and ultimate sugars that usually decorate "mature" complex glycoconjugates present on mammalian plasma and cell surface glycoproteins. This binding specificity is thought to promote recognition of “foreign” surfaces and help protect from “self-activation.” However, MBL does bind with high affinity to clusters of

high-mannose "precursor" glycans on N-linked glycoproteins and glycolipids sequestered in the endoplasmic reticulum and Golgi of mammalian cells (Maynard et al., *J. Biol. Chem.* 257:3788-3794, (1982)). Therefore, damaged cells are potential targets for lectin pathway activation via MBL binding.

5 The ficolins possess a different type of lectin domain than MBL, called the fibrinogen-like domain. Ficolins bind sugar residues in a Ca^{++} -independent manner. In humans, three kinds of ficolins (L-ficolin, M-ficolin and H-ficolin) have been identified. The two serum ficolins, L-ficolin and H-ficolin, have in common a specificity for N-acetyl-D-glucosamine; however, H-ficolin also binds N-acetyl-D-galactosamine. The
10 difference in sugar specificity of L-ficolin, H-ficolin, CL-11, and MBL means that the different lectins may be complementary and target different, though overlapping, glycoconjugates. This concept is supported by the recent report that, of the known lectins in the lectin pathway, only L-ficolin binds specifically to lipoteichoic acid, a cell wall glycoconjugate found on all Gram-positive bacteria (Lynch et al., *J. Immunol.*
15 172:1198-1202, (2004)). The collectins (i.e., MBL) and the ficolins bear no significant similarity in amino acid sequence. However, the two groups of proteins have similar domain organizations and, like C1q, assemble into oligomeric structures, which maximize the possibility of multisite binding.

 The serum concentrations of MBL are highly variable in healthy populations and
20 this is genetically controlled by polymorphisms/mutations in both the promoter and coding regions of the MBL gene. As an acute phase protein, the expression of MBL is further upregulated during inflammation. L-ficolin is present in serum at concentrations similar to those of MBL. Therefore, the L-ficolin branch of the lectin pathway is potentially comparable to the MBL arm in strength. MBL and ficolins can also function
25 as opsonins, which allow phagocytes to target MBL- and ficolin-decorated surfaces (see Jack et al., *J Leukoc Biol.*, 77(3):328-36 (2004), Matsushita and Fujita, *Immunobiology*, 205(4-5):490-7 (2002), Aoyagi et al., *J Immunol*, 174(1):418-25(2005). This opsonization requires the interaction of these proteins with phagocyte receptors (Kuhlman et al., *J. Exp. Med.* 169:1733, (1989); Matsushita et al., *J. Biol. Chem.* 271:2448-54,
30 (1996)), the identity of which has not been established.

 Human MBL forms a specific and high-affinity interaction through its collagen-like domain with unique C1r/C1s-like serine proteases, termed MBL-associated serine proteases (MASPs). To date, three MASPs have been described. First, a single

enzyme "MASP" was identified and characterized as the enzyme responsible for the initiation of the complement cascade (i.e., cleaving C2 and C4) (Matsushita et al., *J Exp Med* 176(6):1497-1502 (1992); Ji et al., *J. Immunol.* 150:571-578, (1993)). It was subsequently determined that the MASP activity was, in fact, a mixture of two proteases:
5 MASP-1 and MASP-2 (Thiel et al., *Nature* 386:506-510, (1997)). However, it was demonstrated that the MBL-MASP-2 complex alone is sufficient for complement activation (Vorup-Jensen et al., *J. Immunol.* 165:2093-2100, (2000)). Furthermore, only MASP-2 cleaved C2 and C4 at high rates (Ambrus et al., *J. Immunol.* 170:1374-1382, (2003)). Therefore, MASP-2 is the protease responsible for activating C4 and C2 to
10 generate the C3 convertase, C4b2a. This is a significant difference from the C1 complex of the classical pathway, where the coordinated action of two specific serine proteases (C1r and C1s) leads to the activation of the complement system. In addition, a third novel protease, MASP-3, has been isolated (Dahl, M.R., et al., *Immunity* 15:127-35, 2001). MASP-1 and MASP-3 are alternatively spliced products of the same gene.

15 MASPs share identical domain organizations with those of C1r and C1s, the enzymatic components of the C1 complex (Sim et al., *Biochem. Soc. Trans.* 28:545, (2000)). These domains include an N-terminal C1r/C1s/sea urchin VEGF/bone morphogenic protein (CUB) domain, an epidermal growth factor-like domain, a second CUB domain, a tandem of complement control protein domains, and a serine protease domain. As in the C1 proteases, activation of MASP-2 occurs through cleavage of an
20 Arg-Ile bond adjacent to the serine protease domain, which splits the enzyme into disulfide-linked A and B chains, the latter consisting of the serine protease domain.

MBL can also associate with an alternatively sliced form of MASP-2, known as MBL-associated protein of 19 kDa (MAp19) or small MBL-associated protein (sMAP),
25 which lacks the catalytic activity of MASP2. (Stover, *J. Immunol.* 162:3481-90, (1999); Takahashi et al., *Int. Immunol.* 11:859-863, (1999)). MAp19 comprises the first two domains of MASP-2, followed by an extra sequence of four unique amino acids. The function of Map19 is unclear (Degn et al., *J Immunol. Methods*, 2011). The MASP-1 and MASP-2 genes are located on human chromosomes 3 and 1, respectively
30 (Schwaebler et al., *Immunobiology* 205:455-466, (2002)).

Several lines of evidence suggest that there are different MBL-MASP complexes and a large fraction of the MASPs in serum is not complexed with MBL (Thiel, et al., *J. Immunol.* 165:878-887, (2000)). Both H- and L-ficolin bind to all MASPs and activate

the lectin complement pathway, as does MBL (Dahl et al., *Immunity* 15:127-35, (2001); Matsushita et al., *J. Immunol.* 168:3502-3506, (2002)). Both the lectin and classical pathways form a common C3 convertase (C4b2a) and the two pathways converge at this step.

5 The lectin pathway is widely thought to have a major role in host defense against infection in the naïve host. Strong evidence for the involvement of MBL in host defense comes from analysis of patients with decreased serum levels of functional MBL (Kilpatrick, *Biochim. Biophys. Acta* 1572:401-413, (2002)). Such patients display susceptibility to recurrent bacterial and fungal infections. These symptoms are usually
10 evident early in life, during an apparent window of vulnerability as maternally derived antibody titer wanes, but before a full repertoire of antibody responses develops. This syndrome often results from mutations at several sites in the collagenous portion of MBL, which interfere with proper formation of MBL oligomers. However, since MBL can function as an opsonin independent of complement, it is not known to what extent the
15 increased susceptibility to infection is due to impaired complement activation.

 In contrast to the classical and lectin pathways, no initiators of the alternative pathway have been found to fulfill the recognition functions that C1q and lectins perform in the other two pathways. Currently it is widely accepted that the alternative pathway spontaneously undergoes a low level of turnover activation, which can be readily
20 amplified on foreign or other abnormal surfaces (bacteria, yeast, virally infected cells, or damaged tissue) that lack the proper molecular elements that keep spontaneous complement activation in check. There are four plasma proteins directly involved in the activation of the alternative pathway: C3, factors B and D, and properdin.

 Although there is extensive evidence implicating both the classical and alternative
25 complement pathways in the pathogenesis of non-infectious human diseases, the role of the lectin pathway is just beginning to be evaluated. Recent studies provide evidence that activation of the lectin pathway can be responsible for complement activation and related inflammation in ischemia/reperfusion injury. Collard et al. (2000) reported that cultured endothelial cells subjected to oxidative stress bind MBL and show deposition of C3 upon
30 exposure to human serum (Collard et al., *Am. J. Pathol.* 156:1549-1556, (2000)). In addition, treatment of human sera with blocking anti-MBL monoclonal antibodies inhibited MBL binding and complement activation. These findings were extended to a rat model of myocardial ischemia-reperfusion in which rats treated with a blocking

antibody directed against rat MBL showed significantly less myocardial damage upon occlusion of a coronary artery than rats treated with a control antibody (Jordan et al., *Circulation* 104:1413-1418, (2001)). The molecular mechanism of MBL binding to the vascular endothelium after oxidative stress is unclear; a recent study suggests that
5 activation of the lectin pathway after oxidative stress may be mediated by MBL binding to vascular endothelial cytokeratins, and not to glycoconjugates (Collard et al., *Am. J. Pathol.* 159:1045-1054, (2001)). Other studies have implicated the classical and alternative pathways in the pathogenesis of ischemia/reperfusion injury and the role of the lectin pathway in this disease remains controversial (Riedermann, N.C., et al., *Am. J.*
10 *Pathol.* 162:363-367, 2003).

A recent study has shown that MASP-1 (and possibly also MASP-3) is required to convert the alternative pathway activation enzyme Factor D from its zymogen form into its enzymatically active form (see Takahashi M. et al., *J Exp Med* 207(1):29-37 (2010)). The physiological importance of this process is underlined by the absence of alternative
15 pathway functional activity in plasma of MASP-1/3-deficient mice. Proteolytic generation of C3b from native C3 is required for the alternative pathway to function. Since the alternative pathway C3 convertase (C3bBb) contains C3b as an essential subunit, the question regarding the origin of the first C3b via the alternative pathway has presented a puzzling problem and has stimulated considerable research.

C3 belongs to a family of proteins (along with C4 and α -2 macroglobulin) that contain a rare posttranslational modification known as a thioester bond. The thioester group is composed of a glutamine whose terminal carbonyl group forms a covalent thioester linkage with the sulfhydryl group of a cysteine three amino acids away. This bond is unstable and the electrophilic glutamyl-thioester can react with nucleophilic
25 moieties such as hydroxyl or amino groups and thus form a covalent bond with other molecules. The thioester bond is reasonably stable when sequestered within a hydrophobic pocket of intact C3. However, proteolytic cleavage of C3 to C3a and C3b results in exposure of the highly reactive thioester bond on C3b and, following nucleophilic attack by adjacent moieties comprising hydroxyl or amino groups, C3b
30 becomes covalently linked to a target. In addition to its well-documented role in covalent attachment of C3b to complement targets, the C3 thioester is also thought to have a pivotal role in triggering the alternative pathway. According to the widely accepted "tick-over theory", the alternative pathway is initiated by the generation of a fluid-phase

convertase, iC3Bb, which is formed from C3 with hydrolyzed thioester (iC3; C3(H₂O)) and factor B (Lachmann, P.J., et al., *Springer Semin. Immunopathol.* 7:143-162, (1984)). The C3b-like C3(H₂O) is generated from native C3 by a slow spontaneous hydrolysis of the internal thioester in the protein (Pangburn, M.K., et al., *J. Exp. Med.* 154:856-867, 5 1981). Through the activity of the C3(H₂O)Bb convertase, C3b molecules are deposited on the target surface thereby initiating the alternative pathway.

Very little is known about the initiators of activation of the alternative pathway. Activators are thought to include yeast cell walls (zymosan), many pure polysaccharides, rabbit erythrocytes, certain immunoglobulins, viruses, fungi, bacteria, animal tumor cells, 10 parasites, and damaged cells. The only feature common to these activators is the presence of carbohydrate, but the complexity and variety of carbohydrate structures has made it difficult to establish the shared molecular determinants which are recognized. It has been widely accepted that alternative pathway activation is controlled through the fine balance between inhibitory regulatory components of this pathway, such as Factor H, 15 Factor I, DAF, and CR1, and properdin, which is the only positive regulator of the alternative pathway (see Schwaebler W.J. and Reid K.B., *Immunol Today* 20(1):17-21 (1999)).

In addition to the apparently unregulated activation mechanism described above, the alternative pathway can also provide a powerful amplification loop for the 20 lectin/classical pathway C3 convertase (C4b2a) since any C3b generated can participate with factor B in forming additional alternative pathway C3 convertase (C3bBb). The alternative pathway C3 convertase is stabilized by the binding of properdin. Properdin extends the alternative pathway C3 convertase half-life six to ten fold. Addition of C3b to the alternative pathway C3 convertase leads to the formation of the alternative pathway 25 C5 convertase.

All three pathways (i.e., the classical, lectin and alternative) have been thought to converge at C5, which is cleaved to form products with multiple proinflammatory effects. The converged pathway has been referred to as the terminal complement pathway. C5a is the most potent anaphylatoxin, inducing alterations in smooth muscle and vascular tone, 30 as well as vascular permeability. It is also a powerful chemotaxin and activator of both neutrophils and monocytes. C5a-mediated cellular activation can significantly amplify inflammatory responses by inducing the release of multiple additional inflammatory mediators, including cytokines, hydrolytic enzymes, arachidonic acid metabolites, and

reactive oxygen species. C5 cleavage leads to the formation of C5b-9, also known as the membrane attack complex (MAC). There is now strong evidence that sublytic MAC deposition may play an important role in inflammation in addition to its role as a lytic pore-forming complex.

5 In addition to its essential role in immune defense, the complement system contributes to tissue damage in many clinical conditions. Thus, there is a pressing need to develop therapeutically effective complement inhibitors to prevent these adverse effects.

SUMMARY

10 This summary is provided to introduce a selection of concepts in a simplified form that are further described below in the Detailed Description. This summary is not intended to identify key features of the claimed subject matter, nor is it intended to be used as an aid in determining the scope of the claimed subject matter.

In one aspect, the present invention provides a method of inhibiting microvascular
15 endothelial cell injury and/or thrombus formation in a subject suffering from a thrombotic microangiopathy (TMA) comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation. In some embodiments, the subject is suffering from, or at risk for developing a TMA selected from the group consisting of hemolytic uremic
20 syndrome (aHUS), thrombotic thrombocytopenic purpura (TTP) and atypical hemolytic uremic syndrome (HUS). In some embodiments, prior to administration of the composition the subject is determined to exhibit one or more symptoms selected from the group consisting of (i) anemia, (ii) thrombocytopenia (iii) renal insufficiency and (iv) rising creatinine, and the composition is administered in an effective amount and for a
25 sufficient time period to improve said one or more symptoms. In some embodiments, the MASP-2 inhibitory agent is an anti-MASP-2 antibody, or fragment thereof. In some embodiments, the MASP-2 inhibitory agent is an anti-MASP-2 monoclonal antibody, or fragment thereof that specifically binds to a portion of SEQ ID NO:6. In some embodiments, the MASP-2 inhibitory agent inhibits microvascular endothelial cell injury.

30 In another aspect, the invention provides a method of inhibiting MASP-2-dependent complement activation in a subject suffering from or at risk for developing atypical hemolytic uremic syndrome (aHUS), comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory agent effective to inhibit

MASP-2 dependent complement activation. In one embodiment, prior to administration of the composition the subject is determined to exhibit one or more symptoms selected from the group consisting of (i) anemia, (ii) thrombocytopenia (iii) renal insufficiency and (iv) rising creatinine, and the composition is administered in an effective amount and
5 for a sufficient time period to improve said one or more symptoms. In one embodiment, the subject is suffering from or at risk for developing non-Factor H-dependent aHUS. In one embodiment, the subject is suffering from aHUS associated with factor I, factor B, or membrane cofactor CD46. In one embodiment, the MASP-2 inhibitory agent is an anti-MASP-2 antibody, or fragment thereof, such as an anti-MASP-2 monoclonal antibody, or
10 fragment thereof that specifically binds to a portion of SEQ ID NO:6. In one embodiment, the MASP-2 inhibitory agent inhibits microvascular endothelial cell injury. In one embodiment, the MASP-2 inhibitory agent inhibits thrombus formation.

In another aspect, the invention provides a method for reducing the likelihood that a subject at risk for developing atypical hemolytic uremic syndrome (aHUS) will suffer
15 clinical symptoms associated with aHUS. The method according to this aspect of the invention comprises (a) determining the presence of a genetic marker in the subject known to be associated with aHUS; (b) periodically monitoring the subject to determine the presence or absence of at least one symptom selected from the group consisting of anemia, thrombocytopenia, renal insufficiency and rising creatinine; and (c)
20 administering to the subject a composition comprising an amount of a MASP-2 inhibitory agent effective to inhibit MASP-2-dependent complement activation upon the determination of the presence of at least one of anemia, thrombocytopenia, renal insufficiency or rising creatinine, wherein the composition is administered in an effective amount and for a sufficient time period to improve said one or more symptoms. In one
25 embodiment, the MASP-2 inhibitory agent is an anti-MASP-2 antibody, or fragment thereof, such as an anti-MASP-2 monoclonal antibody, or fragment thereof that specifically binds to a portion of SEQ ID NO:6. In one embodiment of the method, step (a) comprises performing a genetic screening test on a sample obtained from the subject and identifying the presence of at least one genetic marker associated with aHUS in a
30 gene selected from the group consisting of complement factor H (CFH), factor I (CFI), factor B (CFB), membrane cofactor CD46, C3, complement factor H-related protein (CFHR1), anticoagulant protein thrombomodulin (THBD), complement factor H-related protein 3 (CFHR3) and complement factor H-related protein 4 (CFHR4). In one

embodiment, the method further comprises monitoring the subject for the occurrence of an event known to be associated with triggering aHUS clinical symptoms and administering to the subject the composition comprising the MASP-2 inhibitory agent prior to, during, or after the occurrence of the triggering event. In one embodiment, the event associated with triggering aHUS clinical symptoms is selected from the group consisting of drug exposure, infection, malignancy, injury, organ or tissue transplant and pregnancy. In one embodiment, the infection is a bacterial infection. In one embodiment, the composition is administered subcutaneously. In one embodiment, the MASP-2 inhibitory agent inhibits microvascular endothelial cell injury. In one embodiment, the MASP-2 inhibitory agent inhibits thrombus formation.

In another aspect, the invention provides a method of inhibiting MASP-2-dependent complement activation in a subject suffering from, or at risk for developing, atypical hemolytic uremic syndrome (aHUS) secondary to an infection, comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory agent effective to inhibit MASP-2 complement activation. In one embodiment, the subject is suffering from, or at risk for developing non-enteric aHUS associated with an *S. pneumonia* infection. In one embodiment, the MASP-2 inhibitory agent is an anti-MASP-2 antibody, or fragment thereof, such as an anti-MASP-2 monoclonal antibody, or fragment thereof that specifically binds to a portion of SEQ ID NO:6. In one embodiment, the MASP-2 inhibitory agent inhibits microvascular endothelial cell injury. In one embodiment, the MASP-2 inhibitory agent inhibits thrombus formation.

In another aspect, the invention provides a method of treating a subject suffering from atypical hemolytic uremic syndrome (aHUS) comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory agent effective to inhibit MASP-2 dependent complement activation, wherein the administration of the MASP-2 inhibitory agent is administered via an intravenous catheter or other catheter delivery method. In one embodiment, the method further comprises treating the patient with plasmapheresis. In one embodiment, the composition comprising the MASP-2 inhibitory agent is administered in the absence of plasmapheresis. In one embodiment, the composition comprising the MASP-2 inhibitory agent is administered via a catheter for a first time period, further comprising administering the composition comprising the MASP-2 inhibitory agent for a second time period, wherein the composition is administered subcutaneously during the second time period. In one embodiment, the

method further comprises periodically determining the level of at least one complement factor, wherein the determination of a reduced level of the at least one complement factor in comparison to a standard value or a healthy subject is indicative of the need for continued treatment with the composition. In one embodiment, the MASP-2 inhibitory agent is an anti-MASP-2 antibody, or fragment thereof, such as an anti-MASP-2 monoclonal antibody, or fragment thereof that specifically binds to a portion of SEQ ID NO:6. In one embodiment, the MASP-2 inhibitory agent inhibits microvascular endothelial cell injury. In one embodiment, the MASP-2 inhibitory agent inhibits thrombus formation.

10 In another aspect, the invention provides a method of treating a subject suffering from thrombotic thrombocytopenic purpura (TTP), or exhibiting symptoms consistent with a diagnosis of TTP, comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory agent effective to inhibit MASP-2-dependent complement activation, wherein the administration of the MASP-2 inhibitory agent is administered to the subject via an intravenous catheter or other catheter delivery method. In one embodiment, the subject exhibits at least one or more symptoms selected from the group consisting of central nervous system involvement, thrombocytopenia, severe cardiac involvement, severe pulmonary involvement, gastro-intestinal infarction and gangrene. In one embodiment, the subject tests positive for the presence of an inhibitor of ADAMTS13, and the method further comprises administering an immunosuppressant to the subject. In one embodiment, the composition comprising the MASP-2 inhibitory agent is administered for a first time period in the absence of plasmapheresis. In one embodiment, the subject tests positive for the presence of an inhibitor of ADAMTS-13, and the method further comprises administering ADAMTS-13. In one embodiment, the method further comprises treating the patient with plasmapheresis. In one embodiment, the composition comprising the MASP-2 inhibitory agent is administered in the presence of plasmapheresis. In one embodiment, the composition comprising the MASP-2 inhibitory agent is administered via a catheter for a first time period, further comprising administering the composition comprising the MASP-2 inhibitory agent for a second time period, wherein the composition is administered subcutaneously during the second time period. In one embodiment, the method further comprises periodically determining the level of at least one complement factor, wherein the determination of a reduced level of

the at least one complement factor in comparison to a standard value or a healthy subject is indicative of the need for continued treatment with the composition. In one embodiment, the MASP-2 inhibitory agent is an anti-MASP-2 antibody, or fragment thereof, such as an anti-MASP-2 monoclonal antibody, or fragment thereof that
5 specifically binds to a portion of SEQ ID NO:6. In one embodiment, the MASP-2 inhibitory agent inhibits microvascular endothelial cell injury. In one embodiment, the MASP-2 inhibitory agent inhibits thrombus formation.

In another aspect, the invention provides a method of treating a subject suffering from refractory thrombotic thrombocytopenic purpura (TTP) comprising administering to
10 the subject a composition comprising an amount of a MASP-2 inhibitory agent effective to inhibit MASP-2 dependent complement activation. In one embodiment, the composition is administered subcutaneously. In one embodiment, the method further comprises periodically determining the level of at least one complement factor, wherein the determination of a reduced level of the at least one complement factor in comparison
15 to a standard value or a healthy subject is indicative of the need for continued treatment with the composition. In one embodiment, the MASP-2 inhibitory agent is an anti-MASP-2 antibody, or fragment thereof, such as an anti-MASP-2 monoclonal antibody, or fragment thereof that specifically binds to a portion of SEQ ID NO:6. In one embodiment, the MASP-2 inhibitory agent inhibits microvascular endothelial cell injury.
20 In one embodiment, the MASP-2 inhibitory agent inhibits thrombus formation.

In another aspect, the present invention provides a method of inhibiting MASP-2-dependent complement activation in a subject suffering from, or at risk for developing a thrombotic microangiopathy (TMA), wherein the TMA is at least one of (i) a TMA secondary to cancer; (ii) a TMA secondary to chemotherapy, or (iii) a TMA secondary to
25 transplantation, comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory agent effective to inhibit MASP-2-dependent complement activation. In some embodiments, the subject is suffering from, or is at risk for developing a TMA secondary to cancer, and the MASP-2 inhibitory agent is administered systemically to the subject in an amount effective to reduce the risk of
30 developing TMA, or reduce the severity of TMA. In some embodiments, the subject is suffering from, or is at risk for developing a TMA secondary to chemotherapy, and the MASP-2 inhibitory agent is administered systemically to the subject prior to, during, or after chemotherapy, in an amount effective to reduce the risk of developing TMA, or

reduce the severity of TMA. In some embodiments, the subject is suffering from, or is at risk for developing a TMA secondary to transplantation and the MASP-2 inhibitory agent is administered systemically to the subject prior to, during, or after the transplant procedure, in an amount effective to reduce the risk of developing TMA, or reduce the severity of TMA. In some embodiments the transplant procedure is an allogeneic hematopoietic stem cell transplant. In some embodiments, the subject has previously undergone, or is currently undergoing, treatment with a terminal complement inhibitor that inhibits cleavage of complement protein C5. In some embodiments, the method further comprises administering to the subject a terminal complement inhibitor that inhibits cleavage of complement protein C5, such as a humanized anti-C5 antibody or antigen-binding fragment thereof, such as eculizumab.

In another aspect, the invention provides a method of inhibiting MASP-2-dependent complement activation in a subject suffering from or at risk for developing Upshaw-Schulman Syndrome (USS) comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory agent effective to inhibit MASP-2 dependent complement activation. In some embodiments, the method comprises treating a subject at risk for developing USS, wherein the method comprises administering an amount of a MASP-2 inhibitory agent for a time period effective to ameliorate or prevent one of more clinical symptoms associated with TTP. In some embodiments, the method further comprises periodically monitoring the subject and administering the MASP-2 inhibitory agent upon the presence of an event known to be associated with triggering TTP clinical symptoms. In some embodiments, the method further comprises periodically monitoring the subject and administering the MASP-2 inhibitory agent upon the determination of the presence of anemia, thrombocytopenia or rising creatine. In some embodiments, the subject has previously undergone, or is currently undergoing, treatment with a terminal complement inhibitor that inhibits cleavage of complement protein C5. In some embodiments, the method further comprises administering to the subject a terminal complement inhibitor that inhibits cleavage of complement protein C5, such as a humanized anti-C5 antibody or antigen-binding fragment thereof, such as eculizumab.

In another aspect, the invention provides a method of inhibiting MASP-2-dependent complement activation in a subject suffering from Degos disease, comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory

agent effective to inhibit MASP-2-dependent complement activation. In some embodiments, the subject has previously undergone, or is currently undergoing, treatment with a terminal complement inhibitor that inhibits cleavage of complement protein C5. In some embodiments, the method further comprises administering to the subject a terminal complement inhibitor that inhibits cleavage of complement protein C5, such as a humanized anti-C5 antibody or antigen-binding fragment thereof, such as eculizumab.

In another aspect, the invention provides a method of inhibiting MASP-2-dependent complement activation in a subject suffering from Catastrophic Antiphospholipid Syndrome (CAPS), comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory agent effective to inhibit MASP-2-dependent complement activation. In some embodiments, the subject has previously undergone, or is currently undergoing, treatment with a terminal complement inhibitor that inhibits cleavage of complement protein C5. In some embodiments, the method further comprises administering to the subject a terminal complement inhibitor that inhibits cleavage of complement protein C5, such as a humanized anti-C5 antibody or antigen-binding fragment thereof, such as eculizumab.

In some embodiments of any of the disclosed methods of the invention, the MASP-2 inhibitory agent is a MASP-2 inhibitory antibody or fragment thereof. In some embodiments, the MASP-2 inhibitory antibody has reduced effector function. In some embodiments, the MASP-2 inhibitory antibody does not substantially inhibit the classical pathway. In some embodiments, the MASP-2 inhibitory agent is an anti-MASP-2 monoclonal antibody, or fragment thereof that specifically binds to a portion of SEQ ID NO:6. In some embodiments, the anti-MASP-2 antibody or fragment thereof is selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody and a human antibody. In some embodiments, the MASP-2 inhibitory antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂. In some embodiments, the MASP-2 inhibitory antibody is a single-chain molecule. In some embodiments, the MASP-2 inhibitory antibody is selected from the group consisting of an IgG1 molecule, an IgG2 and an IgG4 molecule. In some embodiments, the MASP-2 inhibitory antibody is an IgG4 molecule comprising a S228P mutation. In some embodiments, the MASP-2 inhibitory antibody binds human MASP-2 with a K_D of 10 nM or less. In some embodiments, the MASP-2 inhibitory antibody binds an epitope in the CCP1 domain of

MASP-2. In some embodiments, the MASP-2 inhibitory antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC₅₀ of 10 nM or less. In some embodiments, the MASP-2 inhibitory antibody inhibits C3b deposition in 90% human serum with an IC₅₀ of 30 nM or less.

5 In some embodiments of any of the disclosed methods of the invention the MASP-2 inhibitory monoclonal antibody, or antigen-binding fragment thereof, comprises: (a) a heavy-chain variable region comprising: i) a heavy chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a
10 heavy-chain CDR-H3 comprising the amino acid sequence from 95-102 of SEQ ID NO:67 and (b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70.
15 In some embodiments, the MASP-2 inhibitory monoclonal antibody comprises a heavy-chain variable region set forth as SEQ ID NO:67 and a light-chain variable region set forth as SEQ ID NO:70. In some embodiments, the MASP-2 inhibitory antibody or antigen binding-fragment thereof specifically recognizes at least part of an epitope recognized by a reference antibody comprising a heavy chain variable region as set forth
20 in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70.

 In another aspect of the invention, methods are provided for inhibiting thrombus formation in a subject suffering from atypical hemolytic uremic syndrome (aHUS), comprising administering to the subject an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, effective to inhibit MASP-2-dependent complement
25 activation. In some embodiments, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a subject suffering from aHUS by at least 40% as compared to untreated serum. In some embodiments, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a subject suffering from aHUS at a level of at least 20% greater (e.g., at least 30% greater, at least 40% greater, or at least 50% greater) than
30 its inhibitory effect on C5b-9 deposition in the serum from the same subject. In some embodiments, the subject is in the acute phase of aHUS. In some embodiments, the subject is in the remission phase of aHUS. In some embodiments, the MASP-2 inhibitory antibody is a monoclonal antibody, or fragment thereof that specifically binds to a portion

of SEQ ID NO:6. In some embodiments, the MASP-2 inhibitory antibody or fragment thereof is selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody and a human antibody. In some embodiments, the MASP-2 inhibitory antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂. In some embodiments, the MASP-2 inhibitory antibody is a single-chain molecule. In some embodiments, the MASP-2 inhibitory antibody is selected from the group consisting of an IgG1 molecule, an IgG2 and an IgG4 molecule. In some embodiments, the MASP-2 inhibitory antibody is an IgG4 molecule comprising a S228P mutation. In some embodiments, the MASP-2 inhibitory antibody binds human MASP-2 with a K_D of 10 nM or less. In some embodiments, the MASP-2 inhibitory antibody binds an epitope in the CCP1 domain of MASP-2. In some embodiments, the MASP-2 inhibitory antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC₅₀ of 10 nM or less. In some embodiments, the MASP-2 inhibitory antibody inhibits C3b deposition in 90% human serum with an IC₅₀ of 30 nM or less. In some embodiments the MASP-2 inhibitory monoclonal antibody, or antigen-binding fragment thereof, comprises: (a) a heavy-chain variable region comprising: i) a heavy chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-102 of SEQ ID NO:67 and (b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70. In some embodiments, the MASP-2 inhibitory monoclonal antibody comprises a heavy-chain variable region set forth as SEQ ID NO:67 and a light-chain variable region set forth as SEQ ID NO:70. In some embodiments, the MASP-2 inhibitory antibody or antigen binding-fragment thereof specifically recognizes at least part of an epitope recognized by a reference antibody comprising a heavy chain variable region as set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70.

In another aspect, the present invention provides a method of treating a subject suffering from plasma therapy-resistant aHUS comprising administering to the subject a

composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation. In one embodiment, the MASP-2 inhibitory antibody is an anti-MASP-2 monoclonal antibody, or fragment thereof that specifically binds to human MASP-2. In one embodiment, the antibody or fragment thereof is
5 selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody, and a human antibody. In one embodiment, the subject has previously undergone, or is currently undergoing, treatment with a terminal complement inhibitor that inhibits cleavage of complement protein C5, such as wherein the terminal complement inhibitor is a
10 humanized anti-C5 antibody or antigen-binding fragment thereof. In one embodiment, the method further comprises treating the patient with plasmapheresis. In one embodiment, the composition comprising the MASP-2 inhibitory antibody is administered in the absence of plasmapheresis.

In another aspect, the present invention provides a method of treating a subject
15 suffering from TMA associated with hematopoietic stem cell transplant comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation. In one embodiment, the MASP-2 inhibitory antibody is an anti-MASP-2 monoclonal antibody, or fragment thereof that specifically binds to human MASP-2. In one embodiment, the
20 antibody or fragment thereof is selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody, and a human antibody. In one embodiment, the subject has previously undergone, or is currently undergoing, treatment with a terminal complement inhibitor that inhibits cleavage of complement protein C5. In one embodiment, the subject is
25 suffering from a TMA associated with hematopoietic stem cell transplant that is resistant to treatment with a platelet transfusion and/or resistant to treatment with plasmapheresis. In one embodiment,
the MASP-2 inhibitory antibody is administered in an amount effective to improve at least one or more clinical parameters associated with TMA associated with hematopoietic
30 stem cell transplant, such as an increase in platelet count (e.g., at least double, at least triple, at least quadruple the platelet count prior to treatment), an increase in haptoglobin, and/or a decrease in lactate dehydrogenase.

In another aspect, the present invention provides a method of treating a human subject suffering from persistent TMA associated with hematopoietic stem cell transplant (HSCT-TMA) comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation. In one embodiment, the method further comprises identifying a human subject having persistent TMA associated with hematopoietic stem cell transplant prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation. In one embodiment, the MASP-2 inhibitory antibody is a monoclonal antibody, or fragment thereof that specifically binds to human MASP-2. In one embodiment, the antibody or antigen-binding fragment thereof is selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody, and a human antibody. In one embodiment, the MASP-2 inhibitory antibody does not substantially inhibit the classical pathway. In one embodiment, the MASP-2 inhibitory antibody inhibits C3b deposition in 90% human serum with an IC₅₀ of 30 nM or less. In one embodiment, the MASP-2 inhibitory antibody or antigen-binding fragment thereof, comprises a heavy chain variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid sequence set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ ID NO:70. In one embodiment, the MASP-2 inhibitory antibody is administered to the patient in the absence of plasmapheresis. In one embodiment, the subject has previously undergone, or is currently undergoing, treatment with a humanized anti-C5 antibody or antigen-binding fragment thereof, such as wherein the terminal complement inhibitor is a humanized anti-C5 antibody or antigen-binding fragment thereof. In one embodiment, the MASP-2 inhibitory antibody is delivered to the subject systemically. In one embodiment, the MASP-2 inhibitory antibody or antigen-binding fragment thereof is administered in an amount effective to improve at least one or more of the following clinical parameters associated with persistent TMA associated with hematopoietic stem cell transplant: (i) an increase in platelet count (e.g., at least double, at least triple, at least quadruple the platelet count prior to treatment)); (ii) an increase in haptoglobin; (iii) a decrease in lactate dehydrogenase (LDH); and/or (iv) a decrease in creatinine.

In another aspect, the present invention provides a method of treating a human subject suffering from, or at risk for developing graft-versus-host disease (GVHD) comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation. In one embodiment, the subject has previously undergone, or is currently undergoing, or will undergo a hematopoietic stem cell transplant. In one embodiment, the method further comprises identifying a human subject suffering from, or at risk for developing graft-versus-host disease prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation. In one embodiment, the MASP-2 inhibitory antibody is a monoclonal antibody, or fragment thereof that specifically binds to human MASP-2. In one embodiment, the MASP-2 inhibitory antibody or fragment thereof is selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody, and a human antibody. In one embodiment, the MASP-2 inhibitory antibody does not substantially inhibit the classical pathway. In one embodiment, the MASP-2 inhibitory antibody inhibits C3b deposition in 90% human serum with an IC₅₀ of 30 nM or less. In one embodiment, the MASP-2 inhibitory antibody is delivered to the subject systemically. In one embodiment, the subject is suffering from acute GVHD. In one embodiment, the subject is suffering from steroid-resistant GVHD. In one embodiment, the MASP-2 inhibitory antibody or antigen-binding fragment thereof, comprises a heavy chain variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid sequence set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ ID NO:70.

In another aspect, the present invention provides a method of treating, preventing or ameliorating one or more neurological symptoms associated with graft-versus-host disease or TMA comprising administering to a subject in need thereof a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation. In one embodiment, the one or more neurological symptoms associated with graft-versus-host disease or TMA is selected from the group consisting of asthenia, paresthesias,

tetraplegia, sensorimotor deficit, dysautonomic polyneuropathy, and/or neurogenic bladder. In one embodiment, the subject has received a hematopoietic stem cell transplant and the subject is suffering from one or more neurological symptoms selected from the group consisting of paresthesias, tetraplegia and neurogenic bladder. In one
5 embodiment, the subject has received a hematopoietic stem cell transplant and is suffering from graft-versus-host disease. In one embodiment, the subject has received a hematopoietic stem cell transplant and is suffering from HSCT-TMA. In one embodiment, the MASP-2 inhibitory antibody is a monoclonal antibody, or fragment thereof that specifically binds to human MASP-2. In one embodiment, the antibody or
10 fragment thereof is selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody, and a human antibody. In one embodiment, the MASP-2 inhibitory antibody does not substantially inhibit the classical pathway. In one embodiment, the MASP-2 inhibitory antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less. In
15 one embodiment, the MASP-2 inhibitory antibody is delivered to the subject systemically. In one embodiment, the method further comprises identifying a human subject suffering from one or more neurological symptoms associated with with hematopoietic stem cell transplant prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding
20 fragment thereof, effective to inhibit MASP-2-dependent complement activation. In one embodiment, the MASP-2 inhibitory antibody or antigen-binding fragment thereof, comprises a heavy chain variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid sequence set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ
25 ID NO:70.

In another aspect, the present invention provides a method of treating a human subject suffering from, or at risk for developing diffuse alveolar hemorrhage (DAH) associated with hematopoietic stem cell transplant comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or
30 antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation. In one embodiment, the subject has previously undergone, or is currently undergoing, or will undergo a hematopoietic stem cell transplant. In one embodiment, the method further comprises identifying a human subject suffering from, or at risk for

developing diffuse alveolar hemorrhage (DAH) prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation. In one embodiment, the MASP-2 inhibitory antibody is a monoclonal
5 antibody, or fragment thereof that specifically binds to human MASP-2. In one embodiment, the MASP-2 inhibitory antibody or fragment thereof is selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody, and a human antibody. In one embodiment, the MASP-2 inhibitory antibody does not substantially inhibit the classical
10 pathway. In one embodiment, the MASP-2 inhibitory antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less. In one embodiment, the MASP-2 inhibitory antibody is delivered to the subject systemically. In one embodiment, the MASP-2 inhibitory antibody or antigen-binding fragment thereof, comprises a heavy chain variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid
15 sequence set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ ID NO:70. In another aspect, the present invention provides compositions for inhibiting the adverse effects of MASP-2-dependent complement activation, comprising a therapeutically effective amount of a MASP-2 inhibitory agent, such as a MASP-2 inhibitory antibody
20 and a pharmaceutically acceptable carrier. Methods are also provided for manufacturing a medicament for use in inhibiting the adverse effects of MASP-2-dependent complement activation in living subjects in need thereof, comprising a therapeutically effective amount of a MASP-2 inhibitory agent in a pharmaceutical carrier. Methods are also provided for manufacturing medicaments for use in inhibiting MASP-2-dependent
25 complement activation for treatment of each of the conditions, diseases and disorders described herein below.

In another aspect, the present invention provides a method of treating a human subject suffering from, or at risk for developing veno-occlusive disease (VOD) associated with hematopoietic stem cell transplant comprising administering to the subject a
30 composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation. In one embodiment, the subject has previously undergone, or is currently undergoing, or will undergo a hematopoietic stem cell transplant. In one embodiment, the method further

comprises identifying a human subject suffering from, or at risk for developing veno-occlusive disease (VOD) prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation. In one
5 embodiment, the MASP-2 inhibitory antibody is a monoclonal antibody, or fragment thereof that specifically binds to human MASP-2. In one embodiment, the MASP-2 inhibitory antibody or fragment thereof is selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody, and a human antibody. In one embodiment, the MASP-2
10 inhibitory antibody does not substantially inhibit the classical pathway. In one embodiment, the MASP-2 inhibitory antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less. In one embodiment, the MASP-2 inhibitory antibody is delivered to the subject systemically. In one embodiment, the MASP-2 inhibitory antibody or antigen-binding fragment thereof, comprises a heavy chain
15 variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid sequence set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ ID NO:70. In another aspect, the present invention provides compositions for inhibiting the adverse effects of MASP-2-dependent complement activation, comprising a therapeutically effective
20 amount of a MASP-2 inhibitory agent, such as a MASP-2 inhibitory antibody and a pharmaceutically acceptable carrier. Methods are also provided for manufacturing a medicament for use in inhibiting the adverse effects of MASP-2-dependent complement activation in living subjects in need thereof, comprising a therapeutically effective amount of a MASP-2 inhibitory agent in a pharmaceutical carrier. Methods are also
25 provided for manufacturing medicaments for use in inhibiting MASP-2-dependent complement activation for treatment of each of the conditions, diseases and disorders described herein below.

The methods, compositions and medicaments of the invention are useful for
30 inhibiting the adverse effects of MASP-2-dependent complement activation *in vivo* in mammalian subjects, including humans suffering from or at risk for developing a thrombotic microangiopathy (TMA), and/or a subject suffering from, or at risk for developing GVHD, and/or a subject suffering from, or at risk for developing post-stem

cell transplant diffuse alveolar hemorrhage and/or a subject suffering from, or at risk for developing veno-occlusive disease (VOD) as further described herein.

DESCRIPTION OF THE DRAWINGS

5 The foregoing aspects and many of the attendant advantages of this invention will become more readily appreciated as the same become better understood by reference to the following detailed description, when taken in conjunction with the accompanying drawings, wherein:

FIGURE 1 is a diagram illustrating the genomic structure of human MASP-2;

10 FIGURE 2A is a schematic diagram illustrating the domain structure of human MASP-2 protein;

FIGURE 2B is a schematic diagram illustrating the domain structure of human MASP-2 protein;

FIGURE 3 is a diagram illustrating the murine MASP-2 knockout strategy;

15 FIGURE 4 is a diagram illustrating the human MASP-2 minigene construct;

FIGURE 5A presents results demonstrating that MASP-2-deficiency leads to the loss of lectin-pathway-mediated C4 activation as measured by lack of C4b deposition on mannan, as described in Example 2;

20 FIGURE 5B presents results demonstrating that MASP-2-deficiency leads to the loss of lectin-pathway-mediated C4 activation as measured by lack of C4b deposition on zymosan, as described in Example 2;

FIGURE 5C presents results demonstrating the relative C4 activation levels of serum samples obtained from MASP-2^{+/-}, MASP-2^{-/-} and wild-type strains as measured by C4b deposition on mannan and on zymosan, as described in Example 2;

25 FIGURE 6 presents results demonstrating that the addition of murine recombinant MASP-2 to MASP-2^{-/-} serum samples recovers lectin-pathway-mediated C4 activation in a protein concentration dependent manner, as measured by C4b deposition on mannan, as described in Example 2;

30 FIGURE 7 presents results demonstrating that the classical pathway is functional in the MASP-2^{-/-} strain, as described in Example 8;

FIGURE 8A presents results demonstrating that anti-MASP-2 Fab2 antibody #11 inhibits C3 convertase formation, as described in Example 10;

FIGURE 8B presents results demonstrating that anti-MASP-2 Fab2 antibody #11 binds to native rat MASP-2, as described in Example 10;

FIGURE 8C presents results demonstrating that anti-MASP-2 Fab2 antibody #41 inhibits C4 cleavage, as described in Example 10;

5 FIGURE 9 presents results demonstrating that all of the anti-MASP-2 Fab2 antibodies tested that inhibited C3 convertase formation also were found to inhibit C4 cleavage, as described in Example 10;

10 FIGURE 10 is a diagram illustrating the recombinant polypeptides derived from rat MASP-2 that were used for epitope mapping of the anti-MASP-2 blocking Fab2 antibodies, as described in Example 11;

FIGURE 11 presents results demonstrating the binding of anti-MASP-2 Fab2 #40 and #60 to rat MASP-2 polypeptides, as described in Example 11;

15 FIGURE 12 presents results demonstrating the blood urea nitrogen clearance for wild type (+/+) and MASP-2 (-/-) mice at 24 and 48 hours after reperfusion in a renal ischemia/reperfusion injury model, as described in Example 12;

FIGURE 13A presents results showing the baseline VEGF protein levels in RPE-choroid complex isolated from wild type (+/+) and MASP-2 (-/-) mice, as described in Example 13;

20 FIGURE 13B presents results showing the VEGF protein levels in RPE-choroid complex at day 3 in wild type (+/+) and MASP-2 (-/-) mice following laser induced injury in a macular degeneration model, as described in Example 13;

FIGURE 14 presents results showing the mean choroidal neovascularization (CNV) volume at day seven following laser induced injury in wild type (+/+) and MASP-2 (-/-) mice, as described in Example 13;

25 FIGURES 15A and 15B present dose response curves for the inhibition of C4b deposition (FIG. 15A) and the inhibition of thrombin activation (FIG 15B) following the administration of a MASP-2 Fab2 antibody in normal rat serum, as described in Example 14;

30 FIGURES 16A and 16B present measured platelet aggregation (expressed as aggregate area) in MASP-2 (-/-) mice (FIG. 16B) as compared to platelet aggregation in untreated wild type mice and wild type mice in which the complement pathway is inhibited by depletory agent cobra venom factor (CVF) and a terminal pathway inhibitor

(C5aR antagonist) (FIGURE 16A) in a localized Schwartzman reaction model of disseminated intravascular coagulation, as described in Example 15;

FIGURE 17 graphically illustrates the blood urea nitrogen (BUN) levels measured in either WT (+/+) (B6) or MASP-2 (-/-) transplant recipient mice of WT (+/+) donor
5 kidneys, as described in Example 16;

FIGURE 18 graphically illustrates the percentage survival of WT (+/+) and MASP-2 (-/-) mice as a function of the number of days after microbial infection in the cecal ligation and puncture (CLP) model, as described in Example 17;

FIGURE 19 graphically illustrates the number of bacteria measured in WT (+/+) and MASP-2 (-/-) after microbial infection in the cecal ligation and puncture (CLP)
10 model, as described in Example 17;

FIGURE 20 is a Kaplan-Mayer plot illustrating the percent survival of WT (+/+), MASP-2 (-/-) and C3 (-/-) mice six days after challenge with intranasal administration of *Pseudomonas aeruginosa*, as described in Example 18;

FIGURE 21 graphically illustrates the level of C4b deposition, measured as % of control, in samples taken at various time points after subcutaneous dosing of either 0.3 mg/kg or 1.0 mg/kg of mouse anti-MASP-2 monoclonal antibody in WT mice, as
15 described in Example 19;

FIGURE 22 graphically illustrates the level of C4b deposition, measured as % of control, in samples taken at various time points after ip dosing of 0.6 mg/kg of mouse anti-MASP-2 monoclonal antibody in WT mice, as described in Example 19;
20

FIGURE 23 graphically illustrates the mean choroidal neovascularization (CNV) volume at day seven following laser induced injury in WT (+/+) mice pre-treated with a single ip injection of 0.3 mg/kg or 1.0 mg/kg mouse anti-MASP-2 monoclonal antibody;
25 as described in Example 20;

FIGURE 24A graphically illustrates the percent survival of MASP-2 (-/-) and WT (+/+) mice after infection with $5 \times 10^8/100 \mu\text{l}$ cfu *N. meningitidis*, as described in Example 21;

FIGURE 24B graphically illustrates the log cfu/ml of *N. meningitidis* recovered at different time points in blood samples taken from the MASP-2 KO (-/-) and WT (+/+) mice infected with 5×10^8 cfu/100 μl *N. meningitidis*, as described in Example 21;
30

FIGURE 25A graphically illustrates the percent survival of MASP-2 KO (-/-) and WT (+/+) mice after infection with 2×10^8 cfu/100 μ l *N. meningitidis*, as described in Example 21;

5 FIGURE 25B graphically illustrates the log cfu/ml of *N. meningitidis* recovered at different time points in blood samples taken from the WT (+/+) mice infected with 2×10^8 cfu/100 μ l *N. meningitidis*, as described in Example 21;

FIGURE 25C graphically illustrates the log cfu/ml of *N. meningitidis* recovered at different time points in blood samples taken from the MASP-2 (-/-) mice infected with 2×10^8 cfu/100 μ l *N. meningitidis*, as described in Example 21;

10 FIGURE 26A graphically illustrates the results of a C3b deposition assay demonstrating that MASP-2 (-/-) mice retain a functional classical pathway, as described in Example 22;

FIGURE 26B graphically illustrates the results of a C3b deposition assay on zymosan coated plates, demonstrating that MASP-2 (-/-) mice retain a functional
15 alternative pathway, as described in Example 22;

FIGURE 27A graphically illustrates myocardial ischemia/reperfusion injury (MIRI)-induced tissue loss following ligation of the left anterior descending branch of the coronary artery (LAD) and reperfusion in C4 (-/-) mice (n=6) and matching WT littermate controls (n=7), showing area at risk (AAR) and infarct size (INF) as described
20 in Example 22;

FIGURE 27B graphically illustrates infarct size (INF) as a function of area at risk (AAR) in C4 (-/-) and WT mice treated as describe in FIGURE 42A, demonstrating that C4 (-/-) mice are as susceptible to MIRI as WT controls (dashed line), as described in Example 22;

25 FIGURE 28A graphically illustrates the results of a C3b deposition assay using serum from WT mice, C4 (-/-) mice and serum from C4 (-/-) mice pre-incubated with mannan, as described in Example 22;

FIGURE 28B graphically illustrates the results of a C3b deposition assay on serum from WT, C4 (-/-), and MASP-2 (-/-) mice mixed with various concentrations of
30 an anti-murine MASP-2 mAb (mAbM11), as described in Example 22;

FIGURE 28C graphically illustrates the results of a C3b deposition assay on human serum from WT (C4 sufficient) and C4 deficient serum, and serum from C4 deficient subjects pre-incubated with mannan, as described in Example 22;

FIGURE 28D graphically illustrates the results of a C3b deposition assay on human serum from WT (C4 sufficient) and C4 deficient subjects mixed with anti-human MASP-2 mAb (mAbH3), as described in Example 22;

5 FIGURE 29A graphically illustrates a comparative analysis of C3 convertase activity in plasma from various complement deficient mouse strains tested either under lectin activation pathway specific assay conditions, or under classical activation pathway specific assay conditions, as described in Example 22;

10 FIGURE 29B graphically illustrates the time-resolved kinetics of C3 convertase activity in plasma from various complement deficient mouse strains tested under lectin activation pathway specific conditions, as described in Example 22;

FIGURE 30 illustrates the results of a Western blot analysis showing activation of human C3, shown by the presence of the a' chain, by thrombin substrates FXIa and FXa, as described in Example 23;

15 FIGURE 31 shows the results of the C3 deposition assay on serum samples obtained from WT, MASP-2 (-/-), F11(-/-), F11(-/-)/C4 (-/-) and C4 (-/-), as described in Example 23;

20 FIGURE 32A is a Kaplan-Meier survival plot showing the percent survival over time after exposure to 7.0 Gy radiation in control mice and in mice treated with anti-murine MASP-2 antibody (mAbM11) or anti-human MASP-2 antibody (mAbH6) as described in Example 29;

FIGURE 32B is a Kaplan-Meier survival plot showing the percent survival over time after exposure to 6.5 Gy radiation in control mice and in mice treated with anti-murine MASP-2 antibody (mAbM11) or anti-human MASP-2 antibody (mAbH6), as described in Example 29;

25 FIGURE 33 is a Kaplan-Meyer plot graphically illustrating the percent survival of MASP-2 KO and WT mice after administration of an infective dose of 2.6×10^7 cfu of *N. meningitidis* serogroup A Z2491, demonstrating that MASP-2 deficient mice are protected from *N. meningitidis* induced mortality, as described in Example 30;

30 FIGURE 34 is a Kaplan-Meyer plot graphically illustrating the percent survival of MASP-2 KO and WT mice after administration of an infective dose of 6×10^6 cfu of *N. meningitidis* serogroup B strain MC58, demonstrating that MASP-2-deficient mice are protected from *N. meningitidis* serogroup B strain MC58 induced mortality, as described in Example 30;

FIGURE 35 graphically illustrates the log cfu/ml of *N. meningitidis* serogroup B strain MC58 recovered at different time points in blood samples taken from the MASP-2 KO and WT mice after i.p. infection with 6×10^6 cfu of *N. meningitidis* serogroup B strain MC58 (n=3 at different time points for both groups of mice, results are expressed as Means $\pm$ SEM) demonstrating that although the MASP-2 KO mice were infected with the same dose of *N. meningitidis* serogroup B strain MC58 as the WT mice, the MASP-2 KO mice have enhanced clearance of bacteraemia as compared to WT, as described in Example 30;

FIGURE 36 graphically illustrates the average illness score of MASP-2 and WT mice at 3, 6, 12 and 24 hours after infection with 6×10^6 cfu/100 μ l *N. meningitidis* Serogroup Serogroup B strain MC58, demonstrating that the MASP-2 deficient mice showed high resistance to the infection, with much lower illness scores at 6 hours, as described in Example 30;

FIGURE 37 is a Kaplan-Meier plot graphically illustrating the percent survival of mice after administration of an infective dose of 4×10^6 /100 μ l cfu *N. meningitidis* Serogroup B strain MC58, followed by administration 3 hours post infection of either inhibitory anti-MASP-2 antibody (1 mg/kg) or control isotype antibody, demonstrating that anti-MASP-2 antibody is effective to treat and improve survival in subjects infected with *N. meningitidis*, as described in Example 31;

FIGURE 38 graphically illustrates the log cfu/ml of viable counts of *N. meningitidis* serogroup B-MC58 recovered at different time points in 20% human serum concentration after i.p. infection with 6.5×10^6 cfu/100 μ l *N. meningitidis* serogroup B strain MC58 at 0, 30, 60 and 90 minutes after incubation in the presence of: (A) normal human serum (NHS) plus human anti-MASP-2 antibody; (B) normal human serum (NHS) plus isotype control antibody; (C) MBL-/- human serum; (D) normal human serum (NHS) and (E) heat inactivated normal human serum (NHS), showing that complement dependent killing of *N. meningitidis* in human serum was significantly enhanced by the addition of the human anti-MASP-2 antibody, as described in Example 32;

FIGURE 39 graphically illustrates the log cfu/ml of viable counts of *N. meningitidis* serogroup B-MC58 recovered at different time points in the mouse sera samples, demonstrating MASP-2 -/- mouse sera has a higher level of bactericidal activity for *N. meningitidis* than WT mouse sera, as described in Example 32;

FIGURE 40 graphically illustrates hemolysis (as measured by hemoglobin release of lysed mouse erythrocytes (Crry/C3-/-) into the supernatant measured by photometry) of mannan-coated murine erythrocytes by human serum over a range of serum concentrations. The sera tested included heat-inactivated (HI) NHS, MBL-/-, NHS +anti-MASP-2 antibody and NHS control, as described in Example 33;

FIGURE 41 graphically illustrates hemolysis (as measured by hemoglobin release of lysed WT mouse erythrocytes into the supernatant measured by photometry) of non-coated murine erythrocytes by human serum over a range of serum concentrations. The sera tested included heat-inactivated (HI) NHS, MBL-/-, NHS +anti-MASP-2 antibody and NHS control, demonstrating that inhibiting MASP-2 inhibits complement-mediated lysis of non-sensitized WT mouse erythrocytes, as described in Example 33;

FIGURE 42 graphically illustrates hemolysis (as measured by hemoglobin release of lysed mouse erythrocytes (CD55/59 -/-) into the supernatant measured by photometry) of non-coated murine erythrocytes by human serum over a range of serum concentrations. The sera tested included heat-inactivated (HI) NHS, MBL-/-, NHS +anti-MASP-2 antibody and NHS control, as described in Example 33;

FIGURE 43 graphically illustrates the percent survival over time (days) after exposure to 8.0 Gy radiation in control mice and in mice treated with anti-human MASP-2 antibody (mAbH6), as described in Example 34;

FIGURE 44 graphically illustrates the time to onset of microvascular occlusion following LPS injection in MASP-2 -/- and WT mice, showing the percentage of mice with thrombus formation measured over 60 minutes, demonstrating that thrombus formation is detected after 15 minutes in WT mice, with up to 80% of the WT mice demonstrated thrombus formation at 60 minutes; in contrast, none of the MASP-2 -/- mice showed any thrombus formation during the 60 minute period (log rank: $p=0.0005$), as described in Example 35;

FIGURE 45 graphically illustrates the percent survival of saline treated control mice ($n=5$) and anti-MASP-2 antibody treated mice ($n=5$) in the STX/LPS-induced model of HUS over time (hours), demonstrating that all of the control mice died by 42 hours, whereas, in contrast, 100 % of the anti-MASP-2 antibody-treated mice survived throughout the time course of the experiment, as described in Example 36;

FIGURE 46 graphically illustrates, as a function of time after injury induction, the percentage of mice with microvascular occlusion in the FITC/Dextran UV model after

treatment with isotype control, or human MASP-2 antibody mAbH6 (10mg/kg) dosed at 16 hours and 1 hour prior to injection of FITC/Dextran, as described in Example 37;

FIGURE 47 graphically illustrates the occlusion time in minutes for mice treated with the human MASP-2 antibody (mAbH6) and the isotype control antibody, wherein the data are reported as scatter-dots with mean values (horizontal bars) and standard error bars (vertical bars). The statistical test used for analysis was the unpaired t test; wherein the symbol “*” indicates $p=0.0129$, as described in Example 37; and

FIGURE 48 graphically illustrates the time until occlusion in minutes for wild-type mice, MASP-2 KO mice, and wild-type mice pre-treated with human MASP-2 antibody (mAbH6) administered i.p. at 10mg/kg 16 hours before, and again 1 hour prior to the induction of thrombosis in the FITC-dextran/light induced endothelial cell injury model of thrombosis with low light intensity (800-1500), as described in Example 37;

FIGURE 49 is a Kaplan-Meier plot showing the percentage of mice with thrombi as a function of time in FITC-Dextran induced thrombotic microangiopathy in mice treated with increasing doses of human MASP-2 inhibitory antibody (mAbH6) or an isotype control antibody, as described in Example 39;

FIGURE 50 graphically illustrates the median time to onset (minutes) of thrombus formation as a function of mAbH6 dose (* $p<0.01$ compared to control), as described in Example 39;

FIGURE 51 is a Kaplan-Meier plot showing the percentage of mice with microvascular occlusion as a function of time in FITC-Dextran induced thrombotic microangiopathy in mice treated with imcreasing doses of human MASP-2 inhibitory antibody (mAbH6) or an isotype control antibody, as described in Example 39;

FIGURE 52 graphically illustrates the median time to microvascular occlusion as a function of mAbH6 dose (* $p<0.05$ compared to control), as described in Example 39;

FIGURE 53A graphically illustrates the level of MAC deposition in the presence or absence of human MASP-2 monoclonal antibody (OMS646) under lectin pathway-specific assay conditions, demonstrating that OMS646 inhibits lectin-mediated MAC deposition with an IC_{50} value of approximately 1 nM, as described in Example 40;

FIGURE 53B graphically illustrates the level of MAC deposition in the presence or absence of human MASP-2 monoclonal antibody (OMS646) under classical pathway-specific assay conditions, demonstrating that OMS646 does not inhibit classical pathway-mediated MAC deposition, as described in Example 40;

FIGURE 53C graphically illustrates the level of MAC deposition in the presence or absence of human MASP-2 monoclonal antibody (OMS646) under alternative pathway-specific assay conditions, demonstrating that OMS646 does not inhibit alternative pathway-mediated MAC deposition, as described in Example 40;

5 FIGURE 54 graphically illustrates the pharmacokinetic (PK) profile of human MASP-2 monoclonal antibody (OMS646) in mice, showing the OMS646 concentration (mean of n=3 animals/groups) as a function of time after administration at the indicated dose, as described in Example 40;

10 FIGURE 55A graphically illustrates the pharmacodynamic (PD) response of human MASP-2 monoclonal antibody (OMS646), measured as a drop in systemic lectin pathway activity in mice following intravenous administration, as described in Example 40;

15 FIGURE 55B graphically illustrates the pharmacodynamic (PD) response of human MASP-2 monoclonal antibody (OMS646), measured as a drop in systemic lectin pathway activity in mice following subcutaneous administration, as described in Example 40;

FIGURE 56 graphically illustrates the inhibitory effect of MASP-2 antibody (OMS646) as compared to sCR1 on aHUS serum-induced C5b-9 deposition on ADP-activated HMEC-1 cells, as described in Example 41;

20 FIGURE 57 graphically illustrates the inhibitory effect of MASP-2 antibody (OMS646) as compared to sCR1 on aHUS serum-induced thrombus formation on ADP-activated HMEC-1 cells, as described in Example 42;

25 FIGURE 58 graphically illustrates the mean change in platelet count from baseline over time (weeks) in subjects suffering from persistent hematopoietic stem cell transplant-associated thrombotic microangiopathy (HSCT-TMA) after treatment with MASP-2 inhibitory antibody (OMS646), as described in Example 46;

FIGURE 59 graphically illustrates the mean change in LDH from baseline over time (weeks) in subjects suffering from persistent (HSCT-TMA) after treatment with MASP-2 inhibitory antibody (OMS646), as described in Example 46;

30 FIGURE 60 graphically illustrates the mean change in haptoglobin from baseline over time (weeks) in subjects suffering from persistent (HSCT-TMA) after treatment with MASP-2 inhibitory antibody (OMS646), as described in Example 46;

FIGURE 61 illustrates the clinical course of a patient post-hematopoietic stem cell transplant (HSCT) that developed HSCT-TMA and graft-versus-host disease (GVHD), demonstrating improvement in HSCT-TMA, GVHD and improvement in neurological symptoms after treatment with MASP-2 inhibitory antibody (OMS646), as described in Example 47;

FIGURE 62A graphically illustrates the level of creatinine over time in the compassionate use patient #1, wherein the vertical line indicates the start of treatment with MASP-2 inhibitory antibody (OMS646), as described in Example 48;

FIGURE 62B graphically illustrates the level of haptoglobin over time in the compassionate use patient #1, wherein the vertical line indicates the start of treatment with MASP-2 inhibitory antibody (OMS646), as described in Example 48;

FIGURE 62C graphically illustrates the level of hemoglobin over time in the compassionate use patient #1, wherein the vertical line indicates the start of treatment with MASP-2 inhibitory antibody (OMS646), as described in Example 48;

FIGURE 62D graphically illustrates the level of LDH over time in the compassionate use patient #1, wherein the vertical line indicates the start of treatment with MASP-2 inhibitory antibody (OMS646), as described in Example 48; and

FIGURE 62E graphically illustrates the level of platelets over time in the compassionate use patient #1, wherein the vertical line indicates the start of treatment with MASP-2 inhibitory antibody (OMS646), as described in Example 48.

25

DESCRIPTION OF THE SEQUENCE LISTING

SEQ ID NO:1 human MASP-2 cDNA

SEQ ID NO:2 human MASP-2 protein (with leader)

SEQ ID NO:3 human MASP-2 protein (mature)

30 SEQ ID NO:4 human MASP-2 cDNA

SEQ ID NO:5 human MASP-2 protein (with leader)

SEQ ID NO:6 human MASP-2 protein (mature)

SEQ ID NO:7 human MASP-2 gDNA (exons 1-6)

ANTIGENS: (IN REFERENCE TO THE MASP-2 MATURE PROTEIN)

- SEQ ID NO:8 CUBI sequence (aa 1-121)
- SEQ ID NO:9 CUBEGF sequence (aa 1-166)
- SEQ ID NO:10 CUBEGFCUBII (aa 1-293)
- 5 SEQ ID NO:11 EGF region (aa 122-166)
- SEQ ID NO:12 serine protease domain (aa 429 – 671)
- SEQ ID NO:13 serine protease domain inactive (aa 610-625 with Ser618
to Ala mutation)
- SEQ ID NO:14 TPLGPKWPEPVFGRL (CUB1 peptide)
- 10 SEQ ID NO:15
- TAPPGYRLRLYFTHFDLELSHLCEYDFVKLSSGAKVLATLC
- GQ (CUBI peptide)
- SEQ ID NO:16 TFRSDYSN (MBL binding region core)
- SEQ ID NO:17 FYSLGSSLDITFRSDYSNEKPFTGF (MBL binding region)
- 15 SEQ ID NO:18 IDECQVAPG (EGF PEPTIDE)
- SEQ ID NO:19 ANMLCAGLESGGKDSCRGDSSGALV (serine protease
binding core)

PEPTIDE INHIBITORS:

- 20 SEQ ID NO:20 MBL full length cDNA
- SEQ ID NO:21 MBL full length protein
- SEQ ID NO:22 OGK-X-GP (consensus binding)
- SEQ ID NO:23 OGKLG
- SEQ ID NO:24 GLR GLQ GPO GKL GPO G
- 25 SEQ ID NO:25 GPO GPO GLR GLQ GPO GKL GPO GPO GPO
- SEQ ID NO:26 GKDGRDGTKGEKGEPGQGLRGLQGPOGKLGPOG
- SEQ ID NO:27 GAOGSOGEEKGAOGPQGPOGPOGKMGPKGEOGDO
- (human h-ficolin)
- SEQ ID NO:28
- 30 GCOGLOGAOGDKGEAGTNGKRGERGPOGPOGKAGPOGPN
- GAOGEO (human ficolin p35)
- SEQ ID NO:29 LQRALEILPNRVTIKANRPFLVFI (C4 cleavage site)

EXPRESSION INHIBITORS:

SEQ ID NO:30 cDNA of CUBI-EGF domain (nucleotides 22-680 of SEQ ID NO:4)

SEQ ID NO:31

5 5' CGGGCACACCATGAGGCTGCTGACCCTCCTGGGC 3'
Nucleotides 12-45 of SEQ ID NO:4 including the MASP-2 translation start site (sense)

SEQ ID NO:32

5'GACATTACCTTCCGCTCCGACTCCAACGAGAAG3'
10 Nucleotides 361-396 of SEQ ID NO:4 encoding a region comprising the MASP-2 MBL binding site (sense)

SEQ ID NO:33

5'AGCAGCCCTGAATACCCACGGCCGTATCCCAA3'
15 Nucleotides 610-642 of SEQ ID NO:4 encoding a region comprising the CUBII domain

CLONING PRIMERS:

SEQ ID NO:34 CGGGATCCATGAGGCTGCTGACCCTC (5' PCR for CUB)

SEQ ID NO:35 GGAATTCCTAGGCTGCATA (3' PCR FOR CUB)

20 SEQ ID NO:36 GGAATTCCTACAGGGCGCT (3' PCR FOR CUBIEGF)

SEQ ID NO:37 GGAATTCCTAGTAGTGGAT (3' PCR FOR CUBIEGFCUBII)

SEQ ID NOS:38-47 are cloning primers for humanized antibody

SEQ ID NO:48 is 9 aa peptide bond

25 EXPRESSION VECTOR:

SEQ ID NO:49 is the MASP-2 minigene insert

SEQ ID NO: 50 is the murine MASP-2 cDNA

SEQ ID NO: 51 is the murine MASP-2 protein (w/leader)

SEQ ID NO: 52 is the mature murine MASP-2 protein

30 SEQ ID NO: 53 the rat MASP-2 cDNA

SEQ ID NO: 54 is the rat MASP-2 protein (w/ leader)

SEQ ID NO: 55 is the mature rat MASP-2 protein

SEQ ID NO: 56-59 are the oligonucleotides for site-directed mutagenesis of human MASP-2 used to generate human MASP-2A

SEQ ID NO: 60-63 are the oligonucleotides for site-directed mutagenesis of murine MASP-2 used to generate murine MASP-2A

5 SEQ ID NO: 64-65 are the oligonucleotides for site-directed mutagenesis of rat MASP-2 used to generate rat MASP-2A

SEQ ID NO: 66 DNA encoding 17D20_dc35VH21N11VL (OMS646) heavy chain variable region (VH) (without signal peptide)

10 SEQ ID NO: 67 17D20_dc35VH21N11VL (OMS646) heavy chain variable region (VH) polypeptide

SEQ ID NO: 68 17N16mc heavy chain variable region (VH) polypeptide

SEQ ID NO: 69: DNA encoding 17D20_dc35VH21N11VL (OMS646) light chain variable region (VL)

15 SEQ ID NO: 70: 17D20_dc35VH21N11VL (OMS646) light chain variable region (VL) polypeptide

SEQ ID NO: 71: 17N16_dc17N9 light chain variable region (VL) polypeptide

20 DETAILED DESCRIPTION

The present invention is based upon the surprising discovery by the present inventors that it is possible to inhibit the lectin mediated MASP-2 pathway while leaving the classical pathway intact. The present invention also describes the use of MASP-2 as a therapeutic target for inhibiting cellular injury associated with lectin-mediated complement pathway activation while leaving the classical (C1q-dependent) pathway component of the immune system intact.

I. DEFINITIONS

25 Unless specifically defined herein, all terms used herein have the same meaning as would be understood by those of ordinary skill in the art of the present invention. The following definitions are provided in order to provide clarity with respect to the terms as they are used in the specification and claims to describe the present invention.

As used herein, the term “MASP-2-dependent complement activation” comprises MASP-2- dependent activation of the lectin pathway, which occurs under physiological

conditions (*i.e.*, in the presence of Ca^{++}) leading to the formation of the lectin pathway C3 convertase C4b2a and upon accumulation of the C3 cleavage product C3b subsequently to the C5 convertase C4b2a(C3b)_n , which has been determined to primarily cause opsonization.

5 As used herein, the term "alternative pathway" refers to complement activation that is triggered, for example, by zymosan from fungal and yeast cell walls, lipopolysaccharide (LPS) from Gram negative outer membranes, and rabbit erythrocytes, as well as from many pure polysaccharides, rabbit erythrocytes, viruses, bacteria, animal tumor cells, parasites and damaged cells, and which has traditionally been thought to
10 arise from spontaneous proteolytic generation of C3b from complement factor C3.

 As used herein, the term "lectin pathway" refers to complement activation that occurs via the specific binding of serum and non-serum carbohydrate-binding proteins including mannan-binding lectin (MBL), CL-11 and the ficolins (H-ficolin, M-ficolin, or L-ficolin).

15 As used herein, the term "classical pathway" refers to complement activation that is triggered by antibody bound to a foreign particle and requires binding of the recognition molecule C1q.

 As used herein, the term "MASP-2 inhibitory agent" refers to any agent that binds to or directly interacts with MASP-2 and effectively inhibits MASP-2-dependent
20 complement activation, including anti-MASP-2 antibodies and MASP-2 binding fragments thereof, natural and synthetic peptides, small molecules, soluble MASP-2 receptors, expression inhibitors and isolated natural inhibitors, and also encompasses peptides that compete with MASP-2 for binding to another recognition molecule (e.g., MBL, H-ficolin, M-ficolin, or L-ficolin) in the lectin pathway, but does not encompass
25 antibodies that bind to such other recognition molecules. MASP-2 inhibitory agents useful in the method of the invention may reduce MASP-2-dependent complement activation by greater than 20%, such as greater than 50%, such as greater than 90%. In one embodiment, the MASP-2 inhibitory agent reduces MASP-2-dependent complement activation by greater than 90% (*i.e.*, resulting in MASP-2 complement activation of only
30 10% or less).

 As used herein, the term "antibody" encompasses antibodies and antibody fragments thereof, derived from any antibody-producing mammal (e.g., mouse, rat, rabbit, and primate including human), or from a hybridoma, phage selection, recombinant

expression or transgenic animals (or other methods of producing antibodies or antibody fragments”), that specifically bind to a target polypeptide, such as, for example, MASP-2, polypeptides or portions thereof. It is not intended that the term “antibody” limited as regards to the source of the antibody or the manner in which it is made (e.g., by

5 hybridoma, phage selection, recombinant expression, transgenic animal, peptide synthesis, etc). Exemplary antibodies include polyclonal, monoclonal and recombinant antibodies; pan-specific, multispecific antibodies (e.g., bispecific antibodies, trispecific antibodies); humanized antibodies; murine antibodies; chimeric, mouse-human, mouse-primate, primate-human monoclonal antibodies; and anti-idiotypic antibodies, and

10 may be any intact antibody or fragment thereof. As used herein, the term “antibody” encompasses not only intact polyclonal or monoclonal antibodies, but also fragments thereof (such as dAb, Fab, Fab', F(ab')₂, Fv), single chain (ScFv), synthetic variants thereof, naturally occurring variants, fusion proteins comprising an antibody portion with an antigen-binding fragment of the required specificity, humanized antibodies, chimeric

15 antibodies, and any other modified configuration of the immunoglobulin molecule that comprises an antigen-binding site or fragment (epitope recognition site) of the required specificity.

A “monoclonal antibody” refers to a homogeneous antibody population wherein the monoclonal antibody is comprised of amino acids (naturally occurring and non-

20 naturally occurring) that are involved in the selective binding of an epitope. Monoclonal antibodies are highly specific for the target antigen. The term “monoclonal antibody” encompasses not only intact monoclonal antibodies and full-length monoclonal antibodies, but also fragments thereof (such as Fab, Fab', F(ab')₂, Fv), single chain (ScFv), variants thereof, fusion proteins comprising an antigen-binding portion,

25 humanized monoclonal antibodies, chimeric monoclonal antibodies, and any other modified configuration of the immunoglobulin molecule that comprises an antigen-binding fragment (epitope recognition site) of the required specificity and the ability to bind to an epitope. It is not intended to be limited as regards the source of the antibody or the manner in which it is made (e.g., by hybridoma, phage selection, recombinant

30 expression, transgenic animals, etc.). The term includes whole immunoglobulins as well as the fragments etc. described above under the definition of “antibody”.

As used herein, the term “antibody fragment” refers to a portion derived from or related to a full-length antibody, such as, for example, an anti-MASP-2 antibody,

generally including the antigen binding or variable region thereof. Illustrative examples of antibody fragments include Fab, Fab', F(ab)₂, F(ab')₂ and Fv fragments, scFv fragments, diabodies, linear antibodies, single-chain antibody molecules and multispecific antibodies formed from antibody fragments.

5 As used herein, a "single-chain Fv" or "scFv" antibody fragment comprises the V_H and V_L domains of an antibody, wherein these domains are present in a single polypeptide chain. Generally, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains, which enables the scFv to form the desired structure for antigen binding.

10 As used herein, a "chimeric antibody" is a recombinant protein that contains the variable domains and complementarity-determining regions derived from a non-human species (e.g., rodent) antibody, while the remainder of the antibody molecule is derived from a human antibody.

As used herein, a "humanized antibody" is a chimeric antibody that comprises a
15 minimal sequence that conforms to specific complementarity-determining regions derived from non-human immunoglobulin that is transplanted into a human antibody framework. Humanized antibodies are typically recombinant proteins in which only the antibody complementarity-determining regions are of non-human origin.

As used herein, the term "mannan-binding lectin" ("MBL") is equivalent to
20 mannan-binding protein ("MBP").

As used herein, the "membrane attack complex" ("MAC") refers to a complex of the terminal five complement components (C5b combined with C6, C7, C8 and C-9) that inserts into and disrupts membranes (also referred to as C5b-9).

As used herein, "a subject" includes all mammals, including without limitation
25 humans, non-human primates, dogs, cats, horses, sheep, goats, cows, rabbits, pigs and rodents.

As used herein, the amino acid residues are abbreviated as follows: alanine (Ala;A), asparagine (Asn;N), aspartic acid (Asp;D), arginine (Arg;R), cysteine (Cys;C), glutamic acid (Glu;E), glutamine (Gln;Q), glycine (Gly;G), histidine (His;H), isoleucine (Ile;I), leucine (Leu;L), lysine (Lys;K), methionine (Met;M), phenylalanine (Phe;F),
30 proline (Pro;P), serine (Ser;S), threonine (Thr;T), tryptophan (Trp;W), tyrosine (Tyr;Y), and valine (Val;V).

In the broadest sense, the naturally occurring amino acids can be divided into groups based upon the chemical characteristic of the side chain of the respective amino acids. By "hydrophobic" amino acid is meant either Ile, Leu, Met, Phe, Trp, Tyr, Val, Ala, Cys or Pro. By "hydrophilic" amino acid is meant either Gly, Asn, Gln, Ser, Thr, 5 Asp, Glu, Lys, Arg or His. This grouping of amino acids can be further subclassed as follows. By "uncharged hydrophilic" amino acid is meant either Ser, Thr, Asn or Gln. By "acidic" amino acid is meant either Glu or Asp. By "basic" amino acid is meant either Lys, Arg or His.

As used herein the term "conservative amino acid substitution" is illustrated by a 10 substitution among amino acids within each of the following groups: (1) glycine, alanine, valine, leucine, and isoleucine, (2) phenylalanine, tyrosine, and tryptophan, (3) serine and threonine, (4) aspartate and glutamate, (5) glutamine and asparagine, and (6) lysine, arginine and histidine.

The term "oligonucleotide" as used herein refers to an oligomer or polymer of 15 ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term also covers those oligonucleobases composed of naturally-occurring nucleotides, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring modifications.

As used herein, an "epitope" refers to the site on a protein (e.g., a human MASP-2 20 protein) that is bound by an antibody. "Overlapping epitopes" include at least one (e.g., two, three, four, five, or six) common amino acid residue(s), including linear and non-linear epitopes.

As used herein, the terms "polypeptide," "peptide," and "protein" are used interchangeably and mean any peptide-linked chain of amino acids, regardless of length 25 or post-translational modification. The MASP-2 protein described herein can contain or be wild-type proteins or can be variants that have not more than 50 (e.g., not more than one, two, three, four, five, six, seven, eight, nine, ten, 12, 15, 20, 25, 30, 35, 40, or 50) conservative amino acid substitutions. Conservative substitutions typically include substitutions within the following groups: glycine and alanine; valine, isoleucine, and 30 leucine; aspartic acid and glutamic acid; asparagine, glutamine, serine and threonine; lysine, histidine and arginine; and phenylalanine and tyrosine.

In some embodiments, the human MASP-2 protein can have an amino acid sequence that is, or is greater than, 70 (e.g., 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82,

83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100) % identical to the human MASP-2 protein having the amino acid sequence set forth in SEQ ID NO: 5.

In some embodiments, peptide fragments can be at least 6 (e.g., at least 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 350, 400, 450, 500, or 600 or more) amino acid residues in length (e.g., at least 6 contiguous amino acid residues of SEQ ID NO: 5). In some embodiments, an antigenic peptide fragment of a human MASP-2 protein is fewer than 500 (e.g., fewer than 450, 400, 350, 325, 300, 275, 250, 225, 200, 190, 180, 170, 160, 150, 140, 130, 120, 110, 100, 95, 90, 85, 80, 75, 70, 65, 60, 55, 50, 49, 48, 47, 46, 45, 44, 43, 42, 41, 40, 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, or 6) amino acid residues in length (e.g., fewer than 500 contiguous amino acid residues in any one of SEQ ID NOS: 5).

Percent (%) amino acid sequence identity is defined as the percentage of amino acids in a candidate sequence that are identical to the amino acids in a reference sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared can be determined by known methods.

II. Overview of the Invention

Lectins (MBL, M-ficolin, H-ficolin, L-ficolin and CL-11) are the specific recognition molecules that trigger the innate complement system and the system includes the lectin initiation pathway and the associated terminal pathway amplification loop that amplifies lectin-initiated activation of terminal complement effector molecules. C1q is the specific recognition molecule that triggers the acquired complement system and the system includes the classical initiation pathway and associated terminal pathway amplification loop that amplifies C1q-initiated activation of terminal complement effector molecules. We refer to these two major complement activation systems as the

lectin-dependent complement system and the C1q-dependent complement system, respectively.

In addition to its essential role in immune defense, the complement system contributes to tissue damage in many clinical conditions. Thus, there is a pressing need to develop therapeutically effective complement inhibitors to prevent these adverse effects. With the recognition that it is possible to inhibit the lectin mediated MASP-2 pathway while leaving the classical pathway intact comes the realization that it would be highly desirable to specifically inhibit only the complement activation system causing a particular pathology without completely shutting down the immune defense capabilities of complement. For example, in disease states in which complement activation is mediated predominantly by the lectin-dependent complement system, it would be advantageous to specifically inhibit only this system. This would leave the C1q-dependent complement activation system intact to handle immune complex processing and to aid in host defense against infection.

The preferred protein component to target in the development of therapeutic agents to specifically inhibit the lectin-dependent complement system is MASP-2. Of all the known protein components of the lectin-dependent complement system (MBL, H-ficolin, M-ficolin, L-ficolin, MASP-2, C2-C9, Factor B, Factor D, and properdin), only MASP-2 is both unique to the lectin-dependent complement system and required for the system to function. The lectins (MBL, H-ficolin, M-ficolin, L-ficolin and CL-11) are also unique components in the lectin-dependent complement system. However, loss of any one of the lectin components would not necessarily inhibit activation of the system due to lectin redundancy. It would be necessary to inhibit all five lectins in order to guarantee inhibition of the lectin-dependent complement activation system. Furthermore, since MBL and the ficolins are also known to have opsonic activity independent of complement, inhibition of lectin function would result in the loss of this beneficial host defense mechanism against infection. In contrast, this complement-independent lectin opsonic activity would remain intact if MASP-2 was the inhibitory target. An added benefit of MASP-2 as the therapeutic target to inhibit the lectin-dependent complement activation system is that the plasma concentration of MASP-2 is among the lowest of any complement protein (≈ 500 ng/ml); therefore, correspondingly low concentrations of high-affinity inhibitors of MASP-2 may be sufficient to obtain full inhibition (Moller-Kristensen, M., et al., *J. Immunol Methods* 282:159-167, 2003).

III. THE ROLE OF MASP-2 IN THROMBOTIC MICROANGIOPATHIES AND THERAPEUTIC METHODS USING MASP-2 INHIBITORY AGENTS

5 Overview

Thrombotic microangiopathy (TMA) is a pathology characterized by blood clots in small blood vessels (Benz, K.; et al., *Curr Opin Nephrol Hypertens* 19(3):242–7 (2010)). Stress or injury to the underlying vascular endothelium is believed to be a primary driver. Clinical and laboratory findings of TMA include thrombocytopenia, anemia, purpura, and renal failure. The classic TMAs are hemolytic uremic syndrome (HUS) and thrombotic thrombocytopenic purpura (TTP). The characteristic underlying pathological feature of TMAs are platelet activation and the formation of microthrombi in the small arterioles and venules. Complement activation initiated, at least in part, by an injury or stress to microvascular endothelium, is also implicated in other TMAs including catastrophic antiphospholipid syndrome (CAPS), systemic Degos disease, and TMAs secondary to cancer, cancer chemotherapy and transplantation.

Direct evidence for a pathological role of complement in a host of nephritides is provided by studies of patients with genetic deficiencies in specific complement components. A number of reports have documented an association of renal injury with deficiencies of complement regulatory factor H (Ault, B.H., *Nephrol.* 14:1045-1053, 2000; Levy, M., et al., *Kidney Int.* 30:949-56, 1986; Pickering, M.C., et al., *Nat. Genet.* 31:424-8, 2002). Factor H deficiency results in low plasma levels of factor B and C3 due to activation-related consumption of these components. Circulating levels of C5b-9 are also elevated in the serum of these patients, implying complement activation. Membranoproliferative glomerulonephritis (MPGN) and idiopathic hemolytic uremic syndrome (HUS) are associated with factor H deficiency or mutations of factor H. Factor H-deficient pigs (Jansen, J.H., et al., *Kidney Int.* 53:331-49, 1998) and factor-H knockout mice (Pickering, M.C., 2002) display MPGN-like symptoms, confirming the importance of factor H in complement regulation. Deficiencies of other complement components are associated with renal disease, secondary to the development of systemic lupus erythematosus (SLE) (Walport, M.J., Davies, et al., *Ann. N.Y. Acad. Sci.* 815:267-81, 1997). Deficiency for C1q, C4 and C2 predispose strongly to the development of SLE via mechanisms relating to defective clearance of immune complexes and apoptotic

material. In many of these SLE patients lupus nephritis occurs, characterized by the deposition of immune complexes throughout the glomerulus.

aHUS

Atypical hemolytic uremic syndrome (aHUS) is part of a group of conditions
5 termed "Thrombotic microangiopathies." In the atypical form of HUS (aHUS), the
disease is associated with defective complement regulation and can be either sporadic or
familial. Familial cases of aHUS are associated with mutations in genes coding for
complement activation or complement regulatory proteins, including complement factor
H, factor I, factor B, membrane cofactor CD46 as well as complement factor H-related
10 protein 1 (CFHR1) and complement factor H-related protein 3 (CFHR3). (Zipfel, P.F.,
et al., *PloS Genetics* 3(3):e41 (2007)). The unifying feature of this diverse array of
genetic mutations associated with aHUS is a predisposition to enhanced complement
activation on cellular or tissue surfaces. Therefore, one aspect of the present invention
comprises treating a patient suffering with aHUS that is associated with a factor H
15 deficiency by administering an effective amount of a MASP-2 inhibitory agent. Another
aspect of the present invention comprises treating a patient suffering with HUS that is
associated with a factor I, factor B, membrane cofactor CD46, CFHR1 or CFHR3
deficiency by administering an effective amount of a MASP-2 inhibitory agent.

Significant progress has been made recently toward the understanding of the
20 molecular pathophysiology underlying enhanced complement activation in aHUS caused
by the diverse set of mutant complement factors. This mechanism is best understood for
factor H mutations. Factor H is an abundant serum protein comprising 20 short
consensus repeat (SCR) domains that acts as a negative regulator of complement
activation both in solution as well as on host cell surfaces. It targets the activated form of
25 C3 and, together with factor I and other cofactors, promotes its inactivation, forestalling
further complement activation. To effectively control complement activation on host cell
surfaces, factor H needs to interact with host cells, which is mediated by SCR domains
16-20. All factor H mutations associated with aHUS described to date are clustered in the
C-terminal region encompassing (SCR) domains 16-20. These mutant factor H proteins
30 are fully functional in controlling C3 activation in solution, but are unable to interact with
host cell surfaces and consequently cannot control C3 activation on cellular surfaces (Exp
Med 204(6):1249-56 (2007)). Thus, certain mutations of factor H are associated with
aHUS because the mutant factor H protein fails to interact with host cell surfaces and thus

cannot effectively down modulate complement activation on host cell surfaces, including the microvascular endothelium. As a result, once initial C3 activation has occurred, subsequent complement activation on microvascular endothelial surfaces proceeds unabated in patients with factor H mutations. This uncontrolled activation of complement ultimately leads to progressive injury to the vascular endothelium, subsequent platelet aggregation and microvascular coagulation, and hemolysis caused by sheer stress of RBC passage through partially occluded microvessels. Thus, aHUS disease manifestations and clinical and laboratory findings are directly linked to a defect in the negative regulation of complement on the surface of the microvascular endothelium.

10 Analogous to factor H mutation, loss-of-function mutations in the negative complement modulators factor I and membrane cofactor protein (CD46) are also linked to aHUS. The opposite has been observed for factor B the C3 protein in that aHUS was found to be associated with gain-of-function mutations in these proteins (Pediatr Nephrol 25(12):2431-42 (2010)). Thus, a host of converging data implicates complement activation in aHUS pathogenesis. This notion is most convincingly supported by the therapeutic efficacy of eculizumab, a monoclonal antibody that blocks the terminal complement protein C5 in the treatment of aHUS.

While the central role of complement as an effector mechanism in aHUS is widely accepted, the triggers initiating complement activation and the molecular pathways involved are unresolved. Not all individuals carrying the above described mutations develop aHUS. In fact, familial studies have suggested that the penetrance of aHUS is only ~50% (Ann Hum Genet 74(1):17-26 (2010)). The natural history of the disease suggests that aHUS most often develops after an initiating event such as an infectious episode or an injury. Infectious agents are well known to activate the complement system. In the absence of pre-existing adaptive immunity, complement activation by infectious agents may be primarily initiated via the lectin pathway. Thus, lectin pathway activation triggered by an infection may represent the initiating trigger for subsequent pathological amplification of complement activation in aHUS-predisposed individuals, which may ultimately lead to disease progression. Accordingly, another aspect of the present invention comprises treating a patient suffering with aHUS secondary to an infection by administering an effective amount of a MASP-2 inhibitory agent.

Other forms of injury to host tissue will activate complement via the lectin pathway, in particular injury to the vascular endothelium. Human vascular endothelial

cells subject to oxidative stress for example respond by expressing surface moieties that bind lectins and activate the lectin pathway of complement (Am J. Pathol 156(6):1549-56 (2000)). Vascular injury following ischemia/reperfusion also activates complement via the lectin pathway *in vivo* (Scand J Immunol 61(5):426-34 (2005)). Lectin pathway activation in this setting has pathological consequences for the host, and inhibition of the lectin pathway by blocking MASP-2 prevents further host tissue injury and adverse outcomes (Schwaeble PNAS 2011).

Thus, other processes that precipitate aHUS are also known to activate the lectin pathway of complement. It is therefore likely that the lectin pathway may represent the initial complement activating mechanism that is inappropriately amplified in a deregulated fashion in individuals genetically predisposed to aHUS, thus initiating aHUS pathogenesis. By inference, agents that block activation of complement via the lectin pathway, including anti-MASP-2 antibodies, are expected to prevent disease progression or reduce exacerbations in aHUS susceptible individuals.

In further support of this concept, recent studies have identified *S. pneumonia* as an important etiological agent in pediatric cases of aHUS. (Nephrology (Carlton), 17:48-52 (2012); Pediatr Infect Dis J. 30(9):736-9 (2011)). This particular etiology appears to have an unfavorable prognosis, with significant mortality and long-term morbidity. Notably, these cases involved non-enteric infections leading to manifestations of microangiopathy, uremia and hemolysis without evidence of concurrent mutations in complement genes known to predispose to aHUS. It is important to note that *S. pneumonia* is particularly effective at activating complement, and does so predominantly through the lectin pathway. Thus, in cases of non-enteric HUS associated with pneumococcal infection, manifestations of microangiopathy, uremia and hemolysis are expected to be driven predominantly by activation of the lectin pathway, and agents that block the lectin pathway, including anti-MASP-2 antibodies, are expected to prevent progression of aHUS or reduce disease severity in these patients. Accordingly, another aspect of the present invention comprises treating a patient suffering with non-enteric aHUS that is associated with *S. pneumonia* infection by administering an effective amount of a MASP-2 inhibitory agent.

In accordance with the foregoing, in some embodiments, in the setting of a subject at risk for developing renal failure associated with aHUS, a method is provided for decreasing the likelihood of developing aHUS, or of developing renal failure associated

- with aHUS, comprising administering an amount of an MASP-2 inhibitory agent for a time period effective to ameliorate or prevent renal failure in the subject. In some embodiments, the method further comprises the step of determining whether a subject is at risk for developing aHUS prior to the onset of any symptoms associated with aHUS.
- 5 In other embodiments, the method comprises determining whether a subject is at risk for developing aHUS upon the onset of at least one or more symptoms indicative of aHUS (e.g., the presence of anemia, thrombocytopenia and/or renal insufficiency) and/or the presence of thrombotic microangiopathy in a biopsy obtained from the subject. The determination of whether a subject is at risk for developing aHUS comprises determining
- 10 whether the subject has a genetic predisposition to developing aHUS, which may be carried out by assessing genetic information (e.g. from a database containing the genotype of the subject), or performing at least one genetic screening test on the subject to determine the presence or absence of a genetic marker associated with aHUS (i.e., determining the presence or absence of a genetic mutation associated with aHUS in the
- 15 genes encoding complement factor H (CFH), factor I (CFI), factor B (CFB), membrane cofactor CD46, C3, complement factor H-related protein 1 (CFHR1), or THBD (encoding the anticoagulant protein thrombomodulin) or complement factor H-related protein 3 (CFHR3), or complement factor H-related protein 4 (CFHR4)) either via genome sequencing or gene-specific analysis (e.g., PCR analysis), and/or determining whether the
- 20 subject has a family history of aHUS. Methods of genetic screening for the presence or absence of a genetic mutation associated with aHUS are well established, for example, see Noris M et al. "Atypical Hemolytic-Uremic Syndrome," 2007 Nov 16 [Updated 2011 Mar 10]. In: Pagon RA, Bird TD, Dolan CR, et al., editors. GeneReviews™, Seattle (WA): University of Washington, Seattle.
- 25 For example, overall the penetrance of the disease in those with mutations of complement factor H (CFH) is 48%, and the penetrance for mutations in CD46 is 53%, for mutations in CFI is 50%, for mutations in C3 is 56% and for mutations in THBD is 64% (Caprioli J. et al., Blood, 108:1267-79 (2006); Noris et al., Clin J Am Soc Nephrol 5:1844-59 (2010)). As described in Caprioli et al., (2006), supra, a substantial number of
- 30 individuals with a mutation in complement Factor H (CFH) never develop aHUS, and it is postulated that suboptimal CFH activity in these individuals is sufficient to protect the host from the effects of complement activation in physiological conditions, however, suboptimal CFH activity is not sufficient to prevent C3b from being deposited on

vascular endothelial cells when exposure to an agent that activates complement produces higher than normal amounts of C3b.

Accordingly, in one embodiment, a method is provided for inhibiting MASP-2-dependent complement activation in a subject suffering from, or at risk for developing non-Factor H-dependent atypical hemolytic uremic syndrome, comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory agent effective to inhibit MASP-2-dependent complement activation. In another embodiment, a method is provided for inhibiting MASP-2-dependent complement activation in a subject at risk for developing Factor H-dependent atypical hemolytic uremic syndrome, comprising periodically monitoring the subject to determine the presence or absence of anemia, thrombocytopenia or rising creatinine, and treating with a MASP-2 inhibitory agent upon the determination of the presence of anemia thrombocytopenia, or rising creatinine. In another embodiment, a method is provided for reducing the likelihood that a subject at risk for developing Factor H-dependent aHUS will suffer clinical symptoms associated with aHUS, comprising administering a MASP-2 inhibitory agent prior to, or during, or after an event known to be associated with triggering aHUS clinical symptoms, for example, drug exposure (e.g., chemotherapy), infection (e.g., bacterial infection), malignancy, an injury, organ or tissue transplant, or pregnancy.

In one embodiment, a method is provided for reducing the likelihood that a subject at risk for developing aHUS will suffer clinical symptoms associated with aHUS, comprising periodically monitoring the subject to determine the presence or absence of anemia, thrombocytopenia or rising creatinine, and treating with a MASP-2 inhibitory agent upon the determination of the presence of anemia, thrombocytopenia, or rising creatinine.

In another embodiment, a method is provided for reducing the likelihood that a subject at risk for developing aHUS will suffer clinical symptoms associated with aHUS comprising administering a MASP-2 inhibitory agent prior to, or during, or after an event known to be associated with triggering aHUS clinical symptoms, for example, drug exposure (e.g., chemotherapy), infection (e.g., bacterial infection), malignancy, an injury, organ or tissue transplant, or pregnancy.

In some embodiments, the MASP-2 inhibitory agent is administered for a time period of at least one, two, three, four days, or longer, prior to, during, or after the event associated with triggering aHUS clinical symptoms and may be repeated as determined

by a physician until the condition has been resolved or is controlled. In a pre-aHUS setting, the MASP-2 inhibitory agent may be administered to the subject systemically, such as by intra-arterial, intravenous, intramuscular, inhalational, nasal, subcutaneous or other parenteral administration.

5 In some embodiments, in the setting of initial diagnosis of aHUS, or in a subject exhibiting one or more symptoms consistent with a diagnosis of aHUS (e.g., the presence of anemia, thrombocytopenia and/or renal insufficiency), the subject is treated with an effective amount of a MASP-2 inhibitory agent (e.g., an anti-MASP-2 antibody) as a first line therapy in the absence of plasmapheresis, or in combination with plasmapheresis. As
10 a first line therapy, the MASP-2 inhibitory agent may be administered to the subject systemically, such as by intra-arterial, intravenous, intramuscular, inhalational, nasal, subcutaneous or other parenteral administration. In some embodiments, the MASP-2 inhibitory agent is administered to a subject as a first line therapy in the absence of plasmapheresis to avoid the potential complications of plasmapheresis including
15 hemorrhage, infection, and exposure to disorders and/or allergies inherent in the plasma donor, or in a subject otherwise averse to plasmapheresis, or in a setting where plasmapheresis is unavailable.

In some embodiments, the method comprises administering a MASP-2 inhibitory agent to a subject suffering from aHUS via a catheter (e.g., intravenously) for a first time
20 period (e.g., at least one day to a week or two weeks) followed by administering a MASP-2 inhibitory agent to the subject subcutaneously for a second time period (e.g., a chronic phase of at least two weeks or longer). In some embodiments, the administration in the first and/or second time period occurs in the absence of plasmapheresis. In some embodiments, the method further comprises determining the level of at least one
25 complement factor (e.g., C3, C5) in the subject prior to treatment, and optionally during treatment, wherein the determination of a reduced level of at least one complement factor in comparison to a standard value or healthy control subject is indicative of the need for continued treatment with the MASP-2 inhibitory agent.

In some embodiments, the method comprises administering a MASP-2 inhibitory
30 agent, such as an anti-MASP-2 antibody, to a subject suffering from, or at risk for developing, aHUS either intravenously, intramuscularly, or preferably, subcutaneously. Treatment may be chronic and administered daily to monthly, but preferably every two

weeks. The anti-MASP-2 antibody may be administered alone, or in combination with a C5 inhibitor, such as eculizumab.

HUS

5 Like atypical HUS, the typical form of HUS displays all the clinical and laboratory findings of a TMA. Typical HUS, however, is often a pediatric disease and usually has no familial component or direct association with mutations in complement genes. The etiology of typical HUS is tightly linked to infection with certain intestinal pathogens. The patients typically present with acute renal failure, hemoglobinuria, and
10 thrombocytopenia, which typically follows an episode of bloody diarrhea. The condition is caused by an enteric infection with *Shigella dysenteriae*, *Salmonella* or shiga toxin-like producing enterohemorrhagic strains of *E. Coli*, such as *E. Coli* O157:H7. The pathogens are acquired from contaminated food or water supply. HUS is a medical emergency and carries a 5-10% mortality. A significant portion of survivors develop chronic kidney
15 disease (Corrigan and Boineau, *Pediatr Rev* 22 (11): 365–9 (2011)) and may require kidney transplantation.

 The microvascular coagulation in typical HUS occurs predominantly, though not exclusively, in the renal microvasculature. The underlying pathophysiology is mediated by Shiga toxin (STX). Excreted by enteropathic microbes into the intestinal lumen, STX
20 crosses the intestinal barrier, enters the bloodstream and binds to vascular endothelial cells via the globotriaosyl ceramide receptor CD77 (Boyd and Lingwood *Nephron* 51:207 (1989)), which is preferentially expressed on glomerular endothelium and mediates the toxic effect of STX. Once bound to the endothelium, STX induces a series of events that damage vascular endothelium, activate leukocytes and cause vWF-dependent thrombus
25 formation (Forsyth et al., *Lancet* 2: 411–414 (1989); Zoja et al., *Kidney Int.* 62: 846–856 (2002); Zanchi et al., *J. Immunol.* 181:1460–1469 (2008); Morigi et al., *Blood* 98: 1828–1835 (2001); Guessou et al., *Infect. Immun.*, 73: 8306–8316 (2005)). These microthrombi obstruct or occlude the arterioles and capillaries of the kidney and other organs. The obstruction of blood flow in arterioles and capillaries by microthrombi increases the shear
30 force applied to RBCs as they squeeze through the narrowed blood vessels. This can result in destruction of RBC by shear force and the formation of RBC fragments called schistocytes. The presence of schistocytes is a characteristic finding in HUS. This mechanism is known as microangiopathic hemolysis. In addition, obstruction of blood

flow results in ischemia, initiating a complement-mediated inflammatory response that causes additional damage to the affected organ.

The lectin pathway of complement contributes to the pathogenesis of HUS by two principle mechanisms: 1) MASP-2-mediated direct activation of the coagulation cascade caused by endothelial injury, and 2) lectin-mediated subsequent complement activation induced by the ischemia resulting from the initial occlusion of microvascular blood flow.

STX injures microvascular endothelial cells, and injured endothelial cells are known to activate the complement system. As detailed above, complement activation following endothelial cell injury is driven predominantly by the lectin pathway. Human vascular endothelial cells subject to oxidative stress respond by expressing surface moieties that bind lectins and activate the lectin pathway of complement (Collard et al., *Am J Pathol.* 156(5):1549-56 (2000)). Vascular injury following ischemia reperfusion also activates complement via the lectin pathway *in vivo* (Scand J Immunol 61(5):426-34 (2005)). Lectin pathway activation in this setting has pathological consequences for the host, and inhibition of the lectin pathway by blockade of MASP-2 prevents further host tissue injury and adverse outcomes (Schwaebler et al., *PNAS* (2011)). In addition to complement activation, lectin-dependent activation of MASP-2 has been shown to result in cleavage of prothrombin to form thrombin and to promote coagulation. Thus, activation of the lectin pathway of complement by injured endothelial cells can directly activate the coagulation system. The lectin pathway of complement, by virtue of MASP-2-mediated prothrombin activation, therefore is likely the dominant molecular pathway linking the initial endothelial injury by STX to the coagulation and microvascular thrombosis that occurs in HUS. It is therefore expected that lectin pathway inhibitors, including, but not limited to, antibodies that block MASP-2 function, will prevent or mitigate microvascular coagulation, thrombosis and hemolysis in patients suffering from HUS. Indeed, administration of anti-MASP-2 antibody profoundly protects mice in a model of typical HUS. As described in Example 36 and shown in FIGURE 45, all control mice exposed to STX and LPS developed severe HUS and became moribund or died within 48 hours. On the other hand, as further shown in FIGURE 45, all mice treated with an anti-MASP-2 antibody and then exposed to STX and LPS survived (Fisher's exact $p < 0.01$; $N=5$). Thus, anti-MASP-2 therapy profoundly protects mice in this model of HUS. It is expected that administration of a MASP-2 inhibitory agent, such as a MASP-2 antibody, will be effective in the treatment of HUS patients and provide

protection from microvascular coagulation, thrombosis, and hemolysis caused by infection with enteropathic *E. coli* or other STX-producing pathogens.

While shown here for HUS caused by STX, it is expected that anti-MASP-2 therapy will also be beneficial for HUS-like syndromes due to endothelial injury caused by other toxic agents. This includes agents such as mitomycin, ticlopidine, cycloplatin, quinine, cyclosporine, bleomycin as well as other chemotherapy drugs and immunosuppressive drugs. Thus, it is expected that anti-MASP-2 antibody therapy, or other modalities that inhibit MASP-2 activity, will effectively prevent or limit coagulation, thrombus formation, and RBC destruction and prevent renal failure in HUS and other TMA related diseases (*i.e.*, aHUS and TTP).

Patients suffering from HUS often present with diarrhea and vomiting, their platelet counts are usually reduced (thrombocytopenia), and RBCs are reduced (anemia). A pre-HUS diarrhea phase typically lasts for about four days, during which subjects at risk for developing HUS typically exhibit one or more of the following symptoms in addition to severe diarrhea: a hematocrit level below 30% with smear evidence of intravascular erythrocyte destruction, thrombocytopenia (platelet count $<150 \times 10^3/\text{mm}^3$), and/or the presence of impaired renal function (serum creatinine concentration greater than the upper limit of reference range for age). The presence of oligoanuria (urine output $\leq 0.5 \text{ mL/kg/h}$ for >1 day) can be used as a measure for progression towards developing HUS (see C. Hickey et al., *Arch Pediatr Adolesc Med* 165(10):884-889 (2011)). Testing is typically carried out for the presence of infection with *E. coli* bacteria (*E. coli* O157:H7), or *Shigella* or *Salmonella* species. In a subject testing positive for infection with enterogenic *E. coli* (e.g., *E. coli* O157:H7), the use of antibiotics is contraindicated because the use of antibiotics may increase the risk of developing HUS through increased STX production (See Wong C. et al., *N Engl J. Med* 342:1930-1936 (2000)). For subjects testing positive for *Shigella* or *Salmonella*, antibiotics are typically administered to clear the infection. Other well established first-line therapy for HUS includes volume expansion, dialysis and plasmapheresis.

In accordance with the foregoing, in some embodiments, in the setting of a subject suffering from one or more symptoms associated with a pre-HUS phase and at risk for developing HUS (*i.e.*, the subject exhibits one or more of the following: diarrhea, a hematocrit level less than 30% with smear evidence of intravascular erythrocyte destruction, thrombocytopenia (platelet count less than $150 \times 10^3/\text{mm}^3$), and/or the

presence of impaired renal function (serum creatinine concentration greater than the upper limit of reference range for age)), a method is provided for decreasing the risk of developing HUS, or of decreasing the likelihood of renal failure in the subject, comprising administering an amount of an MASP-2 inhibitory agent for a time period
5 effective to ameliorate or prevent impaired renal function. In some embodiments, the MASP-2 inhibitory agent is administered for a time period of at least one, two, three, four or more days, and may be repeated as determined by a physician until the condition has been resolved or is controlled. In a pre-HUS setting, the MASP-2 inhibitory agent may be administered to the subject systemically, such as by intra-arterial, intravenous,
10 intramuscular, inhalational, nasal, oral, subcutaneous or other parenteral administration.

The treatment of *E. coli* 0157:H7 infection with bactericidal antibiotics, particularly β -lactams, has been associated with an increased risk of developing HUS (Smith et al., *Pediatr Infect Dis J* 31(1):37-41 (2012)). In some embodiments, in the setting of a subject suffering from symptoms associated with a pre-HUS phase, wherein the
15 subject is known to have an infection with enterogenic *E. coli* for which the use of antibiotics is contra-indicated (e.g., *E. coli* 0157:H7), a method is provided for decreasing the risk of developing HUS, or of decreasing the likelihood of renal failure in the subject, comprising administering an amount of a MASP-2 inhibitory agent for a first time period effective to inhibit or prevent the presence of oligoanuria in the subject (e.g., at least one,
20 two, three or four days), wherein the administration of the MASP-2 inhibitory agent during the first time period occurs in the absence of an antibiotic. In some embodiments, the method further comprises administering the MASP-2 inhibitory agent to the subject in combination with an antibiotic for a second time period (such as at least one to two weeks).

25 In other embodiments, in the setting of a subject suffering from symptoms associated with a pre-HUS phase, wherein the subject is known to have an infection with *Shigella* or *Salmonella*, a method is provided for decreasing the risk of developing HUS, or of decreasing the likelihood of renal failure in the subject, comprising administering an amount of a MASP-2 inhibitory agent and for a time period effective to inhibit or prevent
30 the presence of oligoanuria in the subject, wherein the administration of the MASP-2 inhibitory agent is either in the presence or in the absence of a suitable antibiotic.

In some embodiments, in the setting of an initial diagnosis of HUS, or in a subject exhibiting one or more symptoms consistent with a diagnosis of HUS (e.g., the presence

of renal failure, or microangiopathic hemolytic anemia in the absence of low fibrinogen, or thrombocytopenia) the subject is treated with an effective amount of a MASP-2 inhibitory agent (e.g. a anti-MASP-2 antibody) as a first-line therapy in the absence of plasmapheresis, or in combination with plasmapheresis. As a first-line therapy, the MASP-2 inhibitory agent may be administered to the subject systemically, such as by intra-arterial, intravenous, intramuscular, inhalational, nasal, subcutaneous or other parenteral administration. In some embodiments, the MASP-2 inhibitory agent is administered to a subject as a first line therapy in the absence of plasmapheresis to avoid the complications of plasmapheresis such as hemorrhage, infection, and exposure to disorders and/or allergies inherent in the plasma donor, or in a subject otherwise averse to plasmaphoresis, or in a setting where plasmapheresis is unavailable.

In some embodiments, the method comprises administering a MASP-2 inhibitory agent to a subject suffering from HUS via a catheter (e.g., intravenously) for a first time period (e.g., an acute phase lasting at least one day to a week or two weeks) followed by administering a MASP-2 inhibitory agent to the subject subcutaneously for a second time period (e.g., a chronic phase of at least two weeks or longer). In some embodiments, the administration in the first and/or second time period occurs in the absence of plasmapheresis. In some embodiments, the method further comprises determining the level of at least one complement factor (e.g., C3, C5) in the subject prior to treatment, and optionally during treatment, wherein the determination of a reduced level of the at least one complement factor in comparison to a standard value or healthy control subject is indicative of the need for treatment, and wherein the determination of a normal level is indicative of improvement.

In some embodiments, the method comprises administering a MASP-2 inhibitory agent, such as an anti-MASP-2 antibody, to a subject suffering from, or at risk for developing, HUS either subcutaneously or intravenously. Treatment is preferably daily, but can be as infrequent as weekly or monthly. Treatment will continue for at least one week and as long as 3 months. The anti-MASP-2 antibody may be administered alone, or in combination with a C5 inhibitor, such as eculizumab.

30

TTP:

Thrombotic thrombocytopenic purpura (TTP) is a life threatening disorder of the blood-coagulation system, caused by autoimmune or hereditary dysfunctions that activate

the coagulation cascade or the complement system (George, JN, *N Engl J Med*; 354:1927-35 (2006)). This results in numerous microscopic clots, or thromboses, in small blood vessels throughout the body. Red blood cells are subjected to shear stress which damages their membranes, leading to intravascular hemolysis. The resulting reduced blood flow and endothelial injury results in organ damage, including brain, heart, and kidneys. TTP is clinically characterized by thrombocytopenia, microangiopathic hemolytic anemia, neurological changes, renal failure and fever. In the era before plasma exchange, the fatality rate was 90% during acute episodes. Even with plasma exchange, survival at six months is about 80%.

10 TTP may arise from genetic or acquired inhibition of the enzyme ADAMTS-13, a metalloprotease responsible for cleaving large multimers of von Willebrand factor (vWF) into smaller units. ADAMTS-13 inhibition or deficiency ultimately results in increased coagulation (Tsai, H. *J Am Soc Nephrol* 14: 1072–1081, (2003)). ADAMTS-13 regulates the activity of vWF; in its absence, vWF forms large multimers which are more likely to bind platelets and predisposes patients to platelet aggregation and thrombosis in the microvasculature.

Upshaw-Schulman syndrome (USS, also described as congenital TTP) is a congenital deficiency of ADAMTS13 activity due to ADAMTS13 gene mutations (Schulman et al., *Blood*, 16(1):943-57, 1960; Upshaw et al., *New Engl. J. Med*, 298 (24):1350-2, 1978). Numerous mutations in ADAMTS13 have been identified in individuals with congenital TTP (Kinoshita et al., *International Journal of Hematology*, 74:101-108 (2001); Levy et al., *Nature*, 413 (6855):488-494 (2001); Kokame et al., *PNAS* 99(18):11902-11907 (2002); Savasan et al., *Blood*, 101:4449-4451 (2003); Matsumoto et al., *Blood*, 103:1305-1310 (2004) and Fujimura et al., *Brit. J. Haemat* 144:742-754 (2008)). Subjects with USS typically have 5-10% of normal ADAMTS13 activity (Kokame et al., *PNAS* 99(18):11902-11907, 2002). Although acquired TTP and USS have some similarities, USS has some important differences in clinical features. USS usually presents in infancy or childhood and is characterized by severe hyperbilirubinemia with negative Coombs test soon after birth, response to fresh plasma infusion, and frequent relapses (Savasan et al., *Blood*, 101:4449-4451, 2003). In some cases, patients with this inherited ADAMTS13 deficiency have a mild phenotype at birth and only develop symptoms associated with TTP in clinical situations with increased von Willebrand factor levels, such as infection or pregnancy. For example, Fujimura et al.

reported 9 Japanese women from 6 families with genetically confirmed USS who were diagnosed with the disorder during their first pregnancy. Thrombocytopenia occurred during the second to third trimesters in each of their 15 pregnancies, often followed by TTP. All of these women were found to be severely deficient in ADAMTS13 activity
5 (Fujimura et al., *Brit. J. Haemat* 144:742-754, 2008).

In accordance with the foregoing, in some embodiments, in the setting of a subject with Upshaw-Schulman syndrome (USS) (i.e., the subject is known to be deficient in ADAMTS13 activity and/or the subject is known to have one or more ADAMTS13 gene mutation(s)), a method is provided for decreasing the likelihood of developing clinical
10 symptoms associated with congenital TTP (e.g., thrombocytopenia, anemia, fever, and/or renal failure) comprising administering an amount of a MASP-2 inhibitory agent (e.g., a MASP-2 antibody) for a time period effective to ameliorate or prevent one or more clinical symptoms associated with TTP. In some embodiments, the method further comprises the step of determining whether a subject is at risk for developing symptoms
15 associated with congenital TTP prior to the onset of any symptoms associated with TTP, or upon the onset of at least one or more symptoms indicative of TTP (e.g., the presence of anemia, thrombocytopenia and/or renal insufficiency). The determination of whether a subject is at risk for developing symptoms associated with congenital TTP (i.e., the subject has USS), comprises determining whether the subject has a mutation in the gene
20 encoding ADAMTS13, and/or determining whether the subject is deficient in ADAMTS13 activity, and/or determining whether the subject has a family history of USS. Methods of genetic screening for the presence or absence of a genetic mutation associated with USS are well established, for example see Kinoshita et al., *International Journal of Hematology*, 74:101-108 (2001); Levy et al., *Nature*, 413 (6855):488-494
25 (2001); Kokame et al., *PNAS* 99(18):11902-11907 (2002); Savasan et al., *Blood*, 101:4449-4451 (2003); Matsumoto et al., *Blood*, 103:1305-1310 (2004) and Fujimura et al., *Brit. J. Haemat* 144:742-754 (2008).

In one embodiment, a method is provided for reducing the likelihood that a subject diagnosed with USS will suffer clinical symptoms associated with TTP
30 comprising periodically monitoring the subject to determine the presence or absence of anemia, thrombocytopenia or rising creatinine, and treating with a MASP-2 inhibitory agent (e.g., a MASP-2 antibody) upon the determination of the presence of anemia, thrombocytopenia or rising creatinine, or upon the presence of an event known to be

associated with triggering TTP clinical symptoms, for example, drug exposure (e.g., chemotherapy), infection (e.g. bacterial infection), malignancy, injury, transplant, or pregnancy.

5 In another embodiment, a method is provided for treating a subject with USS and suffering from clinical symptoms associated with TTP comprising administering an amount of a MASP-2 inhibitory agent (e.g., a MASP-2 antibody) for a time period effective to ameliorate or prevent one or more clinical symptoms associated with TTP.

TTP can also develop due to auto-antibodies against ADAMTS-13. In addition, TTP can develop during breast, gastrointestinal tract, or prostate cancer (George JN., 10 *Oncology* (Williston Park). 25:908-14 (2011)), pregnancy (second trimester or postpartum), George JN., *Curr Opin Hematol* 10:339-344 (2003)), or is associated with diseases, such as HIV or autoimmune diseases like systemic lupus erythematosus (Hamasaki K, et al., *Clin Rheumatol*.22:355-8 (2003)). TTP can also be caused by certain drug therapies, including heparin, Quinine, immunemediated ingredient, cancer 15 chemotherapeutic agents (bleomycin, cisplatin, cytosine arabinoside, daunomycin gemcitabine, mitomycin C, and tamoxifen), cyclosporine A, oral contraceptives, penicillin, rifampin and anti-platelet drugs including ticlopidine and clopidogrel (Azarm, T. et al., *J Res Med Sci.*, 16: 353–357 (2011)). Other factors or conditions associated with TTP are toxins such as bee venoms, sepsis, splenic sequestration, transplantation, 20 vasculitis, vascular surgery, and infections like *Streptococcus pneumonia* and cytomegalovirus (Moake JL., *N Engl J Med.*, 347:589–600 (2002)). TTP due to transient functional ADAMTS-13 deficiency can occur as a consequence of endothelial cell injury associated with *S. pneumonia* infection (*Pediatr Nephrol.*, 26:631-5 (2011)).

Plasma exchange is the standard treatment for TTP (Rock GA, et al., *N Engl J 25 Med* 325:393-397 (1991)). Plasma exchange replaces ADAMTS-13 activity in patients with genetic defects and removes ADAMTS-13 autoantibodies in those patients with acquired autoimmune TTP (Tsai, H-M, *Hematol Oncol Clin North Am.*, 21(4): 609–v (2007)). Additional agents such as immunosuppressive drugs are routinely added to therapy (George, JN, *N Engl J Med*, 354:1927-35 (2006)). However, plasma exchange is 30 not successful for about 20% of patients, relapse occurs in more than a third of patients, and plasmapheresis is costly and technically demanding. Furthermore, many patients are unable to tolerate plasma exchange. Consequently there remains a critical need for additional and better treatments for TTP.

Because TTP is a disorder of the blood coagulation cascade, treatment with antagonists of the complement system may aid in stabilizing and correcting the disease. While pathological activation of the alternative complement pathway is linked to aHUS, the role of complement activation in TTP is less clear. The functional deficiency of ADAMTS13 is important for the susceptibility of TTP, however it is not sufficient to cause acute episodes. Environmental factors and/or other genetic variations may contribute to the manifestation of TTP. For example, genes encoding proteins involved in the regulation of the coagulation cascade, vWF, platelet function, components of the endothelial vessel surface, or the complement system may be implicated in the development of acute thrombotic microangiopathy (Galbusera, M. et al., *Haematologica*, 94: 166–170 (2009)). In particular, complement activation has been shown to play a critical role; serum from thrombotic microangiopathy associated with ADAMTS-13 deficiency has been shown to cause C3 and MAC deposition and subsequent neutrophil activation which could be abrogated by complement inactivation (Ruiz-Torres MP, et al., *Thromb Haemost*, 93:443-52 (2005)). In addition, it has recently been shown that during acute episodes of TTP there are increased levels of C4d, C3bBbP, and C3a (M. Réti et al., *J Thromb Haemost*. Feb 28.(2012) doi: 10.1111/j.1538-7836.2012.04674.x. [Epub ahead of print]), consistent with activation of the classical/lectin and alternative pathways. This increased amount of complement activation in acute episodes may initiate the terminal pathway activation and be responsible for further exacerbation of TTP.

The role of ADAMTS-13 and vWF in TTP clearly is responsible for activation and aggregation of platelets and their subsequent role in shear stress and deposition in microangiopathies. Activated platelets interact with and trigger both the classical and alternative pathways of complement. Platelet mediated complement activation increases the inflammatory mediators C3a and C5a (Peerschke E et al., *Mol Immunol*, 47:2170-5 (2010)). Platelets may thus serve as targets of classical complement activation in inherited or autoimmune TTP.

As described above, the lectin pathway of complement, by virtue of MASP-2 mediated prothombin activation, is the dominant molecular pathway linking endothelial injury to the coagulation and microvascular thrombosis that occurs in HUS. Similarly, activation of the lectin pathway of complement may directly drive the coagulation system in TTP. Lectin pathway activation may be initiated in response to the initial endothelium injury caused by ADAMTS-13 deficiency in TTP. It is therefore expected that lectin

pathway inhibitors, including but not limited to antibodies that block MASP-2 function, will mitigate the microangiopathies associated with microvascular coagulation, thrombosis, and hemolysis in patients suffering from TTP.

Patients suffering from TTP typically present in the emergency room with one or more of the following: purpura, renal failure, low platelets, anemia and/or thrombosis, including stroke. The current standard of care for TTP involves intra-catheter delivery (e.g., intravenous or other form of catheter) of replacement plasmapheresis for a period of two weeks or longer, typically three times a week, but up to daily. If the subject tests positive for the presence of an inhibitor of ADAMTS13 (i.e., an endogenous antibody against ADAMTS13), then the plasmapheresis may be carried out in combination with immunosuppressive therapy (e.g., corticosteroids, rituxan, or cyclosporine). Subjects with refractory TTP (approximately 20% of TTP patients) do not respond to at least two weeks of plasmapheresis therapy.

In accordance with the foregoing, in one embodiment, in the setting of an initial diagnosis of TTP or in a subject exhibiting one or more symptoms consistent with a diagnosis of TTP (e.g., central nervous system involvement, severe thrombocytopenia (a platelet count of less than or equal to 5000/ μ L if off aspirin, less than or equal to 20,000/ μ L if on aspirin), severe cardiac involvement, severe pulmonary involvement, gastro-intestinal infarction or gangrene), a method is provided for treating the subject with an effective amount of a MASP-2 inhibitory agent (e.g., a anti-MASP-2 antibody) as a first line therapy in the absence of plasmapheresis, or in combination with plasmapheresis. As a first-line therapy, the MASP-2 inhibitory agent may be administered to the subject systemically, such as by intra-arterial, intravenous, intramuscular, inhalational, nasal, subcutaneous or other parenteral administration. In some embodiments, the MASP-2 inhibitory agent is administered to a subject as a first-line therapy in the absence of plasmapheresis to avoid the potential complications of plasmapheresis, such as hemorrhage, infection, and exposure to disorders and/or allergies inherent in the plasma donor, or in a subject otherwise averse to plasmapheresis, or in a setting where plasmapheresis is unavailable. In some embodiments, the MASP-2 inhibitory agent is administered to the subject suffering from TTP in combination (including co-administration) with an immunosuppressive agent (e.g., corticosteroids, rituxan or cyclosporine) and/or in combination with concentrated ADAMTS-13.

In some embodiments, the method comprises administering a MASP-2 inhibitory agent to a subject suffering from TTP via a catheter (e.g., intravenously) for a first time period (e.g., an acute phase lasting at least one day to a week or two weeks) followed by administering a MASP-2 inhibitory agent to the subject subcutaneously for a second time
5 period (e.g., a chronic phase of at least two weeks or longer). In some embodiments, the administration in the first and/or second time period occurs in the absence of plasmapheresis. In some embodiments, the method is used to maintain the subject to prevent the subject from suffering one or more symptoms associated with TTP.

In another embodiment, a method is provided for treating a subject suffering from
10 refractory TTP (i.e., a subject that has not responded to at least two weeks of plasmaphoresis therapy), by administering an amount of a MASP-2 inhibitor effective to reduce one or more symptoms of TTP. In one embodiment, the MASP-2 inhibitor (e.g., an anti-MASP-2 antibody) is administered to a subject with refractory TTP on a chronic basis, over a time period of at least two weeks or longer via subcutaneous or other
15 parenteral administration. Administration may be repeated as determined by a physician until the condition has been resolved or is controlled.

In some embodiments, the method further comprises determining the level of at least one complement factor (e.g., C3, C5) in the subject prior to treatment, and optionally during treatment, wherein the determination of a reduced level of the at least
20 one complement factor in comparison to a standard value or healthy control subject is indicative of the need for continued treatment with the MASP-2 inhibitory agent.

In some embodiments, the method comprises administering a MASP-2 inhibitory agent, such as an anti-MASP-2 antibody, to a subject suffering from, or at risk for developing, TTP either subcutaneously or intravenously. Treatment is preferably daily,
25 but can be as infrequent as biweekly. Treatment is continued until the subject's platelet count is greater than 150,000/ml for at least two consecutive days. The anti-MASP-2 antibody may be administered alone, or in combination with a C5 inhibitor, such as eculizumab.

In one embodiment, the MASP-2 inhibitory antibody exhibits at least one or more
30 of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or less, said antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM

or less, wherein the antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation, and/or wherein the antibody does not substantially inhibit the classical pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the alternative pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway or the alternative pathway (i.e., inhibits the lectin pathway while leaving the classical and alternative complement pathways intact).

In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a subject suffering from TTP by at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum. In some embodiments, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a subject suffering from TTP at a level of at least 20 percent or greater, (such as at least 30%, at least 40%, at least 50%) more than the inhibitory effect on C5b-9 deposition in serum.

In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a TTP patient by at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum.

In one embodiment, the MASP-2 inhibitory antibody is administered to the subject via an intravenous catheter or other catheter delivery method.

In one embodiment, the invention provides a method of inhibiting thrombus formation in a subject suffering from TTP comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising (I) (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-102 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-

L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70.

10 In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67. In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof, that specifically recognizes at least part of an epitope on human MASP-2 recognized by reference antibody OMS646 comprising a heavy-chain variable region as set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70.

Degos Disease

25 Degos disease, also known as malignant atrophic papulosis, is a very rare TMA affecting the endothelium of small vessels of skin, gastrointestinal tract, and CNS. This vasculopathy causes occlusion of venules and arterioles, resulting in skin lesions, bowel ischemia, and CNS disorders including strokes, epilepsy and cognitive disorders. In the skin, connective tissue necrosis is due to thrombotic occlusion of the small arteries. However, the cause of Degos disease is unknown. Vasculitis, coagulopathy, or primary dysfunction of the endothelial cells have been implicated. Degos disease has a 50% survival of only two to three years. There is no effective treatment for Degos disease

although antiplatelet drugs, anticoagulants, and immunosuppressants are utilized to alleviate symptoms.

While the mechanism of Degos disease is unknown, the complement pathway has been implicated. Margo et al., identified prominent C5b-9 deposits in skin, gastrointestinal tract and brain vessels of four terminal patients with Degos disease (Margo et al., *Am J Clin Pathol* 135(4):599-610, 2011). Experimental treatment with eculizumab was initially effective in the treatment of skin and intestinal lesions, but did not prevent the progression of systemic disease (see Garrett-Bakelman F. et al., "C5b-9 is a potential effector in the pathophysiology of Degos disease; a case report of treatment with eculizumab" (Abstract), Jerusalem: International Society of Hematology; 2010, Poster #156; and Polito J. et al, "Early detection of systemic Degos disease (DD) or malignant atrophic papulosis (MAP) may increase survival" (Abstract), San Antonio, TX: American College of Gastroenterology; 2010, Poster #1205).

Many patients suffering from Degos disease have defects of blood coagulation. Thrombotic occlusion of small arteries in the skin is characteristic of the disease. Because the complement pathway is implicated in this disease, as described herein for other TMAs, it is expected that lectin pathway inhibitors, including but not limited to antibodies that block MASP-2 function, will be beneficial in treating patients suffering from Degos disease.

Accordingly, in another embodiment, the invention provides methods for treating Degos disease by administering a composition comprising a therapeutically effective amount of a MASP-2 inhibitory agent, such as a MASP-2 antibody, in a pharmaceutical carrier to a subject suffering from Degos disease or a condition resulting from Degos disease. The MASP-2 inhibitory agent is administered systemically to the subject suffering from Degos disease or a condition resulting from Degos disease, such as by intra-arterial, intravenous, intramuscular, inhalational, subcutaneous or other parenteral administration, or potentially by oral administration for non-peptidergic agents. The anti-MASP-2 antibody may be administered alone, or in combination with a C5 inhibitor, such as eculizumab.

In one embodiment, the MASP-2 inhibitory antibody exhibits at least one or more of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or

less, said antibody inhibits C3b deposition in 90% human serum with an IC₅₀ of 30 nM or less, wherein the antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation, and/or wherein the antibody does not substantially inhibit the classical pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the alternative pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway or the alternative pathway (i.e., inhibits the lectin pathway while leaving the classical and alternative complement pathways intact).

In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a subject suffering from Degos disease by at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum. In some embodiments, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a subject suffering from Degos disease at a level of at least 20 percent or greater, (such as at least 30%, at least 40%, at least 50%) more than the inhibitory effect on C5b-9 deposition in serum.

In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a Degos disease patient by at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum.

In one embodiment, the MASP-2 inhibitory antibody is administered to the subject via an intravenous catheter or other catheter delivery method.

In one embodiment, the invention provides a method of inhibiting thrombus formation in a subject suffering from Degos disease comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising (I) (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65

of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-102 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67. In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof, that specifically recognizes at least part of an epitope on human MASP-2 recognized by reference antibody OMS646 comprising a heavy-chain variable region as set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70.

Catastrophic antiphospholipid syndrome (CAPS)

Catastrophic antiphospholipid syndrome (CAPS) is an extreme variant of the antiphospholipid antibody (APLA) syndrome. CAPS is characterized by venous and arterial thrombosis due to pathogenic antibodies. CAPS is a TMA with multiple organ thrombosis, ischemia, and organ failure. Like other TMAs, occlusion of small vessels in various organs is characteristic. There is a high mortality rate in CAPS of about 50% and

often it is associated with infection or trauma. Patients have antiphospholipid antibodies, generally IgG.

Clinically, CAPS involves at least three organs or tissues with histopathological evidence of small vessel occlusion. Peripheral thrombosis may involve veins and arteries
5 in the CNS, cardiovascular, renal, or pulmonary systems. Patients are treated with antibiotics, anticoagulants, corticosteroids, plasma exchange, and intravenous immunoglobulin. Nevertheless, death may result from multiple organ failure.

The complement pathway has been implicated in CAPS. For example, studies in animal models indicate that complement inhibition may be an effective means to prevent
10 thrombosis associated with CAPS (Shapira L. et al., *Arthritis Rheum* 64(8):2719-23, 2012). Moreover, as further reported by Shapira et al., administration of eculizumab to a subject suffering from CAPS at doses that blocked complement pathway aborted acute progressive thrombotic events and reversed thrombocytopenia (see also Lim W., *Curr Opin Hematol* 18(5):361-5, 2011). Therefore, as described herein for other TMAs, it is
15 expected that lectin pathway inhibitors, including but not limited to antibodies that block MASP-2 function, will be beneficial in treating patients suffering from CAPS.

Accordingly, in another embodiment, the invention provides methods for treating CAPS by administering a composition comprising a therapeutically effective amount of a MASP-2 inhibitory agent, such as a MASP-2 antibody, in a pharmaceutical carrier to a
20 subject suffering from CAPS or a condition resulting from CAPS. The MASP-2 inhibitory agent is administered systemically to the subject suffering from CAPS or a condition resulting from CAPS, such as by intra-arterial, intravenous, intramuscular, inhalational, subcutaneous or other parenteral administration, or potentially by oral administration for non-peptidergic agents. The anti-MASP-2 antibody may be
25 administered alone, or in combination with a C5 inhibitor, such as eculizumab.

In one embodiment, the MASP-2 inhibitory antibody exhibits at least one or more of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or
30 less, said antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less, wherein the antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂, wherein the antibody is a single-chain molecule,

wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation, and/or wherein the antibody does not substantially inhibit the classical pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the
5 lectin pathway and does not substantially inhibit the alternative pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway or the alternative pathway (i.e., inhibits the lectin pathway while leaving the classical and alternative complement pathways intact).

10 In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a subject suffering from CAPS by at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum. In some embodiments, the MASP-2 inhibitory antibody inhibits thrombus
15 formation in serum from a subject suffering from CAPS at a level of at least 20 percent or greater, (such as at least 30%, at least 40%, at least 50%) more than the inhibitory effect on C5b-9 deposition in serum.

In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a CAPS patient by at least 30%, such as at least 40%, such as at least 50%,
20 such as at least 60%, such as at least 70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum.

In one embodiment, the MASP-2 inhibitory antibody is administered to the subject via an intravenous catheter or other catheter delivery method.

In one embodiment, the invention provides a method of inhibiting thrombus
25 formation in a subject suffering from CAPS comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising (I) (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID
30 NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-102 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and

- iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67. In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof, that specifically recognizes at least part of an epitope on human MASP-2 recognized by reference antibody OMS646 comprising a heavy-chain variable region as set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70.

TMA Secondary to Cancer

Systemic malignancies of any type can lead to clinical and pathologic manifestations of TMA (see e.g., Batts and Lazarus, *Bone Marrow Transplantation* 40:709-719, 2007). Cancer-associated TMA is often found in the lungs and appears to be associated with tumor emboli (Francis KK et al., *Commun Oncol* 2:339-43, 2005). Tumor emboli can reduce blood flow and thus lead to a hypo-perfused state in the affected arterioles and venules. The resulting tissue stress and injury is expected to activate the lectin pathway of complement in a localized fashion. The activated lectin pathway in turn can activate the coagulation cascade via MASP-2 dependent cleavage of prothrombin to thrombin, leading to a pro-thrombotic state characteristic of TMA.

MASP-2 inhibition in this setting is expected to reduce the localized activation of thrombin and thereby alleviate the pro-thrombotic state.

Therefore, as described herein for other TMAs, it is expected that lectin pathway inhibitors, including, but not limited to, antibodies that block MASP-2 function, will be
5 beneficial in treating patients suffering from TMA secondary to cancer.

Accordingly, in another embodiment, the invention provides methods for treating or preventing TMA secondary to cancer by administering a composition comprising a therapeutically effective amount of a MASP-2 inhibitory agent, such as a MASP-2 antibody, in a pharmaceutical carrier to a subject suffering from, or at risk for developing,
10 a TMA secondary to cancer. The MASP-2 inhibitory agent is administered systemically to the subject suffering from, or at risk for developing, a TMA secondary to cancer, such as by intra-arterial, intravenous, intramuscular, inhalational, subcutaneous or other parenteral administration, or potentially by oral administration for non-peptidergic agents. The anti-MASP-2 antibody may be administered alone, or in combination with a C5
15 inhibitor, such as eculizumab.

In one embodiment, the MASP-2 inhibitory antibody exhibits at least one or more of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or
20 less, said antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less, wherein the antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a
25 S228P mutation, and/or wherein the antibody does not substantially inhibit the classical pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the alternative pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway or the alternative pathway (i.e.,
30 inhibits the lectin pathway while leaving the classical and alternative complement pathways intact).

In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation

in serum from a subject suffering from TMA secondary to cancer by at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum.

5 In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a patient suffering TMA secondary to cancer by at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum.

10 In one embodiment, the MASP-2 inhibitory antibody is administered to the subject via an intravenous catheter or other catheter delivery method.

 In one embodiment, the invention provides a method of inhibiting thrombus formation in a subject suffering from TMA secondary to cancer comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or
15 antigen binding fragment thereof, comprising (I) (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-102 of SEQ ID NO:67 and b) a light-chain variable region comprising:
20 i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at
25 least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70).

 In some embodiments, the method comprises administering to the subject a
30 composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67. In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory

antibody, or antigen binding fragment thereof, comprising a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof, that specifically recognizes at least part of an epitope on human MASP-2 recognized by reference antibody OMS646 comprising a heavy-chain variable region as set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70.

10 **TMA Secondary to Cancer Chemotherapy**

Chemotherapy-associated TMA is a condition involving thrombocytopenia, microangiopathic hemolytic anemia, and renal dysfunction that develops in 2-10% of patients with a history of malignant neoplasms treated with chemotherapeutic agents such as gemcytabin, mitomycin, oxaliplatin and others. Chemotherapy-associated TMA is associated with high mortality poor clinical outcomes (see, e.g., Blake-Haskins et al., *Clin Cancer Res* 17(18):5858-5866, 2011).

The etiology of chemotherapy-associated TMA is thought to encompass a non-specific, toxic insult to the microvascular endothelium. A direct injury to endothelial cells has been shown in an animal model of mitomycin-induced TMA (Dlott J. et al., *Ther Apher Dial* 8:102-11, 2004). Endothelial cell injury through a variety of mechanisms has been shown to activate the lectin pathway of complement. For example, Stahl et al. have shown that endothelial cells exposed to oxidative stress activate the lectin pathway of complement both *in vitro* and *in vivo* (Collard et al., *Am J Pathol.* 156(5):1549-56, 2000; La Bonte et al., *J Immunol.* 15;188(2):885-91, 2012). *In vivo*, this process leads to thrombosis, and inhibition of the lectin pathway has been shown to prevent thrombosis (La Bonte et al. *J Immunol.* 15;188(2):885-91, 2012). Furthermore, as demonstrated in Examples 37-39 herein, in the mouse model of TMA where localized photoexcitation of FITC-Dex was used to induce localized injury to the microvasculature with subsequent development of a TMA response, the present inventors have shown that inhibition of MASP-2 can prevent TMA. Thus, microvascular endothelium injury by chemotherapeutic agents may activate the lectin pathway of complement which then creates a localized pro-thrombotic state and promotes a TMA response. Since activation of the lectin pathway and the creation of a pro-thrombotic state is MASP-2-dependent, it is

expected that MASP-2 inhibitors, including, but not limited to, antibodies that block MASP-2 function, will alleviate the TMA response and reduce the risk of cancer chemotherapy-associated TMA.

Accordingly, in another embodiment, the invention provides methods for treating
5 or preventing TMA secondary to chemotherapy by administering a composition comprising a therapeutically effective amount of a MASP-2 inhibitory agent, such as a MASP-2 antibody, in a pharmaceutical carrier to a subject suffering from, or at risk for developing, a TMA secondary to chemotherapy. The MASP-2 inhibitory agent is administered systemically to a subject that has undergone, is undergoing, or will undergo
10 chemotherapy, such as by intra-arterial, intravenous, intramuscular, inhalational, subcutaneous or other parenteral administration, or potentially by oral administration for non-peptidergic agents. The anti-MASP-2 antibody may be administered alone, or in combination with a C5 inhibitor, such as eculizumab.

In one embodiment, the MASP-2 inhibitory antibody exhibits at least one or more
15 of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or less, said antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less, wherein the antibody is an antibody fragment selected from the group consisting
20 of Fv, Fab, Fab', F(ab)₂ and F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation, and/or wherein the antibody does not substantially inhibit the classical pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the
25 lectin pathway and does not substantially inhibit the alternative pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway or the alternative pathway (i.e., inhibits the lectin pathway while leaving the classical and alternative complement pathways intact).

30 In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a subject suffering from TMA secondary to cancer chemotherapy by at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least

70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum.

In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a patient suffering TMA secondary to cancer chemotherapy by at least
5 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum.

In one embodiment, the MASP-2 inhibitory antibody is administered to the subject via an intravenous catheter or other catheter delivery method.

10 In one embodiment, the invention provides a method of inhibiting thrombus formation in a subject suffering from TMA secondary to cancer chemotherapy comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising (I) (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid
15 sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-102 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid
20 sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with
25 at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70).

In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid
30 sequence set forth as SEQ ID NO:67. In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof, that specifically recognizes at least part of an epitope on human MASP-2 recognized by reference antibody OMS646 comprising a heavy-chain variable region as set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70.

TMA Secondary to Transplantation

Transplantation-associated TMA (TA-TMA) is a devastating syndrome that can occur in transplant patients, such as allogeneic hematopoietic stem cell transplant recipients (see e.g., Batts and Lazarus, *Bone Marrow Transplantation* 40:709-719, 2007). The pathogenesis of this condition is poorly understood, but likely involves a confluence of responses that culminate in endothelial cell injury (Laskin B.L. et al., *Blood* 118(6):1452-62, 2011). As discussed above, endothelial cell injury is a prototypic stimulus for lectin pathway activation and the generation of a pro-thrombotic environment.

Recent data further support the role of complement activation via the lectin pathway in the pathogenesis TA-TMA. Laskin et al., have demonstrated that renal arteriolar C4d deposition was much more common in subjects with histologic TA-TMA (75%) compared with controls (8%) (Laskin B.L., et al., *Transplantation*, 27; 96(2):217-23, 2013). Thus, C4d may be a pathologic marker of TA-TMA, implicating localized complement fixation via the lectin or classical pathway.

Since activation of the lectin pathway and the creation of a pro-thrombotic state is MASP-2-dependent, it is expected that MASP-2 inhibitors, including, but not limited to, antibodies that block MASP-2 function, will alleviate the TMA response and reduce the risk of transplantation-associated TMA (TA-TMA).

Accordingly, in another embodiment, the invention provides methods for treating or preventing a TMA secondary to transplantation by administering a composition comprising a therapeutically effective amount of a MASP-2 inhibitory agent, such as a MASP-2 antibody, in a pharmaceutical carrier to a subject suffering from, or at risk for developing a TMA secondary to transplantation. The MASP-2 inhibitory agent is administered systemically to a subject that has undergone, is undergoing, or will undergo a transplant procedure, such as by intra-arterial, intravenous, intramuscular, inhalational,

subcutaneous or other parenteral administration, or potentially by oral administration for non-peptidergic agents. The anti-MASP-2 antibody may be administered alone, or in combination with a C5 inhibitor, such as eculizumab. In some embodiments, the invention provides methods for treating or preventing a TMA secondary to allogeneic stem cell transplant comprising administering a composition comprising an amount of a MASP-2 inhibitory agent, such as a MASP-2 inhibitory antibody, to a subject prior to, during or after undergoing an allogeneic stem cell transplant.

In one embodiment, the MASP-2 inhibitory antibody exhibits at least one or more of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or less, said antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less, wherein the antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation, and/or wherein the antibody does not substantially inhibit the classical pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the alternative pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway or the alternative pathway (i.e., inhibits the lectin pathway while leaving the classical and alternative complement pathways intact).

In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a subject suffering from TMA secondary to transplant by at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum.

In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a patient suffering TMA secondary to transplant by at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as

compared to untreated serum.

In one embodiment, the MASP-2 inhibitory antibody is administered to the subject via an intravenous catheter or other catheter delivery method.

In one embodiment, the invention provides a method of inhibiting thrombus
5 formation in a subject suffering from TMA secondary to transplant comprising
administering to the subject a composition comprising an amount of a MASP-2 inhibitory
antibody, or antigen binding fragment thereof, comprising (I) (a) a heavy-chain variable
region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from
31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid
10 sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the
amino acid sequence from 95-102 of SEQ ID NO:67 and b) a light-chain variable region
comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of
SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from
50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid
15 sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-
chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at
least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least
98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with at
least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least
20 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a
composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding
fragment thereof, comprising a heavy-chain variable region comprising the amino acid
sequence set forth as SEQ ID NO:67. In some embodiments, the method comprises
25 administering to the subject a composition comprising an amount of a MASP-2 inhibitory
antibody, or antigen binding fragment thereof, comprising a light-chain variable region
comprising the amino acid sequence set forth as SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a
composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment
30 thereof, that specifically recognizes at least part of an epitope on human MASP-2
recognized by reference antibody OMS646 comprising a heavy-chain variable region as
set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID
NO:70.

IV. THE ROLE OF MASP-2 IN OTHER DISEASES AND CONDITIONS AND THERAPEUTIC METHODS USING MASP-2 INHIBITORY AGENTS

5 RENAL CONDITIONS

Activation of the complement system has been implicated in the pathogenesis of a wide variety of renal diseases; including, mesangioproliferative glomerulonephritis (IgA-nephropathy, Berger's disease) (Endo, M., et al., *Clin. Nephrology* 55:185-191, 2001), membranous glomerulonephritis (Kerjashki, D., *Arch B Cell Pathol.* 58:253-71, 10 1990; Brenchley, P.E., et al., *Kidney Int.*, 41:933-7, 1992; Salant, D.J., et al., *Kidney Int.* 35:976-84, 1989), membranoproliferative glomerulonephritis (mesangiocapillary glomerulonephritis) (Bartlow, B.G., et al., *Kidney Int.* 15:294-300, 1979; Meri, S., et al., *J. Exp. Med.* 175:939-50, 1992), acute postinfectious glomerulonephritis (poststreptococcal glomerulonephritis), cryoglobulinemic glomerulonephritis (Ohsawa, 15 I., et al., *Clin Immunol.* 101:59-66, 2001), lupus nephritis (Gatenby, P.A., *Autoimmunity* 11:61-6, 1991), and Henoch-Schonlein purpura nephritis (Endo, M., et al., *Am. J. Kidney Dis.* 35:401-407, 2000). The involvement of complement in renal disease has been appreciated for several decades but there is still a major discussion on its exact role in the onset, the development and the resolution phase of renal disease. Under 20 normal conditions the contribution of complement is beneficial to the host, but inappropriate activation and deposition of complement may contribute to tissue damage.

There is substantial evidence that glomerulonephritis, inflammation of the glomeruli, is often initiated by deposition of immune complexes onto glomerular or tubular structures which then triggers complement activation, inflammation and tissue 25 damage. Kahn and Sinniah demonstrated increased deposition of C5b-9 in tubular basement membranes in biopsies taken from patients with various forms of glomerulonephritis (Kahn, T.N., et al., *Histopath.* 26:351-6, 1995). In a study of patients with IgA nephropathy (Alexopoulos, A., et al., *Nephrol. Dial. Transplant* 10:1166-1172, 1995), C5b-9 deposition in the tubular epithelial/basement membrane structures 30 correlated with plasma creatinine levels. Another study of membranous nephropathy demonstrated a relationship between clinical outcome and urinary sC5b-9 levels (Kon, S.P., et al., *Kidney Int.* 48:1953-58, 1995). Elevated sC5b-9 levels were correlated positively with poor prognosis. Lehto et al., measured elevated levels of CD59, a

complement regulatory factor that inhibits the membrane attack complex in plasma membranes, as well as C5b-9 in urine from patients with membranous glomerulonephritis (Lehto, T., et al., *Kidney Int.* 47:1403-11, 1995). Histopathological analysis of biopsy samples taken from these same patients demonstrated deposition of C3 and C9 proteins in the glomeruli, whereas expression of CD59 in these tissues was diminished compared to that of normal kidney tissue. These various studies suggest that ongoing complement-mediated glomerulonephritis results in urinary excretion of complement proteins that correlate with the degree of tissue damage and disease prognosis.

Inhibition of complement activation in various animal models of glomerulonephritis has also demonstrated the importance of complement activation in the etiology of the disease. In a model of membranoproliferative glomerulonephritis (MPGN), infusion of anti-Thy1 antiserum in C6-deficient rats (that cannot form C5b-9) resulted in 90% less glomerular cellular proliferation, 80% reduction in platelet and macrophage infiltration, diminished collagen type IV synthesis (a marker for mesangial matrix expansion), and 50% less proteinuria than in C6+ normal rats (Brandt, J., et al., *Kidney Int.* 49:335-343, 1996). These results implicate C5b-9 as a major mediator of tissue damage by complement in this rat anti-thymocyte serum model. In another model of glomerulonephritis, infusion of graded dosages of rabbit anti-rat glomerular basement membrane produced a dose-dependent influx of polymorphonuclear leukocytes (PMN) that was attenuated by prior treatment with cobra venom factor (to consume complement) (Scandrett, A.L., et al., *Am. J. Physiol.* 268:F256-F265, 1995). Cobra venom factor-treated rats also showed diminished histopathology, decreased long-term proteinuria, and lower creatinine levels than control rats. Employing three models of GN in rats (anti-thymocyte serum, Con A anti-Con A, and passive Heymann nephritis), Couser et al., demonstrated the potential therapeutic efficacy of approaches to inhibit complement by using the recombinant sCR1 protein (Couser, W.G., et al., *J. Am. Soc. Nephrol.* 5:1888-94, 1995). Rats treated with sCR1 showed significantly diminished PMN, platelet and macrophage influx, decreased mesangiolysis, and proteinuria versus control rats. Further evidence for the importance of complement activation in glomerulonephritis has been provided by the use of an anti-C5 MoAb in the NZB/W F1 mouse model. The anti-C5 MoAb inhibits cleavage of C5, thus blocking generation of C5a and C5b-9. Continuous therapy with anti-C5 MoAb for 6 months resulted in significant amelioration of the course of glomerulonephritis. A humanized anti-C5

MoAb monoclonal antibody (5G1.1) that prevents the cleavage of human complement component C5 into its pro-inflammatory components is under development by Alexion Pharmaceuticals, Inc., New Haven, Connecticut, as a potential treatment for glomerulonephritis.

5 Direct evidence for a pathological role of complement in renal injury is provided by studies of patients with genetic deficiencies in specific complement components. A number of reports have documented an association of renal disease with deficiencies of complement regulatory factor H (Ault, B.H., *Nephrol.* 14:1045-1053, 2000; Levy, M., et al., *Kidney Int.* 30:949-56, 1986; Pickering, M.C., et al., *Nat. Genet.* 31:424-8,
10 2002). Factor H deficiency results in low plasma levels of factor B and C3 and in consumption of C5b-9. Both atypical membranoproliferative glomerulonephritis (MPGN) and idiopathic hemolytic uremic syndrome (HUS) are associated with factor H deficiency. Factor H deficient pigs (Jansen, J.H., et al., *Kidney Int.* 53:331-49, 1998) and factor H knockout mice (Pickering, M.C., 2002) display MPGN-like symptoms,
15 confirming the importance of factor H in complement regulation. Deficiencies of other complement components are associated with renal disease, secondary to the development of systemic lupus erythematosus (SLE) (Walport, M.J., Davies, et al., *Ann. N.Y. Acad. Sci.* 815:267-81, 1997). Deficiency for C1q, C4 and C2 predispose strongly to the development of SLE via mechanisms relating to defective clearance of immune
20 complexes and apoptotic material. In many of these SLE patients lupus nephritis occurs, characterized by the deposition of immune complexes throughout the glomerulus.

 Further evidence linking complement activation and renal disease has been provided by the identification in patients of autoantibodies directed against complement components, some of which have been directly related to renal disease (Trouw, L.A., et al., *Mol. Immunol.* 38:199-206, 2001). A number of these autoantibodies show
25 such a high degree of correlation with renal disease that the term nephritic factor (NeF) was introduced to indicate this activity. In clinical studies, about 50% of the patients positive for nephritic factors developed MPGN (Spitzer, R.E., et al., *Clin. Immunol. Immunopathol.* 64:177-83, 1992). C3NeF is an autoantibody directed against the
30 alternative pathway C3 convertase (C3bBb) and it stabilizes this convertase, thereby promoting alternative pathway activation (Daha, M.R., et al., *J. Immunol.* 116:1-7, 1976). Likewise, autoantibody with a specificity for the classical pathway C3 convertase (C4b2a), called C4NeF, stabilizes this convertase and thereby promotes classical pathway

activation (Daha, M.R. et al., *J. Immunol.* 125:2051-2054, 1980; Halbwachs, L., et al., *J. Clin. Invest.* 65:1249-56, 1980). Anti-C1q autoantibodies have been described to be related to nephritis in SLE patients (Hovath, L., et al., *Clin. Exp. Rheumatol.* 19:667-72, 2001; Siegert, C., et al., *J. Rheumatol.* 18:230-34, 1991; Siegert, C., et al., *Clin. Exp. Rheumatol.* 10:19-23, 1992), and a rise in the titer of these anti-C1q autoantibodies was reported to predict a flare of nephritis (Coremans, I.E., et al., *Am. J. Kidney Dis.* 26:595-601, 1995). Immune deposits eluted from postmortem kidneys of SLE patients revealed the accumulation of these anti-C1q autoantibodies (Mannick, M., et al., *Arthritis Rheumatol.* 40:1504-11, 1997). All these facts point to a pathological role for these autoantibodies. However, not all patients with anti-C1q autoantibodies develop renal disease and also some healthy individuals have low titer anti-C1q autoantibodies (Siegert, C.E., et al., *Clin. Immunol. Immunopathol.* 67:204-9, 1993).

In addition to the alternative and classical pathways of complement activation, the lectin pathway may also have an important pathological role in renal disease. Elevated levels of MBL, MBL-associated serine protease and complement activation products have been detected by immunohistochemical techniques on renal biopsy material obtained from patients diagnosed with several different renal diseases, including Henoch-Schonlein purpura nephritis (Endo, M., et al., *Am. J. Kidney Dis.* 35:401-407, 2000), cryoglobulinemic glomerulonephritis (Ohsawa, I., et al., *Clin. Immunol.* 101:59-66, 2001) and IgA neuropathy (Endo, M., et al., *Clin. Nephrology* 55:185-191, 2001). Therefore, despite the fact that an association between complement and renal diseases has been known for several decades, data on how complement exactly influences these renal diseases is far from complete.

BLOOD DISORDERS

Sepsis is caused by an overwhelming reaction of the patient to invading microorganisms. A major function of the complement system is to orchestrate the inflammatory response to invading bacteria and other pathogens. Consistent with this physiological role, complement activation has been shown in numerous studies to have a major role in the pathogenesis of sepsis (Bone, R.C., *Annals. Internal. Med.* 115:457-469, 1991). The definition of the clinical manifestations of sepsis is ever evolving. Sepsis is usually defined as the systemic host response to an infection. However, on many occasions, no clinical evidence for infection (e.g., positive bacterial blood cultures) is found in patients with septic symptoms. This discrepancy was first taken into account at

a Consensus Conference in 1992 when the term "systemic inflammatory response syndrome" (SIRS) was established, and for which no definable presence of bacterial infection was required (Bone, R.C., et al., *Crit. Care Med.* 20:724-726, 1992). There is now general agreement that sepsis and SIRS are accompanied by the inability to regulate the inflammatory response. For the purposes of this brief review, we will consider the clinical definition of sepsis to also include severe sepsis, septic shock, and SIRS.

The predominant source of infection in septic patients before the late 1980s was Gram-negative bacteria. Lipopolysaccharide (LPS), the main component of the Gram-negative bacterial cell wall, was known to stimulate release of inflammatory mediators from various cell types and induce acute infectious symptoms when injected into animals (Haeney, M.R., et al., *Antimicrobial Chemotherapy* 41(Suppl. A):41-6, 1998). Interestingly, the spectrum of responsible microorganisms appears to have shifted from predominantly Gram-negative bacteria in the late 1970s and 1980s to predominantly Gram-positive bacteria at present, for reasons that are currently unclear (Martin, G.S., et al., *N. Eng. J. Med.* 348:1546-54, 2003).

Many studies have shown the importance of complement activation in mediating inflammation and contributing to the features of shock, particularly septic and hemorrhagic shock. Both Gram-negative and Gram-positive organisms commonly precipitate septic shock. LPS is a potent activator of complement, predominantly via the alternative pathway, although classical pathway activation mediated by antibodies also occurs (Fearon, D.T., et al., *N. Engl. J. Med.* 292:937-400, 1975). The major components of the Gram-positive cell wall are peptidoglycan and lipoteichoic acid, and both components are potent activators of the alternative complement pathway, although in the presence of specific antibodies they can also activate the classical complement pathway (Joiner, K.A., et al., *Ann. Rev. Immunol.* 2:461-2, 1984).

The complement system was initially implicated in the pathogenesis of sepsis when it was noted by researchers that anaphylatoxins C3a and C5a mediate a variety of inflammatory reactions that might also occur during sepsis. These anaphylatoxins evoke vasodilation and an increase in microvascular permeability, events that play a central role in septic shock (Schumacher, W.A., et al., *Agents Actions* 34:345-349, 1991). In addition, the anaphylatoxins induce bronchospasm, histamine release from mast cells, and aggregation of platelets. Moreover, they exert numerous effects on granulocytes, such as chemotaxis, aggregation, adhesion, release of lysosomal enzymes, generation of toxic

- super oxide anion and formation of leukotrienes (Shin, H.S., et al., *Science* 162:361-363, 1968; Vogt, W., *Complement* 3:177-86, 1986). These biologic effects are thought to play a role in development of complications of sepsis such as shock or acute respiratory distress syndrome (ARDS) (Hammerschmidt, D.E., et al., *Lancet* 1:947-949, 1980;
- 5 Slotman, G.T., et al., *Surgery* 99:744-50, 1986). Furthermore, elevated levels of the anaphylatoxin C3a is associated with a fatal outcome in sepsis (Hack, C.E., et al., *Am. J. Med.* 86:20-26, 1989). In some animal models of shock, certain complement-deficient strains (e.g., C5-deficient ones) are more resistant to the effects of LPS infusions (Hseuh, W., et al., *Immunol.* 70:309-14, 1990).
- 10 Blockade of C5a generation with antibodies during the onset of sepsis in rodents has been shown to greatly improve survival (Czermak, B.J., et al., *Nat. Med.* 5:788-792, 1999). Similar findings were made when the C5a receptor (C5aR) was blocked, either with antibodies or with a small molecular inhibitor (Huber-Lang, M.S., et al., *FASEB J.* 16:1567-74, 2002; Riedemann, N.C., et al., *J. Clin. Invest.* 110:101-8, 2002). Earlier
- 15 experimental studies in monkeys have suggested that antibody blockade of C5a attenuated *E. coli*-induced septic shock and adult respiratory distress syndrome (Hangen, D.H., et al., *J. Surg. Res.* 46:195-9, 1989; Stevens, J.H., et al., *J. Clin. Invest.* 77:1812-16, 1986). In humans with sepsis, C5a was elevated and associated with significantly reduced survival rates together with multiorgan failure, when compared with that in less
- 20 severely septic patients and survivors (Nakae, H., et al., *Res. Commun. Chem. Pathol. Pharmacol.* 84:189-95, 1994; Nakae, et al., *Surg. Today* 26:225-29, 1996; Bengtson, A., et al., *Arch. Surg.* 123:645-649, 1988). The mechanisms by which C5a exerts its harmful effects during sepsis are yet to be investigated in greater detail, but recent data suggest the generation of C5a during sepsis significantly compromises innate immune functions of
- 25 blood neutrophils (Huber-Lang, M.S., et al., *J. Immunol.* 169:3223-31, 2002), their ability to express a respiratory burst, and their ability to generate cytokines (Riedemann, N.C., et al., *Immunity* 19:193-202, 2003). In addition, C5a generation during sepsis appears to have procoagulant effects (Laudes, I.J., et al., *Am. J. Pathol.* 160:1867-75, 2002). The complement-modulating protein CI INH has also shown efficacy in animal models of
- 30 sepsis and ARDS (Dickneite, G., *Behring Ins. Mitt.* 93:299-305, 1993).

The lectin pathway may also have a role in pathogenesis of sepsis. MBL has been shown to bind to a range of clinically important microorganisms including both Gram-negative and Gram-positive bacteria, and to activate the lectin pathway (Neth, O.,

et al., *Infect. Immun.* 68:688, 2000). Lipoteichoic acid (LTA) is increasingly regarded as the Gram-positive counterpart of LPS. It is a potent immunostimulant that induces cytokine release from mononuclear phagocytes and whole blood (Morath, S., et al., *J. Exp. Med.* 195:1635, 2002; Morath, S., et al., *Infect. Immun.* 70:938, 2002). Recently it was demonstrated that L-ficolin specifically binds to LTA isolated from numerous Gram-positive bacteria species, including *Staphylococcus aureus*, and activates the lectin pathway (Lynch, N.J., et al., *J. Immunol.* 172:1198-02, 2004). MBL also has been shown to bind to LTA from *Enterococcus spp* in which the polyglycerophosphate chain is substituted with glycosyl groups), but not to LTA from nine other species including *S. aureus* (Polotsky, V.Y., et al., *Infect. Immun.* 64:380, 1996).

An aspect of the invention thus provides a method for treating sepsis or a condition resulting from sepsis, by administering a composition comprising a therapeutically effective amount of a MASP-2 inhibitory agent in a pharmaceutical carrier to a subject suffering from sepsis or a condition resulting from sepsis including without limitation severe sepsis, septic shock, acute respiratory distress syndrome resulting from sepsis, and systemic inflammatory response syndrome. Related methods are provided for the treatment of other blood disorders, including hemorrhagic shock, hemolytic anemia, autoimmune thrombotic thrombocytopenic purpura (TTP), hemolytic uremic syndrome (HUS), atypical hemolytic uremic syndrome (aHUS), or other marrow/blood destructive conditions, by administering a composition comprising a therapeutically effective amount of a MASP-2 inhibitory agent in a pharmaceutical carrier to a subject suffering from such a condition. The MASP-2 inhibitory agent is administered to the subject systemically, such as by intra-arterial, intravenous, intramuscular, inhalational (particularly in the case of ARDS), subcutaneous or other parenteral administration, or potentially by oral administration for non-peptidergic agents. The MASP-2 inhibitory agent composition may be combined with one or more additional therapeutic agents to combat the sequelae of sepsis and/or shock. For advanced sepsis or shock or a distress condition resulting therefrom, the MASP-2 inhibitory composition may suitably be administered in a fast-acting dosage form, such as by intravenous or intra-arterial delivery of a bolus of a solution containing the MASP-2 inhibitory agent composition. Repeated administration may be carried out as determined by a physician until the condition has been resolved.

COAGULOPATHIES

Evidence has been developed for the role of the complement system in disseminated intravascular coagulation ("DIC"), such as DIC secondary to significant bodily trauma.

5 Previous studies have shown that C4^{-/-} mice are not protected from renal reperfusion injury. (Zhou, W., et al, "Predominant role for C5b-9 in renal ischemia/reperfusion injury," *J Clin Invest* 105:1363-1371 (2000)) In order to investigate whether C4^{-/-} mice may still be able to activate complement *via* either the classical or the lectin pathway, C3 turn-over in C4^{-/-} plasma was measured in assays specific for either
10 the classical, or the lectin pathway activation route. While no C3 cleavage could be observed when triggering activation *via* the classical, a highly efficient lectin pathway-dependent activation of C3 in C4 deficient serum was observed (FIGURE 30). It can be seen that C3b deposition on mannan and zymosan is severely compromised in MASP-2^{-/-} mice, even under experimental conditions, that according to many previously published
15 papers on alternative pathway activation, should be permissive for all three pathways. When using the same sera in wells coated with immunoglobulin complexes instead of mannan or zymosan, C3b deposition and Factor B cleavage are seen in MASP-2^{+/+} mouse sera and MASP-2^{-/-} sera, but not in C1q depleted sera. This indicates that alternate pathway activation is facilitated in MASP-2^{-/-} sera when the initial C3b is
20 provided via classical activity. FIGURE 30C depicts the surprising finding that C3 can efficiently be activated in a lectin pathway-dependent fashion in C4 deficient plasma.

This "C4 bypass" is abolished by the inhibition of lectin pathway-activation through preincubation of plasma with soluble mannan or mannose.

Aberrant, non-immune, activation of the complement system is potentially
25 hazardous to man and may also play an important role in hematological pathway activation, particularly in severe trauma situations wherein both inflammatory and hematological pathways are activated. In normal health, C3 conversion is <5% of the total plasma C3 protein. In rampant infection, including septicaemia and immune complex disease, C3 conversion re-establishes itself at about 30% with complement
30 levels frequently lower than normal, due to increased utilization and changes in pool distribution. Immediate C3 pathway activation of greater than 30% generally produces obvious clinical evidence of vasodilatation and of fluid loss to the tissues. Above 30% C3 conversion, the initiating mechanisms are predominantly non-immune and the resulting

clinical manifestations are harmful to the patient. Complement C5 levels in health and in controlled disease appear much more stable than C3. Significant decreases and or conversion of C5 levels are associated with the patient's response to abnormal polytrauma (e.g., road traffic accidents) and the likely development of shock lung syndromes. Thus, 5 any evidence of either complement C3 activation beyond 30% of the vascular pool or of any C5 involvement, or both, may be considered likely to be a harbinger of a harmful pathological change in the patient.

Both C3 and C5 liberate anaphylatoxins (C3a and C5a) that act on mast cells and basophils releasing vasodilatory chemicals. They set up chemotactic gradients to guide 10 polymorphonuclear cells (PMN) to the center of immunological disturbances (a beneficial response), but here they differ because C5a has a specific clumping (aggregating) effect on these phagocytic cells, preventing their random movement away from the reaction site. In normal control of infection, C3 activates C5. However, in polytrauma, C5 appears to be widely activated, generating C5a anaphylatoxins systemically. This uncontrolled 15 activity causes polymorphs to clump within the vascular system, and these clumps are then swept into the capillaries of the lungs, which they occlude and generate local damaging effects as a result of superoxide liberation. While not wishing to be limited by theory, the mechanism is probably important in the pathogenesis of acute respiratory distress syndrome (ARDS), although this view has recently been challenged. The C3a 20 anaphylatoxins *in vitro* can be shown to be potent platelet aggregators, but their involvement *in vivo* is less defined and the release of platelet substances and plasmin in wound repair may only secondarily involve complement C3. It is possible that prolonged elevation of C3 activation is necessary to generate DIC.

In addition to cellular and vascular effects of activated complement component 25 outlined above that could explain the link between trauma and DIC, emerging scientific discoveries have identified direct molecular links and functional cross-talk between complement and coagulation systems. Supporting data has been obtained from studies in C3 deficient mice. Because C3 is the shared component for each of the complement pathways, C3 deficient mice are predicted to lack all complement function. Surprisingly, 30 however, C3 deficient mice are perfectly capable of activating terminal complement components. (Huber-Lang, M., et al., "Generation of C5a in the absence of C3: a new complement activation pathway," *Nat. Med* 12:682-687 (2006)) In depth studies revealed that C3-independent activation of terminal complement components is mediated by

thrombin, the rate limiting enzyme of the coagulation cascade. (Huber et al., 2006) The molecular components mediating thrombin activation following initial complement activation remained elusive.

The present inventors have elucidated what is believed to be the molecular basis for cross-talk between complement and clotting cascades and identified MASP-2 as a central control point linking the two systems. Biochemical studies into the substrate specificity of MASP-2 have identified prothrombin as a possible substrate, in addition to the well known C2 and C4 complement proteins. MASP-2 specifically cleaves prothrombin at functionally relevant sites, generating thrombin, the rate limiting enzyme of the coagulation cascade. (Krarup, A., et al., "Simultaneous Activation of Complement and Coagulation by MBL-Associated Serine Protease 2," *PLoS. ONE*. 2:e623 (2007)) MASP-2-generated thrombin is capable of promoting fibrin deposition in a defined reconstituted *in vitro* system, demonstrating the functional relevance of MASP-2 cleavage. (Krarup et al., 2007) As discussed in the examples herein below, the inventors have further corroborated the physiological significance of this discovery by documenting thrombin activation in normal rodent serum following lectin pathway activation, and demonstrated that this process is blocked by neutralizing MASP-2 monoclonal antibodies.

MASP-2 may represent a central branch point in the lectin pathway, capable of promoting activation of both complement and coagulation systems. Because lectin pathway activation is a physiologic response to many types of traumatic injury, the present inventors believe that concurrent systemic inflammation (mediated by complement components) and disseminated coagulation (mediated via the clotting pathway) can be explained by the capacity of MASP-2 to activate both pathways. These findings clearly suggest a role for MASP-2 in DIC generation and therapeutic benefit of MASP-2 inhibition in treating or preventing DIC. MASP-2 may provide the molecular link between complement and coagulation system, and activation of the lectin pathway as it occurs in settings of trauma can directly initiate activation of the clotting system via the MASP-2-thrombin axis, providing a mechanistic link between trauma and DIC. In accordance with an aspect of the present invention, inhibition of MASP-2 would inhibit lectin pathway activation and reduce the generation of both anaphylatoxins C3a and C5a. It is believed that prolonged elevation of C3 activation is necessary to generate DIC.

Microcirculatory coagulation (blot clots in capillaries and small blood vessels) occurs in settings such as septic shock. A role of the lectin pathway in septic shock is established, as evidenced by the protected phenotype of MASP-2 (-/-) mouse models of sepsis, described in Example 17 and FIGURES 18 and 19. Furthermore, as demonstrated in Example 15 and FIGURES 16A and 16B, MASP-2 (-/-) mice are protected in the localized Schwartzman reaction model of disseminated intravascular coagulation (DIC), a model of localized coagulation in microvessels.

V. MASP-2 INHIBITORY AGENTS

In one aspect, the present invention provides methods of inhibiting MASP-2-dependent complement activation in a subject suffering from, or at risk for developing a thrombotic microangiopathy. MASP-2 inhibitory agents are administered in an amount effective to inhibit MASP-2-dependent complement activation in a living subject. In the practice of this aspect of the invention, representative MASP-2 inhibitory agents include: molecules that inhibit the biological activity of MASP-2 (such as small molecule inhibitors, anti-MASP-2 antibodies or blocking peptides which interact with MASP-2 or interfere with a protein-protein interaction), and molecules that decrease the expression of MASP-2 (such as MASP-2 antisense nucleic acid molecules, MASP-2 specific RNAi molecules and MASP-2 ribozymes), thereby preventing MASP-2 from activating the lectin complement pathway. The MASP-2 inhibitory agents can be used alone as a primary therapy or in combination with other therapeutics as an adjuvant therapy to enhance the therapeutic benefits of other medical treatments.

The inhibition of MASP-2-dependent complement activation is characterized by at least one of the following changes in a component of the complement system that occurs as a result of administration of a MASP-2 inhibitory agent in accordance with the methods of the invention: the inhibition of the generation or production of MASP-2-dependent complement activation system products C4b, C3a, C5a and/or C5b-9 (MAC) (measured, for example, as described in Example 2), the reduction of complement activation assessed in a hemolytic assay using unsensitized rabbit or guinea pig red blood cells (measured, for example as described in Example 33), the reduction of C4 cleavage and C4b deposition (measured, for example as described in Example 2), or the reduction of C3 cleavage and C3b deposition (measured, for example, as described in Example 2).

According to the present invention, MASP-2 inhibitory agents are utilized that are effective in inhibiting the MASP-2-dependent complement activation system. MASP-2 inhibitory agents useful in the practice of this aspect of the invention include, for example, anti-MASP-2 antibodies and fragments thereof, MASP-2 inhibitory peptides, 5 small molecules, MASP-2 soluble receptors and expression inhibitors. MASP-2 inhibitory agents may inhibit the MASP-2-dependent complement activation system by blocking the biological function of MASP-2. For example, an inhibitory agent may effectively block MASP-2 protein-to-protein interactions, interfere with MASP-2 dimerization or assembly, block Ca^{2+} binding, interfere with the MASP-2 serine protease 10 active site, or may reduce MASP-2 protein expression.

In some embodiments, the MASP-2 inhibitory agents selectively inhibit MASP-2 complement activation, leaving the C1q-dependent complement activation system functionally intact.

In one embodiment, a MASP-2 inhibitory agent useful in the methods of the 15 invention is a specific MASP-2 inhibitory agent that specifically binds to a polypeptide comprising SEQ ID NO:6 with an affinity of at least ten times greater than to other antigens in the complement system. In another embodiment, a MASP-2 inhibitory agent specifically binds to a polypeptide comprising SEQ ID NO:6 with a binding affinity of at least 100 times greater than to other antigens in the complement system. The binding 20 affinity of the MASP-2 inhibitory agent can be determined using a suitable binding assay.

The MASP-2 polypeptide exhibits a molecular structure similar to MASP-1, MASP-3, and C1r and C1s, the proteases of the C1 complement system. The cDNA molecule set forth in SEQ ID NO:4 encodes a representative example of MASP-2 (consisting of the amino acid sequence set forth in SEQ ID NO:5) and provides the 25 human MASP-2 polypeptide with a leader sequence (aa 1-15) that is cleaved after secretion, resulting in the mature form of human MASP-2 (SEQ ID NO:6). As shown in FIGURE 2, the human *MASP 2* gene encompasses twelve exons. The human MASP-2 cDNA is encoded by exons B, C, D, F, G, H, I, J, K AND L. An alternative splice results in a 20 kDa protein termed MBL-associated protein 19 ("MAp19", also referred to as 30 "sMAP") (SEQ ID NO:2), encoded by (SEQ ID NO:1) arising from exons B, C, D and E as shown in FIGURE 2. The cDNA molecule set forth in SEQ ID NO:50 encodes the murine MASP-2 (consisting of the amino acid sequence set forth in SEQ ID NO:51) and provides the murine MASP-2 polypeptide with a leader sequence that is cleaved after

secretion, resulting in the mature form of murine MASP-2 (SEQ ID NO:52). The cDNA molecule set forth in SEQ ID NO:53 encodes the rat MASP-2 (consisting of the amino acid sequence set forth in SEQ ID NO:54) and provides the rat MASP-2 polypeptide with a leader sequence that is cleaved after secretion, resulting in the mature form of rat MASP-2 (SEQ ID NO:55).

Those skilled in the art will recognize that the sequences disclosed in SEQ ID NO:4, SEQ ID NO:50 and SEQ ID NO:53 represent single alleles of human, murine and rat MASP-2 respectively, and that allelic variation and alternative splicing are expected to occur. Allelic variants of the nucleotide sequences shown in SEQ ID NO:4, SEQ ID NO:50 and SEQ ID NO:53, including those containing silent mutations and those in which mutations result in amino acid sequence changes, are within the scope of the present invention. Allelic variants of the MASP-2 sequence can be cloned by probing cDNA or genomic libraries from different individuals according to standard procedures.

The domains of the human MASP-2 protein (SEQ ID NO:6) are shown in FIGURE 1 and 2A and include an N-terminal C1r/C1s/sea urchin Vegf/bone morphogenic protein (CUBI) domain (aa 1-121 of SEQ ID NO:6), an epidermal growth factor-like domain (aa 122-166), a second CUBI domain (aa 167-293), as well as a tandem of complement control protein domains and a serine protease domain. Alternative splicing of the *MASP 2* gene results in MAP19 shown in FIGURE 1. MAP19 is a nonenzymatic protein containing the N-terminal CUB1-EGF region of MASP-2 with four additional residues (EQSL) derived from exon E as shown in FIGURE 1.

Several proteins have been shown to bind to, or interact with MASP-2 through protein-to-protein interactions. For example, MASP-2 is known to bind to, and form Ca^{2+} dependent complexes with, the lectin proteins MBL, H-ficolin and L-ficolin. Each MASP-2/lectin complex has been shown to activate complement through the MASP-2-dependent cleavage of proteins C4 and C2 (Ikeda, K., et al., *J. Biol. Chem.* 262:7451-7454, 1987; Matsushita, M., et al., *J. Exp. Med.* 176:1497-2284, 2000; Matsushita, M., et al., *J. Immunol.* 168:3502-3506, 2002). Studies have shown that the CUB1-EGF domains of MASP-2 are essential for the association of MASP-2 with MBL (Thielens, N.M., et al., *J. Immunol.* 166:5068, 2001). It has also been shown that the CUB1EGFCUBII domains mediate dimerization of MASP-2, which is required for formation of an active MBL complex (Wallis, R., et al., *J. Biol. Chem.* 275:30962-30969, 2000). Therefore, MASP-2 inhibitory agents can be identified that bind to or interfere

with MASP-2 target regions known to be important for MASP-2-dependent complement activation.

ANTI-MASP-2 ANTIBODIES

In some embodiments of this aspect of the invention, the MASP-2 inhibitory agent comprises an anti-MASP-2 antibody that inhibits the MASP-2-dependent complement activation system. The anti-MASP-2 antibodies useful in this aspect of the invention include polyclonal, monoclonal or recombinant antibodies derived from any antibody producing mammal and may be multispecific, chimeric, humanized, anti-idiotypic, and antibody fragments. Antibody fragments include Fab, Fab', F(ab)₂, F(ab')₂, Fv fragments, scFv fragments and single-chain antibodies as further described herein.

Several anti-MASP-2 antibodies have been described in the literature, some of which are listed below in TABLE 1. These previously described anti-MASP-2 antibodies can be screened for the ability to inhibit the MASP-2-dependent complement activation system using the assays described herein. For example, anti rat MASP-2 Fab2 antibodies have been identified that block MASP-2 dependent complement activation, as described in more detail in Examples 10 and 11 herein. Once an anti-MASP-2 antibody is identified that functions as a MASP-2 inhibitory agent, it can be used to produce anti-idiotypic antibodies and used to identify other MASP-2 binding molecules as further described below.

TABLE 1: MASP-2 SPECIFIC ANTIBODIES FROM THE LITERATURE

ANTIGEN	ANTIBODY TYPE	REFERENCE
Recombinant MASP-2	Rat Polyclonal	Peterson, S.V., et al., <i>Mol. Immunol.</i> 37:803-811, 2000
Recombinant human CCP1/2-SP fragment (MoAb 8B5)	Rat MoAb (subclass IgG1)	Moller-Kristensen, M., et al., <i>J. of Immunol. Methods</i> 282:159-167, 2003
Recombinant human MAp19 (MoAb 6G12) (cross reacts with MASP-2)	Rat MoAb (subclass IgG1)	Moller-Kristensen, M., et al., <i>J. of Immunol. Methods</i> 282:159-167, 2003
hMASP-2	Mouse MoAb (S/P) Mouse MoAb (N-term)	Peterson, S.V., et al., <i>Mol. Immunol.</i> 35:409, April 1998

ANTIGEN	ANTIBODY TYPE	REFERENCE
hMASP-2 (CCP1-CCP2-SP domain)	rat MoAb: Nimoab101, produced by hybridoma cell line 03050904 (ECACC)	WO 2004/106384
hMASP-2 (full length-his tagged)	murine MoAbs: NimoAb104, produced by hybridoma cell line M0545YM035 (DSMZ) NimoAb108, produced by hybridoma cell line M0545YM029 (DSMZ) NimoAb109 produced by hybridoma cell line M0545YM046 (DSMZ) NimoAb110 produced by hybridoma cell line M0545YM048 (DSMZ)	WO 2004/106384

ANTI-MASP-2 ANTIBODIES WITH REDUCED EFFECTOR FUNCTION

In some embodiments of this aspect of the invention, the anti-MASP-2 antibodies have reduced effector function in order to reduce inflammation that may arise from the activation of the classical complement pathway. The ability of IgG molecules to trigger the classical complement pathway has been shown to reside within the Fc portion of the molecule (Duncan, A.R., et al., *Nature* 332:738-740 1988). IgG molecules in which the Fc portion of the molecule has been removed by enzymatic cleavage are devoid of this effector function (see Harlow, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York, 1988). Accordingly, antibodies with reduced effector function can be generated as the result of lacking the Fc portion of the molecule by having a genetically engineered Fc sequence that minimizes effector function, or being of either the human IgG₂ or IgG₄ isotype.

Antibodies with reduced effector function can be produced by standard molecular biological manipulation of the Fc portion of the IgG heavy chains as described in Example 9 herein and also described in Jolliffe et al., *Int'l Rev. Immunol.* 10:241-250, 1993, and Rodrigues et al., *J. Immunol.* 151:6954-6961, 1998. Antibodies with reduced effector function also include human IgG₂ and IgG₄ isotypes that have a reduced ability to activate complement and/or interact with Fc receptors (Ravetch, J.V., et al., *Annu. Rev.*

Immunol. 9:457-492, 1991; Isaacs, J.D., et al., *J. Immunol.* 148:3062-3071, 1992; van de Winkel, J.G., et al., *Immunol. Today* 14:215-221, 1993). Humanized or fully human antibodies specific to human MASP-2 comprised of IgG2 or IgG4 isotypes can be produced by one of several methods known to one of ordinary skilled in the art, as described in Vaughan, T.J., et al., *Nature Biotechnical* 16:535-539, 1998.

PRODUCTION OF ANTI-MASP-2 ANTIBODIES

Anti-MASP-2 antibodies can be produced using MASP-2 polypeptides (e.g., full length MASP-2) or using antigenic MASP-2 epitope-bearing peptides (e.g., a portion of the MASP-2 polypeptide). Immunogenic peptides may be as small as five amino acid residues. For example, the MASP-2 polypeptide including the entire amino acid sequence of SEQ ID NO:6 may be used to induce anti-MASP-2 antibodies useful in the method of the invention. Particular MASP-2 domains known to be involved in protein-protein interactions, such as the CUBI, and CUBIEGF domains, as well as the region encompassing the serine-protease active site, may be expressed as recombinant polypeptides as described in Example 3 and used as antigens. In addition, peptides comprising a portion of at least 6 amino acids of the MASP-2 polypeptide (SEQ ID NO:6) are also useful to induce MASP-2 antibodies. Additional examples of MASP-2 derived antigens useful to induce MASP-2 antibodies are provided below in TABLE 2. The MASP-2 peptides and polypeptides used to raise antibodies may be isolated as natural polypeptides, or recombinant or synthetic peptides and catalytically inactive recombinant polypeptides, such as MASP-2A, as further described in Examples 5-7. In some embodiments of this aspect of the invention, anti-MASP-2 antibodies are obtained using a transgenic mouse strain as described in Examples 8 and 9 and further described below.

Antigens useful for producing anti-MASP-2 antibodies also include fusion polypeptides, such as fusions of MASP-2 or a portion thereof with an immunoglobulin polypeptide or with maltose-binding protein. The polypeptide immunogen may be a full-length molecule or a portion thereof. If the polypeptide portion is hapten-like, such portion may be advantageously joined or linked to a macromolecular carrier (such as keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA) or tetanus toxoid) for immunization.

TABLE 2: MASP-2 DERIVED ANTIGENS

SEQ ID NO:	Amino Acid Sequence
SEQ ID NO:6	Human MASP-2 protein
SEQ ID NO:51	Murine MASP-2 protein
SEQ ID NO:8	CUBI domain of human MASP-2 (aa 1-121 of SEQ ID NO:6)
SEQ ID NO:9	CUBIEGF domains of human MASP-2 (aa 1-166 of SEQ ID NO:6)
SEQ ID NO:10	CUBIEGFCUBII domains of human MASP-2 (aa 1-293 of SEQ ID NO:6)
SEQ ID NO:11	EGF domain of human MASP-2 (aa 122-166 of SEQ ID NO:6)
SEQ ID NO:12	Serine-Protease domain of human MASP-2 (aa 429-671 of SEQ ID NO:6)
SEQ ID NO:13 GKDSCRGDAGGALVFL	Serine-Protease inactivated mutant form (aa 610-625 of SEQ ID NO:6 with mutated Ser 618)
SEQ ID NO:14 TPLGPKWPEPVFGRL	Human CUBI peptide
SEQ ID NO:15: TAPPGYRLRLYFTHFDLEL SHLCEYDFVKLSSGAKVL ATLCGQ	Human CUBI peptide
SEQ ID NO:16: TFRSDYSN	MBL binding region in human CUBI domain
SEQ ID NO:17: FYSLGSSLDITFRSDYSNEK PFTGF	MBL binding region in human CUBI domain
SEQ ID NO:18 IDECQVAPG	EGF peptide
SEQ ID NO:19 ANMLCAGLESGGKDSCRG DSGGALV	Peptide from serine-protease active site

POLYCLONAL ANTIBODIES

Polyclonal antibodies against MASP-2 can be prepared by immunizing an animal with MASP-2 polypeptide or an immunogenic portion thereof using methods well known to those of ordinary skill in the art. See, for example, Green et al., "Production of Polyclonal Antisera," in *Immunochemical Protocols* (Manson, ed.), page 105, and as further described in Example 6. The immunogenicity of a MASP-2 polypeptide can be

increased through the use of an adjuvant, including mineral gels, such as aluminum hydroxide or Freund's adjuvant (complete or incomplete), surface active substances such as lysolecithin, pluronic polyols, polyanions, oil emulsions, keyhole limpet hemocyanin and dinitrophenol. Polyclonal antibodies are typically raised in animals such as horses, cows, dogs, chicken, rats, mice, rabbits, guinea pigs, goats, or sheep. Alternatively, an anti-MASP-2 antibody useful in the present invention may also be derived from a subhuman primate. General techniques for raising diagnostically and therapeutically useful antibodies in baboons may be found, for example, in Goldenberg et al., International Patent Publication No. WO 91/11465, and in Losman, M.J., et al., *Int. J. Cancer* 46:310, 1990. Sera containing immunologically active antibodies are then produced from the blood of such immunized animals using standard procedures well known in the art.

MONOCLONAL ANTIBODIES

In some embodiments, the MASP-2 inhibitory agent is an anti-MASP-2 monoclonal antibody. Anti-MASP-2 monoclonal antibodies are highly specific, being directed against a single MASP-2 epitope. As used herein, the modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogenous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. Monoclonal antibodies can be obtained using any technique that provides for the production of antibody molecules by continuous cell lines in culture, such as the hybridoma method described by Kohler, G., et al., *Nature* 256:495, 1975, or they may be made by recombinant DNA methods (see, e.g., U.S. Patent No. 4,816,567 to Cabilly). Monoclonal antibodies may also be isolated from phage antibody libraries using the techniques described in Clackson, T., et al., *Nature* 352:624-628, 1991, and Marks, J.D., et al., *J. Mol. Biol.* 222:581-597, 1991. Such antibodies can be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof.

For example, monoclonal antibodies can be obtained by injecting a suitable mammal (e.g., a BALB/c mouse) with a composition comprising a MASP-2 polypeptide or portion thereof. After a predetermined period of time, splenocytes are removed from the mouse and suspended in a cell culture medium. The splenocytes are then fused with an immortal cell line to form a hybridoma. The formed hybridomas are grown in cell culture and screened for their ability to produce a monoclonal antibody against MASP-2.

An example further describing the production of anti-MASP-2 monoclonal antibodies is provided in Example 7. (See also *Current Protocols in Immunology*, Vol. 1., John Wiley & Sons, pages 2.5.1-2.6.7, 1991.)

Human monoclonal antibodies may be obtained through the use of transgenic mice that have been engineered to produce specific human antibodies in response to antigenic challenge. In this technique, elements of the human immunoglobulin heavy and light chain locus are introduced into strains of mice derived from embryonic stem cell lines that contain targeted disruptions of the endogenous immunoglobulin heavy chain and light chain loci. The transgenic mice can synthesize human antibodies specific for human antigens, such as the MASP-2 antigens described herein, and the mice can be used to produce human MASP-2 antibody-secreting hybridomas by fusing B-cells from such animals to suitable myeloma cell lines using conventional Kohler-Milstein technology as further described in Example 7. Transgenic mice with a human immunoglobulin genome are commercially available (e.g., from Abgenix, Inc., Fremont, CA, and Medarex, Inc., Annandale, N.J.). Methods for obtaining human antibodies from transgenic mice are described, for example, by Green, L.L., et al., *Nature Genet.* 7:13, 1994; Lonberg, N., et al., *Nature* 368:856, 1994; and Taylor, L.D., et al., *Int. Immun.* 6:579, 1994.

Monoclonal antibodies can be isolated and purified from hybridoma cultures by a variety of well-established techniques. Such isolation techniques include affinity chromatography with Protein-A Sepharose, size-exclusion chromatography, and ion-exchange chromatography (see, for example, Coligan at pages 2.7.1-2.7.12 and pages 2.9.1-2.9.3; Baines et al., "Purification of Immunoglobulin G (IgG)," in *Methods in Molecular Biology*, The Humana Press, Inc., Vol. 10, pages 79-104, 1992).

Once produced, polyclonal, monoclonal or phage-derived antibodies are first tested for specific MASP-2 binding. A variety of assays known to those skilled in the art may be utilized to detect antibodies which specifically bind to MASP-2. Exemplary assays include Western blot or immunoprecipitation analysis by standard methods (e.g., as described in Ausubel et al.), immunoelectrophoresis, enzyme-linked immuno-sorbent assays, dot blots, inhibition or competition assays and sandwich assays (as described in Harlow and Land, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 1988). Once antibodies are identified that specifically bind to MASP-2, the anti-MASP-2 antibodies are tested for the ability to function as a MASP-2 inhibitory agent in one of several assays such as, for example, a lectin-specific C4 cleavage assay

(described in Example 2), a C3b deposition assay (described in Example 2) or a C4b deposition assay (described in Example 2).

The affinity of anti-MASP-2 monoclonal antibodies can be readily determined by one of ordinary skill in the art (see, e.g., Scatchard, A., *NY Acad. Sci.* 51:660-672, 1949).

- 5 In one embodiment, the anti-MASP-2 monoclonal antibodies useful for the methods of the invention bind to MASP-2 with a binding affinity of <100 nM, preferably <10 nM and most preferably <2 nM. In some embodiments, a MASP-2 inhibitory monoclonal antibody useful in the methods of the invention is a MASP-2 inhibitory monoclonal antibody, or antigen binding fragment thereof, comprising (I) (a) a heavy-chain variable
- 10 region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-102 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of
- 15 SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least
- 20 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70.

CHIMERIC/HUMANIZED ANTIBODIES

- 25 Monoclonal antibodies useful in the method of the invention include chimeric antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from
- 30 another species or belonging to another antibody class or subclass, as well as fragments of such antibodies (U.S. Patent No. 4,816,567, to Cabilly; and Morrison, S.L., et al., *Proc. Nat'l Acad. Sci. USA* 81:6851-6855, 1984).

One form of a chimeric antibody useful in the invention is a humanized monoclonal anti-MASP-2 antibody. Humanized forms of non-human (e.g., murine) antibodies are chimeric antibodies, which contain minimal sequence derived from non-human immunoglobulin. Humanized monoclonal antibodies are produced by
5 transferring the non-human (e.g., mouse) complementarity determining regions (CDR), from the heavy and light variable chains of the mouse immunoglobulin into a human variable domain. Typically, residues of human antibodies are then substituted in the framework regions of the non-human counterparts. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor
10 antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the Fv framework regions are those of a human immunoglobulin sequence. The humanized
15 antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones, P.T., et al., *Nature* 321:522-525, 1986; Reichmann, L., et al., *Nature* 332:323-329, 1988; and Presta, *Curr. Op. Struct. Biol.* 2:593-596, 1992.

The humanized antibodies useful in the invention include human monoclonal
20 antibodies including at least a MASP-2 binding CDR3 region. In addition, the Fc portions may be replaced so as to produce IgA or IgM as well as human IgG antibodies. Such humanized antibodies will have particular clinical utility because they will specifically recognize human MASP-2 but will not evoke an immune response in humans against the antibody itself. Consequently, they are better suited for *in vivo* administration
25 in humans, especially when repeated or long-term administration is necessary.

An example of the generation of a humanized anti-MASP-2 antibody from a murine anti-MASP-2 monoclonal antibody is provided herein in Example 6. Techniques for producing humanized monoclonal antibodies are also described, for example, by Jones, P.T., et al., *Nature* 321:522, 1986; Carter, P., et al., *Proc. Nat'l. Acad. Sci. USA* 89:4285, 1992; Sandhu, J.S., *Crit. Rev. Biotech.* 12:437, 1992; Singer, I.I., et al., *J. Immun.* 150:2844, 1993; Sudhir (ed.), *Antibody Engineering Protocols*, Humana Press, Inc., 1995; Kelley, "Engineering Therapeutic Antibodies," in *Protein Engineering: Principles and Practice*, Cleland et al. (eds.), John Wiley & Sons, Inc., pages 399-434,
30

1996; and by U.S. Patent No. 5,693,762, to Queen, 1997. In addition, there are commercial entities that will synthesize humanized antibodies from specific murine antibody regions, such as Protein Design Labs (Mountain View, CA).

5 RECOMBINANT ANTIBODIES

Anti-MASP-2 antibodies can also be made using recombinant methods. For example, human antibodies can be made using human immunoglobulin expression libraries (available for example, from Stratagene, Corp., La Jolla, CA) to produce fragments of human antibodies (V_H , V_L , Fv, Fd, Fab or $F(ab')_2$). These fragments are
10 then used to construct whole human antibodies using techniques similar to those for producing chimeric antibodies.

ANTI-IDIOTYPE ANTIBODIES

Once anti-MASP-2 antibodies are identified with the desired inhibitory activity,
15 these antibodies can be used to generate anti-idiotypic antibodies that resemble a portion of MASP-2 using techniques that are well known in the art. See, e.g., Greenspan, N.S., et al., *FASEB J.* 7:437, 1993. For example, antibodies that bind to MASP-2 and competitively inhibit a MASP-2 protein interaction required for complement activation can be used to generate anti-idiotypes that resemble the MBL binding site on MASP-2
20 protein and therefore bind and neutralize a binding ligand of MASP-2 such as, for example, MBL.

IMMUNOGLOBULIN FRAGMENTS

The MASP-2 inhibitory agents useful in the method of the invention encompass not only intact immunoglobulin molecules but also the well known fragments including
25 Fab, Fab', $F(ab)_2$, $F(ab')_2$ and Fv fragments, scFv fragments, diabodies, linear antibodies, single-chain antibody molecules and multispecific antibodies formed from antibody fragments.

It is well known in the art that only a small portion of an antibody molecule, the paratope, is involved in the binding of the antibody to its epitope (see, e.g., Clark, W.R.,
30 *The Experimental Foundations of Modern Immunology*, Wiley & Sons, Inc., NY, 1986). The pFc' and Fc regions of the antibody are effectors of the classical complement pathway, but are not involved in antigen binding. An antibody from which the pFc' region has been enzymatically cleaved, or which has been produced without the pFc'

region, is designated an F(ab')₂ fragment and retains both of the antigen binding sites of an intact antibody. An isolated F(ab')₂ fragment is referred to as a bivalent monoclonal fragment because of its two antigen binding sites. Similarly, an antibody from which the Fc region has been enzymatically cleaved, or which has been produced without the Fc region, is designated a Fab fragment, and retains one of the antigen binding sites of an intact antibody molecule.

Antibody fragments can be obtained by proteolytic hydrolysis, such as by pepsin or papain digestion of whole antibodies by conventional methods. For example, antibody fragments can be produced by enzymatic cleavage of antibodies with pepsin to provide a 5S fragment denoted F(ab')₂. This fragment can be further cleaved using a thiol reducing agent to produce 3.5S Fab' monovalent fragments. Optionally, the cleavage reaction can be performed using a blocking group for the sulfhydryl groups that result from cleavage of disulfide linkages. As an alternative, an enzymatic cleavage using pepsin produces two monovalent Fab fragments and an Fc fragment directly. These methods are described, for example, U.S. Patent No. 4,331,647 to Goldenberg; Nisonoff, A., et al., *Arch. Biochem. Biophys.* 89:230, 1960; Porter, R.R., *Biochem. J.* 73:119, 1959; Edelman, et al., in *Methods in Enzymology* 1:422, Academic Press, 1967; and by Coligan at pages 2.8.1-2.8.10 and 2.10.-2.10.4.

In some embodiments, the use of antibody fragments lacking the Fc region are preferred to avoid activation of the classical complement pathway which is initiated upon binding Fc to the Fcγ receptor. There are several methods by which one can produce a MoAb that avoids Fcγ receptor interactions. For example, the Fc region of a monoclonal antibody can be removed chemically using partial digestion by proteolytic enzymes (such as ficin digestion), thereby generating, for example, antigen-binding antibody fragments such as Fab or F(ab)₂ fragments (Mariani, M., et al., *Mol. Immunol.* 28:69-71, 1991). Alternatively, the human γ4 IgG isotype, which does not bind Fcγ receptors, can be used during construction of a humanized antibody as described herein. Antibodies, single chain antibodies and antigen-binding domains that lack the Fc domain can also be engineered using recombinant techniques described herein.

30

SINGLE-CHAIN ANTIBODY FRAGMENTS

Alternatively, one can create single peptide chain binding molecules specific for MASP-2 in which the heavy and light chain Fv regions are connected. The Fv fragments may be connected by a peptide linker to form a single-chain antigen binding protein (scFv). These single-chain antigen binding proteins are prepared by constructing a structural gene comprising DNA sequences encoding the V_H and V_L domains which are connected by an oligonucleotide. The structural gene is inserted into an expression vector, which is subsequently introduced into a host cell, such as *E. coli*. The recombinant host cells synthesize a single polypeptide chain with a linker peptide bridging the two V domains. Methods for producing scFvs are described for example, by Whitlow, et al., "Methods: A Companion to Methods in Enzymology" 2:97, 1991; Bird, et al., *Science* 242:423, 1988; U.S. Patent No. 4,946,778, to Ladner; Pack, P., et al., *Bio/Technology* 11:1271, 1993.

As an illustrative example, a MASP-2 specific scFv can be obtained by exposing lymphocytes to MASP-2 polypeptide *in vitro* and selecting antibody display libraries in phage or similar vectors (for example, through the use of immobilized or labeled MASP-2 protein or peptide). Genes encoding polypeptides having potential MASP-2 polypeptide binding domains can be obtained by screening random peptide libraries displayed on phage or on bacteria such as *E. coli*. These random peptide display libraries can be used to screen for peptides which interact with MASP-2. Techniques for creating and screening such random peptide display libraries are well known in the art (U.S. Patent No. 5,223,409, to Lardner; U.S. Patent No. 4,946,778, to Ladner; U.S. Patent No. 5,403,484, to Lardner; U.S. Patent No. 5,571,698, to Lardner; and Kay et al., *Phage Display of Peptides and Proteins* Academic Press, Inc., 1996) and random peptide display libraries and kits for screening such libraries are available commercially, for instance from CLONTECH Laboratories, Inc. (Palo Alto, Calif.), Invitrogen Inc. (San Diego, Calif.), New England Biolabs, Inc. (Beverly, Mass.), and Pharmacia LKB Biotechnology Inc. (Piscataway, N.J.).

Another form of an anti-MASP-2 antibody fragment useful in this aspect of the invention is a peptide coding for a single complementarity-determining region (CDR) that binds to an epitope on a MASP-2 antigen and inhibits MASP-2-dependent complement activation. CDR peptides ("minimal recognition units") can be obtained by constructing genes encoding the CDR of an antibody of interest. Such genes are prepared, for

example, by using the polymerase chain reaction to synthesize the variable region from RNA of antibody-producing cells (see, for example, Larrick et al., *Methods: A Companion to Methods in Enzymology* 2:106, 1991; Courtenay-Luck, "Genetic Manipulation of Monoclonal Antibodies," in *Monoclonal Antibodies: Production, Engineering and Clinical Application*, Ritter et al. (eds.), page 166, Cambridge University Press, 1995; and Ward et al., "Genetic Manipulation and Expression of Antibodies," in *Monoclonal Antibodies: Principles and Applications*, Birch et al. (eds.), page 137, Wiley-Liss, Inc., 1995).

The MASP-2 antibodies described herein are administered to a subject in need thereof to inhibit MASP-2-dependent complement activation. In some embodiments, the MASP-2 inhibitory agent is a high-affinity human or humanized monoclonal anti-MASP-2 antibody with reduced effector function.

PEPTIDE INHIBITORS

In some embodiments of this aspect of the invention, the MASP-2 inhibitory agent comprises isolated MASP-2 peptide inhibitors, including isolated natural peptide inhibitors and synthetic peptide inhibitors that inhibit the MASP-2-dependent complement activation system. As used herein, the term "isolated MASP-2 peptide inhibitors" refers to peptides that inhibit MASP-2 dependent complement activation by binding to, competing with MASP-2 for binding to another recognition molecule (e.g., MBL, H-ficolin, M-ficolin, or L-ficolin) in the lectin pathway, and/or directly interacting with MASP-2 to inhibit MASP-2-dependent complement activation that are substantially pure and are essentially free of other substances with which they may be found in nature to an extent practical and appropriate for their intended use.

Peptide inhibitors have been used successfully *in vivo* to interfere with protein-protein interactions and catalytic sites. For example, peptide inhibitors to adhesion molecules structurally related to LFA-1 have recently been approved for clinical use in coagulopathies (Ohman, E.M., et al., *European Heart J.* 16:50-55, 1995). Short linear peptides (<30 amino acids) have been described that prevent or interfere with integrin-dependent adhesion (Murayama, O., et al., *J. Biochem.* 120:445-51, 1996). Longer peptides, ranging in length from 25 to 200 amino acid residues, have also been used successfully to block integrin-dependent adhesion (Zhang, L., et al., *J. Biol. Chem.* 271(47):29953-57, 1996). In general, longer peptide inhibitors have higher

affinities and/or slower off-rates than short peptides and may therefore be more potent inhibitors. Cyclic peptide inhibitors have also been shown to be effective inhibitors of integrins *in vivo* for the treatment of human inflammatory disease (Jackson, D.Y., et al., *J. Med. Chem.* 40:3359-68, 1997). One method of producing cyclic peptides involves the
5 synthesis of peptides in which the terminal amino acids of the peptide are cysteines, thereby allowing the peptide to exist in a cyclic form by disulfide bonding between the terminal amino acids, which has been shown to improve affinity and half-life *in vivo* for the treatment of hematopoietic neoplasms (e.g., U.S. Patent No. 6,649,592, to Larson).

10 SYNTHETIC MASP-2 PEPTIDE INHIBITORS

MAASP-2 inhibitory peptides useful in the methods of this aspect of the invention are exemplified by amino acid sequences that mimic the target regions important for MASP-2 function. The inhibitory peptides useful in the practice of the methods of the invention range in size from about 5 amino acids to about 300 amino acids. TABLE 3
15 provides a list of exemplary inhibitory peptides that may be useful in the practice of this aspect of the present invention. A candidate MASP-2 inhibitory peptide may be tested for the ability to function as a MASP-2 inhibitory agent in one of several assays including, for example, a lectin specific C4 cleavage assay (described in Example 2), and a C3b deposition assay (described in Example 2).

20 In some embodiments, the MASP-2 inhibitory peptides are derived from MASP-2 polypeptides and are selected from the full length mature MASP-2 protein (SEQ ID NO:6), or from a particular domain of the MASP-2 protein such as, for example, the CUBI domain (SEQ ID NO:8), the CUBIEGF domain (SEQ ID NO:9), the EGF domain (SEQ ID NO:11), and the serine protease domain (SEQ ID NO:12). As previously
25 described, the CUBEGFCUBII regions have been shown to be required for dimerization and binding with MBL (Thielens et al., *supra*). In particular, the peptide sequence TFRSDYN (SEQ ID NO:16) in the CUBI domain of MASP-2 has been shown to be involved in binding to MBL in a study that identified a human carrying a homozygous mutation at Asp105 to Gly105, resulting in the loss of MASP-2 from the MBL complex
30 (Stengaard-Pedersen, K., et al., *New England J. Med.* 349:554-560, 2003).

In some embodiments, MASP-2 inhibitory peptides are derived from the lectin proteins that bind to MASP-2 and are involved in the lectin complement pathway. Several different lectins have been identified that are involved in this pathway, including

mannan-binding lectin (MBL), L-ficolin, M-ficolin and H-ficolin. (Ikeda, K., et al., *J. Biol. Chem.* 262:7451-7454, 1987; Matsushita, M., et al., *J. Exp. Med.* 176:1497-2284, 2000; Matsushita, M., et al., *J. Immunol.* 168:3502-3506, 2002). These lectins are present in serum as oligomers of homotrimeric subunits, each having N-terminal collagen-like
5 fibers with carbohydrate recognition domains. These different lectins have been shown to bind to MASP-2, and the lectin/MASP-2 complex activates complement through cleavage of proteins C4 and C2. H-ficolin has an amino-terminal region of 24 amino acids, a collagen-like domain with 11 Gly-Xaa-Yaa repeats, a neck domain of 12 amino acids, and a fibrinogen-like domain of 207 amino acids (Matsushita, M., et al.,
10 *J. Immunol.* 168:3502-3506, 2002). H-ficolin binds to GlcNAc and agglutinates human erythrocytes coated with LPS derived from *S. typhimurium*, *S. minnesota* and *E. coli*. H-ficolin has been shown to be associated with MASP-2 and MAp19 and activates the lectin pathway. *Id.* L-ficolin/P35 also binds to GlcNAc and has been shown to be associated with MASP-2 and MAp19 in human serum and this complex has been shown
15 to activate the lectin pathway (Matsushita, M., et al., *J. Immunol.* 164:2281, 2000). Accordingly, MASP-2 inhibitory peptides useful in the present invention may comprise a region of at least 5 amino acids selected from the MBL protein (SEQ ID NO:21), the H-ficolin protein (Genbank accession number NM_173452), the M-ficolin protein (Genbank accession number O00602) and the L-ficolin protein (Genbank accession
20 number NM_015838).

More specifically, scientists have identified the MASP-2 binding site on MBL to be within the 12 Gly-X-Y triplets "GKD GRD GTK GEK GEP GQG LRG LQG POG KLG POG NOG PSG SOG PKG QKG DOG KS" (SEQ ID NO:26) that lie between the hinge and the neck in the C-terminal portion of the collagen-like domain of MBP
25 (Wallis, R., et al., *J. Biol. Chem.* 279:14065, 2004). This MASP-2 binding site region is also highly conserved in human H-ficolin and human L-ficolin. A consensus binding site has been described that is present in all three lectin proteins comprising the amino acid sequence "OGK-X-GP" (SEQ ID NO:22) where the letter "O" represents hydroxyproline and the letter "X" is a hydrophobic residue (Wallis et al., 2004, *supra*). Accordingly, in
30 some embodiments, MASP-2 inhibitory peptides useful in this aspect of the invention are at least 6 amino acids in length and comprise SEQ ID NO:22. Peptides derived from MBL that include the amino acid sequence "GLR GLQ GPO GKL GPO G" (SEQ ID NO:24) have been shown to bind MASP-2 *in vitro* (Wallis, et al., 2004, *supra*). To

enhance binding to MASP-2, peptides can be synthesized that are flanked by two GPO triplets at each end ("GPO GPO GLR GLQ GPO GKL GPO GGP OGP O" SEQ ID NO:25) to enhance the formation of triple helices as found in the native MBL protein (as further described in Wallis, R., et al., *J. Biol. Chem.* 279:14065, 2004).

- 5 MASP-2 inhibitory peptides may also be derived from human H-ficolin that include the sequence "GAO GSO GEK GAO GPQ GPO GPO GKM GPK GEO GDO" (SEQ ID NO:27) from the consensus MASP-2 binding region in H-ficolin. Also included are peptides derived from human L-ficolin that include the sequence "GCO GLO GAO GDK GEA GTN GKR GER GPO GPO GKA GPO GPN GAO GEO" (SEQ ID NO:28)
- 10 from the consensus MASP-2 binding region in L-ficolin.

MASP-2 inhibitory peptides may also be derived from the C4 cleavage site such as "LQRALEILPNRVTIKANRPFLVFI" (SEQ ID NO:29) which is the C4 cleavage site linked to the C-terminal portion of antithrombin III (Glover, G.I., et al., *Mol. Immunol.* 25:1261 (1988)).

15

TABLE 3: EXEMPLARY MASP-2 INHIBITORY PEPTIDES

SEQ ID NO	Source
SEQ ID NO:6	Human MASP-2 protein
SEQ ID NO:8	CUBI domain of MASP-2 (aa 1-121 of SEQ ID NO:6)
SEQ ID NO:9	CUBIEGF domains of MASP-2 (aa 1-166 of SEQ ID NO:6)
SEQ ID NO:10	CUBIEGFCUBII domains of MASP-2 (aa 1-293 of SEQ ID NO:6)
SEQ ID NO:11	EGF domain of MASP-2 (aa 122-166)
SEQ ID NO:12	Serine-protease domain of MASP-2 (aa 429-671)
SEQ ID NO:16	MBL binding region in MASP-2
SEQ ID NO:3	Human MAP19
SEQ ID NO:21	Human MBL protein

SEQ ID NO	Source
SEQ ID NO:22 OGK-X-GP, Where "O" = hydroxyproline and "X" is a hydrophobic amino acid residue	Synthetic peptide Consensus binding site from Human MBL and Human ficolins
SEQ ID NO:23 OGKLG	Human MBL core binding site
SEQ ID NO:24 GLR GLQ GPO GKL GPO G	Human MBP Triplets 6-10- demonstrated binding to MASP-2
SEQ ID NO:25 GPOGPOGLRGLQGPO GKLGPOGGPOGPO	Human MBP Triplets with GPO added to enhance formation of triple helices
SEQ ID NO:26 GKDGRDGTKGEKGEP GQGLRGLQGPOGKLG POGNOGPSGSOGPKG QKGDGKS	Human MBP Triplets 1-17
SEQ ID NO:27 GAOGSOGKEKGAOGPQ GPOGPOGKMGPKGEO GDO	Human H-Ficolin (Hataka)
SEQ ID NO:28 GCOGLOGAOGDKGE AGTNGKRGERGPOGP OGKAGPOGPNGAOGE O	Human L-Ficolin P35
SEQ ID NO:29 LQRALEILPNRVTIKA NRPFLVFI	Human C4 cleavage site

Note: The letter "O" represents hydroxyproline. The letter "X" is a hydrophobic residue.

- Peptides derived from the C4 cleavage site as well as other peptides that inhibit the MASP-2 serine protease site can be chemically modified so that they are irreversible protease inhibitors. For example, appropriate modifications may include, but are not necessarily limited to, halomethyl ketones (Br, Cl, I, F) at the C-terminus, Asp or Glu, or

appended to functional side chains; haloacetyl (or other α -haloacetyl) groups on amino groups or other functional side chains; epoxide or imine-containing groups on the amino or carboxy termini or on functional side chains; or imidate esters on the amino or carboxy termini or on functional side chains. Such modifications would afford the advantage of
5 permanently inhibiting the enzyme by covalent attachment of the peptide. This could result in lower effective doses and/or the need for less frequent administration of the peptide inhibitor.

In addition to the inhibitory peptides described above, MASP-2 inhibitory peptides useful in the method of the invention include peptides containing the
10 MASP-2-binding CDR3 region of anti-MASP-2 MoAb obtained as described herein. The sequence of the CDR regions for use in synthesizing the peptides may be determined by methods known in the art. The heavy chain variable region is a peptide that generally ranges from 100 to 150 amino acids in length. The light chain variable region is a peptide that generally ranges from 80 to 130 amino acids in length. The CDR sequences within
15 the heavy and light chain variable regions include only approximately 3-25 amino acid sequences that may be easily sequenced by one of ordinary skill in the art.

Those skilled in the art will recognize that substantially homologous variations of the MASP-2 inhibitory peptides described above will also exhibit MASP-2 inhibitory activity. Exemplary variations include, but are not necessarily limited to, peptides having
20 insertions, deletions, replacements, and/or additional amino acids on the carboxy-terminus or amino-terminus portions of the subject peptides and mixtures thereof. Accordingly, those homologous peptides having MASP-2 inhibitory activity are considered to be useful in the methods of this invention. The peptides described may also include duplicating motifs and other modifications with conservative substitutions.
25 Conservative variants are described elsewhere herein, and include the exchange of an amino acid for another of like charge, size or hydrophobicity and the like.

MASP-2 inhibitory peptides may be modified to increase solubility and/or to maximize the positive or negative charge in order to more closely resemble the segment in the intact protein. The derivative may or may not have the exact primary amino acid
30 structure of a peptide disclosed herein so long as the derivative functionally retains the desired property of MASP-2 inhibition. The modifications can include amino acid substitution with one of the commonly known twenty amino acids or with another amino acid, with a derivatized or substituted amino acid with ancillary desirable characteristics,

such as resistance to enzymatic degradation or with a D-amino acid or substitution with another molecule or compound, such as a carbohydrate, which mimics the natural conformation and function of the amino acid, amino acids or peptide; amino acid deletion; amino acid insertion with one of the commonly known twenty amino acids or with
5 another amino acid, with a derivatized or substituted amino acid with ancillary desirable characteristics, such as resistance to enzymatic degradation or with a D-amino acid or substitution with another molecule or compound, such as a carbohydrate, which mimics the natural conformation and function of the amino acid, amino acids or peptide; or substitution with another molecule or compound, such as a carbohydrate or nucleic acid
10 monomer, which mimics the natural conformation, charge distribution and function of the parent peptide. Peptides may also be modified by acetylation or amidation.

The synthesis of derivative inhibitory peptides can rely on known techniques of peptide biosynthesis, carbohydrate biosynthesis and the like. As a starting point, the artisan may rely on a suitable computer program to determine the conformation of a
15 peptide of interest. Once the conformation of peptide disclosed herein is known, then the artisan can determine in a rational design fashion what sort of substitutions can be made at one or more sites to fashion a derivative that retains the basic conformation and charge distribution of the parent peptide but which may possess characteristics which are not present or are enhanced over those found in the parent peptide. Once candidate
20 derivative molecules are identified, the derivatives can be tested to determine if they function as MASP-2 inhibitory agents using the assays described herein.

SCREENING FOR MASP-2 INHIBITORY PEPTIDES

One may also use molecular modeling and rational molecular design to generate
25 and screen for peptides that mimic the molecular structures of key binding regions of MASP-2 and inhibit the complement activities of MASP-2. The molecular structures used for modeling include the CDR regions of anti-MASP-2 monoclonal antibodies, as well as the target regions known to be important for MASP-2 function including the region required for dimerization, the region involved in binding to MBL, and the serine
30 protease active site as previously described. Methods for identifying peptides that bind to a particular target are well known in the art. For example, molecular imprinting may be used for the *de novo* construction of macromolecular structures such as peptides that bind to a particular molecule. See, for example, Shea, K.J., "Molecular Imprinting of

Synthetic Network Polymers: The De Novo synthesis of Macromolecular Binding and Catalytic Sites," *TRIP* 2(5) 1994.

As an illustrative example, one method of preparing mimics of MASP-2 binding peptides is as follows. Functional monomers of a known MASP-2 binding peptide or the binding region of an anti-MASP-2 antibody that exhibits MASP-2 inhibition (the template) are polymerized. The template is then removed, followed by polymerization of a second class of monomers in the void left by the template, to provide a new molecule that exhibits one or more desired properties that are similar to the template. In addition to preparing peptides in this manner, other MASP-2 binding molecules that are MASP-2 inhibitory agents such as polysaccharides, nucleosides, drugs, nucleoproteins, lipoproteins, carbohydrates, glycoproteins, steroid, lipids and other biologically active materials can also be prepared. This method is useful for designing a wide variety of biological mimics that are more stable than their natural counterparts because they are typically prepared by free radical polymerization of function monomers, resulting in a compound with a nonbiodegradable backbone.

PEPTIDE SYNTHESIS

The MASP-2 inhibitory peptides can be prepared using techniques well known in the art, such as the solid-phase synthetic technique initially described by Merrifield, in *J. Amer. Chem. Soc.* 85:2149-2154, 1963. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Foster City, Calif.) in accordance with the instructions provided by the manufacturer. Other techniques may be found, for example, in Bodanszky, M., et al., *Peptide Synthesis*, second edition, John Wiley & Sons, 1976, as well as in other reference works known to those skilled in the art.

The peptides can also be prepared using standard genetic engineering techniques known to those skilled in the art. For example, the peptide can be produced enzymatically by inserting nucleic acid encoding the peptide into an expression vector, expressing the DNA, and translating the DNA into the peptide in the presence of the required amino acids. The peptide is then purified using chromatographic or electrophoretic techniques, or by means of a carrier protein that can be fused to, and subsequently cleaved from, the peptide by inserting into the expression vector in phase with the peptide encoding sequence a nucleic acid sequence encoding the carrier protein. The fusion protein-peptide may be isolated using chromatographic, electrophoretic or

immunological techniques (such as binding to a resin via an antibody to the carrier protein). The peptide can be cleaved using chemical methodology or enzymatically, as by, for example, hydrolases.

5 The MASP-2 inhibitory peptides that are useful in the method of the invention can also be produced in recombinant host cells following conventional techniques. To express a MASP-2 inhibitory peptide encoding sequence, a nucleic acid molecule encoding the peptide must be operably linked to regulatory sequences that control transcriptional expression in an expression vector and then introduced into a host cell. In addition to transcriptional regulatory sequences, such as promoters and enhancers, expression vectors
10 can include translational regulatory sequences and a marker gene, which are suitable for selection of cells that carry the expression vector.

Nucleic acid molecules that encode a MASP-2 inhibitory peptide can be synthesized with "gene machines" using protocols such as the phosphoramidite method. If chemically synthesized double-stranded DNA is required for an application such as the
15 synthesis of a gene or a gene fragment, then each complementary strand is made separately. The production of short genes (60 to 80 base pairs) is technically straightforward and can be accomplished by synthesizing the complementary strands and then annealing them. For the production of longer genes, synthetic genes (double-stranded) are assembled in modular form from single-stranded fragments that are
20 from 20 to 100 nucleotides in length. For reviews on polynucleotide synthesis, see, for example, Glick and Pasternak, *"Molecular Biotechnology, Principles and Applications of Recombinant DNA"*, ASM Press, 1994; Itakura, K., et al., *Annu. Rev. Biochem.* 53:323, 1984; and Climie, S., et al., *Proc. Nat'l Acad. Sci. USA* 87:633, 1990.

SMALL MOLECULE INHIBITORS

25 In some embodiments, MASP-2 inhibitory agents are small molecule inhibitors including natural and synthetic substances that have a low molecular weight, such as for example, peptides, peptidomimetics and nonpeptide inhibitors (including oligonucleotides and organic compounds). Small molecule inhibitors of MASP-2 can be generated based on the molecular structure of the variable regions of the anti-MASP-2
30 antibodies.

Small molecule inhibitors may also be designed and generated based on the MASP-2 crystal structure using computational drug design (Kuntz I.D., et al., *Science* 257:1078, 1992). The crystal structure of rat MASP-2 has been described

(Feinberg, H., et al., *EMBO J.* 22:2348-2359, 2003). Using the method described by Kuntz et al., the MASP-2 crystal structure coordinates are used as an input for a computer program such as DOCK, which outputs a list of small molecule structures that are expected to bind to MASP-2. Use of such computer programs is well known to one of skill in the art. For example, the crystal structure of the HIV-1 protease inhibitor was used to identify unique nonpeptide ligands that are HIV-1 protease inhibitors by evaluating the fit of compounds found in the Cambridge Crystallographic database to the binding site of the enzyme using the program DOCK (Kuntz, I.D., et al., *J. Mol. Biol.* 161:269-288, 1982; DesJarlais, R.L., et al., *PNAS* 87:6644-6648, 1990).

10 The list of small molecule structures that are identified by a computational method as potential MASP-2 inhibitors are screened using a MASP-2 binding assay such as described in Example 10. The small molecules that are found to bind to MASP-2 are then assayed in a functional assay such as described in Example 2 to determine if they inhibit MASP-2-dependent complement activation.

15

MASP-2 SOLUBLE RECEPTORS

Other suitable MASP-2 inhibitory agents are believed to include MASP-2 soluble receptors, which may be produced using techniques known to those of ordinary skill in the art.

20

EXPRESSION INHIBITORS OF MASP-2

In another embodiment of this aspect of the invention, the MASP-2 inhibitory agent is a MASP-2 expression inhibitor capable of inhibiting MASP-2-dependent complement activation. In the practice of this aspect of the invention, representative MASP-2 expression inhibitors include MASP-2 antisense nucleic acid molecules (such as antisense mRNA, antisense DNA or antisense oligonucleotides), MASP-2 ribozymes and MASP-2 RNAi molecules.

25

Anti-sense RNA and DNA molecules act to directly block the translation of MASP-2 mRNA by hybridizing to MASP-2 mRNA and preventing translation of MASP-2 protein. An antisense nucleic acid molecule may be constructed in a number of different ways provided that it is capable of interfering with the expression of MASP-2. For example, an antisense nucleic acid molecule can be constructed by inverting the coding region (or a portion thereof) of MASP-2 cDNA (SEQ ID NO:4) relative to its normal orientation for transcription to allow for the transcription of its complement.

30

The antisense nucleic acid molecule is usually substantially identical to at least a portion of the target gene or genes. The nucleic acid, however, need not be perfectly identical to inhibit expression. Generally, higher homology can be used to compensate for the use of a shorter antisense nucleic acid molecule. The minimal percent identity is typically greater than about 65%, but a higher percent identity may exert a more effective repression of expression of the endogenous sequence. Substantially greater percent identity of more than about 80% typically is preferred, though about 95% to absolute identity is typically most preferred.

The antisense nucleic acid molecule need not have the same intron or exon pattern as the target gene, and non-coding segments of the target gene may be equally effective in achieving antisense suppression of target gene expression as coding segments. A DNA sequence of at least about 8 or so nucleotides may be used as the antisense nucleic acid molecule, although a longer sequence is preferable. In the present invention, a representative example of a useful inhibitory agent of MASP-2 is an antisense MASP-2 nucleic acid molecule which is at least ninety percent identical to the complement of the MASP-2 cDNA consisting of the nucleic acid sequence set forth in SEQ ID NO:4. The nucleic acid sequence set forth in SEQ ID NO:4 encodes the MASP-2 protein consisting of the amino acid sequence set forth in SEQ ID NO:5.

The targeting of antisense oligonucleotides to bind MASP-2 mRNA is another mechanism that may be used to reduce the level of MASP-2 protein synthesis. For example, the synthesis of polygalacturonase and the muscarine type 2 acetylcholine receptor is inhibited by antisense oligonucleotides directed to their respective mRNA sequences (U.S. Patent No. 5,739,119, to Cheng, and U.S. Patent No. 5,759,829, to Shewmaker). Furthermore, examples of antisense inhibition have been demonstrated with the nuclear protein cyclin, the multiple drug resistance gene (MDG1), ICAM-1, E-selectin, STK-1, striatal GABA_A receptor and human EGF (see, e.g., U.S. Patent No. 5,801,154, to Baracchini; U.S. Patent No. 5,789,573, to Baker; U.S. Patent No. 5,718,709, to Considine; and U.S. Patent No. 5,610,288, to Reubenstein).

A system has been described that allows one of ordinary skill to determine which oligonucleotides are useful in the invention, which involves probing for suitable sites in the target mRNA using Rnase H cleavage as an indicator for accessibility of sequences within the transcripts. Scherr, M., et al., *Nucleic Acids Res.* 26:5079-5085, 1998; Lloyd, et al., *Nucleic Acids Res.* 29:3665-3673, 2001. A mixture of antisense

oligonucleotides that are complementary to certain regions of the MASP-2 transcript is added to cell extracts expressing MASP-2, such as hepatocytes, and hybridized in order to create an RNaseH vulnerable site. This method can be combined with computer-assisted sequence selection that can predict optimal sequence selection for antisense compositions based upon their relative ability to form dimers, hairpins, or other secondary structures that would reduce or prohibit specific binding to the target mRNA in a host cell. These secondary structure analysis and target site selection considerations may be performed using the OLIGO primer analysis software (Rychlik, I., 1997) and the BLASTN 2.0.5 algorithm software (Altschul, S.F., et al., *Nucl. Acids Res.* 25:3389-3402, 1997). The antisense compounds directed towards the target sequence preferably comprise from about 8 to about 50 nucleotides in length. Antisense oligonucleotides comprising from about 9 to about 35 or so nucleotides are particularly preferred. The inventors contemplate all oligonucleotide compositions in the range of 9 to 35 nucleotides (i.e., those of 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35 or so bases in length) are highly preferred for the practice of antisense oligonucleotide-based methods of the invention. Highly preferred target regions of the MASP-2 mRNA are those that are at or near the AUG translation initiation codon, and those sequences that are substantially complementary to 5' regions of the mRNA, e.g., between the -10 and +10 regions of the *MA SP-2* gene nucleotide sequence (SEQ ID NO:4). Exemplary MASP-2 expression inhibitors are provided in TABLE 4.

TABLE 4: EXEMPLARY EXPRESSION INHIBITORS OF MASP-2

SEQ ID NO:30 (nucleotides 22-680 of SEQ ID NO:4)	Nucleic acid sequence of MASP-2 cDNA (SEQ ID NO:4) encoding CUBIEGF
SEQ ID NO:31 5'CGGGCACACCATGAGGCTGCTG ACCCTCCTGGGC3	Nucleotides 12-45 of SEQ ID NO:4 including the MASP-2 translation start site (sense)
SEQ ID NO:32 5'GACATTACCTTCCGCTCCGACTC CAACGAGAAG3'	Nucleotides 361-396 of SEQ ID NO:4 encoding a region comprising the MASP-2 MBL binding site (sense)
SEQ ID NO:33 5'AGCAGCCCTGAATACCCACGGCC GTATCCCAA3'	Nucleotides 610-642 of SEQ ID NO:4 encoding a region comprising the CUBII domain

As noted above, the term "oligonucleotide" as used herein refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term also covers those oligonucleobases composed of naturally occurring nucleotides, sugars and covalent internucleoside (backbone) linkages as well as
5 oligonucleotides having non-naturally occurring modifications. These modifications allow one to introduce certain desirable properties that are not offered through naturally occurring oligonucleotides, such as reduced toxic properties, increased stability against nuclease degradation and enhanced cellular uptake. In illustrative embodiments, the antisense compounds of the invention differ from native DNA by the modification of the
10 phosphodiester backbone to extend the life of the antisense oligonucleotide in which the phosphate substituents are replaced by phosphorothioates. Likewise, one or both ends of the oligonucleotide may be substituted by one or more acridine derivatives that intercalate between adjacent basepairs within a strand of nucleic acid.

Another alternative to antisense is the use of "RNA interference" (RNAi).
15 Double-stranded RNAs (dsRNAs) can provoke gene silencing in mammals *in vivo*. The natural function of RNAi and co-suppression appears to be protection of the genome against invasion by mobile genetic elements such as retrotransposons and viruses that produce aberrant RNA or dsRNA in the host cell when they become active (see, e.g., Jensen, J., et al., *Nat. Genet.* 21:209-12, 1999). The double-stranded RNA molecule may
20 be prepared by synthesizing two RNA strands capable of forming a double-stranded RNA molecule, each having a length from about 19 to 25 (e.g., 19-23 nucleotides). For example, a dsRNA molecule useful in the methods of the invention may comprise the RNA corresponding to a sequence and its complement listed in TABLE 4. Preferably, at least one strand of RNA has a 3' overhang from 1-5 nucleotides. The synthesized RNA
25 strands are combined under conditions that form a double-stranded molecule. The RNA sequence may comprise at least an 8 nucleotide portion of SEQ ID NO:4 with a total length of 25 nucleotides or less. The design of siRNA sequences for a given target is within the ordinary skill of one in the art. Commercial services are available that design siRNA sequence and guarantee at least 70% knockdown of expression (Qiagen, Valencia,
30 Calif).

The dsRNA may be administered as a pharmaceutical composition and carried out by known methods, wherein a nucleic acid is introduced into a desired target cell. Commonly used gene transfer methods include calcium phosphate, DEAE-dextran,

electroporation, microinjection and viral methods. Such methods are taught in Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., 1993.

Ribozymes can also be utilized to decrease the amount and/or biological activity of MASP-2, such as ribozymes that target MASP-2 mRNA. Ribozymes are catalytic
5 RNA molecules that can cleave nucleic acid molecules having a sequence that is completely or partially homologous to the sequence of the ribozyme. It is possible to design ribozyme transgenes that encode RNA ribozymes that specifically pair with a target RNA and cleave the phosphodiester backbone at a specific location, thereby functionally inactivating the target RNA. In carrying out this cleavage, the ribozyme is
10 not itself altered, and is thus capable of recycling and cleaving other molecules. The inclusion of ribozyme sequences within antisense RNAs confers RNA-cleaving activity upon them, thereby increasing the activity of the antisense constructs.

Ribozymes useful in the practice of the invention typically comprise a hybridizing region of at least about nine nucleotides, which is complementary in nucleotide sequence
15 to at least part of the target MASP-2 mRNA, and a catalytic region that is adapted to cleave the target MASP-2 mRNA (see generally, EPA No. 0 321 201; WO88/04300; Haseloff, J., et al., *Nature* 334:585-591, 1988; Fedor, M.J., et al., *Proc. Natl. Acad. Sci. USA* 87:1668-1672, 1990; Cech, T.R., et al., *Ann. Rev. Biochem.* 55:599-629, 1986).

Ribozymes can either be targeted directly to cells in the form of RNA
20 oligonucleotides incorporating ribozyme sequences, or introduced into the cell as an expression vector encoding the desired ribozymal RNA. Ribozymes may be used and applied in much the same way as described for antisense polynucleotides.

Anti-sense RNA and DNA, ribozymes and RNAi molecules useful in the methods of the invention may be prepared by any method known in the art for the synthesis of
25 DNA and RNA molecules. These include techniques for chemically synthesizing oligodeoxyribonucleotides and oligoribonucleotides well known in the art, such as for example solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences may be incorporated into a wide variety
30 of vectors that incorporate suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines.

Various well known modifications of the DNA molecules may be introduced as a means of increasing stability and half-life. Useful modifications include, but are not limited to, the addition of flanking sequences of ribonucleotides or deoxyribonucleotides to the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the oligodeoxyribonucleotide backbone.

VI. PHARMACEUTICAL COMPOSITIONS AND DELIVERY METHODS DOSING

In another aspect, the invention provides compositions for inhibiting the adverse effects of MASP-2-dependent complement activation in a subject suffering from a disease or condition as disclosed herein, comprising administering to the subject a composition comprising a therapeutically effective amount of a MASP-2 inhibitory agent and a pharmaceutically acceptable carrier. The MASP-2 inhibitory agents can be administered to a subject in need thereof, at therapeutically effective doses to treat or ameliorate conditions associated with MASP-2-dependent complement activation. A therapeutically effective dose refers to the amount of the MASP-2 inhibitory agent sufficient to result in amelioration of symptoms associated with the disease or condition.

Toxicity and therapeutic efficacy of MASP-2 inhibitory agents can be determined by standard pharmaceutical procedures employing experimental animal models, such as the murine MASP-2 $-/-$ mouse model expressing the human MASP-2 transgene described in Example 1. Using such animal models, the NOAEL (no observed adverse effect level) and the MED (the minimally effective dose) can be determined using standard methods. The dose ratio between NOAEL and MED effects is the therapeutic ratio, which is expressed as the ratio NOAEL/MED. MASP-2 inhibitory agents that exhibit large therapeutic ratios or indices are most preferred. The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosages for use in humans. The dosage of the MASP-2 inhibitory agent preferably lies within a range of circulating concentrations that include the MED with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized.

For any compound formulation, the therapeutically effective dose can be estimated using animal models. For example, a dose may be formulated in an animal model to achieve a circulating plasma concentration range that includes the MED.

Quantitative levels of the MASP-2 inhibitory agent in plasma may also be measured, for example, by high performance liquid chromatography.

In addition to toxicity studies, effective dosage may also be estimated based on the amount of MASP-2 protein present in a living subject and the binding affinity of the MASP-2 inhibitory agent. It has been shown that MASP-2 levels in normal human subjects is present in serum in low levels in the range of 500 ng/ml, and MASP-2 levels in a particular subject can be determined using a quantitative assay for MASP-2 described in Moller-Kristensen M., et al., *J. Immunol. Methods* 282:159-167, 2003.

Generally, the dosage of administered compositions comprising MASP-2 inhibitory agents varies depending on such factors as the subject's age, weight, height, sex, general medical condition, and previous medical history. As an illustration, MASP-2 inhibitory agents, such as anti-MASP-2 antibodies, can be administered in dosage ranges from about 0.010 to 10.0 mg/kg, preferably 0.010 to 1.0 mg/kg, more preferably 0.010 to 0.1 mg/kg of the subject body weight. In some embodiments the composition comprises a combination of anti-MASP-2 antibodies and MASP-2 inhibitory peptides.

Therapeutic efficacy of MASP-2 inhibitory compositions and methods of the present invention in a given subject, and appropriate dosages, can be determined in accordance with complement assays well known to those of skill in the art. Complement generates numerous specific products. During the last decade, sensitive and specific assays have been developed and are available commercially for most of these activation products, including the small activation fragments C3a, C4a, and C5a and the large activation fragments iC3b, C4d, Bb, and sC5b-9. Most of these assays utilize monoclonal antibodies that react with new antigens (neoantigens) exposed on the fragment, but not on the native proteins from which they are formed, making these assays very simple and specific. Most rely on ELISA technology, although radioimmunoassay is still sometimes used for C3a and C5a. These latter assays measure both the unprocessed fragments and their 'desArg' fragments, which are the major forms found in the circulation. Unprocessed fragments and C5a_{desArg} are rapidly cleared by binding to cell surface receptors and are hence present in very low concentrations, whereas C3a_{desArg} does not bind to cells and accumulates in plasma. Measurement of C3a provides a sensitive, pathway-independent indicator of complement activation. Alternative pathway activation can be assessed by measuring the Bb fragment. Detection of the fluid-phase product of membrane attack pathway activation, sC5b-9, provides evidence that complement is

being activated to completion. Because both the lectin and classical pathways generate the same activation products, C4a and C4d, measurement of these two fragments does not provide any information about which of these two pathways has generated the activation products.

5 The inhibition of MASP-2-dependent complement activation is characterized by at least one of the following changes in a component of the complement system that occurs as a result of administration of a MASP-2 inhibitory agent in accordance with the methods of the invention: the inhibition of the generation or production of MASP-2-dependent complement activation system products C4b, C3a, C5a and/or C5b-9
10 (MAC) (measured, for example, as described in measured, for example, as described in Example 2, the reduction of C4 cleavage and C4b deposition (measured, for example as described in Example 10), or the reduction of C3 cleavage and C3b deposition (measured, for example, as described in Example 10).

15 ADDITIONAL AGENTS

 The compositions and methods comprising MASP-2 inhibitory agents may optionally comprise one or more additional therapeutic agents, which may augment the activity of the MASP-2 inhibitory agent or that provide related therapeutic functions in an additive or synergistic fashion. For example, in the context of treating a subject suffering
20 from TTP, wherein the subject is positive for an inhibitor of ADAM-TS13, one or more MASP-2 inhibitory agents may be administered in combination (including co-administration) with one or more immunosuppressive agents. Suitable immunosuppressive agents include: corticosteroids, rituxan, cyclosporine, and the like. In the context of treating a subject suffering from, or at risk for developing, HUS or
25 aHUS, one or more MASP-2 inhibitory agents may be administered in combination (including co-administration) with a suitable antibiotic. In the context of treating a subject suffering from, or at risk for developing aHUS, one or more MASP-2 inhibitory agents may be administered in combination (including co-administration) with other complement inhibitory agents such as eculizumab (Soliris), TT-30, antibody to factor B,
30 or other agents that inhibit terminal complement components or alternative pathway amplification.

 The inclusion and selection of additional agent(s) will be determined to achieve a desired therapeutic result. In some embodiments, the MASP-2 inhibitory agent may be

administered in combination with one or more anti-inflammatory and/or analgesic agents. Suitable anti-inflammatory and/or analgesic agents include: serotonin receptor antagonists; serotonin receptor agonists; histamine receptor antagonists; bradykinin receptor antagonists; kallikrein inhibitors; tachykinin receptor antagonists, including neurokinin₁ and neurokinin₂ receptor subtype antagonists; calcitonin gene-related peptide (CGRP) receptor antagonists; interleukin receptor antagonists; inhibitors of enzymes active in the synthetic pathway for arachidonic acid metabolites, including phospholipase inhibitors, including PLA₂ isoform inhibitors and PLC γ isoform inhibitors, cyclooxygenase (COX) inhibitors (which may be either COX-1, COX-2, or nonselective COX-1 and -2 inhibitors), lipoxygenase inhibitors; prostanoid receptor antagonists including eicosanoid EP-1 and EP-4 receptor subtype antagonists and thromboxane receptor subtype antagonists; leukotriene receptor antagonists including leukotriene B₄ receptor subtype antagonists and leukotriene D₄ receptor subtype antagonists; opioid receptor agonists, including μ -opioid, δ -opioid, and κ -opioid receptor subtype agonists; purinoceptor agonists and antagonists including P_{2X} receptor antagonists and P_{2Y} receptor agonists; adenosine triphosphate (ATP)-sensitive potassium channel openers; MAP kinase inhibitors; nicotinic acetylcholine inhibitors; and alpha adrenergic receptor agonists (including alpha-1, alpha-2, and nonselective alpha-1 and 2 agonists).

The MASP-2 inhibitory agents of the present invention may also be administered in combination with one or more other complement inhibitors, such as an inhibitor of C5. To date, Eculizumab (Solaris®), an antibody against C5, is the only complement-targeting drug that has been approved for human use. However some pharmacological agents have been shown to block complement *in vivo*. K76COOH and nafamstat mesilate are two agents that have shown some effectiveness in animal models of transplantation (Miyagawa, S., et al., *Transplant Proc.* 24:483-484, 1992). Low molecular weight heparins have also been shown to be effective in regulating complement activity (Edens, R.E., et al., *Complement Today*, pp. 96-120, Basel: Karger, 1993). It is believed that these small molecule inhibitors may be useful as agents to use in combination with the MASP-2 inhibitory agents of the present invention.

Other naturally occurring complement inhibitors may be useful in combination with the MASP-2 inhibitory agents of the present invention. Biological inhibitors of

complement include soluble complement factor 1 (sCR1). This is a naturally-occurring inhibitor that can be found on the outer membrane of human cells. Other membrane inhibitors include DAF, MCP, and CD59. Recombinant forms have been tested for their anti-complement activity *in vitro* and *in vivo*. sCR1 has been shown to be effective in
5 xenotransplantation, wherein the complement system (both alternative and classical) provides the trigger for a hyperactive rejection syndrome within minutes of perfusing blood through the newly transplanted organ (Platt, J.L., et al., *Immunol. Today* 11:450-6, 1990; Marino, I.R., et al., *Transplant Proc.* 1071:6, 1990; Johnstone, P.S., et al., *Transplantation* 54:573-6, 1992). The use of sCR1 protects and extends the survival time
10 of the transplanted organ, implicating the complement pathway in the pathogenesis of organ survival (Leventhal, J.R., et al., *Transplantation* 55:857-66, 1993; Pruitt, S.K., et al., *Transplantation* 57:363-70, 1994).

Suitable additional complement inhibitors for use in combination with the compositions of the present invention also include, by way of example, MoAbs such as
15 an anti-C5 antibody (e.g., eculizumab) being developed by Alexion Pharmaceuticals, Inc., New Haven, Connecticut, and anti-properdin MoAbs.

PHARMACEUTICAL CARRIERS AND DELIVERY VEHICLES

In general, the MASP-2 inhibitory agent compositions of the present invention,
20 combined with any other selected therapeutic agents, are suitably contained in a pharmaceutically acceptable carrier. The carrier is non-toxic, biocompatible and is selected so as not to detrimentally affect the biological activity of the MASP-2 inhibitory agent (and any other therapeutic agents combined therewith). Exemplary pharmaceutically acceptable carriers for peptides are described in U.S. Patent
25 No. 5,211,657 to Yamada. The anti-MASP-2 antibodies and inhibitory peptides useful in the invention may be formulated into preparations in solid, semi-solid, gel, liquid or gaseous forms such as tablets, capsules, powders, granules, ointments, solutions, depositories, inhalants and injections allowing for oral, parenteral or surgical administration. The invention also contemplates local administration of the compositions
30 by coating medical devices and the like.

Suitable carriers for parenteral delivery via injectable, infusion or irrigation and topical delivery include distilled water, physiological phosphate-buffered saline, normal or lactated Ringer's solutions, dextrose solution, Hank's solution, or propanediol. In

addition, sterile, fixed oils may be employed as a solvent or suspending medium. For this purpose any biocompatible oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables. The carrier and agent may be compounded as a liquid, suspension, 5 polymerizable or non-polymerizable gel, paste or salve.

The carrier may also comprise a delivery vehicle to sustain (i.e., extend, delay or regulate) the delivery of the agent(s) or to enhance the delivery, uptake, stability or pharmacokinetics of the therapeutic agent(s). Such a delivery vehicle may include, by way of non-limiting example, microparticles, microspheres, nanospheres or nanoparticles 10 composed of proteins, liposomes, carbohydrates, synthetic organic compounds, inorganic compounds, polymeric or copolymeric hydrogels and polymeric micelles. Suitable hydrogel and micelle delivery systems include the PEO:PHB:PEO copolymers and copolymer/cyclodextrin complexes disclosed in WO 2004/009664 A2 and the PEO and PEO/cyclodextrin complexes disclosed in U.S. Patent Application Publication 15 No. 2002/0019369 A1. Such hydrogels may be injected locally at the site of intended action, or subcutaneously or intramuscularly to form a sustained release depot.

For intra-articular delivery, the MASP-2 inhibitory agent may be carried in above-described liquid or gel carriers that are injectable, above-described sustained-release delivery vehicles that are injectable, or a hyaluronic acid or hyaluronic 20 acid derivative.

For oral administration of non-peptidergic agents, the MASP-2 inhibitory agent may be carried in an inert filler or diluent such as sucrose, cornstarch, or cellulose.

For topical administration, the MASP-2 inhibitory agent may be carried in ointment, lotion, cream, gel, drop, suppository, spray, liquid or powder, or in gel or 25 microcapsular delivery systems via a transdermal patch.

Various nasal and pulmonary delivery systems, including aerosols, metered-dose inhalers, dry powder inhalers, and nebulizers, are being developed and may suitably be adapted for delivery of the present invention in an aerosol, inhalant, or nebulized delivery vehicle, respectively.

30 For intrathecal (IT) or intracerebroventricular (ICV) delivery, appropriately sterile delivery systems (e.g., liquids; gels, suspensions, etc.) can be used to administer the present invention.

The compositions of the present invention may also include biocompatible excipients, such as dispersing or wetting agents, suspending agents, diluents, buffers, penetration enhancers, emulsifiers, binders, thickeners, flavouring agents (for oral administration).

5

PHARMACEUTICAL CARRIERS FOR ANTIBODIES AND PEPTIDES

More specifically with respect to anti-MASP-2 antibodies and inhibitory peptides, exemplary formulations can be parenterally administered as injectable dosages of a solution or suspension of the compound in a physiologically acceptable diluent with a pharmaceutical carrier that can be a sterile liquid such as water, oils, saline, glycerol or ethanol. Additionally, auxiliary substances such as wetting or emulsifying agents, surfactants, pH buffering substances and the like can be present in compositions comprising anti-MASP-2 antibodies and inhibitory peptides. Additional components of pharmaceutical compositions include petroleum (such as of animal, vegetable or synthetic origin), for example, soybean oil and mineral oil. In general, glycols such as propylene glycol or polyethylene glycol are preferred liquid carriers for injectable solutions.

The anti-MASP-2 antibodies and inhibitory peptides can also be administered in the form of a depot injection or implant preparation that can be formulated in such a manner as to permit a sustained or pulsatile release of the active agents.

PHARMACEUTICALLY ACCEPTABLE CARRIERS FOR EXPRESSION INHIBITORS

More specifically with respect to expression inhibitors useful in the methods of the invention, compositions are provided that comprise an expression inhibitor as described above and a pharmaceutically acceptable carrier or diluent. The composition may further comprise a colloidal dispersion system.

Pharmaceutical compositions that include expression inhibitors may include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids. The preparation of such compositions typically involves combining the expression inhibitor with one or more of the following: buffers, antioxidants, low molecular weight polypeptides, proteins, amino acids, carbohydrates including glucose, sucrose or dextrans, chelating agents such as EDTA, glutathione and other stabilizers and excipients. Neutral

buffered saline or saline mixed with non-specific serum albumin are examples of suitable diluents.

5 In some embodiments, the compositions may be prepared and formulated as emulsions which are typically heterogeneous systems of one liquid dispersed in another in the form of droplets (see, Idson, in *Pharmaceutical Dosage Forms*, Vol. 1, Rieger and Banker (eds.), Marcek Dekker, Inc., N.Y., 1988). Examples of naturally occurring emulsifiers used in emulsion formulations include acacia, beeswax, lanolin, lecithin and phosphatides.

10 In one embodiment, compositions including nucleic acids can be formulated as microemulsions. A microemulsion, as used herein refers to a system of water, oil, and amphiphile, which is a single optically isotropic and thermodynamically stable liquid solution (see Rosoff in *Pharmaceutical Dosage Forms*, Vol. 1). The method of the invention may also use liposomes for the transfer and delivery of antisense oligonucleotides to the desired site.

15 Pharmaceutical compositions and formulations of expression inhibitors for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, as well as aqueous, powder or oily bases and thickeners and the like may be used.

20 MODES OF ADMINISTRATION

The pharmaceutical compositions comprising MASP-2 inhibitory agents may be administered in a number of ways depending on whether a local or systemic mode of administration is most appropriate for the condition being treated. Additionally, as
25 described herein above with respect to extracorporeal reperfusion procedures, MASP-2 inhibitory agents can be administered via introduction of the compositions of the present invention to recirculating blood or plasma. Further, the compositions of the present invention can be delivered by coating or incorporating the compositions on or into an implantable medical device.

30 SYSTEMIC DELIVERY

As used herein, the terms "systemic delivery" and "systemic administration" are intended to include but are not limited to oral and parenteral routes including intramuscular (IM), subcutaneous, intravenous (IV), intra-arterial, inhalational,

sublingual, buccal, topical, transdermal, nasal, rectal, vaginal and other routes of administration that effectively result in dispersement of the delivered agent to a single or multiple sites of intended therapeutic action. Preferred routes of systemic delivery for the present compositions include intravenous, intramuscular, subcutaneous and inhalational.

5 It will be appreciated that the exact systemic administration route for selected agents utilized in particular compositions of the present invention will be determined in part to account for the agent's susceptibility to metabolic transformation pathways associated with a given route of administration. For example, peptidergic agents may be most suitably administered by routes other than oral.

10 MASP-2 inhibitory antibodies and polypeptides can be delivered into a subject in need thereof by any suitable means. Methods of delivery of MASP-2 antibodies and polypeptides include administration by oral, pulmonary, parenteral (e.g., intramuscular, intraperitoneal, intravenous (IV) or subcutaneous injection), inhalation (such as via a fine powder formulation), transdermal, nasal, vaginal, rectal, or sublingual routes of

15 administration, and can be formulated in dosage forms appropriate for each route of administration.

By way of representative example, MASP-2 inhibitory antibodies and peptides can be introduced into a living body by application to a bodily membrane capable of absorbing the polypeptides, for example the nasal, gastrointestinal and rectal membranes.

20 The polypeptides are typically applied to the absorptive membrane in conjunction with a permeation enhancer. (See, e.g., Lee, V.H.L., *Crit. Rev. Ther. Drug Carrier Sys.* 5:69, 1988; Lee, V.H.L., *J. Controlled Release* 13:213, 1990; Lee, V.H.L., Ed., *Peptide and Protein Drug Delivery*, Marcel Dekker, New York (1991); DeBoer, A.G., et al., *J. Controlled Release* 13:241, 1990.) For example, STDHF is a synthetic derivative of

25 fusidic acid, a steroidal surfactant that is similar in structure to the bile salts, and has been used as a permeation enhancer for nasal delivery. (Lee, W.A., *Biopharm.* 22, Nov./Dec. 1990.)

The MASP-2 inhibitory antibodies and polypeptides may be introduced in association with another molecule, such as a lipid, to protect the polypeptides from

30 enzymatic degradation. For example, the covalent attachment of polymers, especially polyethylene glycol (PEG), has been used to protect certain proteins from enzymatic hydrolysis in the body and thus prolong half-life (Fuertges, F., et al., *J. Controlled Release* 11:139, 1990). Many polymer systems have been reported for protein delivery

(Bae, Y.H., et al., *J. Controlled Release* 9:271, 1989; Hori, R., et al., *Pharm. Res.* 6:813, 1989; Yamakawa, I., et al., *J. Pharm. Sci.* 79:505, 1990; Yoshihiro, I., et al., *J. Controlled Release* 10:195, 1989; Asano, M., et al., *J. Controlled Release* 9:111, 1989; Rosenblatt, J., et al., *J. Controlled Release* 9:195, 1989; Makino, K., *J. Controlled Release* 12:235, 5 1990; Takakura, Y., et al., *J. Pharm. Sci.* 78:117, 1989; Takakura, Y., et al., *J. Pharm. Sci.* 78:219, 1989).

Recently, liposomes have been developed with improved serum stability and circulation half-times (see, e.g., U.S. Patent No. 5,741,516, to Webb). Furthermore, various methods of liposome and liposome-like preparations as potential drug carriers 10 have been reviewed (see, e.g., U.S. Patent No. 5,567,434, to Szoka; U.S. Patent No. 5,552,157, to Yagi; U.S. Patent No. 5,565,213, to Nakamori; U.S. Patent No. 5,738,868, to Shinkarenko; and U.S. Patent No. 5,795,587, to Gao).

For transdermal applications, the MASP-2 inhibitory antibodies and polypeptides may be combined with other suitable ingredients, such as carriers and/or adjuvants. 15 There are no limitations on the nature of such other ingredients, except that they must be pharmaceutically acceptable for their intended administration, and cannot degrade the activity of the active ingredients of the composition. Examples of suitable vehicles include ointments, creams, gels, or suspensions, with or without purified collagen. The MASP-2 inhibitory antibodies and polypeptides may also be impregnated into 20 transdermal patches, plasters, and bandages, preferably in liquid or semi-liquid form.

The compositions of the present invention may be systemically administered on a periodic basis at intervals determined to maintain a desired level of therapeutic effect. For example, compositions may be administered, such as by subcutaneous injection, every two to four weeks or at less frequent intervals. The dosage regimen will be 25 determined by the physician considering various factors that may influence the action of the combination of agents. These factors will include the extent of progress of the condition being treated, the patient's age, sex and weight, and other clinical factors. The dosage for each individual agent will vary as a function of the MASP-2 inhibitory agent that is included in the composition, as well as the presence and nature of any drug 30 delivery vehicle (e.g., a sustained release delivery vehicle). In addition, the dosage quantity may be adjusted to account for variation in the frequency of administration and the pharmacokinetic behavior of the delivered agent(s).

LOCAL DELIVERY

As used herein, the term "local" encompasses application of a drug in or around a site of intended localized action, and may include for example topical delivery to the skin or other affected tissues, ophthalmic delivery, intrathecal (IT), intracerebroventricular (ICV), intra-articular, intracavity, intracranial or intravesicular administration, placement or irrigation. Local administration may be preferred to enable administration of a lower dose, to avoid systemic side effects, and for more accurate control of the timing of delivery and concentration of the active agents at the site of local delivery. Local administration provides a known concentration at the target site, regardless of interpatient variability in metabolism, blood flow, etc. Improved dosage control is also provided by the direct mode of delivery.

Local delivery of a MASP-2 inhibitory agent may be achieved in the context of surgical methods for treating a disease or condition, such as for example during procedures such as arterial bypass surgery, atherectomy, laser procedures, ultrasonic procedures, balloon angioplasty and stent placement. For example, a MASP-2 inhibitor can be administered to a subject in conjunction with a balloon angioplasty procedure. A balloon angioplasty procedure involves inserting a catheter having a deflated balloon into an artery. The deflated balloon is positioned in proximity to the atherosclerotic plaque and is inflated such that the plaque is compressed against the vascular wall. As a result, the balloon surface is in contact with the layer of vascular endothelial cells on the surface of the blood vessel. The MASP-2 inhibitory agent may be attached to the balloon angioplasty catheter in a manner that permits release of the agent at the site of the atherosclerotic plaque. The agent may be attached to the balloon catheter in accordance with standard procedures known in the art. For example, the agent may be stored in a compartment of the balloon catheter until the balloon is inflated, at which point it is released into the local environment. Alternatively, the agent may be impregnated on the balloon surface, such that it contacts the cells of the arterial wall as the balloon is inflated. The agent may also be delivered in a perforated balloon catheter such as those disclosed in Flugelman, M.Y., et al., *Circulation* 85:1110-1117, 1992. See also published PCT Application WO 95/23161 for an exemplary procedure for attaching a therapeutic protein to a balloon angioplasty catheter. Likewise, the MASP-2 inhibitory agent may be included in a gel or polymeric coating applied to a stent, or may be incorporated into the

material of the stent, such that the stent elutes the MASP-2 inhibitory agent after vascular placement.

5 MASP-2 inhibitory compositions used in the treatment of arthritides and other musculoskeletal disorders may be locally delivered by intra-articular injection. Such compositions may suitably include a sustained release delivery vehicle. As a further example of instances in which local delivery may be desired, MASP-2 inhibitory compositions used in the treatment of urogenital conditions may be suitably instilled intravesically or within another urogenital structure.

10 COATINGS ON A MEDICAL DEVICE

MAASP-2 inhibitory agents such as antibodies and inhibitory peptides may be immobilized onto (or within) a surface of an implantable or attachable medical device. The modified surface will typically be in contact with living tissue after implantation into an animal body. By "implantable or attachable medical device" is intended any device
15 that is implanted into, or attached to, tissue of an animal body, during the normal operation of the device (e.g., stents and implantable drug delivery devices). Such implantable or attachable medical devices can be made from, for example, nitrocellulose, diazocellulose, glass, polystyrene, polyvinylchloride, polypropylene, polyethylene, dextran, Sepharose, agar, starch, nylon, stainless steel, titanium and biodegradable and/or
20 biocompatible polymers. Linkage of the protein to a device can be accomplished by any technique that does not destroy the biological activity of the linked protein, for example by attaching one or both of the N- C-terminal residues of the protein to the device. Attachment may also be made at one or more internal sites in the protein. Multiple attachments (both internal and at the ends of the protein) may also be used. A surface of
25 an implantable or attachable medical device can be modified to include functional groups (e.g., carboxyl, amide, amino, ether, hydroxyl, cyano, nitrido, sulfanamido, acetylinic, epoxide, silanic, anhydric, succinimic, azido) for protein immobilization thereto. Coupling chemistries include, but are not limited to, the formation of esters, ethers, amides, azido and sulfanamido derivatives, cyanate and other linkages to the functional
30 groups available on MASP-2 antibodies or inhibitory peptides. MASP-2 antibodies or inhibitory fragments can also be attached non-covalently by the addition of an affinity tag sequence to the protein, such as GST (D.B. Smith and K.S. Johnson, *Gene* 67:31, 1988),

polyhistidines (E. Hochuli et al., *J. Chromatog.* 411:77, 1987), or biotin. Such affinity tags may be used for the reversible attachment of the protein to a device.

Proteins can also be covalently attached to the surface of a device body, for example, by covalent activation of the surface of the medical device. By way of
5 representative example, matricellular protein(s) can be attached to the device body by any of the following pairs of reactive groups (one member of the pair being present on the surface of the device body, and the other member of the pair being present on the matricellular protein(s)): hydroxyl/carboxylic acid to yield an ester linkage; hydroxyl/anhydride to yield an ester linkage; hydroxyl/isocyanate to yield a urethane
10 linkage. A surface of a device body that does not possess useful reactive groups can be treated with radio-frequency discharge plasma (RFGD) etching to generate reactive groups in order to allow deposition of matricellular protein(s) (e.g., treatment with oxygen plasma to introduce oxygen-containing groups; treatment with propyl amino plasma to introduce amine groups).

15 MASP-2 inhibitory agents comprising nucleic acid molecules such as antisense, RNAi-or DNA-encoding peptide inhibitors can be embedded in porous matrices attached to a device body. Representative porous matrices useful for making the surface layer are those prepared from tendon or dermal collagen, as may be obtained from a variety of commercial sources (e.g., Sigma and Collagen Corporation), or collagen matrices
20 prepared as described in U.S. Patent Nos. 4,394,370, to Jefferies, and 4,975,527, to Koezuka. One collagenous material is termed UltraFiberTM and is obtainable from Norian Corp. (Mountain View, California).

Certain polymeric matrices may also be employed if desired, and include acrylic ester polymers and lactic acid polymers, as disclosed, for example, in U.S. Patent
25 Nos. 4,526,909 and 4,563,489, to Urist. Particular examples of useful polymers are those of orthoesters, anhydrides, propylene-cofumarates, or a polymer of one or more α -hydroxy carboxylic acid monomers, (e.g., α -hydroxy acetic acid (glycolic acid) and/or α -hydroxy propionic acid (lactic acid)).

30 TREATMENT REGIMENS

In prophylactic applications, the pharmaceutical compositions are administered to a subject susceptible to, or otherwise at risk of, a condition associated with MASP-2-dependent complement activation in an amount sufficient to eliminate or reduce

the risk of developing symptoms of the condition. In therapeutic applications, the pharmaceutical compositions are administered to a subject suspected of, or already suffering from, a condition associated with MASP-2-dependent complement activation in a therapeutically effective amount sufficient to relieve, or at least partially reduce, the symptoms of the condition. In both prophylactic and therapeutic regimens, compositions comprising MASP-2 inhibitory agents may be administered in several dosages until a sufficient therapeutic outcome has been achieved in the subject. Application of the MASP-2 inhibitory compositions of the present invention may be carried out by a single administration of the composition, or a limited sequence of administrations, for treatment of an acute condition, e.g., reperfusion injury or other traumatic injury. Alternatively, the composition may be administered at periodic intervals over an extended period of time for treatment of chronic conditions, e.g., arthritides or psoriasis.

The methods and compositions of the present invention may be used to inhibit inflammation and related processes that typically result from diagnostic and therapeutic medical and surgical procedures. To inhibit such processes, the MASP-2 inhibitory composition of the present invention may be applied periprocedurally. As used herein "periprocedurally" refers to administration of the inhibitory composition preprocedurally and/or intraprocedurally and/or postprocedurally, i.e., before the procedure, before and during the procedure, before and after the procedure, before, during and after the procedure, during the procedure, during and after the procedure, or after the procedure. Periprocedural application may be carried out by local administration of the composition to the surgical or procedural site, such as by injection or continuous or intermittent irrigation of the site or by systemic administration. Suitable methods for local perioperative delivery of MASP-2 inhibitory agent solutions are disclosed in US Patent Nos. 6,420,432 to Demopoulos and 6,645,168 to Demopoulos. Suitable methods for local delivery of chondroprotective compositions including MASP-2 inhibitory agent(s) are disclosed in International PCT Patent Application WO 01/07067 A2. Suitable methods and compositions for targeted systemic delivery of chondroprotective compositions including MASP-2 inhibitory agent(s) are disclosed in International PCT Patent Application WO 03/063799 A2.

In one aspect of the invention, the pharmaceutical compositions are administered to a subject suffering from, or at risk for developing a thrombotic microangiopathy (TMA). In one embodiment, the TMA is selected from the group consisting of hemolytic

uremic syndrome (HUS), thrombotic thrombocytopenic purpura (TTP) and atypical hemolytic uremic syndrome (aHUS). In one embodiment, the TMA is aHUS. In one embodiment, the composition is administered to an aHUS patient during the acute phase of the disease. In one embodiment, the composition is administered to an aHUS patient
5 during the remission phase (i.e., in a subject that has recovered or partially recovered from an episode of acute phase aHUS, such remission evidenced, for example, by increased platelet count and/or reduced serum LDH concentrations, for example as described in Loirat C et al., *Orphanet Journal of Rare Diseases* 6:60, 2011, hereby incorporated herein by reference). In one embodiment, the subject is suffering from, or at
10 risk for developing a TMA that is (i) a TMA secondary to cancer; (ii) a TMA secondary to chemotherapy; or (iii) a TMA secondary to transplantation (e.g., organ transplantation, such as kidney transplantation or allogeneic hematopoietic stem cell transplantation). In one embodiment, the subject is suffering from, or at risk for developing Upshaw-Schulman Syndrome (USS). In one embodiment, the subject is suffering from, or at risk
15 for developing Degos disease. In one embodiment, the subject is suffering from, or at risk for developing Catastrophic Antiphospholipid Syndrome (CAPS). In therapeutic applications, the pharmaceutical compositions are administered to a subject suffering from, or at risk for developing a TMA in a therapeutically effective amount sufficient to inhibit thrombus formation, relieve, or at least partially reduce, the symptoms of the
20 condition.

In both prophylactic and therapeutic regimens, compositions comprising MASP-2 inhibitory agents may be administered in several dosages until a sufficient therapeutic outcome has been achieved in the subject. In one embodiment of the invention, the MASP-2 inhibitory agent comprises an anti-MASP-2 antibody, which suitably may be
25 administered to an adult patient (e.g., an average adult weight of 70 kg) in a dosage of from 0.1 mg to 10,000 mg, more suitably from 1.0 mg to 5,000 mg, more suitably 10.0 mg to 2,000 mg, more suitably 10.0 mg to 1,000 mg and still more suitably from 50.0 mg to 500 mg. For pediatric patients, dosage can be adjusted in proportion to the patient's weight. Application of the MASP-2 inhibitory compositions of the present invention may
30 be carried out by a single administration of the composition, or a limited sequence of administrations, for treatment of TMA. Alternatively, the composition may be administered at periodic intervals such as daily, biweekly, weekly, every other week, monthly or bimonthly over an extended period of time for treatment of TMA.

In some embodiments, the subject suffering from or at risk for developing a TMA has previously undergone, or is currently undergoing treatment with a terminal complement inhibitor that inhibits cleavage of complement protein C5. In some embodiments, the method comprises administering to the subject a composition of the invention comprising a MASP-2 inhibitor and further administering to the subject a terminal complement inhibitor that inhibits cleavage of complement protein C5. In some embodiments, the terminal complement inhibitor is a humanized anti-C5 antibody or antigen-binding fragment thereof. In some embodiments, the terminal complement inhibitor is eculizumab.

In one aspect of the invention, the pharmaceutical compositions are administered to a subject susceptible to, or otherwise at risk of, aHUS in an amount sufficient to eliminate or reduce the risk of developing symptoms of the condition. In therapeutic applications, the pharmaceutical compositions are administered to a subject suspected of, or already suffering from, aHUS in a therapeutically effective amount sufficient to relieve, or at least partially reduce, the symptoms of the condition. In one aspect of the invention, prior to administration, the subject may be examined to determine whether the subject exhibits one or more symptoms of aHUS, including (i) anemia, (ii) thrombocytopenia (iii) renal insufficiency and (iv) rising creatinine, and the composition of the present invention is then administered in an effective amount and for a sufficient time period to improve these symptom(s).

In another aspect of the invention, the MASP-2 inhibitory compositions of the present invention may be used to prophylactically treat a subject that has an elevated risk of developing aHUS and thereby reduce the likelihood that the subject will develop aHUS. The presence of a genetic marker in the subject known to be associated with aHUS is first determined by performing a genetic screening test on a sample obtained from the subject and identifying the presence of at least one genetic marker associated with aHUS, complement factor H (CFH), factor I (CFI), factor B (CFB), membrane cofactor CD46, C3, complement factor H-related protein (CFHR1), anticoagulant protein thrombomodulin (THBD), complement factor H-related protein 3 (CFHR3) or complement factor H-related protein 4 (CFHR4). The subject is then periodically monitored (e.g., monthly, quarterly, twice annually or annually) to determine the presence or absence of at least one symptom of aHUS, such as anemia, thrombocytopenia, renal insufficiency and rising creatinine. Upon the determination of the presence of at least one of these symptoms, the

subject can be administered an amount of a MASP-2 inhibitory agent effective to inhibit MASP-2 dependent complement activation, in an effective amount and for a sufficient time period to improve said one or more symptoms. In a still further aspect of the present invention, a subject at increased risk of developing aHUS due to having been screened
5 and determined to have one of the genetic markers associated with aHUS may be monitored for the occurrence of an event associated with triggering aHUS clinical symptoms, including drug exposure, infection (e.g., bacterial infection), malignancy, injury, organ or tissue transplant and pregnancy.

In a still further aspect of the present invention, a composition comprising an
10 amount of a MASP-2 inhibitory agent effective to inhibit MASP-2 dependent complement activation can be administered to a suffering from or at risk of developing atypical hemolytic uremic syndrome (aHUS) secondary to an infection. For example, a patient suffering from or at risk of developing non-enteric aHUS associated with an *S. pneumonia* infection may be treated with the compositions of the present invention.

In a still further aspect of the present invention, a subject suffering from aHUS
15 may initially be treated with a MASP-2 inhibitory composition of the present invention that is administered through a catheter line, such as an intravenous catheter line or a subcutaneous catheter line, for a first period of time such as one hour, twelve hours, one day, two days or three days. The subject may then be treated for a second period of time
20 with the MASP-2 inhibitory composition administered through regular subcutaneous injections, such as daily, biweekly, weekly, every other week, monthly or bimonthly, injections.

In a still further aspect of the present invention, a MASP-2 inhibitory composition
of the present invention may be administered to a subject suffering from aHUS in the
25 absence of plasmapheresis (i.e., a subject whose aHUS symptoms have not been treated with plasmapheresis and are not treated with plasmapheresis at the time of treatment with the MASP-2 inhibitory composition), to avoid the potential complications of plasmapheresis including hemorrhage, infection, and exposure to disorders and/or allergies inherent in the plasma donor, or in a subject otherwise averse to plasmapheresis, or in a
30 setting where plasmapheresis is unavailable.

In a still further aspect of the present invention, a MASP-2 inhibitory composition
of the present invention may be administered to a subject suffering from aHUS coincident
with treating the patient with plasmapheresis. For example, a subject receiving

plasmapheresis treatment can then be administered the MASP-2 inhibitory composition following or alternating with plasma exchange.

5 In a still further aspect of the present invention, a subject suffering from or at risk of developing aHUS and being treated with a MASP-2 inhibitory composition of the present invention can be monitored by periodically determining, such as every twelve hours or on a daily basis, the level of at least one complement factor, wherein the determination of a reduced level of the at least one complement factor in comparison to a standard value or to a healthy subject is indicative of the need for continued treatment with the composition.

10 In both prophylactic and therapeutic regimens, compositions comprising MASP-2 inhibitory agents may be administered in several dosages until a sufficient therapeutic outcome has been achieved in the subject. In one embodiment of the invention, the MASP-2 inhibitory agent comprises an anti-MASP-2 antibody, which suitably may be administered to an adult patient (e.g., an average adult weight of 70 kg) in a dosage of
15 from 0.1 mg to 10,000 mg, more suitably from 1.0 mg to 5,000 mg, more suitably 10.0 mg to 2,000 mg, more suitably 10.0 mg to 1,000 mg and still more suitably from 50.0 mg to 500 mg. For pediatric patients, dosage can be adjusted in proportion to the patient's weight. Application of the MASP-2 inhibitory compositions of the present invention may be carried out by a single administration of the composition, or a limited sequence of
20 administrations, for treatment of aHUS. Alternatively, the composition may be administered at periodic intervals, such as daily, biweekly, weekly, every other week, monthly or bimonthly, over an extended period of time for treatment of aHUS.

In some embodiments, the subject suffering from aHUS has previously undergone, or is currently undergoing treatment with a terminal complement inhibitor
25 that inhibits cleavage of complement protein C5. In some embodiments, the method comprises administering to the subject a composition of the invention comprising a MASP-2 inhibitor and further administering to the subject a terminal complement inhibitor that inhibits cleavage of complement protein C5. In some embodiments, the terminal complement inhibitor is a humanized anti-C5 antibody or antigen-binding
30 fragment thereof. In some embodiments, the terminal complement inhibitor is eculizumab.

In one aspect of the invention, the pharmaceutical compositions are administered to a subject susceptible to, or otherwise at risk of, HUS in an amount sufficient to

eliminate or reduce the risk of developing symptoms of the condition. In therapeutic applications, the pharmaceutical compositions are administered to a subject suspected of, or already suffering from, HUS in a therapeutically effective amount sufficient to relieve, or at least partially reduce, the symptoms of the condition.

5 In another aspect of the present invention, the likelihood of developing impaired renal function in a subject at risk for developing HUS can be reduced by administering to the subject a MASP-2 inhibitory composition of the present invention in an amount effective to inhibit MASP-2 dependent complement activation. For example, a subject at risk for developing HUS and to be treated with a MASP-2 inhibitory composition of the present invention may exhibit one or more symptoms associated with HUS, including
10 diarrhea, a hematocrit level of less than 30% with smear evidence of intravascular erythrocyte destruction, thrombocytopenia and rising creatinine levels. As a further example, a subject at risk for developing HUS and to be treated with the MASP-2 inhibitory compositions of the present invention may be infected with *E. coli*, shigella or
15 salmonella. Such subjects infected with *E. coli*, shigella or salmonella may be treated with a MASP-2 inhibitory composition of the present invention concurrent with antibiotic treatment, or alternately may be treated with a MASP-2 inhibitory composition without concurrent treatment with an antibiotic, particularly for enterogenic *E. coli* for which antibiotic treatment is contra-indicated. A subject infected with enterogenic *E. coli* that
20 has been treated with an antibiotic may be at elevated risk of developing HUS, and may be suitably treated with a MASP-2 inhibitory composition of the present invention to reduce that risk. A subject infected with enterogenic *E. coli* may be treated for a first period of time with a MASP-2 inhibitory composition of the present invention in the absence of an antibiotic and then for a second period of time with both a MASP-2
25 inhibitory composition of the present invention and an antibiotic.

 In a still further aspect of the present invention, a subject suffering from HUS may initially be treated with a MASP-2 inhibitory composition of the present invention that is administered through a catheter line, such as an intravenous catheter line or a subcutaneous catheter line, for a first period of time such as one hour, twelve hours, one
30 day, two days or three days. The subject may then be treated for a second period of time with the MASP-2 inhibitory composition administered through regular subcutaneous injections, such as daily, biweekly, weekly, every other week, monthly or bimonthly, injections.

In a still further aspect of the present invention, a MASP-2 inhibitory composition of the present invention may be administered to a subject suffering from HUS in the absence of plasmapheresis (i.e., a subject whose HUS symptoms have not been treated with plasmapheresis and are not treated with plasmapheresis at the time of treatment with
5 the MASP-2 inhibitory composition), to avoid the potential complications of plasmapheresis including hemorrhage, infection, and exposure to disorders and/or allergies inherent in the plasma donor, or in a subject otherwise averse to plasmapheresis, or in a setting where plasmapheresis is unavailable.

In a still further aspect of the present invention, a MASP-2 inhibitory composition
10 of the present invention may be administered to a subject suffering from HUS coincident with treating the patient with plasmapheresis. For example, a subject receiving plasmapheresis treatment can then be administered the MASP-2 inhibitory composition following or alternating with plasma exchange.

In a still further aspect of the present invention, a subject suffering from or at risk
15 of developing HUS and being treated with a MASP-2 inhibitory composition of the present invention can be monitored by periodically determining, such as every twelve hours or on a daily basis, the level of at least one complement factor, wherein the determination of a reduced level of the at least one complement factor in comparison to a standard value or to a healthy subject is indicative of the need for continued treatment
20 with the composition.

In both prophylactic and therapeutic regimens, compositions comprising MASP-2 inhibitory agents may be administered in several dosages until a sufficient therapeutic outcome has been achieved in the subject. In one embodiment of the invention, the MASP-2 inhibitory agent comprises an anti-MASP-2 antibody, which suitably may be
25 administered to an adult patient (e.g., an average adult weight of 70 kg) in a dosage of from 0.1 mg to 10,000 mg, more suitably from 1.0 mg to 5,000 mg, more suitably 10.0 mg to 2,000 mg, more suitably 10.0 mg to 1,000 mg and still more suitably from 50.0 mg to 500 mg. For pediatric patients, dosage can be adjusted in proportion to the patient's weight. Application of the MASP-2 inhibitory compositions of the present invention may
30 be carried out by a single administration of the composition, or a limited sequence of administrations, for treatment of HUS. Alternatively, the composition may be administered at periodic intervals, such as daily, biweekly, weekly, every other week, monthly or bimonthly, over an extended period of time for treatment of HUS.

In some embodiments, the subject suffering from HUS has previously undergone, or is currently undergoing treatment with a terminal complement inhibitor that inhibits cleavage of complement protein C5. In some embodiments, the method comprises administering to the subject a composition of the invention comprising a MASP-2 inhibitor and further administering to the subject a terminal complement inhibitor that inhibits cleavage of complement protein C5. In some embodiments, the terminal complement inhibitor is a humanized anti-C5 antibody or antigen-binding fragment thereof. In some embodiments, the terminal complement inhibitor is eculizumab.

In one aspect of the invention, the pharmaceutical compositions are administered to a subject susceptible to, or otherwise at risk of, TTP in an amount sufficient to eliminate or reduce the risk of developing symptoms of the condition. In therapeutic applications, the pharmaceutical compositions are administered to a subject suspected of, or already suffering from, TTP in a therapeutically effective amount sufficient to relieve, or at least partially reduce, the symptoms of the condition.

In another aspect of the present invention, a subject exhibiting one or more of the symptoms of TTP, including central nervous system involvement, thrombocytopenia, severe cardiac involvement, severe pulmonary involvement, gastro-intestinal infarction and gangrene, may be treated with a MASP-2 inhibitory composition of the present invention. In another aspect of the present invention, a subject determined to have a depressed level of ADAMTS13 and also testing positive for the presence of an inhibitor of (i.e., an antibody) ADAMTS13 may be treated with a MASP-2 inhibitory composition of the present invention. In a still further aspect of the present invention, a subject testing positive for the presence of an inhibitor of ADAMTS13 may be treated with an immunosuppressant (e.g., corticosteroids, rituxan, or cyclosporine) concurrently with treatment with a MASP-2 inhibitory composition of the present invention. In a still further aspect of the present invention, a subject determined to have a reduced level of ADAMTS13 and testing positive for the presence of an inhibitor of ADAMTS13 may be treated with ADAMTS13 concurrently with treatment with a MASP-2 inhibitory composition of the present invention.

In a still further aspect of the present invention, a subject suffering from TTP may initially be treated with a MASP-2 inhibitory composition of the present invention that is administered through a catheter line, such as an intravenous catheter line or a subcutaneous catheter line, for a first period of time such as one hour, twelve hours, one

day, two days or three days. The subject may then be treated for a second period of time with the MASP-2 inhibitory composition administered through regular subcutaneous injections, such as daily, biweekly, weekly, every other week, monthly or bimonthly, injections.

5 In a still further aspect of the present invention, a MASP-2 inhibitory composition of the present invention may be administered to a subject suffering from HUS in the absence of plasmapheresis (i.e., a subject whose TTP symptoms have not been treated with plasmapheresis and are not treated with plasmapheresis at the time of treatment with the MASP-2 inhibitory composition), to avoid the potential complications of
10 plasmapheresis including hemorrhage, infection, and exposure to disorders and/or allergies inherent in the plasma donor, or in a subject otherwise averse to plasmapheresis, or in a setting where plasmapheresis is unavailable.

In a still further aspect of the present invention, a MASP-2 inhibitory composition of the present invention may be administered to a subject suffering from TTP coincident
15 with treating the patient with plasmapheresis. For example, a subject receiving plasmapheresis treatment can then be administered the MASP-2 inhibitory composition following or alternating with plasma exchange.

In a still further aspect of the present invention, a subject suffering from refractory TTP, i.e., symptoms of TTP that have not responded adequately to other treatment such
20 as plasmapheresis, may be treated with a MASP-2 inhibitory composition of the present invention, with or without additional plasmapheresis.

In a still further aspect of the present invention, a subject suffering from or at risk of developing TTP and being treated with a MASP-2 inhibitory composition of the present invention can be monitored by periodically determining, such as every twelve
25 hours or on a daily basis, the level of at least one complement factor, wherein the determination of a reduced level of the at least one complement factor in comparison to a standard value or to a healthy subject is indicative of the need for continued treatment with the composition.

In both prophylactic and therapeutic regimens, compositions comprising MASP-2
30 inhibitory agents may be administered in several dosages until a sufficient therapeutic outcome has been achieved in the subject. In one embodiment of the invention, the MASP-2 inhibitory agent comprises an anti-MASP-2 antibody, which suitably may be administered to an adult patient (e.g., an average adult weight of 70 kg) in a dosage of

from 0.1 mg to 10,000 mg, more suitably from 1.0 mg to 5,000 mg, more suitably 10.0 mg to 2,000 mg, more suitably 10.0 mg to 1,000 mg and still more suitably from 50.0 mg to 500 mg. For pediatric patients, dosage can be adjusted in proportion to the patient's weight. Application of the MASP-2 inhibitory compositions of the present invention may
5 be carried out by a single administration of the composition, or a limited sequence of administrations, for treatment of TTP. Alternatively, the composition may be administered at periodic intervals, such as daily, biweekly, weekly, every other week, monthly or bimonthly, over an extended period of time for treatment of TTP.

In some embodiments, the subject suffering from TTP has previously undergone,
10 or is currently undergoing treatment with a terminal complement inhibitor that inhibits cleavage of complement protein C5. In some embodiments, the method comprises administering to the subject a composition of the invention comprising a MASP-2 inhibitor and further administering to the subject a terminal complement inhibitor that inhibits cleavage of complement protein C5. In some embodiments, the terminal
15 complement inhibitor is a humanized anti-C5 antibody or antigen-binding fragment thereof. In some embodiments, the terminal complement inhibitor is eculizumab.

VI. EXAMPLES

The following examples merely illustrate the best mode now contemplated for
20 practicing the invention, but should not be construed to limit the invention. All literature citations herein are expressly incorporated by reference.

EXAMPLE 1

This example describes the generation of a mouse strain deficient in MASP-2
25 (MASP-2^{-/-}) but sufficient of MAP19 (MAP19^{+/+}).

Materials and Methods: The targeting vector pKO-NTKV 1901 was designed to disrupt the three exons coding for the C-terminal end of murine MASP-2, including the exon that encodes the serine protease domain, as shown in **FIGURE 3**. PKO-NTKV 1901 was used to transfect the murine ES cell line E14.1a (SV129 Ola).
30 Neomycin-resistant and Thymidine Kinase-sensitive clones were selected. 600 ES clones were screened and, of these, four different clones were identified and verified by southern blot to contain the expected selective targeting and recombination event as shown in **FIGURE 3**. Chimeras were generated from these four positive clones by embryo

transfer. The chimeras were then backcrossed in the genetic background C57/BL6 to create transgenic males. The transgenic males were crossed with females to generate F1s with 50% of the offspring showing heterozygosity for the disrupted *MASP-2* gene. The heterozygous mice were intercrossed to generate homozygous MASP-2 deficient offspring, resulting in heterozygous and wild-type mice in the ration of 1:2:1, respectively.

Results and Phenotype: The resulting homozygous MASP-2^{-/-} deficient mice were found to be viable and fertile and were verified to be MASP-2 deficient by southern blot to confirm the correct targeting event, by Northern blot to confirm the absence of MASP-2 mRNA, and by Western blot to confirm the absence of MASP-2 protein (data not shown). The presence of MASP19 mRNA and the absence of MASP-2 mRNA were further confirmed using time-resolved RT-PCR on a LightCycler machine. The MASP-2^{-/-} mice do continue to express MASP19, MASP-1, and MASP-3 mRNA and protein as expected (data not shown). The presence and abundance of mRNA in the MASP-2^{-/-} mice for Properdin, Factor B, Factor D, C4, C2, and C3 was assessed by LightCycler analysis and found to be identical to that of the wild-type littermate controls (data not shown). The plasma from homozygous MASP-2^{-/-} mice is totally deficient of lectin-pathway-mediated complement activation as further described in **Example 2**.

Generation of a MASP-2^{-/-} strain on a pure C57BL6 Background: The MASP-2^{-/-} mice were back-crossed with a pure C57BL6 line for nine generations prior to use of the MASP-2^{-/-} strain as an experimental animal model.

A transgenic mouse strain that is murine MASP-2^{-/-}, MASP19^{+/+} and that expresses a human MASP-2 transgene (a murine MASP-2 knock-out and a human MASP-2 knock-in) was also generated as follows:

Materials and Methods: A minigene encoding human MASP-2 called "mini hMASP-2" (SEQ ID NO:49) as shown in **FIGURE 4** was constructed which includes the promoter region of the human *MASP 2* gene, including the first 3 exons (exon 1 to exon 3) followed by the cDNA sequence that represents the coding sequence of the following 8 exons, thereby encoding the full-length MASP-2 protein driven by its endogenous promoter. The mini hMASP-2 construct was injected into fertilized eggs of MASP-2^{-/-} in order to replace the deficient murine *MASP 2* gene by transgenically expressed human MASP-2.

EXAMPLE 2

This example demonstrates that MASP-2 is required for complement activation via the lectin pathway.

Methods and Materials:

5 **Lectin pathway specific C4 Cleavage Assay:** A C4 cleavage assay has been described by Petersen, et al., *J. Immunol. Methods* 257:107 (2001) that measures lectin pathway activation resulting from lipoteichoic acid (LTA) from *S. aureus*, which binds L-ficolin. The assay described by Petersen et al., (2001) was adapted to measure lectin pathway activation via MBL by coating the plate with LPS and mannan or zymosan prior
10 to adding serum from MASP-2 ^{-/-} mice as described below. The assay was also modified to remove the possibility of C4 cleavage due to the classical pathway. This was achieved by using a sample dilution buffer containing 1 M NaCl, which permits high affinity binding of lectin pathway recognition components to their ligands but prevents activation of endogenous C4, thereby excluding the participation of the classical pathway by
15 dissociating the C1 complex. Briefly described, in the modified assay serum samples (diluted in high salt (1 M NaCl) buffer) are added to ligand-coated plates, followed by the addition of a constant amount of purified C4 in a buffer with a physiological concentration of salt. Bound recognition complexes containing MASP-2 cleave the C4, resulting in C4b deposition.

20 Assay Methods:

1) Nunc Maxisorb microtiter plates (Maxisorb, Nunc, Cat. No. 442404, Fisher Scientific) were coated with 1 µg/ml mannan (M7504 Sigma) or any other ligand (e.g., such as those listed below) diluted in coating buffer (15 mM Na₂CO₃, 35 mM NaHCO₃, pH 9.6).

25 The following reagents were used in the assay:

- a. mannan (1 µg/well mannan (M7504 Sigma) in 100 µl coating buffer);
 - b. zymosan (1 µg/well zymosan (Sigma) in 100 µl coating buffer);
 - c. LTA (1µg/well in 100 µl coating buffer or 2 µg/well in 20 µl methanol)
 - d. 1 µg of the H-ficolin specific Mab 4H5 in coating buffer
 - 30 e. PSA from *Aerococcus viridans* (2 µg/well in 100 µl coating buffer)
 - f. 100 µl/well of formalin-fixed *S. aureus* DSM20233 (OD₅₅₀=0.5) in coating buffer.
- 2) The plates were incubated overnight at 4°C.

3) After overnight incubation, the residual protein binding sites were saturated by incubated the plates with 0.1% HSA-TBS blocking buffer (0.1% (w/v) HSA in 10 mM Tris-CL, 140 mM NaCl, 1.5 mM NaN₃, pH 7.4) for 1-3 hours, then washing the plates 3X with TBS/tween/Ca²⁺ (TBS with 0.05% Tween 20 and 5 mM CaCl₂,
 5 1 mM MgCl₂, pH 7.4).

4) Serum samples to be tested were diluted in MBL-binding buffer (1 M NaCl) and the diluted samples were added to the plates and incubated overnight at 4°C. Wells receiving buffer only were used as negative controls.

5) Following incubation overnight at 4°C, the plates were washed 3X with
 10 TBS/tween/Ca²⁺. Human C4 (100 µl/well of 1 µg/ml diluted in BBS (4 mM barbital, 145 mM NaCl, 2 mM CaCl₂, 1 mM MgCl₂, pH 7.4)) was then added to the plates and incubated for 90 minutes at 37°C. The plates were washed again 3X with TBS/tween/Ca²⁺.

6) C4b deposition was detected with an alkaline phosphatase-conjugated
 15 chicken anti-human C4c (diluted 1:1000 in TBS/tween/Ca²⁺), which was added to the plates and incubated for 90 minutes at room temperature. The plates were then washed again 3X with TBS/tween/Ca²⁺.

7) Alkaline phosphatase was detected by adding 100 µl of *p*-nitrophenyl phosphate substrate solution, incubating at room temperature for 20 minutes, and reading
 20 the OD₄₀₅ in a microtiter plate reader.

Results: FIGURES 5A-B show the amount of C4b deposition on mannan (FIGURE 5A) and zymosan (FIGURE 5B) in serum dilutions from MASP-2^{+/+} (crosses), MASP-2^{+/-} (closed circles) and MASP-2^{-/-} (closed triangles). FIGURE 5C shows the relative C4 convertase activity on plates coated with zymosan (white bars) or
 25 mannan (shaded bars) from MASP-2^{+/+} mice (n=5) and MASP-2^{-/-} mice (n=4) relative to wild-type mice (n=5) based on measuring the amount of C4b deposition normalized to wild-type serum. The error bars represent the standard deviation. As shown in FIGURES 5A-C, plasma from MASP-2^{-/-} mice is totally deficient in lectin-pathway-mediated complement activation on mannan and on zymosan coated
 30 plates. These results clearly demonstrate that MASP-2 is an effector component of the lectin pathway.

Recombinant MASP-2 reconstitutes Lectin Pathway-Dependent C4 Activation in serum from the MASP-2^{-/-} mice

In order to establish that the absence of MASP-2 was the direct cause of the loss of lectin pathway-dependent C4 activation in the MASP-2^{-/-} mice, the effect of adding recombinant MASP-2 protein to serum samples was examined in the C4 cleavage assay described above. Functionally active murine MASP-2 and catalytically inactive murine MASP-2A (in which the active-site serine residue in the serine protease domain was substituted for the alanine residue) recombinant proteins were produced and purified as described below in **Example 3**. Pooled serum from 4 MASP-2^{-/-} mice was pre-incubated with increasing protein concentrations of recombinant murine MASP-2 or inactive recombinant murine MASP-2A and C4 convertase activity was assayed as described above.

Results: As shown in **FIGURE 6**, the addition of functionally active murine recombinant MASP-2 protein (shown as open triangles) to serum obtained from the MASP-2^{-/-} mice restored lectin pathway-dependent C4 activation in a protein concentration dependent manner, whereas the catalytically inactive murine MASP-2A protein (shown as stars) did not restore C4 activation. The results shown in **FIGURE 6** are normalized to the C4 activation observed with pooled wild-type mouse serum (shown as a dotted line).

EXAMPLE 3

This example describes the recombinant expression and protein production of recombinant full-length human, rat and murine MASP-2, MASP-2 derived polypeptides, and catalytically inactivated mutant forms of MASP-2

Expression of Full-length human, murine and rat MASP-2:

The full length cDNA sequence of human MASP-2 (SEQ ID NO: 4) was also subcloned into the mammalian expression vector pCI-Neo (Promega), which drives eukaryotic expression under the control of the CMV enhancer/promoter region (described in Kaufman R.J. et al., *Nucleic Acids Research* 19:4485-90, 1991; Kaufman, *Methods in Enzymology*, 185:537-66 (1991)). The full length mouse cDNA (SEQ ID NO:50) and rat MASP-2 cDNA (SEQ ID NO:53) were each subcloned into the pED expression vector. The MASP-2 expression vectors were then transfected into the adherent Chinese hamster ovary cell line DXB1 using the standard calcium phosphate transfection procedure

described in Maniatis et al., 1989. Cells transfected with these constructs grew very slowly, implying that the encoded protease is cytotoxic.

In another approach, the minigene construct (SEQ ID NO:49) containing the human cDNA of MASP-2 driven by its endogenous promoter is transiently transfected into Chinese hamster ovary cells (CHO). The human MASP-2 protein is secreted into the culture media and isolated as described below.

Expression of Full-length catalytically inactive MASP-2:

Rationale: MASP-2 is activated by autocatalytic cleavage after the recognition subcomponents MBL or ficolins (either L-ficolin, H-ficolin or M-ficolin) bind to their respective carbohydrate pattern. Autocatalytic cleavage resulting in activation of MASP-2 often occurs during the isolation procedure of MASP-2 from serum, or during the purification following recombinant expression. In order to obtain a more stable protein preparation for use as an antigen, a catalytically inactive form of MASP-2, designed as MASP-2A was created by replacing the serine residue that is present in the catalytic triad of the protease domain with an alanine residue in rat (SEQ ID NO:55 Ser617 to Ala617); in mouse (SEQ ID NO:52 Ser617 to Ala617); or in human (SEQ ID NO:3 Ser618 to Ala618).

In order to generate catalytically inactive human and murine MASP-2A proteins, site-directed mutagenesis was carried out using the oligonucleotides shown in **TABLE 5**. The oligonucleotides in **TABLE 5** were designed to anneal to the region of the human and murine cDNA encoding the enzymatically active serine and oligonucleotide contain a mismatch in order to change the serine codon into an alanine codon. For example, PCR oligonucleotides SEQ ID NOS:56-59 were used in combination with human MASP-2 cDNA (SEQ ID NO:4) to amplify the region from the start codon to the enzymatically active serine and from the serine to the stop codon to generate the complete open reading from of the mutated MASP-2A containing the Ser618 to Ala618 mutation. The PCR products were purified after agarose gel electrophoresis and band preparation and single adenosine overlaps were generated using a standard tailing procedure. The adenosine tailed MASP-2A was then cloned into the pGEM-T easy vector, transformed into *E. coli*.

A catalytically inactive rat MASP-2A protein was generated by kinasing and annealing SEQ ID NO:64 and SEQ ID NO:65 by combining these two oligonucleotides in equal molar amounts, heating at 100°C for 2 minutes and slowly cooling to room temperature. The resulting annealed fragment has PstI and XbaI compatible ends and

was inserted in place of the Pst1-Xba1 fragment of the wild-type rat MASP-2 cDNA (SEQ ID NO:53) to generate rat MASP-2A.

5 'GAGGTGACGCAGGAGGGGCATTAGTGTTT 3' (SEQ ID NO:64)

5' CTAGAAACACTAATGCCCCCTCCTGCGTCACCTCTGCA 3' (SEQ ID NO:65)

The human, murine and rat MASP-2A were each further subcloned into either of the mammalian expression vectors pED or pCI-Neo and transfected into the Chinese Hamster ovary cell line DXB1 as described below.

In another approach, a catalytically inactive form of MASP-2 is constructed using the method described in Chen et al., *J. Biol. Chem.*, 276(28):25894-25902, 2001. Briefly, the plasmid containing the full-length human MASP-2 cDNA (described in Thiel et al., *Nature* 386:506, 1997) is digested with *Xho*I and *Eco*R1 and the MASP-2 cDNA (described herein as SEQ ID NO:4) is cloned into the corresponding restriction sites of the pFastBac1 baculovirus transfer vector (Life Technologies, NY). The MASP-2 serine protease active site at Ser618 is then altered to Ala618 by substituting the double-stranded oligonucleotides encoding the peptide region amino acid 610-625 (SEQ ID NO:13) with the native region amino acids 610 to 625 to create a MASP-2 full length polypeptide with an inactive protease domain. Construction of Expression Plasmids Containing Polypeptide Regions Derived from Human Masp-2.

The following constructs are produced using the MASP-2 signal peptide (residues 1-15 of SEQ ID NO:5) to secrete various domains of MASP-2. A construct expressing the human MASP-2 CUB1 domain (SEQ ID NO:8) is made by PCR amplifying the region encoding residues 1-121 of MASP-2 (SEQ ID NO:6) (corresponding to the N-terminal CUB1 domain). A construct expressing the human MASP-2 CUBIEGF domain (SEQ ID NO:9) is made by PCR amplifying the region encoding residues 1-166 of MASP-2 (SEQ ID NO:6) (corresponding to the N-terminal CUBIEGF domain). A construct expressing the human MASP-2 CUBIEGFCUBII domain (SEQ ID NO:10) is made by PCR amplifying the region encoding residues 1-293 of MASP-2 (SEQ ID NO:6) (corresponding to the N-terminal CUBIEGFCUBII domain). The above mentioned domains are amplified by PCR using Ventr polymerase and pBS-MASP-2 as a template, according to established PCR methods. The 5' primer sequence of the sense primer (5'-CGGGATCCATGAGGCTGCTGACCCTC-3' SEQ ID NO:34) introduces a *Bam*HI restriction site (underlined) at the 5' end of the PCR

products. Antisense primers for each of the MASP-2 domains, shown below in TABLE 5, are designed to introduce a stop codon (boldface) followed by an *EcoRI* site (underlined) at the end of each PCR product. Once amplified, the DNA fragments are digested with *Bam*HI and *Eco*RI and cloned into the corresponding sites of the pFastBac1 vector. The resulting constructs are characterized by restriction mapping and confirmed by dsDNA sequencing.

TABLE 5: MASP-2 PCR PRIMERS

MASP-2 domain	5' PCR Primer	3' PCR Primer
SEQ ID NO:8 CUBI (aa 1–121 of SEQ ID NO:6)	5' <u>CGGGATCC</u> ATGA GGCTGCTGACCCT C-3' (SEQ ID NO:34)	5' <u>GGAATTC</u> CTAGGCTGCAT A (SEQ ID NO:35)
SEQ ID NO:9 CUBIEGF (aa 1–166 of SEQ ID NO:6)	5' <u>CGGGATCC</u> ATGA GGCTGCTGACCCT C-3' (SEQ ID NO:34)	5' <u>GGAATTC</u> CTACAGGGCGC T-3' (SEQ ID NO:36)
SEQ ID NO:10 CUBIEGFCUBII (aa 1-293 of SEQ ID NO:6)	5' <u>CGGGATCC</u> ATGA GGCTGCTGACCCT C-3' (SEQ ID NO:34)	5' <u>GGAATTC</u> CTAGTAGTGGA T 3' (SEQ ID NO:37)
SEQ ID NO:4 human MASP-2	5'ATGAGGCTGCTG ACCCTCCTGGGCC TTC 3' (SEQ ID NO: 56) hMASP-2 forward	5'TTAAAATCACTAATTATG TTCTCGATC 3' (SEQ ID NO: 59) hMASP-2_reverse
SEQ ID NO:4 human MASP-2 cDNA	5'CAGAGGTGACGC AGGAGGGGCAC 3' (SEQ ID NO: 58) hMASP-2_ala_forward	5'GTGCCCCCTCTGCGTCAC CTCTG 3' (SEQ ID NO: 57) hMASP-2_ala_reverse
SEQ ID NO:50 Murine MASP-2 cDNA	5'ATGAGGCTACTC ATCTTCCTGG3' (SEQ ID NO: 60) mMASP-2 forward	5'TTAGAAATTACTTATTAT GTTCTCAATCC3' (SEQ ID NO: 63) mMASP-2_reverse
SEQ ID NO:50 Murine MASP-2 cDNA	5'CCCCCCTGCGT CACCTCTGCAG3' (SEQ ID NO: 62) mMASP-2_ala_forward	5'CTGCAGAGGTGACGCAG GGGGG 3' (SEQ ID NO: 61) mMASP-2_ala_reverse

Recombinant eukaryotic expression of MASP-2 and protein production of enzymatically inactive mouse, rat, and human MASP-2A.

The MASP-2 and MASP-2A expression constructs described above were transfected into DXB1 cells using the standard calcium phosphate transfection procedure (Maniatis et al., 1989). MASP-2A was produced in serum-free medium to ensure that preparations were not contaminated with other serum proteins. Media was harvested from confluent cells every second day (four times in total). The level of recombinant MASP-2A averaged approximately 1.5 mg/liter of culture medium for each of the three species.

MASP-2A protein purification: The MASP-2A (Ser-Ala mutant described above) was purified by affinity chromatography on MBP-A-agarose columns. This strategy enabled rapid purification without the use of extraneous tags. MASP-2A (100-200 ml of medium diluted with an equal volume of loading buffer (50 mM Tris-Cl, pH 7.5, containing 150 mM NaCl and 25 mM CaCl₂) was loaded onto an MBP-agarose affinity column (4 ml) pre-equilibrated with 10 ml of loading buffer. Following washing with a further 10 ml of loading buffer, protein was eluted in 1 ml fractions with 50 mM Tris-Cl, pH 7.5, containing 1.25 M NaCl and 10 mM EDTA. Fractions containing the MASP-2A were identified by SDS-polyacrylamide gel electrophoresis. Where necessary, MASP-2A was purified further by ion-exchange chromatography on a MonoQ column (HR 5/5). Protein was dialysed with 50 mM Tris-Cl pH 7.5, containing 50 mM NaCl and loaded onto the column equilibrated in the same buffer. Following washing, bound MASP-2A was eluted with a 0.05–1 M NaCl gradient over 10 ml.

Results: Yields of 0.25–0.5 mg of MASP-2A protein were obtained from 200 ml of medium. The molecular mass of 77.5 kDa determined by MALDI-MS is greater than the calculated value of the unmodified polypeptide (73.5 kDa) due to glycosylation. Attachment of glycans at each of the *N*-glycosylation sites accounts for the observed mass. MASP-2A migrates as a single band on SDS-polyacrylamide gels, demonstrating that it is not proteolytically processed during biosynthesis. The weight-average molecular mass determined by equilibrium ultracentrifugation is in agreement with the calculated value for homodimers of the glycosylated polypeptide.

PRODUCTION OF RECOMBINANT HUMAN MASP-2 POLYPEPTIDES

Another method for producing recombinant MASP-2 and MASP2A derived polypeptides is described in Thielens, N.M., et al., *J. Immunol.* 166:5068-5077, 2001. Briefly, the *Spodoptera frugiperda* insect cells (Ready-Plaque Sf9 cells obtained from Novagen, Madison, WI) are grown and maintained in Sf900II serum-free medium (Life Technologies) supplemented with 50 IU/ml penicillin and 50 mg/ml streptomycin (Life Technologies). The *Trichoplusia ni* (High Five) insect cells (provided by Jadwiga Chroboczek, Institut de Biologie Structurale, Grenoble, France) are maintained in TC100 medium (Life Technologies) containing 10% FCS (Dominique Dutscher, Brumath, France) supplemented with 50 IU/ml penicillin and 50 mg/ml streptomycin. Recombinant baculoviruses are generated using the Bac-to-Bac system (Life Technologies). The bacmid DNA is purified using the Qiagen midiprep purification system (Qiagen) and is used to transfect Sf9 insect cells using cellfectin in Sf900 II SFM medium (Life Technologies) as described in the manufacturer's protocol. Recombinant virus particles are collected 4 days later, titrated by virus plaque assay, and amplified as described by King and Possee, in *The Baculovirus Expression System: A Laboratory Guide*, Chapman and Hall Ltd., London, pp. 111-114, 1992.

High Five cells (1.75×10^7 cells/175-cm² tissue culture flask) are infected with the recombinant viruses containing MASP-2 polypeptides at a multiplicity of infection of 2 in Sf900 II SFM medium at 28°C for 96 h. The supernatants are collected by centrifugation and diisopropyl phosphorofluoridate is added to a final concentration of 1 mM.

The MASP-2 polypeptides are secreted in the culture medium. The culture supernatants are dialyzed against 50 mM NaCl, 1 mM CaCl₂, 50 mM triethanolamine hydrochloride, pH 8.1, and loaded at 1.5 ml/min onto a Q-Sepharose Fast Flow column (Amersham Pharmacia Biotech) (2.8 x 12 cm) equilibrated in the same buffer. Elution is conducted by applying a 1.2 liter linear gradient to 350 mM NaCl in the same buffer. Fractions containing the recombinant MASP-2 polypeptides are identified by Western blot analysis, precipitated by addition of (NH₄)₂SO₄ to 60% (w/v), and left overnight at 4°C. The pellets are resuspended in 145 mM NaCl, 1 mM CaCl₂, 50 mM triethanolamine hydrochloride, pH 7.4, and applied onto a TSK G3000 SWG column (7.5 x 600 mm) (Tosohaas, Montgomeryville, PA) equilibrated in the same buffer. The purified polypeptides are then concentrated to 0.3 mg/ml by ultrafiltration on Microsep microconcentrators (m.w. cut-off = 10,000) (Filtron, Karlstein, Germany).

EXAMPLE 4

This example describes a method of producing polyclonal antibodies against MASP-2 polypeptides.

5 **Materials and Methods:**

MASP-2 Antigens: Polyclonal anti-human MASP-2 antiserum is produced by immunizing rabbits with the following isolated MASP-2 polypeptides: human MASP-2 (SEQ ID NO:6) isolated from serum; recombinant human MASP-2 (SEQ ID NO:6), MASP-2A containing the inactive protease domain (SEQ ID NO:13), as described in
10 **Example 3**; and recombinant CUBI (SEQ ID NO:8), CUBEGFI (SEQ ID NO:9), and CUBEGFCUBII (SEQ ID NO:10) expressed as described above in **Example 3**.

Polyclonal antibodies: Six-week old Rabbits, primed with BCG (bacillus Calmette-Guerin vaccine) are immunized by injecting 100 µg of MASP-2 polypeptide at 100 µg/ml in sterile saline solution. Injections are done every 4 weeks, with antibody
15 titer monitored by ELISA assay as described in **Example 5**. Culture supernatants are collected for antibody purification by protein A affinity chromatography.

EXAMPLE 5

 This example describes a method for producing murine monoclonal antibodies
20 against rat or human MASP-2 polypeptides.

Materials and Methods:

 Male A/J mice (Harlan, Houston, Tex.), 8-12 weeks old, are injected subcutaneously with 100 µg human or rat rMASP-2 or rMASP-2A polypeptides (made as described in **Example 3**) in complete Freund's adjuvant (Difco Laboratories, Detroit,
25 Mich.) in 200 µl of phosphate buffered saline (PBS) pH 7.4. At two-week intervals the mice are twice injected subcutaneously with 50 µg of human or rat rMASP-2 or rMASP-2A polypeptide in incomplete Freund's adjuvant. On the fourth week the mice are injected with 50 µg of human or rat rMASP-2 or rMASP-2A polypeptide in PBS and are fused 4 days later.

30 For each fusion, single cell suspensions are prepared from the spleen of an immunized mouse and used for fusion with Sp2/0 myeloma cells. 5×10^8 of the Sp2/0 and 5×10^8 spleen cells are fused in a medium containing 50% polyethylene glycol (M.W. 1450) (Kodak, Rochester, N.Y.) and 5% dimethylsulfoxide (Sigma Chemical Co.,

St. Louis, Mo.). The cells are then adjusted to a concentration of 1.5×10^5 spleen cells per 200 μ l of the suspension in Iscove medium (Gibco, Grand Island, N.Y.), supplemented with 10% fetal bovine serum, 100 units/ml of penicillin, 100 μ g/ml of streptomycin, 0.1 mM hypoxanthine, 0.4 μ M aminopterin and 16 μ M thymidine. Two hundred
5 microliters of the cell suspension are added to each well of about twenty 96-well microculture plates. After about ten days culture supernatants are withdrawn for screening for reactivity with purified factor MASP-2 in an ELISA assay.

ELISA Assay: Wells of Immulon 2 (Dynatech Laboratories, Chantilly, Va.) microtest plates are coated by adding 50 μ l of purified hMASP-2 at 50 ng/ml or rat
10 rMASP-2 (or rMASP-2A) overnight at room temperature. The low concentration of MASP-2 for coating enables the selection of high-affinity antibodies. After the coating solution is removed by flicking the plate, 200 μ l of BLOTTO (non-fat dry milk) in PBS is added to each well for one hour to block the non-specific sites. An hour later, the wells are then washed with a buffer PBST (PBS containing 0.05% Tween 20). Fifty microliters
15 of culture supernatants from each fusion well is collected and mixed with 50 μ l of BLOTTO and then added to the individual wells of the microtest plates. After one hour of incubation, the wells are washed with PBST. The bound murine antibodies are then detected by reaction with horseradish peroxidase (HRP) conjugated goat anti-mouse IgG (Fc specific) (Jackson ImmunoResearch Laboratories, West Grove, Pa.) and diluted at
20 1:2,000 in BLOTTO. Peroxidase substrate solution containing 0.1% 3,3',5,5' tetramethyl benzidine (Sigma, St. Louis, Mo.) and 0.0003% hydrogen peroxide (Sigma) is added to the wells for color development for 30 minutes. The reaction is terminated by addition of 50 μ l of 2M H_2SO_4 per well. The Optical Density at 450 nm of the reaction mixture is read with a BioTek ELISA Reader (BioTek Instruments, Winooski, Vt.).

25 **MASP-2 Binding Assay:**

Culture supernatants that test positive in the MASP-2 ELISA assay described above can be tested in a binding assay to determine the binding affinity the MASP-2 inhibitory agents have for MASP-2. A similar assay can also be used to determine if the inhibitory agents bind to other antigens in the complement system.

30 Polystyrene microtiter plate wells (96-well medium binding plates, Corning Costar, Cambridge, MA) are coated with MASP-2 (20 ng/100 μ l/well, Advanced Research Technology, San Diego, CA) in phosphate-buffered saline (PBS) pH 7.4 overnight at 4°C. After aspirating the MASP-2 solution, wells are blocked with PBS

containing 1% bovine serum albumin (BSA; Sigma Chemical) for 2 h at room temperature. Wells without MASP-2 coating serve as the background controls. Aliquots of hybridoma supernatants or purified anti-MASP-2 MoAbs, at varying concentrations in blocking solution, are added to the wells. Following a 2 h incubation at room temperature, the wells are extensively rinsed with PBS. MASP-2-bound anti-MASP-2 MoAb is detected by the addition of peroxidase-conjugated goat anti-mouse IgG (Sigma Chemical) in blocking solution, which is allowed to incubate for 1h at room temperature. The plate is rinsed again thoroughly with PBS, and 100 μ l of 3,3',5,5'-tetramethyl benzidine (TMB) substrate (Kirkegaard and Perry Laboratories, Gaithersburg, MD) is added. The reaction of TMB is quenched by the addition of 100 μ l of 1M phosphoric acid, and the plate is read at 450 nm in a microplate reader (SPECTRA MAX 250, Molecular Devices, Sunnyvale, CA).

The culture supernatants from the positive wells are then tested for the ability to inhibit complement activation in a functional assay such as the C4 cleavage assay as described in **Example 2**. The cells in positive wells are then cloned by limiting dilution. The MoAbs are tested again for reactivity with hMASP-2 in an ELISA assay as described above. The selected hybridomas are grown in spinner flasks and the spent culture supernatant collected for antibody purification by protein A affinity chromatography.

20

EXAMPLE 6

This example describes the generation and production of humanized murine anti-MASP-2 antibodies and antibody fragments.

A murine anti-MASP-2 monoclonal antibody is generated in Male A/J mice as described in **Example 5**. The murine antibody is then humanized as described below to reduce its immunogenicity by replacing the murine constant regions with their human counterparts to generate a chimeric IgG and Fab fragment of the antibody, which is useful for inhibiting the adverse effects of MASP-2-dependent complement activation in human subjects in accordance with the present invention.

1. Cloning of anti-MASP-2 variable region genes from murine hybridoma cells. Total RNA is isolated from the hybridoma cells secreting anti-MASP-2 MoAb (obtained as described in Example 7) using RNazol following the manufacturer's protocol (Biotech, Houston, Tex.). First strand cDNA is synthesized from the total RNA using oligo dT as the primer. PCR is performed using the immunoglobulin

constant C region-derived 3' primers and degenerate primer sets derived from the leader peptide or the first framework region of murine V_H or V_K genes as the 5' primers. Anchored PCR is carried out as described by Chen and Platsucas (Chen, P.F., *Scand. J. Immunol.* 35:539-549, 1992). For cloning the V_K gene, double-stranded cDNA is prepared using a NotI-MAK1 primer (5'-TGCGGCCGCTGTAGGTGCTGTCTTT-3' SEQ ID NO:38). Annealed adaptors AD1 (5'-GGAATTCACCTCGTTATTCTCGGA-3' SEQ ID NO:39) and AD2 (5'-TCCGAGAATAACGAGTG-3' SEQ ID NO:40) are ligated to both 5' and 3' termini of the double-stranded cDNA. Adaptors at the 3' ends are removed by NotI digestion. The digested product is then used as the template in PCR with the AD1 oligonucleotide as the 5' primer and MAK2 (5'-CATTGAAAGCTTTGGGGTAGAAGTTGTTC-3' SEQ ID NO:41) as the 3' primer. DNA fragments of approximately 500 bp are cloned into pUC19. Several clones are selected for sequence analysis to verify that the cloned sequence encompasses the expected murine immunoglobulin constant region. The NotI-MAK1 and MAK2 oligonucleotides are derived from the V_K region and are 182 and 84 bp, respectively, downstream from the first base pair of the C kappa gene. Clones are chosen that include the complete V_K and leader peptide.

For cloning the V_H gene, double-stranded cDNA is prepared using the NotI MAG1 primer (5'-CGCGGCCGCGAGCTGCTCAGAGTGTAGA-3' SEQ ID NO:42). Annealed adaptors AD1 and AD2 are ligated to both 5' and 3' termini of the double-stranded cDNA. Adaptors at the 3' ends are removed by NotI digestion. The digested product are used as the template in PCR with the AD1 oligonucleotide and MAG2 (5'-CGGTAAGCTTCACTGGCTCAGGGAAATA-3' SEQ ID NO:43) as primers. DNA fragments of 500 to 600 bp in length are cloned into pUC19. The NotI-MAG1 and MAG2 oligonucleotides are derived from the murine C γ 7.1 region, and are 180 and 93 bp, respectively, downstream from the first bp of the murine C γ 7.1 gene. Clones are chosen that encompass the complete V_H and leader peptide.

2. Construction of Expression Vectors for Chimeric MASP-2 IgG and Fab. The cloned V_H and V_K genes described above are used as templates in a PCR reaction to add the Kozak consensus sequence to the 5' end and the splice donor to the 3' end of the nucleotide sequence. After the sequences are analyzed to confirm the absence of PCR errors, the V_H and V_K genes are inserted into expression vector cassettes

containing human C_γ1 and C_κ respectively, to give pSV2neoV_H-huC_γ1 and pSV2neoV-huC_γ. CsCl gradient-purified plasmid DNAs of the heavy- and light-chain vectors are used to transfect COS cells by electroporation. After 48 hours, the culture supernatant is tested by ELISA to confirm the presence of approximately 200 ng/ml of chimeric IgG. The cells are harvested and total RNA is prepared. First strand cDNA is synthesized from the total RNA using oligo dT as the primer. This cDNA is used as the template in PCR to generate the Fd and kappa DNA fragments. For the Fd gene, PCR is carried out using 5'-AAGAAGCTTGCCGCCACCATGGATTGGCTGTGGAAC-3' (SEQ ID NO:44) as the 5' primer and a CH1-derived 3' primer (5'-CGGGATCCTCAAACCTTCTTGTCCACCTTGG-3' SEQ ID NO:45). The DNA sequence is confirmed to contain the complete V_H and the CH1 domain of human IgG1. After digestion with the proper enzymes, the Fd DNA fragments are inserted at the HindIII and BamHI restriction sites of the expression vector cassette pSV2dhfr-TUS to give pSV2dhfrFd. The pSV2 plasmid is commercially available and consists of DNA segments from various sources: pBR322 DNA (thin line) contains the pBR322 origin of DNA replication (pBR ori) and the lactamase ampicillin resistance gene (Amp); SV40 DNA, represented by wider hatching and marked, contains the SV40 origin of DNA replication (SV40 ori), early promoter (5' to the dhfr and neo genes), and polyadenylation signal (3' to the dhfr and neo genes). The SV40-derived polyadenylation signal (pA) is also placed at the 3' end of the Fd gene.

For the kappa gene, PCR is carried out using 5'-AAGAAAGCTTGCCGCCACCATGTTCTACTAGCTCT-3' (SEQ ID NO:46) as the 5' primer and a C_κ-derived 3' primer (5'-CGGGATCCTTCTCCCTCTAACACTCT-3' SEQ ID NO:47). DNA sequence is confirmed to contain the complete V_κ and human C_κ regions. After digestion with proper restriction enzymes, the kappa DNA fragments are inserted at the HindIII and BamHI restriction sites of the expression vector cassette pSV2neo-TUS to give pSV2neoK. The expression of both Fd and kappa genes are driven by the HCMV-derived enhancer and promoter elements. Since the Fd gene does not include the cysteine amino acid residue involved in the inter-chain disulfide bond, this recombinant chimeric Fab contains non-covalently linked heavy- and light-chains. This chimeric Fab is designated as cFab.

To obtain recombinant Fab with an inter-heavy and light chain disulfide bond, the above Fd gene may be extended to include the coding sequence for additional 9 amino acids (EPKSCDKTH SEQ ID NO:48) from the hinge region of human IgG1. The BstEII-BamHI DNA segment encoding 30 amino acids at the 3' end of the Fd gene may
5 be replaced with DNA segments encoding the extended Fd, resulting in pSV2dhfrFd/9aa.

3. Expression and Purification of Chimeric Anti-MASP-2 IgG

To generate cell lines secreting chimeric anti-MASP-2 IgG, NSO cells are transfected with purified plasmid DNAs of pSV2neoV_H-huC_γ1 and pSV2neoV-huC
10 kappa by electroporation. Transfected cells are selected in the presence of 0.7 mg/ml G418. Cells are grown in a 250 ml spinner flask using serum-containing medium.

Culture supernatant of 100 ml spinner culture is loaded on a 10-ml PROSEP-A column (Bioprocessing, Inc., Princeton, N.J.). The column is washed with 10 bed volumes of PBS. The bound antibody is eluted with 50 mM citrate buffer, pH 3.0. Equal
15 volume of 1 M Hepes, pH 8.0 is added to the fraction containing the purified antibody to adjust the pH to 7.0. Residual salts are removed by buffer exchange with PBS by Millipore membrane ultrafiltration (M.W. cut-off: 3,000). The protein concentration of the purified antibody is determined by the BCA method (Pierce).

4. Expression and purification of chimeric anti-MASP-2 Fab

To generate cell lines secreting chimeric anti-MASP-2 Fab, CHO cells are
20 transfected with purified plasmid DNAs of pSV2dhfrFd (or pSV2dhfrFd/9aa) and pSV2neokappa, by electroporation. Transfected cells are selected in the presence of G418 and methotrexate. Selected cell lines are amplified in increasing concentrations of methotrexate. Cells are single-cell subcloned by limiting dilution. High-producing single-cell subcloned cell lines are then grown in 100 ml spinner culture using serum-free
25 medium.

Chimeric anti-MASP-2 Fab is purified by affinity chromatography using a mouse anti-idiotypic MoAb to the MASP-2 MoAb. An anti-idiotypic MASP-2 MoAb can be made by immunizing mice with a murine anti-MASP-2 MoAb conjugated with keyhole limpet hemocyanin (KLH) and screening for specific MoAb binding that can be
30 competed with human MASP-2. For purification, 100 ml of supernatant from spinner cultures of CHO cells producing cFab or cFab/9aa are loaded onto the affinity column coupled with an anti-idiotypic MASP-2 MoAb. The column is then washed thoroughly with PBS before the bound Fab is eluted with 50 mM diethylamine, pH 11.5. Residual

salts are removed by buffer exchange as described above. The protein concentration of the purified Fab is determined by the BCA method (Pierce).

The ability of the chimeric MASP-2 IgG, cFab, and cFab/9aa to inhibit MASP-2-dependent complement pathways may be determined by using the inhibitory assays described in **Example 2 or Example 7**.

EXAMPLE 7

This example describes an *in vitro* C4 cleavage assay used as a functional screen to identify MASP-2 inhibitory agents capable of blocking MASP-2-dependent complement activation via L-ficolin/P35, H-ficolin, M-ficolin or mannan.

C4 Cleavage Assay: A C4 cleavage assay has been described by Petersen, S.V., et al., *J. Immunol. Methods* 257:107, 2001, which measures lectin pathway activation resulting from lipoteichoic acid (LTA) from *S. aureus* which binds L-ficolin.

Reagents: Formalin-fixed *S. aureus* (DSM20233) is prepared as follows: bacteria is grown overnight at 37°C in tryptic soy blood medium, washed three times with PBS, then fixed for 1 h at room temperature in PBS/0.5% formalin, and washed a further three times with PBS, before being resuspended in coating buffer (15 mM Na₂CO₃, 35 mM NaHCO₃, pH 9.6).

Assay: The wells of a Nunc MaxiSorb microtiter plate (Nalgene Nunc International, Rochester, NY) are coated with: 100 µl of formalin-fixed *S. aureus* DSM20233 (OD₅₅₀ = 0.5) in coating buffer with 1 µg of L-ficolin in coating buffer. After overnight incubation, wells are blocked with 0.1% human serum albumin (HSA) in TBS (10 mM Tris-HCl, 140 mM NaCl, pH 7.4), then are washed with TBS containing 0.05% Tween 20 and 5 mM CaCl₂ (wash buffer). Human serum samples are diluted in 20 mM Tris-HCl, 1 M NaCl, 10 mM CaCl₂, 0.05% Triton X-100, 0.1% HSA, pH 7.4, which prevents activation of endogenous C4 and dissociates the C1 complex (composed of C1q, C1r and C1s). MASP-2 inhibitory agents, including anti-MASP-2 MoAbs and inhibitory peptides are added to the serum samples in varying concentrations. The diluted samples are added to the plate and incubated overnight at 4°C. After 24 hours, the plates are washed thoroughly with wash buffer, then 0.1 µg of purified human C4 (obtained as described in Dodds, A.W., *Methods Enzymol.* 223:46, 1993) in 100 µl of 4 mM barbital, 145 mM NaCl, 2 mM CaCl₂, 1 mM MgCl₂, pH 7.4 is added to each well. After 1.5 h at

37°C, the plates are washed again and C4b deposition is detected using alkaline phosphatase-conjugated chicken anti-human C4c (obtained from Immunsytem, Uppsala, Sweden) and measured using the colorimetric substrate p-nitrophenyl phosphate.

5 **C4 Assay on mannan:** The assay described above is adapted to measure lectin pathway activation via MBL by coating the plate with LSP and mannan prior to adding serum mixed with various MASP-2 inhibitory agents.

C4 assay on H-ficolin (Hakata Ag): The assay described above is adapted to measure lectin pathway activation via H-ficolin by coating the plate with LPS and H-ficolin prior to adding serum mixed with various MASP-2 inhibitory agents.

10

EXAMPLE 8

The following assay demonstrates the presence of classical pathway activation in wild-type and MASP-2^{-/-} mice.

Methods: Immune complexes were generated *in situ* by coating microtiter plates
15 (Maxisorb, Nunc, cat. No. 442404, Fisher Scientific) with 0.1% human serum albumin in 10 mM Tris, 140 mM NaCl, pH 7.4 for 1 hours at room temperature followed by overnight incubation at 4°C with sheep anti whole serum antiserum (Scottish Antibody Production Unit, Carlisle, Scotland) diluted 1:1000 in TBS/tween/Ca²⁺. Serum samples were obtained from wild-type and MASP-2^{-/-} mice and added to the coated plates.
20 Control samples were prepared in which C1q was depleted from wild-type and MASP-2^{-/-} serum samples. C1q-depleted mouse serum was prepared using protein-A-coupled Dynabeads (Dynal Biotech, Oslo, Norway) coated with rabbit anti-human C1q IgG (Dako, Glostrup, Denmark), according to the supplier's instructions. The plates were incubated for 90 minutes at 37°C. Bound C3b was detected with a
25 polyclonal anti-human-C3c Antibody (Dako A 062) diluted in TBS/tw/ Ca⁺⁺ at 1:1000. The secondary antibody is goat anti-rabbit IgG.

Results: **FIGURE 7** shows the relative C3b deposition levels on plates coated with IgG in wild-type serum, MASP-2^{-/-} serum, C1q-depleted wild-type and C1q-depleted MASP-2^{-/-} serum. These results demonstrate that the classical pathway is
30 intact in the MASP-2^{-/-} mouse strain.

EXAMPLE 9

The following assay is used to test whether a MASP-2 inhibitory agent blocks the classical pathway by analyzing the effect of a MASP-2 inhibitory agent under conditions in which the classical pathway is initiated by immune complexes.

5 **Methods:** To test the effect of a MASP-2 inhibitory agent on conditions of complement activation where the classical pathway is initiated by immune complexes, triplicate 50 µl samples containing 90% NHS are incubated at 37°C in the presence of 10 µg/ml immune complex (IC) or PBS, and parallel triplicate samples (+/-IC) are also included which contain 200 nM anti-properdin monoclonal antibody during the 37°C
10 incubation. After a two hour incubation at 37°C, 13 mM EDTA is added to all samples to stop further complement activation and the samples are immediately cooled to 5°C. The samples are then stored at -70°C prior to being assayed for complement activation products (C3a and sC5b-9) using ELISA kits (Quidel, Catalog Nos. A015 and A009) following the manufacturer's instructions.

15

EXAMPLE 10

This example describes the identification of high affinity anti-MASP-2 Fab2 antibody fragments that block MASP-2 activity.

20 **Background and rationale:** MASP-2 is a complex protein with many separate functional domains, including: binding site(s) for MBL and ficolins, a serine protease catalytic site, a binding site for proteolytic substrate C2, a binding site for proteolytic substrate C4, a MASP-2 cleavage site for autoactivation of MASP-2 zymogen, and two Ca⁺⁺ binding sites. Fab2 antibody fragments were identified that bind with high affinity to MASP-2, and the identified Fab2 fragments were tested in a functional assay to
25 determine if they were able to block MASP-2 functional activity.

To block MASP-2 functional activity, an antibody or Fab2 antibody fragment must bind and interfere with a structural epitope on MASP-2 that is required for MASP-2 functional activity. Therefore, many or all of the high affinity binding anti-MASP-2 Fab2s may not inhibit MASP-2 functional activity unless they bind to structural epitopes
30 on MASP-2 that are directly involved in MASP-2 functional activity.

A functional assay that measures inhibition of lectin pathway C3 convertase formation was used to evaluate the "blocking activity" of anti-MASP-2 Fab2s. It is known that the primary physiological role of MASP-2 in the lectin pathway is to generate

the next functional component of the lectin-mediated complement pathway, namely the lectin pathway C3 convertase. The lectin pathway C3 convertase is a critical enzymatic complex (C4bC2a) that proteolytically cleaves C3 into C3a and C3b. MASP-2 is not a structural component of the lectin pathway C3 convertase (C4bC2a); however, MASP-2 functional activity is required in order to generate the two protein components (C4b, C2a) that comprise the lectin pathway C3 convertase. Furthermore, all of the separate functional activities of MASP-2 listed above appear to be required in order for MASP-2 to generate the lectin pathway C3 convertase. For these reasons, a preferred assay to use in evaluating the "blocking activity" of anti-MASP-2 Fab2s is believed to be a functional assay that measures inhibition of lectin pathway C3 convertase formation.

Generation of High Affinity Fab2s: A phage display library of human variable light and heavy chain antibody sequences and automated antibody selection technology for identifying Fab2s that react with selected ligands of interest was used to create high affinity Fab2s to rat MASP-2 protein (SEQ ID NO:55). A known amount of rat MASP-2 (~1 mg, >85% pure) protein was utilized for antibody screening. Three rounds of amplification were utilized for selection of the antibodies with the best affinity. Approximately 250 different hits expressing antibody fragments were picked for ELISA screening. High affinity hits were subsequently sequenced to determine uniqueness of the different antibodies.

Fifty unique anti-MASP-2 antibodies were purified and 250 µg of each purified Fab2 antibody was used for characterization of MASP-2 binding affinity and complement pathway functional testing, as described in more detail below.

Assays used to Evaluate the Inhibitory (blocking) Activity of Anti-MASP-2 Fab2s

1. Assay to Measure Inhibition of Formation of Lectin Pathway C3 Convertase:

Background: The lectin pathway C3 convertase is the enzymatic complex (C4bC2a) that proteolytically cleaves C3 into the two potent proinflammatory fragments, anaphylatoxin C3a and opsonic C3b. Formation of C3 convertase appears to be a key step in the lectin pathway in terms of mediating inflammation. MASP-2 is not a structural component of the lectin pathway C3 convertase (C4bC2a); therefore anti-MASP-2 antibodies (or Fab2) will not directly inhibit activity of preexisting C3 convertase. However, MASP-2 serine protease activity is required in order to generate the two protein

components (C4b, C2a) that comprise the lectin pathway C3 convertase. Therefore, anti-MASP-2 Fab2 which inhibit MASP-2 functional activity (i.e., blocking anti-MASP-2 Fab2) will inhibit *de novo* formation of lectin pathway C3 convertase. C3 contains an unusual and highly reactive thioester group as part of its structure. Upon cleavage of C3 by C3 convertase in this assay, the thioester group on C3b can form a covalent bond with hydroxyl or amino groups on macromolecules immobilized on the bottom of the plastic wells via ester or amide linkages, thus facilitating detection of C3b in the ELISA assay.

Yeast mannan is a known activator of the lectin pathway. In the following method to measure formation of C3 convertase, plastic wells coated with mannan were incubated for 30 min at 37°C with diluted rat serum to activate the lectin pathway. The wells were then washed and assayed for C3b immobilized onto the wells using standard ELISA methods. The amount of C3b generated in this assay is a direct reflection of the *de novo* formation of lectin pathway C3 convertase. Anti-MASP-2 Fab2s at selected concentrations were tested in this assay for their ability to inhibit C3 convertase formation and consequent C3b generation.

Methods:

96-well Costar Medium Binding plates were incubated overnight at 5°C with mannan diluted in 50 mM carbonate buffer, pH 9.5 at 1 ug/50 TI/well. After overnight incubation, each well was washed three times with 200 TI PBS. The wells were then blocked with 100 TI/well of 1% bovine serum albumin in PBS and incubated for one hour at room temperature with gentle mixing. Each well was then washed three times with 200 TI of PBS. The anti-MASP-2 Fab2 samples were diluted to selected concentrations in Ca^{++} and Mg^{++} containing GVB buffer (4.0 mM barbitol, 141 mM NaCl, 1.0 mM MgCl_2 , 2.0 mM CaCl_2 , 0.1% gelatin, pH 7.4) at 5 C. A 0.5% rat serum was added to the above samples at 5 C and 100 TI was transferred to each well. Plates were covered and incubated for 30 minutes in a 37 C waterbath to allow complement activation. The reaction was stopped by transferring the plates from the 37 C waterbath to a container containing an ice-water mix. Each well was washed five times with 200 TI with PBS-Tween 20 (0.05% Tween 20 in PBS), then washed two times with 200 TI PBS. A 100 TI/well of 1:10,000 dilution of the primary antibody (rabbit anti-human C3c, DAKO A0062) was added in PBS containing 2.0 mg/ml bovine serum albumin and incubated 1 hr at room temperature with gentle mixing. Each well was washed 5 x 200 TI PBS.

TI/well of 1:10,000 dilution of the secondary antibody (peroxidase-conjugated goat anti-rabbit IgG, American Qualex A102PU) was added in PBS containing 2.0 mg/ml bovine serum albumin and incubated for one hour at room temperature on a shaker with gentle mixing. Each well was washed five times with 200 TI with PBS. 100 TI/well of the peroxidase substrate TMB (Kirkegaard & Perry Laboratories) was added and incubated at room temperature for 10 min. The peroxidase reaction was stopped by adding 100 TI/well of 1.0 M H₃PO₄ and the OD₄₅₀ was measured.

2. Assay to Measure Inhibition of MASP-2-dependent C4 Cleavage

Background: The serine protease activity of MASP-2 is highly specific and only two protein substrates for MASP-2 have been identified; C2 and C4. Cleavage of C4 generates C4a and C4b. Anti-MASP-2 Fab2 may bind to structural epitopes on MASP-2 that are directly involved in C4 cleavage (e.g., MASP-2 binding site for C4; MASP-2 serine protease catalytic site) and thereby inhibit the C4 cleavage functional activity of MASP-2.

Yeast mannan is a known activator of the lectin pathway. In the following method to measure the C4 cleavage activity of MASP-2, plastic wells coated with mannan were incubated for 30 minutes at 37 C with diluted rat serum to activate the lectin pathway. Since the primary antibody used in this ELISA assay only recognizes human C4, the diluted rat serum was also supplemented with human C4 (1.0 Tg/ml). The wells were then washed and assayed for human C4b immobilized onto the wells using standard ELISA methods. The amount of C4b generated in this assay is a measure of MASP-2 dependent C4 cleavage activity. Anti-MASP-2 Fab2 at selected concentrations were tested in this assay for their ability to inhibit C4 cleavage.

Methods: 96-well Costar Medium Binding plates were incubated overnight at 5 C with mannan diluted in 50 mM carbonate buffer, pH 9.5 at 1.0 Tg/50 TI/well. Each well was washed 3X with 200 TI PBS. The wells were then blocked with 100 TI/well of 1% bovine serum albumin in PBS and incubated for one hour at room temperature with gentle mixing. Each well was washed 3X with 200 TI of PBS. Anti-MASP-2 Fab2 samples were diluted to selected concentrations in Ca⁺⁺ and Mg⁺⁺ containing GVB buffer (4.0 mM barbital, 141 mM NaCl, 1.0 mM MgCl₂, 2.0 mM CaCl₂, 0.1% gelatin, pH 7.4) at 5 C. 1.0 Tg/ml human C4 (Quidel) was also included in these samples. 0.5% rat serum was added to the above samples at 5 C and 100 TI was transferred to each well.

The plates were covered and incubated for 30 min in a 37 C waterbath to allow complement activation. The reaction was stopped by transferring the plates from the 37 C waterbath to a container containing an ice-water mix. Each well was washed 5 x 200 Tl with PBS-Tween 20 (0.05% Tween 20 in PBS), then each well was washed 5 with 2X with 200 Tl PBS. 100 Tl/well of 1:700 dilution of biotin-conjugated chicken anti-human C4c (Immunsystem AB, Uppsala, Sweden) was added in PBS containing 2.0 mg/ml bovine serum albumin (BSA) and incubated one hour at room temperature with gentle mixing. Each well was washed 5 x 200 Tl PBS. 100 Tl/well of 0.1 Tg/ml of peroxidase-conjugated streptavidin (Pierce Chemical #21126) was added in PBS containing 2.0 mg/ml BSA and incubated for one hour at room temperature on a shaker with gentle mixing. Each well was washed 5 x 200 Tl with PBS. 100 Tl/well of the peroxidase substrate TMB (Kirkegaard & Perry Laboratories) was added and incubated at room temperature for 16 min. The peroxidase reaction was stopped by adding 100 Tl/well of 1.0 M H₃PO₄ and the OD₄₅₀ was measured.

3. Binding Assay of anti-rat MASP-2 Fab2 to 'Native' rat MASP-2

Background: MASP-2 is usually present in plasma as a MASP-2 dimer complex that also includes specific lectin molecules (mannose-binding protein (MBL) and ficolins). Therefore, if one is interested in studying the binding of anti-MASP-2 Fab2 to the physiologically relevant form of MASP-2, it is important to develop a binding assay in which the interaction between the Fab2 and 'native' MASP-2 in plasma is used, rather than purified recombinant MASP-2. In this binding assay the 'native' MASP-2-MBL complex from 10% rat serum was first immobilized onto mannan-coated wells. The binding affinity of various anti-MASP-2 Fab2s to the immobilized 'native' MASP-2 was then studied using a standard ELISA methodology.

Methods: 96-well Costar High Binding plates were incubated overnight at 5°C with mannan diluted in 50 mM carbonate buffer, pH 9.5 at 1 Tg/50 Tl/well. Each well was washed 3X with 200 Tl PBS. The wells were blocked with 100 Tl/well of 0.5% nonfat dry milk in PBST (PBS with 0.05% Tween 20) and incubated for one hour at room temperature with gentle mixing. Each well was washed 3X with 200 Tl of TBS/Tween/Ca⁺⁺ Wash Buffer (Tris-buffered saline, 0.05% Tween 20, containing 5.0 mM CaCl₂, pH 7.4. 10% rat serum in High Salt Binding Buffer (20 mM Tris, 1.0 M NaCl, 10 mM CaCl₂, 0.05% Triton-X100, 0.1% (w/v) bovine serum albumin, pH 7.4)

was prepared on ice. 100 μ l/well was added and incubated overnight at 5°C. Wells were washed 3X with 200 μ l of TBS/Tween/ Ca^{++} Wash Buffer. Wells were then washed 2X with 200 μ l PBS. 100 μ l/well of selected concentration of anti-MASP-2 Fab2 diluted in Ca^{++} and Mg^{++} containing GVB Buffer (4.0 mM barbital, 141 mM NaCl, 1.0 mM MgCl_2 , 2.0 mM CaCl_2 , 0.1% gelatin, pH 7.4) was added and incubated for one hour at room temperature with gentle mixing. Each well was washed 5 x 200 μ l PBS. 100 μ l/well of HRP-conjugated goat anti-Fab2 (Biogenesis Cat No 0500-0099) diluted 1:5000 in 2.0 mg/ml bovine serum albumin in PBS was added and incubated for one hour at room temperature with gentle mixing. Each well was washed 5 x 200 μ l PBS. 100 μ l/well of the peroxidase substrate TMB (Kirkegaard & Perry Laboratories) was added and incubated at room temperature for 70 min. The peroxidase reaction was stopped by adding 100 μ l/well of 1.0 M H_3PO_4 and OD_{450} was measured.

RESULTS:

Approximately 250 different Fab2s that reacted with high affinity to the rat MASP-2 protein were picked for ELISA screening. These high affinity Fab2s were sequenced to determine the uniqueness of the different antibodies, and 50 unique anti-MASP-2 antibodies were purified for further analysis. 250 μ g of each purified Fab2 antibody was used for characterization of MASP-2 binding affinity and complement pathway functional testing. The results of this analysis is shown below in **TABLE 6**.

TABLE 6: ANTI-MASP-2 FAB2 THAT BLOCK LECTIN PATHWAY
COMPLEMENT ACTIVATION

Fab2 antibody #	C3 Convertase (IC_{50} (nM))	K_d	C4 Cleavage IC_{50} (nM)
88	0.32	4.1	ND
41	0.35	0.30	0.81
11	0.46	0.86	<2 nM
86	0.53	1.4	ND
81	0.54	2.0	ND
66	0.92	4.5	ND
57	0.95	3.6	<2 nM
40	1.1	7.2	0.68
58	1.3	2.6	ND
60	1.6	3.1	ND

Fab2 antibody #	C3 Convertase (IC ₅₀ (nM))	K _d	C4 Cleavage IC ₅₀ (nM)
52	1.6	5.8	<2 nM
63	2.0	6.6	ND
49	2.8	8.5	<2 nM
89	3.0	2.5	ND
71	3.0	10.5	ND
87	6.0	2.5	ND
67	10.0	7.7	ND

As shown above in **TABLE 6**, of the 50 anti-MASP-2 Fab2s tested, seventeen Fab2s were identified as MASP-2 blocking Fab2 that potently inhibit C3 convertase formation with IC₅₀ equal to or less than 10 nM Fab2s (a 34% positive hit rate). Eight of the seventeen Fab2s identified have IC₅₀s in the subnanomolar range. Furthermore, all seventeen of the MASP-2 blocking Fab2s shown in **TABLE 6** gave essentially complete inhibition of C3 convertase formation in the lectin pathway C3 convertase assay. **FIGURE 8A** graphically illustrates the results of the C3 convertase formation assay for Fab2 antibody #11, which is representative of the other Fab2 antibodies tested, the results of which are shown in **TABLE 6**. This is an important consideration, since it is theoretically possible that a "blocking" Fab2 may only fractionally inhibit MASP-2 function even when each MASP-2 molecule is bound by the Fab2.

Although mannan is a known activator of the lectin pathway, it is theoretically possible that the presence of anti-mannan antibodies in the rat serum might also activate the classical pathway and generate C3b via the classical pathway C3 convertase. However, each of the seventeen blocking anti-MASP-2 Fab2s listed in this example potently inhibits C3b generation (>95 %), thus demonstrating the specificity of this assay for lectin pathway C3 convertase.

Binding assays were also performed with all seventeen of the blocking Fab2s in order to calculate an apparent K_d for each. The results of the binding assays of anti-rat MASP-2 Fab2s to native rat MASP-2 for six of the blocking Fab2s are also shown in **TABLE 6**. **FIGURE 8B** graphically illustrates the results of a binding assay with the Fab2 antibody #11. Similar binding assays were also carried out for the other Fab2s, the results of which are shown in **TABLE 6**. In general, the apparent K_ds obtained for

binding of each of the six Fab2s to 'native' MASP-2 corresponds reasonably well with the IC_{50} for the Fab2 in the C3 convertase functional assay. There is evidence that MASP-2 undergoes a conformational change from an 'inactive' to an 'active' form upon activation of its protease activity (Feinberg et al., *EMBO J* 22:2348-59 (2003); Gal et al., *J. Biol. Chem.* 280:33435-44 (2005)). In the normal rat plasma used in the C3 convertase formation assay, MASP-2 is present primarily in the 'inactive' zymogen conformation. In contrast, in the binding assay, MASP-2 is present as part of a complex with MBL bound to immobilized mannan; therefore, the MASP-2 would be in the 'active' conformation (Petersen et al., *J. Immunol Methods* 257:107-16, 2001). Consequently, one would not necessarily expect an exact correspondence between the IC_{50} and K_d for each of the seventeen blocking Fab2 tested in these two functional assays since in each assay the Fab2 would be binding a different conformational form of MASP-2. Never-the-less, with the exception of Fab2 #88, there appears to be a reasonably close correspondence between the IC_{50} and apparent K_d for each of the other sixteen Fab2 tested in the two assays (see **TABLE 6**).

Several of the blocking Fab2s were evaluated for inhibition of MASP-2 mediated cleavage of C4. **FIGURE 8C** graphically illustrates the results of a C4 cleavage assay, showing inhibition with Fab2 #41, with an $IC_{50}=0.81$ nM (see **TABLE 6**). As shown in **FIGURE 9**, all of the Fab2s tested were found to inhibit C4 cleavage with IC_{50} s similar to those obtained in the C3 convertase assay (see **TABLE 6**).

Although mannan is a known activator of the lectin pathway, it is theoretically possible that the presence of anti-mannan antibodies in the rat serum might also activate the classical pathway and thereby generate C4b by C1s-mediated cleavage of C4. However, several anti-MASP-2 Fab2s have been identified which potently inhibit C4b generation (>95 %), thus demonstrating the specificity of this assay for MASP-2 mediated C4 cleavage. C4, like C3, contains an unusual and highly reactive thioester group as part of its structure. Upon cleavage of C4 by MASP-2 in this assay, the thioester group on C4b can form a covalent bond with hydroxyl or amino groups on macromolecules immobilized on the bottom of the plastic wells *via* ester or amide linkages, thus facilitating detection of C4b in the ELISA assay.

These studies clearly demonstrate the creation of high affinity FAB2s to rat MASP-2 protein that functionally block both C4 and C3 convertase activity, thereby preventing lectin pathway activation.

5

EXAMPLE 11

This Example describes the epitope mapping for several of the blocking anti-rat MASP-2 Fab2 antibodies that were generated as described in **Example 10**.

Methods:

As shown in **FIGURE 10**, the following proteins, all with N-terminal 6X His tags
10 were expressed in CHO cells using the pED4 vector:

rat MASP-2A, a full length MASP-2 protein, inactivated by altering the serine at the active center to alanine (S613A);

rat MASP-2K, a full-length MASP-2 protein altered to reduce autoactivation (R424K);

15 CUBI-II, an N-terminal fragment of rat MASP-2 that contains the CUBI, EGF-like and CUBII domains only; and

CUBI/EGF-like, an N-terminal fragment of rat MASP-2 that contains the CUBI and EGF-like domains only.

These proteins were purified from culture supernatants by nickel-affinity
20 chromatography, as previously described (Chen et al., *J. Biol. Chem.* 276:25894-02 (2001)).

A C-terminal polypeptide (CCPII-SP), containing CCPII and the serine protease domain of rat MASP-2, was expressed in *E. coli* as a thioredoxin fusion protein using pTrxFus (Invitrogen). Protein was purified from cell lysates using Thiobond affinity
25 resin. The thioredoxin fusion partner was expressed from empty pTrxFus as a negative control.

All recombinant proteins were dialyzed into TBS buffer and their concentrations determined by measuring the OD at 280 nm.

DOT BLOT ANALYSIS:

30 Serial dilutions of the five recombinant MASP-2 polypeptides described above and shown in **FIGURE 10** (and the thioredoxin polypeptide as a negative control for CCPII-serine protease polypeptide) were spotted onto a nitrocellulose membrane. The amount of protein spotted ranged from 100 ng to 6.4 pg, in five-fold steps. In later

experiments, the amount of protein spotted ranged from 50 ng down to 16 pg, again in five-fold steps. Membranes were blocked with 5% skimmed milk powder in TBS (blocking buffer) then incubated with 1.0 µg/ml anti-MASP-2 Fab2s in blocking buffer (containing 5.0 mM Ca^{2+}). Bound Fab2s were detected using HRP-conjugated anti-human Fab (AbD/Serotec; diluted 1/10,000) and an ECL detection kit (Amersham). One membrane was incubated with polyclonal rabbit-anti human MASP-2 Ab (described in Stover et al., *J Immunol* 163:6848-59 (1999)) as a positive control. In this case, bound Ab was detected using HRP-conjugated goat anti-rabbit IgG (Dako; diluted 1/2,000).

MASP-2 Binding Assay

ELISA plates were coated with 1.0 µg/well of recombinant MASP-2A or CUBI-II polypeptide in carbonate buffer (pH 9.0) overnight at 4°C. Wells were blocked with 1% BSA in TBS, then serial dilutions of the anti-MASP-2 Fab2s were added in TBS containing 5.0 mM Ca^{2+} . The plates were incubated for one hour at RT. After washing three times with TBS/tween/ Ca^{2+} , HRP-conjugated anti-human Fab (AbD/Serotec) diluted 1/10,000 in TBS/ Ca^{2+} was added and the plates incubated for a further one hour at RT. Bound antibody was detected using a TMB peroxidase substrate kit (Biorad).

RESULTS:

Results of the dot blot analysis demonstrating the reactivity of the Fab2s with various MASP-2 polypeptides are provided below in **TABLE 7**. The numerical values provided in **TABLE 7** indicate the amount of spotted protein required to give approximately half-maximal signal strength. As shown, all of the polypeptides (with the exception of the thioredoxin fusion partner alone) were recognized by the positive control Ab (polyclonal anti-human MASP-2 sera, raised in rabbits).

TABLE 7: REACTIVITY WITH VARIOUS RECOMBINANT RAT MASP-2 POLYPEPTIDES ON DOT BLOTS

Fab2 Antibody #	MASP-2A	CUBI-II	CUBI/EGF-like	CCPII-SP	Thioredoxin
40	0.16 ng	NR	NR	0.8 ng	NR
41	0.16 ng	NR	NR	0.8 ng	NR
11	0.16 ng	NR	NR	0.8 ng	NR
49	0.16 ng	NR	NR	>20 ng	NR
52	0.16 ng	NR	NR	0.8 ng	NR

Fab2 Antibody #	MASP-2A	CUBI-II	CUBI/EGF-like	CCPII-SP	Thioredoxin
57	0.032 ng	NR	NR	NR	NR
58	0.4 ng	NR	NR	2.0 ng	NR
60	0.4 ng	0.4 ng	NR	NR	NR
63	0.4 ng	NR	NR	2.0 ng	NR
66	0.4 ng	NR	NR	2.0 ng	NR
67	0.4 ng	NR	NR	2.0 ng	NR
71	0.4 ng	NR	NR	2.0 ng	NR
81	0.4 ng	NR	NR	2.0 ng	NR
86	0.4 ng	NR	NR	10 ng	NR
87	0.4 ng	NR	NR	2.0 ng	NR
Positive Control	<0.032 ng	0.16 ng	0.16 ng	<0.032 ng	NR

pathway activity was observed over the second and third weeks, with complete lectin pathway restoration in the mice by 17 days post anti-MASP-2 MoAb administration. NR = No reaction. The positive control antibody is polyclonal anti-human MASP-2 sera, raised in rabbits.

5 All of the Fab2s reacted with MASP-2A as well as MASP-2K (data not shown). The majority of the Fab2s recognized the CCPII-SP polypeptide but not the N-terminal fragments. The two exceptions are Fab2 #60 and Fab2 #57. Fab2 #60 recognizes MASP-2A and the CUBI-II fragment, but not the CUBI/EGF-like polypeptide or the CCPII-SP polypeptide, suggesting it binds to an epitope in CUBII, or spanning the CUBII
10 and the EGF-like domain. Fab2 # 57 recognizes MASP-2A but not any of the MASP-2 fragments tested, indicating that this Fab2 recognizes an epitope in CCP1. Fab2 #40 and #49 bound only to complete MASP-2A. In the ELISA binding assay shown in **FIGURE 11**, Fab2 #60 also bound to the CUBI-II polypeptide, albeit with a slightly lower apparent affinity.

15 These finding demonstrate the identification of unique blocking Fab2s to multiple regions of the MASP-2 protein

EXAMPLE 12

This Example describes the analysis of MASP-2^{-/-} mice in a Murine Renal
20 Ischemia/Reperfusion Model.

Background/Rationale: Ischemia-Reperfusion (I/R) injury in kidney at body temperature has relevance in a number of clinical conditions, including hypovolaemic shock, renal artery occlusion and cross-clamping procedures.

Kidney ischemia-reperfusion (I/R) is an important cause of acute renal failure, associated with a mortality rate of up to 50% (Levy et al., *JAMA* 275:1489-94, 1996; Thadhani et al., *N. Engl. J. Med.* 334:1448-60, 1996). Post-transplant renal failure is a common and threatening complication after renal transplantation (Nicholson et al., *Kidney Int.* 58:2585-91, 2000). Effective treatment for renal I/R injury is currently not available and hemodialysis is the only treatment available. The pathophysiology of renal I/R injury is complicated. Recent studies have shown that the lectin pathway of complement activation may have an important role in the pathogenesis of renal I/R injury (deVries et al., *Am. J. Path.* 165:1677-88, 2004).

Methods:

A MASP-2(-/-) mouse was generated as described in **Example 1** and backcrossed for at least 10 generations with C57Bl/6. Six male MASP-2(-/-) and six wildtype (+/+) mice weighing between 22-25 g were administered an intraperitoneal injection of Hypnovel (6.64 mg/kg; Roche products Ltd. Welwyn Garden City, UK), and subsequently anaesthetized by inhalation of isoflurane (Abbott Laboratories Ltd., Kent, UK). Isoflurane was chosen because it is a mild inhalation anaesthetic with minimal liver toxicity; the concentrations are produced accurately and the animal recovers rapidly, even after prolonged anaesthesia. Hypnovel was administered because it produces a condition of neuroleptanalgesia in the animal and means that less isoflurane needs to be administered. A warm pad was placed beneath the animal in order to maintain a constant body temperature. Next, a midline abdominal incision was performed and the body cavity held open using a pair of retractors. Connective tissue was cleared above and below the renal vein and artery of both right and left kidneys, and the renal pedicle was clamped via the application of microaneurysm clamps for a period of 55 minutes. This period of ischemia was based initially on a previous study performed in this laboratory (Zhou et al., *J. Clin. Invest.* 105:1363-71 (2000)). In addition, a standard ischemic time of 55 minutes was chosen following ischemic titration and it was found that 55 minutes gave consistent injury that was also reversible, with low mortality, less than 5%. After occlusion, 0.4 ml of warm saline (37°C) was placed in the abdominal cavity and then the abdomen was closed for the period of ischemia. Following removal

of the microaneurysm clamps, the kidneys were observed until color change, an indication of blood re-flow to the kidneys. A further 0.4 ml of warm saline was placed in the abdominal cavity and the opening was sutured, whereupon animals were returned to their cages. Tail blood samples were taken at 24 hours after removing the clamps, and at 5 48 hours the mice were sacrificed and an additional blood sample was collected.

Assessment of Renal Injury: Renal function was assessed at 24 and 48 hours after reperfusion in six male MASP-2(-/-) and six WT (+/+) mice. Blood creatinine measurement was determined by mass spectrometry, which provides a reproducible index of renal function (sensitivity < 1.0 $\mu\text{mol/L}$). **FIGURE 12** graphically illustrates the 10 blood urea nitrogen clearance for wildtype C57Bl/6 controls and MASP-2 (-/-) at 24 hours and 48 hours after reperfusion. As shown in **FIGURE 12**, MASP-2(-/-) mice displayed a significant reduction in the amount of blood urea at 24 and 48 hours, in comparison to wildtype control mice, indicating a protective functional effect from renal damage in the ischemia reperfusion injury model.

15 Overall, increased blood urea was seen in both the WT (+/+) and MASP-2 (-/-) mice at 24 and 48 hours following the surgical procedure and ischemic insult. Levels of blood urea in a non-ischemic WT (+/+) surgery animal was separately determined to be 5.8 mmol/L. In addition to the data presented in **FIGURE 12**, one MASP-2 (-/-) animal showed nearly complete protection from the ischemic insult, with values of 6.8 and 20 9.6 mmol/L at 24 and 48 hours, respectively. This animal was excluded from the group analysis as a potential outlier, wherein no ischemic injury may have been present. Therefore, the final analysis shown in **FIGURE 12** included 5 MASP-2(-/-) mice and 6 WT (+/+) mice and a statistically significant reduction in blood urea was seen at 24 and 48 hours in the MASP-2 (-/-) mice (Student t-test $p < 0.05$). These findings indicate 25 inhibition of MASP-2 activity would be expected to have a protective or therapeutic effect from renal damage due to ischemic injury.

EXAMPLE 13

30 This Example describes the results of MASP-2/- in a Murine Macular Degeneration Model.

Background/Rationale: Age-related macular degeneration (AMD) is the leading cause of blindness after age 55 in the industrialized world. AMD occurs in two major forms: neovascular (wet) AMD and atrophic (dry) AMD. The neovascular (wet) form

accounts for 90% of severe visual loss associated with AMD, even though only ~20% of individuals with AMD develop the wet form. Clinical hallmarks of AMD include multiple drusen, geographic atrophy, and choroidal neovascularization (CNV). In December, 2004, the FDA approved Macugen (pegaptanib), a new class of ophthalmic
5 drugs to specifically target and block the effects of vascular endothelial growth factor (VEGF), for treatment of the wet (neovascular) form of AMD (Ng et al., *Nat Rev. Drug Discov* 5:123-32 (2006)). Although Macugen represents a promising new therapeutic option for a subgroup of AMD patients, there remains a pressing need to develop additional treatments for this complex disease. Multiple, independent lines of
10 investigation implicate a central role for complement activation in the pathogenesis of AMD. The pathogenesis of choroidal neovascularization (CNV), the most serious form of AMD, may involve activation of complement pathways.

Over twenty-five years ago, Ryan described a laser-induced injury model of CNV in animals (Ryan, S.J., *Tr. Am. Opth. Soc. LXXVII*:707-745, 1979). The model was
15 initially developed using rhesus monkeys, however, the same technology has since been used to develop similar models of CNV in a variety of research animals, including the mouse (Tobe et al., *Am. J. Pathol.* 153:1641-46, 1998). In this model, laser photocoagulation is used to break Bruch's membrane, an act which results in the formation of CNV-like membranes. The laser-induced model captures many of the
20 important features of the human condition (for a recent review, see Ambati et al., *Survey Ophthalmology* 48:257-293, 2003). The laser-induced mouse model is now well established, and is used as an experimental basis in a large, and ever increasing, number of research projects. It is generally accepted that the laser-induced model shares enough biological similarity with CNV in humans that preclinical studies of pathogenesis and
25 drug inhibition using this model are relevant to CNV in humans.

Methods:

A MASP-2^{-/-} mouse was generated as described in **Example 1** and backcrossed for 10 generations with C57Bl/6. The current study compared the results when MASP-2
30 (-/-) and MASP-2 (+/+) male mice were evaluated in the course of laser-induced CNV, an accelerated model of neovascular AMD focusing on the volume of laser-induced CNV by scanning laser confocal microscopy as a measure of tissue injury and determination of levels of VEGF, a potent angiogenic factor implicated in CNV, in the retinal pigment epithelium (RPE)/choroids by ELISA after laser injury.

Induction of choroidal neovascularization (CNV): Laser photocoagulation (532 nm, 200 mW, 100 ms, 75 μ m; Oculight GL, Iridex, Mountain View, CA) was performed on both eyes of each animal on day zero by a single individual masked to drug group assignment. Laser spots were applied in a standardized fashion around the optic nerve, using a slit lamp delivery system and a coverslip as a contact lens. The morphologic end point of the laser injury was the appearance of a cavitation bubble, a sign thought to correlate with the disruption of Bruch's membrane. The detailed methods and endpoints that were evaluated are as follows.

Fluorescein Angiography: Fluorescein angiography was performed with a camera and imaging system (TRC 50 1A camera; ImageNet 2.01 system; Topcon, Paramus, NJ) at 1 week after laser photocoagulation. The photographs were captured with a 20-D lens in contact with the fundus camera lens after intraperitoneal injection of 0.1 ml of 2.5% fluorescein sodium. A retina expert not involved in the laser photocoagulation or angiography evaluated the fluorescein angiograms at a single sitting in masked fashion.

Volume of choroidal neovascularization (CNV): One week after laser injury, eyes were enucleated and fixed with 4% paraformaldehyde for 30 min at 4°C. Eye cups were obtained by removing anterior segments and were washed three times in PBS, followed by dehydration and rehydration through a methanol series. After blocking twice with buffer (PBS containing 1% bovine serum albumin and 0.5% Triton X-100) for 30 minutes at room temperature, eye cups were incubated overnight at 4°C with 0.5% FITC-isolectin B4 (Vector laboratories, Burlingame, CA), diluted with PBS containing 0.2% BSA and 0.1% Triton X-100, which binds terminal β -D-galactose residues on the surface of endothelial cells and selectively labels the murine vasculature. After two washings with PBS containing 0.1% Triton X-100, the neurosensory retina was gently detached and severed from the optic nerve. Four relaxing radial incisions were made, and the remaining RPE –choroid-sclera complex was flatmounted in antifade medium (Immu-Mount Vectashield Mounting Medium; Vector Laboratories) and cover-slipped.

Flatmounts were examined with a scanning laser confocal microscope (TCS SP; Leica, Heidelberg, Germany). Vessels were visualized by exciting with blue argon wavelength (488 nm) and capturing emission between 515 and 545 nm. A 40X oil-immersion objective was used for all imaging studies. Horizontal optical sections (1 μ m step) were obtained from the surface of the RPE-choroid-sclera complex. The

deepest focal plane in which the surrounding choroidal vascular network connecting to the lesion could be identified was judged to be the floor of the lesion. Any vessel in the laser-targeted area and superficial to this reference plane was judged as CNV. Images of each section were digitally stored. The area of CNV-related fluorescence was measured by computerized image analysis with the microscope software (TCS SP; Leica). The summation of whole fluorescent area in each horizontal section was used as an index for the volume of CNV. Imaging was performed by an operator masked to treatment group assignment.

Because the probability of each laser lesion developing CNV is influenced by the group to which it belongs (mouse, eye, and laser spot), the mean lesion volumes were compared using a linear mixed model with a split plot repeated-measures design. The whole plot factor was the genetic group to which the animal belongs, whereas the split plot factor was the eye. Statistical significance was determined at the 0.05 level. *Post hoc* comparisons of means were constructed with a Bonferroni adjustment for multiple comparisons.

VEGF ELISA. At three days after injury by 12 laser spots, the RPE-choroid complex was sonicated in lysis buffer (20 mM imidazole HCl, 10 mM KCl, 1 mM MgCl₂, 10 mM EGTA, 1% Triton X-100, 10 mM NaF, 1 mM Na molybdate, and 1 mM EDTA with protease inhibitor) on ice for 15 min. VEGF protein levels in the supernatant were determined by an ELISA kit (R&D Systems, Minneapolis, MN) that recognizes all splice variants, at 450 to 570 nm (Emax; Molecular Devices, Sunnyvale, CA), and normalized to total protein. Duplicate measurements were performed in a masked fashion by an operator not involved in photocoagulation, imaging, or angiography. VEGF numbers were represented as the mean $\pm$ SEM of at least three independent experiments and compared using the Mann-Whitney U test. The null hypothesis was rejected at $P < 0.05$.

RESULTS:

Assessment of VEGF Levels:

FIGURE 13A graphically illustrates the VEGF protein levels in RPE-choroid complex isolated from C57Bl6 wildtype and MASP-2(-/-) mice at day zero. As shown in **FIGURE 13A**, the assessment of VEGF levels indicate a decrease in baseline levels for VEGF in the MASP-2 (-/-) mice versus the C57bl wildtype control mice. **FIGURE 13B** graphically illustrates VEGF protein levels measured at day three following laser induced

injury. As shown in **FIGURE 13B** VEGF levels were significantly increased in the wildtype (+/+) mice three days following laser induced injury, consistent with published studies (Nozaki et al., *Proc. Natl. Acad. Sci. USA* 103:2328-33 (2006)). However, surprisingly very low levels of VEGF were seen in the MASP-2 (-/-) mice.

5 **Assessment of choroidal neovascularization (CNV):**

In addition to the reduction in VEGF levels following laser induced macular degeneration, CNV area was determined before and after laser injury. **FIGURE 14** graphically illustrates the CNV volume measured in C57bl wildtype mice and MASP-2(-/-) mice at day seven following laser induced injury. As shown in
10 **FIGURE 14**, the MASP-2 (-/-) mice displayed about a 30% reduction in the CNV area following laser induced damage at day seven in comparison to the wildtype control mice.

These findings indicate a reduction in VEGF and CNV as seen in the MASP (-/-) mice versus the wildtype (+/+) control and that blockade of MASP-2 with an inhibitor would have a preventive or therapeutic effect in the treatment of macular degeneration.

15

EXAMPLE 14

This Example demonstrates that thrombin activation can occur following lectin pathway activation under physiological conditions, and demonstrates the extent of MASP-2 involvement. In normal rat serum, activation of the lectin pathway leads to
20 thrombin activation (assessed as thrombin deposition) concurrent with complement activation (assessed as C4 deposition). As can be seen in **FIGURES 15A** and **15B**, thrombin activation in this system is inhibited by a MASP-2 blocking antibody (Fab2 format), exhibiting an inhibition concentration-response curve (**FIGURE 15B**) that parallels that for complement activation (**FIGURE 15A**). These data suggest that
25 activation of the lectin pathway as it occurs in trauma will lead to activation of both complement and coagulation systems in a process that is entirely dependent on MASP-2. By inference, MASP2 blocking antibodies may prove efficacious in mitigating cases of excessive systemic coagulation, e.g., disseminated intravascular coagulation, which is one of the hallmarks leading to mortality in major trauma cases.

30

EXAMPLE 15

This Example provides results generated using a localized Schwartzman reaction model of disseminated intravascular coagulation ("DIC") in MASP-2 $-/-$ deficient and MASP-2 $+/+$ sufficient mice to evaluate the role of lectin pathway in DIC.

5 **Background/Rationale:**

As described supra, blockade of MASP-2 inhibits lectin pathway activation and reduces the generation of both anaphylatoxins C3a and C5a. C3a anaphylatoxins can be shown to be potent platelet aggregators *in vitro*, but their involvement *in vivo* is less well defined and the release of platelet substances and plasmin in wound repair may only
10 secondarily involve complement C3. In this Example, the role of the lectin pathway was analyzed in MASP-2 $(-/-)$ and WT $(+/+)$ mice in order to address whether prolonged elevation of C3 activation is necessary to generate disseminated intravascular coagulation.

Methods:

15 The MASP-2 $(-/-)$ mice used in this study were generated as described in **Example 1** and backcrossed for at least 10 generations with C57Bl/6.

The localized Schwartzman reaction model was used in this experiment. The localized Schwartzman reaction (LSR) is a lipopolysaccharide (LPS) -induced response with well-characterized contributions from cellular and humoral elements of the innate
20 immune system. Dependent of the LSR on complement is well established (Polak, L., et al., *Nature* 223:738-739 (1969); Fong J.S. et al., *J Exp Med* 134:642-655 (1971)). In the LSR model, the mice were primed for 4 hours with TNF alpha (500 ng, intrascrotal), then the mice were anaesthetized and prepared for intravital microscopy of the cremaster muscle. Networks of post-capillary venules (15-60 μ m diameter) with good blood flow
25 (1-4 mm/s) were selected for observation. Animals were treated with fluorescent antibodies to selectively label neutrophils, or platelets. The network of vessels was sequentially scanned and images of all vessels were digitally recorded of later analysis. After recording the basal state of the microcirculation, mice received a single intravenous injection of LPS (100 μ g), either alone or with the agents listed below. The same
30 network of vessels was then scanned every 10 minutes for 1 hour. Specific accumulation of fluorophores was identified by subtraction of background fluorescence and enhanced by thresholding the image. The magnitude of reactions was measured from recorded images. The primary measure of Schwartzman reactions was aggregate data.

The studies compared the MASP-2 $+/+$ sufficient, or wild type, mice exposed to either a known complement pathway depletory agent, cobra venom factor (CVF), or a terminal pathway inhibitor (C5aR antagonist). The results (**FIGURE 16A**) demonstrate that CVF as well as a C5aR antagonist both prevented the appearance of aggregates in the vasculature. In addition, the MASP-2 $-/-$ deficient mice (**FIGURE 16B**) also demonstrated complete inhibition of the localized Schwartzman reaction, supporting lectin pathway involvement. These results clearly demonstrate the role of MASP-2 in DIC generation and support the use of MASP-2 inhibitors for the treatment and prevention of DIC.

10

EXAMPLE 16

This Example describes the analysis of MASP-2 $(-/-)$ mice in a Murine Renal Transplantation Model.

Background/Rationale:

The role of MASP-2 in the functional outcome of kidney transplantation was assessed using a mouse model.

Methods:

The functional outcome of kidney transplantation was assessed using a single kidney isograft into uninephrectomized recipient mice, with six WT $(+/+)$ transplant recipients (B6), and six MASP-2 $(-/-)$ transplant recipients. To assess the function of the transplanted kidney, the remaining native kidney was removed from the recipient 5 days after transplantation, and renal function was assessed 24 hours later by measurement of blood urea nitrogen (BUN) levels.

Results:

FIGURE 17 graphically illustrates the blood urea nitrogen (BUN) levels of the kidney at 6 days post kidney transplant in the WT $(+/+)$ recipients and the MASP-2 $(-/-)$ recipients. As shown in **FIGURE 17**, strongly elevated BUN levels were observed in the WT $(+/+)$ (B6) transplant recipients (normal BUN levels in mice are < 5 mM), indicating renal failure. In contrast, MASP-2 $(-/-)$ isograft recipient mice showed substantially lower BUN levels, suggesting improved renal function. It is noted that these results were obtained using grafts from WT $(+/+)$ kidney donors, suggesting that the absence of a functional lectin pathway in the transplant recipient alone is sufficient to achieve a therapeutic benefit.

Taken together, these results indicate that transient inhibition of the lectin pathway via MASP-2 inhibition provides a method of reducing morbidity and delayed graft function in renal transplantation, and that this approach is likely to be useful in other transplant settings.

5

EXAMPLE 17

This Example demonstrates that MASP-2 (-/-) mice are resistant to septic shock in a Murine Polymicrobial Septic Peritonitis Model.

Background/Rationale:

10 To evaluate the potential effects of MASP-2 (-/-) in infection, the cecal ligation and puncture (CLP) model, a model of polymicrobial septic peritonitis was evaluated. This model is thought to most accurately mimic the course of human septic peritonitis. The cecal ligation and puncture (CLP) model is a model in which the cecum is ligated and punctured by a needle, leading to continuous leakage of the bacteria into the
15 abdominal cavity which reach the blood through the lymph drainage and are then distributed into all the abdominal organs, leading to multi-organ failure and septic shock (Eskandari et al., *J Immunol* 148(9):2724-2730 (1992)). The CLP model mimics the course of sepsis observed in patients and induces an early hyper-inflammatory response followed by a pronounced hypo-inflammatory phase. During this phase, the animals are
20 highly sensitive to bacterial challenges (Wichterman et al., *J. Surg. Res.* 29(2):189-201 (1980)).

Methods:

The mortality of polymicrobial infection using the cecal ligation and puncture (CLP) model was measured in WT (+/+) (n=18) and MASP-2 (-/-) (n=16) mice. Briefly
25 described, MASP-2 deficient mice and their wild-type littermates were anaesthetized and the cecum was exteriorized and ligated 30% above the distal end. After that, the cecum was punctured once with a needle of 0.4 mm diameter. The cecum was then replaced into the abdominal cavity and the skin was closed with clamps. The survival of the mice subjected to CLP was monitored over a period of 14 days after CLP. A peritoneal lavage
30 was collected in mice 16 hours post CLP to measure bacterial load. Serial dilutions of the peritoneal lavage were prepared in PBS and inoculated in Mueller Hinton plates with subsequent incubation at 37°C under anaerobic conditions for 24 hours after which bacterial load was determined.

The TNF- α cytokine response to the bacterial infection was also measured in the WT (+/+) and MASP-2 (-/-) mice 16 hours after CLP in lungs and spleens via quantitative real time polymerase chain reaction (qRT-PCR). The serum level of TNF- α 16 hours after CLP in the WT (+/+) and MASP-2 (-/-) mice was also quantified by sandwich ELISA.

Results:

FIGURE 18 graphically illustrates the percentage survival of the CLP treated animals as a function of the days after the CLP procedure. As shown in **FIGURE 18**, the lectin pathway deficiency in the MASP-2 (-/-) mice does not increase the mortality of mice after polymicrobial infection using the cecal ligation and puncture model as compared to WT (+/+) mice. However, as shown in **FIGURE 19**, MASP-2 (-/-) mice showed a significantly higher bacterial load (approximately a 1000-fold increase in bacterial numbers) in peritoneal lavage after CLP when compared to their WT (+/+) littermates. These results indicate that MASP-2 (-/-) deficient mice are resistant to septic shock. The reduced bacterial clearance in MASP-2 deficient mice in this model may be due to an impaired C3b mediated phagocytosis, as it was demonstrated that C3 deposition is MASP-2 dependent.

It was determined that the TNF- α cytokine response to the bacterial infection was not elevated in the MASP-2 (-/-) mice as compared to the WT (+/+) controls (data not shown). It was also determined that there was a significantly higher serum concentration of TNF- α in WT (+/+) mice 16 hours after CLP in contrast to MASP-2 (-/-) mice, where the serum level of TNF- α remained nearly unaltered. These results suggest that the intense inflammatory response to the septic condition was tempered in MASP-2 (-/-) mice and allowed the animals to survive in the presence of higher bacterial counts.

Taken together, these results demonstrate the potential deleterious effects of lectin pathway complement activation in the case of septicemia and the increased mortality in patients with overwhelming sepsis. These results further demonstrate that MASP-2 deficiency modulates the inflammatory immune response and reduces the expression levels of inflammatory mediators during sepsis. Therefore, it is believed that inhibition of MASP-2 (-/-) by administration of inhibitory monoclonal antibodies against MASP-2 would be effective to reduce the inflammatory response in a subject suffering from septic shock.

EXAMPLE 18

This Example describes analysis of MASP-2 (-/-) mice in a Murine Intranasal Infectivity Model.

5 **Background/Rationale:**

Pseudomonas aeruginosa is a Gram negative opportunistic human bacterial pathogen that causes a wide range of infections, particularly in immune-compromised individuals. It is a major source of acquired nosocomial infections, in particular hospital-acquired pneumonia. It is also responsible for significant morbidity and mortality in
10 cystic fibrosis (CF) patients. *P. aeruginosa* pulmonary infection is characterized by strong neutrophil recruitment and significant lung inflammation resulting in extensive tissue damage (Palanki M.S. et al., *J. Med. Chem* 51:1546-1559 (2008)).

In this Example, a study was undertaken to determine whether the removal of the lectin pathway in MASP-2 (-/-) mice increases the susceptibility of the mice to bacterial
15 infections.

Methods:

Twenty-two WT (+/+) mice, twenty-two MASP-2 (-/-) mice, and eleven C3 (-/-) mice were challenged with intranasal administration of *P. aeruginosa* bacterial strain. The mice were monitored over the six days post-infection and Kaplan-Mayer plots were
20 constructed showing percent survival.

Results:

FIGURE 20 is a Kaplan-Mayer plot of the percent survival of WT (+/+), MASP-2 (-/-) or C3 (-/-) mice six days post-infection. As shown in **FIGURE 20**, no differences were observed in the MASP-2 (-/-) mice versus the WT (+/+) mice. However, removal of the classical (C1q) pathway in the C3 (-/-) mice resulted in a severe susceptibility to
25 bacterial infection. These results demonstrate that MASP-2 inhibition does not increase susceptibility to bacterial infection, indicating that it is possible to reduce undesirable inflammatory complications in trauma patients by inhibiting MASP-2 without compromising the patient's ability to fight infections using the classical complement
30 pathway.

EXAMPLE 19

This Example describes the pharmacodynamic analysis of representative high affinity anti-MASP-2 Fab2 antibodies that were identified as described in **Example 10**.

Background/Rationale:

5 As described in **Example 10**, in order to identify high-affinity antibodies that block the rat lectin pathway, rat MASP-2 protein was utilized to pan a phage display library. This library was designed to provide for high immunological diversity and was constructed using entirely human immunoglobulin gene sequences. As described in **Example 10**, approximately 250 individual phage clones were identified that bound with
10 high affinity to the rat MASP-2 protein by ELISA screening. Sequencing of these clones identified 50 unique MASP-2 antibody encoding phage. Fab2 protein was expressed from these clones, purified and analyzed for MASP-2 binding affinity and lectin complement pathway functional inhibition.

As shown in **TABLE 6** of **Example 10**, 17 anti-MASP-2 Fab2s with functional
15 blocking activity were identified as a result of this analysis (a 34% hit rate for blocking antibodies). Functional inhibition of the lectin complement pathway by Fab2s was apparent at the level of C4 deposition, which is a direct measure of C4 cleavage by MASP-2. Importantly, inhibition was equally evident when C3 convertase activity was assessed, demonstrating functional blockade of the lectin complement pathway. The 17
20 MASP-2 blocking Fab2s identified as described in **Example 10** potently inhibit C3 convertase formation with IC₅₀ values equal to or less than 10 nM. Eight of the 17 Fab2s identified have IC₅₀ values in the sub-nanomolar range. Furthermore, all 17 of the MASP-2 blocking Fab2s gave essentially complete inhibition of the C3 convertase formation in the lectin pathway C3 convertase assay, as shown in FIGURES 8A-C, and
25 summarized in TABLE 6 of Example 10. Moreover, each of the 17 blocking anti-MASP-2 Fab2s shown in TABLE 6 potently inhibit C3b generation (>95%), thus demonstrating the specificity of this assay for lectin pathway C3 convertase.

Rat IgG2c and mouse IgG2a full-length antibody isotype variants were derived from Fab2 #11. This Example describes the *in vivo* characterization of these isotypes for
30 pharmacodynamic parameters.

Methods:

As described in Example 10, rat MASP-2 protein was utilized to pan a Fab phage display library, from which Fab2#11 was identified. Rat IgG2c and mouse IgG2a full-

length antibody isotype variants were derived from Fab2 #11. Both rat IgG2c and mouse IgG2a full length antibody isotypes were characterized *in vivo* for pharmacodynamic parameters as follows.

In vivo study in mice:

5 A pharmacodynamic study was carried out in mice to investigate the effect of anti-MASP-2 antibody dosing on the plasma lectin pathway activity *in vivo*. In this study, C4 deposition was measured *ex vivo* in a lectin pathway assay at various time points following subcutaneous (sc) and intraperitoneal (ip) administration of 0.3 mg/kg or 1.0 mg/kg of the mouse anti-MASP-2 MoAb (mouse IgG2a full-length antibody isotype
10 derived from Fab2#11).

FIGURE 21 graphically illustrates lectin pathway specific C4b deposition, measured *ex vivo* in undiluted serum samples taken from mice (n=3 mice/group) at various time points after subcutaneous dosing of either 0.3 mg/kg or 1.0 mg/kg of the mouse anti-MASP-2 MoAb. Serum samples from mice collected prior to antibody
15 dosing served as negative controls (100% activity), while serum supplemented *in vitro* with 100 nM of the same blocking anti-MASP-2 antibody was used as a positive control (0% activity).

The results shown in **FIGURE 21** demonstrate a rapid and complete inhibition of C4b deposition following subcutaneous administration of 1.0 mg/kg dose of mouse anti-
20 MASP-2 MoAb. A partial inhibition of C4b deposition was seen following subcutaneous administration of 0.3 mg/kg dose of mouse anti-MASP-2 MoAb.

The time course of lectin pathway recovery was followed for three weeks following a single ip administration of mouse anti-MASP-2 MoAb at 0.6 mg/kg in mice. As shown in **FIGURE 22**, a precipitous drop in lectin pathway activity occurred post
25 antibody dosing followed by complete lectin pathway inhibition that lasted for about 7 days after ip administration. Slow restoration of lectin

These results demonstrate that the mouse anti-MASP-2 Moab derived from Fab2 #11 inhibits the lectin pathway of mice in a dose-responsive manner when delivered systemically.

30

EXAMPLE 20

This Example describes analysis of the mouse anti-MASP-2 Moab derived from Fab2 #11 for efficacy in a mouse model for age-related macular degeneration.

Background/Rationale:

As described in **Example 10**, rat MASP-2 protein was utilized to pan a Fab phage display library, from which Fab2#11 was identified as a functionally active antibody. Full length antibodies of the rat IgG2c and mouse IgG2a isotypes were generated from Fab2 #11. The full length anti-MASP-2 antibody of the mouse IgG2a isotype was characterized for pharmacodynamic parameters as described in **Example 19**. In this Example, the mouse anti-MASP-2 full-length antibody derived from Fab2 #11 was analyzed in the mouse model of age-related macular degeneration (AMD), described by Bora P.S. et al, *J Immunol* 174:491-497 (2005).

Methods:

The mouse IgG2a full-length anti-MASP-2 antibody isotype derived from Fab2 #11 as described in **Example 19**, was tested in the mouse model of age-related macular degeneration (AMD) as described in **Example 13** with the following modifications.

Administration of mouse-anti-MASP-2 MoAbs

Two different doses (0.3 mg/kg and 1.0 mg/kg) of mouse anti-MASP-2 MoAb along with an isotype control MoAb treatment were injected ip into WT (+/+) mice (n= 8 mice per group) 16 hours prior to CNV induction

Induction of choroidal neovascularization (CNV)

The induction of choroidal neovascularization (CNV) and measurement of the volume of CNV was carried out using laser photocoagulation as described in Example 13.

Results:

FIGURE 23 graphically illustrates the CNV area measured at 7 days post laser injury in mice treated with either isotype control MoAb, or mouse anti-MASP-2 MoAb (0.3 mg/kg and 1.0 mg/kg). As shown in **FIGURE 23**, in the mice pre-treated with 1.0 mg/kg anti-MASP-2 MoAb, a statistically significant ($p < 0.01$) approximately 50% reduction in CNV was observed seven days post-laser treatment. As further shown in **FIGURE 23**, it was observed that a 0.3 mg/kg dose of anti-MASP-2 MoAb was not efficacious in reducing CNV. It is noted that the 0.3 mg/kg dose of anti-MASP-2 MoAb was shown to have a partial and transient inhibition of C4b deposition following subcutaneous administration, as described in **Example 19** and shown in **FIGURE 21**.

The results described in this Example demonstrate that blockade of MASP-2 with an inhibitor, such as anti-MASP-2 MoAb, has a preventative and/or therapeutic effect in the treatment of macular degeneration. It is noted that these results are consistent with

the results observed in the study carried out in the MASP-2 (-/-) mice, described in **Example 13**, in which a 30% reduction in the CNV 7 days post-laser treatment was observed in MASP-2 (-/-) mice in comparison to the wild-type control mice. Moreover, the results in this Example further demonstrate that systemically delivered anti-MASP-2 antibody provides local therapeutic benefit in the eye, thereby highlighting the potential for a systemic route of administration to treat AMD patients. In summary, these results provide evidence supporting the use of MASP-2 MoAb in the treatment of AMD.

EXAMPLE 21

This Example demonstrates that MASP-2 deficient mice are protected from *Neisseria meningitidis* induced mortality after infection with *N. meningitidis* and have enhanced clearance of bacteraemia as compared to wild type control mice.

Rationale: *Neisseria meningitidis* is a heterotrophic gram-negative diplococcal bacterium known for its role in meningitis and other forms of meningococcal disease such as meningococemia. *N. meningitidis* is a major cause of morbidity and mortality during childhood. Severe complications include septicaemia, Waterhouse-Friderichsen syndrome, adrenal insufficiency and disseminated intravascular coagulation (DIC). See e.g., Rintala E. et al., *Critical Care Medicine* 28(7):2373-2378 (2000). In this Example, the role of the lectin pathway was analyzed in MASP-2 (-/-) and WT (+/+) mice in order to address whether MASP-2 deficient mice would be susceptible to *N. meningitidis* induced mortality.

Methods:

MASP-2 knockout mice were generated as described in **Example 1** and backcrossed for at least 10 generations with C57Bl/6. 10 week old MASP-2 KO mice (n=10) and wild type C57/B6 mice (n=10) were inoculated by intravenous injection with either a dosage of 5×10^8 cfu/100 μ l, 2×10^8 cfu/100 μ l or 3×10^7 cfu/100 μ l of *Neisseria meningitidis* Serogroup A Z2491 in 400 mg/kg iron dextran. Survival of the mice after infection was monitored over a 72 hour time period. Blood samples were taken from the mice at hourly intervals after infection and analyzed to determine the serum level (log cfu/ml) of *N. meningitidis* in order to verify infection and determine the rate of clearance of the bacteria from the serum.

Results:

FIGURE 24A graphically illustrates the percent survival of MASP-2 KO and WT mice after administration of an infective dose of $5 \times 10^8/100 \mu\text{l}$ cfu *N. meningitidis*. As shown in **FIGURE 24A**, after infection with the highest dose of $5 \times 10^8/100 \mu\text{l}$ cfu *N. meningitidis*, 100% of the MASP-2 KO mice survived throughout the 72 hour period after infection. In contrast, only 20% of the WT mice were still alive 24 hours after infection. These results demonstrate that MASP-2 deficient mice are protected from *N. meningitidis* induced mortality.

FIGURE 24B graphically illustrates the log cfu/ml of *N. meningitidis* recovered at different time points in blood samples taken from the MASP-2 KO and WT mice infected with 5×10^8 cfu/100 μl *N. meningitidis*. As shown in **FIGURE 24B**, in WT mice the level of *N. meningitidis* in the blood reached a peak of about 6.5 log cfu/ml at 24 hours after infection and dropped to zero by 48 hours after infection. In contrast, in the MASP-2 KO mice, the level of *N. meningitidis* reached a peak of about 3.5 log cfu/ml at 6 hours after infection and dropped to zero by 36 hours after infection.

FIGURE 25A graphically illustrates the percent survival of MASP-2 KO and WT mice after infection with 2×10^8 cfu/100 μl *N. meningitidis*. As shown in **FIGURE 25A**, after infection with the dose of 2×10^8 cfu/100 μl *N. meningitidis*, 100% of the MASP-2 KO mice survived throughout the 72 hour period after infection. In contrast, only 80% of the WT mice were still alive 24 hours after infection. Consistent with the results shown in **FIGURE 24A**, these results further demonstrate that MASP-2 deficient mice are protected from *N. meningitidis* induced mortality.

FIGURE 25B graphically illustrates the log cfu/ml of *N. meningitidis* recovered at different time points in blood samples taken from the WT mice infected with 2×10^8 cfu/100 μl *N. meningitidis*. As shown in **FIGURE 25B**, the level of *N. meningitidis* in the blood of WT mice infected with 2×10^8 cfu reached a peak of about 4 log cfu/ml at 12 hours after infection and dropped to zero by 24 hours after infection. **FIGURE 25C** graphically illustrates the log cfu/ml of *N. meningitidis* recovered at different time points in blood samples taken from the MASP-2 KO mice infected with 2×10^8 cfu/100 μl *N. meningitidis*. As shown in **FIGURE 25C**, the level of *N. meningitidis* in the blood of MASP-2 KO mice infected with 2×10^8 cfu reached a peak level of about 3.5 log cfu/ml at 2 hours after infection and dropped to zero at 3 hours after infection. Consistent with the results shown in **FIGURE 24B**, these results demonstrate that although the MASP-2 KO

mice were infected with the same dose of *N. meningitidis* as the WT mice, the MASP-2 KO mice have enhanced clearance of bacteraemia as compared to WT.

The percent survival of MASP-2 KO and WT mice after infection with the lowest dose of 3×10^7 cfu/100 μ l *N. meningitidis* was 100% at the 72 hour time period (data not shown).

Discussion

These results show that MASP-2 deficient mice are protected from *N. meningitidis* induced mortality and have enhanced clearance of bacteraemia as compared to the WT mice. Therefore, in view of these results, it is expected that therapeutic application of MASP-2 inhibitors, such as MASP-2 MoAb, would be expected to be efficacious to treat, prevent or mitigate the effects of infection with *N. meningitidis* bacteria (i.e., sepsis and DIC). Further, these results indicate that therapeutic application of MASP-2 inhibitors, such as MASP-2 MoAb would not predispose a subject to an increased risk to contract *N. meningitidis* infections.

EXAMPLE 22

This Example describes the discovery of novel lectin pathway mediated and MASP-2 dependent C4-bypass activation of complement C3.

Rationale:

The principal therapeutic benefit of utilizing inhibitors of complement activation to limit myocardial ischemia/reperfusion injury (MIRI) was convincingly demonstrated in an experimental rat model of myocardial infarction two decades ago: Recombinant sCR1, a soluble truncated derivative of the cell surface complement receptor type-1 (CR1), was given intravenously and its effect assessed in a rat *in vivo* model of MIRI. Treatment with sCR1 reduced infarct volume by more than 40% (Weisman, H.F., et al., *Science* 249:146-151 (1990)). The therapeutic potential of this recombinant inhibitor was subsequently demonstrated in a clinical trial showing that the administration of sCR1 in patients with MI prevented contractile failure in the post-ischemic heart (Shandelya, S., et al., *Circulation* 87:536-546 (1993)). The primary mechanism leading to the activation of complement in ischemic tissue, however, has not been ultimately defined, mainly due to the lack of appropriate experimental models, the limited understanding of the molecular processes that lead to complement activation of oxygen-deprived cells, and the cross-talk and synergisms between the different complement activation pathways.

As a fundamental component of the immune response, the complement system provides protection against invading microorganisms through both antibody-dependent and -independent mechanisms. It orchestrates many cellular and humoral interactions within the immune response, including chemotaxis, phagocytosis, cell adhesion, and B-cell differentiation. Three different pathways initiate the complement cascade: the classical pathway, the alternative pathway, and the lectin pathway. The classical pathway recognition subcomponent C1q binds to a variety of targets - most prominently immune complexes - to initiate the step-wise activation of associated serine proteases, C1r and C1s, providing a major mechanism for pathogen and immune complex clearance following engagement by the adaptive immune system. Binding of C1q to immune complexes converts the C1r zymogen dimer into its active form to cleave and thereby activate C1s. C1s translates C1q binding into complement activation in two cleavage steps: It first converts C4 into C4a and C4b and then cleaves C4b-bound C2 to form the C3 convertase C4b2a. This complex converts the abundant plasma component C3 into C3a and C3b. Accumulation of C3b in close proximity of the C4b2a complex shifts the substrate specificity for C3 to C5 to form the C5 convertase C4b2a(C3b)_n. The C3 and C5 convertase complexes generated via classical pathway activation are identical to those generated through the lectin pathway activation route. In the alternative pathway, spontaneous low-level hydrolysis of component C3 results in deposition of protein fragments onto cell surfaces, triggering complement activation on foreign cells, while cell-associated regulatory proteins on host tissues avert activation, thus preventing self-damage. Like the alternative pathway, the lectin pathway may be activated in the absence of immune complexes. Activation is initiated by the binding of a multi-molecular lectin pathway activation complex to **Pathogen-Associated Molecular Patterns (PAMPs)**, mainly carbohydrate structures present on bacterial, fungal or viral pathogens or aberrant glycosylation patterns on apoptotic, necrotic, malignant or oxygen-deprived cells (Collard, C.D., et al., *Am. J. Pathol.* 156:1549-1556 (2000); Walport, M.J., *N. Engl. J. Med.* 344:1058-1066 (2001); Schwaebler, W., et al., *Immunobiology* 205:455-466 (2002); and Fujita, T., *Nat. Rev. Immunol.* 2:346-353 (2002)).

Mannan-binding lectin (MBL) was the first carbohydrate recognition subcomponent shown to form complexes with a group of novel serine proteases, named MBL-associated Serine Proteases (MASPs) and numbered according to the sequence of their discovery (i.e., MASP-1, MASP-2 and MASP-3). In man, lectin pathway activation

complexes can be formed with four alternative carbohydrate recognition subcomponents with different carbohydrate binding specificities, i.e., MBL 2, and three different members of the ficolin family, namely L-Ficolin, H-ficolin and M-ficolin and MASPs. Two forms of MBL, MBL A and MBL C, and ficolin-A form lectin activation pathway complexes with MASPs in mouse and rat plasma. We have previously cloned and characterised MASP-2 and an additional truncated MASP-2 gene product of 19 kDa, termed MAp19 or sMAP, in human, mouse and rat (Thiel, S., et al., *Nature* 386:506-510 (1997); Stover, C.M., et al., *J. Immunol.* 162:3481-3490 (1999); Takahashi, M., et al., *Int. Immunol.* 11:859-863 (1999); and Stover, C.M., et al., *J. Immunol.* 163:6848-6859 (1999)). MAp19/ sMAP is devoid of protease activity, but may regulate lectin pathway activation by competing for the binding of MASPs to carbohydrate recognition complexes (Iwaki, D. et al., *J. Immunol.* 177:8626-8632 (2006)).

There is evidence suggesting that of the three MASPs, only MASP-2 is required to translate binding of the lectin pathway recognition complexes into complement activation (Thiel, S., et al. (1997); Vorup-Jensen, T., et al., *J. Immunol.* 165:2093-2100 (2000); Thiel, S., et al., *J. Immunol.* 165:878-887 (2000); Rossi, V., et al., *J. Biol. Chem.* 276:40880-40887 (2001)). This conclusion is underlined by the phenotype of a most recently described mouse strain deficient in MASP-1 and MASP-3. Apart from a delay in the onset of lectin pathway mediated complement activation *in vitro* –MASP-1/3 deficient mice retain lectin pathway functional activity. Reconstitution of MASP-1 and MASP-3 deficient serum with recombinant MASP-1 overcomes this delay in lectin pathway activation implying that MASP-1 may facilitate MASP-2 activation (Takahashi, M., et al., *J. Immunol.* 180:6132-6138 (2008)). A most recent study has shown that MASP-1 (and probably also MASP-3) are required to convert the alternative pathway activation enzyme Factor D from its zymogen form into its enzymatically active form (Takahashi, M., et al., *J. Exp. Med.* 207:29-37 (2010)). The physiological importance of this process is underlined by the absence of alternative pathway functional activity in plasma of MASP-1/3 deficient mice.

The recently generated mouse strains with combined targeted deficiencies of the lectin pathway carbohydrate recognition subcomponents MBL A and MBL C may still initiate lectin pathway activation via the remaining murine lectin pathway recognition subcomponent ficolin A (Takahashi, K., et al., *Microbes Infect.* 4:773-784 (2002)). The absence of any residual lectin pathway functional activity in MASP-2 deficient mice

delivers a conclusive model to study the role of this effector arm of innate humoral immunity in health and disease.

The availability of C4 and MASP-2 deficient mouse strains allowed us to define a novel lectin pathway specific, but MASP-2 dependent, C4-bypass activation route of complement C3. The essential contribution of this novel lectin pathway mediated C4-bypass activation route towards post-ischemic tissue loss is underlined by the prominent protective phenotype of MASP-2 deficiency in MIRI while C4-deficient mice tested in the same model show no protection.

In this Example, we describe a novel lectin pathway mediated and MASP-2 dependent C4-bypass activation of complement C3. The physiological relevance of this new activation route is established by the protective phenotype of MASP-2 deficiency in an experimental model of myocardial ischemia/reperfusion injury (MIRI), where C4 deficient animals were not protected.

Methods:

MASP-2 deficient mice show no gross abnormalities. MASP-2 deficient mice were generated as described in **Example 1**. Both heterozygous ($^{+/-}$) and homozygous ($^{-/-}$) MASP-2 deficient mice are healthy and fertile, and show no gross abnormalities. Their life expectancy is similar to that of their WT littermates (>18 months). Prior to studying the phenotype of these mice in experimental models of disease, our MASP-2 $^{-/-}$ line was backcrossed for eleven generations onto a C57BL/6 background. The total absence of MASP-2 mRNA was confirmed by Northern blotting of poly A⁺ selected liver RNA preparations, while the 1.2kb mRNA encoding MAP19 or sMAP (a truncated alternative splicing product of the MASP2 gene) is abundantly expressed.

qRT-PCR analysis using primer pairs specific for either the coding sequence for the serine protease domain of MASP-2 (B chain) or the remainder of the coding sequence for the A-chain showed that no B chain encoding mRNA is detectable in MASP-2 $^{-/-}$ mice while the abundance of the disrupted A chain mRNA transcript was significantly increased. Likewise, the abundance of MAP19/sMAP encoding mRNA is increased in MASP-2 $^{+/-}$ and MASP-2 $^{-/-}$ mice. Plasma MASP-2 levels, determined by ELISA for 5 animals of each genotype, were 300ng/ml for WT controls (range 260-330ng/ml), 360ng/ml for heterozygous mice (range 330-395ng/ml) and undetectable in MASP-2 $^{-/-}$ mice. Using qRT-PCR, mRNA expression profiles were established demonstrating that MASP-2 $^{-/-}$ mice express mRNA for MBL A, MBL C, ficolin A, MASP-1, MASP-3, C1q,

C1rA, C1sA, Factor B, Factor D, C4, and C3 at an abundance similar to that of their MASP-2 sufficient littermates (data not shown).

Plasma C3 levels of MASP-2^{-/-} (n=8) and MASP-2^{+/+} (n=7) littermates were measured using a commercially available mouse C3 ELISA kit (Kamiya, Biomedical, Seattle, WA). C3 levels of MASP-2 deficient mice (average 0.84 mg/ml, +/- 0.34) were similar to those of the WT controls (average 0.92, +/- 0.37).

Results:

MASP-2 is essential for lectin pathway functional activity.

As described in Example 2 and shown in FIGURE 5, the *in vitro* analyses of MASP-2^{-/-} plasma showed a total absence of lectin pathway functional activity on activating Mannan- and Zymosan-coated surfaces for the activation of C4. Likewise, neither lectin pathway-dependent C4 nor C3 cleavage was detectable in MASP-2^{-/-} plasma on surfaces coated with N-acetyl glucosamine, which binds and triggers activation via MBL A, MBL C and ficolin A (data not shown).

The analyses of sera and plasma of MASP-2^{-/-} mice clearly demonstrated that MASP-2 is essentially required to activate complement via the lectin pathway. The total deficiency of lectin pathway functional activity, however, leaves the other complement activation pathways intact: MASP-2^{-/-} plasma can still activate complement via the classical (FIGURE 26A) and the alternative pathway (FIGURE 26B). In FIGURE 26A and 26B, the symbol "*" symbol indicates serum from WT (MASP-2 (+/+)); the symbol "●" indicates serum from WT (C1q depleted); the symbol "□" indicates serum from MASP-2 (-/-); and the symbol "Δ" indicates serum from MASP-2 (-/-) (C1q depleted).

FIGURE 26A graphically illustrates that MASP-2^{-/-} mice retain a functional classical pathway: C3b deposition was assayed on microtiter plates coated with immune complexes (generated *in situ* by coating with BSA then adding goat anti-BSA IgG). FIGURE 26B graphically illustrates MASP-2 deficient mice retain a functional alternative pathway: C3b deposition was assayed on Zymosan coated microtiter plates under conditions that permit only alternative pathway activation (buffer containing Mg²⁺ and EGTA). Results shown in FIGURE 26A and FIGURE 26B are means of duplicates and are typical of three independent experiments. Same symbols for plasma sources were used throughout. These results show that a functional alternative pathway is present in MASP-2 deficient mice, as evidenced in the results shown in FIGURE 26B under

experimental conditions designed to directly trigger the alternative pathway, while inactivating both the classical pathway and lectin pathway.

The lectin pathway of complement activation critically contributes to inflammatory tissue loss in myocardial ischemia/reperfusion injury (MIRI).

5 In order to study the contribution of lectin pathway functional activity to MIRI, we compared MASP-2^{-/-} mice and WT littermate controls in a model of MIRI following transient ligation and reperfusion of the left anterior descending branch of the coronary artery (LAD). The presence or absence of complement C4 has no impact on the degree of ischemic tissue loss in MIRI. We assessed the impact of C4 deficiency on infarct sizes
10 following experimental MIRI. As shown in **FIGURE 27A** and **FIGURE 27B**, identical infarct sizes were observed in both C4-deficient mice and their WT littermates. **FIGURE 27A** graphically illustrates MIRI-induced tissue loss following LAD ligation and reperfusion in C4^{-/-} mice (n=6) and matching WT littermate controls (n=7). **FIGURE 27B** graphically illustrates INF as a function of AAR, clearly demonstrating that C4^{-/-}
15 mice are as susceptible to MIRI as their WT controls (dashed line).

These results demonstrate that C4 deficient mice are not protected from MIRI. This result was unexpected, as it is in conflict with the widely accepted view that the major C4 activation fragment, C4b, is an essential component of the classical and the lectin pathway C3 convertase C4b2a. We therefore assessed whether a residual lectin
20 pathway specific activation of complement C3 can be detected in C4-deficient mouse and human plasma.

The lectin pathway can activate complement C3 in absence of C4 via a novel MASP-2 dependent C4-bypass activation route.

Encouraged by historical reports indicating the existence of a C4-bypass
25 activation route in C4-deficient guinea pig serum (May, J.E., and M. Frank, *J. Immunol.* 111:1671-1677 (1973)), we analyzed whether C4-deficient mice may have residual classical or lectin pathway functional activity and monitored activation of C3 under pathway-specific assay conditions that exclude contributions of the alternative pathway.

C3b deposition was assayed on Mannan-coated microtiter plates using re-calcified
30 plasma at plasma concentrations prohibitive for alternative pathway activation (1.25% and below). While no cleavage of C3 was detectable in C4-deficient plasma tested for classical pathway activation (data not shown), a strong residual C3 cleavage activity was observed in C4-deficient mouse plasma when initiating complement activation via the

lectin pathway. The lectin pathway dependence is demonstrated by competitive inhibition of C3 cleavage following preincubation of C4-deficient plasma dilutions with soluble Mannan (see **FIGURE 28A**). As shown in **FIGURE 28A-D**, MASP-2 dependent activation of C3 was observed in the absence of C4. **FIGURE 28A** graphically illustrates C3b deposition by C4^{+/+} (crosses) and C4^{-/-} (open circles) mouse plasma. Pre-incubating the C4^{-/-} plasma with excess (1 µg/ml) fluid-phase Mannan prior to the assay completely inhibits C3 deposition (filled circles). Results are typical of 3 independent experiments. **FIGURE 28B** graphically illustrates the results of an experiment in which wild-type, MASP-2 deficient (open squares) and C4^{-/-}-mouse plasma (1%) was mixed with various concentrations of anti-rat MASP-2 mAbM11 (abscissa) and C3b deposition assayed on Mannan-coated plates. Results are means (± SD) of 4 assays (duplicates of 2 of each type of plasma). **FIGURE 28C** graphically illustrates the results of an experiment in which Human plasma: pooled NHS (crosses), C4^{-/-} plasma (open circles) and C4^{-/-} plasma pre-incubated with 1 µg/ml Mannan (filled circles). Results are representative of three independent experiments. **FIGURE 28D** graphically illustrates that inhibition of C3b deposition in C4 sufficient and C4 deficient human plasma (1%) by anti-human MASP-2 mAbH3 (Means ± SD of triplicates). As shown in **FIGURE 28B**, no lectin pathway-dependent C3 activation was detected in MASP-2^{-/-}-plasma assayed in parallel, implying that this C4-bypass activation route of C3 is MASP-2 dependent.

To further corroborate these findings, we established a series of recombinant inhibitory mAbs isolated from phage display antibody libraries by affinity screening against recombinant human and rat MASP-2A (where the serine residue of the active protease domain was replaced by an alanine residue by site-directed mutagenesis to prevent autolytic degradation of the antigen). Recombinant antibodies against MASP-2 (AbH3 and AbM11) were isolated from Combinatorial Antibody Libraries (Knappik, A., et al., *J. Mol. Biol.* 296:57-86 (2000)), using recombinant human and rat MASP-2A as antigens (Chen, C.B. and Wallis, *J. Biol. Chem.* 276:25894-25902 (2001)). An anti-rat Fab2 fragment that potently inhibited lectin pathway-mediated activation of C4 and C3 in mouse plasma (IC₅₀~1 nM) was converted to a full-length IgG2a antibody. Polyclonal anti-murine MASP-2A antiserum was raised in rats. These tools allowed us to confirm MASP-2 dependency of this novel lectin pathway specific C4-bypass activation route of C3, as further described below.

As shown in **FIGURE 28B**, M211, an inhibitory monoclonal antibody which selectively binds to mouse and rat MASP-2 inhibited the C4-bypass activation of C3 in C4-deficient mouse as well as C3 activation of WT mouse plasma via the lectin pathway in a concentration dependent fashion with similar IC₅₀ values. All assays were carried out at high plasma dilutions rendering the alternative pathway activation route dysfunctional (with the highest plasma concentration being 1.25%).

In order to investigate the presence of an analogous lectin pathway specific C4-bypass activation of C3 in humans, we analyzed the plasma of a donor with an inherited deficiency of both human C4 genes (i.e., C4A and C4B), resulting in total absence of C4 (Yang, Y., et al., *J. Immunol.* 173:2803-2814 (2004)). **FIGURE 28C** shows that this patient's plasma efficiently activates C3 in high plasma dilutions (rendering the alternative activation pathway dysfunctional). The lectin pathway specific mode of C3 activation on Mannan-coated plates is demonstrated in murine C4-deficient plasma (**FIGURE 28A**) and human C4 deficient plasma (**FIGURE 28C**) by adding excess concentrations of fluid-phase Mannan. The MASP-2 dependence of this activation mechanism of C3 in human C4-deficient plasma was assessed using AbH3, a monoclonal antibody that specifically binds to human MASP-2 and ablates MASP-2 functional activity. As shown in **FIGURE 28D**, AbH3 inhibited the deposition of C3b (and C3dg) in both C4-sufficient and C4-deficient human plasma with comparable potency.

In order to assess a possible role of other complement components in the C4-bypass activation of C3, we tested plasma of MASP-1/3-/- and Bf/C2-/- mice alongside MASP-2-/-, C4-/- and C1q-/- plasma (as controls) under both lectin pathway specific and classical pathway specific assay conditions. The relative amount of C3 cleavage was plotted against the amount of C3 deposited when using WT plasma.

FIGURE 29A graphically illustrates a comparative analysis of C3 convertase activity in plasma from various complement deficient mouse strains tested either under lectin activation pathway or classical activation pathway specific assay conditions. Diluted plasma samples (1%) of WT mice (n=6), MASP-2-/- mice (n=4), MASP-1/3-/- mice (n=2), C4-/- mice (n=8), C4/MASP-1/3-/- mice (n=8), Bf/C2-/- (n=2) and C1q-/- mice (n=2) were tested in parallel. Reconstitution of Bf/C2-/- plasma with 2.5µg/ml recombinant rat C2 (Bf/C2-/- +C2) restored C3b deposition. Results are means (±SD). **p<0.01 (compared to WT plasma). As shown in **FIGURE 29A**, substantial C3 deposition is seen in C4-/- plasma tested under lectin pathway specific assay conditions,

but not under classical pathway specific conditions. Again, no C3 deposition was seen in MASP-2 deficient plasma via the lectin pathway activation route, while the same plasma deposited C3 via the classical pathway. In MASP-1/3-/- plasma, C3 deposition occurred in both lectin and classical pathway specific assay conditions. No C3 deposition was
 5 seen in plasma with a combined deficiency of C4 and MASP-1/3, either using lectin pathway or classical pathway specific conditions. No C3 deposition is detectable in C2/Bf-/- plasma, either via the lectin pathway, or via the classical pathway. Reconstitution of C2/Bf-/- mouse plasma with recombinant C2, however, restored both lectin pathway and classical pathway-mediated C3 cleavage. The assay conditions were
 10 validated using C1q-/- plasma.

FIGURE 29B graphically illustrates time-resolved kinetics of C3 convertase activity in plasma from various complement deficient mouse strains WT, fB-/-, C4-/-, MASP-1/3-/-, and MASP-2-/-plasma, tested under lectin activation pathway specific assay conditions (1% plasma, results are typical of three independent experiments). As
 15 shown in **FIGURE 29B**, while no C3 cleavage was seen in MASP-2-/-plasma, fB-/- plasma cleaved C3 with similar kinetics to the WT plasma. A significant delay in the lectin pathway-dependent conversion of C3 to C3b (and C3dg) was seen in C4-/-as well as in MASP-1/3 deficient plasma. This delay of C3 activation in MASP-1/3-/- plasma was recently shown to be MASP-1, rather than MASP-3 dependent (Takahashi, M., et al.,
 20 *J. Immunol.* 180:6132-6138 (2008)).

Discussion:

The results described in this Example strongly suggest that MASP-2 functional activity is essential for the activation of C3 via the lectin pathway both in presence and absence of C4. Furthermore, C2 and MASP-1 are required for this novel lectin pathway
 25 specific C4-bypass activation route of C3 to work. The comparative analysis of lectin pathway functional activity in MASP-2-/-as well as C4-/- plasma revealed the existence of a previously unrecognized C4-independent, but MASP-2-dependent activation route of complement C3 and showed that C3 can be activated in a lectin pathway-dependent mode in total absence of C4. While the detailed molecular composition and the sequence of
 30 activation events of this novel MASP-2 dependent C3 convertase remains to be elucidated, our results imply that this C4-bypass activation route additionally requires the presence of complement C2 as well as MASP-1. The loss of lectin pathway-mediated C3 cleavage activity in plasma of mice with combined C4 and MASP-1/3 deficiency may be

explained by a most recently described role of MASP-1 to enhance MASP-2 dependent complement activation through direct cleavage and activation of MASP-2 (Takahashi, M., et al., *J. Immunol.* 180:6132-6138 (2008)). Likewise, MASP-1 may aid MASP-2 functional activity through its ability to cleave C2 (Moller-Kristensen, et al., *Int.* 5 *Immunol.* 19:141-149 (2007)). Both activities may explain the reduced rate by which MASP-1/3 deficient plasma cleaves C3 via the lectin activation pathway and why MASP-1 may be required to sustain C3 conversion via the C4-bypass activation route.

The inability of C2/fB-/- plasma to activate C3 via the lectin pathway was shown to be C2-dependent as the addition of recombinant rat C2 to C2/fB-/- plasma restored the 10 ability of the reconstituted plasma to activate C3 on Mannan-coated plates.

The finding that C4 deficiency specifically disrupts the classical complement activation pathway while the lectin pathway retains a physiologically critical level of C3 convertase activity via a MASP-2 dependent C4-bypass activation route calls for a re-assessment of the role of the lectin pathway in various disease models, including 15 experimental *S. pneumoniae* infection (Brown, J. S., et al., *Proc. Natl. Acad. Sci. U. S. A.* 99:16969-16974 (2002); Experimental Allergic Encephalomyelitis (Boos, L.A., et al., *Glia* 49:158-160 (2005); and models of C3 dependent murine liver regeneration (Clark, A., et al., *Mol. Immunol.* 45:3125-3132 (2008)). The latter group demonstrated that C4-deficient mice can activate C3 in an alternative pathway independent fashion as *in vivo* 20 inhibition of the alternative pathway by an antibody-mediated depletion of factor B functional activity did not effect C3 cleavage-dependent liver regeneration in C4-/- mice (Clark, A., et al. (2008)). This lectin pathway mediated C4-bypass activation route of C3 may also explain the lack of a protective phenotype of C4 deficiency in our model of MIRI as well as in a previously described model of renal allograft rejection (Lin, T., et 25 al., *Am. J. Pathol.* 168:1241-1248 (2006)). In contrast, our recent results have independently demonstrated a significant protective phenotype of MASP-2-/-mice in models of renal transplantation (Farrar, C.A., et al., *Mol. Immunol.* 46:2832 (2009)).

In summary, the results of this Example support the view that MASP-2 dependent C4-bypass activation of C3 is a physiologically relevant mechanism that may be 30 important under conditions where availability of C4 is limiting C3 activation.

EXAMPLE 23

This Example describes activation of C3 by thrombin substrates and C3 deposition on mannan in WT (+/+), MASP-2 (-/-), F11 (-/-), F11/C4 (-/-) and C4 (-/-) mice.

5 **Rationale:**

As described in **Example 14**, it was determined that thrombin activation can occur following lectin pathway activation under physiological conditions, and demonstrates the extent of MASP-2 involvement. C3 plays a central role in the activation of complement system. C3 activation is required for both classical and alternative
10 complement activation pathways. An experiment was carried out to determine whether C3 is activated by thrombin substrates.

Methods:

C3 Activation by thrombin substrates

Activation of C3 was measured in the presence of the following activated forms
15 of thrombin substrates; human FCXIa, human FVIIa, bovine FXa, human FXa, human activated protein C, and human thrombin. C3 was incubated with the various thrombin substrates, then separated under reducing conditions on 10% SDS-polyacrylamide gels. After electrophoretic transfer using cellulose membrane, the membrane was incubated with monoclonal biotin-coupled rat anti-mouse C3, detected with a streptavidin-HRP kit
20 and developed using ECL reagent.

Results:

Activation of C3 involves cleavage of the intact α -chain into the truncated α' chain and soluble C3a (not shown in FIGURE 30). **FIGURE 30** shows the results of a Western blot analysis on the activation of human C3 by thrombin substrates, wherein the
25 uncleaved C3 α chain, and the activation product α' chain are shown by arrows. As shown in **FIGURE 30**, incubation of C3 with the activated forms of human clotting factor XI and factor X, as well as activated bovine clotting factor X, can cleave C3 in vitro in the absence of any complement proteases.

C3 deposition on mannan

C3 deposition assays were carried out on serum samples obtained from WT,
30 MASP-2 (-/-), F11(-/-), F11(-/-)/C4(-/-) and C4(-/-). F11 is the gene encoding coagulation factor XI. To measure C3 activation, microtiter plates were coated with mannan (1 μ g/well), then adding sheep anti-HSA serum (2 μ g/ml) in TBS/tween/ Ca^{2+} .

Plates were blocked with 0.1% HSA in TBS and washed as above. Plasma samples were diluted in 4 mM barbital, 145 mM NaCl, 2 mM CaCl₂, 1 mM MgCl₂, pH 7.4, added to the plates and incubated for 1.5 h at 37°C. After washing, bound C3b was detected using rabbit anti-human C3c (Dako), followed by alkaline phosphatase-conjugated goat anti-rabbit IgG and pNPP.

Results:

FIGURE 31 shows the results of the C3 deposition assay on serum samples obtained from WT, MASP-2 (-/-), F11(-/-), F11(-/-)/C4 (-/-) and C4 (-/-). As shown in **FIGURE 31**, there is a functional lectin pathway even in the complete absence of C4. As further shown in **FIGURE 31**, this novel lectin pathway dependent complement activation requires coagulation factor XI.

Discussion:

Prior to the results obtained in this experiment, it was believed by those in the art that the lectin pathway of complement required C4 for activity. Hence, data from C4 knockout mice (and C4 deficient humans) were interpreted with the assumption that such organisms were lectin pathway deficient (in addition to classical pathway deficiency). The present results demonstrate that this notion is false. Thus, conclusions of past studies suggesting that the lectin pathway was not important in certain disease settings based on the phenotype of C4 deficient animals may be false. The data described in this Example also show that in the physiological context of whole serum the lectin pathway can activate components of the coagulation cascade. Thus, it is demonstrated that there is cross-talk between complement and coagulation involving MASP-2.

EXAMPLE 24

This Example describes methods to assess the effect of an anti-MASP-2 antibody on lysis of red blood cells from blood samples obtained from Paroxysmal nocturnal hemoglobinuria (PNH) patients.

Background/Rationale:

Paroxysmal nocturnal hemoglobinuria (PNH), also referred to as Marchiafava-Micheli syndrome, is an acquired, potentially life-threatening disease of the blood, characterized by complement-induced intravascular hemolytic anemia. The hallmark of PNH is chronic intravascular hemolysis that is a consequence of unregulated activation of the alternative pathway of complement. Lindorfer, M.A., et al., *Blood* 115(11) (2010).

Anemia in PNH is due to destruction of red blood cells in the bloodstream. Symptoms of PNH include red urine, due to appearance of hemoglobin in the urine, and thrombosis. PNH may develop on its own, referred to as "primary PNH" or in the context of other bone marrow disorders such as aplastic anemia, referred to as "secondary PNH".

- 5 Treatment for PNH includes blood transfusion for anemia, anticoagulation for thrombosis and the use of the monoclonal antibody eculizumab (Soliris), which protects blood cells against immune destruction by inhibiting the complement system (Hillmen P. et al., *N. Engl. J. Med.* 350(6):552-9 (2004)). However, a significant portion of PNH patients treated with eculizumab are left with clinically significant immune-mediated hemolytic
10 anemia because the antibody does not block activation of the alternative pathway of complement.

This Example describes methods to assess the effect of an anti-MASP-2 antibody on lysis of red blood cells from blood samples obtained from PNH patients (not treated with Soliris) that are incubated with ABO-matched acidified normal human serum.

15 **Methods:**

Reagents:

- Erythrocytes from normal donors and from patients suffering from PNH (not treated with Soliris) are obtained by venipuncture, and prepared as described in Wilcox, L.A., et al., *Blood* 78:820-829 (1991), hereby incorporated herein by reference. Anti-
20 MASP-2 antibodies with functional blocking activity of the lectin pathway may be generated as described in Example 10.

Hemolysis Analysis:

- The method for determining the effect of anti-MASP-2 antibodies on the ability to block hemolysis of erythrocytes from PNH patients is carried out using the methods
25 described in Lindorfer, M.A., et al., *Blood* 15(11):2283-91 (2010) and Wilcox, L.A., et al., *Blood* 78:820-829 (1991), both references hereby incorporated herein by reference. As described in Lindorfer et al., erythrocytes from PNH patient samples are centrifuged, the buffy coat is aspirated and the cells are washed in gelatin veronal buffer (GVB) before each experiment. The erythrocytes are tested for susceptibility to APC-mediated
30 lysis as follows. ABO-matched normal human sera are diluted with GVB containing 0.15 mM CaCl_2 and 0.5 mM MgCl_2 (GVB^{+2}) and acidified to pH 6.4 (acidified NHS, aNHS) and used to reconstitute the erythrocytes to a hematocrit of 1.6% in 50% aNHS. The mixtures are then incubated at 37°C, and after 1 hour, the erythrocytes are pelleted by

centrifugation. The optical density of an aliquot of the recovered supernate is measured at 405 nM and used to calculate the percent lysis. Samples reconstituted in acidified serum-EDTA are processed similarly and used to define background noncomplement-mediated lysis (typically less than 3%). Complete lysis (100%) is determined after
5 incubating the erythrocytes in distilled water.

In order to determine the effect of anti-MASP-2 antibodies on hemolysis of PNH erythrocytes, erythrocytes from PNH patients are incubated in aNHS in the presence of incremental concentrations of the anti-MASP-2 antibodies, and the presence/amount of hemolysis is subsequently quantified.

10 In view of the fact that anti-MASP-2 antibodies have been shown to block subsequent activation of the alternative complement pathway, it is expected that anti-MASP-2 antibodies will be effective in blocking alternative pathway-mediated hemolysis of PNH erythrocytes, and will be useful as a therapeutic to treat patients suffering from PNH.

15

EXAMPLE 25

This Example describes methods to assess the effect of an anti-MASP-2 blocking antibody on complement activation by cryoglobulins in blood samples obtained from patients suffering from cryoglobulinemia.

20 **Background/Rationale:**

Cryoglobulinemia is characterized by the presence of cryoglobulins in the serum. Cryoglobulins are single or mixed immunoglobulins (typically IgM antibodies) that undergo reversible aggregation at low temperatures. Aggregation leads to classical pathway complement activation and inflammation in vascular beds, particularly in the
25 periphery. Clinical presentations of cryoglobulinemia include vasculitis and glomerulonephritis.

Cryoglobulinemia may be classified as follows based on cryoglobulin composition: Type I cryoglobulinemia, or simple cryoglobulinemia, is the result of a monoclonal immunoglobulin, usually immunoglobulin M (IgM); Types II and III
30 cryoglobulinemia (mixed cryoglobulinemia) contain rheumatoid factors (RFs), which are usually IgM in complexes with the Fc portion of polyclonal IgG.

Conditions associated with cryoglobulinemia include hepatitis C infection, lymphoproliferative disorders and other autoimmune diseases. Cryoglobulin-containing

immune complexes result in a clinical syndrome of systemic inflammation, possibly due to their ability to activate complement. While IgG immune complexes normally activate the classical pathway of complement, IgM containing complexes can also activate complement via the lectin pathway (Zhang, M., et al., *Mol Immunol* 44(1-3):103-110 (2007) and Zhang, M., et al., *J. Immunol.* 177(7):4727-34 (2006)).

Immunohistochemical studies have further demonstrated the cryoglobulin immune complexes contain components of the lectin pathway, and biopsies from patients with cryoglobulinemic glomerulonephritis showed immunohistochemical evidence of lectin pathway activation in situ (Ohsawa, I., et al., *Clin Immunol* 101(1):59-66 (2001)).

These results suggest that the lectin pathway may contribute to inflammation and adverse outcomes in cryoglobulemic diseases.

Methods:

The method for determining the effect of anti-MASP-2 antibodies on the ability to block the adverse effects of Cryoglobulinemia is carried out using the assay for fluid phase C3 conversion as described in Ng Y.C. et al., *Arthritis and Rheumatism* 31(1):99-107 (1988), hereby incorporated herein by reference. As described in Ng et al., in essential mixed cryoglobulinemia (EMC), monoclonal rheumatoid factor (mRF), usually IgM, complexes with polyclonal IgG to form the characteristic cryoprecipitate immune complexes (IC) (type II cryoglobulin). Immunoglobulins and C3 have been demonstrated in vessel walls in affected tissues such as skin, nerve and kidney. As described in Ng et al., ¹²⁵I-labeled mRF is added to serum (normal human serum and serum obtained from patients suffering from cryoglobulinemia), incubated at 37°C, and binding to erythrocytes is measured.

Fluid phase C3 conversion is determined in serum (normal human serum and serum obtained from patients suffering from cryoglobulinemia) in the presence or absence of the following IC: BSA-anti BSA, mRF, mRF plus IgG, or cryoglobulins, in the presence or absence of anti-MASP-2 antibodies. The fixation of C3 and C4 to IC is measured using a coprecipitation assay with F(ab')₂ anti-C3 and F(ab')₂ anti-C4.

In view of the fact that anti-MASP-2 antibodies have been shown to block activation of the lectin pathway it is expected that anti-MASP-2 antibodies will be effective in blocking complement mediated adverse effects associated with cryoglobulinemia, and will be useful as a therapeutic to treat patients suffering from cryoglobulinemia.

EXAMPLE 26

This Example describes methods to assess the effect of an anti-MASP-2 antibody on blood samples obtained from patients with Cold Agglutinin Disease, which manifests
5 as anemia.

Background/Rationale:

Cold Agglutinin Disease (CAD), is a type of autoimmune hemolytic anemia. Cold agglutinins antibodies (usually IgM) are activated by cold temperatures and bind to and aggregate red blood cells. The cold agglutinin antibodies combine with complement
10 and attack the antigen on the surface of red blood cells. This leads to opsoniation of red blood cells (hemolysis) which triggers their clearance by the reticuloendothelial system. The temperature at which the agglutination takes place varies from patient to patient.

CAD manifests as anemia. When the rate of destruction of red blood cell destruction exceeds the capacity of the bone marrow to produce an adequate number of
15 oxygen-carrying cells, then anemia occurs. CAD can be caused by an underlying disease or disorder, referred to as "Secondary CAD", such as an infectious disease (mycoplasma pneumonia, mumps, mononucleosis), lymphoproliferative disease (lymphoma, chronic lymphocytic leukemia), or connective tissue disorder. Primary CAD patients are considered to have a low grade lymphoproliferative bone marrow disorder. Both primary
20 and secondary CAD are acquired conditions.

Methods:**Reagents:**

Erythrocytes from normal donors and from patients suffering from CAD are obtained by venipuncture. Anti-MASP-2 antibodies with functional blocking activity of
25 the lectin pathway may be generated as described in Example 10.

The effect of anti-MASP-2 antibodies to block cold agglutinin-mediated activation of the lectin pathway may be determined as follows. Erythrocytes from blood group I positive patients are sensitized with cold agglutinins (i.e., IgM antibodies), in the presence or absence of anti-MASP-2 antibodies. The erythrocytes are then tested for the
30 ability to activate the lectin pathway by measuring C3 binding.

In view of the fact that anti-MASP-2 antibodies have been shown to block activation of the lectin pathway, it is expected that anti-MASP-2 antibodies will be effective in blocking complement mediated adverse effects associated with Cold

Agglutinin Disease, and will be useful as a therapeutic to treat patients suffering from Cold Agglutinin Disease.

EXAMPLE 27

5 This Example describes methods to assess the effect of an anti-MASP-2 antibody on lysis of red blood cells in blood samples obtained from mice with atypical hemolytic uremic syndrome (aHUS).

Background/Rationale:

Atypical hemolytic uremic syndrome (aHUS) is characterized by hemolytic
10 anemia, thrombocytopenia, and renal failure caused by platelet thrombi in the microcirculation of the kidney and other organs. aHUS is associated with defective complement regulation and can be either sporadic or familial. aHUS is associated with mutations in genes coding for complement activation, including complement factor H, membrane cofactor B and factor I, and well as complement factor H-related 1 (CFHR1)
15 and complement factor H-related 3 (CFHR3). Zipfel, P.F., et al., *PloS Genetics* 3(3):e41 (2007). This Example describes methods to assess the effect of an anti-MASP-2 antibody on lysis of red blood cells from blood samples obtained from aHUS mice.

Methods:

The effect of anti-MASP-2 antibodies to treat aHUS may be determined in a
20 mouse model of this disease in which the endogenous mouse fH gene has been replaced with a human homologue encoding a mutant form of fH frequently found in aHUS patients. See Pickering M.C. et al., *J. Exp. Med.* 204(6):1249-1256 (2007), hereby incorporated herein by reference. As described in Pickering et al., such mice develop an aHUS like pathology. In order to assess the effect of an anti-MASP-2 antibody for the
25 treatment of aHUS, anti-MASP-2 antibodies are administered to the mutant aHUS mice and lysis of red blood cells obtained from anti-MASP-2 ab treated and untreated controls is compared. In view of the fact that anti-MASP-2 antibodies have been shown to block activation of the lectin pathway it is expected that anti-MASP-2 antibodies will be effective in blocking lysis of red blood cells in mammalian subjects suffering from aHUS.

30

EXAMPLE 28

This Example describes methods to assess the effect of an anti-MASP-2 antibody for the treatment of glaucoma.

Rationale/Background:

It has been shown that uncontrolled complement activation contributes to the progression of degenerative injury to retinal ganglion cells (RGCs), their synapses and axons in glaucoma. See Tezel G. et al., *Invest Ophthalmol Vis Sci* 51:5071-5082 (2010).
5 For example, histopathologic studies of human tissues and *in vivo* studies using different animal models have demonstrated that complement components, including C1q and C3, are synthesized and terminal complement complex is formed in the glaucomatous retina (see Stasi K. et al., *Invest Ophthalmol Vis Sci* 47:1024-1029 (2006), Kuehn M.H. et al., *Exp Eye Res* 83:620-628 (2006)). As further described in Kuehn M.H. et al.,
10 *Experimental Eye Research* 87:89-95 (2008), complement synthesis and deposition is induced by retinal I/R and the disruption of the complement cascade delays RGC degeneration. In this study, mice carrying a targeted disruption of the complement component C3 were found to exhibit delayed RGC degeneration after transient retinal I/R when compared to normal animals.

Methods:

The method for determining the effect of anti-MASP-2 antibodies on RGC degeneration is carried out in an animal model of retinal I/R as described in Kuehn M.H. et al., *Experimental Eye Research* 87:89-95 (2008), hereby incorporated herein by reference. As described in Kuehn et al., retinal ischemia is induced by anesthetizing the
20 animals, then inserting a 30-gauge needle connected to a reservoir containing phosphate buffered saline through the cornea into the anterior chamber of the eye. The saline reservoir is then elevated to yield an intraocular pressure of 104 mmHg, sufficient to completely prevent circulation through the retinal vasculature. Elevated intraocular ischemia is confirmed by blanching of the iris and retina and ischemia is maintained for
25 45 minutes in the left eye only; the right eye serves as a control and does not receive cannulation. Mice are then euthanized either 1 or 3 weeks after the ischemic insult. Anti-MASP-2 antibodies are administered to the mice either locally to the eye or systemically to assess the effect of an anti-MASP antibody administered prior to ischemic insult.

Immunohistochemistry of the eyes is carried out using antibodies against C1q and
30 C3 to detect complement deposition. Optic nerve damage can also be assessed using standard electron microscopy methods. Quantitation of surviving retinal RGCs is performed using gamma synuclein labeling.

Results:

As described in Kuehn et al., in normal control mice, transient retinal ischemia results in degenerative changes of the optic nerve and retinal deposits of C1q and C3 detectable by immunohistochemistry. In contrast, C3 deficient mice displayed a marked reduction in axonal degeneration, exhibiting only minor levels of optic nerve damage 1 week after induction. Based on these results, it is expected that similar results would be observed when this assay is carried out in a MASP-2 knockout mouse, and when anti-MASP-2 antibodies are administered to a normal mouse prior to ischemic insult.

10

EXAMPLE 29

This Example demonstrates that a MASP-2 inhibitor, such as an anti-MASP-2 antibody, is effective for the treatment of radiation exposure and/or for the treatment, amelioration or prevention of acute radiation syndrome.

Rationale:

15

Exposure to high doses of ionizing radiation causes mortality by two main mechanisms: toxicity to the bone marrow and gastrointestinal syndrome. Bone marrow toxicity results in a drop in all hematologic cells, predisposing the organism to death by infection and hemorrhage. The gastrointestinal syndrome is more severe and is driven by a loss of intestinal barrier function due to disintegration of the gut epithelial layer and a loss of intestinal endocrine function. This leads to sepsis and associated systemic inflammatory response syndrome which can result in death.

20

The lectin pathway of complement is an innate immune mechanism that initiates inflammation in response to tissue injury and exposure to foreign surfaces (i.e., bacteria). Blockade of this pathway leads to better outcomes in mouse models of ischemic intestinal tissue injury or septic shock. It is hypothesized that the lectin pathway may trigger excessive and harmful inflammation in response to radiation-induced tissue injury. Blockade of the lectin pathway may thus reduce secondary injury and increase survival following acute radiation exposure.

25

The objective of the study carried out as described in this Example was to assess the effect of lectin pathway blockade on survival in a mouse model of radiation injury by administering anti-murine MASP-2 antibodies.

30

Methods and Materials:

Materials. The test articles used in this study were (i) a high affinity anti-murine MASP-2 antibody (mAbM11) and (ii) a high affinity anti-human MASP-2 antibody (mAbH6) that block the MASP-2 protein component of the lectin complement pathway which were produced in transfected mammalian cells. Dosing concentrations were 1 mg/kg of anti-murine MASP-2 antibody (mAbM11), 5mg/kg of anti-human MASP-2 antibody (mAbH6), or sterile saline. For each dosing session, an adequate volume of fresh dosing solutions were prepared.

Animals. Young adult male Swiss-Webster mice were obtained from Harlan Laboratories (Houston, TX). Animals were housed in solid-bottom cages with Alpha-Dri bedding and provided certified PMI 5002 Rodent Diet (Animal Specialties, Inc., Hubbard OR) and water ad libitum. Temperature was monitored and the animal holding room operated with a 12 hour light/12 hour dark light cycle.

Irradiation. After a 2-week acclimation in the facility, mice were irradiated at 6.5 and 7.0 Gy by whole-body exposure in groups of 10 at a dose rate of 0.78 Gy/min using a Therapax X-RAD 320 system equipped with a 320-kV high stability X-ray generator, metal ceramic X-ray tube, variable x-ray beam collimator and filter (Precision X-ray Incorporated, East Haven, CT). Dose levels were selected based on prior studies conducted with the same strain of mice indicating the LD_{50/30} was between 6.5 and 7.0 Gy (data not shown).

Drug Formulation and Administration. The appropriate volume of concentrated stock solutions were diluted with ice cold saline to prepare dosing solutions of 0.2 mg/ml anti-murine MASP-2 antibody (mAbM11) or 0.5 mg/ml anti-human MASP-2 antibody (mAbH6) according to protocol. Administration of anti-MASP-2 antibody mAbM11 and mAbH6 was via IP injection using a 25-gauge needle base on animal weight to deliver 1 mg/kg mAbM11, 5mg/kg mAbH6, or saline vehicle.

Study Design. Mice were randomly assigned to the groups as described in Table 8. Body weight and temperature were measured and recorded daily. Mice in Groups 7, 11 and 13 were sacrificed at post-irradiation day 7 and blood collected by cardiac puncture under deep anesthesia. Surviving animals at post-irradiation day 30 were sacrificed in the same manner and blood collected. Plasma was prepared from collected blood samples according to protocol and returned to Sponsor for analysis.

TABLE 8: Study Groups

Group ID	N	Irradiation Level (Gy)	Treatment	Dose Schedule
1	20	6.5	Vehicle	18 hr prior to irradiation, 2 hr post irradiation, weekly booster
2	20	6.5	anti-murine MASP-2 ab (mAbM11)	18 hr prior to irradiation only
3	20	6.5	anti-murine MASP-2 ab (mAbM11)	18 hr prior to irradiation, 2 hr post irradiation, weekly booster
4	20	6.5	anti-murine MASP-2 ab (mAbM11)	2 hr post irradiation, weekly booster
5	20	6.5	anti-human MASP-2 ab (mAbH6)	18 hr prior to irradiation, 2 hr post irradiation, weekly booster
6	20	7.0	Vehicle	18 hr prior to irradiation, 2 hr post irradiation, weekly booster
7	5	7.0	Vehicle	2 hr post irradiation only
8	20	7.0	anti-murine MASP-2 ab (mAbM11)	18 hr prior to irradiation only
9	20	7.0	anti-murine MASP-2 ab (mAbM11)	18 hr prior to irradiation, 2 hr post irradiation, weekly booster
10	20	7.0	anti-murine MASP-2 ab (mAbM11)	2 hr post irradiation, weekly booster
11	5	7.0	anti-murine MASP-2 ab (mAbM11)	2 hr post irradiation only
12	20	7.0	anti-human MASP-2 ab (mAbH6)	18 hr prior to irradiation, 2 hr post irradiation, weekly booster
13	5	None	None	None

Statistical Analysis. Kaplan-Meier survival curves were generated and used to compare mean survival time between treatment groups using log-Rank and Wilcoxon methods. Averages with standard deviations, or means with standard error of the mean are reported. Statistical comparisons were made using a two-tailed unpaired t-test
5 between controlled irradiated animals and individual treatment groups.

Results

Kaplan-Meier survival plots for 7.0 and 6.5 Gy exposure groups are provided in **FIGURES 32A** and **32B**, respectively, and summarized below in Table 9. Overall, treatment with anti-murine MASP-2 ab (mAbM11) pre-irradiation increased the survival
10 of irradiated mice compared to vehicle treated irradiated control animals at both 6.5 (20% increase) and 7.0 Gy (30% increase) exposure levels. At the 6.5 Gy exposure level, post-irradiation treatment with anti-murine MASP-2 ab resulted in a modest increase in survival (15%) compared to vehicle control irradiated animals.

In comparison, all treated animals at the 7.0 Gy exposure level showed an
15 increase in survival compared to vehicle treated irradiated control animals. The greatest change in survival occurred in animals receiving mAbH6, with a 45% increase compared to control animals. Further, at the 7.0 Gy exposure level, mortalities in the mAbH6 treated group first occurred at post-irradiation day 15 compared to post-irradiation day 8 for vehicle treated irradiated control animals, an increase of 7 days over control animals.
20 Mean time to mortality for mice receiving mAbH6 (27.3 ± 1.3 days) was significantly increased ($p = 0.0087$) compared to control animals (20.7 ± 2.0 days) at the 7.0 Gy exposure level.

The percent change in body weight compared to pre-irradiation day (day -1) was recorded throughout the study. A transient weight loss occurred in all irradiated animals,
25 with no evidence of differential changes due to mAbM11 or mAbH6 treatment compared to controls (data not shown). At study termination, all surviving animals showed an increase in body weight from starting (day -1) body weight.

TABLE 9: Survival rates of test animals exposed to radiation

Test Group	Exposure Level	Survival (%)	Time to Death (Mean $\pm$ SEM, Day)	First/Last Death (Day)
Control Irradiation	6.5 Gy	65 %	24.0 ± 2.0	9/16
mAbM11 pre-	6.5 Gy	85 %	27.7 ± 1.5	13/17

Test Group	Exposure Level	Survival (%)	Time to Death (Mean $\pm$ SEM, Day)	First/Last Death (Day)
exposure				
mAbM11 pre + post-exposure	6.5 Gy	65 %	24.0 $\pm$ 2.0	9/15
mAbM11 post-exposure	6.5 Gy	80 %	26.3 $\pm$ 1.9	9/13
mAbH6 pre+post-exposure	6.5 Gy	65 %	24.6 $\pm$ 1.9	9/19
Control irradiation	7.0 Gy	35 %	20.7 $\pm$ 2.0	8/17
mAbM11 pre-exposure	7.0 Gy	65 %	23.0 $\pm$ 2.3	7/13
mAbM11 pre + post-exposure	7.0 Gy	55 %	21.6 $\pm$ 2.2	7/16
mAbM11 post-exposure	7.0 Gy	70 %	24.3 $\pm$ 2.1	9/14
mAbH6 pre+post-exposure	7.0 Gy	80 %	27.3 $\pm$ 1.3*	15/20

*p = 0.0087 by two-tailed unpaired t-test between controlled irradiated animals and treatment group at the same irradiation exposure level.

Discussion

Acute radiation syndrome consists of three defined subsyndromes: hematopoietic, gastrointestinal, and cerebrovascular. The syndrome observed depends on the radiation dose, with the hematopoietic effects observed in humans with significant partial or whole-body radiation exposures exceeding 1 Gy. The hematopoietic syndrome is characterized by severe depression of bone-marrow function leading to pancytopenia with changes in blood counts, red and white blood cells, and platelets occurring concomitant with damage to the immune system. As nadir occurs, with few neutrophils and platelets present in peripheral blood, neutropenia, fever, complications of sepsis and uncontrollable hemorrhage lead to death.

In the present study, administration of mAbH6 was found to increase survivability of whole-body x-ray irradiation in Swiss-Webster male mice irradiated at 7.0 Gy. Notably, at the 7.0 Gy exposure level, 80% of the animals receiving mAbH6 survived to 30 days compared to 35% of vehicle treated control irradiated animals. Importantly, the first day of death in this treated group did not occur until post-irradiation day 15, a 7-day increase over that observed in vehicle treated control irradiated animals. Curiously, at the

lower X-ray exposure (6.5 Gy), administration of mAbH6 did not appear to impact survivability or delay in mortality compared to vehicle treated control irradiated animals. There could be multiple reasons for this difference in response between exposure levels, although verification of any hypothesis may require additional studies, including interim
5 sample collection for microbiological culture and hematological parameters. One explanation may simply be that the number of animals assigned to groups may have precluded seeing any subtle treatment-related differences. For example, with groups sizes of $n=20$, the difference in survival between 65% (mAbH6 at 6.5 Gy exposure) and 80% (mAbH6 at 7.0 Gy exposure) is 3 animals. On the other hand, the difference
10 between 35% (vehicle control at 7.0 Gy exposure) and 80% (mAbH6 at 7.0 Gy exposure) is 9 animals, and provides sound evidence of a treatment-related difference.

These results demonstrate that anti-MASP-2 antibodies are effective in treating a mammalian subject at risk for, or suffering from the detrimental effects of acute radiation syndrome.

15

EXAMPLE 30

This Example demonstrates that MASP-2 deficient mice are protected from *Neisseria meningitidis* induced mortality after infection with either *N. meningitidis* serogroup A or *Neisseria meningitidis* serogroup B.

20 Methods:

MASP-2 knockout mice (MASP-2 KO mice) were generated as described in Example 1. 10-week-old MASP-2 KO mice ($n=10$) and wild-type (WT) C57/BL6 mice ($n=10$) were inoculated by intraperitoneal (i.p.) injection with a dosage of 2.6×10^7 CFU of *Neisseria meningitidis* serogroup A Z2491 in a volume of 100 μ l. The infective dose
25 was administered to mice in conjunction with iron dextran at a final concentration of 400 mg/kg. Survival of the mice after infection was monitored over a 72-hour time period.

In a separate experiment, 10-week-old MASP-2 KO mice ($n=10$) and wild-type C57/BL6 mice ($n=10$) were inoculated by i.p. injection with a dosage of 6×10^6 CFU of
30 *Neisseria meningitidis* serogroup B strain MC58 in a volume of 100 μ l. The infective dose was administered to mice in conjunction with iron dextran at a final dose of 400 mg/kg. Survival of the mice after infection was monitored over a 72-hour time period. An illness score was also determined for the WT and MASP-2 KO mice during

the 72-hour time period after infection, based on the illness scoring parameters described below in **TABLE 10**, which is based on the scheme of Fransen et al. (2010) with slight modifications.

5 **TABLE 10: Illness Scoring associated with clinical signs in infected mice**

Signs	Score
Normal	0
Slightly ruffled fur	1
Ruffled fur, slow and sticky eyes	2
Ruffled fur, lethargic and eyes shut	3
Very sick and no movement after stimulation	4
Dead	5

Blood samples were taken from the mice at hourly intervals after infection and analyzed to determine the serum level (log cfu/mL) of *N. meningitidis* in order to verify infection and determine the rate of clearance of the bacteria from the serum.

10 **Results:**

FIGURE 33 is a Kaplan-Meyer plot graphically illustrating the percent survival of MASP-2 KO and WT mice after administration of an infective dose of 2.6×10^7 cfu of *N. meningitidis* serogroup A Z2491. As shown in **FIGURE 33**, 100% of the MASP-2 KO mice survived throughout the 72-hour period after infection. In contrast, only 80% of the WT mice ($p=0.012$) were still alive 24 hours after infection, and only 50% of the WT mice were still alive at 72 hours after infection. These results demonstrate that MASP-2-deficient mice are protected from *N. meningitidis* serogroup A Z2491-induced mortality.

FIGURE 34 is a Kaplan-Meyer plot graphically illustrating the percent survival of MASP-2 KO and WT mice after administration of an infective dose of 6×10^6 cfu of *N. meningitidis* serogroup B strain MC58. As shown in **FIGURE 34**, 90% of the MASP-2 KO mice survived throughout the 72-hour period after infection. In contrast, only 20% of the WT mice ($p=0.0022$) were still alive 24 hours after infection. These results demonstrate that MASP-2-deficient mice are protected from *N. meningitidis* serogroup B strain MC58-induced mortality.

FIGURE 35 graphically illustrates the log cfu/mL of *N. meningitidis* serogroup B strain MC58 recovered at different time points in blood samples taken from the MASP-2 KO and WT mice after i.p. infection with 6×10^6 cfu of *N. meningitidis* serogroup B strain MC58 (n=3 at different time points for both groups of mice). The results are expressed as Means $\pm$ SEM. As shown in **FIGURE 35**, in WT mice the level of *N. meningitidis* in the blood reached a peak of about 6.0 log cfu/mL at 24 hours after infection and dropped to about 4.0 log cfu/mL by 36 hours after infection. In contrast, in the MASP-2 KO mice, the level of *N. meningitidis* reached a peak of about 4.0 log cfu/mL at 12 hours after infection and dropped to about 1.0 log cfu/mL by 36 hours after infection (the symbol "*" indicates $p < 0.05$; the symbol "***" indicates $p = 0.0043$). These results demonstrate that although the MASP-2 KO mice were infected with the same dose of *N. meningitidis* serogroup B strain MC58 as the WT mice, the MASP-2 KO mice have enhanced clearance of bacteraemia as compared to WT.

FIGURE 36 graphically illustrates the average illness score of MASP-2 KO and WT mice at 3, 6, 12 and 24 hours after infection with 6×10^6 cfu of *N. meningitidis* serogroup B strain MC58. As shown in **FIGURE 36**, the MASP-2-deficient mice showed high resistance to the infection, with much lower illness scores at 6 hours (symbol "*" indicates $p = 0.0411$), 12 hours (symbol "***" indicates $p = 0.0049$) and 24 hours (symbol "****" indicates $p = 0.0049$) after infection, as compared to WT mice. The results in **FIGURE 36** are expressed as means $\pm$ SEM.

In summary, the results in this Example demonstrate that MASP-2-deficient mice are protected from *Neisseria meningitidis*-induced mortality after infection with either *N. meningitidis* serogroup A or *N. meningitidis* serogroup B.

25

EXAMPLE 31

This Example demonstrates that the administration of anti-MASP-2 antibody after infection with *N. meningitidis* increases the survival of mice infected with *N. meningitidis*.

Background/Rationale:

30

As described in Example 10, rat MASP-2 protein was utilized to pan a Fab phage display library, from which Fab2 #11 was identified as a functionally active antibody. Full-length antibodies of the rat IgG2c and mouse IgG2a isotypes were generated from

Fab2 #11. The full-length anti-MASP-2 antibody of the mouse IgG2a isotype was characterized for pharmacodynamic parameters (as described in Example 19).

In this Example, the mouse anti-MASP-2 full-length antibody derived from Fab2 #11 was analyzed in the mouse model of *N. meningitidis* infection.

5 **Methods:**

The mouse IgG2a full-length anti-MASP-2 antibody isotype derived from Fab2 #11, generated as described above, was tested in the mouse model of *N. meningitidis* infection as follows.

10 ***Administration of mouse-anti-MASP-2 Monoclonal antibodies (MoAb) after infection***

9-week-old C57/BL6 Charles River mice were treated with inhibitory mouse anti-MASP-2 antibody (1.0 mg/kg) (n=12) or control isotype antibody (n=10) at 3 hours after i.p. injection with a high dose (4×10^6 cfu) of *N. meningitidis* serogroup B strain MC58.

15 **Results:**

15 **FIGURE 37** is a Kaplan-Meyer plot graphically illustrating the percent survival of mice after administration of an infective dose of 4×10^6 cfu of *N. meningitidis* serogroup B strain MC58, followed by administration 3 hours post-infection of either inhibitory anti-MASP-2 antibody (1.0 mg/kg) or control isotype antibody. As shown in **FIGURE 37**, 90% of the mice treated with anti-MASP-2 antibody survived throughout
20 the 72-hour period after infection. In contrast, only 50% of the mice treated with isotype control antibody survived throughout the 72-hour period after infection. The symbol "*" indicates $p=0.0301$, as determined by comparison of the two survival curves.

These results demonstrate that administration of anti-MASP-2 antibody is effective to treat and improve survival in subjects infected with *N. meningitidis*.

25 As demonstrated herein, the use of anti-MASP-2 antibody in the treatment of a subject infected with *N. meningitidis* is effective when administered within 3 hours post-infection, and is expected to be effective within 24 hours to 48 hours after infection. Meningococcal disease (either meningococcemia or meningitis) is a medical emergency, and therapy will typically be initiated immediately if meningococcal disease is suspected
30 (i.e., before *N. meningitidis* is positively identified as the etiological agent).

In view of the results in the MASP-2 KO mouse demonstrated in **EXAMPLE 30**, it is believed that administration of anti-MASP-2 antibody prior to infection with *N. meningitidis* would also be effective to prevent or ameliorate the severity of infection.

EXAMPLE 32

This Example demonstrates that administration of anti-MASP-2 antibody is effective to treat *N. meningitidis* infection in human serum.

Rationale:

Patients with decreased serum levels of functional MBL display increased susceptibility to recurrent bacterial and fungal infections (Kilpatrick et al., *Biochim Biophys Acta* 1572:401-413 (2002)). It is known that *N. meningitidis* is recognized by MBL, and it has been shown that MBL-deficient sera do not lyse *Neisseria*.

In view of the results described in Examples 30 and 31, a series of experiments were carried out to determine the efficacy of administration of anti-MASP-2 antibody to treat *N. meningitidis* infection in complement-deficient and control human sera. Experiments were carried out in a high concentration of serum (20%) in order to preserve the complement pathway.

Methods:

1. Serum bactericidal activity in various complement-deficient human sera and in human sera treated with human anti-MASP-2 antibody

The following complement-deficient human sera and control human sera were used in this experiment:

TABLE 11: Human sera samples tested (as shown in FIGURE 38)

Sample	Serum type
A	Normal human sera (NHS) + human anti-MASP-2 Ab
B	NHS + isotype control Ab
C	MBL -/- human serum
D	NHS
E	Heat-Inactivated (HI) NHS

A recombinant antibody against human MASP-2 was isolated from a Combinatorial Antibody Library (Knappik, A., et al., *J. Mol. Biol.* 296:57-86 (2000)), using recombinant human MASP-2A as an antigen (Chen, C.B. and Wallis, *J. Biol. Chem.* 276:25894-25902 (2001)). An anti-human scFv fragment that potently inhibited lectin pathway-mediated activation of C4 and C3 in human plasma (IC₅₀~20 nM) was identified and converted to a full-length human IgG4 antibody.

N. meningitidis serogroup B-MC58 was incubated with the different sera shown in **TABLE 11**, each at a serum concentration of 20%, with or without the addition of inhibitory human anti-MASP-2 antibody (3 µg in 100 µl total volume) at 37°C with shaking. Samples were taken at the following time points: 0-, 30-, 60- and 90-minute intervals, plated out and then viable counts were determined. Heat-inactivated human serum was used as a negative control.

Results:

FIGURE 38 graphically illustrates the log cfu/mL of viable counts of *N. meningitidis* serogroup B-MC58 recovered at different time points in the human sera samples shown in **TABLE 11**. **TABLE 12** provides the Student's t-test results for **FIGURE 38**.

TABLE 12: Student's t-test Results for FIGURE 38 (time point 60 minutes)

	Mean Diff. (Log)	Significant? P<0.05?	P value summary
A vs B	-0.3678	Yes	*** (0.0002)
A vs C	-1.1053	Yes	*** (p<0.0001)
A vs D	-0.2111	Yes	** (0.0012)
C vs D	1.9	Yes	*** (p<0.0001)

As shown in **FIGURE 38** and **TABLE 12**, complement-dependent killing of *N. meningitidis* in human 20% serum was significantly enhanced by the addition of the human anti-MASP-2 inhibitory antibody.

2. Complement-dependent killing of *N. meningitidis* in 20% (v/v) mouse sera deficient of MASP-2.

The following complement-deficient mouse sera and control mouse sera were used in this experiment:

TABLE 13: Mouse sera samples tested (as shown in FIGURE 39)

Sample	Serum Type
A	WT
B	MASP-2 -/-
C	MBL A/C -/-

D	WT heat-inactivated (HIS)
---	---------------------------

N. meningitidis serogroup B-MC58 was incubated with different complement-deficient mouse sera, each at a serum concentration of 20%, at 37°C with shaking. Samples were taken at the following time points: 0-, 15-, 30-, 60-, 90- and 120-minute intervals, plated out and then viable counts were determined. Heat-inactivated human serum was used as a negative control.

Results:

FIGURE 39 graphically illustrates the log cfu/mL of viable counts of *N. meningitidis* serogroup B-MC58 recovered at different time points in the mouse sera samples shown in **TABLE 13**. As shown in **FIGURE 39**, the MASP-2 ^{-/-} mouse sera have a higher level of bactericidal activity for *N. meningitidis* than WT mouse sera. The symbol "***" indicates p=0.0058, the symbol "****" indicates p=0.001. **TABLE 14** provides the Student's t-test results for **FIGURE 39**.

TABLE 14: Student's t-test Results for FIGURE 39

Comparison	Time point	Mean Diff. (LOG)	Significant? (p<0.05)?	P value summary
A vs. B	60 min.	0.39	yes	** (0.0058)
A vs. B	90 min.	0.6741	yes	*** (0.001)

In summary, the results in this Example demonstrate that MASP-2 ^{-/-} sera has a higher level of bactericidal activity for *N. meningitidis* than WT sera.

EXAMPLE 33

This Example demonstrates the inhibitory effect of MASP-2 deficiency on lysis of red blood cells from blood samples obtained from a mouse model of paroxysmal nocturnal hemoglobinuria (PNH).

Background/Rationale:

Paroxysmal nocturnal hemoglobinuria (PNH), also referred to as Marchiafava-Micheli syndrome, is an acquired, potentially life-threatening disease of the blood, characterized by complement-induced intravascular hemolytic anemia. The hallmark of PNH is the chronic complement-mediated intravascular hemolysis that is a consequence

of unregulated activation of the alternative pathway of complement due to the absence of the complement regulators CD55 and CD59 on PNH erythrocytes, with subsequent hemoglobinuria and anemia. Lindorfer, M.A., et al., *Blood* 115(11) (2010), Risitano, A.M., *Mini-Reviews in Medicinal Chemistry*, 11:528-535 (2011). Anemia in PNH is due to destruction of red blood cells in the bloodstream. Symptoms of PNH include red urine, due to appearance of hemoglobin in the urine, back pain, fatigue, shortness of breath and thrombosis. PNH may develop on its own, referred to as "primary PNH" or in the context of other bone marrow disorders such as aplastic anemia, referred to as "secondary PNH". Treatment for PNH includes blood transfusion for anemia, anticoagulation for thrombosis and the use of the monoclonal antibody eculizumab (Soliris®), which protects blood cells against immune destruction by inhibiting the complement system (Hillmen P. et al., *N. Engl. J. Med.* 350(6):552-9 (2004)). Eculizumab (Soliris®) is a humanized monoclonal antibody that targets the complement component C5, blocking its cleavage by C5 convertases, thereby preventing the production of C5a and the assembly of the MAC. Treatment of PNH patients with eculizumab has resulted in a reduction of intravascular hemolysis, as measured by lactate dehydrogenase (LDH), leading to hemoglobin stabilization and transfusion independence in about half of the patients (Hillmen P, et al., *Mini-Reviews in Medicinal Chemistry*, vol 11(6) (2011)). While nearly all patients undergoing therapy with eculizumab achieve normal or almost normal LDH levels (due to control of intravascular hemolysis), only about one third of the patients reach a hemoglobin value above 11gr/dL, and the remaining patients on eculizumab continue to exhibit moderate to severe (*i.e.*,transfusion-dependent) anemia, in about equal proportions (Risitano A.M. et al., *Blood* 113:4094-100 (2009)). As described in Risitano et al., *Mini-Reviews in Medicinal Chemistry* 11:528-535 (2011), it was demonstrated that PNH patients on eculizumab contained C3 fragments bound to a substantial portion of their PNH erythrocytes (while untreated patients did not), leading to the conclusion that membrane-bound C3 fragments work as opsonins on PNH erythrocytes, resulting in their entrapment in the reticuloendothelial cells through specific C3 receptors and subsequent extravascular hemolysis. Therefore, therapeutic strategies in addition to the use of eculizumab are needed for those patients developing C3 fragment-mediated extravascular hemolysis because they continue to require red cell transfusions.

This Example describes methods to assess the effect of MASP-2- deficient serum and serum treated with MASP-2 inhibitory agent on lysis of red blood cells from blood

samples obtained from a mouse model of PNH and demonstrates the efficacy of MASP-2 inhibition to treat subjects suffering from PNH, and also supports the use of inhibitors of MASP-2 to ameliorate the effects of C3 fragment-mediated extravascular hemolysis in PNH subjects undergoing therapy with a C5 inhibitor such as eculizumab.

5 **Methods:**

PNH animal model:

Blood samples were obtained from gene-targeted mice with deficiencies of Crry and C3 (Crry/C3^{-/-}) and CD55/CD59-deficient mice. These mice are missing the respective surface complement regulators and their erythrocytes are, therefore,
10 susceptible to spontaneous complement autolysis as are PNH human blood cells.

In order to sensitize these erythrocytes even more, these cells were used with and without coating by mannan and then tested for hemolysis in WT C56/BL6 plasma, MBL null plasma, MASP-2 ^{-/-} plasma, human NHS, human MBL ^{-/-} plasma, and NHS treated with human anti-MASP-2 antibody.
15

1. Hemolysis assay of Crry/C3 and CD55/CD59 double-deficient murine erythrocytes in MASP-2-deficient/depleted sera and controls

Day 1. Preparation of murine RBC (± mannan coating)

20 **Materials included:** fresh mouse blood, BBS/Mg²⁺/Ca²⁺ (4.4 mM barbituric acid, 1.8 mM sodium barbitone, 145 mM NaCl, pH7.4, 5mM Mg²⁺, 5mM Ca²⁺), chromium chloride, CrCl₃·6H₂O (0.5mg/mL in BBS/Mg²⁺/Ca²⁺) and mannan, 100 µg/mL in BBS /Mg²⁺/Ca²⁺.

Whole blood (2mL) was spun down for 1-2 min at 2000xg in a refrigerated
25 centrifuge at 4°C. The plasma and buffy coat were aspirated off. The sample was then washed three times by re-suspending the RBC pellet in 2 mL ice-cold BBS/gelatin/Mg²⁺/Ca²⁺ and repeating centrifugation step. After the third wash, the pellet was re-suspended in 4mL BBS/Mg²⁺/Ca²⁺. A 2 mL aliquot of the RBC was set aside as an uncoated control. To the remaining 2 mL, 2 mL CrCl₃ and 2 mL mannan
30 were added and the sample was incubated with gentle mixing at room temperature for 5 minutes. The reaction was terminated by adding 7.5mL BBS/gelatin/Mg²⁺/Ca²⁺. The sample was spun down as above, re-suspended in 2 mL BBS/gelatin/Mg²⁺/Ca²⁺ and washed a further two times as above, then stored at 4°C.

Day 2. Hemolysis assay

Materials included BBS/gelatin/Mg²⁺/Ca²⁺ (as above), test sera, 96-well round-bottomed and flat-bottomed plates and a spectrophotometer that reads 96-well plates at 410-414 nm.

- 5 The concentration of the RBC was first determined and the cells were adjusted to 10⁹/mL, and stored at this concentration. Before use, the assay buffer was diluted to 10⁸/mL, and then 100ul per well was used. Hemolysis was measured at 410-414 nm (allowing for greater sensitivity then 541nm). Dilutions of test sera were prepared in ice-cold BBS/gelatin/Mg²⁺/Ca²⁺. 100ul of each serum dilution was pipetted into round-
- 10 bottomed plate (see plate layout). 100ul of appropriately diluted RBC preparation was added (i.e., 10⁸ /mL) (see plate layout), incubated at 37°C for about 1 hour, and observed for lysis. (The plates may be photographed at this point.) The plate was then spun down at maximum speed for 5 minutes. 100ul was aspirated of the fluid-phase, transferred to flat-bottom plates, and the OD was recorded at 410-414 nm. The RBC pellets were
- 15 retained (these can be subsequently lysed with water to obtain an inverse result).

Experiment #1:

- Fresh blood was obtained from CD55/CD59 double-deficient mice and blood of Crry/C3 double-deficient mice and erythrocytes were prepared as described in detail in the above protocol. The cells were split and half of the cells were coated with mannan
- 20 and the other half were left untreated, adjusting the final concentration to 1x 10⁸ per mL, of which 100 µl was used in the hemolysis assay, which was carried out as described above.

Results of Experiment #1: The lectin pathway is involved in erythrocyte lysis in the PNH animal model

- 25 In an initial experiment, it was determined that non-coated WT mouse erythrocytes were not lysed in any mouse serum. It was further determined that mannan-coated Crry^{-/-} mouse erythrocytes were slowly lysed (more than 3 hours at 37 degrees) in WT mouse serum, but they were not lysed in MBL null serum. (Data not shown).

- It was determined that mannan-coated Crry^{-/-} mouse erythrocytes were rapidly
- 30 lysed in human serum but not in heat-inactivated NHS. Importantly, mannan-coated Crry^{-/-} mouse erythrocytes were lysed in NHS diluted down to 1/640 (i.e., 1/40, 1/80, 1/160, 1/320 and 1/640 dilutions all lysed). (Data not shown). In this dilution, the

alternative pathway does not work (AP functional activity is significantly reduced below 8% serum concentration).

Conclusions from Experiment #1

Mannan-coated Crry^{-/-} mouse erythrocytes are very well lysed in highly diluted human serum with MBL but not in that without MBL. The efficient lysis in every serum concentration tested implies that the alternative pathway is not involved or needed for this lysis. The inability of MBL-deficient mouse serum and human serum to lyse the mannan-coated Crry^{-/-} mouse erythrocytes indicates that the classical pathway also has nothing to do with the lysis observed. As lectin pathway recognition molecules are required (i.e., MBL), this lysis is mediated by the lectin pathway.

Experiment #2:

Fresh blood was obtained from the Crry/C3 and CD55/CD59 double-deficient mice and mannan-coated Crry^{-/-} mouse erythrocytes were analyzed in the haemolysis assay as described above in the presence of the following human serum: MBL null; WT; NHS pretreated with human anti-MASP-2 antibody; and heat-inactivated NHS as a control.

Results of Experiment #2: MASP-2 inhibitors prevent erythrocyte lysis in the PNH animal model

With the Mannan-coated Crry^{-/-} mouse erythrocytes, NHS was incubated in the dilutions diluted down to 1/640 (i.e., 1/40, 1/80, 1/160, 1/320 and 1/640), human MBL^{-/-} serum, NHS pretreated with anti-MASP-2 mAb, and heat-inactivated NHS as a control.

The ELISA microtiter plate was spun down and the non-lysed erythrocytes were collected on the bottom of the round-well plate. The supernatant of each well was collected and the amount of hemoglobin released from the lysed erythrocytes was measured by reading the OD415 nm in an ELISA reader.

In the control heat-inactivated NHS (negative control), as expected, no lysis was observed. MBL^{-/-} human serum lysed mannan-coated mouse erythrocytes at 1/8 and 1/16 dilutions. Anti-MASP-2-antibody-pretreated NHS lysed mannan-coated mouse erythrocytes at 1/8 and 1/16 dilutions while WT human serum lysed mannan-coated mouse erythrocytes down to dilutions of 1/32.

FIGURE 40 graphically illustrates hemolysis (as measured by hemoglobin release of lysed mouse erythrocytes (Crry/C3^{-/-}) into the supernatant measured by photometry) of mannan-coated murine erythrocytes by human serum over a range of serum

concentrations in serum from heat-inactivated (HI) NHS, MBL-/-, NHS pretreated with anti-MASP-2 antibody, and NHS control.

From the results shown in FIGURE 40, it is demonstrated that MASP-2 inhibition with anti-MASP-2 antibody significantly shifted the CH₅₀ and inhibited complement-mediated lysis of sensitized erythrocytes with deficient protection from autologous complement activation.

Experiment #3

Fresh blood obtained from the Crry/C3 and CD55/CD59 double-deficient mice in non-coated Crry-/- mouse erythrocytes was analyzed in the hemolysis assay as described above in the presence of the following serum: MBL -/-; WT sera; NHS pretreated with human anti-MASP-2 antibody and heat-inactivated NHS as a control.

Results:

FIGURE 41 graphically illustrates hemolysis (as measured by hemoglobin release of lysed WT mouse erythrocytes into the supernatant measured by photometry) of non-coated murine erythrocytes by human serum over a range of serum concentrations in serum from heat inactivated (HI) NHS, MBL-/-, NHS pretreated with anti-MASP-2 antibody, and NHS control. As shown in FIGURE 41, it is demonstrated that inhibiting MASP-2 inhibits complement-mediated lysis of non-sensitized WT mouse erythrocytes.

FIGURE 42 graphically illustrates hemolysis (as measured by hemoglobin release of lysed mouse erythrocytes (CD55/59 -/-) into the supernatant measured by photometry) of non-coated murine erythrocytes by human serum over a range of serum concentration in serum from heat-inactivated (HI) NHS, MBL-/-, NHS pretreated with anti-MASP-2 antibody, and NHS control.

TABLE 12: CH₅₀ values expressed as serum concentrations

Serum	WT	CD55/59 -/-
Heat-inactivated NHS	No lysis	No lysis
MBL AO/XX donor (MBL deficient)	7.2%	2.1%
NHS + anti-MASP-2 antibody	5.4%	1.5%
NHS	3.1%	0.73%

Note: "CH₅₀" is the point at which complement mediated hemolysis reaches 50%.

In summary, the results in this Example demonstrate that inhibiting MASP-2 inhibits complement-mediated lysis of sensitized and non-sensitized erythrocytes with deficient protection from autologous complement activation. Therefore, MASP-2 inhibitors may be used to treat subjects suffering from PNH, and may also be used to ameliorate (*i.e.*, inhibit, prevent or reduce the severity of) extravascular hemolysis in PNH patients undergoing treatment with a C5 inhibitor such as eculizumab (Soliris®).

10 **EXAMPLE 34**

This Example describes a follow on study to the study described above in Example 29, providing further evidence confirming that a MASP-2 inhibitor, such as a MASP-2 antibody, is effective for the treatment of radiation exposure and/or for the treatment, amelioration or prevention of acute radiation syndrome.

15 **Rationale:** In the initial study described in Example 29, it was demonstrated that pre-irradiation treatment with an anti-MASP-2 antibody in mice increased the survival of irradiated mice as compared to vehicle treated irradiated control animals at both 6.5 Gy and 7.0 Gy exposure levels. It was further demonstrated in Example 29 that at the 6.5 Gy exposure level, post-irradiation treatment with anti-MASP-2 antibody resulted in a modest increase in survival as compared to vehicle control irradiated animals. This Example describes a second radiation study that was carried out to confirm the results of the first study.

Methods:

25 **Design of Study A:**

Swiss Webster mice (n=50) were exposed to ionizing radiation (8.0 Gy). The effect of anti-MASP-2 antibody therapy (mAbH6 5mg/kg), administered 18 hours before and 2 hours after radiation exposure, and weekly thereafter, on mortality was assessed.

Results of Study A:

30 As shown in FIGURE 43, it was determined that administration of the anti-MASP-2 antibody mAbH6 increased survival in mice exposed to 8.0 Gy, with an adjusted median survival rate increased from 4 to 6 days as compared to mice that received vehicle

control, and a mortality reduced by 12% when compared to mice that received vehicle control (log-rank test, $p=0.040$).

Design of Study B:

- 5 Swiss Webster mice ($n=50$) were exposed to ionizing radiation (8.0 Gy) in the following groups (I: vehicle) saline control; (II: low) anti-MASP-2 antibody mAbH6 (5 mg/kg) administered 18 hours before irradiation and 2 hours after irradiation; (III: high) mAbH6 (10 mg/kg) administered 18 hours before irradiation and 2 hours post irradiation; and (IV:high post) mAbH6 (10mg/kg) administered 2 hours post irradiation only.

10 Results of Study B:

Administration of anti-MASP-2 antibody pre- and post-irradiation adjusted the mean survival from 4 to 5 days in comparison to animals that received vehicle control.

Mortality in the anti-MASP-2 antibody-treated mice was reduced by 6-12% in comparison to vehicle control mice. It is further noted that no significant detrimental

- 15 treatment effects were observed (data not shown).

In summary, the results shown in this Example are consistent with the results shown in Example 29 and further demonstrate that anti-MASP-2 antibodies are effective in treating a mammalian subject at risk for, or suffering from the detrimental effects of acute radiation syndrome.

20

EXAMPLE 35

This study investigates the effect of MASP-2-deficiency in a mouse model of LPS (lipopolysaccharide)-induced thrombosis.

Rationale:

- 25 Hemolytic uremic syndrome (HUS), which is caused by Shiga toxin-producing *E. coli* infection, is the leading cause of acute renal failure in children. In this Example, a Schwartzman model of LPS-induced thrombosis (microvascular coagulation) was carried out in MASP-2-/- (KO) mice to determine whether MASP-2 inhibition is effective to inhibit or prevent the formation of intravascular thrombi.

30 Methods:

MASP-2-/- ($n=9$) and WT ($n=10$) mice were analyzed in a Schwartzman model of LPS-induced thrombosis (microvascular coagulation). Mice were administered Serratia

LPS and thrombus formation was monitored over time. A comparison of the incidence of microthrombi and LPS-induced microvascular coagulation was carried out.

Results:

Notably, all MASP-2 $-/-$ mice tested (9/9) did not form intravascular thrombi after Serratia LPS administration. In contrast, microthrombi were detected in 7 of 10 of the WT mice tested in parallel ($p=0.0031$, Fischer's exact). As shown in FIGURE 44, the time to onset of microvascular occlusion following LPS infection was measured in MASP-2 $-/-$ and WT mice, showing the percentage of WT mice with thrombus formation measured over 60 minutes, with thrombus formation detected as early as about 15 minutes. Up to 80% of the WT mice demonstrated thrombus formation at 60 minutes. In contrast, as shown in FIGURE 44, none of the MASP-2 $-/-$ had thrombus formation at 60 minutes (log rank: $p=0.0005$).

These results demonstrate that MASP-2 inhibition is protective against the development of intravascular thrombi in an HUS model.

EXAMPLE 36

This Example describes the effect of anti-MASP-2 antibodies in a mouse model of HUS using intraperitoneal co-injection of purified *Shiga toxin 2* (STX2) plus LPS.

Background:

A mouse model of HUS was developed using intraperitoneal co-injection of purified Shiga toxin 2 (STX2) plus LPS. Biochemical and microarray analysis of mouse kidneys revealed the STX2 plus LPS challenge to be distinct from the effects of either agent alone. Blood and serum analysis of these mice showed neutrophilia, thrombocytopenia, red cell hemolysis, and increased serum creatinine and blood urea nitrogen. In addition, histologic analysis and electron microscopy of mouse kidneys demonstrated glomerular fibrin deposition, red cell congestion, microthrombi formation, and glomerular ultrastructural changes. It was established that this model of HUS induces all clinical symptoms of human HUS pathology in C57BL/6 mice including thrombocytopenia, hemolytic anemia, and renal failure that define the human disease. (J. Immunol 187(1):172-80 (2011))

Methods:

C57BL/6 female mice that weighed between 18 to 20 g were purchased from Charles River Laboratories and divided in to 2 groups (5 mice in each group). One group

of mice was pretreated by intraperitoneal (i.p.) injection with the recombinant anti-MASP-2 antibody mAbM11 (100 µg per mouse; corresponding to a final concentration of 5 mg/kg body weight) diluted in a total volume of 150 µl saline. The control group received saline without any antibody. Six hours after i.p. injection of anti-MASP-2 antibody mAbM11, all mice received a combined i.p. injection of a sublethal dose (3 µg/animal; corresponding to 150 µg/kg body weight) of LPS of *Serratia marcescens* (L6136; Sigma-Aldrich, St. Louis, MO) and a dose of 4.5 ng/animal (corresponding to 225 ng/kg) of STX2 (two times the LD50 dose) in a total volume of 150 µl. Saline injection was used for control.

Survival of the mice was monitored every 6 hours after dosing. Mice were culled as soon as they reached the lethargic stage of HUS pathology. After 36 hours, all mice were culled and both kidneys were removed for immunohistochemistry and scanning electron microscopy. Blood samples were taken at the end of the experiment by cardiac puncture. Serum was separated and kept frozen at -80°C for measuring BUN and serum Creatinine levels in both treated and control groups.

Immunohistochemistry

One-third of each mouse kidney was fixed in 4% paraformaldehyde for 24 h, processed, and embedded in paraffin. Three-micron-thick sections were cut and placed onto charged slides for subsequent staining with H & E stain.

Electron Microscopy

The middle section of the kidneys was cut into blocks of approximately 1 to 2 mm³, and fixed overnight at 4°C in 2.5% glutaraldehyde in 1x PBS. The fixed tissue subsequently was processed by the University of Leicester Electron Microscopy Facility

Cryostat sections

The other third of the kidneys was, cut into blocks approximately 1 to 2 mm³ and snap frozen in liquid nitrogen and kept at -80°C for cryostat sections and mRNA analysis.

Results:

FIGURE 45 graphically illustrates the percent survival of saline-treated control mice (n=5) and anti-MASP-2 antibody-treated mice (n=5) in the STX/LPS-induced model over time (hours). Notably, as shown in FIGURE 45, all of the control mice died by 42 hours. In sharp contrast, 100 % of the anti-MASP-2 antibody-treated mice survived throughout the time course of the experiment. Consistent with the results shown

in FIGURE 45, it was observed that all the untreated mice that either died or had to be culled with signs of severe disease had significant glomerular injuries, while the glomeruli of all anti-MASP-2-treated mice looked normal (data not shown). These results demonstrate that MASP-2 inhibitors, such as anti-MASP-2 antibodies, may be used to treat subjects suffering from, or at risk for developing a thrombotic microangiopathy (TMA), such as hemolytic uremic syndrome (HUS), atypical HUS (aHUS), or thrombotic thrombocytopenic purpura (TTP).

EXAMPLE 37

This Example describes the effect of MASP-2 deficiency and MASP-2 inhibition in a murine FITC-dextran/light induced endothelial cell injury model of thrombosis.

Background/Rationale: As demonstrated in Examples 35 and 36, MASP-2 deficiency (MASP-2 KO) and MASP-2 inhibition (via administration of an inhibitory MASP-2 antibody) protects mice in a model of typical HUS, whereas all control mice exposed to STX and LPS developed severe HUS and became moribund or died within 48 hours. For example, as shown in FIGURE 54, all mice treated with a MASP-2 inhibitory antibody and then exposed to STX and LPS survived (Fisher's exact $p < 0.01$; $N = 5$). Thus, anti-MASP-2 therapy protects mice in this model of HUS.

The following experiments were carried out to analyze the effect of MASP-2 deficiency and MASP-2 inhibition in a fluorescein isothiocyanate (FITC)-dextran-induced endothelial cell injury model of thrombotic microangiopathy (TMA) in order to demonstrate further the benefit of MASP-2 inhibitors for the treatment of HUS, aHUS, TTP, and TMA's with other etiologies.

Methods:

Intravital microscopy

Mice were prepared for intravital microscopy as described by Frommhold et al., *BMC Immunology* 12:56-68, 2011. Briefly, mice were anesthetized with intraperitoneal (i.p.) injection of ketamine (125 mg/kg bodyweight, Ketanest, Pfitzer GmbH, Karlsruhe, Germany) and xylazine (12.5 mg/kg body weight; Rompun, Bayer, Leverkusen, Germany) and placed on a heating pad to maintain body temperature at 37°C. Intravital microscopy was conducted on an upright microscope (Leica, Wetzlar, Germany) with a saline immersion objective (SW 40/0.75 numerical aperture, Zeiss, Jena, Germany). To ease breathing, mice were intubated using PE 90 tubing (Becton Dickson and Company, Sparks, MD, USA). The left carotid artery was cannulated with PE10 tubing (Becton

Dickson and Company, Sparks, MD, USA) for blood sampling and systemic monoclonal antibody (mAb) administration.

Cremaster muscle preparation

The surgical preparation of the cremaster muscle for intravital microscopy was performed as described by Sperandio et al., *Blood*, 97:3812-3819, 2001. Briefly, the scrotum was opened and the cremaster muscle mobilized. After longitudinal incision and spreading of the muscle over a cover glass, the epididymis and testis were moved and pinned to the side, giving full microscopic access to the cremaster muscle microcirculation. Cremaster muscle venules were recorded via a CCD camera (CF8/1; Kappa, Gleichen, Germany) on a Panasonic S-VHS recorder. The cremaster muscle was superfused with thermo-controlled (35°C bicarbonate-buffered saline) as previously described by Frommhold et al., *BMC Immunology* 12:56-68, 2011.

Light excitation FITC dextran injury model

A controlled, light-dose-dependent vascular injury of the endothelium of cremaster muscle venules and arterioles was induced by light excitation of phototoxic (FITC)-dextran (Cat. No. FD150S, Sigma Aldrich, Poole, U.K.). This procedure initiates localized thrombosis. As a phototoxic reagent, 60 µL of a 10% w/v solution of FITC-dextran was injected through the left carotid artery access and allowed to spread homogeneously throughout the circulating blood for 10 minutes. After selecting a well-perfused venule, halogen light of low to midrange intensity (800-1500) was focused on the vessel of interest to induce FITC-dextran fluorescence and mild to moderate phototoxicity to the endothelial surface in order to stimulate thrombosis in a reproducible, controlled manner. The necessary phototoxic light intensity for the excitation of FITC-dextran was generated using a halogen lamp (12V, 100W, Zeiss, Oberkochen, Germany). The phototoxicity resulting from light-induced excitation of the fluorochrome requires a threshold of light intensity and/or duration of illumination and is caused by either direct heating of the endothelial surface or by generation of reactive oxygen radicals as described by Steinbauer et al., *Langenbecks Arch Surg* 385:290-298, 2000.

The intensity of the light applied to each vessel was measured for adjustment by a wavelength-correcting diode detector for low power measurements (Labmaster LM-2, Coherent, Auburn, USA). Off-line analysis of video scans was performed by means of a computer assisted microcirculation analyzing system (CAMAS, Dr. Zeintl, Heidelberg)

and red blood cell velocity was measured as described by Zeintl et al., *Int J Microcirc Clin Exp*, 8(3):293-302, 2000.

Application of monoclonal anti-human MASP-2 inhibitory antibody (mAbH6) and vehicle control prior to induction of thrombosis

5 Using a blinded study design, 9-week-old male C57BL/6 WT littermate mice were given i.p. injections of either the recombinant monoclonal human MASP-2 antibody (mAbH6), an inhibitor of MASP-2 functional activity (given at a final concentration of 10mg/kg body weight), or the same quantity of an isotype control antibody (without MASP-2 inhibitory activity) 16 hours before the phototoxic induction of thrombosis in
10 the cremaster model of intravital microscopy. One hour prior to thrombosis induction, a second dose of either mAbH6 or the control antibody was given. MASP-2 knockout (KO) mice were also evaluated in this model.

 mAbH6 (established against recombinant human MASP-2) is a potent inhibitor of human MASP-2 functional activity, which cross-reacts with, binds to and inhibits mouse
15 MASP-2 but with lower affinity due to its species specificity (data not shown). In order to compensate for the lower affinity of mAbH6 to mouse MASP-2, mAbH6 was given at a high concentration (10mg/kg body weight) to overcome the variation in species specificity, and the lesser affinity for mouse MASP-2, to provide effective blockade of murine MASP-2 functional activity under *in vivo* conditions.

20 In this blinded study, the time required for each individual venuole tested (selection criteria were by comparable diameters and blood flow velocity) to fully occlude was recorded.

 The percentage of mice with microvascular occlusion, the time of onset, and the time to occlusion were evaluated over a 60-minute observation period using intravital
25 microscopy video recordings.

Results:

 FIGURE 46 graphically illustrates, as a function of time after injury induction, the percentage of mice with microvascular occlusion in the FITC/Dextran UV model after treatment with isotype control or human MASP-2 antibody mAbH6 (10mg/kg) dosed at
30 16 hours and 1 hour prior to injection of FITC/Dextran. As shown in FIGURE 46, 85% of the wild-type mice receiving the isotype control antibody occluded within 30 minutes or less, whereas only 19% of the wild-type mice pre-treated with the human MASP-2 antibody (mAbH6) occluded within the same time period, and the time to occlusion was

delayed in the mice that did eventually occlude in the human MASP-2 antibody-treated group. It is further noted that three of the MASP-2 mAbH6 treated mice did not occlude at all within the 60-minute observation period (i.e., were protected from thrombotic occlusion).

5 FIGURE 47 graphically illustrates the occlusion time in minutes for mice treated with the human MASP-2 antibody (mAbH6) and the isotype control antibody. The data are reported as scatter-dots with mean values (horizontal bars) and standard error bars (vertical bars). This figure shows the occlusion time in the mice where occlusion was observable. Thus, the three MASP-2 antibody-treated mice that did not occlude during
10 the 60 minute observation period were not included in this analysis (there was no control treated mouse that did not occlude). The statistical test used for analysis was the unpaired t test; wherein the symbol “*” indicates $p=0.0129$. As shown in FIGURE 47, in the four MASP-2 antibody (mAbH6)-treated mice that occluded, treatment with MASP-2 antibody significantly increased the venous occlusion time in the FITC-dextran/light-
15 induced endothelial cell injury model of thrombosis with low light intensity (800-1500) as compared to the mice treated with the isotype control antibody. The average of the full occlusion time of the isotype control was 19.75 minutes, while the average of the full occlusion time for the MASP-2 antibody treated group was 32.5 minutes.

 FIGURE 48 graphically illustrates the time until occlusion in minutes for wild-
20 type mice, MASP-2 KO mice, and wild-type mice pre-treated with human MASP-2 antibody (mAbH6) administered i.p. at 10mg/kg 16 hours before, and then administered again i.v. 1 hour prior to the induction of thrombosis in the FITC-dextran/light-induced endothelial cell injury model of thrombosis with low light intensity (800-1500). Only the animals that occluded were included in FIGURE 48; $n=2$ for wild-type mice receiving
25 isotype control antibody; $n=2$ for MASP-2 KO; and $n=4$ for wild-type mice receiving human MASP-2 antibody (mAbH6). The symbol “*” indicates $p<0.01$. As shown in FIGURE 48, MASP-2 deficiency and MASP-2 inhibition (mAbH6 at 10mg/kg) increased the venous occlusion time in the FITC-dextran/light-induced endothelial cell injury model of thrombosis with low light intensity (800-1500).

30 Conclusions:

 The results in this Example further demonstrate that a MASP-2 inhibitory agent that blocks the lectin pathway (e.g., antibodies that block MASP-2 function), inhibits microvascular coagulation and thrombosis, the hallmarks of multiple microangiopathic

disorders, in a mouse model of TMA. Therefore, it is expected that administration of a MASP-2 inhibitory agent, such as a MASP-2 inhibitory antibody, will be an effective therapy in patients suffering from HUS, aHUS, TTP, or other microangiopathic disorders and provide protection from microvascular coagulation and thrombosis.

5

EXAMPLE 38

This Example describes a study demonstrating that human MASP-2 inhibitory antibody (mAbH6) has no effect on platelet function in platelet-rich human plasma.

10 Background/Rationale: As described in Example 37, it was demonstrated that MASP-2 inhibition with human MASP-2 inhibitory antibody (mAbH6) increased the venous occlusion time in the FITC-dextran/light-induced endothelial cell injury model of thrombosis. The following experiment was carried out to determine whether the MASP-2 inhibitory antibody (mAbH6) has an effect on platelet function.

15 Methods: The effect of human mAbH6 MASP-2 antibody was tested on ADP-induced aggregation of platelets as follows. Human MASP-2 mAbH6 at a concentration of either 1 µg/ml or 0.1 µg/ml was added in a 40 µL solution to 360 µL of freshly prepared platelet-rich human plasma. An isotype control antibody was used as the negative control. After adding the antibodies to the plasma, platelet activation was induced by
20 adding ADP at a final concentration of 2 µM. The assay was started by stirring the solutions with a small magnet in the 1 mL cuvette. Platelet aggregation was measured in a two-channel Chrono-log Platelet Aggregometer Model 700 Whole Blood/Optical Lumi-Aggregometer.

Results:

25 The percent aggregation in the solutions was measured over a time period of five minutes. The results are shown below in TABLE 13.

TABLE 13: Platelet Aggregation over a time period of five minutes.

Antibody	Amplitude (percent aggregation)	Slope (percent aggregation over time)
MASP-2 antibody (mAbH6)	46%	59

(1 µg/ml)		
Isotype control antibody (1 µg/ml)	49%	64
MASP-2 antibody (mAbH6) (0.1 µg/ml)	52%	63
Isotype control antibody (0.1 µg/ml)	46%	59

As shown above in TABLE 13, no significant difference was observed between the aggregation of the ADP-induced platelets treated with the control antibody or the MASP-2 mAbH6 antibody. These results demonstrate that the human MASP-2 antibody (mAbH6) has no effect on platelet function. Therefore, the results described in Example 37 demonstrating that MASP-2 inhibition with human MASP-2 inhibitory antibody (mAbH6) increased the venous occlusion time in the FITC-dextran/light-induced endothelial cell injury model of thrombosis, were not due to an effect of mAbH6 on platelet function. Thus, MASP-2 inhibition prevents thrombosis without directly impacting platelet function, revealing a therapeutic mechanism that is distinct from existing anti-thrombotic agents.

EXAMPLE 39

This Example describes the effect of MASP-2 inhibition on thrombus formation and vessel occlusion in a murine model of TMA.

Background/Rationale: The lectin pathway plays a dominant role in activating the complement system in settings of endothelial cell stress or injury. This activation is amplified rapidly by the alternative pathway, which is dysregulated in many patients presenting with aHUS. Preventing the activation of MASP-2 and the lectin pathway is thus expected to halt the sequence of enzymatic reactions that lead to the formation of the membrane attack complex, platelet activation, and leukocyte recruitment. This effect limits tissue damage.

In addition, MASP-2 has Factor Xa-like activity and cleaves prothrombin to form

thrombin. This MASP-2-driven activation of the coagulation system may imbalance hemostasis and result in the pathology of TMA. Thus, inhibition of MASP-2 using a MASP-2 inhibitor, such as a MASP-2 inhibitory antibody that blocks activation of the complement and coagulation systems is expected to improve outcomes in aHUS and other TMA-related conditions.

As described in Example 37, it was demonstrated that MASP-2 inhibition with human MASP-2 inhibitory antibody (mAbH6) increased the venous occlusion time in the FITC-dextran/light-induced endothelial cell injury model of thrombosis. In this model of TMA, mice were sensitized by IV injection of FITC- dextran, followed by localized photo-activation of the FITC- dextran in the microvasculature of the mouse cremaster muscle (Thorlacius H et al., *Eur J Clin. Invest* 30(9):804-10, 2000; Agero et al., *Toxicon* 50(5):698-706, 2007).

The following experiment was carried out to determine whether the MASP-2 inhibitory antibody (mAbH6) has a dose-response effect on thrombus formation and vessel occlusion in a murine model of TMA.

Methods: Localized thrombosis was induced by photo-activation of fluorescein isothiocyanate-labeled dextran (FITC-dextran) in the microvasculature of the cremaster muscle of C57 Bl/6 mice and intravital microscopy was used to measure onset of thrombus formation and vessel occlusion using methods described in Example 37, with the following modifications. Groups of mice were dosed with mAbH6 (2mg/kg, 10 mg/kg or 20 mg/kg) or isotype control antibody (20 mg/kg) were administered by intravenous (iv) injection one hour before TMA induction. The time to onset of thrombus formation and time to complete vessel occlusion were recorded. Video playback analysis of intravital microscopy images recorded over 30 to 60 minutes was used to evaluate vessel size, blood flow velocity, light intensity, rate of onset of thrombus formation as equivalent of platelet adhesion, time to onset of thrombus formation, rate of total vessel occlusion and time until total vessel occlusion. Statistical analysis was conducted using SigmaPlot v12.0.

Results:

Initiation of Thrombus Formation

FIGURE 49 is a Kaplan-Meier plot showing the percentage of mice with thrombi as a function of time in FITC-Dextran induced thrombotic microangiopathy in mice treated with increasing doses of human MASP-2 inhibitory antibody (mAbH6 at 2 mg/kg,

10mg/kg or 20 mg/kg) or an isotype control antibody. As shown in FIGURE 49, initiation of thrombus formation was delayed in the mAbH6-treated mice in a dose-dependent manner relative to the control-treated mice.

- FIGURE 50 graphically illustrates the median time to onset (minutes) of thrombus formation as a function of mAbH6 dose (* $p < 0.01$ compared to control). As shown in FIGURE 50, the median time to onset of thrombus formation increased with increasing doses of mAbH6 from 6.8 minutes in the control group to 17.7 minutes in the 20 mg/kg mAbH6 treated group ($p < 0.01$). The underlying experimental data and statistical analysis are provided in TABLES 14 and 15.
- 10 The time to onset of thrombus formation in individual mice recorded based on evaluation of the videographic recording is detailed below in TABLE 14.

TABLE 14: Time to Onset of Thrombus Formation After Light Dye-induced Injury

Control Treatment		mAbH6 Treatment		
Time to Onset (minutes)	Control	2 mg/kg	10 mg/kg	20 mg/kg
	6.07	5.93	12.75	10.00
	1.07	6.95	2.53	10.33
	8.00	8.92	14.00	21.00
	2.40	11.92	3.05	11.50
	8.48	12.75	8.00	19.00
	4.00	12.53	8.17	10.37
	4.00		15.83	22.65
	7.83		11.70	16.37
	6.83		50.67	21.75*
	15.00			32.25*
	15.67			

* vessels did not show onset during the indicated observation period

The statistical analysis comparing time to onset of occlusion between control and mAbH6
5 treated animals is shown below in TABLE 15.

TABLE 15: Time to Onset: data from FITC Dex dose response study

Statistic	Control	mAbH6 (2 mg/kg)	mAbH6 (10 mg/kg)	mAbH6 (20 mg/kg)
Number of events/number of animals (%)	11/11 (100%)	6/6 (100%)	9/9 (100%)	8/10 (80.0%)
Median time (minutes) (95% CI)	6.8 (2.4, 8.5)	10.4 (5.9, 12.8)	11.7 (2.5, 15.8)	17.7 (10.0, 22.7)
Wilcoxon p-value*		0.2364	0.1963	0.0016
Event=Time to onset observed Median (minutes) and its 95% CI were based on Kaplan-Meier estimate NE=not estimable *p-values were adjusted by Dunnett-Hsu multiple comparison				

Microvascular Occlusion

- FIGURE 51 is a Kaplan-Meier plot showing the percentage of mice with microvascular occlusion as a function of time in FITC-Dextran induced thrombotic microangiopathy in mice treated with increasing doses of human MASP-2 inhibitory antibody (mAbH6 at 2 mg/kg, 10mg/kg or 20mg/kg) or an isotype control antibody. As shown in FIGURE 51, complete microvascular occlusion was delayed in the mAbH6 treated groups as compared to the control mice.
- FIGURE 52 graphically illustrates the median time to microvascular occlusion as a function of mAbH6 dose (*p<0.05 compared to control). As shown in FIGURE 52, the median time to complete microvascular occlusion increased from 23.3 minutes in the control group to 38.6 minutes in the 2mg/kg mAbH6 treated group (p<0.05). Doses of 10 mg/kg or 20 mg/kg of mAbH6 performed similarly (median time for complete microvascular occlusion was 40.3 and 38 minutes, respectively) to the 2 mg/kg mAbH6 treated group. The underlying experimental data and statistical analysis are provided in

TABLES 16 and 17.

The time to complete vessel occlusion in individual mice recorded based on primary evaluation of the videographic recording is detailed below in TABLE 16.

5 **TABLE 16: Time to Complete Occlusion After Light Dye-Induced Injury**

Control Treatment		mAbH6 Treatment		
Time to Occlusion (minutes)	Control	2 mg/kg	10 mg/kg	20 mg/kg
	37.50	42.3	30.92	38.00
	29.07	21.91	17.53	28.00
	27.12	24.4	51.38	40.58
	19.38	31.38	36.88	33.00
	19.55	61.17*	26.83	39.10
	18.00	61.55*	40.28	32.03
	16.50		55.83	38.53
	23.33		71.93*	21.75*
	14.83		98.22*	32.25*
	30*			33.17*
	61.8*			

*vessels did not completely occlude during the indicated observation period.

The statistical analysis comparing time to complete occlusion between control and mAbH6 treated animals is shown below in TABLE 17.

TABLE 17: Time to Complete Microvascular Occlusion: data from FITC Dex dose response study

Statistic	Control	mAbH6 (2 mg/kg)	mAbH6 (10 mg/kg)	mAbH6 (20 mg/kg)
Number of events/number of animals (%)	9/11 (81.8%)	4/6 (66.7%)	7/9 (77.8%)	7/10 (70.0%)
Median time (minutes) (95% CI)	23.3 (16.5, 37.5)	36.8 (21.9, NE)	40.3 (17.5, NE)	38.0 (28.0, 40.6)
Wilcoxon p-value*		0.0456	0.0285	0.0260
Event=Time to occlusion observed Median (minutes) and its 95% CI were based on Kaplan-Meier estimate NE=not estimable *p-values were adjusted by Dunnett-Hsu multiple comparison				

Summary

- 5 As summarized in TABLE 18, the initiation of thrombus formation was delayed in the mAbH6 treated mice in a dose-dependent manner relative to the control-treated mice (median time to onset 10.4 to 17.7 minutes vs 6.8 minutes). The median time to complete occlusion was significantly delayed in all mAbH6-treated groups relative to the control-treated groups (Table 18).

TABLE 18: Median Time to Onset of Thrombus Formation and Complete Occlusion

	Control	mAbH6 (2 mg/kg)	mAbH6 (10 mg/kg)	mAbH6 (20 mg/kg)
Median# time to onset of thrombus formation (minutes)	6.8	10.4	11.7	17.7*
Median# time to complete microvascular occlusion (minutes)	23.3	36.8*	40.3*	38.0*

#Median values are based on Kaplan-Meier estimate

*p<0.05 compared to control (Wilcoxon adjusted by Dunnett-Hsu for multiple comparisons)

These results demonstrate that mAbH6, a human monoclonal antibody that binds to MASP-2 and blocks the lectin pathway of the complement system, reduced microvascular thrombosis in a dose-dependent manner in an experimental mouse model of TMA. Therefore, it is expected that administration of a MASP-2 inhibitory agent, such as a MASP-2 inhibitory antibody, will be an effective therapy in patients suffering from HUS, aHUS, TTP, or other microangiopathic disorders such as other TMAs including catastrophic antiphospholipid syndrome (CAPS), systemic Degos disease, and TMAs secondary to cancer, cancer chemotherapy and transplantation and provide protection from microvascular coagulation and thrombosis.

EXAMPLE 40

This Example describes the identification, using phage display, of fully human scFv antibodies that bind to MASP-2 and inhibit lectin-mediated complement activation

while leaving the classical (C1q-dependent) pathway and the alternative pathway components of the immune system intact.

Overview:

Fully human, high-affinity MASP-2 antibodies were identified by screening a phage display library. The variable light and heavy chain fragments of the antibodies were isolated in both a scFv format and in a full-length IgG format. The human MASP-2 antibodies are useful for inhibiting cellular injury associated with lectin pathway-mediated alternative complement pathway activation while leaving the classical (C1q-dependent) pathway component of the immune system intact. In some embodiments, the subject MASP-2 inhibitory antibodies have the following characteristics: (a) high affinity for human MASP-2 (e.g., a K_D of 10 nM or less), and (b) inhibit MASP-2-dependent complement activity in 90% human serum with an IC_{50} of 30 nM or less.

Methods:

Expression of full-length catalytically inactive MASP-2:

The full-length cDNA sequence of human MASP-2 (SEQ ID NO: 4), encoding the human MASP-2 polypeptide with leader sequence (SEQ ID NO:5) was subcloned into the mammalian expression vector pCI-Neo (Promega), which drives eukaryotic expression under the control of the CMV enhancer/promoter region (described in Kaufman R.J. et al., *Nucleic Acids Research* 19:4485-90, 1991; Kaufman, *Methods in Enzymology*, 185:537-66 (1991)).

In order to generate catalytically inactive human MASP-2A protein, site-directed mutagenesis was carried out as described in US2007/0172483, hereby incorporated herein by reference. The PCR products were purified after agarose gel electrophoresis and band preparation and single adenosine overlaps were generated using a standard tailing procedure. The adenosine-tailed MASP-2A was then cloned into the pGEM-T easy vector and transformed into *E. coli*. The human MASP-2A was further subcloned into either of the mammalian expression vectors pED or pCI-Neo.

The MASP-2A expression construct described above was transfected into DXB1 cells using the standard calcium phosphate transfection procedure (Maniatis et al., 1989). MASP-2A was produced in serum-free medium to ensure that preparations were not contaminated with other serum proteins. Media was harvested from confluent cells every second day (four times in total). The level of recombinant MASP-2A averaged

approximately 1.5 mg/liter of culture medium. The MASP-2A (Ser-Ala mutant described above) was purified by affinity chromatography on MBP-A-agarose columns

MASP-2A ELISA on ScFv Candidate Clones identified by panning/scFv conversion and filter screening

5 A phage display library of human immunoglobulin light- and heavy-chain variable region sequences was subjected to antigen panning followed by automated antibody screening and selection to identify high-affinity scFv antibodies to human MASP-2 protein. Three rounds of panning the scFv phage library against HIS-tagged or biotin-tagged MASP-2A were carried out. The third round of panning was eluted first
10 with MBL and then with TEA (alkaline). To monitor the specific enrichment of phages displaying scFv fragments against the target MASP-2A, a polyclonal phage ELISA against immobilized MASP-2A was carried out. The scFv genes from panning round 3 were cloned into a pHOG expression vector and run in a small-scale filter screening to look for specific clones against MASP-2A.

15 Bacterial colonies containing plasmids encoding scFv fragments from the third round of panning were picked, gridded onto nitrocellulose membranes and grown overnight on non-inducing medium to produce master plates. A total of 18,000 colonies were picked and analyzed from the third panning round, half from the competitive elution and half from the subsequent TEA elution. Panning of the scFv phagemid library against
20 MASP-2A followed by scFv conversion and a filter screen yielded 137 positive clones. 108/137 clones were positive in an ELISA assay for MASP-2 binding (data not shown), of which 45 clones were further analyzed for the ability to block MASP-2 activity in normal human serum.

Assay to Measure Inhibition of Formation of Lectin Pathway C3 Convertase

25 A functional assay that measures inhibition of lectin pathway C3 convertase formation was used to evaluate the "blocking activity" of the MASP-2 scFv candidate clones. MASP-2 serine protease activity is required in order to generate the two protein components (C4b, C2a) that comprise the lectin pathway C3 convertase. Therefore, a MASP-2 scFv that inhibits MASP-2 functional activity (i.e., a blocking MASP-2 scFv),
30 will inhibit *de novo* formation of lectin pathway C3 convertase. C3 contains an unusual and highly reactive thioester group as part of its structure. Upon cleavage of C3 by C3 convertase in this assay, the thioester group on C3b can form a covalent bond with

hydroxyl or amino groups on macromolecules immobilized on the bottom of the plastic wells via ester or amide linkages, thus facilitating detection of C3b in the ELISA assay.

Yeast mannan is a known activator of the lectin pathway. In the following method to measure formation of C3 convertase, plastic wells coated with mannan were incubated with diluted human serum to activate the lectin pathway. The wells were then washed and assayed for C3b immobilized onto the wells using standard ELISA methods. The amount of C3b generated in this assay is a direct reflection of the *de novo* formation of lectin pathway C3 convertase. MASP-2 scFv clones at selected concentrations were tested in this assay for their ability to inhibit C3 convertase formation and consequent C3b generation.

Methods:

The 45 candidate clones identified as described above were expressed, purified and diluted to the same stock concentration, which was again diluted in Ca^{++} and Mg^{++} containing GVB buffer (4.0 mM barbital, 141 mM NaCl, 1.0 mM MgCl_2 , 2.0 mM CaCl_2 , 0.1% gelatin, pH 7.4) to assure that all clones had the same amount of buffer. The scFv clones were each tested in triplicate at the concentration of 2 $\mu\text{g/mL}$. The positive control was OMS100 Fab2 and was tested at 0.4 $\mu\text{g/mL}$. C3c formation was monitored in the presence and absence of the scFv/IgG clones.

Mannan was diluted to a concentration of 20 $\mu\text{g/mL}$ (1 $\mu\text{g/well}$) in 50mM carbonate buffer (15mM Na_2CO_3 + 35mM NaHCO_3 + 1.5 mM NaN_3), pH 9.5 and coated on an ELISA plate overnight at 4°C. The next day, the mannan-coated plates were washed 3 times with 200 μl PBS. 100 μl of 1% HSA blocking solution was then added to the wells and incubated for 1 hour at room temperature. The plates were washed 3 times with 200 μl PBS, and stored on ice with 200 μl PBS until addition of the samples.

Normal human serum was diluted to 0.5% in CaMgGVB buffer, and scFv clones or the OMS100 Fab2 positive control were added in triplicates at 0.01 $\mu\text{g/mL}$; 1 $\mu\text{g/mL}$ (only OMS100 control) and 10 $\mu\text{g/mL}$ to this buffer and preincubated 45 minutes on ice before addition to the blocked ELISA plate. The reaction was initiated by incubation for one hour at 37°C and was stopped by transferring the plates to an ice bath. C3b deposition was detected with a Rabbit α -Mouse C3c antibody followed by Goat α -Rabbit HRP. The negative control was buffer without antibody (no antibody = maximum C3b deposition), and the positive control was buffer with EDTA (no C3b deposition). The background was determined by carrying out the same assay except that the wells were

mannan-free. The background signal against plates without mannan was subtracted from the signals in the mannan-containing wells. A cut-off criterion was set at half of the activity of an irrelevant scFv clone (VZV) and buffer alone.

Results: Based on the cut-off criterion, a total of 13 clones were found to block the activity of MASP-2. All 13 clones producing > 50% pathway suppression were selected and sequenced, yielding 10 unique clones. All ten clones were found to have the same light chain subclass, λ 3, but three different heavy chain subclasses: VH2, VH3 and VH6. In the functional assay, five out of the ten candidate scFv clones gave IC₅₀ nM values less than the 25 nM target criteria using 0.5% human serum.

To identify antibodies with improved potency, the three mother scFv clones, identified as described above, were subjected to light-chain shuffling. This process involved the generation of a combinatorial library consisting of the VH of each of the mother clones paired up with a library of naïve, human lambda light chains (VL) derived from six healthy donors. This library was then screened for scFv clones with improved binding affinity and/or functionality.

TABLE 19: Comparison of functional potency in IC₅₀ (nM) of the lead daughter clones and their respective mother clones (all in scFv format)

scFv clone	1% human serum C3 assay (IC ₅₀ nM)	90% human serum C3 assay (IC ₅₀ nM)	90% human serum C4 assay (IC ₅₀ nM)
17D20mc	38	nd	nd
17D20m d3521N11	26	>1000	140
17N16mc	68	nd	nd
17N16m d17N9	48	15	230

Presented below are the heavy-chain variable region (VH) sequences for the mother clones and daughter clones shown above in **TABLE 19**.

The Kabat CDRs (31-35 (H1), 50-65 (H2) and 95-107 (H3)) are bolded; and the Chothia CDRs (26-32 (H1), 52-56 (H2) and 95-101 (H3)) are underlined.

17D20_35VH-21N11VL heavy chain variable region (VH) (SEQ ID NO:67, encoded by SEQ ID NO:66)

QVTLKESGPVLVKPTETLTCTVSGFSL**SRGKMGV**SWIRQPPGKALEW
LAHIFSSDEKSYRTSLKSRLTISKDTSKNQVVLMTNMDPVDAT**YYCARIRRG**
 GIDYWGQGTLVTVSS

5 d17N9 heavy chain variable region (VH) (SEQ ID NO:68)

QVQLQQSGPGLVKPSQTLSTCAISGDSV**STSA**AWNWRQSPSRGLEW**LGRTY**
YRSKWYNDYAVSVKSRITINPDTSKNQFSLQLNSVTPEDT**AVYYCARD**PFGVPPF
 DIWGQGTMTVTVSS

10 Presented below are the light-chain variable region (VL) sequences for the
 mother clones and daughter clones.

The Kabat CDRs (24-34 (L1); 50-56 (L2); and 89-97 (L3) are bolded; and the
 Chothia CDRs (24-34 (L1); 50-56 (L2) and 89-97 (L3) are underlined. These regions are
 the same whether numbered by the Kabat or Chothia system.

15

17D20m_d3521N11 light chain variable region (VL) (SEQ ID NO:70, encoded
 by SEQ ID NO:69)

QPVLTPPPSLSVSPGQTASITCS**G**EKL**GDKYAYWY**QQKPGQSPVLVMYQ
DKQRPSGIPERFSGSNSGNTATLTISGTQAMDEADYYCQ**AWDSSTAVF**GGGTKL

20 TVL

17N16m_d17N9 light chain variable region (VL) (SEQ ID NO:71)

SYELIQPPSVSVAPGQTATITC**G**DNL**GKKRVHWY**QQRPGQAPVLVIYD
DSDRPSGIPDRFSASNSGNTATLTITRGEAGDEADYYCQ**VWDIATDHV**VFGGGT

25 KLTVLAAAGSEQKLISE

The MASP-2 antibodies OMS100 and MoAb_d3521N11VL, (comprising a heavy
 chain variable region set forth as SEQ ID NO:67 and a light chain variable region set
 forth as SEQ ID NO:70, also referred to as “OMS646” and “mAbH6”), which have both
 been demonstrated to bind to human MASP-2 with high affinity and have the ability to
 30 block functional complement activity, were analyzed with regard to epitope binding by
 dot blot analysis. The results show that OMS646 and OMS100 antibodies are highly
 specific for MASP-2 and do not bind to MASP-1/3. Neither antibody bound to MAP19

nor to MASP-2 fragments that did not contain the CCP1 domain of MASP-2, leading to the conclusion that the binding sites encompass CCP1.

The MASP-2 antibody OMS646 was determined to avidly bind to recombinant MASP-2 (K_D 60-250pM) with >5000 fold selectivity when compared to C1s, C1r or MASP-1 (see TABLE 20 below):

TABLE 20: Affinity and Specificity of OMS646 MASP-2 antibody-MASP-2 interaction as assessed by solid phase ELISA studies

Antigen	K_D (pM)
MASP-1	>500,000
MASP-2	62±23*
MASP-3	>500,000
Purified human C1r	>500,000
Purified human C1s	~500,000

*Mean±SD; n=12

OMS646 specifically blocks lectin-dependent activation of terminal complement components

Methods:

The effect of OMS646 on membrane attack complex (MAC) deposition was analyzed using pathway-specific conditions for the lectin pathway, the classical pathway and the alternative pathway. For this purpose, the Wieslab Comp300 complement screening kit (Wieslab, Lund, Sweden) was used following the manufacturer's instructions.

Results:

FIGURE 53A graphically illustrates the level of MAC deposition in the presence or absence of anti-MASP-2 antibody (OMS646) under lectin pathway-specific assay conditions. FIGURE 53B graphically illustrates the level of MAC deposition in the presence or absence of anti-MASP-2 antibody (OMS646) under classical pathway-specific assay conditions. FIGURE 53C graphically illustrates the level of MAC deposition in the presence or absence of anti-MASP-2 antibody (OMS646) under alternative pathway-specific assay conditions.

As shown in FIGURE 53A, OMS646 blocks lectin pathway-mediated activation of MAC deposition with an IC_{50} value of approximately 1nM. However, OMS646 had no effect on MAC deposition generated from classical pathway-mediated activation (FIGURE 53B) or from alternative pathway-mediated activation (FIGURE 53C).

5

Pharmacokinetics and Pharmacodynamics of OMS646 following Intravenous (IV) or Subcutaneous (SC) Administration to Mice

The pharmacokinetics (PK) and pharmacodynamics (PD) of OMS646 were evaluated in a 28 day single dose PK/PD study in mice. The study tested dose levels of 5mg/kg and 15mg/kg of OMS646 administered subcutaneously (SC), as well as a dose level of 5mg/kg OMS646 administered intravenously (IV).

With regard to the PK profile of OMS646, FIGURE 54 graphically illustrates the OMS646 concentration (mean of n=3 animals/groups) as a function of time after administration of OMS646 at the indicated dose. As shown in FIGURE 54, at 5mg/kg SC, OMS646 reached the maximal plasma concentration of 5-6 ug/mL approximately 1-2 days after dosing. The bioavailability of OMS646 at 5 mg/kg SC was approximately 60%. As further shown in FIGURE 54, at 15 mg/kg SC, OMS646 reached a maximal plasma concentration of 10-12 ug/mL approximately 1 to 2 days after dosing. For all groups, the OMS646 was cleared slowly from systemic circulation with a terminal half-life of approximately 8-10 days. The profile of OMS646 is typical for human antibodies in mice.

The PD activity of OMS646 is graphically illustrated in FIGURES 55A and 55B. FIGURES 55A and 55B show the PD response (drop in systemic lectin pathway activity) for each mouse in the 5mg/kg IV (FIGURE 55A) and 5mg/kg SC (FIGURE 55B) groups. The dashed line indicates the baseline of the assay (maximal inhibition; naïve mouse serum spiked in vitro with excess OMS646 prior to assay). As shown in FIGURE 55A, following IV administration of 5mg/kg of OMS646, systemic lectin pathway activity immediately dropped to near undetectable levels, and lectin pathway activity showed only a modest recovery over the 28 day observation period. As shown in FIGURE 55B, in mice dosed with 5mg/kg of OMS646 SC, time-dependent inhibition of lectin pathway activity was observed. Lectin pathway activity dropped to near-undetectable levels within 24 hours of drug administration and remained at low levels for at least 7 days. Lectin pathway activity gradually increased with time, but did not revert to pre-dose

levels within the 28 day observation period. The lectin pathway activity versus time profile observed after administration of 15mg/kg SC was similar to the 5 mg/kg SC dose (data not shown), indicating saturation of the PD endpoint. The data further indicated that weekly doses of 5mg/kg of OMS646, administered either IV or SC, is sufficient to achieve continuous suppression of systemic lectin pathway activity in mice.

EXAMPLE 41

This Example demonstrates that a MASP-2 inhibitory antibody (OMS646) inhibits aHUS serum-induced complement C5b-9 deposition on the surface of activated human microvascular endothelial cells (HMEC-1) after exposure to serum from patients with atypical hemolytic uremic syndrome (aHUS) obtained during the acute phase and the remission phase of the disease.

Background/Rationale: The following study was carried out to analyze aHUS serum-induced complement C5b-9 deposition on the surface of activated HMEC-1 cells after exposure to aHUS patient serum obtained (1) during the acute phase and (2) during the remission phase of the disease in the presence or absence of OMS646, a MASP-2 antibody that specifically binds to MASP-2 and inhibits lectin pathway activation.

Methods:

Patients: Four patients with aHUS, studied both during the acute phase of the disease and in remission, were selected for this study among those included in the International Registry of HUS/TTP and genotyped by the Laboratory of Immunology and Genetics of Transplantation and Rare Diseases of the Mario Negri Institute. One aHUS patient had a heterozygous p.R1210C complement factor H (CFH) mutation and one had anti-CFH autoantibodies, while no mutation or antibodies to CFH were found in the other two aHUS patients.

Tables 21 and 22 summarize the results of screening for complement gene mutations and anti-CFH autoantibodies in the four aHUS patients analyzed in this study along with clinical and biochemical data measured either during the acute phase or at remission.

TABLE 21: Clinical Parameters of the four aHUS patients in this study

Case No.	Mutation or anti-CFH Ab	Disease phase	Platelets (150-400*10 ³ /μl)	LDH (266-500 IU/l)	Hemoglobin (14-18g/dl)	s-Creatinine (0.55-1.25mg/dl)
----------	-------------------------	---------------	---	--------------------	------------------------	-------------------------------

#1	no mutations, no anti-CFH Ab	acute	31,000	1396	12.9	2.37
		remission	267,000	n.a.	11.5	3.76
#2	CFH-R1210C	acute	46,000	1962	7	5.7
		remission	268,000	440	13.4	7.24
#3	anti-CFH Ab	acute	40,000	3362	9.5	1.77
		remission	271,000	338	8.8	0.84
#4	no mutations, no anti-CFH Ab	acute	83,000	1219	7.8	6.8
		remission	222,000	495	12.2	13

Note: n.a. = not available

TABLE 22: Complement Parameters of the four aHUS patients in this study

Case No.	Mutation of anti-CFH Ab	Disease phase	Serum C3 (83-180 mg/dl)	Plasma SC5b-9 (127-400 ng/ml)
#1	no mutations, no anti-CFH Ab	acute	51	69
		remission	n.a.	117
#2	CFH-R1210C	acute	79	421
		remission	119	233
#3	anti-CFH Ab	acute	58	653
		remission	149	591
#4	no mutations, no anti-CFH Ab	acute	108	n.a.
		remission	n.a.	n.a.

Experimental Methods: Cells from a human microvascular endothelial cell line (HMEC-1) of dermal origin were plated on glass slides and used when confluent. Confluent HMEC-1 cells were activated with 10 μ M ADP (adenosine diphosphate) for 10 minutes and then incubated for four hours with serum from the four aHUS patients described above in Tables 23 and 24 collected either during the acute phase of the disease, or from the same aHUS patients at remission, or from 4 healthy control subjects. The serum was diluted 1:2 with test medium (HBSS with 0.5% BSA) in the presence or in the absence of a MASP-2 inhibitory antibody, OMS646 (100 μ g/mL), generated as described above in Example 40, or in the presence of soluble complement receptor 1 (sCR1) (150 μ g/mL), as a positive control of complement inhibition. At the end of the incubation step, the HMEC-1 cells were treated with rabbit anti-human complement C5b-9 followed by FITC-conjugated secondary antibody. In each experiment, serum from one healthy control was tested in parallel with aHUS patient serum (acute phase and remission). A confocal inverted laser microscope was used for acquisition of the fluorescent staining on the endothelial cell surface. Fifteen fields per sample were acquired and the area occupied by the fluorescent staining was evaluated by automatic

edge detection using built-in specific functions of the software Image J and expressed as pixel² per field analyzed. The fields showing the lowest and the highest values were excluded from calculation.

For the statistical analysis (one-way ANOVA followed by Tukey's test for multiple comparisons) results in pixel² of the 13 fields considered in each experimental condition for each patient and control were used.

Results:

The results of the complement deposition analysis with the sera from the four aHUS patients are summarized below in Table 23A, and the results with the sera from the four healthy subjects are summarized below in Table 23B.

TABLE 23A: Effect of complement inhibitors on aHUS serum-induced C5b-9 deposition on ADP-activated HMEC-1 cells

aHUS Patient #	aHUS acute phase			aHUS remission phase		
	untreated	+sCR1	+OMS646	untreated	+sCR1	+OMS646
Patient #1 (no mutation, no anti-CFH ab)	5076 ± 562°	551 ± 80*	3312 ± 422**	4507 ± 533°	598 ± 101§	1650 ± 223§
Patient #2 (CFH-R1210C)	5103 ± 648°	497 ± 67*	2435 ± 394*	3705 ± 570°	420 ± 65§	2151 ± 250§§§
Patient #3 (anti-CFH ab)	3322 ± 421°	353 ± 64*	2582 ± 479	6790 ± 901°	660 ± 83§	2077 ± 353§
Patient #4 (no mutations, no anti-CFH ab)	4267 ± 488°	205 ± 34*	2369 ± 265**	5032 ± 594°	182 ± 29§	3290 ± 552§§

TABLE 23B: Effect of complement inhibitors on sera from four healthy control subjects (not suffering from aHUS) on C5b-9 deposition on ADP-activated HMEC-1 cells

Healthy Control Subject #	Untreated	+sCR1	+OMS646

Control Subject #1 (assayed in parallel with aHUS subject #1)	481 ± 66	375 ± 43	213 ± 57
Control Subject #2 (assayed in parallel with aHUS subject #2)	651 ± 61	240 ± 33	490 ± 69
Control Subject #3 (assayed in parallel with aHUS subject #3)	602 ± 83	234 ± 35	717 ± 109
Control Subject #4 (assayed in parallel with aHUS subject #4)	370 ± 53	144 ± 20	313 ± 36

For Tables 23A and 23B: Data are mean ± SE. °P<0.001 vs control; *P<0.001, **P<0.01 vs aHUS acute phase untreated; §P<0.001, §§P<0.01, §§§P<0.05 vs aHUS remission phase untreated.

5

Figure 56 graphically illustrates the inhibitory effect of MASP-2 antibody (OMS646) and sCR1 on aHUS serum-induced C5b-9 deposition on ADP-activated HMEC-1 cells. In Figure 56, the data are mean ± SE. °P<0.0001 vs control; *P<0.0001 vs aHUS acute phase untreated; ^P<0.0001 vs aHUS acute phase + sCR1; §P<0.0001 vs aHUS remission phase untreated and #P<0.0001 vs aHUS remission phase + sCR1.

10

As shown in Table 23A, 23B and Figure 56, ADP-stimulated HMEC-1 cells exposed to serum from aHUS patients (collected either in the acute phase or in remission) for four hours in static conditions showed an intense deposition of C5b-9 on cell surface as detected by confocal microscopy. By measuring the area covered by C5b-9, a significantly higher amount of C5b-9 deposition was observed on cells exposed to serum from aHUS patients than on cells exposed to serum from healthy control subjects, irrespective of whether aHUS serum was collected in the acute phase or during remission. No difference in serum-induced endothelial C5b-9 deposits was observed between acute phase and remission.

15

As further shown in Table 23A, 23B and Figure 56, addition of the MASP-2

20

antibody OMS646 to aHUS serum (either obtained from patients during acute phase or in remission) led to a significant reduction of C5b-9 deposition on endothelial cell surface as compared to untreated aHUS serum. However, the inhibitory effect of OMS646 on C5b-9 deposition was less profound than the effect exerted by the complement pan-inhibitor sCR1. Indeed, a statistically significant difference was observed between aHUS serum-induced C5b-9 deposits in the presence of OMS646 vs. sCR1 (Figure 56 and Tables 23A and 23B).

When calculated as a mean of the four aHUS patients, the percentages of reduction of C5b-9 deposits (as compared with C5b-9 deposits induced by the untreated serum from the same patients taken as 100%) observed in the presence of the complement inhibitors were as follows:

Acute Phase:

sCR1 (150 µg/ml): 91% reduction in C5b-9 deposits

OMS646 (100 µg/ml): 40% reduction in C5b-9 deposits

Remission Phase:

sCR1 (150 µg/ml): 91% reduction in C5b-9 deposits

OMS646 (100 µg/ml): 54% reduction in C5b-9 deposits

Conclusion: The results described in this Example demonstrate that the lectin pathway of complement is stimulated by activated microvascular endothelial cells and that this stimulation is a significant driver for the exaggerated complement activation response characteristic of aHUS. It is also demonstrated that this stimulation of the lectin pathway and resulting exaggerated complement activation response occurs both during the acute phase and in clinical remission of aHUS. Moreover, this finding does not appear to be limited to any particular complement defect associated with aHUS. As further demonstrated in this Example, selective inhibition of the lectin pathway with a MASP-2 inhibitory antibody such as OMS646 reduces complement deposition in aHUS patients with diverse etiologies.

EXAMPLE 42

This Example demonstrates that a MASP-2 inhibitory antibody (OMS646) inhibits aHUS serum-induced platelet aggregation and thrombus formation on the surface

of activated human microvascular endothelial cells (HMEC-1) after exposure to aHUS patient serum obtained during (1) the acute phase and (2) the remission phase of aHUS.

Methods:

Patients: Three patients (patients #1, #2 and #4 as described in Tables 21, 22, 23A and 23B in Example 41) with aHUS (one patient had a heterozygous p.R1210C CFH mutation, while no mutation or anti-CFH antibodies were found in the other two patients) were studied both during the acute phase of the disease and in remission. The patients were selected for this study among those included in the International Registry of HUS/TTP and genotyped by the Laboratory of Immunology and Genetics of Transplantation and Rare Diseases of the Mario Negri Institute. Five healthy subjects were also selected as blood donors for perfusion experiments.

Methods: Confluent HMEC-1 cells were activated with 10 μ M ADP for 10 minutes and then were incubated for three hours with sera from three aHUS patients (patients #1, 2 and 4 described in Example 41) collected during the acute phase of the disease or from the same patients at remission, or with control sera from healthy subjects. The serum was diluted 1:2 with test medium (HBSS with 0.5% BSA), in the presence or in the absence of a MASP-2 inhibitory antibody, OMS646 (100 μ g/mL), generated as described in Example 40; or with sCR1 (150 μ g/mL), as a positive control of complement inhibition. For patients #1 and #2 additional wells were incubated with sera (from acute phase and remission) diluted 1:2 with test medium containing 100 μ g/mL of irrelevant isotype control antibody or with 20 μ g/mL of OMS646 (for the latter, case #1 was tested only in remission and case #2 both during the acute phase and at remission).

At the end of the incubation step, HMEC-1 cells were perfused in a flow chamber with heparinized whole blood (10 UI/mL) obtained from healthy subjects (containing the fluorescent dye mepacrine that labels platelets) at the shear stress encountered in the microcirculation (60 dynes/cm², three minutes). After three minutes of perfusion, the endothelial-cell monolayers were fixed in acetone. Fifteen images per sample of platelet thrombi on the endothelial cell surface were acquired by confocal inverted laser microscope, and areas occupied by thrombi were evaluated using Image J software. The fields showing the lowest and the highest values were excluded from calculation.

For statistical analysis (one-way ANOVA followed by Tukey's test for multiple comparisons), results in pixel² of the 13 fields considered in each experimental condition for each patient and control were used.

Results:

The results of the thrombus formation experiments with the sera from the three aHUS patients are summarized below in Table 24A, and the results with the sera from the five healthy subjects are summarized below in Table 24B.

5 TABLE 24A: Effect of complement inhibitors on aHUS serum-induced thrombus formation ($\text{pixel}^2 \pm \text{SE}$) on ADP-activated HMEC-1 Cells

Experimental conditions	Disease phase	aHUS Case #1 thrombus formation ($\text{pixel}^2 \pm \text{SE}$) (no mutation, no anti-CFH ab)	aHUS Case #2 thrombus formation ($\text{pixel}^2 \pm \text{SE}$) (CFH-R1210C)	aHUS Case #4 thrombus formation ($\text{pixel}^2 \pm \text{SE}$) (no mutations, no anti-CFH ab)
untreated	acute	5499 $\pm$ 600	22320 $\pm$ 1273°	10291 $\pm$ 1362°
	remission	6468 $\pm$ 1012°	3387 $\pm$ 443°	17676 $\pm$ 1106°
+sCR1 (150 $\mu\text{g/mL}$)	acute	4311 $\pm$ 676	5539 $\pm$ 578*	5336 $\pm$ 1214***
	remission	573 $\pm$ 316§	977 $\pm$ 102§	2544 $\pm$ 498§
+OMS646 (20 $\mu\text{g/mL}$)	acute	not determined	6974 $\pm$ 556*	not determined
	remission	832 $\pm$ 150§	1224 $\pm$ 252§	not determined
+OMS646 (100 $\mu\text{g/mL}$)	acute	3705 $\pm$ 777	9913 $\pm$ 984*	2836 $\pm$ 509*
	remission	3321 $\pm$ 945§§§	733 $\pm$ 102§	1700 $\pm$ 321§
+ irrelevant isotype control antibody (100 $\mu\text{g/mL}$)	acute	5995 $\pm$ 725	18655 $\pm$ 1699	not determined
	remission	10885 $\pm$ 1380	2711 $\pm$ 371	not determined

TABLE 24B: Effect of complement inhibitors on sera from five healthy control subjects (not suffering from aHUS) in the thrombus formation ($\text{pixel}^2 \pm \text{SE}$) assay on ADP-activated HMEC-1 Cells

Experimental conditions	Control #1 thrombus formation ($\text{pixel}^2 \pm \text{SE}$)	Control #2 thrombus formation ($\text{pixel}^2 \pm \text{SE}$)	Control #3 thrombus formation ($\text{pixel}^2 \pm \text{SE}$)	Control #4 thrombus formation ($\text{pixel}^2 \pm \text{SE}$)	Control #5 thrombus formation ($\text{pixel}^2 \pm \text{SE}$)
-------------------------	--	--	--	--	--

untreated	2880±510	1046 ± 172	1144 ± 193	735 ± 124	2811 ± 609
+sCR1 (150 µg/mL)	5192 ± 637	1527 ± 153	1198 ± 138	2239 ± 243	2384 ± 410
+OMS646 (100 µg/mL)	7637 ± 888	1036 ± 175	731 ± 203	2000 ± 356	7177 ± 1477
+irrelevant isotype control antibody (100 µg/mL)	6325 ± 697	1024 ± 235	399 ± 82	45269	not determined
Assayed in parallel with serum from aHUS subject	#1 (acute phase serum)	#1 (remission phase serum)	#2 (acute phase serum)	#2 (remission phase serum)	#5 (acute and remission phase serum)

For Tables 24A and 24B: Data are mean ± SE. °P<0.001 vs control; *P<0.001, ***P<0.05 vs aHUS acute phase untreated; §P<0.001, §§§P<0.05 vs aHUS remission phase untreated.

Figure 57 graphically illustrates the effect of MASP-2 antibody (OMS646) and sCR1 on aHUS serum-induced thrombus formation on ADP-activated HMEC-1 cells. In Figure 57, the data shown are mean ± SE. °P<0.0001, °°P<0.01 vs control; *P<0.0001, **P<0.01 vs aHUS acute phase untreated; §P<0.0001 vs aHUS remission phase untreated.

As shown in Table 24A and Figure 57, a marked increase in the area covered by thrombi was observed on HMEC-1 cells treated with aHUS serum, collected either during acute phase or at remission, in comparison to cells exposed to serum from healthy control subjects (Table 24B and Figure 57). As shown in Figure 57 and Table 24A, OMS646 (at both 100 µg/ml and 20 µg/ml) showed a partial inhibition of thrombus formation on cells pre-exposed to aHUS serum taken during the acute phase. The anti-thrombogenic effect was comparable between the two different doses of OMS646 and was not different from

the effect of sCR1 (Figure 57 and Table 24A). Addition of the irrelevant isotype control antibody had no inhibitory effect on aHUS-serum-induced thrombus formation.

As further shown in Figure 57 and Table 24A, the inhibitory effect of OMS646 was even more evident on aHUS serum collected during remission phase. Indeed, the addition of OMS646, at both 100 µg/ml and 20 µg/ml doses, to aHUS patient serum collected at remission resulted in a nearly complete inhibition of thrombus formation, similar to that observed with the addition of sCR1. The irrelevant isotype control antibody showed no significant inhibitory effect.

When calculated as a mean of the three aHUS patients, the percentages of reduction of the HMEC-1 surface covered by thrombi deposits (as compared with thrombus area induced by the untreated sera from the same patients taken as 100%) recorded with the complement inhibitors were as follows:

Acute Phase:

sCR1 (150 µg/ml): 60% reduction

OMS646 (100 µg/ml): 57% reduction

OMS646 (20 µg/ml): 45% reduction

Remission Phase:

sCR1 (150 µg/ml): 85% reduction

OMS646 (100 µg/ml): 79% reduction

OMS646 (20 µg/ml): 89% reduction

Discussion of Results:

The results in this Example demonstrate that a MASP-2 inhibitory antibody, such as OMS646 (generated as described in Example 40), has a strong inhibitory effect on aHUS serum-induced thrombus formation on HMEC-1 cells. Surprisingly, the inhibitory effect of OMS646 on thrombus formation was greater than its effect on C5b-9 deposits induced on HMEC-1 (as described in Example 41). It is also surprising that the addition of OMS646, at both 100 µg/ml and 20 µg/ml doses, to aHUS patient serum collected at remission resulted in nearly a complete inhibition of thrombus formation. Another surprising finding is the observation that OMS646, in both the acute phase and in remission, was as effective as the positive control sCR1, which is a broad and almost complete inhibitor of the complement system (Weisman H. et al., *Science* 249:146-151,

1990; Lazar H. et al., *Circulation* 100:1438-1442, 1999).

It is noted that the control serum from healthy subjects also induced a modest thrombus formation on HMEC-1 cells. We did not observe a consistent inhibitory effect on control serum induced thrombus formation with either OMS646 or with sCR1. While not wishing to be bound by any particular theory, it is believed that the control-induced thrombi do not depend upon complement, as supported by very low C5b-9 deposits observed on HMEC-1 incubated with control serum (see Example 41).

Conclusion:

In conclusion, the observed anti-thrombotic effect of a MASP-2 inhibitory antibody, such as OMS646, appears substantially greater than one would have expected based on the inhibitory effect of OMS646 on C5b-9 deposition observed in this experimental system (as described in Example 41 and shown in Figure 56). For example, as described in Gastoldi et al., *Immunobiology* 217:1129-1222 Abstract 48 (2012) entitled "C5a/C5aR interaction mediates complement activation and thrombosis on endothelial cells in atypical hemolytic uremic syndrome (aHUS)," it was determined that addition of a C5 antibody inhibiting C5b-9 deposits (60% reduction) limited thrombus formation on HMEC-1 to a comparable extent (60% reduction). In contrast, the MASP-2 inhibitory antibody (OMS646 at 100 µg/mL) inhibited C5b-9 deposits with mean values of (acute phase = 40% reduction; remission phase = 54% reduction); and inhibited thrombus formation at a substantially higher percent (acute phase = 57% reduction; remission phase = 79% reduction). In comparison, OMS646 inhibited complement deposition at a lower percentage than did the positive control complement inhibitor (sCR1 at 150 µg/mL, acute phase inhibition of C5b-9 deposition = 91% reduction; remission phase = 91% reduction) yet was equally effective as the sCR1 positive control in inhibiting thrombus formation (sCR1 at 150 µg/mL, acute phase = 60% reduction; remission phase = 85% reduction). These results demonstrate that a MASP-2 inhibitory antibody (e.g., OMS646) is surprisingly effective at inhibiting thrombus formation in serum obtained from aHUS subjects both in the acute phase and remission phase.

In accordance with the foregoing, in one embodiment, the invention provides a method of inhibiting thrombus formation in a subject suffering from, or at risk for developing, a thrombotic microangiopathy (TMA) comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation. In one embodiment, the TMA is

selected from the group consisting of hemolytic uremic syndrome (HUS), thrombotic thrombocytopenic purpura (TTP) and atypical hemolytic uremic syndrome (aHUS). In one embodiment, the TMA is aHUS. In one embodiment, the composition is administered to an aHUS patient during the acute phase of the disease. In one embodiment, the composition is administered to an aHUS patient during the remission phase (i.e., in a subject that has recovered or partially recovered from an episode of acute phase aHUS, such remission evidenced, for example, by increased platelet count and/or reduced serum LDH concentrations, for example as described in Loirat C et al., *Orphanet Journal of Rare Diseases* 6:60, 2011, hereby incorporated herein by reference).

10 In one embodiment, the MASP-2 inhibitory antibody exhibits at least one or more of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or less, said antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less, wherein the antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation, and/or wherein the antibody does not substantially inhibit the classical pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the alternative pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway or the alternative pathway (i.e., inhibits the lectin pathway while leaving the classical and alternative complement pathways intact).

25 In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a subject suffering from a TMA such as aHUS (acute or remission phase), by at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum. In some embodiments, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a subject suffering from aHUS at a level of at least 20 percent or greater, (such as at least 30%, at

least 40%, at least 50%) more than the inhibitory effect on C5b-9 deposition in serum.

In one embodiment, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from an aHUS patient in remission phase by at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such
5 as at least 85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum. In some embodiments, the MASP-2 inhibitory antibody inhibits thrombus formation in serum in an aHUS patient in remission phase at a level of at least 20 percent or greater, (such as at least 30%, at least 40%, at least 50%) more than the inhibitory effect on C5b-9 deposition in serum.

10 In one embodiment, the MASP-2 inhibitory antibody is administered to the subject via an intravenous catheter or other catheter delivery method.

In one embodiment, the invention provides a method of inhibiting thrombus formation in a subject suffering from a TMA comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding
15 fragment thereof, comprising (I) (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-102 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-
20 L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at least 93%, at least
25 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70.

In one embodiment, the TMA is selected from the group consisting of atypical
30 hemolytic uremic syndrome (aHUS) (either acute or remission phase), HUS and TTP. In one embodiment, the subject is in acute phase of aHUS. In one embodiment, the subject is in remission phase of aHUS.

In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67. In some embodiments, the method comprises
5 administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment
10 thereof, that specifically recognizes at least part of an epitope on human MASP-2 recognized by reference antibody OMS646 comprising a heavy-chain variable region as set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70. Competition between binding members may be assayed easily *in vitro*, for example using ELISA and/or by tagging a specific reporter molecule to one binding
15 member which can be detected in the presence of other untagged binding member(s), to enable identification of specific binding members which bind the same epitope or an overlapping epitope. Thus, there is presently provided a specific antibody or antigen-binding fragment thereof, comprising a human antibody antigen-binding site, which competes with reference antibody OMS646 for binding to human MASP-2.

20

EXAMPLE 43

This Example demonstrates that a human MASP-2 inhibitory antibody (OMS646) is able to inhibit TMA patient plasma-mediated induction of apoptosis in primary human microvascular endothelial cells (MVECs) of dermal origin.

25 Background/Rationale:

The pathophysiology of TMA is known to involve an endothelial cell injury induced by various factors that is followed by occlusions of small vessels (e.g., small arterioles and capillaries) by platelet plugs and/or fibrin thrombi (Hirt-Minkowsk P. et al., *Nephron Clin Pract* 114:c219-c235, 2010; Goldberg R.J. et al., *Am J Kidney Dis*
30 56(6):1168-1174, 2010). It has been shown that MVECs undergo apoptotic injury when exposed *in vitro* to plasma from patients with TMA-related disorders (see Stefanescu et al., *Blood* Vol 112 (2):340-349, 2008; Mitra D. et al., *Blood* 89:1224-1234, 1997). Apoptotic injury associated with TMAs has been documented in MVEC obtained from

tissue biopsies (skin, bone, marrow, spleen, kidney, ileum) of such patients. It has also been shown that apoptotic insults to MVECs reduces the levels of membrane-bound complement regulatory proteins in MVECs (see e.g., Mold & Morris, *Immunology* 102:359-364, 2001; Christmas et al., *Immunology* 119:522, 2006).

5 A positive feedback loop involving terminal complement components is believed to be involved in the pathophysiology of TMAs including atypical hemolytic-uremic syndrome (aHUS), and TMAs associated with catastrophic antiphospholipid syndrome (CAPS), Degos disease, and TMAs secondary to cancer, cancer chemotherapy, autoimmunity and transplantation, each of these conditions are known or thought to be
10 responsive to anti-C5 therapy with the mAb eculizumab (Chapin J. et al., *Brit. J. Hematol* 157:772-774, 2012; Tsai et al., *Br J Haematol* 162(4):558-559, 2013); Magro C. M. et al., *Journal of Rare Diseases* 8:185, 2013).

The following experiment was carried out to analyze the ability of human MASP-2 inhibitory antibody (OMS646) to block TMA patient plasma-mediated induction of
15 apoptosis in primary human dermal MVECs in plasma samples obtained from patients suffering from aHUS, ADAMTS13 deficiency-related thrombotic thrombocytopenic purpura (TTP), CAPS and systemic Degos disease, as well as TMAs secondary to cancer, transplantation, autoimmune disease and chemotherapy.

Methods:

20 An *in vitro* assay was carried out to analyze the efficacy of a MASP-2 inhibitory antibody (OMS646) to block TMA patient plasma-mediated induction of apoptosis in primary human MVECs of dermal origin as described in Stefanescu R. et al., *Blood* Vol 112 (2):340-349, 2008, which is hereby incorporated herein by reference. The plasma samples used in this assay were obtained from a collection of healthy control subjects and
25 from individuals with either acute-phase or convalescent thrombotic microangiopathies. The presence of microangiopathy in the TMA patients was assessed by detecting schistocytes on a peripheral blood smear. In addition, TTP was diagnosed as described in Stefanescu R. et al., *Blood* Vol 112 (2):340-349, 2008.

Endothelial Cell (EC) Culture

30 As described in Stefanescu et al., primary human MVECs of dermal origin were purchased from ScienCell Research Labs (San Diego, CA). MVECs expressed CD34 up through passages 5 and 6 (*Blood* 89:1224-1234, 1997). The MVECs were maintained in polystyrene flasks coated with 0.1% gelatin in water in ECM1001 medium (ScienCell

Research Labs) containing endothelial cell growth supplement, penicillin, streptomycin and 15% fetal bovine serum. All MVECs were used in passages 2 to 6. Subcultures involved a 5 to 10 minute exposure to 0.25% trypsin-EDTA.

Apoptosis Assay

5 Representative primary human MVECs of dermal origin known to be susceptible to TTP/HUS plasma-induced apoptosis were washed with phosphate buffered saline (PBS) and plated in chambers of 12-well plates, coated with 0.1% gelatin in water at 0.15×10^6 viable cells/mL. The plated MVEC cells were starved in complete media for 24 hours then exposed to varying concentrations (2% to 20% v/v) of TMA patient plasma
 10 samples or healthy donor plasma for 18 hours in the presence or absence of MASP-2 mAb OMS646 (150 μ g/mL) and the cells were then harvested by trypsinization. Each TMA patient sample was analyzed in duplicate. The degree of plasma-mediated apoptosis was assessed using propidium iodide (PI) staining, with $>5 \times 10^3$ cells analyzed in a cytofluorograph and A0 peaks defined by computer software (MCycle Av, Phoenix
 15 Flow Systems, San Diego, CA). Enzyme-linked immunosorbent assay (ELISA)-based quantitation of cytoplasmic histone-associated DNA fragments from cell lysate was also performed as per the manufacturer's directions (Roche Diagnostics, Mannheim, Germany).

20 Results:

The results of the TMA patient plasma-induced MVEC apoptosis assay in the presence of MASP-2 mAb (OMS646) are shown below in Table 25.

25 **Table 25:** TMA patient plasma tested on primary human MVEC of dermal origin in the presence of MASP-2 mAb (OMS646)

Subject #	Age/ Sex	Clinical Diagnosis (TMA) and other conditions	MASP-2 ng/ml	Diagnosis based on Cre/LDH	C5a	sC5- b9	ADAM S Activity	Diagnosis based on ADAMS activity	protection with OMS646
#2	41/f	TTP	174	TTP	34.42	772	30%	aHUS	responder
#3	52/f	TTP	150	TTP	48.32	1399	70%	aHUS	non- responder

Subject #	Age/ Sex	Clinical Diagnosis (TMA) and other conditions	MASP-2 ng/ml	Diagnosis based on Cre/LDH	C5a	sC5- b9	ADAM S Activity	Diagnosis based on ADAMS activity	protection with OMS646
#4	20/ m	TTP	224	TTP	36.9	1187	<10%	TTP	responder
#10	60/f	TTP	175.4	TTP	49.5	4406	64%	aHUS	non- responder
#11	59/f	TTP	144.9	TTP	40.3	1352	<10%	TTP	non- responder
#13	49/f	HUS, Cancer, TTP	142.8	TTP	48.6	3843	86%	aHUS	non- responder
#42	27/ m	TTP	341.5	TTP	100.0	5332	<5%	TTP	non- responder
#46	25/f	TTP, Degos, SLE	225.11	TTP	53.9	3426	ND	ND	responder
#48	53/f	TTP, SLE, nephritis s/p renal transplan t	788.5	aHUS	31.2	1066	66%	aHUS	responder
#49	64/f	TTP, APLAs, CVA	494.5		35.4	2100	ND	ND	responder
#51	25/f	aHUS, APLAs	313.1	TTP	26.8	1595	23%	aHUS	responder
#52	56/f	aHUS, SLE	333.1	TTP	18.9	1103	97%	aHUS	non- responder
#53	56/f	aHUS remission	189.9	Remissio n TTP	28.69	344	74%	aHUS	non- responder

Abbreviations used in Table 25:

“APLAs” = antiphospholipid antibodies, associated with Catastrophic antiphospholipid syndrome (CAPS).

“SLE” = systemic lupus erythematosus

“CVA” = cerebrovascular accident (stroke)

5

Consistent with the results reported in Stefanescu R. et al., *Blood* Vol 112 (2):340-349, 2008, significant apoptosis was observed for primary MVECs of dermal origin in the presence of the thirteen TMA patient plasma samples in the absence of MASP-2 antibody. Control plasma samples from healthy human subjects were run in parallel and did not induce apoptosis in the MVECs (data not shown). As shown in Table 25, the MASP-2 inhibitory mAb (OMS646) inhibited TMA patient plasma-mediated induction of apoptosis in primary MVECs (“responders” in Table 25) in 6 of the 13 patient plasma samples tested (46%). In particular, it is noted that MASP-2 inhibitory mAb (OMS646) inhibited apoptosis in plasma obtained from patients suffering from aHUS, TTP, Degos disease, SLE, transplant, and APLAs (CAPS). With regard to the seven patient samples tested in this assay in which the MASP-2 mAb did not block apoptosis (“non-responders” in Table 25), it is noted that apoptosis can be induced by several pathways, not all of which are complement dependent. For example, as noted in Stefanescu R. et al., *Blood* Vol 112 (2):340-349, 2008, apoptosis in an EC assay is dependent on the basal EC activation state which is influenced by plasma factors which may play a role in determining the level of insult required to induce apoptosis. As further noted in Stefanescu R. et al., additional factors capable of modulating apoptosis may be present in the TMA patient plasma, such as cytokines and various components of the complement system. Therefore, due to these complicating factors, it is not surprising that the MASP-2 antibody did not show a blocking effect in all of the plasma samples that exhibited TMA-plasma induced apoptosis.

Further in this regard, it is noted that a similar analysis was carried out using TMA-plasma induced apoptosis assay with the anti-C5 antibody eculizumab and very similar results were observed (see Chapin et al., *Blood* (ASH Annual Meeting Abstracts): Abstract #3342, 120: 2012). Clinical efficacy of eculizumab, a highly successful commercial product, appears greater than the efficacy demonstrated in this model, suggesting that this in vitro model may underestimate the clinical potential of complement inhibitory drugs.

These results demonstrate that a MASP-2 inhibitory antibody such as OMS646 is effective at inhibiting TMA-plasma-induced apoptosis in plasma obtained from patients suffering from a TMA such as aHUS, TTP, Degos disease, SLE, transplant, and APLAs (CAPS). It is known that endothelial damage and apoptosis play a key role in the pathology of TMAs such as idiopathic TTP and sporadic HUS (Kim et al., *Microvascular Research* vol 62(2):83-93, 2001). As described in Dang et al., apoptosis was demonstrated in the splenic red pulp of TTP patients but not in healthy control subjects (Dang et al., *Blood* 93(4):1264-1270, 1999). Evidence of apoptosis has also been observed in renal glomerular cells of MVEC origin in an HUS patient (Arends M. J. et al., *Hum Pathol* 20:89, 1989). Therefore, it is expected that administration of a MASP-2 inhibitory agent, such as a MASP-2 inhibitory antibody (e.g., OMS646) will be an effective therapy in patients suffering from a TMA such as aHUS, TTP or other microangiopathic disorder such as other TMAs including CAPS, systemic Degos disease, and a TMA secondary to cancer; a TMA secondary to chemotherapy, or a TMA secondary to transplantation.

In accordance with the foregoing, in one embodiment, the invention provides a method of inhibiting endothelial cell damage and/or endothelial cell apoptosis, and/or thrombus formation in a subject suffering from, or at risk for developing, a thrombotic microangiopathy (TMA) comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation. In one embodiment, the TMA is selected from the group consisting of atypical hemolytic uremic syndrome (aHUS), thrombotic thrombocytopenic purpura (TTP) and hemolytic uremic syndrome (HUS). In one embodiment, the TMA is aHUS. In one embodiment, the composition is administered to an aHUS patient during the acute phase of the disease. In one embodiment, the composition is administered to an aHUS patient during the remission phase (i.e., in a subject that has recovered or partially recovered from an episode of acute phase aHUS, such remission evidenced, for example, by increased platelet count and/or reduced serum LDH concentrations, for example as described in Loirat C et al., *Orphanet Journal of Rare Diseases* 6:60, 2011, hereby incorporated herein by reference).

In one embodiment, the subject is suffering from, or at risk for developing a TMA that is (i) a TMA secondary to cancer; (ii) a TMA secondary to chemotherapy; or (iii) a TMA secondary to transplantation (e.g., organ transplantation, such as kidney transplantation or

allogeneic hematopoietic stem cell transplantation). In one embodiment, the subject is suffering from, or at risk for developing Upshaw-Schulman Syndrome (USS). In one embodiment, the subject is suffering from, or at risk for developing Degos disease. In one embodiment, the subject is suffering from, or at risk for developing Catastrophic Antiphospholipid Syndrome (CAPS).

In accordance with any of the disclosed embodiments herein, the MASP-2 inhibitory antibody exhibits at least one or more of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or less, said antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less, wherein the antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation, and/or wherein the antibody does not substantially inhibit the classical pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the alternative pathway. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway (i.e., inhibits the lectin pathway while leaving the classical complement pathway intact).

In one embodiment, the MASP-2 inhibitory antibody inhibits plasma induced MVEC apoptosis in serum from a subject suffering from a TMA such as aHUS (acute or remission phase), hemolytic uremic syndrome (HUS), thrombotic thrombocytopenic purpura (TTP), a TMA secondary to cancer; a TMA secondary to chemotherapy; a TMA secondary to transplantation (e.g., organ transplantation, such as kidney transplantation or allogeneic hematopoietic stem cell transplantation), or in serum from a subject suffering from Upshaw-Schulman Syndrome (USS), or in serum from a subject suffering from Degos disease, or in a subject suffering from Catastrophic Antiphospholipid Syndrome (CAPS), wherein the plasma induced MVEC apoptosis is inhibited by at least 5%, such as at least 10%, such as at least 20%, such as at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80% such as at least

85%, such as at least 90%, such as at least 95% up to 99%, as compared to untreated serum. In some embodiments, the MASP-2 inhibitory antibody inhibits thrombus formation in serum from a subject suffering from a TMA (e.g., such as aHUS (acute or remission phase), hemolytic uremic syndrome (HUS), thrombotic thrombocytopenic purpura (TTP), a TMA secondary to cancer; a TMA secondary to chemotherapy; a TMA secondary to transplantation (e.g., organ transplantation, such as kidney transplantation or allogeneic hematopoietic stem cell transplantation), or in serum from a subject suffering from Upshaw-Schulman Syndrome (USS), or in serum from a subject suffering from Degos disease, or in a subject suffering from Catastrophic Antiphospholipid Syndrome (CAPS)), at a level of at least 20 percent or greater, (such as at least 30%, at least 40%, at least 50%) more than the inhibitory effect on C5b-9 deposition in serum.

In one embodiment, the MASP-2 inhibitory antibody is administered to the subject via an intravenous catheter or other catheter delivery method.

In one embodiment, the invention provides a method of inhibiting thrombus formation in a subject suffering from a TMA (such as aHUS (acute or remission phase), hemolytic uremic syndrome (HUS), thrombotic thrombocytopenic purpura (TTP), a TMA secondary to cancer; a TMA secondary to chemotherapy; a TMA secondary to transplantation (e.g., organ transplantation, such as kidney transplantation or allogeneic hematopoietic stem cell transplantation), or in serum from a subject suffering from Upshaw-Schulman Syndrome (USS), or in serum from a subject suffering from Degos disease, or in a subject suffering from Catastrophic Antiphospholipid Syndrome (CAPS)), comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising (I) (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-102 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at

least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70.

In one embodiment, the subject is suffering from a TMA selected from the group
5 consisting of a TMA secondary to cancer; a TMA secondary to chemotherapy; a TMA secondary to transplantation (e.g., organ transplantation, such as kidney transplantation or allogeneic hematopoietic stem cell transplantation), Upshaw-Schulman Syndrome (USS), Degos disease, and Catastrophic Antiphospholipid Syndrome (CAPS).

In some embodiments, the method comprises administering to the subject a
10 composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67. In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a light-chain variable region
15 comprising the amino acid sequence set forth as SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof, that specifically recognizes at least part of an epitope on human MASP-2 recognized by reference antibody OMS646 comprising a heavy-chain variable region as
20 set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70.

EXAMPLE 44

This Example describes the initial results of an ongoing Phase 2 Clinical Trial to
25 Evaluate the Safety and Clinical Efficacy of a fully human monoclonal MASP-2 inhibitory antibody in Adults with Thrombotic Microangiopathies (TMAs).

Background: TMAs are a family of rare, debilitating and life-threatening disorders characterized by excessive thrombi (clots) – aggregations of platelets – in the microcirculation of the body's organs, most commonly the kidney and brain.

30 Methods:

The first stage of an open-label Phase 2 clinical trial was carried out in subjects with primary atypical hemolytic uremic syndrome (aHUS), plasma therapy-resistant

aHUS and thrombotic thrombocytopenic purpura (TTP). This Phase 2 clinical trial has no placebo arm given the life-threatening nature of the disease.

The subjects were age ≥ 18 at screening and were only included in this study if they had a diagnosis of one of the following TMAs:

- 5 1) Primary aHUS, diagnosed clinically and having ADAMTS13 activity $> 10\%$ in plasma. Patients are eligible with or without a documented complement mutation or anti-CFH antibody. Patients are categorized according to their response to plasma therapy (plasma exchange or plasma infusion):
 - 10 ○ Plasma therapy-resistant aHUS patients, for purposes of this study, meet all of the following:
 - (a) screening platelet count $< 150,000/\mu\text{L}$ despite at least four plasma therapy treatments prior to screening;
 - (b) evidence of microangiopathic hemolysis (presence of schistocytes, serum lactate dehydrogenase (LDH) $>$ upper limit of normal (ULN),
15 haptoglobin $<$ LLN); and
 - (c) serum creatinine $>$ ULN.
 - Chronic plasma therapy-responsive aHUS patients (plasma therapy-sensitive) must require at least once-per-week plasma therapy for four weeks before first dose of OMS646 with serum creatinine $>$ ULN.
- 20 2) TTP defined as having all of the following:
 - Platelet count $< 150,000/\mu\text{L}$;
 - Evidence of microangiopathic hemolysis (presence of schistocytes, serum LDH $>$ ULN, or haptoglobin $<$ LLN); and
 - ADAMTS13 activity $\leq 10\%$ during the current episode of TTP or
25 historically.

Note: Subjects were excluded from the study if they had eculizumab therapy within three months prior to screening. The criteria for a patient qualifying as plasma therapy-resistant were for purposes of determining eligibility in this study. In further aspects of
30 the invention, a plasma therapy-resistant patient to be treated with a MASP-2 inhibitor was previously treated with plasma therapy at least once and after such plasma therapy treatment still had one or more clinical markers of aHUS that were not adequately reduced or eliminated by such plasma therapy treatment.

The monoclonal antibody used in this study, OMS646, is a fully human IgG4 mAb directed against human MASP-2. As demonstrated in Example 40, OMS646 avidly binds to recombinant MASP-2 (apparent equilibrium dissociation constant in the range of 100 pM) and exhibits greater than 5,000-fold selectivity over the homologous proteins C1s, C1r, and MASP-1. In functional assays, OMS646 inhibits the human lectin pathway with nanomolar potency (concentration leading to 50% inhibition [IC₅₀] of approximately 3 nM) but has no significant effect on the classical pathway. OMS646 administered either by intravenous (IV) or subcutaneous (SC) injection to mice, non-human primates, and humans resulted in high plasma concentrations that were associated with suppression of lectin pathway activation in an *ex vivo* assay. As further described in Example 42, OMS646 treatment reduced C5b-9 deposition and thrombus formation in *in vitro* models of TMA and thrombus formation in a mouse model of TMA, thus demonstrating that OMS646 has therapeutic utility.

In this study, the OMS646 drug substance was provided at a concentration of 100 mg/mL, which was further diluted for IV administration. The appropriate calculated volume of OMS646 100 mg/mL injection solution was withdrawn from the vial using a syringe for dose preparation. The infusion bag was administered within four hours of preparation.


In Stage 1 of the study, OMS646 was administered to escalating dose cohorts of three subjects per cohort. Each subject in Stage 1 received four weekly doses of OMS646 as shown below in TABLE 26.

TABLE 26: Dosing Schedule For Stage 1

Stage, Cohort	Number of Subjects	OMS646 Dose (mg/kg)
1, Cohort 1	3	0.675, weekly x 4
1, Cohort 2	3	2.0, weekly x 4
1, Cohort 3	3	4.0, weekly x 4

25

Stage 1 Study Design Schematic



The diluted study drug was infused intravenously over a 30-minute period.

5

Primary Endpoints

The co-primary endpoints were:

- Safety as assessed by AEs, vital signs, ECGs, and clinical laboratory tests
- Clinical activity as assessed by change in platelet count

10 Secondary Endpoints

The secondary endpoints were:

- TMA clinical activity
 - Serum LDH
 - Serum haptoglobin
 - Hemoglobin
 - Serum creatinine
 - TMA-related symptoms
 - Need for plasma therapy (plasma exchange or plasma infusion)
 - Need for dialysis
- 15

Allowed Concomitant Therapies

- Plasma therapy-resistant aHUS – plasma therapy during OMS646 treatment was allowed if the investigator considered it to be medically indicated.
- 5 • Chronic plasma therapy-responsive aHUS – investigators were advised that plasma therapy should be continued until there was a sign of improvement in TMA, e.g., increase in platelet count, decrease in LDH, increase in haptoglobin, increase in hemoglobin, decrease in creatinine, at which time the investigator was advised to consider withholding plasma therapy and monitoring TMA parameters to assess whether plasma therapy can be discontinued.
- 10 • TTP – plasma therapy was allowed if the investigator considered it to be medically indicated.
- Investigators were advised that renal dialysis therapy should be managed according to standard of care.
- 15 • Eculizumab was not administered during the study.

Results:

The first cohort of subjects consisted of three aHUS patients treated with the lowest dose of OMS646. Improvements were observed across TMA disease markers in all patients in this study cohort. Platelet count, serum lactate dehydrogenase (LDH) and serum haptoglobin were measured as markers of disease activity. When compared to baseline levels, platelet counts improved in all patients. Serum LDH levels remained normal in one patient, substantially decreased to close to the normal range in another and remained elevated in the third. Haptoglobin improved in two patients, normalizing in one. Creatinine levels in the one patient with independent renal function also improved.

25 As designed, three patients were treated in the second, mid-dose cohort of this clinical trial. Two patients have plasma therapy-resistant aHUS and one patient has thrombotic thrombocytopenic purpura (TTP). Both patients with aHUS were on renal dialysis prior to and at the time of study enrollment. In the second or mid-dose cohort, the two patients with plasma therapy-resistant aHUS demonstrated:

- 30 • 47% increase in mean platelet count, resulting in both patients having counts in the normal range

- 86% decrease in mean schizocyte count, with schistocytes disappearing in one patient
- 71% increase in mean haptoglobin with both patients reaching the normal range during treatment, one slipping slightly below normal at one week following the last dose
- 5% decrease in the mean levels of LDH, with levels in both patients remaining slightly elevated above normal range.

The mid-dose-cohort patient with TTP required repeated plasma infusion therapy prior to entering the study. Laboratory parameters did not show consistent improvement, but the patient did not require plasma therapy while on treatment with OMS646, nor, to date, since completion of treatment.

The drug was well tolerated by all patients throughout the treatment period. Based on the positive data from the second or mid-dose cohort, the third or high-dose cohort was initiated and an aHUS patient has already completed the study treatment period. The data referenced for all patients include measures to one week following the last dose.

The first patient in the third (high-dose) cohort – a plasma therapy-resistant aHUS patient with additional complicating disorders including hepatitis C, cryoglobulinemia and lymphoma – has also completed treatment with OMS646. Prior to OMS646 treatment, the patient required repeated dialysis. Throughout treatment and following completion of the OMS646 course, to date the patient has remained off dialysis. Hematological and renal parameters showed:

- 63% improvement in platelet count, returning to normal levels
- 100% decrease in schistocytes
- Haptoglobin increased from an undetectable level and normalized
- 43% decrease in LDH, resulting in a level just slightly above normal
- 24% reduction in creatinine level

As in the first and second cohorts, the drug was well tolerated.

The primary endpoint in this open-label Phase 2 clinical trial is change in platelet count. Platelet counts in all three aHUS patients in the mid- and high-dose cohorts (two in the mid-dose and one in the high-dose cohort) returned to normal, with a statistically

significant mean increase from baseline of approximately 68,000 platelets/mL (p=0.0055).

In summary, the data obtained so far from this Phase 2 clinical trial show efficacy of OMS646 in patients with primary aHUS, plasma-therapy resistant aHUS and in
5 patients with TTP.

In accordance with the foregoing, in one embodiment, the invention provides a method of treating a subject suffering from plasma therapy-resistant aHUS comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation. In some
10 embodiments, the method further comprises the step of identifying a subject suffering from plasma therapy-resistant aHUS prior to administering to the subject a composition comprising a MASP-2 inhibitory antibody.

In accordance with any of the disclosed embodiments herein, the MASP-2 inhibitory antibody exhibits at least one or more of the following characteristics: said
15 antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or less, said antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less, wherein the antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and
20 F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway (i.e., inhibits the lectin pathway while leaving
25 the classical complement pathway intact).

In one embodiment, the MASP-2 inhibitory antibody is administered in an amount effective to improve at least one or more clinical parameters associated with aHUS such as an increase in platelet count, a decrease in LDH, an increase in haptoglobin, an increase in hemoglobin, and/or a decrease in creatinine.

30 In some embodiments, the method comprises administering a MASP-2 inhibitory antibody to a subject suffering from plasma therapy-resistant aHUS via a catheter (e.g., intravenously) for a first time period (e.g., at least one day to a week or two weeks)

followed by administering a MASP-2 inhibitory antibody to the subject subcutaneously for a second time period (e.g., a chronic phase of at least two weeks or longer). In some embodiments, the administration in the first and/or second time period occurs in the absence of plasma therapy. In some embodiments, the administration in the first and/or
5 second time period occurs in the presence of plasma therapy.

In some embodiments, the method comprises administering a MASP-2 inhibitory antibody, to a subject suffering from plasma therapy-resistant aHUS either intravenously, intramuscularly, or preferably, subcutaneously. Treatment may be chronic and administered daily to monthly, but preferably every two weeks. The MASP-2 inhibitory
10 antibody may be administered alone, or in combination with a C5 inhibitor, such as eculizumab.

In one embodiment, the method comprises treating a subject suffering from plasma therapy-resistant aHUS comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising (I) (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-107 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70.
15
20
25

In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67 and a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70.
30

In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof, that specifically recognizes at least part of an epitope on human MASP-2 recognized by reference antibody OMS646 comprising a heavy-chain variable region as set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70.

EXAMPLE 45

This Example describes additional results obtained in the ongoing Phase 2 Clinical trial to evaluate the Safety and Clinical Efficacy of a fully human monoclonal MASP-2 inhibitory antibody in Adults with Thrombotic Microangiopathies (TMAs) described in Example 44.

Background: As described in Example 44, TMAs are a family of rare, debilitating and life-threatening disorders characterized by excessive thrombi (clots) – aggregations of platelets – in the microcirculation of the body's organs, most commonly the kidney and brain. As described herein, transplantation-associated TMA (TA-TMA) is a devastating syndrome that can occur in transplant patients, such as hematopoietic stem cell transplant (HSCT) recipients. This Example describes the initial results of the Phase 2 Clinical trial to evaluate the safety and clinical efficacy of a fully human monoclonal MASP-2 inhibitory antibody in a patient suffering from hematopoietic stem cell transplant-related TMA.

Methods:

As described in Example 44, the Phase 2 TMA trial consists of a three-level dose-ranging stage, followed by a fixed-dose stage of the MASP-2 inhibitory antibody OMS646. As further described in Example 44, positive results were obtained from aHUS patients and one TTP patient, with consistent and robust improvement in efficacy measures. As in the low-dose cohort, OMS646 was well tolerated by all patients in the mid- and high-dose cohorts throughout the treatment period. Preclinical toxicity studies have been completed and demonstrated no safety concerns, allowing chronic dosing in clinical trials.

As described in Example 44, data from the OMS646 Phase 2 TMA clinical trial were obtained from aHUS and TTP patients. Dosing has now been completed for an

additional hematopoietic stem cell transplant-related TMA patient in the high-dose cohort using the methods described in Example 44. This is a patient with a history of lymphoma for which he underwent hematopoietic stem cell transplant. His post-transplant course has been complicated by a number of life-threatening disorders, including platelet transfusion-requiring TMA. Despite transfusions, his stem cell transplant-related TMA persisted and he was enrolled in the OMS646 Phase 2 Trial.

Results:

Following the four-week dosing period (high-dose cohort) as described in Example 44, the patient with stem cell transplant-related TMA demonstrated:

- Platelet count quadrupled, resulting in a platelet count of more than 100,000;
- Haptoglobin level more than doubled and was normal;
- Plasma lactate dehydrogenase level, a measure of damage within blood vessels, decreased by 35% but was still above normal;
- Shistocyte count remained at only one.

Throughout dosing with OMS646 and since completing OMS646 treatment, the patient has not required any platelet transfusions or plasmapheresis.

In summary, the data obtained so far from this Phase 2 clinical trial (as described in Example 44 and in this Example) show efficacy of OMS646 in patients with primary aHUS, plasma-therapy resistant aHUS, TTP and in a patient with TMA associated with hematopoietic stem cell transplant.

In accordance with the foregoing, in one embodiment, the invention provides a method of treating a human subject suffering from a TMA associated with hematopoietic stem cell transplant comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation. In one embodiment, the subject is suffering from a TMA associated with hematopoietic stem cell transplant that is resistant to treatment with platelet transfusions and/or plasmapheresis. In one embodiment, the method further comprises identifying a human subject suffering from a TMA associated with hematopoietic stem cell transplant that is resistant to treatment with platelet transfusions and/or plasmapheresis prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation.

In accordance with any of the disclosed embodiments herein, the MASP-2 inhibitory antibody exhibits at least one or more of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or less, said antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less, wherein the antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway (i.e., inhibits the lectin pathway while leaving the classical complement pathway intact).

In one embodiment, the MASP-2 inhibitory antibody is administered in an amount effective to improve at least one or more clinical parameters associated with TMA associated with hematopoietic stem cell transplant, such as an increase in platelet count (e.g., at least double, at least triple, at least quadruple the platelet count prior to treatment), an increase in haptoglobin, and/or a decrease in lactate dehydrogenase.

In some embodiments, the method comprises administering a MASP-2 inhibitory antibody to a subject suffering from TMA associated with hematopoietic stem cell transplant via a catheter (e.g., intravenously) for a first time period (e.g., at least one day to a week or two weeks) followed by administering a MASP-2 inhibitory antibody to the subject subcutaneously for a second time period (e.g., a chronic phase of at least two weeks or longer). In some embodiments, the administration in the first and/or second time period occurs in the absence of plasma therapy. In some embodiments, the administration in the first and/or second time period occurs in the presence of plasma therapy.

In some embodiments, the method comprises administering a MASP-2 inhibitory antibody to a subject suffering from TMA associated with hematopoietic stem cell transplant either intravenously, intramuscularly, or preferably, subcutaneously. Treatment may be chronic and administered daily to monthly, but preferably every two

weeks. The MASP-2 inhibitory antibody may be administered alone, or in combination with a C5 inhibitor, such as eculizumab.

In one embodiment, the method comprises treating a subject suffering from TMA associated with hematopoietic stem cell transplant comprising administering to the
5 subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising (I) (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence
10 from 95-107 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at
15 least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70.

20 In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67 and a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70.

25 In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof, that specifically recognizes at least part of an epitope on human MASP-2 recognized by reference antibody OMS646 comprising a heavy-chain variable region as set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID
30 NO:70.

EXAMPLE 46

This Example describes additional results obtained in the ongoing Phase 2 Clinical trial to evaluate the Safety and Clinical Efficacy of a fully human monoclonal MASP-2 inhibitory antibody in adults with Thrombotic Microangiopathies (TMAs) described in Examples 44 and 45.

5 Background: As described in Example 45, transplantation-associated TMA (TA-TMA) is a devastating syndrome that can occur in transplant patients, such as hematopoietic stem cell transplant (HSCT) recipients. Hematopoietic stem cell transplant-associated TMA (HSCT-TMA) is a life-threatening complication that is triggered by endothelial injury. The kidney is the most commonly affected organ, though
10 HSCT-TMA can be a multi-system disease that also involves the lung, bowel, heart and brain. The occurrence of even mild TMA is associated with long-term renal impairment. Development of post-allogeneic HSCT-associated TMA differs in frequency based on varying diagnostic criteria and conditioning and graft-versus-host disease prophylaxis regimens, with calcineurin inhibitors being the most frequent drugs implicated (Ho VT et
15 al., *Biol Blood Marrow Transplant*, 11(8):571-5, 2005). Modification of the immunosuppressive calcineurin inhibitor regimen may result in improvement of the TMA in some patients within a few weeks of the reduction or discontinuation of calcineurin inhibitor administration.

20 TMA is a potentially life-threatening complication of HSCT and is currently managed largely by amelioration of inciting factors including avoidance of immunosuppressive agents (e.g., calcineurin inhibitors) and treatment of ongoing infections, as well as supportive measures such as hemodialysis. Although many HSCT-TMA patients respond well to reduction or discontinuation of immunosuppressive agents, there is a subset of patients that have persistent HSCT-TMA despite conservative
25 treatment measures (i.e., the TMA did not respond to reduction or discontinuation of immunosuppressives). Patients who do not respond to these conservative treatment measures have poor prognosis. Plasma exchange has not shown efficacy and no other therapy is approved.

30 Example 45 describes the initial positive results of the Phase 2 Clinical trial to evaluate the safety and clinical efficacy of a fully human monoclonal MASP-2 inhibitory antibody (OMS646) in a patient suffering from persistent hematopoietic stem cell transplant-related TMA (HSCT-TMA). This Example describes additional results of the ongoing Phase 2 Clinical trial to evaluate the safety and clinical efficacy of OMS646 in

three additional subjects suffering from persistent HSCT-TMA resistant to conservative treatment measures.

Methods:

As described in Example 44, the Phase 2 TMA trial consists of a three-level dose-ranging stage, followed by a fixed-dose stage of the MASP-2 inhibitory antibody OMS646. OMS646 was well tolerated by all patients in the low, mid and high-dose cohorts throughout the treatment period. Preclinical toxicity studies have been completed and demonstrated no safety concerns, allowing chronic dosing in clinical trials.

The Phase 2 TMA trial includes subjects suffering from persistent HSCT-associated TMA resistant to conservative treatment measures, which is defined, for the purposes of this study, as having all of the following at least two weeks following reduction or discontinuation of immunosuppression agent (e.g., calcineurin inhibitor treatment) or at least 30 days after transplant:

- Thrombocytopenia (Platelet count < 150,000/ μ L); and
- Evidence of microangiopathic hemolytic anemia (presence of schistocytes, serum lactate dehydrogenase (LDH) > upper limit of normal (ULN), or haptoglobin < lower limit of normal (LLN).

Allowed Concomitant Therapies for HSCT-associated TMA

- Plasma therapy during OMS646 treatment is allowed if the investigator considers it medically indicated. Patients given plasma therapy could receive additional half doses of OMS646.
- Investigators were advised that renal dialysis therapy should be managed according to standard of care.
- Eculizumab was not administered during the study.

The ongoing TMA study includes subjects suffering from persistent HSCT-TMA that is resistant to standard treatment measures (i.e., persistent TMA at least two weeks after reduction or discontinuation of calcineurin inhibitor treatment).

As described in Example 45, dosing was completed for a patient suffering from persistent HSCT-TMA (patient #1) in the high dose cohort (4 mg/kg OMS646 administered IV once a week for four weeks) using the methods described in Example 44.

Dosing has now been completed for an additional 4 patients suffering from persistent HSCT-TMA, as described below.

Patient #1 (described in Example 44) was treated for four weeks with OMS646 (4mg/kg IV once weekly);

5 Patients #2 and #3 were treated for eight weeks with OMS646 (4mg/kg IV once weekly);

Patients #4 and #5 were treated with OMS646 (4mg/kg IV once weekly) for two and three weeks, respectively. Patients #4 and #5 withdrew from the study after two and three weeks, respectively, did not respond to treatment and deteriorated. It is noted that
10 one of these patients had markedly elevated creatinine at the time of study admission.

Results:

Five patients with persistent HSCT-TMA were enrolled in this study. All subjects were adults and received HSCT for hematological malignancies. FIGURES 58, 59 and
15 60 provide the change from baseline in TMA variables after treatment with MASP-2 inhibitory antibody OMS646. These figures provide data on all five patients, two of whom discontinued the study after 2-3 weeks, with one subsequently relapsing and the other receiving palliative care. As noted in FIGURES 58, 59 and 60, the last possible treatment given occurred at week 7.

20 FIGURE 58 graphically illustrates the mean change in platelet count from baseline over time (weeks) in subjects suffering from persistent hematopoietic stem cell transplant-associated thrombotic microangiopathy (HSCT-TMA) after treatment with MASP-2 inhibitory antibody (OMS646).

FIGURE 59 graphically illustrates the mean change in LDH from baseline over
25 time (weeks) in subjects suffering from persistent (HSCT-TMA) after treatment with MASP-2 inhibitory antibody (OMS646).

FIGURE 60 graphically illustrates the mean change in haptoglobin from baseline over time (weeks) in subjects suffering from persistent (HSCT-TMA) after treatment with MASP-2 inhibitory antibody (OMS646).

30 As shown in FIGURES 59 and 60, statistically significant improvement in LDH and haptoglobin were observed during treatment. As shown in FIGURE 58, platelet count improved but did not reach statistical significance in this small number of patients. Of the three patients who completed treatment, one did not show improvement in

creatinine but was receiving concomitant nephrotoxic agents. Creatinine improved or remained normal in the other two patients. On extended follow up, one patient experienced graft failure and is awaiting a second transplant. The other two patients remain stable.

5 Despite potentially confounding effects of medications associated with bone marrow suppression and nephrotoxicity in two of the subjects, beneficial treatment effects with OMS646 were observed in three of the subjects suffering from persistent HSCT-TMA (one of which is described in Example 45) and further described below. Notably, treatment effects in each of the three responders were initially seen after approximately
10 three weeks of treatment with OMS646.

 Patient #1 (treated for 4 weeks as described in Example 45). Briefly summarized, the platelet count quadrupled, resulting in a platelet count of more than 100,000/ μ L; the haptoglobin level more than doubled and was normal; the plasma lactate dehydrogenase level decreased by 35% but was still above normal; and the schistocyte count remained at
15 only one.

 Patient #2 (treated for 8 weeks): the platelet count did not respond to treatment, although the patient also suffered from marrow suppression secondary to concurrent treatment with valganciclovir and later graft failure; the plasma lactate dehydrogenase level decreased from 712 U/L to as low as 256 U/L; the haptoglobin level increased from
20 undetectable levels to as high as 250 mg/dL.

 Patient #3 (treated for 8 weeks): the platelet count increased from 13,500/ μ L to 133,000/ μ L; the plasma lactate dehydrogenase level decreased from 537 to 225 U/L, the haptoglobin level increased from undetectable levels to as high as 181 mg/dL.

Summary of Results:

25 For the three patients suffering from persistent HSCT-associated TMA who completed dosing with OMS646, the data demonstrate a strong and consistent efficacy signal. Statistically significant improvements in LDH and haptoglobin were observed during treatment. Platelet counts improved but did not reach statistical significance in this small number of patients. Of the three patients who completed treatment, one did not
30 show an improvement in creatinine but was receiving concomitant nephrotoxic agents. Creatinine improved or remained normal in the other two patients.

Conclusions:

OMS646 improved TMA markers in patients suffering from persistent HSCT-TMA who had not responded to conservative treatment measures. The HSCT-TMA patients treated with OMS646 represent some of the most difficult to treat, thereby demonstrating clinical evidence of a therapeutic effect of OMS646 in patients with high-risk persistent HSCT-TMA despite conservative treatment measures.

In accordance with the foregoing, in one embodiment, the invention provides a method of treating a human subject suffering from a TMA associated with hematopoietic stem cell transplant comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation. In one embodiment, the subject is suffering from persistent TMA associated with hematopoietic stem cell transplant that is resistant to conservative treatment measures. In one embodiment, the method further comprises identifying a human subject suffering from persistent TMA associated with hematopoietic stem cell transplant that is resistant to conservative treatment measures prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation.

In accordance with any of the disclosed embodiments herein, the MASP-2 inhibitory antibody exhibits at least one or more of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or less, said antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less, wherein the antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway (i.e., inhibits the lectin pathway while leaving the classical complement pathway intact).

In one embodiment, the MASP-2 inhibitory antibody is administered in an amount effective to improve at least one or more clinical parameters associated with TMA associated with hematopoietic stem cell transplant, such as an increase in platelet

count (e.g., at least double, at least triple, at least quadruple the platelet count prior to treatment), an increase in haptoglobin, and/or a decrease in lactate dehydrogenase.

In some embodiments, the method comprises administering a MASP-2 inhibitory antibody to a subject suffering from TMA associated with hematopoietic stem cell transplant via a catheter (e.g., intravenously) for a first time period (e.g., at least one day to a week or two weeks or three weeks or four weeks or longer) followed by administering a MASP-2 inhibitory antibody to the subject subcutaneously for a second time period (e.g., a chronic phase of at least two weeks or longer). In some embodiments, the administration in the first and/or second time period occurs in the absence of plasma therapy. In some embodiments, the administration in the first and/or second time period occurs in the presence of plasma therapy.

In some embodiments, the method comprises administering a MASP-2 inhibitory antibody to a subject suffering from TMA associated with hematopoietic stem cell transplant either intravenously, intramuscularly, or subcutaneously. Treatment may be chronic and administered daily to monthly, but preferably every at least every two weeks, or at least once a week, such as twice a week or three times a week. The MASP-2 inhibitory antibody may be administered alone, or in combination with a C5 inhibitor, such as eculizumab.

In one embodiment, the method comprises treating a subject suffering from persistent TMA associated with hematopoietic stem cell transplant comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy chain variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid sequence set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ ID NO:70. In some embodiments, the composition comprises a MASP-2 inhibitory antibody comprising (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-107 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the

amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with
5 at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid
10 sequence set forth as SEQ ID NO:67 and a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof, that specifically recognizes at least part of an epitope on human MASP-2
15 recognized by reference antibody OMS646 comprising a heavy-chain variable region as set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70.

In some embodiments, the method comprises administering to a subject suffering from, or at risk for developing TMA associated with HSCT, including a subject suffering
20 from persistent TMA associated with hematopoietic stem cell transplant that is resistant to conservative treatment measures, a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67 and a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70 in a
25 dosage from 1 mg/kg to 10 mg/kg (i.e., 1 mg/kg, 2 mg/kg, 3 mg/kg, 4 mg/kg, 5 mg/kg, 6 mg/kg, 7 mg/kg, 8 mg/kg, 9 mg/kg or 10 mg/kg) at least once weekly (such as at least twice weekly or at least three times weekly) for a period of at least 3 weeks, or for at least 4 weeks, or for at least 5 weeks, or for at least 6 weeks, or for at least 7 weeks, or for at least 8 weeks.

30

EXAMPLE 47

This Example describes a case study demonstrating effective treatment of Graft-versus-Host Disease (GVHD) associated microangiopathy with a MASP-2 inhibitory antibody, OMS646.

Background: Graft-versus-host disease is a common complication of hematopoietic stem cell transplant (HSCT). Both acute and chronic forms exist and result from donor immune cells recognizing the recipient patient as foreign tissue. This triggers an immune response against the recipient patient. Acute GvHD occurs in up to 50% or more of patients who receive allogeneic transplants. Acute GvHD most commonly targets the skin, gastrointestinal tract, and liver, but can also affect the kidney, eye, lung and blood cells. Chronic GvHD occurs in approximately 40% of patients who receive allogeneic transplants and most commonly affects the skin, liver, eye, gastrointestinal tract and lungs. Both acute and chronic GvHD are related to significant morbidity and mortality.

Methods and Results

A 46-year-old man affected by T-cell acute lymphoblastic leukemia (T-ALL) in second complete remission (CR) underwent a full myeloablative conditioning with a cyclophosphamide and total body irradiation regimen (TBI-Cy) and peripheral blood stem cell (PBSC) allogeneic HSCT transplantation from an unrelated 35-year-old female donor 9/10 HLA compatible (antigenic mismatch at locus A). Graft-versus-host disease (GVHD) prophylaxis was based on antithymocyte globulin (ATG, 5 mg/kg) during conditioning, Cyclosporin A (CSA) and short course of Methotrexate (i.e., Methotrexate on day +1, +3, +6 (dose reduction due to low CD34+)). He achieved a robust hematologic reconstitution (engraftment: WBC >1000/mm³, neutrophils >500/mm³ on day +15; platelets >20000/mm³ on day +18, 50000/mm³ on day +21). Chimerism was observed with full donor (bone marrow and peripheral blood, CD3+ and CD3- fraction) on day +30, +60, +90, +180, +354. Complete remission was confirmed (immunophenotype, molecular MRD) on day +30, +60, +90, +180.

At day +35 post-transplant, he developed gastrointestinal (GI) GVHD (presented with emesis, anorexia, diarrhea, abdominal pain; colonoscopy: diffuse erosions; histology: diffuse mucosal ulcerations, lymphogranulocytic inflammation in the lamina propria, gland and crypt distortion, apoptotic bodies), and was diagnosed with acute GvHD, stage 4 (gut), overall grade 4 along with cytomegalovirus (CMV) colitis (determined by positive immunohistochemistry and real-time PCR on gut biopsy; systemic CMV

reactivation (2614 UI/mL on peripheral blood). The two conditions were simultaneously treated with methylprednisolone (1 mg/Kg), foscarnet and ganciclovir with good response and steroid tapered to withdrawal.

At day +121 post-transplant he experienced relapse of steroid refractory GI GVHD (presented with emesis, anorexia, diarrhea, abdominal pain; histological findings (stomach and duodenum), and was diagnosed with late onset acute GI GvHD stage 4 (gut), overall grade 4, which was treated, with good response, by continuing CSA and with a sequential administration of pentostatin and mesenchymal stem cells according to a registered clinical study (ClinicalTrials.Gov Identifier NCT02032446).

At day +210 post-transplant the patient presented with weight loss, marked asthenia, generalized muscular atrophy, tetraparesis, reduced deep tendon reflexes, bilateral calf paresthesia, neurogenic bladder and urinary incontinence in absence of minctional (i.e., micturitional) stimulus. Clinical examination and cerebrospinal fluid, neuroradiological and electrophysiological findings described a picture of axonal, sensorimotor and dysautonomic polyneuropathy. Neuroradiology indicated a normal MRI (brain and spine) and cerebrospinal fluid was normal. Relapse of T-ALL, infections, vascular damage, nutritional deficiencies, demyelinating disorder and posterior reversible encephalopathy syndrome (PRES) were excluded. High dose immunoglobulin was administered with no benefit. A concomitant thrombocytopenia ($6 \times 10^9/L$) with features of macroangiopathic hemolytic anemia (6.7 g/dL) without renal impairment, reticulocytosis ($335 \times 10^9/L$), increased LDH (1635 U/L) and undetectable haptoglobin were also documented. A direct antiglobulin test was negative. Schistocytes were observed on peripheral blood smear (2-3/field 50x), as well as absence of anti-B abs (major incompatibility patient/donor), normal ADAMTS13 activity activity, absence of anti-ADAMTS13 Abs, normal coagulation tests and proteinuria with normal creatinine values. He also presented melena with documented colic ulcerations on endoscopic examination; the histopathological study revealed mucosal edema, fibrosis, ectatic small vessels with intraluminal fibrin deposition, glandular atrophy and apoptotic bodies with no microbiological findings. Thus, although there was evidence of persistent GI GVHD, the clinical and histopathological picture was also consistent with a diagnosis of colic transplant-associated thrombotic microangiopathy. Because of the recurrent GI GVHD, CSA was tapered but not withdrawn (trough level 100-150 ng/mL), without obtaining clinical or hematological changes.

At day +263 post-transplant the patient was enrolled into the clinical trial described above in Examples 45 and 46, which involves treatment with OMS646, a MASP-2 inhibitory antibody that inhibits the lectin pathway of complement activation. As shown in FIGURE 61, after the first 2 weekly doses of the drug a clinical and laboratory response was observed with resolution of melena and hemolysis and rise of platelet count. After 4 doses, interestingly, a remarkable neurological improvement was also documented with both clinical and electrophysiological improvement of the sensorimotor deficit. After 8 doses of OMS646, the GvHD response was maintained despite CSA tapering, with no toxicities observed. By day +354 post-transplant, the patient demonstrated further neurological improvement and initial recovery of micturitional (i.e., micturitional) deficit. The rapid clinical benefit achieved by this patient suggests that effective inhibition of the complement-mediated endothelial damage, resulting from the administration of OMS646, can be highly effective for the treatment of GVHD associated TMA.

In summary, this Example describes a case report of a HSCT patient post-transplant having co-existing HSCT-TMA and GvHD, which both resolved following treatment with a MASP-2 inhibitory antibody (OMS646). Prior to treatment with OMS646 the patient had a difficult post-transplant course complicated by multiple episodes of steroid-refractory GvHD, cytomegalovirus infection and HSCT-TMA. After two prior episodes of GvHD, the patient presented with bloody diarrhea. Intestinal biopsy demonstrated both HSCT-TMA and GvHD. No infections were identified. Notably, the patient also had new onset neurological symptoms of paresthesias, tetraplegia and a neurogenic bladder, which have been reported as neurological manifestations of GvHD and TMA. The patient was unable to walk due to the tetraplegia and required blood transfusions at least once daily. Hematological markers demonstrated HSCT-TMA with thrombocytopenia, elevated lactate dehydrogenase (LDH) and schistocytes. The patient entered the clinical trial and began receiving OMS646. His immunosuppression (cyclosporine) had been decreased 2 weeks earlier and he received only low dose corticosteroids in view of his history of steroid-refractory GvHD. He received no other GvHD treatment. His bloody diarrhea resolved and his hematological markers improved after 2 OMS646 doses. After 4 OMS646 doses he was able to walk. He completed 8 weeks of OMS646 treatment and is doing well at home. All signs and symptoms of HSCT-TMA and GvHD have resolved. His neurological symptoms

continue to improve. Prior to OMS646 treatment this patient was deteriorating and at high risk for early death. Following treatment with OMS646, improvement of GvHD, HSCT-TMA and the neurological symptoms were observed and the patient is doing well at home.

5 In accordance with the foregoing, in one embodiment, the invention provides a method of treating a human subject suffering from, or at risk for developing graft-versus-host-disease (GVHD) comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation and thereby treat, or reduce the risk of developing GVHD, or
10 reduce the severity of one or more symptoms associated with GVHD. In one embodiment, the GVHD is active, acute GVHD. In one embodiment, the GVHD is chronic GVHD. In one embodiment, the GVHD is steroid-resistant (i.e., persistent despite steroid treatment). In one embodiment, the GVHD gastrointestinal (GI) GVHD. In one embodiment, the method further comprises the step of determining the presence of
15 GVHD in a subject prior to treatment with the MASP-2 inhibitory antibody.

In one embodiment, the subject is suffering from, or at risk for developing GVHD associated with a hematopoietic stem cell transplant. In one embodiment, the subject is suffering from, or at risk for developing GVHD associated with a TMA associated with hematopoietic stem cell transplant. In one embodiment, the subject is suffering from
20 leukemia, such as T-cell acute lymphoblastic leukemia.

In one embodiment, the subject is suffering from one or more neurological symptoms associated with GvHD and/or TMA, such as, for example, asthenia, paresthesias, tetraplegia, sensorimotor deficit, dysautonomic polyneuropathy, and/or neurogenic bladder and the MASP-2 inhibitory antibody is administered in an amount
25 and for a time period sufficient to ameliorate one or more of the neurological symptoms.

In accordance with any of the disclosed embodiments herein, the MASP-2 inhibitory antibody exhibits at least one or more of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or less, said antibody inhibits C3b
30 deposition in 90% human serum with an IC_{50} of 30 nM or less, wherein the antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and

F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway (i.e., inhibits the lectin pathway while leaving the classical complement pathway intact).

In one embodiment, the MASP-2 inhibitory antibody is administered in an amount effective to improve at least one or more clinical parameters associated with GVHD.

In some embodiments, the method comprises administering a MASP-2 inhibitory antibody to a subject suffering from GVHD associated with hematopoietic stem cell transplant either intravenously, intramuscularly, or subcutaneously. Treatment may be chronic and administered daily to monthly, but preferably every at least every two weeks, or at least once a week, such as twice a week or three times a week. The MASP-2 inhibitory antibody may be administered alone, or in combination with a C5 inhibitor, such as eculizumab.

In one embodiment, the method comprises treating a subject suffering from GVHD associated with hematopoietic stem cell transplant comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy chain variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid sequence set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ ID NO:70. In some embodiments, the composition comprises a MASP-2 inhibitory antibody comprising (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-107 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at

least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70).

5 In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67 and a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70.

10 In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof, that specifically recognizes at least part of an epitope on human MASP-2 recognized by reference antibody OMS646 comprising a heavy-chain variable region as set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID
15 NO:70.

 In some embodiments, the method comprises administering to a subject suffering from, or at risk for developing GVHD, such as a subject that will undergo, is undergoing, or has undergone HSCT, including a subject suffering from GVHD associated TMA, a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment
20 thereof comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67 and a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70 in a dosage from 1 mg/kg to 10 mg/kg (i.e., 1 mg/kg, 2 mg/kg, 3 mg/kg, 4 mg/kg, 5 mg/kg, 6 mg/kg, 7 mg/kg, 8 mg/kg, 9 mg/kg or 10 mg/kg) at least once weekly (such as at least twice weekly or at least three times weekly)
25 for a period of at least 3 weeks, or for at least 4 weeks, or for at least 5 weeks, or for at least 6 weeks, or for at least 7 weeks, or for at least 8 weeks.

30

EXAMPLE 48

This Example describes a case study demonstrating effective treatment of TMA and diffuse alveolar hemorrhage (DAH) after hematopoietic stem cell transplant with a MASP-2 inhibitory antibody, OMS646.

Background: Hematopoietic stem cell transplantation (HSCT) induces substantial
5 endothelial injury, which contributes to serious post-HSCT complications such as
thrombotic microangiopathy (TMA), graft-versus-host disease (GvHD), diffuse alveolar
hemorrhage (DAH) and veno-occlusive disease (VOD). (see E. Carreras and M. Diaz-
Ricart, *Bone Marrow Transplant* vol 46:1495-1502, 2011; Akil et al., *Biol Blood Marrow*
Transplant 21:1739-1745, 2015; and Vion et al., *Semin Thromb Hemost* 41(06):629-643,
10 2015, each of which is hereby incorporated herein by reference).

Diffuse alveolar hemorrhage (DAH) is a syndrome that can occur in a hematopoietic stem
cell transplant (HSCT) recipient. The diagnostic criteria of DAH in the HSCT recipient
include lung infiltrates, dyspnea and hypoxia (see E. Carreras and M. Diaz-Ricart, *Bone*
Marrow Transplant vol 46:1495-1502, 2011, hereby incorporated herein by reference).
15 The suspected pathogenesis of DAH is lung capillary endothelium damages by HSCT
conditioning plus engrafted neutrophils and silent infections, allowing the leakage of red
blood cells into the pulmonary alveoli (E. Carreras and M. Diaz-Ricart, *Bone Marrow*
Transplant vol 46:1495-1502, 2011). Veno-occlusive disease (VOD), also known as
sinusoidal obstructive syndrome (SOS), is another syndrome that can occur in a
20 hematopoietic stem cell transplant (HSCT) patient. The clinical manifestations of VOD
include marked weight gain due to fluid retention, increased liver size and raised levels of
bilirubin in the blood (see E. Carreras and M. Diaz-Ricart, *Bone Marrow Transplant* vol
46:1495-1502, 2011, hereby incorporated herein by reference).

Methods and Results:

25

Demographics and Past Medical History

The patient was a 14 year-old white female at the time of treatment initiation,
referred to in this Example as “compassionate use patient #1.” The patient had a history
of Diamond-Blackfan anemia and empty sella syndrome.

30 Patient Outcome

The patient is alive on Day 877.

Transplant Requiring Disease

The patient has a history of Diamond-Blackfan anemia. She had required RBC infusions every 3 weeks since infancy and previously developed iron overload.

Transplant

5 The patient underwent her stem cell transplant in September 2015. She received a PBSC transplant from a 9/10 matched unrelated donor. Neutrophil engraftment occurred by approximately Day 35. Platelet engraftment had not been achieved by Day 184. She required platelet and RBC infusions for more than 1 year.

10 Post-transplant Complications

 The patient had a complicated post-transplant course. Shortly after discharge she was readmitted on Day 67 for shortness of breath and presumed pneumonia. She was discharged on Day 162. She was readmitted on Day 167 with varicella zoster reactivation treated with acyclovir and polyserositis treated with corticosteroids. She had persistent
15 thrombocytopenia and anemia, elevated LDH, decreased haptoglobin, schistocytes and a negative direct antibody test. A diagnosis of TMA was made. Immunosuppression was changed to MMF and prednisone. She also experienced HHV6 and EBV reactivation. She was discharged on Day 197.

 On Day 229 she was readmitted with persistent TMA and severe respiratory
20 distress. She was diagnosed with diffuse alveolar hemorrhage that was treated with CPAP and corticosteroids. On Day 231 she had pleural and pericardial effusions with worsening oxygenation. Her blood pressure increased. She required an ACE inhibitor, calcium channel blocker, beta blocker, diuretic, and nitroglycerin for BP control. She experienced a seizure and clonidine was added. Brain MRI showed iron in the brain consistent with
25 secondary hemochromatosis from her RBC transfusion history. Hemodialysis was initiated on Day 231. On Day 259 she began eculizumab treatment due to continued renal dysfunction and worsening pulmonary status. Her TMA improved, but she developed acute pulmonary edema and eculizumab was discontinued. Her TMA worsened and she underwent plasma exchange on Days 269, 276, and 278 without improvement. She was
30 treated with defibrotide that was discontinued due to a coagulopathy and no TMA improvement. She again received lower-dose eculizumab starting on approximately Day 320. The eculizumab was stopped again due to pulmonary edema. The TMA persisted. She also developed C. difficile infection. She was discharged in Day 379 receiving

prednisone, MMF, voriconazole, cotrimazole, amlodipine, erythropoietin, ursodiol and omeprazole.

5 She was readmitted on Day 380 with fever, chills, and dyspnea. She had K. pneumoniae pneumonia and E. faecium sepsis. Her MMF was stopped. She was treated with Tazocin, meropenem, and vancomycin. With persistent bacteremia, her central line was changed and her antibiotics were changed to linezolid, daptomycin, and clindamycin. Her TMA persisted requiring hemodialysis 3 times weekly and daily platelet transfusions. Pulmonary edema diagnosed as diffuse alveolar hemorrhage (treated with corticosteroids without response). On day 404 OMS646 treatment began under a compassionate use
10 protocol. As described herein, OMS646 is a MASP-2 inhibitory antibody that inhibits the lectin pathway of complement activation.

She was treated 3 times weekly with a dose of 4 mg/kg OMS646. Her oxygen was discontinued and her corticosteroid dose was decreased. Her laboratory TMA markers improved. She was discharged on Day 414.

15 She continued outpatient OMS646 treatment 3 times weekly and on Day 416 her hemodialysis was decreased to once weekly. Her antihypertensive medication was also reduced to amlodipine, clonidine and furosemide only. On Day 423 she could discontinue hemodialysis and on Day 428 her OMS646 was decreased to twice weekly with reductions in her corticosteroid and furosemide doses. Her platelet transfusions were
20 substantially decreased during this period.

She continued to do well until Day 486 when she developed an RSV infection. Her creatinine and LDH increased, and her platelet count and haptoglobin fell. She was diagnosed with recurrent TMA and OMS646 treatment was increased to 3 times weekly. Her TMA resolved with the increased OMS646 treatment.

25 She has remained hemodialysis free and her last platelet transfusions were occurred on about Day 630. She is currently doing well and receiving treatment with OMS646, desfuroxamine, levothyroxine, sevelamer, Augmentin, levetiracetam, allopurinol, amlodipine, nebivolol, clonidine and erythropoietin. She is receiving phlebotomies for the hemochromatosis.

30

TMA Episodes

This patient had a difficult course with persisting and/or relapsing TMA. Her course is consistent with multi-organ TMA and she had diffuse alveolar hemorrhage

(DAH), another endothelial injury syndrome. She initially responded to eculizumab, but did not tolerate the treatment. Prior to initiating OMS646 treatment she was on hemodialysis and requiring daily transfusions. Soon after OMS646 treatment her TMA resolved, her DAH resolved, and she was able to discontinue dialysis. She was later able to substantially reduce and later discontinue platelet and RBC transfusions while maintaining stable or increasing platelet count and hemoglobin values. These results are demonstrated in FIGURES 62A-62E as follows:

FIGURE 62A graphically illustrates the level of creatinine over time in the compassionate use patient #1, wherein the vertical line indicates the start of treatment with MASP-2 inhibitory antibody (OMS646).

FIGURE 62B graphically illustrates the level of haptoglobin over time in the compassionate use patient #1, wherein the vertical line indicates the start of treatment with MASP-2 inhibitory antibody (OMS646).

FIGURE 62C graphically illustrates the level of hemoglobin over time in the compassionate use patient #1, wherein the vertical line indicates the start of treatment with MASP-2 inhibitory antibody (OMS646).

FIGURE 62D graphically illustrates the level of LDH over time in the compassionate use patient #1, wherein the vertical line indicates the start of treatment with MASP-2 inhibitory antibody (OMS646).

FIGURE 62E graphically illustrates the level of platelets over time in the compassionate use patient #1, wherein the vertical line indicates the start of treatment with MASP-2 inhibitory antibody (OMS646).

Overall, this patient's course demonstrates successful OMS646 treatment of TMA and DAH allowing the patient to discontinue oxygen therapy, hemodialysis, and transfusions.

Summary

In summary, this Example describes an HCST-TMA case report of an adolescent girl (compassionate use patient #1) who did not tolerate eculizumab treatment, but responded well to compassionate use of OMS646 treatment. Soon after OMS646 treatment her TMA resolved, her DAH resolved, and she was able to discontinue dialysis. As described in Example 47, another HSCT-TMA case report was provided describing a patient who had a difficult post-transplant course, including steroid-resistant GvHD and cytomegalovirus infection. He developed TMA that did not respond to conservative

measures and had co-existing GvHD with multiple neurological complications and was unable to walk. As described in Example 47, following OMS646 treatment, his TMA and GvHD resolved and his neurological complications improved. He was able to return to work and his neurological status has continued to improve.

5 The improvement in overall survival and TMA markers in HSCT recipients combined with resolution of GvHD and resolution of diffuse alveolar hemorrhage in these critically ill patients indicates the role the lectin pathway plays in these syndromes and the efficacy for use of a MASP-2 inhibitory antibody OMS646 in treating and/or preventing post-stem cell transplant TMA, graft-versus-host disease and diffuse alveolar
10 hemorrhage.

 In accordance with the foregoing, in one embodiment, the invention provides a method of treating a human subject suffering from diffuse alveolar hemorrhage associated with hematopoietic stem cell transplant (i.e., in a HSCT recipient) comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory
15 antibody effective to inhibit MASP-2-dependent complement activation. In one embodiment, the method further comprises identifying a human subject suffering from diffuse alveolar hemorrhage associated with HSCT prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation.

20 In accordance with any of the disclosed embodiments herein, the MASP-2 inhibitory antibody exhibits at least one or more of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or less, said antibody inhibits C3b
25 deposition in 90% human serum with an IC_{50} of 30 nM or less, wherein the antibody is an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation. In one embodiment,
30 the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway (i.e., inhibits the lectin pathway while leaving the classical complement pathway intact).

In one embodiment, the MASP-2 inhibitory antibody is administered in an amount effective to improve at least one or more clinical parameters associated with diffuse alveolar hemorrhage in a HSCT recipient. In some embodiments, the method comprises administering a MASP-2 inhibitory antibody to a subject suffering from
5 diffuse alveolar hemorrhage either intravenously, intramuscularly, or subcutaneously. Treatment may be chronic and administered daily to monthly, but preferably every at least every two weeks, or at least once a week, such as twice a week or three times a week. The MASP-2 inhibitory antibody may be administered alone, or in combination with a C5 inhibitor, such as eculizumab.

10 In one embodiment, the method comprises treating a subject suffering from diffuse alveolar hemorrhage associated with hematopoietic stem cell transplant comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy chain variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid sequence
15 set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ ID NO:70. In some embodiments, the composition comprises a MASP-2 inhibitory antibody comprising (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising
20 the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-107 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the
25 amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with
30 at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid

sequence set forth as SEQ ID NO:67 and a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof, that specifically recognizes at least part of an epitope on human MASP-2 recognized by reference antibody OMS646 comprising a heavy-chain variable region as set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70.

In some embodiments, the method comprises administering to a subject suffering from, or at risk for developing diffuse alveolar hemorrhage associated with HSCT, a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67 and a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70 in a dosage from 1 mg/kg to 10 mg/kg (i.e., 1 mg/kg, 2 mg/kg, 3 mg/kg, 4 mg/kg, 5 mg/kg, 6 mg/kg, 7 mg/kg, 8 mg/kg, 9 mg/kg or 10 mg/kg) at least once weekly (such as at least twice weekly or at least three times weekly) for a period of at least 3 weeks, or for at least 4 weeks, or for at least 5 weeks, or for at least 6 weeks, or for at least 7 weeks, or for at least 8 weeks.

In accordance with the foregoing, in another embodiment, the invention provides a method of treating a human subject suffering from veno-occlusive disease (VOD) associated with hematopoietic stem cell transplant (i.e., in a HSCT recipient) comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation. In one embodiment, the method further comprises identifying a human subject suffering from veno-occlusive disease associated with HSCT prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody effective to inhibit MASP-2-dependent complement activation.

In accordance with any of the disclosed embodiments herein, the MASP-2 inhibitory antibody exhibits at least one or more of the following characteristics: said antibody binds human MASP-2 with a K_D of 10 nM or less, said antibody binds an epitope in the CCP1 domain of MASP-2, said antibody inhibits C3b deposition in an *in vitro* assay in 1% human serum at an IC_{50} of 10 nM or less, said antibody inhibits C3b deposition in 90% human serum with an IC_{50} of 30 nM or less, wherein the antibody is

an antibody fragment selected from the group consisting of Fv, Fab, Fab', F(ab)₂ and F(ab')₂, wherein the antibody is a single-chain molecule, wherein said antibody is an IgG2 molecule, wherein said antibody is an IgG1 molecule, wherein said antibody is an IgG4 molecule, wherein the IgG4 molecule comprises a S228P mutation. In one embodiment, the antibody binds to MASP-2 and selectively inhibits the lectin pathway and does not substantially inhibit the classical pathway (i.e., inhibits the lectin pathway while leaving the classical complement pathway intact).

In one embodiment, the MASP-2 inhibitory antibody is administered in an amount effective to improve at least one or more clinical parameters associated with veno-occlusive disease in a HSCT recipient. In some embodiments, the method comprises administering a MASP-2 inhibitory antibody to a subject suffering from veno-occlusive disease either intravenously, intramuscularly, or subcutaneously. Treatment may be chronic and administered daily to monthly, but preferably every at least every two weeks, or at least once a week, such as twice a week or three times a week. The MASP-2 inhibitory antibody may be administered alone, or in combination with a C5 inhibitor, such as eculizumab.

In one embodiment, the method comprises treating a subject suffering from veno-occlusive disease associated with hematopoietic stem cell transplant comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy chain variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid sequence set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ ID NO:70. In some embodiments, the composition comprises a MASP-2 inhibitory antibody comprising (a) a heavy-chain variable region comprising: i) a heavy-chain CDR-H1 comprising the amino acid sequence from 31-35 of SEQ ID NO:67; and ii) a heavy-chain CDR-H2 comprising the amino acid sequence from 50-65 of SEQ ID NO:67; and iii) a heavy-chain CDR-H3 comprising the amino acid sequence from 95-107 of SEQ ID NO:67 and b) a light-chain variable region comprising: i) a light-chain CDR-L1 comprising the amino acid sequence from 24-34 of SEQ ID NO:70; and ii) a light-chain CDR-L2 comprising the amino acid sequence from 50-56 of SEQ ID NO:70; and iii) a light-chain CDR-L3 comprising the amino acid sequence from 89-97 of SEQ ID NO:70, or (II) a variant thereof comprising a heavy-chain variable region with at least 90% identity to SEQ ID NO:67 (e.g., at least

91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:67) and a light-chain variable region with at least 90% identity (e.g., at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identity to SEQ ID NO:70.

5 In some embodiments, the method comprises administering to the subject suffering from veno-occlusive disease associated with HSCT a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen binding fragment thereof, comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67 and a light-chain variable region comprising the amino acid sequence set
10 forth as SEQ ID NO:70.

In some embodiments, the method comprises administering to the subject a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof, that specifically recognizes at least part of an epitope on human MASP-2 recognized by reference antibody OMS646 comprising a heavy-chain variable region as
15 set forth in SEQ ID NO:67 and a light-chain variable region as set forth in SEQ ID NO:70.

In some embodiments, the method comprises administering to a subject suffering from, or at risk for developing veno-occlusive disease associated with HSCT, a composition comprising a MASP-2 inhibitory antibody, or antigen binding fragment thereof
20 comprising a heavy-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:67 and a light-chain variable region comprising the amino acid sequence set forth as SEQ ID NO:70 in a dosage from 1 mg/kg to 10 mg/kg (i.e., 1 mg/kg, 2 mg/kg, 3 mg/kg, 4 mg/kg, 5 mg/kg, 6 mg/kg, 7 mg/kg, 8 mg/kg, 9 mg/kg or 10 mg/kg) at least once weekly (such as at least twice weekly or at least three times weekly) for a period of
25 at least 3 weeks, or for at least 4 weeks, or for at least 5 weeks, or for at least 6 weeks, or for at least 7 weeks, or for at least 8 weeks.

While illustrative embodiments have been illustrated and described, it will be appreciated that various changes can be made therein without departing from the spirit
30 and scope of the invention.

CLAIMS

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A method of treating a human subject suffering from, or at risk for developing
5 graft-versus-host disease (GVHD) comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation.
2. The method of Claim 1, wherein the MASP-2 inhibitory antibody is a monoclonal antibody, or fragment thereof that specifically binds to human MASP-2.
- 10 3. The method of Claim 1, wherein the antibody or fragment thereof is selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody, and a human antibody.
4. The method of Claim 1, wherein the MASP-2 inhibitory antibody does not substantially inhibit the classical pathway.
- 15 5. The method of Claim 1, wherein the MASP-2 inhibitory antibody inhibits C3b deposition in 90% human serum with an IC₅₀ of 30 nM or less.
6. The method of Claim 1, wherein the MASP-2 inhibitory antibody is delivered to the subject systemically.
7. The method of Claim 1, wherein the method further comprises identifying a
20 human subject suffering from, or at risk for developing graft-versus-host disease prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation.
8. The method of Claim 1, wherein the subject has previously undergone, or is
25 currently undergoing, or will undergo a hematopoietic stem cell transplant.
9. The method of claim 1, wherein the subject is suffering from acute GVHD.
10. The method of claim 1, wherein the subject is suffering from chronic GVHD.

11. The method of claim 1, wherein the subject is suffering from steroid-resistant GVHD.
12. The method of Claim 1, wherein the MASP-2 inhibitory antibody or antigen-binding fragment thereof, comprises a heavy chain variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid sequence set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ ID NO:70.
13. The method of any of claims 1-12, wherein the method comprises administering to the subject a composition comprising said MASP-2 inhibitory antibody in a dosage of from 1 mg/kg to 10 mg/kg at least once weekly.
14. A method of treating, preventing or ameliorating one or more neurological symptoms associated with graft-versus-host disease or HSCT-TMA comprising administering to a subject prior to, during, or after receiving a hematopoietic stem cell transplant a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation.
15. The method of claim 14, wherein the one or more neurological symptoms associated with graft-versus-host-disease or HSCT-TMA is selected from the group consisting of asthenia, paresthesias, tetraplegia, sensorimotor deficit, dysautonomic polyneuropathy, and/or neurogenic bladder.
16. The method of claim 14, wherein the subject has received a hematopoietic stem cell transplant and the subject is suffering from one or more neurological symptoms selected from the group consisting of asthenia, paresthesias, tetraplegia, sensorimotor deficit, dysautonomic polyneuropathy, and/or neurogenic bladder.
17. The method of any of claims 14 to 16, wherein the subject has received a hematopoietic stem cell transplant and is suffering from graft-versus-host disease.
18. The method of any of claims 14 to 17, wherein the subject has received a hematopoietic stem cell transplant and is suffering from HSCT-TMA.

19. The method of Claim 14, wherein the MASP-2 inhibitory antibody is a monoclonal antibody, or fragment thereof that specifically binds to human MASP-2.
- 5 20. The method of Claim 14, wherein the antibody or fragment thereof is selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody, and a human antibody.
21. The method of Claim 14, wherein the MASP-2 inhibitory antibody does not substantially inhibit the classical pathway.
- 10 22. The method of Claim 14, wherein the MASP-2 inhibitory antibody inhibits C3b deposition in 90% human serum with an IC₅₀ of 30 nM or less.
23. The method of Claim 14, wherein the MASP-2 inhibitory antibody is delivered to the subject systemically.
- 15 24. The method of Claim 14, wherein the method further comprises identifying a human subject suffering from one or more neurological symptoms associated with hematopoietic stem cell transplant prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation.
- 20 25. The method of Claim 14, wherein the MASP-2 inhibitory antibody or antigen-binding fragment thereof, comprises a heavy chain variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid sequence set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ ID NO:70.
- 25 26. A method of treating a human subject suffering from, or at risk for developing diffuse alveolar hemorrhage comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation.

27. The method of Claim 26, wherein the MASP-2 inhibitory antibody is a monoclonal antibody, or fragment thereof that specifically binds to human MASP-2.
28. The method of Claim 26, wherein the antibody or fragment thereof is selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody, and a human antibody.
29. The method of Claim 26, wherein the MASP-2 inhibitory antibody does not substantially inhibit the classical pathway.
30. The method of Claim 26, wherein the MASP-2 inhibitory antibody inhibits C3b deposition in 90% human serum with an IC₅₀ of 30 nM or less.
31. The method of Claim 26, wherein the MASP-2 inhibitory antibody is delivered to the subject systemically.
32. The method of Claim 26, wherein the method further comprises identifying a human subject suffering from, or at risk for developing diffuse alveolar hemorrhage prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation.
33. The method of Claim 26, wherein the subject has previously undergone, or is currently undergoing, or will undergo a hematopoietic stem cell transplant.
34. The method of Claim 26, wherein the MASP-2 inhibitory antibody or antigen-binding fragment thereof, comprises a heavy chain variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid sequence set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ ID NO:70.
35. The method of any of claims 26-34, wherein the subject has received a hematopoietic stem cell transplant and is suffering from diffuse alveolar hemorrhage.

36. A method of treating a human subject suffering from, or at risk for developing veno-occlusive disease comprising administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation.

5 37. The method of Claim 36, wherein the MASP-2 inhibitory antibody is a monoclonal antibody, or fragment thereof that specifically binds to human MASP-2.

38. The method of Claim 36, wherein the antibody or fragment thereof is selected from the group consisting of a recombinant antibody, an antibody having reduced effector function, a chimeric antibody, a humanized antibody, and a human antibody.

10 39. The method of Claim 36, wherein the MASP-2 inhibitory antibody does not substantially inhibit the classical pathway.

40. The method of Claim 36, wherein the MASP-2 inhibitory antibody inhibits C3b deposition in 90% human serum with an IC₅₀ of 30 nM or less.

15 41. The method of Claim 36, wherein the MASP-2 inhibitory antibody is delivered to the subject systemically.

42. The method of Claim 36, wherein the method further comprises identifying a human subject suffering from, or at risk for developing veno-occlusive disease prior to the step of administering to the subject a composition comprising an amount of a MASP-2 inhibitory antibody, or antigen-binding fragment thereof, effective to inhibit MASP-2-dependent complement activation.

43. The method of Claim 36, wherein the subject has previously undergone, or is currently undergoing, or will undergo a hematopoietic stem cell transplant.

25 44. The method of Claim 36, wherein the MASP-2 inhibitory antibody or antigen-binding fragment thereof, comprises a heavy chain variable region comprising CDR-H1, CDR-H2 and CDR-H3 of the amino acid sequence set forth as SEQ ID NO:67 and a light chain variable region comprising CDR-L1, CDR-L2 and CDR-L3 of the amino acid sequence set forth as SEQ ID NO:70.

45. The method of any of claims 36-44, wherein the subject has received a hematopoietic stem cell transplant and is suffering from veno-occlusive disease.