

ANTI-PDL1 ANTIBODIES, ACTIVATABLE ANTI-PDL1 ANTIBODIES, AND METHODS OF USE THEREOF

Related Applications

[0001] This application claims the benefit of U.S. Provisional Application No. 62/133,231, filed March 13, 2015; U.S. Provisional Application No. 62/139,596, filed March 27, 2015; and U.S. Provisional Application No. 62/218,883, filed September 15, 2015; the contents of each of which are incorporated herein by reference in their entirety.

Field of the Invention

[0002] The invention relates generally to antibodies that bind programmed death ligand 1 (PDL1), activatable antibodies that specifically bind to PDL1 and methods of making and using these anti-PDL1 antibodies and anti-PDL1 activatable antibodies in a variety of therapeutic, diagnostic and prophylactic indications.

Background of the Invention

[0003] Antibody-based therapies have proven effective treatments for several diseases but in some cases, toxicities due to broad target expression have limited their therapeutic effectiveness. In addition, antibody-based therapeutics have exhibited other limitations such as rapid clearance from the circulation following administration.

[0004] In the realm of small molecule therapeutics, strategies have been developed to provide prodrugs of an active chemical entity. Such prodrugs are administered in a relatively inactive (or significantly less active) form. Once administered, the prodrug is metabolized *in vivo* into the active compound. Such prodrug strategies can provide for increased selectivity of the drug for its intended target and for a reduction of adverse effects.

[0005] Accordingly, there is a continued need in the field of antibody-based therapeutics for antibodies that mimic the desirable characteristics of the small molecule prodrug.

Summary of the Invention

[0006] The disclosure provides antibodies or antigen-binding fragments thereof that specifically bind programmed death ligand 1 (PDL1), also known as PD-L1, CD274, B7 homolog 1 and/or B7-H1. The use of the term “PDL1” is intended to cover any variation

thereof, such as, by way of non-limiting example, PD-L1 and/or PDL-1, all variations are used herein interchangeably.

[0007] In some embodiments, the antibody includes an antibody or antigen-binding fragment thereof that specifically binds PDL1. In some embodiments, the antibody or antigen binding fragment thereof that binds PDL1 is a monoclonal antibody, domain antibody, single chain, Fab fragment, a F(ab')₂ fragment, a scFv, a scAb, a dAb, a single domain heavy chain antibody, or a single domain light chain antibody. In some embodiments, such an antibody or antigen binding fragment thereof that binds PDL1 is a mouse, other rodent, chimeric, humanized or fully human monoclonal antibody.

[0008] In some embodiments, the antibody is an isolated antibody or antigen binding fragment thereof (AB) that specifically binds to mammalian PDL1, wherein the AB has one or more of the following characteristics: (a) the AB specifically binds to human PDL1 and murine PDL1; (b) the AB specifically binds to human PDL1 and cynomolgus monkey PDL1; (c) the AB specifically binds to human PDL1, murine PDL1, and cynomolgus monkey PDL1; (d) the AB inhibits binding of human B7-1 and human PD1 to human PDL1 with an EC₅₀ value less than 10 nM; (e) the AB inhibits binding of murine B7-1 and murine PD1 to murine PDL1 with an EC₅₀ value less than 10 nM; and (f) the AB inhibits binding of cynomolgus monkey B7-1 and cynomolgus monkey PD1 to cynomolgus monkey PDL1 with an EC₅₀ value less than 10 nM.

[0009] In some embodiments, the antibody or antigen binding fragment thereof specifically binds to the mammalian PDL1 with a dissociation constant is less than or equal to 0.01 nM, less than or equal to 0.05 nM, less than or equal to 0.1 nM, less than or equal to 0.2 nM, less than or equal to 0.3 nM, less than or equal to 0.4 nM, less than or equal to 0.5 nM, less than or equal to 0.75 nM, and less than or equal to 1 nM.

[00010] In some embodiments, the antibody or antigen binding fragment thereof specifically binds to the mammalian PDL1 with a dissociation constant in the range of 0.01 nM to 1 nM, 0.05 nM to 1 nM, 0.1 nM to 1 nM, 0.2 nM to 1 nM, 0.3 nM to 1 nM, 0.4 nM to 1 nM, 0.5 nM to 1 nM, 0.75 nM to 1 nM, 0.01 nM to 0.75 nM, 0.05 nM to 0.75 nM, 0.1 nM to 0.75 nM, 0.2 nM to 0.75 nM, 0.3 nM to 0.75 nM, 0.4 nM to 0.75 nM, 0.5 nM to 0.75 nM, 0.01 nM to 0.5 nM, 0.05 nM to 0.5 nM, 0.1 nM to 0.5 nM, 0.2 nM to 0.5 nM, 0.3 nM to 0.5 nM, 0.4 nM to 0.5 nM, 0.01 nM to 0.4 nM, 0.05 nM to 0.4 nM, 0.1 nM to 0.4 nM, 0.2 nM to 0.4 nM, 0.3 nM to 0.4 nM, 0.01 nM to 0.3 nM, 0.05 nM to

0.3 nM, 0.1 nM to 0.3 nM, 0.2 nM to 0.3 nM, 0.01 nM to 0.2 nM, 0.05 nM to 0.2 nM, 0.1 nM to 0.2 nM, 0.01 nM to 0.1 nM, 0.05 nM to 0.1 nM, or 0.01 nM to 0.05 nM.

[00011] In some embodiments, the mammalian PDL1 is selected from the group consisting of a human PDL1, a murine PDL1, a rat PDL1, and a cynomolgus monkey PDL1.

[00012] In some embodiments, the antibody or antigen binding fragment thereof specifically binds to human PDL1, murine PDL1 or cynomolgus monkey PDL1 with a dissociation constant of less than 1 nM.

[00013] In some embodiments, the mammalian PDL1 is a human PDL1.

[00014] In some embodiments, the antibody or antigen binding fragment thereof has one or more of the following characteristics: (a) the AB specifically binds human PDL1, murine PDL1, and cynomolgus monkey PDL1; (b) the AB inhibits binding of human B7-1 and human PD1 to human PDL1; (c) the AB inhibits binding of murine B7-1 and murine PD1 to murine PDL1; and (d) the AB inhibits binding of cynomolgus monkey B7-1 and cynomolgus monkey PD1 to cynomolgus monkey PDL1.

[00015] In some embodiments, the antibody or antigen binding fragment thereof blocks the ability of a natural ligand to bind to the mammalian PDL1 with an EC_{50} less than or equal to 0.5 nM, less than or equal to 1 nM, less than or equal to 2 nM, less than or equal to 3 nM, less than or equal to 4 nM, less than or equal to 5 nM, less than or equal to 6 nM, less than or equal to 7 nM, less than or equal to 8 nM, less than or equal to 9 nM, and/or less than or equal to 10 nM.

[00016] In some embodiments, the antibody or antigen binding fragment thereof blocks the ability of a natural ligand to bind to the mammalian PDL1 with an EC_{50} in the range of 0.5 nM to 10 nM, 0.5 nM to 5 nM, 0.5 nM to 3 nM, 0.5 nM to 2 nM, 0.5 nM to 1 nM, 1 nM to 10 nM, 1 nM to 5 nM, 1 nM to 3 nM, 1 nM to 2 nM, 2 nM to 10 nM, 2 nM to 5 nM, 2 nM to 3 nM, 3 nM to 10 nM, 3 nM to 5 nM, or 5 nM to 10 nM. In some embodiments, the natural ligand is a mammalian PD1. In some embodiments, the natural ligand is selected from the group consisting of: a human PD1, a murine PD1, and a cynomolgus monkey PD1.

[00017] In some embodiments, the natural ligand is a mammalian B7-1. In some embodiments, the natural ligand is selected from the group consisting of: a human B7-1, a murine B7-1, and a cynomolgus monkey B7-1.

[00018] In some embodiments, the antibody comprises a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56. In some embodiments, the antibody comprises a light chain amino acid sequence comprising SEQ ID NO: 12 or SEQ ID NO: 58. In some embodiments, the antibody comprises a light chain amino acid sequence comprising SEQ ID NO: 12. In some embodiments, the antibody comprises a light chain amino acid sequence comprising SEQ ID NO: 58.

[00019] In some embodiments, the antibody comprises a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid sequence comprising SEQ ID NO: 58.

[00020] In some embodiments, the antibody comprises a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid sequence comprising SEQ ID NO: 12.

[00021] In some embodiments, the antibody comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56. In some embodiments, the antibody comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, the antibody comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12. In some embodiments, the antibody comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, the antibody comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12.

[00022] In some embodiments, the antibody comprises the heavy chain amino acid sequence of SEQ ID NO: 46 and the light chain amino acid sequence of SEQ ID NO: 58. In some embodiments, the antibody comprises the heavy chain amino acid sequence of SEQ ID NO: 46 and the light chain amino acid sequence of SEQ ID NO: 12.

[00023] In some embodiments, the antibody comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid of SEQ ID NO: 46. In some embodiments, the antibody comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, the antibody comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12. In some embodiments, the antibody comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence selected of SEQ ID NO: 46, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, the antibody comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence selected of SEQ ID NO: 46, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12.

[00024] In some embodiments, the antibody comprises a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3) sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2) sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence shown in Table 16; a VH CDR2 sequence shown in Table 16; a VH CDR3 sequence shown in Table 16; a VL CDR1 sequence shown in Table 16; a VL CDR2 sequence shown in Table 16; and a VL CDR3 sequence shown in Table 16.

[00025] In some embodiments, the antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR1 sequence shown in Table 16; a VH CD2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR2 sequence shown in Table 16; a VH CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR3 sequence shown in Table 16; a VL CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR1 sequence shown in Table 16; a VL CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR2 sequence shown in Table 16; and a VL CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR3 sequence shown in Table 16.

[00026] In some embodiments, the antibody comprises a heavy chain that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein the combination is a combination of the three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 16.

[00027] In some embodiments, the antibody comprises a light chain that comprise a combination of a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the three light chain CDR sequences (VL CDR1, VL CDR2, VL CDR3) shown in a single row in Table 16.

[00028] In some embodiments, the antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 16.

[00029] In some embodiments, the antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%,

99% or more identical to the corresponding CDR sequence in a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 16.

[00030] In some embodiments, the antibody comprises a heavy chain that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 16.

[00031] In some embodiments, the antibody comprises a light chain that comprise a combination of a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of the three light chain CDR sequences (VL CDR1, VL CDR2, VL CDR3) shown in a single row in Table 16.

[00032] In some embodiments, the antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VL CDR1 sequence comprising RASQSISSYLN (SEQ ID NO: 209); a VL CDR2 sequence comprising AASSLQS (SEQ ID NO: 215); a VL CDR3 sequence comprising DNGYPST (SEQ ID NO: 228); a VH CDR1 sequence comprising SYAMS (SEQ ID NO: 212); a VH CDR2 sequence comprising SSIWRNGIVTVYADS (SEQ ID NO: 246); and a VH CDR3 sequence comprising WSAAFDY (SEQ ID NO: 235).

[00033] In some embodiments, the antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR2 sequence comprises SSIWRNGIVTVYADS (SEQ ID NO: 246). In some embodiments, the antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR3 sequence comprises WSAAFDY (SEQ ID NO: 235). In some embodiments, the antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence,

wherein the VH CDR2 sequence comprises SSIWRNGIVTVYADS (SEQ ID NO: 246), and the VH CDR3 sequence comprises WSAAFDY (SEQ ID NO: 235).

[00034] In some embodiments, the antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination comprises a VL CDR1 sequence comprising RASQSISSYLN (SEQ ID NO: 209); a VL CDR2 sequence comprising AASSLQS (SEQ ID NO: 215); a VL CDR3 sequence comprising DNGYPST (SEQ ID NO: 228); a VH CDR1 sequence comprising SYAMS (SEQ ID NO: 212); a VH CDR2 sequence comprising SSIWRNGIVTVYADS (SEQ ID NO: 246); and a VH CDR3 sequence comprising WSAAFDY (SEQ ID NO: 235).

[00035] In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 12 or SEQ ID NO: 58. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 12. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 58. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 58. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 12. In some embodiments, the antibody is encoded by a nucleic

acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 58. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 12.

[00036] In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence

selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12.

[00037] In some embodiments, the antibody is encoded by a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, and 55. In some embodiments, the antibody is encoded by a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 45, 47, 49, 51, 53, and 55. In some embodiments, the antibody is encoded by the light chain nucleic acid sequence of SEQ ID NO: 57. In some embodiments, the antibody is encoded by the light chain nucleic acid sequence of SEQ ID NO: 11. In some embodiments, the antibody is encoded by heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, and 55, and a light chain nucleic acid sequence comprising SEQ ID NO: 57. In some embodiments, the antibody is encoded by heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, and 55, and a light chain nucleic acid sequence comprising SEQ ID NO: 11. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 45, 47, 49, 51, 53, and 55, and a light chain nucleic acid sequence comprising SEQ

ID NO: 57. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 45, 47, 49, 51, 53, and 55, and a light chain nucleic acid sequence comprising SEQ ID NO: 11.

[00038] In some embodiments, the antibody is encoded by a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, and 55. In some embodiments, the antibody is encoded by a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 45, 47, 49, 51, 53, and 55. In some embodiments, the antibody is encoded by a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the light chain nucleic acid sequence of SEQ ID NO: 57. In some embodiments, the antibody is encoded by a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the light chain nucleic acid sequence of SEQ ID NO: 11. In some embodiments, the antibody is encoded by a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, and 55, and a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the light chain nucleic acid sequence comprising SEQ ID NO: 57. In some embodiments, the antibody is encoded by a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, and 55, and a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the light chain nucleic acid sequence comprising SEQ ID NO: 11. In some embodiments, the antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 45, 47, 49, 51, 53, and 55, and a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to light chain nucleic acid sequence comprising SEQ ID NO: 57. In some embodiments, the antibody is

encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 45, 47, 49, 51, 53, and 55, and a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to light chain nucleic acid sequence comprising SEQ ID NO: 11.

[00039] In some embodiments, the antibody or antigen-binding fragment thereof is incorporated in a multispecific antibody or antigen-binding fragment thereof, where at least one arm of the multispecific antibody specifically binds PDL1. In some embodiments, the antibody or antigen-binding fragment thereof is incorporated in a bispecific antibody or antigen-binding fragment thereof, where at least one arm of the bispecific antibody specifically binds PDL1.

[00040] In some embodiments, at least one arm of the multispecific antibody, *e.g.*, a bispecific antibody, comprises a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56. In some embodiments, at least one arm of the multispecific antibody, *e.g.*, a bispecific antibody, comprises a light chain amino acid sequence comprising SEQ ID NO: 12 or SEQ ID NO: 58. In some embodiments, at least one arm of the multispecific antibody, *e.g.*, a bispecific antibody, comprises a light chain amino acid sequence comprising SEQ ID NO: 12. In some embodiments, at least one arm of the multispecific antibody, *e.g.*, a bispecific antibody, comprises a light chain amino acid sequence comprising SEQ ID NO: 58.

[00041] In some embodiments, at least one arm of the multispecific antibody, *e.g.*, a bispecific antibody, a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid sequence comprising SEQ ID NO: 58.

[00042] In some embodiments, at least one arm of the multispecific antibody, *e.g.*, a bispecific antibody, comprises a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid sequence comprising SEQ ID NO: 12.

[00043] In some embodiments, at least one arm of the multispecific antibody, *e.g.*, a bispecific antibody, comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38,

40, 42, 44, 46, 48, 50, 52, 54, and 56. In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12. In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12.

[00044] In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises the heavy chain amino acid sequence of SEQ ID NO: 46 and the light chain amino acid sequence of SEQ ID NO: 58. In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises the heavy chain amino acid sequence of SEQ ID NO: 46 and the light chain amino acid sequence of SEQ ID NO: 12.

[00045] In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid of SEQ ID NO: 46. In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID

NO: 12. In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence selected of SEQ ID NO: 46, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence selected of SEQ ID NO: 46, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12.

[00046] In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3) sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2) sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence shown in Table 16; a VH CDR2 sequence shown in Table 16; a VH CDR3 sequence shown in Table 16; a VL CDR1 sequence shown in Table 16; a VL CDR2 sequence shown in Table 16; and a VL CDR3 sequence shown in Table 16.

[00047] In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR1 sequence shown in Table 16; a VH CD2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR2 sequence shown in

Table 16; a VH CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR3 sequence shown in Table 16; a VL CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR1 sequence shown in Table 16; a VL CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR2 sequence shown in Table 16; and a VL CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR3 sequence shown in Table 16.

[00048] In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 16.

[00049] In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a heavy chain that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein the combination is a combination of the three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 16.

[00050] In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 16.

[00051] In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a heavy chain that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a

combination of three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 16.

[00052] In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VL CDR1 sequence comprising RASQSISSYLN (SEQ ID NO: 209); a VL CDR2 sequence comprising AASSLQS (SEQ ID NO: 215); a VL CDR3 sequence comprising DNGYPST (SEQ ID NO: 228); a VH CDR1 sequence comprising SYAMS (SEQ ID NO: 212); a VH CDR2 sequence comprising SSIWRNGIVTVYADS (SEQ ID NO: 246); and a VH CDR3 sequence comprising WSAAFDY (SEQ ID NO: 235).

[00053] In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR2 sequence comprises SSIWRNGIVTVYADS (SEQ ID NO: 246). In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR3 sequence comprises WSAAFDY (SEQ ID NO: 235). In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR2 sequence comprises SSIWRNGIVTVYADS (SEQ ID NO: 246), and the VH CDR3 sequence comprises WSAAFDY (SEQ ID NO: 235).

[00054] In some embodiments, at least one arm of the multispecific antibody, e.g., a bispecific antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination comprises a VL CDR1 sequence comprising RASQSISSYLN (SEQ ID NO: 209); a VL CDR2 sequence comprising AASSLQS (SEQ ID NO: 215); a VL CDR3 sequence comprising DNGYPST (SEQ ID NO: 228); a VH CDR1 sequence comprising SYAMS (SEQ ID NO: 212); a VH CDR2 sequence comprising SSIWRNGIVTVYADS (SEQ ID NO: 246); and a VH CDR3 sequence comprising WSAAFDY (SEQ ID NO: 235).

[00055] The disclosure also provides activatable antibodies that include an antibody or antigen-binding fragment thereof that specifically binds PDL1 coupled to a masking moiety (MM), such that coupling of the MM reduces the ability of the antibody or antigen-binding fragment thereof to bind PDL1. In some embodiments, the MM is coupled via a sequence that includes a substrate for a protease, for example, a protease that is active in diseased tissue and/or a protease that is co-localized with PDL1 at a treatment site in a subject. The activatable anti-PDL1 antibodies provided herein, also referred to herein as anti-PDL1 activatable antibodies or PDL1 activatable antibodies, are stable in circulation, activated at intended sites of therapy and/or diagnosis but not in normal, *e.g.*, healthy tissue or other tissue not targeted for treatment and/or diagnosis, and, when activated, exhibit binding to PDL1 that is at least comparable to the corresponding, unmodified antibody.

[00056] In some embodiments, the activatable antibody that, in an activated state, specifically binds to mammalian PDL1, includes an antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian PDL1; a masking moiety (MM) that inhibits the binding of the AB to mammalian PDL1 when the activatable antibody is in an uncleaved state; and a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease.

[00057] In some embodiments, the activatable antibody that, in an activated state, has one or more of the following characteristics: (a) specifically binds to mammalian PDL1; and (b) specifically blocks a natural ligand of PDL1 from binding to the mammalian PDL1, wherein the activatable antibody comprises: an antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian PDL1; a masking moiety (MM) that inhibits the binding of the AB to mammalian PDL1 when the activatable antibody is in an uncleaved state; and a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease.

[00058] In some embodiments, the activatable antibody in an uncleaved state specifically binds to the mammalian PDL1 with a dissociation constant less than or equal to 1 nM, less than or equal to 2 nM, less than or equal to 3 nM, less than or equal to 4 nM, less than or equal to 5 nM, less than or equal to 10 nM, less than or equal to 15 nM, less than or equal to 20 nM, and/or less than or equal to 25 nM.

[00059] In some embodiments, the activatable antibody in an uncleaved state specifically binds to the mammalian PDL1 with a dissociation constant greater than or equal to 1 nM, greater than or equal to 2 nM, greater than or equal to 3 nM, greater than or equal

to 4 nM, greater than or equal to 5 nM, greater than or equal to 10 nM, greater than or equal to 15 nM, greater than or equal to 20 nM, and/or greater than or equal to 25 nM.

[00060] In some embodiments, the activatable antibody in an uncleaved state specifically binds to the mammalian PDL1 with a dissociation constant in the range of 1 to 2 nM, 1 to 5 nM, 1 to 10 nM, 1 to 15 nM, 1 to 20 nM, 1 to 25 nM, 2 nM to 5 nM, 2 nM to 10 nM, 2 nM to 15 nM, 2 to 20 nM, 2 to 25 nM, 5 nM to 10 nM, 5 nM to 15 nM, 5 to 20 nM, 5 to 25 nM, 10 nM to 15 nM, 10 to 20 nM, 10 to 25 nM, 15 to 20 nM, 15 to 25 nM, or 20 to 25 nM.

[00061] In some embodiments, the activatable antibody in an activated state specifically binds to the mammalian PDL1 with a dissociation constant is less than or equal to 0.01 nM, less than or equal to 0.05 nM, less than or equal to 0.1 nM, less than or equal to 0.2 nM, less than or equal to 0.3 nM, less than or equal to 0.4 nM, less than or equal to 0.5 nM, less than or equal to 0.75 nM, and less than or equal to 1 nM.

[00062] In some embodiments, the activatable antibody in an activated state specifically binds to the mammalian PDL1 with a dissociation constant in the range of 0.01 nM to 1 nM, 0.05 nM to 1 nM, 0.1 nM to 1 nM, 0.2 nM to 1 nM, 0.3 nM to 1 nM, 0.4 nM to 1 nM, 0.5 nM to 1 nM, 0.75 nM to 1 nM, 0.01 nM to 0.75 nM, 0.05 nM to 0.75 nM, 0.1 nM to 0.75 nM, 0.2 nM to 0.75 nM, 0.3 nM to 0.75 nM, 0.4 nM to 0.75 nM, 0.5 nM to 0.75 nM, 0.01 nM to 0.5 nM, 0.05 nM to 0.5 nM, 0.1 nM to 0.5 nM, 0.2 nM to 0.5 nM, 0.3 nM to 0.5 nM, 0.4 nM to 0.5 nM, 0.01 nM to 0.4 nM, 0.05 nM to 0.4 nM, 0.1 nM to 0.4 nM, 0.2 nM to 0.4 nM, 0.3 nM to 0.4 nM, 0.01 nM to 0.3 nM, 0.05 nM to 0.3 nM, 0.1 nM to 0.3 nM, 0.2 nM to 0.3 nM, 0.01 nM to 0.2 nM, 0.05 nM to 0.2 nM, 0.1 nM to 0.2 nM, 0.01 nM to 0.1 nM, 0.05 nM to 0.1 nM, or 0.01 nM to 0.05 nM.

[00063] In some embodiments, the AB specifically binds to the mammalian PDL1 with a dissociation constant is less than or equal to 0.01 nM, less than or equal to 0.05 nM, less than or equal to 0.1 nM, less than or equal to 0.2 nM, less than or equal to 0.3 nM, less than or equal to 0.4 nM, less than or equal to 0.5 nM, less than or equal to 0.75 nM, and less than or equal to 1 nM.

[00064] In some embodiments, the AB specifically binds to the mammalian PDL1 with a dissociation constant in the range of 0.01 nM to 1 nM, 0.05 nM to 1 nM, 0.1 nM to 1 nM, 0.2 nM to 1 nM, 0.3 nM to 1 nM, 0.4 nM to 1 nM, 0.5 nM to 1 nM, 0.75 nM to 1 nM, 0.01 nM to 0.75 nM, 0.05 nM to 0.75 nM, 0.1 nM to 0.75 nM, 0.2 nM to 0.75 nM, 0.3 nM to 0.75 nM, 0.4 nM to 0.75 nM, 0.5 nM to 0.75 nM, 0.01 nM to 0.5 nM, 0.05 nM to 0.5 nM,

0.1 nM to 0.5 nM, 0.2 nM to 0.5 nM, 0.3 nM to 0.5 nM, 0.4 nM to 0.5 nM, 0.01 nM to 0.4 nM, 0.05 nM to 0.4 nM, 0.1 nM to 0.4 nM, 0.2 nM to 0.4 nM, 0.3 nM to 0.4 nM, 0.01 nM to 0.3 nM, 0.05 nM to 0.3 nM, 0.1 nM to 0.3 nM, 0.2 nM to 0.3 nM, 0.01 nM to 0.2 nM, 0.05 nM to 0.2 nM, 0.1 nM to 0.2 nM, 0.01 nM to 0.1 nM, 0.05 nM to 0.1 nM, or 0.01 nM to 0.05 nM.

[00065] In some embodiments, the mammalian PDL1 is selected from the group consisting of a human PDL1, a murine PDL1, a rat PDL1, and a cynomolgus monkey PDL1. In some embodiments, the AB specifically binds to human PDL1, murine PDL1 or cynomolgus monkey PDL1 with a dissociation constant of less than 1 nM. In some embodiments, the mammalian PDL1 is a human PDL1.

[00066] In some embodiments, the AB has one or more of the following characteristics: (a) the AB specifically binds to human PDL1 and murine PDL1; (b) the AB specifically binds to human PDL1 and cynomolgus monkey PDL1; and (c) the AB specifically binds to human PDL1, murine PDL1, and cynomolgus monkey PDL1.

[00067] In some embodiments, the AB has one or more of the following characteristics: (a) the AB specifically binds human PDL1, murine PDL1, and cynomolgus monkey PDL1; (b) the AB inhibits binding of human B7-1 and human PD1 to human PDL1; (c) the AB inhibits binding of murine B7-1 and murine PD1 to murine PDL1; and (d) the AB inhibits binding of cynomolgus monkey B7-1 and cynomolgus monkey PD1 to cynomolgus monkey PDL1.

[00068] In some embodiments, the AB blocks the ability of a natural ligand to bind to the mammalian PDL1 with an EC_{50} less than or equal to 0.5 nM, less than or equal to 1 nM, less than or equal to 2 nM, less than or equal to 3 nM, less than or equal to 4 nM, less than or equal to 5 nM, less than or equal to 6 nM, less than or equal to 7 nM, less than or equal to 8 nM, less than or equal to 9 nM, and/or less than or equal to 10 nM.

[00069] In some embodiments, the AB blocks the ability of a natural ligand to bind to the mammalian PDL1 with an EC_{50} of 0.5 nM to 10 nM, 0.5 nM to 5 nM, 0.5 nM to 3 nM, 0.5 nM to 2 nM, 0.5 nM to 1 nM, 1 nM to 10 nM, 1 nM to 5 nM, 1 nM to 3 nM, 1 nM to 2 nM, 2 nM to 10 nM, 2 nM to 5 nM, 2 nM to 3 nM, 3 nM to 10 nM, 3 nM to 5 nM, or 5 nM to 10 nM.

[00070] In some embodiments, natural ligand is a mammalian PD1. In some embodiments, the natural ligand is selected from the group consisting of: a human PD1, a murine PD1, and a cynomolgus monkey PD1.

[00071] In some embodiments, the natural ligand is a mammalian B7-1. In some embodiments, the natural ligand is selected from the group consisting of: a human B7-1, a murine B7-1, and a cynomolgus monkey B7-1.

[00072] In some embodiments, the activatable antibody has one or more of the following characteristics (a) the AB induces type 1 diabetes in a non-obese diabetic (NOD) mouse; and (b) the activatable antibody in an uncleaved state inhibits the induction of type 1 diabetes in a NOD mouse.

[00073] In some embodiments, the activatable antibody inhibits the induction of type 1 diabetes in the NOD mouse after administration of the activatable antibody at a single dose of 0.1 mg/kg to 3 mg/kg, 0.5 mg/kg to 3 mg/kg, 1 mg/kg to 3 mg/kg, 2 mg/kg to 3 mg/kg, 0.1 mg/kg to 2 mg/kg, 0.5 mg/kg to 2 mg/kg, 1 mg/kg to 2 mg/kg, 0.1 mg/kg to 1 mg/kg, 0.5 mg/kg to 1 mg/kg, or 0.1 mg/kg to 0.5 mg/kg.

[00074] In some embodiments, the activatable antibody induces type 1 diabetes in a subject after administration of the activatable antibody at a single dose of 1 mg/kg. In some embodiments, the subject is a NOD mouse.

[00075] In some embodiments, the activatable antibody induces type 1 diabetes in a subject after administration of the activatable antibody at single dose less than or equal to 3 mg/kg, less than or equal to 2 mg/kg, less than or equal to 1 mg/kg, less than or equal to 0.5 mg/kg, and/or less than or equal to 0.1 mg/kg.

[00076] In some embodiments, the activatable antibody induces type 1 diabetes in a subject after administration of the activatable antibody at single dose in the range selected from the group consisting of 0.1 mg/kg to 3 mg/kg, 0.5 mg/kg to 3 mg/kg, 1 mg/kg to 3 mg/kg, 2 mg/kg to 3 mg/kg, 0.1 mg/kg to 2 mg/kg, 0.5 mg/kg to 2 mg/kg, 1 mg/kg to 2 mg/kg, 0.1 mg/kg to 1 mg/kg, 0.5 mg/kg to 1 mg/kg, and 0.1 mg/kg to 0.5 mg/kg. In some embodiments, the subject is a NOD mouse.

[00077] In some embodiments, the activatable antibody has one or more of the following characteristics (a) the activatable antibody in an uncleaved state does not induce type 1 diabetes in greater than 50% of a population of non-obese diabetic (NOD) mice, and the AB induces type 1 diabetes in greater than 50% of a population of NOD mice.

[00078] In some embodiments, the activatable antibody does not induce type 1 diabetes in greater than 50% of the population of NOD mice after administration to each mouse in the population a single dose less than or equal to 3 mg/kg, less than or equal to

2 mg/kg, less than or equal to 1 mg/kg, less than or equal to 0.5 mg/kg, and/or less than or equal to 0.1 mg/kg.

[00079] In some embodiments, the activatable antibody does not induce type 1 diabetes in greater than 50% of the population of NOD mice after administration to each mouse in the population a single dose of the activatable antibody at a dosage of: 0.1 mg/kg to 3 mg/kg, 0.5 mg/kg to 3 mg/kg, 1 mg/kg to 3 mg/kg, 2 mg/kg to 3 mg/kg, 0.1 mg/kg to 2 mg/kg, 0.5 mg/kg to 2 mg/kg, 1 mg/kg to 2 mg/kg, 0.1 mg/kg to 1 mg/kg, 0.5 mg/kg to 1 mg/kg, or 0.1 mg/kg to 0.5 mg/kg.

[00080] In some embodiments, the activatable antibody has one or more of the following characteristics (a) the activatable antibody in an uncleaved state does not induce type 1 diabetes in greater than 50% of a population of non-obese diabetic (NOD) mice when administered at a single dose of 1 mg/kg; and (b) the AB induces type 1 diabetes in greater than 50% of a population of NOD mice, when administered at a single dose of 1 mg/kg.

[00081] In some embodiments, the AB induces type 1 diabetes in greater than 50% of the population of the NOD mice after administration to each mouse in the population a single dose of the AB at a dosage less than or equal to 3 mg/kg, less than or equal to 2 mg/kg, less than or equal to 1 mg/kg, less than or equal to 0.5 mg/kg, and/or less than or equal to 0.1 mg/kg.

[00082] In some embodiments, the AB induces type 1 diabetes in greater than 50% of the population of the NOD mice after administration to each mouse in the population a single dose of the AB at a dosage of: 0.1 mg/kg to 3 mg/kg, 0.5 mg/kg to 3 mg/kg, 1 mg/kg to 3 mg/kg, 2 mg/kg to 3 mg/kg, 0.1 mg/kg to 2 mg/kg, 0.5 mg/kg to 2 mg/kg, 1 mg/kg to 2 mg/kg, 0.1 mg/kg to 1 mg/kg, 0.5 mg/kg to 1 mg/kg, or 0.1 mg/kg to 0.5 mg/kg.

[00083] In some embodiments, the NOD mouse is a female NOD/ShiLtJ mouse substrain.

[00084] In some embodiments, the activatable antibody inhibits the induction of type 1 diabetes in a NOD mouse by at least 3-fold compared to the AB.

[00085] In some embodiments, the activatable antibody exhibits a safety margin that is at least a three-fold safety margin relative to the AB.

[00086] In some embodiments, the activatable antibody in an uncleaved state binds to a smaller percentage of a population of peripheral blood CD4⁺ CD8⁺ T lymphocytes than does the AB.

[00087] In some embodiments, the activatable antibody does not bind to greater than 50% of the population of peripheral blood CD4+ CD8+ T lymphocytes after administration of a single dose of the activatable antibody at a dosage less than or equal to 5 mg/kg, less than or equal to 4 mg/kg, less than or equal to 3 mg/kg, less than or equal to 2 mg/kg, less than or equal to 1 mg/kg, less than or equal to 0.5 mg/kg, and/or less than or equal to 0.1 mg/kg.

[00088] In some embodiments, the activatable antibody does not bind to greater than 50% of the population of peripheral blood CD4+ CD8+ T lymphocytes after administration of a single dose of the activatable antibody at a dosage of: 0.1 mg/kg to 5 mg/kg, 0.5 mg/kg to 5 mg/kg, 1 mg/kg to 5 mg/kg, 2 mg/kg to 5 mg/kg, 3 mg/kg to 5 mg/kg, 0.1 mg/kg to 3 mg/kg, 0.5 mg/kg to 3 mg/kg, 1 mg/kg to 3 mg/kg, 2 mg/kg to 3 mg/kg, 0.1 mg/kg to 2 mg/kg, 0.5 mg/kg to 2 mg/kg, 1 mg/kg to 2 mg/kg, 0.1 mg/kg to 1 mg/kg, 0.5 mg/kg to 1 mg/kg, or 0.1 mg/kg to 0.5 mg/kg.

[00089] In some embodiments, the AB binds to greater than 50% of the population of peripheral blood CD4+ CD8+ T lymphocytes after administration of a single dose of the AB at a dosage of: 0.1 mg/kg to 5 mg/kg, 0.5 mg/kg to 5 mg/kg, 1 mg/kg to 5 mg/kg, 2 mg/kg to 5 mg/kg, 3 mg/kg to 5 mg/kg, 0.1 mg/kg to 3 mg/kg, 0.5 mg/kg to 3 mg/kg, 1 mg/kg to 3 mg/kg, 2 mg/kg to 3 mg/kg, 0.1 mg/kg to 2 mg/kg, 0.5 mg/kg to 2 mg/kg, 1 mg/kg to 2 mg/kg, 0.1 mg/kg to 1 mg/kg, 0.5 mg/kg to 1 mg/kg, or 0.1 mg/kg to 0.5 mg/kg.

[00090] In some embodiments, the peripheral blood CD4+ CD8+ T lymphocytes are murine. In some embodiments, the murine peripheral blood CD4+ CD8+ T lymphocytes are derived from a tumor-bearing mouse.

[00091] In some embodiments, the percentage of the population of peripheral blood CD4+CD8+ T lymphocytes to which the AB binds is less than 60% when the activatable antibody is administered at a dose between 1 mg/kg and 5 mg/kg. In some embodiments, the percentage of the population of peripheral blood CD4+CD8+ T lymphocytes to which the AB binds is less than 50% when the activatable antibody is administered at a dose between 1 mg/kg and 5 mg/kg. In some embodiments, the percentage of the population of peripheral blood CD4+CD8+ T lymphocytes to which the AB binds is between 30 and 60% when the activatable antibody is administered at a dose between 1 mg/kg and 5 mg/kg. In some embodiments, the percentage of the population of peripheral blood CD4+CD8+ T lymphocytes to which the AB binds is between 30 and 50% when the activatable antibody is administered at a dose between 1 mg/kg and 5 mg/kg.

[00092] The invention also provides methods of treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with aberrant expression and/or activity of PDL1 in a subject using activatable antibodies that bind PDL1, particularly activatable antibodies that bind and neutralize or otherwise inhibit at least one biological activity of PDL1 and/or PDL1-mediated signaling.

[00093] The activatable antibodies in an activated state bind PDL1 and include (i) an antibody or an antigen binding fragment thereof (AB) that specifically binds to PDL1; (ii) a masking moiety (MM) that, when the activatable antibody is in an uncleaved state, inhibits the binding of the AB to PDL1; and (c) a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease.

[00094] In some embodiments, the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM.

[00095] In some embodiments, the activatable antibody comprises a linking peptide between the MM and the CM.

[00096] In some embodiments, the activatable antibody comprises a linking peptide between the CM and the AB.

[00097] In some embodiments, the activatable antibody comprises a first linking peptide (LP1) and a second linking peptide (LP2), and wherein the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-LP1-CM-LP2-AB or AB-LP2-CM-LP1-MM. In some embodiments, the two linking peptides need not be identical to each other.

[00098] In some embodiments, at least one of LP1 or LP2 comprises an amino acid sequence selected from the group consisting of (GS)_n, (GGS)_n, (GSGGS)_n (SEQ ID NO: 191) and (GGGS)_n (SEQ ID NO: 192), where n is an integer of at least one.

[00099] In some embodiments, at least one of LP1 or LP2 comprises an amino acid sequence selected from the group consisting of GGSG (SEQ ID NO: 193), GGS GG (SEQ ID NO: 194), GSGSG (SEQ ID NO: 195), GSGGG (SEQ ID NO: 196), GGGSG (SEQ ID NO: 197), and GSSSG (SEQ ID NO: 198).

[000100] In some embodiments, LP1 comprises the amino acid sequence GSSGGSGGSGGSG (SEQ ID NO: 199), GSSGGSGGSGG (SEQ ID NO: 200), GSSGGSGGSGGS (SEQ ID NO: 201), GSSGGSGGSGGSGGGS (SEQ ID NO: 202), GSSGGSGGSG (SEQ ID NO: 203), or GSSGGSGGSGS (SEQ ID NO: 204).

[000101] In some embodiments, LP2 comprises the amino acid sequence GSS, GGS, GGGS (SEQ ID NO: 205), GSSGT (SEQ ID NO: 206) or GSSG (SEQ ID NO: 207).

[000102] In some embodiments, the AB has a dissociation constant of about 100 nM or less for binding to PDL1.

[000103] In some embodiments, the activatable antibody includes an antibody or antigen-binding fragment thereof (AB) that specifically binds PDL1. In some embodiments, the antibody or antigen binding fragment thereof that binds PDL1 is a monoclonal antibody, domain antibody, single chain, Fab fragment, a F(ab')₂ fragment, a scFv, a scAb, a dAb, a single domain heavy chain antibody, or a single domain light chain antibody. In some embodiments, such an antibody or antigen binding fragment thereof that binds PDL1 is a mouse, other rodent, chimeric, humanized or fully human monoclonal antibody.

[000104] In some embodiments, the activatable antibody comprises a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56. In some embodiments, the activatable antibody comprises a light chain amino acid sequence comprising SEQ ID NO: 12 or SEQ ID NO: 58. In some embodiments, the activatable antibody comprises a light chain amino acid sequence comprising SEQ ID NO: 12. In some embodiments, the activatable antibody comprises a light chain amino acid sequence comprising SEQ ID NO: 58.

[000105] In some embodiments, the activatable antibody comprises a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid sequence comprising SEQ ID NO: 58.

[000106] In some embodiments, the activatable antibody comprises a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid sequence comprising SEQ ID NO: 12.

[000107] In some embodiments, the activatable antibody comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56. In some embodiments, the activatable antibody comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid

sequence of SEQ ID NO: 58. In some embodiments, the activatable antibody comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12. In some embodiments, the activatable antibody comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, the activatable antibody comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12.

[000108] In some embodiments, the AB of the activatable anti-PDL1 antibody comprises a heavy chain amino acid sequence selected from the group consisting of the heavy chain sequences shown in Table 15. In some embodiments, the AB of the activatable anti-PDL1 antibody comprises a light chain amino acid sequence selected from the group consisting of the light chain sequences shown in Table 15. In some embodiments, the AB of the activatable anti-PDL1 antibody comprises a heavy chain amino acid sequence selected from the group consisting of the heavy chain sequences shown in Table 15 and a light chain amino acid sequence selected from the group consisting of the light chain sequences shown in Table 15.

[000109] In some embodiments, the AB of the activatable anti-PDL1 antibody comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the heavy chain sequences shown in Table 15. In some embodiments, the AB of the activatable anti-PDL1 antibody comprises a light chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the light chain sequences shown in Table 15. In some embodiments, the AB of the activatable anti-PDL1 antibody comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the heavy chain sequences shown in Table 15 and a light chain amino acid sequence that is at

least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of the light chain sequences shown in Table 15.

[000110] In some embodiments, the activatable antibody comprises a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3) sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2) sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence shown in Table 16; a VH CDR2 sequence shown in Table 16; a VH CDR3 sequence shown in Table 16; a VL CDR1 sequence shown in Table 16; a VL CDR2 sequence shown in Table 16; and a VL CDR3 sequence shown in Table 16.

[000111] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR1 sequence shown in Table 16; a VH CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR2 sequence shown in Table 16; a VH CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR3 sequence shown in Table 16; a VL CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR1 sequence shown in Table 16; a VL CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR2 sequence shown in Table 16; and a VL CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR3 sequence shown in Table 16.

[000112] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence,

a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 16.

[000113] In some embodiments, the activatable antibody comprises a heavy chain that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein the combination is a combination of the three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 16.

[000114] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 16.

[000115] In some embodiments, the activatable antibody comprises a heavy chain that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 16.

[000116] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VL CDR1 sequence comprising RASQSISSYLN (SEQ ID NO: 209); a VL CDR2 sequence comprising AASSLQS (SEQ ID NO: 215); a VL CDR3 sequence comprising DNGYPST (SEQ ID NO: 228); a VH CDR1 sequence comprising SYAMS (SEQ ID NO: 212); a VH CDR2 sequence comprising SSIWRNGIVTVYADS (SEQ ID NO: 246); and a VH CDR3 sequence comprising WSAAFDY (SEQ ID NO: 235).

[000117] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR2 sequence comprises SSIWRNGIVTVYADS (SEQ ID NO: 246). In some embodiments, the

activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR3 sequence comprises WSAAFDY (SEQ ID NO: 235). In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR2 sequence comprises SSIWRNGIVTVYADS (SEQ ID NO: 246), and the VH CDR3 sequence comprises WSAAFDY (SEQ ID NO: 235).

[000118] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination comprises a VL CDR1 sequence comprising RASQSISSYLN (SEQ ID NO: 209); a VL CDR2 sequence comprising AASSLQS (SEQ ID NO: 215); a VL CDR3 sequence comprising DNGYPST (SEQ ID NO: 228); a VH CDR1 sequence comprising SYAMS (SEQ ID NO: 212); a VH CDR2 sequence comprising SSIWRNGIVTVYADS (SEQ ID NO: 246); and a VH CDR3 sequence comprising WSAAFDY (SEQ ID NO: 235).

[000119] In some embodiments, the activatable antibody comprises a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3) sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2) sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence shown in Table 17; a VH CDR2 sequence shown in Table 17; a VH CDR3 sequence shown in Table 17; a VL CDR1 sequence shown in Table 17; a VL CDR2 sequence shown in Table 17; and a VL CDR3 sequence shown in Table 17.

[000120] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence that includes a sequence that is

at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR1 sequence shown in Table 17; a VH CD2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR2 sequence shown in Table 17; a VH CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR3 sequence shown in Table 17; a VL CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR1 sequence shown in Table 17; a VL CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR2 sequence shown in Table 17; and a VL CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR3 sequence shown in Table 17.

[000121] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination shown in Table 17.

[000122] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination shown in Table 17.

[000123] In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201. In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56. In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141,

143, 145, 147, 149, 151, 153, 155, 157, 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201, and an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56.

[000124] In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201. In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56. In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201, and an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56.

[000125] In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1145, 1147, 1149, 1151, 1153, 1155, 1157, 1159, 1161, 1163, 1165, 1167, 1169, 1171, 1173, 1175, 1177, 1179, 1181, 1183, 1185, 1187, 1189, 1191, and 1201. In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56. In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 100, 1145, 1147, 1149, 1151, 1153, 1155, 1157, 1159, 1161, 1163, 1165, 1167, 1169, 1171, 1173, 1175, 1177, 1179, 1181, 1183, 1185, 1187, 1189, 1191, and 1201, and an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56.

[000126] In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino

acid sequence selected from the group consisting of SEQ ID NOs: 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201. In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56. In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201, and an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56.

[000127] In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201. In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56. In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201, and an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%

identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56.

[000128] In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1145, 1147, 1149, 1151, 1153, 1155, 1157, 1159, 1161, 1163, 1165, 1167, 1169, 1171, 1173, 1175, 1177, 1179, 1181, 1183, 1185, 1187, 1189, 1191, 1200, and 1201. In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56. In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1145, 1147, 1149, 1151, 1153, 1155, 1157, 1159, 1161, 1163, 1165, 1167, 1169, 1171, 1173, 1175, 1177, 1179, 1181, 1183, 1185, 1187, 1189, 1191, and 1201, and an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56.

[000129] In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201, and the heavy chain amino acid sequence of SEQ ID NO: 46.

[000130] In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201, and the heavy chain amino acid sequence of SEQ ID NO: 46.

[000131] In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1145, 1147, 1149, 1151, 1153, 1155, 1157, 1159, 1161, 1163, 1165, 1167, 1169, 1171, 1173, 1175, 1177, 1179, 1181, 1183, 1185, 1187, 1189, 1191, and 1201, and the heavy chain amino acid sequence of SEQ ID NO: 46.

[000132] In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201, and an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the heavy chain amino acid sequence of SEQ ID NO: 46.

[000133] In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201, and an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the heavy chain amino acid sequence of SEQ ID NO: 46.

[000134] In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1145, 1147, 1149, 1151, 1153, 1155, 1157, 1159, 1161, 1163, 1165, 1167, 1169, 1171, 1173, 1175, 1177, 1179, 1181, 1183, 1185, 1187, 1189, 1191, and 1201, and an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the heavy chain amino acid sequence of SEQ ID NO: 46.

[000135] In some embodiments, the activatable antibody comprises the amino acid sequence of SEQ ID NO: 428. In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 430, 432, 434, and 1202. In some embodiments, the activatable antibody comprises the amino acid sequence of SEQ ID NO: 428, and an amino acid sequence selected from the group consisting of SEQ ID NO: 430, 432, and 434.

[000136] In some embodiments, the activatable antibody comprises the amino acid sequence of SEQ ID NO: 1008. In some embodiments, the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 430, 432, 434, and 1202. In some embodiments, the activatable antibody comprises the amino acid sequence of SEQ ID NO: 1008, and an amino acid sequence selected from the group consisting of SEQ ID NO: 430, 432, 434, and 1202.

[000137] In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 428. In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 430, 432, 434, and 1202. In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 428, and an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 430, 432, 434, and 1202.

[000138] In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 1008. In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 430, 432, 434, and 1202. In some embodiments, the activatable antibody includes an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 1008, and an amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 430, 432, 434, and 1202.

[000139] In some embodiments, the activatable antibody comprises a masking moiety (MM) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 59-81, 208, and 426; a cleavable moiety (CM) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 341, 352, 359, 364, 372, 377, 379, 383, 394, 437, 883-893, 901-920, and 921; and a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination comprises a VL CDR1 sequence comprising RASQSISSYLN (SEQ ID NO: 209); a VL CDR2 sequence comprising AASSLQS (SEQ ID NO: 215); a VL CDR3 sequence comprising DNGYPST (SEQ ID NO: 228); a VH CDR1 sequence comprising SYAMS (SEQ ID NO: 212); a VH CDR2 sequence comprising SSIWRNGIVTVYADS (SEQ ID NO: 246); and a VH CDR3 sequence comprising WSAAFDY (SEQ ID NO: 235).

[000140] In some embodiments, the activatable antibody comprises a masking moiety (MM) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 60, 62, 63, 66-68, 71, 75, and 76; a cleavable moiety (CM) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 341, 364, 377, 394, 437, 883-893, 901-911, and 920; and a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination comprises a VL CDR1 sequence comprising RASQSISSYLN (SEQ ID NO: 209); a VL CDR2 sequence comprising AASSLQS (SEQ ID NO: 215); a VL CDR3 sequence comprising DNGYPST (SEQ ID NO: 228); a VH CDR1 sequence comprising SYAMS (SEQ ID NO: 212); a VH CDR2 sequence comprising SSIWRNGIVTVYADS (SEQ ID NO: 246); and a VH CDR3 sequence comprising WSAAFDY (SEQ ID NO: 235).

[000141] In some embodiments, the activatable antibody comprises a masking moiety (MM) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 59-81, 208, and 426; a cleavable moiety (CM) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 341, 352, 359, 364, 372, 377, 379, 383, 394, 437, 883-893, 901-920, and 921; and of a variable light chain comprising the amino acid sequence of SEQ ID NO: 58 and a variable heavy chain comprising the amino acid sequence of SEQ ID NO: 46.

[000142] In some embodiments, the activatable antibody comprises a masking moiety (MM) comprising an amino acid sequence selected from the group consisting of SEQ ID

NOs: 60, 62, 63, 66-68, 71, 75, and 76; a cleavable moiety (CM) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 341, 364, 377, 394, 437, 883-893, 901-911, and 920; and of a variable light chain comprising the amino acid sequence of SEQ ID NO: 58 and a variable heavy chain comprising the amino acid sequence of SEQ ID NO: 46.

[000143] In some embodiments, the activatable antibody comprises a masking moiety (MM) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 59-81, 208, and 426; a cleavable moiety (CM) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 341, 352, 359, 364, 372, 377, 379, 383, 394, 437, 883-893, 901-920, and 921; and of a variable light chain comprising the amino acid sequence of SEQ ID NO: 12 and a variable heavy chain comprising the amino acid sequence of SEQ ID NO: 46.

[000144] In some embodiments, the activatable antibody comprises a masking moiety (MM) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 60, 62, 63, 66-68, 71, 75, and 76; a cleavable moiety (CM) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 341, 364, 377, 394, 437, 883-893, 901-11, and 920; and of a variable light chain comprising the amino acid sequence of SEQ ID NO: 12 and a variable heavy chain comprising the amino acid sequence of SEQ ID NO: 46.

[000145] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 58. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 12. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42,

44, 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 58. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 12. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 58. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid sequence comprising SEQ ID NO: 12.

[000146] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain

amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence encoding a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a nucleic acid sequence that comprises a nucleic acid sequence encoding a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12.

[000147] In some embodiments, the activatable antibody is encoded by a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, and 55. In some embodiments, the activatable antibody is encoded by a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 45, 47, 49, 51, 53, and 55. In some embodiments, the activatable antibody is encoded by the light chain nucleic acid sequence of SEQ ID NO: 57.

In some embodiments, the activatable antibody is encoded by the light chain nucleic acid sequence of SEQ ID NO: 11. In some embodiments, the activatable antibody is encoded by heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, and 55, and a light chain nucleic acid sequence comprising SEQ ID NO: 57. In some embodiments, the activatable antibody is encoded by heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, and 55, and a light chain nucleic acid sequence comprising SEQ ID NO: 11. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 45, 47, 49, 51, 53, and 55, and a light chain nucleic acid sequence comprising SEQ ID NO: 57. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 45, 47, 49, 51, 53, and 55, and a light chain nucleic acid sequence comprising SEQ ID NO: 11.

[000148] In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, and 55. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 45, 47, 49, 51, 53, and 55. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the light chain nucleic acid sequence of SEQ ID NO: 57. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the light chain nucleic acid sequence of SEQ ID NO: 11. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, and 55, and a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the light chain nucleic acid sequence comprising SEQ ID NO: 57.

In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, and 55, and a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the light chain nucleic acid sequence comprising SEQ ID NO: 11. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 45, 47, 49, 51, 53, and 55, and a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to light chain nucleic acid sequence comprising SEQ ID NO: 57. In some embodiments, the activatable antibody is encoded by a nucleic acid sequence that comprises a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to heavy chain nucleic acid sequence selected from the group consisting of SEQ ID NO: 45, 47, 49, 51, 53, and 55, and a nucleic acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to light chain nucleic acid sequence comprising SEQ ID NO: 11.

[000149] In some embodiments, the MM has a dissociation constant for binding to the AB that is greater than the dissociation constant of the AB to PDL1.

[000150] In some embodiments, the MM has a dissociation constant for binding to the AB that is no more than the dissociation constant of the AB to PDL1.

[000151] In some embodiments, the MM has a dissociation constant for binding to the AB that is less than the dissociation constant of the AB to PDL1.

[000152] In some embodiments, the MM has a dissociation constant for binding to the AB that is approximately equal to the dissociation constant of the AB to PDL1.

[000153] In some embodiments, the MM does not interfere or compete with the AB for binding to PDL1 when the activatable antibody is in a cleaved state.

[000154] In some embodiments, the MM is a polypeptide of about 2 to 40 amino acids in length. In some embodiments, the MM is a polypeptide of up to about 40 amino acids in length.

[000155] In some embodiments, the MM polypeptide sequence is different from that of PDL1. In some embodiments, the MM polypeptide sequence is no more than 50% identical to any natural binding partner of the AB. In some embodiments, the MM

polypeptide sequence is different from that of PDL1 and is no more than 40%, 30%, 25%, 20%, 15%, or 10% identical to any natural binding partner of the AB.

[000156] In some embodiments, the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 59-81, 208, and 426. In some embodiments, the MM comprises an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 59-81, 208, and 426.

[000157] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind PDL1 such that the dissociation constant (K_d) of the AB when coupled to the MM towards PDL1 is at least two times greater than the K_d of the AB when not coupled to the MM towards PDL1.

[000158] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind PDL1 such that the dissociation constant (K_d) of the AB when coupled to the MM towards PDL1 is at least five times greater than the K_d of the AB when not coupled to the MM towards PDL1.

[000159] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind PDL1 such that the dissociation constant (K_d) of the AB when coupled to the MM towards PDL1 is at least 10 times greater than the K_d of the AB when not coupled to the MM towards PDL1.

[000160] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind PDL1 such that the dissociation constant (K_d) of the AB when coupled to the MM towards PDL1 is at least 20 times greater than the K_d of the AB when not coupled to the MM towards PDL1.

[000161] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind PDL1 such that the dissociation constant (K_d) of the AB when coupled to the MM towards PDL1 is at least 40 times greater than the K_d of the AB when not coupled to the MM towards PDL1.

[000162] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind PDL1 such that the dissociation constant (K_d) of the AB when coupled to the MM towards PDL1 is at least 100 times greater than the K_d of the AB when not coupled to the MM towards PDL1.

[000163] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind PDL1 such that the dissociation constant (K_d) of the AB when coupled to

the MM towards PDL1 is at least 1000 times greater than the K_d of the AB when not coupled to the MM towards PDL1.

[000164] In some embodiments, the coupling of the MM to the AB reduces the ability of the AB to bind PDL1 such that the dissociation constant (K_d) of the AB when coupled to the MM towards PDL1 is at least 10,000 times greater than the K_d of the AB when not coupled to the MM towards PDL1.

[000165] In some embodiments, in the presence of PDL1, the MM reduces the ability of the AB to bind PDL1 by at least 90% when the CM is uncleaved, as compared to when the CM is cleaved when assayed *in vitro* using a target displacement assay such as, for example, the assay described in PCT Publication No. WO 2010/081173, the contents of which are hereby incorporated by reference in their entirety.

[000166] In some embodiments, the protease that cleaves the CM is active, e.g., up-regulated, in diseased tissue, and the protease cleaves the CM in the activatable antibody when the activatable antibody is exposed to the protease.

[000167] In some embodiments, the protease is co-localized with PDL1 in a tissue, and the protease cleaves the CM in the activatable antibody when the activatable antibody is exposed to the protease.

[000168] In some embodiments, the protease is present at relatively higher levels in or in close proximity to target-containing tissue of a treatment site or diagnostic site than in tissue of non-treatment sites (for example in healthy tissue), and the protease cleaves the CM in the activatable antibody when the activatable antibody is exposed to the protease.

[000169] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to PDL1 is reduced to occur with a dissociation constant that is at least twofold greater than the dissociation constant of an unmodified AB binding to PDL1, whereas in the cleaved state (i.e., when the activatable antibody is in the cleaved state), the AB binds PDL1.

[000170] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to PDL1 is reduced to occur with a dissociation constant that is at least fivefold greater than the dissociation constant of an unmodified AB binding to PDL1, whereas in the cleaved state (i.e., when the activatable antibody is in the cleaved state), the AB binds PDL1.

[000171] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to PDL1 is reduced to occur with a dissociation constant that is at least 10-fold greater than the dissociation constant of an unmodified AB binding to PDL1, whereas in the cleaved state (i.e., when the activatable antibody is in the cleaved state), the AB binds PDL1.

[000172] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to PDL1 is reduced to occur with a dissociation constant that is at least 20-fold greater than the dissociation constant of an unmodified AB binding to PDL1, whereas in the cleaved state (i.e., when the activatable antibody is in the cleaved state), the AB binds PDL1.

[000173] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to PDL1 is reduced to occur with a dissociation constant that is at least 40-fold greater than the dissociation constant of an unmodified AB binding to PDL1, whereas in the cleaved state, the AB binds PDL1.

[000174] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to PDL1 is reduced to occur with a dissociation constant that is at least 50-fold greater than the dissociation constant of an unmodified AB binding to PDL1, whereas in the cleaved state, the AB binds PDL1.

[000175] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to PDL1 is reduced to occur with a dissociation constant that is at least 100-fold greater than the dissociation constant of an unmodified AB binding to PDL1, whereas in the cleaved state, the AB binds PDL1.

[000176] In some embodiments, the CM is positioned in the activatable antibody such that when the activatable antibody is in the uncleaved state, binding of the activatable antibody to PDL1 is reduced to occur with a dissociation constant that is at least 200-fold greater than the dissociation constant of an unmodified AB binding to PDL1, whereas in the cleaved state, the AB binds PDL1.

[000177] In some embodiments, the CM is a polypeptide of up to 15 amino acids in length.

[000178] In some embodiments, the CM is a polypeptide that includes a first cleavable moiety (CM1) that is a substrate for at least one matrix metalloprotease (MMP) and a second cleavable moiety (CM2) that is a substrate for at least one serine protease (SP). In some embodiments, each of the CM1 substrate sequence and the CM2 substrate sequence of the CM1-CM2 substrate is independently a polypeptide of up to 15 amino acids in length.

[000179] In some embodiments, the CM is a substrate for at least one protease that is or is believed to be up-regulated in cancer. In some embodiments, the CM is a substrate for at least one protease that is or is believed to be up-regulated in inflammation. In some embodiments, the CM is a substrate for at least one protease that is or is believed to be up-regulated in autoimmunity.

[000180] In some embodiments, the CM is a substrate for at least one protease selected from the group consisting of a matrix metalloprotease (MMP), a thrombin, a neutrophil elastase, a cysteine protease, a legumain, and a serine protease, such as a matriptase (MT-SP1), and a urokinase (uPA). Without being bound by theory, it is believed that these proteases are up-regulated in at least one of cancer, inflammation, and/or autoimmunity.

[000181] Exemplary substrates include but are not limited to substrates cleavable by one or more of the following enzymes or proteases listed in Table 12.

[000182] In some embodiments, the CM is selected for use with a specific protease, for example a protease that is known to be co-localized with the target of the activatable antibody.

[000183] In some embodiments, the CM is selected for use with a specific protease, for example a protease that is known to be in close proximity to the target of the activatable antibody.

[000184] In some embodiments, the CM is a substrate for at least one MMP. Examples of MMPs include the MMPs listed in the Table 12. In some embodiments, the CM is a substrate for a protease selected from the group consisting of MMP 9, MMP14, MMP1, MMP3, MMP13, MMP17, MMP11, and MMP19. In some embodiments the CM is a substrate for MMP9. In some embodiments, the CM is a substrate for MMP14.

[000185] In some embodiments, the CM is a substrate that includes the sequence TGRGPSWV (SEQ ID NO: 338); SARGPSRW (SEQ ID NO: 339); TARGPSFK (SEQ ID NO: 340); LSGRSDNH (SEQ ID NO: 341); GGWHTGRN (SEQ ID NO: 342);

HTGRSGAL (SEQ ID NO: 343); PLTGRSGG (SEQ ID NO: 344); AARGPAIH (SEQ ID NO: 345); RGPAFNPM (SEQ ID NO: 346); SSRGPAYL (SEQ ID NO: 347); RGPATPIM (SEQ ID NO: 348); RGPA (SEQ ID NO: 349); GGQPSGMWGW (SEQ ID NO: 350); FPRPLGITGL (SEQ ID NO: 351); VHMPLGFLGP (SEQ ID NO: 352); SPLTGRSG (SEQ ID NO: 353); SAGFSLPA (SEQ ID NO: 354); LAPLGLQRR (SEQ ID NO: 355); SGGPLGVR (SEQ ID NO: 356); PLGL (SEQ ID NO: 357); LSGRSGNH (SEQ ID NO: 883); SGRSANPRG (SEQ ID NO: 884); LSGRSDDH (SEQ ID NO: 885); LSGRSDIH (SEQ ID NO: 886); LSGRSDQH (SEQ ID NO: 887); LSGRSDTH (SEQ ID NO: 888); LSGRSDYH (SEQ ID NO: 889); LSGRSDNP (SEQ ID NO: 890); LSGRSANP (SEQ ID NO: 891); LSGRSANI (SEQ ID NO: 892); and/or LSGRSDNI (SEQ ID NO: 893).

[000186] In some embodiments, the CM comprises the amino acid sequence LSGRSDNH (SEQ ID NO: 341). In some embodiments, the CM comprises the amino acid sequence TGRGPSWV (SEQ ID NO: 338). In some embodiments, the CM comprises the amino acid sequence PLTGRSGG (SEQ ID NO: 344). In some embodiments, the CM comprises the amino acid sequence GGQPSGMWGW (SEQ ID NO: 350). In some embodiments, the CM comprises the amino acid sequence FPRPLGITGL (SEQ ID NO: 351). In some embodiments, the CM comprises the amino acid sequence VHMPLGFLGP (SEQ ID NO: 352). In some embodiments, the CM comprises the amino acid sequence PLGL (SEQ ID NO: 357). In some embodiments, the CM comprises the amino acid sequence SARGPSRW (SEQ ID NO: 339). In some embodiments, the CM comprises the amino acid sequence TARGPSFK (SEQ ID NO: 340). In some embodiments, the CM comprises the amino acid sequence GGWHTGRN (SEQ ID NO: 342). In some embodiments, the CM comprises the amino acid sequence HTGRSGAL (SEQ ID NO: 343). In some embodiments, the CM comprises the amino acid sequence AARGPAIH (SEQ ID NO: 345). In some embodiments, the CM comprises the amino acid sequence RGPAFNPM (SEQ ID NO: 346). In some embodiments, the CM comprises the amino acid sequence SSRGPAYL (SEQ ID NO: 347). In some embodiments, the CM comprises the amino acid sequence RGPATPIM (SEQ ID NO: 348). In some embodiments, the CM comprises the amino acid sequence RGPA (SEQ ID NO: 349). In some embodiments, the CM comprises the amino acid sequence LSGRSGNH (SEQ ID NO: 883). In some embodiments, the CM comprises the amino acid sequence SGRSANPRG (SEQ ID NO: 884). In some embodiments, the CM comprises the amino acid sequence LSGRSDDH (SEQ ID NO: 885).

In some embodiments, the CM comprises the amino acid sequence LSGRSDIH (SEQ ID NO: 886). In some embodiments, the CM comprises the amino acid sequence LSGRSDQH (SEQ ID NO: 887). In some embodiments, the CM comprises the amino acid sequence LSGRSDTH (SEQ ID NO: 888). In some embodiments, the CM comprises the amino acid sequence LSGRSDYH (SEQ ID NO: 889). In some embodiments, the CM comprises the amino acid sequence LSGRSDNP (SEQ ID NO: 890). In some embodiments, the CM comprises the amino acid sequence LSGRSANP (SEQ ID NO: 891). In some embodiments, the CM comprises the amino acid sequence LSGRSANI (SEQ ID NO: 892). In some embodiments, the CM comprises the amino acid sequence LSGRSDNI (SEQ ID NO: 893)

[000187] In some embodiments, the CM is a substrate for an MMP and includes the sequence ISSGLSS (SEQ ID NO: 358); QNQALRMA (SEQ ID NO: 359); AQNLLGMV (SEQ ID NO: 360); STFPFGMF (SEQ ID NO: 361); PVGYTSSL (SEQ ID NO: 362); DWLYWPGI (SEQ ID NO: 363), ISSGLSS (SEQ ID NO: 364), LKAAPRWA (SEQ ID NO: 365); GPSHLVLT (SEQ ID NO: 366); LPGGLSPW (SEQ ID NO: 367); MGLFSEAG (SEQ ID NO: 368); SPLPLRVP (SEQ ID NO: 369); RMHLRSLG (SEQ ID NO: 370); LAAPLGLL (SEQ ID NO: 371); AVGLLAPP (SEQ ID NO: 372); LLAPSHRA (SEQ ID NO: 373); PAGLWLDP (SEQ ID NO: 374); MIAPVAYR (SEQ ID NO: 894); RPSPMWAY (SEQ ID NO: 895); WATPRPMR (SEQ ID NO: 896); FRLLDWQW (SEQ ID NO: 897); ISSGL (SEQ ID NO: 898); ISSGLS (SEQ ID NO: 899); and/or ISSGLL (SEQ ID NO: 900).

[000188] In some embodiments, the CM comprises the amino acid sequence ISSGLSS (SEQ ID NO: 358). In some embodiments, the CM comprises the amino acid sequence QNQALRMA (SEQ ID NO: 359). In some embodiments, the CM comprises the amino acid sequence AQNLLGMV (SEQ ID NO: 360). In some embodiments, the CM comprises the amino acid sequence STFPFGMF (SEQ ID NO: 361). In some embodiments, the CM comprises the amino acid sequence PVGYTSSL (SEQ ID NO: 362). In some embodiments, the CM comprises the amino acid sequence DWLYWPGI (SEQ ID NO: 363). In some embodiments, the CM comprises the amino acid sequence ISSGLSS (SEQ ID NO: 364). In some embodiments, the CM comprises the amino acid sequence LKAAPRWA (SEQ ID NO: 365). In some embodiments, the CM comprises the amino acid sequence GPSHLVLT (SEQ ID NO: 366). In some embodiments, the CM comprises the amino acid sequence LPGGLSPW (SEQ ID NO: 367). In some embodiments, the CM comprises the amino acid sequence MGLFSEAG (SEQ ID NO: 368). In some embodiments, the CM comprises the

amino acid sequence SPLPLRVP (SEQ ID NO: 369). In some embodiments, the CM comprises the amino acid sequence RMHLRSLG (SEQ ID NO: 370). In some embodiments, the CM comprises the amino acid sequence LAAPLGLL (SEQ ID NO: 371). In some embodiments, the CM comprises the amino acid sequence AVGLLAPP (SEQ ID NO: 372). In some embodiments, the CM comprises the amino acid sequence LLAPSHRA (SEQ ID NO: 373). In some embodiments, the CM comprises the amino acid sequence PAGLWLDP (SEQ ID NO: 374). In some embodiments, the CM comprises the amino acid sequence MIAPVAYR (SEQ ID NO: 894). In some embodiments, the CM comprises the amino acid sequence RPSPMWAY (SEQ ID NO: 895). In some embodiments, the CM comprises the amino acid sequence WATPRPMR (SEQ ID NO: 896). In some embodiments, the CM comprises the amino acid sequence FRLLDWQW (SEQ ID NO: 897). In some embodiments, the CM comprises the amino acid sequence ISSGL (SEQ ID NO: 898). In some embodiments, the CM comprises the amino acid sequence ISSGLLS (SEQ ID NO: 899). In some embodiments, the CM comprises the amino acid sequence ISSGLL (SEQ ID NO: 900).

[000189] In some embodiments, the CM is a substrate for thrombin. In some embodiments, the CM is a substrate for thrombin and includes the sequence GPRSFG (SEQ ID NO: 375) or GPRSFG (SEQ ID NO: 376). In some embodiments, the CM comprises the amino acid sequence GPRSFG (SEQ ID NO: 375). In some embodiments, the CM comprises the amino acid sequence GPRSFG (SEQ ID NO: 376).

[000190] In some embodiments, the CM comprises an amino acid sequence selected from the group consisting of NTLSGRSENHSG (SEQ ID NO: 435); NTLSGRSGNHGS (SEQ ID NO: 436); TSTSGRSANPRG (SEQ ID NO: 437); TSGRSANP (SEQ ID NO: 438); VAGRSMRP (SEQ ID NO: 439); VVPEGRRS (SEQ ID NO: 440); ILPRSPAF (SEQ ID NO: 441); MVLGRSLL (SEQ ID NO: 442); QGRAITFI (SEQ ID NO: 443); SPRSIMLA (SEQ ID NO: 444); and SMLRSMPL (SEQ ID NO: 445).

[000191] In some embodiments, the CM comprises the amino acid sequence NTLSGRSENHSG (SEQ ID NO: 435). In some embodiments, the CM comprises the amino acid sequence NTLSGRSGNHGS (SEQ ID NO: 436). In some embodiments, the CM comprises the amino acid sequence TSTSGRSANPRG (SEQ ID NO: 437). In some embodiments, the CM comprises the amino acid sequence TSGRSANP (SEQ ID NO: 438). In some embodiments, the CM comprises the amino acid sequence VAGRSMRP (SEQ ID NO: 439). In some embodiments, the CM comprises the amino acid sequence VVPEGRRS

(SEQ ID NO: 440). In some embodiments, the CM comprises the amino acid sequence ILPRSPAF (SEQ ID NO: 441). In some embodiments, the CM comprises the amino acid sequence MVLGRSLL (SEQ ID NO: 442). In some embodiments, the CM comprises the amino acid sequence QGRAITFI (SEQ ID NO: 443). In some embodiments, the CM comprises the amino acid sequence SPRSIMLA (SEQ ID NO: 444). In some embodiments, the CM comprises the amino acid sequence SMLRSMPL (SEQ ID NO: 445).

[000192] In some embodiments, the CM is a substrate for a neutrophil elastase. In some embodiments, the CM is a substrate for a serine protease. In some embodiments, the CM is a substrate for uPA. In some embodiments, the CM is a substrate for legumain. In some embodiments, the CM is a substrate for matriptase. In some embodiments, the CM is a substrate for a cysteine protease. In some embodiments, the CM is a substrate for a cysteine protease, such as a cathepsin.

[000193] In some embodiments, the CM is a CM1-CM2 substrate and includes the sequence ISSGLLSGRSDNH (SEQ ID NO: 377); ISSGLLSSGGSGGSLSGRSDNH (SEQ ID NO: 378); AVGLLAPPGGTSTSGRSANPRG (SEQ ID NO: 379); TSTSGRSANPRGGGAVGLLAPP (SEQ ID NO: 380); VHMP LGFLGP GGTSTSGRSANPRG (SEQ ID NO: 381); TSTSGRSANPRGGGVHMP LGFLGP (SEQ ID NO: 382); AVGLLAPPGGLSGRSDNH (SEQ ID NO: 383); LSGRSDNHGGAVGLLAPP (SEQ ID NO: 384); VHMP LGFLGP GGLSGRSDNH (SEQ ID NO: 385); LSGRSDNHGGVHMP LGFLGP (SEQ ID NO: 386); LSGRSDNHGGSGGSISSGLLSS (SEQ ID NO: 387); LSGRSGNHGGSGGSISSGLLSS (SEQ ID NO: 388); ISSGLLSSGGSGGSLSGRSGNH (SEQ ID NO: 389); LSGRSDNHGGSGGSGQNQALRMA (SEQ ID NO: 390); QNQALRMAGGSGGSLSGRSDNH (SEQ ID NO: 391); LSGRSGNHGGSGGSGQNQALRMA (SEQ ID NO: 392); QNQALRMAGGSGGSLSGRSGNH (SEQ ID NO: 393); ISSGLLSGRSGNH (SEQ ID NO: 394); ISSGLLSGRSANPRG (SEQ ID NO: 901); AVGLLAPPTSGRSANPRG (SEQ ID NO: 902); AVGLLAPPSGRSANPRG (SEQ ID NO: 903); ISSGLLSGRSDDH (SEQ ID NO: 904); ISSGLLSGRSDIH (SEQ ID NO: 905); ISSGLLSGRSDQH (SEQ ID NO: 906); ISSGLLSGRSDTH (SEQ ID NO: 907); ISSGLLSGRSDYH (SEQ ID NO: 908); ISSGLLSGRSDNP (SEQ ID NO: 909); ISSGLLSGRSANP (SEQ ID NO: 910); ISSGLLSGRSANI (SEQ ID NO: 911); AVGLLAPPGGLSGRSDDH (SEQ ID NO: 912); AVGLLAPPGGLSGRSDIH (SEQ ID NO: 913); AVGLLAPPGGLSGRSDQH (SEQ ID

NO: 914); AVGLLAPPGGLSGRSDTH (SEQ ID NO: 915); AVGLLAPPGGLSGRSDYH (SEQ ID NO: 916); AVGLLAPPGGLSGRSDNP (SEQ ID NO: 917); AVGLLAPPGGLSGRSANP (SEQ ID NO: 918); AVGLLAPPGGLSGRSANI (SEQ ID NO: 919); ISSGLLSGRSDNI (SEQ ID NO: 920); AVGLLAPPGGLSGRSDNI (SEQ ID NO: 921); GLSGRSDNHGGAVGLLAPP (SEQ ID NO: 1009); and/or GLSGRSDNHGGVHMPLGFLGP (SEQ ID NO: 1010).

[000194] In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDNH (SEQ ID NO: 377), which is also referred to herein as substrate 2001. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSSGGSGGSLSGRSDNH (SEQ ID NO: 378), which is also referred to herein as substrate 1001/LP'/0001, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 922). In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGTSTSGRSANPRG (SEQ ID NO: 379), which is also referred to herein as substrate 2015 and/or substrate 1004/LP'/0003, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence TSTSGRSANPRGGGAVGLLAPP (SEQ ID NO: 380), which is also referred to herein as substrate 0003/LP'/1004, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence VHMPLGFLGPGGTSTSGRSANPRG (SEQ ID NO: 381), which is also referred to herein as substrate 1003/LP'/0003, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence TSTSGRSANPRGGGVHMPLGFLGP (SEQ ID NO: 382), which is also referred to herein as substrate 0003/LP'/1003, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDNH (SEQ ID NO: 383), which is also referred to herein as substrate 3001 and/or substrate 1004/LP'/0001, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence LSGRSDNHGGAVGLLAPP (SEQ ID NO: 384), which is also referred to herein as substrate 0001/LP'/1004, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence VHMPLGFLGPGLSGRSDNH (SEQ ID NO: 385), which is also referred to herein as substrate 1003/LP'/0001, wherein LP' as used in this

CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence LSGRSDNHGGVHMP LGFLGP (SEQ ID NO: 386), which is also referred to herein as substrate 0001/LP'/1003, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence LSGRSDNHGGSGGSISSG LLSS (SEQ ID NO: 387), which is also referred to herein as substrate 0001/LP'/1001, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 922). In some embodiments, the CM1-CM2 substrate includes the sequence LSGRSGNHGGSGGSISSG LLSS (SEQ ID NO: 388), which is also referred to herein as substrate 0002/LP'/1001, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 922). In some embodiments, the CM1-CM2 substrate includes the sequence ISSG LLSSGGSGGSLSGRSGNH (SEQ ID NO: 389), which is also referred to herein as substrate 1001/LP'/0002, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 922). In some embodiments, the CM1-CM2 substrate includes the sequence LSGRSDNHGGSGG SQNQALRMA (SEQ ID NO: 390), which is also referred to herein as substrate 0001/LP'/1002, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 922). In some embodiments, the CM1-CM2 substrate includes the sequence QNQALRMAGGSGGSLSGRSDNH (SEQ ID NO: 391), which is also referred to herein as substrate 1002/LP'/0001, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 922). In some embodiments, the CM1-CM2 substrate includes the sequence LSGRSGNHGGSGG SQNQALRMA (SEQ ID NO: 392), which is also referred to herein as substrate 0002/LP'/1002, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 922). In some embodiments, the CM1-CM2 substrate includes the sequence QNQALRMAGGSGGSLSGRSGNH (SEQ ID NO: 393), which is also referred to herein as substrate 1002/LP'/0002, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GGSGGS (SEQ ID NO: 922). In some embodiments, the CM1-CM2 substrate includes the sequence ISSG LLSGRSGNH (SEQ ID NO: 394), which is also referred to herein as substrate 2002. In some embodiments, the CM1-CM2 substrate includes the sequence ISSG LLSGRSANPRG (SEQ ID NO: 901), which is also referred to herein as substrate 2003. In some embodiments, the CM1-CM2 substrate includes the sequence AVG LLAPPTSGRSANPRG (SEQ ID NO: 902), which is also

referred to herein as substrate 2004. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPSGRSANPRG (SEQ ID NO: 903), which is also referred to herein as substrate 2005. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDDH (SEQ ID NO: 904), which is also referred to herein as substrate 2006. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDIH (SEQ ID NO: 905), which is also referred to herein as substrate 2007. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDQH (SEQ ID NO: 906), which is also referred to herein as substrate 2008. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDTH (SEQ ID NO: 907), which is also referred to herein as substrate 2009. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDYH (SEQ ID NO: 908), which is also referred to herein as substrate 2010. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDNP (SEQ ID NO: 909), which is also referred to herein as substrate 2011. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSANP (SEQ ID NO: 910), which is also referred to herein as substrate 2012. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSANI (SEQ ID NO: 911), which is also referred to herein as substrate 2013. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDDH (SEQ ID NO: 912), which is also referred to herein as substrate 3006. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDIH (SEQ ID NO: 913), which is also referred to herein as substrate 3007. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDQH (SEQ ID NO: 914), which is also referred to herein as substrate 3008. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDTH (SEQ ID NO: 915), which is also referred to herein as substrate 3009. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDYH (SEQ ID NO: 916), which is also referred to herein as substrate 3010. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDNP (SEQ ID NO: 917), which is also referred to herein as substrate 3011. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSANP (SEQ ID NO: 918), which is also referred to herein as substrate 3012. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSANI (SEQ ID NO: 919), which is also referred to herein as

substrate 3013. In some embodiments, the CM1-CM2 substrate includes the sequence ISSGLLSGRSDNI (SEQ ID NO: 920), which is also referred to herein as substrate 2014. In some embodiments, the CM1-CM2 substrate includes the sequence AVGLLAPPGGLSGRSDNI (SEQ ID NO: 921), which is also referred to herein as substrate 3014. In some embodiments, the CM1-CM2 substrate includes the sequence GLSGRSDNHGGAVGLLAPP (SEQ ID NO: 1009), which is also referred to herein as substrate 0001/LP'/1004, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG. In some embodiments, the CM1-CM2 substrate includes the sequence GLSGRSDNHGGVHMPLGFLGP (SEQ ID NO: 1010), which is also referred to herein as substrate 0001/LP'/1003, where LP' as used in this CM1-CM2 substrate is the amino acid sequence GG.

[000195] In some embodiments, the CM is a substrate for at least two proteases. In some embodiments, each protease is selected from the group consisting of those shown in Table 12. In some embodiments, the CM is a substrate for at least two proteases, wherein one of the proteases is selected from the group consisting of a MMP, a thrombin, a neutrophil elastase, a cysteine protease, a uPA, a legumain and a matriptase and other serine proteases and the other protease is selected from the group consisting of those shown in Table 12. In some embodiments, the CM is a substrate for at least two proteases selected from the group consisting of a MMP, a thrombin, a neutrophil elastase, a cysteine protease, a uPA, a legumain and a matriptase and other serine proteases.

[000196] In some embodiments, the activatable antibody includes at least a first CM and a second CM. In some embodiments, the first CM and the second CM are each polypeptides of no more than 15 amino acids long. In some embodiments, the first CM and the second CM in the activatable antibody in the uncleaved state have the structural arrangement from N-terminus to C-terminus as follows: MM-CM1-CM2-AB or AB-CM2-CM1-MM. In some embodiments, at least one of the first CM and the second CM is a polypeptide that functions as a substrate for a protease selected from the group consisting of a MMP, a thrombin, a neutrophil elastase, a cysteine protease, a uPA, a legumain, and a matriptase and other serine proteases. In some embodiments, the first CM is cleaved by a first cleaving agent selected from the group consisting of a MMP, a thrombin, a neutrophil elastase, a cysteine protease, a uPA, a legumain, and a matriptase and other serine proteases in a target tissue and the second CM is cleaved by a second cleaving agent in a target tissue. In some embodiments, the other protease is selected from the group consisting of those

shown in Table 12. In some embodiments, the first cleaving agent and the second cleaving agent are the same protease selected from the group consisting of a MMP, a thrombin, a neutrophil elastase, a cysteine protease, a uPA, a legumain, and a matriptase and other serine proteases, and the first CM and the second CM are different substrates for the enzyme. In some embodiments, the first cleaving agent and the second cleaving agent are the same protease selected from the group consisting of those shown in Table 12. In some embodiments, the first cleaving agent and the second cleaving agent are different proteases. In some embodiments, the first cleaving agent and the second cleaving agent are co-localized in the target tissue. In some embodiments, the first CM and the second CM are cleaved by at least one cleaving agent in the target tissue.

[000197] In some embodiments, the activatable antibody is exposed to and cleaved by a protease such that, in the activated or cleaved state, the activated antibody includes a light chain amino acid sequence that includes at least a portion of LP2 and/or CM sequence after the protease has cleaved the CM.

[000198] In some embodiments, the activatable antibody also includes an agent conjugated to the AB. In some embodiments, the agent conjugated to the AB or the AB of an activatable antibody is a therapeutic agent. In some embodiments, the agent is an antineoplastic agent. In some embodiments, the agent is a toxin or fragment thereof. As used herein, a fragment of a toxin is a fragment that retains toxic activity. In some embodiments, the agent is conjugated to the AB via a cleavable linker. In some embodiments, the agent is conjugated to the AB via a linker that includes at least one CM1-CM2 substrate sequence. In some embodiments, the agent is conjugated to the AB via a noncleavable linker. In some embodiments, the agent is conjugated to the AB via a linker that is cleavable in an intracellular or lysosomal environment. In some embodiments, the agent is a microtubule inhibitor. In some embodiments, the agent is a nucleic acid damaging agent, such as a DNA alkylator, a DNA cleaving agent, a DNA cross-linker, or DNA intercalator, or other DNA damaging agent. In some embodiments, the agent is an agent selected from the group listed in Table 11. In some embodiments, the agent is a dolastatin. In some embodiments, the agent is an auristatin or derivative thereof. In some embodiments, the agent is auristatin E or a derivative thereof. In some embodiments, the agent is monomethyl auristatin E (MMAE). In some embodiments, the agent is monomethyl auristatin D (MMAD). In some embodiments, the agent is a maytansinoid or maytansinoid derivative. In some embodiments, the agent is DM1 or DM4. In some embodiments, the

agent is a duocarmycin or derivative thereof. In some embodiments, the agent is a calicheamicin or derivative thereof. In some embodiments, the agent is a pyrrolobenzodiazepine. In some embodiments, the agent is a pyrrolobenzodiazepine dimer.

[000199] In some embodiments, the activatable antibody is conjugated to one or more equivalents of an agent. In some embodiments, the activatable antibody is conjugated to one equivalent of the agent. In some embodiments, the activatable antibody is conjugated to two, three, four, five, six, seven, eight, nine, ten, or greater than ten equivalents of the agent. In some embodiments, the activatable antibody is part of a mixture of activatable antibodies having a homogeneous number of equivalents of conjugated agents. In some embodiments, the activatable antibody is part of a mixture of activatable antibodies having a heterogeneous number of equivalents of conjugated agents. In some embodiments, the mixture of activatable antibodies is such that the average number of agents conjugated to each activatable antibody is between zero to one, between one to two, between two and three, between three and four, between four and five, between five and six, between six and seven, between seven and eight, between eight and nine, between nine and ten, and ten and greater. In some embodiments, the mixture of activatable antibodies is such that the average number of agents conjugated to each activatable antibody is one, two, three, four, five, six, seven, eight, nine, ten, or greater.

[000200] In some embodiments, the activatable antibody comprises one or more site-specific amino acid sequence modifications such that the number of lysine and/or cysteine residues is increased or decreased with respect to the original amino acid sequence of the activatable antibody, thus in some embodiments correspondingly increasing or decreasing the number of agents that can be conjugated to the activatable antibody, or in some embodiments limiting the conjugation of the agents to the activatable antibody in a site-specific manner. In some embodiments, the modified activatable antibody is modified with one or more non-natural amino acids in a site-specific manner, thus in some embodiments limiting the conjugation of the agents to only the sites of the non-natural amino acids.

[000201] In some embodiments, the agent is an anti-inflammatory agent.

[000202] In some embodiments, the activatable antibody also includes a detectable moiety. In some embodiments, the detectable moiety is a diagnostic agent.

[000203] In some embodiments, the activatable antibody also includes a signal peptide. In some embodiments, the signal peptide is conjugated to the activatable antibody via a spacer. In some embodiments, the spacer is conjugated to the activatable antibody in

the absence of a signal peptide. In some embodiments, the spacer is joined directly to the MM of the activatable antibody. In some embodiments, the spacer is joined directly to the MM of the activatable antibody in the structural arrangement from N-terminus to C-terminus of spacer-MM-CM-AB. An example of a spacer joined directly to the N-terminus of MM of the activatable antibody is selected from the group consisting of QGQSGS (SEQ ID NO: 923); GQSGS (SEQ ID NO: 1192); QSGS (SEQ ID NO: 1193); SGS (SEQ ID NO: 1194); GS (SEQ ID NO: 1195); S; QGQSGQG (SEQ ID NO: 924); GQSGQG (SEQ ID NO: 395); QSGQG (SEQ ID NO: 925); SGQG (SEQ ID NO: 926); GQG (SEQ ID NO: 927); QG (SEQ ID NO: 928); G; QGQSGQ (SEQ ID NO: 1196); GQSGQ (SEQ ID NO: 1197); QSGQ (SEQ ID NO: 616); SGQ (SEQ ID NO: 1198); GQ (SEQ ID NO: 1199); and Q. In some embodiments, the spacer includes at least the amino acid sequence QGQSGS (SEQ ID NO: 923). In some embodiments, the spacer includes at least the amino acid sequence GQSGS (SEQ ID NO: 1192). In some embodiments, the spacer includes at least the amino acid sequence QSGS (SEQ ID NO: 1193). In some embodiments, the spacer includes at least the amino acid sequence SGS (SEQ ID NO: 1194). In some embodiments, the spacer includes at least the amino acid sequence GS (SEQ ID NO: 1195). In some embodiments, the spacer includes at least the amino acid sequence S. In some embodiments, the spacer includes at least the amino acid sequence QGQSGQG (SEQ ID NO: 924). In some embodiments, the spacer includes at least the amino acid sequence GQSGQG (SEQ ID NO: 395). In some embodiments, the spacer includes at least the amino acid sequence QSGQG (SEQ ID NO: 925). In some embodiments, the spacer includes at least the amino acid sequence SGQG (SEQ ID NO: 926). In some embodiments, the spacer includes at least the amino acid sequence GQG (SEQ ID NO: 927). In some embodiments, the spacer includes at least the amino acid sequence QG (SEQ ID NO: 928). In some embodiments, the spacer includes at least the amino acid sequence G. In some embodiments, the spacer includes at least the amino acid sequence QGQSGQ (SEQ ID NO: 1196). In some embodiments, the spacer includes at least the amino acid sequence GQSGQ (SEQ ID NO: 1197). In some embodiments, the spacer includes at least the amino acid sequence QSGQ (SEQ ID NO: 616). In some embodiments, the spacer includes at least the amino acid sequence SGQ (SEQ ID NO: 1198). In some embodiments, the spacer includes at least the amino acid sequence GQ (SEQ ID NO: 1199). In some embodiments, the spacer includes at least the amino acid sequence Q. In some embodiments, the activatable antibody does not include a spacer sequence.

[000204] In some embodiments, the AB of the activatable antibody naturally contains one or more disulfide bonds. In some embodiments, the AB can be engineered to include one or more disulfide bonds.

[000205] In some embodiments, the serum half-life of the activatable antibody is longer than that of the corresponding antibody; e.g., the pK of the activatable antibody is longer than that of the corresponding antibody. In some embodiments, the serum half-life of the activatable antibody is similar to that of the corresponding antibody. In some embodiments, the serum half-life of the activatable antibody is at least 15 days when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 12 days when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 11 days when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 10 days when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 9 days when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 8 days when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 7 days when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 6 days when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 5 days when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 4 days when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 3 days when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 2 days when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 24 hours when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 20 hours when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 18 hours when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 16 hours when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 14 hours when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 12 hours when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least

10 hours when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 8 hours when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 6 hours when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 4 hours when administered to an organism. In some embodiments, the serum half-life of the activatable antibody is at least 3 hours when administered to an organism.

[000206] In some embodiments, the activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody is monospecific. In some embodiments, the activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody is multispecific, *e.g.*, by way of non-limiting example, bispecific or trifunctional. In some embodiments, the activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody is formulated as part of a pro-Bispecific T Cell Engager (BITE) molecule, *i.e.*, the BITE includes a masking moiety and a cleavable moiety. In some embodiments, the activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody is formulated as part of a pro-Chimeric Antigen Receptor (CAR) -modified T cell, -modified NK cell, or other -modified immune effector cell. In some embodiments, an activatable anti-PDL1 antibody is formulated as part of another engineered receptor on an immune effector cell; *i.e.*, the pro-CAR or other pro-engineered receptor includes a masking moiety and a cleavable moiety.

[000207] In some embodiments, the activatable antibody or antigen-binding fragment thereof is incorporated in a multispecific activatable antibody or antigen-binding fragment thereof, where at least one arm of the multispecific activatable antibody specifically binds PDL1. In some embodiments, the activatable antibody or antigen-binding fragment thereof is incorporated in a bispecific antibody or antigen-binding fragment thereof, where at least one arm of the bispecific activatable antibody specifically binds PDL1.

[000208] In some embodiments, at least one arm of the multispecific activatable antibody, *e.g.*, a bispecific activatable antibody, comprises a heavy chain amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56. In some embodiments, at least one arm of the multispecific activatable antibody, *e.g.*, a bispecific activatable antibody, comprises a light chain amino acid sequence comprising SEQ ID NO: 12 or SEQ ID NO: 58. In some embodiments, at least one arm of the multispecific activatable antibody, *e.g.*, a bispecific activatable antibody, comprises a light chain amino acid sequence

comprising SEQ ID NO: 12. In some embodiments, at least one arm of the multispecific activatable antibody, *e.g.*, a bispecific activatable antibody, comprises a light chain amino acid sequence comprising SEQ ID NO: 58.

[000209] In some embodiments, at least one arm of the multispecific activatable antibody, *e.g.*, a bispecific activatable antibody, comprises a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid sequence comprising SEQ ID NO: 58.

[000210] In some embodiments, at least one arm of the multispecific activatable antibody, *e.g.*, a bispecific activatable antibody, comprises a heavy chain amino acid sequence comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid sequence comprising SEQ ID NO: 12.

[000211] In some embodiments, at least one arm of the multispecific activatable antibody, *e.g.*, a bispecific activatable antibody, comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56. In some embodiments, at least one arm of the multispecific activatable antibody, *e.g.*, a bispecific activatable antibody, comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, at least one arm of the multispecific activatable antibody, *e.g.*, a bispecific activatable antibody, comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12. In some embodiments, at least one arm of the multispecific activatable antibody, *e.g.*, a bispecific activatable antibody, comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, at least one arm of the multispecific activatable antibody, *e.g.*, a bispecific activatable antibody, comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 46, 48, 50, 52, 54, and 56,

and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12.

[000212] In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises the heavy chain amino acid sequence of SEQ ID NO: 46 and the light chain amino acid sequence of SEQ ID NO: 58. In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises the heavy chain amino acid sequence of SEQ ID NO: 46 and the light chain amino acid sequence of SEQ ID NO: 12.

[000213] In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid of SEQ ID NO: 46. In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12. In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence selected of SEQ ID NO: 46, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 58. In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a heavy chain amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence selected of SEQ ID NO: 46, and a light chain amino acid that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 12.

[000214] In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH

CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3) sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2) sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence shown in Table 16; a VH CDR2 sequence shown in Table 16; a VH CDR3 sequence shown in Table 16; a VL CDR1 sequence shown in Table 16; a VL CDR2 sequence shown in Table 16; and a VL CDR3 sequence shown in Table 16.

[000215] In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR1 sequence shown in Table 16; a VH CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR2 sequence shown in Table 16; a VH CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VH CDR3 sequence shown in Table 16; a VL CDR1 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR1 sequence shown in Table 16; a VL CDR2 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR2 sequence shown in Table 16; and a VL CDR3 sequence that includes a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to a VL CDR3 sequence shown in Table 16.

[000216] In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 16.

[000217] In some embodiments, the antibody comprises a heavy chain that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein the combination is a combination of the three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 16.

[000218] In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 16.

[000219] In some embodiments, the antibody comprises a heavy chain that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein each CDR sequence in the combination comprises a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the corresponding CDR sequence in a combination of three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 16.

[000220] In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein at least one CDR sequence is selected from the group consisting of a VL CDR1 sequence comprising RASQSISSYLN (SEQ ID NO: 209); a VL CDR2 sequence comprising AASSLQS (SEQ ID NO: 215); a VL CDR3 sequence comprising DNGYPST (SEQ ID NO: 228); a VH CDR1 sequence comprising SYAMS (SEQ ID NO: 212); a VH CDR2 sequence comprising SSIWRNGIVTVYADS (SEQ ID NO: 246); and a VH CDR3 sequence comprising WSAAFDY (SEQ ID NO: 235).

[000221] In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR2 sequence comprises SSIWRNGIVTVYADS (SEQ ID NO: 246). In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a

combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR3 sequence comprises WSAAFDY (SEQ ID NO: 235). In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the VH CDR2 sequence comprises SSIWRNGIVTVYADS (SEQ ID NO: 246), and the VH CDR3 sequence comprises WSAAFDY (SEQ ID NO: 235).

[000222] In some embodiments, at least one arm of the multispecific activatable antibody, e.g., a bispecific activatable antibody, comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination comprises a VL CDR1 sequence comprising RASQSISSYLN (SEQ ID NO: 209); a VL CDR2 sequence comprising AASSLQS (SEQ ID NO: 215); a VL CDR3 sequence comprising DNGYPST (SEQ ID NO: 228); a VH CDR1 sequence comprising SYAMS (SEQ ID NO: 212); a VH CDR2 sequence comprising SSIWRNGIVTVYADS (SEQ ID NO: 246); and a VH CDR3 sequence comprising WSAAFDY (SEQ ID NO: 235).

[000223] In some embodiments, the anti-PDL1 antibodies, conjugated anti-PDL1 antibodies, activatable anti-PDL1 antibodies and/or conjugated activatable anti-PDL1 antibodies described herein are used as sole active agents. In some embodiments, the anti-PDL1 antibodies, conjugated anti-PDL1 antibodies, activatable anti-PDL1 antibodies and/or conjugated activatable anti-PDL1 antibodies described herein are used in conjunction with one or more additional agents or a combination of additional agents. Suitable additional agents include current pharmaceutical and/or surgical therapies for an intended application, such as, for example, cancer. For example, the anti-PDL1 antibodies, conjugated anti-PDL1 antibodies, activatable anti-PDL1 antibodies and/or conjugated activatable anti-PDL1 antibodies can be used in conjunction with an additional chemotherapeutic or anti-neoplastic agent.

[000224] In some embodiments, the additional agent(s) is a chemotherapeutic agent, such as a chemotherapeutic agent selected from the group consisting of docetaxel, paclitaxel, abraxane (i.e., albumin-conjugated paclitaxel), doxorubicin, oxaliplatin, carboplatin, cisplatin, irinotecan, and gemcitabine.

[000225] In some embodiments, the additional agent(s) is a checkpoint inhibitor, a kinase inhibitor, an agent targeting inhibitors in the tumor microenvironment, and/or a T cell or NK agonist. In some embodiments, the additional agent(s) is radiation therapy, alone or in combination with another additional agent(s) such as a chemotherapeutic or anti-neoplastic agent. In some embodiments, the additional agent(s) is a vaccine, an oncovirus, and/or a DC-activating agent such as, by way of non-limiting example, a toll-like receptor (TLR) agonist and/or α -CD40. In some embodiments, the additional agent(s) is a tumor-targeted antibody designed to kill the tumor via ADCC or via direct conjugation to a toxin (e.g., an antibody drug conjugate (ADC)).

[000226] In some embodiments, the checkpoint inhibitor is an inhibitor of a target selected from the group consisting of CTLA-4, LAG-3, PD1 (also referred to as PD-1), PDL1, TIGIT, TIM-3, B7H4, and Vista. In some embodiments, the kinase inhibitor is selected from the group consisting of B-RAFi, MEKi, and Btk inhibitors, such as ibrutinib. In some embodiments, the kinase inhibitor is crizotinib. In some embodiments, the tumor microenvironment inhibitor is selected from the group consisting of an IDO inhibitor, an α -CSF1R inhibitor, an α -CCR4 inhibitor, a TGF-beta, a myeloid-derived suppressor cell, or a T-regulatory cell. In some embodiments, the agonist is selected from the group consisting of Ox40, GITR, CD137, ICOS, CD27, and HVEM.

[000227] In some embodiments, the inhibitor is a CTLA-4 inhibitor. In some embodiments, the inhibitor is a LAG-3 inhibitor. In some embodiments, the inhibitor is a PD1 inhibitor. In some embodiments, the inhibitor is a PDL1 inhibitor. In some embodiments, the inhibitor is a TIGIT inhibitor. In some embodiments, the inhibitor is a TIM-3 inhibitor. In some embodiments, the inhibitor is a B7H4 inhibitor. In some embodiments, the inhibitor is a Vista inhibitor. In some embodiments, the inhibitor is a B-RAFi inhibitor. In some embodiments, the B-RAFi inhibitor is vemurafenib. In some embodiments, the inhibitor is a MEKi inhibitor. In some embodiments, the inhibitor is a Btk inhibitor. In some embodiments, the inhibitor is ibrutinib. In some embodiments, the inhibitor is crizotinib. In some embodiments, the inhibitor is an IDO inhibitor. In some embodiments, the inhibitor is an α -CSF1R inhibitor. In some embodiments, the inhibitor is an α -CCR4 inhibitor. In some embodiments, the inhibitor is a TGF-beta. In some embodiments, the inhibitor is a myeloid-derived suppressor cell. In some embodiments, the inhibitor is a T-regulatory cell.

[000228] In some embodiments, the agonist is Ox40. In some embodiments, the agonist is GITR. In some embodiments, the agonist is CD137. In some embodiments, the agonist is ICOS. In some embodiments, the agonist is CD27. In some embodiments, the agonist is HVEM.

[000229] In some embodiments, the anti-PDL1 antibody, conjugated antibody, activatable antibody and/or conjugated activatable antibody is administered before and/or during and/or after treatment in combination with one or more additional agents such as, for example, a chemotherapeutic agent, an anti-inflammatory agent, and/or an immunosuppressive agent. In some embodiments, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and the additional agent are formulated into a single therapeutic composition, and the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and additional agent are administered simultaneously. Alternatively, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and additional agent are separate from each other, *e.g.*, each is formulated into a separate therapeutic composition, and the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and the additional agent are administered simultaneously, or the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and the additional agent are administered at different times during a treatment regimen. For example, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody is administered prior to the administration of the additional agent, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody is administered subsequent to the administration of the additional agent, or the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and the additional agent are administered in an alternating fashion. As described herein, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and additional agent are administered in single doses or in multiple doses.

[000230] In some embodiments, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and the additional agent(s) are administered simultaneously. For example, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and the additional agent(s) can be formulated in a single composition or administered as two or more separate compositions. In some embodiments, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and the additional agent(s) are administered sequentially, or the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and the additional agent are administered at different times during a treatment regimen.

[000231] In some embodiments, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody is administered before and/or during and/or after treatment in combination with one or more additional agents such as, by way of non-limiting example, a chemotherapeutic agent, an anti-inflammatory agent, and/or an immunosuppressive agent, such as an alkylating agent, an anti-metabolite, an anti-microtubule agent, a topoisomerase inhibitor, a cytotoxic antibiotic, and/or any other nucleic acid damaging agent. In some embodiments, the additional agent is a taxane, such as paclitaxel (e.g., Abraxane®). In some embodiments, the additional agent is an anti-metabolite, such as gemcitabine. In some embodiments, the additional agent is an alkylating agent, such as platinum-based chemotherapy, such as carboplatin or cisplatin. In some embodiments, the additional agent is a targeted agent, such as a kinase inhibitor, e.g., sorafenib or erlotinib. In some embodiments, the additional agent is a targeted agent, such as another antibody, e.g., a monoclonal antibody (e.g., bevacizumab), a bispecific antibody, or a multispecific antibody. In some embodiments, the additional agent is a proteasome inhibitor, such as bortezomib or carfilzomib. In some embodiments, the additional agent is an immune modulating agent, such as lenolidomide or IL-2. In some embodiments, the additional agent is radiation. In some embodiments, the additional agent is an agent considered standard of care by those skilled in the art. In some embodiments, the additional agent is a chemotherapeutic agent well known to those skilled in the art.

[000232] In some embodiments, the additional agent is another antibody or antigen-binding fragment thereof, another conjugated antibody or antigen-binding fragment thereof,

another activatable antibody or antigen-binding fragment thereof and/or another conjugated activatable antibody or antigen-binding fragment thereof. In some embodiments the additional agent is another antibody or antigen-binding fragment thereof, another conjugated antibody or antigen-binding fragment thereof, another activatable antibody or antigen-binding fragment thereof and/or another conjugated activatable antibody or antigen-binding fragment thereof against the same target as the first antibody or antigen-binding fragment thereof, the first conjugated antibody or antigen-binding fragment thereof, activatable antibody or antigen-binding fragment thereof and/or a conjugated activatable antibody or antigen-binding fragment thereof, e.g., against PDL1. In some embodiments the additional agent is another antibody or antigen-binding fragment thereof, another conjugated antibody or antigen-binding fragment thereof, another activatable antibody or antigen-binding fragment thereof and/or another conjugated activatable antibody or antigen-binding fragment thereof against a target different than the target of the first antibody or antigen-binding fragment thereof, the first conjugated antibody or antigen-binding fragment thereof, activatable antibody or antigen-binding fragment thereof and/or a conjugated activatable antibody or antigen-binding fragment thereof (i.e., target other than PDL1). In some embodiments, the additional agent is a chimeric antigen receptor-modified T cell, -modified NK cell or other -modified immune effector cell or is another engineered receptor-modified immune effector cell. In some embodiments, the additional agent is a pro-chimeric antigen receptor-modified T cell, -modified NK cell, or other -modified immune effector cell or is another pro-engineered receptor-modified immune effector cell.

[000233] As a non-limiting example, the antibody or antigen-binding fragment and/or the AB of an activatable antibody is a binding partner for any target listed in Table 22. In some embodiments that additional agent is ipilimumab, a CTLA4-binding fragment of ipilimumab, and/or an ipilimumab activatable antibody.

Table 22: Exemplary Targets

1-92-LFA-3	CD52	DL44	HVEM	LIF-R	STEAP1
Alpha-4 integrin	CD56	DLK1	Hyaluronidase	Lewis X	STEAP2
Alpha-V integrin	CD64	DLL4	ICOS	LIGHT	TAG-72
alpha4beta1 integrin	CD70	DPP-4	IFNalpha	LRP4	TAPA1
alpha4beta7 integrin	CD71	DSG1	IFNbeta	LRRC26	TGFbeta

AGR2	CD74	EGFR	IFNgamma	MCSP	TIGIT
Anti-Lewis-Y		EGFRviii	IgE	Mesothelin	TIM-3
Apelin J receptor	CD80	Endothelin B receptor (ETBR)	IgE Receptor (FcεRI)	MRP4	TLR2
APRIL	CD81	ENPP3	IGF	MUC1	TLR4
B7-H4	CD86	EpCAM	IGF1R	Mucin-16 (MUC16, CA-125)	TLR6
BAFF	CD95	EPHA2	IL1B	Na/K ATPase	TLR7
BTLA	CD117	EPHB2	IL1R	Neutrophil elastase	TLR8
C5 complement	CD125	ERBB3	IL2	NGF	TLR9
C-242	CD132 (IL-2RG)	F protein of RSV	IL11	Nicastrin	TMEM31
CA9	CD133	FAP	IL12	Notch Receptors	TNFalpha
CA19-9 (Lewis a)	CD137	FGF-2	IL12p40	Notch 1	TNFR
Carbonic anhydrase 9	CD138	FGF8	IL-12R, IL-12Rbeta1	Notch 2	TNFRS12 A
CD2	CD166	FGFR1	IL13	Notch 3	TRAIL-R1
CD3	CD172A	FGFR2	IL13R	Notch 4	TRAIL-R2
CD6	CD248	FGFR3	IL15	NOV	Transferrin
CD9	CDH6	FGFR4	IL17	OSM-R	Transferrin receptor
CD11a	CEACAM5 (CEA)	Folate receptor	IL18	OX-40	TRK-A
CD19	CEACAM6 (NCA-90)	GAL3ST1	IL21	PAR2	TRK-B
CD20	CLAUDIN-3	G-CSF	IL23	PDGF-AA	uPAR
CD22	CLAUDIN-4	G-CSFR	IL23R	PDGF-BB	VAP1
CD24	cMet	GD2	IL27/IL27R (wsx1)	PDGFRalpha	VCAM-1
CD25	Collagen	GITR	IL29	PDGFRbeta	VEGF
CD27	Cripto	GLUT1	IL-31R	PD-1	VEGF-A
CD28	CSFR	GLUT4	IL31/IL31R	PD-L1	VEGF-B
CD30	CSFR-1	GM-CSF	IL2R	PD-L2	VEGF-C
CD33	CTLA-4	GM-CSFR	IL4	Phosphatidyl-serine	VEGF-D
CD38	CTGF	GP IIb/IIIa receptors	IL4R	P1GF	VEGFR1
CD40	CXCL10	Gp130	IL6, IL6R	PSCA	VEGFR2
CD40L	CXCL13	GP1IB/IIIA	Insulin Receptor	PSMA	VEGFR3
CD41	CXCR1	GPNMB	Jagged Ligands	RAAG12	VISTA

CD44	CXCR2	GRP78	Jagged 1	RAGE	WISP-1
CD44v6		HER2/neu	Jagged 2	SLC44A4	WISP-2
CD47	CXCR4	HGF	LAG-3	Sphingosine 1 Phosphate	WISP-3
CD51	CYR61	hGH			

[000234] As a non-limiting example, the antibody or antigen-binding fragment and/or the AB of an activatable antibody is or is derived from an antibody listed in Table 23.

Table 23: Exemplary sources for Abs

Antibody Trade Name (antibody name)	Target
Avastin™ (bevacizumab)	VEGF
Lucentis™ (ranibizumab)	VEGF
Erbitux™ (cetuximab)	EGFR
Vectibix™ (panitumumab)	EGFR
Remicade™ (infliximab)	TNF α
Humira™ (adalimumab)	TNF α
Tysabri™ (natalizumab)	Integrin α 4
Simulect™ (basiliximab)	IL2R
Soliris™ (eculizumab)	Complement C5
Raptiva™ (efalizumab)	CD11a
Bexxar™ (tositumomab)	CD20
Zevalin™ (ibritumomab tiuxetan)	CD20
Rituxan™ (rituximab)	CD20
(Ocrelizumab)	CD20
Arzerra™ (ofatumumab)	CD20
Gazyva™ (Obinutuzumab)	CD20
Zenapax™ (daclizumab)	CD25
Adcetris™ (brentuximab vedotin)	CD30
Myelotarg™ (gemtuzumab)	CD33
Mylotarg™ (gemtuzumab ozogamicin)	CD33
Campath™ (alemtuzumab)	CD52
ReoPro™ (abixicimab)	Glycoprotein receptor IIb/IIIa
Xolair™ (omalizumab)	IgE
Herceptin™ (trastuzumab)	Her2
Kadcyla™ (trastuzumab emtansine)	Her2
Synagis™ (palivizumab)	F protein of RSV
(ipilimumab)	CTLA-4
(tremelimumab)	CTLA-4
Hu5c8	CD40L
(pertuzumab)	Her2-neu
(ertumaxomab)	CD3/Her2-neu
Orencia™ (abatacept)	CTLA-4
(tanezumab)	NGF
(bavituximab)	Phosphatidylserine
(zalutumumab)	EGFR

(mapatumumab)	EGFR
(matuzumab)	EGFR
(nimotuzumab)	EGFR
ICR62	EGFR
mAb 528	EGFR
CH806	EGFR
MDX-447	EGFR/CD64
(edrecolomab)	EpCAM
RAV12	RAAG12
huJ591	PSMA
Enbrel™ (etanercept)	TNF-R
Amevive™ (alefacept)	1-92-LFA-3
Antril™, Kineret™ (ankinra)	IL-1Ra
GC1008	TGFbeta
	Notch, e.g., Notch 1
	Jagged 1 or Jagged 2
(adecatumumab)	EpCAM
(figitumumab)	IGF1R
(tocilizumab)	IL-6 receptor
Stelara™ (ustekinumab)	IL-12/IL-23
Prolia™ (denosumab)	RANKL

[000235] In some embodiments, the additional antibody or antigen binding fragment thereof, conjugated antibody or antigen binding fragment thereof, activatable antibody or antigen binding fragment thereof, and/or conjugated activatable antibody or antigen binding fragment thereof is a monoclonal antibody, domain antibody, single chain, Fab fragment, a F(ab')₂ fragment, a scFv, a scAb, a dAb, a single domain heavy chain antibody, or a single domain light chain antibody. In some embodiments, the additional antibody or antigen binding fragment thereof, conjugated antibody or antigen binding fragment thereof, activatable antibody or antigen binding fragment thereof, and/or conjugated activatable antibody or antigen binding fragment thereof is a mouse, other rodent, chimeric, humanized or fully human monoclonal antibody.

[000236] The disclosure also provides methods of producing an anti-PDL1 antibody and/or activatable anti-PDL1 antibody polypeptide by culturing a cell under conditions that lead to expression of the polypeptide, wherein the cell comprises an isolated nucleic acid molecule encoding an antibody and/or an activatable antibody described herein, and/or vectors that include these isolated nucleic acid sequences. The disclosure provides methods of producing an antibody and/or activatable antibody by culturing a cell under conditions that lead to expression of the antibody and/or activatable antibody, wherein the cell

comprises an isolated nucleic acid molecule encoding an antibody and/or an activatable antibody described herein, and/or vectors that include these isolated nucleic acid sequences.

[000237] The invention also provides a method of manufacturing activatable antibodies that in an activated state binds PDL1 by (a) culturing a cell comprising a nucleic acid construct that encodes the activatable antibody under conditions that lead to expression of the activatable antibody, wherein the activatable antibody comprises a masking moiety (MM), a cleavable moiety (CM), and an antibody or an antigen binding fragment thereof (AB) that specifically binds PDL1, (i) wherein the CM is a polypeptide that functions as a substrate for a protease; and (ii) wherein the CM is positioned in the activatable antibody such that, when the activatable antibody is in an uncleaved state, the MM interferes with specific binding of the AB to PDL1 and in a cleaved state the MM does not interfere or compete with specific binding of the AB to PDL1; and (b) recovering the activatable antibody. Suitable AB, MM, and/or CM include any of the AB, MM, and/or CM disclosed herein.

[000238] In some embodiments, the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM. In some embodiments, the activatable antibody comprises a linking peptide between the MM and the CM. In some embodiments, the activatable antibody comprises a linking peptide between the CM and the AB. In some embodiments, the activatable antibody comprises a first linking peptide (LP1) and a second linking peptide (LP2), and wherein the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-LP1-CM-LP2-AB or AB-LP2-CM-LP1-MM. In some embodiments, the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: spacer-MM-LP1-CM-LP2-AB or AB-LP2-CM-LP1-MM-spacer. In some embodiments, the two linking peptides need not be identical to each other.

[000239] In some embodiments, at least one of LP1 or LP2 comprises an amino acid sequence selected from the group consisting of (GS)_n, (GGS)_n, (GSGGS)_n (SEQ ID NO: 191) and (GGGS)_n (SEQ ID NO: 192), where n is an integer of at least one.

[000240] In some embodiments, at least one of LP1 or LP2 comprises an amino acid sequence selected from the group consisting of GGSG (SEQ ID NO: 193), GGS GG (SEQ ID NO: 194), GSGSG (SEQ ID NO: 195), GSGGG (SEQ ID NO: 196), GGGSG (SEQ ID NO: 197), and GSSSG (SEQ ID NO: 198).

[000241] In some embodiments, LP1 comprises the amino acid sequence GSSGGSGGSGGSG (SEQ ID NO: 199), GSSGGSGGSGG (SEQ ID NO: 200), GSSGGSGGSGGS (SEQ ID NO: 201), GSSGGSGGSGGSGGGS (SEQ ID NO: 202), GSSGGSGGSG (SEQ ID NO: 203), or GSSGGSGGSGS (SEQ ID NO: 204).

[000242] In some embodiments, LP2 comprises the amino acid sequence GSS, GGS, GGS (SEQ ID NO: 205), GSSGT (SEQ ID NO: 206) or GSSG (SEQ ID NO: 207).

[000243] The invention provides methods of preventing, delaying the progression of, treating, alleviating a symptom of, or otherwise ameliorating an PDL1 mediated disease in a subject by administering a therapeutically effective amount of an anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody described herein to a subject in need thereof.

[000244] The invention also provides methods of preventing, delaying the progression of, treating, alleviating a symptom of, or otherwise ameliorating cancer in a subject by administering a therapeutically effective amount of an anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody described herein to a subject in need thereof. PDL1 is known to be expressed in a variety of cancers, such as, by way of non-limiting example, melanoma, non-small cell lung cancer, nasopharyngeal cancer, glioblastoma/mixed glioma, colon adenocarcinoma, hepatocellular carcinoma, urothelial cancer, multiple myeloma, ovarian cancer, gastric carcinoma, esophageal cancer, pancreatic cancer, renal cell carcinoma (RCC), breast cancer, lymphomas, and leukemias. (See e.g., Chen et al., “Molecular Pathways: Next-Generation Immunotherapy – Inhibiting Programmed Death-Ligand 1 and Programmed Death-1,” Clin. Can. Res., vol. 18: 6580-6587 (2012), the contents of which are hereby incorporated by reference in their entirety).

[000245] In some embodiments, the cancer is a bladder cancer, a bone cancer, a breast cancer, a carcinoid, a cervical cancer, a colon cancer, an endometrial cancer, a glioma, a head and neck cancer, a liver cancer, a lung cancer, a lymphoma, a melanoma, an ovarian cancer, a pancreatic cancer, a prostate cancer, a renal cancer, a sarcoma, a skin cancer, a stomach cancer, a testis cancer, a thyroid cancer, a urogenital cancer, and/or a urothelial cancer.

[000246] In some embodiments, the cancer is selected from the group consisting of melanoma (MEL), renal cell carcinoma (RCC), squamous non-small cell lung cancer (NSCLC), non-squamous NSCLC, colorectal cancer (CRC), castration-resistant prostate

cancer (CRPC), hepatocellular carcinoma (HCC), squamous cell carcinoma of the head and neck, carcinomas of the esophagus, ovary, gastrointestinal tract and breast, or a hematologic malignancy such as multiple myeloma, B-cell lymphoma, T-cell lymphoma, Hodgkin's lymphoma/primary mediastinal B-cell lymphoma, and chronic myelogenous leukemia. In some embodiments, the cancer is due to a PDL1-expressing tumor.

[000247] The invention also provides methods of treating cancer patients with an autoimmune or inflammatory disease by administering a therapeutically effective amount of an anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody described herein to a subject in need thereof. In some embodiments, the autoimmune disease is colitis, RA, pancreatitis, diabetes, or pneumonitis.

[000248] An anti-PDL1 antibody, a conjugated anti-PDL1 antibody, an activatable anti-PDL1 antibody and/or a conjugated activatable anti-PDL1 antibody used in any of the embodiments of these methods and uses can be administered at any stage of the disease. For example, such an anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody can be administered to a patient suffering cancer of any stage, from early to metastatic. The terms subject and patient are used interchangeably herein.

[000249] In some embodiments, the subject is a mammal, such as a human, non-human primate, companion animal (*e.g.*, cat, dog, horse), farm animal, work animal, or zoo animal. In some embodiments, the subject is a human. In some embodiments, the subject is a companion animal. In some embodiments, the subject is an animal in the care of a veterinarian.

[000250] The anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and therapeutic formulations thereof are administered to a subject suffering from or susceptible to a disease or disorder associated with aberrant PDL1 expression and/or activity. A subject suffering from or susceptible to a disease or disorder associated with aberrant PDL1 expression and/or activity is identified using any of a variety of methods known in the art. For example, subjects suffering from cancer or other neoplastic condition are identified using any of a variety of clinical and/or laboratory tests such as, physical examination and blood, urine and/or stool analysis to evaluate health status. For example, subjects suffering from inflammation and/or an inflammatory disorder are identified using any of a variety of

clinical and/or laboratory tests such as physical examination and/or bodily fluid analysis, e.g., blood, urine and/or stool analysis, to evaluate health status.

[000251] Administration of an anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody to a patient suffering from a disease or disorder associated with aberrant PDL1 expression and/or activity is considered successful if any of a variety of laboratory or clinical objectives is achieved. For example, administration of an anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody to a patient suffering from a disease or disorder associated with aberrant PDL1 expression and/or activity is considered successful if one or more of the symptoms associated with the disease or disorder is alleviated, reduced, inhibited or does not progress to a further, *i.e.*, worse, state. Administration of an anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody to a patient suffering from a disease or disorder associated with aberrant PDL1 expression and/or activity is considered successful if the disease or disorder enters remission or does not progress to a further, *i.e.*, worse, state.

[000252] In some embodiments, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody is administered before and/or during and/or after treatment in combination with one or more additional agents such as, for example, a chemotherapeutic agent, an anti-inflammatory agent, and/or an immunosuppressive agent. In some embodiments, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and the additional agent(s) are administered simultaneously. For example, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and the additional agent(s) can be formulated in a single composition or administered as two or more separate compositions. In some embodiments, the anti-PDL1 antibody, conjugated anti-PDL1 antibody, activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody and the additional agent(s) are administered sequentially.

[000253] The invention also provides methods and kits for using the activatable anti-PDL1 antibodies and/or conjugated activatable anti-PDL1 antibodies in a variety of diagnostic and/or prophylactic indications. For example, the invention provides methods and kits for detecting the presence or absence of a cleaving agent and a target of interest in a

subject or a sample by (i) contacting a subject or sample with an anti-PDL1 activatable antibody, wherein the anti-PDL1 activatable antibody comprises a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, and an antigen binding domain or fragment thereof (AB) that specifically binds the target of interest, wherein the anti-PDL1 activatable antibody in an uncleaved, non-activated state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to PDL1, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB and is not a modified form of a natural binding partner of the AB; and (b) wherein, when the AB is in an uncleaved, non-activated state, the MM interferes with specific binding of the AB to PDL1, and when the AB is in a cleaved, activated state the MM does not interfere or compete with specific binding of the AB to PDL1; and (ii) measuring a level of activated anti-PDL1 activatable antibody in the subject or sample, wherein a detectable level of activated anti-PDL1 activatable antibody in the subject or sample indicates that the cleaving agent and PDL1 are present in the subject or sample and wherein no detectable level of activated anti-PDL1 activatable antibody in the subject or sample indicates that the cleaving agent, PDL1 or both the cleaving agent and PDL1 are absent in the subject or sample.

[000254] In some embodiments, the activatable anti-PDL1 antibody is an activatable anti-PDL1 antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable anti-PDL1 antibody is not conjugated to an agent. In some embodiments, the activatable anti-PDL1 antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable anti-PDL1 antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000255] In some embodiments of these methods and kits, the activatable anti-PDL1 antibody includes a detectable label. In some embodiments of these methods and kits, the detectable label includes an imaging agent, a contrasting agent, an enzyme, a fluorescent label, a chromophore, a dye, one or more metal ions, or a ligand-based label. In some embodiments of these methods and kits, the imaging agent comprises a radioisotope. In some embodiments of these methods and kits, the radioisotope is indium or technetium. In some embodiments of these methods and kits, the contrasting agent comprises iodine,

gadolinium or iron oxide. In some embodiments of these methods and kits, the enzyme comprises horseradish peroxidase, alkaline phosphatase, or β -galactosidase. In some embodiments of these methods and kits, the fluorescent label comprises yellow fluorescent protein (YFP), cyan fluorescent protein (CFP), green fluorescent protein (GFP), modified red fluorescent protein (mRFP), red fluorescent protein tdimer2 (RFP tdimer2), HCRED, or a europium derivative. In some embodiments of these methods and kits, the luminescent label comprises an N- methylacrydium derivative. In some embodiments of these methods, the label comprises an Alexa Fluor[®] label, such as Alex Fluor[®] 680 or Alexa Fluor[®] 750. In some embodiments of these methods and kits, the ligand-based label comprises biotin, avidin, streptavidin or one or more haptens.

[000256] In some embodiments of these methods and kits, the subject is a mammal. In some embodiments of these methods, the subject is a human. In some embodiments, the subject is a non-human mammal, such as a non-human primate, companion animal (e.g., cat, dog, horse), farm animal, work animal, or zoo animal. In some embodiments, the subject is a rodent.

[000257] In some embodiments of these methods and kits, the method is an *in vivo* method. In some embodiments of these methods, the method is an *in situ* method. In some embodiments of these methods, the method is an *ex vivo* method. In some embodiments of these methods, the method is an *in vitro* method.

[000258] In some embodiments of the methods and kits, the method is used to identify or otherwise refine a patient population suitable for treatment with an anti-PDL1 activatable antibody of the disclosure, followed by treatment by administering that activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody to a subject in need thereof. For example, patients that test positive for both the target (e.g., PDL1) and a protease that cleaves the substrate in the cleavable moiety (CM) of the anti-PDL1 activatable antibody being tested in these methods are identified as suitable candidates for treatment with such an anti-PDL1 activatable antibody comprising such a CM, and the patient is then administered a therapeutically effective amount of the activatable anti-PDL1 antibody and/or conjugated activatable anti-PDL1 antibody that was tested. Likewise, patients that test negative for either or both of the target (e.g., PDL1) and the protease that cleaves the substrate in the CM in the activatable antibody being tested using these methods might be identified as suitable candidates for another form of therapy. In some embodiments, such patients can be tested with other anti-PDL1 activatable antibodies until a

suitable anti-PDL1 activatable antibody for treatment is identified (e.g., an anti-PDL1 activatable antibody comprising a CM that is cleaved by the patient at the site of disease). In some embodiments, the patient is then administered a therapeutically effective amount of the activatable anti-PDL1 antibody and/or conjugated for which the patient tested positive. Suitable AB, MM, and/or CM include any of the AB, MM, and/or CM disclosed herein.

[000259] Pharmaceutical compositions according to the invention can include an antibody of the invention and a carrier. These pharmaceutical compositions can be included in kits, such as, for example, diagnostic kits.

Brief Description of the Drawings

[000260] Figure 1 is graph depicting by ELISA-based binding that PDL1 c60 ScFv-phage binds specifically to human and mouse PDL1.

[000261] Figure 2 is a graph depicting that hPDL1 inhibits the binding of clones 12, 18 and 60 to hPDL1 in an ELISA format, and, furthermore mPDL1 inhibits the binding of clone 60 to mPDL1.

[000262] Figure 3 is a graph depicting saturation binding data and demonstrates that anti-PDL1 clone 60:IgG binds to human PDL1 with a binding affinity ~ 1 nM, and to mouse PDL1 at 30 nM.

[000263] Figures 4A and 4B are a series of graphs depicting H2 specificity and cross reactivity (Figure 4A) and H3W specificity and cross reactivity (Figure 4B).

[000264] Figures 5A and 5B are a series of graphs depicting that combining the matured heavy variable CDR3 domains with the heavy variable CDR2 domains results in antibodies with enhanced affinity towards human PDL1.

[000265] Figures 6A, 6B, 6C, and 6D are a series of graphs depicting that anti-PDL1 antibody C5H9 binds to human and mouse PDL1 with near equal affinity and that anti-PDL1 antibodies C5B10 and C5E10 only bind human PDL1.

[000266] Figure 7 is a graph depicting the ability of anti-PDL1 antibody C5H9 to bind with high and equal affinity to human, mouse and cyno PDL1.

[000267] Figures 8A and 8B are a series of graphs depicting that anti-PDL1 antibody C5H9v2 has lower immunogenicity potential and enhanced manufacturability. These graphs demonstrate decreased predicted immunogenicity, increased expression levels (3X), and increased monomer percentage in human IgG1 format.

[000268] Figures 9A, 9B, and 9C are a series of graphs depicting ability of anti-PDL1 antibody C5H9v2 to bind with high affinity to (A) mouse, (B) cyno, and (C) human PDL1. Figures 9A-9C show that both anti-PDL1 antibodies C5H9 and C5H9v2 bind human, mouse and cyno PDL1 with near equal affinities.

[000269] Figure 10 is a graph depicting C5H9v2 specificity for a panel of human and mouse proteins. Figure 10 demonstrates that anti-PDL1 antibody C5H9v2 binds only to human and mouse PDL1 demonstrating specificity toward PDL1.

[000270] Figure 11 is a series of graphs depicting B7-1 and PD1 blockade. Figure 11 demonstrates that both anti-PDL1 antibody C5H9 and anti-PDL1 antibody C5H9v2 are potent blockers of either B7-1 or PD1 binding to PDL1, and that blockade includes all three species human, cyno and mouse with single digit nM EC₅₀'s.

[000271] Figure 12 is a graph depicting that anti-PDL1 antibody C5H9 accelerates the onset of diabetes in NOD mice similar to the positive control anti-PDL1 antibody 10F9G2.

[000272] Figure 13 is a graph depicting that anti-PDL1 antibody C5H9v2 accelerates the onset of diabetes in NOD mice in a dose dependent matter.

[000273] Figure 14 is a graph depicting that an anti-PDL1 antibody C5H9v2 inhibits the growth of MC38 syngeneic tumors similar to positive control anti-PDL1 antibody 10F9G2.

[000274] Figure 15 is a graph depicting binding isotherms for anti-PDL1 activatable antibodies that include the anti-PDL1 antibody C5H9v2.

[000275] Figure 16 is a graph depicting binding isotherms for anti-PDL1 activatable antibodies that include the anti-PDL1 antibody C5H9v2.

[000276] Figure 17 is a graph depicting binding isotherms for anti-PDL1 activatable antibodies that include the anti-PDL1 antibody C5H9v2.

[000277] Figures 18A and 18B are a series of graphs depicting that anti-PDL1 activatable antibody PL15-0003-C5H9v2, when administered at a dose of 1 mg/kg, has an absence of diabetes induction and anti-PDL1 activatable antibody PL18-0003-C5H9v2, when administered at a dose of 1 mg/kg, has a delay in diabetes induction compared to the anti-PDL1 antibody C5H9v2 administered at a dose of 1 mg/kg in NOD mice (Figure 18A); in addition, anti-PDL1 activatable antibody PL15-0003-C5H9v2, when administered at a dose of 3 mg/kg induced diabetes in 2 of 8 NOD mice (75% non-diabetic), compared to 1 mg/kg anti-PDL1 antibody C5H9v2 which led to 38% non-diabetic NOD mice (Figure 18B).

[000278] Figure 19 is a graph depicting that anti-PDL1 activatable antibodies PL15-0003-C5H9v2 and PL18-0003-C5H9v2 inhibit the growth of MC38 syngeneic tumors similar to anti-PDL1 antibody C5H9v2.

[000279] Figures 20A, 20B, 20C, and 20D are a series of graphs depicting the percent of anti-PDL1 antibody C5H9v2 and anti-PDL1 activatable antibody PL15-0003-C5H9v2 bound to CD4+ and CD8+ T cells from peripheral blood (Figures 20A, 20B) or spleen (Figures 20C, 20D). In all graphs, the circle represents isotype, the square represents anti-PDL1 antibody C5H9v2, and the triangle represents anti-PDL1 activatable antibody PL15-0003-C5H9v2.

[000280] Figures 21A and 21B are a series of graphs depicting that the presence of tumor-derived proteases capable of cleaving anti-PDL1 activatable antibody PL15-0003-C5H9v2 (depicted by squares) did not lead to high levels of activated activatable antibody in the blood compared to anti-PDL1 antibody C5H9v2 (depicted by circles) or an isotype antibody (depicted by triangles) (Figure 21A) despite the plasma concentrations of the activatable antibody being higher than those of the antibody (Figure 21B).

[000281] Figure 22 is a graph depicting the ability of anti-PDL1 activatable antibody PL07-2001-C5H9v2 to increase CMV-stimulated IFN-gamma secretion as compared to control hIgG4, but with decreased potency relative to anti-PDL1 parental antibody C5H9v2.

[000282] Figures 23A, 23B, 23C, and 23D are a series of images showing the ability of anti-PDL1 activatable antibody PL15-0003-C5H9v2 to be activated and to bind frozen MC38 mouse cancer tissues using an *in situ* imaging method.

[000283] Figure 24 depicts that anti-PDL1 activatable antibody PL15-0003-C5H9v2 exhibited a higher EC₅₀ for PDL1 binding and PD1 blocking than did the anti-PDL1 antibody as measured by ELISA. The figure further demonstrates that activation of anti-PDL1 activatable antibody PL15-0003-C5H9v2 by matriptase fully restored PDL1 binding and PD1 blocking activities to levels comparable to those of anti-PDL1 antibody C5H9v2.

[000284] Figures 25A and 25B are a series of graphs depicting the ability of anti-PDL1 activatable antibodies referred to herein as PL07-2001-C5H9v2 and PF07-3001-C5H9v2 and of the anti-PDL1 antibody referred to herein as C5H9 to bind to human or murine PDL1.

[000285] Figures 26A, 26B, and 26C are a series of graphs depicting the abilities of anti-PDL1 activatable antibody PL07-2001-C5H9v2, uPA-activated anti-PDL1 activatable antibody PL07-2001-C5H9v2, MMP14-activated anti-PDL1 activatable antibody PL07-

2001-C5H9v2, and anti-PDL1 antibody C5H9v2 to block human PD1 (PD-1) to human PDL1 (Figure 26A), cynomolgus PD1 (PD-1) binding to cynomolgus PDL1 (Figure 26B), or rat PD1 (PD-1) binding to rat PDL1 (Figure 26C).

[000286] Figures 27A and 27B are a series of graphs depicting the ability of anti-PDL1 activatable antibody PL07-2001-C5H9v2, uPA-activated anti-PDL1 activatable antibody PL07-2001-C5H9v2, MMP14-activated anti-PDL1 activatable antibody PL07-2001-C5H9v2, or anti-PDL1 antibody C5H9v2 to human B7-1 binding to human PDL1 (Figure 27A) or cynomolgus B7-1 binding to cynomolgus PDL1 (Figure 27B).

[000287] Figure 28 is a graph depicting the ability of the anti-PDL1 activatable antibodies PL15-0003-C5H9v2, PL15-2001-C5H9v2, and PL15-3001-C5H9v2 to inhibit the growth of MC38 syngeneic tumors similar to positive control anti-PDL1 antibody C5H9v2.

[000288] Figures 29A, 29B, and 29C are a series of graphs depicting that the presence of tumor-derived proteases capable of cleaving the activatable antibodies anti-PDL1 antibody C5H9v2 or anti-PDL1 activatable antibody PL15-0003-C5H9v2, PL15-2001-C5H9v2, or PL15-3001-C5H9v2 did not lead to high levels of activated activatable antibodies in the blood.

[000289] Figure 30 is an illustration depicting the ability of proteases in a tumor implanted into a mouse to activate the anti-PDL1 activatable antibody PL07-2001-C5H9v2.

Detailed Description of the Invention

[000290] The present invention provides monoclonal antibodies (mAbs) and activatable monoclonal antibodies that specifically bind human programmed death ligand 1 (PDL1). PDL1 is a 40kDa type I transmembrane protein that forms a complex with its receptor programmed cell death protein 1 (PD1), also known as CD279. Engagement of PDL1 with its receptor PD1 on T cells delivers a signal that inhibits TCR-mediated activation of IL-2 production and T cell proliferation. Aberrant expression and/or activity of PDL1 and PDL1-related signaling has been implicated in the pathogenesis of many diseases and disorders, such as cancer, inflammation, and autoimmunity.

[000291] The activatable anti-PDL1 antibodies are used in methods of treating, preventing, delaying the progression of, ameliorating and/or alleviating a symptom of a disease or disorder associated with aberrant PDL1 expression and/or activity. For example, the activatable anti-PDL1 antibodies are used in methods of treating, preventing, delaying

the progression of, ameliorating and/or alleviating a symptom of a cancer or other neoplastic condition.

[000292] The activatable anti-PDL1 antibodies include an antibody or antigen-binding fragment thereof that specifically binds PDL1 coupled to a masking moiety (MM), such that coupling of the MM reduces the ability of the antibody or antigen-binding fragment thereof to bind PDL1. In some embodiments, the MM is coupled via a sequence that includes a substrate for a protease, for example, a protease that is co-localized with PDL1 at a treatment site in a subject.

[000293] Exemplary activatable anti-PDL1 antibodies of the invention include, for example, activatable antibodies that include a heavy chain and a light chain that are, or are derived from, antibodies described in the Examples, for example in Example 3 and Example 4.

[000294] In some embodiments, the activatable anti-PDL1 antibody includes a heavy chain that comprises or is derived from an amino acid sequence selected from the group consisting of SEQ ID NO: 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, and 56, and a light chain that comprises or is derived from the amino acid sequence of SEQ ID NO: 12 or SEQ ID NO: 58.

[000295] In some embodiments, the activatable anti-PDL1 antibody comprises a combination of a variable heavy chain complementarity determining region 1 (VH CDR1, also referred to herein as CDRH1) sequence, a variable heavy chain complementarity determining region 2 (VH CDR2, also referred to herein as CDRH2) sequence, a variable heavy chain complementarity determining region 3 (VH CDR3, also referred to herein as CDRH3) sequence, a variable light chain complementarity determining region 1 (VL CDR1, also referred to herein as CDRL1) sequence, a variable light chain complementarity determining region 2 (VL CDR2, also referred to herein as CDRL2) sequence, and a variable light chain complementarity determining region 3 (VL CDR3, also referred to herein as CDRL3) sequence, wherein at least one CDR sequence is selected from the group consisting of a VH CDR1 sequence shown in Table 15; a VH CDR2 sequence shown in Table 15; a VH CDR3 sequence shown in Table 15; a VL CDR1 sequence shown in Table 15; a VL CDR2 sequence shown in Table 15; and a VL CDR3 sequence shown in Table 15.

[000296] In some embodiments, the activatable antibody comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, a VH CDR3 sequence, a VL CDR1 sequence,

a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the six CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, and VL CDR3) shown in a single row in Table 15.

[000297] In some embodiments, the antibody comprises a heavy chain that comprise a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein the combination is a combination of the three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 15.

[000298] In some embodiments, the antibody comprises a light chain that comprise a combination of a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the three light chain CDR sequences (VL CDR1, VL CDR2, VL CDR3) shown in a single row in Table 15.

[000299] In some embodiments, the activatable anti-PDL1 antibody includes a heavy chain that comprises or is derived from a heavy chain amino acid sequence shown in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes a light chain that comprises or is derived from a light chain amino acid sequence shown in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes a heavy chain that comprises or is derived from a heavy chain amino acid sequence shown in Table 15, and a light chain that comprises or is derived from a light chain amino acid sequence shown in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain variable region and light chain variable region sequences from the combinations shown in Group A in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group B in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group C in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group D in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group E in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group F in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences

shown in Group G in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group H in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group I in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain and light chain sequences shown in Group J in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group K in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain and light chain sequences shown in Group K in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group L in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group M in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group N in Table 15. In some embodiments, the activatable anti-PDL1 antibody includes the combination of heavy chain variable region and light chain variable region sequences shown in Group O in Table 15.

Table 15. Variable Heavy Chain Region (VH) and Variable Light Chain Region (VL)
Sequences for Activatable Antibodies that Bind PDL1 (CDR sequences are underlined; CDR sequences disclosed herein were identified in accordance with the definitions of Kabat, Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md. (1987 and 1991)))

Group A	
VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTD <u>YGF</u> SWVRQAPGQGLEWMGWITAYNGNTNYAQKLQGRVTM TTDTSTSTVYMELRSLRSDDTAVYYCARDY <u>FYGM</u> DVWGQGT ^{TV} TVSS (SEQ ID NO: 248)
VH	QVQLVQSGAEVKKPGSSVKVCKTSGDTFSTYAI <u>SWVRQAPGQGLEWMGGI</u> IPIFGKAHYAQKFQGRVTI TADESTSTAYMELSSLRSED ^T AVYFCARKFHFVSGSPFGMDVWGQGT ^{TV} TVSS (SEQ ID NO: 249)
VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYDVHWVRQAPGQRLEWMGWLHADTGITKFSQKFQGRVTI TRDTSASTAYMELSSLRSED ^T AVYYCAR <u>ERIQLWFDY</u> WGQGT (SEQ ID NO: 250)

VH	QVQLVQSGAEVKKPGSSVKVSKVSGGIFSTYAINWVRQAPGQGLEWMGGIIPFGTANHAQKFQGRVTI TADESTSTAYMELSSLRSEDTAVYYCARDQGIAAALFDYWGQGLTVTVSS (SEQ ID NO: 251)
VH	EVQLVESGGGLVQPGRSLRLSCAVSGFTFDDYVHVWVRQAPGKGLEWVSGNSGNIGYADSVKGRFTISR NAKNSLYLQMNSLRAEDTALYYCAVPFDYWGQGLTVTVSS (SEQ ID NO: 252)
VH	QVQLVQSGAEVKKPGSSVKVSKTSGDTFSSYAISWVRQAPGQGLEWMGGIIPFGRAHYAQKFQGRVTI TADESTSTAYMELSSLRSEDTAVYFCARKFHFVSGSPFGMDVWGQGTITVTVSS (SEQ ID NO: 253)
VH	QVQLVQSGAEVKKPGSSVKVSKTSGDTFSSYAISWVRQAPGQGLEWMGGIIPFGKAHYAQKFQGRVTI TADESTTTAYMELSSLRSEDTAVYYCARKYDYVSGSPFGMDVWGQGTITVTVSS (SEQ ID NO: 254)
VH	QVQLVQSGAEVKKPGSSVKVSKASGGTFSSYAINWVRQAPGQGLEWMGGIIPFGSANYAQKFQDRVTI TADESTSAAYMELSSLRSEDTAVYYCARDSSGWSRYMDVWGQGTITVTVSS (SEQ ID NO: 255)
VH	QVQLVQSGAEVKEPGSSVKVSKASGGTFNSYAISWVRQAPGQGLEWMGGIIPFGIAHYAQKFQGRVTI TADESTNTAYMDLSSLRSEDTAVYYCARKYSYVSGSPFGMDVWGQGTITVTVSS (SEQ ID NO: 256)
VH	EVQLVESGGGLVQPGRSLRLSCAASGITFDDYGMHWVRQAPGKGLEWVSGISWNRGRIEYADSVKGRFTI SRDNAKNSLYLQMNSLRAEDTALYYCAKGRFRYFDWFLDYWGQGLTVTVSS (SEQ ID NO: 257)
VH	QMQLVQSGGGLVQPGGSLRLSCAASGFTFSSYWMSWVRQAPGKGLEWVANIKQDGSEKYYVDSVKGRFTI SRDNAKNSLYLQMNSLRAEDTAVYYCARDYFWSGFSAFDIWGKGLTVTVS (SEQ ID NO: 449)
VL	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLVWYQQKPGQAPRLLIYDASNRATGIPARFSGSGSGTD FTLTISLLEPEDFAVYYCQQRSNWPRTFGQGTKVEIK (SEQ ID NO: 258)
VL	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDASNRATGIPARFSGSGSGTD FTLTISLLEPEDFAVYYCQQRSNWPTFGQGTKVEIK (SEQ ID NO: 259)
VL	DIQMTQSPSSLSASVGDRVTITCRASQGISSWLAWYQQKPEKAPKSLIYAASSLQSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYNSYPYTFGQGTKLEIK (SEQ ID NO: 260)
VL	EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGT DFTLTISRLEPEDFAVYYCQQYGSSPWTFGQGTKVEIK (SEQ ID NO: 261)
VL	EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGT DFTLTISRLEPEDFAVYYCQQYGSSPFGGGTKVEIK (SEQ ID NO: 262)
VL	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDASNRATGIPARFSGSGSGTD FTLTISLLEPEDFAVYYCQQRSNWPTFGQGTREIK (SEQ ID NO: 263)
VL	AIQLTQSPSSLSASVGDRVTITCRASQGISSALAWYQQKPGKAPKLLIYDASSLESVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQFNYPFTFGPGTKVDIK (SEQ ID NO: 264)
	DIVMTQSPSTLSASVGDRVTITCRASQGISSWLAWYQQKPGRAPKVLIIYKASTLESVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQSYSTPWTFGQGTKLEIK (SEQ ID NO: 450)
Group B	
VH	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWMSWVRQAPGKGLEWVANIKQDGSEKYYVDSVKGRFTI SRDNAKNSLYLQMNSLRAEDTAVYYCAREGGWFGELAFDYWGQGLTVTVSS (SEQ ID NO: 265)
VL	EIVLTQSPGTLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDASSRATGIPDRFSGSGSGT DFTLTISRLEPEDFAVYYCQQYGSLPWTFGQGTKVEIK (SEQ ID NO: 266)

Group C	
VH	EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHVVRQAPGKGLEWVAVISPYGGSTYYADSVKGRFTI SADTSKNTAYLQMNSLRAEDTAVYYCARRHWPGGFDYWGQGLTVTVSA (SEQ ID NO: 267)
VH	EVQLVESGGGLVQPGGSLRLSCAASGFTFSGSWIHVVRQAPGKGLEWVAVILPYGGSSYYADSVKGRFTI SADTSKNTAYLQMNSLRAEDTAVYYCARRHWPGGFDYWGQGLTVTVSA (SEQ ID NO: 451)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYLYHPATFGQGGTKVEIKR (SEQ ID NO: 268)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYINVPWTFGQGGTKVEIKR (SEQ ID NO: 452)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYYAPPWTFGQGGTKVEIKR (SEQ ID NO: 453)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYYTVPWTFGQGGTKVEIKR (SEQ ID NO: 454)
VL	DIQMTQSPSSLSASVGDRVTITCRASQVINTFLAWYQQKPGKAPKLLIYSASTLASGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYYTVPRTFGQGGTKVEIKR (SEQ ID NO: 455)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQGYGVPRTFGQGGTKVEIKR (SEQ ID NO: 456)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYLFTPPTFGQGGTKVEIKR (SEQ ID NO: 457)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYFITPTTFGQGGTKVEIKR (SEQ ID NO: 458)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYYYTPPTFGQGGTKVEIKR (SEQ ID NO: 459)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQFFYTPTTFGQGGTKVEIKR (SEQ ID NO: 460)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQSLFTPPTFGQGGTKVEIKR (SEQ ID NO: 461)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQSLYTPPTFGQGGTKVEIKR (SEQ ID NO: 462)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQSWYHPPTFGQGGTKVEIKR (SEQ ID NO: 463)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYFYIPPTFGQGGTKVEIKR (SEQ ID NO: 464)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYWYTPPTFGQGGTKVEIKR (SEQ ID NO: 465)
VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQSYFIPPTFGQGGTKVEIKR (SEQ ID NO: 466)
Group D	

VH	METGLRWLLLVAVLKGVQCLSVESGGRLVTPGTPLTLTCTASGFTITNYHMFWRQAPGKGLEWIGVIT SSGIGSSSTTTYATWAKGRFTISKSTSTTVNLRITSPTTEDTATYFCARDYFTNTYYALDIWGPGLTVTS S (SEQ ID NO: 467)
VH	QVQLVQSGAEVKKPGSSVKVSKTSGDTFSTYAI <u>SWVRQAPGQGLEWMGGIIPIFGKAHYAQKFQGRVTI</u> TADESTSTAYMELSSLRSED <u>TAVYFCARKFH</u> FVSGSPFGMDVWGQGTTVTVSS (SEQ ID NO: 249)
VH	QVQLVQSGAEVKKPGASVKVSKASGYTFTSYDVHWVRQAPGQRLEWMGWLHADTGITKFSQKFQGRVTI TRDTSASTAYMELSSLRSED <u>TAVYYCARERIQLWFDYWGQGT</u> TLTVTVSS (SEQ ID NO: 468)
VH	QVQLVQSGAEVKKPGSSVKVSKVSGGIFSTYAINWVRQAPGQGLEWMGGIIPIFGTANHAQKFQGRVTI TADESTSTAYMELSSLRSED <u>TAVYYCARDQGIAAALFDYWGQGT</u> TLTVTVSS (SEQ ID NO: 251)
VH	EVQLVESGGGLVQPGRSLRLSCAVSGFTFDYVHVWVRQAPGKGLEWVSGISGNSGNIGYADSVKGRFTI SRDNAKNSLYLQMNSLRAEDTALYYCAVPFDYWGQGTTLTVTVSS (SEQ ID NO: 469)
VH	QVQLVQSGAEVKKPGSSVKVSKTSGDTFSSYAI <u>SWVRQAPGQGLEWMGGIIPIFGRAHYAQKFQGRVTI</u> TADESTSTAYMELSSLRSED <u>TAVYFCARKFH</u> FVSGSPFGMDVWGQGTTVTVTVSS (SEQ ID NO: 253)
VH	QVQLVQSGAEVKKPGSSVKVSKTSGGTFSSYAI <u>SWVRQAPGQGLEWMGGIIPIFGKAHYAQKFQGRVTI</u> TADESTTTTAYMELSSLRSED <u>TAVYYCARKYDYVSGSPFGMDVWGQGT</u> TVTVTVSS (SEQ ID NO: 254)
VH	QVQLVQSGAEVKKPGSSVKVSKASGGTFSSYAINWVRQAPGQGLEWMGGIIPIFGSANYAQKFQDRVTI TADESTSAAYMELSSLRSED <u>TAVYYCARDSSGWSRY</u> YMDVWGQGTTVTVTVSS (SEQ ID NO: 255)
VH	QVQLVQSGAEVKEPGSSVKVSKASGGTFNSYAI <u>SWVRQAPGQGLEWMGGIIPLF</u> GIAHYAQKFQGRVTI TADESTNTAYMDLSSLRSED <u>TAVYYCARKYSYVSGSPFGMDVWGQGT</u> TVTVTVSS (SEQ ID NO: 256)
VH	EVQLVESGGGLVQPGRSLRLSCAASGITFDYGMHWVRQAPGKGLEWVSGISWNRGRIEYADSVKGRFTI SRDNAKNSLYLQMNSLRAEDTALYYCAKGRFRYFDWFLDYWGQGTTLTVTVSS (SEQ ID NO: 257)
VL	MDTRAPTQLLGLLLLWLPGARCALVMTQTPSSTSTAVGGTVTIKCQASQSIISVYLAWYQQKPGQPPLLII YSASTLASGVPSRFKGSRSRGTEYTLTISGVQREDAATYYCLGSAGS (SEQ ID NO: 470)
VL	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLVWYQQKPGQAPRLLIYDASN <u>RATGIPARFSGSGSGTD</u> FTLTISLLEPEDFAVYYCQQRSNWPRTFGQGTKVEIK (SEQ ID NO: 258)
VL	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDASN <u>RATGIPARFSGSGSGTD</u> FTLTISLLEPEDFAVYYCQQRSNWPRTFGQGTKVEIK (SEQ ID NO: 259)
VL	DIQMTQSPSSLSASVGDRVTITCRASQGISSWLAWYQQKPEKAPKSLIYAASSLQSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYNSYPYTFGQGTKLEIK (SEQ ID NO: 260)
VL	EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGT DFTLTISRLEPEDFAVYYCQQYGSSPWFPGQGTKVEIK (SEQ ID NO: 261)
VL	EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGT DFTLTISRLEPEDFAVYYCQQYGSSPFGGGTKVEIK (SEQ ID NO: 262)
VL	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDASN <u>RATGIPARFSGSGSGTD</u> FTLTISLLEPEDFAVYYCQQRSNWPRTFGQGT <u>RLEIK</u> (SEQ ID NO: 263)
VL	AIQLTQSPSSLSASVGDRVTITCRASQGISSALAWYQQKPGKAPKLLIYDASSLESVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQFN <u>SYPTFTFGPGTKVDIK</u> (SEQ ID NO: 264)
Group E	

VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYIMMWVRQAPGKGLEWVSS <u>IYPSGGITFYADTVKGRFTI</u> SRDNSKNTLYLQMNSLRAEDTAVYYCARI <u>KLGTVT</u> TVDYWGQGT <u>LVTVSS</u> (SEQ ID NO: 471)
VL	QSALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMIDVSNRPSGVSNRFSGSKSG NTASLTISGLQAEDEADYYC <u>SSYTSSSTRV</u> FGTGTKVTVL (SEQ ID NO: 472)

Group F	
VH	EVKLQESGP SLVKPSQTL SLTCSVTGYSITSDYWNWIRKFP GNKLEYVGYISYTGSTYYNPSLKS RISIT RDTSKNQYYLQLNSVTSEDTATYYCARYGGWLS PF DYWGQGTTLTVSS (SEQ ID NO: 473)
VH	EVQLQESGPGLVAPSQSL SITCTVSGFSLTTY SINWIRQPPGKLEWL GVMWAGGTNSNSVLKS RLIIS KD NSKSQVFLKMNSLQ TDDTARYYCARYYGNSPYAIDYWGQGTSTVTVSS (SEQ ID NO: 474)
VH	EVKLQESGP SLVKPSQTL SLTCSVTGYSIISDYWNWIRKFP GNKLEYLGYISYTGSTYYNPSLKS RISIT RDTSKNQYYLQLNSVT TED TATYYCARRGGWLL PF DYWGQGTTLTVSS (SEQ ID NO: 475)
VH	EVKLQESGP SLVKPGASVKLSCKASGYTFTSYDINWVKQRP QGLEWIGWIFPRDNN TKY NENFKGKATL TVDTSS TTAYMELHSLTSEDS AVYFCTKENWVGDFDYWGQGTTLTLSS (SEQ ID NO: 476)
VH	EVQLQQSGPDLVTPGASVRIS CQASGYTFPDYYMNWVKQSHGKSLEWIGDIDPNYGGTTYNQKF GKAIL TVDRSS STAYMELRSLTSEDS AVYYCARGALTDWGQGTSLTVSS (SEQ ID NO: 477)
VH	EIVLTQSPATLSLSPGERATLSC RASSSVSYIYWFQ QKPGQSPRPLIYAAFNRATGIPARFSGSGSGTDY TLTIS SLEPEDFAVYYCQQWSNNPLTFG QGTKVEIK (SEQ ID NO: 478)
VH	QVQLVQSGAEVKKPGASVKV SCKASGYTFPDYYMNWVRQAPGQ GLEWMGDIDPNYGGTNYAQKFQGRVTM TRDTSISTAYMEL SRLRSDDTAVYYCARGALTDWGQGT MTVTVSS (SEQ ID NO: 479)
VH	QVQLVQSGAEVKKPGASVKV SCKASGYTFPDYYMNWVRQAPGQ SLEWMGDIDPNYGGTNYNQKFQGRVTM TRDTSISTAYMEL SRLRSDDTAVYYCARGALTDWGQGT MTVTVSS (SEQ ID NO: 480)
VH	EVQLVQSGAEVKKPGASVKV SCKASGYTFPDYYMNWVRQAPGQ SLEWMGDIDPNYGGTNYNQKFQGRVTM TVDRSS STAYMELRSLRSDDTAVYYCARGALTDWGQGT MTVTVSS (SEQ ID NO: 481)
VH	EVQLVESGGGLVQPGRSLRLSCTASGYTFPDYYMNWVRQAPGKLEWVG DIDPNYGGTTYAASVKGRFTI SVDRSKSIAYLQMS LKTEDTAVYYCTR GALTDWGQGTMTVTVSS (SEQ ID NO: 482)
VH	EVQLVESGGGLVQPGRSLRLSCTASGYTFPDYYMNWVRQAPGKLEWVG DIDPNYGGTTYNASVKGRFTI SVDRSKSIAYLQMS LKTEDTAVYYCARGALTDWGQGT MTVTVSS (SEQ ID NO: 483)
VL	DIVMTQSHKLMSTSVGDRVSITCKASQDVGTAVAWYQ QKPGQSPKLLIYWASTRHTGVPDRFTGSGSGTD FTLTISNVQSEDLADYFCQ QDSSYPLTFGAGTK VELK (SEQ ID NO: 484)
VL	DIVTTQSHKLMSTSVGDRVSITCKASQDVGTAVAWYQ QKPGQSPKLLIYWASTRHTGVPDRFTGSGSGTD FTLTISNVQSEDLADYFCQ QDSSYPLTFGAGTK VELK (SEQ ID NO: 485)
VL	DIVMTQSPSSLA VS VG EKVS MGCKSSQSLLYSSNQ KN SLAWYQ QKPGQSPKLLIDWASTRESGVPDRFTG SGSGTDFTLTIS SVKAEDLAVYYCQ YYGYPLTFGAGTKLELK (SEQ ID NO: 486)
VL	DIVMTQSPA IM ASAPGEKVTMTCSASSSIRYMH WYQ QKPGTSPKRWISDTSKLTSGV PARFSGSGSGTSY ALTIS SMEAEDAATYYCHQRSSYPWTFGGG TKLEIK (SEQ ID NO: 487)
VL	QIVLSQSPAILSAS APGEKVTMTCRASSSVSYIYWFQ QKPGSSPKWIYATFNLASGVPARFSGSGSGTSY SLTISR VETEDAATYYCQQWSNNPLTFGAGTK LELK (SEQ ID NO: 488)
VL	EIVLTQSPATLSLSPGERATLSC RASSSVSYIYWFQ QKPGQAPRLLIYAAFNRATGIPARFSGSGSGTDY TLTIS SLEPEDFAVYYCQQWSNNPLTFG QGTKVEIK (SEQ ID NO: 489)
VL	QIVLTQSPATLSLSPGERATLSC RASSSVSYIYWFQ QKPGQSPRPLIYATFNLASGIPARFSGSGSGTSY TLTISR LEPEDFAVYYCQQWSNNPLTFG QGTKVEIK (SEQ ID NO: 490)

VL	DIQLTQSPSSLSASVGDRVTTITCRASSGVSYIYWFQQKPGKAPKLLIYAAFNLASGVPSRFSGSGSGTEY TLTISSLQPEDFATYYCQQWSNNPLTFGQGTKVEIK (SEQ ID NO: 491)
VL	DIQLTQSPSSLSASVGDRVTTITCRASSGVSYIYWFQQKPGKAPKPLIYAAFNLASGVPSRFSGSGSGTEY TLTISSLQPEDFATYYCQQWSNNPLTFGQGTKVEIK (SEQ ID NO: 492)
VL	DIQLTQSPSILSASVGDRVTTITCRASSSVSYIYWFQQKPGKAPKPLIYATFNLASGVPSRFSGSGSGTSY TLTISSLQPEDFATYYCQQWSNNPLTFGQGTKVEIK (SEQ ID NO: 493)

Group G	
VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWVRQAPGQGLEWMGWISAYNGNTNYAQLQGRVTM TTDTSTSTAYMELRSLRSDDTAVYYCARALPSGTILVGGWFDWPWGQGTTLTVTVSS (SEQ ID NO: 494)
VH	EVQLVQSGGGVVPGRSLRLSCAASGFTFSSYALSWVRQAPGKGLEWVSAISGGGGSTYYADSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCAKDVFPETFSMNYGMDVWGQGTTLTVTVSS (SEQ ID NO: 495)
VH	QVQLVQSGGGVVPGGSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSLISGDGGSTYYADSVKGRFTI SRDNSKNSLYLQMNSLRTEDTALYYCAKVLLPCSSTSCYGSVGAFDIWGQGTTLTVTVSS (SEQ ID NO: 496)
VH	QVQLVQSGGSVVRPGESLRLSCVASGFIFDNYMSWVRQVPGKGLEWVSRVNWNGGSTTYADAVKGRFTI SRDNTKNSLYLQMNRLRAEDTAVYYCVREFVGAYDLWGQGTTLTVTVSS (SEQ ID NO: 497)
VH	QVQLVQSGAEVKKPGATVKVSCKVFGDTFRGLYIHWVRQAPGQGLEWMGGIIPIFGTANYAQKFQGRVTI TTDESTSTAYMELSSLRSED ^T AVYYCASGLRWGIWGWFDWPWGQGTTLTVTVSS (SEQ ID NO: 498)
VH	EVQLVQSGAELKKPGSSVKVSCKAFGGTFSDNAISWVRQAPGQGP ^{EW} MGGIIPIFGKPNYAQKFQGRVTI TAD ^{EST} STAYMVLSSLRSED ^T AVYYCARTMVRGFLGMDVWGQGTTLTVTVSS (SEQ ID NO: 499)
VH	QVQLVQSGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCAKDQFVTIFGVPRYGMDVWGQGTTLTVTVSS (SEQ ID NO: 500)
VH	QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGQGLEWMGGIIPIFGTANYAQKFQGRVTI TADKSTSTAYMELSSLRSED ^T AVYYCARGRQMFGAGIDFWGPGTLTVTVSS (SEQ ID NO: 501)
VH	EVQLVESGAEVKKPGSSVKVSCKVSGGTFGTALNWVRQAPGQGLEWMGRIVPLIGLVNYAHNFEGRISI TADKSTGTAYMELSNLRSDDTAVYYCAREVYGGNSDYWGQGTTLTVTVSS (SEQ ID NO: 502)
VH	QVQLVQSGGEVKKPGASVKVSCKASGYTLSSHGITWVRQAPGQGLEWMGWISAHNGHASNAQKVEDRVTM TTDTSTNTAYMELRSLTADDTAVYYCARVHAALYYGMDVWGQGTTLTVTVSS (SEQ ID NO: 503)
VH	QVQLQESGGGVVPGRSLRLSCSASGFTFSRHGMHWVRQAPGKGLEWVAVISHDGSVKYYADSMKGRFSI SRDNSNNTLYLQMDSLRADDTAVYYCARGLSYQVSGWFDWPWGQGTTLTVTVSS (SEQ ID NO: 504)
VH	NFMLTQPHSVSESPGKTVTISCTRSSGSIASNYVQWYQQRPGSSPTTVIYEDNQRP ^{SG} VPDRFSGSIDTS SNSASLTISGLKTKDEADYYCQSYDGITVIFGGGTKLTVL (SEQ ID NO: 505)
VH	NFMLTQPHSVSGSPGKTVTLPC ^{TR} SSGSIASHYVQWYQQRPGSAPT ^{TV} IYEDNKRPSGVPDRFSGSIDSS SNSASLSISGLKTEADYYCQSYDSSNRWVFGGGTKLTVL (SEQ ID NO: 506)

VH	LPVLTQPASLSASPGASASLTCTLRSGLNVSRYIYQYQQKPGSRPQYLLNYKSDSNKQQASGVPSRFSG SKDASANAGILLISGLQSEDEADYYCMIWYSSAVVFGGGTKLTVL (SEQ ID NO: 507)
VL	NFMLTQPHSVSESPGKTVTITISCTRSSGNIASNYVQWYQQRPGSAPT ^{TVIY} <u>EDNQRP</u> SGVPDRFSGSIDSS SNSASLTISGLKTEDEADYYCQSYDSSNLWVFGGGTKLTVL (SEQ ID NO: 508)
VL	SSELTQDPAVSVALGQTVRITCQGD ^{SLRSY} ASWYQQKPGQAPVLVIYGKNRPSGIPDRFSGSSSGNTA SLTITGAQAEDADYYCNSRDSSGNHYVFGTGKVTVL (SEQ ID NO: 509)
VL	LPVLTQAPSVSVAPGKTARITCGGSDIGRKS ^{VH} WYQQKPGQAPALVIYSDRDRPSGISERFSGSNSGNTA TLTISRVEAGDEADYYCQVWDNNSDHVFGAGTELIVL (SEQ ID NO: 510)
VL	QSALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMYDVSNRPSGVSNRFSGSKSG NTASLTISGLQAEDADYYC ^{SSYT} SS ^{TL} PFGGGTKLTVL (SEQ ID NO: 511)
VL	EIVLTQSPATLSLSPGERATLSCRASQSIGNSLAWYQQKPGQAPRLLMYGASSRATGIPDRFSGSGAGTD FTLTIS ^{SLEPED} FATYYCQQHTIPTFSFGPGTKVEVK (SEQ ID NO: 512)
VL	DIVMTQTPSFLSASIGDRV ^{TIT} CRASQIGISYLA ^{WY} QQRPGEAPKLLIYA ^{ASTL} QSGVPSRFSGSGSGTD FTLTISNLQPEDFATYYCQQLNNYPITFGQGTRLEIK (SEQ ID NO: 513)
VL	QSALTQPPSVSVSPGQTANIPCSGDKLGK ^{YAYW} YQQKPGQSPVLLIYQDIKRPSRIPERFSGSNSADTA TLTISGTQAMDEADYYCQ ^{TWDN} SVVFGGGTKLTVL (SEQ ID NO: 514)
VL	NFMLTQPHSVSESPGKTVTITISCTRSSGSIDS ^{NYVQWY} QQRPGSAPT ^{TVIY} <u>EDNQRP</u> SGVPDRFSGSIDSS SNSASLTISGLKTEDEADYYCQSYDSNNRHVIFGGGTKLTVL (SEQ ID NO: 515)
VL	NFMLTQPHSVSESPGKTVTITISCTRSSGNIGTNYVQWYQQRPGSAPVALIYEDYRRPSGVPDRFSGSIDSS SNSASLIISGLKPEDEADYYCQSYHSSGWEFGGGTKLTVL (SEQ ID NO: 516)
VL	QSVLTQPPSVSVAPGQTARITCGGNIGSKGVH ^{WY} QQKPGQAPVLVYDDSDRPSGIPERFSGSNSGNTA TLTISRVEAGDEADYYCQVWDSSSDHWVFGGGTKLTVL (SEQ ID NO: 517)
VL	NFMLTQPHSVSESPGKTVTITISCTRSSGSIASNYVQWYQQRPGSAPT ^{TVIY} <u>EDNQRP</u> SGVPDRFSGSIDSS SNSASLTISGLKTEDEADYYCQSYDSTTPSVFGGGTKLTVL (SEQ ID NO: 518)
VL	QVQLVQSGAEVKKPGASVKV ^{SCKASGYTFTSYGISWVRQAPGQGLEWMGWTS} PHNGLTAF ^{AQILEGRVTM} TTDTSTNTAYMELRNLTFDDTAVYFCAKVHPVFSYALDVWGQGT ^{LVT} VSS (SEQ ID NO: 519)
VL	EVQLVESGA ^{EV} MNP ^{GSSVRV} SCRGSGGDFSTYAF ^{SWVRQAPGQGLEWMGRIIPILGIAN} YAQKFQGRVTI TADKSTSTAYMELSSLRSDDTAVYYCARDGYGSDPVLWGQGT ^{LVT} VSS (SEQ ID NO: 520)
VL	EVQLVQSGAEVKKPGASVKV ^{SCKASGYTFTNYGISWVRQAPGQGLEWMGWISAY} NGNTNYAQKVQGRVTM TTDTSTSTGYMELRSLRSDDTAVYYCARGDFRKPFDYWGQGT ^{LVT} VSS (SEQ ID NO: 521)
Group H	
VH	EVQLVQSGPELKKPGASVKMSCKASGYTFT ^{SYVMHWVKQAPGQRLEWIGYVNP} FNDGTYNEMFKGRATL TSDKSTSTAYMELSSLRSEDSAVYYCARQAWGY ^{PWGQGT} LVTVSS (SEQ ID NO: 522)
VH	EVQLVQSGAEVKKPGASVKMSCKASGYTFT ^{SYVMHWVKQAPGQRLEWIGYVNP} FNDGTYNEMFKGRATL TSDKSTSTAYMELSSLRSEDTAVYYCARQAWGY ^{PWGQGT} LVTVSS (SEQ ID NO: 523)
VH	EVQLVQSGAEVKKPGASVKMSCKASGYTFT ^{SYVMHWVRQAPGQRLEWIGYVNP} FNDGTYNEMFKGRATL TSDKSTSTAYMELSSLRSEDTAVYYCARQAWGY ^{PWGQGT} LVTVSS (SEQ ID NO: 524)
VH	EVQLVQSGAEVKKPGASVKV ^{SCKASGYTFTSYVMHWVRQAPGQRLEWIGYVNP} FNDGTYNEMFKGRATL TSDKSTSTAYMELSSLRSEDTAVYYCARQAWGY ^{PWGQGT} LVTVSS (SEQ ID NO: 525)

VH	EVQLVQSGAEVKKPGASVKVSCKASGYTFTSYVMHWVRQAPGQRLEWIGYVNPFDGTYNEMFKGRATI TSDKSTSTAYMELSSLRSEDTAVYYCARQAWGYPWGQGTTLTVSS (SEQ ID NO: 526)
VL	DIVLTQSPASLALSPGERATLSCRATESVEYYGTSLVQWYQQKPGQPPKLLIYAASSVDSGVPSRFSGSG SGTDFTLTINSLEEEDAAMYFCQQSRRVPYTFGQGTKLEIK (SEQ ID NO: 527)
VL	DIVLTQSPATLSLSPGERATLSCRATESVEYYGTSLVQWYQQKPGQPPKLLIYAASSVDSGVPSRFSGSG SGTDFTLTINSLEAEDAAMYFCQQSRRVPYTFGQGTKLEIK (SEQ ID NO: 528)
VL	EIVLTQSPATLSLSPGERATLSCRATESVEYYGTSLVQWYQQKPGQPPKLLIYAASSVDSGVPSRFSGSG SGTDFTLTINSLEAEDAAMYFCQQSRRVPYTFGQGTKLEIK (SEQ ID NO: 529)
VL	DIVLTQSPATLSLSPGERATLSCRATESVEYYGTSLVQWYQQKPGQPPKLLIYAASSVDSGVPSRFSGSG SGTDFTLTINSLEAEDAATYFCQQSRRVPYTFGQGTKLEIK (SEQ ID NO: 530)
Group I	
VH	EVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGQGLEWMGGIIPFGTANYAQKFQGRVTI TADKSTSTAYMELSSLRSEDTAVYYCAREGTIYDSSGYSFDYWGQGTTLTVSS (SEQ ID NO: 531)
VH	EVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGQGLEWMGIINPSGGSTSYAQKFQGRVSM TRDTSTSTVYMELSSLTSEDTAVYYCARDLFPHIYGNYYGMDIWGQGTTVTVSS (SEQ ID NO: 532)
VH	QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGQGLEWMGGIIPFGTANYAQKFQGRVTI TADKSTSTAYMELSSLRSEDTAVYYCARLAVPGAFDIWGQGTMTVTVSS (SEQ ID NO: 533)
VH	EVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLAVISYDGSNKYYADSVKGRFTISR NSKNTLYLQMNSLRRAEDTAVYYCARGQWLVTLDYWGQGTTLTVSS (SEQ ID NO: 534)
VH	EVQLVESGSEVEKPGSSVKVSCKASGGTFSDSGISWVRQAPGQGLEWMGGIIPMFATPYAQKFQDRVTI TADESTSTVYMELSGLRSDDTAVFYCARDRGRGHLWPYFDLWGRGTLTVTVSS (SEQ ID NO: 535)
VH	EVQLVESGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGQGLEWMGGIIPFGTANYAQKFQGRVTI TADESTSTAYMELSSLRSEDTAVYYCARAPYYYYYMDVWGQGTTVTVSS (SEQ ID NO: 536)
VH	EVQLLESGAEVKKPGSSVKVSCKASGGTSLRYALSWVRQAPGQGPWEVGAIPFGTPHYSKKFQDRVII TVDTSTNTAFMELSSLRFDLTALYFCARGHDEYDISGYHRLDYWGQGTTLTVTVSS (SEQ ID NO: 537)
VH	QVQLVQSGSELKKPGSSVKVSCKASGYSFSGYYIHWVRQAPGQGLEWMGWIDPNSGVTNYVRRFQGRVTM TRDTSLSTAYMELSGLTADDTAVYYCARDENLWQFGYLDYWGQGTTLTVTVSS (SEQ ID NO: 538)
VH	QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSRYGVHWVRQAPGQGLEWMGRLIPIVSMNTNYAQKFQDRVSI TTDKSTGTAYMELRSLTSEDTALYYCASVGQQLPWVFFAWGQGTTLTVTVSS (SEQ ID NO: 539)
VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVRGRFTI SRDNSKNTLYLQMNSLRTEDTAVYYCARGWLDRDIDYWGQGTTLTVTVSS (SEQ ID NO: 540)
VH	EVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVRGRFTI SRDNSKNTLYLQMNSLRTEDTAVYYCARGWLDRDIDYWGQGTTLTVTVSS (SEQ ID NO: 541)
VH	EVQLVQSGGGLVQPGGSLRLSCAASGFTFSDYGMHWVRQPPGKGLEWLAVISYDGSYKIHADSVQGRFTI SRDNAKNSVFLQMNSLKTEDTAVYYCTTDRKWLAWHGMDVWGQGTTVTVTVSS (SEQ ID NO: 542)
VH	EVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGQGLEWMGGIIPFGTANYAQKFQGRVTI TADESTSTAYMELSSLRSEDTAVYYCARDGIVADFQHWGQGTTLTVTVSS (SEQ ID NO: 543)

VH	EVQLVESGAEVKKPGASVKVSCASGDTFSRYGITWVRQAPGRGLEWMGNIVPFFGATNYAQKFQGRLLTI TADKSSYTSYMDLSSLRSDDTAVYYCARDHFYGGGYFDYWGQGTLLTVVSS (SEQ ID NO: 544)
VH	EVQLLESGAEVKKPGASVKVSCASGYTFNSYDINWVRQAPQGQGLEWMGGIIPVFGTANYAESFQGRVTM TADHSTSTAYMELNNLRSEDVAVYYCARDRWHYESRPMDVWGQGTLLTVVSS (SEQ ID NO: 545)
VH	EVQLVESGGGLVLRPGGSLRLACAASGFSFSDDYMTWIRQAPGRGLEWIAVISDSGQTVHYADSVKGRFTI SRDNTKNSLFLQVNTLRAEDTAVYYCAREDLLGYLQSWGQGTLLTVVSS (SEQ ID NO: 546)
VH	QVQLQQSGPGLVKPSQTLSTCAISGDSVSSNSAAWNWIRQSPSRGLEWLGRTYYRSKWYNDYAVSVKSR ITINPDTSKNQFSLQLNSVTPEDTAVYYCARDEPRAVAGSQAYYYYGMDVWGQGTLLTVVSS (SEQ ID NO: 547)
VH	EVQLVQSGAEVKKPGASVKVSCASGYTFTSYMHVVRQAPQGQGLEWMGIINPSDGSTSYAQKFQGRVTM TRDTSTSTVHMESSLRSEDVAVYYCARDLFPFIYGNYYGMDIWGQGTLLTVVSS (SEQ ID NO: 548)
VH	QMQLVQSGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVRGRFTI SRDNSKNTLYLQMNSLRTEDTAVYYCARGWLDRDIDYWGQGTLLTVVSS (SEQ ID NO: 549)
VH	QVQLVQSGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVRGRFTI SRDNSKNTLYLQMNSLRTEDTAVYYCARGWLDRDIDYWGQGTLLTVVSS (SEQ ID NO: 550)
VL	QSVLTQPPSVSAAPGQKVTISCSGNNANIANNVSWYQQLPGTAPKLLIYDNNYRPSGIPDRFSGSKSGT SATLDITGLQTGDEADYYCGVWDGSLTTGVFGGGTKLTVL (SEQ ID NO: 551)
VL	AIQMTQSPSSLSASVGRVTITCRASQGISNYLAWYQQKPGKVPKLLIYAASSTLESQVPSRFRSGSGSGTD FTLTISLQPEDLATYYCQQLHTFPLTFGGGTKVEIK (SEQ ID NO: 552)
VL	QPVLTQPPSASGSPGQSVTISCTGTSSDVGAYNFVSWYRQHPGKAPKLMIEYVNRPSGVPDRFSGSKSG NTASLTVSGLQAEDEADYYCSSYAGTNSLGIFGTGKLTVL (SEQ ID NO: 553)
VL	QSVVTQPPSVSAAPGQKVTISCSGSSSDIGNHYVSWYQQLPGTAPKLLIYDNNQRPSGIPDRFSGSKSGT SATLAITGLQTGDEADYYCGTWDNSLSPHLLFGGGTKLTVL (SEQ ID NO: 554)
VL	QSVLTQPPSVSAAPGQKVTISCSGSSSNMGNVSWYKQVPGTAPKLLIYENDKRPSGIPDRFSGSKSGT SATLGITGLQTGDEADYYCGTWDNSLSGFVFASGTKVTVL (SEQ ID NO: 555)
VL	QSALTQPASVSGSLGQSVTISCTGTSSSDVGSYNLVSWYQQHPGKAPNLMYDVSKRPSGVSNRFSGSKSGN TASLTISGLQAEDEADYYCSSYTGSTVVFSGGKLTVL (SEQ ID NO: 556)
VL	QSVLTQPASVSGSPGQSITISCTGTSSDVGSYNLVSWYQQHPGKAPKLMIEVSKRPSGVSNRFSGSKSG NTASLTISGLQAEDEADYYCSSYGGFNNLLFGGGTKLTVL (SEQ ID NO: 557)
VL	DIVMTQSPSSLSASIGDRVTITCRASQRIAYVNWYQQKPGKAPKVLIIYAASSLRSGVPSRFRSGSGSGTD FTLTISLQPEDFATYYCQQTYSPPWTFGQGTKEIK (SEQ ID NO: 558)
VL	QSVLTQPPSASGSPGQSVTISCTGTSSDIGGYDSVSWYQQHPGKAPKLMYDVSKRPSGVSNRFSGSKSG NTASLTISGLQAEDEADYYCSSTSSIFFYVFGTGKTVL (SEQ ID NO: 559)
VL	LPVLTQPASVSGSPGQSITISCTGTSSDIGGYDVSWSYQQHPGKAPKLMYDVSKRPSGVSNRFSGSKSG NTASLTISGLQAEDEADYYCSSTSSHTVFGTGKLTVL (SEQ ID NO: 560)
VL	QSALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMYDVSNRPSGVSNRFSGSKSG NTASLTISGLQAEDEADYYCSSYRSSTLGPVFGGGTKLTVL (SEQ ID NO: 561)

VL	QAGLTQPPSVSEAPRQRTISCSGSSSNIGNNAVNWYQQLPGKAPKLLIYYDDLPSGVSDRFGSGSKSGT SASLAISGLQSEDEADYYCAAWDDSLNGYVFGTGKLTVL (SEQ ID NO: 562)
VL	QSALTQPRSVSGSPGQSVTISCTGTSSDVGGYNYVSWYQQHPGKAPKLMYDVSKRPSGVPDRFGSGSKSG NTASLTISGLQAEDEADYYCSSYTSSTTHVFGTGKVTVL (SEQ ID NO: 563)
VL	QSVVTQPPSVSAAPGQKVTISCSGSSSNIGNNYVSWYQQLPGTAPKLLIYDNNKRPSGIPDRFGSGSKSGT SATLGITGLQTGDEADYYCGTWDSSLSVWVFGGGTQLTVL (SEQ ID NO: 564)
VL	QSVLTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGRAPRLMIYDVSNRPSGVSNRFGSGSKSG NTASLTISGLQAEDEGDDYYCSSYTSGGTLGPVFGGGTKLTVL (SEQ ID NO: 565)
VL	QAGLTQPPSASGTPGQRTISCSGSSSNIGSNTVNWYQQLPGTAPKLLIYSNNQRPSGVPDRFGSGSKSGT SASLAISGLQSEDEADYYCAAWDDSLNGWVFGGGTKLTVL (SEQ ID NO: 566)
VL	AIRMTQSPSSLSASVGDRVTITCRASQSSNYLNWYQQRPGKAPNLLIYAASSLQSGVPSRFGSGSGT FTLTISLQPEDFATYYCQQTSTPYTFGQGTKLEIK (SEQ ID NO: 567)
VL	QSVLTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYRQHPGKAPKLMYDVSYPSPGVSNRFGSGSKSG NTASLTISGLQAEDEADYYCSSYTDSSSTRYVFGTGKLTVL (SEQ ID NO: 568)
VL	QPVLTPPPSASGTPGQRTISCSGSSSNIEINSVNWYQQLPGTAPKLLIYDNNKRPSGIPDRFGSGSKSGT SATLGITGLQTGDEADYYCGSWDSSLSADVFGTGKLTVL (SEQ ID NO: 569)
VL	QSVLTQPPSVSAAPGKVTISCSGSSSNIGNNYVSWYQQLPGTAPKLLIYRNNQRPSGVPDRFGSGSKSGT SASLAISGLQSEDEADYYCATWDDSLNGWVFGGGTKLTVL (SEQ ID NO: 570)
VL	QSVVTQPPSVSGAPGQRTISCTGSSSNIGAGYDVHWYQQLPGTAPKLLIYGNNNRHSGVPDRFGSGSKSG TSASLAITGLQAEDEAEFFCGTWDSRLTTYVFGSGTKLTVL (SEQ ID NO: 571)
VL	QSVVTQPPSVSAAPGQKVTISCSGSSSNIGNNYVSWYQQLPGTAPKLLIYDNNKRPSGIPDRFGSGSKSGT SATLGITGLQTGDEADYYCGTWDSSLSAVVFGGGTKLTVL (SEQ ID NO: 572)
VL	VIWMTQSPSSLSASVGDRVTITCAASSLQSWYQQKPGKAPKLLIYEASTLESVPSRFGSGSGTEFTLT ISSLQPEDFATYYCQQSYSTPYTFGQGTKLEIK (SEQ ID NO: 573)
VL	QSVVTQPPSVSAAPGQKVTISCSGSSSNIGNNYVSWYQQVPGTAPKLLIYDNNKRPSGIPDRFGSGNSDT SATLGITGLQTGDEADYYCGTWDSSLSAWVFGGGTKLTVL (SEQ ID NO: 574)
VL	QSVVTQPPSVSAAPGQKVTISCSGSSSNIGNNYVSWYQQLPGTAPKLLIYDNNKRPSGIPDRFGSGSKSGT SATLGITGLQTGDEADYYCGTWDSSLSAGSVVFGGGTKLTVL (SEQ ID NO: 575)
VL	SYELMQPPSVSVAPGKTATIACGGENIGRKTVHWYQQKPGQAPVLVIYYDSDRPSGIPERFGSGNSGNTA TLTISRVEAGDEADYYCLVWDSSSDHRIFGGGKLTVL (SEQ ID NO: 576)
VL	SYELMQPPSVSVAPGKTATIACGGENIGRKTVHWYQQKPGQAPVLVIYYDSDRPSGIPERFGSGNSGNTA TLTISRVEAGDEADYYCQVWDSSSDHRIFGGGKLTVL (SEQ ID NO: 577)
VL	SYELMQPPSVSVAPGKTATIACGGENIGRKTVHWYQQKPGQAPVLVIYYDSDRPSGIPERFGSGNSGNTA TLTISRVEAGDEADYYCQVWDSSSDHRIFGGGKLTVL (SEQ ID NO: 578)
VL	SYELMQPPSVSVAPGKTATIACGGENIGRKTVHWYQQKPGQAPVLVIYYDSDRPSGIPERFGSGNSGNTA TLTISRVEAGDEADYYCQVWDSSSDHRIFGGGKLTVL (SEQ ID NO: 579)
Group J*	

HC	QVQLVQSGVEVKKPGASVKVSCKASGYTFTNYYMYWVRQAPGQGLEWMGGINPSNGGTNFNEKFKNRVTL TTDSSTTTAYMELKSLQFDDTAVYYCARRDYRFDMGFDYWGGQTTVTVSSASTKGPSVFPLAPCSRSTSE STAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYTCNVDPKPS NTKVDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDSQEDPEVQFNWYVD GVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYT LPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEG NVFSCSVMEALHNHYTQKSLSLGLGK (SEQ ID NO: 580)
HC	QVQLVESGGGVVQPGSRSLRLDCKASGITFSNSGMHWVRQAPGKGLEWVAVIWDGSKRYYADSVKGRFTI SRDNSKNTLFLQMNSLRAEDTAVYYCATNDYWGQGTLLTVTVSSASTKGPSVFPLAPCSRSTSESTAALGC LVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYTCNVDPKPSNTKVDKR VESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDSQEDPEVQFNWYVDGVEVHNA KTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEE MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSV MEALHNHYTQKSLSLGLGK (SEQ ID NO: 581)
LC	EIVLTQSPATLSLSPGERATLSCRASKGVSTSGYSYLHWYQQKPGQAPRLLIYLAstyleSGVPARFSGSG SGTDFTLTISLLEPEDFAVYYCQHSRDLPLTFGGGTKEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLL NNFYPREAKVQWKVDNALQSGNSQESVTEQDSKSTYSLSSSTLTLSKADYEKHKVYACEVTHQGLSSPVT KSFNRGEC (SEQ ID NO: 582)
LC	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDASNRATGIPARFSGSGSGTD FTLTISLLEPEDFAVYYCQSSNWPRTFGQGTKEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFY PREAKVQWKVDNALQSGNSQESVTEQDSKSTYSLSSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFN RGEC (SEQ ID NO: 583)
Group K	
VH	EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHWRQAPGKGLEWVAWISPYGGSTYYADSVKGRFTI SADTSKNTAYLQMNSLRAEDTAVYYCARRHWPGGFDYWGGQGTLLTVTVSSASTK (SEQ ID NO: 735)
VH	EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHWRQAPGKGLEWVAWISPYGGSTYYADSVKGRFTI SADTSKNTAYLQMNSLRAEDTAVYYCARRHWPGGFDYWGGQGTLLTVTVSS (SEQ ID NO: 736)
HC	EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHWRQAPGKGLEWVAWISPYGGSTYYADSVKGRFTI SADTSKNTAYLQMNSLRAEDTAVYYCARRHWPGGFDYWGGQGTLLTVTVSSASTKGPSVFPLAPSSKSTSGGT AALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHPKPSNT KVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDSHEDPEVKFNWYV DGVEVHNAKTKPREEQYASTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY TLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQ GNVFSCSVMEALHNHYTQKSLSLSPG (SEQ ID NO: 737)
VL	DIQMTQSPSSLSASVGRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSFGSGSGTD FTLTISLQPEDFATYYCQYLYHPATFGQGTKEIKR (SEQ ID NO: 268)

LC	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSGSGTD FTLTISSLQPEDFATYYCQQYLYHPATFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFY PREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFN RGEC (SEQ ID NO: 738)
Group L	
VH	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRFWMSWVRQAPGKGLEWVANINQDGTEKYYVDSVKGRFTI SRDNAKNSLYLQMNSLRAGDTAVYYCANTYYDFWSGHFDYWGQGTLLTVVSS (SEQ ID NO: 739)
VH	QEHLVESGGGVVQPGSLRLSCEASGFTFSNFGMHVVRQAPGKGLEWVAALWSDGSNKYYADSVKGRVTI SRDNSKNTLYLQMNSLRAEDTAVYYCARGRGAPGIPFGYWGQGTLLTVVSS (SEQ ID NO: 740)
VH	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLEWVGRIKRKTDGGTTDYAAPVKGRF TISRDDSKNTLHLQMNSLKTEDTAVYYCTTDDIVVVPVAVMREYYFGMDVWGQGTTLTVVSS (SEQ ID NO: 741)
VH	QVQLVQSGAEVKKPGASVQVSCASGYSTGYIHWVRQAPGQGLEWMGWINPNSGTKKYAHKFQGRVTM TRDTSIDTAYMILSSLISDDTAVYYCARDEDWNFGSWFDSWGQGTLLTVVSS (SEQ ID NO: 742)
VH	QVHLVQSGAEVKKPGASVKVSCASGYTFTGYIHWVRQAPGHGLEWMGWLNPNTGTTKYIQNFQGRVTM TRDTSSSTAYMELTRLRSDDTAVYYCARDEDWNYGSWFDTWGQGTLLTVVSS (SEQ ID NO: 743)
VH	EVQLVESGGGVVRPGGSLRLSCAASGFTFDDYGMTWVRQAPGRGLEWVSGIHWHGKRTGYADSVKGRFTI SRDNAKNSLYLQMNSLKGEDTALYHCVRGGMSTGDWFDWPWGQGTLLVIVSS (SEQ ID NO: 744)
VH	EVQLVESGGGVVRPGGSLRLSCAASGFTFDDYGMTWVRQVPGKGLEWVSGIHWSGRSTGYADSVKGRFTI SRDNAKNSLYLQMNSLRAEDTALYYCARGGMSTGDWFDWPWGQGTLLTVVSS (SEQ ID NO: 745)
VH	EVQLVESGGGLVQPGGSLRLSCAASGFTVGSNYMNVVRQAPGKGLEWVSVIYSGGSTYYADSVKGRFTIS RLTSKNTLYLQMNSLRPEDTAVYYCARGIRGLDVWGQGTTLTVVSS (SEQ ID NO: 746)
VH	EERLVESGGDLVQPGGSLRLSCAASGITVGTNYMNVVRQAPGKGLEWVSVISSGGNTHYADSVKGRFIMS RQTSKNTLYLQMNSLLETEDTAVYYCARGIRGLDVWGQGTMTVTVSS (SEQ ID NO: 747)
VH	QVQLVQSGAEVKMPGSSVRVSCASGGIFSSSTISWVRQAPGQGLEWMGEIIPVFGTVNYAQKFQDRVIF TADESTTTAYMELSSLKSGDTAVYFCARNWGLGSFYIWGQGTMTVTVSS (SEQ ID NO: 748)
VH	EVQLVESGGDLVHPGRSLRLSCAASGFPPFDEYAMHWVRQVPGKGLEWVSGISWSNNNIGYADSVKGRFTI SRDNAKNSLYLQMNSLRPEDTAFYYCAKSGIFDSWGQGTLLTVVSS (SEQ ID NO: 749)
VH	EVQLVESGGGVVQPGSLRLSCAASGFTFSSYGMHWVRQAPGKGLEWVTLLISYEGRNKYYADSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCAKDRTLYGMDVWGQGTTLTVVSS (SEQ ID NO: 750)
VH	QVTLRESGPALVKTTQTTLTCTFSGFSLSTNRMCVTWIRQPPGKALEWLARIDWDGVKYYNTSLKTRLT ISKDTSKNQVVLMTNMDPVDATFYCARSTSLTFYYFDYWGQGTLLTVVSS (SEQ ID NO: 751)
VH	EVQLVESGGGLVQPGGSLRLSCAASEFTVGTNHNWVRQAPGKGLEWVSVIYSGGNTFYADSVKGRFTIS RHTSKNTLYLQMNSLTAEDTAVYYCARGLGMDVWGQGTTLTVVSS (SEQ ID NO: 752)
VH	EVQLVESGGGLVQRGESLRLYCAASGFTFSKYWMNVVRQAPGKGLEWVANIKGDGSEKYYVDSVKGRFTI SRDNAKNSLYLQMNSLRAEDTAVYYCARDYWGSGYYFDWFGQGTLLTVVSS (SEQ ID NO: 753)
VH	EVQLVESGGGLVQSGGSLRLSCAASGFTFSSYWMSWVRQAPGKGLEWVANIKQDGSEKYYVDSVKGRFTI SRDNAKNSLYLQMNSLRADDTAVYYCARDDIVVVPAPMGYYYYYFGMDVWGQGTTLTVVSS (SEQ ID NO: 754)

VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDFAMHWVRQAPGKGLEWVSGISWTGGNMDYANSVKGRFTI SREDAKNSLYLQMNSLRAADTALYYCVKDIRGIVATGGAFDIWGRGTMVTVSS (SEQ ID NO: 755)
VH	EVQLVESGGGLVQPGGSLRLSCAASGFTVGTNYMNVWRQAPGKGLEWISVIYSGGSTFYADSVKGRFTIS RQTSQNTLYLQMNSLRPEDTAVYYCARGIRGFDIWGQGTMVTVSS (SEQ ID NO: 756)
VH	EVQLVESGGGLVQPGGSLRLSCAASGFTISTNYMNVWRQAPGKGLEWVAVIYSSGSTYYIDSVKGRFTIS RLTSKNTVYLMSSLNSED ^T AVYYCARGIRGFDIWGQGTMVTVSS (SEQ ID NO: 757)
VH	EVQLVESGGGLVQPGRSLRLSCAASGFTIDDSAMHWVRQTPGKGLEWVSGISWKSGSIGYADSVRGRFTI SRDNAKNSLYLQMNSLRVEDTALYYCVKDIRGNWNYGGNWFD ^W PWGQGLTVTVSS (SEQ ID NO: 758)
VH	EVQLVESGGGLVQPGGSLRLSCEASGFTVGVNHMNVWRQAPGKGLEWVSVIFSSGRTFYGDYVKGRLTIF RQTSQNTVYLMNSLRSED ^T AIYYCARGIGGLDIWGRGTMVTVSS (SEQ ID NO: 759)
VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYALHWVRQAPGKGLEWVSGISWTGGTIDYADSVKGRFTI SRDNAKNSLYLQMSSLRTED ^T AIYYCTRDIRGNWKYGGWFD ^W PWGQGLTVTVSS (SEQ ID NO: 760)
VH	QVQLVQSGTEVKKPGASVKVSCKASGYTFTAYYMHWVRQAPGQGLDWMGWISPNSGFTNYAQKFQGRVTM TRDTSINTFYMELSGLRSDDTAVYYCAREGSTHHNSFDPWGQGLTVTVSS (SEQ ID NO: 761)
VH	EVQLVESGGGLVQPGGSLRLSCAASGFTVGTNFMNVWRQAPGKGLEWVSAIYSGGTANYADSVKGRFTIS RDTSRNTLYLQMNSLRTE ^D TAVYYCARGGGMDVWGQGT ^T TVTVSS (SEQ ID NO: 762)
VH	QVQLVQSGAEVKKPGSSVKVSCKASGGTFNTYVLSWVRQAPGQGLEWMGEIIPILGAANYAQNFQGRVTF TTDESTNTAYMDLSSLRSED ^T AVYYCARDRTSGGFDPWGQGLTVTVSS (SEQ ID NO: 763)
VH	QVQLVQSGAEVEKPGASVKVSCKASGYIFTHYGISWVRQAPGQGLEWVGWISPYNGYTDYAQKLQGRVTL TTDTSTTTAYMELRNLRSDDTAMYYCSRGRGPYWSFDLWGRGTLTVTVSS (SEQ ID NO: 764)
VL	DIQMTQSPSTLSASVGDRVTITCRASQSISNWLAWYQQKPGKAPKLLIYKASSLES ^G VPSRFSGSGSGTE FTLTIS ^S LQPD ^D FATYYCQQYHSYSYTFGQGTKEIK (SEQ ID NO: 765)
VL	DIQMTQSPSSLSASVGDRVTITCRASQGIRNDLGWYQQKPGKAPKRLIYTASSLQSGVPSRFSGSGSGTE FTLTIS ^S LQPED ^D FATYYCLQHNSYPLTFGGGTKVAIK (SEQ ID NO: 766)
VL	DIQMTQSPSSLSASVGDRVTITCR ^T TSQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVPSRFSGSGSGTE FTLTIS ^S LQPED ^D FATYYCLQHNNYPYTFGQGTKLEIK (SEQ ID NO: 767)
VL	DIVMTQTPLSSPVTLGQPASISCRSSQTLVHGDGNTYLSWIQQRPGQPPRLLIYKVS ^N QFSGVPDRFSGS GAGTDFTLKISRVEADVGLYFCMQATHFPITFGQGTRLEIK (SEQ ID NO: 768)
VL	DIVMTQTPLSSPVTLGQPASISCRSSPSLVHSDGNTYLSWLQQRPGQPPRLLIYKISNRFS ^G VPDRFSGS GAGTDFTLKISRVEADVGVYYCMQATHFPITFGQGTRLEIR (SEQ ID NO: 769)
VL	DIQMTQSPSSLSASLGDRVTITCRASQSINSYLNWYQQKPGKAPKLLIYVASSLQSGVPSRFSGSGSGTE FTLTIS ^N LQPED ^D FATYYCQQSYSTPPITFGQGTRLEIK (SEQ ID NO: 770)
VL	DIQMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYVASSLQSGVPSRFSGSGSGTD FTLTIS ^S LQPED ^D FATYYCQQSYSTPPITFGQGTRLEIK (SEQ ID NO: 771)
VL	DIQMTQSPSSLSASVGDRVTITCRASQTINIYLNWYQQKPGRAPRLLIYAASSLQSGVPSRFSGSGSGTD FTLTIS ^S LQPED ^D FATYYCHQSYSTPPITFGQGTRLEIK (SEQ ID NO: 772)
VL	DIQMTQSPSSLSASVGDRVTITCRASQSMSSYLNWYQQKPGRAPKLLIFAASSLQSGVPSRFSGSGSGTD FTLTIS ^S LQPED ^D FATYYCQQSYSTPPITFGQGTRLEIK (SEQ ID NO: 773)

VL	EIVLTQSPGTL SL SPGERATLSCRASQ S FNFN Y LAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGT DFTLTINRLEPEDFGVFY C Q Q YESAPWTFGQGTKVEIK (SEQ ID NO: 774)
VL	DIQMTQSPSSLSASVGDRVTITCRASQ S ISSYLN W YQQKPGKLLIYAASSLQSGVPSRFRSGGGSGTDFTL TISSLRPEDFATYYC Q QSYCTPPITFGQGTRLEIK (SEQ ID NO: 775)
VL	DIQMTQSPSSLSASVGDRVTITCRASQ S ISSYLN W YQQKPGKAPKLLIYAASSLQSGVPSRFRSGSGSGTD FTLTISLQPEDFATYYC Q QSYSTPPITFGQGTRLEIK (SEQ ID NO: 776)
VL	DRVITITCRASQ V ISN Y LAWYQQKPGKVPRLIYAAS T LQSGVPSRFRSGSGSGTDFTLTISLQPEDVATY YCQKYNAPRTFGQGTKVEIK (SEQ ID NO: 777)
VL	DIQMTQSPSSLSASVGDRVTITCRASQ N INNYLN W YQQKPGKAPKLLIYAASSFQNAVPSRFRSGSGSGTD FTLTISLQPEDFATYYC Q QSYNTPLTFGGGTKVEIK (SEQ ID NO: 778)
VL	DIQMTQSPSSLSASVGDRVTITCRASQ G IRNDL G WYQQKPGKAPKRLIYAASSLQSGVPSRFRSGSGSGTE FTLTISLQPEDFATYYC L QHNSYPYTFGQGTKLEIK (SEQ ID NO: 779)
VL	DIQMTQSPSSLSASVGDRVTITCRASQ S ISSYLN W YQQKPGKAPKLLIYAASSLQSGVPSRFRSGSGSGTD FTLTISLQPEDFATYYC Q QSYSTPPITFGQGTRLEIK (SEQ ID NO: 780)
Group M	
VH	QSLEESGGRLVKPDETLTITCTVSGIDLSSNGLTWVRQAPGEGLEWIGTINKDASAYYASWAKGRLTISK PSSTKVDLKITSPTTEDTATYFCGRIAFKTGT S IWGPGLVTVSS (SEQ ID NO: 1108)
VL	AIVMTQTPSPVSAAVGGTVTINCQASESVYSNNYLSWFQQKPGQPPKLLIYLASTLASGVPSRFRKSGSGG TQFTLTISGVQCDDAATYYCIGGKSSSTDGNAFGGGTEVVVR (SEQ ID NO: 1109)
Group N	
VH	QMQLVQSGAEVKKPGSSSVKVSKASGGTFSSYAISWVRQAPGQGLEWMGGIIPFGTANYAQKFQGRVTI TADSTSTAYMELSSLRSED T AVYYCARGNIVATITPLDYWGQGLVTVSS (SEQ ID NO: 1110)
VH	QPVLTPPPSVSAAPGQKVTISCSGSSSNIANNVSWYQQLPGTAPKLLIFANNKRPSGIPDRFSGSKSGT SAALDITGLQTGDEADYYCGTWDSDLRAGVFVGGGTKLTVL (SEQ ID NO: 1111)
VH	EVQLVQSGAEVKKPGSSSVKVSKASGGTFSSYAISWVRQAPGQGLEWMGGIIPFGTANYAQKFQGRVTI TADKSTSTAYMELSSLRSED T AVYYCAREGTIYDSSGYSFDYWGQGLVTVSS (SEQ ID NO: 1112)
VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVRGRFTI SRDNSKNTLYLQMNSLRTE D AVYYCARGWLD R DIDYWGQGLVTVSS (SEQ ID NO: 1113)
VH	EVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVRGRFTI SRDNSKNTLYLQMNSLRTE D AVYYCARGWLD R DIDYWGQGLVTVSS (SEQ ID NO: 1114)
VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVRGRFTI SRDNSKNTLYLQMNSLRTE D AVYYCARGWLD R DIDYWGQGLVTVSS (SEQ ID NO: 1115)
VH	EVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVRGRFTI SRDNSKNTLYLQMNSLRTE D AVYYCARGWLD R DIDYWGQGLVTVSS (SEQ ID NO: 1116)
VH	EVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVRGRFTI SRDNSKNTLYLQMNSLRTE D AVYYCARGWLD R DIDYWGQGLVTVSS (SEQ ID NO: 1117)

VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVRGRFTI SRDNSKNTLYLQMNSLRTEDTAVYYCARGWLDRLDIDYWGQGLTVTVSS (SEQ ID NO: 1118)
VH	QVQLVQSGAEVKKPGSSVKVSKASGGTFSSRYGVHWVRQAPGQGLEWMGRLIPIVSMNTNYAQKFQDRVSI TTDKSTGTAYMELRSLTSEDALYYCASVGQQLPWVFFAWGQGLTVTVSS (SEQ ID NO: 1119)
VH	QMQLVQSGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVRGRFTI SRDNSKNTLYLQMNSLRTEDTAVYYCARGWLDRLDIDYWGQGLTVTVSS (SEQ ID NO: 1120)
VH	QVQLVQSGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISFDGSNKYYADSVRGRFTI SRDNSKNTLYLQMNSLRTEDTAVYYCARGWLDRLDIDYWGQGLTVTVSS (SEQ ID NO: 1121)
VH	QMQLVQSGAEVKKPGSSVKVSKASGGTFSSYAYSWVRQAPGQGLEWMGGIIPISFGTANYAQKFQGRVTI TADESTSTAYMELSSLRSEDTAVYYCARGPIVATITPLDYWGQGLTVTVSS (SEQ ID NO: 1122)
VH	QMQLVQSGAEVKKPGSSVKVSKASGGTFSSYAYSWVRQAPGQGLEWMGGIIPISFGTANYAQKFQGRVTI TADESTSTAYMELSSLRSEDTAVYYCARGPIVATITPLDYWGQGLTVTVSS (SEQ ID NO: 1123)
VH	QMQLVQSGAEVKKPGSSVKVSKASGGTFSSYAYSWVRQAPGQGLEWMGGIIPISFGTANYAQKFQGRVTI TADESTSTAYMELSSLRSEDTAVYYCARGPIVATITPLDYWGQGLTVTVSS (SEQ ID NO: 1124)
VH	QMQLVQSGAEVKKPGSSVKVSKASGGTFSSYAISWVRQAPGQGLEWMGGIIPAFGTANYAQKFQGRVTI TADESTSTAYMELSSLRSEDTAVYYCARGPIVATITPLDYWGQGLTVTVSS (SEQ ID NO: 1125)
VL	SYELMQPPSVSVAPGKTATIACGGENIGRKTVHWYQQKPGQAPVLVIYYDSDRPSGIPERFSGSNSGNTA TLTISRVEAGDEADYYCQVWDSSSDHRIFGGGTKLTVL (SEQ ID NO: 1126)
VL	AIRMTQSPSSLSASVGRVTITCRASQSISSYLNWYQQKPGKAPKLLIYTTSSSLKSGVPSRFRSGSGSGTD FTLTISRLQPEDFATYYCQQSYSSWTFRGRTKVEIK (SEQ ID NO: 1127)
VL	QSVLTQPPSVSAAPGQKVTISCSGNNIANNVSWYQQLPGTAPKLLIYDNNYRPSGIPDRFSGSKSGT SATLDITGLQTGDEADYYCGVWDGSLTTGVFGGGTKLTVL (SEQ ID NO: 1128)
VL	LPVLTQPASVSGSPGQSITISCTGTSSDIGGYDYVSWYQQHPGKAPKLMYDVSKRPSGVSNRFSGSKSG NTASLTISGLQAEDEADYYCSSYTSSSTHVFGTGTKLTVL (SEQ ID NO: 1129)
VL	QSALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMYDVSNRPSGVSNRFSGSKSG NTASLTISGLQAEDEADYYCSSYRSTLGPVFGGGTKLTVL (SEQ ID NO: 1130)
VL	QAGLTQPPSVSEAPRQRTISCSGSSNIGNNAVNWYQQLPKGKAPKLLIYDDLLPSGVSDRFSGSKSGT SASLAISGLQSEDEADYYCAAWDDSLNGYVFGTGTKLTVL (SEQ ID NO: 1131)
VL	QSALTQPRSVSGSPGQSVTISCTGTSSDVGGYNYVSWYQQHPGKAPKLMYDVSKRPSGVDPDRFSGSKSG NTASLTISGLQAEDEADYYCSSYTSSSTHVFGTGTKVTVL (SEQ ID NO: 1132)
VL	QSVVTQPPSVSAAPGQKVTISCSGSSNIGNNYVSWYQQLPGTAPKLLIYDNNKRPSGIPDRFSGSKSGT SATLGITGLQTGDEADYYCGTWDSSLSVWVFGGGTQLTVL (SEQ ID NO: 1133)
VL	QSVLTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGRAPRLMIYDVSNRPSGVSNRFSGSKSG NTASLTISGLQAEDEADYYCSSYTSGGTLGPVFGGGTKLTVL (SEQ ID NO: 1134)
VL	QSVVTQPPSVSAAPGQKVTISCSGSSNIGNNYVSWYQQLPGTAPKLLIYDNNKRPSGIPDRFSGSKSGT SATLGITGLQTGDEADYYCGTWDSSLSAVVFGGGTKLTVL (SEQ ID NO: 1135)
VL	QSVVTQPPSVSAAPGQKVTISCSGSSNIGNNYVSWYQQVPGTAPKLLIYDNNKRPSGIPDRFSGSNSDT SATLGITGLQTGDEADYYCGTWDSSLSAVVFGGGTKLTVL (SEQ ID NO: 1136)

VL	QSVVTQPPSVSAAPGQKVTISCSGSSSNIGNNYVSWYQQLPGTAPKLLIYDNNKRPSGIPDRFSGSKSGT SATLGITGLQTGDEADYYCGTWDSSLSAGSVVFGGGTKLTVL (SEQ ID NO: 1137)
VL	SYELMQPPSVSVAPGKTATIACGGENIGRKTVHWYQQKPGQAPVLVIYYDSDRPSGIPERFSGSNSGNTA TLTISRVEAGDEADYYCLVWDSSSDHRIFGGGKLTTL (SEQ ID NO: 1138)
VL	SYELMQPPSVSVAPGKTATIACGGENIGRKTVHWYQQKPGQAPVLVIYYDSDRPSGIPERFSGSNSGNTA TLTISRVEAGDEADYYCQVWDSSSDHRIFGGGKLTTL (SEQ ID NO: 1139)
VL	SYELMQPPSVSVAPGKTATIACGGENIGRKTVHWYQQKPGQAPVLVIYYDSDRPSGIPERFSGSNSGNTA TLTISRVEAGDEADYYCQVWDSSSDHRIFGGGKLTTL (SEQ ID NO: 1140)
VL	SYELMQPPSVSVAPGKTATIACGGENIGRKTVHWYQQKPGQAPVLVIYYDSDRPSGIPERFSGSNSGNTA TLTISRVEAGDEADYYCQVWDSSSDHRIFGGGKLTTL (SEQ ID NO: 1141)
Group O	
VH	QVQLVQSGSEVKKSGSSVKVCKTSGGTFSITNYAINWVRQAPGQGLEWMGGILPIFGAAKYAQKFQDRV TITADESTNTAYLELSSLTSEDAMYYCARGKRWLQSDLQYWGQGLVTVSS (SEQ ID NO: 1142)
VL	QPVLTPASVSGSPGQSITISCTGSSSDVGSYDLVSWYQQSPGKVPKLLIYEGVKRPSGVSNRFSGSKSG NTASLTISGLQAEDEADYYCSSYAGTRNFVFGGGTQLTVL (SEQ ID NO: 1143)

Note that the sequences provided for Group J are heavy chain and light chain amino acid sequences; the sequences provided for Group K include variable heavy chain, variable light chain, heavy chain, and light chain amino acid sequences; all other sequences presented in Table 15 are variable heavy chain and variable light chain sequences

[000300] In some embodiments, the activatable anti-PDL1 antibody includes a heavy chain that comprises or is derived from a heavy chain amino acid sequence shown in US Patent Nos. 7,943,743; 8,779,108; 8,217,149; 8,460,927; 7,794,710; 8,741,295; and 8,981,063; in US Patent Application Publication Nos. 2014356353; 20130323249; 2014341917; 20130309250; 20110280877; and 20090317368 (now issued as US Patent No. 8,981,063 listed above); and in PCT Publication Nos. WO2014/100483; WO2012/145493; WO2014/100439; WO2014/195852; WO2014/100079; WO2013/173223; and WO2014/55897, the contents of each of which are hereby incorporated by reference in their entirety. In some embodiments, the activatable anti-PDL1 antibody includes a light chain that comprises or is derived from a light chain amino acid sequence shown in US Patent Nos. 7,943,743; 8,779,108; 8,217,149; 8,460,927; 7,794,710; 8,741,295; and 8,981,063; in US Patent Application Publication Nos. 2014356353; 20130323249; 2014341917; 20130309250; 20110280877; and 20090317368 (now issued as US Patent No. 8,981,063); and in PCT Publication Nos. WO2014/100483; WO2012/145493; WO2014/100439;

WO2014/195852; WO2014/100079; WO2013/173223; and WO2014/55897, the contents of each of which are hereby incorporated by reference in their entirety. In some embodiments, the activatable anti-PDL1 antibody includes a heavy chain and a light chain that comprise or are derived from a heavy chain amino acid sequence and a light chain amino acid sequence shown in US Patent Nos. 7,943,743; 8,779,108; 8,217,149; 8,460,927; 7,794,710; 8,741,295; and 8,981,063; in US Patent Application Publication Nos. 2014356353; 20130323249; 2014341917; 20130309250; 20110280877; and 20090317368 (now issued as US Patent No. 8,981,063); and in PCT Publication Nos. WO2014/100483; WO2012/145493; WO2014/100439; WO2014/195852; WO2014/100079; WO2013/173223; and WO2014/55897, the contents of each of which are hereby incorporated by reference in their entirety.

[000301] In some embodiments, the activatable antibody includes an anti-PDL1 antibody that includes at least one CDR sequence selected from the CDR sequences (VH CDR1, VH CDR2, VH CDR3, VL CDR1, VL CDR2, VL CDR3) from the CDR sequences recited in US Patent No. 7,943,743. In some embodiments, the activatable antibody includes an anti-PDL1 antibody that includes at least one CDR, at least two, at least three, at least four, at least five, and/or six CDR sequences selected from the following sequences in US Patent No. 7,943,743: SEQ ID NO: 21, SEQ ID NO: 31, SEQ ID NO: 41, SEQ ID NO: 51, SEQ ID NO: 61, and SEQ ID NO: 71. In some embodiments, the activatable antibody includes an anti-PDL1 antibody that includes at least one CDR, at least two, at least three, at least four, at least five, and/or six CDR sequences selected from SEQ ID NO: 22, SEQ ID NO: 32, SEQ ID NO: 42, SEQ ID NO: 52, SEQ ID NO: 62, and SEQ ID NO: 72. In some embodiments, the activatable antibody includes an anti-PDL1 antibody that includes at least one CDR, at least two, at least three, at least four, at least five, and/or six CDR sequences selected from SEQ ID NO: 23, SEQ ID NO: 33, SEQ ID NO: 43, SEQ ID NO: 53, SEQ ID NO: 63, and SEQ ID NO: 73.

[000302] In some embodiments, the activatable antibody includes an anti-PDL1 antibody that includes at least one CDR, at least two, at least three, at least four, at least five, and/or six CDR sequences selected from the following sequences in US Patent No. 7,943,743: a VH CDR1 sequence comprising SEQ ID NO: 21, SEQ ID NO: 22, or SEQ ID NO: 23; a VH CDR2 sequence comprising SEQ ID NO: 31, SEQ ID NO: 32, or SEQ ID NO: 33; a VH CDR3 sequence comprising SEQ ID NO: 41, SEQ ID NO: 42, or SEQ ID NO: 43; a VL CDR1 sequence comprising SEQ ID NO: 51, SEQ ID NO: 52, or SEQ ID

NO: 53; a VL CDR2 sequence comprising SEQ ID NO: 61, SEQ ID NO: 62, or SEQ ID NO: 63; and a VL CDR3 sequence comprising SEQ ID NO: 71, SEQ ID NO: 72, or SEQ ID NO: 73.

[000303] In some embodiments, the activatable antibody includes an anti-PDL1 antibody that binds a polypeptide that comprises or is derived from SEQ ID NO: 1 or SEQ ID NO: 3 of US Patent No. 8,460,927.

[000304] In some embodiments, the activatable antibody includes an anti-PDL1 antibody that binds a human PDL1 polypeptide that comprises or is derived from SEQ ID NO: 1 of US Patent No. 8,460,927:

MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVWEMEDKNI IQFVHGEE DL
KVQHSSYRQARLLKQQLSLGNAALQITDVKLQDAGVYRCMISYGGADYKRITVKVNAPYNKINQRILVVD PVT
SEHELTCQAEGYPKAEVIWTSSDHQVLSGKTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTA
ELVIPELPLAHPNERTHLVILGAILLCLGVALTFIFRLRKGRMMDVKKCGIQDTNSKKQSDTHLEET (SEQ
ID NO: 585)

[000305] In some embodiments, the activatable antibody includes an anti-PDL1 antibody that binds a murine PDL1 polypeptide that comprises or is derived from SEQ ID NO: 3 of US Patent No. 8,460,927:

MRIFAGIIFTACCHLLRAFTITAPKDLYVVEYGSNVTMECRFPVERELDLLALVVYWEKEDEQVIQFVAGEEDL
KPQHSNFRGRASLPKDQLKGNALQITDVKLQDAGVYCCIISYGGADYKRITLKVNPYRKINQRISVDPATS
EHELICQAEGYPEAEVIWTNSDHQPVSGKRSVTTSRTEGMLLNVTSSLRVNATANDVFYCTFWRSQPGQNHTAE
LIIPELPATHPPQNRTHWVLLGSILLFLIVVSTVLLFLRKQVRMLDVEKCGVEDTSSKNRNDTQFEET (SEQ
ID NO: 586)

[000306] In some embodiments, the activatable antibody includes an anti-PDL1 antibody that binds a polypeptide that comprises or is derived from SEQ ID NO: 36 or SEQ ID NO: 38 of PCT Publication No. WO 2014/10079.

[000307] In some embodiments, the activatable antibody includes an anti-PDL1 antibody that binds a human PDL1 polypeptide that comprises or is derived from SEQ ID NO: 36 of PCT Publication No. WO 2014/10079.

MRIFAVFIFMTYWHLNNAFTVTVPKDLYVVEYGSNMTIECKFPVEKQLDLAALIVWEMEDKNI IQFVHGEE DL
KVQHSSYRQARLLKQQLSLGNAALQITDVKLQDAGVYRCMISYGGADYKRITVKVNAPYNKINQRILVVD PVT
SEHELTCQAEGYPKAEVIWTSSDHQVLSGKTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTA

ELVIPELPLAHPPNERTHLVILGAILLCLGVALTFIFRLRKGRMMDVKCGIQDTNSKQSDTHLEET (SEQ ID NO: 587)

[000308] In some embodiments, the activatable antibody includes an anti-PDL1 antibody that binds a human PDL1 extracellular domain polypeptide that comprises or is derived from SEQ ID NO: 38 of PCT Publication No. WO 2014/10079.

FTVTVPKDLVVEYGSNMTIECKFPVEKQLDLAALIVWEMEDKNI IQFVHGEEEDLKVQHSSYRQRARLLKDQL
SLGNAALQITDVKLQDAGVYRCMISYGGADYKRITVKVNAPYNKINQRILVVD PVTSEHELTCAEGYPKAEVI
WTSSDHQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPEENHTAELVIPELPLAHPPNERT
(SEQ ID NO: 588)

[000309] In some embodiments, the activatable antibody includes an anti-PDL1 antibody that is or is derived from BMS-936559 (MDX-1 105). In some embodiments, the activatable antibody includes an anti-PDL1 antibody that is or is derived from MPDL3280A (RG7446). In some embodiments, the activatable antibody includes an anti-PDL1 antibody that is or is derived from MEDI4736. In some embodiments, the activatable antibody includes an anti-PDL1 antibody that is or is derived from AMP224. In some embodiments, the activatable antibody includes an anti-PDL1 antibody that is or is derived from MSB0010718C.

[000310] In some embodiments, the activatable anti-PDL1 antibody comprises or is derived from an antibody that is manufactured, secreted or otherwise produced by a hybridoma, such as, for example, the hybridoma(s) disclosed in US Patent No. 8,779,108 and deposited at the National Collections of Industrial and Marine Bacteria (NCIMB) under deposit number 41598, as well as the hybridoma(s) disclosed in US Patent No. 8,741,295 and deposited at the Collection Nationale De Cultures De Microorganismes (CNCM) under deposit number I-4080, I-4081, and/or I-4122.

[000311] In some embodiments, the activatable anti-PDL1 antibody includes a CDR sequence shown in Table 16, a combination of VL CDR sequences (VL CDR1, VL CDR2, VL CDR3) selected from the group consisting of those combinations shown in a single row Table 16, a combination of VH CDR sequences (VH CDR1, VH CDR2, VH CDR3) selected from the group consisting of those combinations shown in Table 16, or a combination of VL CDR and VH CDR sequences (VL CDR1, VL CDR2, VL CDR3, VH CDR1, VH CDR2, VH CDR3) selected the group consisting of those combinations shown in Table 16.

Table 16. CDR Sequences for Antibodies and Activatable Antibodies that Bind PDL1

AB Name	VL			VH		
	CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)	CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)
C8	RASQSISSYLN (209)	KASRLQS (210)	RALKPVT (211)	SYAMS (212)	DITASGQRTTYADS (213)	SKAIFDY (214)
C12	RASQSISSYLN (209)	AASSLQS (215)	SYSTPNT (216)	SYAMS (212)	SINKDGHYTSYADS (217)	NLDEFDY (218)
C16	RASQSISSYLN (209)	SASQLQS (219)	ANSRPST (220)	SYAMS (212)	SIMATGAGTLYADS (221)	DGAGFDY (222)
C20	RASQSISSYLN (209)	NASSLQS (223)	YPYGP (224)	SYAMS (212)	TITSSGAATYYADS (225)	NYTGFDY (226)
C60	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SIYSTGGATAYADS (229)	SSAGFDY (230)
C1	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIYSTGGATAYADS (231)	SSAGQSRP GFDY (232)
D1	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIYSTGGATAYADS (231)	SSAGQSWP GFDY (233)
G7	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIYSTGGATAYADS (231)	SSAGQSFP GFDY (234)
H9	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIYSTGGATAYADS (231)	WSAAFYD (235)
B10	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIYSTGGATAYADS (231)	WSAGYDY (236)
E10	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIYSTGGATAYADS (231)	WSKGFDY (237)
A05	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIWKQGIVTVYDS (238)	SSAGFDY (230)
C05	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIWRNGIVTVYDS (239)	SSAGFDY (230)
C10	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SDIWKQGMVTVYDS (240)	SSAGFDY (230)
D08	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIWRQGLATAYDS (241)	SSAGFDY (230)

G09	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SEIVATGILTSYDS (242)	SSAGFDY (230)
G10	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIGRQGLITVYDS (243)	SSAGFDY (230)
G12	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIWYQGLVTYDS (244)	SSAGFDY (230)
E11	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SDIWKQGFATADS (245)	SSAGFDY (230)
D01	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIWKQGIVTVYDS (238)	SSAGFDY (230)
H06	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIWRQGLATAYDS (241)	SSAGFDY (230)
C5H9	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIWRNGIVTVYADS (246)	WSAAFDY (235)
C5B10	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIWRNGIVTVYADS (246)	WSAGYDY (236)
C5E10	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIWRNGIVTVYADS (246)	WSKGFDY (237)
G12H9	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIWYQGLVTYADS (247)	WSAAFDY (235)
G12B10	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIWYQGLVTYADS (247)	WSAGYDY (236)
G12E10	RASQSISSYLN (209)	YASTLQS (227)	DNGYPST (228)	SYAMS (212)	SSIWYQGLVTYADS (247)	WSKGFDY (237)
C5H9 v2	RASQSISSYLN (209)	AASSLQS (215)	DNGYPST (228)	SYAMS (212)	SSIWRNGIVTVYADS (246)	WSAAFDY (235)

[000312] In some embodiments, the activatable anti-PDL1 antibody includes a CDR sequence shown in Table 17, a combination of VL CDR sequences selected from the group consisting of those combinations shown in Table 17, and/or a combination of VH CDR sequences selected from the group consisting of those combinations shown in Table 17.

[000313] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group A in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group A in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences

selected from the group consisting of the combinations shown in Group A in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group A in Table 17.

[000314] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group B in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group B in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group B in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group B in Table 17.

[000315] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group C in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group C in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group C in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group C in Table 17.

[000316] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group D in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group D in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group D in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group D in Table 17.

[000317] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group E in Table 17. In some embodiments, the activatable anti-

PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group E in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group E in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group E in Table 17.

[000318] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group F in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group F in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group F in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group F in Table 17.

[000319] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group G in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group G in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group G in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group G in Table 17.

[000320] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group H in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group H in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group H in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group H in Table 17.

[000321] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group I in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group I in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group I in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group I in Table 17.

[000322] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group J in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group J in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group J in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group J in Table 17.

[000323] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group K in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group K in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group K in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group K in Table 17.

[000324] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group L in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group L in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences

selected from the group consisting of the combinations shown in Group L in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group L in Table 17.

[000325] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group M in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group M in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group M in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group M in Table 17.

[000326] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group N in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group N in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group N in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group N in Table 17.

[000327] In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group O in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group O in Table 17. In some embodiments, the activatable anti-PDL1 antibody includes a combination of heavy chain CDR sequences selected from the group consisting of the combinations shown in Group O in Table 17, and a combination of light chain CDR sequences selected from the group consisting of the combinations shown in Group O in Table 17.

[000328] In some embodiments, the activatable anti-PDL1 antibody includes a heavy chain CDR1 sequence selected from the group consisting of the sequences shown in Group P in Table 17.

Table 17. Additional CDR Sequences for Antibodies and Activatable Antibodies that Bind PDL1

Group A					
VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
RASQSVSSYLV (304)	DASNRAT (293)	QQRSNWPRT (278)	DYGFS (296)	WITAYNGNTNYAQKLQG (284)	DYFYGMDV (269)
RASQSVSSYLA (305)	AASSLQS (215)	QQRSNWPRT (279)	TYAIS (297)	GIIPIFGKAHYAQKFQG (285)	KFHFVSGSPFG MDV (270)
RASQGISSWLA (306)	GASSRAT (294)	QQYNSYPYT (280)	SYDVH (298)	WLHADTGITKFSQKFQG (286)	ERIQLWFDY (271)
RASQSVSSSYLA (307)	DASSLES (295)	QQYGSSPWT (281)	TYAIN (299)	GIIPIFGTANHAQKFQG (287)	DQGIAAALFDY (272)
RASQGISSALA (308)	KASTLES (447)	QQYGSSP (282)	DYVVH (300)	GISGNSGNIGYADSVKG (288)	PFDY (273)
		QQFNSYPFT (283)	SYAIS (301)	GIIPIFGRAHYAQKFQG (289)	KYDYVSGSPFG MDV (274)
		QQSYSTPWT (448)	SYAIN (302)	GIIPIFGSANYAQKFQD (290)	DSSGWSRYMD V (275)
			DYGMH (303)	GIIPLFGIAHYAQKFQG (291)	KYSYVSGSPFG MDV (276)
				GISWNRGRIEYADSVKG (292)	GRFRYFDWFLD Y (277)
					DYFWSGFSAFD I (446)

Group B	
VL	VH

CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
SGDALPQKYVF (312)	EDSKRPS (313)	YSTDRSGNHRV (314)	TYSMN (309)	SISSSGDYIYYADSVK (310)	DLVTSMVAFDY (311)
RASQRVSSSYLA (318)	DASSRAT (319)	QQYGSLPWT (320)	RYWMS (315)	NIKQDGSEKYYVDSVKG (316)	EGGWFGELAFD Y (317)
RASQSVSSNYLA (324)	GTSSRAT (325)	QQYGSSIFT (326)	SYWMS (321)	NIKQDGGEQYYVDSVK (322)	DWNYGYYDMDV (323)

Group C					
VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
RASQ (D/V) (V/I) (S/N) T (A/F) (V/L) A (327)	SAS (F/T) L (Y/A) S (328)	QQ (Y/G/F/S) (L/Y/F/W) (Y /N/A/T/G/F/ I) (H/V/P/T/ I) P (A/W/R/P /T) T (329)	GFTFS (D/G) SWIH (330)	AWI (S/L) PYGGS (T/S) YYADSVKG (331)	RHWPGGF DY (332)
RASQDVSTAVA (333)	SASFLYS (334)	QQYLYHPAT (335)	GFTFSDSWIH (336)	AWISPYGGSTYYADSVKG (337)	

Group D					
VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
KSSQSLL (H/N) (S/T) (R/S) TRKNYL A (584)	WASTRES (589)	(Q/K) QSYDVV T (590)	SYW (I/M) H (591)	YINPSS (D/G) Y (H/N) E Y (S/N) (E/Q) KF (I/M) D (592)	SGWL (I/ V) HGDYY FD (F/Y) (593)

KSSQSLLSRT RKNYLA (594)		QQSYDVVT (595)	SYWMH (596)	YINPSSDYNEYSEKFMD (597)	SGWLVHG DYYFDY (598)
KSSQSLLTST RKNYLA (599)		KQSYDVVT (600)	GYFTSYWMH (601)	YINPSSGYHEYNQKFID (602)	SGWLIHG DYYFDF (603)
			SYWIIH (604)		
			GTTFTSYWIIH (605)		

Group E					
VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
TGT(N/S) (T/ R/S) DVG(A/G) YNYVS (606)	(E/D) V(I /N/D) (D/ H/N) RPS (607)	SS(F/Y) T(N/ S) (R/T/S) (G /S) (I/T) RV (608)	(K/R/T/Q/G /A/W/M/I/S) Y(V/R/K/L /M/I) M(H/T /N/Q/A/V/Y /W/F/M) (609)	SIYPSGG(F/I) TFYAD(S/T) VKG (610)	IKLGTVT TV(E/D) Y (611)
TGTSSDVGGYN YVS (612)	DVSNRPS (613)	SSYTSSSTRV (614)	SYIMM (615)	SIYPSGGITFYADTVKG (617)	IKLGTVT TVDY (618)

Group F					
VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
			MYMMM (619)	SIYPSGGITFYADSVKG (620)	TGTSSDV GAYNYVS (621)

Group G					
VL			VH		

CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
(R/S/K) (A/S)) S (R/N/D/Q/ E/H/K/M/S) (S LLYSS or Absent) (N/D or Absent) (A/Q /E/L/K/M/P/ S/T/Y/V) X (R /N/S/T) (A/R /N/C/Q/E/H/ I/L/K/M/F/S /T/Y/V) (I/L /M/V) (A/R/N /Q/E/H/I/K/ M/F/S/T/Y) (622)	X (A/C/G/ S/T/V) (A /C/H/I/L /M/F/S/T /Y/V) (R/ N/D/Q/E/ K/S/T) (A /R/N/D/Q /E/G/H/K /M/F/P/S /T/Y) (A/ N/S/T) (623)	(R/N/Q/E/H) QX (A/R/N/C/ Q/E/H/I/L/K /M/F/S/T/Y/ V) (N/G/S) (A /R/N/D/C/Q/ E/G/H/I/L/K /M/F/R/S/T/ W/Y/V) P (A/R /N/D/Q/E/G/ H/K/M/P/S/T /Y) T (624)	G (Y/F) (T/S) (I/L/M/F) (A/R/N/D/C /Q/E/H/I/L /K/M/P/S/T /W/Y/V) (N/ D/Q/E/K/P/ S/T) (A/R/N /D/C/Q/E/G /H/I/L/K/M /F/P/S/T/Y /V) (A/R/N/ D/C/Q/E/G/ H/I/L/K/M/ F/P/S/T/Y/ V) (A/R/C/Q /E/H/I/L/K /M/F/S/T/W /Y/V) N (625)	X (I/L/M/F/V) X (P or Absent) (A/R/N/Q/E/ H/K/M/S/T/Y/V) (A/R /N/D/C/Q/E/G/H/K/M /F/P/S/T/Y/V) (N/G/ S) (N/G/S) T (A/R/N/Q /E/H/K/M/S/T/Y) (A/ R/N/C/Q/E/H/I/L/K/ M/F/S/T/W/Y/V) N (A/ D/Q/E/K/P/S/T) (A/R /N/D/C/Q/E/G/H/I/L /K/M/F/P/S/T/Y/V) (I/L/M/F) K (A/G/S (626)	(A/R/N/ Q/E/H/K /M/S/T/ Y) (A/R/ N/D/C/Q /E/G/H/ K/M/F/P /S/T/Y/ V) (Q/G/ H/T/W) X XXX (F/Y) (AI or Absent) (627)
RASSSVSYIY (628)	ATFNLAS (629)	HQRSSYPWT (630)	GYTFPDYMN (631)	DIDPNYGGTTYNQKFKG (632)	GAL (633)
SASSSIRYMH (634)	DTSKLTS (635)	QQDSSYPLT (636)	GYTFTSYDIN (637)	WIFPRDNNTKYNENFKG (638)	ENWVGDF (639)
KASQDVGTA (640)	WASTRHT (641)	QQYYGYPLT (642)	GYSITSDYWN (643)	YISYTGSTYYNPSLKS (644)	YGGWLSP F (645)
KSSQSLLYSSN QKNSL (646)	WASTRES (589)		GYSIISDYWN (647)		RGGWLLP F (648)
			GFSLTTY SIN (649)	VMWAGGGTNSNSVLKS (650)	YYGNSPY YAI (651)

Group H	
VL	VH

CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
TRSSGSIGSNY VQ (652)	EDNQRP (653)	QSYDSSTWV (654)	SYAIS (301)	WISPIGGSTNYAQK VQG (655)	GLXXXXXXXXXX XXXXXXDV (656)
TRSSGNIASNY VQ (657)	GKNNRPS (658)	QSYDSSNLWV (659)	SYGIS (660)	WISAYNGNTNYAQK LED (661)	ALPSGTILVGG WFDP (662)
QGDSLRSYYAS (663)	SDRDRPS (664)	NSRDSSGNHYV (665)	SYALS (666)	AISGGGGSTYYADS VKD (667)	DVFPETFSMNY GMDV (668)
GGSDIGRKS VH (669)	DVSNRPS (613)	QVWDNNSDHYV (670)	DYAMH (671)	LISGDGGSTYYADS VKD (672)	VLLPCSSTSCY GSVGA FDI (673)
TGTSSDVGGYN YVS (612)	GASSRAT (294)	SSYTSSTLP (674)	NYDMS (675)	RVNWNGGSTTYADA VKD (676)	EFVGAYDL (677)
RASQSIGNSLA (678)	AASTLQS (679)	QQHTIPTFS (680)	GLYIH (681)	WIIPIFGTANYAQK FED (682)	GLRWGIWGWFD P (683)
RASQGIGSYLA (684)	QDIKRPS (685)	QQLNNYPIT (686)	DNAIS (687)	WIIPIFGKPNYAQK FED (688)	TMVRGFLGVMD V (689)
SGDKLGNKYAY (690)	EDYRRPS (691)	QTDNSV V (692)	SYAMS (212)	AISGSGGSTYYADS VKD (693)	DQFVTIFGVPR YGM DV (694)
TRSSGSIDSNY VQ (695)	DDSDRPS (696)	QSYDSNNRHVI (697)	TYALN (698)	RIVPLIGLVNYAHN FED (699)	GRQMFGAGIDF (700)
TRSSGNIGTNY VQ (701)	EDNKRPS (702)	QSYHSSGWE (703)	SHGIT (704)	WISAHNGHASNAQK VED (705)	EVYGGNSDY (706)
GGNNIGSKGVH (707)	YKSDSNKQ QAS (708)	QVWDSSSDHWV (709)	RHGMH (710)	VISHDGSVKYYADS MKD (711)	VHAALYGM DV (712)
TRSSGSIASNY VQ (713)		QSYDSTTPSV (714)		WTSPHNGLTAF AQI (715)	GLSYQVSGWFD P (716)
TRSSGSIASHY VQ (717)		QSYDGITVI (718)		RIIPILGIAN YAQK (719)	VHPVFSYALDV (720)
TLRSGLNVG SY (721)		QSYDSSNRWV (722)	TYAFS (723)	WISAYNGNTNYAQK VED (724)	DGYGSDPVL (725)
		MIWYSSAVV (726)	NYGIS (727)		GDFRKPFDY (728)

Group I

VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
RATESVEYYGT SLVQ (729)	AASSVDS (730)	QQSRRVPYT (731)	SYVMH (732)	YVNPFDGTTYNEMFKG (733)	QAWGYP (734)

Group J					
VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
RASQDVSTAVA (333)	SASFLYS (334)	QQYLYHPAT (335)	GFTFSDSWIH (336)	AWISPYGGSTYYADSVKG (337)	RHWPGGF DY (332)

Group K					
VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
QSIISNW (781)	KAS (782)	QQYHSYSYT (783)	GFTFSRFW (784)	INQDGTEK (785)	ANTYYDFWSGH FDY (786)
QGIRND (787)	TAS (788)	LQHNSYPLT (789)	GFTFSNFG (790)	LWSDGSNK (791)	ARGRGAPGIPI FGY (792)
QGIRND (787)	AAS (793)	LQHNNYPYT (794)	GFTFSNAW (795)	IKRKTDGGTT (796)	TTDDIVVVPVAV MREYYFGMDV (797)
QTLVHGDGNTY (798)	KVS (799)	MQATHFPIT (800)	GYSFTGY (801)	INPNSGTK (802)	ARDEDWNFGSW FDS (803)
PSLVHSDGNTY (804)	KIS (805)	MQATHFPIT (800)	GYTFTGY (806)	LNPNTGTT (807)	ARDEDWNYGSW FDT (808)
QSINSY (809)	VAS (810)	QQSYSTPPIT (811)	GFTFDDYG (812)	IHWHGKRT (813)	VRGGMSTGDWF DP (814)
QSISSY (815)	VAS (810)	QQSYSTPPIT (811)	GFTFDDYG (812)	IHWSGRST (816)	ARGGMSTGDWF DP (817)
QTINIY (818)	AAS (793)	HQSYSTPPIT (819)	GFTVGSNY (820)	IYSGGST (821)	ARGIRGLDV (822)
QSMSSY (1011)	AAS (793)	QQSYSTPPIT (811)	GITVGTNY (1012)	ISSGGNT (1013)	ARGIRGLDV (822)
QSFNFNY (823)	GAS (824)	QQYESAPWT (825)	GGIFSSST (826)	IIPVFGTV (827)	ARNWGLGSFYI (828)
QSISSY (815)	AAS (793)	QQSYCTPPIT (829)	GFPPDEYA (830)	ISWSNNNI (831)	AKSGIFDS (832)
QSISSY (815)	AAS (793)	QQSYSTPPIT (811)	GFTFSSYG (833)	ISYEGRNK (834)	AKDRTLYGMDV (835)
QVISNY (836)	AAS (793)	QKYNAPRT (837)	GFSLSTNRC (838)	IDWDGVK (839)	ARSTSLTFYF DY (840)
QNINNY (841)	AAS (793)	QQSYNTPLT (842)	EFTVGTNH (843)	IYSGGNT (844)	ARGLGMDV (845)
QTISTY (846)	AAS (793)	LQHNSYPYT (847)	GFTFSKYW (848)	IKGDGSEK (849)	ARDYWGSGYYF DF (850)

QSISSY (815)	VVS (851)	QSYSTPFT (852)	GFTFSSYW (853)	IKQDGSEK (854)	ARDDIVVVPAP MGYYYYYFGMD V (855)
	AAS (793)	QSYSTPPIT (811)	GFTFDDFA (856)	ISWTGGNM (857)	VKDIRGIVATG GAFDI (858)
			GFTVGTNY (859)	IYSGGST (821)	ARGIRGFDI (860)
			GFTISTNY (861)	IYSSGST (862)	ARGIRGFDI (860)
			GFTIDDSA (863)	ISWKSISI (864)	VKDIRGNWNYG GNWFDP (865)
			GFTVGVNH (866)	IFSSGRT (867)	ARGIGGLDI (868)
			GFTFDDYA (869)	ISWTGGTI (870)	TRDIRGNWKYG GWFPD (871)
			GYTFTAYY (872)	ISPNSGFT (873)	AREGSTHNSF DP (874)
			GFTVGTNF (875)	IIPILGAA (876)	ARGGGMDV (877)
			GGTFNTYV (878)	ISPYNGYT (879)	ARDRTSGGFDP (880)
			GYIFTHYG (881)		SRGRGPYWSFD L (882)

Group L					
VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
QASESVYSNNY LS (1014)	LASTLAS (1015)	IGGKSSSTDG NA (1016)	SNGLT (1017)	TINKDASAYYASWAK G (1018)	IAFKTGTISI (1019)

Group M					
VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)

RSSKSLHLSNG ITYLY (1020)	QMSNLAS (1021)	AQNLEPPLT (1022)	DYYTH (1023)	WIDPENGKTAYAPKF QG (1024)	GGYDVYFLDY (1025)
--------------------------------	-------------------	---------------------	-----------------	------------------------------	----------------------

Group N					
VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
KASQDVGIVVA (1026)	WASIRHT (1027)	QQYSNYPLYT (1028)	GFSLTSYGVH (1029)	VIWAGGSTNYNSALM S (1030)	AKPYGNSAMDY (1031)
				VIWAGGSTNYVDSVK G (1032)	AKPYGTSAMDY (1033)
				VIWAGGSTNYADSVK G (1034)	

Group O					
VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
ASQSVSTSSSS FMH (1035)	YASNLES (1036)	QHSWEIPYT (1037)	SYGMS (1038)	SISSGGSTYYPDSVK G (1039)	GYDSGFAY (1040)
RASQSVSTSSS SYMH (1041)	YASNLES (1036)	QHSWEIPYT (1037)	SYGMS (1038)	SISSGGTTYYPDSVK G (1042)	GYDSGFAY (1040)
KASQSVSNDVA (1043)	YAANRYT (1044)	QQDYTSPYT (1045)	TYGVH (1046)	VIWRGVTTDYNAAFM AS (1047)	LGFYAMDY (1048)
KASQSVSNDVG (1049)	YASNRYT (1050)	QQDYTSPYT (1045)	SYGVH (1051)	VIWGGVTDYNAAFI S (1052)	LGFYAMDY (1048)
RSSQIIVHSNA NTYLE (1053)	KVSNRFS (1054)	FQGSHPYT (1055)	TYWMH (1056)	QINPDSTTINTAPSL KD (1057)	PGDYGDFDC (1058)
SASSSVSSSYL Y (1059)	NTSNLAS (1060)	HQWRSYPPT (1061)	SGYWN (1062)	YISYSGSTYYNPSLK S (1063)	SLLWFSTGFAY (1064)
SANSSVSIMH (1065)	DTSKLAS (1066)	QQWSSNPWT (1067)	SYGVH (1051)	YIWSGGITDYNAAFK S (1068)	LGFYAMDY (1048)
RASQSVSTSSY SYMH (1069)	YASNLES (1036)	QNSWEIPYT (1070)	STGMS (1071)	SISSGGTTYLGSVQ G (1072)	GYDAGFAY (1073)

KSSQSLLYSSN QKNSLA (1074)	WASNRES (1075)	QQYYSYPLT (1076)	SGYWT (1077)	YIYTGSTYYNPSLKS (1078)	QRDWLGFAY (1079)
RASQSVSTSSY SYVH (1080)	YASNLES (1036)	QHSWEIPYT (1037)	SYGMS (1038)	SISSGGSIIYPDSVK G (1081)	GYDAGFAF (1082)

Group P					
VL			VH		
CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)	CDR1 (SEQ ID NO:)	CDR2 (SEQ ID NO:)	CDR3 (SEQ ID NO:)
			GFTFSMYMMM (1083)		
			GFTFSAYAMA (1084)		
			GFTFSAYRMF (1085)		
			GFTFSAYLMV (1086)		
			GFTFSAYVMF (1087)		
			GFTFSAYVMS (1088)		
			GFTFSGYLMV (1089)		
			GFTFSGYQML (1090)		
			GFTFSGYSMF (1091)		
			GFTFSGYWMA (1092)		
			GFTFSQYLMY (1093)		
			GFTFSQYVMF (1094)		
			GFTFSQYYMY (1095)		

			GFTFSSYLMS (1096)		
			GFTFSSYLMT (1097)		
			GFTFSSYQMV (1098)		
			GFTFSSYSMA (1099)		
			GFTFSSYVMF (1100)		
			GFTFSSYVMS (1101)		
			GFTFSSYVMY (1102)		
			GFTFSSYYMF (1103)		
			GFTFSSYYMV (1104)		
			GFTFSYYSMV (1105)		
			GFTFSWYLMA (1106)		
			GFTFSWYQMS (1107)		

[000329] The ABs in the activatable antibodies of the disclosure specifically bind a PDL1 target, such as, for example, mammalian PDL1, and/or human PDL1. Also included in the disclosure are ABs that bind to the same PDL1 epitope as an antibody of the disclosure and/or an activated activatable antibody described herein. Also included in the disclosure are ABs that compete with an anti-PDL1 antibody and/or an activated anti-PDL1 activatable antibody described herein for binding to a PDL1 target, e.g., human PDL1. Also included in the disclosure are ABs that cross-compete with an anti-PDL1 antibody and/or an activated anti-PDL1 activatable antibody described herein for binding to a PDL1 target, e.g., human PDL1.

[000330] The activatable anti-PDL1 antibodies provided herein include a masking moiety. In some embodiments, the masking moiety is an amino acid sequence that is coupled or otherwise attached to the anti-PDL1 antibody and is positioned within the

activatable anti-PDL1 antibody construct such that the masking moiety reduces the ability of the anti-PDL1 antibody to specifically bind PDL1. Suitable masking moieties are identified using any of a variety of known techniques. For example, peptide masking moieties are identified using the methods described in PCT Publication No. WO 2009/025846 by Daugherty et al., the contents of which are hereby incorporated by reference in their entirety.

[000331] The activatable anti-PDL1 antibodies provided herein include a cleavable moiety. In some embodiments, the cleavable moiety includes an amino acid sequence that is a substrate for a protease, usually an extracellular protease. Suitable substrates are identified using any of a variety of known techniques. For example, peptide substrates are identified using the methods described in U.S. Patent No. 7,666,817 by Daugherty et al.; in U.S. Patent No. 8,563,269 by Stagliano et al.; and in PCT Publication No. WO 2014/026136 by La Porte et al., the contents of each of which are hereby incorporated by reference in their entirety. (See also Boulware et al. “Evolutionary optimization of peptide substrates for proteases that exhibit rapid hydrolysis kinetics.” Biotechnol Bioeng. 106.3 (2010): 339-46).

[000332] Exemplary substrates include but are not limited to substrates cleavable by one or more of the following enzymes or proteases listed in Table 12.

Table 12: Exemplary Proteases and/or Enzymes

ADAMS, ADAMTS, <i>e.g.</i> , ADAM8 ADAM9 ADAM10 ADAM12 ADAM15 ADAM17/TACE ADAMDEC1 ADAMTS1 ADAMTS4 ADAMTS5	Cysteine proteinases, <i>e.g.</i> , Cruzipain Legumain Otubain-2	Serine proteases, <i>e.g.</i> , activated protein C Cathepsin A Cathepsin G Chymase coagulation factor proteases (<i>e.g.</i> , FVIIa, FIXa, FXa, FXIa, FXIIa)
		Elastase
		Granzyme B
		Guanidinobenzoate
		HtrA1
		Human Neutrophil Elastase
		Lactoferrin
		Marapsin
		NS3/4A
		PACE4
Aspartate proteases, <i>e.g.</i> , BACE Renin	KLKs, <i>e.g.</i> , KLK4	Plasmin
	KLK5 KLK6 KLK7 KLK8 KLK10 KLK11 KLK13 KLK14	PSA
Aspartic cathepsins, <i>e.g.</i> , Cathepsin D Cathepsin E	Metallo proteinases, <i>e.g.</i> , Meprin Neprilysin	

	PSMA	tPA
Caspases, <i>e.g.</i> ,	BMP-1	Thrombin
Caspase 1		Tryptase
Caspase 2	MMPs, <i>e.g.</i> ,	uPA
Caspase 3	MMP1	
Caspase 4	MMP2	Type II Transmembrane
Caspase 5	MMP3	Serine Proteases (TTSPs),
		<i>e.g.</i> ,
Caspase 6	MMP7	DESC1
Caspase 7	MMP8	DPP-4
Caspase 8	MMP9	FAP
Caspase 9	MMP10	Hepsin
Caspase 10	MMP11	Matriptase-2
Caspase 14	MMP12	MT-SP1/Matriptase
	MMP13	TMPRSS2
Cysteine cathepsins, <i>e.g.</i> ,	MMP14	TMPRSS3
Cathepsin B	MMP15	TMPRSS4
Cathepsin C	MMP16	
Cathepsin K	MMP17	
Cathepsin L	MMP19	
Cathepsin S	MMP20	
Cathepsin V/L2	MMP23	
Cathepsin X/Z/P	MMP24	
	MMP26	
	MMP27	

[000333] The activatable anti-PDL1 antibodies described herein overcome a limitation of antibody therapeutics, particularly antibody therapeutics that are known to be toxic to at least some degree *in vivo*. Target-mediated toxicity constitutes a major limitation for the development of therapeutic antibodies. The activatable anti-PDL1 antibodies provided herein are designed to address the toxicity associated with the inhibition of the target in normal tissues by traditional therapeutic antibodies. These activatable anti-PDL1 antibodies remain masked until proteolytically activated at the site of disease. Starting with an anti-PDL1 antibody as a parental therapeutic antibody, the activatable anti-PDL1 antibodies of the invention were engineered by coupling the antibody to an inhibitory mask through a linker that incorporates a protease substrate.

[000334] When the AB is modified with a MM and is in the presence of the target, specific binding of the AB to its target is reduced or inhibited, as compared to the specific binding of the AB not modified with an MM or the specific binding of the parental AB to the target.

[000335] The K_d of the AB modified with a MM towards the target is at least 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 50,000, 100,000, 500,000, 1,000,000, 5,000,000, 10,000,000, 50,000,000 or greater, or between 5-10, 10-100, 10-1,000, 10-10,000, 10-100,000, 10-1,000,000, 10-10,000,000, 100-1,000, 100-10,000, 100-100,000, 100-1,000,000, 100-10,000,000, 1,000-10,000, 1,000-100,000, 1,000-1,000,000, 1000-10,000,000, 10,000-100,000, 10,000-1,000,000, 10,000-10,000,000, 100,000-1,000,000, or 100,000-10,000,000 times greater than the K_d of the AB not modified with an MM or of the parental AB towards the target. Conversely, the binding affinity of the AB modified with a MM towards the target is at least 2, 3, 4, 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 50,000, 100,000, 500,000, 1,000,000, 5,000,000, 10,000,000, 50,000,000 or greater, or between 5-10, 10-100, 10-1,000, 10-10,000, 10-100,000, 10-1,000,000, 10-10,000,000, 100-1,000, 100-10,000, 100-100,000, 100-1,000,000, 100-10,000,000, 1,000-10,000, 1,000-100,000, 1,000-1,000,000, 1000-10,000,000, 10,000-100,000, 10,000-1,000,000, 10,000-10,000,000, 100,000-1,000,000, or 100,000-10,000,000 times lower than the binding affinity of the AB not modified with an MM or of the parental AB towards the target.

[000336] In some embodiments, the dissociation constant (K_d) of the MM towards the AB is generally greater than the K_d of the AB towards the target. The K_d of the MM towards the AB can be at least 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 100,000, 1,000,000 or even 10,000,000 times greater than the K_d of the AB towards the target. Conversely, the binding affinity of the MM towards the AB is generally lower than the binding affinity of the AB towards the target. The binding affinity of MM towards the AB can be at least 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 100,000, 1,000,000 or even 10,000,000 times lower than the binding affinity of the AB towards the target.

[000337] In some embodiments, the dissociation constant (K_d) of the MM towards the AB is approximately equal to the K_d of the AB towards the target. In some embodiments, the dissociation constant (K_d) of the MM towards the AB is no more than the dissociation constant of the AB towards the target.

[000338] In some embodiments, the dissociation constant (K_d) of the MM towards the AB is less than the dissociation constant of the AB towards the target.

[000339] In some embodiments, the dissociation constant (K_d) of the MM towards the AB is greater than the dissociation constant of the AB towards the target.

[000340] In some embodiments, the MM has a K_d for binding to the AB that is no more than the K_d for binding of the AB to the target.

[000341] In some embodiments, the MM has a K_d for binding to the AB that is no less than the K_d for binding of the AB to the target.

[000342] In some embodiments, the MM has a K_d for binding to the AB that is approximately equal to the K_d for binding of the AB to the target.

[000343] In some embodiments, the MM has a K_d for binding to the AB that is less than the K_d for binding of the AB to the target.

[000344] In some embodiments, the MM has a K_d for binding to the AB that is greater than the K_d for binding of the AB to the target.

[000345] In some embodiments, the MM has a K_d for binding to the AB that is no more than 2, 3, 4, 5, 10, 25, 50, 100, 250, 500, or 1,000 fold greater than the K_d for binding of the AB to the target. In some embodiments, the MM has a K_d for binding to the AB that is between 1-5, 2-5, 2-10, 5-10, 5-20, 5-50, 5-100, 10-100, 10-1,000, 20-100, 20-1000, or 100-1,000 fold greater than the K_d for binding of the AB to the target.

[000346] In some embodiments, the MM has an affinity for binding to the AB that is less than the affinity of binding of the AB to the target.

[000347] In some embodiments, the MM has an affinity for binding to the AB that is no more than the affinity of binding of the AB to the target.

[000348] In some embodiments, the MM has an affinity for binding to the AB that is approximately equal of the affinity of binding of the AB to the target.

[000349] In some embodiments, the MM has an affinity for binding to the AB that is no less than the affinity of binding of the AB to the target.

[000350] In some embodiments, the MM has an affinity for binding to the AB that is greater than the affinity of binding of the AB to the target.

[000351] In some embodiments, the MM has an affinity for binding to the AB that is 2, 3, 4, 5, 10, 25, 50, 100, 250, 500, or 1,000 less than the affinity of binding of the AB to the target. I In some embodiments, the MM has an affinity for binding to the AB that is between 1-5, 2-5, 2-10, 5-10, 5-20, 5-50, 5-100, 10-100, 10-1,000, 20-100, 20-1000, or 100-1,000 fold less than the affinity of binding of the AB to the target. In some embodiments, the MM has an affinity for binding to the AB that is 2 to 20 fold less than the affinity of binding of the AB to the target. In some embodiments, a MM not covalently linked to the AB and at equimolar concentration to the AB does not inhibit the binding of the AB to the target.

[000352] When the AB is modified with a MM and is in the presence of the target specific binding of the AB to its target is reduced or inhibited, as compared to the specific binding of the AB not modified with an MM or the specific binding of the parental AB to the target. When compared to the binding of the AB not modified with an MM or the binding of the parental AB to the target the AB's ability to bind the target when modified with an MM can be reduced by at least 50%, 60%, 70%, 80%, 90%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% and even 100% for at least 2, 4, 6, 8, 12, 28, 24, 30, 36, 48, 60, 72, 84, or 96 hours, or 5, 10, 15, 30, 45, 60, 90, 120, 150, or 180 days, or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months or more when measured *in vivo* or in an *in vitro* assay.

[000353] The MM inhibits the binding of the AB to the target. The MM binds the antigen binding domain of the AB and inhibits binding of the AB to the target. The MM can sterically inhibit the binding of the AB to the target. The MM can allosterically inhibit the binding of the AB to its target. In these embodiments when the AB is modified or coupled to a MM and in the presence of target there is no binding or substantially no binding of the AB to the target, or no more than 0.001%, 0.01%, 0.1%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, or 50% binding of the AB to the target, as compared to the binding of the AB not modified with an MM, the parental AB, or the AB not coupled to an MM to the target, for at least 2, 4, 6, 8, 12, 28, 24, 30, 36, 48, 60, 72, 84, or 96 hours, or 5, 10, 15, 30, 45, 60, 90, 120, 150, or 180 days, or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months or longer when measured *in vivo* or in an *in vitro* assay.

[000354] When an AB is coupled to or modified by a MM, the MM 'masks' or reduces or otherwise inhibits the specific binding of the AB to the target. When an AB is coupled to or modified by a MM, such coupling or modification can effect a structural change that reduces or inhibits the ability of the AB to specifically bind its target.

[000355] An AB coupled to or modified with an MM can be represented by the following formulae (in order from an amino (N) terminal region to carboxyl (C) terminal region:

(MM)-(AB)

(AB)-(MM)

(MM)-L-(AB)

(AB)-L-(MM)

where MM is a masking moiety, the AB is an antibody or antibody fragment thereof, and the L is a linker. In many embodiments, it may be desirable to insert one or more linkers, *e.g.*, flexible linkers, into the composition so as to provide for flexibility.

[000356] In certain embodiments, the MM is not a natural binding partner of the AB. In some embodiments, the MM contains no or substantially no homology to any natural binding partner of the AB. In some embodiments, the MM is no more than 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% similar to any natural binding partner of the AB. In some embodiments, the MM is no more than 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 25% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 50% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 20% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 10% identical to any natural binding partner of the AB.

[000357] In some embodiments, the activatable antibodies include an AB that is modified by an MM and also includes one or more cleavable moieties (CM). Such activatable antibodies exhibit activatable/switchable binding, to the AB's target. Activatable antibodies generally include an antibody or antibody fragment (AB), modified by or coupled to a masking moiety (MM) and a modifiable or cleavable moiety (CM). In some embodiments, the CM contains an amino acid sequence that serves as a substrate for at least one protease.

[000358] The elements of the activatable antibodies are arranged so that the MM and CM are positioned such that in a cleaved (or relatively active) state and in the presence of a target, the AB binds a target while the activatable antibody is in an uncleaved (or relatively inactive) state in the presence of the target, specific binding of the AB to its target is reduced or inhibited. The specific binding of the AB to its target can be reduced due to the inhibition or masking of the AB's ability to specifically bind its target by the MM.

[000359] The K_d of the AB modified with a MM and a CM towards the target is at least 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 50,000, 100,000, 500,000, 1,000,000, 5,000,000, 10,000,000, 50,000,000 or greater, or between 5-10, 10-100, 10-1,000, 10-10,000, 10-100,000, 10-1,000,000, 10-10,000,000, 100-1,000, 100-10,000, 100-100,000, 100-1,000,000, 100-10,000,000, 1,000-10,000, 1,000-100,000, 1,000-1,000,000,

1000-10,000,000, 10,000-100,000, 10,000-1,000,000, 10,000-10,000,000, 100,000-1,000,000, or 100,000-10,000,000 times greater than the K_d of the AB not modified with an MM and a CM or of the parental AB towards the target. Conversely, the binding affinity of the AB modified with a MM and a CM towards the target is at least 5, 10, 25, 50, 100, 250, 500, 1,000, 2,500, 5,000, 10,000, 50,000, 100,000, 500,000, 1,000,000, 5,000,000, 10,000,000, 50,000,000 or greater, or between 5-10, 10-100, 10-1,000, 10-10,000, 10-100,000, 10-1,000,000, 10-10,000,000, 100-1,000, 100-10,000, 100-100,000, 100-1,000,000, 100-10,000,000, 1,000-10,000, 1,000-100,000, 1,000-1,000,000, 1000-10,000,000, 10,000-100,000, 10,000-1,000,000, 10,000-10,000,000, 100,000-1,000,000, or 100,000-10,000,000 times lower than the binding affinity of the AB not modified with an MM and a CM or of the parental AB towards the target.

[000360] When the AB is modified with a MM and a CM and is in the presence of the target but not in the presence of a modifying agent (for example at least one protease), specific binding of the AB to its target is reduced or inhibited, as compared to the specific binding of the AB not modified with an MM and a CM or of the parental AB to the target. When compared to the binding of the parental AB or the binding of an AB not modified with an MM and a CM to its target, the AB's ability to bind the target when modified with an MM and a CM can be reduced by at least 50%, 60%, 70%, 80%, 90%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% and even 100% for at least 2, 4, 6, 8, 12, 28, 24, 30, 36, 48, 60, 72, 84, or 96 hours or 5, 10, 15, 30, 45, 60, 90, 120, 150, or 180 days, or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months or longer when measured *in vivo* or in an *in vitro* assay.

[000361] As used herein, the term cleaved state refers to the condition of the activatable antibodies following modification of the CM by at least one protease. The term uncleaved state, as used herein, refers to the condition of the activatable antibodies in the absence of cleavage of the CM by a protease. As discussed above, the term "activatable antibodies" is used herein to refer to an activatable antibody in both its uncleaved (native) state, as well as in its cleaved state. It will be apparent to the ordinarily skilled artisan that in some embodiments a cleaved activatable antibody may lack an MM due to cleavage of the CM by protease, resulting in release of at least the MM (*e.g.*, where the MM is not joined to the activatable antibodies by a covalent bond (*e.g.*, a disulfide bond between cysteine residues)).

[000362] By activatable or switchable is meant that the activatable antibody exhibits a first level of binding to a target when the activatable antibody is in a inhibited, masked or

uncleaved state (*i.e.*, a first conformation), and a second level of binding to the target in the uninhibited, unmasked and/or cleaved state (*i.e.*, a second conformation), where the second level of target binding is greater than the first level of binding. In general, the access of target to the AB of the activatable antibody is greater in the presence of a cleaving agent capable of cleaving the CM, *i.e.*, a protease, than in the absence of such a cleaving agent. Thus, when the activatable antibody is in the uncleaved state, the AB is inhibited from target binding and can be masked from target binding (*i.e.*, the first conformation is such the AB cannot bind the target), and in the cleaved state the AB is not inhibited or is unmasked to target binding.

[000363] The CM and AB of the activatable antibodies are selected so that the AB represents a binding moiety for a given target, and the CM represents a substrate for a protease. In some embodiments, the protease is co-localized with the target at a treatment site or diagnostic site in a subject. As used herein, co-localized refers to being at the same site or relatively close nearby. In some embodiments, a protease cleaves a CM yielding an activated antibody that binds to a target located nearby the cleavage site. The activatable antibodies disclosed herein find particular use where, for example, a protease capable of cleaving a site in the CM, *i.e.*, a protease, is present at relatively higher levels in or in close proximity to target-containing tissue of a treatment site or diagnostic site than in tissue of non-treatment sites (for example in healthy tissue). In some embodiments, a CM of the disclosure is also cleaved by one or more other proteases. In some embodiments, it is the one or more other proteases that is co-localized with the target and that is responsible for cleavage of the CM *in vivo*.

[000364] In some embodiments activatable antibodies provide for reduced toxicity and/or adverse side effects that could otherwise result from binding of the AB at non-treatment sites if the AB were not masked or otherwise inhibited from binding to the target.

[000365] In general, an activatable antibody can be designed by selecting an AB of interest and constructing the remainder of the activatable antibody so that, when conformationally constrained, the MM provides for masking of the AB or reduction of binding of the AB to its target. Structural design criteria can be taken into account to provide for this functional feature.

[000366] Activatable antibodies exhibiting a switchable phenotype of a desired dynamic range for target binding in an inhibited versus an uninhibited conformation are provided. Dynamic range generally refers to a ratio of (a) a maximum detected level of a

parameter under a first set of conditions to (b) a minimum detected value of that parameter under a second set of conditions. For example, in the context of an activatable antibody, the dynamic range refers to the ratio of (a) a maximum detected level of target protein binding to an activatable antibody in the presence of at least one protease capable of cleaving the CM of the activatable antibodies to (b) a minimum detected level of target protein binding to an activatable antibody in the absence of the protease. The dynamic range of an activatable antibody can be calculated as the ratio of the dissociation constant of an activatable antibody cleaving agent (*e.g.*, enzyme) treatment to the dissociation constant of the activatable antibodies cleaving agent treatment. The greater the dynamic range of an activatable antibody, the better the switchable phenotype of the activatable antibody. Activatable antibodies having relatively higher dynamic range values (*e.g.*, greater than 1) exhibit more desirable switching phenotypes such that target protein binding by the activatable antibodies occurs to a greater extent (*e.g.*, predominantly occurs) in the presence of a cleaving agent (*e.g.*, enzyme) capable of cleaving the CM of the activatable antibodies than in the absence of a cleaving agent.

[000367] Activatable antibodies can be provided in a variety of structural configurations. Exemplary formulae for activatable antibodies are provided below. It is specifically contemplated that the N- to C-terminal order of the AB, MM and CM may be reversed within an activatable antibody. It is also specifically contemplated that the CM and MM may overlap in amino acid sequence, *e.g.*, such that the CM is contained within the MM.

[000368] For example, activatable antibodies can be represented by the following formula (in order from an amino (N) terminal region to carboxyl (C) terminal region:

(MM)-(CM)-(AB)

(AB)-(CM)-(MM)

where MM is a masking moiety, CM is a cleavable moiety, and AB is an antibody or fragment thereof. It should be noted that although MM and CM are indicated as distinct components in the formulae above, in all exemplary embodiments (including formulae) disclosed herein it is contemplated that the amino acid sequences of the MM and the CM could overlap, *e.g.*, such that the CM is completely or partially contained within the MM. In addition, the formulae above provide for additional amino acid sequences that may be positioned N-terminal or C-terminal to the activatable antibodies elements.

[000369] In certain embodiments, the MM is not a natural binding partner of the AB. In some embodiments, the MM contains no or substantially no homology to any natural binding partner of the AB. In some embodiments, the MM is no more than 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% similar to any natural binding partner of the AB. In some embodiments, the MM is no more than 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 50% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 25% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 20% identical to any natural binding partner of the AB. In some embodiments, the MM is no more than 10% identical to any natural binding partner of the AB.

[000370] In many embodiments it may be desirable to insert one or more linkers, *e.g.*, flexible linkers, into the activatable antibody construct so as to provide for flexibility at one or more of the MM-CM junction, the CM-AB junction, or both. For example, the AB, MM, and/or CM may not contain a sufficient number of residues (*e.g.*, Gly, Ser, Asp, Asn, especially Gly and Ser, particularly Gly) to provide the desired flexibility. As such, the switchable phenotype of such activatable antibody constructs may benefit from introduction of one or more amino acids to provide for a flexible linker. In addition, as described below, where the activatable antibody is provided as a conformationally constrained construct, a flexible linker can be operably inserted to facilitate formation and maintenance of a cyclic structure in the uncleaved activatable antibody.

[000371] For example, in certain embodiments an activatable antibody comprises one of the following formulae (where the formula below represent an amino acid sequence in either N- to C-terminal direction or C- to N-terminal direction):

(MM)-L1-(CM)-(AB)

(MM)-(CM)-L2-(AB)

(MM)-L1-(CM)-L2-(AB)

wherein MM, CM, and AB are as defined above; wherein L1 and L2 are each independently and optionally present or absent, are the same or different flexible linkers that include at least 1 flexible amino acid (*e.g.*, Gly). In addition, the formulae above provide for additional amino acid sequences that may be positioned N-terminal or C-terminal to the activatable antibodies elements. Examples include, but are not limited to, targeting moieties (*e.g.*, a

ligand for a receptor of a cell present in a target tissue) and serum half-life extending moieties (*e.g.*, polypeptides that bind serum proteins, such as immunoglobulin (*e.g.*, IgG) or serum albumin (*e.g.*, human serum albumin (HAS))).

[000372] The CM is specifically cleaved by at least one protease at a rate of about $0.001\text{-}1500 \times 10^4 \text{ M}^{-1}\text{S}^{-1}$ or at least 0.001, 0.005, 0.01, 0.05, 0.1, 0.5, 1, 2.5, 5, 7.5, 10, 15, 20, 25, 50, 75, 100, 125, 150, 200, 250, 500, 750, 1000, 1250, or $1500 \times 10^4 \text{ M}^{-1}\text{S}^{-1}$. In some embodiments, the CM is specifically cleaved at a rate of about $100,000 \text{ M}^{-1}\text{S}^{-1}$. In some embodiments, the CM is specifically cleaved at a rate from about 1×10^2 to about $1 \times 10^6 \text{ M}^{-1}\text{S}^{-1}$ (*i.e.*, from about 1×10^2 to about $1 \times 10^6 \text{ M}^{-1}\text{S}^{-1}$).

[000373] For specific cleavage by an enzyme, contact between the enzyme and CM is made. When the activatable antibody comprising an AB coupled to a MM and a CM is in the presence of target and sufficient enzyme activity, the CM can be cleaved. Sufficient enzyme activity can refer to the ability of the enzyme to make contact with the CM and effect cleavage. It can readily be envisioned that an enzyme may be in the vicinity of the CM but unable to cleave because of other cellular factors or protein modification of the enzyme.

[000374] Linkers suitable for use in compositions described herein are generally ones that provide flexibility of the modified AB or the activatable antibodies to facilitate the inhibition of the binding of the AB to the target. Such linkers are generally referred to as flexible linkers. Suitable linkers can be readily selected and can be of any of a suitable of different lengths, such as from 1 amino acid (*e.g.*, Gly) to 20 amino acids, from 2 amino acids to 15 amino acids, from 3 amino acids to 12 amino acids, including 4 amino acids to 10 amino acids, 5 amino acids to 9 amino acids, 6 amino acids to 8 amino acids, or 7 amino acids to 8 amino acids, and may be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acids in length.

[000375] Exemplary flexible linkers include glycine polymers (G)_n, glycine-serine polymers (including, for example, (GS)_n, (GSGGS)_n (SEQ ID NO: 191) and (GGGS)_n (SEQ ID NO: 192), where n is an integer of at least one), glycine-alanine polymers, alanine-serine polymers, and other flexible linkers known in the art. Glycine and glycine-serine polymers are relatively unstructured, and therefore may be able to serve as a neutral tether between components. Glycine accesses significantly more phi-psi space than even alanine, and is much less restricted than residues with longer side chains (see Scheraga, *Rev. Computational Chem.* 11173-142 (1992)). Exemplary flexible linkers include, but are not

limited to Gly-Gly-Ser-Gly (SEQ ID NO: 193), Gly-Gly-Ser-Gly-Gly (SEQ ID NO: 194), Gly-Ser-Gly-Ser-Gly (SEQ ID NO: 195), Gly-Ser-Gly-Gly-Gly (SEQ ID NO: 196), Gly-Gly-Gly-Ser-Gly (SEQ ID NO: 197), Gly-Ser-Ser-Ser-Gly (SEQ ID NO: 198), and the like. The ordinarily skilled artisan will recognize that design of an activatable antibodies can include linkers that are all or partially flexible, such that the linker can include a flexible linker as well as one or more portions that confer less flexible structure to provide for a desired activatable antibodies structure.

[000376] The disclosure also provides compositions and methods that include an activatable anti-PDL1 antibody that includes an antibody or antibody fragment (AB) that specifically binds PDL1, where the AB is coupled to a masking moiety (MM) that decreases the ability of the AB to bind its target. In some embodiments, the activatable anti-PDL1 antibody further includes a cleavable moiety (CM) that is a substrate for a protease. The compositions and methods provided herein enable the attachment of one or more agents to one or more cysteine residues in the AB without compromising the activity (e.g., the masking, activating or binding activity) of the activatable anti-PDL1 antibody. In some embodiments, the compositions and methods provided herein enable the attachment of one or more agents to one or more cysteine residues in the AB without reducing or otherwise disturbing one or more disulfide bonds within the MM. The compositions and methods provided herein produce an activatable anti-PDL1 antibody that is conjugated to one or more agents, e.g., any of a variety of therapeutic, diagnostic and/or prophylactic agents, for example, in some embodiments, without any of the agent(s) being conjugated to the MM of the activatable anti-PDL1 antibody. The compositions and methods provided herein produce conjugated activatable anti-PDL1 antibodies in which the MM retains the ability to effectively and efficiently mask the AB of the activatable antibody in an uncleaved state. The compositions and methods provided herein produce conjugated activatable anti-PDL1 antibodies in which the activatable antibody is still activated, *i.e.*, cleaved, in the presence of a protease that can cleave the CM.

[000377] The activatable anti-PDL1 antibodies have at least one point of conjugation for an agent, but in the methods and compositions provided herein less than all possible points of conjugation are available for conjugation to an agent. In some embodiments, the one or more points of conjugation are sulfur atoms involved in disulfide bonds. In some embodiments, the one or more points of conjugation are sulfur atoms involved in interchain disulfide bonds. In some embodiments, the one or more points of conjugation are sulfur

atoms involved in interchain sulfide bonds, but not sulfur atoms involved in intrachain disulfide bonds. In some embodiments, the one or more points of conjugation are sulfur atoms of cysteine or other amino acid residues containing a sulfur atom. Such residues may occur naturally in the antibody structure or may be incorporated into the antibody by site-directed mutagenesis, chemical conversion, or mis-incorporation of non-natural amino acids.

[000378] Also provided are methods of preparing a conjugate of an activatable anti-PDL1 antibody having one or more interchain disulfide bonds in the AB and one or more intrachain disulfide bonds in the MM, and a drug reactive with free thiols is provided. The method generally includes partially reducing interchain disulfide bonds in the activatable antibody with a reducing agent, such as, for example, TCEP; and conjugating the drug reactive with free thiols to the partially reduced activatable antibody. As used herein, the term partial reduction refers to situations where an activatable anti-PDL1 antibody is contacted with a reducing agent and less than all disulfide bonds, *e.g.*, less than all possible sites of conjugation are reduced. In some embodiments, less than 99%, 98%, 97%, 96%, 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10% or less than 5% of all possible sites of conjugation are reduced.

[000379] In yet other embodiments, a method of reducing and conjugating an agent, *e.g.*, a drug, to an activatable anti-PDL1 antibody resulting in selectivity in the placement of the agent is provided. The method generally includes partially reducing the activatable anti-PDL1 antibody with a reducing agent such that any conjugation sites in the masking moiety or other non-AB portion of the activatable antibody are not reduced, and conjugating the agent to interchain thiols in the AB. The conjugation site(s) are selected so as to allow desired placement of an agent to allow conjugation to occur at a desired site. The reducing agent is, for example, TCEP. The reduction reaction conditions such as, for example, the ratio of reducing agent to activatable antibody, the length of incubation, the temperature during the incubation, the pH of the reducing reaction solution, etc., are determined by identifying the conditions that produce a conjugated activatable antibody in which the MM retains the ability to effectively and efficiently mask the AB of the activatable antibody in an uncleaved state. The ratio of reduction agent to activatable anti-PDL1 antibody will vary depending on the activatable antibody. In some embodiments, the ratio of reducing agent to activatable anti-PDL1 antibody will be in a range from about 20:1 to 1:1, from about 10:1 to 1:1, from about 9:1 to 1:1, from about 8:1 to 1:1, from about 7:1 to 1:1, from about 6:1 to

1:1, from about 5:1 to 1:1, from about 4:1 to 1:1, from about 3:1 to 1:1, from about 2:1 to 1:1, from about 20:1 to 1:1.5, from about 10:1 to 1:1.5, from about 9:1 to 1:1.5, from about 8:1 to 1:1.5, from about 7:1 to 1:1.5, from about 6:1 to 1:1.5, from about 5:1 to 1:1.5, from about 4:1 to 1:1.5, from about 3:1 to 1:1.5, from about 2:1 to 1:1.5, from about 1.5:1 to 1:1.5, or from about 1:1 to 1:1.5. In some embodiments, the ratio is in a range of from about 5:1 to 1:1. In some embodiments, the ratio is in a range of from about 5:1 to 1.5:1. In some embodiments, the ratio is in a range of from about 4:1 to 1:1. In some embodiments, the ratio is in a range from about 4:1 to 1.5:1. In some embodiments, the ratio is in a range from about 8:1 to about 1:1. In some embodiments, the ratio is in a range of from about 2.5:1 to 1:1.

[000380] In some embodiments, a method of reducing interchain disulfide bonds in the AB of an activatable anti-PDL1 antibody and conjugating an agent, e.g., a thiol-containing agent such as a drug, to the resulting interchain thiols to selectively locate agent(s) on the AB is provided. The method generally includes partially reducing the AB with a reducing agent to form at least two interchain thiols without forming all possible interchain thiols in the activatable antibody; and conjugating the agent to the interchain thiols of the partially reduced AB. For example, the AB of the activatable antibody is partially reduced for about 1 hour at about 37°C at a desired ratio of reducing agent:activatable antibody. In some embodiments, the ratio of reducing agent to activatable antibody will be in a range from about 20:1 to 1:1, from about 10:1 to 1:1, from about 9:1 to 1:1, from about 8:1 to 1:1, from about 7:1 to 1:1, from about 6:1 to 1:1, from about 5:1 to 1:1, from about 4:1 to 1:1, from about 3:1 to 1:1, from about 2:1 to 1:1, from about 20:1 to 1:1.5, from about 10:1 to 1:1.5, from about 9:1 to 1:1.5, from about 8:1 to 1:1.5, from about 7:1 to 1:1.5, from about 6:1 to 1:1.5, from about 5:1 to 1:1.5, from about 4:1 to 1:1.5, from about 3:1 to 1:1.5, from about 2:1 to 1:1.5, from about 1.5:1 to 1:1.5, or from about 1:1 to 1:1.5. In some embodiments, the ratio is in a range of from about 5:1 to 1:1. In some embodiments, the ratio is in a range of from about 5:1 to 1.5:1. In some embodiments, the ratio is in a range of from about 4:1 to 1:1. In some embodiments, the ratio is in a range from about 4:1 to 1.5:1. In some embodiments, the ratio is in a range from about 8:1 to about 1:1. In some embodiments, the ratio is in a range of from about 2.5:1 to 1:1.

[000381] The thiol-containing reagent can be, for example, cysteine or N-acetyl cysteine. The reducing agent can be, for example, TCEP. In some embodiments, the reduced activatable antibody can be purified prior to conjugation, using for example,

column chromatography, dialysis, or diafiltration. Alternatively, the reduced antibody is not purified after partial reduction and prior to conjugation.

[000382] The invention also provides partially reduced activatable anti-PDL1 antibodies in which at least one interchain disulfide bond in the activatable antibody has been reduced with a reducing agent without disturbing any intrachain disulfide bonds in the activatable antibody, wherein the activatable antibody includes an antibody or an antigen binding fragment thereof (AB) that specifically binds to PDL1, a masking moiety (MM) that inhibits the binding of the AB of the activatable antibody in an uncleaved state to the PDL1 target, and a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease. In some embodiments the MM is coupled to the AB via the CM. In some embodiments, one or more intrachain disulfide bond(s) of the activatable antibody is not disturbed by the reducing agent. In some embodiments, one or more intrachain disulfide bond(s) of the MM within the activatable antibody is not disturbed by the reducing agent. In some embodiments, the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM. In some embodiments, reducing agent is TCEP.

[000383] The disclosure also provides partially reduced activatable antibodies in which at least one interchain disulfide bond in the activatable antibody has been reduced with a reducing agent without disturbing any intrachain disulfide bonds in the activatable antibody, wherein the activatable antibody includes an antibody or an antigen binding fragment thereof (AB) that specifically binds to the target, e.g., PDL1, a masking moiety (MM) that inhibits the binding of the AB of the activatable antibody in an uncleaved state to the target, and a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for at least one protease. In some embodiments, the MM is coupled to the AB via the CM. In some embodiments, one or more intrachain disulfide bond(s) of the activatable antibody is not disturbed by the reducing agent. In some embodiments, one or more intrachain disulfide bond(s) of the MM within the activatable antibody is not disturbed by the reducing agent. In some embodiments, the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM. In some embodiments, reducing agent is TCEP.

[000384] In some embodiments, the activatable antibodies described herein also include an agent conjugated to the activatable antibody. In some embodiments, the conjugated agent is a therapeutic agent, such as an anti-inflammatory and/or an antineoplastic agent. In such embodiments, the agent is conjugated to a carbohydrate moiety of the activatable antibody, for example, in some embodiments, where the carbohydrate moiety is located outside the antigen-binding region of the antibody or antigen-binding fragment in the activatable antibody. In some embodiments, the agent is conjugated to a sulfhydryl group of the antibody or antigen-binding fragment in the activatable antibody.

[000385] In some embodiments, the agent is a cytotoxic agent such as a toxin (*e.g.*, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (*i.e.*, a radioconjugate).


[000386] In some embodiments, the agent is a detectable moiety such as, for example, a label or other marker. For example, the agent is or includes a radiolabeled amino acid, one or more biotinyl moieties that can be detected by marked avidin (*e.g.*, streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or calorimetric methods), one or more radioisotopes or radionuclides, one or more fluorescent labels, one or more enzymatic labels, and/or one or more chemiluminescent agents. In some embodiments, detectable moieties are attached by spacer molecules.

[000387] The disclosure also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a toxin (*e.g.*, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (*i.e.*, a radioconjugate). Suitable cytotoxic agents include, for example, dolastatins and derivatives thereof (*e.g.* auristatin E, AFP, MMAF, MMAE, MMAD, DMAF, DMAE). For example, the agent is monomethyl auristatin E (MMAE) or monomethyl auristatin D (MMAD). In some embodiments, the agent is an agent selected from the group listed in Table 11. In some embodiments, the agent is a dolastatin. In some embodiments, the agent is an auristatin or derivative thereof. In some embodiments, the agent is auristatin E or a derivative thereof. In some embodiments, the agent is monomethyl auristatin E (MMAE). In some embodiments, the agent is monomethyl auristatin D (MMAD). In some embodiments, the agent is a maytansinoid or maytansinoid derivative. In some embodiments, the agent is DM1 or DM4. In some embodiments, the agent is a duocarmycin or derivative thereof. In some


embodiments, the agent is a pyrrolobenzodiazepine. In some embodiments, the agent is a pyrrolobenzodiazepine dimer.

[000388] In some embodiments, the agent is linked to the AB using a maleimide caproyl-valine-citrulline linker or a maleimide PEG-valine-citrulline linker. In some embodiments, the agent is linked to the AB using a maleimide caproyl-valine-citrulline linker. In some embodiments, the agent is linked to the AB using a maleimide PEG-valine-citrulline linker. In some embodiments, the agent is monomethyl auristatin D (MMAD) linked to the AB using a maleimide PEG-valine-citrulline-para-aminobenzyloxycarbonyl linker, and this linker payload construct is referred to herein as “vc-MMAD.” In some embodiments, the agent is monomethyl auristatin E (MMAE) linked to the AB using a maleimide PEG-valine-citrulline-para-aminobenzyloxycarbonyl linker, and this linker payload construct is referred to herein as “vc-MMAE.” The structures of vc-MMAD and vc-MMAE are shown below:

vc-MMAD:



vc-MMAE:



[000389] The disclosure also provides conjugated activatable antibodies that include an activatable antibody linked to monomethyl auristatin D (MMAD) payload, wherein the activatable antibody includes an antibody or an antigen binding fragment thereof (AB) that specifically binds to a target, a masking moiety (MM) that inhibits the binding of the AB of the activatable antibody in an uncleaved state to the target, and cleavable moiety (CM)

coupled to the AB, and the CM is a polypeptide that functions as a substrate for at least one MMP protease.

[000390] In some embodiments, the MMAD-conjugated activatable antibody can be conjugated using any of several methods for attaching agents to ABs: (a) attachment to the carbohydrate moieties of the AB, or (b) attachment to sulfhydryl groups of the AB, or (c) attachment to amino groups of the AB, or (d) attachment to carboxylate groups of the AB.

[000391] In some embodiments, the MMAD payload is conjugated to the AB via a linker. In some embodiments, the MMAD payload is conjugated to a cysteine in the AB via a linker. In some embodiments, the MMAD payload is conjugated to a lysine in the AB via a linker. In some embodiments, the MMAD payload is conjugated to another residue of the AB via a linker, such as those residues disclosed herein. In some embodiments, the linker is a thiol-containing linker. In some embodiments, the linker is a cleavable linker. In some embodiments, the linker is a non-cleavable linker. In some embodiments, the linker is selected from the group consisting of the linkers shown in Tables 13 and 14. In some embodiments, the activatable antibody and the MMAD payload are linked via a maleimide caproyl-valine-citrulline linker. In some embodiments, the activatable antibody and the MMAD payload are linked via a maleimide PEG-valine-citrulline linker. In some embodiments, the activatable antibody and the MMAD payload are linked via a maleimide caproyl-valine-citrulline-para-aminobenzyloxycarbonyl linker. In some embodiments, the activatable antibody and the MMAD payload are linked via a maleimide PEG-valine-citrulline-para-aminobenzyloxycarbonyl linker. In some embodiments, the MMAD payload is conjugated to the AB using the partial reduction and conjugation technology disclosed herein.

[000392] In some embodiments, the polyethylene glycol (PEG) component of a linker of the present disclosure is formed from 2 ethylene glycol monomers, 3 ethylene glycol monomers, 4 ethylene glycol monomers, 5 ethylene glycol monomers, 6 ethylene glycol monomers, 7 ethylene glycol monomers, 8 ethylene glycol monomers, 9 ethylene glycol monomers, or at least 10 ethylene glycol monomers. In some embodiments of the present disclosure, the PEG component is a branched polymer. In some embodiments of the present disclosure, the PEG component is an unbranched polymer. In some embodiments, the PEG polymer component is functionalized with an amino group or derivative thereof, a carboxyl group or derivative thereof, or both an amino group or derivative thereof and a carboxyl group or derivative thereof.

[000393] In some embodiments, the PEG component of a linker of the present disclosure is an amino-tetra-ethylene glycol-carboxyl group or derivative thereof. In some embodiments, the PEG component of a linker of the present disclosure is an amino-tri-ethylene glycol-carboxyl group or derivative thereof. In some embodiments, the PEG component of a linker of the present disclosure is an amino-di-ethylene glycol-carboxyl group or derivative thereof. In some embodiments, an amino derivative is the formation of an amide bond between the amino group and a carboxyl group to which it is conjugated. In some embodiments, a carboxyl derivative is the formation of an amide bond between the carboxyl group and an amino group to which it is conjugated. In some embodiments, a carboxyl derivative is the formation of an ester bond between the carboxyl group and an hydroxyl group to which it is conjugated.

[000394] Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolaca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re .

[000395] Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., Science 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. (See WO94/11026).

[000396] Table 11 lists some of the exemplary pharmaceutical agents that may be employed in the herein described disclosure but in no way is meant to be an exhaustive list.

Table 11: Exemplary Pharmaceutical Agents for Conjugation

CYTOTOXIC AGENTS

Auristatins
Auristatin E
Monomethyl auristatin D (MMAD)
Monomethyl auristatin E (MMAE)
Desmethyl auristatin E (DMAE)
Auristatin F
Monomethyl auristatin F (MMAF)
Desmethyl auristatin F (DMAF)
Auristatin derivatives, *e.g.*, amides thereof
Auristatin tyramine
Auristatin quinoline
Dolastatins
Dolastatin derivatives
Dolastatin 16 DmJ
Dolastatin 16 Dpv
Maytansinoids, *e.g.* DM-1; DM-4
Maytansinoid derivatives
Duocarmycin
Duocarmycin derivatives
Alpha-amanitin
Anthracyclines
Doxorubicin

Daunorubicin
Bryostatins
Camptothecin
Camptothecin derivatives
7-substituted Camptothecin
10, 11-
Difluoromethylenedioxcamptothecin
Combretastatins
Debromoaplysiatoxin

Kahalalide-F
Discodermolide
Ecteinascidins

ANTIVIRALS

Acyclovir
Vira A
Symmetrel

ANTIFUNGALS

Nystatin

Turbostatin
Phenstatins
Hydroxyphenstatin
Spongistatin 5
Spongistatin 7
Halistatin 1
Halistatin 2
Halistatin 3
Modified Bryostatins
Halocomstatins
Pyrrolobenzimidazoles (PBI)
Cibrostatin6
Doxaliform
Anthracyclins analogues

Cemadotin analogue (CemCH₂-SH)
Pseudomonas toxin A (PE38) variant
Pseudomonas toxin A (ZZ-PE38) variant
ZJ-101
OSW-1
4-Nitrobenzyloxycarbonyl Derivatives of
O6-Benzylguanine
Topoisomerase inhibitors
Hemiasterlin
Cephalotaxine
Homoharringtonine
Pyrrolobenzodiazepine dimers (PBDs)
Pyrrolobenzodiazepenes

Functionalized pyrrolobenzodiazepenes
Functionalized pyrrolobenzodiazepene
dimers
Calicheamicins
Podophyllotoxins
Taxanes
Vinca alkaloids

**CONJUGATABLE DETECTION
REAGENTS**

Fluorescein and derivatives thereof
Fluorescein isothiocyanate (FITC)

RADIOPHARMACEUTICALS

¹²⁵I
¹³¹I

ADDITIONAL ANTI-NEOPLASTICS

Adriamycin
Cerubidine
Bleomycin
Alkeran
Velban
Oncovin
Fluorouracil
Methotrexate
Thiotepa
Bisantrene
Novantrone
Thioguanine
Procarabazine
Cytarabine

⁸⁹Zr
¹¹¹In
¹²³I
¹³¹I
^{99m}Tc
²⁰¹Tl
¹³³Xe
¹¹C
⁶²Cu
¹⁸F
⁶⁸Ga
¹³N
¹⁵O
³⁸K
⁸²Rb
^{99m}Tc (Technetium)

ANTI-BACTERIALS

Aminoglycosides
Streptomycin
Neomycin
Kanamycin
Amikacin
Gentamicin
Tobramycin
Streptomycin B
Spectinomycin
Ampicillin
Sulfanilamide
Polymyxin
Chloramphenicol

HEAVY METALS

Barium
Gold
Platinum

ANTI-MYCOPLASMA

Tylosine
Spectinomycin

[000397] Those of ordinary skill in the art will recognize that a large variety of possible moieties can be coupled to the resultant antibodies of the disclosure. (*See, for example*, "Conjugate Vaccines", Contributions to Microbiology and Immunology, J. M. Cruse and R. E. Lewis, Jr (eds), Carger Press, New York, (1989), the entire contents of which are incorporated herein by reference).

[000398] Coupling may be accomplished by any chemical reaction that will bind the two molecules so long as the antibody and the other moiety retain their respective activities. This linkage can include many chemical mechanisms, for instance covalent binding, affinity binding, intercalation, coordinate binding and complexation. In some embodiments, the binding is, however, covalent binding. Covalent binding can be achieved either by direct

condensation of existing side chains or by the incorporation of external bridging molecules. Many bivalent or polyvalent linking agents are useful in coupling protein molecules, such as the antibodies of the present disclosure, to other molecules. For example, representative coupling agents can include organic compounds such as thioesters, carbodiimides, succinimide esters, diisocyanates, glutaraldehyde, diazobenzenes and hexamethylene diamines. This listing is not intended to be exhaustive of the various classes of coupling agents known in the art but, rather, is exemplary of the more common coupling agents. (See Killen and Lindstrom, *Jour. Immun.* 133:1335-2549 (1984); Jansen et al., *Immunological Reviews* 62:185-216 (1982); and Vitetta et al., *Science* 238:1098 (1987).

[000399] In some embodiments, in addition to the compositions and methods provided herein, the conjugated activatable antibody can also be modified for site-specific conjugation through modified amino acid sequences inserted or otherwise included in the activatable antibody sequence. These modified amino acid sequences are designed to allow for controlled placement and/or dosage of the conjugated agent within a conjugated activatable antibody. For example, the activatable antibody can be engineered to include cysteine substitutions at positions on light and heavy chains that provide reactive thiol groups and do not negatively impact protein folding and assembly, nor alter antigen binding. In some embodiments, the activatable antibody can be engineered to include or otherwise introduce one or more non-natural amino acid residues within the activatable antibody to provide suitable sites for conjugation. In some embodiments, the activatable antibody can be engineered to include or otherwise introduce enzymatically activatable peptide sequences within the activatable antibody sequence.

[000400] Suitable linkers are described in the literature. (See, for example, Ramakrishnan, S. et al., *Cancer Res.* 44:201-208 (1984) describing use of MBS (M-maleimidobenzoyl-N-hydroxysuccinimide ester). See also, U.S. Patent No. 5,030,719, describing use of halogenated acetyl hydrazide derivative coupled to an antibody by way of an oligopeptide linker. In some embodiments, suitable linkers include: (i) EDC (1-ethyl-3-(3-dimethylamino-propyl) carbodiimide hydrochloride; (ii) SMPT (4-succinimidylloxycarbonyl-alpha-methyl-alpha-(2-pyridyl-dithio)-toluene (Pierce Chem. Co., Cat. (21558G); (iii) SPDP (succinimidyl-6 [3-(2-pyridyldithio) propionamido]hexanoate (Pierce Chem. Co., Cat #21651G); (iv) Sulfo-LC-SPDP (sulfosuccinimidyl 6 [3-(2-pyridyldithio)-propionamide] hexanoate (Pierce Chem. Co. Cat. #2165-G); and (v) sulfo-NHS (N-hydroxysulfo-succinimide: Pierce Chem. Co., Cat. #24510) conjugated to EDC.

Additional linkers include, but are not limited to, SMCC, sulfo-SMCC, SPDB, or sulfo-SPDB.

[000401] The linkers described above contain components that have different attributes, thus leading to conjugates with differing physio-chemical properties. For example, sulfo-NHS esters of alkyl carboxylates are more stable than sulfo-NHS esters of aromatic carboxylates. NHS-ester containing linkers are less soluble than sulfo-NHS esters. Further, the linker SMPT contains a sterically hindered disulfide bond, and can form conjugates with increased stability. Disulfide linkages, are in general, less stable than other linkages because the disulfide linkage is cleaved *in vitro*, resulting in less conjugate available. Sulfo-NHS, in particular, can enhance the stability of carbodimide couplings. Carbodimide couplings (such as EDC) when used in conjunction with sulfo-NHS, forms esters that are more resistant to hydrolysis than the carbodimide coupling reaction alone.

[000402] In some embodiments, the linkers are cleavable. In some embodiments, the linkers are non-cleavable. In some embodiments, two or more linkers are present. The two or more linkers are all the same, *i.e.*, cleavable or non-cleavable, or the two or more linkers are different, *i.e.*, at least one cleavable and at least one non-cleavable.

[000403] The present disclosure utilizes several methods for attaching agents to ABs: (a) attachment to the carbohydrate moieties of the AB, or (b) attachment to sulfhydryl groups of the AB, or (c) attachment to amino groups of the AB, or (d) attachment to carboxylate groups of the AB. According to the disclosure, ABs may be covalently attached to an agent through an intermediate linker having at least two reactive groups, one to react with AB and one to react with the agent. The linker, which may include any compatible organic compound, can be chosen such that the reaction with AB (or agent) does not adversely affect AB reactivity and selectivity. Furthermore, the attachment of linker to agent might not destroy the activity of the agent. Suitable linkers for reaction with oxidized antibodies or oxidized antibody fragments include those containing an amine selected from the group consisting of primary amine, secondary amine, hydrazine, hydrazide, hydroxylamine, phenylhydrazine, semicarbazide and thiosemicarbazide groups. Such reactive functional groups may exist as part of the structure of the linker, or may be introduced by suitable chemical modification of linkers not containing such groups.

[000404] According to the present disclosure, suitable linkers for attachment to reduced ABs include those having certain reactive groups capable of reaction with a sulfhydryl group of a reduced antibody or fragment. Such reactive groups include, but are

not limited to: reactive haloalkyl groups (including, for example, haloacetyl groups), p-mercuribenzoate groups and groups capable of Michael-type addition reactions (including, for example, maleimides and groups of the type described by Mitra and Lawton, 1979, J. Amer. Chem. Soc. 101: 3097-3110).

[000405] According to the present disclosure, suitable linkers for attachment to neither oxidized nor reduced Abs include those having certain functional groups capable of reaction with the primary amino groups present in unmodified lysine residues in the Ab. Such reactive groups include, but are not limited to, NHS carboxylic or carbonic esters, sulfo-NHS carboxylic or carbonic esters, 4-nitrophenyl carboxylic or carbonic esters, pentafluorophenyl carboxylic or carbonic esters, acyl imidazoles, isocyanates, and isothiocyanates.

[000406] According to the present disclosure, suitable linkers for attachment to neither oxidized nor reduced Abs include those having certain functional groups capable of reaction with the carboxylic acid groups present in aspartate or glutamate residues in the Ab, which have been activated with suitable reagents. Suitable activating reagents include EDC, with or without added NHS or sulfo-NHS, and other dehydrating agents utilized for carboxamide formation. In these instances, the functional groups present in the suitable linkers would include primary and secondary amines, hydrazines, hydroxylamines, and hydrazides.

[000407] The agent may be attached to the linker before or after the linker is attached to the AB. In certain applications it may be desirable to first produce an AB-linker intermediate in which the linker is free of an associated agent. Depending upon the particular application, a specific agent may then be covalently attached to the linker. In some embodiments, the AB is first attached to the MM, CM and associated linkers and then attached to the linker for conjugation purposes.

[000408] *Branched Linkers:* In specific embodiments, branched linkers that have multiple sites for attachment of agents are utilized. For multiple site linkers, a single covalent attachment to an AB would result in an AB-linker intermediate capable of binding an agent at a number of sites. The sites may be aldehyde or sulfhydryl groups or any chemical site to which agents can be attached.

[000409] In some embodiments, higher specific activity (or higher ratio of agents to AB) can be achieved by attachment of a single site linker at a plurality of sites on the AB. This plurality of sites may be introduced into the AB by either of two methods. First, one may generate multiple aldehyde groups and/or sulfhydryl groups in the same AB. Second,

one may attach to an aldehyde or sulfhydryl of the AB a "branched linker" having multiple functional sites for subsequent attachment to linkers. The functional sites of the branched linker or multiple site linker may be aldehyde or sulfhydryl groups, or may be any chemical site to which linkers may be attached. Still higher specific activities may be obtained by combining these two approaches, that is, attaching multiple site linkers at several sites on the AB.

[000410] *Cleavable Linkers:* Peptide linkers that are susceptible to cleavage by enzymes of the complement system, such as but not limited to u-plasminogen activator, tissue plasminogen activator, trypsin, plasmin, or another enzyme having proteolytic activity may be used in one embodiment of the present disclosure. According to one method of the present disclosure, an agent is attached via a linker susceptible to cleavage by complement. The antibody is selected from a class that can activate complement. The antibody-agent conjugate, thus, activates the complement cascade and releases the agent at the target site. According to another method of the present disclosure, an agent is attached via a linker susceptible to cleavage by enzymes having a proteolytic activity such as a u-plasminogen activator, a tissue plasminogen activator, plasmin, or trypsin. These cleavable linkers are useful in conjugated activatable antibodies that include an extracellular toxin, e.g., by way of non-limiting example, any of the extracellular toxins shown in Table 11.

[000411] Non-limiting examples of cleavable linker sequences are provided in Table 13.

Table 13: Exemplary Linker Sequences for Conjugation

Types of Cleavable Sequences	Amino Acid Sequence
<u>Plasmin cleavable sequences</u>	
Pro-urokinase	PRFKIIGG (SEQ ID NO: 396) PRFRIIGG (SEQ ID NO: 397)
TGFβ	SSRHRRALD (SEQ ID NO: 398)
Plasminogen	RKSSIIIRMRDVVL (SEQ ID NO: 399)
Staphylokinase	SSSFDKGKYKKGDDA (SEQ ID NO: 400) SSSFDKGKYKRGDDA (SEQ ID NO: 401)
<u>Factor Xa cleavable sequences</u>	
	IEGR (SEQ ID NO: 402) IDGR (SEQ ID NO: 403) GGSIDGR (SEQ ID NO: 404)
<u>MMP cleavable sequences</u>	
Gelatinase A	PLGLWA (SEQ ID NO: 405)

Collagenase cleavable sequences

Calf skin collagen ($\alpha 1$ (I) chain)	GPQGIAGQ (SEQ ID NO: 406)
Calf skin collagen ($\alpha 2$ (I) chain)	GPQGLLGA (SEQ ID NO: 407)
Bovine cartilage collagen ($\alpha 1$ (II) chain)	GIAGQ (SEQ ID NO: 408)
Human liver collagen ($\alpha 1$ (III) chain)	GPLGIAGI (SEQ ID NO: 409)
Human α_2 M	GPEGLRVG (SEQ ID NO: 410)
Human PZP	YGAGLGVV (SEQ ID NO: 411)
	AGLGVVER (SEQ ID NO: 412)
	AGLGISST (SEQ ID NO: 413)
Rat α_1 M	EPQALAMS (SEQ ID NO: 414)
	QALAMSAI (SEQ ID NO: 415)
Rat α_2 M	AAYHLVSQ (SEQ ID NO: 416)
	MDAFLESS (SEQ ID NO: 417)
Rat α_1 I ₃ (2J)	ESLPVVAV (SEQ ID NO: 418)
Rat α_1 I ₃ (27J)	SAPAVESE (SEQ ID NO: 419)
Human fibroblast collagenase (autolytic cleavages)	DVAQFVLT (SEQ ID NO: 420)
	VAQFVLTE (SEQ ID NO: 421)
	AQFVLTEG (SEQ ID NO: 422)
	PVQPIGPQ (SEQ ID NO: 423)

[000412] In addition, agents may be attached via disulfide bonds (for example, the disulfide bonds on a cysteine molecule) to the AB. Since many tumors naturally release high levels of glutathione (a reducing agent) this can reduce the disulfide bonds with subsequent release of the agent at the site of delivery. In some embodiments, the reducing agent that would modify a CM would also modify the linker of the conjugated activatable antibody.

[000413] *Spacers and Cleavable Elements:* In some embodiments, it may be necessary to construct the linker in such a way as to optimize the spacing between the agent and the AB of the activatable antibody. This may be accomplished by use of a linker of the general structure:



wherein

W is either --NH--CH₂-- or --CH₂--;

Q is an amino acid, peptide; and

n is an integer from 0 to 20.

[000414] In some embodiments, the linker may comprise a spacer element and a cleavable element. The spacer element serves to position the cleavable element away from the core of the AB such that the cleavable element is more accessible to the enzyme

responsible for cleavage. Certain of the branched linkers described above may serve as spacer elements.

[000415] Throughout this discussion, it should be understood that the attachment of linker to agent (or of spacer element to cleavable element, or cleavable element to agent) need not be particular mode of attachment or reaction. Any reaction providing a product of suitable stability and biological compatibility is acceptable.

[000416] *Serum Complement and Selection of Linkers:* According to one method of the present disclosure, when release of an agent is desired, an AB that is an antibody of a class that can activate complement is used. The resulting conjugate retains both the ability to bind antigen and activate the complement cascade. Thus, according to this embodiment of the present disclosure, an agent is joined to one end of the cleavable linker or cleavable element and the other end of the linker group is attached to a specific site on the AB. For example, if the agent has an hydroxy group or an amino group, it may be attached to the carboxy terminus of a peptide, amino acid or other suitably chosen linker via an ester or amide bond, respectively. For example, such agents may be attached to the linker peptide via a carbodimide reaction. If the agent contains functional groups that would interfere with attachment to the linker, these interfering functional groups can be blocked before attachment and deblocked once the product conjugate or intermediate is made. The opposite or amino terminus of the linker is then used either directly or after further modification for binding to an AB that is capable of activating complement.

[000417] Linkers (or spacer elements of linkers) may be of any desired length, one end of which can be covalently attached to specific sites on the AB of the activatable antibody. The other end of the linker or spacer element may be attached to an amino acid or peptide linker.

[000418] Thus when these conjugates bind to antigen in the presence of complement the amide or ester bond that attaches the agent to the linker will be cleaved, resulting in release of the agent in its active form. These conjugates, when administered to a subject, will accomplish delivery and release of the agent at the target site, and are particularly effective for the in vivo delivery of pharmaceutical agents, antibiotics, antimetabolites, antiproliferative agents and the like as presented in but not limited to those in Table 11.

[000419] *Linkers for Release without Complement Activation:* In yet another application of targeted delivery, release of the agent without complement activation is desired since activation of the complement cascade will ultimately lyse the target cell.

Hence, this approach is useful when delivery and release of the agent should be accomplished without killing the target cell. Such is the goal when delivery of cell mediators such as hormones, enzymes, corticosteroids, neurotransmitters, genes or enzymes to target cells is desired. These conjugates may be prepared by attaching the agent to an AB that is not capable of activating complement via a linker that is mildly susceptible to cleavage by serum proteases. When this conjugate is administered to an individual, antigen-antibody complexes will form quickly whereas cleavage of the agent will occur slowly, thus resulting in release of the compound at the target site.

[000420] *Biochemical Cross Linkers:* In some embodiments, the activatable antibody may be conjugated to one or more therapeutic agents using certain biochemical cross-linkers. Cross-linking reagents form molecular bridges that tie together functional groups of two different molecules. To link two different proteins in a step-wise manner, hetero-bifunctional cross-linkers can be used that eliminate unwanted homopolymer formation.

[000421] Peptidyl linkers cleavable by lysosomal proteases are also useful, for example, Val-Cit, Val-Ala or other dipeptides. In addition, acid-labile linkers cleavable in the low-pH environment of the lysosome may be used, for example: bis-sialyl ether. Other suitable linkers include cathepsin-labile substrates, particularly those that show optimal function at an acidic pH.

[000422] Exemplary hetero-bifunctional cross-linkers are referenced in Table 14.

Table 14: Exemplary Hetero-Bifunctional Cross Linkers

<u>HETERO-BIFUNCTIONAL CROSS-LINKERS</u>			
Linker	Reactive Toward	Advantages and Applications	Spacer Arm Length after cross-linking (Angstroms)
SMPT	Primary amines Sulfhydryls	Greater stability	11.2 Å
SPDP	Primary amines Sulfhydryls	Thiolation Cleavable cross-linking	6.8 Å
LC-SPDP	Primary amines Sulfhydryls	Extended spacer arm	15.6 Å
Sulfo-LC-SPDP	Primary amines Sulfhydryls	Extender spacer arm Water-soluble	15.6 Å
SMCC	Primary amines Sulfhydryls	Stable maleimide reactive group Enzyme-antibody conjugation Hapten-carrier protein	11.6 Å

Sulfo-SMCC	Primary amines	conjugation Stable maleimide reactive group	11.6 Å
	Sulfhydryls	Water-soluble	
MBS	Primary amines	Enzyme-antibody conjugation	9.9 Å
	Sulfhydryls	Hapten-carrier protein conjugation	
Sulfo-MBS	Primary amines	Water-soluble	9.9 Å
	Sulfhydryls		
SIAB	Primary amines	Enzyme-antibody conjugation	10.6 Å
	Sulfhydryls		
Sulfo-SIAB	Primary amines	Water-soluble	10.6 Å
	Sulfhydryls		
SMPB	Primary amines	Extended spacer arm	14.5 Å
	Sulfhydryls	Enzyme-antibody conjugation	
Sulfo-SMPB	Primary amines	Extended spacer arm	14.5 Å
	Sulfhydryls	Water-soluble	
EDE/Sulfo-NHS	Primary amines	Hapten-Carrier conjugation	0
	Carboxyl groups		
ABH	Carbohydrates	Reacts with sugar groups	11.9 Å
	Nonselective		

[000423] *Non-Cleavable Linkers or Direct Attachment:* In some embodiments of the disclosure, the conjugate may be designed so that the agent is delivered to the target but not released. This may be accomplished by attaching an agent to an AB either directly or via a non-cleavable linker.

[000424] These non-cleavable linkers may include amino acids, peptides, D-amino acids or other organic compounds that may be modified to include functional groups that can subsequently be utilized in attachment to ABs by the methods described herein. A-general formula for such an organic linker could be



wherein

W is either --NH--CH₂-- or --CH₂--;

Q is an amino acid, peptide; and

n is an integer from 0 to 20.

[000425] *Non-Cleavable Conjugates:* In some embodiments, a compound may be attached to ABs that do not activate complement. When using ABs that are incapable of complement activation, this attachment may be accomplished using linkers that are susceptible to cleavage by activated complement or using linkers that are not susceptible to cleavage by activated complement.

[000426] The antibodies disclosed herein can also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang et al., Proc. Natl Acad. Sci. USA, 77: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

[000427] Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present disclosure can be conjugated to the liposomes as described in Martin et al., J. Biol. Chem., 257: 286-288 (1982) via a disulfide-interchange reaction.

Definitions:

[000428] Unless otherwise defined, scientific and technical terms used in connection with the present disclosure shall have the meanings that are commonly understood by those of ordinary skill in the art. The term "a" entity or "an" entity refers to one or more of that entity. For example, a compound refers to one or more compounds. As such, the terms "a", "an", "one or more" and "at least one" can be used interchangeably. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures utilized in connection with, and techniques of, cell and tissue culture, molecular biology, and protein and oligo- or polynucleotide chemistry and hybridization described herein are those well-known and commonly used in the art. Standard techniques are used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (*e.g.*, electroporation, lipofection). Enzymatic reactions and purification techniques are performed according to manufacturer's specifications or as commonly accomplished in the art or as described herein. The foregoing techniques and procedures are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. *See e.g.*, Sambrook *et al.* Molecular Cloning: A Laboratory Manual (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)). The nomenclatures utilized in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well-known and commonly used in the

art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[000429] As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

[000430] As used herein, the term “antibody” refers to immunoglobulin molecules and immunologically active portions of immunoglobulin (Ig) molecules, *i.e.*, molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen. By “specifically bind” or “immunoreacts with” or “immunospecifically bind” is meant that the antibody reacts with one or more antigenic determinants of the desired antigen and does not react with other polypeptides or binds at much lower affinity ($K_d > 10^{-6}$). Antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, domain antibody, single chain, Fab, and F(ab')₂ fragments, scFvs, and an Fab expression library.

[000431] The basic antibody structural unit is known to comprise a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one “light” (about 25 kDa) and one “heavy” chain (about 50-70 kDa). The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. In general, antibody molecules obtained from humans relate to any of the classes IgG, IgM, IgA, IgE and IgD, which differ from one another by the nature of the heavy chain present in the molecule. Certain classes have subclasses as well, such as IgG₁, IgG₂, and others. Furthermore, in humans, the light chain may be a kappa chain or a lambda chain.

[000432] The term “monoclonal antibody” (mAb) or “monoclonal antibody composition”, as used herein, refers to a population of antibody molecules that contain only one molecular species of antibody molecule consisting of a unique light chain gene product and a unique heavy chain gene product. In particular, the complementarity determining regions (CDRs) of the monoclonal antibody are identical in all the molecules of the population. MAbs contain an antigen binding site capable of immunoreacting with a particular epitope of the antigen characterized by a unique binding affinity for it.

[000433] The term “antigen-binding site” or “binding portion” refers to the part of the immunoglobulin molecule that participates in antigen binding. The antigen binding site is formed by amino acid residues of the N-terminal variable (“V”) regions of the heavy (“H”) and light (“L”) chains. Three highly divergent stretches within the V regions of the heavy

and light chains, referred to as “hypervariable regions,” are interposed between more conserved flanking stretches known as “framework regions,” or “FRs”. Thus, the term “FR” refers to amino acid sequences that are naturally found between, and adjacent to, hypervariable regions in immunoglobulins. In an antibody molecule, the three hypervariable regions of a light chain and the three hypervariable regions of a heavy chain are disposed relative to each other in three dimensional space to form an antigen-binding surface. The antigen-binding surface is complementary to the three-dimensional surface of a bound antigen, and the three hypervariable regions of each of the heavy and light chains are referred to as “complementarity-determining regions,” or “CDRs.” The assignment of amino acids to each domain is in accordance with the definitions of Kabat Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md. (1987 and 1991)), or Chothia & Lesk J. Mol. Biol. 196:901-917 (1987), Chothia *et al.* Nature 342:878-883 (1989).

[000434] As used herein, the term “epitope” includes any protein determinant capable of specific binding to an immunoglobulin, an scFv, or a T-cell receptor. The term “epitope” includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor. Epitopic determinants usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics. For example, antibodies may be raised against N-terminal or C-terminal peptides of a polypeptide. An antibody is said to specifically bind an antigen when the dissociation constant is $\leq 1 \mu\text{M}$; in some embodiments, $\leq 100 \text{ nM}$ and in some embodiments, $\leq 10 \text{ nM}$.

[000435] As used herein, the terms “specific binding,” “immunological binding,” and “immunological binding properties” refer to the non-covalent interactions of the type which occur between an immunoglobulin molecule and an antigen for which the immunoglobulin is specific. The strength, or affinity of immunological binding interactions can be expressed in terms of the dissociation constant (K_d) of the interaction, wherein a smaller K_d represents a greater affinity. Immunological binding properties of selected polypeptides can be quantified using methods well known in the art. One such method entails measuring the rates of antigen-binding site/antigen complex formation and dissociation, wherein those rates depend on the concentrations of the complex partners, the affinity of the interaction, and geometric parameters that equally influence the rate in both directions. Thus, both the “on rate constant” (K_{on}) and the “off rate constant” (K_{off}) can be determined by calculation

of the concentrations and the actual rates of association and dissociation. (See Nature 361:186-87 (1993)). The ratio of $K_{\text{off}}/K_{\text{on}}$ enables the cancellation of all parameters not related to affinity, and is equal to the dissociation constant K_d . (See, generally, Davies et al. (1990) Annual Rev Biochem 59:439-473). An antibody of the present disclosure is said to specifically bind to the target, when the binding constant (K_d) is $\leq 1 \mu\text{M}$, in some embodiments $\leq 100 \text{ nM}$, in some embodiments $\leq 10 \text{ nM}$, and in some embodiments $\leq 100 \text{ pM}$ to about 1 pM , as measured by assays such as radioligand binding assays or similar assays known to those skilled in the art.

[000436] The term “isolated polynucleotide” as used herein shall mean a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, which by virtue of its origin the “isolated polynucleotide” (1) is not associated with all or a portion of a polynucleotide in which the “isolated polynucleotide” is found in nature, (2) is operably linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence. Polynucleotides in accordance with the disclosure include the nucleic acid molecules encoding the heavy chain immunoglobulin molecules shown herein, and nucleic acid molecules encoding the light chain immunoglobulin molecules shown herein.

[000437] The term “isolated protein” referred to herein means a protein of cDNA, recombinant RNA, or synthetic origin or some combination thereof, which by virtue of its origin, or source of derivation, the “isolated protein” (1) is not associated with proteins found in nature, (2) is free of other proteins from the same source, *e.g.*, free of murine proteins, (3) is expressed by a cell from a different species, or (4) does not occur in nature.

[000438] The term “polypeptide” is used herein as a generic term to refer to native protein, fragments, or analogs of a polypeptide sequence. Hence, native protein fragments, and analogs are species of the polypeptide genus. Polypeptides in accordance with the disclosure comprise the heavy chain immunoglobulin molecules shown herein, and the light chain immunoglobulin molecules shown herein, as well as antibody molecules formed by combinations comprising the heavy chain immunoglobulin molecules with light chain immunoglobulin molecules, such as kappa light chain immunoglobulin molecules, and vice versa, as well as fragments and analogs thereof.

[000439] The term “naturally-occurring” as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be

isolated from a source in nature and that has not been intentionally modified by man in the laboratory or otherwise is naturally-occurring.

[000440] The term “operably linked” as used herein refers to positions of components so described are in a relationship permitting them to function in their intended manner. A control sequence “operably linked” to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

[000441] The term “control sequence” as used herein refers to polynucleotide sequences that are necessary to effect the expression and processing of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and transcription termination sequence in eukaryotes, generally, such control sequences include promoters and transcription termination sequence. The term “control sequences” is intended to include, at a minimum, all components whose presence is essential for expression and processing, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences. The term “polynucleotide” as referred to herein means nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA.

[000442] The term oligonucleotide referred to herein includes naturally occurring, and modified nucleotides linked together by naturally occurring, and non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide subset generally comprising a length of 200 bases or fewer. In some embodiments, oligonucleotides are 10 to 60 bases in length and in some embodiments, 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides are usually single stranded, *e.g.*, for probes, although oligonucleotides may be double stranded, *e.g.*, for use in the construction of a gene mutant. Oligonucleotides of the disclosure are either sense or antisense oligonucleotides.

[000443] The term “naturally occurring nucleotides” referred to herein includes deoxyribonucleotides and ribonucleotides. The term “modified nucleotides” referred to herein includes nucleotides with modified or substituted sugar groups and the like. The term “oligonucleotide linkages” referred to herein includes oligonucleotide linkages such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoronmidate, and the like. *See e.g.*,

LaPlanche *et al.* Nucl. Acids Res. 14:9081 (1986); Stec *et al.* J. Am. Chem. Soc. 106:6077 (1984), Stein *et al.* Nucl. Acids Res. 16:3209 (1988), Zon *et al.* Anti Cancer Drug Design 6:539 (1991); Zon *et al.* Oligonucleotides and Analogues: A Practical Approach, pp. 87-108 (F. Eckstein, Ed., Oxford University Press, Oxford England (1991)); Stec *et al.* U.S. Patent No. 5,151,510; Uhlmann and Peyman Chemical Reviews 90:543 (1990). An oligonucleotide can include a label for detection, if desired.

[000444] As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. See Immunology - A Synthesis (2nd Edition, E.S. Golub and D.R. Green, Eds., Sinauer Associates, Sunderland, Mass. (1991)). Stereoisomers (*e.g.*, D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as α -, α -disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids may also be suitable components for polypeptides of the present disclosure. Examples of unconventional amino acids include: 4 hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, σ -N-methylarginine, and other similar amino acids and imino acids (*e.g.*, 4-hydroxyproline). In the polypeptide notation used herein, the left-hand direction is the amino terminal direction and the right-hand direction is the carboxy-terminal direction, in accordance with standard usage and convention.

[000445] Similarly, unless specified otherwise, the left-hand end of single-stranded polynucleotide sequences is the 5' end the left-hand direction of double-stranded polynucleotide sequences is referred to as the 5' direction. The direction of 5' to 3' addition of nascent RNA transcripts is referred to as the transcription direction sequence regions on the DNA strand having the same sequence as the RNA and that are 5' to the 5' end of the RNA transcript are referred to as "upstream sequences", sequence regions on the DNA strand having the same sequence as the RNA and that are 3' to the 3' end of the RNA transcript are referred to as "downstream sequences".

[000446] As applied to polypeptides, the term "substantial identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 80 percent sequence identity, in some embodiments, at least 90 percent sequence identity, in some embodiments, at least 95 percent sequence identity, and in some embodiments, at least 99 percent sequence identity.

[000447] In some embodiments, residue positions that are not identical differ by conservative amino acid substitutions.

[000448] As discussed herein, minor variations in the amino acid sequences of antibodies or immunoglobulin molecules are contemplated as being encompassed by the present disclosure, providing that the variations in the amino acid sequence maintain at least 75%, in some embodiments, at least 80%, 90%, 95%, and in some embodiments, 99%. In particular, conservative amino acid replacements are contemplated. Conservative replacements are those that take place within a family of amino acids that are related in their side chains. Genetically encoded amino acids are generally divided into families: (1) acidic amino acids are aspartate, glutamate; (2) basic amino acids are lysine, arginine, histidine; (3) non-polar amino acids are alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan, and (4) uncharged polar amino acids are glycine, asparagine, glutamine, cysteine, serine, threonine, tyrosine. The hydrophilic amino acids include arginine, asparagine, aspartate, glutamine, glutamate, histidine, lysine, serine, and threonine. The hydrophobic amino acids include alanine, cysteine, isoleucine, leucine, methionine, phenylalanine, proline, tryptophan, tyrosine and valine. Other families of amino acids include (i) serine and threonine, which are the aliphatic-hydroxy family; (ii) asparagine and glutamine, which are the amide containing family; (iii) alanine, valine, leucine and isoleucine, which are the aliphatic family; and (iv) phenylalanine, tryptophan, and tyrosine, which are the aromatic family. For example, it is reasonable to expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the binding or properties of the resulting molecule, especially if the replacement does not involve an amino acid within a framework site. Whether an amino acid change results in a functional peptide can readily be determined by assaying the specific activity of the polypeptide derivative. Assays are described in detail herein. Fragments or analogs of antibodies or immunoglobulin molecules can be readily prepared by those of ordinary skill in the art. Suitable amino- and carboxy-termini of fragments or analogs occur near boundaries of functional domains. Structural and functional domains can be identified by comparison of the nucleotide and/or amino acid sequence data to public or proprietary sequence databases. In some embodiments, computerized comparison methods are used to identify sequence motifs or predicted protein conformation domains that occur in other proteins of known structure and/or function. Methods to identify protein sequences that fold into a known three-dimensional structure are known. Bowie *et al.* Science 253:164 (1991). Thus, the foregoing examples demonstrate that those of skill in

the art can recognize sequence motifs and structural conformations that may be used to define structural and functional domains in accordance with the disclosure.

[000449] Suitable amino acid substitutions are those that: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinities, and (5) confer or modify other physicochemical or functional properties of such analogs. Analogs can include various muteins of a sequence other than the naturally-occurring peptide sequence. For example, single or multiple amino acid substitutions (for example, conservative amino acid substitutions) may be made in the naturally- occurring sequence (for example, in the portion of the polypeptide outside the domain(s) forming intermolecular contacts. A conservative amino acid substitution should not substantially change the structural characteristics of the parent sequence (*e.g.*, a replacement amino acid should not tend to break a helix that occurs in the parent sequence, or disrupt other types of secondary structure that characterizes the parent sequence). Examples of art-recognized polypeptide secondary and tertiary structures are described in *Proteins, Structures and Molecular Principles* (Creighton, Ed., W. H. Freeman and Company, New York (1984)); *Introduction to Protein Structure* (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)); and Thornton et al. *Nature* 354:105 (1991).

[000450] The term “polypeptide fragment” as used herein refers to a polypeptide that has an amino terminal and/or carboxy-terminal deletion and/or one or more internal deletion(s), but where the remaining amino acid sequence is identical to the corresponding positions in the naturally-occurring sequence deduced, for example, from a full length cDNA sequence. Fragments typically are at least 5, 6, 8 or 10 amino acids long, in some embodiments, at least 14 amino acids long, in some embodiments, at least 20 amino acids long, usually at least 50 amino acids long, and in some embodiments, at least 70 amino acids long. The term “analog” as used herein refers to polypeptides that are comprised of a segment of at least 25 amino acids that has substantial identity to a portion of a deduced amino acid sequence and that has specific binding to the target, under suitable binding conditions. Typically, polypeptide analogs comprise a conservative amino acid substitution (or addition or deletion) with respect to the naturally- occurring sequence. Analogs typically are at least 20 amino acids long, in some embodiments, at least 50 amino acids long or longer, and can often be as long as a full-length naturally-occurring polypeptide.

[000451] The term “agent” is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials.

[000452] As used herein, the terms “label” or “labeled” refers to incorporation of a detectable marker, *e.g.*, by incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotinyl moieties that can be detected by marked avidin (*e.g.*, streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or calorimetric methods). In certain situations, the label or marker can also be therapeutic. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (*e.g.*, ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I), fluorescent labels (*e.g.*, FITC, rhodamine, lanthanide phosphors), enzymatic labels (*e.g.*, horseradish peroxidase, p-galactosidase, luciferase, alkaline phosphatase), chemiluminescent, biotinyl groups, predetermined polypeptide epitopes recognized by a secondary reporter (*e.g.*, leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance. The term “pharmaceutical agent or drug” as used herein refers to a chemical compound or composition capable of inducing a desired therapeutic effect when properly administered to a patient.

[000453] Other chemistry terms herein are used according to conventional usage in the art, as exemplified by The McGraw-Hill Dictionary of Chemical Terms (Parker, S., Ed., McGraw-Hill, San Francisco (1985)).

[000454] As used herein, “substantially pure” means an object species is the predominant species present (*i.e.*, on a molar basis it is more abundant than any other individual species in the composition), and in some embodiments, a substantially purified fraction is a composition wherein the object species comprises at least about 50 percent (on a molar basis) of all macromolecular species present.

[000455] Generally, a substantially pure composition will comprise more than about 80 percent of all macromolecular species present in the composition, in some embodiments, more than about 85%, 90%, 95%, and 99%. In some embodiments, the object species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species.

[000456] The term patient includes human and veterinary subjects.

[000457] Antibodies and/or activatable antibodies of the disclosure specifically bind a given target, e.g., a human target protein such as human PDL1. Also included in the disclosure are antibodies and/or activatable antibodies that bind to the same epitope as the antibodies and/or activatable antibodies described herein. Also included in the disclosure are antibodies and/or antibodies activatable antibodies that compete with an anti-PDL1 antibody and/or an anti-PDL1 activatable antibody described herein for binding to PDL1, e.g., human PDL1. Also included in the disclosure are antibodies and/or antibodies activatable antibodies that cross-compete with an anti-PDL1 antibody and/or an anti-PDL1 activatable antibody described herein for binding to PDL1, e.g., human PDL1.

[000458] Those skilled in the art will recognize that it is possible to determine, without undue experimentation, if a monoclonal antibody (*e.g.*, a murine monoclonal or humanized antibody) has the same specificity as a monoclonal antibody used in the methods described herein by ascertaining whether the former prevents the latter from binding to the target. If the monoclonal antibody being tested competes with the monoclonal antibody of the disclosure, as shown by a decrease in binding by the monoclonal antibody of the disclosure, then the two monoclonal antibodies bind to the same, or a closely related, epitope. An alternative method for determining whether a monoclonal antibody has the specificity of a monoclonal antibody of the disclosure is to pre-incubate the monoclonal antibody of the disclosure with the target and then add the monoclonal antibody being tested to determine if the monoclonal antibody being tested is inhibited in its ability to bind the target. If the monoclonal antibody being tested is inhibited then, in all likelihood, it has the same, or functionally equivalent, epitopic specificity as the monoclonal antibody of the disclosure.

Multispecific Activatable Antibodies

[000459] The disclosure also provides multispecific anti-PDL1 activatable antibodies. The multispecific activatable antibodies provided herein are multispecific antibodies that recognize PDL1 and at least one or more different antigens or epitopes and that include at least one masking moiety (MM) linked to at least one antigen- or epitope-binding domain of the multispecific antibody such that coupling of the MM reduces the ability of the antigen- or epitope-binding domain to bind its target. In some embodiments, the MM is coupled to the antigen- or epitope-binding domain of the multispecific antibody via a cleavable moiety (CM) that functions as a substrate for at least one protease. The activatable multispecific

antibodies provided herein are stable in circulation, activated at intended sites of therapy and/or diagnosis but not in normal, *i.e.*, healthy tissue, and, when activated, exhibit binding to a target that is at least comparable to the corresponding, unmodified multispecific antibody.

[000460] In some embodiments, the multispecific activatable antibodies are designed to engage immune effector cells, also referred to herein as immune-effector cell engaging multispecific activatable antibodies. In some embodiments, the multispecific activatable antibodies are designed to engage leukocytes, also referred to herein as leukocyte engaging multispecific activatable antibodies. In some embodiments, the multispecific activatable antibodies are designed to engage T cells, also referred to herein as T-cell engaging multispecific activatable antibodies. In some embodiments, the multispecific activatable antibodies engage a surface antigen on a leukocyte, such as on a T cell, on a natural killer (NK) cell, on a myeloid mononuclear cell, on a macrophage, and/or on another immune effector cell. In some embodiments, the immune effector cell is a leukocyte. In some embodiments, the immune effector cell is a T cell. In some embodiments, the immune effector cell is a NK cell. In some embodiments, the immune effector cell is a mononuclear cell, such as a myeloid mononuclear cell. In some embodiments, the multispecific activatable antibodies are designed to bind or otherwise interact with more than one target and/or more than one epitope, also referred to herein as multi-antigen targeting activatable antibodies. As used herein, the terms “target” and “antigen” are used interchangeably.

[000461] In some embodiments, immune effector cell engaging multispecific activatable antibodies of the disclosure include a targeting antibody or antigen-binding fragment thereof that binds PDL1 and an immune effector cell engaging antibody or antigen-binding portion thereof, where at least one of the targeting antibody or antigen-binding fragment thereof and/or the immune effector cell engaging antibody or antigen-binding portion thereof is masked. In some embodiments, the immune effector cell engaging antibody or antigen binding fragment thereof includes a first antibody or antigen-binding fragment thereof (AB1) that binds a first, immune effector cell engaging target, where the AB1 is attached to a masking moiety (MM1) such that coupling of the MM1 reduces the ability of the AB1 to bind the first target. In some embodiments, the targeting antibody or antigen-binding fragment thereof includes a second antibody or fragment thereof that includes a second antibody or antigen-binding fragment thereof (AB2) that binds PDL1, where the AB2 is attached to a masking moiety (MM2) such that coupling of

the MM2 reduces the ability of the AB2 to bind PDL1. In some embodiments, the immune effector cell engaging antibody or antigen binding fragment thereof includes a first antibody or antigen-binding fragment thereof (AB1) that binds a first, immune effector cell engaging target, where the AB1 is attached to a masking moiety (MM1) such that coupling of the MM1 reduces the ability of the AB1 to bind the first target, and the targeting antibody or antigen-binding fragment thereof includes a second antibody or fragment thereof that includes a second antibody or antigen-binding fragment thereof (AB2) that binds PDL1, where the AB2 is attached to a masking moiety (MM2) such that coupling of the MM2 reduces the ability of the AB2 to bind PDL1. In some embodiments, the non-immune effector cell engaging antibody is a cancer targeting antibody. In some embodiments the non-immune cell effector antibody is an IgG. In some embodiments the immune effector cell engaging antibody is a scFv. In some embodiments the PDL1-targeting antibody (e.g., non-immune cell effector antibody) is an IgG and the immune effector cell engaging antibody is a scFv. In some embodiments, the immune effector cell is a leukocyte. In some embodiments, the immune effector cell is a T cell. In some embodiments, the immune effector cell is a NK cell. In some embodiments, the immune effector cell is a myeloid mononuclear cell.

[000462] In some embodiments, T-cell engaging multispecific activatable antibodies of the disclosure include a PDL1-targeting antibody or antigen-binding fragment thereof and a T-cell engaging antibody or antigen-binding portion thereof, where at least one of the PDL1-targeting antibody or antigen-binding fragment thereof and/or the T-cell engaging antibody or antigen-binding portion thereof is masked. In some embodiments, the T-cell engaging antibody or antigen binding fragment thereof includes a first antibody or antigen-binding fragment thereof (AB1) that binds a first, T-cell engaging target, where the AB1 is attached to a masking moiety (MM1) such that coupling of the MM1 reduces the ability of the AB1 to bind the first target. In some embodiments, the targeting antibody or antigen-binding fragment thereof includes a second antibody or fragment thereof that includes a second antibody or antigen-binding fragment thereof (AB2) that binds PDL1, where the AB2 is attached to a masking moiety (MM2) such that coupling of the MM2 reduces the ability of the AB2 to bind PDL1. In some embodiments, the T-cell engaging antibody or antigen binding fragment thereof includes a first antibody or antigen-binding fragment thereof (AB1) that binds a first, T-cell engaging target, where the AB1 is attached to a masking moiety (MM1) such that coupling of the MM1 reduces the ability of the AB1 to

bind the first target, and the targeting antibody or antigen-binding fragment thereof includes a second antibody or fragment thereof that includes a second antibody or antigen-binding fragment thereof (AB2) that binds PDL1, where the AB2 is attached to a masking moiety (MM2) such that coupling of the MM2 reduces the ability of the AB2 to bind PDL1.

[000463] In some embodiments of an immune effector cell engaging multispecific activatable antibody, one antigen is PDL1, and another antigen is typically a stimulatory or inhibitory receptor present on the surface of a T-cell, natural killer (NK) cell, myeloid mononuclear cell, macrophage, and/or other immune effector cell, such as, but not limited to, B7-H4, BTLA, CD3, CD4, CD8, CD16a, CD25, CD27, CD28, CD32, CD56, CD137, CTLA-4, GITR, HVEM, ICOS, LAG3, NKG2D, OX40, PD1, TIGIT, TIM3, or VISTA. In some embodiments, the antigen is a stimulatory receptor present on the surface of a T cell or NK cell; examples of such stimulatory receptors include, but are not limited to, CD3, CD27, CD28, CD137 (also referred to as 4-1BB), GITR, HVEM, ICOS, NKG2D, and OX40. In some embodiments, the antigen is an inhibitory receptor present on the surface of a T-cell; examples of such inhibitory receptors include, but are not limited to, BTLA, CTLA-4, LAG3, PD1, TIGIT, TIM3, and NK-expressed KIRs. The antibody domain conferring specificity to the T-cell surface antigen may also be substituted by a ligand or ligand domain that binds to a T-cell receptor, a NK-cell receptor, a macrophage receptor, and/or other immune effector cell receptor, such as, but not limited to, B7-1, B7-2, B7H3, PDL1, PDL2, or TNFSF9.

[000464] In some embodiments, the T-cell engaging multispecific activatable antibody includes an anti-CD3 epsilon (CD3 ϵ , also referred to herein as CD3e and CD3) scFv and a targeting antibody or antigen-binding fragment thereof, where at least one of the anti-CD3 ϵ scFv and/or the targeting antibody or antigen-binding portion thereof is masked. In some embodiments, the CD3 ϵ scFv includes a first antibody or antigen-binding fragment thereof (AB1) that binds CD3 ϵ , where the AB1 is attached to a masking moiety (MM1) such that coupling of the MM1 reduces the ability of the AB1 to bind CD3 ϵ . In some embodiments, the targeting antibody or antigen-binding fragment thereof includes a second antibody or fragment thereof that includes a second antibody or antigen-binding fragment thereof (AB2) that binds PDL1, where the AB2 is attached to a masking moiety (MM2) such that coupling of the MM2 reduces the ability of the AB2 to bind PDL1. In some embodiments, the CD3 ϵ scFv includes a first antibody or antigen-binding fragment thereof (AB1) that binds CD3 ϵ , where the AB1 is attached to a masking moiety (MM1) such that coupling of the MM1

reduces the ability of the AB1 to bind CD3 ϵ , and the targeting antibody or antigen-binding fragment thereof includes a second antibody or fragment thereof that includes a second antibody or antigen-binding fragment thereof (AB2) that binds PDL1, where the AB2 is attached to a masking moiety (MM2) such that coupling of the MM2 reduces the ability of the AB2 to bind PDL1.

[000465] In some embodiments, the multi-antigen targeting antibodies and/or multi-antigen targeting activatable antibodies include at least a first antibody or antigen-binding fragment thereof that binds a first target and/or first epitope and a second antibody or antigen-binding fragment thereof that binds a second target and/or a second epitope. In some embodiments, the multi-antigen targeting antibodies and/or multi-antigen targeting activatable antibodies bind two or more different targets. In some embodiments, the multi-antigen targeting antibodies and/or multi-antigen targeting activatable antibodies bind two or more different epitopes on the same target. In some embodiments, the multi-antigen targeting antibodies and/or multi-antigen targeting activatable antibodies bind a combination of two or more different targets and two or more different epitopes on the same target.

[000466] In some embodiments, a multispecific activatable antibody comprising an IgG has the IgG variable domains masked. In some embodiments, a multispecific activatable antibody comprising a scFv has the scFv domains masked. In some embodiments, a multispecific activatable antibody has both IgG variable domains and scFv domains, where at least one of the IgG variable domains is coupled to a masking moiety. In some embodiments, a multispecific activatable antibody has both IgG variable domains and scFv domains, where at least one of the scFv domains is coupled to a masking moiety. In some embodiments, a multispecific activatable antibody has both IgG variable domains and scFv domains, where at least one of the IgG variable domains is coupled to a masking moiety and at least one of the scFv domains is coupled to a masking moiety. In some embodiments, a multispecific activatable antibody has both IgG variable domains and scFv domains, where each of the IgG variable domains and the scFv domains is coupled to its own masking moiety. In some embodiments, one antibody domain of a multispecific activatable antibody has specificity for a target antigen and another antibody domain has specificity for a T-cell surface antigen. In some embodiments, one antibody domain of a multispecific activatable antibody has specificity for a target antigen and another antibody domain has specificity for another target antigen. In some embodiments, one antibody

domain of a multispecific activatable antibody has specificity for an epitope of a target antigen and another antibody domain has specificity for another epitope of the target antigen.

[000467] In a multispecific activatable antibody, a scFv can be fused to the carboxyl terminus of the heavy chain of an IgG activatable antibody, to the carboxyl terminus of the light chain of an IgG activatable antibody, or to the carboxyl termini of both the heavy and light chains of an IgG activatable antibody. In a multispecific activatable antibody, a scFv can be fused to the amino terminus of the heavy chain of an IgG activatable antibody, to the amino terminus of the light chain of an IgG activatable antibody, or to the amino termini of both the heavy and light chains of an IgG activatable antibody. In a multispecific activatable antibody, a scFv can be fused to any combination of one or more carboxyl termini and one or more amino termini of an IgG activatable antibody. In some embodiments, a masking moiety (MM) linked to a cleavable moiety (CM) is attached to and masks an antigen binding domain of the IgG. In some embodiments, a masking moiety (MM) linked to a cleavable moiety (CM) is attached to and masks an antigen binding domain of at least one scFv. In some embodiments, a masking moiety (MM) linked to a cleavable moiety (CM) is attached to and masks an antigen binding domain of an IgG and a masking moiety (MM) linked to a cleavable moiety (CM) is attached to and masks an antigen binding domain of at least one scFv.

[000468] The disclosure provides examples of multispecific activatable antibody structures which include, but are not limited to, the following: (VL-CL)₂:(VH-CH1-CH2-CH3-L4-VH*-L3-VL*-L2-CM-L1-MM)₂; (VL-CL)₂:(VH-CH1-CH2-CH3-L4-VL*-L3-VH*-L2-CM-L1-MM)₂; (MM-L1-CM-L2-VL-CL)₂:(VH-CH1-CH2-CH3-L4-VH*-L3-VL*)₂; (MM-L1-CM-L2-VL-CL)₂:(VH-CH1-CH2-CH3-L4-VL*-L3-VH*)₂; (VL-CL)₂:(MM-L1-CM-L2-VL*-L3-VH*-L4-VH-CH1-CH2-CH3)₂; (VL-CL)₂:(MM-L1-CM-L2-VH*-L3-VL*-L4-VH-CH1-CH2-CH3)₂; (MM-L1-CM-L2-VL-CL)₂:(VL*-L3-VH*-L4-VH-CH1-CH2-CH3)₂; (MM-L1-CM-L2-VL-CL)₂:(VH*-L3-VL*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VH*-L3-VL*-L2-CM-L1-MM)₂:(VH-CH1-CH2-CH3)₂; (VL-CL-L4-VL*-L3-VH*-L2-CM-L1-MM)₂:(VH-CH1-CH2-CH3)₂; (MM-L1-CM-L2-VL*-L3-VH*-L4-VL-CL)₂:(VH-CH1-CH2-CH3)₂; (MM-L1-CM-L2-VH*-L3-VL*-L4-VL-CL)₂:(VH-CH1-CH2-CH3)₂; (VL-CL-L4-VH*-L3-VL*-L2-CM-L1-MM)₂:(MM-L1-CM-L2-VL*-L3-VH*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VH*-L3-VL*-L2-CM-L1-MM)₂:(MM-L1-CM-L2-VH*-L3-VL*-L4-VH-CH1-CH2-CH3)₂; (VL-CL-L4-VL*-L3-VH*-L2-CM-

$L1-MM)_2$: $(MM-L1-CM-L2-VL^*-L3-VH^*-L4-VH-CH1-CH2-CH3)_2$; $(VL-CL-L4-VL^*-L3-VH^*-L2-CM-L1-MM)_2$: $(MM-L1-CM-L2-VH^*-L3-VL^*-L4-VH-CH1-CH2-CH3)_2$; $(VL-CL-L4-VH^*-L3-VL^*)_2$: $(MM-L1-CM-L2-VL^*-L3-VH^*-L4-VH-CH1-CH2-CH3)_2$; $(VL-CL-L4-VH^*-L3-VL^*)_2$: $(MM-L1-CM-L2-VH^*-L3-VL^*-L4-VH-CH1-CH2-CH3)_2$; $(VL-CL-L4-VL^*-L3-VH^*)_2$: $(MM-L1-CM-L2-VL^*-L3-VH^*-L4-VH-CH1-CH2-CH3)_2$; $(VL-CL-L4-VL^*-L3-VH^*)_2$: $(MM-L1-CM-L2-VH^*-L3-VL^*-L4-VH-CH1-CH2-CH3)_2$; $(VL-CL-L4-VH^*-L3-VL^*-L2-CM-L1-MM)_2$: $(VL^*-L3-VH^*-L4-VH-CH1-CH2-CH3)_2$; $(VL-CL-L4-VH^*-L3-VL^*-L2-CM-L1-MM)_2$: $(VH^*-L3-VL^*-L4-VH-CH1-CH2-CH3)_2$; $(VL-CL-L4-VL^*-L3-VH^*-L2-CM-L1-MM)_2$: $(VL^*-L3-VH^*-L4-VH-CH1-CH2-CH3)_2$; or $(VL-CL-L4-VL^*-L3-VH^*-L2-CM-L1-MM)_2$: $(VH^*-L3-VL^*-L4-VH-CH1-CH2-CH3)_2$,
 wherein: VL and VH represent the light and heavy variable domains of the first specificity, contained in the IgG; VL* and VH* represent the variable domains of the second specificity, contained in the scFv; L1 is a linker peptide connecting the masking moiety (MM) and the cleavable moiety (CM); L2 is a linker peptide connecting the cleavable moiety (CM), and the antibody; L3 is a linker peptide connecting the variable domains of the scFv; L4 is a linker peptide connecting the antibody of the first specificity to the antibody of the second specificity; CL is the light-chain constant domain; and CH1, CH2, CH3 are the heavy chain constant domains. The first and second specificities may be toward any antigen or epitope.

[000469] In some embodiments of a T-cell engaging multispecific activatable antibody, one antigen is PDL1, and another antigen is typically a stimulatory (also referred to herein as activating) or inhibitory receptor present on the surface of a T-cell, natural killer (NK) cell, myeloid mononuclear cell, macrophage, and/or other immune effector cell, such as, but not limited to, B7-H4, BTLA, CD3, CD4, CD8, CD16a, CD25, CD27, CD28, CD32, CD56, CD137 (also referred to as TNFRSF9), CTLA-4, GITR, HVEM, ICOS, LAG3, NKG2D, OX40, PD1, TIGIT, TIM3, or VISTA. The antibody domain conferring specificity to the T-cell surface antigen may also be substituted by a ligand or ligand domain that binds to a T-cell receptor, a NK-cell receptor, a macrophage receptor, and/or other immune effector cell receptor, such as, but not limited to, PDL1.

[000470] In some embodiments, the targeting antibody is an anti-PDL1 antibody disclosed herein. In some embodiments, the targeting antibody can be in the form an activatable antibody. In some embodiments, the scFv(s) can be in the form of a Pro-scFv (see, e.g., WO 2009/025846, WO 2010/081173).

[000471] In some embodiments, the scFv is specific for binding CD3 ϵ , and comprises or is derived from an antibody or fragment thereof that binds CD3 ϵ , e.g., CH2527, FN18, H2C, OKT3, 2C11, UCHT1, or V9. In some embodiments, the scFv is specific for binding CTLA-4 (also referred to herein as CTLA and CTLA4).

[000472] In some embodiments, the anti-CTLA-4 scFv includes the amino acid sequence:

GGSGGGSGSGGGSGGGSGGGGEIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQA
PRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYYCQQYGSSPLTFGGGTKVEIKRSG
GSTITSYNVYYTKLSSSGTQVQLVQTGGGVVQPGRSLRLSCAASGSTFSSYAMSWVRQAPGKGLEWV
SAISGSGGSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCATNSLYWYFDLWGRGTLVT
VSSAS (SEQ ID NO: 424)

[000473] In some embodiments, the anti-CTLA-4 scFv includes the amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence of SEQ ID NO: 424.

[000474] In some embodiments, the anti-CD3 ϵ scFv includes the amino acid sequence:

GGSGGGSGSGGGSGGGSGGGGQVQLQQSGAELARPGASVKMSCKASGYTFTRYTMHWVKQRPGQG
LEWIGYINPSRGYTNYNQKFKDKATLTDDKSSSTAYMQLSSLTSEDSAVYYCARYYDDHYCLDYWGQ
GTTLTVSSGGGGSGGGSGGGGSQIVLTQSPAISASPGKVTMTCSASSSVSYMNWYQQKSGTSPK
RWIYDTSKLAGVPAHFRGSGSGTSYSLTISGMEAEDAATYYCQQWSSNPFTFGSGTKLEINR
(SEQ ID NO: 425)

[000475] In some embodiments, the anti-CD3 ϵ scFv includes the amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence of SEQ ID NO: 425.

[000476] In some embodiments, the scFv is specific for binding one or more T-cells, one or more NK-cells and/or one or more macrophages. In some embodiments, the scFv is specific for binding a target selected from the group consisting of B7-H4, BTLA, CD3, CD4, CD8, CD16a, CD25, CD27, CD28, CD32, CD56, CD137, CTLA-4, GITR, HVEM, ICOS, LAG3, NKG2D, OX40, PD1, TIGIT, TIM3, or VISTA.

[000477] In some embodiments, the multispecific activatable antibody also includes an agent conjugated to the AB. In some embodiments, the agent is a therapeutic agent. In some embodiments, the agent is an antineoplastic agent. In some embodiments, the agent is a

toxin or fragment thereof. In some embodiments, the agent is conjugated to the multispecific activatable antibody via a linker. In some embodiments, the agent is conjugated to the AB via a cleavable linker. In some embodiments, the linker is a non-cleavable linker. In some embodiments, the agent is a microtubule inhibitor. In some embodiments, the agent is a nucleic acid damaging agent, such as a DNA alkylator or DNA intercalator, or other DNA damaging agent. In some embodiments, the linker is a cleavable linker. In some embodiments, the agent is an agent selected from the group listed in Table 11. In some embodiments, the agent is a dolastatin. In some embodiments, the agent is an auristatin or derivative thereof. In some embodiments, the agent is auristatin E or a derivative thereof. In some embodiments, the agent is monomethyl auristatin E (MMAE). In some embodiments, the agent is monomethyl auristatin D (MMAD). In some embodiments, the agent is a maytansinoid or maytansinoid derivative. In some embodiments, the agent is DM1 or DM4. In some embodiments, the agent is a duocarmycin or derivative thereof. In some embodiments, the agent is a calicheamicin or derivative thereof. In some embodiments, the agent is a pyrrolobenzodiazepine. In some embodiments, the agent is a pyrrolobenzodiazepine dimer.

[000478] In some embodiments, the multispecific activatable antibody also includes a detectable moiety. In some embodiments, the detectable moiety is a diagnostic agent.

[000479] In some embodiments, the multispecific activatable antibody naturally contains one or more disulfide bonds. In some embodiments, the multispecific activatable antibody can be engineered to include one or more disulfide bonds.

[000480] The disclosure also provides an isolated nucleic acid molecule encoding a multispecific activatable antibody described herein, as well as vectors that include these isolated nucleic acid sequences. The disclosure provides methods of producing a multispecific activatable antibody by culturing a cell under conditions that lead to expression of the activatable antibody, wherein the cell comprises such a nucleic acid molecule. In some embodiments, the cell comprises such a vector.

[000481] The disclosure also provides a method of manufacturing multispecific activatable antibodies of the disclosure by (a) culturing a cell comprising a nucleic acid construct that encodes the multispecific activatable antibody under conditions that lead to expression of the multispecific activatable, and (b) recovering the multispecific activatable antibody. Suitable AB, MM, and/or CM include any of the AB, MM, and/or CM disclosed herein.

[000482] The disclosure also provides multispecific activatable antibodies and/or multispecific activatable antibody compositions that include at least a first antibody or antigen-binding fragment thereof (AB1) that specifically binds a first target or first epitope and a second antibody or antigen-binding fragment thereof (AB2) that binds a second target or a second epitope, where at least AB1 is coupled or otherwise attached to a masking moiety (MM1), such that coupling of the MM1 reduces the ability of AB1 to bind its target. In some embodiments, the MM1 is coupled to AB1 via a first cleavable moiety (CM1) sequence that includes a substrate for a protease, for example, a protease that is co-localized with the target of AB1 at a treatment site or a diagnostic site in a subject. The multispecific activatable antibodies provided herein are stable in circulation, activated at intended sites of therapy and/or diagnosis but not in normal, *i.e.*, healthy tissue, and, when activated, exhibit binding to the target of AB1 that is at least comparable to the corresponding, unmodified multispecific antibody. Suitable AB, MM, and/or CM include any of the AB, MM, and/or CM disclosed herein.

[000483] The disclosure also provides compositions and methods that include a multispecific activatable antibody that includes at least a first antibody or antibody fragment (AB1) that specifically binds a target and a second antibody or antibody fragment (AB2), where at least the first AB in the multispecific activatable antibody is coupled to a masking moiety (MM1) that decreases the ability of AB1 to bind its target. In some embodiments, each AB is coupled to a MM that decreases the ability of its corresponding AB to each target. For example, in bispecific activatable antibody embodiments, AB1 is coupled to a first masking moiety (MM1) that decreases the ability of AB1 to bind its target, and AB2 is coupled to a second masking moiety (MM2) that decreases the ability of AB2 to bind its target. In some embodiments, the multispecific activatable antibody comprises more than two AB regions; in such embodiments, AB1 is coupled to a first masking moiety (MM1) that decreases the ability of AB1 to bind its target, AB2 is coupled to a second masking moiety (MM2) that decreases the ability of AB2 to bind its target, AB3 is coupled to a third masking moiety (MM3) that decreases the ability of AB3 to bind its target, and so on for each AB in the multispecific activatable antibody. Suitable AB, MM, and/or CM include any of the AB, MM, and/or CM disclosed herein.

[000484] In some embodiments, the multispecific activatable antibody further includes at least one cleavable moiety (CM) that is a substrate for a protease, where the CM links a MM to an AB. For example, in some embodiments, the multispecific activatable antibody

includes at least a first antibody or antibody fragment (AB1) that specifically binds a target and a second antibody or antibody fragment (AB2), where at least the first AB in the multispecific activatable antibody is coupled via a first cleavable moiety (CM1) to a masking moiety (MM1) that decreases the ability of AB1 to bind its target. In some bispecific activatable antibody embodiments, AB1 is coupled via CM1 to MM1, and AB2 is coupled via a second cleavable moiety (CM2) to a second masking moiety (MM2) that decreases the ability of AB2 to bind its target. In some embodiments, the multispecific activatable antibody comprises more than two AB regions; in some of these embodiments, AB1 is coupled via CM1 to MM1, AB2 is coupled via CM2 to MM2, and AB3 is coupled via a third cleavable moiety (CM3) to a third masking moiety (MM3) that decreases the ability of AB3 to bind its target, and so on for each AB in the multispecific activatable antibody. Suitable AB, MM, and/or CM include any of the AB, MM, and/or CM disclosed herein.

Activatable antibodies Having Non-Binding Steric Moieties or Binding Partners for Non-Binding Steric Moieties

[000485] The disclosure also provides activatable antibodies that include non-binding steric moieties (NB) or binding partners (BP) for non-binding steric moieties, where the BP recruits or otherwise attracts the NB to the activatable antibody. The activatable antibodies provided herein include, for example, an activatable antibody that includes a non-binding steric moiety (NB), a cleavable linker (CL) and antibody or antibody fragment (AB) that binds a target; an activatable antibody that includes a binding partner for a non-binding steric moiety (BP), a CL and an AB; and an activatable antibody that includes a BP to which an NB has been recruited, a CL and an AB that binds the target. Activatable antibodies in which the NB is covalently linked to the CL and AB of the activatable antibody or is associated by interaction with a BP that is covalently linked to the CL and AB of the activatable antibody are referred to herein as “NB-containing activatable antibodies.” By activatable or switchable is meant that the activatable antibody exhibits a first level of binding to a target when the activatable antibody is in an inhibited, masked or uncleaved state (*i.e.*, a first conformation), and a second level of binding to the target when the activatable antibody is in an uninhibited, unmasked and/or cleaved state (*i.e.*, a second conformation, *i.e.*, activated antibody), where the second level of target binding is greater than the first level of target binding. The activatable antibody compositions can exhibit

increased bioavailability and more favorable biodistribution compared to conventional antibody therapeutics.

[000486] In some embodiments, activatable antibodies provide for reduced toxicity and/or adverse side effects that could otherwise result from binding of the at non-treatment sites and/or non-diagnostic sites if the AB were not masked or otherwise inhibited from binding to such a site.

[000487] Anti-PDL1 activatable antibodies that include a non-binding steric moiety (NB) can be made using the methods set forth in PCT Publication No. WO 2013/192546, the contents of which are hereby incorporated by reference in their entirety.

Use Of Antibodies, Conjugated Antibodies, Activatable Antibodies, and Conjugated Activatable Antibodies

[000488] It will be appreciated that administration of therapeutic entities in accordance with the disclosure will be administered with suitable carriers, excipients, and other agents that are incorporated into formulations to provide improved transfer, delivery, tolerance, and the like. A multitude of appropriate formulations can be found in the formulary known to all pharmaceutical chemists: Remington's Pharmaceutical Sciences (15th ed, Mack Publishing Company, Easton, PA (1975)), particularly Chapter 87 by Blaug, Seymour, therein. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles (such as Lipofectin™), DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax. Any of the foregoing mixtures may be appropriate in treatments and therapies in accordance with the present disclosure, provided that the active ingredient in the formulation is not inactivated by the formulation and the formulation is physiologically compatible and tolerable with the route of administration. See also Baldrick P. "Pharmaceutical excipient development: the need for preclinical guidance." Regul. Toxicol Pharmacol. 32(2):210-8 (2000), Wang W. "Lyophilization and development of solid protein pharmaceuticals." Int. J. Pharm. 203(1-2):1-60 (2000), Charman WN "Lipids, lipophilic drugs, and oral drug delivery-some emerging concepts." J Pharm Sci.89(8):967-78 (2000), Powell *et al.* "Compendium of excipients for parenteral formulations" PDA J Pharm Sci Technol. 52:238-311 (1998) and the citations therein for additional information related to formulations, excipients and carriers well known to pharmaceutical chemists.

[000489] Therapeutic formulations of the disclosure, which include an anti-PDL1 antibody and/or activatable anti-PDL1 antibody, such as by way of non-limiting example, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody, are used to prevent, treat or otherwise ameliorate a disease or disorder associated with aberrant target expression and/or activity. For example, therapeutic formulations of the disclosure, which include an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody, are used to treat or otherwise ameliorate a cancer or other neoplastic condition, inflammation, an inflammatory disorder, and/or an autoimmune disease. In some embodiments, the cancer is a solid tumor or a hematologic malignancy where the target is expressed. In some embodiments, the cancer is a solid tumor where the target is expressed. In some embodiments, the cancer is a hematologic malignancy where the target is expressed. In some embodiments, the target is expressed on parenchyma (e.g., in cancer, the portion of an organ or tissue that often carries out function(s) of the organ or tissue). In some embodiments, the target is expressed on a cell, tissue, or organ. In some embodiments, the target is expressed on stroma (i.e., the connective supportive framework of a cell, tissue, or organ). In some embodiments, the target is expressed on an osteoblast. In some embodiments, the target is expressed on the endothelium (vasculature). In some embodiments, the target is expressed on a cancer stem cell. In some embodiments, the agent to which the antibody and/or the activatable antibody is conjugated is a microtubule inhibitor. In some embodiments, the agent to which the antibody and/or the activatable antibody is conjugated is a nucleic acid damaging agent.

[000490] Efficaciousness of prevention, amelioration or treatment is determined in association with any known method for diagnosing or treating the disease or disorder associated with target expression and/or activity, such as, for example, aberrant target expression and/or activity. Prolonging the survival of a subject or otherwise delaying the progression of the disease or disorder associated with target expression and/or activity, e.g., aberrant target expression and/or activity, in a subject indicates that the antibody, conjugated antibody, activatable antibody and/or conjugated activatable antibody confers a clinical benefit.

[000491] An antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody can be administered in the form of pharmaceutical compositions. Principles and considerations involved in preparing such compositions, as well as guidance in the choice of components are provided, for example, in Remington :

The Science And Practice Of Pharmacy 19th ed. (Alfonso R. Gennaro, et al., editors) Mack Pub. Co., Easton, Pa.: 1995; Drug Absorption Enhancement: Concepts, Possibilities, Limitations, And Trends, Harwood Academic Publishers, Langhorne, Pa., 1994; and Peptide And Protein Drug Delivery (Advances In Parenteral Sciences, Vol. 4), 1991, M. Dekker, New York.

[000492] In some embodiments where antibody fragments are used, the smallest fragment that specifically binds to the binding domain of the target protein is selected. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. (See, e.g., Marasco et al., Proc. Natl. Acad. Sci. USA, 90: 7889-7893 (1993)). The formulation can also contain more than one active compounds as necessary for the particular indication being treated, for example, in some embodiments, those with complementary activities that do not adversely affect each other. In some embodiments, or in addition, the composition can comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[000493] The active ingredients can also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions.

[000494] The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

[000495] Sustained-release preparations can be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOTTM (injectable microspheres composed of lactic

acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods.

[000496] In some embodiments, the antibody, the conjugated antibody, activatable antibody and/or conjugated activatable antibody contains a detectable label. An intact antibody, or a fragment thereof (*e.g.*, Fab, scFv, or F(ab)₂) is used. The term “labeled”, with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently-labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently-labeled streptavidin. The term “biological sample” is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. Included within the usage of the term “biological sample”, therefore, is blood and a fraction or component of blood including blood serum, blood plasma, or lymph. That is, the detection method of the disclosure can be used to detect an analyte mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of an analyte mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of an analyte protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations, immunochemical staining, and immunofluorescence. *In vitro* techniques for detection of an analyte genomic DNA include Southern hybridizations. Procedures for conducting immunoassays are described, for example in “ELISA: Theory and Practice: Methods in Molecular Biology”, Vol. 42, J. R. Crowther (Ed.) Human Press, Totowa, NJ, 1995; “Immunoassay”, E. Diamandis and T. Christopoulos, Academic Press, Inc., San Diego, CA, 1996; and “Practice and Theory of Enzyme Immunoassays”, P. Tijssen, Elsevier Science Publishers, Amsterdam, 1985. Furthermore, *in vivo* techniques for detection of an analyte protein include introducing into a subject a labeled anti-analyte protein antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

[000497] The antibodies, conjugated antibodies, activatable antibodies and/or conjugated activatable antibodies of the disclosure are also useful in a variety of diagnostic

and prophylactic formulations. In one embodiment, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody is administered to patients that are at risk of developing one or more of the aforementioned disorders. A patient's or organ's predisposition to one or more of the aforementioned disorders can be determined using genotypic, serological or biochemical markers.

[000498] In some embodiments of the disclosure, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody is administered to human individuals diagnosed with a clinical indication associated with one or more of the aforementioned disorders. Upon diagnosis, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody is administered to mitigate or reverse the effects of the clinical indication.

[000499] An antibody, a conjugated antibody, an activatable antibody, and/or a conjugated activatable antibody of the disclosure is also useful in the detection of a target in patient samples and accordingly are useful as diagnostics. For example, the antibodies and/or activatable antibodies, and conjugated versions thereof, of the disclosure are used in *in vitro* assays, *e.g.*, ELISA, to detect target levels in a patient sample.

[000500] In one embodiment, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody of the disclosure is immobilized on a solid support (*e.g.*, the well(s) of a microtiter plate). The immobilized antibody, conjugated antibody, activatable antibody and/or conjugated activatable antibody serves as a capture antibody for any target that may be present in a test sample. Prior to contacting the immobilized antibody and/or activatable antibody, and/or conjugated versions thereof, with a patient sample, the solid support is rinsed and treated with a blocking agent such as milk protein or albumin to prevent nonspecific adsorption of the analyte.

[000501] Subsequently the wells are treated with a test sample suspected of containing the antigen, or with a solution containing a standard amount of the antigen. Such a sample is, *e.g.*, a serum sample from a subject suspected of having levels of circulating antigen considered to be diagnostic of a pathology. After rinsing away the test sample or standard, the solid support is treated with a second antibody that is detectably labeled. The labeled second antibody serves as a detecting antibody. The level of detectable label is measured, and the concentration of target antigen in the test sample is determined by comparison with a standard curve developed from the standard samples.

[000502] It will be appreciated that based on the results obtained using the antibodies and activatable antibodies of the disclosure, and conjugated versions thereof, in an *in vitro* diagnostic assay, it is possible to stage a disease in a subject based on expression levels of the target antigen. For a given disease, samples of blood are taken from subjects diagnosed as being at various stages in the progression of the disease, and/or at various points in the therapeutic treatment of the disease. Using a population of samples that provides statistically significant results for each stage of progression or therapy, a range of concentrations of the antigen that may be considered characteristic of each stage is designated.

[000503] An antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody can also be used in diagnostic and/or imaging methods. In some embodiments, such methods are *in vitro* methods. In some embodiments, such methods are *in vivo* methods. In some embodiments, such methods are *in situ* methods. In some embodiments, such methods are *ex vivo* methods. For example, activatable antibodies having an enzymatically cleavable CM can be used to detect the presence or absence of an enzyme that is capable of cleaving the CM. Such activatable antibodies can be used in diagnostics, which can include *in vivo* detection (*e.g.*, qualitative or quantitative) of enzyme activity (or, in some embodiments, an environment of increased reduction potential such as that which can provide for reduction of a disulfide bond) through measured accumulation of activated antibodies (*i.e.*, antibodies resulting from cleavage of an activatable antibody) in a given cell or tissue of a given host organism. Such accumulation of activated antibodies indicates not only that the tissue expresses enzymatic activity (or an increased reduction potential depending on the nature of the CM) but also that the tissue expresses target to which the activated antibody binds.

[000504] For example, the CM can be selected to be substrate for at least one protease found at the site of a tumor, at the site of a viral or bacterial infection at a biologically confined site (*e.g.*, such as in an abscess, in an organ, and the like), and the like. The AB can be one that binds a target antigen. Using methods as disclosed herein, or when appropriate, methods familiar to one skilled in the art, a detectable label (*e.g.*, a fluorescent label or radioactive label or radiotracer) can be conjugated to an AB or other region of an antibody and/or activatable antibody. Suitable detectable labels are discussed in the context of the above screening methods and additional specific examples are provided below. Using an AB specific to a protein or peptide of the disease state, along with at least one protease

whose activity is elevated in the disease tissue of interest, activatable antibodies will exhibit an increased rate of binding to disease tissue relative to tissues where the CM specific enzyme is not present at a detectable level or is present at a lower level than in disease tissue or is inactive (e.g., in zymogen form or in complex with an inhibitor). Since small proteins and peptides are rapidly cleared from the blood by the renal filtration system, and because the enzyme specific for the CM is not present at a detectable level (or is present at lower levels in non-disease tissues or is present in inactive conformation), accumulation of activated antibodies in the disease tissue is enhanced relative to non-disease tissues.

[000505] In another example, activatable antibodies can be used to detect the presence or absence of a cleaving agent in a sample. For example, where the activatable antibodies contain a CM susceptible to cleavage by an enzyme, the activatable antibodies can be used to detect (either qualitatively or quantitatively) the presence of an enzyme in the sample. In another example, where the activatable antibodies contain a CM susceptible to cleavage by reducing agent, the activatable antibodies can be used to detect (either qualitatively or quantitatively) the presence of reducing conditions in a sample. To facilitate analysis in these methods, the activatable antibodies can be detectably labeled, and can be bound to a support (*e.g.*, a solid support, such as a slide or bead). The detectable label can be positioned on a portion of the activatable antibody that is not released following cleavage, for example, the detectable label can be a quenched fluorescent label or other label that is not detectable until cleavage has occurred. The assay can be conducted by, for example, contacting the immobilized, detectably labeled activatable antibodies with a sample suspected of containing an enzyme and/or reducing agent for a time sufficient for cleavage to occur, then washing to remove excess sample and contaminants. The presence or absence of the cleaving agent (*e.g.*, enzyme or reducing agent) in the sample is then assessed by a change in detectable signal of the activatable antibodies prior to contacting with the sample *e.g.*, the presence of and/or an increase in detectable signal due to cleavage of the activatable antibody by the cleaving agent in the sample.

[000506] Such detection methods can be adapted to also provide for detection of the presence or absence of a target that is capable of binding the AB of the activatable antibodies when cleaved. Thus, the assays can be adapted to assess the presence or absence of a cleaving agent and the presence or absence of a target of interest. The presence or absence of the cleaving agent can be detected by the presence of and/or an increase in detectable label of the activatable antibodies as described above, and the presence or

absence of the target can be detected by detection of a target-AB complex *e.g.*, by use of a detectably labeled anti-target antibody.

[000507] Activatable antibodies are also useful in *in situ* imaging for the validation of activatable antibody activation, *e.g.*, by protease cleavage, and binding to a particular target. *In situ* imaging is a technique that enables localization of proteolytic activity and target in biological samples such as cell cultures or tissue sections. Using this technique, it is possible to confirm both binding to a given target and proteolytic activity based on the presence of a detectable label (*e.g.*, a fluorescent label).

[000508] These techniques are useful with any frozen cells or tissue derived from a disease site (*e.g.* tumor tissue) or healthy tissues. These techniques are also useful with fresh cell or tissue samples.

[000509] In these techniques, an activatable antibody is labeled with a detectable label. The detectable label may be a fluorescent dye, (*e.g.* a fluorophore, Fluorescein Isothiocyanate (FITC), Rhodamine Isothiocyanate (TRITC), an Alexa Fluor® label), a near infrared (NIR) dye (*e.g.*, Qdot® nanocrystals), a colloidal metal, a hapten, a radioactive marker, biotin and an amplification reagent such as streptavidin, or an enzyme (*e.g.* horseradish peroxidase or alkaline phosphatase).

[000510] Detection of the label in a sample that has been incubated with the labeled, activatable antibody indicates that the sample contains the target and contains a protease that is specific for the CM of the activatable antibody. In some embodiments, the presence of the protease can be confirmed using broad spectrum protease inhibitors such as those described herein, and/or by using an agent that is specific for the protease, for example, an antibody such as A11, which is specific for the protease matriptase and inhibits the proteolytic activity of matriptase; see *e.g.*, International Publication Number WO 2010/129609, published 11 November 2010. The same approach of using broad spectrum protease inhibitors such as those described herein, and/or by using a more selective inhibitory agent can be used to identify a protease that is specific for the CM of the activatable antibody. In some embodiments, the presence of the target can be confirmed using an agent that is specific for the target, *e.g.*, another antibody, or the detectable label can be competed with unlabeled target. In some embodiments, unlabeled activatable antibody could be used, with detection by a labeled secondary antibody or more complex detection system.

[000511] Similar techniques are also useful for *in vivo* imaging where detection of the fluorescent signal in a subject, *e.g.*, a mammal, including a human, indicates that the disease site contains the target and contains a protease that is specific for the CM of the activatable antibody.

[000512] These techniques are also useful in kits and/or as reagents for the detection, identification or characterization of protease activity in a variety of cells, tissues, and organisms based on the protease-specific CM in the activatable antibody.

[000513] The disclosure provides methods of using the antibodies and/or activatable antibodies in a variety of diagnostic and/or prophylactic indications. For example, the disclosure provides methods of detecting presence or absence of a cleaving agent and a target of interest in a subject or a sample by (i) contacting a subject or sample with an activatable antibody, wherein the activatable antibody comprises a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, *e.g.*, a protease, and an antigen binding domain or fragment thereof (AB) that specifically binds the target of interest, wherein the activatable antibody in an uncleaved, non-activated state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB and is not a modified form of a natural binding partner of the AB; and (b) wherein, in an uncleaved, non-activated state, the MM interferes with specific binding of the AB to the target, and in a cleaved, activated state the MM does not interfere or compete with specific binding of the AB to the target; and (ii) measuring a level of activated activatable antibody in the subject or sample, wherein a detectable level of activated activatable antibody in the subject or sample indicates that the cleaving agent and the target are present in the subject or sample and wherein no detectable level of activated activatable antibody in the subject or sample indicates that the cleaving agent, the target or both the cleaving agent and the target are absent and/or not sufficiently present in the subject or sample. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent

comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000514] The disclosure also provides methods of detecting presence or absence of a cleaving agent in a subject or a sample by (i) contacting a subject or sample with an activatable antibody in the presence of a target of interest, *e.g.*, the target, wherein the activatable antibody comprises a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, *e.g.*, a protease, and an antigen binding domain or fragment thereof (AB) that specifically binds the target of interest, wherein the activatable antibody in an uncleaved, non-activated state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB and is not a modified form of a natural binding partner of the AB; and (b) wherein, in an uncleaved, non-activated state, the MM interferes with specific binding of the AB to the target, and in a cleaved, activated state the MM does not interfere or compete with specific binding of the AB to the target; and (ii) measuring a level of activated activatable antibody in the subject or sample, wherein a detectable level of activated activatable antibody in the subject or sample indicates that the cleaving agent is present in the subject or sample and wherein no detectable level of activated activatable antibody in the subject or sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or sample. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000515] The disclosure also provides kits for use in methods of detecting presence or absence of a cleaving agent and the target in a subject or a sample, where the kits include at least an activatable antibody comprises a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, *e.g.*, a protease, and an antigen binding domain or fragment thereof (AB) that specifically binds the target of interest, wherein the activatable

antibody in an uncleaved, non-activated state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB and is not a modified form of a natural binding partner of the AB; and (b) wherein, in an uncleaved, non-activated state, the MM interferes with specific binding of the AB to the target, and in a cleaved, activated state the MM does not interfere or compete with specific binding of the AB to the target; and (ii) measuring a level of activated activatable antibody in the subject or sample, wherein a detectable level of activated activatable antibody in the subject or sample indicates that the cleaving agent is present in the subject or sample and wherein no detectable level of activated activatable antibody in the subject or sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or sample. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000516] The disclosure also provides methods of detecting presence or absence of a cleaving agent in a subject or a sample by (i) contacting a subject or sample with an activatable antibody, wherein the activatable antibody comprises a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, e.g., a protease, an antigen binding domain (AB) that specifically binds the target, and a detectable label, wherein the activatable antibody in an uncleaved, non-activated state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB and is not a modified form of a natural binding partner of the AB; wherein, in an uncleaved, non-activated state, the MM interferes with specific binding of the AB to the target, and in a cleaved, activated state the MM does not interfere or compete with specific binding of the AB to the target; and wherein the detectable label is positioned on a portion of the

activatable antibody that is released following cleavage of the CM; and (ii) measuring a level of detectable label in the subject or sample, wherein a detectable level of the detectable label in the subject or sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or sample and wherein no detectable level of the detectable label in the subject or sample indicates that the cleaving agent is present in the subject or sample. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000517] The disclosure also provides kits for use in methods of detecting presence or absence of a cleaving agent and the target in a subject or a sample, where the kits include at least an activatable antibody and/or conjugated activatable antibody (e.g., an activatable antibody to which a therapeutic agent is conjugated) described herein for use in contacting a subject or biological sample and means for detecting the level of activated activatable antibody and/or conjugated activatable antibody in the subject or biological sample, wherein a detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent and the target are present in the subject or biological sample and wherein no detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent, the target or both the cleaving agent and the target are absent and/or not sufficiently present in the subject or biological sample, such that the target binding and/or protease cleavage of the activatable antibody cannot be detected in the subject or biological sample.

[000518] The disclosure also provides methods of detecting presence or absence of a cleaving agent in a subject or a sample by (i) contacting a subject or biological sample with an activatable antibody in the presence of the target, and (ii) measuring a level of activated activatable antibody in the subject or biological sample, wherein a detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent is present in the subject or biological sample and wherein no detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving

agent is absent and/or not sufficiently present in the subject or biological sample at a detectable level, such that protease cleavage of the activatable antibody cannot be detected in the subject or biological sample. Such an activatable antibody includes a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, e.g., a protease, and an antigen binding domain or fragment thereof (AB) that specifically binds the target, wherein the activatable antibody in an uncleaved (*i.e.*, non-activated) state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB; and (b) wherein the MM of the activatable antibody in an uncleaved state interferes with specific binding of the AB to the target, and wherein the MM of an activatable antibody in a cleaved (*i.e.*, activated) state does not interfere or compete with specific binding of the AB to the target. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the detectable label is attached to the masking moiety. In some embodiments, the detectable label is attached to the cleavable moiety N-terminal to the protease cleavage site. In some embodiments, a single antigen binding site of the AB is masked. In some embodiments wherein an antibody of the disclosure has at least two antigen binding sites, at least one antigen binding site is masked and at least one antigen binding site is not masked. In some embodiments all antigen binding sites are masked. In some embodiments, the measuring step includes use of a secondary reagent comprising a detectable label.

[000519] The disclosure also provides kits for use in methods of detecting presence or absence of a cleaving agent and the target in a subject or a sample, where the kits include at least an activatable antibody and/or conjugated activatable antibody described herein for use in contacting a subject or biological sample with an activatable antibody in the presence of the target, and measuring a level of activated activatable antibody in the subject or biological sample, wherein a detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent is present in the subject or biological sample and wherein no detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or biological sample at a detectable level, such that protease cleavage of the activatable antibody cannot be detected in the subject or biological sample. Such an

activatable antibody includes a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, e.g., a protease, and an antigen binding domain or fragment thereof (AB) that specifically binds the target, wherein the activatable antibody in an uncleaved (*i.e.*, non-activated) state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB; and (b) wherein the MM of the activatable antibody in an uncleaved state interferes with specific binding of the AB to the target, and wherein the MM of an activatable antibody in a cleaved (*i.e.*, activated) state does not interfere or compete with specific binding of the AB to the target. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the detectable label is attached to the masking moiety. In some embodiments, the detectable label is attached to the cleavable moiety N-terminal to the protease cleavage site. In some embodiments, a single antigen binding site of the AB is masked. In some embodiments wherein an antibody of the disclosure has at least two antigen binding sites, at least one antigen binding site is masked and at least one antigen binding site is not masked. In some embodiments all antigen binding sites are masked. In some embodiments, the measuring step includes use of a secondary reagent comprising a detectable label.

[000520] The disclosure also provides kits for use in methods of detecting presence or absence of a cleaving agent in a subject or a sample, where the kits include at least an activatable antibody and/or conjugated activatable antibody described herein for use in contacting a subject or biological sample and means for detecting the level of activated activatable antibody and/or conjugated activatable antibody in the subject or biological sample, wherein the activatable antibody includes a detectable label that is positioned on a portion of the activatable antibody that is released following cleavage of the CM, wherein a detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or biological sample such that the target binding and/or protease cleavage of the activatable antibody cannot be detected in the subject or biological sample, and wherein no detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent is present in the subject or biological sample at a detectable level.

[000521] The disclosure provides methods of detecting presence or absence of a cleaving agent and the target in a subject or a sample by (i) contacting a subject or biological sample with an activatable antibody, wherein the activatable antibody includes a detectable label that is positioned on a portion of the activatable antibody that is released following cleavage of the CM and (ii) measuring a level of activated activatable antibody in the subject or biological sample, wherein a detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent, the target or both the cleaving agent and the target are absent and/or not sufficiently present in the subject or biological sample, such that the target binding and/or protease cleavage of the activatable antibody cannot be detected in the subject or biological sample, and wherein a reduced detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent and the target are present in the subject or biological sample. A reduced level of detectable label is, for example, a reduction of about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95% and/or about 100%. Such an activatable antibody includes a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, and an antigen binding domain or fragment thereof (AB) that specifically binds the target, wherein the activatable antibody in an uncleaved (*i.e.*, non-activated) state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB; and (b) wherein the MM of the activatable antibody in an uncleaved state interferes with specific binding of the AB to the target, and wherein the MM of an activatable antibody in a cleaved (*i.e.*, activated) state does not interfere or compete with specific binding of the AB to the target. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000522] The disclosure also provides kits for use in methods of detecting presence or absence of a cleaving agent and the target in a subject or a sample, where the kits include at least an activatable antibody and/or conjugated activatable antibody described herein for use in contacting a subject or biological sample and means for detecting the level of activated activatable antibody and/or conjugated activatable antibody in the subject or biological sample, wherein a detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent, the target or both the cleaving agent and the target are absent and/or not sufficiently present in the subject or biological sample, such that the target binding and/or protease cleavage of the activatable antibody cannot be detected in the subject or biological sample, and wherein a reduced detectable level of activated activatable antibody in the subject or biological sample indicates that the cleaving agent and the target are present in the subject or biological sample. A reduced level of detectable label is, for example, a reduction of about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95% and/or about 100%.

[000523] The disclosure also provides methods of detecting presence or absence of a cleaving agent in a subject or a sample by (i) contacting a subject or biological sample with an activatable antibody, wherein the activatable antibody includes a detectable label that is positioned on a portion of the activatable antibody that is released following cleavage of the CM; and (ii) measuring a level of detectable label in the subject or biological sample, wherein a detectable level of the detectable label in the subject or biological sample indicates that the cleaving agent is absent and/or not sufficiently present in the subject or biological sample at a detectable level, such that protease cleavage of the activatable antibody cannot be detected in the subject or biological sample, and wherein a reduced detectable level of the detectable label in the subject or biological sample indicates that the cleaving agent is present in the subject or biological sample. A reduced level of detectable label is, for example, a reduction of about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95% and/or about 100%. Such an activatable antibody includes a masking moiety (MM), a cleavable moiety (CM) that is cleaved by the cleaving agent, and an antigen binding domain or fragment thereof (AB) that specifically binds the target, wherein the activatable antibody in

an uncleaved (*i.e.*, non-activated) state comprises a structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; (a) wherein the MM is a peptide that inhibits binding of the AB to the target, and wherein the MM does not have an amino acid sequence of a naturally occurring binding partner of the AB; and (b) wherein the MM of the activatable antibody in an uncleaved state interferes with specific binding of the AB to the target, and wherein the MM of an activatable antibody in a cleaved (*i.e.*, activated) state does not interfere or compete with specific binding of the AB to the target. In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000524] The disclosure also provides kits for use in methods of detecting presence or absence of a cleaving agent of interest in a subject or a sample, where the kits include at least an activatable antibody and/or conjugated activatable antibody described herein for use in contacting a subject or biological sample and means for detecting the level of activated activatable antibody and/or conjugated activatable antibody in the subject or biological sample, wherein the activatable antibody includes a detectable label that is positioned on a portion of the activatable antibody that is released following cleavage of the CM, wherein a detectable level of the detectable label in the subject or biological sample indicates that the cleaving agent, the target, or both the cleaving agent and the target are absent and/or not sufficiently present in the subject or biological sample, such that the target binding and/or protease cleavage of the activatable antibody cannot be detected in the subject or biological sample, and wherein a reduced detectable level of the detectable label in the subject or biological sample indicates that the cleaving agent and the target are present in the subject or biological sample. A reduced level of detectable label is, for example, a reduction of about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95% and/or about 100%.

[000525] In some embodiments of these methods and kits, the activatable antibody includes a detectable label. In some embodiments of these methods and kits, the detectable label includes an imaging agent, a contrasting agent, an enzyme, a fluorescent label, a chromophore, a dye, one or more metal ions, or a ligand-based label. In some embodiments of these methods and kits, the imaging agent comprises a radioisotope. In some embodiments of these methods and kits, the radioisotope is indium or technetium. In some embodiments of these methods and kits, the contrasting agent comprises iodine, gadolinium or iron oxide. In some embodiments of these methods and kits, the enzyme comprises horseradish peroxidase, alkaline phosphatase, or β -galactosidase. In some embodiments of these methods and kits, the fluorescent label comprises yellow fluorescent protein (YFP), cyan fluorescent protein (CFP), green fluorescent protein (GFP), modified red fluorescent protein (mRFP), red fluorescent protein tdimer2 (RFP tdimer2), HCRED, or a europium derivative. In some embodiments of these methods and kits, the luminescent label comprises an N- methylacrydium derivative. In some embodiments of these methods, the label comprises an Alexa Fluor[®] label, such as Alex Fluor[®] 680 or Alexa Fluor[®] 750. In some embodiments of these methods and kits, the ligand-based label comprises biotin, avidin, streptavidin or one or more haptens.

[000526] In some embodiments of these methods and kits, the subject is a mammal. In some embodiments of these methods and kits, the subject is a human. In some embodiments, the subject is a non-human mammal, such as a non-human primate, companion animal (e.g., cat, dog, horse), farm animal, work animal, or zoo animal. In some embodiments, the subject is a rodent.

[000527] In some embodiments of these methods, the method is an *in vivo* method. In some embodiments of these methods, the method is an *in situ* method. In some embodiments of these methods, the method is an *ex vivo* method. In some embodiments of these methods, the method is an *in vitro* method.

[000528] In some embodiments, *in situ* imaging and/or *in vivo* imaging are useful in methods to identify which patients to treat. For example, in *in situ* imaging, the activatable antibodies are used to screen patient samples to identify those patients having the appropriate protease(s) and target(s) at the appropriate location, e.g., at a tumor site.

[000529] In some embodiments *in situ* imaging is used to identify or otherwise refine a patient population suitable for treatment with an activatable antibody of the disclosure. For example, patients that test positive for both the target (e.g., the target) and a protease that

cleaves the substrate in the cleavable moiety (CM) of the activatable antibody being tested (e.g., accumulate activated antibodies at the disease site) are identified as suitable candidates for treatment with such an activatable antibody comprising such a CM. Likewise, patients that test negative for either or both of the target (e.g., the target) and the protease that cleaves the substrate in the CM in the activatable antibody being tested using these methods might be identified as suitable candidates for another form of therapy. In some embodiments, such patients that test negative with respect to a first activatable antibody can be tested with other activatable antibodies comprising different CMs until a suitable activatable antibody for treatment is identified (e.g., an activatable antibody comprising a CM that is cleaved by the patient at the site of disease). In some embodiments, the patient is then administered a therapeutically effective amount of the activatable antibody for which the patient tested positive.

[000530] In some embodiments *in vivo* imaging is used to identify or otherwise refine a patient population suitable for treatment with an activatable antibody of the disclosure. For example, patients that test positive for both the target (e.g., the target) and a protease that cleaves the substrate in the cleavable moiety (CM) of the activatable antibody being tested (e.g., accumulate activated antibodies at the disease site) are identified as suitable candidates for treatment with such an activatable antibody comprising such a CM. Likewise, patients that test negative might be identified as suitable candidates for another form of therapy. In some embodiments, such patients that test negative with respect to a first activatable antibody can be tested with other activatable antibodies comprising different CMs until a suitable activatable antibody for treatment is identified (e.g., an activatable antibody comprising a CM that is cleaved by the patient at the site of disease). In some embodiments, the patient is then administered a therapeutically effective amount of the activatable antibody for which the patient tested positive.

[000531] In some embodiments of the methods and kits, the method or kit is used to identify or otherwise refine a patient population suitable for treatment with an activatable antibody of the disclosure. For example, patients that test positive for both the target (e.g., the target) and a protease that cleaves the substrate in the cleavable moiety (CM) of the activatable antibody being tested in these methods are identified as suitable candidates for treatment with such an activatable antibody comprising such a CM. Likewise, patients that test negative for both of the targets (e.g., the target) and the protease that cleaves the substrate in the CM in the activatable antibody being tested using these methods might be

identified as suitable candidates for another form of therapy. In some embodiments, such patients can be tested with other activatable antibodies until a suitable activatable antibody for treatment is identified (e.g., an activatable antibody comprising a CM that is cleaved by the patient at the site of disease). In some embodiments, patients that test negative for either of the target (e.g., the target) are identified as suitable candidates for treatment with such an activatable antibody comprising such a CM. In some embodiments, patients that test negative for either of the target (e.g., the target) are identified as not being suitable candidates for treatment with such an activatable antibody comprising such a CM. In some embodiments, such patients can be tested with other activatable antibodies until a suitable activatable antibody for treatment is identified (e.g., an activatable antibody comprising a CM that is cleaved by the patient at the site of disease). In some embodiments, the activatable antibody is an activatable antibody to which a therapeutic agent is conjugated. In some embodiments, the activatable antibody is not conjugated to an agent. In some embodiments, the activatable antibody comprises a detectable label. In some embodiments, the detectable label is positioned on the AB. In some embodiments, measuring the level of activatable antibody in the subject or sample is accomplished using a secondary reagent that specifically binds to the activated antibody, wherein the reagent comprises a detectable label. In some embodiments, the secondary reagent is an antibody comprising a detectable label.

[000532] In some embodiments, a method or kit is used to identify or otherwise refine a patient population suitable for treatment with an anti-the target activatable antibody and/or conjugated activatable antibody (e.g., activatable antibody to which a therapeutic agent is conjugated) of the disclosure, followed by treatment by administering that activatable antibody and/or conjugated activatable antibody to a subject in need thereof. For example, patients that test positive for both the targets (e.g., the target) and a protease that cleaves the substrate in the cleavable moiety (CM) of the activatable antibody and/or conjugated activatable antibody being tested in these methods are identified as suitable candidates for treatment with such antibody and/or such a conjugated activatable antibody comprising such a CM, and the patient is then administered a therapeutically effective amount of the activatable antibody and/or conjugated activatable antibody that was tested. Likewise, patients that test negative for either or both of the target (e.g., the target) and the protease that cleaves the substrate in the CM in the activatable antibody being tested using these methods might be identified as suitable candidates for another form of therapy. In some

embodiments, such patients can be tested with other antibody and/or conjugated activatable antibody until a suitable antibody and/or conjugated activatable antibody for treatment is identified (*e.g.*, an activatable antibody and/or conjugated activatable antibody comprising a CM that is cleaved by the patient at the site of disease). In some embodiments, the patient is then administered a therapeutically effective amount of the activatable antibody and/or conjugated activatable antibody for which the patient tested positive.

[000533] In some embodiments of these methods and kits, the MM is a peptide having a length from about 4 to 40 amino acids. In some embodiments of these methods and kits, the activatable antibody comprises a linker peptide, wherein the linker peptide is positioned between the MM and the CM. In some embodiments of these methods and kits, the activatable antibody comprises a linker peptide, where the linker peptide is positioned between the AB and the CM. In some embodiments of these methods and kits, the activatable antibody comprises a first linker peptide (L1) and a second linker peptide (L2), wherein the first linker peptide is positioned between the MM and the CM and the second linker peptide is positioned between the AB and the CM. In some embodiments of these methods and kits, each of L1 and L2 is a peptide of about 1 to 20 amino acids in length, and wherein each of L1 and L2 need not be the same linker. In some embodiments of these methods and kits, one or both of L1 and L2 comprises a glycine-serine polymer. In some embodiments of these methods and kits, at least one of L1 and L2 comprises an amino acid sequence selected from the group consisting of (GS)_n, (GSGGS)_n (SEQ ID NO: 191) and (GGGS)_n (SEQ ID NO: 192), where n is an integer of at least one. In some embodiments of these methods and kits, at least one of L1 and L2 comprises an amino acid sequence having the formula (GGS)_n, where n is an integer of at least one. In some embodiments of these methods and kits, at least one of L1 and L2 comprises an amino acid sequence selected from the group consisting of Gly-Gly-Ser-Gly (SEQ ID NO: 193), Gly-Gly-Ser-Gly-Gly (SEQ ID NO: 194), Gly-Ser-Gly-Ser-Gly (SEQ ID NO: 195), Gly-Ser-Gly-Gly-Gly (SEQ ID NO: 196), Gly-Gly-Gly-Ser-Gly (SEQ ID NO: 197), and Gly-Ser-Ser-Ser-Gly (SEQ ID NO: 198).

[000534] In some embodiments of these methods and kits, the AB comprises an antibody or antibody fragment sequence selected from the cross-reactive antibody sequences presented herein. In some embodiments of these methods and kits, the AB comprises a Fab fragment, a scFv or a single chain antibody (scAb).

[000535] In some embodiments of these methods and kits, the cleaving agent is a protease that is co-localized in the subject or sample with the target and the CM is a polypeptide that functions as a substrate for the protease, wherein the protease cleaves the CM in the activatable antibody when the activatable antibody is exposed to the protease. In some embodiments of these methods and kits, the CM is a polypeptide of up to 15 amino acids in length. In some embodiments of these methods and kits, the CM is coupled to the N-terminus of the AB. In some embodiments of these methods and kits, the CM is coupled to the C-terminus of the AB. In some embodiments of these methods and kits, the CM is coupled to the N-terminus of a VL chain of the AB.

[000536] The antibodies, conjugated antibodies, activatable antibodies and/or conjugated activatable antibodies of the disclosure are used in diagnostic and prophylactic formulations. In one embodiment, an activatable antibody is administered to patients that are at risk of developing one or more of the aforementioned inflammation, inflammatory disorders, cancer or other disorders.

[000537] A patient's or organ's predisposition to one or more of the aforementioned disorders can be determined using genotypic, serological or biochemical markers.

[000538] In some embodiments of the disclosure, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody is administered to human individuals diagnosed with a clinical indication associated with one or more of the aforementioned disorders. Upon diagnosis, an antibody, a conjugated antibody, an activatable antibody and/or a conjugated activatable antibody is administered to mitigate or reverse the effects of the clinical indication.

[000539] Antibodies, conjugated antibodies, activatable antibodies and/or conjugated activatable antibodies of the disclosure are also useful in the detection of the target in patient samples and accordingly are useful as diagnostics. For example, the antibodies, conjugated antibodies, the activatable antibodies and/or conjugated activatable antibodies of the disclosure are used in *in vitro* assays, *e.g.*, ELISA, to detect target levels in a patient sample.

[000540] In one embodiment, an antibody and/or activatable antibody of the disclosure is immobilized on a solid support (*e.g.*, the well(s) of a microtiter plate). The immobilized antibody and/or activatable antibody serves as a capture antibody for any target that may be present in a test sample. Prior to contacting the immobilized antibody and/or activatable

antibody with a patient sample, the solid support is rinsed and treated with a blocking agent such as milk protein or albumin to prevent nonspecific adsorption of the analyte.

[000541] Subsequently the wells are treated with a test sample suspected of containing the antigen, or with a solution containing a standard amount of the antigen. Such a sample is, *e.g.*, a serum sample from a subject suspected of having levels of circulating antigen considered to be diagnostic of a pathology. After rinsing away the test sample or standard, the solid support is treated with a second antibody that is detectably labeled. The labeled second antibody serves as a detecting antibody. The level of detectable label is measured, and the concentration of target antigen in the test sample is determined by comparison with a standard curve developed from the standard samples.

[000542] It will be appreciated that based on the results obtained using the antibodies and/or activatable antibodies of the disclosure in an *in vitro* diagnostic assay, it is possible to stage a disease in a subject based on expression levels of the Target antigen. For a given disease, samples of blood are taken from subjects diagnosed as being at various stages in the progression of the disease, and/or at various points in the therapeutic treatment of the disease. Using a population of samples that provides statistically significant results for each stage of progression or therapy, a range of concentrations of the antigen that may be considered characteristic of each stage is designated.

[000543] Antibodies, conjugated antibodies, activatable antibodies and/or conjugated activatable antibodies can also be used in diagnostic and/or imaging methods. In some embodiments, such methods are *in vitro* methods. In some embodiments, such methods are *in vivo* methods. In some embodiments, such methods are *in situ* methods. In some embodiments, such methods are *ex vivo* methods. For example, activatable antibodies having an enzymatically cleavable CM can be used to detect the presence or absence of an enzyme that is capable of cleaving the CM. Such activatable antibodies can be used in diagnostics, which can include *in vivo* detection (*e.g.*, qualitative or quantitative) of enzyme activity (or, in some embodiments, an environment of increased reduction potential such as that which can provide for reduction of a disulfide bond) through measured accumulation of activated antibodies (*i.e.*, antibodies resulting from cleavage of an activatable antibody) in a given cell or tissue of a given host organism. Such accumulation of activated antibodies indicates not only that the tissue expresses enzymatic activity (or an increased reduction potential depending on the nature of the CM) but also that the tissue expresses target to which the activated antibody binds.

[000544] For example, the CM can be selected to be a protease substrate for a protease found at the site of a tumor, at the site of a viral or bacterial infection at a biologically confined site (*e.g.*, such as in an abscess, in an organ, and the like), and the like. The AB can be one that binds a target antigen. Using methods familiar to one skilled in the art, a detectable label (*e.g.*, a fluorescent label or radioactive label or radiotracer) can be conjugated to an AB or other region of an activatable antibody. Suitable detectable labels are discussed in the context of the above screening methods and additional specific examples are provided below. Using an AB specific to a protein or peptide of the disease state, along with a protease whose activity is elevated in the disease tissue of interest, activatable antibodies will exhibit an increased rate of binding to disease tissue relative to tissues where the CM specific enzyme is not present at a detectable level or is present at a lower level than in disease tissue or is inactive (*e.g.*, in zymogen form or in complex with an inhibitor). Since small proteins and peptides are rapidly cleared from the blood by the renal filtration system, and because the enzyme specific for the CM is not present at a detectable level (or is present at lower levels in non-disease tissues or is present in inactive conformation), accumulation of activated antibodies in the disease tissue is enhanced relative to non-disease tissues.

[000545] In another example, activatable antibodies can be used to detect the presence or absence of a cleaving agent in a sample. For example, where the activatable antibodies contain a CM susceptible to cleavage by an enzyme, the activatable antibodies can be used to detect (either qualitatively or quantitatively) the presence of an enzyme in the sample. In another example, where the activatable antibodies contain a CM susceptible to cleavage by reducing agent, the activatable antibodies can be used to detect (either qualitatively or quantitatively) the presence of reducing conditions in a sample. To facilitate analysis in these methods, the activatable antibodies can be detectably labeled, and can be bound to a support (*e.g.*, a solid support, such as a slide or bead). The detectable label can be positioned on a portion of the activatable antibody that is not released following cleavage, for example, the detectable label can be a quenched fluorescent label or other label that is not detectable until cleavage has occurred. The assay can be conducted by, for example, contacting the immobilized, detectably labeled activatable antibodies with a sample suspected of containing an enzyme and/or reducing agent for a time sufficient for cleavage to occur, then washing to remove excess sample and contaminants. The presence or absence of the cleaving agent (*e.g.*, enzyme or reducing agent) in the sample is then assessed by a change

in detectable signal of the activatable antibodies prior to contacting with the sample *e.g.*, the presence of and/or an increase in detectable signal due to cleavage of the activatable antibody by the cleaving agent in the sample.

[000546] Such detection methods can be adapted to also provide for detection of the presence or absence of a target that is capable of binding the AB of the activatable antibodies when cleaved. Thus, the assays can be adapted to assess the presence or absence of a cleaving agent and the presence or absence of a target of interest. The presence or absence of the cleaving agent can be detected by the presence of and/or an increase in detectable label of the activatable antibodies as described above, and the presence or absence of the target can be detected by detection of a target-AB complex *e.g.*, by use of a detectably labeled anti-target antibody.

[000547] Activatable antibodies are also useful in *in situ* imaging for the validation of activatable antibody activation, *e.g.*, by protease cleavage, and binding to a particular target. *In situ* imaging is a technique that enables localization of proteolytic activity and target in biological samples such as cell cultures or tissue sections. Using this technique, it is possible to confirm both binding to a given target and proteolytic activity based on the presence of a detectable label (*e.g.*, a fluorescent label).

[000548] These techniques are useful with any frozen cells or tissue derived from a disease site (*e.g.* tumor tissue) or healthy tissues. These techniques are also useful with fresh cell or tissue samples.

[000549] In these techniques, an activatable antibody is labeled with a detectable label. The detectable label may be a fluorescent dye, (*e.g.* Fluorescein Isothiocyanate (FITC), Rhodamine Isothiocyanate (TRITC), a near infrared (NIR) dye (*e.g.*, Qdot® nanocrystals), a colloidal metal, a hapten, a radioactive marker, biotin and an amplification reagent such as streptavidin, or an enzyme (*e.g.* horseradish peroxidase or alkaline phosphatase).

[000550] Detection of the label in a sample that has been incubated with the labeled, activatable antibody indicates that the sample contains the target and contains a protease that is specific for the CM of the activatable antibody. In some embodiments, the presence of the protease can be confirmed using broad spectrum protease inhibitors such as those described herein, and/or by using an agent that is specific for the protease, for example, an antibody such as A11, which is specific for the protease matriptase and inhibits the proteolytic activity of matriptase; see *e.g.*, International Publication Number WO 2010/129609, published 11 November 2010. The same approach of using broad spectrum

protease inhibitors such as those described herein, and/or by using a more selective inhibitory agent can be used to identify a protease or class of proteases specific for the CM of the activatable antibody. In some embodiments, the presence of the target can be confirmed using an agent that is specific for the target, *e.g.*, another antibody, or the detectable label can be competed with unlabeled target. In some embodiments, unlabeled activatable antibody could be used, with detection by a labeled secondary antibody or more complex detection system.

[000551] Similar techniques are also useful for *in vivo* imaging where detection of the fluorescent signal in a subject, *e.g.*, a mammal, including a human, indicates that the disease site contains the target and contains a protease that is specific for the CM of the activatable antibody.

[000552] These techniques are also useful in kits and/or as reagents for the detection, identification or characterization of protease activity in a variety of cells, tissues, and organisms based on the protease-specific CM in the activatable antibody.

[000553] In some embodiments, *in situ* imaging and/or *in vivo* imaging are useful in methods to identify which patients to treat. For example, in *in situ* imaging, the activatable antibodies are used to screen patient samples to identify those patients having the appropriate protease(s) and target(s) at the appropriate location, *e.g.*, at a tumor site.

[000554] In some embodiments *in situ* imaging is used to identify or otherwise refine a patient population suitable for treatment with an activatable antibody of the disclosure. For example, patients that test positive for both the target and a protease that cleaves the substrate in the cleavable moiety (CM) of the activatable antibody being tested (*e.g.*, accumulate activated antibodies at the disease site) are identified as suitable candidates for treatment with such an activatable antibody comprising such a CM. Likewise, patients that test negative for either or both of the target and the protease that cleaves the substrate in the CM in the activatable antibody being tested using these methods are identified as suitable candidates for another form of therapy (*i.e.*, not suitable for treatment with the activatable antibody being tested). In some embodiments, such patients that test negative with respect to a first activatable antibody can be tested with other activatable antibodies comprising different CMs until a suitable activatable antibody for treatment is identified (*e.g.*, an activatable antibody comprising a CM that is cleaved by the patient at the site of disease).

[000555] In some embodiments *in vivo* imaging is used to identify or otherwise refine a patient population suitable for treatment with an activatable antibody of the disclosure.

For example, patients that test positive for both the target and a protease that cleaves the substrate in the cleavable moiety (CM) of the activatable antibody being tested (e.g., accumulate activated antibodies at the disease site) are identified as suitable candidates for treatment with such an activatable antibody comprising such a CM. Likewise, patients that test negative are identified as suitable candidates for another form of therapy (*i.e.*, not suitable for treatment with the activatable antibody being tested). In some embodiments, such patients that test negative with respect to a first activatable antibody can be tested with other activatable antibodies comprising different CMs until a suitable activatable antibody for treatment is identified (e.g., an activatable antibody comprising a CM that is cleaved by the patient at the site of disease).

Pharmaceutical compositions

[000556] The antibodies, conjugated antibodies, activatable antibodies and/or conjugated activatable antibodies of the disclosure (also referred to herein as “active compounds”), and derivatives, fragments, analogs and homologs thereof, can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the antibody, the conjugated antibody, activatable antibody and/or conjugated activatable antibody and a pharmaceutically acceptable carrier. As used herein, the term “pharmaceutically acceptable carrier” is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Suitable carriers are described in the most recent edition of Remington’s Pharmaceutical Sciences, a standard reference text in the field, which is incorporated herein by reference. Suitable examples of such carriers or diluents include, but are not limited to, water, saline, ringer’s solutions, dextrose solution, and 5% human serum albumin. Liposomes and non-aqueous vehicles such as fixed oils may also be used. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

[000557] A pharmaceutical composition of the disclosure is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (*i.e.*, topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[000558] Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringeability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In some embodiments, it will be desirable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, aluminum monostearate and gelatin.

[000559] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[000560] Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is

applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

[000561] For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser that contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

[000562] Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

[000563] The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

[000564] In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

[000565] It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the disclosure are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

[000566] The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

[000567] The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

Examples

EXAMPLE 1. Selection of human ScFvs of the embodiments that bind human and/or mouse PDL1

[000568] This Example demonstrates that ScFvs (single-chain variable fragments) of the embodiments that bind PDL1 can be selected from a phage display library of ScFvs with diverse CDR sequences, and that such binding can inhibit PDL1 binding to PD1 and B7-1.

[000569] ScFvs were selected from a fully human ScFv library displayed on M13 bacteriophage; ScFv phage selection was conducted under contract with Creative Biolabs, Shirley, NY. A fusion protein comprised of the extracellular domain (ECD) of human PDL1 or mouse PDL1, with a carboxy terminal His6 tag (Sino Biological, Cat. No. 10084-H02H-200) was used as the antigen in three alternating rounds of selection for ScFvs displayed on M13 bacteriophage that bind PDL1. In the first round, bound phage were released by trypsin digestion, and in subsequent rounds, phage were eluted by human PD1 fusion protein (R&D Systems; Cat. No. 1080-PD-050) competition. Five (5) unique ScFvs that bind PDL1 were isolated. Table 1 lists the 5 ScFvs and SEQ ID NOs of their respective nucleic acid sequences and amino acid sequences.

Table 1. SEQ ID NOs of selected ScFvs

ScFv	Nucleic acid sequence	Amino acid sequence
PDL1 c8	SEQ ID NO: 1	SEQ ID NO: 2
PDL1 c12	SEQ ID NO: 3	SEQ ID NO: 4
PDL1 c16	SEQ ID NO: 5	SEQ ID NO: 6
PDL1 c20	SEQ ID NO: 7	SEQ ID NO: 8
PDL1 c60	SEQ ID NO: 9	SEQ ID NO: 10

[000570] The nucleic acid and amino acid sequences of each of the anti-PDL1 ScFvs (with CDRs underlined) are shown below:

SEQ ID NO: 1

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAGATATTACTGCGTCGGGTTAGAGGACAACGTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAGATCGAAGATTGCTTTTGACTACTGGGGCCAGGGAACCCTGGT
CACCGTCTCGAGCGGTGGAGGCGGTTAGGCGGAGGTGGCAGCGGCGGTGGCGGGTCGACGGACATC
CAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGCCGGG
CAAGTCAGAGCATTAGCAGCTATTTAAATTGGTATCAGCAGAAACCAGGGAAAGCCCCTAAGCTCCT
GATCTATAAGGCATCCCGTTTGCAAAGTGGGGTCCCATCAAGGTTCAAGTGGCAGTGGATCTGGGACA
GATTTCACTCTCACCATCAGCAGTCTGCAACCTGAAGATTTTGCAACTTACTACTGTCAACAGCGTG
CGCTTAAGCCTGTGACGTTCTGGCCAAGGGACCAAGGTGGAAATCAAACGG

SEQ ID NO: 2

EVQLLES^{GGGLVQPGGSLRLS}CAASGFTF^{SSYAMSWVRQAPGKGLEWVSDITASGQRTTYADSVKGR}
^{FTISRDN}SKNTLYLQMN^{SLRAEDTAVYYCAR}^{SKIAFDYWGQ}TLVTVSSGGGGSGGGSGGGGSTD
I^{QMTQSPSSLSASVGRVTITCRASQSISSYLNWYQQKPKAPKLLIYKASRLQSGVPSRFS}SGSGSGT
DFTLT^{ISSLPEDFATYYCQQRAL}KPVTFTGQGTKVEIKR

SEQ ID NO: 3

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTATCCTGTG
CAGCCTCTGGATTACACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTAATAAGGATGGTCATTATACAAGTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAAAATCTTGATGAGTTTGACTACTGGGGCCAGGGAACCCTGGT

CACCGTCTCGAGCGGTGGAGGCGGTTCAGGCGGAGGTGGCAGCGGCGGTGGCGGGTCGACGGACATC
CAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGCCGGG
CAAGTCAGAGCATTAGCAGCTATTTAAATTGGTATCAGCAGAAACCAGGGAAAGCCCCCTAAGCTCCT
GATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCAAGGTTTCAGTGGCAGTGGATCTGGGACA
GATTTCACTCTCACCATCAGCAGTCTGCAACCTGAAGATTTTGCAACTTACTACTGTCAACAGAGTT
ACAGTACCCCTAATACGTTTCGGCCAAGGGACCAAGGTGGAAATCAAACGG

SEQ ID NO: 4

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSINKDGHYTSYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCAKNLDEFDYGQGT LVTVSSGGGSGGGSGGGSGT DI
QMTQSPSSLSASVGDRTITCRASQSISSYLNWYQQKPKAPKLLIYAASSLQSGVPSRFSGSGSGT
DFTLTISSLQPEDFATYYCQQSYSTPNTFGQGTKVEIKR

SEQ ID NO: 5

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCATCTATTATGGCTACTGGTGCTGGTACATTGTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAAGATGGTGCGGGGTTTGACTACTGGGGCCAGGGAACCCTGGT
CACCGTCTCGAGCGGTGGAGGCGGTTCAGGCGGAGGTGGCAGCGGCGGTGGCGGGTCGACGGACATC
CAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGCCGGG
CAAGTCAGAGCATTAGCAGCTATTTAAATTGGTATCAGCAGAAACCAGGGAAAGCCCCCTAAGCTCCT
GATCTATTCTGCATCCCAGTTGCAAAGTGGGGTCCCATCAAGGTTTCAGTGGCAGTGGATCTGGGACA
GATTTCACTCTCACCATCAGCAGTCTGCAACCTGAAGATTTTGCAACTTACTACTGTCAACAGGCGA
ATTCGCGGCCCTTCTACGTTTCGGCCAAGGGACCAAGGTGGAAATCAAACGG

SEQ ID NO: 6

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIMATGAGTLYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCAKD GAGFDYWGQGT LVTVSSGGGSGGGSGGGSGT DI
QMTQSPSSLSASVGDRTITCRASQSISSYLNWYQQKPKAPKLLIYASASQLQSGVPSRFSGSGSGT
DFTLTISSLQPEDFATYYCQQANSRPSTFGQGTKVEIKR

SEQ ID NO: 7

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GTAGTGGGTCTCAACTATTACTTCTTCTGGTGCTGCTACATATTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAAAATTATACTGGTTTTGACTACTGGGGCCAGGGAACCCTGGT

CACCGTCTCGAGCGGTGGAGGCGGTTTCAGGCGGAGGTGGCAGCGGCGGTGGCGGGTCGACGGACATC
CAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGCCGGG
CAAGTCAGAGCATTAGCAGCTATTTAAATTGGTATCAGCAGAAACCAGGGAAAGCCCCTAAGCTCCT
GATCTATAATGCATCCTCCTTGCAAAGTGGGGTCCCATCAAGGTTTCAGTGGCAGTGGATCTGGGACA
GATTTCACTCTCACCATCAGCAGTCTGCAACCTGAAGATTTTGCAACTTACTACTGTCAACAGTATA
CTTATGGTCTCTGGTACGTTTCGGCCAAGGGACCAAGGTGGAAATCAAACGG

SEQ ID NO: 8

EVQLLES^{GGGLVQPGGSLRLS}CAASGFTFSSYAMSWVRQAPGKGLQWVSTITSSGAATYYADSVKGR
FTISRDN^{SKNTLYLQMNSLRAEDTAVYYCAK}NYTGFDYWGQGLVTVSSGGGSGGGSGGGSGGGSTDI
QMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYNASSLQSGVPSRFSSGSGSGT
DFTLTIS^{SLQPEDFATYYCQQY}TGPGTFGQGTKVEIKR

SEQ ID NO: 9

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTATTCTACTGGTGGTGCTACAGCTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTTTTGACTACTGGGGCCAGGGAACCTGGT
CACCGTCTCGAGCGGTGGAGGCGGTTTCAGGCGGAGGTGGCAGCGGCGGTGGCGGGTCGACGGACATC
CAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGCCGGG
CAAGTCAGAGCATTAGCAGCTATTTAAATTGGTATCAGCAGAAACCAGGGAAAGCCCCTAAGCTCCT
GATCTATTATGCATCCACTTTGCAAAGTGGGGTCCCATCAAGGTTTCAGTGGCAGTGGATCTGGGACA
GATTTCACTCTCACCATCAGCAGTCTGCAACCTGAAGATTTTGCAACTTACTACTGTCAACAGGATA
ATGGTTATCCTTCTACGTTTCGGCCAAGGGACCAAGGTGGAAATCAAACGG

SEQ ID NO: 10

EVQLLES^{GGGLVQPGGSLRLS}CAASGFTFSSYAMSWVRQAPGKLEWVSSIYSTGGATAYADSVKGR
FTISRDN^{SKNTLYLQMNSLRAEDTAVYYCAK}SSAGFDYWGQGLVTVSSGGGSGGGSGGGSGGGSTDI
QMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYASTLQSGVPSRFSSGSGSGT
DFTLTIS^{SLQPEDFATYYCQQD}NGYPSTFGQGTKVEIKR

[000571] Figure 1 shows by ELISA-based binding that PDL1 c60 ScFv-phage binds specifically to human and mouse PDL1. Briefly, human PDL1, mouse PDL1, human PD1, mouse PD1, human 41BBLig or Notch1 (R&D Systems) were adsorbed to separate wells of a 96-well ELISA plate. Phage were applied to the plate and allowed to bind. Bound phage were visualized with an anti-M13-HRP conjugate (GE Healthcare, Piscataway, NY) and

developed with the chromogenic substrate tetramethyl benzidine (TMB) (Thermo Scientific, Rockford, IL).

[000572] Figure 2 shows that inclusion of PD1 can inhibit the binding of PDL1 ScFv-phage binding to PDL1 in an ELISA binding assay. Briefly, human PDL1 or mouse PDL1 (R&D Systems) was adsorbed to separate wells of a 96-well ELISA plate. Phage were applied to the plate in the presence or absence of human or mouse PD1-Fc (R&D Systems) and allowed to bind. Bound phage were visualized with an anti-M13-HRP conjugate (GE Healthcare, Piscataway, NY) and developed with the chromogenic substrate tetramethyl benzidine (TMB) (Thermo Scientific, Rockford, IL). Phage binding was reduced in the presence of PD1-Fc, demonstrating that the binding epitope of the phage overlaps with that of PD1 binding.

EXAMPLE 2. Production and testing of fully human anti-PDL1 c60 IgG antibodies

[000573] This Example demonstrates that PDL1 c60 ScFv-phage that bind human and mouse PDL1 can be converted into fully human IgG antibodies that retain human and mouse PDL1 binding.

[000574] Production of fully human IgGs comprising the variable domains of PDL1 c60 was accomplished using techniques similar to those described in PCT Publication No. WO 2010/081173. DNA molecules encoding the variable domains of PDL1 c60 ScFv-phage were cloned into expression vectors for the expression of fully human IgGs. Light chain (Lc) variable domains were amplified from the ScFv templates. Vector (LcpOP (modified from pCDNA3, Invitrogen, Carlsbad, CA)) and amplified DNA were cut with *Bsi*WI and *Eco*RI overnight, combined by ligation and transformed into *E. coli* MC1061 cells. Heavy chain (Hc) variable domains were amplified from ScFv templates. Fully human IgGs anti-PDL1 c60 were expressed from transiently transfected HEK-293 cells and purified from the culture supernatant by Protein A chromatography.

[000575] Figure 3 shows that the fully human anti-PDL1 c60 IgG binds human and mouse PDL1, with affinity for human PDL1 of 1 nM and for mouse PDL1 of 30 nM. Briefly human PDL1-Fc or mouse PDL1-Fc (R & D Systems, Minneapolis, MN) was adsorbed to the wells of a 96-well ELISA plate. Purified anti-PDL1 c60 was applied to the plate and allowed to bind. Bound antibody was visualized with an anti-human IgG-HRP conjugate, Fab specific, (Sigma, St Louis, MO; Cat # A0293-1ML) and developed with the chromogenic substrate TMB.

EXAMPLE 3. Affinity maturation of anti-PDL1 antibodies of the embodiments

[000576] This Example demonstrates the isolation of antibodies of the embodiments with improved binding kinetics and manufacturability profile.

[000577] Anti-PDL1 antibodies were isolated from libraries with CDRs modified from anti-PDL1 c60. Such libraries were designed as shown in Tables 2 and 3. Four libraries of Fab-Phage, based on the sequence of anti-PDL1 c60, were constructed using degenerate synthetic DNA. Residues were either varied by randomization at each indicated nucleotide. In addition, within libraries 3 and 6, additional randomized residues were added to CDR3 of the heavy chain. Libraries were transfected into *E. coli* strain TG1 and phage were prepared following super-infection with M13KO7 (Invitrogen).

[000578] Three rounds of selection were performed for each library with increasing stringency. Three alternating rounds of selection were done by mixing biotinylated Human PDL1 or Biotinylated mouse PDL1 with phage that were blocked with 100 µg/mL pooled human IgG (huIgG, or hIgG) and 2% non-fat dried milk (NFDM) in Tris-buffered saline (TBS; 40 mM Tris, 129 mM NaCl, pH 7.4). Following incubation at room temperature Bio-PDL1 bound to phage were captured with streptavidin magnetic beads which were washed extensively at 37⁰C and the remaining bound phage eluted with 100 mM triethanolamine (TEA) (Sigma, St. Louis, MO) and expanded through *E. coli* TG1. Selected Lc libraries were combined with Hc libraries and each subjected to an additional seven rounds of selection.

Table 2. CDR sequences for affinity maturation variable heavy chain libraries

IMGT	...	32	33	34	39	40	...	55	56	57	58	59	60	61	62	63	66	67	...	107	108	109	110	111	112	113							
VH3-23 germline		S	Y	A	M	S		A	I	S	G	S	G	G	S	T	Y	Y		-	-	-	-	-	-	-							
		(SEQ ID NO: 158)									(SEQ ID NO: 159)																						
Clone 60		S	Y	A	M	S		S	I	Y	S	T	G	G	A	T	A	Y		S	S	A	G	F	D	Y							
		(SEQ ID NO: 158)									(SEQ ID NO: 160)										(SEQ ID NO: 162)												
H2							x			x	x	x			x	x		x		3.36E+07													
(SEQ ID NO: 161)																																	
H3 walk																			(SEQ ID NO: 163)	x	x							1.02E+04					
																			(SEQ ID NO: 164)		x	x											
																			(SEQ ID NO: 165)			x	x										
																			(SEQ ID NO: 166)	x		x											
																			(SEQ ID NO: 167)	x			x										
																			(SEQ ID NO: 168)		x		x										
																			(SEQ ID NO: 169)	x					x								
																			(SEQ ID NO: 170)		x				x								
																			(SEQ ID NO: 171)				x		x								
(SEQ ID NO: 172)					x	x																											
H3 insertions: 4xNNK between G110 and F111																			(SEQ ID NO: 1203)					(SEQ ID NO: 173)					T(x) ₁₋₄ F		1E+06		

193273707 v1

205

[000579] Forty seven (47) individual isolates were chosen from the final pool. Phage were derived from each isolate and assayed for binding to human PDL1-Fc, human B7-1-Fc and human CD28-Fc antigens (R&D Systems). Briefly, the antigens were adsorbed to the wells of a 96-well ELISA plate, each ligand on a separate plate. Phage were applied to correlative wells on each plate and allowed to bind. Bound phage were visualized with an anti-M13-HRP conjugate and developed with the chromogenic substrate TMB. Based on the results of the ELISA and DNA sequence, 6 unique clones with matured heavy variable CDR3 domains and 10 heavy variable CDR2 domains were chosen for further study. Thirteen phage clones, 10 CDR2 and 3 CDR3 were analyzed for their ability to bind mouse and human PDL1, and the results of a phage ELISA are shown in Figure 4A and Figure 4B. All phage clones bind human PDL1 but not B7-1, in addition, CDR2 clones C05 and G12 show strong binding to mouse PDL1 and CDR3 clone H9 showed weak binding to mouse PDL1. Table 4 lists the heavy variable domains encoded by the clones and SEQ ID NOs for the nucleic acid sequences and amino acid sequences of their respective heavy chains. The light variable domains for each are identical to that of c60 (SEQ ID NO 11 and 12).

SEQ ID NO: 11

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTT
GCCGGGCAAGTCAGAGCATTAGCAGCTATTTAAATTGGTATCAGCAGAAACCAGGGAAAGCCCCTAA
GCTCCTGATCTATTATGCATCCACTTTGCAAAGTGGGGTCCCATCAAGGTTTCAGTGGCAGTGGATCT
GGGACAGATTTCACTCTCACCATCAGCAGTCTGCAACCTGAAGATTTTGCAACTTACTACTGTCAAC
AGGATAATGGTTATCCTTCTACGTTTCGGCCAAGGGACCAAGGTGGAAATCAAACGG

SEQ ID NO: 12

DIQMTQSPSSLSASVGDRTTITCRASQSISSYLNWYQQKPGKAPKLLIYYASTLQSGVPSRFSGSGS
GTDFTLTISSLQPEDFATYYCQQDNGYPSTFGQGTKVEIKR

Table 4: Matured anti-PDL1 variable domains

VH	Nucleic acid sequence	Amino acid sequence
PDL1 c1	SEQ ID NO: 13	SEQ ID NO: 14
PDL1 d1	SEQ ID NO: 15	SEQ ID NO: 16
PDL1 g7	SEQ ID NO: 17	SEQ ID NO: 18
PDL1 h9	SEQ ID NO: 19	SEQ ID NO: 20
PDL1 b10	SEQ ID NO: 21	SEQ ID NO: 22

PDL1 E10	SEQ ID NO: 23	SEQ ID NO: 24
PDL1 A05	SEQ ID NO: 25	SEQ ID NO: 26
PDL1 C05	SEQ ID NO: 27	SEQ ID NO: 28
PDL1 C10	SEQ ID NO: 29	SEQ ID NO: 30
PDL1 D08	SEQ ID NO: 31	SEQ ID NO: 32
PDL1 G09	SEQ ID NO: 33	SEQ ID NO: 34
PDL1 G10	SEQ ID NO: 35	SEQ ID NO: 36
PDL1 G12	SEQ ID NO: 37	SEQ ID NO: 38
PDL1 E11	SEQ ID NO: 39	SEQ ID NO: 40
PDL1 D01	SEQ ID NO: 41	SEQ ID NO: 42
PDL1 H06	SEQ ID NO: 43	SEQ ID NO: 44

SEQ ID NO: 13

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTATTCTACTGGTGGTGCTACAGCTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTAGTCGGCCGGGTTTTGACTACTGGGGCCA
GGGAACCCTGGTCACCGTCTCGAGC

SEQ ID NO: 14

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIYSTGGATAYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCAKSSAGQSRPGFDYWGQGTLVTVSS

SEQ ID NO: 15

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTATTCTACTGGTGGTGCTACAGCTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTTCGTGGCCGGGTTTTGACTACTGGGGCCA
GGGAACCCTGGTCACCGTCTCGAGC

SEQ ID NO: 16

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIYSTGGATAYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCAKSSAGQSWPGFDYWGQGTLVTVSS

SEQ ID NO: 17

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTATTCTACTGGTGGTGCTACAGCTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTCAGTCGTTTCCGGGTTTTGACTACTGGGG
CCAGGGAACCCTGGTCACCGTCTCGAGC

SEQ ID NO: 18

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIYSTGGATAYADSVKGR
FTISRDNSKNTLYLQMNSLRAEDTAVYYCAKSSAGQSFPGFDYWQGTLVTVSS

SEQ ID NO: 19

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTATTCTACTGGTGGTGCTACAGCTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAATGGTCTGCTGCTTTTGACTACTGGGGCCAGGGAACCCTGGT
CACCGTCTCGAGC

SEQ ID NO: 20

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIYSTGGATAYADSVKGR
FTISRDNSKNTLYLQMNSLRAEDTAVYYCAKWSAAFDYWQGTLVTVSS

SEQ ID NO: 21

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTATTCTACTGGTGGTGCTACAGCTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAATGGTCTGCTGCTTTTGACTACTGGGGCCAGGGAACCCTGGT
CACCGTCTCGAGC

SEQ ID NO: 22

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIYSTGGATAYADSVKGR
FTISRDNSKNTLYLQMNSLRAEDTAVYYCAKWSAGYDYWGQGLVTVSS

SEQ ID NO: 23

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTATTCTACTGGTGGTGCTACAGCTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAATGGTCTAAGGGTTTTGACTACTGGGGCCAGGGAACCCTGGT
CACCGTCTCGAGC

SEQ ID NO: 24

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIYSTGGATAYADSVKGR
FTISRDNSKNTLYLQMNSLRAEDTAVYYCAKWSKGFQDYWGQGTLVTVSS

SEQ ID NO: 25

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTGGAAGTAGGGTATTGTGACAGTGAGCTTACGCAGACTCCGTGAAGGG
CCGGTTACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCC
GAGGACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTTTTGACTACTGGGGCCAGGGAACCC
TGGTCACCGTCTCGAGC

SEQ ID NO: 26

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWKQGIIVTVYDSVKGRF
TISRDNSKNTLYLQMNSLRAEDTAVYYCAKSSAGFDYWGQGTLVTV

SEQ ID NO: 27

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTGGCGGAATGGTATTGTTACAGTTAGCTTACGCAGACTCCGTGAAGGG
CCGGTTACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCC
GAGGACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTTTTGACTACTGGGGCCAGGGAACCC
TGGTCACCGTCTCGAGC

SEQ ID NO: 28

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWRNGIIVTVYDSVKGRF
TISRDNSKNTLYLQMNSLRAEDTAVYYCAKSSAGFDYWGQGTLVTVSS

SEQ ID NO: 29

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAGATATTTGGAAGTAGGGTATGGTTACAGTGAGCTTACGCAGACTCCGTGAAGGG
CCGGTTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCC
GAGGACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTTTTGACTACTGGGGCCAGGGAACCC
TGGTCACCGTCTCGAGC

SEQ ID NO: 30

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSDIWKQGMVTYDSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCAKSSAGFDYWGQGLVTVSS

SEQ ID NO: 31

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCATCGATTTGGAGGTAGGGTCTGGCGACAGCGAGCTTACGCAGACTCCGTGAAGGG
CCGGTTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCC
GAGGACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTTTTGACTACTGGGGCCAGGGAACCC
TGGTCACCGTCTCGAGC

SEQ ID NO: 32

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWRQGLATAYDSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCAKSSAGFDYWGQGLVTVSS

SEQ ID NO: 33

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAGAGATTGTGGCTACTGGTATTTTGACAAGTAGCTTACGCAGACTCCGTGAAGGG
CCGGTTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCC
GAGGACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTTTTGACTACTGGGGCCAGGGAACCC
TGGTCACCGTCTCGAGC

SEQ ID NO: 34

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSEIVATGILTSYDSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCAKSSAGFDYWGQGLVTVSS

SEQ ID NO: 35

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCATCGATTGGTCGGTAGGGTTTGATTACAGTTAGCTTACGCAGACTCCGTGAAGGG
CCGGTTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCC
GAGGACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTTTTGACTACTGGGGCCAGGGAACCC
TGGTCACCGTCTCGAGC

SEQ ID NO: 36

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIGRQGLITVYDSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCAKSSAGFDYWGQGTLVTVSS

SEQ ID NO: 37

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCATCTATTTGGTATTAGGGTCTGGTGACAGTTAGCTTACGCAGACTCCGTGAAGGG
CCGGTTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCC
GAGGACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTTTTGACTACTGGGGCCAGGGAACCC
TGGTCACCGTCTCGAGC

SEQ ID NO: 38

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWYQGLVTVYDSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCAKSSAGFDYWGQGTLVTVSS

SEQ ID NO: 39

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAGATATTTGGAAGTAGGGTTTTGCTACAGCGAGCTTACGCAGACTCCGTGAAGGG
CCGGTTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCC
GAGGACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTTTTGACTACTGGGGCCAGGGAACCC
TGGTCACCGTCTCGAGC

SEQ ID NO: 40

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSDIWKQGFATADSVKGRFT
ISRDN SKNTLYLQMNSLRAEDTAVYYCAKSSAGFDYWGQGTLVTVSS

SEQ ID NO: 41

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTGGAAGTAGGGTATTGTGACAGTGAGCTTACGCAGACTCCGTGAAGGG
CCGGTTACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCC
GAGGACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTTTTGACTACTGGGGCCAGGGAACCC
TGGTCACCGTCTCGAGC

SEQ ID NO: 42

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWKQGIVTVYDSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCAKSSAGFDYWGQGLVTVSS

SEQ ID NO: 43

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCATCGATTTGGAGGTAGGGTCTGGCGACAGCGAGCTTACGCAGACTCCGTGAAGGG
CCGGTTACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCC
GAGGACACGGCCGTATATTACTGTGCGAAATCTTCTGCTGGTTTTGACTACTGGGGCCAGGGAACCC
TGGTCACCGTCTCGAGC

SEQ ID NO: 44

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWRQGLATAYDSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCAKSSAGFDYWGQGLVTVSS

EXAMPLE 4. Isolation and testing of affinity matured anti-PDL1 antibodies of the embodiments

[000580] This Example demonstrates that matured PDL1 Fab-phage that bind human and mouse PDL1 can be converted into fully human IgG antibodies that retain human and show enhanced mouse PDL1 binding.

[000581] Production of fully human IgGs comprising the variable domains of the matured clones was accomplished using techniques similar to those described in PCT Publication No. WO 2010/081173. DNA encoding the variable domains of anti-PDL1 Fab-Phage clones C1, D1, G7, C05, G12 and combinations of heavy variable CDR3 domains of anti-PDL1 Fab-Phage clones C05 or G12 with anti-PDL1 Fab-Phage clones heavy variable CDR2 domains of B10, E10 or H9 were cloned into expression vectors for the expression of

fully human IgGs. The light variable domains for each are identical to that of c60 (SEQ ID NO: 10). Light chain (Lc) variable domains were amplified from the Fab templates. Vector (LcpOP (modified from pCDNA3, Invitrogen, Carlsbad, CA)) and amplified DNA were cut with *Bsi*WI and *Eco*RI overnight, combined by ligation and transformed into *E. coli* MC1061 cells. Heavy chain (Hc) variable domains were amplified from Fab templates. Fully human IgGs anti-PDL1 antibodies were expressed from transiently transfected HEK-293 cells and purified from the culture supernatant by Protein A chromatography.

[000582] The affinity matured anti-PDL1 antibodies were analyzed for their ability to bind human and mouse PDL1. Briefly human PDL1-Fc or mouse PDL1-Fc (R & D Systems, Minneapolis, MN) was adsorbed to the wells of a 96-well ELISA plate. Purified anti-PDL1 antibodies were applied to the plate and allowed to bind. Bound antibody was visualized with an anti-human IgG-HRP conjugate, Fab specific, (Sigma, St Louis, MO; Cat # A0293-1ML) and developed with the chromogenic substrate TMB. Figure 5 shows that combining the matured heavy variable CDR3 domains with the heavy variable CDR2 domains results in antibodies with enhanced affinity towards human PDL1. In addition, the combination of the matured heavy variable CDR2 domain C05 with the heavy variable CD3 domain H9 results in an antibody with high affinity for both human and mouse PDL1. Table 5 provides the SEQ ID NOs for the nucleic acid sequences and amino acids sequences of the respective variable heavy chains.

[000583] The anti-PDL1 antibodies C5H9, C5B10 and C5E10 were further characterized for their species specificity towards binding to PDL1 and other closely related proteins. Briefly human PDL1-Fc, mouse PDL1-Fc (R & D Systems, Minneapolis, MN) or cynomolgus monkey (cyno) PDL1-Fc (Sino biological) were adsorbed to the wells of a 96-well ELISA plate. Purified anti-PDL1 antibodies were applied to the plate and allowed to bind. Bound antibody was visualized with an anti-human IgG-HRP conjugate, Fab specific, (Sigma, St Louis, MO; Cat # A0293-1ML) and developed with the chromogenic substrate TMB. Figures 6A and 6B show that C5H9 binds to human and mouse PDL1 with near equal affinity and C5B10 and C5E10 only bind human PDL1. None of the antibodies tested bind to either human B7-1 (Figure 6D) or human CD28 (Figure 6C), two closely related proteins. Figure 7 shows that anti-PDL1 antibody C5H9 binds with high and equal affinity to human, mouse and cyno PDL1, but not to human B7H3 (also referred to as B7-H3) (R&D Systems, Minneapolis, MN).

Table 5: Variable heavy domains of PDL1 antibodies of the embodiments

VH	Nucleic acid sequence	Amino acid sequence
PDL1 C5H9	SEQ ID NO: 45	SEQ ID NO: 46
PDL1 C5B10	SEQ ID NO: 47	SEQ ID NO: 48
PDL1 C5E10	SEQ ID NO: 49	SEQ ID NO: 50
PDL1 G12H9	SEQ ID NO: 51	SEQ ID NO: 52
PDL1 G12B10	SEQ ID NO: 53	SEQ ID NO: 54
PDL1 G12E10	SEQ ID NO: 55	SEQ ID NO: 56

SEQ ID NO: 45

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTGGCGGAATGGTATTGTTACAGTTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAATGGTCTGCTGCTTTTGACTIONTGGGGCCAGGGAACCCTGGT
CACCGTCTCGAGC

SEQ ID NO: 46

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWRNGIVTVYADSVKGR
FTISRDNKNTLYLQMNSLRAEDTAVYYCAKWSAAFDYWGQGLVTVSS

SEQ ID NO: 47

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTGGCGGAATGGTATTGTTACAGTTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAATGGTCTGCTGGTTATGACTACTGGGGCCAGGGAACCCTGGT
CACCGTCTCGAGC

SEQ ID NO: 48

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWRNGIVTVYADSVKGR
FTISRDNKNTLYLQMNSLRAEDTAVYYCAKWSAGYDYWGQGLVTVSS

SEQ ID NO: 49

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG

CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTGGCGGAATGGTATTGTTACAGTTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAATGGTCTAAGGGTTTTGACTACTGGGGCCAGGGAACCCTGGT
CACCGTCTCGAGC

SEQ ID NO: 50

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWRNGIVTVYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCAKWSKGFYWGQGTLVTVSS

SEQ ID NO: 51

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCATCTATTTGGTATCAGGGTCTGGTGACAGTTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAATGGTCTGCTGCTTTTACTACTGGGGCCAGGGAACCCTGGT
CACCGTCTCGAGC

SEQ ID NO: 52

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWYQGLVTVYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCAKWSAAFYWGQGTLVTVSS

SEQ ID NO: 53

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
GGAGTGGGTCTCAAGTATTTGGCGGAATGGTATTGTTACAGTTTACGCAGACTCCGTGAAGGGCCGG
TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
ACACGGCCGTATATTACTGTGCGAAATGGTCTGCTGGTTATGACTACTGGGGCCAGGGAACCCTGGT
CACCGTCTCGAGC

SEQ ID NO: 54

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWYQGLVTVYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCAKWSAGYDYWGQGTLVTVSS

SEQ ID NO: 55

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTG
 CAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCT
 GGAGTGGGTCTCAAGTATTTGGCGGAATGGTATTGTTACAGTTTACGCAGACTCCGTGAAGGGCCGG
 TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGG
 ACACGGCCGTATATTACTGTGCGAAATGGTCTAAGGGTTTTGACTACTGGGGCCAGGGAACCCTGGT
 CACCGTCTCGAGC

SEQ ID NO: 56

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWYQGLVTVYADSVKGR
 FTISRDNKNTLYLQMNSLRAEDTAVYYCAKWSKGFQYWGQGLTVTVSS

EXAMPLE 5. Germline revisions in anti-PDL1 antibody C5H9 light variable domain enhances manufacturability and reduces potential immunogenicity.

[000584] This example describes an additional anti-PDL1 antibody embodiment that exhibits reduced predicted immunogenicity, reduced aggregation, and improved expression.

[000585] In silico immunogenicity prediction, using ProPred revealed a strong potential epitope in CDR2 of the light variable domain of C5H9; anti-PDL1 antibody C5H9, also referred to herein as C5H9, comprises a light chain comprising SEQ ID NO: 12 and a heavy chain comprising SEQ ID NO: 46. Alignment of the light variable domain of C5H9 with human germline antibodies revealed two amino acids in CDR2 that when reverted to germline eliminated the predicted epitope. The residues are underlined and in bold text in the sequence shown below in SEQ ID NO: 12.

[000586] Furthermore, framework 4 of the C5H9 light chain was modified as indicated with bold italics in SEQ ID NO: 58 substituting glycine (G) for glutamine (Q) in the J region. This modification was aimed at increasing the flexibility of the junction between variable and constant regions, potentially reducing the tendency of C5H9 to form aggregates. The resultant DNA and amino acid sequences are described by SEQ ID NO: 57 and SEQ ID NO: 58, respectively. The resultant anti-PDL1 antibody, referred to herein as anti-PDL1 antibody C5H9v2, also referred to herein as C5H9v2, comprises a light chain comprising SEQ ID NO: 58 and a heavy chain comprising SEQ ID NO: 46.

SEQ ID NO: 12

DIQMTQSPSSLSASVGDRTTITCRASQSISSYLNWYQQKPGKAPKLLIY**Y****A****S****T**LQSGVPSPRFSGSGS
 GTDFTLTISLQPEDFATYYCQQDNGYPSTFGQGTKVEIKR

SEQ ID NO: 57

GATATTGAGATGACCCAGAGCCCGAGCAGCCTGAGCGCGAGCGTGGGCGATCGCGTGACCATTACCT
GCCGCGCGAGCCAGAGCATTAGCAGCTATCTGAACTGGTATCAGCAGAAACCGGGCAAAGCGCCGAA
ACTGCTGATTTATGCGGCGAGCAGCCTGCAGAGCGGCGTGCCGAGCCGCTTTAGCGGCAGCGGCAGC
GGCACCGATTTTACCCTGACCATTAGCAGCCTGCAGCCGGAAGATTTTGCAGACCTATTATTGCCAGC
AGGATAACGGCTATCCGAGCACCTTTGGCGGCGGCACCAAAGTGGAATTAACGC

SEQ ID NO: 58

DIQMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSSLQSGVPSRFSGSGS
GTDFTLTISSLQPEDFATYYCQQDNGYPSTFGG**G**TKVEIKR

[000587] Figures 8A and 8B show that germline revisions described in SEQ ID NO: 58 reduces the predicted immunogenicity (top panel) and aggregation potential, and increases the expression level by three fold.

EXAMPLE 6. *In vitro* characterization of anti-PDL1 antibodies C5H9 and C5H9v2.

[000588] This Example demonstrates the binding specificities and biological activities of anti-PDL1 antibodies of the disclosure.

[000589] Production of fully human IgGs comprising the variable domains of the matured clones was accomplished using techniques similar to those described in PCT Publication No. WO 2010/081173. DNA encoding the heavy variable domains of anti-PDL1 antibody C5H9 (nucleic acid SEQ ID NO: 45; amino acid SEQ ID NO: 46) were amplified and cloned into vectors for the expression of fully human IgGs. DNA encoding the light variable domains of anti-PDL1 antibody C5H9 (nucleic acid SEQ ID NO: 11; amino acid SEQ ID NO: 12) were amplified and cloned into vectors for the expression of fully human IgGs. Similarly, DNA encoding the heavy variable domains of anti-PDL1 antibody C5H9 (nucleic acid SEQ ID NO: 45; amino acid SEQ ID NO: 46) were amplified and cloned into vectors for the expression of chimeric mouse IgG2as. DNA encoding the light variable domains of anti-PDL1 antibody C5H9v2 (nucleic acid SEQ ID NO: 57; amino acid SEQ ID NO: 58) were amplified and cloned into vectors for the expression of chimeric mouse IgG2as. Fully human IgGs or chimeric mouse IgG2a anti-PDL1 antibodies were expressed from transiently transfected HEK-293 cells and purified from the culture supernatant by Protein A chromatography. Anti-PDL1 IgG antibody C5H9 comprises a

heavy chain comprising a heavy variable domain comprising SEQ ID NO: 46 and a light chain comprising a light variable domain comprising SEQ ID NO: 12. Anti-PDL1 IgG antibody C5H9v2 comprises a heavy chain comprising a heavy variable domain comprising SEQ ID NO: 46 and a light chain comprising a light variable domain comprising SEQ ID NO: 58.

[000590] Anti-PDL1 antibody C5H9v2 was characterized for species specific binding to PDL1 and whether that antibody would bind a panel of human proteins. Briefly human PDL1-Fc, mouse PDL1-Fc (R & D Systems, Minneapolis, MN) or cynomolgus monkey (cyno) PDL1-Fc (Sino biological) was adsorbed to the wells of a 96-well ELISA plate. Purified anti-PDL1 antibodies C5H9 and C5H9v2 were applied to the plate and allowed to bind. Bound antibody was visualized with an anti-human IgG-HRP conjugate, Fab specific, (Sigma, St Louis, MO; Cat # A0293-1ML) and developed with the chromogenic substrate TMB. Figure 9 shows that both anti-PDL1 antibodies C5H9 and C5H9v2 bind human, mouse and cyno PDL1 with near equal affinity. EC₅₀ values for anti-PDL1 antibody C5H9v2 to human, cyno, rat and mouse PDL1 were also similar: 0.25 nM, 0.28 nM, 0.31 nM, and 0.30 nM, respectively.

[000591] In a similar experiment the binding of anti-PDL1 antibody C5H9 was evaluated for binding to a panel of human and mouse proteins. Briefly, each protein, listed in Figure 10, was absorbed to the wells of an ELISA plate. Purified anti-PDL1 antibodies C5H9 and C5H9v2 were applied to the plate and allowed to bind. Bound antibody was visualized with an anti-human IgG-HRP conjugate, Fab specific, (Sigma, St Louis, MO; Cat # A0293-1ML) and developed with the chromogenic substrate TMB. Anti-PDL1 antibody C5H9v2 binds only to human and mouse PDL1 demonstrating specificity toward PDL1.

[000592] The biological activity of PDL1 is mediated through binding to PD1 and/or to B7-1. Blockade of binding is a desired characteristic for a therapeutic antibody; therefore, the potency of blocking the interaction of PDL1 with PD1 and B7-1 was measured. Briefly, human, mouse or cyno PDL1 was adsorbed to the well of an ELISA plate. Biotinylated human, mouse or cyno PD1 or B7-1 was applied to the wells in the absence or the presence of an increasing concentration of either anti-PDL1 antibody C5H9 or anti-PDL1 antibody C5H9v2 and allowed to bind. Bound biotinylated PD1 or B7-1 was visualized with a Streptavidin-HRP conjugate (Thermo Scientific, Cat. No. 21126) and developed with the chromogenic substrate TMB. Figure 11 shows that both anti-PDL1 antibody C5H9 and anti-PDL1 antibody C5H9v2 are potent blockers of either B7-1 or PD1 binding to PDL1, and

that blockade includes all three species human, cyno and mouse with single digit nM EC₅₀'s.

EXAMPLE 7: An anti-PDL1 antibody of the disclosure accelerates induction of diabetes in NOD mice

[000593] In this Example, anti-PDL1 antibody C5H9 was analyzed for the ability to induce diabetes in NOD mice.

[000594] The NOD mice, substrain NOD/ShiLtJ, were obtained from Jackson Laboratory at 6 weeks and acclimated on site for 1 week. At 7 weeks, mice were checked for diabetes prior to enrollment, grouped, and dosed as set forth in Table 6.

Table 6. Groups and Doses for NOD study

Group	Count	Treatment	Dose (mg/kg) Loading/regular	Dose volume (mL/kg)	Schedule	Route
1	6	Control	20/10	10	q2dx6	IP
2	6	Anti-PDL1(C5H9)	20/10	10	q2dx6	IP
3	6	Anti-PDL1 (10F9G2)	20/10	10	q2dx6	IP

[000595] Figure 12, which plots % non-diabetic versus number of days post initial dose, shows that anti-PDL1 antibody C5H9 antibody (Gr2-C5H9) induces diabetes in NOD mice similar to positive control anti-PDL1 antibody 10F9G2 antibody. A control rat IgG2b did not induce diabetes in NOD mice.

[000596] Anti-PDL1 antibody C5H9v2 was also tested for the ability to induce diabetes in NOD mice in a dose dependent matter, using a method similar to that described above, with 9-week old animals using the groups and doses set forth in Table 7.

Table 7. Groups and doses for dose dependent NOD study

Group	Count	Gender	Treatment	Dose (mg/kg)	Dose volume (mL/kg)	Schedule	Route
1	8	F	mIgG2a (C1.18.4)	25	10	Single dose	IP
2	8	F	Anti-PDL1(C5H9v2)	25	10	Single dose	IP
3	8	F	Anti-PDL1(C5H9v2)	5	10	Single dose	IP
4	8	F	Anti-PDL1(C5H9v2)	1	10	Single dose	IP
5	8	F	Anti-PDL1(C5H9v2)	0.2	10	Single dose	IP

[000597] Figure 13, which plots % non-diabetic versus number of days post initial dose, demonstrates that anti-PDL1 antibody C5H9v2 induces diabetes in NOD mice in a dose dependent matter.

EXAMPLE 8. An anti-PDL1 antibody of the embodiments reduces MC38 tumors in mice

[000598] In this Example, anti-PDL1 antibody C5H9v2 was analyzed for the ability to reduce the growth of MC38 syngeneic tumors.

[000599] The mouse colon carcinoma cell line MC38 was obtained from ATCC (American Type Culture Collection, Manassas, VA). MC38 cells were grown in RPMI-1640 supplemented with 10% fetal bovine serum at 37°C in an atmosphere of 5% CO₂ in air. Cells were harvested during the logarithmic growth period, resuspended in PBS, and kept on ice for tumor induction.

[000600] Each mouse was inoculated subcutaneously at the right flank with 1X10⁶ of MC38 cells in PBS for tumor development. The treatments were started when the mean tumor size reached approximately 100-200 mm³ (no more than 200 mm³). Tumor sizes were measured twice weekly in two dimensions using a caliper, and the volume was expressed in mm³ using the formula: $V = 0.5 \times a \times b^2$ where *a* and *b* are the long and short diameters of the tumor, respectively.

[000601] The mice were grouped and dosed as set forth in Table 8.

Table 8. Groups and doses for MC38 syngeneic study

Group	Count	Treatment	Dose (mg/kg)	Dose volume (mL/kg)	Schedule	Route
1	10	Anti-PDL1 (10F9G2)	10	10	t.i.w. for 2 weeks (MWF)	IP
2	10	Anti-PDL1(C5H9v2)	10	10	t.i.w. for 2 weeks (MWF)	IP
3	10	mIgG2a + Hamster IgG1	10	10	t.i.w. for 2 weeks (MWF)	IP

[000602] Figure 14, which plots tumor volume versus number of days post initial dose, demonstrates that anti-PDL1 antibody C5H9v2 inhibits the growth of MC38 syngeneic tumors similar to positive control anti-PDL1 antibody 10F9G2.

EXAMPLE 9. Anti-PDL1 activatable antibody C5H9v2 masking moieties

[000603] This Example describes identification of masking moieties (MM) to reduce binding of activatable antibodies of anti-PDL1 antibody C5H9v2 to their target.

[000604] Anti-PDL1 antibody C5H9v2 and Fab were used to screen the libraries using a method similar to that described in PCT International Publication Number WO 2010/081173, published 15 July 2010. The screening consisted of one round of MACS and three rounds of FACS sorting. The initial MACS was done with protein-A Dynabeads (Invitrogen) and the anti-PDL1 antibody C5H9v2 at a concentration of 100 nM. For MACS, approximately 1×10^{11} cells were screened for binding and 6×10^6 cells were collected. Anti-PDL1 antibody C5H9v2, directly labeled with Alexafluor-488, was used as a probe for all FACS selections. In the first round of FACS, the cells were labeled with 100 nM AlexaFluor- anti-PDL1 antibody C5H9v2. In the second round of FACS, the cells were labeled with 10 nM AlexaFluor- anti-PDL1 antibody C5H9v2. In the third round of FACS, the cells were labeled with 1 nM AlexaFluor- anti-PDL1 antibody C5H9v2. The positive population from the third FACS round was verified to be inhibited by recombinant human PDL1 protein from binding to the anti-PDL1 antibody C5H9v2 and Fab. Individual peptide clones were identified by sequence analysis and subsequently verified for their ability to bind the anti-PDL1 antibody C5H9v2 and Fab.

[000605] The sequences of the anti-PDL1 antibody C5H9v2 masking moieties are listed in Table 9.

Table 9. anti-PDL1 antibody C5H9v2 masking moieties (MM)

MM	Amino Acid Sequence	SEQ ID NO
PL01	YCEVSELFVLPWCMG	SEQ ID NO: 208
PL02	SCLMHPHYAHDYCYV	SEQ ID NO: 426
PL03	LCEVLMLLQHPWCMG	SEQ ID NO: 59
PL04	IACRHFMEQLPFCHH	SEQ ID NO: 60
PL05	FGPRCGEASTCVPYE	SEQ ID NO: 61
PL06	ILYCDSWGAGCLTRP	SEQ ID NO: 62
PL07	GIALCPSHFQCQLPQT	SEQ ID NO: 63
PL08	DGPRCFVSGECSPIG	SEQ ID NO: 64
PL09	LCYKLDYDDRSYCHI	SEQ ID NO: 65

PL10	PCHPHPYDARPYCNV	SEQ ID NO: 66
PL11	PCYWHPFFAYRYCNT	SEQ ID NO: 67
PL12	VCYYMDWLGRNWCSS	SEQ ID NO: 68
PL13	LCDLFLKLREFPYCMG	SEQ ID NO: 69
PL14	YLPCHFVPIGACNNK	SEQ ID NO: 70
PL15	IFCHMGVVVPQCANY	SEQ ID NO: 71
PL16	ACHPHPYDARPYCNV	SEQ ID NO: 72
PL17	PCHPAPYDARPYCNV	SEQ ID NO: 73
PL18	PCHPHAYDARPYCNV	SEQ ID NO: 74
PL19	PCHPHPADARPYCNV	SEQ ID NO: 75
PL20	PCHPHPYAARPYCNV	SEQ ID NO: 76
PL21	PCHPHPYDAAPYCNV	SEQ ID NO: 77
PL22	PCHPHPYDARPACNV	SEQ ID NO: 78
PL23	PCHPHPYDARPYCAV	SEQ ID NO: 79
PL24	PCHAHPYDARPYCNV	SEQ ID NO: 80
PL25	PCHPHPYDARAYCNV	SEQ ID NO: 81

EXAMPLE 10. Activatable antibodies comprising anti-PDL1 antibody C5H9v2

[000606] This Example describes examples of activatable antibodies of the disclosure comprising anti-PDL1 antibody C5H9v2.

[000607] Anti-PDL1 activatable antibodies comprising an anti-PDL1 antibody C5H9v2 masking moiety, a cleavable moiety, and anti-PDL1 antibody C5H9v2 were produced according to methods similar to those described in PCT Publication Nos. WO 2009/025846 and WO 2010/081173. Quality control of the resultant activatable antibodies indicated that most comprised at least 95% monomer. The amino acid and nucleic acid sequences of several activatable antibodies comprising anti-PDL1 antibody C5H9v2 are provided below. Each activatable antibody comprises anti-PDL1 antibody C5H9v2 having the variable heavy chain amino acid sequence shown in SEQ ID NO: 46 and a light chain amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID NO: 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959,

961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, 1144-1191, 1200, and 1201 below.

[000608] While the sequences shown below include the spacer sequence of SEQ ID NO: 923, those of ordinary skill in the art appreciate that the activatable anti-PDL1 antibodies of the disclosure can include any suitable spacer sequence, such as, for example, a spacer sequence selected from the group consisting of QGQSGS (SEQ ID NO: 923); GQSGS (SEQ ID NO: 1192); QSGS (SEQ ID NO: 1193); SGS (SEQ ID NO: 1194); GS (SEQ ID NO: 1195); S; QGQSGQG (SEQ ID NO: 924); GQSGQG (SEQ ID NO: 395); QSGQG (SEQ ID NO: 925); SGQG (SEQ ID NO: 926); GQG (SEQ ID NO: 927); QG (SEQ ID NO: 928); G; QGQSGQ (SEQ ID NO: 1196); GQSGQ (SEQ ID NO: 1197); QSGQ (SEQ ID NO: 1198); SGQ (SEQ ID NO: 616); GQ (SEQ ID NO: 1199); and Q. While the sequences shown below include the spacer sequence of SEQ ID NO: 923, those of ordinary skill in the art will also appreciate that activatable anti-PDL1 antibodies of the disclosure in some embodiments do not include a spacer sequence.

C5H9v2 VH amino acid sequence

EVQLLES GGGVLVQPGGSLRLS CAASGFTFSSYAMSWVRQAPGKGLEWVSSIWRNGIVTVYADSVKGR
FTISRDNSKNTLYLQMNSLRAEDTAVYYCAKWSAAFDYWGQGT LVT VSS (SEQ ID NO: 46)

[spacer (SEQ ID NO: 929)] [PL01-0003 LC (SEQ ID NO: 930)]
[CAAGGTCAGTCTGGATCC] [TATTGCGAGGTTAGTGAGCTGTTTGTCTTCCTTGGTGCATGGGTG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAA ACTGCTGATCTACGCCGCGCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 82)

[spacer (SEQ ID NO: 923)] [PL01-0003 LC (SEQ ID NO: 931)]
[QGQSGS] [YCEVSELFVLPWCMGGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 83)

PL01-0003 LC nucleotide sequence

TATTGCGAGGTTAGTGAGCTGTTTGTTCCTTGGTGCATGGGTGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAACCTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 930)

PL01-0003 LC amino acid sequence

YCEVSELFVLPWCMGGGSSGSGSGSGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 931)

[spacer (SEQ ID NO: 929)] [PL02-0003 LC (SEQ ID NO: 932)]

[CAAGGTCAGTCTGGATCC] [TCTTGCCTTATGCATCCGCATTATGCTCATGATTATTGCTATGTTG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAACCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 84)

[spacer (SEQ ID NO: 923)] [PL02-0003 LC (SEQ ID NO: 933)]

[QGQSGS] [SCLMHPHYAHDYCYVGGGSSGSGSGSGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 85)

PL02-0003 LC nucleotide sequence

TCTTGCCTTATGCATCCGCATTATGCTCATGATTATTGCTATGTTGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAACCTGC

TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 932)

PL02-0003 LC amino acid sequence

SCLMHPHYAHDYCYVGGSSGSGSGSGTSTSGRSANPRGGGSDIQMTQSPSSLASVGDRTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 933)

[spacer (SEQ ID NO: 929)] [PL03-0003 LC (SEQ ID NO: 934)]
[CAAGGTCAGTCTGGATCC] [TTGTGCGAGGTTTTGATGTTGTTGCAGCATCCGTGGTGCATGGGGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 86)

[spacer (SEQ ID NO: 923)] [PL03-0003 LC (SEQ ID NO: 935)]
[QQQSGS] [LCEVLMLLQHPWCMGGGSSGSGSGSGTSTSGRSANPRGGGSDIQMTQSPSSLASV
GDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 87)

PL03-0003 LC nucleotide sequence

TTGTGCGAGGTTTTGATGTTGTTGCAGCATCCGTGGTGCATGGGGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 934)

PL03-0003 LC amino acid sequence

LCEVLMLLQHPWCMGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVTTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 935)

[spacer (SEQ ID NO: 929)] [PL04-0003 LC (SEQ ID NO: 936)]

[CAAGGTCAGTCTGGATCC] [ATTGCGTGCCGGCATTATTTATGGAGCAGTTGCCGTTTTGCCATCATG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 88)

[spacer (SEQ ID NO: 923)] [PL04-0003 LC (SEQ ID NO: 937)]

[QGQSGS] [IACRHFMEQLPFCHHGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVTTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 89)

PL04-0003 LC nucleotide sequence

ATTGCGTGCCGGCATTATTTATGGAGCAGTTGCCGTTTTGCCATCATGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCCGGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 936)

PL04-0003 LC amino acid sequence

IACRHFMEQLPFCHHGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVTTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 937)

[spacer (SEQ ID NO: 929)] [PL05-0003 LC (SEQ ID NO: 938)]
[CAAGGTCAGTCTGGATCC] [TTTGGTCCTAGGTGCGGTGAGGCTTCTACTTGCGTTCCGTATGAGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 90)

[spacer (SEQ ID NO: 923)] [PL05-0003 LC (SEQ ID NO: 939)]
[QGQSGS] [FGPRCGEASTCVPYEGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 91)

PL05-0003 LC nucleotide sequence

TTTGGTCCTAGGTGCGGTGAGGCTTCTACTTGCGTTCCGTATGAGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 938)

PL05-0003 LC amino acid sequence

FGPRCGEASTCVPYEGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLASVGDRVITITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 939)

[spacer (SEQ ID NO: 929)] [PL06-0003 LC (SEQ ID NO: 940)]
[CAAGGTCAGTCTGGATCC] [ATTCTTTATTGCGATAGTTGGGGGCGGGGTGCTTGACGCGGCCGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC

CCGGCAAGGCCCCCAAAGTCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 92)

[spacer (SEQ ID NO: 923)] [PL06-0003 LC (SEQ ID NO: 941)]
[QGQSGS] [ILYCDSWGAGCLTRPGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 93)

PL06-0003 LC nucleotide sequence

ATTCTTTATTGCGATAGTTGGGGGGCGGGGTGCTTGACGCGGCCGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 940)

PL06-0003 LC amino acid sequence

ILYCDSWGAGCLTRPGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLASVGDRVITITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 941)

[spacer (SEQ ID NO: 929)] [PL07-0003 LC (SEQ ID NO: 942)]

[CAAGGTCAGTCTGGATCC] [GGGATTGCGTTGTGCCCGTCTCATTTTTGCCAGCTGCCTCAGACTG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 94)

[spacer (SEQ ID NO: 923)][PL07-0003 LC (SEQ ID NO: 943)]
[QGQSGS][GIALCPSHFCQLPQTGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 95)

PL07-0003 LC nucleotide sequence

GGGATTGCGTTGTGCCCCGTCTCATTTTTGCCAGCTGCCTCAGACTGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAACCTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCTTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 942)

PL07-0003 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVITITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 943)

[spacer (SEQ ID NO: 929)][PL08-0003 LC (SEQ ID NO: 944)]
[CAAGGTCAGTCTGGATCC][GATGGGCCGCGTTGCTTTGTGTGCGGGGAGTGCTCTCCGATTGGTG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGCAAGGCCCCCAAACCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCTTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 96)

[spacer (SEQ ID NO: 923)][PL08-0003 LC (SEQ ID NO: 945)]
[QGQSGS][DGPRCFVSGECSPIGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 97)

PL08-0003 LC nucleotide sequence

GATGGGCCGCGTTGCTTTGTGTCGGGGGAGTGCTCTCCGATTGGTGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 944)

PL08-0003 LC amino acid sequence

DGPRCFVSGECSPIGGGGSSGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRTTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 945)

[spacer (SEQ ID NO: 929)] [PL09-0003 LC (SEQ ID NO: 946)]

[CAAGGTCAGTCTGGATCC] [TTGTGCTATAAGCTGGATTATGATGATAGGTCTTATTGCCATATTG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 98)

[spacer (SEQ ID NO: 923)] [PL09-0003 LC (SEQ ID NO: 947)]

[QGQSGS] [LCYKLDYDDRSYCHIGGGSSGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 99)

PL09-0003 LC nucleotide sequence

TTGTGCTATAAGCTGGATTATGATGATAGGTCTTATTGCCATATTGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC

TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 946)

PL09-0003 LC amino acid sequence

LCYKLDYDDRSYCHIGGGSSGSGSGSGTSTSGRSANPRGGGSDIQMTQSPSSLASVGDRTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 947)

[spacer (SEQ ID NO: 929)] [PL10-0003 LC (SEQ ID NO: 948)]
[CAAGGTCAGTCTGGATCC] [CCGTGCCATCCGCATCCTTATGATGCTCGTCCTTATTGCAATGTGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 100)

[spacer (SEQ ID NO: 923)] [PL10-0003 LC (SEQ ID NO: 949)]
[QQQSGS] [PCHPHPYDARPYCNVGGSSGSGSGSGTSTSGRSANPRGGGSDIQMTQSPSSLASV
GDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 101)

PL10-0003 LC nucleotide sequence

CCGTGCCATCCGCATCCTTATGATGCTCGTCCTTATTGCAATGTGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 948)

PL10-0003 LC amino acid sequence

PCHPHPYDARPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVTTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 949)

[spacer (SEQ ID NO: 929)] [PL11-0003 LC (SEQ ID NO: 950)]

[CAAGGTCAGTCTGGATCC] [CCTTGCTATTGGCATCCTTTTTTTGCGTATAGGTATTGCAATACTG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAACCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 102)

[spacer (SEQ ID NO: 923)] [PL11-0003 LC (SEQ ID NO: 951)]

[QGQSGS] [PCYWHPFFAYRYCNTGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVTTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 103)

PL11-0003 LC nucleotide sequence

CCTTGCTATTGGCATCCTTTTTTTGCGTATAGGTATTGCAATACTGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCCGGCAAGGCCCCCAAACCTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 950)

PL11-0003 LC amino acid sequence

PCYWHPFFAYRYCNTGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVTTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 951)

[spacer (SEQ ID NO: 929)] [PL12-0003 LC (SEQ ID NO: 952)]
[CAAGGTCAGTCTGGATCC] [GTTTGCTATTATATGGATTGGTTGGGGCGGAATTGGTGCTCTTCGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 104)

[spacer (SEQ ID NO: 923)] [PL12-0003 LC (SEQ ID NO: 953)]
[QGQSGS] [VCYYMDWLGRNWCSSGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 105)

PL12-0003 LC nucleotide sequence

GTTTGCTATTATATGGATTGGTTGGGGCGGAATTGGTGCTCTTCGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 952)

PL12-0003 LC amino acid sequence

VCYYMDWLGRNWCSSGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVITITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 953)

[spacer (SEQ ID NO: 929)] [PL13-0003 LC (SEQ ID NO: 954)]
[CAAGGTCAGTCTGGATCC] [CTGTGCGATCTGTTTAAGTTGCGTGAGTTTCCTTATTGCATGGGGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC

CCGGCAAGGCCCCCAAAGTCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 106)

[spacer (SEQ ID NO: 923)] [PL13-0003 LC (SEQ ID NO: 955)]
[QGQSGS] [LCDLFLKREFPYCMGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 107)

PL13-0003 LC nucleotide sequence

CTGTGCGATCTGTTTAAGTTGCGTGAGTTTCCTTATTGCATGGGGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 954)

PL13-0003 LC amino acid sequence

LCDLFLKREFPYCMGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLASVGDRVITITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 955)

[spacer (SEQ ID NO: 929)] [PL14-0003 LC (SEQ ID NO: 956)]
[CAAGGTCAGTCTGGATCC] [TATCTTCCGTGCCATTTTGTTCGATTGGGGCTTGCAATAATAAGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 108)

[spacer (SEQ ID NO: 923)][PL14-0003 LC (SEQ ID NO: 957)]
[QGQSGS][YLPCHFVPIGACNNKGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGGTKVEIKR] (SEQ ID NO: 109)

PL14-0003 LC nucleotide sequence

TATCTTCCGTGCCATTTTGTTCGATTGGGGCTTGCAATAATAAGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAACCTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCTTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 956)

PL14-0003 LC amino acid sequence

YLPCHFVPIGACNNKGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVITITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQD
NGYPSTFGGGGTKVEIKR (SEQ ID NO: 957)

[spacer (SEQ ID NO: 929)][PL15-0003 LC (SEQ ID NO: 958)]
[CAAGGTCAGTCTGGATCC][ATTTTTTGGCATATGGGTGTTGTGGTTCCTCAGTGCGGAATTATG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAACCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCTTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 110)

[spacer (SEQ ID NO: 923)][PL15-0003 LC (SEQ ID NO: 959)]
[QGQSGS][IFCHMGVVVPQCANYGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGGTKVEIKR] (SEQ ID NO: 111)

PL15-0003 LC nucleotide sequence

ATTTTTTGCCATATGGGTGTTGTGGTTCCTCAGTGCGGAATTATGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 958)

PL15-0003 LC amino acid sequence

IFCHMGVVVPQCANYGGSSGSGSGSGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVITITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 959)

[spacer (SEQ ID NO: 929)] [PL16-0003 LC (SEQ ID NO: 960)]

[CAAGGTCAGTCTGGATCC] [GCGTGCCATCCGCATCCTTATGATGCTCGTCCTTATTGCAATGTGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTGCCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 112)

[spacer (SEQ ID NO: 923)] [PL16-0003 LC (SEQ ID NO: 961)]

[QGQSGS] [ACHPHYDARPYCNVGGSSGSGSGSGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 113)

PL16-0003 LC nucleotide sequence

GCGTGCCATCCGCATCCTTATGATGCTCGTCCTTATTGCAATGTGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC

TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 960)

PL16-0003 LC amino acid sequence

ACHPHPYDARPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVITITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 961)

[spacer (SEQ ID NO: 929)] [PL17-0003 LC (SEQ ID NO: 962)]
[CAAGGTCAGTCTGGATCC] [CCGTGCCATCCGGCTCCTTATGATGCTCGTCCTTATTGCAATGTGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 114)

[spacer (SEQ ID NO: 923)] [PL17-0003 LC (SEQ ID NO: 963)]
[QQQSGS] [PCHPAPYDARPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 115)

PL17-0003 LC nucleotide sequence

CCGTGCCATCCGGCTCCTTATGATGCTCGTCCTTATTGCAATGTGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 962)

PL17-0003 LC amino acid sequence

PCHPAPYDARPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVTTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 963)

[spacer (SEQ ID NO: 929)][PL18-0003 LC (SEQ ID NO: 964)]

[CAAGGTCAGTCTGGATCC][CCGTGCCATCCGCATGCTTATGATGCTCGTCCTTATTGCAATGTGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAACCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 116)

[spacer (SEQ ID NO: 923)][PL18-0003 LC (SEQ ID NO: 965)]

[QGQSGS][PCHPHAYDARPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVTTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 117)

PL18-0003 LC nucleotide sequence

CCGTGCCATCCGCATGCTTATGATGCTCGTCCTTATTGCAATGTGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCCGGCAAGGCCCCCAAACCTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 964)

PL18-0003 LC amino acid sequence

PCHPHAYDARPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVTTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 965)

[spacer (SEQ ID NO: 929)] [PL19-0003 LC (SEQ ID NO: 966)]
[CAAGGTCAGTCTGGATCC] [CCGTGCCATCCGCATCCTGCTGATGCTCGTCCTTATTGCAATGTGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 118)

[spacer (SEQ ID NO: 923)] [PL19-0003 LC (SEQ ID NO: 967)]
[QGQSGS] [PCHPHPADARPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 119)

PL19-0003 LC nucleotide sequence

CCGTGCCATCCGCATCCTGCTGATGCTCGTCCTTATTGCAATGTGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 966)

PL19-0003 LC amino acid sequence

PCHPHPADARPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVITITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 967)

[spacer (SEQ ID NO: 929)] [PL20-0003 LC (SEQ ID NO: 968)]
[CAAGGTCAGTCTGGATCC] [CCGTGCCATCCGCATCCTTATGCTGCTCGTCCTTATTGCAATGTGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC

CCGGCAAGGCCCCCAAAGTCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 120)

[spacer (SEQ ID NO: 923)] [PL20-0003 LC (SEQ ID NO: 969)]
[QGQSGS] [PCHPHPYAARPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 121)

PL20-0003 LC nucleotide sequence

CCGTGCCATCCGCATCCTTATGCTGCTCGTCCTTATTGCAATGTGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 968)

PL20-0003 LC amino acid sequence

PCHPHPYAARPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR (SEQ ID NO: 969)

[spacer (SEQ ID NO: 929)] [PL21-0003 LC (SEQ ID NO: 970)]

[CAAGGTCAGTCTGGATCC] [CCGTGCCATCCGCATCCTTATGATGCTGCTCCTTATTGCAATGTGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 122)

[spacer (SEQ ID NO: 923)][PL21-0003 LC (SEQ ID NO: 971)]
[QGQSGS][PCHPHPYDAAPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 123)

PL21-0003 LC nucleotide sequence

CCGTGCCATCCGCATCCTTATGATGCTGCTCCTTATTGCAATGTGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAACCTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCTTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 970)

PL21-0003 LC amino acid sequence

PCHPHPYDAAPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR (SEQ ID NO: 971)

[spacer (SEQ ID NO: 929)][PL22-0003 LC (SEQ ID NO: 972)]

[CAAGGTCAGTCTGGATCC][CCGTGCCATCCGCATCCTTATGATGCTCGTCCTGCTTGCAATGTGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGAAGGCCCCCAAACCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCTTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 124)

[spacer (SEQ ID NO: 923)][PL22-0003 LC (SEQ ID NO: 973)]

[QGQSGS][PCHPHPYDARPCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 125)

PL22-0003 LC nucleotide sequence

CCGTGCCATCCGCATCCTTATGATGCTCGTCCTGCTTGCAATGTGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAACCTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 972)

PL22-0003 LC amino acid sequence

PCHPHPYDARPAENVGGSSGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 973)

[spacer (SEQ ID NO: 929)] [PL23-0003 LC (SEQ ID NO: 974)]

[CAAGGTCAGTCTGGATCC] [CCGTGCCATCCGCATCCTTATGATGCTCGTCCTTATTGCGCTGTGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAACCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 126)

[spacer (SEQ ID NO: 923)] [PL23-0003 LC (SEQ ID NO: 975)]

[QGQSGS] [PCHPHPYDARPYCAVGGSSGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 127)

PL23-0003 LC nucleotide sequence

CCGTGCCATCCGCATCCTTATGATGCTCGTCCTTATTGCGCTGTGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAACCTGC

TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 974)

PL23-0003 LC amino acid sequence

PCHPHPYDARPYCAVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVITITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 975)

[spacer (SEQ ID NO: 929)] [PL24-0003 LC (SEQ ID NO: 976)]
[CAAGGTCAGTCTGGATCC] [CCGTGCCATGCGCATCCTTATGATGCTCGTCCTTATTGCAATGTGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 128)

[spacer (SEQ ID NO: 923)] [PL24-0003 LC (SEQ ID NO: 977)]
[QQQSGS] [PCHAHPYDARPYCNVGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 129)

PL24-0003 LC nucleotide sequence

CCGTGCCATGCGCATCCTTATGATGCTCGTCCTTATTGCAATGTGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 976)

PL24-0003 LC amino acid sequence

PCHAHPYDARPYCNVGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVTTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 977)

[spacer (SEQ ID NO: 929)] [PL25-0003 LC (SEQ ID NO: 978)]

[CAAGGTCAGTCTGGATCC] [CCGTGCCATCCGCATCCTTATGATGCTCGTGCTTATTGCAATGTGG
GAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCC
ACGTGGCGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGAC
AGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGC
CCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATT
TTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGGACAACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCA
AGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTG
T] (SEQ ID NO: 130)

[spacer (SEQ ID NO: 923)] [PL25-0003 LC (SEQ ID NO: 979)]

[QGQSGS] [PCHPHPYDARAYCNVGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASV
GDRVTTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED
FATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 131)

PL25-0003 LC nucleotide sequence

CCGTGCCATCCGCATCCTTATGATGCTCGTGCTTATTGCAATGTGGGAGGTGGCTCGAGCGGTGGCA
GCGGTGGCTCTGGTGGTACTAGCACCTCTGGTCGTTCCGCTAACCCACGTGGCGGCGGTTCTGACAT
CCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGA
GCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAGAAGCCCCGGCAAGGCCCCCAAAGTGC
TGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCAC
CGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGAC
AACGGCTACCCCAGCACCTTTGGCGGAGGTACCAAGGTGGAATCAAGCGTTGTGAGGCCACTCACA
AGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATGAGTGT (SEQ ID NO: 978)

PL25-0003 LC amino acid sequence

PCHPHPYDARAYCNVGGGSSGGSGSGGTSTSGRSANPRGGGSDIQMTQSPSSLSASVGDRVTTITCR
ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQD
NGYPSTFGGGTKVEIKR (SEQ ID NO: 979)

[spacer (SEQ ID NO: 929)][PL03-2001 LC (SEQ ID NO: 980)]
[CAAGGTCAGTCTGGATCC][GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCA
GTGGTCTGTTAAGCGGTCTAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAG
CAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGC
TACCTGAACTGGTATCAGCAGAAGCCCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTC
TGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAG
CTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTT
GGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTG
TCAAGAGCTTCAACAGGAATGAGTGT] (SEQ ID NO: 132)

[spacer (SEQ ID NO: 923)][PL03-2001 LC (SEQ ID NO: 981)]
[QGQSGS][LCEVLMLLQHPWCMGGGSSGGSGSGSGGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 133)

PL03-2001 LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCAGTGGTCTGTTAAGCGGTCTGTA
GCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGG
CGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAG
AAGCCCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCA
GATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTT
CGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAA
ATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATG
AGTGT (SEQ ID NO: 980)

PL03-2001 LC amino acid sequence

LCEVLMLLQHPWCMGGGSSGGSGSGSGGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 981)

[spacer (SEQ ID NO: 923)][PL04-2001 LC (SEQ ID NO: 1183)]
[QGQSGS][IACRHFMEQLPFCHHGGGSSGGSGSGSGGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1182)

PL04-2001 LC amino acid sequence

IACRHFMEQLPFCHHGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVTTITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1183)

[spacer (SEQ ID NO: 929)] [PL06-2001-mk LC (SEQ ID NO: 982)]

[CAAGGTCAGTCTGGATCC] [GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCA
GTGGTCTGTTAAGCGGTCTAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAG
CAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGC
TACCTGAACTGGTATCAGCAGAAGCCCGGCAAGGCCCCCAAAGTGTGATCTACGCCGCCAGCTCTC
TGCAGTCTGGCGTGCCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAG
CTCCCTGCAGCCCAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTT
GGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTG
TCAAGAGCTTCAACAGGAATGAGTGT] (SEQ ID NO: 134)

[spacer (SEQ ID NO: 923)] [PL06-2001-mk LC (SEQ ID NO: 983)]

[QQQSGS] [ILYCDSWGAGCLTRPGGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVTTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 135)

PL06-2001-mk LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCAGTGGTCTGTTAAGCGGTCTGTA
GCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGG
CGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAG
AAGCCCGGCAAGGCCCCCAAAGTGTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCA
GATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCAGGACTT
CGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAA
ATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATG
AGTGT (SEQ ID NO: 982)

PL06-2001-mk LC amino acid sequence

ILYCDSWGAGCLTRPGGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVTTITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 983)

[spacer (SEQ ID NO: 929)][PL07-2001 LC (SEQ ID NO: 984)]
[CAAGGTCAGTCTGGATCC][GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCA
GTGGTCTGTTAAGCGGTCTAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAG
CAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGC
TACCTGAACTGGTATCAGCAGAAGCCCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTC
TGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAG
CTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTT
GGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTG
TCAAGAGCTTCAACAGGAATGAGTGT] (SEQ ID NO: 136)

[spacer (SEQ ID NO: 923)][PL07-2001 LC (SEQ ID NO: 985)]
[QGQSGS][GIALCPSHFCQLPQTGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 137)

PL07-2001 LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCAGTGGTCTGTTAAGCGGTCTGTA
GCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGG
CGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAG
AAGCCCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCA
GATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTT
CGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAA
ATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATG
AGTGT (SEQ ID NO: 984)

PL07-2001 LC amino acid sequence

GIALCPSHFCQLPQTGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 985)

[spacer (SEQ ID NO: 929)][PL10-2001 LC (SEQ ID NO: 986)]
[CAAGGTCAGTCTGGATCC][GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCA
GTGGTCTGTTAAGCGGTCTAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAG
CAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGC
TACCTGAACTGGTATCAGCAGAAGCCCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTC
TGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAG

CTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTT
GGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTG
TCAAGAGCTTCAACAGGAATGAGTGT] (SEQ ID NO: 138)

[spacer (SEQ ID NO: 923)][PL10-2001 LC (SEQ ID NO: 987)]
[QGQSGS][PCHPHPYDARPYCNVGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 139)

PL10-2001 LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCAGTGGTCTGTTAAGCGGTCGTA
GCGATAATCATGGCGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGG
CGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAG
AAGCCCGGCAAGGCCCCCAAACTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCA
GATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTT
CGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAA
ATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATG
AGTGT (SEQ ID NO: 986)

PL10-2001 LC amino acid sequence

PCHPHPYDARPYCNVGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 987)

[spacer (SEQ ID NO: 923)][PL11-2001 LC (SEQ ID NO: 1185)]
[QGQSGS][PCYWHPFFAYRYCNTGGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1184)

PL11-2001 LC amino acid sequence

PCYWHPFFAYRYCNTGGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1185)

[spacer (SEQ ID NO: 923)][PL12-2001 LC (SEQ ID NO: 1187)]

[QGQSGS] [VCYYMDWLGRNWCSSGGGSSGGSGGGSGGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1186)

PL12-2001 LC amino acid sequence

VCYYMDWLGRNWCSSGGGSSGGSGGGSGGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1187)

[spacer (SEQ ID NO: 929)] [PL14-2001 LC (SEQ ID NO: 988)]
[CAAGGTCAGTCTGGATCC] [GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCA
GTGGTCTGTTAAGCGGTCGTAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAG
CAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGC
TACCTGAACTGGTATCAGCAGAAGCCCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTC
TGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAG
CTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTT
GGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTG
TCAAGAGCTTCAACAGGAATGAGTGT] (SEQ ID NO: 140)

[spacer (SEQ ID NO: 923)] [PL14-2001 LC (SEQ ID NO: 989)]
[QGQSGS] [YLPCHFVPIGACNNKGGGSSGGSGGGSGGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 141)

PL14-2001 LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCAGTGGTCTGTTAAGCGGTCGTA
GCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGG
CGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAG
AAGCCCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCA
GATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTT
CGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAA
ATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATG
AGTGT (SEQ ID NO: 988)

PL14-2001 LC amino acid sequence

YLPCHFVPIGACNNKGGGSSGGSGGSGGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVTTTC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 989)

[spacer (SEQ ID NO: 929)][PL15-2001 LC (SEQ ID NO: 990)]
[CAAGGTCAGTCTGGATCC][GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCA
GTGGTCTGTTAAGCGGTCTAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAG
CAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGC
TACCTGAACTGGTATCAGCAGAAGCCCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTC
TGCAGTCTGGCGTGCCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAG
CTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTT
GGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTG
TCAAGAGCTTCAACAGGAATGAGTGT] (SEQ ID NO: 142)

[spacer (SEQ ID NO: 923)][PL15-2001 LC (SEQ ID NO: 991)]
[QGQSGS][IFCHMGVVVPQCANYGGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 143)

PL15-2001 LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCAGTGGTCTGTTAAGCGGTCTGTA
GCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGG
CGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAG
AAGCCCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCAGCA
GATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTT
CGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAA
ATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATG
AGTGT (SEQ ID NO: 990)

PL15-2001 LC amino acid sequence

IFCHMGVVVPQCANYGGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 991)

[spacer (SEQ ID NO: 929)][PL18-2001 LC (SEQ ID NO: 992)]
[CAAGGTCAGTCTGGATCC][GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCA
GTGGTCTGTTAAGCGGTCTAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAG
CAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGC
TACCTGAACTGGTATCAGCAGAAGCCCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTC
TGCAGTCTGGCGTGCCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAG

CTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTT
GGCGGAGGTACCAAGGTGGAAATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTG
TCAAGAGCTTCAACAGGAATGAGTGT] (SEQ ID NO: 144)

[spacer (SEQ ID NO: 923)][PL18-2001 LC (SEQ ID NO: 993)]
[QGQSGS][PCHPHAYDARPYCNVGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 145)

PL18-2001 LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGGCTCTGGTGGTATTAGCAGTGGTCTGTTAAGCGGTCGTA
GCGATAATCATGGCGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGTGGG
CGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAGCAG
AAGCCCGGCAAGGCCCCCAAACTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCAGCA
GATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGACTT
CGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTGGAA
ATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGAATG
AGTGT (SEQ ID NO: 992)

PL18-2001 LC amino acid sequence

PCHPHAYDARPYCNVGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 993)

[spacer (SEQ ID NO: 923)][PL19-2001 LC (SEQ ID NO: 1189)]
[QGQSGS][PCHPHPADARPYCNVGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1188)

PL19-2001 LC amino acid sequence

PCHPHPADARPYCNVGGSSGGSGSGSGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1189)

[spacer (SEQ ID NO: 923)][PL20-2001 LC (SEQ ID NO: 1191)]

[QGQSGS] [PCHPHPYAARPYCNVGGGSSGGSGGGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1190)

PL20-2001 LC amino acid sequence

PCHPHPYAARPYCNVGGGSSGGSGGGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1191)

[spacer (SEQ ID NO: 929)] [PL03-1004/GG/0001 (also referred to
herein as PL03-3001) LC (SEQ ID NO: 994)]

[CAAGGTCAGTCTGGATCC] [GGAGGTGGCTCGAGCGGTGGCAGCGGTGCTGTGGGTCTCCTGGCTC
CCCCGGGCGGCCTGTCCGGCCGCAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCC
CAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGC
AGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAACCTGCTGATCTACGCCGCCAGCT
CTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCAT
CAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACC
TTTGGCGGAGGTACCAAGGTGGAATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCA
TTGTCAAGAGCTTCAACAGGAATGAGTGT] (SEQ ID NO: 146)

[spacer (SEQ ID NO: 923)] [PL03-1004/GG/0001 LC (SEQ ID NO: 995)]
[QGQSGS] [LCEVLMMLQHPWCMGGGSSGGSGAVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSLSA
SVGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 147)

PL03-1004/GG/0001 LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGCTGTGGGTCTCCTGGCTCCCCGGGCGGCCTGTCCGGCC
GCAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGT
GGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAG
CAGAAGCCCGCAAGGCCCCCAAACCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCA
GCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGA
CTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTG
GAAATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGA
ATGAGTGT (SEQ ID NO: 994)

PL03-1004/GG/0001 LC amino acid sequence

LCEVLMLLQHPWCMGGGGSSGGSGAVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSLSASVGDRVTIT
CRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQ
QDNGYPSTFGGGTKVEIKR (SEQ ID NO: 995)

[spacer (SEQ ID NO: 929)] [PL06-1004/GG/0001 (also referred to
herein as PL06-3001) LC (SEQ ID NO: 996)]

[CAAGGTCAGTCTGGATCC] [GGAGGTGGCTCGAGCGGTGGCAGCGGTGCTGTGGGTCTCCTGGCTC
CCCCGGGCGGCCTGTCCGGCCGAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCC
CAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGC
AGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCT
CTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCAT
CAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACC
TTTGGCGGAGGTACCAAGGTGGAATCAAGCGTTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCC
TCACAAAGAGCTTCAACAGGGGAGAGTGT] (SEQ ID NO: 148)

[spacer (SEQ ID NO: 923)] [PL06-1004/GG/0001 LC (SEQ ID NO: 997)]
[QQQSGS] [ILYCDSWGAGCLTRPGGGSSGGSGAVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSLSA
SVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQ
EDFATYYCQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 149)

PL06-1004/GG/0001 LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGCTGTGGGTCTCCTGGCTCCCCCGGGCGGCCTGTCCGGCC
GCAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGT
GGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAG
CAGAAGCCCGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCA
GCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGA
CTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTG
GAAATCAAGCGTTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGG
GAGAGTGT (SEQ ID NO: 996)

PL06-1004/GG/0001 LC amino acid sequence

ILYCDSWGAGCLTRPGGGSSGGSGAVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSLSASVGDRVTIT
CRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQ
QDNGYPSTFGGGTKVEIKR (SEQ ID NO: 997)

[spacer (SEQ ID NO: 929)] [PL07-1004/GG/0001 (also referred to herein as PL07-3001) LC (SEQ ID NO: 998)]
[CAAGGTCAGTCTGGATCC] [GGAGGTGGCTCGAGCGGTGGCAGCGGTGCTGTGGGTCTCCTGGCTC
CCCCGGGCGGCCTGTCCGGCCGACGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCC
CAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGC
AGCTACCTGAACTGGTATCAGCAGAAGCCCGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCT
CTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCAT
CAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACC
TTTGGCGGAGGTACCAAGGTGAAATCAAGCGTTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCC
TCACAAAGAGCTTCAACAGGGGAGAGTGT] (SEQ ID NO: 150)

[spacer (SEQ ID NO: 923)] [PL01-1004/GG/0001 LC (SEQ ID NO: 999)]
[QGQSGS] [GIALCPSHFCQLPQTGGGSSGGSGAVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSLSA
SVGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 151)

PL07-1004/GG/0001 LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGCTGTGGGTCTCCTGGCTCCCCGGGCGGCCTGTCCGGCC
GCAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGT
GGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAG
CAGAAGCCCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCA
GCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGA
CTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTG
GAAATCAAGCGTTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGG
GAGAGTGT (SEQ ID NO: 998)

PL07-1004/GG/0001 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGAVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITIT
CRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQ
QDNGYPSTFGGGTKVEIKR (SEQ ID NO: 999)

[spacer (SEQ ID NO: 929)] [PL14-1004/GG/0001 (also referred to herein as PL14-3001) LC (SEQ ID NO: 1000)]

[CAAGGTCAGTCTGGATCC] [GGAGGTGGCTCGAGCGGTGGCAGCGGTGCTGTGGGTCTCCTGGCTC
CCCCGGGCGGCCTGTCCGGCCGACGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCC
CAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGC

AGCTACCTGAACTGGTATCAGCAGAAGCCCGGCAAGGCCCCCAAAGTCTGATCTACGCCGCCAGCT
CTCTGCAGTCTGGCGTGCCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCAT
CAGCTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACC
TTTGGCGGAGGTACCAAGGTGGAAATCAAGCGTTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCG
TCACAAAGAGCTTCAACAGGGGAGAGTGT] (SEQ ID NO: 152)

[spacer (SEQ ID NO: 923)][PL14-1004/GG/0001 LC (SEQ ID NO: 1001)]
[QGQSGS][YLPCHFVPIGACNNKGGSSGGSGAVGLLAPPGGLSGRSDNHGGSIDIQMTQSPSSLSA
SVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 153)

PL14-1004/GG/0001 LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGCTGTGGGTCTCCTGGCTCCCCGGGCGGCCTGTCCGGCC
GCAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGT
GGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAG
CAGAAGCCCCGCAAGGCCCCCAAAGTGTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCA
GCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCGAGGA
CTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTG
GAAATCAAGCGTTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGG
GAGAGTGT (SEQ ID NO: 1000)

PL14-1004/GG/0001 LC amino acid sequence

YLPCHFVPIGACNNKGGSSGGSGAVGLLAPPGGLSGRSDNHGGSIDIQMTQSPSSLSASVGDRVTIT
CRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQ
QDNGYPSTFGGGTKVEIKR (SEQ ID NO: 1001)

[spacer (SEQ ID NO: 929)][PL15-1004/GG/0001 (also referred to
herein as PL15-3001) LC (SEQ ID NO: 1002)]

[CAAGGTCAGTCTGGATCC][GGAGGTGGCTCGAGCGGTGGCAGCGGTGCTGTGGGTCTCCTGGCTC
CCCCGGGCGGCCTGTCCGGCCGCAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCC
CAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGC
AGCTACCTGAACTGGTATCAGCAGAAGCCCCGCAAGGCCCCCAAAGTGTGATCTACGCCGCCAGCT
CTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCAT
CAGCTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCC
TTTGGCGGAGGTACCAAGGTGGAATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCA
TTGTCAAGAGCTTCAACAGGAATGAGTGT] (SEQ ID NO: 154)

[spacer (SEQ ID NO: 923)][PL15-1004/GG/0001 LC (SEQ ID NO: 1003)]
[QGQSGS][IFCHMGVVVPQCANYGGSSGGSGAVGLLAPPGGLSGRSDNHGGSIDIQMTQSPSSLSA
SVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 155)

PL15-1004/GG/0001 LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGCTGTGGGTCTCCTGGCTCCCCGGGCGGCCTGTCCGGCC
GCAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGT
GGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAG
CAGAAGCCCCGGCAAGGCCCCCAAACCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCA
GCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCATCAGCTCCCTGCAGCCCCGAGGA
CTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTG
GAAATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGA
ATGAGTGT (SEQ ID NO: 1002)

PL15-1004/GG/0001 LC amino acid sequence

IFCHMGVVVPQCANYGGSSGGSGAVGLLAPPGGLSGRSDNHGSDIQMTQSPSSLSASVGDRVITIT
CRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQ
QDNGYPSTFGGGTKVEIKR (SEQ ID NO: 1003)

[spacer (SEQ ID NO: 929)] [PL18-1004/GG/0001 (also referred to
herein as PL18-3001) LC (SEQ ID NO: 1004)]

[CAAGGTCAGTCTGGATCC] [GGAGGTGGCTCGAGCGGTGGCAGCGGTGCTGTGGGTCTCCTGGCTC
CCCCGGGCGGCCTGTCCGGCCGAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCC
CAGCAGCCTGTCTGCTAGCGTGGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGC
AGCTACCTGAACTGGTATCAGCAGAAGCCCCGGCAAGGCCCCCAAACCTGCTGATCTACGCCGCCAGCT
CTCTGCAGTCTGGCGTGCCAGCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCCTGACCAT
CAGCTCCCTGCAGCCGAGGACTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACC
TTTGGCGGAGGTACCAAGGTGAAATCAAGCGTTGTGAGGCCACTCACAAGACATCAACTTCACCCA
TTGTCAAGAGCTTCAACAGGAATGAGTGT] (SEQ ID NO: 156)

[spacer (SEQ ID NO: 923)] [PL18-1004/GG/0001 LC (SEQ ID NO: 1005)]

[QGQSGS] [PCHPHAYDARPYCNVGGSSGGSGAVGLLAPPGGLSGRSDNHGSDIQMTQSPSSLSA
SVGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 157)

PL18-1004/GG/0001 LC nucleotide sequence

GGAGGTGGCTCGAGCGGTGGCAGCGGTGCTGTGGGTCTCCTGGCTCCCCGGGCGGCCTGTCCGGCC
GCAGCGATAATCATGGCGGTTCTGACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCTAGCGT
GGGCGACAGAGTGACCATCACCTGTAGAGCCAGCCAGAGCATCAGCAGCTACCTGAACTGGTATCAG
CAGAAGCCCCGGCAAGGCCCCCAAACCTGCTGATCTACGCCGCCAGCTCTCTGCAGTCTGGCGTGCCCA

GCAGATTTTCCGGCAGCGGCTCTGGCACCGACTTCACCTGACCATCAGCTCCCTGCAGCCCGAGGA
CTTCGCCACCTACTACTGCCAGCAGGACAACGGCTACCCAGCACCTTTGGCGGAGGTACCAAGGTG
GAAATCAAGCGTTGTGAGGCCACTACAAGACATCAACTTCACCCATTGTCAAGAGCTTCAACAGGA
ATGAGTGT (SEQ ID NO: 1004)

PL18-1004/GG/0001 LC amino acid sequence

PCHPHAYDARPYCNVGGSSGGSGAVGLLAPPGGLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITIT
CRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQ
QDNGYPSTFGGGTKVEIKR (SEQ ID NO: 1005)

PL07-0001

[spacer (SEQ ID NO: 923)][PL07-0001 LC (SEQ ID NO: 1145)]
[QGQSGS][GIALCPSHFCQLPQTGGGSSGGSGGSLSGRSDNHGGSDIQMTQSPSSLSASVGDRV
TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATY
YCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1144)

PL07-0001 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGGSLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITITCRASQ
SISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQDNGYP
STFGGGTKVEIKR (SEQ ID NO: 1145)

PL07-0002

[spacer (SEQ ID NO: 923)][PL07-0002 LC (SEQ ID NO: 1147)]
[QGQSGS][GIALCPSHFCQLPQTGGGSSGGSGGSLSGRSGNHGGSDIQMTQSPSSLSASVGDRV
TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATY
YCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1146)

PL07-0002 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGGSLSGRSGNHGGSDIQMTQSPSSLSASVGDRVITITCRASQ
SISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQDNGYP
STFGGGTKVEIKR (SEQ ID NO: 1147)

PL07-1001

[spacer (SEQ ID NO: 923)][PL07-1001 LC (SEQ ID NO: 1149)]

[QGQSGS] [GIALCPSHFCQLPQTGGGSSGGSGSGSGISSGLLSSGGSDIQMTQSPSSLSASVGDRV
TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATY
YCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1148)

PL07-1001 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGSGISSGLLSSGGSDIQMTQSPSSLSASVGDRV
TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATY
YCQQDNGYPSTFGGGTKVEIKR (SEQ ID NO: 1149)

PL07-1002

[spacer (SEQ ID NO: 923)] [PL07-1002 LC (SEQ ID NO: 1151)]
[QGQSGS] [GIALCPSHFCQLPQTGGGSSGGSGSGGQNQALRMAGGSDIQMTQSPSSLSASVGDRV
TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATY
YCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1150)

PL07-1002 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGGQNQALRMAGGSDIQMTQSPSSLSASVGDRV
TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATY
YCQQDNGYPSTFGGGTKVEIKR (SEQ ID NO: 1151)

PL07-1003

[spacer (SEQ ID NO: 923)] [PL07-1003 LC (SEQ ID NO: 1153)]
[QGQSGS] [GIALCPSHFCQLPQTGGGSSGGSGSGGVHMPGLGFLPGGSDIQMTQSPSSLSASVGD
RVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFA
TYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1152)

PL07-1003 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGGVHMPGLGFLPGGSDIQMTQSPSSLSASVGDRV
TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATY
YCQQDNGYPSTFGGGTKVEIKR (SEQ ID NO: 1153)

PL07-1004

[spacer (SEQ ID NO: 923)] [PL07-1004 LC (SEQ ID NO: 1201)]
[QGQSGS] [GIALCPSHFCQLPQTGGGSSGGSGSGGAVGLLAPPGGSDIQMTQSPSSLSASVGDRV
TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATY
YCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1200)

PL07-1004 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGGAVGLLAPPGGSDIQMTQSPSSLSASVGDRVTITCRASQS
ISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQDNGYP
STFGGGTKVEIKR (SEQ ID NO: 1201)

PL07-2002

[spacer (SEQ ID NO: 923)][PL07-2002 LC (SEQ ID NO: 1155)]
[QGQSGS][GIALCPSHFCQLPQTGGGSSGGSGSGGISSGLLSGRSGNHGGSDIQMTQSPSSLSAS
VGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1154)

PL07-2002 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGGISSGLLSGRSGNHGGSDIQMTQSPSSLSASVGDRVTITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1155)

PL07-2003

[spacer (SEQ ID NO: 923)][PL07-2003 LC (SEQ ID NO: 1157)]
[QGQSGS][GIALCPSHFCQLPQTGGGSSGGSGSGGISSGLLSGRSANPRGGGSDIQMTQSPSSLS
ASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQ
PEDFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1156)

PL07-2003 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGGISSGLLSGRSANPRGGGSDIQMTQSPSSLSASVGDRVTI
TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYC
QQDNGYPSTFGGGTKVEIKR (SEQ ID NO: 1157)

PL07-2004

[spacer (SEQ ID NO: 923)][PL07-2004 LC (SEQ ID NO: 1159)]
[QGQSGS][GIALCPSHFCQLPQTGGGSSGGSGSGGAVGLLAPPTSGRSANPRGGGSDIQMTQSPS
SLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTIS
SLQPEDFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1158)

PL07-2004 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGGSGGAVGLLAPPTSGRSANPRGGGSDIQMTQSPSSLASVGDR
VTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFAT
YYCQQDNGYPSTFGGGTKVEIKR (SEQ ID NO: 1159)

PL07-2005

[spacer (SEQ ID NO: 923)][PL07-2005 LC (SEQ ID NO: 1161)]
[QGQSGS][GIALCPSHFCQLPQTGGGSSGGSGSGGAVGLLAPPSGRSANPRGGGSDIQMTQSPSS
LSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISS
LQPEDFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1160)

PL07-2005 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGGAVGLLAPPSGRSANPRGGGSDIQMTQSPSSLSASVGDRV
TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATY
YCQQDNGYPSTFGGGTKVEIKR (SEQ ID NO: 1161)

PL07-2006

[spacer (SEQ ID NO: 923)][PL07-2006 LC (SEQ ID NO: 1163)]
[QGQSGS][GIALCPSHFCQLPQTGGGSSGGSGSGGISSGLLSGRSDDHGGSDIQMTQSPSSLSAS
VGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1162)

PL07-2006 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGGISSGLLSGRSDDHGGSDIQMTQSPSSLSASVGDRVTITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1163)

PL07-2007

[spacer (SEQ ID NO: 923)][PL07-2007 LC (SEQ ID NO: 1165)]
[QGQSGS][GIALCPSHFCQLPQTGGGSSGGSGSGGISSGLLSGRSDIHGGSDIQMTQSPSSLSAS
VGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1164)

PL07-2007 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGGISSGLLSGRSDIHGGSDIQMTQSPSSLSASVGDRVTITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1165)

PL07-2008

[spacer (SEQ ID NO: 923)][PL07-2008 LC (SEQ ID NO: 1167)]

[QGQSGS] [GIALCP SHFCQLPQTGGGSSGGSGSGSGGISSGLLSGRSDQHGGSDIQMTQSPSSLSAS
VGDRV TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1166)

PL07-2008 LC amino acid sequence

GIALCP SHFCQLPQTGGGSSGGSGSGSGGISSGLLSGRSDQHGGSDIQMTQSPSSLSASVGDRV TITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1167)

PL07-2009

[spacer (SEQ ID NO: 923)][PL07-2009 LC (SEQ ID NO: 1169)]

[QGQSGS] [GIALCP SHFCQLPQTGGGSSGGSGSGSGGISSGLLSGRSDTHGGSDIQMTQSPSSLSAS
VGDRV TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1168)

PL07-2009 LC amino acid sequence

GIALCP SHFCQLPQTGGGSSGGSGSGSGGISSGLLSGRSDTHGGSDIQMTQSPSSLSASVGDRV TITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1169)

PL07-2010

[spacer (SEQ ID NO: 923)][PL07-2010 LC (SEQ ID NO: 1171)]

[QGQSGS] [GIALCP SHFCQLPQTGGGSSGGSGSGSGGISSGLLSGRSDYHGGSDIQMTQSPSSLSAS
VGDRV TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1170)

PL07-2010 LC amino acid sequence

GIALCP SHFCQLPQTGGGSSGGSGSGSGGISSGLLSGRSDYHGGSDIQMTQSPSSLSASVGDRV TITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1171)

PL07-2011

[spacer (SEQ ID NO: 923)][PL07-2011 LC (SEQ ID NO: 1173)]
[QGQSGS][GIALCPSHFCQLPQTGGGSSGGSGSGSGISSGLLSGRSDNPGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1172)

PL07-2011 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGSGISSGLLSGRSDNPGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1173)

PL07-2012

[spacer (SEQ ID NO: 923)][PL07-2012 LC (SEQ ID NO: 1175)]
[QGQSGS][GIALCPSHFCQLPQTGGGSSGGSGSGSGISSGLLSGRSANPGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1174)

PL07-2012 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGSGISSGLLSGRSANPGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1175)

PL07-2013

[spacer (SEQ ID NO: 923)][PL07-2013 LC (SEQ ID NO: 1177)]
[QGQSGS][GIALCPSHFCQLPQTGGGSSGGSGSGSGISSGLLSGRSANIGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1176)

PL07-2013 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGSGISSGLLSGRSANIGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1177)

PL07-2014

[spacer (SEQ ID NO: 923)] [PL07-2014 LC (SEQ ID NO: 1179)]
[QGQSGS] [GIALCPSHFCQLPQTGGGSSGGSGSGSGISSGLLSGRSDNIGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1178)

PL07-2014 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGSGISSGLLSGRSDNIGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKR (SEQ ID NO: 1179)

PL07-1004/GG/0003

[spacer (SEQ ID NO: 923)] [PL07-1004/GG/0003 LC (SEQ ID NO: 1181)]
[QGQSGS] [GIALCPSHFCQLPQTGGGSSGGSGSGSGGAVGLLAPPGGTSTSGRSANPRGGGSDIQMT
QSPSSLSASVGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFT
LTISSLQPEDFATYYCQQDNGYPSTFGGGTKVEIKR] (SEQ ID NO: 1180)

PL07-1004/GG/0003 LC amino acid sequence

GIALCPSHFCQLPQTGGGSSGGSGSGSGGAVGLLAPPGGTSTSGRSANPRGGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKR (SEQ ID NO: 1181)

EXAMPLE 11. Anti-PDL1 activatable antibodies of the disclosure

[000609] This example demonstrates that anti-PDL1 activatable antibodies can be made in a variety of combinations of MM, CM, VL, and VH domains as well as in a variety of distinct human isotypes.

Table 18. Anti-PDL1 Activatable Antibody Components

Mask Sequence (MM)	Substrate Sequence (CM)	VL	VH
YCEVSELFVLPWCMG (SEQ ID NO: 208)	LSGRSDNH (SEQ ID NO: 341)	SEQ ID NO: 12	SEQ ID NO: 14

SCLMHPHYAHDYCYV (SEQ ID NO: 426)	TGRGPSWV (SEQ ID NO: 338)	SEQ ID NO: 58	SEQ ID NO: 16
LCEVLMLLQHPWCMG (SEQ ID NO: 59)	PLTGRSGG (SEQ ID NO: 344)		SEQ ID NO: 18
IACRHFMEQLPFCCH (SEQ ID NO: 60)	TARGPSFK (SEQ ID NO: 340)		SEQ ID NO: 20
FGPRCGEASTCVPYE (SEQ ID NO: 61)	NTLSGRSENHSG (SEQ ID NO: 435)		SEQ ID NO: 22
ILYCDSWGAGCLTRP (SEQ ID NO: 62)	NTLSGRSGNHGS (SEQ ID NO: 436)		SEQ ID NO: 24
GIALCPSHFCQLPQT (SEQ ID NO: 63)	TSTSGRSANPRG (SEQ ID NO: 437)		SEQ ID NO: 26
DGPRCFVSGECSPIG (SEQ ID NO: 64)	TSGRSANP (SEQ ID NO: 438)		SEQ ID NO: 28
LCYKLDYDDRSYCHI (SEQ ID NO: 65)	VHMPLGFLGP (SEQ ID NO: 352)		SEQ ID NO: 30
PCHPHPYDARPYCNV (SEQ ID NO: 66)	AVGLLAPP (SEQ ID NO: 372)		SEQ ID NO: 32
PCYWHPFFAYRYCNT (SEQ ID NO: 67)	AQNLLGMV (SEQ ID NO: 360)		SEQ ID NO: 34
VCYYMDWLGRNWCSS (SEQ ID NO: 68)	QNQALRMA (SEQ ID NO: 359)		SEQ ID NO: 36
LCDLFLKLREFPYCMG (SEQ ID NO: 69)	LAAPLGLL (SEQ ID NO: 371)		SEQ ID NO: 38
YLPCHFVPIGACNNK (SEQ ID NO: 70)	STFPFGMF (SEQ ID NO: 361)		SEQ ID NO: 40
IFCHMGVVVPQCANY (SEQ ID NO: 71)	ISSGLLSS (SEQ ID NO: 364)		SEQ ID NO: 42
ACHPHPYDARPYCNV (SEQ ID NO: 72)	PAGLWLDP (SEQ ID NO: 374)		SEQ ID NO: 44
PCHPAPYDARPYCNV (SEQ ID NO: 73)	VAGRSMRP (SEQ ID NO: 439)		SEQ ID NO: 46
PCHPHAYDARPYCNV (SEQ ID NO: 74)	VVPEGRRS (SEQ ID NO: 440)		SEQ ID NO: 48

PCHPHPADARPYCNV (SEQ ID NO: 75)	ILPRSPAF (SEQ ID NO: 441)		SEQ ID NO: 50
PCHPHPYAARPYCNV (SEQ ID NO: 76)	MVLGRSLI (SEQ ID NO: 442)		SEQ ID NO: 52
PCHPHPYDAAPYCNV (SEQ ID NO: 77)	QGRAITFI (SEQ ID NO: 443)		SEQ ID NO: 54
PCHPHPYDARPAACNV (SEQ ID NO: 78)	SPRSIMLA (SEQ ID NO: 444)		SEQ ID NO: 56
PCHPHPYDARPYCAV (SEQ ID NO: 79)	SMLRSMPL (SEQ ID NO: 445)		
PCHAHPYDARPYCNV (SEQ ID NO: 80)	ISSGLLSGRSDNH (SEQ ID NO: 377)		
PCHPHPYDARAYCNV (SEQ ID NO: 81)	AVGLLAPPGGLSGRSDNH (SEQ ID NO: 383)		
	ISSGLLSSGGSGGSLSGRSDNH (SEQ ID NO: 378)		
	LSGRSGNH (SEQ ID NO: 883)		
	SGRSANPRG (SEQ ID NO: 884)		
	LSGRSDDH (SEQ ID NO: 885)		
	LSGRSDIH (SEQ ID NO: 886)		
	LSGRSDQH (SEQ ID NO: 887)		
	LSGRSDTH (SEQ ID NO: 888)		
	LSGRSDYH (SEQ ID NO: 889)		
	LSGRSDNP (SEQ ID NO: 890)		
	LSGRSANP (SEQ ID NO: 891)		

	LSGRSANI (SEQ ID NO: 892)		
	LSGRSDNI (SEQ ID NO: 893)		
	MIAPVAYR (SEQ ID NO: 894)		
	RPSPMWAY (SEQ ID NO: 895)		
	WATPRPMR (SEQ ID NO: 896)		
	FRLLDWQW (SEQ ID NO: 897)		
	ISSGL (SEQ ID NO: 898)		
	ISSGLLS (SEQ ID NO: 899)		
	ISSGLL (SEQ ID NO: 900)		
	ISSGLLSGRSANPRG (SEQ ID NO: 901)		
	AVGLLAPPTSGRSANPRG (SEQ ID NO: 902)		
	AVGLLAPPSGRSANPRG (SEQ ID NO: 903)		
	ISSGLLSGRSDDH (SEQ ID NO: 904)		
	ISSGLLSGRSDIH (SEQ ID NO: 905)		
	ISSGLLSGRSDQH (SEQ ID NO: 906)		
	ISSGLLSGRSDTH (SEQ ID NO: 907)		
	ISSGLLSGRSDYH (SEQ ID NO: 908)		

	ISSGLLSGRSDNP (SEQ ID NO: 909)		
	ISSGLLSGRSANP (SEQ ID NO: 910)		
	ISSGLLSGRSANI (SEQ ID NO: 911)		
	AVGLLAPPGGLSGRSDDH (SEQ ID NO: 912)		
	AVGLLAPPGGLSGRSDIH (SEQ ID NO: 913)		
	AVGLLAPPGGLSGRSDQH (SEQ ID NO: 914)		
	AVGLLAPPGGLSGRSDTH (SEQ ID NO: 915)		
	AVGLLAPPGGLSGRSDYH (SEQ ID NO: 916)		
	AVGLLAPPGGLSGRSDNP (SEQ ID NO: 917)		
	AVGLLAPPGGLSGRSANP (SEQ ID NO: 918)		
	AVGLLAPPGGLSGRSANI (SEQ ID NO: 919)		
	ISSGLLSGRSDNI (SEQ ID NO: 920)		
	AVGLLAPPGGLSGRSDNI (SEQ ID NO: 921)		
	GLSGRSDNHGGAVGLLAPP (SEQ ID NO: 1009)		
	GLSGRSDNHGGVHMPLGFLGP (SEQ ID NO: 1010)		

[000610] Any of the combinations described in Table 18 can be combined with human immunoglobulin constant regions to result in fully human IgGs including IgG1, IgG2, IgG4 or mutated constant regions to result in human IgGs with altered functions such as IgG1 N297A, IgG1 N297Q, or IgG4 S228P. The combinations described in Table 18 are not

limited by the particular combinations shown in any given row and include any mask sequence from column 1 matched with any substrate sequence from column 2 matched with any VL sequence from column 3 matched with any VH sequence from column 4. In addition to the substrate sequences disclosed in column 2, any CM disclosed herein can be used.

[000611] As an example, a spacer sequence (SEQ ID NO: 923) and Mask SEQ ID NO: 63 can be combined with substrate ISSGLLSGRSDNH (SEQ ID NO: 377), VL SEQ ID NO: 58, and combined with human kappa constant domain to give SEQ ID NO: 428; or Mask SEQ ID NO: 63 can be combined with substrate ISSGLLSGRSDNH (SEQ ID NO: 377), VL SEQ ID NO: 58, and combined with human kappa constant domain to give SEQ ID NO: 1008. Furthermore, VH SEQ ID NO: 46 can be combined with human immunoglobulin heavy chain constant domains to give human IgG1 (SEQ ID NO: 430), mutated human IgG4 S228P (SEQ ID NO: 432), mutated human IgG1 N297A (SEQ ID NO: 434), or mutated human IgG1 N297Q (SEQ ID NO: 1202). Co-expression of SEQ ID NO: 427 with SEQ ID NO: 429 will yield a fully human IgG1 anti-PDL1 activatable antibody. Co-expression of SEQ ID NO: 427 with SEQ ID NO: 431 will yield a fully human IgG4S228P anti-PDL1 activatable antibody. Co-expression of SEQ ID NO: 427 with SEQ ID NO: 433 will yield a fully human IgG1 N297A anti-PDL1 activatable antibody. Co-expression of SEQ ID NO: 427 with a nucleic acid sequence encoding the amino acid sequence of SEQ ID NO: 1202 will yield a fully human IgG1 N297Q anti-PDL1 activatable antibody.

[Light chain sequence (SEQ ID NO: 1007) with spacer (SEQ ID NO: 1006)] (SEQ ID NO: 427)

[CAGGGCCAGTCCGGCTCA] [TATCTGCCCTGCCACTTCGTGCCAATCGGGGCCTGTAACAATAAGG
GCGGTGGATCTAGTGGTGGCTCAGGCGGGTCTGGCGGCATTTCCAGTGGACTCTTGTGAGGACGATC
CGATAATCATGGCGGGTCCGACATCCAGATGACACAGAGCCCTTCTTCCCTCTCCGCAAGCGTTGGC
GACAGGGTCACCATTACCTGTAGGGCTTCTCAGAGCATCTCAAGCTATCTGAACTGGTACCAGCAGA
AACCTGGAAAGGCTCCAAAAGTCTGATTTACGCTGCCTCCAGTCTTCAGTCAGGCGTCCCCTCCAG
ATTTAGCGGATCAGGTAGTGGAACTGATTTTACCCTTACAATATCTTCTCTGCAGCCAGAGGACTTC
GCCACATACTATTGTGAGCAAGACAATGGTTACCCAGTACATTTGGCGGAGGGACAAAGGTGCGAGA
TCAAAAGGACCGTAGCAGCACCAAGCGTCTTTATTTTCCCCCCCAGTGACGAACAGCTGAAGAGCGG
AACAGCTTCAGTGGTGTGTCTCCTGAATAACTTCTATCCACGCGAGGCAAAGGTGCAGTGGAAGGTG
GACAATGCACTGCAGTCTGGTAATTCCCAAGAAAGTGTTACTGAGCAGGATTCCAAGGATTCAACTT

ACTCTCTGTCTAGCACCTGACTCTTTCTAAAGCAGATTATGAGAAGCATAAGGTCTACGCTTGCGA
GGTGACCCACCAGGGGCTTTCCTCTCCAGTTACCAAGTCATTCAACCGGGGTGAGTGTTGATGAGAA
TTC] (SEQ ID NO: 427)

Light chain sequence without spacer

TATCTGCCCTGCCACTTCGTGCCAATCGGGGCTGTAACAATAAGGGCGGTGGATCTAGTGGTGGCT
CAGGCGGGTCTGGCGGCATTTCCAGTGGACTCTTGTGAGGACGATCCGATAATCATGGCGGGTCCGA
CATCCAGATGACACAGAGCCCTTCTTCCCTCTCCGCAAGCGTTGGCGACAGGGTCACCATTACCTGT
AGGGCTTCTCAGAGCATCTCAAGCTATCTGAACTGGTACCAGCAGAAACCTGGAAAGGCTCCAAAC
TGCTGATTTACGCTGCCTCCAGTCTTCAGTCAGGCGTCCCCTCCAGATTTAGCGGATCAGGTAGTGG
AACTGATTTTACCCTTACAATATCTTCTCTGCAGCCAGAGGACTTCGCCACATACTATTGTGAGCAA
GACAATGGTTACCCAGTACATTTGGCGGAGGGACAAAGGTGAGATCAAAGGACCGTAGCAGCAC
CAAGCGTCTTTATTTTCCCCCAGTGACGAACAGCTGAAGAGCGGAACAGCTTCAGTGGTGTGTCT
CCTGAATAACTTCTATCCACGCGAGGCAAAGGTGAGTGGAAGGTGGACAATGCACTGCAGTCTGGT
AATTTCCAAGAAAGTGTTACTGAGCAGGATTCCAAGGATTCAACTTACTCTCTGTCTAGCACCTGA
CTCTTTCTAAAGCAGATTATGAGAAGCATAAGGTCTACGCTTGCGAGGTGACCCACCAGGGGCTTTC
CTCTCCAGTTACCAAGTCATTCAACCGGGGTGAGTGTTGATGAGAATTC (SEQ ID NO: 1007)

[Light chain sequence (SEQ ID NO: 1008) with spacer (SEQ ID
NO: 923)] (SEQ ID NO: 428)

[QGQSGS] [GIALCPHFQQLPQTGGGSSGGSGGGISSGLLSGRSDNHGGSDIQMTQSPSSLSAS
VGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE
DFATYYCQQDNGYPSTFGGGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW
KVDNALQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC]
(SEQ ID NO: 428)

Light chain sequence without spacer

GIALCPHFQQLPQTGGGSSGGSGGGISSGLLSGRSDNHGGSDIQMTQSPSSLSASVGDRVITITC
RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQ
DNGYPSTFGGGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSG
NSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID
NO: 1008)

SEQ ID NO: 429

GAAGTGCAGCTGCTCGAAAGCGGCGGAGGCTTGGTGCAGCCAGGAGGGAGCCTGCGACTGTCTTGCG
CAGCCAGCGGATTCACCTTCTCTTCTATGCCATGAGCTGGGTTCGACAGGCACCCGGCAAAGGTCT

CGAGTGGGTGTCTAGCATCTGGCGAAACGGAATAGTTACAGTGTATGCCGATAGCGTGAAGGGTCGC
TTTACTATTTACAGGGATAATTCTAAGAACACCCTCTACCTGCAAATGAATAGCCTTAGGGCAGAAG
ATACCGCCGTGTACTACTGTGCCAAATGGTCCGCAGCCTTTGACTACTGGGGCCAGGGGACACTGGT
GACCGTGTCTCTGTCATCAACCAAGGGGCCATCAGTGTTCCTACTCGCCCCATCTTCCAAGAGTACT
TCCGGCGGAACCGCAGCCCTTGGCTGCCTTGTTAAGGACTATTTCCAGAACCCTGACCGTAAGTT
GGAACTCTGGCGCCCTTACTTCTGGGGTGCACACCTTCCCAGCAGTGTTCAGTCCAGTGGCCTTTA
CTCTCTGTCTAGTGTAGTGACTGTGCCTTCCTCTAGTCTCGGTACCCAGACCTATATTTGTAATGTT
AACCATAAGCCCAGCAATACAAAGGTTGATAAGAAAGTGGAACCCAAGAGCTGCGATAAGACACATA
CCTGCCCACCTTGTCCAGCTCCCGAGCTGCTGGGCGGACCCTCAGTCTTTCTCTTCCACCTAAACC
CAAGGATACCCTTATGATCTCCAGGACTCCTGAGGTGACCTGCGTTGTGGTCGACGTGTCACATGAG
GACCCTGAGGTAAAGTTTAACTGGTACGTGGACGGTGTGGAGGTACATAACGCTAAGACTAAGCCAC
GAGAGGAGCAATACGCTTCCACTTACAGGGTGGTCAGCGTCCTGACCGTTCTCCATCAGGACTGGCT
GAACGGGAAGGAATATAAGTGTAAGGTTAGCAACAAAGCTCTCCCTGCACCAATCGAGAAGACAATC
AGCAAGGCAAAAAGGGCAGCCTCGGGAACCTCAGGTCTACACCCTCCCTCCTAGCAGGGAAGAGATGA
CAAAGAACCAGGTCTCTCTCACCTGCCTGGTGAAAGGCTTCTATCCATCTGACATTGCTGTGGAGTG
GGAATCCAACGGCCAGCCTGAAAATAATTATAAGACCACACCCCCCGTCCTTGATTCCGATGGATCT
TTCTTCCCTGTACAGTAACTCACCGTCGACAAATCACGGTGGCAGCAAGGTAACGTGTTTACAGTGT
CTGTCATGCATGAGGCTCTGCATAACCATTACACACAAAAGTCTTTGTCAATTGTCTCCAGGATGATG
AGAATTCATTGATCATAATCAGCCATAACCAC

SEQ ID NO: 430

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWRNGIVTVYADSVKGR
FTISRDN SKNTLYLQMN SLRAEDTAVYYCAKWSA AFDYWGQGLVTVSSASTKGPSVFPLAPSSKST
SGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNV
NHKPSNTKVDKKVEPKSCDKHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDSHE
DPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI
SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGS
FFLYSKLTVDKSRWQQGNV FSCSV MHEALHNHYTQKSLSLSPG

SEQ ID NO: 431

GAAGTGCAGCTGCTCGAAAGCGGCGGAGGCTTGGTGCAGCCAGGAGGGAGCCTGCGACTGTCTTGCG
CAGCCAGCGGATTCACCTTTCTCTTCTATGCCATGAGCTGGGTTCGACAGGCACCCGGCAAAGGTCT
CGAGTGGGTGTCTAGCATCTGGCGAAACGGAATAGTTACAGTGTATGCCGATAGCGTGAAGGGTCGC
TTTACTATTTACAGGGATAATTCTAAGAACACCCTCTACCTGCAAATGAATAGCCTTAGGGCAGAAG
ATACCGCCGTGTACTACTGTGCCAAATGGTCCGCAGCCTTTGACTACTGGGGCCAGGGGACACTGGT
GACCGTGTCTCTGTCATCAACCAAGGGGCCATCAGTGTTCCTACTCGCCCCATGTAGCAGATCAACA

TCTGAATCCACCGCAGCCCTTGGCTGCCTTGTTAAGGACTATTTCCCAGAACCCGTGACCGTAAGTT
GGA ACTCTGGCGCCCTTACTTCTGGGGTGACACCTTCCCAGCAGTGTTGCAGTCCAGTGGCCTTTA
CTCTCTGTCTAGTGTAGTGACTGTGCCTTCCTCTAGTCTCGGTACCAAGACCTATACCTGCAACGTA
GATCATAAGCCCAGCAATACAAAGGTTGATAAGAGAGTAGAGTCAAAGTACGGCCCACCCTGCCCAC
CTTGTCAGTCTCCGAGTTCCTGGGCGGACCCTCAGTCTTTCTCTTCCCACCTAAACCCAAGGATAC
CCTTATGATCTCCAGGACTCCTGAGGTGACCTGCGTTGTGGTCGACGTGTCACAAGAGGACCCTGAG
GTACAGTTTAACTGGTACGTGGACGGTGTGGAGGTACATAACGCTAAGACTAAGCCACGAGAGGAGC
AATTTAACTCCACTTACAGGGTGGTCAGCGTCCTGACCGTTCTCCATCAGGACTGGCTGAACGGGAA
GGAATATAAGTGTAAGGTTAGCAACAAAGGTCTGCCCAGTTCTATCGAGAAGACAATCAGCAAGGCA
AAAGGGCAGCCTCGGGAACCTCAGGTCTACACCCTCCCTCCTAGCCAGGAAGAGATGACAAAGAACC
AGGTCTCTCTCACCTGCCTGGTGAAAGGCTTCTATCCATCTGACATTGCTGTGGAGTGGGAATCCAA
CGGCCAGCCTGAAAATAATTATAAGACCACACCCCCCGTCCTTGATTCCGATGGATCTTTCTTCCTG
TACAGTCGCCTCACCGTCGACAAATCACGGTGGCAGGAAGGTAACGTGTTTCTGTCATGC
ATGAGGCTCTGCATAACCATTACACACAAAAGTCTTTGTCAATTGTCTCTCGGATGATGAGAATTCAT
TGATCATAATCAGCCATAACCAC

SEQ ID NO: 432

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWRNGIVTVYADSVKGR
FTISRDN SKNTLYLQMN SLRAEDTAVYYCAKWSA AFDYWGQGLVTVSSASTKGPSVFPLAPCSRST
SESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTKTYTCNV
DHKPSNTKVDKRVESKYGPPCP[CPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPE
VQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTI SKA
KGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFL
YSRLTVDKSRWQEGNVFSCSV MHEALHNHYTQKSLSLSLG

SEQ ID NO: 433

GAAGTGCAGCTGCTCGAAAGCGGCGGAGGCTTGGTGCAGCCAGGAGGGAGCCTGCGACTGTCTTGCG
CAGCCAGCGGATTCACTTTCTCTTCCTATGCCATGAGCTGGGTTCGACAGGCACCCGGCAAAGGTCT
CGAGTGGGTGTCTAGCATCTGGCGAAACGGAATAGTTACAGTGTATGCCGATAGCGTGAAGGGTCGC
TTTACTATTTACGGGATAATTCTAAGAACACCCTCTACCTGCAAATGAATAGCCTTAGGGCAGAAG
ATACCGCCGTGTACTACTGTGCCAAATGGTCCGCAGCCTTTGACTACTGGGGCCAGGGGACACTGGT
GACCGTGTCTCTGCATCAACCAAGGGGCCATCAGTGTTCCTACTCGCCCCATCTTCCAAGAGTACT
TCCGGCGGAACCGCAGCCCTTGGCTGCCTTGTTAAGGACTATTTCCCAGAACCCGTGACCGTAAGTT
GGA ACTCTGGCGCCCTTACTTCTGGGGTGACACCTTCCCAGCAGTGTTGCAGTCCAGTGGCCTTTA
CTCTCTGTCTAGTGTAGTGACTGTGCCTTCCTCTAGTCTCGGTACCCAGACCTATATTTGTAATGTT
AACCATAAGCCCAGCAATACAAAGGTTGATAAGAAAGTGGAACCCAAGAGCTGCGATAAGACACATA

CCTGCCCACCTTGTCCAGCTCCCGAGCTGCTGGGCGGACCCTCAGTCTTTCTCTTCCCACCTAAACC
CAAGGATACCTTTATGATCTCCAGGACTCCTGAGGTGACCTGCGTTGTGGTCGACGTGTCACATGAG
GACCCTGAGGTAAAGTTTAACTGGTACGTGGACGGTGTGGAGGTACATAACGCTAAGACTAAGCCAC
GAGAGGAGCAATACGCTTCCACTTACAGGGTGGTCAGCGTCCTGACCGTTCTCCATCAGGACTGGCT
GAACGGGAAGGAATATAAGTGTAAGGTTAGCAACAAAGCTCTCCCTGCACCAATCGAGAAGACAATC
AGCAAGGCAAAAAGGGCAGCCTCGGGAACCTCAGGTCTACACCCTCCCTCCTAGCAGGGAAGAGATGA
CAAAGAACCAGGTCTCTCTCACCTGCCTGGTGAAAGGCTTCTATCCATCTGACATTGCTGTGGAGTG
GGAATCCAACGGCCAGCCTGAAAATAATTATAAGACCACACCCCCCGTCCTTGATTCCGATGGATCT
TTCTTCCTGTACAGTAACTCACCGTCGACAAATCACGGTGGCAGCAAGGTAACGTGTTTACAGCTGTT
CTGTCATGCATGAGGCTCTGCATAACCATTACACACAAAAGTCTTTGTATTGTCTCCAGGATGATG
AGAATTCATTGATCATAATCAGCCATACCAC

SEQ ID NO: 434

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWRNGIVTVYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCAKWSAAFQYWGQGLVTVSSASTKGPSVFPLAPSSKST
SGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNV
NHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDSHE
DPEVKFNWYVDGVEVHNAKTKPREEQY[**A**]STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI
SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGS
FFLYSKLTVDKSRWQQGNVFSFSVMHEALHNHYTQKSLSLSPG

SEQ ID NO: 1202

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSSIWRNGIVTVYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCAKWSAAFQYWGQGLVTVSSASTKGPSVFPLAPSSKST
SGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNV
NHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDSHE
DPEVKFNWYVDGVEVHNAKTKPREEQY[**Q**]STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI
SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGS
FFLYSKLTVDKSRWQQGNVFSFSVMHEALHNHYTQKSLSLSPG

[000612] Additional examples of heavy chain constant domains include, but are not limited to, a heavy chain constant region having a mutation or deletion at one or more of the following amino acids: Ser 228, Leu234, Leu235, Gly236, Gly237, Asp265, Asp270, Asn297, Lys 320, Lys 322, Glu328, Pro329, Pro331 as numbered in the EU index as set forth in Kabat, such that the Fc domain has less affinity for a Fc-gamma receptor.

EXAMPLE 12. Evaluation of efficiency of masking moieties

[000613] This assay measures the ability of the masking peptide to block binding of the activatable antibody to antigen as compared to the binding of the unmodified antibody to antigen.

[000614] The general outline for this assay is as follows: Nunc, Maxisorp plates are coated overnight at 4°C with 100 µl/well of a 1 µg/mL solution of human PDL1 (R & D Systems) in PBS, pH 7.4. Plates are washed 3 times with PBST (PBS, pH 7.4, 0.05% Tween-20), and wells are blocked with 200 µl/well of 10 mg/mL BSA in PBST for 2 hours at RT. Plates are washed 3 times with PBST (PBS, pH 7.4, 0.05% Tween-20). Dilution curves are prepared, in 10 mg/mL BSA in PBST, as shown below in Table 10.

Table 10. Plate layout for masking efficiency assay, one plate for each time point.

	[Antibody] = nM Columns 1-3	[activatable antibody 1] = nM Columns 4-6	[activatable antibody 2] = nM Columns 7-9	[activatable antibody 3] = nM Columns 10-12
A	37	1000	1000	1000
B	12.3	333	333	333
C	4.1	111	111	111
D	1.34	37	37	37
E	0.45	12.3	12.3	12.3
F	0.15	4.1	4.1	4.1
G	0.03	1.34	1.34	1.34
H	0.01	0.45	0.45	Blank

[000615] The binding solutions are added to the plates, which are then incubated for 1 hour at room temperature (RT), and then washed 3 times with PBST (PBS, pH 7.4, 0.05% Tween-20). One hundred (100) µl/well 1:4000 dilution goat-anti-human IgG (Fab specific, Sigma cat # A0293) in 10 mg/mL BSA in PBST is added, and the plate is incubated for 1 hour at RT. The plate is developed with TMB and 1N HCl. Shown in Figures 15-17 are plots of binding isotherms for anti-PDL1 activatable antibodies that include the anti-PDL1 antibody C5H9v2 described herein. Plots are generated in Prizm

(Sigma Plot) and the data are fit to a model of single site saturation and a K_d is determined.

In Figure 17, PL03+ is an effector positive activatable antibody, namely wild type IgG1.

PL03- is an effector negative activatable antibody, namely mutated IgG1 N297Q.

[000616] Masking efficiency is calculated by dividing the K_d for binding of the activatable antibodies by the K_d of the parental antibody. Masking efficiencies are shown in Figures 15-17.

EXAMPLE 13. Anti-PDL1 activatable antibodies of the disclosure delay induction of diabetes in NOD mice

[000617] In this Example, anti-PDL1 activatable antibodies PL15-0003-C5H9v2 and PL18-0003-C5H9v2 were analyzed for the ability to induce diabetes in NOD mice. The NOD mice, substrain NOD/ShiLtJ, were obtained from Jackson Laboratory at 6 weeks and acclimated on site. At 9.5 weeks, mice were checked for diabetes prior to enrollment, grouped, and dosed as set forth in Table 19.

Table 19. Groups and doses for diabetes study with anti-PDL1 activatable antibodies

Group	Count	Treatment	Dose (mg/kg)	Dose volume (mL/kg)	Schedule	Route
1	8	mIgG2a (C1.18.4)	3	10	Day 0	IP
2	8	Anti-PDL1(C5H9v2)	1	10	Day 0	IP
3	8	Anti-PDL1(C5H9v2)	0.3	10	Day 0	IP
4	8	Activatable Antibody PL15-0003-C5H9v2	1	10	Day 0	IP
5	8	Activatable Antibody PL18-0003-C5H9v2	1	10	Day 0	IP

[000618] Figure 18A, which plots % non-diabetic versus number of days post initial dose, shows that anti-PDL1 antibody C5H9v2 induced diabetes in NOD mice at 1 mg/kg while the anti-PDL1 activatable antibody PL15-0003-C5H9v2 (labeled herein as Gr4 - PF15) did not induce diabetes and the anti-PDL1 activatable antibody PL18-0003-C5H9v2 (labeled in Figure 18 as Gr 5 - PL18) exhibited delayed diabetes at 1 mg/kg. Control mIgG2a and anti-PDL1 antibody C5H9v2 at 0.3 mg/kg did not induce diabetes in NOD mice. In an additional group, a single 3 mg/kg dose of anti-PDL1 activatable antibody PL15-0003-C5H9v2 administered IP at a dose volume of 10 mL/kg induced diabetes in 2 of 8 mice (75% non-diabetic), providing a greater than three-fold safety margin relative to anti-PDL1 antibody C5H9v2 (Group 2). Results are shown in Figure 18B.

EXAMPLE 14. Anti-PDL1 activatable antibodies of the disclosure reduce MC38 tumors in mice

[000619] In this Example, anti-PDL1 activatable antibodies PL15-0003-C5H9v2 and PL18-0003-C5H9v2 were analyzed for the ability to reduce the growth of MC38 syngeneic tumors.

[000620] The mouse colon carcinoma cell line MC38 was obtained from ATCC. MC38 were grown in RPMI-1640 supplemented with 10% fetal bovine serum at 37°C in an atmosphere of 5% CO₂ in air. Cells were harvested during the logarithmic growth period, resuspended in PBS with proper cell concentration, and kept on ice for tumor induction.

[000621] Each mouse was inoculated subcutaneously in the right flank with 0.5X10⁶ of MC38 cells in PBS for tumor development. The treatments were started when the mean tumor size reached approximately 100-200 mm³ (no more than 200 mm³). Tumor sizes were measured twice weekly in two dimensions using a caliper, and the volume was expressed in mm³ using the formula: $V = 0.5 \times a \times b^2$ where a and b are the long and short diameters of the tumor, respectively.

[000622] The mice were grouped and dosed as set forth in Table 20.

Table 20. Groups and doses for MC38 syngeneic study with anti-PDL1 activatable antibodies

Group	Count	Treatment	Dose (mg/kg)	Dose volume (mL/kg)	Schedule	Route
1	9	mIgG2a (C1.18.4)	10	10	t.i.w. for 2 weeks	IP
2	9	Anti-PDL1(C5H9v2)	10	10	t.i.w. for 2 weeks	IP
4	9	PL18-0003-C5H9v2	10	10	t.i.w. for 2 weeks	IP
6	9	PL15-0003-C5H9v2	10	10	t.i.w. for 2 weeks	IP

[000623] Figure 19, which plots tumor volume versus number of days post initial dose, demonstrates that anti-PDL1 activatable antibodies PL15-0003-C5H9v2 and PL18-0003-C5H9v2 inhibited the growth of MC38 syngeneic tumors similar to positive control anti-PDL1 antibody C5H9v2.

EXAMPLE 15. An anti-PDL1 activatable antibody of the disclosure demonstrates reduced PDL1 occupancy in blood and spleen from C57Bl/6 mice

[000624] Unlike an antibody that would bind target antigen indiscriminately, an activatable antibody would be a molecule that is inert unless it is activated in the tumor microenvironment. Once activated, the cleaved activatable antibody would bind the targeted antigen within the tumor only, sparing the antigens in the peripheral areas. PDL1 is expressed in tumor cells as well as T cells (CD4+ and CD8+) in circulation and the spleen. This example demonstrates that animals treated with anti-PDL1 activatable antibody did not have detectable activatable antibody on T cells in the periphery, while animals treated with anti-PDL1 antibody showed a detectable dose dependent presence of antibody on T cells in the periphery.

[000625] To measure whether anti-PDL1 activatable antibody PL15-0003-C5H9v2 is protected from binding T cells in circulation and the spleen compared to the anti-PDL1 antibody C5H9v2, 14-week old C57Bl/6 mice (with tumors ranging from 38-671mm³) were dosed as set forth in Table 21. Tumor sizes were measured in two dimensions using a caliper, and the volume was expressed in mm³ using the formula: $V = 0.5 \times a \times b^2$ where *a* and *b* are the long and short diameters of the tumor, respectively. After about 20 hours, animals were euthanized; and blood, spleen, and plasma were harvested from each animal.

Table 21. Groups and doses for PDL1 occupancy study

Group	Count	Treatment	Dose (mg/kg)	Dose volume (mL/kg)	Schedule	Route
1	2	mIgG2a (C1.18.4)	1	10	Day 0	IP
2	2	Anti-PDL1(C5H9v2)	1	10	Day 0	IP
3	2	Anti-PDL1(C5H9v2)	0.3	10	Day 0	IP
4	2	Anti-PDL1(C5H9v2)	0.1	10	Day 0	IP
5	2	PL15-0003--C5H9v2	1	10	Day 0	IP
6	2	PL15-0003-C5H9v2	0.3	10	Day 0	IP

[000626] Whole blood was processed as follows: One volume of blood was added to 10 volumes of RBC Lysis Solution (130-094-183 Miltenyi Biotec) and processed using

manufacturer's protocol. Lymphocytes were resuspended in staining buffer (HBSS 2% FBS). Spleen was processed as follows: Spleen was placed in a 70µM cell strainer (352350 Corning/BD Falcon) with 1 mL of serum free RPMI-1640. The back of a 3-mL syringe was used to mash the spleen in a circular motion. The strainer was washed with media. Splenocytes were pelleted using a centrifuge and blood was lysed using 10 volumes of RBC Lysis Solution (130-094-183 Miltenyi Biotec) according to manufacturer's protocol. Splenocytes were refiltered after resuspending in staining buffer (HBSS 2% FBS). Lymphocytes from whole blood and lymphocytes from spleen were counted and were aliquoted to wells of a 96-well round bottom plates (5e5 cells/well). Cells were pelleted and mouse FcR Reagent was added 1:10 dilution according to manufacturer's protocol (130-092-575 Miltenyi Biotec). One set of cells was stained with a saturating amount, 100nM, of anti-PDL1 antibody C5H9v2 for 60 minutes to obtain a maximum saturation value. Cells were washed 3 times and are stained with a biotinylated 'a' allotype antibody, anti-mIgG2a(a) clone 8.3 (553502 BD Biosciences) at 300ng/mL for 60 minutes. Cells were washed 3 times and then stained with 1 µg/mL (also referred to herein as "µg/mL") of anti-CD4-Pacific Blue (558107 BD Bioscience), 1µg/mL of anti-CD8-APC (553035 BD Bioscience) and 1:500 dilution of SAPE (S-866 Life Technologies) for 30 minutes. Cells were washed 2 times and stained with 7-AAD according to manufacturer's protocol (559925 BD Bioscience). The MACSQuant flow cytometer was used to measure the amount of anti-PDL1 antibody C5H9v2 and anti-PDL1 activatable antibody PL15-0003-C5H9v2 bound to T cells from whole blood and the spleen. Briefly, at the most 30,000 events were collected in the lymphocyte gate. Live cells were gated from the 7AAD negative gate. CD4⁺ and CD8⁺ cells were gated separately and anti-PDL1 mAb and anti-PDL1 activatable antibody MFI were recorded for both CD4 and CD8. Percent occupancy was determined by taking the MFI of the sample compared to the average of the MFI for the maximum binding MFI. Figures 20A-20D demonstrate the percent of anti-PDL1 antibody C5H9v2 and anti-PDL1 activatable antibody PL15-0003-C5H9v2 bound to CD4⁺ and CD8⁺ T cells from peripheral blood (Figures 20A, 20B) or spleen (Figures 20C, 20D). In all graphs, the circle represents isotype, the square represents anti-PDL1 antibody C5H9v2, and the triangle represents anti-PDL1 activatable antibody PL15-0003-C5H9v2. Percent occupancy of the anti-PDL1 antibody C5H9v2 was reflective of the dose titration. Percent occupancy of the anti-PDL1 activatable antibody PL15-0003-C5H9v2 at 1 mg/kg was similar to isotype control mIgG2a at 1 mg/kg.

EXAMPLE 16. An anti-PDL1 activatable antibody of the disclosure demonstrates reduced PDL1 occupancy in blood and spleen from C57Bl/6 mice

[000627] This example demonstrates that animals treated with an anti-PDL1 activatable antibody of the disclosure exhibited reduced binding to blood T cells from tumor bearing mice compared to that exhibited by an anti-PDL1 antibody of the disclosure at a variety of doses.

[000628] The study was conducted in a manner similar to that described in Example 15. Briefly, mice bearing MC38 tumors between 100-200 mm³ were treated with a single dose of anti-PDL1 antibody C5H9v2 or anti-PDL1 activatable antibody PL15-0003-C5H9v2 as indicated in Figures 21A and 21B, and blood was analyzed for surface bound antibody by flow cytometry four days after dosing. Plasma concentrations of the antibody and activatable antibody were determined by ELISA. Figures 21A and 21 B demonstrate that the presence of tumor-derived proteases capable of cleaving the activatable antibody did not lead to high levels of activated activatable antibody in the blood (Figure 21A); i.e., the activatable antibody (depicted by squares) demonstrated reduced peripheral PDL1 binding in tumor-bearing mice compared to the antibody (depicted by circles) or an isotype antibody (depicted by triangles) despite the plasma concentrations of the activatable antibody being higher than those of the antibody (Figure 21B).

EXAMPLE 17. Activity of an anti-PDL1 antibody and an anti-PDL1 activatable antibody of the embodiments in a human T-cell restimulation assay

[000629] In this example, peripheral blood mononuclear cells from a CMV-positive donor were incubated in the presence of CMV viral lysate and an anti-PDL1 antibody or an anti-PDL1 activatable antibody of the disclosure to assess the effect of such anti-PDL1 antibody or anti-PDL1 activatable antibody on interferon gamma (IFN-gamma) cytokine secretion.

[000630] PBMCs from a CMV-positive donor (Hemacare Donor C) were plated at 3.5×10^5 cells per well in the presence of 4 µg/mL CMV viral lysate (Astarte) and either anti-PDL1 antibody C5H9v2, anti-PDL1 activatable antibody PL07-2001-C5H9v2, or an hIgG4 isotype control antibody. After four days, supernatant was removed from each well and IFN-gamma levels were assayed using IFN-gamma ELISA kit (Life Technologies, Carlsbad, CA). Figure 22 demonstrates that anti-PDL1 activatable antibody PL07-2001-

C5H9v2 exhibited increased CMV-stimulated IFN-gamma secretion compared with control hIgG4 but decreased potency relative to anti-PDL1 parental antibody C5H9v2 due to the activatable antibody being masked.

EXAMPLE 18. *In situ* imaging of an anti-PDL1 activatable antibody of the disclosure

[000631] This Example demonstrates the ability of anti-PDL1 activatable antibody PL15-0003-C5H9v2 to be activated and to bind frozen MC38 mouse cancer tissues using an *in situ* imaging method.

[000632] Fluorescently-labeled anti-PDL1 activatable antibody PL15-0003-C5H9v2 or anti-PDL1 antibody C5H9v2 was incubated on frozen PDL1+ MC38 tumor sections for 1 hour in a protease-compatible buffer in the presence or absence of a broad spectrum inhibitor cocktail as described in PCT International publication number WO 2014/107599, published July 10, 2014.

[000633] Results are shown in Figure 23. The tissue image in Figure 23A demonstrates binding of anti-PDL1 antibody C5H9v2 to the tumor section indicating the presence of PDL1 antigen in the tumor section. The tissue image in Figure 23B demonstrates that anti-PDL1 activatable antibody PL15-0003-C5H9v2 was activated by tumor-derived proteolytic cleavage of the anti-PDL1 activatable antibody to yield an anti-PDL1 antibody that bound to the PDL1 target in the tumor section. The tissue image in Figure 23D demonstrates that the fluorescent signal shown in Figure 23B was inhibited by pre-treatment of the tumor section with a 1:100 dilution of broad spectrum inhibitor cocktail set III and 50 mM EDTA, whereas no effect of broad spectrum protease inhibitors was detected on the binding of anti-PDL1 antibody C5H9v2 to the tumor section as demonstrated in the tissue image in Figure 23C. These results demonstrate that the tumor sections comprise sufficient protease activity to activate activatable antibodies of the disclosure.

EXAMPLE 19. Ability of an activatable antibody of the disclosure to demonstrate protease-dependent binding and blocking activity *in vitro*

[000634] This Example demonstrates the ability anti-PDL1 activatable antibody PL15-0003-C5H9v2 to exhibit protease-dependent binding and blocking activity *in vitro*.

[000635] Binding and blocking studies were conducted on anti-PDL1 antibody C5H9v2 and anti-PDL1 activatable antibody PL15-0003-C5H9v2 in a manner similar to

those described elsewhere in the Examples except that plates were coated with 0.5 µg/mL PDL1.

[000636] Figure 24 depicts that anti-PDL1 activatable antibody PL15-0003-C5H9v2 exhibited a higher EC₅₀ for PDL1 binding and PD1 blocking than did the anti-PDL1 antibody as measured by ELISA. The figure further demonstrates that activation of anti-PDL1 activatable antibody PL15-0003-C5H9v2 by matriptase fully restored PDL1 binding and PD1 blocking activities to levels comparable to those of anti-PDL1 antibody C5H9v2.

EXAMPLE 20. Ability of an anti-PDL1 activatable antibody of the disclosure to reduce binding to PDL1 and to demonstrate protease-dependent blocking activity *in vitro*

[000637] This Example measures the ability of a masking peptide to reduce binding of an anti-PDL1 activatable antibody of the disclosure to PDL1 compared to the ability of the parental antibody to bind to PDL1 *in vitro*. This Example also demonstrates the ability of an anti-PDL1 activatable antibody of the disclosure to exhibit protease-dependent blocking activity *in vitro*.

[000638] The abilities of anti-PDL1 activatable antibodies PL07-2001-C5H9v2 and PF07-3001-C5H9v2 and anti-PDL1 antibody C5H9 to bind to human or murine PDL1 were tested in a manner similar to techniques described elsewhere in the Examples. Results are shown in Figure 25.

[000639] The abilities of anti-PDL1 activatable antibody PL07-2001-C5H9v2, uPA-activated anti-PDL1 activatable antibody PL07-2001-C5H9v2, MMP14-activated anti-PDL1 activatable antibody PL07-2001-C5H9v2, and anti-PDL1 antibody C5H9v2 to block PD1 (PD-1) or B7-1 binding to PDL1 were tested in a manner similar to techniques described elsewhere in the Examples.

[000640] Briefly, for the PD1 blockade assay, human, cyno or rat PDL1 was adsorbed to the well of an ELISA plate. Biotinylated human, cyno, or rat PD1 was applied to the wells in the absence or presence of an increasing concentration of either anti-PDL1 activatable antibody PL07-2001-C5H9v2, uPA-activated anti-PDL1 activatable antibody PL07-2001-C5H9v2, MMP14-activated anti-PDL1 activatable antibody PL07-2001-C5H9v2, or anti-PDL1 antibody C5H9v2. Results, shown in Figure 26, indicate that activatable antibody PF07-2001-C5H9v2 exhibited a higher EC₅₀ for PD1 blocking than did the anti-PDL1 antibody as measured by ELISA. However, activation of the activatable

antibody by either uPA or MMP14 restored PD1 blocking activity to levels comparable to those of anti-PDL1 antibody C5H9v2.

[000641] Briefly, for the B7-1 blockade assay, human or cyno PDL1 was adsorbed to the well of an ELISA plate. Biotinylated human or cyno B7-1 was applied to the wells in the absence or presence of an increasing concentration of either anti-PDL1 activatable antibody PL07-2001-C5H9v2, uPA-activated anti-PDL1 activatable antibody PL07-2001-C5H9v2, MMP14-activated anti-PDL1 activatable antibody PL07-2001-C5H9v2, or anti-PDL1 antibody C5H9v2. Results, shown in Figure 27, indicate that activatable antibody PF07-2001-C5H9v2 exhibited a higher EC₅₀ for B7-1 blocking than did the anti-PDL1 antibody as measured by ELISA. However, activation of the activatable antibody by either uPA or MMP14 restored B7-1 blocking activity to levels comparable to those of anti-PDL1 antibody C5H9v2.

EXAMPLE 21. Anti-PDL1 activatable antibodies of the disclosure reduce MC38 tumors in mice

[000642] In this Example, anti-PDL1 activatable antibodies PL15-0003-C5H9v2, PL15-2001-C5H9v2, and PL15-3001-C5H9v2 were analyzed for the ability to reduce the growth of MC38 syngeneic tumors.

[000643] The mouse colon carcinoma cell line MC38 was obtained from ATCC. MC38 were grown in RPMI-1640 supplemented with 10% fetal bovine serum at 37°C in an atmosphere of 5% CO₂ in air. Cells were harvested during the logarithmic growth period, resuspended in PBS with proper cell concentration, and kept on ice for tumor induction.

[000644] Each mouse was inoculated subcutaneously in the right flank with 0.5X10⁶ of MC38 cells in PBS for tumor development. The treatments were started when the mean tumor size reached approximately 100-200 mm³ (no more than 200 mm³). Tumor sizes were measured twice weekly in two dimensions using a caliper, and the volume was expressed in mm³ using the formula: $V = 0.5 \times a \times b^2$ where a and b are the long and short diameters of the tumor, respectively.

[000645] The mice were grouped and dosed as set forth in Table 24.

Table 24. Groups and doses for MC38 syngeneic study with anti-PDL1 activatable antibodies

Group	Count	Treatment	Dose (mg/kg)	Dose volume (mL/kg)	Schedule	Route
1	9	mIgG2a (C1.18.4)	5	10	b.i.w.; 4 doses	IP
2	9	Anti-PDL1 (C5H9v2)	5	10	b.i.w.; 4 doses	IP
3	9	PL15-0003- C5H9v2	5	10	b.i.w.; 4 doses	IP
4	9	PL15-2001- C5H9v2	5	10	b.i.w.; 4 doses	IP
5	9	PL15-3001- C5H9v2	5	10	b.i.w.; 4 doses	IP

[000646] Figure 28, which plots tumor volume versus number of days post initial dose, demonstrates that anti-PDL1 activatable antibodies PL15-0003-C5H9v2, PL15-2001-C5H9v2, and PL15-3001-C5H9v2 inhibited the growth of MC38 syngeneic tumors similar to positive control anti-PDL1 antibody C5H9v2.

EXAMPLE 22. An anti-PDL1 activatable antibody of the disclosure demonstrates reduced PDL1 occupancy in blood from C57Bl/6 mice

[000647] This example demonstrates that animals treated with anti-PDL1 activatable antibodies of the disclosure exhibited reduced binding to blood T cells from tumor bearing mice compared to that exhibited by an anti-PDL1 antibody of the disclosure.

[000648] The study was conducted in a manner similar to that described in Example 15. Briefly, mice bearing MC38 tumors between 100-200 mm³ were treated with a single dose of anti-PDL1 antibody C5H9v2 or anti-PDL1 activatable antibody PL15-0003-C5H9v2, PL15-2001-C5H9v2, or PL15-3001-C5H9v2 as indicated in Table 25, and blood was analyzed for surface bound antibody by flow cytometry four and eight days after dosing. Percent occupancy was calculated as described in Example 15.

Table 25. Groups and doses for PDL1 occupancy study

Group	Count	Treatment	Dose (mg/kg)	Dose volume (mL/kg)	Schedule	Route
-------	-------	-----------	--------------	---------------------	----------	-------

1	9	mIgG2a (C1.18.4)	5	10	b.i.w.; 4 doses	IP
2	9	Anti-PD-L1(C5H9v2)	5	10	b.i.w.; 4 doses	IP
3	9	Anti-PD-L1(C5H9v2)	3	10	b.i.w.; 4 doses	IP
4	9	PL15-0003-C5H9v2	5	10	b.i.w.; 4 doses	IP
5	9	PL15-0003-C5H9v2	3	10	b.i.w.; 4 doses	IP
6	9	PL15-2001-C5H9v2	5	10	b.i.w.; 4 doses	IP
7	9	PL15-2001-C5H9v2	3	10	b.i.w.; 4 doses	IP
8	9	PL15-3001-C5H9v2	5	10	b.i.w.; 4 doses	IP
9	9	PL15-3001-C5H9v2	3	10	b.i.w.; 4 doses	IP

[000649] Plasma concentrations of the antibody and activatable antibodies were determined by ELISA. Figure 29 demonstrates that the presence of tumor-derived proteases capable of cleaving the activatable antibodies did not lead to high levels of activated activatable antibodies in the blood.

EXAMPLE 23. In vivo imaging of an anti-PDL1 activatable antibody of the disclosure

[000650] This Example demonstrates the ability of proteases in a tumor implanted into a mouse to activate an anti-PDL1 activatable antibody of the disclosure.

[000651] A MDA-MB-231-luc2 orthotopic xenograft breast cancer model was used for these studies. Female nu/nu mice were purchased from Charles River Laboratories to arrive at 7 week old. After 1 week acclimation, three million MDA-MD-231-luc2-4D3LN cells (Perkin Elmer) in 30 microliters (ul, also referred to herein as “ μ l”) serum-free RPMI (1:1 matrigel) were injected into the fourth abdominal left mammary fat pad. Tumors were grown to 110-159 mm³. Three mice per group were administered intravenously 5 mg/kg Alexa750-conjugated anti-PDL1 antibody C5H9v2 or anti-PDL1 activatable antibody PL07-2001-C5H9v2. Four mice were administered intravenously 5 mg/kg Alexa750-conjugated anti-RSV antibody control palivizumab (Synagis). Ninety-six hours after administration, optical images were collected from each of the mice. Results are shown in Figure 30. A high-intensity fluorescent signal was detected only in the tumors of mice dosed with the anti-PDL1 antibody or anti-PDL1 activatable antibody PL07-2001-C5H9v2,

suggesting that the anti-PDL1 activatable antibody was activated and accumulated in the tumor through PDL1 binding.

EXAMPLE 24. Activation of an activatable antibody of the disclosure in human plasma samples

[000652] This Example determines whether an activatable antibody of the disclosure is activated in human plasma samples.

[000653] Anti-PDL1 activatable antibody PL07-2001-C5H9v2 was conjugated with Oregon Green dye (ThermoFisher Cat# O6149). The concentration and degree of labeling was determined with a spectrophotometer. Three uM (also referred to herein as “μM”) of labeled anti-PDL1 activatable antibody PL07-2001-C5H9v2 was added into plasma samples obtained from healthy donors, patients with melanoma or lung cancer patients in a final activatable antibody to plasma ratio of 30:70. The samples were incubated for 48 hours in a 37° C humidifying chamber. Reactions of labeled anti-PDL1 activatable antibody PL07-2001-C5H9v2 in plasma were stopped at either 0 or 48 hours by freezing samples in -80° C or denaturing in Wes running buffer (ProteinSimple Cat #PS-MK14). Samples were analyzed by the Wes capillary western blot system (ProteinSimple) using antibodies against Oregon Green dye (ThermoFisher Cat# A-11095 at 100μg/ml) followed by HRP conjugated secondary antibody (Jackson ImmunoResearch Cat# 705-035-147 at 1/40 dilution). Analysis of percent activation of the labeled anti-PDL1 activatable antibody PL07-2001-C5H9v2 was determined using Compass (ProteinSimple) software.

[000654] There was no detectable activatable antibody activation in five normal patient samples, in five melanoma patient samples (three metastases in lymph nodes, two in the brain, and one on the skin of the back), nor in five lung cancer patient samples (one metastasis in the lower lobe of the left lung (stage IV), one in the lower lobe of the right lung (stage IV), one in the upper lobe of the left lung (stage IA), and two in the upper lobe of the right lung (stages IIIA and IB).

Other Embodiments

[000655] While the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following.

What is claimed is:

1. An isolated antibody or antigen binding fragment thereof (AB) that specifically binds to mammalian PDL1, wherein the AB comprises
 - (a) a variable heavy chain complementarity determining region 1 (VH CDR1) comprising the amino acid sequence of SEQ ID NO: 212;
 - (b) a variable heavy chain complementarity region 2 (VH CDR2) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 213, 217, 221, 225, 229, 231, and 238-247;
 - (c) a variable heavy chain complementarity determining region 3 (VH CDR3) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 214, 218, 222, 226, 230, 232, and 233-237;
 - (d) a variable light chain complementarity determining region 1 (VL CDR1) comprising the amino acid sequence of SEQ ID NO: 209;
 - (e) a variable light chain complementarity determining region 2 (VL CDR2) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 210, 215, 219, 223, and 227; and
 - (f) a variable light chain complementarity determining region 3 (VL CDR3) comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 211, 216, 220, 224, and 228.
2. An activatable antibody that, in an activated state, specifically binds to mammalian PDL1, wherein said activatable antibody comprises:
 - an antibody or an antigen binding fragment thereof (AB) that specifically binds to mammalian PDL1, wherein the AB comprises the AB of claim 1;
 - a masking moiety (MM) that inhibits the binding of the AB to mammalian PDL1 when the activatable antibody is in an uncleaved state; and
 - a cleavable moiety (CM) coupled to the AB, wherein the CM is a polypeptide that functions as a substrate for a protease.
3. The activatable antibody of claim 2, that, in an activated state specifically blocks a natural ligand of PDL1 from binding to the mammalian PDL1.

4. The antibody of claim 1 or the activatable antibody of any one of claims 2 to 3, wherein the mammalian PDL1 is a human PDL1.
5. The activatable antibody of any one of claims 2 to 4, wherein:
 - (a) the AB induces type 1 diabetes in a non-obese diabetic (NOD) mouse; and
 - (b) the activatable antibody in an uncleaved state inhibits the induction of type 1 diabetes in a NOD mouse, and/or optionally, wherein
 - (a) the activatable antibody in an uncleaved state does not induce type 1 diabetes in greater than 50% of a population of non-obese diabetic (NOD) mice, and
 - (b) the AB induces type 1 diabetes in greater than 50% of a population of NOD mice, and/or optionally wherein: the NOD mouse is a female NOD/ShiLtJ mouse substrain, and/or optionally, wherein the activatable antibody inhibits the induction of type 1 diabetes in a NOD mouse by at least 3-fold compared to the AB.
6. The activatable antibody of any one of claims 2 to 5, wherein the activatable antibody in an uncleaved state binds to a smaller percentage of a population of peripheral blood CD4⁺ CD8⁺ T lymphocytes than does the AB.
7. The antibody of any one of claims 1 or 4 or the activatable antibody of any one of claims 2-6, wherein
 - the VL CDR2 comprises the amino acid sequence of SEQ ID NO: 227,
 - the VL CDR3 comprises the amino acid sequence of SEQ ID NO: 228,
 - the VH CDR2 comprises the amino acid sequence of SEQ ID NO: 246, and
 - the VH CDR3 comprises the amino acid sequence of SEQ ID NO: 235.
8. The antibody of any one of claims 1 or 4 or the activatable antibody of any one of claims 2 to 6, wherein
 - the VL CDR2 sequence comprises the amino acid sequence of SEQ ID NO: 215;
 - the VL CDR3 sequence comprises the amino acid sequence of SEQ ID NO: 228;
 - the VH CDR2 sequence comprises the amino acid sequence of SEQ ID NO: 246; and
 - the VH CDR3 sequence comprises the amino acid sequence of SEQ ID NO: 235
9. The antibody of claim 1 or the activatable antibody of claim 2, wherein the AB comprises a variable heavy chain (VH) comprising the amino acid sequence of SEQ ID

NO: 46 and a variable light chain (VL) comprising the amino acid sequence of SEQ ID NO: 58, or a VH comprising the amino acid sequence of SEQ ID NO: 46 and a VL comprising the amino acid sequence of SEQ ID NO: 12.

10. The activatable antibody of any one of claims 2 to 9,
 - wherein the MM has a dissociation constant for binding to the AB that is greater than the dissociation constant of the AB to PDL1; and/or optionally,
 - wherein the MM does not interfere or compete with the AB for binding to PDL1 when the activatable antibody is in a cleaved state; and/or optionally,
 - wherein the MM is a polypeptide of no more than 40 amino acids in length; and/or optionally,
 - wherein the MM polypeptide sequence is different from that of human PDL1; and/or optionally, wherein the MM polypeptide sequence is no more than 50% identical to any natural binding partner of the AB; and/or optionally,
 - wherein the MM polypeptide sequence is no more than 25% identical to any natural binding partner of the AB.
11. The activatable antibody of any one of claims 2 to 10, wherein the MM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 59-81, 208, and 426.
12. The activatable antibody of any one of claims 2 to 11, wherein the CM is a substrate for a protease that is active in diseased tissue.
13. The activatable antibody of any one of claims 2 to 11, wherein the CM comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 338-394, and 435-445, 883-921, 1009, and 1010.
14. The antibody of any one of claims 1, 4, or 7-9 or the activatable antibody of any one of claims 2 to 13, wherein the antigen binding fragment thereof is selected from the group consisting of a Fab fragment, a F(ab')₂ fragment, a scFv, a scAb, a dAb, a single domain heavy chain antibody, and a single domain light chain antibody.

15. The activatable antibody of any one of claims 2 to 14, wherein the AB is linked to the CM, wherein the AB is linked directly to the CM or via a linking peptide.
16. The activatable antibody of any one of claims 2 to 15,
wherein the MM is linked to the CM such that the activatable antibody in an uncleaved state comprises the structural arrangement from N-terminus to C-terminus as follows: MM-CM-AB or AB-CM-MM; and/or optionally,
wherein the activatable antibody comprises a linking peptide between the MM and the CM; and/or optionally,
wherein the activatable antibody comprises a linking peptide between the CM and the AB; and/or optionally,
wherein the activatable antibody comprises a first linking peptide (LP1) and a second linking peptide (LP2), and wherein the activatable antibody in the uncleaved state has the structural arrangement from N-terminus to C-terminus as follows: MM-LP1-CM-LP2-AB or AB-LP2-CM-LP1-MM; and/or optionally,
wherein the two linking peptides need not be identical to each other; and/or optionally,
wherein each of LP1 and LP2 is a peptide of about 1 to 20 amino acids in length.
17. The activatable antibody of any one of claims 2 to 20 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 931, 933, 935, 937, 939, 941, 943, 945, 947, 949, 951, 953, 955, 957, 959, 961, 963, 965, 967, 969, 971, 973, 975, 977, 979, 981, 983, 985, 987, 989, 991, 993, 995, 997, 999, 1001, 1003, 1005, and 1144-1191, 1200, and 1201.
18. The activatable antibody of claim 17, wherein the activatable antibody comprises the amino acid sequence of SEQ ID NO: 985.
19. The activatable antibody of claim 17, wherein the activatable antibody comprises the amino acid sequence of SEQ ID NO: 137.

20. The activatable antibody of any one of claims 2 to 19, wherein the activatable antibody comprises the amino acid sequence of SEQ ID NO: 428 or SEQ ID NO: 1008.
21. The activatable antibody of any one of claims 2 to 20, wherein the activatable antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 430, 432, 434, and 1202.
22. The activatable antibody of claim 2, wherein the activatable antibody comprises a light chain comprising the amino acid sequence of SEQ ID NO: 428 and a heavy chain comprising the amino acid sequence of SEQ ID NO: 432.
23. A conjugated antibody or activatable antibody comprising the antibody of any one of claims 1, 4, 7-9, or 14 or the activatable antibody of any one of claims 2 to 22 conjugated to an agent.
24. A pharmaceutical composition comprising the antibody of any one of claims 1, 4, 7-9, or 14 or the activatable antibody of any one of claims 2 to 22, or the conjugated antibody or conjugated activatable antibody of claim 23 and a carrier.
25. The pharmaceutical composition of claim 24 comprising an additional agent.
26. An isolated nucleic acid molecule encoding the antibody of any one of claims 1, 4, 7-9, or 14 or the activatable antibody of any one of claims 2 to 22.
27. A vector comprising the isolated nucleic acid molecule of claim 26.
28. A method of producing an activatable antibody by culturing a cell under conditions that lead to expression of the activatable antibody, wherein the cell comprises the nucleic acid molecule of claim 26, and optionally conjugating an agent to the activatable antibody.

29. The antibody of any of claims 1, 4, 7-9, or 14, or the activatable antibody of any one of claims 2 to 22, the conjugated antibody or activatable antibody of claim 23, or the pharmaceutical composition of any of claims 24-25 for use in the treatment of a cancer.