
What is claimed is:

1. A compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof:



(I)

Wherein:

R_{11}, R_{12}, R_{13}, R_{14} or R_{15} is independently selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl, each hydrogen in R_{11}, R_{12}, R_{13}, R_{14} or R_{15} is independently optionally substituted with 1, 2, 3, 4, 5 or 6 substituent(s) selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl; each heterocyclic or heteroaryl at each occurrence independently contains 1, 2, 3 or 4 heteroatom(s) selected from N, O, S, S=O or $S(=O)_2$;


R_{21} or R_{22} is independently selected from hydrogen, halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl, each hydrogen in R_{21} or R_{22} is independently optionally substituted with 1, 2, 3, 4, 5 or 6 substituent(s) selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl; each heterocyclic or heteroaryl at each occurrence independently contains 1, 2, 3 or 4 heteroatom(s) selected from N, O, S, S=O or $S(=O)_2$;

R_3 is selected from -C_{1-14}alkyl, -C_{2-14}alkenyl, -C_{2-14}alkynyl, -C_{6-10}aryl, 5-10 membered heteroaryl, 3-14 membered heterocyclic, -C_{3-14}carbocyclic, 

or 


, each ring C at each occurrence is independently selected from a C_{3-14} carbocyclic or 3-14 membered heterocyclic ring, each ring D at each occurrence is independently selected from a C_{6-10} aryl or 5-10 membered heteroaryl ring, each hydrogen in R_3 is independently optionally substituted with 1, 2, 3, 4, 5 or 6 R_{31}; each heterocyclic or heteroaryl at each occurrence independently contains 1, 2, 3 or 4 heteroatoms selected from N, O, S, S=O or $S(=O)_2$;



Each R_{31} at each occurrence is independently selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -C(=O)R_5, -C(=O)OR_5, -OC(=O)R_5, -C(=O)NR_5R_6, -NR_5C(=O)R_6, -$NR_5SO_2R_6$, -SO_2R_5, -S(=O)$_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl, each hydrogen in R_{31} is independently optionally substituted with 1, 2, 3, 4, 5 or 6 substituent(s) selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -C(=O)R_5, -C(=O)OR_5, -OC(=O)R_5, -C(=O)NR_5R_6, -NR_5C(=O)R_6, -S(=O)$_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl; each heterocyclic or heteroaryl at each occurrence independently contains 1, 2, 3 or 4 heteroatom(s) selected from N, O, S, S=O or S(=O)$_2$;

R_4 is selected from ;


Each G_1, G_2, G_3 or G_4 at each occurrence is independently selected from N or CH;

Each n1, n2, n3, n4 or n5 at each occurrence is independently selected from 0, 1, 2, 3, 4, 5 or 6, provided that n1 and n2 is not 0 at the same time, n3 and n4 is not 0 at the same time;

Each hydrogen in  is independently optionally substituted with 1R_{42}, 2R_{42}, 3R_{42}, 4R_{42}, 5R_{42} or 6R_{42};

Each R_{41} at each occurrence is independently selected from  or ;

Each of Q at each occurrence is independently selected from C(=O), NR_5C(=O), S(=O)$_2$ or NR_5S(=O)$_2$;

 in is selected from or ;

when is , R_{4a}, R_{4b} or R_{4c} is independently selected from hydrogen, halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -C(=O)R_5, -C(=O)OR_5, -OC(=O)R_5, -C(=O)NR_5R_6, -NR_5C(=O)R_6, -$NR_5SO_2R_6$, -SO_2R_5, -S(=O)$_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl, each hydrogen in R_{4a}, R_{4b} or R_{4c} is independently optionally substituted with 1, 2, 3, 4, 5 or 6 substituents selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -C(=O)R_5, -C(=O)OR_5, -OC(=O)R_5, -C(=O)NR_5R_6, -NR_5C(=O)R_6, -$NR_5SO_2R_6$, -SO_2R_5, -S(=O)$_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl; each

heterocyclic or heteroaryl at each occurrence independently contains 1, 2, 3 or 4 heteroatom(s) selected from N, O, S, S=O or $S(=O)_2$; or

when ⋯ is ═, R_{4a} is selected from hydrogen, halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl, each hydrogen in R_{4a} is independently optionally substituted with 1, 2, 3, 4, 5 or 6 substituents selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl; and R_{4b} and R_{4c} together with the carbon which they both attach to form a C_{3-10} carbocyclic ring or a 3-10 membered heterocyclic ring, each hydrogen in the C_{3-10} carbocyclic ring or the 3-10 membered heterocyclic ring is optionally substituted by 1, 2, 3, 4, 5 or 6 substituents selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl; each heterocyclic or heteroaryl at each occurrence independently contains 1, 2, 3 or 4 heteroatom(s) selected from N, O, S, S=O or $S(=O)_2$; or

when ⋯ is ═, R_{4a} and R_{4c} with the carbon they respectively attach to form a C_{3-10} carbocyclic ring or a 3-10 membered heterocyclic ring, each hydrogen in the C_{3-10} carbocyclic ring or the 3-10 membered heterocyclic ring is optionally substituted by 1, 2, 3, 4, 5 or 6 substituents selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl; and R_{4b} is selected from hydrogen, halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl, each hydrogen in R_{4b} is independently optionally substituted with 1, 2, 3, 4, 5 or 6 substituents selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl; each heterocyclic or heteroaryl at each occurrence independently contains 1, 2, 3 or 4 heteroatom(s) selected from N, O, S, S=O or $S(=O)_2$;

when ⋯ is ═, R_{4a} is absent, R_{4b} is absent, R_{4c} is selected from hydrogen, halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl, each hydrogen in R_{4c} is independently optionally substituted with 1, 2, 3, 4, 5 or 6 substituents selected from halogen,

-C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl; each heterocyclic or heteroaryl at each occurrence independently contains 1, 2, 3 or 4 heteroatom(s) selected from N, O, S, S=O or $S(=O)_2$;

R_{4d} is halogen;

Each R_{42} at each occurrence is independently selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl, each hydrogen in R_{42} is independently optionally substituted with 1, 2, 3, 4, 5 or 6 substituents selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_6$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -POR_5R_6, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl; or

Two R_{42} together with the atom which they both or respectively attach to form a C_{3-6} carbocyclic or 3-6 membered heterocyclic ring, each hydrogen in the C_{3-6} carbocyclic or 3-6 membered heterocyclic ring is independently optionally substituted with 1, 2, 3, 4, 5 or 6 substituents selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_5R_6, -$C(=O)R_5$, -$C(=O)OR_5$, -$OC(=O)R_5$, -$C(=O)NR_5R_6$, -$NR_5C(=O)R_5$, -$NR_5SO_2R_6$, -SO_2R_5, -$S(=O)_2NR_5R_6$, -$PO(R_5)_2$, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl; each heterocyclic and heteroaryl at each occurrence independently contains 1, 2, 3 or 4 heteroatom(s) selected from N, O, S, S=O or $S(=O)_2$;


Each R_5 and R_6 at each occurrence is independently selected from hydrogen or -C_{1-6}alkyl; or

R_5 and R_6 together with the atom which they both or respectively attach to form a 3-10 membered heterocyclic ring, the 3-10 membered heterocyclic ring is optionally further contains 1, 2, 3 or 4 heteroatoms selected from N, O, S, S(=O) or $S(=O)_2$, and each hydrogen in the 3-10 membered heterocyclic ring is independently optionally substituted with $1R_{51}$, $2R_{51}$, $3R_{51}$, $4R_{51}$, $5R_{51}$ or $6R_{51}$;

Each R_{51} is selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_7R_8, -$C(=O)R_7$, -$C(=O)OR_7$, -$OC(=O)R_7$, -$C(=O)NR_7R_8$, -$NR_7C(=O)R_8$, -$NR_7SO_2R_8$, -SO_2R_7, -$S(=O)_2NR_7R_8$, -POR_7R_8, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl, each hydrogen in R_{51} is independently optionally substituted with 1, 2, 3, 4, 5 or 6 substituent(s) selected from halogen, -C_{1-6}alkyl, -C_{2-6}alkenyl, -C_{2-6}alkynyl, heteroC_{2-6}alkyl, -CN, oxo, -OC_{1-6}alkyl, -SC_{1-6}alkyl, -NR_7R_8, -$C(=O)R_7$, -$C(=O)OR_7$, -$OC(=O)R_7$, -$C(=O)NR_7R_8$, -$NR_7C(=O)R_8$, -$NR_7SO_2R_8$, -SO_2R_7, -$S(=O)_2NR_7R_8$, -POR_7R_8, -C_{3-6}carbocyclic, 3-6 membered heterocyclic, -C_{6-10}aryl, or 5-10 membered heteroaryl; each heterocyclic or heteroaryl at each occurrence independently contains 1, 2, 3 or 4 heteroatom(s) selected from N, O, S, S=O or $S(=O)_2$;

Each R_7 and R_8 at each occurrence is independently selected from hydrogen or -C_{1-6}alkyl.

2. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to claim 1:



(I)

Wherein:

R_{11}, R_{12}, R_{13}, R_{14} or R_{15} is independently selected from -OH; halogen; $-NR_aR_b$; -C_{1-6}alkyl; -OC_{1-6}alkyl; -C_{1-6}alkylene-OH; -C_{1-6}alkylene-O-C_{1-6}alkyl; -C_{1-6}alkyl substituted with halogen, $-NH_2$, -CN or -OH; -O-C_{1-6}alkyl substituted with halogen, $-NH_2$, -CN or -OH; -O-C_{1-6}alkyl; $-SO_2R_a$; -CN; $-C(=O)NR_aR_b$; $-C(=O)R_a$; $-OC(=O)R_a$; $-C(=O)OR_a$; or -C_{3-6}carbocyclic;



R_a or R_b is independently selected from hydrogen or -C_{1-6}alkyl;

R_{21} is selected from hydrogen; halogen; -C_{1-6}alkyl; -C_{1-6}alkyl substituted with halogen, $-NH_2$, -CN or -OH; -C_{2-6}alkenyl; or -C_{3-6}carbocyclic;


R_{22} is selected from hydrogen; halogen; -C_{1-6}alkyl; -C_{1-6}alkyl substituted with halogen, $-NH_2$, -CN or -OH; -C_{2-6}alkenyl; or -C_{3-6}carbocyclic;

R_3 is selected from -C_{6-10}aryl or 5-10 membered heteroaryl, each of 5-10 membered heteroaryl at each occurrence independently contains 1, 2, 3 or 4 heteroatoms selected from N, O or S, each hydrogen in the -C_{6-10}aryl or 5-10 membered heteroaryl at each occurrence is independently optionally substituted with $1R_{31}$, $2R_{31}$, $3R_{31}$, $4R_{31}$, $5R_{31}$ or 6 R_{31};

Each R_{31} at each occurrence is independently selected from halogen, -C_{1-6}alkyl, -CN, -OH, -O-C_{1-6}alkyl, $-NH_2$, -NH(C_{1-6}alkyl), -N(C_{1-6}alkyl)$_2$ or -C_{3-6}carbocyclic;

R_4 is , each hydrogen in the  is independently optionally substituted with $1R_{42}$, $2R_{42}$, $3R_{42}$, $4R_{42}$, $5R_{42}$ or 6 R_{42};

n1 or n2 is independently selected from 1, 2, 3, 4, 5 or 6;

R_{41} is selected from ;

R_{4a}, R_{4b} or R_{4c} is independently selected from hydrogen, halogen, -C_{1-6}alkyl or -C_{1-6}alkylene-N(C_{1-6}alkyl)$_2$;

Each R_{42} at each occurrence is independently selected from -C_{1-6}alkyl; -C_{1-6}alkylene-CN or -C_{1-6}alkyl substituted with halogen.

3. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically

acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to claim 2, wherein:

R_{11}, R_{12}, R_{13}, R_{14} or R_{15} is independently selected from -OH; -F; -Cl; -Br; $-NR_aR_b$; $-C_{1-3}$alkyl; $-OC_{1-3}$alkyl; $-C_{1-3}$alkylene-OH; $-C_{1-3}$alkylene-O-C_{1-3}alkyl; $-C_{1-3}$alkyl substituted with -F or -Cl; -O-C_{1-3}alkyl substituted with -F or -Cl; $-SO_2R_a$; -CN; $-C(=O)NR_aR_b$; $-C(=O)R_a$; $-OC(=O)R_a$; $-C(=O)OR_a$; 3-membered carbocyclic; 4-membered carbocyclic; 5-membered carbocyclic or 6-membered carbocyclic;

R_a or R_b is independently selected from hydrogen or $-C_{1-3}$alkyl.


4. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to claim 2 or 3, wherein:


R_{11}, R_{12}, R_{13}, R_{14} or R_{15} is independently selected from -OH, -F, -Cl, $-NH_2$, $-NHCH_3$, $-N(CH_3)_2$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-OCH_3$, $-OCH_2CH_3$, $-OCH_2CH_2CH_3$, $-OCH(CH_3)_2$, $-CH_2OH$, $-CH_2CH_2OH$, $-CH_2OCH_3$, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCH_2F$, $-OCHF_2$, $-OCF_3$, $-SO_2CH_3$, -CN, $-C(=O)NH_2$, $-C(=O)CH_3$, $-OC(=O)CH_3$, $-C(=O)OCH_3$ or 3-membered carbocyclic.


5. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-4, wherein:

R_{11}, R_{12}, R_{13}, R_{14} or R_{15} is independently selected from -OH, -F, -Cl, $-NH_2$, $-CH_3$ or $-CF_3$.

6. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-5, wherein:

the  in the formula (I) is selected from:





or .

7. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-6, wherein:

R_{21} is selected from hydrogen; -F; -Cl; -Br; -C_{1-3}alkyl; -C_{1-3}alkyl substituted with -F or -Cl; -C_{2-3}alkenyl; or -C_{3-6}carbocyclic.

8. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-7, wherein:

R_{21} is selected from hydrogen; -F; -Cl; methyl; ethyl; propyl; isopropyl; methyl substituted with -F; ethyl substituted with -F; propyl substituted with -F; isopropyl substituted with -F; ethenyl; propenyl; 3 membered carbocyclic; 4 membered carbocyclic; 5 membered carbocyclic; or 6 membered carbocyclic.

9. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-8, wherein:

R_{21} is selected from -Cl.

10. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-9, wherein:

R_{22} is selected from hydrogen; -F; -Cl; -Br; -C_{1-3}alkyl; -C_{1-3}alkyl substituted with -F or -Cl; -C_{2-3}alkenyl; or -C_{3-6}carbocyclic.

11. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-10, wherein:

R_{22} is selected from hydrogen; -F; -Cl; methyl; ethyl; propyl; isopropyl; methyl substituted with -F; ethyl substituted with -F; propyl substituted with -F; isopropyl substituted with -F; ethenyl; propenyl; 3 membered carbocyclic; 4 membered carbocyclic; 5 membered carbocyclic; or 6 membered carbocyclic.

12. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-11, wherein:

R_{22} is hydrogen.

13. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-12, wherein:

R_3 is selected from phenyl, naphthyl, 5 membered heteroaryl, 6 membered heteroaryl, 7 membered heteroaryl, 8 membered heteroaryl, 9 membered heteroaryl or 10 membered heteroaryl, each heteroaryl at each occurrence independently contains 1, 2 or 3 heteroatoms selected from N or O, each hydrogen in R_3 at each occurrence is independently optionally substituted with $1R_{31}$, $2R_{31}$, $3R_{31}$, $4R_{31}$ or $5R_{31}$.

14. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-13, wherein:

R_3 is selected from phenyl or 6 membered heteroaryl, the heteroaryl contains 1 or 2 heteroatoms selected from N, each hydrogen in the phenyl or 6 membered heteroaryl at each occurrence is independently optionally substituted by $1R_{31}$, $2R_{31}$, $3R_{31}$ or $4R_{31}$.

15. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-14, wherein:

R_3 is selected from , , , each hydrogen in the R_3 at each occurrence is independently optionally substituted by $1R_{31}$, $2R_{31}$, $3R_{31}$ or $4R_{31}$.

16. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-15, wherein:

Each R_{31} at each occurrence is independently selected from -F, -Cl, -Br, -C_{1-3}alkyl, -CN, -OH, -O-C_{1-3}alkyl, -NH_2, -NH(C_{1-3}alkyl), -N(C_{1-3}alkyl)$_2$ or -C_{3-6}carbocyclic.

17. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-16, wherein:


Each R_{31} at each occurrence is independently selected from -F, -Cl, methyl, ethyl, propyl, isopropyl, -CN, -OH, methoxy, ethoxy, propoxy, isopropoxy, -NH_2, -$NHCH_3$, -$NHCH_2CH_3$, -NH($CH_2CH_2CH_3$), -NH($CH(CH_3)_2$), -N(CH_3)$_2$, -N(CH_2CH_3)$_2$, -N(CH_3)(CH_2CH_3), 3 membered carbocyclic, 4 membered carbocyclic, 5 membered carbocyclic or 6 membered carbocyclic.

18. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-17, wherein:


Each R_{31} at each occurrence is independently selected from methyl or isopropyl.

19. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a



pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-18, wherein:

R_3 is selected from .

20. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-19, wherein:

R_3 is .



21. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-20, wherein:

R_4 is selected from , each hydrogen in the  is independently optionally substituted with $1R_{42}$, $2R_{42}$, $3R_{42}$ or $4R_{42}$;

n1 is selected from 1, 2 or 3;

n2 is selected from 1, 2 or 3.



22. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-21, wherein:

R_4 is selected from , each hydrogen in the  is independently optionally substituted with $1R_{42}$ or $2R_{42}$;


n1 is selected from 1 or 2;

n2 is selected from 1 or 2.

23. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-22, wherein:

R_4 is , each hydrogen in the  is independently optionally substituted with $1R_{42}$ or $2R_{42}$.

24. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-23, wherein:

R_{41} is  ;

R_{4a}, R_{4b}, or R_{4c} is independently selected from hydrogen, -F, -Cl, -Br, -C_{1-3}alkyl or -C_{1-3}alkylene-N(C_{1-3}alkyl)$_2$.

25. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-24, wherein:

R_{4a}, R_{4b} or R_{4c} is independently selected from hydrogen, -F, -Cl, methyl, ethyl, propyl, isopropyl, -CH_2-N(CH_3)$_2$, -CH_2-N(CH_2CH_3)$_2$ or -CH_2-N(CH_3)(CH_2CH_3).

26. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-25, wherein:

R_{4a}, R_{4b} or R_{4c} is independently selected from hydrogen, -F, methyl or -CH_2-N(CH_3)$_2$.

27. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-26, wherein:


R_{4a} is selected from hydrogen or -F;

R_{4b} is hydrogen;



R_{4c} is selected from hydrogen or -CH_2-N(CH_3)$_2$.

28. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-27, wherein:

R_{41} is selected from:

 .

29. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-28, wherein:

R_4 is selected from  , each hydrogen in the  at each occurrence is independently optionally substituted with 1R_{42} or 2R_{42}.

30. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a

pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-29, wherein:

Each R_{42} at each occurrence is independently selected from -C_{1-3}alkyl; -C_{1-3}alkylene-CN; or -C_{1-3}alkyl substituted with -F or -Cl.

31. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-30, wherein:

Each R_{42} at each occurrence is independently selected from methyl; ethyl; propyl; isopropyl; -methylene-CN; -ethylene-CN; -propylene-CN; methyl substituted with -F; ethyl substituted with -F; propyl substituted with -F; or isopropyl substituted with -F.

32. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-31, wherein:


Each R_{42} at each occurrence is independently selected from methyl; ethyl; -methylene-CN or methyl substituted with -F.


33. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-32, wherein:

Each R_{42} at each occurrence is independently selected from -CH_3, -CH_2CH_3, -CH_2CN, -CHF_2 or -CF_3.

34. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-33, wherein:


R_4 is selected from:






35. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-34, wherein:


R_4 is selected from:





36. The compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 2-35, wherein:


the compound is selected from:


























































37. A method for preparing the compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 1-36, wherein the method comprises a coupling reaction between a compound of formula (II) and a compound of formula(III) according to the following reaction Scheme 1 or between a compound of formula (II') and a compound of formula(III') according to the following reaction Scheme 2 catalyzed by a transition metal palladium or nickel reagent:

Scheme 1:




Scheme 2:



Wherein

the L in the compound of formula(III) or formula(II') is a leaving group; preferably, the leaving group is selected from halogen, $-OS(O)_2CF_3$ or -OTs; more preferably, the halogen is selected from -F, -Cl, -Br, or -I; more preferably, the leaving group is -Cl or -Br;

the X in the compound of formula (II) or formula(III') is selected from boronic acid, borate ester or organotin; more preferably, the X is selected from ;

preferably, the coupling reaction is Suzuki coupling reaction or Stille coupling reaction;

preferably, the coupling reaction is catalyzed by the transition metal palladium reagent; more preferably, the transition metal palladium reagent is $Pd(PPh_3)_4$.

38. A pharmaceutical composition comprising the compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 1-36, and at least one pharmaceutically acceptable excipient.

39. The pharmaceutical composition according to claim 38, wherein the said compound in a weight ratio to the said excipient within the range from about 0.0001 to about 10.

40. Use of the compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 1-36; or the pharmaceutical composition according to claim 38 or 39 for the manufacture of a medicament for the treatment of diseases or conditions related to KRAS mutant protein.

41. The use according to claim 40, wherein the diseases or conditions related to KRAS mutant protein is the diseases or conditions related to KRAS G12C mutant protein.

42. The use according to claim 41, wherein the diseases or conditions related to KRAS G12C mutant protein is cancer related to KRAS G12C mutant protein.

43. The use according to claim 42, wherein the cancer is selected from blood cancer, pancreatic cancer, colon cancer, rectal cancer, colorectal cancer or lung cancer.

44. The use according to claim 43, wherein the blood cancer is selected from acute myeloid leukemia or acute lymphocytic leukemia; the lung cancer is selected from non-small cell lung cancer or small cell lung cancer.

45. A method of treating a subject having a diseases or conditions related to KRAS mutant protein, said method comprising administering to the subject a therapeutically effective amount of the compound of formula (I), a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof or a pharmaceutically acceptable salt of the atropisomer thereof according to any one of claims 1-36; or the pharmaceutical composition according to claim 38 or 39.

46. The method according to claim 45, wherein the diseases or conditions related to KRAS mutant protein is the diseases or conditions related to KRAS G12C mutant protein.

47. The method according to claim 46, wherein the diseases or conditions related to KRAS G12C mutant protein is cancer related to KRAS G12C mutant protein.

48. The method according to claim 47, wherein the cancer is selected from blood cancer, pancreatic cancer, colon cancer, rectal cancer, colorectal cancer or lung cancer.

49. The method according to claim 48, wherein the blood cancer is selected from acute myeloid leukemia or acute lymphocytic leukemia; the lung cancer is selected from non-small cell lung cancer or small cell lung cancer.