

PROCESSES FOR MAKING COMPOUNDS USEFUL AS INHIBITORS OF ATR KINASE

BACKGROUND OF THE INVENTION

[0001] ATR (“ATM and Rad3 related”) kinase is a protein kinase involved in cellular responses to DNA damage. ATR kinase acts with ATM (“ataxia telangiectasia mutated”) kinase and many other proteins to regulate a cell’s response to DNA damage, commonly referred to as the DNA Damage Response (“DDR”). The DDR stimulates DNA repair, promotes survival and stalls cell cycle progression by activating cell cycle checkpoints, which provide time for repair. Without the DDR, cells are much more sensitive to DNA damage and readily die from DNA lesions induced by endogenous cellular processes such as DNA replication or exogenous DNA damaging agents commonly used in cancer therapy.

[0002] Healthy cells can rely on a host of different proteins for DNA repair including the DDR kinase ATR. In some cases these proteins can compensate for one another by activating functionally redundant DNA repair processes. On the contrary, many cancer cells harbour defects in some of their DNA repair processes, such as ATM signaling, and therefore display a greater reliance on their remaining intact DNA repair proteins which include ATR.

[0003] In addition, many cancer cells express activated oncogenes or lack key tumour suppressors, and this can make these cancer cells prone to dysregulated phases of DNA replication which in turn cause DNA damage. ATR has been implicated as a critical component of the DDR in response to disrupted DNA replication. As a result, these cancer cells are more dependent on ATR activity for survival than healthy cells. Accordingly, ATR inhibitors may be useful for cancer treatment, either used alone or in combination with DNA damaging agents, because they shut down a DNA repair mechanism that is more important for cellular survival in many cancer cells than in healthy normal cells.

[0004] In fact, disruption of ATR function (e.g. by gene deletion) has been shown to promote cancer cell death both in the absence and presence of DNA damaging agents. This suggests that ATR inhibitors may be effective both as single agents and as potent sensitizers to radiotherapy or genotoxic chemotherapy.

[0005] For all of these reasons, there is a need for the development of potent and selective ATR inhibitors for the treatment of cancer, either as single agents or as combination therapies with radiotherapy or genotoxic chemotherapy. Furthermore, it would be desirable to have a synthetic route to ATR inhibitors that is amenable to large-scale synthesis and improves upon currently known methods.

[0006] ATR peptide can be expressed and isolated using a variety of methods known in the literature (see e.g., Ünsal-Kaçmaz et al, *PNAS* 99: 10, pp6673-6678, May 14, 2002; see also Kumagai et al. *Cell* 124, pp943-955, March 10, 2006; Unsal-Kacmaz et al. *Molecular and Cellular Biology*, Feb 2004, p1292-1300; and Hall-Jackson et al. *Oncogene* 1999, 18, 6707-6713).

DESCRIPTION OF THE FIGURES

FIGURE 1a: XRPD Compound I-2 free base

FIGURE 2a: TGA Compound I-2 free base

FIGURE 3a: DSC Compound I-2 free base

FIGURE 4a: ORTEP plot of the asymmetric unit of the Compound I-2 free form single crystal structure

FIGURE 1b: XRPD Compound I-2 • HCl

FIGURE 2b: TGA Compound I-2 • HCl

FIGURE 3b: DSC Compound I-2 • HCl

FIGURE 4b: ORTEP plot of the asymmetric unit of the Compound I-2 • HCl anhydrous structure.

FIGURE 1c: XRPD Compound I-2 • 2HCl

FIGURE 2c: TGA Compound I-2 • 2HCl

FIGURE 3c: DSC Compound I-2 • 2HCl

FIGURE 1d: XRPD Compound I-2 • HCl monohydrate

FIGURE 2d: TGA Compound I-2 • HCl monohydrate

FIGURE 3d: DSC Compound I-2 • HCl monohydrate

FIGURE 1e: XRPD Compound I-2 • HCl • 2H₂O

FIGURE 2e: TGA Compound I-2 • HCl • 2H₂O

FIGURE 3e: DSC Compound I-2 • HCl • 2H₂O


FIGURE 4a: Solid State Compound I-1 free base

FIGURE 4b: Solid State ¹³CNMR of Compound I-1 • HCl


SUMMARY OF THE INVENTION

[0007] The present invention relates to processes and intermediates for preparing compounds useful as inhibitors of ATR kinase, such as aminopyrazine-isoxazole derivatives and related molecules. Aminopyrazine-isoxazole derivatives are useful as ATR inhibitors and are also useful for preparing ATR inhibitors. The present invention also relates to solid forms of ATR inhibitors as well as deuterated ATR inhibitors.


[0008] One aspect of the invention provides a process for preparing a compound of formula I:



comprising preparing a compound of formula 4:




from a compound of formula 3:




under suitable oxime formation conditions.

[0009] Another aspect comprises preparing a compound of formula 4:




from a compound of formula 3:



under suitable oxime formation conditions.


[0010] Another aspect of the present invention comprises a compound of formula II:



II

or a pharmaceutically acceptable salt thereof, wherein each R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10} , R^{11} , R^{12} , and R^{13} is independently hydrogen or deuterium, and at least one of R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10} , R^{11} , R^{12} , and R^{13} is deuterium.

[0011] Yet another aspect of the invention provides solid forms of a compound of formula I-2:



I-2


[0012] Other aspects of the invention are set forth herein.

[0013] The present invention has several advantages over previously known methods. First, the present process has fewer number of total synthetic steps compared with previously disclosed processes. Second, the present process has improved yields over previously disclosed processes. Third, the present process is effective for compounds wherein R^3 is a wide range of groups, such as alkyl groups or a large, hindered moiety, such as a ring. Fourth,


the present process comprises intermediates which are more stable and have a longer shelf life. In certain embodiments, the non-acidic formation of the oxime group in the present process allows the preservation of acid-sensitive protecting groups such as Boc or CBz during the course of the synthesis. In other embodiments, the process is more easily scaled up to larger quantities due to the elimination of chromatography as a purification step.

DETAILED DESCRIPTION OF THE INVENTION

[0014] One aspect of the invention provides a process for making a compound of preparing a compound of formula 4:



from a compound of formula 3:



under suitable oxime formation conditions;

wherein

R¹ is C₁₋₆alkyl;

R² is C₁₋₆alkyl;


or R¹ and R², together with the oxygen atoms to which they are attached, form an optionally substituted 5 or 6 membered saturated heterocyclic ring having two oxygen atoms;

R³ is hydrogen, C₁₋₆alkyl, or a 3-6 membered saturated or partially unsaturated heterocyclyl having 1-2 heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur; wherein the heterocyclyl is optionally substituted with 1 occurrence of halo or C₁₋₃alkyl;

J¹ is halo, C₁₋₄alkyl, or C₁₋₄alkoxy;

PG is a carbamate protecting group.


[0015] Another aspect provides a process for preparing a compound of formula I:




I

comprising the steps of:

preparing a compound of formula 4:



from a compound of formula 3:



under suitable oxime formation conditions;


wherein

R¹ is C₁₋₆alkyl;

R² is C₁₋₆alkyl;

or R¹ and R², together with the oxygen atoms to which they are attached, form an optionally substituted 5 or 6 membered saturated heterocyclic ring having two oxygen atoms;

R³ is hydrogen, C₁₋₆alkyl, or a 3-6 membered saturated or partially unsaturated heterocyclyl having 1-2 heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur; wherein the heterocyclyl is optionally substituted with 1 occurrence of halo or C₁₋₃alkyl;



Q is phenyl, pyridyl, or an N-alkylated pyridine;


J¹ is H, halo, C₁₋₄alkyl, or C₁₋₄alkoxy;

J^2 is halo; CN; phenyl; oxazolyl; or a C_{1-6} aliphatic group wherein up to 2 methylene units are optionally replaced with O, NR'' , $C(O)$, S, $S(O)$, or $S(O)_2$; said C_{1-6} aliphatic group is optionally substituted with 1-3 fluoro or CN;

q is 0, 1, or 2;


PG is a carbamate protecting group.

[0016] Another embodiment further comprises the step of protecting a compound of formula 2:



under suitable protection conditions to form the compound of formula 3.

[0017] Another embodiment further comprises the step of reacting a compound of formula 1:




with a suitable amine under suitable reductive amination conditions to form a compound of formula 2.


[0018] In some embodiments, the suitable amine is $NHCH_3$. In other embodiments, the

suitable amine is .

[0019] Another embodiment further comprises the step of reacting a compound of formula 4:




under suitable isoxazole formation conditions to form a compound of formula 5:




[0020] Another embodiment further comprises the step of reacting a compound of formula **5** under suitable coupling conditions followed by suitable deprotection conditions to form a compound of formula I.

[0021] In some embodiments, PG is Boc or Cbz. In some embodiments, PG is Boc.

[0022] In other embodiments, R¹ is ethyl and R² ethyl.

[0023] In yet other embodiments, R³ is CH₃ or .

[0024] In some embodiments, R⁴ is ; wherein Q is phenyl. In some embodiments, Q is substituted in the para position with J₂, wherein q is 1.


[0025] In some embodiments, J¹ is H or halo. In some embodiments, J¹ is H. In other embodiments, J¹ is halo.

[0026] In other embodiments, J² is a C₁₋₆aliphatic group wherein up to 1 methylene unit is optionally replaced with S(O)₂. In some embodiments, J² is -S(O)₂-(C₁₋₅alkyl). In some embodiments, q is 1.

[0027] According to another embodiment,


R¹ is ethyl;

R² is ethyl;


R³ is CH₃ or .

PG is Boc or Cbz;

J¹ is H;

R⁴ is  wherein Q is phenyl; J² is -S(O)₂-CH(CH₃)₂;
q is 1;


[0028] In some embodiments, R^3 is CH_3 . In some embodiments, R^3 is CH_3 . In yet

another embodiments, R^3 is CH_3 or .

[0029] According to another embodiment,



R^1 is ethyl;

R^2 is ethyl;

R^3 is  ;


PG is Boc;

J^1 is H;

R^4 is  wherein Q is pyridyl; J^2 is .

q is 1;

[0030] In some embodiments, R^4 is



Reactions Conditions

[0031] In some embodiments, the suitable oxime formation conditions consist of either a single step sequence or a two step sequence.

[0032] In some embodiments, the two step sequence consists of first deprotecting the ketal group in the compound of formula 3 into an aldehyde under suitable deprotection conditions, and then forming the oxime of formula 4 under suitable oxime formation conditions. In some embodiments, suitable deprotection conditions comprise adding catalytic amounts of para-toluenesulfonic acid (pTSA), acetone, and water; and suitable oxime formation conditions comprise mixing together hydroxylamine, a catalytic amount of acid, a dehydrating agent, and an alcoholic solvent. In other embodiments, the acid is pTSA or HCl, the dehydrating agent is molecular sieves or dimethoxyacetone, and the alcoholic solvent is methanol or ethanol.

[0033] In other embodiments, the single step sequence comprises adding $NH_2OH.HCl$ and a mixture of THF and water. In other embodiments, the sequence comprises adding

NH₂OH.HCl with a mixture of 2-methyl tetrahydrofuran and water optionally buffered with Na₂SO₄. In some embodiments, 1 equivalent of the compound of formula **3** is combined with a 1.1 equivalents of NH₂OH.HCl in a 10:1 v/v mixture of THF and water. In some embodiments, 1 equivalent of the compound of formula **3** is combined with a 1.1 equivalents of NH₂OH.HCl in a 10:1 v/v mixture of 2-methyl tetrahydrofuran and water optionally buffered with Na₂SO₄.

[0034] In other embodiments, the protection conditions are selected from the group consisting of

- R-OCOCl, a suitable tertiary amine base, and a suitable solvent; wherein R is C₁₋₆alkyl optionally substituted with phenyl;
- R(CO₂)OR', a suitable solvent, and optionally a catalytic amount of base, wherein R is and R' are each independently C₁₋₆alkyl optionally substituted with phenyl;
- [RO(C=O)]₂O, a suitable base, and a suitable solvent.

[0035] In some embodiments, the suitable base is Et₃N, diisopropylamine, and pyridine; and the suitable solvent is selected from a chlorinated solvent, an ether, or an aromatic hydrocarbon. In other embodiments, the suitable base is Et₃N, the suitable solvent is a chlorinated solvent selected from DCM. In yet other embodiments, the protection conditions comprise adding 1.20 equivalents of (Boc)₂O and 1.02 equivalents of Et₃N in DCM.

[0036] According to another embodiment suitable coupling conditions comprise adding a suitable metal and a suitable base in a suitable solvent. In other embodiments, the suitable metal is Pd[P(tBu)₃]₂; the suitable solvent is a mixture of acetonitrile and water; and the suitable base is sodium carbonate. In yet other embodiments, the suitable coupling conditions comprise adding 0.1 equivalents of Pd[P(tBu)₃]₂; 1 equivalent of boronic acid or ester; and 2 equivalents of sodium carbonate in a 2:1 ratio v/v of acetonitrile/water at 60-70°C.

[0037] According to another embodiment, suitable deprotection conditions comprise combining the compound of formula **5** with a suitable acid in a suitable solvent. In some embodiments, the suitable acid is selected from para-toluenesulfonic acid (pTSA), HCl, TBAF, H₃PO₄, or TFA and the suitable solvent is selected from acetone, methanol, ethanol, CH₂Cl₂, EtOAc, THF, 2-MeTHF, dioxane, toluene, or diethylether.

[0038] According to another embodiment, suitable isoxazole-formation conditions consists of two steps, the first step comprising reacting the compound of formula **4** under suitable chlorooxime formation conditions to form a chlorooxime intermediate; the second

step comprising reacting the chlorooxime intermediate with acetylene under suitable cycloaddition conditions to form a compound of formula 5.

[0039] According to another embodiment, suitable chlorooxime formation conditions are selected from

- N-chlorosuccinimide and suitable solvent or
- potassium peroxymonosulfate, HCl, and dioxane.

[0040] In some embodiments, the suitable solvent is selected from a nonprotic solvent, an aromatic hydrocarbon, or an alkyl acetate. According to another embodiment, the suitable chlorooxime formation conditions are 1.05 equivalents of N-chlorosuccinimide in isopropylacetate at 40-50°C.


[0041] According to another embodiment, suitable cycloaddition conditions consist of a suitable base and a suitable solvent. In some embodiments, the suitable base is selected from pyridine, DIEA, TEA, t-BuONa, and K₂CO₃ and the suitable solvent is selected from acetonitrile, tetrahydrofuran, 2-methyltetrahydrofuran, MTBE, EtOAc, i-PrOAc, DCM, toluene, DMF, and methanol. In other embodiments, the suitable base is selected from Et₃N and the suitable solvent is selected from DCM.

[0042] According to another embodiment, the second step comprises reacting 1 equivalent of acetylene with 1.2 equivalents of the chlorooxime intermediate and 1.3 equivalents of Et₃N in DCM at room temperature.

[0043] According to another embodiment, suitable isoxazole-formation conditions comprise combining the compound of formula 4 with an oxidant in a suitable solvent. In some embodiments, said oxidant is [bis(trifluoroacetoxy)iodo] benzene and said solvent is a 1:1:1 mixture of methanol, water, and dioxane.

Synthesis of Compounds I-2 and I-3


[0044] One embodiment provides a process for preparing a compound of formula I-2:




I-2

comprising one or more of the following steps:


a) Reacting a compound of formula 1b:




with methylamine under suitable reductive amination conditions to form a compound of formula 2b:




b) reacting a compound of formula 2b under suitable Boc protection conditions to form the compound of formula 3b:




c) reacting a compound of formula 3b under suitable oxime formation conditions to form the compound of formula 4-i:



d) reacting a compound of formula 4-i under suitable chlorooxime formation conditions to form the compound of formula 4-ii:




e) reacting the compound of formula 4-ii with a compound of formula 4-iii




4-iii

under suitable cycloaddition conditions to form a compound of formula 4-iv:




4-iv

f) reacting a compound of formula 4-iv with a compound of formula A-5-i:



A-5-i


under suitable coupling conditions to form the compound of formula 5-i:



5-i


g) deprotecting a compound of formula 5-i under suitable Boc deprotection conditions optionally followed by treatment under basic aqueous conditions to form a compound of formula I-2.

[0045] Another embodiment provides a process for preparing a compound of formula I-3:




comprising one or more of the following steps:


- a) Reacting a compound of formula A-1:




with tetrahydro-2*H*-pyran-4-amine under suitable reductive amination conditions to form a compound of formula A-2:




- b) reacting a compound of formula A-2 under suitable Boc protection conditions to form the compound of formula A-3:



- c) reacting a compound of formula A-3 under suitable oxime formation conditions to form the compound of formula A-4:




- d) reacting a compound of formula A-4:




A-4

under suitable chlorooxime formation conditions to form the compound of formula A-4-i:




A-4-i

e) reacting the compound of formula A-4-i with a compound of formula A-4-ii:




A-4-ii

under suitable cycloaddition conditions to form the compound of formula A-5:




A-5

f) reacting a compound of formula A-5 with a compound of formula A-5-i:



A-5-i

under suitable coupling conditions to form the compound of formula A-6:



A-6

- g) deprotecting a compound of formula A-6 under suitable Boc deprotection conditions optionally followed by treatment under basic aqueous conditions to form a compound of formula I-3.

[0046] Suitable coupling conditions comprise combining a suitable palladium catalyst with a suitable base in a suitable solvent. Suitable palladium catalyst include, but are not limited to, Pd[P(tBu)₃]₂, Pd(dtbpf)Cl₂, Pd(PPh₃)₂Cl₂, Pd(PCy₃)₂Cl₂, Pd(dppf)Cl₂, and Pd(dppe)Cl₂. Suitable solvents include, but are not limited to, toluene, MeCN, water, EtOH, IPA, 2-Me-THF, or IPAc. Suitable bases include, but are not limited to, K₂CO₃, Na₂CO₃, or K₃PO₄.

[0047] Suitable oxime formation conditions consist of either a single step sequence or a two step sequence. The two step sequence consists of first deprotecting the ketal group in the compound of formula A-3 into an aldehyde under suitable deprotection conditions, and then forming the oxime of formula A-4 under suitable oxime formation conditions.

[0048] The single step sequence comprises, for example, comprise mixing together hydroxylamine, an acid, an organic solvent, and water. In some embodiments, NH₂OH.HCl is added to a mixture of THF and water. In some embodiments, 1 equivalent of the compound of formula 3-A is combined with a 1.1 equivalents of NH₂OH.HCl in a 10:1 v/v mixture of THF/water.

[0049] Suitable deprotection conditions comprise adding an acid, acetone, and water. Suitable acids include pTSA or HCl, suitable organic solvents include chlorinated solvents (e.g., dichloromethane (DCM), dichloroethane (DCE), CH₂Cl₂, and chloroform); an ether (e.g., THF, 2-MeTHF and dioxane); an aromatic hydrocarbons (e.g., toluene and xylenes, or other aprotic solvents.

[0050] Suitable cycloaddition conditions comprise a suitable base (e.g., pyridine, DIEA, TEA, t-BuONa, or K₂CO₃) and a suitable solvent (e.g., acetonitrile, tetrahydrofuran, 2-methyltetrahydrofuran, MTBE, EtOAc, i-PrOAc, DCM, toluene, DMF, and methanol).


[0051] Suitable chlorooxime formation conditions comprise adding HCl in dioxane to a solution of the oxime in the presence of NCS in a suitable solvent selected from a nonprotic solvents (DCM, DCE, THF, and dioxane), aromatic hydrocarbons (e.g. toluene, xylenes), and alkyl acetates (e.g., isopropyl acetate, ethyl acetate).

[0052] Suitable Boc deprotection conditions comprises adding a suitable Boc deprotecting agent (e.g, TMS-Cl, HCl, TBAF, H₃PO₄, or TFA) and a suitable solvent (e.g., acetone, toluene, methanol, ethanol, 1-propanol, isopropanol, CH₂Cl₂, EtOAc, isopropyl acetate, tetrahydrofuran, 2-methyltetrahydrofuran, dioxane, and diethylether). In some embodiments, the suitable Boc deprotection conditions comprises adding a suitable Boc deprotecting agent selected from HCl, TFA and a suitable solvent selected from acetone, toluene, isopropyl acetate, tetrahydrofuran, or 2-methyltetrahydrofuran.

[0053] Suitable Boc protection conditions include (Boc)₂O, a suitable base, and a suitable solvent. Suitable bases include, but are not limited to, Et₃N, diisopropylamine, and pyridine. Suitable solvents include, but are not limited to, chlorinated solvents (e.g., dichloromethane (DCM), dichloroethane (DCE), CH₂Cl₂, and chloroform); an ether (e.g., THF, 2-MeTHF and dioxane); an aromatic hydrocarbons (e.g., toluene and xylenes, or other aprotic solvents. In some embodiments, the suitable base is Et₃N, the suitable solvent is DCM, tetrahydrofuran or 2-methyltetrahydrofuran. In certain embodiments, the protection conditions comprise adding 1.05 equivalents of (Boc)₂O in 2-methyltetrahydrofuran or DCM.

[0054] Suitable reductive amination conditions comprise adding a reducing agent selected from NaBH₄, NaBH₃CN, or NaBH(OAc)₃ in the presence of a solvent selected from dichloromethane (DCM), dichloroethane (DCE), an alcoholic solvent selected from methanol, ethanol, 1-propanol, isopropanol, or a nonprotic solvent selected from dioxane, tetrahydrofuran, or 2-methyltetrahydrofuran and optionally a base selected from Et₃N or diisopropylethylamine. In some embodiments, the suitable reductive amination conditions comprise adding 1.2 equivalents of NaBH₄ caplets in the presence Et₃N in MeOH.

[0055] Another aspect of the present invention provides a compound of Formula II:



II

or a pharmaceutically acceptable salt thereof,

wherein each R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} is independently hydrogen or deuterium, and

at least one of R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} is deuterium.

[0056] In some embodiments, R^{9a} and R^{9b} are the same. In other embodiments, R^{9a} and R^{9b} are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{10} , R^{11a} , R^{11b} , R^{12a} , R^{12b} , R^{13a} , R^{13b} , R^{14a} , and R^{14b} are deuterium or hydrogen. In yet another embodiment, R^{9a} and R^{9b} are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{10} , R^{11a} , R^{11b} , R^{12a} , R^{12b} , R^{13a} , R^{13b} , R^{14a} , and R^{14b} are hydrogen.

[0057] In one embodiment, R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are the same. In another embodiment, R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , and R^8 are deuterium or hydrogen. In some embodiments, R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , and R^8 are hydrogen.

[0058] In other embodiments, R^{10a} , R^{10b} , and R^{10c} are the same. In one embodiment, R^{10a} , R^{10b} , and R^{10c} are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , and R^{9b} are deuterium or hydrogen. In yet another embodiment, R^{10a} , R^{10b} , and R^{10c} are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , and R^{9b} are hydrogen.

[0059] In some embodiments, R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , and R^{3c} are the same. In another embodiment R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , and R^{3c} are deuterium, and R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are deuterium or hydrogen. In yet another embodiment, R^{1a} , R^{1b} ,

R^{1c} , R^2 , R^{3a} , R^{3b} , and R^{3c} are deuterium, and R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are deuterium.

[0060] In another embodiment, R^6 is deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are deuterium or hydrogen. In yet another embodiment, R^6 is deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are hydrogen.

[0061] In other embodiments, R^2 is deuterium, and R^{1a} , R^{1b} , R^{1c} , R^{3a} , R^{3b} , and R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are deuterium or hydrogen. In another embodiment, R^2 is deuterium, and R^{1a} , R^{1b} , R^{1c} , R^{3a} , R^{3b} , and R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are hydrogen.

[0062] In another embodiment, R^7 is deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are deuterium or hydrogen. In other embodiments, R^7 is deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , R^{10c} are hydrogen.

[0063] In yet another embodiment, R^8 is deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are deuterium or hydrogen. In another embodiment, R^8 is deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , R^{10c} are hydrogen.

[0064] In some embodiments, at least one of R^{10a} , R^{10b} , or R^{10c} are the same. In another embodiment, at least one of R^{10a} , R^{10b} , or R^{10c} are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , and R^{9b} are deuterium or hydrogen. In yet another embodiment, at least one of R^{10a} , R^{10b} , or R^{10c} are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , and R^{9b} are hydrogen.

[0065] In some embodiments, at least two of R^{10a} , R^{10b} , or R^{10c} are the same. In another embodiment, at least two of R^{10a} , R^{10b} , or R^{10c} are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , and R^{9b} are deuterium or hydrogen. In yet another embodiment, at least two of R^{10a} , R^{10b} , or R^{10c} are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , and R^{9b} are hydrogen.

[0066] In another embodiment, R^{1a} , R^{1b} , R^{1c} , R^{3a} , R^{3b} , and R^{3c} are the same. In some embodiments, R^{1a} , R^{1b} , R^{1c} , R^{3a} , R^{3b} , and R^{3c} are deuterium, and R^2 , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are deuterium or hydrogen. In yet another embodiment, R^{1a} , R^{1b} , R^{1c} , R^{3a} , R^{3b} , and R^{3c} is deuterium, and R^2 , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are hydrogen.

[0067] In yet another embodiment, R^4 is deuterium, and $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}, R^5, R^6, R^7, R^8, R^9a, R^9b, R^{10a}, R^{10b}$, and R^{10c} are deuterium or hydrogen. In other embodiments, R^4 is deuterium, and $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}, R^5, R^6, R^7, R^8, R^9a, R^9b, R^{10a}, R^{10b}$, and R^{10c} are hydrogen.

[0068] In another embodiment, R^5 is deuterium, and $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}, R^4, R^6, R^7, R^8, R^9a, R^9b, R^{10a}, R^{10b}$, and R^{10c} are deuterium or hydrogen. In yet another embodiment, R^5 is deuterium, and $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}, R^4, R^6, R^7, R^8, R^9a, R^9b, R^{10a}, R^{10b}$, and R^{10c} are hydrogen.

[0069] In another embodiment, at least one of R^{9a} or R^{9b} are the same. In other embodiments, at least one of R^{9a} or R^{9b} are deuterium, and $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}, R^4, R^5, R^6, R^7, R^8, R^{10a}, R^{10b}$, and R^{10c} are deuterium or hydrogen. In some embodiments, at least one of R^{9a} and R^{9b} are deuterium, and $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}, R^4, R^5, R^6, R^7, R^8, R^{10a}, R^{10b}, R^{10c}$ are hydrogen.

[0070] In one embodiment, R^6, R^{9a} and R^{9b} are the same. In some embodiments, R^6, R^{9a} and R^{9b} are deuterium, and $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}, R^4, R^5, R^7, R^8, R^{10a}, R^{10b}, R^{10c}$ are deuterium or hydrogen. In other embodiments, R^6, R^{9a} and R^{9b} are deuterium, and $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}, R^4, R^5, R^7, R^8, R^{10a}, R^{10b}$, and R^{10c} are hydrogen.

[0071] In some embodiments, R^2, R^{10a}, R^{10b} , and R^{10c} are the same. In another embodiment, R^2, R^{10a}, R^{10b} , and R^{10c} are deuterium, and $R^{1a}, R^{1b}, R^{1c}, R^{3a}, R^{3b}, R^{3c}, R^4, R^5, R^6, R^7, R^8, R^9a$, and R^{9b} are deuterium or hydrogen. In yet another embodiment, R^2, R^{10a}, R^{10b} , and R^{10c} are deuterium, and $R^{1a}, R^{1b}, R^{1c}, R^{3a}, R^{3b}, R^{3c}, R^4, R^5, R^6, R^7, R^8, R^9a$, and R^{9b} are hydrogen.

[0072] In some embodiments, R^7 and at least two of R^{10a}, R^{10b} , or R^{10c} are the same. In another embodiment, R^7 and at least two of R^{10a}, R^{10b} , or R^{10c} are deuterium, and $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}, R^4, R^5, R^6, R^8, R^9a$, and R^{9b} are deuterium or hydrogen. In yet another embodiment, R^7 and at least two of R^{10a}, R^{10b} , or R^{10c} are deuterium, and $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}, R^4, R^5, R^6, R^8, R^9a$, and R^{9b} are hydrogen.

[0073] In some embodiments, $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}$, and at least one of R^{10a}, R^{10b} , or R^{10c} are the same. In another embodiment, $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}$, and at least one of R^{10a}, R^{10b} , or R^{10c} are deuterium, and $R^4, R^5, R^6, R^7, R^8, R^9a$, and R^{9b} are deuterium or hydrogen. In yet another embodiment, $R^{1a}, R^{1b}, R^{1c}, R^2, R^{3a}, R^{3b}, R^{3c}$, and at least one of R^{10a}, R^{10b} , or R^{10c} are deuterium, and $R^4, R^5, R^6, R^7, R^8, R^9a$, and R^{9b} are hydrogen.

[0074] In some embodiments, $R^{1a}, R^{1b}, R^{1c}, R^{3a}, R^{3b}, R^{3c}$, and R^5 are the same. In another embodiment, $R^{1a}, R^{1b}, R^{1c}, R^{3a}, R^{3b}, R^{3c}$, and R^5 are deuterium, and R^2, R^4, R^6, R^7, R^8 ,

R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are deuterium or hydrogen. In yet another embodiment, R^{1a} , R^{1b} , R^{1c} , R^{3a} , R^{3b} , R^{3c} , and R^5 are deuterium, and R^2 , R^4 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are hydrogen.


[0075] In other embodiments, R^4 and R^6 are the same. In another embodiment, R^4 and R^6 are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^5 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are deuterium or hydrogen. In yet another embodiment, R^4 and R^6 are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^5 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are hydrogen.

[0076] In one embodiment, R^2 , R^5 , R^{9a} , and R^{9b} are the same. In some embodiments, R^2 , R^5 , R^{9a} , and R^{9b} are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^{3a} , R^{3b} , R^{3c} , R^4 , R^6 , R^7 , R^8 , R^{10a} , R^{10b} , and R^{10c} are deuterium or hydrogen. In another embodiment, R^2 , R^5 , R^{9a} , and R^{9b} are deuterium, and R^{1a} , R^{1b} , R^{1c} , R^{3a} , R^{3b} , R^{3c} , R^4 , R^6 , R^7 , R^8 , R^{10a} , R^{10b} , and R^{10c} are hydrogen.


[0077] In yet another embodiment, R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^5 , R^6 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are the same. In some embodiments, R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^5 , R^6 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} are deuterium, and R^4 , R^7 , and R^8 are deuterium or hydrogen. In other embodiments, R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^5 , R^6 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and R^{10c} is deuterium, and R^4 , R^7 , and R^8 are hydrogen.

[0078] In some embodiments, the variables are as depicted in the compounds of the disclosure including compounds in the tables below.


Table I



I-1













I-2




I-3

Table II







II-10




II-11




II-12




II-13




II-14




II-15




II-16




II-17




II-18




II-19



II-20



II-21




II-22

[0079] Compounds of this invention include those described generally herein, and are further illustrated by the classes, subclasses, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed. Additionally, general principles of organic chemistry are described in “Organic Chemistry”, Thomas Sorrell, University Science Books, Sausalito: 1999, and “March’s Advanced Organic Chemistry”, 5th Ed., Ed.: Smith, M.B. and March, J., John Wiley & Sons, New York: 2001, the entire contents of which are hereby incorporated by reference.


[0080] As described herein, a specified number range of atoms includes any integer therein. For example, a group having from 1-4 atoms could have 1, 2, 3, or 4 atoms.

[0081] As described herein, compounds of the invention may optionally be substituted with one or more substituents, such as are illustrated generally herein, or as exemplified by particular classes, subclasses, and species of the invention. It will be appreciated that the phrase “optionally substituted” is used interchangeably with the phrase “substituted or unsubstituted.” In general, the term “substituted”, whether preceded by the term “optionally” or not, refers to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. Unless otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. Combinations of substituents envisioned by this invention are preferably those that result in the formation of stable or chemically feasible compounds.

[0082] Unless otherwise indicated, a substituent connected by a bond drawn from the center of a ring means that the substituent can be bonded to any position in the ring. In example **i** below, for instance, J¹ can be bonded to any position on the pyridyl ring. For bicyclic rings, a bond drawn through both rings indicates that the substituent can be bonded from any position of the bicyclic ring. In example **ii** below, for instance, J¹ can be bonded to the 5-membered ring (on the nitrogen atom, for instance), and to the 6-membered ring.



i



ii

[0083] The term “stable”, as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, recovery, purification, and use for one or more of the purposes disclosed herein. In some embodiments, a stable compound or chemically feasible compound is one that is not substantially altered when kept at a temperature of 40°C or less, in the absence of moisture or other chemically reactive conditions, for at least a week.

[0084] The term “aliphatic” or “aliphatic group”, as used herein, means a straight-chain (i.e., unbranched), branched, or cyclic, substituted or unsubstituted hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation that has a single point of attachment to the rest of the molecule.

[0085] Unless otherwise specified, aliphatic groups contain 1-20 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-10 aliphatic carbon atoms. In other embodiments, aliphatic groups contain 1-8 aliphatic carbon atoms. In still other embodiments, aliphatic groups contain 1-6 aliphatic carbon atoms, and in yet other embodiments aliphatic groups contain 1-4 aliphatic carbon atoms. Aliphatic groups may be linear or branched, substituted or unsubstituted alkyl, alkenyl, or alkynyl groups. Specific examples include, but are not limited to, methyl, ethyl, isopropyl, n-propyl, sec-butyl, vinyl, n-butenyl, ethynyl, and tert-butyl. Aliphatic groups may also be cyclic, or have a combination of linear or branched and cyclic groups. Examples of such types of aliphatic groups include, but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexenyl, -CH₂-cyclopropyl, CH₂CH₂CH(CH₃)-cyclohexyl.

[0086] The term “cycloaliphatic” (or “carbocycle” or “carbocyclyl”) refers to a monocyclic C₃-C₈ hydrocarbon or bicyclic C₈-C₁₂ hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic, that has a single point of attachment to the rest of the molecule wherein any individual ring in said bicyclic ring system has 3-7 members. Examples of cycloaliphatic groups include, but are not limited to, cycloalkyl and cycloalkenyl groups. Specific examples include, but are not limited to, cyclohexyl, cyclopropenyl, and cyclobutyl.

[0087] The term “heterocycle”, “heterocyclyl”, or “heterocyclic” as used herein means non-aromatic, monocyclic, bicyclic, or tricyclic ring systems in which one or more ring members are an independently selected heteroatom. In some embodiments, the “heterocycle”, “heterocyclyl”, or “heterocyclic” group has three to fourteen ring members in which one or more ring members is a heteroatom independently selected from oxygen, sulfur, nitrogen, or phosphorus, and each ring in the system contains 3 to 7 ring members.

[0088] Examples of heterocycles include, but are not limited to, 3-1H-benzimidazol-2-one, 3-(1-alkyl)-benzimidazol-2-one, 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydrothiophenyl, 3-tetrahydrothiophenyl, 2-morpholino, 3-morpholino, 4-morpholino, 2-thiomorpholino, 3-thiomorpholino, 4-thiomorpholino, 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 1-tetrahydropiperazinyl, 2-tetrahydropiperazinyl, 3-tetrahydropiperazinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 1-pyrazolinyl, 3-pyrazolinyl, 4-pyrazolinyl, 5-pyrazolinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 2-thiazolidinyl, 3-thiazolidinyl, 4-thiazolidinyl, 1-imidazolidinyl, 2-imidazolidinyl, 4-imidazolidinyl, 5-imidazolidinyl, indolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, benzothiolane, benzodithiane, and 1,3-dihydro-imidazol-2-one.

[0089] Cyclic groups, (e.g. cycloaliphatic and heterocycles), can be linearly fused, bridged, or spirocyclic.

[0090] The term “heteroatom” means one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2*H*-pyrrolyl), NH (as in pyrrolidinyl) or NR⁺ (as in N-substituted pyrrolidinyl)).

[0091] The term “unsaturated”, as used herein, means that a moiety has one or more units of unsaturation. As would be known by one of skill in the art, unsaturated groups can be partially unsaturated or fully unsaturated. Examples of partially unsaturated groups include, but are not limited to, butene, cyclohexene, and tetrahydropyridine. Fully unsaturated groups can be aromatic, anti-aromatic, or non-aromatic. Examples of fully unsaturated groups include, but are not limited to, phenyl, cyclooctatetraene, pyridyl, thienyl, and 1-methylpyridin-2(1*H*)-one.

[0092] The term “alkoxy”, or “thioalkyl”, as used herein, refers to an alkyl group, as previously defined, attached through an oxygen (“alkoxy”) or sulfur (“thioalkyl”) atom.

[0093] The terms “haloalkyl”, “haloalkenyl”, “haloaliphatic”, and “haloalkoxy” mean alkyl, alkenyl or alkoxy, as the case may be, substituted with one or more halogen atoms. This term includes perfluorinated alkyl groups, such as -CF₃ and -CF₂CF₃.


[0094] The terms “halogen”, “halo”, and “hal” mean F, Cl, Br, or I.

[0095] The term “aryl” used alone or as part of a larger moiety as in “aralkyl”, “aralkoxy”, or “aryloxyalkyl”, refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is

aromatic and wherein each ring in the system contains 3 to 7 ring members. The term “aryl” may be used interchangeably with the term “aryl ring”.

[0096] The term “heteroaryl”, used alone or as part of a larger moiety as in “heteroaralkyl” or “heteroarylalkoxy”, refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic, at least one ring in the system contains one or more heteroatoms, and wherein each ring in the system contains 3 to 7 ring members. The term “heteroaryl” may be used interchangeably with the term “heteroaryl ring” or the term “heteroaromatic”. Examples of heteroaryl rings include, but are not limited to, 2-furanyl, 3-furanyl, N-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, benzimidazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, N-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, pyridazinyl (e.g., 3-pyridazinyl), 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, tetrazolyl (e.g., 5-tetrazolyl), triazolyl (e.g., 2-triazolyl and 5-triazolyl), 2-thienyl, 3-thienyl, benzofuryl, benzothiophenyl, indolyl (e.g., 2-indolyl), pyrazolyl (e.g., 2-pyrazolyl), isothiazolyl, 1,2,3-oxadiazolyl, 1,2,5-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,3-triazolyl, 1,2,3-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, purinyl, pyrazinyl, 1,3,5-triazinyl, quinolinyl (e.g., 2-quinolinyl, 3-quinolinyl, 4-quinolinyl), and isoquinolinyl (e.g., 1-isoquinolinyl, 3-isoquinolinyl, or 4-isoquinolinyl).

[0097] It shall be understood that the term “heteroaryl” includes certain types of heteroaryl rings that exist in equilibrium between two different forms. More specifically, for example, species such as hydroxypyridine and pyridinone (and likewise hydroxypyrimidine and pyrimidinone) are meant to be encompassed within the definition of “heteroaryl.”



[0098] The term “protecting group” and “protective group” as used herein, are interchangeable and refer to an agent used to temporarily block one or more desired functional groups in a compound with multiple reactive sites. In certain embodiments, a protecting group has one or more, or preferably all, of the following characteristics: a) is added selectively to a functional group in good yield to give a protected substrate that is b) stable to reactions occurring at one or more of the other reactive sites; and c) is selectively removable in good yield by reagents that do not attack the regenerated, deprotected functional group. As would be understood by one skilled in the art, in some cases, the reagents do not

attack other reactive groups in the compound. In other cases, the reagents may also react with other reactive groups in the compound. Examples of protecting groups are detailed in Greene, T.W., Wuts, P. G in “Protective Groups in Organic Synthesis”, Third Edition, John Wiley & Sons, New York: 1999 (and other editions of the book), the entire contents of which are hereby incorporated by reference. The term “nitrogen protecting group”, as used herein, refers to an agent used to temporarily block one or more desired nitrogen reactive sites in a multifunctional compound. Preferred nitrogen protecting groups also possess the characteristics exemplified for a protecting group above, and certain exemplary nitrogen protecting groups are also detailed in Chapter 7 in Greene, T.W., Wuts, P. G in “Protective Groups in Organic Synthesis”, Third Edition, John Wiley & Sons, New York: 1999, the entire contents of which are hereby incorporated by reference.

[0099] In some embodiments, a methylene unit of an alkyl or aliphatic chain is optionally replaced with another atom or group. Examples of such atoms or groups include, but are not limited to, nitrogen, oxygen, sulfur, $-C(O)-$, $-C(=N-CN)-$, $-C(=NR)-$, $-C(=NOR)-$, $-SO-$, and $-SO_2-$. These atoms or groups can be combined to form larger groups. Examples of such larger groups include, but are not limited to, $-OC(O)-$, $-C(O)CO-$, $-CO_2-$, $-C(O)NR-$, $-C(=N-CN)$, $-NRCO-$, $-NRC(O)O-$, $-SO_2NR-$, $-NRSO_2-$, $-NRC(O)NR-$, $-OC(O)NR-$, and $-NRSO_2NR-$, wherein R is, for example, H or C_{1-6} aliphatic. It should be understood that these groups can be bonded to the methylene units of the aliphatic chain via single, double, or triple bonds. An example of an optional replacement (nitrogen atom in this case) that is bonded to the aliphatic chain via a double bond would be $-CH_2CH=N-CH_3$. In some cases, especially on the terminal end, an optional replacement can be bonded to the aliphatic group via a triple bond. One example of this would be $CH_2CH_2CH_2C\equiv N$. It should be understood that in this situation, the terminal nitrogen is not bonded to another atom.


[00100] It should also be understood that, the term “methylene unit” can also refer to branched or substituted methylene units. For example, in an isopropyl moiety $[-CH(CH_3)_2]$, a nitrogen atom (e.g. NR) replacing the first recited “methylene unit” would result in dimethylamine $[-N(CH_3)_2]$. In instances such as these, one of skill in the art would understand that the nitrogen atom will not have any additional atoms bonded to it, and the “R” from “NR” would be absent in this case.


[00101] Unless otherwise indicated, the optional replacements form a chemically stable compound. Optional replacements can occur both within the chain and/or at either end of the chain; i.e. both at the point of attachment and/or also at the terminal end. Two optional replacements can also be adjacent to each other within a chain so long as it results in a

chemically stable compound. For example, a C₃ aliphatic can be optionally replaced by 2 nitrogen atoms to form -C-N≡N. The optional replacements can also completely replace all of the carbon atoms in a chain. For example, a C₃ aliphatic can be optionally replaced by -NR-, -C(O)-, and -NR- to form -NRC(O)NR- (a urea).

[00102] Unless otherwise indicated, if the replacement occurs at the terminal end, the replacement atom is bound to a hydrogen atom on the terminal end. For example, if a methylene unit of -CH₂CH₂CH₃ were optionally replaced with -O-, the resulting compound could be -OCH₂CH₃, -CH₂OCH₃, or -CH₂CH₂OH. It should be understood that if the terminal atom does not contain any free valence electrons, then a hydrogen atom is not required at the terminal end (e.g., -CH₂CH₂CH=O or -CH₂CH₂C≡N).

[00103] Unless otherwise indicated, structures depicted herein are also meant to include all isomeric (e.g., enantiomeric, diastereomeric, geometric, conformational, and rotational) forms of the structure. For example, the R and S configurations for each asymmetric center, (Z) and (E) double bond isomers, and (Z) and (E) conformational isomers are included in this invention. As would be understood to one skilled in the art, a substituent can freely rotate

around any rotatable bonds. For example, a substituent drawn as  also

represents .

[00104] Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, geometric, conformational, and rotational mixtures of the present compounds are within the scope of the invention.

[00105] Unless otherwise indicated, all tautomeric forms of the compounds of the invention are within the scope of the invention.

[00106] In the compounds of this invention any atom not specifically designated as a particular isotope is meant to represent any stable isotope of that atom. Unless otherwise stated, when a position is designated specifically as "H" or "hydrogen", the position is understood to have hydrogen at its natural abundance isotopic composition. Also unless otherwise stated, when a position is designated specifically as "D" or "deuterium", the position is understood to have deuterium at an abundance that is at least 3340 times greater than the natural abundance of deuterium, which is 0.015% (i.e., at least 50.1% incorporation of deuterium).


[00107] "D" and "d" both refer to deuterium.

[00108] Additionally, unless otherwise indicated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a ^{13}C - or ^{14}C -enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools or probes in biological assays.

Processes

[00109] Processes and compounds described herein are useful for producing ATR inhibitors that contain an aminopyrazine-isoxazole core. The general synthetic procedures shown in schemes herein are useful for generating a wide array of chemical species which can be used in the manufacture of pharmaceutical compounds.

SCHEME A



Step 1

[00110] The compound of formula I can be made according to the steps outlined in Scheme A. Step 1 depicts the use of a readily available aldehyde/ketal as a starting point for the preparation of compounds of formula I, I-A, and I-B. Reductive amination between compound **1** and a suitable primary amine, under conditions known to those skilled in the art leads to compound **2** where a benzylamine motif has been installed. For example, imines can be formed by combining an amine and an aldehyde in a suitable solvent, such as

dichloromethane (DCM), dichloroethane (DCE), an alcoholic solvent (e.g., methanol, ethanol), or a nonprotic solvent (e.g., dioxane or tetrahydrofuran (THF)). These imines can then be reduced by known reducing agents including, but not limited to, NaBH₄, NaBH₃CN, and NaBH(OAc)₃ (see JOC 1996, 3849). In some embodiments, 1.05 equivalents of amine is combined with 1 equivalent of aldehyde in methanol. In other embodiments, 1.2 equivalents of amine is combined with 1 equivalent of aldehyde in methanol. This step is then followed by reduction with 0.6 to 1.4 (such as 1.2) equivalents of NaBH₄. In some cases, if an amine salt is used, base (e.g., Et₃N or diisopropylethylamine) can also be added.

Step 2

[00111] Step 2 depicts the protection of the benzylamine 1 prepared above, using a carbamate-based protecting group, under suitable protection conditions known to those skilled in the art. Various protecting groups, such as Cbz and Boc, can be used. Protection conditions include, but are not limited to the following:

- a) R-OCOCl, a suitable tertiary amine base, and a suitable solvent; wherein R is C₁₋₆alkyl optionally substituted with phenyl;
- b) R(CO₂)OR', a suitable solvent, and optionally a catalytic amount of base, wherein R is and R' are each independently C₁₋₆alkyl optionally substituted with phenyl;
- c) [RO(C=O)]₂O, a suitable base, and a suitable solvent.

[00112] Examples of suitable bases include, but are not limited to, Et₃N, diisopropylamine, and pyridine. Examples of suitable solvents include chlorinated solvents (e.g., dichloromethane (DCM), dichloroethane (DCE), CH₂Cl₂, and chloroform), ethers (e.g., THF, 2-MeTHF, and dioxane), aromatic hydrocarbons (e.g., toluene, xylenes) and other aprotic solvents.

[00113] In some embodiments, protection can be done by reacting the benzylamine with (Boc)₂O and Et₃N in DCM. In some embodiments, 1.02 equivalents of (Boc)₂O and 1.02 equivalents of Et₃N 1.02 are used. . In another embodiment, protection can be done by reacting the benzylamine with (Boc)₂O in 2-MeTHF. In some embodiments, 1.05 equivalents of (Boc)₂O are used.

Step 3

[00114] Step 3 shows how the ketal functional group in 3 is then converted into the oxime 4 in a single step. This direct conversion from ketal to oxime is not extensively described in the literature and it will be appreciated that this step could also be conducted in a

two-step sequence, transiting through the aldehyde after deprotection of the ketal using methodologies known to those skilled in the art.

[00115] Oxime formation conditions comprise mixing together hydroxylamine, acid, optionally a dehydrating agent, and an alcoholic solvent. In some embodiments, the acid is a catalytic amount. In some embodiments, the acid is pTSA or HCl, the dehydrating agent is molecular sieves or dimethoxyacetone, and the alcoholic solvent is methanol or ethanol. In some embodiments, the hydroxylamine hydrochloride is used in which case no additional acid is required. In other embodiments, the desired product is isolated via a biphasic work up and optionally precipitation or crystallization. If a biphasic work up is used, a dehydrating agent is not needed.

[00116] In another embodiment, the oxime formation conditions comprise of mixing together hydroxylamine, an acid, an organic solvent and water. Examples of suitable organic solvents include chlorinated solvents (e.g., dichloromethane (DCM), dichloroethane (DCE), CH_2Cl_2 , and chloroform), ethers (e.g., THF, 2-MeTHF and dioxane), aromatic hydrocarbons (e.g., toluene, xylenes) and other aprotic solvents. In some embodiments, 1.5 equivalents of hydroxylamine hydrochloride are used, the organic solvent is 2-MeTHF and the water is buffered with Na_2SO_4 . In another embodiment, 1.2 equivalents of hydroxylamine hydrochloride are used, the organic solvent is THF.

[00117] In some embodiments, suitable deprotection conditions comprise adding catalytic amounts of para-toluenesulfonic acid (pTSA), acetone, and water; and then forming the oxime using conditions known to one skilled in the art. In other embodiments, a single step sequence is used. In some embodiments, the single step sequence comprises adding $\text{NH}_2\text{OH} \cdot \text{HCl}$ and a mixture of THF and water. In some embodiments, 1 equivalent of the compound of formula **3** is combined with a 1.1 equivalents of $\text{NH}_2\text{OH} \cdot \text{HCl}$ in a 10:1 v/v mixture of THF/water.

Step 4

Step 4 illustrates how the oxime **4** is then transformed and engaged in a [3+2] cycloaddition to form the isoxazole **5**. This transformation can be conducted in one pot but requires two distinct steps. The first step is an oxidation of the oxime functional group into a nitrone, or a similar intermediate with the same degree of oxidation, for example a chlorooxime. This reactive species then reacts with an alkyne in a [3+2] cycloaddition to form the isoxazole adduct.

[00118] In some embodiments, the suitable isoxazole-formation conditions consists of two steps, the first step comprising reacting the compound of formula 4 under suitable chlorooxime formation conditions to form a chlorooxime intermediate; the second step comprising reacting the chlorooxime intermediate with acetylene under suitable cycloaddition conditions to form a compound of formula 5.

[00119] In some embodiments, the chlorooxime formation conditions are selected from

- a) N-chlorosuccinimide and suitable solvent;
- b) potassium peroxydisulfate, HCl, and dioxane; and
- c) Sodium hypochlorite and a suitable solvent

[00120] Examples of suitable solvents include, but are not limited to, nonprotic solvents (e.g., DCM, DCE, THF, 2-MeTHF, MTBE and dioxane), aromatic hydrocarbons (e.g. toluene, xylenes), and alkyl acetates (e.g., isopropyl acetate, ethyl acetate).

[00121] Isolation of the product can be achieved by adding an antisolvent to a solution of a compound of formula 5. Examples of suitable solvents for isolating the chlorooxime intermediate include mixtures of suitable solvents (EtOAc, IPAC) with hydrocarbons (e.g., hexanes, heptane, cyclohexane), or aromatic hydrocarbons (e.g., toluene, xylenes). In some embodiments, heptane is added to a solution of chlorooxime in IPAC.

[00122] Suitable cycloaddition conditions consist of combining the chlorooxime with acetylene with a suitable base and a suitable solvent. Suitable solvents include protic solvents, aprotic solvents, polar solvents, and nonpolar solvents. Examples of suitable solvent include, but are not limited to, acetonitrile, tetrahydrofuran, 2-methyltetrahydrofuran, MTBE, EtOAc, i-PrOAc, DCM, toluene, DMF, and methanol. Suitable bases include, but are not limited to, pyridine, DIEA, TEA, t-BuONa, and K₂CO₃. In some embodiments, suitable cycloaddition conditions comprise adding 1.0 equivalents of chlorooxime, 1.0 equivalents of acetylene, 1.1 equivalents of Et₃N in DCM.

[00123] Isolation of the product can be achieved by adding an antisolvent to a solution of a compound of formula 5. Examples of suitable solvents for isolating the chlorooxime include mixtures of suitable solvents (EtOAc, IPAC) with hydrocarbons (e.g., hexanes, heptane, cyclohexane), or aromatic hydrocarbons (e.g., toluene, xylenes). In some embodiments, heptane is added to a solution of chlorooxime in IPAC.

Step 5

[00124] Step 5 depicts the final step(s) of the preparation of compounds of formula I. When the R₄ group is bromo, intermediate 5 can be subjected to a Suzuki cross-coupling

with boronic acid or esters, under conditions known to those skilled in the art, to form compounds where R₄ an aryl, heteroaryl or alternative moieties resulting from the metal-assisted coupling reaction. When intermediate 5 is suitably functionalised, a deprotection step can be carried out to remove the protecting groups and generate the compounds of formula I.

[00125] Metal assisted coupling reactions are known in the art (see e.g., *Org. Proc. Res. Dev.* 2010, 30-47). In some embodiments, suitable coupling conditions comprise adding 0.1 equivalents of Pd[P(tBu)₃]₂; 1 equivalent of boronic acid or ester; and 2 equivalents of sodium carbonate in a 2:1 ratio v/v of acetonitrile/water at 60-70°C. In other embodiments, suitable coupling conditions comprise adding 0.010-0.005 equivalents Pd(dtbpf)Cl₂, 1 equivalent of boronic acid or ester, and 2 equivalents of potassium carbonate in a 7:2 v/v of toluene and water at 70 °C

[00126] The final product can be treated with a metal scavenger (silica gel, functionalized resins, charcoal) (see e.g., *Org. Proc. Res. Dev.* 2005, 198-205). In some embodiments, the solution of the product is treated with Biotage MP-TMT resin.

[00127] The product can also be isolated by crystallization from an alcoholic solvent (e.g. methanol, ethanol, isopropanol). In some embodiments the solvent is ethanol. In other embodiments the solvent is isopropanol.

[00128] Deprotection of Boc groups is known in the art (see e.g. *Protecting Groups in Organic Synthesis*, Greene and Wuts). In some embodiments, suitable deprotection conditions are hydrochloric acid in acetone at 35-45 °C. In other embodiments, suitable deprotection conditions are TFA in DCM.

Step 6


[00129] Step 6 illustrates how compounds of formula I are converted to compounds of formula I-A using a base under suitable conditions known to those skilled in the art. In some embodiments, isolation of the free-base form of compounds of formula I may be achieved by adding suitable base, such as NaOH to an alcoholic acidic solution of compounds of formula I to precipitate the product.

Step 7

[00130] Step 7 illustrates how compounds of formula I-A are converted to compounds of formula I-B using an acid under suitable conditions known to those skilled in the art.


[00131] In some embodiments suitable conditions involve adding aqueous HCl, to a suspension of compounds of formula I-A in acetone at 35 °C then heating at 50 °C.

SCHEME B: Formation of d1-boronate




[00132] Scheme B shows a general synthetic method for the preparation of d1-boronate intermediates. A suitable 1-halo-(isopropylsulfonyl)benzene is treated with a base such as, but not limited to NaH, LiHMDS or KHMDS followed by quenching of the anion with deuterium source such as D₂O. The halogen is then transformed into a suitable boronate derivative *via*, for example, metal mediated cross-coupling catalyzed by, for instance, Pd(^tBu₃)₂ or Pd(dppf)Cl₂·DCM.

SCHEME C: Formation of d6-boronate




[00133] Scheme C shows a general synthetic method for the preparation of d6-boronate intermediates. A suitable 1-halo-(methylsulfonyl)benzene is treated with a base such as, but not limited to NaH, LiHMDS or KHMDS followed by quenching of the anion with deuterium source such as D₃Cl. This reaction is repeated until the desired amount of deuterium has been incorporated into the molecule. The halogen is then transformed into a suitable boronate derivative *via*, for example, metal mediated cross-coupling catalyzed by, for instance, Pd(^tBu₃)₂ or Pd(dppf)Cl₂·DCM.

SCHEME D: Formation of d7-boronate



[00134] Scheme D shows a general synthetic method for the preparation of d7-boronate intermediates. 4-Bromobenzenethiol is treated with a base such as, but not limited to NaH, LiHMDS or KHMDS followed by quenching of the anion with deuterium source such as 1,1,1,2,3,3,3-heptadeuterio-2-iodo-propane. The sulfide is then oxidized to the corresponding sulfone using, for example, mCPBA or Oxone. The halogen is then transformed into a suitable boronate derivative *via*, for example, metal mediated cross-coupling catalyzed by, for instance, $\text{Pd}(\text{tBu}_3)_2$ or $\text{Pd}(\text{dppf})\text{Cl}_2\cdot\text{DCM}$.


SCHEME E: Formation of aryl ring deuterated boronate



[00135] Scheme E shows a general synthetic method for the preparation of boronate intermediates where the aryl ring is substituted with a deuterium. A suitable 1-iodo-4-bromo-aryl derivative is treated with a substituted thiol such as propane-2-thiol under metal catalyzed coupling conditions using a catalyst such as CuI. The sulfide is then oxidized to the corresponding sulfone using, for example, mCPBA or Oxone. The bromide is then transformed into a suitable boronate derivative *via*, for example, metal mediated cross-coupling catalyzed by, for instance, $\text{Pd}(\text{tBu}_3)_2$ or $\text{Pd}(\text{dppf})\text{Cl}_2\cdot\text{DCM}$. The remaining substituent is then converted into deuterium by, for instance, metal catalyzed halogen-deuterium exchange using a suitable metal catalyst, such as Pd on C under an atmosphere of deuterium gas. In addition, the 1-bromo-(isopropylsulfonyl)benzene can be treated with a


base such as, but not limited to NaH, LiHMDS or KHMDS followed by quenching of the anion with deuterium source such as D₂O. The bromide is then transformed into a suitable boronate derivative *via*, for example, metal mediated cross-coupling catalyzed by, for instance, Pd(^tBu₃)₂ or Pd(dppf)Cl₂·DCM. The remaining substituent is then converted into deuterium by, for instance, metal catalyzed halogen-deuterium exchange using a suitable metal catalyst, such as Pd on C under an atmosphere of deuterium gas.

SCHEME F: Formation of aryl ring deuterated boronate




[00136] Scheme F shows another general synthetic method for the preparation of boronate intermediates where the aryl ring is substituted with a deuterium. A substituted 4-bromobenzenethiol is treated with a base such as, but not limited to NaH, LiHMDS or KHMDS followed by quenching of the anion with deuterium source such as 1,1,1,2,3,3,3-heptadeuterio-2-iodopropane. The sulfide is then oxidized to the corresponding sulfone using, for example, mCPBA or Oxone. The halogen is then transformed into a suitable boronate derivative *via*, for example, metal mediated cross-coupling catalyzed by, for instance, Pd(^tBu₃)₂ or Pd(dppf)Cl₂·DCM. The remaining substituent is then converted into deuterium by, for instance, metal catalyzed halogen-deuterium exchange using a suitable metal catalyst, such as Pd on C under an atmosphere of deuterium gas.

SCHEME G: Formation of aryl ring deuterated boronate



[00137] Scheme G shows another general synthetic method for the preparation of boronate intermediates where the aryl ring is substituted with a deuterium. A substituted 4-bromobenzenethiol is treated with a base such as, but not limited to NaH, LiHMDS or KHMDS followed by quenching of the anion with for instance MeI. The sulfide is then oxidized to the corresponding sulfone using, for example, mCPBA or Oxone. The sulfone is treated with a base such as, but not limited to NaH, LiHMDS or KHMDS followed by quenching of the anion with deuterium source such as D₃Cl. This reaction is repeated until the desired amount of deuterium has been incorporated into the molecule. The halogen is then transformed into a suitable boronate derivative *via*, for example, metal mediated cross-coupling catalyzed by, for instance, Pd(^tBu₃)₂ or Pd(dppf)Cl₂·DCM. The remaining substituent is then converted into deuterium by, for instance, metal catalyzed halogen-deuterium exchange using a suitable metal catalyst, such as Pd on C under an atmosphere of deuterium gas.


SCHEME H: Formation of aryl ring deuterated oxime intermediates



[00138] Scheme H shows a general synthetic method for the preparation of oxime intermediates where the aryl ring is substituted with a deuterium. The methyl group of an appropriately substituted methyl 4-methylbenzoate derivative can be converted into the corresponding dibromide under conditions such as AIBN catalyzed bromination with NBS. This di-bromide is then hydrolysed to the corresponding aldehyde, for instance using AgNO₃ in acetone/water. Protection of the aldehyde as a suitable acetal, for instance the diethyl acetal and subsequent conversion of the remaining substituent into deuterium by, for instance, metal catalyzed halogen-deuterium exchange using a suitable metal catalyst, such as Pd on C under an atmosphere of deuterium gas gives the deuterated ester intermediate. The ester


functionality can be reduced using reagents such as LiAlH_4 , NaBH_4 , NaBD_4 or LiAlD_4 to give corresponding aldehyde. This can be reacted under reductive amination conditions using a suitable amine, such as methylamine or d_3 -methylamine using a reducing agent such as NaBH_4 or NaBD_4 to give the corresponding amine derivative. This can be protected with, for instance a Boc group and the acetal converted into the oxime using, for instance, hydroxylamine hydrochloride in THF/water.

SCHEME I: Formation of aryl ring deuterated oxime intermediates




[00139] Scheme I shows another general synthetic method for the preparation of oxime intermediates where the aryl ring is substituted with a deuterium. The methyl group of an appropriately substituted methyl 4-methylbenzoate derivative can be converted into the corresponding dibromide under conditions such as AIBN catalyzed bromination with NBS. This di-bromide is then hydrolysed to the corresponding aldehyde, for instance using AgNO_3 in acetone/water. Protection of the aldehyde as a suitable acetal, for instance the dimethyl acetal and subsequent conversion of the remaining substituent into deuterium by, for instance, metal catalyzed halogen-deuterium exchange using a suitable metal catalyst, such as Pd on C under an atmosphere of deuterium gas gives the deuterated ester intermediate. The ester functionality can be converted into the corresponding primary amide under standard conditions, such as heating with a solution of ammonia in methanol. The amide can be reduced to the corresponding amine using reagents not limited to LiAlH_4 or LiAlD_4 . This can be protected with, for instance a Boc group. The carbamate NH can be alkylated under basic conditions using for instance NaH, LiHMDS or KHMDS followed by quenching of the anion with deuterium source such MeI or D_3Cl . The acetal can be converted into the oxime using, for instance, hydroxylamine hydrochloride in THF/water.

SCHEME J: Formation of aryl ring deuterated oxime intermediates




[00140] Scheme J shows another general synthetic method for the preparation of oxime intermediates where the aryl ring is substituted with a deuterium. The methyl group of an appropriately substituted methyl 4-methylbenzoate derivative can be converted into the corresponding dibromide under conditions such as AIBN catalyzed bromination with NBS. This di-bromide is then hydrolysed to the corresponding aldehyde, for instance using AgNO_3 in acetone/water. Protection of the aldehyde as a suitable acetal, for instance the dimethyl acetal and subsequent conversion of the remaining substituent into deuterium by, for instance, metal catalyzed halogen-deuterium exchange using a suitable metal catalyst, such as Pd on C under an atmosphere of deuterium gas gives the deuterated ester intermediate. The ester functionality can be converted into the corresponding primary amide under standard conditions, such as heating with a solution of ammonia in methanol. The amide can be reduced to the corresponding amine using reagents not limited to LiAlH_4 or LiAlD_4 . This can be reacted under reductive amination conditions using a suitable amine, such as methylamine, d_3 -methylamine, formaldehyde or d_2 -formaldehyde using a reducing agent such as NaBH_4 or NaBD_4 to give the corresponding amine derivative. This can be protected with, for instance a Boc group. The acetal can be converted into the oxime using, for instance, hydroxylamine hydrochloride in THF/water.

SCHEME K: Formation of aryl ring deuterated oxime intermediates




[00141] Scheme K shows another general synthetic method for the preparation of oxime intermediates where the aryl ring is substituted with a deuterium. The methyl group of an appropriately substituted methyl 4-methylbenzoate derivative can be converted into the corresponding dibromide under conditions such as AIBN catalyzed bromination with NBS. This di-bromide is then hydrolysed to the corresponding aldehyde, for instance using AgNO_3 in acetone/water. Protection of the aldehyde as a suitable acetal, for instance the dimethyl acetal and subsequent conversion of the remaining substituent into deuterium by, for instance, metal catalyzed halogen-deuterium exchange using a suitable metal catalyst, such as Pd on C under an atmosphere of deuterium gas gives the deuterated ester intermediate. This can be reacted under reductive amination conditions using a suitable amine, such as ammonium hydroxide using a reducing agent such as NaBH_4 or NaBD_4 to give the corresponding amine derivative. This can be protected with, for instance a Boc group and the carbamate NH alkylated under basic conditions using for instance NaH , LiHMDS or KHMDS followed by quenching of the anion with deuterium source such as MeI or D_3Cl . The ester can be reduced to the corresponding alcohol using a suitable reducing agent such as LiBH_4 or NaBH_4 . The alcohol can be oxidized to the aldehyde using reagents such as MnO_2 or Dess-Martin periodane. The acetal can be converted into the oxime using, for instance, aqueous hydroxylamine.

SCHEME L: Formation of deuterated oxime intermediates




[00142] Scheme L shows a general synthetic method for the preparation of deuterated oxime intermediates. 4-(diethoxymethyl)benzaldehyde can be reacted under reductive amination conditions using a suitable amine, such as methylamine or d₃-methylamine using a reducing agent such as NaBH₄ or NaBD₄ to give the corresponding amine derivative. This can be protected with, for instance a Boc group and the acetal converted into the oxime using, for instance, hydroxylamine hydrochloride in THF/water.

SCHEME M: Formation of deuterated oxime intermediates




[00143] Scheme M shows another general synthetic method for the preparation of deuterated oxime intermediates. The ester functionality of methyl 4-(dimethoxymethyl)benzoate can be converted into the corresponding primary amide under standard conditions, such as heating with a solution of ammonia in methanol. The amide can be reduced to the corresponding amine using reagents not limited to LiAlH₄ or LiAlD₄. This can be protected with, for instance a Boc group. The carbamate NH can be alkylated under basic conditions using for instance NaH, LiHMDS or KHMDS followed by quenching of the anion with deuterium source such as MeI or D₃Cl. The acetal can be converted into the oxime using, for instance, hydroxylamine hydrochloride in THF/water.

SCHEME N: Formation of deuterated oxime intermediates




[00144] Scheme N shows another general synthetic method for the preparation of deuterated oxime intermediates. The ester functionality of methyl 4-(dimethoxymethyl)benzoate can be converted into the corresponding primary amide under standard conditions, such as heating with a solution of ammonia in methanol. The amide can be reduced to the corresponding amine using reagents not limited to LiAlH_4 or LiAlD_4 . This can be reacted under reductive amination conditions using a suitable amine, such as methylamine, d_3 -methylamine, formaldehyde or d_2 -formaldehyde using a reducing agent such as NaBH_4 or NaBD_4 to give the corresponding amine derivative. This can be protected with, for instance a Boc group. The acetal can be converted into the oxime using, for instance, hydroxylamine hydrochloride in THF/water.

SCHEME O: Formation of deuterated oxime intermediates




[00145] Scheme O shows another general synthetic method for the preparation of deuterated oxime intermediates. A 4-substituted benzylamine can be protected with, for instance a Boc group. The carbamate NH can be alkylated under basic conditions using for instance NaH , LiHMDS or KHMDS followed by quenching of the anion with deuterium source such as MeI or D_3Cl . The ester can be reduced to the corresponding alcohol using a suitable reducing agent such as LiBH_4 or NaBH_4 . The alcohol can be oxidized to the aldehyde using reagents such as MnO_2 or Dess-Martin periodane. The acetal can be converted into the oxime using, for instance, aqueous hydroxylamine.

SCHEME P: Formation of isoxazole derrivatives



[00146] Scheme P shows a general synthetic method for the preparation of deuterated pyrazine-isoxazole derivatives. 3,5-Dibromopyrazin-2-amine is converted into the corresponding silyl-protected alkyne under standard Sonogashira conditions utilizing, for example, Pd(PPh₃)₄ and CuI as catalysts. The pyrazine NH₂ can then be protected as, for example the di-Boc derivative. Coupling of the pyrazine bromide with a boronate, for instance those outlined in Schemes 1 to 6 above, under standard Suzuki cross-coupling conditions followed by removal of the silyl protecting group give the desired alkyne intermediate. Oximes, such as those outline in Schemes 7 to 14 above, can be converted into the corresponding chlorooximes using, for instance, NCS. The alkyne and chlorooxime intermediates can undergo a [3+2] cycloaddition to give corresponding isoxazole under standard conditions, for instance by the addition of Et₃N. The Boc protecting groups can be removed under acid conditions such as TFA in DCM or HCl in MeOH/DCM to give the deuterated pyrazine isoxazole derivatives.

SCHEME Q: Formation of deuterated isoxazole derivatives



[00147] Scheme Q shows a general synthetic method for the preparation of deuterated isoxazole derivatives. The pyrazine NH_2 and benzylamineamine NH can be protected under standard conditions using trifluoroacetic anhydride. Halogenation of the isoxazole ring with, for example NIS followed by removal of the trifluoroacetate protecting group under basic conditions provides the desired halogenated intermediates. The halogen can then be converted into deuterium by, for instance, metal catalyzed halogen-deuterium exchange using a suitable metal catalyst, such as Pd on C under an atmosphere of deuterium gas.

Abbreviations

[00148] The following abbreviations are used:


ATP	adenosine triphosphate
Boc	tert-butyl carbamate
Cbz	Carboxybenzyl
DCM	dichloromethane

DMSO	dimethyl sulfoxide
Et ₃ N	triethylamine
2-MeTHF	2-methyltetrahydrofuran
NMM	N-Methyl morpholine
DMAP	4-Dimethylaminopyridine
TMS	Trimethylsilyl
MTBE	methyl tertbutyl ether
EtOAc	ethyl acetate
i-PrOAc	isopropyl acetate
IPAC	isopropyl acetate
DMF	dimethylformamide
DIEA	diisopropylethylamine
TEA	triethylamine
t-BuONa	sodium tertbutoxide
K ₂ CO ₃	potassium carbonate
PG	Protecting group
pTSA	para-toluenesulfonic acid
TBAF	Tetra-n-butylammonium fluoride
¹ HNMR	proton nuclear magnetic resonance
HPLC	high performance liquid chromatography
LCMS	liquid chromatography-mass spectrometry
TLC	thin layer chromatography
Rt	retention time

SCHEMES AND EXAMPLES

[00149] The compounds of the disclosure may be prepared in light of the specification using steps generally known to those of ordinary skill in the art. Those compounds may be analyzed by known methods, including but not limited to LCMS (liquid chromatography mass spectrometry) and NMR (nuclear magnetic resonance). The following generic schemes and examples illustrate how to prepare the compounds of the present disclosure. The examples are for the purpose of illustration only and are not to be construed as limiting the scope of the invention in any way. ¹H-NMR spectra were recorded at 400 MHz using a Bruker DPX 400 instrument. Mass spec. samples were analyzed on a MicroMass Quattro Micro mass spectrometer operated in single MS mode with electrospray ionization.

Example 1: Synthesis of 2-(4-(5-amino-6-(3-(4-((tetrahydro-2H-pyran-4-ylamino)methyl)phenyl)isoxazol-5-yl)pyrazin-2-yl)pyridin-2-yl)-2-methylpropanenitrile (Compound I-1)



Method 1:

[00150] To a solution of tetrahydropyran-4-amine (100 g, 988.7 mmol) in MeOH (3.922 L) was added 4-(diethoxymethyl)benzaldehyde (196.1 g, 941.6 mmol) over 2 min at RT. The reaction mixture was stirred at RT for 80 min, until the aldimine formation was complete (as seen by NMR). NaBH₄ (44.49 g, 1.176 mol) was carefully added over 45 min, maintaining the temperature between 24 °C and 27 °C by mean of an ice bath. After 75 min at RT, the reaction has gone to completion. The reaction mixture was quenched with 1M NaOH (1 L). The reaction mixture was partitioned between brine (2.5 L) and TBDME (4 L then 2 x 1 L). The organic phase was washed with brine (500 mL) and concentrated in vacuo. The crude mixture was redissolved in DCM (2 L). The aqueous phase was separated, the organic phase was dried over MgSO₄, filtered and concentrated in vacuo to give the title compound as a yellow oil (252.99 g, 91%).

Method 2:

[00151] A solution of N-[[4-(diethoxymethyl)phenyl]methyl] tetrahydropyran-4-amine (252.99 g, 862.3 mmol) and Boc anhydride (191.9 g, 202.0 mL, 879.5 mmol) in DCM (2.530 L) was cooled down to 3.3 °C. Et₃N (89.00 g, 122.6 mL, 879.5 mmol) was added over 4 min, keeping the internal temperature below 5 °C. The bath was removed 45 min after the end of the addition. And the reaction mixture was stirred at RT overnight. The reaction mixture was sequentially washed with 0.5 M citric acid (1 L), saturated NaHCO₃ solution (1 L) and brine (1 L). The organic phase was dried (MgSO₄), filtered and concentrated in vacuo to give a colourless oil (372.38 g, 110%). ¹H NMR (400.0 MHz, DMSO); MS (ES+)

Method 3:

[00152] tert-butyl N-[[4-(diethoxymethyl)phenyl]methyl]-N-tetrahydropyran-4-yl-carbamate (372.38 g, 946.3 mmol) was dissolved in THF (5 L) and water (500 mL). Hydroxylamine hydrochloride (72.34 g, 1.041 mol) was added in one portion and the reaction mixture was stirred overnight at RT. The reaction mixture was partitioned between DCM (5 L) and water. The combined organic extract was washed with water (1L x 2). The organic phase was concentrated in vacuo to a volume of about 2L. The organic layer was dried over MgSO₄, filtered and concentrated in vacuo to give a sticky colourless oil that crystallized on standing under vacuo. (334.42g, 106%). ¹H NMR (400.0 MHz, CDCl₃); MS (ES+)

Method 4:

[00153] tert-butyl N-[[4-[(E)-hydroxyiminomethyl]phenyl]methyl]-N-tetrahydropyran-4-yl-carbamate (334.13 g, 999.2 mmol) was dissolved in isopropyl acetate (3.0 L) (the mixture was warmed to 40 °C to allow all the solids to go into solution). N-chlorosuccinimide (140.1 g, 1.049 mol) was added portionwise over 5 min and the reaction mixture was heated to 55 °C (external block temperature). After 45 min at 55°C The reaction had gone to completion. The reaction mixture was cooled down to RT. The solids were filtered off and rinsed with Isopropyl acetate (1 L). Combined organic extract was sequentially washed with water (1.5 L, 5 times) and brine, dried over MgSO₄, filtered and concentrated in vacuo to give a viscous yellow oil (355.9 g; 96%). ¹H NMR (400.0 MHz, CDCl₃); MS (ES+)

Method 5:

[00154] Et₃N (76.97 g, 106.0 mL, 760.6 mmol) was added over 20 minutes to a solution of tert-butyl N-(5-bromo-3-ethynyl-pyrazin-2-yl)-N-tert-butoxycarbonyl-carbamate (233.0 g, 585.1 mmol) and tert-butyl N-[[4-[(Z)-C-chloro-N-hydroxy-carbonimidoyl]phenyl]methyl]-N-tetrahydropyran-4-yl-carbamate (269.8 g, 731.4 mmol) in DCM (2.330 L) at RT. During

addition of triethylamine, the exotherm was stabilised by cooling the mixture in an ice bath, then the reaction mixture was gradually warmed up to RT and the mixture was stirred at RT overnight. The reaction mixture was sequentially washed with water (1.5 L, 3 times) and brine. The organic extract was dried over MgSO₄, filtered and partially concentrated in vacuo. Heptane (1.5L) was added and the concentration was continued yielding 547.63 g of a yellow-orange solid.

[00155] 542.12 g was taken up into ~2 vol (1 L) of ethyl acetate. The mixture was heated to 74-75 °C internally and stirred until all the solid went into solution. Heptane (3.2 L) was added slowly via addition funnel to the hot solution keeping the internal temperature between 71 °C and 72 °C . At the end of the addition, the dark brown solution was seeded with some recrystallised product, and the reaction mixture was allowed to cool down to RT without any stirring to crystallise O/N. The solid was filtered off and rinsed with heptane (2 x 250 mL), then dried in vacuo to yield 307.38 g of the title product (72 %). 1H NMR (400.0 MHz, CDCl₃); MS (ES+)

Method 6:

[00156] tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]isoxazol-3-yl]phenyl]methyl]-N-tetrahydropyran-4-yl-carbamate (303 g, 414.7 mmol) and 2-methyl-2-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-pyridyl] propanenitrile (112.9 g, 414.7 mmol) were suspended in MeCN (2 L) and H₂O (1 L). Na₂CO₃ (414.7 mL of 2 M, 829.4 mmol) followed by Pd[P(tBu)₃]₂ (21.19 g, 41.47 mmol) were added and the reaction mixture was degassed with N₂ for 1 h. The reaction mixture was placed under a nitrogen atmosphere and heated at 70 °C (block temperature) for 4 h (internal temperature fluctuated between 60 °C and 61 °C). The reaction was cooled down to room temperature and stirred at RT overnight. The reaction mixture was partitioned between EtOAc (2 L) and water (500 mL). The combined organic extract was washed with brine (500 mL), filtered through a short pad of celite and concentrated under reduced pressure to a volume of about 3 L. The solution was dried over MgSO₄, filtered and partially concentrated in vacuo. iPrOH (1.5 L) was added and the solvent was removed in vacuo to yield the desired product as a light brown foam (405 g).


[00157] 400 g was taken up into ~5 vol (2 L) of iPrOH and the mixture was heated to 80 °C until all the solid went into solution. The dark brown solution was seeded, and the reaction mixture was allowed to slowly cool down to RT overnight. The solid was filtered off and rinsed with iPrOH (2 x 250 mL) and Petroleum ether (2x200 mL). The resulting solid was

slurried in petroleum ether (2.5 L), filtered off and dried in vacuo. The resulting solid was dissolved in DCM (2.5 L) and stirred slowly for 1 h with 30 g of SPM32 (3-mercaptopropyl ethyl sulfide silica). The silica was filtered through a pad of florisil and rinsed with DCM. The procedure was repeated twice, then the DCM solution was concentrated in vacuo to give 238.02 g of a light yellow solid.

Method 7:


[00158] tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-[2-(1-cyano-1-methyl-ethyl)-4-pyridyl]pyrazin-2-yl]isoxazol-3-yl]phenyl]methyl]-N-tetrahydropyran-4-yl-carbamate (238 g, 299.0 mmol) was dissolved in DCM (2.380 L). TFA (500 mL, 6.490 mol) was added at RT over 3 min. The reaction mixture was stirred at RT for 3.5 h. The reaction mixture was concentrated under reduced pressure then azeotroped with heptane (2x300ml). The oil was then slurried in abs. EtOH (2.5 L) and filtered. The solid was dissolved in a mixture of ethanol (1.190 L) and water (1.190 L). potassium carbonate (124.0 g, 897.0 mmol) in water (357.0 mL) was added to the solution and the mixture was stirred at RT overnight. **[00159]** The solid was filtered off, was washed with water (2.5 L), and dried at 50 °C in vacuo to give 108.82 g of the title compound (Compound I-1) as a yellow powder. (73 %)

Methods 6a and 7a



[00160] A mixture of tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-bromopyrazin-2-yl]isoxazol-3-yl]phenyl]methyl]-N-tetrahydropyran-4-yl-carbamate (110.0 g, 151 mmol), K₂CO₃ (41.6 g, 301 mmol), and 2-methyl-2-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-pyridyl]propanenitrile (41.0 g, 151 mmol) in toluene (770 mL) and water (220 mL) is stirred and degassed with N₂ for 30 min. at 20 °C. The catalyst Pd(dtbpf)Cl₂ (1.96 g, 3.01 mmol) is added and the mixture is degassed for an additional 10 min. The mixture is heated at 70 °C until the reaction is complete. The mixture is cooled to ambient temperature, diluted with water (220 mL), and filtered through a bed of Celite. The organic phase is concentrated to remove most of the solvent. The concentrate is diluted with *i*-PrOH (550 mL). The resultant suspension is stirred for at least 1 h and then the solid is collected by filtration to afford a tan powder. The solid is dissolved in toluene (990 mL) and stirred with Biotage MP-TMT resin (18.6 g) for 2 h at ambient temperature. The resin is removed by filtration. The filtrate is concentrated then diluted with *i*-PrOH (550 mL) and then re-concentrated. Add *i*-PrOH (550 mL) and stir for 1 h at ambient temperature. Cool the suspension to 5 °C and collect the solid by filtration then dry to afford tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-[2-(1-cyano-1-methyl-ethyl)-4-pyridyl]pyrazin-2-yl]isoxazol-3-yl]phenyl]methyl]-N-tetrahydropyran-4-yl-carbamate (Compound I-1) (81.9 g; 68%, yield, 98.7 area % purity by HPLC) as a cream-colored powder.


Form Change to Compound I-1•HCl•1.5 H₂O




[00161] A suspension of tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-[2-(1-cyano-1-methyl-ethyl)-4-pyridyl]pyrazin-2-yl]isoxazol-3-yl]phenyl]methyl]-N-tetrahydropyran-4-yl-carbamate (Compound I-1) (36.0 g, 72.6 mmol) in CH₃CN (720 mL) is stirred at ambient temperature (20 °C) in a flask equipped with mechanical stirring. A 1 M aqueous solution of HCl (72.6 mL; 72.6 mmol) is added. The suspension is stirred at ambient temperature for 20 h. The solid is collected by filtration. The filter-cake is washed with CH₃CN (3 x 50 mL) then dried under vacuum with high humidity for 2 h to afford

Compound I-1•HCl•1.5 H₂O (30.6 g; 74%) yield, 98.8 area % purity by HPLC) as a yellow powder. ¹H NMR (400 MHz, DMSO) δ 9.63 (d, *J* = 4.7 Hz, 2H), 9.05 (s, 1H), 8.69 (d, *J* = 5.2 Hz, 1H), 8.21 (s, 1H), 8.16 – 8.03 (m, 3H), 7.84 (t, *J* = 4.1 Hz, 3H), 7.34 (br s, 2H), 4.40 – 4.18 (m, 2H), 3.94 (dd, *J* = 11.2, 3.9 Hz, 2H), 3.32 (t, *J* = 11.2 Hz, 3H), 2.17 – 2.00 (m, 2H), 1.81 (s, 6H), 1.75 (dd, *J* = 12.1, 4.3 Hz, 2H).

Example 2: Synthesis of 3-[3-[4-[dideuterio(methylamino)methyl]phenyl]isoxazol-5-yl]-5-(4-isopropylsulfonylphenyl)pyrazin-2-amine (Compound II-1)




Step 1: 5-Bromo-3-((trimethylsilyl)ethynyl)pyrazin-2-amine



[00162] (Trimethylsilyl)acetylene (1.845 g, 2.655 mL, 18.78 mmol) was added dropwise to a solution of 3,5-dibromopyrazin-2-amine (compound **i**) (5 g, 19.77 mmol) in DMF (25


mL). Triethylamine (10.00 g, 13.77 mL, 98.85 mmol), copper(I) iodide (451.7 mg, 2.372 mmol) and Pd(PPh₃)₄ (1.142 g, 0.9885 mmol) were then added and the resulting solution stirred at RT for 30 minutes. The reaction mixture was diluted with EtOAc and water and the layers separated. The aqueous layer was extracted further with EtOAc and the combined organic layers washed with water, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography eluting with 15% EtOAc/Petroleum ether to give the product as a yellow solid (3.99 g, 75% Yield). ¹H NMR (400.0 MHz, DMSO) δ 0.30 (9H, s), 8.06 (1H, s); MS (ES⁺) 271.82.

Step 2: *tert*-Butyl N-*tert*-butoxycarbonyl-N-[5-bromo-3-((trimethylsilyl)ethynyl)pyrazin-2-yl]carbamate




[00163] 5-Bromo-3-(2-trimethylsilylethynyl)pyrazin-2-amine (2.85 g, 10.55 mmol) was dissolved in DCM (89.06 mL) and treated with Boc anhydride (6.908 g, 7.272 mL, 31.65 mmol) followed by DMAP (128.9 mg, 1.055 mmol). The reaction was allowed to stir at ambient temperature for 2 hours. The mixture was then diluted with DCM and NaHCO₃ and the layers separated. The aqueous layer was extracted further with DCM, dried (MgSO₄), filtered and concentrated *in vacuo*. The resultant residue was purified by column chromatography eluting with dichloromethane to give the desired product as a colourless oil (4.95g, 99% Yield). ¹H NMR (400.0 MHz, DMSO) δ 0.27 (9H, s), 1.42 (18H, s), 8.50 (1H, s); MS (ES⁺) 472.09.

Step 3: *tert*-Butyl N-(3-ethynyl-5-(4-(isopropylsulfonyl)phenyl)pyrazin-2-yl)N-*tert*butoxycarbonyl-carbamate *tert*-butyl



[00164] *N*- [5-Bromo-3-(2-trimethylsilylethynyl)pyrazin-2-yl]-*N*-tert-butoxycarbonylcarbamate (3 g, 6.377 mmol) and (4-isopropylsulfonylphenyl)boronic acid (1.491 g, 6.536 mmol) were dissolved in MeCN/water (60/12 mL). K₃PO₄ (2.706 g, 12.75 mmol) was added and the reaction mixture was degassed with a flow of nitrogen (5 cycles). Pd[P(tBu)₃]₂ (162.9 mg, 0.3188 mmol) was added and the resulting mixture was stirred at room temperature for 1h. The reaction mixture was poured quickly into a mixture of ethyl acetate (500 mL), water (90 mL) and 1% aqueous sodium metabisulphite at 4 °C, shaken well and the layer separated. The organic fraction was dried over MgSO₄, filtered and the filtrate was treated with 3-mercaptopropyl ethyl sulphide on silica (0.8mmol/g, 1 g), pre-absorbed onto silica gel then purified by column chromatography on silica gel eluting with 30-40% EtOAc/petroleum ether. The solvents were concentrated in vacuo to leave the product as a yellow viscous oil that was triturated with petroleum ether to yield the product as beige crystals (1.95 g, 61% Yield); ¹H NMR (400 MHz, DMSO) δ 1.20 (m, 6H), 1.39 (s, 18H), 3.50 (m, 1H), 5.01 (s, 1H), 8.03 (m, 2H), 8.46 (m, 2H) and 9.37 (s, 1H).


Step 4: 4-(Dimethoxymethyl)benzamide



[00165] A mixture of methyl 4-(dimethoxymethyl)benzoate (3.8 g, 18.08 mmol) and 7M NH₃ in MeOH (30 mL of 7 M, 210.0 mmol) in a sealed tube was heated at 110 °C for 22 hours. A further portion of 7M NH₃ in MeOH (20 mL of 7 M, 140.0 mmol) was added and the reaction heated at 135 °C for 23 hours. The reaction was cooled to ambient temperature and the solvent removed *in vacuo*. The residue was re-submitted to the reaction conditions (7M NH₃ in MeOH (30 mL of 7 M, 210.0 mmol) at 115 °C) for a further 16 hours. The solvent was removed *in vacuo* and the residue triturated from Et₂O. The resultant precipitate was isolated by filtration to give the sub-title compound as a white solid (590 mg, 17% yield). The filtrate was purified by column chromatography (ISCO Companion, 40 g column, eluting with 0 to 100% EtOAc/Petroleum Ether to 10% MeOH/EtOAc, loaded in EtOAc/MeOH) to give a further portion of the sub-title product as a white solid (225 mg, 6% Yield). Total


isolated (815 mg, 23% Yield); ^1H NMR (400 MHz, DMSO) δ 3.26 (s, 6H), 5.44 (s, 1H), 7.37 (s, 1H), 7.46 (d, J = 8.0 Hz, 2H), 7.84 - 7.91 (m, 2H) and 7.98 (s, 1H) ppm; MS (ES⁺) 196.0.

Steo 5: Dideuterio-[4-(dimethoxymethyl)phenyl]methanamine




LiDH_4 (12.52 mL of 1 M, 12.52 mmol) was added dropwise to a stirred solution of 4-(dimethoxymethyl)benzamide (815 mg, 4.175 mmol) in THF (20 mL) at 0 °C under an atmosphere of nitrogen. The reaction was heated at reflux for 16 hours then cooled to ambient temperature. The reaction was quenched by the sequential addition of D_2O (1 mL), 15% NaOH in D_2O (1 mL) and D_2O (4 mL). The resultant solid was removed by filtration and washed with EtOAc. The filtrate was concentrated *in vacuo* and the residue dried by azeotropic distillation with toluene (x 3) to give the sub-title compound as a yellow oil (819 mg) that was used without further purification; ^1H NMR (400 MHz, DMSO) δ 3.23 (s, 6H), 5.36 (s, 1H) and 7.30 - 7.35 (m, 4H) ppm; MS (ES⁺) 167.0.

Step 6: *tert*-Butyl *N*-[dideuterio-[4-(dimethoxymethyl)phenyl]methyl]carbamate




[00166] Et_3N (633.7 mg, 872.9 μL , 6.262 mmol) was added to a stirred suspension of dideuterio-[4-(dimethoxymethyl)phenyl]methanamine (765 mg, 4.175 mmol) in THF (15 mL) at 0 °C. The reaction was allowed to stir at this temperature for 30 minutes then Boc_2O (956.8 mg, 1.007 mL, 4.384 mmol) was added in portions. The reaction was allowed to warm to ambient temperature and stirred for 18 hours. The solvent was removed *in vacuo* and the residue was purified by column chromatography (ISCO Companion, 120 g column, eluting with 0 to 50% EtOAc/Petroleum Ether, loaded in DCM) to give the sub-title product as a colourless oil (1.04 g, 88% Yield); ^1H NMR (400 MHz, DMSO) δ 1.40 (s, 9H), 3.23 (s, 6H), 5.36 (s, 1H), 7.24 (d, J = 8.2 Hz, 2H), 7.33 (d, J = 8.0 Hz, 2H) and 7.38 (s, 1H) ppm.

Step 7: *tert*-Butyl *N*-[dideuterio-[4-(dimethoxymethyl)phenyl]methyl]-*N*-methylcarbamate



[00167] LiHMDS (1M in THF) (1.377 mL of 1 M, 1.377 mmol) was added dropwise to a stirred solution of *tert*-butyl *N*-[dideuterio-[4-(dimethoxymethyl)phenyl]methyl]carbamate (300 mg, 1.059 mmol) in THF (5 mL) at -78 °C. The solution was stirred at this temperature for 10 minutes then iodomethane (225.4 mg, 98.86 μ L, 1.588 mmol) was added dropwise and the mixture allowed to warm to ambient temperature over 1 hour. The reaction was again cooled to -78 °C and LiHMDS (1M in THF) (635.4 μ L of 1 M, 0.6354 mmol) was added. After 10 minutes iodomethane (105.2 mg, 46.14 μ L, 0.7413 mmol) was added and the reaction allowed to warm to ambient temperature over 6 hours. The mixture was diluted with EtOAc and the organic layer washed with saturated aqueous NaHCO₃ (x 2), brine (x 1), dried (MgSO₄) filtered and concentrated *in vacuo*. The residue was purified by column chromatography (ISCO Companion, 24 g column, eluting with 0 to 30% EtOAc/Petroleum Ether, loaded in DCM) to give the sub-title product as a colourless oil (200 mg, 63% Yield); ¹H NMR (400 MHz, DMSO) δ 1.41 (d, *J* = 27.7 Hz, 9H), 2.76 (s, 3H), 3.24 (s, 6H), 5.37 (s, 1H), 7.23 (d, *J* = 7.9 Hz, 2H) and 7.37 (d, *J* = 8.0 Hz, 2H) ppm.


Step 8: *tert*-Butyl *N*-[dideuterio-[4-[hydroxyiminomethyl]phenyl]methyl]-*N*-(methyl)carbamate



[00168] Hydroxylamine hydrochloride (51.15 mg, 0.7361 mmol) was added to a stirred solution of *tert*-butyl *N*-[dideuterio-[4-(dimethoxymethyl)phenyl]methyl]-*N*-methylcarbamate (199 mg, 0.6692 mmol) in THF (10 mL)/water (1.000 mL) and the reaction allowed to stir at ambient temperature for 4 hours. The reaction was partitioned between DCM and brine and the layers separated. The aqueous layer was extracted with DCM (x 2) and the combined organic extracts washed with brine (x 1), dried (MgSO₄), filtered and


concentrated *in vacuo* to give the sub-title compound as a white solid (180 mg, 100 % Yield); ¹H NMR (400 MHz, DMSO) δ 1.41 (d, J = 24.6 Hz, 9H) 2.76 (s, 3H), 7.25 (d, J = 8.1 Hz, 2H), 7.58 (d, J = 8.0 Hz, 2H), 8.13 (s, 1H) and 11.20 (s, 1H) ppm; MS (ES⁺) 211.0 (M-Boc).

Step 9: *tert*-Butyl N-[[4-[chloro-*N*-hydroxy-carbonimidoyl]phenyl]-dideuterio-methyl]-*N*-methyl-carbamate



[00169] *tert*-Butyl N-[dideuterio-[4-[hydroxyiminomethyl]phenyl]methyl]-*N*-(methyl)carbamate (178 mg, 0.6683 mmol) in DMF (2 mL) was treated with NCS (89.24 mg, 0.6683 mmol) and the reaction warmed to 65 °C for 1 hour. The reaction was cooled to ambient temperature and diluted with water. The mixture was extracted with EtOAc (x 2) and the combined organic extracts washed with brine (x 4), dried (MgSO₄), filtered and concentrated *in vacuo* to give the sub-title compound as a white solid (188 mg, 94% Yield); ¹H NMR (400 MHz, DMSO) δ 1.42 (d, J = 24.7 Hz, 9H), 2.78 (s, 3H), 7.32 (d, J = 8.4 Hz, 2H), 7.78 (d, J = 8.2 Hz, 2H) and 12.36 (s, 1H) ppm.


Step 10: *tert*-Butyl N-[[4-[5-[3-[bis(*tert*-butoxycarbonyl)amino]-6-(4-isopropylsulfonylphenyl)pyrazin-2-yl]isoxazol-3-yl]phenyl]-dideuterio-methyl]-*N*-methyl-carbamate



[00170] Et₃N (36.31 mg, 50.01 μL, 0.3588 mmol) was added dropwise to a stirred solution of *tert*-butyl N-*tert*-butoxycarbonyl-N-[3-ethynyl-5-(4-isopropylsulfonylphenyl)pyrazin-2-yl]carbamate (150 mg, 0.2990 mmol) and *tert*-Butyl N-

[[4-[chloro-*N*-hydroxy-carbonimidoyl]phenyl]-dideuterio-methyl]-*N*-methyl-carbamate (89.93 mg, 0.2990 mmol) in anhydrous THF (3 mL) and the reaction mixture heated at 65 °C for 3 hours. The reaction mixture was cooled to ambient temperature and diluted with EtOAc/brine. Water was added until the aqueous layer became clear and the layers were separated. The aqueous layer was extracted with EtOAc (x 1) and the combined organic extracts were washed with brine (x 1), dried (MgSO₄), filtered and concentrated in vacuo. The residue was purified by column chromatography (ISCO Companion, 40 g column, eluting with 0 to 30% EtOAc/Petroleum Ether, loaded in DCM) to give the sub-title product as a white solid (134 mg, 59% Yield); ¹H NMR (400 MHz, DMSO) δ 1.22 (d, J = 6.8 Hz, 6H) 1.32 (s, 18H), 1.43 (d, J = 23.1 Hz, 9H), 2.82 (s, 3H), 3.56 (pent, 1H), 7.43 (d, J = 8.3 Hz, 3H), 8.02 - 8.03 (m 3H), 8.06 - 8.11 (m, 2H), 8.62 - 8.67 (m, 2H) and 9.51 (s, 1H) ppm; MS (ES⁺) 666.2 (M-Boc).


Step 11: 3-[3-[4-[Dideuterio(methylamino)methyl]phenyl]isoxazol-5-yl]-5-(4-isopropylsulfonylphenyl)pyrazin-2-amine (compound II-1)




II-1

[00171] 3M HCl in MeOH (1.167 mL of 3 M, 3.500 mmol) was added to a stirred solution of *tert*-butyl *N*-[[4-[5-[3-[bis(*tert*-butoxycarbonyl)amino]-6-(4-isopropylsulfonylphenyl)pyrazin-2-yl]isoxazol-3-yl]phenyl]-dideuterio-methyl]-*N*-methyl-carbamate (134 mg, 0.1750 mmol) in DCM (5 mL) and the reaction heated at reflux for 16 hours. The reaction was cooled to ambient temperature and the resultant precipitate was isolated by filtration and dried under vacuum at 40 °C to give the di-HCl salt of the title compound as a yellow solid (58.8 mg, 62% Yield); ¹H NMR (400 MHz, DMSO) δ 1.20 (d, J = 6.8 Hz, 6H), 2.60 (t, J = 5.4 Hz, 3H), 3.48 (hept, J = 6.8 Hz, 1H), 7.22 (br s, 2H), 7.69 - 7.75 (m, 2H), 7.85 (s, 1H), 7.92 - 7.99 (m, 2H), 8.08 - 8.15 (m, 2H) 8.37 - 8.42 (m, 2H), 8.97 (s, 1H) and 9.10 (d, J = 5.8 Hz, 2H) ppm; MS (ES⁺) 466.2.

Example 3: Synthesis of 3-[3-[4-[dideuterio-(trideuteriomethylamino)methyl]phenyl]isoxazol-5-yl]-5-(4-isopropylsulfonylphenyl)pyrazin-2-amine (Compound II-2)




Step 1: *tert*-Butyl *N*-[dideuterio-[4-(dimethoxymethyl)phenyl]methyl]-I-(trideuteriomethyl)carbamate




[00172] LiHMDS (1M in THF) (1.181 mL of 1 M, 1.181 mmol) was added dropwise to a stirred solution of *tert*-butyl *N*-[dideuterio-[4-(dimethoxymethyl)phenyl]methyl]carbamate (300 mg, 1.059 mmol) in THF (5 mL) at -78 °C. The solution was stirred at this temperature for 30 minutes then trideuterio(iodo)methane (198.0 mg, 84.98 μ L, 1.366 mmol) was added dropwise and the mixture allowed to warm to ambient temperature over 21 hours. The reaction was again cooled to -78 °C and a further portion of LiHMDS (1M in THF) (635.4 μ L of 1 M, 0.6354 mmol) was added. After 15 minutes more trideuterio(iodo)methane (76.75 mg, 32.94 μ L, 0.5295 mmol) was added and the reaction allowed to warm to ambient temperature over 5 hours. The mixture was diluted with EtOAc and the organic layer washed with saturated aqueous NaHCO₃ (x 2), brine (x 1), dried (MgSO₄) filtered and concentrated *in vacuo*. The residue was purified by column chromatography (ISCO Companion, 24 g column, eluting with 0 to 30% EtOAc/Petroleum Ether, loaded in DCM) to give the sub-title product as a colourless oil (213 mg, 67% Yield); ¹H NMR (400 MHz, DMSO) δ 1.36 - 1.42 (m, 9H) 3.22 (s, 6H), 5.35 (s, 1H), 7.21 (d, *J* = 7.8 Hz, 2H) and 7.35 (d, *J* = 7.7 Hz, 2H) ppm.

Step 2: *tert*-Butyl *N*-[dideuterio-[4-[hydroxyiminomethyl]phenyl]methyl]-*N*-(trideuteriomethyl)carbamate




[00173] Hydroxylamine hydrochloride (53.95 mg, 0.7763 mmol) was added to a stirred solution of *tert*-butyl *N*-[dideuterio-[4-(dimethoxymethyl)phenyl]methyl]-*N*-(trideuteriomethyl)carbamate (212 mg, 0.7057 mmol) in THF (10 mL)/water (1.000 mL) and the reaction allowed to stir at ambient temperature for 22 hours. The reaction was partitioned between DCM and brine and the layers separated. The aqueous layer was extracted with DCM (x 2) and the combined organic extracts washed with brine (x 1), dried (MgSO₄), filtered and concentrated *in vacuo* to give the sub-title compound as a white solid (190 mg, 100% Yield).); ¹H NMR (400 MHz, DMSO) δ 1.41 (d, J = 24.2 Hz, 9H), 7.25 (d, J = 8.1 Hz, 2H), 7.58 (d, J = 8.0 Hz, 2H), 8.13 (s, 1H) and 11.20 (s, 1H) ppm.

Step 3: *tert*-Butyl *N*-[[4-[chloro-*N*-hydroxy-carbonimidoyl]phenyl]-dideuterio-methyl]-*N*-(trideuteriomethyl)carbamate



[00174] *tert*-Butyl *N*-[dideuterio-[4-[hydroxyiminomethyl]phenyl]methyl]-*N*-(trideuteriomethyl)carbamate (190.0 mg, 0.7054 mmol) in DMF (2 mL) was treated with NCS (94.19 mg, 0.7054 mmol) and the reaction warmed to 65 °C for 1 hour. The reaction was cooled to ambient temperature and diluted with water. The mixture was extracted with EtOAc (x 2) and the combined organic extracts washed with brine (x 4), dried (MgSO₄), filtered and concentrated *in vacuo* to give the sub-title compound as a white solid (198 mg, 93% Yield); ¹H NMR (400 MHz, DMSO) δ 1.41 (d, J = 26.0 Hz, 9H), 7.32 (d, J = 8.3 Hz, 2H), 7.78 (d, J = 8.2 Hz, 2H) and 12.36 (s, 1H) ppm.


Step 4: *tert*-Butyl *N*-[[4-[5-[3-[bis(*tert*-butoxycarbonyl)amino]-6-(4-isopropylsulfonylphenyl)pyrazin-2-yl]isoxazol-3-yl]phenyl]-dideuterio-methyl]-*N*-(trideuteriomethyl)carbamate



xvi

[00175] Et₃N (36.31 mg, 50.01 μ L, 0.3588 mmol) was added dropwise to a stirred solution of *tert*-butyl *N*-*tert*-butoxycarbonyl-*N*-[3-ethynyl-5-(4-isopropylsulfonylphenyl)pyrazin-2-yl]carbamate (150 mg, 0.2990 mmol) and *tert*-butyl *N*-[[4-[chloro-*N*-hydroxy-carbonimidoyl]phenyl]-dideuterio-methyl]-*N*-(trideuteriomethyl)carbamate (90.84 mg, 0.2990 mmol) in anhydrous THF (3 mL) and the reaction mixture heated at 65 °C for 3.5 hours. The reaction mixture was cooled to ambient temperature and diluted with EtOAc/brine. Water was added until the aqueous layer became clear and the layers were separated. The aqueous layer was extracted with EtOAc (x 1) and the combined organic extracts were washed with brine (x 1), dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified by column chromatography (ISCO Companion, 40 g column, eluting with 0 to 35% EtOAc/Petroleum Ether, loaded in DCM) to give the sub-title product as a white solid (158 mg, 69% Yield); ¹H NMR (400 MHz, DMSO) δ 1.22 (d, *J* = 6.8 Hz, 6H), 1.44 (d, *J* = 22.0 Hz, 9H), 3.56 (dt, *J* = 13.5, 6.7 Hz, 2H), 7.43 (d, *J* = 8.2 Hz, 3H), 8.02 (d, *J* = 6.9 Hz, 2H), 8.08 (d, *J* = 8.7 Hz, 2H), 8.65 (d, *J* = 8.8 Hz, 2H) and 9.51 (s, 1H) ppm; MS (ES⁺) 669.3 (M-Boc).


Step 5: 3-[3-[4-[dideuterio-(trideuteriomethylamino)methyl]phenyl]isoxazol-5-yl]-5-(4-isopropylsulfonylphenyl)pyrazin-2-amine (compound II-2)




II-2

[00176] 3M HCl in MeOH (1.361 mL of 3 M, 4.084 mmol) was added to a stirred solution of *tert*-butyl *N*-[[4-[5-[3-[bis(*tert*-butoxycarbonyl)amino]-6-(4-isopropylsulfonylphenyl)pyrazin-2-yl]isoxazol-3-yl]phenyl]-dideuterio-methyl]-*N*-(trideuteriomethyl)carbamate (157 mg, 0.2042 mmol) in DCM (5 mL) and the reaction heated at reflux for 16 hours. The reaction was cooled to ambient temperature and the resultant precipitate was isolated by filtration and dried under vacuum at 40 °C to give the di-HCl salt of the title compound as a yellow solid (72.5 mg, 66% Yield); ¹H NMR (400 MHz, DMSO) δ 1.20 (d, J = 6.8 Hz, 6H), 3.48 (dq, J = 13.6, 6.7 Hz, 1H), 7.21 (s, 2H), 7.68 - 7.78 (m, 2H), 7.85 (s, 1H), 7.91 - 7.99 (m, 2H), 8.08 - 8.13 (m, 2H), 8.36 - 8.42 (m, 2H), 8.96 (s, 1H) and 9.14 (s, 2H) ppm; MS (ES⁺) 469.1.

Example 4: Synthesis of 5-(4-isopropylsulfonylphenyl)-3-[3-[4-[(trideuteriomethylamino)methyl]phenyl]isoxazol-5-yl]pyrazin-2-amine (Compound II-3)




Step 1: Methyl 4-[(*tert*-butoxycarbonylamino)methyl]benzoate



xviii

[00177] Et₃N (1.882 g, 2.592 mL, 18.60 mmol) was added to a stirred suspension of methyl 4-(aminomethyl)benzoate (Hydrochloric Acid (1)) (1.5 g, 7.439 mmol) in THF (20 mL) at 0 °C. The reaction was allowed to stir at this temperature for 30 minutes then Boc₂O (1.705 g, 1.795 mL, 7.811 mmol) was added in portions. The reaction was allowed to warm to ambient temperature and stirred for 18 hours. The mixture was diluted with EtOAc. The organic layer was washed with 1M aqueous HCl (x 2), saturated aqueous NaHCO₃ (x 2) and brine (x 1). The organic layer was dried (MgSO₄), filtered and concentrated *in vacuo* to give the sub-title compound as a white solid that was used without further purification (1.93 g, 98% Yield); ¹H NMR (400 MHz, DMSO) δ 1.40 (s, 9H), 3.85 (s, 3H), 4.20 (d, J = 6.1 Hz, 2H), 7.38 (d, J = 8.2 Hz, 2H), 7.49 (t, J = 6.1 Hz, 1H) and 7.92 (d, J = 8.2 Hz, 2H) ppm; MS (ES⁺) 251.1 (M-Me).

Step 2: Methyl 4-[[*tert*-butoxycarbonyl(trideuteriomethyl)amino]methyl]benzoate




xix

[00178] LiHMDS (1M in THF) (8.112 mL of 1 M, 8.112 mmol) was added dropwise to a stirred solution of methyl 4-[(*tert*-butoxycarbonylamino)methyl]benzoate (1.93 g, 7.275 mmol) in THF (10 mL) at -78 °C. The solution was stirred at this temperature for 30 minutes then trideuterio(iodo)methane (1.360 g, 9.385 mmol) was added dropwise and the mixture allowed to warm to ambient temperature over 3 hours. The reaction was cooled to -78 °C and a further portion of LiHMDS (1M in THF) (2.182 mL of 1 M, 2.182 mmol) was added. After 10 minutes a further portion of trideuterio(iodo)methane (527.4 mg, 3.638 mmol) was added and the reaction allowed to warm to ambient temperature over 17 hours. The mixture was diluted with EtOAc and the organic layer washed with saturated aqueous NaHCO₃ (x 2), brine (x 1), dried (MgSO₄) filtered and concentrated *in vacuo*. The residue was purified by

column chromatography (ISCO Companion, 120 g column, eluting with 0 to 30% EtOAc/Petroleum Ether, loaded in DCM) to give the sub-title product as a pale yellow oil (1.37 g, 67% Yield); ^1H NMR (400 MHz, DMSO) δ 1.38 (d, J = 44.2 Hz, 9H), 3.83 (s, 3H), 4.43 (s, 2H), 7.33 (d, J = 8.2 Hz, 2H) and 7.94 (d, J = 8.1 Hz, 2H) ppm; MS (ES+) 268.1 (M-Me)


Step 3: *tert*-Butyl *N*-[[4-(hydroxymethyl)phenyl]methyl]-*N*-(trideuteriomethyl)carbamate



xx

[00179] LiBH_4 (158.5 mg, 7.278 mmol) was added to a stirred solution of methyl 4-[[*tert*-butoxycarbonyl(trideuteriomethyl)amino]methyl]benzoate (1.37 g, 4.852 mmol) in THF (10 mL) and the reaction warmed to 85 °C for 15 hours. A further portion of LiBH_4 (158.5 mg, 7.278 mmol) was added and the reaction stirred at 65 °C for a further 7 hours. The reaction mixture was cooled to ambient temperature then poured onto crushed ice and whilst stirring, 1M HCl was added dropwise until no effervescence was observed. The mixture was stirred for 10 minutes then saturated aqueous NaHCO_3 was added until the mixture was at pH 8. The aqueous layer was extracted with EtOAc (x 3) and the combined organic extracts dried (MgSO_4), filtered and concentrated *in vacuo*. The residue was purified by column chromatography (ISCO Companion, 120 g column, eluting with 0 to 100% EtOAc/Petroleum Ether, loaded in DCM) to give the sub-title product as a colourless oil (1.03 g, 84% Yield); ^1H NMR (400 MHz, DMSO) δ 1.42 (d, J = 14.6 Hz, 9H), 4.35 (s, 2H), 4.48 (d, J = 5.7 Hz, 2H), 5.15 (t, J = 5.7 Hz, 1H), 7.18 (d, J = 7.9 Hz, 2H) and 7.30 (d, J = 7.7 Hz, 2H) ppm; MS (ES+) 181.1 (M- O^tBu).

Step 4: *tert*-Butyl *N*-[(4-formylphenyl)methyl]-*N*-(trideuteriomethyl)carbamate




xxi

[00180] MnO_2 (5.281 g, 1.051 mL, 60.75 mmol) was added to a stirred solution of *tert*-butyl *N*-[[4-(hydroxymethyl)phenyl]methyl]-*N*-(trideuteriomethyl)carbamate (1.03 g, 4.050


mmol) in DCM (10 mL) and the reaction stirred at ambient temperature for 20 hours. The reaction was filtered through a pad of Celite and washed with DCM. The filtrate was concentrated *in vacuo* to give the sub-title compound as a colourless oil (891 mg, 88% Yield); ¹H NMR (400 MHz, DMSO) δ 1.40 (d, J = 43.4 Hz, 9H), 4.48 (s, 2H), 7.43 (d, J = 8.0 Hz, 2H), 7.91 (d, J = 7.9 Hz, 2H) and 10.00 (s, 1H), ppm.

Step 5: *tert*-butyl *N*-[[4-[hydroxyiminomethyl]phenyl]methyl]-*N*-(trideuteriomethyl)carbamate



[00181] Hydroxylamine (466.0 μL of 50 %w/v, 7.054 mmol) was added to a stirred solution of *tert*-butyl *N*-[(4-formylphenyl)methyl]-*N*-(trideuteriomethyl)carbamate (890 mg, 3.527 mmol) in ethanol (5 mL) and the reaction mixture stirred at ambient temperature for 45 minutes. The reaction mixture was concentrated *in vacuo* and the residue taken up in water and extracted with EtOAc (x 3). The combined organic extracts were washed with brine (x 1), dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was triturated from petroleum ether and the precipitate isolated by filtration to give the sub-title product as a white solid (837 mg, 89% Yield); ¹H NMR (400 MHz, DMSO) δ 1.41 (d, J = 25.8 Hz, 9H), 4.38 (s, 2H), 7.24 (d, J = 8.0 Hz, 2H), 7.58 (d, J = 8.0 Hz, 2H), 8.13 (s, 1H) and 11.20 (s, 1H) ppm; MS (ES⁺) 212.0 (M-^tBu).


Step 6: *tert*-Butyl *N*-[[4-[chloro-*N*-hydroxy-carbonimidoyl]phenyl]methyl]-*N*-(trideuteriomethyl)carbamate



[00182] *tert*-butyl *N*-[[4-[hydroxyiminomethyl]phenyl]methyl]-*N*-(trideuteriomethyl)carbamate (250 mg, 0.9351 mmol) in DMF (2.5 mL) was treated with NCS (124.9 mg, 0.9351 mmol) and the reaction warmed to 65 °C for 1 hour. The reaction was cooled to ambient temperature and diluted with water. The mixture was extracted with

EtOAc (x 2) and the combined organic extracts washed with brine (x 4), dried (MgSO₄), filtered and concentrated *in vacuo* to give the sub-title compound as a white solid (259 mg, 92% Yield); ¹H NMR (400 MHz, DMSO) δ 1.41 (d, J = 29.6 Hz, 9H), 4.42 (s, 2H), 7.31 (d, J = 8.3 Hz, 2H), 7.78 (d, J = 8.0 Hz, 2H), and 12.38 (s, 1H), ppm.


Step 7: *tert*-Butyl *N*-[[4-[5-[3-[bis(*tert*-butoxycarbonyl)amino]-6-(4-isopropylsulfonylphenyl)pyrazin-2-yl]isoxazol-3-yl]phenyl]methyl]-*N*-(trideuteriomethyl)carbamate



xxiv

[00183] Et₃N (48.41 mg, 66.68 μL, 0.4784 mmol) was added dropwise to a stirred solution of *tert*-butyl *N*-*tert*-butoxycarbonyl-*N*-[3-ethynyl-5-(4-isopropylsulfonylphenyl)pyrazin-2-yl]carbamate (200 mg, 0.3987 mmol) and *tert*-butyl *N*-[[4-[chloro-*N*-hydroxy-carbonimidoyl]phenyl]methyl]-*N*-(trideuteriomethyl)carbamate (120.3 mg, 0.3987 mmol) in anhydrous THF (5 mL) and the reaction mixture heated at 65 °C for 2.5 hours. The reaction mixture was cooled to ambient temperature and diluted with EtOAc/brine. Water was added until the aqueous layer became clear and the layers were separated. The aqueous layer was extracted with EtOAc (x 1) and the combined organic extracts were washed with brine (x 1), dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified by column chromatography (ISCO Companion, 40 g column, eluting with 0 to 20% EtOAc/Petroleum Ether, loaded in DCM) to give the sub-title product as a white solid (213.5 mg, 70% Yield); ¹H NMR (400 MHz, DMSO) δ 1.22 (d, J = 6.8 Hz, 6H), 1.31 (s, 18H), 1.43 (d, J = 26.2 Hz, 9H), 3.51 - 3.60 (m, 1H), 4.47 (s, 2H), 7.42 (d, J = 8.1 Hz, 2H), 8.03 (d, J = 5.2 Hz, 3H), 8.08 (d, J = 8.6 Hz, 2H), 8.65 (d, J = 8.6 Hz, 2H) and 9.52 (s, 1H) ppm; MS (ES⁺) 667.4 (M-Boc).


Step 8: 5-(4-Isopropylsulfonylphenyl)-3-[3-[4-[(trideuteriomethylamino)methyl]phenyl]isoxazol-5-yl]pyrazin-2-amine (Compound II-3)




II-3

[00184] 3M HCl in MeOH (1.5 mL of 3 M, 4.500 mmol) was added to a stirred solution of *tert*-butyl *N*-[[4-[5-[3-[bis(*tert*-butoxycarbonyl)amino]-6-(4-isopropylsulfonylphenyl)pyrazin-2-yl]isoxazol-3-yl]phenyl]methyl]-*N*-(trideuteriomethyl)carbamate (213 mg, 0.2777 mmol) in DCM (6 mL) and the reaction heated at reflux for 15 hours. A further portion of 3M HCl in MeOH (0.5 mL of 3 M, 1.500 mmol) was added and the reaction heated at reflux for a further 7 hours. The reaction was cooled to ambient temperature and the resultant precipitate was isolated by filtration and dried under vacuum at 40 °C to give the di-HCl salt of the title compound as a yellow solid (97.6 mg, 65% Yield); ¹H NMR (400 MHz, DMSO) δ 1.20 (d, J = 6.8 Hz, 6H), 3.47 (tt, J = 14.0, 6.9 Hz, 1H), 4.19 - 4.25 (m, 2H), 7.23 (s, 2H), 7.72 (d, J = 8.4 Hz, 2H), 7.85 (s, 1H), 7.95 (d, J = 8.7 Hz, 2H), 8.11 (d, J = 8.4 Hz, 2H), 8.39 (d, J = 8.7 Hz, 2H), 8.97 (s, 1H) and 9.11 (s, 2H) ppm; MS (ES⁺) 467.2.

Example 5: Synthesis of 3-[3-[4-(Methylaminomethyl)phenyl]isoxazol-5-yl]-5-[4-[1,2,2,2-tetradeuterio-1-(trideuteriomethyl)ethyl]sulfonylphenyl]pyrazin-2-amine (Compound II-4)




Step 1: 1-[4-(Diethoxymethyl)phenyl]-N-methyl-methanamine



[00185] 2M methylamine in MeOH (288.1 mL, 576.2 mmol) was diluted with methanol (1.000 L) and stirred at ~20 °C. 4-(Diethoxymethyl)benzaldehyde (100 g, 480.2 mmol) was added dropwise over 1 minute and the reaction stirred at ambient temperature for 1.25 hours. Sodium borohydride (29.07 g, 30.76 mL, 768.3 mmol) was added portionwise over 20 minutes while maintaining the temperature between 20 and 30 °C with an ice-water bath. The reaction solution was stirred at ambient temperature overnight then quenched by the dropwise addition of NaOH (960.4 mL of 1.0 M, 960.4 mmol) over 20 minutes. The reaction was


stirred for 30 minutes and concentrated *in vacuo* to remove MeOH. The reaction was partitioned with MTBE (1.200 L) and the phases separated. The organic phase was washed with water (300.0 mL), dried (Na₂SO₄), and concentrated *in vacuo* to give the title compound as a yellow oil (102.9 g, 96% Yield); ¹H NMR (400 MHz, CDCl₃) δ 1.25 (t, 6H), 2.46 (s, 3H), 3.45 – 3.65 (m, 4H), 3.75 (s, 2H), 5.51 (s, 1H), 7.32 (d, 2H) and 7.44 (d, 2H) ppm.

Step 2: *tert*-Butyl *N*-[[4-(diethoxymethyl)phenyl]methyl]-*N*-methyl-carbamate



[00186] A 1-L glass-jacketed reactor was fitted with an overhead stirrer, thermocouple, and chiller. A solution of 1-[4-(diethoxymethyl)phenyl]-*N*-methyl-methanamine (80.0 g, 358.2 mmol) in DCM (480.0 mL) was stirred at 18 °C. A solution of Boc anhydride (79.75 g, 83.95 mL, 365.4 mmol) in DCM (160.0 mL) was added over 10 minutes and the solution was stirred at 20 - 25 °C overnight. The reaction mixture was filtered, rinsed with DCM (3 x 50 mL) and the filtrate concentrated *in vacuo* to afford give the title compound as a pale yellow liquid (116.6 g, quantitative yield); ¹H NMR (400 MHz, CDCl₃) δ 1.25 (t, 6H), 1.49 – 1.54 (2 x s, 9H), 2.78 – 2.83 (2 x s, 3H), 3.50 – 3.66 (m, 4H), 4.42 (s, 2H), 5.49 (s, 1H), 7.22 (d, 2H) and 7.45 (d, 2H) ppm.


Step 3: *tert*-Butyl *N*-[[4-[hydroxyiminomethyl]phenyl]methyl]-*N*-methyl-carbamate



[00187] A biphasic solution of *tert*-butyl *N*-[[4-(diethoxymethyl)phenyl]methyl]-*N*-methyl-carbamate (50.0 g, 154.6 mmol) in 2-MeTHF (400.0 mL) and Na₂SO₄ (100.0 mL of 10 %w/v, 70.40 mmol) was stirred at 8 - 10 °C in a 1-L, glass-jacketed reactor. Hydroxylamine hydrochloride (46.38 mL of 5.0 M, 231.9 mmol) was added and the biphasic solution was stirred at 30 °C for 16 hours. The reaction was diluted with MTBE (200.0 mL) and the layers separated. The organic phase was washed with water (200.0 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo*. The residue was diluted with heptane (200.0


mL) and the resultant suspension was stirred at ambient temperature for 30 minutes. The solid was collected by filtration to give the title compound as a white solid (36.5 g, 89% Yield); ¹H NMR (400 MHz, CDCl₃) δ 1.50 (s, 9H), 2.88 (br s, 3H), 4.60 (s, 2H), 7.26 (d, 2H), 7.52 (d, 2H) and 8.15 (s, 1H) ppm.

Step 4: *tert*-Butyl *N*-[[4-[chloro-*N*-hydroxy-carbonimidoyl]phenyl]methyl]-*N*-methyl-carbamate



[00188] A suspension of *tert*-butyl *N*-[[4-[hydroxyiminomethyl]phenyl]methyl]-*N*-methyl-carbamate (100 g, 378.3 mmol) in isopropyl acetate (1.000 L) was stirred at ambient temperature. *N*-Chlorosuccinimide (53.04 g, 397.2 mmol) was added and stirred at ambient temperature for 16 hours. The reaction was partitioned with water (500.0 mL) and the phases separated. The organic phase was washed with water (500.0 mL) (2 x), dried (Na₂SO₄), filtered and concentrated *in vacuo* to remove most of the solvent. Heptane (1.000 L) was added and the mixture concentrated *in vacuo* to remove most of the solvent. Heptane (1.000 L) was added and the resultant precipitate isolated by filtration. The filter-cake was washed with heptane (500 mL) and air-dried to give the title compound as an off-white powder (105.45 g, 93% Yield); ¹H NMR (400 MHz, CDCl₃) δ 1.48 (2 x s, 9H), 2.90 (2 x s, 3H), 4.47 (s, 2H), 7.26 (d, 2H), 7.77 (d, 2H) and 8.82 (s, 1H) ppm.


Step 5: *tert*-Butyl *N*-[[4-[5-[3-[bis(*tert*-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]isoxazol-3-yl]phenyl]methyl]-*N*-methyl-carbamate



[00189] A suspension of *tert*-butyl *N*-[[4-[chloro-*N*-hydroxy-carbonimidoyl]phenyl]methyl]-*N*-methyl-carbamate (100.0 g, 334.7 mmol) and *tert*-butyl *N*-tert-butoxycarbonyl-*N*-[3-ethynyl-5-(4-isopropylsulfonylphenyl)pyrazin-2-yl]carbamate


(121.2 g, 304.3 mmol) in DCM (1.212 L) was stirred at ambient temperature. Triethylamine (33.87 g, 46.65 mL, 334.7 mmol) was added in one portion and the reaction stirred at ambient temperature for 16 hours. The reaction was partitioned with water (606.0 mL) and the phases separated. The organic phase was washed with water (606.0 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo* to near dryness. Heptane (363.6 mL) was added and the mixture concentrated to about 300 mL. Further heptane (1.212 L) was added and the mixture heated to 90 °C with stirring. The mixture was slowly cooled to ambient temperature and stirred at this temperature for 1 hour. The resultant precipitate was isolated by filtration and the filter-cake washed with heptane (2 x 363.6 mL) and air-dried to give the title compound as a beige solid (181.8 g, 90% Yield); ¹H NMR (400 MHz, CDCl₃) δ 1.41 (s, 18H), 1.51 (s, 9H), 2.88 (2 x s, 3H), 4.50 (s, 2H), 7.36 – 7.38 (m, 3H), 7.86 (d, 2H) and 8.65 (s, 1H) ppm.

Step 6: 1-Bromo-4-[1,2,2,2-tetradeuterio-1-(trideuteriomethyl)ethyl]sulfanyl-benzene




[00190] Sodium hydride (246.5 mg, 6.163 mmol) was added portionwise to a stirred solution of 4-bromobenzenethiol (compound **xxxi**) (970.9 mg, 5.135 mmol) in DMF (10 mL) at 0 °C. After stirring at this temperature for 15 minutes 1,1,1,2,3,3,3-heptadeuterio-2-iodopropane (1 g, 5.649 mmol) was added and the reaction allowed to warm to ambient temperature over 18 hours. The reaction was quenched by the addition of water and the mixture stirred for 10 minutes. The mixture was extracted with diethyl ether (x 3) and the combined organic extracts washed with water (x 2), brine (x 2), dried (MgSO₄), filtered and concentrated *in vacuo* to give the sub-title compound that was used directly without further purification assuming 100% Yield and purity; ¹H NMR (500 MHz, DMSO) δ 7.25 – 7.37 (m, 2H) and 7.48 - 7.55 (m, 2H) ppm.

Step 7: 1-Bromo-4-[1,2,2,2-tetradeuterio-1-(trideuteriomethyl)ethyl]sulfonyl-benzene



[00191] mCPBA (2.875 g, 12.83 mmol) was added in portions to a stirred solution of 1-bromo-4-[1,2,2,2-tetradeuterio-1-(trideuteriomethyl)ethyl]sulfanyl-benzene (1.223 g, 5.134 mmol) in DCM (20 mL) at 0 °C and the reaction allowed to warm to ambient temperature over 17 hours. The mixture was washed 1M aqueous NaOH (x 2), saturated aqueous Na₂S₂O₃ (x 3), brine (x 1), dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified by column chromatography (ISCO Companion, 80 g column, eluting with 0 to 40% EtOAc/Petroleum Ether, loaded in DCM) to give the sub-title compound as a colourless oil (1.19 g, 86% Yield); ¹H NMR (500 MHz, DMSO) δ 7.77 - 7.81 (m, 2H) and 7.88 - 7.92 (m, 2H) ppm.


Step 8: 4,4,5,5-Tetramethyl-2-[4-[1,2,2,2-tetradeuterio-1-(trideuteriomethyl)ethyl]sulfonylphenyl]-1,3,2-dioxaborolane



[00192] Pd(dppf)Cl₂.DCM (179.8 mg, 0.2202 mmol) was added to a stirred suspension of 1-bromo-4-[1,2,2,2-tetradeuterio-1-(trideuteriomethyl)ethyl]sulfonyl-benzene (1.19 g, 4.404 mmol), bis(dipinacolato)diboron (1.342 g, 5.285 mmol) and KOAc (1.296 g, 13.21 mmol) in dioxane (10 mL). The reaction placed under an atmosphere of nitrogen via 5 x nitrogen/vacuum cycles and the mixture was heated at 80 °C for 4.5 hours. The reaction was cooled to ambient temperature and the solvent removed *in vacuo*. The residue was partitioned between Et₂O and water and the layers separated. The organic layer was dried (MgSO₄),

filtered and concentrated *in vacuo*. The residue was dissolved in 30% EtOAc/Petroleum ether (35 mL) and 1.2 g of Florosil was added. The mixture was stirred for 30 minutes then filtered, washing the solids with further aliquots of 30% EtOAc/Petrol (x 3). The filtrate was concentrated *in vacuo* and triturated from 10% EtOAc/petroleum ether. The resultant solid was isolated by filtration, washed with petroleum ether and dried *in vacuo* to give the sub-title compound as an off-white solid (1052.1 mg, 75% Yield); ¹H NMR (400 MHz, DMSO) δ 1.33 (s, 12H), 7.87 (d, J = 8.4 Hz, 2H) and 7.94 (d, J = 8.4 Hz, 2H) ppm.

Step 9: *tert*-Butyl N-[[4-[5-[3-[bis(*tert*-butoxycarbonyl)amino]-6-[4-[1,2,2,2-tetradeuterio-1-(trideuteriomethyl)ethyl]sulfonylphenyl]pyrazin-2-yl]isoxazol-3-yl]phenyl]methyl]-N-methyl-carbamate




XXXV

[00193] [1,1'-Bis(di-*tert*-butylphosphino)ferrocene]dichloropalladium(II) (106.8 mg, 0.1639 mmol) was added to a mixture of 4,4,5,5-tetramethyl-2-[4-[1,2,2,2-tetradeuterio-1-(trideuteriomethyl)ethyl]sulfonylphenyl]-1,3,2-dioxaborolane (1.3 g, 4.098 mmol), *tert*-butyl N-[[4-[5-[3-[bis(*tert*-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]isoxazol-3-yl]phenyl]methyl]-N-methyl-carbamate (2.707 g, 4.098 mmol) and K₂CO₃ (1.133 g, 8.200 mmol) in toluene (9.100 mL), EtOH (2.600 mL) and water (2.600 mL) and the reaction mixture was degassed with a flow of nitrogen (5 cycles).

The mixture was heated at 75 °C for 1.5 hours. The reaction was cooled to ambient temperature and water (5.2 mL) was added. After stirring the layers were separated and the organic layer dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The residue was triturated with IPA and the resultant precipitate isolated by filtration, washed with IPA (3 x 4 mL) and dried *in vacuo* at 50 °C to give the title compound as a white solid (2.4 g, 76% Yield); ¹H NMR (400 MHz, CDCl₃) δ 1.41 (s, 18H), 1.50 (s, 9H), 2.85 – 2.89 (m, 3H), 4.50 (s, 2H), 7.36 – 7.38 (m, 3H), 7.87 (d, 2H), 8.09 (d, 2H), 8.35 (d, 2H) and 9.06 (s, 1H) ppm.


Step 10: 3-[3-[4-(Methylaminomethyl)phenyl]isoxazol-5-yl]-5-[4-[1,2,2,2-tetradeuterio-1-(trideuteriomethyl)ethyl]sulfonylphenyl]pyrazin-2-amine (compound II-4)




II-4

[00194] Concentrated HCl (3.375 g, 2.812 mL of 37 %w/w, 34.25 mmol) was added to a solution of *tert*-butyl *N*-[[4-[5-[3-[bis(*tert*-butoxycarbonyl)amino]-6-[4-[1,2,2,2-tetradeuterio-1-(trideuteriomethyl)ethyl]sulfonylphenyl]pyrazin-2-yl]isoxazol-3-yl]phenyl]methyl]-*N*-methyl-carbamate (2.2 g, 2.854 mmol) in acetone (28.60 mL) and the reaction heated at reflux for 7 hours. The reaction was cooled to ambient temperature and the resultant precipitate isolated by filtration, washed with acetone (2 x 4.5 mL) and dried *in vacuo* at 50 °C to give the di-HCl salt of the title compound as a yellow solid (1.42 g, 92% Yield); ¹H NMR (400 MHz, DMSO) δ 2.58 (t, 3H), 4.21 (t, 2H), 5.67 (br s, 2H), 7.74 (d, 2H), 7.85 (s, 1H), 7.94 (d, 2H), 8.10 (d, 2H), 8.38 (d, 2H), 8.96 (s, 1H) and 9.33 (br s, 2H) ppm; MS (ES⁺) 471.8.

Example 6: Synthesis of 5-(4-(tert-butylsulfonyl)phenyl)-3-(3-(4-((methylamino)methyl)phenyl)isoxazol-5-yl)pyrazin-2-amine (Compound I-2)




Step 1: Preparation of Compound 4-ii



[00195] A suspension of *tert*-butyl 4-((hydroxyimino)methyl)benzyl (methyl)carbamate (**Compound 4-i**) (650 g, 2.46 mol) in isopropyl acetate (6.5 L) is stirred at ambient temperature. N-Chlorosuccinimide (361 g, 2.71 mol) is added and the reaction temperature maintained overnight at 20–28 °C to ensure complete reaction. The reaction mixture is diluted with water (3.25 L) and EtOAc (1.3 L) and the phases are separated. The organic


phase is washed with water (2 x 3.25 L), dried (Na₂SO₄), and concentrated to a wet-cake. The concentrate is diluted with heptane (9.1 L), ~2 L of solvent removed, and then stirred at ambient temperature for 2–20 h. The solid is collected by filtration. The filter-cake is washed with heptane (2 x 975 mL) and dried to afford Compound 4-ii (692 g; 94% yield, 99.2 area % purity by HPLC) as a colorless powder.

Step 2: Preparation of *tert*-butyl (5-bromo-3-(3-(4-(((*tert*-butoxycarbonyl)(methyl)amino)methyl)phenyl)isoxazol-5-yl)pyrazin-2-yl)(*tert*-butoxycarbonyl)carbamate (**Compound 4-iv**)



[00196] A suspension of *tert*-butyl *N*-(5-bromo-3-ethynylpyrazin-2-yl)-*N*-*tert*-butoxycarbonylcarbamate (**Compound 4-iii**) (1.59 kg, 3.99 mol) and *tert*-butyl 4-(chloro(hydroxyimino)methyl)benzyl(tetrahydro-2*H*-pyran-4-yl)carbamate (1.31 kg, 4.39 mol; 1.10 equiv.) in CH₂Cl₂ (12.7 L) is stirred at ambient temperature. Triethylamine (444 g, 611 mL, 4.39 mol) is added to the suspension and the reaction temperature is maintained between 20–30 °C for 20–48 h to ensure complete reaction. The reaction mixture is diluted with water (8 L) and thoroughly mixed, then the phases are separated. The organic phase is washed with water (8 L), dried (Na₂SO₄), and then concentrated until about 1 L of CH₂Cl₂ remains. The concentrate is diluted with heptane (3.2 L) and re-concentrated at 40 °C/200 torr until no distillate is observed. The concentrate is stirred and further diluted with heptane (12.7 L) to precipitate a solid. The suspension is stirred overnight. The solid is collected by filtration, washed with heptane (2 x 3 L) then dried to afford **Compound 4-iv** (2.42 kg; 92% yield, 100 area % purity by HPLC) as a light tan powder. ¹H NMR (400 MHz, CDCl₃) δ 8.61 (s, 1H), 7.82 (d, *J* = 8.2 Hz, 2H), 7.31 (m, 3H), 4.46 (br s, 2H), 2.84 (br d, 3H), 1.57 (s, 2H), 1.44 (br s, 9H), 1.36 (s, 18H).

Step 3: Preparation of Compound 5-i




[00197] A mixture of *tert*-butyl (5-bromo-3-(3-(4-(((*tert*-butoxycarbonyl)(methyl)amino)methyl)phenyl)isoxazol-5-yl)pyrazin-2-yl)(*tert*-butoxycarbonyl)carbamate (**Compound 4-iv**) (1.00 kg, 1.51 mol), K₂CO₃ (419 g, 3.02 mol), and (4-(isopropylsulfonyl)phenyl)boronic acid (345 g, 1.51 mol) in toluene (7.0 L) and water (2.0 L) was stirred and degassed with N₂ for 30 min. 1,1'-bis(di-*t*-butylphosphino)ferrocene-dichloro-palladium(II) [Pd(dtbpf)Cl₂; 19.7 g, 30.3 mmol] was then added and degassed an additional 20 min. The reaction mixture was warmed at 70 °C for at least 1 h to ensure complete reaction. The reaction mixture was cooled to ambient temperature then filtered through Celite. The reaction vessel and filter pad are rinsed with toluene (2 x 700 mL). The filtrates are combined and the phases are separated. The organic phase is stirred with Biotage MP-TMT resin (170 g) for 4–20 h. The resin is removed by filtration through Celite and the filter pad is washed with toluene (2 x 700 mL). The filtrate and washings are combined and concentrated to near dryness then diluted with *i*-PrOH (5.75 L) and re-concentrated. The concentrate is again dissolved in warm (45 °C) *i*-PrOH (5.75 L) and then cooled to ambient temperature with stirring to induce crystallization then stirred for around 16 – 20 h. The solid is collected by filtration, washed with *i*-PrOH (2 x 1 L), and dried to afford VRT-1018729 (967 g; 84%) as a beige powder. ¹H NMR (400 MHz, CDCl₃) δ 9.04 (s, 1H), 8.33 (d, *J* = 8.6 Hz, 2H), 8.06 (d, *J* = 8.5 Hz, 2H), 7.85 (d, *J* = 8.1 Hz, 2H), 7.34 (m, 3H), 4.47 (br s, 2H), 3.25 (hept, *J* = 7.0 Hz, 1H), 2.85 (br d, 3H), 1.47 (s, 9H), 1.38 (s, 18H), 1.33 (d, *J* = 6.9 Hz, 6H).

Step 4: Preparation of Compound I-2 • 2HCl

[00198] A solution of **Compound 5-i** (950 g, 1.24 mol) in acetone (12.35 L) is warmed to 40 °C then concentrated HCl (1.23 kg, 1.02 L of 37 %w/w, 12.4 mol) is added at a rate to maintain the reaction temperature between 40 – 45 °C for at least 5 h to ensure complete reaction. The suspension is cooled to below 30 °C and the solid collected by filtration. The

filter-cake is washed with acetone (2 x 950.0 mL) then dried to afford Compound I-2 • 2HCl (578 g; 87% yield, 99.5 area % purity by HPLC) as a yellow powder. ¹H NMR (400 MHz, DMSO) δ 9.53 (br d, *J* = 4.8 Hz, 2H), 8.93 (s, 1H), 8.37 (d, *J* = 8.5 Hz, 2H), 8.07 (d, *J* = 8.3 Hz, 2H), 7.92 (d, *J* = 8.6 Hz, 2H), 7.84 (s, 1H), 7.75 (d, *J* = 8.3 Hz, 2H), 4.23 – 4.15 (m, 2H), 3.43 (hept, *J* = 6.8 Hz, 1H), 2.55 (t, *J* = 5.3 Hz, 3H), 1.17 (d, *J* = 6.8 Hz, 6H).

Step 5: Preparation of Compound I-2 • HCl from Compound I-2 • 2HCl



Two-pot process

[00199] A stirred suspension of Compound I-2 • 2HCl (874 g, 1.63 mol) in *i*-PrOH (3.50 L) and water (0.87 L) is warmed at 50 °C for 1–2 h, cooled to ambient temperature, and stirred for 1–20 h. XRPD is performed on a small sample to ensure that Compound I-2 • 2HCl has been converted to another form. The suspension is cooled to 5 °C and stirred for 1 h. The solid is collected by filtration then the filter-cake is washed with 80/20 *i*-PrOH/water (2 x 874 mL), and briefly dried.


[00200] If XRPD shows the Compound I-2 • HCl/anhydrate form, the solid is dried to afford Compound I-2 • HCl/anhydrate (836 g, 99% yield, 99.2 area % purity by HPLC) as a yellow solid. ¹H NMR (400 MHz, DMSO) δ 9.38 (s, 2H), 8.96 (s, 1H), 8.46 – 8.34 (m, 2H), 8.10 (d, *J* = 8.3 Hz, 2H), 7.94 (d, *J* = 8.6 Hz, 2H), 7.85 (s, 1H), 7.75 (d, *J* = 8.3 Hz, 2H), 7.23 (br s, 2H), 4.21 (s, 2H), 3.47 (hept, *J* = 6.7 Hz, 1H), 2.58 (s, 3H), 1.19 (d, *J* = 6.8 Hz, 6H).

[00201] If XRPD shows the Compound I-2 • HCl/hydrate form the solid is stirred in fresh *i*-PrOH (3.50 L) and water (0.87 L) at 50 °C for at least 2 h until XRPD shows complete conversion to Compound I-2 • HCl/anhydrate. The suspension is then cooled to 5 °C and stirred for 1 h. The solid is collected by filtration then the filter-cake is washed with 80/20 *i*-PrOH/water (2 x 874 mL) then dried to afford Compound I-2•HCl/anhydrate.

Alternative procedure (single pot) used


[00202] Compound I-2 • 2HCl (392 g) is charged to the reactor. 4:1 IPA/water (8 L) is charged to a reactor and stirred at ambient temperature overnight. XRPD is used to confirm the conversion to the mono-HCl salt mono-hydrate form. The mixture is heated to 50 °C. Seeds of Compound I-2 • HCl/anhydrate (16 g) are added and the mixture heated at 50 °C until XRPD confirms complete conversion to the desired anhydrate form. The mixture is cooled to ambient, filtered and the solid washed with 4:1 IPA/water (2 x 800 mL) then dried to afford Compound I-2 • HCl/anhydrate (343 g, 94% yield).

Step 4: Alternate Method 1: Preparation of Compound I-2 free base




[00203] A solution of **Compound 5-i** (100 g, 131 mmol) in DCM (200 mL) was stirred at ambient temperature then TFA (299 g, 202 mL, 2.62 mol) was added. After 2 h reaction solution was cooled to 5 °C. The reaction mixture was diluted with EtOH (1.00 L) over about 5 min resulting in a bright yellow suspension. The suspension was cooled to 10 °C then NaOH (1.64 L of 2.0 M, 3.28 mol) was added over 30 min then stirred at ambient temperature overnight. The solid was collected by filtration then washed with water (2 x 400 mL), EtOH (2 x 200 mL) then dried to afford Compound I-2 free-base (57.0 g, 94% yield, 99.7 area % purity by HPLC) as a fine, yellow powder. ¹H NMR (400 MHz, DMSO) δ 8.95 (s, 1H), 8.39 (d, *J* = 8.5 Hz, 2H), 7.95 (dd, *J* = 11.6, 8.4 Hz, 4H), 7.78 (s, 1H), 7.51 (d, *J* = 8.2 Hz, 2H), 7.21 (br s, 2H), 3.72 (s, 2H), 3.47 (hept, *J* = 6.8 Hz, 1H), 2.29 (s, 3H), 1.19 (d, *J* = 6.8 Hz, 6H).

Step 4: Alternate Method 2: Preparation of Compound I-2 • HCl




Step 1: Preparation of *N*-(4-(diethoxymethyl)benzyl)tetrahydro-2*H*-pyran-4-amine(A-2)



[00206] A solution of tetrahydro-2*H*-pyran-4-amine hydrochloride (1.13 kg, 8.21 mol) in MeOH (14.3 L) is stirred at about 20 °C. then Et_3N (1.06 kg, 1.43 L, 8.21 mol) is added. The mixture is stirred for at least 5 min then terephthalaldehyde diethyl acetal (1.43 kg, 6.84 mol) is added while maintaining the reaction temperature between 20–25 °C. The mixture is stirred for at least 45 min to form the imine. NaBH_4 caplets (414 g, 11.0 mol) are added while maintaining the reaction temperature below about 25 °C. The mixture is stirred for 1 h after the addition is completed. The reaction mixture is quenched by adding 1 M NaOH (13.7 L) then extracted with MTBE. The organic solution was washed with brine (7.13 L) then dried (Na_2SO_4) and concentrated to afford Compound A-2 (2197 g; 109% yield, 94.4 area % purity by HPLC) as a hazy oil. ^1H NMR (400 MHz, CDCl_3) δ 7.43 (d, J = 8.1 Hz, 2H), 7.31 (d, J = 8.1 Hz, 2H), 5.49 (s, 1H), 4.66 (br s, 1H), 4.03 – 3.91 (m, 2H), 3.82 (s, 2H), 3.69 – 3.47 (m, 4H), 3.38 (td, J = 11.6, 2.1 Hz, 2H), 2.78 – 2.65 (m, 1H), 1.90 – 1.81 (m, 2H), 1.53 – 1.37 (m, 2H), 1.23 (t, J = 7.1 Hz, 6H).


Step 2: Preparation of *tert*-butyl 4-(diethoxymethyl)benzyl(tetrahydro-2*H*-pyran-4-yl)carbamate (A-3)



[00207] A mixture of *N*-(4-(diethoxymethyl)benzyl)tetrahydro-2*H*-pyran-4-amine (A-2) (2195 g, 7.48 mol) in CH_2Cl_2 (22.0 L) is stirred at 25 °C then di-*t*-butyl dicarbonate (1.71 kg, 7.86 mol) is added. Et_3N (795 g, 1.10 L) is then added while maintaining the reaction temperature between 20 – 25 °C. The reaction mixture is stirred at about 25 °C for 12 – 20 h. After the reaction is completed, the mixture is cooled to about 20 °C and quenched with 0.5 M aqueous citric acid (7.48 L, 3.74 mol) while maintaining the reaction temperature between


20 – 25 °C. The organic phase is collected, washed with sat. NaHCO₃ (6.51 L, 7.48 mol), washed with brine (6.59 L), and dried (Na₂SO₄) then concentrated to afford *tert*-butyl 4-(diethoxymethyl)benzyl(tetrahydro-2*H*-pyran-4-yl)carbamate (A-3) (2801 g; 95% yield, 98.8 area % purity by HPLC) as a thick, amber oil. ¹H NMR (400 MHz, CDCl₃) δ 7.40 (d, *J* = 8.1 Hz, 2H), 7.21 (d, *J* = 7.9 Hz, 2H), 5.49 (s, 1H), 4.39 (br s, 3H), 3.93 (br dd, *J* = 10.8, 3.8 Hz, 2H), 3.67 – 3.47 (m, 4H), 3.40 (br m, 2H), 1.68 - 1.59 (m, 4H), 1.39 (br s, 9H), 1.23 (t, *J* = 7.1 Hz, 6H).

Step 3: Preparation of *tert*-butyl 4-((hydroxyimino)methyl)benzyl(tetrahydro-2*H*-pyran-4-yl)carbamate (A-4)




[00208] A solution of *tert*-butyl 4-(diethoxymethyl)benzyl(tetrahydro-2*H*-pyran-4-yl)carbamate (A-3) (2.80 kg, 7.12 mol) in THF (28.0 L) and water (2.80 L) is stirred at about 20 °C. Hydroxylamine hydrochloride (593 g, 8.54 mol) is added while maintaining the reaction temperature between 20–25 °C. The reaction mixture is stirred at about 20 °C for 16 - 20 h then diluted with CH₂Cl₂ (8.4 L) and 50% brine (11.2 L) and stirred for at least 5 min. The phases are separated then the organic phase is washed with 50% brine (2 x 2.8 L), dried (Na₂SO₄) and concentrated. The concentrate is diluted with MeOH (1.4 L) and re-concentrated. The concentrate is diluted with MeOH (14.0 L) and transferred to a reaction vessel. The solution is warmed to about 25 °C then water (14.0 L) is added over about 1 – 1.5 h; after about 10 L of water is added, the mixture is seeded and a hazy suspension is observed. Additional water (8.4 L) is added over 1.5 h to further precipitate the product. After aging, the solid is collected by filtration. The filter-cake is washed with heptane (5.6 L) and dried to afford *tert*-butyl 4-((hydroxyimino)methyl)benzyl(tetrahydro-2*H*-pyran-4-yl)carbamate (A-4) (1678 g; 71%, 91.5 area % purity by HPLC) as an off-white powder. ¹H NMR (400 MHz, CDCl₃) δ 8.12 (s, 1H), 7.51 (d, *J* = 8.2 Hz, 2H), 7.24 (d, *J* = 7.9 Hz, 2H), 4.40 (br s, 3H), 3.96 (dd, *J* = 10.4, 3.6 Hz, 2H), 3.41 (br m, 2H), 1.69 - 1.61 (m, 4H), 1.39 (br s, 9H).

Step 4: Preparation of (*tert*-butyl 4-(chloro(hydroxyimino)methyl)benzyl (tetrahydro-2*H*-pyran-4-yl)carbamate (A-4-i)



[00209] A suspension of (*E*)-*tert*-butyl 4-((hydroxyimino)methyl)benzyl(tetrahydro-2*H*-pyran-4-yl)carbamate (A-4) (1662 g, 4.97 mol) in *i*-PrOAc (16.6 L) is stirred at 20 °C in a reactor. *N*-chlorosuccinimide (730 g, 5.47 mol) is added maintaining about 20 °C. The suspension is stirred at about 20 °C to complete the reaction. The suspension is diluted with water (8.3 L) and stirred to dissolve the solid. The phases are separated and the organic phase is washed with water (8.3 L). The organic phase is concentrated then diluted with *i*-PrOAc (831 mL). Heptane (13.3 L; 8 V) is slowly added to induce crystallization. The thick suspension is then stirred for 1 h. The solid is collected by filtration; the filter-cake is washed with heptane (2 x 1.6 L; 2 x 1 V) and dried to afford (*Z*)-*tert*-butyl 4-(chloro(hydroxyimino)methyl)benzyl (tetrahydro-2*H*-pyran-4-yl)carbamate (A-4-i) (1628 g; 89%, 98.0 area % purity by HPLC) as a white powder.


Step 5: Preparation of *tert*-butyl (5-bromo-3-(3-(4-(((*tert*-butoxycarbonyl)(tetrahydro-2*H*-pyran-4-yl)amino)methyl)phenyl)isoxazol-5-yl)pyrazin-2-yl)(*tert*-butoxycarbonyl)carbamate (A-5)



[00210] A solution of *tert*-butyl 4-(chloro(hydroxyimino)methyl)benzyl(tetrahydro-2*H*-pyran-4-yl)carbamate (A-4-i) (1.60 kg, 4.34 mol) and *tert*-butyl *N*-(5-bromo-3-ethynylpyrazin-2-yl)-*N*-*tert*-butoxycarbonylcarbamate (**Compound A-4-ii**) (1.73 kg, 4.34 mol) in CH₂Cl₂ (12.8 L) is stirred at 20 °C. Et₃N (483 g, 665 mL; 4.77 mol) is added and the reaction temperature maintained below 30 °C. The suspension stirred at 20 °C to complete

the reaction then diluted with water (8.0 L) and agitated. The phases are separated and the organic phase is washed with water (8.0 L) and then concentrated. *i*-PrOAc (1.6 L) is added and the mixture and heated at 50 °C. Heptane (4.0 L) was slowly added then the suspension is allowed to cool to ambient temperature and stirred overnight. Additional heptane (7.2 L) is added to the suspension and it is stirred for 1 h. The solid is collected by filtration. The filter-cake is washed with heptane (2 x 1.6 L) and dried to afford *tert*-butyl (5-bromo-3-(3-(4-(((*tert*-butoxycarbonyl)(tetrahydro-2*H*-pyran-4-yl)amino)methyl)phenyl)isoxazol-5-yl)pyrazin-2-yl)(*tert*-butoxycarbonyl)carbamate (A-5) (2.478 kg; 78%, 97.8 area % purity by HPLC) as a fine, tan powder. ¹H NMR (400 MHz, CDCl₃) δ 8.60 (s, 1H), 7.78 (d, *J* = 8.3 Hz, 2H), 7.31 (m, 3H), 4.42 (br m, 3H), 4.03 – 3.82 (m, 2H), 3.38 (br s, 2H), 1.60 (m, 4H), 1.36 (s, 27H).


Step 6: Preparation of *tert*-butyl *tert*-butoxycarbonyl(3-(3-(4-(((*tert*-butoxycarbonyl)(tetrahydro-2*H*-pyran-4-yl)amino)methyl)phenyl)isoxazol-5-yl)-5-(4-(isopropylsulfonyl)phenyl)pyrazin-2-yl)carbamate



[00211] A mixture of *tert*-butyl (5-bromo-3-(3-(4-(((*tert*-butoxycarbonyl)(tetrahydro-2*H*-pyran-4-yl)amino)methyl)phenyl)isoxazol-5-yl)pyrazin-2-yl)(*tert*-butoxycarbonyl)carbamate (A-5) (425 g, 582 mmol), K₂CO₃ (161 g, 1.16 mol; 2.0 equiv.), and 4-(isopropylsulfonyl)phenylboronic acid (133 g, 582 mmol) in toluene (2.98 L) and water (850 mL) is stirred and degassed with N₂ at ambient temperature. The catalyst [1,1'-bis(di-*tert*-butylphosphino)ferrocene]dichloropalladium(II), (Pd(dtbpf)Cl₂; 1.90 g, 2.91 mmol) is added and the mixture is degassed for an additional 10 min. The mixture is heated at 70 °C until the reaction is complete. The mixture is cooled to 50 °C, diluted with water (850 mL) and filtered through a bed of Celite. The phases are separated. The organic phase is concentrated then the residue is diluted with EtOH (1.70 L) and re-concentrated. With mixing at 40 °C, the concentrate is diluted with EtOH (1.70 L) to induce crystallization. The

suspension is cooled to 20 °C and stirred for 4 h. The solid is collected by filtration. The filter-cake is washed with EtOH (2 x 425 mL) and air-dried to afford *tert*-butyl *tert*-butoxycarbonyl(3-(3-(4-(((*tert*-butoxycarbonyl)(tetrahydro-2*H*-pyran-4-yl)amino)methyl)phenyl)isoxazol-5-yl)-5-(4-(isopropylsulfonyl)phenyl)pyrazin-2-yl)carbamate (A-6) as a beige powder. The solid is dissolved in THF (2.13 L) and slurried with Biotage MP-TMT resin (48 g) at ambient temperature. The resin is removed by filtration and the filtrate concentrated to remove most of the THF. The concentrate is diluted with EtOH (970 mL) and re-concentrated to about half the original volume. The concentrate is diluted again with EtOH (970 mL) and mixed for 1 h at 40 °C. The suspension is cooled to ambient temperature and the solid is collected by filtration then dried to afford *tert*-butyl *tert*-butoxycarbonyl(3-(3-(4-(((*tert*-butoxycarbonyl)(tetrahydro-2*H*-pyran-4-yl)amino)methyl)phenyl)isoxazol-5-yl)-5-(4-(isopropylsulfonyl)phenyl)pyrazin-2-yl)carbamate (A-6) (416 g; 86% yield, 99.3 area % purity by HPLC) as a white powder. ¹H NMR (400 MHz, CDCl₃) δ 9.04 (s, 1H), 8.38 – 8.28 (m, 2H), 8.10 – 8.01 (m, 2H), 7.82 (d, *J* = 8.2 Hz, 2H), 7.34 (m, 3H), 4.44 (br s, 2H), 3.94 (dd, *J* = 10.5, 3.5 Hz, 2H), 3.40 (br s, 2H), 3.25 (hept, *J* = 6.8 Hz, 1H), 1.65 (m, 4H), 1.38 (br s, 27H), 1.33 (d, *J* = 6.9 Hz, 6H).

Step 7: Preparation of 5-(4-(isopropylsulfonyl)phenyl)-3-(3-(4-(((tetrahydro-2*H*-pyran-4-yl)amino)methyl)phenyl)isoxazol-5-yl)pyrazin-2-amine (I-3) freebase form



below 20 °C. A 5.0 M aqueous solution of NaOH (1.77 L; 8.85 mol) is added while allowing the reaction temperature to rise from about 14°C to about 42 °C. The suspension is heated at 70 – 75 °C for 6 h while removing distillate. The suspension is allowed to cool to ambient temperature. The solid is collected by filtration and the filter-cake is washed with water (4 x 1.64 L). The filter-cake is washed with EtOH (2 x 820 mL) and dried to afford 5-(4-(isopropylsulfonyl)phenyl)-3-(3-(4-(((tetrahydro-2H-pyran-4-yl)amino)methyl)phenyl)isoxazol-5-yl)pyrazin-2-amine (Compound I-1) (257 g; 98% yield, 99.5 area % purity by HPLC) as a yellow powder.

[00213] ¹H NMR (400 MHz, DMSO) δ 8.94 (s, 1H), 8.44 – 8.33 (m, 2H), 7.94 (t, *J* = 8.2 Hz, 4H), 7.76 (s, 1H), 7.53 (d, *J* = 8.2 Hz, 2H), 7.20 (s, 2H), 3.83 (m, 1H), 3.80 (s, 3H), 3.46 (hept, *J* = 6.8 Hz, 1H), 3.25 (td, *J* = 11.4, 2.1 Hz, 2H), 2.66 – 2.54 (m, 1H), 1.79 (br dd, 2H), 1.36 – 1.22 (m, 2H), 1.19 (d, *J* = 6.8 Hz, 6H). ¹³C NMR (101 MHz, DMSO) δ 167.57, 151.76, 141.07, 137.58, 135.75, 129.16, 128.53, 126.57, 126.41, 125.69, 124.52, 102.13, 65.83, 54.22, 52.60, 49.19, 33.18, 15.20.

Compound Analytical Data


Cmpd No.	LCMS ES +	LCMS (Rt min)	HNMR
I-1	---	---	¹ H NMR (400 MHz, DMSO) δ 9.63 (d, <i>J</i> = 4.7 Hz, 2H), 9.05 (s, 1H), 8.69 (d, <i>J</i> = 5.2 Hz, 1H), 8.21 (s, 1H), 8.16 – 8.03 (m, 3H), 7.84 (t, <i>J</i> = 4.1 Hz, 3H), 7.34 (br s, 2H), 4.40 – 4.18 (m, 2H), 3.94 (dd, <i>J</i> = 11.2, 3.9 Hz, 2H), 3.32 (t, <i>J</i> = 11.2 Hz, 3H), 2.17 – 2.00 (m, 2H), 1.81 (s, 6H), 1.75 (dd, <i>J</i> = 12.1, 4.3 Hz, 2H).
I-2	---	---	¹ H NMR (400 MHz, DMSO) δ 8.94 (s, 1H), 8.44 – 8.33 (m, 2H), 7.94 (t, <i>J</i> = 8.2 Hz, 4H), 7.76 (s, 1H), 7.53 (d, <i>J</i> = 8.2 Hz, 2H), 7.20 (s, 2H), 3.83 (m, 1H), 3.80 (s, 3H), 3.46 (hept, <i>J</i> = 6.8 Hz, 1H), 3.25 (td, <i>J</i> = 11.4, 2.1 Hz, 2H), 2.66 – 2.54 (m, 1H), 1.79 (br dd, 2H), 1.36 – 1.22 (m, 2H), 1.19 (d, <i>J</i> = 6.8 Hz, 6H).
II-1	466.2	0.83	¹ H NMR (500 MHz, DMSO) 9.10 (d, <i>J</i> = 5.8 Hz, 2H), 8.97 (s, 1H), 8.42 - 8.37 (m, 2H), 8.15 - 8.08 (m, 2H), 7.99 - 7.92 (m, 2H), 7.85 (s, 1H), 7.75 - 7.69 (m, 2H), 7.22 (br s, 2H), 3.48 (hept, <i>J</i> = 6.8 Hz, 1H), 2.60 (t, <i>J</i> = 5.4 Hz, 3H), 1.20 (d, <i>J</i> = 6.8 Hz, 6H).
II-2	469.1	0.82	¹ H NMR (500 MHz, DMSO) 9.14 (s, 2H), 8.96 (s, 1H), 8.42 - 8.36 (m, 2H), 8.13 - 8.08 (m, 2H), 7.99 - 7.91 (m, 2H), 7.85 (s, 1H), 7.78 - 7.68 (m, 2H), 7.21 (s, 2H), 3.48 (dq, <i>J</i> = 13.6, 6.7 Hz, 1H), 1.20 (d, <i>J</i> = 6.8 Hz, 6H).

Cmpd No.	LCMS ES +	LCMS (Rt min)	HNMR
II-3	467.2	0.78	¹ H NMR (500 MHz, DMSO) 9.11 (s, 2H), 8.97 (s, 1H), 8.39 (d, J = 8.7 Hz, 2H), 8.11 (d, J = 8.4 Hz, 2H), 7.95 (d, J = 8.7 Hz, 2H), 7.85 (s, 1H), 7.72 (d, J = 8.4 Hz, 2H), 7.23 (s, 2H), 4.25 - 4.19 (m, 2H), 3.47 (tt, J = 14.0, 6.9 Hz, 1H), 1.20 (d, J = 6.8 Hz, 6H).
II-4	471.8	0.83	¹ H NMR (400 MHz, DMSO) δ 2.58 (t, 3H), 4.21 (t, 2H), 5.67 (br s, 2H), 7.74 (d, 2H), 7.85 (s, 1H), 7.94 (d, 2H), 8.10 (d, 2H), 8.38 (d, 2H), 8.96 (s, 1H) and 9.33 (br s, 2H) ppm

INTERMEDIATES

Example 8: Preparation of Oxime 5a

SCHEME BB:



Step 1b


[00214] Add MeOH (28.00 L) and 4-(diethoxymethyl)benzaldehyde (Compound **1b**) (3500 g, 16.81 mol) into a reactor at 20 °C. Add methylamine, 33% in EtOH (1.898 kg, 2.511 L of 33 %w/w, 20.17 mol) maintaining 20-30 °C then stir for 1.5 h to form the imine. Add NaBH_4 (381.7 g, 10.09 mol) caplets maintaining the temperature between 20 - 30 °C. Stir at room temperature for at least 30 min to ensure complete reaction. Add aqueous NaOH (16.81 L of 2.0 M, 33.62 mol) maintaining approximately 20 °C. Add MTBE (17.50 L) and brine (7.0 L), stir for at least 5 min then allow the phases to separate. Extract the aqueous layer with MTBE (7.0 L) then combine the organic phases and wash with brine (3.5 L) then dry (Na_2SO_4) then concentrate to 6 L. The biphasic mixture was transferred to a separatory funnel and the aqueous phase removed. The organic phase was concentrated to afford 1-(4-(diethoxymethyl)phenyl)-N-methylmethanamine (Compound **2b**) (3755 g, 16.82 mol, 100%

yield) as an oil. ^1H NMR (400 MHz, CDCl_3) δ 7.43 (d, J = 8.1 Hz, 2H), 7.31 (d, J = 8.1 Hz, 2H), 5.49 (s, 1H), 3.75 (s, 2H), 3.68 – 3.46 (m, 4H), 2.45 (s, 3H), 1.23 (t, J = 7.1 Hz, 6H).

Steps 2b and 3b

[00215] Add 2-MeTHF (15.00 L) and 1-(4-(diethoxymethyl)phenyl)-N-methylmethanamine (Compound 2b) (3750 g, 16.79 mol) to a reactor at 20 °C. Add a solution of Boc anhydride (3.848 kg, 4.051 L, 17.63 mol) in 2-MeTHF (7.500 L) maintaining approximately 25 °C. Stir for at least 30 min to ensure complete conversion to *tert*-butyl 4-(diethoxymethyl)benzyl(methyl)carbamate (Compound 3b), then add a solution of Na_2SO_4 (1.192 kg, 8.395 mol) in water (11.25 L). Heat to 35 °C then add a solution of hydroxylamine hydrochloride (1.750 kg, 25.18 mol) in water (3.75 L) then stir for at least 6 h to ensure complete reaction. Cool to 20 °C, stop the stirring and remove the aqueous phase. Wash the organic layer with brine (3.75 L), dry (Na_2SO_4), filter and concentrate to about 9 L. Add heptane (15.00 L) and crystalline *tert*-butyl 4-((hydroxyimino)methyl)benzyl(methyl)carbamate (Compound 5a) (1.0g portions every 10 min) until nucleation was evident, then concentrate to afford a solid slurry. Add heptane (3.75 L) then cool to room temp and filter. Wash with heptane (5.625 L) then dry to afford *tert*-butyl 4-((hydroxyimino)methyl)benzyl(methyl)carbamate (Compound 5a) (4023 g, 15.22 mol, 91 % yield, 97.2 area % purity by HPLC) as a colorless solid. ^1H NMR (400 MHz, CDCl_3) δ 8.13 (s, 1H), 7.54 (d, J = 8.1 Hz, 2H), 7.25 (br d, 2H), 4.44 (br s, 2H), 2.83 (br d, 3H), 1.47 (br s, 9H).

Scheme CC: Synthesis of Intermediate A-4-ii



[00216] The compound of formula A-4-ii may be made according to the steps outlined in Scheme C. Sonogashira coupling reactions are known in the art (see e.g., Chem. Rev.

2007, 874-922). In some embodiments, suitable Sonogashira coupling conditions comprise adding 1 equivalent of the compound of formula **C-1**, 1 equivalent of TMS-acetylene, 0.010 equivalents of $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$, 0.015 equivalents of CuI and 1.20 equivalents of NMM in isopropanol. The product can be isolated by adding water to the alcoholic reaction mixture.


[00217] Amine salts of a product maybe formed by dissolving the amine in a common organic solvent and adding an acid. Examples of suitable solvents include chlorinated solvents (e.g., dichloromethane (DCM), dichloroethane (DCE), CH_2Cl_2 , and chloroform), ethers (e.g., THF, 2-MeTHF and dioxane), esters (e.g., EtOAc, IPAC) and other aprotic solvents. Examples of suitable acids include but are not limited to HCl , H_3PO_4 , H_2SO_4 , MSA, and PTSA. In some embodiments, the solvent is IPAC and the acid is PTSA. In some embodiments, the acid addition salt is converted back to the free amine base in the presence of a suitable solvent and a suitable base. Suitable solvents include EtOAc, IPAC, dichloromethane (DCM), dichloroethane (DCE), CH_2Cl_2 , chloroform, 2-MeTHF, and suitable bases include NaOH , NaHCO_3 , Na_2CO_3 , KOH , KHCO_3 , K_2CO_3 , and Cs_2CO_3 . In some embodiments, the suitable solvent is EtOAc and the suitable base is KHCO_3 .

[00218] The amine of Compound C-2 may be protected with various amine protecting groups, such as Boc (*tert*-butoxycarbonyl). Introduction of Boc protecting groups is known in the art (see e.g. Protecting Groups in Organic Synthesis, Greene and Wuts). In some embodiments, suitable conditions involve adding 1.00 equivalents of the amine, 2.10 equivalents of di-*tert*-butyl dicarbonate, and 0.03 equivalents of DMAP in EtOAc.


[00219] Reduction in Pd is achieved by treating with a metal scavenger (silica gel, functionalized resins, charcoal). In some embodiments, suitable conditions involve adding charcoal.

[00220] The TMS (trimethylsilyl) protecting group on Compound C-3 may be removed via conditions known to one of skill in the art. In some embodiments, TMS removal conditions comprise reacting the TMS-protected compound with a suitable base in a suitable solvent. Examples of suitable solvents include chlorinated solvents (e.g., dichloromethane (DCM), dichloroethane (DCE), CH_2Cl_2 , and chloroform), ethers (e.g., THF, 2-MeTHF and dioxane), esters (e.g., EtOAc, IPAC), other aprotic solvents and alcohol solvents (e.g., MeOH, EtOH, *i*PrOH). Examples of suitable bases include but are not limited to (e.g., NaOH , KOH , K_2CO_3 , Na_2CO_3). In certain embodiments, suitable conditions comprise adding 1.00 equivalents of the TMS-protected acetylene, 1.10 equivalents of K_2CO_3 , EtOAc and EtOH. In some emboments, the alcoholic solvent, such as EtOH, is added last in the reaction. In some embodiments the product acetylene is isolated by adding water.


Scheme DD: Example Synthesis of Compound A-4-ii



Example 9: Synthesis of Compound A-4-ii




Step 1: Preparation of 5-bromo-3-((trimethylsilyl)ethynyl)pyrazin-2-amine (Compound C-2)



[00221] Charge isopropanol (8.0 L) to a reactor the stir and sparge with a stream of N₂. Add 3,5-dibromopyrazin-2-amine (Compound C-1) (2000 g, 7.91 moles), Pd(PPh₃)₂Cl₂ (56 g, 0.079 moles), CuI (23 g, 0.119 moles), and NMM (1043 mL, 9.49 moles) to the reactor under a N₂ atmosphere. Adjust the reaction temperature to 25 °C. Purge the reactor with N₂ by doing at least three vacuum/N₂ purge cycles. Charge TMS-acetylene (1.12 L, 7.91 moles) to the reaction mixture and maintain the reaction temperature below 30 °C. When the reaction is complete lower the temperature of the reaction mixture to 15 °C then add water (10 L) and

stir for at least 2 h. The solid is collected by filtration washing the solid with 1:1 IPA/water (2 x 6 L). The filter cake is dried under vacuum then charged to a reactor and dissolved in EtOAc (12.5 L). PTSA hydrate (1.28 kg, 6.72 mol) is charged as a solid to the reactor. The mixture is stirred at ambient temperature for at least 5 h then the solid is collected by filtration, washed with 1:1 heptane/EtOAc (3.5 L) followed by heptane (3.5 L). The filter cake is dried to afford 5-bromo-3-((trimethylsilyl)ethynyl)pyrazin-2-amine (Compound C-2) as a PTSA salt (2356 g, 67% yield, 98.9 area % purity by HPLC). ¹H NMR (400 MHz, DMSO) δ 8.12 (s, 1H), 7.48 (d, *J* = 8.1 Hz, 2H), 7.12 (d, *J* = 8.0 Hz, 2H), 2.29 (s, 3H), 0.26 (s, 9H).

Steps 2 and 3



Step 2: Preparation of *tert*-butyl *N*-*tert*-butoxycarbonyl-*N*-[5-bromo-3-((trimethylsilyl)ethynyl)pyrazin-2-yl]carbamate (Compound C-3)

[00222] A solution of 5-bromo-3-((trimethylsilyl)ethynyl)pyrazin-2-amine PTSA salt (Compound C-2) (2350 g, 5.31 mol) in EtOAc (11.5 L) is stirred with a 20% w/w aq. solution of KHCO₃ (4.5 kg, 1.5 eq.) for at least 30 min. The layers are separated and the organic layer is concentrated then dissolved in EtOAc (7 L) and added to a reactor. DMAP (19.5 g, 0.16 mol) is added followed a solution of Boc₂O (2436 g, 11.16 mol) in EtOAc (3 L) is added slowly. The reaction is stirred for at least 30 min to ensure complete reaction then activated charcoal (Darco G-60, 720 g) and Celite (720 g) are added and stirred for at least 2 h. The mixture is filtered washing the solid pad with EtOAc (2 x 1.8 L). The filtrate is concentrated to afford *tert*-butyl *N*-*tert*-butoxycarbonyl-*N*-[5-bromo-3-((trimethylsilyl)ethynyl)pyrazin-2-yl]carbamate (Compound C-3) that is used directly in the next step.

Step 3: Preparation of *tert*-butyl *N*-(5-bromo-3-ethynylpyrazin-2-yl)-*N*-*tert*-butoxycarbonylcarbamate (Compound A-4-ii)

[00223] K_2CO_3 (811 g, 5.87 mol) is charged to a reactor followed by a solution of Compound C-3 (2300 g, 4.89 mol) dissolved in EtOAc (4.6 L) agitation started. EtOH (9.2 L) is added slowly and the mixture stirred for at least 1 h to ensure that the reaction is complete then water (4.6 L) is added and stirred for at least 2 h. The solid is collected by filtration and washed with 1:1 EtOH/water (4.6 L followed by 2.3 L) followed by EtOH (2.3 L). The filter cake is dried to afford *tert*-butyl *N*-(5-bromo-3-ethynylpyrazin-2-yl)-*N*-*tert*-butoxycarbonylcarbamate (Compound A-4-ii) (1568 g, 78% yield, 97.5 area % by HPLC). ^1H NMR (400 MHz, $CDCl_3$) δ 8.54 (s, 1H), 3.52 (s, 1H), 1.42 (s, 18H).

Solid Forms of Compound I-2

[00224] Compound I-2 has been prepared in various solid forms, including salts and co-solvates. The solid forms of the present invention are useful in the manufacture of medicaments for the treatment of cancer. One embodiment provides use of a solid form described herein for treating cancer. In some embodiments, the cancer is pancreatic cancer or non-small cell lung cancer. Another embodiment provides a pharmaceutical composition comprising a solid form described herein and a pharmaceutically acceptable carrier.

[00225] Applicants describe herein five novel solid forms of Compound I-2. The names and stoichiometry for each of these solid forms are provided in Table S-1 below:

Table S-1

Example	Forms	Stoichiometry
Example 10	Compound I-2 free base	N/A
Example 11	Compound I-2 • HCl	1:1
Example 12	Compound I-2 • 2HCl	1:2
Example 13	Compound I-2 • HCl • H_2O	1:1:1
Example 14	Compound I-2 • HCl • $2H_2O$	1:1:2

[00226] Solid state NMR spectra were acquired on the Bruker-Biospin 400 MHz Advance III wide-bore spectrometer equipped with Bruker-Biospin 4mm HFX probe. Samples were packed into 4mm ZrO_2 rotors (approximately 70mg or less, depending on sample availability). Magic angle spinning (MAS) speed of typically 12.5 kHz was applied. The temperature of the probe head was set to 275K to minimize the effect of frictional heating during spinning. The proton relaxation time was measured using ^1H MAS T_1 saturation recovery relaxation experiment in order to set up proper recycle delay of the ^{13}C cross-polarization (CP) MAS experiment. The recycle delay of ^{13}C CPMAS experiment was

adjusted to be at least 1.2 times longer than the measured ^1H T_1 relaxation time in order to maximize the carbon spectrum signal-to-noise ratio. The CP contact time of ^{13}C CPMAS experiment was set to 2 ms. A CP proton pulse with linear ramp (from 50% to 100%) was employed. The Hartmann-Hahn match was optimized on external reference sample (glycine). Carbon spectra were acquired with SPINAL 64 decoupling with the field strength of approximately 100 kHz. The chemical shift was referenced against external standard of adamantane with its upfield resonance set to 29.5 ppm.

[00227] XRPD data for Examples 13-14 were measured on Bruker D8 Advance System (Asset V014333) equipped with a sealed tube Cu source and a Vantec-1 detector (Bruker AXS, Madison, WI) at room temperature. The X-ray generator was operating at a voltage of 40 kV and a current of 40 mA. The powder sample was placed in a shallow silicon holder. The data were recorded in a reflection scanning mode (locked coupled) over the range of 3° - 40° 2 theta with a step size of 0.0144° and a dwell time of 0.25s (105 s per step). Variable divergence slits were used.

Example 10: Compound I-2 (free base)

Compound I-2 free base can be formed according to the methods described in Example 6, Step 4: Alternate Method 1.

XRPD of Compound I-2 (free base)

[00228] Figure 1a shows the X-ray powder diffractogram of the sample which is characteristic of crystalline drug substance.

[00229] Representative XRPD peaks from Compound I-2 free base:

XRPD Peaks	Angle (2-Theta $\pm$ 0.2)	Intensity %
1	23.8	100.0
*2	14.2	43.9
3	22.5	39.3
*4	25.6	31.1
5	19.3	28.6
6	27.2	27.6
7	17.0	25.4
*8	18.1	25.2
9	17.6	19.6
10	20.2	17.2
11	28.3	15.6
12	20.8	14.5
13	29.9	14.5
14	33.2	14.3
15	30.1	13.5

XRPD Peaks	Angle (2-Theta $\pm$ 0.2)	Intensity %
16	26.8	13.4
*17	22.0	12.3
18	36.5	12.3
19	31.8	12.2
20	34.6	11.5
21	31.1	11.2
22	34.0	11.0
23	30.6	10.9
*24	11.1	10.6
25	13.3	10.6

Thermo Analysis of Compound I-2 free base

[00230] A thermal gravimetric analysis of Compound I-2 free base was performed to determine the percent weight loss as a function of time. The sample was heated from ambient temperature to 350°C at the rate of 10°C/min on TA Instrument TGA Q5000 (Asset V014258). Figure 2a shows the TGA result with a one-step weight loss before evaporation or thermal decomposition. From ambient temperature to 215°C, the weight was ~1.9 %.

Differential Scanning Calorimetry of Compound I-2 free base

[00231] The thermal properties of Compound I-2 free base were measured using the TA Instrument DSC Q2000 (Asset V014259). A Compound I-2 free base sample (1.6900 mg) was weighed in a pre-punched pinhole aluminum hermetic pan and heated from ambient temperature to 350°C at 10°C/min. One endothermic peak is observed at 210°C with its onset temperature at 201°C (Figure 3a). The enthalpy associated with the endothermic peak is 78 J/g.

Solid State NMR of Compound I-2 free base

¹³C CP MAS on Compound I-2 free base

275K; ¹H T1=1.30s

12.5 kHz spinning; ref. adamantane 29.5 ppm

For the full spectrum, see Figure 4a.

Representative Peaks

Peak #	Chem Shift [ppm]	Intensity [rel]
1*	171.0	28.7
2	163.7	21.4
3*	152.1	26.3
4	143.1	57.3
5*	141.2	38.8
6	138.8	30.0
7	132.4	62.1
8	130.9	52.3
9	130.0	70.7
10*	126.6	100.0
11	123.5	34.3
12	101.3	34.1
13	57.6	84.3
14*	38.1	42.6
15*	19.2	48.8
16	18.1	53.3

Crystal Structure of Compound I-2 free base

[00232] The free form of Compound I-2 was prepared from the Compound I-2 HCl salt. 200mg Compound I-2 HCl salt was added to 1mL of 6N NaOH solution. 20mL of dichloromethane was used to extract the free form. The dichloromethane layer was dried over K₂CO₃. The solution was filtered off and 5mL of n-heptane was added to it. Crystals were obtained by slow evaporation of the solution at room temperature over night.

[00233] Most crystals obtained were thin plates. Some prismatic shape crystals were found among them.

[00234] A yellow prismatic crystal with dimensions of 0.2× 0.1×0.1 mm³ was selected, mounted on a MicroMount and centered on a Bruker APEX II diffractometer. Three batches of 40 frames separated in reciprocal space were obtained to provide an orientation matrix and initial cell parameters. Final cell parameters were obtained and refined after data collection was completed based on the full data set.

[00235] A diffraction data set of reciprocal space was obtained to a resolution of 116.96° 2θ angle using 0.5° steps with 10 s exposure for each frame. Data were collected at 100 (2) K temperature with a nitrogen flow cryosystem. Integration of intensities and refinement of cell parameters were accomplished using APEXII software.

CRYSTAL DATA

$C_{24}H_{25}N_5O_3S$
$M_r = 463.55$
Monoclinic, $P2_1/n$
$a = 8.9677 (1) \text{ \AA}$
$b = 10.1871 (1) \text{ \AA}$
$c = 24.5914 (3) \text{ \AA}$
$\beta = 100.213 (1)^\circ$
$V = 2210.95 (4) \text{ \AA}^3$
$Z = 4$

Example 11: Compound I-2 • HCl

[00236] Compound I-2 • HCl can be formed according to the methods described in Example 6, Step 4: Alternate Method 2 and Example 6, Step 5.

XRPD of Compound I-2 • HCl

[00237] Figure 1b shows the X-ray powder diffractogram of the sample which is characteristic of crystalline drug substance.

[00238] Representative XRPD peaks from Compound I-2 • HCl

XRPD Peaks	Angle (2-Theta $\pm$ 0.2)	Intensity %
1	14.4	100.0
*2	13.5	89.1
3	20.9	45.8
4	16.4	41.9
5	23.6	33.8
6	27.2	29.6
*7	28.8	26.0
8	16.8	23.3
9	27.6	21.3
*10	15.0	21.1
*11	18.8	20.8
12	30.5	13.3
*13	15.4	13.0
14	26.9	12.7

Thermo Analysis of Compound I-2 • HCl

[00239] A thermal gravimetric analysis of Compound I-2 • HCl was performed to determine the percent weight loss as a function of time. The sample was heated from ambient temperature to 350°C at the rate of 10°C/min on TA Instrument TGA Q5000 (Asset

V014258). Figure 2b shows the TGA result with a two-step weight loss before evaporation or thermal decomposition. From ambient temperature to 100°C, the weight was ~1.1 %, and from 110°C to 240°C the weight loss is ~0.8%.

Differential Scanning Calorimetry of Compound I-2 • HCl

[00240] The thermal properties of Compound I-2 • HCl were measured using the TA Instrument DSC Q2000 (Asset V014259). A Compound I-2 • HCl sample (3.8110 mg) was weighed in a pre-punched pinhole aluminum hermetic pan and heated from ambient temperature to 350°C at 10°C/min. One endothermic peak is observed at 293°C with its onset temperature at 291°C (Figure 3b). The enthalpy associated with the endothermic peak is 160.3 J/g. The second endothermic peak is around 321°C. Both peaks were coupled with sample evaporation and decomposition.

Solid State NMR of Compound I-2 • HCl

¹⁵ CPMAS on Compound I-2 • HCl
275K; 12.5 kHz spinning; ref. adamantane 29.5 ppm
For the full spectrum, see Figure 4b.

Representative Peaks

Peak #	Chem Shift [ppm]	Intensity [rel]
1*	171.7	47.42
2	161.9	28.72
3*	153.4	28.94
4	144.8	42.57
5	142.9	54.14
6	138.7	44.06
7	136.7	60.06
8*	132.9	100
9	131.2	72.62
10	129.8	73.58
11	127.9	63.71
12	125.4	79.5
13	124.1	34.91
14	100.7	53.52
15	54.5	62.56
16	53.9	61.47
17*	31.8	61.15
18	17.0	74.78
19*	15.7	77.79

Crystal Structure of Compound I-2 • HCl

[00241] 180mg Compound I-2 • HCl was added to a vial with 0.8mL 2-propanol and 0.2 mL water. The sealed vial was kept in an oven at 70°C for two weeks. Diffraction quality crystals were observed.

[00242] A yellow needle shape crystal with dimensions of 0.15× 0.02×0.02 mm³ was selected, mounted on a MicroMount and centered on a Bruker APEX II diffractometer (V011510). Three batches of 40 frames separated in reciprocal space were obtained to provide an orientation matrix and initial cell parameters. Final cell parameters were collected and refined was completed based on the full data set.

[00243] A diffraction data set of reciprocal space was obtained to a resolution of 106° 2 θ angle using 0.5° steps with exposure times 20 s each frame for low angle frames and 60s each frame for high angle frames. Data were collected at room temperature.

[00244] To obtain the data in table 1, dry nitrogen was blown to the crystal at 6 Litre/min speed to keep the ambient moisture out. Data in table 2 was obtained without nitrogen. Integration of intensities and refinement of cell parameters were conducted using the APEXII software. The water occupancy can vary between 0 and 1.

Table 1	Table 2
C ₂₄ H ₂₆ ClN ₅ O ₃ S	C ₂₄ H ₂₈ ClN ₅ O ₄ S
<i>M_r</i> = 500.01	<i>Mr</i> = 518.02
Monoclinic, <i>P</i> 2 ₁ / <i>n</i>	Monoclinic, <i>P</i> 2 ₁ / <i>n</i>
<i>a</i> = 5.3332 (2) Å	<i>a</i> = 5.4324 (5) Å
<i>b</i> = 35.4901 (14) Å	<i>b</i> = 35.483 (4) Å
<i>c</i> = 13.5057 (5) Å	<i>c</i> = 13.3478 (12) Å
β = 100.818 (2)°	β = 100.812 (5)°
<i>V</i> = 2510.87 (17) Å ³	<i>V</i> = 2527.2 (4) Å ³

CHN Elemental Analysis

CHN elemental analysis of Compound I-2 • HCl suggest a mono HCl salt.

Element	C	H	N	Cl
% Theory C ₂₄ H ₂₅ N ₅ O ₃ S.HCl	57.60	5.20	14.00	7.10
% Found	56.52	5.38	13.69	7.18

Example 12: Compound I-2 • 2HCl

[00245] Compound I-2 • 2HCl can be formed according to the methods described in Example 6, Step 4.

[00246] XRPD of Compound I-2 • 2HCl

[00247] The XRPD patterns are acquired at room temperature in reflection mode using a Bruker D8 Discover system (Asset Tag V012842) equipped with a sealed tube source and a Hi-Star area detector (Bruker AXS, Madison, WI). The X-Ray generator is operated at a voltage of 40 kV and a current of 35 mA. The powder sample is placed in a nickel holder. Two frames are registered with an exposure time of 120 s each. The data is subsequently integrated over the range of 4.5°-39° 2 theta with a step size of 0.02° and merged into one continuous pattern.

[00248] Figure 1c shows the X-ray powder diffractogram of the sample which is characteristic of crystalline drug substance.

[00249] Representative XRPD peaks from Compound I-2 • 2HCl

XRPD Peaks	Angle (2-Theta ± 0.2)	Intensity %
1	16.8	100.0
*2	15.1	92.2
3	26.2	91.3
4	27.5	91.3
5	27.2	91.2
6	11.8	89.0
7	29.8	89.0
8	22.8	88.8
9	15.7	88.1
*10	18.5	87.5
11	8.4	87.4
12	12.8	86.6
*13	11.5	86.0
14	14.6	86.0
15	20.1	86.0
*16	13.1	85.9
17	16.0	85.9
18	17.3	85.7
19	23.9	85.7
20	19.2	85.4
*21	5.7	85.2
22	21.3	85.2
23	25.3	84.9
24	28.7	84.7
25	24.3	84.1

Thermo Analysis of Compound I-2 • 2HCl

[00250] A thermal gravimetric analysis of Compound I-2 • 2HCl was performed on the TA Instruments TGA model Q5000. Compound I-2 • 2HCl was placed in a platinum sample pan and heated at 10°C/min to 350°C from room temperature. Figure 2c shows the TGA result, which demonstrates a weight loss of 7.0% from room temperature to 188°C, which is consistent with the loss of 1 equivalent of HCl (6.8%). The onset temperature of degradation/melting is 263°C.

Differential Scanning Calorimetry of Compound I-2 • 2HCl

[00251] A DSC thermogram for Compound I-2 • 2HCl drug substance lot 3 was obtained using TA Instruments DSC Q2000. Compound I-2 • 2HCl was heated at 2°C/min to 275°C from -20°C, and modulated at $\pm 1^\circ\text{C}$ every 60 sec. The DSC thermogram (Figure 3c) reveals an endothermic peak below 200°C, which could corresponds to the loss of 1 equivalent of HCl. Melting/recrystallization occurs between 215-245°C, followed by degradation.

Solid State NMR of Compound I-2 • 2HCl

¹³C CPMAS on Compound I-2 • 2HCl

275K; 1H T1=1.7s

12.5 kHz spinning; ref. adamantane 29.5 ppm

For the full spectrum, see Figure 4c.

Peak #	Chem Shift [ppm]	Intensity [rel]
1*	166.5	32.6
2	160.7	24.7
3	145.3	15.0
4*	137.6	56.0
5*	136.1	100.0
6	134.2	22.7
7	132.4	55.9
8	130.0	54.9
9	127.7	70.7
10	125.9	97.1
11	124.7	59.0
12	123.8	91.4
13	123.2	56.0
14	101.6	37.7
15	56.1	60.3
16	50.7	45.6
17*	34.2	56.8
18	18.4	63.5
19*	16.4	70.32

Crystal Structure of Compound I-2 • 2HCl

[00252] 180mg Compound I-2 • HCl was added to a vial with 0.8mL 2-propanol and 0.2 mL water. The sealed vial was kept in an oven at 70°C for two weeks. Diffraction quality crystals were observed.

[00253] A yellow needle shape crystal with dimensions of 0.15× 0.02×0.02 mm³ was selected, mounted on a MicroMount and centered on a Bruker APEX II diffractometer (V011510). Three batches of 40 frames separated in reciprocal space were obtained to provide an orientation matrix and initial cell parameters. Final cell parameters were collected and refined was completed based on the full data set.

[00254] A diffraction data set of reciprocal space was obtained to a resolution of 106° 2 θ angle using 0.5° steps with exposure times 20 s each frame for low angle frames and 60s each frame for high angle frames. Data were collected at room temperature. Dry nitrogen was blown to the crystal at 6 Litre/min speed to keep the ambient moisture out. Integration of intensities and refinement of cell parameters were conducted using the APEXII software.

CRYSTAL DATA

C ₂₄ H ₂₆ ClN ₅ O ₃ S
$M_r = 500.01$
Monoclinic, $P2_1/n$
$a = 5.3332 (2) \text{ \AA}$
$b = 35.4901 (14) \text{ \AA}$
$c = 13.5057 (5) \text{ \AA}$
$\beta = 100.818 (2)^\circ$
$V = 2510.87 (17) \text{ \AA}^3$

Example 13: Compound I-2 • HCl • H₂O

[00255] Compound I-2 • HCl• H₂O can be formed from Compound I-2 • 2 HCl. (E29244-17) A suspension of Compound I-2 • 2 HCl (10.0 g, 18.6 mmol) in isopropyl alcohol (40 mL) and water (10 mL) is warmed at 50 °C for about 1 h and then cooled to below 10 °C. The solid is collected by filtration. The filter-cake is washed with 80/20 isopropyl alcohol/water (2 x 10 mL) and air-dried to afford Compound I-2 • HCl• H₂O as a yellow powder.

XRPD of Compound I-2 • HCl • H₂O

[00256] The XRPD patterns are acquired at room temperature in reflection mode using a Bruker D8 Discover system (Asset Tag V012842) equipped with a sealed tube source and a Hi-Star area detector (Bruker AXS, Madison, WI). The X-Ray generator is operated at a voltage of 40 kV and a current of 35 mA. The powder sample is placed in a nickel holder. Two frames are registered with an exposure time of 120 s each. The data is subsequently integrated over the range of 4.5°-39° 2 theta with a step size of 0.02° and merged into one continuous pattern.

[00257] Figure 1d shows the X-ray powder diffractogram of the sample which is characteristic of crystalline drug substance.

[00258] Representative XRPD peaks from Compound I-2 • HCl • H₂O

[00259]

XRPD Peaks	Angle (2-Theta ± 0.2)	Intensity %
*1	6.6	100.0
*2	19.5	46.5
3	16.8	37.8
4	22.9	36.0
5	13.9	27.0
6	7.3	23.4
7	13.0	22.7
8	16.5	21.2
*9	24.7	20.9
10	17.7	20.8
11	31.1	19.6
12	15.8	19.3
*13	8.1	18.5
14	17.1	18.4
15	12.7	17.2
16	16.0	17.2
17	14.5	16.5
18	20.6	16.0
19	32.7	15.5
*20	11.2	15.2
21	33.9	11.3

Thermo Analysis of Compound I-2 • HCl • H₂O

[00260] Thermogravimetric analysis (TGA) for Compound I-2 • HCl • H₂O was performed on the TA Instruments TGA model Q5000. Compound I-2 • HCl • H₂O was placed in a platinum sample pan and heated at 10°C/min to 400°C from room temperature. The thermogram (Figure 2d) demonstrates a weight loss of 2.9% from room temperature to

100°C, and a weight loss of 0.6% from 100°C to 222°C, which is consistent with theoretical monohydrate (3.5%).

Differential Scanning Calorimetry of Compound I-2 • HCl • H₂O

[00261] A DSC thermogram for Compound I-2 • HCl • H₂O was obtained using TA Instruments DSC Q2000. Compound I-2 • HCl • H₂O was heated at 2°C/min to 275°C from -20°C, and modulated at ± 1°C every 60 sec. The DSC thermogram (Figure 3d) reveals an endothermic peak below 200°C, which could corresponds to the loss of 1 equivalent of HCl. Melting/recrystallization occurs between 215-245°C, followed by degradation.

Example 14: Compound I-2 • HCl • 2H₂O

[00262] Compound I-2 • HCl• 2H₂O can be formed from Compound I-2 • 2 HCl. (E29244-17) A suspension of Compound I-2 • 2 HCl (10.0 g, 18.6 mmol) in isopropyl alcohol (40 mL) and water (10 mL) is warmed at 50 °C for about 1 h and then cooled to below 10 °C. The solid is collected by filtration. The filter-cake is washed with 80/20 isopropyl alcohol/water (2 x 10 mL) and air-dried to afford Compound I-2 • HCl• 2H₂O as a yellow powder.

XRPD of Compound I-2 • HCl • 2H₂O

[00263] The powder x-ray diffraction measurements were performed using PANalytical's X-pert Pro diffractometer at room temperature with copper radiation (1.54060 °Å). The incident beam optic was comprised of a variable divergence slit to ensure a constant illuminated length on the sample and on the diffracted beam side, a fast linear solid state detector was used with an active length of 2.12 degrees 2 theta measured in a scanning mode. The powder sample was packed on the indented area of a zero background silicon holder and spinning was performed to achieve better statistics. A symmetrical scan was measured from 4 – 40 degrees 2 theta with a step size of 0.017 degrees and a scan step time of 15.5s.

[00264] Figure 1e shows the X-ray powder diffractogram of the sample which is characteristic of crystalline drug substance.

[00265]

Representative XRPD peaks from Compound I-2 • HCl • 2H₂O

XRPD Peaks	Angle (2-Theta ± 0.2)	Intensity %
1	16.2	100.0
2	13.4	82.0
*3	26.6	69.2
4	15.9	66.3
5	15.5	63.8
6	17.1	55.3
7	28.3	55.3
8	7.2	51.8
9	20.7	49.7
10	15.3	46.2
11	20.2	44.6
12	28.0	41.2
13	19.9	40.4
14	17.6	39.1
15	26.3	39.0
*16	7.6	36.1
17	27.3	33.6
18	25.6	33.5
19	18.8	32.2
20	27.0	29.1
21	20.8	28.7
22	22.5	28.0
23	13.0	23.8
*24	6.3	22.6
25	25.2	22.6
26	14.3	22.4
27	19.1	20.8
28	25.1	19.7
29	13.7	19.0
30	14.0	17.4
31	33.0	16.2
*32	23.3	15.7
33	16.6	15.1
34	29.6	14.9
35	29.9	14.8
36	27.6	14.8
37	32.1	13.3
*38	24.6	13.1
39	30.8	11.1

Thermo Analysis of Compound I-2 • HCl • 2H₂O

[00266] The TGA (Thermogravimetric Analysis) thermographs were obtained using a TA instrument TGA Q500 respectively at a scan rate of 10°C/min over a temperature range of 25-300°C. For TGA analysis, samples were placed in an open pan. The thermogram demonstrates a weight loss of ~6 from room temperature to 100°C, which is consistent with theoretical dihydrate (6.7%).

Differential Scanning Calorimetry of Compound I-2 • HCl • 2H₂O

[00267] A DSC (Differential Scanning Calorimetry) thermographs were obtained using a TA instruments DSC Q2000 at a scan rate of 10°C/min over a temperature range of 25-300°C. For DSC analysis, samples were weighed into aluminum hermetic T-zero pans that were sealed and punctured with a single hole. The DSC thermogram reveals dehydration between room temperature and 120°C followed by melting/recrystallization between 170-250°C.

Crystal Structure of Compound I-2 • HCl with water

180mg Compound I-2 • HCl was added to a vial with 0.8mL 2-propanol and 0.2 mL water. The sealed vial was kept in an oven at 70°C for two weeks. Diffraction quality crystals were observed.

[00268] A yellow needle shape crystal with dimensions of 0.15× 0.02×0.02 mm³ was selected, mounted on a MicroMount and centered on a Bruker APEX II diffractometer (V011510). Then a kapton tube with water inside covered the pin. The tube was sealed to make sure the crystal is equilibrated with water for two days before the diffraction experiments. Three batches of 40 frames separated in reciprocal space were obtained to provide an orientation matrix and initial cell parameters. Final cell parameters were collected and refined was completed based on the full data set.

[00269] A diffraction data set of reciprocal space was obtained to a resolution of 106° 2θ angle using 0.5° steps with exposure times 20 s each frame for low angle frames and 60s each frame for high angle frames. Data were collected at room temperature. Integration of intensities and refinement of cell parameters were conducted using the APEXII software.

CRYSTAL DATA

$C_{24}H_{28}ClN_5O_4S$
$M_r = 518.02$
Monoclinic, $P2_1/n$
$a = 5.4324 (5) \text{ \AA}$
$b = 35.483 (4) \text{ \AA}$
$c = 13.3478 (12) \text{ \AA}$
$\beta = 100.812 (5)^\circ$
$V = 2527.2 (4) \text{ \AA}^3$

Example 15: Cellular ATR Inhibition Assay:

[00270] Compounds can be screened for their ability to inhibit intracellular ATR using an immunofluorescence microscopy assay to detect phosphorylation of the ATR substrate histone H2AX in hydroxyurea treated cells. HT29 cells are plated at 14,000 cells per well in 96-well black imaging plates (BD 353219) in McCoy's 5A media (Sigma M8403) supplemented with 10% foetal bovine serum (JRH Biosciences 12003), Penicillin/Streptomycin solution diluted 1:100 (Sigma P7539), and 2mM L-glutamine (Sigma G7513), and allowed to adhere overnight at 37°C in 5% CO₂. Compounds are then added to the cell media from a final concentration of 25µM in 3-fold serial dilutions and the cells are incubated at 37°C in 5% CO₂. After 15min, hydroxyurea (Sigma H8627) is added to a final concentration of 2mM.

[00271] After 45min of treatment with hydroxyurea, the cells are washed in PBS, fixed for 10min in 4% formaldehyde diluted in PBS (Polysciences Inc 18814), washed in 0.2% Tween-20 in PBS (wash buffer), and permeabilised for 10min in 0.5% Triton X-100 in PBS, all at room temperature. The cells are then washed once in wash buffer and blocked for 30min at room temperature in 10% goat serum (Sigma G9023) diluted in wash buffer (block buffer). To detect H2AX phosphorylation levels, the cells are then incubated for 1h at room temperature in primary antibody (mouse monoclonal anti-phosphorylated histone H2AX Ser139 antibody; Upstate 05-636) diluted 1:250 in block buffer. The cells are then washed five times in wash buffer before incubation for 1h at room temperature in the dark in a mixture of secondary antibody (goat anti-mouse Alexa Fluor 488 conjugated antibody; Invitrogen A11029) and Hoechst stain (Invitrogen H3570); diluted 1:500 and 1:5000, respectively, in wash buffer. The cells are then washed five times in wash buffer and finally 100ul PBS is added to each well before imaging.

[00272] Cells are imaged for Alexa Fluor 488 and Hoechst intensity using the BD Pathway 855 Bioimager and Attovision software (BD Biosciences, Version 1.6/855) to quantify phosphorylated H2AX Ser139 and DNA staining, respectively. The percentage of phosphorylated H2AX-positive nuclei in a montage of 9 images at 20x magnification is then calculated for each well using BD Image Data Explorer software (BD Biosciences Version 2.2.15). Phosphorylated H2AX-positive nuclei are defined as Hoechst-positive regions of interest containing Alexa Fluor 488 intensity at 1.75-fold the average Alexa Fluor 488 intensity in cells not treated with hydroxyurea. The percentage of H2AX positive nuclei is finally plotted against concentration for each compound and IC50s for intracellular ATR inhibition are determined using Prism software (GraphPad Prism version 3.0cx for Macintosh, GraphPad Software, San Diego California, USA).

[00273] The compounds described herein can also be tested according to other methods known in the art (see Sarkaria et al, "Inhibition of ATM and ATR Kinase Activities by the Radiosensitizing Agent, Caffeine: *Cancer Research* 59: 4375-5382 (1999); Hickson et al, "Identification and Characterization of a Novel and Specific Inhibitor of the Ataxia-Telangiectasia Mutated Kinase ATM" *Cancer Research* 64: 9152-9159 (2004); Kim et al, "Substrate Specificities and Identification of Putative Substrates of ATM Kinase Family Members" *The Journal of Biological Chemistry*, 274(53): 37538-37543 (1999); and Chiang et al, "Determination of the catalytic activities of mTOR and other members of the phosphoinositide-3-kinase-related kinase family" *Methods Mol. Biol.* 281:125-41 (2004)).

Example 16: ATR Inhibition Assay:

[00274] Compounds can be screened for their ability to inhibit ATR kinase using a radioactive-phosphate incorporation assay. Assays are carried out in a mixture of 50mM Tris/HCl (pH 7.5), 10mM MgCl₂ and 1mM DTT. Final substrate concentrations are 10μM [γ -33P]ATP (3mCi 33P ATP/mmol ATP, Perkin Elmer) and 800 μM target peptide (ASELPASQPQPFSAKKK).

[00275] Assays are carried out at 25°C in the presence of 5 nM full-length ATR. An assay stock buffer solution is prepared containing all of the reagents listed above, with the exception of ATP and the test compound of interest. 13.5 μL of the stock solution is placed in a 96 well plate followed by addition of 2 μL of DMSO stock containing serial dilutions of the test compound (typically starting from a final concentration of 15 μM with 3-fold serial dilutions) in duplicate (final DMSO concentration 7%). The plate is pre-incubated for 10 minutes at 25°C and the reaction initiated by addition of 15 μL [γ -33P]ATP (final

concentration 10 μ M).

[00276] The reaction is stopped after 24 hours by the addition of 30 μ L 0.1M phosphoric acid containing 2mM ATP. A multiscreen phosphocellulose filter 96-well plate (Millipore, Cat no. MAPHN0B50) is pretreated with 100 μ L 0.2M phosphoric acid prior to the addition of 45 μ L of the stopped assay mixture. The plate is washed with 5 x 200 μ L 0.2M phosphoric acid. After drying, 100 μ L Optiphase 'SuperMix' liquid scintillation cocktail (Perkin Elmer) is added to the well prior to scintillation counting (1450 Microbeta Liquid Scintillation Counter, Wallac).

[00277] After removing mean background values for all of the data points, K_i (app) data are calculated from non-linear regression analysis of the initial rate data using the Prism software package (GraphPad Prism version 3.0cx for Macintosh, GraphPad Software, San Diego California, USA).

[00278] In general, the compounds of the present invention are effective for inhibiting ATR. Compounds I-1, I-2, II-1, II-2, II-3 and II-4 inhibit ATR at K_i values below 0.001 μ M.

Example 17: Cisplatin Sensitization Assay

[00279] Compounds can be screened for their ability to sensitize HCT116 colorectal cancer cells to Cisplatin using a 96h cell viability (MTS) assay. HCT116 cells, which possess a defect in ATM signaling to Cisplatin (see, Kim et al.; *Oncogene* 21:3864 (2002); see also, Takemura et al.; *JBC* 281:30814 (2006)) are plated at 470 cells per well in 96-well polystyrene plates (Costar 3596) in 150 μ L of McCoy's 5A media (Sigma M8403) supplemented with 10% foetal bovine serum (JRH Biosciences 12003), Penicillin/Streptomycin solution diluted 1:100 (Sigma P7539), and 2mM L-glutamine (Sigma G7513), and allowed to adhere overnight at 37°C in 5% CO₂. Compounds and Cisplatin are then both added simultaneously to the cell media in 2-fold serial dilutions from a top final concentration of 10 μ M as a full matrix of concentrations in a final cell volume of 200 μ L, and the cells are then incubated at 37°C in 5% CO₂. After 96h, 40 μ L of MTS reagent (Promega G358a) is added to each well and the cells are incubated for 1h at 37°C in 5% CO₂. Finally, absorbance is measured at 490nm using a SpectraMax Plus 384 reader (Molecular Devices) and the concentration of compound required to reduce the IC₅₀ of Cisplatin alone by at least 3-fold (to 1 decimal place) can be reported.

Example 18: Single Agent HCT116 Activity

[00280] Compounds can be screened for single agent activity against HCT116 colorectal cancer cells using a 96h cell viability (MTS) assay. HCT116 are plated at 470 cells per well in 96-well polystyrene plates (Costar 3596) in 150µl of McCoy's 5A media (Sigma M8403) supplemented with 10% foetal bovine serum (JRH Biosciences 12003), Penicillin/Streptomycin solution diluted 1:100 (Sigma P7539), and 2mM L-glutamine (Sigma G7513), and allowed to adhere overnight at 37°C in 5% CO₂. Compounds are then added to the cell media in 2-fold serial dilutions from a top final concentration of 10µM as a full matrix of concentrations in a final cell volume of 200µl, and the cells are then incubated at 37°C in 5% CO₂. After 96h, 40µl of MTS reagent (Promega G358a) is added to each well and the cells are incubated for 1h at 37°C in 5% CO₂. Finally, absorbance is measured at 490nm using a SpectraMax Plus 384 reader (Molecular Devices) and IC₅₀ values can be calculated.

Data for Examples 18-21


Cmpd No.	Single agent HT116 IC ₅₀ (nM)	ATR inhibition Ki (nM)	ATR cellular IC ₅₀ (nM)	Cisplatin sensitization (nM)
II-1	62	<1	18	39
II-2	46	<1	-----	29
II-3	66	0.148	10	39
II-4	-----	0.2	-----	-----

[00281] While we have described a number of embodiments of this invention, it is apparent that our basic examples may be altered to provide other embodiments that utilize the compounds, methods, and processes of this invention. Therefore, it will be appreciated that the scope of this invention is to be defined by the appended claims rather than by the specific embodiments that have been represented by way of example herein.


CLAIMS

We claim:

1. A process for preparing a compound of formula 4:



from a compound of formula 3:



under suitable oxime formation conditions;

wherein

R¹ is C₁₋₆alkyl;

R² is C₁₋₆alkyl;


or R¹ and R², together with the oxygen atoms to which they are attached, form an optionally substituted 5 or 6 membered saturated heterocyclic ring having two oxygen atoms;

R³ is hydrogen, C₁₋₆alkyl, or a 3-6 membered saturated or partially unsaturated heterocyclyl having 1-2 heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur; wherein the heterocyclyl is optionally substituted with 1 occurrence of halo or C₁₋₃alkyl;


J¹ is H, halo, C₁₋₄alkyl, or C₁₋₄alkoxy;

PG is a carbamate protecting group.

2. The process of claim 1, wherein the process is used to make a compound of formula I:



wherein



J^1 is halo, C_{1-4} alkyl, or C_{1-4} alkoxy;


Q is phenyl, pyridyl, or an N-alkylated pyridone;

J^2 is halo; CN; phenyl; oxazolyl; or a C_{1-6} aliphatic group wherein up to 2 methylene units are optionally replaced with O, NR'' , C(O), S, S(O), or S(O)₂; said C_{1-6} aliphatic group is optionally substituted with 1-3 fluoro or CN;

q is 0, 1, or 2; and


R^3 is hydrogen, C_{1-6} alkyl, or a 3-6 membered saturated or partially unsaturated heterocyclyl having 1-2 heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur; wherein the heterocyclyl is optionally substituted with 1 occurrence of halo or C_{1-3} alkyl.

3. The process of claim 1 or claim 2, further comprising the step of protecting a compound of formula **2**:



under suitable protection conditions to form the compound of formula **3**.


4. The process of claim 3, further comprising the step of reacting a compound of formula **1**:




with a suitable ligand under suitable reductive amination conditions to form a compound of formula **2**.


5. The process of claim 4, wherein the suitable ligand is an amine.


6. The process of claim 4, further comprising the step of reacting a compound of formula 4:






under suitable isoxazole formation conditions to form a compound of formula 5:



7. The process of claim 6, further comprising the step of reacting a compound of formula 5 under suitable coupling conditions followed by suitable deprotection conditions to form a compound of formula I.
8. The process of claim 1, comprising one or more of the following limitations:
- R^1 is ethyl and R^2 ethyl;
 - R^3 is CH_3 or ; and
 - J^1 is H.




9. The process of claim 2, wherein R^4 is  wherein Q is phenyl, q is 1, and J^2 is a C_{1-6} aliphatic group wherein up to 1 methylene unit is optionally replaced with $\text{S}(\text{O})_2$.
10. The process of claim 9, J^2 is $-\text{S}(\text{O})_2-(\text{C}_{1-5}\text{alkyl})$.
11. The process of claim 2, wherein
- R^1 is ethyl; R^2 is ethyl;
 - R^3 is CH_3 or ;
 - PG is Boc or Cbz;
 - J^1 is H;





12. The process of claim 2, wherein

R^1 is ethyl; R^2 is ethyl;



PG is Boc;

J^1 is H;



13. The process of claim 12, wherein R^4 is

14. The process of any one of claims 1-13, wherein the suitable oxime formation conditions consist of either a single step sequence or a two step sequence.

15. The process of claim 14, wherein the two step sequence consists of first deprotecting the ketal group in the compound of formula 3 into an aldehyde under suitable deprotection conditions, and then forming the oxime of formula 4 under suitable oxime formation conditions.

16. The process of claim 15, wherein

- a) suitable deprotection conditions comprise adding an acid, acetone, and water; and
- b) suitable oxime formation conditions comprise mixing together hydroxylamine, an acid, an organic solvent, and optionally either water or a dehydrating agent.

17. The process of claim 16, wherein the acid is pTSA or HCl, the organic solvent a chlorinated solvents selected from dichloromethane (DCM), dichloroethane (DCE),


CH₂Cl₂, and chloroform; an ether selected from THF, 2-MeTHF and dioxane; or an aromatic hydrocarbons selected from toluene and xylenes.

18. The process of claim 16, wherein the acid is pTSA or HCl, the dehydrating agent is molecular sieves or dimethoxyacetone, and the organic solvent is an alcoholic solvent selected from methanol or ethanol.
19. The process of claim 14, wherein the single step sequence comprises adding NH₂OH.HCl and a mixture of THF and water.
20. The process of claim 19, wherein 1 equivalent of the compound of formula **3** is combined with a 1.1 equivalents of NH₂OH.HCl in a 10:1 v/v mixture of THF/water.
21. The process of any one of claims 1-13, wherein PG is selected from Boc or Cbz.
22. The process of claim 21, wherein the protection conditions are selected from the group consisting of
 - a) R-OCOCl, a suitable tertiary amine base, and a suitable solvent; wherein R is C₁₋₆alkyl optionally substituted with phenyl;
 - b) R(CO₂)OR', a suitable solvent, and optionally a catalytic amount of base, wherein R is and R' are each independently C₁₋₆alkyl optionally substituted with phenyl;
 - c) [RO(C=O)]₂O, a suitable base, and a suitable solvent.
23. The process of claim 22, wherein the suitable base is Et₃N, diisopropylamine, and pyridine; and the suitable solvent is selected from a chlorinated solvent, an ether, or an aromatic hydrocarbon.
24. The process of claim 23, wherein the suitable base is Et₃N, the suitable solvent is a chlorinated solvent selected from DCM.
25. The process of claim 23, wherein the protection conditions comprise adding 1.20 equivalents of (Boc)₂O and 1.02 equivalents of Et₃N in DCM.

26. The process of any one of claims 7-13, wherein the suitable coupling conditions comprise adding a suitable palladium catalyst with a suitable base in a suitable solvent.
27. The process of claim 26, wherein the suitable palladium catalyst is selected from $\text{Pd}[\text{P}(\text{tBu})_3]_2$, $\text{Pd}(\text{dtbpf})\text{Cl}_2$, $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$, $\text{Pd}(\text{PCy}_3)_2\text{Cl}_2$, $\text{Pd}(\text{dppf})\text{Cl}_2$, and $\text{Pd}(\text{dppe})\text{Cl}_2$; one or more of the following: toluene, MeCN, water, EtOH, IPA, 2-Me-THF, or IPAc; and the suitable base is selected from K_2CO_3 , Na_2CO_3 , or K_3PO_4 .
28. The process of claim 26, wherein the suitable palladium catalyst is $\text{Pd}[\text{P}(\text{tBu})_3]_2$, the suitable solvent is a mixture of acetonitrile and water; and the suitable base is sodium carbonate.
29. The process of claim 26, wherein the suitable coupling conditions comprise adding 0.1 equivalents of $\text{Pd}[\text{P}(\text{tBu})_3]_2$; 1 equivalent of boronic acid or ester; and 2 equivalents of sodium carbonate in a 2:1 ratio v/v of acetonitrile/water at 60-70°C.
30. The process of any one of claims 7-13, wherein the suitable deprotection conditions comprise combining the compound of formula 5 with a suitable Boc deprotecting agent in a suitable solvent.
31. The process of claim 30, wherein the suitable Boc deprotecting agent is selected from TMS-Cl, HCl, TBAF, H_3PO_4 , or TFA and the suitable solvent is selected from acetone, toluene, methanol, ethanol, 1-propanol, isopropanol, CH_2Cl_2 , EtOAc, isopropyl acetate, tetrahydrofuran, 2-methyltetrahydrofuran, dioxane, or diethylether.
32. The process of claim 30, wherein the suitable Boc deprotecting agent is selected from HCl, TBAF, H_3PO_4 , or TFA and the suitable solvent is selected from methanol, ethanol, CH_2Cl_2 , EtOAc, THF, dioxane, toluene, or diethylether.
33. The process of claim 30, wherein the suitable Boc deprotecting agent is HCl or TFA and the suitable solvent is acetone, toluene, isopropanol, or isopropyl acetate.

34. The process of claim 6, wherein the suitable isoxazole-formation conditions consists of two steps, the first step comprising reacting the compound of formula 4 under suitable chlorooxime formation conditions to form a chlorooxime intermediate; the second step comprising reacting the chlorooxime intermediate with an acetylene under suitable cycloaddition conditions to form a compound of formula 5.

35. The process of claim 34, wherein the acetylene is a compound of formula 4-iii.



4-iii.

36. The process of claim 34, wherein the suitable chlorooxime formation conditions are selected from

- d) N-chlorosuccinimide and suitable solvent or
- e) potassium peroxymonosulfate, HCl, and dioxane.

37. The process of claim 36, wherein the suitable solvent is selected from a nonprotic solvent, an aromatic hydrocarbon, or an alkyl acetate.


38. The process of claim 37, wherein the suitable chlorooxime formation conditions are 1.05 equivalents of N-chlorosuccinimide in isopropylacetate at 40-50°C.

39. The process of any one of claims 34-38, wherein suitable cycloaddition conditions consist of a suitable base and a suitable solvent.


40. The process of claim 39, wherein the suitable base is selected from pyridine, DIEA, TEA, t-BuONa, and K₂CO₃ and the suitable solvent is selected from acetonitrile, tetrahydrofuran, 2-methyltetrahydrofuran, MTBE, EtOAc, i-PrOAc, DCM, toluene, DMF, and methanol.

41. The process of claim 40, wherein the suitable base is selected from Et₃N and the suitable solvent is selected from DCM.

42. The process of claim 34, wherein the second step comprises reacting 1 equivalent of acetylene with 1.2 equivalents of the chlorooxime intermediate and 1.3 equivalents of Et₃N in DCM at room temperature.
43. The process of claim 6, wherein the suitable isoxazole-formation conditions comprise combining the compound of formula 4 with an oxidant in a suitable solvent.
44. The process of claim 43, wherein said oxidant is [bis(trifluoroacetoxy)iodo] benzene and said solvent is a 1:1:1 mixture of methanol, water, and dioxane.
45. A process for preparing a compound of formula I-3:




comprising the step of deprotecting a compound of formula A-6:




under suitable Boc deprotection conditions optionally followed by treatment under basic aqueous conditions to form a compound of formula I-3.

46. The process of claim 45, further comprising the step of reacting a compound of formula A-5:



A-5

with a compound of formula A-5-i:




A-5-i

under suitable coupling conditions to form the compound of formula A-6.


47. The process of claim 46, further comprising the steps of

a) reacting a compound of formula A-4:




A-4

under suitable chlorooxime formation conditions to form the compound of formula A-4-i:



A-4-i


b) reacting the compound of formula A-4-i with a compound of formula A-4-ii:



A-4-ii

under suitable cycloaddition conditions to form the compound of formula A-5.

48. The process of claim 47, further comprising the step of reacting a compound of formula A-3:




A-3

under suitable oxime formation conditions to form the compound of formula A-4.


49. The process of claim 48, wherein the suitable oxime formation conditions consist of either a single step sequence or a two step sequence.

50. The process of claim 49, wherein the two step sequence consists of first deprotecting the ketal group in the compound of formula A-3:



A-3


into an aldehyde under suitable deprotection conditions, and then forming the oxime of formula A-4:



A-4

under suitable oxime formation conditions.


51. The process of claim 50, further comprising the step of reacting a compound of formula A-2:



A-2

under suitable Boc protection conditions to form the compound of formula A-3.


52. The process of claim 51, further comprising the step of reacting a compound of formula A-1:



A-1

with tetrahydro-2*H*-pyran-4-amine under suitable reductive amination conditions to form the compound of formula A-2.


53. A process for preparing Compound I-2•HCl :



Compound I-2


comprising stirring a suspension of Compound I-2 • 2HCl in *i*-PrOH and water.

54. A process for preparing Compound I-2•HCl :




comprising combining about 1 equivalent of an aqueous solution of HCl with the free base of Compound I-2 in acetone.

55. A process for preparing a compound of formula I-2:




comprising the step of deprotecting a compound of formula 5-i:




under suitable Boc deprotection conditions optionally followed by treatment under basic aqueous conditions to form a compound of formula I-2.

56. The process of claim 55, further comprising the step of reacting a compound of formula 4-iv:



with




under suitable coupling conditions to form the compound of formula 5-i.

57. The process of claim 46 or 56, wherein the suitable coupling conditions comprise combining a suitable palladium catalyst with a suitable base in a suitable solvent.


58. The process of claim 57 wherein the suitable palladium catalyst is selected from Pd[P(tBu)₃]₂, Pd(dtbpf)Cl₂, Pd(PPh₃)₂Cl₂, Pd(PCy₃)₂Cl₂, Pd(dppf)Cl₂, and Pd(dppe)Cl₂; the suitable solvent is selected from one or more of the following: toluene, MeCN, water, EtOH, IPA, 2-Me-THF, or IPAc; and the suitable base is selected from K₂CO₃, Na₂CO₃, or K₃PO₄.

59. The process of claim 53 or 56, comprising the steps of


a) reacting a compound of formula 4-i



under suitable chlorooxime formation conditions to form the compound of formula 4-ii:




b) reacting the compound of formula 4-ii with a compound of formula 4-iii:



under suitable cycloaddition conditions to form the compound of formula 4-iv.

60. The process of claim 47 or 59, wherein the suitable cycloaddition conditions comprises a suitable base selected from pyridine, DIEA, TEA, t-BuONa, or K₂CO₃ and a suitable solvent selected from acetonitrile, tetrahydrofuran, 2-methyltetrahydrofuran, MTBE, EtOAc, i-PrOAc, DCM, toluene, DMF, and methanol.
61. The process of claim 47 or 59, wherein the suitable chlorooxime formation conditions comprise adding HCl in dioxane to a solution of the oxime in the presence of NCS in a suitable solvent selected from a nonprotic solvents (DCM, DCE, THF, and dioxane), aromatic hydrocarbons (e.g. toluene, xylenes), and alkyl acetates (e.g., isopropyl acetate, ethyl acetate).
62. The process of claim 59, further comprising the step of reacting a compound of formula **3b**:




under suitable oxime formation conditions to form the compound of formula 4-i.

63. The process of claim 62, wherein the suitable oxime formation conditions consist of either a single step sequence or a two step sequence.


64. The process of claim 49 or 63, wherein the single step sequence comprises adding $\text{NH}_2\text{OH}\cdot\text{HCl}$ and a mixture of THF and water.

65. The process of claim 64, wherein 1 equivalent of the compound of formula 3b is combined with a 1.1 equivalents of $\text{NH}_2\text{OH}\cdot\text{HCl}$ in a 10:1 v/v mixture of THF/water.

66. The process of claim 63, wherein the two step sequence consists of first deprotecting the ketal group in the compound of formula 3b:



into an aldehyde under suitable deprotection conditions, and then forming the oxime of formula 5a:



under suitable oxime formation conditions.


67. The process of claim 50 or 66, wherein

- a) suitable deprotection conditions comprise adding an acid, acetone, and water; and
- b) suitable oxime formation conditions comprise mixing together hydroxylamine, an optional acid, an organic solvent, and water.

68. The process of claim 67, wherein the acid is pTSA or HCl, the organic solvent a chlorinated solvents selected from dichloromethane (DCM), dichloroethane (DCE), CH_2Cl_2 , and chloroform; an ether selected from THF, 2-MeTHF and dioxane; or an aromatic hydrocarbons selected from toluene and xylenes. [other aprotic solvents]


69. The process of claim 68, wherein suitable oxime formation conditions comprise adding hydroxylamine hydrochloride to a solution of the compound of 3b in THF and water.

70. The process of any one of claims 62-69 further comprising the step of reacting a compound of formula 2b:



under suitable Boc protection conditions to form the compound of formula 3b.

71. The process of claim 70, further comprising the step of reacting a compound of formula 1b:



with methylamine under suitable reductive amination conditions to form the compound of formula **2b**.

72. The process of claim 52 or 71 wherein the suitable reductive amination conditions comprise adding a reducing agent selected from NaBH₄, NaBH₃CN, or NaBH(OAc)₃ in the presence of a solvent selected from dichloromethane (DCM), dichloroethane (DCE), an alcoholic solvent selected from methanol, ethanol, 1-propanol, isopropanol, or a nonprotic solvent selected from dioxane, tetrahydrofuran, or 2-methyltetrahydrofuran and optionally a base selected from Et₃N or diisopropylethylamine.


73. The process of claim 72, wherein the suitable reductive amination conditions comprise adding 1.2 equivalents of NaBH₄ caplets in the presence Et₃N in MeOH.

74. The process of any one of claim 45-73, wherein the suitable Boc deprotection conditions comprises adding a suitable Boc deprotecting agent selected from TMS-Cl, HCl, TBAF, H₃PO₄, or TFA and the suitable solvent is selected from acetone, toluene,

methanol, ethanol, 1-propanol, isopropanol, CH₂Cl₂, EtOAc, isopropyl acetate, tetrahydrofuran, 2-methyltetrahydrofuran, dioxane, or diethylether.


75. The process of claim 74, wherein the suitable Boc deprotecting agent is HCl or TFA and the suitable solvent is acetone or CH₂Cl₂.

76. A process for preparing a compound of formula A-4-ii:




comprising one or more of the following steps:

f) reacting a compound of formula C-1:




under suitable metal-mediated couplings conditions with TMS-acetylene to form a compound of formula C-2:




;

g) reacting a compound of formula C-2 under suitable Boc protection conditions to form a compound of formula C-3:



h) reacting a compound of formula C-3 suitable TMS deprotection conditions to form a compound of formula A-4-ii.

77. The process of claim 70 or claim 76, wherein the Boc protection conditions are to add $(\text{Boc})_2\text{O}$, a suitable base, and a suitable solvent.
78. The process of claim 77, wherein the suitable base is Et_3N , diisopropylamine, and pyridine; and the suitable solvent is selected from a chlorinated solvent, an ether, or an aromatic hydrocarbon.
79. The process of claim 78, wherein the suitable base is Et_3N , the suitable solvent is DCM, tetrahydrofuran or 2-methyltetrahydrofuran.
80. The process of claim 79, wherein the Boc protection conditions comprise adding 1.05 equivalents of $(\text{Boc})_2\text{O}$ in 2-methyltetrahydrofuran or DCM.
81. The process of claim 76, wherein the suitable metal-mediated coupling conditions are Sonogashira coupling conditions; the Boc protection conditions comprise adding $(\text{Boc})_2\text{O}$ in 2-methyltetrahydrofuran or DCM; and the suitable TMS deprotection conditions comprise adding a suitable base, suitable organic solvent, and water.
82. The process of claim 81, wherein the Sonogashira coupling conditions comprise combining 1 equivalent of TMS-acetylene, 0.010 equivalents of $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$, 0.015 equivalents of CuI and 1.20 equivalents of NMM with Compound C-1 in isopropanol; the Boc protection conditions comprise adding 1.05 equivalents of $(\text{Boc})_2\text{O}$ in 2-methyltetrahydrofuran or DCM; and the TMS deprotection conditions comprise adding K_2CO_3 , EtOAc , EtOH , and water.
83. A compound of formula II:



or a pharmaceutically acceptable salt thereof,


wherein each R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and

R^{10c} is independently hydrogen or deuterium, and


at least one of R^{1a} , R^{1b} , R^{1c} , R^2 , R^{3a} , R^{3b} , R^{3c} , R^4 , R^5 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} , R^{10a} , R^{10b} , and

R^{10c} is deuterium.


84. The compound of claim 81, selected from the group consisting of:




II-1




II-2




II-3




II-4




II-5




II-6




II-7




II-8




II-9




II-10




II-11




II-12




II-13




II-14




II-15




II-16




II-17




II-18




II-19



II-20




II-21



and II-22.

85. The compound of claim 81, selected from the group consisting of compound II-1, II-2, II-3, or II-4.

86. A solid form of a compound of formula I-2:



I-2

wherein the form is selected from the group consisting of Compound I-2 free base, Compound I-2 • HCl, Compound I-2 • HCl • H₂O, Compound I-2 • HCl • 2H₂O, and Compound I-2 • 2-hydrochloric acid.

87. The solid form of claim 86, wherein the form is Compound I-2 free base.
88. The solid form of claim 86, wherein the form is crystalline compound I-2 free base.
89. The solid form of claim 88, having a monoclinic crystal system, having a $P2_1/n$ space group, and having the following unit cell dimensions in Å when measured at 120K:
 $a = 8.9677 (1) \text{ Å}$
 $b = 10.1871 (1) \text{ Å}$
 $c = 24.5914 (3) \text{ Å}$
90. The solid form of claim 88, characterized by a weight loss of from about 1.9 % in a temperature range of from about 25 °C to about 215 °C.
91. The solid form of claim 88, characterized by one or more peaks expressed in $2\text{-}\theta \pm 0.2$ at about 14.2, 25.6, 18.1, 22.0, and 11.1 degrees in a X-ray powder diffraction pattern obtained using Cu K alpha radiation.
92. The solid form of claim 88, characterized as having an X-ray powder diffraction pattern substantially the same as that shown in Figure 1a.
93. The solid form of claim 86, wherein the form is Compound I-2 • HCl.
94. The solid form of claim 86, wherein the form is crystalline Compound I-2 • HCl.
95. The solid form of of claim 94, wherein the compound:HCl are in a ratio of 1:1.
96. The solid form of claim 94, having a monoclinic crystal system, having a $P2_1/n$ space group, and having the following unit cell dimensions in Å when measured at 120K:
 $a = 5.3332 (2) \text{ Å}$
 $b = 35.4901 (14) \text{ Å}$
 $c = 13.5057 (5) \text{ Å}$

97. The solid form of claim 94, characterized by a weight loss of from about 1.1 % in a temperature range of from about 25 °C to about 100 °C, and about 0.8% from 110 °C to about 240 °C.
98. The solid form of claim 94, characterized by one or more peaks expressed in 2-theta $\pm$ 0.2 at about 13.5, 28.8, 15.0, 18.8, and 15.4 degrees in a X-ray powder diffraction pattern obtained using Cu K alpha radiation.
99. The solid form of claim 94, characterized by one or more peaks at about 171.7, 153.4, 132.9, 31.8, and 15.7 ppm in solid state ^{13}C NMR.
100. The solid form of claim 94, characterized as having an X-ray powder diffraction pattern substantially the same as that shown in Figure 1b.
101. The solid form of claim 86, wherein the form is Compound I-2•2-hydrochloric acid.
102. The solid form of claim 86, wherein the form is Crystalline Compound I-2• 2HCl.
103. The solid form of claim 102, wherein the compound:HCl are in a ratio of 1:2.
104. The solid form of claim 102, characterized by one or more peaks at about 15.1, 18.5, 11.5, 13.1, and 5.7 in an X-ray powder diffraction pattern.
105. The solid form of claim 102, characterized as having an X-ray powder diffraction pattern substantially the same as that shown in Figure 1c.
106. The solid form of claim 102, having a monoclinic crystal system, having a $P2_1/n$ space group, and having the following unit cell dimensions in Å when measured at 120K:
- $a = 5.3332 (2) \text{ Å}$
 $b = 35.4901 (14) \text{ Å}$
 $c = 13.5057 (5) \text{ Å}$
107. The solid form of claim 102, characterized by a weight loss of from about 7% in a temperature range of from about 25 °C to about 188 °C.
108. The solid form of claim 102, characterized by one or more peaks at about 15.1, 18.5, 11.5, 13.1, and 5.7 degrees in a X-ray powder diffraction pattern obtained using Cu K alpha radiation.
109. The solid form of claim 102, characterized as having an X-ray powder diffraction pattern substantially the same as that shown in Figure 1c.
110. The solid form of claim 102, characterized by one or more peaks at about 166.5, 137.6, 136.1, 34.2, and 16.4 ppm in solid state ^{13}C NMR.
111. The solid form of claim 86, wherein the form is Compound I-2• 2HCl • H₂O.

112. The solid form of claim 86, wherein the form is Crystalline compound I-2 • 2HCl • H₂O.
113. The solid form of claim 112, wherein the compound:HCl:H₂O are in a ratio of 1:2:1.
114. The solid form of claim 112 characterized by a weight loss of from about 2.9% in a temperature range of from about 25 °C to about 100 °C and a weight loss of about 0.6% from about 100 °C to about 222 °C.
115. The solid form of claim 112 characterized by one or more peaks at about 6.6, 19.5, 24.7, 8.1, and 11.2 degrees in a X-ray powder diffraction pattern obtained using Cu K alpha radiation.
116. The solid form of claim 112, characterized as having an X-ray powder diffraction pattern substantially the same as that shown in Figure 1d.
117. The solid form of claim 86, wherein the form is Compound I-2 • HCl • 2H₂O.
118. The solid form of claim 86, wherein the form is Crystalline compound I-2 • HCl • 2H₂O.
119. The solid form of claim 118 wherein the compound:HCl:H₂O are in a ratio of 1:1:2.
120. The solid form of claim 118 characterized by a weight loss of from about 6% in a temperature range of from about 25 °C to about 100 °C
121. The solid form of claim 118 characterized by one or more peaks at about 26.6, 7.6, 6.3, 23.3, and 24.6 degrees in a X-ray powder diffraction pattern obtained using Cu K alpha radiation.
122. A pharmaceutical composition comprising a solid form of claim 86 and a pharmaceutically acceptable carrier.
123. Use of a solid form of claim 86 for the manufacture of a medicament for treating cancer.
124. Use of a solid form of claim 86 for treating cancer.
125. The use according to claim 123 or claim 124, wherein the cancer is pancreatic cancer or non small cell lung cancer.