CLAIMS

1. A wet granulated formulation comprising water, (6-(5-fluoro-2-(3,4,5-trimethoxyphenylamino) pyrimidin-4-ylamino)-2,2-dimethyl-3-oxo-2H-pyrido[3,2-b][1,4]oxazin-4(3H)-yl)methyl phosphate disodium salt (Compound 1) and starch, wherein the wet granulated formulation comprises 10 to 50 weight percent of Compound 1, 100 to 140 weight percent of a water sequestering agent based on the amount of Compound 1, and between 90 to 120 weight percent of water based on the total weight of the dry formulation prior to wet granulation, and wherein when the wet granulated formulation is dried it has a bulk density between 0.35 g/mL and 0.65 g/mL.

2. The formulation of Claim 1, wherein the starch is partially pregelatinized.

3. The formulation of Claim 1, wherein the starch is derived from Maize.

4. The formulation of Claim 1 which further comprises at least one of a filler, a lubricant, a suspending/dispersing agent, a binding agent, and a disintegrant.

5. A tablet comprising the wet granulated formulation of Claim 1.

6. The tablet of Claim 5, wherein said tablet exhibits at least 75% dissolution in less than 45 minutes in an aqueous solution maintained at pH 7.4, a temperature of 37 °C +0.5 °C, and a paddle speed of 75 rpm.

7. The tablet of Claim 5 which further comprises at least one of a filler, a lubricant, a suspending/dispersing agent, a binding agent, and a disintegrant.

8. The tablet of any of Claims 2, 3, or 4 wherein the tablet comprises from greater than 25 mg to 200 mg of (6-(5-fluoro-2-(3,4,5-trimethoxyphenylamino) pyrimidin-4-ylamino)-2,2-dimethyl-3-oxo-2H-pyrido[3,2-b][1,4]oxazin-4(3H)-yl)methyl phosphate disodium salt.

9. The tablet of Claim 5, wherein the tablet comprises 50 mg to 100 mg of (6-(5-fluoro-2-(3,4,5-trimethoxyphenylamino) pyrimidin-4-ylamino)-2,2-dimethyl-3-oxo-2H-pyrido[3,2-b][1,4]oxazin-4(3H)-yl)methyl phosphate disodium salt.

10. The tablet of Claim 6, wherein the tablet comprises 100 mg of (6-(5-fluoro-2-(3,4,5-trimethoxyphenylamino) pyrimidin-4-ylamino)-2,2-dimethyl-3-oxo-2H-pyrido[3,2-b][1,4]oxazin-4(3H)-yl)methyl phosphate disodium salt.

11. A method for formulating (6-(5-fluoro-2-(3,4,5-trimethoxyphenylamino) pyrimidin-4-ylamino)-2,2-dimethyl-3-oxo-2H-pyrido[3,2-b][1,4]oxazin-4(3H)-yl)methyl phosphate disodium salt into a formulation for tablet compression which method comprises:

    a) blending (6-(5-fluoro-2-(3,4,5-trimethoxyphenylamino) pyrimidin-4-ylamino)-2,2-dimethyl-3-oxo-2H-pyrido[3,2-b][1,4]oxazin-4(3H)-yl)methyl phosphate disodium salt with starch and optionally at least one of a filler, a lubricant, a suspending/dispersing agent, a binding agent and a disintegrant, at an impellor speed sufficient to form a homogenous powder mixture having a bulk density, after drying, of between 0.35 to 0.65 g/mL;

    b) spraying between 15 % and 40 % by weight of water with respect to the total weight of the powder mixture plus water and mixing to form wet granules; and

    c) drying the wet granules until a loss on drying of between 5% and 11% is achieved, to provide dried granules.

12. The method of Claim 11, wherein the dried granules are sized between 25 μm to 900 μm in diameter.

13. The method of Claim 11 which further comprises milling the dried granules, to provide milled granules.

14. The method of Claim 10, wherein the dried granules are milled so that 90 weight percent of the milled granules have a particle size between 25 to 900 μm in diameter.

15. The method of Claim 11, further comprising mixing the milled granules with a lubricant until homogenous, and then tabletting the resulting formulation.

16. A wet granulating process, comprising:

a) blending a composition comprising (6-(5-fluoro-2-(3,4,5-trimethoxyphenylamino) pyrimidin-4-ylamino)-2,2-dimethyl-3-oxo-2H-pyrido[3,2-b][1,4]oxazin-4(3H)-yl)methyl phosphate disodium salt and starch to form a blended mixture;

b) granulating the blended mixture of a) while adding water to form wet granules;

c) drying the wet granules of b) at below 65 °C until a loss on drying of between 5% and 11% is achieved to provide dried granules; and

d) blending a lubricant into the dried granules of c) to provide blended granules,

wherein the blended granules comprise about 10 to about 50 weight percent of Compound 1, about 100 to about 140 weight percent of a water sequestering agent based on the amount of Compound 1, and between about 90 to about 120 weight percent of water based on the total weight of the dry formulation prior to wet granulation.

17. The method of claim 13, further comprising:

e) compressing the blended granules to form tablets.