

DESCRIPTION

TITLE OF THE INVENTION: TRIAZINONE COMPOUND AND T-TYPE CALCIUM CHANNEL INHIBITOR

5

TECHNICAL FIELD

[0001] The present invention relates to novel triazinone compounds having an inhibitory activity on a T-type voltage-dependent calcium channel.

10

BACKGROUND ART

[0002] Voltage-dependent calcium channels are transmembrane multisubunit proteins that control inflow of extracellular calcium ions into cells. The voltage-dependent calcium channels are further classified into various types in mammalian cells. Major types of the voltage-dependent calcium channels include L-type, T-type, N-type, P/Q type, and R-type calcium channels, which play respective roles in various tissues including skeletal muscles, cardiac muscles, lungs, smooth muscles, and brain. Among these types, the "T-type" (or "low-voltage activated-type") calcium channel is named after its characteristic that has a shorter (T = transient) opening time than the L-type calcium channel that has a longer (L = long-lasting) opening time [Non-Patent Document 1].

[0003] The T-type calcium channels have channel characteristics, which are known to be a factor to open the L-type calcium channels and a factor to fluctuate the action potential of sodium channels. Here, hyperexcitability of nerves due to an abnormality (abnormal firing) in fluctuations of the action potential of the sodium channels is believed to be a pathogenesis of neuropathic pains. The T-type calcium channels are supposed to relate to the abnormal firing, and blocking of the T-type calcium channels are believed to suppress the abnormal firing itself and to suppress pains

[Non-Patent Document 2].

More specifically, the T-type calcium channels identified in various mammals including humans include three subtypes, $\alpha 1G$ (Cav3.1), $\alpha 1H$ (Cav3.2), and $\alpha 1I$ (Cav3.3). Among the three subtypes of the T-type calcium channels, $\alpha 1H$ is expressed
5 in the dorsal root ganglion (DRG) and the dorsal horn of the spinal cord, which relate to pain transmission [Non-Patent Document 2, Non-Patent Document 12]. In studies using $\alpha 1H$ knockout mice, analgesic action has been disclosed in acute pain models (tail clip, tail flip, and hot plate tests), inflammatory pain models (capsaicin and formalin-induced tests), and visceral pain models (acetic acid and magnesium sulfate inductions). During
10 the tests, no abnormality was observed in general behavior [Non-Patent Document 3].

Analgesic action has also been identified in neuropathic pain model rats (CCD) to which an antisense gene of $\alpha 1H$ is administered to suppress the expression of $\alpha 1H$ in the spinal cord [Non-Patent Document 4]. In addition, in the case of suppressing the expression in the DRG, analgesic action has been identified in neuropathic pain model
15 rats (CCI) [Non-Patent Document 5].

As for the action on pains associated with diabetic neuropathy, in the DRG of pain model rats having diabetic neuropathy prepared by administration of streptozotocin, an increase in gene expression of $\alpha 1H$ [Non-Patent Document 6] and an increase in T-type calcium channel current [Non-Patent Document 7] have been disclosed, and the pain
20 suppressive action has also been identified by the intrathecal administration of an antisense gene of $\alpha 1H$ to the pain model rats [Non-Patent Document 8]. It has been disclosed that the onset of pain has been completely suppressed in $\alpha 1H$ knockout mice to which streptozotocin has been administered, and the expression of $\alpha 1H$ in the DRG has increased and the administration of a T-type calcium channel inhibitor has provided an
25 analgesic action in ob/ob mice as diabetic model mice [Non-Patent Document 9]. From these findings, compounds having the inhibitory activity on the T-type calcium channel should be used as a therapeutic agent for pain.

[0004] The T-type calcium channels is considered to relate to pains such as neuropathic pain, inflammatory pain, and cancer pain, as well as pathology of various

diseases and disorders including epilepsy, essential tremor, schizophrenia, Parkinson's disease, depression, anxiety, sleep disorder, sleep disturbance, mental illness, schizophrenia, cardiac arrhythmia, hypertension, pain, cancer, diabetes, overactive bladder, chronic kidney disease, sterility, and sexual dysfunction [Non-Patent Document 2, Non-Patent Document 3, Non-Patent Document 10, Non-Patent Document 11, Non-Patent Document 13, Non-Patent Document 14].

[0005] Treatment methods for such diseases involve many problems, and thus there is a demand for novel pharmaceutical products. Although some compounds having the T-type calcium channel inhibitory activity have been disclosed (for example, see Patent Documents 1 to 3), there is a demand for development of new medicinal agents.

Prior Art Documents

Patent Documents

[0006] Patent Document 1: International Publication WO 2009/146540
 Patent Document 2: International Publication WO 2011/115813
 Patent Document 3: International Publication WO 2011/035159

Non-Patent Documents

[0007] Non-Patent Document 1: Physiology of Neuron, Kyoto University Press (2009)
 Non-Patent Document 2: British Journal of Pharmacology, Vol. 163, pp. 484-495 (2011)
 Non-Patent Document 3: Genes, Brain and Behavior, Vol. 6, pp. 425-431 (2007)
 Non-Patent Document 4: Acta Pharmacologica Sinica, Vol. 27 (No. 12), pp. 1547-1552 (2006)
 Non-Patent Document 5: The EMBO Journal, Vol. 24, pp. 315-324 (2005)
 Non-Patent Document 6: Journal of Neurochemistry, Vol. 119 (No. 3), pp. 594-603 (2011)
 Non-Patent Document 7: Journal of Neuroscience, Vol. 28 (No. 12), pp. 3305-3316 (2007)

Non-Patent Document 8: Pain, Vol. 145 (No. 1-2), pp. 184-195 (2009)

Non-Patent Document 9: Diabetes, Vol. 58, pp. 2656-2665 (2009)

Non-Patent Document 10: The Journal of Pharmacology and Experimental Therapeutics, Vol. 335, No. 2, pp. 409-417 (2010)

5 Non-Patent Document 11: Journal of Assisted Reproduction and Genetics, Vol. 28, No. 1, pp. 23-30 (2011)

Non-Patent Document 12: Pharmacological Reviews, Vol. 83, pp. 117-161 (2003)

Non-Patent Document 13: Neurourology and Urodynamics, Vol. 26, pp. 870-878 (2007)

10 Non-Patent Document 14: BJU International, Vol. 99 (No. 2), pp. 436-441 (2007)

SUMMARY OF THE INVENTION

Problem to be Solved by the Invention

15 [0008] The present invention provides novel triazinone compounds which are T-type voltage-dependent calcium channel inhibitors. The present invention also provides pharmaceutical compositions containing the compounds of the present invention.


20 Means for Solving the Problem

[0009] As a result of intensive studies for developing inhibitors of a T-type voltage-dependent calcium channel, the inventors of the present invention have found that the compounds of the present invention have a high inhibitory activity on the T-type voltage-dependent calcium channel and have accomplished the present invention.

25 Specifically, the present invention has the following aspects:

[0010] (1)

The present invention provides: a compound of Formula (I):



[wherein

R^1 is

a hydrogen atom,

5 a halogen atom,

a C_{1-6} alkyl group,

a C_{1-6} alkoxy group,

a C_{1-6} alkylthio group,

a mono- C_{1-6} alkylamino group,

10 a di- C_{1-6} alkylamino group

(the C_{1-6} alkyl group, the C_{1-6} alkoxy group, the C_{1-6} alkylthio group, the mono- C_{1-6}

alkylamino group, and the di- C_{1-6} alkylamino group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^8),

or

15 a C_{3-11} cycloalkyl group

(the C_{3-11} cycloalkyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^6);

each of L^1 and L^2 is independently

a single bond,

20 NR^2 ,

O,

S,

SO ,

SO_2 , or

25 a C_{1-6} alkylene group

(the C_{1-6} alkylene group is unsubstituted or substituted with one or more substituents

identically or differently selected from the substituent group V^6 , and a single methylene

group of the C₁₋₆ alkylene group is optionally replaced by O, S, SO₂, C=O, C=S, or NR³);

B is

- a C₃₋₁₁ cycloalkylene group,
 - a C₃₋₁₁ cycloalkenylene group,
 - 5 a 3 to 11-membered heterocyclylene group,
 - a C₆₋₁₄ arylene group,
 - a 5 to 10-membered heteroarylene group
- (the C₃₋₁₁ cycloalkylene group, the C₃₋₁₁ cycloalkenylene group, the 3 to 11-membered heterocyclylene group, the C₆₋₁₄ arylene group, and the 5 to 10-membered heteroarylene
- 10 group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁶, and
- a single methylene group of the C₃₋₁₁ cycloalkylene group or the C₃₋₁₁ cycloalkenylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group),
- a C₁₋₆ alkylene group,
 - 15 a C₂₋₆ alkenylene group, or
 - a C₂₋₆ alkynylene group
- (the C₁₋₆ alkylene group, the C₂₋₆ alkenylene group, and the C₂₋₆ alkynylene group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁸, and
- 20 a single methylene group of the C₁₋₆ alkylene group, the C₂₋₆ alkenylene group, or the C₂₋₆ alkynylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group);
- when L¹ is a single bond, O, or a C₁₋₆ alkylene group and L² is a single bond or a C₁₋₆ alkylene group, B is not a C₁₋₆ alkylene group, a C₂₋₆ alkenylene group, a C₂₋₆ alkynylene group, or a C₆₋₁₄ arylene group;
- 25 when L¹ is a single bond, O, or a C₁₋₆ alkylene group and L² is NR², O, S, SO, or SO₂, B is not a C₁₋₆ alkylene group, a C₂₋₆ alkenylene group, or a C₂₋₆ alkynylene group;

A is

- a C₁₋₆ alkyl group,
- a C₂₋₆ alkenyl group

(the C₁₋₆ alkyl group and the C₂₋₆ alkenyl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁸), a C₃₋₁₁ cycloalkyl group,

a C₃₋₁₁ cycloalkenyl group,

5 a 3 to 11-membered heterocyclyl group,

a C₆₋₁₄ aryl group, or

a 5 to 10-membered heteroaryl group

(the C₃₋₁₁ cycloalkyl group, the C₃₋₁₁ cycloalkenyl group, the 3 to 11-membered heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are

10 unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁶);

L³ is

a C₁₋₆ alkylene group

(the C₁₋₆ alkylene group is unsubstituted or substituted with one or more substituents

15 identically or differently selected from the substituent group V⁸, and a single methylene group of the C₁₋₆ alkylene group is optionally replaced by C=O or C=S);

D is

a C₃₋₁₁ cycloalkyl group,

a C₃₋₁₁ cycloalkenyl group,

20 a 3 to 11-membered heterocyclyl group,

a C₆₋₁₄ aryl group, or

a 5 to 10-membered heteroaryl group

(the C₃₋₁₁ cycloalkyl group, the C₃₋₁₁ cycloalkenyl group, the 3 to 11-membered heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are

25 unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁶);

each of R² and R³ is independently

a hydrogen atom,

a C₁₋₆ alkyl group,

a C₂₋₆ alkenyl group

(the C₁₋₆ alkyl group and the C₂₋₆ alkenyl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁸),

a C₃₋₁₁ cycloalkyl group,

5 a 3 to 11-membered heterocyclyl group,

a C₆₋₁₄ aryl group, or

a 5 to 10-membered heteroaryl group

(the C₃₋₁₁ cycloalkyl group, the 3 to 11-membered heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with one or

10 more substituents identically or differently selected from the substituent group V⁶);

the substituent group V⁶ is a substituent group consisting of substituents constituting the substituent group V⁸, C₁₋₆ alkyl groups, C₂₋₆ alkenyl groups, and C₂₋₆ alkynyl groups (the C₁₋₆ alkyl groups, the C₂₋₆ alkenyl groups, and the C₂₋₆ alkynyl groups are unsubstituted or substituted with one or more substituents identically or differently

15 selected from the substituent group V¹);

the substituent group V⁸ is a substituent group consisting of substituents constituting the substituent group V^a, C₁₋₆ alkoxy groups, C₁₋₆ alkylthio groups, C₁₋₆ alkylcarbonyl groups, C₁₋₆ alkylsulfonyl groups, C₁₋₆ alkoxycarbonyl groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, mono-C₁₋₆ alkylaminocarbonyl groups, di-C₁₋₆

20 alkylaminocarbonyl groups, mono-C₁₋₆ alkylaminosulfonyl groups, di-C₁₋₆

alkylaminosulfonyl groups, C₁₋₆ alkylcarbonylamino groups, C₁₋₆ alkylcarbonyloxy groups, C₁₋₆ alkylsulfonylamino groups, C₁₋₆ alkylsulfonyloxy groups (the C₁₋₆ alkoxy groups, the C₁₋₆ alkylthio groups, the C₁₋₆ alkylcarbonyl groups, the C₁₋₆ alkylsulfonyl groups, the C₁₋₆ alkoxycarbonyl groups, the mono-C₁₋₆ alkylamino groups, the di-C₁₋₆

25 alkylamino groups, the mono-C₁₋₆ alkylaminocarbonyl groups, the di-C₁₋₆

alkylaminocarbonyl groups, the mono-C₁₋₆ alkylaminosulfonyl groups, the di-C₁₋₆

alkylaminosulfonyl groups, the C₁₋₆ alkylcarbonylamino groups, the C₁₋₆

alkylcarbonyloxy groups, the C₁₋₆ alkylsulfonylamino groups, and the C₁₋₆

alkylsulfonyloxy groups are unsubstituted or substituted with one or more substituents

identically or differently selected from the substituent group V^1), C_{3-6} cycloalkoxy groups, mono- C_{3-6} cycloalkylamino groups, di- C_{3-6} cycloalkylamino groups, C_{3-6} cycloalkylcarbonyl groups, C_{3-6} cycloalkylsulfonyl groups, C_{3-6} cycloalkylsulfonylamino groups, C_{3-6} cycloalkylsulfonyloxy groups, C_{3-6} cycloalkylthio groups, C_{3-11} cycloalkyl groups, 3 to 11-membered heterocyclyl groups, C_{6-14} aryl groups, and 5 to 10-membered heteroaryl groups (the C_{3-6} cycloalkoxy groups, the mono- C_{3-6} cycloalkylamino groups, the di- C_{3-6} cycloalkylamino groups, the C_{3-6} cycloalkylcarbonyl groups, the C_{3-6} cycloalkylsulfonyl groups, the C_{3-6} cycloalkylsulfonylamino groups, the C_{3-6} cycloalkylsulfonyloxy groups, the C_{3-6} cycloalkylthio groups, the C_{3-11} cycloalkyl groups, the 3 to 11-membered heterocyclyl groups, the C_{6-14} aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^2);

the substituent group V^a is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, an amino group, a carboxy group, a carbamoyl group, a sulfamoyl group, a phosphono group, a sulfo group, a tetrazolyl group, a formate group, and a formyl group;

the substituent group V^1 is a substituent group consisting of substituents constituting the substituent group V^a , C_{1-6} alkoxy groups, C_{1-3} haloalkoxy groups, mono- C_{1-6} alkylamino groups, di- C_{1-6} alkylamino groups, mono- C_{1-6} alkylaminocarbonyl groups, di- C_{1-6} alkylaminocarbonyl groups, C_{1-6} alkylcarbonylamino groups, C_{1-6} alkylthio groups, C_{1-6} alkylsulfonyl groups, C_{3-6} cycloalkoxy groups, mono- C_{3-6} cycloalkylamino groups, di- C_{3-6} cycloalkylamino groups, C_{3-6} cycloalkylcarbonyl groups, C_{3-6} cycloalkylsulfonyl groups, C_{3-6} cycloalkylthio groups, 3 to 11-membered heterocyclyl groups, C_{6-14} aryl groups, and 5 to 10-membered heteroaryl groups (the C_{3-6} cycloalkoxy groups, the mono- C_{3-6} cycloalkylamino groups, the di- C_{3-6} cycloalkylamino groups, the C_{3-6} cycloalkylcarbonyl groups, the C_{3-6} cycloalkylsulfonyl groups, the C_{3-6} cycloalkylthio groups, the 3 to 11-membered heterocyclyl groups, the C_{6-14} aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more hydroxy groups, one or more halogen atoms, one or more cyano groups, one or more

nitro groups, one or more amino groups, one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one or more phosphinoyl groups, one or more sulfo groups, one or more sulfinio groups, one or more tetrazolyl groups, one or more formyl groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups); and

the substituent group V² is a substituent group consisting of substituents constituting the substituent group V¹, C₁₋₆ alkyl groups, and C₁₋₃ haloalkyl groups], a tautomer of the compound, a pharmaceutically acceptable salt thereof, or a solvate thereof.

(2)

The compound according to (1), wherein

L³ is a C₁₋₃ alkylene group,

the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

(3)

The compound according to (1) or (2), wherein

R¹ is a hydrogen atom or a C₁₋₆ alkoxy group (the C₁₋₆ alkoxy group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁷);

the substituent group V⁷ is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C₁₋₆ alkoxy groups, C₁₋₆ alkylthio groups, C₁₋₆ alkoxycarbonyl groups (the C₁₋₆ alkoxy groups, the C₁₋₆ alkylthio groups, and the C₁₋₆ alkoxycarbonyl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V¹), C₃₋₆ cycloalkoxy groups,

mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, C₃₋₆ cycloalkyl groups, 4 to 7-membered heterocyclyl groups, a phenyl group, and 5 to 6-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆

5 cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the C₃₋₆ cycloalkyl groups, the 4 to 7-membered heterocyclyl groups, the phenyl group, and the 5 to 6-membered heteroaryl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V²);

10 the substituent group V¹ is a substituent group consisting of substituents constituting the substituent group V^a, C₁₋₆ alkoxy groups, C₁₋₃ haloalkoxy groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, mono-C₁₋₆ alkylaminocarbonyl groups, di-C₁₋₆ alkylaminocarbonyl groups, C₁₋₆ alkylcarbonylamino groups, C₁₋₆ alkylthio groups, C₁₋₆ alkylsulfonyl groups, C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, 15 di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, 3 to 11-membered heterocyclyl groups, C₆₋₁₄ aryl groups, and 5 to 10-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio 20 groups, the 3 to 11-membered heterocyclyl groups, the C₆₋₁₄ aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more hydroxy groups, one or more halogen atoms, one or more cyano groups, one or more nitro groups, one or more amino groups, one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one 25 or more phosphinoyl groups, one or more sulfo groups, one or more sulfinio groups, one or more tetrazolyl groups, one or more formyl groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more

di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups);

the substituent group V² is a substituent group consisting of substituents constituting the substituent group V¹, C₁₋₆ alkyl groups, and C₁₋₃ haloalkyl groups; and

5 the substituent group V^a is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, an amino group, a carboxy group, a carbamoyl group, a sulfamoyl group, a phosphono group, a sulfo group, a tetrazolyl group, a formate group, and a formyl group, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

(4)

The compound according to (3), wherein

R¹ is a hydrogen atom,

the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

(5)

The compound according to (1) or (2), wherein

R¹ is a C₁₋₆ alkyl group (the C₁₋₆ alkyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁹);

20 the substituent group V⁹ is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C₁₋₆ alkoxy groups, C₁₋₆ alkylthio groups, C₁₋₆ alkylcarbonyloxy groups, C₁₋₆ alkoxycarbonyl groups (the C₁₋₆ alkoxy groups, the C₁₋₆ alkylthio groups, the C₁₋₆ alkylcarbonyloxy groups, and the C₁₋₆ alkoxycarbonyl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V¹), C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, C₃₋₆ cycloalkyl groups, 4 to 7-membered heterocyclyl groups, a phenyl group, and 5 to 6-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆

cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the C₃₋₆ cycloalkyl groups, the 4 to 7-membered heterocyclyl groups, the phenyl group, and the 5 to 6-membered heteroaryl groups are unsubstituted or substituted with one or more substituents identically or differently

5 selected from the substituent group V²);

the substituent group V¹ is a substituent group consisting of substituents constituting the substituent group V^a, C₁₋₆ alkoxy groups, C₁₋₃ haloalkoxy groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, mono-C₁₋₆ alkylaminocarbonyl groups, di-C₁₋₆ alkylaminocarbonyl groups, C₁₋₆ alkylcarbonylamino groups, C₁₋₆ alkylthio groups, 10 C₁₋₆ alkylsulfonyl groups, C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, 3 to 11-membered heterocyclyl groups, C₆₋₁₄ aryl groups, and 5 to 10-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ 15 cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the 3 to 11-membered heterocyclyl groups, the C₆₋₁₄ aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more hydroxy groups, one or more halogen atoms, one or more cyano groups, one or more nitro groups, one or more amino groups, one or more carboxy groups, one or more 20 carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one or more phosphinoyl groups, one or more sulfo groups, one or more sulfinio groups, one or more tetrazolyl groups, one or more formyl groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino 25 groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups);

the substituent group V² is a substituent group consisting of substituents constituting the substituent group V¹, C₁₋₆ alkyl groups, and C₁₋₃ haloalkyl groups; and

the substituent group V^a is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, an amino group, a carboxy group, a carbamoyl group, a sulfamoyl group, a phosphono group, a sulfo group, a tetrazolyl group, a formate group, and a formyl group,

5 the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

(6)

The compound according to (5), wherein

R^1 is a C_{1-6} alkyl group (the C_{1-6} alkyl group is unsubstituted or substituted with one
10 or more halogen atoms),
the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

(7)

The compound according to any one of (1) to (6), wherein

15 L^1 is a single bond, NR^{2a} , O, S, SO, SO_2 , or a C_{1-6} alkylene group (a single methylene group of the C_{1-6} alkylene group is optionally replaced by O, S, SO_2 , C=O, C=S, or NR^{3a});

L^2 is a single bond and B is a 3 to 11-membered heterocyclylene group or a 5 to 10-membered heteroarylene group (the 3 to 11-membered heterocyclylene group and the
20 5 to 10-membered heteroarylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^b),
or

L^2 is NR^{2b} , O, S, SO, SO_2 , or a C_{1-6} alkylene group (a single methylene group of the C_{1-6} alkylene group is optionally replaced by O, S, SO_2 , C=O, C=S, or NR^{3b}) and B is a
25 C_{3-11} cycloalkylene group, a C_{3-11} cycloalkenylene group, a 3 to 11-membered heterocyclylene group, or a 5 to 10-membered heteroarylene group (the C_{3-11} cycloalkylene group, the C_{3-11} cycloalkenylene group, the 3 to 11-membered heterocyclylene group, and the 5 to 10-membered heteroarylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected

from the substituent group V^5 , and a single methylene group of the C_{3-11} cycloalkylene group and the C_{3-11} cycloalkenylene group is optionally replaced by a 1,1- C_{3-7} cycloalkylene group);

- the substituent group V^5 is a substituent group consisting of a hydroxy group,
- 5 halogen atoms, a cyano group, a nitro group, C_{1-6} alkyl groups, C_{1-6} alkoxy groups, C_{1-6} alkylthio groups, C_{1-6} alkoxycarbonyl groups (the C_{1-6} alkyl groups, the C_{1-6} alkoxy groups, the C_{1-6} alkylthio groups, and the C_{1-6} alkoxycarbonyl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^1), C_{3-6} cycloalkoxy groups, mono- C_{3-6} cycloalkylamino groups,
- 10 di- C_{3-6} cycloalkylamino groups, C_{3-6} cycloalkylcarbonyl groups, C_{3-6} cycloalkylsulfonyl groups, C_{3-6} cycloalkylthio groups, C_{3-6} cycloalkyl groups, 4 to 7-membered heterocyclyl groups, a phenyl group, and 5 to 6-membered heteroaryl groups (the C_{3-6} cycloalkoxy groups, the mono- C_{3-6} cycloalkylamino groups, the di- C_{3-6} cycloalkylamino groups, the C_{3-6} cycloalkylcarbonyl groups, the C_{3-6} cycloalkylsulfonyl groups, the C_{3-6}
- 15 cycloalkylthio groups, the C_{3-6} cycloalkyl groups, the 4 to 7-membered heterocyclyl groups, the phenyl group, and the 5 to 6-membered heteroaryl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^2);

- the substituent group V^1 is a substituent group consisting of substituents constituting
- 20 the substituent group V^a , C_{1-6} alkoxy groups, C_{1-3} haloalkoxy groups, mono- C_{1-6} alkylamino groups, di- C_{1-6} alkylamino groups, mono- C_{1-6} alkylaminocarbonyl groups, di- C_{1-6} alkylaminocarbonyl groups, C_{1-6} alkylcarbonylamino groups, C_{1-6} alkylthio groups, C_{1-6} alkylsulfonyl groups, C_{3-6} cycloalkoxy groups, mono- C_{3-6} cycloalkylamino groups, di- C_{3-6} cycloalkylamino groups, C_{3-6} cycloalkylcarbonyl groups, C_{3-6} cycloalkylsulfonyl
- 25 groups, C_{3-6} cycloalkylthio groups, 3 to 11-membered heterocyclyl groups, C_{6-14} aryl groups, and 5 to 10-membered heteroaryl groups (the C_{3-6} cycloalkoxy groups, the mono- C_{3-6} cycloalkylamino groups, the di- C_{3-6} cycloalkylamino groups, the C_{3-6} cycloalkylcarbonyl groups, the C_{3-6} cycloalkylsulfonyl groups, the C_{3-6} cycloalkylthio groups, the 3 to 11-membered heterocyclyl groups, the C_{6-14} aryl groups, and the 5 to

10-membered heteroaryl groups are unsubstituted or substituted with one or more hydroxy groups, one or more halogen atoms, one or more cyano groups, one or more nitro groups, one or more amino groups, one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one or more phosphinoyl groups, one or more sulfo groups, one or more sulfinoyl groups, one or more tetrazolyl groups, one or more formyl groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups);

the substituent group V² is a substituent group consisting of substituents constituting the substituent group V¹, C₁₋₆ alkyl groups, and C₁₋₃ haloalkyl groups;

the substituent group V^a is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, an amino group, a carboxy group, a carbamoyl group, a sulfamoyl group, a phosphono group, a sulfo group, a tetrazolyl group, a formate group, and a formyl group; and

each of R^{2a}, R^{2b}, R^{3a}, and R^{3b} is independently a hydrogen atom, a C₁₋₃ alkyl group, or a C₁₋₃ haloalkyl group,

the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

(8)

The compound according to (7), wherein

L¹ is a single bond;

L² is a single bond, and B is a 4 to 7-membered heterocyclylene group (the 4 to 7-membered heterocyclylene group is unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V³, and a single methylene group of the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group), or

L^2 is NR^{2c} , O, or a C_{1-6} alkylene group (a single methylene group of the C_{1-6} alkylene group is optionally replaced by O or NR^{3c}), and B is a C_{3-6} cycloalkylene group, a C_{3-6} cycloalkenylene group, or a 4 to 7-membered heterocyclylene group (the C_{3-6} cycloalkylene group, the C_{3-6} cycloalkenylene group, and the 4 to 7-membered heterocyclylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^3 , and a single methylene group of the C_{3-6} cycloalkylene group, the C_{3-6} cycloalkenylene group, and the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1- C_{3-7} cycloalkylene group);

the substituent group V^3 is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C_{1-6} alkyl groups, C_{1-6} haloalkyl groups, C_{3-6} cycloalkyl groups, C_{1-6} alkoxy groups, and C_{1-6} haloalkoxy groups; and

each of R^{2c} and R^{3c} is independently a hydrogen atom, a C_{1-3} alkyl group, or a C_{1-3} haloalkyl group,

the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof

(9)

The compound according to any one of (1) to (6), wherein

L^1 is a single bond;

L^2 is a single bond, and B is a 4 to 7-membered heterocyclylene group (the 4 to 7-membered heterocyclylene group is substituted with one or more substituents selected from an amino group, mono- C_{1-6} alkylamino groups, di- C_{1-6} alkylamino groups, and C_{1-6} alkylsulfonylamino groups and is optionally substituted with one or more substituents identically or differently selected from the substituent group V^3 , and a single methylene group of the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1- C_{3-7} cycloalkylene group), or

L^2 is NR^{2c} , O, or a C_{1-6} alkylene group (a single methylene group of the C_{1-6} alkylene group is optionally replaced by O or NR^{3c}), and B is a C_{3-6} cycloalkylene group, a C_{3-6} cycloalkenylene group, or a 4 to 7-membered heterocyclylene group (the C_{3-6}

cycloalkylene group, the C₃₋₆ cycloalkenylene group, and the 4 to 7-membered heterocyclylene group are substituted with a substituent selected from one or more amino groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, and one or more C₁₋₆ alkylsulfonylamino groups and are optionally substituted with one or more substituents identically or differently selected from the substituent group V³, and a single methylene group of the C₃₋₆ cycloalkylene group, the C₃₋₆ cycloalkenylene group, and the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group);

the substituent group V³ is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C₁₋₆ alkyl groups, C₁₋₆ haloalkyl groups, C₃₋₆ cycloalkyl groups, C₁₋₆ alkoxy groups, and C₁₋₆ haloalkoxy groups; and

each of R^{2c} and R^{3c} is independently a hydrogen atom, a C₁₋₃ alkyl group, or a C₁₋₃ haloalkyl group,

the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

(10)

The compound according to any one of (1) to (9), wherein

D is a 3 to 11-membered heterocyclyl group, a C₆₋₁₄ aryl group, or a 5 to 10-membered heteroaryl group (the 3 to 11-membered heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁵);

the substituent group V⁵ is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C₁₋₆ alkyl groups, C₁₋₆ alkoxy groups, C₁₋₆ alkylthio groups, C₁₋₆ alkoxycarbonyl groups (the C₁₋₆ alkyl groups, the C₁₋₆ alkoxy groups, the C₁₋₆ alkylthio groups, and the C₁₋₆ alkoxycarbonyl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V¹), C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, C₃₋₆ cycloalkyl groups, 4 to 7-membered heterocyclyl

groups, a phenyl group, and 5 to 6-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the C₃₋₆ cycloalkyl groups, the 4 to 7-membered heterocyclyl groups, the phenyl group, and the 5 to 6-membered heteroaryl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V²);

the substituent group V¹ is a substituent group consisting of substituents constituting the substituent group V^a, C₁₋₆ alkoxy groups, C₁₋₃ haloalkoxy groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, mono-C₁₋₆ alkylaminocarbonyl groups, di-C₁₋₆ alkylaminocarbonyl groups, C₁₋₆ alkylcarbonylamino groups, C₁₋₆ alkylthio groups, C₁₋₆ alkylsulfonyl groups, C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, 3 to 11-membered heterocyclyl groups, C₆₋₁₄ aryl groups, and 5 to 10-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the 3 to 11-membered heterocyclyl groups, the C₆₋₁₄ aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more hydroxy groups, one or more halogen atoms, one or more cyano groups, one or more nitro groups, one or more amino groups, one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one or more phosphinoyl groups, one or more sulfo groups, one or more sulfinio groups, one or more tetrazolyl groups, one or more formyl groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups);

the substituent group V^2 is a substituent group consisting of substituents constituting the substituent group V^1 , C_{1-6} alkyl groups, and C_{1-3} haloalkyl groups; and

the substituent group V^a is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, an amino group, a carboxy group, a carbamoyl group, a sulfamoyl group, a phosphono group, a sulfo group, a tetrazolyl group, a formate group, and a formyl group.

the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

(11)

10 The compound according to (10), wherein

D is a 4 to 7-membered heterocyclyl group, a phenyl group, or a 5 to 6-membered heteroaryl group (the 4 to 7-membered heterocyclyl group, the phenyl group, and the 5 to 6-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^4);

15 the substituent group V^4 is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C_{1-6} alkyl groups, C_{1-6} haloalkyl groups, C_{1-6} alkoxy groups, C_{1-6} haloalkoxy groups, C_{3-6} cycloalkyl groups, C_{3-6} cycloalkoxy groups, mono- C_{3-6} cycloalkylamino groups, di- C_{3-6} cycloalkylamino groups, and C_{3-6} cycloalkylthio groups (the C_{1-6} alkyl groups, the C_{1-6} alkoxy groups, the C_{3-6} cycloalkyl groups, the C_{3-6} cycloalkoxy groups, the mono- C_{3-6} cycloalkylamino groups, the di- C_{3-6} cycloalkylamino groups, and the C_{3-6} cycloalkylthio groups are unsubstituted or substituted with one or more hydroxy groups, one or more amino groups, one or more nitro groups, one or more cyano groups, one or more 3 to 11-membered heterocyclyl groups, one or more C_{6-14} aryl groups, or one or more 5 to 10-membered heteroaryl groups (the 3 to 11-membered heterocyclyl groups, the C_{6-14} aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one or more sulfo groups, one or more tetrazolyl groups, one or more formyl groups, one or more nitro groups, one or more cyano groups, one or more

halogen atoms, one or more hydroxy groups, one or more amino groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups)),
 the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

(12)

10 The compound according to (11), wherein

D is a phenyl group or a 5 to 6-membered heteroaryl group (the phenyl group or the 5 to 6-membered heteroaryl group is unsubstituted or substituted with one or more halogen atoms, one or more nitro groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₆ haloalkyl groups, one or more C₁₋₆ alkoxy groups, or one or more C₁₋₆ haloalkoxy groups),
 the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof

(13)

The compound according to any one of (1) to (9), wherein

20 D is a phenyl group or a 5 to 6-membered heteroaryl group (the phenyl group or the 5 to 6-membered heteroaryl group is substituted with one or more C₁₋₆ alkylsulfonylamino groups or one or more C₁₋₆ alkylsulfonyloxy groups (the C₁₋₆ alkylsulfonylamino groups and the C₁₋₆ alkylsulfonyloxy groups are unsubstituted or substituted with one or more halogen atoms, one or more nitro groups, one or more C₁₋₆ alkoxy groups, or one or more C₁₋₆ haloalkoxy groups) and is optionally substituted with
 25 one or more halogen atoms, one or more nitro groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₆ haloalkyl groups, one or more C₁₋₆ alkoxy groups, or one or more C₁₋₆ haloalkoxy groups),
 the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate

thereof.

(14)

The compound according to (13), wherein

D is a 5 to 6-membered heteroaryl group (the 5 to 6-membered heteroaryl group is substituted with one or more C₁₋₆ alkylsulfonyloxy groups (the C₁₋₆ alkylsulfonyloxy groups are unsubstituted or substituted with one or more halogen atoms, one or more nitro groups, one or more C₁₋₆ alkoxy groups, or one or more C₁₋₆ haloalkoxy groups)), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

10 (15)

The compound, according to any one of (1) to (14), wherein

A is a 3 to 11-membered heterocyclyl group, a C₆₋₁₄ aryl group, or a 5 to 10-membered heteroaryl group (the 3 to 11-membered heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁵); and

the substituent group V⁵ is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C₁₋₆ alkyl groups, C₁₋₆ alkoxy groups, C₁₋₆ alkylthio groups, C₁₋₆ alkoxycarbonyl groups (the C₁₋₆ alkyl groups, the C₁₋₆ alkoxy groups, the C₁₋₆ alkylthio groups, and the C₁₋₆ alkoxycarbonyl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V¹), C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, C₃₋₆ cycloalkyl groups, 4 to 7-membered heterocyclyl groups, a phenyl group, and 5 to 6-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the C₃₋₆ cycloalkyl groups, the 4 to 7-membered heterocyclyl groups, the phenyl group, and the 5 to 6-membered heteroaryl groups are unsubstituted or

substituted with one or more substituents identically or differently selected from the substituent group V²);

the substituent group V¹ is a substituent group consisting of substituents constituting the substituent group V^a, C₁₋₆ alkoxy groups, C₁₋₃ haloalkoxy groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, mono-C₁₋₆ alkylaminocarbonyl groups, di-C₁₋₆ alkylaminocarbonyl groups, C₁₋₆ alkylcarbonylamino groups, C₁₋₆ alkylthio groups, C₁₋₆ alkylsulfonyl groups, C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, 3 to 11-membered heterocyclyl groups, C₆₋₁₄ aryl groups, and 5 to 10-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the 3 to 11-membered heterocyclyl groups, the C₆₋₁₄ aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more hydroxy groups, one or more halogen atoms, one or more cyano groups, one or more nitro groups, one or more amino groups, one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one or more phosphinoyl groups, one or more sulfo groups, one or more sulfinio groups, one or more tetrazolyl groups, one or more formyl groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups);

the substituent group V² is a substituent group consisting of substituents constituting the substituent group V¹, C₁₋₆ alkyl groups, and C₁₋₃ haloalkyl groups; and

the substituent group V^a is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, an amino group, a carboxy group, a carbamoyl group, a sulfamoyl group, a phosphono group, a sulfo group, a tetrazolyl

group, a formate group, and a formyl group,

the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

(16)

5 The compound according to (15), wherein

A is a phenyl group or a 5 to 6-membered heteroaryl group (the phenyl group and the 5 to 6-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^3); and

the substituent group V^3 is a substituent group consisting of a hydroxy group,

10 halogen atoms, a cyano group, a nitro group, C_{1-6} alkyl groups, C_{1-6} haloalkyl groups, C_{3-6} cycloalkyl groups, C_{1-6} alkoxy groups, and C_{1-6} haloalkoxy groups,
the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

(17)

15 A T-type calcium channel inhibitor comprising the compound described in any one of (1) to (16), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof, as an active component.

(18)

20 A preventive agent, a therapeutic agent, and/or an improving agent for a disease treatable by a T-type calcium channel inhibitory action comprising the T-type calcium channel inhibitor as described in (17) as an active component.

(19)

A therapeutic agent for neuropathic pain comprising the T-type calcium channel inhibitor as described in (18) as an active component.

25 (20)

A medicine comprising the compound described in any one of (1) to (19), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof, as an active component.

Effects of the Invention

[0011] The present invention can provide novel triazinone compounds that have an excellent T-type calcium channel inhibitory activity and are specifically useful for prevention or treatment of neuropathic pain.

5

MODES FOR CARRYING OUT THE INVENTION

[0012] The present invention will now be described in further detail.

In the present invention, "n-" is normal, "i-" is iso, "s-" and "sec-" are secondary, "t-" and "tert-" are tertiary, "o-" is ortho, "m-" is meta, "p-" is para, "Ph" is phenyl, "Bu" is butyl, "Boc" is tert-butoxycarbonyl, "Z" is benzyloxycarbonyl, "Ts" is p-toluenesulfonyl, and "Bn" is benzyl.

[0013] First, terms used for the explanation of chemical structures in the present specification will be described.

15 [0014] The "halogen atom" means a fluorine atom, a chlorine atom, a bromine atom, and an iodine atom.

[0015] The "C₁₋₃ alkyl group" means a methyl group, an ethyl group, a propyl group, and an isopropyl group.

20 [0016] The "C₁₋₆ alkyl group" means linear or branched alkyl groups having a carbon number of 1 to 6, and specific examples include a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a t-butyl group, an n-pentyl group, and an n-hexyl group.

[0017] The "C₁₋₃ alkylene group" means a methylene group, an ethylene group, a propane-1,3-diyl group, and a propane-1,2-diyl group.

25 [0018] The "C₁₋₆ alkylene group" means divalent substituents formed by removing a single hydrogen atom at any position of the "C₁₋₆ alkyl group" defined above, and specific examples include a methylene group, an ethylene group, a propane-1,3-diyl group, a propane-1,2-diyl group, a 2,2-dimethyl-propane-1,3-diyl group, a hexane-1,6-diyl group, and a 3-methylbutane-1,2-diyl group.

[0019] The "C₁₋₃ haloalkyl group" means substituents formed by substituting one or more hydrogen atoms at any position of the "C₁₋₃ alkyl group" defined above by one or more halogen atoms identically or differently selected from a substituent group consisting of a fluorine atom, a chlorine atom, a bromine atom, and an iodine atom, and specific
 5 examples include a trifluoromethyl group, a 2,2,2-trifluoroethyl group, a pentafluoroethyl group, and a 3-chloro-n-propyl group.

[0020] The "C₁₋₆ haloalkyl group" means substituents formed by substituting one ore more hydrogen atoms at any position of the "C₁₋₆ alkyl group" defined above by one or more halogen atoms identically or differently selected from a substituent group
 10 consisting of a fluorine atom, a chlorine atom, a bromine atom, and an iodine atom, and specific examples include a trifluoromethyl group, a pentafluoroethyl group, a 2,2,2-trifluoro-1,1-dimethyl-ethyl group, a 3-chloro-n-propyl group, and a 4-chloro-n-butyl group.

[0021] The "C₃₋₁₁ cycloalkane" means monocyclic-, condensed-, bridged-, or
 15 spiro-system aliphatic hydrocarbon rings having a ring-constituting carbon number of 3 to 11, and specific examples include cyclopropane, cyclobutane, cyclopentane, cyclohexane, cycloheptane, cyclooctane, adamantane, bicyclo[3.1.0]octane, bicyclo[2.2.1]heptane, and spiro[5.5]undecane.

[0022] The "C₃₋₁₁ cycloalkyl group" means monovalent substituents formed by
 20 removing a single hydrogen atom at any position of the "C₃₋₁₁ cycloalkane" defined above.

[0023] The "C₃₋₁₁ cycloalkylene group" means divalent substituents formed by removing two hydrogen atoms at any positions on different carbons of the "C₃₋₁₁ cycloalkane" defined above.


25 [0024] The "C₃₋₆ cycloalkane" means cycloalkanes having a ring-constituting carbon number of 3 to 6 among the "C₃₋₁₁ cycloalkanes" defined above, and specific examples include cyclopropane, cyclobutane, cyclopentane, and cyclohexane.

[0025] The "C₃₋₆ cycloalkyl group" means monovalent substituents formed by removing a single hydrogen atom at any position of the "C₃₋₆ cycloalkane" defined above.

[0026] The "C₃₋₆ cycloalkylene group" means divalent substituents formed by removing two hydrogen atoms at any positions on different carbons of the "C₃₋₆ cycloalkane" defined above.

[0027] The "C₃₋₇ cycloalkane" means cycloalkanes having a ring-constituting
 5 carbon number of 3 to 7 among the "C₃₋₁₁ cycloalkanes" defined above, and specific examples include cyclopropane, cyclobutane, cyclopentane, cyclohexane, and cycloheptane.

[0028] The "1,1-C₃₋₇ cycloalkylene group" means divalent substituents formed by removing two hydrogen atoms on the same carbon of the "C₃₋₇ cycloalkane" defined
 10 above. The group is specifically exemplified by the structures shown in figures below.



[0029] The "C₄₋₇ cycloalkane" means cycloalkanes having a ring-constituting carbon number of 4 to 7 among the "C₃₋₁₁ cycloalkanes" defined above, and specific examples include cyclobutane, cyclopentane, cyclohexane, and cycloheptane.

15 [0030] The "C₄₋₇ cycloalkylene group" means divalent substituents formed by removing two hydrogen atoms at any positions on different carbons of the "C₄₋₇ cycloalkane" defined above.

[0031] The "C₃₋₁₁ cycloalkene" means non-aromatic rings formed by replacing one or more any bonds of the "C₃₋₁₁ cycloalkane" defined above by double bonds, and
 20 specific examples include cyclopropene, cyclobutene, cyclopentene, cyclohexene, cyclohexa-1,3-diene, cyclohexa-1,4-diene, bicyclo[2.2.1]hepta-2,5-diene, spiro[2.5]oct-4-ene, and 1,2,5,6-tetrahydronaphthalene.

[0032] The "C₃₋₁₁ cycloalkenyl group" means monovalent substituents formed by removing a single hydrogen atom at any position of the "C₃₋₁₁ cycloalkene" defined
 25 above.

[0033] The "C₃₋₁₁ cycloalkenylene group" means divalent substituents formed by removing two hydrogen atoms at any positions on different carbons of the "C₃₋₁₁

cycloalkene" defined above.

[0034] The "C₄₋₇ cycloalkene" means cycloalkenes having a ring-constituting carbon number of 4 to 7 among the "C₃₋₁₁ cycloalkenes" defined above.

[0035] The "C₄₋₇ cycloalkenylene group" means divalent substituents formed by removing two hydrogen atoms at any positions on different carbons of the "C₄₋₇ cycloalkene" defined above.

[0036] The "C₃₋₆ cycloalkene" means cycloalkenes having a ring-constituting carbon number of 3 to 6 among the "C₃₋₁₁ cycloalkenes" defined above, and specific examples include cyclopropene, cyclobutene, cyclopentene, and cyclohexene.

[0037] The "C₃₋₆ cycloalkenylene group" means divalent substituents formed by removing two hydrogen atoms at any positions on different carbons of the "C₃₋₆ cycloalkene" defined above.

[0038] The "C₂₋₆ alkenyl group" means linear or branched alkenyl groups having at least one double bond and a carbon number of 2 to 6, and specific examples include an ethenyl (vinyl) group, a 1-propenyl group, a 2-propenyl (allyl) group, an isopropenyl group, a 1-butenyl group, a 2-butenyl group, a 3-butenyl (homoallyl) group, a 4-pentenyl group, and a 5-hexenyl group.

[0039] The "C₂₋₆ alkenylene group" means divalent substituents formed by removing a single hydrogen atom at any position of the "C₂₋₆ alkenyl group" defined above, and specific examples include an ethenyl group, a prop-1-en-1-yl group, a prop-2-en-1-yl group, a prop-1-en-2-yl group, a but-1-en-1-yl group, a but-2-en-1-yl group, a but-3-en-1-yl group, a pent-1-en-1-yl group, a pent-4-en-1-yl group, a hex-1-en-1-yl group, a hex-5-en-1-yl group, a 4-methylpent-3-en-1-yl group, and a penta-2,4-dien-1-yl group.

[0040] The "C₂₋₆ haloalkenyl group" means substituents formed by substituting one or more hydrogen atoms at any positions of the "C₂₋₆ alkenyl group" defined above by one or more halogen atoms identically or differently selected from a substituent group consisting of a fluorine atom, a chlorine atom, a bromine atom, and an iodine atom.

[0041] The "C₂₋₆ alkynyl group" means linear or branched alkynyl groups having

at least one triple bond and a carbon number of 2 to 6, and specific examples include an ethynyl group, a 1-propynyl group, a 2-propynyl group, a 1-butynyl group, a 2-butynyl group, a 3-butynyl group, a 4-pentynyl group, and a 5-hexynyl group.

[0042] The "C₂₋₆ alkynylene group" means divalent substituents formed by removing a single hydrogen atom at any position of the "C₂₋₆ alkynyl group" defined above. Specific examples of the group include an ethynylene group, a 1-propynylene group, a 2-propynylene group, a 1-butynylene group, a 2-butynylene group, a 3-butynylene group, a 4-pentynylene group, and a 5-hexynylene group.

[0043] The "C₁₋₆ alkoxy group" means linear or branched alkoxy groups having a carbon number of 1 to 6, and specific examples include a methoxy group, an ethoxy group, an n-propoxy group, an isopropoxy group, an n-butoxy group, an isobutoxy group, a t-butoxy group, an n-pentyloxy group, and an n-hexyloxy group.

[0044] The "C₃₋₆ cycloalkoxy group" means groups formed by bonding the single "C₃₋₆ cycloalkyl group" to -O-, and specific examples include a cyclopropoxy group, a cyclobutoxy group, a cyclopentyloxy group, and a cyclohexyloxy group.

[0045] The "C₁₋₃ alkoxy group" means a methoxy group, an ethoxy group, an n-propoxy group, and an isopropoxy group.

[0046] The "C₁₋₆ haloalkoxy group" means substituents formed by substituting one or more hydrogen atoms at any positions of the "C₁₋₆ alkoxy group" defined above by one or more halogen atoms identically or differently selected from a substituent group consisting of a fluorine atom, a chlorine atom, a bromine atom, and an iodine atom, and specific examples include a trifluoromethoxy group, a pentafluoroethoxy group, a 2,2,2-trifluoro-1,1-dimethyl-ethoxy group, a 3-chloro-n-propyloxy group, and a 4-chloro-n-butoxy group.

[0047] The "C₁₋₃ haloalkoxy group" means substituents formed by substituting one or more hydrogen atoms at any positions of the "C₁₋₃ alkoxy group" defined above by one or more halogen atoms identically or differently selected from a substituent group consisting of a fluorine atom, a chlorine atom, a bromine atom, and an iodine atom, and specific examples include a trifluoromethoxy group, a 2,2,2-trifluoroethoxy group, a

pentafluoroethoxy group, and a 3-chloro-n-propyloxy group.

[0048] The "C₆₋₁₄ aromatic hydrocarbon ring" means monocyclic or polycyclic aromatic hydrocarbon rings in which all the atoms constituting the rings are carbon atoms and the number of carbons is 6 to 14 and includes bicyclic condensed rings composed of
 5 a monocyclic aromatic hydrocarbon ring and a monocyclic cycloalkanes or cycloalkenes. Specific examples of the ring include benzene, pentalene, naphthalene, azulene, anthracene, phenanthrene, indene, indane, dihydronaphthalene, and tetrahydronaphthalene.

[0049] The "C₆₋₁₄ aryl group" means monovalent substituents formed by
 10 removing a single hydrogen atom at any position on the aromatic ring of the "C₆₋₁₄ aromatic hydrocarbon ring" defined above. The binding position is not limited, and the group may be bonded at a desired position.

[0050] The "C₆₋₁₄ arylene group" means divalent substituents formed by removing two hydrogen atoms at any positions on the aromatic ring of the "C₆₋₁₄ aromatic
 15 hydrocarbon ring" defined above. The binding positions are not limited, and the groups may be bonded at a desired positions.

[0051] The "5 to 10-membered aromatic heterocycle" means monocyclic or condensed aromatic heterocycles having a ring-constituting atom number of 5 to 10 and containing 1 to 5 hetero atoms as the atoms constituting the ring (the hetero atom is a
 20 nitrogen atom, an oxygen atom, or a sulfur atom), and specific examples include furan, thiophene, pyrrole, imidazole, triazole, tetrazole, thiazole, pyrazole, oxazole, isoxazole, isothiazole, thiadiazole, oxadiazole, pyridine, pyrazine, pyridazine, pyrimidine, triazine, purine, pteridine, quinoline, isoquinoline, naphthyridine, quinoxaline, cinnoline, quinazoline, phthalazine, imidazopyridine, imidazothiazole, imidazooxazole,
 25 benzothiazole, benzoxazole, benzimidazole, indoline, isoindoline, indazoline, pyrrolopyridine, thienopyridine, furopyridine, benzothiadiazole, benzooxadiazole, pyridopyrimidine, benzofuran, benzothiophene, and thienofuran.

[0052] When having a C=N double bond, the "5 to 10-membered aromatic heterocycle" includes N-oxides form.

[0053] The "5 to 10-membered aromatic heterocycle containing NH" means aromatic heterocycles having NH among the "5 to 10-membered aromatic heterocycles" defined above, and specific examples include pyrrole, imidazole, triazole, tetrazole, pyrazole, purine, pteridine, benzimidazole, indoline, isoindoline, indazoline, and pyrrolopyridine.

[0054] The "5 to 10-membered heteroaryl group" means monovalent substituents formed by removing a single hydrogen atom at any position of the "5 to 10-membered aromatic heterocycle" defined above. The binding position is not limited, and the group may be bonded at a desired position.

10 [0055] The "5 to 10-membered heteroaryl group containing NH" means heteroaryl groups having NH among the "5 to 10-membered heteroaryl groups" defined above. The binding position is not limited, and the group may be bonded at a desired position.

[0056] The "5 to 10-membered heteroarylene group" means divalent substituents
15 formed by removing two hydrogen atoms at any positions of the "5 to 10-membered aromatic heterocycle" defined above. The binding positions are not limited, and the groups may be bonded at desired positions.

[0057] The "5 to 10-membered heteroarylene group containing NH" means heteroarylene groups having NH among the "5 to 10-membered heteroarylene groups" defined above. The binding positions are not limited, and the groups may be bonded at
20 desired positions.

[0058] The "5 to 6-membered aromatic heterocycle" means monocyclic aromatic heterocycles having a ring-constituting atom number of 5 to 6 among the "5 to 10-membered aromatic heterocycles" defined above, and specific examples include
25 pyrrole, pyrazole, imidazole, triazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, triazine, furan, thiophene, thiazole, isothiazole, oxazole, isoxazole, oxadiazole, and thiadiazole.

[0059] The "5 to 6-membered heteroaryl group" means monovalent substituents formed by removing a single hydrogen atom at any position of the "5 to 6-membered

aromatic heterocycle" defined above. The binding position is not limited, and the group may be bonded at a desired position.

[0060] The "3 to 11-membered non-aromatic heterocycle" means bridged-, or spiro-system non-aromatic heterocycles that

- 5 1) have a ring-constituting atom number of 3 to 11,
- 2) contain 1 to 5 hetero atoms as the atoms constituting the ring (the hetero atom is a nitrogen atom, an oxygen atom, or a sulfur atom),
- 3) may contain one or more carbonyl group, one or more thiocarbonyl group, one or more double bond, or one or more triple bond in the ring,
- 10 4) when containing sulfur atoms as the atom constituting the ring, the sulfur atom may be replaced by a sulfinyl group or a sulfonyl group, and
- 5) is monocyclic-, condensed-, (in a condensed non-aromatic heterocycle, non-aromatic rings may be condensed with each other or a single non-aromatic ring may be condensed with an aromatic ring), bridged-, or spiro-system non-aromatic heterocycles, and
- 15 specific examples include azetidine, pyrrolidine, piperidine, azepane, azocane, tetrahydrofuran, tetrahydropyran, morpholine, thiomorpholine, piperazine, thiazolidine, 1,4-dioxane, imidazoline, thiazoline, 1,2-benzopyran, isochroman, chroman, indoline, isoindoline, azaindane, azatetrahydronaphthalene, azachroman, tetrahydrobenzofuran, tetrahydrobenzothiophene, 2,3,4,5-tetrahydro-benzo[b]thiophene,
- 20 3,4-dihydro-2H-benzo[b][1,4]dioxepane, indan-1-one, 6,7-dihydro-5H-cyclopentapyrazine, 6,7-dihydro-5H-[1]pyridine, 6,7-dihydro-5H-[1]pyridine, 5,6-dihydro-4H-cyclopenta[b]thiophene, 4,5,6,7-tetrahydro-benzo[b]thiophene, 3,4-dihydro-2H-naphthalen-1-one, 2,3-dihydro-isoindol-1-one, 3,4-dihydro-2H-isoquinolin-1-one,
- 25 3,4-dihydro-2H-benzo[b]oxepin-5-one, pyridone, and 1-H-benzo[d]imidazole-2(3H)-thione.

[0061] The "3 to 11-membered non-aromatic heterocycle containing NH" means non-aromatic heterocycles having NH among the "3 to 11-membered non-aromatic heterocycles" defined above, and specific examples include aziridine, azetidine,

pyrrolidine, piperidine, piperazine, morpholine, thiomorpholine, thiazolidine, imidazoline, thiazoline, indoline, isoindoline, azaindane, azatetrahydronaphthalene, azachroman, 6,7-dihydro-5H-cyclopentapyrazine, 6,7-dihydro-5H-[1]pyridine, 6,7-dihydro-5H-[1]pyridine, 2,3-dihydro-isoindol-1-one, 3,4-dihydro-2H-isoquinolin-1-one, pyridone, and 1-H-benzo[d]imidazole-2(3H)-thione.

[0062] The "3 to 11-membered heterocyclyl group" means monovalent substituents formed by removing a hydrogen atom at any position of the "3 to 11-membered non-aromatic heterocycle" defined above. The binding position is not limited, and the group may be bonded at a desired position.

10 [0063] The "3 to 11-membered heterocyclyl group containing NH" means heterocyclyl groups having NH among the "3 to 11-membered heterocyclyl groups" defined above. The binding position is not limited, and the group may be bonded at a desired position.

[0064] The "3 to 11-membered heterocyclylene group" means divalent substituents formed by removing two hydrogen atoms at any positions on different atoms of the "3 to 11-membered non-aromatic heterocycle" defined above. The binding positions are not limited, and the groups may be bonded at desired positions. In the case of a condensed heterocyclylene group having a non-aromatic ring condensed with an aromatic ring, the heterocyclylene group is substituted on the non-aromatic ring.

20 [0065] The "3 to 11-membered heterocyclylene group containing NH" means heterocyclylene groups having NH among the "3 to 11-membered heterocyclylene groups" defined above. The binding positions are not limited, and the groups may be bonded at desired positions.

[0066] The "4 to 7-membered non-aromatic heterocycle" means monocyclic non-aromatic heterocycles that

- 1) have a ring-constituting atom number of 4 to 7,
- 2) contain 1 to 3 hetero atoms as the atoms constituting the ring (the hetero atom is a nitrogen atom, an oxygen atom, or a sulfur atom),
- 3) may contain a carbonyl group, a double bond, or a triple bond in the ring, and

4) when containing sulfur atoms as the atom constituting the ring, the sulfur atom may be replaced by a sulfinyl group or a sulfonyl group, and

specific examples include azetidine, pyrrolidine, pyrrolidinone, oxazolidine, isoxazolidine, thiazolidine, isothiazolidine, piperazine, piperazinone, piperidine,

5 piperidinone, morpholine, thiomorpholine, azepine, diazepine, oxetane, tetrahydrofuran, 1,3-dioxolane, tetrahydropyran, 1,4-dioxane, oxepane, and homomorpholine.

[0067] The "4 to 7-membered heterocyclcyl group" means monovalent substituents formed by removing a single hydrogen atom at any position of the "4 to 7-membered non-aromatic heterocycle" defined above. The binding position is not limited, and the
10 group may be bonded at a desired position.

[0068] The "4 to 7-membered heterocyclcylene group" means divalent substituents formed by removing two hydrogen atoms at any positions on different atoms of the "4 to 7-membered non-aromatic heterocycle" defined above. The binding positions are not limited, and the group may be bonded at desired positions.

15 [0069] The "C₁₋₆ alkylthio group" means groups formed by bonding the single "C₁₋₆ alkyl group" to -S-, and specific examples include a methylthio group, an ethylthio group, an n-propylthio group, an isopropylthio group, an n-butylthio group, an isobutylthio group, a t-butylthio group, an n-pentylthio group, and an n-hexylthio group.

[0070] The "C₃₋₆ cycloalkylthio group" means groups formed by bonding the
20 single "C₃₋₆ cycloalkyl group" to -S-, and specific examples include a cyclopropylthio group, a cyclobutylthio group, a cyclopentylthio group, and a cyclohexylthio group.

[0071] The "C₁₋₆ haloalkylthio group" means substituents formed by substituting one or more hydrogen atoms at any positions of the "C₁₋₆ alkylthio group" defined above by one or more halogen atoms identically or differently selected from a substituent group
25 consisting of a fluorine atom, a chlorine atom, a bromine atom, and an iodine atom.

[0072] The "C₁₋₆ alkylsulfonyl group" means groups formed by bonding the single "C₁₋₆ alkyl group" to a sulfonyl group, and specific examples include a methylsulfonyl group, an ethylsulfonyl group, an n-propylsulfonyl group, an isopropylsulfonyl group, an n-butylsulfonyl group, an isobutylsulfonyl group, a

t-butylsulfonyl group, an n-pentylsulfonyl group, and an n-hexylsulfonyl group.

[0073] The "C₁₋₆ alkylsulfonylamino group" means groups formed by bonding the single "C₁₋₆ alkylsulfonyl group" to an amino group, and specific examples include a methylsulfonylamino group, an ethylsulfonylamino group, an n-propylsulfonylamino group, an isopropylsulfonylamino group, an n-butylsulfonylamino group, an isobutylsulfonylamino group, a t-butylsulfonylamino group, an n-pentylsulfonylamino group, and an n-hexylsulfonylamino group.

[0074] The "C₁₋₆ alkylsulfonyloxy group" means groups formed by bonding the single "C₁₋₆ alkylsulfonyl group" to an oxy group, and specific examples include a methylsulfonyloxy group, an ethylsulfonyloxy group, an n-propylsulfonyloxy group, an isopropylsulfonyloxy group, an n-butylsulfonyloxy group, an isobutylsulfonyloxy group, a t-butylsulfonyloxy group, an n-pentylsulfonyloxy group, and an n-hexylsulfonyloxy group.

[0075] The "C₃₋₆ cycloalkylsulfonyl group" means groups formed by bonding the single "C₃₋₆ cycloalkyl group" to a sulfonyl group, and specific examples include a cyclopropylsulfonyl group, a cyclobutylsulfonyl group, a cyclopentylsulfonyl group, and a cyclohexylsulfonyl group.

[0076] The "C₃₋₆ cycloalkylsulfonylamino group" means groups formed by bonding the single "C₃₋₆ cycloalkylsulfonyl group" to an amino group, and specific examples include a cyclopropylsulfonylamino group, a cyclobutylsulfonylamino group, a cyclopentylsulfonylamino group, and a cyclohexylsulfonylamino group.

[0077] The "C₃₋₆ cycloalkylsulfonyloxy group" means groups formed by bonding the single "C₃₋₆ cycloalkylsulfonyl group" to an oxy group, and specific examples include a cyclopropylsulfonyloxy group, a cyclobutylsulfonyloxy group, a cyclopentylsulfonyloxy group, and a cyclohexylsulfonyloxy group.

[0078] The "C₁₋₆ haloalkylsulfonyl group" means substituents formed by substituting one or more hydrogen atoms at any positions of the "C₁₋₆ alkylsulfonyl group" defined above by one or more halogen atoms identically or differently selected from a substituent group consisting of a fluorine atom, a chlorine atom, a bromine atom,

and an iodine atom.

[0079] The "C₁₋₆ alkoxy carbonyl group" means groups formed by bonding the single "C₁₋₆ alkoxy group" to a carbonyl group, and specific examples include a methoxy carbonyl group, an ethoxy carbonyl group, an n-propoxy carbonyl group, an isopropoxy carbonyl group, an n-butoxy carbonyl group, an isobutoxy carbonyl group, a t-butoxy carbonyl group, an n-pentyloxy carbonyl group, and an n-hexyloxy carbonyl group.

[0080] The "mono-C₁₋₆ alkylamino group" means groups formed by bonding the single "C₁₋₆ alkyl group" to an amino group, and specific examples include a methylamino group, an ethylamino group, an n-propylamino group, an isopropylamino group, an n-butylamino group, an isobutylamino group, a t-butylamino group, an n-pentylamino group, and an n-hexylamino group.

[0081] The "di-C₁₋₆ alkylamino group" means groups formed by bonding the two "C₁₋₆ alkyl groups", which may be the same as or different from each other, to an amino group, and specific examples include a dimethylamino group, a diethylamino group, a di-n-propylamino group, a diisopropylamino group, a di-n-butylamino group, a diisobutylamino group, a di-t-butylamino group, a di-n-pentylamino group, a di-n-hexylamino group, a N-ethyl-N-methylamino group, a N-methyl-N-n-propylamino group, a N-isopropyl-N-methylamino group, a N-n-butyl-N-methylamino group, a N-isobutyl-N-methylamino group, a N-t-butyl-N-methylamino group, a N-methyl-N-n-pentylamino group, a N-n-hexyl-N-methylamino group, a N-ethyl-N-n-propylamino group, a N-ethyl-N-isopropylamino group, a N-n-butyl-N-ethylamino group, a N-ethyl-N-isobutylamino group, a N-t-butyl-N-ethylamino group, a N-ethyl-N-n-pentylamino group, and a N-ethyl-N-n-hexylamino group.

[0082] The "mono-C₃₋₆ cycloalkylamino group" means groups formed by bonding the single "C₃₋₆ cycloalkyl group" to an amino group, and specific examples include a cyclopropylamino group, a cyclobutylamino group, a cyclopentylamino group, and a cyclohexylamino group.

[0083] The "di-C₃₋₆ cycloalkylamino group" means groups formed by bonding the two "C₃₋₆ cycloalkyl groups", which may be the same as or different from each other, to an amino group, and specific examples include a dicyclopropylamino group, a dicyclobutylamino group, a dicyclopentylamino group, and a dicyclohexylamino group.

5 [0084] The "C₁₋₆ alkylcarbonyl group" means groups formed by bonding the single "C₁₋₆ alkyl group" to a carbonyl group, and specific examples include an acetyl group, a propionyl group, a butyryl group, an isobutyryl group, a pentanoyl group, a 3-methylbutanoyl group, a pivaloyl group, a hexanoyl group, and a heptanoyl group.

10 [0085] The "C₃₋₆ cycloalkylcarbonyl group" means groups formed by bonding the single "C₃₋₆ cycloalkyl group" to a carbonyl group, and specific examples include a cyclopropylcarbonyl group, a cyclobutylcarbonyl group, a cyclopentylcarbonyl group, a cyclohexylcarbonyl group, and a cycloheptylcarbonyl group.

[0086] The "C₁₋₆ haloalkylcarbonyl group" means substituents formed by substituting one or more hydrogen atoms at any position of the "C₁₋₆ alkylcarbonyl group" defined above by one or more halogen atoms identically or differently selected from a substituent group consisting of a fluorine atom, a chlorine atom, a bromine atom, and an iodine atom.

[0087] The "mono-C₁₋₆ alkylaminocarbonyl group" means groups formed by bonding the single "mono-C₁₋₆ alkylamino group" to a carbonyl group, and specific examples include a methylaminocarbonyl group, an ethylaminocarbonyl group, an n-propylaminocarbonyl group, an isopropylaminocarbonyl group, an n-butylaminocarbonyl group, an isobutylaminocarbonyl group, a t-butylaminocarbonyl group, an n-pentylaminocarbonyl group, and an n-hexylaminocarbonyl group.

25 [0088] The "di-C₁₋₆ alkylaminocarbonyl group" means groups formed by bonding the single "di-C₁₋₆ alkylamino group" to a carbonyl group, and specific examples include a dimethylaminocarbonyl group, a diethylaminocarbonyl group, a di-n-propylaminocarbonyl group, a diisopropylaminocarbonyl group, a di-n-butylaminocarbonyl group, a diisobutylaminocarbonyl group, a di-t-butylaminocarbonyl group, a di-n-pentylaminocarbonyl group, a

di-n-hexylaminocarbonyl group, a N-ethyl-N-methylaminocarbonyl group, a N-methyl-N-n-propylaminocarbonyl group, a N-isopropyl-N-methylaminocarbonyl group, a N-n-butyl-N-methylaminocarbonyl group, a N-isobutyl-N-methylaminocarbonyl group, a N-t-butyl-N-methylaminocarbonyl group, a N-methyl-N-n-pentylaminocarbonyl group, a N-n-hexyl-N-methylaminocarbonyl group, a N-ethyl-N-n-propylaminocarbonyl group, a N-ethyl-N-isopropylaminocarbonyl group, a N-n-butyl-N-ethylaminocarbonyl group, a N-ethyl-N-isobutylaminocarbonyl group, a N-t-butyl-N-ethylaminocarbonyl group, a N-ethyl-N-n-pentylaminocarbonyl group, and a N-ethyl-N-n-hexylaminocarbonyl group.

- 10 [0089] The " C_{1-6} alkylcarbonylamino group" means groups formed by bonding the single " C_{1-6} alkylcarbonyl group" to an amino group, and specific examples include a methylcarbonylamino group, an ethylcarbonylamino group, an n-propylcarbonylamino group, an isopropylcarbonylamino group, an n-butylcarbonylamino group, an isobutylcarbonylamino group, a t-butylcarbonylamino group, an n-pentylcarbonylamino group, and an n-hexylcarbonylamino group.

- 15 [0090] The " C_{1-6} alkylcarbonyloxy group" means groups formed by bonding the single " C_{1-6} alkylcarbonyl group" to an oxy group, and specific examples include a methylcarbonyloxy group, an ethylcarbonyloxy group, an n-propylcarbonyloxy group, an isopropylcarbonyloxy group, an n-butylcarbonyloxy group, an isobutylcarbonyloxy group, a t-butylcarbonyloxy group, an n-pentylcarbonyloxy group, and an n-hexylcarbonyloxy group.

- 20 [0091] The "mono- C_{1-6} alkylaminosulfonyl group" means groups formed by bonding the single "mono- C_{1-6} alkylamino group" to a sulfonyl group, and specific examples include a methylaminosulfonyl group, an ethylaminosulfonyl group, an n-propylaminosulfonyl group, an isopropylaminosulfonyl group, an n-butylaminosulfonyl group, an isobutylaminosulfonyl group, a t-butylaminosulfonyl group, an n-pentylaminosulfonyl group, and an n-hexylaminosulfonyl group.

- 25 [0092] The "di- C_{1-6} alkylaminosulfonyl group" means groups formed by bonding the single "di- C_{1-6} alkylamino group" to a sulfonyl group, and specific examples include a

dimethylaminosulfonyl group, a diethylaminosulfonyl group, a di-n-propylaminosulfonyl group, a diisopropylaminosulfonyl group, a di-n-butylaminosulfonyl group, a diisobutylaminosulfonyl group, a di-t-butylaminosulfonyl group, a di-n-pentylaminosulfonyl group, a di-n-hexylaminosulfonyl group, a

5 N-ethyl-N-methylaminosulfonyl group, a N-methyl-N-n-propylaminosulfonyl group, a N-isopropyl-N-methylaminosulfonyl group, a N-n-butyl-N-methylaminosulfonyl group, a N-isobutyl-N-methylaminosulfonyl group, a N-t-butyl-N-methylaminosulfonyl group, a N-methyl-N-n-pentylaminosulfonyl group, a N-n-hexyl-N-methylaminosulfonyl group, a N-ethyl-N-n-propylaminosulfonyl group, a N-ethyl-N-isopropylaminosulfonyl group, a

10 N-n-butyl-N-ethylaminosulfonyl group, a N-ethyl-N-isobutylaminosulfonyl group, a N-t-butyl-N-ethylaminosulfonyl group, a N-ethyl-N-n-pentylaminosulfonyl group, and a N-ethyl-N-n-hexylaminosulfonyl group.

[0093] In the present invention, the binding manner of B will be described.

As obviously shown by Formula (I), in the image on the paper, the left site of the

15 partial formula representing B is bonded to L^1 , and the right site of the partial formula representing B is bonded to L^2 in the present invention.

[0094] Preferred structures for each substituent in the present invention will be described next.

[0095] The substituent R^1 is preferably a hydrogen atom, a halogen atom, a C_{1-6}

20 alkoxy group, a mono- C_{1-6} alkylamino group, a di- C_{1-6} alkylamino group, or a C_{1-6} alkyl group (the C_{1-6} alkoxy group, the mono- C_{1-6} alkylamino group, the di- C_{1-6} alkylamino group, and the C_{1-6} alkyl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^7).

The substituent R^1 is more preferably a hydrogen atom or a C_{1-6} alkoxy group.

25 The substituent R^1 is even more preferably a hydrogen atom.

In another embodiment, the substituent R^1 is more preferably a C_{1-6} alkyl group (the C_{1-6} alkyl group is unsubstituted or substituted with one or more halogen atoms). In the substituent R^1 , the C_{1-6} alkyl group is even more preferably a methyl group.

[0096] Next, preferred structures for the combination of L^1 , L^2 , and B will be

described.

For a preferred combination of L^1 , L^2 , and B,

L^1 and L^2 are single bonds, and B is a C_{4-7} cycloalkylene group or a 4 to 7-membered heterocyclylene group (the C_{4-7} cycloalkylene group and the 4 to 7-membered heterocyclylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^5 , and a single methylene group of the C_{4-7} cycloalkylene group and the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1- C_{3-7} cycloalkylene group);

one of L^1 and L^2 is a single bond, the other of L^1 and L^2 is NR^{2d} (where R^{2d} is a hydrogen atom or a C_{1-6} alkyl group (the C_{1-6} alkyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^7)), O, S, SO, SO_2 , or a C_{1-3} alkylene group (a single methylene group of the C_{1-3} alkylene group is optionally replaced by O, S, SO_2 , C=O, C=S, or NR^{3d} (where R^{3d} has the same definition as R^{2d})), and B is a C_{4-7} cycloalkylene group or a 4 to 7-membered heterocyclylene group (the C_{4-7} cycloalkylene group and the 4 to 7-membered heterocyclylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^5 , and a single methylene group of the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1- C_{3-7} cycloalkylene group);

each of L^1 and L^2 is independently NR^{2e} (where R^{2e} is a hydrogen atom, a C_{1-6} alkyl group (the C_{1-6} alkyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^7)), O, S, SO, SO_2 , or a C_{1-3} alkylene group (a single methylene group of the C_{1-3} alkylene group is optionally replaced by O, S, SO_2 , C=O, C=S, or NR^{3e} (where R^{3e} has the same definition as R^{2e})), and B is a C_{4-7} cycloalkylene group or a 4 to 7-membered heterocyclylene group (the C_{4-7} cycloalkylene group and the 4 to 7-membered heterocyclylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^5 , and a single methylene group of the C_{4-7} cycloalkylene group and the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1- C_{3-7}

cycloalkylene group); or

each of L^1 and L^2 is independently NR^{2f} (where R^{2f} is a hydrogen atom or a C_{1-6} alkyl group (the C_{1-6} alkyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^7)), O, S, SO,
 5 or SO_2 (where L^2 is not NR^{2f} when L^1 is O), and

B is a C_{1-6} alkylene group, (the C_{1-6} alkylene group is unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^5 , and a single methylene group of the C_{1-6} alkylene group is optionally replaced by a 1,1- C_{3-7} cycloalkylene group).

10 [0097] For a more preferred combination of L^1 , L^2 , and B,

L^1 and L^2 are single bonds, and B is a 4 to 7-membered heterocyclylene group (the 4 to 7-membered heterocyclylene group is unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^3);

one of L^1 and L^2 is a single bond, the other of L^1 and L^2 is NR^{2e} (where R^{2e} is a
 15 hydrogen atom or a methyl group) or O, and B is a C_{4-7} cycloalkylene group or a 4 to 7-membered heterocyclylene group (the C_{4-7} cycloalkylene group and the 4 to 7-membered heterocyclylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^3);


each of L^1 and L^2 is independently NR^{2g} (where R^{2g} is a hydrogen atom or a methyl
 20 group), and B is a C_{4-7} cycloalkylene group or a 4 to 7-membered heterocyclylene group (the C_{4-7} cycloalkylene group and the 4 to 7-membered heterocyclylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^3); or

L^1 is NR^{2h} (where R^{2h} is a hydrogen atom or a methyl group), L^2 is NR^{2i} (where R^{2i}
 25 is a hydrogen atom or a methyl group) or an oxygen atom, and B is a C_{1-6} alkylene group (the C_{1-6} alkylene group is unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^3).

[0098] For an even more preferred combination of L^1 , L^2 , and B,


L^1 and L^2 are single bonds, and B is represented by any of Formula (II-1) to Formula

(II-3) shown in Figure (II):




(where m is 0 or 1, and R^b is a substituent selected from the substituent group V^3);

one of L^1 and L^2 is a single bond, the other of L^1 and L^2 is NR^{2j} (where R^{2j} is a
 5 hydrogen atom or a methyl group) or an oxygen atom, and B is represented by any of
 Formula (III-1) to Formula (III-9) shown in Figure (III):



(where m is 0 or 1, and R^b is a substituent selected from the substituent group V^3);

10 each of L^1 and L^2 is independently NR^{2k} (where R^{2k} is a hydrogen atom or a methyl
 group), and B is represented by Formula (IV-1) shown in Figure (IV):



(where m is 0 or 1, and R^b is a substituent selected from the substituent group V^3); or

L^1 is NR^{2l} (where R^{2l} is a hydrogen atom or a methyl group), L^2 is NR^{2m} (where R^{2m}
 15 has the same definition as R^{2k}) or an oxygen atom, and B is an ethylene group.

In another embodiment, for a preferred combination of L^1 , L^2 , and B ,
 L^1 and L^2 are single bonds, and


B is a 4 to 7-membered heterocyclylene group (the 4 to 7-membered heterocyclylene group is substituted with a substituent selected from amino groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, and C₁₋₆ alkylsulfonylamino groups and is optionally substituted with one or more substituents identically or differently selected from the substituent group V³, and a single methylene group of the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group); or L¹ is a single bond, L² is NR^{2c}, O, or a C₁₋₆ alkylene group (a single methylene group of the C₁₋₆ alkylene group is optionally replaced by O or NR^{3c}), and

B is a C₃₋₆ cycloalkylene group, a C₃₋₆ cycloalkenylene group, or a 4 to 7-membered heterocyclylene group (the C₃₋₆ cycloalkylene group, the C₃₋₆ cycloalkenylene group, and the 4 to 7-membered heterocyclylene group are substituted with one or more substituents selected from amino groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, and C₁₋₆ alkylsulfonylamino groups and are optionally substituted with one or more substituents identically or differently selected from the substituent group V³, and a single methylene group of the C₃₋₆ cycloalkylene group, the C₃₋₆ cycloalkenylene group, and the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group),

where each of R^{2c} and R^{3c} is independently a hydrogen atom, a C₁₋₃ alkyl group, or a C₁₋₃ haloalkyl group.

[0099] For a more preferred combination of L¹, L², and B, L¹ and L² are single bonds, and B is a 4 to 7-membered heterocyclylene group (the 4 to 7-membered heterocyclylene group is substituted with a substituent selected from amino groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, and C₁₋₆ alkylsulfonylamino groups and is optionally substituted with one or more substituents identically or differently selected from the substituent group V³).


[0100] For an even more preferred combination of L¹, L², and B, L¹ and L² are single bonds, and B is represented by any of Formula (II-1) to Formula (II-3) shown in Figure (II):



(where m is 1, and R^b is an amino group, a mono- C_{1-6} alkylamino group, a di- C_{1-6} alkylamino group, or a C_{1-6} alkylsulfonylamino group).


[0101] A is preferably a 3 to 11-membered heterocyclyl group, a C_{3-11} cycloalkyl group, a C_{6-14} aryl group, or a 5 to 10-membered heteroaryl group (the 3 to 11-membered heterocyclyl group, the C_{3-11} cycloalkyl group, the C_{6-14} aryl group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^5). A is more preferably a 4 to 7-membered heterocyclyl group, a phenyl group, or a 5 to 6-membered heteroaryl group (the 4 to 7-membered heterocyclyl group, the phenyl group, and the 5 to 6-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^3). A is more preferably a phenyl group or a 5 to 6-membered heteroaryl group (the phenyl group and the 5 to 6-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^3).

A is even more preferably represented by any of Formula (V-1) to Formula (V-3) shown in Figure (V):



(where l is 0 to 3, R^a is a substituent selected from the substituent group V^3 , and R^a may be the same as or different from each other when l is 2 or 3).

[0102] In another embodiment, A is even more preferably represented by any of Formula (V-I-1) to Formula (V-I-3) shown in Figure (V-I):



(where l is 0 to 3, R^a is a substituent selected from the substituent group V^3 , and R^a may be the same as or different from each other when l is 2 or 3).

[0103] L^3 is preferably a C_{1-3} alkylene group (the C_{1-3} alkylene group is
 5 unsubstituted or substituted with one or more substituents identically or differently
 selected from the substituent group V^3 , and a single methylene group of the C_{1-3} alkylene
 group is optionally replaced by $C=O$ or $C=S$).


L^3 is more preferably a methylene group.

[0104] D is preferably a 3 to 11-membered heterocyclyl group, a C_{3-11} cycloalkyl
 10 group, a C_{6-14} aryl group, or a 5 to 10-membered heteroaryl group (the 3 to 11-membered
 heterocyclyl group, the C_{3-11} cycloalkyl group, the C_{6-14} aryl group, and the 5 to
 10-membered heteroaryl group are unsubstituted or substituted with one or more
 substituents identically or differently selected from the substituent group V^5).

D is more preferably a 4 to 7-membered heterocyclyl group, a phenyl group, or a 5
 15 to 10-membered heteroaryl group (the 4 to 7-membered heterocyclyl group, the phenyl
 group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with
 one or more substituents identically or differently selected from the substituent group V^4).

D is even more preferably a phenyl group or a 5 to 10-membered heteroaryl group
 (the phenyl group and the 5 to 10-membered heteroaryl group are unsubstituted or
 20 substituted with one or more substituents identically or differently selected from the
 substituent group V^3).

D is particularly preferably represented by any of Formula (VI-1) to Formula (VI-9)
 shown in Figure (VI):




(where n is 0 to 3, R^c is a substituent selected from the substituent group V^3 , and R^c 's may be the same as or different from each other when n is 2 or 3).

In another embodiment, D is preferably a 3 to 11-membered heterocyclyl group, a C_{6-14} aryl group, or a 5 to 10-membered heteroaryl group (the 3 to 11-membered heterocyclyl group, the C_{6-14} aryl group, and the 5 to 10-membered heteroaryl group are substituted with one or more C_{1-6} alkylsulfonylamino groups or one or more C_{1-6} alkylsulfonyloxy groups (the C_{1-6} alkylsulfonylamino group and the C_{1-6} alkylsulfonyloxy group are unsubstituted or substituted with one or more halogen atoms, one or more nitro groups, one or more C_{1-6} alkoxy groups, or one or more C_{1-6} haloalkoxy groups) and are optionally substituted with one or more substituents identically or differently selected from the substituent group V^5).

D is more preferably a 4 to 7-membered heterocyclyl group, a phenyl group, or a 5 to 10-membered heteroaryl group (the 4 to 7-membered heterocyclyl group, the phenyl group, and the 5 to 10-membered heteroaryl group are substituted with one or more C_{1-6} alkylsulfonylamino groups or one or more C_{1-6} alkylsulfonyloxy groups (the C_{1-6} alkylsulfonylamino group and the C_{1-6} alkylsulfonyloxy group are unsubstituted or substituted with one or more halogen atoms, one or more nitro groups, one or more C_{1-6} alkoxy groups, or one or more C_{1-6} haloalkoxy groups) and are optionally substituted with one or more substituents identically or differently selected from the substituent group V^4).

D is even more preferably a 5 to 6-membered heteroaryl group (the 5 to 6-membered heteroaryl group is substituted with one or more C_{1-6} alkylsulfonyloxy groups (the C_{1-6} alkylsulfonyloxy group is unsubstituted or substituted with one or more halogen atoms)).


D is particularly preferably represented by Formula (VI-1) shown in Figure (VI):



(where n is 1, and R^c is a C_{1-6} alkylsulfonyloxy group (the C_{1-6} alkylsulfonyloxy group is substituted with one or more halogen atoms)).

5 [0105] Compounds preferably used for the T-type calcium channel inhibitor and for the preventive agent, the therapeutic agent, and/or the improving agent for a disease treatable by a T-type calcium channel inhibitory action of the present invention are shown below.

[0106] 1) A compound of Formula (I)



[wherein

10 R^1 is a hydrogen atom, a halogen atom, a C_{1-6} alkoxy group, a mono- C_{1-6} alkylamino group, a di- C_{1-6} alkylamino group, or a C_{1-6} alkyl group (the C_{1-6} alkoxy group, the mono- C_{1-6} alkylamino group, the di- C_{1-6} alkylamino group, and the C_{1-6} alkyl group are unsubstituted or substituted with one or more substituents identically or differently
15 selected from the substituent group V^7);

each of L^1 and L^2 independently is any of a single bond, NR^{2n} (where R^{2n} is a hydrogen atom, a C_{1-6} alkyl group, or a C_{3-11} cycloalkyl group (the C_{1-6} alkyl group and the C_{3-11} cycloalkyl group are unsubstituted or substituted with one or more substituents
20 identically or differently selected from the substituent group V^7)), O, S, SO, SO_2 , or a C_{1-6} alkylene group (the C_{1-6} alkylene group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^3 , and a single methylene group of the C_{1-6} alkylene group is optionally replaced by O, S, SO_2 , C=O, C=S, or NR^{3n} (where the R^{3n} has the same definition as R^{2n}));

25 each of L^1 and L^2 is independently a single bond, NR^{20} (where R^{20} has the same definition as R^{2n}), O, S, SO, SO_2 , or a C_{1-6} alkylene group, and B is a C_{3-11} cycloalkylene

group, a C₃₋₁₁ cycloalkenylene group, a 3 to 11-membered heterocyclylene group, or a 5 to 10-membered heteroarylene group (the C₃₋₁₁ cycloalkylene group, the C₃₋₁₁ cycloalkenylene group, the 3 to 11-membered heterocyclylene group, and the 5 to 10-membered heteroarylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V⁸, and a single methylene group of the C₃₋₁₁ cycloalkylene group and the C₃₋₁₁ cycloalkenylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group); or

L¹ is NR^{2p} (where R^{2p} has the same definition as R²ⁿ), S, SO, or SO₂,

L² is NR^{2q} (where R^{2q} has the same definition as R²ⁿ), O, S, SO, or SO₂, and B is a C₁₋₆ alkylene group, a C₂₋₆ alkenylene group, or a C₂₋₆ alkynylene group (the C₁₋₆ alkylene group, the C₂₋₆ alkenylene group, and the C₂₋₆ alkynylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V⁵, and a single methylene group of the C₁₋₆ alkylene group, the C₂₋₆ alkenylene group, and the C₂₋₆ alkynylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group);

L³ is a C₁₋₆ alkylene group (the C₁₋₆ alkylene group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁵, and a single methylene group of the C₁₋₆ alkylene group is optionally replaced by C=O or C=S);

A is a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group (the C₁₋₆ alkyl group and the C₂₋₆ alkenyl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁷), a C₃₋₁₁ cycloalkyl group, a 3 to 11-membered heterocyclyl group, a C₆₋₁₄ aryl group, or a 5 to 10-membered heteroaryl group (the C₃₋₁₁ cycloalkyl group, the 3 to 11-membered heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁵);

D is a C₃₋₁₁ cycloalkyl group, a C₃₋₁₁ cycloalkenyl group, a 3 to 11-membered heterocyclyl group, a C₆₋₁₄ aryl group, or a 5 to 10-membered heteroaryl group (the C₃₋₁₁ cycloalkyl group, the C₃₋₁₁ cycloalkenyl group, the 3 to 11-membered heterocyclyl group,

the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁸),

a tautomer of the compound, a pharmaceutically acceptable salt thereof, or a solvate thereof.

[0107] 2) The compound according to 1), wherein L³ is a methylene group, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof..

[0108] 3) The compound according to 1) or 2), wherein R¹ is a hydrogen atom or a C₁₋₃ alkoxy group (the C₁₋₃ alkoxy group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁷), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0109] 4) The compound according to 3), wherein R¹ is a hydrogen atom, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

5) The compound according to 1) or 2), wherein R¹ is a C₁₋₆ alkyl group (the C₁₋₆ alkyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁹), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

6) The compound according to 5), wherein R¹ is a C₁₋₃ alkyl group (the C₁₋₃ alkyl group is unsubstituted or substituted with one or more halogen atoms), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

7) The compound according to 6), wherein R¹ is a methyl group, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.


[0110] 8) The compound according to any one of 1) to 7), wherein D is a 3 to 11-membered heterocyclyl group, a C₆₋₁₄ aryl group, or a 5 to 10-membered heteroaryl group (the 3 to 11-membered heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with one or more

substituents identically or differently selected from the substituent group V^8), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0111] 9) The compound according to 8), wherein D is a 4 to 7-membered heterocyclyl group, a phenyl group, or a 5 to 10-membered heteroaryl group (the 4 to 7-membered heterocyclyl group, the phenyl group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^4), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0112] 10) The compound according to 9), wherein D is a phenyl group or a 5 to 6-membered heteroaryl group (the phenyl group and the 5 to 6-membered heteroaryl group are substituted with one or more substituents identically or differently selected from the substituent group V^4), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0113] 11) The compound according to any one of 1) to 9), wherein D has any of the structures of Formulae (VI), n is 0 to 3, R^c is a substituent selected from the substituent group V^3 , and R^c may be the same as or different from each other when n is 2 or 3, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.



[0114] 12) The compound according to 11), wherein n is 1 or 2, R^c is a substituent selected from the substituent group V^3 , and R^c may be the same as or different from each other when n is 2, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.


13) The compound according to 8), wherein D is a 3 to 11-membered heterocyclyl group, a C₆₋₁₄ aryl group, or a 5 to 10-membered heteroaryl group (the 3 to 11-membered heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are substituted with one or more C₁₋₆ alkylsulfonylamino groups or one or more C₁₋₆ alkylsulfonyloxy groups (the C₁₋₆ alkylsulfonylamino group and the C₁₋₆ alkylsulfonyloxy group are unsubstituted or substituted with one or more halogen atoms, one or more nitro groups, one or more C₁₋₆ alkoxy groups, or one or more C₁₋₆ haloalkoxy groups) and are optionally substituted with one or more substituents identically or differently selected from the substituent group V⁵), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0115] 14) The compound according to 13), wherein D is a 4 to 7-membered heterocyclyl group, a phenyl group, or a 5 to 10-membered heteroaryl group (the 4 to 7-membered heterocyclyl group, the phenyl group, and the 5 to 10-membered heteroaryl group are substituted with one or more C₁₋₆ alkylsulfonylamino groups or one or more C₁₋₆ alkylsulfonyloxy groups (the C₁₋₆ alkylsulfonylamino group and the C₁₋₆ alkylsulfonyloxy group are unsubstituted or substituted with one or more halogen atoms, one or more nitro groups, one or more C₁₋₆ alkoxy groups, or one or more C₁₋₆ haloalkoxy groups) and are optionally substituted with one or more substituents identically or differently selected from the substituent group V⁵), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0116] 15) The compound according to 14), wherein D is a 5 to 6-membered heteroaryl group (the 5 to 6-membered heteroaryl group is substituted with one or more C₁₋₆ alkylsulfonyloxy groups (the C₁₋₆ alkylsulfonyloxy group is unsubstituted or substituted with one or more halogen atoms) and is optionally substituted with one or more substituents identically or differently selected from the substituent group V⁵), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0117] 16) The compound according to 15), wherein D has the structure of Formula (VI-I), n is 1, and R^c is a C₁₋₆ alkylsulfonyloxy group (the C₁₋₆ alkylsulfonyloxy

group is substituted with one or more halogen atoms), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.



[0118] 17) The compound according to any one of 1) to 16), wherein A is a 3 to 11-membered heterocyclyl group, a C₃₋₁₁ cycloalkyl group, a C₆₋₁₄ aryl group, or a 5 to 10-membered heteroaryl group (the 3 to 11-membered heterocyclyl group, the C₃₋₁₁ cycloalkyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁵), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0119] 18) The compound according to 17), wherein A is a 4 to 7-membered heterocyclyl group, a phenyl group, or a 5 to 6-membered heteroaryl group (the 4 to 7-membered heterocyclyl group, the phenyl group, and the 5 to 6-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁴), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.


[0120] 19) The compound according to 18), wherein A is a phenyl group (the phenyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁴), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0121] 20) The compound according to 18), wherein A is a 5 to 6-membered heteroaryl group (the 5 to 6-membered heteroaryl group is substituted with one or more substituents identically or differently selected from the substituent group V⁴), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0122] 21) The compound according to any one of 1) to 16), wherein A is a C₁₋₆ alkyl group, a C₃₋₁₁ cycloalkyl group, or a C₂₋₆ alkenyl group (the C₁₋₆ alkyl group, the C₃₋₁₁ cycloalkyl group, and the C₂₋₆ alkenyl group are unsubstituted or substituted with


one or more substituents identically or differently selected from the substituent group V^7), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0123] 22) The compound according to any one of 1) to 18), wherein A has any of the structures of Formulae (V), l is 0 to 2, R^a is a substituent selected from the substituent group V^3 , and R^a 's may be the same as or different from each other when n is 2, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.



[0124] 23) The compound according to 22), wherein l is 1 or 2, R^a is a substituent selected from the substituent group V^3 , and R^a may be the same as or different from each other when n is 2, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0125] 24) The compound according to any one of 1) to 18), wherein A has any of the structures of Formulae (V-I), l is 0 to 2, R^a is a substituent selected from the substituent group V^3 , and R^a may be the same as or different from each other when n is 2, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.




25) The compound according to any one of 1) to 24), wherein L^1 and L^2 are single bonds, B is a C_{3-11} cycloalkylene group, a C_{3-11} cycloalkenylene group, a 3 to 11-membered heterocyclylene group, a C_{6-14} arylene group, or a 5 to 10-membered heteroarylene group (the C_{3-11} cycloalkylene group, the C_{3-11} cycloalkenylene group, the 3 to 11-membered heterocyclylene group, the C_{6-14} arylene group, and the 5 to 10-membered heteroarylene group are unsubstituted or optionally substituted with one or more substituents identically

or differently selected from the substituent group V^5 , and a single methylene group of the C_{3-11} cycloalkylene group and the C_{3-11} cycloalkenylene group is optionally replaced by a 1,1- C_{3-7} cycloalkylene group), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0126] 26) The compound according to 25), wherein B is a 4 to 7-membered heterocyclylene group (the 4 to 7-membered heterocyclylene group is unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^4), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0127] 27) The compound according to any one of 1) to 26), wherein B has any of the structures of Formulae (II), m is 0 to 3, R^b is a substituent selected from the substituent group V^3 , and R^b may be the same as or different from each other when m is 2 or 3, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.



15

[0128] 28) The compound according to 27), wherein m is 0 or 1, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.


[0129] 29) The compound according to any one of 1) to 24), wherein L^1 and L^2 are single bonds, B is a C_{3-11} cycloalkylene group, a C_{3-11} cycloalkenylene group, a 3 to 11-membered heterocyclylene group, or a 5 to 10-membered heteroarylene group (the C_{3-11} cycloalkylene group, the C_{3-11} cycloalkenylene group, the 3 to 11-membered heterocyclylene group, and the 5 to 10-membered heteroarylene group are substituted with one or more substituents selected from amino groups, mono- C_{1-6} alkylamino groups, di- C_{1-6} alkylamino groups, and C_{1-6} alkylsulfonylamino groups and are optionally substituted with one or more substituents identically or differently selected from the substituent group V^3 , and a single methylene group of the 4 to 7-membered

25

heterocyclylene group is optionally replaced by a 1,1- C_{3-7} cycloalkylene group), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0130] 30) The compound according to 29), wherein B is a 4 to 7-membered heterocyclylene group (the 4 to 7-membered heterocyclylene group is substituted with one or more substituents selected from amino groups, mono- C_{1-6} alkylamino groups, di- C_{1-6} alkylamino groups, and C_{1-6} alkylsulfonylamino groups and is optionally substituted with one or more substituents identically or differently selected from the substituent group V^3), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0131] 31) The compound, according to 30), wherein B has any of the structures of Formulae (II), m is 1, R^b is an amino group, a mono- C_{1-6} alkylamino group, a di- C_{1-6} alkylamino group, or a C_{1-6} alkylsulfonylamino group, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.



32) The compound, according to any one of 1) to 24), wherein

L^1 is a single bond,

L^2 is NR^{2r} (where R^{2r} is a hydrogen atom or a C_{1-6} alkyl group (the C_{1-6} alkyl group is unsubstituted or substituted with one or more substituents identically or differently

selected from the substituent group V^7)), O, S, SO, SO_2 , or a C_{1-6} alkylene group (the C_{1-6} alkylene group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^5 , and a single methylene group of the C_{1-6} alkylene group is optionally replaced by O, S, SO_2 , $C=O$, $C=S$, or NR^{3r} (where R^{3r} has the same definition as R^{2r})),

B is a C_{3-11} cycloalkylene group, a C_{3-11} cycloalkenylene group, or a 3 to 11-membered heterocyclylene group (the C_{3-11} cycloalkylene group, the C_{3-11} cycloalkenylene group,

and the 3 to 11-membered heterocyclylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^5 , and a single methylene group of the C_{3-11} cycloalkylene group and the C_{3-11} cycloalkenylene group is optionally replaced by a 1,1- C_{3-7} cycloalkylene group),
 5 the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0132] 33) The compound according to 32), wherein

L^2 is NR^{2s} (where R^{2s} is a hydrogen atom or a methyl group), O, or a C_{1-6} alkylene group (the C_{1-6} alkylene group is unsubstituted or substituted with one or more substituents
 10 identically or differently selected from the substituent group V^7 , and a single methylene group of the C_{1-6} alkylene group is optionally replaced by O, S, SO_2 , $C=O$, $C=S$, or NR^{3s} (where R^{3s} has the same definition as R^{2s})), and

B is a 4 to 7-membered heterocyclylene group (the 4 to 7-membered heterocyclylene group is unsubstituted or optionally substituted with one or more substituents identically
 15 or differently selected from the substituent group V^4), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.


[0133] 34) The compound according to 32), wherein

L^2 is NR^{2t} (where R^{2t} is a hydrogen atom or a methyl group), O, or a C_{1-6} alkylene group (the C_{1-6} alkylene group is unsubstituted or substituted with one or more substituents
 20 identically or differently selected from the substituent group V^7 , and a single methylene group of the C_{1-6} alkylene group is optionally replaced by O, S, SO_2 , $C=O$, $C=S$, or NR^{3t} (where R^{3t} has the same definition as R^{2t})), and

B is a C_{4-7} cycloalkylene group or a C_{4-7} cycloalkenylene group (the C_{4-7} cycloalkylene group and the C_{4-7} cycloalkenylene group are unsubstituted or optionally substituted with
 25 one or more substituents identically or differently selected from the substituent group V^4), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0134] 35) The compound according to any one of 1) to 24), wherein B has any of the structures of Formulae (VII), m is 0 to 3, R^b is a substituent selected from the

substituent group V^3 , and R^b 's may be the same as or different from each other when m is 2 or 3, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.



5 [0135] 36) The compound according to 35), wherein m is 0 or 1, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0136] 37) The compound according to any one of 1) to 24), wherein L^1 is NR^{2u} (where R^{2u} is a hydrogen atom or a C_{1-6} alkyl group (the C_{1-6} alkyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the
10 substituent group V^7)), O, S, SO, SO₂, or a C_{1-6} alkylene group (the C_{1-6} alkylene group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^5 , and a single methylene group of the C_{1-6} alkylene group is optionally replaced by O, S, SO₂, C=O, C=S, or NR^{3u} (where R^{3u} has the same definition as R^{2u})),

15 L^2 is a single bond, and


B is a C_{3-11} cycloalkylene group, a C_{3-11} cycloalkenylene group, or a 3 to 11-membered heterocyclylene group (the C_{3-11} cycloalkylene group, the C_{3-11} cycloalkenylene group, and the 3 to 11-membered heterocyclylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the
20 substituent group V^5 , and a single methylene group of the C_{3-11} cycloalkylene group and the C_{3-11} cycloalkenylene group is optionally replaced by a 1,1- C_{3-7} cycloalkylene group), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0137] 38) The compound according to 37), wherein

L^1 is NR^{2v} (where R^{2v} is a hydrogen atom or a methyl group), O, or a C_{1-6} alkylene group (the C_{1-6} alkylene group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^7 , and a single methylene group of the C_{1-6} alkylene group is optionally replaced by O, S, SO_2 , $C=O$, $C=S$, or NR^{3v} (where R^{3v} has the same definition as R^{2v})), and

B is a 4 to 7-membered heterocyclylene group (the 4 to 7-membered heterocyclylene group is unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^4), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

- 10 [0138] 39) The compound according to 37), wherein B has any of the structures of Formulae (VIII), m is 0 to 3, R^b is a substituent selected from the substituent group V^3 , and R^b may be the same as or different from each other when m is 2 or 3, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.



- 15 [0139] 40) The compound according to 39), wherein m is 0 or 1, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.


- [0140] 41) The compound according to any one of 1) to 24), wherein each of L^1 and L^2 is independently NR^{2w} (where R^w is a hydrogen atom or a C_{1-6} alkyl group (the C_{1-6} alkyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^7)), O, S, SO, SO_2 , or a C_{1-6} alkylene group (the C_{1-6} alkylene group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^5 , and a single methylene group of the C_{1-6} alkylene group is optionally replaced by O, S, SO_2 , $C=O$,
- 20

C=S, or NR^{3w} (where R^{3w} has the same definition as R^{2w})), and

B is a C_{3-11} cycloalkylene group, a C_{3-11} cycloalkenylene group, a 3 to 11-membered heterocyclylene group, a C_{6-14} arylene group, or a 5 to 10-membered heteroarylene group (the C_{3-11} cycloalkylene group, the C_{3-11} cycloalkenylene group, the 3 to 11-membered heterocyclylene group, the C_{6-14} arylene group, and the 5 to 10-membered heteroarylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^5 , and a single methylene group of the C_{3-11} cycloalkylene group and the C_{3-11} cycloalkenylene group is optionally replaced by a 1,1- C_{3-7} cycloalkylene group), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0141] 42) The compound according to 41), wherein each of L^1 and L^2 is independently NR^{2x} (where R^{2x} is a hydrogen atom or a methyl group), O, or a C_{1-6} alkylene group (the C_{1-6} alkylene group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^4 , and a single methylene group of the C_{1-6} alkylene group is optionally replaced by O, S, SO_2 , C=O, C=S, or NR^{3x} (where R^{3x} has the same definition as R^{2x})), and B is a C_{3-11} cycloalkylene group, a C_{3-11} cycloalkenylene group, or a 3 to 11-membered heterocyclylene group (the C_{3-11} cycloalkylene group, the C_{3-11} cycloalkenylene group, and the 3 to 11-membered heterocyclylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^4 , and a single methylene group of the C_{3-11} cycloalkylene group and the C_{3-11} cycloalkenylene group is optionally replaced by a 1,1- C_{3-7} cycloalkylene group), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0142] 43) The compound according to 41), wherein B has the structure of Formula (IV), m is 0 to 3, R^b is a substituent selected from the substituent group V^3 , and R^b may be the same as or different from each other when m is 2 or 3, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.



[0143] 44) The compound according to 43), wherein m is 0 or 1, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0144] 45) The compound according to any one of 1) to 24), wherein

5 L^1 is NR^{2x} (where R^{2x} is a hydrogen atom, a C_{1-6} alkyl group (the C_{1-6} alkyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^7)), S, SO, or SO_2 ,

L^2 is NR^{2y} (where R^{2y} has the same definition as R^{2x}), O, S, SO, or SO_2 , and

10 B is a C_{1-6} alkylene group, a C_{2-6} alkenylene group, or a C_{2-6} alkynylene group (the C_{1-6} alkylene group, the C_{2-6} alkenylene group, and the C_{2-6} alkynylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^5), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0145] 46) The compound according to 45), wherein


15 L^1 is NR^{2z} (where R^{2z} is a hydrogen atom or a methyl group),

L^2 is NR^{2aa} (where R^{2aa} has the same definition as R^z) or O, and

20 B is a C_{1-6} alkylene group (the C_{1-6} alkylene group is unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^4), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0146] 47) The compound according to 46), wherein B is an ethylene group, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0147] 48) A compound of Formula (I):



wherein


R^1 is a hydrogen atom, L^1 and L^2 are single bonds, L^3 is a methylene group, and A, B, and D are a combination listed in Table 1 below,

a tautomer of the compound, a pharmaceutically acceptable salt thereof, or a solvate

5 thereof.

The symbols in Table 1 are the substituents below.

[0148]




[0149]

10 (Table 1)

A	B	D	A	B	D	A	B	D
A 1	B 1	D 1	A 1	B 1	D 2	A 1	B 1	D 3
A 1	B 1	D 4	A 1	B 1	D 5	A 1	B 1	D 6
A 1	B 1	D 7	A 1	B 1	D 8	A 1	B 1	D 9
A 1	B 1	D 1 0	A 1	B 1	D 1 1	A 1	B 1	D 1 2
A 1	B 1	D 1 3	A 1	B 1	D 1 4	A 1	B 1	D 1 5
A 1	B 1	D 1 6	A 2	B 1	D 1	A 2	B 1	D 2
A 2	B 1	D 3	A 2	B 1	D 4	A 2	B 1	D 5
A 2	B 1	D 6	A 2	B 1	D 7	A 2	B 1	D 8
A 2	B 1	D 9	A 2	B 1	D 1 0	A 2	B 1	D 1 1
A 2	B 1	D 1 2	A 2	B 1	D 1 3	A 2	B 1	D 1 4
A 2	B 1	D 1 5	A 2	B 1	D 1 6	A 3	B 1	D 1
A 3	B 1	D 2	A 3	B 1	D 3	A 3	B 1	D 4
A 3	B 1	D 5	A 3	B 1	D 6	A 3	B 1	D 7
A 3	B 1	D 8	A 3	B 1	D 9	A 3	B 1	D 1 0
A 3	B 1	D 1 1	A 3	B 1	D 1 2	A 3	B 1	D 1 3
A 3	B 1	D 1 4	A 1	B 2	D 1 5	A 1	B 2	D 1 6
A 1	B 2	D 1	A 1	B 2	D 2	A 1	B 2	D 3
A 1	B 2	D 4	A 1	B 2	D 5	A 1	B 2	D 6
A 1	B 2	D 7	A 1	B 2	D 8	A 1	B 2	D 9
A 1	B 2	D 1 0	A 1	B 2	D 1 1	A 1	B 2	D 1 2
A 1	B 2	D 1 3	A 1	B 2	D 1 4	A 1	B 2	D 1 5
A 1	B 2	D 1 6	A 2	B 2	D 1	A 2	B 2	D 2
A 2	B 2	D 3	A 2	B 2	D 4	A 2	B 2	D 5
A 2	B 2	D 6	A 2	B 2	D 7	A 2	B 2	D 8
A 2	B 2	D 9	A 2	B 2	D 1 0	A 2	B 2	D 1 1
A 2	B 2	D 1 2	A 2	B 2	D 1 3	A 2	B 2	D 1 4
A 2	B 2	D 1 5	A 2	B 2	D 1 6	A 3	B 2	D 1
A 3	B 2	D 2	A 3	B 2	D 3	A 3	B 2	D 4
A 3	B 2	D 5	A 3	B 2	D 6	A 3	B 2	D 7
A 3	B 2	D 8	A 3	B 2	D 9	A 3	B 2	D 1 0
A 3	B 2	D 1 1	A 3	B 2	D 1 2	A 3	B 2	D 1 3
A 1	B 3	D 1	A 1	B 3	D 2	A 1	B 3	D 3
A 1	B 3	D 4	A 1	B 3	D 5	A 1	B 3	D 6
A 1	B 3	D 7	A 1	B 3	D 8	A 1	B 3	D 9
A 1	B 3	D 1 0	A 1	B 3	D 1 1	A 1	B 3	D 1 2
A 1	B 3	D 1 3	A 1	B 3	D 1 4	A 2	B 3	D 1 5
A 1	B 3	D 1 6	A 2	B 3	D 1	A 2	B 3	D 2
A 2	B 3	D 3	A 2	B 3	D 4	A 2	B 3	D 5
A 2	B 3	D 6	A 2	B 3	D 7	A 2	B 3	D 8
A 2	B 3	D 9	A 2	B 3	D 1 0	A 2	B 3	D 1 1
A 2	B 3	D 1 2	A 2	B 3	D 1 3	A 2	B 3	D 1 4
A 2	B 3	D 1 5	A 2	B 3	D 1 6	A 3	B 3	D 1
A 3	B 3	D 2	A 3	B 3	D 3	A 3	B 3	D 4
A 3	B 3	D 5	A 3	B 3	D 6	A 3	B 3	D 7
A 3	B 3	D 8	A 3	B 3	D 9	A 3	B 3	D 1 0
A 3	B 3	D 1 1	A 3	B 3	D 1 2	A 3	B 3	D 1 3
A 3	B 3	D 1 4	A 3	B 3	D 1 5	A 3	B 3	D 1 6

[0150] 49) The compound according to Formula (I), wherein

R^1 is a hydrogen atom, L^1 and L^2 are single bonds, L^3 is a methylene group, and A, B, and D are a combination listed in Table 1 (where A1 to A3, B1 to B3, and D1 to D16 in Table are the substituents below, in 49)), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.




5

[0151] 50) The compound according to Formula (I), wherein

R^1 is a hydrogen atom, L^1 and L^2 are single bonds, L^3 is a methylene group, and A, B, and D are a combination listed in Table 1 (where A1 to A3, B1 to B3, and D1 to D16 in Table are the substituents below, in 50)), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

10

[0152]




[0153] 51) The compound according to Formula (I), wherein

R^1 is a hydrogen atom, L^1 and L^2 are single bonds, L^3 is a methylene group, and

A, B, and D are a combination listed in Table 1 (where A1 to A3, B1 to B3, and D1 to

5 D16 in Table are the substituents below, in 51)), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.




[0154] 52) The compound according to Formula (I), wherein

R^1 is a hydrogen atom, L^1 and L^2 are single bonds, L^3 is a methylene group, and

A, B, and D are a combination listed in Table 1 (where A1 to A3, B1 to B3, and D1 to

5 D16 in Table are the substituents below, in 52)), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.




[0155] 53) The compound according to Formula (I), wherein

R^1 is a hydrogen atom, L^1 and L^2 are single bonds, L^3 is a methylene group, and

A, B, and D are a combination listed in Table 2 below, the tautomer of the compound, the

5 pharmaceutically acceptable salt thereof, or the solvate thereof.



(Table 2)

A	B	D	A	B	D	A	B	D
A 1	B 1	D 1	A 1	B 1	D 2	A 1	B 1	D 3
A 1	B 1	D 4	A 1	B 1	D 5	A 1	B 1	D 6
A 1	B 1	D 7	A 1	B 1	D 8	A 2	B 1	D 1
A 2	B 1	D 2	A 2	B 1	D 3	A 2	B 1	D 4
A 2	B 1	D 5	A 2	B 1	D 6	A 2	B 1	D 7
A 2	B 1	D 8	A 3	B 1	D 1	A 3	B 1	D 2
A 3	B 1	D 3	A 3	B 1	D 4	A 3	B 1	D 5
A 3	B 1	D 6	A 3	B 1	D 7	A 3	B 1	D 8
A 4	B 1	D 1	A 4	B 1	D 2	A 4	B 1	D 3
A 4	B 1	D 4	A 4	B 1	D 5	A 4	B 1	D 6
A 4	B 1	D 7	A 4	B 1	D 8	A 5	B 1	D 1
A 5	B 1	D 2	A 5	B 1	D 3	A 5	B 1	D 4
A 5	B 1	D 5	A 5	B 1	D 6	A 5	B 1	D 7
A 5	B 1	D 8	A 6	B 1	D 1	A 6	B 1	D 2
A 6	B 1	D 3	A 6	B 1	D 4	A 6	B 1	D 5
A 6	B 1	D 6	A 6	B 1	D 7	A 6	B 1	D 8
A 1	B 2	D 1	A 1	B 2	D 2	A 1	B 2	D 3
A 1	B 2	D 4	A 1	B 2	D 5	A 1	B 2	D 6
A 1	B 2	D 7	A 1	B 2	D 8	A 2	B 2	D 1
A 2	B 2	D 2	A 2	B 2	D 3	A 2	B 2	D 4
A 2	B 2	D 5	A 2	B 2	D 6	A 2	B 2	D 7
A 2	B 2	D 8	A 3	B 2	D 1	A 3	B 2	D 2
A 3	B 2	D 3	A 3	B 2	D 4	A 3	B 2	D 5


A3	B1	D6	A3	B2	D7	A3	B2	D8
A4	B2	D1	A4	B2	D2	A4	B2	D3
A4	B2	D4	A4	B2	D5	A4	B2	D6
A4	B2	D7	A4	B2	D8	A5	B2	D1
A5	B2	D2	A5	B2	D3	A5	B2	D4
A5	B2	D5	A5	B2	D6	A5	B2	D7
A5	B2	D8	A6	B2	D1	A6	B2	D2
A6	B2	D3	A6	B2	D4	A6	B2	D5
A6	B2	D6	A6	B2	D7	A6	B2	D8
A1	B3	D1	A1	B3	D2	A1	B3	D3
A1	B3	D4	A1	B3	D5	A1	B3	D6
A1	B3	D7	A1	B3	D8	A2	B3	D1
A2	B3	D2	A2	B3	D3	A2	B3	D4
A2	B3	D5	A2	B3	D6	A2	B3	D7
A2	B3	D8	A3	B3	D1	A3	B3	D2
A3	B3	D3	A3	B3	D4	A3	B3	D5
A3	B3	D6	A3	B3	D7	A3	B3	D8
A4	B3	D1	A4	B3	D2	A4	B3	D3
A4	B3	D4	A4	B3	D5	A4	B3	D6
A4	B3	D7	A4	B3	D8	A5	B3	D1
A5	B3	D2	A5	B3	D3	A5	B3	D2
A5	B3	D5	A5	B3	D6	A5	B3	D7
A5	B3	D8	A6	B3	D1	A6	B3	D2
A6	B3	D3	A6	B3	D4	A6	B3	D5
A6	B3	D6	A6	B3	D7	A6	B3	D8
A1	B4	D1	A1	B4	D2	A1	B4	D3
A1	B4	D4	A1	B4	D5	A1	B4	D6
A1	B4	D7	A1	B4	D8	A2	B4	D1
A2	B4	D2	A2	B4	D3	A2	B4	D4
A2	B4	D5	A2	B4	D6	A2	B4	D7
A2	B4	D8	A3	B4	D1	A3	B4	D2
A3	B4	D3	A3	B4	D4	A3	B4	D5
A3	B4	D6	A3	B4	D7	A3	B4	D8
A4	B4	D1	A4	B4	D2	A4	B4	D3
A4	B4	D4	A4	B4	D5	A4	B4	D6
A4	B4	D7	A4	B4	D8	A5	B4	D1
A5	B4	D2	A5	B4	D3	A5	B4	D4
A5	B4	D5	A5	B4	D6	A5	B4	D7
A5	B4	D8	A6	B4	D1	A6	B4	D2
A6	B4	D3	A6	B4	D4	A6	B4	D5
A6	B4	D6	A6	B4	D7	A6	B4	D8
A1	B5	D1	A1	B5	D2	A1	B5	D3
A1	B5	D4	A1	B5	D5	A1	B5	D6
A1	B5	D7	A1	B5	D8	A2	B5	D1
A2	B5	D2	A2	B5	D3	A2	B5	D4
A2	B5	D5	A2	B5	D6	A2	B5	D7
A2	B5	D8	A3	B5	D1	A3	B5	D2
A3	B5	D3	A3	B5	D4	A3	B5	D5
A3	B5	D6	A3	B5	D7	A3	B5	D8
A4	B5	D1	A4	B5	D2	A4	B5	D3

A 4	B 5	D 4	A 4	B 5	D 5	A 4	B 5	D 6
A 4	B 5	D 7	A 4	B 5	D 8	A 5	B 5	D 1
A 5	B 5	D 2	A 5	B 5	D 3	A 5	B 5	D 4
A 5	B 5	D 5	A 5	B 5	D 6	A 5	B 5	D 7
A 5	B 5	D 8	A 6	B 5	D 1	A 6	B 5	D 2
A 6	B 5	D 3	A 6	B 5	D 4	A 6	B 5	D 5
A 6	B 5	D 6	A 6	B 5	D 7	A 6	B 5	D 8
A 1	B 6	D 1	A 1	B 6	D 2	A 1	B 6	D 3
A 1	B 6	D 4	A 1	B 6	D 5	A 1	B 6	D 6
A 1	B 6	D 7	A 1	B 6	D 8	A 2	B 6	D 1
A 2	B 6	D 2	A 2	B 6	D 3	A 2	B 6	D 4
A 2	B 6	D 5	A 2	B 6	D 6	A 2	B 6	D 7
A 2	B 6	D 8	A 3	B 6	D 1	A 3	B 6	D 2
A 3	B 6	D 3	A 3	B 6	D 4	A 3	B 6	D 5
A 3	B 6	D 6	A 3	B 6	D 7	A 3	B 6	D 8
A 4	B 6	D 1	A 4	B 6	D 2	A 4	B 6	D 3
A 4	B 6	D 4	A 4	B 6	D 5	A 4	B 6	D 6
A 4	B 6	D 7	A 4	B 6	D 8	A 5	B 6	D 1
A 5	B 6	D 2	A 5	B 6	D 3	A 5	B 6	D 4
A 5	B 6	D 5	A 5	B 6	D 6	A 5	B 6	D 7
A 5	B 6	D 8	A 6	B 6	D 1	A 6	B 6	D 2
A 6	B 6	D 3	A 6	B 6	D 4	A 6	B 6	D 5
A 6	B 6	D 6	A 6	B 6	D 7	A 6	B 6	D 8

[0156] 54) The compound according to Formula (I), wherein

R^1 is a hydrogen atom, L^1 and L^2 are single bonds, L^3 is a methylene group, and

- 5 A, B, and D are a combination listed in Table 2 (where A1 to A6, B1 to B6, and D1 to D8 in Table are the substituents below, in 54), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.




[0157] 55) The compound according to Formula (I), wherein

R^1 is a hydrogen atom, L^1 and L^2 are single bonds, L^3 is a methylene group, and

A, B, and D are a combination listed in Table 2 (where A1 to A6, B1 to B6, and D1 to D8

5 in Table are the substituents below, in 55), the tautomer of the compound; the pharmaceutically acceptable salt thereof, or the solvate thereof.




[0158] 56) The compound including the combinations according to 48) to 55), wherein L^1 is SO_2 , the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

5 [0159] 57) The compound including the combinations according to 48) to 55), wherein L^1 is CO, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0160] 58) The compound according to Formula (I), wherein

R^1 is a hydrogen atom, L^1 and L^2 are single bonds, L^3 is a methylene group, and

10 A, B, and D are a combination listed in Table 1 (where A1 to A3, B1 to B3, and D1 to D16 in Table are the substituents below, in 58)), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.




[0161] 59) The compound including the combination according to 58), wherein L^1 is NH, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

5 [0162] 60) The compound including the combination according to 58), wherein L^1 is O, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0163] 61) The compound according to Formula (I), wherein

R^1 is a hydrogen atom, L^1 and L^2 are single bonds, L^3 is a methylene group, and

10 A, B, and D are a combination listed in Table 1 (where A1 to A3, B1 to B3, and D1 to D16 in Table are the substituents below, in 61)), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.



62) The compound including the combination according to 61), wherein L^2 is NH, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

5 [0164] 63) The compound including the combination according to 61), wherein L^2 is O, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

64) The compound including the combination according to 61), wherein L^2 is NHCH_2 , the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

10

[0165] 65) The compound including the combination according to any one of 48) to 64), wherein R^1 is a methoxy group, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

[0166] 66) The compound including the combination according to any one of 48) to 64), wherein R^1 is a n-butoxy group the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

15

[0167] 67) The compound including the combination according to any one of 48) to 64), wherein R¹ is a trifluoromethoxy group, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

68) The compound including the combination according to any one of 48) to 64),
5 wherein R¹ is a trifluoromethyl group, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

69) The compound including the combination according to any one of 48) to 64), wherein R¹ is a methyl group, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

10 [0168] 70) The compound including the combination according to 48) to 69), wherein L³ is 1,2-ethylene, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

71)

15 A T-type calcium channel inhibitor comprising the compound, as described in any one of 1) to 70), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof, as an active component.

72)

20 A preventive agent, a therapeutic agent, and/or an improving agent for a disease treatable by a T-type calcium channel inhibitory action comprising the T-type calcium channel inhibitor as described in 71) as an active component.

73)

A therapeutic agent for neuropathic pain comprising the T-type calcium channel inhibitor as described in 71) as an active component.

74)

25 A medicine comprising the compound, as described in any one of 1) to 70), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof, as an active component.

[0169] The present invention also includes compounds obtained through, for example, tautomerization or geometrical isomerization of the compound of Formula (I) of

the present invention regardless of endocyclic or exocyclic isomerization, mixtures of them, and mixtures of respective isomers. If the compound has an asymmetric center or an asymmetric center is generated by isomerization, the present invention includes respective optical isomers and mixtures of optical isomers at any ratio. If having two or
5 more asymmetric centers, the compound includes diastereomers due to optical isomerism of the respective asymmetric centers. The compound of the present invention includes mixtures containing all isomers at any ratio. For example, diastereomers can be separated by a method well-known to a person skilled in the art, such as fractional crystallization and column chromatography, and an optically active compound can be
10 produced by an organic chemical technique well-known for the purpose.

[0170] The compound of Formula (I) of the present invention or pharmaceutically acceptable salts thereof can be present in any crystal form depending on production conditions and can be present as any hydrate. These crystal forms, hydrates, and mixtures of them are also included in the scope of the present invention. The compound
15 may be present as a solvate containing an organic solvent such as acetone, ethanol, 1-propanol, and 2-propanol, and such solvates are also included in the scope of the present invention.

[0171] The present invention includes pharmaceutically acceptable salts of Formula (I) of the present invention.

20 The compound of Formula (I) of the present invention can be converted into a pharmaceutically acceptable salt, as necessary, or a free form of the compound can be converted from such a salt. Examples of the pharmaceutically acceptable salt of the present invention include

alkali metal salts (such as lithium, sodium, and potassium salts),
25 alkaline earth metal salts (such as magnesium and calcium salts),
ammonium salts,
organic base salts,
amino acid salts,
inorganic acid salts (such as hydrochlorates, hydrobromides, phosphates, and sulfates),

and

organic acid salts (such as acetates, citrates, maleates, fumarates, tartarates, benzenesulfonates, methanesulfonates, and p-toluenesulfonates).

[0172] The present invention also includes prodrugs of the compound of Formula (I) of the present invention.

Prodrugs are derivatives that are derived from a pharmaceutical compound and have a chemically or metabolically degradable group and are compounds that are degraded by solvolysis or under physiological conditions in vivo into a pharmacologically active, pharmaceutical compound. Methods for selecting and producing an appropriate prodrug derivative are described in Design of Prodrugs (Elsevier, Amsterdam 1985), for example. For the present invention, if having a hydroxy group, the compound is reacted with an appropriate acyl halide, an appropriate acid anhydride, or an appropriate halogenated alkyloxycarbonyl compound to produce an acyloxy derivative as a prodrug, for example. Particularly preferred structures for the prodrug are exemplified by -O-COC₂H₅, -O-CO(t-Bu), -O-COC₁₅H₃₁, -O-CO(m-CO₂Na-Ph), -O-COCH₂CH₂CO₂Na-OCOCH(NH₂)CH₃, -O-COCH₂N(CH₃)₂, and -O-CH₂OC(=O)CH₃. If the compound included in the present invention has an amino group, the compound having an amino group is reacted with an appropriate acid halide, an appropriate mixed acid anhydride, or an appropriate halogenated alkyloxycarbonyl compound to produce a prodrug, for example. Particularly preferred structures for the prodrug are exemplified by -N-CO(CH₂)₂₀OCH₃, -N-COCH(NH₂)CH₃, and -N-CH₂OC(=O)CH₃.

[0173] The preventive agent, the therapeutic agent, and/or the improving agent that are used for a disease treatable by a T-type calcium channel inhibitory action and comprise the T-type calcium channel inhibitor of the present invention as an active component can be typically administered in oral administration forms such as tablets, capsules, powdered drugs, granules, pills, and syrups, rectal administration forms, transdermal systems, or injection forms. The agent can be administered as a single therapeutic agent or as a mixture with other therapeutic agents. Such an agent can be singly administered but is typically administered in the form of a pharmaceutical

composition. Such a formulation can be produced by adding pharmacologically, pharmaceutically acceptable additives in usual ways. In other words, the oral formulations may contain common additives such as diluents, lubricants, bonding agents, disintegrants, wetting agents, plasticizers, and coating agents. Liquid formulations for oral administration may be aqueous suspensions, oily suspensions, solutions, emulsions, syrups, elixirs, or other forms, or may be produced as dry syrups, to which water or another appropriate solvent is added before use. The liquid formulations may contain common additives such as suspending agents, flavors, diluents, and emulsifiers. For rectal administration, the agent can be administered as suppositories. The suppositories can contain appropriate substances such as cacao butter, laurin butter, macrogol, glycerogelatin, Witepsol, sodium stearate, and mixtures of them as a base and, as necessary, emulsifiers, suspending agents, preservatives, and other additives. For the injections, pharmaceutical components such as distilled water for injection, physiological saline, 5% glucose solution, propylene glycol, other solvents or solubilizing agents, pH regulators, tonicity agents, and stabilizing agents may be used to form aqueous dosage forms or use-time dissolution type dosage forms.

[0174] The pharmaceutical composition of the present invention comprises the compound (or a pharmaceutically acceptable salt of the compound) of the present invention as an active component, pharmaceutically acceptable carriers, and, if needed, one or a plurality of additional therapeutic agents or adjuvants. Examples of such an additional therapeutic agent include i) cannabinoid receptor agonists or antagonists, ii) narcotic analgesics (opioid analgesics), iii) serotonin (5-HT) receptor agonists or antagonists, iv) sodium channel blockers, v) NMDA receptor agonists or antagonists, vi) COX-2 selective inhibitors, vii) NK1 antagonists, viii) non-steroidal anti-inflammatory drugs ("NSAIDs"), ix) selective serotonin reuptake inhibitors ("SSRIs") and/or selective serotonin and norepinephrine reuptake inhibitors ("SSNRIs"), x) tricyclic antidepressants, xi) GABA receptor agonists, xii) lithium, xiii) valproates, xiv) Neurontin (gabapentin), xv) pregabalin, xvi) adrenergic receptor agonists or antagonists, xvii) neurotrophin, xviii) capsaicin (TRPV1) receptor agonists or antagonists, xix) CGRP receptor antagonists, xx)

steroids, xxi) bisphosphonates, and xxii) histamine receptor antagonists. The composition contains compositions suitable for oral, rectal, local, and parenteral (including subcutaneous, intramuscular, and intravenous) administrations. An optimum route for any administration is determined depending on a specific host and specific conditions and seriousness of the host to which the active component is to be administered. The pharmaceutical composition can be favorably administered in a single unit dosage form and can be prepared by any method well-known in the field of pharmaceutics.

[0175] To administer the medicinal agent of the present invention to a human, the dose is determined depending on the age and conditions of a patient. The dose for adults is typically about 0.1 to 1,000 mg/human/day through oral or rectal administration and about 0.05 mg to 500 mg/human/day through injection. These numerical values are merely illustrative values, and the dose should be determined depending on the conditions of a patient.

[0176] The compound or the pharmaceutical composition of the present invention are intended to be used for all diseases to which T-type calcium channels relate to.

The primary target disease is pain.

The target disease is specifically chronic pain and more specifically neuropathic pain.

[0177] In more detail, the pain is classified into chronic pains and acute pains including neuropathic pain, inflammatory pain, cancer pain, and visceral pain, of which primary diseases are exemplified by diabetic neuropathy, traumatic neurological disorder, nerve compression, strangulation, spinal cord injury, cerebral apoplexy, fibromyalgia syndrome, carpal tunnel syndrome, osteoarthritis, rheumatoid arthritis and multiple sclerosis, herpes zoster, herpes simplex, syphilis, nerve disorders induced by cancer chemotherapy, HIV, and HIV treatment, chronic joint pain, postherpetic neuralgia, neuroma pain, trigeminal neuralgia, phantom limb pain, postoperative pain, stump pain, tooth pain, plexus neuropathy, glossopharyngeal neuralgia, laryngeal neuralgia, migraine, carcinomatous neuropathy, polyneuropathy, causalgia, low back pain, complex regional

pain syndrome (CRPS), and thalamic pain. Pains derived from other primary diseases except the above are also included in the target disease of the present invention.

[0178] Examples of other diseases except the pain include diseases associated with central nervous system (CNS) disorders, diseases associated with bladder function disorders, cerebral apoplexy, itching, atopic dermatitis, hypertension, ischemic heart diseases, atrial fibrillation, age-related macular degeneration, cancer, diabetes mellitus, chronic kidney disease, sterility, and sexual dysfunction. Examples of the diseases associated with central nervous system (CNS) disorders include epilepsy, essential tremor, schizophrenia, Parkinson's disease, manic-depressive illness, bipolar disorder, depression, anxiety, dementia, drug dependence, Huntington's disease, and sleep disturbance. Examples of the diseases associated with bladder function disorders include overactive bladder.

[0179] The compound is also clinically used for the treatment of epilepsy and partial and generalized tonic seizure. The compound is also useful for neuroprotection under ischemic conditions caused by cerebral apoplexy or neurotrauma and is useful for the treatment of multiple sclerosis. The compound is useful for the treatment of tachyarrhythmia. The compound is useful for the treatment of depression, more specifically, depressive disorders, for example, sudden or recurrent major depressive disorder, and mood disorders such as dysthymic disorder and bipolar disorders, for example, bipolar I disorder, bipolar II disorder, and cyclothymic disorder; and neuropsychiatric disorders including panic disorder with or without agoraphobia, agoraphobia with no panic disorder history, specific phobias, for example, phobia specific to animals and anthropophobia, obsessive-compulsive disorder, stress disorders such as posttraumatic stress disorder and acute stress disorder, and anxiety disorders such as generalized anxiety disorder.

[0180] In addition to primates including human beings, various other mammals can be treated by the method of the present invention. Examples of the treatable mammals include, but are not limited to, rodents (for example, mice), cattle, sheep, goats, horses, dogs, and cats. The method can be performed on other species such as birds (for

example, chickens).

[0181] When specifically used for the treatment of depression or anxiety, the compound of the present invention can be used in combination with other antidepressants or antianxiety drugs such as norepinephrine reuptake inhibitors, selective serotonin reuptake inhibitors (SSRIs), monoamine oxidase inhibitors (MAOIs), reversible inhibitors of monoamine oxidase (RIMAs), serotonin and noradrenaline reuptake inhibitors (SNRIs), α -adrenergic receptor antagonists, atypical antidepressants, benzodiazepines, 5-HT_{1A} agonists or antagonists, specifically, 5-HT_{1A} partial agonists, neurokinin-1 receptor antagonists, corticotropin-releasing factor (CRF) antagonists, and pharmaceutically acceptable salts of them.

[0182] In addition, the compound of the present invention can be administered in order to prevent the conditions and the disorders described above and to prevent other conditions and disorders to which calcium channel activity relates, in a dose level effective in the prevention.

[0183] Creams, ointments, jellies, solutions, or suspensions containing the compound can be locally used. Mouthwashes and gargles are included within the local applications for the present invention.

[0184] The compound of the present invention can be synthesized by the methods shown below, but the production methods below are typical examples of the production method and are not intended to limit the production method.

[0185] In a typical method for producing the compound of the present invention, for smooth reactions, a reaction in the presence of an acid or a base may efficiently proceed, and a reaction under microwave irradiation may efficiently proceed.

Among the typical production methods shown below, in schemes (1) to (3), (6) to (8), (11) to (15), and (18) to (20), a reaction particularly in the presence of a base such as potassium carbonate and triethylamine may be efficient for a smooth reaction.

Among the typical production methods shown below, in schemes (4), (10), (15), and (16), a reaction particularly in the presence of a base such as sodium hydroxide, potassium hydroxide, potassium tert-butoxide, sodium tert-butoxide, cesium carbonate,

potassium carbonate, and triethylamine may be efficient for a smooth reaction.

Among the typical production methods shown below, in formulae (5) and (12), a reaction particularly with a Bronsted acid catalyst such as acetic acid, trifluoroacetic acid, hydrochloric acid, sulfuric acid, and p-toluenesulfonic acid or with a Lewis acid such as titanium tetrachloride, a trifluoroborane-diethyl ether complex, and scandium triflate may
5 be efficient for a smooth reaction.

[0186] In the typical production methods shown below, each of reagents and raw material compounds can be appropriately used in an equimolar amount or an excess molar amount relative to one compound of raw material compounds.

10 [0187] In the typical production methods of the compound of the present invention shown below, general formulae of intermediates and a final product are shown in each step, but the general formulae of these intermediates and final product also generally includes derivatives protected with protective groups. A derivative protected with a protective group means a compound that can yield a target compound by
15 hydrolysis, reduction, oxidation, dehydration, halogenation, alkylation, or a similar reaction, as necessary, and includes a compound protected with a protective group that is acceptable for organic synthetic chemistry, for example.

Protection and deprotection can be carried out with well-known protective groups through protection and deprotection reactions (see Protective Groups in Organic
20 Synthesis, Fourth edition, by T.W. Greene, John Wiley & Sons Inc, 2006, for example).

Hydrolysis, reduction, oxidation, dehydration, and halogenation can be carried out by well-known transformation methods of functional groups (see Comprehensive Organic Transformations, Second Edition, by R.C. Larock, Wiley-VCH, 1999, for example).

[0188] (Symbol in typical production method)

25 In the typical production methods shown below, symbols in the drawings are as follows unless otherwise noted:


In formulae, R^1 , L^1 , L^2 , L^3 , A, B, and D are the same as in General Formula (I).

R^1 is a leaving group such as halogen atoms, a methanesulfonyloxy group, and a p-toluenesulfonyloxy group.

X is a halogen atom.


R^{PR} is a hydrogen atom or a protective group such as a Boc group and a Z group.

[0189] Of the compounds of Formula (I), compound (1)-3 can be produced by the production method of scheme (1) below, for example.




The compound (1)-3 can be synthesized using compound (1)-1 and compound (1)-2 in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

[0190] Of the compounds of Formula (I), compound (2)-3 can be produced by the production method of scheme (2) below, for example. (In the scheme, D is a 3 to 11-membered non-aromatic heterocycle containing NH or a 5 to 10-membered aromatic heterocycle containing NH)



The compound (2)-3 can be synthesized using compound (2)-1 and compound (2)-2 in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.


[0191] Of the compounds of Formula (I), compound (3)-3 can be produced by the production method of scheme (3) below, for example. (In the scheme, L¹ is a single bond, S, NR², or O, and B is a 3 to 11-membered heterocyclylene group containing NH or a 5 to 10-membered heteroarylene group containing NH when L¹ is a single bond)



The compound (3)-3 can be synthesized using compound (3)-1 and an equal amount or

excess amount of halogenated derivative (3)-2 in the presence of copper powder or a copper salt in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.


[0192] Of the compounds of Formula (I), compound (4)-3 can be produced by the production method of scheme (4) below, for example. (In the scheme, L¹ is a single bond, S, NR², or O, and B is a 3 to 11-membered heterocyclylene group containing NH or a 5 to 10-membered heteroarylene group containing NH when L¹ is a single bond)



The compound (4)-3 can be synthesized using compound (4)-1 and compound (4)-2 in the presence of a palladium catalyst such as tetrakis(triphenylphosphine)palladium(0), bis(triphenylphosphine)palladium(II) dichloride, and bis(acetonitrile)palladium(II) dichloride in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature. Alternatively, the reaction can be carried out under reaction conditions used for Buchwald-Hartwig reaction (see Advanced Synthesis & Catalysis, 2000, 346, pp. 1599-1626, for example). Although the conditions are not limited, tris(dibenzylideneacetone)dipalladium(0), tetrakis(triphenylphosphine)palladium(0), palladium(II) acetate, or a similar catalyst may be appropriately combined with 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (Xantphos), 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (SPhos), or 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl(XPhos), for example.


[0193] Of the compounds of Formula (I), compound (5)-2 can be produced by the production method of scheme (5) below, for example. (In the scheme, A is a C₃₋₁₁ cycloalkyl group, a C₃₋₁₁ cycloalkenyl group, a 3 to 11-membered heterocyclyl group, a C₁₋₆ alkyl group, a C₁₋₆ haloalkyl group, a C₃₋₁₁ cycloalkyl group, a C₂₋₆ alkenyl group, or a C₂₋₆ haloalkenyl group, L¹ is a single bond or NR², and B is a 3 to 11-membered heterocyclylene group containing NH or a 5 to 10-membered heteroarylene group

containing NH when L^1 is a single bond)



The compound (5)-2 can be synthesized using compound (5)-1 and a ketone or an aldehyde in the presence of a reducing agent such as sodium borohydride and triacetoxyborohydride in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.


[0194] Of the compounds of Formula (I), compound (6)-3 can be produced by the production method of scheme (6) below, for example. (In the scheme, L^2 is a single bond, S, NR^2 , or O, B is a 3 to 11-membered heterocyclylene group containing NH or a 5 to 10-membered heteroarylene group containing NH when L^2 is a single bond, and R^A is a C_{1-6} alkyl group (the C_{1-6} alkyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^6) or a dodecyl group)



The compound (6)-3 can be synthesized using compound (6)-1, (6)-4, or (6)-5 and compound (6)-2 in an appropriate solvent or without solvent at from 0°C to a heat-reflux


temperature.

[0195] Of the compounds of Formula (I), compound (7)-3 can be produced by the production method of scheme (7) below, for example. (In the scheme, B is a 3 to 11-membered heterocyclylene group containing NH or a 5 to 10-membered heteroarylene group containing NH)



The compound (7)-3 can be synthesized using compound (7)-1 and compound (7)-2 in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.


[0196] Of the compounds of Formula (I), compound (8)-3 can be produced by the production method of scheme (8) below, for example. (In the scheme, R¹ is a C₁₋₆ alkoxy group, and other symbols are the same as the respective definitions above)



The compound (8)-3 can be synthesized from compound (8)-1 with alkyl halide (8)-2 in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

[0197] Raw material synthesis 1

Compound (9)-2 can be produced by the production method of scheme (9) below, for example. (In the scheme, X is a halogen atom)




The compound (9)-2 can be synthesized by catalytic hydrogenation of compound

(9)-1 with palladium or a similar catalyst in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

[0198] Raw material synthesis 2


Compound (10)-2 can be produced by the production method of scheme (10) below,
5 for example. (In the scheme, X is a halogen atom)



The compound (10)-2 can be synthesized by hydrolysis of compound (10)-1 with a base such as sodium hydroxide in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

10 [0199] Raw material synthesis 3

Compound (11)-3 can be produced by the production method of scheme (11) below, for example. (In the scheme, X is a halogen atom, L^2 is a single bond, NR^2 , O, S, SO, or SO_2 , and B is a 3 to 11-membered heterocyclylene group containing NH or a 5 to 10-membered heteroarylene group containing NH when L^2 is a single bond)




15

The compound (11)-3 can be synthesized from compound (11)-1 and compound (11)-2 in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

[0200] Raw material synthesis 4

20 Compound (12)-3 can be produced by the production method of scheme (12) below, for example. (In the scheme, L^3 is CH_2)




Compound (12)-2 can be obtained by reaction of compound (12)-1 with formaldehyde or paraformaldehyde in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

- 5 The compound (12)-3 can be synthesized by halogenation of the compound (12)-2 with thionyl chloride or a similar reagent or by sulfonyl esterification of the compound (12)-2 with p-toluenesulfonyl chloride, methanesulfonyl chloride, or a similar reagent.

[0201] Raw material synthesis 5

- 10 Compound (13)-3 can be produced by the production method of scheme (13) below, for example.




The compound (13)-3 can be obtained by reaction of compound (13)-1 and compound (13)-2 in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

- 15 When R^{PR} is a protective group in the compound (13)-3, deprotection can yield a compound of which R^{PR} is a hydrogen atom.

[0202] Raw material synthesis 6

Compound (14)-4 can be produced by the production method of scheme (14) below, for example.




Compound (14)-3 can be synthesized by reaction of compound (14)-1 and compound (14)-2 in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

The compound (14)-4 can be synthesized by reaction of the compound (14)-3 with an equal amount or excess amount of phosphorus oxychloride in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

[0203] Raw material synthesis 7

Compound (15)-4 can be produced by the production method of scheme (15) below, for example.




The compound (15)-4 can be obtained by hydrolysis of compound (15)-1 and subsequent reaction with compound (15)-3.

Compound (15)-2 can be synthesized by hydrolysis of the compound (15)-1 with a base such as sodium hydroxide and potassium hydroxide in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

The compound (15)-4 can be synthesized by reaction of the compound (15)-2 and the compound (15)-3 in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

[0204] Raw material synthesis 8

Compound (16)-3 can be produced by the production method of scheme (16) below, for example.




The compound (16)-3 can be obtained by reaction of compound (16)-1 and compound (16)-2 in the presence of a palladium catalyst such as tetrakis(triphenylphosphine)palladium(0), bis(triphenylphosphine)palladium(II)

dichloride, and bis(acetonitrile)palladium(II) dichloride in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature. Alternatively, the reaction can be carried out under reaction conditions used for Buchwald-Hartwig reaction (see Advanced Synthesis & Catalysis, 2000, 346, pp. 1599-1626, for example). Although the conditions are not limited, tris(dibenzylideneacetone)dipalladium(0), tetrakis(triphenylphosphine)palladium(0), palladium(II) acetate, or a similar catalyst may be appropriately combined with 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (Xantphos), 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (SPhos), or 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl(XPhos), for example.

- 10 When R^{PR} is a protective group in the compound (16)-3, deprotection can yield a compound of which R^{PR} is a hydrogen atom.

[0205] Raw material synthesis 9

- Compound (17)-3 can be produced by the production method of scheme (17) below, for example (in the scheme, A is a 3 to 11-membered non-aromatic heterocycle containing NH or a 5 to 10-membered aromatic heterocycle containing NH).



- The compound (17)-3 can be synthesized by reaction of compound (17)-1 and compound (17)-2 using a Mitsunobu reagent and a phosphine reagent in an appropriate solvent or without solvent at from -78°C to a heat-reflux temperature. Examples of the Mitsunobu reagent include diethyl azodicarboxylate and diisopropyl azodicarboxylate, and examples of the phosphine reagent include triphenylphosphine and tributylphosphine.

When R^{PR} is a protective group in the compound (17)-3, deprotection can yield a compound of which R^{PR} is a hydrogen atom.

[0206] Raw material synthesis 10

- 25 Compound (18)-3 can be produced by the production method of scheme (18) below, for example (in the scheme, L¹ is S, NR², or O).




The compound (18)-3 can be synthesized by reaction of compound (18)-1 and compound (18)-2 with an equal amount or excess amount of a base such as sodium hydride and lithium hydride in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

When R^{PR} is a protective group in the compound (18)-3, deprotection can yield a compound of which R^{PR} is a hydrogen atom.

[0207] Raw material synthesis 11

Compound (19)-3 can be produced by the production method of scheme (19) below, for example (in the scheme, L¹ is S, NR², or O).




The compound (19)-3 can be obtained by reaction of compound (19)-1 and compound (19)-2.

The compound (19)-3 can be synthesized by reaction of the compound (19)-1 and the compound (19)-2 in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

When R^{PR} is a protective group in the compound (19)-3, deprotection can yield a compound of which R^{PR} is a hydrogen atom.

[0208] Raw material synthesis 12

Compound (20)-3 can be produced by the production method of scheme (20) below, for example (in the scheme, E is a C₁₋₆ alkoxy carbonyl group or a carboxy group).




Compound (20)-2 can be synthesized from compound (20)-1 with an equal amount

or excess amount of a reducing agent such as a borane-THF complex, sodium borohydride, and lithium aluminum hydride in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

The compound (20)-3 can be synthesized from the compound (20)-2 with an equal amount or excess amount of a chlorinating agent such as thionyl chloride in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature or synthesized from the compound (20)-2 with a sulfonyl chloride such as p-toluenesulfonyl chloride and methanesulfonyl chloride in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

[0209] Raw material synthesis 13

Compounds (21)-8 and (21)-9 can be produced by the production method of scheme (21) below, for example. (In the scheme, R^A is a C₁₋₆ alkyl group (the C₁₋₆ alkyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁶) or a dodecyl group).



Compound (21)-2 can be synthesized by reaction of compound (21)-1 with 1,1'-carbonyldiimidazole in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature.

Compound (21)-4 can be synthesized by reaction of the compounds (21)-2 and (21)-3 in an appropriate solvent or without solvent at from 0°C to a heat-reflux

temperature.

The compound (21)-8 can be synthesized by reaction of the compound (21)-4 with an orthoester as shown by the compound (21)-5 in an appropriate solvent or without solvent at from 0°C to a heat-reflux temperature or by reaction of the compound (21)-4
 5 with an acid chloride or an acid anhydride shown by the compound (21)-6 or (21)-7 in an appropriate solvent or without solvent using a dehydrating reagent such as phosphorus oxychloride, as necessary, at from 0°C to a heat-reflux temperature.

The compound (21)-9 can be synthesized by reaction of the compound (21)-8 with an oxidizing agent such as m-chloroperbenzoic acid, sodium periodate, and oxone in an
 10 appropriate solvent or without solvent at from 0°C to a hear-reflux temperature.

[0210] The production method is not limited to the above, and the compound of Formula (1) can be synthesized by a common method for synthesizing a triazine compound. The common method for synthesizing a triazine compound is described in the following document: Heterocyclic Compounds, New Edition, Applications
 15 (Kodansha Ltd., 2004) pp. 167 to 195.

[0211] Synthesis method:

The compound of the present invention can be prepared in accordance with the scheme provided below and the procedures provided in Examples. The substituents are the same as the above unless otherwise defined or obvious to a person skilled in the art.

20 [0212] Novel compounds of the present invention can be easily synthesized through techniques known to a person skilled in the art, for example, described in Advanced Organic Chemistry, March, the fifth edition, John Wiley and Sons, New York, NY, 2001; Advanced Organic Chemistry, Carey and Sundberg, Vols. A and B, the third edition, Plenum Press, Inc., New York, NY, 1990; Protective groups in Organic Synthesis,
 25 Green and Wuts, the second edition, John Wiley and Sons, New York, NY, 1991; Comprehensive Organic Transformations, Larock, VCH Publishers, Inc., New York, NY, 1988; Handbook of Heterocyclic Chemistry, Katritzky and Pozharskii, the second edition, Pergamon, New York, NY, 2000; and reference documents cited therein. Other reference documents referred for the synthesis of novel compounds in the present

invention include Buckwald et al., *Tetrahedron*, 2004, Vol. 60, pp. 7397-7403; Li et al., *Tetrahedron Lett.*, 2004, Vol. 45, pp. 4257-4260; and Jean et al., *J. Org. Chem.*, 2004, Vol. 69, pp. 8893-8902. Starting materials for the compound can be prepared by standard synthetic conversions from chemical precursors that are easily available from commercial suppliers including Aldrich Chemical Co. (Milwaukee, WI); Sigma Chemical Co. (St. Louis, MO); Lancaster Synthesis (Windham, N.H.); Ryan Scientific (Columbia, S.C.); Maybridge (Cornwall, UK); Matrix Scientific (Columbia, S.C.); Arcos (Pittsburgh, PA); and Trans World Chemicals (Rockville, MD).

[0213] The procedures for synthesizing compounds described in the present specification can include one or a plurality of steps of protection and deprotection of functional groups and purification (for example, recrystallization, distillation, column chromatography, flash chromatography, medium-pressure column chromatography, thin-layer chromatography (TLC), radial chromatography, and high-pressure liquid chromatography (HPLC)). A product can be characterized by various techniques well-known in the chemical field, including proton and carbon-13 nuclear magnetic resonance (^1H and ^{13}C NMR), infrared and ultraviolet spectroscopy (IR and UV), X-ray crystallography, elementary analysis, and HPLC-mass analysis (HPLC-MS). The procedures for the protection and deprotection of functional groups and the methods of purification, structure identification, and quantitative determination are well-known to a person skilled in the chemical synthesis.

[0214] Solvents preferably used for the synthesis of the compound by the reactions dissolve one or all of reactants, at least partially, and do not disadvantageously react with any of reactants or reaction products. Specific examples of the preferred solvent include aromatic hydrocarbons (for example, toluene and xylene), halogenated solvents (for example, methylene chloride, chloroform, carbon tetrachloride, and chlorobenzene), ethers (for example, diethyl ether, diisopropyl ether, tert-butyl methyl ether, diglyme, tetrahydrofuran, dioxane, and anisole), nitriles (for example, acetonitrile and propionitrile), ketones (for example, 2-butanone, diethyl ketone, and tert-butyl methyl ketone), alcohols (for example, methanol, ethanol, n-propanol, isopropanol,

n-butanol, and t-butanol), N,N-dimethylformamide (DMF), dimethyl sulfoxide (DMSO), and water. Mixtures of two or more of the solvents may be used.

[0215] Examples of the base preferably used for the synthesis of the compound of the present invention typically include

- 5 alkali metal hydrides and alkaline earth metal hydrides (for example, lithium hydride, sodium hydride, potassium hydride, and calcium hydride),
alkali metal amides (for example, lithium diisopropylamide (LDA), lithium hexamethyldisilazide (LHMDS), potassium hexamethyldisilazide (KHMDs), lithium amide, sodium amide, and potassium amide),
10 metal carbonates (for example, lithium carbonate, sodium carbonate, cesium carbonate, sodium hydrogen carbonate, and cesium hydrogen carbonate),
alkali metal alkoxides and alkaline earth metal alkoxides (for example, sodium methoxide, sodium ethoxide, potassium tert-butoxide, and magnesium ethoxide),
alkali metal alkyls (for example, methyllithium, n-butyllithium, sec-butyllithium,
15 t-butyllithium, and phenyllithium), alkyl magnesium halides, organic bases (for example, trimethylamine, triethylamine, triisopropylamine, N,N-diisopropylethylamine, piperidine, N-methylpiperidine, morpholine, N-methylmorpholine, pyridine, collidine, lutidine, and 4-dimethylaminopyridine), and
bicyclic amines (for example, 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) and
20 1,4-diazabicyclo[2.2.2]octane (DABCO)).

[0216] Functional groups of the compounds shown in the schemes below can be treated by a standard functional group conversion technique operable by a person skilled in the art, yielding an intended compound according to the present invention, if appropriate.

- 25 [0217] Other variations and modifications are obvious to a person skilled in the art and are included in the scope and teaching of the present invention. The present invention is not limited except the description in the claims below.

[0218] The present invention will be described in detail with reference to Reference Synthesis Examples, Synthesis Examples, Test Examples, and Formulation

Examples below. The present invention, however, is not limiting to these Examples.

In Examples, NMR means a nuclear magnetic resonance spectrum, LC/MS means liquid chromatography/mass spectrometry, (v/v) means (volume/volume), and expression of Rf and expression of Ex in the drawings mean Reference Synthesis Example and

5 Synthesis Example, respectively.

Morphology means a shape.

When ¹H-NMR data is described, the data is measured at 300 MHz and represents chemical shifts δ (unit: ppm) (split patterns and integral values) of signals determined by using tetramethyl silane as an internal standard. "s" means a singlet, "d" means a
10 doublet, "t" means a triplet, "q" means a quartet, "quint" means a quintet, "sextet" means a sextet, "septet" means a septet, "dd" means a double doublet, "ddd" means a double double doublet, "m" means a multiplet, "br" means broad, "J" means a coupling constant, CDCl₃ means deuterated chloroform, and "DMSO-d₆" means deuterated dimethyl sulfoxide.

15 As a micro-wave reactor, Initiator sixty manufactured by Biotage Gb Ltd was used.

In the purification by silica gel column chromatography, one of Hi-Flash column manufactured by Yamazen Ltd., Silica gel 60 manufactured by Merck & Co., Inc., and PSQ60B manufactured by Fuji Silysia Chemical Ltd. was used, unless otherwise stated.

In the purification by silica gel (amino-based) column chromatography, Hi-Flash
20 Amino Column manufactured by Yamazen Ltd. or DM1020 manufactured by Fuji Silysia Chemical Ltd. was used, unless otherwise stated.

In the purification by silica gel thin-layer chromatography, PLC Plate manufactured by Merck & Co., Inc. was used, unless otherwise stated.

In the purification by amino-based thin-layer chromatography, PLCP5 Plates NH
25 manufactured by Fuji Silysia Chemical Ltd. was used, unless otherwise stated.

In the purification by preparative high-performance liquid chromatography, 6A Preparative High-performance Liquid Chromatography System manufactured by SHIMADZU CORPORATION was used, unless otherwise stated.

LC/MS was measured using an ESI (electrospray ionization) method under

following conditions. "ESI+" means an ESI positive ion mode, "ESI-" means an ESI negative ion mode, "LC/MS: cond" means analysis conditions of LC/MS, and "RT" means a retention time.

[0219] LC/MS Conditions 1

5 Apparatus: Waters Micromass ZQ

Column: Waters SunFire C18 (3.5 μ m, 4.6 $\times$ 20 mm)

Column temperature: 40°C

Solvents used

Solution A: 0.1% formic acid aqueous solution

10 Solution B: 0.1% formic acid - acetonitrile solution

Elution conditions used:

The measurement was started at a flow rate of 0.4 mL/min and a mixing ratio of the solution A and the solution B of 90/10 (v/v), and then the mixing ratio of the solution A and the solution B was linearly changed to 15/85 (v/v) for 3 minutes.

15 Thereafter, the mixing ratio of the solution A and the solution B was fixed at 15/85 (v/v) for 2 minutes, and then the mixing ratio of the solution A and the solution B and the flow rate were linearly changed to 90/10 (v/v) and 0.5 mL/min, respectively, for 0.5 minutes. Thereafter these conditions were fixed for 2.5 minutes.

[0220] LC/MS Conditions 2

20 Apparatus: Thermo LTQ XL

Column: Waters AQUITY UPLC BEH C18 (1.7 μ m, 2.1 $\times$ 50 mm)

Column temperature: 40°C

Solvents used

Solution A: 0.1% formic acid aqueous solution

25 Solution B: 0.1% formic acid - acetonitrile solution

Elution conditions used:

Conditions were fixed at a flow rate of 0.6 mL/min and a mixing ratio of the solution A and the solution B of 90/10 (v/v) and the measurement was started, and then, after 0.5 minutes, the mixing ratio of the solution A and the solution B was linearly

changed to 10/90 (v/v) for 2.5 minutes.

Thereafter, the mixing ratio of the solution A and the solution B was fixed at 10/90 (v/v) for 0.7 minutes, and then the mixing ratio of the solution A and the solution B and the flow rate were linearly changed to 90/10 (v/v) and 0.8 mL/min, respectively, for 0.1 minutes. Thereafter these conditions were fixed for 1.0 minute.

Thereafter, the mixing ratio of the solution A and the solution B and the flow rate were linearly changed to 90/10 (v/v) and 0.6 mL/min, respectively, for 0.1 minutes.

[0221] LC/MS Conditions 3

Apparatus: Waters Aquity SQD

10 Column: Waters AQUITY UPLC BEH C18 (1.7 μ m, 2.1×50 mm)

Column temperature: 40°C

Solvents used

Solution A: 0.1% formic acid aqueous solution

Solution B: 0.1% formic acid - acetonitrile solution

15 Elution conditions used:

Conditions were fixed at a flow rate of 0.6 mL/min and a mixing ratio of the solution A and the solution B of 90/10 (v/v) and the measurement was started, and then, after 0.5 minutes, the mixing ratio of the solution A and the solution B was linearly changed to 10/90 (v/v) for 1.5 minutes.

20 Thereafter, the mixing ratio of the solution A and the solution B was fixed at 10/90 (v/v) for 0.3 minutes, and then the mixing ratio of the solution A and the solution B and the flow rate were linearly changed to 90/10 (v/v) and 0.8 mL/min, respectively for 0.1 minutes. Thereafter these conditions were fixed for 1.0 minutes.

Thereafter, the mixing ratio of the solution A and the solution B and the flow rate were linearly changed to 90/10 (v/v) and 0.6 mL/min, respectively, for 0.1 minutes.

[0222] Reference Synthesis Example 1

2,4-Dichloro-6-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazine

To a tetrahydrofuran solution (150 mL) of 1,3,5-trichlorotriazine (19.6 g, 107 mmol),

sodium carbonate (24.5 g, 232 mmol) was added at 0°C and the resultant solution was stirred at 0°C for 5 minutes. To the reaction solution, 1-(4-fluorophenyl) piperazine dihydrochloride (15.8 g, 62.6 mmol) was added in two portions and the resultant mixture was stirred at room temperature for 3 days. After completion of the reaction, water (20 mL) was added and the pH was adjusted to 7 with 1 M hydrochloric acid. The obtained solid was collected by filtration, washed with water, and dried under reduced pressure to obtain the title compound (20.8 g, quantitative).

[0223] Reference Synthesis Example 2

6-Chloro-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

To a tetrahydrofuran solution (200 mL) of 2,4-dichloro-6-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazine (21.0 g, 63 mmol) synthesized in Reference Synthesis Example 1, 1 M sodium hydroxide aqueous solution (128 mL, 128 mmol) was added at room temperature and the resultant solution was stirred for 1 day. To the reaction solution, 1 M sodium hydroxide aqueous solution (32 mL, 32 mmol) was added and the resultant mixture was stirred for 6 hours. To the reaction solution, 1 M sodium hydroxide aqueous solution (19 mL, 19 mmol) was further added and the resultant solution was stirred for 2 hours. After completion of the reaction, pH of the reaction solution was adjusted to 4 by adding 1 M hydrochloric acid. After removing impurities, the resultant mixture was concentrated under reduced pressure to obtain a solid. The solid was collected by filtration, washed with water, and dried under reduced pressure to obtain the title compound (15.2 g, yield 79%).

[0224] Reference Synthesis Example 3

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

An acetic acid (15 mL) - water (35mL) mixed solution of 6-chloro-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one (1.00 g, 3.23 mmol) synthesized in Reference Synthesis Example 2 and palladium hydroxide-activated carbon catalyst (100 mg) was stirred under hydrogen atmosphere at room temperature for

3 days. After completion of the reaction, the reaction solution was filtered with Celite and the filtered residue was washed with methanol, followed by concentrating the obtained solution under reduced pressure. Toluene was added to the residue and the mixture was azeotropically dehydrated four times. The obtained residue was dried under reduced pressure to obtain the title compound (876 mg, yield 99%).

[0225] Reference Synthesis Example 4

{4-[4-(4-Fluorophenyl)piperazin-1-yl]-2-oxo-1,3,5-triazin-1(2H)-yl}methyl
4-methylbenzenesulfonate

10 To an ethanol/water solution (2/1 (v/v)) of
4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one (94 mg, 0.34 mmol)
synthesized in Reference Synthesis Example 3, a formaldehyde aqueous solution (42 mg,
0.51 mmol) was added and the resultant solution was stirred at 50°C for 5 hours. To the
reaction solution, a formaldehyde aqueous solution (83 mg, 1.02 mmol) was added and
15 the resultant mixture was stirred at 50°C for 3 hours. After completion of the reaction,
the reaction solution was concentrated under reduced pressure to obtain a crude product
of 4-[4-(4-fluorophenyl)piperazin-1-yl]-1-(hydroxymethyl)-1,3,5-triazin-2(1H)-one.
Subsequently, to a dichloromethane solution (2 mL) of
4-[4-(4-fluorophenyl)piperazin-1-yl]-1-(hydroxymethyl)-1,3,5-triazin-2(1H)-one,
20 triethylamine (57 mg, 0.41 mmol) and p-toluenesulfonic acid anhydride (123 mg, 0.38
mmol) were added and the resultant mixture was stirred at room temperature. After
completion of the reaction, the reaction solution was concentrated under reduced pressure
to obtain a crude product of the title compound and the crude product was used in the
next reaction as it was.

25 [0226] Reference Synthesis Example 5

tert-Butyl 4-(4,6-dichloro-1,3,5-triazin-2-yl)piperazine-1-carboxylate

tert-Butyl piperazine-1-carboxylate (9.09 g, 48.8 mmol), 1,3,5-trichlorotriazine (10.0 g, 54.2 mmol), and sodium carbonate (11.4 g, 108 mmol) were used to obtain the title

compound (14.9 g, yield 91%) by synthesis in a similar manner to Reference Synthesis Example 1.

[0227] Reference Synthesis Example 6

tert-Butyl 4-(6-chloro-4-oxo-4,5-dihydro-1,3,5-triazin-2-yl)piperazine-1-carboxylate

5

tert-Butyl 4-(4,6-dichloro-1,3,5-triazin-2-yl)piperazine-1-carboxylate (14.9 g, 44.5 mmol) synthesized in Reference Synthesis Example 5 and 1 M sodium hydroxide aqueous solution (111 mL, 111 mmol) were used to obtain the title compound (14.2 g, quantitative) by synthesis in a similar manner to Reference Synthesis Example 2.

10 [0228] Reference Synthesis Example 7

tert-Butyl 4-(4-oxo-4,5-dihydro-1,3,5-triazin-2-yl)piperazine-1-carboxylate

tert-Butyl 4-(6-chloro-4-oxo-4,5-dihydro-1,3,5-triazin-2-yl)piperazine-1-carboxylate (14.2 g, 45.0 mmol) synthesized in Reference Synthesis Example 6 and palladium hydroxide-activated carbon catalyst (142 mg) were used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 3. The crude product was used in the next reaction as it was.

15 [0229] Reference Synthesis Example 8

tert-Butyl 4-[5-(4-chlorobenzyl)-4-oxo-4,

20 5-dihydro-1,3,5-triazin-2-yl]piperazine-1-carboxylate

To an N,N-dimethylformamide solution (200 mL) of the crude product of tert-butyl 4-(4-oxo-4,5-dihydro-1,3,5-triazin-2-yl)piperazine-1-carboxylate synthesized in Reference Synthesis Example 7 and potassium carbonate (7.46 g, 54.0 mmol), 4-chlorobenzyl bromide (10.2 g, 49.5 mmol) was added at room temperature and the resultant reaction solution was stirred at room temperature for 1 day. After completion of the reaction, water was added to the reaction solution and a deposited solid was collected by filtration. The filtered residue was suspended in ethyl acetate and the suspended solid was collected by filtration and dried under reduced pressure to obtain the

title compound (12.4 g, two step yield 68%).

[0230] Reference Synthesis Example 9

1-(4-Chlorobenzyl)-4-(piperazin-1-yl)-1,3,5-triazin-2(1H)-one

5 To a dichloromethane solution (92 mL) of tert-butyl 4-[5-(4-chlorobenzyl)-4-oxo-4,5-dihydro-1,3,5-triazin-2-yl]piperazine-1-carboxylate (9.20 g, 22.7 mmol) synthesized in Reference Synthesis Example 8, trifluoroacetic acid (40 mL) was added and the resultant solution was stirred at room temperature for 4 hours. After completion of the reaction, the reaction solution was concentrated under reduced pressure and pH was adjusted to 7
10 by adding water and saturated sodium bicarbonate aqueous solution. Extraction with ethyl acetate from the resultant liquid was performed three times, and the organic layer was washed with brine. A deposited solid was collected by filtration, washed with water and ethyl acetate, and dried under reduced pressure to obtain the title compound (3.18 g, yield 46%).

15 [0231] Reference Synthesis Example 10

1-(4-Chlorobenzyl)-4-hydroxy-1,3,5-triazin-2(1H)-one

To an N,N-dimethylformamide solution (200 mL) of 1,3,5-triazine-2,4-diol (5.00 g, 44.2 mmol, synthesized according to the method described in Angew. Chem., 74, 354;
20 1962), sodium hydride (2.12 g, 48.6 mmol, purity 55%) was added at 0°C and the resultant solution was stirred for 1 hour. To the reaction solution, 4-chlorobenzyl bromide (9.99 g, 48.6 mmol) was added and the resultant solution was stirred at room temperature for 2 hours. After completion of the reaction, saturated ammonium chloride aqueous solution was added thereto, and extraction with ethyl acetate from the resultant
25 mixture was performed. The organic layer was washed with saturated ammonium chloride aqueous solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. To the obtained residue, toluene was added and the solid was collected by filtration, washed with toluene, and dried under reduced pressure to obtain the title compound (1.50 g, yield 14%).

[0232] Reference Synthesis Example 11

4-Chloro-1-(4-chlorobenzyl)-1,3,5-triazin-2(1H)-one

A toluene solution (25 mL) of 1-(4-chlorobenzyl)-4-hydroxy-1,3,5-triazin-2(1H)-one
 5 (600 mg, 2.52 mmol) synthesized in Reference Synthesis Example 10,
 N,N-diisopropylethylamine (0.52 mL, 4.63 mmol), and phosphorus oxychloride (0.94
 mL) was stirred at 70°C for 30 minutes. After completion of the reaction, the reaction
 solution was concentrated under reduced pressure and toluene was added, followed by
 concentrating the resultant mixture under reduced pressure again. To the obtained crude
 10 product of 4-chloro-1-(4-chlorobenzyl)-1,3,5-triazin-2(1H)-one, chloroform was added
 and the resultant solution was used in the next reaction as the chloroform solution.

[0233] Reference Synthesis Example 12

1-(4,6-Dichloro-1,3,5-triazin-2-yl)-4-(4-fluorophenyl)piperidin-4-ol

15 4-(4-Fluorophenyl)piperidin-4-ol (195 mg, 1.00 mmol), 1,3,5-trichlorotriazine (313
 mg, 1.70 mmol), and sodium carbonate (530 mg, 5.00 mmol) were used to obtain the title
 compound (304 mg, yield 89%) by synthesis in a similar manner to Reference Synthesis
 Example 1.

[0234] Reference Synthesis Example 13

20 6-Chloro-4-[4-(4-fluorophenyl)-4-hydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one

1-(4,6-Dichloro-1,3,5-triazin-2-yl)-4-(4-fluorophenyl)piperidin-4-ol (304 mg, 0.89
 mmol) synthesized in Reference Synthesis Example 12 and 1 M sodium hydroxide
 aqueous solution (1.80 mL, 1.80 mmol) were used to obtain the title compound (169 mg,
 25 yield 59%) by synthesis in a similar manner to Reference Synthesis Example 2.

[0235] Reference Synthesis Example 14

4-[4-(4-Fluorophenyl)-4-hydroxy-piperidin-1-yl]-1,3,5-triazin-2(1H)-one6-Chloro-4-[4-(4-fluorophenyl)-4-hydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one

(239 mg, 0.74 mmol) synthesized in Reference Synthesis Example 13 and palladium hydroxide-activated carbon catalyst (24 mg) were used to obtain the title compound (264 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 3.

[0236] Reference Synthesis Example 15

5 [3-(Trifluoromethyl)-1H-pyrazol-1-yl]methyl 4-methylbenzenesulfonate

A tetrahydrofuran solution (50 mL) of [3-(trifluoromethyl)-1H-pyrazol-1-yl]methanol (6.23 g, 37.5 mmol), p-toluenesulfonic acid anhydride (14.7 g, 45.0 mmol), and triethylamine (7.84 mL, 56.2 mmol) was stirred at room temperature for 19 hours. After completion of the reaction, water was added to the reaction solution and extraction with ethyl acetate from the resultant mixture was performed three times. The obtained organic layer was washed with a saturated sodium bicarbonate aqueous solution and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to obtain a crude product (12.8 g) of the title compound.

[0237] Reference Synthesis Example 16

tert-Butyl

4-(4-oxo-5-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)piperazine-1-carboxylate

20

An N,N-dimethylformamide solution (16 mL) of tert-butyl 4-(4-oxo-4,5-dihydro-1,3,5-triazin-2-yl)piperazine-1-carboxylate (862 mg, 3.07 mmol) synthesized in Reference Synthesis Example 7, potassium carbonate (1.02 g, 7.37 mmol), sodium iodide (46 mg, 0.31 mmol), 1-(chloromethyl)-3-(trifluoromethyl)-1H-pyrazole (536 mg, 2.90 mmol), and [3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl 4-methylbenzenesulfonate (1.03 g, 3.23 mmol) was stirred at 70°C for 2 hours. After completion of the reaction, the reaction solution was cooled to room temperature and water was added to the reaction solution, followed by extraction from the resultant mixture with ethyl acetate three times. The obtained organic layer was washed with

brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure.

The obtained residue was purified with silica gel column chromatography to obtain the title compound (228 mg, yield 15%).

[0238] Reference Synthesis Example 17

5 4-(Piperazin-1-yl)-1-([3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl)-1,3,5-triazin-2(1H)-one

tert-Butyl

4-(4-oxo-5-([3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl)-4,5-dihydro-1,3,5-triazin-2-yl)
10)piperazine-1-carboxylate (1.77 g, 4.12 mmol) synthesized in Reference Synthesis Example 16 was used to obtain a crude product (601 mg) of the title compound by synthesis in a similar manner to Reference Synthesis Example 9.

[0239] Reference Synthesis Example 18

6-[4-(4-Fluorophenyl)piperazin-1-yl]-1,3,5-triazine-2,4(1H,3H)-dione
15

An acetic acid solution of

2,4-dichloro-6-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazine (2.40 g, 7.31 mmol) synthesized in Reference Synthesis Example 1 and sodium acetate (1.80 g) was stirred at 80°C for 8 hours. After completion of the reaction, the reaction solution was poured
20 into water and the obtained solid was collected by filtration. Extraction with ethyl acetate was performed from the filtrate and the resultant ethyl acetate phase was concentrated under reduced pressure. The residue and the solid collected by filtration were combined and the combination was suspended into ethyl acetate, collected by filtration, washed with ethyl acetate, and dried under reduced pressure to obtain the title
25 compound (1.82 g, yield 86%).

[0240] Reference Synthesis Example 19

3-(4-Chlorobenzyl)-6-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazine-2,4(1H,3H)-dione

To an N,N-dimethylformamide solution of

6-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazine-2,4(1H,3H)-dione (1.20 g, 4.12 mmol) synthesized in Reference Synthesis Example 18, lithium hydride (75 mg, 4.94 mmol), 4-chlorobenzyl chloride (0.70 g, 4.12 mmol), and sodium iodide (catalytic amount) were added at room temperature and the resultant solution was stirred at 50°C

for 8 hours. After completion of the reaction, the reaction solution was poured into water and the resultant mixture was washed with ethyl acetate. Diluted hydrochloric acid was added to the water phase to adjust pH to 3 and the obtained solid was collected by filtration. Extraction with a mixed solution of ethyl acetate and tetrahydrofuran from the filtrate was performed and the organic layer was concentrated under reduced pressure.

The residue and the solid collected by filtration were combined to obtain the title compound (280 mg, yield 15%).

[0241] Reference Synthesis Example 20

4-(4-Chlorophenyl)-1-(4,6-dichloro-1,3,5-triazin-2-yl)piperidin-4-ol

4-(4-Chlorophenyl)piperidin-4-ol (2.12 g, 10.0 mmol), 1,3,5-trichlorotriazine (2.77 g, 15.0 mmol) and sodium carbonate (2.12 g, 20.0 mmol) were used to obtain a crude product of title compound by synthesis in a similar manner to Reference Synthesis Example 1.

[0242] Reference Synthesis Example 21

6-Chloro-4-[4-(4-chlorophenyl)-4-hydroxy-piperidin-1-yl]-1,3,5-triazin-2(1H)-one

The crude product of 4-(4-chlorophenyl)-1-(4,6-dichloro-1,3,5-triazin-2-yl)piperidin-4-ol (10.0 mmol) synthesized in Reference Synthesis Example 20 and 1 M sodium hydroxide aqueous solution (120.0 mL, 120 mmol) were used to obtain the title compound (3.19 g, yield 93%) by synthesis in a similar manner to Reference Synthesis Example 2.

[0243] Reference Synthesis Example 22a

4-[4-(4-Chlorophenyl)-4-hydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one

[0244] Reference Synthesis Example 22b

4-(4-Hydroxy-4-phenylpiperidin-1-yl)-1,3,5-triazin-2(1H)-one

6-Chloro-4-[4-(4-chlorophenyl)-4-hydroxy-piperidin-1-yl]-1,3,5-triazin-2(1H)-one (341 mg, 1.00 mmol) synthesized in Reference Synthesis Example 21 and palladium hydroxide-activated carbon catalyst (34.1 mg) were used to obtain a mixture of compounds of Reference Synthesis Example 22a and Reference Synthesis Example 22b being the title compounds (424 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 3.

[0245] Reference Synthesis Example 23

10 tert-Butyl 4-[(4-fluorophenyl)amino]piperidine-1-carboxylate

Under a nitrogen atmosphere, a xylene solution (65 mL) of tert-butyl 4-amino-piperidine-1-carboxylate (1.00 g, 5.00 mmol), 4-bromo-1-fluorobenzene (0.500 mL, 4.57 mmol), 2'-(dicyclohexylphosphino)-N,N-dimethyl-[1,1'-biphenyl]-2-amine (360 mg, 1.00 mmol), and tris(dibenzylideneacetone) dipalladium (0) (209 mg, 0.250 mmol) was stirred at 140°C for 4 hours. After completion of the reaction, the reaction solution was concentrated under reduced pressure and purified by silica gel column chromatography (hexane/ethyl acetate = 100/0 → 70/30 (v/v)) to obtain the title compound (980 mg, yield 66%).

20 [0246] Reference Synthesis Example 24

N-(4-Fluorophenyl)-N-methylpiperidin-4-amine

To an N,N-dimethylformamide solution (5 ml) of tert-butyl 4-[(4-fluorophenyl)amino]piperidine-1-carboxylate (250 mg, 0.850 mmol) synthesized in Reference Synthesis Example 23, sodium hydride (37 mg, 0.93 mmol, purity 60%) and methyl iodide (58 μ L, 0.93 mmol) were added at room temperature and the resultant solution was stirred at room temperature for 15 hours. After completion of the reaction, saturated ammonium chloride aqueous solution was added, and extraction with ethyl acetate was performed from the resultant mixture. The resultant organic layer was dried

over anhydrous magnesium sulfate and concentrated under reduced pressure to obtain a crude product of tert-butyl 4-[(4-fluorophenyl) (methyl)amino]piperidine-1-carboxylate (286 mg). Subsequently, to the obtained tert-butyl

4-[(4-fluorophenyl)(methyl)amino]piperidine-1-carboxylate (130 mg, 0.420 mmol), 4 M hydrogen chloride/1,4-dioxane solution (2 mL) was added and the resultant mixture was stirred at room temperature for 1 day. After completion of the reaction, the resultant solid was collected by filtration and washed with dichloromethane. To the obtained solid, water was added and pH of the resultant solution was adjusted to 9 with 1 M sodium hydroxide aqueous solution, followed by extraction from the resultant mixture with ethyl acetate and chloroform. The obtained organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to obtain the title compound (62 mg, yield 72%).

[0247] Reference Synthesis Example 25

tert-Butyl (R)-[1-(4-fluorophenyl)piperidin-3-yl]carbamate

Under a nitrogen atmosphere, a toluene solution (4 mL) of tert-butyl (R)-piperidin-3-yl-carbamate (500 mg, 2.50 mmol), 4-bromo-1-fluorobenzene (0.250 mL, 2.28 mmol), 2-(di-tert-butylphosphino)biphenyl (136 mg, 0.500 mmol), tris(dibenzylideneacetone) dipalladium (0) (104 mg, 0.130 mmol), and sodium tert-butoxide (307 mg, 3.50 mmol) was stirred at 100°C for 4 hours. After completion of the reaction, the reaction solution was concentrated under reduced pressure and purified by silica gel column chromatography (hexane/ethyl acetate = 100/0 → 70/30 (v/v)) to obtain the title compound (550 mg, yield 82%).

[0248] Reference Synthesis Example 26

(R)-1-(4-Fluorophenyl)piperidin-3-amine hydrochloride

To tert-butyl (R)-[1-(4-fluorophenyl)piperidin-3-yl]carbamate (250 mg, 0.85 mmol) synthesized in Reference Synthesis Example 25, 4 M hydrogen chloride/1,4-dioxane solution (2.5 mL) was added and the resultant mixture was stirred at room temperature

for 18 hours. After completion of the reaction, the resultant solid was collected by filtration to obtain the title compound (220 mg, quantitative).

[0249] Reference Synthesis Example 27

tert-Butyl (S)-[1-(4-fluorophenyl)piperidin-3-yl]carbamate

5

tert-Butyl (S)-piperidin-3-yl-carbamate (500 mg, 2.50 mmol) was used to obtain the title compound (484 mg, yield 72%) by synthesis in a similar manner to Reference Synthesis Example 25.

[0250] Reference Synthesis Example 28

10 (S)-1-(4-Fluorophenyl)piperidin-3-amine hydrochloride

To tert-butyl (S)-[1-(4-fluorophenyl)piperidin-3-yl]carbamate (250 mg, 0.85 mmol) synthesized in Reference Synthesis Example 27 was used to obtain the title compound (220 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 26.

15

[0251] Reference Synthesis Example 29

tert-Butyl (R)-[1-(4-fluorophenyl)pyrrolidin-3-yl]carbamate

tert-Butyl (R)-pyrrolidin-3-yl-carbamate (1.00 g, 5.36 mmol) was used to obtain the title compound (1.20 g, yield 80%) by synthesis in a similar manner to Reference Synthesis Example 25.

20

[0252] Reference Synthesis Example 30

(R)-1-(4-Fluorophenyl)pyrrolidin-3-amine hydrochloride

tert-Butyl (R)-[1-(4-fluorophenyl)pyrrolidin-3-yl]carbamate (250 mg, 0.89 mmol) synthesized in Reference Synthesis Example 29 was used to obtain the title compound (220 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 26.

25

[0253] Reference Synthesis Example 31

tert-Butyl (S)-[1-(4-fluorophenyl)pyrrolidin-3-yl]carbamate

tert-Butyl (S)-pyrrolidin-3-yl-carbamate (1.00 g, 5.36 mmol) was used to obtain the title compound (669 mg, yield 45%) by synthesis in a similar manner to Reference

5 Synthesis Example 25.

[0254] Reference Synthesis Example 32

(S)-1-(4-Fluorophenyl)pyrrolidin-3-amine hydrochloride

tert-Butyl (S)-[1-(4-fluorophenyl)pyrrolidin-3-yl]carbamate (250 mg, 0.89 mmol) synthesized in Reference Synthesis Example 31 was used to obtain the title compound (208 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 26.

[0255] Reference Synthesis Example 33

tert-Butyl 4-[3-(trifluoromethyl)-1H-pyrazol-1-yl]piperidine-1-carboxylate

15

To a tetrahydrofuran solution (1.0 mL) of 3-(trifluoromethyl)-1H-pyrazole (272 mg, 2.00 mmol), tert-butyl 4-hydroxy-piperidine-1-carboxylate (402 mg, 2.00 mmol), and triphenylphosphine (629 mg, 2.40 mmol), diisopropyl azodicarboxylate (1.36 mL, 2.60 mmol, toluene solution) was added at room temperature and the resultant solution was stirred at room temperature for 22 hours. After completion of the reaction, the reaction solution was concentrated under reduced pressure and purified by preparative liquid chromatography to obtain the title compound (289 mg, yield 45%).

20

[0256] Reference Synthesis Example 34

4-[3-(Trifluoromethyl)-1H-pyrazol-1-yl]piperidine

25

To a dichloromethane solution (1.5 mL) of tert-butyl 4-[3-(trifluoromethyl)-1H-pyrazol-1-yl]piperidine-1-carboxylate (150 mg, 0.50 mmol) synthesized in Reference Synthesis Example 33, trifluoroacetic acid (0.7 mL) was added at room temperature, and the resultant solution was stirred for 3 hours. After

completion of the reaction, the reaction solution was concentrated under reduced pressure and water was added to the concentrated reaction solution, followed by extraction from the resultant mixture with ethyl acetate. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to obtain the
5 title compound (140 mg, quantitative).

[0257] Reference Synthesis Example 35

tert-Butyl 3-[(4-fluorobenzyl)oxy]azetidine-1-carboxylate

To a tetrahydrofuran solution (2.0 mL) of tert-butyl
10 3-hydroxyazetidine-1-carboxylate (87 mg, 0.50 mmol), sodium hydride (33 mg, 0.75 mmol, purity 55%) was added at 0°C and the resultant solution was stirred for 30 minutes. Subsequently, 4-fluorobenzyl bromide (62.3 µL, 0.50 mmol) was added to the reaction solution and the resultant mixture was stirred at room temperature for 19 hours. After completion of the reaction, water was added to the reaction solution and extraction with
15 ethyl acetate from the resultant mixture was performed three times. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate) to obtain the title compound (68.5 mg, yield 49%).

20 [0258] Reference Synthesis Example 36

3-[(4-Fluorobenzyl)oxy]azetidine

tert-Butyl 3-[(4-fluorobenzyl)oxy]azetidine-1-carboxylate (68.5 mg, 0.24 mmol) synthesized in Reference Synthesis Example 35 was used to obtain the title compound
25 (53.3 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 34.

[0259] Reference Synthesis Example 37

tert-Butyl 3-(4-fluorophenoxy)azetidine-1-carboxylate

An N,N-dimethylformamide solution (1.5 mL) of tert-butyl 3-(tosyloxy)azetidine-1-carboxylate (164 mg, 0.50 mmol), 4-fluorophenol (67.3 mg, 0.60 mmol), and cesium carbonate (116 mg, 0.60 mmol) was stirred at 80°C for 16 hours. After completion of the reaction, water was added to the reaction solution and extraction with ethyl acetate from the resultant mixture was performed three times. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate) to obtain the title compound (97.4 mg, yield 73%).

10 [0260] Reference Synthesis Example 38

3-(4-Fluorophenoxy)azetidine

tert-Butyl 3-(4-fluorophenoxy)azetidine-1-carboxylate (97.4 mg, 0.36 mmol) synthesized in Reference Synthesis Example 37 was used to obtain the title compound (84.6 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 34.

[0261] Reference Synthesis Example 39

tert-Butyl 4-(4-fluorophenyl)-3,4-dihydroxypiperidine-1-carboxylate

20 To an acetone-water mixed solution of microencapsulated osmium oxide (753 mg, 0.30 mmol, 10% by weight, manufactured by Wako Pure Chemical Industries, Ltd.) and N-methylmorpholine-N-oxide (1.58 g, 13.5 mmol), tert-butyl 4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (2.50 g, 9.01 mmol) was added and the resultant solution was stirred at room temperature for 6 hours. After completion of the reaction, the reaction solution was filtered and saturated sodium thiosulfate aqueous solution was added to the filtrate, followed by extraction from the resultant mixture with ethyl acetate three times. The obtained organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography

25

(hexane/ethyl acetate) to obtain the title compound (2.45 g, yield 87%).

[0262] Reference Synthesis Example 40a

4-(4-Fluorophenyl)piperidine-3,4-diol

[0263] Reference Synthesis Example 40b

5 4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol

To a dichloromethane solution (14 mL) of tert-butyl
4-(4-fluorophenyl)-3,4-dihydroxypiperidine-1-carboxylate (1.45 g, 4.66 mmol)
synthesized in Reference Synthesis Example 39, trifluoroacetic acid (14 mL) was added
10 and the resultant solution was stirred at room temperature for 4 hours. After completion
of the reaction, the reaction solution was concentrated under reduced pressure by 4 mL,
and saturated sodium bicarbonate aqueous solution was added to the concentrated
solution, followed by extraction from the resultant mixture with ethyl acetate three times.
The obtained organic layer was washed with brine, dried over anhydrous sodium sulfate,
15 and concentrated under reduced pressure. The obtained residue was purified by silica
gel (amino-based) column chromatography to obtain the title compound of Reference
Synthesis Example 40a (189 mg, yield 48%).

The reaction solution in Reference Synthesis Example 40a was concentrated under
reduced pressure by 6 mL and p-toluenesulfonic acid hydrate (420 mg) and toluene (5
20 mL) were added to the concentrated solution, followed by stirring the resultant mixture at
100°C for 4 hours. After completion of the reaction, the reaction solution was
concentrated under reduced pressure and saturated sodium bicarbonate aqueous solution
was added to the concentrated solution, followed by extraction from the resultant mixture
with ethyl acetate three times. The organic layer was washed with brine, dried over
25 anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained
residue was purified by silica gel (amino-based) column chromatography to obtain the
title compound of Reference Synthesis Example 40b (82.6 mg, yield 15%).

[0264] Reference Synthesis Example 41

1-Amino-3-(4-fluorophenyl)propan-2-ol hydrochloride

To a dichloromethane solution (20 mL) of 1-allyl-4-fluorobenzene (1.00 g, 7.34 mmol), meta-chloroperoxybenzoic acid (1.80 g, 7.71 mmol) was added at 0°C and the resultant solution was stirred at room temperature for 3 days. After completion of the reaction, saturated sodium carbonate aqueous solution was added thereto and extraction with chloroform from the resultant mixture was performed. The organic layer was dried over anhydrous magnesium sulfate and filtered. To aqueous ammonia-methanol mixed solution, the chloroform solution of the product was added dropwise and the resultant mixture was stirred at 100°C for 30 minutes with a microwave reactor. To the reaction solution, concentrated hydrochloric acid was added and the resultant mixture was concentrated under reduced pressure. The obtained residue was recrystallized with diethyl ether-methanol mixed solution to obtain the title compound (0.75 g, yield 50%).

[0265] Reference Synthesis Example 42

2-Bromo-N-[3-(4-fluorophenyl)-2-hydroxypropyl]acetamide

To a dichloromethane suspension (16 mL) of 1-amino-3-(4-fluorophenyl)propan-2-ol hydrochloride (0.75 g, 3.64 mmol) synthesized in Reference Synthesis Example 41, triethylamine (0.92 g, 9.10 mmol) and bromoacetyl bromide (0.88 g, 4.37 mmol) were added at 0°C and the resultant solution was stirred for 0.5 hours. After completion of the reaction, water was added thereto and extraction with chloroform from the resultant mixture was performed. The obtained organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to obtain the title compound (1.03 g, yield 97%).

[0266] Reference Synthesis Example 43

6-(4-Fluorobenzyl)morpholin-3-one

To an ethanol solution (16 mL) of 2-bromo-N-[3-(4-fluorophenyl)-2-hydroxypropyl]acetamide (1.00 g, 3.45 mmol) synthesized in Reference Synthesis Example 42, potassium carbonate (715 mg, 5.17

mmol) was added and the resultant solution was stirred at 80°C for 2.5 hours. After completion of the reaction, the reaction solution was filtered with Celite and the filtered residue was washed with ethanol. The filtrate was concentrated under reduced pressure and chloroform was added to the concentrated filtrate, followed by concentrating the filtrate under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 1/20 → ethyl acetate/methanol = 20/1 (v/v)) to obtain the title compound (100 mg, yield 14%).

[0267] Reference Synthesis Example 44

2-(4-Fluorobenzyl)morpholine

To a tetrahydrofuran solution (4 mL) of 6-(4-fluorobenzyl)morpholin-3-one (100 mg, 0.48 mmol) synthesized in Reference Synthesis Example 43, lithium aluminum hydride (21.8 mg, 0.57 mmol) was added at 0°C and the resultant solution was stirred for 4 hours. To the reaction solution, lithium aluminum hydride (20.0 mg) was added and the resultant mixture was stirred at room temperature for 1 hour. After completion of the reaction, saturated sodium sulfate aqueous solution was added to the reaction solution at 0°C and the resultant mixture was stirred at room temperature for 20 minutes. The reaction solution was dried over added anhydrous magnesium sulfate and concentrated under reduced pressure to obtain a crude product (84.9 mg) of the title compound.

[0268] Reference Synthesis Example 45

(R)-1-(4-Fluorobenzyl)pyrrolidin-3-amine hydrochloride

To a chloroform solution (2 mL) of tert-butyl (R)-pyrrolidin-3-yl-carbamate (500 mg, 2.68 mmol), 4-fluorobenzyl bromide (329 µL, 2.68 mmol) was added and the resultant solution was stirred at room temperature for 1 day. Subsequently, 4 M hydrogen chloride/1,4-dioxane solution (2 mL) was added to the reaction solution and the resultant mixture was stirred for 1 day. After completion of the reaction, solidified product was collected by filtration and dried under reduced pressure to obtain a crude product of the title compound.

[0269] Reference Synthesis Example 46

1-(4-Fluorophenyl)pyrrolidin-3-ol

Under nitrogen atmosphere, a 1,4-dioxane suspension of 1-fluoro-4-iodobenzene
 5 (2.00 g, 9.01mmol), pyrrolidin-3-ol (785 mg, 9.01 mmol), potassium phosphate (3.83 g,
 18.0 mmol), copper (I) iodide (343 mg, 1.80 mmol), copper (114 mg, 1.80 mmol), and
 N,N-dimethylaminoethanol (100mg) was stirred at 90°C for 1 day and at 110°C for 5
 hours. After completion of the reaction, the reaction solution was filtered with Celite
 and the filtrate was concentrated under reduced pressure. To the residue, water was
 10 added and extraction with ethyl acetate from the resultant mixture was performed. The
 organic layer was dried over anhydrous sodium sulfate and concentrated under reduced
 pressure to obtain a crude product of the title compound.

[0270] Reference Synthesis Example 47

(S)-1-(4-Fluorobenzyl)pyrrolidin-3-amine hydrochloride

15

tert-Butyl (S)-pyrrolidin-3-yl-carbamate (500 mg, 2.68 mmol) was used to obtain a
 crude product of the title compound by synthesis in a similar manner to Reference
 Synthesis Example 45.

[0271] Reference Synthesis Example 48

20 3-Methyl-1-([(1,1,1-trifluoro-2-methylpropan-2-yl)oxy]carbonyl]-1H-imidazol-3-ium
 iodide

To a chloroform solution (10 mL) of 1,1,1-trifluoro-2-methylpropan-2-ol (500 mg,
 3.90 mmol), 1,1'-carbonyldiimidazole (696 mg, 4.29 mmol) was added at room
 25 temperature and the resultant solution was stirred for 1 day. To the reaction solution,
 water was added and extraction with chloroform from the resultant mixture was
 performed twice. The organic layer was dried over anhydrous sodium sulfate and
 concentrated under reduced pressure. To an acetonitrile solution of the obtained residue,
 methyl iodide (2.21 g, 15.6 mmol) was added and the resultant mixture was stirred for 1

week. The reaction solution was concentrated under reduced pressure to obtain a crude product of the title compound and the crude product was used in the next reaction as it was.

[0272] Reference Synthesis Example 49

5 1-(4-Fluorophenyl)-3-hydroxypyrrolidin-2-one

A mixture of 3-hydroxydihydrofuran-2(3H)-one (5.00 g, 49.0 mmol) and 4-fluoroaniline (6.53 g, 59.0 mmol) was stirred at 150°C for 1 day. After completion of the reaction, dichloromethane and 10 M sodium hydroxide aqueous solution were added
10 to the reaction solution and the resultant mixture was filtered. To the filtrate, dichloromethane and concentrated hydrochloric acid were added to adjust pH to 1. The obtained solid was collected by filtration, washed with water, and dried under reduced pressure to obtain the title compound (6.78 g, yield 71%).

[0273] Reference Synthesis Example 50

15 3-Amino-1-(4-fluorophenyl)pyrrolidin-2-one

To a dichloromethane (10 mL) - acetonitrile (2 mL) mixed solution of 1-(4-fluorophenyl)-3-hydroxy-pyrrolidin-2-one (500 mg, 2.56 mmol) obtained in Reference Synthesis Example 49, triphenylphosphine (805 mg, 3.07 mmol), and
20 di-tert-butyl iminodicarboxylate (667 mg, 3.07 mmol), diethyl azodicarboxylate (1.34 g, 3.07 mmol, toluene solution) was added at room temperature and the resultant mixture was stirred for 1 day. Di-tert-butyl iminodicarboxylate (667 mg, 3.07 mmol) and diethyl azodicarboxylate (1.34 g, 3.07 mmol, toluene solution) were added and the resultant mixture was further stirred for 3 hours. After completion of the reaction, the reaction
25 solution was concentrated under reduced pressure and 4 M hydrogen chloride/dioxane solution (15 mL) was added to the obtained residue, followed by stirring the resultant mixture for 2 hours. Ethyl acetate was added thereto and extraction with water from the resultant mixture was performed twice. To the water phase, saturated sodium bicarbonate aqueous solution and 1 M sodium hydroxide aqueous solution were added to

adjust pH to 13 and extraction with chloroform from the resultant mixture was performed twice. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the title compound (145 mg, yield 25%).

[0274] Reference Synthesis Example 51

5 [5-(trifluoromethyl)thiophen-2-yl]methyl 4-methylbenzenesulfonate

To a tetrahydrofuran solution (4 mL) of 5-(trifluoromethyl)thiophene-2-carboxylic acid (196 mg, 1.00 mmol), lithium aluminum hydride (49.5 mg, 1.20 mmol) was added at room temperature and the resultant solution was stirred at room temperature for 3 days.

10 After completion of the reaction, water, ethyl acetate and 1 M hydrochloric acid were added thereto and extraction with ethyl acetate from the resultant mixture was performed three times. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to obtain a crude product (411 mg, yellow oily product) of [5-(trifluoromethyl)thiophen-2-yl]methanol. Subsequently, to a
15 tetrahydrofuran solution (4 mL) of the crude product of [5-(trifluoromethyl)thiophen-2-yl]methanol (411 mg, 1.00 mmol), triethylamine (279 μ L, 2.00 mmol) and p-toluenesulfonic acid anhydride (392 mg, 1.20 mmol) were added at room temperature and the resultant mixture was stirred at room temperature for 24 hours. The reaction solution was concentrated under reduced pressure to obtain a crude product
20 (828 mg) of the title compound. The crude product was used in the next reaction as it was.

[0275] Reference Synthesis Example 52

[4-(Trifluoromethyl)-1H-pyrazol-1-yl]methyl 4-methylbenzenesulfonate

25 To a tetrahydrofuran solution (2 mL) of 4-(trifluoromethyl)-1H-pyrazole, 1,8-diazabicyclo[5.4.0]undec-7-ene (10.9 μ L, 0.07 mmol) and paraformaldehyde (43.8 mg, 1.46 mmol) were added at room temperature and the resultant mixture was stirred at 60°C for 12 hours and at room temperature for 1 day. The reaction solution was filtered and triethylamine (153 μ L, 1.10 mmol) and p-toluenesulfonic acid anhydride (286 mg,

0.88 mmol) were added to the filtrate, following by stirring the resultant mixture for 30 minutes. After completion of the reaction, saturated sodium carbonate aqueous solution was added to the reaction solution and extraction with ethyl acetate from the resultant mixture was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. To the obtained residue, ethyl acetate was added and the resultant mixture was filtered and concentrated under reduced pressure to obtain a crude product (232 mg) of the title compound.

[0276] Reference Synthesis Example 53

Ethyl 2-[3-(trifluoromethyl)-1H-pyrazol-1-yl]acetate

To a chloroform solution (2 mL) of 3-(trifluoromethyl)-1H-pyrazole (68 mg, 0.50 mmol), potassium carbonate (104 mg, 0.75 mmol) and ethyl bromoacetate (100 mg, 0.60 mmol) were added and the resultant mixture was stirred at 70°C for 2 hours. To the reaction solution, adequate amount of sodium iodide was added and the resultant mixture was further stirred for 1 day. To the reaction solution, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (147 mg, 0.77 mmol), 1-hydroxybenzotriazole (catalytic amount) and ethanol (500 µL) were added and the resultant mixture was stirred for 2 hours. After completion of the reaction, water was added to the reaction solution and extraction with chloroform from the resultant mixture was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product of the title compound. The crude product was used in the next reaction as it was.

[0277] Reference Synthesis Example 54

2-[3-(Trifluoromethyl)-1H-pyrazol-1-yl]ethyl 4-methylbenzenesulfonate

To a tetrahydrofuran solution (4 mL) of the crude product of ethyl 2-[3-(trifluoromethyl)-1H-pyrazol-1-yl]acetate synthesized in Reference Synthesis Example 53, lithium aluminum hydride (40 mg, 1.05 mmol) was added at 0°C and the resultant solution was stirred for 3 hours. After completion of the reaction, saturated

sodium sulfate aqueous solution was added to the reaction solution and the resultant mixture was dried over anhydrous magnesium sulfate and filtered. The obtained filtrate was concentrated under reduced pressure. Subsequently, to a dichloromethane solution (2 mL) of the obtained residue, triethylamine (80 μ L, 0.57 mmol) and p-toluenesulfonyl chloride (300 mg, 1.57 mmol) were added and the resultant mixture was stirred at room temperature for 1 hour. The reaction solution was concentrated under reduced pressure and used in the next reaction.

[0278] Reference Synthesis Example 55

5-(Chloromethyl)-3-(trifluoromethyl)-1H-pyrazole

10

To a tetrahydrofuran solution (4 mL) of ethyl 3-(trifluoromethyl)-1H-pyrazol-5-carboxylate (104 mg, 0.50 mmol), diisobutylaluminum hydride (1.75 mL, 1.75 mmol, toluene solution) was added at 0°C and the resultant solution was stirred for 2 hours. To the reaction solution, saturated sodium sulfate aqueous solution was added and the resultant mixture was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. Subsequently, to a dichloromethane solution of the obtained residue (28 mg), thionyl chloride (39.4 mg, 0.33 mmol) was added and the resultant mixture was stirred at room temperature for 1 day. After completion of the reaction, the reaction solution was concentrated under reduced pressure to obtain a crude product of the title compound. The crude product was used in the next reaction as it was.

[0279] Reference Synthesis Example 56

1-(Chloromethyl)-5-phenyl-3-(trifluoromethyl)-1H-pyrazole

25 To a tetrahydrofuran solution (10 mL) of 5-phenyl-3-(trifluoromethyl)-1H-pyrazole (500 mg, 2.36 mmol), 1,8-diazabicyclo[5.4.0]undec-7-ene (35.2 μ L, 0.24 mmol) and paraformaldehyde (142 mg, 4.71 mmol) were added at room temperature and the resultant solution was stirred at room temperature for 1 day. The reaction solution was filtered and thionyl chloride (1.26 mL, 17.2 mmol) was added at room temperature to the

filtrate, followed by stirring the resultant mixture for 1 day. After completion of the reaction, the reaction solution was concentrated under reduced pressure to obtain a mixed solution of the title compound. The mixed solution was used in the next reaction as it was.

5 [0280] Reference Synthesis Example 57

[3-(Trifluoromethyl)-1H-pyrazol-1-yl]methyl 4-methylbenzenesulfonate

To an N,N-dimethylformamide solution (2 mL) of 3-(trifluoromethyl)-1H-pyrazole (100 mg, 0.73 mmol), sodium hydride (35.3 mg, 0.88 mmol, purity 55%) was added at
10 room temperature and the resultant solution was stirred for 1 hour. To the reaction solution, paraformaldehyde (26.4 mg, 0.88 mmol) was added and the resultant mixture was stirred for 1 day. Subsequently, triethylamine (204 μ L, 1.46 mmol) and p-toluenesulfonyl chloride (230 mg, 1.21 mmol) were added at 0°C to the reaction solution and the resultant mixture was stirred for 40 minutes. After completion of the
15 reaction, water was added to the reaction solution and extraction with ethyl acetate from the resultant mixture was performed. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to obtain the title compound (181 mg, yield 78%).

[0281] Reference Synthesis Example 58

20 [5-(2,2,2-Trifluoroethoxy)pyridin-2-yl]methyl methanesulfonate

To a dichloromethane solution (2.8 mL) of
[5-(2,2,2-trifluoroethoxy)pyridin-2-yl]methanol (138 mg, 0.67 mmol), triethylamine (468 μ L, 3.33 mmol) and methanesulfonyl chloride (77.4 μ L, 1.00 mmol) were added at 0°C
25 and the resultant solution was stirred for 30 minutes. After completion of the reaction, the reaction solution was washed with saturated ammonium chloride aqueous solution and brine. The obtained organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to obtain a crude product (129 mg) of the title compound.

[0282] Reference Synthesis Example 59

3-(tert-Butyl)-5-(chloromethyl)-1H-pyrazole

To a chloroform (20 mL) - dichloromethane (10 mL) mixed solution of
5 [3-(tert-butyl)-1H-pyrazol-5-yl]methanol (617 mg, 4.00 mmol), thionyl chloride (577 μ L, 8.01 mmol) was added at room temperature and the resultant solution was stirred at room temperature for 1 day. After completion of the reaction, the reaction solution was concentrated under reduced pressure to obtain a crude product (931 mg) of the title compound.

10 [0283] Reference Synthesis Example 60

[2-(Trifluoromethyl)thiazol-4-yl]methyl 4-methylbenzenesulfonate

To a dichloromethane solution (1.7 mL) of [2-(trifluoromethyl)thiazol-4-yl]methanol
(42.0 mg, 0.23 mmol), triethylamine (47.9 μ L, 0.34 mmol) and p-toluenesulfonic acid
15 anhydride (89.8 mg, 0.28 mmol) were added at 0°C and the resultant solution was stirred at room temperature for 4 hours. The reaction solution was concentrated under reduced pressure to obtain a crude product (191 mg) of the title compound.

[0284] Reference Synthesis Example 61

3-(tert-Butyl)-5-(chloromethyl)-1-methyl-1H-pyrazole

20

[3-(tert-butyl)-1-methyl-1H-pyrazol-5-yl]methanol (120 mg, 0.71 mmol) was used to obtain a crude product (50 mg) of the title compound by synthesis in a similar manner to Reference Synthesis Example 59.

[0285] Reference Synthesis Example 62

25 [2-(Trifluoromethyl)thiazol-5-yl]methyl 4-methylbenzenesulfonate

[2-(Trifluoromethyl)thiazol-5-yl]methanol (49.7 mg, 0.27 mmol) was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 60.

[0286] Reference Synthesis Example 63

1-(Chloromethyl)-4-(trifluoromethyl)pyridin-2(1H)-one

4-(Trifluoromethyl)pyridin-2-ol (100 mg, 0.61 mmol) was used to obtain a crude
5 product (134 mg) of the title compound by synthesis in a similar manner to Reference
Synthesis Example 56

[0287] Reference Synthesis Example 64

tert-Butyl 3-[(4-fluorophenyl)thio]azetidine-1-carboxylate

10 4-Fluorothiophenol (94.0 mg, 0.73 mmol) was used to obtain the title compound
(139 mg, yield 80%) by synthesis in a similar manner to Reference Synthesis Example
37.

[0288] Reference Synthesis Example 65

3-[(4-Fluorophenyl)thio]azetidine hydrochloride

15

To tert-butyl 3-[(4-fluorophenyl)thio]azetidine-1-carboxylate (139 mg, 0.49 mmol)
synthesized in Reference Synthesis Example 64, 4 M hydrogen chloride/1,4-dioxane
solution (1.0 mL) was added and the resultant solution was stirred at 0°C for 30 minutes
and at room temperature for 2 hours. After completion of the reaction, the reaction
20 solution was concentrated under reduced pressure to obtain a crude product (119 mg) of
the title compound.

[0289] Reference Synthesis Example 66

tert-Butyl 3-[(4-fluorobenzyl)thio]azetidine-1-carboxylate

25 4-Fluorobenzylmercaptan (68.8 μ L, 0.56 mmol) was used to obtain the title
compound (57.4 mg, yield 41%) by synthesis in a similar manner to Reference Synthesis
Example 35.

[0290] Reference Synthesis Example 67

3-[(4-Fluorobenzyl)thio]azetidine hydrochloride

tert-Butyl 3-[(4-fluorobenzyl)thio]azetidine-1-carboxylate (57.4 mg, 0.19 mmol) synthesized in Reference Synthesis Example 66 was used to obtain a crude product (51.2 mg) of the title compound by synthesis in a similar manner to Reference Synthesis

5 Example 65.

[0291] Reference Synthesis Example 68

6-(Chloromethyl)benzo[d]thiazole

10 To a dichloromethane solution (1.2 mL) of benzo[d]thiazol-6-yl-methanol (60 mg, 0.36 mmol), thionyl chloride (53 μ L) was added at room temperature and the resultant solution was stirred for 1 day. After completion of the reaction, the reaction solution was concentrated under reduced pressure to obtain a crude product of the title compound.

[0292] Reference Synthesis Example 69

(5-Chloro-benzofuran-2-yl)methanol

15

To a tetrahydrofuran solution (12 mL) of 5-chlorobenzofuran-2-carboxylic acid (620 mg, 3.15 mmol), borane-tetrahydrofuran complex (7.88 mL, 7.88 mmol, tetrahydrofuran solution) was added at 0°C and the resultant solution was stirred at room temperature for 3 hours, at 60°C for 1 day, and at room temperature for 1 day. After completion of the
20 reaction, concentration under reduced pressure was carried out three times with adding methanol to the reaction solution and the concentrated solution was purified by silica gel column chromatography (hexane/ethyl acetate = 3/1 $\rightarrow$ 1/1 (v/v)) to obtain the title compound (268 mg, yield 47%).

[0293] Reference Synthesis Example 70

25 5-Chloro-2-(chloromethyl)benzofuran

(5-Chloro-benzofuran-2-yl)methanol (60 mg, 0.33 mmol) synthesized in Reference Synthesis Example 69 was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 68.

[0294] Reference Synthesis Example 71

tert-Butyl (3R,4R)-3-(dibenzylamino)-4-hydroxypyrrolidine-1-carboxylate

To an N,N-dimethylformamide solution (2.5 mL) of tert-butyl
5 (3R,4R)-3-(benzylamino)-4-hydroxypyrrolidine-1-carboxylate (500 mg, 1.70 mmol),
benzyl bromide (203 μ L, 1.70 mmol) was added at room temperature and the resultant
solution was stirred for 1 day. To the reaction solution, tetrabutylammonium iodide
(188 mg, 0.51 mmol) and potassium carbonate (470 mg, 3.40 mmol) were added and the
resultant mixture was stirred for 1 day. After completion of the reaction, water was
10 added to the reaction solution and extraction with ethyl acetate from the resultant mixture
was performed. The organic layer was washed with brine and concentrated under
reduced pressure. The obtained residue was purified by silica gel column
chromatography (hexane/ethyl acetate = 1/0 $\rightarrow$ 1/1 (v/v)) to obtain the title compound
(698 mg, quantitative).

15 [0295] Reference Synthesis Example 72

(3R,4R)-4-(Dibenzylamino)pyrrolidin-3-ol

To tert-butyl (3R,4R)-3-(dibenzylamino)-4-hydroxy-pyrrolidine-1-carboxylate (350
mg, 0.91 mmol) synthesized in Reference Synthesis Example 71, 4 M hydrogen
20 chloride/1,4-dioxane solution (3 mL) was added and the resultant solution was stirred at
room temperature. The reaction solution was concentrated under reduced pressure and
chloroform was added to the obtained residue, followed by washing the resultant mixture
with saturated sodium bicarbonate aqueous solution. The organic layer was dried over
anhydrous magnesium sulfate and concentrated under reduced pressure to obtain a crude
25 product (258 mg) of the title compound.

[0296] Reference Synthesis Example 73

(3R,4R)-4-(Dibenzylamino)-1-(4-fluorophenyl)pyrrolidin-3-ol

(3R,4R)-4-(Dibenzylamino)pyrrolidin-3-ol (461 mg, 1.63 mmol) synthesized in

Reference Synthesis Example 72 was used to obtain the title compound (380 mg, yield 47%) by synthesis in a similar manner to Reference Synthesis Example 25.

[0297] Reference Synthesis Example 74

(3R,4R)-4-Amino-1-(4-fluorophenyl)pyrrolidin-3-ol

5

A methanol suspension (4.6 mL) of (3R,4R)-4-(dibenzylamino)-1-(4-fluorophenyl)pyrrolidin-3-ol (230 mg, 0.61 mmol) synthesized in Reference Synthesis Example 73 and palladium hydroxide-activated carbon catalyst (23 mg) was stirred under hydrogen atmosphere at 50°C for 1 day. After completion of the reaction, the suspension was filtered with Celite and the filtrate was concentrated under reduced pressure to obtain a crude product (146 mg) of the title compound.

10

[0298] Reference Synthesis Example 75

tert-Butyl (3R,4R)-3-(dibenzylamino)-4-fluoropyrrolidine-1-carboxylate

15

To a dichloromethane solution (8.7 mL) of tert-butyl (3R,4R)-3-(dibenzylamino)-4-hydroxypyrrolidine-1-carboxylate (320 mg, 0.92 mmol) synthesized in Reference Synthesis Example 71, (diethylamino)sulfur trifluoride (295 mg, 1.84 mmol) was added at 0°C and the resultant solution was stirred at room temperature. After completion of the reaction, the reaction solution was washed with water and saturated sodium bicarbonate aqueous solution. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to obtain a crude product (240 mg) of the title compound. The crude product was used in the next reaction as it was.

20

[0299] Reference Synthesis Example 76

25 (3R,4R)-N,N-Dibenzyl-4-fluoropyrrolidin-3-amine

tert-Butyl (3R,4R)-3-(dibenzylamino)-4-fluoropyrrolidine-1-carboxylate (240 mg) synthesized in Reference Synthesis Example 75 was used to obtain crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 72.

The crude product was used in the next process as it was.

[0300] Reference Synthesis Example 77

(3R,4R)-N,N-Dibenzyl-4-fluoro-1-(4-fluorophenyl)pyrrolidin-3-amine

- 5 (3R,4R)-N,N-Dibenzyl-4-fluoropyrrolidin-3-amine (177 mg, 0.62 mmol) synthesized in Reference Synthesis Example 76 was used to obtain the title compound (190 mg, 97%yield) by synthesis in a similar manner to Reference Synthesis Example 25.

[0301] Reference Synthesis Example 78

(3R,4R)-4-Fluoro-1-(4-fluorophenyl)pyrrolidin-3-amine

10

(3R,4R)-N,N-Dibenzyl-4- fluoro-1-(4-fluorophenyl)pyrrolidin-3-amine (190 mg) synthesized in Reference Synthesis Example 77 was used to obtain the title compound (57 mg, yield 57%) by synthesis in a similar manner to Reference Synthesis Example 74.

[0302] Reference Synthesis Example 79

- 15 tert-Butyl [1-(4-fluorophenyl)pyrrolidin-3-yl]carbamate

tert-Butyl pyrrolidin-3-yl-carbamate (254 mg, 1.37 mmol) was used to obtain the title compound (217 mg, yield 68%) by synthesis in a similar manner to Reference Synthesis Example 25.

- 20 [0303] Reference Synthesis Example 80

tert-Butyl [1-(4-fluorophenyl)pyrrolidin-3-yl](methyl)carbamate

- To an N,N-dimethylformamide solution (3.9 mL) of tert-butyl [1-(4-fluorophenyl)pyrrolidin-3-yl]carbamate (195 mg, 0.70 mmol) synthesized in Reference Synthesis Example 79, sodium hydride (28 mg, 0.70 mmol, purity 60%) and methyl iodide (44 μ L, 0.70 mmol) were added at 0°C and the resultant solution was stirred at room temperature for 1 day. To the reaction solution, the same amounts of sodium hydride and methyl iodide were further added and the resultant mixture was stirred for 1 day. After completion of the reaction, ethyl acetate was added to the
- 25

reaction solution and the resultant mixture was washed with water and brine. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel (amino-based) column chromatography (hexane/ethyl acetate = 9/1) to obtain the title compound (181 mg, yield 88%).

[0304] Reference Synthesis Example 81

1-(4-Fluorophenyl)-N-methyl-pyrrolidin-3-amine

tert-Butyl [1-(4-fluorophenyl)pyrrolidin-3-yl](methyl)carbamate (84 mg, 0.28 mmol) synthesized in Reference Synthesis Example 80 was used to obtain crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 72.

[0305] Reference Synthesis Example 82

tert-Butyl [1-(4-fluorophenyl)piperidin-4-yl]carbamate

tert-Butyl piperidin-4-yl-carbamate (1.00 g, 5.00 mmol) was used to obtain the title compound (495 mg, yield 37%) by synthesis in a similar manner to Reference Synthesis Example 25.

[0306] Reference Synthesis Example 83

1-(4-Fluorophenyl)piperidin-4-amine

tert-Butyl [1-(4-fluorophenyl)piperidin-4-yl]carbamate (130 mg, 0.440 mmol) synthesized in Reference Synthesis Example 82 was used to obtain the title compound (67 mg, yield 78%) by synthesis in a similar manner to Reference Synthesis Example 72.

[0307] Reference Synthesis Example 84

2,4-Dichloro-6-[4-(2,4-difluorophenyl)piperazin-1-yl]-1,3,5-triazine

1,3,5-Trichlorotriazine (1.25 g, 6.80 mmol) and 1-(2,4-difluorophenyl)piperazine (0.40 g, 2.00 mmol) were used to obtain a crude product (1.00 g) of the title compound by synthesis in a similar manner to Reference Synthesis Example 1.

[0308] Reference Synthesis Example 85

6-Chloro-4-[4-(2,4-difluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

The crude product of

- 5 2,4-dichloro-6-[4-(2,4-difluorophenyl)piperazin-1-yl]-1,3,5-triazine (1.00 g, 2.00 mmol) synthesized in Reference Synthesis Example 84 was used to obtain the title compound (0.59 g, two step yield 90%) by synthesis in a similar manner to Reference Synthesis Example 2.

[0309] Reference Synthesis Example 86

- 10 4-[4-(2,4-Difluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

- 6-Chloro-4-[4-(2,4-difluorophenyl)-piperazin-1-yl]-1,3,5-triazin-2(1H)-one (0.59 g, 1.81 mmol) synthesized in Reference Synthesis Example 85 was used to obtain the title compound (0.19 g, yield 36%) by synthesis in a similar manner to Reference Synthesis
15 Example 3.

[0310] Reference Synthesis Example 87

1-tert-Butyl 3-methyl 4-

[5-(4-chlorobenzyl)-4-oxo-4,5-dihydro-1,3,5-triazin-2-yl]piperazine-1,3-dicarboxylate

- 20 To tetrahydrofuran solution (2.3 mL) of 1-tert-butyl 3-methyl piperazine-1,3-dicarboxylate (43 mg, 0.17 mmol), sodium carbonate (37 mg, 0.35 mmol) and 4-chloro-1-(4-chlorobenzyl)-1,3,5-triazin-2(1H)-one (45 mg, 0.17 mmol) synthesized in Reference Synthesis Example 11 were added and the resultant solution was stirred at room temperature for 1 day. After completion of the reaction, water was added thereto
25 and extraction with chloroform from the resultant mixture was performed three times. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure and ethyl acetate was added to the obtained residue. The obtained solid was collected by filtration and dried under reduced pressure to obtain the title compound (34 mg, yield 42%).

[0311] Reference Synthesis Example 88

Methyl

1-[5-(4-chlorobenzyl)-4-oxo-4,5-dihydro-1,3,5-triazin-2-yl]piperazine-2-carboxylate

5 1-tert-Butyl 3-methyl

4-[5-(4-chlorobenzyl)-4-oxo-4,5-dihydro-1,3,5-triazin-2-yl]piperazine-1,3-dicarboxylate (34 mg, 0.07 mmol) synthesized in Reference Synthesis Example 87 was used to obtain crude product (42 mg) of the title compound by synthesis in a similar manner to Reference Synthesis Example 9. The crude product was used in the next reaction as it was.

[0312] Reference Synthesis Example 89

4-[4-(4-Fluorophenyl)-4-hydroxypiperidin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

15 4-[4-(4-Fluorophenyl)-4-hydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one (426 mg, 1.18 mmol) synthesized in Reference Synthesis Example 14 and 1-(chloromethyl)-3-(trifluoromethyl)-1H-pyrazole (219 mg, 1.18 mmol) were used to obtain the title compound (158 mg, yield 30%) by synthesis in a similar manner to Reference Synthesis Example 8.

20 [0313] Reference Synthesis Example 90

4-[4-(4-Fluorophenyl)-4-hydroxypiperidin-1-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25 4-[4-(4-Fluorophenyl)-4-hydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one (100 mg, 0.34 mmol) synthesized in Reference Synthesis Example 14 and [5-(trifluoromethyl)thiophen-2-yl]methyl 4-methylbenzenesulfonate (558 mg, 0.67 mmol) synthesized in Reference Synthesis Example 51 were used to obtain the title compound (20 mg, yield 13%) by synthesis in a similar manner to Reference Synthesis Example 8.

[0314] Reference Synthesis Example 91a

4-[4-(4-Chlorophenyl)-4-hydroxypiperidin-1-yl]-1-{{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Reference Synthesis Example 91b

5 4-(4-Hydroxy-4-phenyl)piperidin-1-yl)-1-{{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

An N,N-dimethylformamide suspension (1.4 mL) of a mixture of 4-[4-(4-chlorophenyl)-4-hydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 22a and 4-(4-hydroxy-4-phenylpiperidin-1-yl)-1,3,5-triazin-2(1H)-one (140 mg), [3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl 4-methylbenzenesulfonate (282 mg, 0.91 mmol) synthesized in Reference Synthesis Example 15, and potassium carbonate (252 mg, 1.83 mmol) was stirred at 80°C for 3 hours. After completion of the reaction, water was added thereto and extraction with ethyl acetate from the resultant mixture was performed. The obtained organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 1/4 (v/v)) and preparative high-performance liquid chromatography to obtain the title compound (14 mg, yield 6.8%) in Reference Synthesis Example 91a and the title compound (7.4 mg, yield 3.9%) in Reference Synthesis Example 91b.

[0315] Reference Synthesis Example 92a

4-(4-Hydroxy-4-phenyl)piperidin-1-yl)-1-{{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25

Reference Synthesis Example 92 b

4-[4-(4-Chlorophenyl)-4-hydroxy-piperidin-1-yl]-1-{{[5-(trifluoromethyl)thiophen-2-yl]methyl}1,3,5-triazin-2(1H)-one

A mixture (100 mg) of

4-[4-(4-Chlorophenyl)-4-hydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 22a and

4-(4-hydroxy-4-phenylpiperidin-1-yl)-1,3,5-triazin-2(1H)-one,

5 [5-(trifluoromethyl)thiophen-2-yl]methyl 4-methylbenzenesulfonate (558 mg, 0.65 mmol) synthesized in Reference Synthesis Example 51, and potassium carbonate (180 mg, 1.30 mmol) were used to obtain a mixture of compounds in Reference Synthesis Examples 92a and 92b being the title compounds (21 mg) by synthesis in a similar manner to Reference Synthesis Example 91.

10 [0316] Reference Synthesis Example 93

5-(Hydroxymethyl)pyridin-2-yl trifluoromethanesulfonate

To a tetrahydrofuran solution (2.0 mL) of lithium aluminum hydride (19.0 mg, 0.450 mmol), ethyl 6-{{(trifluoromethyl)sulfonyl}oxy}nicotinate (112 mg, 0.375 mmol) was
15 added and the resultant solution was stirred at 0°C for 5 minutes. After completion of the reaction, 1 M sodium hydroxide aqueous solution, Celite, and ethyl acetate were added to the reaction solution and the resultant mixture was filtered with Celite, followed by extraction from the filtrate with ethyl acetate. The organic layer was washed with
20 obtain a crude product of the title compound (62.9 mg).

[0317] Reference Synthesis Example 94

5-(Chloromethyl)pyridin-2-yl trifluoromethanesulfonate

5-(Hydroxymethyl)pyridin-2-yl trifluoromethanesulfonate (62.9 mg, 0.245 mmol)
25 synthesized in Reference Synthesis Example 93 was used to obtain a crude product of the title compound (64.4 mg), by synthesis in a similar manner to Reference Synthesis Example 68.

[0318] Reference Synthesis Example 95

2-(Trifluoromethyl)benzofuran-5-carbaldehyde

To a tetrahydrofuran solution (12 mL) of 5-bromo-2(trifluoromethyl)benzofuran (587 mg, 2.21 mmol), n-butyl lithium (hexane solution, 1.59 mol/L) (1.70 mL, 2.66 mmol) was added at -78°C and the resultant solution was stirred for 40 minutes. To the reaction solution, N,N-dimethylformamide (257 µL, 3.32 mmol) was added and the resultant mixture was stirred for 3 hours while the temperature of the mixture was raised to room temperature. After completion of the reaction, saturated ammonium chloride aqueous solution was added to the reaction solution and extraction with ethyl acetate from the resultant mixture was performed. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate) to obtain the title compound (180 mg, yield 39%).

[0319] Reference Synthesis Example 96

[2-(Trifluoromethyl)benzofuran-5-yl]methanol

To an ethanol solution (2.0 mL) of 2-(trifluoromethyl)benzofuran-5-carbaldehyde (220 mg, 1.03 mmol) synthesized in Reference Synthesis Example 95, sodium borohydride (115 mg, 3.05 mmol) was added and the resultant solution was stirred at 70°C for 1 hour and 30 minutes. After completion of the reaction, the reaction solution was cooled to room temperature and acetic acid was added to the cooled solution, followed by concentrating the resultant mixture under reduced pressure. To the obtained residue, water was added and extraction with dichloromethane from the resultant mixture was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the title compound (219 mg, yield 99%).

[0320] Reference Synthesis Example 97

5-(Chloromethyl)-2-(trifluoromethyl)benzofuran

[2-(Trifluoromethyl)benzofuran-5-yl]methanol (30.0 mg, 0.139 mmol) synthesized in Reference Synthesis Example 96 was used to obtain a crude product of the title

compound by synthesis in a similar manner to Reference Synthesis Example 68.

[0321] Reference Synthesis Example 98

[2-(Trifluoromethyl)-2,3-dihydrobenzofuran-5-yl]methanol

5 To an ethanol solution (1.0 mL) of [2-(trifluoromethyl)benzofuran-5-yl]methanol (30.0 mg, 0.139 mmol) synthesized in Reference Synthesis Example 96, palladium-activated carbon (15.0 mg) was added and the resultant solution was stirred under hydrogen atmosphere at room temperature for 30 hours. After completion of the reaction, the reaction solution was filtered with Celite and the filtrate was concentrated
10 under reduced pressure to obtain a crude product of the title compound (15.9 mg).

[0322] Reference Synthesis Example 99

5-(Chloromethyl)-2-(trifluoromethyl)-2,3-dihydrobenzofuran

The crude product of [2-(trifluoromethyl)-2,3-dihydrobenzofuran-5-yl]methanol
15 (15.9 mg) synthesized in Reference Synthesis Example 98 was used to obtain a crude product of the title compound (15.8 mg) by synthesis in a similar manner to Reference Synthesis Example 68.

[0323] Reference Synthesis Example 100

(5-Bromo-4-methylthiophen-2-yl)methanol

20

Methyl 5-bromo-4-methylthiophene-2-carboxylate (100 mg, 0.425 mmol) was used to obtain the title compound (33.7 mg, yield 38%) by synthesis in a similar manner to Reference Synthesis Example 96.

[0324] Reference Synthesis Example 101

25 2-Bromo-5-(bromomethyl)-3-methylthiophene

To a tetrahydrofuran solution (1.0 mL) of (5-bromo-4-methylthiophen-2-yl)methanol (33.7 mg, 0.163 mmol) synthesized in Reference Synthesis Example 100, carbon tetrabromide (59.4 mg, 0.179 mmol) and

triphenylphosphine (47.0 mg, 0.179 mmol) were added and the resultant solution was stirred at room temperature for 20 hours. After completion of the reaction, the reaction solution was filtered and the obtained filtrate was concentrated under reduced pressure to obtain the title compound as a crude product.

5 [0325] Reference Synthesis Example 102

[4-Bromo-5-(trifluoromethyl)thiophen-2-yl]methanol

To a tetrahydrofuran solution (5.0 mL) of 3-bromo-2-(trifluoromethyl)thiophene (500 μ L, 2.08 mmol), lithium diisopropylamide (2.06 mL, 2.29 mmol) was added under
 10 nitrogen atmosphere at -40°C, and the resultant mixture was stirred for 30 minutes. To the reaction solution, paraformaldehyde (68.8 mg, 2.29 mmol) was added and the resultant mixture was stirred for 30 minutes, followed by raising the temperature of the mixture to -10°C. The mixture was stirred at -10°C for 4 hours and at 0°C for 1 hour. After completion of the reaction, a 1 M hydrochloric acid was added to the reaction
 15 solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was washed with a saturated sodium chloride aqueous solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate) to obtain the title compound (106 mg, yield 20%).

20 [0326] Reference Synthesis Example 103

3-Bromo-5-(bromomethyl)-2-(trifluoromethyl)thiophene

[4-Bromo-5-(trifluoromethyl)thiophen-2-yl]methanol (106 mg, 0.407 mmol) synthesized in Reference Synthesis Example 102 was used to obtain a crude product of
 25 the title compound by synthesis in a similar manner to Reference Synthesis Example 101.

[0327] Reference Synthesis Example 104

2-(Bromomethyl)-5-fluorothiophene

(5-Fluorothiophen-2-yl)methanol (29.8 mg, 0.225 mmol) was used to obtain a crude

product of the title compound by synthesis in a similar manner to Reference Synthesis Example 101.

[0328] Reference Synthesis Example 105

[5-(Tetrahydrofuran-2-yl)thiophen-2-yl]methanol

5

5-(Tetrahydrofuran-2-yl)thiophene-2-carbaldehyde, (50.0 mg, 0.274 mmol) was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 96.

[0329] Reference Synthesis Example 106

10 2-[5-(Chloromethyl)thiophen-2-yl]tetrahydrofuran

[5-(Tetrahydrofuran-2-yl)thiophen-2-yl]methanol synthesized in Reference Synthesis Example 105 was used to obtain the title compound by synthesis in a similar manner to Reference Synthesis Example 68. The title compound was used in the next reaction as it was.

15

[0330] Reference Synthesis Example 107

Ethyl 5-(2,2,2-trifluoroethoxy)thiophene-2-carboxylate

20

To an N,N-dimethylformamide solution (30 mL) of 2,2,2,-trifluoroethanol (1.50 g, 15.0 mmol), sodium hydride (655 mg, 15.0 mmol) and ethyl 5-chlorothiophene-2-carboxylate (1.91 g, 10.0 mmol) were added and the resultant mixture was stirred at 65°C for 6 hours, at room temperature for 15.5 hours, and at 65°C for 4 hours. After completion of the reaction, water and 1 M hydrochloric acid were added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was washed with a saturated sodium chloride aqueous solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate) to obtain the title compound (464 mg, yield 18%).

25

[0331] Reference Synthesis Example 108

[5-(2,2,2-Trifluoroethoxy)thiophen-2-yl]methanol

Ethyl 5-(2,2,2-trifluoroethoxy)thiophene-2-carboxylate (254 mg, 1.00 mmol) synthesized in Reference Synthesis Example 107 was used to obtain the title compound (95.2 mg, yield 45%) by synthesis in a similar manner to Reference Synthesis Example 93.

[0332] Reference Synthesis Example 109

2-(Bromomethyl)-5-(2,2,2-trifluoroethoxy)thiophene

[5-(2,2,2-Trifluoroethoxy)thiophen-2-yl]methanol (95.0 mg, 0.448 mmol) synthesized in Reference Synthesis Example 108 was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 101.

[0333] Reference Synthesis Example 110

[5-(2,2,2-Trifluoroethoxy)pyrazin-2-yl]methanol

Methyl 5-(2,2,2-trifluoroethoxy)pyrazine-2-carboxylate (582 mg, 2.46 mmol) was used to obtain the title compound (73.9 mg, yield 14%) by synthesis in a similar manner to Reference Synthesis Example 93.

[0334] Reference Synthesis Example 111

2-(Chloromethyl)-5-(2,2,2-trifluoroethoxy)pyrazine

[5-(2,2,2-Trifluoroethoxy)pyrazin-2-yl]methanol (66.3 mg, 0.319 mmol) synthesized in Reference Synthesis Example 110 was used to obtain the title compound (39.5 mg, yield 55%) by synthesis in a similar manner to Reference Synthesis Example 68.

[0335] Reference Synthesis Example 112

Methyl 2-(2,2,2-trifluoroethoxy)thiazole-5-carboxylate

To a tetrahydrofuran solution (5.0 mL) of sodium hydride (465 mg, 11.6 mmol) and 2,2,2-trifluoroethanol (1.70 mL, 23.3 mmol), methyl 2-bromothiazole-5-carboxylate (517

mg, 2.33 mmol) was added and the resultant mixture was stirred at room temperature for 2 hours. After completion of the reaction, 1 M sodium hydroxide aqueous solution was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was washed with water, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to obtain a crude product of the title compound (444 mg).

[0336] Reference Synthesis Example 113

[2-(2,2,2-Trifluoroethoxy)thiazol-5-yl]methanol

10 Methyl 2-(2,2,2-trifluoroethoxy)thiazole-5-carboxylate (413 mg, 1.71 mmol) synthesized in Reference Synthesis Example 112 was used to obtain a crude product of the title compound (405 mg) by synthesis in a similar manner to Reference Synthesis Example 93.

[0337] Reference Synthesis Example 114

15 5-(Chloromethyl)-2-(2,2,2-trifluoroethoxy)thiazole

[2-(2,2,2-trifluoroethoxy)thiazol-5-yl]methanol (50.0 mg, 0.235 mmol) synthesized in Reference Synthesis Example 113 was used to obtain a crude product of the title compound (50.3 mg) by synthesis in a similar manner to Reference Synthesis Example 20 68.

[0338] Reference Synthesis Example 115

Methyl 6-[(2,2,2-trifluoroethoxy)methyl]nicotinate

To an N,N-dimethylformamide solution (10 mL) of methyl 25 6-(hydroxymethyl)nicotinate (500 mg, 2.99 mmol) and 2,2,2-trifluoroethyl trifluoromethanesulfonate (518 μ L, 3.59 mmol), sodium hydride (179 mg, 4.49 mmol) was added at 0°C and the resultant mixture was stirred at room temperature for 4 hours. After completion of the reaction, water was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was

dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 5/1) to obtain the title compound (171 mg, yield 23%).

[0339] Reference Synthesis Example 116

5 {6-[(2,2,2-Trifluoroethoxy)methyl]pyridin-3-yl}methanol

Methyl 6-[(2,2,2-trifluoroethoxy)methyl]nicotinate (171 mg, 0.685 mmol)

synthesized in Reference Synthesis Example 115 was used to obtain the title compound (134 mg, yield 88%) by synthesis in a similar manner to Reference Synthesis Example

10 93.

[0340] Reference Synthesis Example 117

5-(Chloromethyl)-2-[(2,2,2-trifluoroethoxy)methyl]pyridine

{6-[(2,2,2-Trifluoroethoxy)methyl]pyridin-3-yl}methanol (134 mg, 0.604 mmol)

15 synthesized in Reference Synthesis Example 116 was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 68.

[0341] Reference Synthesis Example 118

3-(1,1-Difluoroethyl)-1H-pyrazole

20 To a dichloroethane solution (6.0 mL) of 1-(1H-pyrazol-3-yl)ethanone (300 mg, 2.73 mmol) and bis(2-methoxyethyl)amino-sulfur trifluoride (1.11 mL, 5.99 mmol), one drop of ethanol was added at 0°C and the resultant mixture was stirred at room temperature for 1 hour and 30 minutes. After completion of the reaction, the reaction solution was added dropwise to saturated sodium bicarbonate aqueous solution at 0°C. After adding
25 dichloromethane, the resultant mixture was washed with saturated sodium chloride aqueous solution. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was purified by column chromatography (hexane/ethyl acetate = 3/1) to obtain the title compound (145 mg, yield 41%).

[0342] Reference Synthesis Example 119

[3-(1,1-Difluoroethyl)-1H-pyrazol-1-yl]methanol

To an ethanol solution (1.0 mL) of 3-(1,1-difluoroethyl)-1H-pyrazole (145 mg, 1.09 mmol) synthesized in Reference Synthesis Example 118, a formalin aqueous solution (1.0 mL) was added and the resultant mixture was stirred at 70°C for 1 hour. After completion of the reaction, the reaction solution was concentrated under reduced pressure to obtain the title compound as a crude product.

[0343] Reference Synthesis Example 120

10 1-(Chloromethyl)-3-(1,1-difluoroethyl)-1H-pyrazole

The crude product of [3-(1,1-difluoroethyl)-1H-pyrazol-1-yl]methanol synthesized in Reference Synthesis Example 119 was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 68.

15 [0344] Reference Synthesis Example 121

1-[5-(Trifluoromethyl)thiophen-2-yl]ethanol

To a tetrahydrofuran solution (2.0 mL) of 5-(trifluoromethyl)thiophene-2-carbaldehyde (80.0 mg, 0.444 mmol), methylmagnesium bromide (533 μ L, 0.444 mmol) was added at 0°C and the resultant mixture was stirred at room temperature for 5 minutes. After completion of the reaction, acetic acid was added to the reaction solution and the resultant mixture was concentrated under reduced pressure. To the obtained residue, water was added and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was concentrated under reduced pressure to obtain a crude product of the title compound.

[0345] Reference Synthesis Example 122

2-(1-Chloroethyl)-5-(trifluoromethyl)thiophene

The crude product of 1-[5-(trifluoromethyl)thiophen-2-yl]ethanol (30.0 mg)

synthesized in Reference Synthesis Example 121 was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 68.

[0346] Reference Synthesis Example 123

Ethyl 5-(1,1-difluoroethyl)thiophene-2-carboxylate

5

To a toluene (630 μ L) - ethanol (150 μ L) solution of ethyl 5-acetylthiophene-2-carboxylate (74.1 mg, 0.374 mmol), bis(2-methoxyethyl)amino-sulfur trifluoride (689 μ L, 3.74 mmol) was added and the resultant mixture was stirred at 70°C for 4 hours. After completion of the reaction, the reaction solution was added dropwise to saturated sodium bicarbonate aqueous solution at 0°C and extraction from the resultant mixture with dichloromethane was performed. The organic layer was concentrated under reduced pressure to obtain a crude product of the title compound. The crude product was used in the next step as it was.

[0347] Reference Synthesis Example 124

15 5-(1,1-Difluoroethyl)thiophene-2-carboxylic acid

To a methanol solution (1.0 mL) of ethyl 5-(1,1-difluoroethyl)thiophene-2-carboxylate (28.6 mg, 0.130 mmol) synthesized in Reference Synthesis Example 123, 1 M sodium hydroxide aqueous solution (312 μ L, 0.312 mmol) was added and the resultant mixture was stirred at room temperature for 1 day. After completion of the reaction, the reaction solution was concentrated under reduced pressure and separated with toluene-water. To the water phase, 12 M hydrochloric acid was added until pH reached 2 and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was concentrated under reduced pressure to obtain a crude product of the title compound. The crude product was used in the next step as it was.

[0348] Reference Synthesis Example 125

[5-(1,1-Difluoroethyl)thiophen-2-yl]methanol

5-(1,1-Difluoroethyl)thiophene-2-carboxylic acid synthesized in Reference Synthesis Example 124 was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 217.

[0349] Reference Synthesis Example 126

5 2-(Chloromethyl)-5-(1,1-difluoroethyl)thiophene

[5-(1,1-Difluoroethyl)thiophen-2-yl]methanol (100 mg, 0.561 mmol) synthesized in Reference Synthesis Example 125 was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 68.

10 [0350] Reference Synthesis Example 127

4-(Chloromethyl)phenyl trifluoromethanesulfonate

4-(Hydroxymethyl)phenyl trifluoromethanesulfonate (843 mg, 3.34 mmol) was used to obtain a crude product of the title compound (1.50 g) by synthesis in a similar manner to Reference Synthesis Example 68.

[0351] Reference Synthesis Example 128

[(2-Bromo-5-fluorobenzyl)oxy] (tert-butyl) dimethyl silane

To an acetonitrile solution (2.0 mL) of (2-bromo-5-fluorophenyl)methanol (820 mg, 4.00 mmol), triethylamine (835 μ L, 6.00 mmol) and tert-butyl dimethyl silyl trifluoromethanesulfonate (1.27 g, 4.80 mmol) were added and the resultant mixture was stirred at room temperature for 1 day. After completion of the reaction, saturated ammonium chloride aqueous solution was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 95/5 $\rightarrow$ 85/15) to obtain the title compound (1.01 g, yield 79%).

[0352] Reference Synthesis Example 129

tert-Butyl ((S,4S)-4-(4-fluorophenyl)-3,4-dihydropiperidine-1-carboxylate

To a tert-butanol (29 mL) - water (29 mL) solution of AD-mix- α (8.10 g, manufactured by Aldrich Chemical Company, Inc.) and methanesulfonamide (522 mg, 5.49 mmol), tert-butyl 4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine-1(2H)-carboxylate (1.50 g, 5.49 mmol) was added at 0°C and the resultant mixture was stirred at 0°C for 30 minutes and at room temperature for 2 hours and 30 minutes. After completion of the reaction, water was added to the reaction solution and extraction from the resultant mixture with dichloromethane was performed. The organic layer was washed with 1 M sodium hydroxide aqueous solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1 $\rightarrow$ 1/1) to obtain the title compound (1.66 g, yield 97%).

[0353] Reference Synthesis Example 130

tert-Butyl (3S,4S)-4-(4-fluorophenyl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate

To a dichloromethane solution (17 mL) of tert-butyl (3S,4S)-4-(4-fluorophenyl)-3,4-dihydroxypiperidine-1-carboxylate (1.66 g, 5.33 mmol) synthesized in Reference Synthesis Example 129 and 4-dimethylaminopyridine (33.0 mg, 0.270 mmol), pivalic acid anhydride (1.29 g, 6.93 mmol) and triethylamine (1.49 mL, 10.7 mmol) were added and the resultant mixture was stirred at room temperature for 18 hours and 30 minutes. After completion of the reaction, ethyl acetate was added to the reaction solution and the resultant mixture was washed with each of saturated ammonium chloride aqueous solution and saturated sodium chloride aqueous solution. The obtained organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 9/1 $\rightarrow$ 3/1) to obtain the title compound (2.07 g, yield 98%).

[0354] Reference Synthesis Example 131

tert-Butyl (R)-4-(4-fluorophenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate

To a toluene solution (40 mL) of tert-butyl (3S,4S)-4-(4-fluorophenyl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate (2.07 g, 5.23 mmol) synthesized in Reference Synthesis Example 130, methyl

- 5 N-(triethylammoniosulfonyl)carbamate (1.74 g, 7.32 mmol) was added and the resultant mixture was stirred at 60°C for 3 hours. After completion of the reaction, ethyl acetate was added to the reaction solution and the resultant mixture was washed with each of saturated sodium bicarbonate aqueous solution and saturated sodium chloride aqueous solution. The obtained organic layer was dried over anhydrous sodium sulfate and
- 10 concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 9/1 → 3/1) to obtain the title compound (1.47 g, yield 75%).

[0355] Reference Synthesis Example 132

(R)-4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl pivalate

15

To a dichloromethane solution (3.0 mL) of tert-butyl (R)-4-(4-fluorophenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate (1.69 g, 4.48 mmol) synthesized in Reference Synthesis Example 131, trifluoroacetic acid (1.4 mL) was added and the resultant mixture was stirred at room temperature for 2 hours.

- 20 After completion of the reaction, saturated sodium bicarbonate aqueous solution was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the title compound as a crude product.

[0356] Reference Synthesis Example 133

- 25 (R)-4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol

To a 1,4-dioxane (4.5 mL) - water (2.5 mL) solution of the crude product of (R)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl pivalate synthesized in Reference Synthesis Example 132, lithium hydroxide monohydrate (229 mg, 5.45 mmol) was added

and the resultant mixture was stirred at 115°C for 3 hours. After completion of the reaction, water was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was concentrated under reduced pressure and the obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 49/1 → 7/3) to obtain the title compound (640 mg, two step yield 74%).

[0357] Reference Synthesis Example 134

tert-Butyl (3R,4R)-4-(4-fluorophenyl)-3,4-dihydroxypiperidine-1-carboxylate

10 To a tert-butanol (29 mL) - water (29 mL) solution of AD-mix-β (8.10 g, manufactured by Aldrich Chemical Company, Inc.) and methanesulfonamide (515 mg, 5.41 mmol), tert-butyl 4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine-1(2H)-carboxylate (1.50 g, 5.49 mmol) was added at 0°C and the resultant mixture was stirred at room temperature for 1 hour and 30 minutes. After completion of the reaction, water was
15 added to the reaction solution and extraction from the resultant mixture with dichloromethane was performed. The organic layer was washed with 1 M sodium hydroxide aqueous solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1 → 1/1) to obtain the title compound (1.69 g,
20 yield 100%).

[0358] Reference Synthesis Example 135

tert-Butyl

(3R,4R)-4-(4-fluorophenyl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate

25 To a dichloromethane solution (17 mL) of tert-butyl (3R,4R)-4-(4-fluorophenyl)-3,4-dihydroxypiperidine-1-carboxylate (1.69 g, 5.43 mmol) synthesized in Reference Synthesis Example 134 and 4-dimethylaminopyridine (33.0 mg, 0.270 mmol), pivalic acid anhydride (1.31 g, 7.06 mmol) and triethylamine (1.52 mL, 10.9 mmol) were added and the resultant mixture was stirred at room temperature for 22

hours and 30 minutes. After completion of the reaction, ethyl acetate was added to the reaction solution and the resultant mixture was washed with each of saturated ammonium chloride aqueous solution and saturated sodium chloride aqueous solution. The obtained organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 9/1 → 3/1) to obtain the title compound (2.04 g, yield 95%).

[0359] Reference Synthesis Example 136

tert-Butyl (S)-4-(4-fluorophenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate

To a toluene solution (40 mL) of tert-butyl (3R,4R)-4-(4-fluorophenyl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate (2.04 g, 5.16 mmol) synthesized in Reference Synthesis Example 135, methyl N-(triethylammoniosulfonyl)carbamate (1.72 g, 7.22 mmol) was added and the resultant mixture was stirred at 60°C for 3 hours. After completion of the reaction, ethyl acetate was added to the reaction solution and the resultant mixture was washed with each of saturated sodium bicarbonate aqueous solution and saturated sodium chloride aqueous solution. The obtained organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 9/1 → 3/1) to obtain the title compound (1.82 g, yield 93%).

[0360] Reference Synthesis Example 137

(S)-4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl pivalate

To a dichloromethane solution (32 mL) of tert-butyl (S)-4-(4-fluorophenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate (1.62 g, 4.29 mmol) synthesized in Reference Synthesis Example 136, trifluoroacetic acid (6.5 mL) was added and the resultant mixture was stirred at room temperature for 20 hours. After completion of the reaction, saturated sodium bicarbonate aqueous solution was

added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the title compound as a crude product.

[0361] Reference Synthesis Example 138

5 (S)-4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol

To a 1,4-dioxane (11 mL) - water (6.5 mL) solution of the crude product of (S)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl pivalate synthesized in Reference Synthesis Example 137, lithium hydroxide monohydrate (1.54 g, 36.7 mmol) was added
10 and the resultant mixture was stirred at 115°C for 3 hours. After completion of the reaction, water was added to the reaction solution and extraction from the resultant mixture with chloroform was performed. The organic layer was concentrated under reduced pressure and the obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 32/1 → 13/7) to obtain the title compound (383
15 mg, two step yield 46%).

[0362] Reference Synthesis Example 139

tert-Butyl 5-chloro-6-methoxy-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate

To a 1,4-dioxane (50 mL) - water (10 mL) solution of
20 6-bromo-3-chloro-2-methoxypyridine (1.98 g, 8.90 mmol), tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (2.50 g, 8.09 mmol), and sodium carbonate (1.71 g, 16.1 mmol), tetrakis(triphenylphosphine) palladium (0) (467 mg, 0.400 mmol) was added under argon atmosphere and the resultant mixture was stirred at 100°C for 3 hours. After completion
25 of the reaction, saturated ammonium chloride aqueous solution was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1) to obtain the title compound (2.55 g,

yield 97%).

[0363] Reference Synthesis Example 140

tert-Butyl

(3S,4S)-4-(5-chloro-6-methoxypyridin-2-yl)-3,4-dihydroxypiperidine-1-carboxylate

5

tert-Butyl 5-chloro-6-methoxy-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate (2.10 g, 6.47 mmol) synthesized in Reference Synthesis Example 139 was used to obtain the title compound (2.19 g, yield 94%) by synthesis in a similar manner to Reference Synthesis Example 129.

10 [0364] Reference Synthesis Example 141

tert-Butyl

(3S,4S)-4-(5-chloro-6-methoxypyridin-2-yl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate

15 tert-Butyl

(3S,4S)-4-(5-chloro-6-methoxypyridin-2-yl)-3,4-dihydroxypiperidine-1-carboxylate (2.19 g, 6.10 mmol) synthesized in Reference Synthesis Example 140 was used to obtain the title compound (2.98 g, quantitative) by synthesis in a similar manner to Reference Synthesis Example 130.

20 [0365] Reference Synthesis Example 142

tert-Butyl

(R)-5-chloro-6-methoxy-5'-(pivaloyloxy)-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate

25 tert-Butyl

(3S,4S)-4-(5-chloro-6-methoxypyridin-2-yl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate (2.80 g, 6.10 mmol) synthesized in Reference Synthesis Example 141 was used to obtain the title compound (1.35 g, yield 52%) by synthesis in a similar manner to Reference Synthesis Example 131.

[0366] Reference Synthesis Example 143

(R)-5-Chloro-6-methoxy-1',2',3',6'-tetrahydro-[2,4'-bipyridine]-3'-yl pivalate

To a methanol solution (1.0 mL) of tert-butyl

- 5 (R)-5-chloro-6-methoxy-5'-(pivaloyloxy)-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate (1.35 g, 3.18 mmol) synthesized in Reference Synthesis Example 142, 4 M hydrogen chloride/1,4-dioxane (15 mL) was added and the resultant mixture was stirred at room temperature for 5 minutes. After completion of the reaction, the reaction solution was concentrated under reduced pressure and saturated sodium bicarbonate aqueous solution
10 and water were added to the concentrated solution, followed by extraction from the resultant mixture with ethyl acetate. The organic layer was concentrated under reduced pressure to obtain a crude product of the title compound (1.32 g).

[0367] Reference Synthesis Example 144

(R)-5-Chloro-6-methoxy-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-ol

15

The crude product of

- (R)-5-chloro-6-methoxy-1',2',3',6'-tetrahydro-[2,4'-bipyridine]-3'-yl pivalate (1.32 g) synthesized in Reference Synthesis Example 143 was used to obtain the title compound (160 mg, two step yield 21%) by synthesis in a similar manner to Reference Synthesis
20 Example 133.

[0368] Reference Synthesis Example 145

tert-Butyl 5-fluoro-6-methyl-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate

- 6-Bromo-3-fluoro-2-methylpyridine (1.40 g, 7.37 mmol) was used to obtain the title
25 compound (1.89 g, yield 97%) by synthesis in a similar manner to Reference Synthesis Example 139.

[0369] Reference Synthesis Example 146

tert-Butyl

(3S,4S)-4-(5-fluoro-6-methylpyridin-2-yl)-3,4-dihydropiperidine-1-carboxylate

tert-Butyl 5-fluoro-6-methyl-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate (1.38 g, 4.72 mmol) synthesized in Reference Synthesis Example 145 was used to obtain the title compound (1.08 g, yield 70%) by synthesis in a similar manner to Reference Synthesis Example 129.

[0370] Reference Synthesis Example 147

tert-Butyl

(3S,4S)-4-(5-fluoro-6-methylpyridin-2-yl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate

10

tert-Butyl

(3S,4S)-4-(5-fluoro-6-methylpyridin-2-yl)-3,4-dihydroxypiperidine-1-carboxylate (1.08 g, 3.31 mmol) synthesized in Reference Synthesis Example 146 was used to obtain the title compound (1.26 g, yield 93%) by synthesis in a similar manner to Reference Synthesis Example 130.

15

[0371] Reference Synthesis Example 148

tert-Butyl

(R)-5-fluoro-6-methyl-5'-(pivaloyloxy)-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate

20

tert-Butyl

(3S,4S)-4-(5-fluoro-6-methylpyridin-2-yl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate (1.26 g, 3.07 mmol) synthesized in Reference Synthesis Example 147 was used to obtain the title compound (488 mg, yield 40%) by synthesis in a similar manner to Reference Synthesis Example 131.

25

[0372] Reference Synthesis Example 149

(R)-5-Fluoro-6-methyl-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-yl pivalate

tert-Butyl

(R)-5-fluoro-6-methyl-5'-(pivaloyloxy)-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate

(488 mg, 1.24 mmol) synthesized in Reference Synthesis Example 148 was used to obtain a crude product of the title compound (390 mg) by synthesis in a similar manner to Reference Synthesis Example 143.

[0373] Reference Synthesis Example 150

5 (R)-5-Fluoro-6-methyl-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-ol

The crude product of

(R)-5-fluoro-6-methyl-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-yl pivalate (390 mg) synthesized in Reference Synthesis Example 149 was used to obtain the title compound
10 (208 mg, two step yield 80%) by synthesis in a similar manner to Reference Synthesis Example 133.

[0374] Reference Synthesis Example 151

tert-Butyl (3S,4S)-4-(6-chloropyridin-2-yl)-3,4-dihydropiperidine-1-carboxylate

15 tert-Butyl 6-chloro-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate (918 mg, 3.11 mmol) was used to obtain the title compound (933 mg, yield 93%) by synthesis in a similar manner to Reference Synthesis Example 129.

[0375] Reference Synthesis Example 152

tert-Butyl

20 (3S,4S)-4-(6-chloropyridin-2-yl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate

tert-Butyl (3S,4S)-4-(6-chloropyridin-2-yl)-3,4-dihydropiperidine-1-carboxylate (930 mg, 2.83 mmol) synthesized in Reference Synthesis Example 151 was used to obtain the title compound (1.14 g, yield 99%) by synthesis in a similar manner to
25 Reference Synthesis Example 130.

[0376] Reference Synthesis Example 153

tert-Butyl

(R)-6-chloro-5'-(pivaloyloxy)-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate

tert-Butyl

- (3S,4S)-4-(6-chloropyridin-2-yl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate (1.14 g, 2.76 mmol) synthesized in Reference Synthesis Example 152 was used to obtain the title compound (340 mg, yield 31%) by synthesis in a similar manner to Reference Synthesis Example 131.

[0377] Reference Synthesis Example 154

(R)-6-Chloro-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-yl pivalate

tert-Butyl

- 10 (R)-6-chloro-5'-(pivaloyloxy)-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate (340 mg, 0.861 mmol) synthesized in Reference Synthesis Example 153 was used to obtain a crude product of the title compound (248 mg) by synthesis in a similar manner to Reference Synthesis Example 132.

[0378] Reference Synthesis Example 155

- 15 (R)-6-Chloro-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-ol

- The crude product of (R)-6-chloro-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-yl pivalate (248 mg) synthesized in Reference Synthesis Example 154 was used to obtain the title compound (68.0 mg, two step yield 38%) by synthesis in a similar manner to Reference Synthesis Example 133.

[0379] Reference Synthesis Example 156

tert-Butyl

(3S,4S)-4-hydroxy-4-(6-methylpyridin-2-yl)-3-(pivaloyloxy)piperidine-1-carboxylate

- 25 A 1,4-dioxane (46 mL) - water (12 mL) solution of tert-butyl (3S,4S)-4-(6-chloropyridin-2-yl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate (2.30 g, 5.57 mmol) synthesized in Reference Synthesis Example 152, 2,4,6-trimethyl-1,3,5,2,4,6-trioxatriborinane (1.2 mL, 13.4 mmol), 1,1'-bis(diphenylphosphino)ferrocene-palladium (II) dichloride - dichloromethane

complex (915 mg, 1.12 mmol), and sodium carbonate (1.20 g, 11.3 mmol) was stirred under argon atmosphere at 88°C for 3 hours. After completion of the reaction, saturated ammonium chloride aqueous solution was added to the reaction solution and extraction from the resultant mixture with chloroform was performed. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1) to obtain the title compound (1.50 g, yield 69%).

[0380] Reference Synthesis Example 157

10 tert-Butyl

(R)-6-methyl-5'-(pivaloyloxy)-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate

tert-Butyl

(3S,4S)-4-hydroxy-4-(6-methylpyridin-2-yl)-3-(pivaloyloxy)piperidine-1-carboxylate (1.50 g, 3.82 mmol) synthesized in Reference Synthesis Example 156 was used to obtain the title compound (790 mg, yield 56%) by synthesis in a similar manner to Reference Synthesis Example 131.

[0381] Reference Synthesis Example 158

(R)-6-Methyl-1',2',3',6'-tetrahydro-[2,4'-bipyridine]-3'-yl pivalate

20

tert-Butyl

(R)-6-methyl-5'-(pivaloyloxy)-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate (790 mg, 2.11 mmol) synthesized in Reference Synthesis Example 157 was used to obtain a crude product of the title compound (576 mg) by synthesis in a similar manner to Reference Synthesis Example 132.

[0382] Reference Synthesis Example 159

(R)-6-Methyl-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-ol

The crude product of (R)-6-methyl-1',2',3',6'-tetrahydro-[2,4'-bipyridine]-3'-yl

pivalate (576 mg) synthesized in Reference Synthesis Example 158 was used to obtain the title compound (226 mg, two step yield 56%) by synthesis in a similar manner to Reference Synthesis Example 133.

[0383] Reference Synthesis Example 160

5 tert-Butyl 4-(4-fluoro-2-methylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate

1-Bromo-4-fluoro-2-methylbenzene (1.20 mL, 9.70 mmol) was used to obtain the title compound (1.95 g, yield 83%) by synthesis in a similar manner to Reference Synthesis Example 139.

10 [0384] Reference Synthesis Example 161

tert-Butyl (3S,4S)-4-(4-fluoro-2-methylphenyl)-3,4-dihydroxypiperidine-1-carboxylate

tert-Butyl 4-(4-fluoro-2-methylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (1.48 g, 5.08 mmol) synthesized in Reference Synthesis Example 160 was used to obtain the title compound (695 mg, yield 42%) by synthesis in a similar manner to Reference Synthesis Example 129.

[0385] Reference Synthesis Example 162

tert-Butyl

(3S,4S)-4-(4-fluoro-2-methylphenyl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate

20

tert-Butyl

(3S,4S)-4-(4-fluoro-2-methylphenyl)-3,4-dihydroxypiperidine-1-carboxylate (695 mg, 2.14 mmol) synthesized in Reference Synthesis Example 161 was used to obtain the title compound (856 mg, yield 98%) by synthesis in a similar manner to Reference Synthesis Example 130.

25

[0386] Reference Synthesis Example 163

tert-Butyl

(R)-4-(4-fluoro-2-methylphenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate

tert-Butyl

(3S,4S)-4-(4-fluoro-2-methylphenyl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate (856 mg, 2.09 mmol) synthesized in Reference Synthesis Example 162 was used to obtain the title compound (649 mg, yield 79%) by synthesis in a similar manner to Reference Synthesis Example 131.

[0387] Reference Synthesis Example 164

(R)-4-(4-Fluoro-2-methylphenyl)-1,2,3,6-tetrahydropyridin-3-yl pivalate

tert-Butyl

10 (R)-4-(4-fluoro-2-methylphenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate (649 mg, 1.66 mmol) synthesized in Reference Synthesis Example 163 was used to obtain the title compound (455 mg, yield 94%) by synthesis in a similar manner to Reference Synthesis Example 132.

[0388] Reference Synthesis Example 165

15 (R)-4-(4-Fluoro-2-methylphenyl)-1,2,3,6-tetrahydropyridin-3-ol

(R)-4-(4-Fluoro-2-methylphenyl)-1,2,3,6-tetrahydropyridin-3-yl pivalate (455 mg, 1.56 mmol) synthesized in Reference Synthesis Example 164 was used to obtain the title compound (295 mg, yield 91%) by synthesis in a similar manner to Reference Synthesis Example 133.

[0389] Reference Synthesis Example 166

tert-Butyl 6-methoxy-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate

25 2-Bromo-6-methoxypyridine (1.34 g, 7.12 mmol) was used to obtain the title compound (1.86 g, yield 99%) by synthesis in a similar manner to Reference Synthesis Example 139.

[0390] Reference Synthesis Example 167

tert-Butyl (3S,4S)-3,4-dihydroxy-4-(6-methoxypyridin-2-yl)piperidine-1-carboxylate

tert-Butyl 6-methoxy-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate (1.20 g, 4.13 mmol) synthesized in Reference Synthesis Example 166 was used to obtain the title compound (1.13 g, yield 84%) by synthesis in a similar manner to Reference Synthesis Example 129.

5 [0391] Reference Synthesis Example 168

tert-Butyl

(3S,4S)-4-hydroxy-4-(6-methoxypyridin-2-yl)-3-(pivaloyloxy)piperidine-1-carboxylate

tert-Butyl

10 (3S,4S)-3,4-dihydroxy-4-(6-methoxypyridin-2-yl)piperidine-1-carboxylate (1.12 g, 3.45 mmol) synthesized in Reference Synthesis Example 167 was used to obtain the title compound (1.12 g, yield 79%) by synthesis in a similar manner to Reference Synthesis Example 130.

[0392] Reference Synthesis Example 169

15 tert-Butyl

(R)-6-methoxy-5'-(pivaloyloxy)-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate

tert-Butyl

(3S,4S)-4-hydroxy-4-(6-methoxypyridin-2-yl)-3-(pivaloyloxy)piperidine-1-carboxylate
20 (1.12 g, 2.74 mmol) synthesized in Reference Synthesis Example 168 was used to obtain the title compound (616 mg, yield 58%) by synthesis in a similar manner to Reference Synthesis Example 131.

[0393] Reference Synthesis Example 170

(R)-6-Methoxy-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-yl pivalate

25 tert-Butyl

(R)-6-methoxy-5'-(pivaloyloxy)-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate (615 mg, 1.57 mmol) synthesized in Reference Synthesis Example 169 was used to obtain a crude product of the title compound (530 mg) by synthesis in a similar manner to Reference Synthesis Example 132.

[0394] Reference Synthesis Example 171

(R)-6-Methoxy-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-ol

The crude product of (R)-6-methoxy-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-yl
 5 pivalate (530 mg) synthesized in Reference Synthesis Example 170 was used to obtain
 the title compound (217 mg, two step yield 66%) by synthesis in a similar manner to
 Reference Synthesis Example 133.

[0395] Reference Synthesis Example 172

tert-Butyl 4-(3-fluoro-4-methylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate

10

4-Bromo-2-fluoro-1-methylbenzene (1.35 g, 7.12 mmol) was used to obtain the title
 compound (1.91 g, quantitative) by synthesis in a similar manner to Reference Synthesis
 Example 139.

[0396] Reference Synthesis Example 173

15 tert-Butyl (3S,4S)-4-(3-fluoro-4-methylphenyl)-3,4-dihydroxypiperidine-1-carboxylate

tert-Butyl 4-(3-fluoro-4-methylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (1.20
 g, 4.12 mmol) synthesized in Reference Synthesis Example 172 was used to obtain the
 title compound (1.06 g, yield 79%) by synthesis in a similar manner to Reference
 20 Synthesis Example 129.

[0397] Reference Synthesis Example 174

tert-Butyl

(3S,4S)-4-(3-fluoro-4-methylphenyl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate

25

tert-Butyl

(3S,4S)-4-(3-fluoro-4-methylphenyl)-3,4-dihydroxypiperidine-1-carboxylate (1.05 g,
 3.23 mmol) synthesized in Reference Synthesis Example 173 was used to obtain the title
 compound (1.27 g, yield 96%) by synthesis in a similar manner to Reference Synthesis
 Example 130.

[0398] Reference Synthesis Example 175

tert-Butyl

(R)-4-(3-fluoro-4-methylphenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate

5 tert-Butyl

(3S,4S)-4-(3-fluoro-4-methylphenyl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate (1.26 g, 3.08 mmol) synthesized in Reference Synthesis Example 174 was used to obtain the title compound (861 mg, yield 71%) by synthesis in a similar manner to Reference Synthesis Example 131.

10 [0399] Reference Synthesis Example 176

(R)-4-(3-Fluoro-4-methylphenyl)-1,2,3,6-tetrahydropyridin-3-yl pivalate

tert-Butyl

(R)-4-(3-fluoro-4-methylphenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate (850 mg, 2.17 mmol) synthesized in Reference Synthesis Example 175 was used to obtain a crude product of the title compound (643 mg) by synthesis in a similar manner to Reference Synthesis Example 132.

[0400] Reference Synthesis Example 177

(R)-4-(3-Fluoro-4-methylphenyl)-1,2,3,6-tetrahydropyridin-3-ol

20

The crude product of

(R)-4-(3-fluoro-4-methylphenyl)-1,2,3,6-tetrahydropyridin-3-yl pivalate (643 mg) synthesized in Reference Synthesis Example 176 was used to obtain the title compound (398 mg, two step yield 88%) by synthesis in a similar manner to Reference Synthesis Example 133.

25

[0401] Reference Synthesis Example 178

tert-Butyl 4-(3-chloro-4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate

4-Bromo-2-chloro-1-fluorobenzene (1.49 g, 7.12 mmol) was used to obtain the title

compound (2.14 g, quantitative) by synthesis in a similar manner to Reference Synthesis Example 139.

[0402] Reference Synthesis Example 179

tert-Butyl (3S,4S)-4-(3-chloro-4-fluorophenyl)-3,4-dihydroxypiperidine-1-carboxylate

5

tert-Butyl 4-(3-chloro-4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (1.20 g, 3.85 mmol) synthesized in Reference Synthesis Example 178 was used to obtain the title compound (1.25 g, yield 94%) by synthesis in a similar manner to Reference Synthesis Example 129.

10 [0403] Reference Synthesis Example 180

tert-Butyl

(3S,4S)-4-(3-chloro-4-fluorophenyl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate

tert-Butyl

15 (3S,4S)-4-(3-chloro-4-fluorophenyl)-3,4-dihydroxypiperidine-1-carboxylate (1.24 g, 3.59 mmol) synthesized in Reference Synthesis Example 179 was used to obtain the title compound (1.31 g, yield 85%) by synthesis in a similar manner to Reference Synthesis Example 130.

[0404] Reference Synthesis Example 181

20 tert-Butyl

(R)-1-(3-chloro-4-fluorophenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate

tert-Butyl

(3S,4S)-4-(3-chloro-4-fluorophenyl)-4-hydroxy-3-(pivaloyloxy)piperidine-1-carboxylate
25 (1.30 g, 3.02 mmol) synthesized in Reference Synthesis Example 180 was used to obtain the title compound (845 mg, yield 68%) by synthesis in a similar manner to Reference Synthesis Example 131.

[0405] Reference Synthesis Example 182

(R)-4-(3-Chloro-4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl pivalate

tert-Butyl

(R)-4-(3-chloro-4-fluorophenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate (830 mg, 2.02 mmol) synthesized in Reference Synthesis Example 181 was used to

5 obtain a crude product of the title compound (649 mg) by synthesis in a similar manner to Reference Synthesis Example 132.

[0406] Reference Synthesis Example 183

(R)-4-(3-chloro-4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol

10 The crude product of (R)-4-(3-chloro-4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl pivalate (649 mg) synthesized in Reference Synthesis Example 182 was used to obtain the title compound (370 mg, two step yield 78%) by synthesis in a similar manner to Reference Synthesis Example 133.

[0407] Reference Synthesis Example 184

15 tert-Butyl 6-methoxy-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate

2-Bromo-6-methoxypyridine (1.34 g, 7.12 mmol) was used to obtain the title compound (1.89 g, quantitative) by synthesis in a similar manner to Reference Synthesis Example 139.

20 [0408] Reference Synthesis Example 185

tert-Butyl 3,4-dihydroxy-4-(6-methoxypyridin-2-yl)piperidine-1-carboxylate

To an acetone (10 mL) - water (1.0 mL) solution of tert-butyl 6-methoxy-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate (1.00 g, 3.44 mmol) 25 synthesized in Reference Synthesis Example 184, supported osmium tetroxide (276 mg, 0.102 mmol, osmium content 9.4%, manufactured by Wako Pure Chemical Industries, Ltd.) and N-methylmorpholine-N-oxide (1.20 g, 10.3 mmol) were added and the resultant mixture was stirred at room temperature for 6 days. After completion of the reaction, the reaction solution was filtered and ethyl acetate was added to the filtrate, followed by

washing the resultant mixture with saturated ammonium chloride aqueous solution and brine. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the title compound (1.01 g, yield 91%).

[0409] Reference Synthesis Example 186

5 tert-Butyl 4-hydroxy-4-(6-methoxypyridin-2-yl)-3-(pivaloyloxy)piperidine-1-carboxylate

tert-Butyl 3,4-dihydroxy-4-(6-methoxypyridin-2-yl)piperidine-1-carboxylate (1.01 g, 3.11 mmol) synthesized in Reference Synthesis Example 185 was used to obtain the title compound (950 mg, yield 75%) by synthesis in a similar manner to Reference Synthesis
10 Example 130.

[0410] Reference Synthesis Example 187

tert-Butyl 6-methoxy-5'-(pivaloyloxy)-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate

tert-Butyl

15 4-hydroxy-4-(6-methoxypyridin-2-yl)-3-(pivaloyloxy)piperidine-1-carboxylate (844 mg, 2.06 mmol) synthesized in Reference Synthesis Example 186 was used to obtain the title compound (538 mg, yield 67%) by synthesis in a similar manner to Reference Synthesis Example 131.

[0411] Reference Synthesis Example 188

20 6-Methoxy-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-yl pivalate

tert-Butyl

6-methoxy-5'-(pivaloyloxy)-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate (50.0 mg, 0.128 mmol) synthesized in Reference Synthesis Example 187 was used to obtain a crude
25 product of the title compound by synthesis in a similar manner to Reference Synthesis Example 132. The crude product was used in the next reaction as it was.

[0412] Reference Synthesis Example 189

6-Methoxy-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-ol

The crude product of 6-methoxy-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-yl pivalate synthesized in Reference Synthesis Example 188 was used to obtain the title compound (20.0 mg, two step yield 76%) by synthesis in a similar manner to Reference Synthesis Example 133.

5 [0413] Reference Synthesis Example 190

tert-Butyl (3S,4R)-3,4-dihydroxy-4-(4-methylthiophen-2-yl)piperidine-1-carboxylate

tert-Butyl 4-(4-methylthiophen-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (700 mg, 2.51 mmol) was used to obtain the title compound (570 mg, yield 73%) by synthesis in a similar manner to Reference Synthesis Example 129.

[0414] Reference Synthesis Example 191

tert-Butyl

(3S,4R)-4-hydroxy-4-(4-methylthiophen-2-yl)-3-(pivaloyloxy)piperidine-1-carboxylate

15 tert-Butyl

(3S,4R)-3,4-dihydroxy-4-(4-methylthiophen-2-yl)piperidine-1-carboxylate (570 mg, 1.81 mmol) synthesized in Reference Synthesis Example 190 was used to obtain the title compound (615 mg, yield 85%) by synthesis in a similar manner to Reference Synthesis Example 130.

20 [0415] Reference Synthesis Example 192

tert-Butyl

(R)-4-(4-methylthiophen-2-yl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate

tert-Butyl

25 (3S,4R)-4-hydroxy-4-(4-methylthiophen-2-yl)-3-(pivaloyloxy)piperidine-1-carboxylate (615 mg, 1.54 mmol) synthesized in Reference Synthesis Example 191 was used to obtain the title compound (338 mg, yield 58%) by synthesis in a similar manner to Reference Synthesis Example 131.

[0416] Reference Synthesis Example 193

(R)-4-(4-Methylthiophen-2-yl)-1,2,3,6-tetrahydropyridin-3-yl pivalate

tert-Butyl

(R)-4-(4-methylthiophen-2-yl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate

5 (338 mg, 0.891 mmol) synthesized in Reference Synthesis Example 192 was used to obtain a crude product of the title compound (219 mg) by synthesis in a similar manner to Reference Synthesis Example 132.

[0417] Reference Synthesis Example 194

(R)-4-(4-Methylthiophen-2-yl)-1,2,3,6-tetrahydropyridin-3-ol

10

The crude product of (R)-4-(4-methylthiophen-2-yl)-1,2,3,6-tetrahydropyridin-3-yl pivalate (219 mg) synthesized in Reference Synthesis Example 193 was used to obtain the title compound (112 mg, two step yield 64%) by synthesis in a similar manner to Reference Synthesis Example 133.

15 [0418] Reference Synthesis Example 195

tert-Butyl 4-(5-methylthiophen-3-yl)-5,6-dihydropyridine-1(2H)-carboxylate

tert-Butyl

4-{[(trifluoromethyl)sulfonyl]oxy}-5,6-dihydropyridine-1(2H)-carboxylate (1.66 g, 5.00
20 mmol) and a crude product of 4,4,5,5-tetramethyl-2-(5-methylthiophen-3-yl)-1,3,2-dioxaborolane (3.40 g) were used to obtain a crude product of the title compound (970 mg) by synthesis in a similar manner to Reference Synthesis Example 139.

[0419] Reference Synthesis Example 196

25 tert-Butyl (3S,4S)-3,4-dihydroxy-4-(5-methylthiophen-3-yl)piperidine-1-carboxylate

The crude product of tert-butyl

4-(5-methylthiophen-3-yl)-5,6-dihydropyridine-1(2H)-carboxylate (970 mg) synthesized in Reference Synthesis Example 195 was used to obtain a crude product of the title

compound (415 mg) by synthesis in a similar manner to Reference Synthesis Example 129.

[0420] Reference Synthesis Example 197

tert-Butyl

5 (3S,4S)-4-hydroxy-4-(5-methylthiophen-3-yl)-3-(pivaloyloxy)piperidine-1-carboxylate

The crude product of tert-butyl

(3S,4S)-3,4-dihydroxy-4-(5-methylthiophen-3-yl)piperidine-1-carboxylate (415 mg)

synthesized in Reference Synthesis Example 196 was used to obtain a crude product of

10 the title compound (310 mg) by synthesis in a similar manner to Reference Synthesis Example 130.

[0421] Reference Synthesis Example 198

tert-Butyl

(R)-4-(5-methylthiophen-3-yl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate

15

The crude product of tert-butyl

(3S,4S)-4-hydroxy-4-(5-methylthiophen-3-yl)-3-(pivaloyloxy)piperidine-1-carboxylate

(310 mg) synthesized in Reference Synthesis Example 197 was used to obtain a crude

product of the title compound (233 mg) by synthesis in a similar manner to Reference

20 Synthesis Example 131.

[0422] Reference Synthesis Example 199

(R)-4-(5-Methylthiophen-3-yl)-1,2,3,6-tetrahydropyridin-3-yl pivalate

The crude product of tert-butyl

25 (R)-4-(5-methylthiophen-3-yl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate

(233 mg) synthesized in Reference Synthesis Example 198 was used to obtain a crude

product of the title compound (185 mg) by synthesis in a similar manner to Reference

Synthesis Example 132.

[0423] Reference Synthesis Example 200

(R)-4-(5-Methylthiophen-3-yl)-1,2,3,6-tetrahydropyridin-3-ol

The crude product of (R)-4-(5-methylthiophen-3-yl)-1,2,3,6-tetrahydropyridin-3-yl pivalate (185 mg) synthesized in Reference Synthesis Example 199 was used to obtain
 5 the title compound (101 mg, five step yield 10%) by synthesis in a similar manner to Reference Synthesis Example 133.

[0424] Reference Synthesis Example 201

tert-Butyl5-(1,3-dioxoisindolin-2-yl)-4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate

10

To a tetrahydrofuran solution (10 mL) of tert-butyl
 4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridine-1(2H)-carboxylate (1.00 g, 3.41
 mmol) synthesized in Reference Synthesis Example 354, triphenylphosphine (0.894 g,
 3.41 mmol), and phthalimide (0.501 g, 3.41 mmol), diisopropyl azodicarboxylate (1.9 M,
 15 toluene solution) (1.97 mL, 3.75 mmol) was added dropwise and the resultant mixture
 was stirred at room temperature for 1 day. After completion of the reaction, the reaction
 solution was concentrated and the resultant residue was purified by silica gel column
 chromatography to obtain the title compound (1.21 g, yield 84%).

[0425] Reference Synthesis Example 202

20 tert-Butyl 5-amino-4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate

To an ethanol solution (5.0 mL) of tert-butyl
 5-(1,3-dioxoisindolin-2-yl)-4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate
 (500 mg, 2.36 mmol) synthesized in Reference Synthesis Example 201, hydrazine
 25 monohydrate (177 mg, 7.08 mmol) was added and the resultant mixture was stirred under
 reflux by heating for 5 hours. After completion of the reaction, the reaction solution
 was filtered and the filtrate was concentrated under reduced pressure. Chloroform was
 added to the obtained residue and the resultant mixture was filtered again, followed by
 concentrating the resultant filtrate under reduced pressure. The obtained residue was

purified by silica gel column chromatography to obtain the title compound (150 mg, yield 22%).

[0426] Reference Synthesis Example 203

tert-Butyl

5 4-(4-fluorophenyl)-5-(methylsulfonamido)-5,6-dihydropyridine-1(2H)-carboxylate

To a tetrahydrofuran solution (2.0 mL) of tert-butyl
5-amino-4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (100 mg, 0.342
mmol) synthesized in Reference Synthesis Example 202, triethylamine (52.0 μ L, 0.374
10 mmol), and N,N-dimethyl-4-aminopyridine (2.00 mg, 0.0170 mmol) were added, and
then methanesulfonyl chloride (29.0 μ L, 0.374 mmol) was added at 0°C with stirring,
followed by stirring the resultant reaction solution for 2 hours. After completion of the
reaction, saturated ammonium chloride aqueous solution was added to the reaction
solution and extraction from the resultant mixture with ethyl acetate was performed.
15 The organic layer was dried over anhydrous magnesium sulfate and concentrated under
reduced pressure to obtain a crude product of the title compound.

[0427] Reference Synthesis Example 204

N-[4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl]methanesulfonamide

20 tert-Butyl

4-(4-fluorophenyl)-5-(methylsulfonamido)-5,6-dihydropyridine-1(2H)-carboxylate (140
mg, 0.378 mmol) synthesized in Reference Synthesis Example 203 was used to obtain a
crude product of the title compound by synthesis in a similar manner to Reference
Synthesis Example 132.

25 [0428] Reference Synthesis Example 205

2-[4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl]isoindoline-1,3-dione

tert-Butyl

5-(1,3-dioxoisindolin-2-yl)-4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate

(350 mg, 0.829 mmol) synthesized in Reference Synthesis Example 201 was used to obtain the title compound (90.0 mg, yield 34%) by synthesis in a similar manner to Reference Synthesis Example 132.

[0429] Reference Synthesis Example 206

5 4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-amine

2-[4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl]isoindolin-1,3-dione (90.0 mg., 0.278 mmol) synthesized in Reference Synthesis Example 205 was used to obtain the title compound (60.0 mg, quantitative) by synthesis in a similar manner to Reference

10 Synthesis Example 202.

[0430] Reference Synthesis Example 207

tert-Butyl 5-acetamido-4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate

To a tetrahydrofuran solution (2.0 ml) of tert-butyl

15 5-amino-4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (100 mg, 0.342 mmol) synthesized in Reference Synthesis Example 202, triethylamine (52.0 μ L, 0.374 mmol) and 4-dimethylaminopyridine (2.08 mg, 0.0170 mmol) were added and then acetic anhydride (26.0 μ L, 0.374 mmol) was added with stirring at 0°C, followed by stirring the resultant reaction solution at 0°C for 2 hours. After completion of the reaction, the
20 reaction solution was purified by silica gel column chromatography to obtain the title compound (119 mg, yield 87%).

[0431] Reference Synthesis Example 208

N-[4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl]acetamide

25 tert-Butyl 5-acetamido-4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (119 mg, 0.356 mmol) synthesized in Reference Synthesis Example 207 was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 132.

[0432] Reference Synthesis Example 209

tert-Butyl

(S)-5-(1,3-dioxoisindolin-2-yl)-4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate

5 tert-Butyl (R)-4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridine-1(2H)-carboxylate (387 mg, 1.32 mmol) synthesized in Reference Synthesis Example 364 was used to obtain the title compound (250 mg, yield 45%) by synthesis in a similar manner to Reference Synthesis Example 201.

[0433] Reference Synthesis Example 210

10 (S)-2-[4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl]isoindoline-1,3-dione

tert-Butyl

(S)-5-(1,3-dioxoisindolin-2-yl)-4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (0.250 g, 0.592 mmol) synthesized in Reference Synthesis Example 209 was used to
15 obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 132.

[0434] Reference Synthesis Example 211

(S)-4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-amine

20 To an ethanol solution (1.0 mL) of (S)-2-[4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl]isoindoline-1,3-dione (170 mg, 0.402 mmol) synthesized in Reference Synthesis Example 210, hydrazine monohydrate (150 mg, 3.00 mmol) was added and the resultant mixture was stirred under reflux by heating for 2 hours. After completion of the reaction, the reaction solution was filtered
25 and the filtrate was concentrated under reduced pressure. Ethyl acetate was added to the obtained residue and the resultant mixture was filtered again, followed by concentrating the resultant filtrate under reduced pressure to obtain the title compound (46.0 mg, yield 59%).

[0435] Reference Synthesis Example 212

tert-Butyl

(R)-5-(1,3-dioxoisindolin-2-yl)-4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate

5 tert-Butyl (S)-4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridine-1(2H)-carboxylate (387 mg, 1.32 mmol) synthesized in Reference Synthesis Example 365 was used to obtain the title compound (100 mg, yield 20%) by synthesis in a similar manner to Reference Synthesis Example 201.

[0436] Reference Synthesis Example 213

10 (R)-2-[4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl]isoindoline-1,3-dione

Tert-butyl

(R)-5-(1,3-dioxoisindolin-2-yl)-4-(4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (100 mg, 0.237 mmol) synthesized in Reference Synthesis Example 212 was used to
15 obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 132.

[0437] Reference Synthesis Example 214

(R)-4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-amine

20 (R)-2-[4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-yl]isoindoline-1,3-dione (72.0 mg, 0.224 mmol) synthesized in Reference Synthesis Example 213 was used to obtain the title compound (23.0 mg, yield 53%) by synthesis in a similar manner to Reference Synthesis Example 211.

[0438] Reference Synthesis Example 215

25 (R)-1-(4-Fluorophenyl)-5-(hydroxymethyl)pyrrolidin-2-one

To (S)-5-(hydroxymethyl)dihydrofuran-2(3H)-one (500 mg, 4.31 mmol), 4-fluoroaniline (574 mg, 5.17 mmol) was added and the resultant mixture was stirred at 150°C for 3 days. After completion of the reaction, chloroform and hydrochloric acid

were added to the reaction solution and extraction from the resultant mixture with chloroform was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product of the title compound (990 mg).

5 [0439] Reference Synthesis Example 216

(R)-5-(Aminomethyl)-1-(4-fluorophenyl)pyrrolidin-2-one

To a dichloromethane (10 mL) - acetonitrile (2.0 mL) solution of (R)-1-(4-fluorophenyl)-5-(hydroxymethyl)pyrrolidin-2-one (500 mg, 2.39 mmol) synthesized in Reference Synthesis Example 215, triphenylphosphine (753 mg, 2.87 mmol), di-tert-butyl iminodicarboxylate (626 mg, 2.87 mmol), and diethyl azodicarboxylate (40% toluene solution) (1.25 g, 2.87 mmol) were added at room temperature and the resultant mixture was stirred for 1 hour and 30 minutes. After completion of the reaction, the reaction solution was concentrated under reduced pressure and 4 M hydrogen chloride/1,4-dioxane solution (7.5 mL) was added to the obtained residue, followed by stirring the resultant mixture for 1 day. After completion of the reaction, water and ethyl acetate were added to the reaction solution and the resultant mixture was washed with ethyl acetate. Thereafter, 1 M sodium hydroxide aqueous solution was added to the water phase and extraction from the resultant mixture with chloroform was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product of the title compound (120 mg).

[0440] Reference Synthesis Example 217

(R)-[1-(4-Fluorophenyl)pyrrolidin-2-yl]methanamine

25

To a tetrahydrofuran solution (1.0 mL) of (R)-5-(aminomethyl)-1-(4-fluorophenyl)pyrrolidin-2-one (50.0 mg, 0.240 mmol) synthesized in Reference Synthesis Example 216, borane-tetrahydrofuran complex (8.5% tetrahydrofuran solution) (0.760 mL, 0.720 mmol) was added and the resultant mixture

was stirred at room temperature for 1 hour 30 minutes. Thereafter, the borane-tetrahydrofuran complex (1.0 mL) was further added and the resultant mixture was stirred at 70°C for 1 day. After completion of the reaction, methanol was added to the reaction solution and the resultant mixture was stirred at 70°C for 3 hours, followed by concentrating the reaction solution under reduced pressure to obtain a crude product of the title compound.

[0441] Reference Synthesis Example 218

(S)-1-(4-Fluorophenyl)-5-(hydroxymethyl)pyrrolidin-2-one

(R)-5-(Hydroxymethyl)dihydrofuran-2(3H)-one (500 mg, 4.31 mmol) was used to obtain a crude product of the title compound (1.15 g) by synthesis in a similar manner to Reference Synthesis Example 215.

[0442] Reference Synthesis Example 219

(S)-5-(Aminomethyl)-1-(4-fluorophenyl)pyrrolidin-2-one

(S)-1-(4-Fluorophenyl)-5-(hydroxymethyl)pyrrolidin-2-one (500 mg, 2.39 mmol) synthesized in Reference Synthesis Example 218 was used to obtain a crude product of the title compound (114 mg) by synthesis in a similar manner to Reference Synthesis Example 216.

[0443] Reference Synthesis Example 220

(S)-[1-(4-Fluorophenyl)pyrrolidin-2-yl]methanamine

(S)-5-(Aminomethyl)-1-(4-fluorophenyl)pyrrolidin-2-one (50.0 mg, 0.240 mmol) synthesized in Reference Synthesis Example 219 was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 217.

[0444] Reference Synthesis Example 221

3-(1,3-Dioxoisindolin-2-yl)cyclopenta-1-en-1-yl trifluoromethanesulfonate

To a dichloromethane solution (30 mL) of

2-(3-oxocyclopentyl)isoindoline-1,3-dione (1.08 g, 4.71 mmol), trifluoromethanesulfonic acid anhydride (1.16 mL, 7.06 mmol) and N,N-diisopropylethylamine (1.16 mL, 9.42 mmol) were added at 0°C and the resultant mixture was stirred at 0°C for 1 day. After completion of the reaction, the reaction solution was concentrated under reduced pressure and the obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1) to obtain the title compound (780 mg, yield 46%).

[0445] Reference Synthesis Example 222

2-[3-(4-Fluorophenyl)cyclopent-2-en-1-yl]isoindoline-1,3-dione

To a dimethoxyethane (2.0 mL) - water (1.0 mL) solution of 3-(1,3-dioxoisindolin-2-yl)cyclopenta-1-en-1-yl trifluoromethanesulfonate (100 mg, 2.77 mmol) obtained in Reference Synthesis Example 221, 4-fluorophenyl boronic acid (46.5 mg, 3.32 mmol), tetrakis(triphenylphosphine) palladium (0) (16.0 mg, 1.38 mmol), and cesium fluoride (84.1 mg, 5.54 mmol) were added and the resultant mixture was stirred at 70°C for 1 day. After completion of the reaction, the reaction solution was concentrated under reduced pressure and the obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 3/1) to obtain the title compound (64.0 mg, yield 75%).

[0446] Reference Synthesis Example 223

3-(4-Fluorophenyl)cyclopent-2-enamine

To an ethanol solution (0.50 mL) of 2-[3-(4-fluorophenyl)cyclopent-2-en-1-yl]isoindoline-1,3-dione (10.0 mg, 0.0325 mmol) synthesized in Reference Synthesis Example 222, hydrazine monohydrate (3.16 µL, 0.0651 mmol) was added and the resultant mixture was stirred at room temperature for 1 day. Further, the reaction solution was stirred at 70°C for 6 hours and then stirred at room temperature for 1 day. After completion of the reaction, the reaction solution was concentrated under reduced pressure and was washed with chloroform to obtain a crude product of the title compound.

[0447] Reference Synthesis Example 224

5-(4-Fluorophenyl)bicyclo[3.1.0]hexan-2-amine

To a dichloromethane solution (0.28 mL) of 3-(4-fluorophenyl)cyclopent-2-enamine
5 (55.8 mg, 0.315 mmol) synthesized in Reference Synthesis Example 223, diethylzinc
(15% solution in toluene) (162 μ L, 1.78 mmol) was added and then diiodomethane (127
 μ L, 1.58 mmol) was added at 0°C with stirring and the resultant mixture was stirred at
room temperature for 1 day. After completion of the reaction, saturated ammonium
chloride aqueous solution was added to the reaction solution and extraction from the
10 resultant mixture with chloroform was performed. The organic layer was concentrated
under reduced pressure to obtain a crude product of the title compound.

[0448] Reference Synthesis Example 225

2-Benzyl-5-(4-fluorophenyl)octahydropyrrolo[3,4-c]pyrrole

15 2-Benzyl-5-(4-fluorophenyl)octahydropyrrolo[3,4-c]pyrrole (50.0 mg, 0.247 mmol),
1-bromo-4-fluorobenzene (33.0 μ L, 0.303 mmol), and sodium tert-butoxide (144 mg,
1.50 mmol) were used to obtain the title compound (52.0 mg, yield 70%) by synthesis in
a similar manner to Reference Synthesis Example 23.

[0449] Reference Synthesis Example 226

20 2-(4-Fluorophenyl)octahydropyrrolo[3,4-c]pyrrole

To an ethanol solution (3.0 mL) of
2-benzyl-5-(4-fluorophenyl)octahydropyrrolo[3,4-c]pyrrole (52.0 mg, 0.175 mmol)
synthesized in Reference Synthesis Example 225, palladium hydroxide-activated carbon
25 catalyst (catalytic amount) was added and the resultant mixture was stirred under
hydrogen atmosphere at room temperature for 2 days. After completion of the reaction,
the reaction solution was filtered with Celite and the filtrate was concentrated under
reduced pressure to obtain a crude product of the title compound (25.0 mg).

[0450] Reference Synthesis Example 227

rac-(3S,5S)-5-(4-Fluorophenyl)tetrahydrofuran-3-yl acetate

Reference Synthesis Example 228

rac-(3R,5S)-5-(4-Fluorophenyl)tetrahydrofuran-3-yl acetate

- 5 Iodobenzene diacetate (4.18 g, 13.0 mmol), acetic acid (2.0 mL), and trifluoromethanesulfonic acid (44.2 μ L, 0.500 mmol) were added to 1-(4-fluorophenyl)but-3-en-1-ol (1.66 g, 10.0 mmol) and the resultant mixture was stirred at 50°C for 4 hours. After completion of the reaction, the reaction solution was concentrated and saturated sodium bicarbonate aqueous solution was added to the residue,
- 10 followed by extraction with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 3/1) to obtain a compound of Reference Synthesis Example 227 as the title compound (645 mg, yield 29%) and a compound of Reference Synthesis Example 228 as the title compound (503
- 15 mg, yield 22%).

[0451] Reference Synthesis Example 229

2-[rac-(3R,5S)-5-(4-Fluorophenyl)tetrahydrofuran-3-yl]isoindoline-1,3-dione

- To a 1,4-dioxane solution (2.0 mL) of
- 20 rac-(3S,5S)-5-(4-fluorophenyl)tetrahydrofuran-3-yl acetate (645 mg, 2.88 mmol) synthesized in Reference Synthesis Example 227, 1 M sodium hydroxide aqueous solution (5.76 mL) was added and the resultant mixture was stirred under reflux by heating for 8 hours. After completion of the reaction, extraction from the reaction solution with chloroform was performed and the organic layer was dried over anhydrous
- 25 magnesium sulfate and concentrated under reduced pressure. The obtained residue was used to obtain the title compound (623 mg, yield 70%) by synthesis in a similar manner to Reference Synthesis Example 201.

[0452] Reference Synthesis Example 230

rac-(3R,5S)-5-(4-Fluorophenyl)tetrahydrofuran-3-amine

2-[rac-(3R,5S)-5-(4-Fluorophenyl)tetrahydrofuran-3-yl]isoindoline-1,3-dione (611 mg, 1.96 mmol) synthesized in Reference Synthesis Example 229 was used to obtain the title compound (344 mg, yield 97%) by synthesis in a similar manner to Reference

5 Synthesis Example 202.

[0453] Reference Synthesis Example 231

2-[rac-(3S, 5S)-5-(4-Fluorophenyl)tetrahydrofuran-3-yl]isoindoline-1,3-dione

10 rac-(3R,5S)-5-(4-Fluorophenyl)tetrahydrofuran-3-yl acetate (593 mg, 2.24 mmol) synthesized in Reference Synthesis Example 228 was used to obtain the title compound (404 mg, yield 58%) by synthesis in a similar manner to Reference Synthesis Example 229.

[0454] Reference Synthesis Example 232

rac-(3S,5S)-5-(4-Fluorophenyl)tetrahydrofuran-3-amine

15

2-[rac-(3S, 5S)-5-(4-Fluorophenyl)tetrahydrofuran-3-yl]isoindoline-1,3-dione (393 mg, 1.26 mmol) synthesized in Reference Synthesis Example 231 was used to obtain the title compound (212 mg, yield 93%) by synthesis in a similar manner to Reference Synthesis Example 202.

20 [0455] Reference Synthesis Example 233

2-[rac-(2S,4S)-2-(4-Fluorophenyl)tetrahydro-2H-pyran-4-yl]isoindoline-1,3-dione

25 rac-(2S,4R)-2-(4-Fluorophenyl)tetrahydro-2H-pyran-4-ol (1.96 g, 10.0 mmol) was used to obtain the title compound (2.63 g, yield 81%) by synthesis in a similar manner to Reference Synthesis Example 201.

[0456] Reference Synthesis Example 234

rac-(2S,4S)-2-(4-Fluorophenyl)tetrahydro-2H-pyran-4-amine

2-[rac-(2S,4S)-2-(4-Fluorophenyl)tetrahydro-2H-pyran-4-yl]isoindoline-1,3-dione

(2.62 g, 8.08 mmol) synthesized in Reference Synthesis Example 233 was used to obtain the title compound (1.55 g, yield 99%) by synthesis in a similar manner to Reference Synthesis Example 202.

[0457] Reference Synthesis Example 235

5 N-[rac-(2S,4R)-2-(4-Fluorophenyl)tetrahydro-2H-pyran-4-yl]acetamide

To an acetonitrile solution (4.0 mL) of 4-fluorobenzaldehyde (421 μ L, 4.00 mmol) and 3-buten-1-ol (679 μ L, 8.00 mmol), sulfuric acid (427 μ L) was added and the resultant mixture was stirred at room temperature for 2 hours. After completion of the reaction, ethyl acetate and 1 M sodium hydroxide aqueous solution were added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (ethyl acetate) to obtain the title compound (382 mg, yield 40%).

15 [0458] Reference Synthesis Example 236

tert-Butyl [rac-(2S,4R)-2-(4-fluorophenyl)tetrahydro-2H-pyran-4-yl]carbamate

To a tetrahydrofuran solution (2.0 mL) of N-[rac-(2S,4R)-2-(4-fluorophenyl)tetrahydro-2H-pyran-4-yl]acetamide (382 mg, 1.61 mmol) synthesized in Reference Synthesis Example 235, di-tert-butyl dicarbonate (703 mg, 3.22 mmol) and 4-dimethylaminopyridine (7.80 mg, 80.0 μ mol) were added and the resultant mixture was stirred under reflux by heating for 2 hours. To the residue obtained by concentrating the reaction solution under reduced pressure, methanol (5.0 mL) and hydrazine monohydrate (403 mg, 8.05 mmol) were added and the resultant mixture was stirred at room temperature for 2 hours. After completion of the reaction, the reaction solution was concentrated under reduced pressure and saturated ammonium chloride aqueous solution was added to the resultant residue, followed by extraction from the resultant mixture with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was

purified by silica gel column chromatography (hexane/ethyl acetate = 6/1) to obtain the title compound (408 mg, yield 86%).

[0459] Reference Synthesis Example 237

rac-(2S,4R)-2-(4-Fluorophenyl)tetrahydro-2H-pyran-4-amine

5

tert-Butyl [rac-(2S,4R)-2-(4-fluorophenyl)tetrahydro-2H-pyran-4-yl]carbamate (408 mg, 1.38 mmol) synthesized in Reference Synthesis Example 236 was used to obtain the title compound (260 mg, yield 96%) by synthesis in a similar manner to Reference Synthesis Example 132.

10 [0460] Reference Synthesis Example 238

3-Benzyl-6-(4-fluorophenyl)-3-azabicyclo[3.1.0]hexan-2-one

To a dichloromethane solution (40 mL) of (E)-1-(3-chloroprop-1-en-1-yl)-4-fluorobenzene (2.40 g, 14.0 mmol), rhodium (II) acetate (57.7 mg, 0.140 mmol) was added and then a dichloromethane solution (20 mL) of ethyl diazoacetate (1.92 g, 16.8 mmol) was added dropwise at room temperature over 18 hours with stirring. After completion of the reaction, the reaction solution was concentrated and N, N-dimethylformamide (20 mL), benzylamine (1.50 g, 14.0 mmol), and sodium bicarbonate (1.18 g, 14.0 mmol) were added to the resultant residue and the resultant mixture was stirred at 150°C for 2 hours. After completion of the reaction, ethyl acetate was added to the reaction solution and the resultant mixture was washed with 1 M hydrochloric acid and a saturated sodium chloride aqueous solution. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. To the obtained residue, ethyl acetate was added and the resultant mixture was filtered. The filtrate was concentrated under reduced pressure and purified by silica gel column chromatography (hexane/ethyl acetate = 3/1) to obtain the title compound (956 mg, yield 24%).

[0461] Reference Synthesis Example 239

3-Benzyl-6-(4-fluorophenyl)-3-azabicyclo[3.1.0]hexane

To a tetrahydrofuran solution (40 mL) of
 3-benzyl-6-(4-fluorophenyl)-3-azabicyclo[3.1.0]hexan-2-one (956 mg, 3.40 mmol)
 synthesized in Reference Synthesis Example 238, lithium aluminum hydride (258 mg,
 5 6.80 mmol) was added at 0°C and the resultant mixture was stirred under reflux by
 heating for 30 minutes. After completion of the reaction, 1 M sodium hydroxide
 aqueous solution was added to the reaction solution at 0°C and extraction from the
 resultant mixture with dichloromethane was performed. The organic layer was purified
 by silica gel (amino-based) column chromatography (hexane/ethyl acetate = 15/1 →
 10 10/1) to obtain the title compound (593 mg, yield 66%).

[0462] Reference Synthesis Example 240

6-(4-Fluorophenyl)-3-azabicyclo[3.1.0]hexane

3-Benzyl-6-(4-fluorophenyl)-3-azabicyclo[3.1.0]hexane (593 mg, 2.26 mmol)
 15 synthesized in Reference Synthesis Example 239 was used to obtain the title compound
 (331 mg, yield 83%) by synthesis in a similar manner to Reference Synthesis Example
 226.

[0463] Reference Synthesis Example 241

2-{[trans-1-Benzyl-4-(4-fluorophenyl)pyrrolidin-3-yl]methyl}isoindoline-1,3-dione

20

[trans-1-Benzyl-4-(4-fluorophenyl)pyrrolidin-3-yl]methanol was used to obtain the
 title compound (4.83 g, quantitative) by synthesis in a similar manner to Reference
 Synthesis Example 201.

[0464] Reference Synthesis Example 242

25 tert-Butyl

trans-3-[(1,3-dioxoisoindolin-2-yl)methyl]-4-(4-fluorophenyl)pyrrolidine-1-carboxylate

To an ethanol solution (50 mL) of
 2-{[trans-1-benzyl-4-(4-fluorophenyl)pyrrolidin-3-yl]methyl}isoindoline-1,3-dione

synthesized in Reference Synthesis Example 241, di-tert-butyl dicarbonate (5.06 g, 23.2 mmol) and palladium-activated carbon (1.00 g) were added and the resultant mixture was stirred under hydrogen atmosphere at room temperature for 16 hours. After completion of the reaction, the reaction solution was filtered with Celite and the filtrate was concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1) and recrystallized using isopropyl ether to obtain the title compound (1.31 g, yield 43%).

[0465] Reference Synthesis Example 243

tert-Butyl trans-3-(aminomethyl)-4-(4-fluorophenyl)pyrrolidine-1-carboxylate

tert-Butyl

trans-3-[(1,3-dioxoisindolin-2-yl)methyl]-4-(4-fluorophenyl)pyrrolidine-1-carboxylate (849 mg, 2.00 mmol) synthesized in Reference Synthesis Example 242 was used to obtain the title compound (648 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 223.

[0466] Reference Synthesis Example 244

2-Bromo-4-nitro-1H-pyrrole

To a tetrahydrofuran solution (10 mL) of 3-nitro-1H-pyrrole (520 mg, 4.64 mmol), 1,3-dibromo-5,5-dimethylhydantoin (730 mg, 2.55 mmol) was added at -78°C and the resultant mixture was stirred at room temperature for 1 day. After completion of the reaction, the reaction solution was concentrated under reduced pressure and water was added to the resultant residue, followed by extraction from the resultant mixture with ethyl acetate. The organic layer was concentrated under reduced pressure to obtain the title compound as a crude product.

[0467] Reference Synthesis Example 245

2-(4-Fluorophenyl)-4-nitro-1H-pyrrole

2-Bromo-4-nitro-1H-pyrrole (886 mg, 4.64 mmol) synthesized in Reference

Synthesis Example 244 and 4-fluorophenylboronic acid (779 mg, 5.57 mmol) were used to obtain the title compound (383 mg, yield 40%) by synthesis in a similar manner to Reference Synthesis Example 139.

[0468] Reference Synthesis Example 246

5 2-(4-Fluorophenyl)-1-methyl-4-nitro-1H-pyrrole

To an N,N-dimethylformamide solution (0.50 mL) of 2-(4-fluorophenyl)-4-nitro-1H-pyrrole (10.0 mg, 0.0485 mmol) synthesized in Reference Synthesis Example 245, sodium hydride (2.50 mg, 0.0582 mmol) was added at 0°C and the resultant mixture was stirred for 1 hour. Thereafter, methyl iodide (18.8 µL, 0.0582 mmol) was added to the reaction solution and the resultant mixture was stirred at room temperature for 7 hours and 30 minutes. After completion of the reaction, water was added to the reaction solution and extraction from the resultant mixture with toluene was performed. The organic layer was concentrated under reduced pressure to obtain the title compound as a crude product.

[0469] Reference Synthesis Example 247

5-(4-Fluorophenyl)-1-methylpyrrolidin-3-amine

To 2-(4-fluorophenyl)-1-methyl-4-nitro-1H-pyrrole (200 mg, 1.03 mmol) synthesized in Reference Synthesis Example 246, acetic acid (2.0 mL) and platinum oxide (20.0 mg) were added and the resultant solution was stirred under hydrogen atmosphere at room temperature for 3 days. After completion of the reaction, the reaction solution was filtered with Celite and the filtrate was concentrated under reduced pressure. The obtained residue was purified by silica gel (amino-based) column chromatography (chloroform/methanol = 1/0 → 1/1) to obtain the title compound (15.0 mg, yield 7%).

[0470] Reference Synthesis Example 248

1-(4-Fluorophenyl)-3-nitro-1H-pyrrole

To a dichloromethane solution (4.0 mL) of 3-nitropyrrole (200 mg, 1.78 mmol), 4-fluorophenylboronic acid (499 mg, 3.57 mmol), and copper acetate (II) (486 mg, 2.68 mmol), N,N-diisopropylethylamine (606 μ L, 3.57 mmol) was added and the resultant mixture was stirred at room temperature for 1 day. After completion of the reaction, the reaction solution was filtered with Celite and the filtrate was concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 1/1) to obtain the title compound (131 mg, yield 36%).

[0471] Reference Synthesis Example 249

1-(4-Fluorophenyl)-1H-pyrrol-3-amine

To an ethanol solution (2.6 mL) of 1-(4-fluorophenyl)-3-nitro-1H-pyrrole (131 mg, 0.635 mmol) synthesized in Reference Synthesis Example 248, palladium-activated carbon (13.1 mg) was added and the resultant mixture was stirred under hydrogen atmosphere at room temperature for 1 day. After completion of the reaction, the reaction solution was filtered with Celite and the filtrate was concentrated under reduced pressure to obtain the title compound as a crude product.

[0472] Reference Synthesis Example 250

2-[cis-3-(4-Fluorophenyl)cyclobutyl]isoindoline-1,3-dione

trans-3-(4-Fluorophenyl)cyclobutanol (239 mg, 1.44 mmol) was used to obtain the title compound (266 mg, yield 62%) by synthesis in a similar manner to Reference Synthesis Example 201.

[0473] Reference Synthesis Example 251

cis-3-(4-Fluorophenyl)cyclobutanamine

2-[cis-3-(4-Fluorophenyl)cyclobutyl]isoindoline-1,3-dione (266 mg, 0.899 mmol) synthesized in Reference Synthesis Example 250 was used to obtain the title compound (51.3 mg, yield 35%) by synthesis in a similar manner to Reference Synthesis Example 223.

[0474] Reference Synthesis Example 252

2-[trans-3-(4-Fluorophenyl)cyclobutyl]isoindoline-1,3-dione

cis-3-(4-Fluorophenyl)cyclobutanol (166 mg, 1.00 mmol) was used to obtain the
 5 title compound (140 mg, yield 48%) by synthesis in a similar manner to Reference
 Synthesis Example 201.

[0475] Reference Synthesis Example 253

trans-3-(4-Fluorophenyl)cyclobutanamine

10 2-[trans-3-(4-Fluorophenyl)cyclobutyl]isoindoline-1,3-dione (140 mg, 0.475 mmol)
 synthesized in Reference Synthesis Example 252 was used to obtain the title compound
 (59.2 mg, yield 75%) by synthesis in a similar manner to Reference Synthesis Example
 223.

[0476] Reference Synthesis Example 254

15 Methyl cis-1-(4-fluorophenyl)-3-hydroxycyclobutane-carboxylate

To a tetrahydrofuran solution (5.5 mL) of 2-(4-fluorophenyl)acetic acid (1.90 g, 12.3
 mmol), 2 M isopropyl magnesium chloride (13.5 mL, 27.1 mmol) was added at room
 temperature and the resultant mixture was stirred at 40°C for 70 minutes. Thereafter,
 20 2-(chloromethyl)oxirane (1.74 mL, 22.1 mmol) was added to the reaction solution and
 the resultant mixture was stirred at room temperature for 3 hours and 30 minutes.
 Further, 2 M isopropyl magnesium chloride - tetrahydrofuran solution (12.3 mmol, 24.6
 mmol) was added and the resultant mixture was stirred for 1 day. After completion of
 the reaction, 1 M hydrochloric acid was added to the reaction solution to acidify the
 25 liquid and extraction from the resultant mixture with ethyl acetate was performed. The
 organic layer was concentrated under reduced pressure and methanol (30 mL) and
 concentrated sulfuric acid (2.0 mL) were added to the obtained residue, followed by
 stirring the resultant mixture under reflux by heating for 1 hour. After completion of the
 reaction, the reaction solution was concentrated under reduced pressure and ice water was

added to the residue, followed by extraction from the resultant mixture with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 3/2) to obtain the title compound (1.83 g, yield 66%).

[0477] Reference Synthesis Example 255

Methyl trans-3-(1,3-dioxoisindolin-2-yl)-1-(4-fluorophenyl)cyclobutane-carboxylate

Methyl cis-1-(4-fluorophenyl)-3-hydroxycyclobutane-carboxylate (1.83 g, 8.16 mmol) synthesized in Reference Synthesis Example 254 was used to obtain the title compound (2.21 g, yield 77%) by synthesis in a similar manner to Reference Synthesis Example 201.

[0478] Reference Synthesis Example 256

trans-3-(1,3-Dioxoisindolin-2-yl)-1-(4-fluorophenyl)cyclobutane carboxylic acid

To a tetrahydrofuran solution (5.0 mL) of methyl trans-3-(1,3-dioxoisindolin-2-yl)-1-(4-fluorophenyl)cyclobutane-carboxylate (353 mg, 1.00 mmol) synthesized in Reference Synthesis Example 255, potassium trimethylsilanolate (257 mg, 2.00 mmol) was added and the resultant mixture was stirred at room temperature for 2 hours. Thereafter, 4 M hydrogen chloride/1,4-dioxane (10 mL) was added and the resultant mixture was stirred at room temperature for 1 day, and then under reflux by heating for 2 hours. After completion of the reaction, saturated sodium chloride aqueous solution was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the title compound as a crude product.

[0479] Reference Synthesis Example 257

trans-3-(1,3-Dioxoisindolin-2-yl)-1-(4-fluorophenyl)cyclobutanecarboxamide

To a dichloromethane solution (5.0 mL) of the crude product of trans-3-(1,3-dioxoisindolin-2-yl)-1-(4-fluorophenyl)cyclobutane carboxylic acid synthesized in Reference Synthesis Example 256, oxalyl chloride (171 μ L, 2.00 mmol) and a catalytic amount of N,N-dimethylformamide were added and the resultant mixture was stirred at room temperature for 2 hours. Thereafter, saturated ammonium aqueous solution (3.0 mL) was added to the reaction solution and the resultant mixture was stirred for 30 minutes. After completion of the reaction, 1 M hydrochloric acid was added to the reaction solution and extraction from the resultant mixture with chloroform was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/3) to obtain the title compound (234 mg, yield 69%).

[0480] Reference Synthesis Example 258

trans-3-(1,3-Dioxoisindolin-2-yl)-1-(4-fluorophenyl)cyclobutanecarbonitrile

To trans-3-(1,3-dioxoisindolin-2-yl)-1-(4-fluorophenyl)cyclobutanecarboxamide (102 mg, 0.300 mmol) synthesized in Reference Synthesis Example 257, thionyl chloride (2.0 mL) was added and the resultant mixture was stirred under reflux by heating for 2 hours. After completion of the reaction, toluene was added to the reaction solution and the resultant mixture was concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1) to obtain the title compound (86.0 mg, yield 89%).

[0481] Reference Synthesis Example 259

trans-3-Amino-1-(4-fluorophenyl)cyclobutanecarbonitrile

trans-3-(1,3-Dioxoisindolin-2-yl)-1-(4-fluorophenyl)cyclobutanecarbonitrile (86.0 mg, 0.268 mol) synthesized in Reference Synthesis Example 258 was used to obtain the title compound (50.3 mg, yield 99%) by synthesis in a similar manner to Reference Synthesis Example 223.

[0482] Reference Synthesis Example 260

tert-Butyl 6,6-difluoro-4-(4-fluorophenyl)-1,4-diazepane-1-carboxylate

Reference Synthesis Example 261

tert-Butyl 6,6-difluoro-4-phenyl-1,4-diazepane-1-carboxylate

5

To a toluene solution (2.0 mL) of 1-fluoro-4-iodobenzene (62.6 mg, 0.282 mmol), tert-butyl 6,6-difluoro-1,4-diazepane-1-carboxylate (100 mg, 0.423 mmol), cesium carbonate (184 mg, 0.564 mmol), 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (23.3 mg, 0.0423 mmol), and palladium acetate (II) (6.30 mg, 0.0282 mmol) were added and the resultant mixture was stirred under nitrogen atmosphere at 110°C for 4 hours. After completion of the reaction, the reaction solution was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1) to obtain a compound (10.2 mg, yield 11%) in Reference Synthesis Example 260 and a compound (6.30 mg, yield 7%) in Reference Synthesis Example 261 as the title compounds.

15 [0483] Reference Synthesis Example 262

6,6-Difluoro-1-(4-fluorophenyl)-1,4-diazepane trifluoroacetate

tert-Butyl 6,6-difluoro-4-(4-fluorophenyl)-1,4-diazepane-1-carboxylate (10.2 mg, 0.0309 mmol) synthesized in Reference Synthesis Example 260 was used to obtain the title compound as a crude product (6.60 mg) by synthesis in a similar manner to Reference Synthesis Example 359.

[0484] Reference Synthesis Example 263

6,6-Difluoro-1-phenyl-1,4-diazepane trifluoroacetate

25 tert-Butyl 6,6-difluoro-4-phenyl-1,4-diazepane-1-carboxylate (6.30 mg, 0.0202 mmol) synthesized in Reference Synthesis Example 261 was used to obtain the title compound as a crude product (6.40 mg) by synthesis in a similar manner to Reference Synthesis Example 359.

[0485] Reference Synthesis Example 264

tert-Butyl 4-(4-fluorophenyl)-4,7-diazaspiro[2.5]octane-7-carboxylate

To a toluene solution (2.0 mL) of tert-butyl 4,7-diazaspiro[2.5]octane-7-carboxylate hydrochloride (100 mg, 0.402 mmol), 1-fluoro-4-iodobenzene (446 mg, 2.01 mmol),
 5 sodium tert-butoxide (77.3 mg, 0.804 mmol), and bis (tri-tert-butylphosphine) palladium (0) (41.1 mg, 0.0804 mmol) were added and the resultant mixture was stirred at 110°C for 1 hour and 30 minutes. After completion of the reaction, the reaction solution was purified by silica gel column chromatography (hexane/ethyl acetate = 1/0 → 4/1) to obtain the title compound (116 mg, yield 94%).

10 [0486] Reference Synthesis Example 265

4-(4-Fluorophenyl)-4,7-diazaspiro[2.5]octane

tert-Butyl 4-(4-fluorophenyl)-4,7-diazaspiro[2.5]octane-7-carboxylate (20.0 mg, 65.3 μmol) synthesized in Reference Synthesis Example 264 was used to obtain a crude
 15 product (13.3 mg) of the title compound by synthesis in a similar manner to Reference Synthesis Example 132.

[0487] Reference Synthesis Example 266

6-Methoxy-1',2',3',6'-tetrahydro-2,4'-bipyridine hydrochloride

20 tert-Butyl 6-methoxy-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate (100 mg, 0.344 mmol) synthesized in Reference Synthesis Example 184 was used to obtain a crude product (49.3 mg) of the title compound by synthesis in a similar manner to Reference Synthesis Example 26.

[0488] Reference Synthesis Example 267

25 5-Chloro-6-methoxy-1',2',3',6'-tetrahydro-2,4'-bipyridine hydrochloride

tert-Butyl 5-chloro-6-methoxy-5',6'-dihydro-[2,4'-bipyridine]-1'(2'H)-carboxylate (100 mg, 0.308 mmol) synthesized in Reference Synthesis Example 139 was used to obtain a crude product (70.4 mg) of the title compound by synthesis in a similar manner

to Reference Synthesis Example 26.

[0489] Reference Synthesis Example 268

4-(3-Fluoro-4-methylphenyl)-1,2,3,6-tetrahydropyridine hydrochloride

5 tert-Butyl 4-(3-fluoro-4-methylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (100 mg, 0.343 mmol) synthesized in Reference Synthesis Example 172 was used to obtain a crude product (57.9 mg) of the title compound by synthesis in a similar manner to Reference Synthesis Example 26.

[0490] Reference Synthesis Example 269

10 4-(3-Chloro-4-fluorophenyl)-1,2,3,6-tetrahydropyridine hydrochloride

tert-Butyl 4-(3-chloro-4-fluorophenyl)-5,6 -dihydropyridine-1(2H)-carboxylate (100 mg, 0.321 mmol) synthesized in Reference Synthesis Example 178 was used to obtain a crude product (77.5 mg) of the title compound by synthesis in a similar manner to

15 Reference Synthesis Example 26.

[0491] Reference Synthesis Example 270

4-(4-Methylthiophen-2-yl)-1,2,3,6-tetrahydropyridine

To a crude product synthesized by using tert-butyl

20 4-(4-methylthiophen-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (10.0 mg, 0.358 mmol) in a similar manner to Reference Synthesis Example 26, saturated sodium bicarbonate aqueous solution was added and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was washed with water, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to obtain a crude product of the title
25 compound.

[0492] Reference Synthesis Example 271

tert-Butyl

4-(2-{[(tert-butyldimethylsilyl)oxy]methyl}-4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate

[(2-Bromo-5-fluorobenzyl)oxy] (tert-butyl) dimethyl silane (115 mg, 0.360 mmol) synthesized in Reference Synthesis Example 128 was used to obtain the title compound (68.0 mg, yield 54%) by synthesis in a similar manner to Reference Synthesis Example 139.

[0493] Reference Synthesis Example 272

[5-Fluoro-2-(1,2,3,6-tetrahydropyridin-4-yl)phenyl]methanol

To a crude product obtained by synthesis in a similar manner to Reference Synthesis Example 132 using tert-butyl 4-(2-{[(tert-butyldimethylsilyl)oxy]methyl}-4-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (68.0 mg, 0.161 mmol) synthesized in Reference Synthesis Example 271, ethanol (2.0 mL) and 1 M sodium hydroxide aqueous solution (1.0 mL) were added and the resultant mixture was stirred at room temperature for 2 hours. After completion of the reaction, 1 M hydrochloric acid was added to the reaction solution and extraction from the resultant mixture with chloroform was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product of the title compound.

[0494] Reference Synthesis Example 273

20 3-Methyl 1-tert-butyl

3-methyl-4-{[(trifluoromethyl)sulfonyl]oxy}-2,3-dihydropyridine-1,3(6H)-dicarboxylate

To a tetrahydrofuran solution of lithium diisopropylamide (6.47 mL, 7.19 mmol), 3-methyl 1-tert-butyl 3-methyl- 4-oxopiperidine-1,3-dicarboxylate (1.63 g, 5.99 mmol) was added under nitrogen atmosphere at -78°C and the resultant mixture was stirred for 30 minutes. To the reaction solution, trifluoromethanesulfonic acid anhydride (1.28 mL, 7.78 mmol) was added and the resultant mixture was stirred for 30 minutes. Thereafter, the temperature of the mixture was raised to room temperature and the mixture was stirred for 15 hours. After completion of the reaction, saturated ammonium chloride

aqueous solution was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1) to obtain the title compound (714 mg, yield 30%).

[0495] Reference Synthesis Example 274

3-Methyl 1-tert-butyl

4-(4-fluorophenyl)-3-methyl-2,3-dihydropyridine-1,3(6H)-dicarboxylate

To a dimethoxyethane solution (6.0 mL) of 3-methyl 1-tert-butyl 3-methyl-4-{[(trifluoromethyl)sulfonyl]oxy}-2,3-dihydropyridine-1,3(6H)-dicarboxylate (300 mg, 0.743 mmol) synthesized in Reference Synthesis Example 273, 4-fluorophenylboronic acid (347 mg, 2.14 mmol), tetrakis(triphenylphosphine) palladium (0) (102 mg, 0.0880 mmol), lithium chloride (321 mg, 7.57 mmol), and 2 M sodium carbonate aqueous solution (2.5 mL) were added and the resultant mixture was stirred at 90°C for 3 hours. After completion of the reaction, water was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 5/1) to obtain the title compound (207 mg, yield 80%).

[0496] Reference Synthesis Example 275

1-(tert-Butoxycarbonyl)-4-(4-fluorophenyl)-3-methyl-1,2,3,6-tetrahydropyridine-3-carboxylic acid

25

To a 1,4-dioxane solution (2.0 mL) of 3-methyl 1-tert-butyl 4-(4-fluorophenyl)-3-methyl-2,3-dihydropyridine-1,3(6H)-dicarboxylate (207 mg, 0.590 mmol) obtained in Reference Synthesis Example 274, lithium hydroxide (75.0 mg, 1.77 mmol) and water (0.50 mL) were added and the resultant mixture was stirred overnight at

90°C. After completion of the reaction, saturated ammonium chloride aqueous solution was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed three times. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to obtain the title compound (130 mg, yield 66%).

[0497] Reference Synthesis Example 276

tert-Butyl

4-(4-fluorophenyl)-5-[(methoxycarbonyl)amino]-5-methyl-5,6-dihydropyridine-1(2H)-carboxylate

10

To a toluene solution (15 mL) of 1-(tert-butoxycarbonyl)-4-(4-fluorophenyl)-3-methyl-1,2,3,6-tetrahydropyridine-3-carboxylic acid (130 mg, 0.387 mmol) obtained in Reference Synthesis Example 275, diphenyl phosphorazide (500 μ L, 2.32 mmol) and triethylamine (809 μ L 5.80 mmol) were added and the resultant mixture was stirred at 100°C for 30 minutes. To the reaction solution, methanol (15 mL) was added and the resultant mixture was stirred for 2 hours. After completion of the reaction, the reaction solution was concentrated under reduced pressure and the obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1) to obtain the title compound (120 mg, yield 85%).

20

[0498] Reference Synthesis Example 277

Methyl [4-(4-fluorophenyl)-3-methyl-1,2,3,6-tetrahydropyridin-3-yl]carbamate

A residue obtained by synthesis in a similar manner to in Reference Synthesis Example 132 using tert-butyl 4-(4-fluorophenyl)-5-[(methoxycarbonyl)amino]-5-methyl-5,6-dihydropyridine-1(2H)-carboxylate (120 mg, 0.329 mmol) obtained in Reference Synthesis Example 276 was purified by silica gel column chromatography (chloroform/methanol = 4/1) to obtain a crude product (120 mg) of the title compound.

25

[0499] Reference Synthesis Example 278

4-(4-Fluorophenyl)-3-methyl-1,2,3,6-tetrahydropyridin-3-amine

To the crude product of methyl

- 5 [4-(4-fluorophenyl)-3-methyl-1,2,3,6-tetrahydropyridin-3-yl]carbamate (120 mg) obtained in Reference Synthesis Example 277, isopropyl alcohol (10 mL) and potassium hydroxide (0.38 g, 6.70 mmol) were added and the resultant mixture was stirred overnight at 90°C. After completion of the reaction, a sodium chloride aqueous solution was added to the reaction solution and extraction from the resultant mixture with
- 10 dichloromethane was performed. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to obtain the title compound (10.0 mg, yield 11%).

[0500] Reference Synthesis Example 279

tert-Butyl 4-(4-fluorophenyl)-4-hydroxy-2-methylpiperidine-1-carboxylate

15

- To a tetrahydrofuran solution (10 mL) of 1-bromo-4-fluorobenzene (492 mg, 2.81 mmol), n-butyl lithium (1.62 M, hexane solution) (1.73 mL, 2.81 mmol) was added at -78°C and the resultant mixture was stirred for 40 minutes. To the reaction solution, a tetrahydrofuran solution of tert-butyl 2-methyl-4-oxopiperidine-1-carboxylate (500 mg,
- 20 2.34 mmol) was added and the resultant mixture was stirred at room temperature for 1 day. After completion of the reaction, water was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product (729 mg) of the title compound.

- 25 [0501] Reference Synthesis Example 280

4-(4-Fluorophenyl)-2-methyl-1,2,3,6-tetrahydropyridine trifluoroacetate4-(4-Fluorophenyl)-6-methyl-1,2,3,6-tetrahydropyridine trifluoroacetate

tert-Butyl 4-(4-fluorophenyl)-4-hydroxy-2-methylpiperidine-1-carboxylate (50.0 mg,

0.162 mmol) synthesized in Reference Synthesis Example 279 was used to obtain a mixture of 4-(4-fluorophenyl)-2-methyl-1,2,3,6-tetrahydropyridine trifluoroacetate and 4-(4-fluorophenyl)-6-methyl-1,2,3,6-tetrahydropyridine trifluoroacetate as the title compounds by synthesis in a similar manner to Reference Synthesis Example 359.

5 [0502] Reference Synthesis Example 281

tert-Butyl 4-fluoro-4-(4-fluorophenyl)-2-methylpiperidine-1-carboxylate

To a dichloromethane solution (2.0 mL) of tert-butyl 4-(4-fluorophenyl)-4-hydroxy-2-methylpiperidine-1-carboxylate (60.0 mg, 0.194 mmol) synthesized in Reference Synthesis Example 279, bis(2-methoxyethyl)amino-sulfur trifluoride (46.9 mg, 0.291 mmol) was added at -78°C and the resultant mixture was stirred at room temperature for 1 day. After completion of the reaction, saturated sodium carbonate aqueous solution was added to the reaction solution and extraction from the resultant mixture with chloroform was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product (54.4 mg) of the title compound.

[0503] Reference Synthesis Example 282

4-Fluoro-4-(4-fluorophenyl)-2-methylpiperidine

20 tert-Butyl 4-fluoro-4-(4-fluorophenyl)-2-methylpiperidine-1-carboxylate (54.4 mg, 0.175 mmol) synthesized in Reference Synthesis Example 281 was used to obtain the title compound as a crude product by synthesis in a similar manner to Reference Synthesis Example 143.

[0504] Reference Synthesis Example 283

25 tert-Butyl 4-(4-fluorophenyl)-4-hydroxy-3-methylpiperidine-1-carboxylate

tert-Butyl 3-methyl-4-oxopiperidine-1-carboxylate (100 mg, 0.469 mmol) was used to obtain the title compound (148 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 279.

[0505] Reference Synthesis Example 284

4-(4-Fluorophenyl)-3-methyl-1,2,3,6-tetrahydropyridine hydrochloride

4-(4-Fluorophenyl)-5-methyl-1,2,3,6-tetrahydropyridine hydrochloride

5 A 4 M hydrogen chloride/1,4-dioxane (2.0 mL) solution of tert-butyl
4-(4-fluorophenyl)-4-hydroxy-3-methylpiperidine-1-carboxylate (37.6 mg, 0.122 mmol)
synthesized in Reference Synthesis Example 283 was stirred at room temperature for 1
hour and thereafter 12 M hydrochloric acid (0.60 mL) was added to the solution and the
resultant mixture was stirred at room temperature for 2 days. Further, 12 M
10 hydrochloric acid (1.0 mL) was added and the resultant mixture was stirred for 2 days.
After completion of the reaction, the reaction solution was concentrated under reduced
pressure to obtain a mixture of 4-(4-fluorophenyl)-3-methyl-1,2,3,6-tetrahydropyridine
hydrochloride and 4-(4-fluorophenyl)-5-methyl-1,2,3,6-tetrahydropyridine hydrochloride
as the title compounds.

15 [0506] Reference Synthesis Example 285

4-(4-Fluorophenyl)-3-methylpiperidin-4-ol

tert-Butyl 4-(4-fluorophenyl)-4-hydroxy-3-methylpiperidine-1-carboxylate (30.0 mg,
0.0970 mmol) synthesized in Reference Synthesis Example 283 was used to obtain the
20 title compound as a crude product by synthesis in a similar manner to Reference
Synthesis Example 132.

[0507] Reference Synthesis Example 286

1-(5-Methylthiophen-3-yl)piperazine

25 A toluene solution (3.0 mL) of tert-butyl piperazine-1-carboxylate (559 mg, 3.00
mmol), 4-bromo-2-methylthiophene (354 mg, 2.00 mmol), 2,2'-bis(diphenylphosphino)-
1,1'-binaphthyl (37.4 mg, 0.0600 mmol), tris(dibenzylideneacetone) dipalladium (0)
(36.6 mg, 0.0400 mmol), and sodium tert-butoxide (481 mg, 5.00 mmol) was stirred at
120°C with a microwave for 30 minutes. After completion of the reaction, the reaction

solution was concentrated and the residue was purified by silica gel column chromatography (hexane/ethyl acetate = 10/1). To the obtained crude product, trifluoroacetic acid (2.0 mL) was added and the resultant mixture was stirred at room temperature for 5 minutes. After completion of the reaction, the reaction solution was concentrated under reduced pressure and 1 M sodium hydroxide aqueous solution was added to the resultant residue, followed by extraction from the resultant mixture with chloroform. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to obtain the title compound (111 mg, yield 30%).

[0508] Reference Synthesis Example 287

10 1-(5-Chloro-6-methoxypyridin-2-yl)piperazine

6-Bromo-3-chloro-2-methoxypyridine (134 mg, 0.600 mmol) was used to obtain the title compound (113 mg, yield 83%) by synthesis in a similar manner to Reference Synthesis Example 286.

15 [0509] Reference Synthesis Example 288

1-(5-Fluoro-6-methylpyridin-2-yl)piperazine

6-Bromo-3-fluoro-2-methylpyridine (38.0 mg, 0.200 mmol) was used to obtain the title compound (41.5 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 286.

[0510] Reference Synthesis Example 289

Methyl 1H-imidazole-1-carbonylcarbamimidothioate

To S-methylisothiourrea sulfate (8.35 g, 30.0 mmol), 1 M sodium hydroxide aqueous solution (60 mL) was added and then 1,1'-carbonyldiimidazole (9.73 g, 60.0 mmol) was slowly added at 0°C with stirring, followed by stirring the resultant mixture at 0°C for 30 minutes. After completion of the reaction, the precipitated solid was collected by filtration and the obtained solid was washed with ice water and dried to obtain the title compound (7.89 g, yield 72%).

[0511] Reference Synthesis Example 290

Methyl

N-[(5-(trifluoromethyl)thiophen-2-yl)methyl]amino)carbonyl]-carbamimidothioate

- 5 A tetrahydrofuran solution (30 mL) of methyl
1H-imidazole-1-carboxylcarbamimidothioate (5.70 g, 31.0 mmol) synthesized in
Reference Synthesis Example 289 was refluxed. A tetrahydrofuran solution (26 mL) of
[5-(trifluoromethyl)-thiophen-2-yl]methanamine (2.81 g, 15.5 mmol) was added
dropwise to the solution and the resultant mixture was refluxed for 30 minutes. After
10 completion of the reaction, the reaction solution was concentrated under reduced pressure
and the obtained residue was purified by silica gel column chromatography (hexane/ethyl
acetate = 1/1) to obtain the title compound (3.28 g, yield 71%).

[0512] Reference Synthesis Example 291

4-(Methylthio)-1-[(5-(trifluoromethyl)thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

- 15 To methyl
N-[(5-(trifluoromethyl)thiophen-2-yl)methyl]amino)carbonyl]-carbamimidothioate
(2.68 g, 9.03 mmol) synthesized in Reference Synthesis Example 290, triethyl
orthoformate (10 mL) was added and the resultant mixture was stirred at 150°C for 1
20 hour. After completion of the reaction, the reaction solution was cooled to room
temperature and ethanol (20 mL) was added, followed by cooling the resultant mixture to
0°C. The precipitated solid was collected by filtration and the solid was washed with
ethanol to obtain the title compound (1.69 g, yield 61%).

[0513] Reference Synthesis Example 292

- 25 6-Methyl-4-(methylthio)-1-[(5-(trifluoromethyl)thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

To methyl

N-[(5-(trifluoromethyl)thiophen-2-yl)methyl]amino)carbonyl]-carbamimidothioate

(59.5 mg, 0.200 mmol) synthesized in Reference Synthesis Example 290, triethyl orthoacetate (0.60 mL) was added and the resultant mixture was stirred at 150°C for 1 hour. After completion of the reaction, the reaction solution was cooled to room temperature and purified by silica gel column chromatography (hexane/ethyl acetate = 3/2) to obtain the title compound (45.8 mg, yield 71%).

[0514] Reference Synthesis Example 293

6-Ethyl-4-(methylthio)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10 Triethyl orthopropionate was used to obtain the title compound (27.0 mg, yield 81%) by synthesis in a similar manner to Reference Synthesis Example 292.

[0515] Reference Synthesis Example 294

4-(Methylthio)-6-propyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

15

Triethyl orthobutyrate was used to obtain the title compound (24.8 mg, yield 71%) by synthesis in a similar manner to Reference Synthesis Example 292.

[0516] Reference Synthesis Example 295

Methyl N-({[(5-bromothiophen-2-yl)methyl]amino}carbonyl)-carbamimidothioate

20

(5-Bromothiophen-2-yl)methanamine (1.12 g, 5.64 mmol) was used to obtain the title compound (1.45 g, yield 83%) by synthesis in a similar manner to Reference Synthesis Example 290.

[0517] Reference Synthesis Example 296

25 1-[(5-Bromothiophen-2-yl)methyl]-4-(methylthio)-1,3,5-triazin-2(1H)-one

Methyl N-({[(5-bromothiophen-2-yl)methyl]amino}carbonyl)-carbamimidothioate (700 mg, 2.27 mmol) synthesized in Reference Synthesis Example 295 and triethyl orthoformate were used to obtain the title compound (547 mg, yield 76%) by synthesis in

a similar manner to Reference Synthesis Example 292.

[0518] Reference Synthesis Example 297

1-[(5-Bromothiophen-2-yl)methyl]-6-methyl-4-(methylthio)-1,3,5-triazin-2(1H)-one

5 Methyl N-({[(5-bromothiophen-2-yl)methyl]amino}carbonyl)-carbamimidothioate (749 mg, 2.43 mmol) synthesized in Reference Synthesis Example 295 was used to obtain the title compound (576 mg, yield 71%) by synthesis in a similar manner to Reference Synthesis Example 292.

[0519] Reference Synthesis Example 298

10 Methyl N-({[(5-chlorothiophen-2-yl)methyl]amino}carbonyl)-carbamimidothioate

(5-Chlorothiophen-2-yl)methanamine (1.37 g, 9.32 mmol) was used to obtain the title compound (2.34 g, yield 95%) by synthesis in a similar manner to Reference Synthesis Example 290.

15 [0520] Reference Synthesis Example 299

1-[(5-Chlorothiophen-2-yl)methyl]-4-(methylthio)-1,3,5-triazin-2(1H)-one

Methyl N-({[(5-chlorothiophen-2-yl)methyl]amino}carbonyl)-carbamimidothioate (1.00 g, 3.79 mmol) synthesized in Reference Synthesis Example 298 was used to obtain
20 the title compound (662 mg, yield 64%) by synthesis in a similar manner to Reference Synthesis Example 296.

[0521] Reference Synthesis Example 300

1-[(5-Chlorothiophen-2-yl)methyl]-6-methyl-4-(methylthio)-1,3,5-triazin-2(1H)-one

25 Methyl N-({[(5-chlorothiophen-2-yl)methyl]amino}carbonyl)-carbamimidothioate (1.34 g, 5.07 mmol) synthesized in Reference Synthesis Example 298 was used to obtain the title compound (1.20 g, yield 82%) by synthesis in a similar manner to Reference Synthesis Example 292.

[0522] Reference Synthesis Example 301

Methyl N-[(5-(trifluoromethyl)furan-2-yl)methyl]amino)carbonyl]-carbamimidothioate

[5-(trifluoromethyl)furan-2-yl]methanamine (628 mg, 3.80 mmol) was used to obtain the title compound (870 mg, yield 81%) by synthesis in a similar manner to

5 Reference Synthesis Example 290.

[0523] Reference Synthesis Example 302

6-Methyl-4-(methylthio)-1-{[5-(trifluoromethyl)furan-2-yl]methyl}-1,3,5-triazin-2(1H)-o
ne

10 Methyl

N-[(5-(trifluoromethyl)furan-2-yl)methyl]amino)carbonyl]-carbamimidothioate (300 mg, 1.07 mmol) synthesized in Reference Synthesis Example 301 was used to obtain the title compound (204 mg, yield 63%) by synthesis in a similar manner to Reference Synthesis Example 292.

15 [0524] Reference Synthesis Example 303

4-(Methylthio)-1-{[5-(trifluoromethyl)furan-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Methyl

20 N-[(5-(trifluoromethyl)furan-2-yl)methyl]amino)carbonyl]-carbamimidothioate (569 mg, 2.02 mmol) synthesized in Reference Synthesis Example 301 was used to obtain the title compound (268 mg, yield 42%) by synthesis in a similar manner to Reference Synthesis Example 296.

[0525] Reference Synthesis Example 304

Methyl N-[(5-(tert-butyl)thiophen-2-yl)methyl]amino)carbonyl]-carbamimidothioate

25

[5-(tert-Butyl)thiophen-2-yl]methanamine (327 mg, 1.94 mmol) was used to obtain the title compound (547 mg, yield 99%) by synthesis in a similar manner to Reference Synthesis Example 290.

[0526] Reference Synthesis Example 305

1-([5-(tert-Butyl)thiophen-2-yl]methyl)-6-methyl-4-(methylthio)-1,3,5-triazin-2(1H)-one

Methyl

N-([5-(tert-butyl)thiophen-2-yl]methyl)amino)carbonyl]-carbamimidothioate (191 mg, 0.67 mmol) synthesized in Reference Synthesis Example 304 was used to obtain the title compound (109 mg, yield 55%) by synthesis in a similar manner to Reference Synthesis Example 292.

[0527] Reference Synthesis Example 306

1-([5-(tert-Butyl)thiophen-2-yl]methyl)-4-(methylthio)-1,3,5-triazin-2(1H)-one

10

Methyl

N-([5-(tert-butyl)thiophen-2-yl]methyl)amino)carbonyl]-carbamimidothioate (348 mg, 1.22 mmol) synthesized in Reference Synthesis Example 304 was used to obtain the title compound (275 mg, yield 73%) by synthesis in a similar manner to Reference Synthesis Example 296.

[0528] Reference Synthesis Example 307

5-[(1,3-Dioxoisindolin-2-yl)methyl]pyridin-2-yl trifluoromethanesulfonate

5-(Hydroxymethyl)pyridin-2-yl trifluoromethanesulfonate (1.68 g, 6.55 mmol) was used to obtain a crude product (3.39 g) of the title compound by synthesis in a similar manner to Reference Synthesis Example 201.

[0529] Reference Synthesis Example 308

5-(Aminomethyl)pyridin-2-yl trifluoromethanesulfonate

5-[(1,3-Dioxoisindolin-2-yl)methyl]pyridin-2-yl trifluoromethanesulfonate (3.39 g, 6.55 mmol) synthesized in Reference Synthesis Example 307 was used to obtain a crude product (2.84 g) of the title compound by synthesis in a similar manner to Reference Synthesis Example 202.

[0530] Reference Synthesis Example 309

5-(-{3-[Imino(methylthio)methyl]ureido}methyl)pyridin-2-yl trifluoromethanesulfonate

5-(Aminomethyl)pyridin-2-yl trifluoromethanesulfonate (1.81 g, 9.83 mmol) synthesized in Reference Synthesis Example 308 was used to obtain the title compound (2.55 g, quantitative) by synthesis in a similar manner to Reference Synthesis Example 290.

[0531] Reference Synthesis Example 310

5-{[6-Methyl-4-(methylthio)-2-oxo-1,3,5-triazin-1(2H)-yl]methyl}pyridin-2-yl trifluoromethanesulfonate

10

5-({3-[imino(methylthio)methyl]ureido}methyl)pyridin-2-yl trifluoromethanesulfonate (1.00 g, 2.69 mmol) synthesized in Reference Synthesis Example 309 was used to obtain the title compound (568 mg, yield 53%) by synthesis in a similar manner to Reference Synthesis Example 292.

15 [0532] Reference Synthesis Example 311

5-{[4-(Methylthio)-2-oxo-1,3,5-triazin-1(2H)-yl]methyl}pyridin-2-yl trifluoromethanesulfonate

20 5-({3-[Imino(methylthio)methyl]ureido}methyl)pyridin-2-yl trifluoromethanesulfonate (1.00 g, 2.69 mmol) synthesized in Reference Synthesis Example 309 was used to obtain the title compound (431 mg, yield 42%) by synthesis in a similar manner to Reference Synthesis Example 296.

[0533] Reference Synthesis Example 312

Methyl

25 N-[(1-[5-(trifluoromethyl)thiophen-2-yl]ethyl)amino]carbonyl]-carbamimidothioate

1-[5-(Trifluoromethyl)thiophen-2-yl]ethanamine (50.0 mg, 0.256 mmol) was used to obtain the title compound (93.2 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 290.

[0534] Reference Synthesis Example 313

6-Methyl-4-(methylthio)-1-{1-[5-(trifluoromethyl)thiophen-2-yl]ethyl}-1,3,5-triazin-2(1H)-one

Methyl

- 5 N-[(1-[5-(trifluoromethyl)thiophen-2-yl]ethyl)amino]carbonyl-carbamimidothioate (93.2 mg, 0.299 mmol) synthesized in Reference Synthesis Example 312 was used to obtain the title compound (54.4 mg, yield 41%) by synthesis in a similar manner to Reference Synthesis Example 292.

[0535] Reference Synthesis Example 314

- 10 1-(4-Methoxybenzyl)-4-(methylthio)-1,3,5-triazin-2(1H)-one

- A tetrahydrofuran solution (5.0 mL) of methyl
1H-imidazole-1-carboxylcarbamidithioate (921 mg, 5.00 mmol) synthesized in
Reference Synthesis Example 289 was refluxed and a tetrahydrofuran solution (5.0 mL)
15 of 4-methoxybenzylamine (343 mg, 2.50 mmol) was added dropwise to the refluxed
solution, followed by refluxing the resultant mixture for 30 minutes. After completion
of the reaction, the reaction solution was concentrated under reduced pressure and the
obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate
= 1/1) to obtain a compound. The obtained compound was used to obtain the title
20 compound (355 mg, two step yield 48%) by synthesis in a similar manner to Reference
Synthesis Example 291.

[0536] Reference Synthesis Example 315

1-(4-Methoxybenzyl)-6-methyl-4-(methylthio)-1,3,5-triazin-2(1H)-one

- 25 A crude product of methyl
N-[(4-methoxyphenyl)methyl]amino]carbonyl-carbamimidothioate was used to obtain
the title compound (631 mg, quantitative) by synthesis in a similar manner to Reference
Synthesis Example 292.

[0537] Reference Synthesis Example 316

[5-(Perfluoroethyl)thiophen-2-yl]methanol

A crude product of 5-(perfluoroethyl)thiophen-2-carboxylic acid was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference

5 Synthesis Example 217.

[0538] Reference Synthesis Example 317

2-{[5-(Perfluoroethyl)thiophen-2-yl]methyl}isoindoline-1,3-dione

The crude product of [5-(perfluoroethyl)thiophen-2-yl]methanol synthesized in
10 Reference Synthesis Example 316 was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 201.

[0539] Reference Synthesis Example 318

[5-(Perfluoroethyl)thiophen-2-yl]methanamine

15 The crude product of 2-{[5-(perfluoroethyl)thiophen-2-yl]methyl}isoindoline-1,3-dione synthesized in Reference Synthesis Example 317 was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 202.

[0540] Reference Synthesis Example 319

20 Methyl

N-[(1-[5-(perfluoroethyl)thiophen-2-yl]methyl)amino]carbonyl]-carbamimidothioate

The crude product of [5-(perfluoroethyl)thiophen-2-yl]methanamine synthesized in
Reference Synthesis Example 318 was used to obtain the title compound (5.00 mg, five
25 step yield 7%) by synthesis in a similar manner to Reference Synthesis Example 290.

[0541] Reference Synthesis Example 320

6-Methyl-4-(methylthio)-1-{[5-(perfluoroethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Methyl

N-([5-(perfluoroethyl)thiophen-2-yl]methyl)amino)carbonyl]-carbamimidothioate (5.00 mg, 0.0144 mmol) synthesized in Reference Synthesis Example 319 was used to obtain the title compound (20.0 mg, yield 37%) by synthesis in a similar manner to

5 Reference Synthesis Example 292.

[0542] Reference Synthesis Example 321

Methyl

N-([4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl]methyl)amino)carbonyl]-carbamimidothi
oate

10

(4,5,6,7-Tetrahydrobenzo[b]thiophen-2-yl)methanamine (355 mg, 2.12 mmol) was used to obtain the title compound (410 mg, yield 68%) by synthesis in a similar manner to Reference Synthesis Example 290.

[0543] Reference Synthesis Example 322

15 4-(Methylthio)-1-[(4,5,6,7-tetrahydro-benzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)
-one

Methyl

20 N-([4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]amino)carbonyl)-carbamimidothi
oate (200 mg, 0.706 mmol) synthesized in Reference Synthesis Example 321 was used to obtain the title compound (130 mg, yield 63%) by synthesis in a similar manner to Reference Synthesis Example 296.

[0544] Reference Synthesis Example 323

25 6-Methyl-4-(methylthio)-1-[(4,5,6,7-tetrahydrobenzo[b]thiophene-2-yl)methyl]-1,3,5-tria
zin-2(1H)-one

Methyl

N-([4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]amino)carbonyl)-carbamimidothi
oate (210 mg, 0.741 mmol) synthesized in Reference Synthesis Example 321 was used to

obtain the title compound (100 mg, yield 44%) by synthesis in a similar manner to Reference Synthesis Example 292.

[0545] Reference Synthesis Example 324

Methyl N-({[(adamantan-1-yl)methyl]amino}carbonyl)-carbamimidothioate

5

Adamantan-1-ylmethanamine (118 μ L, 0.666 mmol) was used to obtain a crude product (250 mg) of the title compound by synthesis in a similar manner to Reference Synthesis Example 290.

[0546] Reference Synthesis Example 325

10 1-(Adamantan-1-ylmethyl)-6-methyl-4-(methylthio)-1,3,5-triazin-2(1H)-one

The crude product of methyl

N-({[(adamantan-1-yl)methyl]amino}carbonyl)-carbamimidothioate (250 mg) synthesized in Reference Synthesis Example 324 was used to obtain the title compound (103 mg, two step yield 51%) by synthesis in a similar manner to Reference Synthesis Example 292.

15

[0547] Reference Synthesis Example 326

Methyl N-({[(4-chlorobenzen-1-yl)methyl]amino}carbonyl)-carbamimidothioate

20 4-Chlorobenzylamine (142 mg, 1.00 mmol) was used to obtain the title compound (232 mg, yield 90%) by synthesis in a similar manner to Reference Synthesis Example 290.

[0548] Reference Synthesis Example 327

2-({[3-(4-Chlorobenzyl)ureido](methylthio)methylene}amino)-2-oxoethyl acetate

25

To a dichloromethane solution (2.0 mL) of methyl N-({[(4-chlorobenzen-1-yl)methyl]amino}carbonyl)-carbamimidothioate (257 mg, 1.00 mmol) synthesized in Reference Synthesis Example 326, triethylamine (279 μ L, 2.00 mmol) and 2-chloro-2-oxoethyl acetate (162 μ L, 1.50 mmol) were added and the

resultant mixture was stirred at room temperature for 1 hour. After completion of the reaction, the reaction solution was purified by silica gel column chromatography (hexane/ethyl acetate = 1/1) to obtain the title compound (221 mg, yield 62%).

[0549] Reference Synthesis Example 328

5 [1-(4-Chlorobenzyl)-4-(methylthio)-6-oxo-1,6-dihydro-1,3,5-triazin-2-yl]methyl acetate

A phosphorus oxychloride solution (3.0 mL) of
2-([3-(4-chlorobenzyl)ureido](methylthio)methylene)amino)-2-oxoethyl acetate (201
mg, 0.616 mmol) synthesized in Reference Synthesis Example 327 was stirred at 80°C
10 for 3 hours. After completion of the reaction, the reaction solution was added to
chloroform - saturated sodium bicarbonate aqueous solution and extraction from the
resultant mixture with chloroform was performed. The organic layer was dried over
anhydrous sodium sulfate and concentrated under reduced pressure. The obtained
residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1 →
15 0/1) to obtain the title compound (219 mg, quantitative).

[0550] Reference Synthesis Example 329

2-[(Methylthio)(3-{[5-(trifluoromethyl)thiophen-2-yl]methyl}ureido)methylene]amino]-2-oxoethyl acetate

20 Methyl

N-[(3-{[5-(trifluoromethyl)thiophen-2-yl]methyl}amino)carbonyl]-carbamimidothioate
(500 mg, 1.68 mmol) synthesized in Reference Synthesis Example 290 was used to
obtain the title compound (382 mg, yield 57%) by synthesis in a similar manner to
Reference Synthesis Example 327.

25 [0551] Reference Synthesis Example 330

[4-(Methylthio)-6-oxo-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,6-dihydro-1,3,5-triazin-2-yl]methyl acetate

2-[(Methylthio)(3-{[5-(trifluoromethyl)thiophen-2-yl]methyl}ureido)methylene]a

mino]-2-oxoethyl acetate (382 mg, 0.961 mmol) synthesized in Reference Synthesis Example 329 was used to obtain the title compound (276 mg, yield 76%) by synthesis in a similar manner to Reference Synthesis Example 328.

[0552] Reference Synthesis Example 331

5 6-Methoxy-4-(methylthio)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Tetramethoxymethane was used to obtain the title compound (64.4 mg, yield 95%) by synthesis in a similar manner to Reference Synthesis Example 292.

10 [0553] Reference Synthesis Example 332a

(R)-1-(4-Chloro-1,3,5-triazin-2-yl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol

Reference Synthesis Example 332b

(S)-1-(4-Chloro-1,3,5-triazin-2-yl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol

15 To a tetrahydrofuran (1.2 mL) - water (2.0 mL) solution of (R)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol (158 mg, 0.817 mmol) synthesized in Reference Synthesis Example 133, sodium carbonate (104 mg, 0.981 mmol) was added at 0°C and the resultant mixture was stirred at room temperature for 2 hours. After completion of the reaction, ethyl acetate was added to the reaction solution and the
20 resultant mixture was washed with water. Thereafter, the organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 7/3 → 1/1) to obtain a compound (156 mg, yield 62%) of Reference Synthesis Example 332a as the title compound.

25 Alternatively, (S)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol (188 mg, 0.972 mmol) synthesized in Reference Synthesis Example 138 was used to obtain a compound (189 mg, yield 63%) of 332b as the title compound by synthesis in a similar manner.

[0554] Reference Synthesis Example 333a

(R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-o

ne

Reference Synthesis Example 333b

(S)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

5

To an acetic acid (1.6 mL) - water (0.30 mL) solution of (R)-1-(4-chloro-1,3,5-triazin-2-yl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol (155 mg, 0.505 mmol) synthesized in Reference Synthesis Example 332a, sodium acetate (155 mg, 1.89 mmol) was added and the resultant mixture was stirred at 90°C for 2 hours.

10 After completion of the reaction, water was added to the reaction solution and the resultant mixture was cooled to 0°C. The precipitated solid was collected by filtration and the solid was washed with water to obtain a compound (122 mg, yield 77%) of Reference Synthesis Example 333a as the title compound.

Alternatively,

15 (S)-1-(4-chloro-1,3,5-triazin-2-yl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol (189 mg, 0.616 mmol) synthesized in Reference Synthesis Example 332b was used to obtain a compound (108 mg, yield 61%) of 333b as the title compound by synthesis in a similar manner.

[0555] Reference Synthesis Example 334

20 (R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

To 1-(4-methoxybenzyl)-6-methyl-4-(methylthio)-1,3,5-triazin-2(1H)-one (1.39 g, 5.0 mmol) synthesized in Reference Synthesis Example 315, trifluoroacetic acid (10 mL) was added and the resultant mixture was stirred for 1 hour under reflux. After completion of the reaction, the reaction solution was concentrated under reduced pressure and 1,4-dioxane (10 mL), (R)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol (966 mg, 5.00 mmol) synthesized in Reference Synthesis Example 133, and N,N-diisopropylethylamine (1.92 g, 10.0 mmol) were added to the concentrated solution,

followed by refluxing the resultant mixture for 2 hours. After completion of the reaction, water was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The obtained residue was

5 purified by silica gel column chromatography (ethyl acetate/methanol = 10/1) to obtain the title compound (1.41 g, yield 94%).

[0556] Reference Synthesis Example 335

1-(4-Chloro-1,3,5-triazin-2-yl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol

10 4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol (300 mg, 1.55 mmol) synthesized in Reference Synthesis Example 40b was used to obtain the title compound (445 mg, yield 94%) by synthesis in a similar manner to Reference Synthesis Example 341.

[0557] Reference Synthesis Example 336

4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

15

1-(4-Chloro-1,3,5-triazin-2-yl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol (235 mg, 0.766 mmol) synthesized in Reference Synthesis Example 335 was used to obtain the title compound (127 mg, yield 58%) by synthesis in a similar manner to Reference Synthesis Example 342.

20 [0558] Reference Synthesis Example 337

(R)-1-(4-Chloro-N-[1-(4-fluorophenyl)pyrrolidin-3-yl]-1,3,5-triazin-2-amine

To a tetrahydrofuran solution (2.0 mL) - water (0.80 mL) solution of 2,4-dichloro-1,3,5-triazine (166 mg, 1.11 mmol) and sodium carbonate (141 mg, 1.33

25 mmol), (R)-1-(4-fluorophenyl)pyrrolidin-3-amine hydrochloride (200 mg, 1.11 mmol) synthesized in Reference Synthesis Example 30 was added at 0°C and the resultant mixture was stirred for 3 hours. After completion of the reaction, water was added to the reaction solution and extraction from the resultant mixture with ethyl acetate was performed. The organic layer was dried over anhydrous magnesium sulfate and

concentrated under reduced pressure to obtain the title compound as a crude product.

[0559] Reference Synthesis Example 338

(R)-4-{[1-(4-Fluorophenyl)pyrrolidin-3-yl]amino}-1,3,5-triazin-2(1H)-one

5 To (R)-4-chloro-N-[1-(4-fluorophenyl)pyrrolidin-3-yl]-1,3,5-triazin-2-amine synthesized in Reference Synthesis Example 337, sodium acetate (153 mg, 1.11 mmol), acetic acid (2.0 mL), and water (0.40 mL) were added and the resultant mixture was stirred at 90°C for 5 hours. After completion of the reaction, the reaction solution was added to saturated sodium bicarbonate aqueous solution and extraction from the resultant
10 mixture with ethyl acetate was performed. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure and the obtained solid was washed with ethyl acetate to obtain the title compound (65.0 mg, two step yield 15%).

[0560] Reference Synthesis Example 339

6-Methyl-4-[4-(p-tolyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

15

1-(p-Tolyl)piperazine (250 mg, 1.42 mmol) was used instead of (R)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol to obtain the title compound (312 mg, yield 77%) by synthesis in a similar manner to Reference Synthesis Example 334.

[0561] Reference Synthesis Example 340

20 4-[4-(4-Fluorophenyl)piperazin-1-yl]-6-methyl-1,3,5-triazin-2(1H)-one

1-(4-Fluorophenyl)piperazine (1.25 g, 6.94 mmol) was used instead of (R)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol to obtain the title compound (1.50 g, yield 82%) by synthesis in a similar manner to Reference Synthesis Example 334.

25 [0562] Reference Synthesis Example 341

2-Chloro-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazine

To a tetrahydrofuran (10 mL) - water (4.0 mL) solution of 4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine hydrochloride (1.43 g, 6.67 mmol) and

sodium carbonate (1.56 g, 14.7 mol), 2,4-dichloro-1,3,5-triazine (1.00 g, 6.67 mmol) was added at 0°C and the resultant mixture was stirred for 1 hour. After completion of the reaction, ethyl acetate was added to the reaction solution and the resultant mixture was washed with each of saturated ammonium chloride aqueous solution and saturated sodium chloride aqueous solution. The obtained organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (chloroform) to obtain the title compound (1.64 g, yield 85%).

[0563] Reference Synthesis Example 342

10 4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

To an acetic acid (9.0 mL) - water (1.8 mL) solution of 2-chloro-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazine (900 mg, 3.10 mmol) synthesized in Reference Synthesis Example 341, sodium acetate (900 mg, 11.0 mmol) was added and the resultant mixture was stirred at 90°C for 1 hour. After completion of the reaction, the reaction solution was concentrated under reduced pressure and water was added to the obtained residue, followed by stirring the resultant mixture at 0°C. The precipitated solid was collected by filtration and the solid was dried under reduced pressure to obtain the title compound (750 mg, yield 89%).

20 [0564] Reference Synthesis Example 343

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

4-(4-Fluorophenyl)-1,2,3,6-tetrahydropyridine hydrochloride (250 mg, 1.42 mmol) was used instead of (R)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol to obtain the title compound (85.0 mg, two step yield 60%) by synthesis in a similar manner to Reference Synthesis Example 334.

[0565] Reference Synthesis Example 344

2-Chloro-4-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazine

4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2,3,6-tetrahydropyridine 2,2,2-trifluoroacetate (5.94 g, 19.4 mmol) was used to obtain the title compound (4.21 g, yield 96%) by synthesis in a similar manner to Reference Synthesis Example 341.

5 [0566] Reference Synthesis Example 345

4-[4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2-ol

2-Chloro-4-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazine (4.21 g, 13.1 mmol) synthesized in Reference Synthesis Example 344 was used to obtain the title compound (5.17 g, quantitative) by synthesis in a similar manner to Reference Synthesis Example 342.

[0567] Reference Synthesis Example 346

15 4-[4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

4-[4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2-ol (3.32 g, 10.9 mmol) synthesized in Reference Synthesis Example 345 and 2-(chloromethyl)-5-(trifluoromethyl)thiophene were used to obtain the title compound (2.36 g, yield 46%) by synthesis in a similar manner to Reference Synthesis Example 16.

[0568] Reference Synthesis Example 347

1-(4-Chlorobenzyl)-4-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

25 To a chloroform solution (3.0 mL) of

4-chloro-1-(4-chlorobenzyl)-1,3,5-triazin-2(1H)-one (128 mg, 0.500 mmol) synthesized in Reference Synthesis Example 11, triethylamine (70.0 μ L, 0.500 mmol) and 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2,3,6-tetrahydropyridine (156 mg, 0.750 mmol) were added and the resultant mixture was stirred at room temperature for 4

hours. After completion of the reaction, water was added to the reaction solution and extraction from the resultant mixture with chloroform was performed. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column

5 chromatography (hexane/ethyl acetate = 1/0 → 4/1) to obtain the title compound (86.0 mg, yield 40%).

[0569] Reference Synthesis Example 348

4-(4-Hydroxypiperidin-1-yl)-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10

To a 1,4-dioxane solution (1.0 mL) of 6-methyl-4-(methylthio)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (321 mg, 1.00 mmol) synthesized in Reference Synthesis Example 292, 4-piperidinol (202 mg, 2.00 mmol) was added and the resultant mixture was stirred under
15 reflux by heating for 10 hours. After completion of the reaction, the reaction solution was concentrated under reduced pressure and the obtained residue was purified by silica gel column chromatography (ethyl acetate/methanol = 1/0 → 10/1) to obtain the title compound (374 mg, yield 100%).

[0570] Reference Synthesis Example 349

20 1-(6-Methyl-4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)piperidin-4-yl methanesulfonate

To a dichloromethane solution (2.0 mL) of 4-(4-hydroxypiperidin-1-yl)-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (374 mg, 1.00 mmol) synthesized in Reference Synthesis Example
25 348 and triethylamine (557 µL, 4.00 mmol), methanesulfonyl chloride (155 µL, 2.00 mmol) was added dropwise at 0°C and the resultant mixture was stirred at room temperature for 1 hour. After the reaction, water was added to the reaction solution and extraction from the resultant mixture with dichloromethane was performed. The

obtained organic layer was dried over anhydrous sodium sulfate and thereafter concentrated under reduced pressure to obtain the title compound (453 mg, yield 100%).

[0571] Reference Synthesis Example 350

tert-Butyl

5 4-(6-methyl-4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)piperazine-1-carboxylate

6-Methyl-4-(methylthio)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (309 mg, 0.962 mmol) synthesized in Reference Synthesis Example 292 and
10 tert-butyl piperazine-1-carboxylate (269 mg, 1.44 mmol) were used to obtain the title compound (429 mg, yield 97%) by synthesis in a similar manner to Reference Synthesis Example 348.

[0572] Reference Synthesis Example 351

6-Methyl-4-(piperazin-1-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2
15 (1H)-one

tert-Butyl

4-(6-methyl-4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)piperazine-1-carboxylate (429 mg, 0.933 mmol) synthesized in Reference
20 Synthesis Example 350 was used to obtain the title compound (335 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 132.

[0573] Reference Synthesis Example 352

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-(4-methoxybenzyl)-1,3,5-triazin-2
25 (1H)-one

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one (294 mg, 1.08 mmol) synthesized in Reference Synthesis Example 342 and
1-(chloromethyl)-4-methoxybenzene (162 μ L, 1.19 mmol) were used to obtain the title compound (346 mg, yield 82%) by synthesis in a similar manner to Reference Synthesis

Example 16.

[0574] Reference Synthesis Example 353

4-[5-Chloro-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-[5-(trifluoromethyl)thiophen-2-yl]methyl]-1,3,5-triazin-2(1H)-one

5

4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (10.0 mg, 0.0212 mmol) synthesized in Synthesis Example 190 was used to obtain the title compound as a crude product by synthesis in a similar manner to Reference Synthesis Example 68.

10

[0575] Reference Synthesis Example 354

tert-Butyl 4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridine-1(2H)-carboxylate

A mixture containing equal quantity of both enantiomers of tert-butyl 4-(4-fluorophenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate (700 mg, 1.85 mmol) synthesized in Reference Synthesis Example 131 and Reference Synthesis Example 136 was used to obtain the title compound (500 mg, yield 92%) by synthesis in a similar manner to Reference Synthesis Example 133.

15

[0576] Reference Synthesis Example 355

tert-Butyl 3,3-bis(hydroxymethyl)-4-oxo-piperidine-1-carboxylate

20

To tert-butyl 4-oxopiperidine-1-carboxylate (1.00 g, 5.02 mmol), water (2.0 mL) and potassium carbonate (10.4 mg, 0.0750 mmol) were added at room temperature and thereafter the mixture was heated to 40°C, followed by adding formaldehyde (36% aqueous solution) (0.795 mg, 9.54 mmol) and stirring the resultant mixture for 2 hours. After completion of the reaction, the reaction solution was purified by silica gel column chromatography (hexane/ethyl acetate = 1/2) to obtain the title compound (52.0 mg, yield 4%).

25

[0577] Reference Synthesis Example 356

tert-Butyl 3,3-bis[(methoxymethoxy)methyl]-4-oxopiperidine-1-carboxylate

To a dichloromethane solution (7.0 mL) of tert-butyl 3,3-bis(hydroxymethyl)-4-oxopiperidine-1-carboxylate (376 mg, 1.45 mmol) synthesized in Reference Synthesis Example 355, N,N-diisopropylpropylethylamine (469 mg, 3.63 mmol) and chloro(methoxy)methane (245 mg, 3.05 mmol) were added and the resultant mixture was stirred at room temperature for 1 day. After completion of the reaction, water was added to the reaction solution and extraction from the resultant reaction mixture with chloroform was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 1/1 → 1/2) to obtain the title compound (233 mg, yield 46%).

[0578] Reference Synthesis Example 357

tert-Butyl

5,5-bis[(methoxymethoxy)methyl]-4-[[[(trifluoromethyl)sulfonyl]oxy]-5,6-dihydropyridine-1(2H)-carboxylate

To a tetrahydrofuran solution (6.0 mL) of tert-butyl 3,3-bis[(methoxymethoxy)methyl]-4-oxopiperidine-1-carboxylate (233 mg, 0.671 mmol) synthesized in Reference Synthesis Example 356, sodium bis(trimethylsilyl)amide (1.0 M tetrahydrofuran solution) (740 μ L, 0.740 mmol) was added at 0°C and the resultant mixture was stirred for 5 minutes. Thereafter, 1,1,1-trifluoro-N-phenyl-N-[(trifluoromethyl)sulfonyl]methanesulfonamide (287 mg, 0.804 mmol) was added to the mixture and the resultant mixture was stirred at room temperature for 1 day. After completion of the reaction, the reaction solution was concentrated under reduced pressure and the obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 1/0 → 1/1) to obtain the title compound (211 mg, yield 66%).

[0579] Reference Synthesis Example 358

tert-Butyl

4-(4-fluorophenyl)-5,5-bis[(methoxymethoxy)methyl]-5,6-dihydropyridine-1(2H)-carboxylate

tert-Butyl

- 5 5,5-bis[(methoxymethoxy)methyl]-4-[[trifluoromethyl)sulfonyl]oxy}-5,6-dihydropyridine-1(2H)-carboxylate (211 mg, 0.439 mmol) synthesized in Reference Synthesis Example 357 was used to obtain the title compound (104 mg, yield 56%) by synthesis in a similar manner to Reference Synthesis Example 139.

[0580] Reference Synthesis Example 359

- 10 4-(4-fluorophenyl)-3,3-bis[(methoxymethoxy)methyl]-1,2,3,6-tetrahydropyridine trifluoroacetate

To a dichloromethane solution (0.80 mL) of tert-butyl

- 15 4-(4-fluorophenyl)-5,5-bis[(methoxymethoxy)methyl]-5,6-dihydropyridine-1(2H)-carboxylate (40.0 mg, 0.0940 mmol) synthesized in Reference Synthesis Example 358, trifluoroacetic acid (40.0 μ L, 0.538 mmol) was added at room temperature and the resultant mixture was stirred for 1 day. After completion of the reaction, the reaction solution was concentrated under reduced pressure to obtain a crude product of the title compound.

- 20 [0581] Reference Synthesis Example 360

tert-Butyl 4-(4-fluorophenyl)-3,3-bis[(methoxymethoxy)methyl]piperidine-1-carboxylate

tert-Butyl

- 25 4-(4-fluorophenyl)-5,5-bis[(methoxymethoxy)methyl]-5,6-dihydropyridine-1(2H)-carboxylate (50.0 mg, 0.118 mmol) synthesized in Reference Synthesis Example 358 was used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 98. The crude product was used in the next process as it was.

[0582] Reference Synthesis Example 361

4-(4-Fluorophenyl)-3,3-bis[(methoxymethoxy)methyl]piperidine

The crude product of tert-butyl

4-(4-fluorophenyl)-3,3-bis[(methoxymethoxy)methyl]piperidine-1-carboxylate

5 synthesized in Reference Synthesis Example 360 was used to obtain the title compound (58.8 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 132.

[0583] Reference Synthesis Example 362

4-{4-(4-Fluorophenyl)-3,3-bis[(methoxymethoxy)methyl]piperidin-1-yl}-6-methyl-1-[5
10 -(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

4-(4-Fluorophenyl)-3,3-bis[(methoxymethoxy)methyl]piperidine (58.8 mg, 0.180 mmol) synthesized in Reference Synthesis Example 361 was used to obtain the title compound (14.0 mg, yield 16%) by synthesis in a similar manner to Reference Synthesis
15 Example 348.

[0584] Reference Synthesis Example 363

4-[11-(4-Fluorophenyl)-2,4-dioxo-8-azaspiro[5.5]undecan-8-yl]-6-methyl-1-[5-(trifluoro
methyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

20 To a 1,4-dioxane solution (600 μ L) of
4-{4-(4-fluorophenyl)-3,3-bis[(methoxymethoxy)methyl]piperidin-1-yl}-6-methyl-1-[5-
(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (10.0 mg, 16.6 μ mol)
synthesized in Reference Synthesis Example 362, 12 M hydrochloric acid (200 μ L) was
added and the resultant mixture was stirred at 60°C for 2 hours. To saturated sodium
25 carbonate - chloroform solution, the reaction solution was added dropwise and extraction
from the resultant mixture with chloroform was performed. The organic layer was dried
over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained
residue was purified by silica gel column chromatography (hexane/ethyl acetate = 1/4 $\rightarrow$
0/1) to obtain the title compound (6.60 mg, yield 76%).

[0585] Reference Synthesis Example 364

tert-Butyl (R)-4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridine-1(2H)-carboxylate

To a 1,4-dioxane (500 μ L) - water (100 μ L) solution of tert-butyl

- 5 (R)-4-(4-fluorophenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate (50.0 mg, 0.132 mmol) synthesized in Reference Synthesis Example 131, lithium hydroxide monohydrate (17.0 mg, 0.405 mmol) was added and the resultant mixture was stirred at 115°C for 10 hours. After completion of the reaction, the reaction solution was analyzed with chiral column chromatography to measure optical purity.

10 [0586] Reference Synthesis Example 365

tert-Butyl (S)-4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridine-1(2H)-carboxylate

To a 1,4-dioxane (500 μ L) - water (100 μ L) solution of tert-butyl

- 15 (S)-4-(4-fluorophenyl)-5-(pivaloyloxy)-5,6-dihydropyridine-1(2H)-carboxylate (50.0 mg, 0.132 mmol) synthesized in Reference Synthesis Example 136, lithium hydroxide monohydrate (17.0 mg, 0.405 mmol) was added and the resultant mixture was stirred at 115°C for 10 hours. After completion of the reaction, the reaction solution was analyzed with chiral column chromatography to measure optical purity.

[0587] Reference Synthesis Example 366

- 20 1-(4-Chloro-1,3,5-triazin-2-yl)-4-(4-chlorophenyl)piperidin-4-ol

4-(4-Chlorophenyl)piperidin-4-ol (705 mg, 3.33 mmol) was used to obtain the title compound (370 mg, yield 34%) by synthesis in a similar manner to Reference Synthesis Example 341.

25 [0588] Reference Synthesis Example 367

4-[4-(4-Chlorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

1-(4-Chloro-1,3,5-triazin-2-yl)-4-(4-chlorophenyl)piperidin-4-ol (200 mg, 0.615 mmol) synthesized in Reference Synthesis Example 366 was used to obtain the title

compound (400 mg) as a crude product by synthesis in a similar manner to Reference Synthesis Example 342.

[0589] Reference Synthesis Example 368

7-Benzyl-9-(4-fluorophenyl)-3-oxa-7-azabicyclo[3.3.1]nonan-9-ol

5

To 4-fluorobenzene magnesium bromide (2M diethyl ether solution) (5.0 mL, 10.0 mmol), (1R, 5S)-7-benzyl-3-oxa-7-azabicyclo[3.3.1]nonan-9-one (462 mg, 2.00 mmol) was added dropwise at 0°C and the resultant mixture was stirred at room temperature for 1 minute. After completion of the reaction, water and 1 M sodium hydroxide aqueous solution were added to the reaction solution and extraction from the resultant reaction mixture with ethyl acetate was performed. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel (amino-based) column chromatography (hexane/ethyl acetate = 5/1) to obtain the title compound (156 mg, yield 24%).

10

[0590] Reference Synthesis Example 369

9-(4-Fluorophenyl)-3-oxa-7-azabicyclo[3.3.1]nonan-9-ol

7-Benzyl-9-(4-fluorophenyl)-3-oxa-7-azabicyclo[3.3.1]nonan-9-ol (72.5 mg, 0.231 mmol) synthesized in Reference Synthesis Example 368 was used to obtain the title compound (35.6 mg, yield 65%) by synthesis in a similar manner to Reference Synthesis Example 226.

20

[0591] Reference Synthesis Example 370

[trans-3-Amino-1-(4-fluorophenyl)cyclobutyl]methanol

25

To a tetrahydrofuran solution (5.0 mL) of methyl trans-1-(4-fluorophenyl)-3-hydroxycyclobutane-carboxylate (177 mg, 5.00 mmol) synthesized in Reference Synthesis Example 255, lithium aluminum hydride (38.0 mg, 1.00 mmol) was added and the resultant mixture was stirred at room temperature for 10 minutes. After completion of the reaction, 1 M sodium hydroxide aqueous solution and

tetrahydrofuran were added to the reaction solution and the resultant mixture was filtered with Celite, followed by concentrating the filtrate under reduced pressure. To the obtained crude product, dichloromethane was added and the resultant mixture was dried over anhydrous sodium sulfate and pressure was reduced to obtain the title compound (95.3 mg, yield 98%).

[0592] Reference Synthesis Example 371

Methyl

N-(trifluoroacetyl)-N'-([5-(trifluoromethyl)thiophen-2-yl]methyl)carbamoyl)carbamimidothioate

10

Methyl

N-([5-(trifluoromethyl)thiophen-2-yl]methyl)amino)carbonyl]-carbamimidothioate (29.7 mg, 0.100 mmol) synthesized in Reference Synthesis Example 290 and trifluoroperacetic acid (1.0 mL) were used to obtain a crude product of the title compound by synthesis in a similar manner to Reference Synthesis Example 327.

15

[0593] Reference Synthesis Example 372

4-(Methylthio)-6-(trifluoromethyl)-1-[5-(trifluoromethyl)thiophen-2-yl]methyl]-1,3,5-triazin-2(1H)-one

20

The crude product of methyl

N-(trifluoroacetyl)-N'-([5-(trifluoromethyl)thiophen-2-yl]methyl)carbamoyl)carbamimidothioate synthesized in Reference Synthesis Example 371 was used to obtain the title compound (21.3 mg, yield 57%) by synthesis in a similar manner to Reference Synthesis Example 328.

25

[0594] Reference Synthesis Example 373

Methyl

N-([2-(trifluoromethyl)thiazol-5-yl]methyl)amino)carbonyl]-carbamimidothioate

[2-(Trifluoromethyl)thiazol-5-yl]methanamine (370 mg, 2.03 mmol) was used to

obtain the title compound (607 mg, quantitative) by synthesis in a similar manner to Reference Synthesis Example 289.

[0595] Reference Synthesis Example 374

6-Methyl-4-(methylthio)-1-{{2-(trifluoromethyl)thiazol-5-yl}methyl}-1,3,5-
 5 triazin-2(1H)-one

Methyl

10 N-[({{2-(trifluoromethyl)thiazol-5-yl}methyl}amino)carbonyl]-carbamimidothioate (607 mg, 2.03 mmol) synthesized in Reference Synthesis Example 373 was used to obtain the title compound (630 mg, yield 97%) by synthesis in a similar manner to Reference Synthesis Example 292.

[0596] Synthesis Example 1

1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

15 To an N,N-dimethylformamide solution (170 mL) of 4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one (8.48 g, 30.8 mmol) synthesized in Reference Synthesis Example 3, 4-chlorobenzyl bromide (6.32 g, 30.9 mmol) and potassium carbonate (5.11 g, 37.0 mmol) were added and the resultant mixture was stirred at 50°C for 1 hour and at 70°C for 3 hours. To the reaction solution,
 20 4-chlorobenzyl bromide (1.90 g, 9.25 mmol) and potassium carbonate (2.13 g, 15.4 mmol) were added and the resultant reaction solution was stirred at 70°C further for 2 hours. After the completion of the reaction, water was added to the reaction solution and a resultant solid was filtered and was dried under reduced pressure. The resultant solid was suspended in ethyl acetate and the suspension was stirred at room temperature
 25 for 10 minutes. The solid was filtered and was dried under reduced pressure. These operations from suspending of the solid to the second drying of the solid were performed twice to obtain the title compound (6.49 g, yield: 53%).

[0597] In Synthesis Examples 2 to 10, each title compound was synthesized in a similar manner to Synthesis Example 1. The names of the synthesized compound and

the synthesis yields are indicated below.

[0598] Synthesis Example 2

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-(4-nitrobenzyl)-1,3,5-triazin-2(1H)-one

5 Yield: 18%

[0599] Synthesis Example 3

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-[4-(trifluoromethyl)benzyl]-1,3,5-triazin-2(1H)-one

10 Yield: 23%

[0600] Synthesis Example 4

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-(4-methylbenzyl)-1,3,5-triazin-2(1H)-one

Yield: 24%

15 [0601] Synthesis Example 5

1-[4-(tert-Butyl)benzyl]-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 25%

[0602] Synthesis Example 6

20 4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-[4-(trifluoromethoxy)benzyl]-1,3,5-triazin-2(1H)-one

Yield: 12%

[0603] Synthesis Example 7

25 1-(3-Chlorobenzyl)-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 30%

[0604] Synthesis Example 8

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-[3-(trifluoromethyl)benzyl]-1,3,5-triazin-2(1H)-one

ne

Yield: 8.5%

[0605] Synthesis Example 9

5 4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-(3-methylbenzyl)-1,3,5-triazin-2(1H)-one

Yield: 37%

[0606] Synthesis Example 10

1-(3-Fluorobenzyl)-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

10

Yield: 23%

[0607] Synthesis Example 11

1-{[3-(tert-Butyl)-1-methyl-1H-pyrazol-5-yl]methyl}-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

15

To an N,N-dimethylformamide solution (2 mL) of 4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one (50.0 mg, 0.18 mmol) synthesized in Reference Synthesis Example 3, 3-(tert-butyl)-5-(chloromethyl)-1-methyl-1H-pyrazole (50 mg, 0.21 mmol) synthesized in Reference Synthesis Example 61, potassium carbonate (75.3 mg, 0.54 mmol), and sodium iodide (2.7 mg, 0.02 mmol) were added and the resultant mixture was stirred at 80°C for 8 hours. To the reaction solution, water was added and extraction with chloroform from the resultant reaction solution was performed three times. The organic layer was dried over anhydrous magnesium sulfate and was concentrated under reduced pressure. The resultant residue was purified by preparative thin-layer chromatography (ethyl acetate/chloroform=1/1) and the purified product was recrystallized from chloroform/hexane to obtain the title compound (8.3 mg, yield: 11%).

25

[0608] In Synthesis Examples 12 to 24, each title compound was synthesized in a similar manner to Synthesis Example 11. The names of the synthesized compound and

the synthesis yields are indicated below.

[0609] Synthesis Example 12

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-(4-methoxybenzyl)-1,3,5-triazin-2(1H)-one

5 Yield: 25%

[0610] Synthesis Example 13

1-(2-Fluorobenzyl)-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 21%

10 [0611] Synthesis Example 14

Ethyl

5-({4-[4-(4-fluorophenyl)piperazin-1-yl]-2-oxo-1,3,5-triazin-1(2H)-yl}methyl)furan-2-carboxylate

15 Yield: 4.7%

[0612] Synthesis Example 15

1-{[3-(tert-Butyl)-1H-pyrazol-5-yl]methyl}-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

20 Yield: 13%

[0613] Synthesis Example 16

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[2-oxo-4-(trifluoromethyl)pyridin-1(2H)-yl]methyl}-1,3,5-triazin-2(1H)-one

25 Yield: 19%

[0614] Synthesis Example 17

1-(Benzo[d]thiazol-6-ylmethyl)-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 10%

[0615] Synthesis Example 18

1-[(5-Chlorobenzofuran-2-yl)methyl]-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

5

Yield: 26%

[0616] Synthesis Example 19

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

10

Yield: 36%

[0617] Synthesis Example 20

1-[(5-Chlorothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

15

Yield: 26%

[0618] Synthesis Example 21

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[5-(trifluoromethyl)furan-2-yl]methyl}-1,3,5-triazin-2(1H)-one

20

Yield: 29%

[0619] Synthesis Example 22

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[6-(trifluoromethyl)pyridin-3-yl]methyl}-1,3,5-triazin-2(1H)-one

25

Yield: 2.3%

[0620] Synthesis Example 23

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-5-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 2.1%

[0621] Synthesis Example 24

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[5-phenyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]
5 methyl}-1,3,5-triazin-2(1H)-one

Yield: 40%

[0622] Synthesis Example 25

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5
10 -triazin-2(1H)-one

To an N,N-dimethylformamide solution (1 mL) of

4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one (50.0 mg, 0.18 mmol)
synthesized in Reference Synthesis Example 3, [5-(trifluoromethyl)thiophen-2-yl]methyl
15 4-methylbenzenesulfonate (121 mg, 0.36 mmol) synthesized in Reference Synthesis
Example 51 and potassium carbonate (99 mg, 0.72 mmol) were added and the resultant
mixture was stirred at 80°C for 2 hours. To the reaction solution,
[5-(trifluoromethyl)thiophen-2-yl]methyl 4-methylbenzenesulfonate (500 mg, 0.48
mmol) was added and the resultant reaction solution was stirred further for 2 hours.
20 After the completion of the reaction, water was added to the reaction solution and
extraction with ethyl acetate from the resultant reaction solution was performed three
times. The organic layer was washed with brine, was dried over anhydrous magnesium
sulfate, and was concentrated under reduced pressure. The resultant residue was
purified by silica gel column chromatography (hexane/ethyl acetate=4/1) to obtain the
25 title compound (9.4 mg, yield: 12%).

[0623] In Synthesis Examples 26 to 30 and 32 to 33, each title compound was
synthesized in a similar manner to Synthesis Example 25. The names of the synthesized
compound and the synthesis yields are indicated below.

[0624] Synthesis Example 26

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 4.4%

5 [0625] Synthesis Example 27

1-[(3-Cyclohexyl-1H-pyrazol-1-yl)methyl]-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 2.0%

10 [0626] Synthesis Example 28

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[5-(2,2,2-trifluoroethoxy)pyridin-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 8.3%

15 [0627] Synthesis Example 29

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[2-(trifluoromethyl)thiazol-4-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 8.9%

20 [0628] Synthesis Example 30

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[2-(trifluoromethyl)thiazol-5-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 3.0%

25 [0629] Synthesis Example 31

1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)piperazin-1-yl]-6-methoxy-1,3,5-triazin-2(1H)-one

The synthesis was performed using

3-(4-chlorobenzyl)-6-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazine-2,4(1H,3H)-dione (1.11 g, 2.67 mmol) synthesized in Reference Synthesis Example 19 in a similar manner to Synthesis Example 66 to obtain the title compound (180 mg, yield: 16%).

[0630] Synthesis Example 32

5 4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[4-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 16%

[0631] Synthesis Example 33

10 4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{2-[3-(trifluoromethyl)-1H-pyrazol-1-yl]ethyl}-1,3,5-triazin-2(1H)-one

Yield: 9.0%

[0632] Synthesis Example 34

15 Ethyl
5-(tert-butyl)-1-({4-[4-(4-fluorophenyl)piperazin-1-yl]-2-oxo-1,3,5-triazin-1(2H)-yl}methyl)-1H-pyrazole-3-carboxylate

Synthesis Example 35

20 Ethyl
3-(tert-butyl)-1-({4-[4-(4-fluorophenyl)piperazin-1-yl]-2-oxo-1,3,5-triazin-1(2H)-yl}methyl)-1H-pyrazole-5-carboxylate

A tetrahydrofuran solution (1 mL) of ethyl 3-(tert-butyl)-1H-pyrazole-5-carboxylate (196 mg, 1.00 mmol), paraformaldehyde (36 mg, 1.20 mmol), and
25 1,8-diazabicyclo[5.4.0] undeca-7-ene (15 μ L, 0.1 mmol) was stirred at room temperature for 1 day. After the completion of the reaction, the reaction solution was filtered and to the filtrate, triethylamine (209 μ L, 1.50 mmol) and p-toluenesulfonyl chloride (229 mg, 1.20 mmol) were added, followed by stirring the resultant reaction solution at room

temperature for 1 hour. To the reaction solution, water was added and the resultant reaction solution was extracted with ethyl acetate, followed by concentrating the resultant organic layer under reduced pressure. To an N,N-dimethylformamide solution (1 mL) of the above-resultant residue and

- 5 4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one (138 mg, 0.50 mmol) synthesized in Reference Synthesis Example 3, potassium carbonate (138 mg, 1.00 mmol) and sodium iodide (7.5 mg, 0.05 mmol) were added and the resultant mixture was stirred at 90°C for 1 day. After the completion of the reaction, water was added to the reaction solution and extraction with ethyl acetate from the resultant reaction solution
- 10 was performed. The organic layer was concentrated under reduced pressure and the resultant residue was purified by preparative high performance liquid chromatography to obtain the compound of Synthesis Example 34 (yield: 2%) and the compound of Synthesis Example 35 (yield: 2%).

[0633] Synthesis Example 36

- 15 1-(4-Chlorobenzyl)-6-(dimethylamino)-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

- To an N,N-dimethylformamide solution (5 mL) of 4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one (100 mg, 0.36 mmol)
- 20 synthesized in Reference Synthesis Example 3, 4-chlorobenzyl chloride (69 mg, 0.43 mmol), potassium carbonate (59 mg, 0.43 mmol), and sodium iodide (54 mg, 0.36 mmol) were added and the resultant mixture was stirred at 90°C for 5 hours. The reaction solution was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography to obtain the title compound (1 mg, yield: 1%).

- 25 [0634] Synthesis Example 37

4-[4-(3-Methoxyphenyl)piperazin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

A 1,4-dioxane solution (1.0 mL) of

4-(piperazin-1-yl)-1-{{3-(trifluoromethyl)-1H-pyrazol-1-yl}methyl}-1,3,5-triazin-2(1H)-one (50 mg, 0.07 mmol) synthesized in Reference Synthesis Example 17,

1-bromo-3-methoxybenzene (8.6 μ L, 0.07 mmol), tris(dibenzylideneacetone) dipalladium (0) (6.2 mg, 0.007 mmol),

- 5 dicyclohexyl(2',4',6'-triisopropyl-[1,1'-biphenyl]-2-yl)phosphine (6.5 mg, 0.01 mmol), and cesium carbonate (44 mg, 0.14 mmol) was stirred in a nitrogen atmosphere at 80°C for 1 day. After the completion of the reaction, water was added to the reaction solution and extraction with ethyl acetate from the resultant reaction solution was performed three times. The organic layer was washed with brine, was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure. The resultant residue was
10 purified by preparative thin-layer chromatography (hexane/ethyl acetate=1/1 (v/v)) to obtain the title compound (9.4 mg, yield: 32%).

[0635] In Synthesis Examples 38 to 63, each title compound was synthesized in a similar manner to Synthesis Example 37. The names of the synthesized compound and
15 the synthesis yields are indicated below.

[0636] Synthesis Example 38

4-[4-(4-Chlorophenyl)piperazin-1-yl]-1-{{3-(trifluoromethyl)-1H-pyrazol-1-yl}methyl}-1,3,5-triazin-2(1H)-one

20 Yield: 6.8%

[0637] Synthesis Example 39

4-[4-(m-Tolyl)piperazin-1-yl]-1-{{3-(trifluoromethyl)-1H-pyrazol-1-yl}methyl}-1,3,5-triazin-2(1H)-one

25 Yield: 4.5%

[0638] Synthesis Example 40

4-[4-(3-Cyclopropylphenyl)piperazin-1-yl]-1-{{3-(trifluoromethyl)-1H-pyrazol-1-yl}methyl}-1,3,5-triazin-2(1H)-one

Yield: 36%

[0639] Synthesis Example 41

1-{{[3-(Trifluoromethyl)-1H-pyrazol-1-yl]methyl}-4-{4-[3-(trifluoromethyl)phenyl]piperazin-1-yl}}-1,3,5-triazin-2(1H)-one

5

Yield: 36%

[0640] Synthesis Example 42

1-{{[3-(Trifluoromethyl)-1H-pyrazol-1-yl]methyl}-4-{4-[4-(trifluoromethyl)phenyl]piperazin-1-yl}}-1,3,5-triazin-2(1H)-one

10

Yield: 15%

[0641] Synthesis Example 43

1-{{[3-(Trifluoromethyl)-1H-pyrazol-1-yl]methyl}-4-(4-{4-[(trifluoromethyl)thio]phenyl}piperazin-1-yl))-1,3,5-triazin-2(1H)-one

15

Yield: 14%

[0642] Synthesis Example 44

4-[4-(p-tolyl)piperazin-1-yl]-1-{{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

20

Yield: 19%

[0643] Synthesis Example 45

4-{4-[4-(Trifluoromethoxy)phenyl]piperazin-1-yl}-1-{{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

25

Yield: 7.5%

[0644] Synthesis Example 46

4-[4-(4-Ethylphenyl)piperazin-1-yl]-1-{{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 5.6%

[0645] Synthesis Example 47

4-{4-[4-(1,1,2,2-Tetrafluoroethoxy)phenyl]piperazin-1-yl}-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 5.5%

[0646] Synthesis Example 48

4-[4-(4-Fluoro-3-methylphenyl)piperazin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 6.0%

[0647] Synthesis Example 49

4-[4-(4-Oxo-5-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)piperazin-1-yl]benzonitrile

Yield: 17%

[0648] Synthesis Example 50

Methyl

3-[4-(4-Oxo-5-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)piperazin-1-yl]benzoate

Yield: 12%

[0649] Synthesis Example 51

2-Fluoro-5-[4-(4-oxo-5-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)piperazin-1-yl]benzonitrile

Yield: 6.9%

[0650] Synthesis Example 52

4-{4-[4-Fluoro-3-(trifluoromethyl)phenyl]piperazin-1-yl}-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 12%

5 [0651] Synthesis Example 53

4-[4-(3-Chloro-4-fluorophenyl)piperazin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 9.2%

10 [0652] Synthesis Example 54

3-[4-(4-Oxo-5-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)piperazin-1-yl]benzonitrile

Yield: 17%

15 [0653] Synthesis Example 55

4-[4-(3-Chlorophenyl)piperazin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 16%

20 [0654] Synthesis Example 56

4-{4-[4-Chloro-3-(trifluoromethyl)phenyl]piperazin-1-yl}-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 17%

25 [0655] Synthesis Example 57

4-[4-(4-Fluoro-3-nitrophenyl)piperazin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 24%

[0656] Synthesis Example 58

1-{[3-(Trifluoromethyl)-1H-pyrazol-1-yl]methyl}-4-[4-(3,4,5-trifluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

5 Yield: 20%

[0657] Synthesis Example 59

4-[4-(o-Tolyl)piperazin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

10 Yield: 3.9%

[0658] Synthesis Example 60

4-[4-(4-Chloro-2-fluorophenyl)piperazin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

15 Yield: 7.0%

[0659] Synthesis Example 61

Ethyl

2-Chloro-5-[4-(4-oxo-5-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)piperazin-1-yl]benzoate

20

Yield: 15%

[0660] Synthesis Example 62

2-Chloro-5-[4-(4-oxo-5-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)piperazin-1-yl]benzonitrile

25

Yield: 12%

[0661] Synthesis Example 63

4-[4-(2-Methyl-4-nitrophenyl)piperazin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 18%

[0662] Synthesis Example 64

4-[4-(2,4-Difluorophenyl)piperazin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl
 5 }-1,3,5-triazin-2(1H)-one

The synthesis was performed using

4-[4-(2,4-difluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one (189 mg, 0.65 mmol)
 synthesized in Reference Synthesis Example 86 and
 10 [3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl 4-methylbenzenesulfonate (413 mg, 1.29
 mmol) synthesized in Reference Synthesis Example 15 in a similar manner to Synthesis
 Example 1 to obtain the title compound (21 mg, yield: 7.4%).

[0663] Synthesis Example 65

6-Butoxy-1-(4-chlorobenzyl)-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-on
 15 e

To a solution of

6-chloro-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one (50 mg, 0.16
 mmol) synthesized in Reference Synthesis Example 2 in a solvent mixture of
 20 N,N-dimethylformamide (1 mL) and 1,2-dimethoxyethane (1 mL), lithium hydride (1.3
 mg, 0.16 mmol) was added at 0°C and the resultant mixture was stirred at room
 temperature for 15 minutes. To the mixture, 4-chlorobenzyl bromide (40 mg, 0.19
 mmol) was added and the resultant mixture was stirred at room temperature for 30
 minutes and then at 60°C for 2 hours. Next, at room temperature, 1-butanol (30 µL,
 25 0.32 mmol) and lithium hydride (2.6 mg, 0.32 mmol) were added to the resultant mixture
 and the resultant mixture was stirred for 30 hours. After the completion of the reaction,
 water was added to the reaction solution and extraction with ethyl acetate from the
 resultant reaction solution was performed three times. The organic layer was washed
 with brine, was dried over anhydrous sodium sulfate, and was concentrated under

reduced pressure. The resultant residue was purified by amino-type thin-layer chromatography (hexane/ethyl acetate=1/1 (v/v)) to obtain the title compound (18 mg, yield: 21%).

[0664] Synthesis Example 66

5 1-{4-[(4-Chlorobenzyl)oxy]benzyl}-4-[4-(4-fluorophenyl)piperazin-1-yl]-6-methoxy-1,3,5-triazin-2(1H)-one

3-{4-[(4-Chlorobenzyl)oxy]benzyl}-6-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazine-2,4(1H,3H)-dione

10

In argon atmosphere, to an N,N-dimethylformamide solution (5 mL) of 6-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazine-2,4(1H,3H)-dione (200 mg, 0.69 mmol) synthesized in Reference Synthesis Example 18, lithium hydride (12 mg, 1.51 mmol) was added at 0°C and the resultant mixture was stirred for 1 hour. To the mixture, 1-chloro-4-{[4-(chloromethyl)phenoxy]methyl}benzene (183 mg, 0.69 mmol) was added and the resultant mixture was stirred at 50°C for 2 hours and at 70°C for 3 days. After the completion of the reaction, water and ethanol were added to the reaction solution and a resultant solid was filtered to obtain a crude product of a precursor (66St) of the title compound, which was used as it was in the next reaction below.

20

1-{4-[(4-chlorobenzyl)oxy]benzyl}-4-[4-(4-fluorophenyl)piperazin-1-yl]-6-methoxy-1,3,5-triazin-2(1H)-one

To an N,N-dimethylformamide solution (4 mL) of a crude product (0.69 mmol) of 3-{4-[(4-chlorobenzyl)oxy]benzyl}-6-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazine-2,4(1H,3H)-dione, potassium carbonate (332 mg, 2.40 mmol) and methyl iodide (137 μ L, 2.20 mmol) were added at room temperature and the resultant mixture was stirred at room temperature for 1 day. After the completion of the reaction, the reaction solution was concentrated under reduced pressure and the resultant residue was purified by silica

25

gel column chromatography to obtain the title compound (26 mg, two step yield: 7%).

[0665] Synthesis Example 67

1-[(2H-Indazol-2-yl)methyl]-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

5

To a dichloromethane solution of

{4-[4-(4-fluorophenyl)piperazin-1-yl]-2-oxo-1,3,5-triazin-1(2H)-yl}methyl

4-methylbenzenesulfonate (165 mg, 0.36 mmol) synthesized in Reference Synthesis

Example 4, 1,8-diazabicyclo[5.4.0] undeca-7-ene (65 μ L, 0.44 mmol) and 1H-indazole

10 (43 mg, 0.40 mmol) were added at room temperature and the resultant mixture was

stirred at room temperature for 30 minutes. After the completion of the reaction, the

reaction solution was concentrated under reduced pressure and the resultant residue was

purified by silica gel column chromatography (ethyl acetate) to obtain the title compound

(8.5 mg, yield: 5.2%).

15 [0666] Synthesis Example 68

4-[4-(4-Chlorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

To a dichloromethane solution (1.0 mL) of

20 4-[4-(4-chlorophenyl)-4-hydroxypiperidin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]

methyl}-1,3,5-triazin-2(1H)-one (14 mg, 0.03 mmol) synthesized in Reference Synthesis

Example 91a, trifluoroacetic acid (0.5 mL) was added and the resultant mixture was

stirred at room temperature for 3 hours. After the completion of the reaction, the

reaction solution was concentrated under reduced pressure and to the resultant

25 concentrate, water and saturated sodium hydrogen carbonate aqueous solution were

added. A deposited solid was washed with water and was dried under reduced pressure

to obtain the title compound (7.4 mg, yield: 55%).

[0667] Synthesis Example 69

4-(4-Phenyl-5,6-dihydropyridin-1(2H)-yl)-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]meth

yl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using

4-(4-hydroxy-4-phenylpiperidin-1-yl)-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one (7.4 mg, 0.02 mmol) synthesized in Reference Synthesis Example 91b in a similar manner to Synthesis Example 68 to obtain the title compound (6.9 mg, yield: 98%).

[0668] Synthesis Example 70

4-(4-Phenyl-5,6-dihydropyridin-1(2H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Synthesis Example 71

4-[4-(4-Chlorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using the mixture (21 mg) of

4-(4-hydroxy-4-phenylpiperidin-1-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one and 4-[4-(4-chlorophenyl)-4-hydroxypiperidin-1-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one that were synthesized in Reference Synthesis Example 91 in a similar manner to Synthesis Example 68. The resultant residue was purified by preparative high performance chromatography to obtain the title compound of Synthesis Example 70 (3.0 mg, yield: 16%) and the title compound of Synthesis Example 71 (7.5 mg, yield: 37%).

[0669] Synthesis Example 72

1-(4-Chlorobenzyl)-4-[4-(3,3-dimethylbutyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

To a dichloromethane solution (1 mL) of

1-(4-chlorobenzyl)-4-(piperazin-1-yl)-1,3,5-triazin-2(1H)-one (31 mg, 0.10 mmol)

synthesized in Reference Synthesis Example 9 and 3,3-dimethylbutanal (13 μ L, 0.10 mmol), sodium triacetoxymethylborohydride (42 mg, 0.20 mmol) was added and the resultant mixture was stirred at room temperature for 4 hours. After the completion of the reaction, saturated sodium bicarbonate water was added to the reaction solution and extraction with ethyl acetate from the resultant reaction solution was performed. The organic layer was concentrated under reduced pressure and to the resultant residue, ethyl acetate was added. A resultant solid was filtered and was dried under reduced pressure to obtain the title compound (11 mg, yield: 28%).

[0670] Synthesis Example 73

10 1-(4-Chlorobenzyl)-4-[4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

The synthesis was performed using

1-(4-chlorobenzyl)-4-(piperazin-1-yl)-1,3,5-triazin-2(1H)-one (31 mg, 0.10 mmol) synthesized in Reference Synthesis Example 9 and dihydro-2H-pyran-4(3H)-one (10 mg, 0.10 mmol) in a similar manner to Synthesis Example 72 to obtain the title compound (2.6 mg, yield: 6.6%).

[0671] Synthesis Example 74

20 1-(4-Chlorobenzyl)-4-[4-(4-fluorobenzoyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

To an N,N-dimethylformamide solution (1 mL) of

1-(4-chlorobenzyl)-4-(piperazin-1-yl)-1,3,5-triazin-2(1H)-one (31 mg, 0.10 mmol) synthesized in Reference Synthesis Example 9, triethylamine (28 μ L, 0.20 mmol) and 4-fluorobenzoyl chloride (18 μ L, 0.15 mmol) were added at room temperature and the resultant mixture was stirred at room temperature for 4 hours. After the completion of the reaction, saturated ammonium chloride aqueous solution was added to the reaction solution and extraction with ethyl acetate from the resultant reaction solution was performed. The organic layer was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography (hexane/ethyl

acetate=1/4→0/1 (v/v)) to obtain the title compound (27 mg, yield: 63%).

[0672] Synthesis Example 75

1,1,1-Trifluoro-2-methylpropan-2-yl

4-[5-(4-chlorobenzyl)-4-oxo-4,5-dihydro-1,3,5-triazin-2-yl]piperazine-1-carboxylate

5

The synthesis was performed using

1-(4-chlorobenzyl)-4-(piperazin-1-yl)-1,3,5-triazin-2(1H)-one (50 mg, 0.16 mmol)

synthesized in Reference Synthesis Example 9 and

3-methyl-1-{[(1,1,1-trifluoro-2-methylpropan-2-yl)oxy]carbonyl}-1H-imidazol-3-ium

10 iodide (89 mg, 0.25 mmol) synthesized in Reference Synthesis Example 48 in a similar manner to Synthesis Example 74 to obtain the title compound (42 mg, yield: 56%).

[0673] Synthesis Example 76

1-(4-Chlorobenzyl)-4-[4-(5-nitropyridin-2-yl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

15 An N,N-dimethylformamide suspension (1 mL) of

1-(4-chlorobenzyl)-4-(piperazin-1-yl)-1,3,5-triazin-2(1H)-one (31 mg, 0.10 mmol)

synthesized in Reference Synthesis Example 9, 2-chloro-5-nitropyridine (16 mg, 0.10 mmol) and potassium carbonate (18 mg, 0.20 mmol) was stirred at 100°C for 1 hour.

After the completion of the reaction, water was added to the reaction solution and

20 extraction with ethyl acetate from the resultant reaction solution was performed three

times. The organic layer was washed with brine, was dried over anhydrous sodium

sulfate, and was concentrated under reduced pressure. The resultant residue was

purified by preparative silica gel thin-layer chromatography to obtain the title compound (2.1 mg, yield: 4.9%).

25 [0674] Synthesis Example 77

1-(4-Chlorobenzyl)-4-{4-[5-(trifluoromethyl)pyridin-2-yl]piperazin-1-yl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using

1-(4-chlorobenzyl)-4-(piperazin-1-yl)-1,3,5-triazin-2(1H)-one (31 mg, 0.10 mmol) synthesized in Reference Synthesis Example 9 and 2-chloro-5-trifluoromethyl pyridine (18 mg, 0.10 mmol) in a similar manner to Synthesis Example 76 to obtain the title compound (1.7 mg, yield: 3.7%).

5 [0675] Synthesis Example 78

1-(4-Chlorobenzyl)-4-[4-(5-chlorothiophen-2-yl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

An N,N-dimethylaminoethanol suspension (1 mL) of 1-(4-chlorobenzyl)-4-(piperazin-1-yl)-1,3,5-triazin-2(1H)-one (30 mg, 0.10 mmol) synthesized in Reference Synthesis Example 9, 2-bromo-5-chlorothiophene (110 μ L, 0.10 mmol), copper (3.0 mg, 0.05 mmol), copper (I) iodide (9 mg, 0.05 mmol), and potassium phosphate (43 mg, 0.20 mmol) was stirred at 80°C for 3 days. After the completion of the reaction, water was added to the reaction solution and extraction with ether and ethyl acetate from the resultant reaction solution was performed. The organic layer was washed with brine, was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure. The resultant residue was purified by silica gel column chromatography (ethyl acetate/methanol) to obtain the title compound (0.5 mg, yield: 1.2%).

[0676] Synthesis Example 79

20 1-(4-Chlorobenzyl)-4-[[1-(4-fluorophenyl)piperidin-4-yl]amino]-1,3,5-triazin-2(1H)-one

To a chloroform solution (1.1 mL) of 4-chloro-1-(4-chlorobenzyl)-1,3,5-triazin-2(1H)-one (59 mg, 0.23 mmol) synthesized in Reference Synthesis Example 11, potassium carbonate (80 mg, 0.58 mmol) and 1-(4-fluorophenyl)piperidine-4-amine (66 mg, 0.34 mmol) synthesized in Reference Synthesis Example 83 were added and the resultant mixture was stirred at room temperature for 30 minutes. After the completion of the reaction, water was added to the reaction solution and extraction with chloroform from the resultant reaction solution was performed. The organic layer was concentrated under reduced pressure and the

resultant residue was purified by silica gel column chromatography (hexane/ethyl acetate=1/5→0/1 and next, ethyl acetate/methanol=19/1 (v/v)) to obtain the title compound (50 mg, yield: 52%).

- [0677] In Synthesis Examples 80 to 83, 85 to 97, and 100 to 106, each title
5 compound was synthesized in a similar manner to Synthesis Example 79. The names of the synthesized compounds and the synthesis yields are indicated below.

[0678] Synthesis Example 80

1-(4-Chlorobenzyl)-4-{4-[(4-fluorophenyl)amino]piperidin-1-yl}-1,3,5-triazin-2(1H)-one

- 10 Yield: 47%

[0679] Synthesis Example 81

1-(4-Chlorobenzyl)-4-{[1-(4-fluorophenyl)piperidin-4-yl][methylamino]}-1,3,5-triazin-2(1H)-one

- 15 Yield: 34%

[0680] Synthesis Example 82

(R)-1-(4-Chlorobenzyl)-4-{[1-(4-fluorophenyl)piperidin-3-yl]amino}-1,3,5-triazin-2(1H)-one

- 20 Yield: 47%

[0681] Synthesis Example 83

(S)-1-(4-Chlorobenzyl)-4-{[1-(4-fluorophenyl)piperidin-3-yl]amino}-1,3,5-triazin-2(1H)-one

- 25 Yield: 42%

[0682] Synthesis Example 84

1-(4-Chlorobenzyl)-4-[4-(5-fluoro-2-thioxo-2,3-dihydro-1H-benzo[d]imidazol-1-yl)piperidin-1-yl]-1,3,5-triazin-2(1H)-one

To an acetonitrile solution (1 mL) of
 1-(4-chlorobenzyl)-4-hydroxy-1,3,5-triazin-2(1H)-one (20 mg, 0.08 mmol) synthesized in
 Reference Synthesis Example 10, (benzotriazol-1-yloxy)tripyrrolidino-phosphonium (47
 mg, 0.10 mmol) and 1,8-diazabicyclo[5.4.0] undeca-7-ene (15 μ L, 0.10 mmol) were
 5 added at room temperature and the resultant mixture was stirred for 20 minutes. To the
 reaction solution, 5-fluoro-1H-benzo[d]imidazole-2(3H)-thione (25 mg, 0.10 mmol) was
 added and the resultant reaction solution was stirred at 80°C for 10 minutes. After the
 completion of the reaction, the reaction solution was concentrated under reduced pressure
 and the resultant residue was purified by silica gel column chromatography
 10 (chloroform/ethyl acetate=1/1 (v/v)) to obtain the title compound (0.8 mg, yield: 2.0%).

[0683] Synthesis Example 85

1-(4-Chlorobenzyl)-4-{[3-(4-fluorophenyl)cyclopentyl]amino}-1,3,5-triazin-2(1H)-one

Yield: 31%

15 [0684] Synthesis Example 86

(R)-1-(4-Chlorobenzyl)-4-([1-(4-fluorophenyl)pyrrolidin-3-yl]methyl)amino)-1,3,5-triazin-2(1H)-one

Yield: 3.4%

20 [0685] Synthesis Example 87

1-(4-Chlorobenzyl)-4-{3-[(4-fluorobenzyl)oxy]azetidin-1-yl}-1,3,5-triazin-2(1H)-one

Yield: 11%

[0686] Synthesis Example 88

25 1-(4-Chlorobenzyl)-4-{4-[3-(trifluoromethyl)-1H-pyrazol-1-yl]piperidin-1-yl}-1,3,5-triazin-2(1H)-one

Yield: 3.1%

[0687] Synthesis Example 89

1-(4-Chlorobenzyl)-4-[3-(4-fluorophenoxy)azetidin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 4.9%

[0688] Synthesis Example 90

5 1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 18%

[0689] Synthesis Example 91

10 1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-3,4-dihydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 36%

[0690] Synthesis Example 92

15 (R)-1-(4-Chlorobenzyl)-4-{[1-(4-fluorophenyl)pyrrolidin-3-yl]amino}-1,3,5-triazin-2(1H)-one

Yield: 37%

[0691] Synthesis Example 93

20 1-(4-Chlorobenzyl)-4-{[1-(4-fluorophenyl)pyrrolidin-3-yl][methyl]amino}-1,3,5-triazin-2(1H)-one

Yield: 38%

[0692] Synthesis Example 94

25 (3R,4R)-1-(4-Chlorobenzyl)-4-{[1-(4-fluorophenyl)-4-hydroxypyrrolidin-3-yl]amino}-1,3,5-triazin-2(1H)-one

Yield: 2.4%

[0693] Synthesis Example 95

(3R)-1-(4-Chlorobenzyl)-4-{[4-fluoro-1-(4-fluorophenyl)pyrrolidin-3-yl]amino}-1,3,5-triazin-2(1H)-one

Yield: 8.2%

5 [0694] Synthesis Example 96

(R)-1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-2-methylpiperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 19%

10 [0695] Synthesis Example 97

1-(4-Chlorobenzyl)-4-({2-[(4-fluorophenyl)amino]ethyl}amino)-1,3,5-triazin-2(1H)-one

Yield: 8.9%

[0696] Synthesis Example 98

15 4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using

4-[4-(4-fluorophenyl)-4-hydroxypiperidin-1-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one (76 mg, 0.17 mmol) synthesized in Reference Synthesis Example 89 in a similar manner to Synthesis Example 114 to obtain the title compound (70 mg, yield: 84%).

[0697] Synthesis Example 99

25 4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using

4-[4-(4-fluorophenyl)-4-hydroxypiperidin-1-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (20 mg, 0.04 mmol) synthesized in Reference Synthesis

Example 90 in a similar manner to Synthesis Example 114 to obtain the title compound (18 mg, yield: 97%).

[0698] Synthesis Example 100

1-(4-Chlorobenzyl)-4-{3-[(4-fluorophenyl)thio]azetidin-1-yl}-1,3,5-triazin-2(1H)-one

5

Yield: 28%

[0699] Synthesis Example 101

1-(4-Chlorobenzyl)-4-{3-[(4-fluorobenzyl)thio]azetidin-1-yl}-1,3,5-triazin-2(1H)-one

10 Yield: 54%

[0700] Synthesis Example 102

(S)-1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-3-methylpiperazin-1-yl]-1,3,5-triazin-2(1H)-one

15 Yield: 7.5%

[0701] Synthesis Example 103

1-(4-Chlorobenzyl)-4-[2-(4-fluorobenzyl)morpholino]-1,3,5-triazin-2(1H)-one

Yield: 0.6%

20 [0702] Synthesis Example 104

1-(4-Chlorobenzyl)-4-{[1-(4-fluorophenyl)-2-oxopyrrolidin-3-yl]amino}-1,3,5-triazin-2(1H)-one

Yield: 4.0%

25 [0703] Synthesis Example 105

(R)-1-(4-Chlorobenzyl)-4-{[1-(4-fluorobenzyl)pyrrolidin-3-yl]amino}-1,3,5-triazin-2(1H)-one

Yield: 5.0%

[0704] Synthesis Example 106

(S)-1-(4-Chlorobenzyl)-4-[[1-(4-fluorobenzyl)pyrrolidin-3-yl]amino]-1,3,5-triazin-2(1H)-one

5 Yield: 9.0%

[0705] Synthesis Example 107

1-(4-Chlorobenzyl)-4-[5-fluoro-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

10 To a dichloromethane solution (1.0 mL) of
1-(4-chlorobenzyl)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one (15 mg, 0.04 mmol) synthesized in Synthesis Example 90,
(diethylamino)sulfur trifluoride (10 μ L, 0.04 mmol) was added at 0°C and the resultant mixture was stirred at room temperature for 1.5 hours. After the completion of the
15 reaction, dichloromethane was added to the reaction solution and the resultant reaction solution was washed with saturated sodium bicarbonate water and brine. The resultant organic layer was dried over anhydrous sodium sulfate and was concentrated under reduced pressure. To the resultant residue, silica gel was added and the mixture was filtered, followed by washing the mixture with ethyl acetate. The filtrate was
20 concentrated under reduced pressure to obtain the title compound (7 mg, yield: 46%).

[0706] Synthesis Example 108

1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-3-hydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one

25 A solution of
1-(4-chlorobenzyl)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one (10 mg, 0.03 mmol) synthesized in Reference Synthesis Example 90 and palladium-activated carbon (3 mg) in a solvent mixture of methanol (5 mL) and dichloromethane (5 mL) was stirred in a hydrogen atmosphere for 3 days. After the

completion of the reaction, the reaction solution was filtered through Celite and the filtrate was concentrated under reduced pressure to obtain the title compound (8.9 mg, yield: 90%).

[0707] Synthesis Example 109

5 Methyl

1-[5-(4-chlorobenzyl)-4-oxo-4,5-dihydro-1,3,5-triazin-2-yl]-4-(4-fluorophenyl)piperazine-2-carboxylate

The synthesis was performed using methyl

10 1-[5-(4-chlorobenzyl)-4-oxo-4,5-dihydro-1,3,5-triazin-2-yl]piperazine-2-carboxylate (26 mg, 0.07 mmol) synthesized in Reference Synthesis Example 88 in a similar manner to Synthesis Example 37 to obtain the title compound (12 mg, yield: 37%).

[0708] Synthesis Example 110

1-(4-Chlorobenzyl)-4-[[1-(4-fluorophenyl)pyrrolidin-3-yl]oxy]-1,3,5-triazin-2(1H)-one

15

The synthesis was performed using a chloroform solution (2 mL) of 4-chloro-1-(4-chlorobenzyl)-1,3,5-triazin-2(1H)-one (67 mg, 0.26 mmol) synthesized in Reference Synthesis Example 11 and 1-(4-fluorophenyl)pyrrolidin-3-ol (71 mg, 0.39 mmol) synthesized in Reference Synthesis Example 46 in a similar manner to Synthesis Example 79 to obtain the title compound (4.0 mg, yield: 4%).

[0709] Synthesis Example 111

1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-4-hydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one

25

The synthesis was performed using 4-[4-(4-fluorophenyl)-4-hydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one (273 mg, 0.94 mmol) synthesized in Reference Synthesis Example 14 in a similar manner to Synthesis Example 1 to obtain the title compound (83 mg, yield: 21%).

[0710] Synthesis Example 112

1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

To

- 5 1-(4-chlorobenzyl)-4-[4-(4-fluorophenyl)-4-hydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one (240 mg, 0.58 mmol) synthesized in Synthesis Example 111, trifluoroacetic acid (2.3 mL) was added and the resultant mixture was stirred at room temperature for 1 hour. After the completion of the reaction, the reaction solution was concentrated under reduced pressure and the resultant residue was purified by silica gel (amino-type) column chromatography (hexane/ethyl acetate=1/1) to obtain the title compound (44 mg, yield: 19%).

[0711] Synthesis Example 113

1-(4-Chlorobenzyl)-4-[4-fluoro-4-(4-fluorophenyl)piperidin-1-yl]-1,3,5-triazin-2(1H)-one

- 15 The synthesis was performed using 1-(4-chlorobenzyl)-4-[4-(4-fluorophenyl)-4-hydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one (62 mg, 0.15 mmol) synthesized in Synthesis Example 111 in a similar manner to Synthesis Example 107 to obtain the title compound (35 mg, yield: 57%).

[0712] Synthesis Example 114

- 20 1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)piperidin-1-yl]-1,3,5-triazin-2(1H)-one

A methanol (1 mL) solution of

- 1-(4-chlorobenzyl)-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one (30 mg, 0.08 mmol) synthesized in Synthesis Example 112 and
25 palladium-activated carbon (3 mg) was stirred in a hydrogen atmosphere at room temperature for 1 day and at 80°C for 7 hours. To the resultant mixture, chloroform was added and the resultant mixture was stirred further for 1 day. After the completion of the reaction, the reaction solution was filtered through Celite and the filtrate was concentrated under reduced pressure to obtain the title compound (28 mg, yield: 92%).

[0713] Synthesis Example 115

1-(4-Chlorobenzyl)-4-[4-(4-chlorobenzyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

The filtrate obtained in Reference Synthesis Example 9 was concentrated under
 5 reduced pressure and the resultant residue was purified by silica gel (amino-type) column chromatography (hexane/ethyl acetate) to obtain the title compound.

[0714] Synthesis Example 116

(1R*,3S*)-1-(4-Chlorobenzyl)-4-{[3-(4-fluorophenyl)cyclopentyl]amino}-1,3,5-triazin-2(1H)-one

10

A mixture of stereoisomers of
 1-(4-chlorobenzyl)-4-{[3-(4-fluorophenyl)cyclopentyl]amino}-1,3,5-triazin-2(1H)-one
 synthesized in Synthesis Example 85 was subjected to separation by preparative high
 performance liquid chromatography (gradient: hexane/ethanol 80/20→71/29→50/50,
 15 Chiralpak IA) and a fraction containing a single diastereomer that was eluted at a
 retention time of 7.00 minutes was concentrated to obtain the title compound.

[0715] Synthesis Example 117

(1S*,3S*)-1-(4-Chlorobenzyl)-4-{[3-(4-fluorophenyl)cyclopentyl]amino}-1,3,5-triazin-2(1H)-one

20

The separation of stereoisomers was performed in the same manner as in Synthesis
 Example 116 and a fraction containing a single diastereomer that was eluted at a retention
 time of 7.72 minutes was concentrated to obtain the title compound.

[0716] Synthesis Example 118

(1R*,3R*)-1-(4-Chlorobenzyl)-4-{[3-(4-fluorophenyl)cyclopentyl]amino}-1,3,5-triazin-2(1H)-one

25

The separation of stereoisomers was performed in the same manner as in Synthesis
 Example 116 and a fraction containing a single diastereomer that was eluted at a retention

time of 8.75 minutes was concentrated to obtain the title compound.

[0717] Synthesis Example 119

(1S*,3R*)-1-(4-Chlorobenzyl)-4-{[3-(4-fluorophenyl)cyclopentyl]amino}-1,3,5-triazin-2(1H)-one

5

The separation of stereoisomers was performed in the same manner as in Synthesis Example 116 and a fraction containing a single diastereomer that was eluted at a retention time of 9.30 minutes was concentrated to obtain the title compound.

[0718] Synthesis Example 120

10 1-(4-Chlorobenzyl)-4-{4-[5-(trifluoromethyl)thiophen-2-yl]piperazin-1-yl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using 2-bromo-5-(trifluoromethyl)thiophene in a similar manner to Synthesis Example 78 (yield: 0.2%).

15 [0719] In Synthesis Examples 121 to 139, each title compound was synthesized in a similar manner to Synthesis Example 79. The names of the synthesized compounds and the synthesis yields are indicated below.

[0720] Synthesis Example 121

20 1-(4-Chlorobenzyl)-4-{[3-(4-fluorophenyl)cyclopent-2-en-1-yl]amino}-1,3,5-triazin-2(1H)-one

Yield: 3%

[0721] Synthesis Example 122

25 1-(4-Chlorobenzyl)-4-{[5-(4-fluorophenyl)-1-methylpyrrolidin-3-yl]amino}-1,3,5-triazin-2(1H)-one

Yield: 10%

[0722] Synthesis Example 123

1-(4-Chlorobenzyl)-4-[3-fluoro-4-(4-fluorophenyl)piperidin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 22%

[0723] Synthesis Example 124

1-(4-Chlorobenzyl)-4-{[1-(4-fluorobenzyl)azetidin-3-yl]amino}-1,3,5-triazin-2(1H)-one

5

Yield: 17%

[0724] Synthesis Example 125

1-(4-Chlorobenzyl)-4-[(4'-fluoro-[1,1'-biphenyl]-4-yl)amino]-1,3,5-triazin-2(1H)-one

10 Yield: 41%

[0725] Synthesis Example 126

1-(4-Chlorobenzyl)-4-[(4'-fluoro-[1,1'-biphenyl]-3-yl)amino]-1,3,5-triazin-2(1H)-one

Yield: 26%

15 [0726] Synthesis Example 127

1-(4-Chlorobenzyl)-4-{[5-(4-fluorophenyl)bicyclo[3.1.0]hexan-2-yl]amino}-1,3,5-triazin-2(1H)-one

Yield: 10%

20 [0727] Synthesis Example 128

(S)-1-(4-Chlorobenzyl)-4-([1-(4-fluorophenyl)pyrrolidin-2-yl]methyl)amino)-1,3,5-triazin-2(1H)-one

Yield: 13%

25 [0728] Synthesis Example 129

(R)-1-(4-Chlorobenzyl)-4-([1-(4-fluorophenyl)pyrrolidin-2-yl]methyl)amino)-1,3,5-triazin-2(1H)-one

Yield: 11%

[0729] Synthesis Example 130

1-(4-Chlorobenzyl)-4-{[4-(3,4-difluorophenyl)thiazol-2-yl]amino}-1,3,5-triazin-2(1H)-one

5 Yield: 2%

[0730] Synthesis Example 131

1-(4-Chlorobenzyl)-4-{[1-(4-fluorophenyl)-1H-pyrrol-3-yl]amino}-1,3,5-triazin-2(1H)-one

10 Yield: 1%

[0731] Synthesis Example 132

1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-3-methyl-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-5-methyl-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

The synthesis was performed using the mixture of

4-(4-fluorophenyl)-3-methyl-1,2,3,6-tetrahydropyridine hydrochloride and

4-(4-fluorophenyl)-5-methyl-1,2,3,6-tetrahydropyridine hydrochloride that was

20 synthesized in Reference Synthesis Example 284 to obtain a mixture of

1-(4-chlorobenzyl)-4-[4-(4-fluorophenyl)-3-methyl-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one and

1-(4-chlorobenzyl)-4-[4-(4-fluorophenyl)-5-methyl-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one that are the title compounds.

25 Yield: 26%

[0732] Synthesis Example 133

1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-2-methyl-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-6-methyl-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

riazin-2(1H)-one

The synthesis was performed using the mixture of
 4-(4-fluorophenyl)-2-methyl-1,2,3,6-tetrahydropyridine trifluoroacetate and
 4-(4-fluorophenyl)-6-methyl-1,2,3,6-tetrahydropyridine trifluoroacetate that was
 synthesized in Reference Synthesis Example 280 to obtain a mixture of
 1-(4-chlorobenzyl)-4-[4-(4-fluorophenyl)-2-methyl-5,6-dihydropyridin-1(2H)-yl]-1,3,5-tri-
 iazin-2(1H)-one and
 1-(4-chlorobenzyl)-4-[4-(4-fluorophenyl)-6-methyl-5,6-dihydropyridin-1(2H)-yl]-1,3,5-tri-
 iazin-2(1H)-one that are the title compounds.

Yield: 40%

[0733] Synthesis Example 134

1-(4-Chlorobenzyl)-4-[4-fluoro-4-(4-fluorophenyl)-2-methylpiperidin-1-yl]-1,3,5-triazin-
 2(1H)-one

Yield: 19%

[0734] Synthesis Example 135

1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-4-hydroxy-3-methylpiperidin-1-yl]-1,3,5-triazin-
 2(1H)-one

Yield: 5%

[0735] Synthesis Example 136

1-(4-Chlorobenzyl)-4-{[1-(4-fluorophenyl)-3-methylpyrrolidin-3-yl]amino}-1,3,5-triazin-
 2(1H)-one

Yield: 5%

[0736] Synthesis Example 137

1-(4-Chlorobenzyl)-4-{[1-(4-fluorophenyl)-4-methylpyrrolidin-3-yl]amino}-1,3,5-triazin-
 2(1H)-one

Yield: 3%

[0737] Synthesis Example 138

1-(4-Chlorobenzyl)-4-[4-(2,4-difluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

5

Yield: 33%

[0738] Synthesis Example 139

1-(4-Chlorobenzyl)-4-{[1-(4-fluorophenyl)-1H-pyrazol-4-yl]amino}-1,3,5-triazin-2(1H)-one

10

Yield: 22%

[0739] In Synthesis Examples 140 to 142, each title compound was synthesized in a similar manner to Synthesis Example 1. The names of the synthesized compounds and the synthesis yields are indicated below.

15 [0740] Synthesis Example 140

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-[4-(hexyloxy)benzyl]-1,3,5-triazin-2(1H)-one

Yield: 30%

[0741] Synthesis Example 141

20 1-(4-Chloro-3-methoxybenzyl)-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 55%

[0742] Synthesis Example 142

25 1-[4-(Cyclopropylmethoxy)benzyl]-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 42%

[0743] In Synthesis Examples 143 to 158, each title compound was synthesized in

a similar manner to Synthesis Example 11. The names of the synthesized compounds and the synthesis yields are indicated below.

[0744] Synthesis Example 143

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[5-(trifluoromethyl)pyridin-2-yl]methyl}-1,3,5-t
5 riazin-2(1H)-one

Yield: 50%

[0745] Synthesis Example 144

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]methyl}-1
10 ,3,5-triazin-2(1H)-one

Yield: 27%

[0746] Synthesis Example 145

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[5-(2,2,2-trifluoroethoxy)pyrazin-2-yl]methyl}-
15 1,3,5-triazin-2(1H)-one

Yield: 20%

[0747] Synthesis Example 146

5-({4-[4-(4-Fluorophenyl)piperazin-1-yl]-2-oxo-1,3,5-triazin-1(2H)-yl}methyl)thiophene
20 -2-carbonitrile

Yield: 31%

[0748] Synthesis Example 147

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-[3-(2,2,2-trifluoroethoxy)benzyl]-1,3,5-triazin-2(
25 1H)-one

Yield: 50%

[0749] Synthesis Example 148

1-[(5-Bromothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1

H)-one

Yield: 26%

[0750] Synthesis Example 149

5 1-[(4,5-Dibromothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 30%

[0751] Synthesis Example 150

10 4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[1-methyl-5-(trifluoromethyl)-1H-pyrazol-3-yl]-methyl}-1,3,5-triazin-2(1H)-one

Yield: 49%

[0752] Synthesis Example 151

15 1-[(3,5-Dibromothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 38%

[0753] Synthesis Example 152

20 1-[(4-Bromothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 23%

[0754] Synthesis Example 153

25 1-[(5-Bromo-4-methylthiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 13%

[0755] Synthesis Example 154

1-{[4-Bromo-5-(trifluoromethyl)thiophen-2-yl]methyl}-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 11%

5 [0756] Synthesis Example 155

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-[(5-methylthiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 10%

10 [0757] Synthesis Example 156

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-{[3-(perfluoroethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 30%

15 [0758] Synthesis Example 157

4-[4-(4-Fluorophenyl)piperazin-1-yl]-1-[(5-fluorothiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 27%

20 [0759] Synthesis Example 158

1-{[5-(Difluoromethyl)thiophen-2-yl]methyl}-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 28%

25 [0760] In Synthesis Examples 159 to 187, each title compound was synthesized using 4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 342 in a similar manner to Synthesis Example 11. The names of the synthesized compounds and the synthesis yields are indicated below.

[0761] Synthesis Example 159

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)furan-2-yl]methyl}-1,3,5-triazin-2(1H)-one

5 Yield: 56%

[0762] Synthesis Example 160

1-[(4-Bromothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

10 Yield: 56%

[0763] Synthesis Example 161

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[2-(trifluoromethyl)pyrimidin-5-yl]methyl}-1,3,5-triazin-2(1H)-one

15 Yield: 60%

[0764] Synthesis Example 162

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[6-(2,2,2-trifluoroethoxy)pyridazin-3-yl]methyl}-1,3,5-triazin-2(1H)-one

20 Yield: 99%

[0765] Synthesis Example 163

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[2-(2,2,2-trifluoroethoxy)thiazol-5-yl]methyl}-1,3,5-triazin-2(1H)-one

25 Yield: 68%

[0766] Synthesis Example 164

1-{[5-(Difluoromethyl)thiophen-2-yl]methyl}-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 57%

[0767] Synthesis Example 165

1-{[4-Bromo-5-(trifluoromethyl)thiophen-2-yl]methyl}-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

5

Yield: 12%

[0768] Synthesis Example 166

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[6-(hexyloxy)pyridin-3-yl]methyl}-1,3,5-triazin-2(1H)-one

10

Yield: 13%

[0769] Synthesis Example 167

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{(5-methylthiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

15

Yield: 22%

[0770] Synthesis Example 168

1-{[6-(Difluoromethoxy)pyridin-3-yl]methyl}-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

20

Yield: 29%

[0771] Synthesis Example 169

5-({4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-2-oxo-1,3,5-triazin-1(2H)-yl}methyl)pyridin-2-yl trifluoromethanesulfonate

25

Yield: 42%

[0772] Synthesis Example 170

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)furan-3-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 39%

[0773] Synthesis Example 171

1-[(5-Chlorothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1
5 ,3,5-triazin-2(1H)-one

Yield: 41%

[0774] Synthesis Example 172

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[6-(2,2,2-trifluoroethoxy)pyridin
10 -3-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 30%

[0775] Synthesis Example 173

1-[(5-Chlorobenzofuran-2-yl)methyl]-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]
15]-1,3,5-triazin-2(1H)-one

Yield: 35%

[0776] Synthesis Example 174

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[3-(trifluoromethyl)-1,2,4-oxadia
20 zol-5-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 29%

[0777] Synthesis Example 175

1-[(5-Bromothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1
25 ,3,5-triazin-2(1H)-one

Yield: 1%

[0778] Synthesis Example 176

1-[(2,2-Dimethylchroman-6-yl)methyl]-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-

yl]-1,3,5-triazin-2(1H)-one

Yield: 72%

[0779] Synthesis Example 177

5 1-[(2,3-Dihydrobenzofuran-5-yl)methyl]-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 86%

[0780] Synthesis Example 178

10 1-(Chroman-6-ylmethyl)-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 9%

[0781] Synthesis Example 179

15 4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(2,2,2-trifluoroethoxy)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 2%

[0782] Synthesis Example 180

20 4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-[(5-fluorothiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 47%

[0783] Synthesis Example 181

25 1-[(2,3-Dihydrobenzo[b][1,4]dioxin-6-yl)methyl]-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 63%

[0784] Synthesis Example 182

1-(Benzo[d][1,3]dioxol-5-ylmethyl)-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 53%

5 [0785] Synthesis Example 183

1-[4-(tert-Butyl)benzyl]-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 92%

10 [0786] Synthesis Example 184

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[2-(trifluoromethyl)benzofuran-5-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 76%

15 [0787] Synthesis Example 185

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 44%

20 [0788] Synthesis Example 186

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[3-(perfluoroethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 29%

25 [0789] Synthesis Example 187

1-[(5-Acetylthiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 2%

[0790] In Synthesis Example 188 and Synthesis Example 189, each title compound was synthesized using

4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 336 in a similar manner to Synthesis

5 Example 1. The names of the synthesized compounds and the synthesis yields are indicated below.

[0791] Synthesis Example 188

4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

10

Yield: 55%

[0792] Synthesis Example 189

5-({4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-2-oxo-1,3,5-triazin-1(2H)-yl}methyl)pyridin-2-yl trifluoromethanesulfonate

15

Yield: 47%

[0793] In Synthesis Examples 190 to 202, each title compound was synthesized using

4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 336 in a similar manner to Synthesis

20

Example 11. The names of the synthesized compounds and the synthesis yields are indicated below.

[0794] Synthesis Example 190

4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25

Yield: 28%

[0795] Synthesis Example 191

1-[4-(tert-Butyl)benzyl]-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1

,3,5-triazin-2(1H)-one

Yield: 62%

[0796] Synthesis Example 192

- 5 1-{{6-(Difluoromethoxy)pyridin-3-yl}methyl}-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 77%

[0797] Synthesis Example 193

- 10 4-{{4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl}}-1-{{5-(trifluoromethyl)furan-2-yl}methyl}-1,3,5-triazin-2(1H)-one

Yield: 31%

[0798] Synthesis Example 194

- 15 4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{{3-(perfluoroethyl)-1H-pyrazol-1-yl}methyl}-1,3,5-triazin-2(1H)-one

Yield: 32%

[0799] Synthesis Example 195

- 20 4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{{2-(trifluoromethyl)-2,3-dihydrobenzofuran-5-yl}methyl}-1,3,5-triazin-2(1H)-one

Yield: 28%

[0800] Synthesis Example 196

- 25 4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{{5-(tetrahydrofuran-2-yl)thiophen-2-yl}methyl}-1,3,5-triazin-2(1H)-one

Yield: 20%

[0801] Synthesis Example 197

1-[(5-Chlorothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 57%

5 [0802] Synthesis Example 198

4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-[(5-fluorothiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 12%

10 [0803] Synthesis Example 199

1-[(5-Bromothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 50%

15 [0804] Synthesis Example 200

1-{[5-(tert-Butyl)thiophen-2-yl]methyl}-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 9%

20 [0805] Synthesis Example 201

4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 30%

25 [0806] Synthesis Example 202

4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{1-[5-(trifluoromethyl)thiophen-2-yl]ethyl}-1,3,5-triazin-2(1H)-one

Yield: 21%

[0807] In Synthesis Example 203 and Synthesis Example 205, each title compound was synthesized using

(R)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 333a in a similar manner to Synthesis Example 1. The names of the synthesized compounds and the synthesis yields are indicated below.

[0808] Synthesis Example 203

(R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-1,3,5-triazin-2(1H)-one

10

Yield: 3%

[0809] Synthesis Example 204

(S)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-1,3,5-triazin-2(1H)-one

15

The synthesis was performed using

(S)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 333b in a similar manner to Synthesis Example 1.

20 Yield: 4%

[0810] Synthesis Example 205

(R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

25 Yield: 27%

[0811] In Synthesis Examples 206 to 212, each title compound was synthesized using

(R)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 333a in a similar manner to Synthesis

Example 11. The names of the synthesized compounds and the synthesis yields are indicated below.

[0812] Synthesis Example 206

(R)-1-([3-(1,1-Difluoroethyl)-1H-pyrazol-1-yl]methyl)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 10%

[0813] Synthesis Example 207

(R)-1-([5-(1,1-Difluoroethyl)thiophen-2-yl]methyl)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 70%

[0814] Synthesis Example 208

(R)-5-([4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-2-oxo-1,3,5-triazin-1(2H)-yl]methyl)pyridin-2-yl trifluoromethanesulfonate

Yield: 47%

[0815] Synthesis Example 209

(R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-([5-(tetrahydrofuran-2-yl)thiophen-2-yl]methyl)-1,3,5-triazin-2(1H)-one

Yield: 7%

[0816] Synthesis Example 210

(R)-1-([5-(tert-Butyl)thiophen-2-yl]methyl)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 16%

[0817] Synthesis Example 211

(R)-4-([4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-2-oxo-1,3,5-triazin-1(2H)-yl]methyl)pyridin-2-yl trifluoromethanesulfonate

in-1(2H)-yl)methyl)phenyl trifluoromethanesulfonate

Yield: 50%

[0818] Synthesis Example 212

5 (R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-({6-[(2,2,2-trifluoroethoxy)methyl]pyridin-3-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 57%

[0819] Synthesis Example 213

10 (R)-4-({4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-2-oxo-1,3,5-triazin-1(2H)-yl)methyl}phenyl trifluoromethanesulfonate

The synthesis was performed using

(R)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 334 and
15 4-(chloromethyl)phenyl trifluoromethanesulfonate synthesized in Reference Synthesis Example 127 in a similar manner to Synthesis Example 1 (yield: 22%).

[0820] In Synthesis Examples 214 to 219, each title compound was synthesized using

20 (R)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 334 in a similar manner to Synthesis Example 11. The names of the synthesized compounds and the synthesis yields are indicated below.

[0821] Synthesis Example 214

25 (R)-1-{[6-(Difluoromethoxy)pyridin-3-yl)methyl]-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 55%

[0822] Synthesis Example 215

(R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[3-(perfluoroethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 16%

5 [0823] Synthesis Example 216

(R)-5-({4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-2-oxo-1,3,5-triazin-1(2H)-yl}methyl)pyridin-2-yl trifluoromethanesulfonate

Yield: 44%

10 [0824] Synthesis Example 217

(R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-({6-[(2,2,2-trifluoroethoxy)methyl]pyridin-3-yl}methyl)-1,3,5-triazin-2(1H)-one

Yield: 31%

15 [0825] Synthesis Example 218

(R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 11%

20 [0826] Synthesis Example 219

(R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[2-(trifluoromethyl)thiazol-5-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 6%

25 [0827] Synthesis Example 220

4-[4-(4-Chlorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)furan-2-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using

4-[4-(4-chlorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 367 and 2-(bromomethyl)-5-(trifluoromethyl)furan in a similar manner to Synthesis Example 1 (yield: 33%).

[0828] Synthesis Example 221

5 4-[4-(4-fluorophenyl)piperazin-1-yl]-6-methyl-1-([2-(trifluoromethyl)thiazol-5-yl]methyl)-1,3,5-triazin-2(1H)-one

The synthesis was performed using

10 4-[4-(4-fluorophenyl)piperazin-1-yl]-6-methyl-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 340 and 5-(chloromethyl)-2-(trifluoromethyl)thiazole in a similar manner to Synthesis Example 11 (yield: 22%).

[0829] Synthesis Example 222

15 1-({3-[1-(4-fluorophenyl)cyclopropyl]-1,2,4-oxadiazol-5-yl}methyl)-4-[4-(4-fluorophenyl)piperazin-1-yl]-6-methyl-1,3,5-triazin-2(1H)-one

The synthesis was performed using

20 4-[4-(4-fluorophenyl)piperazin-1-yl]-6-methyl-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 340 and 5-(chloromethyl)-3-[1-(4-fluorophenyl)cyclopropyl]-1,2,4-oxadiazole in a similar manner to Synthesis Example 11 (yield: 36%).

[0830] Synthesis Example 223

6-Methyl-1-([3-(perfluoroethyl)-1H-pyrazol-1-yl]methyl)-4-[4-(p-tolyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

25 The synthesis was performed using

6-methyl-4-[4-(p-tolyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 339 and 1-(chloromethyl)-3-(perfluoroethyl)-1H-pyrazole in a similar manner to Synthesis Example 11 (yield: 25%).

[0831] Synthesis Example 224

(R)-4-{{[1-(4-fluorophenyl)pyrrolidin-3-yl]amino}-1-}[[5-(trifluoromethyl)thiophen-3-yl]methyl]-1,3,5-triazin-2(1H)-one

The synthesis was performed using

- 5 (R)-4-{{[1-(4-fluorophenyl)pyrrolidin-3-yl]amino}-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 338 and 2-(bromomethyl)-5-(trifluoromethyl)thiophene in a similar manner to Synthesis Example 1 (yield: 24%).

[0832] Synthesis Example 225

- 10 4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-}[[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-1,3,5-triazin-2(1H)-one

The synthesis was performed using

- 4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 343 and
 15 [3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl 4-methylbenzenesulfonate synthesized in Reference Synthesis Example 57 in a similar manner to Synthesis Example 25 (yield: 23%).

[0833] In Synthesis Example 226 and Synthesis Example 227, each title compound was synthesized using

- 20 1-(4-chlorobenzyl)-4-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 347 in a similar manner to Reference Synthesis Example 139. The names of the synthesized compounds and the synthesis yields are indicated below.

[0834] Synthesis Example 226

- 25 1-(4-Chlorobenzyl)-1-[4-(thiophen-3-yl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 30%

[0835] Synthesis Example 227

1-(4-Chlorobenzyl)-4-(6-fluoro-5',6'-dihydro-[3,4'-bipyridin]-1'(2'H)-yl)-1,3,5-triazin-2(1H)-one

Yield: 54%

- 5 [0836] In Synthesis Examples 228 to 302, each title compound was synthesized using

4-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 346 in a similar manner to Reference Synthesis Example 139. The
10 names of the synthesized compounds and the synthesis yields are indicated below.

 [0837] Synthesis Example 228

4-[4-(3-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

- 15 Yield: 80%

 [0838] Synthesis Example 229

4-[4-(4-Fluoro-2-methoxyphenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

- 20 Yield: 64%

 [0839] Synthesis Example 230

4-[4-(Thiophen-3-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

- 25 Yield: 50%

 [0840] Synthesis Example 231

4-[4-(2-Chloropyrimidin-5-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 68%

[0841] Synthesis Example 232

4-(5,6-Dihydro-[4,4'-bipyridin]-1(2H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

5

Yield: 76%

[0842] Synthesis Example 233

4-[4-(5-Chlorothiophen-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10

Yield: 26%

[0843] Synthesis Example 234

4-[4-(2-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

15

Yield: 44%

[0844] Synthesis Example 235

4-[4-(5-Chloropyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

20

Yield: 61%

[0845] Synthesis Example 236

4-(4-{4-[(Trifluoromethyl)thio]phenyl}-5,6-dihydropyridin-1(2H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25

Yield: 64%

[0846] Synthesis Example 237

4-[4-(m-Tolyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 74%

[0847] Synthesis Example 238

Ethyl

- 5 3-[1-(4-Oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)-1,2,3,6-tetrahydropyridin-4-yl]benzoate

Yield: 39%

[0848] Synthesis Example 239

- 10 4-{4-[4-(Trifluoromethyl)phenyl]-5,6-dihydropyridin-1(2H)-yl}-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 82%

[0849] Synthesis Example 240

- 15 4-(6-Methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 98%

[0850] Synthesis Example 241

- 20 4-(6-Fluoro-5',6'-dihydro-[3,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 64%

[0851] Synthesis Example 242

- 25 4-[4-(4-Fluoro-2-methylphenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 89%

[0852] Synthesis Example 243

4-[4-(3-Cyclopropylphenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 96%

5 [0853] Synthesis Example 244

4-(5',6'-Dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 80%

10 [0854] Synthesis Example 245

4-(5',6'-Dihydro-[3,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 67%

15 [0855] Synthesis Example 246

4-[4-(3-Fluoro-4-methylphenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 63%

20 [0856] Synthesis Example 247

4-[4-(3-Chloro-4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 61%

25 [0857] Synthesis Example 248

4-[4-(4-Morpholinophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 42%

[0858] Synthesis Example 249

4-[5-(Trifluoromethyl)-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

5 Yield: 45%

[0859] Synthesis Example 250

4-[4-(Thiazol-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10 Yield: 28%

[0860] Synthesis Example 251

4-[4-(4-Fluoro-3-methoxyphenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

15 Yield: 94%

[0861] Synthesis Example 252

4-(6-Methoxy-5',6'-dihydro-[3,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

20 Yield: 93%

[0862] Synthesis Example 253

4-[4-(3-Chloro-2-methylphenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25 Yield: 69%

[0863] Synthesis Example 254

4-[4-(3-Chloro-5-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 30%

[0864] Synthesis Example 255

4-{4-[4-Fluoro-3-(2,2,2-trifluoroethoxy)phenyl]-5,6-dihydropyridin-1(2H)-yl}-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

5

Yield: 64%

[0865] Synthesis Example 256

4-(5-Chloro-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10

Yield: 11%

[0866] Synthesis Example 257

4-[6-(2,2,2-Trifluoroethoxy)-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

15

Yield: 66%

[0867] Synthesis Example 258

4-(6-Chloro-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

20

Yield: 57%

[0868] Synthesis Example 259

4-{4-[1-(2,2,2-Trifluoroethyl)-1H-pyrazol-4-yl]-5,6-dihydropyridin-1(2H)-yl}-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25

Yield: 29%

[0869] Synthesis Example 260

4-[4-(Benzofuran-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 43%

[0870] Synthesis Example 261

4-(5-Fluoro-5',6'-dihydro-[3,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]
5 l)methyl}-1,3,5-triazin-2(1H)-one

Yield: 38%

[0871] Synthesis Example 262

4-(2'-Fluoro-5,6-dihydro-[4,4'-bipyridin]-1(2H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]
10 methyl}-1,3,5-triazin-2(1H)-one

Yield: 34%

[0872] Synthesis Example 263

4-(5-Fluoro-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]
15 l)methyl}-1,3,5-triazin-2(1H)-one

Yield: 36%

[0873] Synthesis Example 264

4-[4-(3,4-Difluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-
20 2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 57%

[0874] Synthesis Example 265

4-[4-(Thiazol-5-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]m
25 ethyl}-1,3,5-triazin-2(1H)-one

Yield: 17%

[0875] Synthesis Example 266

4-{4-[3-Chloro-5-(trifluoromethyl)phenyl]-5,6-dihydropyridin-1(2H)-yl}-1-{[5-(trifluoro

methyl)thiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 28%

[0876] Synthesis Example 267

- 5 4-{4-[4-Fluoro-3-(trifluoromethyl)phenyl]-5,6-dihydropyridin-1(2H)-yl}-1-{[5-(trifluoro
methyl)thiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 96%

[0877] Synthesis Example 268

- 10 4-(4-Fluoro-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-y
l)methyl}-1,3,5-triazin-2(1H)-one

Yield: 48%

[0878] Synthesis Example 269

- 15 4-[4-(1-Oxo-2,3-dihydro-1H-inden-5-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluorom
ethyl)thiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 55%

[0879] Synthesis Example 270

- 20 4-(6-Fluoro-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-y
l)methyl}-1,3,5-triazin-2(1H)-one

Yield: 37%

[0880] Synthesis Example 271

- 25 4-(4-Methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2
-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 20%

[0881] Synthesis Example 272

4-[4-(6-Methoxypyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 49%

5 [0882] Synthesis Example 273

4-(5-Chloro-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 70%

10 [0883] Synthesis Example 274

4-[6-(Trifluoromethyl)-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 33%

15 [0884] Synthesis Example 275

4-[4-(3-Methylisothiazol-5-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 82%

20 [0885] Synthesis Example 276

4-[4-(4-Fluoro-3-methylphenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 77%

25 [0886] Synthesis Example 277

4-[4-(5-Fluoro-2-methylphenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 80%

[0887] Synthesis Example 278

2-Fluoro-5-[1-(4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)-1,2,3,6-tetrahydropyridin-4-yl]benzonitrile

5 Yield: 72%

[0888] Synthesis Example 279

4-[4-(3-Methoxyphenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10 Yield: 77%

[0889] Synthesis Example 280

4-(3-Chloro-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

15 Yield: 36%

[0890] Synthesis Example 281

4-[4-(5-Methylthiophen-3-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

20 Yield: 15%

[0891] Synthesis Example 282

4-[4-(5-Methylthiazol-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25 Yield: 39%

[0892] Synthesis Example 283

4-[4-(6,7-Dihydro-5H-pyrrolo[1,2-a]imidazol-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 80%

[0893] Synthesis Example 284

4-(6-Methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

5

Yield: 86%

[0894] Synthesis Example 285

4-[4-(3,5-Difluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10

Yield: 79%

[0895] Synthesis Example 286

4-(5-Methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

15

Yield: 17%

[0896] Synthesis Example 287

4-[4-(3-Fluoro-4-methoxyphenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]ethyl}-1,3,5-triazin-2(1H)-one

20

Yield: 66%

[0897] Synthesis Example 288

4-[6-Methoxy-5-(prop-1-en-2-yl)-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25

Yield: 63%

[0898] Synthesis Example 289

4-(2'-Methoxy-5,6-dihydro-[4,4'-bipyridin]-1(2H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 68%

[0899] Synthesis Example 290

4-(5-Methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 28%

[0900] Synthesis Example 291

4-{4-[4-(Difluoromethyl)phenyl]-5,6-dihydropyridin-1(2H)-yl}-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 57%

[0901] Synthesis Example 292

2-Fluoro-4-[1-(4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)-1,2,3,6-tetrahydropyridin-4-yl]benzonitrile

Yield: 6%

[0902] Synthesis Example 293

4-[6-(Difluoromethoxy)-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 74%

[0903] Synthesis Example 294

4-[4-(p-Tolyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 52%

[0904] Synthesis Example 295

4-[4-(Benzo[d]oxazol-5-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

n-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 35%

[0905] Synthesis Example 296

5 1'-(4-Oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)-1',2',3',6'-tetrahydro-[2,4'-bipyridine]-5-carbonitrile

Yield: 45%

[0906] Synthesis Example 297

10 6-Chloro-1'-(4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)-1',2',3',6'-tetrahydro-[2,4'-bipyridine]-5-carbonitrile

Yield: 33%

[0907] Synthesis Example 298

15 1'-(4-Oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)-1',2',3',6'-tetrahydro-[2,4'-bipyridine]-6-carbonitrile

Yield: 36%

[0908] Synthesis Example 299

20 4-(5-Fluoro-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2-(1H)-one

Yield: 39%

[0909] Synthesis Example 300

25 4-[4-(5-Methoxythiophen-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 46%

[0910] Synthesis Example 301

1'-(4-Oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)-1',2',3',6'-tetrahydro-[2,4'-bipyridine]-6-carbaldehyde

Yield: 89%

5 [0911] Synthesis Example 302

5-Fluoro-2-[1-(4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)-1,2,3,6-tetrahydropyridin-4-yl]benzaldehyde

Yield: 80%

10 [0912] Synthesis Example 303

4-[4-(3-Fluoro-4-hydroxyphenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using a crude product obtained by synthesis using
15 4-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 346 and 4-bromo-2-fluorophenyl acetate in a similar manner to Reference Synthesis Example 139, in a similar manner to Reference Synthesis Example 2 (two step yield 41%).

20 [0913] Synthesis Example 304

4-[4-(4-Fluoro-3-hydroxyphenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using a crude product obtained by synthesis using
25 4-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 346 and 5-bromo-2-fluorophenyl acetate in a similar manner to Reference Synthesis Example 139, in a similar manner to Reference Synthesis Example 2 (two step yield 35%).

[0914] Synthesis Example 305

4-{4-[4-Fluoro-2-(hydroxymethyl)phenyl]-5,6-dihydropyridin-1(2H)-yl}-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

- 5 The synthesis was performed using a crude product obtained by synthesis using 4-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 346 and [(2-bromo-5-fluorobenzyl)oxy](tert-butyl)dimethylsilane synthesized in Reference Synthesis Example 128 in a similar manner to Reference
- 10 Synthesis Example 139, in a similar manner to Reference Synthesis Example 272 (two step yield 40%).

[0915] In Synthesis Examples 306 to 338, each title compound was synthesized using

- 4-(methylthio)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 291 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.
- 15

[0916] Synthesis Example 306

- 4-{[rac-(2S,4S)-2-(4-Fluorophenyl)tetrahydro-2H-pyran-4-yl]amino}-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one
- 20

Yield: 62%

[0917] Synthesis Example 307

- 4-{[rac-(2S,4R)-2-(4-Fluorophenyl)tetrahydro-2H-pyran-4-yl]amino}-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one
- 25

Yield: 79%

[0918] Synthesis Example 308

4-{[rac-(3R,5S)-5-(4-Fluorophenyl)tetrahydrofuran-3-yl]amino}-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

hiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 63%

[0919] Synthesis Example 309

5 4-{[rac-(3S,5S)-5-(4-Fluorophenyl)tetrahydrofuran-3-yl]amino}-1-{[5-(trifluoromethyl)t
hiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 73%

[0920] Synthesis Example 310

10 4-([rac-(1S,2S)-2-(4-Fluorophenyl)cyclopropyl)methyl]amino)-1-{[5-(trifluoromethyl)t
hiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 56%

[0921] Synthesis Example 311

15 4-[6-(4-Fluorophenyl)-3-azabicyclo[3.1.0]hexan-3-yl]-1-{[5-(trifluoromethyl)thiophen-2-
yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 80%

[0922] Synthesis Example 312

20 tert-Butyl
trans-3-(4-fluorophenyl)-4-{[(4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl)methyl}-4,5-di
hydro-1,3,5-triazin-2-yl)amino]methyl}pyrrolidine-1-carboxylate

Yield: 89%

25 [0923] Synthesis Example 313

4-[5-(4-Fluorophenyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl]-1-{[5-(trifluoromethyl)thi
ophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 58%

[0924] Synthesis Example 314

N-[4-(4-Fluorophenyl)-1-(4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)-1,2,3,6-tetrahydropyridin-3-yl]methanesulfonamide

5 Yield: 12%

[0925] Synthesis Example 315

(R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10 Yield: 31%

[0926] Synthesis Example 316

(S)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

15 Yield: 20%

[0927] Synthesis Example 317

N-[4-(4-Fluorophenyl)-1-(4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)-1,2,3,6-tetrahydropyridin-3-yl]acetamide

20 Yield: 35%

[0928] Synthesis Example 318

cis-4-{[3-(4-Fluorophenyl)cyclobutyl]amino}-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25 Yield: 58%

[0929] Synthesis Example 319

trans-4-{[3-(4-Fluorophenyl)cyclobutyl]amino}-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

in a quantitative yield

[0930] Synthesis Example 320

trans-1-(4-Fluorophenyl)-3-[(4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)amino]cyclobutanecarbonitrile

5

Yield: 43%

[0931] Synthesis Example 321

4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10

Yield: 42%

[0932] Synthesis Example 322

4-(5'-Hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

15

Yield: 20%

[0933] Synthesis Example 323

(S)-4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

20

Yield: 15%

[0934] Synthesis Example 324

(R)-4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25

Yield: 32%

[0935] Synthesis Example 325

(R)-4-[4-(3-Fluoro-4-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 76%

[0936] Synthesis Example 326

(R)-4-[4-(3-Chloro-4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 76%

[0937] Synthesis Example 327

(R)-4-[5-Hydroxy-4-(4-methylthiophen-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 50%

[0938] Synthesis Example 328

(R)-4-(5'-Hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 31%

[0939] Synthesis Example 329

(R)-4-(5-Chloro-5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 15%

[0940] Synthesis Example 330

(R)-4-(6-Chloro-5'-hydroxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 19%

[0941] Synthesis Example 331

(R)-4-[4-(4-Fluoro-2-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

uoromethyl)thiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 51%

[0942] Synthesis Example 332

5 (R)-4-(5'-Hydroxy-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 73%

[0943] Synthesis Example 333

10 (R)-4-[5-Hydroxy-4-(5-methylthiophen-3-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 37%

[0944] Synthesis Example 334

15 4-[5-Amino-4-(4-fluorophenyl)-5-methyl-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 40%

[0945] Synthesis Example 335

20 (R)-4-(5-Fluoro-5'-hydroxy-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 77%

[0946] Synthesis Example 336

25 4-[4-(4-Fluorophenyl)-5-(hydroxymethyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl)methyl}-1,3,5-triazin-2(1H)-one

Yield: 60%

[0947] Synthesis Example 337

4-[[[(1r,3r)-3-(4-Fluorophenyl)-3-(hydroxymethyl)cyclobutyl]amino]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 73%

5 [0948] Synthesis Example 338

4-[(1R,5S,9s)-9-(4-Fluorophenyl)-9-hydroxy-3-oxa-7-azabicyclo[3.3.1]nonan-7-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 88%

10 [0949] Synthesis Example 339

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-(trifluoromethyl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using

4-(methylthio)-6-(trifluoromethyl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 372 and
15 4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine in a similar manner to Reference Synthesis Example 348 (yield: 40%).

[0950] In Synthesis Examples 340 to 383, each title compound was synthesized using

20 6-methyl-4-(methylthio)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 292 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

[0951] Synthesis Example 340

25 4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 31%

[0952] Synthesis Example 341

4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 93%

5 [0953] Synthesis Example 342

(R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 88%

10 [0954] Synthesis Example 343

(S)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 75%

15 [0955] Synthesis Example 344

4-[4-(4-Fluorophenyl)piperazin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 94%

20 [0956] Synthesis Example 345

4-[4-(6-Methoxypyridin-2-yl)piperazin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 95%

25 [0957] Synthesis Example 346

4-[4-(5-Chloro-6-methoxypyridin-2-yl)piperazin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 90%

[0958] Synthesis Example 347

(S)-4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{{5-(trifluoromethyl)thiophen-2-yl}methyl}-1,3,5-triazin-2(1H)-one

5 Yield: 66%

[0959] Synthesis Example 348

(R)-4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{{5-(trifluoromethyl)thiophen-2-yl}methyl}-1,3,5-triazin-2(1H)-one

10 Yield: 76%

[0960] Synthesis Example 349

(R)-4-[4-(3-Fluoro-4-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{{5-(trifluoromethyl)thiophen-2-yl}methyl}-1,3,5-triazin-2(1H)-one

15 Yield: 87%

[0961] Synthesis Example 350

(R)-4-[4-(3-Chloro-4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{{5-(trifluoromethyl)thiophen-2-yl}methyl}-1,3,5-triazin-2(1H)-one

20 Yield: 71%

[0962] Synthesis Example 351

4-[4-(3-Fluoro-4-methylphenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{{5-(trifluoromethyl)thiophen-2-yl}methyl}-1,3,5-triazin-2(1H)-one

25 Yield: 67%

[0963] Synthesis Example 352

4-[4-(3-Chloro-4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{{5-(trifluoromethyl)thiophen-2-yl}methyl}-1,3,5-triazin-2(1H)-one

Yield: 64%

[0964] Synthesis Example 353

4-(6-Methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

5

Yield: 87%

[0965] Synthesis Example 354

4-(5-Chloro-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10

Yield: 62%

[0966] Synthesis Example 355

6-Methyl-4-[4-(4-methylthiophen-2-yl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

15

Yield: 73%

[0967] Synthesis Example 356

(R)-4-(5'-Hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

20

Yield: 37%

[0968] Synthesis Example 357

6-Methyl-4-[4-(5-methylthiophen-3-yl)piperazin-1-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25

Yield: 88%

[0969] Synthesis Example 358

4-[4-(3-Fluoro-4-methylphenyl)piperazin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 94%

[0970] Synthesis Example 359

4-[4-(3-Chloro-4-fluorophenyl)piperazin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 89%

[0971] Synthesis Example 360

cis-4-{[3-(4-Fluorophenyl)cyclobutyl]amino}-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 94%

[0972] Synthesis Example 361a

(R)-4-(5-Chloro-5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

[0973] Synthesis Example 361b

4-[(3'R)-5-Chloro-3'-hydroxy-6-methoxy-3',4'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

20 The synthesis was performed using

(R)-5-chloro-6-methoxy-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-ol synthesized in Reference Synthesis Example 144 to obtain the title compound (yield: 30%) of Synthesis Example 361a and the title compound (yield: 48%) of Synthesis Example 361b.

[0974] Synthesis Example 362

25 (R)-4-[5-Hydroxy-4-(4-methylthiophen-2-yl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 99%

[0975] Synthesis Example 363

6-Methyl-4-[4-(p-tolyl)piperazin-1-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 89%

5 [0976] Synthesis Example 364

6-Methyl-4-[4-(6-methylpyridin-2-yl)piperazin-1-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 99%

10 [0977] Synthesis Example 365

4-[4-(5-Fluoro-6-methylpyridin-2-yl)piperazin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 92%

15 [0978] Synthesis Example 366

4-[4-(6-Chloropyridin-2-yl)piperazin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 90%

20 [0979] Synthesis Example 367

4-[4-(3,5-Difluorophenyl)piperazin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 94%

25 [0980] Synthesis Example 368

6-Methyl-4-[4-(5-methylpyridin-2-yl)piperazin-1-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 95%

[0981] Synthesis Example 369

(R)-4-[4-(4-Fluoro-2-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

5 Yield: 56%

[0982] Synthesis Example 370

4-[2-(4-Fluorophenyl)-2,8-diazaspiro[4.5]decan-8-yl]-6-methyl-1-[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10 Yield: 95%

[0983] Synthesis Example 371

(R)-4-(5'-Hydroxy-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-6-methyl-1-[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

15 Yield: 79%

[0984] Synthesis Example 372

(R)-4-[5-Hydroxy-4-(5-methylthiophen-3-yl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

20 Yield: 59%

[0985] Synthesis Example 373

4-{4-[4-Fluoro-2-(hydroxymethyl)phenyl]-5,6-dihydropyridin-1(2H)-yl}-6-methyl-1-[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25 Yield: 25%

[0986] Synthesis Example 374

4-{4-[4-Fluoro-2-(hydroxymethyl)phenyl]piperazin-1-yl}-6-methyl-1-[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 80%

[0987] Synthesis Example 375

(R)-4-(5-Fluoro-5'-hydroxy-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

5

Yield: 56%

[0988] Synthesis Example 376

4-[6,6-Difluoro-4-(4-fluorophenyl)-1,4-diazepan-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10

Yield: 85%

[0989] Synthesis Example 377

4-(6,6-Difluoro-4-phenyl-1,4-diazepan-1-yl)-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

15

Yield: 99%

[0990] Synthesis Example 378

4-[4-(6-Methyl-4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)piperazin-1-yl]benzonitrile

20

Yield: 69%

[0991] Synthesis Example 379

4-[4-(4-Fluorophenyl)-5-(hydroxymethyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25

Yield: 60%

[0992] Synthesis Example 380

4-[8-(4-Fluorophenyl)-3,8-diazabicyclo[3.2.1]octan-3-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 40%

[0993] Synthesis Example 381

4-[4-(4-Fluorophenyl)-4,7-diazaspiro[2.5]octan-7-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 49%

[0994] Synthesis Example 382

4-[4-(Adamantan-1-yl)piperazin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 86%

[0995] Synthesis Example 383

4-{[4-(3,4-Difluorophenyl)thiazol-2-yl]amino}-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 25%

[0996] In Synthesis Examples 384 to 389, each title compound was synthesized using 1-[(5-bromothiophen-2-yl)methyl]-4-(methylthio)-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 296 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

[0997] Synthesis Example 384

(R)-4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-[(5-bromothiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 58%

[0998] Synthesis Example 385

(R)-1-[(5-Bromothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyri

din-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 86%

[0999] Synthesis Example 386

- 5 (R)-1-[(5-Bromothiophen-2-yl)methyl]-4-[4-(3-fluoro-4-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 36%

[1000] Synthesis Example 387

- 10 (R)-1-[(5-Bromothiophen-2-yl)methyl]-4-[4-(3-chloro-4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Yield: 93%

[1001] Synthesis Example 388

- 15 (R)-1-[(5-Bromothiophen-2-yl)methyl]-4-(5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1' (2'H)-yl)-1,3,5-triazin-2(1H)-one

Yield: 59%

[1002] Synthesis Example 389

- 20 (R)-1-[(5-Bromothiophen-2-yl)methyl]-4-(5-chloro-5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1' (2'H)-yl)-1,3,5-triazin-2(1H)-one

Yield: 13%

[1003] In Synthesis Examples 390 to 396, each title compound was synthesized

- 25 using

1-[(5-bromothiophen-2-yl)methyl]-6-methyl-4-(methylthio)-1,3,5-triazin-2(1H)-one

synthesized in Reference Synthesis Example 297 in a similar manner to Reference

Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

[1004] Synthesis Example 390

(R)-4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-[(5-bromothiophen-2-yl)methyl]-6-methyl-1,3,5-triazin-2(1H)-one

5 Yield: 63%

[1005] Synthesis Example 391

(R)-1-[(5-Bromothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

10 Yield: 96%

[1006] Synthesis Example 392

(R)-1-[(5-Bromothiophen-2-yl)methyl]-4-[4-(3-fluoro-4-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

15 Yield: 95%

[1007] Synthesis Example 393

(R)-1-[(5-Bromothiophen-2-yl)methyl]-4-[4-(3-chloro-4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

20 Yield: 99%

[1008] Synthesis Example 394

(R)-1-[(5-Bromothiophen-2-yl)methyl]-4-(5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1' (2'H)-yl)-6-methyl-1,3,5-triazin-2(1H)-one

25 Yield: 22%

[1009] Synthesis Example 395

(R)-1-[(5-Bromothiophen-2-yl)methyl]-4-(5-chloro-5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1' (2'H)-yl)-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 13%

[1010] Synthesis Example 396

(R)-1-[(5-Bromothiophen-2-yl)methyl]-4-[5-hydroxy-4-(5-methylthiophen-3-yl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

5

in a quantitative yield

[1011] In Synthesis Examples 397 to 402, each title compound was synthesized using 1-[(5-chlorothiophen-2-yl)methyl]-4-(methylthio)-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 299 in a similar manner to Reference

10 Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

[1012] Synthesis Example 397

(R)-1-[(5-Chlorothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

15

Yield: 98%

[1013] Synthesis Example 398

(R)-4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-[(5-chlorothiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

20

Yield: 73%

[1014] Synthesis Example 399

(R)-1-[(5-Chlorothiophen-2-yl)methyl]-4-[4-(3-fluoro-4-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

25

Yield: 34%

[1015] Synthesis Example 400

(R)-4-[4-(3-Chloro-4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-[(5-chlorothiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 62%

[1016] Synthesis Example 401

(R)-1-[(5-Chlorothiophen-2-yl)methyl]-4-(5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1' (2'H)-yl)-1,3,5-triazin-2(1H)-one

Yield: 41%

[1017] Synthesis Example 402

(R)-4-(5-Chloro-5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-[(5-chlorothiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 22%

[1018] In Synthesis Examples 403 to 410, each title compound was synthesized using

1-[(5-chlorothiophen-2-yl)methyl]-6-methyl-4-(methylthio)-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 300 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

[1019] Synthesis Example 403

(R)-1-[(5-Chlorothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

in a quantitative yield

[1020] Synthesis Example 404

(R)-1-[(5-Chlorothiophen-2-yl)methyl]-4-[4-(3-fluoro-4-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 85%

[1021] Synthesis Example 405

(R)-4-[4-(3-Chloro-4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-[(5-chlorothiophen-2-yl)methyl]-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 70%

5 [1022] Synthesis Example 406

(R)-1-[(5-Chlorothiophen-2-yl)methyl]-4-(5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1' (2'H)-yl)-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 36%

10 [1023] Synthesis Example 407

(R)-4-(5-Chloro-5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-[(5-chlorothiophen-2-yl)methyl]-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 13%

15 [1024] Synthesis Example 408

(R)-1-[(5-Chlorothiophen-2-yl)methyl]-4-[5-hydroxy-4-(4-methylthiophen-2-yl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 88%

20 [1025] Synthesis Example 409

(R)-4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-[(5-chlorothiophen-2-yl)methyl]-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 77%

25 [1026] Synthesis Example 410

(S)-1-[(5-Chlorothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 93%

[1027] In Synthesis Examples 411 to 415, each title compound was synthesized using 1-([5-(tert-butyl)thiophen-2-yl]methyl)-4-(methylthio)-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 306 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

[1028] Synthesis Example 411

(R)-1-([5-(tert-Butyl)thiophen-2-yl]methyl)-4-[4-(3-fluoro-4-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

10 Yield: 88%

[1029] Synthesis Example 412

(R)-1-([5-(tert-Butyl)thiophen-2-yl]methyl)-4-[4-(3-chloro-4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

15 Yield: 96%

[1030] Synthesis Example 413

(R)-1-([5-(tert-Butyl)thiophen-2-yl]methyl)-4-(5-chloro-5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1' (2'H)-yl)-1,3,5-triazin-2(1H)-one

20 Yield: 69%

[1031] Synthesis Example 414

(R)-1-([5-(tert-Butyl)thiophen-2-yl]methyl)-4-[4-(4-fluoro-2-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

25 Yield: 90%

[1032] Synthesis Example 415

(R)-1-([5-(tert-Butyl)thiophen-2-yl]methyl)-4-(5-fluoro-5'-hydroxy-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1' (2'H)-yl)-1,3,5-triazin-2(1H)-one

Yield: 91%

[1033] In Synthesis Examples 416 to 426, each title compound was synthesized using

1-{[5-(tert-Butyl)thiophen-2-yl]methyl}-6-methyl-4-(methylthio)-1,3,5-triazin-2(1H)-one
 5 synthesized in Reference Synthesis Example 305 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

[1034] Synthesis Example 416

(R)-1-{[5-(tert-Butyl)thiophen-2-yl]methyl}-4-[4-(3-fluoro-4-methylphenyl)-5-hydroxy-5
 10 ,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 82%

[1035] Synthesis Example 417

(R)-1-{[5-(tert-Butyl)thiophen-2-yl]methyl}-4-[4-(3-chloro-4-fluorophenyl)-5-hydroxy-5
 15 ,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 97%

[1036] Synthesis Example 418

(R)-1-{[5-(tert-Butyl)thiophen-2-yl]methyl}-4-(5-chloro-5'-hydroxy-6-methoxy-5',6'-dih
 20 ydro-[2,4'-bipyridin]-1'(2'H)-yl)-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 64%

[1037] Synthesis Example 419

1-{[5-(tert-Butyl)thiophen-2-yl]methyl}-4-[4-(3-fluoro-4-methylphenyl)piperazin-1-yl]-6
 25 -methyl-1,3,5-triazin-2(1H)-one

Yield: 84%

[1038] Synthesis Example 420

1-{[5-(tert-Butyl)thiophen-2-yl]methyl}-4-[4-(3-chloro-4-fluorophenyl)piperazin-1-yl]-6-

methyl-1,3,5-triazin-2(1H)-one

Yield: 93%

[1039] Synthesis Example 421

5 1- {[5-(tert-Butyl)thiophen-2-yl]methyl} -6-methyl-4-[4-(5-methylthiophen-3-yl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

Yield: 74%

[1040] Synthesis Example 422

10 (R)-1- {[5-(tert-Butyl)thiophen-2-yl]methyl} -4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 97%

[1041] Synthesis Example 423

15 (R)-1- {[5-(tert-Butyl)thiophen-2-yl]methyl} -4-[4-(4-fluoro-2-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 81%

[1042] Synthesis Example 424

20 (R)-1- {[5-(tert-Butyl)thiophen-2-yl]methyl} -4-(5'-hydroxy-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1' (2'H)-yl)-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 67%

[1043] Synthesis Example 425

25 (R)-1- {[5-(tert-Butyl)thiophen-2-yl]methyl} -4-[5-hydroxy-4-(5-methylthiophen-3-yl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 46%

[1044] Synthesis Example 426

(R)-1-{[5-(tert-Butyl)thiophen-2-yl]methyl}-4-(5-fluoro-5'-hydroxy-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-6-methyl-1,3,5-triazin-2(1H)-one

Yield: 49%

5 [1045] In Synthesis Examples 427 to 431, each title compound was synthesized using

4-(methylthio)-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 322 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis

10 yields are indicated below.

 [1046] Synthesis Example 427

(R)-4-[4-(3-Fluoro-4-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

15 Yield: 90%

 [1047] Synthesis Example 428

(R)-4-[4-(3-Chloro-4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

20 Yield: 88%

 [1048] Synthesis Example 429a

(R)-4-(5-Chloro-5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

 [1049] Synthesis Example 429b

25 4-[(3'R)-5-Chloro-3'-hydroxy-6-methoxy-3',4'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl]-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

The synthesis was performed using

(R)-5-chloro-6-methoxy-1',2',3',6'-tetrahydro-[2,4'-bipyridin]-3'-ol synthesized in

Reference Synthesis Example 144 to obtain the title compound (yield: 45%) of Synthesis Example 429a and the title compound (yield: 29%) of Synthesis Example 429b.

[1050] Synthesis Example 430

(R)-4-(5'-Hydroxy-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 91%

[1051] Synthesis Example 431

(R)-4-(5-Fluoro-5'-hydroxy-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 76%

[1052] In Synthesis Example 432 and Synthesis Example 433, each title compound was synthesized using

6-methyl-4-(methylthio)-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 323 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

[1053] Synthesis Example 432

(R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 80%

[1054] Synthesis Example 433

(R)-4-(5-Fluoro-5'-hydroxy-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-6-methyl-1-[(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]-1,3,5-triazin-2(1H)-one

Yield: 44%

[1055] In Synthesis Examples 434 to 438, each title compound was synthesized

using 4-(methylthio)-1-{[5-(trifluoromethyl)furan-2-yl]methyl}-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 303 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

5 [1056] Synthesis Example 434

4-[4-(3-Fluoro-4-methylphenyl)piperazin-1-yl]-1-{[5-(trifluoromethyl)furan-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 46%

10 [1057] Synthesis Example 435

4-[4-(3-Chloro-4-fluorophenyl)piperazin-1-yl]-1-{[5-(trifluoromethyl)furan-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 71%

15 [1058] Synthesis Example 436

4-[4-(5-Methylthiophen-3-yl)piperazin-1-yl]-1-{[5-(trifluoromethyl)furan-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 92%

20 [1059] Synthesis Example 437

4-[4-(5-Chloro-6-methoxypyridin-2-yl)piperazin-1-yl]-1-{[5-(trifluoromethyl)furan-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 100%

25 [1060] Synthesis Example 438

(R)-4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)furan-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 69%

[1061] In Synthesis Examples 439 to 444, each title compound was synthesized using

6-methyl-4-(methylthio)-1- $\{[5-(\text{trifluoromethyl})\text{furan-2-yl}]\text{methyl}\}$ -1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 302 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

[1062] Synthesis Example 439

(R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1- $\{[5-(\text{trifluoromethyl})\text{furan-2-yl}]\text{methyl}\}$ -1,3,5-triazin-2(1H)-one

10

Yield: 48%

[1063] Synthesis Example 440

(R)-4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1- $\{[5-(\text{trifluoromethyl})\text{furan-2-yl}]\text{methyl}\}$ -1,3,5-triazin-2(1H)-one

15

Yield: 72%

[1064] Synthesis Example 441

4-[4-(6-Methoxypyridin-2-yl)piperazin-1-yl]-6-methyl-1- $\{[5-(\text{trifluoromethyl})\text{furan-2-yl}]\text{methyl}\}$ -1,3,5-triazin-2(1H)-one

20

Yield: 62%

[1065] Synthesis Example 442

4-[4-(5-Chloro-6-methoxypyridin-2-yl)piperazin-1-yl]-6-methyl-1- $\{[5-(\text{trifluoromethyl})\text{furan-2-yl}]\text{methyl}\}$ -1,3,5-triazin-2(1H)-one

25

Yield: 71%

[1066] Synthesis Example 443

4-[4-(3-Fluoro-4-methylphenyl)piperazin-1-yl]-6-methyl-1- $\{[5-(\text{trifluoromethyl})\text{furan-2-yl}]\text{methyl}\}$ -1,3,5-triazin-2(1H)-one

Yield: 96%

[1067] Synthesis Example 444

6-Methyl-4-[4-(5-methylthiophen-3-yl)piperazin-1-yl]-1-{[5-(trifluoromethyl)furan-2-yl]
5 methyl}-1,3,5-triazin-2(1H)-one

in a quantitative yield

[1068] In Synthesis Examples 445 to 447, each title compound was synthesized using 5-{[4-(methylthio)-2-oxo-1,3,5-triazin-1(2H)-yl]methyl}pyridin-2-yl

10 trifluoromethanesulfonate synthesized in Reference Synthesis Example 311 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

[1069] Synthesis Example 445

(R)-5-({4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-2-oxo-1,3,5-triazin
15 -1(2H)-yl}methyl)pyridin-2-yl trifluoromethanesulfonate

Yield: 16%

[1070] Synthesis Example 446

(R)-5-({4-[4-(3-Fluoro-4-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-2-oxo-
20 1,3,5-triazin-1(2H)-yl}methyl)pyridin-2-yl trifluoromethanesulfonate

Yield: 14%

[1071] Synthesis Example 447

(R)-5-({4-[4-(3-Chloro-4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-2-oxo-1
25 ,3,5-triazin-1(2H)-yl}methyl)pyridin-2-yl trifluoromethanesulfonate

Yield: 71%

[1072] In Synthesis Examples 448 to 451, each title compound was synthesized using 5-{[6-methyl-4-(methylthio)-2-oxo-1,3,5-triazin-1(2H)-yl]methyl}pyridin-2-yl

trifluoromethanesulfonate synthesized in Reference Synthesis Example 310 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

[1073] Synthesis Example 448

- 5 (R)-5-({4-[4-(3-Fluoro-4-methylphenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-2-oxo-1,3,5-triazin-1(2H)-yl}methyl)pyridin-2-yl trifluoromethanesulfonate

Yield: 71%

[1074] Synthesis Example 449

- 10 (R)-5-({4-[4-(3-Chloro-4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-2-oxo-1,3,5-triazin-1(2H)-yl}methyl)pyridin-2-yl trifluoromethanesulfonate

Yield: 74%

[1075] Synthesis Example 450

- 15 (R)-5-{{4-[5-Chloro-5'-hydroxy-6-methoxy-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl]-6-methyl-2-oxo-1,3,5-triazin-1(2H)-yl]methyl}pyridin-2-yl trifluoromethanesulfonate

Yield: 22%

[1076] Synthesis Example 451

- 20 (R)-5-({4-[5-Amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-2-oxo-1,3,5-triazin-1(2H)-yl}methyl)pyridin-2-yl trifluoromethanesulfonate

Yield: 19%

[1077] Synthesis Example 452

- 25 4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methoxy-1-{{5-(trifluoromethyl)thiophen-2-yl}methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using
6-methoxy-4-(methylthio)-1-{{5-(trifluoromethyl)thiophen-2-yl}methyl}-1,3,5-triazin-2(1

H)-one synthesized in Reference Synthesis Example 331 and 4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine in a similar manner to Reference Synthesis Example 348 (yield: 38%).

[1078] Synthesis Example 453

5 (R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{1-[5-(trifluoromethyl)thiophen-2-yl]ethyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using

6-methyl-4-(methylthio)-1-{1-[5-(trifluoromethyl)thiophen-2-yl]ethyl}-1,3,5-triazin-2(1

10 H)-one synthesized in Reference Synthesis Example 313 and

(R)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol synthesized in Reference Synthesis Example 133 in a similar manner to Reference Synthesis Example 348 (yield: 64%).

[1079] Synthesis Example 454

15 (R)-4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{5-(perfluoroethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using

6-methyl-4-(methylthio)-1-{5-(perfluoroethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 320 and

20 (R)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol synthesized in Reference Synthesis Example 133 in a similar manner to Synthesis Example 348 (yield: 73%).

[1080] Synthesis Example 455

(R)-{1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-oxo-1,6-dihydro-1,3,5-triazin-2-yl}methyl acetate

25 The synthesis was performed using

[1-(4-chlorobenzyl)-4-(methylthio)-6-oxo-1,6-dihydro-1,3,5-triazin-2-yl]methyl acetate synthesized in Reference Synthesis Example 328 and

(R)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol synthesized in Reference Synthesis Example 133 in a similar manner to Reference Synthesis Example 348 (yield: 17%).

[1081] Synthesis Example 456

{4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-oxo-1-[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,6-dihydro-1,3,5-triazin-2-yl}methyl acetate

5 The synthesis was performed using

[4-(methylthio)-6-oxo-1-[5-(trifluoromethyl)thiophen-2-yl]methyl]-1,6-dihydro-1,3,5-triazin-2-yl]methyl acetate synthesized in Reference Synthesis Example 330 and 4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin in a similar manner to Reference Synthesis Example 348 (yield: 59%).

10 [1082] Synthesis Example 457

(R)-{4-[4-(4-Fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-oxo-1-[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,6-dihydro-1,3,5-triazin-2-yl}methyl acetate

The synthesis was performed using

15 [4-(methylthio)-6-oxo-1-[5-(trifluoromethyl)thiophen-2-yl]methyl]-1,6-dihydro-1,3,5-triazin-2-yl]methyl acetate synthesized in Reference Synthesis Example 330 and (R)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin-3-ol synthesized in Reference Synthesis Example 133 in a similar manner to Reference Synthesis Example 348 (yield: 46%).

[1083] Synthesis Example 458a

20 {4-[4-(4-Fluorophenyl)piperazin-1-yl]-6-oxo-1-[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,6-dihydro-1,3,5-triazin-2-yl}methyl acetate

Synthesis Example 458b

4-[4-(4-Fluorophenyl)piperazin-1-yl]-6-(hydroxymethyl)-1-[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25

The synthesis was performed using

[4-(methylthio)-6-oxo-1-[5-(trifluoromethyl)thiophen-2-yl]methyl]-1,6-dihydro-1,3,5-triazin-2-yl]methyl acetate synthesized in Reference Synthesis Example 330 and 1-(4-fluorophenyl)piperazine in a similar manner to Reference Synthesis Example 348 to

obtain the title compound (yield: 15%) of Synthesis Example 458a and the title compound (yield: 65%) of Synthesis Example 458b.

[1084] Synthesis Example 459

6-Ethyl-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-[[5-(trifluoromethyl)thiophen-2-yl]methyl]-1,3,5-triazin-2(1H)-one

The synthesis was performed using

6-ethyl-4-(methylthio)-1-[[5-(trifluoromethyl)thiophen-2-yl]methyl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 293 and 4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin in a similar manner to Reference Synthesis Example 348 (yield: 84%).

[1085] Synthesis Example 460

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-propyl-1-[[5-(trifluoromethyl)thiophen-2-yl]methyl]-1,3,5-triazin-2(1H)-one

The synthesis was performed using

4-(methylthio)-6-propyl-1-[[5-(trifluoromethyl)thiophen-2-yl]methyl]-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 294 and 4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine in a similar manner to Reference Synthesis Example 348 (yield: 84%).

[1086] Synthesis Example 461

1-(Adamantan-1-ylmethyl)-4-[4-(4-fluorophenyl)piperazin-1-yl]-6-methyl-1,3,5-triazin-2(1H)-one

The synthesis was performed using

1-(adamantan-1-ylmethyl)-6-methyl-4-(methylthio)-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 325 and 1-(4-fluorophenyl)piperazine in a similar manner to Reference Synthesis Example 348 (yield: 95%).

[1087] Synthesis Example 462

6-[(Dimethylamino)methyl]-4-[4-(4-fluorophenyl)piperazin-1-yl]-1-[[5-(trifluoromethyl)thiophen-2-yl]methyl]-1,3,5-triazin-2(1H)-one

To a dichloromethane solution (1.5 mL) of

- 5 4-[4-(4-fluorophenyl)piperazin-1-yl]-6-(hydroxymethyl)-1-[[5-(trifluoromethyl)thiophen-2-yl]methyl]-1,3,5-triazin-2(1H)-one (10.0 mg, 21.3 μ mol) synthesized in Synthesis Example 458b, triethylamine (6.0 μ L, 42.6 μ mol) and methanesulfonyl chloride (2.4 μ L, 32.0 μ mol) were added and the resultant mixture was stirred at room temperature for 1 hour and 30 minutes. The volume of the reaction solution was divided in half. To one
10 of the divided reaction solution, dimethylamine (20 μ L) and 1,4-dioxane (4.0 mL) were added and the resultant mixture was stirred at 90°C for 20 minutes. After the completion of the reaction, the reaction solution was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography (ethyl acetate) to obtain the title compound (1.05 mg, yield: 20%).

- 15 [1088] Synthesis Example 463

4-[4-(4-Fluorophenyl)piperazin-1-yl]-6-(morpholinomethyl)-1-[[5-(trifluoromethyl)thiophen-2-yl]methyl]-1,3,5-triazin-2(1H)-one

- The synthesis was performed using morpholine (20 μ L) in a similar manner to
20 Synthesis Example 462 to obtain the title compound (0.960 mg, yield: 17%).

[1089] Synthesis Example 464

1-(4-Chlorobenzyl)-4-[4-(4-fluorophenyl)-3-oxopiperidin-1-yl]-1,3,5-triazin-2(1H)-one

A toluene solution (10 mL) of

- 25 1-(4-chlorobenzyl)-4-[4-(4-fluorophenyl)-3,4-dihydroxypiperidin-1-yl]-1,3,5-triazin-2(1H)-one (202 mg, 0.468 mmol) synthesized in Synthesis Example 91 and p-toluenesulfonic acid monohydrate (341 mg, 1.99 mmol) was stirred under reflux by heating for 4 hours. After the completion of the reaction, the reaction solution was concentrated under reduced pressure and to the resultant residue, saturated sodium

bicarbonate aqueous solution was added, and extraction with ethyl acetate was performed. The organic layer was concentrated under reduced pressure and the resultant residue was purified by reverse-phase column chromatography (water/acetonitrile) to obtain the title compound (15.0 mg, yield: 8%).

5 [1090] Synthesis Example 465

1-(4-Chlorobenzyl)-4-{[3-(4-fluorophenyl)-2,3-dihydroxycyclopentyl]amino}-1,3,5-triazin-2(1H)-one

To an acetone(2.0 mL)-water(0.20 mL) solution of

10 1-(4-chlorobenzyl)-4-{[3-(4-fluorophenyl)cyclopent-2-en-1-yl]amino}-1,3,5-triazin-2(1H)-one (100 mg, 0.252 mmol) synthesized in Synthesis Example 121, immobilized osmium tetroxide (32.0 mg, 0.0126 mmol, osmium content: 9.4%) was added and the resultant mixture was stirred at room temperature for 3 hours. Then, to the reaction solution, 4-methylmorpholine N-oxide (44.3 mg, 0.378 mmol) was added and the resultant mixture
15 was stirred further for 3 days. After the completion of the reaction, to the reaction solution, saturated sodium thiosulfate aqueous solution was added and extraction with ethyl acetate from the reaction solution was performed. The organic layer was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography (chloroform-ethyl acetate) to obtain the title compound (15.8 mg,
20 yield: 12%).

[1091] Synthesis Example 466a

1-(4-Chlorobenzyl)-4-[3-fluoro-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

Synthesis Example 466b

25 1-(4-Chlorobenzyl)-4-[3,3-difluoro-4-(4-fluorophenyl)piperidin-1-yl]-1,3,5-triazin-2(1H)-one

The synthesis was performed using

1-(4-chlorobenzyl)-4-[4-(4-fluorophenyl)-3-oxopiperidin-1-yl]-1,3,5-triazin-2(1H)-one

synthesized in Synthesis Example 464 in a similar manner to Reference Synthesis Example 75 to obtain the title compound (2.40 mg, yield: 6%) of Synthesis Example 466a and the title compound (9.00 mg, yield: 6%) of Synthesis Example 466b.

[1092] Synthesis Example 467

5 1-(4-Chloro-3-hydroxybenzyl)-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one

To a dichloromethane solution (1.0 mL) of
1-(4-chloro-3-methoxybenzyl)-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-o
10 ne (10.0 mg, 0.0232 mmol) synthesized in Synthesis Example 141, 1M boron tribromide (1.00 mL, 1.00 mmol) was added at -78°C and while elevating gradually the temperature of the resultant mixture to room temperature, the mixture was stirred for 3 hours. After the completion of the reaction, to the reaction solution, water was added and extraction with ethyl acetate from the reaction solution was performed. The organic layer was
15 dried and was concentrated under reduced pressure and the resultant solid was washed with ethyl acetate to obtain the title compound (6.00 mg, yield: 62%).

[1093] Synthesis Example 468

4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-[4-(2,2,2-trifluoroethoxy)benzyl]-1,3,5-triazin-2(1H)-one
20

To an N,N-dimethylformamide solution of
4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-(4-hydroxybenzyl)-1,3,5-triazin-2(1H)-one (10.0 mg, 0.0260 mmol) synthesized in Synthesis Example 483a, cesium carbonate (10.3 mg, 0.0320 mmol) and 2,2,2-trifluoroethyl trifluoromethanesulfonate
25 (4.57 µL, 0.0320 mmol) were added and the resultant mixture was stirred at 70°C for 2 and a half hours. After the completion of the reaction, the reaction solution was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography (chloroform/ethyl acetate=1/1) to obtain the title compound (2.43 mg, yield: 14%).

[1094] Synthesis Example 469

4-[4-(4-(4-fluorophenyl)piperazin-1-yl)-1-{[4-methyl-5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

5 A 1,4-dioxane (2.0 mL)-water (0.25 mL) solution of
1-{[4-bromo-5-(trifluoromethyl)thiophen-2-yl]methyl}-4-[4-(4-fluorophenyl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one (50.9 mg, 0.0982 mmol) synthesized in Synthesis Example 154, tetrakis(triphenylphosphine) palladium (0) (11.6 mg, 0.0100 mmol), trimethyl borate (41 µL, 0.295 mmol), and sodium carbonate (20.8 mg, 0.196 mmol) was stirred in
10 nitrogen atmosphere at 100°C for 3 hours. After the completion of the reaction, to the reaction solution, water was added and a deposited solid was smeared on silica gel. The resultant mixture was eluted with ethyl acetate and the resultant eluate was concentrated under reduced pressure. The resultant solid was purified by silica gel column chromatography to obtain the title compound (5.82 mg, yield: 13%).

[1095] Synthesis Example 470

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[4-methyl-5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using

20 1-{[4-bromo-5-(trifluoromethyl)thiophen-2-yl]methyl}-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one (74.8 mg, 0.150 mmol) synthesized in Synthesis Example 165 in a similar manner to Synthesis Example 469 to obtain the title compound (1.50 mg, yield: 2%).

[1096] Synthesis Example 471

25 1-(4-chlorobenzyl)-4-{[3-(4-fluorophenyl)-2,3-dihydroxycyclopent-3-en-1-yl]amino}-1,3,5-triazin-2(1H)-one

A toluene solution (200 µL) of

1-(4-chlorobenzyl)-4-{[3-(4-fluorophenyl)-2,3-dihydroxycyclopentyl]amino}-1,3,5-triazin-

n-2(1H)-one (14.0 mg, 0.0325 mmol) synthesized in Synthesis Example 465 and p-toluenesulfonic acid monohydrate (6.18 mg, 0.0325 mmol) was stirred at 100°C for 8 hours. After the completion of the reaction, to the reaction solution, triethylamine was added and the reaction solution was concentrated under reduced pressure, followed by
 5 purifying the resultant residue by silica gel column chromatography (chloroform/ethyl acetate=1/2) to obtain the title compound (7.00 mg, yield: 52%).

[1097] Synthesis Example 472

1-[4-(difluoromethoxy)benzyl]-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

10

To an acetonitrile (1.0 mL)-water (1.0 mL) solution of 4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-(4-hydroxybenzyl)-1,3,5-triazin-2(1H)-one (15.0 mg, 0.0396 mmol) synthesized in Synthesis Example 483a, potassium hydroxide (44.5 mg, 0.793 mmol) was added and while stirring the resultant reaction
 15 solution at -78°C, to the reaction solution, diethyl (bromodifluoromethyl)phosphonate (14.1 µL, 0.0794 mmol) was added, followed by stirring the resultant reaction solution at room temperature for 2 and a half hours. After the completion of the reaction, extraction with diethyl ether from the reaction solution was performed. The organic layer was dried over anhydrous sodium sulfate and was concentrated under reduced
 20 pressure. The resultant residue was purified by silica gel column chromatography to obtain the title compound (2.10 mg, yield: 11%).

[1098] Synthesis Example 473

4-(1-[4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-2-oxo-1,3,5-triazin-1(2H)-yl]methyl)phenyl trifluoromethanesulfonate

25

To a dichloromethane solution of 4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-(4-hydroxybenzyl)-1,3,5-triazin-2(1H)-one (50.0 mg, 0.132 mmol) synthesized in Synthesis Example 483a, N,N-diisopropylethylamine (27.0 µL, 0.158 mmol) was added and while stirring the

resultant mixture at 0°C, to the mixture, trifluoromethanesulfonic acid anhydride (23.8 µL, 0.145 mmol) was added, followed by stirring the resultant mixture at room temperature for 1 day. After the completion of the reaction, the reaction solution was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography (chloroform/ethyl acetate=2/1) to obtain the title compound (2.47 mg, yield: 4%).

[1099] Synthesis Example 474

4-[5-Fluoro-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using

4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (44.0 mg, 0.973 mmol) synthesized in Synthesis Example 190 in a similar manner to Synthesis Example 107 to obtain the title compound (28.0 mg, yield: 63%).

[1100] Synthesis Example 475

4-[4-(4-fluorophenyl)-5-methoxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

To an N,N-dimethylformamide (2.0 mL)-dichloromethane (2.0 mL) solution of 4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (50.0 mg, 0.111 mmol) synthesized in Synthesis Example 190, triethylamine (23.2 µL, 0.167 mmol) and methyl iodide (20.6 µL, 0.332 mmol) were added at room temperature and the resultant mixture was stirred at room temperature for 1 day. Next, to the reaction solution, potassium carbonate (50.0 mg, 0.362 mmol) and methyl iodide (40.0 µL, 0.643 mmol) were added and the resultant mixture was stirred at 50°C for 1 hour. Then, to the reaction solution, sodium hydride (12.0 mg, 0.500 mmol) and methyl iodide (40.0 µM, 0.643 mmol) were added and the resultant mixture was stirred at room temperature for 1 day. After the completion of the

reaction, to the reaction solution, water was added and extraction with chloroform from the reaction solution was performed. The organic layer was dried over anhydrous magnesium sulfate and was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography to obtain the title compound (7.80 mg, yield: 15%).

[1101] Synthesis Example 476

4-(4-Fluorophenyl)-1-(4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)-1,2,3,6-tetrahydropyridin-3-yl acetate

10 To

4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (20.0 mg, 0.0442 mmol) synthesized in Synthesis Example 190, at room temperature, acetic anhydride (2.0 mL) and sodium acetate (9.00 mg, 0.110 mmol) were added at room temperature and the resultant mixture was stirred at 100°C for 40 minutes. After the completion of the reaction, the reaction solution was concentrated and to the resultant residue, water was added, followed by filtering a deposited colorless solid and by drying the solid to obtain the title compound (20.6 mg, yield: 94%).

[1102] Synthesis Example 477

20 4-[4-(4-fluorophenyl)-5-oxo-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

To

4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (20.0 mg, 0.0442 mmol) synthesized in Synthesis Example 190, dichloromethane (1.0 mL), sodium bicarbonate (11.1 mg, 0.132 mmol), and Dess-Martin periodinane (28.1 mg, 0.0663 mmol; manufactured by Tokyo Chemical Industry Co., Ltd.) were added and the resultant mixture was stirred at room temperature for 1 day. Then, to the reaction solution, dichloromethane (4.0 mL) and

Dess-Martin periodinane (30.0 mg, 0.0707 mmol) were added and the resultant mixture was stirred at room temperature for 2 hours. After the completion of the reaction, the reaction solution was purified by silica gel column chromatography (ethyl acetate) to obtain the title compound (19.4 mg, yield: 98%).

5 [1103] Synthesis Example 478

4-[4-(4-fluorophenyl)-1-(4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)-1,2,3,6-tetrahydropyridin-3-yl]formate

To a crude product synthesized using

10 4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (50.0 mg, 0.111 mmol) synthesized in Synthesis Example 190 in a similar manner to Reference Synthesis Example 68, N,N-dimethylformamide (1.0 mL) and copper cyanide (49.3 mg, 0.550 mmol) were added and the resultant mixture was stirred at 100°C for 1 day. After the completion of
15 the reaction, to the reaction solution, saturated sodium chloride aqueous solution was added and extraction with ethyl acetate from the reaction solution was performed. The organic layer was dried over anhydrous sodium sulfate and was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography (hexane/ethyl acetate=1/1→0/1) to obtain the title compound (0.8 mg,
20 yield: 2%).

[1104] Synthesis Example 479

1-[(5-Cyclopropylthiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one

25 The synthesis was performed using

1-[(5-bromothiophen-2-yl)methyl]-4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1,3,5-triazin-2(1H)-one (120 mg, 0.268 mmol) synthesized in Synthesis Example 175 in a similar manner to Synthesis Example 469 to obtain the title compound (2.92 mg, yield: 3%).

[1105] Synthesis Example 480a

4-[3,5-Difluoro-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

Synthesis Example 480b

5 4-[5,5-Difluoro-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

To

4-[4-(4-fluorophenyl)-5-oxo-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (60.0 mg, 0.133 mmol) synthesized in Synthesis Example 477, bis(2-methoxyethyl)aminosulfur trifluoride (1.5 mL) was added and the resultant mixture was stirred at 80°C for 1 hour. After the completion of the reaction, the reaction solution was ice-cooled and was added to sodium carbonate aqueous solution dropwise and extraction with chloroform from the reaction solution was performed twice.
15 The organic layer was dried over anhydrous sodium sulfate and was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography. The resultant mixture was purified by preparative reverse-phase high performance liquid chromatography (gradient: acetonitrile/water (0.1% formic acid)=40/60→60/40→100/0, column: Waters X Bridge C18 5 μm Φ 19×100 mm) to
20 obtain the title compound (6.80 mg, yield: 11%) of Synthesis Example 480a and the title compound (7.30 mg, yield: 12%) of Synthesis Example 480b.

[1106] Synthesis Example 481

4-[5-(Dimethylamino)-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

25

To

4-[5-amino-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (10.0 mg, 0.0222 mmol) synthesized in Synthesis Example 321, a formalin aqueous solution (37%) (0.80 mL) and formic acid

(1.2 mL) were added and the resultant mixture was stirred at 90°C for 1 hour. After the completion of the reaction, to the reaction solution, saturated potassium carbonate aqueous solution was added and extraction with ethyl acetate from the reaction solution was performed. The organic layer was dried over anhydrous magnesium sulfate and
 5 was concentrated under reduced pressure and the resultant residue was washed with hexane to obtain the title compound (7.44 mg, yield: 70%).

[1107] Synthesis Example 482

4-[5-Fluoro-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10

The synthesis was performed using

(R)-4-[4-(4-fluorophenyl)-5-hydroxy-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (44.0 mg, 0.0943 mmol)

synthesized in Synthesis Example 342 in a similar manner to Reference Synthesis

15 Example 75 to obtain a mixture of optical isomers of the title compound (22.4 mg, yield: 43%).

[1108] Synthesis Example 483a

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-(4-hydroxybenzyl)-1,3,5-triazin-2(1H)-one

20

Synthesis Example 483b

4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-[4-[(4-hydroxybenzyl)oxy]benzyl]-1,3,5-triazin-2(1H)-one

To a dichloromethane solution (8.0 mL) of

25 4-[4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-(4-methoxybenzyl)-1,3,5-triazin-2(1H)-one (200 mg, 0.510 mmol) synthesized in Reference Synthesis Example 352, boron tribromide (1.53 mL, 1.53 mmol) was added at 0°C and the resultant mixture was stirred at room temperature for 1 day. After the completion of the reaction, to the reaction solution, water was added and extraction with chloroform from the reaction solution was

performed. The organic layer was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography (chloroform/ethyl acetate → methanol/ethyl acetate) to obtain the title compound (35.0 mg, yield: 18%) of Synthesis Example 483a and the title compound (7.15 mg, yield: 3%) of Synthesis Example 483b.

5 [1109] Synthesis Example 484

4-[4-(4-Fluorophenyl)pyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

To an N,N-dimethylformamide solution (1.0 mL) of

10 4-[5-chloro-4-(4-fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (10.0 mg, 0.0212 mmol) synthesized in Reference Synthesis Example 353, triethylamine (10.0 μ L, 0.0717 mmol) and potassium cyanide (10.0 mg, 0.154 mmol) were added and the resultant mixture was stirred at room temperature for 2.5 hours and at 80°C for 40 minutes. After the completion of the
15 reaction, to the reaction solution, saturated sodium bicarbonate aqueous solution was added and extraction with ethyl acetate from the reaction solution was performed. The organic layer was dried over anhydrous sodium sulfate and was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography (hexane/ethyl acetate=1/1→0/1) to obtain the title compound (0.7 mg,
20 yield: 8%).

[1110] Synthesis Example 485

4-[4-(4-Fluorophenyl)-3,3-bis(hydroxymethyl)piperidin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

To a 1,4-dioxane solution (500 μ L) of

25 4-[11-(4-fluorophenyl)-2,4-dioxo-8-azaspiro[5.5]undecan-8-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (5.00 mg, 9.53 μ mol) synthesized in Reference Synthesis Example 363, 12 M hydrochloric acid (500 μ L) was added and the resultant mixture was stirred at 110°C for 2 hours. To the resultant reaction solution, 12 M hydrochloric acid (500 μ L) was added and the resultant mixture was stirred at

110°C for 2.5 hours. After the completion of the reaction, to the reaction solution, saturated sodium carbonate aqueous solution was added and extraction with chloroform from the reaction solution was performed. The organic layer was dried over anhydrous sodium sulfate and was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography (ethyl acetate/methanol=1/0→20/1) to obtain the title compound (1.59 mg, yield: 33%).

[1111] Synthesis Example 486

4-[11-(4-Fluorophenyl)-2,4-dioxo-8-azaspiro[5.5]undeca-10-en-8-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

10

The synthesis was performed using a product obtained by synthesizing using 6-methyl-4-(methylthio)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 292 and 4-(4-fluorophenyl)-3,3-bis[(methoxymethoxy)methyl]-1,2,3,6-tetrahydropyridine synthesized in Reference Synthesis Example 359 in a similar manner to Reference Synthesis Example 348, in a similar manner to Synthesis Example 485 (yield: 37%).

15

[1112] Synthesis Example 487

4-[4-[2-(Adamantan-1-yl)-2-oxoethyl]piperazin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

20

To an acetonitrile solution (4.0 mL) of 6-methyl-4-(piperazin-1-yl)-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one (28.7 mg, 80.0 μmol) synthesized in Reference Synthesis Example 351, 1-(adamantan-1-yl)-2-bromoethanone (23.1 mg, 90.0 μmol), potassium carbonate (13.8 mg, 100 μmol), and sodium iodide (15.0 mg, 100 μmol) were added and the resultant mixture was stirred at room temperature for 1 day. After the completion of the reaction, to the reaction solution, saturated sodium bicarbonate aqueous solution was added and extraction with ethyl acetate from the reaction solution was performed. The organic layer was concentrated under reduced pressure and the resultant residue was purified by

25

silica gel column chromatography (chloroform/methanol=10/0→9/1) to obtain the title compound (20.8 mg, yield: 49%).

[1113] Synthesis Example 488

4-[4-({3-[1-(4-Fluorophenyl)cyclopropyl]-1,2,4-oxadiazol-5-yl}methyl)piperazin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using

5-(chloromethyl)-3-[1-(4-fluorophenyl)cyclopropyl]-1,2,4-oxadiazole (22.7 mg, 90.0 μ mol) in a similar manner to Synthesis Example 487 to obtain the title compound (27.2 mg, yield: 59%).

[1114] Synthesis Example 489

4-[4-(4-Bromo-1H-pyrazol-1-yl)piperidin-1-yl]-6-methyl-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

To an N,N-dimethylformamide solution (1.0 mL) of 1-(6-methyl-4-oxo-5-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-4,5-dihydro-1,3,5-triazin-2-yl)piperidin-4-yl methanesulfonate (45.2 mg, 0.100 mmol) synthesized in Reference Synthesis Example 349, 4-bromo-1H-pyrazole (29.4 mg, 0.200 mmol) and potassium carbonate (27.6 mg, 0.200 mmol) were added and the resultant mixture was stirred at 100°C for 1 hour. After the completion of the reaction, to the reaction solution, water was added and extraction with ethyl acetate from the reaction solution was performed. The organic layer was dried over anhydrous sodium sulfate and was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography (ethyl acetate) to obtain the title compound (24.4 mg, yield: 48%).

[1115] Synthesis Example 490

6-methyl-4-{4-[4-(Trifluoromethyl)-1H-pyrazol-1-yl]piperidin-1-yl}-1-{[5-(trifluoromethyl)thiophen-2-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using 4-trifluoromethyl-1H-pyrazole (27.2 mg, 0.200

mmol) in a similar manner to Synthesis Example 489 to obtain the title compound (23.9 mg, yield: 49%).

[1116] In Synthesis Examples 491 to 493, each title compound was synthesized using

- 5 6-methyl-4-(methylthio)-1-{[2-(trifluoromethyl)thiazol-5-yl]methyl}-1,3,5-triazin-2(1H)-one synthesized in Reference Synthesis Example 374 in a similar manner to Reference Synthesis Example 348. The names of the synthesized compounds and the synthesis yields are indicated below.

[1117] Synthesis Example 491

- 10 (R)-4-(5'-Hydroxy-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-6-methyl-1-{[2-(trifluoromethyl)thiazol-5-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 69%

[1118] Synthesis Example 492

- 15 (R)-4-(5-Fluoro-5'-hydroxy-6-methyl-5',6'-dihydro-[2,4'-bipyridin]-1'(2'H)-yl)-6-methyl-1-{[2-(trifluoromethyl)thiazol-5-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 78%

[1119] Synthesis Example 493

- 20 (R)-4-[5-Hydroxy-4-(5-methylthiophen-3-yl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[2-(trifluoromethyl)thiazol-5-yl]methyl}-1,3,5-triazin-2(1H)-one

Yield: 59%

[1120] Synthesis Example 494

- 25 4-[4-(4-Fluorophenyl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[2-(trifluoromethyl)thiazol-5-yl]methyl}-1,3,5-triazin-2(1H)-one

To a 1,4-dioxane solution (500 μ L) of
6-methyl-4-(methylthio)-1-{[2-(trifluoromethyl)thiazol-5-yl]methyl}-1,3,5-triazin-2(1H)-

one (38.7 mg, 0.120 mmol) synthesized in Reference Synthesis Example 374, 4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine hydrochloride (30.8 mg, 0.144 mmol) and N,N-diisopropylethylamine (41.1 μ L, 0.240 mmol) were added and the resultant mixture was stirred under reflux by heating for 1 day. After the completion of the reaction, the reaction solution was purified by silica gel column chromatography (hexane/ethyl acetate=1/1 $\rightarrow$ 0/1) to obtain the title compound (33.3 mg, yield: 61%).

[1121] Synthesis Example 495

6-Methyl-4-[4-(5-methylthiophen-3-yl)-5,6-dihydropyridin-1(2H)-yl]-1-[2-(trifluoromethyl)thiazol-5-yl]methyl}-1,3,5-triazin-2(1H)-one

To tert-Butyl 4-(5-methylthiophen-3-yl)-5,6-dihydropyridine-1(2H)-carboxylate (50.0 mg, 0.179 mmol), 4M hydrogen chloride/1,4-dioxane (500 μ L) was added and the resultant mixture was stirred at room temperature for 1 day. After the completion of the reaction, using a crude product obtained by concentrating the resultant reaction solution, the synthesis was performed in a similar manner to Synthesis Example 494 to obtain the title compound (30.0 mg, yield: 44%).

[1122] Synthesis Example 496

4-[4-(5-Methylthiophen-3-yl)piperazin-1-yl]-1-[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl}-1,3,5-triazin-2(1H)-one

The synthesis was performed using 1-(5-methylthiophen-3-yl)piperazine (401 mg, 2.20 mmol) synthesized in Reference Synthesis Example 286 by the same method as the total method of the method in Reference Synthesis Example 341 and the method in Reference Synthesis Example 342 to obtain a crude product of

4-[4-(5-methylthiophen-3-yl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one. Using the obtained crude product (50.0 mg) of

4-[4-(5-methylthiophen-3-yl)piperazin-1-yl]-1,3,5-triazin-2(1H)-one and

[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl 4-methylbenzenesulfonate (70.0 mg, 0.219 mmol) synthesized in Reference Synthesis Example 57, the synthesis was performed in a

similar manner to Synthesis Example 1 to obtain the title compound (32.8 mg, yield: 43%).

[1123] Synthesis Example 497


4-[5-Fluoro-4-(5-methylthiophen-3-yl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[2-(trifluoromethyl)thiazol-5-yl]methyl}-1,3,5-triazin-2(1H)-one



































The synthesis was performed using


























(R)-4-[5-hydroxy-4-(5-methylthiophen-3-yl)-5,6-dihydropyridin-1(2H)-yl]-6-methyl-1-{[2-(trifluoromethyl)thiazol-5-yl]methyl}-1,3,5-triazin-2(1H)-one (15.8 mg, 0.0337 mmol) synthesized in Synthesis Example 493 in a similar manner to Reference Synthesis Example 75 to obtain the title compound (4.80 mg, yield: 30%) as a mixture of optical isomers.


































[1124] The chemical structural formulae of the compounds synthesized in Reference Synthesis Examples and Synthesis Examples will be shown hereinafter.
























Tables 2 to 6 show the data of physical properties of the compounds synthesized in Reference Synthesis Examples, and Tables 7 to 13 show the data of physical properties of the compounds synthesized in Synthesis Examples. As described above, the sign Rf in the drawings means Reference Synthesis Example, and the sign Ex means Synthesis Example.

































































































Rf 261




Rf 262




Rf 263




Rf 264




Rf 265




Rf 266




Rf 267




Rf 268




Rf 269




Rf 270




Rf 271




Rf 272




Rf 273




Rf 274




Rf 275




Rf 276




Rf 277




Rf 278



Rf 279



Rf 280




Rf 281




Rf 282




Rf 283



Rf 284




Rf 285




Rf 286




Rf 287




Rf 288




Rf 289




Rf 290




Rf 291




Rf 292







Rf 323




Rf 324




Rf 325




Rf 326




Rf 327




Rf 328




Rf 329




Rf 330




Rf 331




Rf 332a




Rf 332b




Rf 333a




Rf 333b




Rf 334




Rf 335




Rf 336




Rf 337




Rf 338




Rf 339



Rf 340




Rf 341




Rf 342




Rf 343




Rf 344



Rf 345




























Rf 346















Rf 347

345






























































































































































































































































































































[Table 3]

Rf	Data
1	Morphology: colorless solid, $^1\text{H-NMR}$ (CDCl_3) δ : 3.18 (t, J = 5.1 Hz, 4H), 4.07 (t, J = 5.1 Hz, 4H), 6.96–7.04 (m, 4H).
2	Morphology: colorless solid
3	Morphology: colorless solid LC/MS: cond. 1 RT 2.97 min LC/MS(ESI^+) m/z ; 276 $[\text{M}+\text{H}]^+$
5	Morphology: colorless solid LC/MS: cond. 1 RT 4.56 min LC/MS(ESI^+) m/z ; 334 $[\text{M}+\text{H}]^+$
6	Morphology: colorless solid LC/MS: cond. 1 RT 3.44 min LC/MS(ESI^+) m/z ; 316 $[\text{M}+\text{H}]^+$
7	Morphology: colorless solid LC/MS: cond. 1 RT 3.06 min LC/MS(ESI^+) m/z ; 282 $[\text{M}+\text{H}]^+$ LC/MS(ESI^-) m/z ; 280 $[\text{M}-\text{H}]^-$
8	Morphology: colorless solid $^1\text{H-NMR}$ (CDCl_3) δ : 1.48 (s, 9H), 3.43–3.50 (m, 4H), 3.80–3.90 (m, 4H), 4.94 (s, 2H), 7.25–7.39 (m, 4H), 7.92 (s, 1H). LC/MS: cond. 1 RT 4.19 min LC/MS(ESI^+) m/z ; 406 $[\text{M}+\text{H}]^+$
9	Morphology: colorless solid LC/MS: cond. 2 RT 1.31 min LC/MS(ESI^+) m/z ; 306 $[\text{M}+\text{H}]^+$
10	Morphology: white solid $^1\text{H-NMR}$ (CDCl_3) δ : 4.90 (s, 2H), 7.30–7.45 (m, 4H), 8.56 (s, 1H), 11.6 (s, 1H). LC/MS: cond. 2 RT 1.61 min LC/MS(ESI^+) m/z ; 238 $[\text{M}+\text{H}]^+$
11	LC/MS: cond. 2 RT 1.97 min LC/MS(ESI^+) m/z ; 256 $[\text{M}+\text{H}]^+$
12	Morphology: brown amorphous LC/MS: cond. 1 RT 4.39 min LC/MS(ESI^+) m/z ; 343 $[\text{M}+\text{H}]^+$
13	Morphology: colorless solid LC/MS: cond. 1 RT 3.32 min LC/MS(ESI^+) m/z ; 325 $[\text{M}+\text{H}]^+$
14	Morphology: pale yellow solid LC/MS: cond. 1 RT 2.74 min LC/MS(ESI^+) m/z ; 291 $[\text{M}+\text{H}]^+$
15	$^1\text{H-NMR}$ (CDCl_3) δ : 2.41 (s, 3H), 6.05 (s, 2H), 6.46 (d, J = 2.5 Hz, 1H), 7.26 (d, J = 8.2 Hz, 2H), 7.60–7.62 (m, 1H), 7.65 (d, J = 8.2 Hz, 2H).
16	Morphology: colorless solid $^1\text{H-NMR}$ (CDCl_3) δ : 1.47 (s, 9H), 3.43–3.50 (m, 4H), 3.82–3.91 (m, 4H), 6.01 (s, 2H), 6.54 (d, J = 2.3 Hz, 1H), 7.90–7.94 (m, 1H), 8.30 (s, 1H). LC/MS: cond. 1 RT 4.14 min LC/MS(ESI^+) m/z ; 430 $[\text{M}+\text{H}]^+$
17	Morphology: colorless amorphous LC/MS: cond. 1 RT 0.35 min LC/MS(ESI^+) m/z ; 330 $[\text{M}+\text{H}]^+$
18	Morphology: white solid LC/MS: cond. 1 RT 2.60 min LC/MS(ESI^+) m/z ; 292 $[\text{M}+\text{H}]^+$
19	Morphology: white solid $^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ : 3.16 (m, 4H), 3.78 (m, 4H), 4.86 (s, 2H), 6.97–7.10 (m, 4H), 7.28–7.38 (m, 4H). LC/MS: cond. 1 RT 4.00 min LC/MS(ESI^+) m/z ; 416 $[\text{M}+\text{H}]^+$
20	Morphology: pale yellow solid
21	Morphology: colorless solid LC/MS: cond. 1 RT 3.40 min LC/MS(ESI^+) m/z ; 341 $[\text{M}+\text{H}]^+$
22a	Morphology: colorless amorphous
22b	Morphology: colorless amorphous

[Table 4]

Rf	Data
23	Morphology: pale brown solid LC/MS: cond. 1 RT 3.85 min LC/MS (ESI ⁺) m/z; 295 [M+H] ⁺
24	Morphology: pale brown solid ¹ H-NMR (CDCl ₃) δ : 1.39–1.57 (m, 2H), 2.00 (d, J = 13.0 Hz, 2H), 2.44–2.54 (m, 4H), 2.72 (dt, J = 10.5, 2.0 Hz, 2H), 3.46–3.57 (m, 2H), 6.84–6.98 (m, 4H). LC/MS: cond. 1 RT 0.39 min LC/MS (ESI ⁺) m/z; 209 [M+H] ⁺
25	Morphology: pale yellow solid ¹ H-NMR (CDCl ₃) δ : 1.54 (s, 9H), 1.61–1.90 (m, 4H), 2.84–3.10 (m, 3H), 3.20–3.30 (m, 1H), 3.78–3.92 (m, 1H), 4.92 (br. s, 1H), 6.83–7.00 (m, 4H). LC/MS: cond. 1 RT 4.15 min LC/MS (ESI ⁺) m/z; 295 [M+H] ⁺
26	Morphology: colorless solid ¹ H-NMR (DMSO-d ₆) δ : 1.55–2.05 (m, 4H), 2.90–3.15 (m, 2H), 3.22–3.58 (m, 3H), 7.15 (d, J = 6.6 Hz, 4H), 8.49 (br. s, 3H). LC/MS: cond. 1 RT 0.57 min LC/MS (ESI ⁺) m/z; 195 [M+H] ⁺
27	Morphology: pale yellow solid ¹ H-NMR (CDCl ₃) δ : 1.54 (s, 9H), 1.61–1.90 (m, 4H), 2.84–3.10 (m, 3H), 3.20–3.30 (m, 1H), 3.78–3.92 (m, 1H), 4.92 (br. s, 1H), 6.83–7.00 (m, 4H).
28	Morphology: colorless solid LC/MS: cond. 1 RT 0.50 min LC/MS (ESI ⁺) m/z; 195 [M+H] ⁺
29	Morphology: orange solid LC/MS: cond. 1 RT 4.40 min LC/MS (ESI ⁺) m/z; 281 [M+H] ⁺ LC/MS (ESI [−]) m/z; 325 [M+HCO ₂] [−]
30	Morphology: colorless solid LC/MS: cond. 1 RT 0.42 min LC/MS (ESI ⁺) m/z; 181 [M+H] ⁺
31	Morphology: pale orange solid ¹ H-NMR (CDCl ₃) δ : 1.45 (s, 9H), 1.86–2.00 (m, 1H), 2.23–2.35 (m, 1H), 3.12 (dd, J = 9.5, 3.8 Hz, 1H), 3.21–3.55 (m, 3H), 4.35 (br. s, 1H), 4.72 (br. s, 1H), 6.42–6.52 (m, 2H), 6.93 (t, J = 8.9 Hz, 2H).
32	Morphology: colorless solid LC/MS: cond. 1 RT 0.47 min LC/MS (ESI ⁺) m/z; 181 [M+H] ⁺
33	Morphology: colorless oil ¹ H-NMR (CDCl ₃) δ : 1.48 (s, 9H), 1.84–2.04 (m, 2H), 2.08–2.20 (m, 2H), 2.80–2.96 (m, 2H), 4.20–4.40 (m, 3H), 6.56 (dd, J = 17.1, 2.0 Hz, 1H), 7.49 (d, J = 20.1 Hz, 1H).
34	Morphology: colorless oil
36	Morphology: brown oil
37	Morphology: colorless oil
39	LC/MS: cond. 2 RT 2.22 min LC/MS (ESI ⁺) m/z; 312 [M+H] ⁺
40a	LC/MS: cond. 2 RT 0.60 min LC/MS (ESI ⁺) m/z; 212 [M+H] ⁺
40b	Morphology: brown solid ¹ H-NMR (DMSO-d ₆) δ : 2.93–3.01 (m, 1H), 3.19–3.27 (m, 1H), 3.48–3.80 (m, 2H), 4.39 (s, 1H), 6.15 (t, J = 3.3 Hz, 1H), 7.00–7.08 (m, 2H), 7.46–7.54 (m, 2H). LC/MS: cond. 2 RT 0.97 min LC/MS (ESI ⁺) m/z; 194 [M+H] ⁺
41	Morphology: colorless solid LC/MS: cond. 1 RT 0.52 min LC/MS (ESI ⁺) m/z; 170 [M+H] ⁺

[Table 5]

Rf	Data
42	Morphology: brown oil LC/MS : cond. 1 RT 3.10 min LC/MS (ESI ⁺) m/z; 290 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 288 [M-H] ⁻
43	Morphology: colorless solid LC/MS : cond. 1 RT 3.12 min LC/MS (ESI ⁺) m/z; 210 [M+H] ⁺
44	Morphology: colorless oil LC/MS : cond. 1 RT 0.54 min LC/MS (ESI ⁺) m/z; 196 [M+H] ⁺
45	Morphology: colorless solid LC/MS : cond. 1 RT 0.34 min LC/MS (ESI ⁺) m/z; 195 [M+H] ⁺
46	Morphology: brown oil LC/MS : cond. 2 RT 1.85 min LC/MS (ESI ⁺) m/z; 182 [M+H] ⁺
47	Morphology: colorless solid LC/MS : cond. 2 RT 0.37 min LC/MS (ESI ⁺) m/z; 195 [M+H] ⁺
48	Morphology: black oil
49	Morphology: pink solid LC/MS : cond. 1 RT 1.74 min LC/MS (ESI ⁺) m/z; 196 [M+H] ⁺
50	Morphology: yellow oil LC/MS : cond. 2 RT 0.86 min LC/MS (ESI ⁺) m/z; 195 [M+H] ⁺
51	Morphology: brown oil
52	Morphology: yellow oil LC/MS : cond. 1 RT 4.25 min LC/MS (ESI ⁺) m/z; 321 [M+H] ⁺
53	LC/MS : cond. 1 RT 3.68 min LC/MS (ESI ⁺) m/z; 223 [M+H] ⁺
54	Morphology: orange oil LC/MS : cond. 1 RT 4.21 min LC/MS (ESI ⁺) m/z; 335 [M+H] ⁺
55	LC/MS : cond. 2 RT 1.85 min LC/MS (ESI ⁻) m/z; 183 [M-H] ⁻
56	Morphology: brown oil LC/MS : cond. 2 RT 2.67 min LC/MS (ESI ⁺) m/z; 261 [M+H] ⁺
57	Morphology: colorless solid
58	Morphology: red oil LC/MS : cond. 1 RT 3.56 min LC/MS (ESI ⁺) m/z; 286 [M+H] ⁺
59	Morphology: brown solid LC/MS : cond. 1 RT 3.67 min LC/MS (ESI ⁺) m/z; 173 [M+H] ⁺
60	Morphology: pale yellow oil LC/MS : cond. 1 RT 4.30 min LC/MS (ESI ⁺) m/z 338 [M+H] ⁺
61	Morphology: brown solid LC/MS : cond. 1 RT 3.88 min LC/MS (ESI ⁺) m/z; 187 [M+H] ⁺
62	Morphology: brown oil LC/MS : cond. 1 RT 4.34 min LC/MS (ESI ⁺) m/z; 338 [M+H] ⁺
63	Morphology: yellow oil LC/MS : cond. 1 RT 3.09 min LC/MS (ESI ⁺) m/z; 212 [M+H] ⁺
64	Morphology: yellow oil LC/MS : cond. 1 RT 4.50 min LC/MS (ESI ⁺) m/z; 284 [M+H] ⁺
65	Morphology: pale yellow solid LC/MS : cond. 1 RT 0.42 min LC/MS (ESI ⁺) m/z; 184 [M+H] ⁺
66	Morphology: yellow oil LC/MS : cond. 1 RT 4.50 min LC/MS (ESI ⁺) m/z; 298 [M+H] ⁺

[Table 6]

Rf	Data
67	Morphology: pale yellow oil LC/MS : cond. 1 RT 0.47 min LC/MS (ESI ⁺) m/z; 198 [M+H] ⁺
68	Morphology: yellow oil
69	Morphology: white solid ¹ H-NMR (CDCl ₃) δ : 1.92 (t, J = 6.0 Hz, 1H), 4.47 (d, J = 6.0 Hz, 2H), 6.62 (s, 1H), 7.23 (dd, J = 2.2, 8.4 Hz, 1H), 7.37 (d, J = 8.4 Hz, 1H), 7.52 (d, J = 2.2 Hz, 1H). LC/MS : cond. 2 RT 1.71 min LC/MS (ESI ⁺) m/z; 283 [M+H] ⁺
70	Morphology: pale yellow oil
71	Morphology: colorless solid LC/MS : cond. 1 RT 3.94 min LC/MS (ESI ⁺) m/z; 383 [M+H] ⁺
72	Morphology: brown oil LC/MS : cond. 2 RT 1.71 min LC/MS (ESI ⁺) m/z; 283 [M+H] ⁺
73	Morphology: brown amorphous ¹ H-NMR (CDCl ₃) δ : 1.81 (d, J = 3.9 Hz, 1H), 3.06 (dd, J = 6.5 Hz, 9.5 Hz, 1H), 3.28-3.38 (m, 1H), 3.38-3.49 (m, 2H) 3.59 (dd, J = 7.1 Hz, 9.2 Hz, 1H), 3.68 (d, J = 13.5 Hz, 2H), 3.84 (d, J = 13.8 Hz, 2H), 4.45-4.54 (m, 1H), 6.40-6.49 (m, 2H), 6.93 (t, J = 8.9 Hz, 2H), 7.29-7.41 (m, 10H). LC/MS : cond. 2 RT 2.41 min LC/MS (ESI ⁺) m/z; 377 [M+H] ⁺
74	Morphology: brown oil ¹ H-NMR (CDCl ₃) δ : 3.03 (dd, J = 4.7 Hz, 9.5 Hz, 1H) 3.20 (dd, J = 4.2 Hz, 9.6 Hz, 1H), 3.43-3.54 (m, 1H), 3.61-3.72 (m, 2H), 4.10-4.17 (m, 1H), 6.43-6.48 (m, 2H), 6.94 (t, J = 8.9 Hz, 2H). LC/MS : cond. 2 RT 1.13 min LC/MS (ESI ⁺) m/z; 197 [M+H] ⁺
75	Morphology: colorless solid LC/MS : cond. 1 RT 5.08 min LC/MS (ESI ⁺) m/z; 385 [M+H] ⁺
76	Morphology: pale yellow amorphous
77	Morphology: colorless amorphous LC/MS : cond. 2 RT 3.43 min LC/MS (ESI ⁺) m/z; 379 [M+H] ⁺
78	Morphology: colorless solid LC/MS : cond. 2 RT 1.39 min LC/MS (ESI ⁺) m/z; 199 [M+H] ⁺
79	Morphology: yellow solid LC/MS : cond. 1 RT 0.37 min LC/MS (ESI ⁺) m/z; 281 [M+H] ⁺
80	Morphology: pale yellow solid LC/MS : cond. 1 RT 4.63 min LC/MS (ESI ⁺) m/z; 295 [M+H] ⁺
81	Morphology: brown oil LC/MS : cond. 2 RT 1.21 min LC/MS (ESI ⁺) m/z; 195 [M+H] ⁺
82	Morphology: pale brown solid LC/MS : cond. 1 RT 3.42 min LC/MS (ESI ⁺) m/z; 295 [M+H] ⁺
83	Morphology: colorless solid LC/MS : cond. 1 RT 0.39 min LC/MS (ESI ⁺) m/z; 195 [M+H] ⁺
84	Morphology: colorless solid LC/MS : cond. 1 RT 4.70 min LC/MS (ESI ⁺) m/z; 346 [M+H] ⁺

[Table 7]

Rf	Data
85	Morphology: colorless solid LC/MS: cond. 1 RT 3.59 min LC/MS(ESI ⁺) m/z; 328 [M+H] ⁺
86	Morphology: pale yellow amorphous LC/MS: cond. 1 RT 3.22 min LC/MS(ESI ⁺) m/z; 294 [M+H] ⁺
87	Morphology: colorless solid LC/MS: cond. 1 RT 4.24 min LC/MS(ESI ⁺) m/z; 464 [M+H] ⁺
88	Morphology: colorless oil LC/MS: cond. 1 RT 0.94 min LC/MS(ESI ⁺) m/z; 364 [M+H] ⁺
89	Morphology: colorless amorphous LC/MS: cond. 1 RT 3.99 min LC/MS(ESI ⁺) m/z; 439 [M+H] ⁺
90	Morphology: colorless solid LC/MS: cond. 2 RT 2.39 min LC/MS(ESI ⁺) m/z; 455 [M+H] ⁺
91 a	Morphology: colorless amorphous ¹ H-NMR (CDCl ₃) δ : 1.76-2.10 (m, 4H), 3.30-3.52 (m, 2H), 4.72-4.90 (m, 2H), 5.97 (d, J = 13.9 Hz, 1H), 6.05 (d, J = 13.9 Hz, 1H), 6.54 (d, J = 2.3 Hz, 1H), 7.32 (d, J = 2.3 Hz, 2H), 7.33 (d, J = 2.3 Hz, 2H), 7.90-7.95 (m, 1H), 8.29 (s, 1H). LC/MS: cond. 2 RT 1.78 min LC/MS(ESI ⁺) m/z; 455 [M+H] ⁺
91 b	Morphology: colorless amorphous ¹ H-NMR (CDCl ₃) δ : 1.70-2.10 (m, 4H), 3.30-3.60 (m, 2H), 4.70-4.90 (m, 2H), 5.48 (d, J = 13.9 Hz, 1H), 6.06 (d, J = 13.9 Hz, 1H), 6.54 (d, J = 2.0 Hz, 1H), 7.10-7.50 (m, 5H), 7.90-8.00 (m, 1H), 8.28 (s, 1H) LC/MS: cond. 2 RT 1.66 min LC/MS(ESI ⁺) m/z; 421 [M+H] ⁺
92 a	Morphology: colorless solid
92 b	Morphology: colorless solid

[Table 8]

Rf	Data
93	Morphology: yellow oil LC/MS: cond. 2 RT 1.90 min LC/MS (ESI ⁺) m/z; 258 [M+H] ⁺
94	Morphology: brown oil 1H-NMR (CDCl ₃) δ : 4.61 (s, 2H), 7.19 (d, J = 8.6 Hz, 1H), 7.94 (dd, J = 8.2, 2.5 Hz, 1H), 8.39 (d, J = 2.4 Hz, 1H).
95	Morphology: yellow solid 1H-NMR (CDCl ₃) δ : 7.30 (s, 1H), 7.71 (d, J = 8.9 Hz, 1H), 8.02 (dd, J = 8.6, 1.5 Hz, 1H), 8.21 – 8.24 (m, 1H), 10.09 (s, 1H).
96	Morphology: pale yellow solid 1H-NMR (CDCl ₃) δ : 1.50 – 2.20 (brs, 1H), 4.81 (s, 2H), 7.16 (s, 1H), 7.42 – 7.48 (m, 1H), 7.56 (d, J = 8.4 Hz, 1H), 7.68 (s, 1H).
97	Morphology: pale yellow oil 1H-NMR (CDCl ₃) δ : 4.70 (s, 2H), 7.13 – 7.20 (m, 1H), 7.47 (dd, J = 8.6, 1.5 Hz, 1H), 7.56 (d, J = 8.6 Hz, 1H), 7.69 (d, J = 1.5 Hz, 1H).
98	Morphology: colorless solid
99	Morphology: yellow oil
100	Morphology: colorless oil 1H-NMR (CDCl ₃) δ : 1.75–1.96 (m, 1H), 2.15 (s, 3H), 4.69 (d, J = 5.7 Hz, 2H), 6.69 (s, 1H).
101	Morphology: brown oil
102	Morphology: black oil 1H-NMR (CDCl ₃) δ : 2.01 (t, J = 6.2 Hz, 1H), 4.84 (d, J = 6.0 Hz, 2H), 6.96 (s, 1H).
103	Morphology: brown oil
104	Morphology: yellow oil
105	Morphology: brown oil
106	Morphology: brown oil
107	Morphology: colorless oil
108	Morphology: colorless solid 1H-NMR (CDCl ₃) δ : 1.67 (t, J = 5.9 Hz, 1H), 4.36 (q, J = 8.1 Hz, 2H), 4.68 (d, J = 5.6 Hz, 2H), 6.21 (d, J = 3.6 Hz, 1H), 6.64 (d, J = 3.6 Hz, 1H).
109	Morphology: brown oil
110	Morphology: brown oil LC/MS: cond. 2 RT 1.52 min LC/MS (ESI ⁺) m/z; 209 [M+H] ⁺
111	Morphology: yellow oil LC/MS: cond. 2 RT 2.20 min LC/MS (ESI ⁺) m/z; 227 [M+H] ⁺
112	Morphology: yellow oil LC/MS: cond. 2 RT 2.27 min LC/MS (ESI ⁺) m/z; 242 [M+H] ⁺
113	Morphology: yellow oil LC/MS: cond. 1 RT 2.95 min LC/MS (ESI ⁺) m/z; 214 [M+H] ⁺
114	Morphology: yellow oil 1H-NMR (CDCl ₃) δ : 4.67 (s, 2H), 4.80 (q, J = 3.0 Hz, 2H), 7.08 (s, 1H).
115	Morphology: yellow solid LC/MS: cond. 2 RT 2.00 min LC/MS (ESI ⁺) m/z; 250 [M+H] ⁺
116	Morphology: yellow oil LC/MS: cond. 2 RT 0.85 min LC/MS (ESI ⁺) m/z; 222 [M+H] ⁺
117	Morphology: yellow oil LC/MS: cond. 2 RT 2.03 min LC/MS (ESI ⁺) m/z; 240 [M+H] ⁺
118	Morphology: yellow oil 1H-NMR (CDCl ₃) δ : 2.03 (t, J = 18.5 Hz, 3H), 6.51 (d, J = 2.4 Hz, 1H), 7.59 (d, J = 2.4 Hz, 1H).
119	Morphology: yellow oil

[Table 9]

Rf	Data
120	Morphology: brown oil
121	Morphology: brown oil
122	Morphology: pale brown oil
123	Morphology: pale brown oil
124	Morphology: colorless oil
125	Morphology: pale yellow oil
126	Morphology: pale yellow oil
127	¹ H-NMR (CDCl ₃) δ : 4.59(s, 2H), 7.26–7.29(m, 2H), 7.47–7.50(m, 2H).
128	Morphology: colorless liquid LC/MS: cond. 2 RT 3.02 min LC/MS (ESI ⁺) m/z; 320 [M+H] ⁺
129	Morphology: white amorphous ¹ H-NMR (CDCl ₃) δ : 1.48(s, 9H), 1.82(brs, 2H), 2.79(brs, 1H), 2.94–3.14(m, 2H), 3.89–4.19(m, 3H), 7.06(m, 2H), 7.45(m, 2H). LC/MS: cond. 1 RT 3.77 min LC/MS (ESI ⁻) m/z; 356[M+HCO ₂] ⁻
130	Morphology: colorless amorphous ¹ H-NMR (CDCl ₃) δ : 0.96(s, 9H), 1.49 (s, 9H), 1.84–1.89(m, 2H), 2.27(brs, 1H), 3.05–3.23(m, 2H), 3.98(brs, 1H), 4.14(brs, 1H), 5.24(brs, 1H), 6.98–7.04(m, 2H), 7.36–7.43(m, 2H). LC/MS: cond. 2 RT 2.74 min LC/MS (ESI ⁺) m/z; 413 [M+NH ₄] ⁺
131	Morphology: colorless oil ¹ H-NMR (CDCl ₃) δ : 1.08(s, 9H), 1.49 (s, 9H), 3.31(brs, 1H), 3.81(brs, 1H), 4.19(brs, 1H), 4.50(brs, 1H), 5.75(s, 1H), 6.23(brs, 1H), 7.00(t, 2H), 7.28(t, 2H). LC/MS: cond. 2 RT 3.06 min LC/MS (ESI ⁺) m/z; 278 [M+H-Boc] ⁺
132	¹ H-NMR (CDCl ₃) δ : 1.05 (s, 9H), 3.07–3.21 (m, 2H), 3.42–3.61 (m, 2H), 5.69 (brs, 1H), 6.32 (dd, J = 4.2, 3.0 Hz, 1H), 6.99 (t, J = 9.0 Hz, 2H), 7.25–7.31 (m, 2H). LC/MS: cond. 2 RT 1.72 min LC/MS (ESI ⁺) m/z; 278 [M+H] ⁺
133	Morphology: pale yellow amorphous ¹ H-NMR (CDCl ₃) δ : 3.24 (dd, J = 12.9, 2.7 Hz, 1H), 3.42 (dd, J = 12.6, 2.7 Hz, 1H), 3.52 (d, J = 3.9 Hz, 2H), 4.40(s, 1H), 6.16 (t, J = 3.6 Hz, 1H), 7.04 (t, J = 8.7 Hz, 2H), 7.48–7.53 (m, 2H). LC/MS: cond. 2 RT 0.37 min LC/MS (ESI ⁺) m/z; 194 [M+H] ⁺
134	Morphology: white amorphous ¹ H-NMR (CDCl ₃) δ : 1.48(s, 9H), 1.82(brs, 2H), 2.79(brs, 1H), 2.94–3.14(m, 2H), 3.89–4.19(m, 3H), 7.06(t, 2H), 7.45(t, 2H). LC/MS: cond. 1 RT 3.77 min LC/MS (ESI ⁻) m/z; 356[M+HCO ₂] ⁻
135	Morphology: colorless amorphous ¹ H-NMR (CDCl ₃) δ : 0.96(s, 9H), 1.49 (s, 9H), 1.84–1.89(m, 2H), 2.27(brs, 1H), 3.05–3.23(m, 2H), 3.98(brs, 1H), 4.14(brs, 1H), 5.24(brs, 1H), 6.98–7.04(m, 2H), 7.36–7.43(m, 2H). LC/MS: cond. 2 RT 2.74 min LC/MS (ESI ⁺) m/z; 413 [M+NH ₄] ⁺
136	Morphology: colorless amorphous ¹ H-NMR (CDCl ₃) δ : 1.08(s, 9H), 1.49 (s, 9H), 3.31(brs, 1H), 3.81(brs, 1H), 4.19(brs, 1H), 4.50(brs, 1H), 5.75(s, 1H), 6.23(br, 1H), 7.00(t, 2H), 7.28(t, 2H). LC/MS: cond. 2 RT 3.06 min LC/MS (ESI ⁺) m/z; 278 [M+H-Boc] ⁺
137	¹ H-NMR (CDCl ₃) δ : 1.05 (s, 9H), 3.07–3.21 (m, 2H), 3.42–3.61 (m, 2H), 5.69 (brs, 1H), 6.32 (dd, J = 4.2, 3.0 Hz, 1H), 6.99 (t, J = 9.0 Hz, 2H), 7.25–7.31 (m, 2H). LC/MS: cond. 2 RT 1.72 min LC/MS (ESI ⁺) m/z; 278 [M+H] ⁺
138	Morphology: pale yellow amorphous ¹ H-NMR (CDCl ₃) δ : 3.24 (dd, J = 12.9, 2.7 Hz, 1H), 3.42 (dd, J = 12.6, 2.7 Hz, 1H), 3.52 (d, J = 3.9 Hz, 2H), 4.40(s, 1H), 6.16 (t, J = 3.6 Hz, 1H), 7.04 (t, J = 8.7 Hz, 2H), 7.48–7.53 (m, 2H). LC/MS: cond. 2 RT 0.37 min LC/MS (ESI ⁺) m/z; 194 [M+H] ⁺

[Table 10]

Rf	Data
139	Morphology: pale yellow oil LC/MS: cond. 2 RT 3.01 min LC/MS (ESI ⁺) m/z; 325 [M+H] ⁺
140	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.31 min LC/MS (ESI ⁺) m/z; 359 [M+H] ⁺
141	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.90 min LC/MS (ESI ⁺) m/z; 443 [M+H] ⁺
142	Morphology: colorless oil LC/MS: cond. 2 RT 3.18 min LC/MS (ESI ⁺) m/z; 425 [M+H] ⁺
143	Morphology: pale red oil
144	Morphology: yellow solid LC/MS: cond. 2 RT 1.34 min LC/MS (ESI ⁺) m/z; 241 [M+H] ⁺
145	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.72 min LC/MS (ESI ⁺) m/z; 293 [M+H] ⁺
146	Morphology: colorless oil LC/MS: cond. 2 RT 2.10 min LC/MS (ESI ⁺) m/z; 327 [M+H] ⁺
147	Morphology: colorless oil LC/MS: cond. 2 RT 2.88 min LC/MS (ESI ⁺) m/z; 411 [M+H] ⁺
148	Morphology: colorless solid LC/MS: cond. 2 RT 3.03 min LC/MS (ESI ⁺) m/z; 393 [M+H] ⁺
149	Morphology: pale yellow oil
150	Morphology: pale yellow solid LC/MS: cond. 2 RT 0.81 min LC/MS (ESI ⁺) m/z; 209 [M+H] ⁺
151	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.11 min LC/MS (ESI ⁺) m/z; 329 [M+H] ⁺
152	Morphology: colorless amorphous
153	Morphology: colorless amorphous LC/MS: cond. 2 RT 3.01 min LC/MS (ESI ⁺) m/z; 395 [M+H] ⁺
154	Morphology: orange oil LC/MS: cond. 2 RT 1.76 min LC/MS (ESI ⁺) m/z; 295 [M+H] ⁺
155	Morphology: yellow oil LC/MS: cond. 2 RT 0.72 min LC/MS (ESI ⁺) m/z; 211 [M+H] ⁺
156	Morphology: yellow oil LC/MS: cond. 2 RT 2.54 min LC/MS (ESI ⁺) m/z; 393 [M+H] ⁺
157	Morphology: colorless oil LC/MS: cond. 2 RT 2.55 min LC/MS (ESI ⁺) m/z; 375 [M+H] ⁺
158	Morphology: colorless solid LC/MS: cond. 2 RT 1.52 min LC/MS (ESI ⁺) m/z; 275 [M+H] ⁺
159	Morphology: yellow solid LC/MS: cond. 2 RT 0.35 min LC/MS (ESI ⁺) m/z; 191 [M+H] ⁺
160	Morphology: colorless oil
161	Morphology: white amorphous LC/MS: cond. 2 RT 2.30 min LC/MS (ESI ⁺) m/z; 326 [M+H] ⁺
162	Morphology: colorless amorphous LC/MS: cond. 1 RT 4.82 min LC/MS (ESI ⁺) m/z; 310 [M+H-Boc] ⁺
163	Morphology: colorless oil LC/MS: cond. 1 RT 5.12 min LC/MS (ESI ⁺) m/z; 392 [M+H] ⁺
164	Morphology: pale yellow oil
165	Morphology: colorless oil 1H-NMR (CDCl ₃) δ : 2.29(s, 3H), 3.03-3.16(m, 2H), 3.46(s, 2H), 4.12(s, 1H), 5.72(s, 2H), 6.83-6.89(m, 2H), 7.11(m, 1H).

[Table 11]

Rf	Data
166	Morphology: colorless oil LC/MS: cond. 2 RT 2.80 min LC/MS (ESI ⁺) m/z; 291 [M+H] ⁺
167	Morphology: white amorphous LC/MS: cond. 2 RT 2.07 min LC/MS (ESI ⁺) m/z; 325 [M+H] ⁺
168	Morphology: colorless oil LC/MS: cond. 2 RT 2.78 min LC/MS (ESI ⁺) m/z; 409 [M+H] ⁺
169	Morphology: colorless oil LC/MS: cond. 2 RT 3.02 min LC/MS (ESI ⁺) m/z; 391 [M+H] ⁺
171	Morphology: brown oil LC/MS: cond. 2 RT 0.68 min LC/MS (ESI ⁺) m/z; 207 [M+H] ⁺
172	Morphology: colorless oil 1H-NMR (CDCl ₃) δ : 1.55(s, 9H), 1.57(s, 3H), 2.48(brs, 2H), 3.62(t, J = 5, 79 Hz, 2H), 4.06(s, 2H), 6.02(brs, 1H), 6.98-7.17(m, 3H).
173	Morphology: white amorphous LC/MS: cond. 2 RT 2.27 min LC/MS (ESI ⁺) m/z; 326 [M+H] ⁺
174	Morphology: white amorphous LC/MS: cond. 2 RT 2.87 min LC/MS (ESI ⁺) m/z; 310 [M+H-Boc] ⁺
175	Morphology: colorless oil LC/MS: cond. 2 RT 3.19 min LC/MS (ESI ⁺) m/z; 292 [M+H-Boc] ⁺
176	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.01 min LC/MS (ESI ⁺) m/z; 292 [M+H] ⁺
177	Morphology: colorless oil LC/MS: cond. 2 RT 1.25 min LC/MS (ESI ⁺) m/z; 208 [M+H] ⁺
178	Morphology: colorless oil 1H-NMR (CDCl ₃) δ : 1.49(s, 9H), 2.47(brs, 2H), 3.63(t, J = 5, 76 Hz, 2H), 4.06(brs, 2H), 6.00 (brs, 1H), 7.09(t, J = 8.64 Hz, 1H), 7.21-7.26(m, 1H), 7.39(d, 1H)
179	Morphology: white amorphous LC/MS: cond. 2 RT 2.31 min LC/MS (ESI ⁺) m/z; 246 [M+H-Boc] ⁺
180	Morphology: white amorphous LC/MS: cond. 2 RT 2.89 min LC/MS (ESI ⁺) m/z; 330 [M+H-Boc] ⁺
181	Morphology: colorless oil LC/MS: cond. 2 RT 3.21 min LC/MS (ESI ⁺) m/z; 312 [M+H-Boc] ⁺
182	Morphology: pale brown oil LC/MS: cond. 2 RT 2.03 min LC/MS (ESI ⁺) m/z; 312 [M+H] ⁺
183	Morphology: pale yellow solid LC/MS: cond. 2 RT 1.30 min LC/MS (ESI ⁺) m/z; 228 [M+H] ⁺
184	Morphology: colorless oil
185	Morphology: pale brown oil LC/MS: cond. 1 RT 3.70 min LC/MS (ESI ⁺) m/z; 325 [M+H] ⁺
186	Morphology: colorless oil LC/MS: cond. 2 RT 2.78 min LC/MS (ESI ⁺) m/z; 409 [M+H] ⁺
187	Morphology: colorless oil LC/MS: cond. 1 RT 4.87 min LC/MS (ESI ⁺) m/z; 391 [M+H] ⁺
188	Morphology: brown oil
189	Morphology: brown oil 1H-NMR (CDCl ₃) δ : 2.99-3.31(m, 2H), 3.51-3.63(m, 2H), 3.92(s, 3H), 4.55(brs, 1H), 6.61-6.66(m, 2H), 7.09(d, J = 7.5 Hz, 1H), 7.57(t, J = 7.8 Hz, 1H)
190	Morphology: colorless oil
191	Morphology: colorless oil
192	Morphology: colorless oil LC/MS: cond. 2 RT 2.81 min LC/MS (ESI ⁺) m/z; 380 [M+H] ⁺

[Table 12]

Rf	Data
193	Morphology: brown oil LC/MS: cond. 2 RT 1.92 min LC/MS (ESI ⁺) m/z; 280 [M+H] ⁺
194	Morphology: brown oil LC/MS: cond. 2 RT 1.05 min LC/MS (ESI ⁺) m/z; 196 [M+H] ⁺
195	Morphology: yellow oil LC/MS: cond. 2 RT 2.91 min LC/MS (ESI ⁺) m/z; 280 [M+H] ⁺
196	Morphology: yellow oil LC/MS: cond. 2 RT 2.12 min LC/MS (ESI ⁺) m/z; 314 [M+H] ⁺
197	Morphology: yellow oil
198	Morphology: yellow oil LC/MS: cond. 2 RT 3.14 min LC/MS (ESI ⁺) m/z; 380 [M+H] ⁺
199	Morphology: yellow oil LC/MS: cond. 2 RT 1.99 min LC/MS (ESI ⁺) m/z; 280 [M+H] ⁺
200	Morphology: pale yellow solid LC/MS: cond. 2 RT 1.08 min LC/MS (ESI ⁺) m/z; 196 [M+H] ⁺
201	Morphology: colorless oil LC/MS: cond. 2 RT 2.77 min LC/MS (ESI ⁺) m/z; 423 [M+H] ⁺
202	Morphology: colorless oil LC/MS: cond. 2 RT 1.66 min LC/MS (ESI ⁺) m/z; 293 [M+H] ⁺
203	Morphology: colorless oil LC/MS: cond. 2 RT 2.30 min LC/MS (ESI ⁺) m/z; 371 [M+H] ⁺
204	Morphology: colorless solid LC/MS: cond. 2 RT 1.08 min LC/MS (ESI ⁺) m/z; 271 [M+H] ⁺
205	Morphology: colorless oil LC/MS: cond. 2 RT 1.69 min LC/MS (ESI ⁺) m/z; 323 [M+H] ⁺
206	Morphology: pale yellow oil LC/MS: cond. 2 RT 0.36 min LC/MS (ESI ⁺) m/z; 193 [M+H] ⁺
207	Morphology: colorless solid LC/MS: cond. 2 RT 2.12 min LC/MS (ESI ⁺) m/z; 335 [M+H] ⁺
208	Morphology: colorless oil LC/MS: cond. 2 RT 1.02 min LC/MS (ESI ⁺) m/z; 235 [M+H] ⁺
209	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.77 min LC/MS (ESI ⁺) m/z; 423 [M+H] ⁺
210	Morphology: colorless oil LC/MS: cond. 2 RT 1.68 min LC/MS (ESI ⁺) m/z; 323 [M+H] ⁺
211	Morphology: colorless oil LC/MS: cond. 2 RT 0.38 min LC/MS (ESI ⁺) m/z; 193 [M+H] ⁺
212	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.77 min LC/MS (ESI ⁺) m/z; 423 [M+H] ⁺
213	Morphology: colorless oil LC/MS: cond. 2 RT 1.70 min LC/MS (ESI ⁺) m/z; 323 [M+H] ⁺
214	Morphology: colorless oil LC/MS: cond. 2 RT 0.39 min LC/MS (ESI ⁺) m/z; 193 [M+H] ⁺
215	Morphology: black oil LC/MS: cond. 2 RT 1.13 min LC/MS (ESI ⁺) m/z; 210 [M+H] ⁺
216	Morphology: dark brown oil LC/MS: cond. 2 RT 0.39 min LC/MS (ESI ⁺) m/z; 209 [M+H] ⁺
217	LC/MS: cond. 2 RT 1.40 min LC/MS (ESI ⁺) m/z; 195 [M+H] ⁺
218	Morphology: black oil LC/MS: cond. 2 RT 1.13 min LC/MS (ESI ⁺) m/z; 210 [M+H] ⁺

[Table 13]

Rf	Data
219	Morphology: dark brown oil LC/MS: cond. 2 RT 0.39 min LC/MS (ESI ⁺) m/z; 209 [M+H] ⁺
220	LC/MS: cond. 2 RT 1.40 min LC/MS (ESI ⁺) m/z; 195 [M+H] ⁺
221	Morphology: pale brown oil
222	Morphology: colorless solid LC/MS: cond. 2 RT 2.74 min LC/MS (ESI ⁺) m/z; 308 [M+H] ⁺
223	Morphology: pale brown solid
224	Morphology: pale brown solid
225	Morphology: yellow solid LC/MS: cond. 2 RT 1.84 min LC/MS (ESI ⁺) m/z; 297 [M+H] ⁺
226	Morphology: yellow oil LC/MS: cond. 2 RT 1.34 min LC/MS (ESI ⁺) m/z; 207 [M+H] ⁺
227	Morphology: colorless oil 1H-NMR (CDCl ₃) δ : 1.94-2.05(m, 1H), 2.11(s, 3H), 2.40(q, J = 5.4 Hz, 1H), 3.92(dd, J = 12.0, 10.5 Hz, 1H), 4.34(q, J = 5.1 Hz, 1H), 5.03(q, J = 5.4 Hz, 1H), 5.37-5.40(m, 1H), 7.00-7.05(m, 2H), 7.28-7.33(m, 2H).
228	Morphology: colorless oil 1H-NMR (CDCl ₃) δ : 1.90-1.97(m, 1H), 2.03(s, 3H), 2.68-2.77(m, 1H), 3.95(dd, J = 11.1, 5.1 Hz, 1H), 4.13(d, J = 10.8 Hz, 1H), 4.84(t, J = 7.8 Hz, 1H), 5.33-5.36(m, 1H), 7.00-7.06(m, 2H), 7.33-7.37(m, 2H).
229	Morphology: colorless solid LC/MS: cond. 1 RT 4.14 min LC/MS (ESI ⁺) m/z; 312 [M+H] ⁺
230	Morphology: colorless oil 1H-NMR (CDCl ₃) δ : 1.56-1.66(m, 1H), 2.59-2.68(m, 1H), 3.70-3.79(m, 2H), 3.99-4.06(m, 1H), 4.88(t, J = 7.8 Hz, 1H), 6.99-7.07(m, 2H), 7.31-7.38(m, 2H).
231	Morphology: colorless solid LC/MS: cond. 1 RT 4.14 min LC/MS (ESI ⁺) m/z; 312 [M+H] ⁺
232	Morphology: colorless oil 1H-NMR (CDCl ₃) δ : 1.94-2.14(m, 2H), 3.59-3.68(m, 1H), 3.70-3.79(m, 1H), 4.20-4.28(m, 1H), 5.08-5.15(m, 1H), 6.99-7.07(m, 2H), 7.31-7.38(m, 2H).
233	Morphology: colorless solid LC/MS: cond. 1 RT 4.29 min LC/MS (ESI ⁺) m/z; 326 [M+H] ⁺
234	Morphology: colorless oil 1H-NMR (CDCl ₃) δ : 1.41-1.48(m, 1H), 1.64-1.71(m, 1H), 1.86-2.02(m, 2H), 3.48(t, J = 3.6 Hz, 1H), 3.84-3.90(m, 1H), 4.00-4.09(m, 1H), 4.82(dd, J = 10.8, 2.7 Hz, 1H), 6.99-7.04(m, 2H), 7.29-7.35(m, 2H).
235	Morphology: colorless solid LC/MS: cond. 1 RT 2.93 min LC/MS (ESI ⁺) m/z; 238 [M+H] ⁺
236	Morphology: colorless solid 1H-NMR (CDCl ₃) δ : 1.25-1.40(m, 1H), 1.45-1.58(m, 1H), 1.47(s, 9H), 1.93-1.98(m, 1H), 2.20(brs, 1H), 3.59-3.68(m, 1H), 3.81(brs, 1H), 4.13-4.18(m, 1H), 4.33-4.36(m, 1H), 4.45(brs, 1H), 7.00-7.34(m, 2H), 7.25-7.31(m, 2H).
237	Morphology: colorless oil LC/MS: cond. 1 RT 0.37 min LC/MS (ESI ⁺) m/z; 196 [M+H] ⁺
238	Morphology: colorless oil LC/MS: cond. 1 RT 4.10 min LC/MS (ESI ⁺) m/z; 282 [M+H] ⁺
239	Morphology: colorless oil LC/MS: cond. 1 RT 2.45 min LC/MS (ESI ⁺) m/z; 268 [M+H] ⁺
240	Morphology: colorless oil LC/MS: cond. 1 RT 0.40 min LC/MS (ESI ⁺) m/z; 178 [M+H] ⁺
241	Morphology: colorless oil LC/MS: cond. 1 RT 3.04 min LC/MS (ESI ⁺) m/z; 415 [M+H] ⁺

[Table 14]

Rf	Data
242	Morphology: colorless solid LC/MS: cond. 1 RT 4.55 min LC/MS (ESI ⁺) m/z; 425 [M+H] ⁺
243	Morphology: colorless oil 1H-NMR (CDCl ₃) δ : 1.47(s, 9H), 2.27-2.38(m, 1H), 2.64-2.76(m, 1H), 2.75-2.81(m, 1H), 2.93-3.05(m, 1H), 3.12-3.19(m, 1H), 3.23-3.36(m, 1H), 3.75-3.88(m, 2H), 6.99-7.06(m, 2H), 7.18-7.23(m, 2H).
244	Morphology: brown oil LC/MS: cond. 2 RT 1.91 min LC/MS (ESI ⁺) m/z; 191 [M+H] ⁺
245	Morphology: brown solid LC/MS: cond. 2 RT 2.37 min LC/MS (ESI ⁻) m/z; 205 [M-H] ⁻
246	Morphology: brown oil
247	Morphology: pale yellow oil
248	Morphology: pale yellow solid 1H-NMR (CDCl ₃) δ : 6.90-6.91 (m, 2H), 7.18-7.24(m, 2H), 7.38-7.42 (m, 2H), 7.81-7.83(m, 1H).
249	Morphology: purple oil LC/MS: cond. 2 RT 1.13 min LC/MS (ESI ⁺) m/z; 177 [M+H] ⁺
250	Morphology: colorless solid 1H-NMR (CDCl ₃) δ : 2.64-2.73(m, 2H), 3.04-3.12(m, 2H), 3.16-3.25m, 1H), 3.73-3.83(m, 1H), 6.98-7.06(m, 2H), 7.40-7.47(m, 2H), 7.70-7.75(m, 2H), 7.82-7.87(m, 2H).
251	Morphology: colorless oil
252	Morphology: colorless solid LC/MS: cond. 1 RT 4.54 min LC/MS (ESI ⁺) m/z; 296 [M+H] ⁺
253	Morphology: colorless oil
254	Morphology: colorless oil LC/MS: cond. 2 RT 1.78 min LC/MS (ESI ⁺) m/z; 225 [M+H] ⁺
255	Morphology: colorless solid LC/MS: cond. 1 RT 4.35 min LC/MS (ESI ⁺) m/z; 354 [M+H] ⁺
256	Morphology: colorless oil LC/MS: cond. 1 RT 3.94 min LC/MS (ESI ⁺) m/z; 340 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 338 [M-H] ⁻
257	Morphology: colorless solid LC/MS: cond. 1 RT 3.70 min LC/MS (ESI ⁺) m/z; 339 [M+H] ⁺
258	Morphology: colorless solid 1H-NMR (CDCl ₃) δ : 3.08-3.17(m, 2H), 3.54-3.63(m, 2H), 5.21-5.33(m, 1H), 7.12-7.19(m, 2H), 7.69-7.79(m, 4H), 7.85-7.90(m, 2H).
259	Morphology: colorless solid LC/MS: cond. 2 RT 1.21 min LC/MS (ESI ⁺) m/z; 191 [M+H] ⁺
260	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.76 min LC/MS (ESI ⁺) m/z; 331 [M+H] ⁺
261	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.76 min LC/MS (ESI ⁺) m/z; 313 [M+H] ⁺
262	Morphology: brown amorphous LC/MS: cond. 2 RT 1.45 min LC/MS (ESI ⁺) m/z; 231 [M+H] ⁺
263	Morphology: brown amorphous LC/MS: cond. 2 RT 1.35 min LC/MS (ESI ⁺) m/z; 213 [M+H] ⁺
264	Morphology: yellow oil LC/MS: cond. 2 RT 2.81 min LC/MS (ESI ⁺) m/z; 307 [M+H] ⁺
265	Morphology: brown amorphous LC/MS: cond. 2 RT 1.36 min LC/MS (ESI ⁺) m/z; 207 [M+H] ⁺

[Table 15]

Rf	Data
266	Morphology: yellow solid LC/MS: cond. 2 RT 1.09 min LC/MS (ESI ⁺) m/z; 191 [M+H] ⁺
267	Morphology: pale yellow solid LC/MS: cond. 2 RT 1.52 min LC/MS (ESI ⁺) m/z; 225 [M+H] ⁺
268	Morphology: colorless solid LC/MS: cond. 2 RT 1.52 min LC/MS (ESI ⁺) m/z; 192 [M+H] ⁺
269	Morphology: colorless solid LC/MS: cond. 2 RT 1.54 min LC/MS (ESI ⁺) m/z; 212 [M+H] ⁺
270	Morphology: brown oil LC/MS: cond. 2 RT 1.35 min LC/MS (ESI ⁺) m/z; 180 [M+H] ⁺
271	Morphology: colorless oil LC/MS: cond. 2 RT 3.65 min LC/MS (ESI ⁺) m/z; 322 [M+H-Boc] ⁺
272	Morphology: colorless liquid LC/MS: cond. 2 RT 0.98 min LC/MS (ESI ⁺) m/z; 208 [M+H] ⁺
273	Morphology: pale brown oil 1H-NMR (CDCl ₃) δ : 1.40(s, 3H), 1.44(s, 9H), 3.13(brs, 1H), 3.73(s, 3H), 3.90(dd, J = 18.9, 3.3 Hz, 1H), 4.26-4.32(m, 2H), 5.87(s, 1H)
274	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.79 min LC/MS (ESI ⁺) m/z; 350 [M+H] ⁺
275	Morphology: colorless oil 1H-NMR (CDCl ₃) δ : 1.20(s, 3H), 1.46(s, 9H), 3.13(d, J = 12 Hz, 1H), 3.90(dd, J = 19.2, 2.7 Hz, 1H), 4.23-4.37(m, 2H), 5.80(s, 1H), 6.88-6.98(m, 2H), 7.18-7.25(m, 2H)
276	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.56 min LC/MS (ESI ⁺) m/z; 365 [M+H] ⁺
277	Morphology: colorless oil LC/MS: cond. 2 RT 1.42 min LC/MS (ESI ⁺) m/z; 265 [M+H] ⁺
278	Morphology: colorless oil LC/MS: cond. 2 RT 0.53 min LC/MS (ESI ⁺) m/z; 207 [M+H] ⁺
279	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.48 min LC/MS (ESI ⁺) m/z; 310 [M+H] ⁺
280	LC/MS: cond. 2 RT 1.33 min LC/MS (ESI ⁺) m/z; 192 [M+H] ⁺
281	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.85, 2.97 min LC/MS (ESI ⁺) m/z; 312 [M+H] ⁺
282	Morphology: colorless oil
283	Morphology: colorless oil LC/MS: cond. 2 RT 2.53 min LC/MS (ESI ⁺) m/z; 310 [M+H] ⁺
284	LC/MS: cond. 2 RT 1.40 min LC/MS (ESI ⁺) m/z; 192 [M+H] ⁺
285	Morphology: pale yellow oil LC/MS: cond. 2 RT 1.04, 1.14 min LC/MS (ESI ⁺) m/z; 210 [M+H] ⁺
286	Morphology: yellow oil LC/MS: cond. 2 RT 1.10 min LC/MS (ESI ⁺) m/z; 183 [M+H] ⁺
287	Morphology: colorless solid LC/MS: cond. 2 RT 1.49 min LC/MS (ESI ⁺) m/z; 228 [M+H] ⁺
288	Morphology: yellow solid LC/MS: cond. 2 RT 1.15 min LC/MS (ESI ⁺) m/z; 196 [M+H] ⁺
289	Morphology: colorless solid 1H-NMR (DMSO-D ₆) δ : 2.49 (s, 3H), 7.02 (s, 1H), 7.60 (s, 1H), 8.28 (s, 1H), 9.26 (brs, 2H) LC/MS: cond. 2 RT 0.40 min LC/MS (ESI ⁺) m/z; 185 [M+H] ⁺

[Table 16]

Rf	Data
290	Morphology: colorless solid LC/MS: cond. 2 RT 1.68 min LC/MS (ESI ⁺) m/z; 298 [M+H] ⁺
291	Morphology: colorless solid 1H-NMR (CDCl ₃) δ : 2.43 (s, 3H), 5.22 (s, 2H), 7.32 (s, 1H), 7.61 (s, 1H), 8.77 (s, 1H). LC/MS: cond. 1 RT 3.72 min LC/MS (ESI ⁺) m/z; 308 [M+H] ⁺
292	Morphology: yellow solid LC/MS: cond. 1 RT 3.84 min LC/MS (ESI ⁺) m/z; 322 [M+H] ⁺
293	Morphology: colorless oil LC/MS: cond. 1 RT 2.41 min LC/MS (ESI ⁺) m/z; 336 [M+H] ⁺
294	Morphology: colorless oil LC/MS: cond. 1 RT 2.56 min LC/MS (ESI ⁺) m/z; 350 [M+H] ⁺
295	Morphology: colorless solid 1H-NMR (CDCl ₃) δ : 2.37(s, 3H), 4.47(d, J = 6.1 Hz, 2H), 5.68(brs, 1H), 6.73(d, J = 3.7 Hz, 1H), 6.88(d, J = 3.7 Hz, 1H).
296	Morphology: colorless solid LC/MS: cond. 2 RT 2.01 min LC/MS (ESI ⁺) m/z; 318 [M+H] ⁺
297	Morphology: pale yellow solid 1H-NMR (CDCl ₃) δ : 2.51(s, 3H), 2.55(s, 3H), 5.20(s, 2H), 6.87(d, J = 3.7 Hz, 1H), 6.92(d, J = 3.7 Hz, 1H).
298	Morphology: colorless oil LC/MS: cond. 2 RT 1.55 min LC/MS (ESI ⁺) m/z; 264 [M+H] ⁺
299	Morphology: colorless solid LC/MS: cond. 2 RT 1.97 min LC/MS (ESI ⁺) m/z; 274 [M+H] ⁺
300	Morphology: pale yellow solid 1H-NMR (CDCl ₃) δ : 2.51(s, 3H), 2.55(s, 3H), 5.19(s, 2H), 6.78(d, J=3.7 Hz, 1H), 6.89(d, J = 3.3 Hz, 1H).
301	Morphology: colorless oil LC/MS: cond. 2 RT 1.55 min LC/MS (ESI ⁺) m/z; 282 [M+H] ⁺
302	Morphology: colorless oil LC/MS: cond. 2 RT 2.04 min LC/MS (ESI ⁺) m/z; 306 [M+H] ⁺
303	Morphology: colorless solid
304	Morphology: colorless oil LC/MS: cond. 1 RT 3.09 min LC/MS (ESI ⁺) m/z; 286 [M+H] ⁺
305	Morphology: yellow solid
306	Morphology: colorless solid
307	Morphology: colorless solid
308	Morphology: colorless amorphous
309	Morphology: yellow oil LC/MS: cond. 2 RT 1.78 min LC/MS (ESI ⁺) m/z; 373 [M+H] ⁺
310	Morphology: colorless oil
311	Morphology: colorless solid
312	Morphology: colorless oil LC/MS: cond. 2 RT 1.92 min LC/MS (ESI ⁺) m/z; 312 [M+H] ⁺
313	Morphology: colorless oil LC/MS: cond. 2 RT 2.28 min LC/MS (ESI ⁺) m/z; 336 [M+H] ⁺
314	Morphology: colorless solid LC/MS: cond. 2 RT 1.73 min LC/MS (ESI ⁺) m/z; 264 [M+H] ⁺
315	Morphology: yellow amorphous LC/MS: cond. 2 RT 1.84 min LC/MS (ESI ⁺) m/z; 278 [M+H] ⁺
316	Morphology: colorless oil

[Table 17]

Rf	Data
317	Morphology: pale brown oil
318	Morphology: pale brown oil
319	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.11 min LC/MS (ESI ⁺) m/z; 348 [M+H] ⁺
320	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.41 min LC/MS (ESI ⁺) m/z; 372 [M+H] ⁺
321	LC/MS: cond. 2 RT 1.97 min LC/MS (ESI ⁺) m/z; 284 [M+H] ⁺
322	LC/MS: cond. 2 RT 2.30 min LC/MS (ESI ⁺) m/z; 294 [M+H] ⁺
323	LC/MS: cond. 2 RT 2.39 min LC/MS (ESI ⁺) m/z; 308 [M+H] ⁺
324	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.14 min LC/MS (ESI ⁺) m/z; 282 [M+H] ⁺
325	Morphology: orange solid LC/MS: cond. 2 RT 2.51 min LC/MS (ESI ⁺) m/z; 306 [M+H] ⁺
326	Morphology: colorless solid LC/MS: cond. 1 RT 1.09 min LC/MS (ESI ⁺) m/z; 258 [M+H] ⁺
327	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.39 min LC/MS (ESI ⁺) m/z; 358 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 356 [M-H] ⁻
328	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.14 min LC/MS (ESI ⁺) m/z; 340 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 338 [M-H] ⁻
329	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.46 min LC/MS (ESI ⁺) m/z; 398 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 396 [M-H] ⁻
330	Morphology: colorless solid LC/MS: cond. 2 RT 2.27 min LC/MS (ESI ⁺) m/z; 380 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 378 [M-H] ⁻
331	Morphology: colorless solid LC/MS: cond. 1 RT 4.05 min LC/MS (ESI ⁺) m/z; 338 [M+H] ⁺
332a	Morphology: colorless solid LC/MS: cond. 1 RT 3.70 min LC/MS (ESI ⁺) m/z; 307 [M+H] ⁺
332b	Morphology: colorless amorphous LC/MS: cond. 1 RT 3.67 min LC/MS (ESI ⁺) m/z; 307 [M+H] ⁺
333a	Morphology: colorless solid LC/MS: cond. 1 RT 2.95 min LC/MS (ESI ⁺) m/z; 289 [M+H] ⁺
333b	Morphology: colorless solid LC/MS: cond. 1 RT 2.77 min LC/MS (ESI ⁺) m/z; 289 [M+H] ⁺
334	Morphology: colorless solid LC/MS: cond. 1 RT 3.05 min LC/MS (ESI ⁺) m/z; 303 [M+H] ⁺
335	Morphology: white amorphous LC/MS: cond. 2 RT 2.08 min LC/MS (ESI ⁺) m/z; 307 [M+H] ⁺
336	Morphology: pale brown solid LC/MS: cond. 2 RT 1.51 min LC/MS (ESI ⁺) m/z; 289 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 287 [M-H] ⁻
337	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.36 min LC/MS (ESI ⁺) m/z; 294 [M+H] ⁺
338	Morphology: colorless solid LC/MS: cond. 2 RT 1.63 min LC/MS (ESI ⁺) m/z; 276 [M+H] ⁺
339	Morphology: colorless solid LC/MS: cond. 2 RT 1.73 min LC/MS (ESI ⁺) m/z; 286 [M+H] ⁺
340	Morphology: colorless solid LC/MS: cond. 2 RT 1.71 min LC/MS (ESI ⁺) m/z; 290 [M+H] ⁺
341	Morphology: colorless oil LC/MS: cond. 2 RT 2.66 min LC/MS (ESI ⁺) m/z; 291 [M+H] ⁺

[Table 18]

Rf	Data
342	Morphology: white solid LC/MS: cond. 2 RT 1.84 min LC/MS (ESI ⁺) m/z; 273 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 271 [M-H] ⁻
343	Morphology: colorless solid LC/MS: cond. 2 RT 1.93 min LC/MS (ESI ⁺) m/z; 287 [M+H] ⁺
344	Morphology: white solid LC/MS: cond. 2 RT 2.65 min LC/MS (ESI ⁺) m/z; 323 [M+H] ⁺
345	Morphology: white solid LC/MS: cond. 2 RT 1.76 min LC/MS (ESI ⁺) m/z; 305 [M+H] ⁺
346	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.64 min LC/MS (ESI ⁺) m/z; 469 [M+H] ⁺
347	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.55 min LC/MS (ESI ⁺) m/z; 429 [M+H] ⁺
348	Morphology: colorless amorphous LC/MS: cond. 2 RT 1.89 min LC/MS (ESI ⁺) m/z; 375 [M+H] ⁺
349	Morphology: colorless amorphous LC/MS: cond. 1 RT 3.79 min LC/MS (ESI ⁺) m/z; 453 [M+H] ⁺
350	Morphology: colorless solid LC/MS: cond. 2 RT 2.55 min LC/MS (ESI ⁺) m/z; 460 [M+H] ⁺
351	Morphology: colorless amorphous LC/MS: cond. 2 RT 1.55 min LC/MS (ESI ⁺) m/z; 360 [M+H] ⁺
352	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.42 min LC/MS (ESI ⁺) m/z; 393 [M+H] ⁺
353	Morphology: brown amorphous LC/MS: cond. 2 RT 2.60 min LC/MS (ESI ⁺) m/z; 471 [M+H] ⁺
354	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.30 min LC/MS (ESI ⁺) m/z; 294 [M+H] ⁺
355	Morphology: yellow amorphous
356	Morphology: colorless liquid LC/MS: cond. 2 RT 2.17 min LC/MS (ESI ⁺) m/z; 348 [M+H] ⁺
357	Morphology: colorless oil LC/MS: cond. 2 RT 2.82 min LC/MS (ESI ⁺) m/z; 480 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 478 [M-H] ⁻
358	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.88 min LC/MS (ESI ⁺) m/z; 426 [M+H] ⁺
359	Morphology: yellow oil LC/MS: cond. 2 RT 1.70 min LC/MS (ESI ⁺) m/z; 326 [M+H] ⁺
361	Morphology: yellow amorphous LC/MS: cond. 2 RT 1.72 min LC/MS (ESI ⁺) m/z; 328 [M+H] ⁺
362	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.82 min LC/MS (ESI ⁺) m/z; 601 [M+H] ⁺
363	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.60 min LC/MS (ESI ⁺) m/z; 525 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 523 [M-H] ⁻
364	LC/MS: cond. 2 RT 2.29 min LC/MS (ESI ⁺) m/z; 294 [M+H] ⁺ Enantiomeric excess: 99% ee, ChiralPak IA(3um 4.6x150mm), 1.5mL/min, Hex/EtOH = 99/1(v/v), 40°C, 11.1min (Rf 364), 14.1min(Rf 364)
365	LC/MS: cond. 2 RT 2.29 min LC/MS (ESI ⁺) m/z; 294 [M+H] ⁺ Enantiomeric excess: 98% ee, ChiralPak IA(3um 4.6x150mm), 1.5mL/min, Hex/EtOH = 99/1(v/v), 40°C, 11.1min (Rf 364), 14.1min(Rf 365)

[Table 19]

Rf	Data
366	Morphology: white solid
367	Morphology: white solid LC/MS: cond. 2 RT 2.01 min LC/MS (ESI ⁺) m/z; 289 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 287 [M-H] ⁻
368	Morphology: colorless solid LC/MS: cond. 2 RT 0.74, 0.88 min LC/MS (ESI ⁺) m/z; 328 [M+H] ⁺
369	Morphology: colorless solid LC/MS: cond. 2 RT 0.79 min LC/MS (ESI ⁺) m/z; 238 [M+H] ⁺
370	Morphology: colorless solid LC/MS: cond. 2 RT 0.62 min LC/MS (ESI ⁺) m/z; 196 [M+H] ⁺
371	LC/MS: cond. 2 RT 2.32 min LC/MS (ESI ⁺) m/z; 394 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 392 [M-H] ⁻
372	Morphology: colorless oil LC/MS: cond. 1 RT 4.37 min LC/MS (ESI ⁺) m/z; 376 [M+H] ⁺
373	Morphology: yellow oil LC/MS: cond. 2 RT 1.37 min LC/MS (ESI ⁺) m/z; 299 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 297 [M-H] ⁻
374	Morphology: orange oil LC/MS: cond. 2 RT 1.93 min LC/MS (ESI ⁺) m/z; 323 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 321 [M-H] ⁻

[Table 20]

Ex	Data
1	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 3.11 (m, 4H), 4.03 (m, 4H), 4.95 (s, 2H), 6.89 (m, 2H), 6.97 (m, 2H), 7.32 (m, 4H), 7.93 (s, 1H). LC/MS:cond.1 RT 4.40 min LC/MS (ESI ⁺) m/z; 400 [M+H] ⁺
2	Morphology: yellow solid LC/MS:cond.1 RT 4.05 min LC/MS (ESI ⁺) m/z; 411 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 409 [M-H] ⁻
3	Morphology: colorless solid LC/MS:cond.1 RT 4.44 min LC/MS (ESI ⁺) m/z; 434 [M+H] ⁺
4	Morphology: colorless solid LC/MS:cond.1 RT 4.22 min LC/MS (ESI ⁺) m/z; 380 [M+H] ⁺
5	Morphology: colorless solid LC/MS:cond.1 RT 4.72 min LC/MS (ESI ⁺) m/z; 422 [M+H] ⁺
6	Morphology: colorless solid LC/MS:cond.1 RT 4.49 min LC/MS (ESI ⁺) m/z; 450 [M+H] ⁺
7	Morphology: colorless solid LC/MS:cond.1 RT 4.29 min LC/MS (ESI ⁺) m/z; 400 [M+H] ⁺
8	Morphology: colorless solid LC/MS:cond.1 RT 4.39 min LC/MS (ESI ⁺) m/z; 434 [M+H] ⁺
9	Morphology: colorless solid LC/MS:cond.1 RT 4.26 min LC/MS (ESI ⁺) m/z; 380 [M+H] ⁺
10	Morphology: colorless solid LC/MS:cond.1 RT 4.14 min LC/MS (ESI ⁺) m/z; 384 [M+H] ⁺
11	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 1.28 (s, 9H), 3.13 (m, 4H), 3.84 (s, 3H), 4.02-4.07 (m, 4H), 4.99 (s, 2H), 6.11 (s, 1H), 6.86-7.01 (m, 4H), 7.86 (s, 1H) LC/MS:cond.1 RT 3.99 min LC/MS (ESI ⁺) m/z; 426 [M+H] ⁺
12	Morphology: colorless solid LC/MS:cond.1 RT 4.05 min LC/MS (ESI ⁺) m/z; 396 [M+H] ⁺
13	Morphology: pale yellow solid LC/MS:cond.1 RT 4.09 min LC/MS (ESI ⁺) m/z; 384 [M+H] ⁺
14	Morphology: colorless solid LC/MS:cond.1 RT 3.90 min LC/MS (ESI ⁺) m/z; 428 [M+H] ⁺
15	Morphology: colorless solid LC/MS:cond.1 RT 3.90 min LC/MS (ESI ⁺) m/z; 412 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 410 [M-H] ⁻
16	Morphology: colorless solid LC/MS:cond.1 RT 3.90 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
17	Morphology: pale yellow solid LC/MS:cond.2 RT 2.08 min LC/MS (ESI ⁺) m/z; 423 [M+H] ⁺
18	Morphology: pale yellow solid LC/MS:cond.2 RT 2.60 min LC/MS (ESI ⁺) m/z; 440 [M+H] ⁺
19	Morphology: colorless solid LC/MS:cond.1 RT 4.22 min LC/MS (ESI ⁺) m/z; 424 [M+H] ⁺
20	Morphology: colorless solid LC/MS:cond.1 RT 4.32 min LC/MS (ESI ⁺) m/z; 406 [M+H] ⁺
21	Morphology: colorless solid LC/MS:cond.1 RT 2.40 min LC/MS (ESI ⁺) m/z; 424 [M+H] ⁺
22	Morphology: colorless solid LC/MS:cond.1 RT 4.07 min LC/MS (ESI ⁺) m/z; 435 [M+H] ⁺

[Table 21]

Ex	Data
23	Morphology: colorless solid LC/MS:cond.1 RT 3.85 min LC/MS(ESI ⁺) m/z; 424 [M+H] ⁺ LC/MS(ESI ⁻) m/z; 422 [M-H] ⁻
24	Morphology: colorless solid LC/MS:cond.2 RT 2.72 min LC/MS(ESI ⁺) m/z; 500 [M+H] ⁺
25	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 3.09-3.30 (m, 4H), 4.00-4.09 (m, 4H), 5.12 (s, 2H), 6.84-7.02 (m, 4H), 7.10 (d, J = 3.7 Hz, 1H), 7.31 (d, J = 3.7 Hz, 1H), 8.00 (s, 1H). LC/MS:cond.2 RT 2.51 min LC/MS(ESI ⁺) m/z; 440 [M+H] ⁺
26	Morphology: colorless solid LC/MS:cond.1 RT 4.10 min LC/MS(ESI ⁺) m/z; 424 [M+H] ⁺
27	Morphology: colorless solid LC/MS:cond.1 RT 4.40 min LC/MS(ESI ⁺) m/z; 438 [M+H] ⁺
28	Morphology: pale brown solid LC/MS:cond.1 RT 4.16 min LC/MS(ESI ⁺) m/z; 465 [M+H] ⁺
29	Morphology: colorless solid LC/MS:cond.1 RT 4.07 min LC/MS(ESI ⁺) m/z; 441 [M+H] ⁺
30	Morphology: colorless solid LC/MS:cond.1 RT 4.02 min LC/MS(ESI ⁺) m/z; 441 [M+H] ⁺
31	Morphology: colorless solid LC/MS:cond.1 RT 4.47 min LC/MS(ESI ⁺) m/z; 430 [M+H] ⁺
32	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 3.11 (t, J = 5.0 Hz, 4H), 4.02-4.07 (m, 4H), 6.00 (s, 2H), 6.86-7.01 (m, 4H), 7.72 (s, 1H), 8.17 (s, 1H), 8.29 (s, 1H). LC/MS:cond.1 RT 4.39 min LC/MS(ESI ⁺) m/z; 424 [M+H] ⁺
33	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 3.07-3.13 (m, 4H), 3.95-4.04 (m, 4H), 4.29 (dd, J = 4.0 Hz, 6.0 Hz, 2H), 4.56 (dd, J = 4.0 Hz, 6.0 Hz, 2H), 6.50 (4.29 (d, J = 3.0 Hz, 1H), 6.86-7.01 (m, 4H), 7.37 (m, 1H), 7.42 (s, 1H). LC/MS:cond.2 RT 2.27 min LC/MS(ESI ⁺) m/z; 438 [M+H] ⁺
34	Morphology: colorless solid LC/MS:cond.1 RT 4.62 min LC/MS(ESI ⁺) m/z; 484 [M+H] ⁺
35	Morphology: colorless solid LC/MS:cond.1 RT 3.09 min LC/MS(ESI ⁺) m/z; 484 [M+H] ⁺
36	Morphology: colorless oil ¹ H-NMR (CDCl ₃) δ : 2.90 (s, 6H), 3.11 (m, 4H), 4.01 (m, 4H), 5.06 (s, 2H), 6.88-6.98 (m, 4H), 7.08 (m, 2H), 7.28 (m, 2H) LC/MS:cond.1 RT 4.14 min LC/MS(ESI ⁺) m/z; 443 [M+H] ⁺
37	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 3.16-3.25 (m, 4H), 3.79 (s, 3H), 4.00-4.07 (m, 4H), 6.02 (s, 2H), 6.44-6.50 (m, 2H), 6.50-6.57 (m, 2H), 7.15-7.25 (m, 1H), 7.93 (d, J = 1.6 Hz, 1H), 8.30 (s, 1H). LC/MS:cond.1 RT 4.04 min LC/MS(ESI ⁺) m/z; 436 [M+H] ⁺
38	Morphology: colorless amorphous LC/MS:cond.1 RT 4.30 min LC/MS(ESI ⁺) m/z; 440 [M+H] ⁺
39	Morphology: colorless solid LC/MS:cond.1 RT 4.18 min LC/MS(ESI ⁺) m/z; 420 [M+H] ⁺
40	Morphology: colorless solid LC/MS:cond.1 RT 4.35 min LC/MS(ESI ⁺) m/z; 446 [M+H] ⁺

[Table 22]

Ex	Data
41	Morphology: colorless solid LC/MS:cond.1 RT 4.40 min LC/MS (ESI ⁺) m/z; 474 [M+H] ⁺
42	Morphology: colorless solid LC/MS:cond.1 RT 4.39 min LC/MS (ESI ⁺) m/z; 474 [M+H] ⁺
43	Morphology: colorless solid LC/MS:cond.1 RT 4.60 min LC/MS (ESI ⁺) m/z; 506 [M+H] ⁺
44	Morphology: colorless solid LC/MS:cond.1 RT 4.15 min LC/MS (ESI ⁺) m/z; 420 [M+H] ⁺
45	Morphology: colorless solid LC/MS:cond.1 RT 4.43 min LC/MS (ESI ⁺) m/z; 490 [M+H] ⁺
46	Morphology: colorless solid LC/MS:cond.1 RT 4.32 min LC/MS (ESI ⁺) m/z; 434 [M+H] ⁺
47	Morphology: colorless solid LC/MS:cond.1 RT 4.34 min LC/MS (ESI ⁺) m/z; 522 [M+H] ⁺
48	Morphology: colorless solid LC/MS:cond.1 RT 4.27 min LC/MS (ESI ⁺) m/z; 438 [M+H] ⁺
49	Morphology: colorless solid LC/MS:cond.1 RT 3.94 min LC/MS (ESI ⁺) m/z; 431 [M+H] ⁺
50	Morphology: colorless solid LC/MS:cond.1 RT 4.07 min LC/MS (ESI ⁺) m/z; 464 [M+H] ⁺
51	Morphology: colorless solid LC/MS:cond.1 RT 4.07 min LC/MS (ESI ⁺) m/z; 449 [M+H] ⁺
52	Morphology: colorless solid LC/MS:cond.1 RT 4.43 min LC/MS (ESI ⁺) m/z; 492 [M+H] ⁺
53	Morphology: colorless solid LC/MS:cond.1 RT 4.34 min LC/MS (ESI ⁺) m/z; 458 [M+H] ⁺
54	Morphology: colorless solid LC/MS:cond.1 RT 3.99 min LC/MS (ESI ⁺) m/z; 431 [M+H] ⁺
55	Morphology: colorless solid LC/MS:cond.1 RT 4.32 min LC/MS (ESI ⁺) m/z; 440 [M+H] ⁺
56	Morphology: colorless solid LC/MS:cond.1 RT 4.57 min LC/MS (ESI ⁺) m/z; 508 [M+H] ⁺
57	Morphology: yellow solid LC/MS:cond.1 RT 4.10 min LC/MS (ESI ⁺) m/z; 469 [M+H] ⁺
58	Morphology: colorless solid LC/MS:cond.1 RT 4.34 min LC/MS (ESI ⁺) m/z; 460 [M+H] ⁺
59	Morphology: colorless solid LC/MS:cond.1 RT 4.35 min LC/MS (ESI ⁺) m/z; 420 [M+H] ⁺
60	Morphology: yellow solid LC/MS:cond.1 RT 4.40 min LC/MS (ESI ⁺) m/z; 458 [M+H] ⁺
61	Morphology: yellow solid LC/MS:cond.1 RT 4.34 min LC/MS (ESI ⁺) m/z; 512 [M+H] ⁺
62	Morphology: colorless solid LC/MS:cond.1 RT 4.18 min LC/MS (ESI ⁺) m/z; 465 [M+H] ⁺
63	Morphology: yellow solid LC/MS:cond.1 RT 4.25 min LC/MS (ESI ⁺) m/z; 465 [M+H] ⁺
64	Morphology: colorless solid LC/MS:cond.1 RT 4.20 min LC/MS (ESI ⁺) m/z; 442 [M+H] ⁺

[Table 23]

Ex	Data
65	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 0.94 (t, J = 7.4 Hz, 3H), 1.30–1.40 (m, 2H), 1.64–1.77 (m, 2H), 3.04–3.18 (m, 4H), 3.90–4.00 (m, 2H), 4.00–4.10 (m, 2H), 4.36 (t, J = 6.4 Hz, 2H), 5.04 (s, 2H), 6.86–7.04 (m, 4H), 7.24–7.36 (m, 4H). LC/MS:cond.1 RT 4.92 min LC/MS(ESI ⁺) m/z; 472 [M+H] ⁺
66st	LC/MS:cond.1 RT 4.34 min LC/MS(ESI ⁻) m/z; 520 [M-H] ⁻
66	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 3.10 (m, 4H), 4.01 (m, 7H), 5.00 (s, 2H), 5.02 (s, 2H), 6.87 (m, 4H), 7.34 (m, 6H), 7.97 (m, 2H). LC/MS:cond.1 RT 4.85 min LC/MS(ESI ⁺) m/z; 536 [M+H] ⁺
67	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 3.09 (m, 4H), 4.02 (m, 4H), 6.29 (s, 2H), 6.84–6.91 (m, 2H), 6.96 (m, 2H), 7.09 (m, 1H), 7.29 (m, 1H), 7.63 (m, 1H), 7.66 (m, 1H), 8.38 (s, 1H), 8.45 (s, 1H) LC/MS:cond.2 RT 2.19 min LC/MS(ESI ⁺) m/z; 406 [M+H] ⁺
68	Morphology: colorless solid LC/MS:cond.1 RT 4.43 min LC/MS(ESI ⁺) m/z; 437 [M+H] ⁺
69	Morphology: colorless solid LC/MS:cond.1 RT 4.24 min LC/MS(ESI ⁺) m/z; 403 [M+H] ⁺
70	Morphology: colorless solid LC/MS:cond.2 RT 2.01 min LC/MS(ESI ⁺) m/z; 419 [M+H] ⁺
71	Morphology: colorless solid LC/MS:cond.2 RT 2.12 min LC/MS(ESI ⁺) m/z; 453 [M+H] ⁺
72	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 0.91 (s, 9H), 1.37–1.45 (m, 2H), 2.31–2.39 (m, 2H), 2.41–2.52 (m, 4H), 3.82–3.98 (m, 4H), 4.92 (s, 2H), 7.28–7.32 (m, 4H), 7.88 (s, 1H). LC/MS:cond.1 RT 2.80 min LC/MS(ESI ⁺) m/z; 390 [M+H] ⁺ LC/MS(ESI ⁻) m/z; 434 [M+HCO ₂] ⁻
73	Morphology: colorless solid LC/MS:cond.2 RT 1.37 min LC/MS(ESI ⁺) m/z; 390 [M+H] ⁺
74	Morphology: colorless solid LC/MS:cond.1 RT 3.70 min LC/MS(ESI ⁺) m/z; 428 [M+H] ⁺ LC/MS(ESI ⁻) m/z; 472 [M+HCO ₂] ⁻

Ex	Data
75	Morphology: colorless solid LC/MS:cond.2 RT 2.47 min LC/MS(ESI ⁺) m/z; 460 [M+H] ⁺
76	Morphology: yellow solid ¹ H-NMR (CDCl ₃) δ : 3.76-3.90 (m, 4H), 3.60-4.80 (m, 4H), 4.96 (s, 2H), 6.60 (d, J = 9.6 Hz, 1H), 7.26-7.37 (m, 4H), 7.96 (s, 1H), 8.26 (dd, J = 2.6, 9.2 Hz, 1H), 9.05 (d, J = 2.6 Hz, 1H). LC/MS:cond.1 RT 3.97 min LC/MS(ESI ⁺) m/z; 428 [M+H] ⁺
77	Morphology: colorless solid LC/MS:cond.1 RT 4.30 min LC/MS(ESI ⁺) m/z; 451 [M+H] ⁺
78	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 3.04-3.10 (m, 4H), 3.95-4.05 (m, 4H), 4.94 (s, 2H), 5.96 (d, J = 4.3 Hz, 1H), 6.58 (d, J = 4.3 Hz, 1H), 7.27-7.40 (m, 4H), 7.93 (s, 1H). LC/MS:cond.1 RT 4.42 min LC/MS(ESI ⁺) m/z; 422 [M+H] ⁺
79	Morphology: colorless solid LC/MS:cond.1 RT 3.40 min LC/MS(ESI ⁺) m/z; 414 [M+H] ⁺ LC/MS(ESI ⁻) m/z; 412 [M-H] ⁻
80	Morphology: colorless solid LC/MS:cond.1 RT 3.77 min LC/MS(ESI ⁺) m/z; 414 [M+H] ⁺ LC/MS(ESI ⁻) m/z; 458 [M+HCO ₂] ⁻
81	Morphology: pale brown amorphous LC/MS:cond.1 RT 3.85 min LC/MS(ESI ⁺) m/z; 428 [M+H] ⁺ LC/MS(ESI ⁻) m/z; 472 [M+HCO ₂] ⁻
82	Morphology: colorless solid LC/MS:cond.1 RT 3.90 min LC/MS(ESI ⁺) m/z; 414 [M+H] ⁺ LC/MS(ESI ⁻) m/z; 458 [M+HCO ₂] ⁻
83	Morphology: colorless solid LC/MS:cond.1 RT 3.88 min LC/MS(ESI ⁺) m/z; 414 [M+H] ⁺ LC/MS(ESI ⁻) m/z; 458 [M+HCO ₂] ⁻
84	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 1.95-2.07 (m, 2H), 2.20-2.41 (m, 2H), 2.99-3.19 (m, 2H), 4.97 (s, 2H), 5.05 (m, 1H), 5.17 (m, 1H), 5.50 (m, 1H), 6.84-6.98 (m, 2H), 7.20 (d, 1H), 7.35 (m, 4H) 7.95 (s, 1H). LC/MS:cond.1 RT 4.05 min LC/MS(ESI ⁺) m/z; 471 [M+H] ⁺
85	Morphology: colorless solid LC/MS:cond.2 RT 2.41 min LC/MS(ESI ⁺) m/z; 399 [M+H] ⁺
86	Morphology: colorless solid LC/MS:cond.2 RT 2.65 min LC/MS(ESI ⁺) m/z; 414 [M+H] ⁺
87	Morphology: pale yellow solid LC/MS:cond.2 RT 2.29 min LC/MS(ESI ⁺) m/z; 401 [M+H] ⁺
88	Morphology: colorless solid LC/MS:cond.2 RT 2.45 min LC/MS(ESI ⁺) m/z; 439 [M+H] ⁺
89	Morphology: colorless solid LC/MS:cond.2 RT 2.32 min LC/MS(ESI ⁺) m/z; 387 [M+H] ⁺
90	Morphology: colorless solid LC/MS:cond.2 RT 2.39 min LC/MS(ESI ⁺) m/z; 413 [M+H] ⁺
91	Morphology: colorless solid LC/MS:cond.2 RT 2.24 min LC/MS(ESI ⁺) m/z; 431 [M+H] ⁺
92	Morphology: colorless solid LC/MS:cond.2 RT 2.43 min LC/MS(ESI ⁺) m/z; 400 [M+H] ⁺
93	Morphology: colorless amorphous LC/MS:cond.2 RT 2.59 min LC/MS(ESI ⁺) m/z; 414 [M+H] ⁺

[Table 25]

Ex	Data
94	Morphology: colorless solid LC/MS:cond. 2 RT 2.59 min LC/MS(ESI ⁺) m/z; 416 [M+H] ⁺
95	Morphology: colorless solid LC/MS:cond. 2 RT 2.59 min LC/MS(ESI ⁺) m/z; 418 [M+H] ⁺
96	Morphology: colorless solid LC/MS:cond. 1 RT 4.52 min LC/MS(ESI ⁺) m/z; 414 [M+H] ⁺
97	Morphology: colorless solid LC/MS:cond. 1 RT 3.62 min LC/MS(ESI ⁺) m/z; 374 [M+H] ⁺ LC/MS(ESI ⁻) m/z; 372 [M-H] ⁻
98	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 2.58 (brs, 2H), 4.00-4.20 (m, 2H), 4.35-4.55 (m, 2H), 6.00 (brs, 1H), 6.04 (s, 2H), 6.54 (d, J = 2.3 Hz, 1H), 7.02 (t, J = 8.6 Hz, 2H), 7.20-7.40 (m, 2H), 7.94 (s, 1H), 8.34 (s, 1H). LC/MS:cond. 1 RT 4.42 min LC/MS(ESI ⁺) m/z; 421 [M+H] ⁺
99	Morphology: pale brown solid ¹ H-NMR (CDCl ₃) δ : 2.58 (brs, 2H), 4.09 (q, J = 5.3 Hz, 2H), 4.45 (q, J = 2.9 Hz, 2H), 5.12 (s, 2H), 6.02 (brs, 1H), 6.97-7.13 (m, 1H), 7.02 (t, J = 8.8 Hz, 2H), 7.25-7.40 (m, 3H), 8.00 (s, 1H). LC/MS:cond. 2 RT 2.79 min LC/MS(ESI ⁺) m/z; 437 [M+H] ⁺
100	Morphology: colorless solid LC/MS:cond. 1 RT 4.09 min LC/MS(ESI ⁺) m/z; 403 [M+H] ⁺
101	Morphology: colorless solid LC/MS:cond. 1 RT 4.14 min LC/MS(ESI ⁺) m/z; 417 [M+H] ⁺
102	Morphology: colorless solid LC/MS:cond. 1 RT 4.31 min LC/MS(ESI ⁺) m/z; 414 [M+H] ⁺
103	Morphology: colorless solid LC/MS:cond. 1 RT 4.36 min LC/MS(ESI ⁺) m/z; 415 [M+H] ⁺
104	Morphology: colorless solid LC/MS:cond. 2 RT 2.30 min LC/MS(ESI ⁺) m/z; 414 [M+H] ⁺
105	Morphology: colorless solid LC/MS:cond. 2 RT 1.95 min LC/MS(ESI ⁺) m/z; 414 [M+H] ⁺
106	Morphology: colorless solid LC/MS:cond. 2 RT 1.90 min LC/MS(ESI ⁺) m/z; 414 [M+H] ⁺
107	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 3.35-3.60 (m, 1H), 3.90-4.10 (m, 1H), 4.95 (s, 2H), 4.90-5.55 (m, 3H), 6.25-6.35 (m, 1H), 7.00-7.10 (m, 2H), 7.25-7.50 (m, 6H), 7.98 (s, 1H). LC/MS:cond. 2 RT 2.69 min LC/MS(ESI ⁺) m/z; 415 [M+H] ⁺
108	Morphology: colorless solid LC/MS:cond. 2 RT 2.48 min LC/MS(ESI ⁺) m/z; 415 [M+H] ⁺
109	Morphology: pale yellow solid LC/MS:cond. 1 RT 4.44 min LC/MS(ESI ⁺) m/z; 458 [M+H] ⁺
110	Morphology: colorless solid LC/MS:cond. 2 RT 2.64 min LC/MS(ESI ⁺) m/z; 401 [M+H] ⁺

[Table 26]

Ex	Data
111	Morphology: colorless solid LC/MS:cond.1 RT 4.05 min LC/MS(ESI ⁺) m/z; 415 [M+H] ⁺
112	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 2.53-2.65 (m, 2H), 4.08-4.13 (m, 2H), 4.41-4.50 (m, 2H), 4.95 (s, 2H), 6.00-6.03 (m, 1H), 7.02 (t, J = 8.6 Hz, 2H), 7.25-7.40 (m, 6H), 7.93 (s, 1H). LC/MS:cond.1 RT 4.51 min LC/MS(ESI ⁺) m/z; 397 [M+H] ⁺
113	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 2.20-2.85 (m, 4H), 3.20-3.44 (m, 2H), 4.76-5.10 (m, 4H) 7.06 (t, J = 8.8 Hz, 2H), 7.20-7.50 (m, 6H), 7.93 (s, 1H). LC/MS:cond.1 RT 4.51 min LC/MS(ESI ⁺) m/z; 417 [M+H] ⁺
114	Morphology: colorless solid ¹ H-NMR (CDCl ₃) δ : 1.60-1.80 (m, 2H), 1.85-2.00 (m, 2H), 2.70-3.10 (m, 3H), 4.85-5.15 (m, 2H), 4.94 (s, 2H), 6.99 (t, J = 8.6 Hz, 2H), 7.10-8.20 (m, 2H), 7.26-7.39 (m, 4H), 7.91 (s, 1H). LC/MS:cond.1 RT 4.51 min LC/MS(ESI ⁺) m/z; 399 [M+H] ⁺
115	LC/MS:cond.1 RT 2.99 min LC/MS(ESI ⁺) m/z; 430 [M+H] ⁺

[Table 27]

Ex	Data
116	Morphology: colorless solid
117	Morphology: colorless solid
118	Morphology: colorless solid
119	Morphology: colorless solid
120	Morphology: yellow solid LC/MS: cond. 3 RT 2.04 min LC/MS (ESI ⁺) m/z; 456 [M+H] ⁺
121	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.42 min LC/MS (ESI ⁺) m/z; 397 [M+H] ⁺
122	Morphology: pale brown oil LC/MS: cond. 2 RT 1.76 min LC/MS (ESI ⁺) m/z; 414 [M+H] ⁺
123	Morphology: colorless solid LC/MS: cond. 2 RT 2.51 min LC/MS (ESI ⁺) m/z; 417 [M+H] ⁺
124	Morphology: colorless solid LC/MS: cond. 1 RT 2.70 min LC/MS (ESI ⁺) m/z; 400 [M+H] ⁺
125	Morphology: colorless solid LC/MS: cond. 2 RT 2.55 min LC/MS (ESI ⁺) m/z; 407 [M+H] ⁺
126	Morphology: colorless solid LC/MS: cond. 2 RT 2.56 min LC/MS (ESI ⁺) m/z; 407 [M+H] ⁺
127	Morphology: colorless solid LC/MS: cond. 2 RT 2.51 min LC/MS (ESI ⁺) m/z; 411 [M+H] ⁺
128	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.46 min LC/MS (ESI ⁺) m/z; 414 [M+H] ⁺
129	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.46 min LC/MS (ESI ⁺) m/z; 414 [M+H] ⁺
130	Morphology: white solid LC/MS: cond. 2 RT 2.53 min LC/MS (ESI ⁻) m/z; 430 [M-H] ⁻
131	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.36 min LC/MS (ESI ⁺) m/z; 396 [M+H] ⁺
132	Morphology: colorless solid LC/MS: cond. 2 RT 2.66 min, 2.70 min LC/MS (ESI ⁺) m/z; 411 [M+H] ⁺
133	Morphology: colorless solid LC/MS: cond. 2 RT 2.69 min LC/MS (ESI ⁺) m/z; 411 [M+H] ⁺
134	Morphology: colorless solid LC/MS: cond. 2 RT 2.68 min LC/MS (ESI ⁺) m/z; 431 [M+H] ⁺
135	Morphology: colorless solid LC/MS: cond. 1 RT 4.10 min LC/MS (ESI ⁺) m/z; 429 [M+H] ⁺
136	Morphology: colorless solid LC/MS: cond. 2 RT 2.48 min LC/MS (ESI ⁺) m/z; 414 [M+H] ⁺
137	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.49 min LC/MS (ESI ⁺) m/z; 414 [M+H] ⁺
138	Morphology: colorless solid LC/MS: cond. 2 RT 2.51 min LC/MS (ESI ⁺) m/z; 418 [M+H] ⁺
139	Morphology: white solid LC/MS: cond. 2 RT 2.23 min LC/MS (ESI ⁺) m/z; 397 [M+H] ⁺
140	Morphology: colorless solid LC/MS: cond. 1 RT 4.84 min LC/MS (ESI ⁺) m/z; 466 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 510 [M+HCO ₂] ⁻
141	Morphology: colorless solid LC/MS: cond. 1 RT 4.12 min LC/MS (ESI ⁺) m/z; 430 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 474 [M+HCO ₂] ⁻

[Table 28]

Ex	Data
142	Morphology: colorless solid LC/MS: cond. 1 RT 4.27 min LC/MS (ESI ⁺) m/z; 436 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 480 [M+HCO ₂] ⁻
143	Morphology: pale yellow solid LC/MS: cond. 1 RT 3.99 min LC/MS (ESI ⁺) m/z; 435 [M+H] ⁺
144	Morphology: pale yellow solid LC/MS: cond. 1 RT 4.17 min LC/MS (ESI ⁺) m/z; 465 [M+H] ⁺
145	Morphology: colorless solid LC/MS: cond. 1 RT 4.07 min LC/MS (ESI ⁺) m/z; 466 [M+H] ⁺
146	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.15 min LC/MS (ESI ⁺) m/z; 397 [M+H] ⁺
147	Morphology: colorless solid LC/MS: cond. 1 RT 4.27 min LC/MS (ESI ⁺) m/z; 464 [M+H] ⁺
148	Morphology: colorless solid LC/MS: cond. 2 RT 2.44 min LC/MS (ESI ⁺) m/z; 450 [M+H] ⁺
149	Morphology: colorless solid LC/MS: cond. 2 RT 2.63 min LC/MS (ESI ⁺) m/z; 528 [M+H] ⁺
150	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.26 min LC/MS (ESI ⁺) m/z; 438 [M+H] ⁺
151	Morphology: colorless solid LC/MS: cond. 2 RT 2.67 min LC/MS (ESI ⁺) m/z; 528 [M+H] ⁺
152	Morphology: colorless solid LC/MS: cond. 2 RT 2.40 min LC/MS (ESI ⁺) m/z; 450 [M+H] ⁺
153	Morphology: colorless solid LC/MS: cond. 2 RT 2.59 min LC/MS (ESI ⁺) m/z; 464 [M+H] ⁺
154	Morphology: colorless solid LC/MS: cond. 2 RT 2.67 min LC/MS (ESI ⁺) m/z; 518 [M+H] ⁺
155	Morphology: pale yellow solid LC/MS: cond. 1 RT 4.00 min LC/MS (ESI ⁺) m/z; 386 [M+H] ⁺
156	Morphology: colorless solid LC/MS: cond. 2 RT 2.52 min LC/MS (ESI ⁺) m/z; 474 [M+H] ⁺
157	Morphology: colorless solid LC/MS: cond. 2 RT 2.30 min LC/MS (ESI ⁺) m/z; 390 [M+H] ⁺
158	Morphology: colorless solid LC/MS: cond. 2 RT 2.31 min LC/MS (ESI ⁺) m/z; 422 [M+H] ⁺
159	Morphology: colorless solid LC/MS: cond. 1 RT 4.29 min LC/MS (ESI ⁺) m/z; 421 [M+H] ⁺
160	Morphology: colorless solid LC/MS: cond. 2 RT 2.57 min LC/MS (ESI ⁺) m/z; 447 [M+H] ⁺
161	Morphology: colorless solid LC/MS: cond. 1 RT 4.04 min LC/MS (ESI ⁺) m/z; 433 [M+H] ⁺
162	Morphology: colorless solid LC/MS: cond. 1 RT 4.12 min LC/MS (ESI ⁺) m/z; 463 [M+H] ⁺
163	Morphology: colorless solid LC/MS: cond. 1 RT 4.32 min LC/MS (ESI ⁺) m/z; 468 [M+H] ⁺
164	Morphology: colorless solid LC/MS: cond. 2 RT 2.47 min LC/MS (ESI ⁺) m/z; 419 [M+H] ⁺
165	Morphology: colorless solid LC/MS: cond. 2 RT 2.82 min LC/MS (ESI ⁺) m/z; 515 [M+H] ⁺
166	Morphology: colorless solid LC/MS: cond. 1 RT 4.80 min LC/MS (ESI ⁺) m/z; 464 [M+H] ⁺

[Table 29]

Ex	Data
167	Morphology: colorless solid LC/MS: cond. 2 RT 2.52 min LC/MS (ESI ⁺) m/z; 383 [M+H] ⁺
168	Morphology: colorless solid LC/MS: cond. 2 RT 2.43 min LC/MS (ESI ⁺) m/z; 430 [M+H] ⁺
169	Morphology: colorless solid LC/MS: cond. 2 RT 2.60 min LC/MS (ESI ⁺) m/z; 512 [M+H] ⁺
170	Morphology: colorless solid LC/MS: cond. 2 RT 2.56 min LC/MS (ESI ⁺) m/z; 421 [M+H] ⁺
171	Morphology: colorless solid LC/MS: cond. 2 RT 2.59 min LC/MS (ESI ⁺) m/z; 403 [M+H] ⁺
172	Morphology: colorless solid LC/MS: cond. 2 RT 2.57 min LC/MS (ESI ⁺) m/z; 462 [M+H] ⁺
173	Morphology: colorless solid LC/MS: cond. 2 RT 2.72 min LC/MS (ESI ⁺) m/z; 437 [M+H] ⁺
174	Morphology: colorless solid LC/MS: cond. 2 RT 2.45 min LC/MS (ESI ⁺) m/z; 423 [M+H] ⁺
175	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.61 min LC/MS (ESI ⁺) m/z; 447 [M+H] ⁺
176	Morphology: colorless solid LC/MS: cond. 2 RT 2.74 min LC/MS (ESI ⁺) m/z; 447 [M+H] ⁺
177	Morphology: colorless solid LC/MS: cond. 2 RT 2.41 min LC/MS (ESI ⁺) m/z; 405 [M+H] ⁺
178	Morphology: colorless solid LC/MS: cond. 2 RT 2.52 min LC/MS (ESI ⁺) m/z; 419 [M+H] ⁺
179	Morphology: yellow solid LC/MS: cond. 2 RT 1.96 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
180	Morphology: colorless solid LC/MS: cond. 2 RT 2.47 min LC/MS (ESI ⁺) m/z; 387 [M+H] ⁺
181	Morphology: colorless solid LC/MS: cond. 2 RT 2.38 min LC/MS (ESI ⁺) m/z; 421 [M+H] ⁺
182	Morphology: colorless solid LC/MS: cond. 2 RT 2.38 min LC/MS (ESI ⁺) m/z; 407 [M+H] ⁺
183	Morphology: colorless solid LC/MS: cond. 2 RT 2.88 min LC/MS (ESI ⁺) m/z; 419 [M+H] ⁺
184	Morphology: colorless solid LC/MS: cond. 2 RT 2.77 min LC/MS (ESI ⁺) m/z; 471 [M+H] ⁺
185	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.81 min LC/MS (ESI ⁺) m/z; 423 [M+H] ⁺
186	Morphology: colorless solid LC/MS: cond. 2 RT 2.66 min LC/MS (ESI ⁺) m/z; 471 [M+H] ⁺
187	Morphology: colorless solid LC/MS: cond. 2 RT 2.25 min LC/MS (ESI ⁺) m/z; 411 [M+H] ⁺
188	Morphology: colorless solid LC/MS: cond. 2 RT 2.16 min LC/MS (ESI ⁺) m/z; 437 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 481 [M+HCO ₂] ⁻
189	Morphology: white solid LC/MS: cond. 2 RT 2.29 min LC/MS (ESI ⁺) m/z; 528 [M+H] ⁺
190	Morphology: colorless solid LC/MS: cond. 1 RT 3.99 min LC/MS (ESI ⁺) m/z; 453 [M+H] ⁺
191	Morphology: colorless solid LC/MS: cond. 2 RT 2.51 min LC/MS (ESI ⁺) m/z; 435 [M+H] ⁺

[Table 30]

Ex	Data
192	Morphology: colorless solid LC/MS: cond. 2 RT 2.09 min LC/MS (ESI ⁺) m/z; 446 [M+H] ⁺
193	Morphology: colorless solid LC/MS: cond. 2 RT 2.20 min LC/MS (ESI ⁺) m/z; 437 [M+H] ⁺
194	Morphology: colorless solid LC/MS: cond. 2 RT 2.33 min LC/MS (ESI ⁺) m/z; 487 [M+H] ⁺
195	Morphology: colorless solid LC/MS: cond. 2 RT 2.28 min LC/MS (ESI ⁺) m/z; 489 [M+H] ⁺
196	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.09 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
197	Morphology: colorless solid LC/MS: cond. 2 RT 2.20 min LC/MS (ESI ⁺) m/z; 419 [M+H] ⁺
198	Morphology: colorless solid LC/MS: cond. 2 RT 2.08 min LC/MS (ESI ⁺) m/z; 403 [M+H] ⁺
199	Morphology: colorless solid LC/MS: cond. 2 RT 2.22 min LC/MS (ESI ⁺) m/z; 463 [M+H] ⁺
200	Morphology: colorless solid LC/MS: cond. 2 RT 2.48 min LC/MS (ESI ⁺) m/z; 441 [M+H] ⁺
201	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.42 min LC/MS (ESI ⁺) m/z; 439 [M+H] ⁺
202	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.38 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
203	Morphology: colorless solid LC/MS: cond. 2 RT 2.15 min LC/MS (ESI ⁺) m/z; 437 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 481 [M+HCO ₂] ⁻
204	Morphology: colorless solid LC/MS: cond. 2 RT 2.15 min LC/MS (ESI ⁺) m/z; 437 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 481 [M+HCO ₂] ⁻
205	Morphology: yellow solid LC/MS: cond. 2 RT 2.43 min LC/MS (ESI ⁺) m/z; 439 [M+H] ⁺
206	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.08 min LC/MS (ESI ⁺) m/z; 433 [M+H] ⁺
207	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.22 min LC/MS (ESI ⁺) m/z; 449 [M+H] ⁺
208	Morphology: white solid LC/MS: cond. 2 RT 2.30 min LC/MS (ESI ⁺) m/z; 528 [M+H] ⁺
209	Morphology: brown solid LC/MS: cond. 2 RT 2.10 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
210	Morphology: brown solid LC/MS: cond. 2 RT 2.48 min LC/MS (ESI ⁺) m/z; 441 [M+H] ⁺
211	Morphology: white solid LC/MS: cond. 2 RT 2.39 min LC/MS (ESI ⁺) m/z; 527 [M+H] ⁺
212	Morphology: colorless solid LC/MS: cond. 2 RT 2.00 min LC/MS (ESI ⁺) m/z; 492 [M+H] ⁺
213	Morphology: white solid LC/MS: cond. 2 RT 2.48 min LC/MS (ESI ⁺) m/z; 541 [M+H] ⁺
214	Morphology: colorless solid LC/MS: cond. 2 RT 2.19 min LC/MS (ESI ⁺) m/z; 460 [M+H] ⁺
215	Morphology: colorless solid LC/MS: cond. 2 RT 2.49 min LC/MS (ESI ⁺) m/z; 501 [M+H] ⁺
216	Morphology: colorless oil LC/MS: cond. 2 RT 2.38 min LC/MS (ESI ⁺) m/z; 542 [M+H] ⁺

[Table 31]

Ex	Data
217	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.10 min LC/MS (ESI ⁺) m/z; 506 [M+H] ⁺
218	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.32 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
219	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.24 min LC/MS (ESI ⁺) m/z; 468 [M+H] ⁺
220	Morphology: colorless solid LC/MS: cond. 1 RT 4.49 min LC/MS (ESI ⁺) m/z; 437 [M+H] ⁺
221	Morphology: colorless solid LC/MS: cond. 2 RT 2.46 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
222	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.65 min LC/MS (ESI ⁺) m/z; 506 [M+H] ⁺
223	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.80 min LC/MS (ESI ⁺) m/z; 484 [M+H] ⁺
224	Morphology: colorless solid LC/MS: cond. 2 RT 2.45 min LC/MS (ESI ⁺) m/z; 440 [M+H] ⁺
225	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.69 min LC/MS (ESI ⁺) m/z; 435 [M+H] ⁺
226	Morphology: white solid LC/MS: cond. 1 RT 4.22 min LC/MS (ESI ⁺) m/z; 385 [M+H] ⁺
227	Morphology: white solid LC/MS: cond. 1 RT 3.90 min LC/MS (ESI ⁺) m/z; 398 [M+H] ⁺
228	Morphology: colorless solid LC/MS: cond. 2 RT 2.66 min LC/MS (ESI ⁺) m/z; 437 [M+H] ⁺
229	Morphology: colorless solid LC/MS: cond. 2 RT 2.68 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
230	Morphology: colorless solid LC/MS: cond. 2 RT 2.57 min LC/MS (ESI ⁺) m/z; 425 [M+H] ⁺
231	Morphology: colorless solid LC/MS: cond. 2 RT 2.27 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
232	Morphology: colorless solid LC/MS: cond. 2 RT 1.70 min LC/MS (ESI ⁺) m/z; 420 [M+H] ⁺
233	Morphology: colorless solid LC/MS: cond. 2 RT 2.84 min LC/MS (ESI ⁺) m/z; 459 [M+H] ⁺
234	Morphology: colorless solid LC/MS: cond. 1 RT 2.65 min LC/MS (ESI ⁺) m/z; 437 [M+H] ⁺
235	Morphology: colorless solid LC/MS: cond. 2 RT 2.46 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
236	Morphology: colorless solid LC/MS: cond. 2 RT 2.98 min LC/MS (ESI ⁺) m/z; 519 [M+H] ⁺
237	Morphology: colorless solid LC/MS: cond. 2 RT 2.78 min LC/MS (ESI ⁺) m/z; 433 [M+H] ⁺
238	Morphology: colorless solid LC/MS: cond. 2 RT 2.72 min LC/MS (ESI ⁺) m/z; 491 [M+H] ⁺
239	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.83 min LC/MS (ESI ⁺) m/z; 487 [M+H] ⁺
240	Morphology: colorless solid LC/MS: cond. 2 RT 2.61 min LC/MS (ESI ⁺) m/z; 450 [M+H] ⁺
241	Morphology: colorless solid LC/MS: cond. 2 RT 2.33 min LC/MS (ESI ⁺) m/z; 438 [M+H] ⁺

[Table 32]

Ex	Data
242	Morphology: colorless solid LC/MS: cond. 2 RT 2.76 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
243	Morphology: colorless solid LC/MS: cond. 2 RT 2.88 min LC/MS (ESI ⁺) m/z; 459 [M+H] ⁺
244	Morphology: colorless solid LC/MS: cond. 2 RT 1.86 min LC/MS (ESI ⁺) m/z; 420 [M+H] ⁺
245	Morphology: colorless solid LC/MS: cond. 2 RT 1.74 min LC/MS (ESI ⁺) m/z; 420 [M+H] ⁺
246	Morphology: colorless solid LC/MS: cond. 2 RT 2.80 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
247	Morphology: colorless solid LC/MS: cond. 1 RT 4.60 min LC/MS (ESI ⁺) m/z; 470 [M+H] ⁺
248	Morphology: pale gray solid LC/MS: cond. 1 RT 4.20 min LC/MS (ESI ⁺) m/z; 504 [M+H] ⁺
249	Morphology: colorless solid LC/MS: cond. 2 RT 2.63 min LC/MS (ESI ⁺) m/z; 488 [M+H] ⁺
250	Morphology: colorless solid LC/MS: cond. 2 RT 2.25 min LC/MS (ESI ⁺) m/z; 426 [M+H] ⁺
251	Morphology: colorless solid LC/MS: cond. 2 RT 2.61 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
252	Morphology: colorless solid LC/MS: cond. 2 RT 2.40 min LC/MS (ESI ⁺) m/z; 450 [M+H] ⁺
253	Morphology: colorless solid LC/MS: cond. 2 RT 2.92 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
254	Morphology: colorless solid LC/MS: cond. 2 RT 2.85 min LC/MS (ESI ⁺) m/z; 471 [M+H] ⁺
255	Morphology: colorless solid LC/MS: cond. 2 RT 2.77 min LC/MS (ESI ⁺) m/z; 535 [M+H] ⁺
256	Morphology: colorless solid LC/MS: cond. 2 RT 2.56 min LC/MS (ESI ⁺) m/z; 454 [M+H] ⁺
257	Morphology: colorless solid LC/MS: cond. 2 RT 2.79 min LC/MS (ESI ⁺) m/z; 518 [M+H] ⁺
258	Morphology: colorless solid LC/MS: cond. 2 RT 2.58 min LC/MS (ESI ⁺) m/z; 454 [M+H] ⁺
259	Morphology: colorless solid LC/MS: cond. 2 RT 2.33 min LC/MS (ESI ⁺) m/z; 491 [M+H] ⁺
260	Morphology: pale orange solid LC/MS: cond. 2 RT 2.80 min LC/MS (ESI ⁺) m/z; 459 [M+H] ⁺
261	Morphology: colorless solid LC/MS: cond. 2 RT 2.24 min LC/MS (ESI ⁺) m/z; 438 [M+H] ⁺
262	Morphology: colorless solid LC/MS: cond. 2 RT 2.31 min LC/MS (ESI ⁺) m/z; 438 [M+H] ⁺
263	Morphology: colorless solid LC/MS: cond. 1 RT 4.12 min LC/MS (ESI ⁺) m/z; 438 [M+H] ⁺
264	Morphology: colorless solid LC/MS: cond. 1 RT 4.47 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
265	Morphology: pale yellow solid LC/MS: cond. 1 RT 3.80 min LC/MS (ESI ⁺) m/z; 426 [M+H] ⁺
266	Morphology: colorless solid LC/MS: cond. 2 RT 3.01 min LC/MS (ESI ⁺) m/z; 521 [M+H] ⁺

[Table 33]

Ex	Data
267	Morphology: colorless solid LC/MS: cond. 2 RT 2.83 min LC/MS (ESI ⁺) m/z; 505 [M+H] ⁺
268	Morphology: colorless solid LC/MS: cond. 2 RT 2.28 min LC/MS (ESI ⁺) m/z; 438 [M+H] ⁺
269	Morphology: colorless solid LC/MS: cond. 2 RT 2.37 min LC/MS (ESI ⁺) m/z; 473 [M+H] ⁺
270	Morphology: colorless solid LC/MS: cond. 2 RT 2.45 min LC/MS (ESI ⁺) m/z; 438 [M+H] ⁺
271	Morphology: colorless solid LC/MS: cond. 2 RT 1.77 min LC/MS (ESI ⁺) m/z; 450 [M+H] ⁺
272	Morphology: colorless solid LC/MS: cond. 2 RT 2.37 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
273	Morphology: colorless solid LC/MS: cond. 2 RT 2.79 min LC/MS (ESI ⁺) m/z; 484 [M+H] ⁺
274	Morphology: colorless solid LC/MS: cond. 2 RT 2.68 min LC/MS (ESI ⁺) m/z; 488 [M+H] ⁺
275	Morphology: colorless solid LC/MS: cond. 2 RT 2.37 min LC/MS (ESI ⁺) m/z; 440 [M+H] ⁺
276	Morphology: colorless solid LC/MS: cond. 2 RT 2.79 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
277	Morphology: colorless solid LC/MS: cond. 2 RT 2.76 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
278	Morphology: colorless solid LC/MS: cond. 2 RT 2.53 min LC/MS (ESI ⁺) m/z; 462 [M+H] ⁺
279	Morphology: colorless solid LC/MS: cond. 2 RT 2.61 min LC/MS (ESI ⁺) m/z; 449 [M+H] ⁺
280	Morphology: colorless solid LC/MS: cond. 2 RT 2.73 min LC/MS (ESI ⁺) m/z; 484 [M+H] ⁺
281	Morphology: colorless solid LC/MS: cond. 2 RT 2.72 min LC/MS (ESI ⁺) m/z; 439 [M+H] ⁺
282	Morphology: colorless solid LC/MS: cond. 2 RT 2.38 min LC/MS (ESI ⁺) m/z; 440 [M+H] ⁺
283	Morphology: colorless solid LC/MS: cond. 2 RT 1.72 min LC/MS (ESI ⁺) m/z; 449 [M+H] ⁺
284	Morphology: colorless solid LC/MS: cond. 2 RT 1.81 min LC/MS (ESI ⁺) m/z; 434 [M+H] ⁺
285	Morphology: colorless solid LC/MS: cond. 2 RT 2.70 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
286	Morphology: colorless solid LC/MS: cond. 2 RT 2.12 min LC/MS (ESI ⁺) m/z; 450 [M+H] ⁺
287	Morphology: colorless solid LC/MS: cond. 2 RT 2.58 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
288	Morphology: colorless solid LC/MS: cond. 2 RT 2.99 min LC/MS (ESI ⁺) m/z; 490 [M+H] ⁺
289	Morphology: colorless solid LC/MS: cond. 2 RT 2.33 min LC/MS (ESI ⁺) m/z; 450 [M+H] ⁺
290	Morphology: colorless solid LC/MS: cond. 2 RT 1.91 min LC/MS (ESI ⁺) m/z; 434 [M+H] ⁺
291	Morphology: colorless solid LC/MS: cond. 2 RT 2.63 min LC/MS (ESI ⁺) m/z; 469 [M+H] ⁺

[Table 34]

Ex	Data
292	Morphology: colorless solid LC/MS: cond. 1 RT 4.27 min LC/MS (ESI ⁺) m/z; 462 [M+H] ⁺
293	Morphology: colorless solid LC/MS: cond. 2 RT 2.64 min LC/MS (ESI ⁺) m/z; 486 [M+H] ⁺
294	Morphology: colorless solid LC/MS: cond. 2 RT 2.78 min LC/MS (ESI ⁺) m/z; 433 [M+H] ⁺
295	Morphology: colorless solid LC/MS: cond. 2 RT 2.36 min LC/MS (ESI ⁺) m/z; 460 [M+H] ⁺
296	Morphology: colorless solid LC/MS: cond. 2 RT 2.31 min LC/MS (ESI ⁺) m/z; 445 [M+H] ⁺
297	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.52 min LC/MS (ESI ⁺) m/z; 479 [M+H] ⁺
298	Morphology: colorless solid LC/MS: cond. 2 RT 2.38 min LC/MS (ESI ⁺) m/z; 445 [M+H] ⁺
299	Morphology: colorless solid LC/MS: cond. 2 RT 2.56 min LC/MS (ESI ⁺) m/z; 452 [M+H] ⁺
300	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.63 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
301	Morphology: colorless solid LC/MS: cond. 2 RT 2.37 min LC/MS (ESI ⁺) m/z; 448 [M+H] ⁺
302	Morphology: colorless solid LC/MS: cond. 2 RT 2.52 min LC/MS (ESI ⁺) m/z; 465 [M+H] ⁺
303	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.29 min LC/MS (ESI ⁺) m/z; 453 [M+H] ⁺
304	Morphology: colorless solid LC/MS: cond. 2 RT 2.33 min LC/MS (ESI ⁺) m/z; 453 [M+H] ⁺
305	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.35 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
306	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.37 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
307	Morphology: colorless solid LC/MS: cond. 2 RT 2.36 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
308	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.30 min LC/MS (ESI ⁺) m/z; 441 [M+H] ⁺
309	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.31 min LC/MS (ESI ⁺) m/z; 441 [M+H] ⁺
310	Morphology: colorless solid LC/MS: cond. 2 RT 2.45 min LC/MS (ESI ⁺) m/z; 425 [M+H] ⁺
311	Morphology: colorless solid LC/MS: cond. 2 RT 2.56 min LC/MS (ESI ⁺) m/z; 437 [M+H] ⁺
312	Morphology: colorless solid LC/MS: cond. 2 RT 2.62 min LC/MS (ESI ⁺) m/z; 554 [M+H] ⁺
313	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.55 min LC/MS (ESI ⁺) m/z; 466 [M+H] ⁺
314	Morphology: colorless solid LC/MS: cond. 2 RT 2.32 min LC/MS (ESI ⁺) m/z; 530 [M+H] ⁺
315	Morphology: colorless solid LC/MS: cond. 2 RT 2.29 min LC/MS (ESI ⁺) m/z; 453 [M+H] ⁺
316	Morphology: colorless solid LC/MS: cond. 2 RT 2.29 min LC/MS (ESI ⁺) m/z; 453 [M+H] ⁺
317	Morphology: colorless solid LC/MS: cond. 2 RT 2.21 min LC/MS (ESI ⁺) m/z; 494 [M+H] ⁺

[Table 35]

Ex	Data
318	Morphology: colorless solid LC/MS: cond. 1 RT 4.30 min LC/MS (ESI ⁺) m/z; 425 [M+H] ⁺
319	Morphology: colorless solid LC/MS: cond. 1 RT 4.27 min LC/MS (ESI ⁺) m/z; 425 [M+H] ⁺
320	Morphology: colorless solid LC/MS: cond. 1 RT 4.14 min LC/MS (ESI ⁺) m/z; 450 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 448 [M-H] ⁻
321	Morphology: colorless solid LC/MS: cond. 2 RT 1.90 min LC/MS (ESI ⁺) m/z; 452 [M+H] ⁺
322	Morphology: colorless solid
323	Morphology: brown solid LC/MS: cond. 2 RT 1.91 min LC/MS (ESI ⁺) m/z; 452 [M+H] ⁺
324	Morphology: colorless solid LC/MS: cond. 2 RT 1.88 min LC/MS (ESI ⁺) m/z; 452 [M+H] ⁺
325	Morphology: colorless solid LC/MS: cond. 2 RT 2.42 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
326	Morphology: colorless solid LC/MS: cond. 2 RT 2.44 min LC/MS (ESI ⁺) m/z; 487 [M+H] ⁺
327	Morphology: colorless solid LC/MS: cond. 2 RT 2.35 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
328	Morphology: colorless solid LC/MS: cond. 2 RT 2.25 min LC/MS (ESI ⁺) m/z; 466 [M+H] ⁺
329	Morphology: colorless solid LC/MS: cond. 2 RT 2.43 min LC/MS (ESI ⁺) m/z; 500 [M+H] ⁺
330	Morphology: pale brown solid LC/MS: cond. 2 RT 2.26 min LC/MS (ESI ⁺) m/z; 470 [M+H] ⁺
331	Morphology: colorless solid LC/MS: cond. 2 RT 2.40 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
332	Morphology: colorless solid LC/MS: cond. 2 RT 1.68 min LC/MS (ESI ⁺) m/z; 450 [M+H] ⁺
333	Morphology: colorless solid LC/MS: cond. 2 RT 2.34 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
334	Morphology: colorless solid LC/MS: cond. 2 RT 1.98 min LC/MS (ESI ⁺) m/z; 466 [M+H] ⁺
335	Morphology: colorless solid LC/MS: cond. 2 RT 2.23 min LC/MS (ESI ⁺) m/z; 468 [M+H] ⁺
336	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.39 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
337	Morphology: colorless solid LC/MS: cond. 1 RT 3.87 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
338	Morphology: colorless solid LC/MS: cond. 1 RT 3.75 min LC/MS (ESI ⁺) m/z; 497 [M+H] ⁺
339	Morphology: colorless solid LC/MS: cond. 1 RT 4.79 min LC/MS (ESI ⁺) m/z; 505 [M+H] ⁺
340	Morphology: yellow solid LC/MS: cond. 1 RT 4.55 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
341	Morphology: colorless solid LC/MS: cond. 1 RT 4.15 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 465 [M-H] ⁻

[Table 36]

Ex	Data
342	Morphology: colorless amorphous ¹ H-NMR (CDCl ₃) δ : 2.51(d, J = 6.6 Hz, 3H), 3.41-3.46(m, 1H), 3.92-3.99(m, 1H), 4.70(d, J = 26.4 Hz, 1H), 4.87-5.07(m, 2H), 5.16-5.31(m, 2H), 6.09-6.15(m, 1H), 7.00-7.06(m, 3H), 7.27(m, 1H), 7.45-7.50(m, 2H) LC/MS: cond. 2 RT 2.39 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
343	Morphology: colorless amorphous ¹ H-NMR (CDCl ₃) δ : 2.51(d, J = 7.5 Hz, 3H), 3.41-3.47(m, 1H), 3.92-3.99(m, 1H), 4.70(d, J = 26.7, 1H), 4.87-5.05(m, 2H), 5.16-5.31(m, 2H), 6.09-6.14(m, 1H), 7.00-7.08(m, 3H), 7.28(m, 1H), 7.45-7.50(m, 2H)
344	Morphology: colorless solid LC/MS: cond. 2 RT 2.61 min LC/MS (ESI ⁺) m/z; 454 [M+H] ⁺
345	Morphology: colorless solid LC/MS: cond. 2 RT 2.65 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
346	Morphology: colorless solid LC/MS: cond. 2 RT 2.82 min LC/MS (ESI ⁺) m/z; 501 [M+H] ⁺
347	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.01 min LC/MS (ESI ⁺) m/z; 466 [M+H] ⁺
348	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.01 min LC/MS (ESI ⁺) m/z; 466 [M+H] ⁺
349	Morphology: colorless solid LC/MS: cond. 2 RT 2.53 min LC/MS (ESI ⁺) m/z; 481 [M+H] ⁺
350	Morphology: colorless solid LC/MS: cond. 2 RT 2.55 min LC/MS (ESI ⁺) m/z; 501 [M+H] ⁺
351	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.94 min LC/MS (ESI ⁺) m/z; 465 [M+H] ⁺
352	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.93 min LC/MS (ESI ⁺) m/z; 485 [M+H] ⁺
353	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.74 min LC/MS (ESI ⁺) m/z; 464 [M+H] ⁺
354	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.92 min LC/MS (ESI ⁺) m/z; 498 [M+H] ⁺
355	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.87 min LC/MS (ESI ⁺) m/z; 453 [M+H] ⁺
356	Morphology: yellow amorphous LC/MS: cond. 2 RT 2.36 min LC/MS (ESI ⁺) m/z; 480 [M+H] ⁺
357	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.60 min LC/MS (ESI ⁺) m/z; 456 [M+H] ⁺
358	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.79 min LC/MS (ESI ⁺) m/z; 468 [M+H] ⁺
359	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.80 min LC/MS (ESI ⁺) m/z; 488 [M+H] ⁺
360	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.57 min LC/MS (ESI ⁺) m/z; 439 [M+H] ⁺
361a	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.53 min LC/MS (ESI ⁺) m/z; 514 [M+H] ⁺
361b	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.62 min LC/MS (ESI ⁺) m/z; 514 [M+H] ⁺
362	Morphology: red amorphous LC/MS: cond. 2 RT 2.46 min LC/MS (ESI ⁺) m/z; 469 [M+H] ⁺
363	Morphology: colorless solid LC/MS: cond. 2 RT 2.70 min LC/MS (ESI ⁺) m/z; 450 [M+H] ⁺

[Table 37]

Ex	Data
364	Morphology: colorless solid LC/MS: cond. 2 RT 1.89 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
365	Morphology: colorless solid LC/MS: cond. 2 RT 2.64 min LC/MS (ESI ⁺) m/z; 469 [M+H] ⁺
366	Morphology: colorless solid LC/MS: cond. 2 RT 2.70 min LC/MS (ESI ⁺) m/z; 471 [M+H] ⁺
367	Morphology: colorless solid LC/MS: cond. 2 RT 2.74 min LC/MS (ESI ⁺) m/z; 472 [M+H] ⁺
368	Morphology: colorless solid LC/MS: cond. 2 RT 1.92 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
369	Morphology: colorless solid LC/MS: cond. 2 RT 2.50 min LC/MS (ESI ⁺) m/z; 481 [M+H] ⁺
370	Morphology: colorless solid LC/MS: cond. 2 RT 2.95 min LC/MS (ESI ⁺) m/z; 508 [M+H] ⁺
371	Morphology: yellow amorphous LC/MS: cond. 2 RT 1.77 min LC/MS (ESI ⁺) m/z; 464 [M+H] ⁺
372	Morphology: colorless solid LC/MS: cond. 2 RT 2.45 min LC/MS (ESI ⁺) m/z; 469 [M+H] ⁺
373	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.45 min LC/MS (ESI ⁺) m/z; 481 [M+H] ⁺
374	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.41 min LC/MS (ESI ⁺) m/z; 484 [M+H] ⁺
375	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.33 min LC/MS (ESI ⁺) m/z; 482 [M+H] ⁺
376	Morphology: colorless solid LC/MS: cond. 2 RT 2.73 min LC/MS (ESI ⁺) m/z; 504 [M+H] ⁺
377	Morphology: colorless solid LC/MS: cond. 2 RT 2.73 min LC/MS (ESI ⁺) m/z; 486 [M+H] ⁺
378	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.47 min LC/MS (ESI ⁺) m/z; 461 [M+H] ⁺
379	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.51 min LC/MS (ESI ⁺) m/z; 481 [M+H] ⁺
380	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.69 min LC/MS (ESI ⁺) m/z; 480 [M+H] ⁺
381	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 3.20 min LC/MS (ESI ⁺) m/z; 480 [M+H] ⁺
382	Morphology: colorless solid LC/MS: cond. 2 RT 2.14 min LC/MS (ESI ⁺) m/z; 494 [M+H] ⁺
383	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.69 min LC/MS (ESI ⁺) m/z; 486 [M+H] ⁺
384	Morphology: colorless solid LC/MS: cond. 2 RT 1.83 min LC/MS (ESI ⁺) m/z; 462 [M+H] ⁺
385	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.22 min LC/MS (ESI ⁺) m/z; 463 [M+H] ⁺
386	Morphology: colorless solid LC/MS: cond. 2 RT 2.36 min LC/MS (ESI ⁺) m/z; 477 [M+H] ⁺
387	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.38 min LC/MS (ESI ⁺) m/z; 497 [M+H] ⁺
388	Morphology: colorless solid LC/MS: cond. 2 RT 2.17 min LC/MS (ESI ⁺) m/z; 476 [M+H] ⁺
389	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.36 min LC/MS (ESI ⁺) m/z; 510 [M+H] ⁺

[Table 38]

Ex	Data
390	Morphology: colorless solid LC/MS: cond. 2 RT 1.95 min LC/MS (ESI ⁺) m/z; 476 [M+H] ⁺
391	Morphology: yellow amorphous LC/MS: cond. 2 RT 2.34 min LC/MS (ESI ⁺) m/z; 477 [M+H] ⁺
392	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.48 min LC/MS (ESI ⁺) m/z; 491 [M+H] ⁺
393	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.49 min LC/MS (ESI ⁺) m/z; 511 [M+H] ⁺
394	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.29 min LC/MS (ESI ⁺) m/z; 490 [M+H] ⁺
395	Morphology: yellow amorphous LC/MS: cond. 2 RT 2.47 min LC/MS (ESI ⁺) m/z; 524 [M+H] ⁺
396	Morphology: orange colored solid LC/MS: cond. 2 RT 2.41 min LC/MS (ESI ⁺) m/z; 479 [M+H] ⁺
397	Morphology: yellow solid LC/MS: cond. 2 RT 2.20 min LC/MS (ESI ⁺) m/z; 419 [M+H] ⁺
398	Morphology: colorless solid LC/MS: cond. 2 RT 1.78 min LC/MS (ESI ⁺) m/z; 418 [M+H] ⁺
399	Morphology: colorless solid LC/MS: cond. 2 RT 2.34 min LC/MS (ESI ⁺) m/z; 433 [M+H] ⁺
400	Morphology: colorless solid LC/MS: cond. 2 RT 2.36 min LC/MS (ESI ⁺) m/z; 453 [M+H] ⁺
401	Morphology: colorless solid LC/MS: cond. 2 RT 2.14 min LC/MS (ESI ⁺) m/z; 432 [M+H] ⁺
402	Morphology: colorless solid LC/MS: cond. 2 RT 2.32 min LC/MS (ESI ⁺) m/z; 466 [M+H] ⁺
403	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.31 min LC/MS (ESI ⁺) m/z; 433 [M+H] ⁺
404	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.46 min LC/MS (ESI ⁺) m/z; 447 [M+H] ⁺
405	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.47 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
406	Morphology: yellow solid LC/MS: cond. 2 RT 2.27 min LC/MS (ESI ⁺) m/z; 446 [M+H] ⁺
407	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.45 min LC/MS (ESI ⁺) m/z; 480 [M+H] ⁺
408	Morphology: yellow solid LC/MS: cond. 2 RT 2.38 min LC/MS (ESI ⁺) m/z; 435 [M+H] ⁺
409	Morphology: yellow amorphous LC/MS: cond. 2 RT 1.92 min LC/MS (ESI ⁺) m/z; 432 [M+H] ⁺
410	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.31 min LC/MS (ESI ⁺) m/z; 433 [M+H] ⁺
411	Morphology: colorless solid LC/MS: cond. 2 RT 2.61 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
412	Morphology: colorless solid LC/MS: cond. 2 RT 2.63 min LC/MS (ESI ⁺) m/z; 475 [M+H] ⁺
413	Morphology: colorless solid LC/MS: cond. 2 RT 2.61 min LC/MS (ESI ⁺) m/z; 488 [M+H] ⁺
414	Morphology: colorless solid LC/MS: cond. 2 RT 2.58 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺

[Table 39]

Ex	Data
415	Morphology: colorless solid LC/MS: cond. 2 RT 2.43 min LC/MS (ESI ⁺) m/z; 456 [M+H] ⁺
416	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.73 min LC/MS (ESI ⁺) m/z; 469 [M+H] ⁺
417	Morphology: colorless solid LC/MS: cond. 2 RT 2.75 min LC/MS (ESI ⁺) m/z; 489 [M+H] ⁺
418	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.73 min LC/MS (ESI ⁺) m/z; 502 [M+H] ⁺
419	Morphology: colorless solid LC/MS: cond. 2 RT 3.01 min LC/MS (ESI ⁺) m/z; 456 [M+H] ⁺
420	Morphology: colorless solid LC/MS: cond. 2 RT 3.01 min LC/MS (ESI ⁺) m/z; 476 [M+H] ⁺
421	Morphology: yellow solid LC/MS: cond. 2 RT 2.84 min LC/MS (ESI ⁺) m/z; 444 [M+H] ⁺
422	Morphology: colorless solid LC/MS: cond. 2 RT 2.60 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
423	Morphology: colorless solid LC/MS: cond. 2 RT 2.70 min LC/MS (ESI ⁺) m/z; 469 [M+H] ⁺
424	Morphology: colorless solid LC/MS: cond. 2 RT 1.98 min LC/MS (ESI ⁺) m/z; 452 [M+H] ⁺
425	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.65 min LC/MS (ESI ⁺) m/z; 457 [M+H] ⁺
426	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.54 min LC/MS (ESI ⁺) m/z; 470 [M+H] ⁺
427	Morphology: colorless solid LC/MS: cond. 2 RT 2.56 min LC/MS (ESI ⁺) m/z; 453 [M+H] ⁺
428	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.58 min LC/MS (ESI ⁺) m/z; 473 [M+H] ⁺
429a	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.55 min LC/MS (ESI ⁺) m/z; 486 [M+H] ⁺
429b	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.67 min LC/MS (ESI ⁺) m/z; 486 [M+H] ⁺
430	Morphology: colorless solid LC/MS: cond. 2 RT 1.81 min LC/MS (ESI ⁺) m/z; 436 [M+H] ⁺
431	Morphology: colorless solid LC/MS: cond. 2 RT 2.36 min LC/MS (ESI ⁺) m/z; 454 [M+H] ⁺
432	Morphology: yellow solid LC/MS: cond. 2 RT 2.55 min LC/MS (ESI ⁺) m/z; 453 [M+H] ⁺
433	Morphology: colorless solid LC/MS: cond. 2 RT 2.49 min LC/MS (ESI ⁺) m/z; 468 [M+H] ⁺
434	Morphology: colorless solid LC/MS: cond. 2 RT 2.58 min LC/MS (ESI ⁺) m/z; 438 [M+H] ⁺
435	Morphology: colorless solid LC/MS: cond. 2 RT 2.59 min LC/MS (ESI ⁺) m/z; 458 [M+H] ⁺
436	Morphology: colorless solid LC/MS: cond. 2 RT 2.36 min LC/MS (ESI ⁺) m/z; 426 [M+H] ⁺
437	Morphology: colorless solid LC/MS: cond. 2 RT 2.61 min LC/MS (ESI ⁺) m/z; 471 [M+H] ⁺
438	Morphology: pale yellow solid LC/MS: cond. 2 RT 1.83 min LC/MS (ESI ⁺) m/z; 436 [M+H] ⁺
439	Morphology: colorless solid LC/MS: cond. 2 RT 2.30 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺

[Table 40]

Ex	Data
440	Morphology: pale yellow solid LC/MS: cond. 2 RT 1.89 min LC/MS (ESI ⁺) m/z; 450 [M+H] ⁺
441	Morphology: colorless solid LC/MS: cond. 2 RT 2.54 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
442	Morphology: colorless solid LC/MS: cond. 2 RT 2.72 min LC/MS (ESI ⁺) m/z; 485 [M+H] ⁺
443	Morphology: colorless solid LC/MS: cond. 2 RT 2.69 min LC/MS (ESI ⁺) m/z; 452 [M+H] ⁺
444	Morphology: colorless solid LC/MS: cond. 2 RT 2.49 min LC/MS (ESI ⁺) m/z; 440 [M+H] ⁺
445	Morphology: pale brown solid LC/MS: cond. 2 RT 1.95 min LC/MS (ESI ⁺) m/z; 527 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 571 [M+HCO ₂] ⁻
446	Morphology: colorless solid LC/MS: cond. 2 RT 2.42 min LC/MS (ESI ⁺) m/z; 542 [M+H] ⁺
447	Morphology: colorless solid LC/MS: cond. 2 RT 2.44 min LC/MS (ESI ⁺) m/z; 562 [M+H] ⁺
448	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.50 min LC/MS (ESI ⁺) m/z; 556 [M+H] ⁺
449	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.52 min LC/MS (ESI ⁺) m/z; 576 [M+H] ⁺
450	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.50 min LC/MS (ESI ⁺) m/z; 589 [M+H] ⁺
451	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.02 min LC/MS (ESI ⁺) m/z; 541 [M+H] ⁺
452	Morphology: colorless solid LC/MS: cond. 2 RT 4.42 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
453	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.39 min LC/MS (ESI ⁺) m/z; 481 [M+H] ⁺
454	Morphology: pale yellow oil LC/MS: cond. 2 RT 2.55 min LC/MS (ESI ⁺) m/z; 517 [M+H] ⁺
455	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.32 min LC/MS (ESI ⁺) m/z; 485 [M+H] ⁺
456	Morphology: colorless solid LC/MS: cond. 2 RT 2.72 min LC/MS (ESI ⁺) m/z; 509 [M+H] ⁺
457	Morphology: colorless solid LC/MS: cond. 2 RT 2.40 min LC/MS (ESI ⁺) m/z; 525 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 523 [M-H] ⁻
458a	Morphology: colorless solid LC/MS: cond. 2 RT 2.61 min LC/MS (ESI ⁺) m/z; 512 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 510 [M-H] ⁻
458b	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.37 min LC/MS (ESI ⁺) m/z; 470 [M+H] ⁺
459	Morphology: colorless solid LC/MS: cond. 1 RT 4.77 min LC/MS (ESI ⁺) m/z; 465 [M+H] ⁺
460	Morphology: colorless solid LC/MS: cond. 1 RT 4.89 min LC/MS (ESI ⁺) m/z; 479 [M+H] ⁺
461	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.93 min LC/MS (ESI ⁺) m/z; 438 [M+H] ⁺
462	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.06 min LC/MS (ESI ⁺) m/z; 497 [M+H] ⁺

[Table 41]

Ex	Data
463	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.30 min LC/MS (ESI ⁺) m/z; 539 [M+H] ⁺
464	Morphology: colorless solid LC/MS: cond. 1 RT 4.00 min LC/MS (ESI ⁺) m/z; 413 [M+H] ⁺
465	Morphology: colorless solid LC/MS: cond. 2 RT 2.02 min LC/MS (ESI ⁺) m/z; 431 [M+H] ⁺
466a	Morphology: white solid LC/MS: cond. 1 RT 4.42 min LC/MS (ESI ⁺) m/z; 415 [M+H] ⁺
466b	Morphology: white solid LC/MS: cond. 2 RT 2.53 min LC/MS (ESI ⁺) m/z; 435 [M+H] ⁺
467	Morphology: colorless solid LC/MS: cond. 1 RT 3.82 min LC/MS (ESI ⁺) m/z; 416 [M+H] ⁺
468	Morphology: colorless solid LC/MS: cond. 2 RT 2.61 min LC/MS (ESI ⁺) m/z; 461 [M+H] ⁺
469	Morphology: yellow solid LC/MS: cond. 2 RT 2.62 min LC/MS (ESI ⁺) m/z; 454 [M+H] ⁺
470	Morphology: colorless solid LC/MS: cond. 2 RT 2.77 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
471	Morphology: pale yellow solid LC/MS: cond. 2 RT 2.25 min LC/MS (ESI ⁺) m/z; 413 [M+H] ⁺
472	Morphology: colorless solid LC/MS: cond. 2 RT 2.50 min LC/MS (ESI ⁺) m/z; 429 [M+H] ⁺
473	Morphology: colorless solid LC/MS: cond. 2 RT 2.70 min LC/MS (ESI ⁺) m/z; 511 [M+H] ⁺
474	Morphology: colorless solid LC/MS: cond. 2 RT 2.54 min LC/MS (ESI ⁺) m/z; 455 [M+H] ⁺
475	Morphology: colorless solid LC/MS: cond. 2 RT 2.51 min LC/MS (ESI ⁺) m/z; 467 [M+H] ⁺
476	Morphology: colorless solid LC/MS: cond. 2 RT 2.53 min LC/MS (ESI ⁺) m/z; 495 [M+H] ⁺
477	Morphology: colorless solid LC/MS: cond. 2 RT 2.43 min LC/MS (ESI ⁺) m/z; 451 [M+H] ⁺
478	Morphology: colorless solid LC/MS: cond. 2 RT 2.49 min LC/MS (ESI ⁺) m/z; 481 [M+H] ⁺
479	Morphology: brown solid LC/MS: cond. 2 RT 2.66 min LC/MS (ESI ⁺) m/z; 409 [M+H] ⁺
480a	Morphology: colorless solid LC/MS: cond. 2 RT 2.61 min LC/MS (ESI ⁺) m/z; 473 [M+H] ⁺
480b	Morphology: colorless solid LC/MS: cond. 2 RT 2.66 min LC/MS (ESI ⁺) m/z; 473 [M+H] ⁺
481	Morphology: colorless solid LC/MS: cond. 2 RT 1.96 min LC/MS (ESI ⁺) m/z; 480 [M+H] ⁺
482	Morphology: white solid LC/MS: cond. 2 RT 2.65 min LC/MS (ESI ⁺) m/z; 469 [M+H] ⁺
483a	Morphology: colorless solid LC/MS: cond. 2 RT 2.14 min LC/MS (ESI ⁺) m/z; 379 [M+H] ⁺
483b	Morphology: colorless solid LC/MS: cond. 2 RT 2.24 min LC/MS (ESI ⁺) m/z; 485 [M+H] ⁺
484	Morphology: colorless solid LC/MS: cond. 2 RT 2.75 min LC/MS (ESI ⁺) m/z; 435 [M+H] ⁺

[Table 42]

Ex	Data
485	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.39 min LC/MS (ESI ⁺) m/z; 513 [M+H] ⁺ LC/MS (ESI ⁻) m/z; 511 [M-H] ⁻
486	Morphology: colorless solid LC/MS: cond. 2 RT 2.68 min LC/MS (ESI ⁺) m/z; 523 [M+H] ⁺
487	Morphology: yellow amorphous LC/MS: cond. 2 RT 2.33 min LC/MS (ESI ⁺) m/z; 536 [M+H] ⁺
488	Morphology: colorless oil LC/MS: cond. 2 RT 2.67 min LC/MS (ESI ⁺) m/z; 576 [M+H] ⁺
489	Morphology: colorless solid LC/MS: cond. 2 RT 2.48 min LC/MS (ESI ⁺) m/z; 503 [M+H] ⁺
490	Morphology: colorless solid LC/MS: cond. 2 RT 2.55 min LC/MS (ESI ⁺) m/z; 493 [M+H] ⁺
491	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 1.50 min LC/MS (ESI ⁺) m/z; 465 [M+H] ⁺
492	Morphology: colorless amorphous LC/MS: cond. 2 RT 2.13 min LC/MS (ESI ⁺) m/z; 483 [M+H] ⁺
493	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.26 min LC/MS (ESI ⁺) m/z; 470 [M+H] ⁺
494	Morphology: pale brown solid LC/MS: cond. 2 RT 2.60 min LC/MS (ESI ⁺) m/z; 452 [M+H] ⁺
495	Morphology: pale yellow amorphous LC/MS: cond. 2 RT 2.67 min LC/MS (ESI ⁺) m/z; 454 [M+H] ⁺
496	Morphology: colorless solid LC/MS: cond. 2 RT 2.32 min LC/MS (ESI ⁺) m/z; 426 [M+H] ⁺
497	Morphology: colorless solid LC/MS: cond. 2 RT 2.53 min LC/MS (ESI ⁺) m/z; 472 [M+H] ⁺

[1125] Pharmacological analysis

5 The pharmacological analysis on the compound of the present invention will be described next.

1. Calcium influx inhibition assay

Human T-type calcium channel (Cav 3.2) expressing HEK293 cells were obtained from Prof. Edward Perez-Reyes, University of Virginia, USA. The calcium-sensitive
10 fluorescent dye used was FLIPR Calcium 4 Assay Kit (Molecular Devices). To a black 96-well plate with a clear bottom coated with type I collagen, the human T-type calcium channel (Cav 3.2) expressing HEK293 cells were seeded and cultured overnight, and the culture medium was removed. A solution of the calcium-sensitive fluorescent dye was

added, and the plate was incubated at 37°C in an atmosphere of 5% carbon dioxide for 30 minutes. To the plate, a diluted solution of a compound was added, and the whole was further incubated at 37°C in an atmosphere of 5% carbon dioxide for 30 minutes. While fluorescence was continuously analyzed from the bottom with a FlexStation 3 (Molecular
 5 Devices), a calcium solution was added. The increase in the fluorescence due to calcium influx caused by the stimulus was observed for 70 seconds. From the rise in the fluorescence from the base line, the inhibition percentage was calculated. The logarithms of compound concentrations were plotted with respect to the inhibition activities to give an IC₅₀ value.

10 The IC₅₀ values of all the compounds of Synthesis Examples showed 10 µM or less.

Table 43 shows the resulting T-type calcium channel inhibition concentrations of the compounds of Synthesis Examples.

Table 43

	Table 43	
	Synthesis Example No.	IC ₅₀ value (µM)
15	1	0.17
	3	0.10
	5	0.056
	6	0.019
	8	0.14
	15	0.28
	16	0.58
20	18	0.050
	21	0.061
	25	0.025
	27	0.15
	29	0.32

	31	0.23
	34	0.52
	38	0.075
	39	0.21
5	43	0.44
	49	0.86
	58	0.17
	59	0.80
	60	0.42
10	65	0.61
	66	0.088
	70	0.019
	71	0.024
	82	0.12
15	85	0.057
	90	0.019
	92	0.047
	96	0.11
	107	0.076
20	114	0.13
	127	0.030
	129	0.06
	130	0.077
	131	0.20
25	134	0.32
	140	0.055
	144	0.096
	153	0.051
	162	0.60

	178	0.060
	182	0.10
	183	0.041
	184	0.043
5	188	0.042
	189	0.012
	190	0.0033
	194	0.0099
	195	0.0036
10	196	0.020
	201	0.0057
	202	0.047
	207	0.0075
	208	0.0062
15	211	0.0020
	213	0.0034
	219	0.0055
	224	0.057
	230	0.022
20	256	0.042
	259	0.60
	260	0.14
	264	0.028
	265	0.57
25	269	0.075
	272	0.35
	275	0.030
	278	0.045
	283	0.51

	285	0.056
	295	0.18
	300	0.16
	304	0.064
5	307	0.27
	309	0.52
	310	0.17
	311	0.16
	312	0.55
10	314	0.046
	315	0.00046
	316	0.0023
	317	0.056
	318	0.033
15	321	0.0036
	326	0.0019
	327	0.0040
	329	0.0048
	333	0.0018
20	340	0.013
	342	0.00094
	344	0.012
	348	0.0045
	349	0.0011
25	350	0.0011
	357	0.013
	360	0.083
	362	0.0015
	363	0.0087

	365	0.013
	370	0.63
	377	0.41
	379	0.054
5	380	0.31
	382	0.86
	383	0.089
	392	0.0013
	395	0.0017
10	400	0.0044
	403	0.0020
	413	0.0069
	415	0.016
	423	0.040
15	428	0.0055
	430	0.16
	436	0.33
	439	0.0069
	448	0.016
20	450	0.046
	453	0.024
	454	0.0019
	457	0.013
	458b	0.053
25	460	0.061
	461	0.40
	462	0.61
	465	0.28
	466b	0.57

	468	0.068
	475	0.039
	476	0.061
	479	0.066
5	481	0.47
	484	0.24
	485	0.41
	486	0.41
	487	0.50
10	489	0.51
	493	0.0037

[1126] Formulation Example 1

15 Granules containing the following components are produced.

Component	Compound of Formula (I)	10 mg
	Lactose	700 mg
	Cornstarch	274 mg
	HPC-L	16 mg

20	-----	
	Total	1,000 mg

The compound of Formula (I) and lactose are sieved through a 60-mesh sieve. Cornstarch is sieved through a 120-mesh sieve. The sieved materials are mixed in a V-type mixer. To the mixed powder, an aqueous solution of low-viscosity

25 hydroxypropylcellulose (HPC-L) is added, then the whole is kneaded and granulated (extruding granulation, a pore size of 0.5 to 1 mm), and the granules are dried. The resulting dried granules are sieved through a vibrating screen (12/60 mesh) to give granules.

[1127] Formulation Example 2

Powders that are to be filled into capsules and contain the following components are produced.

5	Component	Compound of Formula (I)	10 mg
		Lactose	79 mg
		Cornstarch	10 mg
		Magnesium stearate	1 mg

		Total	100 mg

The compound of Formula (I) and lactose are sieved through a 60-mesh sieve.

- 10 Cornstarch is sieved through a 120-mesh sieve. The sieved materials and magnesium stearate are mixed in a V-type mixer. Into a No. 5 hard gelatin capsule, 100 mg of the 10% mixture is filled.

[1128] Formulation Example 3

- 15 Granules that are to be filled into capsules and contain the following components are produced.

	Component	Compound of Formula (I)	15 mg
		Lactose	90 mg
		Cornstarch	42 mg
		HPC-L	3 mg

20		Total	150 mg

The compound of Formula (I) and lactose are sieved through a 60-mesh sieve.

- Cornstarch is sieved through a 120-mesh sieve. The sieved materials are mixed in a V-type mixer. To the mixed powder, an aqueous solution of low-viscosity
 25 hydroxypropylcellulose (HPC-L) is added, then the whole is kneaded and granulated, and the granules are dried. The resulting dried granules are sieved through a vibrating screen (12/60 mesh), and 150 mg of the sieved granules are filled into a No. 4 hard gelatin capsule.

[1129] Formulation Example 4

Tablets containing the following components are produced.

5	Component	Compound of Formula (I)	10 mg
		Lactose	90 mg
		Microcrystalline cellulose	30 mg
		Magnesium stearate	5 mg
		CMC-Na	15 mg

		Total	150 mg

The compound of Formula (I), lactose, microcrystalline cellulose, and CMC-Na (sodium carboxymethylcellulose) are sieved through a 60-mesh sieve and mixed. To the mixed powder, magnesium stearate is added to give a mixed powder for formulation. The mixed powder is directly compressed into 150-mg tablets.

[1130] Formulation Example 5

An intravenous formulation is prepared as follows:

15	Compound of Formula (I)	100 mg
	Saturated fatty acid glyceride	1,000 mL

The formulated solution is intravenously administered to a patient at a speed of 1 mL/min.


20 INDUSTRIAL APPLICABILITY

[1131] The compound of the present invention has an excellent T-type calcium channel inhibitory activity and is specifically useful for prevention or treatment of pains, such as chronic pains and acute pains including neuropathic pain, inflammatory pain, cancer pain, and visceral pain, which are caused by diabetic neuropathy, postherpetic neuralgia, trigeminal neuralgia, phantom limb pain, postoperative pain, stump pain, traumatic neurological disorder, carpal tunnel syndrome, plexus neuropathy, glossopharyngeal neuralgia, laryngeal neuralgia, migraine, fibromyalgia syndrome, rheumatoid arthritis, multiple sclerosis, HIV, herpes simplex, syphilis, carcinomatous

neuropathy, polyneuropathy, causalgia, low back pain, complex regional pain syndrome (CRPS), thalamic pain, osteoarthritis, spinal cord injury, and cerebral apoplexy.

CLAIMS

1. A compound of Formula (I):



5 [wherein

R¹ is

a hydrogen atom,

a halogen atom,

a C₁₋₆ alkyl group,

10 a C₁₋₆ alkoxy group,

a C₁₋₆ alkylthio group,

a mono-C₁₋₆ alkylamino group,

a di-C₁₋₆ alkylamino group

(the C₁₋₆ alkyl group, the C₁₋₆ alkoxy group, the C₁₋₆ alkylthio group, the mono-C₁₋₆

15 alkylamino group, and the di-C₁₋₆ alkylamino group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁸),

or

a C₃₋₁₁ cycloalkyl group

(the C₃₋₁₁ cycloalkyl group is unsubstituted or substituted with one or more substituents

20 identically or differently selected from the substituent group V⁶);

each of L¹ and L² is independently

a single bond,

NR²,

O,

25 S,

SO,

SO₂, or

a C₁₋₆ alkylene group

(the C₁₋₆ alkylene group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁶, and a single methylene group of the C₁₋₆ alkylene group is optionally replaced by O, S, SO₂, C=O, C=S, or NR³);

5 B is

a C₃₋₁₁ cycloalkylene group,

a C₃₋₁₁ cycloalkenylene group,

a 3 to 11-membered heterocyclylene group,

a C₆₋₁₄ arylene group,

10 a 5 to 10-membered heteroarylene group

(the C₃₋₁₁ cycloalkylene group, the C₃₋₁₁ cycloalkenylene group, the 3 to 11-membered heterocyclylene group, the C₆₋₁₄ arylene group, and the 5 to 10-membered heteroarylene group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁶, and

15 a single methylene group of the C₃₋₁₁ cycloalkylene group or the C₃₋₁₁ cycloalkenylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group),

a C₁₋₆ alkylene group,

a C₂₋₆ alkenylene group, or

a C₂₋₆ alkynylene group

20 (the C₁₋₆ alkylene group, the C₂₋₆ alkenylene group, and the C₂₋₆ alkynylene group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁸, and

a single methylene group of the C₁₋₆ alkylene group, the C₂₋₆ alkenylene group, or the C₂₋₆ alkynylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group);

25 when L¹ is a single bond, O, or a C₁₋₆ alkylene group and L² is a single bond or a C₁₋₆ alkylene group, B is not a C₁₋₆ alkylene group, a C₂₋₆ alkenylene group, a C₂₋₆ alkynylene group, or a C₆₋₁₄ arylene group; when L¹ is a single bond, O, or a C₁₋₆ alkylene group and L² is NR², O, S, SO, or SO₂, B is not a C₁₋₆ alkylene group, a C₂₋₆ alkenylene group, or a C₂₋₆ alkynylene group;

A is

a C₁₋₆ alkyl group,

a C₂₋₆ alkenyl group

(the C₁₋₆ alkyl group and the C₂₋₆ alkenyl group are unsubstituted or substituted with one

5 or more substituents identically or differently selected from the substituent group V⁸), a

C₃₋₁₁ cycloalkyl group,

a C₃₋₁₁ cycloalkenyl group,

a 3 to 11-membered heterocyclyl group,

a C₆₋₁₄ aryl group, or

10 a 5 to 10-membered heteroaryl group

(the C₃₋₁₁ cycloalkyl group, the C₃₋₁₁ cycloalkenyl group, the 3 to 11-membered

heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are

unsubstituted or substituted with one or more substituents identically or differently

selected from the substituent group V⁶);

15 L³ is

a C₁₋₆ alkylene group

(the C₁₋₆ alkylene group is unsubstituted or substituted with one or more substituents

identically or differently selected from the substituent group V⁸, and a single methylene

group of the C₁₋₆ alkylene group is optionally replaced by C=O or C≡S);

20 D is

a C₃₋₁₁ cycloalkyl group,

a C₃₋₁₁ cycloalkenyl group,

a 3 to 11-membered heterocyclyl group,

a C₆₋₁₄ aryl group, or

25 a 5 to 10-membered heteroaryl group

(the C₃₋₁₁ cycloalkyl group, the C₃₋₁₁ cycloalkenyl group, the 3 to 11-membered

heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are

unsubstituted or substituted with one or more substituents identically or differently

selected from the substituent group V⁶);

each of R^2 and R^3 is independently

a hydrogen atom,

a C₁₋₆ alkyl group,

a C₂₋₆ alkenyl group

5 (the C₁₋₆ alkyl group and the C₂₋₆ alkenyl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁸),

a C₃₋₁₁ cycloalkyl group,

a 3 to 11-membered heterocyclcyl group,

a C₆₋₁₄ aryl group, or

10 a 5 to 10-membered heteroaryl group

(the C₃₋₁₁ cycloalkyl group, the 3 to 11-membered heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁶);

the substituent group V^6 is a substituent group consisting of substituents

15 constituting the substituent group V⁸, C₁₋₆ alkyl groups, C₂₋₆ alkenyl groups, and C₂₋₆ alkynyl groups (the C₁₋₆ alkyl groups, the C₂₋₆ alkenyl groups, and the C₂₋₆ alkynyl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V¹);

the substituent group V^8 is a substituent group consisting of substituents

constituting the substituent group V^a, C₁₋₆ alkoxy groups, C₁₋₆ alkylthio groups, C₁₋₆ alkylcarbonyl groups, C₁₋₆ alkylsulfonyl groups, C₁₋₆ alkoxy carbonyl groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, mono-C₁₋₆ alkylaminocarbonyl groups, di-C₁₋₆ alkylaminocarbonyl groups, mono-C₁₋₆ alkylaminosulfonyl groups, di-C₁₋₆ alkylaminosulfonyl groups, C₁₋₆ alkylcarbonylamino groups, C₁₋₆ alkylcarbonyloxy groups, C₁₋₆ alkylsulfonylamino groups, C₁₋₆ alkylsulfonyloxy groups (the C₁₋₆ alkoxy groups, the C₁₋₆ alkylthio groups, the C₁₋₆ alkylcarbonyl groups, the C₁₋₆ alkylsulfonyl groups, the C₁₋₆ alkoxy carbonyl groups, the mono-C₁₋₆ alkylamino groups, the di-C₁₋₆ alkylamino groups, the mono-C₁₋₆ alkylaminocarbonyl groups, the di-C₁₋₆ alkylaminocarbonyl groups, the mono-C₁₋₆ alkylaminosulfonyl groups, the di-C₁₋₆

alkylaminosulfonyl groups, the C₁₋₆ alkylcarbonylamino groups, the C₁₋₆ alkylcarbonyloxy groups, the C₁₋₆ alkylsulfonylamino groups, and the C₁₋₆ alkylsulfonyloxy groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V¹), C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylsulfonylamino groups, C₃₋₆ cycloalkylsulfonyloxy groups, C₃₋₆ cycloalkylthio groups, C₃₋₁₁ cycloalkyl groups, 3 to 11-membered heterocyclyl groups, C₆₋₁₄ aryl groups, and 5 to 10-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylsulfonylamino groups, the C₃₋₆ cycloalkylsulfonyloxy groups, the C₃₋₆ cycloalkylthio groups, the C₃₋₁₁ cycloalkyl groups, the 3 to 11-membered heterocyclyl groups, the C₆₋₁₄ aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V²);

the substituent group V^a is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, an amino group, a carboxy group, a carbamoyl group, a sulfamoyl group, a phosphono group, a sulfo group, a tetrazolyl group, a formate group, and a formyl group;

the substituent group V¹ is a substituent group consisting of substituents constituting the substituent group V^a, C₁₋₆ alkoxy groups, C₁₋₃ haloalkoxy groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, mono-C₁₋₆ alkylaminocarbonyl groups, di-C₁₋₆ alkylaminocarbonyl groups, C₁₋₆ alkylcarbonylamino groups, C₁₋₆ alkylthio groups, C₁₋₆ alkylsulfonyl groups, C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, 3 to 11-membered heterocyclyl groups, C₆₋₁₄ aryl groups, and 5 to 10-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆

cycloalkylthio groups, the 3 to 11-membered heterocyclyl groups, the C₆₋₁₄ aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more hydroxy groups, one or more halogen atoms, one or more cyano groups, one or more nitro groups, one or more amino groups, one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one or more phosphinoyl groups, one or more sulfo groups, one or more sulfinio groups, one or more tetrazolyl groups, one or more formyl groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups); and

the substituent group V² is a substituent group consisting of substituents constituting the substituent group V¹, C₁₋₆ alkyl groups, and C₁₋₃ haloalkyl groups], a tautomer of the compound, a pharmaceutically acceptable salt thereof, or a solvate thereof.

2. The compound according to claim 1, wherein

L³ is a C₁₋₃ alkylene group,

the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

3. The compound according to claim 1 or 2, wherein

R¹ is a hydrogen atom or a C₁₋₆ alkoxy group (the C₁₋₆ alkoxy group is unsubstituted or substituted with one or more substituents identically or differently selected from a substituent group V⁷);

the substituent group V⁷ is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C₁₋₆ alkoxy groups, C₁₋₆ alkylthio groups, C₁₋₆ alkoxycarbonyl groups (the C₁₋₆ alkoxy groups, the C₁₋₆ alkylthio groups, and the

- C_{1-6} alkoxycarbonyl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^1), C_{3-6} cycloalkoxy groups, mono- C_{3-6} cycloalkylamino groups, di- C_{3-6} cycloalkylamino groups, C_{3-6} cycloalkylcarbonyl groups, C_{3-6} cycloalkylsulfonyl groups, C_{3-6} cycloalkylthio groups,
- 5 C_{3-6} cycloalkyl groups, 4 to 7-membered heterocyclyl groups, a phenyl group, and 5 to 6-membered heteroaryl groups (the C_{3-6} cycloalkoxy groups, the mono- C_{3-6} cycloalkylamino groups, the di- C_{3-6} cycloalkylamino groups, the C_{3-6} cycloalkylcarbonyl groups, the C_{3-6} cycloalkylsulfonyl groups, the C_{3-6} cycloalkylthio groups, the C_{3-6} cycloalkyl groups, the 4 to 7-membered heterocyclyl groups, the phenyl group, and the 5
- 10 to 6-membered heteroaryl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V^2);

- the substituent group V^1 is a substituent group consisting of substituents constituting the substituent group V^a , C_{1-6} alkoxy groups, C_{1-3} haloalkoxy groups, mono- C_{1-6} alkylamino groups, di- C_{1-6} alkylamino groups, mono- C_{1-6} alkylaminocarbonyl
- 15 groups, di- C_{1-6} alkylaminocarbonyl groups, C_{1-6} alkylcarbonylamino groups, C_{1-6} alkylthio groups, C_{1-6} alkylsulfonyl groups, C_{3-6} cycloalkoxy groups, mono- C_{3-6} cycloalkylamino groups, di- C_{3-6} cycloalkylamino groups, C_{3-6} cycloalkylcarbonyl groups, C_{3-6} cycloalkylsulfonyl groups, C_{3-6} cycloalkylthio groups, 3 to 11-membered heterocyclyl groups, C_{6-14} aryl groups, and 5 to 10-membered heteroaryl groups (the C_{3-6}
- 20 cycloalkoxy groups, the mono- C_{3-6} cycloalkylamino groups, the di- C_{3-6} cycloalkylamino groups, the C_{3-6} cycloalkylcarbonyl groups, the C_{3-6} cycloalkylsulfonyl groups, the C_{3-6} cycloalkylthio groups, the 3 to 11-membered heterocyclyl groups, the C_{6-14} aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more hydroxy groups, one or more halogen atoms, one or more cyano groups, one or
- 25 more nitro groups, one or more amino groups, one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one or more phosphinoyl groups, one or more sulfo groups, one or more sulfinio groups, one or more tetrazolyl groups, one or more formyl groups, one or more C_{1-6} alkyl groups, one or more C_{1-3} haloalkyl groups, one or more C_{1-6} alkoxy groups, one or more C_{1-3} haloalkoxy

groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups);

5 the substituent group V² is a substituent group consisting of substituents constituting the substituent group V¹, C₁₋₆ alkyl groups, and C₁₋₃ haloalkyl groups; and

 the substituent group V^a is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, an amino group, a carboxy group, a carbamoyl group, a sulfamoyl group, a phosphono group, a sulfo group, a tetrazolyl
10 group, a formate group, and a formyl group,

the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

4. The compound according to claim 3, wherein

15 R¹ is a hydrogen atom,
the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

5. The compound according to claim 1 or 2, wherein

20 R¹ is a C₁₋₆ alkyl group (the C₁₋₆ alkyl group is unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁹);

 the substituent group V⁹ is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C₁₋₆ alkoxy groups, C₁₋₆ alkylthio groups, C₁₋₆ alkylcarbonyloxy groups, C₁₋₆ alkoxycarbonyl groups (the C₁₋₆ alkoxy groups, the
25 C₁₋₆ alkylthio groups, the C₁₋₆ alkylcarbonyloxy groups, and the C₁₋₆ alkoxycarbonyl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V¹), C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, C₃₋₆ cycloalkyl groups, 4 to

7-membered heterocyclyl groups, a phenyl group, and 5 to 6-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the C₃₋₆ cycloalkyl groups, the 4 to 7-membered heterocyclyl groups, the phenyl group, and the 5 to 6-membered heteroaryl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V²);

the substituent group V¹ is a substituent group consisting of substituents constituting the substituent group V^a, C₁₋₆ alkoxy groups, C₁₋₃ haloalkoxy groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, mono-C₁₋₆ alkylaminocarbonyl groups, di-C₁₋₆ alkylaminocarbonyl groups, C₁₋₆ alkylcarbonylamino groups, C₁₋₆ alkylthio groups, C₁₋₆ alkylsulfonyl groups, C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, 3 to 11-membered heterocyclyl groups, C₆₋₁₄ aryl groups, and 5 to 10-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the 3 to 11-membered heterocyclyl groups, the C₆₋₁₄ aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more hydroxy groups, one or more halogen atoms, one or more cyano groups, one or more nitro groups, one or more amino groups, one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one or more phosphinoyl groups, one or more sulfo groups, one or more sulfinio groups, one or more tetrazolyl groups, one or more formyl groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups);

the substituent group V^2 is a substituent group consisting of substituents constituting the substituent group V^1 , C_{1-6} alkyl groups, and C_{1-3} haloalkyl groups; and

the substituent group V^a is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, an amino group, a carboxy group, a carbamoyl group, a sulfamoyl group, a phosphono group, a sulfo group, a tetrazolyl group, a formate group, and a formyl group,
 5 the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

10 6. The compound according to claim 5, wherein

R^1 is a C_{1-6} alkyl group (the C_{1-6} alkyl group is unsubstituted or substituted with one or more halogen atoms),
 the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

15

7. The compound according to any one of claims 1 to 6, wherein

L^1 is a single bond, NR^{2a} , O, S, SO, SO_2 , or a C_{1-6} alkylene group (a single methylene group of the C_{1-6} alkylene group is optionally replaced by O, S, SO_2 , C=O, C=S, or NR^{3a});

20 L^2 is a single bond and B is a 3 to 11-membered heterocyclylene group or a 5 to 10-membered heteroarylene group (the 3 to 11-membered heterocyclylene group and the 5 to 10-membered heteroarylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^3),
 or

25 L^2 is NR^{2b} , O, S, SO, SO_2 , or a C_{1-6} alkylene group (a single methylene group of the C_{1-6} alkylene group is optionally replaced by O, S, SO_2 , C=O, C=S, or NR^{3b}) and B is a C_{3-11} cycloalkylene group, a C_{3-11} cycloalkenylene group, a 3 to 11-membered heterocyclylene group, or a 5 to 10-membered heteroarylene group (the C_{3-11} cycloalkylene group, the C_{3-11} cycloalkenylene group, the 3 to 11-membered

heterocyclylene group, and the 5 to 10-membered heteroarylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V^5 , and a single methylene group of the C_{3-11} cycloalkylene group and the C_{3-11} cycloalkenylene group is optionally replaced by a 1,1- C_{3-7}

5 cycloalkylene group);

the substituent group V^5 is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C_{1-6} alkyl groups, C_{1-6} alkoxy groups, C_{1-6} alkylthio groups, C_{1-6} alkoxycarbonyl groups (the C_{1-6} alkyl groups, the C_{1-6} alkoxy groups, the C_{1-6} alkylthio groups, and the C_{1-6} alkoxycarbonyl groups are unsubstituted or
 10 substituted with one or more substituents identically or differently selected from the substituent group V^1), C_{3-6} cycloalkoxy groups, mono- C_{3-6} cycloalkylamino groups, di- C_{3-6} cycloalkylamino groups, C_{3-6} cycloalkylcarbonyl groups, C_{3-6} cycloalkylsulfonyl groups, C_{3-6} cycloalkylthio groups, C_{3-6} cycloalkyl groups, 4 to 7-membered heterocyclyl groups, a phenyl group, and 5 to 6-membered heteroaryl groups (the C_{3-6} cycloalkoxy
 15 groups, the mono- C_{3-6} cycloalkylamino groups, the di- C_{3-6} cycloalkylamino groups, the C_{3-6} cycloalkylcarbonyl groups, the C_{3-6} cycloalkylsulfonyl groups, the C_{3-6} cycloalkylthio groups, the C_{3-6} cycloalkyl groups, the 4 to 7-membered heterocyclyl groups, the phenyl group, and the 5 to 6-membered heteroaryl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the
 20 substituent group V^2);

the substituent group V^1 is a substituent group consisting of substituents constituting the substituent group V^a , C_{1-6} alkoxy groups, C_{1-3} haloalkoxy groups, mono- C_{1-6} alkylamino groups, di- C_{1-6} alkylamino groups, mono- C_{1-6} alkylaminocarbonyl groups, di- C_{1-6} alkylaminocarbonyl groups, C_{1-6} alkylcarbonylamino groups, C_{1-6}
 25 alkylthio groups, C_{1-6} alkylsulfonyl groups, C_{3-6} cycloalkoxy groups, mono- C_{3-6} cycloalkylamino groups, di- C_{3-6} cycloalkylamino groups, C_{3-6} cycloalkylcarbonyl groups, C_{3-6} cycloalkylsulfonyl groups, C_{3-6} cycloalkylthio groups, 3 to 11-membered heterocyclyl groups, C_{6-14} aryl groups, and 5 to 10-membered heteroaryl groups (the C_{3-6} cycloalkoxy groups, the mono- C_{3-6} cycloalkylamino groups, the di- C_{3-6} cycloalkylamino

groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the 3 to 11-membered heterocyclyl groups, the C₆₋₁₄ aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more hydroxy groups, one or more halogen atoms, one or more cyano groups, one or more nitro groups, one or more amino groups, one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one or more phosphinoyl groups, one or more sulfo groups, one or more sulfinio groups, one or more tetrazolyl groups, one or more formyl groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups);

the substituent group V² is a substituent group consisting of substituents constituting the substituent group V¹, C₁₋₆ alkyl groups, and C₁₋₃ haloalkyl groups;

the substituent group V^a is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, an amino group, a carboxy group, a carbamoyl group, a sulfamoyl group, a phosphono group, a sulfo group, a tetrazolyl group, a formate group, and a formyl group; and

each of R^{2a}, R^{2b}, R^{3a}, and R^{3b} is independently a hydrogen atom, a C₁₋₃ alkyl group, or a C₁₋₃ haloalkyl group,

the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

8. The compound according to claim 7, wherein

L¹ is a single bond;

L² is a single bond, and B is a 4 to 7-membered heterocyclylene group (the 4 to 7-membered heterocyclylene group is unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V³, and a

single methylene group of the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group), or

L^2 is NR^{2c}, O, or a C₁₋₆ alkylene group (a single methylene group of the C₁₋₆ alkylene group is optionally replaced by O or NR^{3c}), and B is a C₃₋₆ cycloalkylene group, a C₃₋₆ cycloalkenylene group, or a 4 to 7-membered heterocyclylene group (the C₃₋₆ cycloalkylene group, the C₃₋₆ cycloalkenylene group, and the 4 to 7-membered heterocyclylene group are unsubstituted or optionally substituted with one or more substituents identically or differently selected from the substituent group V³, and a single methylene group of the C₃₋₆ cycloalkylene group, the C₃₋₆ cycloalkenylene group, and the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group);

the substituent group V³ is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C₁₋₆ alkyl groups, C₁₋₆ haloalkyl groups, C₃₋₆ cycloalkyl groups, C₁₋₆ alkoxy groups, and C₁₋₆ haloalkoxy groups; and

each of R^{2c} and R^{3c} is independently a hydrogen atom, a C₁₋₃ alkyl group, or a C₁₋₃ haloalkyl group,

the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

9. The compound according to any one of claims 1 to 6, wherein

L^1 is a single bond;

L^2 is a single bond, and B is a 4 to 7-membered heterocyclylene group (the 4 to 7-membered heterocyclylene group is substituted with one or more substituents selected from an amino group, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, and C₁₋₆ alkylsulfonylamino groups and is optionally substituted with one or more substituents identically or differently selected from the substituent group V³, and a single methylene group of the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group), or

L^2 is NR^{2c}, O, or a C₁₋₆ alkylene group (a single methylene group of the C₁₋₆

alkylene group is optionally replaced by O or NR^{3c}), and B is a C₃₋₆ cycloalkylene group, a C₃₋₆ cycloalkenylene group, or a 4 to 7-membered heterocyclylene group (the C₃₋₆ cycloalkylene group, the C₃₋₆ cycloalkenylene group, and the 4 to 7-membered heterocyclylene group are substituted with a substituent selected from one or more amino groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, and one or more C₁₋₆ alkylsulfonylamino groups and are optionally substituted with one or more substituents identically or differently selected from the substituent group V³, and a single methylene group of the C₃₋₆ cycloalkylene group, the C₃₋₆ cycloalkenylene group, and the 4 to 7-membered heterocyclylene group is optionally replaced by a 1,1-C₃₋₇ cycloalkylene group);

the substituent group V³ is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C₁₋₆ alkyl groups, C₁₋₆ haloalkyl groups, C₃₋₆ cycloalkyl groups, C₁₋₆ alkoxy groups, and C₁₋₆ haloalkoxy groups; and

each of R^{2c} and R^{3c} is independently a hydrogen atom, a C₁₋₃ alkyl group, or a C₁₋₃ haloalkyl group, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

10. The compound according to any one of claims 1 to 9, wherein

D is a 3 to 11-membered heterocyclyl group, a C₆₋₁₄ aryl group, or a 5 to 10-membered heteroaryl group (the 3 to 11-membered heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁵);

the substituent group V⁵ is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C₁₋₆ alkyl groups, C₁₋₆ alkoxy groups, C₁₋₆ alkylthio groups, C₁₋₆ alkoxycarbonyl groups (the C₁₋₆ alkyl groups, the C₁₋₆ alkoxy groups, the C₁₋₆ alkylthio groups, and the C₁₋₆ alkoxycarbonyl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V¹), C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups,

di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, C₃₋₆ cycloalkyl groups, 4 to 7-membered heterocyclyl groups, a phenyl group, and 5 to 6-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the C₃₋₆ cycloalkyl groups, the 4 to 7-membered heterocyclyl groups, the phenyl group, and the 5 to 6-membered heteroaryl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V²);

10 the substituent group V¹ is a substituent group consisting of substituents constituting the substituent group V^a, C₁₋₆ alkoxy groups, C₁₋₃ haloalkoxy groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, mono-C₁₋₆ alkylaminocarbonyl groups, di-C₁₋₆ alkylaminocarbonyl groups, C₁₋₆ alkylcarbonylamino groups, C₁₋₆ alkylthio groups, C₁₋₆ alkylsulfonyl groups, C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, 3 to 11-membered heterocyclyl groups, C₆₋₁₄ aryl groups, and 5 to 10-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the 3 to 11-membered heterocyclyl groups, the C₆₋₁₄ aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more hydroxy groups, one or more halogen atoms, one or more cyano groups, one or more nitro groups, one or more amino groups, one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one or more phosphinoyl groups, one or more sulfo groups, one or more sulfinio groups, one or more tetrazolyl groups, one or more formyl groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆

alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups);

the substituent group V² is a substituent group consisting of substituents constituting the substituent group V¹, C₁₋₆ alkyl groups, and C₁₋₃ haloalkyl groups; and

5 the substituent group V^a is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, an amino group, a carboxy group, a carbamoyl group, a sulfamoyl group, a phosphono group, a sulfo group, a tetrazolyl group, a formate group, and a formyl group,
the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate
10 thereof.

11. The compound according to claim 10, wherein

D is a 4 to 7-membered heterocyclyl group, a phenyl group, or a 5 to 6-membered heteroaryl group (the 4 to 7-membered heterocyclyl group, the phenyl group, and the 5 to
15 6-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from a substituent group V⁴);

the substituent group V⁴ is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C₁₋₆ alkyl groups, C₁₋₆ haloalkyl groups, C₁₋₆ alkoxy groups, C₁₋₆ haloalkoxy groups, C₃₋₆ cycloalkyl groups, C₃₋₆ cycloalkoxy groups,
20 mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, and C₃₋₆ cycloalkylthio groups (the C₁₋₆ alkyl groups, the C₁₋₆ alkoxy groups, the C₃₋₆ cycloalkyl groups, the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, and the C₃₋₆ cycloalkylthio groups are unsubstituted or substituted with one or more hydroxy groups, one or more amino groups, one or more
25 nitro groups, one or more cyano groups, one or more 3 to 11-membered heterocyclyl groups, one or more C₆₋₁₄ aryl groups, or one or more 5 to 10-membered heteroaryl groups (the 3 to 11-membered heterocyclyl groups, the C₆₋₁₄ aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or

more phosphono groups, one or more sulfo groups, one or more tetrazolyl groups, one or more formyl groups, one or more nitro groups, one or more cyano groups, one or more halogen atoms, one or more hydroxy groups, one or more amino groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups)), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

12. The compound according to claim 11, wherein

D is a phenyl group or a 5 to 6-membered heteroaryl group (the phenyl group or the 5 to 6-membered heteroaryl group is unsubstituted or substituted with one or more halogen atoms, one or more nitro groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₆ haloalkyl groups, one or more C₁₋₆ alkoxy groups, or one or more C₁₋₆ haloalkoxy groups), the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

13. The compound according to any one of claims 1 to 9, wherein

D is a phenyl group or a 5 to 6-membered heteroaryl group (the phenyl group or the 5 to 6-membered heteroaryl group is substituted with one or more C₁₋₆ alkylsulfonylamino groups or one or more C₁₋₆ alkylsulfonyloxy groups (the C₁₋₆ alkylsulfonylamino groups and the C₁₋₆ alkylsulfonyloxy groups are unsubstituted or substituted with one or more halogen atoms, one or more nitro groups, one or more C₁₋₆ alkoxy groups, or one or more C₁₋₆ haloalkoxy groups) and is optionally substituted with one or more halogen atoms, one or more nitro groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₆ haloalkyl groups, one or more C₁₋₆ alkoxy groups, or one or more C₁₋₆

haloalkoxy groups),

the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

5 14. The compound according to claim 13, wherein

D is a 5 to 6-membered heteroaryl group (the 5 to 6-membered heteroaryl group is substituted with one or more C₁₋₆ alkylsulfonyloxy groups (the C₁₋₆ alkylsulfonyloxy groups are unsubstituted or substituted with one or more halogen atoms, one or more nitro groups, one or more C₁₋₆ alkoxy groups, or one or more C₁₋₆ haloalkoxy groups)),

10 the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof.

15. The compound according to any one of claims 1 to 14, wherein

A is a 3 to 11-membered heterocyclyl group, a C₆₋₁₄ aryl group, or a 5 to
15 10-membered heteroaryl group (the 3 to 11-membered heterocyclyl group, the C₆₋₁₄ aryl group, and the 5 to 10-membered heteroaryl group are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V⁵);
and

the substituent group V⁵ is a substituent group consisting of a hydroxy group,
20 halogen atoms, a cyano group, a nitro group, C₁₋₆ alkyl groups, C₁₋₆ alkoxy groups, C₁₋₆ alkylthio groups, C₁₋₆ alkoxycarbonyl groups (the C₁₋₆ alkyl groups, the C₁₋₆ alkoxy groups, the C₁₋₆ alkylthio groups, and the C₁₋₆ alkoxycarbonyl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V¹), C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups,
25 di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, C₃₋₆ cycloalkyl groups, 4 to 7-membered heterocyclyl groups, a phenyl group, and 5 to 6-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆

cycloalkylthio groups, the C₃₋₆ cycloalkyl groups, the 4 to 7-membered heterocyclyl groups, the phenyl group, and the 5 to 6-membered heteroaryl groups are unsubstituted or substituted with one or more substituents identically or differently selected from the substituent group V²);

- 5 the substituent group V¹ is a substituent group consisting of substituents constituting the substituent group V^a, C₁₋₆ alkoxy groups, C₁₋₃ haloalkoxy groups, mono-C₁₋₆ alkylamino groups, di-C₁₋₆ alkylamino groups, mono-C₁₋₆ alkylaminocarbonyl groups, di-C₁₋₆ alkylaminocarbonyl groups, C₁₋₆ alkylcarbonylamino groups, C₁₋₆ alkylthio groups, C₁₋₆ alkylsulfonyl groups, C₃₋₆ cycloalkoxy groups, mono-C₃₋₆ cycloalkylamino groups, di-C₃₋₆ cycloalkylamino groups, C₃₋₆ cycloalkylcarbonyl groups, C₃₋₆ cycloalkylsulfonyl groups, C₃₋₆ cycloalkylthio groups, 3 to 11-membered heterocyclyl groups, C₆₋₁₄ aryl groups, and 5 to 10-membered heteroaryl groups (the C₃₋₆ cycloalkoxy groups, the mono-C₃₋₆ cycloalkylamino groups, the di-C₃₋₆ cycloalkylamino groups, the C₃₋₆ cycloalkylcarbonyl groups, the C₃₋₆ cycloalkylsulfonyl groups, the C₃₋₆ cycloalkylthio groups, the 3 to 11-membered heterocyclyl groups, the C₆₋₁₄ aryl groups, and the 5 to 10-membered heteroaryl groups are unsubstituted or substituted with one or more hydroxy groups, one or more halogen atoms, one or more cyano groups, one or more nitro groups, one or more amino groups, one or more carboxy groups, one or more carbamoyl groups, one or more sulfamoyl groups, one or more phosphono groups, one or more phosphinoyl groups, one or more sulfo groups, one or more sulfinio groups, one or more tetrazolyl groups, one or more formyl groups, one or more C₁₋₆ alkyl groups, one or more C₁₋₃ haloalkyl groups, one or more C₁₋₆ alkoxy groups, one or more C₁₋₃ haloalkoxy groups, one or more mono-C₁₋₆ alkylamino groups, one or more di-C₁₋₆ alkylamino groups, one or more mono-C₁₋₆ alkylaminocarbonyl groups, one or more di-C₁₋₆ alkylaminocarbonyl groups, one or more C₁₋₆ alkylcarbonylamino groups, one or more C₁₋₆ alkylthio groups, or one or more C₁₋₆ alkylsulfonyl groups);

 the substituent group V² is a substituent group consisting of substituents constituting the substituent group V¹, C₁₋₆ alkyl groups, and C₁₋₃ haloalkyl groups; and

 the substituent group V^a is a substituent group consisting of a hydroxy group,

halogen atoms, a cyano group, a nitro group, an amino group, a carboxy group, a carbamoyl group, a sulfamoyl group, a phosphono group, a sulfo group, a tetrazolyl group, a formate group, and a formyl group,
the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate
5 thereof.

16. The compound according to claim 15, wherein

A is a phenyl group or a 5 to 6-membered heteroaryl group (the phenyl group and the 5 to 6-membered heteroaryl group are unsubstituted or substituted with one or more
10 substituents identically or differently selected from the substituent group V^3); and

the substituent group V^3 is a substituent group consisting of a hydroxy group, halogen atoms, a cyano group, a nitro group, C_{1-6} alkyl groups, C_{1-6} haloalkyl groups, C_{3-6} cycloalkyl groups, C_{1-6} alkoxy groups, and C_{1-6} haloalkoxy groups,
the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate
15 thereof.

17. A T-type calcium channel inhibitor comprising:

the compound, the tautomer of the compound, the pharmaceutically acceptable salt thereof, or the solvate thereof as claimed in any one of claims 1 to 16, as an active
20 component.

18. A preventive agent, a therapeutic agent, and/or an improving agent for a disease treatable by a T-type calcium channel inhibitory action, the preventive agent, the therapeutic agent, and/or the improving agent comprising:

25 the T-type calcium channel inhibitor as claimed in claim 17 as an active component.

19. A therapeutic agent for neuropathic pain, the therapeutic agent comprising:

the T-type calcium channel inhibitor as claimed in claim 18 as an active

component.

20. A medicine comprising:

the compound, the tautomer of the compound, the pharmaceutically acceptable
5 salt thereof, or the solvate thereof as claimed in any one of claims 1 to 19, as an active
component.