

ARYL HYDROCARBON RECEPTOR ANTAGONISTS AND USES THEREOF

5

Cross-Reference to Related Applications

This application claims priority to, and the benefit of, U.S. Application Nos. 62/484,692, filed April 12, 2017, 62/613,382, filed January 3, 2018, and 62/625,896, filed February 2, 2018, the entire contents of each of which are incorporated herein by reference.

10

Field

The present disclosure relates to aryl hydrocarbon receptor antagonists useful, for example, for ex vivo expansion, enrichment, and maintenance of hematopoietic stem cells, as well as methods of treating various hematopoietic pathologies by administration of the expanded hematopoietic stem cells.

15

Background

While hematopoietic stem cells have significant therapeutic potential, a limitation that has hindered their clinical use has been the difficulty associated with obtaining sufficient numbers of these cells. In particular, hematopoietic stem cells are resistant to maintenance, propagation, and expansion ex vivo. Another challenge to be overcome in order to further develop the use of hematopoietic stem cells (HSCs) as a therapeutic modality is the loss of multi-potency that can occur when these cells are cultured ex vivo.

20

There is currently a need for compositions and methods for the ex vivo maintenance, propagation, and expansion of HSCs that preserve the multi-potency and hematopoietic functionality of such cells.

25

Summary

The present disclosure features aryl hydrocarbon receptor antagonists, such as substituted imidazopyridines and imidazopyrazines, as well as methods of expanding hematopoietic stem cells by culturing hematopoietic stem cells in the presence of such agents. Additionally described herein are methods of chemically synthesizing aryl hydrocarbon receptor antagonists, such as substituted imidazopyridines and imidazopyrazines, as well as kits containing aryl hydrocarbon receptor antagonists that can be used for the expansion of hematopoietic stem cells. Additionally, the disclosure provides methods of treating various hematopoietic pathologies in a patient by administration of expanded hematopoietic stem cells. The patient may be suffering, for example, from a hemoglobinopathy or another disease of a cell in the hematopoietic lineage, and is thus in need of hematopoietic stem cell transplantation. As described herein, hematopoietic stem cells are capable of differentiating into a multitude of cell types in the hematopoietic family, and can be administered to a patient in order to populate or reconstitute a blood cell type that is deficient in the patient. The disclosure thus provides methods of treating a variety of hematopoietic conditions, such as sickle cell anemia, thalassemia,


30

35

40

Fanconi anemia, Wiskott-Aldrich syndrome, adenosine deaminase deficiency-severe combined immunodeficiency, metachromatic leukodystrophy, Diamond-Blackfan anemia and Schwachman-Diamond syndrome, human immunodeficiency virus infection, and acquired immune deficiency syndrome, among others.

- 5 In a first aspect, the disclosure features a compound represented by formula (I)



- wherein L is a linker selected from the group consisting of $-NR_{7a}(CR_{8a}R_{8b})_n-$, $-O(CR_{8a}R_{8b})_n-$, $-C(O)(CR_{8a}R_{8b})_n-$, $-C(S)(CR_{8a}R_{8b})_n-$, $-S(O)_{0-2}(CR_{8a}R_{8b})_n-$, $-(CR_{8a}R_{8b})_n-$, $-NR_{7a}C(O)(CR_{8a}R_{8b})_n-$, $-NR_{7a}C(S)(CR_{8a}R_{8b})_n-$, $-OC(O)(CR_{8a}R_{8b})_n-$, $-OC(S)(CR_{8a}R_{8b})_n-$, $-C(O)NR_{7a}(CR_{8a}R_{8b})_n-$, $-C(S)NR_{7a}(CR_{8a}R_{8b})_n-$, $-C(O)O(CR_{8a}R_{8b})_n-$, $-C(S)O(CR_{8a}R_{8b})_n-$, $-S(O)_2NR_{7a}(CR_{8a}R_{8b})_n-$, $-NR_{7a}S(O)_2(CR_{8a}R_{8b})_n-$, $-NR_{7a}C(O)NR_{7b}(CR_{8a}R_{8b})_n-$, $-NR_{7a}(CR_{8a}R_{8b})_nNR_{7a}-$, $-NR_{7a}(CR_{8a}R_{8b})_nO-$, $-NR_{7a}(CR_{8a}R_{8b})_nS-$, $-O(CR_{8a}R_{8b})_nNR_{7a}-$, $-O(CR_{8a}R_{8b})_nO-$, $-O(CR_{8a}R_{8b})_nS-$, $-S(CR_{8a}R_{8b})_nNR_{7a}-$, $-S(CR_{8a}R_{8b})_nO-$, $-S(CR_{8a}R_{8b})_nS-$, and $-NR_{7a}C(O)O(CR_{8a}R_{8b})_n-$, wherein R_{7a} , R_{7b} , R_{8a} , and R_{8b} are each independently selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl, and each n is independently an integer from 2 to 6;

- R_1 is selected from the group consisting of $-S(O)_2NR_{9a}R_{9b}$, $-NR_{9a}C(O)R_{9b}$, $-NR_{9a}C(S)R_{9b}$, $-NR_{9a}C(O)NR_{9b}R_{9c}$, $-C(O)R_{9a}$, $-C(S)R_{9a}$, $-S(O)_{0-2}R_{9a}$, $-C(O)OR_{9a}$, $-C(S)OR_{9a}$, $-C(O)NR_{9a}R_{9b}$, $-C(S)NR_{9a}R_{9b}$, $-NR_{9a}S(O)_2R_{9b}$, $-NR_{9a}C(O)OR_{9b}$, $-OC(O)CR_{9a}R_{9b}R_{9c}$, $-OC(S)CR_{9a}R_{9b}R_{9c}$, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein R_{9a} , R_{9b} , and R_{9c} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

- R_2 is selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl;

- R_3 is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

R_4 is selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl;

- R_5 is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R_6 is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;


or a salt thereof.

- As used herein to describe linkers (represented by "L" in formulas (I), (II), and the like), the notation "- (Linker) -" (wherein "linker" is represented using chemical symbols such as $NR_{7a}(CR_{8a}R_{8b})_n$, $O(CR_{8a}R_{8b})_n$, $C(O)(CR_{8a}R_{8b})_n$, $C(S)(CR_{8a}R_{8b})_n$, $S(O)_{0-2}(CR_{8a}R_{8b})_n$, $(CR_{8a}R_{8b})_n$, $NR_{7a}C(O)(CR_{8a}R_{8b})_n$,


- $\text{NR}_{7a}\text{C}(\text{S})(\text{CR}_{8a}\text{R}_{8b})_n$, $\text{OC}(\text{O})(\text{CR}_{8a}\text{R}_{8b})_n$, $\text{OC}(\text{S})(\text{CR}_{8a}\text{R}_{8b})_n$, $\text{C}(\text{O})\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n$, $\text{C}(\text{S})\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n$,
 $\text{C}(\text{O})\text{O}(\text{CR}_{8a}\text{R}_{8b})_n$, $\text{C}(\text{S})\text{O}(\text{CR}_{8a}\text{R}_{8b})_n$, $\text{S}(\text{O})_2\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n$, $\text{NR}_{7a}\text{S}(\text{O})_2(\text{CR}_{8a}\text{R}_{8b})_n$, and
 $\text{NR}_{7a}\text{C}(\text{O})\text{NR}_{7b}(\text{CR}_{8a}\text{R}_{8b})_n$ designates that the left hyphen represents a covalent bond to the indicated
 position on the imidazopyridine or imidazopyrazine ring system, while the right hyphen represents a
 5 covalent bond to R_1 .
- In some embodiments, R_1 is selected from the group consisting of $-\text{S}(\text{O})_2\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{R}_{9b}$,
 $-\text{NR}_{9a}\text{C}(\text{S})\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{NR}_{9b}\text{R}_{9c}$, $-\text{C}(\text{O})\text{R}_{9a}$, $-\text{C}(\text{S})\text{R}_{9a}$, $-\text{S}(\text{O})_{0-2}\text{R}_{9a}$, $-\text{C}(\text{O})\text{OR}_{9a}$, $-\text{C}(\text{S})\text{OR}_{9a}$, $-\text{C}(\text{O})\text{NR}_{9a}\text{R}_{9b}$,
 $-\text{C}(\text{S})\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{S}(\text{O})_2\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{OR}_{9b}$, $-\text{OC}(\text{O})\text{CR}_{9a}\text{R}_{9b}\text{R}_{9c}$, $-\text{OC}(\text{S})\text{CR}_{9a}\text{R}_{9b}\text{R}_{9c}$, phenyl, 1H-
 pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-
 10 oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, and 1H-indazolyl, wherein the
 phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-
 oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, or 1H-indazolyl is optionally
 substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of
 cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy,
 15 amino, $-\text{O}(\text{CH}_2)_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{S}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{OS}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, and $-\text{NR}_{10a}\text{S}(\text{O})_2\text{R}_{10b}$, wherein R_{10a} and
 R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl,
 optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally
 substituted cycloalkyl, and optionally substituted heterocycloalkyl.
- In some embodiments, R_1 is selected from the group consisting of $-\text{S}(\text{O})_2\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{R}_{9b}$,
 20 $-\text{NR}_{9a}\text{C}(\text{S})\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{NR}_{9b}\text{R}_{9c}$, $-\text{C}(\text{O})\text{R}_{9a}$, $-\text{C}(\text{S})\text{R}_{9a}$, $-\text{S}(\text{O})_{0-2}\text{R}_{9a}$, $-\text{C}(\text{O})\text{OR}_{9a}$, $-\text{C}(\text{S})\text{OR}_{9a}$, $-\text{C}(\text{O})\text{NR}_{9a}\text{R}_{9b}$,
 $-\text{C}(\text{S})\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{S}(\text{O})_2\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{OR}_{9b}$, $-\text{OC}(\text{O})\text{CR}_{9a}\text{R}_{9b}\text{R}_{9c}$, and $-\text{OC}(\text{S})\text{CR}_{9a}\text{R}_{9b}\text{R}_{9c}$.
- In some embodiments, R_1 is selected from the group consisting of phenyl, 1H-pyrrolopyridinyl,
 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl,
 2-oxo-2,3-dihydro-1H-benzoimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-
 25 pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-
 oxo-2,3-dihydro-1H-benzoimidazolyl, or 1H-indazolyl is optionally substituted, for example, with from 1 to
 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy,
 halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{O}(\text{CH}_2)_2\text{NR}_{10a}\text{R}_{10b}$, -
 $\text{S}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{OS}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, and $-\text{NR}_{10a}\text{S}(\text{O})_2\text{R}_{10b}$.
- In some embodiments, R_1 is selected from the group consisting of phenyl, 1H-indol-2-yl, 1H-indol-
 30 3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-
 oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl,
 wherein the phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-
 1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, or 2-oxo-
 35 2,3-dihydro-1H-benzo[d]imidazol-5-yl is optionally substituted, for example, with from 1 to 3 substituents
 independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-
 substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{O}(\text{CH}_2)_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{S}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, -
 $\text{OS}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, and $-\text{NR}_{10a}\text{S}(\text{O})_2\text{R}_{10b}$.
- In some embodiments, R_1 is selected from the group consisting of phenyl, phenol-4-yl, 1H-indol-
 40 2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-

triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl.

In some embodiments, R_1 is selected from the group consisting of:



In some embodiments, R_1 is selected from the group consisting of:



In some embodiments, R_1 is selected from the group consisting of phenol-4-yl and 1H-indol-3-yl.

In some embodiments, L is selected from the group consisting of $-NR_{7a}(CR_{8a}R_{8b})_{n-}$ and

10 $O(CR_{8a}R_{8b})_{n-}$.

In some embodiments, L is selected from the group consisting of $-NH(CH_2)_2-$ and $-O(CH_2)_2-$.

In some embodiments, R_2 is hydrogen.

In some embodiments, R_3 is selected from the group consisting of optionally substituted aryl and optionally substituted heteroaryl.


15 In some embodiments, R_3 is selected from the group consisting of phenyl, thiophenyl, furanyl, 1H-benzoimidazolyl, quinolinyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl, wherein the phenyl, thiophenyl, furanyl, 1H-benzoimidazolyl, quinolinyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted, for example, with from 1
20 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, $-C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$, and wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl.

25 In some embodiments, R_3 is selected from the group consisting of thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, imidazo[1,2-a]pyridin-3-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl and thiazol-5-yl, wherein the thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl, or
30 thiazol-5-yl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄

alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$.

In some embodiments, R_3 is selected from the group consisting of thiophen-3-yl, benzo[b]thiophen-3-yl, pyridin-3-yl, pyrimidin-5-yl, 1H-imidazol-1-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, and imidazo[1,2-a]pyridin-3-yl, wherein the thiophen-3-yl, benzo[b]thiophen-3-yl, pyridin-3-yl, pyrimidin-5-yl, 1H-imidazol-1-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, or imidazo[1,2-a]pyridin-3-yl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$.


In some embodiments, R_3 is selected from the group consisting of optionally substituted:



In some embodiments, R_3 is pyridin-3-yl, wherein the pyridin-3-yl is optionally substituted at C5, for example, with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, $-C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$.

In some embodiments, the pyridin-3-yl is substituted at C5 with a substituent selected from the group consisting of ethoxycarbonyl, methoxy, cyano, methyl, methylsulfonyl, fluoro, chloro, trifluoromethyl, ethynyl, and cyclopropyl.

In some embodiments, R_3 is selected from the group consisting of:



In some embodiments, R_3 is imidazo[1,2-a]pyridin-3-yl, wherein the imidazo[1,2-a]pyridin-3-yl is optionally substituted, for example, with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, $-C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$.

In some embodiments, R_3 is benzo[b]thiophen-3-yl, wherein the benzo[b]thiophen-3-yl is optionally substituted, for example, with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, $-C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$.

In some embodiments, R_3 is 1H-imidazo[4,5-b]pyridin-1-yl, wherein the 1H-imidazo[4,5-b]pyridin-1-yl is optionally substituted, for example, with a substituent selected from the group consisting of C1-4

alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, $C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$.

In some embodiments, R_3 is isoquinolin-4-yl, wherein the isoquinolin-4-yl is optionally substituted, for example, with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, $C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$.

In some embodiments, R_4 is hydrogen.

In some embodiments, R_5 is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4 alkyl.

In some embodiments, R_5 is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-hydroxypropan-2-yl, and nonan-2-yl.


In some embodiments, R_5 is (S)-1-hydroxypropan-2-yl.

In some embodiments, R_5 is (R)-1-hydroxypropan-2-yl

In some embodiments, R_5 is (S)-sec-butyl.


In some embodiments, R_5 is (R)-sec-butyl.

In some embodiments, R_5 is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-C(O)R_{12a}$, $-S(O)_{0-2}R_{12a}$, $-C(O)OR_{12a}$, and $-C(O)NR_{12a}R_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl.

In some embodiments, R_5 is selected from the group consisting of:



In some embodiments, R₅ is (ii).

In some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl.

In some embodiments, R₅ is (S)-4-methoxybutan-2-yl.

In some embodiments, R₅ is (R)-4-methoxybutan-2-yl.

In some embodiments, R₅ is (S)-5-methoxypentan-2-yl.


In some embodiments, R₅ is (R)-5-methoxypentan-2-yl.

In some embodiments, R₅ is (S)-4-ethoxybutan-2-yl.

In some embodiments, R₅ is (R)-4-ethoxybutan-2-yl.

In some embodiments, R₆ is hydrogen.

In some embodiments, the disclosure features a compound represented by formula (I-a)



wherein L is a linker selected from the group consisting of -NR_{7a}(CR_{8a}R_{8b})_n-, -O(CR_{8a}R_{8b})_n-, -C(O)(CR_{8a}R_{8b})_n-, -C(S)(CR_{8a}R_{8b})_n-, -S(O)₀₋₂(CR_{8a}R_{8b})_n-, -(CR_{8a}R_{8b})_n-, -NR_{7a}C(O)(CR_{8a}R_{8b})_n-, -NR_{7a}C(S)(CR_{8a}R_{8b})_n-, -OC(O)(CR_{8a}R_{8b})_n-, -OC(S)(CR_{8a}R_{8b})_n-, -C(O)NR_{7a}(CR_{8a}R_{8b})_n-, -C(S)NR_{7a}(CR_{8a}R_{8b})_n-, -C(O)O(CR_{8a}R_{8b})_n-, -C(S)O(CR_{8a}R_{8b})_n-, -S(O)₂NR_{7a}(CR_{8a}R_{8b})_n-, -NR_{7a}S(O)₂(CR_{8a}R_{8b})_n-, -NR_{7a}C(O)NR_{7b}(CR_{8a}R_{8b})_n-, and -NR_{7a}C(O)O(CR_{8a}R_{8b})_n-, wherein R_{7a}, R_{7b}, R_{8a}, and R_{8b} are each independently selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl, and each n is independently an integer from 2 to 6;

R₁ is selected from the group consisting of -S(O)₂NR_{9a}R_{9b}-, -NR_{9a}C(O)R_{9b}-, -NR_{9a}C(S)R_{9b}-, -NR_{9a}C(O)NR_{9b}R_{9c}-, -C(O)R_{9a}-, -C(S)R_{9a}-, -S(O)₀₋₂R_{9a}-, -C(O)OR_{9a}-, -C(S)OR_{9a}-, -C(O)NR_{9a}R_{9b}-, -C(S)NR_{9a}R_{9b}-, -NR_{9a}S(O)₂R_{9b}-, -NR_{9a}C(O)OR_{9b}-, -OC(O)CR_{9a}R_{9b}R_{9c}-, -OC(S)CR_{9a}R_{9b}R_{9c}-, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein R_{9a}, R_{9b}, and R_{9c} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl (for example, R₁ may be selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy,

amino, $-\text{O}(\text{CH}_2)_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{S}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{OS}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, and $-\text{NR}_{10a}\text{S}(\text{O})_2\text{R}_{10b}$, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl);

5 Ar is selected from the group consisting of optionally substituted monocyclic aryl and heteroaryl, such as optionally substituted thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl;


R_5 is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R_6 is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

15 In some embodiments, Ar is pyridin-3-yl, wherein the pyridin-3-yl is optionally substituted at C5, for example, with a substituent selected from the group consisting of ethoxycarbonyl, methoxy, cyano, methyl, methylsulfonyl, fluoro, chloro, trifluoromethyl, ethynyl, and cyclopropyl.

In some embodiments, the disclosure features a compound represented by formula (I-b)



20 wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{O}(\text{CH}_2)_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{S}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{OS}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, and $-\text{NR}_{10a}\text{S}(\text{O})_2\text{R}_{10b}$, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

30 Ar is selected from the group consisting of optionally substituted monocyclic aryl and heteroaryl, such as optionally substituted thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl;


R_5 is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

- 5 R_6 is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;
or a salt thereof.

- In some embodiments, A is selected from the group consisting of phenyl, phenol-4-yl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl.

In some embodiments, A is selected from the group consisting of phenol-4-yl and 1H-indol-3-yl.

In some embodiments, the disclosure features a compound represented by formula (I-c)



- 15 wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;


- 25 B is an optionally substituted ring system selected from the group consisting of thiophenyl, furanyl, 1H-benzoimidazolyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl, wherein the thiophenyl, furanyl, 1H-benzoimidazolyl, isoquinolinyl, 1H-imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;

R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

- 5 R₆ is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;
or a salt thereof.

- In some embodiments, B is pyridin-3-yl, wherein the pyridin-3-yl is optionally substituted at C5, for example, with a substituent selected from the group consisting of ethoxycarbonyl, methoxy, cyano,
10 methyl, methylsulfonyl, fluoro, chloro, trifluoromethyl, ethynyl, and cyclopropyl.

In some embodiments, the disclosure features a compound represented by formula (I-d)




- wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -
15 O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

- B is an optionally substituted ring system selected from the group consisting of thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl, wherein the thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, 1H-imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)_{0.2}R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each
25 independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and
30

R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

5 In some embodiments, the disclosure features a compound represented by formula (I-e)




wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl, wherein the phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, or 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

B is an optionally substituted ring system selected from the group consisting of thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, imidazo[1,2-a]pyridin-3-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl and thiazol-5-yl, wherein the thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl, or thiazol-5-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and


30 R₅ is selected from the group consisting of C₁₋₁₀ alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C₁₋₁₀ alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-

pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;


In some embodiments, R₅ is selected from the group consisting of:



15 in some embodiments, R₅ is (ii);

in some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl; or a salt thereof.

In some embodiments, the disclosure features a compound represented by formula (I-f)




25 wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

q is an integer from 0 to 4;


each Z is independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

- 5 R₅ is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-hydroxypropan-2-yl, and nonan-2-yl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



- 10 wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;

- 15 In some embodiments, R₅ is selected from the group consisting of:




in some embodiments, R₅ is (ii);

- 20 in some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;

or a salt thereof.

In some embodiments, each Z is independently a substituent selected from the group consisting of ethoxycarbonyl, methoxy, cyano, methyl, methylsulfonyl, fluoro, chloro, trifluoromethyl, ethynyl, and cyclopropyl.


- 30 In some embodiments, the disclosure features a compound represented by formula (I-g)



wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;


5 Z is a substituent selected from the group consisting of C₁₋₄ alkyl, halo, halo-substituted-C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

R₅ is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-hydroxypropan-2-yl, and nonan-2-yl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;

In some embodiments, R₅ is selected from the group consisting of:




in some embodiments, R_5 is (ii);

in some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-

methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;

or a salt thereof.

In some embodiments, the disclosure features a compound represented by formula (I-h)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

q is an integer from 0 to 4;

r is 0 or 1;


W and V are each independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;

In some embodiments, R₅ is selected from the group consisting of:




in some embodiments, R₅ is (ii);

- 5 in some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;
- 10 ethoxyhexan-2-yl;

or a salt thereof.

In some embodiments, the disclosure features a compound represented by formula (I-i)




- wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;
- 15

q is an integer from 0 to 4;

r is 0 or 1;


- W and V are each independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and
- 20

- R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)
- 25
- 30



- wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;

In some embodiments, R₅ is selected from the group consisting of:




in some embodiments, R₅ is (ii);

- in some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;

or a salt thereof.

In some embodiments, the disclosure features a compound represented by formula (I-j)



- wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;


q is an integer from 0 to 4;

r is 0 or 1;

- W and V are each independently a substituent selected from the group consisting of C₁₋₄ alkyl, halo, halo-substituted-C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, cyano, amino,


$C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C_{1-4} alkyl; and

- R_5 is selected from the group consisting of C_{1-10} alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C_{1-10} alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C_{1-4} alkyl, and halo-substituted- C_{1-4} alkyl, or R_5 is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



- wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C_{1-4} alkyl, C_{1-4} alkenyl, C_{1-4} alkynyl, C_{2-6} cycloalkyl, C_{1-4} alkoxy, halo, halo-substituted- C_{1-4} alkyl, halo-substituted- C_{1-4} alkoxy, amino, $-C(O)R_{12a}$, $-S(O)_{0-2}R_{12a}$, $-C(O)OR_{12a}$, and $-C(O)NR_{12a}R_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C_{1-4} alkyl;

In some embodiments, R_5 is selected from the group consisting of:




in some embodiments, R_5 is (ii);

- in some embodiments, R_5 is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;

or a salt thereof.

- In some embodiments, the disclosure features a compound represented by formula (I-k)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

q is an integer from 0 to 4;

5 r is 0 or 1;


W and V are each independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and


10 R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



20 wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;

25 In some embodiments, R₅ is selected from the group consisting of:






in some embodiments, R_5 is (ii);

- in some embodiments, R_5 is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;


or a salt thereof.

- 10 In some embodiments, the compound is compound (1)




or a salt thereof.

In some embodiments, the compound is compound (2)




- 15 or a salt thereof.

In some embodiments, the compound is compound (3)




or a salt thereof.

In some embodiments, the compound is compound (4)



or a salt thereof.


In some embodiments, the compound is compound (5)



5


or a salt thereof.

In some embodiments, the compound is compound (6)



or a salt thereof.


In some embodiments, the compound is compound (7)



10


or a salt thereof.

In some embodiments, the compound is compound (8)



or a salt thereof.


In some embodiments, the compound is compound (9)



5


or a salt thereof.

In some embodiments, the compound is compound (10)



or a salt thereof.


In some embodiments, the compound is compound (11)



10


or a salt thereof.

In some embodiments, the compound is compound (23)



or a salt thereof.


In some embodiments, the compound is compound (25)



5


or a salt thereof.

In some embodiments, the compound is compound (26)



or a salt thereof.

In another aspect, the disclosure features a compound represented by formula (II)



10

wherein L is a linker selected from the group consisting of -NR_{7a}(CR_{8a}R_{8b})_n-, -O(CR_{8a}R_{8b})_n-, -C(O)(CR_{8a}R_{8b})_n-, -C(S)(CR_{8a}R_{8b})_n-, -S(O)₀₋₂(CR_{8a}R_{8b})_n-, -(CR_{8a}R_{8b})_n-, -NR_{7a}C(O)(CR_{8a}R_{8b})_n-, -NR_{7a}C(S)(CR_{8a}R_{8b})_n-, -OC(O)(CR_{8a}R_{8b})_n-, -OC(S)(CR_{8a}R_{8b})_n-, -C(O)NR_{7a}(CR_{8a}R_{8b})_n-, -C(S)NR_{7a}(CR_{8a}R_{8b})_n-, -C(O)O(CR_{8a}R_{8b})_n-, -C(S)O(CR_{8a}R_{8b})_n-, -S(O)₂NR_{7a}(CR_{8a}R_{8b})_n-, -NR_{7a}S(O)₂(CR_{8a}R_{8b})_n-, -NR_{7a}C(O)NR_{7b}(CR_{8a}R_{8b})_n-, -NR_{7a}(CR_{8a}R_{8b})_nNR_{7a}-, -NR_{7a}(CR_{8a}R_{8b})_nO-, -NR_{7a}(CR_{8a}R_{8b})_nS-, -O(CR_{8a}R_{8b})_nNR_{7a}-, -O(CR_{8a}R_{8b})_nO-, -O(CR_{8a}R_{8b})_nS-, -S(CR_{8a}R_{8b})_nNR_{7a}-, -

15

$S(CR_{8a}R_{8b})_nO-$, $-S(CR_{8a}R_{8b})_nS-$, and $-NR_{7a}C(O)O(CR_{8a}R_{8b})_n-$, wherein R_{7a} , R_{7b} , R_{8a} , and R_{8b} are each independently selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl, and each n is independently an integer from 2 to 6;

R_1 is selected from the group consisting of $-S(O)_2NR_{9a}R_{9b}$, $-NR_{9a}C(O)R_{9b}$, $-NR_{9a}C(S)R_{9b}$, $-NR_{9a}C(O)NR_{9b}R_{9c}$, $-C(O)R_{9a}$, $-C(S)R_{9a}$, $-S(O)_{0-2}R_{9a}$, $-C(O)OR_{9a}$, $-C(S)OR_{9a}$, $-C(O)NR_{9a}R_{9b}$, $-C(S)NR_{9a}R_{9b}$, $-NR_{9a}S(O)_2R_{9b}$, $-NR_{9a}C(O)OR_{9b}$, $-OC(O)CR_{9a}R_{9b}R_{9c}$, $-OC(S)CR_{9a}R_{9b}R_{9c}$, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein R_{9a} , R_{9b} , and R_{9c} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

R_3 is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

R_4 is selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl;

R_5 is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R_6 is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; or a salt thereof.

In some embodiments, R_1 is selected from the group consisting of $-S(O)_2NR_{9a}R_{9b}$, $-NR_{9a}C(O)R_{9b}$, $-NR_{9a}C(S)R_{9b}$, $-NR_{9a}C(O)NR_{9b}R_{9c}$, $-C(O)R_{9a}$, $-C(S)R_{9a}$, $-S(O)_{0-2}R_{9a}$, $-C(O)OR_{9a}$, $-C(S)OR_{9a}$, $-C(O)NR_{9a}R_{9b}$, $-C(S)NR_{9a}R_{9b}$, $-NR_{9a}S(O)_2R_{9b}$, $-NR_{9a}C(O)OR_{9b}$, $-OC(O)CR_{9a}R_{9b}R_{9c}$, $-OC(S)CR_{9a}R_{9b}R_{9c}$, phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-O(CH_2)_2NR_{10a}R_{10b}$, $-S(O)_2NR_{10a}R_{10b}$, $-OS(O)_2NR_{10a}R_{10b}$, and $-NR_{10a}S(O)_2R_{10b}$; wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl.

In some embodiments, R_1 is selected from the group consisting of $-S(O)_2NR_{9a}R_{9b}$, $-NR_{9a}C(O)R_{9b}$, $-NR_{9a}C(S)R_{9b}$, $-NR_{9a}C(O)NR_{9b}R_{9c}$, $-C(O)R_{9a}$, $-C(S)R_{9a}$, $-S(O)_{0-2}R_{9a}$, $-C(O)OR_{9a}$, $-C(S)OR_{9a}$, $-C(O)NR_{9a}R_{9b}$, $-C(S)NR_{9a}R_{9b}$, $-NR_{9a}S(O)_2R_{9b}$, $-NR_{9a}C(O)OR_{9b}$, $-OC(O)CR_{9a}R_{9b}R_{9c}$, and $-OC(S)CR_{9a}R_{9b}R_{9c}$.


In some embodiments, R_1 is selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-

oxo-2,3-dihydro-1H-benzoimidazolyl, or 1H-indazolyl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-O(CH_2)_2NR_{10a}R_{10b}$, $-S(O)_2NR_{10a}R_{10b}$, $-OS(O)_2NR_{10a}R_{10b}$, and $-NR_{10a}S(O)_2R_{10b}$.


- 5 In some embodiments, R_1 is selected from the group consisting of phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl, wherein the phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, or 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-O(CH_2)_2NR_{10a}R_{10b}$, $-S(O)_2NR_{10a}R_{10b}$, $-OS(O)_2NR_{10a}R_{10b}$, and $-NR_{10a}S(O)_2R_{10b}$.

- 15 In some embodiments, R_1 is selected from the group consisting of phenyl, phenol-4-yl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl.

In some embodiments, R_1 is selected from the group consisting of:



In some embodiments, R_1 is selected from the group consisting of:



In some embodiments, R_1 is selected from the group consisting of phenol-4-yl and 1H-indol-3-yl.

In some embodiments, L is selected from the group consisting of $-NR_{7a}(CR_{8a}R_{8b})_{n-}$ and

- 25 $O(CR_{8a}R_{8b})_{n-}$.

In some embodiments, L is selected from the group consisting of $-NH(CH_2)_2-$ and $-O(CH_2)_2-$.

In some embodiments, R_3 is selected from the group consisting of optionally substituted aryl and optionally substituted heteroaryl.


- 30 In some embodiments, R_3 is selected from the group consisting of phenyl, thiophenyl, furanyl, 1H-benzoimidazolyl, quinoliny, isoquinoliny, imidazopyridiny, benzothiophenyl, pyrimidiny, pyridiny, 1H-imidazolyl, pyraziny, pyridaziny, 1H-pyrrolyl, and thiazolyl, wherein the phenyl, thiophenyl, furanyl, 1H-benzoimidazolyl, quinoliny, isoquinoliny, imidazopyridiny, benzothiophenyl, pyrimidiny, pyridiny, 1H-

imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, and wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl.

In some embodiments, R₃ is selected from the group consisting of thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, imidazo[1,2-a]pyridin-3-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl and thiazol-5-yl, wherein the thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl, or thiazol-5-yl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

In some embodiments, R₃ is selected from the group consisting of thiophen-3-yl, benzo[b]thiophen-3-yl, pyridin-3-yl, pyrimidin-5-yl, 1H-imidazol-1-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, and imidazo[1,2-a]pyridin-3-yl, wherein the thiophen-3-yl, benzo[b]thiophen-3-yl, pyridin-3-yl, pyrimidin-5-yl, 1H-imidazol-1-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, or imidazo[1,2-a]pyridin-3-yl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.


In some embodiments, R₃ is selected from the group consisting of optionally substituted:



In some embodiments, R₃ is pyridin-3-yl, wherein the pyridin-3-yl is optionally substituted at C5, for example, with a substituent selected from the group consisting of C₁₋₄ alkyl, halo, halo-substituted-C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

30 In some embodiments, the pyridin-3-yl is substituted at C5 with a substituent selected from the group consisting of ethoxycarbonyl, methoxy, cyano, methyl, methylsulfonyl, fluoro, chloro, trifluoromethyl, ethynyl, and cyclopropyl.

In some embodiments, R₃ is selected from the group consisting of:



In some embodiments, R₃ is imidazo[1,2-a]pyridin-3-yl, wherein the imidazo[1,2-a]pyridin-3-yl is optionally substituted, for example, with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

In some embodiments, R₃ is benzo[b]thiophen-3-yl, wherein the benzo[b]thiophen-3-yl is optionally substituted, for example, with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

In some embodiments, R₃ is 1H-imidazo[4,5-b]pyridin-1-yl, wherein the 1H-imidazo[4,5-b]pyridin-1-yl is optionally substituted, for example, with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

In some embodiments, R₃ is isoquinolin-4-yl, wherein the isoquinolin-4-yl is optionally substituted, for example, with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

In some embodiments, R₄ is hydrogen.

In some embodiments, R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl.

In some embodiments, R₅ is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-hydroxypropan-2-yl, and nonan-2-yl.


In some embodiments, R₅ is (S)-1-hydroxypropan-2-yl.

In some embodiments, R₅ is (R)-1-hydroxypropan-2-yl.

In some embodiments, R₅ is (S)-sec-butyl.


In some embodiments, R₅ is (R)-sec-butyl.

In some embodiments, R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl.

In some embodiments, R₅ is selected from the group consisting of:



In some embodiments, R_5 is (ii).

In some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl.

In some embodiments, R₅ is (S)-4-methoxybutan-2-yl.

In some embodiments, R₅ is (R)-4-methoxybutan-2-yl.

20 In some embodiments, R₅ is (S)-5-methoxypentan-2-yl.


In some embodiments, R₅ is (R)-5-methoxypentan-2-yl.

In some embodiments, R₅ is (S)-4-ethoxybutan-2-yl.

In some embodiments, R₅ is (R)-4-ethoxybutan-2-yl.

In some embodiments, R₆ is hydrogen.

25 In some embodiments, the disclosure features a compound represented by formula (II-a)



wherein L is a linker selected from the group consisting of -NR_{7a}(CR_{8a}R_{8b})_n-, -O(CR_{8a}R_{8b})_n-, -C(O)(CR_{8a}R_{8b})_n-, -C(S)(CR_{8a}R_{8b})_n-, -S(O)₀₋₂(CR_{8a}R_{8b})_n-, -(CR_{8a}R_{8b})_n-, -NR_{7a}C(O)(CR_{8a}R_{8b})_n-, -NR_{7a}C(S)(CR_{8a}R_{8b})_n-, -OC(O)(CR_{8a}R_{8b})_n-, -OC(S)(CR_{8a}R_{8b})_n-, -C(O)NR_{7a}(CR_{8a}R_{8b})_n-, -C(S)NR_{7a}(CR_{8a}R_{8b})_n-, -C(O)O(CR_{8a}R_{8b})_n-, -C(S)O(CR_{8a}R_{8b})_n-, -S(O)₂NR_{7a}(CR_{8a}R_{8b})_n-, -

$\text{NR}_{7a}\text{S}(\text{O})_2(\text{CR}_{8a}\text{R}_{8b})_{n-}$, $-\text{NR}_{7a}\text{C}(\text{O})\text{NR}_{7b}(\text{CR}_{8a}\text{R}_{8b})_{n-}$, and $-\text{NR}_{7a}\text{C}(\text{O})\text{O}(\text{CR}_{8a}\text{R}_{8b})_{n-}$, wherein R_{7a} , R_{7b} , R_{8a} , and R_{8b} are each independently selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl, and each n is independently an integer from 2 to 6;

- R_1 is selected from the group consisting of $-\text{S}(\text{O})_2\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{S})\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{NR}_{9b}\text{R}_{9c}$, $-\text{C}(\text{O})\text{R}_{9a}$, $-\text{C}(\text{S})\text{R}_{9a}$, $-\text{S}(\text{O})_{0-2}\text{R}_{9a}$, $-\text{C}(\text{O})\text{OR}_{9a}$, $-\text{C}(\text{S})\text{OR}_{9a}$, $-\text{C}(\text{O})\text{NR}_{9a}\text{R}_{9b}$, $-\text{C}(\text{S})\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{S}(\text{O})_2\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{OR}_{9b}$, $-\text{OC}(\text{O})\text{CR}_{9a}\text{R}_{9b}\text{R}_{9c}$, $-\text{OC}(\text{S})\text{CR}_{9a}\text{R}_{9b}\text{R}_{9c}$, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein R_{9a} , R_{9b} , and R_{9c} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl (for example, R_1 may be selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{O}(\text{CH}_2)_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{S}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{OS}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, and $-\text{NR}_{10a}\text{S}(\text{O})_2\text{R}_{10b}$, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl);

Ar is selected from the group consisting of optionally substituted monocyclic aryl and heteroaryl, such as optionally substituted thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl;


- R_5 is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

- R_6 is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

In some embodiments, Ar is pyridin-3-yl, wherein the pyridin-3-yl is optionally substituted at C5, for example, with a substituent selected from the group consisting of ethoxycarbonyl, methoxy, cyano, methyl, methylsulfonyl, fluoro, chloro, trifluoromethyl, ethynyl, and cyclopropyl.

- In some embodiments, the disclosure features a compound represented by formula (II-b)



wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

Ar is selected from the group consisting of optionally substituted monocyclic aryl and heteroaryl, such as optionally substituted thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl;

R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and


R₆ is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

In some embodiments, A is selected from the group consisting of phenyl, phenol-4-yl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl.

In some embodiments, A is selected from the group consisting of phenol-4-yl and 1H-indol-3-yl.

In some embodiments, the disclosure features a compound represented by formula (II-c)



wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

B is an optionally substituted ring system selected from the group consisting of thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl, wherein the thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, 1H-imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;


R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R₆ is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

In some embodiments, B is pyridin-3-yl, wherein the pyridin-3-yl is optionally substituted at C5, for example, with a substituent selected from the group consisting of ethoxycarbonyl, methoxy, cyano, methyl, methylsulfonyl, fluoro, chloro, trifluoromethyl, ethynyl, and cyclopropyl.

In some embodiments, the disclosure features a compound represented by formula (II-d)




wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

B is an optionally substituted ring system selected from the group consisting of thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl, wherein the thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, 1H-imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.


In some embodiments, the disclosure features a compound represented by formula (II-e)



wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl, wherein the phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, or 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;


B is an optionally substituted ring system selected from the group consisting of thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, imidazo[1,2-a]pyridin-3-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl and thiazol-5-yl, wherein the thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl, or thiazol-5-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

R₅ is selected from the group consisting of C₁₋₁₀ alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C₁₋₁₀ alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C₁₋₄ alkyl, and halo-substituted-C₁₋₄alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;

In some embodiments, R₅ is selected from the group consisting of:




5 in some embodiments, R₅ is (ii);

in some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;

or a salt thereof.

In some embodiments, the disclosure features a compound represented by formula (II-f)




15 wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

q is an integer from 0 to 4;


each Z is independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a},
20 -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

R₅ is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-hydroxypropan-2-yl, and nonan-2-yl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and
25 (v)



- wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are
- 5 each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;


In some embodiments, R₅ is selected from the group consisting of:



- 10 in some embodiments, R₅ is (ii);
- in some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;
- 15 or a salt thereof.

- In some embodiments, each Z is independently a substituent selected from the group consisting of ethoxycarbonyl, methoxy, cyano, methyl, methylsulfonyl, fluoro, chloro, trifluoromethyl, ethynyl, and
- 20 cyclopropyl.


In some embodiments, the disclosure features a compound represented by formula (II-g)



- wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;
- 25 Z is a substituent selected from the group consisting of C₁₋₄ alkyl, halo, halo-substituted-C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and


- R₅ is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-
- 30

hydroxypropan-2-yl, and nonan-2-yl, or R_5 is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



- wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-C(O)R_{12a}$, $-S(O)_{0-2}R_{12a}$, $-C(O)OR_{12a}$, and $-C(O)NR_{12a}R_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl;

In some embodiments, R_5 is selected from the group consisting of:




in some embodiments, R_5 is (ii);

- in some embodiments, R_5 is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;

- or a salt thereof.

In some embodiments, the disclosure features a compound represented by formula (II-h)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

- q is an integer from 0 to 4;
 r is 0 or 1;


W and V are each independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

- 5 R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



- 15 wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;

- 20 In some embodiments, R₅ is selected from the group consisting of:




in some embodiments, R₅ is (ii);

- 25 in some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;

or a salt thereof.

In some embodiments, the disclosure features a compound represented by formula (II-i)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

q is an integer from 0 to 4;

5 r is 0 or 1;


W and V are each independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and


10 R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



20 wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;

25 In some embodiments, R₅ is selected from the group consisting of:






in some embodiments, R_5 is (ii);

in some embodiments, R_5 is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;

or a salt thereof.

In some embodiments, the disclosure features a compound represented by formula (II-j)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

q is an integer from 0 to 4;

r is 0 or 1;

W and V are each independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, $C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl; and


R_5 is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R_5 is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl,

C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;

In some embodiments, R₅ is selected from the group consisting of:




in some embodiments, R_5 is (ii);

in some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;

15 or a salt thereof.

In some embodiments, the disclosure features a compound represented by formula (II-k)



wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;


20 q is an integer from 0 to 4;

r is 0 or 1;

W and V are each independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-


pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C₁₋₄ alkyl, and halo-substituted-C₁₋₄alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;

10

In some embodiments, R₅ is selected from the group consisting of:




15 in some embodiments, R₅ is (ii);

in some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl;

20


or a salt thereof.

In some embodiments, the compound is compound (12)




25 or a salt thereof.

In some embodiments, the compound is compound (13)



or a salt thereof.


In some embodiments, the compound is compound (14)



5


or a salt thereof.

In some embodiments, the compound is compound (15)



or a salt thereof.


In some embodiments, the compound is compound (16)



10


or a salt thereof.

In some embodiments, the compound is compound (17)



or a salt thereof.


In some embodiments, the compound is compound (18)



5


or a salt thereof.

In some embodiments, the compound is compound (19)



or a salt thereof.


In some embodiments, the compound is compound (20)



10


or a salt thereof.

In some embodiments, the compound is compound (21)



or a salt thereof.


In some embodiments, the compound is compound (22)



5


or a salt thereof.

In some embodiments, the compound is compound (24)



or a salt thereof.


In some embodiments, the compound is compound (27)



10

or a salt thereof.

In some embodiments, the compound is compound (28)



or a salt thereof.

In another aspect, the disclosure features a method of producing an expanded population of hematopoietic stem cells ex vivo, the method including contacting a population of hematopoietic stem cells with the compound of any one of the above aspects or embodiments in an amount sufficient to produce an expanded population of hematopoietic stem cells.

In another aspect, the disclosure features a method of enriching a population of cells with hematopoietic stem cells ex vivo, the method including contacting a population of hematopoietic stem cells with the compound of any one of the above aspects or embodiments in an amount sufficient to produce a population of cells enriched with hematopoietic stem cells.

In another aspect, the disclosure features a method of maintaining the hematopoietic stem cell functional potential of a population of hematopoietic stem cells ex vivo for two or more days, the method including contacting a first population of hematopoietic stem cells with the compound of any one of the above aspects or embodiments, wherein the first population of hematopoietic stem cells exhibits a hematopoietic stem cell functional potential after two or more days that is greater than that of a control population of hematopoietic stem cells cultured under the same conditions and for the same time as the first population of hematopoietic stem cells but not contacted with the compound.

In some embodiments, the first population of hematopoietic stem cells exhibits a hematopoietic stem cell functional potential after three or more days (for example, three days, ten days, thirty days, sixty days, or more) of culture that is greater than that of the control population of hematopoietic stem cells.

In some embodiments, the hematopoietic stem cells are mammalian cells, such as human cells. In some embodiments, the human cells are CD34+ cells, such as CD34+ cells are CD34+, CD34+CD38-, CD34+CD38-CD90+, CD34+CD38-CD90+CD45RA-, CD34+CD38-CD90+CD45RA-CD49F+, or CD34+CD90+CD45RA- cells.

In some embodiments, the hematopoietic stem cells are CD34+ hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD90+ hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD45RA- hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD34+CD90+ hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD34+CD45RA- hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD90+CD45RA- hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD34+CD90+CD45RA- hematopoietic stem cells.

In some embodiments, the hematopoietic stem cells are obtained from human cord blood, mobilized human peripheral blood, or human bone marrow. The hematopoietic stem cells may, for example, be freshly isolated from the human or may have been previously cryopreserved.

In some embodiments, the hematopoietic stem cells or progeny thereof maintain hematopoietic stem cell functional potential after two or more days upon transplantation of the hematopoietic stem cells into a human subject.

5 In some embodiments, the hematopoietic stem cells or progeny thereof are capable of localizing to hematopoietic tissue and reestablishing hematopoiesis upon transplantation of the hematopoietic stem cells into a human subject.

In some embodiments, upon transplantation into a human subject, the hematopoietic stem cells give rise to a population of cells selected from the group consisting of megakaryocytes, thrombocytes, platelets, erythrocytes, mast cells, myoblasts, basophils, neutrophils, eosinophils, microglia, granulocytes, 10 monocytes, osteoclasts, antigen-presenting cells, macrophages, dendritic cells, natural killer cells, T-lymphocytes, and B-lymphocytes.

In another aspect, the disclosure features a method of treating a patient (e.g., a human patient) suffering from a stem cell disorder, the method including administering to the patient a population of hematopoietic stem cells, wherein the hematopoietic stem cells were produced by contacting the 15 hematopoietic stem cells or progenitors thereof with a compound of any of the above aspects or embodiments.

In another aspect, the disclosure features a method of preparing an expanded population of hematopoietic stem cells for transplantation into a patient (e.g., a human patient) suffering from a stem cell disorder, the method including contacting a first population of hematopoietic stem cells with a 20 compound of any of the above aspects or embodiments for a time sufficient to produce the expanded population of hematopoietic stem cells.

In another aspect, the disclosure features a method of treating a patient (e.g., a human patient) suffering from a stem cell disorder, the method including:

- 25 a. preparing an expanded population of hematopoietic stem cells by contacting a first population of hematopoietic stem cells with a compound of any of the above aspects or embodiments; and
- b. administering the expanded population of hematopoietic stem cells to the patient.

In yet another aspect, provided herein is a method of treating a stem cell disorder in a patient (e.g., a human patient) in need thereof, comprising administering an expanded population of 30 hematopoietic stem cells to the patient, wherein the expanded population of hematopoietic stem cells is prepared by contacting a first population of hematopoietic stem cells with a compound of any of the above aspects or embodiments for a time sufficient to produce the expanded population of hematopoietic stem cells.

In some embodiments, the stem cell disorder is a hemoglobinopathy.

35 In some embodiments, the stem cell disorder is selected from the group consisting of sickle cell anemia, thalassemia, Fanconi anemia, and Wiskott-Aldrich syndrome.

In some embodiments, the stem cell disorder is Fanconi anemia.

In some embodiments, the stem cell disorder is a myelodysplastic disorder.

In some embodiments, the stem cell disorder is an immunodeficiency disorder, such as a 40 congenital immunodeficiency or an acquired immunodeficiency. The acquired immunodeficiency may be, for example, human immunodeficiency virus (HIV) or acquired immune deficiency syndrome (AIDS).

In some embodiments, the stem cell disorder is a metabolic disorder, such as a glycogen storage disease, a mucopolysaccharidose, Gaucher's Disease, Hurlers Disease, a sphingolipidose, or metachromatic leukodystrophy.

In some embodiments, the stem cell disorder is cancer, such as a hematological cancer. The cancer may be, for example, leukemia, lymphoma, multiple myeloma, or neuroblastoma. In some
5 embodiments, the cancer is acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia, chronic lymphoid leukemia, multiple myeloma, diffuse large B-cell lymphoma, or non-Hodgkin's lymphoma.

In some embodiments, the stem cell disorder is a disorder selected from the group consisting of
10 adenosine deaminase deficiency and severe combined immunodeficiency, hyper immunoglobulin M syndrome, Chediak-Higashi disease, hereditary lymphohistiocytosis, osteopetrosis, osteogenesis imperfecta, storage diseases, thalassemia major, systemic sclerosis, systemic lupus erythematosus, multiple sclerosis, and juvenile rheumatoid arthritis.

In some embodiments, the stem cell disorder is an autoimmune disorder. For example, the stem
15 cell disorder may be multiple sclerosis, human systemic lupus, rheumatoid arthritis, inflammatory bowel disease, treating psoriasis, Type 1 diabetes mellitus, acute disseminated encephalomyelitis, Addison's disease, alopecia universalis, ankylosing spondylitis, antiphospholipid antibody syndrome, aplastic anemia, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune inner ear disease, autoimmune lymphoproliferative syndrome, autoimmune oophoritis, Balo disease, Behcet's disease,
20 bullous pemphigoid, cardiomyopathy, Chagas' disease, chronic fatigue immune dysfunction syndrome, chronic inflammatory demyelinating polyneuropathy, Crohn's disease, cicatricial pemphigoid, coeliac sprue-dermatitis herpetiformis, cold agglutinin disease, CREST syndrome, Degos disease, discoid lupus, dysautonomia, endometriosis, essential mixed cryoglobulinemia, fibromyalgia-fibromyositis, Goodpasture's syndrome, Grave's disease, Guillain-Barre syndrome, Hashimoto's thyroiditis,
25 Hidradenitis suppurativa, idiopathic and/or acute thrombocytopenic purpura, idiopathic pulmonary fibrosis, IgA neuropathy, interstitial cystitis, juvenile arthritis, Kawasaki's disease, lichen planus, Lyme disease, Meniere disease, mixed connective tissue disease, myasthenia gravis, neuromyotonia, opsoclonus myoclonus syndrome, optic neuritis, Ord's thyroiditis, pemphigus vulgaris, pernicious anemia, polychondritis, polymyositis and dermatomyositis, primary biliary cirrhosis, polyarteritis nodosa,
30 polyglandular syndromes, polymyalgia rheumatica, primary agammaglobulinemia, Raynaud phenomenon, Reiter's syndrome, rheumatic fever, sarcoidosis, scleroderma, Sjögren's syndrome, stiff person syndrome, Takayasu's arteritis, temporal arteritis, ulcerative colitis, uveitis, vasculitis, vitiligo, vulvodynia, or Wegener's granulomatosis.

In some embodiments, the stem cell disorder is a neurological disorder, such as Parkinson's
35 disease, Alzheimer's disease, multiple sclerosis, Amyotrophic lateral sclerosis, Huntington's disease, mild cognitive impairment, amyloidosis, AIDS-related dementia, encephalitis, stroke, head trauma, epilepsy, mood disorders, or dementia.

In another aspect, provided herein is a method of producing microglia in the central nervous system of a patient (e.g., a human patient) in need thereof, comprising administering an expanded
40 population of hematopoietic stem cells to the patient, wherein the expanded population of hematopoietic stem cells is prepared by contacting a first population of hematopoietic stem cells with a compound of any

of the above aspects or embodiments for a time sufficient to produce the expanded population of hematopoietic stem cells, and wherein administration of the expanded population of hematopoietic stem cells results in formation of microglia in the central nervous system of the patient.

In another aspect, the disclosure features a composition comprising a population of
5 hematopoietic stem cells, wherein the hematopoietic stem cells or progenitors thereof have been contacted with the compound of any one of the above aspects or embodiments, thereby expanding the hematopoietic stem cells or progenitors thereof.

In another aspect, the disclosure features a kit including the compound of any one of the above aspects or embodiments and a package insert, wherein the package insert instructs a user of the kit to
10 contact a population of hematopoietic stem cells with the compound for a time sufficient to produce an expanded population of hematopoietic stem cells.

In another aspect, the disclosure features a kit including the compound of any one of the above aspects or embodiments and a package insert, wherein the package insert instructs a user of the kit to contact a population of cells including hematopoietic stem cells with the compound for a time sufficient to
15 produce a population of cells enriched with hematopoietic stem cells.

In another aspect, the disclosure features a kit including the compound of any one of the above aspects or embodiments and a package insert, wherein the package insert instructs a user of the kit to contact a population of hematopoietic stem cells with the compound for a time sufficient to maintain the hematopoietic stem cell functional potential of the population of hematopoietic stem cells ex vivo for two
20 or more days.

In some embodiments, the kit further includes a population of cells including hematopoietic stem cells.

In another aspect, the disclosure features a compound of any of the above aspects or embodiments for use in producing an expanded population of hematopoietic stem cells ex vivo.

25 In another aspect, the disclosure features a compound of any of the above aspects or embodiments for use in enriching a population of cells with hematopoietic stem cells ex vivo.

In another aspect, the disclosure features a compound of any of the above aspects or embodiments for use in maintaining the hematopoietic stem cell functional potential of a population of hematopoietic stem cells ex vivo for two or more days.

30 In another aspect, the disclosure features use of a compound of any of the above aspects or embodiments in producing an expanded population of hematopoietic stem cells ex vivo.

In another aspect, the disclosure features use of a compound of any of the above aspects or embodiments in enriching a population of cells with hematopoietic stem cells ex vivo.

35 In another aspect, the disclosure features use of a compound of any of the above aspects or embodiments in maintaining the hematopoietic stem cell functional potential of a population of hematopoietic stem cells ex vivo for two or more days.

In another aspect, the disclosure features a composition for use in treating a human patient suffering from a disorder, said composition comprising a population of hematopoietic stem cells, wherein said hematopoietic stem cells were produced by contacting said hematopoietic stem cells or progenitors
40 thereof with a compound of any of the above aspects or embodiments.

In another aspect, the disclosure features a composition for use in treating a human patient suffering from a disorder, said composition comprising an expanded population of hematopoietic stem cells prepared by contacting a first population of hematopoietic stem cells with a compound of any of the above aspects or embodiments for a time sufficient to produce said expanded population of
 5 hematopoietic stem cells.

In another aspect, the disclosure features use of a compound of any of the above aspects or embodiments in preparing a medicament for treating a human patient suffering from a disorder.

In some embodiments, the disorder is selected from hemoglobinopathy, sickle cell anemia, thalassemia, Fanconi anemia, Wiskott-Aldrich syndrome, a myelodysplastic disorder, an
 10 immunodeficiency disorder, a metabolic disorder, cancer, adenosine deaminase deficiency and severe combined immunodeficiency, hyper immunoglobulin M syndrome, Chediak-Higashi disease, hereditary lymphohistiocytosis, osteopetrosis, osteogenesis imperfecta, storage diseases, thalassemia major, systemic sclerosis, systemic lupus erythematosus, multiple sclerosis, juvenile rheumatoid arthritis, an autoimmune disorder, and a neurological disorder.

15 In some embodiments, the immunodeficiency disorder is a congenital immunodeficiency.

In some embodiments, the immunodeficiency disorder is an acquired immunodeficiency.

In some embodiments, the acquired immunodeficiency is human immunodeficiency virus or acquired immune deficiency syndrome.

In some embodiments, the metabolic disorder is selected from the group consisting of glycogen
 20 storage diseases, mucopolysaccharidoses, Gaucher's Disease, Hurlers Disease, sphingolipidoses, and metachromatic leukodystrophy.

In some embodiments, the cancer is a hematological cancer.

In some embodiments, the cancer is selected from the group consisting of leukemia, lymphoma, multiple myeloma, and neuroblastoma.

25 In some embodiments, the cancer is acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia, chronic lymphoid leukemia, multiple myeloma, diffuse large B-cell lymphoma, or non-Hodgkin's lymphoma.

In some embodiments, the autoimmune disorder is selected from the group consisting of multiple sclerosis, human systemic lupus, rheumatoid arthritis, inflammatory bowel disease, treating psoriasis,
 30 Type 1 diabetes mellitus, acute disseminated encephalomyelitis, Addison's disease, alopecia universalis, ankylosing spondylitis, antiphospholipid antibody syndrome, aplastic anemia, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune inner ear disease, autoimmune lymphoproliferative syndrome, autoimmune oophoritis, Balo disease, Behcet's disease, bullous pemphigoid, cardiomyopathy, Chagas' disease, chronic fatigue immune dysfunction syndrome, chronic inflammatory demyelinating
 35 polyneuropathy, Crohn's disease, cicatricial pemphigoid, coeliac sprue-dermatitis herpetiformis, cold agglutinin disease, CREST syndrome, Degos disease, discoid lupus, dysautonomia, endometriosis, essential mixed cryoglobulinemia, fibromyalgia-fibromyositis, Goodpasture's syndrome, Grave's disease, Guillain-Barre syndrome, Hashimoto's thyroiditis, Hidradenitis suppurativa, idiopathic and/or acute thrombocytopenic purpura, idiopathic pulmonary fibrosis, IgA neuropathy, interstitial cystitis, juvenile
 40 arthritis, Kawasaki's disease, lichen planus, Lyme disease, Meniere disease, mixed connective tissue disease, myasthenia gravis, neuromyotonia, opsoclonus myoclonus syndrome, optic neuritis, Ord's

thyroiditis, pemphigus vulgaris, pernicious anemia, polychondritis, polymyositis and dermatomyositis, primary biliary cirrhosis, polyarteritis nodosa, polyglandular syndromes, polymyalgia rheumatica, primary agammaglobulinemia, Raynaud phenomenon, Reiter's syndrome, rheumatic fever, sarcoidosis, scleroderma, Sjögren's syndrome, stiff person syndrome, Takayasu's arteritis, temporal arteritis, ulcerative colitis, uveitis, vasculitis, vitiligo, vulvodynia, and Wegener's granulomatosis.

In some embodiments, the neurological disorder is selected from the group consisting of Parkinson's disease, Alzheimer's disease, multiple sclerosis, Amyotrophic lateral sclerosis, Huntington's disease, mild cognitive impairment, amyloidosis, AIDS-related dementia, encephalitis, stroke, head trauma, epilepsy, mood disorders, and dementia.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. In the specification, the singular forms also include the plural unless the context clearly dictates otherwise. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below. All publications, patent applications, patents and other references mentioned herein are incorporated by reference. The references cited herein are not admitted to be prior art to the claimed invention. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods and examples are illustrative only and are not intended to be limiting. In the case of conflict between the chemical structures and names of the compounds disclosed herein, the chemical structures will control.

Other features and advantages of the disclosure will be apparent from the following detailed description and claims.

Brief Description of the Figures

FIG. 1 is a graph demonstrating the effect of compound (5) and compound (16) on the aryl hydrocarbon receptor-driven expression of luciferase in the absence of the aryl hydrocarbon receptor agonist VAF347 in transiently transfected HepG2 cells in vitro. Experimental details for this experiment are reported in Example 3, below.

FIG. 2 is a graph demonstrating the effect of compound (5) and compound (16) on the aryl hydrocarbon receptor-driven expression of luciferase in the presence of the aryl hydrocarbon receptor agonist VAF347 in transiently transfected HepG2 cells in vitro. Experimental details for this experiment are reported in Example 3, below.

FIG. 3 is a graph demonstrating the effect of compound (5) and compound (16) on the quantity of CD34+ cells in a hematopoietic stem cell population over the course of a seven-day experiment. Experimental details for this experiment are reported in Example 3, below.

FIG. 4 is a scheme showing the design of experiments, described in Example 5, below, aimed at examining the ability of hematopoietic stem cells to migrate to central nervous system tissue and promote the engraftment of microglial cells in the brain.

FIGS. 5A and 5B are graphs showing the ability of hematopoietic stem cells expanded, ex vivo, in the presence of compound (16) to increase the frequency of CD45+ cells in peripheral blood of NSG

mice, and to promote the engraftment of microglial cells in the brains of NSG mice. Each bar graph shows the median value obtained upon examination of n=8 NSG mice.

FIGS. 6A and 6B are graphs showing the ability of hematopoietic stem cells expanded, ex vivo, in the presence of compound (24) to increase the frequency of CD45+ cells in peripheral blood of NSG mice, and to promote the engraftment of microglial cells in the brains of NSG mice. Each bar graph shows the median value obtained upon examination of n=6-8 NSG mice.

FIG. 7 sets forth a ¹H-NMR spectra of compound (24) in DMSO-d₆.

FIG. 8 sets forth a ¹H-NMR spectra of an isolated compound (24) enantiomer peak in DMSO-d₆.

FIG. 9 sets forth a ¹H-NMR spectra of an isolated compound (24) enantiomer peak in DMSO-d₆.

FIG. 10 sets forth a ¹H-NMR spectra of compound (27) in chloroform-d.

FIG. 11 sets forth a ¹H-NMR spectra of compound (28) in chloroform-d.

FIG. 12 is a graph demonstrating expansion capabilities of AHR antagonists in terms of CD34+ frequency. Experimental details for this experiment are reported in Example 9, below.

FIG. 13 is a graph demonstrating expansion capabilities of AHR antagonists in terms of CD34+ number. Experimental details for this experiment are reported in Example 9, below.

FIG. 14 is a graph demonstrating endogenous AHR antagonist activity. Experimental details for this experiment are reported in Example 9, below.

FIG. 15 is a graph demonstrating AHR antagonist activity in the presence of VAF347.

Experimental details for this experiment are reported in Example 9, below.

Detailed Description

The compositions and methods described herein provide tools for expanding hematopoietic stem cells, for instance, by culturing hematopoietic stem cells ex vivo in the presence of an aryl hydrocarbon receptor antagonist represented by formula (I) or (II) described herein. It has presently been discovered that aryl hydrocarbon receptor antagonists of the formula (I) or (II) described herein are capable of inducing the proliferation of hematopoietic stem cells while maintaining the hematopoietic stem cell functional potential of the ensuing cells. As hematopoietic stem cells exhibit the ability to differentiate into a multitude of cell types within the hematopoietic lineage, the aryl hydrocarbon receptor antagonists described herein can be used to amplify a population of hematopoietic stem cells prior to transplantation of the hematopoietic stem cells to a patient in need thereof. Exemplary patients in need of a hematopoietic stem cell transplant are those suffering from a hemoglobinopathy, immunodeficiency, or metabolic disease, such as one of the various pathologies described herein.

Despite the promise of hematopoietic stem cell transplant therapy, methods of expanding hematopoietic stem cells ex vivo to produce quantities sufficient for transplantation has been challenging due to the propensity of hematopoietic stem cells to differentiate upon proliferation. The aryl hydrocarbon receptor antagonists described herein represent a solution to this long-standing difficulty, as the compounds set forth herein are capable of inducing the expansion of hematopoietic stem cells while preserving their capacity for reconstituting various populations of cells in the hematopoietic family. The compositions described herein therefore provide useful tools for the proliferation of hematopoietic stem cells prior to hematopoietic stem cell transplant therapy, and thus constitute methods of treating a variety of hematopoietic conditions, such as sickle cell anemia, thalassemia, Fanconi anemia, Wiskott-Aldrich

syndrome, adenosine deaminase deficiency-severe combined immunodeficiency, metachromatic leukodystrophy, Diamond-Blackfan anemia and Schwachman-Diamond syndrome, human immunodeficiency virus infection, and acquired immune deficiency syndrome, among others.

5 Definitions

Listed below are definitions of various terms used in this application. These definitions apply to terms as they are used throughout this specification and claims, unless otherwise limited in specific instances, either individually or as part of a larger group.

As used herein, the term "about" refers to a value that is within 10% above or below the value being described. For example, the term "about 5 nM" indicates a range of from 4.5 nM to 5.5 nM.

As used herein, the term "donor" refers to a human or animal from which one or more cells are isolated prior to administration of the cells, or progeny thereof, into a recipient. The one or more cells may be, for example, a population of hematopoietic stem cells.

As used herein, the term "endogenous" describes a substance, such as a molecule, cell, tissue, or organ (for example, a hematopoietic stem cell or a cell of hematopoietic lineage, such as a megakaryocyte, thrombocyte, platelet, erythrocyte, mast cell, myoblast, basophil, neutrophil, eosinophil, microglial cell, granulocyte, monocyte, osteoclast, antigen-presenting cell, macrophage, dendritic cell, natural killer cell, T-lymphocyte, or B-lymphocyte) that is found naturally in a particular organism, such as a human patient.

As used herein, the term "engraftment potential" is used to refer to the ability of hematopoietic stem and progenitor cells to repopulate a tissue, whether such cells are naturally circulating or are provided by transplantation. The term encompasses all events surrounding or leading up to engraftment, such as tissue homing of cells and colonization of cells within the tissue of interest. The engraftment efficiency or rate of engraftment can be evaluated or quantified using any clinically acceptable parameter as known to those of skill in the art and can include, for example, assessment of competitive repopulating units (CRU); incorporation or expression of a marker in tissue(s) into which stem cells have homed, colonized, or become engrafted; or by evaluation of the progress of a subject through disease progression, survival of hematopoietic stem and progenitor cells, or survival of a recipient. Engraftment can also be determined by measuring white blood cell counts in peripheral blood during a post-transplant period. Engraftment can also be assessed by measuring recovery of marrow cells by donor cells in a bone marrow aspirate sample.

As used herein, the term "exogenous" describes a substance, such as a molecule, cell, tissue, or organ (for example, a hematopoietic stem cell or a cell of hematopoietic lineage, such as a megakaryocyte, thrombocyte, platelet, erythrocyte, mast cell, myoblast, basophil, neutrophil, eosinophil, microglial cell, granulocyte, monocyte, osteoclast, antigen-presenting cell, macrophage, dendritic cell, natural killer cell, T-lymphocyte, or B-lymphocyte) that is not found naturally in a particular organism, such as a human patient. Exogenous substances include those that are provided from an external source to an organism or to cultured matter extracted therefrom.

As used herein, the term "expanding amount" refers to a quantity or concentration of an agent, such as an aryl hydrocarbon receptor antagonist described herein, sufficient to induce the proliferation of a population of CD34+ cells (e.g., a CD34+ CD90+ cells), for example, by from about 1.1-fold to about

1,000-fold, or more (e.g., about 1.1-fold, 1.2-fold, 1.3-fold, 1.4-fold, 1.5-fold, 1.6-fold, 1.7-fold, 1.8-fold, 1.9-fold, 2-fold, 2.1-fold, 2.2-fold, 2.3-fold, 2.4-fold, 2.5-fold, 2.6-fold, 2.7-fold, 2.8-fold, 2.9-fold, 3-fold, 3.1-fold, 3.2-fold, 3.3-fold, 3.4-fold, 3.5-fold, 3.6-fold, 3.7-fold, 3.8-fold, 3.9-fold, 4-fold, 4.1-fold, 4.2-fold, 4.3-fold, 4.4-fold, 4.5-fold, 4.6-fold, 4.7-fold, 4.8-fold, 4.9-fold, 5-fold, 5.1-fold, 5.2-fold, 5.3-fold, 5.4-fold, 5.5-fold, 5.6-fold, 5.7-fold, 5.8-fold, 5.9-fold, 6-fold, 6.1-fold, 6.2-fold, 6.3-fold, 6.4-fold, 6.5-fold, 6.6-fold, 6.7-fold, 6.8-fold, 6.9-fold, 7-fold, 7.1-fold, 7.2-fold, 7.3-fold, 7.4-fold, 7.5-fold, 7.6-fold, 7.7-fold, 7.8-fold, 7.9-fold, 8-fold, 8.1-fold, 8.2-fold, 8.3-fold, 8.4-fold, 8.5-fold, 8.6-fold, 8.7-fold, 8.8-fold, 8.9-fold, 9-fold, 9.1-fold, 9.2-fold, 9.3-fold, 9.4-fold, 9.5-fold, 9.6-fold, 9.7-fold, 9.8-fold, 9.9-fold, 10-fold, 50-fold, 100-fold, 200-fold, 300-fold, 400-fold, 500-fold, 600-fold, 700-fold, 800-fold, 900-fold, 1,000-fold, or more).

As used herein, the term "hematopoietic stem cells" ("HSCs") refers to immature blood cells having the capacity to self-renew and to differentiate into mature blood cells comprising diverse lineages including but not limited to granulocytes (e.g., promyelocytes, neutrophils, eosinophils, basophils), erythrocytes (e.g., reticulocytes, erythrocytes), thrombocytes (e.g., megakaryoblasts, platelet producing megakaryocytes, platelets), monocytes (e.g., monocytes, macrophages), dendritic cells, microglia, osteoclasts, and lymphocytes (e.g., NK cells, B-cells and T-cells). Such cells may include CD34⁺ cells. CD34⁺ cells are immature cells that express the CD34 cell surface marker. In humans, CD34⁺ cells are believed to include a subpopulation of cells with the stem cell properties defined above, whereas in mice, HSCs are CD34⁻. In addition, HSCs also refer to long term repopulating HSCs (LT-HSC) and short term repopulating HSCs (ST-HSC). LT-HSCs and ST-HSCs are differentiated, based on functional potential and on cell surface marker expression. For example, human HSCs are CD34⁺, CD38⁻, CD45RA⁻, CD90⁺, CD49F⁺, and lin⁻ (negative for mature lineage markers including CD2, CD3, CD4, CD7, CD8, CD10, CD11B, CD19, CD20, CD56, CD235A). In mice, bone marrow LT-HSCs are CD34⁻, SCA-1⁺, C-kit⁺, CD135⁻, Slamf1/CD150⁺, CD48⁻, and lin⁻ (negative for mature lineage markers including Ter119, CD11b, Gr1, CD3, CD4, CD8, B220, IL7ra), whereas ST-HSCs are CD34⁺, SCA-1⁺, C-kit⁺, CD135⁻, Slamf1/CD150⁺, and lin⁻ (negative for mature lineage markers including Ter119, CD11b, Gr1, CD3, CD4, CD8, B220, IL7ra). In addition, ST-HSCs are less quiescent and more proliferative than LT-HSCs under homeostatic conditions. However, LT-HSC have greater self renewal potential (i.e., they survive throughout adulthood, and can be serially transplanted through successive recipients), whereas ST-HSCs have limited self renewal (i.e., they survive for only a limited period of time, and do not possess serial transplantation potential). Any of these HSCs can be used in the methods described herein. ST-HSCs are particularly useful because they are highly proliferative and thus, can more quickly give rise to differentiated progeny.

As used herein, the term "hematopoietic progenitor cells" includes pluripotent cells capable of differentiating into several cell types of the hematopoietic system, including, without limitation, granulocytes, monocytes, erythrocytes, megakaryocytes, B-cells and T-cells, among others. Hematopoietic progenitor cells are committed to the hematopoietic cell lineage and generally do not self-renew. Hematopoietic progenitor cells can be identified, for example, by expression patterns of cell surface antigens, and include cells having the following immunophenotype: CD34⁺ or CD34⁺CD90⁻. Hematopoietic progenitor cells include short-term hematopoietic stem cells, multi-potent progenitor cells, common myeloid progenitor cells, granulocyte-monocyte progenitor cells, and megakaryocyte-erythrocyte progenitor cells. The presence of hematopoietic progenitor cells can be determined functionally, for

instance, by detecting colony-forming unit cells, e.g., in complete methylcellulose assays, or phenotypically through the detection of cell surface markers using flow cytometry and cell sorting assays described herein and known in the art.

As used herein, the term "hematopoietic stem cell functional potential" refers to the functional properties of hematopoietic stem cells which include 1) multi-potency (which refers to the ability to differentiate into multiple different blood lineages including, but not limited to, granulocytes (e.g., promyelocytes, neutrophils, eosinophils, basophils), erythrocytes (e.g., reticulocytes, erythrocytes), thrombocytes (e.g., megakaryoblasts, platelet producing megakaryocytes, platelets), monocytes (e.g., monocytes, macrophages), dendritic cells, microglia, osteoclasts, and lymphocytes (e.g., NK cells, B-cells and T-cells), 2) self-renewal (which refers to the ability of hematopoietic stem cells to give rise to daughter cells that have equivalent potential as the mother cell, and further that this ability can repeatedly occur throughout the lifetime of an individual without exhaustion), and 3) the ability of hematopoietic stem cells or progeny thereof to be reintroduced into a transplant recipient whereupon they home to the hematopoietic stem cell niche and re-establish productive and sustained hematopoiesis.

As used herein, patients that are "in need of" a hematopoietic stem cell transplant include patients that exhibit a defect or deficiency in one or more blood cell types, as well as patients having a stem cell disorder. Hematopoietic stem cells generally exhibit 1) multi-potency, and can thus differentiate into multiple different blood lineages including, but not limited to, granulocytes (e.g., promyelocytes, neutrophils, eosinophils, basophils), erythrocytes (e.g., reticulocytes, erythrocytes), thrombocytes (e.g., megakaryoblasts, platelet producing megakaryocytes, platelets), monocytes (e.g., monocytes, macrophages), dendritic cells, microglia, osteoclasts, and lymphocytes (e.g., NK cells, B-cells and T-cells), 2) self-renewal, and can thus give rise to daughter cells that have equivalent potential as the mother cell, and 3) the ability to be reintroduced into a transplant recipient whereupon they home to the hematopoietic stem cell niche and re-establish productive and sustained hematopoiesis. Hematopoietic stem cells can thus be administered to a patient defective or deficient in one or more cell types of the hematopoietic lineage in order to re-constitute the defective or deficient population of cells in vivo. For example, the patient may be suffering from cancer, and the deficiency may be caused by administration of a chemotherapeutic agent or other medicament that depletes, either selectively or non-specifically, the cancerous cell population. Additionally or alternatively, the patient may be suffering from a non-malignant hemoglobinopathy, such as sickle cell anemia, thalassemia, Fanconi anemia, and Wiskott-Aldrich syndrome. The subject may be one that is suffering from adenosine deaminase severe combined immunodeficiency (ADA SCID), HIV/AIDS, metachromatic leukodystrophy, Diamond-Blackfan anemia, and Schwachman-Diamond syndrome. The subject may have or be affected by an inherited blood disorder (e.g., sickle cell anemia) or an autoimmune disorder. Additionally or alternatively, the subject may have or be affected by a malignancy, such as a malignancy selected from the group consisting of hematologic cancers (e.g., leukemia, lymphoma, multiple myeloma, or myelodysplastic syndrome) and neuroblastoma. In some embodiments, the subject has or is otherwise affected by a metabolic disorder. For example, the subject may suffer or otherwise be affected by a metabolic disorder selected from the group consisting of glycogen storage diseases, mucopolysaccharidoses, Gaucher's Disease, Hurlers Disease, sphingolipidoses, metachromatic leukodystrophy, or any other diseases or disorders which may benefit from the treatments and therapies disclosed herein and including, without limitation, severe

combined immunodeficiency, Wiscott-Aldrich syndrome, hyper immunoglobulin M (IgM) syndrome, Chediak-Higashi disease, hereditary lymphohistiocytosis, osteopetrosis, osteogenesis imperfecta, storage diseases, thalassemia major, sickle cell disease, systemic sclerosis, systemic lupus erythematosus, multiple sclerosis, juvenile rheumatoid arthritis and those diseases, or disorders described in "Bone Marrow Transplantation for Non-Malignant Disease," ASH Education Book, 1:319-338 (2000), the disclosure of which is incorporated herein by reference in its entirety as it pertains to pathologies that may be treated by administration of hematopoietic stem cell transplant therapy. Additionally or alternatively, a patient "in need of" a hematopoietic stem cell transplant may be one that is or is not suffering from one of the foregoing pathologies, but nonetheless exhibits a reduced level (e.g., as compared to that of an otherwise healthy subject) of one or more endogenous cell types within the hematopoietic lineage, such as megakaryocytes, thrombocytes, platelets, erythrocytes, mast cells, myoblasts, basophils, neutrophils, eosinophils, microglia, granulocytes, monocytes, osteoclasts, antigen-presenting cells, macrophages, dendritic cells, natural killer cells, T-lymphocytes, and B-lymphocytes. One of skill in the art can readily determine whether one's level of one or more of the foregoing cell types, or other blood cell type, is reduced with respect to an otherwise healthy subject, for instance, by way of flow cytometry and fluorescence activated cell sorting (FACS) methods, among other procedures, known in the art.

As used herein, the term "recipient" refers to a patient that receives a transplant, such as a transplant containing a population of hematopoietic stem cells. The transplanted cells administered to a recipient may be, e.g., autologous, syngeneic, or allogeneic cells.

As used herein, the term "sample" refers to a specimen (e.g., blood, blood component (e.g., serum or plasma), urine, saliva, amniotic fluid, cerebrospinal fluid, tissue (e.g., placental or dermal), pancreatic fluid, chorionic villus sample, and cells) taken from a subject.

As used herein, the terms "subject" and "patient" refer to an organism, such as a human, that receives treatment for a particular disease or condition as described herein. For instance, a patient, such as a human patient, that is in need of hematopoietic stem cell transplantation may receive treatment that includes a population of hematopoietic stem cells so as to treat a stem cell disorder, such as a cancer, autoimmune disease, or metabolic disorder described herein. For instance, a patient, such as a human patient suffering from a stem cell disorder, may receive treatment in the form of a population of hematopoietic stem cells, such as a population of from about 1×10^6 to about 1×10^9 hematopoietic stem cells.

As used herein, the phrase "stem cell disorder" broadly refers to any disease, disorder, or condition that may be treated or cured by engrafting or transplanting a population of hematopoietic stem or progenitor cells in a target tissue within a patient. For example, Type I diabetes has been shown to be cured by hematopoietic stem cell transplant, along with various other disorders. Diseases that can be treated by infusion of hematopoietic stem or progenitor cells into a patient include, sickle cell anemia, thalassemias, Fanconi anemia, aplastic anemia, Wiskott-Aldrich syndrome, ADA SCID, HIV/AIDS, metachromatic leukodystrophy, Diamond-Blackfan anemia, and Schwachman-Diamond syndrome. Additional diseases that may be treated by transplantation of hematopoietic stem and progenitor cells as described herein include blood disorders (e.g., sickle cell anemia) and autoimmune disorders, such as scleroderma, multiple sclerosis, ulcerative colitis, and Crohn's disease. Additional diseases that may be treated using hematopoietic stem and progenitor cell transplant therapy include cancer, such as a cancer

described herein. Stem cell disorders include a malignancy, such as a neuroblastoma or a hematologic cancers, such as leukemia, lymphoma, and myeloma. For instance, the cancer may be acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia, chronic lymphoid leukemia, multiple myeloma, diffuse large B-cell lymphoma, or non-Hodgkin's lymphoma. Disorders that may be treated by transplanting a population of hematopoietic stem cells to a patient include neurological disorders, such as Parkinson's disease, Alzheimer's disease, multiple sclerosis, Amyotrophic lateral sclerosis, Huntington's disease, mild cognitive impairment, amyloidosis, AIDS-related dementia, encephalitis, stroke, head trauma, epilepsy, mood disorders, and dementia. As described herein, without being limited by mechanism, the ability of hematopoietic stem cell transplantation to treat such disorders may be due, in part, to the capacity of hematopoietic stem cells to migrate to the central nervous system and differentiate into microglial cells, thereby repopulating a hematopoietic cell line that may be damaged or deficient in patients having a neurological disorder. Additional diseases treatable using hematopoietic stem or progenitor cell transplant therapy include myelodysplastic syndrome. In some embodiments, the patient has or is otherwise affected by a metabolic storage disorder. For example, the patient may suffer or otherwise be affected by a metabolic disorder selected from the group consisting of glycogen storage diseases, mucopolysaccharidoses, Gaucher's Disease, Hurlers Disease, sphingolipidoses, metachromatic leukodystrophy, or any other diseases or disorders which may benefit from the treatments and therapies disclosed herein and including, without limitation, severe combined immunodeficiency, Wiscott-Aldrich syndrome, hyper immunoglobulin M (IgM) syndrome, Chediak-Higashi disease, hereditary lymphohistiocytosis, osteopetrosis, osteogenesis imperfecta, storage diseases, thalassemia major, sickle cell disease, systemic sclerosis, systemic lupus erythematosus, multiple sclerosis, juvenile rheumatoid arthritis and those diseases, or disorders described in "Bone Marrow Transplantation for Non-Malignant Disease," ASH Education Book, 1:319-338 (2000), the disclosure of which is incorporated herein by reference in its entirety as it pertains to pathologies that may be treated by administration of hematopoietic stem or progenitor cell transplant therapy.

As used herein, the terms "treat", "treating" or "treatment" refer to a method of alleviating or abating a disease and/or its attendant symptoms. As used herein, the terms "preventing" or "prevent" describes reducing or eliminating the onset of the symptoms or complications of the disease, condition, or disorder. As used herein, the terms "disease(s)", "disorder(s)", and "condition(s)" are used interchangeably, unless the context clearly dictates otherwise.

"Treating" may refer to therapeutic treatment, in which the object is to prevent or slow down (lessen) an undesired physiological change or disorder or to promote a beneficial phenotype in the patient being treated. Beneficial or desired clinical results include, but are not limited to, the observation of an increase in the cell count or relative concentration of hematopoietic stem cells in a patient in need of a hematopoietic stem cell transplant following administration of an exogenous hematopoietic stem cell graft to the patient. Beneficial results of therapy described herein may also include an increase in the cell count or relative concentration of one or more cells of hematopoietic lineage, such as a megakaryocyte, thrombocyte, platelet, erythrocyte, mast cell, myoblast, basophil, neutrophil, eosinophil, microglial cell, granulocyte, monocyte, osteoclast, antigen-presenting cell, macrophage, dendritic cell, natural killer cell, T-lymphocyte, or B-lymphocyte, following conditioning therapy and subsequent hematopoietic stem cell

transplant therapy. Additional beneficial results may include the reduction in quantity of a disease-causing cell population, such as a population of cancer cells or auto-immune cells.

As used herein, the terms "variant" and "derivative" are used interchangeably and refer to naturally-occurring, synthetic, and semi-synthetic analogues of a compound, peptide, protein, or other substance described herein. A variant or derivative of a compound, peptide, protein, or other substance described herein may retain or improve upon the biological activity of the original material.

As used herein, the term "alkyl" refers to a straight- or branched-chain alkyl group having, for example, from 1 to 20 carbon atoms in the chain, or, in certain embodiments, from 1 to 6 carbon atoms in the chain. Examples of alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, tert-pentyl, neopentyl, isopentyl, tert-pentyl, hexyl, isohexyl, and the like.

As used herein, the term "alkylene" refers to a straight- or branched-chain divalent alkyl group. The divalent positions may be on the same or different atoms within the alkyl chain. Examples of alkylene include methylene, ethylene, propylene, isopropylene, and the like.

As used herein, the term "heteroalkyl" refers to a straight or branched-chain alkyl group having, for example, from 1 to 20 carbon atoms in the chain, and further containing one or more heteroatoms (e.g., oxygen, nitrogen, or sulfur, among others) in the chain.

As used herein, the term "heteroalkylene" refers to a straight- or branched-chain divalent heteroalkyl group. The divalent positions may be on the same or different atoms within the heteroalkyl chain. The divalent positions may be one or more heteroatoms.

As used herein, the term "alkenyl" refers to a straight- or branched-chain alkenyl group having, for example, from 2 to 20 carbon atoms in the chain. It denotes a monovalent group derived from a hydrocarbon moiety containing, for example, from two to six carbon atoms having at least one carbon-carbon double bond. The double bond may or may not be the point of attachment to another group. Examples of alkenyl groups include, but are not limited to, vinyl, propenyl, isopropenyl, butenyl, tert-butenyl, 1-methyl-2-buten-1-yl, hexenyl, and the like.

As used herein, the term "alkenylene" refers to a straight- or branched-chain divalent alkenyl group. The divalent positions may be on the same or different atoms within the alkenyl chain. Examples of alkenylene include ethenylene, propenylene, isopropenylene, butenylene, and the like.

As used herein, the term "heteroalkenyl" refers to a straight- or branched-chain alkenyl group having, for example, from 2 to 20 carbon atoms in the chain, and further containing one or more heteroatoms (e.g., oxygen, nitrogen, or sulfur, among others) in the chain.

As used herein, the term "heteroalkenylene" refers to a straight- or branched-chain divalent heteroalkenyl group. The divalent positions may be on the same or different atoms within the heteroalkenyl chain. The divalent positions may be one or more heteroatoms.

As used herein, the term "alkynyl" refers to a straight- or branched-chain alkynyl group having, for example, from 2 to 20 carbon atoms in the chain and at least one carbon-carbon triple bond. Examples of alkynyl groups include, but are not limited to, propargyl, butynyl, pentynyl, hexynyl, and the like.

As used herein, the term "alkynylene" refers to a straight- or branched-chain divalent alkynyl group. The divalent positions may be on the same or different atoms within the alkynyl chain.

As used herein, the term "heteroalkynyl" refers to a straight- or branched-chain alkynyl group having, for example, from 2 to 20 carbon atoms in the chain, and further containing one or more heteroatoms (e.g., oxygen, nitrogen, or sulfur, among others) in the chain.

As used herein, the term "heteroalkynylene" refers to a straight- or branched-chain divalent heteroalkynyl group. The divalent positions may be on the same or different atoms within the heteroalkynyl chain. The divalent positions may be one or more heteroatoms.

As used herein, the term "cycloalkyl" refers to a monocyclic, or fused, bridged, or spiro polycyclic ring structure that is saturated and has, for example, from 3 to 12 carbon ring atoms. Examples of cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, bicyclo[3.1.0]hexane, and the like. Also contemplated is a monovalent group derived from a monocyclic or polycyclic carbocyclic ring compound having at least one carbon-carbon double bond by the removal of at least one or two hydrogen atoms. Examples of such groups include, but are not limited to, cyclopropenyl, cyclobutenyl, cyclopentenyl,

As used herein, the term "cycloalkylene" refers to a divalent cycloalkyl group. The divalent positions may be on the same or different atoms within the ring structure. Examples of cycloalkylene include cyclopropylene, cyclobutylene, cyclopentylene, cyclohexylene, and the like.

As used herein, the term "heterocycloalkyl" or "heterocyclyl" refers to a monocyclic, or fused, bridged, or spiro polycyclic ring structure that is saturated and has, for example, from 3 to 12 ring atoms per ring structure selected from carbon atoms and heteroatoms selected from, e.g., nitrogen, oxygen, and sulfur, among others. The ring structure may contain, for example, one or more oxo groups on carbon, nitrogen, or sulfur ring members. Exemplary heterocycloalkyl groups include, but are not limited to, [1,3] dioxolane, pyrrolidinyl, pyrazolinyl, pyrazolidinyl, imidazolinyl, imidazolidinyl, piperazinyl, piperidinyl, oxazolidinyl, isooxazolidinyl, morpholinyl, thiazolidinyl, isothiazolidinyl, and tetrahydrofuryl.

As used herein, the term "heterocycloalkylene" refers to a divalent heterocycloalkyl group. The divalent positions may be on the same or different atoms within the ring structure.

As used herein, the term "aryl" refers to a monocyclic or multicyclic aromatic ring system containing, for example, from 6 to 19 carbon atoms. Aryl groups include, but are not limited to, phenyl, fluorenyl, naphthyl, tetrahydronaphthyl, indanyl, indenyl, and the like. The divalent positions may be one or more heteroatoms.

As used herein, the term "arylene" refers to a divalent aryl group. The divalent positions may be on the same or different atoms.

As used herein, the term "heteroaryl" refers to a monocyclic heteroaromatic, or a bicyclic or a tricyclic fused-ring heteroaromatic group. In certain embodiments, the heteroaryl group contains five to ten ring atoms of which one ring atom is selected from S, O, and N; zero, one, or two ring atoms are additional heteroatoms independently selected from S, O, and N; and the remaining ring atoms are carbon. Heteroaryl groups include, but are not limited to, pyridyl, pyrrolyl, furyl, thienyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,3,4-triazinyl, 1,2,3-triazinyl, benzofuryl, [2,3-dihydro]benzofuryl, isobenzofuryl, benzothienyl, benzotriazolyl, isobenzothienyl, indolyl, isoindolyl, 3H-indolyl, benzimidazolyl, imidazo[1,2-a]pyridyl, benzothiazolyl, benzoxazolyl, quinoliziny, quinazolinyl, phthalazinyl, quinoxalinyl, cinnolinyl, naphthyridinyl, pyrido[3,4-b]pyridyl, pyrido[3,2-b]pyridyl, pyrido[4,3-

b]pyridyl, quinolyl, isoquinolyl, tetrazolyl, 5,6,7,8-tetrahydroquinolyl, 5,6,7,8-tetrahydroisoquinolyl, purinyl, pteridinyl, carbazolyl, xanthenyl, benzoquinolyl, and the like.

As used herein, the term "heteroarylene" refers to a divalent heteroaryl group. The divalent positions may be on the same or different atoms. The divalent positions may be one or more
5 heteroatoms.

Unless otherwise constrained by the definition of the individual substituent, the foregoing chemical moieties, such as "alkyl", "alkylene", "heteroalkyl", "heteroalkylene", "alkenyl", "alkenylene", "heteroalkenyl", "heteroalkenylene", "alkynyl", "alkynylene", "heteroalkynyl", "heteroalkynylene", "cycloalkyl", "cycloalkylene", "heterocycloalkyl", heterocycloalkylene, "aryl", "arylene", "heteroaryl", and
10 "heteroarylene" groups can optionally be substituted. As used herein, the term "optionally substituted" refers to a compound or moiety containing one or more (for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) substituents, as permitted by the valence of the compound or moiety or a site thereof, such as a substituent selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, alkyl aryl, alkyl heteroaryl, alkyl cycloalkyl, alkyl heterocycloalkyl, amino, ammonium, acyl, acyloxy, acylamino,
15 aminocarbonyl, alkoxy, carbamate, ureido, carbamate, aryl, heteroaryl, sulfinyl, sulfonyl, alkoxy, sulfanyl, halogen, carboxy, trihalomethyl, cyano, hydroxy, mercapto, nitro, and the like. The substitution may include situations in which neighboring substituents have undergone ring closure, such as ring closure of vicinal functional substituents, to form, for instance, lactams, lactones, cyclic anhydrides, acetals, hemiacetals, thioacetals, amins, and hemiaminals, formed by ring closure, for example, to furnish a
20 protecting group.

As used herein, the term "optionally substituted" refers to a chemical moiety that may have one or more chemical substituents, as valency permits, such as C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-10 cycloalkyl, C2-10 heterocycloalkyl, C2-10 aryl, C2-10 alkylaryl, C2-10 heteroaryl, C2-10 alkylheteroaryl, amino, ammonium, acyl, acyloxy, acylamino, aminocarbonyl, alkoxy, carbamate, sulfinyl,
25 sulfonyl, alkoxy, sulfanyl, halogen, carboxy, trihalomethyl, cyano, hydroxy, mercapto, nitro, and the like. An optionally substituted chemical moiety may contain, e.g., neighboring substituents that have undergone ring closure, such as ring closure of vicinal functional substituents, thus forming, e.g., lactams, lactones, cyclic anhydrides, acetals, thioacetals, or amins formed by ring closure, for instance, in order to generate protecting group.

30 In accordance with the application, any of the aryls, substituted aryls, heteroaryls and substituted heteroaryls described herein, can be any aromatic group.


The terms "hal," "halo," and "halogen," as used herein, refer to an atom selected from fluorine, chlorine, bromine and iodine.

As described herein, compounds of the application and moieties present in the compounds may
35 optionally be substituted with one or more substituents, such as are illustrated generally above, or as exemplified by particular classes, subclasses, and species of the application. It will be appreciated that the phrase "optionally substituted" is used interchangeably with the phrase "substituted or unsubstituted." In general, the term "substituted", whether preceded by the term "optionally" or not, refers to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. Unless
40 otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than

- one substituent selected from a specified group, the substituent may be either the same or different at every position. The terms "optionally substituted", "optionally substituted alkyl", "optionally substituted alkenyl", "optionally substituted alkynyl", "optionally substituted cycloalkyl", "optionally substituted cycloalkenyl", "optionally substituted aryl", "optionally substituted heteroaryl", "optionally substituted aralkyl", "optionally substituted heteroaralkyl", "optionally substituted heterocycloalkyl", and any other optionally substituted group as used herein, refer to groups that are substituted or unsubstituted by independent replacement of one, two, or three or more of the hydrogen atoms thereon with substituents including, but not limited to:
- F, -Cl, -Br, -I, -OH, protected hydroxy, -NO₂, -CN, -NH₂, protected amino, -NH-C₁-C₁₂-alkyl, -NH-C₂-C₁₂-alkenyl, -NH-C₂-C₁₂-alkenyl, -NH-C₃-C₁₂-cycloalkyl, -NH-aryl, -NH-heteroaryl, -NH-heterocycloalkyl, -dialkylamino, -diarylamino, -diheteroarylamino, -O-C₁-C₁₂-alkyl, -O-C₂-C₁₂-alkenyl, -O-C₂-C₁₂-alkenyl, -O-C₃-C₁₂-cycloalkyl, -O-aryl, -O-heteroaryl, -O-heterocycloalkyl, -C(O)-C₁-C₁₂-alkyl, -C(O)-C₂-C₁₂-alkenyl, -C(O)-C₂-C₁₂-alkenyl, -C(O)-C₃-C₁₂-cycloalkyl, -C(O)-aryl, -C(O)-heteroaryl, -C(O)-heterocycloalkyl, -CONH₂, -CONH-C₁-C₁₂-alkyl, -CONH-C₂-C₁₂-alkenyl, -CONH-C₂-C₁₂-alkenyl, -CONH-C₃-C₁₂-cycloalkyl, -CONH-aryl, -CONH-heteroaryl, -CONH-heterocycloalkyl, -OCO₂-C₁-C₁₂-alkyl, -OCO₂-C₂-C₁₂-alkenyl, -OCO₂-C₃-C₁₂-cycloalkyl, -OCO₂-aryl, -OCO₂-heteroaryl, -OCO₂-heterocycloalkyl, -OCONH₂, -OCONH-C₁-C₁₂-alkyl, -OCONH-C₂-C₁₂-alkenyl, -OCONH-C₂-C₁₂-alkenyl, -OCONH-C₃-C₁₂-cycloalkyl, -OCONH-aryl, -OCONH-heteroaryl, -OCONH-heterocycloalkyl, -NHC(O)-C₁-C₁₂-alkyl, -NHC(O)-C₂-C₁₂-alkenyl, -NHC(O)-C₂-C₁₂-alkenyl, -NHC(O)-C₃-C₁₂-cycloalkyl, -NHC(O)-aryl, -NHC(O)-heteroaryl, -NHC(O)-heterocycloalkyl, -NHCO₂-C₁-C₁₂-alkyl, -NHCO₂-C₂-C₁₂-alkenyl, -NHCO₂-C₂-C₁₂-alkenyl, -NHCO₂-C₃-C₁₂-cycloalkyl, -NHCO₂-aryl, -NHCO₂-heteroaryl, -NHCO₂-heterocycloalkyl, -NHC(O)NH₂, -NHC(O)NH-C₁-C₁₂-alkyl, -NHC(O)NH-C₂-C₁₂-alkenyl, -NHC(O)NH-C₂-C₁₂-alkenyl, -NHC(O)NH-C₃-C₁₂-cycloalkyl, -NHC(O)NH-aryl, -NHC(O)NH-heteroaryl, -NHC(O)NH-heterocycloalkyl, -NHC(S)NH₂, -NHC(S)NH-C₁-C₁₂-alkyl, -NHC(S)NH-C₂-C₁₂-alkenyl, -NHC(S)NH-C₂-C₁₂-alkenyl, -NHC(S)NH-C₃-C₁₂-cycloalkyl, -NHC(S)NH-aryl, -NHC(S)NH-heteroaryl, -NHC(S)NH-heterocycloalkyl, -NHC(NH)NH₂, -NHC(NH)NH-C₁-C₁₂-alkyl, -NHC(NH)NH-C₂-C₁₂-alkenyl, -NHC(NH)NH-C₂-C₁₂-alkenyl, -NHC(NH)NH-C₃-C₁₂-cycloalkyl, -NHC(NH)NH-aryl, -NHC(NH)NH-heteroaryl, -NHC(NH)NH-heterocycloalkyl, -NHC(NH)NH-C₁-C₁₂-alkyl, -NHC(NH)NH-C₂-C₁₂-alkenyl, -NHC(NH)NH-C₂-C₁₂-alkenyl, -NHC(NH)NH-C₃-C₁₂-cycloalkyl, -C(NH)NH-C₁-C₁₂-alkyl, -C(NH)NH-C₂-C₁₂-alkenyl, -C(NH)NH-C₂-C₁₂-alkenyl, -C(NH)NH-C₃-C₁₂-cycloalkyl, -C(NH)NH-aryl, -C(NH)NH-heteroaryl, -C(NH)NH-heterocycloalkyl, -S(O)-C₁-C₁₂-alkyl, -S(O)-C₂-C₁₂-alkenyl, -S(O)-C₂-C₁₂-alkenyl, -S(O)-C₃-C₁₂-cycloalkyl, -S(O)-aryl, -S(O)-heteroaryl, -S(O)-heterocycloalkyl, -SO₂NH₂, -SO₂NH-C₁-C₁₂-alkyl, -SO₂NH-C₂-C₁₂-alkenyl, -SO₂NH-C₂-C₁₂-alkenyl, -SO₂NH-C₃-C₁₂-cycloalkyl, -SO₂NH-aryl, -SO₂NH-heteroaryl, -SO₂NH-heterocycloalkyl, -NHSO₂-C₁-C₁₂-alkyl, -NHSO₂-C₂-C₁₂-alkenyl, -NHSO₂-C₂-C₁₂-alkenyl, -NHSO₂-C₃-C₁₂-cycloalkyl, -NHSO₂-aryl, -NHSO₂-heteroaryl, -NHSO₂-heterocycloalkyl, -CH₂NH₂, -CH₂SO₂CH₃, -aryl, -arylalkyl, -heteroaryl, -heteroarylalkyl, -heterocycloalkyl, -C₃-C₁₂-cycloalkyl, polyalkoxyalkyl, polyalkoxy, -methoxymethoxy, -methoxyethoxy, -SH, -S-C₁-C₁₂-alkyl, -S-C₂-C₁₂-alkenyl, -S-C₂-C₁₂-alkenyl, -S-C₃-C₁₂-cycloalkyl, -S-aryl, -S-heteroaryl, -S-heterocycloalkyl, or methylthiomethyl.

Compounds of Formula (I) and Formula (II)

The aryl hydrocarbon receptor antagonists described herein include compounds represented by formula (I) or formula (II):



- wherein L is a linker selected from the group consisting of -NR_{7a}(CR_{8a}R_{8b})_n-, -O(CR_{8a}R_{8b})_n-, -C(O)(CR_{8a}R_{8b})_n-, -C(S)(CR_{8a}R_{8b})_n-, -S(O)₀₋₂(CR_{8a}R_{8b})_n-, -(CR_{8a}R_{8b})_n-, -NR_{7a}C(O)(CR_{8a}R_{8b})_n-, -NR_{7a}C(S)(CR_{8a}R_{8b})_n-, -OC(O)(CR_{8a}R_{8b})_n-, -OC(S)(CR_{8a}R_{8b})_n-, -C(O)NR_{7a}(CR_{8a}R_{8b})_n-, -C(S)NR_{7a}(CR_{8a}R_{8b})_n-, -C(O)O(CR_{8a}R_{8b})_n-, -C(S)O(CR_{8a}R_{8b})_n-, -S(O)₂NR_{7a}(CR_{8a}R_{8b})_n-, -NR_{7a}S(O)₂(CR_{8a}R_{8b})_n-, -NR_{7a}C(O)NR_{7b}(CR_{8a}R_{8b})_n-, and -NR_{7a}C(O)O(CR_{8a}R_{8b})_n-, wherein R_{7a}, R_{7b}, R_{8a}, and R_{8b} are each independently selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl, and each n is independently an integer from 2 to 6;

- R₁ is selected from the group consisting of -S(O)₂NR_{9a}R_{9b}-, -NR_{9a}C(O)R_{9b}-, -NR_{9a}C(S)R_{9b}-, -NR_{9a}C(O)NR_{9b}R_{9c}-, -C(O)R_{9a}-, -C(S)R_{9a}-, -S(O)₀₋₂R_{9a}-, -C(O)OR_{9a}-, -C(S)OR_{9a}-, -C(O)NR_{9a}R_{9b}-, -C(S)NR_{9a}R_{9b}-, -NR_{9a}S(O)₂R_{9b}-, -NR_{9a}C(O)OR_{9b}-, -OC(O)CR_{9a}R_{9b}R_{9c}-, -OC(S)CR_{9a}R_{9b}R_{9c}-, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein R_{9a}, R_{9b}, and R_{9c} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

- R₂ is selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl;

- R₃ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

R₄ is selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl;

- R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R₆ is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

- In some embodiments, R₁ is selected from the group consisting of -S(O)₂NR_{9a}R_{9b}-, -NR_{9a}C(O)R_{9b}-, -NR_{9a}C(S)R_{9b}-, -NR_{9a}C(O)NR_{9b}R_{9c}-, -C(O)R_{9a}-, -C(S)R_{9a}-, -S(O)₀₋₂R_{9a}-, -C(O)OR_{9a}-, -C(S)OR_{9a}-, -C(O)NR_{9a}R_{9b}-, -C(S)NR_{9a}R_{9b}-, -NR_{9a}S(O)₂R_{9b}-, -NR_{9a}C(O)OR_{9b}-, -OC(O)CR_{9a}R_{9b}R_{9c}-, -OC(S)CR_{9a}R_{9b}R_{9c}-, phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted, for example, with from 1 to 3 substituents independently selected from the group consisting of

cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{O}(\text{CH}_2)_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{S}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{OS}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, and $-\text{NR}_{10a}\text{S}(\text{O})_2\text{R}_{10b}$, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl. For instance, R_1 may be selected from the group consisting of $-\text{S}(\text{O})_2\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{S})\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{NR}_{9b}\text{R}_{9c}$, $-\text{C}(\text{O})\text{R}_{9a}$, $-\text{C}(\text{S})\text{R}_{9a}$, $-\text{S}(\text{O})_2\text{R}_{9a}$, $-\text{C}(\text{O})\text{OR}_{9a}$, $-\text{C}(\text{S})\text{OR}_{9a}$, $-\text{C}(\text{O})\text{NR}_{9a}\text{R}_{9b}$, $-\text{C}(\text{S})\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{S}(\text{O})_2\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{OR}_{9b}$, $-\text{OC}(\text{O})\text{CR}_{9a}\text{R}_{9b}\text{R}_{9c}$, and $-\text{OC}(\text{S})\text{CR}_{9a}\text{R}_{9b}\text{R}_{9c}$. R_1 may be selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{O}(\text{CH}_2)_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{S}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{OS}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, and $-\text{NR}_{10a}\text{S}(\text{O})_2\text{R}_{10b}$.

In some embodiments, R_2 is hydrogen.

In some embodiments, R_3 is selected from the group consisting of phenyl, thiophenyl, furanyl, 1H-benzimidazolyl, quinolinyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl, wherein the phenyl, thiophenyl, furanyl, 1H-benzimidazolyl, quinolinyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{C}(\text{O})\text{R}_{11a}$, $-\text{S}(\text{O})_2\text{R}_{11a}$, $-\text{C}(\text{O})\text{OR}_{11a}$, and $-\text{C}(\text{O})\text{NR}_{11a}\text{R}_{11b}$, and wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl.

In some embodiments, R_4 is hydrogen.

In some embodiments, R_5 is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl.

In some embodiments, R_5 is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-hydroxypropan-2-yl, and nonan-2-yl.


In some embodiments, R_5 is (S)-1-hydroxypropan-2-yl.

In some embodiments, R_5 is (R)-1-hydroxypropan-2-yl.

In some embodiments, R_5 is (S)-sec-butyl.


In some embodiments, R₅ is (R)-sec-butyl.

In some embodiments, R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



- wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl.

In some embodiments, R₅ is selected from the group consisting of:



in some embodiments, R₅ is (ii);

- In some embodiments, R₅ is selected from the group consisting of 4-methoxybutan-2-yl, (S)-4-methoxybutan-2-yl, (R)-4-methoxybutan-2-yl, 4-ethoxybutan-2-yl, (S)-4-ethoxybutan-2-yl, (R)-4-ethoxybutan-2-yl, 5-methoxypentan-2-yl, (S)-5-methoxypentan-2-yl, (R)-5-methoxypentan-2-yl, 5-ethoxypentan-2-yl, (S)-5-ethoxypentan-2-yl, (R)-5-ethoxypentan-2-yl, 6-methoxyhexan-2-yl, (S)-6-methoxyhexan-2-yl, (R)-6-methoxyhexan-2-yl, 6-ethoxyhexan-2-yl, (S)-6-ethoxyhexan-2-yl, and (R)-6-ethoxyhexan-2-yl.

- 20 In some embodiments, R₅ is (S)-4-methoxybutan-2-yl.

In some embodiments, R₅ is (R)-4-methoxybutan-2-yl.


In some embodiments, R₅ is (S)-5-methoxypentan-2-yl.


In some embodiments, R₅ is (R)-5-methoxypentan-2-yl.

In some embodiments, R₅ is (S)-4-ethoxybutan-2-yl.


- 25 In some embodiments, R₅ is (R)-4-ethoxybutan-2-yl.

Particular aryl hydrocarbon receptor antagonists described herein include:







(4);




(5);




(6);




(7);




(8);




(9);




(10);




(11);




(12);




(13);




(14);




(15);




(16);




(17);




(18);




(19);




(20);




(21);



(22);



(23);



as well as salts thereof.

Where the number of any given substituent is not specified, there may be one or more substituents present. For example, "halo-substituted C1-4 alkyl" may include one or more of the same or different halogens.

When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both *E* and *Z* geometric isomers. Likewise, all tautomeric forms of carbonyl-containing compounds are also intended to be included.

It is to be understood that the compounds provided herein may contain chiral centers. Such chiral centers may be of either the (*R*) or (*S*) configuration, or may be a mixture thereof. Thus, the compounds provided herein may be enantiomerically pure, or may be stereoisomeric or diastereomeric mixtures. As such, one of skill in the art will recognize that administration of a compound in its (*R*) form is equivalent, for compounds that undergo epimerization *in vivo*, to administration of the compound in its (*S*) form.

Compounds described herein include, but are not limited to, those set forth above, as well as any of their isomers, such as diastereomers and enantiomers, as well as salts, esters, amides, thioesters, solvates, and polymorphs thereof, as well as racemic mixtures and pure isomers of the compounds set forth above.

Synthesis


Substituent protecting groups


The synthesis of aryl hydrocarbon receptor antagonists described herein may involve the selective protection and deprotection of alcohols, amines, ketones, sulfhydryls or carboxyl functional groups of a precursor. For example, commonly used protecting groups for amines include carbamates, such as *tert*-butyl, benzyl, 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 9-fluorenylmethyl, allyl, and *m*-nitrophenyl. Other commonly used protecting groups for amines include amides, such as formamides, acetamides, trifluoroacetamides, sulfonamides, trifluoromethanesulfonyl amides, trimethylsilylethanesulfonamides, and *tert*-butylsulfonyl amides. Examples of commonly used protecting groups for carboxyls include esters, such as methyl, ethyl, *tert*-butyl, 9-fluorenylmethyl, 2-(trimethylsilyl)ethoxy methyl, benzyl, diphenylmethyl, *O*-nitrobenzyl, ortho-esters, and halo-esters. Examples of commonly used protecting groups for alcohols include ethers, such as methyl, methoxymethyl, methoxyethoxymethyl, methylthiomethyl, benzyloxymethyl, tetrahydropyranyl, ethoxyethyl, benzyl, 2-naphthylmethyl, *O*-nitrobenzyl, *P*-nitrobenzyl, *P*-methoxybenzyl, 9-phenylxanthyl, trityl (including methoxy-trityls), and silyl ethers. Examples of commonly used protecting groups for sulfhydryls include many of the same protecting groups used for hydroxyls. In addition, sulfhydryls can be protected in a reduced form (e.g., as disulfides) or an oxidized form (e.g., as sulfonic acids, sulfonic esters, or sulfonic amides). Protecting groups can be chosen such that selective conditions (e.g., acidic conditions, basic conditions, catalysis by a nucleophile, catalysis by a Lewis acid, or hydrogenation) are required to remove each, exclusive of other protecting groups in a compound. The conditions required for the addition of protecting groups to amine, alcohol, sulfhydryl, and carboxyl functionalities and the conditions required for their removal are provided in detail, for example, in T.W. Green and P.G.M. Wuts, *Protective Groups in Organic Synthesis* (2nd Ed.), John Wiley & Sons, 1991 and P.J. Kocienski, *Protecting Groups*, Georg Thieme Verlag, 1994.

Exemplary synthetic methods

Aryl hydrocarbon receptor antagonists represented by formula (I) or (II) may be synthesized, for instance, by way of a palladium-catalyzed coupling reaction, such as a process depicted in Scheme 1, below.

Scheme 1. Pd-catalyzed coupling of halogenated imidazopyridine and imidazopyrazine precursors with amines, alcohols, alkynes, boronic esters and other L-R₁, R₃, and/or R₅ synthons






wherein X is C or N, with the proviso that R₂ is absent when X is N;

- L is selected from the group consisting of -NR_{7a}(CR_{8a}R_{8b})_n-, -O(CR_{8a}R_{8b})_n-, -S(O)₀₋₂(CR_{8a}R_{8b})_n-, and -(CR_{8a}R_{8b})_n-, wherein R_{7a}, R_{8a}, and R_{8b} are each independently selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl, and each n is independently an integer from 2 to 6;
- R₁ is selected from the group consisting of -S(O)₂NR_{9a}R_{9b}-, -NR_{9a}C(O)R_{9b}-, -NR_{9a}C(S)R_{9b}-, -NR_{9a}C(O)NR_{9b}R_{9c}-, -C(O)R_{9a}-, -C(S)R_{9a}-, -S(O)₀₋₂R_{9a}-, -C(O)OR_{9a}-, -C(S)OR_{9a}-, -C(O)NR_{9a}R_{9b}-, -C(S)NR_{9a}R_{9b}-, -NR_{9a}S(O)₂R_{9b}-, -NR_{9a}C(O)OR_{9b}-, -OC(O)CR_{9a}R_{9b}R_{9c}-, -OC(S)CR_{9a}R_{9b}R_{9c}-, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein R_{9a}, R_{9b}, and R_{9c} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;
- R₂ is selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl;
- R₃ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;
- R₄ is selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl;
- R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and
- R₆ is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl.
- For instance, when an organoboron species is employed as a synthon such as, for example, R₃-B(pin) and/or R₅-B(pin), Suzuki or Suzuki-Miyaura reaction conditions may be used to join the halogenated imidazopyridine or imidazopyrazine precursor with the organoboron R₃ or R₅ synthon species in the presence of a Pd catalyst, such as, for example Pd(dppf)Cl₂.
- For instance, when L is an amino- or hydroxy-containing linker, Hartwig-Buchwald conditions may be used to join the halogenated precursor with the amine or alcohol in the presence of a Pd catalyst, as shown in Scheme 2, below.

Scheme 2.




Exemplary Hartwig-Buchwald conditions include the use of a Pd catalyst, such as $\text{Pd}_2(\text{dba})_3$, in the presence of dicyclohexylphosphino-2'-(*N,N*-dimethylamino)biphenyl, *t*BuONa, dioxane, 120 °C, microwave irradiation. Hartwig-Buchwald reaction conditions are known in the art and are described, for instance, in Bailey et al., *Bioorganic and Medicinal Chemistry Letters* 19:3602-3606 (2009), the disclosure of which is incorporated herein by reference as it pertains to conditions useful for the Hartwig-Buchwald amination or etherification.

When L is a thiol, nucleophilic aromatic substitution conditions may be employed to join the linker to a halogenated precursor, as shown in Scheme 3, below.

10

Scheme 3.




Nucleophilic aromatic substitution conditions include the use of a base to deprotonate the thiol shown in the above linker-R₁ pair. This reactive modality is particularly useful when the halogenated aryl precursor is activated by the presence of one or more electron-withdrawing substituents, (such as nitro, cyano, trifluoromethyl, trichloromethyl, and the like) and/or when X is nitrogen.

Additional amine, hydroxyl, and thiol arylation techniques that may be used to produce the compounds described herein include those described in Burke, A. J. and Marques, C. S. (eds) (2014) *Amine, Phenol, Alcohol, and Thiol Arylation, in Catalytic Arylation Methods: From the Academic Lab to Industrial Processes*, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, the disclosure of which is incorporated herein by reference as it pertains to processes for chemical synthesis.

A Pd-catalyzed Heck reaction can be used to join olefinic linkers to a halogenated aryl precursor, as shown in Scheme 4, below.

25

Scheme 4.




wherein "=" denotes a carbon-carbon double bond, which can subsequently be reduced to yield a saturated alkylene linker using olefin reduction methods known in the art.

For instances in which L is selected from the group consisting of $-C(O)(CR_{8a}R_{8b})_{n-}$, $-C(S)(CR_{8a}R_{8b})_{n-}$, $-NR_{7a}C(O)(CR_{8a}R_{8b})_{n-}$, $-NR_{7a}C(S)(CR_{8a}R_{8b})_{n-}$, $-OC(O)(CR_{8a}R_{8b})_{n-}$, $-OC(S)(CR_{8a}R_{8b})_{n-}$, $-C(O)NR_{7a}(CR_{8a}R_{8b})_{n-}$, $-C(S)NR_{7a}(CR_{8a}R_{8b})_{n-}$, $-C(O)O(CR_{8a}R_{8b})_{n-}$, $-C(S)O(CR_{8a}R_{8b})_{n-}$, $-S(O)_2NR_{7a}(CR_{8a}R_{8b})_{n-}$, $-NR_{7a}S(O)_2(CR_{8a}R_{8b})_{n-}$, $-NR_{7a}C(O)NR_{7b}(CR_{8a}R_{8b})_{n-}$, and $-NR_{7a}C(O)O(CR_{8a}R_{8b})_{n-}$,

- 5 techniques such as acylation, sulfonamidation, and thionation reactions can be employed in order to produce compounds represented by formulas (I) and (II). Exemplary acylation, sulfonamidation, and thionation processes that may be used to synthesize the aryl hydrocarbon receptor antagonists described herein are depicted in Schemes 5-7, below.


10 Scheme 5.



- In Scheme 5, LG denotes a nucleofugal leaving group, such as a halogen (for instance, chlorine or bromine), a sulfonate (for instance, tosylate, brosylate, triflate, mesylate, and the like), and other leaving groups known in the art. In this way, linkers ("L") containing an amide, ester, ketone, urea, carbamate, or the like in which the carbonyl carbon is bound directly to the imidazopyridine or imidazopyrazine ring system can be synthesized.


- Similarly, to linkers ("L") containing a thioketone, thioamide, thioester, and the like can be synthesized by reacting the corresponding ketone, amide, or ester with a thionating reagent, as shown in Scheme 6, below.

Scheme 6.



wherein $C(O)Y$ denotes an amide, ester, ketone, or the like.


- Exemplary thionating reagents are known in the art and include, for instance, Lawesson's reagent, which is described, for example, Jesberger, et al., Synthesis 13:1929-1258 (2003), the disclosure of which is incorporated herein by reference as it pertains to thionation techniques.



When the linker ("L") contains a sulfonamide moiety bound to the imidazopyridine or

imidazopyrazine ring system, for instance, at the sulfur or nitrogen of the sulfonamide functionality, sulfonamidation techniques known in the art can be used to produce the corresponding compound of formula (I) or (II). An exemplary sulfonamidation process is depicted in Scheme 7, below.

5 Scheme 7.



In Scheme 7, LG denotes a nucleofugal leaving group, such as a halogen (for instance, chlorine or bromine), a sulfonate (for instance, tosylate, brosylate, triflate, mesylate, and the like), and other leaving groups known in the art. In this way, linkers ("L") containing a sulfonamide in which the sulfur or nitrogen of the sulfonamide moiety is bound directly to the imidazopyridine or imiadzopyrazine ring system can be synthesized.

Stem Cells

15 In some embodiments, the stem cells of which the population is modified (e.g., expanded) with the compositions and methods described are capable of being expanded upon contacting the aryl hydrocarbon receptor antagonist. In some embodiments, the stem cells are not genetically modified stem cells.

20 In some embodiments, the stem cells are embryonic stem cells or adult stem cells. In some embodiments, the stem cells are totipotent stem cells, pluripotent stem cells, multipotent stem cells, oligopotent stem cells, or unipotent stem cells. In some embodiments, the stem cells are tissue-specific stem cells.

25 In some embodiments, the stem cells are hematopoietic stem cells, intestinal stem cells, osteoblastic stem cells, mesenchymal stem cells (i.e., lung mesenchymal stem cells, bone marrow-derived mesenchymal stromal cells, or bone marrow stromal cells), neural stem cells (i.e., neuronal dopaminergic stem cells or motor-neuronal stem cells), epithelial stem cells (i.e., lung epithelial stem cells, breast epithelial stem cells, vascular epithelial stem cells, or intestinal epithelial stem cells), cardiac myocyte progenitor stem cells, skin stem cells (i.e., epidermal stem cells or follicular stem cells (hair follicle stem cells)), skeletal muscle stem cells, adipose stem cells, liver stem cells, induced pluripotent stem cells, umbilical cord stem cells, amniotic fluid stem cells, limbal stem cells, dental pulp stem cells, placental stem cells, myoblasts, endothelial progenitor cells, exfoliated teeth derived stem cells, or hair

follicle stem cells.

In some embodiments, the stem cells are hematopoietic stem cells.

In some embodiments, the stem cells are primary stem cells. For example, the stem cells are obtained from bone marrow, adipose tissue, or blood. In some embodiments, the the stem cells are
5 cultured stem cells.

In some embodiments, the stem cells are CD34+ cells. In some embodiments, the stem cells are CD90+ cells. In some embodiments, the stem cells are CD45RA- cells. In some embodiments, the stem cells are CD34+CD90+ cells. In some embodiments, the stem cells are CD34+CD45RA- cells. In some
10 embodiments, the stem cells are CD90+CD45RA- cells. In some embodiments, the stem cells are CD34+CD90+CD45RA- cells.

In some embodiments, the hematopoietic stem cells are extracted from the bone marrow, mobilized into the peripheral blood and then collected by apheresis, or isolated from umbilical cord blood units.

In some embodiments, the hematopoietic stem cells are CD34+ hematopoietic stem cells. In
15 some embodiments, the hematopoietic stem cells are CD90+ hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD45RA- hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD34+CD90+ hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD34+CD45RA- hematopoietic stem cells. In some
20 embodiments, the hematopoietic stem cells are CD90+CD45RA- hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD34+CD90+CD45RA- hematopoietic stem cells.

Methods for Expanding Hematopoietic Stem Cells

In another aspect, the disclosure features a method of producing an expanded population of hematopoietic stem cells ex vivo, the method including contacting a population of hematopoietic stem
25 cells with the compound of any one of the above aspects or embodiments in an amount sufficient to produce an expanded population of hematopoietic stem cells.

In another aspect, the disclosure features a method of enriching a population of cells with hematopoietic stem cells ex vivo, the method including contacting a population of hematopoietic stem
30 cells with the compound of any one of the above aspects or embodiments in an amount sufficient to produce a population of cells enriched with hematopoietic stem cells.

In another aspect, the disclosure features a method of maintaining the hematopoietic stem cell functional potential of a population of hematopoietic stem cells ex vivo for two or more days, the method
35 including contacting a first population of hematopoietic stem cells with the compound of any one of the above aspects or embodiments, wherein the first population of hematopoietic stem cells exhibits a hematopoietic stem cell functional potential after two or more days that is greater than that of a control population of hematopoietic stem cells cultured under the same conditions and for the same time as the first population of hematopoietic stem cells but not contacted with the compound.

In one embodiment, said method for expanding hematopoietic stem cells, comprises (a) providing a starting cell population comprising hematopoietic stem cells and (b) culturing said starting cell
40 population ex vivo in the presence of an AHR antagonist agent compound of any one of the above aspects or embodiments.

The starting cell population comprising hematopoietic stem cells will be selected by the person skilled in the art depending on the envisaged use. Various sources of cells comprising hematopoietic stem cells have been described in the art, including bone marrow, peripheral blood, neonatal umbilical cord blood, placenta or other sources such as liver, particularly fetal liver.

- 5 The cell population may first be subjected to enrichment or purification steps, including negative and/or positive selection of cells based on specific cellular markers in order to provide the starting cell population. Methods for isolating said starting cell population based on specific cellular markers may use fluorescent activated cell sorting (FACS) technology also called flow cytometry or solid or insoluble substrate to which is bound antibodies or ligands that interact with specific cell surface markers. For
10 example, cells may be contacted with a solid substrate (e.g., column of beads, flasks, magnetic particles) containing the antibodies and any unbound cells are removed. When a solid substrate comprising magnetic or paramagnetic beads is used, cells bound to the beads can be readily isolated by a magnetic separator.

- In one embodiment, said starting cell population is enriched in a desirable cell marker phenotype
15 (e.g., CD34+, CD133+, CD90+) or based on efflux of dyes such as rhodamine, Hoechst or aldehyde dehydrogenase activity. In one specific embodiment, said starting cell population is enriched in CD34+ cells. Methods for enriching blood cell population in CD34+ cells include kits commercialized by Miltenyi Biotec (CD34+ direct isolation kit, Miltenyi Biotec, Bergisch, Gladbach, Germany) or by Baxter (Isolex 3000).

- 20 In some embodiments, the hematopoietic stem cells are CD34+ hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD90+ hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD45RA- hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD34+CD90+ hematopoietic stem cells. In some
25 embodiments, the hematopoietic stem cells are CD34+CD45RA- hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD90+CD45RA- hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD34+CD90+CD45RA- hematopoietic stem cells.

- In some embodiments, the hematopoietic stem cells are mammalian cells, such as human cells. In some embodiments, the human cells are CD34+ cells, such as CD34+ cells are CD34+, CD34+CD38-, CD34+CD38-CD90+, CD34+CD38-CD90+CD45RA-, CD34+CD38-CD90+CD45RA-CD49F+, or
30 CD34+CD90+CD45RA- cells.

 In some embodiments, the hematopoietic stem cells are obtained from human cord blood, mobilized human peripheral blood, or human bone marrow. The hematopoietic stem cells may, for example, be freshly isolated from the human or may have been previously cryopreserved.

- The amount of cord blood from a single birth is often inadequate to treat an adult or an older
35 child. One advantage of the expansion methods using the compounds of the invention, or an agent capable of down-regulating the activity and/or expression of aryl hydrocarbon receptor and/or a downstream effector of aryl hydrocarbon receptor pathway, is that it enables the production of a sufficient amount of hematopoietic stem cells from only one cord blood unit.

- Accordingly, in one embodiment, the starting cell population is derived from neonatal umbilical
40 cord blood cells which have been enriched in CD34+ cells. In one related embodiment, said starting cell population is derived from one or two umbilical cord blood units.

In another embodiment, the starting cell population is derived from human mobilized peripheral blood cells which have been enriched in CD34+ cells. In one related embodiment, said starting cell population is derived from human mobilized peripheral blood cells isolated from only one patient.

Said starting cell population enriched in CD34+ cells may preferably contain at least about 50% CD34+ cells, in some embodiments, more than about 90% CD34+ cells, and may comprise between 10^5 and 10^9 nucleated cells.

The starting cell population may be used directly for expansion or frozen and stored for use at a later date.

Conditions for culturing the starting cell population for hematopoietic stem cell expansion will vary depending, inter alia, on the starting cell population, the desired final number of cells, and desired final proportion of HSCs.

In one embodiment, the culturing conditions comprises the use of other cytokines and growth factors, generally known in the art for hematopoietic stem cell expansion. Such cytokines and growth factors include without limitation IL-1, IL-3, IL-6, IL-11, G-CSF, GM-CSF, SCF, Flt3-L, thrombopoietin (TPO), erythropoietin, and analogs thereof. As used herein, "analogs" include any structural variants of the cytokines and growth factors having the biological activity as the naturally occurring forms, including without limitation, variants with enhanced or decreased biological activity when compared to the naturally occurring forms or cytokine receptor agonists such as an agonist antibody against the TPO receptor (for example, VB22B sc(Fv)2 as detailed in patent publication WO 2007/145227, and the like). Cytokine and growth factor combinations are chosen to expand HSC and progenitor cells while limiting the production of terminally differentiated cells. In one specific embodiment, one or more cytokines and growth factors are selected from the group consisting of SCF, Flt3-L and TPO. In one specific embodiment, at least TPO is used in a serum-free medium under suitable conditions for HSC expansion. In one related embodiment, a mixture of IL6, SCF, Flt3-L and TPO is used in the method for expanding HSCs in combination with the compound of the present disclosure.

The expansion of HSC may be carried out in a basal medium, which may be supplemented with mixtures of cytokines and growth factors. A basal medium typically comprises amino acids, carbon sources, vitamins, serum proteins (e.g. albumin), inorganic salts, divalent cations, buffers and any other element suitable for use in expansion of HSC. Examples of such basal medium appropriate for a method of expanding HSC include, without limitation, StemSpan® SFEM—Serum-Free Expansion Medium (StemCell Technologies, Vancouver, Canada), StemSpan® H3000—Defined Medium (StemCell Technologies, Vancouver, Canada), CellGro® SCGM (CellGenix, Freiburg Germany), StemPro®-34 SFM (Invitrogen).

In one embodiment, the compound of the present disclosure is administered during the expansion method of said starting cell population under a concentration appropriate for HSC expansion. In one specific embodiment, said compound or AHR modulating agent is administered at a concentration comprised between 1 pM and 100 μ M, for example between 10 pM and 10 μ M, or between 100 pM and 1 μ M.

In one embodiment where starting cell population essentially consists of CD34+ enriched cells from one or two cord blood units, the cells are grown under conditions for HSC expansion from about 3 days to about 90 days, for example between 7 and 28 days and/or until the indicated fold expansion and

the characteristic cell populations are obtained. In one specific embodiment, the cells are grown under conditions for HSC expansion not more than 21 days, 14 days or 7 days.

In one embodiment, the starting cell population is cultured during a time sufficient to reach an absolute number of CD34+ cells of at least 10^5 , 10^6 , 10^7 , 10^8 or 10^9 cells. In another embodiment, said
5 starting cell population is cultured during a time sufficient for a 10 to 50000 fold expansion of CD34+ cells, for example between 100 and 10000 fold expansion, for examples between 50 and 1000 fold expansion.

The cell population obtained after the expansion method may be used without further purification or may be subject to further purification or selection steps.

The cell population may then be washed to remove the compound of the present disclosure
10 and/or any other components of the cell culture and resuspended in an appropriate cell suspension medium for short term use or in a long-term storage medium, for example a medium suitable for cryopreservation.

Cell Population with Expanded Hematopoietic Stem Cells as Obtained by the Expansion Method 15 and Therapeutic Compositions

In another aspect, the disclosure features a composition comprising a population of hematopoietic stem cells, wherein the hematopoietic stem cells or progenitors thereof have been contacted with the compound of any one of the above aspects or embodiments, thereby expanding the hematopoietic stem cells or progenitors thereof.

The invention further provides a cell population with expanded hemapoetic stem cells obtainable
20 or obtained by the expansion method described above. In one embodiment, such cell population is resuspended in a pharmaceutically acceptable medium suitable for administration to a mammalian host, thereby providing a therapeutic composition.

The compound as defined in the present disclosure enables the expansion of HSCs, for example
25 from only one or two cord blood units, to provide a cell population quantitatively and qualitatively appropriate for efficient short and long term engraftment in a human patient in need thereof. In one embodiment, the present disclosure relates to a therapeutic composition comprising a cell population with expanded HSCs derived from not more than one or two cord blood units. In one embodiment, the present disclosure relates to a therapeutic composition containing a total amount of cells of at least about
30 10^5 , at least about 10^6 , at least about 10^7 , at least about 10^8 or at least about 10^9 cells with about 20% to about 100%, for example between about 43% to about 80%, of total cells being CD34+ cells. In certain embodiments, said composition contains between 20-100%, for example between 43-80%, of total cells being CD34+CD90+CD45RA-.

In some embodiments, the hematopoietic stem cells are CD34+ hematopoietic stem cells. In
35 some embodiments, the hematopoietic stem cells are CD90+ hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD45RA- hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD34+CD90+ hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD34+CD45RA- hematopoietic stem cells. In some embodiments, the hematopoietic stem cells are CD90+CD45RA- hematopoietic stem cells. In some
40 embodiments, the hematopoietic stem cells are CD34+CD90+CD45RA- hematopoietic stem cells.

In some embodiments, the hematopoietic stem cells of the therapeutic composition are

mammalian cells, such as human cells. In some embodiments, the human cells are CD34+ cells, such as CD34+ cells are CD34+, CD34+CD38-, CD34+CD38-CD90+, CD34+CD38-CD90+CD45RA-, CD34+CD38-CD90+CD45RA-CD49F+, or CD34+CD90+CD45RA- cells.

In some embodiments, the hematopoietic stem cells of the therapeutic composition are obtained from human cord blood, mobilized human peripheral blood, or human bone marrow. The hematopoietic stem cells may, for example, be freshly isolated from the human or may have been previously cryopreserved.

Methods of Treatment

As described herein, hematopoietic stem cell transplant therapy can be administered to a subject in need of treatment so as to populate or repopulate one or more blood cell types, such as a blood cell lineage that is deficient or defective in a patient suffering from a stem cell disorder. Hematopoietic stem and progenitor cells exhibit multi-potency, and can thus differentiate into multiple different blood lineages including, but not limited to, granulocytes (e.g., promyelocytes, neutrophils, eosinophils, basophils), erythrocytes (e.g., reticulocytes, erythrocytes), thrombocytes (e.g., megakaryoblasts, platelet producing megakaryocytes, platelets), monocytes (e.g., monocytes, macrophages), dendritic cells, microglia, osteoclasts, and lymphocytes (e.g., NK cells, B-cells and T-cells). Hematopoietic stem cells are additionally capable of self-renewal, and can thus give rise to daughter cells that have equivalent potential as the mother cell, and also feature the capacity to be reintroduced into a transplant recipient whereupon they home to the hematopoietic stem cell niche and re-establish productive and sustained hematopoiesis. Thus, hematopoietic stem and progenitor cells represent a useful therapeutic modality for the treatment of a wide array of disorders in which a patient has a deficiency or defect in a cell type of the hematopoietic lineage. The deficiency or defect may be caused, for example, by depletion of a population of endogenous cells of the hematopoietic system due to administration of a chemotherapeutic agent (e.g., in the case of a patient suffering from a cancer, such as a hematologic cancer described herein). The deficiency or defect may be caused, for example, by depletion of a population of endogenous hematopoietic cells due to the activity of self-reactive immune cells, such as T lymphocytes or B lymphocytes that cross-react with self antigens (e.g., in the case of a patient suffering from an autoimmune disorder, such as an autoimmune disorder described herein). Additionally or alternatively, the deficiency or defect in cellular activity may be caused by aberrant expression of an enzyme (e.g., in the case of a patient suffering from various metabolic disorders, such as a metabolic disorder described herein).

Thus, hematopoietic stem cells can be administered to a patient defective or deficient in one or more cell types of the hematopoietic lineage in order to re-constitute the defective or deficient population of cells in vivo, thereby treating the pathology associated with the defect or depletion in the endogenous blood cell population. Hematopoietic stem and progenitor cells can be used to treat, e.g., a non-malignant hemoglobinopathy (e.g., a hemoglobinopathy selected from the group consisting of sickle cell anemia, thalassemia, Fanconi anemia, aplastic anemia, and Wiskott-Aldrich syndrome). In these cases, for example, a population of hematopoietic stem cells may be expanded ex vivo by culturing the cells in the presence of an aryl hydrocarbon receptor antagonist described herein. The hematopoietic stem cells

thus expanded may then be administered to a patient, where the cells may home to a hematopoietic stem cell niche and re-constitute a population of cells that are damaged or deficient in the patient.

Hematopoietic stem or progenitor cells mobilized to the peripheral blood of a subject may be withdrawn (e.g., harvested or collected) from the subject by any suitable technique. For example, the
5 hematopoietic stem or progenitor cells may be withdrawn by a blood draw. In some embodiments, hematopoietic stem or progenitor cells mobilized to a subject's peripheral blood as contemplated herein may be harvested (i.e., collected) using apheresis. In some embodiments, apheresis may be used to enrich a donor's blood with mobilized hematopoietic stem or progenitor cells.

Additionally or alternatively, hematopoietic stem and progenitor cells can be used to treat an
10 immunodeficiency, such as a congenital immunodeficiency. Additionally or alternatively, the compositions and methods described herein can be used to treat an acquired immunodeficiency (e.g., an acquired immunodeficiency selected from the group consisting of HIV and AIDS). In these cases, for example, a population of hematopoietic stem cells may be expanded ex vivo by culturing the cells in the presence of an aryl hydrocarbon receptor antagonist described herein. The hematopoietic stem cells thus expanded
15 may then be administered to a patient, where the cells may home to a hematopoietic stem cell niche and re-constitute a population of immune cells (e.g., T lymphocytes, B lymphocytes, NK cells, or other immune cells) that are damaged or deficient in the patient.

Hematopoietic stem and progenitor cells can also be used to treat a metabolic disorder (e.g., a metabolic disorder selected from the group consisting of glycogen storage diseases,
20 mucopolysaccharidoses, Gaucher's Disease, Hurlers Disease, sphingolipidoses, Sly Syndrome, alpha-Mannosidosis, X-ALD, Aspartylglucosaminuria, Wolman Disease, late infantile metachromatic leukodystrophy, Niemann Pick Type C disease, Niemann Pick Type B disease, Juvenile Tay Sachs, Infantile Tay Sachs, Juvenile Sandhoff, Infantile Sandhoff, GM1 gangliosidosis, MPSIV (Morquio), Presymptomatic or milder forms of globoid cell leukodystrophy, infantile Krabbe when newborn and asymptomatic, early diagnosis fucosidosis, Fabry, MPSIS, MPSIH/S, MPSII, MPSVI in conjunction with
25 ERT or where alloantibodies attenuate efficacy of ERT, Pompe where alloantibodies attenuate efficacy of ERT, Mucopolidosis II, and metachromatic leukodystrophy). In these cases, for example, a population of hematopoietic stem cells may be expanded ex vivo by culturing the cells in the presence of an aryl hydrocarbon receptor antagonist described herein. The hematopoietic stem cells thus expanded may
30 then be administered to a patient, where the cells may home to a hematopoietic stem cell niche and re-constitute a population of hematopoietic cells that are damaged or deficient in the patient.

Additionally or alternatively, hematopoietic stem or progenitor cells can be used to treat a malignancy or proliferative disorder, such as a hematologic cancer or myeloproliferative disease. In the case of cancer treatment, for example, a population of hematopoietic stem cells may be expanded ex vivo
35 by culturing the cells in the presence of an aryl hydrocarbon receptor antagonist described herein. The hematopoietic stem cells thus expanded may then be administered to a patient, where the cells may home to a hematopoietic stem cell niche and re-constitute a population of cells that are damaged or deficient in the patient, such as a population of hematopoietic cells that is damaged or deficient due to the administration of one or more chemotherapeutic agents to the patient. In some embodiments,
40 hematopoietic stem or progenitor cells may be infused into a patient in order to repopulate a population of cells depleted during cancer cell eradication, such as during systemic chemotherapy. Exemplary

hematological cancers that can be treated by way of administration of hematopoietic stem and progenitor cells in accordance with the compositions and methods described herein are acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia, chronic lymphoid leukemia, multiple myeloma, diffuse large B-cell lymphoma, and non-Hodgkin's lymphoma, as well as other cancerous conditions, including neuroblastoma.

Additional diseases that can be treated by the administration of hematopoietic stem and progenitor cells to a patient include, without limitation, adenosine deaminase deficiency and severe combined immunodeficiency, hyper immunoglobulin M syndrome, Chediak-Higashi disease, hereditary lymphohistiocytosis, osteopetrosis, osteogenesis imperfecta, storage diseases, thalassemia major, systemic sclerosis, systemic lupus erythematosus, multiple sclerosis, and juvenile rheumatoid arthritis.

In addition, administration of hematopoietic stem and progenitor cells can be used to treat autoimmune disorders. In some embodiments, upon infusion into a patient, transplanted hematopoietic stem and progenitor cells may home to a stem cell niche, such as the bone marrow, and establish productive hematopoiesis. This, in turn, can re-constitute a population of cells depleted during autoimmune cell eradication, which may occur due to the activity of self-reactive lymphocytes (e.g., self-reactive T lymphocytes and/or self-reactive B lymphocytes). Autoimmune diseases that can be treated by way of administering hematopoietic stem and progenitor cells to a patient include, without limitation, psoriasis, psoriatic arthritis, Type 1 diabetes mellitus (Type 1 diabetes), rheumatoid arthritis (RA), human systemic lupus (SLE), multiple sclerosis (MS), inflammatory bowel disease (IBD), lymphocytic colitis, acute disseminated encephalomyelitis (ADEM), Addison's disease, alopecia universalis, ankylosing spondylitis, antiphospholipid antibody syndrome (APS), aplastic anemia, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune inner ear disease (AIED), autoimmune lymphoproliferative syndrome (ALPS), autoimmune oophoritis, Balo disease, Behcet's disease, bullous pemphigoid, cardiomyopathy, Chagas' disease, chronic fatigue immune dysfunction syndrome (CFIDS), chronic inflammatory demyelinating polyneuropathy, Crohn's disease, cicatricial pemphigoid, coeliac sprue-dermatitis herpetiformis, cold agglutinin disease, CREST syndrome, Degos disease, discoid lupus, dysautonomia, endometriosis, essential mixed cryoglobulinemia, fibromyalgia-fibromyositis, Goodpasture's syndrome, Grave's disease, Guillain-Barre syndrome (GBS), Hashimoto's thyroiditis, Hidradenitis suppurativa, idiopathic and/or acute thrombocytopenic purpura, idiopathic pulmonary fibrosis, IgA neuropathy, interstitial cystitis, juvenile arthritis, Kawasaki's disease, lichen planus, Lyme disease, Meniere disease, mixed connective tissue disease (MCTD), myasthenia gravis, neuromyotonia, opsoclonus myoclonus syndrome (OMS), optic neuritis, Ord's thyroiditis, pemphigus vulgaris, pernicious anemia, polychondritis, polymyositis and dermatomyositis, primary biliary cirrhosis, polyarteritis nodosa, polyglandular syndromes, polymyalgia rheumatica, primary agammaglobulinemia, Raynaud phenomenon, Reiter's syndrome, rheumatic fever, sarcoidosis, scleroderma, Sjögren's syndrome, stiff person syndrome, Takayasu's arteritis, temporal arteritis (also known as "giant cell arteritis"), ulcerative colitis, collagenous colitis, uveitis, vasculitis, vitiligo, vulvodynia ("vulvar vestibulitis"), and Wegener's granulomatosis.

Hematopoietic stem cell transplant therapy may additionally be used to treat neurological disorders, such as Parkinson's disease, Alzheimer's disease, multiple sclerosis, Amyotrophic lateral sclerosis, Huntington's disease, mild cognitive impairment, amyloidosis, AIDS-related dementia, encephalitis, stroke, head trauma, epilepsy, mood disorders, and dementia. As described herein, upon

transplantation into a patient, hematopoietic stem cells may migrate to the central nervous system and differentiate into, for example, microglial cells, thereby re-constituting a population of cells that may be damaged or deficient in a patient suffering from a neurological disorder. In these cases, for example, a population of hematopoietic stem cells may be expanded ex vivo by culturing the cells in the presence of
5 an aryl hydrocarbon receptor antagonist described herein. The hematopoietic stem cells thus expanded may then be administered to a patient suffering from a neurological disorder, where the cells may home to the central nervous system, such as the brain of the patient, and re-constitute a population of hematopoietic cells (e.g., microglial cells) that are damaged or deficient in the patient.

As described herein, hematopoietic stem cell transplant therapy can be administered to a subject
10 in need of treatment so as to populate or repopulate one or more blood cell types, such as a blood cell lineage that is deficient or defective in a patient suffering from a stem cell disorder. Hematopoietic stem and progenitor cells exhibit multi-potency, and can thus differentiate into multiple different blood lineages including, in one embodiment, microglia.

The methods disclosed herein for treating disorders in a subject in need thereof comprise the
15 administration of an expanded population of hematopoietic stem cells to a subject in need thereof. In one embodiment, the number of expanded hematopoietic stem cells administered to the subject is equal to or greater than the amount of hematopoietic stem cells needed to achieve a therapeutic benefit. In one embodiment, the number of expanded hematopoietic stem cells administered to the subject is greater than the amount of hematopoietic stem cells needed to achieve a therapeutic benefit. In one
20 embodiment, the therapeutic benefit achieved is proportional to the number of expanded hematopoietic stem cells that are administered.

A dose of the expanded hematopoietic stem cell composition of the disclosure is deemed to have achieved a therapeutic benefit if it alleviates a sign or a symptom of the disease. The sign or symptom of the disease may comprise one or more biomarkers associated with the disease, or one or more clinical
25 symptoms of the disease.

For example, administration of the expanded hematopoietic stem cell composition may result in the reduction of a biomarker that is elevated in individuals suffering from the disease, or elevate the level of a biomarker that is reduced in individuals suffering from the disease.

For example, administering the expanded hematopoietic stem cell composition of the disclosure
30 may elevate the level of an enzyme that is reduced in an individual suffering from a metabolic disorder. This change in biomarker level may be partial, or the level of the biomarker may return to levels normally seen in healthy individuals.

Selection of donors and patients

35 In some embodiments, the patient is the donor. In such cases, withdrawn hematopoietic stem or progenitor cells may be re-infused into the patient, such that the cells may subsequently home hematopoietic tissue and establish productive hematopoiesis, thereby populating or repopulating a line of cells that is defective or deficient in the patient (e.g., a population of megakaryocytes, thrombocytes, platelets, erythrocytes, mast cells, myeloblasts, basophils, neutrophils, eosinophils, microglia,
40 granulocytes, monocytes, osteoclasts, antigen-presenting cells, macrophages, dendritic cells, natural killer cells, T-lymphocytes, and B-lymphocytes). In this scenario, the transplanted hematopoietic stem or

progenitor cells are least likely to undergo graft rejection, as the infused cells are derived from the patient and express the same HLA class I and class II antigens as expressed by the patient.

Alternatively, the patient and the donor may be distinct. In some embodiments, the patient and the donor are related, and may, for example, be HLA-matched. As described herein, HLA-matched donor-recipient pairs have a decreased risk of graft rejection, as endogenous T cells and NK cells within the transplant recipient are less likely to recognize the incoming hematopoietic stem or progenitor cell graft as foreign, and are thus less likely to mount an immune response against the transplant. Exemplary HLA-matched donor-recipient pairs are donors and recipients that are genetically related, such as familial donor-recipient pairs (e.g., sibling donor-recipient pairs).

In some embodiments, the patient and the donor are HLA-mismatched, which occurs when at least one HLA antigen, in particular with respect to HLA-A, HLA-B and HLA-DR, is mismatched between the donor and recipient. To reduce the likelihood of graft rejection, for example, one haplotype may be matched between the donor and recipient, and the other may be mismatched.

Administration and Dosing of Hematopoietic Stem or Progenitor Cells

Hematopoietic stem and progenitor cells described herein may be administered to a subject, such as a mammalian subject (e.g., a human subject) suffering from a disease, condition, or disorder described herein, by one or more routes of administration. For instance, hematopoietic stem cells described herein may be administered to a subject by intravenous infusion. Hematopoietic stem cells may be administered at any suitable dosage. Non-limiting examples of dosages include about 1×10^5 CD34+ cells/kg of recipient to about 1×10^8 CD34+ cells/kg (e.g., from about 2×10^5 CD34+ cells/kg to about 9×10^7 CD34+ cells/kg, from about 3×10^5 CD34+ cells/kg to about 8×10^7 CD34+ cells/kg, from about 4×10^5 CD34+ cells/kg to about 7×10^7 CD34+ cells/kg, from about 5×10^5 CD34+ cells/kg to about 6×10^7 CD34+ cells/kg, from about 5×10^5 CD34+ cells/kg to about 1×10^8 CD34+ cells/kg, from about 6×10^5 CD34+ cells/kg to about 1×10^8 CD34+ cells/kg, from about 7×10^5 CD34+ cells/kg to about 1×10^8 CD34+ cells/kg, from about 8×10^5 CD34+ cells/kg to about 1×10^8 CD34+ cells/kg, from about 9×10^5 CD34+ cells/kg to about 1×10^8 CD34+ cells/kg, from about 1×10^7 CD34+ cells/kg to about 1×10^8 CD34+ cells/kg, or from about 1×10^8 CD34+ cells/kg to about 1×10^7 CD34+ cells/kg, among others).

Hematopoietic stem or progenitor cells and pharmaceutical compositions described herein may be administered to a subject in one or more doses. When multiple doses are administered, subsequent doses may be provided one or more days, weeks, months, or years following the initial dose.


Examples

The following examples are put forth so as to provide those of ordinary skill in the art with a description of how the compositions and methods described herein may be used, made, and evaluated, and are intended to be purely exemplary and are not intended to limit the scope of what the inventors regard as their invention.

Example 1. Synthesis of Compound (5)

Compound (5) can be synthesized, for example, using a Hartwig-Buchwald amination process that includes coupling a halogenated imidazopyridine precursor to a protected 2-aminoethyl indole, as shown in Scheme 8, below.


5 Scheme 8.



Deprotection of the ensuing adduct, for instance, using conventional deprotection methods known in the art, can yield compound (5).

Alternatively, the tandem amination-hydrolysis procedure outline in Scheme 9, below, may be used to synthesize compound (5).


Scheme 9.



15 Example 2. Synthesis of Compound (16)

In a manner similar to that described in Example 1, compound (16) can be synthesized by way of a Hartwig-Buchwald process, followed by a Suzuki coupling and indole deprotection, as shown in Scheme 10, below.

5 Scheme 10.



Example 3. Capacity of Compounds (5) and (16) to expand hematopoietic stem cells

10 To determine the ability of compounds (5) and (16) to inhibit the activity of the aryl hydrocarbon receptor and to induce the proliferation of hematopoietic stem cells, a series of HSC expansion experiments were conducted. In the first experiment, compounds (5) and (16) were assessed for their capacity to attenuate aryl hydrocarbon receptor signaling. To this end, HepG2 hepatocytes were transiently transfected with a luciferase reporter construct under the control of a promoter responsive to
 15 aryl hydrocarbon receptor signal transduction. The cells were plated at a density of 25,000 cells per well in a microtiter plate. The HepG2 cells were immediately treated with compound (5) or (16) in the absence (FIG. 1) or presence (FIG. 2) of the aryl hydrocarbon receptor agonist, VAF347 (80 nM). Luciferase activity was subsequently analyzed six hours after plating.

As shown in FIGS. 1 and 2, compounds (5) and (16) were capable of suppressing aryl
 20 hydrocarbon receptor activity even in the presence of the activator VAF347.

To assess the ability of compounds (5) and (16) to induce the proliferation of hematopoietic stem cells, a population of mononuclear peripheral blood cells enriched in CD34+ cells were plated at a density of 2,350 cells per well (50 μ L) in a microtiter plate in the presence of each compound. The percentage of CD34+ hematopoietic stem cells was assessed seven days following the initial plating. The results of this
 25 experiment are reported in FIG. 3. As shown therein, compounds (5) and (16) were capable of potentiating hematopoietic stem cell growth in a dose-dependent manner. The compounds of formula (I) and (II) described herein can thus be used to expand hematopoietic stem cells ex vivo in order to obtain sufficient quantities of such cells for in vivo applications.

Surprisingly, compounds (5) and (16) were capable of promoting hematopoietic stem cell
 30 expansion with a potency greater than that reported for StemRegenin1 (SR1), which is described, for example, in US Patent No. 8,927,281, which is incorporated herein by reference. This difference in biological activity is expected to have a significant clinical benefit, as a reduced quantity of aryl hydrocarbon receptor antagonists according to formulas (I) and (II) described herein relative to SR1 may be used to prepare an amplified population of hematopoietic stem cells suitable for transplantation to a
 35 patient in need thereof (for instance, as described in Example 4, below).

Example 4. Administration of hematopoietic stem cells to a human patient in need thereof

Using the methods disclosed herein, a population of hematopoietic stem cells that have been expanded ex vivo using the aryl hydrocarbon receptor antagonists of formula (I) or (II) can be administered to a human patient in need of hematopoietic stem cell transplant therapy. Prior to the transplantation, a population of hematopoietic stem cells may be cultured in the presence of the aryl hydrocarbon receptor antagonist for one or more days (e.g., for one, two, three, four, five, six, seven, eight, nine, ten, or more days, replenishing culture medium as needed). The hematopoietic stem cell population may be expanded to 1×10^6 to 1×10^{12} hematopoietic stem cells prior to infusion into the patient in need of transplant therapy.

Following the conclusion of the expansion process, the patient may receive an infusion (e.g., an intravenous infusion) of the expanded, exogenous hematopoietic stem cells, such as from a practitioner that performed the ex vivo expansion or from a different physician. The patient may then be administered an infusion of autologous, syngeneic, or allogeneic hematopoietic stem cells, for instance, at a dosage of from 1×10^3 to 1×10^9 hematopoietic stem cells/kg. The engraftment of the hematopoietic stem cell transplant may be monitored, for example, by detecting an increase in concentration of hematopoietic stem cells or cells of the hematopoietic lineage (such as megakaryocytes, thrombocytes, platelets, erythrocytes, mast cells, myoblasts, basophils, neutrophils, eosinophils, microglia, granulocytes, monocytes, osteoclasts, antigen-presenting cells, macrophages, dendritic cells, natural killer cells, T-lymphocytes, and B-lymphocytes) in a blood sample isolated from the patient following administration of the transplant. This analysis may be conducted, for example, from 1 hour to 6 months, or more, following hematopoietic stem cell transplant therapy (e.g., 1 hour, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, 7 hours, 8 hours, 9 hours, 10 hours, 11 hours, 12 hours, 13 hours, 14 hours, 15 hours, 16 hours, 17 hours, 18 hours, 19 hours, 20 hours, 21 hours, 22 hours, 23 hours, 24 hours, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 17 weeks, 18 weeks, 19 weeks, 20 weeks, 21 weeks, 22 weeks, 23 weeks, 24 weeks, or more). A finding that the concentration of hematopoietic stem cells or cells of the hematopoietic lineage has increased (e.g., by 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 200%, 500%, or more) following the transplant therapy relative to the concentration of the corresponding cell type prior to transplant therapy provides one indication that the transplantation therapy is successful.

Example 5. Engraftment of microglial cells in the central nervous system following hematopoietic stem cell transplant

To investigate the ability of hematopoietic stem cells to differentiate into microglial cells and subsequently engraft in central nervous system tissue, such as the brain of a hematopoietic stem cell transplant recipient, a series of experiments were conducted in which human hematopoietic stem cells were first expanded ex vivo in the presence of an aryl hydrocarbon receptor antagonist (compound (16) or compound (24)) and were subsequently transplanted into NSG mice, in accordance with the scheme shown in FIG. 4. The frequency of human CD45+ cells in the peripheral blood of the mice was then

determined, as well as the profile of microglial cells in the brain tissue using flow cytometry and immunohistochemistry techniques.

As shown in FIGS. 5A and 5B, upon transplantation of hematopoietic stem cells expanded ex vivo in the presence of compound (16), NSG mice exhibited an increased frequency of hCD45+ cells in peripheral blood, as well as an increased engraftment of hCD45+CD11b+ microglial cells in the brain. Similar results were obtained upon transplantation of hematopoietic stem cells expanded ex vivo in the presence of compound (24), as shown in FIGS. 6A and 6B.

Collectively, these data demonstrate the ability of hematopoietic stem cells expanded in the presence of aryl hydrocarbon receptor antagonists described herein to promote the engraftment of microglial cells in central nervous system tissue of a hematopoietic stem cells transplant recipient. These findings provide further evidence that hematopoietic stem cell transplantation can be used to treat a wide array of neurological disorders, including Parkinson's disease, Alzheimer's disease, multiple sclerosis, Amyotrophic lateral sclerosis, Huntington's disease, mild cognitive impairment, amyloidosis, AIDS-related dementia, encephalitis, stroke, head trauma, epilepsy, mood disorders, and dementia, among others.

Materials and Methods

Cord Blood Expansion and Transplantation

Approximately 60,000 cord blood CD34+ cells were seeded in T25 flasks at a final volume of 12 mL in HSC growth media (SFEM supplemented with Pen/Strep, 50 ng/mL FLT3L, TPO, SCF, and IL-6). Flasks were incubated for 10 days at 37°C/5% CO₂. Cells were cultured in the presence of 500 nM of AHR antagonist, where indicated. Cells were transferred to a larger flask when needed to maintain cells at a density less than 1x10⁶ cells/mL throughout the culture period.

At the time of thaw, an equal number of cells to the starting cell cultures were injected into NSG mice, sublethally irradiated (200 cGy) 24 hours prior to injection. After 10 days of culture, the entire progeny of the cultures was injected into NSG mice. Peripheral blood was harvested by retro-orbital bleeding at approximately weeks 4 and 8 or by cardiac puncture at week 12 and chimerism was assessed by flow cytometry using antibodies against hCD45, mCD45, hCD33, hCD19, hCD3 and a viability dye.

Brain Harvesting and Processing

At 3 months, brains were harvested. 1 hemisphere was fixed in formalin, embedded, and used for immunohistochemistry. The other hemisphere was crushed in Dounce buffer (15 mM HEPES/0.5% glucose in phenol red-free HBSS) and filtered through a 40 µm filter to create a single cell suspension and resuspended in 900 µL 0.5% BSA/PBS. Myelin was depleted from brain samples, per manufacturer's instructions, by incubating with 100 µL myelin removal beads (Miltenyi Biotec), incubating for 15 minutes at 4°C, washing with PBS, and resuspending in 1 mL MACS Buffer prior to deletion on an AutoMACs Pro.

Flow Cytometric Detection of Microglia

Myelin-depleted samples were resuspended in 100 µL PBS and stained with antibodies against hCD45, mCD45, CD11b, CD19, CD3, and 7-AAD viability dye. Cells were washed once in PBS and resuspended in 300 µL final volume. The entire sample was acquired by flow cytometry (BD Celesta) to quantitate the number of microglia per brain hemisphere.

Immunohistochemical Detection of Microglia

Embedded brains were sectioned at approximately 5 microns and stained with Ku80 (brown) and Iba-1 (red) primary antibodies). Mouse brains were analyzed from each transplanted mouse and five levels were analyzed each. Glass slides were scanned at 20X using an Aperio AT2 whole slide scanner.

- 5 Image analysis was performed on the digital slide images using Visiopharm software.

Example 6. Synthesis of Compound (24)


- In a manner similar to that described in Example 1 and Example 2, compound (24) can be synthesized by way of an arene iodination and Hartwig-Buchwald process, followed by a first Suzuki
- 10 coupling, a second Suzuki coupling and alkene reduction by catalytic hydrogenation over palladium as shown in Scheme 11, below.

- 6-(5-fluoropyridin-3-yl)-N-[2-(1H-indol-3-yl)ethyl]-3-(3-methoxy-1-methylpropyl)imidazo[1,2-a]pyrazin-8-amine, Compound (24) was isolated as a white solid. FIG. 7 sets forth a ¹H-NMR spectrum (d₆-DMSO) consistent with the structure. HPLC analysis gave 97.93% Area at 254 nm and 97.65% Area
- 15 at 210 nm; Retention time: 3.765 min; HPLC conditions: Agilent 1100 HPLC. Zorbax Eclipse XDBC18 50 x 4.6 mm 1.8 micron column. Solvent A: Water (0.1% TFA); Solvent B: Acetonitrile (0.07% TFA). Gradient: 95% A to 95% B over 5 min; hold for 1 min; recycle over 1 min; 30 s hold. UV Detection: 210 and 254 nm with no reference. Column temperature: 30 °C. Mass spectrum was consistent with structure MS (ESI+) for C₂₆H₂₇FN₆O *m/z* 459.1 (M+H)⁺; MS (ESI-) for C₂₆H₂₇FN₆O *m/z* 457.2 (M-H)⁻.

- 20 Preparative HPLC AD-H (2 x 25 cm) 25% methanol/CO₂, 100 bar, 65 mL/min, 220 nm, inj vol.: 0.5 mL, 20 mg/mL methanol:DCM and analytical HPLC AD-H (25 x 0.46 cm) 35% methanol/CO₂, 100 bar 3 mL/min, 220, 254 and 280 nm allowed for the resolution of distinct enatiomeric peaks for Compound (24). FIG. 8 sets forth a ¹H-NMR spectrum (d₆-DMSO) consistent with the structure isolated as peak 1. FIG. 9 sets forth a ¹H-NMR spectrum (d₆-DMSO) consistent with the structure isolated as peak 2.

25


Scheme 11.

**Example 7. Synthesis of Compound (27)**

In a manner similar to that described in Example 6, compound (27) can be synthesized by way of a first Suzuki coupling, and second Suzuki coupling and alkene reduction by catalytic hydrogenation over palladium as shown in Scheme 12, below.

- 6-(5-fluoropyridin-3-yl)-N-[2-(1H-indol-3-yl)ethyl]-3-(4-methoxy-1-methylbutyl)imidazo[1,2-a]pyrazin-8-amine, Compound (27) was isolated as a white solid. FIG. 10 sets forth a ¹H-NMR spectrum (CDCl₃) consistent with the structure. HPLC analysis gave 98.0 %Area at 254 nm and 97.2 %Area at 210 nm; Retention time: 3.670 min; HPLC conditions: Agilent 1100 HPLC. Zorbax Eclipse XDBC18; 50 x 4.6 mm 1.8 micron column. Solvent A: Water (0.1% TFA); Solvent B: Acetonitrile (0.07% TFA). Gradient: 95% A to 95% B over 5 min; hold for 1 min; recycle over 1 min; 30 s hold. UV Detection: 210 and 254 nm with no reference. Column temperature: 30 °C. Mass Spectrum was consistent with structure MS (ESI+) for C₂₇H₂₉FN₆O *m/z* 473.1 (M+H)⁺; MS (ESI-) for C₂₇H₂₉FN₆O *m/z* 471.2 (M-H)⁻.

Scheme 12.



15

Example 8. Synthesis of Compound (28)


In a manner similar to that described in Example 6 and Example 7, compound (28) can be synthesized by way of a first Suzuki coupling, and second Suzuki coupling and alkene reduction by catalytic hydrogenation over palladium as shown in Scheme 13, below.

- 3-(3-ethoxy-1-methylpropyl)-6-(5-fluoropyridin-3-yl)-N-[2-(1H-indol-3-yl)ethyl]imidazo[1,2-a]pyrazin-8-amine, Compound (28) was isolated as an off-white solid. FIG. 11 sets forth a ¹H-NMR spectrum (CDCl₃) consistent with the structure. HPLC analysis gave 99.3 %Area at 254 nm and 99.0 %Area at 210 nm; Retention time: 3.757 min; HPLC conditions: Agilent 1100 HPLC. Zorbax Eclipse XDBC18; 50 x 4.6 mm 1.8 micron column. Solvent A: Water (0.1% TFA); Solvent B: Acetonitrile (0.07% TFA). Gradient: 95% A to 95% B over 5 min; hold for 1 min; recycle over 1 min; 30 s hold. UV Detection:

210 and 254 nm with no reference. Column temperature: 30 °C. Mass spectrum was consistent with structure MS (ESI+) for $C_{27}H_{29}FN_6O$ m/z 473.1 (M+H)+; MS (ESI-) for $C_{27}H_{29}FN_6O$ m/z 471.2 (M-H)-.

5


Scheme 13.



Suitable substrates may be provided for the first Suzuki coupling in the synthesis of compound (24), compound (27), and compound (28) in Example 6, Example 7, and Example 8, respectively, by alkylation of a suitable alkynyl alcohol and borylation of the alkyne as shown in Scheme 14, below.

10

Scheme 14.



15

Example 9. Evaluation of AHR Antagonists

To determine the activity of a scale-up batch of Compound (24), individual enantiomers, and two derivatives Compound (27) and Compound (28) were evaluated for i) the ability to expand CD34+

hematopoietic stem/progenitor cells and ii) AHR antagonist activity. The compounds were compared to control AHR antagonist SR1, an additional control AHR antagonist, and stored sample of Compound (24) CD34+ Frequency and Number

- 5 *Methods* – mobilized peripheral blood CD34+ cells were cultured in the presence of compound, in dose response beginning at 10 μ M. CD34+ cell frequency and number was assessed at day 7 by flow cytometry. All synthesized compounds show equivalent expansion capabilities if not greater than SR1. Table 1. CD34+ Frequency and Number

Compound	EC50 (μ M)
SR1	0.112
Control AHR Antagonist	0.001
Compound (24) – storage (racemic)	0.007
Compound (24) – scale-up (racemic)	0.007
Compound (24) – Enantiomer Peak 1	0.011
Compound (24) – Enantiomer Peak 2	0.004
Compound (28)	0.010
Compound (27)	0.008

AHR Antagonist Assay

- 10 *Methods* – HepG2 cells transiently transfected with the pGudLuc6.1 plasmid were thawed and 25,000 cells were plated per well and immediately treated the AHR agonist VAF347 (fixed at 40 nM) and/or the indicated AHR antagonists. Luciferase activity was measured 24 hours post-culture, corresponding to endogenous AHR antagonist activity (without VAF347, FIG. 14) or in the presence of VAF347 (FIG. 15). All synthesized compounds show equivalent AHR antagonist activity if not greater than SR1.
- 15

Table 2. Endogenous AHR Antagonist Activity

Compound	EC50 (μ M)
SR1	0.033
Control AHR Antagonist	0.0009
Compound (24) – storage (racemic)	0.001
Compound (24) – scale-up (racemic)	ND

Compound (24) – Enantiomer Peak 1	0.001
Compound (24) – Enantiomer Peak 2	0.0004
Compound (28)	0.002
Compound (27)	0.002

Table 3. AHR Antagonist Activity in the Presence of VAF347

Compound	EC50 (μM)
SR1	0.193
Control AHR Antagonist	0.014
Compound (24) – storage (racemic)	0.013
Compound (24) – scale-up (racemic)	0.011
Compound (24) – Enantiomer Peak 1	0.012
Compound (24) – Enantiomer Peak 2	0.006
Compound (28)	0.030
Compound (27)	0.009

Other Embodiments


All publications, patents, and patent applications mentioned in this specification are incorporated herein by reference to the same extent as if each independent publication or patent application was specifically and individually indicated to be incorporated by reference.

- 5 While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the invention that come within known or customary practice within the art to which the invention pertains and may be applied to the essential features hereinbefore set forth, and
- 10 follows in the scope of the claims.

Other embodiments are within the claims.

CLAIMS

1. A compound represented by formula (I)



wherein L is selected from the group consisting of $-\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{O}(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{C}(\text{O})(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{C}(\text{S})(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{S}(\text{O})_{0-2}(\text{CR}_{8a}\text{R}_{8b})_n-$, $-(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{NR}_{7a}\text{C}(\text{O})(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{NR}_{7a}\text{C}(\text{S})(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{OC}(\text{O})(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{OC}(\text{S})(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{C}(\text{O})\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{C}(\text{S})\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{C}(\text{O})\text{O}(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{C}(\text{S})\text{O}(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{S}(\text{O})_2\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{NR}_{7a}\text{S}(\text{O})_2(\text{CR}_{8a}\text{R}_{8b})_n-$, $-\text{NR}_{7a}\text{C}(\text{O})\text{NR}_{7b}(\text{CR}_{8a}\text{R}_{8b})_n-$, and $-\text{NR}_{7a}\text{C}(\text{O})\text{O}(\text{CR}_{8a}\text{R}_{8b})_n-$, wherein R_{7a} , R_{7b} , R_{8a} , and R_{8b} are each independently selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl, and each n is independently an integer from 2 to 6;

R_1 is selected from the group consisting of $-\text{S}(\text{O})_2\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{S})\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{NR}_{9b}\text{R}_{9c}$, $-\text{C}(\text{O})\text{R}_{9a}$, $-\text{C}(\text{S})\text{R}_{9a}$, $-\text{S}(\text{O})_{0-2}\text{R}_{9a}$, $-\text{C}(\text{O})\text{OR}_{9a}$, $-\text{C}(\text{S})\text{OR}_{9a}$, $-\text{C}(\text{O})\text{NR}_{9a}\text{R}_{9b}$, $-\text{C}(\text{S})\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{S}(\text{O})_2\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{OR}_{9b}$, $-\text{OC}(\text{O})\text{CR}_{9a}\text{R}_{9b}\text{R}_{9c}$, $-\text{OC}(\text{S})\text{CR}_{9a}\text{R}_{9b}\text{R}_{9c}$, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein R_{9a} , R_{9b} , and R_{9c} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

R_2 is selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl;

R_3 is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

R_4 is selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl;

R_5 is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R_6 is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

2. The compound of claim 1, wherein R_1 is selected from the group consisting of - $S(O)_2NR_{9a}R_{9b}$, - $NR_{9a}C(O)R_{9b}$, - $NR_{9a}C(S)R_{9b}$, - $NR_{9a}C(O)NR_{9b}R_{9c}$, - $C(O)R_{9a}$, - $C(S)R_{9a}$, - $S(O)_2R_{9a}$, - $C(O)OR_{9a}$, - $C(S)OR_{9a}$, - $C(O)NR_{9a}R_{9b}$, - $C(S)NR_{9a}R_{9b}$, - $NR_{9a}S(O)_2R_{9b}$, - $NR_{9a}C(O)OR_{9b}$, - $OC(O)CR_{9a}R_{9b}R_{9c}$, - $OC(S)CR_{9a}R_{9b}R_{9c}$, phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein said phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, - $O(CH_2)_2NR_{10a}R_{10b}$, - $S(O)_2NR_{10a}R_{10b}$, - $OS(O)_2NR_{10a}R_{10b}$, and - $NR_{10a}S(O)_2R_{10b}$, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl.

3. The compound of claim 2, wherein R_1 is selected from the group consisting of - $S(O)_2NR_{9a}R_{9b}$, - $NR_{9a}C(O)R_{9b}$, - $NR_{9a}C(S)R_{9b}$, - $NR_{9a}C(O)NR_{9b}R_{9c}$, - $C(O)R_{9a}$, - $C(S)R_{9a}$, - $S(O)_2R_{9a}$, - $C(O)OR_{9a}$, - $C(S)OR_{9a}$, - $C(O)NR_{9a}R_{9b}$, - $C(S)NR_{9a}R_{9b}$, - $NR_{9a}S(O)_2R_{9b}$, - $NR_{9a}C(O)OR_{9b}$, - $OC(O)CR_{9a}R_{9b}R_{9c}$, and - $OC(S)CR_{9a}R_{9b}R_{9c}$.

4. The compound of claim 2, wherein R_1 is selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein said phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, - $O(CH_2)_2NR_{10a}R_{10b}$, - $S(O)_2NR_{10a}R_{10b}$, - $OS(O)_2NR_{10a}R_{10b}$, and - $NR_{10a}S(O)_2R_{10b}$.

5. The compound of claim 4, wherein R_1 is selected from the group consisting of phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzodimidazol-5-yl, wherein said phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-

triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, or 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{O}(\text{CH}_2)_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{S}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{OS}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, and $-\text{NR}_{10a}\text{S}(\text{O})_2\text{R}_{10b}$.

6. The compound of claim 5, wherein R_1 is selected from the group consisting of phenyl, phenol-4-yl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl.

7. The compound of claim 6, wherein R_1 is selected from the group consisting of phenol-4-yl and 1H-indol-3-yl.

8. The compound of any one of claims 1-7, wherein L is selected from the group consisting of $-\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n-$ and $-\text{O}(\text{CR}_{8a}\text{R}_{8b})_n-$.

9. The compound of any one of claims 1-8, wherein R_2 is hydrogen.

10. The compound of any one of claims 1-9, wherein R_3 is selected from the group consisting of phenyl, thiophenyl, furanyl, 1H-benzoimidazolyl, quinolinyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl, wherein said phenyl, thiophenyl, furanyl, 1H-benzoimidazolyl, quinolinyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{C}(\text{O})\text{R}_{11a}$, $-\text{S}(\text{O})_0-2\text{R}_{11a}$, $-\text{C}(\text{O})\text{OR}_{11a}$, and $-\text{C}(\text{O})\text{NR}_{11a}\text{R}_{11b}$, and wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl.

11. The compound of claim 10, wherein R_3 is selected from the group consisting of thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, imidazo[1,2-a]pyridin-3-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl and thiazol-5-yl, wherein said thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl, or thiazol-5-yl is optionally substituted with from 1 to 3 substituents

independently selected from the group consisting of cyano, hydroxy, C₁-4 alkyl, C₁-4 alkenyl, C₁-4 alkynyl, C₂-6 cycloalkyl, C₁-4 alkoxy, halo, halo-substituted-C₁-4 alkyl, halo-substituted-C₁-4 alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

12. The compound of claim 11, wherein R₃ is selected from the group consisting of thiophen-3-yl, benzo[b]thiophen-3-yl, pyridin-3-yl, pyrimidin-5-yl, 1H-imidazol-1-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, and imidazo[1,2-a]pyridin-3-yl, wherein said thiophen-3-yl, benzo[b]thiophen-3-yl, pyridin-3-yl, pyrimidin-5-yl, 1H-imidazol-1-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, or imidazo[1,2-a]pyridin-3-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁-4 alkyl, C₁-4 alkenyl, C₁-4 alkynyl, C₂-6 cycloalkyl, C₁-4 alkoxy, halo, halo-substituted-C₁-4 alkyl, halo-substituted-C₁-4 alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

13. The compound of claim 12, wherein R₃ is pyridin-3-yl, wherein said pyridin-3-yl is optionally substituted at C5 with a substituent selected from the group consisting of C₁-4 alkyl, halo, halo-substituted-C₁-4 alkyl, C₁-4 alkenyl, C₁-4 alkynyl, C₂-6 cycloalkyl, C₁-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

14. The compound of claim 13, wherein said pyridin-3-yl is substituted at C5 with a substituent selected from the group consisting of ethoxycarbonyl, methoxy, cyano, methyl, methylsulfonyl, fluoro, chloro, trifluoromethyl, ethynyl, and cyclopropyl.

15. The compound of claim 12, wherein R₃ is imidazo[1,2-a]pyridin-3-yl, wherein said imidazo[1,2-a]pyridin-3-yl is optionally substituted with a substituent selected from the group consisting of C₁-4 alkyl, halo, halo-substituted-C₁-4 alkyl, C₁-4 alkenyl, C₁-4 alkynyl, C₂-6 cycloalkyl, C₁-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

16. The compound of claim 12, wherein R₃ is benzo[b]thiophen-3-yl, wherein said benzo[b]thiophen-3-yl is optionally substituted with a substituent selected from the group consisting of C₁-4 alkyl, halo, halo-substituted-C₁-4 alkyl, C₁-4 alkenyl, C₁-4 alkynyl, C₂-6 cycloalkyl, C₁-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

17. The compound of claim 12, wherein R₃ is 1H-imidazo[4,5-b]pyridin-1-yl, wherein said 1H-imidazo[4,5-b]pyridin-1-yl is optionally substituted with a substituent selected from the group consisting of C₁-4 alkyl, halo, halo-substituted-C₁-4 alkyl, C₁-4 alkenyl, C₁-4 alkynyl,

C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.




18. The compound of claim 12, wherein R₃ is isoquinolin-4-yl, wherein said isoquinolin-4-yl is optionally substituted with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.



19. The compound of any one of claims 1-18, wherein R₄ is hydrogen.

20. The compound of any one of claims 1-19, wherein R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein said C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl.

21. The compound of claim 20, wherein R₅ is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-hydroxypropan-2-yl, and nonan-2-yl.

22. The compound of any one of claims 1-19, wherein R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)






wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-C(O)R_{12a}$, $-S(O)_{0-2}R_{12a}$, $-C(O)OR_{12a}$, and $-C(O)NR_{12a}R_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl.

23. The compound of any one of claims 1-22, wherein R_6 is hydrogen.

24. The compound of claim 1, wherein said compound is represented by formula (I-a)



wherein L is a linker selected from the group consisting of $-NR_{7a}(CR_{8a}R_{8b})_n$, $-O(CR_{8a}R_{8b})_n$, $-C(O)(CR_{8a}R_{8b})_n$, $-C(S)(CR_{8a}R_{8b})_n$, $-S(O)_{0-2}(CR_{8a}R_{8b})_n$, $-(CR_{8a}R_{8b})_n$, $-NR_{7a}C(O)(CR_{8a}R_{8b})_n$, $-NR_{7a}C(S)(CR_{8a}R_{8b})_n$, $-OC(O)(CR_{8a}R_{8b})_n$, $-OC(S)(CR_{8a}R_{8b})_n$, $-C(O)NR_{7a}(CR_{8a}R_{8b})_n$, $-C(S)NR_{7a}(CR_{8a}R_{8b})_n$, $-C(O)O(CR_{8a}R_{8b})_n$, $-C(S)O(CR_{8a}R_{8b})_n$, $-S(O)_2NR_{7a}(CR_{8a}R_{8b})_n$, $-NR_{7a}S(O)_2(CR_{8a}R_{8b})_n$, $-NR_{7a}C(O)NR_{7b}(CR_{8a}R_{8b})_n$, and $-NR_{7a}C(O)O(CR_{8a}R_{8b})_n$, wherein R_{7a} , R_{7b} , R_{8a} , and R_{8b} are each independently selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl, and each n is independently an integer from 2 to 6;

R_1 is selected from the group consisting of $-S(O)_2NR_{9a}R_{9b}$, $-NR_{9a}C(O)R_{9b}$, $-NR_{9a}C(S)R_{9b}$, $-NR_{9a}C(O)NR_{9b}R_{9c}$, $-C(O)R_{9a}$, $-C(S)R_{9a}$, $-S(O)_{0-2}R_{9a}$, $-C(O)OR_{9a}$, $-C(S)OR_{9a}$, $-C(O)NR_{9a}R_{9b}$, $-C(S)NR_{9a}R_{9b}$, $-NR_{9a}S(O)_2R_{9b}$, $-NR_{9a}C(O)OR_{9b}$, $-OC(O)CR_{9a}R_{9b}R_{9c}$, $-OC(S)CR_{9a}R_{9b}R_{9c}$, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein R_{9a} , R_{9b} , and R_{9c} are each independently selected from the group consisting of hydrogen, optionally substituted

aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

Ar is selected from the group consisting of optionally substituted monocyclic aryl and heteroaryl;


R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R₆ is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

25. The compound of claim 24, wherein R₁ is selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl.

26. The compound of claim 1, wherein said compound is represented by formula (I-b)



wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{O}(\text{CH}_2)_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{S}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{OS}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, and $-\text{NR}_{10a}\text{S}(\text{O})_2\text{R}_{10b}$, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;


Ar is selected from the group consisting of optionally substituted monocyclic aryl and heteroaryl;

R_5 is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R_6 is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

27. The compound of claim 1, wherein said compound is represented by formula (I-c)



wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{O}(\text{CH}_2)_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{S}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{OS}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, and $-\text{NR}_{10a}\text{S}(\text{O})_2\text{R}_{10b}$, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;


B is an optionally substituted ring system selected from the group consisting of thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl, wherein the thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, 1H-imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{C}(\text{O})\text{R}_{11a}$, $-\text{S}(\text{O})_2\text{R}_{11a}$, $-\text{C}(\text{O})\text{OR}_{11a}$, and $-\text{C}(\text{O})\text{NR}_{11a}\text{R}_{11b}$, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl;

R_5 is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R_6 is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

28. The compound of claim 1, wherein said compound is represented by formula (I-d)




wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

B is an optionally substituted ring system selected from the group consisting of thiophenyl, furanyl, 1H-benzoimidazolyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl, wherein the thiophenyl, furanyl, 1H-benzoimidazolyl, isoquinolinyl, 1H-imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

29. The compound of claim 1, wherein said compound is represented by formula (I-e)




wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl, wherein the phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, or 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

B is an optionally substituted ring system selected from the group consisting of thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, imidazo[1,2-a]pyridin-3-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl and thiazol-5-yl, wherein the thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl, or thiazol-5-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and


R₅ is selected from the group consisting of C₁₋₁₀ alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-

pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R_5 is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-C(O)R_{12a}$, $-S(O)_{0-2}R_{12a}$, $-C(O)OR_{12a}$, and $-C(O)NR_{12a}R_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl; or a salt thereof.

30. The compound of claim 1, wherein said compound is represented by formula (I-f)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

q is an integer from 0 to 4;

each Z is independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl; and


R₅ is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-hydroxypropan-2-yl, and nonan-2-yl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4

alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{C}(\text{O})\text{R}_{12a}$, $-\text{S}(\text{O})_{0-2}\text{R}_{12a}$, $-\text{C}(\text{O})\text{OR}_{12a}$, and $-\text{C}(\text{O})\text{NR}_{12a}\text{R}_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl;
or a salt thereof.


31. The compound of claim 1, wherein said compound is represented by formula (I-g)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

Z is a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, $\text{C}(\text{O})\text{R}_{11a}$, $-\text{S}(\text{O})_{0-2}\text{R}_{11a}$, $-\text{C}(\text{O})\text{OR}_{11a}$, and $-\text{C}(\text{O})\text{NR}_{11a}\text{R}_{11b}$, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl; and


R_5 is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-hydroxypropan-2-yl, and nonan-2-yl, or R_5 is selected from the group consisting of (i), (ii), (iii), (iv), and (v)





wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;
or a salt thereof.

32. The compound of claim 1, wherein said compound is represented by formula (I-h)



wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;


q is an integer from 0 to 4;

r is 0 or 1;

W and V are each independently a substituent selected from the group consisting of C₁₋₄ alkyl, halo, halo-substituted-C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and


R₅ is selected from the group consisting of C₁₋₁₀ alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C₁₋₁₀ alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-

pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R_5 is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-C(O)R_{12a}$, $-S(O)_{0-2}R_{12a}$, $-C(O)OR_{12a}$, and $-C(O)NR_{12a}R_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl; or a salt thereof.

33. The compound of claim 1, wherein said compound is represented by formula (I-i)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

q is an integer from 0 to 4;

r is 0 or 1;

W and V are each independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and


R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -

$C(O)NR_{12a}R_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;
or a salt thereof.

34. The compound of claim 1, wherein said compound is represented by formula (I-j)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;


q is an integer from 0 to 4;

r is 0 or 1;

W and V are each independently a substituent selected from the group consisting of C₁₋₄ alkyl, halo, halo-substituted-C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, cyano, amino, $C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and


R_5 is selected from the group consisting of C₁₋₁₀ alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C₁₋₁₀ alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C₁₋₄ alkyl, and halo-substituted-C₁₋₄alkyl, or R_5 is selected from the group consisting of (i), (ii), (iii), (iv), and (v)





wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;
or a salt thereof.

35. The compound of claim 1, wherein said compound is represented by formula (I-k)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

q is an integer from 0 to 4;

r is 0 or 1;


W and V are each independently a substituent selected from the group consisting of C₁₋₄ alkyl, halo, halo-substituted-C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)




wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; or a salt thereof.

36. The compound of claim 1, wherein said compound is compound (1)




or a salt thereof.

37. The compound of claim 1, wherein said compound is compound (2)




or a salt thereof.

38. The compound of claim 1, wherein said compound is compound (3)




or a salt thereof.

39. The compound of claim 1, wherein said compound is compound (4)




or a salt thereof.

40. The compound of claim 1, wherein said compound is compound (5)




or a salt thereof.

41. The compound of claim 1, wherein said compound is compound (6)




or a salt thereof.

42. The compound of claim 1, wherein said compound is compound (7)




or a salt thereof.

43. The compound of claim 1, wherein said compound is compound (8)




or a salt thereof.

44. The compound of claim 1, wherein said compound is compound (9)




or a salt thereof.

45. The compound of claim 1, wherein said compound is compound (10)




or a salt thereof.

46. The compound of claim 1, wherein said compound is compound (11)




or a salt thereof.

47. The compound of claim 1, wherein said compound is compound (23)




or a salt thereof.

48. The compound of claim 1, wherein said compound is compound (25)




or a salt thereof.

49. The compound of claim 1, wherein said compound is compound (26)



or a salt thereof.

50. A compound represented by formula (II)



wherein L is selected from the group consisting of $-\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{O}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{C}(\text{O})(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{C}(\text{S})(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{S}(\text{O})_{0-2}(\text{CR}_{8a}\text{R}_{8b})_n$, $-(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{NR}_{7a}\text{C}(\text{O})(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{NR}_{7a}\text{C}(\text{S})(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{OC}(\text{O})(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{OC}(\text{S})(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{C}(\text{O})\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{C}(\text{S})\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{C}(\text{O})\text{O}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{C}(\text{S})\text{O}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{S}(\text{O})_2\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{NR}_{7a}\text{S}(\text{O})_2(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{NR}_{7a}\text{C}(\text{O})\text{NR}_{7b}(\text{CR}_{8a}\text{R}_{8b})_n$, and $-\text{NR}_{7a}\text{C}(\text{O})\text{O}(\text{CR}_{8a}\text{R}_{8b})_n$, wherein R_{7a} , R_{7b} , R_{8a} , and R_{8b} are each independently selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl, and each n is independently an integer from 2 to 6;

R_1 is selected from the group consisting of $-\text{S}(\text{O})_2\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{S})\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{NR}_{9b}\text{R}_{9c}$, $-\text{C}(\text{O})\text{R}_{9a}$, $-\text{C}(\text{S})\text{R}_{9a}$, $-\text{S}(\text{O})_{0-2}\text{R}_{9a}$, $-\text{C}(\text{O})\text{OR}_{9a}$, $-\text{C}(\text{S})\text{OR}_{9a}$,

C(O)NR_{9a}R_{9b}, -C(S)NR_{9a}R_{9b}, -NR_{9a}S(O)₂R_{9b}, -NR_{9a}C(O)OR_{9b}, -OC(O)CR_{9a}R_{9b}R_{9c}, -OC(S)CR_{9a}R_{9b}R_{9c}, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein R_{9a}, R_{9b}, and R_{9c} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

R₃ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

R₄ is selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl;

R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R₆ is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; or a salt thereof.

51. The compound of claim 50, wherein R₁ is selected from the group consisting of -S(O)₂NR_{9a}R_{9b}, -NR_{9a}C(O)R_{9b}, -NR_{9a}C(S)R_{9b}, -NR_{9a}C(O)NR_{9b}R_{9c}, -C(O)R_{9a}, -C(S)R_{9a}, -S(O)₂R_{9a}, -C(O)OR_{9a}, -C(S)OR_{9a}, -C(O)NR_{9a}R_{9b}, -C(S)NR_{9a}R_{9b}, -NR_{9a}S(O)₂R_{9b}, -NR_{9a}C(O)OR_{9b}, -OC(O)CR_{9a}R_{9b}R_{9c}, -OC(S)CR_{9a}R_{9b}R_{9c}, phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein said phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl.

52. The compound of claim 51, wherein R₁ is selected from the group consisting of -S(O)₂NR_{9a}R_{9b}, -NR_{9a}C(O)R_{9b}, -NR_{9a}C(S)R_{9b}, -NR_{9a}C(O)NR_{9b}R_{9c}, -C(O)R_{9a}, -C(S)R_{9a}, -S(O)₂R_{9a},

$2R_{9a}$, $-C(O)OR_{9a}$, $-C(S)OR_{9a}$, $-C(O)NR_{9a}R_{9b}$, $-C(S)NR_{9a}R_{9b}$, $-NR_{9a}S(O)_2R_{9b}$, $-NR_{9a}C(O)OR_{9b}$, $-OC(O)CR_{9a}R_{9b}R_{9c}$, and $-OC(S)CR_{9a}R_{9b}R_{9c}$.

53. The compound of claim 51, wherein R_1 is selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, and 1H-indazolyl, wherein said phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-O(CH_2)_2NR_{10a}R_{10b}$, $-S(O)_2NR_{10a}R_{10b}$, $-OS(O)_2NR_{10a}R_{10b}$, and $-NR_{10a}S(O)_2R_{10b}$.

54. The compound of claim 53, wherein R_1 is selected from the group consisting of phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl, wherein said phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, or 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-O(CH_2)_2NR_{10a}R_{10b}$, $-S(O)_2NR_{10a}R_{10b}$, $-OS(O)_2NR_{10a}R_{10b}$, and $-NR_{10a}S(O)_2R_{10b}$.

55. The compound of claim 54, wherein R_1 is selected from the group consisting of phenyl, phenol-4-yl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl.

56. The compound of claim 55, wherein R_1 is selected from the group consisting of phenol-4-yl and 1H-indol-3-yl.

57. The compound of any one of claims 50-56, wherein L is selected from the group consisting of $-NR_{7a}(CR_{8a}R_{8b})_n-$ and $-O(CR_{8a}R_{8b})_n-$.

58. The compound of any one of claims 50-57, wherein R_3 is selected from the group consisting of phenyl, thiophenyl, furanyl, 1H-benzoimidazolyl, quinolinyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl, wherein said phenyl, thiophenyl, furanyl, 1H-benzoimidazolyl,

quinolinyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, and wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl.

59. The compound of claim 58, wherein R₃ is selected from the group consisting of thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, imidazo[1,2-a]pyridin-3-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl and thiazol-5-yl, wherein said thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl, or thiazol-5-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

60. The compound of claim 59, wherein R₃ is selected from the group consisting of thiophen-3-yl, benzo[b]thiophen-3-yl, pyridin-3-yl, pyrimidin-5-yl, 1H-imidazol-1-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, and imidazo[1,2-a]pyridin-3-yl, wherein said thiophen-3-yl, benzo[b]thiophen-3-yl, pyridin-3-yl, pyrimidin-5-yl, 1H-imidazol-1-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, or imidazo[1,2-a]pyridin-3-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

61. The compound of claim 60, wherein R₃ is pyridin-3-yl, wherein said pyridin-3-yl is optionally substituted at C5 with a substituent selected from the group consisting of C₁₋₄ alkyl, halo, halo-substituted-C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

62. The compound of claim 61, wherein said pyridin-3-yl is substituted at C5 with a substituent selected from the group consisting of ethoxycarbonyl, methoxy, cyano, methyl, methylsulfonyl, fluoro, chloro, trifluoromethyl, ethynyl, and cyclopropyl.

63. The compound of claim 60, wherein R₃ is imidazo[1,2-a]pyridin-3-yl, wherein said imidazo[1,2-a]pyridin-3-yl is optionally substituted with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

64. The compound of claim 60, wherein R₃ is benzo[b]thiophen-3-yl, wherein said benzo[b]thiophen-3-yl is optionally substituted with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

65. The compound of claim 60, wherein R₃ is 1H-imidazo[4,5-b]pyridin-1-yl, wherein said 1H-imidazo[4,5-b]pyridin-1-yl is optionally substituted with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.

66. The compound of claim 60, wherein R₃ is isoquinolin-4-yl, wherein said isoquinolin-4-yl is optionally substituted with a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}.






67. The compound of any one of claims 50-66, wherein R₄ is hydrogen.

68. The compound of any one of claims 50-67, wherein R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein said C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl.

69. The compound of claim 68, wherein R₅ is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-

butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-hydroxypropan-2-yl, and nonan-2-yl.


70. The compound of any one of claims 50-67, wherein R_5 is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-C(O)R_{12a}$, $-S(O)_{0-2}R_{12a}$, $-C(O)OR_{12a}$, and $-C(O)NR_{12a}R_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl.

71. The compound of any one of claims 50-70, wherein R_6 is hydrogen.

72. The compound of claim 50, wherein said compound is represented by formula (II-a)



wherein L is selected from the group consisting of $-\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{O}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{C}(\text{O})(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{C}(\text{S})(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{S}(\text{O})_{0-2}(\text{CR}_{8a}\text{R}_{8b})_n$, $-(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{NR}_{7a}\text{C}(\text{O})(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{NR}_{7a}\text{C}(\text{S})(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{OC}(\text{O})(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{OC}(\text{S})(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{C}(\text{O})\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{C}(\text{S})\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{C}(\text{O})\text{O}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{C}(\text{S})\text{O}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{S}(\text{O})_2\text{NR}_{7a}(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{NR}_{7a}\text{S}(\text{O})_2(\text{CR}_{8a}\text{R}_{8b})_n$, $-\text{NR}_{7a}\text{C}(\text{O})\text{NR}_{7b}(\text{CR}_{8a}\text{R}_{8b})_n$, and $-\text{NR}_{7a}\text{C}(\text{O})\text{O}(\text{CR}_{8a}\text{R}_{8b})_n$, wherein R_{7a} , R_{7b} , R_{8a} , and R_{8b} are each independently selected from the group consisting of hydrogen and optionally substituted C1-4 alkyl, and each n is independently an integer from 2 to 6;

R_1 is selected from the group consisting of $-\text{S}(\text{O})_2\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{S})\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{NR}_{9b}\text{R}_{9c}$, $-\text{C}(\text{O})\text{R}_{9a}$, $-\text{C}(\text{S})\text{R}_{9a}$, $-\text{S}(\text{O})_{0-2}\text{R}_{9a}$, $-\text{C}(\text{O})\text{OR}_{9a}$, $-\text{C}(\text{S})\text{OR}_{9a}$, $-\text{C}(\text{O})\text{NR}_{9a}\text{R}_{9b}$, $-\text{C}(\text{S})\text{NR}_{9a}\text{R}_{9b}$, $-\text{NR}_{9a}\text{S}(\text{O})_2\text{R}_{9b}$, $-\text{NR}_{9a}\text{C}(\text{O})\text{OR}_{9b}$, $-\text{OC}(\text{O})\text{CR}_{9a}\text{R}_{9b}\text{R}_{9c}$, $-\text{OC}(\text{S})\text{CR}_{9a}\text{R}_{9b}\text{R}_{9c}$, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein R_{9a} , R_{9b} , and R_{9c} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

Ar is selected from the group consisting of optionally substituted monocyclic aryl and heteroaryl;


R_5 is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R_6 is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

73. The compound of claim 72, wherein R_1 is selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{O}(\text{CH}_2)_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{S}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, $-\text{OS}(\text{O})_2\text{NR}_{10a}\text{R}_{10b}$, and $-\text{NR}_{10a}\text{S}(\text{O})_2\text{R}_{10b}$, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl.

74. The compound of claim 50, wherein said compound is represented by formula (II-b)



wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;


Ar is selected from the group consisting of optionally substituted monocyclic aryl and heteroaryl;

R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R₆ is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

75. The compound of claim 50, wherein said compound is represented by formula (II-c)



wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzoimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;


B is an optionally substituted ring system selected from the group consisting of thiophenyl, furanyl, 1H-benzoimidazolyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl, wherein the thiophenyl, furanyl, 1H-benzoimidazolyl, isoquinolinyl, 1H-imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;

R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl; and

R₆ is selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.

76. The compound of claim 50, wherein said compound is represented by formula (II-d)




wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, and 1H-indazolyl, wherein the phenyl, 1H-pyrrolopyridinyl, 1H-pyrrolopyridinyl, 1H-indolyl, thiophenyl, pyridinyl, 1H-1,2,4-triazolyl, 2-oxoimidazolidinyl, 1H-pyrazolyl, 2-oxo-2,3-dihydro-1H-benzimidazolyl, or 1H-indazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -O(CH₂)₂NR_{10a}R_{10b}, -S(O)₂NR_{10a}R_{10b}, -OS(O)₂NR_{10a}R_{10b}, and -NR_{10a}S(O)₂R_{10b}, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

B is an optionally substituted ring system selected from the group consisting of thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, and thiazolyl, wherein the thiophenyl, furanyl, 1H-benzimidazolyl, isoquinolinyl, 1H-imidazopyridinyl, benzothiophenyl, pyrimidinyl, pyridinyl, 1H-imidazolyl, pyrazinyl, pyridazinyl, 1H-pyrrolyl, or thiazolyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

R₅ is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;

or a salt thereof.


77. The compound of claim 50, wherein said compound is represented by formula (II-e)



wherein A is an optionally substituted ring system selected from the group consisting of phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, and 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl, wherein the phenyl, 1H-indol-2-yl, 1H-indol-3-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 2-oxoimidazolidin-1-yl, 1H-pyrazol-3-yl, 1H-pyrazol-4-yl, or 2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-O(CH_2)_2NR_{10a}R_{10b}$, $-S(O)_2NR_{10a}R_{10b}$, $-OS(O)_2NR_{10a}R_{10b}$, and $-NR_{10a}S(O)_2R_{10b}$, wherein R_{10a} and R_{10b} are each independently selected from the group consisting of hydrogen, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkyl, optionally substituted heteroalkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl;


B is an optionally substituted ring system selected from the group consisting of thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, imidazo[1,2-a]pyridin-3-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl and thiazol-5-yl, wherein the thiophen-2-yl, thiophen-3-yl, furan-3-yl, 1H-benzo[d]imidazol-1-yl, isoquinolin-4-yl, 1H-imidazo[4,5-b]pyridin-1-yl, benzo[b]thiophen-3-yl, pyrimidin-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 1H-imidazol-1-yl, pyrazin-2-yl, pyridazin-4-yl, 1H-pyrrol-2-yl, or thiazol-5-yl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl; and

R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkenyl, C₁₋₄ alkynyl, C₂₋₆ cycloalkyl, C₁₋₄ alkoxy, halo, halo-substituted-C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; or a salt thereof.

78. The compound of claim 50, wherein said compound is represented by formula (II-f)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

q is an integer from 0 to 4;

each Z is independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl; and


R₅ is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-hydroxypropan-2-yl, and nonan-2-yl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4

alkyl, halo-substituted-C1-4 alkoxy, amino, $-C(O)R_{12a}$, $-S(O)_{0-2}R_{12a}$, $-C(O)OR_{12a}$, and $-C(O)NR_{12a}R_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;
or a salt thereof.





79. The compound of claim 50, wherein said compound is represented by formula (II-g)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

Z is a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, $C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and


R_5 is selected from the group consisting of isopropyl, methyl, ethyl, prop-1-en-2-yl, isobutyl, cyclohexyl, sec-butyl, (S)-sec-butyl, (R)-sec-butyl, 1-hydroxypropan-2-yl, (S)-1-hydroxypropan-2-yl, (R)-1-hydroxypropan-2-yl, and nonan-2-yl, or R_5 is selected from the group consisting of (i), (ii), (iii), (iv), and (v)





wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;
or a salt thereof.

80. The compound of claim 50, wherein said compound is represented by formula (II-h)



wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;


q is an integer from 0 to 4;

r is 0 or 1;

W and V are each independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and


R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl,

tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R_5 is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-C(O)R_{12a}$, $-S(O)_{0-2}R_{12a}$, $-C(O)OR_{12a}$, and $-C(O)NR_{12a}R_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; or a salt thereof.

81. The compound of claim 50, wherein said compound is represented by formula (II-i)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

q is an integer from 0 to 4;

r is 0 or 1;

W and V are each independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C1-4 alkyl; and


R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4

alkyl, halo-substituted-C1-4 alkoxy, amino, $-C(O)R_{12a}$, $-S(O)_{0-2}R_{12a}$, $-C(O)OR_{12a}$, and $-C(O)NR_{12a}R_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;
or a salt thereof.

82. The compound of claim 50, wherein said compound is represented by formula (II-j)




wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;

q is an integer from 0 to 4;

r is 0 or 1;


W and V are each independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, $C(O)R_{11a}$, $-S(O)_{0-2}R_{11a}$, $-C(O)OR_{11a}$, and $-C(O)NR_{11a}R_{11b}$, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

R_5 is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4alkyl, or R_5 is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, $-\text{C}(\text{O})\text{R}_{12a}$, $-\text{S}(\text{O})_{0-2}\text{R}_{12a}$, $-\text{C}(\text{O})\text{OR}_{12a}$, and $-\text{C}(\text{O})\text{NR}_{12a}\text{R}_{12b}$, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; or a salt thereof.


83. The compound of claim 50, wherein said compound is represented by formula (II-k)



wherein A is an optionally substituted ring system selected from the group consisting of phenol-4-yl and 1H-indol-3-yl;
 q is an integer from 0 to 4;
 r is 0 or 1;

W and V are each independently a substituent selected from the group consisting of C1-4 alkyl, halo, halo-substituted-C1-4 alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, cyano, amino, C(O)R_{11a}, -S(O)₀₋₂R_{11a}, -C(O)OR_{11a}, and -C(O)NR_{11a}R_{11b}, wherein R_{11a} and R_{11b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and


R₅ is selected from the group consisting of C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, and 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl, wherein the C1-10 alkyl, prop-1-en-2-yl, cyclohexyl, cyclopropyl, 2-(2-oxopyrrolidin-1-yl)ethyl, oxetan-2-yl, oxetan-3-yl, benzhydryl, tetrahydro-2H-pyran-2-yl, tetrahydro-2H-pyran-3-yl, phenyl, tetrahydrofuran-3-yl, benzyl, (4-pentylphenyl)(phenyl)methyl, or 1-(1-(2-oxo-6,9,12-trioxa-3-azatetradecan-14-yl)-1H-1,2,3-triazol-4-yl)ethyl is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C1-4 alkyl, and halo-substituted-C1-4 alkyl, or R₅ is selected from the group consisting of (i), (ii), (iii), (iv), and (v)



wherein n is an integer from 1 to 6, m is an integer from 0 to 6, p is an integer from 0 to 5, and each R is independently selected from the group consisting of cyano, hydroxy, C₁₋₄ alkyl, C1-4 alkenyl, C1-4 alkynyl, C2-6 cycloalkyl, C1-4 alkoxy, halo, halo-substituted-C1-4 alkyl, halo-substituted-C1-4 alkoxy, amino, -C(O)R_{12a}, -S(O)₀₋₂R_{12a}, -C(O)OR_{12a}, and -C(O)NR_{12a}R_{12b}, and wherein R_{12a} and R_{12b} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl;


or a salt thereof.

84. The compound of claim 50, wherein said compound is compound (12)




or a salt thereof.

85. The compound of claim 50, wherein said compound is compound (13)




or a salt thereof.

86. The compound of claim 50, wherein said compound is compound (14)




or a salt thereof.

87. The compound of claim 50, wherein said compound is compound (15)




or a salt thereof.

88. The compound of claim 50, wherein said compound is compound (16)




or a salt thereof.

89. The compound of claim 50, wherein said compound is compound (17)



or a salt thereof.


90. The compound of claim 50, wherein said compound is compound (18)



(18)

or a salt thereof.


91. The compound of claim 50, wherein said compound is compound (19)



(19)

or a salt thereof.


92. The compound of claim 50, wherein said compound is compound (20)



(20)


or a salt thereof.

93. The compound of claim 50, wherein said compound is compound (21)




or a salt thereof.

94. The compound of claim 50, wherein said compound is compound (22)




or a salt thereof.

95. The compound of claim 50, wherein said compound is compound (24)




or a salt thereof.

96. The compound of claim 50, wherein said compound is compound (27)



or a salt thereof.

97. The compound of claim 50, wherein said compound is compound (28)



or a salt thereof.

98. A method of producing an expanded population of hematopoietic stem cells ex vivo, said method comprising contacting a population of hematopoietic stem cells with the compound of any one of claims 1-97 in an amount sufficient to produce an expanded population of hematopoietic stem cells.

99. A method of enriching a population of cells with hematopoietic stem cells ex vivo, said method comprising contacting a population of hematopoietic stem cells with the compound of any one of claims 1-97 in an amount sufficient to produce a population of cells enriched with hematopoietic stem cells.

100. A method of maintaining the hematopoietic stem cell functional potential of a population of hematopoietic stem cells ex vivo for two or more days, said method comprising contacting a first population of hematopoietic stem cells with the compound of any one of claims 1-97, wherein the first population of hematopoietic stem cells exhibits a hematopoietic stem cell functional potential after two or more days that is greater than that of a control population of hematopoietic stem cells cultured under the same conditions and for the same

time as said first population of hematopoietic stem cells but not contacted with said compound.

101. The method of claim 100, wherein said first population of hematopoietic stem cells exhibits a hematopoietic stem cell functional potential after three or more days of culture that is greater than that of said control population of hematopoietic stem cells.

102. The method of claim 100, wherein said first population of hematopoietic stem cells exhibits a hematopoietic stem cell functional potential after ten or more days of culture that is greater than that of said control population of hematopoietic stem cells.

103. The method of claim 100, wherein said first population of hematopoietic stem cells exhibits a hematopoietic stem cell functional potential after thirty or more days of culture that is greater than that of said control population of hematopoietic stem cells.

104. The method of claim 100, wherein said first population of hematopoietic stem cells exhibits a hematopoietic stem cell functional potential after sixty or more days of culture that is greater than that of said control population of hematopoietic stem cells.

105. The method of any one of claims 98-104, wherein said hematopoietic stem cells are mammalian cells.

106. The method of claim 105, wherein said mammalian cells are human cells.

107. The method of claim 106, wherein said human cells are CD34+ cells.

108. The method of claim 107, wherein said CD34+ cells are CD34+, CD34+CD38-, CD34+CD38-CD90+, CD34+CD38-CD90+CD45RA-, CD34+CD38-CD90+CD45RA-CD49F+, or CD34+CD90+CD45RA- cells.

109. The method of any one of claims 98-108, wherein said hematopoietic stem cells are obtained from human cord blood.

110. The method of any one of claims 98-108, wherein said hematopoietic stem cells are obtained from mobilized human peripheral blood.

111. The method of any one of claims 98-108, wherein said hematopoietic stem cells are obtained from human bone marrow.

112. The method of any one of claims 98-111, wherein said hematopoietic stem cells are freshly isolated from said human.

113. The method of any one of claims 98-111, wherein said hematopoietic stem cells have been previously cryopreserved.

114. The method of any one of claims 98-113, wherein said hematopoietic stem cells or progeny thereof maintain hematopoietic stem cell functional potential after two or more days upon transplantation of said hematopoietic stem cells into a human subject.

115. The method of any one of claims 98-114, wherein said hematopoietic stem cells or progeny thereof are capable of localizing to hematopoietic tissue and reestablishing hematopoiesis upon transplantation of said hematopoietic stem cells into a human subject.

116. The method of any one of claims 98-115, wherein upon transplantation into a human subject, said hematopoietic stem cells give rise to a population of cells selected from the group consisting of megakaryocytes, thrombocytes, platelets, erythrocytes, mast cells, myoblasts, basophils, neutrophils, eosinophils, microglia, granulocytes, monocytes, osteoclasts, antigen-presenting cells, macrophages, dendritic cells, natural killer cells, T-lymphocytes, and B-lymphocytes.

117. A method of treating a human patient suffering from a disorder, said method comprising administering to said patient a population of hematopoietic stem cells, wherein said hematopoietic stem cells were produced by contacting said hematopoietic stem cells or progenitors thereof with the compound of any one of claims 1-97.

118. A method of preparing an expanded population of hematopoietic stem cells for transplantation into a human patient suffering from a disorder, said method comprising contacting a first population of hematopoietic stem cells with the compound of any one of claims 1-97 for a time sufficient to produce said expanded population of hematopoietic stem cells.

119. A method of treating a human patient suffering from a disorder, said method comprising:

- a. preparing an expanded population of hematopoietic stem cells by contacting a first population of hematopoietic stem cells with the compound of any one of claims 1-97 for a time sufficient to produce said expanded population of hematopoietic stem cells; and
- b. administering said expanded population of hematopoietic stem cells to said patient.

120. A method of treating a disorder in a human patient in need thereof, comprising administering an expanded population of hematopoietic stem cells to the patient,

wherein the expanded population of hematopoietic stem cells is prepared by contacting a first population of hematopoietic stem cells with the compound of any one of claims 1-97 for a time sufficient to produce the expanded population of hematopoietic stem cells.

121. The method of any one of claims 117-120, wherein said disorder is a hemoglobinopathy.

122. The method of any one of claims 117-120, wherein said disorder is selected from the group consisting of sickle cell anemia, thalassemia, Fanconi anemia, and Wiskott-Aldrich syndrome.

123. The method of any one of claims 117-120, wherein the disorder is a myelodysplastic disorder

124. The method of any one of claims 117-120, wherein said disorder is an immunodeficiency disorder.

125. The method of claim 124, wherein said immunodeficiency disorder is a congenital immunodeficiency.

126. The method of claim 124, wherein said immunodeficiency disorder is an acquired immunodeficiency.

127. The method of claim 126, wherein said acquired immunodeficiency is human immunodeficiency virus or acquired immune deficiency syndrome.

128. The method of any one of claims 117-120, wherein said disorder is a metabolic disorder.

129. The method of claim 128, wherein said metabolic disorder is selected from the group consisting of glycogen storage diseases, mucopolysaccharidoses, Gaucher's Disease, Hurlers Disease, sphingolipidoses, and metachromatic leukodystrophy.

130. The method of any one of claims 117-120, wherein the disorder is cancer.

131. The method of claim 130, wherein the cancer is a hematological cancer.

132. The method of claim 130, wherein the cancer is selected from the group consisting of leukemia, lymphoma, multiple myeloma, and neuroblastoma.

133. The method of claim 130, wherein the cancer is acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia, chronic lymphoid leukemia, multiple myeloma, diffuse large B-cell lymphoma, or non-Hodgkin's lymphoma.

134. The method of any one of claims 117-120, wherein the disorder is a disorder selected from the group consisting of adenosine deaminase deficiency and severe combined immunodeficiency, hyper immunoglobulin M syndrome, Chediak-Higashi disease, hereditary lymphohistiocytosis, osteopetrosis, osteogenesis imperfecta, storage diseases, thalassemia major, systemic sclerosis, systemic lupus erythematosus, multiple sclerosis, and juvenile rheumatoid arthritis.

135. The method of any one of claims 117-120, wherein the disorder is an autoimmune disorder.

136. The method of claim 135, wherein the autoimmune disorder is selected from the group consisting of multiple sclerosis, human systemic lupus, rheumatoid arthritis, inflammatory bowel disease, treating psoriasis, Type 1 diabetes mellitus, acute disseminated encephalomyelitis, Addison's disease, alopecia universalis, ankylosing spondylitis, antiphospholipid antibody syndrome, aplastic anemia, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune inner ear disease, autoimmune lymphoproliferative syndrome, autoimmune oophoritis, Balo disease, Behcet's disease, bullous pemphigoid, cardiomyopathy, Chagas' disease, chronic fatigue immune dysfunction syndrome, chronic inflammatory demyelinating polyneuropathy, Crohn's disease, cicatricial pemphigoid, coeliac sprue-dermatitis herpetiformis, cold agglutinin disease, CREST syndrome, Degos disease, discoid lupus, dysautonomia, endometriosis, essential mixed cryoglobulinemia, fibromyalgia-fibromyositis, Goodpasture's syndrome, Grave's disease, Guillain-Barre syndrome, Hashimoto's thyroiditis, Hidradenitis suppurativa, idiopathic and/or acute thrombocytopenic purpura, idiopathic pulmonary fibrosis, IgA neuropathy, interstitial cystitis, juvenile arthritis, Kawasaki's disease, lichen planus, Lyme disease, Meniere disease, mixed connective tissue disease, myasthenia gravis, neuromyotonia, opsoclonus myoclonus syndrome, optic neuritis, Ord's thyroiditis, pemphigus vulgaris, pernicious anemia, polychondritis, polymyositis and dermatomyositis, primary biliary cirrhosis, polyarteritis nodosa, polyglandular syndromes, polymyalgia rheumatica, primary agammaglobulinemia, Raynaud phenomenon, Reiter's syndrome, rheumatic fever, sarcoidosis, scleroderma, Sjögren's syndrome, stiff person syndrome, Takayasu's arteritis, temporal arteritis, ulcerative colitis, uveitis, vasculitis, vitiligo, vulvodynia, and Wegener's granulomatosis.

137. The method of any one of claims 117-120, wherein the disorder is a neurological disorder.

138. The method of claim 137, wherein the neurological disorder is selected from the group consisting of Parkinson's disease, Alzheimer's disease, multiple sclerosis, Amyotrophic lateral sclerosis, Huntington's disease, mild cognitive impairment, amyloidosis, AIDS-related dementia, encephalitis, stroke, head trauma, epilepsy, mood disorders, and dementia.

139. A method of producing microglia in the central nervous system of a human patient in need thereof, comprising administering an expanded population of hematopoietic stem cells to the patient, wherein the expanded population of hematopoietic stem cells is prepared by contacting a first population of hematopoietic stem cells with the compound of any one of claims 1-97 for a time sufficient to produce the expanded population of hematopoietic stem cells, and wherein administration of the expanded population of hematopoietic stem cells results in formation of microglia in the central nervous system of the patient.

140. A composition comprising a population of hematopoietic stem cells, wherein said hematopoietic stem cells or progenitors thereof have been contacted with the compound of any one of claims 1-97, thereby expanding said hematopoietic stem cells or progenitors thereof.

141. A kit comprising the compound of any one of claims 1-97 and a package insert, wherein said package insert instructs a user of said kit to contact a population of hematopoietic stem cells with said compound for a time sufficient to produce an expanded population of hematopoietic stem cells.

142. A kit comprising the compound of any one of claims 1-97 and a package insert, wherein said package insert instructs a user of said kit to contact a population of cells comprising hematopoietic stem cells with said compound for a time sufficient to produce a population of cells enriched with hematopoietic stem cells.

143. A kit comprising the compound of any one of claims 1-97 and a package insert, wherein said package insert instructs a user of said kit to contact a population of hematopoietic stem cells with said compound for a time sufficient to maintain the hematopoietic stem cell functional potential of said population of hematopoietic stem cells ex vivo for two or more days.

144. The kit of any one of claims 141-143, wherein said kit further comprises a population of cells comprising hematopoietic stem cells.

145. The compound of any one of claims 1-97 for use in producing an expanded population of hematopoietic stem cells ex vivo.

146. The compound of any one of claims 1-97 for use in enriching a population of cells with hematopoietic stem cells ex vivo.

147. The compound of any one of claims 1-97 for use in maintaining the hematopoietic stem cell functional potential of a population of hematopoietic stem cells ex vivo for two or more days.

148. Use of the compound of any one of claims 1-97 in producing an expanded population of hematopoietic stem cells ex vivo.

149. Use of the compound of any one of claims 1-97 in enriching a population of cells with hematopoietic stem cells ex vivo.

150. Use of the compound of any one of claims 1-97 in maintaining the hematopoietic stem cell functional potential of a population of hematopoietic stem cells ex vivo for two or more days.

151. A composition for use in treating a human patient suffering from a disorder, said composition comprising a population of hematopoietic stem cells, wherein said hematopoietic stem cells were produced by contacting said hematopoietic stem cells or progenitors thereof with the compound of any one of claims 1-97.

152. A composition for use in treating a human patient suffering from a disorder, said composition comprising an expanded population of hematopoietic stem cells prepared by contacting a first population of hematopoietic stem cells with the compound of any one of claims 1-97 for a time sufficient to produce said expanded population of hematopoietic stem cells.

153. Use of the compound of any one of claims 1-97 in preparing a medicament for treating a human patient suffering from a disorder.

154. The composition of claim 151 or 152 or the use of claim 153, wherein said disorder is selected from hemoglobinopathy, sickle cell anemia, thalassemia, Fanconi anemia, Wiskott-Aldrich syndrome, a myelodysplastic disorder, an immunodeficiency disorder, a metabolic disorder, cancer, adenosine deaminase deficiency and severe combined immunodeficiency, hyper immunoglobulin M syndrome, Chediak-Higashi disease, hereditary lymphohistiocytosis, osteopetrosis, osteogenesis imperfecta, storage diseases,

thalassemia major, systemic sclerosis, systemic lupus erythematosus, multiple sclerosis, juvenile rheumatoid arthritis, an autoimmune disorder, and a neurological disorder.

155. The composition or use of claim 154, wherein said immunodeficiency disorder is a congenital immunodeficiency.

156. The composition or use of claim 154, wherein said immunodeficiency disorder is an acquired immunodeficiency.

157. The composition or use of claim 156, wherein said acquired immunodeficiency is human immunodeficiency virus or acquired immune deficiency syndrome.

158. The composition or use of claim 154, wherein said metabolic disorder is selected from the group consisting of glycogen storage diseases, mucopolysaccharidoses, Gaucher's Disease, Hurlers Disease, sphingolipidoses, and metachromatic leukodystrophy.

159. The composition or use of claim 154, wherein the cancer is a hematological cancer.

160. The composition or use of claim 154, wherein the cancer is selected from the group consisting of leukemia, lymphoma, multiple myeloma, and neuroblastoma.

161. The composition or use of claim 154, wherein the cancer is acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia, chronic lymphoid leukemia, multiple myeloma, diffuse large B-cell lymphoma, or non-Hodgkin's lymphoma.

162. The composition or use of claim 154, wherein the autoimmune disorder is selected from the group consisting of multiple sclerosis, human systemic lupus, rheumatoid arthritis, inflammatory bowel disease, treating psoriasis, Type 1 diabetes mellitus, acute disseminated encephalomyelitis, Addison's disease, alopecia universalis, ankylosing spondylitis, antiphospholipid antibody syndrome, aplastic anemia, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune inner ear disease, autoimmune lymphoproliferative syndrome, autoimmune oophoritis, Balo disease, Behcet's disease, bullous pemphigoid, cardiomyopathy, Chagas' disease, chronic fatigue immune dysfunction syndrome, chronic inflammatory demyelinating polyneuropathy, Crohn's disease, cicatricial pemphigoid, coeliac sprue-dermatitis herpetiformis, cold agglutinin disease, CREST syndrome, Degos disease, discoid lupus, dysautonomia, endometriosis, essential mixed cryoglobulinemia, fibromyalgia-fibromyositis, Goodpasture's syndrome, Grave's disease, Guillain-Barre syndrome, Hashimoto's thyroiditis, Hidradenitis suppurativa, idiopathic and/or

acute thrombocytopenic purpura, idiopathic pulmonary fibrosis, IgA neuropathy, interstitial cystitis, juvenile arthritis, Kawasaki's disease, lichen planus, Lyme disease, Meniere disease, mixed connective tissue disease, myasthenia gravis, neuromyotonia, opsoclonus myoclonus syndrome, optic neuritis, Ord's thyroiditis, pemphigus vulgaris, pernicious anemia, polychondritis, polymyositis and dermatomyositis, primary biliary cirrhosis, polyarteritis nodosa, polyglandular syndromes, polymyalgia rheumatica, primary agammaglobulinemia, Raynaud phenomenon, Reiter' s syndrome, rheumatic fever, sarcoidosis, scleroderma, Sjögren's syndrome, stiff person syndrome, Takayasu's arteritis, temporal arteritis, ulcerative colitis, uveitis, vasculitis, vitiligo, vulvodynia, and Wegener's granulomatosis.

163. The composition or use of claim 154, wherein the neurological disorder is selected from the group consisting of Parkinson's disease, Alzheimer's disease, multiple sclerosis, Amyotrophic lateral sclerosis, Huntington's disease, mild cognitive impairment, amyloidosis, AIDS-related dementia, encephalitis, stroke, head trauma, epilepsy, mood disorders, and dementia.