




[図1]




[図2]




[図3]




[図4]




[図5]



[図6]




[図7]




[図8]


(A)

投与前
(Apr-14)

(B)


投与後
(Mar-15)

[図9]




[図10]


(A)

投与前
(Jul-14)

(B)

投与後
(Nov-14)


[図11]



[図12]


(A)

投与前
(Aug-13)




(B)

投与後
(May-14)



[図13]




[図14]


(A)

投与前
(Apr-14)


(B)

投与後
(Feb-15)


[図15]




[図16]



[図17]




[図18]




[図19]


(A)

投与前
(Jun-14)

(B)

投与後
(Jun-15)

[図20]



- G1: PBS(コントロール)
- ◆ G2: ゲフィチニブ 10 mg/kg
- ▲ G3: ベスタチン 30 mg/kg
- G4: ゲフィチニブ 10 mg/kg + ベスタチン 0.1 mg/kg
- ✱ G5: ゲフィチニブ 10 mg/kg + ベスタチン 0.3 mg/kg
- G6: ゲフィチニブ 10 mg/kg + ベスタチン 1 mg/kg
- ◇ G7: ゲフィチニブ 10 mg/kg + ベスタチン 3 mg/kg
- △ G8: ゲフィチニブ 10 mg/kg + ベスタチン 10 mg/kg
- G9: ゲフィチニブ 10 mg/kg + ベスタチン 30 mg/kg