

each of R_{72} , R_{82} and R_{92} is independently hydrogen, $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, $-C_{1-10}$ heterocyclyl, $-OH$, $-CF_3$, $-C(O)R^{31}$, $-CO_2R^{31}$, $-C(=O)NR^{31}$, $-S(O)_{0.2}R^{31}$, $-C(=S)OR^{31}$, $-C(=O)SR^{31}$;

each of R_{10} and R_{14} is independently $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, $-C_{1-10}$ heterocyclyl, optionally substituted by one or more independent R_{11} substituents;

each of R_{11} , R_{12} , R_{13} and R_{15} is independently hydrogen, halogen, $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, $-C_{1-10}$ heterocyclyl, $-OH$, $-CF_3$, $-OCF_3$, $-OR^{31}$, $-NR^{31}R^{32}$, $-C(O)R^{31}$, $-CO_2R^{31}$, $-C(=O)NR^{31}$, $-NO_2$, $-CN$, $-S(O)_{0.2}R^{31}$, $-SO_2NR^{31}R^{32}$, $-NR^{31}C(=O)R^{32}$, $-NR^{31}C(=O)OR^{32}$, $-NR^{31}C(=O)NR^{32}R^{33}$, $-NR^{31}S(O)_{0.2}R^{32}$, $-C(=S)OR^{31}$, $-C(=O)SR^{31}$, $-NR^{31}C(=NR^{32})NR^{32}R^{33}$, $-NR^{31}C(=NR^{32})OR^{33}$, $-NR^{31}C(=NR^{32})SR^{33}$, $-OC(=O)OR^{33}$, $-OC(=O)NR^{31}R^{32}$, $-OC(=O)SR^{31}$, $-SC(=O)SR^{31}$, $-P(O)OR^{31}OR^{32}$, or $-SC(=O)NR^{31}NR^{32}$;


each of R^{31} , R^{32} , R^{33} and R^{34} is independently hydrogen, halogen, $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, $-C_{1-10}$ heterocyclyl, or wherein R^{31} together with R^{32} form a heterocyclic ring;

wherein ring A comprises one or more heteroatoms selected from N, O, or S;

wherein if X_7 is O or X_2-X_3 is $R_1C=CR_3$, ring A comprises at least two heteroatoms selected from N, O, or S; and

wherein if X_3 is O or N, or X_2-X_3 is $R_1C=N$, at least one of X_7 or X_9 is not N.

[0006a] In one embodiment, there is provided a compound of Formula II-E:



or a pharmaceutically acceptable salt thereof; wherein:

R_1 is $-C_{1-10}$ heterocyclyl, $-C_{1-10}$ heterocyclyl- C_{1-10} alkyl, or $-C_{1-10}$ heterocyclyl- C_{3-10} aryl, each of which is unsubstituted or substituted by one or more independent R_{10} or R_{11} substituents;

R_{21} is $-L-C_{1-10}$ heteroalkyl, $-L-C_{3-10}$ aryl, $-L-C_{1-10}$ hetaryl, $-L-C_{3-10}$ cycloalkyl, or $-L-C_{1-10}$ heterocyclyl, each of which is unsubstituted or substituted by one or more independent R_{12} substituents;

L is a bond, $-O-$, $-N(R^{31})-$, $-S(O)_{0.2}-$, $-C(=O)-$, $-C(=O)O-$, $-OC(=O)-$, $-C(=O)N(R^{31})-$, $-N(R^{31})C(=O)-$, $-NR^{31}C(=O)O-$, $-NR^{31}C(=O)NR^{32}-$, $-NR^{31}S(O)_{0.2}-$, $-S(O)_{0.2}N(R^{31})-$, $-C(=S)O-$, $-C(=O)S-$, $-NR^{31}C(=NR^{32})NR^{32}-$, $-NR^{31}C(=NR^{32})O-$, $-NR^{31}C(=NR^{32})S-$, $-OC(=O)O-$, $-OC(=O)NR^{31}-$, $-OC(=O)S-$, $-SC(=O)S-$, $-P(O)OR^{31}O-$, or $-SC(=O)NR^{31}-$;

R_{72} is hydrogen;


R_{10} is $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, or $-C_{1-10}$ heterocyclyl, each of which is unsubstituted or substituted by one or more independent R_{11} substituents;

R_{11} and R_{12} are independently hydrogen, halogen, $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, $-C_{1-10}$ heterocyclyl, $-OH$, $-CF_3$, $-OCF_3$, $-OR^{31}$, $-NR^{31}R^{32}$, $-C(O)R^{31}$, $-CO_2R^{31}$, $-C(=O)NR^{31}$, $-NO_2$, $-CN$, $-S(O)_{0.2}R^{31}$, $-SO_2NR^{31}R^{32}$, $-NR^{31}C(=O)R^{32}$, $-NR^{31}C(=O)OR^{32}$, $-NR^{31}C(=O)NR^{32}R^{33}$, $-NR^{31}S(O)_{0.2}R^{32}$, $-C(=S)OR^{31}$, $-C(=O)SR^{31}$, $-NR^{31}C(=NR^{32})NR^{32}R^{33}$, $-NR^{31}C(=NR^{32})OR^{33}$, $-NR^{31}C(=NR^{32})SR^{33}$, $-OC(=O)OR^{33}$, $-OC(=O)NR^{31}R^{32}$, $-OC(=O)SR^{31}$, $-SC(=O)SR^{31}$, $-P(O)OR^{31}OR^{32}$, or $-SC(=O)NR^{31}NR^{32}$; each of R^{31} , R^{32} , and R^{33} is independently hydrogen, halogen, $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, or $-C_{1-10}$ heterocyclyl, or R^{31} together with R^{32} form a heterocyclic ring.

[0006b] In one embodiment, there is provided use of the compound as described herein in the manufacture of a medicament for inhibiting activity of a protein kinase present in a cell.


[0006c] In one embodiment, there is provided use of the compound as described herein in the manufacture of a medicament for treatment of cancer, optionally wherein the cancer is selected from the group consisting of breast cancer, pancreatic cancer, non-small cell lung cancer (NSCLC), thyroid cancer, seminomas, melanoma, bladder cancer, liver cancer, kidney cancer, myelodysplastic syndrome (MDS), acute myelogenous leukemia (AML), and colorectal cancer.

[0007] In some embodiments, the invention provides a compound of Formula I-A:




or a pharmaceutically acceptable salt or prodrug thereof.

[0008] In some embodiments, the invention provides a compound of Formula I-B:



or a pharmaceutically acceptable salt or prodrug thereof.

[0009] In some cases, the invention provides a compound of Formula II:




or a pharmaceutically acceptable salt or prodrug thereof.

[0010] In some embodiments, X_1 is $C=O$. In some embodiments, X_2 is NR_1 . In some embodiments, X_5 is N and X_6 is C. In other embodiments, X_5 is C and X_6 is C. In some embodiments, X_9 is CR_{21} .

CLAIMS

1. A compound of Formula II-E:



or a pharmaceutically acceptable salt thereof; wherein:

R_1 is $-C_{1-10}$ heterocyclyl, $-C_{1-10}$ heterocyclyl- C_{1-10} alkyl, or $-C_{1-10}$ heterocyclyl- C_{3-10} aryl, each of which is unsubstituted or substituted by one or more independent R_{10} or R_{11} substituents;

R_{21} is $-L-C_{1-10}$ heteroalkyl, $-L-C_{3-10}$ aryl, $-L-C_{1-10}$ hetaryl, $-L-C_{3-10}$ cycloalkyl, or $-L-C_{1-10}$ heterocyclyl, each of which is unsubstituted or substituted by one or more independent R_{12} substituents;

L is a bond, $-O-$, $-N(R^{31})-$, $-S(O)_{0-2}-$, $-C(=O)-$, $-C(=O)O-$, $-OC(=O)-$, $-C(=O)N(R^{31})-$, $-N(R^{31})C(=O)-$, $-NR^{31}C(=O)O-$, $-NR^{31}C(=O)NR^{32}-$, $-NR^{31}S(O)_{0-2}-$, $-S(O)_{0-2}N(R^{31})-$, $-C(=S)O-$, $-C(=O)S-$, $-NR^{31}C(=NR^{32})NR^{32}-$, $-NR^{31}C(=NR^{32})O-$, $-NR^{31}C(=NR^{32})S-$, $-OC(=O)O-$, $-OC(=O)NR^{31}-$, $-OC(=O)S-$, $-SC(=O)S-$, $-P(O)OR^{31}O-$, or $-SC(=O)NR^{31}-$;

R₇₂ is hydrogen;

R₁₀ is -C₁₋₁₀ alkyl, -C₂₋₁₀ alkenyl, -C₂₋₁₀ alkynyl, -C₁₋₁₀ heteroalkyl, -C₃₋₁₀ aryl, -C₁₋₁₀ hetaryl, -C₃₋₁₀ cycloalkyl, or -C₁₋₁₀ heterocyclyl, each of which is unsubstituted or substituted by one or more independent R₁₁ substituents;

R₁₁ and R₁₂ are independently hydrogen, halogen, -C₁₋₁₀ alkyl, -C₂₋₁₀ alkenyl, -C₂₋₁₀ alkynyl, -C₁₋₁₀ heteroalkyl, -C₃₋₁₀ aryl, -C₁₋₁₀ hetaryl, -C₃₋₁₀ cycloalkyl, -C₁₋₁₀ heterocyclyl, -OH, -CF₃, -OCF₃, -OR³¹, -NR³¹R³², -C(O)R³¹, -CO₂R³¹, -C(=O)NR³¹, -NO₂, -CN, -S(O)₀₋₂R³¹, -SO₂NR³¹R³², -NR³¹C(=O)R³², -NR³¹C(=O)OR³², -NR³¹C(=O)NR³²R³³, -NR³¹S(O)₀₋₂R³², -C(=S)OR³¹, -C(=O)SR³¹, -NR³¹C(=NR³²)NR³²R³³, -NR³¹C(=NR³²)OR³³, -NR³¹C(=NR³²)SR³³, -OC(=O)OR³³, -OC(=O)NR³¹R³², -OC(=O)SR³¹, -SC(=O)SR³¹, -P(O)OR³¹OR³², or -SC(=O)NR³¹NR³²;

each of R³¹, R³², and R³³ is independently hydrogen, halogen, -C₁₋₁₀ alkyl, -C₂₋₁₀ alkenyl, -C₂₋₁₀ alkynyl, -C₁₋₁₀ heteroalkyl, -C₃₋₁₀ aryl, -C₁₋₁₀ hetaryl, -C₃₋₁₀ cycloalkyl, or -C₁₋₁₀ heterocyclyl, or R³¹ together with R³² form a heterocyclic ring.

2. The compound of claim 1, wherein R₂₁ is selected from the group consisting of -L-C₁₋₁₀ heteroalkyl, -L-C₃₋₁₀ aryl, -L-C₁₋₁₀ hetaryl, -L-C₃₋₁₀ cycloalkyl, and -L-C₁₋₁₀ heterocyclyl, each of which is unsubstituted or substituted by one or more independent R₁₂ substituents; and wherein L is a bond.

3. The compound of claim 2, wherein R₂₁ is -L-C₁₋₁₀ hetaryl unsubstituted or substituted by one or more independent R₁₂ substituents; and wherein L is a bond.

4. The compound of claim 3, wherein the C₁₋₁₀ hetaryl of R₂₁ comprises one or more nitrogen atoms.


5. The compound of claim 4, wherein the C₁₋₁₀ hetaryl of R₂₁ is selected from the group consisting of pyrazolyl, pyridinyl, pyrazinyl, pyrimidinyl, and pyridazinyl.

6. The compound of claim 5, wherein the C₁₋₁₀ hetaryl of R₂₁ is unsubstituted.

7. The compound of claim 5, wherein the C₁₋₁₀ hetaryl or R₂₁ is substituted with one, two, or three independent R₁₂ substituents.

8. The compound of any one of claims 2-5 or 7, wherein each R₁₂ substituent is independently selected from the group consisting of -C₁₋₁₀ alkyl, -C₂₋₁₀ alkenyl, -C₂₋₁₀ alkynyl, -C₁₋₁₀ heteroalkyl, -C₃₋₁₀ aryl, -C₁₋₁₀ hetaryl, -C₃₋₁₀ cycloalkyl, -C₁₋₁₀ heterocyclyl, -OH, -CF₃, -OCF₃, and -OR³¹; wherein each R³¹ is independently hydrogen or -C₁₋₁₀ alkyl.

9. The compound of claim 8, wherein each R₁₂ substituent is independently selected from the group consisting -Me, -Et, -i-Pr, -n-Pr, -OH, -OMe, -OEt, and -OPr.



11. Use of the compound of any one of claims 1-10 in the manufacture of a medicament for inhibiting activity of a protein kinase present in a cell.

12. The use of claim 11, wherein the protein kinase is ERK.

13. Use of the compound of any one of claims 1-10 in the manufacture of a medicament for treatment of cancer, optionally wherein the cancer is selected from the group consisting of breast cancer, pancreatic cancer, non-small cell lung cancer (NSCLC), thyroid cancer, seminomas, melanoma, bladder cancer, liver cancer, kidney cancer, myelodysplastic syndrome (MDS), acute myelogenous leukemia (AML), and colorectal cancer.

14. A pharmaceutical composition comprising the compound of any one of claims 1-10 and a pharmaceutically acceptable excipient, optionally wherein the composition is formulated in an oral dosage.

15. The compound of claim 1, wherein the compound is selected from:

