REPLACED SHEETS SG
☑ Description : paged ...1.... to paged ...31....
☑ Claim(s) : paged ...32... to paged ...34....
☐ Drawing(s) : paged ..... to paged .....
☑ Abstract : paged ...41... to paged .....
DATE OF RECEIPT ..... 1 .....

9607958-7
IPOS

RTM Copy

WO 93/08470

This Sheet Has Been Replaced

Method and composition for determining the immunological activity of bioactive substances.

The present invention relates to a method and composition for assaying certain bioactive substances and determining their immunological activity with respect to their ability to induce production of cytokines. Cytokines, such as interferons (IFNs) and tumor necrosis factors (TNFs), are hormone-like proteins which play an important role in virtually all immunological reactions, as well as in the regulatory process responsible for the maintenance of homeostasis. The production of such cytokines can be induced by certain substances which, on account of their bioactive and immunomodulating activity, are useful in the therapy of immunodeficiencies and related diseases.

There is, of course, a substantial need of methods for properly and easily assaying the immunological activity of such bioactive substances. It is the object of the present invention to provide such a method.

The method according to the present invention allows to determine whether a tested substance is able to induce production of different cytokines and permits a quick and efficient check of the properties of certain substances, namely their immunological activity, at each stage of production, separation, purification and formulation into final compositions. The present invention provides also a composition and kits suitable for carrying out said method.

Tests used until now to determine immunological activity are numerous but difficult to carry out. When a new therapeutical composition having an immunological effect is clinically tested, its immunological effectiveness is evaluated by monitoring the numerous, well known and analytically determinable features of immunological systems employing standard methods. Once the activity is proven,

This Sheet Has Been Replaced

there emerges the necessity for representative single tests enabling a quick check of the status of a patient in terms of his or her immunological response.

In clinical studies, evaluating IFN serum levels may be difficult, unprecise and misleading. Interferons are often produced locally in various tissues, they are short-lived, their action is paracrine (in the immediate vicinity only) and they are strongly bound to cellular receptors and carrier proteins.

On the other hand, technological processes for manufacturing, purification and/or separation of immunologically active substances such as in particular, extracts from raw peat are usually multi-step processes, and there is always the need for establishing a reliable and fast test enabling a thorough control of the production. Moreover, since the substances are frequently of a complex nature, as it is the case with peat extracts, it is important to have a similar test for determining the activity of individual fractions of such substances. It is also necessary to find out a microscale testing method, since the products to be analysed are very costly.

The following four basic findings allowed to solve the above described problems.

1. Some immunoactive peat extracts induce the production of cytokines, such as interferons and tumor necrosis factors, in in-vitro cultures of peripheral blood leukocytes, the induction being dependant on the dose.

2. Interferon-β (and TNF-α) is produced in significantly higher quantities by BALB/c mice resident peritoneal cells (RPC) when treated with immunoactive peat extracts as compared to a spontaneous release of these cytokines in non-treated samples.

3. Certain peat extracts, when tested in-vitro as mentioned above show a synergistic effect in combination with known immunomodulators, such as organoselenium compounds, the p-chlorophenylamide of 3-methyl-5-benzoylamino-isothiazol-4-carboxylic acid and indomethacin. This allows the use of much smaller quantities of immunologically active substances to show their activity.

4. PBL of healthy volunteers treated with TTP (a certain peat-derived product) administered orally at a dosage of 5 mg/day, tested for induction of certain cytokines, IFN-α, IFN-γ and TNF-α, lose the ability to respond to induction of the cytokines with TTP solution after prolonged, uninterrupted administration of TTP orally and regain this ability after approximately two weeks of hiatus.


The method according to the present invention comprises the steps of treating a human peripheral blood leukocyte (PBL) culture or a suspension of BALB/c mice resident peritoneal cells (RPC), with a solution of the substance to be tested in order to induce production of cytokines, and then determining said cytokines according to standard identification methods.

The peripheral blood leukocyte (PBL) culture used according to the first variant or embodiment of the instant method is a short term culture prepared from fresh leukocytes of healthy humans with a nutrient medium suitable for tissue culture. The preferred density of a ready-for-use culture is about 8×10^6 leukocytes/ml. The solution to be tested is preferably used in a concentration of 0,1 - 200 µg/ml. The method is, in particular, suitable for testing peat extracts.


This Sheet Has Been Replaced

A micro-scale determination is possible when the solution to be tested, e.g. a peat extract or a fraction thereof, is admixed with an immunomodulating agent, such as a non-steroidal anti-inflammatory drug, preferably a substance selected from the group comprising organoselenium compounds, such as ebselen, certain isothiazole derivatives, such as the p-chlorophenylamide of 3-methyl-5-benzoylamino-isothiazole-4-carboxylic acid, and indomethacin, as well as analogs, homologs and metabolites of such substances.


Examples of organoselenium compounds which can be used for this purpose are compounds of the following formulae 1-3; the compound of formula 2 (Ebselen) being preferred:



(1) Bis-[2-(N-phenylcarboxamido)]-phenyldiselenide




(2) 2-Phenyl-1,2-benzisoselenazol-3(2H)-one (Ebselen PZ 51)



(3) Bis-[2-(N-(2-pyridyl)carboxamido)]-phenyldiselenide

This Sheet Has Been Replaced

Examples of isothiazole derivatives which can be used for the above purpose are compounds of the general formula I



(I)

wherein R_1 is halophenyl, preferably chlorophenyl, R_2 is phenyl and R_3 is lower alkyl, preferably methyl. The halogen atom in the halophenyl group, chlorophenyl group R_1 is preferably in p-position of the phenyl ring.

The preferred compound of the above formula I is the p-chlorophenylamide of 3-methyl-5-benzoylamino-isothiazole-4-carboxylic acid, i.e. the compound of formula I wherein R_1 is p-chlorophenyl, R_2 is phenyl and R_3 is methyl. For brevity sake, this compound will be referred to below as 'Compound ITCL'. There exists a Trade Mark registration "VRATIZOLIN^R" for this compound in Poland. Its synthesis and properties are described in Arch. Immunol. et Ther. Exp. 1973, 21, 891.

All the immunomodulating agents referred to above for being useful for micro-scale determinations, i.e. the organoselenium compounds, the isothiazole derivatives and indomethacin, have in common that they are non-steroidal anti-inflammatories which inhibit the synthesis of prostaglandins. They amplify the test results in the method according to the present invention. For this reason, they are particularly useful for micro-determinations. Amplification of the results may be 5-20 times.


Out of all the various non-steroidal anti-inflammatory compounds useful for the purpose of amplification, seleno organic compound (1, 2 and 3) and compound ITCL, but especially indomethacin, are the preferred ones.

In the instant invention, different antisera, which recognise individual induced cytokines, are employed. These sera are known and used for testing and standardisation of different cytokines. Identification of the induced cytokines should take place at the moment of completion of the cytokine formation and prior to its proteolysis. Among different cytokines induced, some are formed slowly and remain stable in the solution, as for example interferon γ, while others are formed rapidly and are susceptible to proteolysis, as for example tumor necrosis factors.

The method as described above may be employed also for the determination of an immunological response of a human individual to a certain immunoactive substance, provided that such immunological response is not stimulated by other factors, as for example by viral or bacterial infections. It may be applied for monitoring of the drug response in individual patients, for the determination of the optimal dose of a therapeutic cytokine inducer, such as TTP, as well as for the establishment of the effective scheme of the drug administration. The method is based on the assessment of the hyporeactivity state to IFN induction.

Thus, the present invention also relates to a method for determining the immunological response of a human individual to a therapy using a cytokine inducing immunomodulator substance. Said method is characterised in that a peripheral blood leukocyte (PBL) culture of the human individual treated with such immunomodulator substance, is treated in defined time intervals with a solution of the substance administered in order to induce production of cytokines, which are then determined according to standard methods (including ELISA assays of cytokines), in order to determine the moment of development of hyporeactivity to the substance after prolonged administration of the substance and the moment of regaining the ability to respond to an additional dose of the substance after a certain period of hiatus.

This Sheet Has Been Replaced



The time intervals are preferably about 7-14 days.

In a preferred embodiment, the above method is applied in the course of a therapy using a peat extract as cytokine inducing immunomodulator substance.

According to a particularly preferred embodiment of the invention, this method is applied in the course of a therapy using TTP as cytokine inducing immunomodulator substance, said TTP being, inter alia, the subject of PCT application No. PCT/EP92/00491, wherein said TTP, its characteristics and its production are described in detail.

In view of the above, the instant invention also relates to TTP, whenever used, tested or determined according to the methods of the instant invention.

The present invention also provides, in its second variant, for a testing method which is suitable for quick monitoring of technological processes for the production, purification, separation or the like of the immmunoactive substances. For instance, when a suspension of BALB/c mice resident peritoneal cells (RPC) is treated with the solution to be tested, mice interferon-β as well as tumor necrosis factor are induced. Both may be detected and qualitatively determined according to standard identification methods. In this method, a suspension of fresh mouse resident peritoneal cells (RPC), containing approx. 1 x 10^6 cells/ml is used. The solution to be tested is prepared in a nutrient medium Eagle, or RPMI-1640 with the addition of 10% of fetal calf serum. The results obtained are evaluated against a calibration curve prepared with a model substance. As a model substance, a sample of the same immunologically active substance tested in a traditional way may be used. Again, the method is particularly suitable for monitoring the process for obtaining peat extracts as well as analysing their fractions.

The instant method is very simple and efficient in comparison with other known methods, because it is based on a direct induction of interferon β instead - as is the case in a number of conventional tests - on a secondary effect of formulation of antibodies. Until now, it was not possible to find any similar method due to the fact that the immunosystem of mice significantly differs from the human immunosystem and tests recognised as indicative for human immunologic response - such as the presence of interferons in the tissues of spleen or lymphonodes etc. - do not work with the best laboratory test animals, i.e. with mice of BALB/c type.

It was now found that a clear immunological response of the BALB/c mice can be obtained when resident peritoneal cells (RPC) are chosen as a biological material for testing, fresh suspensions of such cells being used. For each test it is sufficient to sacrifice just three animals. Results are obtained within several hours. The biological material may also be cultured and a corresponding line of macrophages be established.

The method according to the present invention employs the use of biological material, namely a peripheral blood leukocyte (PBL) culture or a suspension of BALB/c mice resident peritoneal cells (RPC), according to the information given above. Presence and concentration of each cytokine is determined by methods for testing and standardisation of interferons known from "Methods in Enzymology", 1986, vol.119, part C: "Interferon", edited by Sidney Pestka. Determination of the whole spectrum of cytokines induced identifies the immunological status of a patient.

For the proper evaluation of the results, it is essential that the biological material for carrying out both variants of the method according to the present invention is selected

This Sheet Has Been Replaced

from the proper donors. In case of human leukocytes, samples taken from individuals who show a hypo-reactivity (rare cases), i.e. a low level of immunological response, should be excluded from the evaluation. In case of the BALB/c mice RPC suspension, samples taken from additionally stimulated animals (for example infected ones) should also be excluded from the evaluation. In case of the BALB/c mice RPC suspensions, samples should not be taken from too young or from too old animals, as there is an age-dependent deficiency in cytokine production in such mice. 5-8 weeks old healthy animals should be selected. Samples showing very high spontaneous production of cytokines should be eliminated, especially when immunoregulatory substances are tested, since no potentiation of cytokine induction or sometimes even reduction of such induction may be observed.

That embodiment of the method according to the present invention, wherein amplification of the results is achieved by admixing a non-steroidal anti-inflammatory drug (one of the substances referred to above, such as compound ITCL or seleno organic compound 1, 2 and 3, but preferably indomethacin) to the solution to be tested, is a microscale test, particularly suited for determining the induced TNF and IFN activity. This test is economic, fast, applicable for assaying a great number of samples simultaneously and suitable for standardization and, at least partial, automatization.

The two variants of the method according to the present invention are described by way of example in more detail below:

Preparation of the solution of the substance to be tested: The substance to be tested, e.g. a peat extract, is dissolved in sterile bidistilled water at a concentration of 10 mg/ml. The samples are sterilized by filtration through

This Sheet Has Been Replaced

0.45 µm, 600 kPa max Millipore(R), antibacterial filters. Next, solutions are made in a complete RPMI-1640 medium containing heat-inactivated fetal bovine serum (FBS).

A) The PBL variant:

Cytokine induction

Buffy coats from healthy blood donors may be obtained from the regional transfusion center. Alternatively, peripheral blood leukocytes (PBL) may be isolated from heparinized venous blood of healthy volunteers by Ficoll-Hypaque(R) (or Histopaque(R)) density gradient (g = 1.077) centrifugation, with subsequent twice washing of the cells. The erythrocytes were lysed with NH_4Cl treatment according to Cantell et al. (Cantell, K., Hirvonen, S., Kauppinen, H.L.: Production and Partial Purification of Human Immune Interferon. Meth. Enzymol. 119, 54, 1988). The leukocytes from a single donor containing approximately $8x10^6$ leukocytes/ml in RPMI 1640 medium supplemented with 10% fetal bovine serum (FBS), L-glutamine and antibiotics were used. All lots of FBS were pretested. Only non-mitogenic FBS for PBL cultures was used. The cytokine inducers were added to 200-1000 µl volumes of the cultures. The reference cytokine inducers were phytohemagglutinin (PHA) (Pharmacia Fine Chemicals, Sweden or Sigma, USA). The induced cultures of PBL were incubated in an atmosphere of 5% CO_2 in air at 37°C for 20 h and centrifuged. Supernatants were stored at 4°C and assayed for IFN activity within one week. Supernatants for TNF activity determination have to be stored at -90°C or in liquid nitrogen to avoid inactivation due to proteolysis.

Interferon assay

Confluent monolayers of A549 cells were prepared in microplates in Dulbecco-modified Minimum Essential Medium (DMEM) with 10% FBS, L-glutamine, and antibiotics (penicillin 100 units/ml and streptomycin 100 µg/ml). IFN

This Sheet Has Been Replaced

samples diluted in the plates were added to the cell monolayer and incubated at 37°C for 20 h in 5% CO_2 in air. The cells were then washed and challenged with encephalomyocarditis virus (EMCV). The titer of IFN was defined as the dilution of IFN sample that reduced virus cytopathogenic effect by 50% after 48 h of incubation. The MTT (3-[4.5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) method (Hansen, M.B., Nielsen, S.E. and Berg. K.: Re-examination and Further Development of a Precise and Rapid Dye Method for Measuring Cell Growth/Cell Kill. J. Immunol. Meth. 1989, 119, 203-210) to measure the cell kill in the ELISA scanner was also used. Laboratory standards of IFN have to be included in all assays e.g. recombinant human IFN-γ (specific activity $2x10^6$ units/mg), the natural human leukocyte IFN-α ($3x10^6$ IU/ml) and IFN-γ ($2x10^6$ IU/ml).

TNF assay

The cytotoxic activity of TNF was measured in L_{929} cells according to Flick and Gifford (Flick, D.A., Gifford, G.E.: Comparison of in Vitro Cell Cytotoxic Assays for Tumor Necrosis Factor. J. Immunol. Meth. 68, 1667, 1984). The samples and actinomycin D (5µg/ml) solution were added to monolayer cultures of the cells. After incubation at 37°C for 20 h, cytotoxic effects of TNF were determined either by the microscopic examination of the cultures or by using the MTT method. The amount causing approximately 50% destruction of the cell cultures was defined as one unit of TNF activity. Comparison with a preparation of TNF-α (Genentech Inc., USA) showed that 1 unit in the assays was equal to 100 - 200 pg/ml TNF.

Cytokine neutralization assays

The cytokines produced by PBL treated with the examined preparations may be identified by neutralization assays with the specific antibodies (Inglot et al., Organoselenides as

This Sheet Has Been Replaced

potential immunostimulants and inducers of interferon gamma and other cytokines in human peripheral blood leukocytes, Experientia 1990, 46, 308-311). Furthermore, various ELISA kits for the determination of the immuno-activity of a defined cytokine may be applied.

Comment

In the supernatants obtained from the cultured lymphoid cells, other than TNF or IFN cytokines may be found and identified by other standard methods, e.g. Interleukins (IL-1-10), GM-CSF, TGF-β etc.

B) The RPC variant: Cytokine induction.

Mouse resident peritoneal cells (RPC) are obtained from approximately 6 weeks old BALB/c mice killed with ether, by injecting into the peritoneal cavity 5 ml of complete RPMI-1640 medium at room temperature. The washing medium containing RPC is collected into ice-cooled 50 ml centrifuge tubes. The RPC are neither washed nor centrifuged. The cells are counted in a Bürker hemocytometer and suspended to a density of 1-1.5x10^6 cells/ml. Such cell density is required for all of the assays measuring the effects of various concentrations of the preparation (from 0.1-500 µg/ml). The negative and positive controls are included in all of the tests. The negative control measures the spontaneous release of TNF or IFN by RPC incubated with the complete RPMI-1640 medium without the inducers, whereas the positive control shows the effect of a standard lipo-polysaccharide inducer (LPS from E. coll 055:85, Sigma 1-10 µg/ml). All of the cultures are incubated at 26°C for 20 h. Thereafter, the cultures are centrifuged at 1000 rpm for 10 min and supernatants are collected. Using an automatic pipette, the supernatants are diluted from 1:2 to 1:256.

This Sheet Has Been Replaced



Bioassays of the cytokines

To determine TNF activity in supernatants from RPC cultures, the 20 h old monolayer of L_{929} cell cultures are used.

The mouse fibroblast-like L_{929} cells, $2x10^4$ cells per well in 100 µl of complete medium, are seeded in 96-well flat bottom plates and incubated for 20 h at 37°C in order to obtain a monolayer.

The transfer of supernatants must be very accurate. The incubation of the L_{929} cultures is carried on for 24 h at 37°C in an atmosphere of 5% CO_2 in air.

Evaluations

1) Reading of the cytotoxic effects under the reverse microscope.
2) MTT test (3-[4.5-dimethylthiazol-2-yl]-2,5-diphenyl-tetrazolium bromide, Sigma).

To measure cell killing with L_{929} cell cultures, 25 µl of MTT dye solution at a concentration of 5 mg/ml are added into every well of the microplates.

Next, the plates are incubated for 2 h at 37°C in an atmosphere of 5% CO_2 in air.

Thereafter, 100 µl of the solvent solution containing 45 ml of dimethylformamide, 13.5 g of SDS (sodium dodecyl sulphate), and 55 ml of distilled water, are added to every well.

Incubation is carried on for 12 hrs at 37°C in an CO_2 incubator. The results of MTT color test are read in a Multiskan 340/CC reader (Labsystem) using a 570 nm filter. Reference recombinant TNF-α has to be used.

This Sheet Has Been Replaced



Interferon bioassay

IFN is assayed by inhibition of the cytopathic effect caused by mouse encephalomyocarditis virus (EMCV in mouse L_{929} cells, reference Mu IFN α/β, standard from National Institute of Health Bethesda, MD, USA is included). The cytopathic effect is observed under a reversed microscope and also it is measured by the MTT method as described below:

MTT method according to Berg et al.
Berg K., Hansen M.B. and Nielsen S.E. (1990): A new sensitive bioassay for precise quantification of interferon activity as measured via the mitichondrial dehydrogenase function in cells (MTT-method). APMIS, 98, 156-162.

MTT (3/4,5- dimethyl-thiazol-2-yl/2,5-diphenyl tetrazolium bromide, Sigma) is diluted in PBS at a concentration of 5mg/ml. To measure cell killing with L_{929} cultures, 25 µl of MTT dye solution at a concentration of 5 mg/ml is added to every well of the microplates. The control reference sample of TNF or IFN is included. Next, the plates are incubated for 2 hrs at 37° C in an atmosphere of 5% CO_2 in air. Thereafter, 100 µl of a solution containing 45 ml dimethylformamide, 13,5 g of SDS (sodium dodecyl sulfate) and 55 ml of distilled water are added to every well. After overnight incubation at 37° C the optical densities at 570 nm are measured, using a microplate reader Start Fax Awareness Technology Inc. 2100, employing the extraction buffer as the blank.

Both TNF-α, IFN-β, and other cytokines, may be also assayed using commercial ELISA kits.

A composition for carrying out the method of the present invention is essentially characterized in that it contains double processed tissue culture water, a culture medium and

This Sheet Has Been Replaced



sera completing the culture medium as well as the human PBL or mice RPC.

A diagnostic kit (cytokine inducer kit) for use in the PBL variant of the present method (for 10-20 manual assays) may contain the following items:

1) Double processed tissue culture water, 100 ml
2) Histopaque(R) or Ficoll-Hypaque(R) - 1077, (density gradient medium), 100 ml
3) RPMI-1640-Medium, 2x100 ml (with sodium bicarbonate and L-glutamine, sterile filtered, endotoxin tested)
4) Fetal bovine serum (FBS), 20 ml, low in endotoxin and hemoglobin
5) Lectin from Phaseolus vulgaris 0.5 mg (Phytohemagglutinin - PHA-P)
6) Round bottom tubes with caps, polypropylene sterile for tissue cultures 17x100 mm, capacity 14 ml, 25 per bag
7) Tissue culture plates with lids, 2 plates, 96 well, flatbottom, sterile.

Reagents available from: Sigma Chemical Co., St. Louis, Mo., USA or other companies.

Such a PBL kit may be used as so-called second phase screening system, to confirm results obtained with RPC cells or sometimes as a direct screening system for several substances which may be inactive as immunomodulators in rodents, e.g. organoselenium compounds.

A diagnostic kit (cytokine inducer kit) for use in the RPC variant of the present method (for 10-20 manual assays) may contain the following items:

1) Double processed tissue culture water, 100 ml
2) RPMI-1640-medium, 2x100 ml, (with sodium bicarbonate and L-glutamine), sterile, filtered, endotoxin tested)

This Sheet Has Been Replaced

3) Fetal bovine serum (FBS), 20 ml, low in endotoxin and hemoglobin
4) Lipopolysaccharide (LPS) 0.5 mg, from E. coli 0127:B8
5) Round bottom tubes with caps, sterile, 25 per bag, Polypropylene, for tissue culture, 12 x 75 mm, 6 ml capacity
6) Tissue culture plates with lids, 5 plates, 96 well, flat bottom, sterile.

Reagents are available from Sigma Chemical Co. St. Louis, Mo., USA or other companies.

A composition for carrying out the method of the present invention involving amplification of the results, which composition is particularly suited for micro-scale tests, is essentially characterised in that it contains on the one hand a PBL culture or an RPC suspension, and on the other hand a non-steroidal anti-inflammatory drug, preferably a substance selected from the group of organoselenium compounds, such as ebselen, indomethacin or Compound ITCL, and derivatives, analogs, homologs and metabolites of these compounds, most preferably indomethacin.

The following specific Examples illustrate the present invention. It is understood that they do not limit the scope of the invention in any way.

Example 1:

Individual solutions of RPC were obtained from 36 female, 7-8 weeks old BALB/c mice killed with ether, by injecting into the peritoneal cavity 5 ml of Eagle's minimum essential medium (EMEM), supplemented with 10% heat-inactivated calf serum. Washings from the individual mice contained 1-2 x 10^6 cells/ml. Suspensions of the cells from each mouse were divided into two samples distributed to two tubes. One

This Sheet Has Been Replaced

sample was treated with 100 µg TTP/ml and the other served as a negative control showing spontaneous release of cytokines.

All cultures were incubated at 26°C for 20h and thereafter they were centrifugated at 1000 rpm for 10 min. The supernatants were collected and assayed for IFN-β activities in a bioassay using a micromethod of inhibition of cytopathic effect caused by EMCV (encephalomyocarditis virus) in mouse L_{929} cells. In each assay an internal laboratory standard was included which had been calibrated against the international reference preparation of Mu IFN, G002-904-511, NIH, Bethesda, USA. The cytopathic effect was measured by the MTT method according to Berg et al, APMIS, 98, 156-162. The results obtained were as follows: for the controls the IFN-β level was 1-8 U/ml average being <2, for the samples treated with TTP, the IFN-β level was 4-127 U/ml, the average being 16 U/ml. The results are graphically presented on the accompanying drawing Fig.1 (Comparison of IFN-β production by RPC treated with TTP and nontreated. 1-limit of detection, 2-mediana. According to Mediana test P=0,0000. Every point shows the level of IFN in RPC isolated from individual mice).

Example 2:

The procedure as described in Example 1 was repeated with a group of 7 mice, however, instead of dividing each individual mouse RPC suspension into two samples, only one suspension was divided into four samples. The samples were distributed to 4 tubes (1 ml in each). One of the samples was a negative control and the remaining three were treated with 100, 10 and 1 µg TTP/ml to determine a TTP dose-dependance of the IFN-β release. After incubation of the cultures for 24h at 26°C the tubes were centrifugated. In the collected supernatants IFN-β was determined as described in Example 1. The results obtained are presented on the

This Sheet Has Been Replaced

accompanying drawing Fig.2 (Dependence of IFN production on TTP concentration). Each curve relates to the results observed with a single mouse derived RPC suspension.

Example 3:

The procedure as described in Example 1 was followed, except that TNF-α was determined instead of IFN-β.

Supernatants from the RPC cultures, non-treated and treated with 100 µg TTP/ml were assayed for TNF activity in a bioassay. The mouse fibroblast-like L_{929} cells (4×10^4 cells per well in 100 µl of complete EMEM) were seeded in 96-well flat bottom plates (Falcon, Linbro, Flow) and incubated for 4 h at 37°C in an atmosphere of 5% CO_2 in air. The samples were diluted in EMEM with actinomycin D (end concentration 2,4 µg/ml) in the additional plate. Next, the culture medium above the L_{929} cells monolayer was removed and the prepared dilutions were transferred to this culture using a multichannel pipette. After incubation for 20 h at 37°C, the cultures were assayed for cell kill by the MTT method as described above. The results obtained were as follows: for negative controls the TNF-α level was within a range of 1-256 U/ml the average being 8 U/ml, and for the samples treated with TTP the TNF-α level was 2-256 also, but the average was 64 U/ml. The results obtained are shown on the accompanying drawing Fig.3 (Comparison of TNF-α production by RPC treated with TTP and nontreated. 1-limit of detection, 2-mediana. According to Mediana test P=0,0030. Every point shows the level of TNF in RPC isolated from individual mice).

Example 4:

The procedure as described in Example 2 was followed, except that instead of IFN-β, TNF-α was determined essentially as described in Example 3 above.

This Sheet Has Been Replaced



The results obtained are shown on the accompanying drawing Fig.4 (Dependence of TNF production on TTP concentration).

Example 5:

A. Biological material. Over 115 buffy coats from individual healthy blood donors, obtained from Wroclaw Regional Transfusion Center, have been used for the assays. The characteristics of the donors are given in a Table 1, below.

Table 1: Characteristics of donors of blood used as a source of PBL

Age (years)	No.	Male	Female	Donation	
				Multiple	Single
21-30	24	22	2	24	-
31-40	45	44	1	45	-
41-59	46	44	2	40	6
Total	115	110	5	109	6
%	100	96	4	95	5

The majority of them were young males who donated blood many times. Considerable variations in the response of PBL from the individual donors have been observed and are reported in Table 2 below.

As it results from Table 2, 10-30% of PBL cultures may not respond to a 100 µg/ml TTP dose, whereas only 7% cannot be stimulated by a 10 µg/ml PHA dose, and 20% did not respond to a 10 µg/ml LPS dose with IFN production, and 50% did not respond to the same dose of LPS with TNF production. These data are essential for proper selection of biological material. Non-responding PBL cultures are detected by evaluation of the negative (non-treated, spontaneous release of cytokine) and positive (treated with a standard inducer, such as PHA or LPS) controls.

This Sheet Has Been Replaced



Table 2: IFN and TNF response of PBL from individual blood donors after stimulation either with various batches of TTP or with standard inducers

Inducer	Dose (µg/ml)	No.PBL tested	No. responders to: IFN (%)	TNF (%)
TTP 010391	100	19	13 (68)	13 (68)
	30	13	6 (46)	9 (69)
	10	15	14 (93)	13 (87)
TTP 020391	100	14	10 (71)	11 (79)
	30	12	8 (67)	8 (67)
	10	10	5 (50)	5 (50)
TTP 101091	100	26	21 (81)	20 (77)
	30	20	9 (45)	8 (40)
	10	24	14 (58)	15 (63)
TTP 010991	100	5	5 (100)	4 (80)
	30	4	2 (50)	0 (0)
	10	5	2 (40)	3 (60)
TTP 021091	100	5	4 (80)	4 (80)
	30	4	3 (75)	4 (100)
	10	5	3 (60)	3 (60)
TTP 111191	100	5	3 (60)	4 (80)
	30	4	2 (50)	3 (75)
	10	5	4 (80)	2 (40)
TTP (8x)	100	29	19 (66)	10 (34)
	30	24	14 (58)	11 (46)
	10	21	16 (76)	2 (10)
PHA	10	69	64 (93)	54 (78)
LPS	10	41	33 (80)	21 (51)
None	-	69	36 (52)	20 (29)

This Sheet Has Been Replaced



Table 3: Induction of IFN or TNF by different batches of 100 µ/ml peat extract in International Activity Units

	IFN / units/ml			TNF / units/ml				
Series No.	Number of assays	Range	Medi-ana	Number of assays	Time of assays 24h Range	Medi-ana	6 days Range	Medi-ana
Control without induces	333	<10-100	10	41	9-750	27	9-80	9
31090	3	5-30	30	3	ND	ND	9-80	80
291290	21	10-3000	100	14	ND	ND	9-750	40
010391	23	10-2000	60	17	ND	ND	9-160	9
020391	19	10-1000	30	32	9-750	200	9-250	27

The conditioned media were stored at 4°C before assaying

Human peripheral blood leukocytes from healthy blood donors ($8x10^6$ cells/ml) were used.

B. Experiments performed.

Peat extract samples taken from different production batches numbered as shown in Table 3 have been tested using the PBL test as described above. In the experiments, interferon (IFN) and tumor necrosis factor (TNF) have been determined qualitatively and according to the above mentioned standard

This Sheet Has Been Replaced

procedure; their activity, as expressed in International Activity Units, was determined. Experiments were repeated a number of times quoted in Table 3. Result ranges and medians are also given.

For illustration purposes, a similar test was carried out with a standardised peat preparation. At a concentration of 30 µg/ml, interferon activity was within a range of 30 - 300 International Activity Units, while at a concentration of 100 µg/ml, it was 300 - 1000 International Activity Units.

Intermediate concentrations showed a linear relation between dose and response.

The immunoactive peat extracts tested in the above Example are products described in PCT Application No. PCT/EP92/00491.

From the data given in Table 3 it is clear that in PBL cultures negative and positive control is essential for evaluation of the results.

Further experiments show that, when non-responding PBL cultures are not taken into consideration and statistical evaluation of the large number of experiments is performed, the effectiveness of the tests according to the invention cannot be questioned.

For over two years the cytokine inducing activity of over 20 different batches of TTP, including 10 standard commercial batches of the drug, the activity of which has been determined in biological assays, 2) batches rejected by the manufacturer due to the inadequate biological activity determined in mice which was below the established standard and 3) laboratory peat extracts prepared on a small scale. The results obtained are presented in Table 4 below in the

This Sheet Has Been Replaced

form of mean levels of IFN and TNF induced (with calculated standard deviations - SD and statistical significance).

Table 4: Effects of different batches of TTP on IFN and TNF production by human PBL

Inducer	Dose (µg/ml)	IFN Log_{10} units/ml (± SD)	TNF units/ml (± SD)
TTP 010391	100	1.41 ± 1.07[b]	1.26 ± 0.91[c]
	30	0.96 ± 1.07	1.25 ± 0.88[b]
	10	1.81 ± 0.58[c]	1.45 ± 0.63[c]
TTP 020391	100	1.23 ± 0.81[a]	1.35 ± 0.75[c]
	30	1.12 ± 0.90	1.14 ± 0.83[b]
	10	0.75 ± 0.76	0.86 ± 0.90
TTP 101091	100	1.70 ± 0.91[c]	1.38 ± 0.81[c]
	30	0.99 ± 1.12	0.62 ± 0.80
	10	1.22 ± 1.07[a]	0.94 ± 0.76[b]
TTP 010991	100	2.24 ± 0.30[c]	1.35 ± 0.73[b]
	30	1.00 ± 1.00	0.0
	10	0.56 ± 0.68	0.88 ± 0.73
TTP 021091	100	1.72 ± 0.87[b]	1.49 ± 0.84[b]
	30	1.39 ± 0.81	1.79 ± 0.20[c]
	10	1.11 ± 0.91	1.01 ± 0.84
TTP 111191	100	1.36 ± 1.12	1.18 ± 0.62[b]
	30	1.00 ± 1.00	1.21 ± 0.71[a]
	10	1.56 ± 0.86[a]	0.54 ± 0.66
TTP (8x)	100	1.04 ± 0.82[a]	0.63 ± 0.91
	30	0.88 ± 0.81	0.75 ± 0.84
	10	1.10 ± 0.70[b]	0.13 ± 0.40[a]
PHA	10	2.08 ± 0.79[c]	1.67 ± 0.99[c]
LPS	10	1.36 ± 0.80[c]	0.97 ± 1.01[b]
None	-	0.64 ± 0.63	0.42 ± 0.67

Mean values were calculated from all results of IFN or TNF titrations. The non-responders were regarded as 0.

This Sheet Has Been Replaced

[a-c] Different from "None" (spontaneous production of the cytokine) at [a] $p<0.05$, [b] $p<0.01$ or [c] $p<0.001$.

Mediana values corresponding to the results presented in Table 4 are given in the next Table 5, while selected data are presented in a graphic form on Fig.5 and 6 showing the effects of different batches of TTP on IFN and TNF production, respectively.

Table 5: Effect of different batches of TTP on IFN and TNF production by human PBL

Inducer	Dose (µg/ml)	IFN Units/ml[1]	TNF Units/ml[1]
TTP 010391	100	30	27
	30	<10	27
	10	100	27
TTP 020391	100	30	50
	30	30	34
	10	<10	<9
TTP 101091	100	100	34
	30	<10	<9
	10	45	18
TTP 010991	100	300	27
	30	<10	<9
	10	<10	18
TTP 021091	100	100	27
	30	60	80
	10	60	27
TTP 111191	100	100	18
	30	<10	40
	10	100	<9
TTP (8x)	100	20	<9
	30	10	<9
	10	20	<9
PHA	10	200	80
LPS	10	30	27
None	-	10	<9

Mediana values were calculated from all results of IFN or TNF titrations.

Identification of the induced cytokines has been performed by means of neutralisation of IFN and TNF induced by TTP.

The above results of the experiments are explained by Fig. 5 and 6 respectively.

Effects of different batches of TTP on IFN production by human PBL (Fig.5). Seven different batches of TTP were used. Every point of the graph refers to the PBL of an individual healthy blood donor. IFN activity refers to antiviral units. The shaded area shows the limit of the non-significant data. The horizontal lines show mean values. The induction of IFN by 30 and 100 µg/ml of TTP is statistically significant (at $p < 0.05$).

Effects of different batches of TTP on TNF production by human PBL. TNF activity is expressed in the cytotoxic units for the mouse L_{929} cells. The response to 30, 100 and 200 µg/ml of TTP is significant (at $p < 0.05$) (Fig. 6).

In the experiments, potent polyclonal antisera have been used, namely: anti-natural IFN-α, anti-lymphoblastoid (Namalwa) IFN-α and anti-natural IFN-γ to neutralize antiviral activity in supernatants of PBL cultures treated for 20 h with three different batches of TTP. Results, as shown on accompanying drawing Fig.7 (Results of neutralization of IFNs by polyclonal Anti-IFNs sera) indicate that portions of IFN types α and γ produced by PBL from the individual blood donors varied considerably. It is suggested that the released IFN pattern of the each PBL donor is an individual characteristic of the donor.

This Sheet Has Been Replaced

Neutralisation assays carried out with a potent polyclonal rabbit anti-TNF-α serum (Genzyme Inc.) showed that the TNF induced with TTP is mainly TNF type α. It has been confirmed by the results of similar neutralisations carried out with rabbit polyclonal anti-TNF-β serum (Genzyme Inc.) which did not neutralise the TTP induced TNF activity.

Fig.7 shows the results of neutralization of IFNs by polyclonal anti-IFNs sera. IFN containing media from PBL cultures incubated with TTP (three different batches of TTP and seven different PBL were used) were treated with the indicated anti-IFN sera. 1 - Non treated preparations; 2 - Treated with anti-HuIFN-α; 3 - Treated with anti-HuIFN-Ly; 4 - Treated with anti-HuIFN-γ.

Example 6: The procedure described in Example 5 was followed in order to show the amplifying effect of the presence of an amplifyer, viz. indomethacin, according to the present invention.

Each PBL culture was divided into several samples. One of them was a negative control showing spontaneous release of cytokines, the other one was a positive control treated with a standard potent cytokine inducer of a nature indicated below and the remaining samples were treated with 1-100 µg/ml TTP, batch no. 010391 or 020391, as well as TTP I and TTP II which appear in Tables 6b and 6c respectively, and 5 µg/ml indomethacin + TTP of the same batches (Table 6a) or 10 µg/ml compound ITCL + TTP of the same batches (Table 6b), or else 10 or 20 µg/ml seleno-organic compounds (1), (2) or (3) as defined above + TTP of the same batches. In the supernatants of the incubated cultures IFN and TNF were determined.

This Sheet Has Been Replaced



The results obtained are shown in Tables 6a, b and c below.

Table 6a:

The effect of amplification of the cytokine induction in the presence of indomethacin

Exp.	Inducer	Indo-methacin	Concen-tration µg/ml	Cytokines (units/ml) IFN	TNF
1.a)	TTP 010391	-	100	300	750
	-"-	-	10	30	250
	-"-	-	1	20	80
	-"-	+	100+5	1000	2200
	-"-	+	10+5	200	750
	-"-	+	1+5	<10	250
	LPS	-	10	200	500
	none	-	-	<10	9
	none	+	5	20	27
2.b)	TTP 020391	-	100	600	50
	-"-	-	40	20	9
	-"-	-	20	10	9
	-"-	-	10	<10	<9
	-"-	+	100+5	600	80
	-"-	+	40+5	100	27
	-"-	+	20+5	100	9
	-"-	+	10+5	30	9
	PHA	-	5	600	50
	PHA	+	5+5	2000	80
	none	-	-	60	<9
	none	+	5	10	<9

a) PBL were prepared from fresh heparinized blood by Ficoll-Hypaque separation technique. PBL cultures contained $7x10^6$ cells/ml,

b) PBL were prepared from the buffy coat obtained from the Regional Transfusion Center. The cells were processed according to the Cantell et al. method (Meth. Enzymol. 1981, 78, 29-38).

PBL cultures contained $8x10^6$ cells/ml.

Table 6b:

The effect of amplification of the cytokine induction in the presence of Compound ITCL

Exp.	Inducer	Compound ITCL	Concentration	Cytokines (units/ml) IFN	TNF
1.	TTP 010391	-	100	100	18
	TTP 010391	-	30	100	9
	TTP 010391	+	100+10	3000	80
	TTP 010391	+	30+10	300	80
	PHA	-	10	3000	250
	none	-	-	10	<9
	none	+	10	<10	<9
2.	TTP I	-	100	3000	27
	TTP I	-	30	30	<9
	TTP I	+	100+10	3000	250
	TTP I	+	30+10	300	9
	PHA	-	10	100	18
	none	-	-	10	<9
	none	+	10	10	<9

This Sheet Has Been Replaced



Table 6c:

The effect of amplification of the cytokine induction in the presence of the seleno-organic compounds

Exp.	Inducer	Seleno-organic compounds	Concentration µg/ml	Cytokines (units/ml) IFN	TNF
1.	TTPs	-	10	100	<9
	TTPs	-	5	10	<9
	TTPs	Comp. (1)	10+20	300	27
	TTPs	Comp. (1)	5+20	100	27
	TTPs	Comp. (2)	10+20	300	50
	TTPs	Comp. (2)	5+20	100	9
	PHA	-	10	100	<9
	none	Comp. (1)	20	200	9
	none	Comp. (2)	20	100	27
	none	-	-	10	<9
2.	TTP II	-	50	10	<9
	TTP II	-	5	10	<9
	TTP II	Comp. (3)	50+10	30	27
	TTP II	Comp. (3)	10+10	30	27
	PHA	-	-	30	<9
	none	Comp. (3)	10	10	<9
	none	-	-	10	<9

PBL cultures contained $8x10^6$ cells/ml. TTP is a peat preparation produced on a small scale. The relatively low TNF response is probably due to the fact that the buffy coats were stored for 20 hrs at 4°C before the preparation of the cultures. The seleno-organic compounds are inducers of the cytokines as well as TTP.

This Sheet Has Been Replaced


Example 7: A diagnostic test reflecting the effect of administration of TTP orally in daily doses of 5 mg in the form of commercially available tablets on PBL in vitro culture response to an additional dose of TTP induction of cytokines was performed according to the particulars given below:

The test was performed with 4 healthy female volunteers, age 43-58, who donated 20 ml venous blood taken from v.ulnaris weekly, before, during and after the administration of TTP at a daily dose of 5 mg in tablets. Two different regimen of administration have been performed. Two of the volunteers (referred to further as B.K. and J.Z.J.) were subjected to a treatment with TTP consisting of three seven-day cycles: the drug was administered per os, one 5 mg tablet daily for one week with a seven day rest followed by a seven-day period of TTP oral administration. The total dosage was 21 tablets. The other two volunteers (referred to further as A.D.I. and W.F.) were treated with one 5 mg TTP tablet daily administered per os continuously for three weeks. The total dosage was also 21 tablets. After two weeks of rest, the treatment was repeated.

The 5 mg. TTP tablets used were a standard commercial product of TORF CORPORATION Pharmaceutical Factory of Wroclaw, Poland. Each tablet contained 5 mg TTP, 43 mf Lactose and 2 mg MYVATEX. To induce cytokines in PBL in vitro cultures, powdered TTP as a pure substance was used in form of a stock solutiom containing 20 mg TTP/ml of pyrogen free redistilled water stored at 4°C.

Blood samples taken from the volunteers were heparinised with Heparin POLFA solution without preservatives at a final concentration of 10 units/ml and treated as described earlier to prepare a PBL in vitro culture, which was then treated with cytokine inducers PHA, LPS and TTP in a manner described in the preceding examples. 200 µg cultures were

This Sheet Has Been Replaced



used in each treatment. The supernatants were stored at 4°C and assayed for IFN within one week, and stored at -20°C and assayed, respectively for TNF to avoid inactivation due to spontaneous proteolysis.

IFN and TNF were assayed in the same manner as described above. The results obtained are presented on the accompanying drawings Figs.8-13 (response of PBL cultures of different patients to IFN inducers and to TNF inducers during the oral administration of TTP).

As may be seen from the data presented on Figs.8 and 9 showing the IFN response of the first group of volunteers during administration cycle, the response declined and returned to the initial value after two weeks rest. In contrast, TNF response shown on Figs.10 and 11 increased during administration of TTP and returned to the normal level after two weeks rest.

Data shown on Figs.12 and 13 reflecting the response to the cytokine induction in the other way of administration of volunteer A.D.I. prove that the hyporeactivity state to IFN induction is reached after 3 weeks of continuous administration of TTP orally. The hyporeactivity state shown as loss of ability of PBL cultures to respond to the induction of IFN with TTP solution disappears after two weeks of rest. In contrast, during three weeks of TTP administration, the hyporeactivity to TNF induction did not develop (Fig.12).

The above results indicate that the optimum dose of immunomodulator being a cytokine inducer can be established individually for each patient treated with a simple PBL test according to the present invention.

CLAIMS

1. Method for determining the immunological activity of bioactive, cytokine inducing immunomodulator substances, characterised in that a human peripheral blood leukocyte (PBL) culture or a suspension of BALB/c mice resident peritoneal cells (RPC) is treated with a solution of the substance to be tested in order to induce production of cytokines which then are determined according to standard identification methods.

2. Method according to claim 1, characterised in that a human PBL culture is used, wherein the production of different cytokines is induced, whereafter each of the induced cytokines is determined qualitatively and quantitatively using a set of different - preferably known - sera which recognize individual cytokines.

3. Method according to claim 1 or 2, characterised in that the human PBL culture is prepared with a nutrient medium suitable for tissue culture and that the density of the ready-for-use PBL culture is approx. 8×10^6 leukocytes/ml.

4. Method according to any one of claims 1 to 3, characterised in that the solution to be tested is used in a concentration of 0,1 - 200 µg/ml.

5. Method according to any one of claims 1 to 4, characterised in that each cytokine is identified quickly enough to avoid proteolysis of the cytokine.

6. Method according to claim 1, characterised in that a PBL culture or an RPC suspension is used, and the bioactive substances induce the production of interferons and tumor necrosis factors.

This Sheet Has Been Replaced

7. Method according to claim 6, characterised in that the RPC suspension contains about 1 - 2 x 10^6 cells/ml.

8. Method according to claim 6 or 7, characterised in that the solution to be tested is prepared in a nutrient medium Eagle or RPMI-1640 with an addition of 10% of fetal calf serum.

9. Method according to any one of the preceding claims, characterised in that the results obtained are evaluated against a calibration curve.

10. Method according to any one of the preceding claims, characterised in that peat extracts, such as TTP, are tested.

11. Method according to any one of claims 1-6, 8, 9 or 10, characterised in that, in the case of the PBL variant, amplification of the results is achieved by admixing to the solution to be tested a non-steroidal anti-inflammatory drug, preferably compound ITCL and seleno organic compounds (1-3), and most preferably indomethacin.

12. Method for determining the immunological response of a human individual to a therapy using a cytokine inducing immunomodulator substance, characterised in that a peripheral blood leukocyte (PBL) culture of the human individual, treated with such immunomodulator substance, is treated in defined time intervals with a solution of the substance administered, in order to induce production of cytokines, which are then determined according to standard methods, in order to determine the moment of development of hyporeactivity to the substance after prolonged administration of the substance and the moment of regaining the ability to respond to an additional dose of the substance after a certain period of hiatus.

This Sheet Has Been Replaced

13. Method according to claim 12, characterised in that the time intervals are about 7 to 14 days.

14. Method according to claim 12 or 13, characterised in that it is applied in the course of a therapy using a peat extract as cytokine inducing immunomodulator substance.

15. Method according to claim 12, 13 or 14, characterized in that it is applied in the course of a therapy using TTP as cytokine inducing immunomodulator substance.

16. TTP, whenever used, tested or determined by means of a method as claimed in any one of claims 1-15.

17. Composition for carrying out the method according to any one of claims 1 to 15, characterised in that it contains, on one hand, a PBL culture or a RPC suspension, and, on the other hand, a non-steroidal anti-inflammatory drug, preferably indomethacin.

This Sheet Has Been Replaced





Abstract

METHOD AND COMPOSITION FOR DETERMINING THE IMMUNOLOGICAL ACTIVITY OF BIOACTIVE SUBSTANCES

The immunological activity of certain bioactive substances is determined by treating a human peripheral blood leukocyte (PBL) culture or a suspension of BALB/c mice resident peritoneal cells (RPC), with a solution of the substance to be tested in order to induce production of cytokines which then are determined according to standard identification methods; amplification of the results may be achieved by admixing - to the solution to be tested - a non-steroidal anti-inflammatory and prostaglandin synthesis inhibiting drug, preferably indomethacin. The method is also useful for determining the immunological response of a human individual to a therapy using a cytokine inducing substance.

REPLACEMENT SHEET
13 FEB 1998

9807958-7 IPOS

REPLACEMENT SHEETS SG
☑ Description : paged 1... to paged 32.
☑ Claim(s) : paged 33. to paged 35.
☐ Drawing(s) : paged ..... to paged .....
☑ Abstract : paged 36... to paged ....

DATE OF RECEIPT 13/2/98

METHOD AND COMPOSITION FOR DETERMINING THE IMMUNOLOGICAL ACTIVITY OF BIOACTIVE SUBSTANCES.

The present invention relates to a method and composition for assaying certain bioactive substances and determining their immunological activity with respect to their ability to induce production of cytokines. Cytokines, such as interferons (IFNs) and tumor necrosis factors (TNFs), are hormone-like proteins which play an important role in virtually all immunological reactions, as well as in the regulatory process responsible for the maintenance of homeostasis. The production of such cytokines can be induced by certain substances which, on account of their bioactive and immunomodulating activity, are useful in the therapy of immunodeficiencies and related diseases.

There is, of course, a substantial need of methods for properly and easily assaying the immunological activity of such bioactive substances. It is the object of the present invention to provide such a method.

EP-A-238 851 discloses *in vitro* test methods for determining the magnitude of immune capacity of an animal or human, based on exposure of PBL to a mitogen or an antigen such as PHA, tetanus toxoid etc. in combination with an additional immunoamplifier capable of increasing the IL-2 production, such as amplifier Beta, tyrosylglycine, tyrosylglycylglycine, etc. Whether or not the immune response may be amplified is determined by assaying the production of IL-2 and, possibly, of gamma-interferon.

The method according to the present invention, however, allows to determine whether a tested substance is able to induce production of different cytokines and permits a quick and efficient check of the properties of certain substances, namely their immunological activity, at each stage of production, separation, purification and formulation into

REPLACEMENT SHEET
13 FEB 1998

final compositions. The present invention also provides a composition and kits suitable for carrying out said method.

Tests used until now to determine immunological activity are numerous but difficult to carry out. When a new therapeutical composition having an immunological effect is clinically tested, its immunological effectiveness is evaluated by monitoring the numerous, well known and analytically determinable features of immunological systems employing standard methods. Once the activity is proven, there emerges the necessity for representative single tests enabling a quick check of the status of a patient in terms of his or her immunological response.

In clinical studies, evaluating IFN serum levels may be difficult, unprecise and misleading. Interferons are often produced locally in various tissues, they are short-lived, their action is paracrine (in the immediate vicinity only) and they are strongly bound to cellular receptors and carrier proteins.

On the other hand, technological processes for manufacturing, purification and/or separation of immunologically active substances such as in particular, extracts from raw peat are usually multi-step processes, and there is always the need for establishing a reliable and fast test enabling a thorough control of the production. Moreover, since the substances are frequently of a complex nature, as it is the case with peat extracts, it is important to have a similar test for determining the activity of individual fractions of such substances. It is also necessary to find out a microscale testing method, since the products to be analysed are very costly.

The following four basic findings allowed to solve the above described problems.

REPLACEMENT SHEET
13 FEB 1998

1. Some immunoactive peat extracts induce the production of cytokines, such as interferons and tumor necrosis factors, in in-vitro cultures of peripheral blood leukocytes (PBL), the induction being dependent on the dose.

2. Interferon-ß (and TNF-α) is produced in significantly higher quantities by BALB/c mice resident peritoneal cells (RPC) when treated with immunoactive peat extracts as compared to a spontaneous release of these cytokines in non-treated samples.

3. Certain peat extracts, when tested in-vitro as mentioned above show a synergistic effect in combination with known immunomodulators, such as organoselenium compounds, the p-chlorophenylamide of 3-methyl-5-benzoylamino-isothiazol-4-carboxylic acid and indomethacin. This allows the use of much smaller quantities of immunologically active substances to show their activity.

4. PBL of healthy volunteers treated with TTP (a certain peat-derived product) administered orally at a dosage of 5 mg/day, tested for induction of certain cytokines, IFN-α, IFN-γ and TNF-α, lose the ability to respond to induction of the cytokines with TTP solution after prolonged, uninterrupted administration of TTP orally and regain this ability after approximately two weeks of rest, i.e., of interrupted TTP administration.

The method according to the present invention comprises basically the steps of treating a human peripheral blood leukocyte (PBL) culture or a suspension of BALB/c mice resident peritoneal cells (RPC), with a solution of the substance to be tested in order to induce production of cytokines, and then determining said cytokines according to standard identification methods.

REPLACEMENT SHEET
13 FEB 1998

The method according to the present invention is defined as follows:

Method for testing bio-active, cytokine inducing immunomodulator substances by determining their immunological activity, characterized in that

a) a cell culture, comprising human peripheral blood leukocytes (PBL) obtained from donors that are not hyporeactive to interferon (IFN) induction, is prepared with a nutrient medium suitable for tissue culture; or

b) a suspension of fresh resident peritoneal cells (RPC) obtained from - preferably 5 to 8 weeks old - healthy "virgin" BALB/c mice that are not immunologically stimulated or that do not show very high spontaneous cytokine production, is prepared with a suitable medium; and

c) non-hyporeactiveness under a) or lack of immunological stimulation or lack of very high spontaneous cytokine production under b) is ensured by positive and negative control; whereupon

d) the cell culture under a) or the cell suspension under b) is treated with a solution of the substance to be tested in order to induce the production of cytokines which then are determined according to standard identification methods.


The peripheral blood leukocyte (PBL) culture used according to the first variant or embodiment of the present method is a short term culture prepared from fresh leukocytes of healthy humans with a nutrient medium suitable for tissue culture. The preferred density of a ready-for-use culture is about 8×10^6 leukocytes/ml. The solution to be tested is preferably used in a concentration of 0.1 - 200 µg/ml. The method is, in particular, suitable for testing peat extracts.

A micro-scale determination is possible when the solution to be tested, e.g. a peat extract or a fraction thereof, is


REPLACEMENT SHEET
13 FEB 1998

admixed with an immunomodulating agent, such as a non-steroidal anti-inflammatory drug, preferably a substance selected from the group comprising organoselenium compounds, such as Ebselen, certain isothiazole derivatives, such as the p-chlorophenylamide of 3-methyl-5-benzoylamino-isothiazole-4-carboxylic acid, and indomethacin, as well as analogues, homologues and metabolites of such substances.


Examples of organoselenium compounds which can be used for this purpose are compounds of the following formulae 1-3; the compound of formula 2 (Ebselen) being preferred:



(1) Bis-[2-(N-phenylcarboxamido)]-phenyldiselenide



(2) 2-Phenyl-1,2-benzisoselenazol-3(2H)-one (Ebselen PZ 51)




REPLACEMENT SHEET
13 FEB 1998

(3) Bis-[2-(N-(2-pyridyl)carboxamido)]-phenyldiselenide

REPLACEMENT SHEET
13 FEB 1998

Examples of isothiazole derivatives which can be used for the above purpose are compounds of the general formula I



(I)

wherein R_1 is halophenyl, preferably chlorophenyl, R_2 is phenyl and R_3 is lower alkyl, preferably methyl. The halogen atom in the halophenyl group, chlorophenyl group R_1 is preferably in p-position of the phenyl ring.

The preferred compound of the above formula I is the p-chlorophenylamide of 3-methyl-5-benzoylamino-isothiazole-4-carboxylic acid, i.e. the compound of formula I wherein R_1 is p-chlorophenyl, R_2 is phenyl and R_3 is methyl. For brevity sake, this compound will be referred to below as 'Compound ITCL'. There exists a Trade Mark registration "VRATIZOLIN^R" for this compound in Poland. Its synthesis and properties are described in Arch. Immunol. et Ther. Exp. 1973, 21, 891.

All the immunomodulating agents referred to above for being useful for micro-scale determinations, i.e. the organoselenium compounds, the isothiazole derivatives and indomethacin, have in common that they are non-steroidal anti-inflammatories which inhibit the synthesis of prostaglandins. They amplify the test results in the method according to the present invention. For this reason, they are particularly useful for micro-determinations. Amplification of the results may be 5-20 times.

Out of all the various non-steroidal anti-inflammatory compounds useful for the purpose of amplification, seleno

13 FEB 1998

organic compound (1, 2 and 3) and compound ITCL, but especially indomethacin, are the preferred ones.
In the present invention, different antisera, which recognise individual induced cytokines, are employed. These sera are known and used for testing and standardisation of different cytokines. Identification of the induced cytokines should take place at the moment of completion of the cytokine formation and prior to its proteolysis. Among different cytokines induced, some are formed slowly and remain stable in the solution, as for example interferon γ, while others are formed rapidly and are susceptible to proteolysis, as for example tumor necrosis factors.

The method as described above may be employed also for the determination of an immunological response of a human individual to a certain immunoactive substance, provided that such immunological response is not stimulated by other factors, as for example by viral or bacterial infections. It may be applied for monitoring of the drug response in individual patients, for the determination of the optimal dose of a therapeutic cytokine inducer, such as TTP, as well as for the establishment of the effective scheme of the drug administration. The method is based on the assessment of the hyporeactivity state to IFN induction.

Thus, the present invention also relates to a method for monitoring the immunological response of a patient to a therapy using a cytokine inducing substance, characterized in that

a) cultures of peripheral blood leukocytes (PBL) obtained from blood samples that are taken from said patient in defined time intervals are prepared with a nutrient medium suitable for tissue culture;
b) said PBL cultures are treated with a cytokine inducing substance in order to induce production of cytokines, preferably interferon and tumor necrosis factor; and

REPLACEMENT SHEET
13 FEB 1998

c) said cytokines are then determined according to standard methods,

in order to assess development and disappearance of hyporeactivity of said PBL-cultures to interferon induction.

The time intervals are preferably about 7-14 days.

In a preferred embodiment, the above method is applied in the course of a therapy using a peat extract as cytokine inducing immunomodulator substance.

According to a particularly preferred embodiment of the invention, this method is applied in the course of a therapy using TTP as cytokine inducing immunomodulator substance.

In view of the above, the present invention also relates to TTP, whenever used, tested or determined according to the methods of the present invention.

The present invention also provides, in its second variant, for a testing method which is suitable for quick monitoring of technological processes for the production, purification, separation or the like of the immmunoactive substances. For instance, when a suspension of BALB/c mice resident peritoneal cells (RPC) is treated with the solution to be tested, mice interferon-ß as well as tumor necrosis factor are induced. Both may be detected and qualitatively determined according to standard identification methods.

In this method, a suspension of fresh mouse resident peritoneal cells (RPC), containing approx. 1 x 10^6 cells/ml, is used. The solution to be tested is prepared in a nutrient medium Eagle, or RPMI-1640 with the addition of 10% of fetal calf serum. The results obtained are evaluated against a calibration curve prepared with a model substance. As a model substance, a sample of the same immunologically active substance tested in a traditional way may be used. Again,

REPLACEMENT SHEET
13 FEB 1998

the method is particularly suitable for monitoring the process for obtaining peat extracts as well as analysing their fractions.

The present method is very simple and efficient in comparison with other known methods, because it is based on a direct induction of interferon ß instead - as is the case in a number of conventional tests - on a secondary effect of formulation of antibodies. Until now, it was not possible to find any similar method due to the fact that the immunosystem of mice significantly differs from the human immunosystem and tests recognised as indicative for human immunologic response - such as the presence of interferons in the tissues of spleen or lymphonodes etc. - do not work with the best laboratory test animals, i.e. with mice of BALB/c type.

It was now found that a clear immunological response of the BALB/c mice can be obtained when resident peritoneal cells (RPC) are chosen as a biological material for testing, fresh suspensions of such cells being used. For each test it is sufficient to sacrifice just three animals. Results are obtained within several hours. The biological material may also be cultured and a corresponding line of macrophages be established.

The method according to the present invention employs the use of biological material, namely a peripheral blood leukocyte (PBL) culture or a suspension of BALB/c mice resident peritoneal cells (RPC), according to the information given above. Presence and concentration of each cytokine is determined by methods for testing and standardisation of interferons known from "Methods in Enzymology", 1986, vol.119, part C: "Interferon", edited by Sidney Pestka. Determination of the whole spectrum of cytokines induced identifies the immunological status of a patient.

REPLACEMENT SHEET
13 FEB 1998

For the proper evaluation of the results, it is essential that the biological material for carrying out both variants of the method according to the present invention is selected from the proper donors. In case of human leukocytes, samples taken from individuals who show a hypo-reactivity (rare cases), i.e. a low level of immunological response, should be excluded from the evaluation. In case of the BALB/c mice RPC suspension, samples taken from additionally stimulated animals (for example infected ones) should also be excluded from the evaluation. In case of the BALB/c mice RPC suspensions, samples should not be taken from too young or from too old animals, as there is an age-dependent deficiency in cytokine production in such mice. 5-8 weeks old healthy animals should be selected. Samples showing very high spontaneous production of cytokines should be eliminated, especially when immunoregulatory substances are tested, since no potentiation of cytokine induction or sometimes even reduction of such induction may be observed.

That embodiment of the method according to the present invention, wherein amplification of the results is achieved by admixing a non-steroidal anti-inflammatory drug (one of the substances referred to above, such as compound ITCL or seleno organic compound 1, 2 and 3, but preferably indomethacin) to the solution to be tested, is a microscale test, particularly suited for determining the induced TNF and IFN activity. This test is economic, fast, applicable for assaying a great number of samples simultaneously and suitable for standardization and, at least partial, automatization.

The two variants of the method according to the present invention are described by way of example in more detail below:

Preparation of the solution of the substance to be tested:

REPLACEMENT SHEET
13 FEB 1998

The substance to be tested, e.g. a peat extract, is dissolved in sterile bidistilled water at a concentration of 10 mg/ml. The samples are sterilized by filtration through 0.45 µm, 600 kPa max Millipore(R), antibacterial filters. Next, solutions are made in a complete RPMI-1640 medium containing heat-inactivated fetal bovine serum (FBS).

A) The PBL variant:

Cytokine induction

Buffy coats from healthy blood donors may be obtained from the regional transfusion center. Alternatively, peripheral blood leukocytes (PBL) may be isolated from heparinized venous blood of healthy volunteers by Ficoll-Hypaque(R) (or Histopaque(R)) density gradient (g = 1.077) centrifugation, with subsequent twice washing of the cells. The erythrocytes were lysed with NH_4Cl treatment according to Cantell et al. (Cantell, K., Hirvonen, S., Kauppinen, H.L.: Production and Partial Purification of Human Immune Interferon. Meth. Enzymol. 119, 54, 1988). The leukocytes from a single donor containing approximately $8x10^6$ leukocytes/ml in RPMI 1640 medium supplemented with 10% fetal bovine serum (FBS), L-glutamine and antibiotics were used. All lots of FBS were pretested. Only non-mitogenic FBS for PBL cultures was used. The cytokine inducers were added to 200-1000 µl volumes of the cultures. The reference cytokine inducers were phytohemagglutinin (PHA) (Pharmacia Fine Chemicals, Sweden or Sigma, USA). The induced cultures of PBL were incubated in an atmosphere of 5% CO_2 in air at 37°C for 20 h and centrifuged. Supernatants were stored at 4°C and assayed for IFN activity within one week. Supernatants for TNF activity determination have to be stored at -90°C or in liquid nitrogen to avoid inactivation due to proteolysis.

Interferon assay

Confluent monolayers of A549 cells were prepared in microplates in Dulbecco-modified Minimum Essential Medium

REPLACEMENT SHEET
13 FEB 1998

(DMEM) with 10% FBS, L-glutamine, and antibiotics (penicillin 100 units/ml and streptomycin 100 µg/ml). IFN samples diluted in the plates were added to the cell monolayer and incubated at 37°C for 20 h in 5% CO_2 in air. The cells were then washed and challenged with encephalomyocarditis virus (EMCV). The titer of IFN was defined as the dilution of IFN sample that reduced virus cytopathogenic effect by 50% after 48 h of incubation. The MTT (3-[4.5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) method (Hansen, M.B., Nielsen, S.E. and Berg. K.: Re-examination and Further Development of a Precise and Rapid Dye Method for Measuring Cell Growth/Cell Kill. J. Immunol. Meth. 1989, 119, 203-210) to measure the cell kill in the ELISA scanner was also used. Laboratory standards of IFN have to be included in all assays e.g. recombinant human IFN-γ (specific activity $2x10^6$ units/mg), the natural human leukocyte IFN-α ($3x10^6$ IU/ml) and IFN-γ ($2x10^6$ IU/ml).

TNF assay

The cytotoxic activity of TNF was measured in L_{929} cells according to Flick and Gifford (Flick, D.A., Gifford, G.E.: Comparison of in Vitro Cell Cytotoxic Assays for Tumor Necrosis Factor. J. Immunol. Meth. 68, 1667, 1984). The samples and actinomycin D (5µg/ml) solution were added to monolayer cultures of the cells. After incubation at 37°C for 20 h, cytotoxic effects of TNF were determined either by the microscopic examination of the cultures or by using the MTT method. The amount causing approximately 50% destruction of the cell cultures was defined as one unit of TNF activity. Comparison with a preparation of TNF-α (Genentech Inc., USA) showed that 1 unit in the assays was equal to 100 - 200 pg/ml TNF.

Cytokine neutralization assays

The cytokines produced by PBL treated with the examined preparations may be identified by neutralization assays with the specific antibodies (Inglot et al., Organoselenides as

13 FEB 1998

potential immunostimulants and inducers of interferon gamma and other cytokines in human peripheral blood leukocytes, Experientia 1990, 46, 308-311). Furthermore, various ELISA kits for the determination of the immuno-activity of a defined cytokine may be applied.

Comment

In the supernatants obtained from the cultured lymphoid cells, other than TNF or IFN cytokines may be found and identified by other standard methods, e.g. Interleukins (IL-1-10), GM-CSF, TGF-ß etc.

B) The RPC variant: Cytokine induction.

Mouse resident peritoneal cells (RPC) are obtained from approximately 6 weeks old BALB/c mice killed with ether, by injecting into the peritoneal cavity 5 ml of complete RPMI-1640 medium at room temperature. The washing medium containing RPC is collected into ice-cooled 50 ml centrifuge tubes. The RPC are neither washed nor centrifuged. The cells are counted in a Bürker hemocytometer and suspended to a density of 1-1.5x10^6 cells/ml. Such cell density is required for all of the assays measuring the effects of various concentrations of the preparation (from 0.1-500 µg/ml). The negative and positive controls are included in all of the tests. The negative control measures the spontaneous release of TNF or IFN by RPC incubated with the complete RPMI-1640 medium without the inducers, whereas the positive control shows the effect of a standard lipo-polysaccharide inducer (LPS from E. coli 055:85, Sigma 1-10 µg/ml). All of the cultures are incubated at 26°C for 20 h. Thereafter, the cultures are centrifuged at 1000 rpm for 10 min and supernatants are collected. Using an automatic pipette, the supernatants are diluted from 1:2 to 1:256.

REPLACEMENT SHEET
13 FEB 1998

Bioassays of the cytokines

To determine TNF activity in supernatants from RPC cultures, the 20 h old monolayer of L_{929} cell cultures are used.

The mouse fibroblast-like L_{929} cells, $2x10^4$ cells per well in 100 µl of complete medium, are seeded in 96-well flat bottom plates and incubated for 20 h at 37°C in order to obtain a monolayer.

The transfer of supernatants must be very accurate. The incubation of the L_{929} cultures is carried on for 24 h at 37°C in an atmosphere of 5% CO_2 in air.

Evaluations

1) Reading of the cytotoxic effects under the reverse microscope.
2) MTT test (3-[4.5-dimethylthiazol-2-yl]-2,5-diphenyl-tetrazolium bromide, Sigma).

To measure cell killing with L_{929} cell cultures, 25 µl of MTT dye solution at a concentration of 5 mg/ml are added into every well of the microplates.

Next, the plates are incubated for 2 h at 37°C in an atmosphere of 5% CO_2 in air.

Thereafter, 100 µl of the solvent solution containing 45 ml of dimethylformamide, 13.5 g of SDS (sodium dodecyl sulphate), and 55 ml of distilled water, are added to every well.

Incubation is carried on for 12 hrs at 37°C in an CO_2 incubator. The results of MTT color test are read in a Multiskan 340/CC reader (Labsystem) using a 570 nm filter. Reference recombinant TNF-α has to be used.

REPLACEMENT SHEET
13 FEB 1998

Interferon bioassay

IFN is assayed by inhibition of the cytopathic effect caused by mouse encephalomyocarditis virus (EMCV in mouse L_{929} cells, reference Mu IFN α/β, standard from National Institute of Health Bethesda, MD, USA is included). The cytopathic effect is observed under a reversed microscope and also it is measured by the MTT method as described below:

MTT method according to Berg et al.
Berg K., Hansen M.B. and Nielsen S.E. (1990): A new sensitive bioassay for precise quantification of interferon activity as measured via the mitochondrial dehydrogenase function in cells (MTT-method). APMIS, 98, 156-162.

MTT (3/4,5- dimethyl-thiazol-2-yl/2,5-diphenyl tetrazolium bromide, Sigma) is diluted in PBS at a concentration of 5mg/ml. To measure cell killing with L_{929} cultures, 25 µl of MTT dye solution at a concentration of 5 mg/ml is added to every well of the microplates. The control reference sample of TNF or IFN is included. Next, the plates are incubated for 2 hrs at 37° C in an atmosphere of 5% CO_2 in air. Thereafter, 100 µl of a solution containing 45 ml dimethylformamide, 13.5 g of SDS (sodium dodecyl sulfate) and 55 ml of distilled water are added to every well. After overnight incubation at 37° C the optical densities at 570 nm are measured, using a microplate reader Start Fax Awareness Technology Inc. 2100, employing the extraction buffer as the blank.

Both TNF-α, IFN-ß, and other cytokines, may be also assayed using commercial ELISA kits.

A composition for carrying out the method of the present invention is essentially characterized in that it contains double processed tissue culture water, a culture medium and

REPLACEMENT SHEET
13 FEB 1998

sera completing the culture medium as well as the human PBL or mice RPC.

A diagnostic kit (cytokine inducer kit) for use in the PBL variant of the present method (for 10-20 manual assays) may contain the following items:

1) Double processed tissue culture water, 100 ml
2) Histopaque(R) or Ficoll-Hypaque(R) - 1077, (density gradient medium), 100 ml
3) RPMI-1640-Medium, 2x100 ml (with sodium bicarbonate and L-glutamine, sterile filtered, endotoxin tested)
4) Fetal bovine serum (FBS), 20 ml, low in endotoxin and hemoglobin
5) Lectin from Phaseolus vulgaris 0.5 mg (Phytohemagglutinin - PHA-P)
6) Round bottom tubes with caps, polypropylene sterile for tissue cultures 17x100 mm, capacity 14 ml, 25 per bag
7) Tissue culture plates with lids, 2 plates, 96 well, flatbottom, sterile.

Reagents available from: Sigma Chemical Co., St. Louis, Mo., USA or other companies.

Such a PBL kit may be used as so-called second phase screening system, to confirm results obtained with RPC cells or sometimes as a direct screening system for several substances which may be inactive as immunomodulators in rodents, e.g. organoselenium compounds.

A diagnostic kit (cytokine inducer kit) for use in the RPC variant of the present method (for 10-20 manual assays) may contain the following items:

1) Double processed tissue culture water, 100 ml
2) RPMI-1640-medium, 2x100 ml, (with sodium bicarbonate and L-glutamine), sterile, filtered, endotoxin tested)

REPLACEMENT SHEET
13 FEB 1998

3) Fetal bovine serum (FBS), 20 ml, low in endotoxin and hemoglobin
4) Lipopolysaccharide (LPS) 0.5 mg, from E. coli 0127:B8
5) Round bottom tubes with caps, sterile, 25 per bag, Polypropylene, for tissue culture, 12 x 75 mm, 6 ml capacity
6) Tissue culture plates with lids, 5 plates, 96 well, flat bottom, sterile.

Reagents are available from Sigma Chemical Co. St. Louis, Mo., USA or other companies.

A composition according to the invention for carrying out the method of the present invention involving amplification of the results, which composition is particularly suited for micro-scale tests, is essentially characterised in that it contains on the one hand a PBL culture or an RPC suspension, and on the other hand a non-steroidal, anti-inflammatory and prostaglandin synthesis inhibiting drug, preferably a substance selected from the group of organoselenium compounds, such as Ebselen, indomethacin or Compound ITCL, and derivatives, analogues, homologues and metabolites of these compounds, most preferably indomethacin.

The following specific Examples illustrate the present invention. It is intended that they do not limit the scope of the invention in any way.

Example 1:

Individual solutions of RPC were obtained from 36 female, 7-8 weeks old BALB/c mice killed with ether, by injecting into the peritoneal cavity 5 ml of Eagle's minimum essential medium (EMEM), supplemented with 10% heat-inactivated calf serum. Washings from the individual mice contained 1-2 x 10^6 cells/ml. Suspensions of the cells from each mouse were divided into two samples distributed to two tubes. One

REPLACEMENT SHEET
13 FEB 1998

sample was treated with 100 µg TTP/ml and the other served as a negative control showing spontaneous release of cytokines.

All cultures were incubated at 26°C for 20h and thereafter they were centrifuged at 1000 rpm for 10 min. The supernatants were collected and assayed for IFN-ß activities in a bioassay using a micromethod of inhibition of cytopathic effect caused by EMCV (encephalomyocarditis virus) in mouse L_{929} cells. In each assay an internal laboratory standard was included which had been calibrated against the international reference preparation of Mu IFN, G002-904-511, NIH, Bethesda, USA. The cytopathic effect was measured by the MTT method according to Berg et al, APMIS, 98, 156-162. The results obtained were as follows: for the controls the IFN-ß level was 1-8 U/ml average being <2, for the samples treated with TTP, the IFN-ß level was 4-127 U/ml, the average being 16 U/ml. The results are graphically presented on the accompanying drawing Fig.1 (Comparison of IFN-ß production by RPC treated with TTP and non-treated. 1-limit of detection, 2-mediana. According to Mediana test P=0,0000. Every point shows the level of IFN in RPC isolated from individual mice).

Example 2:

The procedure as described in Example 1 was repeated with a group of 7 mice, however, instead of dividing each individual mouse RPC suspension into two samples, only one suspension was divided into four samples. The samples were distributed to 4 tubes (1 ml in each). One of the samples was a negative control and the remaining three were treated with 100, 10 and 1 µg TTP/ml to determine a TTP dose-dependence of the IFN-ß release. After incubation of the cultures for 24h at 26°C the tubes were centrifuged. In the collected supernatants IFN-ß was determined as described in Example 1. The results obtained are presented on the

REPLACEMENT SHEET
13 FEB 1998

accompanying drawing Fig.2 (Dependence of IFN production on TTP concentration). Each curve relates to the results observed with a single mouse derived RPC suspension.

Example 3:
The procedure as described in Example 1 was followed, except that TNF-α was determined instead of IFN-ß.

Supernatants from the RPC cultures, non-treated and treated with 100 μg TTP/ml were assayed for TNF activity in a bioassay. The mouse fibroblast-like L_{929} cells (4 x 10^4 cells per well in 100 μl of complete EMEM) were seeded in 96-well flat bottom plates (Falcon, Linbro, Flow) and incubated for 4 h at 37°C in an atmosphere of 5% CO_2 in air. The samples were diluted in EMEM with actinomycin D (end concentration 2.4 μg/ml) in the additional plate. Next, the culture medium above the L_{929} cells monolayer was removed and the prepared dilutions were transferred to this culture using a multichannel pipette. After incubation for 20 h at 37°C, the cultures were assayed for cell kill by the MTT method as described above. The results obtained were as follows: for negative controls the TNF-α level was within a range of 1-256 U/ml the average being 8 U/ml, and for the samples treated with TTP the TNF-α level was 2-256 also, but the average was 64 U/ml. The results obtained are shown on the accompanying drawing Fig.3 (Comparison of TNF-α production by RPC treated with TTP and non-treated. 1-limit of detection, 2-mediana. According to Mediana test P= 0.0030. Every point shows the level of TNF in RPC isolated from individual mice).

Example 4:
The procedure as described in Example 2 was followed, except that instead of IFN-ß, TNF-α was determined essentially as described in Example 3 above.

REPLACEMENT SHEET
13 FEB 1998

The results obtained are shown on the accompanying drawing Fig.4 (Dependence of TNF production on TTP concentration).

Example 5:

A. Biological material. Over 115 buffy coats from individual healthy blood donors, obtained from Wroclaw Regional Transfusion Center, have been used for the assays. The characteristics of the donors are given in a Table 1, below.

Table 1: Characteristics of donors of blood used as a source of PBL

Age (years)	No.	Male	Female	Donation	
				Multiple	Single
21-30	24	22	2	24	-
31-40	45	44	1	45	-
41-59	46	44	2	40	6
Total	115	110	5	109	6
%	100	96	4	95	5

The majority of them were young males who donated blood many times. Considerable variations in the response of PBL from the individual donors have been observed and are reported in Table 2 below.

As it results from Table 2, 10-30% of PBL cultures may not respond to a 100 μg/ml TTP dose, whereas only 7% cannot be stimulated by a 10 μg/ml PHA dose, and 20% did not respond to a 10 μg/ml LPS dose with IFN production, and 50% did not respond to the same dose of LPS with TNF production. These data are essential for proper selection of biological material. Non-responding PBL cultures are detected by evaluation of the negative (non-treated, spontaneous release of cytokine) and positive (treated with a standard inducer, such as PHA or LPS) controls.

13 FEB 1998

Table 2: IFN and TNF response of PBL from individual blood donors after stimulation either with various batches of TTP or with standard inducers

Inducer	Dose (µg/ml)	No.PBL tested	No. responders to: IFN (%)	TNF (%)
TTP 010391	100	19	13 (68)	13 (68)
	30	13	6 (46)	9 (69)
	10	15	14 (93)	13 (87)
TTP 020391	100	14	10 (71)	11 (79)
	30	12	8 (67)	8 (67)
	10	10	5 (50)	5 (50)
TTP 101091	100	26	21 (81)	20 (77)
	30	20	9 (45)	8 (40)
	10	24	14 (58)	15 (63)
TTP 010991	100	5	5 (100)	4 (80)
	30	4	2 (50)	0 (0)
	10	5	2 (40)	3 (60)
TTP 021091	100	5	4 (80)	4 (80)
	30	4	3 (75)	4 (100)
	10	5	3 (60)	3 (60)
TTP 111191	100	5	3 (60)	4 (80)
	30	4	2 (50)	3 (75)
	10	5	4 (80)	2 (40)
TTP (8x)	100	29	19 (66)	10 (34)
	30	24	14 (58)	11 (46)
	10	21	16 (76)	2 (10)
PHA	10	69	64 (93)	54 (78)
LPS	10	41	33 (80)	21 (51)
None	-	69	36 (52)	20 (29)

REPLACEMENT SHEET
13 FEB 1998

Table 3: Induction of IFN or TNF by different batches of 100 µ/ml peat extract in International Activity Units

	IFN / units/ml			TNF / units/ml				
Series No.	Number of assays	Range	Medi ana	Number of assays	Time of assays 24h Range	Medi ana	6 days Range	Medi ana
Control without induces	333	<10-100	10	41	9-750	27	9-80	9
31090	3	5-30	30	3	ND	ND	9-80	80
291290	21	10-3000	100	14	ND	ND	9-750	40
010391	23	10-2000	60	17	ND	ND	9-160	9
020391	19	10-1000	30	32	9-750	200	9-250	27

The conditioned media were stored at 4°C before assaying

Human peripheral blood leukocytes from healthy blood donors (8x106 cells/ml) were used.

B. Experiments performed.

Peat extract samples taken from different production batches numbered as shown in Table 3 have been tested using the PBL test as described above. In the experiments, interferon (IFN) and tumor necrosis factor (TNF) have been determined qualitatively and according to the above mentioned standard procedure; their activity, as expressed in International Activity Units, was determined. Experiments were repeated a number of times quoted in Table 3. Result ranges and medians are also given.

REPLACEMENT SHEET
13 FEB 1998

For illustration purposes, a similar test was carried out with a standardised peat preparation. At a concentration of 30 µg/ml, interferon activity was within a range of 30 - 300 International Activity Units, while at a concentration of 100 µg/ml, it was 300 - 1000 International Activity Units.

Intermediate concentrations showed a linear relation between dose and response.

From the data given in Table 3 it is clear that in PBL cultures negative and positive control is essential for evaluation of the results.

Further experiments show that, when non-responding PBL cultures are not taken into consideration and statistical evaluation of the large number of experiments is performed, the effectiveness of the tests according to the invention cannot be questioned.

For over two years the cytokine inducing activity of over 20 different batches of TTP, including 10 standard commercial batches of the drug, the activity of which has been determined in biological assays, 2) batches rejected by the manufacturer due to the inadequate biological activity determined in mice which was below the established standard and 3) laboratory peat extracts prepared on a small scale.

The results obtained are presented in Table 4 below in the form of mean levels of IFN and TNF induced (with calculated standard deviations - SD and statistical significance).

REPLACEMENT SHEET
13 FEB 1998

Table 4: Effects of different batches of TTP on IFN and TNF production by human PBL

Inducer	Dose (µg/ml)	IFN Log_{10} units/ml (± SD)	TNF units/ml (± SD)
TTP 010391	100	1.41 ± 1.07[b]	1.26 ± 0.91[c]
	30	0.96 ± 1.07	1.25 ± 0.88[b]
	10	1.81 ± 0.58[c]	1.45 ± 0.63[c]
TTP 020391	100	1.23 ± 0.81[a]	1.35 ± 0.75[c]
	30	1.12 ± 0.90	1.14 ± 0.83[b]
	10	0.75 ± 0.76	0.86 ± 0.90
TTP 101091	100	1.70 ± 0.91[c]	1.38 ± 0.81[c]
	30	0.99 ± 1.12	0.62 ± 0.80
	10	1.22 ± 1.07[a]	0.94 ± 0.76[b]
TTP 010991	100	2.24 ± 0.30[c]	1.35 ± 0.73[b]
	30	1.00 ± 1.00	0.0
	10	0.56 ± 0.68	0.88 ± 0.73
TTP 021091	100	1.72 ± 0.87[b]	1.49 ± 0.84[b]
	30	1.39 ± 0.81	1.79 ± 0.20[c]
	10	1.11 ± 0.91	1.01 ± 0.84
TTP 111191	100	1.36 ± 1.12	1.18 ± 0.62[b]
	30	1.00 ± 1.00	1.21 ± 0.71[a]
	10	1.56 ± 0.86[a]	0.54 ± 0.66
TTP (8x)	100	1.04 ± 0.82[a]	0.63 ± 0.91
	30	0.88 ± 0.81	0.75 ± 0.84
	10	1.10 ± 0.70[b]	0.13 ± 0.40[a]
PHA	10	2.08 ± 0.79[c]	1.67 ± 0.99[c]
LPS	10	1.36 ± 0.80[c]	0.97 ± 1.01[b]
None	-	0.64 ± 0.63	0.42 ± 0.67

Mean values were calculated from all results of IFN or TNF titrations. The non-responders were regarded as 0.

[a-c] Different from "None" (spontaneous production of the cytokine) at [a] $p<0.05$, [b] $p<0.01$ or [c] $p<0.001$.
Mediana values corresponding to the results presented in Table 4 are given in the next Table 5, while selected data

13 FEB 1998

are presented in a graphic form on Fig.5 and 6 showing the effects of different batches of TTP on IFN and TNF production, respectively.

Table 5: Effect of different batches of TTP on IFN and TNF production by human PBL

Inducer	Dose (µg/ml)	IFN Units/ml[1]	TNF Units/ml[1]
TTP 010391	100	30	27
	30	<10	27
	10	100	27
TTP 020391	100	30	50
	30	30	34
	10	<10	<9
TTP 101091	100	100	34
	30	<10	<9
	10	45	18
TTP 010991	100	300	27
	30	<10	<9
	10	<10	18
TTP 021091	100	100	27
	30	60	80
	10	60	27
TTP 111191	100	100	18
	30	<10	40
	10	100	<9
TTP (8x)	100	20	<9
	30	10	<9
	10	20	<9
PHA	10	200	80
LPS	10	30	27
None	-	10	<9

Mediana values were calculated from all results of IFN or TNF titrations.

Identification of the induced cytokines has been performed by means of neutralisation of IFN and TNF induced by TTP.

REPLACEMENT SHEET
13 FEB 1998

The above results of the experiments are explained by Fig. 5 and 6 respectively.

Effects of different batches of TTP on IFN production by human PBL (Fig.5). Seven different batches of TTP were used. Every point of the graph refers to the PBL of an individual healthy blood donor. IFN activity refers to antiviral units. The shaded area shows the limit of the non-significant data. The horizontal lines show mean values. The induction of IFN by 30 and 100 µg/ml of TTP is statistically significant (at $p < 0.05$).

Effects of different batches of TTP on TNF production by human PBL. TNF activity is expressed in the cytotoxic units for the mouse L_{929} cells. The response to 30, 100 and 200 µg/ml of TTP is significant (at $p < 0.05$) (Fig. 6).

In the experiments, potent polyclonal antisera have been used, namely: anti-natural IFN-α, anti-lymphoblastoid (Namalwa) IFN-α and anti-natural IFN-γ to neutralize antiviral activity in supernatants of PBL cultures treated for 20 h with three different batches of TTP. Results, as shown on accompanying drawing Fig.7 (Results of neutralization of IFNs by polyclonal Anti-IFNs sera) indicate that portions of IFN types α and γ produced by PBL from the individual blood donors varied considerably. It is suggested that the released IFN pattern of the each PBL donor is an individual characteristic of the donor.

Neutralisation assays carried out with a potent polyclonal rabbit anti-TNF-α serum (Genzyme Inc.) showed that the TNF induced with TTP is mainly TNF type α. It has been confirmed by the results of similar neutralisations carried out with rabbit polyclonal anti-TNF-ß serum (Genzyme Inc.) which did not neutralise the TTP induced TNF activity.

Fig.7 shows the results of neutralization of IFNs by polyclonal anti-IFNs sera. IFN containing media from PBL

REPLACEMENT SHEET
13 FEB 1998

cultures incubated with TTP (three different batches of TTP and seven different PBL were used) were treated with the indicated anti-IFN sera. 1 - Non treated preparations; 2 - Treated with anti-HuIFN-α; 3 - Treated with anti-HuIFN-Ly; 4 - Treated with anti-HuIFN-γ.

Example 6: The procedure described in Example 5 was followed in order to show the amplifying effect of the presence of an amplifier, viz. indomethacin, according to the present invention.

Each PBL culture was divided into several samples. One of them was a negative control showing spontaneous release of cytokines, the other one was a positive control treated with a standard potent cytokine inducer of a nature indicated below and the remaining samples were treated with 1-100 μg/ml TTP, batch no. 010391 or 020391, as well as TTP I and TTP II which appear in Tables 6b and 6c respectively, and 5 μg/ml indomethacin + TTP of the same batches (Table 6a) or 10 μg/ml compound ITCL + TTP of the same batches (Table 6b), or else 10 or 20 μg/ml seleno-organic compounds (1), (2) or (3) as defined above + TTP of the same batches. In the supernatants of the incubated cultures IFN and TNF were determined.
The results obtained are shown in Tables 6a, b and c below.

Table 6a:
The effect of amplification of the cytokine induction in the presence of indomethacin

Exp.	Inducer	Indo-methacin	Concen-tration μg/ml	Cytokines (units/ml) IFN	TNF
1.a)	TTP 010391	-	100	300	750
	TTP 010391	-	10	30	250
	TTP 010391	-	1	20	80
	TTP 010391	+	100+5	1000	2200
	TTP 010391	+	10+5	200	750

REPLACEMENT SHEET
13 FEB 1998

	TTP 010391	+	1+5	<10	250
	LPS	-	10	200	500
	none	-	-	<10	9
	none	+	5	20	27
2.b)	TTP 020391	-	100	600	50
	TTP 020391	-	40	20	9
	TTP 020391	-	20	10	9
	TTP 020391	-	10	<10	<9
	TTP 020391	+	100+5	600	80
	TTP 020391	+	40+5	100	27
	TTP 020391	+	20+5	100	9
	TTP 020391	+	10+5	30	9
	PHA	-	5	600	50
	PHA	+	5+5	2000	80
	none	-	-	60	<9
	none	+	5	10	<9

a) PBL were prepared from fresh heparinized blood by Ficoll-Hypaque separation technique. PBL cultures contained $7x10^6$ cells/ml,

b) PBL were prepared from the buffy coat obtained from the Regional Transfusion Center. The cells were processed according to the Cantell et al. method (Meth. Enzymol. 1981, 78, 29-38).

PBL cultures contained $8x10^6$ cells/ml.

Table 6b:

The effect of amplification of the cytokine induction in the presence of Compound ITCL

Exp.	Inducer	Compound ITCL	Concen-tration	Cytokines (units/ml) IFN	TNF
1.	TTP 010391	-	100	100	18
	TTP 010391	-	30	100	9
	TTP 010391	+	100+10	3000	80
	TTP 010391	+	30+10	300	80

REPLACEMENT SHEET
13 FEB 1998

	PHA	-	10	3000	250
	none	-	-	10	<9
	none	+	10	<10	<9
2.	TTP I	-	100	3000	27
	TTP I	-	30	30	<9
	TTP I	+	100+10	3000	250
	TTP I	+	30+10	300	9
	PHA	-	10	100	18
	none	-	-	10	<9
	none	+	10	10	<9

Table 6c:

The effect of amplification of the cytokine induction in the presence of the seleno-organic compounds

Exp.	Inducer	Seleno-organic compounds	Concentration µg/ml	Cytokines (units/ml) IFN	TNF
1.	TTPs	-	10	100	<9
	TTPs	-	5	10	<9
	TTPs	Comp. (1)	10+20	300	27
	TTPs	Comp. (1)	5+20	100	27
	TTPs	Comp. (2)	10+20	300	50
	TTPs	Comp. (2)	5+20	100	9
	PHA	-	10	100	<9
	none	Comp. (1)	20	200	9
	none	Comp. (2)	20	100	27
	none	-	-	10	<9
2.	TTP II	-	50	10	<9
	TTP II	-	5	10	<9
	TTP II	Comp. (3)	50+10	30	27
	TTP II	Comp. (3)	10+10	30	27
	PHA	-	-	30	<9
	none	Comp. (3)	10	10	<9
	none	-	-	10	<9

REPLACEMENT SHEET
13 FEB 1998

PBL cultures contained $8x10^6$ cells/ml. TTP is a peat preparation produced on a small scale. The relatively low TNF response is probably due to the fact that the buffy coats were stored for 20 hrs at 4°C before the preparation of the cultures. The seleno-organic compounds are inducers of the cytokines as well as TTP.

Example 7: A diagnostic test reflecting the effect of administration of TTP orally in daily doses of 5 mg in the form of commercially available tablets on PBL in vitro culture response to an additional dose of TTP induction of cytokines was performed according to the particulars given below:

The test was performed with 4 healthy female volunteers, age 43-58, who donated 20 ml venous blood taken from v.ulnaris weekly, before, during and after the administration of TTP at a daily dose of 5 mg in tablets. Two different regimen of administration have been performed. Two of the volunteers (referred to further as B.K. and J.Z.J.) were subjected to a treatment with TTP consisting of three seven-day cycles: the drug was administered per os, one 5 mg tablet daily for one week with a seven day rest followed by a seven-day period of TTP oral administration. The total dosage was 21 tablets. The other two volunteers (referred to further as A.D.I. and W.F.) were treated with one 5 mg TTP tablet daily administered per os continuously for three weeks. The total dosage was also 21 tablets. After two weeks of rest, the treatment was repeated.

The 5 mg. TTP tablets used were a standard commercial product of TORF CORPORATION Pharmaceutical Factory of Wroclaw, Poland. Each tablet contained 5 mg TTP, 43 mg Lactose and 2 mg MYVATEX. To induce cytokines in PBL in vitro cultures, powdered TTP as a pure substance was used in form of a stock solution containing 20 mg TTP/ml of pyrogen free redistilled water stored at 4°C.

This Sheet Has Been Replaced

Blood samples taken from the volunteers were heparinised with Heparin POLFA solution without preservatives at a final concentration of 10 units/ml and treated as described earlier to prepare a PBL in vitro culture, which was then treated with cytokine inducers PHA, LPS and TTP in a manner described in the preceding examples. 200 µl cultures were used in each treatment. The supernatants were stored at 4°C and assayed for IFN within one week, and stored at -20°C and assayed, respectively for TNF to avoid inactivation due to spontaneous proteolysis.

IFN and TNF were assayed in the same manner as described above. The results obtained are presented on the accompanying drawings Figs.8-13 (response of PBL cultures of different patients to IFN inducers and to TNF inducers during the oral administration of TTP).

As may be seen from the data presented on Figs.8 and 9 showing the IFN response of the first group of volunteers during administration cycle, the response declined and returned to the initial value after two weeks rest. In contrast, TNF response shown on Figs.10 and 11 increased during administration of TTP and returned to the normal level after two weeks rest.

Data shown on Figs.12 and 13 reflecting the response to the cytokine induction in the other way of administration of volunteer A.D.I. prove that the hyporeactivity state to IFN induction is reached after 3 weeks of continuous administration of TTP orally. The hyporeactivity state shown as loss of ability of PBL cultures to respond to the induction of IFN with TTP solution disappears after two weeks of rest. In contrast, during three weeks of TTP administration, the hyporeactivity to TNF induction did not develop (Fig.12).

The above results indicate that the optimum dose of immunomodulator being a cytokine inducer can be established

individually for each patient treated with a simple PBL test according to the present invention.

REPLACEMENT SHEET
13 FEB 1998

REPLACEMENT SHEET
13 FEB 1998

* C L M S *

CLAIMS

1. Method for testing bio-active, cytokine inducing immunomodulator substances by determining their immunological activity, characterized in that

a) a cell culture, comprising human peripheral blood leukocytes (PBL) obtained from donors that are not hyporeactive to interferon (IFN) induction, is prepared with a nutrient medium suitable for tissue culture; or

b) a suspension of fresh resident peritoneal cells (RPC) obtained from - preferably 5 to 8 weeks old - healthy "virgin" BALB/c mice that are not immunologically stimulated or that do not show very high spontaneous cytokine production, is prepared with a suitable medium; and

c) non-hyporeactiveness under a) or lack of immunological stimulation or lack of very high spontaneous cytokine production under b) is ensured by positive and negative control; whereupon

d) the cell culture under a) or the cell suspension under b) is treated with a solution of the substance to be tested in order to induce the production of cytokines which then are determined according to standard identification methods.

2. Method according to claim 1, characterized in that the solution to be tested is used in a concentration of 0,1 - 200 µg/ml.

3. Method according to Claim 1, characterized in that in case of the PBL variant the cell density of the ready-for-use culture is approximately 8×10^6 leukocytes/ml, and in case of the RPC variant the cell density is about $1 - 2 \times 10^6$ cells/ml.

4. Method according to any one of the preceding claims, characterized in that the nutrient medium is prepared with

REPLACEMENT SHEET
13 FEB 1998

* C L M S *

- preferably sterile filtered and endotoxin tested - RPMI-1640 medium or EMEM (Eagle's Minimum Essential Medium), plus 10% of non-mitogenic fetal bovine serum (FBS), and optionally further ingredients.

5. Method according to any one of the preceding claims, characterized in that the results obtained are evaluated against a positive and a negative control.

6. Method according to any one of the preceding claims, characterized in that peat extracts or fractions thereof are tested.

7. Method according to claim 6, characterized in that a said peat extract is TTP.

8. Method according to any one of the preceding claims, characterized in that for rendering said method a micro-scale method, amplification of the results is achieved by admixing to the solution to be tested an immunomodulating agent selected from the group consisting of non-steroidal anti-inflammatory and prostaglandin-synthesis inhibiting drugs, organoselenium compounds, isothiazole derivatives, and analogues, homologues and metabolites thereof.

9. Method according to claim 8, characterized in that said agent for amplification is selected from the group consisting of Bis-[2-(N-phenylcarboxamido)]-phenyldiselenide, 2-Phenyl-1,2-benzisoselenazol-3(2H)-one (=Ebselen PZ 51), Bis-[2-(N-(2-pyridyl)carboxamido)]-phenyldiselenide, p-chlorophenylamide of 3-methyl-5-benzoylamino-isothiazole-4-carboxylic acid, and indomethacin.

10. Method for monitoring the immunological response of a patient to a therapy using a cytokine inducing substance, characterized in that

a) cultures of peripheral blood leukocytes (PBL) obtained from blood samples that are taken from said

REPLACEMENT SHEET
13 FEB 1998

* C L M S *

patient in defined time intervals are prepared with a nutrient medium suitable for tissue culture;

b) said PBL cultures are treated with a cytokine inducing substance in order to induce production of cytokines, preferably interferon and tumor necrosis factor; and

c) said cytokines are then determined according to standard methods,

in order to assess development and disappearance of hyporeactivity of said PBL-cultures to interferon induction.

11. Method according to claim 10, characterized in that said time intervals are weekly intervals.

12. Method according to Claim 10 or 11, characterized in that it is applied in the course of a therapy using a cytokine inducing immunomodulator substance, preferably peat extract or a fraction thereof.

13. Method according to Claim 12, characterized in that the peat extract is TTP.

14. Use of TTP in a method as claimed in any one of the preceding claims.

15. Composition for carrying out the method according to any one of claims 1 to 13, characterized in that it contains, on one hand, a PBL culture or a RPC suspension as defined in claim 1, and, on the other hand, a non-steroidal anti-inflammatory and prostaglandin-synthesis inhibiting drug as an amplifier.

16. The composition according to claim 15, characterized in that said amplifier is selected from the group consisting of a seleno-organic compound, an isothiazole derivative, indomethacin and analogues, homologues and metabolites of such substances.