

DESCRIPTION

Title of Invention: CYCLOPROPANAMINE COMPOUND AND USE THEREOF

Technical Field

[0001]

5 The present invention relates to a cyclopropanamine compound having a lysine-specific demethylase-1 (sometimes to be abbreviated as LSD1 in the present specification) inhibitory action, and useful as a medicament such as a prophylactic or therapeutic agent for cancer, schizophrenia,
10 developmental disorders, particularly diseases having intellectual disability (e.g., autistic spectrum disorders, Rett syndrome, Down's syndrome, Kabuki syndrome, fragile X syndrome, Kleeftstra syndrome, neurofibromatosis type 1, Noonan syndrome, tuberous sclerosis), neurodegenerative diseases
15 (e.g., Alzheimer's disease, Parkinson's disease, spinocerebellar degeneration (e.g., dentatorubral pallidoluysian atrophy) and Huntington's disease), epilepsy (e.g., Dravet syndrome) or drug dependence, and the like.

[0002]

20 (Background of the Invention)

 LSD1 is a demethylation enzyme of histone, catalyzes a demethylation reaction of a monomethylated product and a dimethylated product of the 4th lysine residue of histone H3 (H3K4me1/2), and forms formaldehyde as a by-product. In
25 addition, LSD1 forms a complex with flavin adenine dinucleotide (FAD) which is a kind of coenzyme, and FAD promotes oxidation of lysine residue by enzymes as a redox mediator.

[0003]

30 When a compound having a LSD1 inhibitory activity is applied to neuronal cells, histone H3 methylation especially H3K4 methylation around GAD1 gene promoter is increased through inhibition of histone demethylation activity of LSD1 (Experimental Example 3 to be mentioned later). There are many
35 publications with analyses of relationship between gene

expression level and histone H3K4 methylation status, which conclude that promotion of histone H3K4 methylation at a gene promoter leads to an activated transcription of the gene (Becker, Nature 2006, 442: 31-32; Ruthenburg et al., Nature Reviews Molecular Cell Biology 2007, 8: 983-994). Therefore, it is assumed that administration of a compound having an LSD1 inhibitory activity accumulates histone H3K4 methylation in neurons in the brain, which in turn results in the GAD1 mRNA expression in the brain. It is widely known that the induction of GAD1 mRNA expression in the brain is effective for the treatment of central nervous system diseases. For example, intracerebral injection of a GAD1 gene expression vector to Parkinson's disease patients is known to induce GAD1 mRNA expression and improve the symptoms of Parkinson's disease patients (LeWitt et al. Lancet Neurol. 2011, 10: 309-319; Carlson, Physiology of Behavior 11th edition 2013). From the above, it is considered that the administration of an LSD1 inhibitor increases the histone H3K4 methylation and results in increase of the GAD1 expression level in the brain, which may be effective for the treatment of central nervous system diseases.

[0004]


On the other hand, LSD1 also catalyzes a demethylation reaction of a methylated product of the 9th lysine residue of histone H3 (H3K9me). A decrease of the methylation of H3K9 in animal models of some diseases, for example, animal models of cocaine dependence and Kleefstra syndrome has been reported (Science 8 Jan. 2010, 327, 213-216, Genes Dev. Apr. 2005, 19, 815-826). A decrease of the H3K9 methylation is known to cause abnormally enhanced expression of some genes. Therefore, it is assumed that administration of a compound having an LSD1 inhibitory activity accelerates histone H3K9 methylation in neurons in the brain, which in turn decreases the expression of genes abnormally expressed in the brain. From the above, it is considered that LSD1 inhibitor is possibly effective for

the treatment of some diseases with a decrease of methylated H3K9.

[0005]

WO 2010/084160 (patent document 1) discloses a compound of the following formula or a pharmaceutically acceptable salt thereof as an LSD1 inhibitor:

[0006]



FORMULA I


[0007]

wherein R1-R5 are H, halo and the like; R6 is H or alkyl; R7 is H, alkyl or cycloalkyl; R8 is -L-heterocyclyl or -L-aryl wherein L is $-(CH_2)_n-(CH_2)_n-$, $-(CH_2)_nO(CH_2)_n-$ or $-(CH_2)_nS(CH_2)_n-$, and n is 0, 1, 2 or 3.

[0008]

WO 2010/043721 (patent document 2) discloses a compound of the following formula or a pharmaceutically acceptable salt thereof as an LSD1 inhibitor:

[0009]



FORMULA I

[0010]

wherein R1-R5 are H, halo and the like; R6 is H or alkyl; R7

is H, alkyl or cycloalkyl; R8 is -C(=O)NRxRy or -C(=O)Rz wherein Rx and Ry are each independently H, alkyl and the like, and Rz is H, alkoxy and the like.

[0011]

5 WO 2011/035941 (patent document 3) discloses a compound of the following formula or a pharmaceutically acceptable salt thereof as an LSD1 inhibitor:

[0012]



I

10 [0013]

wherein (A') is aryl, arylalkoxy, heterocyclyl and the like; (A) is heteroaryl or aryl; X is 0, 1, 2 or 3; (B) is a cyclopropyl ring; (Z) is -NH-; (L) is -CH₂CH₂- and the like; (D) is -N(-R₁)-R₂, -O-R₃ or -S-R₃ wherein R₁ and R₂ are each
15 independently H, alkyl and the like; and R₃ is H, alkyl and the like.

[0014]

WO 2011/042217 (patent document 4) discloses a compound of the following formula or a pharmaceutically acceptable salt
20 thereof as an LSD1 inhibitor:

[0015]



(I)

[0016]


wherein (A') is aryl, arylalkoxy, arylalkyl, heterocyclyl and
25 the like; (A) is heteroaryl or aryl; X is 0, 1, 2 or 3; (B) is a cyclopropyl ring; (Z) is -NH-; (L) is -(CH₂)_mCR₁R₂- wherein m is 0, 1, 2, 3, 4, 5 or 6; and R₁ and R₂ are each independently H or C₁-6 alkyl.

[0017]

30 US 2010/0324147 (patent document 5) discloses a compound of the following formula or a salt thereof as an LSD1

inhibitor:

[0018]




[0019]

5 wherein X is a bond, O, S or NH; and R_A , R_B , R_C , R_D and R_E are each independently H, C1-7 alkyl and the like.


[0020]

WO 2010/143582 (patent document 6) discloses a compound of the following formula or a pharmaceutically acceptable salt
10 thereof as an LSD1 inhibitor:

[0021]



(I)



(II)

[0022]


wherein R^1 is H, an alkyl group optionally having a substituent
15 attached thereto and the like; R^2 is an alkylene group

optionally having a substituent attached thereto; R³ is an alkyl group optionally having a substituent attached thereto, a phenyl group optionally having a substituent attached thereto and the like; R⁴ is an alkyl group optionally having a substituent attached thereto, a phenyl group optionally having a substituent attached thereto and the like; and X is O, NH₂, NHCO, CONH, S or CH₂.

[0023]


J. Am. Chem. Soc. 2010, 132, 6827-6833 (non-patent document 1) discloses compounds of the following formulas as an LSD 1/2 inhibitor:

[0024]




• HCl

(13b)



(15e)



• HCl

(14e)

[0025]

WO 2012/156531 (patent document 7) discloses use of an LSD1 inhibitor for the prophylaxis or treatment of inflammatory diseases.


[0026]

5 WO 2012/156537 (patent document 8) discloses use of an LSD1 inhibitor for the prophylaxis or treatment of thrombosis, thrombus formation or circulatory diseases.

[0027]

WO 2012/135113 (patent document 9) discloses, as an LSD1
10 inhibitor, a compound of the following formula or a pharmaceutically acceptable salt thereof:

[0028]



[0029]

15 wherein

R₁ is selected from the group consisting of C₁₋₆ alkyl, -NSO₂Me, -NSO₂Ph, arylalkoxy, C₃₋₇ cycloalkyl, -NC(O)Ra, 1-methyl-1H-pyrazol-4-yl, hydroxy, C₁₋₄ alkoxy, halogen, amide, amino, substituted amino and -C(O)ORa;

20 R₂ is H or COOH;

each R₃ is independently selected from the group consisting of aryl, heteroaryl, H, C₁₋₆ alkyl, -SO₂Ra, -NC(O)Ra, -CH₂C(O)ORa, -C(O)ORa, -C(O)Ra, -C(O)NRaRb, substituted amino, amino, urea, amide, sulfonamide, arylalkyl and heteroarylalkyl;

25 each Ra is independently H, phenyl, phenylmethyl, 3,5-dimethylisoxazol-4-yl, 1,2-dimethyl-1H-imidazol-4-yl, C₃₋₇ cycloalkyl, C₁₋₆ alkyl, C₁₋₄ alkoxy, C₁₋₃ alkylamino or -NHPh;
Rb is H or C₁₋₃ alkyl, or when attached to the same atom,

Ra and Rb together form a 5- or 6-membered heterocycloalkyl ring;

R₄ is C₁₋₄ alkyl, acyl, -C(O)CF₃ or H;

W is -(CH₂)₁₋₄ or -CH(Rc) (CH₂)₀₋₃ wherein Rc is CN or C₁₋₄ alkyl;

5 Y is N or C;

X is N or C;

Z is O or (CH₂)_q wherein q is 0-2, and when q is 0, Z is a bond;

m is 0-3, n is 0-3;


10 provided that when Z is O, Y is N and X is C;

also provided that when X is C, at least one of the R₃ groups attached to X is not H.

[0030]

WO 2013/022047 (patent document 10) discloses, as an LSD1
15 inhibitor, the compound of the following formula or a salt thereof.

[0031]



[0032]

20 wherein

A is a hydrocarbon group optionally having substituent(s), or a heterocyclic group optionally having substituent(s);

R is a hydrogen atom, a hydrocarbon group optionally having substituent(s), or a heterocyclic group optionally having

25 substituent(s); or

A and R are optionally bonded to each other to form a ring optionally having substituent(s);

Q¹, Q², Q³ and Q⁴ are each independently a hydrogen atom or a substituent; Q¹ and Q², and Q³ and Q⁴, are each optionally

30 bonded to each other to form a ring optionally having

substituent(s);

X is a hydrogen atom, an acyclic hydrocarbon group optionally having substituent(s), or a saturated cyclic group optionally having substituent(s);

5 Y¹, Y² and Y³ are each independently a hydrogen atom, a hydrocarbon group optionally having substituent(s), or a heterocyclic group optionally having substituent(s);


X and Y¹, and Y¹ and Y², are each optionally bonded to each other to form a ring optionally having substituent(s); and

10 Z¹, Z² and Z³ are each independently a hydrogen atom or a substituent.

[0033]


WO 2012/013727 (patent document 11) discloses, as an LSD1 inhibitor, the compound of the following formula or a salt thereof.

[0034]



(I)

[0035]



(II)

20 [0036]


wherein each symbol is as defined in patent document 11.

[0037]

WO 2013/057322 (patent document 12) discloses, as an LSD1 inhibitor, the compound of the following formula or a salt thereof.

25

[0038]




[0039]

wherein each symbol is as defined in patent document 12.

5 [0040]

WO 2013/057320 (patent document 13) discloses, as an LSD1 inhibitor, the compound of the following formula or a salt thereof.

[0041]



10


[0042]

wherein each symbol is as defined in patent document 13.

[0043]

WO 2014/058071 (patent document 14) discloses, as an LSD1
15 inhibitor, the compound of the following formula or a salt thereof.

[0044]



[0045]

20 wherein each symbol is as defined in patent document 14.

[0046]

The Journal of Neuroscience, October 17, 2007, 27(42):

11254-11262 (non-patent document 2) discloses that a decrease in histone H3K4 methylation and a decrease in Gad1 mRNA expression are observed in the brain of schizophrenia patients. In addition, Nature Neuroscience, Feb. 2015, 18, 199-209 (non-
5 patent document 3) discloses that H3K4 methylation pathway is deeply involved in mental diseases such as schizophrenia and the like.

[0047]

MOLECULAR AND CELLULAR BIOLOGY, Aug. 2011, 31(16), 3298-
10 3311 (non-patent document 4) discloses that LSD1 is a component of a protein complex that regulates transcription of beta globin and potentially involved in the suppression of transcription of beta globin. Activation of beta globin transcription is known to be useful for the treatment of
15 sickle cell anaemia and beta thalassemia, from which it is assumed that LSD1 inhibition results in disinhibition of beta globin transcription, and provides a treatment effect..

Document List

20 patent documents

[0048]

patent document 1: WO 2010/084160

patent document 2: WO 2010/043721

patent document 3: WO 2011/035941

25 patent document 4: WO 2011/042217

patent document 5: US 2010/0324147

patent document 6: WO 2010/143582

patent document 7: WO 2012/156531

patent document 8: WO 2012/156537

30 patent document 9: WO 2012/135113

patent document 10: WO 2013/022047

patent document 11: WO 2012/013727

patent document 12: WO 2013/057322

patent document 13: WO 2013/057320

35 patent document 14: WO 2014/058071

non-patent documents

[0049]

non-patent document 1: J. Am. Chem. Soc. 2010, 132, 6827-6833

non-patent document 2: The Journal of Neuroscience, October 17,
5 2007, 27(42):11254-11262

non-patent document 3: Nature Neuroscience, Feb. 2015, 18,
199-209

non-patent document 4: MOLECULAR AND CELLULAR BIOLOGY, Aug.
2011, 31(16), 3298-3311

10 Summary of the Invention

Problems to be Solved by the Invention

[0050]

An object of the present invention is to provide a
cyclopropanamine compound having a superior LSD1 inhibitory
15 action and high LSD1 selectivity, and useful as a medicament
such as a prophylactic or therapeutic agent for cancer,
schizophrenia, developmental disorders, particularly diseases
having intellectual disability (e.g., autistic spectrum
disorders, Rett syndrome, Down's syndrome, Kabuki syndrome,
20 fragile X syndrome, Kleeftstra syndrome, neurofibromatosis type
1, Noonan syndrome, tuberous sclerosis), neurodegenerative
diseases (e.g., Alzheimer's disease, Parkinson's disease,
spinocerebellar degeneration (e.g., dentatorubral
pallidoluysian atrophy) and Huntington's disease (Huntington
25 chorea)), epilepsy (e.g., Dravet syndrome) or drug dependence
such as cocaine dependence, and the like.

Means of Solving the Problems

[0051]

The present inventors have conducted intensive studies in
30 an attempt to solve the aforementioned problems and found that
a compound represented by the following formula (I) has a
superior LSD1 inhibitory action and high LSD1 selectivity,
which resulted in the completion of the present invention.


[0052]

35 Accordingly, the present invention relates to the

following.

[1] A compound represented by the formula

[0053]



5 [0054]

wherein

A is an optionally substituted heterocyclic group, or an optionally substituted hydrocarbon group;


B is a ring selected from

- 10 (1) a 5- or 6-membered aromatic heterocycle optionally fused with an optionally substituted 5- or 6-membered ring, and
(2) a benzene ring fused with an optionally substituted 5- or 6-membered ring,

wherein the ring represented by B is optionally substituted,

- 15 and binds, via two adjacent carbon atoms with one atom in between, to a group represented by the formula

[0055]




, and

[0056]

- 20 a group represented by the formula

[0057]



;

[0058]

- R¹, R², R³ and R⁴ are each independently a hydrogen atom, an
25 optionally substituted hydrocarbon group or an optionally

substituted heterocyclic group;

A and R¹ are optionally bonded with each other to form, together with the adjacent nitrogen atom, an optionally substituted cyclic group; and

- 5 R² and R³ are optionally bonded with each other to form, together with the adjacent nitrogen atom, an optionally substituted cyclic group,
or a salt thereof (hereinafter to be also referred to as compound (I)).

10 [0059]

[2] The compound of [1], wherein A is

- (1) an optionally substituted heterocyclic group, or
(2) an optionally substituted C₃₋₁₀ cycloalkyl group,
or a salt thereof.

15 [0060]

[3] The compound of [1] or [2], wherein B is a ring selected from

- (1) a 5- or 6-membered aromatic heterocycle, and
(2) a benzene ring fused with an optionally substituted 5- or
20 6-membered ring,


wherein the ring represented by B is optionally substituted, and binds, via two adjacent carbon atoms with one atom in between, to a group represented by the formula (II), and a group represented by the formula (III),

25 or a salt thereof.

[0061]

[4] The compound of [1], [2] or [3], wherein, in the formula


[0062]



30 [0063]

B is a ring selected from

[0064]



[0065]

wherein

X¹, X², X³ and X⁴ are each independently a carbon atom or a
 5 nitrogen atom;

at least one of X¹, X², X³ and X⁴ is a nitrogen atom;

Y¹, Y² and Y³ are each independently a carbon atom, a nitrogen
 atom, an oxygen atom or a sulfur atom;

at least one of Y¹, Y² and Y³ is a nitrogen atom, an oxygen atom
 10 or a sulfur atom; and

Z¹, Z², Z³ and Z⁴ are each independently a carbon atom or a
 nitrogen atom, which ring is optionally substituted, or a salt
 thereof.

[0066]


15 [5] The compound of [1], [2], [3] or [4], wherein R¹, R² and R⁴
 are each independently a hydrogen atom or an optionally
 substituted C₁₋₆ alkyl group,
 or a salt thereof.

[0067]

20 [6] The compound of [1], [2], [3], [4] or [5], wherein R³ is
 (1) a hydrogen atom,
 (2) an optionally substituted C₁₋₆ alkyl group,
 (3) an optionally substituted C₃₋₁₀ cycloalkyl group, or
 (4) an optionally substituted heterocyclic group,
 25 or a salt thereof.

[0068]

[7] The compound of [1], wherein, in the formula
[0069]



[0070]


5 A is

(1) (i) a 5- or 6-membered aromatic heterocyclic group or (ii) a 4- to 6-membered non-aromatic heterocyclic group, each of which is optionally substituted by C₁₋₆ alkyl group(s) optionally substituted by halogen atom(s), or

10 (2) a C₃₋₁₀ cycloalkyl group optionally substituted by halogen atom(s);

B is a ring selected from

[0071]



15 [0072]

wherein

X¹, X², X³ and X⁴ are each independently a carbon atom or a nitrogen atom;

at least one of X¹, X², X³ and X⁴ is a nitrogen atom;

20 Y¹, Y² and Y³ are each independently a carbon atom, a nitrogen atom, an oxygen atom or a sulfur atom;

at least one of Y¹, Y² and Y³ is a nitrogen atom, an oxygen atom or a sulfur atom; and

Z¹, Z², Z³ and Z⁴ are each independently a carbon atom or a nitrogen atom, which ring is optionally substituted by C₁₋₆ alkyl group(s);

R¹ is a hydrogen atom;

5 R² is a hydrogen atom;

R³ is

(1) a hydrogen atom,

(2) a C₁₋₆ alkyl group optionally substituted by substituent(s) selected from

10 (a) a C₃₋₁₀ cycloalkyl group,

(b) a C₆₋₁₄ aryl group optionally substituted by carboxy group(s),

(c) a 4- to 6-membered non-aromatic heterocyclic group optionally substituted by C₁₋₆ alkyl group(s) optionally

15 substituted by substituent(s) selected from a carboxy group, and a C₆₋₁₄ aryl group optionally substituted by carboxy group(s), and

(d) a 5- or 6-membered aromatic heterocyclic group optionally substituted by amino group(s),

20 (3) a C₃₋₁₀ cycloalkyl group optionally substituted by substituent(s) selected from an amino group and a halogen atom or

(4) a 4- to 6-membered non-aromatic heterocyclic group optionally substituted by substituent(s) selected from

25 (a) a C₁₋₆ alkyl group optionally substituted by halogen atom(s),

(b) a C₃₋₁₀ cycloalkyl group,

(c) a C₁₋₆ alkyl-carbonyl group, and

(d) a C₃₋₁₀ cycloalkyl-carbonyl group; and

30 R⁴ is a hydrogen atom,

or a salt thereof.

[0073]

[7A] The compound of [1], wherein A is

(1) (i) a 5- or 6-membered aromatic heterocyclic group or (ii)

35 a 4- to 6-membered non-aromatic heterocyclic group, each of

which is optionally substituted by C₁₋₆ alkyl group(s)

optionally substituted by halogen atom(s), or

(2) a C₃₋₁₀ cycloalkyl group optionally substituted by halogen atom(s);

5 B is a ring selected from

thiophene, thiazole, pyrazole, pyridine, naphthalene and 2,3-dihydrobenzofuran, wherein the ring is optionally substituted by C₁₋₆ alkyl group(s);

R¹ is a hydrogen atom;

10 R² is a hydrogen atom;

R³ is

(1) a hydrogen atom,

(2) a C₁₋₆ alkyl group optionally substituted by substituent(s) selected from

15 (a) a C₃₋₁₀ cycloalkyl group,

(b) a C₆₋₁₄ aryl group optionally substituted by carboxy group(s),

(c) a 4- to 6-membered non-aromatic heterocyclic group optionally substituted by C₁₋₆ alkyl group(s) optionally

20 substituted by substituent(s) selected from a carboxy group,

and a C₆₋₁₄ aryl group optionally substituted by carboxy group(s), and

(d) a 5- or 6-membered aromatic heterocyclic group optionally substituted by amino group(s),

25 (3) a C₃₋₁₀ cycloalkyl group optionally substituted by substituent(s) selected from an amino group and a halogen atom or

(4) a 4- to 6-membered non-aromatic heterocyclic group optionally substituted by substituent(s) selected from

30 (a) a C₁₋₆ alkyl group optionally substituted by halogen atom(s),

(b) a C₃₋₁₀ cycloalkyl group,

(c) a C₁₋₆ alkyl-carbonyl group, and

(d) a C₃₋₁₀ cycloalkyl-carbonyl group; and

35 R⁴ is a hydrogen atom,

or a salt thereof.

[0074]

[8] The compound of [1], wherein A is

- (1) a piperidinyl group, an isoxazolyl group, a pyrazolyl
5 group, a thiadiazolyl group, a thiazolyl group, a
tetrahydropyranyl group, an oxetanyl group, an oxadiazolyl
group, a thienyl group, a pyridyl group or an oxazolyl group,
each of which is optionally substituted by C₁₋₆ alkyl group(s)
optionally substituted by halogen atom(s), or
10 (2) a C₃₋₁₀ cycloalkyl group optionally substituted by halogen
atom(s);

B is a ring selected from

- thiophene, thiazole, pyrazole, pyridine, naphthalene and 2,3-
dihydrobenzofuran, wherein the ring is optionally substituted
15 by C₁₋₆ alkyl group(s);

R¹ is a hydrogen atom;

R² is a hydrogen atom;

R³ is

- (1) a hydrogen atom,
20 (2) a C₁₋₆ alkyl group optionally substituted by substituent(s)
selected from

(a) a C₃₋₁₀ cycloalkyl group,

(b) a C₆₋₁₄ aryl group optionally substituted by carboxy
group(s),

- 25 (c) a tetrahydropyranyl group or a piperidinyl group
each optionally substituted by C₁₋₆ alkyl group(s) optionally
substituted by substituent(s) selected from a carboxy group,
and a C₆₋₁₄ aryl group optionally substituted by carboxy
group(s), and

- 30 (d) an oxadiazolyl group optionally substituted by an
amino group,

(3) a C₃₋₁₀ cycloalkyl group optionally substituted by
substituent(s) selected from an amino group and a halogen atom
or

- 35 (4) a tetrahydropyranyl group or a piperidinyl group each

optionally substituted by substituent(s) selected from

(a) a C₁₋₆ alkyl group optionally substituted by halogen atom(s),

(b) a C₃₋₁₀ cycloalkyl group,

5 (c) a C₁₋₆ alkyl-carbonyl group, and

(d) a C₃₋₁₀ cycloalkyl-carbonyl group; and

R⁴ is a hydrogen atom,

or a salt thereof.

[0075]

10 [8A] The compound of [1], wherein A is

(1) a piperidinyl group, an isoxazolyl group, a pyrazolyl group, a thiadiazolyl group, a thiazolyl group, a tetrahydropyranyl group, an oxetanyl group, an oxadiazolyl group, a thienyl group, a pyridyl group or an oxazolyl group,

15 each of which is optionally substituted by C₁₋₆ alkyl group(s) (preferably, 1 or 2 C₁₋₆ alkyl groups) optionally substituted by halogen atom(s) (preferably, 1 to 3 halogen atoms), or

(2) a cyclopropyl group, a cyclobutyl group, a cyclopentyl group or a cyclohexyl group, each of which is optionally
20 substituted by halogen atom(s) (preferably, 1 or 2 halogen atoms);

B is a ring selected from

thiophene, thiazole, pyrazole, pyridine, naphthalene and 2,3-dihydrobenzofuran, wherein the ring is optionally substituted
25 by C₁₋₆ alkyl group(s) (preferably, one C₁₋₆ alkyl group);

R¹ is a hydrogen atom;

R² is a hydrogen atom;

R³ is

(1) a hydrogen atom,

30 (2) a C₁₋₆ alkyl group optionally substituted by substituent(s) (preferably, one substituent) selected from

(a) a cyclopropyl group or cyclobutyl group,

(b) a phenyl group optionally substituted by carboxy group(s) (preferably, one carboxy group),

35 (c) a tetrahydropyranyl group or a piperidinyl group

each optionally substituted by C₁₋₆ alkyl group(s) (preferably, one C₁₋₆ alkyl group) optionally substituted by substituent(s) (preferably, one substituent) selected from a carboxy group, and a phenyl group optionally substituted by carboxy group(s) (preferably, one carboxy group), and

(d) an oxadiazolyl group optionally substituted by an amino group (one amino group),

(3) a cyclobutyl group or a cyclohexyl group each optionally substituted by substituent(s) (preferably, 1 or 2 substituents) selected from an amino group and a halogen atom, or

(4) a tetrahydropyranyl group or a piperidinyl group each optionally substituted by substituent(s) (preferably, one substituent) selected from

(a) a C₁₋₆ alkyl group optionally substituted by halogen atom(s) (preferably, 1 to 3 halogen atoms),

(b) a cyclopropyl group,

(c) a C₁₋₆ alkyl-carbonyl group, and

(d) a cyclopropyl-carbonyl group; and

R⁴ is a hydrogen atom, or a salt thereof.

[0076]

[9] The compound of [1], wherein A is

(1) a pyrazolyl group, a thiadiazolyl group or a tetrahydropyranyl group, each optionally substituted by C₁₋₆ alkyl group(s) (preferably, 1 or 2 C₁₋₆ alkyl groups), or

(2) a cyclohexyl group optionally substituted by halogen atom(s) (preferably, 1 or 2 halogen atoms);

B is

a thiophene ring, which ring is optionally substituted by C₁₋₆ alkyl group(s) (preferably, one C₁₋₆ alkyl group);

R¹ is a hydrogen atom;

R² is a hydrogen atom;

R³ is a cyclopropylmethyl group, a tetrahydropyranylmethyl group, a cyclobutylmethyl group, a cyclobutyl group or a

tetrahydropyranyl group; and

R⁴ is a hydrogen atom,

or a salt thereof.

[0077]

5 [9A] The compound of [1], wherein A is
a pyrazolyl group, a thiadiazolyl group or a tetrahydropyranyl
group, each optionally substituted by C₁₋₆ alkyl group(s)
(preferably, one C₁₋₆ alkyl group),

B is

10 a thiophene ring, which ring is optionally substituted by C₁₋₆
alkyl group(s) (preferably, one C₁₋₆ alkyl group);

R¹ is a hydrogen atom;

R² is a hydrogen atom;

R³ is a cyclopropylmethyl group, a tetrahydropyranylmethyl
15 group or a cyclobutyl group; and

R⁴ is a hydrogen atom,

or a salt thereof.

[0078]

[10] The compound of [1], wherein A is

20 a pyrazolyl group, a thiadiazolyl group or a tetrahydropyranyl
group, each optionally substituted by C₁₋₆ alkyl group(s)
(preferably, one C₁₋₆ alkyl group),

B is

a thiophene ring, which ring is optionally substituted by C₁₋₆
25 alkyl group(s) (preferably, one C₁₋₆ alkyl group);

R¹ is a hydrogen atom;

R² is a hydrogen atom;

R³ is a cyclopropylmethyl group or a cyclobutyl group; and

R⁴ is a hydrogen atom,

30 or a salt thereof.

[0079]

[11] 5-((1R,2R)-2-((Cyclopropylmethyl)amino)cyclopropyl)-N-
(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide or a salt
thereof.

35 [0080]

[12] 4-((1S,2R)-2-(Cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide or a salt thereof.

[0081]

5 [13] 4-((1S,2R)-2-(Cyclobutylamino)cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide or a salt thereof.

[0082]

[14] A medicament comprising the compound of any one of [1] to 10 [13], [7A], [8A] and [9A] or a salt thereof.

[0083]

[15] The medicament of [14], which is an LSD1 inhibitor.

[0084]

[15A] The medicament of [14], which is a prophylactic or 15 therapeutic agent for schizophrenia, Alzheimer's disease, Parkinson's disease or Huntington's disease (Huntington chorea).

[0085]

[16] The medicament of [14], which is a prophylactic or 20 therapeutic agent for schizophrenia, developmental disorders, autistic spectrum disorders, Rett syndrome, Down's syndrome, Kabuki syndrome, fragile X syndrome, Kleefstra syndrome, neurofibromatosis type 1, Noonan syndrome, tuberous sclerosis, Alzheimer's disease, Parkinson's disease, spinocerebellar 25 degeneration, Huntington's disease, epilepsy or drug dependence.

[0086]

[17] The compound of any one of [1] to [13], [7A], [8A] and [9A] or a salt thereof for use in the prophylaxis or treatment 30 of schizophrenia, developmental disorders, autistic spectrum disorders, Rett syndrome, Down's syndrome, Kabuki syndrome, fragile X syndrome, Kleefstra syndrome, neurofibromatosis type 1, Noonan syndrome, tuberous sclerosis, Alzheimer's disease, Parkinson's disease, spinocerebellar degeneration, 35 Huntington's disease, epilepsy or drug dependence.

[0087]

[18] A method of inhibiting LSD1 in a mammal, comprising administering an effective amount of the compound of any one of [1] to [13], [7A], [8A] and [9A] or a salt thereof to the
5 mammal.

[0088]

[19] A method for the prophylaxis or treatment of schizophrenia, developmental disorders, autistic spectrum disorders, Rett syndrome, Down's syndrome, Kabuki syndrome,
10 fragile X syndrome, Kleefstra syndrome, neurofibromatosis type 1, Noonan syndrome, tuberous sclerosis, Alzheimer's disease, Parkinson's disease, spinocerebellar degeneration, Huntington's disease, epilepsy or drug dependence in a mammal, comprising administering an effective amount of the compound
15 of any one of [1] to [13], [7A], [8A] and [9A] or a salt thereof to the mammal.


[0089]

[20] Use of the compound of any one of [1] to [13], [7A], [8A] and [9A] or a salt thereof in the production of a prophylactic
20 or therapeutic agent for schizophrenia, developmental disorders, autistic spectrum disorders, Rett syndrome, Down's syndrome, Kabuki syndrome, fragile X syndrome, Kleefstra syndrome, neurofibromatosis type 1, Noonan syndrome, tuberous sclerosis, Alzheimer's disease, Parkinson's disease,
25 spinocerebellar degeneration, Huntington's disease, epilepsy or drug dependence.

[0090]

[7B] The compound of [4], wherein B is a ring represented by the formula

30 [0091]



[0092]

wherein

Y¹, Y² and Y³ are each independently a carbon atom, a nitrogen atom, an oxygen atom or a sulfur atom; and at least one of Y¹, Y² and Y³ is a nitrogen atom, an oxygen atom or a sulfur atom, which ring is optionally substituted, or a salt thereof.

[0093]

[8B] The compound of [1], wherein B is a ring selected from thiophene and thiazole, and the ring is optionally substituted, or a salt thereof.

10 [0094]

[9B] The compound of [1], [7B] or [8B], wherein R² is a hydrogen atom and R³ is a cyclopropylmethyl group, or a salt thereof.

[0095]

15 [10B] The compound of [1], [7B] or [8B], wherein R² is a hydrogen atom and R³ is a cyclobutyl group, or a salt thereof.

[0096]

[11B] The compound of [1], [7B] or [8B], wherein R² is a hydrogen atom and R³ is a tetrahydropyranyl group, or a salt

20 thereof.

[0097]

[12B] The compound of [1], [7B] or [8B], wherein R² is a hydrogen atom and R³ is a tetrahydropyranylmethyl group, or a salt thereof.

25 [0098]

[13B] The compound of [1], [7B] or [8B], wherein R² is a hydrogen atom and R³ is an optionally substituted piperidinyl group (preferably, a piperidinyl group optionally substituted by one C₁₋₆ alkyl group optionally substituted by 1 to 3 halogen

30 atoms), or a salt thereof.

[0099]

[14B] The compound of any one of [1] and [7B] to [13B], wherein R¹ is a hydrogen atom, or a salt thereof.

[0100]

35 [15B] The compound of any one of [1] and [7B] to [14B],

wherein A is

- (1) a piperidinyl group, an isoxazolyl group, a pyrazolyl group, a thiadiazolyl group, a thiazolyl group or a tetrahydropyranyl group, each optionally substituted by C₁₋₆ alkyl group(s) (preferably, one C₁₋₆ alkyl group) optionally substituted by halogen atom(s) (preferably, 1 to 3 halogen atoms), or
- (2) a cyclopentyl group or a cyclohexyl group, each optionally substituted by halogen atom(s) (preferably, 1 or 2 halogen atoms), or a salt thereof.

[0101]

The definition of each substituent used in the present specification is described in detail in the following. Unless otherwise specified, each substituent has the following definition.

In the present specification, examples of the "halogen atom" include fluorine, chlorine, bromine and iodine.

In the present specification, examples of the "C₁₋₆ alkyl group" include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, 1-ethylpropyl, hexyl, isohexyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl and 2-ethylbutyl.

In the present specification, examples of the "optionally halogenated C₁₋₆ alkyl group" include a C₁₋₆ alkyl group optionally having 1 to 7, preferably 1 to 5, halogen atoms. Specific examples thereof include methyl, chloromethyl, difluoromethyl, trichloromethyl, trifluoromethyl, ethyl, 2-bromoethyl, 2,2,2-trifluoroethyl, tetrafluoroethyl, pentafluoroethyl, propyl, 2,2-difluoropropyl, 3,3,3-trifluoropropyl, isopropyl, butyl, 4,4,4-trifluorobutyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, 5,5,5-trifluoropentyl, hexyl and 6,6,6-trifluorohexyl.

In the present specification, examples of the "C₂₋₆ alkenyl group" include ethenyl, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 3-methyl-

2-butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 4-methyl-3-pentenyl, 1-hexenyl, 3-hexenyl and 5-hexenyl.

In the present specification, examples of the "C₂₋₆ alkynyl group" include ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-butynyl, 3-butynyl, 1-pentynyl, 2-pentynyl, 3-pentynyl, 4-pentynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl and 4-methyl-2-pentynyl.

In the present specification, examples of the "C₃₋₁₀ cycloalkyl group" include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, bicyclo[2.2.1]heptyl, bicyclo[2.2.2]octyl, bicyclo[3.2.1]octyl and adamantyl.

In the present specification, examples of the "optionally halogenated C₃₋₁₀ cycloalkyl group" include a C₃₋₁₀ cycloalkyl group optionally having 1 to 7, preferably 1 to 5, halogen atoms. Specific examples thereof include cyclopropyl, 2,2-difluorocyclopropyl, 2,3-difluorocyclopropyl, cyclobutyl, difluorocyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

In the present specification, examples of the "C₃₋₁₀ cycloalkenyl group" include cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl and cyclooctenyl.

In the present specification, examples of the "C₆₋₁₄ aryl group" include phenyl, 1-naphthyl, 2-naphthyl, 1-anthryl, 2-anthryl and 9-anthryl.

In the present specification, examples of the "C₇₋₁₆ aralkyl group" include benzyl, phenethyl, naphthylmethyl and phenylpropyl.

[0102]

In the present specification, examples of the "C₁₋₆ alkoxy group" include methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentyloxy and hexyloxy.

In the present specification, examples of the "optionally halogenated C₁₋₆ alkoxy group" include a C₁₋₆ alkoxy group optionally having 1 to 7, preferably 1 to 5, halogen atoms. Specific examples thereof include methoxy, difluoromethoxy,

trifluoromethoxy, ethoxy, 2,2,2-trifluoroethoxy, propoxy, isopropoxy, butoxy, 4,4,4-trifluorobutoxy, isobutoxy, sec-butoxy, pentyloxy and hexyloxy.

In the present specification, examples of the "C₃₋₁₀ cycloalkyloxy group" include cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy, cycloheptyloxy and cyclooctyloxy.

In the present specification, examples of the "C₁₋₆ alkylthio group" include methylthio, ethylthio, propylthio, isopropylthio, butylthio, sec-butylthio, tert-butylthio, pentylthio and hexylthio.

In the present specification, examples of the "optionally halogenated C₁₋₆ alkylthio group" include a C₁₋₆ alkylthio group optionally having 1 to 7, preferably 1 to 5, halogen atoms. Specific examples thereof include methylthio, difluoromethylthio, trifluoromethylthio, ethylthio, propylthio, isopropylthio, butylthio, 4,4,4-trifluorobutylthio, pentylthio and hexylthio.

In the present specification, examples of the "C₁₋₆ alkyl-carbonyl group" include acetyl, propanoyl, butanoyl, 2-methylpropanoyl, pentanoyl, 3-methylbutanoyl, 2-methylbutanoyl, 2,2-dimethylpropanoyl, hexanoyl and heptanoyl.

In the present specification, examples of the "optionally halogenated C₁₋₆ alkyl-carbonyl group" include a C₁₋₆ alkyl-carbonyl group optionally having 1 to 7, preferably 1 to 5, halogen atoms. Specific examples thereof include acetyl, chloroacetyl, trifluoroacetyl, trichloroacetyl, propanoyl, butanoyl, pentanoyl and hexanoyl.

In the present specification, examples of the "C₁₋₆ alkoxy-carbonyl group" include methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl, tert-butoxycarbonyl, pentyloxycarbonyl and hexyloxycarbonyl.

In the present specification, examples of the "C₆₋₁₄ aryl-carbonyl group" include benzoyl, 1-naphthoyl and 2-naphthoyl.

In the present specification, examples of the "C₇₋₁₆ aralkyl-carbonyl group" include phenylacetyl and phenylpropionyl.

In the present specification, examples of the "5- to 14-membered aromatic heterocyclylcarbonyl group" include nicotinoyl, isonicotinoyl, thenoyl and furoyl.

In the present specification, examples of the "3- to 14-membered non-aromatic heterocyclylcarbonyl group" include morpholinylcarbonyl, piperidinylcarbonyl and pyrrolidinylcarbonyl.

[0103]

In the present specification, examples of the "mono- or di-C₁₋₆ alkyl-carbamoyl group" include methylcarbamoyl, ethylcarbamoyl, dimethylcarbamoyl, diethylcarbamoyl and N-ethyl-N-methylcarbamoyl.

In the present specification, examples of the "mono- or di-C₇₋₁₆ aralkyl-carbamoyl group" include benzylcarbamoyl and phenethylcarbamoyl.

In the present specification, examples of the "C₁₋₆ alkylsulfonyl group" include methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, butylsulfonyl, sec-butylsulfonyl and tert-butylsulfonyl.

In the present specification, examples of the "optionally halogenated C₁₋₆ alkylsulfonyl group" include a C₁₋₆ alkylsulfonyl group optionally having 1 to 7, preferably 1 to 5, halogen atoms. Specific examples thereof include methylsulfonyl, difluoromethylsulfonyl, trifluoromethylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, butylsulfonyl, 4,4,4-trifluorobutylsulfonyl, pentylsulfonyl and hexylsulfonyl.

In the present specification, examples of the "C₆₋₁₄ arylsulfonyl group" include phenylsulfonyl, 1-naphthylsulfonyl and 2-naphthylsulfonyl.

[0104]

In the present specification, examples of the

"substituent" include a halogen atom, a cyano group, a nitro group, an optionally substituted hydrocarbon group, an optionally substituted heterocyclic group, an acyl group, an optionally substituted amino group, an optionally substituted carbamoyl group, an optionally substituted thiocarbamoyl group, 5 an optionally substituted sulfamoyl group, an optionally substituted hydroxy group, an optionally substituted sulfanyl (SH) group and an optionally substituted silyl group.

In the present specification, examples of the 10 "hydrocarbon group" (including "hydrocarbon group" of "optionally substituted hydrocarbon group") include a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₂₋₆ alkynyl group, a C₃₋₁₀ cycloalkyl group, a C₃₋₁₀ cycloalkenyl group, a C₆₋₁₄ aryl group and a C₇₋₁₆ aralkyl group.

15 [0105]

In the present specification, examples of the "optionally substituted hydrocarbon group" include a hydrocarbon group optionally having substituent(s) selected from the following substituent group A.

20 [substituent group A]

- (1) a halogen atom,
- (2) a nitro group,
- (3) a cyano group,
- (4) an oxo group,
- 25 (5) a hydroxy group,
- (6) an optionally halogenated C₁₋₆ alkoxy group,
- (7) a C₆₋₁₄ aryloxy group (e.g., phenoxy, naphthoxy),
- (8) a C₇₋₁₆ aralkyloxy group (e.g., benzyloxy),
- (9) a 5- to 14-membered aromatic heterocyclyloxy group (e.g., 30 pyridyloxy),
- (10) a 3- to 14-membered non-aromatic heterocyclyloxy group (e.g., morpholinylloxy, piperidinylloxy),
- (11) a C₁₋₆ alkyl-carbonyloxy group (e.g., acetoxy, propanoyloxy),
- 35 (12) a C₆₋₁₄ aryl-carbonyloxy group (e.g., benzoyloxy, 1-

- naphthoyloxy, 2-naphthoyloxy),
- (13) a C₁₋₆ alkoxy-carbonyloxy group (e.g., methoxycarbonyloxy, ethoxycarbonyloxy, propoxycarbonyloxy, butoxycarbonyloxy),
- (14) a mono- or di-C₁₋₆ alkyl-carbamoyloxy group (e.g.,
- 5 methylcarbamoyloxy, ethylcarbamoyloxy, dimethylcarbamoyloxy, diethylcarbamoyloxy),
- (15) a C₆₋₁₄ aryl-carbamoyloxy group (e.g., phenylcarbamoyloxy, naphthylcarbamoyloxy),
- (16) a 5- to 14-membered aromatic heterocyclylcarbonyloxy
- 10 group (e.g., nicotinoyloxy),
- (17) a 3- to 14-membered non-aromatic heterocyclylcarbonyloxy group (e.g., morpholinylcarbonyloxy, piperidinylcarbonyloxy),
- (18) an optionally halogenated C₁₋₆ alkylsulfonyloxy group (e.g., methylsulfonyloxy, trifluoromethylsulfonyloxy),
- 15 (19) a C₆₋₁₄ arylsulfonyloxy group optionally substituted by a C₁₋₆ alkyl group (e.g., phenylsulfonyloxy, toluenesulfonyloxy),
- (20) an optionally halogenated C₁₋₆ alkylthio group,
- (21) a 5- to 14-membered aromatic heterocyclic group,
- (22) a 3- to 14-membered non-aromatic heterocyclic group,
- 20 (23) a formyl group,
- (24) a carboxy group,
- (25) an optionally halogenated C₁₋₆ alkyl-carbonyl group,
- (26) a C₆₋₁₄ aryl-carbonyl group,
- (27) a 5- to 14-membered aromatic heterocyclylcarbonyl group,
- 25 (28) a 3- to 14-membered non-aromatic heterocyclylcarbonyl group,
- (29) a C₁₋₆ alkoxy-carbonyl group,
- (30) a C₆₋₁₄ aryloxy-carbonyl group (e.g., phenyloxycarbonyl, 1-naphthyloxycarbonyl, 2-naphthyloxycarbonyl),
- 30 (31) a C₇₋₁₆ aralkyloxy-carbonyl group (e.g., benzyloxycarbonyl, phenethyloxycarbonyl),
- (32) a carbamoyl group,
- (33) a thiocarbamoyl group,
- (34) a mono- or di-C₁₋₆ alkyl-carbamoyl group,
- 35 (35) a C₆₋₁₄ aryl-carbamoyl group (e.g., phenylcarbamoyl),

- (36) a 5- to 14-membered aromatic heterocyclylcarbamoyl group (e.g., pyridylcarbamoyl, thienylcarbamoyl),
- (37) a 3- to 14-membered non-aromatic heterocyclylcarbamoyl group (e.g., morpholinylcarbamoyl, piperidinylcarbamoyl),
- 5 (38) an optionally halogenated C₁₋₆ alkylsulfonyl group,
- (39) a C₆₋₁₄ arylsulfonyl group,
- (40) a 5- to 14-membered aromatic heterocyclylsulfonyl group (e.g., pyridylsulfonyl, thienylsulfonyl),
- (41) an optionally halogenated C₁₋₆ alkylsulfinyl group,
- 10 (42) a C₆₋₁₄ arylsulfinyl group (e.g., phenylsulfinyl, 1-naphthylsulfinyl, 2-naphthylsulfinyl),
- (43) a 5- to 14-membered aromatic heterocyclylsulfinyl group (e.g., pyridylsulfinyl, thienylsulfinyl),
- (44) an amino group,
- 15 (45) a mono- or di-C₁₋₆ alkylamino group (e.g., methylamino, ethylamino, propylamino, isopropylamino, butylamino, dimethylamino, diethylamino, dipropylamino, dibutylamino, N-ethyl-N-methylamino),
- (46) a mono- or di-C₆₋₁₄ arylamino group (e.g., phenylamino),
- 20 (47) a 5- to 14-membered aromatic heterocyclylamino group (e.g., pyridylamino),
- (48) a C₇₋₁₆ aralkylamino group (e.g., benzylamino),
- (49) a formylamino group,
- (50) a C₁₋₆ alkyl-carbonylamino group (e.g., acetylamino,
- 25 propanoylamino, butanoylamino),
- (51) a (C₁₋₆ alkyl) (C₁₋₆ alkyl-carbonyl)amino group (e.g., N-acetyl-N-methylamino),
- (52) a C₆₋₁₄ aryl-carbonylamino group (e.g., phenylcarbonylamino, naphthylcarbonylamino),
- 30 (53) a C₁₋₆ alkoxy-carbonylamino group (e.g., methoxycarbonylamino, ethoxycarbonylamino, propoxycarbonylamino, butoxycarbonylamino, tert-butoxycarbonylamino),
- (54) a C₇₋₁₆ aralkyloxy-carbonylamino group (e.g.,
- 35 benzyloxycarbonylamino),

(55) a C₁₋₆ alkylsulfonylamino group (e.g., methylsulfonylamino, ethylsulfonylamino),

(56) a C₆₋₁₄ arylsulfonylamino group optionally substituted by a C₁₋₆ alkyl group (e.g., phenylsulfonylamino,

5 toluenesulfonylamino),

(57) an optionally halogenated C₁₋₆ alkyl group,

(58) a C₂₋₆ alkenyl group,

(59) a C₂₋₆ alkynyl group,

(60) a C₃₋₁₀ cycloalkyl group,

10 (61) a C₃₋₁₀ cycloalkenyl group and

(62) a C₆₋₁₄ aryl group.

[0106]

The number of the above-mentioned substituents in the "optionally substituted hydrocarbon group" is, for example, 1
15 to 5, preferably 1 to 3. When the number of the substituents is two or more, the respective substituents may be the same or different.

In the present specification, examples of the "heterocyclic group" (including "heterocyclic group" of
20 "optionally substituted heterocyclic group") include (i) an aromatic heterocyclic group, (ii) a non-aromatic heterocyclic group and (iii) a 7- to 10-membered bridged heterocyclic group, each containing, as a ring-constituting atom besides carbon atom, 1 to 4 hetero atoms selected from a nitrogen atom, a
25 sulfur atom and an oxygen atom.

[0107]

In the present specification, examples of the "aromatic heterocyclic group" (including "5- to 14-membered aromatic heterocyclic group") include a 5- to 14-membered (preferably
30 5- to 10-membered) aromatic heterocyclic group containing, as a ring-constituting atom besides carbon atom, 1 to 4 hetero atoms selected from a nitrogen atom, a sulfur atom and an oxygen atom.

Preferable examples of the "aromatic heterocyclic group"
35 include 5- or 6-membered monocyclic aromatic heterocyclic

groups such as thienyl, furyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, triazolyl, 5 tetrazolyl, triazinyl and the like; and 8- to 14-membered fused polycyclic (preferably bi or tricyclic) aromatic heterocyclic groups such as benzothiophenyl, benzofuranyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, 10 benzotriazolyl, imidazopyridinyl, thienopyridinyl, furopyridinyl, pyrrolopyridinyl, pyrazolopyridinyl, oxazolopyridinyl, thiazolopyridinyl, imidazopyrazinyl, imidazopyrimidinyl, thienopyrimidinyl, furopyrimidinyl, pyrrolopyrimidinyl, pyrazolopyrimidinyl, oxazolopyrimidinyl, 15 thiazolopyrimidinyl, pyrazolotriazinyl, naphtho[2,3-b]thienyl, phenoxathiinyl, indolyl, isoindolyl, 1H-indazolyl, purinyl, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, carbazolyl, β -carbolinyl, phenanthridinyl, acridinyl, phenazinyl, 20 phenothiazinyl, phenoxazinyl and the like.

[0108]

In the present specification, examples of the "non-aromatic heterocyclic group" (including "3- to 14-membered non-aromatic heterocyclic group") include a 3- to 14-membered 25 (preferably 4- to 10-membered) non-aromatic heterocyclic group containing, as a ring-constituting atom besides carbon atom, 1 to 4 hetero atoms selected from a nitrogen atom, a sulfur atom and an oxygen atom.

Preferable examples of the "non-aromatic heterocyclic 30 group" include 3- to 8-membered monocyclic non-aromatic heterocyclic groups such as aziridinyl, oxiranyl, thiranyl, azetidiny, oxetanyl, thietanyl, tetrahydrothienyl, tetrahydrofuranyl, pyrrolinyl, pyrrolidinyl, imidazolinyl, imidazolidinyl, oxazolinyl, oxazolidinyl, pyrazolinyl, 35 pyrazolidinyl, thiazolinyl, thiazolidinyl,

tetrahydroisothiazolyl, tetrahydrooxazolyl,
tetrahydroisooxazolyl, piperidinyl, piperazinyl,
tetrahydropyridinyl, dihydropyridinyl, dihydrothiopyranyl,
tetrahydropyrimidinyl, tetrahydropyridazinyl, dihydropyranyl,
5 tetrahydropyranyl, tetrahydrothiopyranyl, morpholinyl,
thiomorpholinyl, azepanyl, diazepanyl, azepinyl, oxepanyl,
azocanyl, diazocanyl and the like; and
9- to 14-membered fused polycyclic (preferably bi or
tricyclic) non-aromatic heterocyclic groups such as
10 dihydrobenzofuranyl, dihydrobenzimidazolyl,
dihydrobenzoxazolyl, dihydrobenzothiazolyl,
dihydrobenzisoctiazolyl, dihydronaphtho[2,3-b]thienyl,
tetrahydroisoquinolyl, tetrahydroquinolyl, 4H-quinolizinyl,
indolinyl, isoindolinyl, tetrahydrothieno[2,3-c]pyridinyl,
15 tetrahydrobenzazepinyl, tetrahydroquinoxalinyl,
tetrahydrophenanthridinyl, hexahydrophenothiazinyl,
hexahydrophenoxazinyl, tetrahydrophthalazinyl,
tetrahydronaphthyridinyl, tetrahydroquinazolinyl,
tetrahydrocinnolinyl, tetrahydrocarbazolyl, tetrahydro- β -
20 carbolinyl, tetrahydroacrydinyl, tetrahydrophenazinyl,
tetrahydrothioxanthenyl, octahydroisoquinolyl and the like.
[0109]

In the present specification, preferable examples of the
"7- to 10-membered bridged heterocyclic group" include
25 quinuclidinyl and 7-azabicyclo[2.2.1]heptanyl.

In the present specification, examples of the "nitrogen-
containing heterocyclic group" include a "heterocyclic group"
containing at least one nitrogen atom as a ring-constituting
atom.

30 In the present specification, examples of the "optionally
substituted heterocyclic group" include a heterocyclic group
optionally having substituent(s) selected from the
aforementioned substituent group A.

The number of the substituents in the "optionally
35 substituted heterocyclic group" is, for example, 1 to 3. When

the number of the substituents is two or more, the respective substituents may be the same or different.

[0110]

In the present specification, examples of the "acyl group" include a formyl group, a carboxy group, a carbamoyl group, a thiocarbamoyl group, a sulfinio group, a sulfo group, a sulfamoyl group and a phosphono group, each optionally having "1 or 2 substituents selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₃₋₁₀ cycloalkenyl group, a C₆₋₁₄ aryl group, a C₇₋₁₆ aralkyl group, a 5- to 14-membered aromatic heterocyclic group and a 3- to 14-membered non-aromatic heterocyclic group, each of which optionally has 1 to 3 substituents selected from a halogen atom, an optionally halogenated C₁₋₆ alkoxy group, a hydroxy group, a nitro group, a cyano group, an amino group and a carbamoyl group".

Examples of the "acyl group" also include a hydrocarbon-sulfonyl group, a heterocyclylsulfonyl group, a hydrocarbon-sulfinyl group and a heterocyclylsulfinyl group.

Here, the hydrocarbon-sulfonyl group means a hydrocarbon group-bonded sulfonyl group, the heterocyclylsulfonyl group means a heterocyclic group-bonded sulfonyl group, the hydrocarbon-sulfinyl group means a hydrocarbon group-bonded sulfinyl group and the heterocyclylsulfinyl group means a heterocyclic group-bonded sulfinyl group.

Preferable examples of the "acyl group" include a formyl group, a carboxy group, a C₁₋₆ alkyl-carbonyl group, a C₂₋₆ alkenyl-carbonyl group (e.g., crotonoyl), a C₃₋₁₀ cycloalkyl-carbonyl group (e.g., cyclobutanecarbonyl, cyclopentanecarbonyl, cyclohexanecarbonyl, cycloheptanecarbonyl), a C₃₋₁₀ cycloalkenyl-carbonyl group (e.g., 2-cyclohexenecarbonyl), a C₆₋₁₄ aryl-carbonyl group, a C₇₋₁₆ aralkyl-carbonyl group, a 5- to 14-membered aromatic heterocyclylcarbonyl group, a 3- to 14-membered non-aromatic heterocyclylcarbonyl group, a C₁₋₆ alkoxy-carbonyl group, a C₆₋₁₄

aryloxy-carbonyl group (e.g., phenyloxycarbonyl,
 naphthyloxycarbonyl), a C₇₋₁₆ aralkyloxy-carbonyl group (e.g.,
 benzyloxycarbonyl, phenethyloxycarbonyl), a carbamoyl group, a
 mono- or di-C₁₋₆ alkyl-carbamoyl group, a mono- or di-C₂₋₆
 5 alkenyl-carbamoyl group (e.g., diallylcarbamoyl), a mono- or
 di-C₃₋₁₀ cycloalkyl-carbamoyl group (e.g., cyclopropylcarbamoyl),
 a mono- or di-C₆₋₁₄ aryl-carbamoyl group (e.g., phenylcarbamoyl),
 a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, a 5- to 14-membered
 aromatic heterocyclylcarbamoyl group (e.g., pyridylcarbamoyl),
 10 a thiocarbamoyl group, a mono- or di-C₁₋₆ alkyl-thiocarbamoyl
 group (e.g., methylthiocarbamoyl, N-ethyl-N-
 methylthiocarbamoyl), a mono- or di-C₂₋₆ alkenyl-thiocarbamoyl
 group (e.g., diallylthiocarbamoyl), a mono- or di-C₃₋₁₀
 cycloalkyl-thiocarbamoyl group (e.g., cyclopropylthiocarbamoyl,
 15 cyclohexylthiocarbamoyl), a mono- or di-C₆₋₁₄ aryl-thiocarbamoyl
 group (e.g., phenylthiocarbamoyl), a mono- or di-C₇₋₁₆ aralkyl-
 thiocarbamoyl group (e.g., benzylthiocarbamoyl,
 phenethylthiocarbamoyl), a 5- to 14-membered aromatic
 heterocyclylthiocarbamoyl group (e.g., pyridylthiocarbamoyl),
 20 a sulfinio group, a C₁₋₆ alkylsulfinyl group (e.g.,
 methylsulfinyl, ethylsulfinyl), a sulfo group, a C₁₋₆
 alkylsulfonyl group, a C₆₋₁₄ arylsulfonyl group, a phosphono
 group and a mono- or di-C₁₋₆ alkylphosphono group (e.g.,
 dimethylphosphono, diethylphosphono, diisopropylphosphono,
 25 dibutylphosphono).

[0111]

In the present specification, examples of the "optionally
 substituted amino group" include an amino group optionally
 having "1 or 2 substituents selected from a C₁₋₆ alkyl group, a
 30 C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group,
 a C₇₋₁₆ aralkyl group, a C₁₋₆ alkyl-carbonyl group, a C₆₋₁₄ aryl-
 carbonyl group, a C₇₋₁₆ aralkyl-carbonyl group, a 5- to 14-
 membered aromatic heterocyclylcarbonyl group, a 3- to 14-
 membered non-aromatic heterocyclylcarbonyl group, a C₁₋₆ alkoxy-
 35 carbonyl group, a 5- to 14-membered aromatic heterocyclic

group, a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group, a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, a C₁₋₆ alkylsulfonyl group and a C₆₋₁₄ arylsulfonyl group, each of which optionally has 1 to 3 substituents selected from
5 substituent group A".

Preferable examples of the optionally substituted amino group include an amino group, a mono- or di-(optionally halogenated C₁₋₆ alkyl)amino group (e.g., methylamino, trifluoromethylamino, dimethylamino, ethylamino, diethylamino,
10 propylamino, dibutylamino), a mono- or di-C₂₋₆ alkenylamino group (e.g., diallylamino), a mono- or di-C₃₋₁₀ cycloalkylamino group (e.g., cyclopropylamino, cyclohexylamino), a mono- or di-C₆₋₁₄ arylamino group (e.g., phenylamino), a mono- or di-C₇₋₁₆ aralkylamino group (e.g., benzylamino, dibenzylamino), a mono-
15 or di-(optionally halogenated C₁₋₆ alkyl)-carbonylamino group (e.g., acetylamino, propionylamino), a mono- or di-C₆₋₁₄ aryl-carbonylamino group (e.g., benzoylamino), a mono- or di-C₇₋₁₆ aralkyl-carbonylamino group (e.g., benzylcarbonylamino), a mono- or di-5- to 14-membered aromatic
20 heterocyclylcarbonylamino group (e.g., nicotinoylamino, isonicotinoylamino), a mono- or di-3- to 14-membered non-aromatic heterocyclylcarbonylamino group (e.g., piperidinylcarbonylamino), a mono- or di-C₁₋₆ alkoxy-carbonylamino group (e.g., tert-butoxycarbonylamino), a 5- to
25 14-membered aromatic heterocyclylamino group (e.g., pyridylamino), a carbamoylamino group, a (mono- or di-C₁₋₆ alkyl-carbamoyl)amino group (e.g., methylcarbamoylamino), a (mono- or di-C₇₋₁₆ aralkyl-carbamoyl)amino group (e.g., benzylcarbamoylamino), a C₁₋₆ alkylsulfonylamino group (e.g.,
30 methylsulfonylamino, ethylsulfonylamino), a C₆₋₁₄ arylsulfonylamino group (e.g., phenylsulfonylamino), a (C₁₋₆ alkyl)(C₁₋₆ alkyl-carbonyl)amino group (e.g., N-acetyl-N-methylamino) and a (C₁₋₆ alkyl)(C₆₋₁₄ aryl-carbonyl)amino group (e.g., N-benzoyl-N-methylamino).

35 [0112]

In the present specification, examples of the "optionally substituted carbamoyl group" include a carbamoyl group optionally having "1 or 2 substituents selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group, a C₇₋₁₆ aralkyl group, a C₁₋₆ alkyl-carbonyl group, a C₆₋₁₄ aryl-carbonyl group, a C₇₋₁₆ aralkyl-carbonyl group, a 5- to 14-membered aromatic heterocyclylcarbonyl group, a 3- to 14-membered non-aromatic heterocyclylcarbonyl group, a C₁₋₆ alkoxy-carbonyl group, a 5- to 14-membered aromatic heterocyclic group, a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group and a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, each of which optionally has 1 to 3 substituents selected from substituent group A".

Preferable examples of the optionally substituted carbamoyl group include a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group, a mono- or di-C₂₋₆ alkenyl-carbamoyl group (e.g., diallylcarbamoyl), a mono- or di-C₃₋₁₀ cycloalkyl-carbamoyl group (e.g., cyclopropylcarbamoyl, cyclohexylcarbamoyl), a mono- or di-C₆₋₁₄ aryl-carbamoyl group (e.g., phenylcarbamoyl), a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbonyl-carbamoyl group (e.g., acetylcarbamoyl, propionylcarbamoyl), a mono- or di-C₆₋₁₄ aryl-carbonyl-carbamoyl group (e.g., benzoylcarbamoyl) and a 5- to 14-membered aromatic heterocyclylcarbamoyl group (e.g., pyridylcarbamoyl).

[0113]

In the present specification, examples of the "optionally substituted thiocarbamoyl group" include a thiocarbamoyl group optionally having "1 or 2 substituents selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group, a C₇₋₁₆ aralkyl group, a C₁₋₆ alkyl-carbonyl group, a C₆₋₁₄ aryl-carbonyl group, a C₇₋₁₆ aralkyl-carbonyl group, a 5- to 14-membered aromatic heterocyclylcarbonyl group, a 3- to 14-membered non-aromatic heterocyclylcarbonyl group, a C₁₋₆ alkoxy-carbonyl group, a 5- to 14-membered aromatic

heterocyclic group, a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group and a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, each of which optionally has 1 to 3 substituents selected from substituent group A".

5 Preferable examples of the optionally substituted thiocarbamoyl group include a thiocarbamoyl group, a mono- or di-C₁₋₆ alkyl-thiocarbamoyl group (e.g., methylthiocarbamoyl, ethylthiocarbamoyl, dimethylthiocarbamoyl, diethylthiocarbamoyl, N-ethyl-N-methylthiocarbamoyl), a mono-
10 or di-C₂₋₆ alkenyl-thiocarbamoyl group (e.g., diallylthiocarbamoyl), a mono- or di-C₃₋₁₀ cycloalkyl-thiocarbamoyl group (e.g., cyclopropylthiocarbamoyl, cyclohexylthiocarbamoyl), a mono- or di-C₆₋₁₄ aryl-thiocarbamoyl group (e.g., phenylthiocarbamoyl), a mono- or di-C₇₋₁₆ aralkyl-
15 thiocarbamoyl group (e.g., benzylthiocarbamoyl, phenethylthiocarbamoyl), a mono- or di-C₁₋₆ alkyl-carbonyl-thiocarbamoyl group (e.g., acetylthiocarbamoyl, propionylthiocarbamoyl), a mono- or di-C₆₋₁₄ aryl-carbonyl-thiocarbamoyl group (e.g., benzoylthiocarbamoyl) and a 5- to
20 14-membered aromatic heterocyclylthiocarbamoyl group (e.g., pyridylthiocarbamoyl).

[0114]

In the present specification, examples of the "optionally substituted sulfamoyl group" include a sulfamoyl group
25 optionally having "1 or 2 substituents selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group, a C₇₋₁₆ aralkyl group, a C₁₋₆ alkyl-carbonyl group, a C₆₋₁₄ aryl-carbonyl group, a C₇₋₁₆ aralkyl-carbonyl group, a 5- to 14-membered aromatic heterocyclylcarbonyl group, a 3- to
30 14-membered non-aromatic heterocyclylcarbonyl group, a C₁₋₆ alkoxy-carbonyl group, a 5- to 14-membered aromatic heterocyclic group, a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group and a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, each of which optionally has 1 to 3 substituents selected from
35 substituent group A".

Preferable examples of the optionally substituted sulfamoyl group include a sulfamoyl group, a mono- or di-C₁₋₆ alkyl-sulfamoyl group (e.g., methylsulfamoyl, ethylsulfamoyl, dimethylsulfamoyl, diethylsulfamoyl, N-ethyl-N-methylsulfamoyl), a mono- or di-C₂₋₆ alkenyl-sulfamoyl group (e.g., diallylsulfamoyl), a mono- or di-C₃₋₁₀ cycloalkyl-sulfamoyl group (e.g., cyclopropylsulfamoyl, cyclohexylsulfamoyl), a mono- or di-C₆₋₁₄ aryl-sulfamoyl group (e.g., phenylsulfamoyl), a mono- or di-C₇₋₁₆ aralkyl-sulfamoyl group (e.g., benzylsulfamoyl, phenethylsulfamoyl), a mono- or di-C₁₋₆ alkyl-carbonyl-sulfamoyl group (e.g., acetylsulfamoyl, propionylsulfamoyl), a mono- or di-C₆₋₁₄ aryl-carbonyl-sulfamoyl group (e.g., benzoylsulfamoyl) and a 5- to 14-membered aromatic heterocyclylsulfamoyl group (e.g., pyridylsulfamoyl).

15 [0115]

In the present specification, examples of the "optionally substituted hydroxy group" include a hydroxyl group optionally having "a substituent selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group, a C₇₋₁₆ aralkyl group, a C₁₋₆ alkyl-carbonyl group, a C₆₋₁₄ aryl-carbonyl group, a C₇₋₁₆ aralkyl-carbonyl group, a 5- to 14-membered aromatic heterocyclylcarbonyl group, a 3- to 14-membered non-aromatic heterocyclylcarbonyl group, a C₁₋₆ alkoxy-carbonyl group, a 5- to 14-membered aromatic heterocyclic group, a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group, a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, a C₁₋₆ alkylsulfonyl group and a C₆₋₁₄ arylsulfonyl group, each of which optionally has 1 to 3 substituents selected from substituent group A".

Preferable examples of the optionally substituted hydroxy group include a hydroxy group, a C₁₋₆ alkoxy group, a C₂₋₆ alkenyloxy group (e.g., allyloxy, 2-butenyloxy, 2-pentenylloxy, 3-hexenyloxy), a C₃₋₁₀ cycloalkyloxy group (e.g., cyclohexyloxy), a C₆₋₁₄ aryloxy group (e.g., phenoxy, naphthyloxy), a C₇₋₁₆ aralkyloxy group (e.g., benzyloxy, phenethylloxy), a C₁₋₆ alkyl-

carbonyloxy group (e.g., acetyloxy, propionyloxy, butyryloxy, isobutyryloxy, pivaloyloxy), a C₆₋₁₄ aryl-carbonyloxy group (e.g., benzoyloxy), a C₇₋₁₆ aralkyl-carbonyloxy group (e.g., benzylcarbonyloxy), a 5- to 14-membered aromatic
5 heterocyclylcarbonyloxy group (e.g., nicotinoyloxy), a 3- to 14-membered non-aromatic heterocyclylcarbonyloxy group (e.g., piperidinylcarbonyloxy), a C₁₋₆ alkoxy-carbonyloxy group (e.g., tert-butoxycarbonyloxy), a 5- to 14-membered aromatic heterocyclyloxy group (e.g., pyridyloxy), a carbamoyloxy group,
10 a C₁₋₆ alkyl-carbamoyloxy group (e.g., methylcarbamoyloxy), a C₇₋₁₆ aralkyl-carbamoyloxy group (e.g., benzylcarbamoyloxy), a C₁₋₆ alkylsulfonyloxy group (e.g., methylsulfonyloxy, ethylsulfonyloxy) and a C₆₋₁₄ arylsulfonyloxy group (e.g., phenylsulfonyloxy).

15 [0116]

In the present specification, examples of the "optionally substituted sulfanyl group" include a sulfanyl group optionally having "a substituent selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄
20 aryl group, a C₇₋₁₆ aralkyl group, a C₁₋₆ alkyl-carbonyl group, a C₆₋₁₄ aryl-carbonyl group and a 5- to 14-membered aromatic heterocyclic group, each of which optionally has 1 to 3 substituents selected from substituent group A" and a halogenated sulfanyl group.

25 Preferable examples of the optionally substituted sulfanyl group include a sulfanyl (-SH) group, a C₁₋₆ alkylthio group, a C₂₋₆ alkenylthio group (e.g., allylthio, 2-butenylthio, 2-pentenylthio, 3-hexenylthio), a C₃₋₁₀ cycloalkylthio group (e.g., cyclohexylthio), a C₆₋₁₄ arylthio group (e.g., phenylthio,
30 naphthylthio), a C₇₋₁₆ aralkylthio group (e.g., benzylthio, phenethylthio), a C₁₋₆ alkyl-carbonylthio group (e.g., acetylthio, propionylthio, butyrylthio, isobutyrylthio, pivaloylthio), a C₆₋₁₄ aryl-carbonylthio group (e.g., benzoylthio), a 5- to 14-membered aromatic heterocyclylthio
35 group (e.g., pyridylthio) and a halogenated thio group (e.g.,

pentafluorothio).

[0117]

In the present specification, examples of the "optionally substituted silyl group" include a silyl group optionally
5 having "1 to 3 substituents selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group and a C₇₋₁₆ aralkyl group, each of which optionally has 1 to 3 substituents selected from substituent group A".

Preferable examples of the optionally substituted silyl
10 group include a tri-C₁₋₆ alkylsilyl group (e.g., trimethylsilyl, tert-butyl(dimethyl)silyl).

[0118]

In the present specification, examples of the "5- or 6-membered aromatic heterocycle" include a 5- or 6-membered
15 aromatic heterocycle containing, as a ring-constituting atom besides carbon atoms, 1 to 4 hetero atoms selected from a nitrogen atom, a sulfur atom and an oxygen atom. Preferable examples of the "5- or 6-membered aromatic heterocycle" include 5- or 6-membered aromatic heterocycles such as
20 thiophene, furan, pyrrole, imidazole, pyrazole, thiazole, isothiazole, oxazole, isoxazole, pyridine, pyrazine, pyrimidine, pyridazine, 1,2,4-oxadiazole, 1,3,4-oxadiazole, 1,2,4-thiadiazole, 1,3,4-thiadiazole, triazole, triazine and the like.

25 [0119]

A is an optionally substituted heterocyclic group, or an optionally substituted hydrocarbon group.

[0120]

A is preferably

- 30 (1) an optionally substituted heterocyclic group, or
(2) an optionally substituted C₃₋₁₀ cycloalkyl group.

[0121]

A is more preferably

- (1) a heterocyclic group (preferably, (i) a 5- or 6-membered
35 aromatic heterocyclic group or (ii) a 4- to 6-membered non-

aromatic heterocyclic group, each containing, as a ring-constituting atom besides carbon atoms, 1 to 4 hetero atoms selected from a nitrogen atom, a sulfur atom and an oxygen atom) (e.g., piperidinyl group, isoxazolyl group, pyrazolyl group, thiadiazolyl group, thiazolyl group, tetrahydropyranyl group, oxetanyl group, oxadiazolyl group, thienyl group, pyridyl group, oxazolyl group) optionally substituted by C₁₋₆ alkyl group(s) (preferably, 1 or 2 C₁₋₆ alkyl groups) optionally substituted by halogen atom(s) (preferably, 1 to 3 halogen atoms), or
(2) a C₃₋₁₀ cycloalkyl group (e.g., cyclopropyl group, cyclobutyl group, cyclopentyl group, cyclohexyl group) optionally substituted by halogen atom(s) (preferably, 1 or 2 halogen atoms).

[0122]

A is more preferably

(1) a piperidinyl group, an isoxazolyl group, a pyrazolyl group, a thiadiazolyl group, a thiazolyl group or a tetrahydropyranyl group, each optionally substituted by C₁₋₆ alkyl group(s) (preferably, one C₁₋₆ alkyl group) optionally substituted by halogen atom(s) (preferably, 1 to 3 halogen atoms), or
(2) a cyclopentyl group or a cyclohexyl group, each optionally substituted by halogen atom(s) (preferably, 1 or 2 halogen atoms).

[0123]

A is particularly preferably

(1) a pyrazolyl group, a thiadiazolyl group or a tetrahydropyranyl group, each optionally substituted by C₁₋₆ alkyl group(s) (preferably, one C₁₋₆ alkyl group), or
(2) a cyclopentyl group or a cyclohexyl group, each optionally substituted by halogen atom(s) (preferably, 1 or 2 halogen atoms).


[0124]

B is a ring selected from

(1) a 5- or 6-membered aromatic heterocycle optionally fused with an optionally substituted 5- or 6-membered ring, and
(2) a benzene ring fused with an optionally substituted 5- or 6-membered ring,

5 and the ring represented by B is optionally substituted and binds, via two adjacent carbon atoms with one atom in between, to a group represented by the formula


[0125]



10 [0126]

and a group represented by the formula

[0127]



[0128]

15 Examples of the "optionally substituted 5- or 6-membered ring" include a 5- or 6-membered ring optionally having substituent(s) selected from the aforementioned substituent group A.

[0129]

20 The number of the substituents for the "optionally substituted 5- or 6-membered ring" is, for example, 1 to 3. When the number of the substituents is two or more, respective substituents may be the same or different.

[0130]

25 The substituent that B optionally has is, for example, a substituent selected from the aforementioned substituent group A.

[0131]

The number of the substituents that B optionally has is, for example, 1 to 3, preferably 1 or 2, more preferably 1. When the number of the substituents is two or more, respective substituents may be the same or different.

5 [0132]


B is preferably a ring selected from
 (1) a 5- or 6-membered aromatic heterocycle, and
 (2) a benzene ring fused with an optionally substituted 5- or 6-membered ring,

10 and the ring represented by B is optionally substituted, via two adjacent carbon atoms with one atom in between, and binds to a group represented by the formula (II), and a group represented by the formula (III).

[0133]

15 In the formula


[0134]



[0135]

20 B is more preferably, a ring selected from

[0136]



[0137]

wherein

X¹, X², X³ and X⁴ are each independently a carbon atom or a nitrogen atom;

at least one of X¹, X², X³ and X⁴ is a nitrogen atom;

5 Y¹, Y² and Y³ are each independently a carbon atom, a nitrogen atom, an oxygen atom or a sulfur atom;


at least one of Y¹, Y² and Y³ is a nitrogen atom, an oxygen atom or a sulfur atom; and

10 Z¹, Z², Z³ and Z⁴ are each independently a carbon atom or a nitrogen atom, which ring is optionally substituted.

[0138]

Preferable examples of B include a ring selected from

[0139]




15 [0140]

The ring is optionally substituted by 1 or 2 (preferably 1) C₁₋₆ alkyl groups.

[0141]

B is more preferably a ring represented by the formula

[0142]



[0143]

wherein

- 5 Y¹, Y² and Y³ are each independently a carbon atom, a nitrogen atom, an oxygen atom or a sulfur atom; and at least one of Y¹, Y² and Y³ is a nitrogen atom, an oxygen atom or a sulfur atom, and the ring is optionally substituted.

[0144]

- 10 B is particularly preferably a ring selected from thiophene and thiazole, and the ring is optionally substituted.

[0145]

Another preferable embodiment of B is thiophene or thiazole each optionally substituted by 1 or 2 (preferably 1)

- 15 C₁₋₆ alkyl groups.


[0146]

A still another preferable embodiment of B is thiophene optionally substituted by one C₁₋₆ alkyl group.

[0147]

- 20 A yet another preferable embodiment of B is

[0148]



[0149]

wherein R⁷ is a hydrogen atom or a C₁₋₆ alkyl group.

- 25 [0150]

R¹, R², R³ and R⁴ are each independently a hydrogen atom, an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group.

[0151]

- 30 R¹ is preferably a hydrogen atom or an optionally

substituted C₁₋₆ alkyl group, more preferably, a hydrogen atom or a C₁₋₆ alkyl group, particularly preferably a hydrogen atom.
[0152]

A and R¹ are optionally bonded with each other to form,
5 together with the adjacent nitrogen atom optionally substituted cyclic group.
[0153]

Examples of the "cyclic group" of the "optionally substituted cyclic group", which is formed, together with the
10 adjacent nitrogen atom, by A and R¹ bonded to each other include a 4- to 10-membered heterocyclic group containing one nitrogen atom as a ring-constituting atom besides carbon atom, and optionally further containing one hetero atom selected from a nitrogen atom, a sulfur atom and an oxygen atom, and
15 the like. Examples of the "4- to 10-membered heterocyclic group" include azetidin-1-yl, pyrrolidin-1-yl, pyrrolin-1-yl, pyrazolin-1-yl, imidazolin-1-yl, thiazolidin-3-yl, oxazolidin-3-yl, isothiazolidin-2-yl, isoxazolidin-2-yl, thiazolin-3-yl, oxazolin-3-yl, isothiazolin-2-yl, isoxazolin-2-yl, piperidin-
20 1-yl, piperazin-1-yl, morpholin-4-yl, thiomorpholin-4-yl, azepan-1-yl, diazepan-1-yl, 1,4-oxazepan-4-yl, 1,4-thiazepan-4-yl, 5,7-dihydro-6H-pyrrolo[3,4-b]pyridin-6-yl and the like.

Examples of the substituent of the "optionally substituted cyclic group", which is formed, together with the
25 adjacent nitrogen atom, by A and R¹ bonded to each other, include substituents selected from the aforementioned substituent group A.

The number of the substituents in "optionally substituted cyclic group", which is formed, together with the adjacent
30 nitrogen atom, by A and R¹ bonded to each other, is, for example, 1 to 3. When the number of the substituents is two or more, the respective substituents may be the same or different.
[0154]

Preferable examples of the "optionally substituted cyclic
35 group", which is formed, together with the adjacent nitrogen

atom, by A and R¹ bonded to each other include a 4 to 10-membered heterocyclic group containing one nitrogen atom as a ring-constituting atom besides carbon atoms, and optionally further containing one hetero atom selected from a nitrogen atom, a sulfur atom and an oxygen atom (e.g., azetidin-1-yl, pyrrolidin-1-yl, piperidin-1-yl, 5,7-dihydro-6H-pyrrolo[3,4-b]pyridin-6-yl), and the heterocyclic group is optionally substituted by 1 to 3 substituents selected from a hydroxy group and a cyano group, .

10 [0155]

As regards -N(R¹)(A) moiety,
R¹ is preferably a hydrogen atom.

A is preferably

- (1) a piperidinyl group, an isoxazolyl group, a pyrazolyl group, a thiadiazolyl group, a thiazolyl group or a tetrahydropyranyl group each optionally substituted by C₁₋₆ alkyl group(s) (preferably, one C₁₋₆ alkyl group) optionally substituted by halogen atom(s) (preferably, 1 to 3 halogen atoms), or
- 20 (2) a cyclopentyl group or a cyclohexyl group each optionally substituted by halogen atom(s) (preferably, 1 or 2 halogen atoms).

[0156]

R² is preferably a hydrogen atom or an optionally substituted C₁₋₆ alkyl group, more preferably, a hydrogen atom or a C₁₋₆ alkyl group, particularly preferably a hydrogen atom.

[0157]

R³ is preferably

- (1) a hydrogen atom,
- 30 (2) an optionally substituted C₁₋₆ alkyl group,
- (3) an optionally substituted C₃₋₁₀ cycloalkyl group, or
- (4) an optionally substituted heterocyclic group.

[0158]

R³ is more preferably

- 35 (1) a hydrogen atom,

(2) a C₁₋₆ alkyl group substituted by one C₃₋₁₀ cycloalkyl group (e.g., cyclopropylmethyl group),

(3) a C₁₋₆ alkyl group substituted by one 4- to 6-membered non-aromatic heterocyclic group (preferably, a 4- to 6-membered non-aromatic heterocyclic group containing, as a ring-constituting atom besides carbon atoms, 1 or 2 hetero atoms selected from a nitrogen atom, a sulfur atom and an oxygen atom) (e.g., tetrahydropyranylmethyl group),

(4) a C₃₋₁₀ cycloalkyl group (e.g., cyclobutyl group, cyclohexyl group) optionally substituted by one amino group, or

(5) a 4- to 6-membered non-aromatic heterocyclic group (preferably, a 4- to 6-membered non-aromatic heterocyclic group containing, as a ring-constituting atom besides carbon atoms, 1 or 2 hetero atoms selected from a nitrogen atom, a sulfur atom and an oxygen atom) (e.g., tetrahydropyranyl group, piperidiny group), and the heterocyclic group is optionally substituted by one substituent selected from

(a) a C₁₋₆ alkyl group optionally substituted by halogen atom(s) (preferably, 1 to 3 halogen atoms), and

(b) a C₃₋₁₀ cycloalkyl group (e.g., cyclopropyl group).

[0159]

R³ is more preferably a hydrogen atom, a cyclopropylmethyl group, a tetrahydropyranylmethyl group, a cyclobutyl group or a tetrahydropyranyl group.

[0160]

R³ is particularly preferably a cyclopropylmethyl group, a tetrahydropyranylmethyl group, a cyclobutyl group or a tetrahydropyranyl group.

[0161]

R² and R³ are optionally bonded with each other to form, together with the adjacent nitrogen atom, an optionally substituted cyclic group.

[0162]

Examples of the "cyclic group" of the "optionally substituted cyclic group", which is formed, together with the

adjacent nitrogen atom, by R^2 and R^3 bonded to each other include a 4- to 10-membered heterocyclic group containing one nitrogen atom as a ring-constituting atom besides carbon atom, and optionally further containing one hetero atom selected
5 from a nitrogen atom, a sulfur atom and an oxygen atom, and the like. Examples of the "4- to 10-membered heterocyclic group" include azetidin-1-yl, pyrrolidin-1-yl, pyrrolin-1-yl, pyrazolin-1-yl, imidazolin-1-yl, thiazolidin-3-yl, oxazolidin-3-yl, isothiazolidin-2-yl, isoxazolidin-2-yl, thiazolin-3-yl,
10 oxazolin-3-yl, isothiazolin-2-yl, isoxazolin-2-yl, piperidin-1-yl, piperazin-1-yl, morpholin-4-yl, thiomorpholin-4-yl, azepan-1-yl, diazepan-1-yl, 1,4-oxazepan-4-yl, 1,4-thiazepan-4-yl, 5,7-dihydro-6H-pyrrolo[3,4-b]pyridin-6-yl and the like.

Examples of the substituent of the "optionally
15 substituted cyclic group", which is formed, together with the adjacent nitrogen atom, by R^2 and R^3 bonded to each other include substituents selected from the aforementioned substituent group A.

The number of the substituents in "optionally substituted
20 cyclic group", which is formed, together with the adjacent nitrogen atom, by R^2 and R^3 bonded to each other is, for example, 1 to 3. When the number of the substituents is two or more, the respective substituents may be the same or different.
[0163]

25 Preferable examples of the "optionally substituted cyclic group", which is formed, together with the adjacent nitrogen atom, by R^2 and R^3 bonded to each other include a 4- to 10-membered heterocyclic group containing one nitrogen atom as a ring-constituting atom besides carbon atom, and optionally
30 further containing one hetero atom selected from a nitrogen atom, a sulfur atom and an oxygen atom (e.g., pyrrolidin-1-yl, piperidin-1-yl, morpholin-4-yl, thiomorpholin-4-yl), and the heterocyclic group is optionally substituted by 1 to 3 oxo groups.

35 [0164]

A preferable combination of R^2 and R^3 is
 a hydrogen atom and a cyclopropylmethyl group,
 a hydrogen atom and a cyclobutyl group,
 a hydrogen atom and a tetrahydropyranyl group,
 5 a hydrogen atom and a tetrahydropyranylmethyl group, or
 a hydrogen atom and an optionally substituted piperidinyl
 group (preferably, a piperidinyl group optionally substituted
 by one C_{1-6} alkyl group optionally substituted by 1 to 3 halogen
 atoms).


10 [0165]

R^4 is preferably a hydrogen atom or an optionally
 substituted C_{1-6} alkyl group, more preferably, a hydrogen atom
 or a C_{1-6} alkyl group, particularly preferably a hydrogen atom.
 [0166]

15 In the formula (I), the relative configuration of the
 substituent represented by the formula $(R^1)(A)N-CO-B-$ and the
 substituent represented by the formula $-N(R^2)(R^3)$ on the
 cyclopropane ring is cis or trans, preferably trans.
 [0167]

20 A compound of the formula (I) having a preferable
 relative configuration is shown by the following formula (IA).
 [0168]

relative configuration





25 [0169]

wherein A, B, R^1 , R^2 , R^3 and R^4 are as defined above.

Compound (I) encompasses a compound having the absolute
 configuration represented by the following formula (IB) or
 (IC), and a mixture thereof at any ratio.

30 [0170]

absolute configuration



[0171]

wherein A, B, R¹, R², R³ and R⁴ are as defined above.

5 [0172]

Preferable examples of compound (I) include the following compounds.

[Compound A]

Compound (I) wherein

10 B is a ring selected from thiophene, thiazole, pyrazole, pyridine, naphthalene and 2,3-dihydrobenzofuran, and the ring is optionally substituted by C₁₋₆ alkyl group(s) (preferably, one C₁₋₆ alkyl group);

A is

15 (1) a heterocyclic group (preferably, (i) a 5- or 6-membered aromatic heterocyclic group or (ii) a 4- to 6-membered non-aromatic heterocyclic group, each containing, as a ring-constituting atom besides carbon atoms, 1 to 4 hetero atoms selected from a nitrogen atom, a sulfur atom and an oxygen
20 atom) (e.g., piperidinyl group, isoxazolyl group, pyrazolyl group, thiadiazolyl group, thiazolyl group, tetrahydropyranyl group, oxetanyl group, oxadiazolyl group, thienyl group, pyridyl group, oxazolyl group) optionally substituted by C₁₋₆ alkyl group(s) (preferably, 1 or 2 C₁₋₆ alkyl groups) optionally
25 substituted by halogen atom(s) (preferably, 1 to 3 halogen atoms) or

(2) a C₃₋₁₀ cycloalkyl group (e.g., cyclopropyl group,

cyclobutyl group, cyclopentyl group, cyclohexyl group)
optionally substituted by halogen atom(s) (preferably, 1 or 2
halogen atoms);

R¹ is a hydrogen atom;

5 R² is a hydrogen atom;

R³ is

(1) a hydrogen atom,

(2) a C₁₋₆ alkyl group substituted by one C₃₋₁₀ cycloalkyl group
(e.g., cyclopropylmethyl group),

10 (3) a C₁₋₆ alkyl group substituted by one 4- to 6-membered non-
aromatic heterocyclic group (preferably, a 4- to 6-membered
non-aromatic heterocyclic group containing, as a ring-
constituting atom besides carbon atoms, 1 or 2 hetero atoms
selected from a nitrogen atom, a sulfur atom and an oxygen
15 atom) (e.g., tetrahydropyranylmethyl group),

(4) a C₃₋₁₀ cycloalkyl group (e.g., cyclobutyl group, cyclohexyl
group) optionally substituted by one amino group, or

(5) a 4- to 6-membered non-aromatic heterocyclic group
(preferably, a 4- to 6-membered non-aromatic heterocyclic
20 group containing, as a ring-constituting atom besides carbon
atoms, 1 or 2 hetero atoms selected from a nitrogen atom, a
sulfur atom and an oxygen atom) (e.g., tetrahydropyranyl group,
piperidinyll group) optionally substituted by one substituent
selected from

25 (a) a C₁₋₆ alkyl group optionally substituted by halogen
atom(s) (preferably, 1 to 3 halogen atoms), and

(b) a C₃₋₁₀ cycloalkyl group (e.g., cyclopropyl group);

and

R⁴ is a hydrogen atom.

30 [0173]

[Compound B]

Compound (I) wherein

B is a ring selected from

thiophene, thiazole, pyrazole, pyridine, naphthalene and 2,3-
35 dihydrobenzofuran, wherein the ring is optionally substituted

by C₁₋₆ alkyl group(s) (preferably, one C₁₋₆ alkyl group);

A is

- (1) a pyrazolyl group, a thiadiazolyl group or a tetrahydropyranyl group, each optionally substituted by C₁₋₆ alkyl group(s) (preferably, one C₁₋₆ alkyl group), or
(2) a cyclopentyl group or a cyclohexyl group, each optionally substituted by halogen atom(s) (preferably, 1 or 2 halogen atoms);

R¹ is a hydrogen atom;

10 R² is a hydrogen atom;

R³ is a hydrogen atom, a cyclopropylmethyl group, a tetrahydropyranylmethyl group, a cyclobutyl group or a tetrahydropyranyl group; and

R⁴ is a hydrogen atom.

15 [0174]

[Compound C]

Compound (I) wherein

B is a ring selected from thiophene and thiazole, and the ring is optionally substituted by C₁₋₆ alkyl group(s) (preferably,
20 one C₁₋₆ alkyl group);

A is

- (1) a pyrazolyl group, a thiadiazolyl group or a tetrahydropyranyl group, each optionally substituted by C₁₋₆ alkyl group(s) (preferably, one C₁₋₆ alkyl group, or
25 (2) a cyclopentyl group or a cyclohexyl group, each optionally substituted by halogen atom(s) (preferably, 1 or 2 halogen atoms);

R¹ is a hydrogen atom;

R² is a hydrogen atom;


30 R³ is a cyclopropylmethyl group, a tetrahydropyranylmethyl group, a cyclobutyl group or a tetrahydropyranyl group; and
R⁴ is a hydrogen atom.

[0175]

[Compound D]

35 A compound represented by the formula (IA)

[0176]



[0177]


wherein A, B, R¹, R², R³ and R⁴ are as defined for compound A,
5 and the steric configuration shows a relative configuration,
or a salt thereof.

[0178]

[Compound E]

A compound represented by the formula (IA)

10 [0179]



[0180]


wherein A, B, R¹, R², R³ and R⁴ are as defined for compound B,
and the steric configuration shows a relative configuration,
15 or a salt thereof.

[0181]

[Compound F]

A compound represented by the formula (IA)

[0182]



20

[0183]

wherein A, B, R¹, R², R³ and R⁴ are as defined for compound C,
and the steric configuration shows a relative configuration,

or a salt thereof.

[0184]

The salt of compound (I) is preferably a pharmacologically acceptable salt. Examples of such salt
5 include salts with inorganic bases, salts with organic bases, salts with inorganic acids, salts with organic acids, and salts with basic or acidic amino acids.

Preferable examples of salts with inorganic bases include alkali metal salts such as sodium salts, potassium salts and
10 the like; alkali earth metal salts such as calcium salts, magnesium salts and the like; aluminum salts; and ammonium salts.

Preferable examples of salts with organic bases include salts with trimethylamine, triethylamine, pyridine, picoline,
15 ethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, N,N-dibenzylethylenediamine and the like.

Preferable examples of salts with inorganic acids include salts with hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid and the like.

20 Preferable examples of salts with organic acids include salts with formic acid, acetic acid, trifluoroacetic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, *p*-toluenesulfonic acid and the like.

25 Preferable examples of salts with basic amino acids include salts with arginine, lysine, ornithine and the like.

Preferable examples of salts with acidic amino acids include salts with aspartic acid, glutamic acid and the like.

The salt of compound (I) is preferably a salt with an
30 inorganic acid (preferably, hydrochloric acid) or an organic acid (preferably, trifluoroacetic acid).

[0185]

Compound (I) may also be used as a prodrug. A prodrug of compound (I) means a compound which is converted to compound
35 (I) due to a reaction due to an enzyme, gastric acid, etc.

under the physiological condition in the living body, that is, a compound which is converted to compound (I) with oxidation, reduction, hydrolysis, and the like according to an enzyme; a compound which is converted to compound (I) by hydrolysis etc. due to gastric acid, and the like. A prodrug for compound (I) may be a compound obtained by subjecting an amino group in compound (I) to an acylation, alkylation or phosphorylation (e.g., a compound obtained by subjecting an amino group in compound (I) to an eicosanoylation, alanylation, pentylaminocarbonylation, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methoxycarbonylation, tetrahydrofuranylation, pyrrolidylmethylation, pivaloyloxymethylation and tert-butylation, etc.); a compound obtained by subjecting a hydroxy group in compound (I) to an acylation, alkylation, phosphorylation or boration (e.g., a compound obtained by subjecting a hydroxy group in compound (I) to an acetylation, palmitoylation, propanoylation, pivaloylation, succinylation, fumarylation, alanylation and dimethylaminomethylcarbonylation); a compound obtained by subjecting a carboxyl group in compound (I) to an esterification or amidation (e.g., a compound obtained by subjecting a carboxyl group in compound (I) to an ethyl esterification, phenyl esterification, carboxymethyl esterification, dimethylaminomethyl esterification, pivaloyloxymethyl esterification, ethoxycarbonyloxyethyl esterification, phthalidyl esterification, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl esterification, cyclohexyloxycarbonylethyl esterification and methylamidation) and the like. Any of these compounds can be produced from compound (I) by a method known per se.

[0186]

The prodrug of compound (I) may be a compound that converts to compound (I) under physiological conditions as described in Development of Pharmaceutical Products, vol. 7, Molecular Design, 163-198, Hirokawa Shoten (1990).

Compound (I) may be labeled with an isotope (e.g., ^2H , ^3H , ^{14}C , ^{35}S , ^{125}I , ^{11}C , ^{18}F) and the like.

Compound (I) labeled with or substituted by an isotope can be used, for example, as a tracer used for Positron
5 Emission Tomography (PET) (PET tracer), and is useful in the field of medical diagnosis and the like.

Compound (I) may be an anhydrate or a hydrate. Compound (I) may be a solvate or a non-solvate. Furthermore, compound (I) may be a deuterated compound.

10 Compound (I) may be a crystal, and both a single crystal and crystal mixtures are encompassed in compound (I). Crystals can be produced by crystallization according to crystallization methods known per se.

In addition, compound (I) may be a pharmaceutically
15 acceptable cocrystal or cocrystal salt. Here, the cocrystal or cocrystal salt means a crystalline substance consisting of two or more particular substances which are solids at room temperature, each having different physical properties (e.g., structure, melting point, heat of melting, hygroscopicity, and
20 stability). The cocrystal and cocrystal salt can be produced by cocrystallization method known per se.

When compound (I) includes isomers such as optical isomers, stereoisomers, regioisomers, rotational isomers, geometrical isomers, and the like, one of the isomers and
25 mixture are also encompassed in compound (I).

[0187]

Compound (I) or a prodrug thereof (hereinafter sometimes to be simply abbreviated as the compound of the present invention) has low toxicity (e.g., acute toxicity, chronic
30 toxicity, genetic toxicity, reproductive toxicity, cardiotoxicity, carcinogenicity), and can be used as it is or in the form of a pharmaceutical composition (in the present specification, sometimes to be abbreviated as "medicament of the present invention") after mixing with a pharmacologically
35 acceptable carrier etc. to mammals (e.g., human, mouse, rat,

rabbit, dog, cat, bovine, horse, swine, monkey) as an agent for the prophylaxis or treatment of various diseases mentioned below.

[0188]

5 As a pharmaceutical acceptable carrier here, common organic or inorganic carrier substances are used as formulation raw materials. Carriers are added as vehicles, lubricants, binders and disintegrants in the solid formulations; and as solvents, solubilizing agents, suspending
10 agents, isotonization agents, buffering agents, soothing agents etc. in the liquid formulations. If desired, formulation additives such as preservatives, antioxidants, colorants, sweeteners, etc. can be used.

[0189]

15 Preferable examples of the vehicles are as follows: lactose, sucrose, D-mannitol, D-sorbitol, starch, α -starch, dextrin, crystalline cellulose, low-substituted hydroxypropyl cellulose, sodium carboxymethylcellulose, gum Arabic, pullulan, light anhydrous silicic acid, synthetic aluminum silicate, and
20 magnesium metasilicic aluminate.

[0190]

Preferable examples of the lubricant include magnesium stearate, calcium stearate, talc, colloidal silica, and the like.

25 [0191]

Preferable examples of the binders are as follows: α -starch, sucrose, gelatin, gum Arabic, methylcellulose, carboxymethylcellulose, sodium carboxymethylcellulose, crystalline cellulose, sucrose, D-mannitol, trehalose, dextrin,
30 pullulan, hydroxypropylcellulose, hydroxypropyl methylcellulose, and polyvinylpyrrolidone.

[0192]

Preferable examples of the disintegrants are as follows: lactose, sucrose, starch, carboxymethylcellulose, calcium
35 carboxymethylcellulose, croscarmellose sodium, sodium

carboxymethyl starch, light anhydrous silicic acid, and low-substituted hydroxypropylcellulose.

[0193]

Preferable examples of the solvents are as follows: water
5 for injection, physiological saline, Linger solution, alcohol, propylene glycol, polyethylene glycol, sesame oil, corn oil, olive oil, and cottonseed oil.

[0194]

Preferable examples of the solubilizing agents are as
10 follows: polyethylene glycol, propylene glycol, D-mannitol, trehalose, benzyl benzoate, ethanol, tris-aminomethane, cholesterol, triethanolamine, sodium carbonate, sodium citrate, sodium salicylate, and sodium acetate.

[0195]

15 Preferable examples of the suspending agent include surfactants such as stearyl triethanolamine, sodium lauryl sulfate, laurylaminopropionic acid, lecithin, benzalkonium chloride, benzetonium chloride, glycerin monostearate and the like; hydrophilic polymers such as polyvinyl alcohol,
20 polyvinylpyrrolidone, carboxymethylcellulose sodium, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose and the like; polysorbates, and polyoxyethylene hydrogenated castor oil.

[0196]

25 Preferable examples of the isotonization agents include sodium chloride, glycerin, D-mannitol, D-sorbitol, and glucose.

[0197]

Preferable examples of the buffering agent include buffer
30 solutions such as phosphates, acetates, carbonates, and citrates.

[0198]

Preferable examples of the soothing agent include benzyl alcohol.

[0199]

35 Preferable examples of the preservative include p-

hydroxybenzoates, chlorobutanol, benzyl alcohol, phenethyl alcohol, dehydroacetic acid, and sorbic acid.

[0200]

Preferable examples of antioxidants include sulfites and
5 ascorbates.

[0201]

Preferable examples of the colorants include water soluble edible tar dyes (e.g., edible dyes such as Food Red No. 2 and No. 3, Food Yellow No. 4 and No. 5, Food Blue No. 1 and
10 2, etc.); water insoluble lake dyes (e.g., aluminum salts of the aforementioned water soluble edible tar dyes), and natural dyes (e.g., β -carotene, chlorophyll, ferric oxide red).

[0202]

Preferable examples of the sweeteners include sodium
15 saccharin, dipotassium glycyrrhizinate, aspartame and stevia.

[0203]

Examples of the dosage form of the medicament of the present invention include oral preparations such as tablet (including sublingual tablet, orally disintegrating tablet),
20 capsule (including soft capsule, microcapsule), granule, powder, troche, syrup, emulsion, suspension and the like; and parenteral agents such as injection (e.g., subcutaneous injection, intravenous injection, intramuscular injection, intraperitoneal injection, drip infusion), external
25 preparation (e.g., dermal preparation, ointment), suppository (e.g., rectal suppository, vaginal suppository), pellet, nasal preparation, pulmonary preparation (inhalant), eye drop and the like, which can be respectively safely administered orally or parenterally.

30 [0204]

These preparations may be a release control preparation (e.g., sustained-release microcapsule) such as an immediate-release preparation, a sustained-release preparation and the like.

35 [0205]

The medicament of the present invention can be manufactured by the common methods in the field of formulation technology, for example, methods listed in the Japanese Pharmacopoeia, and the like.

5 [0206]

The content of the compound of the present invention in the medicament of the present invention varies based on the dosage forms, dosages of the compound of the present invention, and the like. For example, it is approximately about 0.1 to
10 100 wt%.

[0207]

The compound of the present invention has a superior LSD1 inhibitory action and can be used as a prophylactic or therapeutic agent for various diseases in mammals (e.g., human,
15 mouse, rat, rabbit, dog, cat, bovine, horse, swine, monkey). Moreover, since the compound of the present invention shows low monoamine oxidase A (MAO-A) and monoamine oxidase B (MAO-B) inhibitory activity and high LSD1 selectivity, it causes fewer side effects.

20 [0208]

In addition, the compound of the present invention is expected to show, after transfer into the brain, suppression of a decrease in histone H3K4 methylation and suppression of a decrease in Gad1 mRNA expression, which are derived from the
25 inhibition of LSD1. As a result, it is also useful as a medicament based on superior actions such as neuronal function, enhancement of neural plasticity, promotion of neurogenesis, and promotion of BDNF production.

In addition, the compound of the present invention can,
30 after transfer into the brain, suppress a decrease in histone H3K9 methylation derived from LSD1 inhibition. As a result, it is effective for the treatment of some diseases associated with a decrease in the methylation of H3K9.

[0209]

35 The compound of the present invention can be used as a

prophylactic or therapeutic agent for cancer. Examples of the cancer include breast cancer, prostate cancer, pancreatic cancer, gastric cancer, lung cancer, colorectal cancer, rectal cancer, esophagus cancer, duodenal cancer, tongue cancer, 5 pharyngeal cancer, brain tumor, schwannoma, non-small cell lung cancer, small cell lung cancer, liver cancer, kidney cancer, bile duct cancer, uterine body cancer, cervical cancer, ovarian cancer, urinary bladder cancer, skin cancer, hemangioma, malignant lymphoma, malignant melanoma, thyroid 10 cancer, bone tumor, angiofibroma, retinoblastoma, penile cancer, childhood solid cancer, Kaposi's sarcoma, AIDS-related Kaposi's sarcoma, tumor of maxillary sinus, fibrous histiocytoma, leiomyosarcoma, rhabdomyosarcoma, and leukemia. Among these, the compound can be preferably used for prostate 15 cancer, leukemia, and malignant lymphoma.

[0210]

It is known that the level of H3K4me2, which is a substrate of LSD1, and memory improvement are correlated (Nature 2007, Vol. 447, page 175), and the compound of the 20 present invention having a superior LSD1 inhibitory action can also be used as a prophylactic or therapeutic agent for neurodegenerative diseases.

[0211]

The compound of the present invention can be used as a 25 therapeutic agent for anemia. Examples of anemia include sickle cell anaemia and beta thalassemia.

[0212]

The compound of the present invention can be used as a prophylactic or therapeutic agent for central nervous system 30 diseases. It is useful as a prophylactic or therapeutic agent for diseases such as

(1) psychiatric diseases [e.g., depression, major depression, bipolar depression, dysthymic disorder, emotional disorder (seasonal affective disorder and the like), recurrent 35 depression, postpartum depression, stress disorder, depression

symptom, mania, anxiety, generalized anxiety disorder, anxiety syndrome, mood disorder, panic disorder, phobia, social phobia, social anxiety disorder, obsessive disorder, post-traumatic stress syndrome, post-traumatic stress disorder, Tourette syndrome, adjustment disorder, bipolar disorder, neurosis, schizophrenia (e.g., positive symptom, negative symptom, cognitive impairment), chronic fatigue syndrome, anxiety neurosis, compulsive neurosis, panic disorder, anxiety symptom, anxious mental state, emotional abnormality, cyclothymia, nervous erethism, faint, addiction, low sex drive, attention deficit hyperactivity disorder (ADHD), psychotic major depression, refractory major depression, treatment-resistant depression],

(2) neurodegenerative diseases [e.g., Alzheimer's disease, Alzheimer-type senile dementia, Parkinson's disease, Huntington's disease (Huntington chorea), multi-infarct dementia, frontotemporal dementia, frontotemporal dementia with Parkinsonism, progressive supranuclear palsy, Pick's syndrome, Niemann-Pick syndrome, corticobasal degeneration, vascular dementia, postencephalitic parkinsonism, dementia with Lewy body, HIV dementia, amyotrophic lateral sclerosis (ALS), motor neurogenesis disease (MND), Creutzfeldt-Jakob disease or prion disease, cerebral palsy, progressive supranuclear palsy, multiple sclerosis, spinocerebellar degeneration (e.g., dentatorubral pallidoluysian atrophy)],

(3) developmental disorders, particularly, diseases with intellectual disability (e.g., autistic spectrum disorders, Rett syndrome, Down's syndrome, Kabuki syndrome, fragile X syndrome, Kleeftstra syndrome, neurofibromatosis type 1, Noonan syndrome, tuberous sclerosis),

(4) age-related cognition and memory disorders [e.g., age-related memory disorders, senile dementia],

(5) sleep disorders [e.g., intrinsic sleep disorders (e.g., psychophysiological insomnia and the like), extrinsic sleep disorder, circadian rhythm disorders (e.g., time zone change

syndrome (jet lag), shift work sleep disorder, irregular sleep-wake pattern, delayed sleep phase syndrome, advanced sleep phase syndrome, non-24-hour sleep-wake disorder and the like), parasomnia, sleep disorders associated with internal
5 medical or psychiatric disorder (e.g., chronic obstructive pulmonary disease, Alzheimer's disease, Parkinson's disease, cerebrovascular dementia, schizophrenia, depression, anxiety neurosis), stress insomnia, insomnia, insomniac neurosis, sleep apnea syndrome],
10 (6) respiratory depression caused by anesthetics, traumatic disease, or neurodegenerative disease and the like,
(7) epilepsy (e.g., Dravet syndrome), traumatic brain injury, cerebral apoplexy, neurotic anorexia, eating disorder, anorexia nervosa, hyperorexia, other eating disorder, alcohol
15 dependence, alcohol abuse, alcoholic amnesia, alcohol paranoia, alcohol preference, alcohol withdrawal, alcoholic insanity, alcohol intoxication, alcoholic jealousy, alcoholic mania, alcohol-dependent psychiatric disorder, alcoholic insanity, drug dependence, drug intoxication, pharmacophilia,
20 pharmacophobia, pharmacomania, drug withdrawal, migraine, stress headache, catatonic headache, diabetic neuropathy, obesity, diabetes, muscular spasm, Meniere's disease, autonomic ataxia, alopecia, glaucoma, hearing loss, hypertension, cardiac disease, tachycardia, congestive cardiac
25 failure, hyperventilation, bronchial asthma, apnea, sudden infant death syndrome, inflammatory disease, allergic disease, impotence, climacteric disorder, infertility, cancer, immunodeficiency syndrome caused by HIV infection, immunodeficiency syndrome caused by stress, cerebrospinal
30 meningitis, acromegaly, incontinence, metabolic syndrome, osteoporosis, peptic ulcer, irritable bowel syndrome, inflammatory bowel disease, ulcerative colitis, Crohn's disease, stress gastrointestinal disorder, stress vomiting, stress ulcer, diarrhea, constipation, postoperative ileus, and
35 the like.

[0213]

The compound of the present invention is particularly useful as an agent for the prophylaxis or treatment of diseases such as schizophrenia, developmental disorders, particularly diseases having intellectual disability (e.g., autistic spectrum disorders, Rett syndrome, Down's syndrome, Kabuki syndrome, fragile X syndrome, Kleefstra syndrome, neurofibromatosis type 1, Noonan syndrome, tuberous sclerosis), neurodegenerative diseases (e.g., Alzheimer's disease, Parkinson's disease, spinocerebellar degeneration (e.g., dentatorubral pallidoluysian atrophy) and Huntington's disease (Huntington chorea)), epilepsy (e.g., Dravet syndrome) or drug dependence, and the like.

[0214]

Since the compound of the present invention has a superior LSD1 inhibitory activity, it is expected to show a superior treatment effects for the above-mentioned diseases.

[0215]

The dosage of the compound of the present invention varies depending on the administration subjects, administration routes, target diseases, symptoms, and the like. For example, for oral administration to adult patients with cancer, generally a single dose is about 0.01 to 100 mg/kg body weight, preferably 0.1 to 50 mg/kg body weight, further preferably 0.5 to 20 mg/kg body weight, and this dosage is preferably administered 1 to 3 times daily.

[0216]

The compound of the present invention can be used in combination with a medicament such as chemotherapeutic agent, immunotherapeutic agent, medicament inhibiting actions of cell growth factor and receptor thereof (hereinafter to be abbreviated as a concomitant drug).

By combining the compound of the present invention and a concomitant drug, a superior effect such as (1) the dose can be reduced as compared to single

administration of the compound of the present invention or a concomitant drug,

(2) the drug to be combined with the compound of the present invention can be selected according to the condition of
5 patients (mild case, severe case and the like),

(3) the period of treatment can be set longer by selecting a concomitant drug having different action and mechanism from the compound of the present invention,

(4) a sustained treatment effect can be designed by selecting
10 a concomitant drug having different action and mechanism from the compound of the present invention,

(5) a synergistic effect can be afforded by a combined use of the compound of the present invention and a concomitant drug, and the like, can be achieved.

15 [0217]

Hereinafter the compound of the present invention and a concomitant drug used in combination are referred to as the "combination agent of the present invention".

When using the combination agent of the present invention,
20 the administration time of the compound of the present invention and the concomitant drug is not restricted, and the compound of the present invention or a pharmaceutical composition thereof and the concomitant drug or a pharmaceutical composition thereof can be administered to an
25 administration subject simultaneously, or may be administered at different times. The dosage of the concomitant drug may be determined according to the dose clinically used, and can be appropriately selected depending on an administration subject, administration route, disease, combination and the like.

30 The administration mode of the combination agent of the present invention is not particularly restricted, and it is sufficient that the compound of the present invention and the concomitant drug are combined in administration. Examples of such administration mode include the following methods:

35 (1) administration of a single preparation obtained by

simultaneously processing the compound of the present invention and the concomitant drug, (2) simultaneous administration of two kinds of preparations of the compound of the present invention and the concomitant drug, which have
5 been separately produced, by the same administration route, (3) administration of two kinds of preparations of the compound of the present invention and the concomitant drug, which have been separately produced, by the same administration route in a staggered manner, (4) simultaneous
10 administration of two kinds of preparations of the compound of the present invention and the concomitant drug, which have been separately produced, by different administration routes, (5) administration of two kinds of preparations of the compound of the present invention and the concomitant drug,
15 which have been separately produced, by different administration routes in a staggered manner (e.g., administration in the order of the compound of the present invention and the concomitant drug, or in the reverse order) and the like.

20 The dose of the concomitant drug can be appropriately determined based on the dose employed in clinical situations. The mixing ratio of the compound of the present invention and a concomitant drug can be appropriately determined depending on the administration subject, administration route, target
25 disease, symptom, combination and the like.

For example, the content of the compound of the present invention in the combination agent of the present invention differs depending on the form of a preparation, and usually from about 0.01 to about 100 wt%, preferably from about 0.1 to
30 about 50 wt%, further preferably from about 0.5 to about 20 wt%, based on the preparation.

The content of the concomitant drug in the combination agent of the present invention differs depending on the form of a preparation, and usually from about 0.01 to about 100 wt%,
35 preferably from about 0.1 to about 50 wt%, further preferably

from about 0.5 to about 20 wt%, based on the preparation.

The content of additives such as a carrier and the like in the combination agent of the present invention differs depending on the form of a preparation, and usually from about 1 to about 99.99 wt%, preferably from about 10 to about 90 wt%, based on the preparation.

When the compound of the present invention and a concomitant drug are separately formulated into preparations, the contents thereof are similar to the above.

10 [0218]

Examples of the chemotherapeutic agent include alkylating agents (e.g., nitrogen mustard, nitrogen mustard-N-oxide hydrochloride, chlorambutyl, cyclophosphamide, ifosfamide, thiotepa, carboquone, improsulfan tosylate, busulfan, 15 nimustine hydrochloride, mitobronitol, melphalan, dacarbazine, ranimustine, estramustine phosphate sodium, triethylenemelamine, carmustine, lomustine, streptozocin, pipobroman, etoglucid, carboplatin, cisplatin, miboplatin, nedaplatin, oxaliplatin, altretamine, ambamustine, 20 dibrospidium hydrochloride, fotemustine, prednimustine, pumitepa, ribomustin, temozolomide, treosulphan, trophosphamide, zinostatin stimalamer, adozelesin, systemustine, bizelesin), metabolic antagonists (e.g., mercaptopurine, 6-mercaptopurine riboside, thioinosine, 25 methotrexate, pemetrexed, enocitabine, cytarabine, cytarabine ocfosphate, ancitabine hydrochloride, 5-FU drug (e.g., fluorouracil, tegafur, UFT, doxifluridine, carmofur, gallocitabine, emitefur, capecitabine), aminopterin, nelzarabine, leucovorin calcium, tabloid, butocine, folinate 30 calcium, levofolinate calcium, cladribine, emitefur, fludarabine, gemcitabine, hydroxycarbamide, pentostatin, piritrexim, idoxuridine, mitoguazone, tiazofurin, ambamustine, bendamustine), antitumor antibiotics (e.g., actinomycin D, actinomycin C, mitomycin C, chromomycin A3, bleomycin 35 hydrochloride, bleomycin sulfate, peplomycin sulfate,

daunorubicin hydrochloride, doxorubicin hydrochloride, aclarubicin hydrochloride, pirarubicin hydrochloride, epirubicin hydrochloride, neocarzinostatin, mithramycin, sarkomycin, carzinophilin, mitotane, zorubicin hydrochloride, mitoxantrone hydrochloride, idarubicin hydrochloride), and plant-derived antitumor agents (e.g., etoposide, etoposide phosphate, vinblastine sulfate, vincristine sulfate, vindesine sulfate, teniposide, paclitaxel, docetaxel, vinorelbine).

[0219]

10 Examples of the immunotherapeutic agent include picibanil, krestin, schizophyllan, lentinan, ubenimex, interferon, interleukin, macrophage colony stimulating factor, granulocyte colony stimulating factor, erythropoietin, lymphotoxin, BCG vaccine, corynebacterium parvum, levamisole, polysaccharide K, 15 procodazole and anti-CTLA4 antibody.

[0220]

 Examples of the "medicament inhibiting actions of cell growth factor and receptor thereof" include anti-VEGF antibody (e.g., Bevacizumab), anti-HER2 antibody (e.g., Trastuzumab, 20 Pertuzumab), anti-EGFR antibody (e.g., Cetuximab, Panitumumab, Matuzumab, Nimotuzumab), anti-VEGFR antibody, anti-HGF antibody, Imatinib mesylate, Erlotinib, Gefitinib, Sorafenib, Sunitinib, Dasatinib, Lapatinib, Vatalanib, 4-(4-fluoro-2-methyl-1H-indol-5-yloxy)-6-methoxy-7-[3-(1-pyrrolidinyl)propoxy]quinazoline (AZD-2171), Lestaurtinib, 25 Pazopanib, Canertinib, Tandutinib, 3-(4-bromo-2,6-difluorobenzyloxy)-5-[3-[4-(1-pyrrolidinyl)butyl]ureido]isothiazole-4-carboxamide (CP-547632), Axitinib, N-(3,3-dimethyl-2,3-dihydro-1H-indol-6-yl)-2-(pyridin-4-ylmethylamino)pyridine-3-carboxamide (AMG-706), 30 Nilotinib, 6-[4-(4-ethylpiperazin-1-ylmethyl)phenyl]-N-[1(R)-phenylethyl]-7H-pyrrolo[2,3-d]pyrimidin-4-amine (AEE-788), Vandetanib, Temsirolimus, Everolimus, Enzastaurin, N-[4-[4-(4-methylpiperazin-1-yl)-6-(3-methyl-1H-pyrazol-5-ylamino)pyrimidin-2-ylsulfanyl]phenyl]cyclopropanecarboxamide 35

(VX-680), 2-[N-[3-[4-[5-[N-(3-fluorophenyl) carbamoylmethyl]-1H-pyrazol-3-ylamino]quinazolin-7-yloxy]propyl]-N-ethylamino]ethyl phosphate (AZD-1152), 4-[9-chloro-7-(2,6-difluorophenyl)-5H-primido[5,4-d][2]benzazepin-2-ylamino]benzoic acid (MLN-8054), N-[2-methoxy-5-[(E)-2-(2,4,6-trimethoxyphenyl)vinylsulfonylmethyl]phenyl]glycine sodium salt (ON-1910Na), 4-[8-cyclopentyl-7(R)-ethyl-5-methyl-6-oxo-5,6,7,8-tetrahydropteridin-2-ylamino]-3-methoxy-N-(1-methylpiperidin-4-yl)benzamide (BI-2536), 5-(4-bromo-2-chlorophenylamino)-4-fluoro-1-methyl-1H-benzimidazole-6-carbohydroxamic acid 2-hydroxyethyl ester (AZD-6244), N-[2(R),3-dihydroxypropoxy]-3,4-difluoro-2-(2-fluoro-4-iodophenylamino)benzamide (PD-0325901) and everolimus (RAD001). [0221]

15 Examples of the concomitant drug for the central nervous system diseases include the following.

benzodiazepine (chlordiazepoxide, diazepam, potassium clorazepate, lorazepam, clonazepam, alprazolam etc.), L-type calcium channel inhibitor (pregabalin etc.), tricyclic or

20 tetracyclic antidepressant (imipramine hydrochloride, amitriptyline hydrochloride, desipramine hydrochloride, clomipramine hydrochloride etc.), selective serotonin reuptake inhibitor (fluvoxamine maleate, fluoxetine hydrochloride, citalopram hydrobromide, sertraline hydrochloride, paroxetine

25 hydrochloride, escitalopram oxalate etc.), serotonin-noradrenaline reuptake inhibitor (venlafaxine hydrochloride, duloxetine hydrochloride, desvenlafaxine hydrochloride etc.), noradrenaline reuptake inhibitor (reboxetine mesylate etc.), mirtazapine, trazodone hydrochloride, nefazodone hydrochloride,

30 bupropion hydrochloride, setiptiline maleate, 5-HT_{1A} agonist, (buspirone hydrochloride, tandospirone citrate, osetozotan hydrochloride etc.), 5-HT₃ antagonist (cyamemazine etc.), non-cardioselective β blocker (propranolol hydrochloride, oxprenolol hydrochloride etc.), histamine H₁ antagonist

35 (hydroxyzine hydrochloride etc.), therapeutic drug for

schizophrenia (chlorpromazine, haloperidol, sulpiride, clozapine, trifluoperazine hydrochloride, fluphenazine hydrochloride, olanzapine, quetiapine fumarate, risperidone, aripiprazole etc.), CRF antagonist, other antianxiety drug
5 (meprobamate etc.), tachykinin antagonist (MK-869, saredutant etc.), medicament that acts on metabotropic glutamate receptor, CCK antagonist, β 3 adrenaline antagonist (amibegron hydrochloride etc.), GAT-1 inhibitor (tiagabine hydrochloride etc.), N-type calcium channel inhibitor, carbonic anhydrase II
10 inhibitor, NMDA glycine site agonist, NMDA antagonist (memantine etc.), peripheral benzodiazepine receptor agonist, vasopressin antagonist, vasopressin V1b antagonist, vasopressin V1a antagonist, phosphodiesterase inhibitor, opioid antagonist, opioid agonist, uridine, nicotinic acid
15 receptor agonist, thyroid hormone (T3, T4), TSH, TRH (Hirtonin, Ceredist), MAO inhibitor (phenelzine sulfate, tranlycypromine sulfate, moclobemide etc.), 5-HT_{2A} antagonist, 5-HT_{2A} inverse agonist, COMT inhibitor (entacapone etc.), therapeutic drug for bipolar disorder (lithium carbonate, sodium valproate,
20 lamotrigine, riluzole, felbamate etc.), cannabinoid CB1 antagonist (rimonabant etc.), FAAH inhibitor, sodium channel inhibitor, anti-ADHD drug (methylphenidate hydrochloride, methamphetamine hydrochloride etc.), therapeutic drug for alcohol dependence, therapeutic drug for autism, therapeutic
25 drug for chronic fatigue syndrome, therapeutic drug for spasm, therapeutic drug for fibromyalgia syndrome, therapeutic drug for headache, therapeutic drug for insomnia (etizolam, zopiclone, triazolam, zolpidem, ramelteon, indiplon etc.), therapeutic drug for quitting smoking, therapeutic drug for
30 myasthenia gravis, therapeutic drug for cerebral infarction, therapeutic drug for mania, therapeutic drug for hypersomnia, therapeutic drug for pain, therapeutic drug for dysthymia, therapeutic drug for autonomic ataxia, therapeutic drug for male and female sexual dysfunction, therapeutic drug for
35 migraine, therapeutic drug for pathological gambler,

therapeutic drug for restless legs syndrome, therapeutic drug for substance dependence (varenicline etc.), therapeutic drug for alcohol-related syndrome, therapeutic drug for irritable bowel syndrome, therapeutic drug for Alzheimer's disease (donepezil, galanthamine, memantine etc.), therapeutic drug for Parkinson's disease, therapeutic drug for Huntington's disease (Huntington chorea), therapeutic drug for ALS (riluzole etc., neurotrophic factor etc.), antiepileptic drug (phenobarbital, phenytoin, valproic acid, clonazepam, topiramate, gabapentin, phenobarbital ethosuximide etc.), therapeutic drug for lipid abnormality such as cholesterol-lowering drug (statin series (pravastatin sodium, atorvastatin, simvastatin, rosuvastatin etc.), fibrate (clofibrate etc.), squalene synthetase inhibitor), therapeutic drug for abnormal behavior or suppressant of dromomania due to dementia (sedatives, antianxiety drug etc.), apoptosis inhibitor, antiobesity drug, therapeutic drug for diabetes, therapeutic drug for hypertension, therapeutic drug for hypotension, therapeutic drug for rheumatism (DMARD), anti-cancer agent, therapeutic drug for parathyroid (PTH), calcium receptor antagonist, sex hormone or a derivative thereof (progesterone, estradiol, estradiol benzoate etc.), neuronal differentiation promoter, nerve regeneration promoter, non-steroidal anti-inflammatory drug (meloxicam, tenoxicam, indomethacin, ibuprofen, celecoxib, rofecoxib, aspirin, indomethacin etc.), steroid (dexamethasone, cortisone acetate etc.), anti-cytokine drug (TNF inhibitor, MAP kinase inhibitor etc.), antibody medicament, nucleic acid or nucleic acid derivative, aptamer drug, and the like.

[0222]

The above-mentioned concomitant drug may be used in a combination of two or more kinds at an appropriate ratio.

When the compound of the present invention is used in combination with a concomitant drug, the respective dosages can be reduced within a safe range in consideration of the

opposite effects of the respective drugs. As a result, the opposite effect caused by these agents can be prevented safely.

The compound of the present invention can also be used in combination with a non-medication therapy. Specific examples of the non-medication therapy include (1) operation; (2) hypertensive chemical therapy using angiotensin II and the like; (3) gene therapy; (4) hyperthermic therapy; (5) cryotherapy; (6) laser ablation method; (7) radiation therapy; and (8) immunotherapy.

10 [0223]

[Production method]

The production method of the compound of the present invention is explained below. Those of ordinary skill in the art can produce compound (I) according to the methods shown in each step of the following production methods and Examples, or a method analogous thereto.

[0224]

The starting materials and reagents used in each step of the following production methods, as well as the obtained compounds may each form a salt. Examples of such salt include those similar to the salts of the aforementioned compounds of the present invention and the like.

[0225]

When the compound obtained in each step is a free compound, it can be converted to a desired salt by a method known per se. Conversely, when the compound obtained in each step is a salt, it can be converted to a free form or a desired other kind of salt by a method known per se.

[0226]

While the compounds obtained in each step can be directly used for the next reaction in the form of a reaction mixture or as a crude product, the compound obtained in each step can be isolated and/or purified according to a conventional method from the reaction mixture by separation means such as concentration, crystallization, recrystallization,

distillation, solvent extraction, fractionation, chromatography and the like.

[0227]

When the compound of a starting material or a reagent for
5 each step is commercially available, the commercially available product can be directly used.

[0228]

In the reaction of each step, while the reaction time varies depending on the reagents and solvents to be used,
10 unless particularly indicated, it is generally 1 min to 72 hr, preferably 10 min to 8 hr.

[0229]

In the reaction of each step, while the reaction temperature varies depending on the reagents and solvents to
15 be used, unless particularly indicated, it is generally -78°C to 300°C, preferably -78°C to 150°C.

[0230]

In the reaction of each step, while the pressure varies depending on the reagents and solvents to be used, unless
20 particularly indicated, it is generally 1 atm to 20 atm, preferably 1 atm to 3 atm.

[0231]

In the reaction of each step, microwave synthesizers such as Initiator manufactured by Biotage and the like are
25 sometimes used. While the reaction temperature varies depending on the reagents and solvents to be used, unless particularly indicated, it is generally room temperature to 300°C, preferably 50°C to 250°C. While the reaction time varies depending on the reagents and solvents to be used, unless
30 particularly indicated, it is generally 1 min to 48 hr, preferably 1 min to 8 hr.

[0232]

In the reaction of each step, unless particularly indicated, the reagents are used in 0.5 equivalent to 20
35 equivalents, preferably 0.8 equivalent to 5 equivalents,

relative to the substrate. When a reagent is used as a catalyst, the reagent is used in 0.001 equivalent to 1 equivalent, preferably 0.01 equivalent to 0.2 equivalent, relative to the substrate. When the reagent is also used as a reaction solvent, the reagent is used in a solvent amount.

[0233]

In the reaction of each step, unless particularly indicated, the reaction is performed without solvent, or by dissolving or suspending a starting material in a suitable solvent. Specific examples of the solvent include the solvents described in Examples, and the following.

alcohols: methanol, ethanol, tert-butyl alcohol, 2-methoxyethanol and the like;

ethers: diethyl ether, diphenyl ether, tetrahydrofuran, 1,2-dimethoxyethane and the like;

aromatic hydrocarbons: chlorobenzene, toluene, xylene and the like;

saturated hydrocarbons: cyclohexane, hexane and the like ;

amides: N,N-dimethylformamide, N-methylpyrrolidone and the like;

hydrocarbon halides: dichloromethane, carbon tetrachloride and the like;

nitriles: acetonitrile and the like;

sulfoxides: dimethyl sulfoxide and the like;

aromatic organic bases: pyridine and the like;

acid anhydrides: acetic anhydride and the like;

organic acids: formic acid, acetic acid, trifluoroacetic acid and the like;

inorganic acids: hydrochloric acid, sulfuric acid and the like;

esters: ethyl acetate and the like;

ketones: acetone, methyl ethyl ketone and the like;

water.

Two or more kinds of the above-mentioned solvents may be mixed and used at appropriate ratios.

[0234]

When a base is used in the reaction of each step, for example, the bases shown below or the bases described in the Examples are used.

- 5 inorganic bases: sodium hydroxide, magnesium hydroxide, lithium hydroxide and the like;
basic salts: sodium carbonate, calcium carbonate, sodium hydrogen carbonate and the like;
organic bases: triethylamine, diethylamine, pyridine, 4-
10 dimethylaminopyridine, N,N-dimethylaniline, 1,4-diazabicyclo[2.2.2]octane, 1,8-diazabicyclo[5.4.0]-7-undecene, imidazole, piperidine and the like;
metal alkoxides: sodium ethoxide, potassium tert-butoxide and the like;
15 alkali metal hydrides: sodium hydride and the like;
metal amides: sodium amide, lithium diisopropylamide, lithium hexamethyldisilazide and the like;
organic lithiums: n-butyllithium and the like.

[0235]

- 20 When an acid or an acidic catalyst is used in the reaction of each step, for example, the acids or acidic catalysts shown below or those described in the Examples are used.

- inorganic acids: hydrochloric acid, sulfuric acid, nitric acid,
25 hydrobromic acid, phosphoric acid and the like;
organic acids: acetic acid, trifluoroacetic acid, citric acid, p-toluenesulfonic acid, 10-camphorsulfonic acid and the like;
Lewis acids: boron trifluoride diethyl ether complex, zinc iodide, anhydrous aluminum chloride, anhydrous zinc chloride,
30 anhydrous iron chloride and the like.

[0236]

- Unless otherwise specified, the reaction of each step is performed by a method known per se, for example, the methods described in Jikken Kagaku Koza (Courses in Experimental
35 Chemistry), 5th Edition, Volumes 13-19 (The Chemical Society

of Japan ed.); Shin Jikken Kagaku Koza (New Experimental Chemistry Course), Volumes 14-15 (The Chemical Society of Japan ed.); Seimitsu Yuki Gosei (Reactions and Syntheses: In the Organic Chemistry Laboratory), Revised 2nd Edition (L. F. Tietze, Th. Eicher, Nankodo Co., Ltd.); Revised Edition Organic Name Reactions; The Reaction Mechanism and Essence (Hideo Togo, Kodansha Ltd.); ORGANIC SYNTHESSES Collective Volumes I-VII (John Wiley & Sons, Inc.); Modern Organic Synthesis in the Laboratory A Collection of Standard Experimental Procedures (Jie Jack Li, OXFORD UNIVERSITY PRESS); Comprehensive Heterocyclic Chemistry III, Vol.1 - Vol.14 (Elsevier Japan KK); Strategic Applications of Named Reactions in Organic Synthesis (translated by Kiyoshi Tomioka, Kagaku-Dojin Publishing Company, INC); Comprehensive Organic Transformations (VCH Publishers Inc.) 1989 and the like, or the methods described in the Examples.

[0237]

In each step, protection or deprotection of functional groups is performed according to a method known per se, for example, the methods described in Wiley-Interscience, 2007, "Protective Groups in Organic Synthesis, 4th Ed." (Theodora W. Greene, Peter G. M. Wuts); Thieme, 2004, "Protecting Groups 3rd Ed." (P. J. Kocienski) and the like, or the methods described in the Examples.

Examples of the protecting group for hydroxyl group of alcohol and the like and phenolic hydroxyl group include ether-type protecting groups such as methoxymethyl ether, benzyl ether, t-butyldimethylsilyl ether, tetrahydropyranyl ether and the like; carboxylate-type protecting groups such as acetate and the like; sulfonate-type protecting groups such as methanesulfonate and the like; carbonate-type protecting groups such as t-butyl carbonate and the like; and the like.

Examples of the protecting group for carbonyl group of aldehyde include acetal-type protecting groups such as dimethyl acetal and the like; cyclic acetal-type protecting

groups such as cyclic 1,3-dioxane and the like; and the like.

Examples of the protecting group for carbonyl group of ketone include ketal-type protecting groups such as dimethyl ketal and the like; cyclic ketal-type protecting groups such as cyclic 1,3-dioxane and the like; oxime-type protecting groups such as O-methyloxime and the like; hydrazone-type protecting groups such as N,N-dimethylhydrazone and the like; and the like.

Examples of the protecting group for carboxyl group include ester-type protecting groups such as methyl ester and the like; amide-type protecting groups such as N,N-dimethylamide and the like; and the like.

Examples of the protecting group for thiol include ether-type protecting groups such as benzyl thioether and the like; ester-type protecting groups such as thioacetate, thiocarbonate, thiocarbamate and the like; and the like.

Examples of the protecting group for amino group, and aromatic heterocycle such as imidazole, pyrrole, indole and the like include carbamate-type protecting groups such as benzyl carbamate, tert-butyl carbamate and the like; amide-type protecting groups such as acetamide and the like; alkylamine-type protecting groups such as N-triphenylmethylamine and the like; sulfonamide-type protecting groups such as methanesulfonamide and the like; and the like.

Protecting groups can be removed by a method known per se, for example, methods using acid, base, ultraviolet rays, hydrazine, phenylhydrazine, sodium N-methyldithiocarbamate, tetrabutylammonium fluoride, palladium acetate, trialkylsilyl halide (e.g., trimethylsilyl iodide, trimethylsilyl bromide), reduction methods and the like.

[0238]

When a reduction reaction is performed in each step, the reducing agent to be used includes metal hydrides such as lithium aluminum hydride, sodium triacetoxyborohydride, sodium cyanoborohydride, diisobutylaluminum hydride (DIBAL-H), sodium

borohydride, tetramethylammonium triacetoxyborohydride and the like; boranes such as borane-tetrahydrofuran complex and the like; Raney-nickel; Raney-cobalt; hydrogen; formic acid and the like. When a carbon-carbon double bond or triple bond is
5 reduced, a method using a catalyst such as palladium-carbon, Lindlar catalyst and the like can be used.

[0239]

When oxidation reaction is performed in each step, the oxidizing agent to be used includes peracids such as m-
10 chloroperbenzoic acid (MCPBA), hydrogen peroxide, t-butyl hydroperoxide and the like; perchlorates such as tetrabutylammonium perchlorate and the like; chlorates such as sodium chlorate and the like; chlorites such as sodium chlorite and the like; periodic acids such as sodium periodate
15 and the like; hypervalent iodine reagents such as iodosylbenzene and the like; reagents having manganese such as manganese dioxide, potassium permanganate and the like; lead compounds such as lead tetraacetate and the like; reagents having chrome such as pyridinium chlorochromate (PCC),
20 pyridinium dichromate (PDC), Jones reagent and the like; halogen compounds such as N-bromosuccinimide (NBS) and the like; oxygen; ozone; sulfur trioxide-pyridine complex; osmium tetroxide; selenium dioxide; 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) and the like.

25 [0240]

When radical cyclization reaction is performed in each step, the radical initiator to be used includes azo compounds such as azobisisobutyronitrile (AIBN) and the like; water-soluble radical initiators such as 4,4'-azobis-4-
30 cyanopentanoic acid (ACPA) and the like; triethylboron in the presence of air or oxygen; benzoyl peroxide and the like. The radical reaction reagent to be used includes tributylstannane, tris(trimethylsilyl)silane, 1,1,2,2-tetraphenyldisilane, diphenylsilane, samarium iodide and the like.

35 [0241]

When the Wittig reaction is performed in each step, the Wittig reagent to be used includes alkylidene phosphoranes and the like. Alkylidene phosphoranes can be prepared by a method known per se, for example, by reacting a phosphonium salt with
5 a strong base.

[0242]

When the Horner-Emmons reaction is performed in each step, the reagent to be used includes phosphonoacetates such as methyl dimethylphosphonoacetate, ethyl diethylphosphonoacetate,
10 tert-butyl diethylphosphonoacetate and the like; bases such as alkali metal hydrides, organic lithiums, 1,8-diazabicyclo[5.4.0]undec-7-ene and the like; lithium chloride, sodium iodide and the like.

[0243]

15 When the Friedel-Crafts reaction is performed in each step, the reagent to be used includes Lewis acid and acid chloride or alkylating agents (e.g., alkyl halides, alcohol, olefins and the like). Alternatively, organic acid and inorganic acid can also be used instead of Lewis acid, and
20 acid anhydrides such as acetic anhydride and the like can also be used instead of acid chloride.

[0244]

When an aromatic nucleophilic substitution reaction is performed in each step, nucleophilic agents (e.g., amines,
25 imidazole and the like) and bases (e.g., basic salts, organic bases and the like) are used as the reagents.

[0245]

When a nucleophilic addition reaction by a nucleophilic agent, a nucleophilic addition reaction by carbanion, a
30 nucleophilic 1,4-addition reaction by carbanion (Michael addition) or a nucleophilic substitution reaction by carbanion is performed in each step, a base used to generate carbanion includes organic lithiums, metal alkoxides, inorganic bases, organic bases and the like.

35 [0246]

When the Grignard reaction is performed in each step, the Grignard reagent includes arylmagnesium halides such as phenylmagnesium bromide and the like; alkylmagnesium halides such as methylmagnesium bromide and the like. Grignard reagents can be prepared by a method known per se, for example, by reacting alkyl halide or aryl halide with metal magnesium in ether or tetrahydrofuran as a solvent.

[0247]

When the Knoevenagel condensation is performed in each step, active methylene compounds (e.g., malonic acid, diethyl malonate, malononitrile and the like) and bases (e.g., organic bases, metal alkoxides, inorganic bases) sandwiched between two electron withdrawing groups are used as the reagents.

[0248]

When the Vilsmeier-Haack reaction is performed in each step, phosphoryl chloride and amide derivative (e.g., N,N-dimethylformamide and the like) are used as the reagents.

[0249]

When an azidation reaction of alcohols, alkyl halides or sulfonates is performed in each step, the azidating agent to be used includes diphenylphosphoryl azide (DPPA), trimethylsilyl azide, sodium azide and the like. For example, when alcohols are azidated, a method using diphenylphosphoryl azide and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), a method using trimethylsilyl azide and Lewis acid, and the like can be used.

[0250]

When a reductive amination reaction is performed in each step, the reducing agent to be used includes sodium triacetoxymethylborohydride, sodium cyanoborohydride, hydrogen, formic acid, sodium borohydride, 2-picoline-borane complex and the like. When the substrate is an amine compound, a carbonyl compound to be used includes paraformaldehyde, aldehydes such as acetaldehyde and the like, and ketones such as cyclohexanone and the like. When the substrate is a carbonyl

compound, amines to be used includes primary amines such as ammonia, methylamine and the like; secondary amines such as dimethylamine and the like, and the like. An additive such as trimethyl orthoformate and the like may be added to the

5 reaction.

[0251]

When the Mitsunobu reaction is performed in each step, azodicarboxylates (e.g., diethyl azodicarboxylate (DEAD), diisopropyl azodicarboxylate (DIAD) and the like) and

10 triphenylphosphine are used as the reagents.

[0252]

When an esterification reaction, an amidation reaction or a ureation reaction is performed in each step, the reagent to be used includes acyl halides such as acid chloride, acid

15 bromide and the like; and activated carboxylic acids such as acid anhydride, active ester form, sulfuric acid ester form and the like. An activator of carboxylic acid includes

carbodiimide-type condensing agents such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (WSCD) and the

20 like; triazine-type condensing agents such as 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride-n-hydrate (DMT-MM) and the like; carbonate-type condensing agents such as 1,1-carbonyldiimidazole (CDI) and the like; diphenylphosphoryl azide (DPPA); benzotriazol-1-yloxy-

25 trisdimethylaminophosphonium salt (BOP reagent); 2-chloro-1-methyl-pyridinium iodide (Mukaiyama reagent); thionyl chloride; lower alkyl haloformates such as ethyl chloroformate and the like; O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU); sulfuric acid;

30 or combinations of these and the like. When a carbodiimide-type condensing agent is used, an additive such as 1-hydroxybenzotriazole (HOBt), N-hydroxysuccinimide (HOSu), dimethylaminopyridine (DMAP) and the like may be further added to the reaction.

35 [0253]

When a coupling reaction is performed in each step, the metal catalyst to be used includes palladium compounds such as palladium(II) acetate, tetrakis(triphenylphosphine)palladium(0),
5 dichlorobis(triphenylphosphine)palladium(II),
dichlorobis(triethylphosphine)palladium(II),
tris(dibenzylideneacetone)dipalladium(0), 1,1'-bis(diphenylphosphino)ferrocene palladium(II) chloride, palladium(II) acetate and the like; nickel compounds such as
10 tetrakis(triphenylphosphine)nickel(0) and the like; rhodium compounds such as tris(triphenylphosphine)rhodium(III) chloride and the like; cobalt compound; copper compounds such as copper oxide, copper(I) iodide and the like; platinum compound and the like. Furthermore, a base may be added to the
15 reaction, and examples of such base include inorganic bases, basic salts and the like.

[0254]

When a thiocarbonylation reaction is performed in each step, diphosphorus pentasulfide is typically used as a
20 thiocarbonylating agent. However, a reagent having a 1,3,2,4-dithiadiphosphetane-2,4-disulfide structure such as 2,4-bis(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane-2,4-disulfide (Lawesson's reagent) and the like may be used besides diphosphorus pentasulfide.

25 [0255]

When the Wohl-Ziegler reaction is performed in each step, the halogenating agent to be used includes N-iodosuccinimide, N-bromosuccinimide (NBS), N-chlorosuccinimide (NCS), bromine, sulfonyl chloride and the like. Furthermore, the reaction can
30 be accelerated by adding a radical initiator such as heat, light, benzoyl peroxide, azobisisobutyronitrile and the like to the reaction.

[0256]

When halogenation of hydroxy group is performed in each
35 step, the halogenating agent to be used includes hydrohalic

acid and acid halide of inorganic acid, specifically, hydrochloric acid, thionyl chloride, phosphorus oxychloride and the like for chlorination, and 48% hydrobromic acid and the like for bromination. In addition, a method for obtaining
5 alkyl halide from alcohol by a reaction of triphenylphosphine and carbon tetrachloride or carbon tetrabromide and the like may be used. Alternatively, a method for synthesizing alkyl halide by a two-step reaction including converting alcohol to sulfonate and reacting same with lithium bromide, lithium
10 chloride or sodium iodide may be used.

[0257]

When the Arbuzov reaction is performed in each step, the reagent to be used includes alkyl halides such as ethyl bromoacetate and the like; and phosphites such as triethyl
15 phosphite, tri(isopropyl)phosphite and the like.

[0258]

When a sulfonylation reaction is performed in each step, the sulfonylating agent to be used includes methanesulfonyl chloride, p-toluenesulfonyl chloride, methanesulfonic
20 anhydride, p-toluenesulfonic anhydride and the like.

[0259]

When hydrolysis is performed in each step, an acid or a base is used as a reagent. When acid hydrolysis of t-butyl ester is performed, formic acid, triethylsilane and the like
25 are sometimes added to reductively trap the by-produced t-butyl cation.

[0260]

When a dehydration reaction is performed in each step, the dehydrating agent to be used includes sulfuric acid,
30 phosphorus pentoxide, phosphorus oxychloride, N,N'-dicyclohexylcarbodiimide, alumina, polyphosphoric acid and the like.

[0261]

When the Corey-Chaykovsky cyclopropanation reaction is
35 performed in each step, the reagent to be used includes

trimethylsulfoxonium iodide, potassium tert-butoxide, sodium hydride and the like.

[0262]

When a carbon monoxide insertion reaction and a
5 subsequent nucleophilic substitution reaction by alcohol are performed in each step, the metal catalyst to be used includes palladium compounds such as palladium(II) acetate, tetrakis(triphenylphosphine)palladium(0), dichlorobis(triphenylphosphine)palladium(II),
10 dichlorobis(triethylphosphine)palladium(II), tris(dibenzylideneacetone)dipalladium(0), 1,1'-bis(diphenylphosphino)ferrocene palladium(II) chloride, palladium(II) acetate and the like; and examples of the phosphine ligand include triphenylphosphine, 2,2'-
15 bis(diphenylphosphino)-1,1'-binaphthyl, 2-(di-tert-butylphosphino)biphenyl, 2-(dicyclohexylphosphino)biphenyl, 2-(dicyclohexylphosphino)-2',6'-dimethoxy-1,1'-biphenyl, 2-(dicyclohexylphosphino)-2',4',6'-triisopropyl-1,1'-biphenyl, 2-(dicyclohexylphosphino)-2'-(N,N-dimethylamino)biphenyl,
20 1,1'-bis(diphenylphosphino)ferrocene, tri-tert-butylphosphine, tricyclohexylphosphine, (9,9-dimethyl-9H-xanthene-4,5-diyl)bis(diphenylphosphine) and the like.

Examples of the alcohol include methanol, ethanol, 2-propanol, benzyl alcohol and the like.

25 This reaction is performed using, as necessary, a phosphine ligand generally under a carbon monoxide atmosphere from normal pressure to about 10 atm in an inert solvent.

[0263]

When the Curtius rearrangement reaction is performed in
30 each step, the reagents to be used includes diphenylphosphoryl azide and tert-butyl alcohol and the like. As reagents, ethyl chloroformate, a base, sodium azide, tert-butyl alcohol and the like can also be used.

When a diastereomeric salt resolution method is performed
35 in each step, the optically active organic base to be used

includes, for example, ((1R,4aS,10aR)-7-isopropyl-1,4a-dimethyl-1,2,3,4,4a,9,10,10a-octahydrophenanthren-1-yl)methanamine, (R)-2-amino-3-phenylpropan-1-ol and the like.

The compound obtained in each step may be used after
5 optical resolution by a known means such as chiral column chromatography, optical fractional crystallization, diastereomer derivatization and the like.

[0264]


In the schemes, R⁵ and R⁶ are alkyl groups such as methyl
10 group, ethyl group, tert-butyl group and the like, and other symbols are each as defined above. Compound (Ia) to compound (Ik), compound (Iaa) and compound (Iga) are included in the aforementioned compound (I), and each of them shows a compound group of compound (I) wherein R³=H. The thiophene ring,
15 pyrazole ring, naphthalene ring, dihydrobenzofuran ring, pyridine ring and thiazole ring each optionally has substituent(s) on the ring.

[0265]

<Reaction scheme 1>

20 Compound (Ia) can be produced from compound (1) by the following method.

[0266]




25 [0267]

<Reaction scheme 2>

Compound (Ib) can be produced from compound (10) by the following method.

[0268]



[0269]

Compound (11) can be produced by reacting compound (10) with N-phenylbis(trifluoromethanesulfonimide) and a base. Examples of the base include sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, cesium carbonate, tripotassium phosphate, sodium methoxide, sodium ethoxide, triethylamine, diisopropylethylamine, pyridine, 1,8-diazabicyclo[5.4.0]undec-7-ene and the like.

Compound (12) can be produced by reacting compound (11) with acrylates. This reaction is performed using a base and a palladium catalyst and, where necessary, a phosphine ligand may also be used. The acrylate to be used includes tert-butyl acrylate, methyl acrylate, ethyl acrylate and the like. The palladium catalyst to be used includes palladium(II) acetate, tetrakis(triphenylphosphine)palladium(0), dichlorobis(triphenylphosphine)palladium(II), dichlorobis(triethylphosphine)palladium(II), tris(dibenzylideneacetone)dipalladium(0), 1,1'-bis(diphenylphosphino)ferrocene palladium(II) chloride, palladium(II) acetate and the like. Examples of the phosphine ligand include triphenylphosphine, 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, 2-(di-tert-butylphosphino)biphenyl, 2-(dicyclohexylphosphino)biphenyl, 2-(dicyclohexylphosphino)-2',6'-dimethoxy-1,1'-biphenyl, 2-(dicyclohexylphosphino)-2',4',6'-triisopropyl-1,1'-biphenyl,

2-(dicyclohexylphosphino)-2'-(N,N-dimethylamino)biphenyl,
1,1'-bis(diphenylphosphino)ferrocene, tri-tert-butylphosphine,
tricyclohexylphosphine, (9,9-dimethyl-9H-xanthene-4,5-
diyl)bis(diphenylphosphine), tri(o-tolyl)phosphine and the


5 like. Examples of the base include sodium hydroxide, potassium
hydroxide, sodium carbonate, potassium carbonate, cesium
carbonate, tripotassium phosphate, sodium methoxide, sodium
ethoxide, triethylamine, diisopropylethylamine, pyridine, 1,8-
diazabicyclo[5.4.0]undec-7-ene and the like. In this step, for
10 example, a microwave synthesizer such as Initiator
manufactured by Biotage and the like may also be used.

[0270]

<Reaction scheme 3>

Compound (Ic) can be produced from compound (19) by the
15 following method.

[0271]




[0272]

<Reaction scheme 4>

20 Compound (Id) can be produced from compound (31) by the
following method.

[0273]



[0274]


Compound (32) can be produced by reacting compound (31) with acrylates. This reaction is performed using a base and a palladium catalyst, as necessary, and a phosphine ligand may also be used. The acrylate to be used includes tert-butyl acrylate, methyl acrylate, ethyl acrylate and the like. The palladium catalyst to be used includes palladium(II) acetate, tetrakis(triphenylphosphine)palladium(0), dichlorobis(triphenylphosphine)palladium(II), dichlorobis(triethylphosphine)palladium(II), tris(dibenzylideneacetone)dipalladium(0), 1,1'-bis(diphenylphosphino)ferrocene palladium(II) chloride, palladium(II) acetate and the like. Examples of the phosphine ligand include triphenylphosphine, 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, 2-(di-tert-butylphosphino)biphenyl, 2-(dicyclohexylphosphino)biphenyl, 2-(dicyclohexylphosphino)-2',6'-dimethoxy-1,1'-biphenyl, 2-(dicyclohexylphosphino)-2',4',6'-triisopropyl-1,1'-biphenyl, 2-(dicyclohexylphosphino)-2'-(N,N-dimethylamino)biphenyl, 1,1'-bis(diphenylphosphino)ferrocene, tri-tert-butylphosphine, tricyclohexylphosphine, (9,9-dimethyl-9H-xanthene-4,5-diyl)bis(diphenylphosphine), tri(o-tolyl)phosphine and the like. Examples of the base include sodium hydroxide, potassium

[0279]

<Reaction scheme 7>

Compound (Ig) can be produced from compound (58) by the following method.

5 [0280]




[0281]

<Reaction scheme 8>

Compound (Ih) can be produced from compound (71) by the following method.

10 [0282]




[0283]

<Reaction scheme 9>

Compound (Ii) can be produced from compound (81) by the following method.

[0284]




5 [0285]

<Reaction scheme 10>

Compound (Ij) can be produced from compound (70) by the following method.

[0286]




[0287]

<Reaction scheme 11>

Compound (Ik) can be produced from compound (89) by the following method.

15 [0288]




[0289]

<Reaction scheme 12>

Compound (Iaa) can be produced from compound (6) or compound (4) by the following method.

[0290]




5

[0291]

<Reaction scheme 13>

Compound (Iga) can be produced from compound (61) or compound (63) by the following method.

[0292]



Examples

[0293]

The present invention is explained in detail in the following by referring to Examples, Experimental Examples and Formulation Examples. However, the examples do not limit the present invention and the present invention can be modified within the scope of the present invention.

[0294]

The "room temperature" in the following Examples is generally about 10°C to about 35°C. The ratio for a mixed solvent is, unless otherwise specified, a volume mixing ratio and % means wt% unless otherwise specified.

[0295]

In silica gel column chromatography, the indication of NH means use of aminopropylsilane-bonded silica gel. In HPLC (high performance liquid chromatography), the indication of C18 means use of octadecyl-bonded silica gel. Unless otherwise specified, the ratio of elution solvents is a volume mixing

ratio.

[0296]

In the following Examples, the following abbreviations are used.

5 mp: melting point

MS: mass spectrum

[M+H]⁺, [M-H]⁻: molecular ion peak

M: molar concentration

N: normality

10 CDCl₃: deuterated chloroform

DMSO-d₆: deuterated dimethyl sulfoxide

¹H NMR: proton nuclear magnetic resonance

LC/MS: liquid chromatography mass spectrometer

ESI: ElectroSpray Ionization

15 APCI: Atmospheric Pressure Chemical Ionization

THF: tetrahydrofuran

DMF: N,N-dimethylformamide

DMSO: dimethyl sulfoxide

TFA: trifluoroacetic acid

20 SFC: supercritical fluid chromatography

TEA: triethylamine

AA: acetic acid

DMAP: 4-dimethylaminopyridine

[0297]

25 ¹H NMR was measured by Fourier-transform NMR. For the analyses, ACD/SpecManager (trade name) and the like were used. A peak showing very mild proton of hydroxyl group, amino group or the like is not described.

[0298]

30 MS was measured by LC/MS. As the ionization method, ESI method or APCI method was used. The data indicates measured values (found). Generally, a molecular ion peak is observed. However, when a compound has a tert-butoxycarbonyl group, a peak free of a tert-butoxycarbonyl group or tert-butyl group
35 may be observed as a fragment ion. When a compound has a

hydroxyl group, a peak free of H₂O may be observed as a fragment ion. In the case of a salt, a molecular ion peak or a fragment ion peak of a free form is generally observed.

[0299]

5 The unit of the sample concentration (c) in optical rotation ($[\alpha]_D$) is g/100 mL.

As the elemental analytical values (Anal.), Calculated (Calcd) and measured values (Found) are described.

[0300]

10 Example 1

4-(trans-2-aminocyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride
A) (E)-ethyl 3-(5-bromo-2-methylthiophene-3-yl)acrylate

To a mixture of potassium tert-butoxide (4.59 g) and dry
15 THF (60 mL) was added ethyl diethylphosphonoacetate (7.5 mL) at -5°C, and the mixture was stirred for 20 min. To the reaction mixture was added a mixture of 5-bromo-2-methylthiophene-3-carbaldehyde (7.0 g) and dry THF (10 mL) at -5°C, and the mixture was stirred for 30 min. To the reaction
20 mixture was added water (100 mL), and the mixture was extracted twice with ethyl acetate (100 mL each). The extracts were combined, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (7.2 g).

25 ¹H-NMR (400 MHz, DMSO-d₆) δ 1.17-1.35 (3H, m), 2.45 (3H, s), 4.19 (2H, q, J = 5.8 Hz), 6.40 (1H, d, J = 15.7 Hz), 7.48 (1H, d, J = 15.7 Hz), 7.78 (1H, s).

[0301]

B) ethyl trans-2-(5-bromo-2-methylthiophen-3-yl)cyclopropanecarboxylate
30

To a suspension of sodium hydride (50% in oil, 2.26 g) in DMSO (50 mL) was added trimethylsulfoxonium iodide (10.4 g), and the mixture was stirred at room temperature for 1 hr. To the reaction mixture was added a mixture of ethyl (E)-ethyl 3-(5-bromo-2-methylthiophen-3-yl)acrylate (6.5 g) and DMSO (30
35

mL), and the mixture was stirred at room temperature for 4 hr. To the reaction mixture was added ice water (200 mL), and the mixture was extracted with ethyl acetate (100 mL, twice). The extracts were combined, washed with saturated brine (100 mL),
5 dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (3.5 g).

¹H NMR (400 MHz, DMSO-d₆) δ 1.23 (3H, t, J = 5.2 Hz), 1.24-1.29
10 (1H, m), 1.30-1.40 (1H, m), 1.86-1.90 (1H, m), 2.23-2.28 (1H, m), 2.33 (3H, s), 4.09 (2H, q, J = 7.1 Hz), 6.82 (1H, s).

[0302]

C) trans-2-(5-bromo-2-methylthiophen-3-yl)cyclopropanecarboxylic acid

15 Ethyl trans-2-(5-bromo-2-methylthiophen-3-yl)cyclopropanecarboxylate (500 mg) was dissolved in methanol (8 mL), a mixture of sodium hydroxide (138 mg) and water (2 mL) was added at 0°C, and the mixture was stirred at room temperature for 2 hr. The reaction mixture was concentrated
20 under reduced pressure, and adjusted to pH 6 with 2 mol/L hydrochloric acid at an inside temperature of not more than 10°C. The precipitated solid was collected by filtration and dried under reduced pressure to give the title compound (350 mg).

25 ¹H NMR (400 MHz, DMSO-d₆) δ 1.22-1.27 (1H, m), 1.31-1.36 (1H, m), 1.72-1.76 (1H, m), 2.19-2.24 (1H, m), 2.34 (3H, s), 6.79 (1H, s), 12.31 (1H, brs).

[0303]

30 D) tert-butyl (trans-2-(5-bromo-2-methylthiophen-3-yl)cyclopropyl)carbamate

To a mixture of trans-2-(5-bromo-2-methylthiophen-3-yl)cyclopropanecarboxylic acid (1.5 g) and tert-butyl alcohol (70 mL) were added triethylamine (8.3 mL) and diphenylphosphoryl azide (3.7 mL) at room temperature, and the
35 mixture was stirred at 90°C for 16 hr. The reaction mixture

was cooled to room temperature, and concentrated under reduced pressure. To the residue was added water (100 mL), and the mixture was extracted twice with ethyl acetate (200 mL each). The extracts were combined, washed with saturated brine (100
5 mL), dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (0.75 g).

¹H NMR (400 MHz, DMSO-d₆) δ 0.99-1.05 (2H, m), 1.38 (9H, s),
10 1.72-1.76 (1H, m), 2.35 (3H, s), 2.52-2.54 (1H, m), 6.72 (1H, s), 7.20 (1H, s).

[0304]

E) methyl 4-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylate

15 A mixture of tert-butyl (trans-2-(5-bromo-2-methylthiophen-3-yl)cyclopropyl)carbamate (1.3 g), diisopropylethylamine (3.42 mL), dichloro(1,1'-bis(diphenylphosphino)ferrocene)palladium (0.96 g) and methanol (45 mL) was heated under a carbon monoxide atmosphere
20 (10 atm) at 80°C for 16 hr. After cooling to room temperature, the insoluble material was filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (0.90 g).

25 ¹H NMR (400 MHz, DMSO-d₆) δ 1.05-1.15 (2H, m), 1.36 (9H, s), 1.73-1.78 (1H, m), 2.47 (3H, s), 2.50-2.55 (1H, m), 3.76 (3H, s), 7.25 (1H, brs), 7.34 (1H, s).

[0305]

F) 4-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylic acid
30

Methyl 4-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylate (900 mg) was dissolved in methanol (45 mL), a mixture of sodium hydroxide (1160 mg) and water (15 mL) was
35 added at 0°C, and the mixture was stirred at room temperature

for 16 hr. The reaction mixture was concentrated under reduced pressure, and water (10 mL) was added to the residue. The mixture was adjusted to pH 6 with 2 mol/L hydrochloric acid at an inside temperature of not more than 10°C, and extracted with
5 20% methanol-dichloromethane solution (100 mL each). The extracts were combined, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (660 mg).

¹H NMR (400 MHz, DMSO-d₆) δ 1.02-1.07 (2H, m), 1.38 (9H, s),
10 1.72-1.76 (1H, m), 2.45 (3H, s), 2.49-2.54 (1H, m), 7.24 (2H, s), 12.79 (1H, s).

[0306]

G) tert-butyl (trans-2-(2-methyl-5-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-3-yl)cyclopropyl)carbamate
15 4-(trans-2-((tert-Butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylic acid (100 mg), 5-methyl-1,3,4-thiadiazol-2-amine (44.8 mg) and triethylamine (0.187 mL) were dissolved in DMF (3.0 mL), and the mixture was cooled to 0°C. O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
20 hexafluorophosphate (153 mg) was added, and the mixture was stirred at room temperature overnight. The reaction mixture was poured into water, and the mixture was extracted with ethyl acetate. The extract was successively washed with saturated aqueous sodium hydrogen carbonate solution and
25 saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate) and concentrated under reduced pressure. The obtained residue was diluted with ethyl acetate-THF, successively
30 washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (64 mg).

MS: [M+H]⁺ 395.2.

[0307]

35 H) 4-(trans-2-aminocyclopropyl)-5-methyl-N-(5-methyl-1,3,4-

thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

To a mixture of tert-butyl (trans-2-(2-methyl-5-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-3-yl)cyclopropyl)carbamate (64 mg), ethyl acetate (5 mL) and
5 methanol (2 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (0.811 mL) at room temperature, and the mixture was stirred at room temperature for 3 days. The precipitated solid was collected by filtration and dried under reduced pressure to give the title compound (45 mg).

10 [0308]

Example 2

4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

15 To a mixture of 4-(trans-2-aminocyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride (42 mg), sodium hydrogen carbonate (38.4 mg), THF (0.5 mL) and methanol (0.5 mL) was added cyclopropanecarbaldehyde (13 µL) at room temperature. Under a
20 nitrogen atmosphere, the reaction mixture was stirred at 50°C for 1.5 hr and at room temperature for 1 hr. To the reaction mixture was added cyclopropanecarbaldehyde (8.8 µL) at room temperature, and the mixture was stirred at 50°C for 1.5 hr under a nitrogen atmosphere and at room temperature for 30 min.
25 Sodium borohydride (8.7 mg) was added under ice-cooling and the reaction mixture was stirred for 30 min under a nitrogen atmosphere. Under ice-cooling, to the reaction mixture were added ethyl acetate (5 mL), water and saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted
30 with ethyl acetate. The extract was washed with saturated brine, and the aqueous layer was extracted with a mixture of ethyl acetate and THF. The combined organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was washed with a mixture
35 of ethyl acetate and diisopropyl ether, and dissolved in

methanol. A 2 mol/L hydrogen chloride/methanol solution (2 mL) was added, and the mixture was concentrated under reduced pressure. The residue was crystallized from ethanol/diisopropyl ether to give the title compound (25.9 mg).

5 ¹H NMR (300 MHz, DMSO-d₆) δ 0.33-0.43 (2H, m), 0.55-0.65 (2H, m), 1.06-1.14 (1H, m), 1.16-1.27 (1H, m), 1.53-1.62 (1H, m), 2.40-2.47 (1H, m), 2.52 (3H, s), 2.62 (3H, s), 2.85-3.02 (3H, m), 7.85 (1H, brs), 9.24 (2H, brs), 12.66 (1H, brs).

[0309]

10 Example 3

5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-1-methyl-1H-pyrazole-3-carboxamide hydrochloride

A) methyl 1-methyl-5-(((trifluoromethyl)sulfonyl)oxy)-1H-pyrazole-3-carboxylate

To a mixture of methyl 5-hydroxy-1-methyl-1H-pyrazole-3-carboxylate (2.404 g), diisopropylethylamine (5.38 mL) and DMF (24 mL) was added N-phenylbis(trifluoromethanesulfonimide) (6.05 g) at 0°C, and the mixture was stirred at room temperature overnight under a nitrogen atmosphere. The reaction mixture was poured into water, extracted with ethyl acetate, washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate) and concentrated under reduced pressure to give the title compound (3.92 g).

¹H NMR (300 MHz, CDCl₃) δ 3.92 (3H, s), 3.94 (3H, s), 6.69 (1H, s).

[0310]

30 B) methyl 5-((1E)-3-tert-butoxy-3-oxoprop-1-en-1-yl)-1-methyl-1H-pyrazole-3-carboxylate

A mixture of methyl 1-methyl-5-(((trifluoromethyl)sulfonyl)oxy)-1H-pyrazole-3-carboxylate (890 mg), tert-butyl acrylate (1.35 mL), tri(o-tolyl)phosphine (188 mg), palladium acetate (69.3 mg), triethylamine (1.29 mL)

and DMF (8 mL) was stirred at 80°C for 3 hr under a nitrogen atmosphere. The reaction mixture was poured into water at room temperature, and the mixture was extracted with ethyl acetate, washed with saturated brine, dried over anhydrous sodium

5 sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate) and concentrated under reduced pressure to give the title compound (144 mg).

¹H NMR (300 MHz, CDCl₃) δ 1.53 (9H, s), 3.93 (3H, s), 4.02 (3H, s), 6.34 (1H, d, J = 15.9 Hz), 7.07 (1H, s), 7.45 (1H, d, J = 15.9 Hz).

[0311]

C) methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)-1-methyl-1H-pyrazole-3-carboxylate

15 By a method similar to that of Example 1, step B, the title compound was obtained.

MS: [M+H]⁺ 281.2.

[0312]

D) 2-(3-(methoxycarbonyl)-1-methyl-1H-pyrazol-5-yl)cyclopropanecarboxylic acid

To methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)-1-methyl-1H-pyrazole-3-carboxylate (135 mg) was added trifluoroacetic acid (2 mL) at 0°C, and the mixture was stirred at room temperature for 4 hr and concentrated under reduced pressure. Water was added to the residue, and the mixture was extracted with ethyl acetate, successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was diluted with ethyl acetate and saturated aqueous sodium hydrogen carbonate solution. The aqueous layer was separated, acidified with 2 mol/L hydrochloric acid, and extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was diluted with toluene and

concentrated under reduced pressure to give the title compound (100 mg).

MS: [M+H]⁺225.1.

[0313]

5 E) 5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-1-methyl-1H-pyrazole-3-carboxamide hydrochloride

By a method similar to that of Example 1, steps D, F, G and H and Example 2, the title compound was obtained.

10 [0314]

Example 4

5-(trans-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)-1-methyl-1H-pyrazole-3-carboxamide hydrochloride

By a method similar to that of Example 3, steps A, B, C and D and Example 1, steps D, F, G and H, the title compound was obtained.

[0315]

Example 5

4-(trans-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)-2-naphthamide dihydrochloride

A) 3-bromo-N-methoxy-N-methyl-1-naphthamide

3-Bromo-1-naphthoic acid (3.00 g), N,O-dimethylhydroxylamine hydrochloride (1.32 g), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (2.80 g) and
25 anhydrous 1-hydroxybenzotriazole (1.94 g) were dissolved in anhydrous DMF (60.0 mL), triethylamine (4.16 mL) was added, and the mixture was stirred at room temperature for 16 hr. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The extract was successively
30 washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (2.69 g).

MS: [M+H]⁺ 293.9.

35 [0316]

B) 3-bromo-1-naphthaldehyde

To a solution of 3-bromo-N-methoxy-N-methyl-1-naphthamide (2.69 g) in THF (50.0 mL) was slowly added 1.5 mol/L diisobutylaluminum hydride toluene solution (9.15 mL) at 0°C, and the mixture was stirred at 0°C for 2 hr under a nitrogen atmosphere. To the reaction mixture was added 1 mol/L hydrochloric acid at 0°C, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.52 g).
¹H NMR (300 MHz, CDCl₃) δ 7.57-7.66 (1H, m), 7.67-7.74 (1H, m), 7.84 (1H, d, J = 8.3 Hz), 8.06 (1H, d, J = 2.3 Hz), 8.26 (1H, s), 9.16 (1H, d, J = 8.7 Hz), 10.36 (1H, s).

[0317]

C) (E)-tert-butyl 3-(3-bromonaphthalen-1-yl)acrylate

Lithium chloride (0.28 g) was dried in vacuo, and acetonitrile (20.0 mL) was added. To the mixture were added 3-bromo-1-naphthaldehyde (1.52 g) and tert-butyl diethylphosphonoacetate (1.71 g) at 0°C, and the mixture was stirred at 0°C for 5 min under a nitrogen atmosphere. To the reaction mixture was added 1,8-diazabicyclo[5.4.0]undec-7-ene (1.02 mL) at 0°C, and the mixture was stirred at room temperature for 18 hr under a nitrogen atmosphere. Water was added to the reaction mixture at room temperature, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.93 g).
¹H NMR (300 MHz, CDCl₃) δ 1.57 (9H, s), 6.46 (1H, d, J = 15.9 Hz), 7.51-7.61 (2H, m), 7.75-7.82 (2H, m), 8.02 (1H, d, J = 1.5 Hz), 8.10-8.16 (1H, m), 8.32 (1H, d, J = 15.9 Hz).

[0318]

D) 4-(trans-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)-2-naphthamide dihydrochloride

By a method similar to that of Example 1, steps B, D, E, F, G and H and Example 3, step D, the title compound was obtained.

[0319]

Example 6

N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-2-naphthamide hydrochloride

A) tert-butyl (trans-2-(3-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)naphthalen-1-yl)cyclopropyl)((tetrahydro-2H-pyran-4-yl)methyl)carbamate

4-(trans-2-Aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)-2-naphthamide dihydrochloride (56.6 mg) and sodium hydrogen carbonate (35.9 mg) were dissolved in THF (5.00 mL)/methanol (5.00 mL), and tetrahydro-2H-pyran-4-carbaldehyde (19.5 mg) was added. Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 3 hr, and sodium borohydride (10.8 mg) was added at 0°C. The reaction mixture was stirred at room temperature for 2 hr, to the reaction mixture was added di-tert-butyl dicarbonate (0.099 mL), and the mixture was stirred at room temperature for 16 hr. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (33.5 mg). MS: [M+H]⁺ 523.1.

[0320]

B) N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-2-naphthamide hydrochloride

By a method similar to that of Example 1, step H, the title compound was obtained.

Anal. Calcd for $C_{23}H_{26}N_4O_2S \cdot HCl$: C, 60.19; H, 5.93; N, 12.21.

Found: C, 59.52; H, 5.87; N, 11.88.

5 [0321]

Example 7

4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)-2-naphthamide dihydrochloride

By a method similar to that of Example 6, the title
10 compound was obtained.

Anal. Calcd for $C_{21}H_{22}N_4OS \cdot 2HCl$: C, 55.88; H, 5.36; N, 12.41.

Found: C, 56.60; H, 5.56; N, 12.39.

[0322]

Example 8

15 N-(4,4-difluorocyclohexyl)-5-methyl-4-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide hydrochloride

A) tert-butyl (trans-2-(5-((4,4-difluorocyclohexyl)carbamoyl)-2-methylthiophen-3-yl)cyclopropyl)carbamate

20 4-(trans-2-((tert-Butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylic acid (100 mg), 4,4-difluorocyclohexanamine hydrochloride (63.5 mg) and triethylamine (0.187 mL) were dissolved in DMF (1.0 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
25 hexafluorophosphate (153 mg) was added at room temperature, and the mixture was stirred overnight. Water was added to the reaction mixture, and the resulting precipitate was collected by filtration to give the title compound (140 mg).

MS: $[M+H]^+$ 415.3.

30 [0323]

B) 4-(trans-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)-5-methylthiophene-2-carboxamide hydrochloride

To a solution (2.0 mL) of tert-butyl (trans-2-(5-((4,4-difluorocyclohexyl)carbamoyl)-2-methylthiophen-3-yl)cyclopropyl)carbamate (140 mg) in ethyl acetate was added 4
35

mol/L hydrogen chloride/ethyl acetate solution (2.0 mL) at room temperature, and the mixture was stirred at room temperature for 3 days. The resulting precipitate was collected by filtration to give the title compound (90 mg).

5 MS: $[M-HCl+H]^+$ 315.2.

[0324]

C) N-(4,4-difluorocyclohexyl)-5-methyl-4-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide hydrochloride

10 To a mixture of 4-(trans-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)-5-methylthiophene-2-carboxamide hydrochloride (48.9 mg), sodium hydrogen carbonate (46.8 mg), THF (1 mL) and methanol (1 mL) was added tetrahydro-2H-pyran-4-carbaldehyde (19.1 mg) at room temperature. The reaction
15 mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, and sodium borohydride (7.91 mg) was added at 0°C. Under a nitrogen atmosphere, the reaction mixture was stirred at room temperature for 1 hr. To the reaction mixture were added ethyl acetate, water and saturated aqueous sodium
20 hydrogen carbonate solution under ice-cooling, and the mixture was extracted with ethyl acetate. The extract was washed with water and saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The obtained residue was purified by silica
25 gel column chromatography (ethyl acetate/methanol). To the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution and the mixture was concentrated under reduced pressure to give the title compound (42.3 mg).

1H NMR (300 MHz, DMSO- d_6) δ 1.11-1.34 (3H, m), 1.54-2.12 (13H, m), 2.83-3.04 (4H, m), 3.21-3.34 (2H, m), 3.86 (3H, dd, J = 11.3, 2.3 Hz), 7.40 (1H, s), 8.12 (1H, d, J = 7.9 Hz), 8.56 (1H, brs), 9.39 (2H, brs).

[0325]

Example 9

35 7-(trans-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-

yl)-2,3-dihydro-1-benzofuran-5-carboxamide dihydrochloride

By a method similar to that of Example 1, the title compound was obtained.

[0326]

5 Example 10

7-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)-2,3-dihydro-1-benzofuran-5-carboxamide dihydrochloride

By a method similar to that of Example 2, the title
10 compound was obtained.

[0327]

Example 11

7-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-2,3-dihydro-1-benzofuran-5-carboxamide
15 hydrochloride

By a method similar to that of Example 1, steps G and H and Example 2, the title compound was obtained.

[0328]

Example 12

20 N-(4,4-difluorocyclohexyl)-7-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-2,3-dihydro-1-benzofuran-5-carboxamide hydrochloride

By a method similar to that of Example 2, the title compound was obtained.

25 [0329]

Example 13

5-(trans-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)-6-methylnicotinamide dihydrochloride

By a method similar to that of Example 1, the title
30 compound was obtained.

[0330]

Example 14

5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-6-methylnicotinamide dihydrochloride

35 By a method similar to that of Example 2, the title

compound was obtained.

[0331]

Example 15

3-(trans-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)-1-naphthamide dihydrochloride

A) (E)-methyl 3-(3-(tert-butoxy)-3-oxoprop-1-en-1-yl)-1-naphthoate

A mixture of methyl 3-bromo-1-naphthoate (500 mg), tert-butyl acrylate (0.422 mL), tri(o-tolyl)phosphine (114 mg), palladium acetate (43.2 mg), triethylamine (0.781 mL) and anhydrous DMF (3.00 mL) was heated at 100°C for 1 hr under microwave irradiation. The reaction mixture was poured into water and the mixture was extracted with ethyl acetate. The extract was washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate) and concentrated under reduced pressure to give the title compound (575 mg).

¹H NMR (300 MHz, CDCl₃) δ 1.56 (9H, s), 4.03 (3H, s), 6.55 (1H, d, J = 15.8 Hz), 7.52-7.59 (1H, m), 7.60-7.67 (1H, m), 7.74 (1H, d, J = 15.8 Hz), 7.89 (1H, d, J = 7.5 Hz), 8.07 (1H, s), 8.37 (1H, d, J = 1.9 Hz), 8.87 (1H, d, J = 8.7 Hz).

[0332]

B) 3-(trans-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)-1-naphthamide dihydrochloride

By a method similar to that of Example 1, steps B, D, F, G and H and Example 3, step D, the title compound was obtained. Anal. Calcd for C₁₇H₁₆N₄OS 2HCl H₂O: C, 49.16; H, 4.85; N, 13.49. Found: C, 49.12; H, 5.47; N, 13.12.

[0333]

Example 16

N-(5-methyl-1,3,4-thiadiazol-2-yl)-3-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-1-naphthamide dihydrochloride

By a method similar to that of Example 6, the title

compound was obtained.

Anal. Calcd for $C_{23}H_{26}N_4O_2S \cdot 2HCl \cdot 1.4H_2O$: C, 53.06; H, 5.96; N, 10.76. Found: C, 52.93; H, 5.96; N, 10.31.

[0334]

5 Example 17

4-(trans-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide 1/2 fumarate

To a mixture of 4-(trans-2-aminocyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide
10 dihydrochloride (45 mg), triethylamine (0.034 mL), trimethyl orthoformate (0.027 mL) and methanol (2.0 mL) was added cyclobutanone (0.014 mL) at room temperature, and the mixture was stirred at room temperature overnight. Sodium borohydride (9.3 mg) was added under ice-cooling, and the reaction mixture
15 was stirred at 0°C for 10 min. After stirring, water was added under ice-cooling, and the mixture was extracted with ethyl acetate. The extract was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The
20 obtained residue was purified by silica gel column chromatography (ethyl acetate/methanol) and concentrated under reduced pressure. The obtained residue was suspended in ethyl acetate, and a solution of fumaric acid (14.2 mg) in ethanol was added at room temperature. The reaction mixture was
25 stirred for 30 min, and the resulting precipitate was collected by filtration to give the title compound (23 mg).
 1H NMR (300 MHz, DMSO- d_6) δ 0.78-0.88 (1H, m), 0.91-1.02 (1H, m), 1.52-1.83 (5H, m), 2.05-2.19 (2H, m), 2.20-2.30 (2H, m), 2.47 (3H, s), 2.61 (3H, s), 6.57-6.62 (1H, m), 7.70-7.77 (1H,
30 m).

[0335]

Example 18

4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-5-methylthiophene-2-carboxamide fumarate

35 By a method similar to that of Example 19, the title

compound was synthesized.

[0336]

Example 19

4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-
5 (tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide fumarate

A) tert-butyl (trans-2-(2-methyl-5-((tetrahydro-2H-pyran-4-yl)carbamoyl)thiophen-3-yl)cyclopropyl)carbamate

4-(trans-2-((tert-Butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylic acid (100 mg), tetrahydro-2H-
10 pyran-4-amine (37.4 mg) and triethylamine (0.187 mL) were dissolved in DMF (1.0 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (153 mg) was added at room temperature, and the mixture was stirred overnight. The reaction mixture was poured into water, and the
15 resulting precipitate was collected by filtration to give the title compound (122 mg).

MS: [M+H]⁺ 381.2.

[0337]

B) 4-(trans-2-aminocyclopropyl)-5-methyl-N-(tetrahydro-2H-
20 pyran-4-yl)thiophene-2-carboxamide hydrochloride

To a mixture of tert-butyl (trans-2-(2-methyl-5-((tetrahydro-2H-pyran-4-yl)carbamoyl)thiophen-3-yl)cyclopropyl)carbamate (122 mg) and ethyl acetate (2 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (1.60
25 mL) at room temperature, and the mixture was stirred at room temperature overnight. The resulting precipitate was collected by filtration to give the title compound (95 mg).

MS: [M-HCl+H]⁺ 281.2.

[0338]

30 C) 4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide fumarate

To a mixture of 4-(trans-2-aminocyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide hydrochloride (45 mg), triethylamine (0.040 mL) and methanol
35 (2.0 mL) was added cyclopropanecarbaldehyde (14.9 mg) at room

temperature, and the reaction mixture was stirred at room temperature overnight. Sodium borohydride (10.8 mg) was added under ice-cooling, and the reaction mixture was stirred at 0°C for 10 min. After stirring, water was added under ice-cooling, and the mixture was extracted with ethyl acetate. The extract was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography (ethyl acetate/methanol) and concentrated under reduced pressure. The obtained residue was suspended in ethyl acetate, a solution of fumaric acid (13.5 mg) in ethanol was added at room temperature, and the reaction mixture was stirred for 30 min. After stirring, the resulting precipitate was collected by filtration to give the title compound (40 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.09-0.18 (2H, m), 0.38-0.47 (2H, m), 0.82-0.98 (2H, m), 1.02-1.13 (1H, m), 1.41-1.59 (2H, m), 1.66-1.77 (2H, m), 1.79-1.90 (1H, m), 2.36-2.45 (4H, m), 2.57 (2H, dd, J = 6.82, 1.89 Hz), 3.28-3.42 (2H, m), 3.80-3.96 (3H, m), 6.57 (2H, s), 7.26 (1H, s), 8.04 (1H, d, J = 7.57 Hz).

[0339]

Example 20

5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide dihydrochloride

By a method similar to that of Example 2, the title compound was synthesized.

[0340]

Example 21

N-(4,4-difluorocyclohexyl)-5-methyl-4-(trans-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-2-carboxamide fumarate 4-(trans-2-Aminocyclopropyl)-N-(4,4-difluorocyclohexyl)-5-methylthiophene-2-carboxamide hydrochloride (49 mg) and borane-2-methylpyridine complex (44.8 mg) were dissolved in methanol (2.0 mL)/acetic acid (0.20 mL), tetrahydro-4H-pyran-

4-one (21.0 mg) was added, and the mixture was stirred at room temperature overnight. Under ice-cooling, to the reaction mixture was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate.

5 The extract was washed with water and saturated brine, dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) and concentrated under reduced pressure. The obtained residue was dissolved in
10 ethyl acetate, a solution of fumaric acid (9.32 mg) in ethanol was added at room temperature, and the reaction mixture was stirred for 30 min. After stirring, the mixture was concentrated under reduced pressure to give the title compound (12 mg).

15 ¹H NMR (300 MHz, METHANOL-d₄) δ 1.04-1.15 (1H, m), 1.21-1.33 (1H, m), 1.41-2.16 (14H, m), 2.40 (3H, s), 2.64-2.74 (1H, m), 3.30-3.42 (2H, m), 3.77-3.96 (3H, m), 6.58-6.64 (2H, m), 7.14-7.17 (1H, m).

[0341]

20 Example 22

4-(trans-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide dihydrochloride

To a mixture of 4-(trans-2-aminocyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide
25 dihydrochloride (50 mg), triethylamine (0.040 mL), trimethyl orthoformate (0.032 mL) and methanol (2.0 mL) was added cyclobutanone (15.1 mg) at room temperature, and the mixture was stirred at room temperature overnight. Sodium borohydride (10.8 mg) was added under ice-cooling, and the reaction
30 mixture was stirred for 10 min. Water was added under ice-cooling, and the mixture was extracted with ethyl acetate. The extract was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The obtained
35 residue was purified by silica gel column chromatography

(ethyl acetate/methanol). To the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution and the reaction mixture was concentrated under reduced pressure. The obtained residue was crystallized from ethanol/heptane to give
5 the title compound (10 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.12-1.22 (1H, m), 1.47-1.58 (1H, m), 1.75-1.91 (2H, m), 2.14-2.32 (4H, m), 2.35-2.44 (1H, m), 2.47-2.49 (3H, m), 2.71-2.85 (1H, m), 3.80 (3H, s), 3.82-3.92 (1H, m), 7.47-7.50 (1H, m), 7.50-7.53 (1H, m), 7.89-7.95 (1H, m),
10 m), 9.47-9.67 (2H, m) 10.25-10.32 (1H, m).

[0342]

Example 23

5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide dihydrochloride

15 A) ethyl (2E)-3-(4-bromothiophen-2-yl)acrylate

Lithium chloride (453 mg) was dried in high vacuo, and acetonitrile (40 mL) was added. To this mixture were added 4-bromothiophene-2-carbaldehyde (2.00 g) and ethyl diethylphosphonoacetate (2.46 g) at 0°C, and the mixture was
20 stirred at 0°C for 5 min under a nitrogen atmosphere. To the reaction mixture was added 1,8-diazabicyclo[5.4.0]undec-7-ene (1.66 mL) at 0°C, and the mixture was stirred at room temperature overnight under a nitrogen atmosphere. Water was added to the reaction mixture, and the mixture was extracted
25 with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (2.52 g).

30 ¹H NMR (300 MHz, DMSO-d₆) δ 1.18-1.31 (3H, m), 4.18 (2H, q, J = 7.07 Hz), 6.37 (1H, d, J = 15.90 Hz), 7.58-7.89 (3H, m).

[0343]

B) ethyl trans-2-(4-bromothiophen-2-yl)cyclopropanecarboxylate

To a suspension of trimethylsulfoxonium iodide (2.54 g)
35 in DMSO (20 mL) was added sodium hydride (60% in oil, 461 mg),

and the mixture was stirred at room temperature for 30 min. To the reaction mixture was added a solution of ethyl (2E)-3-(4-bromothiophen-2-yl)acrylate (2.51 g) in DMSO (10 mL), and the mixture was stirred at room temperature overnight under a nitrogen atmosphere. Water was added to the reaction mixture at 0°C, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.27 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.20 (3H, t, J = 7.19 Hz), 1.35-1.54 (2H, m), 1.97 (1H, ddd, J = 8.42, 5.40, 3.98 Hz), 2.58-2.67 (1H, m), 4.05-4.16 (2H, m), 6.98 (1H, dd, J = 1.51, 0.76 Hz), 7.46 (1H, d, J = 1.51 Hz).

[0344]

C) trans-2-(4-bromothiophen-2-yl)cyclopropanecarboxylic acid

Ethyl trans-2-(4-bromothiophen-2-yl)cyclopropanecarboxylate (1.27 g) was dissolved in THF (5 mL) and ethanol (5 mL), 8 mol/L aqueous sodium hydroxide solution (1.44 mL) was added at 0°C, and the mixture was stirred at room temperature overnight. The reaction mixture was acidified with 1 mol/L hydrochloric acid at 0°C, and extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.11 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.29-1.50 (2H, m), 1.78-1.87 (1H, m), 2.54-2.63 (1H, m), 6.97 (1H, dd, J = 1.51, 0.76 Hz), 7.45 (1H, d, J = 1.51 Hz), 12.49 (1H, brs).

[0345]

D) tert-butyl (trans-2-(4-bromothiophen-2-yl)cyclopropyl)carbamate

To a mixture of trans-2-(4-bromothiophen-2-

yl)cyclopropanecarboxylic acid (1.10 g) and tert-butyl alcohol (10 mL) were added triethylamine (0.745 mL) and diphenylphosphoryl azide (1.15 mL) at room temperature, and the mixture was stirred at room temperature for 2 hr, and then
5 at 80°C overnight. The reaction mixture was cooled to room temperature, water was added, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was
10 purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (930 mg).
¹H NMR (300 MHz, DMSO-d₆) δ 1.04-1.23 (2H, m), 1.38 (9H, s), 2.07 (1H, ddd, J = 9.09, 6.06, 3.03 Hz), 2.61 (1H, brs), 6.80 (1H, d, J = 0.76 Hz), 7.22-7.34 (1H, m), 7.38 (1H, d, J = 1.51
15 Hz).

[0346]

E) methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate

A mixture of tert-butyl (trans-2-(4-bromothiophen-2-yl)cyclopropyl)carbamate (925 mg), triethylamine (0.810 mL),
20 dichloro(1,1'-bis(diphenylphosphino)ferrocene)palladium (425 mg) and methanol (100 mL) was stirred at 90°C for 8 hr under a carbon monoxide atmosphere (3 atm). The insoluble material was filtered off and washed with methanol, and the filtrate was
25 concentrated under reduced pressure. The residue was extracted with ethyl acetate and water, and the organic layer was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to
30 give the title compound (813 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.08-1.23 (2H, m), 1.39 (9H, s), 2.08 (1H, ddd, J = 9.18, 6.34, 3.03 Hz), 2.60 (1H, brs), 3.76 (3H, s), 7.08-7.15 (1H, m), 7.29 (1H, brs), 8.07 (1H, d, J = 1.51 Hz).

35 [0347]

F) methyl 5-(trans-2-aminocyclopropyl)thiophene-3-carboxylate hydrochloride

To a mixture of methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (450 mg) and ethyl acetate (5 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (3.78 mL) at 0°C, and the mixture was stirred at room temperature overnight. The reaction mixture was concentrated under reduced pressure, the residue was dissolved in methanol, and ethyl acetate was added. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (320 mg).
¹H NMR (300 MHz, DMSO-d₆) δ 1.29 (1H, dt, J = 7.76, 6.15 Hz), 1.47 (1H, ddd, J = 10.13, 5.96, 4.73 Hz), 2.56 (1H, ddd, J = 9.75, 6.15, 3.79 Hz), 2.81-2.94 (1H, m), 3.73-3.80 (3H, m), 7.23 (1H, d, J = 0.76 Hz), 8.12-8.18 (1H, m), 8.51 (3H, brs).
[0348]

G) methyl 5-(trans-2-((tert-butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiophene-3-carboxylate

A mixture of methyl 5-(trans-2-aminocyclopropyl)thiophene-3-carboxylate hydrochloride (50.0 mg), sodium hydrogen carbonate (44.9 mg), THF (2 mL) and methanol (2 mL) was stirred at room temperature for 30 min, and cyclopropanecarbaldehyde (18.0 mg) was added. Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 2 hr, and sodium borohydride (12.1 mg) was added at 0°C. Under a nitrogen atmosphere, the reaction mixture was stirred at room temperature for 1.5 hr, and to the reaction mixture were added ethyl acetate and water under ice-cooling. The mixture was extracted with ethyl acetate, and the extract was washed with water and saturated brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure, and the obtained residue was dissolved in THF (3 mL). Triethylamine (0.037 mL) and di-tert-butyl dicarbonate (0.077 mL) were added, and the reaction mixture was stirred at room

temperature overnight. Water was added, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The

5 residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (75.7 mg).
¹H NMR (300 MHz, DMSO-d₆) δ 0.09-0.30 (2H, m), 0.34-0.52 (2H, m), 0.91-1.01 (1H, m), 1.20-1.28 (1H, m), 1.28-1.43 (10H, m), 2.23-2.38 (1H, m), 2.70-2.81 (1H, m), 2.92 (1H, dd, J = 14.39, 7.19 Hz), 3.22 (1H, dd, J = 14.20, 7.00 Hz), 3.73-3.80 (3H, m), 7.19 (1H, dd, J = 1.51, 0.76 Hz), 8.08 (1H, d, J = 1.51 Hz).

[0349]

H) 5-(trans-2-((tert-butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiophene-3-carboxylic acid

Methyl 5-(trans-2-((tert-butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiophene-3-carboxylate (73.0 mg) was dissolved in THF (1.5 mL) and methanol (1.5 mL), 2 mol/L aqueous sodium hydroxide solution
20 (0.260 mL) was added at 0°C, and the mixture was stirred at room temperature overnight. The reaction mixture was acidified with 1 mol/L hydrochloric acid, and extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and
25 concentrated under reduced pressure to give the title compound (66.5 mg).

MS: [M+2H-Boc]⁺ 238.2.

[0350]

I) tert-butyl (cyclopropylmethyl)(trans-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate

5-(trans-2-((tert-butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiophene-3-carboxylic acid (33.0 mg), 5-methyl-1,3,4-thiadiazol-2-amine (12.4 mg) and triethylamine (0.027 mL) were dissolved in DMF
35 (2 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-

tetramethyluronium hexafluorophosphate (44.6 mg) was added, and the mixture was stirred at room temperature overnight. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed
5 with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (27.4 mg).

MS: [M+H]⁺ 435.3.

10 [0351]

J) 5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide dihydrochloride

To a mixture of tert-butyl (cyclopropylmethyl)(trans-2-
15 (4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate (27.0 mg) and ethyl acetate (2 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (0.310 mL) at 0°C, and the mixture was stirred at room temperature overnight. The reaction mixture was concentrated
20 under reduced pressure, and the residue was fractionated with LC/MS (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)). The obtained fraction was extracted with ethyl acetate and saturated aqueous sodium hydrogen carbonate solution. The organic layer was successively washed with water and saturated
25 brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was dissolved in ethyl acetate, 4 mol/L hydrogen chloride/ethyl acetate solution was added, and the mixture was stirred at room temperature for 2 hr. The precipitated solid was collected by filtration and
30 washed with ethyl acetate to give the title compound (22.9 mg).
¹H NMR (300 MHz, DMSO-d₆) δ 0.35-0.44 (2H, m), 0.53-0.64 (2H, m), 1.28-1.39 (1H, m), 1.63-1.75 (1H, m), 2.63 (3H, s), 2.80 (1H, ddd, J = 9.75, 6.15, 3.79 Hz), 2.90-2.99 (2H, m), 7.51 (1H, s), 8.45 (1H, d, J = 1.51 Hz), 9.57 (2H, brs), 10.08 (1H,
35 brs), 12.36-13.00 (1H, m).

[0352]

Example 24

N-(4,4-difluorocyclohexyl)-5-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide

5 hydrochloride

A) 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylic acid

Methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (450
10 mg) was dissolved in methanol (5 mL) and THF (5 mL), 2 mol/L aqueous sodium hydroxide solution (1.89 mL) was added at 0°C, and the mixture was stirred at room temperature overnight. The reaction mixture was acidified with 1 mol/L hydrochloric acid at 0°C and extracted with ethyl acetate. The extract was
15 washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (385 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.02-1.23 (2H, m), 1.39 (9H, s), 2.07 (1H, ddd, J = 9.09, 6.25, 3.22 Hz), 2.59 (1H, brs), 7.07
20 (1H, d, J = 0.76 Hz), 7.28 (1H, brs), 7.97 (1H, d, J = 1.51 Hz), 12.61 (1H, brs).

[0353]

B) N-(4,4-difluorocyclohexyl)-5-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide
25 hydrochloride

By a method similar to that of Example 1, steps G and H and Example 2, the title compound was obtained.

[0354]

Example 25

30 4-(trans-2-aminocyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide dihydrochloride

4-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylic acid (50 mg), 1-methyl-1H-pyrazol-4-amine hydrochloride (27 mg) and triethylamine (0.094
35 mL) were dissolved in DMF (1.0 mL), O-(7-azabenzotriazol-1-

yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (77 mg) was added at room temperature, and the mixture was stirred at room temperature overnight. The reaction mixture was poured into water, extracted with ethyl acetate, washed with water
5 and saturated brine, dried over anhydrous magnesium sulfate and concentrated under reduced pressure. To a mixture of the obtained residue and ethyl acetate (2.0 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (2.0 mL) at room temperature, and the mixture was stirred at room temperature
10 overnight. The resulting precipitate was collected by filtration to give the title compound (55 mg).

MS: [M-2HCl+H]⁺ 277.1.

[0355]

Example 26

15 4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide fumarate

To a mixture of 4-(trans-2-
((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(5-methyl-
20 1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride (15 mg) and ethyl acetate was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed with water and saturated brine, dried over anhydrous magnesium sulfate and
25 concentrated under reduced pressure. To a mixture of the obtained residue and ethyl acetate was added a solution of fumaric acid (4.13 mg) in ethanol at room temperature, and the reaction mixture was stirred for 30 min. The resulting precipitate was collected by filtration to give the title
30 compound (9.0 mg).

¹H NMR (300 MHz, METHANOL-d₄) δ 0.26-0.36 (2H, m), 0.58-0.69 (2H, m), 1.01-1.12 (1H, m), 1.13-1.22 (1H, m), 1.30-1.40 (1H, m), 2.10-2.20 (1H, m), 2.55 (3H, s), 2.67 (3H, s), 2.72-2.79 (1H, m), 2.83-2.90 (2H, m), 6.63-6.72 (1H, m), 7.52-7.65 (1H,
35 m).

[0356]

Example 27

4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

5 A) ethyl (E)-3-(5-bromothiophen-3-yl)acrylate

To a suspension of lithium chloride (174 mg) and acetonitrile (15 mL) were added 5-bromothiophene-3-carbaldehyde (770 mg) and ethyl diethylphosphonoacetate (0.840 mL) at 0°C, and the mixture was stirred at 0°C for 10 min. To
10 the reaction mixture was added 1,8-diazabicyclo[5.4.0]undec-7-ene (0.638 mL), and the mixture was stirred at room temperature overnight. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried
15 over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.01 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.19-1.29 (3H, m), 4.11-4.23 (2H, m), 6.48 (1H, d, J = 15.90 Hz), 7.53 (1H, d, J = 15.0 Hz),
20 7.74 (1H, d, J = 1.51 Hz), 7.98 (1H, d, J = 1.51 Hz).

[0357]

B) ethyl trans-2-(5-bromothiophen-3-yl)cyclopropanecarboxylate

To a suspension of trimethylsulfoxonium iodide (1.01 g)
25 in DMSO (10 mL) was added sodium hydride (60% in oil, 184 mg) under a nitrogen atmosphere, and the mixture was stirred at room temperature for 30 min. To the reaction mixture was added a mixture of ethyl (E)-3-(5-bromothiophen-3-yl)acrylate (1.00 g) and DMSO (5 mL), and the mixture was stirred at room
30 temperature overnight under a nitrogen atmosphere. Water was added to the reaction mixture at room temperature, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced
35 pressure. The residue was purified by silica gel column

chromatography (hexane/ethyl acetate) to give the title compound (263 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.16-1.24 (3H, m), 1.27-1.42 (2H, m), 1.92 (1H, ddd, J = 8.33, 5.30, 4.16 Hz), 2.44 (1H, ddd, J = 9.18, 6.72, 4.16 Hz), 4.03-4.14 (2H, m), 7.09 (1H, d, J = 1.89 Hz), 7.32 (1H, d, J = 1.51 Hz).

[0358]

C) trans-2-(5-bromothiophen-3-yl)cyclopropanecarboxylic acid

Ethyl trans-2-(5-bromothiophen-3-yl)cyclopropanecarboxylate (260 mg) was dissolved in THF (2 mL) and ethanol (2 mL), 2 mol/L aqueous sodium hydroxide solution (1.18 mL) was added at 0°C, and the mixture was stirred at room temperature overnight. The reaction mixture was acidified with 1 mol/L hydrochloric acid, and extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (215 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.22-1.40 (2H, m), 1.78 (1H, ddd, J = 8.33, 5.30, 4.16 Hz), 2.39 (1H, ddd, J = 9.09, 6.44, 4.16 Hz), 7.08 (1H, d, J = 1.51 Hz), 7.30 (1H, d, J = 1.51 Hz), 12.35 (1H, brs).

[0359]

D) tert-butyl (trans-2-(5-bromothiophen-3-yl)cyclopropyl)carbamate

To a mixture of trans-2-(5-bromothiophen-3-yl)cyclopropanecarboxylic acid (230 mg) and tert-butyl alcohol (3 mL) were added triethylamine (0.156 mL) and diphenylphosphoryl azide (0.241 mL) at room temperature, and the mixture was stirred at room temperature for 2 hr and then at 80°C overnight. The reaction mixture was cooled to room temperature, water was added, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was

purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (205 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.92-1.10 (2H, m), 1.34-1.41 (9H, m), 1.87 (1H, ddd, J = 9.28, 6.25, 3.41 Hz), 2.53-2.65 (1H, m),
5 7.00 (1H, d, J = 1.51 Hz), 7.12 (1H, d, J = 1.51 Hz), 7.20 (1H, brs).

[0360]

E) methyl 4-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-2-carboxylate

10 A mixture of tert-butyl (trans-2-(5-bromothiophen-3-yl)cyclopropyl)carbamate (330 mg), triethylamine (0.289 mL), dichloro(1,1'-bis(diphenylphosphino)ferrocene)palladium (152 mg) and methanol (30 mL) was stirred at 90°C for 8 hr under a carbon monoxide atmosphere (3 atm). The insoluble material was
15 filtered off, and the filtrate was concentrated under reduced pressure. The residue was extracted with ethyl acetate and water, and the organic layer was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel
20 column chromatography (hexane/ethyl acetate) to give the title compound (253 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.99-1.12 (2H, m), 1.38 (9H, s), 1.88-1.98 (1H, m), 2.60 (1H, brs), 3.80 (3H, s), 7.22 (1H, brs), 7.50 (1H, d, J = 1.51 Hz), 7.59 (1H, d, J = 1.51 Hz).

25 [0361]

F) 4-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-2-carboxylic acid

Methyl 4-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-2-carboxylate (250
30 mg) was dissolved in THF (4 mL) and methanol (4 mL), 2 mol/L aqueous sodium hydroxide solution (1.05 mL) was added at 0°C, and the mixture was stirred at room temperature overnight. The reaction mixture was acidified with 1 mol/L hydrochloric acid at 0°C, and the mixture was extracted with ethyl acetate. The
35 extract was successively washed with water and saturated brine,

dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (235 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.96-1.12 (2H, m), 1.38 (9H, s), 1.92 (1H, ddd, J = 9.18, 6.34, 3.03 Hz), 2.58 (1H, brs), 7.22 (1H, brs), 7.39-7.46 (1H, m), 7.50 (1H, d, J = 1.51 Hz), 12.99 (1H, brs).

[0362]

G) tert-butyl (trans-2-(5-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-3-yl)cyclopropyl)carbamate

10 4-(trans-2-((tert-Butoxycarbonyl)amino)cyclopropyl)thiophene-2-carboxylic acid (80.0 mg), 5-methyl-1,3,4-thiadiazol-2-amine (35.8 mg) and triethylamine (0.079 mL) were dissolved in DMF (3 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
15 hexafluorophosphate (129 mg) was added, and the mixture was stirred at room temperature overnight. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under
20 reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (58.2 mg).

MS: [M+H]⁺ 381.2.

[0363]

25 H) 4-(trans-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

To a mixture of tert-butyl (trans-2-(5-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-3-yl)cyclopropyl)carbamate (55.0 mg) and ethyl acetate (3 mL) was added 4 mol/L hydrogen
30 chloride/ethyl acetate solution (0.361 mL) at 0°C, and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (26.3 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.11-1.23 (1H, m), 1.44 (1H, ddd, J = 10.03, 5.87, 4.54 Hz), 2.43 (1H, ddd, J = 9.75, 6.15, 3.41

Hz), 2.62 (3H, s), 2.81 (1H, dd, $J = 7.76, 3.98$ Hz), 7.73 (1H, s), 7.92-8.19 (1H, m), 8.54 (3H, brs).

[0364]

I) 4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

To a mixture of 4-(trans-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride (24.0 mg), sodium hydrogen carbonate (14.3 mg), THF (1.5 mL) and methanol (1.5 mL) was added cyclopropanecarbaldehyde (5.71 mg) at room temperature. Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, and sodium borohydride (3.86 mg) was added at 0°C. The reaction mixture was stirred at room temperature for 1 hr, and ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture. The mixture was extracted with ethyl acetate, and the extract was successively washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (ethyl acetate/methanol). To the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution and the reaction mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (17.6 mg).

^1H NMR (300 MHz, DMSO- d_6) δ 0.30-0.44 (2H, m), 0.50-0.68 (2H, m), 1.05-1.17 (1H, m), 1.19-1.32 (1H, m), 1.52-1.66 (1H, m), 2.55-2.67 (4H, m), 2.86-3.00 (3H, m), 7.75 (1H, s), 8.07 (1H, brs), 9.37-9.60 (2H, m).

[0365]

Example 28

4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-2-carboxamide hydrochloride

A) tert-butyl (trans-2-(5-((4,4-

difluorocyclohexyl) carbamoyl) thiophen-3-yl)cyclopropyl) carbamate

4-(trans-2-((tert-

Butoxycarbonyl) amino) cyclopropyl) thiophene-2-carboxylic acid

5 (150 mg), 4,4-difluorocyclohexanamine hydrochloride (100 mg) and triethylamine (0.295 mL) were dissolved in DMF (6 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (242 mg) was added at room temperature, and the mixture was stirred overnight. To the reaction mixture
10 were added ethyl acetate and water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to
15 give the title compound (211 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.92-1.12 (2H, m), 1.33-1.42 (9H, m), 1.48-1.69 (2H, m), 1.79-2.14 (7H, m), 2.62 (1H, brs), 3.92 (1H, d, J = 7.57 Hz), 7.22 (1H, brs), 7.31 (1H, s), 7.55 (1H, d, J = 1.14 Hz), 8.20 (1H, d, J = 7.95 Hz).

20 [0366]

B) 4-(trans-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-2-carboxamide hydrochloride

To a mixture of tert-butyl (trans-2-(5-((4,4-difluorocyclohexyl) carbamoyl) thiophen-3-

25 yl)cyclopropyl) carbamate (210 mg) and ethyl acetate (3 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (1.31 mL) at 0°C, and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration, washed with ethyl acetate, and dried under reduced pressure to
30 give the title compound (108 mg).

MS: [M-HCl+H]⁺ 301.2.

[0367]

C) 4-(trans-2-((cyclopropylmethyl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-2-carboxamide hydrochloride

35 To a mixture of 4-(trans-2-aminocyclopropyl)-N-(4,4-

difluorocyclohexyl)thiophene-2-carboxamide hydrochloride (30.0 mg), sodium hydrogen carbonate (18.7 mg), THF (1.5 mL) and methanol (1.5 mL) was added cyclopropanecarbaldehyde (7.49 mg) at room temperature. Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, and sodium borohydride (5.05 mg) was added at 0°C. The reaction mixture was stirred at room temperature for 1 hr, and ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture. The mixture was extracted with ethyl acetate, and the extract was successively washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and to the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The reaction mixture was stirred at room temperature overnight, and the precipitated solid was collected by filtration, washed with ethyl acetate and dried under reduced pressure to give the title compound (18.1 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.36 (2H, brs), 0.59 (2H, brs), 0.97-1.34 (2H, m), 1.45-1.68 (3H, m), 1.80-2.13 (6H, m), 2.94 (3H, d, J = 7.57 Hz), 3.92 (2H, brs), 7.44-7.75 (2H, m), 8.23 (1H, d, J = 8.33 Hz), 9.19 (1H, brs).

[0368]

Example 29

N-(4,4-difluorocyclohexyl)-4-(trans-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-2-carboxamide hydrochloride

To a mixture of 4-(trans-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-2-carboxamide hydrochloride (30.0 mg), sodium hydrogen carbonate (18.7 mg), THF (1.5 mL) and methanol (1.5 mL) was added dihydro-2H-pyran-4(3H)-one (10.7 mg). Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, and sodium borohydride (5.05 mg) was added at 0°C. The reaction mixture was stirred at room temperature for 1 hr, and

ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture. The mixture was extracted with ethyl acetate, and the extract was successively washed with water and saturated brine, and dried
5 over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (methanol/ethyl acetate), and to the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The reaction mixture was stirred at room
10 temperature overnight, and the precipitated solid was collected by filtration, and washed with ethyl acetate to give the title compound (14.8 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.11-1.33 (1H, m), 1.42-2.20 (15H, m), 2.93 (1H, brs), 3.38-3.55 (2H, m), 3.92 (3H, d, J = 10.60
15 Hz), 7.43-7.71 (2H, m), 8.23 (1H, d, J = 6.82 Hz), 9.43 (1H, brs).

[0369]

Example 30

5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-
20 difluorocyclohexyl)thiophene-3-carboxamide hydrochloride
A) tert-butyl (trans-2-(4-((4,4-
difluorocyclohexyl)carbamoyl)thiophen-2-
yl)cyclopropyl)carbamate

5-(trans-2-((tert-
25 Butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylic acid
(100 mg), 4,4-difluorocyclohexanamine hydrochloride (66.6 mg)
and triethylamine (0.197 mL) were dissolved in DMF (4 mL), O-
(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
hexafluorophosphate (161 mg) was added, and the mixture was
30 stirred at room temperature overnight. To the reaction mixture
were added ethyl acetate and water, and the mixture was
extracted with ethyl acetate. The extract was washed with
saturated brine, dried over anhydrous sodium sulfate, and
concentrated under reduced pressure. The residue was purified
35 by silica gel column chromatography (hexane/ethyl acetate) to

give the title compound (92.1 mg).

MS: $[M+H]^+$ 401.2.

[0370]

B) 5-(trans-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide hydrochloride

To a mixture of tert-butyl (trans-2-(4-((4,4-difluorocyclohexyl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate (90.0 mg) and ethyl acetate (3 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (0.562 mL) at 0°C, and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (76.8 mg).

MS: $[M-HCl+H]^+$ 301.2.

[0371]

C) 5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide hydrochloride

To a mixture of 5-(trans-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide hydrochloride (30.0 mg), sodium hydrogen carbonate (18.7 mg), THF (1.5 mL) and methanol (1.5 mL) was added cyclopropanecarbaldehyde (7.49 mg) at room temperature. Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, and sodium borohydride (5.05 mg) was added at 0°C. Under a nitrogen atmosphere, the reaction mixture was stirred at room temperature for 1 hr, and ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture. The mixture was extracted with ethyl acetate, and the extract was successively washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and to the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The reaction mixture was stirred at room temperature overnight, and the

precipitated solid was collected by filtration, and washed with ethyl acetate to give the title compound (14.6 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.38 (2H, m, J = 3.41 Hz), 0.58 (2H, d, J = 6.44 Hz), 1.07 (1H, brs), 1.20-1.38 (1H, m), 1.45-1.68 (3H, m), 1.86 (3H, m), 2.03 (3H, m), 2.69 (2H, m), 2.95 (2H, d, J = 7.19 Hz), 3.93 (1H, brs), 7.28 (1H, s), 7.95 (1H, s), 8.05 (1H, d, J = 7.95 Hz), 9.26 (2H, brs).

[0372]

Example 31

10 N-(4,4-difluorocyclohexyl)-5-(trans-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide hydrochloride

To a mixture of 5-(trans-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide hydrochloride (30.0 mg), sodium hydrogen carbonate (18.7 mg), THF (1.5 mL) and 15 methanol (1.5 mL) was added dihydro-2H-pyran-4(3H)-one (10.7 mg) at room temperature. Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, and sodium borohydride (5.05 mg) was added at 0°C. The reaction mixture was stirred at room 20 temperature for 1 hr, and ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture. The mixture was extracted with ethyl acetate, and the extract was successively washed with water and saturated brine, and dried over anhydrous sodium sulfate. The 25 solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (methanol/ethyl acetate), and to the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The reaction mixture was stirred at room temperature overnight, and the 30 precipitated solid was collected by filtration, and washed with ethyl acetate to give the title compound (18.1 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.25-1.41 (1H, m), 1.62 (5H, d, J = 10.60 Hz), 1.77-2.15 (9H, m), 2.62-2.80 (1H, m), 2.89-3.08 (1H, m), 3.39-3.59 (1H, m), 3.92 (3H, d, J = 9.47 Hz), 7.28 (1H, s), 35 7.95 (1H, s), 8.04 (1H, d, J = 7.95 Hz), 9.44 (2H, brs).

[0373]

Example 32

5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide

5 hydrochloride

A) tert-butyl (2-(4-((tetrahydro-2H-pyran-4-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate

5-(trans-2-((tert-

Butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylic acid

10 (50.0 mg), tetrahydro-2H-pyran-4-amine (19.6 mg) and triethylamine (0.049 mL) were dissolved in DMF (2 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (81.0 mg) was added at room temperature, and the mixture was stirred overnight. To the reaction mixture
15 were added ethyl acetate and water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to
20 give the title compound (61.8 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.99-1.11 (1H, m), 1.12-1.21 (1H, m), 1.39 (9H, s), 1.44-1.62 (2H, m), 1.72 (2H, dd, J = 12.49, 2.27 Hz), 2.06 (1H, ddd, J = 8.99, 5.77, 3.41 Hz), 2.62 (1H, brs), 3.33-3.42 (2H, m), 3.80-4.00 (3H, m), 7.17 (1H, s), 7.28
25 (1H, brs), 7.83 (1H, d, J = 1.51 Hz), 8.00 (1H, d, J = 7.95 Hz).

[0374]

B) 5-(trans-2-aminocyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride

30 To a mixture of tert-butyl (2-(4-((tetrahydro-2H-pyran-4-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate (59.0 mg) and ethyl acetate (3 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (0.402 mL) at 0°C, and the mixture was stirred at room temperature overnight. The precipitated solid was
35 collected by filtration and washed with ethyl acetate to give

the title compound (45.5 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.17-1.31 (1H, m), 1.40-1.62 (3H, m), 1.64-1.79 (2H, m), 2.55 (1H, td, J = 6.44, 3.41 Hz), 2.75-2.88 (1H, m), 3.29-3.45 (2H, m), 3.79-4.01 (3H, m), 7.24-7.32
5 (1H, m), 7.94 (1H, d, J = 1.51 Hz), 8.08 (1H, d, J = 7.95 Hz), 8.55 (3H, d, J = 2.65 Hz).

[0375]

C) 5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide

10 hydrochloride

To a mixture of 5-(trans-2-aminocyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride (43.0 mg), sodium hydrogen carbonate (29.8 mg), THF (2 mL) and methanol (2 mL) was added
15 cyclopropanecarbaldehyde (11.9 mg). Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, and sodium borohydride (8.06 mg) was added at 0°C. Under a nitrogen atmosphere, the reaction mixture was stirred at room temperature for 1 hr, and
20 ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture. The mixture was extracted with ethyl acetate, and the extract was washed successively with water and saturated brine. The extract was dried over anhydrous sodium sulfate, and the
25 solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and to the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution under ice-cooling. The reaction mixture was stirred at room temperature for 2 hr.
30 The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (29.8 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.31-0.44 (2H, m), 0.51-0.66 (2H, m), 0.98-1.15 (1H, m), 1.21-1.38 (1H, m), 1.42-1.66 (3H, m), 1.66-1.81 (2H, m), 2.70 (1H, ddd, J = 9.66, 6.25, 3.41 Hz),
35 2.91-3.01 (3H, m), 3.33-3.45 (2H, m), 3.78-4.04 (3H, m), 7.28

(1H, s), 7.95 (1H, d, J = 1.51 Hz), 8.06 (1H, d, J = 7.57 Hz), 9.30 (2H, brs).

[0376]

Example 33

5 5-(trans-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide dihydrochloride

A) methyl 5-(trans-2-((tert-butoxycarbonyl)(cyclobutyl)amino)cyclopropyl)thiophene-3-carboxylate

10 To a mixture of methyl 5-(trans-2-aminocyclopropyl)thiophene-3-carboxylate hydrochloride (100 mg), sodium hydrogen carbonate (90.0 mg), THF (4 mL) and methanol (4 mL) was added cyclobutanone (36.0 mg). Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C
15 for 1.5 hr and at room temperature for 30 min, and sodium borohydride (24.3 mg) was added at 0°C. The reaction mixture was stirred at room temperature for 1 hr, and ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture. The mixture was extracted
20 with ethyl acetate, and the extract was successively washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The obtained residue was dissolved in THF (3 mL), and triethylamine (0.090 mL) and di-tert-butyl dicarbonate
25 (0.120 mL) were added. The reaction mixture was stirred at room temperature overnight, and the mixture was extracted with ethyl acetate and water. The organic layer was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue
30 was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (98.9 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.23-1.33 (2H, m), 1.37 (9H, s), 1.45-1.68 (2H, m), 2.01-2.32 (5H, m), 2.62 (1H, ddd, J = 7.10, 5.02, 3.41 Hz), 3.77 (3H, s), 3.92-4.09 (1H, m), 7.14-7.21 (1H,
35 m), 8.08 (1H, d, J = 1.51 Hz).

[0377]

B) 5-(trans-2-((tert-butoxycarbonyl)(cyclobutyl)amino)cyclopropyl)thiophene-3-carboxylic acid

5 Methyl 5-(trans-2-((tert-butoxycarbonyl)(cyclobutyl)amino)cyclopropyl)thiophene-3-carboxylate (95.0 mg) was dissolved in THF (2 mL) and methanol (2 mL), 2 mol/L aqueous sodium hydroxide solution (0.338 mL) was added at 0°C, and the mixture was stirred at room
10 temperature overnight. The reaction mixture was acidified with 1 mol/L hydrochloric acid at 0°C, and extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound
15 (90.1 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.22-1.32 (2H, m), 1.38 (9H, s), 1.48-1.67 (2H, m), 2.04-2.30 (5H, m), 2.61 (1H, ddd, J = 7.29, 4.45, 3.41 Hz), 3.94-4.07 (1H, m), 7.11-7.15 (1H, m), 7.99 (1H, d, J = 1.51 Hz), 12.60 (1H, brs).

20 [0378]

C) tert-butyl cyclobutyl(trans-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate

5-(trans-2-((tert-butoxycarbonyl)(cyclobutyl)amino)cyclopropyl)thiophene-3-
25 carboxylic acid (45.0 mg), 5-methyl-1,3,4-thiadiazol-2-amine (18.4 mg) and triethylamine (0.028 mL) were dissolved in DMF (3 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (60.8 mg) was added, and the mixture was stirred at room temperature overnight. The
30 reaction mixture was extracted with ethyl acetate and water, and the organic layer was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title
35 compound (38.3 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.14-1.43 (11H, m), 1.48-1.69 (2H, m), 2.04-2.34 (5H, m), 2.58-2.67 (4H, m), 4.03 (1H, quin, J = 8.52 Hz), 7.47 (1H, s), 8.36 (1H, d, J = 1.14 Hz), 12.64 (1H, brs).

5 [0379]

D) 5-(trans-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide dihydrochloride

To a mixture of tert-butyl cyclobutyl(trans-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate (36.0 mg) and ethyl acetate (2 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (0.207 mL) at 0°C, and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title
10 compound (26.5 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.17 (1H, t, J = 7.00 Hz), 1.26-1.42 (1H, m), 1.56-1.68 (1H, m), 1.74-1.92 (2H, m), 2.29 (4H, d, J = 13.63), 2.69-3.11 (3H, m), 3.60-4.26 (2H, m), 7.42 (1H, s), 8.45 (1H, brs), 9.67-10.18 (1H, m), 13.03 (1H, brs).

20 [0380]

Example 34

5-(trans-2-(cyclobutylamino)cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-3-carboxamide hydrochloride

A) tert-butyl cyclobutyl(trans-2-(4-((1-methyl-1H-pyrazol-4-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate
25

5-(trans-2-((tert-butoxycarbonyl)(cyclobutyl)amino)cyclopropyl)thiophene-3-carboxylic acid (45.0 mg), 1-methyl-1H-pyrazol-4-amine (15.5 mg) and triethylamine (0.028 mL) were dissolved in DMF (3 mL),
30 O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (60.8 mg) was added, and the mixture was stirred at room temperature overnight. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine,
35 dried over anhydrous sodium sulfate, and concentrated under

reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (24.0 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.20-1.35 (2H, m), 1.36-1.42 (9H, m), 1.47-1.70 (2H, m), 2.01-2.34 (5H, m), 2.57-2.67 (1H, m), 3.76-3.85 (3H, m), 3.93-4.15 (1H, m), 7.30 (1H, d, J = 0.76 Hz), 7.50 (1H, s), 7.93 (2H, s), 10.11 (1H, s).

[0381]

B) 5-(trans-2-(cyclobutylamino)cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-3-carboxamide hydrochloride

To a mixture of tert-butyl cyclobutyl(trans-2-(4-((1-methyl-1H-pyrazol-4-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate (22.0 mg) and ethyl acetate (1 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (0.132 mL) at 0°C, and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (11.2 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.24-1.38 (1H, m), 1.51-1.66 (1H, m), 1.72-1.90 (2H, m), 2.12-2.40 (4H, m), 2.74 (1H, ddd, J = 9.94, 6.15, 3.22 Hz), 2.82-2.96 (1H, m), 3.75-3.95 (4H, m), 7.29-8.12 (4H, m), 9.76 (2H, brs), 10.26 (1H, s).

[0382]

Example 35

4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiazole-2-carboxamide hydrochloride

A) ethyl (E)-3-(2-bromothiazol-4-yl)acrylate

To a suspension of lithium chloride (450 mg) in acetonitrile (30 mL) were added 2-bromothiazole-4-carbaldehyde (2.00 g) and ethyl diethylphosphonoacetate (2.17 mL) at 0°C, and the mixture was stirred under a nitrogen atmosphere at 0°C for 5 min. To the reaction mixture was added 1,8-diazabicyclo[5.4.0]undec-7-ene (1.65 mL) at 0°C, and the mixture was stirred under a nitrogen atmosphere at room temperature overnight. Water was added to the reaction mixture,

and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column

5 chromatography (hexane/ethyl acetate) to give the title compound (2.34 g).

^1H NMR (300 MHz, DMSO-d_6) δ 1.25 (3H, t, $J = 7.00$ Hz), 4.19 (2H, q, $J = 6.94$ Hz), 6.54 (1H, d, $J = 15.52$ Hz), 7.60 (1H, d, $J = 15.52$ Hz), 8.14 (1H, s).

10 [0383]

B) ethyl trans-2-(2-bromothiazol-4-yl)cyclopropanecarboxylate

To a suspension of trimethylsulfoxonium iodide (2.35 g) in DMSO (20 mL) was added sodium hydride (50% in oil, 512 mg) under a nitrogen atmosphere, and the mixture was stirred at
15 room temperature for 30 min. To the reaction mixture was added a mixture of ethyl (E)-3-(2-bromothiazol-4-yl)acrylate (2.33 g) and DMSO (10 mL), and the mixture was stirred at room temperature overnight. Water was added to the reaction mixture at 0°C , and the mixture was extracted with ethyl acetate. The
20 extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (912 mg).

25 ^1H NMR (300 MHz, DMSO-d_6) δ 1.16-1.23 (3H, m), 1.39-1.48 (2H, m), 1.93-2.03 (1H, m), 2.56-2.65 (1H, m), 4.09 (2H, q, $J = 6.94$ Hz), 7.56 (1H, s).

[0384]

C) trans-2-(2-bromothiazol-4-yl)cyclopropanecarboxylic acid

30 Ethyl trans-2-(2-bromothiazol-4-yl)cyclopropanecarboxylate (905 mg) was dissolved in THF (5 mL) and ethanol (5 mL), 2 mol/L aqueous sodium hydroxide solution (4.10 mL) was added at 0°C , and the mixture was stirred at room temperature overnight. The reaction mixture
35 was acidified with 1 mol/L hydrochloric acid, and extracted

with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (809 mg).

5 ¹H NMR (300 MHz, DMSO-d₆) δ 1.32-1.45 (2H, m), 1.88 (1H, ddd, J = 8.24, 5.40, 4.16 Hz), 2.53-2.60 (1H, m), 7.54 (1H, s), 12.42 (1H, brs).

[0385]

D) tert-butyl (trans-2-(2-bromothiazol-4-yl)cyclopropyl)carbamate

To a mixture of trans-2-(2-bromothiazol-4-yl)cyclopropanecarboxylic acid (805 mg) and tert-butyl alcohol (10 mL) were added triethylamine (0.543 mL) and diphenylphosphoryl azide (0.838 mL) at room temperature, and
15 the mixture was stirred at room temperature for 2 hr and then at 80°C overnight. The reaction mixture was cooled to room temperature, water was added, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate,
20 and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (514 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.05-1.14 (2H, m), 1.37 (9H, s), 2.06 (1H, td, J = 7.57, 3.03 Hz), 2.79 (1H, brs), 7.24 (1H, brs), 7.36 (1H, s).

[0386]

E) trans-2-(2-bromothiazol-4-yl)cyclopropanamine hydrochloride

To a mixture of tert-butyl (trans-2-(2-bromothiazol-4-yl)cyclopropyl)carbamate (200 mg) and ethyl acetate (3 mL) was
30 added 4 mol/L hydrogen chloride/ethyl acetate solution (1.57 mL) at 0°C, and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (158 mg).

35 ¹H NMR (300 MHz, DMSO-d₆) δ 1.24 (1H, dt, J = 7.95, 6.06 Hz),

1.36-1.46 (1H, m), 2.56 (1H, ddd, J = 9.75, 6.15, 3.41 Hz),
2.81-2.95 (1H, m), 7.56 (1H, s), 8.54 (3H, brs).

[0387]

F) trans-2-(2-bromothiazol-4-yl)-N-

5 (cyclopropylmethyl)cyclopropanamine

A mixture of trans-2-(2-bromothiazol-4-yl)cyclopropanamine hydrochloride (100 mg), sodium hydrogen carbonate (131 mg), THF (3 mL) and methanol (3 mL) was stirred at room temperature for 30 min, and cyclopropanecarbaldehyde
10 (32.9 mg) was added. Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 2 hr, and sodium borohydride (22.2 mg) was added at 0°C. Under a nitrogen atmosphere, the reaction mixture was stirred at room temperature for 1.5 hr, and ethyl acetate and water were added to the reaction mixture
15 under ice-cooling. The mixture was extracted with ethyl acetate, and the extract was successively washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (NH,
20 hexane/ethyl acetate) to give the title compound (84.4 mg).
¹H NMR (300 MHz, DMSO-d₆) δ 0.02-0.09 (2H, m), 0.33-0.41 (2H, m), 0.76-0.89 (1H, m), 0.90-1.03 (2H, m), 1.87-2.02 (1H, m), 2.36-2.47 (4H, m), 7.27 (1H, s).

[0388]

25 G) tert-butyl (trans-2-(2-bromothiazol-4-

yl)cyclopropyl)(cyclopropylmethyl)carbamate

trans-2-(2-Bromothiazol-4-yl)-N-(cyclopropylmethyl)cyclopropanamine (81.0 mg) was dissolved in THF (4 mL), and triethylamine (0.062 mL) and di-tert-butyl
30 dicarbonate (0.103 mL) were added. The reaction mixture was stirred at room temperature overnight, and extracted with ethyl acetate and water. The organic layer was washed with water and saturated brine, dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The
35 residue was purified by silica gel column chromatography.

(hexane/ethyl acetate) to give the title compound (106 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.06-0.28 (2H, m), 0.33-0.53 (2H, m), 0.87-1.03 (1H, m), 1.24 (2H, dd, J = 7.57, 6.44 Hz), 1.35 (9H, s), 2.18-2.33 (1H, m), 2.79-2.99 (2H, m), 3.21 (1H, dd, J = 14.39, 6.82 Hz), 7.44 (1H, s).

[0389]

H) methyl 4-(trans-2-((tert-butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiazole-2-carboxylate

10 A mixture of tert-butyl (trans-2-(2-bromothiazol-4-yl)cyclopropyl)(cyclopropylmethyl)carbamate (80.0 mg), triethylamine (0.060 mL), dichloro(1,1'-bis(diphenylphosphino)ferrocene)palladium (31.4 mg) and methanol (15 mL) was stirred at 90°C for 8 hr under a carbon
15 monoxide atmosphere (3 atm). After cooling to room temperature, the insoluble material was filtered off, and the filtrate was concentrated under reduced pressure. The residue was extracted with ethyl acetate and water, and the organic layer was washed with saturated brine, dried over anhydrous sodium sulfate, and
20 concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (76.0 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.05-0.28 (2H, m), 0.32-0.53 (2H, m), 0.87-1.05 (1H, m), 1.27-1.37 (11H, m), 2.36 (1H, td, J = 7.86, 3.22 Hz), 2.87-3.03 (2H, m), 3.22 (1H, dd, J = 14.39, 6.82 Hz), 3.82-3.93 (3H, m), 7.85 (1H, s).

[0390]

I) 4-(trans-2-((tert-butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiazole-2-carboxylic acid
30

Methyl 4-(trans-2-((tert-butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiazole-2-carboxylate (93.0 mg) was dissolved in methanol (5 mL), 2 mol/L aqueous sodium hydroxide solution (0.330 mL) was added
35 at 0°C, and the mixture was stirred at room temperature for 1

hr. The reaction mixture was acidified with 1 mol/L hydrochloric acid at 0°C, and extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (78.9 mg).
MS: [M+H]⁺ 339.2.

[0391]

J) tert-butyl (cyclopropylmethyl)(trans-2-(2-((4,4-difluorocyclohexyl)carbamoyl)thiazol-4-yl)cyclopropyl)carbamate
4-(trans-2-((tert-butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiazole-2-carboxylic acid (38.0 mg), 4,4-difluorocyclohexanamine hydrochloride (21.2 mg) and triethylamine (0.063 mL) were dissolved in DMF (3 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (51.2 mg) was added, and the mixture was stirred at room temperature overnight. To the reaction mixture were added ethyl acetate and water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (17.6 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.07-0.28 (2H, m), 0.32-0.54 (2H, m), 0.90-1.04 (1H, m), 1.23-1.42 (11H, m), 1.64-1.95 (5H, m), 1.99-2.12 (3H, m), 2.29 (1H, ddd, J = 9.47, 6.25, 3.22 Hz), 2.94-3.07 (2H, m), 3.10-3.25 (1H, m), 3.96 (1H, d, J = 7.95 Hz), 7.66 (1H, s), 8.54 (1H, d, J = 8.71 Hz).

[0392]

K) 4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiazole-2-carboxamide hydrochloride

To a mixture of tert-butyl (cyclopropylmethyl)(trans-2-(2-((4,4-difluorocyclohexyl)carbamoyl)thiazol-4-yl)cyclopropyl)carbamate (16.0 mg) and ethyl acetate (1.5 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution.

(0.088 mL) at 0°C, and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (10.1 mg).

5 ¹H NMR (300 MHz, DMSO-d₆) δ 0.36 (2H, d, J = 4.16 Hz), 0.57 (2H, d, J = 6.44 Hz), 1.00-1.18 (1H, m), 1.33-1.49 (1H, m), 1.55-1.67 (1H, m), 1.68-1.87 (4H, m), 2.05 (3H, brs), 2.76-2.85 (1H, m), 2.94 (2H, d, J = 6.44 Hz), 3.13 (1H, brs), 3.97 (2H, brs), 7.78 (1H, s), 8.59 (1H, d, J = 8.71 Hz), 9.33-9.67 (2H, m).

10 [0393]

Example 36

5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-2-carboxamide hydrochloride

A) trans-2-(5-bromothiophen-2-yl)cyclopropanecarboxylic acid

15 To a mixture of lithium hydroxide monohydrate (12.7 g), water (100 mL) and methanol (100 mL) was added ethyl trans-2-(5-bromothiophen-2-yl)cyclopropanecarboxylate (16.7 g) at room temperature. After stirring at room temperature for 24 hr, methanol was evaporated under reduced pressure. The aqueous
20 layer was washed with dichloromethane, acidified with 2 mol/L hydrochloric acid, and extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (14.7 g).

25 ¹H NMR (400 MHz, CDCl₃) δ 1.35-1.40 (1H, m), 1.65-1.69 (1H, m), 1.90-1.94 (1H, m), 2.67-2.72 (1H, m), 6.61 (1H, d, J = 3.6 Hz), 6.87 (1H, d, J = 3.6 Hz).

[0394]

B) tert-butyl (2-(5-bromothiophen-2-yl)cyclopropyl)carbamate

30 trans-2-(5-Bromothiophen-2-yl)cyclopropanecarboxylic acid (14.7 g) was dissolved in acetone (150 mL), triethylamine (9.03 g) was added, and then ethyl chloroformate (9.68 g) was added at -20°C. After stirring at -10°C to -20°C for 2 hr, a mixture of sodium azide (6.57 g) and water (20 mL) was added,
35 and the mixture was stirred at -10°C to -20°C for 3 hr.

Acetone was evaporated by concentration under reduced pressure, and water (150 mL) and toluene (150 mL) were added. The aqueous layer was extracted twice with toluene (100 mL). The combined extract was dried over anhydrous sodium sulfate and concentrated to 150 mL under reduced pressure. To the obtained mixture was added tert-butyl alcohol (25 mL), and the mixture was heated under reflux for 18 hr. The mixture was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (8.66 g).

^1H NMR (400 MHz, CDCl_3) δ 1.15 (2H, t, $J = 6.8$ Hz), 1.46 (9H, s), 2.08-2.13 (1H, m), 2.67-2.74 (1H, m), 4.81 (1H, s), 6.57 (1H, d, $J = 3.2$ Hz), 6.82 (1H, d, $J = 3.2$ Hz).

[0395]

C) trans-2-(5-bromothiophen-2-yl)cyclopropanamine hydrochloride

To a mixture of tert-butyl (2-(5-bromothiophen-2-yl)cyclopropyl)carbamate (200 mg) and ethyl acetate (5 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (1.57 mL) at 0°C , and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (135 mg).

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 1.23 (1H, dt, $J = 7.95, 6.25$ Hz), 1.36-1.51 (1H, m), 2.47-2.54 (1H, m), 2.73-2.85 (1H, m), 6.75 (1H, dd, $J = 3.79, 1.14$ Hz), 6.99-7.10 (1H, m), 8.50 (2H, brs).

[0396]

D) tert-butyl (2-trans-(5-bromothiophen-2-yl)cyclopropyl)(cyclopropylmethyl)carbamate

To a mixture of trans-2-(5-bromothiophen-2-yl)cyclopropanamine hydrochloride (130 mg), sodium hydrogen carbonate (107 mg), THF (4 mL) and methanol (4 mL) was added cyclopropanecarbaldehyde (43.0 mg) at room temperature. Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and then at room temperature for 30 min, and sodium

borohydride (29.0 mg) was added at 0°C. The reaction mixture was stirred at room temperature for 1 hr, and ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture. The mixture was extracted
5 with ethyl acetate, and the extract was successively washed with water and saturated brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The obtained residue was dissolved in THF (5 mL), triethylamine (0.094 mL) and di-tert-butyl dicarbonate (0.156
10 mL) were added, and the mixture was stirred at room temperature overnight. To the reaction mixture were added ethyl acetate and water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under
15 reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (160 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.08-0.27 (2H, m), 0.35-0.52 (2H, m), 0.88-1.06 (1H, m), 1.14-1.24 (1H, m), 1.25-1.34 (1H, m),
20 1.39 (9H, s), 2.20-2.32 (1H, m), 2.65-2.77 (1H, m), 2.93 (1H, dd, J = 14.39, 6.82 Hz), 3.18 (1H, dd, J = 14.39, 6.82 Hz), 6.71 (1H, dd, J = 3.79, 0.76 Hz), 7.02 (1H, d, J = 3.79 Hz).
[0397]

E) methyl 5-(trans-2-((tert-butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiophene-
25 2-carboxylate

A mixture of tert-butyl (2-trans-(5-bromothiophen-2-yl)cyclopropyl)(cyclopropylmethyl)carbamate (155 mg), triethylamine (0.116 mL), dichloro(1,1'-
30 bis(diphenylphosphino)ferrocene)palladium (60.9 mg) and methanol (15 mL) was stirred at 90°C for 8 hr under a carbon monoxide atmosphere (3 atm). After cooling to room temperature, the insoluble material was filtered off, and the filtrate was concentrated under reduced pressure. The residue was extracted
35 with ethyl acetate and water, and the organic layer was washed

with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (138 mg).

5 ¹H NMR (300 MHz, DMSO-d₆) δ 0.06-0.30 (2H, m), 0.33-0.52 (2H, m), 0.84-1.05 (1H, m), 1.23-1.33 (1H, m), 1.35-1.49 (10H, m), 2.39 (1H, ddd, J = 9.47, 6.25, 3.22 Hz), 2.76-2.87 (1H, m), 2.88-3.02 (1H, m), 3.20 (1H, dd, J = 14.20, 7.00 Hz), 3.73-3.82 (3H, m), 6.96 (1H, d, J = 3.79 Hz), 7.63 (1H, d, J = 3.79
10 Hz).

[0398]

F) 5-(trans-2-((tert-butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiophene-2-carboxylic acid

15 Methyl 5-(trans-2-((tert-butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiophene-2-carboxylate (130 mg) was dissolved in methanol (5 mL), 2 mol/L aqueous sodium hydroxide solution (0.462 mL) was added at 0°C, and the mixture was stirred at room temperature
20 overnight. The reaction mixture was acidified with 1 mol/L hydrochloric acid at 0°C, and extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (125 mg).

25 MS: [M+2H-(Boc)]⁺ 238.2.

[0399]

G) tert-butyl (cyclopropylmethyl)(trans-2-(5-((4,4-difluorocyclohexyl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate

30 5-(trans-2-((tert-butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiophene-2-carboxylic acid (35.0 mg), 4,4-difluorocyclohexanamine hydrochloride (19.6 mg) and triethylamine (0.058 mL) were dissolved in DMF (3 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-
35 tetramethyluronium hexafluorophosphate (47.3 mg) was added,

and the mixture was stirred at room temperature overnight. To the reaction mixture were added ethyl acetate and water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, dried over anhydrous sodium

5 sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (42.3 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.08-0.28 (2H, m), 0.34-0.52 (2H, m), 0.88-1.04 (1H, m), 1.18-1.27 (1H, m), 1.32-1.42 (10H, m),
10 1.50-1.69 (2H, m), 1.77-1.96 (3H, m), 1.97-2.13 (3H, m), 2.32 (1H, m), 2.70-2.80 (1H, m), 2.89-3.04 (1H, m), 3.10-3.26 (1H, m), 3.91 (1H, d, J = 7.57 Hz), 6.86 (1H, d, J = 3.79 Hz), 7.58 (1H, d, J = 3.79 Hz), 8.16 (1H, d, J = 7.57 Hz).

[0400]

15 H) 5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-2-carboxamide hydrochloride

tert-Butyl (cyclopropylmethyl)(trans-2-(5-((4,4-difluorocyclohexyl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate (40.0 mg) was dissolved in ethyl
20 acetate (3 mL), 4 mol/L hydrogen chloride/ethyl acetate solution (0.220 mL) was added at 0°C, and the mixture was stirred at room temperature for 4 hr. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (22.1 mg).

25 ¹H NMR (300 MHz, DMSO-d₆) δ 0.31-0.44 (2H, m), 0.51-0.64 (2H, m), 0.97-1.15 (1H, m), 1.23-1.40 (1H, m), 1.50-1.72 (3H, m), 1.78-2.10 (6H, m), 2.74 (1H, ddd, J = 9.84, 6.25, 3.60 Hz), 2.87-3.05 (3H, m), 3.82-4.00 (1H, m), 6.93 (1H, d, J = 3.79 Hz), 7.63 (1H, d, J = 3.79 Hz), 8.25 (1H, d, J = 7.95 Hz),
30 9.39 (2H, brs).

[0401]

Example 37

5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

35 A) tert-butyl (cyclopropylmethyl)(trans-2-(5-((5-methyl-1,3,4-

thiadiazol-2-yl)carbamoyl)thiophen-2-yl)cyclopropyl) carbamate
5-(trans-2-((tert-

Butoxycarbonyl)(cyclopropylmethyl)amino)cyclopropyl)thiophene-
2-carboxylic acid (35.0 mg), 5-methyl-1,3,4-thiadiazol-2-amine

5 (13.1 mg) and triethylamine (0.029 mL) were dissolved in DMF
(3 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-
tetramethyluronium hexafluorophosphate (47.3 mg) was added,
and the mixture was stirred at room temperature overnight. To
the reaction mixture were added ethyl acetate and water, and
10 the mixture was extracted with ethyl acetate. The extract was
washed with saturated brine, dried over anhydrous sodium
sulfate, and concentrated under reduced pressure. The residue
was purified by silica gel column chromatography (hexane/ethyl
acetate) to give the title compound (23.7 mg).

15 ¹H NMR (300 MHz, DMSO-d₆) δ 0.08-0.31 (2H, m), 0.35-0.53 (2H,
m), 0.90-1.07 (1H, m), 1.20-1.50 (11H, m), 2.33-2.45 (1H, m),
2.62 (3H, s), 2.78-2.88 (1H, m), 2.90-3.07 (1H, m), 3.19 (1H,
dd, J = 14.39, 6.82 Hz), 6.98 (1H, d, J = 3.79 Hz), 8.08 (1H,
brs), 12.85 (1H, brs).

20 [0402]

B) 5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-
methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide
dihydrochloride

To a mixture of tert-butyl (cyclopropylmethyl)(trans-2-
25 (5-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-
yl)cyclopropyl)carbamate (22.0 mg) and ethyl acetate (2 mL)
was added 4 mol/L hydrogen chloride/ethyl acetate solution
(0.127 mL) at 0°C, and the mixture was stirred at room
temperature overnight. The precipitated solid was collected by
30 filtration, and washed with ethyl acetate and methanol to give
the title compound (10.2 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.32-0.44 (2H, m), 0.51-0.65 (2H,
m), 0.97-1.19 (1H, m), 1.30-1.45 (1H, m), 1.65-1.81 (1H, m),
2.62 (3H, s), 2.78-2.90 (1H, m), 2.96 (2H, d, J = 5.68 Hz),
35 3.01-3.11 (1H, m), 3.86 (2H, s), 7.06 (1H, d, J = 3.79 Hz),

8.09 (1H, s), 9.55 (2H, brs).

[0403]

Example 38

N-cyclopentyl-5-(trans-2-(tetrahydro-2H-pyran-4-

5 yl amino)cyclopropyl)thiophene-3-carboxamide hydrochloride

[optical isomer, compound derived from methyl 5-(trans-2-

((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-

carboxylate (optical isomer, retention time long)]

A) methyl 5-(trans-2-((tert-

10 butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate

(optical isomer, retention time long)

B) methyl 5-(trans-2-((tert-

butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate

(optical isomer, retention time short)

15 A racemate (16.9 g) of methyl 5-(trans-2-((tert-
butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate was
fractionated by SFC (column: CHIRALPAK AD, 20 mmID×250 mmL,
manufactured by Daicel Corporation, mobile phase: carbon
dioxide/methanol = 820/180), and the obtained fraction was
20 concentrated under reduced pressure to give the title compound
(7.23 g) having a short retention time and the title compound
(7.39 g) having a long retention time.

[0404]

methyl 5-(trans-2-((tert-

25 butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate
(optical isomer, retention time long)

¹H NMR (300 MHz, DMSO-d₆) δ 1.07-1.25 (2H, m), 1.39 (9H, s),
2.08 (1H, ddd, J = 9.09, 6.25, 3.22 Hz), 2.60 (1H, d, J = 5.30
Hz), 3.76 (3H, s), 7.08-7.14 (1H, m), 7.29 (1H, brs), 8.07 (1H,
30 d, J = 1.14 Hz).

HPLC retention time 2.859 min (column: CHIRALPAK ADH (trade
name), 4.6 mmID×150 mmL, mobile phase: carbon dioxide/methanol
= 820/180, flow rate: 4.0 mL/min, temperature: 35°C, detection:
UV 220 nm, concentration: 0.5 mg/mL, injection volume: 0.005
35 mL).

[0405]

methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time short)

5 ¹H NMR (300 MHz, DMSO-d₆) δ 1.08-1.24 (2H, m), 1.39 (9H, s), 2.08 (1H, ddd, J = 8.99, 6.15, 3.03 Hz), 2.60 (1H, brs), 3.77 (3H, s), 7.12 (1H, dd, J = 1.51, 0.76 Hz), 7.29 (1H, brs), 8.07 (1H, d, J = 1.14 Hz).

HPLC retention time 1.950 min (column: CHIRALPAK ADH (trade
10 name), 4.6 mmIDx150 mmL, mobile phase: carbon dioxide/methanol = 820/180, flow rate: 4.0 mL/min, temperature: 35°C, detection: UV 220 nm, concentration: 0.5 mg/mL, injection volume: 0.005 mL).

Methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate
15 (optical isomer, retention time long) was clarified to be methyl 5-((1R,2R)-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate.

Methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate
20 (optical isomer, retention time short) was clarified to be methyl 5-((1S,2S)-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate.

[0406]

25 C) 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylic acid [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)]

30 Methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long) (458 mg) was dissolved in methanol (5 mL), 2 mol/L aqueous sodium hydroxide solution (1.93 mL) was added at 0°C, and the mixture was stirred at room
35 temperature overnight. The reaction mixture was acidified with

1 mol/L hydrochloric acid at 0°C, and extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound

5 (435 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.05-1.22 (2H, m), 1.39 (9H, s), 2.07 (1H, ddd, J = 9.09, 6.06, 3.03 Hz), 2.59 (1H, d, J = 7.57 Hz), 7.05-7.09 (1H, m), 7.28 (1H, brs), 7.97 (1H, d, J = 1.51 Hz), 12.61 (1H, brs).

10 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylic acid [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] was
15 clarified to be 5-((1R,2R)-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylic acid. [0407]

D) tert-butyl (trans-2-(4-(cyclopentylcarbamoyl)thiophen-2-yl)cyclopropyl)carbamate [optical isomer, compound derived
20 from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)]

5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylic acid
25 [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] (70.0 mg), cyclopentanamine (23.1 mg) and triethylamine (0.069 mL) were dissolved in DMF (3 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-
30 tetramethyluronium hexafluorophosphate (113 mg) was added, and the mixture was stirred at room temperature overnight. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was successively washed with water and saturated brine, dried over
35 anhydrous sodium sulfate, and concentrated under reduced

pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (77.8 mg).

MS: $[M+H]^+$ 351.3.

5 tert-Butyl (trans-2-(4-(cyclopentylcarbamoyl)thiophen-2-yl)cyclopropyl)carbamate [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] was clarified to be
10 tert-butyl ((1R,2R)-2-(4-(cyclopentylcarbamoyl)thiophen-2-yl)cyclopropyl)carbamate.

[0408]

E) 5-(trans-2-aminocyclopropyl)-N-cyclopentylthiophene-3-carboxamide hydrochloride [optical isomer, compound derived
15 from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)]

To a mixture of tert-butyl (trans-2-(4-(cyclopentylcarbamoyl)thiophen-2-yl)cyclopropyl)carbamate
20 [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] (75.0 mg) and ethyl acetate (2 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (0.535 mL) at 0°C, and the
25 mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (56.2 mg).

MS: $[M+H-(HCl)]^+$ 251.2.

5-(trans-2-Aminocyclopropyl)-N-cyclopentylthiophene-3-carboxamide hydrochloride [optical isomer, compound derived
30 from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] was clarified to be 5-((1R,2R)-2-aminocyclopropyl)-N-cyclopentylthiophene-3-
35 carboxamide hydrochloride.

[0409]

F) N-cyclopentyl-5-(trans-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide hydrochloride [optical isomer, compound derived from methyl 5-(trans-2-
5 ((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)]

To a mixture of 5-(trans-2-aminocyclopropyl)-N-cyclopentylthiophene-3-carboxamide hydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-
10 butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] (53.0 mg), sodium hydrogen carbonate (38.8 mg), THF (3 mL) and methanol (3 mL) was added dihydro-2H-pyran-4(3H)-one (22.2 mg) at room temperature. Under a nitrogen atmosphere, the reaction mixture
15 was stirred at 60°C for 1.5 hr and at room temperature for 30 min, and sodium borohydride (10.5 mg) was added at 0°C. The reaction mixture was stirred at room temperature for 1 hr, and ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture. The
20 mixture was extracted with ethyl acetate, and the extract was successively washed with water and saturated brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/methanol), and to the
25 obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The reaction mixture was stirred at room temperature overnight, and the precipitated solid was collected by filtration, and washed with ethyl acetate to give the title compound (38.9 mg).

30 ¹H NMR (300 MHz, DMSO-d₆) δ 1.28-1.39 (1H, m), 1.39-1.75 (9H, m), 1.78-1.92 (2H, m), 1.93-2.06 (2H, m), 2.73 (1H, ddd, J = 9.75, 6.34, 3.60 Hz), 2.99 (1H, brs), 3.24-3.34 (2H, m), 3.47 (1H, brs), 3.85-4.02 (2H, m), 4.15 (1H, sxt, J = 6.74 Hz), 7.29 (1H, s), 7.94 (1H, d, J = 1.14 Hz), 8.02 (1H, d, J = 7.19
35 Hz), 9.54 (2H, brs).

N-Cyclopentyl-5-(trans-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide hydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] was clarified to be N-cyclopentyl-5-((1R,2R)-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide hydrochloride [0410]

10 Example 39

N-cyclopentyl-5-(trans-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide hydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time short)]

By a method similar to that of Example 38, the title compound was obtained.

N-Cyclopentyl-5-(trans-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide hydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time short)] was clarified to be N-cyclopentyl-5-((1S,2S)-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide. [0411]

Example 40

N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide dihydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)]

A) tert-butyl (trans-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate

(optical isomer, retention time long)]

5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylic acid [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] (70.0 mg), 5-methyl-1,3,4-thiadiazol-2-amine (31.3 mg) and triethylamine (0.069 mL) were dissolved in DMF (3 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (113 mg) was added, and the mixture was stirred at room temperature overnight. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (37.6 mg).

MS: [M+H]⁺ 381.2.

tert-Butyl (trans-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] was clarified to be tert-butyl ((1R,2R)-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate.

[0412]

B) 5-(trans-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide dihydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)]

To a mixture of tert-butyl (trans-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-

carboxylate (optical isomer, retention time long)] (37.6 mg), ethyl acetate (2 mL) and methanol (2 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (0.247 mL) at 0°C, and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (34.5 mg).
MS: [M+H-(2HCl)]⁺ 281.1.

5-(trans-2-Aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide dihydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] was clarified to be 5-((1R,2R)-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide dihydrochloride.
[0413]

C) N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide dihydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)]

To a mixture of 5-(trans-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide dihydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] (33.0 mg), sodium hydrogen carbonate (19.6 mg), THF (2 mL) and methanol (2 mL) was added tetrahydro-2H-pyran-4-carbaldehyde (12.8 mg) at room temperature. Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, and sodium borohydride (5.30 mg) was added at 0°C. The reaction mixture was stirred at room temperature for 1 hr, and ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture. The mixture was extracted with ethyl acetate, and the extract was successively washed with water and saturated brine and dried

over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/methanol), and to the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The reaction mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (19.6 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.16-1.41 (3H, m), 1.60-1.77 (3H, m), 1.84-2.06 (1H, m), 2.63 (3H, s), 2.70-2.84 (1H, m), 2.94-3.11 (3H, m), 3.29 (2H, t, J = 11.93 Hz), 3.90-4.00 (2H, m), 7.51 (1H, s), 8.46 (1H, s), 9.26 (2H, brs).

N-(5-Methyl-1,3,4-thiadiazol-2-yl)-5-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide dihydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] was clarified to be N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide dihydrochloride.

[0414]

Example 41

N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide dihydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time short)]

By a method similar to that of Example 40, the title compound was obtained.

N-(5-Methyl-1,3,4-thiadiazol-2-yl)-5-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide dihydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate

(optical isomer, retention time short)] was clarified to be N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-((1S,2S)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide dihydrochloride.

5 [0415]

Example 42

5-(trans-2-(cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)]

10 A) tert-butyl (trans-2-(4-((tetrahydro-2H-pyran-4-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)]

5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylic acid [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] (70.0 mg) was dissolved in DMF (3 mL). Tetrahydro-2H-pyran-4-amine (27.5 mg), triethylamine (0.069 mL) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (113 mg) were
25 added, and the mixture was stirred at room temperature overnight. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced
30 pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (71.8 mg).

MS: [M+H]⁺ 367.2.

tert-Butyl (trans-2-(4-((tetrahydro-2H-pyran-4-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate [optical
35

isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] was clarified to be tert-butyl ((1R,2R)-2-(4-((tetrahydro-2H-pyran-4-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate.

[0416]

B) 5-(trans-2-aminocyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide dihydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)]

To a mixture of tert-butyl (trans-2-(4-((tetrahydro-2H-pyran-4-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] (69.5 mg) and ethyl acetate (2 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (0.474 mL) at 0°C, and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (53.6 mg).

MS: [M+H-(2HCl)]⁺ 267.2.

5-(trans-2-Aminocyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide dihydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] was clarified to be 5-((1R,2R)-2-aminocyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide dihydrochloride.

[0417]

C) 5-(trans-2-(cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)]

To a mixture of 5-(trans-2-aminocyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide dihydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] (51.0 mg), sodium hydrogen carbonate (35.4 mg), THF (2 mL) and methanol (2 mL) was added cyclobutanone (14.2 mg) at room temperature. Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, and sodium borohydride (9.56 mg) was added at 0°C. The reaction mixture was stirred at room temperature for 1 hr, and ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture. The mixture was extracted with ethyl acetate, and the extract was successively washed with water and saturated brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/methanol), and the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The reaction mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (28.7 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.03 (2H, d, J = 6.06 Hz), 1.22-1.36 (1H, m), 1.43-1.62 (3H, m), 1.65-1.91 (4H, m), 2.11-2.32 (4H, m), 2.65 (1H, brs), 2.85 (1H, d, J = 3.41 Hz), 3.77-4.00 (4H, m), 7.26 (1H, s), 7.95 (1H, s), 8.05 (1H, d, J = 7.95 Hz), 9.49 (2H, brs).

5-(trans-2-(Cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time long)] was clarified to be 5-((1R,2R)-2-(cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride.

[0418]

Example 43

5-(trans-2-(cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time short)]

By a method similar to that of Example 42, the title compound was obtained.

5-(trans-2-(Cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride [optical isomer, compound derived from methyl 5-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate (optical isomer, retention time short)] was clarified to be 5-((1S,2S)-2-(cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride.

[0419]

Example 44

N-(4,4-difluorocyclohexyl)-5-((1R,2R)-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide hydrochloride
A) tert-butyl ((1R,2R)-2-(4-((4,4-difluorocyclohexyl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate

5-((1R,2R)-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylic acid (400 mg) was dissolved in DMF (10 mL), 4,4-difluorocyclohexanamine hydrochloride (291 mg), triethylamine (0.59 mL) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (644 mg) were added, and the mixture was stirred at room temperature overnight. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (680 mg). The title

compound was used for the next reaction without further purification.

MS: $[M-tBu(C_4H_9)+2H]^+$ 345.1.

[0420]

- 5 B) 5-((1R,2R)-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide hydrochloride
tert-Butyl ((1R,2R)-2-(4-((4,4-difluorocyclohexyl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate (680 mg) and 4 mol/L hydrogen
10 chloride/ethyl acetate solution (1.7 mL) were stirred at room temperature overnight. The precipitated solid was collected by filtration to give the title compound (500 mg).

MS: $[M+H-(HCl)]^+$ 301.1.

[0421]

- 15 C) N-(4,4-difluorocyclohexyl)-5-((1R,2R)-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide hydrochloride

To a mixture of 5-((1R,2R)-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide hydrochloride (62.0
20 mg), borane-2-picoline complex (29.5 mg), methanol (2.0 mL) and acetic acid (0.2 mL) was added dihydro-2H-pyran-4(3H)-one (58.2 mg) at room temperature. The reaction mixture was stirred at room temperature for 18 hr, and saturated aqueous sodium hydrogen carbonate solution was added at 0°C. The
25 mixture was extracted with ethyl acetate, and the extract was successively washed with water and saturated brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by NH silica gel column chromatography (ethyl acetate/hexane), and to the
30 obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The reaction mixture was concentrated under reduced pressure, and the obtained residue was crystallized from ethanol/heptane to give the title compound (40.0 mg).

1H NMR (300 MHz, DMSO- d_6) δ 1.27-1.41 (1H, m), 1.49-1.72 (5H, m), 1.77-2.16 (8H, m), 2.61-2.73 (1H, m), 2.89-3.08 (1H, m),

3.26-3.39 (2H, m), 3.40-3.55 (1H, m), 3.83-4.02 (3H, m), 7.24-7.31 (1H, m), 7.92-7.98 (1H, m), 7.99-8.08 (1H, m), 8.99-9.48 (2H, m).

[0422]

5 The compounds of Examples 45 to 59 could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

Example 45

10 5-((1R,2R)-2-(cis-(4-aminocyclohexyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide dihydrochloride

or

15 5-((1R,2R)-2-(trans-(4-aminocyclohexyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide dihydrochloride

Example 46

20 5-((1R,2R)-2-(trans-(4-aminocyclohexyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide dihydrochloride

or

25 5-((1R,2R)-2-(cis-(4-aminocyclohexyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide dihydrochloride

Example 47

30 5-((1R,2R)-2-((4,4-difluorocyclohexyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride

Example 48

35 5-((1R,2R)-2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide dihydrochloride

Example 49

40 N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-((1R,2R)-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide

hydrochloride

Example 50

N-(2-methyl-1,3-thiazol-5-yl)-5-((1R,2R)-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide

5 hydrochloride

Example 51

N-(4,4-difluorocyclohexyl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide

hydrochloride

10 Example 52

5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-3-carboxamide

hydrochloride

15 Example 53

N-(2-methyl-1,3-thiazol-5-yl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide

dihydrochloride

Example 54

20 N-(1,3-dimethyl-1H-pyrazol-4-yl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide

dihydrochloride

Example 55

N-(1,5-dimethyl-1H-pyrazol-4-yl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide

25 dihydrochloride

Example 56

N-cyclopropyl-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide

30 hydrochloride

Example 57

4-(trans-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide hydrochloride

Example 58

35 N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-(trans-2-((tetrahydro-2H-

pyran-4-ylmethyl) amino) cyclopropyl) thiophene-2-carboxamide
hydrochloride

Example 59

4-(trans-2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl) -N-
5 (5-methyl-1,3,4-thiadiazol-2-yl) thiophene-2-carboxamide
dihydrochloride

[0423]

Example 60

3-(4-(((1R,2R)-2-(4-((5-methyl-1,3,4-thiadiazol-2-
10 yl) carbamoyl) -2-thienyl) cyclopropyl) amino) methyl) piperidin-1-
yl) propanoic acid trihydrochloride

A) tert-butyl 4-(((1R,2R)-2-(4-((5-methyl-1,3,4-thiadiazol-2-
yl) carbamoyl) thiophen-2-
yl) cyclopropyl) amino) methyl) piperidine-1-carboxylate

15 By a method similar to that of Example 2, the title
compound was obtained.

MS: [M+H]⁺ 478.3.

[0424]

B) tert-butyl 4-((2,2,2-trifluoro-N-((1R,2R)-2-(4-((5-methyl-
20 1,3,4-thiadiazol-2-yl) carbamoyl) thiophen-2-
yl) cyclopropyl) acetamido) methyl) piperidine-1-carboxylate

To a mixture of tert-butyl 4-(((1R,2R)-2-(4-((5-methyl-
1,3,4-thiadiazol-2-yl) carbamoyl) thiophen-2-
yl) cyclopropyl) amino) methyl) piperidine-1-carboxylate (301 mg),
25 triethylamine (127.0 mg) and THF (4.0 mL) was added
trifluoroacetic anhydride (355.0 mg) at room temperature. The
reaction mixture was stirred at room temperature for 18 hr,
and saturated aqueous sodium hydrogen carbonate solution was
added at 0°C. The mixture was extracted with ethyl acetate,
30 and the extract was successively washed with water and
saturated brine and dried over anhydrous sodium sulfate. The
solvent was evaporated under reduced pressure. The residue was
purified by NH silica gel column chromatography (ethyl
acetate/hexane) to give the title compound (213.0 mg).

35 MS: [M+H]⁺ 574.1.

[0425]

C) N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-((1R,2R)-2-(2,2,2-trifluoro-N-(piperidin-4-ylmethyl)acetamido)cyclopropyl)thiophene-3-carboxamide
5 hydrochloride

By a method similar to that of Example 1, step H, the title compound was obtained.

MS: $[M-HCl+H]^+$ 474.1.

[0426]

10 D) tert-butyl 3-(4-((2,2,2-trifluoro-N-((1R,2R)-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-yl)cyclopropyl)acetamido)methyl)piperidin-1-yl)propanoate

To a mixture of N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-((1R,2R)-2-(2,2,2-trifluoro-N-(piperidin-4-ylmethyl)acetamido)cyclopropyl)thiophene-3-carboxamide
15 hydrochloride (50.0 mg), triethylamine (10.9 mg) and ethanol (2.0 mL) was added tert-butyl acrylate (12.6 mg) at room temperature. The reaction mixture was stirred at room temperature overnight, and the solvent was evaporated under
20 reduced pressure. To the residue were added ethyl acetate and saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced
25 pressure and the residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (44.0 mg).

MS: $[M+H]^+$ 602.1.

[0427]

30 E) tert-butyl 3-(4-(((1R,2R)-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-yl)cyclopropyl)amino)methyl)piperidin-1-yl)propanoate

To a mixture of tert-butyl 3-(4-((2,2,2-trifluoro-N-((1R,2R)-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-

35

yl)cyclopropyl)acetamido)methyl)piperidin-1-yl)propanoate (44.0 mg), methanol (1.0 mL) and THF (1.0 mL) was added 1 mol/L aqueous sodium hydroxide solution (1.0 mL) at room temperature. The reaction mixture was stirred at room
5 temperature for 10 min, and water was added at room temperature. The mixture was extracted with ethyl acetate, and the extract was washed with saturated brine and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure to give the title compound (30.0 mg).

10 MS: [M+H]⁺ 506.2.

[0428]

F) 3-(4-(((1R,2R)-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)-2-thienyl)cyclopropyl)amino)methyl)piperidin-1-yl)propanoic acid trihydrochloride

15 A mixture of tert-butyl 3-(4-(((1R,2R)-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-2-yl)cyclopropyl)amino)methyl)piperidin-1-yl)propanoate (30.0 mg) and 4 mol/L hydrogen chloride/ethyl acetate solution (1.0 mL) was stirred at room temperature overnight, and the solvent
20 was evaporated under reduced pressure to give the title compound (33.0 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.29-1.41 (1H, m), 1.44-1.64 (2H, m), 1.72 (1H, brs), 2.01 (3H, d, J = 12.49), 2.63 (3H, s), 2.76-3.08 (7H, m), 3.17-3.35 (3H, m), 3.42-3.56 (2H, m), 7.51
25 (1H, s), 8.46 (1H, d, J = 1.14), 9.36-9.73 (2H, m), 9.96-12.92 (2H, m).

[0429]

The compounds of Examples 61 to 66 could be produced according to the production methods described in the present
30 specification, a method shown in the Examples, or a method analogous thereto.

Example 61

4-(trans-2-(cyclobutylamino)cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide dihydrochloride

35 Example 62

4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide dihydrochloride

Example 63

4-(trans-2-(cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide hydrochloride

Example 64

4-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide hydrochloride

10 Example 65

5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-2-methylthiophene-3-carboxamide hydrochloride

Example 66

15 N-(4,4-difluorocyclohexyl)-2-methyl-5-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide hydrochloride

[0430]

Example 67

20 N-(4,4-difluorocyclohexyl)-2-methyl-5-(trans-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide hydrochloride

To a mixture of 5-(trans-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)-2-methylthiophene-3-carboxamide hydrochloride (30.0 mg), sodium hydrogen carbonate (10.8 mg), THF (1.5 mL) and methanol (1.5 mL) was added dihydro-2H-pyran-4(3H)-one (9.48 μ L). Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, and sodium borohydride (4.85 mg) was added under ice-cooling. The reaction mixture was stirred at room temperature for 1 hr, and diluted with ethyl acetate. Water and saturated aqueous sodium hydrogen carbonate solution were added, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and the solvent was

evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography (ethyl acetate/methanol). To the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution, and the mixture was stirred overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (15.6 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.22-1.36 (1H, m), 1.49-1.75 (5H, m), 1.78-2.09 (9H, m), 2.53 (3H, s), 2.64 (1H, brs), 2.89 (1H, brs), 3.45 (1H, brs), 3.93 (3H, d, J = 10.98 Hz), 7.06 (1H, s), 7.83 (1H, d, J = 7.95 Hz), 9.50 (2H, brs).

[0431]

Example 68

5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride

The title compound could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

[0432]

Example 69

5-(trans-2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide dihydrochloride

To a mixture of 5-(trans-2-aminocyclopropyl)-2-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide dihydrochloride (30.0 mg), sodium hydrogen carbonate (17.2 mg), THF (1.5 mL) and methanol (1.5 mL) was added cyclopropanecarbaldehyde (7.32 μL). Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min. Sodium borohydride (4.63 mg) was added under ice-cooling, and the reaction mixture was stirred at room temperature for 1 hr and diluted with ethyl acetate. Water and saturated aqueous sodium hydrogen carbonate solution were added, and the mixture was extracted with ethyl

acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography

5 (ethyl acetate/methanol). To the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution, and the mixture was stirred overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (22.1 mg).

10 ¹H NMR (300 MHz, DMSO-d₆) δ 0.35-0.45 (2H, m), 0.55-0.65 (2H, m), 1.03-1.20 (1H, m), 1.24-1.36 (1H, m), 1.59-1.70 (1H, m), 2.60-2.75 (8H, m), 2.86-3.02 (3H, m), 7.47 (1H, s), 9.51 (2H, d, J = 4.54 Hz).

[0433]

15 Example 70

2-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide dihydrochloride

The title compound could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

[0434]

Example 71

5-((1R,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride

A) ((1R,4aS,10aR)-7-isopropyl-1,4a-dimethyl-1,2,3,4,4a,9,10,10a-octahydrophenanthren-1-yl)methanamine (1R,2R)-2-(4-bromothiophen-2-yl)cyclopropanecarboxylate

30 ((1R,4aS,10aR)-7-Isopropyl-1,4a-dimethyl-1,2,3,4,4a,9,10,10a-octahydrophenanthren-1-yl)methanamine (1.040 g) was dissolved in methanol (3.2 mL) and diisopropyl ether (15.7 mL) at 40°C, and trans-2-(4-bromothiophen-2-yl)cyclopropanecarboxylic acid (900 mg) was added. The

35 reaction mixture was stirred for 5 min at the same temperature,

and methanol (8 mL) was added. The reaction mixture was heated to 53°C to dissolve the precipitate. The reaction mixture was cooled to 46°C and stirred at 46-47°C for 2 hr. The reaction mixture was slowly cooled to room temperature (28°C) and stirred overnight. The precipitate was collected by filtration, and the obtained solid was washed with a mixed solvent of ethyl acetate/hexane (1/5 = ethyl acetate/hexane (v/v), 4 mL) to give the title compound (594 mg, 95.7%d.e.).

MS: [M-H-(C₂₀H₃₁N)]⁻ 245.0.

10 Column: CHIROBIOTIC R 4.6 mmID×250 mmL

Eluent solvent: Methanol/TEA/AA=1000/3/1 (v/v/v)

Flow rate: 1.0 mL/min

Retention time: 4.4 min.

Temperature: 30°C

15 Detection: UV 254 nm

Concentration: 0.5 mg/mL

Injection volume: 0.010 mL

[0435]

B) ((1R,4aS,10aR)-7-isopropyl-1,4a-dimethyl-

20 1,2,3,4,4a,9,10,10a-octahydrophenanthren-1-yl)methanamine

(1R,2R)-2-(4-bromothiophen-2-yl)cyclopropanecarboxylate

((1R,4aS,10aR)-7-Isopropyl-1,4a-dimethyl-

1,2,3,4,4a,9,10,10a-octahydrophenanthren-1-yl)methanamine

(1R,2R)-2-(4-bromothiophen-2-yl)cyclopropanecarboxylate (290

25 mg, 95.7%d.e.) was dissolved in ethanol (3.77 mL) and

diisopropyl ether (1.45 mL) at 62°C, and diisopropyl ether

(8.12 mL) was further added dropwise at 62°C. The mixture was

cooled to 46°C and stirred at 45-47°C for 2 hr. The mixture was slowly cooled to room temperature (28°C) and stirred

30 overnight. Furthermore, the mixture was stirred at 0°C for 1

hr. The precipitate was collected by filtration, and the

obtained solid was washed with ethyl acetate/hexane (1/5 =

ethyl acetate/hexane (v/v), 2 mL) to give the title compound

(246 mg, 99.6%d.e.).

35 ¹H NMR (300 MHz, DMSO-d₆) δ 0.84 (3H, s), 1.00-1.76 (19H, m),

2.21-2.38 (2H, m), 2.39-2.47 (2H, m), 2.71-2.87 (2H, m), 3.38-3.50 (2H, m), 6.81-6.86 (1H, m), 6.88-6.92 (1H, m), 6.92-6.98 (1H, m), 7.10-7.19 (1H, m), 7.40 (1H, d, J = 1.5 Hz).

Column: CHIROBIOTIC R 4.6 mmID×250 mmL

5 Eluent solvent: Methanol/TEA/AA=1000/3/1 (v/v/v)

Flow rate: 1.0 mL/min

Retention time: 4.3 min.

Temperature: 30°C

Detection: UV 254 nm

10 Concentration: 0.5 mg/mL

Injection volume: 0.010 mL

[0436]

C) (1R,2R)-2-(4-bromothiophen-2-yl)cyclopropanecarboxylic acid

To a suspension of ((1R,4aS,10aR)-7-isopropyl-1,4a-dimethyl-1,2,3,4,4a,9,10,10a-octahydrophenanthren-1-yl)methanamine (1R,2R)-2-(4-bromothiophen-2-yl)cyclopropanecarboxylate (26.8 g) in ethyl acetate (300 mL) were added 1 mol/L aqueous sodium hydroxide solution (60.4 mL) and water (200 mL) at room temperature, and the mixture was
20 stirred at room temperature for 5 min. The aqueous layer was separated and washed with ethyl acetate (200 mL, twice). To the aqueous layer was added 2 mol/L hydrochloric acid (35.2 mL) under ice-cooling at 0°C to adjust the mixture from pH 2 to pH 3, and the mixture was extracted with ethyl acetate (100 mL,
25 twice). The organic layer was separated, and the extract was washed with saturated brine, and dried over magnesium sulfate. The solvent was evaporated under reduced pressure to give the title compound (11.7 g).

¹H NMR (300 MHz, CDCl₃) δ 1.39 (1H, ddd, J = 8.5, 6.4, 4.7 Hz),
30 1.64-1.73 (1H, m), 1.90-1.98 (1H, m), 2.66-2.78 (1H, m), 6.75 (1H, dd, J = 1.5, 0.8 Hz), 7.02 (1H, d, J = 1.5 Hz).

[0437]

D) tert-butyl ((1R,2R)-2-(4-bromothiophen-2-yl)cyclopropyl)carbamate

35 To a solution of (1R,2R)-2-(4-bromothiophen-2-

yl)cyclopropanecarboxylic acid (15 g) and tert-butyl alcohol (150 mL) were added triethylamine (10.15 mL) and diphenylphosphoryl azide (13.07 mL) at room temperature, and the mixture was stirred at room temperature for 1 hr, and then
5 at 80°C overnight. The reaction mixture was cooled to room temperature, saturated aqueous sodium hydrogen carbonate solution was added, and the mixture was extracted with ethyl acetate. The organic layer was separated, successively washed with water and saturated brine, dried over anhydrous sodium
10 sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (14.6 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.04-1.24 (2H, m), 1.38 (9H, s), 2.07 (1H, ddd, J = 9.18, 6.15, 3.22 Hz), 2.61 (1H, brs), 6.80
15 (1H, d, J = 0.76 Hz), 7.27 (1H, brs), 7.38 (1H, d, J = 1.51 Hz).

[0438]

E) methyl 5-((1R,2R)-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylate

20 To a solution of tert-butyl ((1R,2R)-2-(4-bromothiophen-2-yl)cyclopropyl)carbamate (7.2 g) in methanol (180 mL) were added triethylamine (6.31 mL) and dichloro(1,1'-bis(diphenylphosphino)ferrocene)palladium (828 mg) at room temperature, and the mixture was stirred at 90°C for 6 hr under
25 a carbon monoxide atmosphere (3 atm). The reaction mixture was filtered through celite and washed with methanol. The filtrate was concentrated under reduced pressure, water was added to the residue, and the mixture was extracted with ethyl acetate. The contaminants in the organic layer were removed by
30 filtration, and the filtrate was concentrated under reduced pressure to give residue A.

To a solution of tert-butyl ((1R,2R)-2-(4-bromothiophen-2-yl)cyclopropyl)carbamate (7.2 g) in methanol (180 mL) were added triethylamine (6.31 mL) and dichloro(1,1'-
35 bis(diphenylphosphino)ferrocene)palladium (828 mg) at room

temperature, and the mixture was stirred at 90°C for 6 hr under a carbon monoxide atmosphere (3 atm). The reaction mixture was filtered through celite, and washed with methanol. The filtrate was concentrated under reduced pressure, water was added to the residue, and the mixture was extracted with ethyl acetate. The contaminants in the organic layer were removed by filtration, and the filtrate was concentrated under reduced pressure to give residue B.

The residues A and B were combined and purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (11.1 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.05-1.25 (2H, m), 1.35-1.42 (9H, m), 2.08 (1H, ddd, J = 9.18, 6.34, 3.03 Hz), 2.60 (1H, brs), 3.76 (3H, s), 7.12 (1H, d, J = 0.76 Hz), 7.28 (1H, brs), 8.07 (1H, d, J = 1.51 Hz).

[0439]

F) 5-((1R,2R)-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-3-carboxylic acid

By a method similar to that of Example 38, step C, the title compound was obtained.

[0440]

G) tert-butyl ((1R,2R)-2-(4-((tetrahydro-2H-pyran-4-yl)carbamoyl)thiophen-2-yl)cyclopropyl)carbamate

By a method similar to that of Example 42, step A, the title compound was obtained.

[0441]

H) 5-((1R,2R)-2-aminocyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide dihydrochloride

By a method similar to that of Example 42, step B, the title compound was obtained.

[0442]

I) 5-((1R,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride

To a mixture of 5-((1R,2R)-2-aminocyclopropyl)-N-

(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide dihydrochloride (1.54 g), triethylamine (2.12 mL), THF (20 mL) and methanol (20 mL) was added cyclopropanecarbaldehyde (0.428 g) at room temperature. The reaction mixture was stirred at
5 room temperature for 2 hr, and sodium borohydride (0.231 g) was added at 0°C. The reaction mixture was stirred at 0°C for 10 min and added to water. The methanol and tetrahydrofuran were evaporated under reduced pressure, and the mixture was extracted with ethyl acetate. The extract was washed with
10 saturated brine and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (ethyl acetate/methanol). To the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution and the mixture
15 was concentrated under reduced pressure. The obtained residue was crystallized from ethanol/heptane to give the title compound (1.04 g).

¹H NMR (300 MHz, DMSO-d₆) δ 0.33-0.43 (2H, m), 0.51-0.64 (2H, m), 0.99-1.15 (1H, m), 1.24-1.37 (1H, m), 1.42-1.65 (3H, m),
20 1.67-1.78 (2H, m), 2.64-2.75 (1H, m), 2.90-3.04 (3H, m), 3.32-3.42 (2H, m), 3.79-4.00 (3H, m), 7.28 (1H, s), 7.95 (1H, d, J = 1.5 Hz), 8.05 (1H, d, J = 8.0 Hz), 9.25 (2H, brs).

[0443]

The compounds of Examples 72 to 80 could be produced
25 according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

Example 72

N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-((1S,2R)-2-((tetrahydro-
30 2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide hydrochloride

Example 73

4-((1R,2S)-2-(cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide hydrochloride

35 Example 74

N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-((1R,2S)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide hydrochloride

Example 75

5 4-((1R,2S)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide hydrochloride

Example 76

4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide hydrochloride

Example 77

4-((1S,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide hydrochloride

15 Example 78

4-((1S,2R)-2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

Example 79

20 4-((1R,2S)-2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

Example 80

5-((1R,2R)-2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide hydrochloride

[0444]

Example 81

30 N-(5-methyl-1,2-oxazol-3-yl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide hydrochloride

A) tert-butyl ((1R,2R)-2-(4-((5-methylisoxazol-3-yl)carbamoyl)thiophen-2-yl)cyclopropyl)((tetrahydro-2H-pyran-4-yl)methyl)carbamate

35 To a mixture of 5-((1R,2R)-2-(tert-

butoxycarbonyl)((tetrahydro-2H-pyran-4-yl)methyl)amino)cyclopropyl)thiophene-3-carboxylic acid (50.0 mg) and THF (2 mL) was added oxalyl chloride (0.014 mL) at 0°C, and the mixture was stirred at 0°C for 1 hr. The solvent was
5 evaporated under reduced pressure to give an acid chloride intermediate.

To a mixture of 5-methylisoxazol-3-amine (19.3 mg), DMAP (8.01 mg) and pyridine (2 mL) was added a mixture of the aforementioned acid chloride and THF (1.5 mL) at 60°C, and the
10 mixture was stirred at 60°C for 2 hr. The reaction mixture was neutralized with saturated aqueous sodium hydrogen carbonate solution, and extracted with ethyl acetate. The extract was successively washed with water and saturated brine, and dried over magnesium sulfate, and the solvent was evaporated under
15 reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (17.2 mg).

MS: [M-Boc+H]⁺ 362.3.

[0445]

20 B) N-(5-methyl-1,2-oxazol-3-yl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide hydrochloride

By a method similar to that of Example 1, step H, the title compound was obtained.

25 [0446]

Example 82

4-((1S,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

30 A) (R)-2-amino-3-phenylpropan-1-ol (1R,2R)-2-(5-bromothiophen-3-yl)cyclopropanecarboxylate

trans-2-(5-Bromothiophen-3-yl)cyclopropanecarboxylic acid (47.8 g) and (R)-2-amino-3-phenylpropan-1-ol (29.2 g) were dissolved in ethanol (480 mL) at 57°C, diisopropyl ether (480
35 mL) was added, and the mixture was cooled to 53°C. The

reaction mixture was stirred at 53°C for 1 hr, cooled to room temperature, and stirred overnight. The reaction mixture was cooled to 0°C and stirred for 1 hr. The precipitate was collected by filtration, and washed with a mixed solvent of
5 ethyl acetate/hexane (1/2 = ethyl acetate/hexane (v/v)) to give the title compound (32.6 g, >99% d.e.).

MS: [M-H-(C₉H₁₃NO)]⁻ 246.8.

[0447]

B) (R)-2-amino-3-phenylpropan-1-ol (1R,2R)-2-(5-bromothiophen-
10 3-yl)cyclopropanecarboxylate

(R)-2-Amino-3-phenylpropan-1-ol (1R,2R)-2-(5-bromothiophen-3-yl)cyclopropanecarboxylate (32.5 g) was dissolved in ethanol (430 mL) at 60°C, diisopropyl ether (850 mL) was added, and the mixture was cooled to 55°C. The
15 reaction mixture was stirred at 55°C for 1 hr, cooled to room temperature, and stirred overnight. The reaction mixture was cooled to 0°C, stirred for 1 hr, and the precipitate was collected by filtration and washed with a mixed solvent of ethyl acetate/hexane (1/2 = ethyl acetate/hexane (v/v)) to
20 give the title compound (27.5 g, >99% d.e.).

MS: [M-H-(C₉H₁₃NO)]⁻ 246.8.

[0448]

C) (1R,2R)-2-(5-bromothiophen-3-yl)cyclopropanecarboxylic acid

To a mixture of (R)-2-amino-3-phenylpropan-1-ol (1R,2R)-
25 2-(5-bromothiophen-3-yl)cyclopropanecarboxylate (26.0 g) and ethyl acetate was added 1 mol/L aqueous hydrochloric acid solution, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent
30 was evaporated under reduced pressure. Toluene was added to the residue, and the mixture was concentrated under reduced pressure to give the title compound (16.1 g).

MS: [M-H]⁻ 246.8.

[0449]

35 D) tert-butyl ((1R,2S)-2-(5-bromothiophen-3-

yl)cyclopropyl)carbamate

To a mixture of (1R,2R)-2-(5-bromothiophen-3-yl)cyclopropanecarboxylic acid (16.0 g) and tert-butyl alcohol (200 mL) were added triethylamine (10.8 mL) and
5 diphenylphosphoryl azide (16.7 mL) at room temperature, and the mixture was stirred at room temperature for 2 hr and at 80°C overnight. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, and dried
10 over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure, and the residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (18.1 g).

¹H NMR (300 MHz, DMSO-d₆) δ 0.91-1.11 (2H, m), 1.37 (9H, s),
15 1.87 (1H, ddd, J = 9.28, 6.25, 3.41 Hz), 2.56 (1H, s), 7.00 (1H, d, J = 1.89 Hz), 7.12 (1H, d, J = 1.89 Hz), 7.18 (1H, brs).

[0450]

E) methyl 4-((1S,2R)-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-2-carboxylate
20

To a mixture of tert-butyl ((1R,2S)-2-(5-bromothiophen-3-yl)cyclopropyl)carbamate (9.00 g) and methanol (200 mL) were added triethylamine (7.88 mL) and 1,1'-bis(diphenylphosphino)ferrocene-palladium(II) dichloride (1.04
25 g), and the mixture was heated at 90°C for 6 hr under a carbon monoxide atmosphere (3 atm). The insoluble material was filtered off by celite and washed with methanol, and the filtrate was concentrated under reduced pressure and extracted with ethyl acetate and water. The extract was washed with
30 saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give residue A.

To a mixture of tert-butyl ((1R,2S)-2-(5-bromothiophen-3-yl)cyclopropyl)carbamate (9.00 g) and methanol (200 mL) were
35 added triethylamine (7.88 mL) and 1,1'-

bis(diphenylphosphino)ferrocene-palladium(II) dichloride (1.04 g), and the mixture was stirred at 90°C for 6 hr under a carbon monoxide atmosphere (3 atm). The insoluble material was filtered off by celite and washed with methanol, and the filtrate was concentrated under reduced pressure and extracted with ethyl acetate and water. The extract and the residues A were combined, washed with 5% ammonia water (twice), N-acetyl-L-cysteine aqueous solution (twice), water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (15.2 g).

MS: [M-Boc+2H]⁺ 198.1.

[0451]

F) 4-((1S,2R)-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-2-carboxylic acid

To a solution of methyl 4-((1S,2R)-2-((tert-butoxycarbonyl)amino)cyclopropyl)thiophene-2-carboxylate (15.2 g) in THF (50 mL)/methanol (50 mL) was added 2 mol/L aqueous sodium hydroxide solution (63.9 mL) at 0°C, and the mixture was stirred at room temperature overnight. The reaction mixture was acidified with 1 mol/L hydrochloric acid at 0°C, and extracted with ethyl acetate. The extract was successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was washed with diisopropyl ether to give the title compound (13.4 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.00-1.11 (2H, m), 1.34-1.42 (9H, m), 1.92 (1H, ddd, J = 9.09, 6.25, 3.22 Hz), 2.54-2.66 (1H, m), 7.21 (1H, br. s.), 7.42 (1H, d, J = 1.14 Hz), 7.50 (1H, d, J = 1.51 Hz), 12.99 (1H, brs).

[0452]

G) tert-butyl ((1R,2S)-2-(5-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-3-yl)cyclopropyl)carbamate

To a solution of 4-((1S,2R)-2-((tert-

butoxycarbonyl)amino)cyclopropyl)thiophene-2-carboxylic acid (13.3 g) in DMF (200 mL) were added 5-methyl-1,3,4-thiadiazol-2-amine (6.49 g), triethylamine (13.08 mL) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium

5 hexafluorophosphate (21.42 g), and the mixture was stirred at room temperature overnight. To the reaction mixture were added water and ethyl acetate, and the precipitate was collected by filtration and washed with ethyl acetate to give the title compound (12.1 g). The filtrate was extracted with ethyl
10 acetate. The organic layer was successively washed with 0.1 mol/L hydrochloric acid, saturated aqueous sodium hydrogen carbonate solution, water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. To the residue was added ethyl acetate, and the
15 precipitate was collected by filtration. The obtained solid was washed with ethyl acetate to give the title compound (2.86 g).

MS: [M+H]⁺ 381.1.

[0453]

20 H) 4-((1S,2R)-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

To a suspension of tert-butyl ((1R,2S)-2-(5-((5-methyl-1,3,4-thiadiazol-2-yl)carbamoyl)thiophen-3-yl)cyclopropyl)carbamate (1.67 g) in ethyl acetate (10 mL) and
25 methanol (10 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (10.97 mL) at 0°C, and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration, and washed with ethyl acetate to give the title compound (1.55 g).

30 MS: [M+H-(2HCl)]⁺ 281.0.

[0454]

I) 4-((1S,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

35 To a mixture of 4-((1S,2R)-2-aminocyclopropyl)-N-(5-

methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide
dihydrochloride (1.00 g), sodium hydrogen carbonate (549 mg),
THF (15 mL) and methanol (15 mL) was added
cyclopropanecarbaldehyde (254 μ L). Under a nitrogen atmosphere,
5 the reaction mixture was stirred at 60°C for 1.5 hr and at room
temperature for 30 min. Sodium borohydride (161 mg) was added
under ice-cooling, and the reaction mixture was stirred at
room temperature for 1 hr. The reaction mixture was diluted
with ethyl acetate, water and saturated aqueous sodium
10 hydrogen carbonate solution were added, and the mixture was
extracted with ethyl acetate. The extract was successively
washed with water and saturated brine, and dried over
anhydrous sodium sulfate. The solvent was evaporated under
reduced pressure, and the obtained residue was purified by
15 silica gel column chromatography (ethyl acetate/methanol). To
the obtained fraction was added 4 mol/L hydrogen
chloride/ethyl acetate solution, and the mixture was stirred
overnight and concentrated under reduced pressure. The residue
was crystallized from methanol/diisopropyl ether to give the
20 title compound (723 mg).

^1H NMR (300 MHz, DMSO- d_6) δ 0.33-0.42 (2H, m), 0.54-0.65 (2H,
m), 1.01-1.17 (1H, m), 1.20-1.32 (1H, m), 1.53-1.66 (1H, m),
2.56-2.65 (4H, m), 2.86-3.02 (3H, m), 7.75 (1H, s), 8.07 (1H,
brs), 9.26-9.56 (2H, m).

25 [0455]

Example 83

4-((1R,2S)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-
methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide
dihydrochloride

30 The title compound could be produced according to the
production methods described in the present specification, a
method shown in the Examples, or a method analogous thereto.

[0456]

Example 84

35 4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-

thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

To a mixture of (1S,2R)-4-(2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride (3.36 g), THF (30 mL) and methanol (30 mL) was added triethylamine (3.31 mL), triethyl orthoformate (2.10 mL) and cyclobutanone (0.856 mL) at room temperature, and the mixture was stirred at room temperature overnight. Sodium borohydride (540 mg) was added under ice-cooling, and the reaction mixture was stirred at room temperature for 1 hr. The reaction mixture was diluted with ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the obtained residue was purified by silica gel column chromatography (ethyl acetate/methanol). To the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution, and the mixture was stirred overnight and concentrated under reduced pressure. The residue was crystallized from ethanol/water/ethyl acetate to give the title compound (2.17 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.18-1.29 (1H, m), 1.48-1.60 (1H, m), 1.72-1.89 (2H, m), 2.12-2.39 (4H, m), 2.54-2.67 (4H, m), 2.74-2.86 (1H, m), 3.74-3.92 (1H, m), 7.74 (1H, d, J = 1.14 Hz), 8.06 (1H, s), 9.79 (2H, brs).

[0457]

The compounds of Examples 85 to 92 could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

Example 85

4-((1R,2S)-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride

Example 86

N-(1-ethyl-1H-pyrazol-4-yl)-5-((1R,2R)-2-((tetrahydro-2H-

pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide dihydrochloride

Example 87

4-((1S,2R)-2-((cyclopropylmethyl) amino) cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl) thiophene-2-carboxamide dihydrochloride

Example 88

4-((1R,2S)-2-((cyclopropylmethyl) amino) cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl) thiophene-2-carboxamide dihydrochloride

Example 89

4-((1S,2R)-2-(cyclobutylamino) cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl) thiophene-2-carboxamide dihydrochloride

Example 90

4-((1R,2S)-2-(cyclobutylamino) cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl) thiophene-2-carboxamide dihydrochloride

Example 91

5-((1R,2R)-2-((1-acetylpiperidin-4-yl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl) thiophene-3-carboxamide hydrochloride

Example 92

5-((1R,2R)-2-((1-(cyclopropylcarbonyl) piperidin-4-yl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl) thiophene-3-carboxamide hydrochloride

[0458]

Example 93

N-(4,4-difluorocyclohexyl)-5-methyl-4-((1S,2R)-2-(tetrahydro-2H-pyran-4-ylamino) cyclopropyl) thiophene-2-carboxamide fumarate

A) 4-((1S,2R)-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)-5-methylthiophene-2-carboxamide hydrochloride

4-((1S,2R)-2-((tert-Butoxycarbonyl) amino) cyclopropyl)-5-methylthiophene-2-carboxylic acid (50 mg), 4,4-difluorocyclohexanamine (27.3 mg) and triethylamine (0.094 mL) were dissolved in DMF (1 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (77 mg) was added, and the mixture was stirred at room temperature overnight. Water was added to the reaction mixture, and the

mixture was extracted with ethyl acetate. The organic layer was successively washed with water and saturated brine, dried over magnesium sulfate, and concentrated under reduced pressure. The obtained residue (70.5 mg) and 4 mol/L hydrogen chloride/ethyl acetate solution (4.0 mL) were stirred at room temperature overnight. The precipitated solid was collected by filtration and washed with ethyl acetate to give the title compound (51 mg).

MS: $[M-HCl+H]^+$ 315.1.

10 [0459]

B) N-(4,4-difluorocyclohexyl)-5-methyl-4-((1S,2R)-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-2-carboxamide fumarate

To a mixture of 4-((1S,2R)-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)-5-methylthiophene-2-carboxamide hydrochloride (40.0 mg), borane-2-picoline complex (36.7 mg), methanol (2.0 mL) and acetic acid (0.2 mL) was added dihydro-2H-pyran-4(3H)-one (17.1 mg) at room temperature. The reaction mixture was stirred at room temperature for 18 hr, and saturated aqueous sodium hydrogen carbonate solution was added at 0°C. The mixture was extracted with ethyl acetate, and the extract was successively washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane), the obtained residue was dissolved in ethyl acetate, and a solution of fumaric acid (9.3 mg) in ethanol was added at room temperature. The reaction mixture was stirred at room temperature for 30 min, and the solvent was evaporated under reduced pressure to give the title compound (40.0 mg).

1H NMR (300 MHz, DMSO- d_6) δ 0.76-0.91 (1H, m), 0.95-1.04 (1H, m), 1.21-1.40 (2H, m), 1.48-1.66 (2H, m), 1.69-2.11 (9H, m), 2.26-2.34 (1H, m), 2.40 (3H, s), 2.75-2.87 (1H, m), 3.23-3.34 (2H, m), 3.76-3.96 (3H, m), 6.60 (2H, s), 7.21 (1H, s), 7.94-8.05 (1H, m).

[0460]

Example 94

N-(4,4-difluorocyclohexyl)-5-methyl-4-((1R,2S)-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-2-carboxamide

5 fumarate

The title compound could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

[0461]

10 Example 95

4-((1S,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide
fumarate

A) 4-((1S,2R)-2-aminocyclopropyl)-5-methyl-N-(5-methyl-1,3,4-
15 thiadiazol-2-yl)thiophene-2-carboxamide

4-((1S,2R)-2-((tert-Butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylic acid (150 mg), 5-methyl-1,3,4-thiadiazol-2-amine (69.7 mg) and triethylamine (0.281 mL) were dissolved in DMF (5 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-
20 tetramethyluronium hexafluorophosphate (230 mg) was added, and the mixture was stirred at room temperature overnight. Water was added to the reaction mixture, and the precipitate was collected by filtration to give solid A. The filtrate was extracted with ethyl acetate. The organic layer was
25 successively washed with water and saturated brine, dried over magnesium sulfate, and concentrated under reduced pressure to give solid B. The solid (197 mg) combined with the obtained solids A and B, and 4 mol/L hydrogen chloride/ethyl acetate solution (4 mL) were stirred at room temperature overnight.
30 The precipitated solid was collected by filtration, and washed with ethyl acetate to give the title compound (120 mg).

MS: [M-HCl+H]⁺ 295.1.

[0462]

B) 4-((1S,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-
35 methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-

carboxamide fumarate

To a mixture of 4-((1S,2R)-2-aminocyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide hydrochloride (40.0 mg), triethylamine (22.0 mg), THF (1.0 mL) and methanol (1.0 mL) was added cyclopropanecarbaldehyde (11.8 mg) at room temperature. The reaction mixture was stirred at 50°C for 1.5 hr, and stirred at room temperature for 1 hr. Sodium borohydride (8.2 mg) was added at 0°C, the reaction mixture was stirred at 0°C for 30 min, and ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture. The mixture was extracted with ethyl acetate/THF, and the extract was washed with saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (methanol/ethyl acetate), the obtained residue was suspended in ethyl acetate, and a solution of fumaric acid (8.4 mg) in ethanol was added at room temperature. The reaction mixture was stirred at room temperature for 30 min, and the solid was collected by filtration to give the title compound (28.0 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.10-0.21 (2H, m), 0.37-0.49 (2H, m), 0.85-1.00 (2H, m), 1.04-1.16 (1H, m), 1.80-1.92 (1H, m), 2.40-2.46 (1H, m), 2.48 (3H, s), 2.56-2.60 (2H, m), 2.61 (3H, s), 6.58 (2H, s), 7.76 (1H, s).

[0463]

Example 96

4-((1R,2S)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide fumarate

The title compound could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

[0464]

Example 97

5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-((1S,2R)-2-

((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide hydrochloride

To a mixture of 4-((1S,2R)-2-aminocyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide
5 hydrochloride (40.0 mg), triethylamine (22.0 mg), THF (1.0 mL) and methanol (1.0 mL) was added tetrahydro-2H-pyran-4-carbaldehyde (18.6 mg) at room temperature. The reaction mixture was stirred at room temperature overnight, sodium borohydride (8.2 mg) was added at 0°C, and the reaction mixture
10 was stirred at 0°C for 10 min. To the reaction mixture were added ethyl acetate and water, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column
15 chromatography (methanol/ethyl acetate), and to the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The reaction mixture was concentrated under reduced pressure, and the obtained residue was crystallized from ethanol/heptane to give the title compound (31.0 mg).

20 ¹H NMR (300 MHz, DMSO-d₆) δ 1.14-1.35 (4H, m), 1.52-1.63 (1H, m), 1.64-1.76 (2H, m), 1.86-2.06 (1H, m), 2.53 (3H, s), 2.62 (3H, s), 2.86-3.09 (3H, m), 3.21-3.30 (2H, m), 3.79-3.95 (2H, m), 7.70-8.03 (1H, m), 8.94-9.41 (2H, m), 12.34-12.97 (1H, m).
[0465]

25 Example 98

5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-((1R,2S)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide hydrochloride

The title compound could be produced according to the
30 production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.
[0466]

Example 99

4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(5-
35 methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide

hydrochloride

To a mixture of 4-((1S,2R)-2-aminocyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride (40.0 mg), triethylamine (22.0 mg), trimethyl orthoformate (17.3 mg), methanol (1.0 mL) and THF (1.0 mL) was added cyclobutanone (11.8 mg) at room temperature, and the mixture was stirred at room temperature for 3 hr. Sodium borohydride (8.2 mg) was added under ice-cooling, the reaction mixture was stirred for 30 min, and ethyl acetate and saturated aqueous sodium hydrogen carbonate solution were added. The mixture was extracted with ethyl acetate/THF, and the extract was washed with saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography (ethyl acetate/methanol), and to the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The reaction mixture was concentrated under reduced pressure, and the obtained residue was washed with ethyl acetate to give the title compound (18.0 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.12-1.24 (1H, m), 1.45-1.57 (1H, m), 1.74-1.92 (2H, m), 2.17-2.31 (4H, m), 2.34-2.46 (1H, m), 2.62 (3H, s), 2.71-2.86 (1H, m), 3.31 (3H, s), 3.76-3.94 (1H, m), 7.70-7.91 (1H, m), 9.06-9.71 (2H, m), 12.20-13.03 (1H, m).

[0467]

Example 100

4-((1R,2S)-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide hydrochloride

The title compound could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

[0468]

Example 101

4-((4-(((1R,2R)-2-(4-((4,4-difluorocyclohexyl)carbamoyl)-2-

thienyl)cyclopropyl)amino)methyl)piperidin-1-yl)methyl)benzoic acid dihydrochloride

The title compound could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

[0469]

Example 102

4-((1S,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide

10 dihydrochloride

To a mixture of 4-((1S,2R)-2-aminocyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide dihydrochloride (35.0 mg), sodium hydrogen carbonate (21.0 mg), THF (1.5 mL) and methanol (1.5 mL) was added

15 cyclopropanecarbaldehyde (8.99 μ L). Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, sodium borohydride (5.69 mg) was added under ice-cooling, and the reaction mixture was stirred at room temperature for 1 hr. The reaction mixture was
20 diluted with ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under
25 reduced pressure. The obtained residue was purified by silica gel column chromatography (ethyl acetate/methanol), and to the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The mixture was stirred overnight, and the precipitated solid was collected by filtration, and washed
30 with ethyl acetate to give the title compound (32.6 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.33-0.45 (2H, m), 0.50-0.66 (2H, m), 1.00-1.26 (2H, m), 1.50-1.65 (1H, m), 2.49 (3H, s), 2.82-3.05 (3H, m), 3.80 (3H, s), 7.49-7.58 (2H, m), 7.92 (1H, s), 9.41 (2H, brs), 10.34 (1H, s).

35 MS: [M-2HCl+H]⁺ 331.2.

[0470]

Example 103

4-((1R,2S)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide

5 dihydrochloride

The title compound could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

[0471]

10 Example 104

4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide dihydrochloride

To a mixture of 4-((1S,2R)-2-aminocyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide

15 dihydrochloride (35.0 mg), sodium hydrogen carbonate (21.0 mg), THF (1.5 mL) and methanol (1.5 mL) was added cyclobutanone (9.01 μ L). Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, sodium borohydride (5.69 mg) was added under ice-cooling, 20 and the reaction mixture was stirred at room temperature for 1 hr. The reaction mixture was diluted with ethyl acetate, water and saturated aqueous sodium hydrogen carbonate solution were added, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, 25 and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography (ethyl acetate/methanol), and to the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The mixture 30 was stirred overnight, and the precipitated solid was collected by filtration, and washed with ethyl acetate to give the title compound (29.6 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.10-1.24 (1H, m), 1.49-1.60 (1H, m), 1.72-1.89 (2H, m), 2.15-2.35 (4H, m), 2.39-2.48 (4H, m), 35 2.73-2.84 (1H, m), 3.76-3.91 (4H, m), 7.50-7.56 (2H, m), 7.92

(1H, s), 9.72 (2H, d, J = 3.79 Hz), 10.33 (1H, s).

[0472]

The compounds of Examples 105 to 107 could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

Example 105

4-((1R,2S)-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide dihydrochloride
10 [optical isomer, compound derived from methyl 4-(trans-2-((tert-butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylate (optical isomer, retention time long)]

Example 106

4-((1S,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide
15 hydrochloride

Example 107

4-((1R,2S)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide
20 hydrochloride

[0473]

Example 108

4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide
25 hydrochloride

A) tert-butyl ((1R,2S)-2-(2-methyl-5-((tetrahydro-2H-pyran-4-yl)carbamoyl)thiophen-3-yl)cyclopropyl)carbamate

4-((1S,2R)-2-((tert-Butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylic acid (100 mg), tetrahydro-2H-pyran-4-amine (0.042 mL) and triethylamine (0.117 mL) were
30 dissolved in DMF (5 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (153 mg) was added, and the mixture was stirred at room temperature overnight. Water was added to the reaction mixture, and the mixture was
35 extracted with ethyl acetate. The organic layer was

successively washed with water and saturated brine, dried over anhydrous sodium sulfate, and purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (114 mg).

5 MS: $[M-HCl+H]^+$ 381.3.

[0474]

B) 4-((1S,2R)-2-aminocyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide hydrochloride

To a solution of tert-butyl ((1R,2S)-2-(2-methyl-5-
10 ((tetrahydro-2H-pyran-4-yl)carbamoyl)thiophen-3-yl)cyclopropyl)carbamate (110 mg) in ethyl acetate (3 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (0.723 mL), and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration and washed
15 with ethyl acetate to give the title compound (92 mg).

MS: $[M-HCl+H]^+$ 281.2.

[0475]

C) 4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide
20 hydrochloride

To a mixture of 4-((1S,2R)-2-aminocyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide hydrochloride (35.0 mg), sodium hydrogen carbonate (13.9 mg), THF (1.5 mL) and methanol (1.5 mL) was added cyclobutanone
25 (9.94 μ L). Under a nitrogen atmosphere, the reaction mixture was stirred at 60°C for 1.5 hr and at room temperature for 30 min, sodium borohydride (6.27 mg) was added under ice-cooling, and the reaction mixture was stirred at room temperature for 1 hr. The reaction mixture was diluted with ethyl acetate, water
30 and saturated aqueous sodium hydrogen carbonate solution were added, and the mixture was extracted with ethyl acetate. The extract was successively washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The obtained residue was
35 purified by silica gel column chromatography (ethyl

acetate/methanol), and to the obtained fraction was added 4 mol/L hydrogen chloride/ethyl acetate solution. The mixture was stirred overnight, and the precipitated solid was collected by filtration and washed with ethyl acetate to give
5 the title compound (29.7 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.10-1.19 (1H, m), 1.43-1.59 (3H, m), 1.67-1.87 (4H, m), 2.14-2.30 (4H, m), 2.34-2.43 (1H, m), 2.44 (3H, s), 2.69-2.81 (1H, m), 3.35-3.40 (1H, m), 3.76-3.97 (4H, m), 7.30-7.39 (1H, m), 8.08 (1H, d, J = 7.95 Hz), 9.56
10 (2H, brs).

[0476]

Example 109

4-((1R,2S)-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide

15 hydrochloride

The title compound could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

[0477]

20 The compounds of Examples 110 and 111 could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

Example 110

25 4-(((1R,2R)-2-(4-((4,4-difluorocyclohexyl)carbamoyl)-2-thienyl)cyclopropyl)amino)methyl)benzoic acid

Example 111

5-((1R,2R)-2-(((5-amino-1,3,4-oxadiazol-2-yl)methyl)amino)cyclopropyl)-N-(4,4-

30 difluorocyclohexyl)thiophene-3-carboxamide

[0478]

Example 112

5-((1R,2R)-2-((cyclobutylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide

35 To a mixture of 5-((1R,2R)-2-aminocyclopropyl)-N-

(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide dihydrochloride (40 mg), triethylamine (0.055 mL), THF (1 mL) and methanol (1 mL) was added cyclobutanecarbaldehyde (13.33 mg) at room temperature. The reaction mixture was stirred at
5 room temperature for 5 min, and sodium borohydride (7.5 mg) was added at 0°C. The reaction mixture was stirred at room temperature for 10 min. To the reaction mixture were added water, ethyl acetate and saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl
10 acetate. The extract was washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/methanol), and to the obtained fraction was added 4 mol/L hydrogen
15 chloride/ethyl acetate solution. The reaction mixture was concentrated under reduced pressure, and the obtained residue was washed with ethyl acetate to give the title compound (31 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.21-1.35 (1H, m), 1.44-1.65 (3H, m), 1.82 (6H, d, J = 7.2 Hz), 1.98-2.14 (2H, m), 2.57-2.79 (2H, m), 2.88-2.98 (1H, m), 3.06-3.16 (2H, m), 3.35-3.42 (2H, m), 3.80-4.00 (3H, m), 7.28 (1H, s), 7.95 (1H, d, J = 1.5 Hz), 8.06 (1H, d, J = 7.6 Hz), 9.01-9.34 (2H, m).

[0479]

25 Example 113

5-((1R,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide malate

To a suspension of L-(-)-malic acid (50.2 mg, 0.37 mmol) in ethyl acetate (2 mL) was added a solution of 5-((1R,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide (120 mg, 0.37 mmol) in ethanol (1 mL) at 60°C. The reaction mixture was stirred at room temperature for 1 hr, and concentrated under reduced pressure. To the residue were added ethyl acetate (2 mL) and ethanol
35 (0.5 mL), and ethyl acetate (2 mL) was further added. The

precipitate was collected by filtration to give the title compound (106 mg). The obtained title compound (106 mg) was recrystallized from ethanol (0.3 mL) and ethyl acetate (1.2 mL) to give the title compound (65.0 mg).

5 ¹H NMR (300 MHz, DMSO-d₆) δ 0.12-0.25 (2H, m), 0.37-0.54 (2H, m), 0.82-1.12 (2H, m), 1.23 (1H, dt, J = 9.6, 4.9 Hz), 1.41-1.62 (2H, m), 1.72 (2H, dd, J = 12.9, 2.3 Hz), 2.14-2.29 (1H, m), 2.33-2.43 (1H, m), 2.52-2.73 (4H, m), 3.17-3.46 (2H, m), 3.81-4.00 (3H, m), 4.06 (1H, dd, J = 7.6, 6.1 Hz), 7.18 (1H, d, 10 J = 0.8 Hz), 7.84 (1H, d, J = 1.1 Hz), 7.99 (1H, d, J = 8.0 Hz).

[0480]

Example 114

5-((1R,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide succinate

To a solution of succinic acid (36.9 mg, 0.31 mmol) in ethanol (3 mL) was added 5-((1R,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide (100 mg, 0.31 mmol) at 60°C. The 20 reaction mixture was stirred at room temperature for 1 hr and concentrated under reduced pressure. To the residue were added ethyl acetate (2 mL) and ethanol (0.5 mL), and ethyl acetate (2 mL) was further added. The precipitate was collected by filtration to give the title compound (105 mg). The obtained 25 title compound (105 mg) was recrystallized from ethanol (0.8 mL) and ethyl acetate (1.2 mL) to give the title compound (62.0 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.06-0.19 (2H, m), 0.34-0.49 (2H, m), 0.78-1.15 (3H, m), 1.41-1.61 (2H, m), 1.64-1.78 (2H, m), 30 1.97-2.07 (1H, m), 2.32-2.43 (5H, m), 2.50-2.54 (2H, m), 3.30-3.52 (2H, m), 3.75-4.01 (3H, m), 7.13 (1H, d, J = 0.8 Hz), 7.79 (1H, d, J = 1.1 Hz), 7.97 (1H, d, J = 8.0 Hz).

[0481]

Example 115

35 5-((1R,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-

(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide fumarate

To a solution of fumaric acid (42.7 mg, 0.37 mmol) in ethanol (1 mL) was added a solution of 5-((1R,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide (118 mg, 0.37 mmol) in ethyl acetate (3 mL) at 60°C. To the reaction mixture was added ethyl acetate (2 mL), the mixture was stirred at room temperature, and the precipitate was collected by filtration to give the title compound (122 mg). The obtained title compound (122 mg) was recrystallized from ethanol (1.08 mL) and ethyl acetate (1.8 mL) to give the title compound (85 mg). ¹H NMR (300 MHz, DMSO-d₆) δ 0.06-0.17 (2H, m), 0.36-0.48 (2H, m), 0.82-1.18 (3H, m), 1.52 (2H, qd, J = 11.9, 4.4 Hz), 1.66-1.80 (2H, m), 2.00-2.12 (1H, m), 2.35-2.44 (1H, m), 2.50-2.57 (2H, m), 3.36 (2H, td, J = 11.5, 1.9 Hz), 3.75-4.01 (3H, m), 6.59 (2H, s), 7.13 (1H, s), 7.80 (1H, d, J = 1.5 Hz), 7.98 (1H, d, J = 8.0 Hz).

[0482]

Example 116

4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide phosphate

To a mixture of 4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide (100 mg) and ethanol (4 mL) was added 1 mol/l aqueous phosphoric acid solution (0.329 mL) at room temperature, and the mixture was stirred at room temperature for 3 hr. The precipitated solid was collected by filtration, and crystallized from ethanol/water/diisopropyl ether to give the title compound (97.0 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.90-1.12 (2H, m), 1.52-1.74 (2H, m), 1.77-1.94 (2H, m), 1.95-2.21 (3H, m), 2.32-2.39 (1H, m), 2.62 (3H, s), 3.34-3.50 (1H, m), 7.56 (1H, s), 7.98-8.04 (1H, m).

[0483]

Example 117

4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide succinate

To a mixture of 4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide (100 mg) and ethyl acetate (4 mL) was added a mixture of succinic acid (35.3 mg) and ethanol (1 mL) at 70°C, and the mixture was stirred at room temperature for 2 hr. The precipitated solid was collected by filtration, and crystallized from ethanol/water/diisopropyl ether to give the title compound (78.2 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.83-1.02 (2H, m), 1.49-1.82 (4H, m), 1.82-1.91 (1H, m), 2.03-2.19 (2H, m), 2.20-2.30 (1H, m), 2.40 (4H, s), 2.62 (3H, s), 3.31 (2H, dt, J = 15.52, 7.76 Hz), 7.51 (1H, d, J = 1.14 Hz), 7.96 (1H, d, J = 1.14 Hz), 10.00 (2H, brs).

[0484]

Example 118

4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide sulfate

A) 4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide

To a suspension of 4-((1S,2R)-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride (13.3 g), THF (120 mL) and methanol (120 mL) were added triethylamine (15.7 mL), triethyl orthoformate (8.32 mL) and cyclobutanone (3.39 mL) at room temperature, and the mixture was stirred at room temperature overnight. Sodium borohydride (2.14 g) was added under ice-cooling, and the reaction mixture was stirred at room temperature for 1 hr. To the reaction mixture was added saturated aqueous ammonium chloride solution at 0°C, and the mixture was stirred at 0°C for 30 min. The organic solvent was evaporated under reduced pressure. To the residue were added saturated aqueous sodium hydrogen carbonate solution and THF, and the mixture was extracted with ethyl acetate. The extract was successively

washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography (ethyl acetate/methanol) to give the
5 title compound (6.35 g).

MS: [M+H]⁺ 335.0.

[0485]

B) 4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide sulfate

10 To a mixture of 4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide (9.51 g) and THF (150 mL) was added 10% aqueous sulfuric acid solution (27.4 mL) at room temperature, and the mixture was stirred at room temperature
15 overnight. To the reaction mixture was added ethyl acetate (100 mL), and the mixture was further stirred for 1 hr. The precipitated solid was collected by filtration, and washed with ethyl acetate. The obtained solid (10.8 g) was dissolved in ethanol (100 mL) and water (65 mL) at 65°C, and the mixture
20 was stirred at 65°C for 30 min. To the reaction mixture was added dropwise ethyl acetate (250 mL). The reaction mixture was slowly cooled to room temperature, and stirred at room temperature overnight. The precipitated solid was collected by filtration, and washed with ethyl acetate to give the title
25 compound (9.46 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.21-1.32 (1H, m), 1.42 (1H, ddd, J = 10.22, 6.06, 4.54 Hz), 1.74-1.92 (2H, m), 2.06-2.31 (4H, m), 2.39-2.48 (1H, m), 2.63 (3H, s), 2.84 (1H, dt, J = 7.48, 4.02 Hz), 3.87 (1H, quin, J = 8.05 Hz), 7.74 (1H, d, J = 1.14 Hz),
30 8.04 (1H, d, J = 1.14 Hz).

[0486]

Example 119

4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide

35 To 4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(5-

methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride (650 mg) were added ethyl acetate and saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was
5 washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (523 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 0.82-1.00 (2H, m), 1.48-1.88 (5H, m), 2.01-2.17 (2H, m), 2.18-2.25 (1H, m), 2.62 (3H, s), 3.20-
10 3.41 (3H, m), 7.48 (1H, d, J = 1.14 Hz), 7.94 (1H, d, J = 1.14 Hz).

[0487]

Example 120

5-((1R,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-
15 (tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide

To 5-((1R,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride (600 mg) were added ethyl acetate and saturated aqueous sodium hydrogen carbonate solution, and the mixture
20 was extracted with ethyl acetate. The extract was washed with saturated brine, dried over magnesium sulfate, and concentrated under reduced pressure to give the title compound (520 mg).

¹H NMR (300 MHz, CDCl₃) δ 0.06-0.17 (2H, m), 0.42-0.55 (2H, m),
25 0.86-1.04 (2H, m), 1.10-1.22 (1H, m), 1.44-1.57 (2H, m), 1.91-2.13 (3H, m), 2.37-2.50 (1H, m), 2.53-2.66 (2H, m), 3.45-3.61 (2H, m), 3.93-4.05 (2H, m), 4.06-4.25 (1H, m), 5.62-5.77 (1H, m), 6.92-6.98 (1H, m), 7.52 (1H, d, J = 1.5 Hz).

[0488]

30 The compounds of Examples 121 and 122 could be produced according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

Example 121

35 4-((1S,2R)-2-((cyclobutylmethyl)amino)cyclopropyl)-5-methyl-N-

(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide
dihydrochloride

Example 122

4-((1S,2R)-2-((cyclobutylmethyl)amino)cyclopropyl)-5-methyl-N-
5 (tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide
hydrochloride
[0489]

Example 123

4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(1-
10 methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide hydrochloride
A) methyl 4-((1S,2R)-2-((tert-
butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-
carboxylate

Methyl 4-(trans-2-((tert-
15 butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-
carboxylate (8 g) was fractionated by SFC (column: CHIRALPAK
AD, 20 mmID×250 mmL, manufactured by Daicel Corporation,
mobile phase: carbon dioxide/methanol = 900/100), and the
obtained fraction was concentrated under reduced pressure to
20 give methyl 4-((1S,2R)-2-((tert-
butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-
carboxylate (optical isomer, retention time short) (3.68 g)
and methyl 4-((1R,2S)-2-((tert-
butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-
25 carboxylate (optical isomer, retention time long) (3.72 g).
methyl 4-((1S,2R)-2-((tert-butoxycarbonyl)amino)cyclopropyl)-
5-methylthiophene-2-carboxylate (optical isomer, retention
time short)

HPLC retention time 4.834 min (column: CHIRALPAK ADH, 4.6
30 mmID×150 mmL, mobile phase: carbon dioxide/methanol= 900/100,
flow rate: 2.5 mL/min, temperature: 35°C, detection: UV 220 nm,
concentration: 0.5 mg/mL, injection volume: 0.005 mL).

MS: [M-tBu+2H]⁺ 256.1.

methyl 4-((1R,2S)-2-((tert-butoxycarbonyl)amino)cyclopropyl)-
35 5-methylthiophene-2-carboxylate (optical isomer, retention

time long)

HPLC retention time 6.885 min (column: CHIRALPAK ADH, 4.6 mmIDx150 mmL, mobile phase: carbon dioxide/methanol= 900/100, flow rate: 2.5 mL/min, temperature: 35°C, detection: UV 220 nm, concentration: 0.5 mg/mL, injection volume: 0.005 mL).

MS: [M-tBu+2H]⁺ 256.1.

[0490]

B) 4-((1S,2R)-2-((tert-butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylic acid

10 Methyl 4-((1S,2R)-2-((tert-butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylate (3.2 g) was dissolved in methanol (20 mL) and tetrahydrofuran (10 mL), 2 mol/L aqueous sodium hydroxide solution (12.9 mL) was added at 0°C, and the mixture was
15 stirred at room temperature overnight. The reaction mixture was acidified with 1 mol/L hydrochloric acid, and extracted with ethyl acetate. The extract was washed with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound
20 (3.06 g).

MS: [M-H]⁻ 296.0.

[0491]

C) tert-butyl ((1R,2S)-2-(2-methyl-5-((1-methyl-1H-pyrazol-4-yl)carbamoyl)thiophen-3-yl)cyclopropyl)carbamate

25 4-((1S,2R)-2-((tert-Butoxycarbonyl)amino)cyclopropyl)-5-methylthiophene-2-carboxylic acid (3.06 g), 1-methyl-1H-pyrazol-4-amine hydrochloride (1.37 g) and triethylamine (5.73 mL) were dissolved in DMF (30 mL), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (4.69 g) was
30 added, and the mixture was stirred at 40°C overnight. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The extract was washed with saturated aqueous sodium hydrogen carbonate solution, water and saturated brine, dried over anhydrous sodium sulfate, and
35 concentrated under reduced pressure. The residue was purified

by silica gel column chromatography (hexane/ethyl acetate and methanol/ethyl acetate) to give the title compound (3.7 g).

MS: $[M+H]^+$ 377.1

[0492]

- 5 D) 4-((1S,2R)-2-aminocyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide dihydrochloride

To a suspension of tert-butyl ((1R,2S)-2-(2-methyl-5-((1-methyl-1H-pyrazol-4-yl)carbamoyl)thiophen-3-yl)cyclopropyl)carbamate (3.7 g) in ethyl acetate (45 mL) and
10 methanol (25 mL) was added 4 mol/L hydrogen chloride/ethyl acetate solution (49.1 mL) at 0°C, and the mixture was stirred at room temperature overnight. The precipitated solid was collected by filtration, and washed with ethyl acetate to give the title compound (3 g).

15 MS: $[M+H-2HCl]^+$ 277.1.

[0493]

- E) 4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide

To a suspension of 4-((1S,2R)-2-aminocyclopropyl)-5-
20 methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide dihydrochloride (3 g) in methanol (40 mL) were added triethylamine (2.99 mL, 21.5 mmol), trimethoxymethane (1.90 mL, 17.2 mmol) and cyclobutanone (0.77 mL, 10.3 mmol) at room temperature. The reaction mixture was stirred at room
25 temperature overnight, and sodium borohydride (487 mg) was added under cooling at -40°C. The reaction mixture was stirred at -40°C for 1 hr, saturated aqueous ammonium chloride solution was added, and the mixture was concentrated under reduced pressure. To the residue was added saturated aqueous sodium
30 hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (ethyl
35 acetate/methanol) to give the title compound (2 g).

MS: [M+H]⁺ 331.1.

[0494]

F) 4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide hydrochloride

5 4N Hydrogen chloride/ethyl acetate solution (6.08 mL, 24.33 mmol) was added to a solution of 4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide (2.68 g, 8.11 mmol) in ethyl acetate (20 mL) at 0°C. The precipitate was collected by
10 filtration, and washed with ethyl acetate to give a white solid. The obtained white solid was recrystallized from ethanol, water and ethyl acetate to give the title compound (1.920 g) as a white solid.

¹H NMR (300 MHz, DMSO-d₆) δ 1.10-1.26 (1H, m), 1.51-1.55 (1H, m), 1.66-1.94 (2H, m), 2.14-2.36 (4H, m), 2.39-2.50 (4H, m),
15 2.76-2.81 (1H, m), 3.70-3.93 (4H, m), 7.50-7.57 (2H, m), 7.92 (1H, s), 9.68 (2H, brs), 10.33 (1H, s).

[0495]

The compounds of Examples 124 and 125 could be produced
20 according to the production methods described in the present specification, a method shown in the Examples, or a method analogous thereto.

Example 124

4-((1S,2R)-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide dihydrochloride
25

Example 125

5-((1S,2S)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide hydrochloride

30 [0496]

Example compounds are shown in the following Tables. MS in the Tables shows measured values.

[0497]

[Table 1-1]

Ex. No.	IUPAC name	structure	salt	MS
1	4-(trans-2-aminocyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		2HCl	295.0
2	4-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		2HCl	349.1
3	5-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(4,4-difluorocyclohexyl)-1-methyl-1H-pyrazole-3-carboxamide		HCl	353.1
4	5-(trans-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)-1-methyl-1H-pyrazole-3-carboxamide		HCl	299.1
5	4-(trans-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)-2-naphthamide		2HCl	325.1
6	N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-((tetrahydro-2H-pyran-4-ylmethyl)amino)-cyclopropyl)-2-naphthamide		HCl	421.1

[0498]

[Table 1-2]

Ex. No.	IUPAC name	structure	salt	MS
7	4-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)-2-naphthamide		2HCl	379.2
8	N-(4,4-difluorocyclohexyl)-5-methyl-4-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-thiophene-2-carboxamide		HCl	413.2
9	7-(trans-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)-2,3-dihydro-1-benzofuran-5-carboxamide		2HCl	317.0
10	7-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)-2,3-dihydro-1-benzofuran-5-carboxamide		2HCl	371.1
11	7-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(4,4-difluorocyclohexyl)-2,3-dihydro-1-benzofuran-5-carboxamide		HCl	391.2
12	N-(4,4-difluorocyclohexyl)-7-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-2,3-dihydro-1-benzofuran-5-carboxamide		HCl	435.2

[0499]

[Table 1-3]

Ex. No.	IUPAC name	structure	salt	MS
13	5-(trans-2-aminocyclopropyl)-N-(4,4-difluorocyclohexyl)-6-methylnicotinamide		2HCl	310.2
14	5-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(4,4-difluorocyclohexyl)-6-methylnicotinamide		2HCl	364.2
15	3-(trans-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)-1-naphthamide		2HCl	325.1
16	N-(5-methyl-1,3,4-thiadiazol-2-yl)-3-((tetrahydro-2H-pyran-4-ylmethyl)amino)-cyclopropyl)-1-naphthamide		2HCl	423.2
17	4-(trans-2-(cyclobutylamino)-cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		1/2 fumarate	349.1
18	4-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(4,4-difluorocyclohexyl)-5-methylthiophene-2-carboxamide		fumarate	369.1

[0500]

[Table 1-4]

Ex. No.	IUPAC name	structure	salt	MS
19	4-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide		fuma-rate	335.1
20	5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-((trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-thiophene-2-carboxamide		2HCl	391.1
21	N-(4,4-difluorocyclohexyl)-5-methyl-4-(trans-2-(tetrahydro-2H-pyran-4-ylamino)-cyclopropyl)thiophene-2-carboxamide		fuma-rate	399.2
22	4-(trans-2-(cyclobutylamino)-cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	331.3
23	5-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide		2HCl	335.1
24	N-(4,4-difluorocyclohexyl)-5-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-thiophene-3-carboxamide		HCl	399.2

[0501]

[Table 1-5]

Ex. No.	IUPAC name	structure	salt	MS
25	4-(trans-2-aminocyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	277.1
26	4-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		fuma- rate	349.1
27	4-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		2HCl	335.1
28	4-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(4,4-difluorocyclohexyl)-thiophene-2-carboxamide		HCl	355.2
29	N-(4,4-difluorocyclohexyl)-4-(trans-2-(tetrahydro-2H-pyran-4-ylamino)-cyclopropyl)thiophene-2-carboxamide		HCl	385.1
30	5-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(4,4-difluorocyclohexyl)-thiophene-3-carboxamide		HCl	355.2







[0502]

[Table 1-6]

Ex. No.	IUPAC name	structure	salt	MS
31	N-(4,4-difluorocyclohexyl)-5-(trans-2-(tetrahydro-2H-pyran-4-ylamino)-cyclopropyl)thiophene-3-carboxamide		HCl	385.1
32	5-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide		HCl	321.1
33	5-(trans-2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide		2HCl	335.1
34	5-(trans-2-(cyclobutylamino)cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-3-carboxamide		HCl	317.2
35	4-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(4,4-difluorocyclohexyl)-thiazole-2-carboxamide		HCl	356.2
36	5-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(4,4-difluorocyclohexyl)-thiophene-2-carboxamide		HCl	355.1





[0503]

[Table 1-7]

Ex. No.	IUPAC name	structure	salt	MS
37	5-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		2HCl	335.1
38	N-cyclopentyl-5-((1R,2R)-2-(tetrahydro-2H-pyran-4-ylamino)-cyclopropyl)thiophene-3-carboxamide		HCl	335.1
39	N-cyclopentyl-5-((1S,2S)-2-(tetrahydro-2H-pyran-4-ylamino)-cyclopropyl)thiophene-3-carboxamide		HCl	335.1
40	N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-((1R,2R)-2-(tetrahydro-2H-pyran-4-ylmethyl)-amino)cyclopropyl)thiophene-3-carboxamide		2HCl	379.1
41	N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-((1S,2S)-2-(tetrahydro-2H-pyran-4-ylmethyl)-amino)cyclopropyl)thiophene-3-carboxamide		2HCl	379.1
42	5-((1R,2R)-2-(cyclobutylamino)-cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide		HCl	321.1







[0504]

[Table 1-8]

Ex. No.	IUPAC name	structure	salt	MS
43	5-((1S,2S)-2-(cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide		HCl	321.1
44	N-(4,4-difluorocyclohexyl)-5-((1R,2R)-2-(tetrahydro-2H-pyran-4-ylamino)-cyclopropyl)thiophene-3-carboxamide		HCl	385.0
45	5-((1R,2R)-2-(cis-(4-aminocyclohexyl)-amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-thiophene-3-carboxamide or 5-((1R,2R)-2-(trans-(4-aminocyclohexyl)-amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-thiophene-3-carboxamide		2HCl	398.1
46	5-((1R,2R)-2-(cis-(4-aminocyclohexyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-thiophene-3-carboxamide or 5-((1R,2R)-2-(trans-(4-aminocyclohexyl)-amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-thiophene-3-carboxamide		2HCl	398.1

[0505]

[Table 1-9]

Ex. No.	IUPAC name	structure	salt	MS
47	5-((1R,2R)-2-((4,4-difluorocyclohexyl)-amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide		HCl	385.0
48	5-((1R,2R)-2-((1-cyclopropylpiperidin-4-yl)amino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide		2HCl	403.9
49	N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-((1R,2R)-2-(tetrahydro-2H-pyran-4-ylamino)-cyclopropyl)thiophene-3-carboxamide		HCl	364.9
50	N-(2-methyl-1,3-thiazol-5-yl)-5-((1R,2R)-2-(tetrahydro-2H-pyran-4-ylamino)-cyclopropyl)thiophene-3-carboxamide		HCl	364.0
51	N-(4,4-difluorocyclohexyl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)-amino)cyclopropyl)-thiophene-3-carboxamide		HCl	399.0
52	5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)-amino)cyclopropyl)-N-(1-(2,2,2-trifluoroethyl)-piperidin-4-yl)thiophene-3-carboxamide		HCl	446.1

[0506]

[Table 1-10]

Ex. No.	IUPAC name	structure	salt	MS
53	N-(2-methyl-1,3-thiazol-5-yl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)-amino)cyclopropyl)-thiophene-3-carboxamide		2HCl	378.0
54	N-(1,3-dimethyl-1H-pyrazol-4-yl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)-amino)cyclopropyl)-thiophene-3-carboxamide		2HCl	375.0
55	N-(1,5-dimethyl-1H-pyrazol-4-yl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)-amino)cyclopropyl)-thiophene-3-carboxamide		2HCl	375.1
56	N-cyclopropyl-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)-amino)cyclopropyl)-thiophene-3-carboxamide		HCl	321.0
57	4-(trans-2-(cyclobutylamino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		HCl	335.0
58	N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-((trans-2-((tetrahydro-2H-pyran-4-ylmethyl)-amino)cyclopropyl)-thiophene-2-carboxamide		HCl	379.1

[0507]

[Table 1-11]

Ex. No.	IUPAC name	structure	salt	MS
59	4-(trans-2-((1-cyclopropyl-piperidin-4-yl)-amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		2HCl	404.2
60	3-(4-(((1R,2R)-2-(4-((5-methyl-1,3,4-thiadiazol-2-yl)-carbamoyl)-2-thienyl)cyclopropyl)amino)methyl)-piperidin-1-yl)propanoic acid		3HCl	450.0
61	4-(trans-2-(cyclobutylamino)-cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	317.1
62	4-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	317.0
63	4-(trans-2-(cyclobutylamino)-cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide		HCl	321.0
64	4-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide		HCl	321.0

[0508]

[Table 1-12]

Ex. No.	IUPAC name	structure	salt	MS
65	5-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(4,4-difluorocyclohexyl)-2-methylthiophene-3-carboxamide		HCl	369.1
66	N-(4,4-difluorocyclohexyl)-2-methyl-5-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)-amino)cyclopropyl)-thiophene-3-carboxamide		HCl	413.1
67	N-(4,4-difluorocyclohexyl)-2-methyl-5-(trans-2-(tetrahydro-2H-pyran-4-ylamino)-cyclopropyl)thiophene-3-carboxamide		HCl	399.0
68	5-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-2-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide		HCl	335.1
69	5-(trans-2-((cyclopropylmethyl)amino)-cyclopropyl)-2-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide		2HCl	349.0
70	2-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-(trans-2-((tetrahydro-2H-pyran-4-ylmethyl)-amino)cyclopropyl)-thiophene-3-carboxamide		2HCl	393.0

[0509]

[Table 1-13]

Ex. No.	IUPAC name	structure	salt	MS
71	5-((1R,2R)-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide		HCl	321.0
72	N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-((1S,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-thiophene-2-carboxamide		HCl	377.0
73	4-((1R,2S)-2-(cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide		HCl	321.0
74	N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-((1R,2S)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-thiophene-2-carboxamide		HCl	379.0
75	4-((1R,2S)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide		HCl	321.0
76	4-((1S,2R)-2-(cyclobutylamino)-cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide		HCl	321.0

[0510]

[Table 1-14]

Ex. No.	IUPAC name	structure	salt	MS
77	4-((1S,2R)-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide		HCl	321.0
78	4-((1S,2R)-2-((1-cyclopropylpiperidin-4-yl)-amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		2HCl	404.0
79	4-((1R,2S)-2-((1-cyclopropylpiperidin-4-yl)-amino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		2HCl	404.0
80	5-((1R,2R)-2-((1-cyclopropylpiperidin-4-yl)-amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide		HCl	424.0
81	N-(5-methyl-1,2-oxazol-3-yl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-3-carboxamide		HCl	362.0
82	4-((1S,2R)-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		2HCl	335.0






[0511]

[Table 1-15]

Ex. No.	IUPAC name	structure	salt	MS
83	4-((1R,2S)-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		2HCl	335.0
84	4-((1S,2R)-2-(cyclobutylamino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		2HCl	335.0
85	4-((1R,2S)-2-(cyclobutylamino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		2HCl	335.0
86	N-(1-ethyl-1H-pyrazol-4-yl)-5-((1R,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-thiophene-3-carboxamide		2HCl	375.1
87	4-((1S,2R)-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	317.1
88	4-((1R,2S)-2-((cyclopropylmethyl)amino)-cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	317.1

[0512]

[Table 1-16]

Ex. No.	IUPAC name	structure	salt	MS
89	4-((1S,2R)-2-(cyclobutylamino)cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	317.1
90	4-((1R,2S)-2-(cyclobutylamino)cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	317.0
91	5-((1R,2R)-2-((1-acetylpiperidin-4-yl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-thiophene-3-carboxamide		HCl	426.1
92	5-((1R,2R)-2-((1-(cyclopropylcarbonyl)piperidin-4-yl)amino)-cyclopropyl)-N-(4,4-difluorocyclohexyl)-thiophene-3-carboxamide		HCl	452.1
93	N-(4,4-difluorocyclohexyl)-5-methyl-4-((1S,2R)-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-2-carboxamide		fuma- rate	399.0





[0513]

[Table 1-17]

Ex. No.	IUPAC name	structure	salt	MS
94	N-(4,4-difluorocyclohexyl)-5-methyl-4-((1R,2S)-2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-2-carboxamide		fuma-rate	399.0
95	4-((1S,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		fuma-rate	349.0
96	4-((1R,2S)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		fuma-rate	349.0
97	5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-((1S,2R)-2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-thiophene-2-carboxamide		HCl	391.0





[0514]

[Table 1-18]

Ex. No.	IUPAC name	structure	salt	MS
98	5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-((1R,2S)-2-((tetrahydro-2H-pyran-4-ylmethyl)-amino)cyclopropyl)-thiophene-2-carboxamide		HCl	391.0
99	4-((1S,2R)-2-(cyclobutylamino)-cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		HCl	349.0
100	4-((1R,2S)-2-(cyclobutylamino)-cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		HCl	349.0
101	4-((4-(((1R,2R)-2-(4-((4,4-difluoro-cyclohexyl)-carbamoyl)-2-thienyl)cyclopropyl)amino)methyl)piperidin-1-yl)methyl)benzoic acid		2HCl	532.1





[0515]

[Table 1-19]

Ex. No.	IUPAC name	structure	salt	MS
102	4-((1S,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	331.1
103	4-((1R,2S)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	331.1
104	4-((1S,2R)-2-(cyclobutylamino)-cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	331.1
105	4-((1R,2S)-2-(cyclobutylamino)-cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	331.1







[0516]

[Table 1-20]

Ex. No.	IUPAC name	structure	salt	MS
106	4-((1S,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide		HCl	335.0
107	4-((1R,2S)-2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide		HCl	335.0
108	4-((1S,2R)-2-(cyclobutylamino)-cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide		HCl	335.0
109	4-((1R,2S)-2-(cyclobutylamino)-cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide		HCl	335.0

[0517]

[Table 1-21]

Ex. No.	IUPAC name	structure	salt	MS
110	4-((((1R,2R)-2-(4-((4,4-difluorocyclohexyl)-carbamoyl)-2-thienyl)cyclopropyl)amino)methyl)benzoic acid		HCl	435.0
111	5-((1R,2R)-2-(((5-amino-1,3,4-oxadiazol-2-yl)-methyl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-thiophene-3-carboxamide		2HCl	398.0
112	5-((1R,2R)-2-(((cyclobutylmethyl)-amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide		HCl	335.1
113	5-((1R,2R)-2-(((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide		L-malate	321.2
114	5-((1R,2R)-2-(((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide		succinate	321.2
115	5-((1R,2R)-2-(((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide		fumarate	321.1

[0518]

[Table 1-22]

Ex. No.	IUPAC name	structure	salt	MS
116	4-((1S,2R)-2-(cyclobutylamino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		H ₃ PO ₄	335.1
117	4-((1S,2R)-2-(cyclobutylamino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		succinate	335.1
118	4-((1S,2R)-2-(cyclobutylamino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		H ₂ SO ₄	335.1
119	4-((1S,2R)-2-(cyclobutylamino)-cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide			335.0
120	5-((1R,2R)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide			321.1

[0519]

[Table 1-23]

Ex. No.	IUPAC name	structure	salt	MS
121	4-((1S,2R)-2-((cyclobutylmethyl)-amino)cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		2HCl	345.1
122	4-((1S,2R)-2-((cyclobutylmethyl)-amino)cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide		HCl	349.1
123	4-((1S,2R)-2-(cyclobutylamino)-cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide		HCl	331.2
124	4-((1S,2R)-2-aminocyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide		2HCl	281.1
125	5-((1S,2S)-2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide		HCl	321.2

[0520]

Other preferable specific examples of the compound represented by the formula (I) include the following compounds shown in the Examples, an optically active form thereof, and a mixture of optically active forms thereof. The following compounds shown in the Examples, an optically active form thereof, and a mixture of optically active forms thereof can be produced according to the aforementioned production methods, a method shown in the Examples, or a method analogous thereto.

10 The relative configuration of the substituent on the cyclopropane ring is cis or trans, preferably trans.

[0521]

Example A1

4-(2-(cyclobutylamino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)thiophene-2-carboxamide

15

Example A2

4-(2-(cyclobutylamino)cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl)thiophene-2-carboxamide

Example A3

4-(2-(cyclobutylamino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)thiophene-2-carboxamide

20

Example A4

4-(2-(cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide

25 Example A5

4-(2-(cyclobutylamino)cyclopropyl)-N-cyclopentylthiophene-2-carboxamide

Example A6

4-(2-(cyclobutylamino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-2-carboxamide

30

Example A7

4-(2-(cyclobutylamino)cyclopropyl)-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-2-carboxamide

Example A8

35 4-(2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,2-oxazol-3-

yl)thiophene-2-carboxamide

Example A9

4-(2-(cyclobutylamino)cyclopropyl)-N-(3-methyl-1,2-oxazol-5-yl)thiophene-2-carboxamide

5 Example A10

4-(2-(cyclobutylamino)cyclopropyl)-N-(2-methyl-1,3-thiazol-5-yl)thiophene-2-carboxamide

[0522]

Example A11

10 4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide

Example A12

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)thiophene-2-carboxamide

15 Example A13

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl)thiophene-2-carboxamide

Example A14

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)thiophene-2-carboxamide

Example A15

N-cyclopentyl-4-(2-((cyclopropylmethyl)amino)cyclopropyl)thiophene-2-carboxamide

Example A16

25 4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-2-carboxamide

Example A17

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,2-oxazol-3-yl)thiophene-2-carboxamide

30 Example A18

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(3-methyl-1,2-oxazol-5-yl)thiophene-2-carboxamide

Example A19

35 4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(2-methyl-1,3-thiazol-5-yl)thiophene-2-carboxamide

Example A20

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(oxetan-3-yl)thiophene-2-carboxamide
[0523]

5 Example A21

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(3,3-difluorocyclobutyl)thiophene-2-carboxamide

Example A22

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,3,4-oxadiazol-2-yl)thiophene-2-carboxamide
10

Example A23

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-2-thienyl)thiophene-2-carboxamide

Example A24

15 4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(pyridin-4-yl)thiophene-2-carboxamide

Example A25

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(2-methyl-1,3-oxazol-5-yl)thiophene-2-carboxamide

20 Example A26

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-(trifluoromethyl)-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide

Example A27

25 4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(3-methyloxetan-3-yl)thiophene-2-carboxamide

Example A28

N-cyclopropyl-4-(2-((cyclopropylmethyl)amino)cyclopropyl)thiophene-2-carboxamide

30 Example A29

N-cyclopentyl-4-(2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-2-carboxamide

Example A30

N-(2-methyl-1,3-thiazol-5-yl)-4-(2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-2-carboxamide
35

[0524]

Example A31

4-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl) thiophene-2-carboxamide

5 Example A32

4-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl) thiophene-2-carboxamide

Example A33

4-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-N-(1,3-
10 dimethyl-1H-pyrazol-5-yl) thiophene-2-carboxamide

Example A34

N-cyclopentyl-4-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl) thiophene-2-carboxamide

Example A35

15 4-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl) thiophene-2-carboxamide

Example A36

4-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-N-(1-(2,2,2-trifluoroethyl) piperidin-4-yl) thiophene-2-carboxamide

20 Example A37

4-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-N-(5-methyl-1,2-oxazol-3-yl) thiophene-2-carboxamide

Example A38

4-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-N-(3-
25 methyl-1,2-oxazol-5-yl) thiophene-2-carboxamide

Example A39

4-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-N-(2-methyl-1,3-thiazol-5-yl) thiophene-2-carboxamide

Example A40

30 N-(1,3-dimethyl-1H-pyrazol-4-yl)-4-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-2-carboxamide

[0525]

Example A41

N-(1-ethyl-1H-pyrazol-4-yl)-4-(2-((tetrahydro-2H-pyran-4-
35 ylmethyl) amino) cyclopropyl) thiophene-2-carboxamide

Example A42

N-(1,3-dimethyl-1H-pyrazol-5-yl)-4-(2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide

Example A43

5 N-cyclopentyl-4-(2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide

Example A44

4-(2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-2-carboxamide

10 Example A45

N-(5-methyl-1,2-oxazol-3-yl)-4-(2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide

Example A46

15 N-(3-methyl-1,2-oxazol-5-yl)-4-(2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide

Example A47

N-(2-methyl-1,3-thiazol-5-yl)-4-(2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide

Example A48

20 N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

Example A49

25 N-(1-methyl-1H-pyrazol-4-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

Example A50

30 N-(1-ethyl-1H-pyrazol-4-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

[0526]

Example A51

35 N-(1,3-dimethyl-1H-pyrazol-5-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

Example A52

N-cyclopentyl-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

Example A53

5 N-(4,4-difluorocyclohexyl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

Example A54

10 N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

Example A55

15 N-(5-methyl-1,2-oxazol-3-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

Example A56

N-(3-methyl-1,2-oxazol-5-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

20 Example A57

N-(2-methyl-1,3-thiazol-5-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

Example A58

25 4-(2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide

Example A59

4-(2-(cyclobutylamino)cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide

30 Example A60

N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-(2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide

[0527]

Example A61

35 N-(4,4-difluorocyclohexyl)-4-(2-((tetrahydro-2H-pyran-4-

ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide

Example A62

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide

5 [0528]

Example B1

4-(2-(cyclobutylamino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)-5-methylthiophene-2-carboxamide

Example B2

10 4-(2-(cyclobutylamino)cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl)-5-methylthiophene-2-carboxamide

Example B3

4-(2-(cyclobutylamino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)-5-methylthiophene-2-carboxamide

15 Example B4

4-(2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide

Example B5

20 4-(2-(cyclobutylamino)cyclopropyl)-N-cyclopentyl-5-methylthiophene-2-carboxamide

Example B6

4-(2-(cyclobutylamino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-5-methylthiophene-2-carboxamide

Example B7

25 4-(2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-2-carboxamide

Example B8

4-(2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(5-methyl-1,2-oxazol-3-yl)thiophene-2-carboxamide

30 Example B9

4-(2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(3-methyl-1,2-oxazol-5-yl)thiophene-2-carboxamide

Example B10

35 4-(2-(cyclobutylamino)cyclopropyl)-5-methyl-N-(2-methyl-1,3-thiazol-5-yl)thiophene-2-carboxamide

[0529]

Example B11

4-(2-((cyclopropylmethyl) amino) cyclopropyl)-5-methyl-N-(1-methyl-1H-pyrazol-4-yl) thiophene-2-carboxamide

5 Example B12

4-(2-((cyclopropylmethyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)-5-methylthiophene-2-carboxamide

Example B13

4-(2-((cyclopropylmethyl) amino) cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl)-5-methylthiophene-2-carboxamide

Example B14

4-(2-((cyclopropylmethyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)-5-methylthiophene-2-carboxamide

Example B15

15 N-cyclopentyl-4-(2-((cyclopropylmethyl) amino) cyclopropyl)-5-methylthiophene-2-carboxamide

Example B16

4-(2-((cyclopropylmethyl) amino) cyclopropyl)-5-methyl-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl) thiophene-2-carboxamide

20 Example B17

4-(2-((cyclopropylmethyl) amino) cyclopropyl)-5-methyl-N-(5-methyl-1,2-oxazol-3-yl) thiophene-2-carboxamide

Example B18

25 4-(2-((cyclopropylmethyl) amino) cyclopropyl)-5-methyl-N-(3-methyl-1,2-oxazol-5-yl) thiophene-2-carboxamide

Example B19

4-(2-((cyclopropylmethyl) amino) cyclopropyl)-5-methyl-N-(2-methyl-1,3-thiazol-5-yl) thiophene-2-carboxamide

Example B20

30 4-(2-((cyclopropylmethyl) amino) cyclopropyl)-5-methyl-N-(oxetan-3-yl) thiophene-2-carboxamide

[0530]

Example B21

4-(2-((cyclopropylmethyl) amino) cyclopropyl)-N-(3,3-difluorocyclobutyl)-5-methylthiophene-2-carboxamide

Example B22

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-oxadiazol-2-yl)thiophene-2-carboxamide

Example B23

5 4-(2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(5-methyl-2-thienyl)thiophene-2-carboxamide

Example B24

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(pyridin-4-yl)thiophene-2-carboxamide

10 Example B25

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(2-methyl-1,3-oxazol-5-yl)thiophene-2-carboxamide

Example B26

15 4-(2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(5-(trifluoromethyl)-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide

Example B27

4-(2-((cyclopropylmethyl)amino)cyclopropyl)-5-methyl-N-(3-methyloxetan-3-yl)thiophene-2-carboxamide

20 Example B28

N-cyclopropyl-4-(2-((cyclopropylmethyl)amino)cyclopropyl)-5-methylthiophene-2-carboxamide

Example B29

25 N-cyclopentyl-5-methyl-4-(2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-2-carboxamide

Example B30

5-methyl-N-(2-methyl-1,3-thiazol-5-yl)-4-(2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-2-carboxamide

[0531]

30 Example B31

4-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide

Example B32

35 4-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(1-

ethyl-1H-pyrazol-4-yl)-5-methylthiophene-2-carboxamide

Example B33

4-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)-5-methylthiophene-2-carboxamide

5 Example B34

N-cyclopentyl-4-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-5-methylthiophene-2-carboxamide

Example B35

4-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-5-methylthiophene-2-carboxamide

Example B36

4-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-5-methyl-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-2-carboxamide

15 Example B37

4-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-5-methyl-N-(5-methyl-1,2-oxazol-3-yl)thiophene-2-carboxamide

Example B38

4-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-5-methyl-N-(3-methyl-1,2-oxazol-5-yl)thiophene-2-carboxamide

Example B39

4-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-5-methyl-N-(2-methyl-1,3-thiazol-5-yl)thiophene-2-carboxamide

Example B40

25 N-(1,3-dimethyl-1H-pyrazol-4-yl)-5-methyl-4-(2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide
[0532]

Example B41

N-(1-ethyl-1H-pyrazol-4-yl)-5-methyl-4-(2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide

Example B42

N-(1,3-dimethyl-1H-pyrazol-5-yl)-5-methyl-4-(2-((tetrahydro-2H-pyran-4-ylmethyl)amino)cyclopropyl)thiophene-2-carboxamide

Example B43

35 N-cyclopentyl-5-methyl-4-(2-((tetrahydro-2H-pyran-4-

ylmethyl) amino) cyclopropyl) thiophene-2-carboxamide

Example B44

5-methyl-4-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl)-N-(1-(2,2,2-

5 trifluoroethyl)piperidin-4-yl) thiophene-2-carboxamide

Example B45

5-methyl-N-(5-methyl-1,2-oxazol-3-yl)-4-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-2-carboxamide

Example B46

10 5-methyl-N-(3-methyl-1,2-oxazol-5-yl)-4-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-2-carboxamide

Example B47

5-methyl-N-(2-methyl-1,3-thiazol-5-yl)-4-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-2-carboxamide

15 Example B48

5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-2-carboxamide

Example B49

20 5-methyl-N-(1-methyl-1H-pyrazol-4-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-2-carboxamide

Example B50

N-(1-ethyl-1H-pyrazol-4-yl)-5-methyl-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-2-carboxamide

[0533]

Example B51

N-(1,3-dimethyl-1H-pyrazol-5-yl)-5-methyl-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-2-carboxamide

Example B52

N-cyclopentyl-5-methyl-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-2-carboxamide

Example B53

N-(4,4-difluorocyclohexyl)-5-methyl-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

5 Example B54

5-methyl-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

Example B55

10 5-methyl-N-(5-methyl-1,2-oxazol-3-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

Example B56

15 5-methyl-N-(3-methyl-1,2-oxazol-5-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

Example B57

20 5-methyl-N-(2-methyl-1,3-thiazol-5-yl)-4-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-2-carboxamide

[0534]

Example C1

5-(2-(cyclobutylamino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)thiophene-3-carboxamide

25 Example C2

5-(2-(cyclobutylamino)cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl)thiophene-3-carboxamide

Example C3

30 5-(2-(cyclobutylamino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)thiophene-3-carboxamide

Example C4

5-(2-(cyclobutylamino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide

Example C5

35 5-(2-(cyclobutylamino)cyclopropyl)-N-cyclopentylthiophene-3-

carboxamide

Example C6

5-(2-(cyclobutylamino)cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide

5 Example C7

5-(2-(cyclobutylamino)cyclopropyl)-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-3-carboxamide

Example C8

5-(2-(cyclobutylamino)cyclopropyl)-N-(5-methyl-1,2-oxazol-3-yl)thiophene-3-carboxamide

Example C9

5-(2-(cyclobutylamino)cyclopropyl)-N-(3-methyl-1,2-oxazol-5-yl)thiophene-3-carboxamide

Example C10

15 5-(2-(cyclobutylamino)cyclopropyl)-N-(2-methyl-1,3-thiazol-5-yl)thiophene-3-carboxamide

[0535]

Example C11

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-3-carboxamide

Example C12

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)thiophene-3-carboxamide

Example C13

25 5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl)thiophene-3-carboxamide

Example C14

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)thiophene-3-carboxamide

30 Example C15

N-cyclopentyl-5-(2-((cyclopropylmethyl)amino)cyclopropyl)thiophene-3-carboxamide

Example C16

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-3-carboxamide

Example C17

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,2-oxazol-3-yl)thiophene-3-carboxamide

Example C18

5 5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(3-methyl-1,2-oxazol-5-yl)thiophene-3-carboxamide

Example C19

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(2-methyl-1,3-thiazol-5-yl)thiophene-3-carboxamide

10 Example C20

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(oxetan-3-yl)thiophene-3-carboxamide

[0536]

Example C21

15 5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(3,3-difluorocyclobutyl)thiophene-3-carboxamide

Example C22

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-1,3,4-oxadiazol-2-yl)thiophene-3-carboxamide

20 Example C23

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-methyl-2-thienyl)thiophene-3-carboxamide

Example C24

25 5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(pyridin-4-yl)thiophene-3-carboxamide

Example C25

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(2-methyl-1,3-oxazol-5-yl)thiophene-3-carboxamide

Example C26

30 5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(5-(trifluoromethyl)-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide

Example C27

35 5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(3-methyloxetan-3-yl)thiophene-3-carboxamide

Example C28

N-cyclopropyl-5-(2-
((cyclopropylmethyl)amino)cyclopropyl)thiophene-3-carboxamide

Example C29

5 N-cyclopentyl-5-(2-(tetrahydro-2H-pyran-4-
ylamino)cyclopropyl)thiophene-3-carboxamide

Example C30

N-(2-methyl-1,3-thiazol-5-yl)-5-(2-(tetrahydro-2H-pyran-4-
ylamino)cyclopropyl)thiophene-3-carboxamide

10 [0537]

Example C31

5-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(5-
methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide

Example C32

15 5-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(1-
ethyl-1H-pyrazol-4-yl)thiophene-3-carboxamide

Example C33

5-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(1,3-
dimethyl-1H-pyrazol-5-yl)thiophene-3-carboxamide

20 Example C34

N-cyclopentyl-5-(2-((1-cyclopropylpiperidin-4-
yl)amino)cyclopropyl)thiophene-3-carboxamide

Example C35

25 5-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(4,4-
difluorocyclohexyl)thiophene-3-carboxamide

Example C36

5-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(1-
(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-3-carboxamide

Example C37

30 5-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(5-
methyl-1,2-oxazol-3-yl)thiophene-3-carboxamide

Example C38

5-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(3-
methyl-1,2-oxazol-5-yl)thiophene-3-carboxamide

35 Example C39

5-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-N-(2-methyl-1,3-thiazol-5-yl) thiophene-3-carboxamide

Example C40

N-(1,3-dimethyl-1H-pyrazol-4-yl)-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide

[0538]

Example C41

N-(1-ethyl-1H-pyrazol-4-yl)-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide

10 Example C42

N-(1,3-dimethyl-1H-pyrazol-5-yl)-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide

Example C43

N-cyclopentyl-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide

Example C44

5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl)-N-(1-(2,2,2-trifluoroethyl) piperidin-4-yl) thiophene-3-carboxamide

Example C45

20 N-(5-methyl-1,2-oxazol-3-yl)-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide

Example C46

N-(3-methyl-1,2-oxazol-5-yl)-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide

25 Example C47

N-(2-methyl-1,3-thiazol-5-yl)-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide

Example C48

30 N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-(2-((1-(2,2,2-trifluoroethyl) piperidin-4-yl) amino) cyclopropyl) thiophene-3-carboxamide

Example C49

N-(1-methyl-1H-pyrazol-4-yl)-5-(2-((1-(2,2,2-trifluoroethyl) piperidin-4-yl) amino) cyclopropyl) thiophene-3-carboxamide

Example C50

N-(1-ethyl-1H-pyrazol-4-yl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-3-carboxamide

5 [0539]

Example C51

N-(1,3-dimethyl-1H-pyrazol-5-yl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-3-carboxamide

10 Example C52

N-cyclopentyl-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-3-carboxamide

Example C53

15 N-(4,4-difluorocyclohexyl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-3-carboxamide

Example C54

20 N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-3-carboxamide

Example C55

N-(5-methyl-1,2-oxazol-3-yl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-3-carboxamide

25 Example C56

N-(3-methyl-1,2-oxazol-5-yl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-3-carboxamide

Example C57

30 N-(2-methyl-1,3-thiazol-5-yl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)amino)cyclopropyl)thiophene-3-carboxamide

[0540]

Example D1

35 5-(2-(cyclobutylamino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-

4-yl)-2-methylthiophene-3-carboxamide

Example D2

5-(2-(cyclobutylamino)cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl)-2-methylthiophene-3-carboxamide

5 Example D3

5-(2-(cyclobutylamino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)-2-methylthiophene-3-carboxamide

Example D4

10 5-(2-(cyclobutylamino)cyclopropyl)-2-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide

Example D5

5-(2-(cyclobutylamino)cyclopropyl)-N-cyclopentyl-2-methylthiophene-3-carboxamide

Example D6

15 5-(2-(cyclobutylamino)cyclopropyl)-N-(4,4-difluorocyclohexyl)-2-methylthiophene-3-carboxamide

Example D7

5-(2-(cyclobutylamino)cyclopropyl)-2-methyl-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-3-carboxamide

20 Example D8

5-(2-(cyclobutylamino)cyclopropyl)-2-methyl-N-(5-methyl-1,2-oxazol-3-yl)thiophene-3-carboxamide

Example D9

25 5-(2-(cyclobutylamino)cyclopropyl)-2-methyl-N-(3-methyl-1,2-oxazol-5-yl)thiophene-3-carboxamide

Example D10

5-(2-(cyclobutylamino)cyclopropyl)-2-methyl-N-(2-methyl-1,3-thiazol-5-yl)thiophene-3-carboxamide

[0541]

30 Example D11

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-3-carboxamide

Example D12

35 5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)-2-methylthiophene-3-carboxamide

Example D13

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl)-2-methylthiophene-3-carboxamide

Example D14

5 5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)-2-methylthiophene-3-carboxamide

Example D15

N-cyclopentyl-5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methylthiophene-3-carboxamide

10 Example D16

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-3-carboxamide

Example D17

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(5-methyl-1,2-oxazol-3-yl)thiophene-3-carboxamide

Example D18

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(3-methyl-1,2-oxazol-5-yl)thiophene-3-carboxamide

Example D19

20 5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(2-methyl-1,3-thiazol-5-yl)thiophene-3-carboxamide

Example D20

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(oxetan-3-yl)thiophene-3-carboxamide

25 [0542]

Example D21

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-N-(3,3-difluorocyclobutyl)-2-methylthiophene-3-carboxamide

Example D22

30 5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(5-methyl-1,3,4-oxadiazol-2-yl)thiophene-3-carboxamide

Example D23

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(5-methyl-2-thienyl)thiophene-3-carboxamide

35 Example D24

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(pyridin-4-yl)thiophene-3-carboxamide

Example D25

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(2-methyl-1,3-oxazol-5-yl)thiophene-3-carboxamide

Example D26

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(5-(trifluoromethyl)-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide

10 Example D27

5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methyl-N-(3-methyloxetan-3-yl)thiophene-3-carboxamide

Example D28

15 N-cyclopropyl-5-(2-((cyclopropylmethyl)amino)cyclopropyl)-2-methylthiophene-3-carboxamide

Example D29

N-cyclopentyl-2-methyl-5-(2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide

Example D30

20 2-methyl-N-(2-methyl-1,3-thiazol-5-yl)-5-(2-(tetrahydro-2H-pyran-4-ylamino)cyclopropyl)thiophene-3-carboxamide
[0543]

Example D31

25 5-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-2-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)thiophene-3-carboxamide

Example D32

5-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl)-2-methylthiophene-3-carboxamide

30 Example D33

5-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)-2-methylthiophene-3-carboxamide

Example D34

35 N-cyclopentyl-5-(2-((1-cyclopropylpiperidin-4-yl)amino)cyclopropyl)-2-methylthiophene-3-carboxamide

Example D35

5-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl)-2-methylthiophene-3-carboxamide

Example D36

5 5-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-2-methyl-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-3-carboxamide

Example D37

5-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-2-methyl-N-(5-methyl-1,2-oxazol-3-yl)thiophene-3-carboxamide
10

Example D38

5-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-2-methyl-N-(3-methyl-1,2-oxazol-5-yl)thiophene-3-carboxamide

Example D39

15 5-(2-((1-cyclopropylpiperidin-4-yl) amino) cyclopropyl)-2-methyl-N-(2-methyl-1,3-thiazol-5-yl)thiophene-3-carboxamide

Example D40

N-(1,3-dimethyl-1H-pyrazol-4-yl)-2-methyl-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl)thiophene-3-carboxamide
20 [0544]

Example D41

N-(1-ethyl-1H-pyrazol-4-yl)-2-methyl-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl)thiophene-3-carboxamide

Example D42

25 N-(1,3-dimethyl-1H-pyrazol-5-yl)-2-methyl-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl)thiophene-3-carboxamide

Example D43

N-cyclopentyl-2-methyl-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl)thiophene-3-carboxamide

30 Example D44

2-methyl-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl)-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)thiophene-3-carboxamide

Example D45

35 2-methyl-N-(5-methyl-1,2-oxazol-3-yl)-5-(2-((tetrahydro-2H-

pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide

Example D46

2-methyl-N-(3-methyl-1,2-oxazol-5-yl)-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide

5 Example D47

2-methyl-N-(2-methyl-1,3-thiazol-5-yl)-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide

Example D48

2-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-3-carboxamide

Example D49

2-methyl-N-(1-methyl-1H-pyrazol-4-yl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-3-carboxamide

Example D50

N-(1-ethyl-1H-pyrazol-4-yl)-2-methyl-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-3-carboxamide

20 [0545]

Example D51

N-(1,3-dimethyl-1H-pyrazol-5-yl)-2-methyl-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-3-carboxamide

25 Example D52

N-cyclopentyl-2-methyl-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-3-carboxamide

Example D53

30 N-(4,4-difluorocyclohexyl)-2-methyl-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-3-carboxamide

Example D54

2-methyl-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-

yl) amino) cyclopropyl) thiophene-3-carboxamide

Example D55

2-methyl-N-(5-methyl-1,2-oxazol-3-yl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-3-

5 carboxamide

Example D56

2-methyl-N-(3-methyl-1,2-oxazol-5-yl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-3-carboxamide

10 Example D57

2-methyl-N-(2-methyl-1,3-thiazol-5-yl)-5-(2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl) amino) cyclopropyl) thiophene-3-carboxamide

Example D58

15 5-(2-(cyclobutylamino) cyclopropyl)-2-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl) thiophene-3-carboxamide

Example D59

5-(2-((cyclopropylmethyl) amino) cyclopropyl)-2-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl) thiophene-3-carboxamide

20 Example D60

2-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl)-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide
[0546]

Example D61

25 5-(2-((cyclopropylmethyl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl)-2-methylthiophene-3-carboxamide

Example D62

N-(4,4-difluorocyclohexyl)-2-methyl-5-(2-(tetrahydro-2H-pyran-4-ylamino) cyclopropyl) thiophene-3-carboxamide

30 Example D63

N-(4,4-difluorocyclohexyl)-2-methyl-5-(2-((tetrahydro-2H-pyran-4-ylmethyl) amino) cyclopropyl) thiophene-3-carboxamide

Example D64

5-(2-((cyclopropylmethyl) amino) cyclopropyl)-2-methyl-N-
35 (tetrahydro-2H-pyran-4-yl) thiophene-3-carboxamide

[0547]

Example E1

4-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl) thiophene-2-carboxamide

5 Example E2

4-(2-((3-aminocyclobutyl) amino) cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl) thiophene-2-carboxamide

Example E3

5-(2-((3-aminocyclobutyl) amino) cyclopropyl)-2-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl) thiophene-3-carboxamide

Example E4

5-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl) thiophene-3-carboxamide

Example E5

15 4-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl) thiophene-2-carboxamide

Example E6

4-(2-((4-aminocyclohexyl) amino) cyclopropyl)-5-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl) thiophene-2-carboxamide

20 Example E7

5-(2-((4-aminocyclohexyl) amino) cyclopropyl)-2-methyl-N-(5-methyl-1,3,4-thiadiazol-2-yl) thiophene-3-carboxamide

Example E8

5-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(5-methyl-1,3,4-thiadiazol-2-yl) thiophene-3-carboxamide

Example E9

4-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl) thiophene-2-carboxamide

Example E10

30 4-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl)-5-methylthiophene-2-carboxamide

[0548]

Example E11

5-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(1-ethyl-1H-pyrazol-4-yl)-2-methylthiophene-3-carboxamide

Example E12

5- (2- ((3-aminocyclobutyl) amino) cyclopropyl) -N- (1-ethyl-1H-pyrazol-4-yl) thiophene-3-carboxamide

Example E13

5 4- (2- ((4-aminocyclohexyl) amino) cyclopropyl) -N- (1-ethyl-1H-pyrazol-4-yl) thiophene-2-carboxamide

Example E14

4- (2- ((4-aminocyclohexyl) amino) cyclopropyl) -N- (1-ethyl-1H-pyrazol-4-yl) -5-methylthiophene-2-carboxamide

10 Example E15

5- (2- ((4-aminocyclohexyl) amino) cyclopropyl) -N- (1-ethyl-1H-pyrazol-4-yl) -2-methylthiophene-3-carboxamide

Example E16

15 5- (2- ((4-aminocyclohexyl) amino) cyclopropyl) -N- (1-ethyl-1H-pyrazol-4-yl) thiophene-3-carboxamide

Example E17

4- (2- ((3-aminocyclobutyl) amino) cyclopropyl) -N- (1-methyl-1H-pyrazol-4-yl) thiophene-2-carboxamide

Example E18

20 4- (2- ((3-aminocyclobutyl) amino) cyclopropyl) -5-methyl-N- (1-methyl-1H-pyrazol-4-yl) thiophene-2-carboxamide

Example E19

5- (2- ((3-aminocyclobutyl) amino) cyclopropyl) -2-methyl-N- (1-methyl-1H-pyrazol-4-yl) thiophene-3-carboxamide

25 Example E20

5- (2- ((3-aminocyclobutyl) amino) cyclopropyl) -N- (1-methyl-1H-pyrazol-4-yl) thiophene-3-carboxamide

[0549]

Example E21

30 4- (2- ((4-aminocyclohexyl) amino) cyclopropyl) -N- (1-methyl-1H-pyrazol-4-yl) thiophene-2-carboxamide

Example E22

4- (2- ((4-aminocyclohexyl) amino) cyclopropyl) -5-methyl-N- (1-methyl-1H-pyrazol-4-yl) thiophene-2-carboxamide

35 Example E23

5-(2-((4-aminocyclohexyl) amino) cyclopropyl)-2-methyl-N-(1-methyl-1H-pyrazol-4-yl)thiophene-3-carboxamide

Example E24

5-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(1-methyl-1H-pyrazol-4-yl)thiophene-3-carboxamide

Example E25

4-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)thiophene-2-carboxamide

Example E26

4-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)-5-methylthiophene-2-carboxamide

Example E27

5-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)-2-methylthiophene-3-carboxamide

Example E28

5-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)thiophene-3-carboxamide

Example E29

4-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)thiophene-2-carboxamide

Example E30

4-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)-5-methylthiophene-2-carboxamide

[0550]

Example E31

5-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)-2-methylthiophene-3-carboxamide

Example E32

5-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-4-yl)thiophene-3-carboxamide

Example E33

4-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)thiophene-2-carboxamide

Example E34

4-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(1,3-dimethyl-

1H-pyrazol-5-yl)-5-methylthiophene-2-carboxamide

Example E35

5-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)-2-methylthiophene-3-carboxamide

5 Example E36

5-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)thiophene-3-carboxamide

Example E37

4-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)thiophene-2-carboxamide

Example E38

4-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)-5-methylthiophene-2-carboxamide

Example E39

15 5-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)-2-methylthiophene-3-carboxamide

Example E40

5-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(1,3-dimethyl-1H-pyrazol-5-yl)thiophene-3-carboxamide

20 [0551]

Example E41

4-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-2-carboxamide

Example E42

25 4-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl)-5-methylthiophene-2-carboxamide

Example E43

5-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl)-2-methylthiophene-3-carboxamide

30 Example E44

5-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-3-carboxamide

Example E45

35 4-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl)thiophene-2-carboxamide

Example E46

4-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl)-5-methylthiophene-2-carboxamide

Example E47

5 5-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl)-2-methylthiophene-3-carboxamide

Example E48

5-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(4,4-difluorocyclohexyl) thiophene-3-carboxamide

10 Example E49

4-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(5-methyl-1,2-oxazol-3-yl) thiophene-2-carboxamide

Example E50

4-(2-((3-aminocyclobutyl) amino) cyclopropyl)-5-methyl-N-(5-methyl-1,2-oxazol-3-yl) thiophene-2-carboxamide

[0552]

Example E51

5-(2-((3-aminocyclobutyl) amino) cyclopropyl)-2-methyl-N-(5-methyl-1,2-oxazol-3-yl) thiophene-3-carboxamide

20 Example E52

5-(2-((3-aminocyclobutyl) amino) cyclopropyl)-N-(5-methyl-1,2-oxazol-3-yl) thiophene-3-carboxamide

Example E53

4-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(5-methyl-1,2-oxazol-3-yl) thiophene-2-carboxamide

Example E54

4-(2-((4-aminocyclohexyl) amino) cyclopropyl)-5-methyl-N-(5-methyl-1,2-oxazol-3-yl) thiophene-2-carboxamide

Example E55

30 5-(2-((4-aminocyclohexyl) amino) cyclopropyl)-2-methyl-N-(5-methyl-1,2-oxazol-3-yl) thiophene-3-carboxamide

Example E56

5-(2-((4-aminocyclohexyl) amino) cyclopropyl)-N-(5-methyl-1,2-oxazol-3-yl) thiophene-3-carboxamide

35 Example E57

4-(2-((3-aminocyclobutyl)amino)cyclopropyl)-N-(2-methyl-1,3-thiazol-5-yl)thiophene-2-carboxamide

Example E58

4-(2-((3-aminocyclobutyl)amino)cyclopropyl)-5-methyl-N-(2-methyl-1,3-thiazol-5-yl)thiophene-2-carboxamide

Example E59

5-(2-((3-aminocyclobutyl)amino)cyclopropyl)-2-methyl-N-(2-methyl-1,3-thiazol-5-yl)thiophene-3-carboxamide

Example E60

5-(2-((3-aminocyclobutyl)amino)cyclopropyl)-N-(2-methyl-1,3-thiazol-5-yl)thiophene-3-carboxamide

[0553]

Example E61

4-(2-((4-aminocyclohexyl)amino)cyclopropyl)-N-(2-methyl-1,3-thiazol-5-yl)thiophene-2-carboxamide

Example E62

4-(2-((4-aminocyclohexyl)amino)cyclopropyl)-5-methyl-N-(2-methyl-1,3-thiazol-5-yl)thiophene-2-carboxamide

Example E63

5-(2-((4-aminocyclohexyl)amino)cyclopropyl)-2-methyl-N-(2-methyl-1,3-thiazol-5-yl)thiophene-3-carboxamide

Example E64

5-(2-((4-aminocyclohexyl)amino)cyclopropyl)-N-(2-methyl-1,3-thiazol-5-yl)thiophene-3-carboxamide

Example E65

4-(2-((4,4-difluorocyclohexyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide

Example E66

4-(2-((4,4-difluorocyclohexyl)amino)cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide

Example E67

5-(2-((4,4-difluorocyclohexyl)amino)cyclopropyl)-2-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide

Example E68

5-(2-((4,4-difluorocyclohexyl)amino)cyclopropyl)-N-

(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide

Example E69

4-(2-((4,4-difluorocyclohexyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide

5 Example E70

4-(2-((4,4-difluorocyclohexyl)amino)cyclopropyl)-5-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-2-carboxamide

[0554]

Example E71

10 5-(2-((4,4-difluorocyclohexyl)amino)cyclopropyl)-2-methyl-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide

Example E72

5-(2-((4,4-difluorocyclohexyl)amino)cyclopropyl)-N-(tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide

15 [0555]

Experimental Example 1

The genetic engineering method described below was performed according to the method described in a book (Maniatis et al., Molecular Cloning, Cold Spring Harbor Laboratory, 1989) or the method described in the protocol attached to the reagent.

(1) Construction of GST-tagged expression vector having TEV Protease cleavage sequence

A GST-tagged expression vector having TEV Protease cleavage sequence was constructed by successive 2 times of PCR method. Firstly, PCR was performed using pGEX6P1 (GE Healthcare) as a template, two primers

GST-Sw-F:

5'-AGAATCATTTAAATGGTGATCATGTAACCCATCCT-3'

30 [SEQ ID NO: 1]

GST-Tv-R1:

5'-CGCCCTGAAAGTACAGGTTCTCATCCGATTTTGGAGGATGGTCG-3'

[SEQ ID NO: 2]

and PrimeStar GXL DNA Polymerase (Takara Bio Inc.). Template DNA 0.5 µL, PrimeStar GXL DNA Polymerase Buffer 10 µL, 2.5 mM

dNTP solution 4 μ L, 10 μ M primer solution each 1.5 μ L, PrimeStar GXL DNA Polymerase 1 μ L, and sterilized distilled water 31.5 μ L were mixed. After a treatment at 98°C for 1 min, the PCR was started with 35 repeats of reactions at 98°C for 10
5 seconds, at 65°C for 5 seconds, and at 72°C for 25 seconds, followed by a reaction at 72°C for 1 min. Then, PCR was performed using the obtained PCR product as a template, two primers

GST-Sw-F:

10 5'-AGAATCATTTAAATGGTGATCATGTAACCCATCCT-3'

[SEQ ID NO: 1]

GST-Tv-R2:

5'-ATAATAGGATCCGCCCTGAAAGTACAGGTTCTC-3'

[SEQ ID NO: 3]

15 and PrimeStar GXL DNA Polymerase. Template DNA 0.5 μ L, PrimeStar GXL DNA Polymerase Buffer 10 μ L, 2.5 mM dNTP solution 4 μ L, 10 μ M primer solution each 1.5 μ L, PrimeStar GXL DNA Polymerase 1 μ L, and sterilized distilled water 31.5 μ L were mixed. After a treatment at 98°C for 1 min, the PCR
20 was started with 25 repeats of reactions at 98°C for 10 seconds, at 65°C for 5 seconds, and at 72°C for 25 seconds, followed by a reaction at 72°C for 1 min. The obtained PCR product was electrophoresed on agarose gel (1%), and an about 0.3 kbp DNA fragment containing a part of the GST gene was recovered from
25 the gel. The recovered DNA fragment was cleaved with restriction enzymes Swa I (New England Biolabs) and Bam HI (Takara Bio Inc.), and inserted into the Swa I/Bam HI site of pGEX6P1 to construct an expression vector pGEX7V1.

[0556]

30 (2) Cloning of human LSD1 (AOF2) cDNA

Human LSD1 cDNA was cloned by PCR method using brain cDNA Library (Takara Bio Inc.) as a template, two primers hLSD1-NheI-ko-F:

5'-TATTATGCTAGCGCCACCATGTTATCTGGGAAGAAGGCGGCAGC-3'

35 [SEQ ID NO: 4]

hLSD1-St-NotI-R:

5'-TATTATGCGGCCGCTCACATGCTTGGGGACTGCTGTGC-3'

[SEQ ID NO: 5]

and Pyrobest DNA Polymerase (Takara Bio Inc.). Template DNA
5 0.5 µL, Pyrobest DNA Polymerase Buffer 5 µL, 2.5 mM dNTP
solution 4 µL, 10 µM primer solution each 2.5 µL, Pyrobest DNA
Polymerase 0.5 µL, and sterilized distilled water 35 µL were
mixed. After a reaction at 98°C for 1 min, the PCR was started
with 35 repeats of reactions at 98°C for 10 seconds, at 68°C
10 for 5 seconds, and at 72°C for 2.5 min, followed by a reaction
at 72°C for 1 min. The obtained PCR product was
electrophoresed on agarose gel (1%), and an about 2.5 kbp DNA
fragment containing the human LSD1 cDNA was recovered from the
gel. The recovered DNA fragment was cleaved with restriction
15 enzymes Nhe I and Not I (Takara Bio Inc.), and inserted into
the Nhe I/Not I site of pcDNA3.1(+) (Invitrogen) to construct
an expression plasmid pcDNA3.1/hLSD1.

[0557]

(3) Construction of expression plasmid for human LSD1(172-833)
20 in Escherichia coli

A plasmid for expression of human LSD1(172-833) in
Escherichia coli was constructed by PCR method using
pcDNA3.1/hLSD1 as a template, two primers
hLSD1-172aa-Bgl2-F:

25 5'-ATAATAAGATCTTCGGGTGTGGAGGGCGCAGCTT-3'

[SEQ ID NO: 6]

hLSD1-833aa-St-NotI-R:

5'-ATAATAGCGGCCGCCATGGCCCCCAAACTGGTCTGCA-3'

[SEQ ID NO: 7]

30 and PrimeStar MAX DNA Polymerase (Takara Bio Inc.). Template
DNA 1 µL, PrimeStar MAX DNA Polymerase Enzyme PreMix 25 µL, 10
µM primer solution each 1.5 µL, and sterilized distilled water
21 µL were mixed. After a reaction at 98°C for 1 min, the PCR
was started with 25 repeats of reactions at 98°C for 10 seconds
35 and at 68°C for 8 seconds, followed by a reaction at 72°C for 1

min. The obtained PCR product was electrophoresed on agarose gel (1%), and an about 2 kbp DNA fragment containing human LSD1(172-833) cDNA was recovered from the gel. The recovered DNA fragment was cleaved with restriction enzymes Bgl II and Not I (Takara Bio Inc.), and inserted into the Bam HI/Not I site of pGEX7V1 to construct an expression plasmid pGEX7V1/GST-hLSD1(172-833).

[0558]

(4) Preparation of LSD1

10 Escherichia coli C43(DE3) pLysS was transformed with the expression plasmid pGEX7V1/GST-hLSD1(172-833). The obtained recombinant Escherichia coli was inoculated in a TB medium (1.2% tryptone, 2.4% yeast extract, 0.4% glycerol, 17 mM potassium dihydrogen phosphate and 72 mM dipotassium hydrogen 15 phosphate) added with 100 mg/L ampicillin and 35 mg/L chloramphenicol, and cultured at 37°C. When the turbidity reached 500 Klett units, the culture temperature was changed to 16°C, IPTG (Isopropyl β -D-1-thiogalactopyranoside) having a final concentration of 0.5 mM was added to induce expression, 20 and the cells were cultured further for 14 hr. The culture medium was centrifuged at 6,000 g for 15 min, and Escherichia coli pellets were recovered.

[0559]

Escherichia coli pellets for 12 L of the culture medium 25 were suspended in 1000 mL of PBS (Immuno-Biological Laboratories Co., Ltd.), 0.15 M NaCl, 5%(V/V) Glycerol (Buffer A), and 5000 units Benzonase (Merck), 1000 mg Lysozyme, and 10 tablets of Protease Inhibitor (Roche) were added. Using Branson ultrasonic disintegrator, the suspension was disrupted 30 by ultrasonication for 3 min, and centrifuged at 33,000 g for 60 min, and the supernatant was recovered. The supernatant was applied to two GSTrap 4B 5 mL columns (GE Healthcare) equilibrated in advance with 0.1 M Tris (pH 8.0), 0.15 M NaCl, 5%(V/V) Glycerol (Buffer B), and the columns were each washed 35 with 30 mL of Buffer B. GST-hLSD1(172-833) was eluted from

each column with Buffer B added with 13 mL of GSH with the final concentration 20 mM, applied to two HiLoad 26/60 Superdex 200 pg columns (GE Healthcare) equilibrated in advance with Buffer B, and eluted with 380 mL of Buffer B.

5 Total 60 mL of GST-hLSD1(172-833)-containing fraction was diluted 5-fold with 20 mM Tris (pH 8.0) (Buffer C), applied to Mono Q 10/100 GL column (GE Healthcare) equilibrated in advance with Buffer C, and 0-500 mM NaCl gradient elution was performed to give purified GST-hLSD1(172-833). 3.4 mg of His-
10 TEV protease was added to about 34 mg of GST-hLSD1(172-833), and the mixture was treated with 50 mM Tris (pH 8.0), 0.5 mM EDTA, 1 mM DTT at 4°C for 16 hr to cleave the GST tag. The reaction mixture after the cleavage reaction was applied to two series-coupled columns with Ni-NTA Superflow Cartridges 1
15 mL (QIAGEN), and GSTrap 4B 5 mL column (GE Healthcare) equilibrated in advance with Buffer A added with Imidazole at a final concentration of 20 mM, and a flow-through fraction containing hLSD1(172-833) free of GST-tag was recovered. It was concentrated to 10 mL with AmiconUltra 15 (NWC0 30K)
20 (Millipore Japan), and purified with HiLoad 26/60 Superdex 200 pg column (GE Healthcare) equilibrated with Buffer A to give hLSD1 purified product (8.4 mg). The protein concentration of hLSD1 was measured by BCA Protein Assay Kit (Thermo Fisher Scientific K.K.) using bovine serum albumin as the standard.

25 [0560]

(5) Measurement of LSD1 inhibitory activity

A test compound dissolved in DMSO was added by to a reaction solution (50 mM Tris-HCl (pH 8.0), 0.1% BSA, 1 mM DTT) containing LSD1 enzyme, and the mixture was reacted at
30 room temperature for 60 min. Biotin-histone H3 mono methylated K4 peptide solution (NH₂-ART(me-
K)QTARKSTGGKAPRKQLAGGK(Biotin)-CONH₂) was added to start the reaction. After reaction at room temperature for 5 min, 2-PCPA solution was added to terminate the reaction. A detection
35 solution (800 mM potassium fluoride, 0.1% BSA) containing

europium-labeled anti-histone H3 antibody (Wako Pure Chemical Industries, Ltd.) and Streptavidin-XL665 (Cisbio) was further added, and the mixture was left standing for 60 min. A time-resolved fluorescence (excitation 320 nm, emission 615 nm, 665 nm) was measured by Envision (PerkinElmer). The LSD1 inhibitory rate (%) of the test compound was calculated by the following formula.

$$\text{inhibitory rate (\%)} = (1 - (\text{test compound count} - \text{blank}) \div (\text{control} - \text{blank})) \times 100$$

10 The count of the LSD1 enzyme reaction mixture under compound non-addition conditions is indicated as control, and the count under compound non-addition and LSD1 enzyme non-addition conditions is indicated as blank. A concentration necessary for achieving 50% inhibitory rate was taken as IC₅₀ value. The results are shown in Table 2.

[0561]

Experimental Example 2

(1) Measurement of MAO-A inhibitory activity

20 The MAO-A inhibitory activity evaluation described below followed the protocol of MAO-Glo (registered trademark) Assay of Promega KK.

A test compound dissolved in DMSO was added to a reaction solution (100 mM HEPES (pH 7.5), 5% glycerol) containing MAO-A enzyme (Sigma-Aldrich Co. LLC.), and the mixture was reacted at room temperature for 15 min. MAO substrate (Promega KK) was added to start the reaction. After reaction at room temperature for 60 min, Luciferine detection reagent (Promega KK) was added to terminate the reaction. After reaction at room temperature for 20 min with stirring, the luminescence was measured by Envision (PerkinElmer). The MAO-A inhibitory rate (%) of the test compound was calculated by the following formula.

$$\text{inhibitory rate (\%)} = (1 - (\text{test compound count} - \text{blank}) \div (\text{control} - \text{blank})) \times 100$$

35 The count of the MAO-A enzyme reaction mixture under

compound non-addition conditions is indicated as control, and the count under compound non-addition and MAO-A enzyme non-addition conditions is indicated as blank. A concentration necessary for achieving 50% inhibitory rate was taken as IC₅₀ value. The results are shown in Table 2.

[0562]

(2) Measurement of MAO-B inhibitory activity

The MAO-B inhibitory activity evaluation described below followed the protocol of MAO-Glo (registered trademark) Assay of Promega KK.

A test compound dissolved in DMSO was added to a reaction solution (100 mM HEPES (pH 7.5), 5% glycerol, 10% DMSO) containing MAO-B enzyme (Sigma-Aldrich Co. LLC.), and the mixture was reacted at room temperature for 15 min. MAO substrate (Promega KK) was added to start the reaction. After reaction at room temperature for 60 min, Luciferine detection reagent (Promega KK) (50 µL) was added to terminate the reaction. After reaction at room temperature for 20 min with stirring, the luminescence was measured by Envision (PerkinElmer). The MAO-B inhibitory rate (%) of the test compound was calculated by the following formula.

$$\text{inhibitory rate (\%)} = (1 - (\text{test compound count} - \text{blank}) \div (\text{control} - \text{blank})) \times 100$$

The count of the MAO-B enzyme reaction mixture under compound non-addition conditions is indicated as control, and the count under compound non-addition and MAO-B enzyme non-addition conditions is indicated as blank. A concentration necessary for achieving 50% inhibitory rate was taken as IC₅₀ value. The results are shown in Table 2.

30

[0563]

[Table 2-1]

Example No.	LSD1 IC ₅₀ value (μM)	MAO-A IC ₅₀ value (μM)	MAO-B IC ₅₀ value (μM)
1	1.3	>100	>100
2	0.33	>100	>100
3	0.29	>100	>100
4	1.7	>100	>100
5	4.6	>100	>100
6	0.11	>100	>100
7	0.26	82	>100
8	0.25	>100	>100
9	3.0	>100	>100
10	0.23	>100	>100
11	0.18	>100	>100
12	<0.1	>100	>100
13	12	>100	93
14	2.4	>100	28
15	0.73	89	66
16	<0.1	>100	>100
17	0.22	>100	>100
18	0.26	>100	>100
19	0.65	>100	>100
20	<0.1	>100	>100
21	0.23	>100	88
22	0.14	>100	>100
23	<0.1	>100	>100
24	<0.1	>100	>100
25	3.0	>100	>100
26	0.50	>100	>100
27	0.33	>100	>100
28	<0.1	>100	>100
29	0.40	>100	>100
30	<0.1	>100	>100

[0564]

[Table 2-2]

Example No.	LSD1 IC ₅₀ value (μM)	MAO-A IC ₅₀ value (μM)	MAO-B IC ₅₀ value (μM)
31	<0.1	>100	>100
32	<0.1	>100	>100
33	<0.1	>100	>100
34	<0.1	>100	>100
35	0.68	>100	>100
36	3.8	22	6.0
37	99	85	90
38	<0.1	>100	>100
39	<0.1	23	>100
40	<0.1	>100	>100
41	<0.1	67	>100
42	<0.1	>100	>100
43	0.13	>100	>100
44	<0.1	>100	100
45	<0.1	>100	>100
46	<0.1	>100	>100
47	<0.1	>100	>100
48	<0.1	31	>100
49	<0.1	>100	>100
50	<0.1	>100	>100
51	<0.1	>100	>100
52	<0.1	>100	>100
53	<0.1	61	96
54	<0.1	>100	>100
55	<0.1	>100	>100
56	<0.1	>100	>100
57	0.44	>100	>100
58	<0.1	>100	>100
59	<0.1	>100	>100
60	<0.1	>100	>100

[0565]

[Table 2-3]

Example No.	LSD1 IC ₅₀ value (μM)	MAO-A IC ₅₀ value (μM)	MAO-B IC ₅₀ value (μM)
61	<0.1	>100	>100
62	<0.1	>100	>100
63	0.32	>100	>100
64	0.30	>100	>100
65	<0.1	25	>100
66	<0.1	98	>100
67	0.10	76	>100
68	0.14	>100	>100
69	<0.1	33	>100
70	<0.1	10	23
71	<0.1	>100	>100
72	<0.1	>100	>100
73	0.58	>100	>100
74	<0.1	>100	>100
75	0.34	>100	>100
76	0.20	>100	>100
77	0.36	>100	>100
78	<0.1	>100	>100
79	0.27	>100	>100
80	<0.1	>100	>100
81	<0.1	36	>100
82	0.24	>100	>100
83	0.28	>100	>100
84	0.23	>100	>100
85	0.81	>100	>100
86	<0.1	87	>100
87	0.18	>100	>100
88	<0.1	>100	>100
89	<0.1	>100	>100
90	0.10	>100	>100

[0566]

[Table 2-4]

Example No.	LSD1 IC ₅₀ value (μM)	MAO-A IC ₅₀ value (μM)	MAO-B IC ₅₀ value (μM)
91	<0.1	>100	>100
92	<0.1	>100	>100
93	0.16	>100	>100
94	0.21	>100	>100
95	0.18	>100	>100
96	0.59	>100	>100
97	<0.1	>100	>100
98	0.27	>100	>100
99	<0.1	>100	>100
100	0.82	>100	>100
101	<0.1	>100	>100
102	0.21	>100	>100
103	0.15	>100	>100
104	<0.1	>100	>100
105	0.21	>100	>100
106	0.44	>100	>100
107	1.40	>100	>100
108	0.34	>100	>100
109	2.30	>100	>100
110	<0.1	49	>100
111	0.10	61	>100
112	<0.1	>100	>100
113	<0.1	>100	>100
114	<0.1	>100	>100
115	<0.1	>100	>100
116	0.29	>100	>100
117	0.26	>100	>100
118	0.24	>100	>100
119	0.19	>100	>100
120	<0.1	>100	>100

[0567]

[Table 2-5]

Example No.	LSD1 IC ₅₀ value (μM)	MAO-A IC ₅₀ value (μM)	MAO-B IC ₅₀ value (μM)
121	0.12	>100	>100
122	0.17	>100	>100
123	0.10	>100	>100
124	1.90	>100	>100
125	<0.1	>100	>100

[0568]

5 As shown in Table 2, the compound of the present invention has a superior LSD1 inhibitory activity. In addition, the MAO-A inhibitory activity and MAO-B inhibitory activity of the compound of the present invention are low, and the compound of the present invention has a selective LSD1
10 inhibitory activity.

[0569]

Experimental Example 3

Gad1 H3K4 methylation induction activity in rat primary culture neurons

15 Experimental method

Hippocampus and cerebral cortex were isolated from fetal SD rat at embryonic day 19, a cell suspension was prepared using Nerve Cell Dissociation Medium (SUMITOMO BAKELITE, MS-0006L), and plated on a poly L-lysine-coated 6-well plate
20 (SUMITOMO BAKELITE, MS-0006L) at a density of 900000 cells/well. Under the conditions of 37°C and 5% CO₂, the cells were cultured in a neurobasal medium (Invitrogen, #211103049) containing B27 supplement (Invitrogen, #17504044, 1:50 dilution), 2 mM L-glutamine (Lonza, #B76053), 100 U/mL
25 penicillin/100 μg/mL streptomycin (Lonza, #17-602E), and 20 μg/mL gentamicin sulfate (Lonza, #17-519Z) for 10 days.

[0570]

Thereafter, the compound was added to final

concentrations of 10 μ M, the cells were further cultured for 3 days, and chromatin immunoprecipitation was performed.

Chromatin immunoprecipitation was performed by using ChIP-IT Express Enzymatic (Active Motif, #53009) and H3K4me2 antibody

5 (Millipore, #07-030). The culture supernatant was aspirated, ice-cold PBS was added, and the cell suspension was collected on ice using CELL SCRAPER (IWAKI). The supernatant was removed by centrifugation at 3000 rpm, 4°C for 5 min. Lysis buffer (500 μ L) was added to the precipitate, and the cells were
10 lysed by incubating for 30 min on ice. Thereafter, the suspension was centrifuged at 2400 g, 4°C for 10 min, the supernatant was removed, the pellet was suspended in cell lysis buffer [60 mM KCl, 15 mM NaCl, 5 mM MgCl₂, 0.1 mM EGTA, 15 mM Tris-HCl (pH 7.6), 1.2 M sucrose, 0.5 mM DTT, protease
15 inhibitor (Roche, #4693132)] (500 μ L), and the suspension was centrifuged at 10000 g, 4°C for 10 min. The supernatant was removed, and the pellet was suspended in Digestion buffer (120 μ L) and preincubated at 37°C for 5 min. Shearing cocktail was added, and the mixture was incubated at 37°C for 20 min. 0.5 M
20 EDTA (2.4 μ L) was added, and the mixture was incubated for 10 min on ice and centrifuged at 18000 rpm, 4°C for 10 min. The supernatant was collected as a chromatin fraction and subjected to immunoprecipitation.

[0571]

25 Using the DNA obtained by the chromatin immunoprecipitation as a template, quantitative PCR of the Gad1 gene upstream genomic region was performed, and the measurement value was taken as the Gad1 H3K4me2 level. The quantitative PCR was performed by ABI PRISM 7900HT Sequence
30 Detection System (Applied Biosystems) using forward primer: 5'-TGATCTTTTCCCTGCTGTCA-3' (SEQ ID NO: 8), reverse primer: 5'-TCCCATGAGTAATCCAGAACG-3' (SEQ ID NO: 9), and SYBR Green Realtime PCR Master Mix-Plus- (TOYOBO, #QPK-212). The Gad1 H3K4me2 induction by the compound was expressed by the H3K4me2
35 level when the compound was added, as compared to the H3K4me2

level of the control (without addition of the compound) as 100%.

Gad1 H3K4me2 induction activity (%) (H3K4me2 induction
5 expressed as percentage of control) = (H3K4me2 level with
addition of compound ÷ H3K4me2 level without addition of
compound) ×100

The Gad1 H3K4me2 induction activity by each compound as
measured by the above-mentioned method is shown in Table 3.

10 [0572]

[Table 3]

Example No.	Gad1 H3K4me2 induction activity (%)
	10 μM
102	153
82	115
123	143
99	146
97	184
71	182
118	233

[0573]

15 From the results of Table 3, it was clarified that the
compound of the present invention has an inductive effect on
H3K4 methylation.

[0574]

Experimental Example 4

20 Evaluation of blood cell number in mouse

Experimental method

Male ICR mice (hereinafter mice) were acclimated for at
least one week in a rearing facility. The mice were raised in
a rearing room with controlled temperature and humidity under
25 a 12:12 hour light-dark cycle, and allowed free ingestion of
feed and water.

The compounds were suspended in 0.5% methylcellulose/0.5% citric acid/distilled water and orally administered. All compounds were repeatedly administered to the mice at a dose of 1 mg/kg, 10 mg/kg or 100 mg/kg (body weight) for 7 or 9 days. One day after the final administration of the compound, the whole blood was collected.

Using Sysmex XT-1800i (Sysmex Corporation), the white blood cell number, red blood cell number, and platelet number in the collected whole blood per unit volume were measured. The influence of each compound on each blood cell number was determined by calculating the number with the mean of each blood cell number of the mouse without administration of the compound (0 mg/kg group), and the mean of each blood cell number in the whole blood of a mouse with the administration of the compound. The values of the blood cell numbers measured by the above-mentioned method are shown in Table 4.

[0575]

[Table 4-1]

Example No.	White blood cells (10 cells)							
	0 mg/kg		1 mg/kg		10 mg/kg		100 mg/kg	
	Mean	S.E.	Mean	S.E.	Mean	S.E.	Mean	S.E.
102	457	71	335	40	425	34	361	50
99	340	30	399	46	378	40	347	28
97	397	59	377	43	366	44	332	34
71	275	15	275	29	276	20	323	16
82	322	17	403	47	424	51	329	21
84	312	44	370	30	339	41	313	38
118	275	15	299	51	312	34	245	12
123	389	42	390	47	368	22	380	24
104	375	22	373	44	397	28	-	-
95	375	22	422	46	408	27	-	-

[0576]

[Table 4-2]

Example No.	Red blood cells (10^4 cells)							
	0 mg/kg		1 mg/kg		10 mg/kg		100 mg/kg	
	Mean	S.E.	Mean	S.E.	Mean	S.E.	Mean	S.E.
102	888	16	903	12	929	18	929	16
99	863	17	892	17	915	20	876	19
97	902	12	902	12	890	25	929	18
71	830	32	873	23	817	20	853	8
82	872	11	899	20	926	21	903	19
84	884	25	910	18	880	24	900	12
118	830	32	853	17	907	19	881	13
123	829	13	846	14	854	13	866	7
104	883	8	870	16	872	16	-	-
95	883	8	888	15	898	12	-	-

[0577]

5 [Table 4-3]

Example No.	Platelets (10^3 cells)							
	0 mg/kg		1 mg/kg		10 mg/kg		100 mg/kg	
	Mean	S.E.	Mean	S.E.	Mean	S.E.	Mean	S.E.
102	1401	53	1387	56	1499	43	1417	61
99	1468	71	1442	65	1380	49	1519	99
97	1466	22	1462	65	1483	78	1533	55
71	1148	63	1178	76	1197	67	1322	91
82	1389	35	1399	44	1445	38	1411	58
84	1515	60	1533	43	1627	58	1665	107
118	1148	63	1051	71	1158	66	1176	60
123	1130	58	1086	79	1234	68	1333	75
104	1345	33	1370	32	1404	77	-	-
95	1345	33	1352	66	1482	66	-	-

[0578]

From the results of Table 4, it was clarified that the

compound of the present invention reduces an influence on the white blood cell number, red blood cell number and platelet number.

[0579]

5 Experimental Example 5

Evaluation of hippocampal distribution in mouse

Experimental method

Male ICR mice (hereinafter mice) were acclimated for at least one week in a rearing facility. The mice were raised in
10 a rearing room with controlled temperature and humidity under a 12:12 hour light-dark cycle, and allowed free ingestion of feed and water.

The compounds were suspended in 0.5% methylcellulose/0.5% citric acid/distilled water and orally administered. All
15 compounds were administered to the mice at a dose of 10 mg/kg (body weight).

Blood samples were collected at 0.5 hr or 1 hr from compound administration, and the hippocampus was isolated simultaneously. The plasma concentration and hippocampus
20 concentration of each test compound were measured by the LC/MS/MS method, the ratio (hippocampus/plasma concentration ratio) was calculated, and distribution into hippocampus was evaluated. The results are shown in Table 5.

[0580]

[Table 5]

Example No.	time after administration (h)	Concentration ($\mu\text{g/mL}$ or $\mu\text{g/g}$)				Ratio (hippocampus/plasma)	
		Plasma		Hippocampus			
		Mean	S.D.	Mean	S.D.	Mean	S.D.
102	1	0.348	0.008	0.104	0.021	0.298	0.062
99	1	2.948	0.934	0.640	0.168	0.223	0.047
97	1	0.978	0.182	0.194	0.012	0.205	0.051
71	1	0.229	0.027	0.082	0.015	0.358	0.063
82	1	1.327	0.115	0.170	0.021	0.128	0.015
84	1	0.523	0.120	0.177	0.040	0.339	0.014
118	1	0.405	0.113	0.151	0.022	0.384	0.061
123	0.5	1.107	0.123	0.369	0.089	0.333	0.070

5 [0581]

As shown in Table 5, the compounds of the present invention were all confirmed to have distributed into the hippocampus.

[0582]

10 Formulation Example 1

A medicament containing the compound of the present invention as an active ingredient can be produced, for example, according to the following formulation.

1. capsule

15	(1) compound obtained in Example 1	10 mg
	(2) lactose	90 mg
	(3) crystalline cellulose	70 mg
	(4) magnesium stearate	10 mg
	1 capsule	180 mg

20 The total amount of the above-mentioned (1), (2) and (3) and 5 mg of (4) are blended, and the mixture is granulated. Thereto is added the remaining 5 mg of (4), and the whole is

sealed in a gelatin capsule.

[0583]

2. tablet

(1) compound obtained in Example 1 10 mg

5 (2) lactose 35 mg

(3) cornstarch 150 mg

(4) crystalline cellulose 30 mg

(5) magnesium stearate 5 mg

1 tablet 230 mg

10 The total amount of the above-mentioned (1), (2) and (3), 20 mg of (4) and 2.5 mg of (5) are blended, and the mixture is granulated. Thereto are added the remaining 10 mg of (4) and 2.5 mg of (5), and the mixture is compression-molded to give a tablet.

15 Industrial Applicability

[0584]

The compound of the present invention has a superior LSD1 inhibitory action, and is useful as a medicament such as a prophylactic or therapeutic agent for cancer, schizophrenia, 20 developmental disorders, particularly diseases having intellectual disability (e.g., autistic spectrum disorders, Rett syndrome, Down's syndrome, Kabuki syndrome, fragile X syndrome, Kleeftstra syndrome, neurofibromatosis type 1, Noonan syndrome, tuberous sclerosis), neurodegenerative diseases 25 (e.g., Alzheimer's disease, Parkinson's disease, spinocerebellar degeneration (e.g., dentatorubral pallidoluysian atrophy) and Huntington's disease (Huntington chorea)), epilepsy (e.g., Dravet syndrome) or drug dependence, and the like.

30 [0585]

This application is based on patent application No. 2014-82057 filed in Japan, the entire contents of which are incorporated by reference herein.

SEQUENCE LISTING

<110> TAKEDA PHARMACEUTICAL COMPANY LIMITED

<120> CYCLOPROPANAMINE COMPOUND AND USE THEREOF

<130> 092297

<150> JP 2014-082057

<151> 2014-04-11

<160> 9

<170> PatentIn version 3.5

<210> 1

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> primer

<400> 1

agaatcattt aaatggtgat catgtaacc atcct

35

<210> 2

<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> primer

<400> 2

cgccctgaaa gtacaggttc tcatccgatt ttggaggatg gtcg

44

<210> 3

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> primer

<400> 3

ataataggat ccgccctgaa agtacaggtt ctc

33

<210> 4

<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> primer
 <400> 4
 tattatgcta gcgccaccat gttatctggg aagaaggcgg cagc 44

 <210> 5
 <211> 38
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> primer

 <400> 5
 tattatgcgg ccgctcacat gcttggggac tgctgtgc 38

 <210> 6
 <211> 34
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> primer

 <400> 6
 ataataagat cttcgggtgt ggagggcgca gctt 34

 <210> 7
 <211> 39
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> primer

 <400> 7
 ataatagcgg ccgccatggc ccccaaaaac tggctctgca 39

 <210> 8
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> primer

 <400> 8
 tgatcttttc cctgctgtca 20

 <210> 9
 <211> 21
 <212> DNA

<213> Artificial Sequence

<220>

<223> primer


<400> 9

tcccatgagt aatccagaac g

21

CLAIMS

1. A compound represented by the formula



wherein


A is an optionally substituted heterocyclic group, or an optionally substituted hydrocarbon group;

10 B is a ring selected from

(1) a 5- or 6-membered aromatic heterocycle optionally fused with an optionally substituted 5- or 6-membered ring, and

(2) a benzene ring fused with an optionally substituted 5- or 6-membered ring,


15 wherein the ring represented by B is optionally substituted, and binds, via two adjacent carbon atoms with one atom in between, to a group represented by the formula



, and

20

a group represented by the formula



25 R¹, R², R³ and R⁴ are each independently a hydrogen atom, an

optionally substituted hydrocarbon group or an optionally substituted heterocyclic group;

A and R¹ are optionally bonded with each other to form, together with the adjacent nitrogen atom, an optionally substituted cyclic group; and

R² and R³ are optionally bonded with each other to form, together with the adjacent nitrogen atom, an optionally substituted cyclic group, or a salt thereof.

10

2. The compound according to claim 1, wherein A is

- (1) an optionally substituted heterocyclic group, or
- (2) an optionally substituted C₃₋₁₀ cycloalkyl group, or a salt thereof.

15

3. The compound according to claim 1, wherein B is a ring selected from

- (1) a 5- or 6-membered aromatic heterocycle, and
- (2) a benzene ring fused with an optionally substituted 5- or 6-membered ring,


20

wherein the ring represented by B is optionally substituted, and binds, via two adjacent carbon atoms with one atom in between, to a group represented by the formula (II), and a group represented by the formula (III),

25


or a salt thereof.

4. The compound according to claim 1, wherein, in the formula



30

B is a ring selected from



wherein

X¹, X², X³ and X⁴ are each independently a carbon atom or a
 5 nitrogen atom;

at least one of X¹, X², X³ and X⁴ is a nitrogen atom;

Y¹, Y² and Y³ are each independently a carbon atom, a nitrogen
 atom, an oxygen atom or a sulfur atom;


at least one of Y¹, Y² and Y³ is a nitrogen atom, an oxygen atom
 10 or a sulfur atom; and

Z¹, Z², Z³ and Z⁴ are each independently a carbon atom or a
 nitrogen atom, which ring is optionally substituted, or a salt
 thereof.

15 5. The compound according to claim 1, wherein R¹, R² and R⁴ are
 each independently a hydrogen atom or an optionally
 substituted C₁₋₆ alkyl group,
 or a salt thereof.

20 6. The compound according to claim 1, wherein R³ is
 (1) a hydrogen atom,
 (2) an optionally substituted C₁₋₆ alkyl group,
 (3) an optionally substituted C₃₋₁₀ cycloalkyl group, or
 (4) an optionally substituted heterocyclic group,
 25 or a salt thereof.

7. The compound according to claim 1, wherein, in the formula




5 A is

(1) (i) a 5- or 6-membered aromatic heterocyclic group or (ii) a 4- to 6-membered non-aromatic heterocyclic group, each of which is optionally substituted by C₁₋₆ alkyl group(s) optionally substituted by halogen atom(s), or

10 (2) a C₃₋₁₀ cycloalkyl group optionally substituted by halogen atom(s);

B is a ring selected from



15

wherein

X¹, X², X³ and X⁴ are each independently a carbon atom or a nitrogen atom;

at least one of X¹, X², X³ and X⁴ is a nitrogen atom;

20 Y¹, Y² and Y³ are each independently a carbon atom, a nitrogen atom, an oxygen atom or a sulfur atom;

at least one of Y¹, Y² and Y³ is a nitrogen atom, an oxygen atom or a sulfur atom; and

Z¹, Z², Z³ and Z⁴ are each independently a carbon atom or a nitrogen atom, which ring is optionally substituted by C₁₋₆ alkyl group(s);

R¹ is a hydrogen atom;

5 R² is a hydrogen atom;

R³ is

(1) a hydrogen atom,

(2) a C₁₋₆ alkyl group optionally substituted by substituent(s) selected from

10 (a) a C₃₋₁₀ cycloalkyl group,

(b) a C₆₋₁₄ aryl group optionally substituted by carboxy group(s),

(c) a 4- to 6-membered non-aromatic heterocyclic group optionally substituted by C₁₋₆ alkyl group(s) optionally substituted by substituent(s) selected from a carboxy group, and a C₆₋₁₄ aryl group optionally substituted by carboxy group(s), and

(d) a 5- or 6-membered aromatic heterocyclic group optionally substituted by amino group(s),

20 (3) a C₃₋₁₀ cycloalkyl group optionally substituted by substituent(s) selected from an amino group and a halogen atom or

(4) a 4- to 6-membered non-aromatic heterocyclic group optionally substituted by substituent(s) selected from

25 (a) a C₁₋₆ alkyl group optionally substituted by halogen atom(s),

(b) a C₃₋₁₀ cycloalkyl group,

(c) a C₁₋₆ alkyl-carbonyl group, and

(d) a C₃₋₁₀ cycloalkyl-carbonyl group; and

30 R⁴ is a hydrogen atom,

or a salt thereof.

8. The compound according to claim 1, wherein A is

(1) a piperidinyl group, an isoxazolyl group, a pyrazolyl group, a thiadiazolyl group, a thiazolyl group, a

tetrahydropyranyl group, an oxetanyl group, an oxadiazolyl group, a thienyl group, a pyridyl group or an oxazolyl group, each of which is optionally substituted by C₁₋₆ alkyl group(s) optionally substituted by halogen atom(s), or

- 5 (2) a C₃₋₁₀ cycloalkyl group optionally substituted by halogen atom(s);

B is a ring selected from

thiophene, thiazole, pyrazole, pyridine, naphthalene and 2,3-dihydrobenzofuran, wherein the ring is optionally substituted

- 10 by C₁₋₆ alkyl group(s);

R¹ is a hydrogen atom;

R² is a hydrogen atom;

R³ is

(1) a hydrogen atom,

- 15 (2) a C₁₋₆ alkyl group optionally substituted by substituent(s) selected from

(a) a C₃₋₁₀ cycloalkyl group,

(b) a C₆₋₁₄ aryl group optionally substituted by carboxy group(s),

- 20 (c) a tetrahydropyranyl group or a piperidinyl group each optionally substituted by C₁₋₆ alkyl group(s) optionally substituted by substituent(s) selected from a carboxy group, and a C₆₋₁₄ aryl group optionally substituted by carboxy group(s), and

- 25 (d) an oxadiazolyl group optionally substituted by an amino group,

(3) a C₃₋₁₀ cycloalkyl group optionally substituted by substituent(s) selected from an amino group and a halogen atom or

- 30 (4) a tetrahydropyranyl group or a piperidinyl group each optionally substituted by substituent(s) selected from

(a) a C₁₋₆ alkyl group optionally substituted by halogen atom(s),

(b) a C₃₋₁₀ cycloalkyl group,

- 35 (c) a C₁₋₆ alkyl-carbonyl group, and

(d) a C₃₋₁₀ cycloalkyl-carbonyl group; and
R⁴ is a hydrogen atom,
or a salt thereof.

5 9. The compound according to claim 1, wherein A is
(1) a pyrazolyl group, a thiadiazolyl group or a
tetrahydropyranyl group, each optionally substituted by C₁₋₆
alkyl group(s), or
(2) a cyclohexyl group optionally substituted by halogen
10 atom(s);
B is
a thiophene ring, which ring is optionally substituted by C₁₋₆
alkyl group(s);
R¹ is a hydrogen atom;
15 R² is a hydrogen atom;
R³ is a cyclopropylmethyl group, a tetrahydropyranylmethyl
group, a cyclobutylmethyl group, a cyclobutyl group or a
tetrahydropyranyl group; and
R⁴ is a hydrogen atom,
20 or a salt thereof.

10. The compound according to claim 1, wherein A is
a pyrazolyl group, a thiadiazolyl group or a tetrahydropyranyl
group, each optionally substituted by C₁₋₆ alkyl group(s),
25 B is
a thiophene ring, which ring is optionally substituted by C₁₋₆
alkyl group(s);
R¹ is a hydrogen atom;
R² is a hydrogen atom;
30 R³ is a cyclopropylmethyl group or a cyclobutyl group; and
R⁴ is a hydrogen atom,
or a salt thereof.

11. 5-((1R,2R)-2-((Cyclopropylmethyl)amino)cyclopropyl)-N-
35 (tetrahydro-2H-pyran-4-yl)thiophene-3-carboxamide or a salt

thereof.

12. 4-((1S,2R)-2-(Cyclobutylamino)cyclopropyl)-N-(5-methyl-
1,3,4-thiadiazol-2-yl)thiophene-2-carboxamide or a salt
5 thereof.

13. 4-((1S,2R)-2-(Cyclobutylamino)cyclopropyl)-5-methyl-N-(1-
methyl-1H-pyrazol-4-yl)thiophene-2-carboxamide or a salt
thereof.

10

14. A medicament comprising the compound according to claim 1
or a salt thereof.

15. The medicament according to claim 14, which is an LSD1
15 inhibitor.

16. The medicament according to claim 14, which is a
prophylactic or therapeutic agent for schizophrenia,
developmental disorders, autistic spectrum disorders, Rett
20 syndrome, Down's syndrome, Kabuki syndrome, fragile X syndrome,
Kleefstra syndrome, neurofibromatosis type 1, Noonan syndrome,
tuberous sclerosis, Alzheimer's disease, Parkinson's disease,
spinocerebellar degeneration, Huntington's disease, epilepsy
or drug dependence.

25

17. The compound according to claim 1 or a salt thereof for
use in the prophylaxis or treatment of schizophrenia,
developmental disorders, autistic spectrum disorders, Rett
syndrome, Down's syndrome, Kabuki syndrome, fragile X syndrome,
30 Kleefstra syndrome, neurofibromatosis type 1, Noonan syndrome,
tuberous sclerosis, Alzheimer's disease, Parkinson's disease,
spinocerebellar degeneration, Huntington's disease, epilepsy
or drug dependence.

35 18. A method of inhibiting LSD1 in a mammal, comprising

administering an effective amount of the compound according to claim 1 or a salt thereof to the mammal.

19. A method for the prophylaxis or treatment of schizophrenia,
5 developmental disorders, autistic spectrum disorders, Rett
syndrome, Down's syndrome, Kabuki syndrome, fragile X syndrome,
Kleefstra syndrome, neurofibromatosis type 1, Noonan syndrome,
tuberous sclerosis, Alzheimer's disease, Parkinson's disease,
spinocerebellar degeneration, Huntington's disease, epilepsy
10 or drug dependence in a mammal, comprising administering an
effective amount of the compound according to claim 1 or a
salt thereof to the mammal.

20. Use of the compound according to claim 1 or a salt thereof
15 in the production of a prophylactic or therapeutic agent for
schizophrenia, developmental disorders, autistic spectrum
disorders, Rett syndrome, Down's syndrome, Kabuki syndrome,
fragile X syndrome, Kleefstra syndrome, neurofibromatosis type
1, Noonan syndrome, tuberous sclerosis, Alzheimer's disease,
20 Parkinson's disease, spinocerebellar degeneration,
Huntington's disease, epilepsy or drug dependence.