

ACT 115A

Novel Thiophene Derivatives

5 Field of the invention

The present invention relates to S1P1/EDG1 receptor agonists of Formula (I) and their use as active ingredients in the preparation of pharmaceutical compositions. The invention also concerns related aspects including processes for the preparation
10 of the compounds, pharmaceutical compositions containing a compound of the Formula (I), and their use as compounds improving vascular function and as immunomodulating agents, either alone or in combination with other active compounds or therapies.

15 Background of the invention

The human immune system is designed to defend the body against foreign micro-organisms and substances that cause infection or disease. Complex regulatory mechanisms ensure that the immune response is targeted against the intruding
20 substance or organism and not against the host. In some cases, these control mechanisms are unregulated and autoimmune responses can develop. A consequence of the uncontrolled inflammatory response is severe organ, cell, tissue or joint damage. With current treatment, the whole immune system is usually suppressed and the body's ability to react to infections is also severely
25 compromised. Typical drugs in this class include azathioprine, chlorambucil, cyclophosphamide, cyclosporin, or methotrexate. Corticosteroids which reduce inflammation and suppress the immune response, may cause side effects when used in long term treatment. Nonsteroidal anti-inflammatory drugs (NSAIDs) can reduce pain and inflammation, however, they exhibit considerable side effects.
30 Alternative treatments include agents that activate or block cytokine signaling.


Orally active compounds with immunomodulating properties, without compromising immune responses and with reduced side effects would significantly improve current treatments of uncontrolled inflammatory disease.

In the field of organ transplantation the host immune response must be suppressed to prevent organ rejection. Organ transplant recipients can experience some rejection even when they are taking immunosuppressive drugs. Rejection occurs most frequently in the first few weeks after transplantation, but rejection episodes can also happen months or even years after transplantation. Combinations of up to three or four medications are commonly used to give maximum protection against rejection while minimizing side effects. Current standard drugs used to treat the rejection of transplanted organs interfere with discrete intracellular pathways in the activation of T-type or B-type white blood cells. Examples of such drugs are cyclosporin, daclizumab, basiliximab, everolimus, or FK506, which interfere with cytokine release or signaling; azathioprine or leflunomide, which inhibit nucleotide synthesis; or 15-deoxyspergualin, an inhibitor of leukocyte differentiation.

The beneficial effects of broad immunosuppressive therapies relate to their effects; however, the generalized immunosuppression which these drugs produce diminishes the immune system's defense against infection and malignancies. Furthermore, standard immunosuppressive drugs are often used at high dosages and can cause or accelerate organ damage.

Summary of Invention


In a first aspect, there is provided a compound of the Formula (I),



Formula (I)

wherein

A represents $^*\text{-CO-CH}_2\text{CH}_2\text{-}$, $^*\text{-CO-CH=CH-}$,



wherein the asterisks indicate the bond that is linked to the thiophene group of Formula (I);

R^1 represents C_{2-5} -alkyl;

R^2 represents hydrogen, methyl or ethyl;

R^3 represents hydrogen;

R^4 represents methyl, ethyl, or methoxy;

R^5 represents $-\text{CH}_2-(\text{CH}_2)_n-\text{NHSO}_2R^{51}$, $-(\text{CH}_2)_n\text{CH}(\text{OH})-\text{CH}_2-\text{NHSO}_2R^{51}$, $-\text{CH}_2-(\text{CH}_2)_n-\text{NHCOR}^{52}$, $-(\text{CH}_2)_n\text{CH}(\text{OH})-\text{CH}_2-\text{NHCOR}^{52}$, $-\text{CH}_2-(\text{CH}_2)_n-\text{CONR}^{53}R^{54}$, $-\text{CO-NHR}^{53}$, 1-(3-carboxy-azetidiny)-2-acetyl, 1-(2-carboxy-pyrrolidiny)-2-acetyl, 1-(3-carboxy-pyrrolidiny)-2-acetyl, 1-(3-carboxy-azetidiny)-3-propionyl, 1-(2-carboxy-pyrrolidiny)-3-propionyl, 1-(3-carboxy-pyrrolidiny)-3-propionyl, hydroxy, hydroxy- C_{2-5} -alkoxy, di-(hydroxy- C_{1-4} -alkyl)- C_{1-4} -alkoxy, 2,3-dihydroxypropoxy, 2-hydroxy-

2b

3-methoxy-propoxy, $-\text{OCH}_2-(\text{CH}_2)_m-\text{NHSO}_2\text{R}^{51}$, $-\text{OCH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{NHSO}_2\text{R}^{51}$, $-\text{OCH}_2-(\text{CH}_2)_m-\text{NHCOR}^{52}$, or $-\text{OCH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{NHCOR}^{52}$;

R^{51} represents C_{1-3} -alkyl, methylamino, ethylamino, or dimethylamino;

R^{52} represents hydroxymethyl, 2-hydroxyethyl, 2-hydroxy-1-hydroxymethyl-ethyl, or 2,3-dihydroxypropyl;

R^{53} represents hydrogen, C_{1-3} -alkyl, 2-hydroxyethyl, 2-hydroxy-1-hydroxymethyl-ethyl, 2,3-dihydroxypropyl, carboxymethyl, 1-(C_{1-5} -alkylcarboxy)methyl, 2-carboxyethyl, or 2-(C_{1-5} -alkylcarboxy)ethyl;

R^{54} represents hydrogen, or methyl;

m represents the integer 1 or 2;

n represents 0, 1, or 2; and

R^6 represents methyl, ethyl or halogen;

and salts thereof.

In a second aspect, there is provided a pharmaceutical composition comprising a compound as defined above, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

In a third aspect, there is provided a compound as defined above, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition as defined above, for use as a medicament.

In a fourth aspect, there is provided the use of a compound as defined above, or a pharmaceutically acceptable salt thereof, for the preparation of a

pharmaceutical composition for the prevention or treatment of diseases or disorders associated with an activated immune system.

In a fifth aspect, there is provided the use of a compound as defined above, or a pharmaceutically acceptable salt thereof, for the preparation of a pharmaceutical composition for use in combination with one or several agents selected from the group consisting of immunosuppressants, corticosteroids, NSAID's, cytotoxic drugs, adhesion molecule inhibitors, cytokines, cytokine inhibitors, cytokine receptor antagonists and recombinant cytokine receptors, for the prevention or treatment of diseases or disorders associated with an activated immune system.

Description of the invention

The present invention provides novel compounds of Formula (I) that are agonists for the G protein-coupled receptor S1P1/EDG1 and have a powerful and long-lasting immunomodulating effect which is achieved by reducing the number of circulating and infiltrating T- and B-lymphocytes, without affecting their maturation, memory, or expansion. The reduction of circulating T- / B-lymphocytes as a result of S1P1/EDG1 agonism, possibly in combination with the observed improvement of endothelial cell layer function associated with S1P1/EDG1 activation, makes such compounds useful to treat uncontrolled inflammatory disease and to improve vascular functionality.

The compounds of the present invention can be utilized alone or in combination with standard drugs inhibiting T-cell activation, to provide a new immunomodulating therapy with a reduced propensity for infections when compared to standard

immunosuppressive therapy. Furthermore, the compounds of the present invention can be used in combination with reduced dosages of traditional immunosuppressant therapies, to provide on the one hand effective immunomodulating activity, while on the other hand reducing end organ damage associated with higher doses of standard immunosuppressive drugs. The observation of improved endothelial cell layer function associated with S1P1/EDG1 activation provides additional benefits of compounds to improve vascular function.

The nucleotide sequence and the amino acid sequence for the human S1P1/EDG1 receptor are known in the art and are published in e.g.: Hla, T., and Maciag, T. *J. Biol Chem.* **265** (1990), 9308-9313; WO 91/15583 published 17 October 1991; WO 99/46277 published 16 September 1999. The potency and efficacy of the compounds of Formula (I) are assessed using a GTP γ S assay to determine EC₅₀ values and by measuring the circulating lymphocytes in the rat after oral administration, respectively (see in Examples).

The general terms used hereinbefore and hereinafter preferably have, within this disclosure, the following meanings, unless otherwise indicated:

Where the plural form is used for compounds, salts, pharmaceutical compositions, diseases and the like, this is intended to mean also a single compound, salt, or the like.

Any reference hereinbefore or hereinafter to a compound of Formula (I) is to be understood as referring also to salts, especially pharmaceutically acceptable salts, of a compound of Formula (I), as appropriate and expedient.

The term **C₁₋₅-alkyl**, alone or in combination with other groups, means saturated, branched or preferably straight chain groups with one to five carbon atoms. Preferred examples of C₁₋₅-alkyl groups are methyl, ethyl, n-propyl, n-butyl, isobutyl, and n-pentyl.

The term **C₁₋₄-alkyl**, alone or in combination with other groups, means saturated, branched or preferably straight chain groups with one to four carbon atoms, preferably one to three carbon atoms, i.e. C₁₋₃-alkyl. Preferred examples of C₁₋₄-alkyl groups are methyl, ethyl, and n-propyl.

5

Likewise, the term **C₂₋₅-alkyl**, alone or in combination with other groups, means saturated, branched or straight chain groups with two to five carbon atoms. Preferred examples of C₂₋₅-alkyl groups are ethyl, n-propyl, n-butyl, iso-butyl, n-pentyl, and iso-pentyl.

10

The term **C₁₋₅-alkoxy**, alone or in combination with other groups, means an R-O group, wherein R is a C₁₋₅-alkyl.

15

The term **C₁₋₄-alkoxy**, alone or in combination with other groups, means an R-O group, wherein R is a C₁₋₄-alkyl. Preferred examples of C₁₋₄-alkoxy groups are methoxy, ethoxy, propoxy, iso-propoxy, and iso-butoxy.

The term **C₂₋₅-alkoxy**, alone or in combination with other groups, means an R-O group, wherein R is a C₂₋₅-alkyl.

20

The term **halogen** means fluoro, chloro, bromo or iodo, preferably fluoro or chloro.

Salts are preferably the pharmaceutically acceptable salts of the compounds of Formula (I).

25


The term "pharmaceutically acceptable salts" refers to non-toxic, inorganic or organic acid and/or base addition salts, Lit.: e.g. "Salt selection for basic drugs", *Int. J. Pharm.* (1986), **33**, 201-217.

30

The compounds of Formula (I) may contain one or more stereogenic or asymmetric centers, such as one or more asymmetric carbon atoms. Substituents at a double bond or a ring may be present in cis- (= Z-) or trans (= E-) form unless indicated otherwise. The compounds of Formula (I) may thus be present as mixtures of

stereoisomers or preferably as pure stereoisomers. Mixtures of stereoisomers may be separated in a manner known to a person skilled in the art.


i) The invention relates to novel thiophene compounds of the Formula (I),



Formula (I)

wherein

10 A represents *-CO-CH₂CH₂-, *-CO-CH=CH-,



wherein the asterisks indicate the bond that is linked to the thiophene group of Formula (I);

15 R¹ represents C₂₋₅-alkyl;

R² represents hydrogen, methyl or ethyl;

R³ represents hydrogen, C₁₋₄-alkyl, C₁₋₄-alkoxy, or halogen;

R^4 represents hydrogen, C_{1-4} -alkyl, C_{1-4} -alkoxy, halogen, trifluoromethyl or trifluoromethoxy;

5 R^5 represents hydrogen, hydroxy- C_{1-5} -alkyl, 2,3-dihydroxypropyl, di-(hydroxy- C_{1-4} -alkyl)- C_{1-4} -alkyl, $-\text{CH}_2-(\text{CH}_2)_n-\text{NHSO}_2R^{51}$, $-(\text{CH}_2)_n\text{CH}(\text{OH})-\text{CH}_2-\text{NHSO}_2R^{51}$, $-\text{CH}_2-(\text{CH}_2)_n-\text{NHCOR}^{52}$, $-(\text{CH}_2)_n\text{CH}(\text{OH})-\text{CH}_2-\text{NHCOR}^{52}$, $-\text{CH}_2-(\text{CH}_2)_n-\text{CONR}^{53}R^{54}$, $-\text{CO}-\text{NHR}^{53}$, 1-(3-carboxy-azetidiny)-2-acetyl, 1-(2-carboxy-pyrrolidiny)-2-acetyl, 1-(3-carboxy-pyrrolidiny)-2-acetyl, 1-(3-carboxy-azetidiny)-3-propionyl, 1-(2-carboxy-pyrrolidiny)-3-propionyl, 1-(3-carboxy-pyrrolidiny)-3-propionyl, hydroxy, C_{1-5} -alkoxy, hydroxy- C_{2-5} -alkoxy, di-(hydroxy- C_{1-4} -alkyl)- C_{1-4} -alkoxy, 2,3-dihydroxypropoxy, 2-hydroxy-3-methoxy-propoxy, $-\text{OCH}_2-(\text{CH}_2)_m-\text{NHSO}_2R^{51}$, $-\text{OCH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{NHSO}_2R^{51}$, $-\text{OCH}_2-(\text{CH}_2)_m-\text{NHCOR}^{52}$, or $-\text{OCH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{NHCOR}^{52}$;

15 R^{51} represents C_{1-3} -alkyl, methylamino, ethylamino, or dimethylamino;

R^{52} represents hydroxymethyl, 2-hydroxyethyl, 2-hydroxy-1-hydroxymethyl-ethyl, or 2,3-dihydroxypropyl;

20 R^{53} represents hydrogen, C_{1-3} -alkyl, 2-hydroxyethyl, 2-hydroxy-1-hydroxymethyl-ethyl, 2,3-dihydroxypropyl, carboxymethyl, 1-(C_{1-5} -alkylcarboxy)methyl, 2-carboxyethyl, or 2-(C_{1-5} -alkylcarboxy)ethyl;

R^{54} represents hydrogen, or methyl;

25

m represents the integer 1 or 2;


n represents 0, 1, or 2; and

R^6 represents hydrogen, C_{1-4} -alkyl or halogen;

30


and salts thereof.

ii) A particular embodiment of the invention relates to thiophene derivatives according to embodiment i), wherein **A** represents




wherein the asterisks indicate the bond that is linked to the thiophene group of
5 Formula (I), and salts thereof.

iii) Another particular embodiment of the invention relates to thiophene derivatives according to embodiment i), wherein **A** represents




10 wherein the asterisks indicate the bond that is linked to the thiophene group of Formula (I), and salts thereof.

iv) Another particular embodiment of the invention relates to thiophene derivatives according to embodiment i), wherein **A** represents




wherein the asterisks indicate the bond that is linked to the thiophene group of Formula (I), and salts thereof.

- v) Another particular embodiment of the invention relates to thiophene derivatives according to embodiment i), wherein **A** represents



wherein the asterisks indicate the bond that is linked to the thiophene group of Formula (I), and salts thereof.

- 10 vi) Another particular embodiment of the invention relates to thiophene derivatives according to embodiment i), wherein **A** represents



and salts thereof.

- 15 vii) Another particular embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to vi), wherein **R**¹ represents n-propyl or isobutyl, and salts thereof.

- viii) Another particular embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to vii), wherein **R**² represents hydrogen or methyl, and salts thereof.
- 20

- ix) A preferred embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to vii), wherein **R**² represents hydrogen, and salts thereof.
- 25

- x) Another preferred embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to ix), wherein **R**³ represents methoxy, and **R**⁴ and **R**⁶ represent hydrogen, and salts thereof.

xi) Another preferred embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to ix), wherein R^3 represents hydrogen, R^4 represents methyl, ethyl, or methoxy, and R^6 represents methyl, ethyl or halogen, and salts thereof.

xii) Another preferred embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to ix), wherein R^3 represents hydrogen, and R^4 and R^6 represent a methyl group, and salts thereof.

xiii) A particularly preferred embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to ix), wherein R^3 represents hydrogen, R^4 represents a methyl group, and R^6 represents an ethyl group, and salts thereof.

xiv) Another preferred embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to ix), wherein R^3 represents hydrogen, R^4 represents a methoxy group, and R^6 represents a chlorine atom, and salts thereof.


xv) Another preferred embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to ix), wherein R^3 represents hydrogen, R^4 represents a methyl group, and R^6 represents a chlorine atom, and salts thereof.

xvi) Another preferred embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to xv), wherein R^5 represents 2,3-dihydroxypropyl, $-\text{CH}_2-(\text{CH}_2)_n-\text{NHCOR}^{52}$, $-(\text{CH}_2)_n\text{CH}(\text{OH})-\text{CH}_2-\text{NHCOR}^{52}$, hydroxy, hydroxy- C_{2-5} -alkoxy, di-(hydroxy- C_{1-4} -alkyl)- C_{1-4} -alkoxy, 2,3-dihydroxypropoxy, 2-hydroxy-3-methoxy-propoxy, $-\text{OCH}_2-(\text{CH}_2)_m-\text{NHCOR}^{52}$, or $-\text{OCH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{NHCOR}^{52}$, and salts thereof.

xvii) Another preferred embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to xv), wherein R^5 represents hydroxy,


hydroxy-C₂₋₅-alkoxy, di-(hydroxy-C₁₋₄-alkyl)-C₁₋₄-alkoxy, 2,3-dihydroxypropoxy, 2-hydroxy-3-methoxy-propoxy, -OCH₂-(CH₂)_m-NHCOR⁵², or -OCH₂-CH(OH)-CH₂-NHCOR⁵², and salts thereof.

- 5 xviii) Another preferred embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to xv), wherein R⁵ represents 3-hydroxy-2-hydroxymethyl-propoxy, 2,3-dihydroxypropoxy, or -OCH₂-CH(OH)-CH₂-NHCOR⁵², and salts thereof.
- 10 xix) Another preferred embodiment of the invention relates to thiophene derivatives according to any one of the embodiments i) to xv), wherein R⁵ represents 2,3-dihydroxypropoxy, or -OCH₂-CH(OH)-CH₂-NHCOR⁵², wherein R⁵² represents hydroxymethyl, and salts thereof.
- 15 xx) An especially preferred embodiment of the invention relates to thiophene derivatives according to embodiment i), wherein A represents



- wherein the asterisks indicate the bond that is linked to the thiophene group of Formula (I); R¹ represents n-propyl or isobutyl; R² represents hydrogen; R³ represents hydrogen or methoxy; R⁴ represents hydrogen, methyl, ethyl or methoxy; R⁵ represents 3-hydroxy-2-hydroxymethyl-propoxy, 2,3-dihydroxypropoxy, or -OCH₂-CH(OH)-CH₂-NHCOR⁵²; and R⁶ represents hydrogen, methyl, ethyl or chlorine; and salts thereof.
- 20

- 25 xxi) Another especially preferred embodiment of the invention relates to thiophene derivatives according to embodiment i), wherein A represents *-CO-CH₂CH₂-, *-CO-CH=CH-,



wherein the asterisks indicate the bond that is linked to the thiophene group of Formula (I);

R^1 represents C_{2-5} -alkyl;

R^2 and R^3 both represent hydrogen;

5 R^4 represents C_{1-4} -alkyl;

R^5 represents hydroxy, 2,3-dihydroxypropoxy, or $-OCH_2-CH(OH)-CH_2-NHCOR^{52}$;

R^{52} represents hydroxymethyl; and

R^6 represents C_{1-4} -alkyl;

and salts thereof.

10

xxii) Especially preferred thiophene compounds according to Formula (I) are:

(2R)-N-(3-{4-[5-(5-ethyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-2,6-dimethyl-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

15 (2S)-N-(3-{4-[5-(5-ethyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-2,6-dimethyl-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

(2R)-N-(3-{2,6-dimethyl-4-[5-(5-butyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

(2S)-N-(3-{2,6-dimethyl-4-[5-(5-butyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

20 (2R)-N-(3-{2,6-dimethyl-4-[5-(5-isobutyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

(2S)-N-(3-{2,6-dimethyl-4-[5-(5-isobutyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

25 (2R)-N-(3-{2-ethyl-6-methyl-4-[5-(5-isobutyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

(2S)-N-(3-{2-ethyl-6-methyl-4-[5-(5-isobutyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

(2R)-N-(3-{2,6-dimethyl-4-[5-(5-propyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

30 (2S)-N-(3-{2,6-dimethyl-4-[5-(5-propyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

(2R)-N-(3-{2-ethyl-6-methyl-4-[5-(5-propyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

(2S)-N-(3-{2-ethyl-6-methyl-4-[5-(5-propyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

(2R)-N-(3-{2-ethyl-4-[5-(5-isobutyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-6-methyl-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

5 (2S)-N-(3-{2-ethyl-4-[5-(5-isobutyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-6-methyl-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

(2R)-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propane-1,2-diol,

10 (2S)-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propane-1,2-diol,

(2R)-2-hydroxy-N-(2-hydroxy-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propyl)-acetamide,

(2S)-2-hydroxy-N-(2-hydroxy-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propyl)-acetamide,

15 (2R)-2-hydroxy-N-(2-hydroxy-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2-ethyl-6-methyl-phenoxy}-propyl)-acetamide, and

(2S)-2-hydroxy-N-(2-hydroxy-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2-ethyl-6-methyl-phenoxy}-propyl)-acetamide,

and salts of these compounds.

20

xxiii) Further especially preferred thiophene compounds according to Formula (I) are:

2-hydroxy-N-((2S)-2-hydroxy-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propyl)-acetamide,

25 N-((2S)-3-{2-ethyl-6-methyl-4-[5-(5-propyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,

(2R)-3-{2-ethyl-4-[5-(5-isobutyl-4-methyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-6-methyl-phenoxy}-propane-1,2-diol,

30 (2S)-3-{2-ethyl-4-[5-(5-isobutyl-4-methyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-6-methyl-phenoxy}-propane-1,2-diol,

N-((2R)-3-{2-ethyl-4-[5-(5-isobutyl-4-methyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-6-methyl-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide, and

N-((2S)-3-{2-ethyl-4-[5-(5-isobutyl-4-methyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-6-methyl-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide,
and salts of these compounds.

- 5 The compounds of Formula (I) and their pharmaceutically acceptable salts can be used as medicaments, e.g. in the form of pharmaceutical compositions for enteral or parental administration.

10 The production of the pharmaceutical compositions can be effected in a manner which will be familiar to any person skilled in the art (see for example Mark Gibson, Editor, *Pharmaceutical Preformulation and Formulation*, IHS Health Group, Englewood, CO, USA, 2001; Remington, *The Science and Practice of Pharmacy*, 20th Edition, Philadelphia College of Pharmacy and Science) by bringing the described compounds of Formula (I) or their pharmaceutically acceptable salts,
15 optionally in combination with other therapeutically valuable substances, into a galenical administration form together with suitable, non-toxic, inert, pharmaceutically acceptable solid or liquid carrier materials and, if desired, usual pharmaceutical adjuvants.

- 20 The pharmaceutical compositions comprising a compound of Formula (I) are useful for the prevention and/or treatment of diseases or disorders associated with an activated immune system.

25 Such diseases or disorders are selected from the group consisting of rejection of transplanted organs, tissue or cells; graft-versus-host diseases brought about by transplantation; autoimmune syndromes including rheumatoid arthritis; systemic lupus erythematosus; antiphospholipid syndrome; Hashimoto's thyroiditis; lymphocytic thyroiditis; multiple sclerosis; myasthenia gravis; type I diabetes; uveitis; episcleritis; scleritis; Kawasaki's disease, uveo-retinitis; posterior uveitis;
30 uveitis associated with Behcet's disease; uveomeningitis syndrome; allergic encephalomyelitis; chronic allograft vasculopathy; post-infectious autoimmune diseases including rheumatic fever and post-infectious glomerulonephritis; inflammatory and hyperproliferative skin diseases; psoriasis; psoriatic arthritis;

atopic dermatitis; myopathy; myositis; osteomyelitis; contact dermatitis; eczematous dermatitis; seborrhoeic dermatitis; lichen planus; pemphigus; bullous pemphigoid; epidermolysis bullosa; urticaria; angioedema; vasculitis; erythema; cutaneous eosinophilia; acne; scleroderma; alopecia areata; keratoconjunctivitis; vernal

5 conjunctivitis; keratitis; herpetic keratitis; dystrophia epithelialis corneae; corneal leukoma; ocular pemphigus; Mooren's ulcer; ulcerative keratitis; scleritis; Graves' ophthalmopathy; Vogt-Koyanagi-Harada syndrome; sarcoidosis; pollen allergies; reversible obstructive airway disease; bronchial asthma; allergic asthma; intrinsic asthma; extrinsic asthma; dust asthma; chronic or inveterate asthma; late asthma

10 and airway hyper-responsiveness; bronchiolitis; bronchitis; endometriosis; orchitis; gastric ulcers; ischemic bowel diseases; inflammatory bowel diseases; necrotizing enterocolitis; intestinal lesions associated with thermal burns; coeliac disease; proctitis; eosinophilic gastroenteritis; mastocytosis; Crohn's disease; ulcerative colitis; vascular damage caused by ischemic diseases and thrombosis;

15 atherosclerosis; fatty heart; myocarditis; cardiac infarction; aortitis syndrome; cachexia due to viral disease; vascular thrombosis; migraine; rhinitis; eczema; interstitial nephritis; IgA-induced nephropathy; Goodpasture's syndrome; hemolytic-uremic syndrome; diabetic nephropathy; glomerulosclerosis; glomerulonephritis; tubulointerstitial nephritis; interstitial cystitis; multiple myositis; Guillain-Barré

20 syndrome; Meniere's disease; polyneuritis; multiple neuritis; myelitis; mononeuritis; radiculopathy; hyperthyroidism; Basedow's disease; thyrotoxicosis; pure red cell aplasia; aplastic anemia; hypoplastic anemia; idiopathic thrombocytopenic purpura; autoimmune hemolytic anemia; autoimmune thrombocytopenia; agranulocytosis; pernicious anemia; megaloblastic anemia; anerythroplasia; osteoporosis; fibroid

25 lung; idiopathic interstitial pneumonia; dermatomyositis; leukoderma vulgaris; ichthyosis vulgaris; photoallergic sensitivity; cutaneous T cell lymphoma; polyarteritis nodosa; Huntington's chorea; Sydenham's chorea; myocardosis; myocarditis; scleroderma; Wegener's granuloma; Sjogren's syndrome; adiposis; eosinophilic fascitis; lesions of gingiva, periodontium, alveolar bone, substantia

30 ossea dentis; male pattern alopecia or alopecia senilis; muscular dystrophy; pyoderma; Sezary's syndrome; hypophysitis; chronic adrenal insufficiency; Addison's disease; ischemia-reperfusion injury of organs which occurs upon preservation; endotoxin shock; pseudomembranous colitis; colitis caused by drug or

radiation; ischemic acute renal insufficiency; chronic renal insufficiency; lung cancer; malignancy of lymphoid origin; acute or chronic lymphocytic leukemias; lymphoma; pulmonary emphysema; cataracta; siderosis; retinitis pigmentosa; senile macular degeneration; vitreal scarring; corneal alkali burn; dermatitis erythema; 5 ballous dermatitis; cement dermatitis; gingivitis; periodontitis; sepsis; pancreatitis; peripheral artery disease; carcinogenesis; solid cancer tumors; metastasis of carcinoma; hypobaropathy; autoimmune hepatitis; primary biliary cirrhosis; sclerosing cholangitis; partial liver resection; acute liver necrosis; cirrhosis; alcoholic cirrhosis; hepatic failure; fulminant hepatic failure; late-onset hepatic failure; and 10 "acute-on-chronic" liver failure.

Preferred diseases or disorders to be treated and/or prevented with the compounds of Formula (I) are selected from the group consisting of rejection of transplanted organs such as kidney, liver, heart, lung, pancreas, cornea, and skin; graft-versus- 15 host diseases brought about by stem cell transplantation; autoimmune syndromes including rheumatoid arthritis, multiple sclerosis, inflammatory bowel diseases such as Crohn's disease and ulcerative colitis, psoriasis, psoriatic arthritis, thyroiditis such as Hashimoto's thyroiditis, uveo-retinitis; atopic diseases such as rhinitis, conjunctivitis, dermatitis; asthma; type I diabetes; post-infectious autoimmune 20 diseases including rheumatic fever and post-infectious glomerulonephritis; solid cancers and tumor metastasis.

Particularly preferred diseases or disorders to be treated and/or prevented with the compounds of Formula (I) are selected from the group consisting of rejection of 25 transplanted organs selected from kidney, liver, heart and lung; graft-versus-host diseases brought about by stem cell transplantation; autoimmune syndromes selected from rheumatoid arthritis, multiple sclerosis, psoriasis, psoriatic arthritis, Crohn's disease, and Hashimoto's thyroiditis; and atopic dermatitis.


30 The present invention also relates to a method for the prevention or treatment of a disease or disorder mentioned herein comprising administering to a subject a pharmaceutically active amount of a compound of Formula (I).

Furthermore, compounds of the Formula (I) are also useful, in combination with one or several immunomodulating agents, for the prevention and/or treatment of the diseases and disorders mentioned herein. According to a preferred embodiment of the invention, said agents are selected from the group consisting of immunosuppressants, corticosteroids, NSAID's, cytotoxic drugs, adhesion molecule inhibitors, cytokines, cytokine inhibitors, cytokine receptor antagonists and recombinant cytokine receptors.


The present invention also relates to the use of a compound of Formula (I) for the preparation of a pharmaceutical composition, optionally for use in combination with one or several immunomodulating agents, for the prevention or treatment of the diseases and disorders mentioned herein.

The compounds of Formula (I) can be manufactured by the methods given below, by the methods given in the Examples or by analogous methods. Optimum reaction conditions may vary with the particular reactants or solvents used, but such conditions can be determined by a person skilled in the art by routine optimisation procedures.

Compounds of the Formula (I) of the present invention can be prepared according to the general sequence of reactions outlined below. Only a few of the synthetic possibilities leading to compounds of Formula (I) are described.




Structure 1




Structure 2

In case **A** represents $-\text{CO}-\text{CH}=\text{CH}-$, the compounds of Formula (I) may be prepared by reacting a compound of Structure 1 with a compound of Structure 2 in the presence of a base or an acid. The functional groups present in the residues R^3 to

R^6 may require temporary protection or may even be introduced in additional steps that follow the condensation reaction. Compounds of Formula (I) wherein **A** represents $-\text{CO}-\text{CH}_2-\text{CH}_2-$ may be prepared by reacting a compound of Formula (I) wherein **A** represents $-\text{CO}-\text{CH}=\text{CH}-$ with hydrogen in the presence of a catalyst such as Pd/C, Pt/C, PtO_2 , etc. in a solvent such as ethanol, methanol, THF, etc. or mixtures thereof.




Structure 3



Structure 4


- 10 A compound of Structure 1 may be prepared by reacting a compound of Structure 3 with a methyl Grignard reagent or by treating a compound of Structure 4 with 2 equivalents of methyllithium in a solvent such as diethyl ether, THF, etc. at temperatures between -20 and 50°C . The Weinreb amide compound of Structure 3 is prepared by treating a compound of Structure 4 with N,O-dimethylhydroxylamine hydrochloride in the presence of coupling reagent such as EDC, DCC, etc. (M. Mentzel, H. M. R. Hoffmann, N-Methoxy N-methyl amides (Weinreb amides) in modern organic synthesis, *Journal fuer Praktische Chemie/Chemiker-Zeitung* **339** (1997), 517-524; J. Singh, N. Satyamurthi, I. S. Aidhen, The growing synthetic utility of Weinreb's amide, *Journal fuer Praktische Chemie* (Weinheim, Germany) **342** (2000) 340-347; V. K. Khlestkin, D. G. Mazhukin, Recent advances in the application of N,O-dialkylhydroxylamines in organic chemistry, *Current Organic Chemistry* **7** (2003), 967-993).



Structure 5


Compounds of Formula (I) which represent a 5-thienyl-[1,2,4]oxadiazole derivative, are prepared by reacting a compound of Structure 5 in a solvent such as xylene, toluene, benzene, pyridine, DMF, dichloromethane, acetic acid, trifluoroacetic acid, etc. at rt or elevated temperatures in the presence or absence of auxiliaries such as acids (e.g. TFA, acetic acid, HCl, etc.), bases (e.g. NaH, NaOAc, Na₂CO₃, K₂CO₃, triethylamine, etc.), tetraalkylammonium salts, or water removing agents (e.g. oxalyl chloride, a carboxylic acid anhydride, POCl₃, PCl₅, P₄O₁₀, molecular sieves, methoxycarbonylsulfamoyl triethylammonium hydroxide (Burgess reagent), etc.) (Lit.: e.g. A. R. Gangloff, J. Litvak, E. J. Shelton, D. Sperandio, V. R. Wang, K. D. Rice, *Tetrahedron Lett.* **42** (2001), 1441-1443; T. Suzuki, K. Iwaoka, N. Imanishi, Y. Nagakura, K. Miyta, H. Nakahara, M. Ohta, T. Mase, *Chem. Pharm. Bull.* **47** (1999), 120-122; R. F. Poulain, A. L. Tartar, B. P. Déprez, *Tetrahedron Lett.* **42** (2001), 1495-1498; R. M. Srivastava, F. J. S. Oliveira, D. S. Machado, R. M. Souto-Maior, *Synthetic Commun.* **29** (1999), 1437-1450; E. O. John, J. M. Shreeve, *Inorganic Chemistry* **27** (1988), 3100-3104; B. Kaboudin, K. Navaee, *Heterocycles* **60** (2003), 2287-2292).

Compounds of Structure 5 may be prepared by reacting a compound of Structure 4 with a compound of Structure 6 in a solvent such as DMF, THF, DCM, etc. in the presence or absence of one or more coupling agents such as TBTU, DCC, EDC, HBTU, HOBt, CDI, etc. and in the presence or absence of a base such as triethylamine, Hünig's base, NaH, K₂CO₃, etc. (Lit.: e.g. A. Hamze, J.-F. Hernandez, P. Fulcrand, J. Martinez, *J. Org. Chem.* **68** (2003) 7316-7321; and the literature cited above).




Compounds of Formula (I) which represent a 3-thienyl-[1,2,4]oxadiazole derivative are prepared in an analogous fashion (Lit.: e.g. C. T. Brain, J. M. Paul, Y. Loong, P. J. Oakley, *Tetrahedron Lett.* **40** (1999) 3275-3278) by reacting a compound of

Structure 7 with a compound of Structure 8 and subsequent cyclisation of the corresponding hydroxyamidine ester intermediate.



Structure 7




Structure 8


5

Compounds of Structure 6 and 7 may be prepared by reacting a compound of Structure 9 and 10, respectively, with hydroxylamine or one of its salts in a solvent such as methanol, ethanol, pyridine, etc. in the presence or absence of a base such as Na_2CO_3 , K_2CO_3 , potassium tert.-butoxide, triethylamine, etc. (Lit.: e.g. T. Suzuki, K. Iwaoka, N. Imanishi, Y. Nagakura, K. Miyata, H. Nakahara, M. Ohta, T. Mase, *Chem. Pharm. Bull.* **47** (1999), 120-122; J. Cui, D. Crich, D. Wink, M. Lam, A. L. Rheingold, D. A. Case, W. T. Fu, Y. Zhou, M. Rao, A. J. Olson, M. E. Johnson, *Bioorg. Med. Chem.* **11** (2003), 3379-3392; R. Miller, F. Lang, Z. J. Song, D. Zewge, WO 2004/035538 (Merck & Co., Inc., USA); B. Kaboudin, K. Navaee, *Heterocycles* **60** (2003), 2287-2292).

15



Structure 9




Structure 10

20


Depending on the nature of the functionalities present in the residues R^3 to R^6 in Structures 2, 5, 6, 8, and 9, these functionalities may require temporary protection. Appropriate protecting groups are known to a person skilled in the art and include e.g. a benzyl or a trialkylsilyl group to protect an alcohol, a ketal to protect a diol, etc. These protecting groups may be employed according to standard methodology

(e.g. T. W. Greene, P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 3rd Edition, Wiley New York, 1991; P. J. Kocienski, *Protecting Groups*, Thieme Stuttgart, 1994). Alternatively, the desired residues R^3 to R^6 , in particular R^5 , may also be introduced in later steps that follow the coupling of the thiophene compounds of Structure 1, 4, 7 or 10 with the phenyl derivatives of Structure 2, 6, 8 or 9 by using a suitable precursor of a compound of Structure 2, 6, 8 and 9. The phenyl compounds of Structure 2, 6, 8 and 9 or their precursors are either commercially available or are prepared according to procedures known to a person skilled in the art.




Compounds of Formula (I) which represent a 2-thienyl-[1,3,4]oxadiazole or a 2-thienyl-[1,3,4]thiadiazole derivative are prepared similarly by reacting a compound of Structure 4 with hydrazine (by using a coupling reagent such as TBTU, DCC, EDC, HBTU, PyBOP, HOBt, CDI, etc.) to form a compound of Structure 11 which is then coupled with a compound of Structure 8 to give a compound of Structure 12. A compound of Structure 12 can also be prepared by following the reverse reaction order i.e. by first coupling a compound of Structure 8 with hydrazine followed by reacting the corresponding hydrazide intermediate with a compound of Structure 4. Dehydration of a compound of Structure 12 to form the desired 2-thienyl-[1,3,4]oxadiazole derivative is affected by treating a compound of Structure 12 with a reagent such as POCl_3 , CCl_4 or CBr_4 in combination with triphenylphosphine, P_2O_5 , Burgess reagent, etc. in a solvent such as toluene, acetonitrile, dioxane, THF, or CHCl_3 at temperatures between 20 and 120°C in the presence or absence of microwave irradiation. (Lit.: e.g. M. A. Garcia, S. Martin-Santamaria, M. Cacho, F. Moreno de la Llave, M. Julian, A. Martinez, B. De Pascual-Teresa, A. Ramos, *J. Med. Chem.* **48** (2005) 4068-4075; C. T. Brain, J. M. Paul, Y. Loong, P. J. Oakley, *Tetrahedron Lett.* **40** (1999) 3275-3278). Likewise, 2-thienyl-[1,3,4]thiadiazole


derivatives are obtained by cyclising a compound of Structure 12 with Lawesson's reagent optionally in combination with P_2S_5 in the absence or presence of a solvent such as pyridine, toluene, THF, or acetonitrile at elevated temperatures with or without microwave irradiation (Lit.: e.g. A. A. Kiryanov, P. Sampson, A. J. Seed, *J. Org. Chem.* **66** (2001) 7925-7929).



Compounds of Formula (I) which represent a 5-thienyl-oxazole or a 5-thienyl-thiazole derivative are prepared by treating a compound of Structure 13 either with $POCl_3$, PCl_5 , I_2 in combination with triphenylphosphine and triethylamine, Burgess reagent, trifluoroacetic anhydride, etc. in a solvent such as toluene, benzene, dioxane, or THF at temperatures between 20 and 120°C, or with Lawesson's reagent, optionally in combination with P_2S_5 , in the presence or absence of a solvent such as pyridine, toluene, THF, or acetonitrile at elevated temperatures with or without microwave irradiation as mentioned above (Lit.: e.g. N. Sato, T. Shibata, M. Jitsuoka, T. Ohno, T. Takahashi, T. Hirohashi, T. Kanno, H. Iwaasa, A. Kanatani, T. Fukami, Takehiro *Bioorg. & Med. Chem. Lett.* **14** (2004) 1761-1764). The compounds of Structure 13 are prepared by reacting a compound of Structure 14 with a compound of Structure 9. The aminoketon of Structure 14 can be prepared from a compound of Structure 1 by procedures given in the literature (e.g. J. L. LaMattina, *J. Heterocyclic Chem.* **20** (1983) 533-538; M. Pesson, M. Antoine, P. Girard, J. L. Benichon, S. Chabassier, P. De Lajudie, S. Patte, F. Roquet, G. Montay, *Eur. J. Med. Chem.* **15** (1980) 263-268). Compounds of Formula (I) which represent a 2-thienyl-oxazole or a 2-thienyl-thiazole derivative are prepared in an analogous fashion from a compound of Structure 15 and a compound of Structure 4.




Structure 14




Structure 15

Alternatively, the bonds between the thiophene or the phenylring and the central 5-membered heteroaromatic ring can also be formed by applying palladium catalysed cross coupling reactions.




Structure 16

The compounds of Structure 4 can be prepared by alkylating a thiophene-2-carboxylic acid of Structure 16 according to literature procedures (e.g. D. W. Knight, A. P. Nott, *J. Chem. Soc. Perkin Trans. 1* **1983** 791-794). The thiophene-2-carboxylic acids of Structure 16 or their methyl or ethyl esters are either commercially available or can be prepared following literature procedures (e.g. S. Gronowitz, T. Klingstedt, L. Svensson, U. Hansson, *Lipids* **28** (1993) 889-897).




Structure 17



Structure 18

The compounds of Structure 4 may also be prepared by reacting a compound of Structure 17, wherein R represents methyl, ethyl, tert. butyl, etc. with a 2,4,6-trialkenylcyclotriboroxane under Suzuki conditions (Lit.: e.g. F. Kerins, D. F. O'Shea, *J. Org. Chem.* **67** (2002) 4968-4971) to give a compound of Structure 18, wherein R represents methyl, ethyl, tert.-butyl, etc. and R^a and R^b both independently represent hydrogen, methyl, ethyl, etc., which upon hydrogenation

and subsequent ester cleavage furnishes the desired compound of Structure 4. Compounds of Structure 17 are either commercially available or are prepared according to literature procedures.



5

Compounds of Structure 4, wherein R² represents methyl or ethyl, may also be prepared by reacting a compound of Structure 19, wherein R^c represents ethyl or n-propyl, with POCl₃ in DMF under Vilsmeier conditions to give a compound of Structure 20 (Lit.: e.g. G. Alvernhe, D. Greif, B. Langlois, A. Laurent, I. Le Dréan, M. Pulst, A. Selmi, M. Weissenfels, *Bull. Soc. Chim. Fr.* **131** (1994) 167-172). The compound of Structure 20 is treated with a mercaptoacetic acid ester in the presence of a base such as NaH, NaOEt, NaOMe, K tert.-butoxide, etc., in THF, dioxane, DMF, ethanol, methanol, etc., or mixtures thereof, to form an intermediate of Structure 21. Cyclisation under basic conditions using a non aqueous base such as NaOMe, NaOEt, K^tBu, DBU, etc., in a solvent such as methanol, ethanol, THF, DMF, etc., or mixtures thereof, preferably at elevated temperatures, followed by saponification with an aqueous base such as aq. NaOH, aq. LiOH, aq. KOH, etc., or an acid such as aq. HCl, TFA, etc., in a solvent such as water, ethanol, methanol, THF, etc., or mixtures thereof, furnishes the compounds of Structure 4. Treating a compound of Structure 20 with mercaptoacetonitrile, which can be generated *in situ* from thioacetic acid S-cyanomethyl ester, may furnish a compound of Structure 10.

15

20

Methods that effect the transformation of a compound of Structure 4 into a compound of Structure 10, or the opposite, are known to a person skilled in the art.

25

Examples

The following examples illustrate the invention but do not at all limit the scope thereof.


All temperatures are stated in °C. Compounds are characterized by ¹H-NMR (300
 5 MHz) or ¹³C-NMR (75 MHz) (Varian Oxford; chemical shifts are given in ppm relative to the solvent used; multiplicities: s = singlet, d = doublet, t = triplet; p = pentuplet, hex = hexet, hept = heptet, m = multiplet, br = broad, coupling constants are given in Hz); by LC-MS (Finnigan Navigator with HP 1100 Binary Pump and DAD, column: 4.6x50 mm, Zorbax SB-AQ, 5 μm, 120 Å, gradient: 5-95%
 10 acetonitrile in water, 1 min, with 0.04% trifluoroacetic acid, flow: 4.5 mL/min), t_R is given in min; by TLC (TLC-plates from Merck, Silica gel 60 F₂₅₄); or by melting point. Compounds are purified by preparative HPLC (column: X-terra RP18, 50x19 mm, 5 μm, gradient: 10-95% acetonitrile in water containing 0.5 % of formic acid) or by MPLC (Labomatic MD-80-100 pump, Linear UVIS-201 detector, column: 350x18
 15 mm, Labogel-RP-18-5s-100, gradient: 10% methanol in water to 100% methanol).

Abbreviations (as used herein):

	aq.	aqueous
	BSA	bovine serum albumin
20	Bu	butyl
	CC	column chromatography
	CDI	carbonyl diimidazole
	DBU	1,8-diazabicyclo[5.4.0]undec-7-en
	DCC	dicyclohexyl carbodiimide
25	DCM	dichloromethane
	DIPEA	diisopropyl-ethylamine, Hünig's base, ethyl-diisopropylamine
	DME	1,2-dimethoxyethane
	DMF	dimethylformamide
	DMSO	dimethylsulfoxide
30	EA	ethyl acetate
	EDC	N-(3-dimethylaminopropyl)-N'-ethyl-carbodiimide
	Et	ethyl
	EtOH	ethanol


	h	hour(s)
	HBTU	O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
	HOBt	1-hydroxybenzotriazole
5	HPLC	high performance liquid chromatography
	KOtBu	potassium tert-butoxide
	LC-MS	liquid chromatography – mass spectrometry
	LDA	lithium diisopropyl amide
	Lit.	literature
10	Me	methyl
	min	minute(s)
	MPLC	medium pressure liquid chromatography
	NaOAc	sodium acetate
	NMO	N-methyl-morpholine-N-oxide
15	org.	organic
	Ph	phenyl
	prep.	preparative
	PyBOP	benzotriazol-1-yl-oxy-tris-pyrrolidino-phosphonium- hexafluoro-phosphate
20	rt	room temperature
	sat.	saturated
	S1P	sphingosine 1-phosphate
	TBME	tert.-butyl methyl ether
	TBTU	2-(1H-benzotriazole-1-yl)-1,2,3,3-tetramethyluronium
25		tetrafluoroborate
	TFA	trifluoroacetic acid
	THF	tetrahydrofuran
	TLC	thin layer chromatography
	t _R	retention time

30

5-Ethyl-thiophene-2-carboxylic acid


- A solution of thiophencarboxylic acid (4.00 g, 30.9 mmol) in THF (24 mL) is added dropwise via a syringe to a stirred solution of LDA (32.5 mL, 2 M in toluene) in THF (40 mL) cooled to -78°C . The temperature of the reaction is maintained at -78°C for 10 min before iodoethane (4.87 g, 30.9 mmol) is added. The mixture is stirred at -78°C for 1 h and is then allowed to warm to rt overnight. The reaction is quenched by the addition of water. The mixture is acidified and extracted three times with diethyl ether. The org. extracts are dried over MgSO_4 , filtered and evaporated. The crude product is purified by MPLC on reversed phase silica gel to give the title compound (1.10 g) as a brownish solid; LC-MS: $t_{\text{R}} = 0.80$ min, ^1H NMR (CDCl_3): δ 1.34 (t, $J = 7.3$ Hz, 3 H), 2.90 (q, $J = 7.6$ Hz, 2 H), 6.84 (d, $J = 3.5$ Hz, 1 H), 7.73 (d, $J = 3.8$ Hz, 1 H).

5-n-Propyl-thiophene-2-carboxylic acid




- The title compound is prepared in analogy to 5-ethyl-thiophene-2-carboxylic acid starting from 2-thiophenecarboxylic acid and 1-iodopropane; LC-MS: $t_{\text{R}} = 0.87$ min, ^1H NMR (CDCl_3): δ 0.99 (t, $J = 7.0$ Hz, 3 H), 1.74 (hex, $J = 7.3$ Hz, 2 H), 2.83 (t, $J = 7.6$ Hz, 2 H), 6.82 (d, $J = 3.5$ Hz, 1 H), 7.73 (d, $J = 3.8$ Hz, 1 H)

20 5-n-Butyl-thiophene-2-carboxylic acid



The title compound is prepared in analogy to 5-ethyl-thiophene-2-carboxylic acid starting from 2-thiophenecarboxylic acid and 1-iodobutane; LC-MS: $t_{\text{R}} = 0.92$ min.


25 5-Isobutyl-thiophene-2-carboxylic acid



To a solution of 2-thiophene-carboxylic acid (4.16 g, 32.1 mmol) in THF (200 mL) tert. butyllithium (49 mL, 1.7 M solution in pentane, 83.6 mmol) is slowly added at -


78°C. The mixture is stirred at -78°C for 30 min before isobutylbromide (22.7 g, 160.7 mmol) is carefully added. The mixture is stirred at -78°C for 5 h, then at rt for 16 h. The reaction is quenched by the addition of water (400 mL). The mixture is acidified and extracted with EA. The org. extract is dried over MgSO₄, filtered and evaporated. The crude product is purified by MPLC on reverse phase silica gel to give the title compound (1.67 g) as a brownish oil; LC-MS: t_R = 0.91 min, ¹H NMR (CDCl₃): δ 0.96 (d, J = 6.7 Hz, 6 H), 1.94 (hept, J = 6.7 Hz, 1H), 2.72 (d, J = 7.0 Hz, 2 H), 6.80 (d, J = 3.8 Hz, 1 H), 7.73 (d, J = 3.8 Hz, 1 H).

10 **5-(2-methyl-propenyl)-4-methyl-thiophene-2-carboxylic acid**




a) To a solution of methyl 5-bromo-4-methyl-thiophene-2-carboxylate (3.65 g, 15.53 mmol) in DME (30 mL), 2,4,6-tris-(2-methyl-propenyl)-cyclotriboroxane ((5.04 g, 15.53 mmol) F. Kerins, D. F. O'Shea, *J. Org. Chem.* **67** (2002), 4968-4971) followed by 2 M aq. K₂CO₃ (12 mL) is added. The solution is degassed and put under argon before Pd(PPh₃)₄ (366 mg, 0.317 mmol) is added. The mixture is stirred at 80°C for 10 h before it is cooled to rt, diluted with diethyl ether (50 mL) and washed with sat. aq. NaHCO₃-solution (2x30 mL). The organic extract is dried over MgSO₄, filtered and concentrated to give 4-methyl-5-(2-methyl-propenyl)-thiophene-2-carboxylic acid methyl ester (4.08 g) as a yellow oil; LC-MS: t_R = 1.04 min, $[M+1]^+$ = 211.04.


b) A solution of 4-methyl-5-(2-methyl-propenyl)-thiophene-2-carboxylic acid methyl ester (4.08 g, 19.4 mmol) in methanol (33 mL) is treated with 2 N aq. LiOH solution (10 mL). The mixture is stirred at rt for 2 h, then at 45°C for 3 h. The reaction mixture is diluted with diethyl ether. The organic layer is separated and extracted with water. The combined aq. phase is acidified with 1 M aq. HCl, and extracted twice with diethyl ether. The combined second organic extracts are dried over MgSO₄, filtered and concentrated to give 4-methyl-5-(2-methyl-propenyl)-thiophene-2-carboxylic acid (3.05 g) as an off-white solid; LC-MS: t_R = 0.92 min, $[M+1]^+$ = 196.98.

1-(5-Isobutyl-thiophen-2-yl)-ethanone

To a solution of 5-isobutyl-thiophene-2-carboxylic acid (550 mg, 2.99 mmol) in diethyl ether (20 mL), a solution of methyl lithium (3.75 mL, 1.6 M in diethyl ether) is slowly added at rt. The reaction mixture is stirred at rt for 1 h before it is carefully washed twice with water, dried over MgSO_4 , filtered and evaporated to give the title compound (336 mg) as a slightly yellow oil, LC-MS: $t_R = 0.91$ min, $[M+1]^+ = 183.07$.

4,N-Dihydroxy-3,5-dimethyl-benzamidine


The title compound is prepared from commercially available 4-hydroxy-3,5-dimethyl-benzonitrile according to literature procedures (e.g. E. Meyer, A. C. Joussef, H. Gallardo, *Synthesis* **2003**, 899-905); ^1H NMR (CD_3OD): δ 7.20 (s, 2H), 2.20 (s, 6H).

4-Allyloxy-N-hydroxy-3,5-dimethyl-benzamidine

The title compound is prepared by allylating commercially available 4-hydroxy-3,5-dimethyl-benzonitrile with allylbromide in the presence of NaOH in isopropanol at rt.


The nitrile is then transformed to the hydroxyamidine according to literature procedures (e.g. E. Meyer, A. C. Joussef, H. Gallardo, *Synthesis* **2003**, 899-905); ^1H NMR (CD_3OD): δ 7.27 (s, 2 H), 6.10 (m, 1 H), 5.42 (m, 1 H), 5.26 (m, 1 H), 4.31 (dt, $J = 5.6, 1.5$ Hz, 2 H), 2.29 (s, 6 H).

3-Ethyl-4,N-dihydroxy-5-methyl-benzamidine




The title compound is prepared from commercially available 2-ethyl-6-methyl-phenol following literature procedures (G. Trapani, A. Latrofa, M. Franco, C. Altomare, E. Sanna, M. Usala, G. Biggio, G. Liso, *J. Med. Chem.* **41** (1998) 1846-1854; A. K. Chakraborti, G. Kaur, *Tetrahedron* **55** (1999) 13265-13268; E. Meyer, A. C. Joussef, H. Gallardo, *Synthesis* **2003**, 899-905); LC-MS: t_R = 0.55 min; 1H NMR (D_6 -DMSO): δ 9.25 (s br, 1H), 7.21 (s, 2H), 5.56 (s, 2H), 2.55 (q, J = 7.6 Hz, 2H), 2.15 (s, 3H), 1.10 (t, J = 7.6 Hz, 3H).

10 4-Allyloxy-3-ethyl-N-hydroxy-5-methyl-benzamidine



The title compound is prepared by allylating 3-ethyl-4-hydroxy-5-methyl-benzaldehyde which is prepared from 2-ethyl-6-methyl-phenol following literature procedures (see 3-ethyl-4,N-dihydroxy-5-methyl-benzamidine). The aldehyde is then transformed into the corresponding hydroxyamidine according to literature procedures (see 3-ethyl-4,N-dihydroxy-5-methyl-benzamidine); LC-MS: t_R = 0.72 min, $[M+1]^+$ = 235.09; 1H NMR(CD_3OD): δ 7.31 (s, 1 H), 7.29 (s, 1 H), 6.10 (m, 1 H), 5.43 (dd, J = 17.0, 1.5 Hz, 1 H), 5.27 (dd, J = 10.3, 1.2 Hz, 1 H), 4.81 (s br, 3H), 4.31 (d, J = 5.6 Hz, 2 H), 2.67 (q, J = 7.6 Hz, 2 H), 2.30 (s, 3 H), 1.23 (t, J = 7.6 Hz, 4 H).


3,5-Diethyl-4,N-dihydroxy-benzamidine



The title compound is prepared from commercially available 2,6-diethylaniline following literature procedures (G. G. Ecke, J. P. Napolitano, A. H. Filbey, A. J. Kolka, J. Org. Chem. **22** (1957) 639-642; and literature cited for 3-ethyl-4,N-dihydroxy-5-methyl-benzamidine).


5

4-Allyloxy-N-hydroxy-2-methoxy-benzamidine



The title compound is prepared from commercially available 4-hydroxy-2-methoxy-benzaldehyde following literature procedures (references cited for 3-ethyl-4,N-dihydroxy-5-methyl-benzamidine); LC-MS: $t_R = 0.64$ min; $[M+1]^+ = 223.24$; ^1H NMR ($\text{D}_6\text{-DMSO}$): δ 9.33 (s br, 1H), 7.30 (d, $J = 8.2$ Hz, 1H), 6.60 (d, $J = 2.3$ Hz, 1H), 6.50 (dd, $J = 2.3, 8.2$ Hz, 1H), 6.10-5.94 (m, 1H), 5.50 (s, 2H), 5.40 (d, $J = 17.0$ Hz, 1H), 5.24 (d, $J = 10.6$ Hz, 1H), 4.57 (d, $J = 4.7$ Hz, 2H), 3.76 (s, 3H).

15 4-Allyloxy-3,5-dimethyl-benzoic acid hydrazide




a) A mixture of 4-bromo-2,6-dimethyl-phenol (20.1 g, 100 mmol) and allylchloride (32.7 g, 428 mmol) in 3 N aq. NaOH (100 mL) and isopropanol (250 mL) is stirred at 60°C for 15 h before it is diluted with 1 N aq. NaOH (100 mL). The mixture is extracted with diethyl ether (300 mL, 150 mL) and the combined org. extracts are washed with 1 N aq. NaOH (2x100 mL), 1 M aq. NaH_2PO_4 (50 mL), dried over Na_2SO_4 , filtered and concentrated to give 2-allyloxy-5-bromo-1,3-dimethyl-benzene (23.6 g) as a yellow oil, LC-MS: $t_R = 1.08$ min, $[M+1]^+ = 241.20$.

b) To a solution of 2-allyloxy-5-bromo-1,3-dimethyl-benzene (23.6 g, 98.0 mmol) in THF (150 mL) is added at -75°C a solution of n-BuLi (90 mL, 1.5 M in diethyl ether). The temperature remains at -75°C. The mixture is stirred for 30 min and then
 5 transferred via double-tip canula into a cooled (0°C) solution of dimethylcarbonate (21.4 g, 238 mmol) in THF (90 mL). The mixture is stirred for 2 h at 0°C, then warmed to rt during 15 h. The solvent of the mixture is evaporated and re-evaporated from ethanol (200 mL) to remove most of the butylacetate side product. The mixture is taken up in 2 N aq. LiOH (150 mL) and EtOH (200 mL) and stirred at
 10 rt for 2 h, then at 60°C for 1 h. The ethanol is evaporated and the remaining mixture is diluted with 0.5 N aq. NaOH and extracted with diethyl ether (200 mL). The org. extract is washed with 1M aq. NaOH (5 x 50 mL) and the combined aq. washings are re-extracted with ether (100 mL). The aq. phase is acidified with 25% aq. HCl and extracted with DCM (5 x 50 mL). The combined org. extracts are dried over
 15 Na₂SO₄, filtered, evaporated and dried in vacuo at 60°C for 15 h to give 4-allyloxy-3,5-dimethyl-benzoic acid (8.0) as yellow-brown solid. LC-MS: *t_R* = 0.90 min.

c) To a solution of 4-allyloxy-3,5-dimethyl-benzoic acid (5.26 g, 25.5 mmol) in CHCl₃ (75 mL), thionylchloride (7.5 mL, 103 mmol) is added at rt. The mixture is
 20 refluxed for 2 h before the solvent is evaporated to give crude 4-allyloxy-3,5-dimethyl-benzoic acid chloride as a brownish oil. To a solution of the acid chloride in DCM (50 mL), hydrazine (75 mL of a 1 M solution in THF) in DCM (250 mL) is added at 0°C. The mixture is stirred at rt for 15 h before it is diluted with diethyl ether and extracted with 1 N aq. HCl (75 mL, then 5x50 mL). The combined aq.
 25 extracts are basified by adding 33% aq. KOH solution and extracted with DCM (5x50 mL). The combined DCM extracts are dried over Na₂SO₄, filtered and evaporated to give 4-allyloxy-3,5-dimethyl-benzoic acid hydrazide (5.39 g) as a white solid; LC-MS: *t_R* = 0.71 min; [M+1]⁺ = 221.20; ¹H NMR (D₆-DMSO): δ 2.22 (s, 6 H), 4.28-4.37 (m, 2 H), 4.39 (s, 2 H), 5.19-5.28 (m, 1 H), 5.36-5.47 (m, 1 H), 6.00-
 30 6.15 (m, 1 H), 7.49 (s, 2 H), 9.55 (s, 1 H).


4-Benzyloxy-3,5-dimethyl-benzoic acid hydrazide



4-Benzyloxy-3,5-dimethylbenzoic acid hydrazide is prepared in analogy to 4-allyloxy-3,5-dimethylbenzoic acid hydrazide starting from 4-benzyloxy-3,5-dimethylbenzoic acid; LC-MS: $t_R = 0.81$ min; $[M+1]^+ = 271.41$.

5

4-Benzyloxy-3-ethyl-5-methylbenzoic acid hydrazide




- a) To a solution of 3-ethyl-4-hydroxy-5-methylbenzaldehyde (34.9 g, 0.213 mol, prepared from 2-ethyl-6-methylphenol according to the literature cited for 3-ethyl-4,N-dihydroxy-5-methylbenzamidine) in acetonitrile (350 mL), K_2CO_3 (58.7 g, 0.425 mol) and benzylbromide (36.4 g, 0.213 mol) is added. The mixture is stirred at 60°C for 2 h before it is cooled to rt, diluted with water and extracted twice with EA. The organic extracts are washed with water and concentrated to give crude 4-benzyloxy-3-ethyl-5-methylbenzaldehyde (45 g) as an orange oil. 1H NMR ($CDCl_3$):
- 10 δ 1.29 (t, $J = 7.5$ Hz, 3 H), 2.40 (s, 3 H), 2.77 (q, $J = 7.8$ Hz, 2 H), 4.90 (s, 2 H), 7.31-7.52 (m, 5 H), 7.62 (d, $J = 1.5$ Hz, 1 H), 7.66 (d, $J = 1.8$ Hz, 1 H), 9.94 (s, 1 H).

- b) To a mixture of 4-benzyloxy-3-ethyl-5-methylbenzaldehyde (132 g, 0.519 mol) and 2-methyl-2-butene (364 g, 5.19 mol) in tert.-butanol (1500 mL), a solution of NaH_2PO_4 dihydrate (249 g, 2.08 mol) in water (1500 mL) is added. To this mixture, $NaClO_2$ (187.8 g, 2.08 mol) is added in portions. The temperature of the reaction mixture is kept below 30°C, and evolution of gas is observed. Upon completion of the addition, the orange bi-phasic mixture is stirred well for 3 h before it is diluted with TBME (1500 mL). The organic layer is separated and washed with 20% aq.
- 20

NaHS solution (1500 mL) and water (500 mL). The organic phase is then extracted three times with 0.5 N aq. NaOH (1000 mL), the aqueous phase is acidified with 25 % aq. HCl (500 mL) and extracted twice with TBME (1000 mL). These organic extracts are combined and evaporated to dryness to give 4-benzyloxy-3-ethyl-5-methyl-benzoic acid; ^1H NMR (D_6 -DMSO): δ 1.17 (t, $J = 7.5$ Hz, 3 H), 2.31 (s, 3 H), 2.67 (q, $J = 7.5$ Hz, 2 H), 4.86 (s, 2 H), 7.34-7.53 (m, 5 H), 7.68 (s, 2 H), 12.70 (s, 1 H).


c) 4-Benzyloxy-3-ethyl-5-methyl-benzoic acid is converted to 4-benzyloxy-3-ethyl-5-methyl-benzoic acid hydrazide following step c) of the preparation of 4-allyloxy-3,5-dimethyl-benzoic acid hydrazide; LC-MS: $t_{\text{R}} = 0.82$ min, $[\text{M}+1]^+ = 285.44$.

Methanesulfonic acid 2,2-dimethyl-[1,3]dioxan-5-ylmethyl ester



The title compound is prepared following the procedures given in B. Xu, A. Stephens, G. Kirschenheuter, A. F. Greslin, X. Cheng, J. Sennelo, M. Cattaneo, M. L. Zighetti, A. Chen, S.-A. Kim, H. S. Kim, N. Bischofberger, G. Cook, K. A. Jacobson, *J. Med. Chem.* **45** (2002) 5694-5709.

Example 1




a) To a solution of 5-ethyl-thiophene-2-carboxylic acid (502 mg, 3.21 mmol) and DIPEA (1.04 g, 8.03 mmol) in DCM (16 mL), TBTU (1.13 g, 3.53 mmol) is added at 0°C . The mixture is stirred at 0°C for 1 h before the reaction is quenched with water (2 mL). The DCM is evaporated and the remaining residue is diluted with EA, washed with sat. aq. NaHCO_3 solution and water, dried over MgSO_4 , filtered and evaporated. The crude product was purified by crystallisation from acetonitrile to

give 5-ethyl-thiophene-2-carboxylic acid (4,N-dihydroxy-3,5-dimethyl-benzamidine) ester (600 mg) as a white solid; LC-MS: $t_R = 0.94$ min, $[M+1]^+ = 319.02$.


b) A solution of 5-ethyl-thiophene-2-carboxylic acid (4,N-dihydroxy-3,5-dimethyl-benzamidine) ester (600 mg, 1.88 mmol) in dioxane (40 mL) is stirred at 100°C for 18 h. The solvent is evaporated and the residue is separated by CC on silica gel eluting with heptane:EA 4:1 to give 4-[5-(5-ethyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-2,6-dimethyl-phenol (525 mg) as a pale yellow oil; LC-MS: $t_R = 1.09$ min, $[M+1]^+ = 301.11$.

Example 2




2,6-Dimethyl-4-[5-(5-propyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenol is prepared in analogy to Example 1; LC-MS: $t_R = 1.10$ min, $[M+1]^+ = 315.35$.

Example 3




4-[5-(5-Butyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-2,6-dimethyl-phenol is prepared in analogy to Example 1; LC-MS: $t_R = 1.15$ min, $[M+1]^+ = 329.15$.


Example 4



4-[5-(5-Isobutyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-2,6-dimethyl-phenol is prepared in analogy to Example 1; LC-MS: $t_R = 1.15$ min, $[M+1]^+ = 329.09$.







Example 5

To a solution of 4-[5-(5-ethyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-2,6-dimethyl-phenol (79 mg, 0.263 mmol) in isopropanol (3 mL) and 3 N aq. NaOH (0.6 mL), (R)-3-chloro-propane-1,2-diol (148 mg, 1.31 mmol) is added. The mixture is stirred at rt for 2 h, then at 65°C for 16 h before another portion of (R)-3-chloro-propane-1,2-diol (119 mg, 1.05 mmol) is added. Stirring is continued at 65°C for 24 h. The reaction mixture is diluted with water and extracted with diethyl ether. The org. extract is dried over MgSO₄, filtered and evaporated. The crude product is purified by chromatography on prep. TLC plates with DCM containing 4% of 7 N NH₃ in methanol to give (2R)-3-{4-[5-(5-ethyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-2,6-dimethyl-phenoxy}-propane-1,2-diol as an off-white solid; LC-MS: *t_R* = 0.99 min; [M+1]⁺ = 375.13; ¹H NMR (CDCl₃): δ 7.83 (s, 2H), 7.80 (d, *J* = 3.8 Hz, 1H), 6.93 (d, *J* = 3.8 Hz, 1H), 4.20-4.11 (m, 1H), 3.97-3.93 (m, 1H), 3.91 (dd, *J* = 11.5, 4.3 Hz, 1H), 3.85 (dd, *J* = 11.5, 5.5 Hz, 1H), 2.97 (q, *J* = 7.5 Hz, 2 H), 2.39 (s, 6 H), 1.41 (t, *J* = 7.5 Hz, 3H).

Examples 6 to 12


The following Examples are prepared in analogy to Example 5 starting from the Examples indicated using either (R)- or (S)-3-chloro-propane-1,2-diol:

Example	starting from Example	R ^a	R ^b	LC-MS	
				<i>t_R</i> (min)	[M+H] ⁺
6	1	ethyl		0.99	375.01

7	2	n-propyl		1.03	389.20
8	2	n-propyl		1.00	389.38
9	3	n-butyl		1.04	403.40
10	3	n-butyl		1.06	403.11
11	4	isobutyl		1.03	403.40
12	4	isobutyl		1.06	403.05

Example 12

^1H NMR(CDCl_3): δ 0.98 (d, J = 6.4 Hz, 6 H), 1.96 (hept, J = 6.5 Hz, 1H), 2.35 (s, 6 H), 2.76 (d, J = 7.0 Hz, 2 H), 3.76-3.94 (m, 4 H), 4.09-4.18 (m, 1 H), 6.86 (d, J = 3.5 Hz, 1 H), 7.77 (d, J = 3.5 Hz, 1 H), 7.80 (s, 2 H).


Example 13

a) A mixture of 4-[5-(5-ethyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-2,6-dimethyl-phenol (367 mg, 1.22 mmol) and epichlorohydrine (565 mg, 6.10 mmol) in isopropanol (20 mL) and 3 N aq. NaOH (6 mL) is stirred at 40°C for 15 h. The mixture is diluted with diethyl ether, washed with sat. aq. NaHCO_3 and water, dried over MgSO_4 , filtered and evaporated. The crude product is purified by CC on silica gel eluting with heptane:EA 9:1 to give 3-(3,5-dimethyl-4-oxiranylmethoxy-phenyl)-5-(5-ethyl-thiophen-2-yl)-[1,2,4]oxadiazole (123 mg) as a pale yellow oil; LC-MS: t_R = 1.15 min, $[\text{M}+1]^+$ = 357.11.

b) A solution of 3-(3,5-dimethyl-4-oxiranylmethoxy-phenyl)-5-(5-ethyl-thiophen-2-yl)-[1,2,4]oxadiazole (123 mg, 0.345 mmol) in 7 N NH_3 in methanol (10 mL) is stirred at 45°C for 16 h. The solvent is evaporated and the residue is separated by chromatography on prep. TLC plates with DCM containing 6% of 7 N NH_3 in methanol to give 1-amino-3-{4-[5-(5-ethyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-2,6-dimethyl-phenoxy}-propan-2-ol (35 mg) as a white solid; LC-MS: t_R = 0.84 min, $[\text{M}+1]^+ = 374.14$.

c) To a cold solution (0°C) of 1-amino-3-{4-[5-(5-ethyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-2,6-dimethyl-phenoxy}-propan-2-ol (32 mg, 86 μmol) in DCM (1 mL), DIPEA (45 mg, 345 μmol), glycolic acid (13 mg, 172 μmol) and finally TBTU (33 mg, 101 μmol) is added. The mixture is stirred at rt for 1 h before it is diluted with EA and washed with water. The aq. phase is extracted back with EA. The combined org. extracts are dried over MgSO_4 , filtered and evaporated. The crude product is purified by crystallisation from acetonitrile to give N-(3-{4-[5-(5-ethyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-2,6-dimethyl-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide (7 mg) as a white solid; LC-MS: t_R = 0.95 min, $[\text{M}+1]^+ = 432.16$.

Examples 14 to 16



The following Examples are prepared in analogy to Example 13 starting from the Example indicated:

Example	starting from Example	R^a	LC-MS	
			t_R (min)	$[\text{M}+\text{H}]^+$
14	2	n-propyl	0.99	446.14
15	3	n-butyl	1.02	460.26
16	4	isobutyl	1.01	460.18


Example 15

¹H NMR(D₆-DMSO): δ 0.93 (t, *J* = 7.3 Hz, 3 H), 1.33-1.44 (m, 2 H), 1.63-1.72 (m, 2 H), 2.33 (s, 6 H), 2.93 (t, *J* = 7.5 Hz, 2 H), 3.20-3.30 (m, 1 H), 3.37-3.48 (m, 1 H), 3.69-3.80 (m, 2 H), 3.84 (d, *J* = 5.8 Hz, 2 H), 3.91-3.99 (m, 1 H), 5.30 (d, *J* = 5.3 Hz, 1 H), 5.56 (t, *J* = 5.8 Hz, 1 H), 7.12 (d, *J* = 3.8 Hz, 1 H), 7.69 (t, *J* = 6.0 Hz, 1 H), 7.72 (s, 2 H), 7.90 (d, *J* = 3.5 Hz, 1 H).

Example 16


¹H NMR (D₆-DMSO): δ 0.92 (d, *J* = 6.4 Hz, 5 H), 1.83-2.00 (m, 1 H), 2.30 (s, 6 H), 2.78 (d, *J* = 7.0 Hz, 2 H), 3.14-3.27 (m, 2 H), 3.36-3.48 (m, 2 H), 3.65-3.78 (m, 2 H), 3.81 (d, *J* = 5.6 Hz, 2 H), 3.87-3.99 (m, 1 H), 5.26 (d, *J* = 5.0 Hz, 1 H), 5.52 (t, *J* = 5.6 Hz, 1 H), 7.07 (d, *J* = 3.5 Hz, 1 H), 7.66 (m, 3 H), 7.89 (d, *J* = 3.5 Hz, 1 H).

Example 17



(2R)-N-(3-{2,6-Dimethyl-4-[5-(5-propyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide is prepared in analogy to Example 13 starting from Example 2 and using (R)-2-chloromethyl-oxirane as the alkylating agent; LC-MS: *t_R* = 0.98 min, [M+1]⁺ = 446.20.


Example 18



(2S)-N-(3-{2,6-Dimethyl-4-[5-(5-propyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide is prepared in analogy to Example 13 starting from Example 2 and using (S)-2-chloromethyl-oxirane as the alkylating agent; LC-MS: *t_R* = 0.98 min, [M+1]⁺ = 446.19, ¹H NMR (CDCl₃): δ 1.04 (t, *J* = 7.3 Hz, 3 H), 1.72-1.84 (m, 2 H), 2.36 (s, 6 H), 2.89 (t, *J* = 7.5 Hz, 2 H), 3.19 (s

br, 1H), 3.47-3.56 (m, 1 H), 3.60 (s br, 1 H), 3.73-3.92 (m, 3 H), 4.18 (s, 2 H), 6.90 (d, $J = 3.8$ Hz, 1 H), 7.12 (t, $J = 5.8$ Hz, 1 H), 7.78 (d, $J = 3.8$ Hz, 1 H), 7.80 (s, 2 H)


Example 19



5


2-Ethyl-4-[5-(5-isobutyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-6-methyl-phenol is prepared in analogy to Example 1 using 3-ethyl-4,N-dihydroxy-5-methyl-benzamidine; LC-MS: $t_R = 1.16$ min, $[M+1]^+ = 343.02$.

10 Example 20



N-(3-{2-Ethyl-4-[5-(5-isobutyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-6-methyl-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide is prepared in analogy to Example 13 starting from Example 19; LC-MS: $t_R = 1.01$ min; 1H NMR ($CDCl_3$): δ 0.99 (d, $J = 6.7$ Hz, 3 H), 1.29 (t, $J = 7.6$ Hz, 3 H), 1.90-2.05 (m, 1 H), 2.35 (s, 3 H), 2.57 (s br, 1H), 2.67-2.80 (m, 4 H), 3.23 (s br, 1H), 3.44-3.56 (m, 1 H), 3.72-3.92 (m, 3 H), 4.14-4.22 (m, 3H), 6.87 (d, $J = 3.5$ Hz, 1 H), 6.97 (t br, 1 H), 7.78 (d, $J = 3.8$ Hz, 1 H), 7.82 (s, 1H), 7.84 (s, 1H).

20 Example 21




a) To a solution of 5-isobutyl-thiophene-2-carboxylic acid (830 mg, 4.51 mmol) and DIPEA (680 mg, 5.27 mmol) in DCM (20 mL) is added TBTU (1.59 g, 4.96 mmol) at rt. The mixture is stirred at rt for 45 min before 4-allyloxy-3,5-dimethyl-benzoic acid

hydrazide (993 mg, 4.50 mmol) is added. Stirring is continued for 2 h. The mixture is diluted with ether (200 mL) and washed with 1M aq. HCl (3 x 50 mL), 1M aq. NaOH (3 x 50 mL) and brine (50 mL). The org. extract is dried over MgSO_4 , filtered and evaporated to give the crude 4-allyloxy-3,5-dimethyl-benzoic acid N'-(5-isobutyl-thiophene-2-carbonyl)-hydrazide (1.40 g) as a yellow oil that slowly solidifies; LC-MS: $t_R = 1.02$ min, $[\text{M}+1]^+ = 387.10$.

b) A solution of 4-allyloxy-3,5-dimethyl-benzoic acid N'-(5-isobutyl-thiophene-2-carbonyl)-hydrazide (1.40 g, 3.62 mmol) and Burgess reagent (1.12 g, 4.71 mmol) in THF (15 mL) is stirred at 110°C for 3 min under microwave irradiation. The mixture is diluted with diethyl ether, washed with water, dried over MgSO_4 , filtered and evaporated. The crude product is purified by CC on silica gel eluting with heptane:EA 9:1 to give 2-(4-allyloxy-3,5-dimethyl-phenyl)-5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazole (1.02 g) as a colourless oil; LC-MS: $t_R = 1.21$ min, $[\text{M}+1]^+ = 369.15$.

c) To a solution of 2-(4-allyloxy-3,5-dimethyl-phenyl)-5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazole (1.02 g, 2.77 mmol) in acetone (24 mL) and water (2.4 mL), NMO (1.85 g, 13.7 mmol) and OsO_4 (128 mg, $13\ \mu\text{mol}$, as a 2.5% solution in butanol) is added at rt. The mixture is stirred at rt for 16 h before it is diluted with water (50 mL) and extracted with EA (2x 100 mL). The combined org. extracts are dried over Na_2SO_4 , filtered and evaporated. The crude product is purified by CC on silica gel eluting with EA to give 3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propane-1,2-diol (680 mg) as a colourless oil; LC-MS: $t_R = 0.99$ min, $[\text{M}+1]^+ = 403.14$.

Example 22



a) To a solution of 3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propane-1,2-diol (627 mg, 1.56 mmol) and DIPEA (311 mg, 2.41


mmol) in DCM (10 mL), methanesulfonyl chloride (207 mg, 1.81 mmol) is added. The mixture is stirred at rt before another portion of DIPEA (40 mg, 0.31 mmol) and methanesulfonyl chloride (36 mg, 0.31 mmol) is added. Stirring is continued for 1 h before the reaction is quenched by adding water. The mixture is extracted with
 5 DCM. The org. extract is washed with water, dried over MgSO_4 , filtered and evaporated. The crude product is purified by CC on silica gel eluting with heptane:EA 7:3 to give methanesulfonic acid 2-hydroxy-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propyl ester (220 mg) as a colourless oil; LC-MS: $t_R = 1.08$ min, $[\text{M}+1]^+ = 481.09$.

10

b) A solution of methanesulfonic acid 2-hydroxy-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propyl ester (220 mg, 0.458 mmol) in THF (10 mL) and 7 N NH_3 in methanol (10 mL) is stirred at 65°C for 15 h. The solvent is evaporated and the residue is separated by chromatography on prep.
 15 TLC plates with DCM containing 6% of 7 N NH_3 in methanol to give 1-amino-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propan-2-ol (110 mg) as a yellow oil; LC-MS: $t_R = 0.83$ min, $[\text{M}+1]^+ = 402.48$.


c) To a solution of 1-amino-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propan-2-ol (109 mg, 271 μmol) in DCM (10 mL), DIPEA (140 mg, 1.086 mmol) and glycolic acid (41 mg, 543 μmol) followed by TBTU (102 mg, 319 μmol) is added at 0°C . The mixture is stirred at rt for 1 h before it is diluted with EA and washed with water. The water phase is separated and extracted once more with EA. The combined org. extracts are dried over MgSO_4 , filtered and
 25 evaporated. The crude product is purified by chromatography on prep. TLC plates with DCM containing 4% of 7 N NH_3 in methanol to give 2-hydroxy-N-(2-hydroxy-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propyl)-acetamide (31 mg) as a white solid; LC-MS: $t_R = 0.94$ min, $[\text{M}+1]^+ = 460.51$.

30 Example 23




To a solution of 1-(5-isobutyl-thiophen-2-yl)-ethanone (336 mg, 1.84 mmol) and 3,5-dimethyl-4-hydroxy-benzaldehyde (277 mg, 1.84 mmol) in ethanol (6 mL), 5 N HCl in isopropanol (2.5 mL) is added. The dark solution is stirred at rt for 6 h before the solvent is removed in vacuo. The residue is dissolved in EA (80 mL) and washed with sat. aq. NaHCO₃ solution (25 mL) and brine. The washings are extracted back with EA. The combined org. extracts are dried over MgSO₄, filtered and evaporated to give 3-(4-hydroxy-3,5-dimethyl-phenyl)-1-(5-isobutyl-thiophen-2-yl)-propenone (682 mg) as a yellow powder; LC-MS: t_R = 1.10 min, $[M+1]^+$ = 315.15.

Example 24




To a solution of 3-(4-hydroxy-3,5-dimethyl-phenyl)-1-(5-isobutyl-thiophen-2-yl)-propanone (660 mg) in THF (10 mL) and ethanol (5 mL), Pd/C (150 mg, 10% Pd) is added and the suspension is stirred at rt under 5 bar of H₂ for 23 h. Another portion of Pd/C (150 mg) is added and stirring is continued at rt under 5 bar of H₂ for 8 h. The catalyst is filtered off and the filtrate is evaporated to give 3-(4-hydroxy-3,5-dimethyl-phenyl)-1-(5-isobutyl-thiophen-2-yl)-propan-1-one (554 mg) as an orange oil; LC-MS: t_R = 1.10 min, $[M+1]^+$ = 317.04.

Example 25




2-Hydroxy-N-(2-hydroxy-3-{4-[3-(5-isobutyl-thiophen-2-yl)-3-oxo-propyl]-2,6-dimethyl-phenoxy}-propyl)-acetamide is prepared from Example 24 in analogy to Example 13; LC-MS: $t_R = 0.97$ min, $[M+1]^+ = 448.21$.

5 Example 26




2-Ethyl-6-methyl-4-[5-(5-propyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenol is prepared in analogy to Example 1 starting from 5-propyl-thiophene-2-carboxylic acid and 3-ethyl-4,N-dihydroxy-5-methyl-benzamidine; LC-MS: $t_R = 1.12$ min, $[M+1]^+ = 329.14$.

Example 27




(2R)-N-(3-{2-Ethyl-6-methyl-4-[5-(5-propyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide is prepared in analogy to Example 13 starting from Example 26; LC-MS: $t_R = 0.95$ min, $[M+1]^+ = 460.15$.


Example 28




(2S)-N-(3-{2-Ethyl-6-methyl-4-[5-(5-propyl-thiophen-2-yl)-[1,2,4]oxadiazol-3-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide is prepared in analogy to Example 13 starting from Example 26; LC-MS: $t_R = 0.95$ min, $[M+1]^+ = 460.13$.

Example 29

2-(4-Benzyloxy-3,5-dimethyl-phenyl)-5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazole is prepared in analogy to Example 21 step a) and b) by coupling and cyclising 5-isobutyl-thiophene-2-carboxylic acid with 4-benzyloxy-3,5-dimethyl-benzoic acid hydrazide, LC-MS: $t_R = 1.06$ min, $[M+1]^+ = 437.17$. To a solution of this compound (2.83 g, 6.75 mmol) in ethanol (50 mL) and THF (50 mL) is added Pd/C (10% Pd, 400 mg) and the slurry is stirred at rt for 48 h under 5 bar of H_2 . The mixture is filtered, the filtrate is concentrated and the crude product is purified by CC on silica gel eluting with heptane:EA 8:25 to give 4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenol (943 mg) as a white powder; LC-MS: $t_R = 1.09$ min, $[M+1]^+ = 329.36$.

Example 30


Starting from 4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenol, 2-hydroxy-N-((2R)-2-hydroxy-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propyl)-acetamide is prepared in analogy to Example 13; LC-MS: $t_R = 0.95$ min, $[M+1]^+ = 460.18$.

Example 31

Starting from 4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenol,


2-hydroxy-N-((2S)-2-hydroxy-3-{4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-2,6-dimethyl-phenoxy}-propyl)-acetamide is prepared in analogy to Example 13; LC-MS: $t_R = 0.95$ min, $[M+1]^+ = 460.18$, 1H NMR δ 0.99 (d, $J = 6.5$ Hz, 6 H), 1.90-2.01 (m, 1 H), 2.31 (s, 6 H), 2.75 (d, $J = 7.0$ Hz, 2 H), 3.46-3.54 (m, 1 H), 3.73-3.91 (m, 3 H), 4.15-4.23 (m, 3 H), 4.26 (s br, 1 H), 4.45 (s br, 1 H), 6.84 (d, $J = 3.3$ Hz, 1 H), 7.42 (t br, $J = 5.5$ Hz, 1 H), 7.64 (d, $J = 3.5$ Hz, 1 H), 7.68 (s, 2 H).

10 Example 32




2-Ethyl-6-methyl-4-[5-(5-propyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-phenol is prepared in analogy to Example 29 starting from 4-benzyloxy-3-ethyl-5-methylbenzoic acid hydrazide and 5-propyl-thiophene-2-carboxylic acid; LC-MS: $t_R = 0.99$ min, $[M+1]^+ = 329.13$.

Example 33




N-((2R)-3-{2-Ethyl-6-methyl-4-[5-(5-propyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide is prepared in analogy to Example 13 from Example 32; LC-MS: $t_R = 0.85$ min, $[M+1]^+ = 460.08$.

Example 34




N-((2S)-3-{2-Ethyl-6-methyl-4-[5-(5-propyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide is prepared in analogy to Example 13 from Example 32; LC-MS: $t_R = 0.85$ min, $[M+1]^+ = 460.08$.

5 Example 35




2-Ethyl-4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-6-methyl-phenol is prepared in analogy to Example 29 from 5-isobutyl-thiophene-2-carboxylic acid and 4-benzyloxy-3-ethyl-5-methyl-benzoic acid hydrazide; LC-MS: $t_R = 1.03$ min, $[M+1]^+ = 343.22$.

Example 36




N-((2R)-3-{2-Ethyl-6-methyl-4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide is prepared in analogy to Example 13 from Example 35; LC-MS: $t_R = 0.91$ min, $[M+1]^+ = 474.18$.

Example 37



N-((2S)-3-{2-Ethyl-6-methyl-4-[5-(5-isobutyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide is prepared in analogy to Example 13 from Example 35; LC-MS: $t_R = 0.91$ min, $[M+1]^+ = 474.17$.


Example 38



2-Ethyl-4-[5-(5-isobutyl-4-methyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-6-methylphenol is prepared from 4-methyl-5-(2-methyl-propenyl)-thiophene-2-carboxylic acid and 4-benzyloxy-3-ethyl-5-methyl-benzoic acid hydrazide in analogy to Example 29;

5 LC-MS: $t_R = 1.13$ min, $[M+1]^+ = 357.49$.


Example 39



10 (2R)-3-{2-Ethyl-4-[5-(5-isobutyl-4-methyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-6-methyl-phenoxy}-propane-1,2-diol is obtained as a white solid from Example 38 in analogy to Example 5; LC-MS: $t_R = 1.05$ min, $[M+1]^+ = 431.13$; 1H NMR ($CDCl_3$): δ 1.01 (d, $J = 6.8$ Hz, 6 H), 1.31 (t, $J = 7.5$ Hz, 3 H), 1.91-2.02 (m, 1 H), 2.22 (s, 3 H), 2.39 (s, 3 H), 2.68 (d, $J = 7.0$ Hz, 2 H), 2.75 (q, $J = 7.3$ Hz, 2 H), 3.82-3.97 (m, 4 H), 4.14-4.22 (m, 1 H), 7.55 (s, 1 H), 7.78 (s, 1 H), 7.80 (s, 1 H).


15

Example 40



20 (2S)-3-{2-Ethyl-4-[5-(5-isobutyl-4-methyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-6-methyl-phenoxy}-propane-1,2-diol is obtained as a white solid from Example 38 in analogy to Example 5; LC-MS: $t_R = 1.05$ min, $[M+1]^+ = 431.14$.


Example 41



N-((2R)-3-{2-Ethyl-4-[5-(5-isobutyl-4-methyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-6-methyl-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide is prepared from Example 38 in analogy to Example 13; LC-MS: $t_R = 1.00$ min, $[M+1]^+ = 488.19$.

5

Example 42



N-((2S)-3-{2-Ethyl-4-[5-(5-isobutyl-4-methyl-thiophen-2-yl)-[1,3,4]oxadiazol-2-yl]-6-methyl-phenoxy}-2-hydroxy-propyl)-2-hydroxy-acetamide is prepared from Example 38 in analogy to Example 13; LC-MS: $t_R = 1.00$ min, $[M+1]^+ = 488.17$; ^1H NMR δ 1.00 (d, $J = 6.5$ Hz, 6 H), 1.27 (t, $J = 7.5$ Hz, 3 H), 1.90-2.01 (m, 1 H), 2.21 (s, 3 H), 2.34 (s, 3 H), 2.67 (d, $J = 7.3$ Hz, 2 H), 2.71 (q, $J = 7.5$ Hz, 2H), 3.46-3.56 (m, 1 H), 3.75-3.91 (m, 3 H), 4.08 (s br, 2 H), 4.20 (s, 3 H), 7.33 (t br, $J = 5.5$ Hz, 1 H), 7.54 (s, 1 H), 7.72 (s, 1 H), 7.74 (s, 1 H).

15

Example 43: GTP γ S assay to determine EC₅₀ values

GTP γ S binding assays are performed in 96 well microtiter plates (Nunc, 442587) in a final volume of 200 μ l, using membrane preparations of CHO cells expressing recombinant human S1P1 receptor. Assay conditions are 20 mM Hepes (Fluka, 54461), 100 mM NaCl (Fluka, 71378), 5 mM MgCl₂ (Fluka, 63064), 0.1% BSA (Calbiochem, 126609), 1 μ M GDP (Sigma, G-7127), 2.5% DMSO (Fluka, 41644), 50 pM ^{35}S -GTP γ S (Amersham Biosciences, SJ1320). The pH is 7.4. Test compounds are dissolved and diluted in 100% DMSO and pre-incubated at room temperature for 30 min in 150 μ l of the above assay buffer, in the absence of ^{35}S -GTP γ S. After addition of 50 μ l of ^{35}S -GTP γ S, the assay is incubated for 1 h at rt. The assay is terminated by transfer of the reaction mixture to a Multiscreen plate (Millipore, MAHFC1H60) using a cell harvester from Packard Biosciences, and the

25

plates are washed with ice-cold 10 mM Na₂HPO₄/NaH₂PO₄ (70%/30%), dried, sealed at the bottom and, after addition of 25 µl MicroScint20 (Packard Biosciences, order# 6013621), sealed on the top. Membrane-bound ³⁵S-GTPγS is measured with a TopCount from Packard Biosciences.

5

EC₅₀ is the concentration of agonist inducing 50 % of the maximal specific ³⁵S-GTPγS binding. Specific binding is determined by subtracting non-specific binding from maximal binding. Maximal binding is the amount of cpm bound to the Multiscreen plate in the presence of 10 µM of S1P. Non-specific binding is the

10

amount of binding in the absence of an agonist in the assay.

Table 1 shows the EC₅₀ value of some compounds of the present invention. The EC₅₀ values were determined according to the method described above:

15 **Table 1:**

Compound of Example	EC ₅₀ [nM]
16	2.6
17	0.6
22	2.8
28	2.4
31	3.3
42	0.6

Example 44: Assessment of *In vivo* Efficacy

The efficacy of the compounds of Formula (I) is assessed by measuring the circulating lymphocytes after oral administration of 3 to 30 mg/kg of a compound of Formula (I) to normotensive male Wistar rats. The animals are housed in climate-controlled conditions with a 12 h-light/dark cycle, and have free access to normal rat chow and drinking water. Blood is collected before and 3, 6 and 24 h after drug administration. Full blood is subjected to hematology using Advia Hematology system (Bayer Diagnostics, Zürich, Switzerland).

25

All data are presented as mean $\pm$ SEM. Statistical analyses are performed by analysis of variance (ANOVA) using Statistica (StatSoft) and the Student-Newman-Keuls procedure for multiple comparisons. The null hypothesis is rejected when $p < 0.05$.

5

As an example, Table 2 shows the effect on lymphocyte counts 6 h after oral administration of 10 mg/kg of some compounds of the present invention to normotensive male Wistar rats as compared to a group of animals treated with vehicle only.


10

Table 2:

Compound of Example	Lymphocyte counts
17	-59. $\pm$ 4%
37	-67 $\pm$ 3%
40	-58 $\pm$ 3%

Claims


1. A compound of the Formula (I),



Formula (I)

wherein

A represents *-CO-CH₂CH₂-, *-CO-CH=CH-,



wherein the asterisks indicate the bond that is linked to the thiophene group of Formula (I);

R¹ represents C₂₋₅-alkyl;

R² represents hydrogen, methyl or ethyl;

R³ represents hydrogen;

R^4 represents methyl, ethyl, or methoxy;

R^5 represents $-\text{CH}_2-(\text{CH}_2)_n-\text{NHSO}_2R^{51}$, $-(\text{CH}_2)_n\text{CH}(\text{OH})-\text{CH}_2-\text{NHSO}_2R^{51}$, $-\text{CH}_2-(\text{CH}_2)_n-\text{NHCOR}^{52}$, $-(\text{CH}_2)_n\text{CH}(\text{OH})-\text{CH}_2-\text{NHCOR}^{52}$, $-\text{CH}_2-(\text{CH}_2)_n-\text{CONR}^{53}R^{54}$, $-\text{CO}-$
 5 NHR^{53} , 1-(3-carboxy-azetidiny)-2-acetyl, 1-(2-carboxy-pyrrolidiny)-2-acetyl, 1-(3-carboxy-pyrrolidiny)-3-propionyl, 1-(2-carboxy-pyrrolidiny)-3-propionyl, 1-(3-carboxy-pyrrolidiny)-3-propionyl, hydroxy, hydroxy- C_{2-5} -alkoxy, di-(hydroxy- C_{1-4} -alkyl)- C_{1-4} -alkoxy, 2,3-dihydroxypropoxy, 2-hydroxy-3-methoxy-propoxy, $-\text{OCH}_2-(\text{CH}_2)_m-\text{NHSO}_2R^{51}$, $-\text{OCH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{NHSO}_2R^{51}$, $-\text{OCH}_2-(\text{CH}_2)_m-\text{NHCOR}^{52}$, or $-\text{OCH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{NHCOR}^{52}$;

R^{51} represents C_{1-3} -alkyl, methylamino, ethylamino, or dimethylamino;

R^{52} represents hydroxymethyl, 2-hydroxyethyl, 2-hydroxy-1-hydroxymethyl-ethyl, or
 15 2,3-dihydroxypropyl;

R^{53} represents hydrogen, C_{1-3} -alkyl, 2-hydroxyethyl, 2-hydroxy-1-hydroxymethyl-ethyl, 2,3-dihydroxypropyl, carboxymethyl, 1-(C_{1-5} -alkylcarboxy)methyl, 2-carboxyethyl, or 2-(C_{1-5} -alkylcarboxy)ethyl;

20

R^{54} represents hydrogen, or methyl;

m represents the integer 1 or 2;


n represents 0, 1, or 2; and

25

R^6 represents methyl, ethyl or halogen;


and salts thereof.

30 2. A compound according to claim 1, wherein **A** represents




wherein the asterisks indicate the bond that is linked to the thiophene group of Formula (I), and salts thereof.

- 5 3. A compound according to claim 1, wherein **A** represents




wherein the asterisks indicate the bond that is linked to the thiophene group of Formula (I), and salts thereof.

- 10 4. A compound according to claim 1, wherein **A** represents




wherein the asterisks indicate the bond that is linked to the thiophene group of Formula (I), and salts thereof.

- 15 5. A compound according to claim 1, wherein **A** represents



wherein the asterisks indicate the bond that is linked to the thiophene group of Formula (I), and salts thereof.

- 5 6. A compound according to claim 1, wherein **A** represents



and salts thereof.

7. A compound according to any one of claims 1 to 6, wherein **R**¹ represents n-propyl or isobutyl, and salts thereof.
- 10 8. A compound according to any one of claims 1 to 7, wherein **R**² represents hydrogen or methyl, and salts thereof.
- 15 9. A compound according to any one of claims 1 to 7, wherein **R**² represents hydrogen, and salts thereof.

10. A compound according to any one of claims 1 to 9, wherein R^5 represents $-CH_2-(CH_2)_n-NHCOR^{52}$, $-(CH_2)_nCH(OH)-CH_2-NHCOR^{52}$, hydroxy, hydroxy- C_{2-5} -alkoxy, di-(hydroxy- C_{1-4} -alkyl)- C_{1-4} -alkoxy, 2,3-dihydroxypropoxy, 2-hydroxy-3-methoxy-propoxy, $-OCH_2-(CH_2)_m-NHCOR^{52}$, or $-OCH_2-CH(OH)-CH_2-NHCOR^{52}$, and salts thereof.

11. A compound according to any one of claims 1 to 9, wherein R^5 represents hydroxy, hydroxy- C_{2-5} -alkoxy, di-(hydroxy- C_{1-4} -alkyl)- C_{1-4} -alkoxy, 2,3-dihydroxypropoxy, 2-hydroxy-3-methoxy-propoxy, $-OCH_2-(CH_2)_m-NHCOR^{52}$, or $-OCH_2-CH(OH)-CH_2-NHCOR^{52}$, and salts thereof.

12. A compound according to any one of claims 1 to 9, wherein R^5 represents 3-hydroxy-2-hydroxymethyl-propoxy, 2,3-dihydroxypropoxy, or $-OCH_2-CH(OH)-CH_2-NHCOR^{52}$, and salts thereof.

13. A compound according to any one of claims 1 to 9, wherein R^5 represents 2,3-dihydroxypropoxy, or $-OCH_2-CH(OH)-CH_2-NHCOR^{52}$, wherein R^{52} represents hydroxymethyl, and salts thereof.

14. A compound according to claim 1, wherein A represents