# SALTS OF N-(4-FLUOROBENZYL)-N-(1-METHYLPIPERIDIN-4-YL)-N'-(4-(2-METHYLPROPYLOXY)PHENYLMETHYL)CARBAMIDE AND THEIR PREPARATION

## BACKGROUND OF THE INVENTION

### Field of the Invention


[0001] The present invention relates to the fields of medicine and chemistry. More particularly, the present invention relates to N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)-phenylmethyl)carbamide, its salts, and syntheses and uses thereof.

### Description of the Related Art

[0002] WO 01/66521 describes N-azacycloalkyl-N-aralkyl carbamides and carboxylic acid amides, which constitute a new class of compounds effective in inhibiting an activity of monoamine receptors, including the serotonin receptor of the 5-HT2A subclass. Examples of disease conditions for which such compounds can be used include, but are not limited to, neuropsychiatric diseases such as schizophrenia and related idiopathic psychoses, depression, anxiety, sleep disorders, appetite disorders, affective disorders such as major depressions, bipolar disorder, depression with psychotic features and Tourette's Syndrome. Other beneficial treatments may be drug-induced psychoses and side-effects of Parkinson's disease as well as psychoses secondary to neurodegenerative disorders such as Alzheimer's or Huntington's Disease, hypertension, migraine, vasospasm, ischemia and the primary treatment and secondary prevention of various thrombotic conditions including myocardial infarction, thrombotic or ischemic stroke, idiopathic and thrombotic thrombocytopenic purpura and peripheral vascular disease.


## SUMMARY OF THE INVENTION

[0003] One embodiment disclosed herein includes a salt of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide of formula I,




(I),


that includes an anion selected from the group consisting of phosphate, sulphate, nitrate, diphosphate, bicarbonate, carbonate, clavulanate, isothionate, borate, halide, nitrate, acetate, succinate, lactate, lactobionate, laurate, mandelate, malate, citrate, fumarate, maleate, oleate, oxalate, ascorbate, nicotinate, benzoate, mesylate, salicylate, stearate, tannate, tosylate, valerate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluensulfonate, 2-ethane disulfonate, and naphthalenesulfonate. In some embodiments, the anion is selected from the group consisting of citrate, fumarate, maleate, malate, phosphate, succinate, sulphate, and edisylate. In one embodiment, when the anion is selected from the group consisting of citrate, maleate, malate, phosphate, succinate, and sulphate, the stoichiometry is 1:1 and when the anion is selected from the group consisting of edisylate and fumarate, the stoichiometry is 2:1. In one embodiment, the salt is a citrate of formula IV,



(IV).

One embodiment of the citrate salt exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 31.8, about 15.9, about 7.9, about 6.3, about 5.96, about 5.23, and about 4.68.


[0004] In another embodiment, the salt is a fumarate of formula V,



(V).

[0005] In one embodiment, the fumarate salt exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 21.7, about 18.3, about 15.7, about 14.5, about 12.6, about 12.3, about 10.9, about 5.52, about 4.72, and about 4.47. In another embodiment, the fumarate saltexhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 18.4, about 15.7, about 12.6, about 9.2, about 5.50, about 4.93, about 4.70, about 4.51, about 4.17, and about 4.06.


[0006] In one embodiment, the salt is a maleate of formula VI,



(VI).

In one embodiment, the maleate salt exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 13.0, about 5.71, about 5.24, about 4.77, about 4.37, and about 4.19 .


[0007] In another embodiment, the salt is a malate of formula VII,



(VII).

In one embodiment, the malate salt exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 13.1, about 12.0, about 5.35, about 5.05, about 4.83, about 4.75, about 4.71, about 4.37, about 4.29, about 4.17, about 4.00, about 3.87, and about 3.83.


[0008] In another embodiment, the salt is a phosphate of formula VIII,



(VIII).

In one embodiment, the phosphate salt exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 17.3, about 5.91, about 4.80, about 4.27, about 4.14, and about 3.86.


[0009] In another embodiment, the salt is a succinate of formula IX,



(IX).

In one embodiment, the succinate salt exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 12.8, about 7.6, about 5.51, about 5.19, about 4.79, about 4.16, and about 4.05.


**[0010]** In another embodiment, the salt is a sulphate of formula X,



(X).

In one embodiment, the sulphate salt exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 17.0, about 9.6, about 5.49, about 4.79, about 4.65, about 4.53, about 4.30, about 4.15, about 4.04, and about 3.89.

**[0011]** In another embodiment, the salt is an edisylate (ethanedisulfonate) of formula XI,



(XI).

In one embodiment, the edisylate salt exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 10.0, about 6.05, about 5.31, about 4.97, about 4.68, about 4.26, and about 4.12.

**[0012]** Another embodiment disclosed herein includes a process for the preparation of a salt disclosed above, including:


a) forming a solution of the compound of formula I in an organic solvent;

b) adding an acid selected from the group consisting of citric acid, fumaric acid, maleic acid, L-(-)-malic acid, phosphoric acid, succinic acid, sulphuric acid, or 1,2-ethane disulfonic acid to said solution; and

c) isolating the salt.

In one embodiment, the isolating includes separating the salt from a suspension formed after step b). In another embodiment, the isolating includes precipitating the salt from a solution formed after step b) by one or more of cooling, solvent removal, or adding a non-solvent.

**[0013]** Another embodiment disclosed herein includes a salt of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide of formula I,



(I),


produced by a process comprising:

a) forming a solution of the compound of formula I in an organic solvent;

b) adding an acid selected from the group consisting of citric acid, fumaric acid, maleic acid, L-(-)-malic acid, phosphoric acid, succinic acid, sulphuric acid, or 1,2-ethane disulfonic acid to said solution; and

c) isolating the salt.

**[0014]** Another embodiment disclosed herein includes a pharmaceutical composition comprising a salt of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide of formula I,




(I),

that includes an anion selected from the group consisting of phosphate, sulphate, nitrate, diphosphate, bicarbonate, carbonate, clavulanate, isothionate, borate, halide, nitrate, acetate, succinate, lactate, lactobionate, laurate, mandelate, malate, citrate, fumarate, maleate, oleate, oxalate, ascorbate, nicotinate, benzoate, mesylate, salicylate, stearate, tannate, tosylate, valerate,

methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluensulfonate, 2-ethane disulfonate, and naphthalenesulfonate and a pharmaceutically acceptable carrier.

[0015] Another embodiment disclosed herein includes a method for the treatment of a neuropsychiatric disease, comprising administering to a subject at least one salt of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide of formula I,



(I),

wherein the salt comprises an anion selected from the group consisting of phosphate, sulphate, nitrate, diphosphate, bicarbonate, carbonate, clavulanate, isothionate, borate, halide, nitrate, acetate, succinate, lactate, lactobionate, laurate, mandelate, malate, citrate, fumarate, maleate, oleate, oxalate, ascorbate, nicotinate, benzoate, mesylate, salicylate, stearate, tannate, tosylate, valerate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluensulfonate, 2-ethane disulfonate, and naphthalenesulfonate. In one embodiment, the neuropsychiatric disease is selected from the group consisting of psychosis, schizophrenia, schizoaffective disorders, mania, psychotic depression, affective disorders, dementia, anxiety, sleep disorders, appetite disorders, bipolar disorder, psychosis secondary to hypertension, migraine, vasospasm, and ischemia, motor tics, tremor, psychomotor slowing, bradykinesia, and neuropathic pain.

[0016] Another embodiment disclosed herein includes a method of inhibiting an activity of a monoamine receptor, comprising administering to a subject at least one salt as described above.

[0017] Another embodiment disclosed herein includes a method for the treatment of neurodegenerative diseases, comprising administering to a subject at least one salt as described above. In some embodiments, the neurodegenerative disease is selected from the group consisting Parkinson's disease, Huntington's disease, Alzheimer's disease, Spinocerebellar Atrophy, Tourette's Syndrome, Friedrich's Ataxia, Machado- Joseph's disease, Lewy Body Dementia, Dystonia, Progressive Supranuclear Palsy, and Frontotemporal Dementia.

[0018] Another embodiment disclosed herein includes a method for treating dyskinesia associated with dopaminergic therapy, comprising administering to a subject at least one salt as described above.

[0019] Another embodiment disclosed herein includes a method for treating dystonia, myoclonus, or tremor associated with dopaminergic therapy, comprising administering to a subject at least one salt as described above.

[0020] Another embodiment disclosed herein includes a method for treating a thrombotic condition, comprising administering to a subject at least one salt as described above. In some embodiments, the thrombotic condition is selected from the group consisting of myocardial infarction, thrombotic or ischemic stroke, idiopathic and thrombotic thrombocytopenic purpura, peripheral vascular disease, and Raynaud's disease.

## BRIEF DESCRIPTION OF THE DRAWINGS

[0021] Figure 1 is a X-ray powder diffraction pattern of the crystalline citrate salt of the compound of formula IV.

[0022] Figure 2 is a X-ray powder diffraction pattern of form A of the crystalline fumarate salt of the compound of formula V.

[0023] Figure 3 is a X-ray powder diffraction pattern of form B of the crystalline fumarate salt of the compound of formula V.

[0024] Figure 4 is a X-ray powder diffraction pattern of the crystalline maleate salt of the compound of formula VI.

[0025] Figure 5 is a X-ray powder diffraction pattern of the crystalline malate salt of the compound of formula VII.

[0026] Figure 6 is a X-ray powder diffraction pattern of the crystalline phosphate salt of the compound of formula VIII.


[0027] Figure 7 is a X-ray powder diffraction pattern of the crystalline succinate salt of the compound of formula IX.

[0028] Figure 8 is a X-ray powder diffraction pattern of the crystalline sulphate salt of the compound of formula X.

[0029] Figure 9 is a X-ray powder diffraction pattern of the crystalline edisylate salt of the compound of formula XI.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT


[0030] One useful N-azacycloalkyl-N-aralkyl carbamide is N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide of formula I:



(I)


Synthesis of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy) phenylmethyl)carbamide

**[0031]** One embodiment is a method of synthesizing the compound of formula (I) comprises reacting the compound of formula II ((4-fluorobenzyl)-(1-methylpiperidin-4-yl)amine)



(II)

with the compound of formula III (4-(2-methylpropyloxy)phenylmethyl-isocyanate)



(III)

**[0032]** In one embodiment, about 0.9 to about 1.1 equivalents of (4-fluorobenzyl)-(1-methylpiperidin-4-yl)amine per equivalent of 4-(2-methylpropyloxy)phenylmethyl-isocyanate is used. In some embodiments, the resulting compound of formula I is isolated from the reaction mixture. In one embodiment, a salt-forming acid is added after the reaction. The formed salt may be isolated by solvent removal, precipitation, or both solvent removal and precipitation, followed by deliberation of the compound of formula I under alkaline aqueous conditions through dissolution in an organic solvent in a two phase system, and separating the compound of formula I from the organic solution. In a preferred embodiment, 1.0 equivalent of (4-fluorobenzyl)-(1-methylpiperidin-4-yl)amine per equivalent of 4-(2-methylpropyloxy)phenylmethyl-isocyanate is used in the reaction. The reaction may be carried out in the presence of Lewis acids as catalysts such as metal salts or more preferably metal alkoxylates. Some examples are $MgCl_2$, $FeCl_2$, $FeCl_3$, $FeBr_2$, $Fe(SO_4)_2$, $NiCl_2$, $BCl_3$, $AlCl_3$, $BBr_3$, $TiCl_4$, $TiBr_4$, $ZrCl_4$, $BCl_3$, $Al(O\text{-}C_1\text{-}C_4\text{-Alkyl})_3$, and $Ti(O\text{-}C_1\text{-}C_4\text{-Alkyl})_3$. The amount of catalyst may be from about 0.0001 to about 5 percent by weight and preferably about 0.01 to about 3 percent by weight relative to the compound of formula II.

[0033] The reaction is preferably carried out in the presence of an inert organic solvent such as aliphatic ethers (e.g., diethyl ether, methyl propyl ether, dibutyl ether, ethylene glycol dimethyl ether, tetrahydrofuran or dioxane), esters of aliphatic carboxylic acids or alcohols (e.g., $C_2$-$C_4$alkyl esters of acetic acid), lactones (e.g., valerolactone), halogenated hydrocarbons (e.g., di- or trichloromethane, tetrachloroethane), or aliphatic $C_3$-$C_8$ketones (e.g., acetone, methyl propyl ketone, diethyl ketone, or methyl i- or t-butyl ketone).


[0034] The reaction temperature is preferably in the range of about -30 °C to about 60 °C and more preferably in the range of about 5 °C to about 30 °C. The reaction time may be controlled by monitoring the consumption of the compound of formula II or formula III either by on-line process analytics, or by recovering and analyzing samples off-line.

[0035] Isolation of the compound of formula I may be performed by any suitable method including removal of the solvent by distillation of the reaction residue under reduced pressure and lower temperatures, such as up to about 100 °C, preferably up to about 80 °C. Isolation may also occur by partial removal of solvent to increase the concentration, filtering of impurities, precipitating the solid compound of formula I either by further concentration or addition of a non-solvent such as an aliphatic hydrocarbon (e.g., pentane, hexane, heptane, octane, cyclohexane, methylcyclohexane, or water), filtering of the solid, and drying. The isolated compound of formula I may be purified by known methods such as distillation or chromatographic methods.


[0036] It was found that removal of impurities such as formed side-products prior to the isolation is a convenient route to produce the compound of formula I with high purity. It was further found that purification can be effectively improved by forming salts of the carbamide, which can be precipitated as crystalline compounds and re-crystallized from solvents to remove impurities. The free carbamide of formula I is then deliberated by dissolution of the salt in water, addition of a base, and extraction of the carbamide with an organic solvent. The organic solutions may be washed with water and aqueous sodium chloride before removal of the solvent by distillation, optionally under reduced pressure. Impurities may be removed in this method by precipitation or dissolution in water in then use of a two phase systems. When precipitation of the salt is desired for easy isolation by filtration or centrifugation, partial removal of the organic solvent and addition of fresh solvent may be carried out. Suitable solvents with low salt solubility are aprotic organic solvents such as hydrocarbons, halogenated hydrocarbons, ethers, ketones, carboxylic acid esters and lactones, acetonitrile, and alcohols having at least 3 carbon atoms.

[0037] The starting materials for the above-described reaction can be obtained by known and analogous methods. Specifically, the compound of formula II may be obtained by the

reaction of N-methylpiperid-4-one with 4-fluorobenzylamine in the presence of a metal hydride, for example according to the scheme




**[0038]** Compounds of formula III may be prepared by reacting 4-hydroxybenzaldehyde with isobutylhalogenide (e.g., isobutylbromogenide) to form 4-isobutoxybenzaldehyde, which may be converted with hydroxylamine to the aldoxime form:





This oxime may be catalytically hydrogenated with a palladium catalyst to the corresponding 4-isobutoxybenzylamine, from which the isocyanate of formula III may be obtained by reaction with phosgene.

Salts of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy) phenylmethyl)carbamide

**[0039]** Some embodiments are salts of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide comprising an anion selected from the group consisting of phosphate, sulphate, nitrate, diphosphate, bicarbonate, carbonate, clavulanate, isothionate, borate, halide, nitrate, acetate, succinate, lactate, lactobionate, laurate, mandelate, malate, citrate, fumarate, maleate, oleate, oxalate, ascorbate, nicotinate, benzoate, mesylate, salicylate, stearate, tannate, tosylate, valerate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluensulfonate, 2-ethane disulfonate, or naphthalenesulfonate.

**[0040]** The salts may be obtained as crystalline solids. When the anion is citrate, maleate, malate, phosphate, succinate, or sulphate, the salt has a 1:1 stoichiometry. The edisylate shows a 2:1 stoichiometry of the free base to the acid and the fumarate also presumably has a 2:1 stoichiometry. In some embodiments, the salts may form hydrates or other solvates. Specifically, it has been found that the malate and succinate can form hydrates. In some embodiments, various polymorphic forms of the salts are provided. In some embodiments, the salts are amorphous.


**[0041]** In one embodiment, the salt is the citrate of formula IV,



(IV).

[0042] One embodiment provides a crystalline form of the citrate of formula IV, which exhibits the X-ray powder diffraction pattern depicted in Figure 1; hereinafter designated as crystalline citrate. Specifically the X-ray powder diffraction pattern exhibits the following characteristic peaks expressed in d-values (Å): 31.8 (vs), 15.9 (m), 7.9 (m), 6.9 (w), 6.3 (m), 5.96 (m), 5.83 (w), 5.23 (m), 4.68 (m), 4.56 (m), 4.17 (m), 4.05 (w), 3.95 (m), 3.91 (m), 3.79 (w), 3.49 (w), and 3.13 (w). The abbreviations in parenthesis are used herein as follows: (vs) = very strong intensity, (s) = strong intensity, (m) = medium intensity, (w) = weak intensity, and (vw) = very weak intensity. In various embodiments, crystalline citrate is present in amounts of at least about 50%, 70%, 80%, 90%, 95%, or 98%, with the remainder being other salt or crystalline forms (including hydrates and solvates) and/or amorphous forms of the compound of formula I.

[0043] Another embodiment is the fumarate salt of formula V,




(V).

[0044] One embodiment provides a crystalline form of the fumarate of formula V, which exhibits the X-ray powder diffraction pattern depicted in Figure 2; hereinafter referred to as crystalline fumarate form A. Specifically the X-ray powder diffraction pattern exhibits the following characteristic peaks expressed in d-values (Å): 21.7 (m), 18.3 (s), 15.7 (s), 14.5 (s), 12.6 (s), 12.3 (m), 10.9 (w), 9.1 (w), 6.8 (w), 6.40 (w), 5.87 (w), 5.52 (m), 5.26 (m), 5.12 (w), 4.72 (s), 4.66 (s), 4.51 (m), 4.47 (s), 4.24 (m), 3.64 (m). In various embodiments, crystalline fumarate form A is present in amounts of at least about 50%, 70%, 80%, 90%, 95%, or 98%, with the remainder being other salt or crystalline forms (including hydrates and solvates) and/or amorphous forms of the compound of formula I.

[0045] Crystalline fumarate form A can also exist in mixtures with the amorphous form and/or with crystalline fumarate form B. Crystalline fumarate form B exhibits the X-ray power diffraction pattern depicted in Figure 3. Specifically the X-ray powder diffraction pattern exhibits the following characteristic peaks expressed in d-values (Å): 18.4 (vs), 15.7 (vs), 12.6 (vs), 10.0 (w), 9.2 (m), 6.8 (m), 6.37 (m), 6.12 (m), 5.68 (m), 5.50 (vs), 5.13 (m), 4.93 (s), 4.70 (s), 4.51 (s), 4.39 (m), 4.30 (m), 4.17 (s), 4.06 (s), 3.88 (m), 3.81 (w), 3.66 (m), 3.64 (m), 3.42 (m). In various embodiments, crystalline fumarate form B is present in amounts of at least about 50%, 70%, 80%, 90%, 95%, or 98%, with the remainder being other salt or crystalline forms (including hydrates and solvates) and/or amorphous forms of the compound of formula I.

[0046] Crystalline fumarate form A may be obtained from rapid crystallization procedures with cooling rates from about 10-100°C per hour, more preferably about 30-60°C per hour, and recovering the solid shortly after the suspension is cooled from about 60°C to 23±2°C, or below. Crystalline fumarate form B may be obtained from slower crystallization procedures with cooling rates from 1-60°C per hour, more preferably form 5-20°C per hour, with subsequent stirring of the obtained suspension at temperatures from 5°C to 40°C for at least one, but up to 60 hours, more preferably for about 24 hours at 23±2°C.


[0047] Another embodiment is the maleate salt of formula VI,



(VI).

[0048] One embodiment provides a crystalline form of the maleate of formula VI, which exhibits the X-ray powder diffraction pattern depicted in Figure 4; hereinafter referred to as crystalline maleate. Specifically the X-ray powder diffraction pattern exhibits the following characteristic peaks expressed in d-values (Å): 17.1 (w), 13.0 (vs), 10.0 (w), 8.6 (w), 7.9 (w), 5.71 (vs), 5.24 (m), 4.98 (m), 4.86 (w), 4.77 (m), 4.70 (w), 4.37 (m), 4.29 (w), 4.19 (vs), 3.92 (w), 3.76 (w), 3.67 (w), 3.62 (m), 3.52 (w), 3.38 (m), 3.27 (m), 3.05 (m). In various embodiments, crystalline maleate is present in amounts of at least about 50%, 70%, 80%, 90%, 95%, or 98%, with the remainder being other salt or crystalline forms (including hydrates and solvates) and/or amorphous forms of the compound of formula I.


[0049] Another embodiment is the malate salt of formula VII,



(VII).

[0050] One embodiment provides a crystalline form of the malate of formula VII, which exhibits the X-ray powder diffraction pattern depicted in Figure 5; hereinafter referred to as crystalline malate. Specifically the X-ray powder diffraction pattern exhibits the following characteristic peaks expressed in d-values (Å): 19.8 (m), 16.2 (w), 13.1 (vs), 12.0 (s), 7.7 (m), 7.2 (m), 6.1 (m), 5.35 (s), 5.05 (s), 4.89 (m), 4.83 (s), 4.75 (vs), 4.71 (vs), 4.63 (m), 4.55 (m), 4.37 (vs), 4.29 (vs), 4.17 (s), 4.00 (s), 3.97 (m), 3.87 (s), 3.83 (s), 3.61 (m). Without being bound by any particular theory, this crystalline form of the malate of formula VII may be a sesquihydrate. In various embodiments, crystalline malate is present in amounts of at least about 50%, 70%, 80%, 90%, 95%, or 98%, with the remainder being other salt or crystalline forms (including hydrates and solvates) and/or amorphous forms of the compound of formula I.


[0051] Another embodiment is the phosphate salt of formula VIII,



(VIII).

[0052] One embodiment provides a crystalline form of the phosphate of formula V, which exhibits the X-ray powder diffraction pattern depicted in Figure 6; hereinafter referred to as crystalline phosphate. Specifically the X-ray powder diffraction pattern exhibits the following characteristic peaks expressed in d-values (Å): 17.3 (vs), 10.1 (m), 8.9 (m), 6.7 (w), 6.5 (m), 5.91 (s), 5.74 (m), 5.16 (w), 4.93 (m), 4.80 (m), 4.75 (w), 4.56 (m), 4.27 (m), 4.14 (m), 3.86 (m), 3.55 (m). In various embodiments, crystalline phosphate is present in amounts of at least about 50%, 70%, 80%, 90%, 95%, or 98%, with the remainder being other salt or crystalline forms (including hydrates and solvates) and/or amorphous forms of the compound of formula I.


[0053] Another embodiment is the succinate salt of formula IX,



(IX).

[0054] One embodiment provides a crystalline form of the succinate of formula IX, which exhibits the X-ray powder diffraction pattern depicted in Figure 7; hereinafter referred to as crystalline succinate. Specifically the X-ray powder diffraction pattern exhibits the following characteristic peaks expressed in d-values (Å): 12.8 (vs), 8.6 (w), 7.6 (m), 6.4 (w), 5.51 (s), 5.27 (w), 5.19 (m), 4.79 (m), 4.42 (w), 4.32 (m), 4.16 (s), 4.05 (s), 3.91 (m), 3.69 (w), 3.31 (w), 3.27 (w), 3.14 (w), 2.97 (w), 2.76 (w). In various embodiments, crystalline succinate is present in amounts of at least about 50%, 70%, 80%, 90%, 95%, or 98%, with the remainder being other salt or crystalline forms (including hydrates and solvates) and/or amorphous forms of the compound of formula I.


[0055] Another embodiment is the sulphate salt of formula X,



(X).

[0056] One embodiment provides a crystalline form of the sulphate of formula X, which exhibits the X-ray powder diffraction pattern depicted in Figure 8; hereinafter referred to as crystalline sulphate. Specifically the X-ray powder diffraction pattern exhibits the following characteristic peaks expressed in d-values (Å): 17.0 (vs), 9.6 (m), 8.3 (w), 6.8 (m), 6.4 (m), 5.49 (vs), 5.29 (w), 4.79 (s), 4.65 (m), 4.53 (s), 4.42 (m), 4.30 (vs), 4.18 (m), 4.15 (s), 4.04 (m), 3.89 (w), 3.60 (m), 3.56 (w). In various embodiments, crystalline sulphate is present in amounts of at least about 50%, 70%, 80%, 90%, 95%, or 98%, with the remainder being other salt or crystalline forms (including hydrates and solvates) and/or amorphous forms of the compound of formula I.

[0057] Another embodiment is the edisylate (ethanedisulfonate) salt of formula XI,



(XI).

[0058] One embodiment provides a crystalline form of the edisylate of formula XI, which exhibits the X-ray powder diffraction pattern depicted in Figure 9; hereinafter referred to as crystalline edisylate. Specifically the X-ray powder diffraction pattern exhibits the following characteristic peaks expressed in d-values (Å): 12.1 (m), 10.0 (s), 9.3 (m), 8.1 (m), 6.6 (m), 6.05 (vs), 5.31 (s), 5.18 (m), 4.97 (vs), 4.81 (w), 4.68 (s), 4.57 (m), 4.46 (m), 4.35 (m), 4.26 (s), 4.12 (s), 3.96 (m), 3.88 (w), 3.75 (m), 3.62 (m), 3.53 (w), 3.48 (m), 3.42 (w), 3.31 (m), 3.15 (w), 3.07 (w). In various embodiments, crystalline edisylate is present in amounts of at least about 50%, 70%, 80%, 90%, 95%, or 98%, with the remainder being other salt or crystalline forms (including hydrates and solvates) and/or amorphous forms of the compound of formula I.

[0059] Salts of the compound of formula I described herein may be prepared by the reaction of equivalent amounts of the base of formula I with an acid in a suitable inert organic solvent. Accordingly, one embodiment is a process for the preparation of a salt of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2methylpropyloxy)phenylmethyl)carbamide of formula I having anions selected from the group consisting of citrate, fumarate, maleate, malate, phosphate, succinate, sulphate and edisylate, that includes:

a) forming a solution of the compound of formula I in an organic solvent;

b) adding a suitable organic or inorganic acid to the solution; and

c) isolating the salt of the compound of formula I from an obtained suspension, or precipitating the salt by cooling, solvent removal, adding a non-solvent, or a combination of these methods.

[0060] Suitable solvents include, but are not limited to, hydrocarbons such as toluene, halogenated hydrocarbons such as di- or trichloromethane, tetrachloroethane, esters of aliphatic carboxylic acids and alcohols ($C_2$-$C_4$alkyl esters of acetic acid) (ethyl acetate), lactones (valerolactone), acetonitrile, ethers (diethylether, methyl propyl ether, t–butyl-methyl-ether, dibutyl ether, ethylene glycol dimethyl ether, tetrahydrofuran or dioxane), aliphatic $C_3$-$C_8$ketones (acetone, methyl propyl ketone, diethyl ketone or methyl i- or t-butyl ketone), and alcohols (methanol, ethanol, n- or i-propanol and butanol).

[0061] Suitable salt-forming acids include but are not limited to phosphoric acid, sulphuric acid, nitric, diphosphate, bicarbonate, carbonic acid, clavulanic acid, isothionic acid, boric acid, hydrohalic acid (e.g., hydrochloric acid or hydrobromic acid), nitric acid, and aliphatic or aromatic carboxylic or sulfonic acids (e.g., acetic, succinic, lactic, lactobionic, lauric, mandelic, malic, tartaric, citric, fumaric, maleic, oleic, oxalic, ascorbic, nicotinic, benzoic, mesylic, salicylic, stearic, tannic, tosylic, valeric, methanesulfonic, ethanesulfonic, benzenesulfonic, p-toluensulfonic, salicylic, 2-ethane disulfonic, or naphthalenesulfonic acid).

[0062] When used, the non-solvent may be an aliphatic hydrocarbon such as petrol ether, pentane, hexane, heptane, octane, cyclopentane, cyclohexane or methylcyclohexane. Other non-solvents may be determined with solubility tests of a selected salt in various solvents.

[0063] Various crystallization techniques may be used to form and isolate crystalline compounds, such as stirring of a suspension (phase equilibration), precipitation, re-crystallization, solvent evaporation, cooling to initiate crystallization, and cooling down to -100 °C (e.g., down to -30 °C). Diluted or saturated solutions may be used for crystallization, with or without seeding with suitable nucleating agents. Obtained crystalline solids may be purified by crystallization techniques well known in the art. Temperatures up to 100 °C may be applied to form solutions.

[0064] The salts described herein may be obtained in good yields. Re-crystallization produces purified forms suitable for use in pharmaceutical compositions. The salts may occur in more than one crystalline form. For example, some of the salts may form hydrates or solvates.

[0065] The salts described herein are especially suitable as active compounds or pro-drugs in pharmaceutical formulations to inhibit an activity of a monoamine receptor, preferably a serotonin receptor of the 5-HT2A subclass. The salts of formula IV are very soluble in aqueous systems and the free base is liberated at physiological pH ranges, providing a high bioavailability. The salts of formula IV and formula XI, in the crystalline forms disclosed herein, possess good storage stability. The crystalline compounds facilitate processing and handling for the manufacture of the salts and their formulation.

[0066] Accordingly, one embodiment is a pharmaceutical composition comprising at least one salt described herein and a pharmaceutically acceptable carrier or diluent. The amount of the salts used depends on type of formulation and desired dosages during administration time periods. The amount in an oral formulation may be from 0.1 to 500 mg, preferably from 0.5 to 300 mg, and more preferably from 1 to 100 mg.

[0067] Oral formulations may be solid formulations such as capsules, tablets, pills and troches, or liquid formulations such as aqueous suspensions, elixirs and syrups. Solid and liquid formulations encompass also incorporation of the salts into liquid or solid food. Liquids also encompass solutions of the salts for parenteral applications such as infusion or injection.

[0068] The crystalline solid salts described herein may directly be used as powder (micronized particles), granules, suspensions or solutions, or they may be combined together with other pharmaceutically acceptable ingredients in admixing the components and optionally finely divide them, and then filling capsules, composed for example from hard or soft gelatine, compressing tablets, pills or troches, or suspend or dissolve them in carriers for suspensions, elixirs, and syrups. Coatings may be applied after compression to form pills.

[0069] Pharmaceutically acceptable ingredients are well known for the various types of formulations and may be for example binders such as natural or synthetic polymers, excipients, lubricants, surfactants, sweetening and flavouring agents, coating materials, preservatives, dyes, thickeners, adjuvants, antimicrobial agents, antioxidants and carriers for the various formulation types.

[0070] Examples for binders are gum tragacanth, acacia, starch, gelatine, and biological degradable polymers such as homo- or co-polyesters of dicarboxylic acids, alkylene glycols, polyalkylene glycols and/or aliphatic hydroxyl carboxylic acids; homo- or co-polyamides of dicarboxylic acids, alkylene diamines, and/or aliphatic amino carboxylic acids; corresponding polyester-polyamide-co-polymers, polyanhydrides, polyorthoesters, polyphosphazene and polycarbonates. The biological degradable polymers may be linear, branched or crosslinked. Specific examples are poly-glycolic acid, poly-lactic acid, and poly-d,l-lactide/glycolide. Other examples for polymers are water-soluble polymers such as polyoxaalkylenes (e.g., polyoxaethylene, polyoxapropylene and mixed polymers thereof), poly-acrylamides and hydroxylalkylated polyacrylamides, poly-maleic acid and esters or -amides thereof, poly-acrylic acid and esters or -amides thereof, poly-vinylalcohol und esters or -ethers thereof, poly-vinylimidazole, poly-vinylpyrrolidon, and natural polymers like chitosan.

[0071] Examples for excipients are phosphates such as dicalcium phosphate.

[0072] Examples for lubricants are natural or synthetic oils, fats, waxes, or fatty acid salts like magnesium stearate.

[0073] Surfactants may be anionic, cationic, amphoteric, or neutral. Examples for surfactants are lecithin, phospholipids, octyl sulfate, decyl sulfate, dodecyl sulfate, tetradecyl sulfate, hexadecyl sulfate and octadecyl sulfate, Na oleate or Na caprate, 1-acylaminoethane-2-sulfonic acids, such as 1-octanoylaminoethane-2-sulfonic acid, 1-decanoylaminoethane-2-sulfonic acid, 1-dodecanoylaminoethane-2-sulfonic acid, 1-tetradecanoylaminoethane-2-sulfonic acid, 1-hexadecanoylaminoethane-2-sulfonic acid, and 1-octadecanoylaminoethane-2-sulfonic acid, and taurocholic acid and taurodeoxycholic acid, bile acids and their salts, such as cholic acid, deoxycholic acid and sodium glycocholates, sodium caprate or sodium laurate, sodium oleate, sodium lauryl sulphate, sodium cetyl sulphate, sulfated castor oil and sodium dioctyl-sulfosuccinate, cocamidopropylbetaine and laurylbetaine, fatty alcohols, cholesterols, glycerol

mono- or -distearate, glycerol mono- or -dioleate and glycerol mono- or -dipalmitate, and polyoxyethylene stearate.

[0074] Examples for sweetening agents are sucrose, fructose, lactose or aspartam.

[0075] Examples for flavouring agents are peppermint, oil of wintergreen or fruit flavours like cherry or orange flavour.

[0076] Examples for coating materials are gelatine, wax, shellac, sugar or biological degradable polymers.

[0077] Examples for preservatives are methyl or propylparabens, sorbic acid, chlorobutanol, phenol and thimerosal.

[0078] Examples for adjuvants are fragrances.

[0079] Examples for thickeners are synthetic polymers, fatty acids and fatty acid salts and esters and fatty alcohols.

[0080] Examples for antioxidants are vitamins, such as vitamin A, vitamin C, vitamin D or vitamin E, vegetable extracts or fish oils.

[0081] Examples for liquid carriers are water, alcohols such as ethanol, glycerol, propylene glycol, liquid polyethylene glycols, triacetin and oils. Examples for solid carriers are talc, clay, microcrystalline cellulose, silica, alumina and the like.

[0082] The pharmaceutical formulations may also contain isotonic agents, such as sugars, buffers or sodium chloride.

[0083] The salts described herein may also be formulated as effervescent tablet or powder, which disintegrate in an aqueous environment to provide a drinking solution.

[0084] A syrup or elixir may contain the salts described herein, sucrose or fructose as a sweetening agent, a preservative like methylparaben, a dye, and a flavouring agent.

[0085] Slow release formulations may also be prepared from the salts described herein in order to achieve a controlled release of the active agent in contact with the body fluids in the gastro intestinal tract, and to provide a substantial constant and effective level of the active agent in the blood plasma. Any of the compounds of formula IV to XI may be embedded for this purpose in a polymer matrix of a biological degradable polymer, a water-soluble polymer or a mixture of both, and optionally suitable surfactants. Embedding can mean in this context the incorporation of micro-particles in a matrix of polymers. Controlled release formulations are also obtained through encapsulation of dispersed micro-particles or emulsified micro-droplets via known dispersion or emulsion coating technologies.

[0086] The salts described herein may also be useful for administering a combination of therapeutic agents to an animal. Such a combination therapy can be carried out in using at least one further therapeutic agent which can be additionally dispersed or dissolved in a formulation.

[0087] The salts described herein and formulations containing the salts can also be administered in combination with other therapeutic agents that are effective to treat a given condition to provide a combination therapy.

[0088] In some embodiments, the crystalline salts and the pharmaceutical composition disclosed herein are used to treat neuropsychiatric diseases including psychosis, schizophrenia, schizoaffective disorders, mania, psychotic depression, affective disorders, dementia, anxiety, sleep disorders, appetite disorders, bipolar disorder, psychosis secondary to hypertension, migraine, vasospasm, and ischemia, motor tics, tremor, psychomotor slowing, bradykinesia, and neuropathic pain. In one embodiment, the salts and compositions are used to inhibit an activity of a monoamine receptor, preferably a serotonin receptor of the 5-HT2A subclass.

[0089] Another embodiment is a method for the treatment of neurodegenerative diseases, including Parkinson's disease, Huntington's disease, Alzheimer's disease, Spinocerebellar Atrophy, Tourette's Syndrome, Friedrich's Ataxia, Machado- Joseph's disease, Lewy Body Dementia, Dystonia, Progressive Supranuclear Palsy, and Frontotemporal Dementia by administering a salt described herein.

[0090] Another embodiment is a method for treating dyskinesia associated with dopaminergic therapy, by administering a salt described herein.

[0091] Another embodiment is a method for treating dystonia, myoclonus, or tremor associated with dopaminergic therapy, by administering a salt described herein.

[0092] Another embodiment is a method for treating a thrombotic condition including myocardial infarction, thrombotic or ischemic stroke, idiopathic and thrombotic thrombocytopenic purpura, peripheral vascular disease, and Raynaud's disease, by administering a salt described herein.

[0093] Another embodiment is a method of treating addiction, including alcohol addiction, opioid addiction, and nicotine addiction, by administering a salt described herein.

[0094] Another embodiment is a method of treating a decrease in libido or ejaculatory problems by administering a salt described herein.

[0095] One embodiment includes a method of delivering a compound of formula I to a subject, comprising administering to the subject an effective amount of the salt chosen from compounds of formula IV, V, VI, VII, VIII, IX, X, or XI.

## EXAMPLES

Experimental Procedures

[0096] Powder X-ray Diffraction (PXRD): PXRD was performed on a Philips 1710 powder X-ray diffractometer using $CuK_\alpha$ radiation. d-spacings were calculated from the 2θ

values using the wavelength of 1.54060 Å. Generally, 2θ values were within an error of ±0.1-0.2°. The experimental error on the d-spacing values was therefore dependent on the peak location.

**[0097]** Differential Scanning Calorimetry (DSC): Perkin Elmer DSC 7 in gold sample pan sealed under nitrogen. Heating rate 10 K/min.

**[0098]** FT-Raman Spectroscopy: Bruker RFS100. Nd:YAG 1064 nm excitation, 100 mW laser power, Ge-detector, 64 scans, range 25-3500 $cm^{-1}$, 2 $cm^{-1}$ resolution.


**[0099]** TG-FTIR: Thermogravimetric measurements were carried out with a Netzsch Thermo-Microbalance TG 209 coupled to a Bruker FTIR Spectrometer Vector 22 (sample pans with pinhole, nitrogen atmosphere, heating rate 10K/min).

**[0100]** HPLC: HPLC measurement were carried out with a HP LC1090M, Column Symmetry C18, 3.0·150 mm.

**[0101]** Solubility: The approximate solubility in water was determined by adding double distilled water in steps of 5 µl to 5 mg substance and sonification of the suspension for 2 minutes. The limit value of completely dissolved amount was determined. Determination of solubilities below 20 mg/l occurred with stirring suspensions in water, filtering off the excess, and measuring of the amount of substance in the filtrate.


Example 1: Preparation of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide

a) Preparation of





**[0102]** Triacetoxy borohydride (6.5 kg) was added over 1.5 h to a solution of N-methylpiperid-4-one (3.17 kg) and 4-fluorobenzylamine (3.50 kg) in methanol (30 l), maintaining the temperature under 27 °C. The reaction mixture was stirred for 15 h at 22 °C. The residual amine was checked by gel chromatography (4-fluorobenzylamine: < 5%). A solution of 30% sodium hydroxide (12.1 kg) in water (13.6 kg) was added in 75 minutes (min) maintaining the temperature under 20 °C. Methanol was distilled off to a residual volume of 26 litters. Ethyl acetate was added (26 L), the solution was stirred for 15 min, the phases were decanted over 15 min and the lower aqueous phase was discarded. Ethyl acetate was distilled under reduced pressure from the organic phase at 73-127 °C. At this stage the residue was mixed with a second crude batch prepared according to this method. The combined products were then distilled at 139-140 °C / 20 mbar to yield 11.2 kg product (> 82%).

b) Preparation of




[0103] 4-Hydroxybenzaldehyde (4.0 kg) and ethanol (20 l) were added to a solution of isobutyl bromide (9.0 kg) in ethanol (15 l). Potassium carbonate (13.6 kg) was added and the suspension was refluxed (74-78 °C) for 5 days. The residual 4-hydroxybenzaldehyde was checked by HPLC (< 10%). The suspension was cooled to 20 °C and used in the next step.

c) Preparation of



[0104] Hydroxylamine (50% in water, 8.7 kg) was added to the product from previous step b)(174 l, 176 kg) and ethanol (54 l). The suspension was refluxed (77 °C) for 3 h. Unreacted residual amounts of the compound of step b was checked by HPLC (< 5%). The suspension was cooled to 30 °C, filtered and the filter was washed with ethanol (54 l). The solution was concentrated by distillation under reduced pressure at 30 °C to a residual volume of 67 litters. The solution was cooled to 25 °C and water (110 l) was added. The suspension was concentrated by distillation under reduced pressure at 30 °C to a residual volume of 102 litters. Petrol ether (60-90 fraction, 96 l) was added and the mixture was heated to reflux (70 °C). The solution was cooled to 40 °C and crystallization was initiated by seeding. The suspension was cooled to 5 °C and stirred for 4h. The product was centrifuged and the cake was washed with petrol ether (60-90 fraction, 32 l). The wet cake was dried at about 40 °C to yield 16kg product (63%).


d) Preparation of



[0105] The product from previous step c) (15.7 kg) was dissolved in ethanol (123 l). Acetic acid (8.2 kg) and palladium on charcoal 5% wet (1.1 kg) were added. The oxime was hydrogenated at 22 °C and 1.5 bar for 4h. Consumption of oxime was checked by HPLC (for information). The catalyst was filtered and the solvent was distilled under reduced pressure at 36

°C to a final volume of 31 l. Ethyl acetate (63 l) was added and the mixture was heated to reflux (75 °C) until dissolution. The solution was cooled to 45 °C and the crystallization was initiated by seeding. The suspension was cooled to 6-10 °C and stirred for 2.5h. The product was centrifuged and the cake was washed with 2 portions of ethyl acetate (2 x 0.8 l). The wet cake was dried at a temperature of about 40 °C to yield 8 kg (41%).

e) Preparation of



[0106] Aqueous sodium hydroxide (30%, 5.0 kg) was added to a suspension of the product from previous step d) (7.9 kg) in heptane (41 l). The solution was heated to 47 °C, stirred for 15 min and decanted over 15 min. The pH was checked (pH>12) and the aqueous phase was separated. The solvent was removed by distillation under reduced pressure at 47-65 °C. Heptane was added (15 l) and it was removed by distillation under reduced pressure at 58-65 °C. Heptane was added (7 l), the solution was filtered and the filter was washed with heptane (7 l). The solvent was removed by distillation under reduced pressure at 28-60 °C. Tetrahydrofuran (THF, 107 l) and triethylamine (TEA, 6.8 kg) were added and the temperature was fixed at 22 °C. In another reactor, phosgene (5.0 kg) was introduced in tetrahydrofuran (88 l) previously cooled to -3 °C. The THF and TEA solution was added to the solution of phosgene in 3h 50 min maintaining the temperature at -3 °C. The reactor was washed with tetrahydrofuran (22 l). The mixture was stirred for 45 min at 20 °C and then for 90 min at reflux (65 °C). The solvent was distilled under reduced pressure at 25-30 °C to a residual volume of 149 l. The absence of phosgene was controlled. At this stage, there still was phosgene and the suspension was degassed by bubbling nitrogen through it. After this operation the level of phosgene above the solution was below 0.075 ppm. The suspension was filtered and washed with tetrahydrofuran (30 l). The solvent was distilled under reduced pressure at 20-25 °C to a residual volume of 40 l. Tetrahydrofuran (51 l) was added and the solvent was distilled under reduced pressure at 20-25 °C to a residual volume of 40 l. The final volume was adjusted to about 52 litters by addition of tetrahydrofuran (11 l). The solution was analysed and used in the next step.

solvent was removed under reduced pressure at 20-25 °C. Heptane (49 l) was added and the suspension was stirred for 15 min at 40 °C. Then, 8 l of solvent was removed by distillation under reduced pressure at 38-41 °C. The slurry was cooled to 20 °C and stirred for 1 h. The product was centrifuged and the cake was washed with heptane (5 l). The wet compound of Formula I (5.5 kg) was dissolved in ethanol (28 l) at 45 °C. A solution of tartaric acid (0.72 kg) in ethanol (11 l) was added at 45 °C and the line was washed with ethanol (9 l). The solution was cooled to 43 °C, seeded with the tartrate salt of the compound of Formula I, then the slurry was cooled to 35 °C in 30 min, stirred at this temperature for 1 h and cooled to -5 °C. After 14 h at this temperature the product was centrifuged and washed with two portions of ethanol (2x6 l). The wet cake was dried at about 45 °C for 76 h to yield 4 kg of the hemi-tartrate.

i) Re-crystallization

[0110] 150.0 g of hemi-tartrate obtained in h) was dissolved under stirring at 65 °C in 112 ml absolute ethanol and then cooled under stirring to 48 °C at a cooling rate of 1 °C/min. Crystallization started after a few minutes at this temperature and the suspension turned to a thick paste within 1 h. The suspension was heated again to 60 °C and then cooled to 48 °C at a rate of 1 °C/min. The obtained suspension was stirred and was cooled to 15 °C at a cooling rate of 3 °C/h. The crystalline precipitate was separated by filtration and the bottle was washed with 10 ml absolute ethanol cooled to 5 °C. The crystalline residue was dried under vacuum and 40 °C for 50 hours to yield 146 g crystalline pure hemi-tartrate.

j) Second purification

[0111] 15.78 g of the tartrate salt prepared from step i) was dissolved in 130 ml water. 500 ml TBME was added and the pH was adjusted to 9.8 by addition of 2 N NaOH solution. After precipitation of a white solid, the aqueous phase was extracted 5 times by 500 ml TBME. The organic phases were concentrated until a volume of about 400 ml remained. The solution was stored at 6°C. The precipitate was filtered, washed with TBME and finally dried in vacuum for 5 hours. Yield: 8.24 g of a white powder. The mother liquor was concentrated to a fourth and stored at 6°C. The precipitate was filtered and dried in vacuum for 18 hours. Yield: 1.6 g of a white powder.

[0112] PXRD revealed a crystalline compound of formula I. No Raman peaks from tartaric acid were found. The first scan of DSC (-50°C to 210°C, 10°K/min) revealed a melting point at 123.6°C. Above about 190°C, the sample started to decompose.

Example 2: Preparation of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide citrate of formula IV

a)

[0113] 90 mg of the product from Example 1 and 40 mg citric acid were suspended in 5.0 ml ethylacetate. The suspension was stirred at 60 °C for 15 minutes (min), cooled to 23±2 °C, and then stored for 30 min at 23±2 °C. The precipitate was filtered off and dried in air for 30 min to yield 52 mg of a crystalline white powder. Optical microscopy shows that the obtained solid was crystalline.

b)

[0114] 182 mg of the product from Example 2 and 78.4 mg citric acid were suspended in 10.0 ml ethyl acetate. The suspension was stirred at 60 °C for 30 min, then stirred at 40 °C for 90 min, and finally stirred for 60 min at 23 °C. The suspension was filtered and washed with heptane, yielding 237 mg of a white crystalline powder with an endothermic peak near 153 °C (enthalpy of fusion of about 87 J/g), determined by differential scanning calorimetry at a rate of 10K/min (DSC). Thermogravimetry (TG-FTIR) showed a mass loss of about 0.7% between 60 and 160 °C, which was attributed to absorbed water. Decomposition started at about 170 °C. Solubility in water was about 14 mg/ml. The crystalline powder remained substantially unchanged when stored for 1 week at 60 °C and about 75% r.h. in an open container (HPLC area was 99.4% compared to reference value of 99.9%). Elemental analysis and $^1$H-NMR complies with an 1:1 stoichiometry.

[0115] The powder X-ray diffraction pattern (PXRD) of the obtained citrate salt is shown in Figure 1 and the characteristic peaks in 2 theta with the corresponding d-spacing values in Å are given in Table 1.

Table 1: d-Spacings for the compound of formula IV

| Angle [°2θ] | d-spacings [Å] | Intensity (qualitative) |
|---|---|---|
| 2.8 | 31.8 | vs |
| 5.6 | 15.9 | m |
| 11.2 | 7.9 | m |
| 12.6 | 7.0 | vw |

| | | |
|---|---|---|
| 12.9 | 6.9 | w |
| 14.0 | 6.3 | m |
| 14.9 | 5.96 | m |
| 15.2 | 5.83 | w |
| 16.9 | 5.23 | m |
| 17.9 | 4.94 | vw |
| 18.1 | 4.89 | vw |
| 18.9 | 4.68 | m |
| 19.5 | 4.56 | m |
| 21.3 | 4.17 | m |
| 21.9 | 4.05 | w |
| 22.5 | 3.95 | m |
| 22.7 | 3.91 | m |
| 23.4 | 3.79 | w |
| 24.1 | 3.70 | vw |
| 24.5 | 3.62 | vw |
| 25.5 | 3.49 | w |
| 28.5 | 3.13 | w |
| 29.9 | 2.99 | vw |
| 31.0 | 2.89 | w |

Example 3: Preparation of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide fumarate of formula V

a)

**[0116]** 90 mg of the product from Example 1 and 24.3 mg fumaric acid were suspended in 5.0 ml ethyl acetate. The suspension was stirred at 60 °C for 15 min, then stored for 75 min at 23±2 °C. Optical microscopy revealed a crystalline substance. The suspension was filtered and washed with t-butyl methyl ether (TBME). Yield: 83 mg of a white powder. PXRD and Raman spectroscopy indicate a crystalline form A, containing amorphous parts.

**[0117]** The powder X-ray diffraction pattern (PXRD) is shown in Figure 2 and the characteristic peaks in 2 theta with the corresponding d-spacing values in Å are given in Table 2.

Table 2: d-Spacings for the compound of formula V form A

| Angle [°2θ] | d-spacings [Å] | Intensity (qualitative) |
|---|---|---|
| 4.1 | 21.7 | m |
| 4.8 | 18.3 | s |
| 5.6 | 15.7 | s |
| 6.1 | 14.5 | s |

| | | |
|---|---|---|
| 7.0 | 12.6 | s |
| 7.2 | 12.3 | m |
| 8.1 | 10.9 | w |
| 9.7 | 9.1 | w |
| 13.1 | 6.8 | w |
| 13.8 | 6.40 | w |
| 15.1 | 5.87 | w |
| 16.0 | 5.52 | m |
| 16.8 | 5.26 | m |
| 17.3 | 5.12 | w |
| 18.8 | 4.72 | s |
| 19.0 | 4.66 | s |
| 19.7 | 4.51 | m |
| 19.8 | 4.47 | s |
| 20.9 | 4.24 | m |
| 24.5 | 3.64 | m |

b)

**[0118]** 180 mg of the product from Example 2 and 48.2 mg fumaric acid were suspended in 10.0 ml ethyl acetate. The suspension was stirred at 60 °C for 30 min, then for 90 min at 40 °C and finally for 70 min at 23 °C. The precipitate was filtered off and washed with heptane, yielding 167 mg of a crystalline white powder. TG-FTIR showed a mass loss of about 8.6% between 60 and 170 °C, which was attributed to absorbed water, ethyl acetate and $CO_2$. Decomposition started at about 160 °C. $^1$H-NMR complies with an 1:0.75 stoichiometry (base : fumaric acid). PXRD and Raman spectroscopy indicate a crystalline form B.

c)

**[0119]** 48.6 mg of the product from Example 2 was suspended in 10.0 ml ethyl acetate. 180 mg fumaric acid was dissolved in 1 ml ethanol and added to the suspension. The resulting mixture was stirred at 50 °C for 1 hour and then at 23 °C for 21 hours. Thereafter, 12 ml ethylacetate was added and the solution was further stirred for 24 hours at 23 °C. The solvent volume was reduced to half by a nitrogen flow and 9 ml heptane was then added. The formed suspension was further stirred for 24 hours at 23±2 °C. The precipitate was filtered off to yield 191 mg of a crystalline white powder. PXRD and Raman spectroscopy indicate a crystalline form B. The solubility in water was > 500 mg/ml. TG-FTIR shows a mass loss of about 0.9% between 70 and 140 °C, which was attributed to ethyl acetate. Storage at 75% r.h. in an open container reveals changes of the substance after 3 days, detected with Raman spectroscopy. $^1$H-NMR complies with an 1:0.75 stoichiometry (base : fumaric acid). The crystalline powder

remains substantially unchanged when stored for 1 week at 60 °C and about 75% r.h. in an open container (HPLC area was 96.7% compared to reference value of 99.4%). It is possible that the crystalline powder is a mixture of a fumarate and a hemi-fumarate.

**[0120]** The powder X-ray diffraction pattern (PXRD) is shown in Figure 3 and the characteristic peaks in 2 theta with the corresponding d-spacing values in Å are given in Table 3.

Table 3: d-Spacings for the compound of formula V form B

| Angle [°2θ] | d-spacings [Å] | Intensity (qualitative) |
|---|---|---|
| 4.8 | 18.4 | vs |
| 5.6 | 15.7 | vs |
| 7.0 | 12.6 | vs |
| 8.8 | 10.0 | w |
| 9.6 | 9.2 | m |
| 10.5 | 8.4 | m |
| 10.9 | 8.1 | w |
| 11.3 | 7.8 | vw |
| 11.8 | 7.5 | w |
| 13.1 | 6.8 | m |
| 13.9 | 6.37 | m |
| 14.5 | 6.12 | m |
| 15.6 | 5.68 | m |
| 16.1 | 5.50 | vs |
| 17.3 | 5.13 | m |
| 18.0 | 4.93 | s |
| 18.9 | 4.70 | s |
| 19.7 | 4.51 | s |
| 20.2 | 4.39 | m |
| 20.6 | 4.30 | m |
| 21.3 | 4.17 | s |
| 21.9 | 4.06 | s |
| 22.9 | 3.88 | m |
| 23.3 | 3.81 | w |
| 24.3 | 3.66 | m |
| 24.4 | 3.64 | m |
| 26.1 | 3.42 | m |
| 28.7 | 3.11 | w |

Example 4: Preparation of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide maleate of formula VI

a)

**[0121]** 181 mg of the product from Example 1 and 48.2 mg maleic acid were dissolved in 10.0 ml ethyl acetate. The solution was stirred at 60 °C for 15 min, then for 20 min at 23±2 °C. Precipitation of a white solid started after this time. The suspension was stored at 5 °C for 48 hours and then the solvent volume was reduced to a fourth by a nitrogen flow. Storage at 5 °C was continued for 72 hours. The white solid was filtered off yielding 113 mg of a crystalline powder. PXRD and Raman spectroscopy indicate a crystalline form. TG-FTIR shows a mass loss of about 7.2% between 60 and 160 °C, which was attributed to absorbed water and ethyl acetate. Decomposition starts at about 160 °C.

**[0122]** The powder X-ray diffraction pattern (PXRD) is shown in Figure 4 and the characteristic peaks in 2 theta with the corresponding d-spacing values in Å are given in Table 4.

Table 4: d-Spacings for the compound of formula VI

| Angle [°2θ] | d-spacings [Å] | Intensity (qualitative) |
|---|---|---|
| 5.2 | 17.1 | w |
| 6.8 | 13.0 | vs |
| 8.9 | 10.0 | w |
| 10.3 | 8.6 | w |
| 11.2 | 7.9 | w |
| 12.5 | 7.1 | vw |
| 14.7 | 6.03 | vw |
| 15.5 | 5.71 | vs |
| 16.9 | 5.24 | m |
| 17.8 | 4.98 | m |
| 18.2 | 4.86 | w |
| 18.6 | 4.77 | m |
| 18.9 | 4.70 | w |
| 20.3 | 4.37 | m |
| 20.7 | 4.29 | w |
| 21.2 | 4.19 | vs |
| 21.8 | 4.08 | vw |
| 22.7 | 3.92 | w |
| 23.7 | 3.76 | w |
| 24.2 | 3.67 | w |
| 24.6 | 3.62 | m |
| 25.3 | 3.52 | w |
| 26.0 | 3.42 | vw |

| Angle [°2θ] | d-spacings [Å] | Intensity (qualitative) |
|---|---|---|
| 5.2 | 17.1 | w |
| 6.8 | 13.0 | vs |
| 8.9 | 10.0 | w |
| 26.3 | 3.38 | m |
| 26.8 | 3.32 | vw |
| 27.3 | 3.27 | m |
| 28.4 | 3.14 | vw |
| 28.7 | 3.10 | w |
| 29.3 | 3.05 | m |
| 30.1 | 2.97 | w |
| 32.6 | 2.75 | w |

b)

**[0123]** 181 mg of the product from Example 2 and 48.0 mg maleic acid were dissolved in 3.0 ml acetone. The solution was stored at 5 °C for 5 days. The solvent volume was reduced to a fourth by a nitrogen flow and storage at 5 ° continued for 48 hours. The solvent was evaporated at ambient condition and 2 ml heptane and 100 µl acetone were added under stirring. Stirring was continued for 24 hours. The precipitated solid was filtered off to yield 182 mg of a crystalline white powder. PXRD and Raman spectroscopy indicates a crystalline maletae, which was possibly admixed with another crystalline form. TG-FTIR shows a mass loss of about 5.9% between 60 and 160 °C, which was attributed to absorbed water, acetone and heptane. Decomposition started at about 170 °C. $^1$H-NMR complies with an 1:1 stoichiometry. The solubility in water was > 500 mg/ml.

Example 5: Preparation of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide malate of formula VII

**[0124]** 181 mg of the product from Example 1 and 56.0 mg L-(-)-malic acid were suspended in 10.0 ml ethyl acetate. The suspension was stirred at 60 °C for 30 min to form a clear solution. The solution was stored at 5 °C for 1 day. The solid was filtered off from the formed suspension yielding 155 mg of a crystalline white powder. PXRD and Raman spectroscopy indicate a crystalline form A. TG-FTIR shows a mass loss of about 5.5% between 50 and 160 °C, which was attributed to water and $CO_2$. Decomposition started at about 160 °C. Elemental analysis and $^1$H-NMR complies with an 1:1 stoichiometry. Solubility in water was > 500 mg/ml.

**[0125]** The powder X-ray diffraction pattern (PXRD) is shown in Figure 5 and the characteristic peaks in 2 theta with the corresponding d-spacing values in Å are given in Table 5.

Table 5: d-Spacings for the compound of formula VII

| Angle [°2θ] | d-spacings [Å] | Intensity (qualitative) |
|---|---|---|
| 4.4 | 19.8 | m |
| 5.5 | 16.2 | w |
| 6.8 | 13.1 | vs |
| 7.4 | 12.0 | s |
| 10.1 | 8.8 | w |
| 10.3 | 8.6 | w |
| 11.5 | 7.7 | m |
| 12.2 | 7.2 | m |
| 14.5 | 6.1 | m |
| 15.0 | 5.92 | w |
| 16.5 | 5.35 | s |
| 17.5 | 5.05 | s |
| 18.1 | 4.89 | m |
| 18.4 | 4.83 | s |
| 18.7 | 4.75 | vs |
| 18.8 | 4.71 | vs |
| 19.2 | 4.63 | m |
| 19.5 | 4.55 | m |
| 20.3 | 4.37 | vs |
| 20.7 | 4.29 | vs |
| 21.3 | 4.17 | s |
| 22.2 | 4.00 | s |
| 22.4 | 3.97 | m |
| 23.0 | 3.87 | s |
| 23.2 | 3.83 | s |
| 23.7 | 3.75 | vw |
| 24.7 | 3.61 | m |
| 25.0 | 3.56 | vw |
| 27.5 | 3.24 | m |
| 29.2 | 3.05 | w |
| 29.9 | 2.98 | w |
| 30.5 | 2.93 | w |

Example 6: Preparation of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide phosphate of formula VIII

**[0126]** 181 mg of the product from Example 1 was dissolved in 3 ml 2-propanol. 842 µl phosphoric acid (0.5. molar) was added and a clear solution was formed. The sample was stored for at 5 °C for 1 day. The precipitate was filtered off and dried in vacuum for 15 hours.

Yield was 60 mg of white crystalline powder. PXRD and Raman spectroscopy indicate a crystalline form A. TG-FTIR shows a mass loss of about 3.9% between 80 and 160 °C, which was attributed to 2-propanol. Decomposition started at about 170 °C. $^1$H-NMR complies with an 1:1 stoichiometry. Solubility in water was > 250 mg/ml.

**[0127]** The powder X-ray diffraction pattern (PXRD) is shown in Figure 6 and the characteristic peaks in 2 theta with the corresponding d-spacing values in Å are given in Table 6.

Table 6: d-Spacings for the compound of formula VIII

| Angle [°2θ] | d-spacings [Å] | Intensity (qualitative) |
|---|---|---|
| 5.1 | 17.3 | vs |
| 8.7 | 10.1 | m |
| 10.0 | 8.9 | m |
| 13.3 | 6.7 | w |
| 13.7 | 6.5 | m |
| 15.0 | 5.91 | s |
| 15.4 | 5.74 | m |
| 16.3 | 5.44 | vw |
| 17.2 | 5.16 | w |
| 18.0 | 4.93 | m |
| 18.5 | 4.80 | m |
| 18.7 | 4.75 | w |
| 19.4 | 4.56 | m |
| 20.8 | 4.27 | m |
| 21.5 | 4.14 | m |
| 23.0 | 3.86 | m |
| 23.5 | 3.78 | vw |
| 25.0 | 3.55 | m |
| 30.9 | 2.89 | w |

Example 7: Preparation of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide succinate of formula IX

a)

**[0128]** 90 mg of the product from Example 1 and 24.7 mg succinic acid were suspended in 5.0 ml ethyl acetate. The mixture was stirred at 60 °C for 15 min forming a clear solution. The solution was stored for 30 min at 23±2 °C and then cooled to 5 °C. Precipitation occurs after 30 min. The suspension was stored for 16 hours at 5 °C and the precipitate was

filtered off, washed with TBME and heptane to yield 55 mg of a crystalline white solid. PXRD and Raman spectroscopy indicate a crystalline form.

b)

[0129] 179 mg of the product from Example 1 and 48.9 mg succinic acid were suspended in 10.0 ml ethyl acetate. The mixture was stirred at 60 °C for 15 min forming a clear solution. The solution was stored for 40 min at 23±2 °C and then cooled to 5 °C. Precipitation occurs after 30 min. The suspension was stirred for 1 hour at 23 °C and the precipitate was filtered off, washed with heptane to yield 147 mg of a crystalline white powder. PXRD and Raman spectroscopy indicate a crystalline form. TG-FTIR shows a mass loss of about 18.8% between 60 and 250 °C, which was attributed to mostly CO2 and water. Elemental analysis indicates the formation of a dihydrate. $^1$H-NMR complies with an 1:1 stoichiometry. Solubility in water was > 500 mg/ml.

[0130] The powder X-ray diffraction pattern (PXRD) is shown in Figure 7 and the characteristic peaks in 2 theta with the corresponding d-spacing values in Å are given in table 7.

Table 7: d-Spacings for the compound of formula IX

| Angle [°2θ] | d-spacings [Å] | Intensity (qualitative) |
|---|---|---|
| 5.4 | 16.7 | vw |
| 6.9 | 12.8 | vs |
| 10.3 | 8.6 | w |
| 11.6 | 7.6 | m |
| 13.8 | 6.4 | w |
| 16.1 | 5.51 | s |
| 16.8 | 5.27 | w |
| 17.1 | 5.19 | m |
| 17.7 | 5.00 | vw |
| 18.5 | 4.79 | m |
| 19.1 | 4.65 | vw |
| 19.4 | 4.58 | vw |
| 20.1 | 4.42 | w |
| 20.5 | 4.32 | m |
| 21.4 | 4.16 | s |
| 21.9 | 4.05 | s |
| 22.7 | 3.91 | m |
| 23.2 | 3.83 | vw |
| 24.1 | 3.69 | w |
| 24.7 | 3.60 | vw |

| Angle [°2θ] | d-spacings [Å] | Intensity (qualitative) |
|---|---|---|
| 5.4 | 16.7 | vw |
| 6.9 | 12.8 | vs |
| 10.3 | 8.6 | w |
| 11.6 | 7.6 | m |
| 13.8 | 6.4 | w |
| 26.3 | 3.38 | vw |
| 26.9 | 3.31 | w |
| 27.3 | 3.27 | w |
| 28.0 | 3.19 | vw |
| 28.4 | 3.14 | w |
| 30.1 | 2.97 | w |
| 32.4 | 2.76 | w |
| 33.6 | 2.66 | w |
| 34.1 | 2.62 | w |

Example 8: Preparation of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide sulphate of formula X

[0131] 180 mg of the product from Example 1 was dissolved in 5 ml ethanol. 842 µl sulphuric acid (0.5 molar) was added and the formed clear solution was stored at 5 °C for 48 hours. The solvent was evaporated by a nitrogen flow. The solid residue was suspended in 3 ml TBME and 0.1 ml ethanol and the suspension was stirred for 17 hours at 23±2 °C. Filtration yields 80 mg of a crystalline white powder. PXRD and Raman spectroscopy indicate a crystalline form.

[0132] The powder X-ray diffraction pattern (PXRD) is shown in Figure 7 and the characteristic peaks in 2 theta with the corresponding d-spacing values in Å are given in table 8.

Table 8: d-Spacings for the compound of formula IX

| Angle [°2θ] | d-spacings [Å] | Intensity (qualitative) |
|---|---|---|
| 2.9 | 30.8 | w |
| 5.2 | 17.0 | vs |
| 9.2 | 9.6 | m |
| 10.7 | 8.3 | w |
| 11.5 | 7.7 | vw |
| 13.1 | 6.8 | m |
| 13.9 | 6.4 | m |
| 16.1 | 5.49 | vs |
| 16.7 | 5.29 | w |

| Angle [°2θ] | d-spacings [Å] | Intensity (qualitative) |
|---|---|---|
| 2.9 | 30.8 | w |
| 5.2 | 17.0 | vs |
| 9.2 | 9.6 | m |
| 10.7 | 8.3 | w |
| 11.5 | 7.7 | vw |
| 13.1 | 6.8 | m |
| 18.5 | 4.79 | s |
| 19.1 | 4.65 | m |
| 19.6 | 4.53 | s |
| 20.1 | 4.42 | m |
| 20.6 | 4.30 | vs |
| 21.2 | 4.18 | m |
| 21.4 | 4.15 | s |
| 22.0 | 4.04 | m |
| 22.9 | 3.89 | w |
| 24.7 | 3.60 | m |
| 25.0 | 3.56 | w |
| 26.3 | 3.38 | vw |
| 27.0 | 3.30 | w |
| 28.0 | 3.19 | vw |
| 28.5 | 3.13 | vw |
| 29.2 | 3.05 | vw |
| 31.6 | 2.83 | vw |
| 32.7 | 2.74 | w |

Example 9: Preparation of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide edisylate of formula XI

**[0133]** 180 mg of the product from Example 1 was dissolved in 2 ml dioxane and a solution of 48 mg 1,2-ethane disulphonic acid dihydrate in 4 ml dioxane was then added. The solution was stored at 8 °C for 10 days. The precipitated solid was filtered off to yield 206 mg of a crystalline white powder. PXRD and Raman spectroscopy indicate a crystalline form. TG-FTIR shows a mass loss of about 1.2% between 60 and 160 °C, which was attributed to dioxane. Decomposition starts at about 170 °C. Elemental analysis indicates a 2:1 stoichiometry (compound of formula I : 1,2-ethane disulphonic acid. $^1$H-NMR complies with both a 2:1 or a 1:1 stoichiometry. Solubility in water was 4 mg/ml. The crystalline powder remains a white powder when stored for 1 week at 60 °C and about 75% r.h. in a closed container (HPLC area was 97.4% compared to reference value of 96.8%). Storage for 1 week at 100 °C in a closed ampoule does

not decompose the crystalline product and the white powder remains substantially unchanged (HPLC area 97.4%).


[0134] The powder X-ray diffraction pattern (PXRD) is shown in Figure 9 and the characteristic peaks in 2 theta with the corresponding d-spacing values in Å are given in table 9.

Table 9: d-Spacings for the compound of formula XI

| Angle [°2θ] | d-spacings [Å] | Intensity (qualitative) |
|---|---|---|
| 7.3 | 12.1 | m |
| 8.1 | 10.9 | vw |
| 8.9 | 10.0 | s |
| 9.5 | 9.3 | m |
| 10.9 | 8.1 | m |
| 13.3 | 6.6 | m |
| 14.1 | 6.3 | vw |
| 14.6 | 6.05 | vs |
| 16.7 | 5.31 | s |
| 17.1 | 5.18 | m |
| 17.8 | 4.97 | vs |
| 18.4 | 4.81 | w |
| 18.9 | 4.68 | s |
| 19.4 | 4.57 | m |
| 19.9 | 4.46 | m |
| 20.4 | 4.35 | m |
| 20.8 | 4.26 | s |
| 21.5 | 4.12 | s |
| 21.9 | 4.05 | vw |
| 22.5 | 3.96 | m |
| 22.9 | 3.88 | w |
| 23.7 | 3.75 | m |
| 24.6 | 3.62 | m |
| 25.2 | 3.53 | w |
| 25.6 | 3.48 | m |
| 26.0 | 3.42 | w |
| 26.9 | 3.31 | m |
| 28.3 | 3.15 | w |
| 29.1 | 3.07 | w |
| 29.6 | 3.01 | vw |
| 35.9 | 2.49 | w |

**Claims:**

1. A salt of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide of formula I,



(I),

wherein the anion is selected from the group consisting of phosphate, sulphate, succinate, malate, fumarate and maleate.

2. The salt of claim 1, wherein the anion is selected from the group consisting of fumarate, maleate, malate, phosphate, succinate and sulphate.

3. The salt of claim 2, wherein when the anion is selected from the group consisting of maleate, malate, phosphate, succinate, and sulphate, the stoichiometry is 1: 1, and wherein when the anion is fumarate, the stoichiometry is 2: 1.

4. The salt of claim 1, wherein the salt is a fumarate salt.

5. The salt of claim 4 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 18.3, about 15.7, about 14.5, about 12.6, about 4.72, and about 4.47.


6. The salt of claim 4 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 15.7, about 12.6, about 4.72, and about 4.47.

7. The salt of claim 4 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 18.4, about 15.7, about 12.6, and about 5.50.

8. The salt of claim 4 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 18.4, about 15.7, about 12.6, about 5.50 and about 4.51.

9. The salt of claim 1, wherein the salt is a maleate salt.

10. The of salt of claim 9, wherein the salt is of formula VI:



(VI).

11. The salt of claim 9, wherein the salt is in the form of a hydrate.

12. The salt of claim 9 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 13.0, about 5.71, about 4.77 and about 4.19.

13. The salt of claim 9 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 13.0, about 5.71, and about 4.19.


14. The salt of claim 1, wherein the salt is a malate salt.

15. The salt of claim 14 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 13.1, about 12.0, about 5.35, about 4.75, about 4.37, about 4.29, about 4.00 and about 3.87.

16. The salt of claim 14 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 13.1, about 4.75, and about 4.29.

17. The salt of claim 1, wherein the salt is a phosphate salt.

18. The of salt of claim 17, wherein the salt is of formula VIII:



(VIII).

19. The salt of claim 17 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 17.3, about 5.91, about 4.80, about 4.27, about 4.14, and about 3.86.

20. The salt of claim 17 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 17.3, about 5.91, about 4.27, and about 4.14.

21. The salt of claim 1, wherein the salt is a succinate salt.

22. The succinate salt of claim 21, wherein the salt is in the form of a hydrate.

22. The salt of claim 21 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 12.8, about 5.51, about 4.79, about 4.16, and about 4.05.

23. The salt of claim 21 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 12.8, about 5.51, and about 4.05.

24. The salt of claim 1, wherein the salt is a sulphate salt.

25. The salt of claim 24 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 17.0, about 5.49, about 4.53, and about 4.30.

26. The salt of claim 24 that exhibits a X-ray powder diffraction pattern comprising peaks having d-values in angstroms of about 17.0, about 5.49, and about 4.30.

27. A process for the preparation of the salt of claim 1, comprising

a) forming a solution of the compound of formula I in an organic solvent;


b) adding an acid selected from the group consisting of fumaric acid, maleic acid, L-(-)-malic acid, phosphoric acid, succinic acid or sulphuric acid to said solution; and

c) isolating the salt.

28. The process of claim 27, wherein said isolating comprises separating the salt from a suspension formed after step b).

29. The process of claim 27, wherein said isolating comprises precipitating the salt from a solution formed after step b) by one or more of cooling, solvent removal, or adding a non-solvent.

30. A salt of N-(4-fluorobenzyl)-N-(1-methylpiperidin-4-yl)-N'-(4-(2-methylpropyloxy)phenylmethyl)carbamide of formula I,



(I),

produced by a process comprising:

a) forming a solution of the compound of formula I in an organic solvent;

b) adding an acid selected from the group consisting of fumaric acid, maleic acid, L-(-)-malic acid, phosphoric acid, succinic acid or sulphuric acid to said solution; and

c) isolating the salt.

31. A pharmaceutical composition comprising one or more salts of claim 1 and a pharmaceutically acceptable carrier.

32. The pharmaceutical composition of claim 31, wherein the composition is in a solid dosage form.

33. The pharmaceutical composition of claim 32, wherein the solid dosage form is suitable for oral administration.

34. The pharmaceutical composition of claim 31, wherein the amount of the salt of formula IV is from about 1 to 100 mg.

35. A salt or pharmaceutical composition of any one of claims 1-34 for use in the treatment of a neuropsychiatric disease.

36. The use of claim 35, wherein the neuropsychiatric disease is selected from the group consisting of psychosis, schizophrenia, schizoaffective disorders, mania, psychotic depression, affective disorders, dementia, anxiety, sleep disorders, appetite disorders, bipolar disorder, psychosis secondary to hypertension, migraine, vasospasm, and ischemia, motor tics, tremor, psychomotor slowing, bradykinesia, and neuropathic pain.

37. A salt or pharmaceutical composition of any one of claims 1-34 for use in the treatment of a neurodegenerative disease.

38. The use of claim 37, wherein the neurodegenerative disease is selected from the group consisting Parkinson's disease, Huntington's disease, Alzheimer's disease, Spinocerebellar Atrophy, Tourette's Syndrome, Fredrich's Ataxia, Machado-Joseph's disease, Lewy Body Dementia, Dystonia, Progressive Supranuclear Palsy, and Frontotemporal Dementia.

39. A salt or pharmaceutical composition of any one of claims 1-34 for use in the treatment of dyskinesia associated with dopaminergic therapy.

40. A salt or pharmaceutical composition of any one of claims 1-34 for use in the treatment of dystonia, myoclonus, or tremor associated with dopaminergic therapy.

41. A salt or pharmaceutical composition of any one of claims 1-34 for use in the treatment of a thrombotic condition.

42. The use of claim 41, wherein the thrombotic condition is selected from the group consisting of myocardial infarction, thrombotic or ischemic stroke, idiopathic and thrombotic thrombocytopenic purpura, peripheral vascular disease, and Raynaud's disease.

43. Use of a salt of any one of claims 1-30 in the manufacture of a medicament for the treatment of a neuropsychiatric disease.

44. The use of claim 43, wherein wherein the neuropsychiatric disease is selected from the group consisting of psychosis, schizophrenia, schizoaffective disorders, mania, psychotic depression, affective disorders, dementia, anxiety, sleep disorders, appetite disorders, bipolar

disorder, psychosis secondary to hypertension, migraine, vasospasm, and ischemia, motor tics, tremor, psychomotor slowing, bradykinesia, and neuropathic pain.

45. Use of a salt of any one of claims 1-30 in the manufacture of a medicament for the treatment of a neurodegenerative disease.

46. (New) The use of claim 45, wherein the neurodegenerative disease is selected from the group consisting Parkinson's disease, Huntington's disease, Alzheimer's disease, Spinocerebellar Atrophy, Tourette's Syndrome, Fredrich's Ataxia, Machado-Joseph's disease, Lewy Body Dementia, Dystonia, Progressive Supranuclear Palsy, and Frontotemporal Dementia.

47. Use of a salt of any one of claims 1-30 in the manufacture of a medicament for the treatment of dyskinesia associated with dopaminergic therapy.

48. Use of a salt of any one of claims 1-30 in the manufacture of a medicament for the treatment of dystonia, myoclonus, or tremor associated with dopaminergic therapy.

49. Use of a salt of any one of claims 1-30 in the manufacture of a medicament for the treatment of a thrombotic condition.

50. The use of claim 49, wherein the thrombotic condition is selected from the group consisting of myocardial infarction, thrombotic or ischemic stroke, idiopathic and thrombotic thrombocytopenic purpura, peripheral vascular disease, and Raynaud's disease.