


- 1 -

N, N'-BIS (QUINOLIN-4-YL)-DIAMINE DERIVATIVES, THEIR PREPARATION AND THEIR USE AS ANTIMALARIALS

The invention is concerned with novel diamines, namely N,N'-bis(quinolin-4-yl)-diamine derivatives of the general formula



10 wherein R¹ signifies halogen or trifluoromethyl, R² signifies hydrogen or halogen, A signifies cyclohexane-1,3-diyl, 2-methyl-cyclohexane-1,3-diyl, cyclohexane-1,4-diyl, dicyclohexylmethane-4,4'-diyl, cyclopentane-1,3-diyl, phenylene-1,4, phenylene-1,3, phenylene-1,2, ethene-1,2-
 15 diyl or ethyne-1,2-diyl and n and m can be the same or different and signify 0, 1 or 2,
 as well as their pharmaceutically acceptable salts.

20 These novel compounds have very good activities against not only chloroquine-sensitive, but also against chloroquine-resistant malaria pathogens. For this reason they are very well suited for preventing malaria and for treating it, especially where the pathogens are resistant to chloroquine.

25 Objects of the present invention are the novel bis-quinoline derivatives of general formula I as well as pharmaceutically usable salts thereof per se and as pharmaceutically active substances, the manufacture of these compounds and salts, medicaments containing a compound of formula I or a
 30 pharmaceutically acceptable acid addition salt thereof, the production of such medicaments and the use of the compounds of formula I and their pharmaceutically acceptable salts in the control or prevention of illnesses, especially of malaria, and, respectively, for the production of such medicaments.

In formula I R¹ preferably signifies chlorine, A preferably signifies cyclohexane-1,3-diyl, 2-methyl-cyclohexane-1,3-diyl, cyclohexane-1,4-diyl, dicyclohexylmethane-4,4'-diyl, cyclopentane-1,3-diyl, phenylene-1,4, phenylene-1,3 or
5 phenylene-1,2 and n and m preferably signify 0 or 1.

Compounds of formula I in which R¹ signifies chlorine, A signifies ethene-1,2-diyl and n and m signify 1 are also preferred.

10


Particularly preferred compounds of general formula I are:

- (7-Chloroquinolin-4-yl)-[2-(7-chloroquinolin-4-yl-aminomethyl)-benzyl]-amine,
15 (7-chloroquinolin-4-yl)-[3-(7-chloroquinolin-4-yl-aminomethyl)-benzyl]-amine
(7-chloroquinolin-4-yl)-[4-(7-chloroquinolin-4-yl-aminomethyl)-benzyl]-amine,
±-trans-N,N'-bis-(7-chloroquinolin-4-yl)-cyclohexane-1,4-
20 diamine,
cis-N,N'-bis-(7-chloroquinolin-4-yl)-cyclohexane-1,4-diamine,
cis-N,N'-bis-(7-chloroquinolin-4-yl)-cyclohexane-1,3-diamine,
25 ±-trans-N,N'-bis-(7-chloroquinolin-4-yl)-cyclohexane-1,3-diamine,
cis,cis-N,N'-bis-(7-chloroquinolin-4-yl)-2-methyl-cyclohexane-1,3-diamine,
N,N'-bis-(7-chloroquinolin-4-yl)-phenylene-1,3-diamine,
30 cis-N,N'-bis-(7-chloroquinolin-4-yl)-cyclopentane-1,3-diamine,
(7-chloroquinolin-4-yl)-[3-(7-chloroquinolin-4-yl-aminomethyl)-cyclohexylmethyl]-amine,
N,N'-bis(7-chloroquinolin-4-yl)-dicyclohexylmethane-4,4'-
35 diamine and
N,N'-bis-(7-chloroquinolin-4-yl)-2-butene-1,4-diamine.

The novel compounds of formula I can be manufactured in accordance with the invention by

a) reacting a compound of the general formula

5



wherein R¹ and R² have the significance set forth above and X signifies a leaving group,

10 with a diamine of the general formula



wherein n, m and A have the significance set forth above,
15 or an acid addition salt thereof, and

b) if desired, converting the compound obtained into a pharmaceutically acceptable acid addition salt.

20 Bis-quinoline derivatives of general formula I are manufactured according to variant a) of the process in accordance with the invention by reacting a corresponding compound of general formula II with the corresponding diamines of general formula III. α,α' -Diamino-o-xylene, α,α' -diamino-m-xylene, α,α' -diamino-p-xylene, $\pm$ -trans-cyclohexane-1,4-diamine, cis-cyclohexane-1,4-diamine, cis-cyclohexane-1,3-diamine, $\pm$ -trans-cyclohexane-1,3-diamine, cis,cis-2-methyl-cyclohexane-1,3-diamine, phenylene-1,3-diamine, cis-cyclopentane-1,3-diamine, 1,3-bis-(aminomethyl)-cyclohexane, 4,4'-diamino-dicyclohexyl-
25 methane or trans-2-butene-1,4-diamine can be used, for example, as the diamine. In place of the free diamines, their acid addition salts can also be used.

Suitable compounds of formula II are, for example, 4,7-
35 dichloroquinoline or 4-chloro-7-trifluoromethyl-quinoline.

The reaction is conveniently effected under a protective gas atmosphere in a temperature range of 100-200°C and in a solvent, with phenol, N-methyl-2-pyrrolidone, N,N-dimethylacetamide, ethoxyethanol or acetonitrile being preferred. The presence of a
5 tertiary amine such as triethylamine influences the reaction favourably, but is not essential. When the acid addition salt of a diamine is used, the presence of a tertiary amine such as triethylamine is required. The reaction time can vary between 2 and 28 hours.

10

The conversion into a pharmaceutically acceptable acid addition salt according to process variant b) is conveniently effected by adding an acid. Hydrochloric acid, methanesulphonic acid or acetic acid is especially preferred because of the
15 physiological compatibility of the corresponding salts. Convenient solvents which are especially suitable are: water, methanol, ethanol, isopropanol, diethyl ether, N,N-dimethylformamide or dioxan.

20 The diamines and quinoline derivatives which are required as starting materials for process variant a) are commercial products or can be prepared according to methods known per se.

As mentioned earlier, the bis-quinoline derivatives of
25 general formula I in accordance with the invention and their pharmaceutically usable salts have extremely valuable pharmacological properties.

The following Tables 1-3 show their activity against not
30 only chloroquine-resistant, but also chloroquine-sensitive malaria pathogens.

Test method for the determination of the activity against
Plasmodium falciparum in vitro

35

The preparations are tested on intraerythrocytary stages of *Plasmodium falciparum* from asynchronous cultures according to the method of Desjardin et al. (Desjardins, R.E. et al: Quantitative

assessment of antimalarial activity in vitro by a semiautomated microdilution technique. Antimicrob. Agents Chemother. 16, 710-718, (1979)).

- 5 The culture medium consists of RPMI 1640 with the addition of 25 mM HEPES, 25 mM NaHCO₃, 100 µg/ml neomycin and 105 human serum (A⁺). Human-A⁺ erythrocytes are used as the *Plasmodium falciparum* host cells. The parasites are maintained at 37°C in an atmosphere of 3% O₂, 4% CO₂, 93% N₂
10 and 95% relative humidity.

- In order to determine the activity, the preparations are dissolved in dimethyl sulphoxide, pre-diluted in the culture medium to a suitable starting concentration and subsequently
15 titrated-out on to microtitre plates in the 2nd stage over 6-7 steps. After the addition of the parasite culture (0.7% parasitemia in 2.5% erythrocyte suspension) the test plates are incubated under the conditions given above for 48 h - 72 h. The parasite growth in the different preparation concentrations is
20 determined using [G-³H]-hypoxanthin incorporation compared to untreated control cultures on the same test plates. The 50% growth inhibition (IC₅₀) is calculated according to logit regression analysis from the resulting dosage-activity curve.

- 25 The preparations are tested on at least one chloroquine-resistant and one chloroquine-sensitive *Plasmodium falciparum* strain. Additional sensitive and resistant strains are included for further characterization.

- 30 Test method for the determination of the activity against *Plasmodium berghei* in vivo

- The preparations are tested on mice infected with malaria pathogens (*Plasmodium berghei*). Male albino mice
35 (IBM:MORO(SPF), FUELLINS DORF) weighing about 25 g are used as the test animals. They are kept in climatized rooms at 21-22°C in groups of 5 animals per cage. They receive ad libitum a diet feed with a low PABA content (NAFAG FUTTER © No. 9009 PAB-45,

PABA content 45 mg/kg) and drinking water. On the first day of the test (D0) the test animals are infected with *Plasmodium berghei* (strain ANKA). For this there is used heparinized blood of a donor mouse with about 30% parasitemia, which is diluted with physiological saline such that it contains 10^8 parasitized erythrocytes per ml. 0.2 ml of this suspension is injected intravenously (i.v.) into the mice to be treated and into the control mice. In untreated control animals the parasitemia normally reaches 30-40% on the third day after the infection (D+3) and the test animals die between days +5 and +7.

The substances to be tested are dissolved or suspended in distilled water or in a mixture of 7% Tween 80, 3% alcohol (96%) and water. Usually, 0.25 ml of this solution or suspension is administered once subcutaneously and perorally to groups of 5 test animals. Treatment is effected 24 hours after the infection. 10 control animals are treated in the same manner with solvent or suspension medium per test.

All substances are tested in a first test in a single dosage of 10 mg/kg. Only those substances which in this test (10 mg/kg) have shown a parasitaemia reduction of 90% are used for the titration. Suitable dilutions of the test substance can be used to obtain an accurate titration of the activity.

48 hours after the treatment (D+3) blood smears are prepared from all animals using blood from tail veins and are stained with giemsa. The average erythrocyte infection rate (parasitemia in %) in the control groups as well as in the groups which have been treated with the test compounds is determined by counting under a microscope. The difference in the average values of the infection rates of control group (100%) and treated groups is calculated and expressed as a percentage reduction (GI%). The ED₅₀ or ED₉₀ is determined mathematically by means of the JMP programme (nonlinear fit). The ED₅₀ (ED₉₀) in mg/kg is that dose which after single administration reduces the average erythrocyte infection rate by 50% (90%) in comparison to the control group.

Table 1: Values measured in vitro for the growth inhibition of chloroquine-sensitive strains of the human-pathogenenic *Plasmodium falciparum* (IC50 values in ng/ml)

5

	Strain designation				
Example	NF54	FCH5C2	HB3	RFMEF3	Ro73
1	10.2	6.6	7.5	6.5	3.6
2	10.3	12.5	8.9	7.8	13.3
3	9.6	12.7	4.6	12.5	7.4
4	7.2	3.6	3.3	3.3	1.6
5	2.6	3.9	3.2	3.4	2.3
6	3.8	6.4	5.2	4.8	3.0
7	10.7				
8	7.7	6.1	6.1	6.3	1.6
9	23.7	15.6	9.5	10.8	10.4
11	17.2	14.5	12.7	16.8	4.9
12	11.6	19.4	26.3	13.7	17.8
13	30.8				
Chloroquine	7.6	12.2	11.3	14.2	8.0

Table 2: Value measured in vitro for the growth inhibition of chloroquine-resistant strains of the human-pathogenic *Plasmodium falciparum* (IC₅₀ values in ng/ml) and the ratio of average IC₅₀ values for resistant strains and average values for sensitive strains as a measurement of resistance towards the test substance ("resistance index").

	Strain designation								
Example	RFCF3	ItG2F6	Indo	K1	W2	7G8	W2 Mef	Resistance index	
1	5.4	4.7	6.9	6.9	5.8	7.0	4.9	0.9	
2	26.3	19.0	15.8	17.0	100.0	13.9	17.0	2.8	
3	5.7	7.9	2.9	9.1	5.0	6.8	9.7	0.7	
4	9.5	5.3	6.0	11.6	13.1	2.8	7.0	2.1	
5	5.1	6.8	4.4	7.7	4.8	9.0	6.5	2.1	
6	4.2	4.1	3.0	4.4	4.0	3.8	7.6	1.0	
7	10.5								
8	4.6	6.4	6.4	8.8	5.6	9.4	9.1	1.3	
9	12.5	12.9	7.7	17.1	12.1	17.1	12.1	0.9	
11	14.1	11.6	14.4	17.8	9.9	20.5	15.9	1.1	
12	7.2	13.3	13.7	13.0	25.0	17.1	21.6	0.9	
13	18.1								
Chloroquine	130.0	68.0	52.0	114.0	123.0	81.0	79.0	8.7	

Table 3: Activity measured in vivo against *Plasmodium berghei* in mice. GI % is the percentage parasitemia reduction after single, perorally (po) or subcutaneously (sc) administered dose of 10 mg/kg of test substance; ED₅₀ is the effective perorally administered dose of test substance

Example	GI % at 10 mg/kg po	GI % at 10 mg/kg sc	ED ₅₀ po mg/kg
1	62.0	97.0	
2	87.0	99.0	
3	99.9	99.9	2.5
4	73.0	37.0	
5	98.0	99.9	1.5
6	98.0	99.9	1.7
7	97.0	99.0	
8	66.0	67.0	
9	86.0	83.0	
11	28.0	99.6	
Chloroquine	99.9	99.9	2.0

The compounds of formula I and the pharmaceutically acceptable acid addition salts of the compounds of formula I can be used as medicaments, e.g. in the form of pharmaceutical preparations. The pharmaceutical preparations can be administered orally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions. The administration can, however, also be effected rectally, e.g. in the form of suppositories, or parenterally, e.g. in the form of injection solutions.

The compounds of formula I and the pharmaceutically acceptable acid addition salts of the compounds of formula I can be processed with pharmaceutically inert, inorganic or organic carriers for the production of pharmaceutical preparations. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts and the like can be used, for example, as such carriers for tablets, coated tablets, dragées and hard gelatine capsules. Suitable carriers for soft gelatine capsules are, for example,

vegetable oils, waxes, fats, semi-solid and liquid polyols and the like. Depending on the nature of the active ingredient no carriers are, however, usually required in the case of soft gelatine capsules. Suitable carriers for the production of solutions and
5 syrups are, for example, water, polyols, glycerol, vegetable oils and the like. Suitable carriers for suppositories are, for example, natural or hardened oils, waxes, fats, semi-liquid or liquid polyols and the like.

10 The pharmaceutical preparations can, moreover, contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, coating agents or antioxidants. They can also contain still other therapeutically valuable
15 substances.

Medicaments containing a compound of formula I or a pharmaceutically acceptable acid addition salt thereof and a therapeutically inert carrier are also an object of the present
20 invention, as is a process for their manufacture which comprises bringing one or more compounds of formula I and/or pharmaceutically acceptable acid addition salts thereof and, if desired, one or more other therapeutically valuable substances into a galenical administration form together with one or more
25 therapeutically inert carriers.

In accordance with the invention compounds of general formula I as well as their pharmaceutically acceptable acid addition salts can be used for the treatment or prevention of
30 malaria and, respectively, for the production of corresponding medicaments. The dosage can vary within wide limits and will, of course, be fitted to the individual requirements in each particular case. In the case of oral administration the dosage lies in a range of about 10 mg to about 2.5 g per day of a
35 compound of general formula I or the corresponding amount of a pharmaceutically acceptable acid addition salt thereof, although the upper limit can also be exceeded when this is found to be indicated.

In the following Examples, which illustrate the present invention but are not intended to limit its scope in any manner, all temperatures are given in degrees Celsius. The 250 MHz-¹H-NMR spectra were measured at room temperature; chemical shifts δ (ppm) relative to δ (TMS) = 0.0 ppm.

Example 1

(7-Chloroquinolin-4-yl)-[2-(7-chloroquinolin-4-yl-amino-methyl)-benzyl]-amine

2.67 g of 4,7-dichloroquinoline, 1.47 g of α,α' -diamino-o-xylene dihydrochloride and 3.8 ml of triethylamine are reacted at 140°C under argon for 5 hours. After cooling 100 ml of 2N NaOH and 100 ml of ethyl acetate are added, a small amount (0.68 g) of insoluble crystalline product is filtered off and the organic phase is separated. The residue remaining after evaporation of the solvent is combined with the filtered-off crystals and recrystallized from 25 ml of methanol. There is obtained 0.45 g of the amine as a colourless product which is dissolved in 5 ml of hot methanol. The dimethanesulphonate crystallizes out after the addition of 0.13 ml of methanesulphonic acid. There is obtained 0.56 g, m.p.: > 250°C.

Elementary analysis: $C_{26}H_{20}Cl_2N_4 \cdot 2(CH_3SO_3H) \cdot 0.5 H_2O$

Calculated:: C 51.35%, H 4.39%, Cl 10.64%, N 8.40%, S 9.62%

Found: C 51.14%, H 4.32%, Cl 10.68%, N 8.20%, S 9.64%

Example 2

(7-Chloroquinolin-4-yl)-[3-(7-chloroquinolin-4-yl-amino-methyl)-benzyl]-amine

3.96 g of 4,7-dichloroquinoline and 1.36 g of α,α' -diamino-m-xylene are reacted at 140°C in 4 g of phenol under argon for 24 hours. After cooling the mixture is treated with 20 ml of ethyl acetate and 30 ml of water and adjusted to pH 1 by adding a

small amount of concentrated hydrochloric acid. The product which thereby crystallizes out is filtered off under suction and triturated in 200 ml of 2N NaOH and 50 ml of ethyl acetate. The separated crystalline amine is suspended in 80 ml of 0.5N HCl and brought into solution on a steam bath by the addition of ethanol (75 ml). 3.14 g of the dihydrochloride crystallize out upon cooling, m.p.: >250°C

Elementary analysis: $C_{26}H_{20}Cl_2N_4 \cdot 2HCl \cdot H_2O$

10 Calculated:: C 56.75%, H 4.40%, Cl 25.77%, N 10.18%

Found: C 56.54%, H 4.23%, Cl 25.79%, N 9.89%

1H -NMR in DMSO- d_6 , δ (ppm): 3.35 (s, 2H [H_2O]), 4.79 (d, 4H, $J = 6$ Hz), 6.58 (d, 2H, $J = 7.5$ Hz), 7.24 (s, 1H), 7.37 (m, 3H), 7.69 (dd, 2H, $J = 2.5$ and 9 Hz), 8.04 (d, 2H, $J = 2.5$ Hz), 8.38 (d, 2H, $J = 7.5$ Hz), 8.59 (d, 2H, $J = 9$ Hz), 10.23 (t, 2H, $J = 6$ Hz).

Example 3

20 (7-Chloroquinolin-4-yl)-[4-(7-chloroquinolin-4-yl-amino-methyl)-benzyl]-amine

2.9 g of 4,7-dichloroquinoline, 1 g of α, α' -diamino-p-xylene and 1.3 g of triethylamine are reacted at 120°C in 10 ml of ethoxy-ethanol under argon for 16 hrs. After cooling 40 ml of water and 40 ml of ethyl acetate are added and the mixture is adjusted to pH 1 by means of concentrated hydrochloric acid. The product which thereby separates is triturated in 150 ml of 2N NaOH with the addition of 100 ml of ethyl acetate, 0.8 g of crystalline amine being obtained. 0.84 g of crystalline dihydrochloride, m.p.: >250°C, is obtained therefrom by boiling in 15 ml of 1N HCl and 10 ml of ethanol.

Elementary analysis: $C_{26}H_{20}Cl_2N_4 \cdot 2HCl$

35 Calculated:: C 58.67%, H 4.17%, Cl 26.64%, N 10.53%

Found: C 56.40%, H 4.22%, Cl 26.93%, N 10.38%

¹H-NMR in DMSO-d₆ + D₂O, δ (ppm): 4.57 (s, 4H), 5.54 (d, 2H, J = 7.5 Hz), 7.26 (s, 1H), 7.56 (dd, 2H, J = 2 and 9 Hz), 7.75 (d, 2H, J = 2 Hz), 8.15 (d, 2H, J = 7.5 Hz), 8.23 (d, 2H, J = 9 Hz)

5

Example 4

±-trans-N,N'-Bis-(7-chloroquinolin-4-yl)-cyclohexane-1,4-diamine

10 Analogously to Example 3, from 2.28 g of ±-trans-cyclohexane-1,4-diamine and 7.92 g of 4,7-dichloroquinoline there are obtained 2 g of the dihydrochloride; colourless crystals from methanol-water, m.p.: >250°C.

15 Elementary analysis: C₂₄H₂₂Cl₂N₄·2HCl

Calculated:: C 56.49%, H 4.74%, Cl 27.79%, N 10.98%

Found: C 56.51%, H 4.63%, Cl 28.03%, N 10.98%

Example 5

20

cis-N,N'-Bis-(7-chloroquinolin-4-yl)-cyclohexane-1,4-diamine

The dihydrochloride, colourless crystals from methanol-water, m.p.: >260°C, is obtained analogously to Example 3 from cis-
25 cyclohexane-1,4-diamine and 4,7-dichloroquinoline.

Elementary analysis: C₂₄H₂₂Cl₂N₄·2HCl

Calculated:: C 56.49%, H 4.74%, Cl 27.79%, N 10.98%

Found: C 56.76%, H 4.73%, Cl 27.57%, N 10.83%

30

¹H-NMR in DMSO-d₆; δ(ppm): 1.86 (m, 4H); 2.00 (m, 4H); 3.81 (m, 2H); 6.63 (d, 2H); 7.14 (d, 2H); 7.50 (dd, 2H); 7.82 (d, 2H); 8.44 (d, 2H); 8.48 (d, 2H).

Example 6cis-N,N'-Bis-(7-chloroquinolin-4-yl)-cyclohexane-1,3-diamine

- 5 23.2 g of cis-cyclohexane-1,3-diamine and 80.5 g of 4,7-dichloroquinoline are reacted at 150°C in 400 ml of N-methyl-2-pyrrolidone under argon for 24 hours. After cooling insoluble material is filtered off and the filtrate is treated with 600 ml of water. The separated crude product is introduced into 400 ml
10 of water and 100 ml of glacial acetic acid. The mixture is extracted three times with 200 ml of ethyl acetate each time and then adjusted to pH 12 by the addition of 600 ml of 2N NaOH. The separated crystalline diamine is recrystallized from 270 ml of dimethylformamide, 20.1 g of pure product being obtained.
15 This is converted into the dimethanesulphonate in 200 ml of boiling acetonitrile with 6.4 ml of methanesulphonic acid. 19.5 g of colourless crystalline salt, m.p.: > 250°C, are obtained.

Elementary analysis: $C_{24}H_{22}Cl_2N_4 \cdot 2(CH_3SO_3H)$

- 20 Calculated:: C 49.60%, H 4.80%, Cl 11.26%, N 8.90%, S 10.18%
Found: C 49.50%, H 5.00%, Cl 11.34%, N 8.77%, S 10.21%

- 1H -NMR in DMSO- d_6 + D_2O , δ (ppm): 1.6 -2.45 (multiple m, 8H), 2.52 (s, 6H), 4.15 (m, 2H), 7.08 (d, 2H, $J = 7.5$ Hz), 7.76 (dd, 2H, $J = 2$
25 and 8.5 Hz), 7.94 (d, 2H, $J = 2$ Hz), 8.46 (d, 2H, $J = 7.5$ Hz), 8.56 (d, 2H, $J = 8.5$ Hz)

Example 7

- 30 $\pm$ -trans-N,N'-Bis-(7-chloroquinolin-4-yl)-cyclohexane-1,3-diamine

- Analogously to Example 6, from 8.5 g of $\pm$ -trans-cyclohexane-1,3-diamine bis-trifluoroacetate, 10.1 g of triethylamine and
35 9.9 g of 4,7-dichloroquinoline there are obtained 1.6 g of pure diamine. This is dissolved in 16 ml of hot isopropanol. The dihydrochloride crystallizes out after the addition of 1.6 ml of

4.8N isopropanolic hydrochloric acid. 1.04 g of colourless crystallize are obtained, m.p.: >250°.

¹H-NMR in DMSO-d₆ + D₂O, δ(ppm): 1.84 (broad m, 6H), 2.24 (m, 2H),
5 4.52 (m, 2H), 7.02 (d, 2H, J = 7.5 Hz), 7.78 (dd, 2H, J = 2 und 9.5 Hz), 8.02 (d, 2H, J = 2 Hz), 8.55 (d, 2H, J = 7.5 Hz), 8.86 (d, 2H, J = 9.5 Hz)..

Example 8

10

cis,cis-N,N'-Bis-(7-chloroquinolin-4-yl)-2-methyl-cyclohexane-1,3-diamine

Analogously to Example 3, from 3.36 of cis,cis-2-methyl-
15 cyclohexane-1,3-diamine dihydrochloride, 10.38 g of 4,7-dichloroquinoline and 5.3 g of triethylamine there are obtained 1.23 g of pure diamine. From a hot solution in 10 ml of glacial acetic acid there crystallize, after the addition of 0.7 ml of methanesulphonic acid, 1.23 g of colourless dimethanesulphonate
20 which is recrystallized from methanol; m.p.: >250°C.

Elementary analysis: C₂₅H₂₄Cl₂N₄.2(CH₃SO₃H)

Calculated:: C 50.39%, H 5.01%, Cl 11.02%, N 8.71%, S 09.96%

Found: C 50.28%, H 4.97%, Cl 11.11%, N 8.53%, S 10.03%

25

¹H-NMR in DMSO-d₆, δ(ppm): 0.90 (d, 3H, J = 6 Hz), 1.5 - 2.2
(multiple m, 6H), 2.44 (s, 6H), 2.64 (m, 1H), 4.402 (m, 2H), 7.44 (d,
2H, J = 8 Hz), 7.80 (dd, 2H, J = 2 and 9.5 Hz), 7.97 (d, 2H, J = 2 Hz),
8.62 (d, 2H, J = 8 Hz), 8.80 (d, 2H, J = 9.5 Hz), 8.87 (d, 2H, J = 8
30 Hz), 13.9 (breit, 2H).

Example 9

N,N'-Bis-(7-chloroquinolin-4-yl)-phenylene-1,3-diamine

35

0.54 g of phenylene-1,3-diamine and 2 g of 4,7-dichloro-quinoline are dissolved in 20 ml of 1-methyl-2-pyrrolidone and heated to 140°C while stirring for 16 hours. After cooling the reaction

solution is added to 60 ml of ethyl acetate. The crystals which thereby separate are filtered off under suction and recrystallized from HCl-saturated methanol solution. There is obtained 0.77 g: m.p.: >280°C

5

Elementary analysis: $C_{24}H_{16}Cl_2N_4 \cdot 2(HCl)$

Calculated:: C 57.17%, H 3.60%, Cl 28.12%, N 11.11%

Found: C 57.08%, H 3.61%, Cl 27.98%, N 10.90%

10 1H -NMR in DMSO- d_6 , δ (ppm): 7.04 (d, 2H); 7.51 (dd, 2H); 7.59 (d, 1H); 7.75 (t, 1H); 7.90 (dd, 2H); 8.10 (d, 2H); 8.63 (d, 2H); 8.78 (d, 2H); 10.92 (s, 2H).

Example 10

15

cis-N,N'-Bis-(7-chloroquinolin-4-yl)-cyclopentane-1,3-diamine

3.6 g of pure diamine are obtained from 2.62 g of cis-cyclopentane-1,3-diamine, 4 g of 4,7-dichloroquinoline and 2 g of triethylamine analogously to Example 6. 2.5 g of di-methanesulphonate, m.p.: >25°C, crystallize from a solution of 2 g of the base in 20 ml of glacial acetic acid upon adding 0.5 ml of methanesulphonic acid and 20 ml of diethyl ether.

25 Elementary analysis: $C_{23}H_{20}Cl_2N_4 \cdot 2(CH_3SO_3H)$

Calculated:: C 48.78%, H 4.59%, Cl 11.52%, N 9.10%, S 10.42%

Found: C 48.62%, H 4.37%, Cl 11.13%, N 8.86%, S 10.23%

30 1H -NMR in DMSO- d_6 , δ (ppm): 1.95-2.3 (multiple m, 5H), 2.39 (s, 6H), 2.85 (m, 1H), 4.42 (m, 2H), 7.01 (d, 2H, J = 7.5 Hz), 7.81 (dd, 2H, J = 2 and 9 Hz), 7.97 (d, 2H, J = 2 Hz), 8.62 (d, 2H, J = 7.5 Hz), 8.74 (d, 2H, J = 9 Hz), 9.16 (d, 2H, J = 8 Hz), 13.87 (broad, 2H).

Example 11(7-Chloroquinolin-4-yl)-[3-(7-chloroquinolin-4-yl-amino-methyl)-cyclohexylmethyl]-amine

5

Analogously to Example 3, from 10 g of 1,3-bis-(aminomethyl)-cyclohexane, 27.9 g of 4,7-dichloroquinoline and 19.6 g of triethylamine there are obtained 6.8 g of crystalline diamine.

This is converted in 175 ml of methanol using 4.9 ml of
10 methanesulphonic acid into the di-methanesulphonate which is recrystallized from 43 ml of water, 17 ml of acetone and 175 ml of diethyl ether. There are obtained 5.6 g of colourless crystalline product, m.p.: 92-95°C.

15 Elementary analysis: $C_{26}H_{26}Cl_2N_4 \cdot 2.2(CH_3SO_3H)$

Calculated:: C 50.05%, H 5.18%, Cl 10.48%, N 8.28%, S 10.42%

Found: C 50.15%, H 5.36%, Cl 10.70%, N 8.34%, S 10.16%

Example 12

20

N,N'-Bis-(7-chloroquinolin-4-yl)-dicyclohexylmethane-4,4'-diamine

6.3 g of colourless, crystalline dihydrochloride, m.p.: >250°C, are
25 obtained from 4.2 g of 4,4'-diaminodicyclohexylmethane and 7.92 g of 4,7-dichloroquinoline.

Elementary analysis: $C_{31}H_{34}Cl_2N_4 \cdot 2HCl \cdot 0.5H_2O$

Calculated:: C 59.62%, H 6.13%, Cl 22.71%, N 8.97%

30 Found: C 59.60%, H 5.86%, Cl 23.22%, N 9.05%

Example 13N,N'-Bis-(7-chloroquinolin-4-yl)-2-butene-1,4-diamine

35

In analogy to Example 6, from 1.6 g of trans-2-butene-1,4-diamine dihydrochloride, 4 g of 4,7-dichloroquinoline and 4 g of triethylamine there are obtained 1.5 g of the base. This is

converted into the bis-methanesulphonate by adding 0.5 ml of methanesulphonic acid to a boiling solution in 28 ml of methanol and 12 ml of glacial acetic acid. There is obtained 0.86 g of colourless crystals, m.p.: >250°C

5

Elementary analysis: $C_{22}H_{20}Cl_2N_4 \cdot 2(CH_3SO_3H)$

Calculated:: C 47.92%, H 4.36%, Cl 11.79%, N 9.31%, S 10.66%

Found: C 47.98%, H 4.36%, Cl 11.77%, N 9.15%, S 10.58%

10 1H -NMR in DMSO- d_6 , δ (ppm): 2.41 (s, 6H), 4.24 (s, broad, 4H), 5.86 (s, broad, 2H), 6.83 (d, 2H, $J = 7.5$ Hz), 7.79 (dd, 2H, $J = 2$ and 9 Hz), 7.98 (d, 2H, $J = 2$ Hz), 8.52 (d, 2H, $J = 9$ Hz), 8.56 (d, 2H, $J = 7.5$ Hz), 9.63 (t, 2H), 13.85 (broad, 2H).

15 The intermediates used in the foregoing Examples, insofar as they are not commercially available, can be prepared in the following manner:

Example 14

20

α,α' -Diamino-ortho-xylene (see Example 1)

12.3 g of α,α' -dibromoxylene and 17.3 g of potassium phthalimide are stirred in 390 ml of N,N-dimethylformamide at a bath
25 temperature of 150°C for 24 hours. After cooling the resulting intermediate is filtered off under suction, washed with ethyl acetate, dried and then reacted in 200 ml of ethanol with 4.6 ml of hydrazine hydrate under reflux. After cooling the resulting suspension is treated with 50 ml of 25% hydrochloric acid,
30 filtered and concentrated to 1/3 of its volume. After repeated filtration the mixture is adjusted to pH 14 by the addition of concentrated sodium hydroxide solution and evaporated. The product is dissolved out from the residue with 300 ml of methylene chloride. After evaporation of the solvent the residue
35 is taken up in 25 ml of isopropanol and the dihydrochloride of the product is crystallized with 20 ml of 3.26N isopropanolic hydrochloric acid. There are obtained 2.64 g, colourless crystals; m.p.: >250°C.

¹H-NMR in DMSO-d₆, δ(ppm): 4.16 (s, 4H), 7.44 (m, 2H), 7.50 (m, 2H), 8.67 (s, 6H)

5

Example 15

±-trans-Cyclohexyl-1,3-diamine and cis-cyclohexyl-1,3-diamine
(see Examples 6 and 7)

- 10 50 g of phenylene-1,3-diamine in 508 ml of 1N NaOH are treated with 222 g of di-tert.-butyl dicarbonate dissolved in 500 ml of dioxan. The mixture is stirred at RT for 24 hours and thereafter evaporated in a vacuum. 700 ml of water are added to the residue and the mixture is adjusted to pH 1 with conc.
- 15 hydrochloric acid and extracted 5 times with 500 ml of ethyl acetate each time. After evaporation of the solvent the residue is recrystallized from 1200 ml of hexane and 850 ml of tert.-butyl methyl ether. 88.1 g of bis-BOC derivative, m.p.: 145-148°C, are obtained. This is hydrogenated on 30 g of Rh/C in 4 l
- 20 of methanol at 50 bar and 65°C. After filtering off from the catalyst the solvent is removed in a vacuum and the residue is recrystallized from 1800 ml of methanol. There is obtained a first crystal fraction of 33 g and, after concentration of the filtrate to about 800 ml, a second crystal fraction of 5.6 g. A
- 25 further 14.4 g are obtained after boiling the filtered-off catalyst residue with 900 ml of methanol. The three crystallizates consist of pure cis-N,N'-bis-BOC-cyclohexane-1,3-diamine; m.p.: 232°C (dec.); total 53 g.
- 30 From the mother liquor of the above crystallization there are obtained, after evaporation and recrystallization from 160 ml of hexane and 17 ml of ethyl acetate, 10 g of trans-N,N'-bis-BOC-cyclohexane-1,3-diamine as colourless crystals: m.p.: 131-132°C.
- 35 cis-Cyclohexane-1,3-diamine: 93.4 g of the BOC-derivative are introduced into 500 ml of trifluoroacetic acid while cooling with ice. After stirring at RT for 4 hours the trifluoroacetic acid is evaporated, the residue is dissolved in 200 ml of water,

concentrated sodium hydroxide solution is added until the pH has reached 14 and the mixture is again evaporated to dryness. The cis-cyclohexane-1,3-diamine is then taken up with 500 ml of dichloromethane and, after evaporation of the solvent, is isolated
5 as an oil (23.2 g).

¹H-NMR in DMSO-d₆, δ(ppm): 0.70 - 0.91 (m, 3H), 1.14 (m, 1H), 1.57 (m, 1H), 1.67 (m, 2H), 1.84 (m, 1H), 2.5 (m, 2H)

10

±-trans-Cyclohexane-1,3-diamine: 22.9 g of the BOC derivative are introduced into 100 ml of trifluoroacetic acid while cooling with ice. The mixture is left at RT for 4 hours, evaporated and the residue is recrystallized from 200 ml of acetonitrile. 8.5 g
15 of ±-trans-cyclohexane-1,3-diamine trifluoroacetate, m.p. 200°C (sintering) are obtained.

¹H-NMR in DMSO-d₆, δ(ppm): 1.5 - 1.8 (m, 6H), 1.88 (t, 2H), 3.49 (m, 2H)

20

Example 16

cis,cis-2-Methyl-cyclohexane-1,3-diamine (see Example 8)

25 cis,cis-2-Methyl-cyclohexane-1,3-diamine is obtained as an oil in analogy to Example 15 from 2,6-diaminotoluene after conversion into the bis-BOC derivative, hydrogenation on Rh/C and cleavage of the BOC protecting groups.

30 ¹H-NMR in DMSO-d₆, δ(ppm): 0.76 (d, 3H, J = 7.5 Hz), 1.21 (m, 3H), 1.43 (m, 2H), 1.61 (m, 1H), 1.89 (m, 1H), 2.65 (m, 2H)

Example 17

35 cis-Cyclopentane-1,3-diamine (see Example 10)

88.7 g of diethyl 2,3-diazabicyclo[2.2.1]heptane-2,3-dicarboxylate (prepared according to P.G. Gassman and K.T. Mansfield,

Organic Synthesis 49, 1-6) are reduced with 16.8 g of sodium in 150 ml of tetrahydrofuran and 1.5 l of liquid ammonia. After the addition of 39.2 g of ammonium chloride the ammonia is evaporated and the residue is taken up with 250 ml of dichloromethane and 250 ml of 2N hydrochloric acid. The aqueous phase is separated, back-washed with two 50 ml portions of methylene chloride and the organic phases are combined and evaporated to dryness. The residue is recrystallized from 600 ml of toluene and 300 ml of hexane. 78 g of cis-cyclopentane-1,3-diamine bis-ethylcarbamate (m.p.: 110-111°C) are obtained.

33 g of the bis-carbamate are boiled under reflux in 1 l of 33% HBr in glacial acetic acid for 5 hours. The separated product is filtered off under suction and washed with five 50 ml portions of acetone. 27.8 g of cis-cyclopentane-1,3-diamine dihydrobromide (m.p. > 250°C) are obtained.

Elementary analysis: $C_5H_{12}N_2 \cdot 2HBr$

Calculated:: C 22.92%, H 5.39%, Br 61.00%, N 10.69%

Found: C 23.08%, H 5.35%, Br 61.14%, N 10.63%

Example A

7-Chloroquinolin-4-yl)-[2-(7-chloroquinolin-4-yl-aminomethyl)-benzyl]-amine can be formulated as the active ingredient according to methods known per se to give pharmaceutical preparations of the following composition:

1. 500 mg tablets

30

Active ingredient	500 mg
Lactose powd.	149 mg
Polyvinylpyrrolidone	15 mg
Diocetyl sodiumsulphosuccinate	1 mg
Na carboxymethylstarch	30 mg
Magnesium stearate	<u>5 mg</u>
	700 mg

2. 50 mg tablets

	Active ingredient	50 mg
	Lactose powd.	50 mg
5	Microcrystalline cellulose	82 mg
	Na carboxymethylstarch	<u>15 mg</u>
		200 mg

3. 100 mg capsules

10	Active ingredient	100.0 mg
	Lactose powd.	104.7 mg
	Corn starch	70.0 mg
	Hydroxypropylmethylcellulose	10.0 mg
15	Diethyl sodiumsulphosuccinate	0.3 mg
	Talc	12.0 mg
	Magnesiumstearate	<u>3.0 mg</u>
		300.0 mg

4. 500 mg suppositories

	Active ingredient	500 mg
	Suppository mass	ad 2000 mg

5. 100 mg suppository


	Active ingredient	100 mg
	Medium chain triglyceride	<u>300 mg</u>
		400 mg

Claims



1. N,N'-Bis(quinolin-4-yl)-diamine derivatives of the general formula

5



wherein R¹ signifies halogen or trifluoromethyl, R² signifies hydrogen or halogen, A signifies cyclohexane-1,3-diyl, 2-methyl-cyclohexane-1,3-diyl, cyclohexane-1,4-diyl, dicyclohexylmethane-4,4'-diyl, cyclopentane-1,3-diyl, phenylene-1,4, phenylene-1,3, phenylene-1,2, ethene-1,2-diyl or ethyne-1,2-diyl and n and m can be the same or different and signify 0, 1 or 2,

as well as their pharmaceutically acceptable acid addition salts.

2. Compounds according to claim 1, wherein R¹ signifies chlorine, A signifies cyclohexane-1,3-diyl, 2-methyl-cyclohexane-1,3-diyl, cyclohexane-1,4-diyl, dicyclohexylmethane-4,4'-diyl, cyclopentane-1,3-diyl, phenylene-1,4, phenylene-1,3 or phenylene-1,2 and n and m signify 0 or 1.

3. Compounds according to claim 1, wherein R¹ signifies chlorine, A signifies ethene-1,2-diyl and n and m signify 1.

25

4. (7-Chloroquinolin-4-yl)-[2-(7-chloroquinolin-4-yl-aminomethyl)-benzyl]-amine,

(7-chloroquinolin-4-yl)-[3-(7-chloroquinolin-4-yl-aminomethyl)-benzyl]-amine

(7-chloroquinolin-4-yl)-[4-(7-chloroquinolin-4-yl-aminomethyl)-benzyl]-amine,

±-trans-N,N'-bis-(7-chloroquinolin-4-yl)-cyclohexane-1,4-diamine,

30

cis-N,N'-bis-(7-chloroquinolin-4-yl)-cyclohexane-1,4-diamine,

cis-N,N'-bis-(7-chloroquinolin-4-yl)-cyclohexane-1,3-diamine,

5 ±-trans-N,N'-bis-(7-chloroquinolin-4-yl)-cyclohexane-1,3-diamine,

cis,cis-N,N'-bis-(7-chloroquinolin-4-yl)-2-methyl-cyclohexane-1,3-diamine,

N,N'-bis-(7-chloroquinolin-4-yl)-phenylene-1,3-diamine,

10 cis-N,N'-bis-(7-chloroquinolin-4-yl)-cyclopentane-1,3-diamine,

(7-chloroquinolin-4-yl)-[3-(7-chloroquinolin-4-yl-aminomethyl)-cyclohexylmethyl]-amine,

15 N,N'-bis(7-chloroquinolin-4-yl)-dicyclohexylmethane-4,4'-diamine and

N,N'-bis-(7-chloroquinolin-4-yl)-2-butene-1,4-diamine


as well as their pharmaceutically acceptable acid addition salts.

20 5. Compounds according to any one of claims 1-4 for use as therapeutically active substances.

6. Compounds according to any one of claims 1-4 for use as therapeutically active substances against not only chloro-
25 quine-resistant, but also chloroquine-sensitive malaria pathogens.

7. A process for the manufacture of compounds of general formula I according to any one of claims 1-4, which
30 process comprises

a) reacting a compound of the general formula



II





5



10

15

30 14. Compounds according to any one of claims 1-4, insofar as they are manufactured according to the process claimed in claim 7 or a process equivalent thereto.

* * *




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 215/44, 215/46, A61K 31/47		A1	(11) International Publication Number: WO 95/35287
			(43) International Publication Date: 28 December 1995 (28.12.95)
(21) International Application Number: PCT/EP95/02123		(81) Designated States: AU, CA, CN, JP, KE, LK, MX, NZ, <u>SG</u> , US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 3 June 1995 (03.06.95)			
(30) Priority Data: 1928/94-0 17 June 1994 (17.06.94) CH		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(71) Applicant (for all designated States except US): F.HOFFMANN-LA ROCHE AG [CH/CH]; Gren- zacherstrasse 124, CH-4002 Basle (CH).		SG 9612089-4	
(72) Inventors; and (75) Inventors/Applicants (for US only): HOFHEINZ, Werner [DE/CH]; Talmattweg 7, CH-4103 Bottmingen (CH). LE- UPIN, Werner [CH/CH]; Bündtenstrasse 8, CH-4410 Liestal (CH).		SG's copy	
(74) Agents: POPPE, Regina; P.O. Box 3255, CH-4002 Basle (CH) et al.			

(54) Title: N,N'-BIS(QUINOLIN-4-YL)-DIAMINE DERIVATIVES, THEIR PREPARATION AND THEIR USE AS ANTIMALARIALS

(57) Abstract

Disclosed are N,N'-bis(quinolin-4-yl)-diamine derivatives of general formula (I) wherein R¹ signifies halogen or trifluoromethyl, R² signifies hydrogen or halogen, A signifies cyclohexane-1,3-diyl, 2-methyl-cyclohexane-1,3-diyl, cyclohexane-1,4-diyl, dicyclohexylmethane-4,4'-diyl, cyclopentane-1,3-diyl, phenylene-1,4, phenylene-1,3, phenylene-1,2, ethene-1,2-diyl or ethyne-1,2-diyl and n and m can be the same or different and signify 0, 1 or 2, as well as their pharmaceutically acceptable salts. These products are useful as agents for preventing malaria and for treating it, especially where the pathogens are resistant to chloroquine.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

9612089-4