

**Pharmaceutical Formulations Containing Relacorilant, a Heteroaryl-Ketone Fused
Azadecalin Compound**

CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application claims priority to, and the benefit of, U.S. Application Serial No. 62/781,983, filed December 19, 2018, the entire contents of which application is hereby incorporated by reference in its entirety.

BACKGROUND

[0002] Many heteroaryl-ketone fused azadecalin compounds bind to glucocorticoid receptors (GRs) and may thereby have therapeutic activity. Such compounds and their activities are disclosed, for example, in U.S. Patent 8,859,774; U.S. Patent 9,273,047; U.S. Patent 9,707,223; U.S. Patent 9,943,505; U.S. Patent 9,956,216; and others. Pharmaceutical formulations containing heteroaryl-ketone fused azadecalin compounds may be used for administration of those compounds to humans or animals for therapeutic purposes.

[0003] GRs play an important role in normal physiology, and the disruption or alteration of GR function is implicated in many diseases and disorders. The most important GR ligand in humans is cortisol. Cortisol binding to GR activates GR. Compounds which bind GRs and which affect GR activity may be termed glucocorticoid receptor modulators (GRMs). GRMs which reduce GR activity, e.g. by interfering with or competing for ligand binding to GR, are termed GR antagonists (GRAs). Diseases and disorders of abnormal GR activation include Cushing's syndrome (typically caused by excess cortisol), Addison's disease (typically caused by inadequate levels of cortisol), and others. Therapeutic uses for GRM compounds, whether as single agents or in combination with other agents, include treatments for endocrine disorders including Cushing's syndrome and Cushing's disease, cancer, immune system disorders, cardiovascular disorders including hypertension, diabetes, hyperglycemia, liver diseases, bone diseases, obesity, antipsychotic-induced weight gain, psychiatric disorders, addictive behaviors and disorders, and other diseases and disorders.


[0004] However, many GRM compounds are difficult to formulate in compositions suitable for administration to subjects in need of the compounds. Many such compounds are only sparingly soluble in water, may be incompatible with other ingredients typically used in pharmaceutical compositions, or may have other properties which make it difficult to prepare therapeutic formulations. Many GRM compounds are unstable, degrading over time, or with exposure to light or heat. Such characteristics of one heteroaryl-ketone fused azadecalin compound may differ from those of another heteroaryl-ketone fused azadecalin compound, so that a formulation suitable for one heteroaryl-ketone fused azadecalin compound may not be suitable for a different heteroaryl-ketone fused azadecalin compound.

[0005] For example, many heteroaryl-ketone fused azadecalin compounds are poorly soluble in water, and may have poor bioavailability in some formulations. Poor solubility or bioavailability necessitates the administration of large volumes of solution, large pills or tablets, or large numbers of pills or tablets in order to provide sufficient active compound for effective treatments. In addition, some heteroaryl-ketone fused azadecalin compounds, and formulations containing these compounds, may interact with capsule materials, or be incompatible with capsule materials. Such interactions, incompatibilities, and instabilities may lead to degradation of the heteroaryl-fused azadecalin compounds, reduce their bioavailability, or otherwise reduce their utility for therapeutic purposes. For these reasons, and others, pharmaceutical formulations for use in pills, tablets, solutions for injection, and other pharmaceutical formulations often may be unsuitable for therapeutic use.

[0006] Accordingly, improved formulations including heteroaryl-ketone fused azadecalin compounds having improved stability and/or bioavailability are needed.

SUMMARY

[0007] Applicant discloses herein formulations comprising the heteroaryl-ketone fused azadecalin (HKFA) compound (R)-(1-(4-fluorophenyl)-6-((1-methyl-1H-pyrazol-4-yl)sulfonyl)-4,4a,5,6,7,8-hexahydro-1H-pyrazolo[3,4-g]isoquinolin-4a-yl)(4-(trifluoromethyl)pyridin-2-yl)methanone (Example 18 of U.S. 8,859,774), also known as “relacorilant” and as “CORT125134”, which has the following structure:



[0008] In embodiments, the formulations disclosed herein are suitable for pharmaceutical use, and have improved stability and/or bioavailability as compared to prior or alternative formulations. In embodiments, the formulations may include relacorilant and a pharmaceutically acceptable excipient. In embodiments, the formulations contain relacorilant and a pharmaceutically acceptable excipient and are suitable for use in pharmaceutical compositions for oral administration of relacorilant to human patients for treating a disease or disorder, or to animals for veterinary therapeutic purposes.

[0009] In embodiments, the pharmaceutical formulation is suitable for the administration of an effective amount of relacorilant, e.g., in a unit dose formulation containing between about 1 and about 1000 milligrams (mg) of relacorilant, or between about 1 and about 500 mg of relacorilant. In embodiments, unit dose formulations contain 10 mg, or 20 mg, or 25 mg, or 50 mg, or 100 mg, or 200 mg, or 300 mg, or 400 mg, or 500 mg relacorilant. In embodiments, the pharmaceutical formulation is suitable for the administration of an effective amount of relacorilant, e.g., a daily dose of relacorilant of between about 1 and 100 mg/kg/day, or a daily dose of relacorilant of between about 1 and 20 mg/kg/day.

[0010] In embodiments, the pharmaceutical formulation comprising relacorilant is suitable for administration when encapsulated in a gelatin capsule. In embodiments, the pharmaceutical formulation comprising relacorilant is suitable for administration when encapsulated in a soft gelatin ("softgel") capsule.

[0011] The formulation comprising relacorilant may contain, for example, about 5 to about 35% relacorilant, about 45% to about 75% surfactant, about 5 to about 35% solubilizer, and may include an antioxidant. In embodiments, the formulation comprising relacorilant may contain, for example, about 10 to about 30% relacorilant, about 50% to about 70% surfactant, about 10 to

about 30% solubilizer, and may include an antioxidant. In embodiments, the formulation comprising relacorilant may contain, for example, about 12 to about 28% relacorilant, about 52% to about 68% surfactant, about 12 to about 28% solubilizer, and may include an antioxidant. In further embodiments the formulation may have the following composition: about 15% to 25% relacorilant, about 55% to 65% surfactant, and about 15% to 25% solubilizer, optionally including an antioxidant. In yet further embodiments, the formulation may have the following composition: about 20% relacorilant, about 60% surfactant, about 20% solubilizer, and about 0.02% antioxidant. In particular embodiments, the formulation may have the following composition: about 20% relacorilant, about 60% polyoxyl glyceride, about 20% propylene glycol compound, and about 0.02% antioxidant. In further particular embodiments, the formulation may have the following composition: about 20% relacorilant, about 60% lauryl polyoxyl-32 glycerides, about 20% propylene glycol monocaprylate, and about 0.02% antioxidant (e.g., butylated hydroxytoluene (BHT), butylated hydroxyanisole (BHA), tert-butyl hydroquinone (TBHQ), a tocopherol, a gallate, or other antioxidant).

[0012] In embodiments, the capsule containing the relacorilant-containing formulation may be a softgel capsule. Softgel capsules for containing a relacorilant-containing formulation may be made of about 50% to about 75% gelatin and about 25% to about 50% of a plasticizer (e.g., a sorbitol/glycerin blend), with purified water and/or colorants included as needed; and may be made of about 58% to about 68% gelatin and about 30% to about 42% plasticizer, or may be made of about 60% to about 70% gelatin and about 30% to about 40% of a plasticizer, or about 63% to about 65% gelatin and about 35% to about 37% of a plasticizer, or about 64% gelatin and about 36% plasticizer, with purified water and/or colorants included as needed. In embodiments, a batch of material for making softgel capsules for containing a relacorilant-containing formulation may be made, for example, of 195.05 g gelatin and 111.46 g of a sorbitol/glycerin blend, with purified water and/or colorants included as needed.

[0013] In further exemplary embodiments, Applicant discloses softgel capsules containing relacorilant; the softgel capsules containing relacorilant may be any softgel capsule as disclosed herein containing any relacorilant formulation disclosed herein. In embodiments, the softgel capsules containing relacorilant formulations may be made of about 63% to about 65% gelatin and about 35% to about 37% of a plasticizer, or about 64% gelatin and about 36% plasticizer, with purified water and/or colorants included as needed. The plasticizer may be, for example, a

sorbitol/glycerin blend. The relacorilant held in the softgel capsule may be in any formulation described herein. In embodiments, the softgel capsule containing relacorilant contains a relacorilant formulation of about 15 to about 25% relacorilant, about 50% to about 70% surfactant, about 15 to about 25% solubilizer, and may include an antioxidant. In embodiments, the softgel capsule containing relacorilant contains a relacorilant formulation comprising about 20 relacorilant, about 60% surfactant, about 20% solubilizer, and may include an antioxidant.

[0014] In embodiments, the formulations disclosed herein are suitable for use in the treatment of disorders characterized by cortisol excess. In embodiments, the formulations disclosed herein are suitable for use in the treatment of Cushing's syndrome, including Cushing's Disease. In embodiments, the formulations disclosed herein are suitable for use in the treatment of other disorders. In embodiments of such treatment methods, pharmaceutical compositions comprising relacorilant may be administered alone, or may be administered in combination with other treatments for such diseases and disorders.

[0015] Many HKFA compounds are difficult to dissolve, difficult to formulate with pharmaceutically acceptable excipients, and are unstable (i.e., degrade by unacceptable amounts over time) in many formulations. Applicant discloses herein novel relacorilant-containing formulations and pharmaceutical compositions, with improved stability as compared to alternative formulations and compositions which are suitable for administration to human subjects. The novel formulations disclosed herein are believed to allow the safe and effective administration of relacorilant to patients suffering from diseases and disorders amenable to treatment by GRMs such as relacorilant. The formulations disclosed herein provide advantages including improved stability and compatibility with capsules suitable for oral administration, providing improved relacorilant formulations, and treatments using such improved relacorilant-containing formulations, as compared to alternative formulations and compositions.

BRIEF DESCRIPTION OF THE DRAWINGS

[0016] FIG. 1 shows the plasma concentration of relacorilant in healthy, fasted human subjects at times after ingestion of 150 mg relacorilant (provided in softgel capsules containing the formulation of Table III). The $C_{\max}$ was nearly 200 ng/mL at its peak about 3 hours after administration (open circles; eight subjects were tested with this composition; the subjects were fasted when they were orally administered the relacorilant formulation).


[0017] FIG. 2 illustrates an idealized side view of a container used for *in vitro* digestion testing of test formulations. The container holds three phases into which the active pharmaceutical ingredient may partition.

[0018] FIG. 3 shows the results of *in vitro* digestion testing per Fig. 2, indicating that up to about 15% of the formulation of Table I partitioned into the aqueous phase during the early phases of the *in vitro* testing, and about 10% partitioned into the aqueous phase at time zero of the testing.

DETAILED DESCRIPTION

[0019] Many fused azadecalin compounds are difficult to formulate, and many fused azadecalin compounds are unstable over time in formulations that are suitable for oral administration. Thus, development of stable and bioavailable formulations containing fused azadecalin compounds is an important challenge that it is necessary to overcome in order to provide improved fused azadecalin formulations suitable for human administration. Preferred embodiments entail the development of stable and bioavailable formulations containing fused azadecalin compounds suitable for oral administration to human patients.

[0020] Applicant has surprisingly developed multiple formulations which allow the incorporation of the heteroaryl-ketone fused azadecalin (HKFA) compound “relacorilant” into pharmaceutical compositions suitable for oral administration to patients. Relacorilant is (R)-(1-(4-fluorophenyl)-6-((1-methyl-1H-pyrazol-4-yl)sulfonyl)-4,4a,5,6,7,8-hexahydro-1H-pyrazolo[3,4-g]isoquinolin-4a-yl)(4-(trifluoromethyl)pyridin-2-yl)methanone, and has the following structure:



These relacorilant-containing formulations provide improved stability and improved compatibility with capsules as compared to other, prior or alternative, relacorilant-containing formulations.

[0021] Disclosed herein are novel formulations and novel pharmaceutical compositions containing relacorilant. In embodiments, the formulations may include relacorilant and a pharmaceutically acceptable excipient, such as, e.g., a solubilizer, a surfactant, an antioxidant, a filler, or one or more other excipient. In embodiments, solubilizers such as, e.g., glycerols including glycerol caprylates, polyethylene or polypropylene glycerols, (commercially available as, e.g., Capmul ECM (glycerol caprate)) and surfactants such as polyethoxylated castor or other oils, (commercially available as, e.g., Kolliphor EL), may provide superior pharmaceutical formulations as compared to prior or alternative formulations. In embodiments, such formulations comprising HKFA compounds may also include antioxidants, e.g., butylated hydroxytoluene (BHT), butylated hydroxyanisole (BHA), tert-butyl hydroquinone (TBHQ), a gallate such as propyl gallate, a tocopherol such as alpha tocopherol, or other antioxidant.

[0022] In embodiments, Applicant discloses herein gelatin (“softgel”) capsule formulations suitable for use in encapsulating relacorilant-containing formulations. Softgel capsules for containing a relacorilant-containing formulation may be made of about 50% to about 75% gelatin and about 25% to about 50% of a plasticizer, with purified water and/or colorants included as needed; and may be made of about 60% to about 70% gelatin and about 30% to about 40% of a plasticizer, or about 63% to about 66% gelatin and about 34% to about 37% of a plasticizer, with purified water and/or colorants included as needed. In embodiments, a batch of material for making softgel capsules for containing a relacorilant-containing formulation may be made, for example, of 195.05 g gelatin and 111.46 g of a sorbitol/glycerin blend plasticizer. In embodiments, the gelatin capsule may contain purified water as needed, and may further contain colorants as needed, if desired.

[0023] In embodiments, Applicant discloses herein relacorilant-containing formulations, including without limitation the following formulations (“%” is taken as weight per cent herein):

a formulation comprising about 10 to about 30% relacorilant, about 50% to about 70% surfactant, about 10 to about 30% solubilizer, and optionally including an antioxidant;

a formulation comprising about 15 to about 25% relacorilant, about 55% to about 65% surfactant, about 15 to about 25% solubilizer, and optionally including an antioxidant;

a formulation comprising about 18% to 22% relacorilant, about 56% to 64% surfactant, and about 18% to 22% solubilizer, optionally including an antioxidant;

a formulation comprising about 20% relacorilant, about 60% surfactant, about 20% solubilizer, and about 0.1 to about 0.5% antioxidant;

a formulation comprising about 20% relacorilant, about 60% polyoxyl glyceride compounds, about 20% propylene glycol compounds, and about 0.02% BHT or BHA; and

a formulation comprising about 20% relacorilant, about 60% Lauroyl polyoxyl-32 glycerides, about 20% propylene glycol monocaprylate, and about 0.02% BHT or BHA.

[0024] In embodiments, Applicant discloses herein relacorilant-containing formulations, including without limitation the following formulations, provided below in the amounts contained in a single capsule (e.g., a softgel capsule):

A formulation containing 25 mg relacorilant, 74.975 mg Lauroyl poloxyl-32 glycerides, 25 mg Propylene glycol monocaprylate, type I, and 0.025 mg Butylated hydroxytoluene (BHT);

A formulation containing 50 mg relacorilant, 149.95 mg Lauroyl poloxyl-32 glycerides, 50 mg Propylene glycol monocaprylate, type I, and 0.05 mg Butylated hydroxytoluene (BHT);

A formulation containing 100 mg relacorilant, 299.9 mg Lauroyl poloxyl-32 glycerides, 100 mg Propylene glycol monocaprylate, type I, and 0.1 mg Butylated hydroxytoluene (BHT);

A formulation containing 200 mg relacorilant, 599.8 mg Lauroyl poloxyl-32 glycerides, 200 mg Propylene glycol monocaprylate, type I, and 0.2 mg Butylated hydroxytoluene (BHT);

A formulation containing 300 mg relacorilant, 899.7 mg Lauroyl poloxyl-32 glycerides, 300 mg Propylene glycol monocaprylate, type I, and 0.3 mg Butylated hydroxytoluene (BHT);

A formulation containing 400 mg relacorilant, 1199.6 mg Lauroyl poloxyl-32 glycerides, 400 mg Propylene glycol monocaprylate, type I, and 0.4 mg Butylated hydroxytoluene (BHT);

A formulation containing 500 mg relacorilant, 1499.5 mg Lauroyl poloxyl-32 glycerides, 500 mg Propylene glycol monocaprylate, type I, and 0.5 mg Butylated hydroxytoluene (BHT);

A formulation containing 750 mg relacorilant, 2249.25 mg Lauroyl poloxyl-32 glycerides, 750 mg Propylene glycol monocaprylate, type I, and 0.75 mg Butylated hydroxytoluene (BHT);

A formulation containing 1000 mg relacorilant, 2999 mg Lauroyl poloxyl-32 glycerides, 1000 mg Propylene glycol monocaprylate, type I, and 1 mg Butylated hydroxytoluene (BHT); and other formulations with the same proportions. In order to determine the amounts of a further formulation with the same proportions having a different amount of relacorilant *X*, one must determine the ratio of *X* to 100 mg relacorilant, and multiply the ingredient amounts of the formulation above containing 100 mg relacorilant by that ratio. For example, where the further formulation includes 800 mg relacorilant, the ratio of 800 mg relacorilant:100 mg relacorilant is 8, and the amounts for each ingredient of the formulation including 100 mg relacorilant would be multiplied by 8.

[0025] In embodiments, Applicant discloses herein single unit dosage forms containing relacorilant, including without limitation the following single unit dosage forms:

a single unit dosage form consisting of a capsule containing a uniform admixture of between about 10 milligrams (mg) and about 70 mg of relacorilant and pharmaceutically acceptable excipients, wherein said uniform admixture is a formulation as disclosed herein, including, e.g., a formulation disclosed above; and

a single unit dosage form consisting of a capsule containing a uniform admixture of between about 20 milligrams (mg) and about 50 mg of relacorilant and pharmaceutically acceptable excipients, wherein said uniform admixture is a formulation as disclosed herein, including, e.g., a formulation disclosed above.

a single unit dosage form consisting of a capsule containing a uniform admixture of between about 50 milligrams (mg) and about 200 mg of relacorilant and pharmaceutically acceptable excipients, wherein said uniform admixture is a formulation as disclosed herein, including, e.g., a formulation disclosed above.

a single unit dosage form consisting of a capsule containing a uniform admixture of between about 100 milligrams (mg) and about 400 mg of relacorilant and pharmaceutically acceptable excipients, wherein said uniform admixture is a formulation as disclosed herein, including, e.g., a formulation disclosed above.

a single unit dosage form consisting of a capsule containing a uniform admixture of between about 200 milligrams (mg) and about 500 mg of relacorilant and pharmaceutically acceptable excipients, wherein said uniform admixture is a formulation as disclosed herein, including, e.g., a formulation disclosed above.

In embodiments, the contents of a single unit dosage form as disclosed herein, including as disclosed above, weighs between about 300 milligrams (mg) and about 600 mg. In embodiments, the contents of a single unit dosage form as disclosed herein, including as disclosed above, has a total weight of about 300 mg; or about 400 mg; or about 500 mg; or about 600 mg; or about 700 mg; or about 800 mg; or about 900 mg; or about 1 gram (g); or about 1.5 g; or about 2 g. In embodiments, the single unit dosage form is enclosed in a softgel capsule. In yet other embodiments, the single unit dosage form is enclosed in a hardshell capsule, which may have a size selected from, e.g., size 5, size 4, size 3, size 2, size 1, size 0, size 00, and size 000.

[0026] Relacorilant binds to the glucocorticoid receptor (GR), and is a GR modulator (GRM). Relacorilant GR modulation comprises antagonizing binding of other ligands, such as, e.g., cortisol, to a GR, and thus relacorilant is a GR antagonist (GRA). Since relacorilant binds well to GR, but binds only poorly to the progesterone receptor (PR) and to other steroid hormone receptors, relacorilant may be termed a selective GR modulator (SGRM). The novel pharmaceutical compositions disclosed herein are useful, for example, by providing pharmaceutical compositions for modulating a GR while not significantly affecting a PR.

[0027] In embodiments, the relacorilant-containing formulations may, as one of their effects, reduce harmful effects of excessive cortisol levels in subjects to which they have been administered. In embodiments, the relacorilant-containing formulations may, as one of their effects, be effective to treat disorders in subjects to which they have been administered. In embodiments, the relacorilant-containing formulations may, as one of their effects, be effective to enhance the action of co-administered drugs or therapies in the treatment of disorders in subjects to which they have been administered, where co-administered drugs or therapies may be administered close in time to, at a time or times prior to, at the same time as, or following administration of the relacorilant-containing formulations.

[0028] In embodiments, Applicant discloses herein methods of treating conditions amenable to treatment with relacorilant, including without limitation the following methods:

a method of treating a condition amenable to treatment by relacorilant comprising administering relacorilant in a single unit dosage form, wherein said single unit dosage form contains a uniform admixture of said relacorilant and pharmaceutical excipients, and wherein said uniform admixture is a formulation as disclosed herein. In embodiments of the methods disclosed herein, the single unit dosage form consists of a capsule containing between about 10 milligrams (mg) and about 500 mg of relacorilant.

[0029] In embodiments of the methods disclosed herein, the single unit dosage form consists of a capsule containing a uniform admixture of between about 5 milligrams (mg) and about 500 mg of relacorilant. In embodiments of the methods disclosed herein, the single unit dosage form consists of a capsule containing a uniform admixture of between about 10 milligrams (mg) and about 250 mg of relacorilant. In embodiments of the methods disclosed herein, the single unit dosage form consists of a capsule containing a uniform admixture of between about 20 milligrams (mg) and about 150 mg of relacorilant.

[0030] In embodiments, the pharmaceutical composition comprising relacorilant is formulated for oral administration. In embodiments, the oral administration comprises sublingual administration. In embodiments the HKFA compound is formulated for administration as a suppository, and may be administered rectally or intravaginally.

B. DEFINITIONS

[0031] The abbreviations used herein have their conventional meaning within the chemical and biological arts.

[0032] As used herein, the term "compound" is used to denote a molecular moiety of unique, identifiable chemical structure. A compound may exist in a free species form, in which it is not associated with other molecules. A compound may also exist as part of a larger aggregate, in which it is associated with other molecule(s), but nevertheless retains its chemical identity. A solvate, in which the compound is associated with a molecule(s) of a solvent, is an example of such an associated form. A hydrate is a solvate in which the associated solvent is water. The recitation of a "compound" refers to the molecular moiety itself (of the recited structure), regardless of whether it exists in a free form or an associated form.

[0033] “Patient”, “subject”, “subject in need” and the like refer to a person having, or suspected of having, a disease or condition which may be treated by administration of a therapeutic drug.

[0034] As used herein, the terms “administer,” “administering,” “administered” or “administration” refer to providing a compound or a composition (*e.g.*, one described herein), to a subject or patient.

[0035] As used herein, the term “administration” refers to the delivery of a drug or other therapeutic into the body of a subject in need of treatment by the drug or therapeutic, effective to achieve a therapeutic effect. Administration may be by any suitable route of administration, including, for example, oral administration; intravenous administration; subcutaneous administration; parenteral administration; intra-arterial administration; nasal administration; topical administration; and other routes of administration.

[0036] “Treat”, “treating” and “treatment” refers to providing a drug or other therapeutic agent to a patient. Such treatments are intended to, and typically result in reduction of symptoms, or amelioration of symptoms, or abatement of symptoms, or abolition of symptoms, of the disease or disorder to be treated. Such treatments are intended to, and typically result in, an indication of success in the treatment or amelioration of a pathology or condition. Indicia of success include, *e.g.*, any objective or subjective parameter such as abatement; remission; diminishing of symptoms or making the pathology or condition more tolerable to the patient; slowing in the rate of degeneration or decline; making the final point of degeneration less debilitating; or improving a patient's physical or mental well-being. The treatment or amelioration of symptoms can be based on objective or subjective parameters; including the results of a physical examination; histopathological examination (*e.g.*, analysis of biopsied tissue); laboratory analysis of urine, saliva, tissue samples, serum, plasma, or blood; or imaging.

[0037] As used herein, the term “Cushing’s syndrome” refers to an array of symptoms caused by excess cortisol (excess cortisol is termed “hypercortisolism”). “Cushing’s Disease” refers to pituitary-dependent Cushing’s syndrome, *e.g.*, excess cortisol caused by pituitary abnormality (typically a pituitary tumor). Symptoms associated with, and indicative of, Cushing’s syndrome include, for example, elevated blood pressure (hypertension), elevated blood glucose (hyperglycemia), increased weight (typically in the mid-section, and in the face causing a

characteristic “moon-face”), immune suppression, thin skin, acne, depression, hirsutism, and other symptoms. The excess cortisol that leads to Cushing’s syndrome may be due to exogenous cortisol (or cortisol analog such as prednisone, cortisone, dexamethasone, or other cortisol analog or cortisol mimetic); may be due to a pituitary tumor; may be due to an adrenal tumor or other adrenal disease; may be due to a tumor not located in or near to the pituitary or an adrenal gland; or may be due to another cause. In some cases, a Cushing’s syndrome patient may refuse surgery, or surgery may have failed to cure the disease or relieve the symptoms; a GRM, such as relacorilant, is often useful in treating such patients.

[0038] A condition associated with Cushing’s syndrome may be, without limitation, a condition associated with endogenous Cushing’s syndrome; hyperglycemia secondary to hypercortisolism; hypercortisolism associated with, or causing, type 2 diabetes mellitus or glucose intolerance; hyperglycemia secondary to hypercortisolism associated with, or caused by, type 2 diabetes mellitus or glucose intolerance.

[0039] The term “glucocorticosteroid” or “glucocorticoid” (both abbreviated GC) refers to a steroid hormone or synthetic analog that binds to a glucocorticoid receptor. Glucocorticosteroids are typically characterized by having 21 carbon atoms, an α,β -unsaturated ketone in ring A, and an α -ketol group attached to ring D. They differ in the extent of oxygenation or hydroxylation at C-11, C-17, and C-19; see Rawn, “Biosynthesis and Transport of Membrane Lipids and Formation of Cholesterol Derivatives,” in *Biochemistry*, Daisy *et al.* (eds.), 1989, pg. 567.

[0040] As used herein, the term “glucocorticoid receptor” (“GR”) refers to a family of intracellular receptors which specifically bind to cortisol and/or cortisol analogs. The glucocorticoid receptor is also referred to as the cortisol receptor. The term includes isoforms of GR, recombinant GR and mutated GR. “Glucocorticoid receptor” (“GR”) refers to the type II GR which specifically binds to cortisol and/or cortisol analogs such as dexamethasone (*See, e.g.*, Turner & Muller, *J. Mol. Endocrinol.* October 1, 2005 35 283-292).

[0041] As used herein, the terms “effective amount,” “amounts effective,” “therapeutic amount,” and “therapeutically effective amount” refer to an amount or amounts of one or more pharmacological agents effective to treat, eliminate, or mitigate at least one symptom of the disease being treated. In some cases, “effective amount,” “amounts effective,” “therapeutic amount,” and “therapeutically effective amount” can refer to an amount of a functional agent or

of a pharmaceutical composition useful for exhibiting a detectable therapeutic or inhibitory effect. The effect can be detected by any assay method known in the art.

[0042] As used herein, the term “consisting essentially of”, as related to a list of ingredients in a formulation, refers to the recited ingredients, where the formulation may also include any unlisted ingredients that would not materially affect the utility of the formulation.

[0043] As used herein, the terms “excipient”, “pharmaceutically acceptable excipient”, and “pharmaceutically acceptable carrier” are used interchangeably to refer to any compound or material which is included in a formulation or in a pharmaceutical composition other than the active ingredient (or active ingredients where more than one compound may have the desired pharmaceutical activity). An excipient may serve as, e.g., a solvent, solubilizer or solubility enhancer; an emulsifier; a bulking agent; a stabilizer; a diluent; a surfactant; a preservative; a colorant; a flavor; a filler; a lubricant; or other agent which may serve other functions. Non-limiting examples of excipients include water, sodium chloride (NaCl), normal saline solutions, lactated Ringer's, normal sucrose, normal glucose, binders, fillers, disintegrants, lubricants, coatings, sweeteners, flavors and colors, and the like. As used herein, these terms are intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. One of skill in the art will recognize that other pharmaceutical excipients are useful in the present invention, and their use is well known in the art. Any such pharmaceutically-acceptable excipients may be used in the formulations disclosed herein is contemplated except insofar as an excipient is incompatible with the active compound. It will be understood that supplementary active compounds can also be incorporated into the compositions. It is understood that the pharmaceutically acceptable excipients are non-toxic. Additional information on suitable pharmaceutically acceptable excipients can be found in Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, Pa., 1985, which is incorporated herein by reference.

[0044] As used herein, “active pharmaceutical ingredient”, “API”, and variants thereof refer to the compound in a formulation or pharmaceutical composition that has therapeutic activity when administered to a patient. Although other ingredients may also have beneficial effects, the formulation or pharmaceutical composition is configured to, and has been developed for,

delivery of the API in order to provide the therapeutic benefit of the API to the patient to whom it is administered, while any beneficial effects of other ingredients is incidental.

[0045] As used herein, the terms “formulation”, “pharmaceutical formulation”, “composition”, “pharmaceutical composition”, and the like are interchangeable and refer to compositions suitable for administration to a patient for treatment of a medical condition or for amelioration of symptoms of a medical condition. A pharmaceutical formulation as disclosed herein comprises an active ingredient (e.g., relacorilant), and an inert ingredient (e.g., a pharmaceutically acceptable excipient). In embodiments, a pharmaceutical formulation includes one or more active ingredients and one or more pharmaceutically acceptable excipients. A pharmaceutical formulation is suitable for administration to a patient; in embodiments, a pharmaceutical formulation is suitable for oral administration to a patient.

[0046] As used herein, the terms “per cent” and “%” refer to a percentage taken by comparing a first value to a second value, and multiplying the resulting decimal fraction by 100. As used herein, the first value may be the weight of an ingredient in a formulation containing multiple ingredients, and the second value may be the weight of all ingredients (i.e., the total weight of the formulation). Thus, for example, where the weight of a relacorilant dose in a pharmaceutical formulation containing that relacorilant is 100 milligrams (mg), and the total weight of all the ingredients in the formulation is 400 mg, then relacorilant makes up 25% of the formulation.

[0047] As used herein, the terms “weight percent”, “weight %”, “(%w/w)” and the like refer to the percentage of an ingredient of a composition with respect to the total weight of a composition containing at least two ingredients.

[0048] As used herein, the phrase “between about x% and about y%” (where x and y may be any number) is used inclusively, so that the range include the number values stated and all values between those minimum and maximum values.

[0049] As used herein, the phrases “unit dosage form” and “single unit dosage form” refer to formulations contained in pills, capsules, and other enclosures that are suitable for administration to a patient. The dosage (i.e., the amount of API) in a unit dosage form is fixed, and allows standardization of administration, without requiring measurement of the API or formulation by a patient or healthcare practitioner. Doses administered to a patient may be increased by


administration of multiple pills, capsules, etc. In some cases, fractional doses may be administered, e.g., by cutting a pill in two or more pieces.

[0050] In embodiments, relacorilant-containing formulations disclosed herein may be suitable for oral administration.

[0051] Pharmaceutical formulations may be contained in capsules for administration of pharmaceutical ingredients, such as relacorilant, to patients for therapeutic purposes. In embodiments, such capsules are suitable for administration pharmaceutical ingredients, such as relacorilant, in unit doses to patients.

FORMULATIONS

[0052] Heteroaryl-ketone fused azadecalin (HKFA) compounds may be useful in formulations as disclosed herein, including in formulations suitable for oral administration (see, e.g., U.S. Patent 8,859,774, the entire contents of which is hereby incorporated by reference in its entirety). Relacorilant is an exemplary HKFA compound, and formulations which provide relacorilant in a form suitable for oral administration to a patient are useful. Relacorilant is (R)-(1-(4-fluorophenyl)-6-((1-methyl-1H-pyrazol-4-yl)sulfonyl)-4,4a,5,6,7,8-hexahydro-1H-pyrazolo[3,4-g]isoquinolin-4a-yl)(4-(trifluoromethyl)pyridin-2-yl)methanone (Example 18 of U.S. 8,859,774), also known as “CORT125134”, and has the following structure:



[0053] Formulations suitable for administration to patients, and, in embodiments, suitable for oral administration to patients, may include relacorilant in proportions of between about 1% and about 50% by weight; in embodiments, such formulations may include relacorilant in proportions of between about 5% and about 35% by weight; in embodiments, such formulations may include relacorilant in proportions of between about 10% and about 30% by weight; in

embodiments, such formulations may include relacorilant in proportions of between about 15% and about 25% by weight; in further embodiments, such formulations may include relacorilant in proportions of between about 17% and about 23% by weight; and in further embodiments, such formulations may include relacorilant in proportions of between about 19% and about 21% by weight; and in embodiments, such formulations contain about 20% relacorilant by weight.

[0054] Accordingly, Applicant discloses herein compositions comprising relacorilant may be useful in treating patients suffering from a condition amenable to treatment with relacorilant, and may be used in the treatment of a patient in need of such treatment. Conditions amenable to treatment with relacorilant may include without limitation, any of the conditions, disorders, and diseases disclosed herein (e.g., see *supra*), including for example, Cushing's syndrome, Cushing's Disease, diabetes, hyperglycemia and other disorders.

[0055] For preparing pharmaceutical compositions from the compounds of the present invention, pharmaceutically acceptable excipients can be either solid or liquid. Preparations may include formulations incorporated into, or contained by, tablets, pills, capsules, cachets, or suppositories. An excipient can be one or more substances, which may also act as diluents, flavoring agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material, as disclosed herein (e.g., *supra*). Details on techniques for formulation and administration are well described in the scientific and patent literature, see, e.g., the latest edition of Remington's Pharmaceutical Sciences, Mack Publishing Co, Easton Pa. ("Remington's").

[0056] Suitable solid excipients include, but are not limited to, magnesium carbonate; magnesium stearate; talc; pectin; dextrin; starch; tragacanth; a low melting wax; cocoa butter; carbohydrates; sugars including, but not limited to, lactose, sucrose, mannitol, or sorbitol, starch from corn, wheat, rice, potato, or other plants; cellulose such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; and gums including arabic and tragacanth; as well as proteins including, but not limited to, gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

[0057] Formulations as disclosed herein may comprise excipients and other components known in the art to be suitable for use in pharmaceutical compositions for administration to humans. Such excipients and components include, without limitation, surfactants, solubilizers,

antioxidants, fillers, and other components. For example, suitable commercially available excipients and components include, without limitation, Capmul ECM (glycerol caprate); Kolliphor EL (polyethoxylated castor oil); Kolliphor EM; Gelucire 44/14 (lauroyl polyoxyl-32 glycerides); Capryol PGMC (propylene glycol monocaprylates); polysorbate 20; polysorbate 80; sorbitol/glycerin blend (e.g., the “50/50 Blend Sorbitol Special™” commercially available from SPI Pharma, Septemes, FR); corn oil; lauroglycol 90; butylated hydroxytoluene (BHT); butylated hydroxyanisole (BHA); tert-butyl hydroquinone (TBHQ); propyl gallate; alpha-tocopherol (vitamin E); myristic acid; and others.

Surfactants

[0058] As used herein, the term “surfactant” refers to, without limitation, polysorbate (polyoxyethylene sorbitan monolaureate) surfactants and detergents (such as, e.g., Tween® 20, Tween® 80, and others); may include polyoxyl glycerides, e.g., lauryl polyoxyl glycerides and other polyoxyl glycerides, and may include ionic, and non-ionic surfactants.

[0059] Formulations suitable for administration, including oral administration, to a patient and containing relacorilant may include a surfactant, including a non-ionic surfactant, in proportions of between about 10% and about 90% by weight; in embodiments, such formulations and compositions containing relacorilant may include surfactant, including a non-ionic surfactant, in proportions of between about 15% and about 80% by weight; in embodiments, such formulations and compositions containing relacorilant may include surfactant, including a non-ionic surfactant, in proportions of between about 20% and about 75% by weight; and in embodiments, such formulations and compositions containing relacorilant may include surfactant, including a non-ionic surfactant, in proportions of between about 25% and about 70% by weight. In embodiments, such formulations and compositions containing relacorilant may not include a non-ionic surfactant. In embodiments, such formulations and compositions containing relacorilant may not include a surfactant.

[0060] Surfactants, including non-ionic surfactants, are used as excipients. Surfactant excipients suitable for use in formulations as disclosed herein may include non-ionic surfactants, including sorbitan surfactants such as sorbitan monooleate; polysorbates (of many sizes, including e.g., polysorbate 20, polysorbate 80, also known as Tween® 20 and Tween® 80 as discussed above); and other non-ionic surfactants including, e.g., those sold as Cremophor® or Kolliphor® (e.g.,

Cremophor[®] EL, Kolliphor[®] EL, Cremophor[®] RH 40, Cremophor[®] RH 60, etc.), *d*- α -tocopherol, Solutol HS 15, poloxamer 407, Labrafil[®] M-1944CS (Gattefosse), Labrafil M-2125CS (Gattefosse), Labrasol[®] (Gattefosse, Saint-Priest, Lyon, France), Softigen[®] 767), and other surfactants known in the art.

[0061] For example, polysorbate (polyoxyethylene sorbitan monolaureate) surfactants and detergents (such as, e.g., Tween[®] 20, Tween[®] 80, and others, where numbers such as “20” and “80” indicate the numbers of repeating polyethylene glycol units in the polymers) are used as excipients in formulating pharmaceuticals, and aid in the emulsification and/or solubilization of active compounds in addition to their actions as surfactants. Such nonionic detergents and surfactants are available, e.g., from SIGMA-Aldrich, St. Louis, MO, USA.

[0062] In embodiments, relacorilant-containing formulations as disclosed herein may include Gelucire (e.g., Gelucire[®] 44/14, available from Gattefosse, Saint-Priest, Lyon, France) as a surfactant.

[0063] In embodiments, relacorilant-containing formulations as disclosed herein may include a “Kolliphor” as a surfactant. Kolliphor[®] EL (also known as cremophor; see above) is a polyethoxylated castor oil used as a non-ionic oil-in-water emulsifier. Kolliphor P188 is a poloxamer. Kolliphor[®] HS 15 (polyethylene glycol (15)-hydroxystearate is a solubilizer/emulsifier. Kolliphor[®] RH40 is Macrogolglycerol hydroxystearate. These materials are available from SIGMA-Aldrich, St. Louis, MO, USA.

Solubilizers

[0064] Formulations suitable for administration, including oral administration, to a patient and containing relacorilant may include a solvent, solubilizer, or solubility enhancer, or a plurality of solvents, solubilizers, or solubility enhancers in proportions of between about 2% and about 90% by weight; in embodiments, such formulations and compositions containing relacorilant may include a solvent, solubilizer, or solubility enhancer, or a plurality thereof, in proportions of between about 3% and about 80% by weight; or, in embodiments, in proportions of between about 5% and about 70% by weight; or in embodiments, in proportions of between about 8% and about 60% by weight; or, in further embodiments, in proportions of between about 10% and about 50% by weight. In embodiments, such formulations and compositions containing relacorilant may not include a solvent.

[0065] As used herein, the term “solubilizer” refers, without limitation, to solubilizers such as, e.g., polyethylene glycols (PEG), ethanol, propylene glycols, glycerols, glycerin, *N*-methyl-2-pyrrolidone, dimethylacetamide, dimethylsulfoxide, other water-soluble organic solvents, and other solubilizers known in the art. In embodiments, a solubilizer may be, e.g., a propylene glycol or propylene glycol ester, such as, e.g., a propylene glycol caprylate (e.g., a propylene glycol monocaprylate).

[0066] Polyethylene glycols of many forms, and derivatives thereof, are useful as solubilizers, and are useful in other ways, in the formulations disclosed herein. For example, PEG may have mean molecular weight of 300 kD or 400 kD (e.g., polyethylene glycol 300, polyethylene glycol 400); may be used in succinate form as polyethylene glycol (PEG) succinates (e.g., polyethylene glycol 1000 succinate) and in the form of mono- and di-fatty acid esters of PEG (e.g., PEG 300, 400, and 1750 mono- and di-fatty acid esters); and in other forms. Polyethylene glycols of a wide range of molecular weights (e.g., PEG300 which has an average molecular weight of about 300 grams per mole, and PEG400 which has an average molecular weight of about 400 grams per mole) are available, e.g., from SIGMA-Aldrich, St. Louis, MO, USA.

[0067] Solvents, solubilizers, and solubility enhancers used as excipients in formulations as disclosed herein may also include ethanol, glycerin, *N*-methyl-2-pyrrolidone, dimethylacetamide, dimethylsulfoxide, and other water-soluble organic solvents. In embodiments, formulations as disclosed herein containing a HKFA may include diethylene glycol monoethyl ether (sold under the names Transcutol[®], Carbitol[®], dioxitol, and other names); Transcutol[®], e.g., Transcutol[®] HP, is available from Gattefosse, Saint-Priest, Lyon, France.

Further Excipients

[0068] Excipients used in formulations as disclosed herein may include lipids and phospholipids, including naturally occurring lipid compositions such as, e.g., castor oil, corn oil, cottonseed oil, olive oil, peanut oil, peppermint oil, safflower oil, sesame oil, soybean oil, hydrogenated vegetable oils, hydrogenated soybean oil, and medium-chain triglycerides of coconut oil and palm seed oil). Excipients used in formulations as disclosed herein may also include phospholipids such as, e.g., phosphatidylcholine, distearoylphosphatidylglycerol L- α -dimyristoylphosphatidylcholine, L- α -dimyristoylphosphatidylglycerol, and other phospholipids.

Lipids and phospholipids may be used, e.g., as emulsifiers, bulking agents, fillers, lubricants, and for other purposes.

[0069] In embodiments, formulations as disclosed herein containing a GRM may include triglycerides, including medium chain triglycerides. Medium chain triglycerides are triglycerides with fatty acid chain lengths of between about 6 to about 12 carbons long. Triglycerides may include, e.g., 1,2,3-triacetoxyp propane (also known as triacetin or as glycerin triacetate). Triacetin is available, e.g., from SIGMA-Aldrich, St. Louis, MO, USA. Triglyceride excipients such as, e.g., triacetin, may serve, e.g., as solvents, as plasticizers, and as humectants.

[0070] Excipients used in formulations as disclosed herein may include various cyclodextrins such as, e.g., α -cyclodextrin, β -cyclodextrin, hydroxypropyl- β -cyclodextrin, sulfobutylether- β -cyclodextrin, and others. Cyclodextrins may be used, e.g., as emulsifiers, solubilizers, and for other purposes.

Formulations Containing Relacorilant

[0071] In embodiments, relacorilant-containing formulations as disclosed herein may include Propylene Glycol Monocaprylate (sold, e.g., as Capryol™ 90 by Gattefosse, Saint-Priest, Lyon, France). In embodiments, relacorilant-containing formulations as disclosed herein may include Caprylocaproyl polyoxylglycerides such as Labrasol® (sold by Gattefosse, Saint-Priest, Lyon, France; see above).

[0072] In embodiments, formulations suitable for administration, including oral administration, to a patient and containing relacorilant may include a polyethylene glycol (PEG) in proportions of between about 5% and about 50% by weight; in embodiments, such formulations and compositions containing a GRM may include a PEG in proportions of between about 10% and about 40% by weight; and in embodiments, such formulations and compositions containing relacorilant may include a PEG in proportions of between about 15% and about 30% by weight.

[0073] In embodiments, formulations suitable for administration, including oral administration, to a patient and containing relacorilant may include a polysorbate (such as, e.g., Tween® 20 or Tween® 80) in proportions of between about 5% and about 75% by weight; and in embodiments, such formulations and compositions containing relacorilant may include a polysorbate in

proportions of between about 20% and about 60% by weight. In embodiments, such formulations and compositions containing relacorilant may not include a polysorbate.

[0074] Excipients used in formulations as disclosed herein may include organic materials (which may be liquid or may be semi-solid at room temperature) such as, e.g., beeswax, a tocopherol (e.g., α -tocopherol (also termed vitamin E, or vitE)), oleic acid, gum Arabic, lanolin, starch, syrup, honey, and medium-chain mono- and diglycerides. Such organic materials may be used, e.g., as bulking agents, fillers, lubricants, and for other purposes.

[0075] In embodiments, relacorilant-containing formulations as disclosed herein may include Vitamin E (e.g., α -tocopherol and other tocopherols, including, e.g., α -tocopherol polyethylene glycol succinate (Vitamin E TPGS)). Vitamin E, including vitamin E TPGS, are available, e.g., from SIGMA-Aldrich, St. Louis, MO, USA. Vitamin E may serve, e.g., as a solubilizer; as an emulsifier; and as an antioxidant.

[0076] Further excipients and materials useful in formulations and in the manufacture of pharmaceuticals for administration, including pharmaceutical for oral administration, include sterile water for irrigation; ethanol; gelatin (e.g., edible gelatin, having a jelly strength measured in “bloom” units, such as Gelatin 220 Bloom); coatings such as hydroxypropyl methylcellulose (HPMC, “hypromellose”, e.g., Pharmacoat[®] 603 and Pharmacoat[®] 615) and polyvinyl alcohol coating materials; and other materials.

Exemplary Formulations

[0077] Many GRMs, including many HKFA GRMs, are difficult to formulate in pharmaceutically acceptable formulations. Problems include poor solubility in water; poor solubility in solvents that may be used in pharmaceutical formulations; poor stability; light sensitivity; precipitation in physiological environments; low bioavailability; incompatibility with pharmaceutically acceptable capsules and coatings; and other difficulties. Poor stability is a particular problem in developing formulations suitable for administration to patients in clinical practice.

[0078] Applicant discloses herein formulations which are believed to overcome such obstacles. Some embodiments of suitable formulations are presented in the Examples, and are also more generally described herein. For example, a formulation suitable for oral administration to humans

and containing relacorilant may contain, for example, about 15% to about 25% relacorilant, about 50% to about 70% surfactant, about 15% to about 25% solubilizer, and may include a small but effective amount of antioxidant (e.g., less than about 0.1%), where % indicates %w/w. In another example, a formulation suitable for oral administration to humans and containing relacorilant may contain, for example, about 20% relacorilant, about 60% surfactant, about 20% solubilizer, and may optionally include a small but effective amount of antioxidant (e.g., less than about 0.1%), where % indicates %w/w.

[0079] In embodiments of such formulations, the surfactant may be a glyceride, such as a polyoxyl glyceride, such as, e.g., lauroyl polyoxyl-32 glyceride. In embodiments, the surfactant may be a polysorbate, such as, e.g., TWEEN 20 or TWEEN 80. It will be understood that other suitable surfactants are known to those of skill in the art, and may be used with, or used in place of, a polysorbate. It will be understood that other suitable surfactants are known to those of skill in the art.

[0080] In embodiments of such formulations, the solubilizer may include glycerol, or polyethylene glycol, or polypropylene glycol. In embodiments of such formulations, the surfactant may be a propylene glycol, such as propylene glycol monocaprylate, e.g., propylene glycol monocaprylate Type I. In embodiments of such formulations, the surfactant may be a glycerol caprate. In further embodiments of such formulations and pharmaceutical compositions, without limitation, the solubilizer may be one or more of an oil, such as corn oil, castor oil, and other oils; may be one or more of a polyethylene glycol (PEG), ethanol, propylene glycol, glycerin, *N*-methyl-2-pyrrolidone, dimethylacetamide, dimethylsulfoxide; and may include combinations thereof. It will be understood that other suitable solubilizers are known to those of skill in the art, and may be used with, or used in place of, the solubilizers recited herein.

[0081] In embodiments, an antioxidant is included in formulations containing relacorilant suitable for administration to humans. The antioxidant may be, for example, BHT, BHA, TBHQ, alpha-tocopherol (vitamin E), a gallate (such as, e.g., propyl gallate), or other antioxidant, and may comprise combinations thereof.

[0082] For example, embodiments of formulations containing relacorilant suitable for administration to humans may include one or more of the following commercially available

ingredients: Capmul MCM; Kolliphor EL; Kolliphor ELP; Gelucire 44/14; Capryol PGMC; polysorbate 80; propyl gallate; BHT; BHA; alpha-tocopherol; myristic acid; and others.

[0083] In embodiments, a unit dosage of relacorilant is contained in a capsule, e.g., a softgel capsule, or a hardshell capsule. Formulations may be encapsulated in gelatin capsules; for example, in preferred embodiments, the capsules may be softgel capsules. Softgel capsules are typically prepared in a size suitable to accommodate their contents. Capsules are typically sized so as to accommodate from one or a few milligrams up to a several hundred milligrams of the active pharmaceutical ingredient (API), such as relacorilant. In embodiments, the amount of relacorilant contained in a softgel capsule may be about 10 mg, about 20 mg, about 30 mg, about 40 mg, about 50 mg, about 60 mg, about 70 mg, about 80 mg, about 100 mg, about 120 mg, about 140 mg, about 150 mg, about 170 mg, about 175 mg, about 180 mg, about 200 mg, about 225 mg, about 250 mg, about 300 mg, about 325 mg, about 350 mg, about 375 mg, about 400 mg, about 425 mg, about 450 mg, about 475 mg, about 500 mg, about 550 mg, about 600 mg, about 650 mg, about 700 mg, about 750 mg, about 800 mg, or more. Hardshell capsules may be chosen for a size that is suitable for holding the desired formulation, and may be, e.g., size 5 capsules, size 4 capsules; size 3 capsules; size 2 capsules; size 1 capsules; size 0 capsules; size 00 capsules; and other size capsules.

ADMINISTRATION

[0084] In embodiments, the formulation is suitable for the administration of an effective amount of relacorilant, e.g., in a unit dose formulation containing between about 1 and about 500 milligrams (mg) of relacorilant. In some embodiments, a unit dose relacorilant formulation is suitable for oral administration. In some embodiments, a unit dose relacorilant formulation may contain up to about 750 mg of relacorilant. In some embodiments, the unit dose of relacorilant is 1, 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450, 500, or 750 mg. In embodiments, the formulation comprising relacorilant is suitable for administration on a daily basis, or on a twice daily basis, or on a three times per day basis. In embodiments, the formulation comprising relacorilant is suitable for administration every other day, or every third day, or every fourth day, or every fifth day, or every sixth day, or on a once per week basis. In embodiments, the formulation comprising relacorilant is suitable for

administration for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, or 80 weeks.

[0085] In embodiments, the pharmaceutical formulation is suitable for the administration of an effective amount of relacorilant, e.g., a daily dose of relacorilant of between about 1 and 100 mg/kg/day, preferably between about 1 mg/kg/day and about 20 mg/kg/day. In some embodiments, the daily dose of relacorilant is 1, 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 30, 40, 50, 60, 70, 80, 90 or 100 mg/kg/day. In embodiments, the relacorilant is administered for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, or 80 weeks.

[0086] In embodiments, the pharmaceutical formulation comprising relacorilant is suitable for administration when encapsulated in a gelatin capsule. In embodiments, the pharmaceutical formulation comprising relacorilant is suitable for administration when encapsulated in a soft gelatin (“softgel”) capsule. In embodiments, such a softgel capsule is composed of about 50% to about 75% gelatin and about 25% to about 50% sorbitol/glycerin blend, and may also include purified water and/or colorants. In some embodiments, a unit dose relacorilant formulation is contained in a softgel capsule. An exemplary softgel capsule may be composed of the following ingredients:

Component	Composition (mg)
Gelatin	195.05
Purified Water	As needed for gel mass preparation
Sorbitol/Glycerin Blend	111.46
Colorants	As needed

Sorbitol/glycerin blend contains a 50:50 blend of sorbitol and glycerin.

[0087] In embodiments, the pharmaceutical formulation comprising relacorilant is suitable for administration with another pharmaceutical formulation, e.g., with a pill, tablet, oral solution, injectable, or other formulation including another active ingredient.

[0088] The formulations and methods disclosed herein may be useful in treating patients suffering from any condition amenable to treatment with a GRM, as discussed *supra*.

[0089] In embodiments, the pharmaceutical composition comprising relacorilant is formulated for oral administration. In embodiments, the oral administration comprises sublingual administration. In embodiments the relacorilant is formulated for administration as a suppository, and may be administered rectally or intravaginally.

[0090] Methods of treating diseases including without limitation, e.g., Cushing's syndrome, by administration of a GRM in such pharmaceutical compositions are also provided. In embodiments, a patient suffering from a disease or condition amenable to treatment by the GRM is administered a pharmaceutical composition disclosed herein; in embodiments, the administration comprises oral administration.

[0091] The relacorilant-containing pharmaceutical preparation is preferably in unit dosage form. In such form the preparation is subdivided into unit doses containing appropriate quantities of relacorilant. A unit dosage form may be, for example, a pill or capsule including or containing relacorilant.

[0092] In embodiments, a unit dosage form contains between about 0.1 milligram (mg) of relacorilant and about 750 mg of relacorilant; in further embodiments, a unit dosage form contains between about 1 mg of relacorilant and about 600 mg of relacorilant; in further embodiments, a unit dosage form contains between about 5 mg of relacorilant and about 500 mg of relacorilant; in further embodiments, a unit dosage form contains between about 10 mg of relacorilant and about 400 mg of relacorilant; in further embodiments, a unit dosage form contains between about 15 mg of relacorilant and about 350 mg of relacorilant; in further embodiments, a unit dosage form contains between about 20 mg of relacorilant and about 300 mg of relacorilant; in further embodiments, a unit dosage form contains between about 25 mg of relacorilant and about 250 mg of relacorilant.

[0093] In particular embodiments, the unit dosage form may contain between about 5 milligrams (mg) and about 200 mg relacorilant. In further embodiments, the unit dosage form may contain between about 10 mg and about 150 mg relacorilant. In yet further embodiments, the unit dosage form may contain between about 15 mg and about 100 mg relacorilant. In still further embodiments, the unit dosage form may contain between about 20 mg and about 80 mg relacorilant. In embodiments, the unit dosage form may contain between about 25 mg and about

75 mg relacorilant. In yet further embodiments, the unit dosage form may contain between about 30 mg and about 60 mg relacorilant.

[0094] In particular embodiments, the unit dosage form may contain about 10 mg of relacorilant; or about 25 mg of relacorilant; or about 40 mg of relacorilant; or about 50 mg of relacorilant; or about 75 mg of relacorilant; or about 100 mg of relacorilant; or about 125 mg of relacorilant; or about 150 mg of relacorilant; or about 200 mg of relacorilant; or about 250 mg of relacorilant; or about 300 mg of relacorilant; or about 350 mg of relacorilant; or about 400 mg of relacorilant; or about 450 mg of relacorilant; or about 500 mg of relacorilant; or about 550 mg of relacorilant; or about 600 mg of relacorilant; or about 650 mg of relacorilant; or about 700 mg of relacorilant; or about 750 mg of relacorilant; or about 800 mg of relacorilant.

[0095] The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

[0096] The single unit dosage form containing relacorilant may be administered once a day, or two, three, or more times per day so as to provide the preferred dosage level per day. Multiple units of the single unit dosage form containing relacorilant may be administered at the same time, or on the same day, so as to achieve the desired dosage (amount of relacorilant) per day.

[0097] The composition can also contain other compatible therapeutic agents. The compounds described herein can be used in combination with one another, or with adjunctive agents that may not be effective alone, but may contribute to the efficacy of the active agent.

USES

[0098] In embodiments, relacorilant is suitable for use in the treatment of disorders characterized by cortisol excess. For example, in embodiments, pharmaceutical formulations containing relacorilant are suitable for use in the treatment of Cushing's syndrome, including Cushing's Disease, and other diseases and disorders. Such treatments may be administered alone, or in combination with other treatments for such diseases and disorders.

[0099] The pharmaceutical formulations containing relacorilant and the methods disclosed herein comprising administration of pharmaceutical formulations containing relacorilant may be

useful in treating patients suffering from any condition amenable to treatment with a GRM. Conditions amenable to treatment with a GRM may include without limitation, for example, obesity, diabetes, hyperglycemia, cardiovascular disease, hypertension, Syndrome X, depression, anxiety, glaucoma, human immunodeficiency virus (HIV) or acquired immunodeficiency syndrome (AIDS), neurodegeneration, Alzheimer's disease, Parkinson's disease, cognition enhancement, Cushing's Syndrome, Addison's Disease, osteoporosis, frailty, muscle frailty, inflammatory diseases, osteoarthritis, rheumatoid arthritis, asthma and rhinitis, adrenal function-related ailments, viral infection, immunodeficiency, immunomodulation, autoimmune diseases, allergies, wound healing, compulsive behavior, multi-drug resistance, addiction, psychosis, anorexia, cachexia, post-traumatic stress syndrome, post-surgical bone fracture, medical catabolism, major psychotic depression, mild cognitive impairment, psychosis, dementia, stress disorders, antipsychotic induced weight gain, delirium, cognitive impairment in depressed patients, cognitive deterioration in individuals with Down's syndrome, psychosis associated with interferon-alpha therapy, chronic pain, pain associated with gastroesophageal reflux disease, postpartum psychosis, postpartum depression, neurological disorders in premature infants, and migraine headaches. In some embodiments, the disorder or condition can be major psychotic depression, a stress disorder or antipsychotic induced weight gain. In embodiments, the disorder or condition can be selected from obesity, diabetes, hyperglycemia, hypertension, depression, anxiety, and Cushing's Syndrome. In embodiments, the disorder or condition can be Cushing's Syndrome, and may be, in particular embodiments, Cushing's Disease, and may include hyperglycemia secondary to hypercortisolism.

[0100] Other conditions that can be treated using the formulations disclosed herein include alcohol dependence, symptoms of alcohol withdrawal, and cognitive deficits associated with excess alcohol consumption. The formulations disclosed herein can also be used to treat cancer, such as cancer of the bone, breast, prostate, ovary, skin, brain, bladder, cervix, liver, lung, and other cancers, including, without limitation, leukemia, lymphoma, neuroblastoma, among others.

EXAMPLES

[0101] The following examples are presented by way of illustration of embodiments of the methods disclosed herein, and serve to illustrate, but not to limit, the present disclosure of formulations containing relacorilant.

[0102] In the following examples, excipients included: glycerides, e.g., Lauroyl polyoxyl glycerides, such as Lauroyl polyoxyl-32 glycerides and diglycerides; including propylene glycols, such as propylene glycol monocaprylate; sorbitol; glycerin; and including blends of sorbitol and glycerin; and may have included colorants and anti-oxidants such as, e.g. butylated hydroxytoluene (BHT). Purified water was also used as needed.

[0103] In further embodiments, formulations may include surfactants, including without limitation polysorbate (polyoxyethylene sorbitan monolaureate) surfactants and detergents (such as, e.g., Tween[®] 20, Tween[®] 80, and others); lipids, such as, e.g., corn oil, castor oil, cottonseed oil, olive oil, peanut oil, and other oils, including hydrogenated vegetable oils, medium-chain triglycerides of coconut oil and palm seed oil, and other oils suitable for use in pharmaceutical compositions for human administration; solubilizers (including solvents and solubility enhancers), such as, e.g., polyethylene glycols (PEG), ethanol, propylene glycol, glycerin, *N*-methyl-2-pyrrolidone, dimethylacetamide, dimethylsulfoxide, and other water-soluble organic solvents without limitation; and antioxidants such as BHT, butylated hydroxyanisole (BHA), tert-butyl hydroquinone (TBHQ), vitamin E (alpha tocopherol), and others without limitation.

[0104] In the following examples, the gel capsules used were softgel capsules, and included gelatin, sorbitol, and glycerin. In addition, the softgel capsules typically further included purified water (as needed for gel preparation), and colorants if desired. An exemplary gel capsule shell included 195.05 g gelatin; and 111.46 g of a sorbitol-glycerin blend containing equal amounts of sorbitol and glycerin, with smaller amounts of D-glucitol. The sorbitol-glycerin blend is commercially available as "Sorbitol Special[™] from SPI Pharma (Septemes, FR).

[0105] As noted above, many GRMs, including many HKFA GRMs such as relacorilant, are poorly soluble in pharmaceutically acceptable compositions and solvents; in addition, pharmaceutical formulations of many such GRMs, including many such HKFA GRMs, often provide only poor bioavailability. In addition, many GRMs, including many HKFA GRMs, are difficult to formulate so as to provide acceptable stability. Many GRMs, such as relacorilant, are not stable over time in some formulations, particularly in basic (low pH) formulations, or when exposed to light, or under other storage conditions. Due to the requirements for solubilization, and the solvents needed for such solubilization, many GRMs, including many HKFA GRMs, are difficult to formulate so as to provide acceptable biocompatibility of the pharmaceutical

formulation (e.g., to provide formulations which do not include, or only include minimal amounts of, excipients which may have uncomfortable, adverse, or toxic effects on the subject to which they are administered).

[0106] The following examples discuss several exemplary relacorilant-containing formulations prepared as discussed herein. All of the formulations discussed herein are believed to provide useful amounts of solubilization of the active ingredient, and to provide sufficient formulation stability, including sufficient stability of the active ingredient, so as to be suitable for use in the administration of pharmaceutically active compounds to human subjects. Suitable stability includes the ability of a formulation to maintain effective amounts of relacorilant over time (over a period of months, or potentially even years, of storage). In particular, all of the formulations discussed herein are believed to provide useful amounts of solubilization of relacorilant, and to provide sufficient formulation stability, including sufficient stability of the relacorilant, so as to be suitable for use in the administration of relacorilant to human subjects. In addition, the formulations disclosed herein are believed to provide improved bioavailability of relacorilant as compared to prior or alternative formulations. The formulations including relacorilant disclosed herein are further believed to provide improved biocompatibility as compared to prior or alternative formulations. The formulations disclosed herein, being suitable for use in the administration of relacorilant to human subjects, are believed to be suitable for use in the treatment of human subjects suffering from disorders amenable to treatment by HKFA compounds such as relacorilant. The formulations disclosed herein are believed to provide improved pharmaceutical formulations which solve the problems of poor solubility, or poor bioavailability, or poor biocompatibility, or poor stability, previously encountered with prior or alternative formulations of relacorilant for administration to human subjects.

[0107] The following examples are provided by way of illustration only and not by way of limitation. Those of skill will readily recognize a variety of noncritical parameters which could be changed or modified to yield essentially similar results.

EXAMPLE 1 “CLINICAL FORMULATION”

[0108] Relacorilant was found to be stable in acidic aqueous solutions; however, relacorilant was found to degrade over time in neutral solutions at elevated temperatures, and was found to be sensitive to basic conditions. Relacorilant stability and compatibility with various excipients was

tested and suitable excipients were identified. An initial formulation comprising a polyethoxylated castor oil (Kolliphor EL, available from BASF, Florham Park, New Jersey) and a monodiglyceride mixture (Capmul MCM, a monodiglyceride of medium chain, mainly caprylic and capric, fatty acids, available from ABITEC Corporation, Columbus, Ohio) demonstrated significant decomposition (i.e., instability) when stability was tested for two weeks at 40 °C at 75% relative humidity, and at 60 °C. Other formulations including propylene glycol monocaprylates (Capryol PGMC, available from Gattefosse USA, New Jersey) and stearyl polyoxyl-32 glycerides (Gelucire 50/13, available from Gattefosse USA, New Jersey) failed to provide desirable dissolution properties. Lauryl polyoxyl-32 glycerides (Gelucire 44/14, available from Gattefosse USA, New Jersey) in place of the stearyl polyoxyl-32 glycerides were then used. As discussed below, the formulation of Table III surprisingly provided improved stability, suitable for use in relacorilant formulations for administration to human patients, including oral administration.

[0109] Several test formulations were prepared and subjected to stability and other testing. The compositions of these test formulations are presented in Table I, in which “API” is relacorilant; “MDC” indicates medium-chain monodiglycerides (mainly caprylic and capric); “PEC” indicates polyethoxylated castor oil; “LPG” indicates lauroyl polyoxyl-32 glycerides; “PGM” indicates propylene glycol monocaprylates; “BHT” indicates butylated hydroxytoluene; “BHA” indicates butylated hydroxyanisole; “GLT” indicates propyl gallate; “MYR” indicates myristic acid; and “AT” indicates α -Tocopherol.

TABLE I. TEST FORMULATIONS

Formulation	Composition (weight %)	pH
A	20 API, 40 MDC, 40 PEC	7.0
B	20 API, 39.96 MDC, 40 PEC, 0.04 BHT	
C	20 API, 39.91 MDC, 40 PEC, 0.04 BHT, 0.04 BHA	
D	20 API, 39.76 MDC, 40 PEC, 0.04 BHT, 0.2 GLT	7.0
E	20 API, 39.8 MDC, 40 PEC, 0.2 AT	
F	20 API, 39.26 MDC, 40 PEC, 0.04 BHT, 0.2 GLT, 0.5 MYR	

G	20 API, 59.98 LPG, 20 PGM, 0.02 BHT	4.0
---	-------------------------------------	-----

[0110] The stability of relacorilant in the several test formulations was examined for two weeks at 50 °C. The degree of degradation seen with the preferred formulation (formulation G in Table I above) was suggestive of a good compatibility with these reagents during manufacturing and long term storage. Amounts of relacorilant and relacorilant-derived impurities were measure by high pressure liquid chromatography (HPLC). There was no significant reduction in the amount of relacorilant in formulation G after two weeks storage at 50 °C. The highest percentage of relacorilant, the least amount of impurities, and the smallest increase in impurities after two weeks storage at 50 °C was found with formulation G, as compared to the other formulations.

[0111] The stability of relacorilant in a series of potential excipients was also examined for two weeks at 40 °C at 75% relative humidity, and at 60 °C, including in formulations of Table I. The predominant degradation path was through formation of the aromatic decarbonylation of relacorilant with both propylene glycol monocaprylate and lauroyl polyoxyl-32 glyceride excipients. Relacorilant stability was good with both these excipients. In the formulation of Table I, relacorilant showed good stability at 40°C/ 75%RH. Degradation was more pronounced in lauroyl polyoxyl-32 glycerides than propylene glycol monocaprylates at 60°C; however, the degree of degradation seen was suggestive of a good compatibility with these reagents during manufacturing and long term storage. These results are presented in Table II (showing purity, as measured by % of (relacorilant at the end of the test period) compared to (the total of relacorilant plus relacorilant-derived impurities).

Table II. Compatibility of Relacorilant with Excipients

Ingredient	Control	T ₀	40°C/75%RH		60°C	
			T (1 week)	T (2 week)	T (1 week)	T (2 week)
Propylene glycol monocaprylate	98.03%	98.09%	98.00%	97.84%	97.51%	96.65%
Lauroyl polyoxyl-32 Glycerides	98.09%	98.13%	97.64%	97.55%	97.22%	96.77%

[0112] As a result of such testing, the final, improved formulation was determined.

[0113] The following Table III provides an improved formulation suitable for incorporation of relacorilant in a softgel capsule with improved dissolution and stability. The formulation may be used to provide, e.g., 100 mg relacorilant per capsule (which may be suitable, for example, for administration to patients suffering from Cushing's syndrome or Cushing's Disease). The formulation may also be used to provide 150 mg relacorilant per capsule, or other amounts of relacorilant, with appropriately scaled percentages of other ingredients (retaining the same relative proportions of all ingredients).

[0114] The formulation of Table III included the following ingredients (percentages are weight per cent):

TABLE III "Clinical Formulation"

Ingredients	Function	Composition (% w/w)	Per 100 mg Relacorilant Dosage Form
Relacorilant	API	20.0	100 mg
Lauryl polyoxyl-32 glycerides	Surfactant	59.98	299.9 mg
propylene glycol monocaprylates	Solubilizer	20.0	100 mg
BHT, BHA	Antioxidant	0.02	0.1 mg
Total		100%	500 mg

[0115] Relacorilant (the API) solubilizes well in this formulation, and is stable in this formulation. This formulation is semi-solid. The formulation provides suitable bioavailability for clinical use, as shown in the pharmacokinetic profile illustrated in Fig. 1. Eight fasted human volunteers were administered 150 milligrams of relacorilant in the formulation of Table III, and the plasma levels of relacorilant were measured over time after administration. Mean (with

standard deviations) of the eight subjects are shown. Peak plasma concentrations of about 175 ng/mL were measured in these subjects.

[0116] This formulation was tested for stability. Stability testing included testing of some formulations for up to one year. Applicant has determined that relacorilant is sensitive to both moisture and light. In addition, Applicant has determined that relacorilant is more stable in formulations having neutral or acidic pH, while it is significantly less stable over time in formulations having basic pH (e.g., particularly at pH greater than 8).

[0117] An *in vitro* model was also used to model the effects of digestion of the formulation (see Williams HD et al., *J. Pharm Sci.* 101:3360-3380 (2012)). A schematic diagram illustrating the *in vitro* model is shown in Fig. 2. The results of this testing is believed to be indicative of the bioavailability of the drug product when administered *in vivo*. The *in vitro* digestive model includes three phases in a single container: i) a pellet that contains precipitated drug and insoluble (e.g., calcium) soaps of fatty acids; ii) an aqueous phase containing drug solubilized in micellar and vesicular structure, which comprises bile salts (BS, such as, e.g., sodium taurodeoxycholate as used by Williams et al.) and phospholipids (PL) and exogenous lipid digestion products (e.g., fatty acids (FA), monoglycerides (MG), and diglycerides (DG)); and iii) an undispersed oil phase, typically containing drug, remaining triglycerides (TG), and some diglycerides (DG). The experimental configuration used when performing an *in vitro* digestion model is shown in FIG. 2, which illustrates an idealized side view of a container used for *in vitro* digestion testing of test formulations. The container holds three phases into which the active pharmaceutical ingredient may partition. The amount of API which is found in each of the three phases provides a measure of the expected behavior of the API *in vivo*, and allows comparison of the expected bioavailability that would result from the formulations being compared. For example, the amount of API found in the aqueous phase provides an estimate of the amount (or proportion) of the API that would be found in the plasma of a subject following administration of the API in that formulation.

[0118] FIG. 3 shows the results of *in vitro* digestion testing for the formulation of Table III. The data presented in FIG. 3 illustrate the performance of the formulation of Table III in the *in vitro* digestion model over time. Approximately 10 – 15% percent of relacorilant partitioned into the aqueous phase early on during the testing, and then that percentage was reduced over time. These

results further indicate that there is a correlation between in vitro digestion model and human PK profile for the formulation (see FIG. 1).

[0119] The formulation of Table III was also administered to monkeys for in vivo testing (100 mg of the API, relacorilant). These results show considerable plasma exposure (AUC_{inf}) in monkeys. These results are presented in Table IV.

TABLE IV 100 mg Dose

Parameter	Plasma Levels (Clinical Formulation)
C_{max} (ng/mL)	1266
AUC_{inf}	11653
AUC_{0-24}	10347

[0120] In embodiments, it will be understood by those of skill in the art that modifications to these formulations may include: using purified commercial excipients (for example, purified Kolliphor ELP significantly reduces impurities as compared to Kolliphor products not labeled as “purified”); and lower pH (e.g., pH of less than pH 8, and preferably about pH 7 or less) may help to further reduce impurities. It is believed that reducing impurities improves the stability and other performance parameters of the formulation.

[0121] Other relacorilant-containing formulation have been prepared, some prior to the preparation of Table III, some prepared after its testing in attempts to improve on the formulation of Table III. For example, an initial formulation comprising a polyethoxylated castor oil with a monodiglyceride mixture (a monodiglyceride of medium chain, mainly caprylic and capric, fatty acids) demonstrated significant decomposition (i.e., instability) when stability was tested for two weeks at at 40 °C at 75% relative humidity, and at 60 °C. An attempt to improve the formulation of Table III (a formulation of 20% relacorilant, 40% polyoxylated castor oil surfactant, and 40% glycerol caprylate solubilizer, where per cent is weight per cent) was found to be unstable (impurities increased 4.5% in one month at 40 °C at 75 % relative humidity), and proved to be unsuitable for clinical use for at least that reason. Accordingly, the formulation of Table III was found to be surprisingly stable while providing suitable bioavailability, in contrast to all other formulations tested. Thus, the formulation of Table III provides surprising and unexpected advantages over other formulations.

EXAMPLE 2 FORMULATIONS FOR A RANGE OF RELACORILANT AMOUNTS

[0122] Relacorilant formulations may be provided which include multiple amounts of relacorilant. For example, single unit dosage forms each containing 25 mg, 50 mg, 100 mg, 200 mg, 300 mg, 400 mg, and 500 mg of relacorilant may be prepared according to the methods provided herein. This example provides instructions for preparing relacorilant formulations for the preparation of formulations for use in single unit dosage forms of relacorilant containing a range of relacorilant amounts. The single unit dosage forms may be softgel capsules containing these formulations.

TABLE V Relacorilant Formulations (for Several Dose Amounts)

Component	Amount per capsule (mg)						
Capsule Fill Formulation:							
Relacorilant	25.00	50.00	100.00	200.00	300.00	400.00	500.00
Lauroyl polyoxyl-32 glycerides	74.975	149.95	299.90	599.80	899.70	1199.60	1499.50
Propylene glycol monocaprylate	25.00	50.00	100.00	200.00	300.00	400.00	500.00
Butylated hydroxytoluene (BHT)	0.025	0.05	0.10	0.20	0.30	0.40	0.50
Total Weight	125	250	500	1000	1500	2000	2500

[0123] Softgel capsules may be filled with the formulations described in Table V above to provide single unit dosage forms containing the desired amounts of relacorilant. Softgel capsules containing the relacorilant formulations described in Table V may be made of about 50% to about 75% gelatin and about 25% to about 50% of a sorbitol/glycerin blend (by weight), with purified water and/or colorants included as needed. In some embodiments, softgel capsules containing these relacorilant formulations may be made of about 60% to about 70% gelatin and about 30% to about 40% of a plasticizer, or about 63% to about 66% gelatin and about 34% to about 37% of a plasticizer, with purified water and/or colorants included as needed. In particular

embodiments, softgel capsules containing these relaxorilant formulations may be made of 64% gelatin and 36% of a sorbitol/glycerin blend (by weight), or of 63.64% gelatin and 36.36% of a sorbitol/glycerin blend (by weight), these percentages not including the weight of any purified water and/or colorants that may be included as needed.

[0124] All patents, patent publications, publications, and patent applications cited in this specification are hereby incorporated by reference herein in their entireties as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. In addition, although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.