

Biofilm Disruption

Field of the Invention

The present invention relates to methods of dispersing biofilms comprising Gram-negative bacteria, the methods comprising exposing the biofilm to an epoxytiglicenone
5 compound or a salt thereof. Methods of treating infections comprising the localised administration, for example, topically or by injection, of an epoxytiglicenone compound into or onto an established biofilm comprising Gram-negative bacteria to disrupt the structure of that biofilm and methods of preventing biofilms comprising Gram-negative bacteria forming or dispersing biofilms comprising Gram-negative biofilms that have
10 formed on medical devices are also described.

Background of the Invention

Gram-negative bacteria are associated with a range of infections of the respiratory, urinogenital and gastro-intestinal tracts and of the circulatory system. They are also a major factor in the persistence of chronic wounds, of osteomyelitis, and of surgical site,
15 medical device and post-implant infections. Serious infections with Gram-negative bacteria can cause significant morbidity and mortality, especially in immunocompromised patients. The most important clinical Gram-negative pathogens are *Pseudomonas aeruginosa*, *Acinetobacter baumannii* and members of the Enterobacteriaceae, especially *Escherichia coli* and *Klebsiella pneumoniae*. Examples
20 of other medically relevant Gram-negative bacteria include *Neisseria* species (which cause gonorrhoea and meningitis), *Hemophilus influenzae*, *Legionella pneumophila*, *Yersinia pestis*, *Proteus mirabilis* and *Salmonella* species.

The treatment of infections of Gram-negative bacteria has become increasingly problematic due both to (a) the emergence and spread, amongst many of these
25 microbes, of multidrug resistance to most, or all, conventional antibiotics (Ho *et al.* 2010; Doi *et al.* 2017) and (b) their ability to form recalcitrant, aggregated communities called biofilms (Cepas *et al.* 2019). The management of recalcitrant Gram-negative biofilms in particular poses a complex and challenging clinical problem with limited treatment options. Current strategies in many clinical situations involve removal
30 (where feasible) of the biofilm by aggressive physical methods (e.g. debridement) coupled with high-dose and often prolonged antibacterial chemotherapy as standard of

- 2 -

care (Hoiby *et al.* 2015). However, complete eradication of the biofilm with these methods is often unsuccessful leading to recurrence and selection for increased levels of bacterial resistance. This vicious cycle of chronicity further erodes the efficacy of antibiotics and facilitates the more widespread use of ‘last resort’ antibiotics, many of which have significant side effects.

There is an urgent need for new agents to treat Gram-negative biofilm infections. While traditional approaches to treating bacterial infections have been antibiotic-based and focused on directly targeting and inhibiting or killing the causal pathogens (using concepts of minimal inhibitory concentration (MIC) and minimal bactericidal concentration (MBC)), the complex nature of biofilms means that new therapeutic strategies are required.


Biofilms are bacterial communities embedded in a protective, self-produced matrix of extracellular polymeric substances (EPS) formed from polysaccharides, proteins, lipids and nucleic acids (RNA and extracellular DNA). Living in the biofilm, the resident bacteria are significantly more tolerant to antibiotics compared to their planktonic free-living counterparts (Hoiby *et al.* 2010) and are largely protected from the host’s immune system. Much of this resistance of biofilms can be attributed to the EPS, which presents a formidable physical barrier to penetration of antibiotics and cellular effectors of immunity (Gunn *et al.* 2016). The structure of biofilms as a dense polymeric ‘mesh’ of EPS and bacteria also means they are highly resistant to removal from tissue and material surfaces to which they strongly adhere.

In contrast to the current standards of care, which are critically dependent on antibiotics, a range of ‘non-antibiotic’ strategies to treat Gram-negative biofilms have been proposed and are currently under investigation. These include interfering with signalling networks in biofilms, targeting biofilm adhesion, and disrupting the EPS matrix of biofilms (Koo *et al.* 2017). Unlike conventional antibiotics, these strategies also have the potential benefit of minimising the development of resistance as they target bacterial ‘virulence factors’ rather than bacterial growth and hence induce low selection pressure.

- Of these ‘non-antibiotic’ strategies, disrupting the integrity of the EPS matrix of established biofilms is a particularly attractive. By making the EPS matrix more permeable, disaggregating the bacteria and interfering with the pathogenic signalling network, this strategy theoretically makes the individual bacteria more accessible to the immune system and/or to the potentiating the action of antibiotics (Gunn *et al.* 2016; Fleming & Rumbaugh 2017). Using this approach, a range of therapies involving enzymes (proteases, DNases, glycoside hydrolases), peptides, monoclonal antibodies and polymers that target specific components of the matrix (e.g. exopolysaccharides and eDNA) are under investigation (Koo *et al.* 2017).
- 10 Epoxytiglenones are small molecules with a range of bioactive properties. They are potent anti-tumour compounds, have antiparasitic properties, and stimulate immune and other cellular responses that promote healing of acute and chronic wounds. Previously, epoxytiglenones have been reported to have direct antibiotic activity against a range of Gram-positive bacteria (WO2007/070985 and WO2014/169356). However, in recent
- 15 studies to more specifically evaluate the antibiotic properties of epoxytiglenone compounds against pathogenic strains of three Gram-negative pathogens (*E. coli*, *P. aeruginosa* and *A. baumannii*), no antibiotic effects have been found and it was not possible to establish a minimal inhibitory concentration (MIC) in conventional planktonic culture systems used for routine antibiotic screening.
- 20 The present invention is predicated, at least in part, on the subsequent, unexpected discovery that despite the absence of any direct antibiotic activity against planktonic Gram-negative bacteria, epoxytiglenones have potent, ‘non-antibiotic’ effects on established biofilms of a range of Gram-negative bacteria.

Summary

- 25 In an example of the disclosure there is provided a method of dispersing a biofilm comprising Gram-negative bacteria comprising exposing the biofilm to an epoxytiglenone compound for formula (I):



wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

R₂ is selected from -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -
 5 OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₃ is selected from -OH, -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -
 OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;


R₆ is selected from hydrogen, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C(O)C<sub>1-
 10 6</sub>alkyl, -C(O)C₂₋₆alkenyl, -C(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈
 cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈
 cycloalkyl, -C(O)aryl, -C(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -OC(O)C₂₋₆
 alkynylaryl;

R₇ is selected from hydroxy, -OC₁₋₆alkyl, -OC₂₋₆alkenyl, -OC₂₋₆alkynyl, -OC(O)C<sub>1-
 15 6</sub>alkyl, -OC(O)C₂₋₆alkenyl, -OC(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈
 cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈
 cycloalkyl, -OC(O)aryl, -OC(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -C(O)C₂₋₆
 alkynylaryl;

R₈ is selected from hydrogen and C₁₋₆alkyl; or a salt thereof.

20 In an aspect, there is provided an *ex vivo* method of dispersing a biofilm comprising
 Gram-negative bacteria comprising exposing the biofilm to an epoxytiglenone
 compound for formula (II):

- 4a -



(II)

wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

R₂ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

5 R₃ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;


R₆ is selected from hydrogen and -C(O)C₁₋₆alkyl;

R₇ is hydroxy; and

R₈ is selected from hydrogen or C₁₋₆alkyl; or a pharmaceutically acceptable salt thereof.

10

In an example of the disclosure there is provided a method of treating a bacterial infection comprising a biofilm comprising Gram-negative bacteria, said method comprising locally administering to the bacterial infection an epoxytiglenone compound of formula (I):



wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

R₂ is selected from -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -
 5 OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₃ is selected from -OH, -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -
 OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;


R₆ is selected from hydrogen, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C(O)C<sub>1-
 10 6</sub>alkyl, -C(O)C₂₋₆alkenyl, -C(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈
 cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈
 cycloalkyl, -C(O)aryl, -C(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -OC(O)C₂₋₆
 alkynylaryl;

R₇ is selected from hydroxy, -OC₁₋₆alkyl, -OC₂₋₆alkenyl, -OC₂₋₆alkynyl, -OC(O)C<sub>1-
 15 6</sub>alkyl, -OC(O)C₂₋₆alkenyl, -OC(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈
 cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈
 cycloalkyl, -OC(O)aryl, -OC(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -C(O)C₂₋₆
 alkynylaryl;

R₈ is selected from hydrogen and C₁₋₆alkyl; or a pharmaceutically acceptable salt thereof.

20 In a further aspect of the disclosure there is provided a use of an epoxytiglienone
 compound of formula (II):

- 5a -



wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

R₂ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

5 R₃ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;


R₆ is selected from hydrogen and -C(O)C₁₋₆alkyl;

R₇ is hydroxy;

R₈ is selected from hydrogen and C₁₋₆alkyl; or a pharmaceutically acceptable salt thereof, in
 10 the manufacture of a medicament for the treatment of a bacterial infection comprising a biofilm
 comprising Gram-negative bacteria wherein the compound of formula (II) is to be administered
 locally to the bacterial infection.

In a further aspect of the disclosure there is provided an *ex vivo* method of preventing a biofilm
 comprising Gram-negative bacteria forming on a medical device or dispersing a biofilm
 15 comprising Gram-negative bacteria on a medical device, said method comprising


applying an epoxytiglienone compound of formula (I) to the medical device: the compound of formula (I) comprising



wherein

- 5 R₁ is selected from hydrogen and C₁₋₆alkyl;
R₂ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;
R₃ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;
R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;
R₆ is selected from hydrogen and -C(O)C₁₋₆alkyl;
10 R₇ is hydroxy; and
R₈ is selected from hydrogen and C₁₋₆alkyl; or a pharmaceutically acceptable salt thereof

- In a further example of the disclosure there is provided a method of preventing a biofilm comprising Gram-negative bacteria forming or dispersing a biofilm comprising Gram-negative bacteria on a medical device, said method comprising applying an epoxytiglienone compound
- 15 of formula (I) to the medical device, the compound of formula (I) being:



wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

R₂ is selected from -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -
 5 OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₃ is selected from -OH, -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -
 OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;

R₆ is selected from hydrogen, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C(O)C<sub>1-
 10 6</sub>alkyl, -C(O)C₂₋₆alkenyl, -C(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈
 cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈
 cycloalkyl, -C(O)aryl, -C(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -OC(O)C₂₋₆
 alkynylaryl;

R₇ is selected from hydroxy, -OC₁₋₆alkyl, -OC₂₋₆alkenyl, -OC₂₋₆alkynyl, -OC(O)C<sub>1-
 15 6</sub>alkyl, -OC(O)C₂₋₆alkenyl, -OC(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈
 cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈
 cycloalkyl, -OC(O)aryl, -OC(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -C(O)C₂₋₆
 alkynylaryl;

R₈ is selected from hydrogen and C₁₋₆alkyl; or a pharmaceutically acceptable salt thereof.

Detailed Description

Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art to which the invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, preferred methods and materials are described. For the purposes of the present invention, the following terms are defined below.

- 7 -

The articles "a" and "an" are used herein to refer to one or to more than one (i.e. to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

As used herein, the term "about" refers to a quantity, level, value, dimension, size, or amount that varies by as much as 25%, 20%, 15% or 10% to a reference quantity, level, value, dimension, size, or amount.

Throughout this specification, unless the context requires otherwise, the words "comprise", "comprises" and "comprising" will be understood to imply the inclusion of a stated step or element or group of steps or elements but not the exclusion of any other step or element or group of steps or elements.

The term "alkyl" refers to optionally substituted linear and branched hydrocarbon groups having 1 to 8 carbon atoms. Where appropriate, the alkyl group may have a specified number of carbon atoms, for example, -C₁-C₆ alkyl which includes alkyl groups having 1, 2, 3, 4, 5 or 6 carbon atoms in linear or branched arrangements. Non-limiting examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, 2-methylpropyl, 1-methylpropyl, 2,2-dimethylethyl, pentyl, 2-methylbutyl, 3-methylbutyl, 1,2-dimethylpropyl, 2,3-dimethylpropyl, 3,3-dimethylpropyl, hexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl and 3-ethylbutyl, heptyl, 1-methylhexyl, 2-methylhexyl, 3-methylhexyl, 4-methylhexyl, 5-methylhexyl, 1-ethylpentyl, 2-ethylpentyl, 3-ethylpentyl, 1,1-dimethylpentyl, 1,2-dimethylpentyl, 1,3-dimethylpentyl, 1,4-dimethylpentyl, 1,5-dimethylpentyl, 2,3-dimethylpentyl, 2,4-dimethylpentyl, 1,1-dimethylpentyl, 2,2-dimethylpentyl, octyl, 1-methylheptyl, 2-methylheptyl, 3-methylheptyl, 4-methylheptyl, 5-methylheptyl, 1-ethylhexyl, 2-ethylhexyl, 3-ethylhexyl, 1,1-dimethylhexyl, 1,2-dimethylhexyl, 1,3-dimethylhexyl, 1,4-dimethylhexyl, 1,5-dimethylhexyl, 2,3-dimethylhexyl, 2,4-dimethylhexyl, 1,1-dimethylhexyl, 2,2-dimethylhexyl, and the like.

The term "alkenyl" refers to optionally substituted, unsaturated linear or branched hydrocarbons, having 2 to 8 carbon atoms and having at least one double bond. Where appropriate, the alkenyl group may have a specified number of carbon atoms, for

- example, C₂-C₆ alkenyl which includes alkenyl groups having 2, 3, 4, 5 or 6 carbon atoms in linear or branched arrangements. Non-limiting examples of alkenyl groups include, ethenyl, propenyl, 1-methylethenyl, butenyl, 1-methylprop-1-enyl, 1-methylprop-1-enyl, 1-ethylethenyl, pentenyl, 1-methylbut-1-enyl, 2-methylbut-1-enyl, 2-methylbut-2-enyl, 1,2-dimethylprop-1-enyl, 1,2-dimethylprop-2-enyl, hexenyl, 1-methylpent-1-enyl, 2-methylpent-1-enyl, 3-methylpent-1-enyl, 1-ethylbut-1-enyl, 2-ethylbut-1-enyl, 1-methylpent-2-enyl, 2-methylpent-2-enyl, 3-methylpent-2-enyl, 4-methylpent-2-enyl, 1-ethylbut-2-enyl, 2-ethylbut-2-enyl, 1,2-dimethylbut-2-enyl, 1,3-dimethylbut-2-enyl, 2,3-dimethylbut-2-enyl, 1-methylpent-3-enyl, 2-methylpent-3-enyl, 3-methylpent-3-enyl, 4-methylpent-3-enyl, 1-ethylbut-3-enyl, 2-ethylbut-3-enyl, 3-ethylbut-3-enyl, 1,2-dimethylbut-3-enyl, 1,3-dimethylbut-3-enyl, 1,1-dimethylbut-3-enyl, 2,3-dimethylbut-3-enyl, 2,2-dimethylbut-3-enyl, hex-1,3-diene, 1-heptenyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 5-heptenyl, 6-heptenyl, 2,4-heptadienyl, 2,6-heptadienyl, 2,4,6-heptatrienyl, 1-methylhexa-1-enyl, 2-methylhexa-1-enyl, 3-methylhexa-1-enyl, 4-methylhexa-1-enyl, 5-methylhexa-1-enyl, 1-methylhexa-2-enyl, 2-methylhexa-2-enyl, 3-methylhexa-2-enyl, 4-methylhexa-2-enyl, 5-methylhexa-2-enyl, 1-methylhexa-3-enyl, 2-methylhexa-3-enyl, 3-methylhexa-3-enyl, 4-methylhexa-3-enyl, 5-methylhexa-3-enyl, 1-methylhexa-4-enyl, 2-methylhexa-4-enyl, 3-methylhexa-4-enyl, 4-methylhexa-4-enyl, 5-methylhexa-4-enyl, 1-methylhexa-5-enyl, 2-methylhexa-5-enyl, 3-methylhexa-5-enyl, 4-methylhexa-5-enyl, 5-methylhexa-5-enyl, 1-octenyl, 2-octenyl, 3-octenyl, 4-octenyl, 5-octenyl, 6-octenyl, 2,4-octadienyl, 2,6-octadienyl, 2,4,6-octatrienyl, 1-methylhepta-1-enyl, 2-methylhepta-1-enyl, 3-methylhepta-1-enyl, 4-methylhepta-1-enyl, 5-methylhepta-1-enyl, 1-methylhepta-2-enyl, 2-methylhepta-2-enyl, 3-methylhepta-2-enyl, 4-methylhepta-2-enyl, 5-methylhepta-2-enyl, 1-methylhepta-3-enyl, 2-methylhepta-3-enyl, 3-methylhepta-3-enyl, 4-methylhepta-3-enyl, 5-methylhepta-3-enyl, 1-methylhepta-4-enyl, 2-methylhepta-4-enyl, 3-methylhepta-4-enyl, 4-methylhepta-4-enyl, 5-methylhepta-4-enyl, 1-methylhepta-5-enyl, 2-methylhepta-5-enyl, 3-methylhepta-5-enyl, 4-methylhepta-5-enyl, 5-methylhepta-5-enyl and the like.
- The term "alkynyl" refers to optionally substituted unsaturated linear or branched hydrocarbons, having 2 to 8 carbon atoms, having at least one triple bond. Where appropriate, the alkynyl group may have a specified number of carbon atoms, for

example, C₂-C₆ alkynyl which includes alkynyl groups having 2, 3, 4, 5 or 6 carbon atoms in linear or branched arrangements. Non-limiting examples include ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 1-methylprop-2-ynyl, 1-pentynyl, 2-pentynyl, 3-pentynyl, 4-pentynyl, 1-methylbut-2-ynyl, 1,1-dimethylprop-2-ynyl, 3-methylbut-1-ynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl, 3-methylpent-1-ynyl, 4-methylpent-1-ynyl, 1-methylpent-2-ynyl, 4-methylpent-2-ynyl, 3,3-dimethylbut-1-ynyl, 1,1-dimethylbut-2-ynyl, 1,2-dimethylbut-3-ynyl, 1,1-dimethylbut-3-ynyl, 2,2-dimethylbut-3-ynyl, 1-heptynyl, 2-heptynyl, 3-heptynyl, 4-heptynyl, 5-heptynyl, 6-heptynyl, 3-methylhex-1-ynyl, 4-methylhex-1-ynyl, 1-methylhex-2-ynyl, 4-methylhex-2-ynyl, 3,3-dimethylpent-1-ynyl, 1,1-dimethylpent-2-ynyl, 1,2-dimethylpent-3-ynyl, 1,1-dimethylpent-3-ynyl, 2,2-dimethylpent-3-ynyl, 1-octynyl, 2-octynyl, 3-octynyl, 4-octynyl, 5-octynyl, 6-octynyl, 7-octynyl, 3-methylhept-1-ynyl, 4-methylhept-1-ynyl, 1-methylhept-2-ynyl, 4-methylhept-2-ynyl, 3,3-dimethylhex-1-ynyl, 1,1-dimethylhex-2-ynyl, 1,2-dimethylhex-3-ynyl, 1,1-dimethylhex-3-ynyl, 2,2-dimethylhex-3-ynyl, and the like

The terms "cycloalkyl" and "carbocyclic" refer to optionally substituted saturated or unsaturated mono-cyclic hydrocarbon groups. Where appropriate, the cycloalkyl group may have a specified number of carbon atoms, for example, C₃-C₆ cycloalkyl is a carbocyclic group having 3, 4, 5 or 6 carbon atoms. Non-limiting examples may include cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cyclohexadienyl and the like.

"Aryl" means a C₆-C₁₄ membered monocyclic, bicyclic or tricyclic carbocyclic ring system having up to 7 atoms in each ring, wherein at least one ring is aromatic. Examples of aryl groups include, but are not limited to, phenyl, naphthyl, tetrahydronaphthyl, indanyl and biphenyl. The aryl may comprise 1-3 benzene rings. If two or more aromatic rings are present, then the rings may be fused together, so that adjacent rings share a common bond.

Each alkyl, alkenyl, alkynyl, cycloalkyl or aryl whether an individual entity or as part of a larger entity may be optionally substituted with one or more optional substituents selected from the group consisting of C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₆cycloalkyl, oxo (=O), -OH, -SH, C₁₋₆alkylO-, C₂₋₆alkenylO-, C₃₋₆cycloalkylO-, C₁₋₆alkylS-, C₂₋₆alkenylS-,

- 10 -

C₃₋₆cycloalkylS-, -CO₂H, -CO₂C₁₋₆alkyl, -NH₂, -NH(C₁₋₆alkyl), -N(C₁₋₆alkyl)₂, -NH(phenyl), -N(phenyl)₂, -CN, -NO₂, -halogen, -CF₃, -OCF₃, -SCF₃, -CHF₂, -OCHF₂, -SCHF₂, -phenyl, -C₁₋₆alkylphenyl, -Ophenyl, -C(O)phenyl, -C(O)C₁₋₆alkyl. Examples of suitable substituents, particularly for the cycloalkyl and aryl groups in R₆ and R₇,
5 include, but are not limited to, methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl, *tert*-butyl, vinyl, methoxy, ethoxy, propoxy, isopropoxy, butoxy, methylthio, ethylthio, propylthio, isopropylthio, butylthio, hydroxy, hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, fluoro, chloro, bromo, iodo, cyano, nitro, -CO₂H, -CO₂CH₃, -C(O)CH₃, trifluoromethyl, trifluoromethoxy, trifluoromethylthio,
10 difluoromethyl, difluoromethoxy, difluoromethylthio, amino, methylamino, dimethylamino, phenyl, phenoxy, phenylcarbonyl, benzyl and acetyl.


The epoxytiglicenone compounds may be in the form of pharmaceutically acceptable salts. It will be appreciated however that non-pharmaceutically acceptable salts also fall within the scope of the invention since these may be useful as intermediates in the
15 preparation of pharmaceutically acceptable salts or may be useful during storage or transport, or may be useful in non-pharmaceutical settings. Suitable pharmaceutically acceptable salts include, but are not limited to, salts of pharmaceutically acceptable inorganic acids such as hydrochloric, sulphuric, phosphoric, nitric, carbonic, boric, sulfamic, and hydrobromic acids, or salts of pharmaceutically acceptable organic acids
20 such as acetic, propionic, butyric, tartaric, maleic, hydroxymaleic, fumaric, maleic, citric, lactic, mucic, gluconic, benzoic, succinic, oxalic, phenylacetic, methanesulphonic, toluenesulphonic, benzenesulphonic, salicylic, salicylanilic, aspartic, glutamic, edetic, stearic, palmitic, oleic, lauric, pantothenic, tannic, ascorbic and valeric acids.

25 Base salts include, but are not limited to, those formed with pharmaceutically acceptable cations, such as sodium, potassium, lithium, calcium, magnesium, ammonium and alkylammonium.

Basic nitrogen-containing groups may be quarternised with such agents as lower alkyl halide, such as methyl, ethyl, propyl, and butyl chlorides, bromides and iodides; dialkyl
30 sulfates like dimethyl and diethyl sulfate; and others.

- It will also be recognised that the epoxytiglienone compounds may possess asymmetric centres and are therefore capable of existing in more than one stereoisomeric form. The invention thus also relates to compounds in substantially pure isomeric form at one or more asymmetric centres e.g., greater than about 90% ee, such as about 95% or 97% ee or greater than 99% ee, as well as mixtures, including racemic mixtures, thereof. Such isomers may be obtained by isolation from natural sources, by asymmetric synthesis, for example using chiral intermediates, or by chiral resolution. The compounds of the invention may exist as geometrical isomers. The invention also relates to compounds in substantially pure *cis* (Z) or *trans* (E) forms or mixtures thereof.
- 10 The compounds of the present invention may be obtained by isolation from a plant or plant part, or by derivatisation of the isolated compound, or by derivatisation of a related compound. Isolation procedures and derivatisation procedures may be found in WO 2007/070985 and WO2014/169356.


- 15 The term "6,7-epoxytiglienone compound" refers to a compound having the following carbon cyclic structure:



- The compounds have a tricyclo[9.3.0.0]tetradecane system with a fused cyclopropane ring appended to the six membered ring. The epoxide is fused to the seven-membered ring in the 6,7-position and the five membered ring has a 1,2-ene-3-one structure.
- 20 **Methods of the Invention**

The present invention provides a method of dispersing a biofilm comprising Gram-negative bacteria, said method comprising exposing the biofilm to an epoxytiglienone compound for formula (I):

- 12 -



wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

R₂ is selected from -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl,
5 -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₃ is selected from -OH, -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl,
-OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;

R₆ is selected from hydrogen, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C(O)C₁₋₆alkyl,
10 -C(O)C₂₋₆alkenyl, -C(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈cycloalkyl,
-C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈cycloalkyl,
-C(O)aryl, -C(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -OC(O)C₂₋₆alkynylaryl;

R₇ is selected from hydroxy, -OC₁₋₆alkyl, -OC₂₋₆alkenyl, -OC₂₋₆alkynyl, -OC(O)C₁₋₆alkyl,
-OC(O)C₂₋₆alkenyl, -OC(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈cycloalkyl,
15 -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈cycloalkyl,
-OC(O)aryl, -OC(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -C(O)C₂₋₆alkynylaryl;

R₈ is selected from hydrogen and C₁₋₆alkyl; or a salt thereof.

The biofilm comprising the Gram-negative bacteria may be in any situation suitable for treatment. The biofilm may be on an inanimate surface or may be on or in a biological
20 system. In some embodiments, the biofilm is on an inanimate surface, for example, a bench such as a medical, laboratory, food preparation or manufacturing surface such as a surgery bed or floor, a laboratory bench, a kitchen bench or floor, manufacturing equipment for pharmaceuticals, nutraceuticals, cosmetics or personal care products. In some embodiments, the inanimate surface may be a medical device such as medical
25 instruments, surgical instruments, catheters, prostheses and implants. In some embodiments, the biofilm is infecting a subject, for example, a wound such as a surgical

wound or a burn. In some embodiments, the biofilm is infecting the subject at the site of the use of a medical device such as a catheter or an implant.

In some embodiments, the biofilm comprises only Gram-negative bacteria. In some embodiments, the biofilm only comprises one species of Gram-negative bacteria. In
5 other embodiments, the biofilm comprises more than one species of Gram-negative bacteria. In some embodiments, the biofilm comprises populations of Gram-negative and Gram-positive bacteria.


The term “dispersing” as used herein means that the biofilm is degraded such that at least a portion of the bacteria are released from the matrix of extracellular polymeric
10 substances (EPS) and assume a planktonic state. Once in a planktonic state, the bacteria are accessible to the immune system and/or to the action of antibiotics.

In some embodiments, the biofilm comprising Gram-negative bacteria comprises one or more Gram-negative bacteria selected from *Pseudomonas* species, *Acinetobacter* species, *Aeromonas* species, *Bacteroides* species, *Bordetella* species, *Borrelia* species,
15 *Burkholderia* species, *Citrobacter* species, *Compylobacter* species, *Escherichia* species, *Enterobacter* species, *Flavobacterium* species, *Fusobacterium* species, *Klebsiella* species, *Leptospira* species, *Neisseria* species, *Helicobacter* species, *Hemophilus* species, *Legionella* species, *Moraxella* species, *Yersinia* species, *Oligella* species, *Pantoea* species, *Porphyromonas* species, *Prevotella* species, *Proteus* species,
20 *Raoutella* species, *Salmonella* species, *Serratia* species, *Shigella* species, *Sphingomonas* species, *Stenotrophomonas* species, *Treponema* species, *Veillonella* species and *Vibrio* species, especially *Pseudomonas* species, *Acinetobacter* species, *Escherichia* species, *Klebsiella* species, *Neisseria* species, *Hemophilus* species, *Legionella* species, *Yersinia* species, *Proteus* species and *Salmonella* species, more
25 especially *Pseudomonas* species, *Acinetobacter* species, *Escherichia* species and *Klebsiella* species. In some embodiments, the Gram-negative bacteria are particularly good biofilm forming bacteria and are selected from *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, *Escherichia coli*, *Klebsiella pneumoniae*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Hemophilus influenzae*, *Legionella pneumophila*,
30 *Yersinia pestis*, *Yersinia enterocolitica*, *Salmonella enterica*, *Salmonella bongori*, *Proteus mirabilis*, *Enterobacter cloacae*, *Serratia marcescens*, *Bacteroides fragilis*,

- 14 -

Fusobacterium necrophorum, *Burkholderia cepacia* and *Prevotella intermedia*, especially *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, *Escherichia coli*, *Klebsiella pneumoniae*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Hemophilus influenzae*, *Legionella pneumophila*, *Yersinia pestis*, *Yersinia enterocolitica*,
 5 *Salmonella enterica* and *Salmonella bongori*, more especially *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, *Escherichia coli* and *Klebsiella pneumoniae*.

In another aspect of the invention the biofilm described above dispersed by the method is present in a bacterial infection on or in a subject. Therefore, the invention includes a method of treating a bacterial infection comprising a biofilm comprising Gram-negative
 10 bacteria, said method comprising locally administering to the bacterial infection an epoxytiglenone compound of formula (I):



wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

15 R₂ is selected from -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₃ is selected from -OH, -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;

20 R₆ is selected from hydrogen, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C(O)C₁₋₆alkyl, -C(O)C₂₋₆alkenyl, -C(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈cycloalkyl, -C(O)aryl, -C(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -OC(O)C₂₋₆alkynylaryl;

25 R₇ is selected from hydroxy, -OC₁₋₆alkyl, -OC₂₋₆alkenyl, -OC₂₋₆alkynyl, -OC(O)C₁₋₆alkyl, -OC(O)C₂₋₆alkenyl, -OC(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋

- 15 -

₈cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈cycloalkyl, -OC(O)aryl, -OC(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -C(O)C₂₋₆alkynylaryl;

R₈ is selected from hydrogen and C₁₋₆alkyl; or a pharmaceutically acceptable salt thereof.

- 5 In some embodiments, the infection is an acute infection. In other embodiments, the infection is a chronic infection. In some embodiments, the infection is a post-surgical infection or an infection at the site of insertion or use of a medical device, for example, at the site of insertion of a catheter, such as a venous catheter or a urinary catheter or the site of an implant such as a dental implant or a cochlear implant.
- 10 By the term "local administration" is meant that the compound of formula (I) is applied directly to the biofilm. In some embodiments, the local administration will be topical administration such as by use of a lotion, cream, ointment, foam, suspension, liquid wash or spray. In other embodiments, local administration may be achieved during surgery where the infected tissue is exposed, or delivered by injection guided by an
- 15 imaging technique, for example, guided by endoscopic ultrasound or stereotactic imaging. In particular embodiments, the local administration is to a biofilm containing infection located at a site accessible from outside of the body, for example, on the skin, on an external wound, in the mouth or nose, in the ear or ear canal, around the anus/rectum, the vagina, the urethra or the milk canal of domestic animals.
- 20 In some embodiments, the dispersal of the biofilm is accompanied by stimulation of the immune system such that the immune system is able to resolve the infection without the need for antibiotic administration, for example, by stimulating the expression in stromal and dermal cell types of host-defence peptides and of chemokines/cytokines involved in neutrophil/granulocyte recruitment, and by inducing the potent antimicrobial defences
- 25 of resident and recruited innate immune cells, including release of reactive oxygen species, extracellular nets and broad-spectrum antimicrobial peptides.

In other embodiments, the dispersal of the biofilm is accompanied by co-administration of an antibiotic. In some embodiments, the compound of formula (I) or salt thereof is administered in combination with an antibiotic to which the Gram-negative bacteria is

30 susceptible when in planktonic state. Thus, once the biofilm disperses, the planktonic

Gram-negative bacteria are exposed to the antibacterial activity of the antibiotic.

Suitable antibiotics for Gram-negative bacteria include, but are not limited to, 3rd generation cephalosporins such as ceftazidime and ceftiofur; 4th generation cephalosporins such as cefepime; aminoglycosides such as streptomycin, neomycin, gentamycin, amikacin, tobramycin and plazomicin; monocyclic beta lactams such as aztreonam; beta lactam inhibitor combinations such as piperacillin/tazobactam, amoxicillin/clavulanic acid, cefepime/AAI101, aztreonam/avibactam, ceftaroline/avibactam, imipenem/relebactam and meropenem/RPX7009; beta-lactams such as penicillin, cephalosporin, S649266 and BAL30072; Carbapenems such as imipenem, doripenem, ertapenem and meropenem; polymyxin-E antibiotics such as colistin; quinolones/fluoroquinolones such as ciprofloxacin, fleroxacin, norfloxacin, enrofloxacin, marbofloxacin, finafloxacin, lascufloxacin, avarofloxacin, nadifloxacin, delafloxacin, nemonoxacin and zabofloxacin; sulphonamides such as cotrimoxazole, tetracycline/glycylcyclines such as tigecycline, eravacycline and omadacycline; topoisomerase inhibitors such as ETX0914 and GSK2140944; and other antibiotics such as chloramphenicol and fosfomycin.

By the term “combination” is intended that the compound of formula (I) and the antibiotic are active in the biofilm such that the biofilm is disrupted and the planktonic bacteria exposed to the antibiotic. In some embodiments, the combination of compound of formula (I) and antibiotic are administered in a single composition. In other embodiments, the compound of formula (I) and the antibiotic are administered simultaneously or sequentially in separate compositions.

The subject that has the bacterial infection and may be treated is a mammal, a bird, an aquatic animal such as a fish, or a reptile. In some embodiments, the subject is a human, a laboratory animal such as a primate, mouse, rat or rabbit, a companion animal such as a dog or cat, a working animal such as a horse, donkey and the like, a livestock animal such as a cow, bull, pig, sheep, goat, deer, llama, alpaca and the like, or a captive wild animal such as those in zoos or wildlife parks including lions, leopards, cheetah, elephant, zebra, antelope, giraffe, koala, kangaroo and reptiles such as crocodiles, lizards, snakes and the like, a bird, especially a poultry bird such as a chicken, duck, goose, quail or pheasant, or a captive bird, such as a budgerigar or


canary, cockatoo, parakeet, macaw, parrot and the like, or a fish, especially a captive fish such as an aquaculture fish (salmon, trout, barramundi and the like) or tropical fish (zebra fish, guppy, Siamese fighting fish, clown fish, cardinal tetra and the like), dolphins, whales, and the like. In particular embodiments, the subject is a human or a companion animal.

An "effective amount" means an amount necessary at least partly to attain the desired response, such as biofilm dispersement within the infected site. The amount varies depending upon the health and physical condition of the individual to be treated, the taxonomic group of individual to be treated, the formulation of the composition, the assessment of the medical situation, and other relevant factors. It is expected that the effective amount of 6,7-epoxytiglicone compound will fall in a relatively broad range that can be determined through routine trials. An effective amount in relation to a human patient, for example, may lie in the range of about 0.1 ng per kg of body weight to 1 g per kg of body weight per dosage or 0.1 ng to 1 g per cm² of body surface per dosage. The dosage is preferably in the range of 1 µg to 1 g per kg of body weight per dosage or 1 µg to 1 g per cm² of body surface area, such as is in the range of 1 mg to 1 g per kg of body weight per dosage, or 1 mg to 1 g per cm² per body surface area per dosage. In one embodiment, the dosage is in the range of 1 ng to 500 mg per kg of body weight per dosage or 0.1 ng to 500 mg per cm² of surface area per dosage. In another embodiment, the dosage is in the range of 1 mg to 250 mg per kg of body weight per dosage or 1 mg to 250 mg per cm² of body surface area per dosage. In yet another embodiment, the dosage is in the range of 1 mg to 100 mg per kg of body weight per dosage or 1 mg to 100 mg per cm² of body surface area per dosage, such as up to 50 mg per kg of body weight per dosage or 50 mg per cm² of body surface area per dosage. In yet another embodiment, the dosage is in the range of 1 µg to 1 mg per kg of body weight per dosage or 1 µg to 1 mg per cm² of body surface area per dosage.

In a further aspect of the invention the biofilm described above dispersed by the method is present or potentially present on a medical device. Therefore, the invention includes a method of preventing a biofilm comprising Gram-negative bacteria forming on a medical device or dispersing a biofilm comprising Gram-negative bacteria on a medical device; said method comprising applying an epoxytiglicone compound of formula (I)

- 18 -

to the medical device; the compound of formula (I) being:



wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

5 R₂ is selected from -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₃ is selected from -OH, -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;

10 R₆ is selected from hydrogen, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C(O)C₁₋₆alkyl, -C(O)C₂₋₆alkenyl, -C(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈cycloalkyl, -C(O)aryl, -C(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -OC(O)C₂₋₆alkynylaryl;

15 R₇ is selected from hydroxy, -OC₁₋₆alkyl, -OC₂₋₆alkenyl, -OC₂₋₆alkynyl, -OC(O)C₁₋₆alkyl, -OC(O)C₂₋₆alkenyl, -OC(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈cycloalkyl, -OC(O)aryl, -OC(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -C(O)C₂₋₆alkynylaryl;

R₈ is selected from hydrogen and C₁₋₆alkyl; or a pharmaceutically acceptable salt thereof.


20 In some embodiments the medical device is coated with a composition comprising compound of formula (I) or a salt thereof to prevent the formation of a biofilm comprising Gram-negative bacteria. In other embodiments, the medical device is washed with or coated with a composition comprising a compound of formula (I) to disperse biofilm comprising Gram-negative bacteria and remove them from the device.

- 19 -

As used herein, the term “medical device” refers to a device that is used on human or animal bodies for therapeutic benefit and has a physical or mechanical effect on the body or is used to measure or monitor functions in the body. Suitable medical devices include indwelling medical devices such as urinary catheters, vascular access devices, endotracheal tubes, tracheostomies, enteral feeding tubes and wound drains; invasive medical devices such as central lines, mechanical heart valves, pacemakers, prosthetic or replacement joints, pins, rods, screws and plates for anchoring fractured bones, catheters such as urinary catheters, venous catheters, Swan-Ganz catheters, Quinton catheters, intrauterine catheters, drainage catheters and pigtail catheters; prosthetics such as limb prosthesis, dental prosthesis including dentures, obturators and dental implants; implants such as cochlear implants coronary stents, contraceptive implants, cosmetic implants and dental implants. Medical devices as used herein also include medical equipment such as syringes, specula, blood pressure monitors, scanners, ultrasound probes and the like; as well as surgical instruments such as scalpels, forceps, clamps, retractors, lancets, endoscopes and calipers.

In some embodiments, the wash or coating comprises an antiseptic, disinfectant or antibiotic.

The invention also relates to a use of compound of formula (I):



wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

R₂ is selected from -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₃ is selected from -OH, -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

- 20 -


R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;

R₆ is selected from hydrogen, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C(O)C₁₋₆alkyl, -C(O)C₂₋₆alkenyl, -C(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈cycloalkyl, -C(O)aryl, -C(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -OC(O)C₂₋₆alkynylaryl;

R₇ is selected from hydroxy, -OC₁₋₆alkyl, -OC₂₋₆alkenyl, -OC₂₋₆alkynyl, -OC(O)C₁₋₆alkyl, -OC(O)C₂₋₆alkenyl, -OC(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈cycloalkyl, -OC(O)aryl, -OC(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -C(O)C₂₋₆alkynylaryl;

R₈ is selected from hydrogen and C₁₋₆alkyl; or a pharmaceutically acceptable salt thereof; in the manufacture of a medicament for treating a bacterial infection comprising a biofilm comprising Gram-negative bacteria, wherein the medicament is adapted for local administration to the bacterial infection.

In another aspect, the present invention provides a compound of Formula (I) for use in treating a bacterial infection comprising a biofilm comprising Gram-negative bacteria, wherein the compound of formula (I) is:



wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

R₂ is selected from -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₃ is selected from -OH, -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;

- 21 -


R₆ is selected from hydrogen, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C(O)C₁₋₆alkyl, -C(O)C₂₋₆alkenyl, -C(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈cycloalkyl, -C(O)aryl, -C(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -OC(O)C₂₋₆alkynylaryl;

- 5 R₇ is selected from hydroxy, -OC₁₋₆alkyl, -OC₂₋₆alkenyl, -OC₂₋₆alkynyl, -OC(O)C₁₋₆alkyl, -OC(O)C₂₋₆alkenyl, -OC(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈cycloalkyl, -OC(O)aryl, -OC(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -C(O)C₂₋₆alkynylaryl;

10 R₈ is selected from hydrogen and C₁₋₆alkyl; or a pharmaceutically acceptable salt thereof; and the compound is for local administration to the bacterial infection.

6,7-epoxytiglienone compounds

The compounds that are useful in the methods of the present invention are compounds of formula (I):



15 wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

R₂ is selected from -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

20 R₃ is selected from -OH, -OC₁₋₈alkyl, -OC₂₋₈alkenyl, -OC₂₋₈alkynyl, -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;

R₆ is selected from hydrogen, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C(O)C₁₋₆alkyl, -C(O)C₂₋₆alkenyl, -C(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈cycloalkyl,

25 -C(O)aryl, -C(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -OC(O)C₂₋₆alkynylaryl;

- 22 -

R₇ is selected from hydroxy, -OC₁₋₆alkyl, -OC₂₋₆alkenyl, -OC₂₋₆alkynyl, -OC(O)C₁₋₆alkyl, -OC(O)C₂₋₆alkenyl, -OC(O)C₂₋₆alkynyl, -C(O)C₃₋₈cycloalkyl, -C(O)C₁₋₆alkylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkenylC₃₋₈cycloalkyl, -C(O)C₂₋₆alkynylC₃₋₈cycloalkyl, -OC(O)aryl, -OC(O)C₁₋₆alkylaryl, -C(O)C₂₋₆alkenylaryl and -C(O)C₂₋₆alkynylaryl;

- 5 R₈ is selected from hydrogen and C₁₋₆alkyl; or a salt thereof, especially a pharmaceutically acceptable salt thereof.

In particular embodiments of formula (I), one or more of the following applies:

R₁ is selected from -C₁₋₃alkyl, especially -CH₃;

- 10 R₂ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl; especially -OC(O)C₃₋₆alkyl and -OC(O)C₃₋₆alkenyl;

R₃ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl; especially -OC(O)C₃₋₆alkyl, -OC(O)C₃₋₆alkenyl and -OC(O)C₃₋₆alkynyl;


R₄ and R₅ are independently selected from -C₁₋₃alkyl; especially where both are methyl;

- 15 R₆ is selected from hydrogen, -C(O)C₁₋₆alkyl, -C(O)C₂₋₆alkenyl, -C(O)C₂₋₆alkynyl and -C(O)aryl; especially hydrogen, -C(O)CH₃, -C(O)CH₂CH₃, -C(O)CH(CH₃)₂ and -C(O)CH₂CH₂CH₃;

R₇ is selected from hydroxyl, -OC(O)C₁₋₆alkyl, -OC(O)C₂₋₆alkenyl and -OC(O)C₂₋₆alkynyl; especially hydroxyl;

R₈ is C₁₋₃alkyl, especially methyl.

- 20 In some embodiments, the compounds of formula (I) have stereochemistry as shown in formula (II) below:



In some embodiments, the epoxide in the 6,7-position is above the plane of the ring

system. In other embodiments, the epoxide in the 6,7-position is below the plane of the ring system. In some embodiments, the R₂ group in the 12 position is *S* and in other embodiments, the R₂ group in the 12 position is *R*.

In some embodiments of formula (I), the alkyl or alkenyl group of R₂ and/or R₃ are
5 branched alkyl or alkenyl groups. In other embodiments of formula (I), the alkyl or alkenyl group of R₂ and/or R₃ are linear alkyl or alkenyl groups.

In some embodiments, the alkyl or alkenyl group of R₂ and/or R₃ has a chain length of intermediate hydrophobicity, for example, C₄, C₅ or C₆.

In some embodiments, R₆ is an acyl group, for example an acetyl (-C(O)CH₃),
10 -C(O)CH₂CH₃, -C(O)CH(CH₃)₂ or -C(O)CH₂CH₂CH₃. In some embodiments, R₆ is hydrogen.

In particular embodiments, the epoxytiglienone compound is selected from:

- 12-tigloyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 1);
- 15 12,13-di-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 2);
- 12-hexanoyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 3);
- 12,13-dihexanoyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one
20 (Compound 4);
- 12-tigloyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13-pentahydroxy-20-acetyloxy-1-tiglien-3-one (Compound 5);
- 12-propanoyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 6);
- 25 12,13-ditigloyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 7);
- 12-(2-methylbutanoyl)-13-tigloyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 8);
- 30 12-butanoyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 9);

- 12-(3,3-dimethylbut-2-enoyl)-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 10);
- 12-hex-2,4-dienoyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 11);
- 5 12-tigloyl-13-(2-methylpropanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 12);
- 12-but-2-enoyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 13);
- 12-tigloyl-13-butanoyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one
10 (Compound 14);
- 12,13-dibutanoyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 15);
- 12,13-dipentanoyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 16);
- 15 12,13-di-(2*E*,4*E*)-hexa-2,4-dienoyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 17);
- 12,13-di-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 18);
- 12-(2-methylprop-2-enoyl)-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tigliaen-3-one (Compound 19);
- 20 12,13-di-heptanoyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tigliaen-3-one (Compound 20);
- 12,13-di-(3-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tigliaen-3-one (Compound 21);
- 25 or a salt thereof, especially pharmaceutically acceptable salts thereof.

In particular embodiments, the 6,7-epoxytiglienone compound is selected from compounds 1, 2, 3, 4 and 6.

The epoxytiglienone compounds may be obtained by extraction or may be obtained semisynthetically from extracted compounds. Methods for obtaining the
30 epoxytiglienone compounds are provided in WO2007/070985 and WO2014/169356, the contents of which are incorporated by reference.

Compositions

While the 6,7-epoxytiglienone compounds or salts thereof may be used neat, it may be more convenient to use them in the form of a composition, together with a carrier, diluent and/or excipient. In some embodiments, the composition may be a solution
5 useful to soak medical devices. In other embodiments, the composition may be a coating composition useful to coat medical devices. In yet other embodiments, the composition may be a pharmaceutical composition suitable for administration to a patient.

Dosage form and rates for pharmaceutical use and compositions are readily
10 determinable by a person of skill in the art.

The 6,7-epoxytiglienone compound is formulated for localized administration directly onto or into the biofilm. In some embodiments, the 6,7-epoxytiglienone compound is formulated for topical administration in the form of a gel, ointment, lotion, cream or transdermal patch that may be applied directly onto the biofilm. In other embodiments,
15 the epoxytiglianone compound is formulated for injection, for example, where the composition is injected internally to locally contact the biofilm.

In some embodiments, suitably, the composition is a pharmaceutical composition(s) and comprises a pharmaceutically acceptable excipient or an acceptable excipient. By “pharmaceutically acceptable excipient” is meant a solid or liquid filler, diluent or
20 encapsulating substance that may be safely used. Depending upon the particular route of administration, a variety of carriers, well known in the art may be used. These carriers or excipients may be selected from a group including sugars, starches, cellulose and its derivatives, cyclodextrins, malt, gelatine or other gelling agents, polymers, talc, calcium sulphate, vegetable oils, synthetic oils, alcohols and/or polyols, alginic acid,
25 phosphate buffered solutions, emulsifiers, isotonic saline, and pyrogen-free water.

Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water-propylene glycol solutions. For example, injectable liquid preparations can be formulated as solutions in aqueous 1,2-propanediol, dimethylsulfoxide (DMSO), aqueous solutions of gamma cyclodextrin or 2-hydroxypropyl-beta-cyclodextrin, saline

- 26 -

solution or polyethylene glycol solution, with or without buffer. A preferred range of pH is 3.0-4.5. Suitable buffers buffer the preparation at pH 3.5-4.5 and include, but are not limited to, acetate buffer and citrate buffer.

The compositions of 6,7-epoxytiglicenone compound may thus be formulated administration (e.g. by injection, for example bolus injection at the site of a biofilm infection) and may be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The compositions may take such forms as suspensions, solutions, gels or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising and/or dispersing agents. Alternatively, the active ingredient may be in powder form, obtained by aseptic isolation of sterile solid or by lyophilisation from solution, for constitution with a suitable vehicle, e.g. sterile, pyrogen-free water, before use.

Pharmaceutical compositions of 6,7-epoxytiglicenone compound suitable for administration may be presented in discrete units such as syringes, vials, tubes or sachets each containing a predetermined amount of one or more pharmaceutically active 6,7-epoxytiglicenone compounds, as a powder or granules or as a solution or a suspension in an aqueous liquid, a cyclodextrin solution, a non-aqueous liquid, an oil-in-water emulsion or a water-in-oil emulsion or as a solution or suspension in a cream or gel or as a suspension of micro- or nano-particles incorporating a 6,7-epoxytiglicenone compound, including but not limited to silica or polylactide micro- or nano-particles. Such compositions may be prepared by any of the methods of pharmacy but all methods include the step of bringing into association one or more pharmaceutically active compounds of the invention with the carrier which constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the agents of the invention with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product in to the desired presentation.

For topical administration to the epidermis or other organ, the compounds according to the invention may be formulated as gels, ointments, emulsions, pastes, creams or lotions, or as a transdermal patch. Gels may be prepared using suitable thickening

- 27 -

agents and adding them to aqueous/alcoholic compositions of compound. Suitable thickening or gelling agents are known in the art, such as the polyvinyl carboxy polymer Carbomer 940. Ointments and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents.

- 5 Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilising agents, dispersing agents, suspending agents, thickening agents, or colouring agents.

Formulations suitable for topical administration also include solutions or suspensions that may be administered topically in the form of a bath or soak solution or a spray or
10 may be absorbed into a dressing.

Liquid formulations suitable for soaking or washing medical devices may also include solutions or suspensions in solvents such as water and alcohols or mixtures thereof. The solutions or suspensions may also contain other components such as emulsifiers, surfactants, antiseptics, disinfectants, antibiotics, colouring agents and the like.

- 15 Coatings are suitably polymeric coatings known in the art that may incorporate a compound of formula (I) such that it is accessible at the surface of the coating and is sufficient to prevent a biofilm comprising Gram-negative bacteria forming. Suitable coatings include hydroxyapatite, calcium phosphate, biophosphonates, bioactive ceramics, polyhydroxyalkanoates, mesoporous materials, hydrogels and drug-eluting
20 coatings such as polymer coatings, calcium phosphate, chitosan, collagen and bone grafts/ cancellous bone.

Brief Description of the Figures

Figure 1 provides confocal laser scanning microscopy (CLSM) images showing biofilm disruption of *E. coli* IR57 following treatment with Compounds 1, 4 and 6
25 compared to the comparator compound CC-1, untreated and ethanol equivalent (blank) controls. Upper panel in each image group is overhead view, small lower panel is cross-sectional view.

Figure 2 provides **A:** biofilm biomass volume or bio-volume ($\mu\text{m}^3/\mu\text{m}^2$) that was quantified and confirmed with COMSTAT image analysis after treatment of the *E. coli*

IR57 biofilm with Compounds 1, 4 and 6, compared to the comparator compound CC-1, untreated and ethanol equivalent (blank) controls. Decreases in bio-volume were evident in biofilms treated with these compounds, which were significant with Compound 1 treatment ($p < 0.05$). **B:** there was no significant differences between any of the compounds and control treatments in the DEAD/LIVE bacterial ratio, demonstrating disruption to the *E. coli* IR57 biofilm density and bio-volume in these experiments was not related to a direct antibiotic activity.

Figure 3 provides a graph of the mean squared displacement (MSD) of 200 nm FluoSpheres® over time (in seconds) within *E. coli* IR57 biofilm structures. The disruption and alteration of the *E. coli* IR57 biofilm structure is illustrated by substantially higher mean squared displacement of the FluoSphere® particles following treatment with Compounds 1, 4 and 6, compared with an untreated and ethanol equivalent (blank) controls.

Figure 4 provides a graph of increased biofilm creep compliance (demonstrating) decreased resistance to mechanical deformation of the biofilm medium derived from the MSD vs lag time as seen in Figure 3) following treatment *E. coli* IR57 biofilm with Compounds 1, 4 and 6, compared to the untreated and ethanol equivalent (blank) controls.

Figure 5 provides **A:** the CLSM images when Compounds 1, 4, 6 and the comparator compound CC-1 are applied to biofilms of *A. baumannii* 7789 compared to an untreated and ethanol equivalent (blank) control. Upper panel in each image is overhead view, small lower panel is cross-sectional view. **B:** the CLSM images when compounds 1, 4, 6 and the comparator compound CC-1 are applied to biofilms of *P. aeruginosa* PAO1 compared to an untreated and ethanol equivalent (blank) control. Upper panel in each image is overhead view, small lower panel is cross-sectional view.

Figure 6 provides **A:** biofilm biomass volume or bio-volume ($\mu\text{m}^3/\mu\text{m}^2$) that was quantified and confirmed with COMSTAT image analysis after treatment of the *A. baumannii* 7789 biofilms with Compounds 1, 4, 6 and the comparator compound CC-1 compared to the untreated and ethanol equivalent (blank) controls. Significant decreases in bio-volume were evident in biofilms treated with compounds 1, 4 and comparator

compound CC-1 ($p < 0.05$). **B:** biofilm biomass volume or bio-volume ($\mu\text{m}^3/\mu\text{m}^2$) that was quantified and confirmed with COMSTAT image analysis after treatment of the *P. aeruginosa* PAO1 biofilms with Compounds 1, 4, 6 and the comparator compound CC-1 compared to the untreated and ethanol equivalent (blank) controls. Only Compound 4 had a significant effect on biofilm bio-volume compared to other treatments ($P < 0.05$). **C & D:** there were no significant differences in the DEAD/LIVE bacterial ratio, that was quantified and confirmed with COMSTAT image analysis, after treatment of the *A. baumannii* 7789 and *P. aeruginosa* PAO1 biofilms with Compounds 1, 4, 6 and the comparator compound CC-1 compared to the untreated and ethanol equivalent (blank) controls.

Figure 7 provides cell membrane permeabilization data when Compounds 1, 4 and 6 were applied to planktonic cells of *E. coli* IR57, *P. aeruginosa* PAO1 and *S. aureus* (1004A; MRSA) compared to the untreated control and positive control of 70% Isopropanol. The Gram-negative strains (*E. coli* IR57 and *P. aeruginosa* PAO1) only showed significant increases in cell permeabilization when treated with compounds 1 and 4 at concentration $\geq 512 \mu\text{g/mL}$ unlike the Gram-positive strain (MRSA 1004A) which demonstrated significant cell permeabilization at concentrations as low as $32 \mu\text{g/mL}$.

Figure 8 provides representative images showing the induction of NETosis and necrosis of neutrophils treated with six concentrations of Compound 4 at two times after treatment, 3 and 6 hours. NETosis/necrosis was first observed at 3 hours after treatment in the two highest concentrations (50 and $500 \mu\text{M}$) and at 6 hours in the four lower concentrations. At $500 \mu\text{M}$ the chromatin in the cells is condensed and indicates necrosis has occurred. At $50 \mu\text{M}$ the chromatin is more diffuse indicating that NETosis has occurred.

Figure 9 provides a graph with mean values and standard deviations for *in vitro* release of human defence peptide LL-37 from neutrophils at 3 hours after application of four therapeutically relevant concentrations of Compound 4. Release of LL-37 from neutrophils increased in a concentration dependent manner.

Examples

The compounds of the present invention may be obtained by isolation from a plant or plant part, or by derivatisation of the isolated compound, or by derivatisation of a related compound. Isolation procedures and derivatisation procedures may be found in WO 2007/070985 and WO2014/169356.

5 **Example 1: Epoxy-tigilanes have no direct antibiotic activity against Gram-negative bacteria in planktonic culture systems.**

The effects of five epoxytiglienones (Compounds 1, 2, 3, 4 and 6) and two comparator
comparator compounds (epoxytiglienones having longer carbon chains at the C12
position) on six human pathogenic bacteria (two Gram-positive and four Gram-negative
10 species) were measured in conventional planktonic culture systems. A minimum
inhibitory concentrations (MIC) assay (the standard assay employed to define and
quantify antibiotic activity) was performed to determine antibacterial activity for each
epoxytiglienone against each bacterial strain.

The Gram-positive bacteria used in this study were methicillin resistant *Staphylococcus*
15 *aureus* (MRSA) 1004A and *Streptococcus pyogenes*. The Gram-negative bacteria used
in this study were *Pseudomonas aeruginosa* PAO1, *Escherichia coli* IR57 (V7),
Klebsiella pneumoniae, and *Acinetobacter baumannii* 7789 (V19). MIC assays were
performed on planktonic cultures of each bacteria grown in Mueller-Hinton broth using
the standard broth dilution method described by Jorgensen *et al.* (1999) using
20 epoxytiglienones dissolved in ethanol. The results are shown in Table 1 below.

Table 1. Minimum inhibitory concentrations ($\mu\text{g/mL}$) determined for Compounds 1, 2,
3, 4 and 6 and comparator compounds CC-1 and CC-2 against two Gram-positive
bacteria and four Gram-negative bacteria. C = confluent growth, i.e. no MIC could be
determined at any concentration tested. Data presented represent the mean of three
25 replicate tests.

Bacteria	Compound number					Comparator*	
	1	2	3	4	6	CC-1	CC-2
Gram-positive species							

- 31 -

<i>S. aureus</i>	256	256	256	512	C	8	16
<i>S. pyogenes</i>	128	128	128	256	C	8	8
Gram-negative species							
<i>E. coli</i>	C	C	C	C	C	C	C
<i>P. aeruginosa</i>	C	C	C	C	C	C	C
<i>K. pneumoniae</i>	C	C	C	C	C	C	C
<i>A. baumannii</i>	C	C	C	C	C	C	C

* Comparator compounds are epoxytiglienones having longer carbon chains at the C12 position. CC-1 is 12-(2,4-decadienoyl)-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one and CC-2 12R-(2,4-decadienoyl)-13R-(2-methylbutanoyl)-6,7-epoxy-4R,5R,9S,12R,13R,20-hexahydroxy-1-tiglien-3-one.

- 5 The results demonstrated that the direct antimicrobial properties of the epoxytiglienones in planktonic culture were restricted to the Gram-positive bacteria. No MIC value could be determined for any of the compounds with any of the Gram-negative bacteria tested, all of which were unaffected (with confluent growth of bacteria at concentrations of the test compounds exceeding 512 µg/mL).

10 **Example 2: Epoxytiglienones disrupt established biofilm of Gram-negative *E. coli***

The effects of administration of Compounds 1, 4 and 6 and comparator compound CC-1 on disruption of established *in vitro* biofilms of *Escherichia coli* (*E. coli*) was investigated using methods described in Powell *et al.* (2018).

- 15 Biofilms of *E. coli* IR57 were grown on 96-well glass-bottomed plates in Mueller-Hinton (MH) broth for 24 h, before 50% of the supernatant was replaced with fresh MH broth with or without the epoxytiglienone compound added to a final concentration of 256 µg/mL (vehicle is ethanol). An ethanol equivalent blank was used as a further control treatment. The plates were then incubated for a further 24 h at 37°C.

Confocal laser scanning microscopy (CLSM) was then used to image the biofilms, following staining with Live/Dead® BacLight stain with phosphate buffered saline (PBS) added to each well before imaging with z-stack CLSM. The resultant images were analysed by COMSTAT software to produce measurements of (i) biofilm biomass
5 volume or bio-volume ($\mu\text{m}^3/\mu\text{m}^2$) and (ii) DEAD/LIVE bacterial ratio.

CLSM demonstrated marked differences between treatments in the distribution of the bacteria in the biofilms. Compounds 1, 4 and 6 induced significant changes the distribution and density of bacteria in the biofilm compared to the untreated and ethanol equivalent (blank) controls (Figure 1); this was quantified and confirmed with
10 COMSTAT image analysis. Significant decreases in bio-volume were evident in biofilms treated with Compound 1 ($p < 0.05$; Figure 2A). The comparator compound CC-1 had no apparent effect on biofilm distribution and density or bio-volume. Interestingly, there were no significant differences between any of the epoxytiglenones and the two control treatments in the DEAD/LIVE bacterial ratios (Figure 2B),
15 demonstrating the effect of the epoxytiglenones on biofilm density and bio-volume in these experiments was unrelated to direct antibiotic activity.

Example 3: Epoxytiglenones disrupt the extracellular matrix and increase particle diffusion through an established biofilm of the Gram-negative *E. coli* bacteria

The effect of three epoxytiglenones (Compounds 1, 4 and 6) on the assembly and
20 permeability of established biofilms of *E. coli* was assessed using multiple particle tracking (MPT). MPT is a recently described technique, allowing simultaneous tracking of micron-size particles through biofilms using microscopy, from which the diffusion-based parameters of embedded particles within the extracellular polymeric matrix (EPS) of the biofilm can be determined (Cao *et al.* 2016). MPT measurements
25 also allows for calculation of micro-rheological properties of the biofilm structure following treatment with test compounds.

E. coli biofilms were established and epoxytiglenone and control treatments applied as described in Example 2 above. Twenty-four hours after application of the treatments, the biofilms were stained with SYTO9® and 0.0025% of negatively-charged,
30 carboxylate-modified FluoSpheres® (200 nm) added onto the biofilms and incubated

for a further 2 hours. FluoSphere® particle movements within the biofilms were then captured on video using epifluorescence microscopy with a high frame-rate camera (33 ms). Particle trajectories were tracked using ImageJ software (Mosaic) before calculation of three parameters of the 200nm FluoSphere® particles: (i) ensemble
 5 diffusion coefficient (Deff), (ii) ensemble mean squared displacement (MSD) and (iii) creep compliance (a measure of resistance to mechanical deformation derived from the MSD vs lag time). Each treatment was replicated three times.

All three epoxytiglenone compounds that were tested increased particle diffusion through the established *E. coli* biofilms by between 80 and 420 times compared to the
 10 control treatments (Table 2).

Table 2. Diffusion coefficients for 200 nm negatively-charged, carboxylate-modified FluoSpheres® particles in *E. coli* IR57 biofilms treated with epoxytiglenones.

Compounds	Diffusion coefficient (Deff)
	(cm ² . s ⁻¹ x10 ⁻⁹) ± standard deviation
Compound 4	0.8449 ± 0.1250
Compound 1	0.6018 ± 0.0901
Compound 6	0.1607 ± 0.0104
No treatment controls	0.002 ± 0.0014
Ethanol only control	0.0155 ± 0.0052

Significant disruption and alteration to the structure of the *E. coli* biofilm matrix by the
 15 epoxytiglenones were illustrated by substantially higher mean squared displacement values of the FluoSphere® particles following treatment (Figure 3) and decreased resistance to mechanical deformation in the treated biofilms was evident in the increase of the creep compliance of the epoxytiglenone treated biofilms versus controls (Figure 4).

Example 4: Epoxytiglienones significantly decrease biomass in established biofilms of two other Gram-negative pathogens *P. aeruginosa* and *A. baumannii*

The effects of the epoxytiglienones (Compounds 1, 4, 6 and the comparator compound CC-1) on disruption of established *in vitro* biofilms were further investigated with two other species of Gram-negative bacteria (*Pseudomonas aeruginosa* PA01 and *Acinetobacter baumannii* 7789) using methods described in Example 2 above. Cell permeabilisation following Compound 1, 4 and 6 treatment was also determined using SYTOX™ Green Nucleic Acid Stain with untreated control and positive control of 70% isopropanol.

Consistent with the results of the *E. coli* study, the epoxytiglienone compounds significantly changed the distribution and density of bacteria in the *A. baumannii* biofilms compared to the untreated and ethanol equivalent (blank) controls (Figure 5A). This was also reflected in their effects as measured by bio-volume for *A. baumannii* biofilms (Figure 6A), where significant reductions were seen for Compounds 1, 4 and CC-1. For *P. aeruginosa* only Compound 4 had a significant effect on biofilm distribution (Figure 5B) and on bio-volume compared to other treatments (Figure 6B). DEAD/LIVE bacterial ratios were also assessed and, as in Example 2 with *E. coli*, there were no differences between treatments in either *A. baumannii* or *P. aeruginosa* (Figure 6C & D). Cell permeabilization studies on *E. coli* and *P. aeruginosa* bacteria (Figure 7) only showed significant increases in permeabilization when treated with Compounds 1 and 4 at concentrations $\geq 512 \mu\text{g/mL}$, which is greater than the compound concentration used in the biofilm disruption assays ($256 \mu\text{g/mL}$). This further confirms that the effects of the epoxytiglienones on biofilm density and bio-volume in these experiments was unrelated to any direct, antibiotic activity in killing bacteria within the biofilms.

Example 5: Epoxytiglienones resolve established biofilm *in vivo* in a mouse model of chronic biofilm infection

The effects of administration of Compound 4 on biofilm infection *in vivo* was studied in a diabetic murine model of chronic biofilm infection (Zhao *et al.* 2010). In this study we used methods described by Dhall *et al.* (2014) in which bacterial biofilm infections develop spontaneously following wound creation.

- 35 -

Briefly, *db/db* diabetic mice (> 6 months old) were housed for 4-5 weeks in non-sterile conditions prior to the creation of wound on the back of each mouse with a 6 mm diameter excisional punch biopsy. Mice were then administered a catalase inhibitor intra-peritoneally (1 g/kg aminotriazole) and a glutathione peroxidase (GPx) inhibitor
5 topically around the edge of the wound site (1 g/kg mercaptosuccinic acid) prior to dressing with Tegaderm. After 24 h, biofilms were evident at the wound site on all mice. The mice were divided into two groups (7 mice per group) and treated with either Compound 4 (0.3 mg/mL in a hydrogel vehicle) or a Control treatment (hydrogel vehicle only). The Tegaderm dressing was then replaced. Two further
10 administrations of Compound 4 or the Control (vehicle only) were performed at 8 and 15 days. No antibiotics or other antibacterial treatments were applied during the study.

The presence of biofilm and the surface area of the wound were assessed over the course of the study. Biopsies were taken from Compound 4-treated and vehicle-only (control) treated mice between day 21 and 28 post wounding, for histological and
15 histochemical analysis of the wound site. Only 1 of the 7 wounds in the Control group healed within this period, with the presence of a fibrinous slough and bacterial infiltration of the wound site clearly evident. In epoxytiglicenone-treated infected wounds, complete wound healing was evident (with complete re-epithelialisation, resolution of inflammation and an absence of bacteria within the dermis) in 6 of the 7
20 mice treated with Compound 4.

Example 6: Epoxytiglicenone treatment of human adult keratinocytes and fibroblasts *in vitro* induces upregulation of chemokines/cytokines involved in neutrophil recruitment

In addition to disrupting the structure of biofilms comprising Gram-negative bacteria, effects of epoxytiglicenones on regulation of genes involved in host response to bacterial
25 infection were investigated in microarray studies with human adult epidermal keratinocytes (HEKa) and human adult dermal fibroblasts (HDF) *in vitro*.

For these studies, HEKa were cultured in EpiLife® Medium supplemented with Supplement S7 (both from Life Technologies, Carlsbad, CA, USA), while HDF were
30 cultured in Medium 106 supplemented with LSGS (Low Serum Growth Supplement)

- 36 -

(both from Life Technologies). Cells were then treated with vehicle or 170 nM Compound 4 for 0, 0.5, 1, 2, 4, 8, 24, 48 and 72 hours. RNA was extracted using a Qiagen RNeasy mini kit and biotinylated using an Illumina TotalPrep RNA Amplification Kit (Ambion, Austin, TX, USA). Labelled RNA was hybridised to
5 HumanHT-12 v4 BeadChip Arrays (Illumina Inc, San Diego, CA, USA) and scanned according to standard Illumina protocols. Data were extracted in GenomeStudio (Illumina) using default analysis settings and no normalisation method. Resulting data were imported into GeneSpring GX (Agilent, Santa Clara, CA, USA). Expression values were normalised using quantile normalisation with default settings.

10 In both HDF and HEKa *in vitro*, treatment with Compound 4 (170 nM) induced significant upregulation of two critical chemokines/cytokines involved in PMNL recruitment (*IL8*, *CXCL1*) within 2 to 4 hours. In HEKa, Compound 4 significantly upregulated production of host defence peptides (*DEFB2*, *DEFB3*, *DEFB4*, *RNASE7*) from 8 to 72 hours post treatment.

15 **Example 7: Epoxytiglicenone treatment induces NETosis/necrosis and release of the antimicrobial peptide cathelicidin LL-37 *in vitro* from isolated human neutrophils**

Neutrophils are the most abundant leukocyte in the blood and constitute the first line of host defense against infectious pathogens. Central to their function is their ability to be recruited to sites of infection, to recognize microbes, and then be activated to kill
20 pathogens through a combination of phagocytotic and cytotoxic mechanisms. Amongst these mechanisms used by neutrophils to kill pathogens are: (i) the production of reactive oxygen species, (ii) the expulsion of their nuclear chromatin contents (coated with histones, proteases and granular and cytosolic proteins) to immobilise and catch pathogens (the process of NETosis), and (iii) the release of antimicrobial peptides.

25 Subsequent to identifying the effect of epoxytiglicenone compounds in upregulating production in HDF and HEKa of chemokines and cytokines involved in neutrophil recruitment (see Example 6 above), the effect of Compound 4 on two aspects of neutrophil function, induction of NETosis/necrosis and antimicrobial peptide release, were examined.

- 37 -

For these assays, neutrophils were isolated from fresh blood of a healthy human donor by lysis of a red blood cell pellet that had been obtained by Ficoll-Paque sedimentation. The neutrophils ($\sim 4 \times 10^6$ cells/mL) were incubated with 10 μ g/mL dihydroethidium (DHE) (Sigma-Aldrich) in complete culture medium at 37°C for 15 min alongside an aliquot of unstained cells to be tested as unstained control.

NETosis/necrosis assays followed methods fully described by Brinkmann *et al.* 2010. Isolated neutrophils were plated into 96-well plates (RPMI1640, 10% FCS) and incubated with a 1:50,000 dilution of Hoechst (10 mg/mL) and Sytox® Green (5 mM). Compound 4 was added at six concentrations (0.005, 0.05, 0.5, 5, 50, 500 μ M) and the cells incubated at 37°C, 5% CO₂. Vehicle treated controls were included in all assays and there were three replicates of each assay. Hoechst/Sytox® Green fluorescence images were recorded for each well at 3 and 6 hours.

NETosis and necrosis of neutrophils was initially observed at 3 hours after treatment in the two highest concentrations (50 μ M and 500 μ M) of Compound 4 and was evident at 6 hours in the four lower concentrations (0.005 μ M – 5 μ M; Figure 8).

To examine antimicrobial peptide release by neutrophils in response to treatment with Compound 4, isolated neutrophils were incubated with either vehicle or Compound 4 at the four concentrations (62.5, 125, 250 and 500 μ M). At 3 h, cell culture supernatants were removed and tested for LL-37 content using a LL-37-directed ELISA kit (Hycult Biotech). ELISA readings were normalised to vehicle only controls to determine fold increases in LL-37 release.

Release of LL-37 from neutrophils at 3 hours after treatment with Compound 4 increased in a concentration dependent manner (Figure 9). At concentrations of Compound 4 of between 125 and 500 μ M, LL-37 release was 3- to 5- fold higher than the control treatment (Figure 9).

The data from this Example show that at therapeutically relevant concentrations *in vitro*, Compound 4 induces suicidal neutrophil NETosis transitions to necrosis, leading to the release of the potent antimicrobial defence peptide LL-37.

The data from Examples 6 and 7 above also demonstrate that in addition to their direct effects in disrupting the structure of Gram-negative biofilms, epoxytiglenones can also induce local innate immune responses in both migratory/resident myeloid cells (e.g. neutrophils) and in dermal and stromal cell types. Such responses suggest that the
5 effect of epoxytiglenones alone may be adequate to resolve many biofilm infections without the need for conventional antibiotics.

References

It is to be understood that, if any prior art publication is referred to herein, such reference does not constitute an admission that the publication forms a part of the common general knowledge in the art, in Australia or any other country.


- 5 Brinkmann V *et al.* 2010. Neutrophil extracellular traps: How to generate and visualize. *Them. J. Vis. Exp.* 36: e 1724.
- Cao, H. *et al.* (2016) Revealing region-specific biofilm viscoelastic properties by means of a micro-rheological approach. *npj Biofilms and Microbiomes*, 2(1), pp. 1–7.
- Cepas V *et al.* 2019. Relationship between biofilm formation and antimicrobial
10 resistance in Gram-negative bacteria. *Microb. Drug Resist.* 25: 72-79.
- Dhall S *et al.* 2014. “Generating and Reversing Chronic Wounds in Diabetic Mice by Manipulating Wound Redox Parameters.” *J. Diabetes Res.* 562625.
- Doi *et al.* 2017. Gram-negative bacterial infections: Research priorities, accomplishments and future directions. *Clin Infect Dis.* 64 (S1): S30-S35
- 15 Fleming D & Rumbaugh KP 2017. Approaches to dispersing medical biofilms. *Microorganisms* 2017 5, 15.
- Gunn JS *et al.* 2016. What’s on the outside matters: the role of extracellular polymeric substance of Gram-negative biofilms in evading host immunity and as a target for therapeutic intervention. *J. Biol. Chem.* 291: 12538-12546.
- 20 Hoiby *et al.* 2010. Antibiotic resistance of bacterial biofilms. *Int. J. Antimicrob. Agents* 35: 322-332.
- Hoiby N *et al.* 2015. ESCMID guideline for the diagnosis and treatment of biofilm infections. *Clin. Microbiol. Infect.* 21 (Suppl. 1): S1-S25.
- Ho J *et al.* 2010. Multiresistant Gram-negative infections: a global perspective. *Curr.*
25 *Opin. Infect. Dis.* 23: 546-53.
- Jorgensen JJH *et al.* 1999. Antibacterial susceptibility tests: dilution and disk diffusion methods. In Murray PR *et al.* (Eds.), *Manual of Clinical Microbiology* (pp 1526-1543). Washington, DC: ASM Press.
- Koo H *et al.* 2017. Targeting microbial biofilms: current and prospective therapeutic
30 strategies. *Nat. Rev. Microbiol.* 15: 740-755.

Powell *et al.* 2018 Targeted disruption of the extracellular polymeric network of *Pseudomonas aeruginosa* biofilms by alginate oligosaccharides. *NPJ Biofilms Microbiomes*. 4: 13.

5 Zhao *et al.* 2010. Delayed wound healing in diabetic (db/db) mice with *Pseudomonas aeruginosa* biofilm challenge: a model for the study of chronic wounds. *Wound Repair Regen*. 18: 467-477.

Claims

1. An *ex vivo* method of dispersing a biofilm comprising Gram-negative bacteria comprising exposing the biofilm to an epoxytiglenone compound for formula (II):



wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

R₂ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₃ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;

R₆ is selected from hydrogen and -C(O)C₁₋₆alkyl;

R₇ is hydroxy; and

R₈ is selected from hydrogen or C₁₋₆alkyl; or a pharmaceutically acceptable salt thereof.

2. The method according to claim 1 wherein R₁ is C₁₋₃ alkyl.
3. The method according to claim 1 or claim 2, wherein R₂ is selected from -OC(O)C₃₋₆alkyl and -OC(O)C₃₋₆alkenyl.
4. The method according to any one of claims 1 to 3, wherein R₃ is selected from -OC(O)C₃₋₆alkyl, -OC(O)C₃₋₆alkenyl and -OC(O)C₃₋₆alkynyl.
5. The method according to any one of claims 1 to 4, wherein R₄ and R₅ are each methyl.
6. The method according to any one of claims 1 to 5, wherein R₆ is selected from hydrogen, -C(O)CH₃, -C(O)CH₂CH₃, -C(O)CH(CH₃)₂ or -C(O)CH₂CH₂CH₃.
7. The method according to any one of claims 1 to 6, wherein R₈ is C₁₋₃alkyl.

8. The method according to any one of claims 1 to 7, wherein the alkyl or alkenyl group of R₂ and/or R₃ are branched alkyl or branched alkenyl groups.
9. The method according to any one of claims 1 to 7, wherein the alkyl or alkenyl group of R₂ and/or R₃ are linear alkyl or linear alkenyl groups.
10. The method according to claim 1, wherein the compound of formula (II) is selected from:

12-tigloyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 1);
12,13-di-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 2);
12-hexanoyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 3);
12,13-dihexanoyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 4);
12-tigloyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13-pentahydroxy-20-acetyloxy-1-tiglien-3-one (Compound 5);
12-propanoyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 6);
12,13-ditigloyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 7);
12-(2-methylbutanoyl)-13-tigloyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 8);
12-butanoyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 9);
12-(3,3-dimethylbut-2-enoyl)-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 10);
12-hex-2,4-dienoyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 11);
12-tigloyl-13-(2-methylpropanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 12);
12-but-2-enoyl-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-

tiglien-3-one (Compound 13);

12-tigloyl-13-butanoyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 14);

12,13-dibutanoyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 15);

12,13-dipentanoyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 16);

12,13-di-(2*E*,4*E*)-hexa-2,4-dienoyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tiglien-3-one (Compound 17);

12-(2-methylprop-2-enoyl)-13-(2-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tigliaen-3-one (Compound 19);

12,13-di-heptanoyl-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tigliaen-3-one (Compound 20); and

12,13-di-(3-methylbutanoyl)-6,7-epoxy-4,5,9,12,13,20-hexahydroxy-1-tigliaen-3-one (Compound 21);


or a pharmaceutically acceptable salt thereof.

11. The method according to any one of claims 1 to 10, wherein the biofilm comprising Gram-negative bacteria comprises at least one Gram-negative bacteria selected from *Pseudomonas* species, *Acinetobacter* species, *Aeromonas* species, *Bacteroides* species, *Bordetella* species, *Borrelia* species, *Burkholderia* species, *Citrobacter* species, *Compylobacter* species, *Escherichia* species, *Enterobacter* species, *Flavobacterium* species, *Fusobacterium* species, *Klebsiella* species, *Leptospira* species, *Neisseria* species, *Helicobacter* species, *Hemophilus* species, *Legionella* species, *Moraxella* species, *Yersinia* species, *Oligella* species, *Pantoea* Species, *Porphyromonas* species, *Prevotella* species, *Proteus* species, *Raoutella* species, *Salmonella* species, *Serratia* species, *Shigella* species, *Sphingomonas* species, *Stenotrophomonas* species, *Treponema* species, *Veillonella* species and *Vibrio* species.

12. The method according to claim 11, wherein the Gram-negative bacteria is selected from *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, *Escherichia coli*, *Klebsiella pneumoniae*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Hemophilus influenzae*, *Legionella pneumophila*, *Yersinia pestis*, *Yersinia enterocolitica*, *Salmonella*

enterica, *Salmonella bongori*, *Proteus mirabilis*, *Enterobacter cloacae*, *Serratia marcescens*, *Bacteroides fragilis*, *Fusobacterium necrophorum*, *Burkholderia cepacia* and *Prevotella intermedia*.

13. Use of an epoxytiglienone compound of formula (II):



wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

R₂ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₃ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;

R₆ is selected from hydrogen and -C(O)C₁₋₆alkyl;

R₇ is hydroxy; and

R₈ is selected from hydrogen and C₁₋₆alkyl; or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the treatment of a bacterial infection comprising a biofilm comprising Gram-negative bacteria wherein the compound of formula (II) is to be administered locally to the bacterial infection.

14. The use according to claim 13, wherein the compound of formula (II) is to be administered topically.


15. The use according to claim 13 or claim 14, wherein the infection is a post-surgical infection or an infection at the site of insertion of a medical device or implantation of an implant.

16. The use according to any one of claims 13 to 15, wherein the bacterial infection

is a chronic infection.

17. The use according to any one of claims 13 to 16, wherein the compound of formula (II) is to be administered in combination with an antibiotic to which the Gram-negative bacteria is susceptible when in planktonic state.

18. An *ex vivo* method of preventing a biofilm comprising Gram-negative bacteria forming on a medical device or dispersing a biofilm comprising Gram-negative bacteria on a medical device, said method comprising applying an epoxytiglenone compound of formula (II) to the medical device, the compound of formula (II) comprising



wherein

R₁ is selected from hydrogen and C₁₋₆alkyl;

R₂ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₃ is selected from -OC(O)C₁₋₇alkyl, -OC(O)C₂₋₇alkenyl and -OC(O)C₂₋₇alkynyl;

R₄ and R₅ are independently selected from hydrogen and C₁₋₆alkyl;

R₆ is selected from hydrogen and -C(O)C₁₋₆alkyl;

R₇ is hydroxy; and

R₈ is selected from hydrogen and C₁₋₆alkyl; or a pharmaceutically acceptable salt thereof.

19. A method according to claim 18, wherein the medical device is a surgical instrument, catheter or medical implant.

20. A method according to claim 18 or claim 19, wherein the medical device is a catheter.

21. A method according to claim 19, wherein the medical device is a dental implant.