Description

Technical Field

The present invention relates to the production of recombinant proteins in yeast. More particularly, the present invention is directed to an improved α-factor expression system which provides for the secretion of heterologous proteins from yeast host cells.

Background

Kurjan et al. (1982) Cell 30:933-943 discloses the first cloning and sequencing of a gene encoding a yeast α-factor precursor gene. Kurjan et al., U.S. Patent No. 4,546,082, also reports the cloning of this gene, and suggests that the α-factor leader sequence can be employed to direct the secretion of heterologous proteins in yeast. The patent, however, does not contain data which would indicate that the patentees ever successfully employed the α-factor leader to express and secrete a heterologous protein in yeast.

EPO Publication No. 116,201 discloses the first successful application of the α-factor leader to direct the expression and secretion of an heterologous protein, epidermal growth factor, from a transformed yeast. Subsequent to this work, there have been additional reports of the expression of heterologous proteins in yeast employing the α-factor leader. See, e.g., Elliott et al. (1983) Proc. Nat'l Acad. Sci. USA 80:7080-7084; Bitter et al. (1984) Proc. Nat'l Acad. Sci. USA 81:5330-5334; Smith et al. (1985) Science 229:1219-1229; EPO Publication Nos. 114,695; 123,228; 123,294; 123,544; 128,773; and 206,783; and WO86/01515, in the name of the present applicants. See also Brake et al. in Protein Transport and Secretion, p.103 (J.M. Gething ed. 1984).

The expression systems in the above reports produce a full-length α-factor leader fused to a heterologous protein. While the above work demonstrates that the α-factor expression system is widely useful, it is not generally predictable prior to performing the experiment whether a particular heterologous protein will be successfully secreted, processed and biologically active. See, e.g., EPO 206,783, supra, pp. 2-5; Rothblatt et al. (1987) EMBO J. 6:3455-3463; V.L. MacKay, "Secretion of Heterologous Proteins in Yeast" (in press).

There have been several reports, based on unpublished data, that deletions from the "pro" region of the α-factor leader (between the signal peptide and the first spacer) causes substantial declines in the amount of non-yeast protein secreted by yeast transformed by the heterologous constructs. Sidhu et al. (1987) Gene 54:175-184, reports that yeast acid phosphatase (PHO5) is secreted into the medium from a heterologous construct employing both a full-length α-factor leader, and a truncated a-factor leader, but that expression levels are 3-4X less than for the PHO5 gene under the control of its homologous leader. It has also been reported that deletions in the prepro- α-factor precursor gene results in substantial declines in secretion of the native α-factor peptide. See, e.g., V.L. MacKay, supra; Rothblatt et al., supra.

A need exists, therefore, to improve the α-factor expression system, particularly for applications to non-yeast proteins that are not efficiently produced by current α-factor expression constructs.

Summary of the Invention

It has been surprisingly discovered that a truncated form of the α-factor leader sequence can efficiently direct the expression and secretion of heterologous polypeptides in yeast. Particularly surprising is the discovery that truncated α-factor leader sequences can substantially improve the efficiency of expression of such heterologous proteins in relation to expression systems using the full-length α-factor leader; i.e., higher levels of correct N-terminal processing, secretion of heterologous proteins wherein a greater percentage of the molecules are biologically active, etc. These results are particularly surprising in view of reports that deletions from the leader sequence of the α-factor precursor result in decreased levels of secretion of active α-factor.

The present invention provides, therefore, for alternative α-factor-based expression constructs, which are particularly useful for the expression of heterologous polypeptides which are either inefficiently expressed by full-length α-factor leader constructs, or are not expressed at all by such full-length constructs.

In one embodiment, the present invention is directed to a yeast cell comprising a DNA construct that provides for the expression and secretion of a non-yeast protein, said DNA construct comprising a coding sequence under the control of yeast-recognized transcription initiation and termination sequences, said coding sequence encoding a precursor polypeptide comprised of a leader sequence and said non-yeast

protein linked by a processing site that provides for the cleavage of said non-yeast protein from said precursor polypeptide, wherein said leader sequence is about 25 to about 50 N-terminal residues of said precursor polypeptide and comprises the signal peptide of a Saccharomyces or Kluveromyces α-factor precursor and a single glycosylation site.

In another embodiment, the present invention is directed to a double-stranded DNA molecule comprising a region encoding a precursor polypeptide secretable by a yeast host, said region, with reference to one of the DNA strands, having the structure:

5'-AF-CHO-X_n-S-Gene*-3'

wherein AF-CHO-X_n comprises a single functional glycosylation site in which :

AF	encodes a Saccharomyces or Kluveromyces α-factor signal peptide; CHO encodes a glycosylation site;
X_n	encodes a polypeptide of n amino acids in length that does not contain a glycosylation site or a processing site that provides for cleavage of said precursor polypeptide in vivo by yeast;
n	is an integer from 0 to about 30;
	and wherein
Gene*	encodes a non-yeast protein; and
S	encodes a processing site that provides for cleavage of said precursor polypeptide.

The present invention is also directed to methods of employing the above cells and DNA constructs to produce recombinant proteins, as well as the compositions of recombinant proteins produced by the above methods. Other embodiments will also be readily apparent to those of ordinary skill in the art.

Description of the Figures

Figure 1 is a flow diagram showing the construction of pYBCA5, and both the nucleotide and amino acid sequences of the synthetic proinsulin gene employed in Example I. The various synthetic oligonucleotides used in construct are delineated by black dots. Arrows above the sequence show the beginning and end of the B, C and A proinsulin chains. The boxes mark the dibasic processing sites.

Figure 2 is the nucleotide and amino acid sequence of a synthetic oligonucleotide encoding a modified α-factor leader and the first 13 amino acids of the proinsulin gene used to construct pYGAI3 in Example I. The modified a-factor leader has had the glycosylation sites removed by changing the codons for ^{Asn}23,57,67 to encode Gln (boxed). The arrow denotes the junction between the sequence encoding the KEX2 endopeptidase site and the N-terminus of human proinsulin.

Figure 3 shows the synthetic gene of fragment pYGAIC3 encoding the proinsulin analog, where a KEX2 endopeptidase site has replaced the C peptide (boxed). The synthetic 133 bp fragment referred to in Example II is defined by the vertical and horizontal lines through the nucleotide sequence.

Figure 4 is a restriction map of yeast shuttle vector pAB24.

Figure 5 shows the DNA sequence of the synthetic gene encoding IGF1 described in Example III.

Figure 6 is a restriction map of pYLUIGF1-55, an expression vector described in Example III encoding IGF1 under the control of a truncated α-factor leader.

Figure 7 is a restriction map of pYLUIGF1-24, an expression vector described in Example III encoding IGF1 under the control of a full-length α-factor leader with three glycosylation sites.

Detailed Description

The practice of the present invention will employ, unless otherwise indicated, conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are fully explained in the literature. See, e.g., Maniatis, Fritsch & Sambrook, Molecular Cloning: A Laboratory Manual (1982); DNA Cloning, Vols. I & II (D.N. Glover, ed. 1985); Oligonucleotide Synthesis (M.J. Gait, ed. 1984); Transcription and Translation (B.D. Hames & S.J. Higgins, eds. 1984); Immobilized Cells and Enzymes (IRL Press, 1986); B. Perbal, A Practical Guide to Molecular Cloning (1984).

In describing the present invention, the following terminology will be used in accordance with the definitions set out below.

A "replicon" is any genetic element (e.g., plasmid, cosmid, chromosome, virus) that functions as an autonomous unit of DNA replication in vivo--i.e., capable of replication under its own control.

A "vector" is a replicon such as a plasmid, phage, or cosmid to which another DNA segment may be attached so as to bring about the replication of the attached segment.

A "double-stranded DNA molecule" refers to the polymeric form of deoxyribonucleotides (adenine, guanine, thymidine, or cytosine) in its normal, double-stranded helix. This term refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, inter alia, in linear DNA molecules (e.g., restriction fragments), viruses, plasmids, and chromosomes. In discussing the structure of a particular double-stranded DNA molecule, sequences will be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the nontranscribed strand of DNA, i.e., the strand having a sequence homologous to the mRNA produced from a particular coding sequence.

A DNA "coding sequence" is DNA sequence which can be transcribed and translated into a polypeptide in vivo when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by and include the translation start codon at the 5' (amino) terminus, and a translation stop codon at the 3' (carboxy) terminus. A coding sequence can include, but is not limited to, procaryotic DNA sequences, viral DNA sequences, cDNA or genomic DNA sequences from eucaryotic sources (e.g., mammalian), and even synthetic DNA sequences.

"Yeast-recognized transcription initiation and termination sequences" refer to DNA regulatory regions which flank a coding sequence and are responsible for the transcription in yeast of an mRNA homologous to the coding sequence which can then be translated into the polypeptide encoded by the coding sequence. Transcription initiation sequences include yeast promoter sequences, which are DNA regulatory sequences capable of binding yeast RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For the purposes of defining the present invention, the promoter sequence is bounded (and excludes) at its 3' terminus by the translation start codon of a coding sequence and extends upstream (5' direction) to include the minimum number of nucleotides or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined by mapping with nuclease Sl), as well as protein-binding domains (consensus sequences) responsible for the binding of the yeast RNA polymerase. Promoters useful in the present invention include the wild-type α-factor promoter, as well as other yeast promoters. Particularly preferred are promoters involved with the enzymes in the glycolytic pathway, e.g., phosphoglucoisomerase, phosphofructokinase, phosphotrioseisomerase, phosphoglucomutase, enolase, pyruvic kinase, glyceraldehyde-3-phosphate dehydrogenase, alcohol dehydrogenase, as well as hybrids of these promoters. See, e.g., EPO Publication Nos. 120,551; 164,556. Transcription initiation sequences can also include other regulatory regions responsible for promoter regulation or enhancement. In like manner, a transcription terminator sequence located 3' to the translation stop codon can be either the wild-type a-factor transcription termination sequence, or another yeast-recognized termination sequence, such as those from the genes for the above glycolytic enzymes.

A coding sequence is "under the control" of transcription initiation and termination sequences when RNA polymerase binds the transcription initiation sequence and transcribes the coding sequence into mRNA terminating at the transcription termination sequence, and the mRNA is then translated into the polypeptide encoded by the coding sequence (i.e., "expression"). The precursor polypeptide encoded by the coding sequences of the present invention is "secreted" when at least a portion (usually the non-yeast protein in the absence of the leader sequence) is transported extracellularly where it is found in the cell growth medium. Usually, only the portion of precursor protein downstream from the leader sequence is secreted, and this downstream portion may also be subjected to additional processing during secretion, such as proteolytic cleavage, glycosylation, folding, disulfide bond formation, etc.

A cell has been "transformed" by exogenous DNA when such exogenous DNA has been introduced inside the cell wall. Exogenous DNA may or may not be integrated (covalently linked) to chromosomal DNA making up the genome of the cell. The exogenous DNA may be maintained extrachromosomally on a replicon such as a plasmid. When the exogenous DNA has become integrated to the chromosome, it is inherited by daughter cells through chromosome replication. A cell which has been transformed by exogenous DNA which is integrated into the chromosome is referred to as a "stably" transformed cell. A "clone" or "clonal population" is a population of cells derived from a single cell or common ancestor by mitosis.

Two DNA sequences are "substantially homologous" when at least about 60% (preferably at least about 75%, and most preferably at least about 90%) of the nucleotides match over a defined length of the molecules. Sequences that are substantially homologous can be identified in a Southern hybridization experiment under conditions of a selected stringency as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Maniatis et al., supra; DNA Cloning, supra; Nucleic Acid Hybridization, supra.

A "heterologous region" of a DNA molecule is an identifiable segment of DNA within a larger DNA molecule that is not found in association with the larger molecule in nature. Thus, when the heterologous region encodes a mammalian protein, the heterologous region will usually be flanked by DNA that does not flank the mammalian DNA sequence in the genome of the source organism. Another example of a heterologous coding sequence is a construct where the coding sequence itself is not found in nature (e.g., a cDNA where the genomic coding sequence contains introns, or synthetic sequences having codons different from organisms which encode the same or similar protein). Allelic variations or naturally occurring mutational events do not give rise to a "heterologous" region of DNA as used herein.

As used herein, "yeast" includes ascosporogenous yeasts (Endomycetales), basidiosporogenous yeasts and yeast belonging to the Fungi imperfecti (Blastomycetes). The ascosporogenous yeasts are divided into two families, Spermophthoraceae and Saccharomycetaceae. The latter is comprised of four subfamilies, Schizosaccharomycoideae (e.g., genus Schizosaccharomyces), Nadsonioideae, Lipomycoideae and Saccharomycoideae (e.g., genera Pichia, Kluyveromyces and Saccharomyces). The basidiosporogenous yeasts include the genera Leucosporidium, Rhodosporidium, Sporidiobolus, Filobasidium and Filobasidiella. Yeast belonging to the Fungi Imperfecti are divided into two families, Sporobolomycetaceae (e.g., genera Sporobolomyces, Bullera) and Cryptococcaceae (e.g., genus Candida). Of particular interest to the present invention are species within the genera Pichia, Kluyveromyces, Saccharomyces, Schizosaccharomyces and Candida. Of particular interest are the Saccharomyces species S. cerevisiae, S. carlsbergensis, S. diastaticus, S. douglasii, S. kluyveri, S. norbensis and S. oviformis. Species of particular interest in the genus Kluyveromyces include K. lactis. Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in Biology and Activities of Yeast (F.A. Skinner, S.M. Passmore & R.R. Davenport eds. 1980) (Soc. App. Bacteriol. Symp. Series No. 9). In addition to the foregoing, those of ordinary skill in the art are presumably familiar with the biology of yeast and the manipulation of yeast genetics. See, e.g., Biochemistry and Genetics of Yeast (M. Bacila, B.L. Horecker & A.O.M. Stoppani eds. 1978); The Yeasts (A.H. Rose & J.S. Harrison eds., 2nd ed., 1987); The Molecular Biology of the Yeast Saccharomyces (Strathern et al. eds. 1981). The disclosures of the foregoing references are incorporated herein by reference.

The present invention employs truncated leader sequences from a yeast α-factor gene. α-factor is an oligopeptide mating pheromone about 13 residues in length produced from a large precursor polypeptide between about 100 and 200 residues (typically about 120-160) in length (prepro-a-factor). The precursor is comprised of a hydrophobic "signal sequence" of about 20 residues (e.g., about 19-23, typically about 20-22) followed by an additional leader region of about 60 hydrophilic residues (the "pro" region), which is then linked to several tandem repeats of the mature pheromone sequence (typically about 2-6) separated by short oligopeptide spacer regions (typically about 6-8 residues) which provide for proteolytic processing to the mature pheromone.

The cloning of various prepro-α-factor genes has been reported. See, e.g., Kurjan et al., U.S. Patent No. 4,546,082; Singh et al. (1983) Nucleic Acids Res. 11:4049-4063; commonly owned U.S. Patent Application Serial No. 078,551, filed 28 July 1987 (EP-A-301669). In addition, DNA sequences encoding the prepro-α-factor gene can be identified by hybridization with probes from known prepro-α-factor sequences. See, e.g., Brake et al. (1983) Molec. & Cell Biol. 3:1440-1450. α-factor may also be purified from a yeast species, sequenced and probes designed to clone the prepro-α-factor gene. See, e.g., McCullough et al. (1979) J. Bacteriol. 138:146-154; Sato et al. (1981) Agric. Biol. Chem. 44:1451-1453; Singh et al. (1983), supra. It has also been determined that the α-factor leader sequence from one yeast species can be functional in another yeast species. See, e.g., U.S.S.N. 078,551 (EP-A-301669), supra. Thus, the present invention contemplates not only the use of α-factor leader sequences from yeast in general, but the use of such leader sequences in heterologous yeast species. For ease of presentation, however, the invention will be discussed in terms of the prepro-α-factor gene MFa1 from S. cerevisiae. See, e.g., Kurjan et al., U.S. Patent No. 4,546,083; Singh et al. (1983), supra.

The present invention employs chimeric DNA constructs encoding hybrid precursor polypeptides comprised of a leader sequence and a non-yeast polypeptide. For purposes of this invention, the leader sequence DNA is defined as beginning at the N-terminal start codon (methionine) of the precursor polypeptide through the codon encoding the last amino acid residue before the processing site that intervenes between the leader sequence and the sequence encoding the non-yeast protein. The leader sequence of the present invention is comprised of a truncated form of a yeast α-factor leader sequence, typically about 25 to about 50 amino acid residues in length. Thus, the leader sequence of the present invention is approximately 30 amino acid residues shorter than the typical full-length α-factor leader. MFα1, for example, contains a leader sequence of 83 amino acid residues followed by a hexapeptide spacer sequence which is cleaved by yeast processing enzymes. In making deletions from the leader sequence, it

is important that at least one glycosylation site (-Asn-Y-Thr/Ser-) is retained to provide for efficient secretion.

It is also necessary that the leader retain a functional α-factor signal sequence. As indicated above, the signal peptide is usually about 20 amino acids in length, and characteristic features including a hydrophobic core. See, e.g., von Heijne, (1984) J. Mol. Biol. 173:243-251. All of the prepro-α-factor sequences examined today encode for a hydrophobic peptide of about 20 residues in length. While the exact length of a signal peptide necessary to direct the precursor polypeptide to the secretory pathway is not defined, it will usually require between about 19 and about 23 residues, the minimum sequence required being readily definable by the testing of deletion mutants.

Thus, with reference to MFα1, deletions within the range of about 30 to about 60 residues, typically between about 33 and about 58 residues, and more typically between about 48 and about 58 residues, are contemplated by the present invention. These deletions will generally occur in the region between and including residues 26 through 83. It is preferred that the deletions include the glycosylation sites at residues 57-59 and 67-69. The deleted α-factor leader sequences may be replaced, in part, by non-α-factor leader sequences, if desired. The sequences should generally encode hydrophilic amino acid residues, should not encode glycosylation or processing sites, and preferably should be selected to maintain the overall length of the leader to be about 50 residues or less, preferably about 23 to 40 residues, and most preferably about 25 to 35 residues.

As indicated above, the leader sequence of the present invention has immediately 3' thereto a processing site which allows for the cleavage of the leader from the non-yeast protein sequence to which it is fused in the precursor polypeptide. The processing site employed in the present invention is defined as the codons defining the minimum number of amino acid residues which are specifically recognized for cleavage by the selected process (e.g., chemical, enzymatic, etc.). Various processing sites are known in the art, including both those active in vivo and in vitro. For example, the processing site may provide for in vitro processing by encoding a cleavage site for a proteolytic enzyme which does not occur in the yeast host. The recovered precursor polypeptide would then be treated with the enzyme to cleave the non-yeast protein from the precursor. Another in vitro processing site is a methionine codon which can be cleaved by post-expression treatment with cyanogen bromide. See, e.g., U.S. Patent No. 4,366,246.

In vivo processing sites can be selected from any peptide signals recognized by a yeast proteolytic enzyme which will provide for expression of the desired non-yeast protein sequence. Particularly preferred processing sites are those for the enzymes involved in processing of native prepro-α-factor. For example, dipeptidyl aminopeptidase A (DPAPase A) removes terminal -X-Ala- sequences, where X is Glu or Asp. See, e.g., Julius et al. (1983) Cell 32:839-852. The endopeptidase encoded by the KEX2 gene cleaves basic dipeptides comprised of Lys and Arg residues; i.e., Lys-Arg, Arg-Arg, Arg-Lys and Lys-Lys. Fuller et al., Microbiology 1986, pp. 273-278 (1986). In yeast, the α-factor precursor is first cleaved by the KEX2 endopeptidase, and then the N-termini are trimmed by DPAPase A to provide mature α-factor pheromone. Since it appears that the latter proteolytic process is a rate limiting step, it is preferred to eliminate the signals for DPAPase A, such that the processing site is comprised only of the signal for the KEX2 endopeptidase. In such an embodiment, therefore, the leader sequence will be joined to the non-yeast protein sequence by the dibasic peptide recognition site for KEX2 endopeptidase, such as Lys-Arg or Arg-Arg.

The carboxy-terminal portion of the precursor polypeptide of the present invention is a non-yeast protein. The DNA sequence encoding this portion is defined herein as beginning with the first codon downstream (3' direction) from the last codon of the processing site through to the translation stop codon which defines the carboxy-terminal of the precursor polypeptide. This DNA sequence will be considered to encode a "non-yeast protein" when, over its entire sequence, it defines a polypeptide that is not substantially homologous to a polypeptide expressed by yeast. In general, the preferred non-yeast proteins will be mammalian protein sequence (including their analogs; i.e., "muteins", fragments, etc.). As defined herein, non-yeast proteins can include, therefore, a fusion protein comprised of mammalian and yeast sequences, as well as "pro" forms of the mature mammalian protein.

DNA sequences encoding the non-yeast proteins can be sequences cloned from non-yeast organisms, or they can be synthetic sequences, usually prepared using yeast-preferred codons. Usually, the non-yeast proteins will be at least about 8 amino acids in length and can include polypeptides up to about 100,000 daltons or higher. Usually, the non-yeast polypeptide sequence will be less than about 300,000 daltons, and more usually less than about 150,000 daltons. Of particular interest are polypeptides of from about 5,000 to about 150,000 daltons, more particularly of about 5,000 to about 100,000 daltons. Illustrative non-yeast proteins of interest include hormones and factors, such as growth hormone, somatomedins, epidermal growth factor, luteinizing hormone, thyroid-stimulating hormone, oxytocin, insulin, vasopressin, renin, calcitonin, follicle-stimulating hormone, prolactin, erythropoietin, colony-stimulating factors, lymphokines such

as interleukin-2, globins, immunoglobulins, interferons (e.g., α, β or γ), enzymes, β-endorphin, enkephalin, dynorphin, insulin-like growth factors, etc.

In a preferred embodiment of the present invention, DNA constructs encoding the above-described precursor polypeptides have the structure:

5'-AF-CHO-X_n-S-Gene*-3'

wherein AF encodes a yeast alpha-factor signal peptide; CHO encodes a glycosylation site; X_n encodes a polypeptide of n amino acids in length that does not contain a glycosylation site or processing site that will cause the precursor polypeptide to be cleaved in vivo by the yeast host; n is an integer from 0 to about 30; Gene* encodes a non-yeast protein; and S encodes a processing site that provides for cleavage of said precursor polypeptide.

The signal peptide encoded by AF is the same α-factor signal peptide described above. It is approximately 20 residues in length (e.g., about 19-23) and is of sufficient length to direct the precursor polypeptide into the yeast secretory pathway. The precise minimum or maximum length can be determined for a particular α-factor by screening a series of deletion constructs.

The DNA sequence defined by CHO encodes a glycosylation site. It will generally be nine nucleotides in length, including three codons for the amino acids Asn-Y-Y'-, wherein Y is any amino acid residue, and Y' is Thr or Ser.

X_n, if present, encodes, for example, portions of the α-factor leader which are not deleted or unrelated amino acid sequences. In general, X_n will be a maximum of about 30 amino acid residues, more preferably a maximum of about 20 residues, and most preferably a maximum of about 10 residues. While it may not be necessary for X_n to encode any polypeptides (i.e., n=0), it may be desired to provide some spacing between the glycosylation site CHO, and the processing site S in the event that carbohydrate additions at the glycosylation site sterically hinder access of the agent which cleaves the processing sites. In such case, n will usually be a minimum of about 1, more preferably a minimum of about 2, while most preferably a minimum of about 3.

It is preferred that X_n, if present, not contain any functional glycosylation sites or processing sites recognized and cleaved by the yeast host. Further, when departing from sequences found in an α-factor leader, it is preferred to select hydrophilic amino acid residues. It is possible that the length of X_n will affect the efficiency of expression and secretion of the non-yeast protein. Selection of the appropriate length of X_n to optimize expression can be done through screening constructs of various sizes.

The non-yeast protein encoded by Gene* and the processing site encoded by S are as described above.

The DNA constructs of the present invention will normally be maintained in a replicon capable of stable maintenance in a host, particularly a yeast host. The replicons, usually plasmids, will include one or more replication systems, desirably two replication systems, allowing for maintenance of the replicon in both a yeast host for expression, and in a procaryotic host for cloning. Examples of such yeast-bacteria shuttle vectors include YEp24 [Botstein et al. (1979) Gene 8:17-24], pC1/1 [Brake et al. (1984) Proc. Natl. Acad. Sci. USA 81:4642-4646], and YRpl7 [Stnichomb et al. (1982) J. Mol. Biol. 158:157]. Furthermore, a plasmid expression vector may be a high or low copy number plasmid, the copy number generally ranging from about 1 to about 200. With high copy number yeast vectors, there will generally be at least 10, preferably at least 20, and usually not exceeding about 150 copies in a single host. Depending upon the non-yeast protein selected, either a high or low copy number vector may be desirable, depending upon the effect of the vector and the foreign protein on the host. See, e.g., Brake et al., supra. DNA constructs of present invention can also be integrated into the yeast genome by an integrating vector. Examples of such vectors are known in the art. See, e.g., Botstein et al., supra.

The selection of suitable yeast and other microorganism hosts for the practice of the present invention is within the skill of the art. When selecting yeast hosts for expression, suitable hosts may include those shown to have, inter alia, good secretion capacity, low proteolytic activity, and overall robustness. Yeast and other microorganisms are generally available from a variety of sources, including the Yeast Genetic Stock Center, Department of Biophysics and Medical Physics, University of California, Berkeley, California; and the American Type Culture Collection, Rockville, Maryland.

Methods of introducing exogenous DNA into yeast hosts are well known in the art. There is a wide variety of ways to transform yeast. For example, spheroplast transformation is taught, for example, by Hinnen et al. (1978) Proc. Natl. Acad. Sci. USA 75:1919-1933, and Stinchcomb et al., EPO Publication No. 45,573. Transformants are grown in an appropriate nutrient medium, and, where appropriate, maintained under selective pressure to insure retention of endogenous DNA. Where expression is inducible, growth can

be permitted of the yeast host to yield a high density of cells, and then expression is induced. The secreted, processed non-yeast protein can be harvested by any conventional means, and purified by chromatography, electrophoresis, dialysis, solvent-solvent extraction, and the like.

Examples

The following examples are provided for illustrative purposes only, and are not intended to limit the scope of the present invention. It is believed that the deposit of the starting biological materials is not necessary for the practice of the present invention since either the same or equivalent materials are publicly available.

I.

The following example provides a comparison of the levels of expression and secretion obtained with modified α-factor constructs used to express human proinsulin. Three constructs employ full-length α-factor leaders; one having α-factor leader with the three native glycosylation sites, one having all three of the glycosylation sites eliminated, and one having all of the sites, except the one at Asn_{23}, removed. The fourth construct is a truncated α-factor leader which retains a single glycosylation site at Asn_{23}.

A. pYGAI1

This plasmid encodes an α-factor leader [Brake et al. (1984) Proc. Natl. Acad. Sci. USA 81:4642-4646; EPO Publication No. 116,201] linked to human proinsulin. The proinsulin is encoded by a synthetic gene made with yeast preferred codons (Figure 1). The α-factor leader sequence, the synthetic proinsulin gene and the α-factor terminator sequence are from pYBCA5, the construction of which is shown in Figure 1. Transcription is mediated by the 404 bp BamHI-NcoI GAPDH promoter fragment. Travis et al. (1985) J. Biol. Chem. 260:4384-4389. The 1206 bp BamHI expression cassette consisting of the GAPDH promoter, the sequence encoding the α-factor leader linked to proinsulin and the α-factor terminator was cloned into the unique BamHI site of the yeast shuttle vector pAB24 (below) or pC1/1 such that the GAPDH promoter sequence was proximal to the SalI site of the vector to give the plasmids pYGAI1-AB24 or pYGAI1-C1/1, respectively. The 1206 bp BamHI expression cassette was also subcloned into the unique BamHI site of a derivative of pBR322 [pBR322(WEcoRI-SalI)BamHI; Travis et al. supra.] This plasmid was called pGAI1.

Plasmid pAB24 (Figure 4) is a yeast shuttle vector which contains the complete 2μm sequence [Broach, in: Molecular Biology of the Yeast Saccharomyces, Vol., 1, p. 445 (1981)] and pBR322 sequences. It also contains the yeast URA3 gene derived from plasmid YEp24 [Botstein et al. (1979) Gene 8:17) and the yeast LEU2d gene derived from plasmid pC1/1. EPO Publication No. 116,201. Plasmid pAB24 was constructed by digesting YEp24 with EcoRI and religating the vector to remove the partial 2m sequences. The resulting plasmid, YEp24WRI, was linearized by digestion with ClaI and ligated with the complete 2m plasmid which had been linearized with ClaI. The resulting plasmid, pCBou, was then digested with XbaI and the 8605 bp vector fragment was gel isolated. This isolated XbaI fragment was ligated with a 4460 bp XbaI fragment containing the LEU2d gene isolated from pC1/1; the orientation of the LEU2d gene is in the same direction as the URA3 gene.

B. pYGAI3

Plasmid pYGAI3 differs from pYGAI1 in that it encodes a modified α-factor leader wherein the codons for Asn at residues 23, 57 and 67 have been changed to encode Gln, thereby eliminating all three signals for N-linked glycosylation.

The α-factor leader and the N-terminal 13 amino acids of proinsulin encoded by this plasmid were constructed by ligation of synthetic oligonucleotides to give a 294 bp fragment with a 5' NcoI overhang and a 3' HindIII overhang, the sequence which is shown in Figure 2. The sequence of appropriate oligonucleotides was altered during the synthesis so that codons which specified Asn at positions 23, 57 and 67 of the natural α-factor leader now specified Gln at the same positions. The DNA sequence specifying the N-terminal 13 amino acids of proinsulin was identical to that in pYGAI1. The 294 bp synthetic DNA (NcoI-HindIII) fragment was substituted for the comparable fragment of pGAI1 and pYGAI1-C1/1 which gave the plasmids pGAI3 and pYGAI3, respectively.

C. pYGAI8

Plasmid pYGAI8 contains DNA encoding an α-factor leader which eliminates two of the three glycosylation sites. $Asn_{57,67}$ have been modified to $Gln_{57,67}$. The resulting plasmid has only a single glycosylation site at position Asn_{23}. pYGAI8 was prepared as follows.

First, a 5' fragment was isolated from the expression cassette of pGAI1 by cutting with HpaII, followed by cutting with BamHI, and then gel isolating a 504 bp fragment containing the GAPDH promoter and the sequence encoding residues 1-33 of the α-factor leader. Next, plasmid pYGAI3 encoding an α-factor leader lacking glycosylation sites was also sequentially cut with HpaII and BamHI, and a 702 bp fragment isolated containing sequences encoding modified α-factor leader residues 34-83, the LysArg processing site, the proinsulin sequence and the a-factor termination sequence. This fragment was then ligated to the 504 bp fragment from pGAI1, cut with BamHI and a 1206 bp fragment isolated.

The above 1.2 kb BamHI fragment which contained a complete GAPDH promoter/α-factor leader/proinsulin/α-factor terminator expression cassette was then ligated into BamHI-cut and phosphatase-treated pBR322(WEcoRI-SalI)BamHI to give plasmid pGAI8, which was cloned in E. coli.

The 1.2 kb expression cassette from pGAI8 was removed by cutting with BamHI and then gel isolating the fragment. It was ligated into BamHI-cut and phosphatase-treated yeast shuttle vector pAB24. Insertion of the expression cassette was in the unique BamHI site of the pBR322 sequences such that the GAPDH promoter was proximal to the unique SalI site of the vector. This plasmid was pYGAI8.

D. pYGAI7

Plasmid pYGAI7 contains the DNA encoding a truncated α-factor leader and the synthetic gene for human proinsulin. The α-factor leader has been truncated so that it encodes only amino acids 1-35 of the α-factor leader and therefore contains a single site for glycosylation at Asn_{23}. This yeast expression vector was constructed as follows.

First, pGAI1 was cut with HindIII. An HpaII-HindIII linker was added of the following structure:

```
5'-CGGCTAAAAGATTCGTTAACCAACACTTGTGTGGTTCTCACTTGGTTGA
   CGATTTTCTAAGCAATTGGTTGTGAACACACCAAGAGTGAACCAACTTCGA-5'
```

After adding the linker, the linearized plasmid was cut with BamHI, and a 558 bp HpaII-BamHI fragment was gel isolated. This fragment contains the codons for residues 34-35 of the α-factor leader linked directly to a Lys-Arg processing site and the proinsulin sequence. There are no intervening sequences between the codon for residue 35 of the α-factor leader and the processing site directly adjacent to the proinsulin sequence.

Second, pGAI1 was cut with HpaII and BamHI, and a 504 bp fragment gel isolated. This fragment contains the GAPDH promoter and nucleotides encoding amino acids 1-33 of the α-factor leader, the 3' end terminating in an HpaII overhand complementary to the 5' end of the above-described 558 bp HpaII-BamHI fragment. These two fragments were ligated together and cut with BamHI to provide an expression cassette containing the GAPDH promoter, sequences encoding a modified α-factor leader containing residues 1-35 directly linked to a Lys-Arg processing site, the proinsulin gene, and the α-factor terminator. The cassette was then ligated into a BamHI site of pBR322(WEcoRI-SalI)BamHI, as described above, to give plasmid pGAI7 and cloned in E.coli.

pGAI7 was then cut with BamHI, and the 1062 bp expression cassette gel isolated. The expression cassette was then ligated into the BamHI site of pAB24 to give plasmid pYGAI7.

E. Comparative Expression

Plasmids pYGAI1-C1/1, pYGAI3, pYGAI1-AB24, pYGAI7 and pYGAI8 were transformed into Saccharomyces cerevisiae strain AB103.1 (Mata, leu2-3,112,ura3-52, his4-580, pep4-3[cir°]) essentially as described by Hinnen et al. (1978) Proc. Natl. Acad. Sci. USA 75:1929-1933. Transformants of pYGAI1-C1/1 and pYGAI3 were selected for leucine prototrophy, transformants of the other plasmids were selected for ura prototrophy.

Data shown in Table 1 compares secretion of proinsulin mediated by the natural α-factor leader (pYGAI1-C1/1) or the a-factor leader with Gln substituted for Asn at positions 23, 57 and 67 (pYGAI3).

Inoculum cultures (.2 ml of individual transformants) were grown for 48 hr in synthetic complete medium lacking leucine [SD-leu; Sherman et al., Methods in Yeast Genetics, p. 62 (Cold Spring Harbor Laboratory, 1982)] and diluted 20-fold into the same medium. Cultures were grown 48-72 hrs, culture supernatants were prepared by centrifugation and were assayed for immunoreactive cross-reacting insulin-like material (ILM) in a competition radioimmune assay with ^{125}I-labeled insulin. As can be seen in Table 1, elimination of the three glycosylation sites from the α-factor leader resulted in essentially no secretion of insulin-like-material compared to that mediated by the native α-factor leader.

Data presented in Table 2 compares transformants of pYGAI1-pAB24 (full-length native α-factor leader, pYGAI8 (full-length α-factor leader with only one glycosylation site at Asn_{23}) and pYGAI7 (truncated α-factor leader containing a single glycosylation site at Asn_{23}) for their ability to secrete insulin-like-material. Inoculum cultures of the indicated transformants (.2 ml) in SD-Leu grown for .48 hr at 30°C were pelleted by centrifugation, washed and diluted 20-50 fold into ura$^-$ medium. This medium contains 0.67% yeast nitrogen base, 1% succinic acid, 0.35% NaOH, 0.5% casamino acids, 2% glucose, 0.005% adenine, 0.01% tryptophan and 0.02% threonine. Cultures were grown at 30°C for 48-72 hr, and culture supernatants prepared and assayed as described above. Data presented in Table 2 show that the transformants carrying the construct employing the truncated α-factor leader retaining a single glycosylation site at Asn_{23} secreted generally more immunoreactive insulin-like-material than did transformants bearing the construct with the full-length native α-factor leader. Transformants bearing the construct with the full-length α-factor leader with the same single glycosylation site (Asn_{23}) secreted much less insulin cross-reactive material than did transformants bearing the full-length native α-factor leader or the truncated α-factor leader.

Table 1

Effect of Elimination of α-Factor Leader Glycosylation Sites on Secretion of Insulin-Like-Material

Transformant	OD_{650}	ILM[1]	
		mg/ml	mg/ml,OD_{650}
AB103.1[pYGAI1-C1/1]$_{-1}$	5.9	.24	.04
$_{-2}$	5.9	.24	.04
$_{-3}$	2.1	.08	.04
AB103.1[pYGAI3] $_{-1}$	ND	.003	-
$_{-2}$	ND	.007	-

1) Cross-reactive insulin-like-material (ILM) as determined by a competition radioimmune assay with ^{125}I-labeled insulin and insulin standards. Data is reported as ILM secreted per ml of culture and in some cases as ILM secreted per ml normalized to a culture cell density with an absorbance at wavelength 650mm of 1.

Table 2

Effect of Truncated or Full-Length α-Factor Leader with a Single Glycosylation Site at Asn_{23} on Secretion of Insulin-Like-Material

Transformant	No. of Tests[1]	ILM[2]					
		mg/ml			mg/ml OD_{650}		
		Range	Mean	Std. dev.	Range	Mean	Std. dev.
AB103.1 [pYGAI1-AB24]	7	.22-.55	.37	.12	.02-.04	.026	.01
AB103.1 [pYGAI7)	8	.38-1.0	.62	.24	.03-.06	.043	.01
AB103.1 (pYGAI8]	8	.03-.18	.09	.05	.003-.01	.007	.003

1) A minimum of three independent transformants were tested.
2) Secreted cross-reactive insulin-like-material (ILM) was determined by a competition radioimmune assay with ^{125}I-labeled insulin and insulin standards. Data is reported as ILM secreted per ml of culture and as ILM secreted per ml normalized to a culture cell density with an absorbance at wavelength 650 mm of 1.

II.

This example compares the expression of a full-length α-factor leader construct, retaining all glycosylation sites, to an expression construct employing a truncated α-factor sequence retaining only a single glycosylation site at Asn_{23}. The non-yeast protein employed in this example is a human proinsulin analog wherein the connecting "C" peptide has been replaced by a yeast KEX2 endopeptidase cleavage site.

A. pYGAIC3

The plasmid pGAIC3 was made by replacing the 231 bp HindIII-SalI fragment of pGAI1 which encodes amino acids 14 through 30 of the B chain, the C-peptide, the A chain and 2 translation stop codons with a 132 bp synthetic HindIII-SalI gene fragment (shown in Figure 3) which encodes amino acids 14 through 30 of the B chain, a Lys-Arg KEX2 endopeptidase cleavage site, the A chain, and translation stop codons. The plasmid pYGAIC3 was prepared from pGAIC3 as follows.

Plasmid pGAIC3 was digested with BamHI, and the 1107 bp BamHI expression cassette containing the GAPDH promoter, the sequence encoding α-factor leader linked to proinsulin analog, and the α-factor transcription terminator was isolated and ligated into BamHI digested and phosphatase-treated pAB24, and then cloned in E. coli. Plasmid pYGAIC3 was obtained, in which the expression cassette was oriented such that the GAPDH promoter was proximal to the unique SalI site of the vector.

B. $PYaf_L7C3$

Plasmid $pYaF_L7C3$ contains DNA encoding the truncated a-factor leader described above for pYGAI7 linked to the sequence for the proinsulin analog, also described above (pYGAIC3). This plasmid was constructed as follows.

First, pGAIC3 was cut with HindIII and SalI, and a 132 bp fragment was gel isolated. This fragment contains sequences encoding all but the first 12 codons of the proinsulin analog. It was ligated into a gel isolated 4640 bp fragment from HindIII- and SalI-digested pGAI7 to provide plasmid paf_L7C3. After cloning in E. coli, this plasmid was cut with BamHI and a 1062 bp BamHI fragment was gel isolated. This expression cassette contains the truncated α-factor leader construct of pGAI7 with the proinsulin analog in place of the normal proinsulin sequence. The expression cassette was then ligated into the BamHI site of pAB24, as described above, to give $pYaf_L7C3$.

Comparative Expression

Expression levels were determined for pYGAIC3 and pYaf$_L$7C3 in two strains of S. cerevisiae. Strain AB103.1 has been described in Example I. Strain AB110-4 is a derivative of Saccharomyces cerevisiae strain AB110 (Mata, leu2, ura3-52, pep4-3 his4-580[cir°]) in which a deletion has been engineered into the pep4 gene. These strains were transformed as described above with plasmids pYGAIC3 and pYaf$_L$7C3, and ura prototrophs were selected. Inoculum cultures were grown in SD-leu [Sherman et al., supra.] at 30°C for 24-48 hours then pelleted by centrifugation, washed and diluted 20 fold into ura$^-$ medium (described above) and grown for 48-72 hours at 30°C. Cell-free conditioned culture medium was prepared by centrifugation for assay in a competition insulin radioimmune assay.

The results are shown in Table 3. As can be seen the truncated α-factor construct mediates increased secretion of immunoreactive proinsulin analog, compared to the natural α-factor leader sequence.

Table 3

Secretion of ILM from a Proinsulin Analog Construct Mediated by a Truncated a-Factor Leader or Natural a-Factor Leader

Transformant	No. of Tests[1]	ILM[2]					
		mg/ml			mg/ml, OD_{650}		
		Range	Mean	Std. dev.	Range	Mean	Std. dev.
AB103.1 [pYGAIC3]	6	1-2.75	1.66	.65	.11-.20	.14	.04
AB103.1 [pYaf$_L$7C3]	6	1.5-6.63	4.46	2.15	.14-.60	.40	.19
AB110.4 [pYGAIC3]	3	1.15-1.4	1.28	.13	.10-.12	.11	.01
AB110.4 [pYaf$_L$7C3]	3	2.15-4.12	3.46	1.14	.19-.38	.31	.10

1) A minimum of three independent transformants were tested.
2) Secreted cross-reactive insulin-like-material (ILM) was determined by a competition radioimmune assay with ^{125}I-labeled insulin and insulin standards. Results are reported as amounts of ILM per ml of culture and as amount secreted per ml normalized to a cell density with an absorbance at 650 mm wavelength = 1.

III.

This example described the construction of a truncated a-factor expression vector which mediates increased expression levels of active insulin-like growth factor-1.

First, a DNA sequence encoding a truncated a-factor leader and a coding sequence for IGF1 was prepared. A synthetic sequence was prepared by standard procedures employing an Applied Biosystems 380A DNA synthesis machine according to manufacturer's direction. Fourteen DNA sequences were synthesized ranging from 22 to 57 bases in length, purified by PAGE, and phosphorylated individually by T4 kinase in the presence of ATP. The sequences were then annealed and ligated by standard procedures.

The sequence of the synthetic gene is shown in Figure 5. The purified synthetic gene fragment was cloned into NcoI/SalI digested pBS100 (described below). The resulting plasmid was called pBS100 Taf$_L$ IGF1.

Plasmid pBS100 contains a yeast expression cassette cloned into a pBR322 derivative, pAB12. The expression cassette contains the hybrid ADH-2/GAPDH promoter and the GAPDH terminator flanking a non-essential gene segment. The ADH-2/GADPH promoter is a 1200 bp BamHI-NcoI fragment isolated from pJS103 (see below) and the GAPDH terminator is a 900 bp SalI-BamHI fragment isolated from plasmid pPAG1. EPO Publication No. 164,556. Plasmid pBS100 also contains a non-essential fragment between NcoI and SalI sites which is replaced by gene fragments of interest. The expression cassette can be removed from pBS100 by digestion with BamHI and cloned into yeast shuttle vectors for introduction into yeast cells.

Plasmid pJS103, which contains the hybrid ADH-2/GAPDH promoter employed above, was constructed as follows. The ADH-2 portion of the promoter was constructed by cutting a plasmid containing the wild-type ADH2 gene from plasmid pADR2 [Beier et al. (1982) Nature 300:724-728] with restriction enzyme EcoR5, which cuts at position +66 relative to the ATG start codon, as well as in two other sites in pADR2, outside of the ADH2 region. The resulting mixture of a vector fragment and two smaller fragments was reacted with Bal31 exonuclease to remove about 300 bp. Synthetic XhoI linkers were ligated onto the Bal31-treated DNA. The resulting DNA linker vector fragment (about 5 kb) was separated from the linkers by column chromatography, cut with restriction enzyme XhoI, religated, and used to transform E. coli to ampicillin resistance. The positions of the XhoI linker were determined by DNA sequencing. One plasmid which contained an XhoI linker within the 5' nontranscribed region of the ADH2 gene (position -232 from ATG) was cut with the restriction enzyme XhoI, treated with nuclease SI, and subsequently treated with the restriction enzyme EcoRI to create a linear vector molecule having 1 blunt end at the site of the XhoI linker and an EcoRI end. The GAPDH portion of the promoter was constructed by cutting plasmid pPGAP [EPO Publication No. 164,556] with the enzymes BamHI and EcoRI, followed by the isolation of the 0.4 Kbp DNA fragment. This purified fragment was then completely digested with the enzyme AluI and an approximately 200 bp fragment was isolated. This GAPDH promoter fragment was ligated to the ADH-2 fragment present on the linear vector described above to give plasmid pJS103.

A BamHI fragment was then isolated from pBS100 Ta_{fL} IGF1. This fragment contains the ADH2/GAPDH promoter, a truncated α-factor leader (AA 1-25, 81-83) a LysArg processing site, a coding sequence for IGF1, and the GAPDH terminator sequence. This BamHI fragment was then cloned into pAB24, previously digested with BamHI. A positive clone was selected, and while initially called plasmid 18.5, it was subsequently named pYLUIGF1-55 . (See Figure 6.)

A second expression vector, pYLUIGF1-24 was also prepared by analogous methods. A restriction map is shown in Figure 7. This vector is similar to pYLUIGF1-55, except that it has a full-length α-factor leader directing secretion with three glycosylation sites (compare Example I.A.) and the α-factor terminator.

Yeast strain AB110 (EPO Publication No. 164,556) was transformed with pYLUIGF1-55 and pYLUIGF1-24 by conventional spheroplasting techniques [Hinnen et al. (1978) Proc. Natl. Acad. Sci. USA 75:1919-1933], and expression compared.

The expression of IGF1 from AB110 (pYLUIGF1-55) and AB110 (pYLUIGF1-24) is non-constitutive. Induction of IGF1 expression was achieved by bringing about a low concentration of glucose in the growth medium. Under standard conditions, shake flask cultures (25 ml) fully utilize the glucose in the medium by 18-24 hours post inoculation. Thus, 25 ml cultures of AB110 (pYLUIGF1-55) and AB110 (pXLUIGF1-24) were grown under standard conditions for 72 hours. Supernatant samples were taken at 49 and 72 hours post inoculation and assayed for IGF1 biological activity (RRA) and for immunoreactivity (RIA) with anti-IGF1 antibodies. As can be seen, pYLUIGF1-55, with a truncated α-factor leader, secreted protein of which a substantially greater fraction was biologically active. Although pYLUIGF1-24 secreted more protein that showed reactivity with IGF1 antibodies, relatively little of this protein was biologically active.

The results are shown in Table 4. The radioreceptor assay (RRA) measures the ability of IGF-1 to bind to its receptor. This is a measure of the biological activity of recombinant polypeptide since it is believed that IGF-1 exerts all of its activity through its receptor. The receptor assay is described in Marshall et al. (1974) J. Clin. Endorinol. Metab. 19:283-292. The radioimmunoassay (RIA) is a competitive assay that measures the amount of protein antigenically cross-reactive with native IGF-1, whether or not it is biologically active. The assay is described in Zapf et al. (1981) J. Clin. Invest. 68:1321-1330.

Table 4. Secretion of IGF1 Mediated by a Truncated a-Factor Leader or a Natural a-Factor Leader

Transformant	49 hrs		72 hrs	
	RRA[1]	RIA[2]	RRA	RIA
AB110 (pYLUIGF1-55)	1.0	14	1.3	14
AB110 (pYLUIFG1-24)	1.3	54	2.7	66

1) mg/ml

2) mg/ml

Deposit of Biological Materials

The following expression vectors were deposited with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland, U.S.A., and will be maintained under the provisions of the Budapest Treaty. The accession numbers and dates of deposit are listed below.

Deposited Material	ATCC Number	Deposit Date
E. coli (pYGAI7)	67597	12/29/87
E. coli (pYafL7C3)	67596	12/29/87
E. coli (pYLUIGFI-55)	67595	12/29/87

These deposits are provided for the convenience of those skilled in the art. These deposits are neither an admission that such deposits are required to practice the present invention nor that equivalent embodiments are not within the skill of the art in view of the present disclosure. The public availability of these deposits is not a grant of a license to make, use or sell the deposited materials under this or any other patent. The nucleic acid sequences of the deposited materials are incorporated in the present disclosure by reference, and are controlling if in conflict with any sequences described herein.

Although the foregoing invention has been described in some detail for the purpose of illustration, it will be obvious that changes and modifications may be practiced within the scope of the appended claims by those of ordinary skill in the art.

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, IT, LU, NL, SE

1. A yeast cell comprising a DNA construct that provides for the expression and secretion of a non-yeast protein, said DNA construct comprising a coding sequence under the control of yeast-recognized transcription initiation and termination sequences, said coding sequence encoding a precursor polypeptide comprised of a leader sequence and said non-yeast protein linked by a processing site that provides for the cleavage of said non-yeast protein from said precursor polypeptide, wherein said leader sequence is about 25 to about 50 N-terminal residues of said precursor polypeptide and comprises the signal peptide of a Saccharomyces or Kluveromyces alpha-factor precursor and a single glycosylation site.

2. The cell according to claim 1 wherein said non-yeast protein is a mammalian protein, and said alpha-factor is Saccharomyces alpha factor.

Table 4. Secretion of IGF1 Mediated by a Truncated a-Factor Leader or a Natural a-Factor Leader

Transformant	49 hrs			
72 hrs	RRA[1]	RIA[2]	RRA	RIA
AB110 (pYLUIGF1-55)	1.0	14	1.3	14
AB110 (pYLUIFG1-24)	1.3	54	2.7	66

1) mg/ml

2) mg/ml

Deposit of Biological Materials

The following expression vectors were deposited with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland, U.S.A., and will be maintained under the provisions of the Budapest Treaty. The accession numbers and dates of deposit are listed below.

Deposited Material	ATCC Number	Deposit Date
E. coli (pYGAI7)	67597	12/29/87
E. coli (pYaf$_L$7C3)	67596	12/29/87
E. coli (pYLUIGFI-55)	67595	12/29/87

These deposits are provided for the convenience of those skilled in the art. These deposits are neither an admission that such deposits are required to practice the present invention nor that equivalent embodiments are not within the skill of the art in view of the present disclosure. The public availability of these deposits is not a grant of a license to make, use or sell the deposited materials under this or any other patent. The nucleic acid sequences of the deposited materials are incorporated in the present disclosure by reference, and are controlling if in conflict with any sequences described herein.

Although the foregoing invention has been described in some detail for the purpose of illustration, it will be obvious that changes and modifications may be practiced within the scope of the appended claims by those of ordinary skill in the art.

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, IT, LU, NL, SE

1. A yeast cell comprising a DNA construct that provides for the expression and secretion of a non-yeast protein, said DNA construct comprising a coding sequence under the control of yeast-recognized transcription initiation and termination sequences, said coding sequence encoding a precursor polypeptide comprised of a leader sequence and said non-yeast protein linked by a processing site that provides for the cleavage of said non-yeast protein from said precursor polypeptide, wherein said leader sequence is about 25 to about 50 N-terminal residues of said precursor polypeptide and comprises the signal peptide of a Saccharomyces or Kluveromyces alpha-factor precursor and a single glycosylation site.

2. The cell according to claim 1 wherein said non-yeast protein is a mammalian protein, and said alpha-factor is Saccharomyces alpha factor.

* C L M S *

3. The cell according to claim 2 wherein said mammalian protein is human insulin-like growth factor I or a precursor of human insulin.

4. The cell according to claim 3 wherein said precursor of human insulin is human proinsulin or wherein said precursor of human insulin comprises insulin a chain and insulin b chain linked by a yeast-recognised processing site cleaved in vivo.

5. The cell according claim 4 wherein said processing site is cleaved by the KEX2 gene product of Saccharomyces.

6. The cell according to any one of claims 1 to 5 therein said yeast cell is from the genus Saccharomyces.

7. The cell according to claim 6 wherein said yeast cell is S. cerevisiae.

8. The cell according claim 7 wherein said yeast α-factor precursor is S. cerevisiae MFα1.

9. A double-stranded DNA molecule comprising a region encoding a precursor polypeptide secretable by a yeast host, said region, with reference to one of the strands, having the structure:

5'-AF-CHO-X_n-S-Gene*-3'

wherein AF-CHO-X_n comprises a single functional glycosylation site in which :

AF	encodes a Saccharomyces or Kluveromyces alpha-factor signal peptide;
CHO	encodes a glycosylation site;
X_n	encodes a polypeptide of n amino acids in length that does not contain a glycosylation site or a processing site that provices for cleavage of said precursor polypeptide in vivo by yeast;
n	is an integer from 0 to about 30;
	and wherein
Gene*	encodes a non-yeast protein; and
S	encodes a processing site that provides for cleavage of said precursor polypeptide.

10. The DNA molecule according to claim 9 wherein AF encodes a polypeptide of about 19 to 23 amino acids in length.

11. The DNA molecule according claim 9 or claim 10 wherein n is an integer from about 0 to about 20.

12. The DNA molecule according to claim 11 wherein n is an integer from about 0 to about 10.

13. The DNA molecule according to claim 12 wherein n is an integer from about 3 to about 10.

14. The DNA molecule according any one of claims 9 to 13 wherein said yeast host is a Saccharomyces.

15. The DNA molecule according to any one of claims 9 to 14 wherein said yeast α-factor signal peptide is a Saccharomyces signal peptide.

16. The DNA molecule according to any one of claims 9 to 15 wherein S encodes a processing site recognised in vivo by said yeast host.

17. The DNA according to claim 16 wherein S encodes a dipeptide recognised by the KEX2 endopeptidase.

18. The DNA molecule according to claim 17 wherein said dipeptide is 5'-Lys-Arg-3' or 5'-Arg-Arg-3'.

19. The DNA molecule according to any of claims 9 to 18 comprising a replicon.

* C L M S *

20. The DNA molecule according to claim 19 wherein said region encoding said precursor polypeptide is under the control of yeast-recognised transcription initiation and termination sequences, and said replicon is a yeast replicon.

21. The DNA molecule according to claim 20 wherein said replicon is a plasmid or a chromosome.

22. A method for the production of a non-yeast protein which comprises growing a yeast cell according to any one of claims 1 to 8 under conditions whereby the non-yeast protein is expressed.

23. A method according to claim 22 wherein the yeast cell comprises a DNA molecule according to any one of claims 9 to 21.

24. A method according to claim 22 or 23 which further comprises recovery of the non-yeast protein in purified form.

Claims for the following Contracting States : ES, GR

1. A method for the production of a non-yeast protein which comprises growing a yeast cell comprising a DNA construct that provides for the expression and secretion of a non-yeast protein, said DNA construct comprising a coding sequence under the control of yeast-recognized transcription initiation and termination sequences, said coding sequence encoding a precursor polypeptide comprised of a leader sequence and said non-yeast protein linked by a processing site that provides for the cleavage of said non-yeast protein from said precursor polypeptide, wherein said leader sequence is about 25 to about 50 N-terminal residues of said precursor polypeptide and comprises the signal peptide of a Saccharomyces or Kluveromyces alpha factor precursor and a single glycosylation site, under conditions whereby the non-yeast protein is expressed.

2. The method according to claim 1 wherein said non-yeast protein is a mammalian protein, and said alpha-factor is Saccharomyces alpha factor.

3. The method according to claim 2 wherein said mammalian protein is human insulin-like growth factor I or a precursor of human insulin.

4. The method according to claim 3 wherein said precursor of human insulin is human proinsulin or wherein said precursor of human insulin comprises insulin a chain and insulin b chain linked by a yeast-recognised processing site cleaved *in vivo*.

5. The method according to claim 4 wherein said processing site is cleaved by the KEX2 gene product of Saccharomyces.

6. The method according to any one of claims 1 to 5 wherein said yeast cell is from the genus Saccharomyces.

7. The method according to claim 6 wherein said yeast cell is S. cerevisiae.

8. The method according to claim 7 wherein said yeast α-factor precursor is S. cerevisiae MFα1.

9. The method according to any one of claims 1 to 8 wherein the DNA construct comprises a double-stranded DNA molecule comprising a region encoding a precursor polypeptide secretable by a yeast host, said region, with reference to one of the strands, having the structure:

5'-AF-CHO-X_n-S-Gene*-3'

wherein AF-CHO-X_n comprises a single functional glycosylation site in which:
AF encodes a Saccharomyces or Kluveromyces alpha-factor signal peptide; CHO encodes a glycosylation site;
X_n encodes a polypeptide of n amino acids in length that does not contain a glycosylation site or a processing site that provides for cleavage of said precursor polypeptide *in vivo* by yeast;

n is an integer from 0 to about 30;
Gene* encodes a non-yeast protein; and
S encodes a processing site that provides for cleavage of said precursor polypeptide.

10. The method according to claim 9 wherein AF encodes a polypeptide of about 19 to 23 amino acids in length.

11. The method according to claim 10 or claim 11 wherein n is an integer from about 0 to about 20.

12. The method according to claim 11 wherein n is an integer from about 0 to about 10.

13. The method according to claim 12 wherein n is an integer from about 3 to about 10.

14. The method according any one of claims 9 to 13 wherein said yeast host is a Saccharomyces.

15. The method according to any one of claims 9 to 14 wherein said yeast α-factor signal peptide is a Sacchromyces signal peptide.

16. The method according to any one of claims 9 to 15 wherein S encodes a processing site recognised in vivo by said yeast host.

17. The method to claim 16 wherein S encodes a dipeptide recognised by the KEX2 endopeptidase.

18. The method according to claim 17 wherein said dipeptide is 5'-Lys-Arg-3' or 5'-Arg-Arg-3'.

19. The method according to any of claims 9 to 18 comprising a replicon.

20. The method according to claim 19 wherein said region encoding said precursor polypeptide is under the control of yeast-recognised transcription initiation and termination sequences, and said replicon is a yeast replicon.

21. The method according to claim 20 wherein said replicon is a plasmid or a chromosome.

22. A method according to any one of claims 1 to 21 which further comprises recovery of the non-yeast protein in purified form.

Patentansprüche
Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, IT, LU, NL, SE

1. Hefezelle, umfassend eine DNA-Konstruktion, die die Expression und Sekretion eines Nicht-Hefeproteins gewährleistet, wobei die DNA-Konstruktion eine codierende Sequenz unter der Kontrolle von durch die Hefe erkannten Transkriptionsinitiations- und -terminationssequenzen umfaßt, wobei die codierende Sequenz ein Vorläuferpolypeptid aus einer Leadersequenz und dem Nicht-Hefeprotein, verknüpft durch eine Processing-Stelle, die die Spaltung des Nicht-Hefeproteins von dem Vorläuferpolypeptid gewährleistet, codiert, wobei die Leadersequenz etwa 25 bis etwa 50 N-terminale Reste des Vorläuferpolypeptids ausmacht, und das Signalpeptid eines Vorläufers des α-Faktors von Saccharomyces oder Kluveromyces und eine einzige Glycosylierungsstelle umfaßt.

2. Zelle nach Anspruch 1, dadurch **gekennzeichnet,** daß das Nicht-Hefeprotein ein Säugerprotein ist, und daß der α-Faktor der Saccharomyces-α-Faktor ist.

3. Zelle nach Anspruch 2, dadurch **gekennzeichnet,** daß das Säugerprotein ein menschlicher insulinartiger Wachstumsfaktor I oder ein Vorläufer von menschlichem Insulin ist.

4. Zelle nach Anspruch 3, dadurch **gekennzeichnet,** daß der Vorläufer von menschlichem Insulin menschliches Proinsulin ist, oder daß der Vorläufer von menschlichem Insulin die Insulin A-Kette und die Insulin B-Kette in Verknüpfung durch eine von der Hefe erkannte Processing-Stelle, die in vivo gespalten wird, umfaßt.

* C L M S *

5. Zelle nach Anspruch 4, dadurch **gekennzeichnet,** daß die Processing-Stelle durch das KEX2-Genprodukt von Saccharomyces gespalten wird.

6. Zelle nach einem der Ansprüche 1 bis 5, dadurch **gekennzeichnet,** daS die Hefezelle der Gattung Saccharomyces angehört.

7. Zelle nach Anspruch 6, dadurch **gekennzeichnet,** daß die Hefezelle S. cerevisiae ist.

8. Zelle nach Anspruch 7, dadurch **gekennzeichnet,** daß der Vorläufer des Hefe-α-Faktors S. cerevisiae MFα1 ist.

9. Doppelsträngiges DNA-Molekül umfassend eine Region, die ein Vorläuferpolypeptid, das von einer Wirtshefe sezernierbar ist, codiert, wobei die Region unter Bezugnahme auf einen der Stränge die Struktur

5'-AF-CHO-X_n-S-Gen*-3'

besitzt, worin
AF-CHO-X_n eine einzige funktionelle Glycosylierungsstelle umfaßt, worin:

AF	ein α-Faktor-Signal von Saccharomyces oder Kluveromyces codiert;
CHO	eine Glycosylierungsstelle codiert;
X_n	ein Polypeptid mit n Aminosäuren in einer Länge, die keine Glycosylierungsstelle oder keine Processing-Stelle enthält, die die Spaltung des Vorläuferpolypeptids in vivo durch Hefe gewährleistet, codiert;
n	eine ganze Zahl von 0 bis etwa 30 ist; und worin
Gen*	ein Nicht-Hefeprotein codiert; und
S	eine Processing-Stelle, die die Spaltung des Vorläuferpolypeptids gewährleistet, codiert.

10. DNA-Molekül nach Anspruch 9, dadurch **gekennzeichnet,** daß AF ein Polypeptid mit einer Länge von etwa 19 bis 23 Aminosäuren codiert.

11. DNA-Molekül nach Anspruch 9 oder 10, dadurch **gekennzeichnet,** daß n eine ganze Zahl von etwa 0 bis etwa 20 ist.

12. DNA-Molekül nach Anspruch 11, dadurch **gekennzeichnet,** daß n eine ganze Zahl von etwa 0 bis etwa 10 ist.

13. DNA-Molekül nach Anspruch 12, dadurch **gekennzeichnet,** daß n eine ganze Zahl von etwa 3 bis etwa 10 ist.

14. DNA-Molekül nach einem der Ansprüche 9 bis 13, dadurch **gekennzeichnet,** daß die Wirtshefe Saccharomyces ist.

15. DNA-Molekül nach einem der Ansprüche 9 bis 14, dadurch **gekennzeichnet,** daß das α-Faktor-Signalpeptid der Hefe ein Saccharomyces-Signalpeptid ist.

16. DNA-Molekül nach einem der Ansprüche 9 bis 15, dadurch **gekennzeichnet,** daß S eine Processing-Stelle, die in vivo von der Wirtshefe erkannt wird, codiert.

17. DNA nach Anspruch 16, dadurch **gekennzeichnet,** daß S ein Dipeptid, das von der KEX2-Endopeptidase erkannt wird, codiert.

18. DNA-Molekül nach Anspruch 17, dadurch **gekennzeichnet,** daß das Dipeptid 5'-Lys-Arg-3' oder 5'-Arg-Arg-3' ist.

19. DNA-Molekül nach einem der Ansprüche 9 bis 18, umfassend ein Replicon.

* C L M S *

20. DNA-Molekül nach Anspruch 19, dadurch **gekennzeichnet,** daß die Region, die das Vorläuferpolypeptid codiert, unter der Kontrolle von durch die Hefe erkannten Transkriptionsinitiations- und -terminationssequenzen ist, und daß das Replicon ein Hefereplicon ist.

21. DNA-Molekül nach Anspruch 20, dadurch **gekennzeichnet,** daß das Replicon ein Plasmid oder ein Chromosom ist.

22. Verfahren zur Produktion eines Nicht-Hefeproteins, dadurch **gekennzeichnet,** daß man eine Hefezelle nach einem der Ansprüche 1 bis 8 unter Bedingungen, bei denen das Nicht-Hefeprotein exprimiert wird, züchtet.

23. Verfahren nach Anspruch 22, dadurch **gekennzeichnet,** daß die Hefezelle ein DNA-Molekül nach einem der Ansprüche 9 bis 21 umfaßt.

24. Verfahren nach Anspruch 22 oder 23, dadurch **gekennzeichnet,** daß man weiterhin das Nicht-Hefeprotein in gereinigter Form isoliert.

Patentansprüche für folgende Vertragsstaaten : ES, GR

1. Verfahren zur Produktion eines Nicht-Hefeproteins, dadurch **gekennzeichnet,** daß man eine Hefezelle umfassend eine DNA-Konstruktion, die die Expression und Sekretion eines Nicht-Hefeproteins gewährleistet, wobei die DNA-Konstruktion eine codierende Sequenz unter der Kontrolle von durch die Hefe erkannten Transkriptionsinitiations- und -terminationssequenzen umfaßt, wobei die codierende Sequenz ein Vorläuferpolypeptid aus einer Leadersequenz und dem Nicht-Hefeprotein, verknüpft durch eine Processing-Stelle, die die Spaltung des Nicht-Hefeproteins von dem Vorläuferpolypeptid gewährleistet, worin die Leadersequenz etwa 25 bis etwa 50 N-terminale Reste des Vorläuferpolypeptids ausmacht und das Signal peptid eines Vorläufers des α-Faktors vom Saccharomyces oder Kluveromyces und eine einzige Glycosylierungsstelle umfaßt, unter Bedingungen, bei denen das Nicht-Hefeprotein exprimiert wird, züchtet.

2. Verfahren nach Anspruch 1, dadurch **gekennzeichnet,** daß das Nicht-Hefeprotein ein Säugerprotein ist und daß der α-Faktor der Saccharomyces-α-Faktor ist.

3. Verfahren nach Anspruch 2, dadurch **gekennzeichnet,** daß das Säugerprotein ein menschlicher insulinartiger Wachstumsfaktor I oder ein Vorläufer von menschlichem Insulin ist.

4. Verfahren nach Anspruch 3, dadurch **gekennzeichnet,** daß der Vorläufer von menschlichem Insulin menschliches Proinsulin ist, oder daß der Vorläufer von menschlichem Insulin die Insulin A-Kette und die Insulin B-Kette in Verknüpfung durch eine von der Hefe erkannte Processing-Stelle, die in vivo gespalten wird, umfaßt.

5. Verfahren nach Anspruch 4, dadurch **gekennzeichnet,** daß die Processing-Stelle durch das KEX2-Genprodukt von Saccharomyces gespalten wird.

6. Verfahren nach einem der Ansprüche 1 bis 5, dadurch **gekennzeichnet,** daß die Hefezelle der Gattung Saccharomyces angehört.

7. Verfahren nach Anspruch 6, dadurch **gekennzeichnet,** daß die Hefezelle S. cerevisiae ist.

8. Verfahren nach Anspruch 7, dadurch **gekennzeichnet,** daß der Vorläufer des α-Faktors der Hefe S. cerevisiae-MFα1 ist.

9. Verfahren nach einem der Ansprüche 1 bis 8, dadurch **gekennzeichnet,** daß die DNA-Konstruktion ein doppelsträngiges DNA-Molekül umfassend eine Region, die ein Vorläuferpolypeptid codiert, das durch eine Wirtshefe sezernierbar ist, umfaßt, wobei die Region unter Bezug auf einen der Stränge die Struktur

5'-AF-CHO-X_n-S-Gen*-3'

besitzt, worin
$AF\text{-}CHO\text{-}X_n$ eine einzige funktionelle Glycosylierungsstelle umfaßt, worin:

AF	ein α-Faktor-Signal von Saccharomyces oder Kluveromyces codiert;
CHO	eine Glycosylierungsstelle codiert;
X_n	ein Polypeptid mit n Aminosäuren in einer Länge, die keine Glycosylierungsstelle oder keine Processing-Stelle enthält, die die Spaltung des Vorläuferpolypeptids in vivo durch Hefe gewährleistet, codiert;
n	eine ganze Zahl von 0 bis etwa 30 ist; und worin
Gen*	ein Nicht-Hefeprotein codiert; und
S	eine Processing-Stelle, die die Spaltung des Vorläuferpolypeptids gewährleistet, codiert.

10. Verfahren nach Anspruch 9, dadurch **gekennzeichnet,** daß AF ein Polypeptid mit einer Länge von etwa 19 bis 23 Aminosäuren codiert.

11. Verfahren nach Anspruch 9 oder 10, dadurch **gekennzeichnet,** daß n eine ganze Zahl von etwa 0 bis etwa 20 ist.

12. Verfahren nach Anspruch 11, dadurch **gekennzeichnet,** daß n eine ganze Zahl von etwa 0 bis etwa 10 ist.

13. Verfahren nach Anspruch 12, dadurch **gekennzeichnet,** daß n eine ganze Zahl von etwa 3 bis etwa 10 ist.

14. Verfahren nach einem der Ansprüche 9 bis 13, dadurch **gekennzeichnet,** daß die Wirtshefe Saccharomyces ist.

15. Verfahren nach einem der Ansprüche 9 bis 14, dadurch **gekennzeichnet,** daß das α-Faktor-Signalpeptid der Hefe ein Saccharomyces-Signalpeptid ist.

16. Verfahren nach einem der Ansprüche 9 bis 15, dadurch **gekennzeichnet,** daß S eine Processing-Stelle, die in vivo von der Wirtshefe erkannt wird, codiert.

17. Verfahren nach Anspruch 16, dadurch **gekennzeichnet,** daß S ein Dipeptid, das von der KEX2-Endopeptidase erkannt wird, codiert.

18. Verfahren nach Anspruch 17, dadurch **gekennzeichnet,** daß das Dipeptid 5'-Lys-Arg-3' oder 5'-Arg-Arg-3' ist.

19. Verfahren nach einem der Ansprüche 9 bis 18, umfassend ein Replicon.

20. Verfahren nach Anspruch 19, dadurch **gekennzeichnet,** daß die Region, die das Vorläuferpolypeptid codiert, unter der Kontrolle von durch die Hefe erkannten Transkriptionsinitiations- und -terminationssequenzen ist, und daß das Replicon ein Hefereplicon ist.

21. Verfahren nach Anspruch 20, dadurch **gekennzeichnet,** daß das Replicon ein Plasmid oder ein Chromosom ist.

22. Verfahren nach einem der Ansprüche 1 bis 21, dadurch **gekennzeichnet,** daß man weiterhin das Nicht-Hefeprotein in gereinigter Form isoliert.

Revendications
Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, IT, LU, NL, SE

1. Cellule de levure comprenant une construction d'ADN qui pourvoit à l'expression et à la sécretion d'une protéine autre que de levure, ladite construction d'ADN comprenant une séquence codante sous le contrôle de séquences d'initiation et de terminaison de transcription reconnues par la levure, ladite séquence codante codant pour un polypeptide précurseur comprenant une séquence leader et ladite

protéine autre que de levure liée par un site de transformation qui participe au clivage de ladite protéine autre que de levure dudit polypeptide précurseur, dans laquelle ladite séquence leader correspond environ aux 25 à 50 résidus N terminaux dudit polypeptide précurseur et comprend le peptide signal d'un précurseur de facteur alpha de Saccharomyces ou de Kluveromyces et un seul site de glycosylation.

2. Cellule selon la revendication 1 dans laquelle ladite protéine autre que de levure est une protéine mammifère, et ledit facteur alpha est un facteur alpha de Saccharomyces.

3. Cellule selon la revendication 2, dans laquelle la protéine mammifère est le facteur I de croissance de type insuline humaine ou un précurseur d'insuline humaine.

4. Cellule selon la revendication 3, dans laquelle ledit précurseur d'insuline humaine est de la pro-insuline humaine ou dans laquelle ledit précurseur d'insuline humaine comprend une chaîne a d'insuline et une chaîne b d'insuline liées par un site de transformation reconnu par la levure clivé in vivo.

5. Cellule selon la revendication 4, dans laquelle ledit site de transformation est clivé par le produit du gène KEX2 de Saccharomyces.

6. Cellule selon l'une des revendications 1 à 5, dans laquelle ladite cellule de levure est du genre Saccharomyces.

7. Cellule selon la revendication 6, dans laquelle ladite cellule de levure est S. cerevisiae.

8. Cellule selon la revendication 7, dans laquelle ledit précurseur du facteur α de levure est MFα1 de S. cerevisiae.

9. Molecule d'ADN double brins comprenant une région codant pour un polypeptide précurseur pouvant être secreté par une levure-hôte, ladite région, en référence à l'un des brins, présentant la structure suivante :

5'-AF-CHO-X_n-S-Gène*-3'

dans laquelle AF-CHO-X_n comprend un seul site de glycosylation fonctionnel dans lequel : AF code pour un signal de facteur α de Saccharomyces ou de Kluveromyces ;
CHO code pour un site de glycosylation ;
X_n code pour un polypeptide d'une longueur de n acides aminés qui ne contient pas de site de glycosylation ou de site de transformation participant au clivage dudit polypeptide précurseur in vivo par la levure ;
n est un nombre entier compris entre 0 et environ 30 ;
Gène* code pour une protéine autre que de levure ; et
S code pour un site de transformation qui participe au clivage dudit polypeptide précurseur.

10. Molécule d'ADN selon la revendication 9, dans laquelle AF code pour un polypeptide d'une longueur d'environ 19 à 23 acides aminés.

11. Molécule d'ADN selon la revendication 9 ou 10, dans laquelle n est un nombre entier compris entre 0 et environ 20.

12. Molécule d'ADN selon la revendication 11, dans laquelle n est un nombre entier compris environ entre 0 et environ 10.

13. Molécule d'ADN selon la revendication 12, dans laquelle n est un nombre entier compris entre environ 3 et environ 10.

14. Molécule d'ADN selon l'une des revendications 9 à 13 dans laquelle ladite levure-hôte est un Saccharomyces.

* C L M S *

15. Molécule d'ADN selon l'une des revendications 9 à 14 dans laquelle ledit peptide signal de facteur α de levure est un peptide signal de Saccharomyces.

16. Molécule d'ADN selon l'une des revendications 9 à 15, dans laquelle S code pour un site de transformation reconnu in vivo par ladite levure-hôte.

17. Molécule d'ADN selon la revendication 16, dans laquelle S code pour un dipeptide reconnu par l'endopeptidase KEX2.

18. Molécule d'ADN selon la revendication 17, dans laquelle ledit dipeptide est 5'-Lys-Arg-3' ou 5'-Arg-Arg-3'.

19. Molécule d'ADN selon l'une des revendications 9 à 18 comprenant un réplicon.

20. Molécule d'ADN selon la revendication 19, dans laquelle ladite région codant pour ledit polypeptide précurseur est sous le contrôle de séquences d'initiation et de terminaison de transcription reconnues par la levure, et ledit réplicon est un réplicon de levure.

21. Molécule d'ADN selon la revendication 20, dans laquelle ledit réplicon est un plasmide ou un chromosome.

22. Méthode de fabrication d'une protéine autre que de levure, comprenant la culture d'une cellule de levure selon l'une des revendications 1 à 8 sous des conditions dans lesquelles la protéine autre que de levure est exprimée.

23. Méthode selon la revendication 22, dans laquelle la cellule de levure comprend une molécule d'ADN selon l'une des revendications 9 à 21.

24. Méthode selon la revendication 22 ou 23, comprenant également la récupération de la protéine autre que de levure sous une forme purifiée.

Revendications pour les Etats contractants suivants : ES, GR

1. Méthode de fabrication d'une protéine autre que de levure qui comprend la culture d'une cellule de levure comprenant une construction d'ADN qui pourvoit à l'expression et à la sécretion d'une protéine autre que de levure, ladite construction d'ADN comprenant une séquence codante sous le contrôle de séquences d'initiation et de terminaison de transcription reconnues par la levure, ladite séquence codante codant pour un polypeptide précurseur comprenant une séquence leader et ladite protéine autre que de levure liée par un site de transformation qui participe au clivage de ladite protéine autre que de levure dudit polypeptide précurseur, dans laquelle ladite séquence leader correspond environ aux 25 à 50 résidus N terminaux dudit polypeptide précurseur et comprend le peptide signal d'un précurseur de facteur alpha de Saccharomyces ou de Kluveromyces et un seul site de glycosylation, sous des conditions dans lesquelles la protéine autre que de levure est exprimée.

2. Méthode selon la revendication 1, dans laquelle ladite protéine autre que de levure est une protéine mammifère, et ledit facteur alpha est un facteur alpha de Saccharomyces.

3. Méthode selon la revendication 2, dans laquelle la protéine mammifère est le facteur I de croissance de type insuline humaine ou un précurseur d'insuline humaine.

4. Méthode selon la revendication 3, dans laquelle ledit précurseur d'insuline humaine est de la pro-insuline humaine ou dans laquelle ledit précurseur d'insuline humaine comprend une chaîne a d'insuline et une chaîne b d'insuline liées par un site de transformation reconnu par la levure clivé in vivo.

5. Méthode selon la revendication 4, dans laquelle ledit site de transformation est clivé par le produit du gène KEX2 de Saccharomyces.

* C L M S *

6. Méthode selon l'une des revendications 1 à 5, dans laquelle ladite cellule de levure est du genre Saccharomyces.

7. Méthode selon la revendication 6, dans laquelle ladite cellule de levure est S. cerevisiae.

8. Méthode selon la revendication 7, dans laquelle ledit précurseur du facteur α de levure est MFα1 de S. cerevisiae.

9. Méthode selon l'une des revendications 1 à 8 dans laquelle la construction d'ADN comprend une molécule d'ADN double brins comprenant une région codant pour un polypeptide précurseur pouvant être secreté par une levure-hôte, ladite région, en référence à l'un des brins, présentant la structure suivante :

5'-AF-CHO-X_n-S-Gène*-3'

dans laquelle AF-CHO-X_n comprend un seul site de glycosylation fonctionnel dans lequel : AF code pour un signal de facteur α de Saccharomyces ou de Kluveromyces ;
CHO code pour un site de glycosylation ;
X_n code pour un polypeptide d'une longueur de n acides aminés qui ne contient pas de site de glycosylation ou de site de transformation qui participe au clivage dudit polypeptide précurseur in vivo par la levure ;
n est un nombre entier compris entre 0 et environ 30 ;
Gène* code pour une protéine autre que de levure ; et
S code pour un site de transformation qui participe au clivage dudit polypeptide précurseur.

10. Méthode selon la revendication 9, dans laquelle AF code pour un polypeptide d'une longueur de 19 à 23 acides aminés environ.

11. Méthode selon la revendication 9 ou 10, dans laquelle n est un nombre entier compris entre 0 et environ 20.

12. Méthode selon la revendication 11, dans laquelle n est un nombre entier compris environ entre 0 et environ 10.

13. Méthode selon la revendication 12, dans laquelle n est un nombre entier compris entre environ 3 et environ 10.

14. Méthode selon l'une des revendications 9 à 13 dans laquelle ladite levure-hôte est un Saccharomyces.

15. Méthode selon l'une des revendications 9 à 14 dans laquelle ledit peptide signal de facteur α de levure est un peptide signal de Saccharomyces.

16. Méthode selon l'une des revendications 9 à 15, dans laquelle S code pour un site de transformation reconnu in vivo par ladite levure-hôte.

17. Méthode selon la revendication 16, dans laquelle S code pour un dipeptide reconnu par l'endopeptidase KEX2.

18. Méthode selon la revendication 17, dans laquelle ledit dipeptide est 5'-Lys-Arg-3' ou 5'-Arg-Arg-3'.

19. Méthode selon l'une des revendications 9 à 18 comprenant un réplicon.

20. Méthode selon la revendication 19, dans laquelle ladite région codant pour ledit polypeptide précurseur est sous le contrôle de séquences d' initiation et de terminaison de transcription reconnues par la levure, et ledit réplicon et un réplicon de levure.

21. Méthode selon la revendication 20, dans laquelle ledit réplicon est un plasmide ou un chromosome.

* C L M S *

22. Méthode selon l'une des revendications 1 à 21 comprenant également la récupération de la protéine autre que de levure sous une forme purifiée.