



- (51) International Patent Classification:
A61L 27/24 (2006.01) *A61L 27/56* (2006.01)
- (21) International Application Number:
PCT/US2012/035361
- (22) International Filing Date:
27 April 2012 (27.04.2012)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
61/479,937 28 April 2011 (28.04.2011) US
- (71) Applicant (for all designated States except US): **LIFE-CELL CORPORATION** [US/US]; One Millennium Way, Branchburg, NJ 08876 (US).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): **CHEN, Yi** [US/US]; 62 Canal View Drive, Lawrenceville, NJ 08648 (US).
- (74) Agents: **RACITI, Eric, P.** et al.; Finnegan, Henderson, Farabow, Garrett & Dunner LLP, 901 New York Avenue, Washington, DC 20001-4413 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: METHOD FOR ENZYMATIC TREATMENT OF TISSUE PRODUCTS




FIG. 1A

(57) Abstract: Methods for treating tissue matrices and tissue matrices produced according to the methods are provided. The methods can include treating a tissue matrix with a proteolytic enzyme to produce a desired pliability of the tissue matrix.