



(51) International Patent Classification:

C07H 15/04 (2006.01) C11D 1/66 (2006.01)

(21) International Application Number:

PCT/EP2014/063551

(22) International Filing Date:

26 June 2014 (26.06.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

13174915.2 3 July 2013 (03.07.2013) EP

(71) Applicant: BASF SE [DE/DE]; 67056 Ludwigshafen (DE).

(72) Inventors: BAUER, Frederic; Im Linsenbusch 25, 67146 Deidesheim (DE). ESKUCHEN, Rainer; Alt Langenfeld 42, 40764 Langenfeld (DE). TROPSCHE, Jürgen; Im oberen Berg 81, 67354 Römerberg (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,




BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: MIXTURES OF COMPOUNDS, THEIR PREPARATION, AND USES



(57) Abstract: The current invention is directed towards mixtures of compounds, comprising (A) in the range of from 93 to 97 % by weight compound of general formula (I) (B) in the range of from 3 to 6.5 % by weight compound of general formula (II) (C) in the range of from 0.2 to 0.5 % by weight compound of general formula (III) wherein the integers are defined as follows: R¹ is -(CH₂)_nCH₃, R² is -(CH₂)_{n+2}CH₃, R³ is -(CH₂)_{n+1}CH(CH₃)₂, R⁴ is -(CH₂)_{n-1}CH(CH₃)₂ G¹ selected from monosaccharides with 4 to 6 carbon atoms, x in the range of from 1.1 to 10, n is a number in the range of from 1 to 4.