

**PROCESS FOR PREPARING *N*-(5-(3-(7-(3-FLUOROPHENYL)-3*H*-IMIDAZO[4,5-
C]PYRIDIN-2-YL)-1*H*-INDAZOL-5-YL)PYRIDIN-3-YL)-3-METHYLBUTANAMIDE**

RELATED APPLICATIONS


5 This application claims the benefit of U.S. Provisional Application Nos. 62/344,170, filed June 1, 2016 and 62/418,657, filed November 7, 2016, which are incorporated herein by reference in their entirety.

TECHNICAL FIELD

10 Provided herein is a process for preparing *N*-(5-(3-(7-(3-fluorophenyl)-3*H*-imidazo[4,5-
imidazo[4,5-*c*]pyridin-2-yl)-1*H*-indazol-5-yl)pyridin-3-yl)-3-methylbutanamide, including
amorphous and polymorph forms thereof. Also provided herein are intermediates and
pharmaceutically acceptable salts of *N*-(5-(3-(7-(3-fluorophenyl)-3*H*-imidazo[4,5-*c*]pyridin-2-
yl)-1*H*-indazol-5-yl)pyridin-3-yl)-3-methylbutanamide. This compound is useful for the
15 treatment of various diseases including cancer, abnormal cellular proliferation, angiogenesis,
Alzheimer's disease, osteoarthritis and other Wnt-related diseases.

BACKGROUND

The compound of Formula 1:




(1)

20 *N*-(5-(3-(7-(3-fluorophenyl)-3*H*-imidazo[4,5-*c*]pyridin-2-yl)-1*H*-indazol-5-yl)pyridin-3-yl)-
3-methylbutanamide, is a Wnt inhibitor. The compound of Formula (1) can be prepared as
25 disclosed in U.S. Patent No. 8,252,812, incorporated by reference herein in its entirety. There
exists a need for alternative synthetic procedures for the preparation of the compound of
Formula (1). Such alternative synthetic procedures are disclosed herein.

SUMMARY

Provided herein is a process for preparing a compound of Formula (1)



(1)

5 or a pharmaceutically acceptable salt thereof, the process comprising:

(a) reacting a compound of Formula (2)




(2)

or a salt thereof, wherein:

10 R^1 is a nitrogen protecting group, and

X^1 is a first leaving group;


with a compound of Formula (4)



(4)

15 or a salt thereof, wherein X^2 is a second leaving group;


to prepare a compound of Formula (5)



(5)


or a salt thereof,

- (b) reacting the compound of Formula (5), or the salt thereof, with a compound of
 5 Formula (6)



(6)

or a salt thereof, to prepare a compound of Formula (7)




(7)

10

or a salt thereof; and

- (c) deprotecting the compound of Formula (7), or the salt thereof, to prepare the
 compound of Formula (1), or the pharmaceutically acceptable salt thereof.


Also provided herein is a process for preparing a compound of Formula (1):



(1)

or a pharmaceutically acceptable salt thereof, the process comprising:

(a) reacting a compound of Formula (2)




(2)

or salt thereof, wherein:

R¹ is a nitrogen protecting group, and

X¹ is selected from the group consisting of -Cl, -Br, -I, and -OTf;


10 with a boron reagent to prepare a compound of Formula (3)



(3)

15 or salt thereof, wherein A is selected from the group consisting of a boronic acid, a boronic ester, a boronate, a borinate, a boranate, a boranamide, an N-coordinated boronate, and a trifluoroborate;


(b) reacting the compound of Formula (3), or the salt thereof, with a compound of Formula (4)



(4)

20 or a salt thereof, wherein X² is selected from the group consisting of -Cl, -Br, -I, and -OTf;


to prepare a compound of Formula (5)



(5)


5 or a salt thereof;

(c) reacting the compound of Formula (5), or the salt thereof, with a compound of Formula (6)



(6)

10 or a salt thereof, to prepare a compound of Formula (7)




(7)

or a salt thereof; and

(d) deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1), or the pharmaceutically acceptable salt thereof.


15 Further provided herein is a process for preparing a compound of Formula (1):



(1)


or a salt thereof, the process comprising:

(a) reacting a compound of Formula (8)



(8)


or a salt thereof, with bis(pinacolato)diboron to produce a compound of Formula (9)



(9)


or a salt thereof;

(b) reacting the compound of Formula (9), or the salt thereof, with a compound of Formula (10)



(10)


or a salt thereof, to prepare a compound of Formula (11)



(11)


or a salt thereof;

- (c) reacting the compound of Formula (11), or the salt thereof, with a compound of
 5 Formula (6)



(6)

or a salt thereof, to prepare a compound of Formula (12)




(12)

10

or a salt thereof; and

- (d) deprotecting the compound of Formula (12), or the salt thereof, to prepare the
 compound of Formula (1), or a salt thereof.


The disclosure also provides a process for preparing a compound of Formula (1)



(1)


or a salt thereof, the process comprising:

(a) reacting a compound of Formula (8)



(8)


or a salt thereof, with bis(pinacolato)diboron and Pd(dppf)Cl₂ to produce a compound of Formula (9)



(9)


or a salt thereof;

(b) reacting the compound of Formula (9), or the salt thereof, with a compound of Formula (10)



(10)


or a salt thereof, with $\text{Pd(PPh}_3)_4$ and K_3PO_4 to prepare a compound of Formula (11)



(11)


or a salt thereof;

- 5 (c) reacting the compound of Formula (11), or the salt thereof, with a compound of Formula (6)



(6)

or a salt thereof, to prepare a compound of Formula (12)



(12)

or a salt thereof; and

- (d) deprotecting the compound of Formula (12), or the salt thereof, to prepare the compound of Formula (1), or a salt thereof, wherein deprotecting the compound of Formula
15 (12) to prepare the compound of Formula (1) comprises reacting the compound of Formula (12) with TFA. In some embodiments, the TFA is neat TFA. In some embodiments, the ratio

of mass equivalents of TFA to the compound of Formula (12), or a salt thereof, is about 2:1 to about 16:1. In some embodiments, the ratio of mass equivalents of TFA to the compound of Formula (12), or a salt thereof, is about 7:1 to about 9:1. In some embodiments, the ratio of mass equivalents of TFA to the compound of Formula (12), or a salt thereof, is about 8:1.

5 In some embodiments of the process provided herein, deprotecting the compound of Formula (12), or a salt thereof, is performed at a temperature of about 15°C to about 25°C. In some embodiments, deprotecting the compound of Formula (12), or a salt thereof, is performed at a temperature of about 20°C.

In some embodiments of the process provided herein, deprotecting the compound of Formula (12), or a salt thereof, is performed for a time of about 2 hours to about 7 hours. In some embodiments, deprotecting the compound of Formula (12), or a salt thereof, is performed for a time of about 3 hours to about 7 hours. In some embodiments, deprotecting the compound of Formula (12), or a salt thereof, is performed for a time of about 2 hours to about 4 hours. In some embodiments, deprotecting the compound of Formula (12), or a salt thereof, is performed for a time of about 5 hours. In some embodiments, deprotecting the compound of Formula (12), or a salt thereof, is performed for a time of about 3 hours.

In some embodiments of the process provided herein, deprotecting the compound of Formula (12), or a salt thereof, comprises forming a first mixture; and adding water to the first mixture at a temperature of about 0°C to about 10°C to form a second mixture. In some embodiments, water is added at a temperature of about 5°C to form the second mixture. In some embodiments, the process comprises reslurrying the second mixture for a time of about 0.5 hours to about 1 hour. In some embodiments, the process comprises reslurrying the second mixture for a time of about 0.75 hours. In some embodiments, the process comprises reslurrying the second mixture at a temperature of about 0°C to about 10°C. In some embodiments, the process comprises reslurrying the second mixture at a temperature of about 5°C.

In some embodiments, the process further comprises filtering the second mixture to provide a filtrate. In some embodiments, the process further comprises adding water to the filtrate at a temperature of about 0°C to about 10°C to form a third mixture. In some embodiments, water is added to the filtrate at a temperature of about 5°C to form the third mixture. In some embodiments, the process comprises reslurrying the third mixture at a temperature of about 5°C to about 15°C. In some embodiments, the process comprises reslurrying the third mixture at a temperature of about 10°C. In some embodiments, the process comprises reslurrying the third mixture for a time of about 1 hour to about 2 hours. In some

embodiments, the process comprises reslurrying the third mixture for a time of about 1.5 hours. In some embodiments, the process further comprises filtering the third mixture to provide a first residual solid.

In some embodiments of the process provided herein, the process further comprises
5 adding ethanol to the first residual solid to form a fourth mixture. In some embodiments, the process comprises reslurrying the fourth mixture at a temperature of about 25°C to about 35°C. In some embodiments, the process comprises reslurrying the fourth mixture at a temperature of about 30°C. In some embodiments, the process comprises reslurrying the fourth mixture for a time of about 2 hours to about 4 hours. In some embodiments, the process comprises
10 reslurrying the fourth mixture for a time of about 3 hours. In some embodiments, the process comprises filtering the fourth mixture to provide a second residual solid.

In some embodiments of the process provided herein, the process comprises adding water to the second residual solid to form a fifth mixture. In some embodiments, the process comprises reslurrying the fifth mixture at a temperature of about 20°C to about 30°C. In some
15 embodiments, the process comprises reslurrying the fifth mixture at a temperature of about 25°C. In some embodiments, the process comprises reslurrying the fifth mixture for a time of about 0.5 hour to about 1.5 hours. In some embodiments, the process comprises reslurrying the fifth mixture for a time of about 1 hour.

In some embodiments of the process provided herein, the process comprises adding a
20 base to the fifth mixture to form a sixth mixture. In some embodiments, the base is selected from the group consisting of: lithium carbonate, sodium carbonate, potassium carbonate, cesium carbonate, sodium hydrogen carbonate, potassium hydrogen carbonate, sodium sulfate, potassium sulfate, cesium sulfate, lithium phosphate, sodium phosphate, potassium phosphate, and cesium phosphate. In some embodiments, the base is sodium carbonate. In some
25 embodiments, the process comprises reslurrying the sixth mixture at a temperature of about 20°C to about 30°C. In some embodiments, the process comprises reslurrying the sixth mixture at a temperature of about 25°C. In some embodiments, the process comprises reslurrying the sixth mixture for a time of about 5 hours to about 7 hours. In some embodiments, the process comprises reslurrying the sixth mixture for a time of about 6 hours. In some embodiments, the
30 process further comprises filtering the sixth mixture to provide a third residual solid.


In some embodiments of the process provided herein, the process further comprises adding water to the third residual solid to form a seventh mixture. In some embodiments, the process comprises reslurrying the seventh mixture at a temperature of about 20°C to about 30°C. In some embodiments, the process comprises reslurrying the seventh mixture at a

temperature of about 25°C. In some embodiments, the process comprises reslurrying the seventh mixture for a time of about 5 hours to about 8 hours. In some embodiments, the process comprises reslurrying the seventh mixture for a time of about 6.5 hours. In some embodiments, the process further comprises filtering the seventh mixture to provide a fourth residual solid.

- 5 In some embodiments of the process provided herein, the process further comprises (a) adding water to the fourth residual solid to provide an eighth mixture; (b) reslurrying the eighth mixture; and (c) filtering the eighth mixture to provide a fifth residual solid. In some embodiments, steps (a)-(c) are performed an additional one or more times.

- 10 In some embodiments of the process provided herein, the process further comprises adding isopropanol to the fifth residual solid to form a ninth mixture. In some embodiments, the process comprises reslurrying the ninth mixture at a temperature of about 20°C to about 30°C. In some embodiments, the process comprises reslurrying the ninth mixture at a temperature of about 25°C. In some embodiments, the process comprises reslurrying the ninth mixture for a time of about 1 hour to about 3 hours. In some embodiments, the process comprises reslurrying the ninth mixture for a time of about 2 hours. In some embodiments, the process further comprises filtering the ninth mixture to provide a sixth residual solid.


Also provided herein is a process for preparing a compound of Formula (1)



(1)


- 20 or a salt thereof, the process comprising:

(a) reacting the compound of Formula (5)




(5)

or a salt thereof, wherein R¹ is a nitrogen protecting group, with a compound of Formula (6)



(6)


or a salt thereof, to prepare a compound of Formula (7)



(7)

or a salt thereof.


Further provided herein is a process for preparing a compound of Formula (1)



(1)


or a salt thereof, the process comprising:

reacting a compound of Formula (2)



(2)

or a salt thereof, wherein X^1 is selected from the group consisting of $-Cl$, $-Br$, $-I$, and $-OTf$, and R^1 is a nitrogen protecting group, with a boron reagent to prepare a compound of Formula (3)




5

(3)

or a salt thereof, wherein:

A is selected from the group consisting of a boronic acid, a boronic ester, a boronate, a borinate, a boranate, a boranamide, an N-coordinated boronate, and a trifluoroborate.


This disclosure also provides a process for preparing a compound of Formula (7)



10

(7)


or a salt thereof, wherein R^1 is nitrogen protecting group, comprising reacting a compound of Formula (5)



15

(5)


or a salt thereof, with a compound of Formula (6)



(6)

or a salt thereof, to prepare a compound of Formula (7), or the salt thereof.

Also provided herein is a process for preparing a compound of Formula (3)




(3)

or a salt thereof, wherein:

R^1 is a nitrogen protecting group, and

A is selected from the group consisting of a boronic acid, a boronic ester, a boronate, a borinate, a boranate, a boranamide, an N-coordinated boronate, and a trifluoroborate;


comprising reacting a compound of Formula (2)



(2)


or a salt thereof, wherein X^1 is selected from the group consisting of $-Cl$, $-Br$, $-I$, and $-OTf$ with a boron reagent to prepare a compound of Formula (3), or the salt thereof.

Further provided herein is a process for preparing a compound of Formula (1)



(1)


or a salt thereof, the process comprising deprotecting a compound of Formula (12)



(12)

or a salt thereof, to prepare the compound of Formula (1) or a salt thereof.


This disclosure provides a process for preparing a pharmaceutical composition,
 5 including mixing (i) a compound of Formula (1)



(1)

or salt thereof, prepared by any of the processes described herein, and (ii) a pharmaceutically acceptable carrier (excipient), to form the composition.

10 Also provided is a process for preparing a polymorph form of a compound of Formula
 (1)




the process comprising: (a) preparing the compound of Formula (1) according to any one of the processes provided herein; and (b) converting the compound of Formula (1) to the polymorph form. In some embodiments, step (b) comprises reslurrying the compound of Formula (1) or a composition comprising the compound of Formula (1) in a solvent or mixture of solvents to generate the polymorph form. In some embodiments, reslurrying is performed at room temperature. In some embodiments, reslurrying is performed at a temperature of about 50°C. In some embodiments, reslurrying is performed at a temperature of about 30°C to about 35°C. In some embodiments, reslurrying is performed for a time of about 10 hours to about 80 hours. In some embodiments, reslurrying is performed for a time of about 58 hours to about 80 hours.

In some embodiments, the process further comprises a filtering step to provide the polymorph form as a residual solid.

In some embodiments, the reslurrying comprises a solvent or mixture of solvents selected from methanol, water, or a mixture thereof.

In some embodiments, the polymorph form of the compound of Formula (1) is a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water.

Also provided is a process for preparing a polymorph form of a compound of Formula (1)



(1),

the process comprising converting the compound of Formula (1) to the polymorph form. In some embodiments, the process comprises reslurrying the compound of Formula (1) or a composition comprising the compound of Formula (1) in a solvent or mixture of solvents to generate the polymorph form. In some embodiments, the reslurrying is performed at room temperature. In some embodiments, the reslurrying is performed at a temperature of about 50°C. In some embodiments, the reslurrying is performed at a temperature of about 30°C to about 35°C. In some embodiments, the reslurrying is performed for a time of about 10 hours

to about 80 hours. In some embodiments, the reslurrying is performed for a time of about 58 hours to about 80 hours. In some embodiments, the process further comprises a filtering step to provide the polymorph form as a residual solid.

In some embodiments, the reslurrying comprises a solvent or mixture of solvents
5 selected from methanol, water, or a mixture thereof.

In some embodiments, the polymorph form of the compound of Formula (1) is a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water.

10 DESCRIPTION OF DRAWINGS

FIGS. 1A-1D are scans of polymorph Form 1 of the compound of Formula (I). **FIG. 1A** is an x-ray powder diffraction scan of fully dried Form 1. **FIG. 1B** is a differential scanning calorimetry scan of Form 1. **FIG. 1C** is a thermal gravimetric analysis scan of Form 1. **FIG. 1D** is a dynamic vapor sorption scan of Form 1.

15 **FIGS. 2A-2H** are scans of polymorph Forms 2, 2*, and 2** of the compound of Formula (I). **FIG. 2A** is an x-ray powder diffraction scan of fully dried Form 2. **FIG. 2B** is a differential scanning calorimetry scan of Form 2. **FIG. 2C** is a thermal gravimetric analysis scan of Form 2. **FIG. 2D** is an x-ray powder diffraction scan of fully dried Form 2*. **FIG. 2E** is a differential scanning calorimetry scan of Form 2*. **FIG. 2F** is a thermal gravimetric
20 analysis scan of Form 2*. **FIG. 2G** is an x-ray powder diffraction scan of Form 2*. **FIG. 2H** is a differential scanning calorimetry scan of Form 2**.

FIGS. 3A-3C are scans of polymorph Form 3 of the compound of Formula (I). **FIG. 3A** is an x-ray powder diffraction scan of fully dried Form 3. **FIG. 3B** is a differential scanning calorimetry scan of Form 3. **FIG. 3C** is a thermal gravimetric analysis scan of Form 3.

25 **FIGS. 4A-4I** are scans of polymorph Forms 4, 4*, and 4** of the compound of Formula (I). **FIG. 4A** is an x-ray powder diffraction scan of fully dried Form 4. **FIG. 4B** is a differential scanning calorimetry scan of Form 4. **FIG. 4C** is a thermal gravimetric analysis scan of Form 4. **FIG. 4D** is an x-ray powder diffraction scan of fully dried Form 4*. **FIG. 4E** is a differential scanning calorimetry scan of Form 4*. **FIG. 4F** is a thermal gravimetric analysis scan of Form
30 4*. **FIG. 4G** is an x-ray powder diffraction scan of Form 4*. **FIG. 4H** is a differential scanning calorimetry scan of Form 4*. **FIG. 4I** is a thermal gravimetric analysis scan of Form 4**.

FIGS. 5A-5D are scans of polymorph Forms 5 and 5* of the compound of Formula (I). **FIG. 5A** is an x-ray powder diffraction scan of fully dried Form 5. **FIG. 5B** is a differential

scanning calorimetry scan of Form 5. **FIG. 5C** is a thermal gravimetric analysis scan of Form 5. **FIG. 5D** is an x-ray powder diffraction scan of Form 5*.

FIGS. 6A and 6B are scans of polymorph Form 6 of the compound of Formula (I). **FIG. 6A** is an x-ray powder diffraction scan of Form 6. **FIG. 6B** is a differential scanning calorimetry scan of Form 6.

FIGS. 7A-7C are scans of polymorph Form 7 of the compound of Formula (I). **FIG. 7A** is an x-ray powder diffraction scan of fully dried Form 7. **FIG. 7B** is a differential scanning calorimetry scan of Form 7. **FIG. 7C** is a thermal gravimetric analysis scan of Form 7.

FIGS. 8A-8C are scans of polymorph Form 8 of the compound of Formula (I). **FIG. 8A** is an x-ray powder diffraction scan of fully dried Form 8. **FIG. 8B** is a differential scanning calorimetry scan of Form 8. **FIG. 8C** is a thermal gravimetric analysis scan of Form 8.

FIGS. 9A-9D are scans of polymorph Form 9 of the compound of Formula (I). **FIG. 9A** is an x-ray powder diffraction scan of fully dried Form 9. **FIG. 9B** is a differential scanning calorimetry scan of Form 9. **FIG. 9C** is a thermal gravimetric analysis scan of Form 9. **FIG. 9D** is a dynamic vapor sorption scan of Form 9.

FIGS. 10A-10E are scans of polymorph Forms 10 and 10* of the compound of Formula (I). **FIG. 10A** is an x-ray powder diffraction scan of fully dried Form 10. **FIG. 10B** is a differential scanning calorimetry scan of Form 10. **FIG. 10C** is a thermal gravimetric analysis scan of Form 10. **FIG. 10D** is an x-ray powder diffraction scan of Form 10*. **FIG. 10E** is a differential scanning calorimetry scan of Form 10*.

FIGS. 11A-11F are scans of polymorph Forms 11 and 11* of the compound of Formula (I). **FIG. 11A** is an x-ray powder diffraction scan of fully dried Form 11. **FIG. 11B** is a differential scanning calorimetry scan of Form 11. **FIG. 11C** is a thermal gravimetric analysis scan of Form 11. **FIG. 11D** is an x-ray powder diffraction scan of fully dried Form 11*. **FIG. 11E** is a differential scanning calorimetry scan of Form 11*. **FIG. 11F** is a thermal gravimetric analysis scan of Form 11*.

FIGS. 12A-12C are scans of Form 12, an example of a non-stoichiometric hydrate of polymorph Form 1 of the compound of Formula (I). **FIG. 12A** is an x-ray powder diffraction scan of Form 12. **FIG. 12B** is a differential scanning calorimetry scan of Form 12. **FIG. 12C** is a thermal gravimetric analysis scan of Form 12.

FIGS. 13A-13D are scans of Form 13, an example of a non-stoichiometric hydrate of polymorph Form 1 of the compound of Formula (I). **FIG. 13A** is an x-ray powder diffraction scan of Form 13. **FIG. 13B** is a differential scanning calorimetry scan of Form 13. **FIG. 13C**

is a thermal gravimetric analysis scan of Form 13. **FIG. 13D** is a dynamic vapor sorption scan of Form 13.

DETAILED DESCRIPTION

5 1. Definitions

It is appreciated that certain features of the disclosure, which are, for clarity, described in the context of separate embodiments, can also be provided in combination in a single embodiment. Conversely, various features of the disclosure which are, for brevity, described in the context of a single embodiment, can also be provided separately or in any suitable
10 subcombination.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art to which this disclosure belongs. In the event that there is a plurality of definitions for terms cited herein, those in this section prevail unless otherwise stated. All patents, applications, published applications, and
15 other publications cited herein are incorporated by reference in their entirety.

For the terms “for example” and “such as,” and grammatical equivalences thereof, the phrase “and without limitation” is understood to follow unless explicitly stated otherwise. As used herein, the term “about” is meant to account for variations due to experimental error. As used herein, the singular forms “a,” “an,” and “the” include plural referents unless the
20 context clearly dictates otherwise.

The term “salt” includes any ionic form of a compound and one or more counter-ionic species (cations and/or anions). Salts also include zwitterionic compounds (*i.e.*, a molecule containing one more cationic and anionic species, *e.g.*, zwitterionic amino acids). Counter ions present in a salt can include any cationic, anionic, or zwitterionic species. Exemplary anions
25 include, but are not limited to: chloride, bromide, iodide, nitrate, sulfate, bisulfate, sulfite, bisulfite, phosphate, acid phosphate, perchlorate, chlorate, chlorite, hypochlorite, periodate, iodate, iodite, hypoiodite, carbonate, bicarbonate, isonicotinate, acetate, trichloroacetate, trifluoroacetate, lactate, salicylate, citrate, tartrate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucaronate, saccharate, formate,
30 benzoate, glutamate, methanesulfonate, trifluoromethanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, *p*-trifluoromethylbenzenesulfonate, hydroxide, aluminates, and borates. Exemplary cations include, but are not limited to: monovalent alkali metal cations, such as lithium, sodium, potassium, and cesium, and divalent alkaline earth metals, such as beryllium, magnesium, calcium, strontium, and barium. Also included are

transition metal cations, such as gold, silver, copper and zinc, as well as non-metal cations, such as ammonium salts.

Compounds provided herein can also include all isotopes of atoms occurring in the intermediates or final compounds. Isotopes include those atoms having the same atomic number but different mass numbers. For example, isotopes of hydrogen include hydrogen, 5 deuterium, and tritium.

The term “compound,” as used herein, is meant to include all stereoisomers, geometric isomers, tautomers, and isotopes of the structures depicted. Compounds herein identified by name or structure as one particular tautomeric form are intended to include other tautomeric 10 forms unless otherwise specified.

The term “polymorph,” as used herein, refers to crystals of the same molecule having different physical properties as a result of the order of the molecules in the crystal lattice. Polymorphs of a single compound have one or more different chemical, physical, mechanical, electrical, thermodynamic, and/or biological properties from each other. Differences in 15 physical properties exhibited by polymorphs can affect pharmaceutical parameters such as storage stability, compressibility, density (important in composition and product manufacturing), dissolution rates (an important factor in determining bio-availability), solubility, melting point, chemical stability, physical stability, powder flowability, water sorption, compaction, and particle morphology. Differences in stability can result from changes 20 in chemical reactivity (*e.g.* differential oxidation, such that a dosage form discolors more rapidly when comprised of one polymorph than when comprised of another polymorph) or mechanical changes (*e.g.*, crystal changes on storage as a kinetically favored polymorph converts to a thermodynamically more stable polymorph) or both (*e.g.*, one polymorph is more hygroscopic than the other). As a result of solubility/dissolution differences, some transitions 25 affect potency and/or toxicity. In addition, the physical properties of the crystal may be important in processing; for example, one polymorph might be more likely to form solvates or might be difficult to filter and wash free of impurities (*i.e.*, particle shape and size distribution might be different between one polymorph relative to the other). “Polymorph” does not include amorphous forms of the compound. As used herein, “amorphous” refers to a noncrystalline 30 form of a compound which may be a solid state form of the compound or a solubilized form of the compound. For example, “amorphous” refers to a compound without a regularly repeating arrangement of molecules or external face planes.

The term “anhydrous,” as used herein, refers to a crystal form of the compound of Formula (I) that has 1% or less by weight water. For example, 0.5% or less, 0.25% or less, or 0.1% or less by weight water.

5 The term “solvate” as used herein refers to a crystalline form of a compound of Formula (I), such as a polymorph form of the compound, where the crystal lattice comprises one or more solvents of crystallization.

The term “non-stoichiometric hydrate” refers to a crystalline form of a compound of Formula I that comprises water, but wherein variations in the water content do not cause significant changes to the crystal structure. In some embodiments, a non-stoichiometric hydrate
10 can refer to a crystalline form of a compound of Formula I that has channels or networks throughout the crystal structure into which water molecules can diffuse. During drying of non-stoichiometric hydrates, a considerable proportion of water can be removed without significantly disturbing the crystal network, and the crystals can subsequently rehydrate to give the initial non-stoichiometric hydrated crystalline form. Unlike stoichiometric hydrates, the
15 dehydration and rehydration of non-stoichiometric hydrates is not accompanied by a phase transition, and thus all hydration states of a non-stoichiometric hydrate represent the same crystal form. In some embodiments, a non-stoichiometric hydrate can have up to about 20% by weight water, such as, about 20%, about 19%, about 18%, about 17%, about 16%, about 15%, about 14%, about 13%, about 12%, about 11%, about 10%, about 9%, about 8%, about 7%,
20 about 6%, about 5%, about 4%, about 3%, about 2%, or greater than 1% water by weight. In some embodiments, a non-stoichiometric hydrate can have between 1% and about 20% by weight water, such as between 1% and about 5%, 1% and about 10%, 1% and about 15%, about 2% and about 5%, about 2% and about 10%, about 2% and about 15%, about 2% and about 20%, about 5% and about 10%, about 5% and about 15%, about 5% and about 20%, about 10%
25 and about 15%, about 10% and about 20%, or about 15% and about 20% by weight water.

In some embodiments the % water by weight in a crystal form, such as a non-stoichiometric hydrate, is determined by the Karl Fischer titration method. In some embodiments, the crystal form is dried prior to Karl Fischer titration.

30 “Purity,” when used in reference to a composition including a polymorph of the compound of Formula (1), refers to the percentage of one specific polymorph form relative to another polymorph form or an amorphous form of the compound of Formula (1) in the referenced composition. For example, a composition comprising polymorph Form 1 having a purity of 90% would comprise 90 weight parts Form 1 and 10 weight parts of other polymorph and/or amorphous forms of the compound of Formula (1).

As used herein, a compound or composition is “substantially free of” one or more other components if the compound or composition contains no significant amount of such other components. Such components can include starting materials, residual solvents, or any other impurities that can result from the preparation of and/or isolation of the compounds and compositions provided herein. In some embodiments, a polymorph form provided herein is substantially free of other polymorph forms. In some embodiments, a particular polymorph of the compound of Formula (1) is “substantially free” of other polymorphs if the particular polymorph constitutes at least about 95% by weight of the compound of Formula (1) present. In some embodiments, a particular polymorph of the compound of Formula (1) is “substantially free” of other polymorphs if the particular polymorph constitutes at least about 97%, about 98%, about 99%, or about 99.5% by weight of the compound of Formula (1) present. In certain embodiments, a particular polymorph of the compound of Formula (1) is “substantially free” of water if the amount of water constitutes no more than about 2%, about 1%, or about 0.5% by weight of the polymorph.

As used herein, a compound is “substantially present” as a given polymorph if at least about 50% by weight of the compound is in the form of that polymorph. In some embodiments, at least about 60% by weight of the compound is in the form of that polymorph. In some embodiments, at least about 70% by weight of the compound is in the form of that polymorph. In some embodiments, at least about 80% by weight of the compound is in the form of that polymorph. In some embodiments, at least about 90% by weight of the compound is in the form of that polymorph. In some embodiments, at least about 95% by weight of the compound is in the form of that polymorph. In some embodiments, at least about 96% by weight of the compound is in the form of that polymorph. In some embodiments, at least about 97% by weight of the compound is in the form of that polymorph. In some embodiments, at least about 98% by weight of the compound is in the form of that polymorph. In some embodiments, at least about 99% by weight of the compound is in the form of that polymorph. In some embodiments, at least about 99.5% by weight of the compound is in the form of that polymorph.

In some embodiments, the compounds provided herein, including salts and amorphous and polymorph forms thereof, are substantially isolated. By “substantially isolated” is meant that the compound is at least partially or substantially separated from the environment in which it was formed or detected. Partial separation can include, for example, a composition enriched in the compounds provided herein. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the compounds

provided herein, or salt or amorphous or polymorph form thereof. Methods for isolating compounds and their salts are routine in the art.

As used herein, the term “inert atmosphere” refers to a substantially oxygen free environment and primarily consists of non-reactive gases. Exemplary inert atmospheres
5 include a nitrogen atmosphere or an argon atmosphere.


The phrase “pharmaceutically acceptable” is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication,
10 commensurate with a reasonable benefit/risk ratio.

The terms “ambient temperature” and “room temperature” or “RT”, as used herein, are understood in the art, and refer generally to a temperature, *e.g.* a reaction temperature, that is about the temperature of the room in which the reaction is carried out, for example, a temperature from about 20°C to about 30°C, typically around 25°C.

Also provided herein are pharmaceutically acceptable salts of the compounds described herein. As used herein, “pharmaceutically acceptable salts” refers to derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic
20 salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts of the compounds provided herein include the conventional non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of the compounds provided herein can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical
25 methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; in some embodiments, a non-aqueous media like ether, ethyl acetate, alcohol (*e.g.*, methanol, ethanol, iso-propanol, or butanol), or acetonitrile (ACN) can be used. Lists of suitable salts can be found in *Remington's Pharmaceutical*
30 *Sciences*, 17th ed., Mack Publishing Company, Easton, Pa., 1985, p. 1418 and *Journal of Pharmaceutical Science*, 66, 2 (1977), each of which is incorporated herein by reference in its entirety. Conventional methods for preparing salt forms are described, for example, in *Handbook of Pharmaceutical Salts: Properties, Selection, and Use*, Wiley-VCH, 2002.

2. Processes for preparing a compound of Formula (1) and intermediates thereof

Provided herein is a process for preparing a compound of Formula (1)




5

(1)


including salts and amorphous and polymorph forms thereof. The process includes:

(a) reacting a compound of Formula (2)




(2)

10 or a salt thereof, wherein R¹ is a nitrogen protecting group, and X¹ is a first leaving group, with a compound of Formula (4)



(4)

or a salt thereof, wherein X² is a second leaving group; to prepare a compound of Formula (5)




15


(5)

or a salt thereof;

(b) reacting the compound of Formula (5), or the salt thereof, with a compound of Formula (6)



5 or a salt thereof, to prepare a compound of Formula (7)



or a salt thereof; and

(c) deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1). In some embodiments, the process further comprises forming a salt of the compound of Formula (1). In some embodiments, the process further comprises preparing a polymorph form of the compound of Formula (1). In some embodiments, the polymorph form of the compound of Formula (1) is Form 1. In some embodiment, the polymorph form of the compound of Formula (1) is a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water.


As used herein, the term “leaving group” refers to an atom or group of atoms that departs with a pair of electrons in heterolytic bond cleavage. A leaving group can be a stable anion of a strong acid, such as a halide (e.g., fluoride, chloride, bromide, iodide) or a sulfonate ester (e.g., methanesulfonate, trifluoromethanesulfonate, 4-methylbenzenesulfonate). A leaving group can also be a group, such as a chloride, a bromide, an iodide, or a trifluoromethanesulfonate group, that is capable of undergoing oxidative addition to palladium during a palladium-catalyzed reaction, such as a palladium-catalyzed Suzuki-Miyaura cross-coupling reaction.

As used herein, the term “nitrogen protecting group” refers to any group that is capable of reversibly protecting a nitrogen functionality (e.g., an amine). Suitable nitrogen protecting groups can be found in the relevant chapters of standard reference works such as *Protective Groups in Organic Chemistry*, J.F.W. McOmie, Plenum Press, London and New York, 1973; 5 *Greene's Protective Groups in Organic Synthesis*, P.G.M. Wuts, fourth edition, Wiley, New York, 2006; and *Protecting Groups*, P.J. Kocienski, third edition, Thieme, New York, 2005.

Non-limiting examples of nitrogen protecting groups include acetyl, benzyl, cumyl, benzhydryl, trityl (Trt), benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), benzyloxymethyl (BOM), pivaloyl-oxy-methyl (POM), trichloroethoxycarbonyl (Troc), 1- 10 adamantyloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, trimethylsilylethoxymethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyridinyl, and pivaloyl.

As an example, the compound of Formula (1), or a salt or amorphous or polymorph 15 form thereof, can be prepared as shown in Scheme 1, wherein X^1 , X^2 , and R^1 are as defined above.

Scheme 1



In some embodiments, reacting a compound of Formula (2), or a salt thereof, with a 20 compound of Formula (4), or a salt thereof, to prepare a compound of Formula (5), or a salt thereof, is performed in the presence of a palladium catalyst, a nickel catalyst, a tin catalyst, a

copper catalyst, or combinations thereof. In some embodiments, reacting a compound of Formula (2), or a salt thereof, with a compound of Formula (4), or a salt thereof, to prepare a compound of Formula (5), or a salt thereof, is performed in the presence of a palladium catalyst.

As used herein, the term “palladium catalyst” refers to a catalyst that contains palladium, such as Pd(0) and Pd(II), and any necessary ligands required for the catalyst to react. The term “palladium catalyst” can also refer to a palladium catalyst on a solid support.

As used herein, the term “nickel catalyst” refers to a catalyst that contains nickel, such as Ni(0) and Ni(II), and any necessary ligands required for the catalyst to react. The term “nickel catalyst” can also refer to a nickel catalyst on a solid support.

As used herein, the term “copper catalyst” refers to a catalyst that contains copper, such as Cu(0) and Cu(II), and any necessary ligands required for the catalyst to react. The term “copper catalyst” can also refer to a copper catalyst on a solid support.

In some embodiments, a compound of Formula (2), or a salt thereof, can be converted into a more reactive intermediate prior to reacting with a compound of Formula (4), or a salt thereof. In some embodiments, a compound of Formula (2), or a salt thereof, can be converted into a more reactive intermediate via borylation prior to reacting with a compound of Formula (4), or a salt thereof.

Reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, can be performed in the presence of Na₂SO₃, elemental sulfur, or a combination thereof. In some embodiments, reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed in the presence of Na₂SO₃. The Na₂SO₃ can be ground Na₂SO₃. For example, the ground Na₂SO₃ can have a particle size of about 100 microns to about 300 microns or about 150 microns to about 250 microns. In some embodiments, the ground Na₂SO₃ has a particle size of about 150 microns to about 250 microns. In some embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed in the presence of SO₂.

Reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, can be performed in the presence of an organic solvent. In some embodiments, the organic solvent is *N*-methyl-2-pyrrolidone, dimethylacetamide, *n*-butanol, or a combination thereof. In some embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of

Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed in the presence of *N*-methyl-2-pyrrolidone.

In some embodiments, reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed under an inert atmosphere. For example, the inert atmosphere can be a nitrogen (N₂) atmosphere or an argon atmosphere. In some embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed under a N₂ atmosphere.

In some embodiments, reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed at a temperature of about 75°C to about 150°C, about 100°C to about 120°C, or about 110°C to about 115°C. In some embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed at a temperature of about 100°C, 105°C, 110°C, 115°C, or about 120°C.

In some embodiments, reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed for a time of about 5 hours to about 10 hours or about 7 hours to about 9 hours. In some embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed for a time of about 5 hours, 6 hours, 7 hours, 8 hours, 9 hours, or about 10 hours.

Reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, can be performed at a temperature of about 75°C to about 150°C or about 100°C to about 120°C for a time of about 5 hours to about 10 hours. In some embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed at a temperature of about 110°C to about 115°C for a time of about 7 hours to about 9 hours.

In some embodiments, the process further includes preparing a salt of the compound of Formula (7) prior to deprotecting the compound of Formula (7) to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof. In some embodiments, the process further includes preparing an oxalate salt, a citrate salt, or an acetate salt of the compound of Formula (7) prior to deprotecting the compound of Formula (7), or the salt

thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof. In some embodiments, the process further includes preparing an oxalate salt of the compound of Formula (7) prior to deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1), or salt or amorphous or polymorph form thereof.

Preparing a salt of a compound can include preparing the salt of the compound from the free base form of the compound; for example by reacting the compound with an acid or preparing it from another salt, by anion exchange.

In some embodiments, the process further includes preparing a free base form of the compound of Formula (7) prior to deprotecting the compound of Formula (7) to prepare the compound of Formula (1), or salt or amorphous or polymorph form thereof.

In some embodiments, the process further includes precipitating the compound of Formula (7), or a salt thereof, in an organic solvent prior to deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof. In some embodiments, the process further includes precipitating the compound of Formula (7), or a salt thereof, in a non-polar organic solvent prior to deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof. In some embodiments, the organic solvent is an aromatic hydrocarbon, such as benzene, toluene, and xylene, or an aliphatic organic solvent, such as hexane, heptane and octane. In some embodiments, the process further includes precipitating the compound of Formula (7), or a salt thereof, in n-heptane prior to deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof.

In some embodiments, prior to deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof, the process further includes one or more of: (a) preparing a salt of the compound of Formula (7); (b) preparing a free base form of the compound of Formula (7) from the salt of the compound of Formula (7); and (c) precipitating the free base form of the compound of Formula (7) in a non-polar organic solvent.


In some embodiments, prior to deprotecting a compound of Formula (7), or a salt thereof, to prepare a compound of Formula (1), or a salt or amorphous or polymorph form thereof, the process further includes one or more of: (a) preparing an oxalate salt, a citrate salt, or an acetate salt of a compound of Formula (7); (b) preparing a free base form of a compound of Formula (7) from the salt of a compound of Formula (7); and (c) precipitating the free base

form of a compound of Formula (7) in an aliphatic organic solvent such as hexane, heptane and octane.

In some embodiments, prior to deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof, the process further includes one or more of: (a) preparing an oxalate salt of the compound of Formula (7); (b) preparing a free base form of the compound of Formula (7) from the oxalate salt of the compound of Formula (7); and (c) precipitating the free base form compound of Formula (7) in n-heptane, prior to deprotecting the compound of Formula (7) to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof.

In some embodiments, the compound of Formula (7), or a salt thereof, is isolated prior to preparing the compound of Formula (1), or a salt or amorphous or polymorph form thereof. In some embodiments, the compound of Formula (7), or a salt thereof, is isolated with a purity of greater than about 90%, 92%, 94%, 96%, 98%, or greater than about 99%. In some embodiments, the compound of Formula (7), or a salt thereof, is isolated with a purity of greater than about 99%. In some embodiments, the purity is determined by chromatography. In some embodiments, the purity is determined by high-performance liquid chromatography (HPLC) analysis.

Also provided herein is a process for preparing a compound of Formula (1):



(1)


including salts and amorphous and polymorph forms thereof. The process includes:

(a) reacting a compound of Formula (2)



(2)


or salt thereof, wherein R^1 is a nitrogen protecting group, and X^1 is a halide or a sulfonate (e.g., $-Cl$, $-Br$, $-I$, or $-OTf$), with a boron reagent to prepare a compound of Formula (3)



(3)

- 5 or salt thereof, wherein A is selected from the group consisting of a boronic acid, a boronic ester, a boronate, a borinate, a boranate, a boranamide, an N-coordinated boronate, and a trifluoroborate;


(b) reacting the compound of Formula (3), or the salt thereof, with a compound of Formula (4)



(4)

10

or salt thereof, wherein X^2 is a halide or a sulfonate (e.g., $-Cl$, $-Br$, $-I$, or $-OTf$); to prepare a compound of Formula (5)




(5)

15

or a salt thereof;


(c) reacting the compound of Formula (5), or the salt thereof, with a compound of Formula (6)



(6)

20

or a salt thereof, to prepare a compound of Formula (7)




(7)

or a salt thereof; and

- (d) deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1). In some embodiments, the process further comprises forming a salt of the compound of Formula (1). In some embodiments, the process further comprises forming a polymorph form of the compound of Formula (1) thereof. In some embodiments, the polymorph form of the compound of Formula (1) is Form 1. In some embodiments, the polymorph form of the compound of Formula (1) is a non-stoichiometric hydrate of polymorph
- 5
- 10

In some embodiments, a compound of Formula (1), or a salt or amorphous or polymorph form thereof, can be prepared as shown in Scheme 2, wherein X¹, X², A, and R¹ are as defined above.

Scheme 2




In some embodiments, the nitrogen protecting group is an acetyl, benzyl, cumyl, benzhydryl, trityl, benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc),
 5 benzyloxymethyl (BOM), pivaloyl-oxy-methyl (POM), trichloroethoxycarbonyl (Troc), 1-adamantyloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, trimethylsilylethoxymethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyridinyl, and pivaloyl. In some embodiments, the nitrogen protecting group is a trityl (triphenylmethyl)
 10 group.

In some embodiments, X^1 is selected from the group consisting of $-Cl$, $-Br$, $-I$, and $-OTf$. In some embodiments, X^1 is $-OTf$.

In some embodiments, X^2 is selected from the group consisting of $-Cl$, $-Br$, $-I$, and $-OTf$. In some embodiments, X^2 is $-OTf$.


15 In some embodiments, the nitrogen protecting group is not a tetrahydropyran-2-yl.

In some embodiments, A is selected from the group consisting of:



of attachment of A.

5 In some embodiments, A is



In some embodiments, X^1 is $-\text{Br}$. In some embodiments, X^1 is $-\text{I}$. In some embodiments, X^1 is not $-\text{I}$.

In some embodiments, X^2 is $-\text{Br}$. In some embodiments, X^2 is $-\text{I}$. In some embodiments, X^2 is not $-\text{I}$.

10 Reacting a compound of Formula (2), or a salt thereof, with the boron reagent to prepare a compound of Formula (3), or a salt thereof, can be performed in the presence of a palladium catalyst. The palladium catalyst can be a $\text{Pd}(0)$ catalyst. In some embodiments, the palladium catalyst is selected from the group consisting of palladium(II)acetate, $\text{Pd}(\text{dppf})\text{Cl}_2$, $\text{Pd}(\text{dba})_2$, tetrakis(triphenylphosphine)palladium(0), $(\text{MeCN})_2\text{PdCl}_2$, and

15 tris(dibenzylideneacetone)dipalladium(0). In some embodiments, the palladium catalyst is $\text{Pd}(\text{dppf})\text{Cl}_2$.

In some embodiments, the ratio of molar equivalents of the palladium catalyst to the compound of Formula (2) is about 0.01:1 to about 0.1:1 or about 0.02:1 to about 0.5:1. The ratio of molar equivalents of the palladium catalyst to the compound of Formula (2) can be

20 about 0.01:1, 0.02:1, 0.03:1, 0.04:1, 0.05:1, or about 0.1:1. In some embodiments, the ratio of molar equivalents of the palladium catalyst to the compound of Formula (2) is 0.03:1.

In some embodiments, reacting a compound of Formula (2), or a salt thereof, with the boron reagent to prepare a compound of Formula (3), or a salt thereof, is performed in the

presence of a polar aprotic solvent. In some embodiments, reacting a compound of Formula (2), or a salt thereof, with the boron reagent to prepare a compound of Formula (3) is performed in the presence of *N,N*-dimethylformamide (DMF), 1,4-dioxane, *N,N*-dimethylacetamide, or combinations thereof.

5 Reacting a compound of Formula (2), or a salt thereof, with the boron reagent to prepare a compound of Formula (3), or a salt thereof, can be performed at a temperature of about 80°C to about 110°C, about 85°C to about 100°C, or about 90°C to about 95°C. For example, reacting the compound of Formula (2), or a salt thereof, with the boron reagent to prepare the compound of Formula (3), or a salt thereof, can be performed at a temperature of about 80°C,
 10 85°C, 90°C, 95°C, 100°C, 105°C, or about 110°C. In some embodiments, reacting the compound of Formula (2), or a salt thereof, with the boron reagent to prepare the compound of Formula (3), or a salt thereof, is performed at a temperature of about 92°C.

 Reacting a compound of Formula (2), or a salt thereof, with the boron reagent to prepare a compound of Formula (3) can be performed for a time of about 10 hours to about 25 hours
 15 or about 16 hours to about 20 hours. For example, reacting the compound of Formula (2), or a salt thereof, with the boron reagent to prepare the compound of Formula (3), or a salt thereof, can be performed for a time of about 10 hours, 11 hours, 12 hours, 13 hours, 14 hours, 15 hours, 16 hours, 17 hours, 18 hours, 19 hours, 20 hours, 21 hours, 22 hours, 23 hours, 24 hours, or about 25 hours. In some embodiments, the period of time is between about 15 hours and about
 20 25 hours.

 In some embodiments, reacting a compound of Formula (2), or a salt thereof, with the boron reagent to prepare a compound of Formula (3), or a salt thereof, is performed at a temperature of about 80°C to about 110°C for a time of about 15 hours to about 25 hours. In some embodiments, reacting the compound of Formula (2), or a salt thereof, with the boron
 25 reagent to prepare the compound of Formula (3), or a salt thereof, is performed at a temperature of about 85°C to about 95°C for a time of about 16 hours to about 20 hours.

 Reacting a compound of Formula (3), or a salt thereof, with a compound of Formula (4), or a salt thereof, to prepare a compound of Formula (5), or a salt thereof, can be performed in the presence of a palladium catalyst. In some embodiments, a Suzuki-Miyaura cross-
 30 coupling reaction is used to prepare the compound of Formula (5), or a salt thereof, from the compound of Formula (3), or a salt thereof, and Formula (4), or a salt thereof. In some embodiments, the palladium catalyst is selected from the group consisting of PdCl₂(PPh₃)₂, Pd(*t*-Bu)₃, PdCl₂dppf, CH₂Cl₂, Pd(PPh₃)₄, Pd(OAc)/PPh₃, Cl₂Pd[(Pet₃)₂], Pd(DIPHOS)₂, Cl₂Pd(Bipy), [PdCl (Ph₂PCH₂PPh₂)]₂, Cl₂Pd[P(*o*-tolyl)₃]₂, Pd₂(dba)₃/P(*o*-tolyl)₃,

$\text{Pd}_2(\text{dba})_3/\text{P}(\text{furyl})_3$, $\text{Cl}_2\text{Pd}[\text{P}(\text{furyl})_3]_2$, $\text{Cl}_2\text{Pd}(\text{PMePh}_2)_2$, $\text{Cl}_2\text{Pd}[\text{P}(4\text{-F-Ph})_3]_2$, $\text{Cl}_2\text{Pd}[\text{P}(\text{C}_6\text{F}_5)_3]_2$, $\text{Cl}_2\text{Pd}[\text{P}(2\text{-COOH-Ph})(\text{Ph})_2]_2$, $\text{Pd}[\text{P}(t\text{-Bu})_3]_2$, $\text{PdCl}_2(\text{dppe})$, $\text{PdCl}_2(\text{dppp})$, $\text{PdCl}_2[\text{PCy}_3]_2$, and $\text{Cl}_2\text{Pd}[\text{P}(4\text{-COOH-Ph})(\text{Ph})_2]_2$. In some embodiments, the palladium catalyst is $\text{Pd}(\text{PPh}_3)_4$.

The ratio of molar equivalents of the palladium catalyst to the compound of Formula (3), or a salt thereof, can be about 0.01:1 to about 0.1:1 or about 0.02:1 to about 0.5:1. The ratio of molar equivalents of the palladium catalyst to the compound of Formula (3) or a salt thereof, can be about 0.01:1, 0.02:1, 0.03:1, 0.04:1, 0.05:1, or about 0.1:1. In some embodiments, the ratio of molar equivalents of the palladium catalyst to the compound of Formula (3) is 0.03:1.

Reacting a compound of Formula (3), or a salt thereof, with a compound of Formula (4), or a salt thereof, can be performed in the presence of a base. In some embodiments, the base is selected from the group consisting of Na_2CO_3 , $\text{Ba}(\text{OH})_2$, K_3PO_4 , Cs_2CO_3 , K_2CO_3 , TIOH , KF , CsF , KOtBu , NEt_3 , Bu_4F , and NaOH . In some embodiments, the base is K_3PO_4 . The ratio of base to the compound of Formula (4) can be about 0.5:1, 1:1, 1.5:1, 2:1, 3.0:1, 4:1 or about 5:1. In some embodiments, the ratio of base to the compound of Formula (4), or a salt thereof, is about 3.0:1.

Reacting a compound of Formula (3), or a salt thereof, with a compound of Formula (4), or a salt thereof, to prepare a compound of Formula (5), or a salt thereof, can be performed in the presence of an organic solvent. In some embodiments, the organic solvent is selected from the group consisting of toluene, dimethylsulfoxide (DMSO), dimethylformamide (DMF), acetone, acetonitrile, 1,4-dioxane, dimethylacetamide (DMA), tetrahydrofuran (THF), and dimethoxyethane, or a combination thereof. In some embodiments, reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed in the presence of 1,4-dioxane. In some embodiments, reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed in the presence 1,4-dioxane, DMA, THF, dimethoxyethane, or a combination thereof and water. In some embodiments, reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed in the presence of 1,4-dioxane and water.

In some embodiments, reacting a compound of Formula (3), or a salt thereof, with a compound of Formula (4), or a salt thereof, to prepare a compound of Formula (5), or a salt thereof, is performed under an inert atmosphere. For example, the inert atmosphere can be a nitrogen (N_2) atmosphere or an argon atmosphere. In some embodiments, reacting the

compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed under an N₂ atmosphere.

In some embodiments, reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed at a temperature of about 70°C to about 110°C, 80°C to about 100°C, or about 85°C to about 95°C. In some embodiments, reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed at a temperature of about 70°C, 80°C, 85°C, 90°C, 95°C, 100°C, or about 110°C. In some embodiments, reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed at a temperature of about 90°C.

In some embodiments, reacting a compound of Formula (3), or a salt thereof, with a compound of Formula (4), or a salt thereof, to prepare a compound of Formula (5), or a salt thereof, is performed for a time of about 1 hour to about 5 hours or about 2 to about 3 hours. For example, reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, can be performed for a time of about 1 hour, 2 hours, 3 hours, 4 hours, or about 5 hours.

Reacting a compound of Formula (3) or a salt thereof, with a compound of Formula (4), or a salt thereof, to prepare a compound of Formula (5), or a salt thereof, can be performed at a temperature of about 80°C to about 100°C for a time of about 1 hour to about 5 hours. In some embodiments, reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed at a temperature of about 85°C to about 95°C for a time of about 2 hours to about 3 hours.

In some embodiments, the process further includes precipitating a compound of Formula (5), or a salt thereof, prior to reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof. In some embodiments, the process further includes precipitating a compound of Formula (5) in water prior to reacting a compound of Formula (5) with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof.

In some embodiments, the process further includes preparing a salt of a compound of Formula (5) prior to reacting a compound of Formula (5), or the salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof. For example, the process can further include preparing an oxalate salt, citrate salt, or acetate salt of

the compound of Formula (5) prior to reacting the compound of Formula (5), or the salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof. In some embodiments, the process further includes preparing an oxalate salt of the compound of Formula (5) prior to reacting the compound of Formula (5), or the salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof.

In some embodiments, the process can further include preparing a free base form of a compound of Formula (5) prior to reacting a compound of Formula (5) with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof.

In some embodiments, the process further includes precipitating a compound of Formula (5), or a salt thereof, in an organic solvent prior to reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof. In some embodiments, the process further includes precipitating the compound of Formula (5), or a salt thereof, in a non-polar organic solvent prior to reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof. In some embodiments, the non-polar organic solvent is an aromatic hydrocarbon, such as benzene, toluene and xylene, or an aliphatic organic solvent, such as hexane, heptane and octane. In some embodiments, the process further includes precipitating the compound of Formula (5), or a salt thereof, in n-heptane prior to reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof.

In some embodiments, prior to reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, the process further includes one or more of: (a) precipitating the compound of Formula (5); (b) preparing a salt of the compound of Formula (5) from the precipitated compound of Formula (5); (c) preparing a free base form of the compound of Formula (5) from the salt of the compound of Formula (5); and (d) precipitating the free base form of the compound of Formula (5).

In some embodiments, prior to reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, the process further includes one or more of: (a) precipitating the compound of Formula (5) in water; (b) preparing an oxalate salt, a citrate salt, or an acetate salt of the compound of Formula (5) from the precipitated compound of Formula (5); (c) preparing a free

base form of the compound of Formula (5) from the salt of the compound of Formula (5); and
(d) precipitating the free base form of the compound of Formula (5) in a non-polar organic
solvent. In some embodiments, the non-polar organic solvent is an aliphatic organic solvent.
In some embodiments, the aliphatic organic solvent is selected from the group consisting of
5 hexane, heptane and octane.

In some embodiments, prior to reacting a compound of Formula (5), or a salt thereof,
with the compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or
a salt thereof, the process further includes one or more of: (a) precipitating the compound of
Formula (5) in water; (b) preparing an oxalate salt of the compound of Formula (5) from the
10 precipitated compound of Formula (5); (c) preparing a free base form of the compound of
Formula (5) from the salt of the compound of Formula (5); and (d) precipitating the free base
form of the compound of Formula (5) in n-heptane.

In some embodiments, a compound of Formula (5), or a salt thereof, is isolated prior to
preparing a compound of Formula (7), or a salt thereof. In some embodiments, a compound of
15 Formula (5), or a salt thereof, is isolated with a purity of greater than about 90%, 92%, 94%,
96%, 98%, or greater than about 99%. In some embodiments, the compound of Formula (5),
or a salt thereof, can be isolated with a purity of greater than about 98%. In some embodiments,
the purity is determined by chromatography. In some embodiments, the purity is determined
by high-performance liquid chromatography (HPLC) analysis.

20 Reacting a compound of Formula (5), or a salt thereof, with a compound of Formula
(6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, can be performed
in the presence of Na_2SO_3 , elemental sulfur, or a combination thereof. In some embodiments,
reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a
salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed in the
25 presence of Na_2SO_3 . The Na_2SO_3 can be ground Na_2SO_3 . For example, the ground Na_2SO_3 can
have a particle size of about 100 microns to about 300 microns. In some embodiments, the
ground Na_2SO_3 has a particle size of about 150 microns to about 250 microns. In some
embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of
Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is
30 performed in the presence of SO_2 .

Reacting a compound of Formula (5), or a salt thereof, with a compound of Formula
(6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, can be performed
in the presence of an organic solvent. In some embodiments, the organic solvent is selected
from the group consisting of *N*-methyl-2-pyrrolidone, dimethylacetamide, n-butanol, or a

combination thereof. In some embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed in the presence of *N*-methyl-2-pyrrolidone.

5 In some embodiments, reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed under an inert atmosphere. For example, the inert atmosphere can be a nitrogen (N₂) atmosphere or an argon atmosphere. In some embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed under an N₂ atmosphere.

10 In some embodiments, reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed at a temperature of about 75°C to about 150°C, about 100°C to about 120°C, or about 110°C to about 115°C. For example, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, can be performed at a temperature of about 100°C, 105°C, 110°C, 115°C, or about 120°C.

In some embodiments, reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed for a time of about 5 hours to about 10 hours or about 7 hours to about 9 hours. For example, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, can be performed for a time of about 5 hours, 6 hours, 7 hours, 8 hours, 9 hours, or about 10 hours.

25 Reacting the compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, can be performed at a temperature of about 75°C to about 150°C, such as about 100°C to about 120°C, for a time of about 5 hours to about 10 hours. In some embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed at a temperature of about 110°C to about 115°C for a time of about 7 hours to about 9 hours.

30 In some embodiments, the process further includes preparing a salt of a compound of Formula (7) prior to deprotecting a compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1), or a salt thereof. For example, the process can further include preparing an oxalate salt, a citrate salt, or an acetate salt of the compound of Formula (7) prior to deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of

Formula (1), or a salt thereof. In some embodiments, the process further includes preparing an oxalate salt of the compound of Formula (7) prior to deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof.

5 In some embodiments, the process further includes preparing a free base form of a compound of Formula (7) prior to deprotecting a compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof.

In some embodiments, the process further includes precipitating a compound of Formula (7), or a salt thereof, prior to deprotecting a compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof. In some embodiments, the process further includes precipitating a compound of Formula (7), or a salt thereof, in an organic solvent prior to deprotecting a compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof. For example, the process can further include precipitating the compound of Formula (7), or a salt thereof, in a non-polar organic solvent prior to deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof. In some embodiments, the non-polar organic solvent is an aromatic hydrocarbon, such as benzene, toluene and xylene, or an aliphatic organic solvent, such as hexane, heptane and octane. In some embodiments, the process further includes precipitating the compound of Formula (7), or a salt thereof, in n-heptane prior to deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof.


In some embodiments, prior to deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof, the process further includes one or more of: (a) preparing a salt of the compound of Formula (7); (b) preparing a free base form of the compound of Formula (7) from the salt of the compound of Formula (7); and (c) precipitating the free base form compound of Formula (7) in a non-polar organic solvent.

In some embodiments, prior to deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof, the process further includes one or more of: (a) preparing an oxalate salt, a citrate salt, or an acetate salt of the compound of Formula (7); (b) preparing a free base form of the compound of Formula (7) from the salt of the compound of Formula (7); and (c) precipitating the free base form of the compound of Formula (7) in an aliphatic organic solvent, such as hexane, heptane and octane.

In some embodiments, prior to deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof, the process further includes one or more of: (a) preparing an oxalate salt of the compound of Formula (7); (b) preparing a free base form of the compound of Formula (7) from the oxalate salt of the compound of Formula (7); and (c) precipitating the free base form compound of Formula (7) in n-heptane.

In some embodiments, the compound of Formula (7), or a salt thereof, is isolated prior to preparing the compound of Formula (1), or a salt or amorphous or polymorph form thereof. In some embodiments, the compound of Formula (7), or a salt thereof, is isolated with a purity of greater than about 90%, 92%, 94%, 96%, 98%, or greater than about 99%. In some embodiments, the compound of Formula (7), or a salt thereof, is isolated with a purity of greater than about 99%. In some embodiments, the purity is determined by chromatography. In some embodiments, the purity is determined by high-performance liquid chromatography (HPLC) analysis.


Also provided herein is a process for preparing a compound of Formula (1):



(1)


including salts and amorphous and polymorph forms thereof. The process includes:

(a) reacting a compound of Formula (8)



(8)


or a salt thereof, with bis(pinacolato)diboron to produce a compound of Formula (9)



(9)


or a salt thereof;

- (b) reacting the compound of Formula (9), or the salt thereof, with a compound of
 5 Formula (10)



(10)

or a salt thereof, to prepare a compound of Formula (11)




(11)

10

or a salt thereof;


- (c) reacting the compound of Formula (11), or the salt thereof, with a compound of
 Formula (6)



(6)

15

or a salt thereof, to prepare a compound of Formula (12)




(12)

or a salt thereof; and

- (d) deprotecting the compound of Formula (12), or the salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof.

In some embodiments, a compound of Formula (1), or a salt or amorphous or polymorph form thereof, is prepared as shown in Scheme 3.

Scheme 3



10

In some embodiments, reacting the compound of Formula (8), or a salt thereof, with the boron reagent to prepare the compound of Formula (9), or a salt thereof, is performed in the presence of a palladium catalyst. The palladium catalyst can be selected from the group

consisting of palladium(II)acetate, Pd(dppf)Cl₂, Pd(dba)₂, tetrakis(triphenylphosphine)palladium(0), (MeCN)₂PdCl₂, and tris(dibenzylideneacetone)dipalladium(0). In some embodiments, the palladium catalyst is Pd(dppf)Cl₂.

5 The ratio of molar equivalents of the palladium catalyst to the compound of Formula (8), or a salt thereof, can be about 0.01:1 to about 0.1:1 or about 0.02:1 to about 0.5:1. The ratio of molar equivalents of the palladium catalyst to the compound of Formula (8), or a salt thereof, can be about 0.01:1, 0.02:1, 0.03:1, 0.04:1, 0.05:1, or about 0.1:1. In some embodiments, the ratio of molar equivalents of the palladium catalyst to the compound of Formula (8), or a salt thereof, is 0.03:1.

10 Reacting the compound of Formula (8), or a salt thereof, with the boron reagent to prepare the compound of Formula (9), or a salt thereof, can be performed at a temperature of about 80°C to about 110°C, about 85°C to about 100°C, or about 90°C to about 95°C. For example, reacting the compound of Formula (8), or a salt thereof, with the boron reagent to prepare the compound of Formula (9), or a salt thereof, can be performed at a temperature of about 80°C, 85°C, 90°C, 95°C, 100°C, 105°C, or about 110°C. In some embodiments, reacting the compound of Formula (8), or a salt thereof, with the boron reagent to prepare the compound of Formula (9), or a salt thereof, is performed at a temperature of about 92°C.

15 Reacting the compound of Formula (8), or a salt thereof, with the boron reagent to prepare the compound of Formula (9), or a salt thereof, can be performed for a time of about 15 hours to about 25 hours or about 16 hours to about 20 hours. For example, reacting the compound of Formula (8), or a salt thereof, with the boron reagent to prepare the compound of Formula (9), or a salt thereof, can be performed for a time of about 15 hours, 16 hours, 17 hours, 18 hours, 19 hours, 20 hours, 21 hours, 22 hours, 23 hours, 24 hours, or about 25 hours.

20 In some embodiments, reacting the compound of Formula (8), or a salt thereof, with the boron reagent to prepare the compound of Formula (9), or a salt thereof, is performed at a temperature of about 80°C to about 110°C for a time of about 15 hours to about 25 hours. In some embodiments, reacting the compound of Formula (8), or a salt thereof, with the boron reagent to prepare the compound of Formula (9), or a salt thereof, is performed at a temperature of about 85°C to about 95°C for a time of about 16 hours to about 20 hours.

25 Reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, can be performed in the presence of a palladium catalyst. In some embodiments, a Suzuki-Miyaura cross-coupling reaction is used to prepare the compound of Formula (11), or a salt thereof,

from the compounds of Formula (9), or a salt thereof, and Formula (10), or a salt thereof. In some embodiments, the palladium catalyst is selected from the group consisting of $\text{PdCl}_2(\text{PPh}_3)_2$, $\text{Pd}(\text{t-Bu})_3$, PdCl_2dppf , CH_2Cl_2 , $\text{Pd}(\text{PPh}_3)_4$, $\text{Pd}(\text{OAc})/\text{PPh}_3$, $\text{Cl}_2\text{Pd}[(\text{Pet}_3)_2]$, $\text{Pd}(\text{DIPHOS})_2$, $\text{Cl}_2\text{Pd}(\text{Bipy})$, $[\text{PdCl}(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]_2$, $\text{Cl}_2\text{Pd}[\text{P}(\text{o-tolyl})_3]_2$, $\text{Pd}_2(\text{dba})_3/\text{P}(\text{o-tolyl})_3$, $\text{Pd}_2(\text{dba})_3/\text{P}(\text{furyl})_3$, $\text{Cl}_2\text{Pd}[\text{P}(\text{furyl})_3]_2$, $\text{Cl}_2\text{Pd}(\text{PMePh}_2)_2$, $\text{Cl}_2\text{Pd}[\text{P}(4\text{-F-Ph})_3]_2$, $\text{Cl}_2\text{Pd}[\text{P}(\text{C}_6\text{F}_5)_3]_2$, $\text{Cl}_2\text{Pd}[\text{P}(2\text{-COOH-Ph})(\text{Ph})_2]_2$, $\text{Pd}[\text{P}(t\text{-Bu})_3]_2$, $\text{PdCl}_2(\text{dppe})$, $\text{PdCl}_2(\text{dppp})$, $\text{PdCl}_2[\text{PCy}_3]_2$, and $\text{Cl}_2\text{Pd}[\text{P}(4\text{-COOH-Ph})(\text{Ph})_2]_2$. In some embodiments, the palladium catalyst is $\text{Pd}(\text{PPh}_3)_4$.

The ratio of molar equivalents of the palladium catalyst to the compound of Formula (9), or a salt thereof, can be about 0.01:1 to about 0.1:1 or about 0.02:1 to about 0.5:1. The ratio of molar equivalents of the palladium catalyst to the compound of Formula (9), or a salt thereof, can be about 0.01:1, 0.02:1, 0.03:1, 0.04:1, 0.05:1, or about 0.1:1. In some embodiments, the ratio of molar equivalents of the palladium catalyst to the compound of Formula (9), or a salt thereof, is 0.03:1.

In some embodiments, reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed in the presence of a base. The base can be selected from the group consisting of Na_2CO_3 , $\text{Ba}(\text{OH})_2$, K_3PO_4 , Cs_2CO_3 , K_2CO_3 , TlOH , KF , CsF , Bu_4F , and NaOH . In some embodiments, the base is K_3PO_4 .

Reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, can be performed in the presence of an organic solvent. For example, the organic solvent can be toluene, dimethylsulfoxide (DMSO), dimethylformamide (DMF), acetone, acetonitrile, 1,4-dioxane, dimethylacetamide (DMA), tetrahydrofuran (THF), dimethoxyethane, or a combination thereof. In some embodiments, reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed in the presence of 1,4-dioxane. In some embodiments, reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed in the presence 1,4-dioxane, DMA, THF, dimethoxyethane, or a combination thereof and water. In some embodiments, reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed in the presence of 1,4-dioxane and water.

In some embodiments, reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed under an inert atmosphere. For example, the inert atmosphere can be a nitrogen (N₂) atmosphere or an argon atmosphere. In some embodiments, reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed under a N₂ atmosphere.

In some embodiments, reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed at a temperature of about 70°C to about 110°C, 80°C to about 100°C, or about 85°C to about 95°C. For example, reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, can be performed at a temperature of about 70°C, 80°C, 85°C, 90°C, 95°C, 100°C, or about 110°C. In some embodiments, reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed at a temperature of about 90°C.

Reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, can be performed at a temperature of about 80°C to about 100°C for a time of about 1 hour to about 5 hours. In some embodiments, reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed at a temperature of about 85°C to about 95°C for a time of about 2 hours to about 3 hours.

In some embodiments, reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed in the presence of a palladium catalyst and a base; under an inert atmosphere; and performed at a temperature of about 85°C to about 95°C for a time of about 2 hours to about 3 hours.

In some embodiments, the process further includes precipitating the compound of Formula (11) in water prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.

In some embodiments, the process further includes preparing a salt of the compound of Formula (11) prior to reacting the compound of Formula (11), or the salt thereof, with the

compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof. For example, the process can further include preparing an oxalate salt, citrate salt, or acetate salt of the compound of Formula (11) prior to reacting the compound of Formula (11), or the salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof. In some embodiments, the process further includes preparing an oxalate salt of the compound of Formula (11), or a salt thereof, prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.

In some embodiments, the process further includes preparing a free base form of the compound of Formula (11) prior to reacting the compound of Formula (11) with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.

In some embodiments, the process further includes precipitating the compound of Formula (11), or a salt thereof, in an organic solvent prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof. For example, the process can further include precipitating the compound of Formula (11), or a salt thereof, in a non-polar organic solvent prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof. In some embodiments, the non-polar organic solvent is an aromatic hydrocarbon, such as benzene, toluene and xylene, or an aliphatic organic solvent, such as hexane, heptane and octane. In some embodiments, the process further includes precipitating the compound of Formula (11), or a salt thereof, in n-heptane prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.

In some embodiments, prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, the process further includes one or more of: (a) precipitating the compound of Formula (11) in water; (b) preparing a salt of the compound of Formula (11) from the precipitated compound of Formula (11); (c) preparing a free base form of the compound of Formula (11) from the salt of the compound of Formula (11); and (d) precipitating the free base form of the compound of Formula (11) in a non-polar organic solvent prior to reacting the compound of Formula (11) with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.

In some embodiments, prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, the process further includes one or more of (a) precipitating the compound of Formula (11) in water; (b) preparing an oxalate salt of the compound of Formula (11) from the precipitated compound of Formula (11); (c) preparing a free base form of the compound of Formula (11) from the salt of the compound of Formula (11); and (d) precipitating the free base form of the compound of Formula (11) in n-heptane.

In some embodiments, the compound of Formula (11), or a salt thereof, is isolated prior to preparing the compound of Formula (11), or a salt thereof. In some embodiments, the compound of Formula (11), or a salt thereof, is isolated with a purity of greater than about 90%, 92%, 94%, 96%, 98%, or greater than about 99%. In some embodiments, the compound of Formula (11), or a salt thereof, is isolated with a purity of greater than about 98%. In some embodiments, the purity is determined by chromatography. In some embodiments, the purity is determined by high-performance liquid chromatography (HPLC) analysis.

Reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, can be performed in the presence of Na_2SO_3 , elemental sulfur, or a combination thereof. In some embodiments, reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, is performed in the presence of Na_2SO_3 . The Na_2SO_3 can be ground Na_2SO_3 . For example, the ground Na_2SO_3 can have a particle size of about 100 microns to about 300 microns or about 150 microns to about 250 microns. In some embodiments, the ground Na_2SO_3 has a particle size of about 150 microns to about 250 microns. In some embodiments, reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, is performed in the presence of SO_2 .

Reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, can be performed in the presence of an organic solvent. In some embodiments, the organic solvent is *N*-methyl-2-pyrrolidone, dimethylacetamide, n-butanol, or a combination thereof. In some embodiments, reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, is performed in the presence of *N*-methyl-2-pyrrolidone.

In some embodiments, reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a

salt thereof, is performed under an inert atmosphere. For example, the inert atmosphere can be a nitrogen (N₂) atmosphere or an argon atmosphere. In some embodiments, reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, is performed under an N₂ atmosphere.

In some embodiments, reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, is performed at a temperature of about 100°C to about 120°C or about 110°C to about 115°C. For example, reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, can be performed at a temperature of about 100°C, 105°C, 110°C, 115°C, or about 120°C.

In some embodiments, reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, is performed for a time of about 5 hours to about 10 hours or about 7 hours to about 9 hours. For example, reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, can be performed for a time of about 5 hours, 6 hours, 7 hours, 8 hours, 9 hours, or about 10 hours.

Reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, can be performed at a temperature of about 75°C to about 150°C, such as 100°C to about 120°C, for a time of about 5 hours to about 10 hours. In some embodiments, reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, is performed at a temperature of about 110°C to about 115°C for a time of about 7 hours to about 9 hours.

In some embodiments, the process further includes preparing a salt of the compound of Formula (12) prior to deprotecting the compound of Formula (12), or the salt thereof, to prepare the compound of Formula (1), or a salt thereof. For example, the process can further include preparing an oxalate salt, a citrate salt, or an acetate salt of the compound of Formula (12) prior to deprotecting the compound of Formula (12), or the salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof. In some embodiments, the process further includes preparing an oxalate salt of the compound of Formula (12) prior to

deprotecting the compound of Formula (12), or the salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof.

In some embodiments, the process further includes preparing a free base form of the compound of Formula (12) prior to deprotecting the compound of Formula (12) to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof.

In some embodiments, the process further includes precipitating the compound of Formula (12), or a salt thereof, in an organic solvent prior to deprotecting the compound of Formula (12), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof. For example, the process can further include precipitating the compound of Formula (12), or a salt thereof, in an organic solvent prior to deprotecting the compound of Formula (12), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof. In some embodiments, the process further includes precipitating the compound of Formula (12), or a salt thereof, in n-heptane prior to deprotecting the compound of Formula (12), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof.

In some embodiments, prior to deprotecting the compound of Formula (12), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof, the process further includes one or more of: (a) preparing a salt of the compound of Formula (12); (b) preparing a free base form of the compound of Formula (12) from the salt of the compound of Formula (12); and (c) precipitating the free base form compound of Formula (12) in a non-polar organic solvent.


In some embodiments, prior to deprotecting the compound of Formula (12), or a salt thereof, to prepare the compound of Formula (1), or a salt or amorphous or polymorph form thereof, the process further includes one or more of: (a) preparing an oxalate salt of the compound of Formula (12); (b) preparing a free base form of the compound of Formula (12) from the oxalate salt of the compound of Formula (12); and (c) precipitating the free base form compound of Formula (12) in n-heptane.

In some embodiments, the compound of Formula (12), or a salt thereof, is isolated prior to preparing the compound of Formula (1), or a salt or amorphous or polymorph form thereof. In some embodiments, the compound of Formula (12), or a salt thereof, is isolated with a purity of greater than about 90%, 92%, 94%, 96%, 98%, or greater than about 99%. In some embodiments, the compound of Formula (12), or a salt thereof, is isolated with a purity of greater than about 99%. In some embodiments, the purity is determined by chromatography.

In some embodiments, the purity is determined by high-performance liquid chromatography (HPLC) analysis.

In some embodiments, the process further includes preparing the compound of Formula (8), or a salt thereof, by a process including protecting the compound (14)


5



(14)


or a salt thereof, with a trityl group. The process can further include preparing compound (14) from compound (13)

10



(13)


or a salt thereof, by reacting compound (13) with sodium nitrite to prepare the compound of Formula (14)



(14)


15 or a salt thereof.

In some embodiments, the process further includes preparing the compound of Formula (8), or a salt thereof, by a process including: (i) reacting a compound of Formula (13)



(13)


20 or a salt thereof, with a nitrite to prepare the compound of Formula (14)




(14)

or a salt thereof; and (ii) protecting the compound of Formula (14), or the salt thereof, with a trityl group to prepare the compound of Formula (8), or the salt thereof. In some embodiments, the nitrite is selected from the group consisting of sodium nitrite, silver nitrite, calcium nitrite, and potassium nitrite.


5 For example, a compound of Formula (8), or a salt thereof, can be prepared as shown in Scheme 4.



In some embodiments, the process further includes preparing the compound of Formula (10) by a process including: (i) reacting a compound of Formula (15)




or a salt thereof, with a compound of Formula (20)



or a salt thereof, wherein X^3 is a leaving group, to prepare a compound of Formula (10), or the salt thereof.

For example, a compound of Formula (10) can be prepared as shown in Scheme 5.




In some embodiments, X^3 can be a halide. In some embodiments, X^3 is $-\text{Cl}$. For example, the compound of Formula (20) can be isovaleryl chloride (i.e., where X^3 is $-\text{Cl}$).

The ratio of molar equivalents of the compound of Formula (10), or a salt thereof, to the compound of Formula (20), or a salt thereof, can be at least 1:1, 1:1.1, 1:1.2, 1:1.3, 1:1.5, or at least 1:2. The ratio of molar equivalents of the compound of Formula (10), or a salt thereof,


to the compound of Formula (20), or a salt thereof, can be about 1:1, 1:1.1, 1:1.2, 1:1.3, 1:1.5, or about 1:2. In some embodiments, the ratio of molar equivalents of the compound of Formula (10), or a salt thereof, to the compound of Formula (20), or a salt thereof, is about 1:1.3.

In some embodiments, the compound of Formula (6), or a salt thereof, is prepared by a process including: (i) reacting a compound of Formula (16)




(16)

or a salt thereof, with Br₂ to prepare a compound of Formula (17)




(17)

or a salt thereof; (ii) reacting the compound of Formula (17), or the salt thereof, with a compound of Formula (18)



(18)

or a salt thereof, to prepare a compound of Formula (19)




(19)

or a salt thereof; and (iii) selectively reducing the nitro group of the compound of Formula (19), or the salt thereof, to prepare a compound of Formula (6), or the salt thereof.

For example, a compound of Formula (6), or a salt thereof, can be prepared as shown in Scheme 6.


Scheme 6




Reacting the compound of Formula (17), or a salt thereof, with the compound of Formula (18), or a salt thereof, to prepare the compound of Formula (19), or a salt thereof, can be performed in the presence of a palladium catalyst. In some embodiments, reacting the compound of Formula (17), or a salt thereof, with the compound of Formula (18), or a salt thereof, to prepare the compound of Formula (19), or a salt thereof, is carried out under an N₂ atmosphere. In some embodiments, reacting the compound of Formula (17), or a salt thereof, with the compound of Formula (18), or a salt thereof, to prepare the compound of Formula (19), or a salt thereof, further comprises a palladium catalyst and is carried out under an N₂ atmosphere.

In some embodiments, deprotecting the compound of Formula (12), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof, is performed in the presence of an acid. In some embodiments, the acid is TFA. The TFA can be neat TFA. In some embodiments, deprotecting the compound of Formula (12), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof, is accomplished by hydrogenolysis.

Also provided herein is a process for preparing a compound of Formula (1)




including salts and amorphous and polymorph forms thereof. The process includes: (a) reacting a compound of Formula (8)




(8)

- 5 or a salt thereof, with bis(pinacolato)diboron and Pd(dppf)Cl₂ to produce a compound of Formula (9)




(9)

- 10 or a salt thereof; (b) reacting the compound of Formula (9), or the salt thereof, with a compound of Formula (10)




(10)

or a salt thereof, with Pd(PPh₃)₄ and K₃PO₄ to prepare a compound of Formula (11)




(11)

or a salt thereof; (c) reacting the compound of Formula (11), or the salt thereof, with a compound of Formula (6)



(6)

5 or a salt thereof, to prepare a compound of Formula (12)




(12)

or a salt thereof; and (d) deprotecting the compound of Formula (12), or the salt thereof, to prepare the compound of Formula (1), wherein deprotecting the compound of Formula (12) to prepare the compound of Formula (1) comprises reacting the compound of Formula (12) with TFA. In some embodiments, the process further comprises forming a salt of the compound of Formula (1). In some embodiments, the process further comprises forming a polymorph form of the compound of Formula (1). In some embodiments, the polymorph form of the compound of Formula (1) is Form 1. In some embodiments, the polymorph form of the compound of Formula (1) is a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water.

In some embodiments, the ratio of molar equivalents of Pd(dppf)Cl₂ to the compound of Formula (8), or a salt thereof, is about 0.01:1 to about 0.1:1 or about 0.02:1 to about 0.5:1. For example, the ratio of molar equivalents of Pd(dppf)Cl₂ to the compound of Formula (8), or a salt thereof, can be about 0.01:1 to about 0.1:1. The ratio of Pd(dppf)Cl₂ to the compound of Formula (8), or a salt thereof, can be about 0.01:1, 0.02:1, 0.03:1, 0.04:1, 0.05:1, or about 0.1:1. In some embodiments, the ratio of molar equivalents of Pd(dppf)Cl₂ to the compound of Formula (8), or a salt thereof, is about 0.03:1.


In some embodiments, the ratio of molar equivalents of $\text{Pd}(\text{PPh}_3)_4$ to the compound of Formula (9), or a salt thereof, is about 0.01:1 to about 0.1:1 or about 0.02:1 to about 0.5:1. For example, the ratio of molar equivalents of $\text{Pd}(\text{PPh}_3)_4$ to the compound of Formula (9), or a salt thereof, can be about 0.01:1 to about 0.1:1. The ratio of $\text{Pd}(\text{PPh}_3)_4$ to the compound of Formula (9), or a salt thereof, can be about 0.01:1, 0.02:1, 0.03:1, 0.04:1, 0.05:1, or about 0.1:1. In some embodiments, the ratio of molar equivalents of $\text{Pd}(\text{PPh}_3)_4$ to the compound of Formula (9), or a salt thereof, is about 0.03:1.

Further provided herein is a process for preparing a compound of Formula (1)




(1)

including salts and amorphous and polymorph forms thereof. The process includes reacting the compound of Formula (5)




(5)

or a salt thereof, with a compound of Formula (6)



(6)

or a salt thereof, wherein R^1 is a nitrogen protecting group, to prepare a compound of Formula (7)




(7)

or a salt thereof.

In some embodiments, the nitrogen protecting group is an acetyl, benzyl, cumyl, benzhydryl, trityl, benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), benzyloxymethyl (BOM), pivaloyl-oxy-methyl (POM), trichloroethoxycarbonyl (Troc), 1-adamantyloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, trimethylsilylethoxymethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyridinyl, and pivaloyl. In some embodiments, the nitrogen protecting group is a trityl group.

In some embodiments, the nitrogen protecting group is not a tetrahydropyran-2-yl.

In some embodiments, the process further includes deprotecting the compound of Formula (7) or a salt thereof to prepare the compound of Formula (1)




(1)

including salts and amorphous and polymorph forms thereof.


In some embodiments, the nitrogen protecting group is a trityl group and deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof, includes reacting the compound of Formula (7), or a salt thereof, with TFA.

Also provided herein is a process for preparing a compound of Formula (1)




(1)

including salts and amorphous and polymorph forms thereof. The process includes reacting a compound of Formula (2)



(2)

or a salt thereof, wherein X^1 is selected from the group consisting of $-Cl$, $-Br$, $-I$, and $-OTf$, and R^1 is an nitrogen protecting group, with a boron reagent to prepare a compound of Formula (3)




(3)

or a salt thereof, wherein A is selected from the group consisting of a boronic acid, a boronic ester, a boronate, a borinate, a boranate, a boranamide, an N-coordinated boronate, and a trifluoroborate.


In some embodiments, the nitrogen protecting group is selected from the group consisting of acetyl, benzyl, cumyl, benzhydryl, trityl, benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), benzyloxymethyl (BOM), pivaloyl-oxy-methyl (POM), trichloroethoxycarbonyl (Troc), 1-adamantyloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, trimethylsilylethoxymethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyridinyl, and pivaloyl. In some embodiments, the nitrogen protecting group is a trityl.

In some embodiments, the nitrogen protecting group is not a tetrahydropyran-2-yl.

In some embodiments, A is selected from the group consisting of:



In some embodiments, A is:




In some embodiments, X^1 is $-\text{Br}$. In some embodiments, X^1 is $-\text{I}$.

Reacting a compound of Formula (2), or a salt thereof, with the boron reagent to prepare
10 a compound of Formula (3), or a salt thereof, can be performed in the presence of a palladium
catalyst. The palladium catalyst can be selected from the group consisting of
palladium(II)acetate, Pd(dppf)Cl₂, Pd(dba)₂, tetrakis(triphenylphosphine)palladium(0),
(MeCN)₂PdCl₂, and tris(dibenzylideneacetone)dipalladium(0). In some embodiments, the
palladium catalyst is Pd(dppf)Cl₂.


15 In some embodiments, the ratio of molar equivalents of the palladium catalyst to a compound of Formula (2), or a salt thereof, is about 0.01:1 to about 0.1:1 or about 0.02:1 to about 0.5:1. The ratio of molar equivalents of the palladium catalyst to the compound of Formula (2), or a salt thereof, can be about 0.01:1, 0.02:1, 0.03:1, 0.04:1, 0.05:1, or about 0.1:1. In some embodiments, the ratio of molar equivalents of the palladium catalyst to the compound
20 of Formula (2), or a salt thereof, is 0.03:1.

In some embodiments, the process further includes isolating the compound of Formula (3), or a salt thereof.

In some embodiments, the process further includes reacting the compound of Formula (3), or the salt thereof, with a compound of Formula (4)



5 or a salt thereof, wherein X² is a leaving group; to prepare a compound of Formula (5)



or a salt thereof.

In some embodiments, X² is selected from the group consisting of -Cl, -Br, -I, and -OTf. In some embodiments, X² is -Br.

In some embodiments, reacting a compound of Formula (3), or a salt thereof, with a compound of Formula (4), or a salt thereof, to prepare a compound of Formula (5), or a salt thereof, is performed in the presence of a palladium catalyst. The palladium catalyst can be selected from the group consisting of PdCl₂(PPh₃)₂, Pd(t-Bu)₃, PdCl₂ dppf CH₂Cl₂, Pd(PPh₃)₄, Pd(OAc)₂/PPh₃, Cl₂Pd[(Pet₃)₂], Pd(DIPHOS)₂, Cl₂Pd(Bipy), [PdCl (Ph₂PCH₂PPh₂)]₂, Cl₂Pd[P(o-tolyl)₃]₂, Pd₂(dba)₃/P(o-tolyl)₃, Pd₂(dba)₃/P(furyl)₃, Cl₂Pd[P(furyl)₃]₂, Cl₂Pd(PMePh₂)₂, Cl₂Pd[P(4-F-Ph)₃]₂, Cl₂Pd[P(C₆F₅)₃]₂, Cl₂Pd[P(2-COOH-Ph)(Ph)₂]₂, Pd[P(t-Bu)₃]₂, PdCl₂(dppe), PdCl₂(dppp), PdCl₂(PCy₃)₂, and Cl₂Pd[P(4-COOH-Ph)(Ph)₂]₂. In some


embodiments, the palladium catalyst is Pd(PPh₃)₄.

The ratio of molar equivalents of the palladium catalyst to the compound of Formula (3), or a salt thereof, can be about 0.01:1 to about 0.1:1 or about 0.02:1 to about 0.5:1. The ratio of molar equivalents of the palladium catalyst to the compound of Formula (3), or a salt thereof, can be about 0.01:1, 0.02:1, 0.03:1, 0.04:1, 0.05:1, or about 0.1:1. In some embodiments, the ratio of molar equivalents of the palladium catalyst to the compound of Formula (3), or a salt thereof, is 0.03:1.

In some embodiments, reacting a compound of Formula (3), or a salt thereof, with a compound of Formula (4), or a salt thereof, is performed in the presence of a base. The base


can be selected from the group consisting of Na_2CO_3 , $\text{Ba}(\text{OH})_2$, K_3PO_4 , Cs_2CO_3 , K_2CO_3 , TlOH , KF , CsF , KOtBu , NEt_3 , Bu_4F , and NaOH . In some embodiments, the base is K_3PO_4 . The ratio of base to the compound of Formula (4), or a salt thereof, can be about 0.5:1, 1:1, 1.5:1, 2:1, 3.0:1, 4:1 or about 5:1. In some embodiments, the ratio of base to the compound of Formula (4), or a salt thereof, is about 3.0:1. In some embodiments, the ratio K_3PO_4 to the compound of Formula (4), or a salt thereof, is about 3.0:1.

Also provided herein is a process for preparing a compound of Formula (7)




(7)

or a salt thereof, wherein R^1 is a nitrogen protecting group. The process includes reacting a compound of Formula (5)



(5)

or a salt thereof, with a compound of Formula (6)



(6)

or a salt thereof, to prepare a compound of Formula (7). In some embodiments, the process further comprises forming a salt of the compound of Formula (7). In some embodiments, the process comprises performing the reaction between a compound of Formula (5) and a compound of Formula (6) in the presence of an acid. In some embodiments, the acid is

hydrochloric acid (HCl). In some embodiments, up to 1 molar equivalent of an acid (with respect to a compound of Formula (6)) is added.

In some embodiments, the nitrogen protecting group is selected from the group consisting of acetyl, benzyl, cumyl, benzhydryl, trityl, benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), benzyloxymethyl (BOM), pivaloyl-oxy-methyl (POM),
5 trichloroethoxycarbonyl (Troc), 1-adamantyloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, trimethylsilylethoxymethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyridinyl, and pivaloyl. In some embodiments, the
10 nitrogen protecting group is a trityl.

In some embodiments, the nitrogen protecting group is not a tetrahydropyran-2-yl.

In some embodiments, reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed in the presence of Na₂SO₃, elemental sulfur, or a combination thereof. In
15 some embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed in the presence of Na₂SO₃. In some embodiments, the Na₂SO₃ is ground Na₂SO₃. For example, the ground Na₂SO₃ can have a particle size of about 100 microns to about 300 microns or about 150 microns to about 250 microns. In some embodiments, the
20 ground Na₂SO₃ has a particle size of about 150 microns to about 250 microns. In some embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed in the presence of SO₂.

In some embodiments, reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed in the presence of an organic solvent. In some embodiments, the organic solvent is *N*-methyl-2-pyrrolidone, dimethylacetamide, n-butanol, or a combination thereof. In
25 some embodiments, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed in the presence of *N*-methyl-2-pyrrolidone.
30

In some embodiments, reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed under an inert atmosphere. For example, the inert atmosphere can be a nitrogen (N₂) atmosphere or an argon atmosphere. In some embodiments, reacting the


compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed under a N₂ atmosphere.

In some embodiments, reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed at a temperature of about 75°C to about 150°C, about 100°C to about 120°C, or about 110°C to about 115°C. For example, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), can be performed at a temperature of about 100°C, 105°C, 110°C, 115°C, or about 120°C.

In some embodiments, reacting a compound of Formula (5), or a salt thereof, with a compound of Formula (6), or a salt thereof, to prepare a compound of Formula (7), or a salt thereof, is performed for a time of about 5 hours to about 10 hours or about 7 hours to about 9 hours. For example, reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, can be performed for a time of about 5 hours, 6 hours, 7 hours, 8 hours, 9 hours, or about 10 hours.


Reacting a compound of Formula (5) or a salt thereof, with a compound of Formula (6) or a salt thereof, to prepare a compound of Formula (7) or a salt thereof, can be performed at a temperature of about 75°C to about 150°C, about 100°C to about 120°C for a time of about 5 hours to about 10 hours. For example, reacting the compound of Formula (5) or a salt thereof, with the compound of Formula (6) or a salt thereof, to prepare the compound of Formula (7) or a salt thereof, can be performed at a temperature of about 110°C to about 115°C for a time of about 7 hours to about 9 hours.

Also provided herein is a process for preparing a compound of Formula (3)



(3)

or a salt thereof, wherein R¹ is a nitrogen protecting group and A is selected from the group consisting of a boronic acid, a boronic ester, a boronate, a borinate, a boranate, a boranamide, an N-coordinated boronate, and a trifluoroborate. The process includes reacting a compound of Formula (2)



(2)



or a salt thereof, with a boron reagent to prepare a compound of Formula (3), or the salt thereof. In some embodiments, X¹ is selected from the group consisting of -Cl, -Br, -I, and -OTf.

5 In some embodiments, the nitrogen protecting group is selected from the group consisting of acetyl, benzyl, cumyl, benzhydryl, trityl, benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), benzyloxymethyl (BOM), pivaloyl-oxy-methyl (POM), trichloroethoxycarbonyl (Troc), 1-adamantylloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, 10 trimethylsilylethoxymethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyridinyl, and pivaloyl.

In some embodiments, the nitrogen protecting group is a trityl group.

In some embodiments, the nitrogen protecting group is not a tetrahydropyran-2-yl.

In some embodiments, A is selected from the group consisting of:



In some embodiments, A is

In some embodiments, X^1 is $-\text{Br}$. In some embodiments, X^1 is $-\text{I}$. In some embodiments, X^1 is not $-\text{I}$.

In some embodiments, X^2 is $-\text{Br}$. In some embodiments, X^2 is $-\text{I}$. In some embodiments, X^2 is not $-\text{I}$.

5 In some embodiments, reacting a compound of Formula (2), or a salt thereof, with the boron reagent to prepare a compound of Formula (3), or a salt thereof, can be performed in the presence of a palladium catalyst. In some embodiments, the palladium catalyst is a $\text{Pd}(0)$ catalyst. In some embodiments, the palladium catalyst is selected from the group consisting of palladium(II)acetate, $\text{Pd}(\text{dppf})\text{Cl}_2$, $\text{Pd}(\text{dba})_2$, tetrakis(triphenylphosphine)palladium(0),
10 $(\text{MeCN})_2\text{PdCl}_2$, and tris(dibenzylideneacetone)dipalladium(0). In some embodiments, the palladium catalyst is $\text{Pd}(\text{dppf})\text{Cl}_2$.

In some embodiments, the ratio of molar equivalents of the palladium catalyst to the compound of Formula (2), or a salt thereof, is about 0.01:1 to about 0.1:1 or about 0.02:1 to about 0.5:1. The ratio of molar equivalents of the palladium catalyst to the compound of
15 Formula (2), or a salt thereof, can be about 0.01:1, 0.02:1, 0.03:1, 0.04:1, 0.05:1, or about 0.1:1. In some embodiments, the ratio of molar equivalents of the palladium catalyst to the compound of Formula (2), or a salt thereof, is 0.03:1.

In some embodiments, reacting a compound of Formula (2), or a salt thereof, with the boron reagent to prepare a compound of Formula (3), or a salt thereof, is performed at a
20 temperature of about 80°C to about 110°C , about 85°C to about 100°C , or about 90°C to about 95°C . In some embodiments, reacting the compound of Formula (2), or a salt thereof, with the boron reagent to prepare the compound of Formula (3) is performed at a temperature of about 80°C , 85°C , 90°C , 95°C , 100°C , 105°C , or about 110°C . Reacting the compound of Formula (2), or a salt thereof, with the boron reagent to prepare the compound of Formula (3), or a salt
25 thereof, can be performed at a temperature of about 92°C .


Reacting a compound of Formula (2), or a salt thereof, with the boron reagent to prepare a compound of Formula (3), or a salt thereof, can be performed for a time of about 15 hours to about 25 hours or about 16 hours to about 20 hours. For example, reacting the compound of Formula (2), or a salt thereof, with the boron reagent to prepare the compound of Formula (3)
30 can be performed for a time of about 15 hours, 16 hours, 17 hours, 18 hours, 19 hours, 20 hours, 21 hours, 22 hours, 23 hours, 24 hours, or about 25 hours.

In some embodiments, reacting a compound of Formula (2), or a salt thereof, with the boron reagent to prepare a compound of Formula (3), or a salt thereof, can be performed at a temperature of about 80°C to about 110°C for a time of about 15 hours to about 25 hours. In

some embodiments, reacting a compound of Formula (2), or a salt thereof, with the boron reagent to prepare a compound of Formula (3), or a salt thereof, is performed at a temperature of about 85°C to about 95°C for a time of about 16 hours to about 20 hours.


In some embodiments, the process further includes isolating the compound of Formula (3), or a salt thereof. In some embodiments, the compound of Formula (3), or a salt thereof, is isolated by filtration.

Also provided herein is a process for preparing a compound of Formula (4)




(4)

or a salt thereof, wherein X^2 is selected from the group consisting of $-Cl$, $-Br$, $-I$, and $-OTf$. The process includes reacting a compound of Formula (21)



(21)


or a salt thereof, wherein X^2 is as defined above, with a compound of Formula (22)



(22)


wherein X^4 is a leaving group, to prepare the compound of Formula (4), or a salt thereof. In some embodiments, the leaving group is a halide or an activated ester.

Also provided herein is a process for preparing a compound of Formula (5)




(5)

or a salt thereof, wherein R^1 is a nitrogen protecting group. The process includes reacting the compound of Formula (3)



(3)


or the salt thereof, wherein A is selected from the group consisting of a boronic acid, a boronic ester, a boronate, a borinate, a boranate, a boranamide, an N-coordinated boronate, and a trifluoroborate, with a compound of Formula (4)



(4)


or salt thereof, wherein X² is selected from the group consisting of -Cl, -Br, -I, and -OTf, to prepare the compound of Formula (5).

Also provide herein is a process for preparing a compound of Formula (1)



(1)

or a salt or amorphous or polymorph form thereof. The process includes deprotecting a compound of Formula (12)




15

(12)

or a salt thereof, to prepare the compound of Formula (1). In some embodiments, the process further comprises forming a salt of the compound of Formula (1). In some embodiments, the process further comprises forming a polymorph form of the compound of Formula (1). In some
 5 embodiments, the polymorph is Form 1. In some embodiments, the polymorph is a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water.


Also provided herein is a process for preparing a polymorph form of a compound of Formula (1)



10

(1).

The process includes deprotecting a compound of Formula (12)



(12)

15 or a salt thereof, to prepare the compound of Formula (1), and preparing a polymorph form of the compound of Formula (1). In some embodiments, the polymorph form of the compound of Formula (1) is Form 1. In some embodiments, the polymorph form of the compound of Formula (1) is a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water.

In some embodiments, deprotecting the compound of Formula (12) to prepare the compound of Formula (1) comprises reacting the compound of Formula (12) with TFA to obtain a compound of Formula (1). In some embodiments, the process further comprises preparing a polymorph form of the compound of Formula (1). In some embodiments, the process comprises converting the compound of Formula (1) to a polymorph form. In some
5 embodiments, the process comprises reslurrying a compound of Formula (1) or a composition comprising the compound of Formula (1) in a solvent or mixture of solvents to generate a polymorph form of the compound of Formula (1) as a residual solid. In some embodiments, the composition comprises a non-stoichiometric hydrate of polymorph Form 1 having between
10 1% and about 20% by weight water. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C. In some embodiments, the process further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

15 The compounds and intermediates provided herein, including salts thereof, can be prepared using known organic synthesis techniques and can be synthesized according to any of numerous possible synthetic routes.


The reactions for preparing the compounds provided herein can be carried out in suitable solvents which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially non-reactive with the starting materials (reactants), the
20 intermediates, or products, at the temperatures at which the reactions are carried out, *e.g.*, temperatures that can range from the solvent's freezing temperature to the solvent's boiling temperature. A given reaction can be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction
25 step can be selected by the skilled artisan.

Preparation of the compounds provided herein can involve the protection and deprotection of various chemical groups. The chemistry of protecting groups can be found, for example, in *Protecting Group Chemistry*, 1st Ed., Oxford University Press, 2000; *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 6th Ed., Wiley-
30 Interscience Publication, 2006; and Petursson, S. *et al.*, "Protecting Groups in Carbohydrate Chemistry," *J. Chem. Educ.*, 74(11), 1297 (1997) (each of which is incorporated herein by reference in its entirety).


Reactions can be monitored according to any suitable method known in the art. For example, product formation can be monitored by spectroscopic means, such as nuclear

magnetic resonance spectroscopy (*e.g.*, ^1H or ^{13}C), infrared spectroscopy, spectrophotometry (*e.g.*, UV-visible), mass spectrometry, or by chromatographic methods such as high performance liquid chromatography (HPLC), liquid chromatography-mass spectrometry (LCMS), or thin layer chromatography (TLC). Compounds can be purified by those skilled in the art by a variety of methods, including high performance liquid chromatography (HPLC) (*“Preparative LC-MS Purification: Improved Compound Specific Method Optimization”* K.F. Blom, *et al.*, *J. Combi. Chem.* 6(6), 874 (2004), which is incorporated herein by reference in its entirety) and normal phase silica chromatography.

In some embodiments, provided herein is a compound, or a pharmaceutically acceptable salt thereof, of Formula **20**:




where:


X^3 is selected from the group consisting of $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, and ; wherein each R^2 is independently selected from alkyl or cycloalkyl.

In some embodiments of Formula **20**, X^3 is Cl .

In some embodiments, a compound, or a pharmaceutically acceptable salt thereof, of Formula **20** has the structure of Formula **22**:



In some embodiments, provided herein is a compound, or a pharmaceutically acceptable salt thereof, of Formula **21**:




where:

X^2 is selected from the group consisting of $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, and $-\text{OTf}$.


In some embodiments of Formula **21**, X^2 is Br .

In some embodiments, a compound, or a pharmaceutically acceptable salt thereof, of Formula **21** has the structure of Formula **15**:



(15)

5 In some embodiments, provided herein is a compound, or a pharmaceutically acceptable salt thereof, of Formula **4**:




(4)

10 where:

X² is selected from the group consisting of -Cl, -Br, -I, and -OTf.

In some embodiments of Formula **4**, X² is Br.


In some embodiments, a compound, or a pharmaceutically acceptable salt thereof, of Formula **4** has the structure of Formula **10**:



(10)

15

In some embodiments, provided herein is a compound, or a pharmaceutically acceptable salt thereof, of Formula **23**:



(23)

20


where:

X² is selected from the group consisting of -Cl, -Br, -I, and -OTf.

In some embodiments of Formula **23**, X² is Br.


In some embodiments, a compound, or a pharmaceutically acceptable salt thereof, of Formula **23** has the structure of Formula **17**:

25



(17)

In some embodiments, provided herein is a compound, or a pharmaceutically acceptable salt thereof, of Formula **24**:




(24)

where:

A is selected from the group consisting of a boronic acid, a boronic ester, a boronate, a borinate, a boranate, a boranamide, an N-coordinated boronate, and a trifluoroborate.


In some embodiments of Formula **24**, A is selected from the group consisting of:




In some embodiments of Formula **24**, A is



In some embodiments of Formula **24**, A is




In some embodiments, a compound, or a pharmaceutically acceptable salt thereof, of Formula **24** has the structure of Formula **18**:



(18)

In some embodiments, provided herein is a compound, or a pharmaceutically acceptable salt thereof, of Formula 2:



(2)

where:


X¹ is selected from the group consisting of -Cl, -Br, -I, and -OTf; and

R¹ is selected from the group consisting of an acetyl, benzyl, cumyl, benzhydryl, trityl, benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), benzyloxymethyl (BOM), pivaloyl-oxy-methyl (POM), trichloroethoxycarbonyl (Troc), 1-adamantyloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, trimethylsilylethoxymethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyridinyl, and pivaloyl.

In some embodiments of Formula 2, X¹ is Br.


In some embodiments of Formula 2, R¹ is a trityl group.

In some embodiments, a compound, or a pharmaceutically acceptable salt thereof, of Formula 2 has the structure of Formula 8:



(8)

In some embodiments, provided herein is a compound, or a pharmaceutically acceptable salt thereof, of Formula 3:




(3)

where:

A is selected from the group consisting of a boronic acid, a boronic ester, a boronate, a
5 borinate, a boranate, a boranamide, an N-coordinated boronate, and a trifluoroborate; and

R¹ is selected from the group consisting of an acetyl, benzyl, cumyl, benzhydryl, trityl, benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), benzyloxymethyl (BOM), pivaloyl-oxy-methyl (POM), trichloroethoxycarbonyl (Troc), 1-adamantyloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, trimethylsilylethoxymethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyridinyl, and pivaloyl.

In some embodiments of Formula 3, A is selected from the group consisting of:




In some embodiments of Formula 3, A is 

In some embodiments of Formula 3, A is




In some embodiments of Formula 3, R¹ is a trityl group.

In some embodiments, a compound, or a pharmaceutically acceptable salt thereof, of Formula 3 has the structure of Formula 9:



(9)

- 5 In some embodiments, provided herein is a compound, or a pharmaceutically acceptable salt thereof, of Formula 5:




(5)

where:

- 10 R¹ is selected from the group consisting of an acetyl, benzyl, cumyl, benzhydryl, trityl, benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), benzyloxymethyl (BOM), pivaloyl-oxy-methyl (POM), trichloroethoxycarbonyl (Troc), 1-adamantyloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, trimethylsilylethoxymethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-
15 methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyridinyl, and pivaloyl.


In some embodiments of Formula 5, R¹ is a trityl group.

In some embodiments, a compound, or a pharmaceutically acceptable salt thereof, of Formula 5 has the structure of Formula 11:



(11)

In some embodiments, provided herein is a compound, or a pharmaceutically acceptable salt thereof, of Formula 7:




(7)

wherein:

R¹ is selected from the group consisting of an acetyl, benzyl, cumyl, benzhydryl, trityl, benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), benzyloxymethyl (BOM),
 10 pivaloyl-oxy-methyl (POM), trichloroethoxycarbonyl (Troc), 1-adamantyloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, trimethylsilylethoxymethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyridinyl, and pivaloyl.

In some embodiments of Formula 7, R¹ is a trityl group.


15 In some embodiments, a compound, or a pharmaceutically acceptable salt thereof, of Formula 7 has the structure of Formula 12:



(12)

3. Process for preparing polymorph forms of the compound of Formula (1)

5 Provided is a process for preparing polymorph forms of the compound of Formula (1):



(1).

The polymorph forms of the compound of Formula (1) include, *e.g.*, solvates, hydrates, non-stoichiometric hydrates, and non-solvated forms of the compound of Formula (1),
10 including, for example, polymorph Forms 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, and 13.

One such polymorph is a polymorph known as Form 1. Form 1 is an anhydrous polymorph of the compound of Formula (I). In one embodiment, Form 1 has an X-ray powder diffraction (XRPD or XRD) pattern, obtained with CuK α 1-radiation, with at least peaks at $^{\circ}2\theta$ values of 6.8 ± 0.2 , 12.4 ± 0.2 , and 18.5 ± 0.2 . In some embodiments, Form 1 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.8 ± 0.2 , 12.4 ± 0.2 , 16.5 ± 0.2 , 18.5 ± 0.2 , and 19.2 ± 0.2 . In some embodiments, Form 1 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.8 ± 0.2 , 9.3 ± 0.2 , 12.4 ± 0.2 , 13.9 ± 0.2 , 16.5 ± 0.2 , 18.5 ± 0.2 , 19.2 ± 0.2 , and 24.6 ± 0.2 . For example, in some
15 embodiments, Form 1 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.8 ± 0.2 , 9.3 ± 0.2 , 12.4 ± 0.2 , 13.9 ± 0.2 , 14.5 ± 0.2 , 16.5 ± 0.2 , 18.5 ± 0.2 , 19.2 ± 0.2 , 20.3 ± 0.2 , and 24.6 ± 0.2 .

20 In some embodiments, a composition comprising polymorph Form 1 is substantially pure. For example, the composition has a purity of at least about 90%. In some embodiments,

the composition has a purity of at least about 95%. In some embodiments, the composition has a purity of at least about 98%. For example, the composition can have a purity of at least 98.5%, 98.6%, 98.7%, 98.8%, 98.9%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or 99.9%. In some embodiments, the composition is substantially free of other forms of the compound of Formula (1). For example, in some embodiments, the composition is substantially free of other anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than about 15% by weight of other forms of the compound of Formula (1). For example, the composition can contain less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of other anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than about 15% by weight of the polymorph Form 9. For example, the composition can contain less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of the polymorph of Form 9. In some embodiments, the composition contains less than about 15% by weight of one or more other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of one or more other forms of the compound of Formula (1). For example, the composition can contain less than about 15% of Form 2, Form 3, Form 4, Form 5, Form 6, Form 7, Form 8, Form 9, Form 10, Form 11, a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water, or a combination of two or more thereof. In some embodiments, the composition contains less than 1% by weight water.

In some embodiments, polymorph Form 1 exhibits an endotherm between about 50-100°C as measured by differential scanning calorimetry (DSC) related to sorbed water. In some embodiments, polymorph Form 1 exhibits a recrystallization event that is observed between about 270-290°C, e.g., around 280°C. In some embodiments, the endotherm and exotherm are observed when using a scan rate of 10°C per minute.

In some embodiments, polymorph Form 1 recrystallizes into polymorph Form 9 with a melting point of around 363°C. In some embodiments, polymorph Form 1 undergoes a total mass loss of about 0.33% before around 100°C, e.g., from about 39°C to about 100°C, as measured by thermal gravimetric analysis (TGA).

Provided herein is a process of preparing polymorph Form 1. In some embodiments, the process comprises drying a composition comprising the compound of Formula (1), including amorphous and polymorph forms thereof, to generate polymorph Form 1. In some embodiments, the composition comprises a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water. In some embodiments, the process

comprises reslurrying a composition comprising the compound of Formula (1), including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate polymorph Form 1 as a residual solid. In some embodiments, the reslurrying takes place at room temperature (RT). In some embodiments, the reslurrying takes place at around 50°C. In some embodiments, the process further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

In some embodiments, the process comprises reslurrying a composition comprising the compound of Formula (1), including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate polymorph Form 1 as a residual solid. In some embodiments, the compound of Formula (1) is a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water. In some embodiments, the solvent is methanol. In some embodiments, the solvent is toluene. In some embodiments, the solvent is heptane. In some embodiments, the solvent is dichloromethane (DCM). In some embodiments, the solvent is water. In some embodiments, the solvent is in a mixture with water, for example the solvent can be a mixture of water and acetonitrile, methanol, ethyl acetate (EA), methyl tert-butyl ether (MtBE), isopropyl alcohol (IPAc), methyl acetate (MA), methyl isobutyl ketone (MIBK), DCM, n-butyl acetate, heptane, toluene, or n-butanol. In some embodiments, the water is present in an amount of about 5% by weight. In some embodiments, the reslurrying takes place at room temperature. In some embodiments, the reslurrying takes place at around 50°C. In some embodiments, the process further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

In some embodiments, the polymorph form of the compound of Formula (1) is a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water. In some embodiments, for example, above 30% relative humidity (RH), Form 1 readily sorbs water and shows a distinctive shift in Form 1 peaks from 6.8 ± 0.2 to 6.2 ± 0.2 and 12.6 ± 0.2 to 11 ± 0.2 . In some embodiments, a non-stoichiometric hydrate of Form 1 comprises up to about 20% by weight water. For example, up to about 20%, about 19%, about 18%, about 17%, about 16%, about 15%, about 14%, about 13%, about 12%, about 11%, about 10%, about 9%, about 8%, about 7%, about 6%, about 5%, about 4%, about 3%, about 2%, or greater than 1% water by weight. In some embodiments, a non-stoichiometric hydrate of Form 1 has between 1 to about 20% water by weight, *e.g.*, between 1% and about 10%, about 5% and about 15%, about

10% and about 20%, 1% and about 5%, about 5% and about 10%, about 10% and about 15%, about 15% and about 20%, or about 17% and about 20% water by weight.

In some embodiments, a composition comprising a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water is substantially pure.

5 For example, the composition can have a purity of at least about 90%. In some embodiments, the composition has a purity of at least about 95%. In some embodiments, the composition has a purity of at least about 98%. For example, the composition can have a purity of at least 98.5%, 98.6%, 98.7%, 98.8%, 98.9%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or 99.9%. In some embodiments, the composition is substantially free of other forms

10 of the compound of Formula (1). For example, in some embodiments, the composition is substantially free of anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of other forms of the compound of Formula (1) (e.g., anhydrous forms of the

15 compound of Formula (1)). In some embodiments, the composition contains less than 20% by weight of polymorph Form 9 having X-ray powder diffraction pattern comprising peaks at 2θ values of 4.9 ± 0.2 , 18.6 ± 0.2 , and 21.1 ± 0.2 . For example, the composition contains less than 15% by weight of Form 9, such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of other forms of the compound of Form 9. In some

20 embodiments, the composition contains less than 15% of one or more other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less of one or more other forms of the compound of Formula (1). For example, the composition can contain less than about 15% of Form 1, Form 2, Form 3, Form 4, Form 5, Form 6, Form 7, Form 8, Form 9, Form 10, Form 11, or a combination of two or

25 more thereof.

One example of a non-stoichiometric hydrate of polymorph Form 1 is referred to as Form 12.

In one embodiment, provided herein is a polymorph Form 12 having an XRPD pattern, obtained with $\text{CuK}\alpha 1$ -radiation, with at least peaks at 2θ positions 6.4 ± 0.2 ,

30 11.0 ± 0.2 , and 18.4 ± 0.2 . In some embodiments, Form 12 has an XRPD pattern with at least peaks at 2θ positions 6.4 ± 0.2 , 9.2 ± 0.2 , 11.0 ± 0.2 , 18.4 ± 0.2 , and 19.7 ± 0.2 . In some embodiments, Form 12 has an XRPD pattern with at least peaks at 2θ positions 6.4 ± 0.2 , 9.2 ± 0.2 , 11.0 ± 0.2 , 15.6 ± 0.2 , 18.4 ± 0.2 , 19.7 ± 0.2 , 24.4 ± 0.2 , and 25.2 ± 0.2 . For example, in some embodiments, Form 12 has an XRPD pattern with at least peaks at 2θ positions

6.4±0.2, 9.2±0.2, 11.0±0.2, 15.6±0.2, 16.1±0.2, 18.4±0.2, 19.7±0.2, 20.8±0.2, 24.4±0.2, and 25.2±0.2.

In some embodiments, provided herein is polymorph Form 12 that exhibits an endotherm between about 50-100°C as measured by DSC. In some embodiments, polymorph Form 12 exhibits an exotherm at around 283°C. In some embodiments, the endotherms and exotherms are observed when using a scan rate of 10°C per minute.

In some embodiments, provided herein is polymorph Form 12 that has a melting point of around 364°C. In some embodiments, polymorph Form 12 undergoes a weight loss of about 1.4% before around 100°C, e.g., from about 30°C to about 100°C, as measured by TGA.

Another example of a non-stoichiometric hydrate of polymorph Form 1 is referred to as Form 13.

In one embodiment, polymorph Form 13 has an XRPD pattern, obtained with CuKα1-radiation, with at least peaks at $^{\circ}2\theta$ values of 6.4±0.2, 11.0±0.2, and 18.4±0.2. In some embodiments, Form 13 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.4±0.2, 9.2±0.2, 11.0±0.2, 18.4±0.2, and 19.7±0.2. In some embodiments, Form 13 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.4±0.2, 9.2±0.2, 11.0±0.2, 15.6±0.2, 18.4±0.2, 19.7±0.2, 24.4±0.2, and 25.2±0.2. For example, in some embodiments, Form 13 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.4±0.2, 9.2±0.2, 11.0±0.2, 15.6±0.2, 16.1±0.2, 18.4±0.2, 19.7±0.2, 20.8±0.2, 24.4±0.2, and 25.2±0.2.

In some embodiments, polymorph Form 13 exhibits an endotherm between about 50-100°C as measured by DSC. In some embodiments, polymorph Form 13 exhibits an exotherm at between about 265-285°C, e.g., around 278°C. For example, in some embodiments, the endotherms and exotherms are observed when using a scan rate of 10°C per minute.

In some embodiments, polymorph Form 13 has a melting point of around 363°C. In some embodiments, polymorph Form 13 undergoes a weight loss of about 1.9% before around 100°C as measured by TGA.

Provided herein are methods of preparing a non-stoichiometric hydrate of polymorph Form 1. In some embodiments, the method comprises reslurrying a composition comprising the compound of Formula (I), including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate a non-stoichiometric hydrate of polymorph Form 1 as a residual solid. In some embodiments, the composition comprising the compound of Formula (I) is a mixture of a non-stoichiometric hydrate of polymorph Form 1 and Form 1. In

some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C. In some embodiments, the method further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

5 In some embodiments, the method comprises reslurrying a composition comprising a mixture of a non-stoichiometric hydrate of polymorph Form 1 and Form 1 in a solvent or mixture of solvents to generate a non-stoichiometric hydrate of polymorph Form 1 as a residual solid. In some embodiments, the solvent is in a mixture with water, for example the solvent can be a mixture of water and acetonitrile, methanol, MtBE, MA, MIBK, DCM, IPAc, n-butyl
10 acetate, heptane, toluene, or n-butanol. In some embodiments, the water is present in an amount of about 5% by weight. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C.

In some embodiments, the polymorph form of the compound of Formula (1) is a polymorph known as Form 2. Form 2 is an anhydrous polymorph of the compound of Formula
15 (1). In one embodiment, polymorph Form 2 has an XRPD pattern, obtained with CuK α 1-radiation, with at least peaks at $^{\circ}2\theta$ values of 7.0 ± 0.2 , 21.5 ± 0.2 , and 22.0 ± 0.2 . In some embodiments, Form 2 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 7.0 ± 0.2 , 18.9 ± 0.2 , 21.5 ± 0.2 , 22.0 ± 0.2 , and 24.2 ± 0.2 . In some embodiments, Form 2 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 7.0 ± 0.2 , 14.1 ± 0.2 , 18.9 ± 0.2 , 19.2 ± 0.2 , 21.5 ± 0.2 ,
20 22.0 ± 0.2 , 24.2 ± 0.2 , and 26.4 ± 0.2 . For example, in some embodiments, Form 2 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 7.0 ± 0.2 , 10.4 ± 0.2 , 14.1 ± 0.2 , 17.6 ± 0.2 , 18.9 ± 0.2 , 19.2 ± 0.2 , 21.5 ± 0.2 , 22.0 ± 0.2 , 24.2 ± 0.2 , and 26.4 ± 0.2 .

In some embodiments, a composition comprising polymorph Form 2 is substantially pure. For example, the composition can have a purity of at least about 90%. In some
25 embodiments, the composition has a purity of at least about 95%. In some embodiments, the composition has a purity of at least about 98%. For example, the composition can have a purity of at least 98.5%, 98.6%, 98.7%, 98.8%, 98.9%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or 99.9%. In some embodiments, the composition is substantially free of other forms of the compound of Formula (1). For example, in some embodiments, the
30 composition is substantially free of other anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of other forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of one or more other

forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of one or more other forms of the compound of Formula (1). For example, the composition can contain less than about 15% of Form 1, Form 3, Form 4, Form 5, Form 6, Form 7, Form 8, Form 9, Form 10, Form 11, a non-stoichiometric hydrate of Form 1, or a combination of two or more thereof.

In some embodiments, polymorph Form 2 exhibits an endotherm between about 50-100°C as measured by DSC. In some embodiments, polymorph Form 2 exhibits an endotherm between about 220-230°C. In some embodiments, polymorph Form 2 exhibits an exotherm between about 233-238°C. In some embodiments, polymorph Form 2 exhibits an exotherm between about 290-295°C. In some embodiments, the endotherms and exotherms are observed when using a scan rate of 10°C per minute.

In some embodiments, polymorph Form 2 has a melting point of around 363°C. In some embodiments, polymorph Form 2 undergoes a weight loss of about 2.7% before around 116°C, *e.g.*, from about 36°C to about 116°C, as measured by TGA.

Provided herein is a process of preparing polymorph Form 2. In some embodiments, the process comprises reslurrying a composition comprising the compound of Formula (1), including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate Form 2 as a residual solid. In some embodiments, the composition comprises a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C. In some embodiments, the process further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

In some embodiments, the process comprises reslurrying a composition comprising a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water in a solvent or mixture of solvents to generate polymorph Form 2 as a residual solid. In some embodiments, the solvent is acetonitrile. In some embodiments, the solvent is ethanol. In some embodiments, the solvent is in a mixture with water, for example the solvent can be a mixture of water and ethanol or water and n-propanol. In some embodiments, the water is present in an amount of about 5% by weight. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C.

In some embodiments, the polymorph form of the compound of Formula (1) is a polymorph known as Form 3. Form 3 is an anhydrous polymorph of the compound of Formula (1). In one embodiment, polymorph Form 3 has an XRPD pattern, obtained with CuK α 1-

radiation, with at least peaks at $^{\circ}2\theta$ values of 7.2 ± 0.2 , 22.2 ± 0.2 , and 24.4 ± 0.2 . In some embodiments, Form 3 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.3 ± 0.2 , 7.2 ± 0.2 , 21.6 ± 0.2 , 22.2 ± 0.2 , and 24.4 ± 0.2 . In some embodiments, Form 3 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.3 ± 0.2 , 7.2 ± 0.2 , 11.0 ± 0.2 , 18.4 ± 0.2 , 19.0 ± 0.2 , 21.6 ± 0.2 , 22.2 ± 0.2 , and 24.4 ± 0.2 . For example, in some embodiments, Form 3 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.3 ± 0.2 , 7.2 ± 0.2 , 11.0 ± 0.2 , 14.2 ± 0.2 , 17.8 ± 0.2 , 18.4 ± 0.2 , 19.0 ± 0.2 , 21.6 ± 0.2 , 22.2 ± 0.2 , and 24.4 ± 0.2 .

In some embodiments, a composition comprising polymorph Form 3 is substantially pure. For example, the composition can have a purity of at least about 90%. In some embodiments, the composition has a purity of at least about 95%. In some embodiments, the composition has a purity of at least about 98%. For example, the composition can have a purity of at least 98.5%, 98.6%, 98.7%, 98.8%, 98.9%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or 99.9%. In some embodiments, the composition is substantially free of other forms of the compound of Formula (1). For example, in some embodiments, the composition is substantially free of other anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of other forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of one or more other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of one or more other forms of the compound of Formula (1). For example, the composition can contain less than about 15% of Form 1, Form 2, Form 4, Form 5, Form 6, Form 7, Form 8, Form 9, Form 10, Form 11, a non-stoichiometric hydrate of Form 1, or a combination of two or more thereof.

In some embodiments, polymorph Form 3 exhibits an exotherm between about 190-220°C, as measured by DSC. In some embodiments, polymorph Form 3 exhibits an exotherm at between about 225-235°C, *e.g.*, around 230°C, as measured by DSC. In some embodiments, polymorph Form 3 exhibits an exotherm at between about 292-300°C, *e.g.*, around 297°C, as measured by DSC. For example, in some embodiments, the endotherms and exotherms are observed when using a scan rate of 10°C per minute.

In some embodiments, polymorph Form 3 has a melting point of around 365°C. In some embodiments, polymorph Form 3 undergoes a weight loss of about 1.6% before around 81°C and a weight loss of about 1.7% between about 81-169°C as measured by TGA.

Provided herein is a process of preparing polymorph Form 3. In some embodiments, the process comprises reslurrying a composition comprising the compound of Formula (1), including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate Form 3 as a residual solid. In some embodiments, the composition comprises a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water. In some
5 embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C. In some embodiments, the process further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

10 In some embodiments, the process comprises reslurrying a composition comprising a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water in a solvent or mixture of solvents to generate polymorph Form 3 as a residual solid. In some embodiments, the solvent is IPAc. In some embodiments, the solvent is n-butyl acetate. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying
15 takes place at around 50°C.

In some embodiments, the polymorph form of the compound of Formula (1) is a polymorph known as Form 4. Form 4 is an anhydrous polymorph of the compound of Formula (1). In one embodiment, polymorph Form 4 has an XRPD pattern, obtained with CuK α 1-radiation, with at least peaks at $^{\circ}2\theta$ values of 7.0 ± 0.2 , 21.8 ± 0.2 , and 25.1 ± 0.2 . In some
20 embodiments, Form 4 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 7.0 ± 0.2 , 19.5 ± 0.2 , 21.8 ± 0.2 , 23.2 ± 0.2 , and 25.1 ± 0.2 . In some embodiments, Form 4 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 7.0 ± 0.2 , 17.6 ± 0.2 , 18.3 ± 0.2 , 19.5 ± 0.2 , 21.8 ± 0.2 , 23.2 ± 0.2 , 25.1 ± 0.2 , and 25.8 ± 0.2 . For example, in some embodiments, Form 4 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 7.0 ± 0.2 , 9.6 ± 0.2 , 17.6 ± 0.2 , 18.3 ± 0.2 , 19.5 ± 0.2 ,
25 21.8 ± 0.2 , 23.2 ± 0.2 , 25.1 ± 0.2 , 25.8 ± 0.2 , and 29.3 ± 0.2 .

In some embodiments, a composition comprising polymorph Form 4 is substantially pure. For example, the composition can have a purity of at least about 90%. In some embodiments, the composition has a purity of at least about 95%. In some embodiments, the composition has a purity of at least about 98%. For example, the composition can have a purity
30 of at least 98.5%, 98.6%, 98.7%, 98.8%, 98.9%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or 99.9%. In some embodiments, the composition is substantially free of other forms of the compound of Formula (1). For example, in some embodiments, the composition is substantially free of other anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of other forms of the

compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of other forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of one or more other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of one or more other forms of the compound of Formula (1). For example, the composition can contain less than about 15% of Form 1, Form 2, Form 3, Form 5, Form 6, Form 7, Form 8, Form 9, Form 10, Form 11, a non-stoichiometric hydrate of Form 1, or a combination of two or more thereof.

In some embodiments, polymorph Form 4 exhibits an endotherm between about 50-100°C as measured by DSC. In some embodiments, polymorph Form 4 exhibits an endotherm at between about 180-215°C. In some embodiments, polymorph Form 4 exhibits an endotherm between about 220-230°C. In some embodiments, polymorph Form 4 exhibits an exotherm at between about 230-240°C, *e.g.*, around 235°C. In some embodiments, polymorph Form 4 exhibits an exotherm at between about 300-310°C. For example, in some embodiments, the endotherms and exotherms are observed when using a scan rate of 10°C per minute.

In some embodiments, polymorph Form 4 has a melting point of between about 366-369°C, *e.g.*, around 367°C. In some embodiments, polymorph Form 4 undergoes a weight loss of about 8.3% before around 200°C, *e.g.*, from about 42°C to about 200°C, as measured by TGA.

Provided herein is a process of preparing polymorph Form 4. In some embodiments, the process comprises reslurrying a composition comprising the compound of Formula (1), including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate Form 4 as a residual solid. In some embodiments, the composition comprises a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C. In some embodiments, the process further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

In some embodiments, the process comprises reslurrying a composition comprising a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water in a solvent or mixture of solvents to generate polymorph Form 4 as a residual solid. In some embodiments, the solvent is EA. In some embodiments, the solvent is MA. In some embodiments, the solvent is MtBE. In some embodiments, the solvent is n-propanol. In some embodiments, the solvent is acetone. In some embodiments, the solvent is in a mixture with

water, for example the solvent can be a mixture of water and MA, EA, or acetone. In some embodiments, the water is present in an amount of about 5% by weight. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C.

5 In some embodiments, the polymorph form of the compound of Formula (1) is a polymorph known as Form 5. Form 5 is an anhydrous polymorph of the compound of Formula (1). In one embodiment, polymorph Form 5 has an XRPD pattern, obtained with CuK α 1-radiation, with at least peaks at $^{\circ}2\theta$ values of 7.3 ± 0.2 , 22.3 ± 0.2 , and 24.5 ± 0.2 . In some
10 embodiments, Form 5 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.3 ± 0.2 , 7.3 ± 0.2 , 21.7 ± 0.2 , 22.3 ± 0.2 , and 24.5 ± 0.2 . In some embodiments, Form 5 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.3 ± 0.2 , 7.3 ± 0.2 , 11.0 ± 0.2 , 19.1 ± 0.2 , 19.5 ± 0.2 , 21.7 ± 0.2 , 22.3 ± 0.2 , and 24.5 ± 0.2 . For example, in some embodiments, Form 5 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.3 ± 0.2 , 7.3 ± 0.2 , 11.0 ± 0.2 , 14.3 ± 0.2 , 19.1 ± 0.2 , 19.5 ± 0.2 , 21.7 ± 0.2 , 22.3 ± 0.2 , 24.5 ± 0.2 , and 26.5 ± 0.2 .

15 In some embodiments, a composition comprising polymorph Form 5 is substantially pure. For example, the composition can have a purity of at least about 90%. In some embodiments, the composition has a purity of at least about 95%. In some embodiments, the composition has a purity of at least about 98%. For example, the composition can have a purity of at least 98.5%, 98.6%, 98.7%, 98.8%, 98.9%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%,
20 99.6%, 99.7%, 99.8%, or 99.9%. In some embodiments, the composition is substantially free of other forms of the compound of Formula (1). For example, in some embodiments, the composition is substantially free of other anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%,
25 5%, 4%, 3%, 2%, 1% or less by weight of other forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of one or more other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of one or more other forms of the compound of Formula (1). For example, the composition can contain less than about 15% of Form 1, Form
30 2, Form 3, Form 4, Form 6, Form 7, Form 8, Form 9, Form 10, Form 11, a non-stoichiometric hydrate of Form 1, or a combination of two or more thereof.

In some embodiments, polymorph Form 5 exhibits an endotherm between about 50-100°C as measured by DSC. In some embodiments, polymorph Form 5 exhibits an endotherm at between about 210-235°C, *e.g.*, around 222°C. In some embodiments, polymorph Form 5

exhibits an exotherm at between about 227-240°C, *e.g.*, around 235°C. In some embodiments, polymorph Form 5 exhibits an exotherm at between about 280-300°C, *e.g.*, around 293°C. For example, in some embodiments, the endotherms and exotherms are observed when using a scan rate of 10°C per minute.

5 In some embodiments, polymorph Form 5 has a melting point of around 363°C. In some embodiments, polymorph Form 5 undergoes a weight loss of about 3.1% before around 100°C and about 1.7% between about 100-250°C as measured by TGA.

Provided herein is a process of preparing polymorph Form 5. In some embodiments, the process comprises reslurrying a composition comprising the compound of Formula (1),
10 including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate Form 5 as a residual solid. In some embodiments, the composition comprises a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C. In some embodiments, the process further comprises drying the residual
15 solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

In some embodiments, the process comprises reslurrying a composition comprising a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water in a solvent or mixture of solvents to generate polymorph Form 5 as a residual solid. In some
20 embodiments, the solvent is MtBE. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C.

In some embodiments, the polymorph form of the compound of Formula (1) is a polymorph known as Form 6. Form 6 is an anhydrous polymorph of the compound of Formula (1).

25 In some embodiments, a composition comprising polymorph Form 6 is substantially pure. For example, the composition can have a purity of at least about 90%. In some embodiments, the composition has a purity of at least about 95%. In some embodiments, the composition has a purity of at least about 98%. For example, the composition can have a purity of at least 98.5%, 98.6%, 98.7%, 98.8%, 98.9%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%,
30 99.6%, 99.7%, 99.8%, or 99.9%. In some embodiments, the composition is substantially free of other forms of the compound of Formula (1). For example, in some embodiments, the composition is substantially free of other anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%,

5%, 4%, 3%, 2%, 1% or less by weight of other forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of one or more other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of one or more other forms of the compound of Formula (1). For example, the composition can contain less than about 15% of Form 1, Form 2, Form 3, Form 4, Form 5, Form 7, Form 8, Form 9, Form 10, Form 11, a non-stoichiometric hydrate of Form 1, or a combination of two or more thereof.

In some embodiments, polymorph Form 6 exhibits an exotherm between about 245-260°C as measured by DSC. For example, in some embodiments, the endotherms and exotherms are observed when using a scan rate of 10°C per minute. In some embodiments, polymorph Form 6 has a melting point of around 364°C.

Provided herein is a process of preparing polymorph Form 6. In some embodiments, the process comprises reslurrying a composition comprising the compound of Formula (1), including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate Form 6 as a residual solid. In some embodiments, the composition comprises a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C. In some embodiments, the process further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

In some embodiments, the process comprises reslurrying a composition comprising a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water in a solvent or mixture of solvents to generate polymorph Form 6 as a residual solid. In some embodiments, the solvent is IPAc. In some embodiments, the solvent is in a mixture with water, for example the solvent can be a mixture of water and IPAc. In some embodiments, the water is present in an amount of about 5% by weight. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C.

In some embodiments, the polymorph of the compound of Formula (1) is a polymorph known as Form 7. Form 7 is an anhydrous polymorph of the compound of Formula (1). In one embodiment, polymorph Form 7 has an XRPD pattern, obtained with CuK α 1-radiation, with at least peaks at $^{\circ}2\theta$ values of 7.1 ± 0.2 , 21.6 ± 0.2 , and 23.2 ± 0.2 . In some embodiments, Form 7 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 4.9 ± 0.2 , 7.1 ± 0.2 , 18.5 ± 0.2 , 21.6 ± 0.2 , and 23.2 ± 0.2 . In some embodiments, Form 7 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 4.9 ± 0.2 , 7.1 ± 0.2 , 10.9 ± 0.2 , 18.5 ± 0.2 , 19.4 ± 0.2 , 21.6 ± 0.2 , 23.2 ± 0.2 , and 30.3 ± 0.2 .

For example, in some embodiments, Form 7 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 4.9 ± 0.2 , 7.1 ± 0.2 , 8.8 ± 0.2 , 10.9 ± 0.2 , 18.5 ± 0.2 , 19.4 ± 0.2 , 21.6 ± 0.2 , 22.1 ± 0.2 , 23.2 ± 0.2 , and 30.3 ± 0.2 .

In some embodiments, a composition comprising polymorph Form 7 is substantially pure. For example, the composition can have a purity of at least about 90%. In some embodiments, the composition has a purity of at least about 95%. In some embodiments, the composition has a purity of at least about 98%. For example, the composition can have a purity of at least 98.5%, 98.6%, 98.7%, 98.8%, 98.9%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or 99.9%. In some embodiments, the composition is substantially free of other forms of the compound of Formula (1). For example, in some embodiments, the composition is substantially free of other anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of other forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of one or more other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of one or more other forms of the compound of Formula (1). For example, the composition can contain less than about 15% of Form 1, Form 2, Form 3, Form 4, Form 5, Form 6, Form 8, Form 9, Form 10, Form 11, a non-stoichiometric hydrate of Form 1, or a combination of two or more thereof.

In some embodiments, polymorph Form 7 exhibits an exotherm between about 227-235°C, *e.g.*, around 232°C, as measured by DSC. In some embodiments, polymorph Form 7 exhibits an exotherm between about 299-305°C, *e.g.*, around 303°C. For example, in some embodiments, the endotherms and exotherms are observed when using a scan rate of 10°C per minute.

In some embodiments, polymorph Form 7 has a melting point of around 365°C. In some embodiments, polymorph Form 7 undergoes a weight loss of about 12% before around 200°C, *e.g.*, from about 36°C to about 200°C, as measured by TGA.

Provided herein is a process of preparing polymorph Form 7. In some embodiments, the process comprises reslurrying a composition comprising the compound of Formula (1), including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate Form 7 as a residual solid. In some embodiments, the composition comprises a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes

place at around 50°C. In some embodiments, the process further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

In some embodiments, the process comprises reslurrying a composition comprising a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water in a solvent or mixture of solvents to generate polymorph Form 7 as a residual solid. In some embodiments, the solvent is methyl ethyl ketone (MEK). In some embodiments, the solvent is in a mixture with water, for example the solvent can be a mixture of water and MEK. In some embodiments, the water is present in an amount of about 5% by weight. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C.

In some embodiments, the polymorph form of the compound of Formula (1) is a polymorph known as Form 8. Form 8 is an anhydrous polymorph of the compound of Formula (1). In one embodiment, polymorph Form 8 has an XRPD pattern, obtained with CuK α 1-radiation, with at least peaks at $^{\circ}2\theta$ values of 6.9 ± 0.2 , 17.7 ± 0.2 , and 21.5 ± 0.2 . In some embodiments, Form 8 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.9 ± 0.2 , 11.5 ± 0.2 , 17.7 ± 0.2 , 21.5 ± 0.2 , and 27.6 ± 0.2 . In some embodiments, Form 8 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.9 ± 0.2 , 11.5 ± 0.2 , 15.3 ± 0.2 , 16.9 ± 0.2 , 17.7 ± 0.2 , 21.5 ± 0.2 , 27.6 ± 0.2 , and 28.9 ± 0.2 . For example, in some embodiments, Form 8 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.9 ± 0.2 , 11.5 ± 0.2 , 12.7 ± 0.2 , 14.2 ± 0.2 , 15.3 ± 0.2 , 16.9 ± 0.2 , 17.7 ± 0.2 , 21.5 ± 0.2 , 27.6 ± 0.2 , and 28.9 ± 0.2 .

In some embodiments, a composition comprising polymorph Form 8 is substantially pure. For example, the composition can have a purity of at least about 90%. In some embodiments, the composition has a purity of at least about 95%. In some embodiments, the composition has a purity of at least about 98%. For example, the composition can have a purity of at least 98.5%, 98.6%, 98.7%, 98.8%, 98.9%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or 99.9%. In some embodiments, the composition is substantially free of other forms of the compound of Formula (1). For example, in some embodiments, the composition is substantially free of other anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of other forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of one or more other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%,

7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of one or more other forms of the compound of Formula (1). For example, the composition can contain less than about 15% of Form 1, Form 2, Form 3, Form 4, Form 5, Form 6, Form 7, Form 9, Form 10, Form 11, a non-stoichiometric hydrate of Form 1, or a combination of two or more thereof.

5 In some embodiments, polymorph Form 8 exhibits an endotherm between about 41-60°C as measured by DSC. In some embodiments, polymorph Form 8 exhibits an exotherm at between about 221-235°C, *e.g.*, around 231°C. In some embodiments, polymorph Form 8 exhibits an endotherm between about 279-290°C, *e.g.*, around 285°C. For example, in some embodiments, the endotherms and exotherms are observed when using a scan rate of 10°C per
10 minute.

In some embodiments, polymorph Form 8 has a melting point of around 364°C. In some embodiments, polymorph Form 8 undergoes a weight loss of about 4.2% before around 190°C and about 3.9% between about 190-261°C as measured by TGA.

Provided herein is a process of preparing polymorph Form 8. In some embodiments,
15 the process comprises reslurrying a composition comprising the compound of Formula (1), including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate Form 8 as a residual solid. In some embodiments, the composition comprises a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water. In some
20 embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C. In some embodiments, the process further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

In some embodiments, the process comprises reslurrying a composition comprising a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water in
25 a solvent or mixture of solvents to generate polymorph Form 8 as a residual solid. In some embodiments, the solvent is MIBK. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C.

In some embodiments, the polymorph form of the compound of Formula (1) is a polymorph known as Form 9. Form 9 is an anhydrous polymorph of the compound of Formula
30 (1). In one embodiment, polymorph Form 9 has an XRPD pattern, obtained with CuK α 1-radiation, with at least peaks at $^{\circ}2\theta$ values of 4.9 ± 0.2 , 18.6 ± 0.2 , and 21.1 ± 0.2 . In some embodiments, Form 9 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 4.9 ± 0.2 , 18.6 ± 0.2 , 21.1 ± 0.2 , 24.1 ± 0.2 , and 25.2 ± 0.2 . In some embodiments, Form 9 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 4.9 ± 0.2 , 15.3 ± 0.2 , 16.5 ± 0.2 , 18.6 ± 0.2 , 21.1 ± 0.2 ,

22.4±0.2, 24.1±0.2, and 25.2±0.2. For example, in some embodiments, Form 9 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 4.9±0.2, 10.1±0.2, 15.3±0.2, 16.5±0.2, 18.6±0.2, 21.1±0.2, 22.4±0.2, 24.1±0.2, 25.2±0.2, and 28.6±0.2.

In some embodiments, a composition comprising polymorph Form 9 is substantially pure. For example, the composition can have a purity of at least about 90%. In some embodiments, the composition has a purity of at least about 95%. In some embodiments, the composition has a purity of at least about 98%. For example, the composition can have a purity of at least 98.5%, 98.6%, 98.7%, 98.8%, 98.9%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or 99.9%. In some embodiments, the composition is substantially free of other forms of the compound of Formula (1). For example, in some embodiments, the composition is substantially free of other anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of other forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of one or more other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of one or more other forms of the compound of Formula (1). For example, the composition can contain less than about 15% of Form 1, Form 2, Form 3, Form 4, Form 5, Form 6, Form 7, Form 8, Form 10, Form 11, a non-stoichiometric hydrate of Form 1, or a combination of two or more thereof.

In some embodiments, polymorph Form 9 exhibits a single melting endotherm at around 364°C as measured by DSC. For example, in some embodiments, the endotherm is observed when using a scan rate of 10°C per minute. In some embodiments, other polymorph forms provided herein, such as, *e.g.*, Form 1 and Form 2, can convert to Form 9 when heated to just before melting (*i.e.*, around 364°C).

In some embodiments, polymorph Form 9 has a melting point of around 364°C. In some embodiments, polymorph Form 9 undergoes a weight loss of about 0.28% before around 100°C, *e.g.*, from about 30.5°C to about 100°C, as measured by TGA.

Provided herein is a process of preparing polymorph Form 9. In some embodiments, the process comprises reslurrying a composition comprising the compound of Formula (1), including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate Form 9 as a residual solid. In some embodiments, the composition comprises a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes

place at around 50°C. In some embodiments, the process further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

In some embodiments, the process comprises reslurrying a composition comprising a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water in a solvent or mixture of solvents to generate polymorph Form 9 as a residual solid. In some embodiments, the solvent is n-butanol. In some embodiments, the solvent is IPAc. In some embodiments, the solvent is n-butyl acetate. In some embodiments, the solvent is in a mixture with water, for example the solvent can be a mixture of water and ethanol or water and n-propanol. In some embodiments, the water is present in an amount of about 5% by weight. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C.

In some embodiments, the polymorph form of the compound of Formula (1) is a polymorph known as Form 10. Polymorph Form 10 is associated with DMSO. For example, DMSO is on the surface of the polymorph. In one embodiment, polymorph Form 10 has an XRPD pattern, obtained with CuK α 1-radiation, with at least peaks at $^{\circ}2\theta$ values of 20.7 ± 0.2 , 21.7 ± 0.2 , and 24.2 ± 0.2 . In some embodiments, Form 10 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 18.2 ± 0.2 , 19.0 ± 0.2 , 20.7 ± 0.2 , 21.7 ± 0.2 , and 24.2 ± 0.2 . In some embodiments, Form 10 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 17.8 ± 0.2 , 18.2 ± 0.2 , 19.0 ± 0.2 , 20.7 ± 0.2 , 21.7 ± 0.2 , 23.4 ± 0.2 , 24.2 ± 0.2 , and 27.9 ± 0.2 . For example, in some embodiments, Form 10 has an XRPD pattern with at least peaks at $^{\circ}2\theta$ values of 6.7 ± 0.2 , 17.8 ± 0.2 , 18.2 ± 0.2 , 19.0 ± 0.2 , 19.9 ± 0.2 , 20.7 ± 0.2 , 21.7 ± 0.2 , 23.4 ± 0.2 , 24.2 ± 0.2 , and 27.9 ± 0.2 .

In some embodiments, a composition comprising polymorph Form 10 is substantially pure. For example, the composition can have a purity of at least about 90%. In some embodiments, the composition has a purity of at least about 95%. In some embodiments, the composition has a purity of at least about 98%. For example, the composition can have a purity of at least 98.5%, 98.6%, 98.7%, 98.8%, 98.9%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or 99.9%. In some embodiments, the composition is substantially free of other forms of the compound of Formula (1). For example, in some embodiments, the composition is substantially free of other anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of other forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of one or more other

forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of one or more other forms of the compound of Formula (1). For example, the composition can contain less than about 15% of Form 1, Form 2, Form 3, Form 4, Form 5, Form 6, Form 7, Form 8, Form 9, Form 11, a non-stoichiometric hydrate of Form 1, or a combination of two or more thereof.

In some embodiments, polymorph Form 10 exhibits an endotherm between about 212-237°C as measured by DSC. In some embodiments, polymorph Form 10 exhibits an endotherm at between about 234-245°C, *e.g.*, around 237°C. In some embodiments, polymorph Form 10 exhibits an exotherm between about 300-325°C, *e.g.*, around 308°C. For example, in some
10 embodiments, the endotherms and exotherms are observed when using a scan rate of 10°C per minute.

In some embodiments, polymorph Form 10 has a melting point of between about 364-372°C, such as, *e.g.*, around 369°C. In some embodiments, polymorph Form 10 undergoes a weight loss of about 0.6% before around 100°C, a weight loss of about 3.8% between about
15 100-170°C, and a weight loss of about 7.1% between about 170-260°C as measured by TGA.

Provided herein is a process of preparing polymorph Form 10. In some embodiments, the process comprises reslurrying a composition comprising the compound of Formula (1), including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate Form 10 as a residual solid. In some embodiments, the composition comprises a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water. In some
20 embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C. In some embodiments, the process further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

In some embodiments, the process comprises reslurrying a composition comprising a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water in a solvent or mixture of solvents to generate polymorph Form 10 as a residual solid. In some
25 embodiments, the solvent is DMSO. In some embodiments, the solvent is in a mixture with water, for example the solvent can be a mixture of water and DMSO. In some embodiments, the water is present in an amount of about 5% by weight. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C.

In some embodiments, the polymorph form of the compound of Formula (1) is a polymorph known as Form 11. Form 11 is an anhydrous polymorph of the compound of Formula (1). In one embodiment, polymorph Form 11 has an XRPD pattern, obtained with

CuK α 1-radiation, with at least peaks at 2θ values of 6.4 ± 0.2 , 18.5 ± 0.2 , and 22.4 ± 0.2 . In some embodiments, Form 11 has an XRPD pattern with at least peaks at 2θ values of 6.4 ± 0.2 , 17.8 ± 0.2 , 18.5 ± 0.2 , 19.9 ± 0.2 , and 22.4 ± 0.2 . In some embodiments, Form 11 has an XRPD pattern with at least peaks at 2θ values of 6.4 ± 0.2 , 8.4 ± 0.2 , 17.8 ± 0.2 , 18.5 ± 0.2 , 19.9 ± 0.2 , 22.4 ± 0.2 , 24.5 ± 0.2 , and 26.8 ± 0.2 . For example, in some embodiments, Form 11 has an XRPD pattern with at least peaks at 2θ values of 6.4 ± 0.2 , 8.4 ± 0.2 , 17.8 ± 0.2 , 18.5 ± 0.2 , 19.9 ± 0.2 , 20.3 ± 0.2 , 22.4 ± 0.2 , 22.9 ± 0.2 , 24.5 ± 0.2 , and 26.8 ± 0.2 .

In some embodiments, a composition comprising polymorph Form 11 is substantially pure. For example, the composition can have a purity of at least about 90%. In some embodiments, the composition has a purity of at least about 95%. In some embodiments, the composition has a purity of at least about 98%. For example, the composition can have a purity of at least 98.5%, 98.6%, 98.7%, 98.8%, 98.9%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or 99.9%. In some embodiments, the composition is substantially free of other forms of the compound of Formula (1). For example, in some embodiments, the composition is substantially free of other anhydrous forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of other forms of the compound of Formula (1). In some embodiments, the composition contains less than 15% by weight of one or more other forms of the compound of Formula (1), such as less than 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less by weight of one or more other forms of the compound of Formula (1). For example, the composition can contain less than about 15% of Form 1, Form 2, Form 3, Form 4, Form 5, Form 6, Form 7, Form 8, Form 9, Form 10, a non-stoichiometric hydrate of Form 1, or a combination of two or more thereof.

In some embodiments, polymorph Form 11 exhibits an endotherm between about 215-230°C as measured by DSC. In some embodiments, polymorph Form 11 exhibits an exotherm at between about 230-240°C, *e.g.*, around 235°C. In some embodiments, polymorph Form 11 exhibits an exotherm between about 300-315°C, *e.g.*, around 310°C. For example, in some embodiments, the endotherms and exotherms are observed when using a scan rate of 10°C per minute.

In some embodiments, polymorph Form 11 has a melting point of around 368°C. In some embodiments, polymorph Form 11 undergoes a weight loss of about 0.8% before around 100°C and a weight loss of about 7.0% between about 100-249°C, as measured by TGA.

Provided herein is a process of preparing polymorph Form 11. In some embodiments, the process comprises reslurrying a composition comprising the compound of Formula (1), including amorphous and polymorph forms thereof, in a solvent or mixture of solvents to generate Form 11 as a residual solid. In some embodiments, the composition comprises a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water. In some
5 embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C. In some embodiments, the process further comprises drying the residual solid, for example, under vacuum. In some embodiments, the drying is at a temperature of between about 60°C and 90°C, such as, *e.g.*, around 75°C.

10 In some embodiments, the process comprises reslurrying a composition comprising a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water in a solvent or mixture of solvents to generate polymorph Form 11 as a residual solid. In some embodiments, the solvent is dimethylformamide (DMF). In some embodiments, the solvent is in a mixture with water, for example the solvent can be a mixture of water and DMF. In some
15 embodiments, the water is present in an amount of about 5% by weight. In some embodiments, the reslurrying takes place at RT. In some embodiments, the reslurrying takes place at around 50°C.

4. Processes for preparing compositions including the compound of Formula

(1)

20 Also provided herein is a process for preparing a pharmaceutical composition, including mixing (i) a compound of Formula (1), or salt or amorphous or polymorph form thereof, prepared according to any of the processes described herein, and (ii) a pharmaceutically acceptable carrier (excipient), to form the composition.

25 In making the compositions provided herein, the compound of Formula (1), or salt or amorphous or polymorph form thereof, can be mixed with an excipient, diluted by an excipient, or enclosed within such a carrier in the form of, for example, a capsule, sachet, paper, or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier, or medium for the active ingredient. Thus, the
30 compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders. In some

embodiments, the composition is formulated for oral administration. In some embodiments, the composition is formulated as a tablet or capsule.

As used herein, “pharmaceutically acceptable carrier” includes any material which, when combined with an active ingredient, allows the ingredient to retain biological activity and is non-reactive with the subject’s immune system. Examples include, but are not limited to, any of the standard pharmaceutical carriers, such as a phosphate buffered saline solution, water, emulsions such as oil/water emulsion, and various types of wetting agents. Compositions comprising such carriers can be formulated by well-known conventional methods (see, for example, *Remington’s Pharmaceutical Sciences*, 18th edition, A. Gennaro, ed., Mack Publishing Co., Easton, Pa., 1990; and Remington, *The Science and Practice of Pharmacy*, 20th Ed., Mack Publishing, 2000).

The compositions comprising the compound of Formula (1), or a pharmaceutically acceptable salt or amorphous or polymorph form thereof, can be formulated in a unit dosage form. In some embodiments, each dosage contains from about 1 mg to about 1,000 mg (1 g). The term “unit dosage form” refers to physically discrete units suitable as unitary dosages for human subjects and other patients, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient. In some embodiments, the active material is the compound of Formula (1), or a salt or amorphous or polymorph form thereof.

In some embodiments, the compositions provided herein contain from about 5 mg to about 50 mg of the compound of Formula (1), or a pharmaceutically acceptable salt or amorphous or polymorph form thereof. For example, the compositions provided herein can contain about 5 mg to about 10 mg, about 10 mg to about 15 mg, about 15 mg to about 20 mg, about 20 mg to about 25 mg, about 25 mg to about 30 mg, about 30 mg to about 35 mg, about 35 mg to about 40 mg, about 40 mg to about 45 mg, or about 45 mg to about 50 mg of the compound of Formula (1), or a pharmaceutically acceptable salt or amorphous or polymorph form thereof.

In some embodiments, the compositions provided herein contain from about 50 mg to about 500 mg of the compound of Formula (1), or a pharmaceutically acceptable salt or amorphous or polymorph form thereof. For example, the compositions provided herein can contain about 50 mg to about 100 mg, about 100 mg to about 150 mg, about 150 mg to about 200 mg, about 200 mg to about 250 mg, about 250 mg to about 300 mg, about 350 mg to about 400 mg, or about 450 mg to about 500 mg of the compound of Formula (1), or a pharmaceutically acceptable salt or amorphous or polymorph form thereof.


In some embodiments, the compositions provided herein contain from about 500 mg to about 1,000 mg of Formula (1), or a pharmaceutically acceptable salt or amorphous or polymorph form thereof. For example, the compositions provided herein can contain about 500 mg to about 550 mg, about 550 mg to about 600 mg, about 600 mg to about 650 mg, about 650 mg to about 700 mg, about 700 mg to about 750 mg, about 750 mg to about 800 mg, about 800 mg to about 850 mg, about 850 mg to about 900 mg, about 900 mg to about 950 mg, or about 950 mg to about 1,000 mg of Formula (1), or a pharmaceutically acceptable salt or amorphous or polymorph form thereof.


EXAMPLES

EXAMPLE 1: Synthesis of *N*-(5-(3-(7-(3-fluorophenyl)-3H-imidazo[4,5-c]pyridin-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-3-methylbutanamide (1)

The compound of Formula (1) was synthesized as detailed in Scheme 7 and described below.

Scheme 7






(9)

Synthesis of 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-trityl-1H-indazole-3-carbaldehyde (9).

- 5 To a multi-neck, 10 L, cylindrical jacketed reactor was added (8) (300.0 g, 1.0 equiv., 87.5% potency assumed), bis(pinacolato)diboron (171.2 g, 1.2 equiv.), and KOAc (110.2 g, 2.0 equiv.). DMF (1575 mL, 6 volumes) was added to the flask and the flask was evacuated and refilled with N₂ three times. Next, Pd(dppf)Cl₂ (12.3 g, 0.03 equiv.) was added to the flask and the vessel was evacuated and refilled with N₂ three times. The flask was heated to 90 –
- 10 95°C using a temperature control unit (TCU) overnight (e.g., over 20 hours). The mixture was cooled to ambient temperature and H₂O (2100 mL) was added drop-wise to the vessel over 104 minutes while maintaining an internal temperature between 15 – 25°C. The precipitated solids were stirred for an additional 30 min and then the reaction mixture was filtered through filter paper on a Buchner funnel. Subsequently, the cake was rinsed with water (1575 mL, 6 vol).
- 15 The cake was conditioned until it ceased dripping. The wet cake weighed 840 g. The cake was kept under hi-vac at ambient temperature overnight. The weight was reduced to 685 g.



(11)


- 20 *Synthesis of N-(5-(3-formyl-1-trityl-1H-indazol-5-yl)pyridin-3-yl)-3-methylbutanamide (11).*

The crude boronate ester (9) was charged to a 5-L, 3-neck, round bottomed flask and (10) (*N*-(5-bromopyridin-3-yl)-3-methylbutanamide) (159.9 g, 1.0 equiv., 90.3% potency

assumed), K_3PO_4 monohydrate (388.2 g, 3.00 equiv.), 1,4-dioxane (1269 mL, 5 v), and H_2O (390 mL, 1.5 v) were charged to the vessel. Agitation was started and the internal temperature increased from 17°C to 26.7°C. The flask underwent a vacuum/ N_2 cycle 3 times. Next, $Pd(PPh_3)_4$ (19.7 g, 0.03 equiv.) was added to the flask. A vacuum/ N_2 cycle was completed 3
5 times. The flask was heated by a heating mantle to 85 – 95°C in 70 min. After 180 minutes, the mixture was cooled to ambient temperature and no remaining 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-trityl-1H-indazole-3-carbaldehyde (9) was detected by HPLC.

The reaction mixture was cooled to room temperature and added drop-wise, via addition funnel, to a multi-neck, jacketed cylindrical reactor containing H_2O (4.8 L) over 126 minutes.
10 The residual was rinsed with dioxane (200 mL) into the 10-L reactor. It was then stirred at 15 – 25°C for 90 minutes and then filtered through filter paper over a Buchner funnel. The filtration took about 20 minutes, and the cake was rinsed with water (1.6 L). This took 25 minutes, and the cake was conditioned for 45 minutes. The cake was transferred to a drying dish and kept under hi-vac at room temperature overnight. The solids were charged back into
15 the 10-L reactor after losing approximately 150 g weight. 2-methyl THF (1.69 L) was added to the flask. A dark brown solution was obtained after stirring for approximately 30 minutes at 40 – 50°C. A 9% oxalic acid solution (1348 g) was added drop-wise to the reactor in 39 minutes while keeping the temperature at 40 – 50 °C. A slurry was formed, and it was aged for 4 h at 40 – 50°C and then stirred at 0 – 10°C for 16 h. The batch was filtered (17 minutes), and the
20 reactor and the brown solids were rinsed with 615 mL of cold 2-methyl THF, which removed a significant amount of color from the solids. The pH of the filtrate was 1. The solids were charged back to the 10-L reactor, followed by the addition of 2-methyl-THF (1.69 L). A solution of 10% K_2CO_3 (1 L, 1067 g) was added to the slurry to adjust the pH to 9 at 10 – 20°C in 24 minutes. The aqueous layer (lower, colorless) was removed and the organic layer was
25 cooled to 5 – 10°C. Aqueous oxalic acid (1360 g) was added to the solution in 60 minutes. Solids precipitated from solution and, after aging overnight at 5 – 10°C, the solids (light brown) were collected by filtration (94 minutes) and washed with cold 2-methyl-THF (615 mL, 68 minutes), which removed a significant amount of color from the solids. The solids were charged back to the 10-L reactor and suspended in 2-methyl-THF (1.84 L). The pH of the
30 mixture was adjusted to 9 with 10% K_2CO_3 (1.5 L) to form a biphasic homogeneous solution. The aqueous layer was removed and the organic layer was washed with 25% NaCl (400 mL) and H_2O (1575 mL) twice. The pH of the aqueous phase was approximately 7 as determined by pH paper. The organic layer was transferred to heptanes (3860 mL) dropwise while stirring at a rate to keep the temperature below 25°C. The addition took 3.5 hours and a nice flowing

slurry was obtained. The slurry was concentrated to approximately 5 vol on a rotovap under vacuum and then filtered through a Buchner funnel (2 minutes). The cake was washed with heptanes (2.32 L) and dried under hi-vac at 40 – 50°C overnight to afford 225 g of N-(5-(3-formyl-1-trityl-1H-indazol-5-yl)pyridin-3-yl)-3-methylbutanamide (compound (11)) as an off-white powder in 71% yield. The crude compound (11), the first isolated compound (11) oxalate salt, the second isolated compound (11) oxalate salt, and the final product of compound (11) were analyzed. The purities were: 90.89%, 97.54%, 97.94%, and 98.27%, respectively.



10


(12)

Synthesis of N-(5-(3-(7-(3-fluorophenyl)-3H-imidazo[4,5-c]pyridin-2-yl)-1-trityl-1H-indazol-5-yl)pyridin-3-yl)-3-methylbutanamide (12).

To a 2 L, multi-neck, round bottomed flask was charged compound (11) (70 g, 1.0 equiv., 98% purity), compound (6) (5-(3-fluorophenyl)pyridine-3,4-diamine) (34.37 g, 1.00 equiv., 97.6% potency), grinded Na₂SO₃ (30.6 g, 2.0 equiv.) and NMP (1065 mL). A 0.1 mL portion of the reaction was taken for Karl Fischer (KF) titration analysis, which showed 3129 ppm of H₂O content. The reaction mixture was heated to 110 – 115°C by a heating mantle. The reaction went to completion after 24 hours with 14.1% of compound (1), 78.3% of compound (12), and 0.42% of compound (11). The reaction mixture was quenched into a mixture of water (2135 mL), NaHCO₃ (103 g), and 1030 mL EtOAc in a 10-L reactor over 27 minutes. The flask was rinsed with NMP (67 mL) into the quench mixture. The mixture was filtered through a Celite pad and the filtration took 16 minutes. The reactor and the cake were rinsed with EtOAc (200 mL). The filtrate was transferred back to the 10-L reactor and the phases were separated. The aqueous layer was extracted with EtOAc (760 mL). The combined organic layer was washed with a mixture of water (525 mL) and 25% NaCl solution (125 mL). A 9% oxalic acid solution (308 g) was added dropwise to the organic layer in 45 minutes to obtain a pH of 1.

25

The slurry was filtered through filter paper over 21 minutes. The reactor and cake were rinsed with EtOAc (380 mL) and this took another 10 minutes. An aliquot of the wet cake was analyzed for purity and the result showed a 87.89% by area under curve (AUC) purity for compound (12). The wet cake was charged back to a 5-L 4-neck RBF followed by the addition of MeOH (693 mL) and DCM (1654 mL). The mixture was heated up to reflux (35°C) and Celite (47 g) was added. A Buchner funnel was set up with a Celite pad on the bottom and a pad of activated carbon (60 g, Darco™ G-60) on the top. The mixture was stirred at reflux for 70 minutes and then filtered hot through the Buchner funnel. The filtration took 13 minutes. The reactor and the cake were rinsed with a mixed solvent of MeOH/DCM (173 mL / 414 mL). It took 10 minutes to complete. The dark brown solution was concentrated on a rotovap to approximately 4 vol and acetone (870 mL) was added. The mixture was concentrated to approximately 4 vol and once again acetone (2 × 870 mL) was added. The mixture was concentrated to approximately 3 vol and diluted with acetone (260 mL). ¹H NMR analysis of the solvent showed 2.8% by weight of MeOH to acetone and the solvent swap was deemed complete. The slurry was filtered and the cake was rinsed with acetone (430 mL). An aliquot of the wet cake was analyzed and it had a purity of 99.10% AUC. The wet cake was charged back to a 5-L multi-neck RBF followed by the addition of 2-methyl-THF (878 mL). A 10% K₂CO₃ solution (440 mL) was added to the mixture over 20 minutes to obtain a pH of approximately 10. The mixture was transferred to a 2-L separatory funnel and the phases were separated. The organic layer was washed with 4.8% a Na₂SO₄ solution (2 × 526 mL) and then diluted with tert-butyl methyl ether (MTBE; 927 mL). The organic solution was added to heptanes over 1 hour and 51 minutes; keeping the temperature at 15 – 25°C. The resulting slurry was concentrated to approximately 3 vol on a rotovap to afford a very thick slurry. It was filtered through filter paper (total 15 minutes) and the cake was dried under hi-vac at 40 – 50°C for 15 h. Compound (12) was obtained (61.1 g) in 66% yield. HPLC analysis showed 99.16% purity.



(1)

Synthesis of N-(5-(3-(7-(3-fluorophenyl)-3H-imidazo[4,5-c]pyridin-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-3-methylbutanamide (1).


To a 1-L 3-neck round bottomed flask was charged TFA (250 mL). Compound (12) (55 g) was added portion-wise over 12 minutes to the flask while using an ice/water bath to control the temperature below 20°C. After 2 h, a sample was analyzed by HPLC and showed that no compound (12) was detectable. TFA (37 mL) was added to the brown solution to make up the rinse volume. H₂O (275 mL) was added dropwise over 1 hour and 37 minutes to the TFA mixture at a rate to keep the temperature below 10°C. Solids precipitated during the addition. A free-flowing slurry was obtained at the beginning and turned into an extremely thick slurry at the end of the addition. The slurry was stirred at 0 – 10°C for 84 minutes. The batch was filtered. The cake was rinsed with a 1.2:1 TFA/H₂O solution (110 mL). The wet cake was conditioned at room temperature overnight. The wet cake was slurried in 90% EtOH/water (672 mL) for 3 hours and 40 minutes. The material was isolated by filtration through a tight weave filter cloth. The cake was rinsed with 90% EtOH/water (134 mL) and it took another 1 hour and 12 minutes. The wet cake was charged back to a 1-L RBF, followed by the addition of water (385 mL). A 5% Na₂CO₃ solution (19.25 g Na₂CO₃ was dissolved in 366 mL of water) was added to the slurry over 29 minutes. The slurry was stirred at approximately 20°C for 5 h and then filtered through a tight weave filter cloth. The cake was rinsed with water (110 mL, it took 64 minutes) and then kept at room temperature overnight. The wet cake was reslurried in water (550 mL) at approximately 25°C twice (5 hours and 2 hours, respectively). The filtrations took 47 min and 54 min, respectively. The wet cake was reslurried in 90% IPA/water (669 mL) at approximately 25°C for 67 min and then filtered through a tight weave filter cloth. The filtration took 2 hours and 9 minutes. The cake was rinsed with 90% IPA/water (200 mL) and conditioned at RT overnight. The solid was reslurried in 90% MeOH/water (400 mL) at 30 – 35°C for 46 hours and then filtered through a tight weave filter cloth. The cake was rinsed with 90% MeOH/water (134 mL). After drying, 18.48 g of compound (1) was obtained in 50% yield. HPLC analysis showed 99.33% purity.

EXAMPLE 2: Synthesis of N-(5-(3-(7-(3-fluorophenyl)-3H-imidazo[4,5-c]pyridin-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-3-methylbutanamide (1).


The synthesis of compound (1) (Scheme 8) started from the borylation of compound (8) with bis(pinacolato)diboron to form compound (9), followed by the *in situ* Suzuki-Miyaura cross-coupling with compound (10) to form compound (11). The cyclization of compound (11)

with compound (6) gave compound (12). The synthesis was finalized with the deprotection of the trityl group of compound (12) to afford compound (1).

Scheme 8.



- 5 *Borylation and Suzuki-Miyaura Cross-Coupling Reaction.* Compound (11) was prepared and purified by first borylating compound (8) and subsequently performing a Suzuki-Miyaura cross-coupling reaction as described in the process below.



Process	
1.	Charge 50.0 g (1.0 X, 1.00 eq.) active (8) into R1. (R1= reactor).
2.	Charge 32.6 g (0.652 X, 1.20 eq.) (0.625 ~ 0.679 X) bis(pinacolato)diboron into R1.
3.	Charge 21.0 g (0.42 X, 2.00 eq.) (0.40 ~ 0.44 X) KOAc into R1.
4.	Charge 275.0 g (5.5 X, 5.8 vol.) (5.0 ~ 6.0 X) DMF into R1.


5. Turn on R1 stir.
6. Degas R1 with N ₂ for 3 times.
7. Charge 2.35 g (0.047 X, 0.03 eq.) (0.039 ~ 0.055 X) Pd(dppf)Cl ₂ into R1.
8. Degas R1 with N ₂ for 3 times.
9. Heat R1 to 92°C (87 ~ 97°C).
10. Stir R1 for 18 h (16 ~ 20 h) at 92°C (87 ~ 97°C).
11. Cool the material in R1 to 20°C (15 ~ 25°C).
12. Add 375.0 g (7.5 X, 7.5 vol.) (7.0 ~ 8.0 X) process water at 20°C (15 ~ 25°C) into R1 dropwise via headtank NLT1.5 h.
13. Stir the material in R1 for 45 min (30 ~ 60 min) at 20°C (15 ~ 25°C).
14. Filter the material. Transfer the mother liquid into T1.
15. Rinse R1 with 300.0 g (6.0 X, 6.0 vol.) (5.5 ~ 6.5 X) process water and then transfer to funnel to slurry the cake. Transfer the mother liquid into T1.
16. Test the purity by HPLC: report.
17. Charge (9) into R1.
18. Charge 27.5 g (0.550 X, 1.00 eq.) (0.539 ~ 0.561X) active (10) into R1.
19. Charge 85.5 g (1.71 X, 3.0 eq.) (1.60 ~ 1.82 X) K ₃ PO ₄ ·3H ₂ O into R1.
20. Charge 275.0 g (5.5 X, 5.3 vol.) (5.0 ~ 6.0 X) 1,4-dioxane into R1.
21. Charge 40.0 g (0.8 X, 0.8 vol.) (0.7 ~ 0.9 X) H ₂ O into R1.
22. Turn on R1 stir.
23. Degas R1 with N ₂ three times.
24. Charge 3.71 g (0.0741 X, 0.03 eq.) (0.0692 ~ 0.0790 X) Pd(PPh ₃) ₄ into R1 under N ₂ protect.
25. Degas R1 with N ₂ three times.
26. Heat R1 to 90°C (85 ~ 95°C).
27. Stir R1 at 90°C (85 ~ 95°C) for 2.5 h (2.0 ~ 3.0 h).
28. Cool R1 to 20°C (15 ~ 25°C).

29. Transfer the material in R1 into drum.
30. Charge 775.0 g (15.5 X, 15.5 vol.) (15.0 ~ 16.0 X) process water into R1.
31. Add the material in step 29 drum into R1 slowly by dropwise while keeping R1 at 20°C (15 ~ 25°C) about 3.0 h (2.0 ~ 4.0 h).
32. Stir R1 for 45 min (30 ~ 60 min) at 20°C (15 ~ 25°C).
33. Filter the material in R1. Transfer the mother liquid into T2 (T2= tank).
34. Rinse R1 with 300.0 g (6.0 X, 6.0 vol.) (5.5 ~ 6.5 X) process water and then transfer to funnel to slurry the solid. Transfer the mother liquid into T2. Test the wet cake of purity by HPLC: report. Test the residual F1 in T2: report.
35. Charge the solid in funnel into R1.
36. Charge 275.0 g (5.5 X, 6.4 vol.) (5.0 ~ 6.0 X) 2-Me-THF into R1.
37. Stir R1 for 45 min (30 ~ 60 min) at 20°C (15 ~ 25°C).
38. Heat R1 to 45°C (40 ~ 50°C).
39. Add 275.0 g (5.5 X, 2.85 eq.) (5.0 ~ 6.0 X) 10% aqueous oxalic acid slowly by dropwise into R1 at 45°C (40 ~ 50°C).
40. Stir R1 at 45°C (40 ~ 50°C) for 5 h (4 ~ 6 h).
41. Cool R1 to 7°C (5 ~ 10°C).
42. Stir R1 at 7°C (5 ~ 10°C) for 18 h (16 ~ 20 h).
43. Filter the material in R1, and transfer the mother liquid into T2.
44. Rinse R1 with 125.0 g (2.5 X, 2.9 vol.) (2.0 ~ 3.0 X) 2-Me-THF and then transfer to funnel to slurry the solid. Transfer the mother liquid into T3.
45. Charge the cake into R1.
46. Charge 275.0 g (5.5 X, 6.4 vol.) (5.0 ~ 6.0 X) 2-Me-THF into R1.
47. Stir R1 for 45 min (30 ~ 60 min) at 20°C (15 ~ 25°C).
48. Charge 300.0 g (6.0 X, 2.0 eq.) (5.0 ~ 7.0 X) 10% K ₂ CO ₃ aqueous into R1.
49. Stir R1 for 45 min (30 ~ 60 min) at 20°C (15 ~ 25°C).
50. Transfer the bottom layer into T4.
51. Cool R1 to 7°C (5 ~ 10°C).

52. Add 275.0 g (5.5 X, 2.85 eq.) (5.0 ~ 6.0 X) 10% aqueous oxalic acid slowly by dropwise into R1 at 7°C (5 ~ 10°C).
53. Stir R1 at 7°C (5 ~ 10°C) for 5 h (4 ~ 6 h).
54. Filter the material in R1, and transfer the mother liquid into T2.
55. Rinse R1 with 125.0 g (2.5 X, 2.9 vol.) (2.0 ~ 3.0 X) 2-Me-THF and then transfer to funnel to slurry the solid. Transfer the mother liquid into T3.
56. Charge the cake into R1.
57. Charge 350.0 g (7.0 X, 8.1 vol.) (6.5 ~ 7.5 X) 2-Me-THF into R1.
58. Stir for 45 min (30 ~ 60 min) at 20°C (15 ~ 25°C).
59. Charge 300.0 g (6.0 X, 2.0 eq.) (5.0 ~ 7.0 X) 10% K ₂ CO ₃ aqueous into R1.
60. Stir R1 for 45 min (30 ~ 60 min) at 20°C (15 ~ 25°C).
61. Transfer the bottom layer into T4.
62. Charge 350.0 g (7.0 X) (6.0 ~ 8.0 X) 5% NaCl aqueous into R1.
63. Stir R1 for 25 min (15 ~ 35 min) at 20°C (15 ~ 25°C).
64. Transfer the bottom layer into T3.
65. Charge 350.0 g (7.0 X) (6.0 ~ 8.0 X) 5% NaCl aqueous into R1.
66. Stir R1 for 25 min (15 ~ 35 min) at 20°C (15 ~ 25°C).
67. Transfer the bottom layer into T4.
68. Charge 350.0 g (7.0 X) (6.0 ~ 8.0 X) 5% NaCl aqueous into R1.
69. Stir R1 for 25 min (15 ~ 35 min) at 20°C (15 ~ 25°C).
70. Transfer the bottom layer into T4.
71. Transfer the material in R1 into drum.
72. Charge 475.0 g (9.5 X, 14.0 vol.) (9.0 ~ 10.0 X) n-heptane into R1.
73. Add the material in drum of step 71 into R1 slowly at 20°C (15 ~ 25°C).
74. Stir R1 for 45 min (30 ~ 60 min) at 20°C (15 ~ 25°C).
75. Concentrated R1 to 275.0 mL (5.5 vol.) (5.0 ~ 6.0 vol.) Under vacuum below 40°C.

76. Adjust R1 to 20°C (15 ~ 25°C).
77. Filter the material in R1. The filtrate is transferred to T5.
78. Rinse R1 with 300.0 g (6.0 X, 8.8 vol.) (5.5 ~ 6.5 X) n-heptane and then transfer to filter. Transfer the mother liquid into T5.
79. Dry the product (11) at 45°C (40 ~ 50°C) for 18 h (16 ~ 20 h).

Imidazopyridine formation. The compound of Formula (11) can be converted to the compound of Formula (12) and purified as described in the process below.




Process
1) Charge 50.0 g (1.0 X, 1.00 eq.) active (11) into R1. (R1 = reactor).
2) Charge 24.5 g (0.49 X, 1.00 eq.) (0.48 ~ 0.50 X) active (6) bis hydrochloride salt into R1.
3) Charge 12.5 g (0.25 X, 1.10 eq.) (0.24 ~ 0.26 X) Na ₂ SO ₃ into R1.
4) Charge 775.0 g (15.5 X, 15.5 vol.) (15.0 ~ 16.0 X) NMP in R1.
5) Stir R1 for 20 min (10 ~ 30 min).
6) Degas with N ₂ twice.
7) Heat R1 to 112°C (110 ~ 115°C).
8) Stir R1 at 112°C (110 ~ 115°C) for 8 h (7 ~ 9 h).
9) Cool R1 to 25°C (20 ~ 30°C).
10) Transfer the material in R1 into drum.
11) Charge 1600.0 g (32 X) (31 ~ 33 X) 5% NaHCO ₃ aqueous into R1 at 25°C

(20 ~ 30°C).
12) Charge 675.0 g (13.5 X, 15.0 vol.) (13.0 ~ 14.0 X) EA (ethyl acetate) into R1.
13) Add the material in step 11 drum into R1 slowly while keeping R1 at 25°C (20 ~ 30°C).
14) Stir R1 for 45 min (30 ~ 60 min) at 25°C (20 ~ 30°C).
15) Stay R1 for 25 min (15 ~ 35 min) at 25°C (20 ~ 30°C).
16) Split the material in R1. Transfer the bottom layer into T1.
17) Filter the material in R1 through diatomite pad, transfer the filtrate into T2.
18) Rinse diatomite pad with 100.0 g (2.0 X, 2.2 vol.) (1.5 ~ 2.5 X) EA, transfer the filtrate into T2.
19) Transfer the filtrate in T1 into R1.
20) Charge 475.0 g (9.5 X, 10.6 vol.) (9.0 ~ 10.0 X) EA into R1 at 25°C (20 ~ 30°C).
21) Stir R1 for 25 min (15 ~ 35 min) at 25°C (20 ~ 30°C).
22) Stay R1 for 25 min (15 ~ 35 min) at 25°C (20 ~ 30°C).
23) Split the material in R1, transfer the aqueous layer into T1.
24) Transfer the material in T2 into R1.
25) Charge 475.0 g (9.5 X) (9.0 ~ 10.0 X) 5%NaCl solution into R1 at 25°C (20 ~ 30°C).
26) Stir R1 for 25 min (15 ~ 35 min) at 25°C (20 ~ 30°C).
27) Stay R1 for 25 min (15 ~ 35 min) at 25°C (20 ~ 30°C).
28) Split the material in R1. Transfer the bottom layer into T1.
29) Add 225.0 g (4.5 X, 2.5 eq.) (4.2 ~ 4.7 X) 9% oxalic acid solution into R1 slowly at 25°C (20 ~ 30°C).
30) Stir R1 for 1.0 h (0.5 ~ 1.5 h) at 25°C (20 ~ 30°C).
31) Filter the material in R1. Transfer the mother liquid into T3.

32) Rinse R1 with 175.0 g (3.5 X, 3.9 vol.) (3.0 ~ 4.0 X) EA and then transfer to funnel to slurry the solid. Transfer the mother liquid into T3. Residual K1 in EA solution.
33) Charge the filter cake into R1.
34) Charge 4000.0 g (80.0 X) (78.0 ~ 82.0 X) MeOH/DCM (m:m = 1:4) into R1.
35) Heat R1 to 38°C (35 ~ 40°C).
36) Stir R1 for 45 min (30 ~ 60 min) to clear at 38°C (35 ~ 40°C).
37) Filter the material in R1 through diatomite pad and active carbon (0.15 X) (0.13 ~ 0.17 X). The filtrate is transferred to R1.
38) Rinse filter cake with 500.0 g (10.0 X) (9.0 ~ 11.0 X) MeOH/DCM (m:m=1:4) and then transfer to filter. Transfer the mother liquid into R1.
39) Concentrate to 275.0 mL (5.5 vol.) (5.0 ~ 6.0 vol.) Under vacuum below 45°C.
40) Charge 475.0 g (9.5 X, 12.0 vol.) (9.0 ~ 10.0 X) acetone into R1.
41) Concentrate to 225.0 mL (4.5 vol.) (4.0 ~ 5.0 vol.) Under vacuum below 45°C.
42) Charge 475.0 g (9.5 X, 12.0 vol.) (9.0 ~ 10.0 X) acetone into R1.
43) Concentrate to 275.0 mL (5.5 vol.) (5.0 ~ 6.0 vol.) Under vacuum below 45°C.
44) Charge 125.0 g (2.5 X, 3.2 vol.) (2.0 ~ 3.0 X) acetone into R1. Residual MeOH report.
45) Cool R1 to 25°C (20 ~ 30°C).
46) Filter the material in R1. Transfer the mother liquid into T3.
47) Rinse R1 with 125.0 g (2.5 X, 3.2 vol.) (2.0 ~ 3.0 X) acetone and then transfer to centrifuge to slurry the solid. Transfer the mother liquid into T3. Report: residual K1 in acetone solution; Check purity of K1 wet $\geq$ 98.5%, if pass proceed to following steps, otherwise repeat step 35-47, Residual Pd $\leq$ 20 ppm.

48) Charge the cake of STEP 47 into R1.
49) Charge 525.0 g (10.5 X, 12.2 vol.) (10.0 ~ 11.0 X) 2-Me-THF into R1.
50) Stir for 25 min (15 ~ 35 min) at 25°C (20 ~ 30°C).
51) Charge 310.0 g (6.2 X, 2.5 eq.) (6.0 ~ 6.4 X) 10% K ₂ CO ₃ aqueous into R1 at 25°C (20 ~ 30°C).
52) Stir R1 at 25°C (20 ~ 30°C) for 45 min (30 ~ 60 min).
53) Transfer the bottom layer into T4.
54) Charge 350.0 g (7.0 X, 1.38 eq.) (6.5 ~ 7.5 X) 5% Na ₂ SO ₄ solution into R1 at 25°C (20 ~ 30°C).
55) Stir R1 for 25 min (15 ~ 35 min) at 25°C (20 ~ 30°C).
56) Transfer the bottom layer into T4.
57) Charge 350.0 g (7.0 X, 1.38 eq.) (6.5 ~ 7.5 X) 5% Na ₂ SO ₄ solution into R1 at 25°C (20 ~ 30°C).
58) Stir R1 for 25 min (15 ~ 35 min) at 25°C (20 ~ 30°C).
59) Transfer the bottom layer into T4.
60) Charge 475.0 g (9.5 X, 12.5 vol.) (9.0 ~ 10.0 X) MTBE into R1.
61) Transfer the material in R1 into drum.
62) Charge 850.0 g (17.0 X, 25.0 vol.) (16.5 ~ 17.5 X) n-heptane into R1.
63) Add the material in drum of step 61) into R1 slowly by dropwise at 20°C (15 ~ 25°C).
64) Concentrate to 275 mL (5.5 vol.) (5.0 ~ 6.0 vol.) Under vacuum below 45°C.
65) Cool R1 to 25°C (20 ~ 30°C).
66) Filter the material in R1. Transfer the mother liquid into T5.
67) Rinse R1 with 200.0 g (4.0 X, 5.9 vol.) (3.5 ~ 4.5 X) n-heptane and then transfer to funnel to slurry the solid. Transfer the mother liquid into T5. Residual (12) in n-heptane solution.
68) Dry the product (12) at 45°C (40 ~ 50°C) for 18 h (16 ~ 20 h).

Deprotection of the trityl group. The deprotection of the trityl group to yield a compound of Formula (1) was accomplished as described in the procedure below.




5

Process
1) Charge TFA into R1.
2) Charge (12) in portions into R1 maintaining temperature at 15-25°C.
3) Stir the solution at 15-25°C for 2 h.
4) Charge purified water dropwise by maintaining the temperature of solution 0-10°C.
5) Charge crystal seed of TFA salt of (1) into the reactor.
6) Charge 2nd lot of purified water dropwise by maintaining the temperature of solution 0-10°C.
7) Stir the materials in R1 at 5-15°C for 2 h.
8) Filtered the solids and washed with aqueous TFA solution.
9) Transfer the wet cake of step 8 to R1.
10) Reslurry the solid in 90% EtOH/water at 20-30°C for 4 h.
11) Filter the solids.
12) Suspend the solids in purified water.
13) Basify the mixture to pH 9-10 by adding 5% Na ₂ CO ₃ solution.
14) Stir the mixture at 20-30°C for 5 h.

15) Filter the solids and rinse with water.
16) Transfer the solids into clean R1 again.
17) Reslurry the solids in purified water at 20-30°C for 6 h.
18) Filter the solids.
19) Transfer the solids into R1 again.
20) Reslurry the solids in purified water at 20-30°C for 20 h.
21) Filter the solids and rinse with water.
22) Transfer the solids into R1 again.
23) Reslurry the solids in 90% isopropanol/water at 20-30°C for 2 h.
24) Filter the solids and rinse with 90% isopropanol/water.
25) Transfer the solids into R1 again.
26) Charge 90% methanol/water into R1.
27) Charge crystal seed of (1) into R1.
28) Reslurry the solids in 90% methanol/water at 30-35°C for 48 h.
29) Filter the solids and rinse with 90% methanol/water.
30) Dry the product under vacuum at 40-50°C.

EXAMPLE 3: Synthesis of polymorph Form 1 of N-(5-(3-(7-(3-fluorophenyl)-3H-imidazo[4,5-c]pyridin-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-3-methylbutanamide (1)

The synthesis of polymorph Form 1 of compound (1) from compound (12) involved
 5 the deprotection of the trityl group followed by crystallization as described below.



A. Synthesis of compound (1)

A 3-L 3-neck round bottomed flask was charged with trifluoroacetic acid (TFA, 800 g). Compound (12) (100 g) was slowly added to the flask while maintaining the temperature of the reaction mixture at 20°C. The reaction was stirred until at least 99% conversion to compound (1) was observed (typically 3 to 5 h). The reaction mixture was cooled to 5°C. Water (250 g) was slowly added to the flask while maintaining the temperature of the reaction mixture at 5°C, and stirred for 45 minutes at 5°C. The resulting mixture was filtered. The reaction flask was rinsed with 8:2.5 m:m TFA:water (150 g) and the resulting solution was passed through the filter cake.

A round-bottomed flask was charged with the combined mother liquors of the previous two filtrations, and cooled to 5°C. Water (350 g) was added to the reaction mixture at 5°C, stirred for 1.5 h at 10°C, and the resulting mixture was filtered. The reaction flask was rinsed with 4:3 m/m TFA:water (200 g), the resulting mixture was slurried with the filter cake, and the slurry was filtered. Residual compound (1) was tested for in the mother liquors of the previous two filtrations, as well as the purity of compound (1) in the filter cake.

The filter cake was added to a round-bottomed flask, followed by 90% ethanol (950 g), and was stirred at 30°C for 3 h. The resulting mixture was stirred at 30°C for 2-4 hours, filtered, and the filter cake was rinsed with 90% ethanol (200 g). The mother liquors of the previous two filtrations were tested for residual compound (1), and the purity of compound (1) in the filter cake was tested.

The filter cake was added to a round-bottomed flask, followed by water (700 g), and was stirred at 25°C for 1 h. Aqueous 5% Na₂CO₃ (740 g) was then added and the solution was

stirred at 25°C for 6 h. The resulting mixture was filtered, and the filter cake was rinsed with water (200 g).

The filter cake was added to a round-bottomed flask, followed by water (950 g), and was stirred at 25°C for 6.5 h. The resulting mixture was filtered, and the filter cake was rinsed with water (200 g). The filter cake was added to a round-bottomed flask, followed by water (950 g), and was stirred at 25°C for 3.5 h. The resulting mixture was filtered, and the filter cake was rinsed with water (200 g). The purity of compound (1) was tested, as well as the quantity of residual Pd in the filter cake. The residual TFA and residue on ignition of filter cake was also tested. If the residual TFA was not 0.07% or lower, and/or residue on ignition (after drying) was not 0.4% or lower, the filter cake was added to a round bottomed flask, followed by water (950 g). The resulting mixture was stirred at 25°C for 3.5 h, filtered, and rinsed with water (200 g). The residual TFA of the filter cake was again tested. If the residual TFA concentration was 0.07% or lower and the residual on ignition (after drying) was 0.4% or lower, the filter cake was added to a round bottomed flask, followed by 90% isopropyl alcohol (950 g). The resulting mixture was stirred at 25°C for 2 h. The resulting mixture was filtered and the filter cake was rinsed with 90% isopropyl alcohol (200 g).

B. Preparation of polymorph Form 1 of compound (1)

The filter cake was added to a round-bottomed flask, followed by 90% methanol (600 g), and the resulting mixture was heated to 25°C. 4.5 g of polymorph Form 1 of compound (1) was added to the mixture, the mixture was heated to 33°C, and the solution was reslurried at 33°C for 13 h. Formation of polymorph Form 1 of compound (1) was monitored by taking an X-ray powder diffractogram (XRPD) of a small sample, then reslurrying continued at 33°C for another 13 h. Another sample was obtained. XRPD analysis was performed to test whether complete conversion of compound (1) to the polymorph Form 1 of compound (1) had occurred. Particle size distribution (PSD) was measured to test whether the crystals had a distribution of $2.0\ \mu\text{m} \leq D_{50} \leq 5.0\ \mu\text{m}$, $D_{90} \leq 15.0\ \mu\text{m}$. If complete conversion to polymorph Form 1 had not occurred or the PSD was outside of the range, reslurrying continued at 33°C. XPRD and particle size distribution analysis occurred every 13-15 hours until both tests yielded a positive result.

The mixture was then filtered, and the filter cake was rinsed with 90% methanol (200 g). Tests were performed to determine whether (1) the purity of compound (1) was at least 98% (positive result), (2) residual palladium had a concentration of, at most, 10 ppm (positive result), (3) the XRPD indicated complete conversion to polymorph Form 1 (positive result), and (4) the particle size distribution was $2.0\ \mu\text{m} \leq D_{50} \leq 5.0\ \mu\text{m}$, $D_{90} \leq 15.0\ \mu\text{m}$ (positive

result). When each of the four aforementioned tests yielded a positive result, polymorph Form 1 of compound (1) was dried at 55°C for 27 h. If any of the four aforementioned tests yielded a negative result, the filter cake was added to a round-bottomed flask, followed by 90% methanol (600 g), and was reslurried at 33°C for 4 h. The mixture was filtered, and the filter cake was rinsed with 90% methanol (200 g). The filter cake was added to a round-bottomed flask, followed by 90% methanol (600 g), and reslurrying at 33°C for 18 h. The mixture was filtered, and the filter cake was rinsed with 90% methanol (200 g) to obtain polymorph Form 1 of compound (1). Polymorph Form 1 of compound (1) was dried at 55°C for 27 h.

10 **EXAMPLE 4: Polymorph screen**

A polymorph screen was performed on the compound of Formula (1) to determine solubility, polymorphism, and thermodynamic stability.

A. Analysis of the starting solid

X-ray powder diffraction (XRD), differential scanning calorimetry (DSC), and thermal gravimetric analysis (TGA) scans of the starting solid compound of Formula (1) obtained from Example 3A, above, indicated that the starting solid was a crystalline material and was a mixture of Form 1 and a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water. According to the DSC scan (FIG. 12B), the solid showed an endotherm between 50°C-100°C; it also showed an exotherm at 284°C; and the solid eventually melted at 364°C. According to the TGA scan (FIG. 12C), a 1.4% weight loss was observed before 100°C.

The solubility of the starting solid was measured by the gravimetric method and indicated that the compound had low solubility at RT and at 50°C in all solvents tested except DMF and DMSO. Results from the solubility data test at RT and at 50°C are shown in Table 1.

25 **Table 1. Solubility data of the starting solid obtained from Example 3A**

Solvents	Solubility at RT (mg/mL)	Solubility at 50°C (mg/mL)
Acetone	1	1
Acetonitrile	~0	0
MeOH	1	1
Toluene	1	1
EtOH	2	2
IPAc	~0	~0

Solvents	Solubility at RT (mg/mL)	Solubility at 50°C (mg/mL)
EA	1	1
MtBE	~0	~0
IPA	2	5
MEK	1	1
MA	~0	~0
n-Propanol	1	2
MIBK	1	1
n-Butyl acetate	~0	~0
water	1	1
Heptane	~0	~0
n-Butanol	1	2
DMSO	n/a	n/a
DMF	12	16
DCM	2	2
Acetic acid	~0	3

Reslurry experiments in various solvents were performed. Approximately 30-80 mg of the starting solid (a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water) was slurried in 39 different solvents (pure and binary solvents; the ratio of organic solvent/water (V/V) was 95%/5%) at RT and 50°C for 5 days. Three solvates, one non-stoichiometric hydrate, and eleven non-solvated forms were identified. A “*” after a particular Form, *e.g.*, Form 2*, indicates that the forms had similar XRD scans with minor differences and were considered to belong to the same class. Generally, the identified forms showed multiple endotherms/exotherms on differential scanning calorimetry (DSC) scans; Form 9 showed a single endotherm. XRD of both wet and dry samples were scanned (FIG. 12A (dry sample)). The data is shown in Tables 2 and 3 below.

Table 2. Results of reslurry experiments at RT

Solvent	Crystalline Form (wet/dry)		Solvent	Crystalline Form (wet/dry)	
Acetone	Solvate 1	Form 2	Acetone/water	Solvate 2	Form 4**
Acetonitrile	Form 2	Form 1	Acetonitrile/water	Form 12	Form 1

MeOH	Form 13	Form 1	MeOH/water	Form 12	Form 1
Toluene	Form 1	Form 2*	Toluene/water	Form 13	Form 1
EtOH	Form 2*	Form 3	EtOH/water	Solvate 3	Form 2
IPAc	Form 3	Form 4	IPAc/water	Form 12	Form 1
EA	Form 4*	Form 5	EA/water	Form 12	Form 1
MtBE	Form 5*	Form 6	MtBE/water	Form 12	Form 1
IPA	Form 6	Form 7	IPA/water	Form 6	Form 6
MEK	Form 7	Form 4	MEK/water	Form 7	Form 7
MA	Form 4	Form 4*	MA/water	Form 13	Form 1
n-Propanol	Form 4*	Form 8	n-Propanol/water	Form 2**	Form 2**
MIBK	Form 8	Form 3	MIBK/water	Form 12	Form 1
n-Butyl acetate	Form 3*	Form 1	n-Butyl acetate/water	Form 13	Form 12
Water	Form 13	Form 1	Heptane/water	Form 13	Form 12
Heptane	Form 1	Form 9	n-Butanol/water	Form 13	Form 13
n-Butanol	Form 9	Form 10	DMSO/water	amorphous	Form 10
DMSO	amorphous	Form 11	DMF/water	Form 11	Form 11
DMF	Form 11	Form 1	DCM/water	Form 13	Form 1
DCM	Form 1	Form 2			

Table 3. Results of reslurry experiments at 50°C

Solvent	Crystalline Form (wet/dry)		Solvent	Crystalline Form (wet/dry)	
Acetone	Solvate 2	Form 4**	Acetone/water	Form 4**	Form 4**
Acetonitrile	Form 2*	Form 2	Acetonitrile/water	Form 13	Form 13
MeOH	Form 1	Form 1	MeOH/water	Form 13	Form 13
Toluene	Form 1	Form 1	Toluene/water	Form 13	Form 13
EtOH	Form 2*	Form 2*	EtOH/water	Form 9	Form 9
IPAc	Form 9	Form 9	IPAc/water	Form 13	Form 13
EA	Form 4*	Form 4	EA/water	Form 4*	Form 4*
MtBE	Form 5*	Form 4	MtBE/water	Form 13	Form 13
IPA	Form 6	Form 6	IPA/water	Form 6	Form 6
MEK	Form 7	Form 7	MEK/water	Form 7	Form 7

MA	Form 4	Form 4	MA/water	Form 12	Form 4
n-Propanol	Form 4	Form 4**	n-Propanol/water	Form 9	Form 9
MIBK	Form 8	Form 8	MIBK/water	Form 13	Form 1
n-Butyl acetate	Form 9	Form 9	n-Butyl acetate/water	Form 13	Form 1
water	Form 13	Form 13	Heptane/water	Form 13	Form 1
Heptane	Form 13	Form 13	n-Butanol/water	Form 13	Form 1
n-Butanol	Form 9	Form 9	DMSO/water	Amorphous	Form 10
DMSO	Amorphous	Form 10*	DMF/water	Form 11	Form 11
DMF	Form 11	Form 11*	DCM/water	Form 13	Form 1
DCM	Form 13	Form 13			

The slurry experiments identified 3 solvated forms from wet samples (Solvates 1, 2, and 3); 2 non-stoichiometric hydrates of Form 1 (Forms 12 and 13); and 11 non-solvated forms (Forms 1-11). In some instances, similar XRD scans with minor differences were obtained. These were considered to be part of the same class (*e.g.*, the same form). For example, XRD scans of Form 2 and Form 2* were similar and were considered to belong to the same class. The solvated forms were obtained from wet sample analysis; after drying, the sample indicated a different XRD.

Solvate 1 was obtained from acetone at RT, and after drying, a low crystallinity solid was generated. Solvate 2 was obtained from acetone (at RT) and acetone/water (at RT), and after drying, Form 4** was generated. Solvate 3 was obtained from EtOH/water at RT, and after drying, Form 2 was generated.

B. Form 1

The experiments that generated Form 1 are shown in Table 4, below. Form 1 was generally obtained from drying Form 13 or Form 12. Form 1 can be considered a dehydrated hydrate. Reslurry in many binary solvents (with 5% water) generated Form 1. Purity of the residual solid was 98.9%. KF of Form 1 (one sample) solid was 5.8%; residual MeOH of Form 1 solid was 0.01%. A TGA scan of fully dried Form 1 solid was performed (FIG. 1C). A 0.33% weight loss was observed before 100°C.

Form 1 showed sharp crystalline peaks on the XRD scan (FIG. 1A). The XRD peaks of Form 1 are shown in Table 5, below. According to the DSC scan (FIG. 1B), the solid showed an endotherm between 50-100°C; it showed an exotherm at 281°C; and the melting point was 363°C.

The Form 1 solid was dried at 75°C under vacuum overnight, and XRD, DSC, and TGA scans were performed. Comparison of the first and the second XRD scans (after drying at 75°C under vacuum overnight), showed no change. However, the DSC scans indicated the absence of endotherm. The loss of the early peak on the DSC scan had no effect on the XRD trace, showing that the endotherm between 50-100°C on DSC scan was due to the free solvent.

The Form 1 solid was heated in a DSC chamber to 305°C (past the endotherm/exotherm around 280°C), and then scanned by XRD. Comparison of the first and the third XRD and DSC scans showed that after heating to 305°C, Form 1 converted to Form 9. Thus, the endotherm/exotherm around 280°C might be due to melting/crystallization events.

Form 1 tended to convert to a non-stoichiometric hydrate of Form 1 having between 1% and about 20% by weight water (e.g., Form 13) at a relative humidity (RH) above 40~50%. The hydrate lost its water below 30% RH. Form 1 converted to a non-stoichiometric hydrate of Form 1 when exposed to air.

The dynamic vapor sorption (DVS) scan of Form 1 solid showed a 17% water absorption at 90% RH (FIG. 1D). The XRD data indicated that the solid used in the DVS test converted to the hydrate form before the start of the DVS test. However, at 0% RH, water was lost, perhaps indicating that the solid was Form 1.

Table 4. Summary of experiments that generated Form 1

Form	Solvent	Temperature	Wet	Dry
Form 1	MeOH	RT	Form 13	Form 1
	MeOH	50°C	Form 1	Form 1
	Toluene	RT	Form 1	Form 1
	Toluene	50°C	Form 1	Form 1
	water	RT	Form 13	Form 1
	Heptane	RT	Form 1	Form 1
	DCM	RT	Form 1	Form 1
	Acetonitrile/water	RT	Form 12	Form 1
	MeOH/water	RT	Form 12	Form 1
	Toluene/water	RT	Form 13	Form 1
	IPAc/water	RT	Form 13	Form 1
	EA/water	RT	Form 12	Form 1
	MtBE/water	RT	Form 12	Form 1
	MA/water	RT	Form 13	Form 1

	MIBK/water	RT	Form 12	Form 1
	MIBK/water	50°C	Form 13	Form 1
	DCM/water	RT	Form 13	Form 1
	DCM/water	50°C	Form 13	Form 1
	n-Butyl acetate/water	50°C	Form 13	Form 1
	Heptane/water	50°C	Form 13	Form 1
	n-Butanol/water	50°C	Form 13	Form 1

*Amount of water in binary solvents is 5%

Table 5. XRD peaks of Form 1

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
5.778	15.2835	57	97	28.3	1765	18.5	0.309
6.801	12.9871	19	343	100	8306	87.1	0.412
9.26	9.5427	20	178	51.9	3884	40.7	0.371
12.421	7.1203	30	231	67.3	4862	51	0.358
13.919	6.357	35	147	42.9	3668	38.5	0.424
14.501	6.1033	40	133	38.8	3439	36.1	0.44
16.5	5.3681	47	196	57.1	4286	44.9	0.372
17.26	5.1333	53	46	13.4	560	5.9	0.207
18.52	4.7868	68	342	99.7	9539	100	0.474
19.161	4.6282	54	215	62.7	4130	43.3	0.327
20.302	4.3706	49	133	38.8	2823	29.6	0.361
20.619	4.304	43	80	23.3	2047	21.5	0.435
23.056	3.8543	41	38	11.1	765	8	0.342
24.642	3.6098	33	175	51	7235	75.8	0.703
25.302	3.5171	86	80	23.3	2345	24.6	0.498
26.1	3.4113	83	69	20.1	1545	16.2	0.381
27.46	3.2453	52	46	13.4	872	9.1	0.322
28.739	3.1038	39	84	24.5	2146	22.5	0.434
30.444	2.9337	34	32	9.3	1080	11.3	0.54
33.302	2.6882	30	27	7.9	683	7.2	0.405

C. Forms 2, 2*, and 2***

The experiments that generated Forms 2, 2*, and 2** are shown in Table 6, below.

- 5 XRD scans of Forms 2, 2* and 2** were performed (FIGS. 2A, 2D, and 2G show the XRD

scans of Forms 2, 2*, and 2**, respectively). The XRD peaks of Forms 2 and 2* are shown in Tables 7 and 8, below, respectively. DSC scans were also performed (FIGS. 2B, 2E, and 2H show the DSC scans of Forms 2, 2*, and 2**, respectively). According to the DSC scans, Forms 2, 2* and 2** each showed an endotherm between 50°C-100°C, and multiple endotherms and exotherms before melting at 363°C. The endotherm before 100°C may be due to the containment of water/solvent in the solid. Form 2 was obtained from acetonitrile; Form 2* from ethanol; Form 2** from n-propanol/5% water.

A TGA scan of Form 2 (FIG. 2C) showed a 2.7% weight loss before 116°C. FIG. 2F shows the TGA scan of Form 2*

A PLM photo of Form 2 was taken (not shown), indicating that the particle size of this solid was around 50um.

The Form 2 solid was heated in a DSC machine to 90°C (past the endotherm between 50-100°C); to 270°C (past the endotherm/exotherm around 240°C); and finally to 330°C (past the exotherm around 330°C). The residual solid was analyzed by XRD. According to the first and second XRD and DSC scans, the form did not change before and after heating to 90°C. The endotherm between 50-100°C might be free solvent or hydrate. According to the first and third XRD and DSC scans, after heating a Form 2 sample to 270°C, the solid converted to low crystalline solids. According to the first and fourth XRD and DSC scans, after heating the sample to 330°C, the solid converted to Form 9. Thus, the exotherm around 290°C was a re-crystallization event. According to an XRD and DSC overlay, the behavior of Form 2* was similar to Form 2.

Residual acetonitrile and EtOH in Form 2 and 2* was not detected.

Table 6. Summary of experiments that generated Forms 2, 2*, and 2**

Form	Solvent	Temperature	Wet	Dry
Form 2	Acetonitrile	RT	Form 2	Form 2
	Acetonitrile	50°C	Form 2*	Form 2
	EtOH/water	RT	Solvate 3	Form 2
Form 2*	EtOH	RT	Form 2*	Form 2*
	EtOH	50°C	Form 2*	Form 2*
	Acetonitrile	50°C	Form 2*	Form 2
Form 2**	n-Propanol/water	RT	Form 2**	Form 2**

*Amount of water in binary solvents is 5%

Table 7. XRD peaks of Form 2

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
7.021	12.5802	164	2202	54.1	36151	38.2	0.279
8.298	10.6462	156	194	4.8	2332	2.5	0.204
10.399	8.5	193	397	9.8	6246	6.6	0.267
11.258	7.8531	206	151	3.7	1407	1.5	0.158
12.239	7.2259	181	287	7	5980	6.3	0.354
14.1	6.2759	186	648	15.9	14147	15	0.371
14.597	6.0632	195	182	4.5	7983	8.4	0.746
16.18	5.4734	235	201	4.9	4033	4.3	0.341
16.561	5.3484	251	280	6.9	8382	8.9	0.509
17.033	5.2013	288	160	3.9	1810	1.9	0.192
17.639	5.0238	295	366	9	3542	3.7	0.165
18.878	4.6968	316	1210	29.7	29303	31	0.412
19.22	4.614	333	585	14.4	21169	22.4	0.615
19.863	4.4662	340	95	2.3	437	0.5	0.078
20.411	4.3474	385	86	2.1	671	0.7	0.133
21.48	4.1335	532	1944	47.8	61345	64.8	0.536
22.04	4.0297	647	4071	100	94605	100	0.395
23.036	3.8576	634	142	3.5	1478	1.6	0.177
24.24	3.6686	497	1688	41.5	28976	30.6	0.292
25.561	3.482	422	120	2.9	2545	2.7	0.361
25.918	3.4349	365	271	6.7	11426	12.1	0.717
26.379	3.3759	349	497	12.2	15133	16	0.518
26.739	3.3313	387	181	4.4	2845	3	0.267
27.979	3.1863	297	235	5.8	4050	4.3	0.293
29.043	3.072	338	347	8.5	4584	4.8	0.225

29.661	3.0094	321	310	7.6	7879	8.3	0.432
30.204	2.9565	355	135	3.3	1501	1.6	0.189
31.58	2.8308	232	206	5.1	3991	4.2	0.329
32.602	2.7443	193	63	1.5	1129	1.2	0.305

Table 8. XRD peaks of Form 2*

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
4.859	18.1701	127	87	1.2	1714	1.9	0.335
7.119	12.4067	148	3587	48.4	44853	50.4	0.213
8.321	10.6166	149	407	5.5	4871	5.5	0.203
10.439	8.4669	186	1184	16	13629	15.3	0.196
11.319	7.8109	190	413	5.6	4673	5.3	0.192
12.3	7.1899	179	1010	13.6	13220	14.9	0.223
12.803	6.9089	182	140	1.9	1587	1.8	0.193
14.121	6.2667	179	1966	26.5	27290	30.7	0.236
14.559	6.0791	199	169	2.3	4381	4.9	0.441
16.236	5.4546	244	436	5.9	5696	6.4	0.222
16.62	5.3297	271	674	9.1	7919	8.9	0.2
17.059	5.1935	313	629	8.5	6279	7.1	0.17
17.699	5.0071	303	1094	14.7	12619	14.2	0.196
18.858	4.7018	359	2334	31.5	31734	35.7	0.231
19.321	4.5903	325	1650	22.2	28313	31.8	0.292
19.823	4.4751	412	127	1.7	582	0.7	0.078
20.321	4.3665	327	333	4.5	3361	3.8	0.172
21.479	4.1336	451	3245	43.8	56365	63.3	0.295
22.119	4.0154	612	7417	100	89000	100	0.204
22.782	3.9	536	327	4.4	11890	13.4	0.618
23.098	3.8475	466	638	8.6	11127	12.5	0.296
24.3	3.6597	361	4873	65.7	61170	68.7	0.213
25.599	3.4769	487	475	6.4	7278	8.2	0.26
25.88	3.4399	541	562	7.6	10968	12.3	0.332
26.361	3.3782	372	1289	17.4	20859	23.4	0.275
26.739	3.3312	266	660	8.9	13196	14.8	0.34
27.938	3.1909	284	560	7.6	9888	11.1	0.3
28.641	3.1142	319	210	2.8	2324	2.6	0.188

29.398	3.0357	357	100	1.3	2376	2.7	0.404
29.779	2.9977	295	708	9.5	13168	14.8	0.316
30.3	2.9473	283	451	6.1	6600	7.4	0.249
31.658	2.8239	239	667	9	9228	10.4	0.235
32.519	2.7511	221	191	2.6	2896	3.3	0.258
33.903	2.6419	213	72	1	876	1	0.207
34.82	2.5744	229	110	1.5	3822	4.3	0.591
35.504	2.5264	230	97	1.3	3876	4.4	0.679

D. Form 3

The experiments that generated Form 3 are shown in Table 9, below. XRD and DSC scans of Form 3 were taken (FIGS. 3A and 3B, respectively). Table 10, below, shows the XRD peaks of Form 3. Multiple exotherms and endotherms were observed from the DSC scan of Form 3.

A TGA scan of Form 3 was taken (FIG. 3C) and showed a 1.6% weight loss of the solid before 81°C, followed by a 1.7% weight loss between 81°C and 169°C.

Form 3 was obtained from IPAc at RT, while Form 3* was obtained from reslurry in n-butyl acetate.

10 **Table 9. Summary of experiments that generated Form 3 and Form 3***

Form	Solvent	Temperature	Wet	Dry
Form 3	IPAc	RT	Form 3	Form 3
	n-Butyl acetate	RT	Form 3*	Form 3
Form 3*	n-Butyl acetate	RT	Form 3*	Form 3

Table 10. XRD peaks of Form 3

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
5.024	17.5739	231	87	4.4	845	1.9	0.165
6.34	13.9294	368	1030	52.5	12361	27.5	0.204
7.219	12.2357	182	1962	100	36491	81.1	0.316
8.441	10.4665	188	159	8.1	3261	7.2	0.349
9.237	9.5659	207	320	16.3	3365	7.5	0.179
10.561	8.37	240	278	14.2	6270	13.9	0.383
10.998	8.0381	217	849	43.3	17119	38.1	0.343
11.46	7.715	256	87	4.4	662	1.5	0.129
12.439	7.11	215	311	15.9	6502	14.5	0.355
12.865	6.8756	209	92	4.7	1599	3.6	0.295

14.22	6.2233	231	522	26.6	12265	27.3	0.399
15.524	5.7034	273	311	15.9	2957	6.6	0.162
16.021	5.5276	309	218	11.1	2669	5.9	0.208
16.78	5.2792	368	330	16.8	3780	8.4	0.195
17.181	5.1567	384	99	5	2614	5.8	0.449
17.782	4.9837	428	496	25.3	6264	13.9	0.215
18.381	4.8227	509	551	28.1	5102	11.3	0.157
19.02	4.6622	447	589	30	20513	45.6	0.592
19.758	4.4896	487	423	21.6	14362	31.9	0.577
20.8	4.267	520	214	10.9	1518	3.4	0.121
21.19	4.1893	408	418	21.3	4581	10.2	0.186
21.6	4.1107	553	1017	51.8	41986	93.3	0.702
22.181	4.0044	662	1736	88.5	44981	100	0.44
23.185	3.8333	508	259	13.2	3327	7.4	0.218
24.44	3.6392	467	1441	73.4	29510	65.6	0.348
25.198	3.5313	551	232	11.8	1362	3	0.1
25.618	3.4745	557	79	4	365	0.8	0.079
26.103	3.4109	512	180	9.2	7374	16.4	0.696
26.479	3.3634	475	306	15.6	11652	25.9	0.647
27.3	3.264	455	133	6.8	1016	2.3	0.13
28.04	3.1796	378	93	4.7	1485	3.3	0.271
28.82	3.0953	372	201	10.2	3455	7.7	0.292
29.258	3.0499	362	76	3.9	2580	5.7	0.577
29.88	2.9878	334	191	9.7	4011	8.9	0.357
31.802	2.8115	251	205	10.4	4094	9.1	0.34
32.62	2.7429	231	87	4.4	1109	2.5	0.217
32.943	2.7167	215	52	2.7	1107	2.5	0.362
33.961	2.6375	217	101	5.1	1686	3.7	0.284

E. Form 4

The experiments that generated Forms 4, 4*, and 4** are shown in Table 11, below. XRD of Forms 4, 4*, and 4** were taken (FIGS. 4A, 4D, and 4G, respectively). Tables 12 and 13, below, show the XRD peaks of Form 4 and Form 4*, respectively. DSC scans of Forms 4, 4*, and 4** were also performed (FIGS. 4B, 4E, and 4H, respectively). According to the DSC scans, Form 4 showed an endotherm between 50°C-100°C, followed by multiple

endotherms/exotherms, and then melted at around 367°C. Forms 4* and 4** showed similar DSC patterns as Form 4.

TGA scans of Form 4, Form 4*, and Form 4** were taken (FIGS. 4C, 4F, and 4I, respectively). For Form 4, there was an 8.3% weight loss before 200°C; for Form 4*, there was a 4.4% weight loss before 102°C, followed by a 0.5% weight loss between 102°C and 250°C; and for Form 4**, there were three stages of weight loss, which were 2.8%, 1.9%, and 1.3%, respectively.

These solid forms were obtained from methyl acetate, n-propanol, MIBK, MtBE, ethyl acetate, acetone/water, and ethyl acetate/water.

10 **Table 11. Summary of experiments that generated Forms 4, 4*, and 4****

Form	Solvent	Temperature	Wet	Dry
Form 4	EA	RT	Form 4*	Form 4
	EA	50°C	Form 4*	Form 4
	MA	RT	Form 4	Form 4
	MA	50°C	Form 4	Form 4
	MA/water	50°C	Form 12	Form 4
	MtBE	50°C	Form 5*	Form 4
	n-Propanol	RT	Form 4	Form 4*
Form 4*	EA	RT	Form 4*	Form 4*
	EA	50°C	Form 4*	Form 4
	EA/water	50°C	Form 4*	Form 4*
	n-Propanol	RT	Form 4	Form 4*
Form 4**	Acetone/water	RT	Solvate 2	Form 4**
	Acetone	50°C	Solvate 2	Form 4**
	n-Propanol	50°C	Form 4	Form 4**
	Acetone/water	50°C	Form 4**	Form 4**

*Amount of water in binary solvents is 5%

Table 12. XRD peaks of Form 4

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
3.433	25.7129	197	48	1	697	0.7	0.247
7.019	12.5829	222	3897	77.3	66968	69.4	0.292
8.659	10.203	242	448	8.9	8198	8.5	0.311
8.98	9.8395	223	219	4.3	7649	7.9	0.594

9.64	9.1672	251	516	10.2	6969	7.2	0.23
10.917	8.0978	210	77	1.5	1041	1.1	0.23
12.339	7.1673	220	465	9.2	9572	9.9	0.35
13.82	6.4023	268	501	9.9	11493	11.9	0.39
14.278	6.1981	271	192	3.8	7288	7.6	0.645
14.923	5.9314	288	172	3.4	1636	1.7	0.162
16.462	5.3804	310	329	6.5	3066	3.2	0.158
17.041	5.199	375	105	2.1	942	1	0.153
17.638	5.0241	435	1073	21.3	13511	14	0.214
18.281	4.8488	487	772	15.3	9782	10.1	0.215
19.52	4.5437	504	1590	31.5	31949	33.1	0.342
21.759	4.081	677	5040	100	96504	100	0.326
23.22	3.8275	693	1457	28.9	28109	29.1	0.328
25.12	3.5421	710	3091	61.3	69330	71.8	0.381
25.76	3.4556	455	827	16.4	22029	22.8	0.453
27.221	3.2733	419	180	3.6	2915	3	0.275
28.638	3.1145	409	210	4.2	4338	4.5	0.351
29.259	3.0498	461	568	11.3	11998	12.4	0.359
30.137	2.9629	409	149	3	1946	2	0.222
31.817	2.8102	253	110	2.2	4034	4.2	0.623
32.319	2.7677	245	137	2.7	3829	4	0.475

Table 13. XRD peaks of Form 4*

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
4.981	17.7282	270	684	15.8	12231	12.6	0.304
7.22	12.2329	244	3416	79	65744	67.8	0.327
8.459	10.4447	202	335	7.7	4814	5	0.244
10.56	8.3707	219	629	14.5	10739	11.1	0.29
11.42	7.7419	240	203	4.7	2908	3	0.244
12.42	7.1209	221	614	14.2	11445	11.8	0.317
13.019	6.7947	238	59	1.4	423	0.4	0.122
14.26	6.2057	227	1052	24.3	20787	21.4	0.336
16.318	5.4274	409	85	2	665	0.7	0.133
16.722	5.2973	332	496	11.5	8980	9.3	0.308
17.199	5.1515	393	226	5.2	3448	3.6	0.259
17.82	4.9733	402	725	16.8	8502	8.8	0.199

18.98	4.672	432	1352	31.3	36895	38.1	0.464
19.44	4.5623	439	990	22.9	28546	29.4	0.49
20.46	4.3371	444	119	2.8	1163	1.2	0.166
21.58	4.1144	458	1982	45.8	71568	73.8	0.614
22.22	3.9974	837	4325	100	96937	100	0.381
23.16	3.8373	758	114	2.6	1085	1.1	0.162
24.42	3.6421	522	2466	57	48977	50.5	0.338
25.679	3.4663	590	252	5.8	5211	5.4	0.352
26.5	3.3607	470	671	15.5	23177	23.9	0.587
26.95	3.3056	356	313	7.2	3645	3.8	0.198
28.118	3.1709	385	255	5.9	5045	5.2	0.336
29.9	2.9858	360	383	8.9	13112	13.5	0.582
30.421	2.9359	346	239	5.5	5602	5.8	0.398
31.779	2.8134	293	336	7.8	5905	6.1	0.299
32.618	2.743	267	124	2.9	1934	2	0.265

F. Forms 5 and 5*

The experiments that generated Forms 5 and 5* are shown in Table 14, below. XRD scans of Forms 5 and 5* were taken (FIGS. 5A and 5D, respectively). The XRD peaks of Form 5 are shown in Table 15, below. A DSC scan of Form 5 was also performed and showed an endotherm between 50°C-100°C, and multiple endotherms and exotherms before melting at 363°C (FIG. 5B).

A TGA scan of Form 5 solid showed a 3.1% weight loss before 100°C, followed by a 1.7% weight loss between 100°C and 250°C (FIG. 5C).

Forms 5 and 5* were obtained from reslurrying Form 12 in MtBE at RT and 50°C. Wet solid showed Form 5*, while dry solid indicated Form 5.

Table 14. Summary of experiments that generated Forms 5 and 5*

Form	Solvent	Temperature	Wet	Dry
Form 5	MtBE	RT	Form 5*	Form 5
Form 5*	MtBE	RT	Form 5*	Form 5
	MtBE	50°C	Form 5*	Form 4

Table 15. XRD peaks of Form 5

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
5.098	17.3185	260	155	2.4	2464	2.1	0.27

6.38	13.8428	256	1778	27.7	34733	29.6	0.332
7.28	12.1332	214	3964	61.6	78158	66.5	0.335
8.518	10.3715	234	241	3.7	3170	2.7	0.224
9.24	9.5627	227	472	7.3	6614	5.6	0.238
10.639	8.3083	266	765	11.9	20508	17.5	0.456
11.019	8.0226	242	1596	24.8	37620	32	0.401
11.483	7.6998	398	133	2.1	949	0.8	0.121
12.44	7.1091	246	584	9.1	11910	10.1	0.347
12.94	6.8358	249	152	2.4	4189	3.6	0.469
14.301	6.1883	279	1114	17.3	22226	18.9	0.339
14.839	5.9648	300	167	2.6	5989	5.1	0.61
15.581	5.6827	404	376	5.8	4045	3.4	0.183
16.08	5.5073	452	459	7.1	9013	7.7	0.334
16.357	5.4146	509	260	4	11967	10.2	0.782
16.839	5.2606	521	473	7.4	7195	6.1	0.259
17.254	5.1351	550	258	4	4373	3.7	0.288
17.839	4.968	562	414	6.4	4207	3.6	0.173
18.439	4.8078	667	590	9.2	5946	5.1	0.171
19.059	4.6527	616	1603	24.9	35964	30.6	0.381
19.5	4.5486	671	1163	18.1	30384	25.9	0.444
20.882	4.2506	850	305	4.7	2860	2.4	0.159
21.679	4.0959	935	2272	35.3	66194	56.4	0.495
22.28	3.9867	1083	6430	100	117449	100	0.311
23.221	3.8273	856	564	8.8	9429	8	0.284
24.461	3.6361	697	4250	66.1	74709	63.6	0.299
25.276	3.5206	726	170	2.6	1349	1.1	0.135
26.081	3.4137	756	442	6.9	17518	14.9	0.674
26.52	3.3582	689	1014	15.8	34615	29.5	0.58
28.139	3.1686	528	306	4.8	4846	4.1	0.269
28.821	3.0952	533	463	7.2	7067	6	0.259
29.94	2.9819	499	755	11.7	15565	13.3	0.35
30.458	2.9324	435	467	7.3	9861	8.4	0.359
31.86	2.8065	343	648	10.1	13697	11.7	0.359
32.642	2.741	314	125	1.9	2403	2	0.327
34.002	2.6344	298	123	1.9	1956	1.7	0.27

G. Form 6

The experiments that generated Form 6 are shown in Table 16, below. XRD and DSC scans of Form 6 were taken (FIGS. 6A and 6B, respectively). According to the DSC scan, the solid showed a small exotherm at 250°C and a melting endotherm at 358°C.

- 5 Form 6 was obtained by reslurrying starting material in IPA and IPA/5% water at RT and 50°C.

Table 16. Summary of experiments that generated Form 6

Form	Solvent	Temperature	Wet	Dry
Form 6	IPA	RT	Form 6	Form 6
	IPA	50°C	Form 6	Form 6
	IPA/water	RT	Form 6	Form 6
	IPA/water	50°C	Form 6	Form 6

*Amount of water in binary solvents is 5%

H. Form 7

- 10 The experiments that generated Form 7 are shown in Table 17, below. XRD and DSC scans of Form 7 were taken (FIGS. 7A and 7B, respectively). The XRD peaks of Form 7 are shown in Table 18, below. According to the DSC scan, the solid showed two exotherms at 227°C and 299°C, followed by a melting endotherm at 365°C. Form 7 showed a low degree of crystallinity on XRD. The double exotherm on the DSC scans may be associated with the low
- 15 crystallinity observed on the XRD scan.

A TGA scan of Form 7 solid showed a 12% weight loss before 200°C (FIG. 7C).

Form 7 was obtained from MEK and MEK/5% water at RT and 50°C.

Table 17. Summary of experiments that generated Form 7

Form	Solvent	Temperature	Wet	Dry
Form 7	MEK	RT	Form 7	Form 7
	MEK	50°C	Form 7	Form 7
	MEK/water	RT	Form 7	Form 7
	MEK/water	50°C	Form 7	Form 7

*Amount of water in binary solvents is 5%

- 20 **Table 18. XRD peaks of Form 7**

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
4.94	17.8745	362	1384	23.3	50829	29.2	0.624
7.06	12.5111	286	3171	53.3	69159	39.8	0.371

8.759	10.0876	370	628	10.6	9606	5.5	0.26
9.9	8.9272	429	537	9	11110	6.4	0.352
10.881	8.1241	546	879	14.8	16425	9.4	0.318
11.84	7.4681	588	413	6.9	7187	4.1	0.296
12.997	6.8061	463	135	2.3	1351	0.8	0.17
14.404	6.1442	604	126	2.1	3331	1.9	0.449
15.1	5.8626	791	596	10	8819	5.1	0.252
15.92	5.5622	792	593	10	24460	14.1	0.701
16.581	5.3421	739	641	10.8	14919	8.6	0.396
18.5	4.7919	1066	1555	26.1	43174	24.8	0.472
19.4	4.5717	1087	930	15.6	17521	10.1	0.32
20.382	4.3535	1178	154	2.6	867	0.5	0.096
21.56	4.1183	1424	5949	100	173972	100	0.497
22.098	4.0192	1830	692	11.6	17678	10.2	0.434
23.22	3.8275	1749	1971	33.1	42151	24.2	0.364
24.203	3.6743	1776	351	5.9	11935	6.9	0.578
24.884	3.5751	1658	271	4.6	2378	1.4	0.149
25.759	3.4556	1416	492	8.3	19894	11.4	0.687
26.3	3.3858	1335	499	8.4	23631	13.6	0.805
27.34	3.2594	1192	307	5.2	4494	2.6	0.249
28.641	3.1142	1004	382	6.4	18030	10.4	0.802
29.078	3.0684	979	324	5.4	14234	8.2	0.747
30.28	2.9492	759	711	12	16004	9.2	0.383
31.985	2.7959	551	111	1.9	4816	2.8	0.738
33.402	2.6804	509	102	1.7	2060	1.2	0.343
34.24	2.6167	474	92	1.5	1901	1.1	0.351

I. Form 8

The experiments that generated Form 8 are shown in Table 19, below. XRD and DSC scans of Form 8 were taken (FIGS. 8A and 8B, respectively). The XRD peaks of Form 8 are shown in Table 20, below. According to the DSC scan, the solid showed two endotherms at 205°C and 231°C, followed by an exotherm at 279°C, followed by a melting endotherm at 362°C. Form 8 showed a low degree of crystallinity on the XRD scan. The double exotherm on the DSC scan may confirm the low crystallinity seen on XRD (low crystalline material convert to higher crystallinity solid).

A TGA scan of Form 8 showed a 4.2% weight loss before 190°C, followed by a 3.9% weight loss between 190°C and 261°C (FIG. 8C).

Form 8 was obtained from MIBK at RT and 50°C. An MIBK/5% water reslurry does not produce the same form.

5 **Table 19. Summary of experiments that generated Form 8**

Form	Solvent	Temperature	Wet	Dry
Form 8	MIBK	RT	Form 8	Form 8
	MIBK	50°C	Form 8	Form 8

Table 20. XRD peaks of Form 8

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
6.88	12.8368	318	2815	80.8	71578	51.7	0.432
10.699	8.2619	380	70	2	722	0.5	0.175
11.48	7.7016	344	466	13.4	9513	6.9	0.347
12.66	6.9866	348	136	3.9	1759	1.3	0.22
14.16	6.2496	435	166	4.8	3298	2.4	0.338
15.259	5.8017	483	269	7.7	6267	4.5	0.396
16.879	5.2484	669	333	9.6	7638	5.5	0.39
17.681	5.0121	780	1959	56.2	76035	54.9	0.66
19.618	4.5213	833	134	3.8	2110	1.5	0.268
21.5	4.1296	1116	3484	100	138450	100	0.676
24.244	3.6682	899	99	2.8	2643	1.9	0.454
27.559	3.234	753	366	10.5	11182	8.1	0.519
28.881	3.0889	636	279	8	8137	5.9	0.496
30.878	2.8935	403	87	2.5	1890	1.4	0.369
31.221	2.8624	386	69	2	1898	1.4	0.468

J. Form 9

The experiments that generated Form 9 are shown in Table 21, below. XRD and DSC scans of Form 9 were taken (FIGS. 9A and 9B, respectively). The XRD peaks of Form 9 are shown in Table 22, below. According to the DSC scan, the solid showed a single melting endotherm at 364°C.

A TGA scan of Form 9 showed a 0.28% weight loss before 100°C (FIG. 9C).

Other forms, when heated to just before melting at 364°C, seemed to convert to Form 9. This has been confirmed for Forms 1 and 2.

A DVS scan of Form 9 showed a 0.8% water absorption at 90% RH. Form 9 did not change its form before and after the DVS scan (FIG. 9D).

Table 21. Summary of experiments that generated Form 9

Form	Solvent	Temperature	Wet	Dry
	n-Butanol	RT	Form 9	Form 9
Form 9	IPAc	50°C	Form 9	Form 9
	n-Butyl acetate	50°C	Form 9	Form 9
	n-Butanol	50°C	Form 9	Form 9
	EtOH/water	50°C	Form 9	Form 9
	n-Propanol/water	50°C	Form 9	Form 9

*Amount of water in binary solvents is 5%

5 **Table 22. XRD peaks of Form 9**

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
4.94	17.8746	21	895	100	23398	100	0.444
6.26	14.1076	21	34	3.8	513	2.2	0.257
10.099	8.7516	28	66	7.4	1172	5	0.302
11.883	7.4413	30	46	5.1	828	3.5	0.306
13.16	6.7221	27	37	4.1	400	1.7	0.184
15.341	5.771	39	71	7.9	1541	6.6	0.369
16.518	5.3622	40	93	10.4	1728	7.4	0.316
18.622	4.7608	46	260	29.1	7069	30.2	0.462
19.74	4.4938	80	138	15.4	1937	8.3	0.239
21.101	4.2068	64	342	38.2	8314	35.5	0.413
22.42	3.9622	56	77	8.6	1721	7.4	0.38
24.1	3.6897	58	198	22.1	3904	16.7	0.335
25.2	3.5311	63	157	17.5	3615	15.5	0.391
26.897	3.312	46	44	4.9	1307	5.6	0.505
28.577	3.121	35	54	6	1754	7.5	0.552
29.884	2.9874	32	30	3.4	477	2	0.254
30.926	2.8891	35	32	3.6	682	2.9	0.341

K. Forms 10 and 10*

The experiments that generated Forms 10 and 10* are shown in Table 23, below. XRD scans of Forms 10 and 10* were taken (FIGS. 10A and 10D, respectively). The XRD peaks of

Form 10 are shown in Table 24, below. DSC scans of Forms 10 and 10* were also taken and indicated multiple endotherms/exotherms, followed by melting at 367°C (FIGS. 10B and 10E, respectively).

Forms 10 and 10* were produced by drying of amorphous solids (obtained from DMSO and DMSO/water reslurry at RT and 50°C). Both Form 10 and 10* are associated with DMSO.

A TGA scan of Form 10 solid showed a 0.6% weight loss before 100°C, followed by a 3.8% weight loss between 100°C and 170°C, followed by a 7.1% weight loss between 170°C and 260°C (FIG. 10C).

Table 23. Summary of experiments that generated Forms 10 and 10*

Form	Solvent	Temperature	Wet	Dry
Form 10	DMSO	RT	amorphous	Form 10
	DMSO/water	RT	amorphous	Form 10
	DMSO/water	50°C	amorphous	Form 10
Form 10*	DMSO	50°C	amorphous	Form 10*

*Amount of water in binary solvents is 5%

Table 24. XRD peaks of Form 10

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
6.701	13.1792	148	1553	32.1	31364	34.4	0.343
8.3	10.6444	207	1026	21.2	17914	19.6	0.297
9.38	9.4203	212	1352	27.9	21528	23.6	0.271
10.819	8.1705	223	514	10.6	8714	9.6	0.288
11.919	7.4192	271	635	13.1	9435	10.3	0.253
12.919	6.8469	266	1160	24	22094	24.2	0.324
13.718	6.45	242	81	1.7	856	0.9	0.18
14.84	5.9646	271	244	5	4716	5.2	0.329
15.536	5.6988	312	147	3	1304	1.4	0.151
16.58	5.3424	392	1813	37.5	30451	33.4	0.286
17.821	4.9731	434	2208	45.6	58342	64	0.449
18.16	4.881	434	2862	59.2	89029	97.6	0.529
19.001	4.6667	1021	3215	66.5	45840	50.2	0.242
19.88	4.4623	1163	1454	30.1	19014	20.8	0.222
20.701	4.2873	1514	4838	100	78140	85.7	0.275
21.66	4.0994	596	4067	84.1	91229	100	0.381
23.38	3.8017	596	2251	46.5	64928	71.2	0.49

24.22	3.6717	663	4578	94.6	84228	92.3	0.313
26	3.4242	595	430	8.9	11172	12.2	0.442
27.12	3.2853	639	146	3	1986	2.2	0.231
27.88	3.1974	642	2073	42.8	48132	52.8	0.395
28.88	3.089	638	477	9.9	14155	15.5	0.504
29.867	2.9891	544	205	4.2	4572	5	0.379
30.32	2.9454	528	568	11.7	11936	13.1	0.357
31.098	2.8735	517	443	9.2	5841	6.4	0.224
31.661	2.8236	433	118	2.4	953	1	0.137
33.379	2.6822	433	311	6.4	9235	10.1	0.505
34.22	2.6181	444	281	5.8	6059	6.6	0.367
34.822	2.5743	460	84	1.7	2707	3	0.548
35.438	2.5309	465	89	1.8	858	0.9	0.164

L. Forms 11 and 11*

The experiments that generated Forms 11 and 11* are shown in Table 25, below. XRD scans of Forms 11 and 11* were taken (FIGS. 11A and 11D, respectively). The XRD peaks of Form 11 and Form 11* are shown in Tables 26 and 27, below, respectively. DSC scans of Forms 11 and 11* were also taken (FIGS. 11B and 11E, respectively). According to the DSC scans, the solid showed multiple endotherms/exotherms and eventually melted at 368°C. Amorphous halo was observed in the XRD of both Forms. The double exotherm on the DSC of both forms may also be associated with the amorphous halo observed on XRD scans.

TGA scans of Form 11 and 11* were taken (FIGS. 11C and 11F, respectively). Form 11 solids showed a 0.8% weight loss before 100°C, followed by a 7.0% weight loss between 100°C and 249°C. Form 11* solids showed a 1.0% weight loss before 100°C, followed by a 7.0% weight loss before 250°C.

Forms 11 and 11* were obtained from DMF and DMF/5% water at RT and 50°C.

Table 25. Summary of experiments that generated Forms 11 and 11*

Form	Solvent	Temperature	Wet	Dry
Form 11	DMF	RT	Form 11	Form 11
	DMF	50°C	Form 11	Form 11*
	DMF/water	RT	Form 11	Form 11
	DMF/water	50°C	Form 11	Form 11
Form 11*	DMF	50°C	Form 11	Form 11*

*Amount of water in binary solvents is 5%

Table 26. XRD peaks of Form 11

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
6.42	13.7554	19	496	81.7	9502	100	0.326
8.421	10.4908	20	335	55.2	5775	60.8	0.293
8.86	9.9726	24	166	27.3	4268	44.9	0.437
10.859	8.1404	21	91	15	1292	13.6	0.241
12.479	7.0871	44	83	13.7	1004	10.6	0.206
12.977	6.8165	29	51	8.4	1542	16.2	0.514
14.519	6.0957	28	91	15	1421	15	0.265
16.801	5.2727	57	104	17.1	2226	23.4	0.364
17.801	4.9787	103	358	59	5109	53.8	0.243
18.519	4.7871	101	607	100	8460	89	0.237
18.861	4.7011	102	125	20.6	1763	18.6	0.24
19.922	4.453	85	383	63.1	7376	77.6	0.327
20.258	4.38	79	180	29.7	5778	60.8	0.546
20.899	4.247	76	105	17.3	1291	13.6	0.209
21.738	4.085	86	55	9.1	757	8	0.234
22.441	3.9585	94	471	77.6	7125	75	0.257
22.859	3.8871	78	167	27.5	3724	39.2	0.379
24.458	3.6365	60	298	49.1	4544	47.8	0.259
26.82	3.3213	45	195	32.1	4777	50.3	0.416
29	3.0764	43	99	16.3	3112	32.8	0.534
29.524	3.023	63	37	6.1	190	2	0.087
31.04	2.8788	38	46	7.6	826	8.7	0.305
31.825	2.8095	36	56	9.2	737	7.8	0.224
32.456	2.7563	31	40	6.6	857	9	0.364

Table 27. XRD peaks of Form 11*

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
6.441	13.7116	24	424	93.4	8643	100	0.347
6.944	12.7196	20	84	18.5	2078	24	0.421
8.518	10.3718	22	227	50	4871	56.4	0.365
8.86	9.9721	23	147	32.4	3581	41.4	0.414
10.859	8.141	26	107	23.6	1695	19.6	0.269

12.519	7.0648	34	90	19.8	2165	25	0.409
13.021	6.7935	31	54	11.9	1517	17.6	0.478
14.618	6.0547	32	76	16.7	1605	18.6	0.359
16.638	5.3238	55	115	25.3	2410	27.9	0.356
17.838	4.9684	71	368	81.1	6709	77.6	0.31
18.522	4.7864	130	454	100	7473	86.5	0.28
19.96	4.4447	109	315	69.4	6433	74.4	0.347
20.26	4.3795	109	146	32.2	5359	62	0.624
20.904	4.2461	127	58	12.8	559	6.5	0.164
21.639	4.1034	142	194	42.7	4690	54.3	0.411
22.441	3.9586	161	368	81.1	5409	62.6	0.25
22.94	3.8735	78	150	33	6057	70.1	0.686
23.398	3.7988	78	116	25.6	2330	27	0.341
24.44	3.6391	75	305	67.2	5097	59	0.284
26.819	3.3215	68	206	45.4	4795	55.5	0.396
29.018	3.0745	56	109	24	4093	47.4	0.638
29.566	3.0188	82	43	9.5	341	3.9	0.135
31.022	2.8804	58	55	12.1	509	5.9	0.157
31.881	2.8047	49	48	10.6	482	5.6	0.171
32.338	2.7661	42	50	11	1360	15.7	0.462

M. Form 13 and Form 12

The experiments that generated Form 13 and Form 12 are shown in Tables 28 and 30, below, respectively. Forms 12 and 13 are examples of non-stoichiometric hydrate forms of Form 1 that have between 1% and about 20% by weight water. XRD scans of Form 13 and Form 12 were taken (FIGS. 13A and 12A, respectively). The XRD peaks of Form 13 are shown in Table 29, below. DSC scans of Form 13 and Form 12 were also taken (FIGS. 13B and 12B, respectively). According to the DSC scan, Form 13 solids showed an endotherm between 50°C-100°C, followed by a small exotherm at 278°C; and a melting endotherm at 363°C. According to the DSC scan, Form 12 solids showed an endotherm between 50°C-100°C, followed by an exotherm at 283°C; and a melting endotherm at 364°C.

The purity of the Form 13 sample was 98.8%; the KF of an undried Form 13 sample was 35.7%. A DVS scan of Form 13 solid showed a 17% water sorption at 90% RH (FIG. 13D). Form 13 converted to Form 1 upon drying.

A TGA scan of Form 13 solid showed a 1.9% weight loss before 100°C (FIG. 13C).

Form 13 solid was heated in a DSC chamber to 170°C (past the endotherm between 50-100°C), and then scanned by XRD. A comparison of the first and the second XRD and DSC scans, after heating to 170°C, showed that Form 13 converted to Form 1. It can be concluded
5 that the endotherm between 50-100°C is due to bonded water.

Form 13 solid was heated in a DSC chamber to 330°C (past the endotherm/exotherm around 300°C), and then scanned by XRD. A comparison of the first and the third XRD and DSC scans, after heating to 170°C, showed that Form 13 converted to Form 9. It can be concluded that the endotherm/exotherm is due to melting/crystallization events.

10 **Table 28. Summary of experiments that generated Form 13**

Form	Solvent	Temperature	Wet	Dry
Form 13	MeOH	RT	Form 13	Form 1
	MeOH/water	50°C	Form 13	Form 13
	water	RT	Form 13	Form 1
	water	50°C	Form 13	Form 13
	Toluene/water	RT	Form 13	Form 1
	Toluene/water	50°C	Form 13	Form 13
	MA/water	RT	Form 13	Form 1
	n-Butyl acetate/water	RT	Form 13	Form 12
	n-Butyl acetate/water	50°C	Form 13	Form 1
	Heptane	50°C	Form 13	Form 13
	Heptane/water	RT	Form 13	Form 12
	Heptane/water	50°C	Form 13	Form 1
	n-Butanol/water	RT	Form 13	Form 13
	n-Butanol/water	50°C	Form 13	Form 1
	DCM	50°C	Form 13	Form 13
	DCM/water	RT	Form 13	Form 1
	DCM/water	50°C	Form 13	Form 1
	Acetonitrile/water	50°C	Form 13	Form 13
	IPAc/water	50°C	Form 13	Form 13
	MtBE/water	50°C	Form 13	Form 13

	MIBK/water	50°C	Form 13	Form 1
--	------------	------	---------	--------

*Amount of water in binary solvents is 5%

Table 29. XRD peaks of Form 13

2-Theta	d(A)	BG	Height	I%	Area	I%	FWHM
5.06	17.45	278	309	6.5	3685	4.8	0.203
6.379	13.8451	223	4743	100	76110	100	0.273
9.24	9.5632	164	1370	28.9	20018	26.3	0.248
11	8.0364	173	3445	72.6	51777	68	0.256
12.899	6.8574	195	173	3.6	3114	4.1	0.306
13.462	6.572	199	204	4.3	2376	3.1	0.198
14.159	6.2498	202	390	8.2	5424	7.1	0.236
15.56	5.6901	262	1335	28.1	19295	25.4	0.246
16.059	5.5145	302	1002	21.1	17561	23.1	0.298
16.841	5.26	313	774	16.3	7797	10.2	0.171
17.46	5.075	322	314	6.6	3863	5.1	0.209
18.419	4.8128	339	2354	49.6	29374	38.6	0.212
19.3	4.5951	357	210	4.4	8112	10.7	0.657
19.741	4.4935	329	1566	33	30236	39.7	0.328
20.202	4.3919	342	210	4.4	2880	3.8	0.233
20.84	4.2589	300	1054	22.2	18033	23.7	0.291
21.201	4.1873	284	964	20.3	15700	20.6	0.277
22.121	4.015	259	197	4.2	2208	2.9	0.191
23.2	3.8307	268	482	10.2	7844	10.3	0.277
24.42	3.642	280	1101	23.2	16244	21.3	0.251
24.839	3.5816	303	468	9.9	9306	12.2	0.338
25.219	3.5284	385	1093	23	16646	21.9	0.259
26.164	3.4032	359	357	7.5	5064	6.7	0.241
26.499	3.3609	402	317	6.7	7316	9.6	0.392
26.798	3.324	346	179	3.8	8025	10.5	0.762
27.339	3.2594	394	720	15.2	13063	17.2	0.308
27.639	3.2247	341	318	6.7	5673	7.5	0.303
28.799	3.0974	256	805	17	16756	22	0.354
29.902	2.9857	262	234	4.9	3508	4.6	0.255
31.234	2.8613	230	106	2.2	1473	1.9	0.236
31.96	2.798	226	308	6.5	3908	5.1	0.216

32.939	2.717	208	117	2.5	1444	1.9	0.21
33.962	2.6375	199	266	5.6	4617	6.1	0.295
34.917	2.5675	217	73	1.5	736	1	0.171

Table 30. Summary of experiments that generated Form 12

Form	Solvent	Temperature	Wet	Dry
Form 12	Acetonitrile/water	RT	Form 12	Form 1
	MeOH/water	RT	Form 12	Form 1
	IPAc/water	RT	Form 12	Form 1
	EA/water	RT	Form 12	Form 1
	MtBE/water	RT	Form 12	Form 1
	MIBK/water	RT	Form 12	Form 1
	n-Butyl acetate/water	RT	Form 13	Form 12
	Heptane/water	RT	Form 13	Form 12
	MA/water	50°C	Form 12	Form 4

*Amount of water in binary solvents is 5%

N. Solvates 1-3

The experiments that generated Solvates 1, 2, and 3 are shown in Table 31, below.

- 5 Solvates 1 and 2 solids were exposed to air overnight, and then analyzed by XRD. After the analysis, the solids were dried at 50°C under vacuum, and then analyzed by XRD again.

After exposure to air overnight, Solvate 1 converted to low crystallinity; after drying at 50°C, the sample was still a low crystallinity solid. After exposure to air overnight, the XRD pattern of Solvate 2 changed a little; after drying at 50°C, the form remained the same as the solid exposed to air overnight.

10

Table 31. Summary of experiments that generated solvates 1-3

Form	Solvent	Temperature	Wet	Dry
Solvate 1	Acetone	RT	Solvate 1	Low crystallinity
Solvate 2	Acetone/water	RT	Solvate 2	Form 4**
	Acetone	50°C	Solvate 2	Form 4**
Solvate 3	EtOH/water	RT	Solvate 3	Form 2

*Amount of water in binary solvent is 5%

EXAMPLE 5: Competitive reslurry experiments between polymorph forms

In order to find out the thermodynamic stability between the different forms, several competitive reslurry experiments were carried out. Form 1, Form 2, Form 2*, Form 3, Form 4, Form 4*, Form 4**, Form 5, Form 7, Form 8, Form 9, Form 10, Form 11, Form 11*, and Form 13 (10 mg for each) were mixed and slurried in 2 mL of solvent at both RT and 50°C. The solids were slurried for 3-5 days and then analyzed by XRD. According to the analytical data, Form 2* was the most stable form in a MeOH, EtOH, and acetone system at both RT and 50°C. Form 4 or 4* was most stable in EA at RT and 50°C. Form 13 was most stable in water at RT and 50°C. Table 32 shows the XRD scan results from the competitive reslurry experiments.

10 **Table 32. XRD scan results of competitive reslurry experiments**

Temperature	Solvent	Form after 3 days; wet/dry	Form after 5 days; wet/dry
RT	MeOH	Form 2*/Form 2*	Form 2*/Form 2*
	EtOH	Form 2*/Form 2*	Form 2*/Form 2*
	Acetone	Form 2*/Form 2*	Form 2*/Form 2*
	EA	Form 4/ Form 4	Form 4/ Form 4
	water	Form 13/ Form 13	Form 13/Form 1&Form 13
50°C	MeOH	Form 2*/Form 2*	Form 2*/Form 2*
	EtOH	Form 2*/Form 2*	Form 2*/Form 2*
	Acetone	Form 2*/Form 2*	Form 2*/Form 2*
	EA	Form 4/ Form 4	Form 4*/ Form 4*
	water	Form 13/ Form 13	Form 13/ Form 13

In order to find out the thermodynamic stability between Form 13 and Form 9, several competitive reslurry experiments were carried out. 15 mg of Form 1, Form 9 and Form 13 solids were mixed in 1 mL of toluene, IPAc, and n-butyl acetate, and slurried for 3 days at RT and 50°C.

15 The residual solid was analyzed by XRD. After a three-day reslurry, it was difficult to tell which one was more stable between Form 13 and Form 9. The XRD scan results of the experiment are shown in Table 33, below.


Table 33. XRD scan results competitive reslurry experiments

Temperature	Solvent	Form after 3 days; wet/dry
RT	Toluene	Form 13/Form 1
	IPAc	Form 9+Form 13/Form 9+Form 1

	n-Butyl acetate	Form 9+Form 13/Form 9+Form 1
50°C	Toluene	Form 9+Form 13/Form 9+Form 1
	IPAc	Form 9/Form 9
	n-Butyl acetate	Form 9+Form 13/Form 9+Form 1

WHAT IS CLAIMED IS:


1. A process for preparing a compound of Formula (1)



(1)


or a salt thereof, the process comprising:

- (a) reacting a compound of Formula (8)



(8)


or a salt thereof, with bis(pinacolato)diboron and Pd(dppf)Cl₂ to produce a compound of Formula (9)



(9)


or a salt thereof;

- (b) reacting the compound of Formula (9), or the salt thereof, with a compound of Formula (10)



(10)


or a salt thereof, with $\text{Pd}(\text{PPh}_3)_4$ and K_3PO_4 to prepare a compound of Formula (11)



(11)


or a salt thereof,

(c) reacting the compound of Formula (11), or the salt thereof, with a compound of Formula (6)



(6)

or a salt thereof, to prepare a compound of Formula (12)



(12)

or a salt thereof, and

(d) deprotecting the compound of Formula (12), or the salt thereof, to prepare the compound of Formula (1), or a salt thereof, wherein deprotecting the compound of Formula (12) to prepare the compound of Formula (1) comprises reacting the compound of Formula (12) with TFA.


2. The process of claim 1, wherein the ratio of molar equivalents of Pd(dppf)Cl_2 to the compound of Formula (9), or a salt thereof, is about 0.01:1 to about 0.1:1.

3. The process of any one of claims 1-2, wherein the ratio of molar equivalents of Pd(dppf)Cl_2 to the compound of Formula (9), or a salt thereof, is about 0.03:1.

4. The process of claim 1, wherein the ratio of molar equivalents of $\text{Pd(PPh}_3)_4$ to the compound of Formula (9), or a salt thereof, is about 0.01:1 to about 0.1:1.

5. The process of any one of claims 1 and 4, wherein the ratio of molar equivalents of $\text{Pd(PPh}_3)_4$ to the compound of Formula (9), or a salt thereof, is about 0.03:1.


6. A process for preparing a compound of Formula (1)



(1)

or a pharmaceutically acceptable salt thereof, the process comprising:

(a) reacting a compound of Formula (2)



(2)


or a salt thereof,

wherein:

R^1 is a nitrogen protecting group, and

X^1 is a first leaving group,


with a compound of Formula (4)



(4)

or a salt thereof, wherein X^2 is a second leaving group;


to prepare a compound of Formula (5)



(5)


or a salt thereof;

(b) reacting the compound of Formula (5), or the salt thereof, with a compound of Formula (6)



(6)

or a salt thereof, to prepare a compound of Formula (7)




(7)

or a salt thereof; and

(c) deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1), or the pharmaceutically acceptable salt thereof.


7. A process for preparing a compound of Formula (1):



(1)

or a pharmaceutically acceptable salt thereof, the process comprising:

(a) reacting a compound of Formula (2)



(2)


or salt thereof,

wherein:

R¹ is an nitrogen protecting group, and

X¹ is selected from the group consisting of -Cl, -Br, -I, and -OTf,

with a boron reagent to prepare a compound of Formula (3)




(3)

or salt thereof,

wherein A is selected from the group consisting of a boronic acid, a boronic ester, a boronate, a borinate, a boranate, a boranamide, an N-coordinated boronate, and a trifluoroborate;


(b) reacting the compound of Formula (3), or the salt thereof, with a compound of Formula (4)



(4)

or salt thereof,


wherein X^2 is selected from the group consisting of $-Cl$, $-Br$, $-I$, and $-OTf$,
to prepare a compound of Formula (5)



(5)


or a salt thereof,

(c) reacting the compound of Formula (5), or the salt thereof, with a compound of Formula (6)



(6)

or a salt thereof, to prepare a compound of Formula (7)



(7)


or a salt thereof; and

(d) deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1), or the pharmaceutically acceptable salt thereof.


8. The process of claim 6 or 7, wherein the nitrogen protecting group is selected from the group consisting of acetyl, benzyl, cumyl, benzhydryl, trityl, benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), benzyloxymethyl (BOM), pivaloyl-oxy-methyl (POM), trichloroethoxycarbonyl (Troc), 1-adamantyloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, trimethylsilyethoxymethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyrridiny, and pivaloyl.

9. The process of any one of claims 6-8, wherein the nitrogen protecting group is trityl.

10. The process of any one of claims 6-9, wherein A is selected from the group consisting of:



11. The process of any one of claims 6-10, wherein A is:



12. The process of any one of claims 6-11, wherein X¹ is -Br.

13. The process of any one of claims 6-11, wherein X^1 is $-I$.
14. The process of any one of claims 6-13, wherein X^2 is $-Br$.
15. The process of any one of claims 6-13, wherein X^2 is $-I$.
16. The process of any one of claims 6-15, wherein reacting the compound of Formula (2), or a salt thereof, with the boron reagent to prepare the compound of Formula (3), or a salt thereof, is performed in the presence of a palladium catalyst.
17. The process of claim 16, wherein the palladium catalyst is selected from the group consisting of palladium(II)acetate, $Pd(dppf)Cl_2$, $Pd(dba)_2$, tetrakis(triphenylphosphine)palladium(0), $(MeCN)_2PdCl_2$, and tris(dibenzylideneacetone)dipalladium(0).
18. The process of any one of claims 16-17, wherein the palladium catalyst is $Pd(dppf)Cl_2$.
19. The process of any one of claims 6 and 16-18, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (2), or a salt thereof, is about 0.01:1 to about 0.1:1.
20. The process of any one of claims 6 and 16-19, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (2), or a salt thereof, is about 0.03:1.
21. The process of any one of claims 6-20, wherein reacting the compound of Formula (2), or a salt thereof, with the boron reagent to prepare the compound of Formula (3), or a salt thereof, is performed at a temperature of about $80^\circ C$ to about $110^\circ C$ for a time of about 15 hours to about 25 hours.
22. The process of any one of claims 6-21, wherein reacting the compound of Formula (2), or a salt thereof, with the boron reagent to prepare the compound of Formula (3), or a salt thereof,

thereof, is performed at a temperature of about 85°C to about 95°C for a time of about 16 hours to about 20 hours.

23. The process of any one of claims 6-22, wherein reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed in the presence of a palladium catalyst.

24. The process of claim 23, wherein the palladium catalyst is selected from the group consisting of PdCl₂(PPh₃)₂, Pd(*t*-Bu)₃, PdCl₂ dppf CH₂Cl₂, Pd(PPh₃)₄, Pd(OAc)/PPh₃, Cl₂Pd[(Pet₃)]₂, Pd(DIPHOS)₂, Cl₂Pd(Bipy), [PdCl (Ph₂PCH₂PPh₂)]₂, Cl₂Pd[P(*o*-tolyl)₃]₂, Pd₂(dba)₃/P(*o*-tolyl)₃, Pd₂(dba)/P(furyl)₃, Cl₂Pd[P(furyl)₃]₂, Cl₂Pd(PMePh₂)₂, Cl₂Pd[P(4-F-Ph)₃]₂, Cl₂Pd[P(C₆F₅)₃]₂, Cl₂Pd[P(2-COOH-Ph)(Ph)₂]₂, Pd[P(*t*-Bu)₃]₂, PdCl₂(dppe), PdCl₂(dppp), PdCl₂[PCy₃]₂, and Cl₂Pd[P(4-COOH-Ph)(Ph)₂]₂.

25. The process of any one of claims 23-24, wherein the palladium catalyst is Pd(PPh₃)₄.

26. The process of any one of claims 23-25, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (3), or a salt thereof, is about 0.01:1 to about 0.1:1.

27. The process of any one of claims 23-26, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (3), or a salt thereof, is about 0.03:1.

28. The process of any one of claims 23-27, wherein reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, is performed in the presence of a base.

29. The process of claim 28, wherein the base is selected from the group consisting of Na₂CO₃, Ba(OH)₂, K₃PO₄, Cs₂CO₃, K₂CO₃, TIOH, KF, CsF, KO^tBu, NEt₃, Bu₄F, and NaOH.

30. The process of any one of claims 28-29, wherein the base is K₃PO₄.

31. The process of any one of claims 28-30, wherein the ratio of base to the compound of Formula (4), a salt thereof, is about 3.0:1.
32. The process of any one of claims 6-31, wherein reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed in the presence of 1,4-dioxane.
33. The process of any one of claims 6-32, wherein reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed under an inert atmosphere.
34. The process of any one of claims 6-33, wherein reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed under an N₂ atmosphere.
35. The process of any one of claims 6-34, wherein reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed at a temperature of about 80°C to about 100°C for a time of about 1 hour to about 5 hours.
36. The process of any one of claims 6-35, wherein reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed at a temperature of about 85°C to about 95°C for a time of about 2 hours to about 3 hours.
37. The process of any one of claims 6-36, further comprising precipitating the compound of Formula (5) in water prior to reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof.
38. The process of any one of claims 6-37, further comprising preparing a salt of the compound of Formula (5) prior to reacting the compound of Formula (5), or the salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof.

39. The process of any one of claims 6-38, further comprising preparing an oxalate salt of the compound of Formula (5) prior to reacting the compound of Formula (5), or the salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof.

40. The process of any one of claims 6-39, further comprising preparing a free base form of the compound of Formula (5) prior to reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof.

41. The process of any one of claims 6-40, further comprising precipitating the compound of Formula (5), or a salt thereof, in a non-polar organic solvent prior to reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof.

42. The process of any one of claims 6-40, further comprising precipitating the compound of Formula (5), or a salt thereof, in n-heptane prior to reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof.

43. The process of any one of claims 6-36, further comprising one or more of:
(a) precipitating the compound of Formula (5) in water;
(b) preparing a salt of the compound of Formula (5) from the precipitated compound of Formula (5);
(c) preparing a free base form of the compound of Formula (5) from the salt of the compound of Formula (5); and
(d) precipitating the free base form of the compound of Formula (5) in a nonpolar organic solvent,

prior to reacting the compound of Formula (5), or the salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof.

44. The process of any one of claims 6-36, further comprising one or more of:
(a) precipitating the compound of Formula (5) in water;

- (b) preparing an oxalate salt of the compound of Formula (5) from the precipitated compound of Formula (5);
- (c) preparing a free base form of the compound of Formula (5) from the salt of the compound of Formula (5); and
- (d) precipitating the free base form of the compound of Formula (5) in n-heptane, prior to reacting the compound of Formula (5), or the salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof.

45. The process of any one of claims 6-44, wherein reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed in the presence of Na_2SO_3 .

46. The process of claim 45, wherein the Na_2SO_3 is ground Na_2SO_3 .

47. The process of any one of claims 6-46, wherein reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed in the presence of *N*-methyl-2-pyrrolidone.

48. The process of any one of claims 6-47, wherein reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed under an inert atmosphere.

49. The process of any one of claims 6-48, wherein reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed under an N_2 atmosphere.

50. The process of any one of claims 6-49, wherein reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (7), or a salt thereof, is performed at a temperature of about 100°C to about 120°C for a time of about 5 hours to about 10 hours.

51. The process of any one of claims 6-50, wherein reacting the compound of Formula (5), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the

compound of Formula (7), or a salt thereof, is performed at a temperature of about 110°C to about 115°C for a time of about 7 hours to about 9 hours.

52. The process of any one of claims 6-51, further comprising preparing a salt of the compound of Formula (7) prior to deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1), or a salt thereof.

53. The process of any one of claims 6-51, further comprising preparing an oxalate salt of the compound of Formula (7) prior to deprotecting the compound of Formula (7), or the salt thereof, to prepare the compound of Formula (1), or a salt thereof.

54. The process of any one of claims 6-53, further comprising preparing a free base form of the compound of Formula (7) prior to deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof.

55. The process of any one of claims 6-54, further comprising precipitating the compound of Formula (7), or a salt thereof, in a nonpolar organic solvent prior to deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof.

56. The process of any one of claims 6-54, further comprising precipitating the compound of Formula (7), or a salt thereof, in n-heptane prior to deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof.

57. The process of any one of claims 6-51, further comprising one or more of:


- (a) preparing a salt of the compound of Formula (7);
- (b) preparing a free base form of the compound of Formula (7) from the salt of the compound of Formula (7); and
- (c) precipitating the free base form compound of Formula (7) in a nonpolar organic solvent,

prior to deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof.

58. The process of any one of claims 6-51, further comprising one or more of:

- (a) preparing an oxalate salt of the compound of Formula (7);
- (b) preparing a free base form of the compound of Formula (7) from the oxalate salt of the compound of Formula (7); and
- (c) precipitating the free base form compound of Formula (7) in n-heptane, prior to deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof.


59. A process for preparing a compound of Formula (1):



(1)


or a salt thereof, the process comprising:

- (a) reacting a compound of Formula (8)



(8)


or a salt thereof, with bis(pinacolato)diboron to produce a compound of Formula (9)



(9)


or a salt thereof;

(b) reacting the compound of Formula (9), or the salt thereof, with a compound of Formula (10)



(10)


or a salt thereof, to prepare a compound of Formula (11)



(11)


or a salt thereof,

(c) reacting the compound of Formula (11), or the salt thereof, with a compound of Formula (6)



(6)

or a salt thereof, to prepare a compound of Formula (12)



(12)

or a salt thereof; and

(d) deprotecting the compound of Formula (12), or the salt thereof, to prepare the compound of Formula (1), or a salt thereof.

60. The process of claim 59, wherein reacting the compound of Formula (8), or a salt thereof, with the boron reagent to prepare the compound of Formula (9), or a salt thereof, is performed in the presence of a palladium catalyst.

61. The process of any one of claims 59-60, wherein the palladium catalyst is selected from the group consisting of palladium(II)acetate, Pd(dppf)Cl₂, Pd(dba)₂, tetrakis(triphenylphosphine)palladium(0), (MeCN)₂PdCl₂, and tris(dibenzylideneacetone)dipalladium(0).

62. The process of any one of claims 59-61, wherein the palladium catalyst is Pd(dppf)Cl₂.

63. The process of any one of claims 60-62, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (8), or a salt thereof, is about 0.01:1 to about 0.1:1.

64. The process of any one of claims 60-63, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (8), or a salt thereof, is about 0.03:1.

65. The process of any one of claims 59-64, wherein reacting the compound of Formula (8), or a salt thereof, with the boron reagent to prepare the compound of Formula (9), or a salt thereof, is performed at a temperature of about 80°C to about 110°C for a time of about 15 hours to about 25 hours.
66. The process of any one of claims 59-65, wherein reacting the compound of Formula (8), or a salt thereof, with the boron reagent to prepare the compound of Formula (9), or a salt thereof, is performed at a temperature of about 85°C to about 95°C for a time of about 16 hours to about 20 hours.
67. The process of any one of claims 59-66, wherein reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11) is performed in the presence of a palladium catalyst.
68. The process of claim 67, wherein the palladium catalyst is selected from the group consisting of $\text{PdCl}_2(\text{PPh}_3)_2$, $\text{Pd}(\text{t-Bu})_3$, $\text{PdCl}_2 \text{ dppf } \text{CH}_2\text{Cl}_2$, $\text{Pd}(\text{PPh}_3)_4$, $\text{Pd}(\text{OAc})/\text{PPh}_3$, $\text{Cl}_2\text{Pd}[(\text{Pet}_3)]_2$, $\text{Pd}(\text{DIPHOS})_2$, $\text{Cl}_2\text{Pd}(\text{Bipy})$, $[\text{PdCl}(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]_2$, $\text{Cl}_2\text{Pd}[\text{P}(\text{o-tol})_3]_2$, $\text{Pd}_2(\text{dba})_3/\text{P}(\text{o-tol})_3$, $\text{Pd}_2(\text{dba})_3/\text{P}(\text{furyl})_3$, $\text{Cl}_2\text{Pd}[\text{P}(\text{furyl})_3]_2$, $\text{Cl}_2\text{Pd}(\text{PMePh}_2)_2$, $\text{Cl}_2\text{Pd}[\text{P}(4\text{-F-Ph})_3]_2$, $\text{Pd}[\text{P}(t\text{-Bu})_3]_2$, $\text{PdCl}_2(\text{dppe})$, $\text{PdCl}_2(\text{dppp})$, $\text{PdCl}_2[\text{PCy}_3]_2$, $\text{Cl}_2\text{Pd}[\text{P}(\text{C}_6\text{F}_5)_3]_2$, $\text{Cl}_2\text{Pd}[\text{P}(2\text{-COOH-Ph})(\text{Ph})_2]_2$, and $\text{Cl}_2\text{Pd}[\text{P}(4\text{-COOH-Ph})(\text{Ph})_2]_2$.
69. The process of any one of claims 67-68, wherein the palladium catalyst is $\text{Pd}(\text{PPh}_3)_4$.
70. The process of any one of claims 67-69, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (9) or a salt thereof, is about 0.01:1 to about 0.1:1.
71. The process of any one of claims 67-70, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (9), or a salt thereof, is about 0.03:1.
72. The process of any one of claims 67-71, wherein the reacting is performed in the presence of a base.

73. The process of claim 72, wherein the base is selected from the group consisting of Na_2CO_3 , $\text{Ba}(\text{OH})_2$, K_3PO_4 , Cs_2CO_3 , K_2CO_3 , TlOH , KF , CsF , KOtBu , NEt_3 , Bu_4F , and NaOH .
74. The process of any one of claims 72-73, wherein the base is K_3PO_4 .
75. The process of any one of claims 59-74, wherein reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed in the presence of 1,4-dioxane.
76. The process of any one of claims 59-75, wherein reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed under an inert atmosphere.
77. The process of any one of claims 59-76, wherein reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed under an N_2 atmosphere.
78. The process of any one of claims 59-77, wherein reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed at a temperature of about 80°C to about 100°C for a time of about 1 hour to about 5 hours.
79. The process of any one of claims 59-78, wherein reacting the compound of Formula (9), or a salt thereof, with the compound of Formula (10), or a salt thereof, to prepare the compound of Formula (11), or a salt thereof, is performed at a temperature of about 85°C to about 95°C for a time of about 2 hours to about 3 hours.
80. The process of any one of claims 59-79, further comprising precipitating the compound of Formula (11) in water prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.
81. The process of any one of claims 59-80, further comprising preparing a salt of the compound of Formula (11) prior to reacting the compound of Formula (11), or the salt

thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.

82. The process of any one of claims 59-81, further comprising preparing an oxalate salt of the compound of Formula (11) prior to reacting the compound of Formula (11), or the salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.

83. The process of any one of claims 59-82, further comprising preparing a free base form of the compound of Formula (11) prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.

84. The process of any one of claims 59-83, further comprising precipitating the compound of Formula (11), or a salt thereof, in a nonpolar organic solvent prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.

85. The process of any one of claims 59-84, further comprising precipitating the compound of Formula (11), or a salt thereof, in n-heptane prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.

86. The process of any one of claims 59-85, further comprising one or more of:

- (a) precipitating the compound of Formula (11) in water;
- (b) preparing a salt of the compound of Formula (11) from the precipitated compound of Formula (5);
- (c) preparing a free base form of the compound of Formula (11) from the salt of the compound of Formula (5); and
- (d) precipitating the free base form of the compound of Formula (11) in a nonpolar organic solvent,

prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.


87. The process of any one of claims 59-86, further comprising one or more of:
- (a) precipitating the compound of Formula (11) in water;
 - (b) preparing an oxalate salt of the compound of Formula (11) from the precipitated compound of Formula (11);
 - (c) preparing a free base form of the compound of Formula (11) from the salt of the compound of Formula (11); and
 - (d) precipitating the free base form of the compound of Formula (11) in *n*-heptane, prior to reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof.
88. The process of any one of claims 59-87, wherein reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, further comprises Na_2SO_3 .
89. The process of claim 88, wherein the Na_2SO_3 is ground Na_2SO_3 .
90. The process of any one of claims 59-89, wherein reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, is performed in the presence *N*-methyl-2-pyrrolidone.
91. The process of any one of claims 59-90, wherein reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, is performed under an inert atmosphere.
92. The process of any one of claims 59-91, wherein reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, is performed under an N_2 atmosphere.
93. The process of any one of claims 59-92, wherein reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, is performed at a temperature of about 100°C to about 120°C for a time of about 5 hours to about 10 hours.

94. The process of any one of claims 59-93, wherein reacting the compound of Formula (11), or a salt thereof, with the compound of Formula (6), or a salt thereof, to prepare the compound of Formula (12), or a salt thereof, is performed at a temperature of about 110°C to about 115°C for a time of about 7 hours to about 9 hours.
95. The process of any one of claims 59-94, further comprising preparing a salt of the compound of Formula (12) prior to deprotecting the compound of Formula (12), or the salt thereof, to prepare the compound of Formula (1), or a salt thereof.
96. The process of any one of claims 59-95, further comprising preparing an oxalate salt of the compound of Formula (12) prior to deprotecting the compound of Formula (12), or the salt thereof, to prepare the compound of Formula (1), or a salt thereof.
97. The process of any one of claims 59-96, further comprising preparing a free base form of the compound of Formula (12) prior to deprotecting the compound of Formula (12), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof.
98. The process of any one of claims 59-97, further comprising precipitating the compound of Formula (12), or a salt thereof, in a nonpolar organic solvent prior to deprotecting the compound of Formula (12), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof.
99. The process of any one of claims 59-98, further comprising precipitating the compound of Formula (12), or a salt thereof, in n-heptane prior to deprotecting the compound of Formula (12), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof.
100. The process of any one of claims 59-99, further comprising one or more of:
- (a) preparing a salt of the compound of Formula (12);
 - (b) preparing a free base form of the compound of Formula (12) from the salt of the compound of Formula (12); and
 - (c) precipitating the free base form compound of Formula (12) in a nonpolar organic solvent,
- prior to deprotecting the compound of Formula (12), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof.

101. The process of any one of claims 59-100, further comprising one or more of:
- (a) preparing an oxalate salt of the compound of Formula (12);
 - (b) preparing a free base form of the compound of Formula (12) from the oxalate salt of the compound of Formula (12); and
 - (c) precipitating the free base form compound of Formula (12) in n-heptane, prior to deprotecting the compound of Formula (12), to prepare the compound of Formula (1).


102. The process of claim any one of claims 59-101, further comprising preparing the compound of Formula (8), or a salt thereof, by a process comprising:

- (i) reacting a compound of Formula (13)



(13)

or a salt thereof, with sodium nitrite to prepare a compound of Formula (14)




(14)

or a salt thereof; and

- (ii) protecting the compound of Formula (14), or the salt thereof, with a trityl group to prepare the compound of Formula (8), or the salt thereof.


103. The process of any one of claims 59-102, further comprising preparing the compound of Formula (10) by a process comprising:

- (i) reacting a compound of Formula (15)



(15)

or a salt thereof, with a compound of Formula (20)



or a salt thereof,

wherein X^3 is a leaving group,

to prepare a compound of Formula (10), or the salt thereof.


104. The process of claim 103, wherein the compound of Formula (20) is isovaleryl chloride.

105. The process of claim 103 or 104, wherein the ratio of molar equivalents of the compound of Formula (10), or a salt thereof, to the compound of Formula (20), or a salt thereof, is at least 1:1.3.


106. The process of claim 103 or 104, wherein the ratio of molar equivalents of the compound of Formula (10), or a salt thereof, to the compound of Formula (20), or a salt thereof, is 1:1.3.

107. The process of any one of claims 59-106, wherein the compound of Formula (6), or a salt thereof, is prepared by a process comprising:

(i) reacting a compound of Formula (16)




or a salt thereof, with Br_2 to prepare a compound of Formula (17)




or a salt thereof,

(ii) reacting the compound of Formula (17), or the salt thereof, with a compound of Formula (18)



(18)

or a salt thereof, to prepare a compound of Formula (19)



(19)

or a salt thereof; and

(iii) selectively reducing the nitro group of the compound of Formula (19), or the salt thereof, to prepare a compound of Formula (6), or the salt thereof.

108. The process of claim 107, wherein the reacting the compound of Formula (17), or a salt thereof, with the compound of Formula (18), or a salt thereof, to prepare the compound of Formula (19), or a salt thereof, further comprises a palladium catalyst.


109. The process of claim 107 or 108, wherein the reacting the compound of Formula (17), or a salt thereof, with the compound of Formula (18), or a salt thereof, to prepare the compound of Formula (19), or a salt thereof, is carried out under an N₂ atmosphere.

110. The process of claim 107, wherein the reacting the compound of Formula (17), or a salt thereof, with the compound of Formula (18), or a salt thereof, to prepare the compound of Formula (19), or a salt thereof, further comprises a palladium catalyst and is carried out under an N₂ atmosphere.

111. The process of any one of claims 59-110, wherein deprotecting the compound of Formula (12), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof, is performed in the presence of TFA.

112. The process of any one of claims 6-58, wherein deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1), or a salt thereof, is performed in the presence of TFA.


113. A process for preparing a compound of Formula (1)



(1)


or a salt thereof, the process comprising:

reacting the compound of Formula (5)




(5)

or a salt thereof, wherein R¹ is nitrogen protecting group, with a compound of Formula (6),



(6)


or a salt thereof, to prepare a compound of Formula (7)



(7)

or a salt thereof.


114. The process of claim 113, the process further comprising deprotecting the compound of Formula (7), or a salt thereof, to prepare the compound of Formula (1)



(1)

or a salt thereof.


115. A process for preparing a compound of Formula (1)



(1)

or a salt thereof, the process comprising:


reacting a compound of Formula (2)



(2)

or a salt thereof,

wherein X^1 is selected from the group consisting of $-Cl$, $-Br$, $-I$, and $-OTf$, and R^1 is a nitrogen protecting group, with a boron reagent to prepare a compound of Formula (3)



(3)

or a salt thereof,


wherein:

A is selected from the group consisting of a boronic acid, a boronic ester, a boronate, a borinate, a boranate, a boranamide, an N-coordinated boronate, and a trifluoroborate.


116. The process of claim 115, wherein the nitrogen protecting group is selected from the group consisting of acetyl, benzyl, cumyl, benzhydryl, trityl, benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), benzyloxymethyl (BOM), pivaloyl-oxy-methyl (POM), trichloroethoxycarbonyl (Troc), 1-adamantyloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, trimethylsilyloxyethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyrrolidinyl, and pivaloyl.

117. The process of claim 115 or 116, wherein the nitrogen protecting group is trityl.

118. The process of any one of claims 115-117, wherein A is selected from the group consisting of:



119. The process of any one of claims 115-118, wherein A is:



120. The process of any one of claims 115-119, wherein X^1 is $-\text{Br}$.

121. The process of any one of claims 115-120, wherein X^1 is $-\text{I}$.

122. The process of any one of claims 115-121, wherein reacting the compound of Formula (2), or a salt thereof, with the boron reagent to prepare the compound of Formula (3), or a salt thereof, is performed in the presence of a palladium catalyst.

123. The process of claim 122, wherein the palladium catalyst is selected from the group consisting of palladium(II)acetate, $\text{Pd}(\text{dppf})\text{Cl}_2$, $\text{Pd}(\text{dba})_2$, tetrakis(triphenylphosphine)palladium(0), $(\text{MeCN})_2\text{PdCl}_2$, and tris(dibenzylideneacetone)dipalladium(0).


124. The process of claim 122, wherein the palladium catalyst is $\text{Pd}(\text{dppf})\text{Cl}_2$.

125. The process of any one of claims 122-124, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (2), or a salt thereof, is about 0.01:1 to about 0.1:1.

126. The process of any one of claims 122-125, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (2), or a salt thereof, is about 0.03:1.

127. The process of any one of claims 115-126, further comprising isolating the compound of Formula (3), or a salt thereof.

128. The process of any one of claims 115-127, further comprising reacting the compound of Formula (3), or the salt thereof, with a compound of Formula (4)




(4)

or a salt thereof,

wherein X^2 is a leaving group;

to prepare a compound of Formula (5)



(5)

or a salt thereof.

129. The process of claim 128, wherein X^2 is selected from the group consisting of $-Cl$, $-Br$, $-I$, and $-OTf$.

130. The process of claim 129, wherein X^2 is $-Br$.

131. The process of any one of claims 115-130, wherein reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, to prepare the compound of Formula (5), or a salt thereof, is performed in the presence of a palladium catalyst.

132. The process of claim 131, wherein the palladium catalyst is selected from the group consisting of $PdCl_2(PPh_3)_2$, $Pd(t-Bu)_3$, $PdCl_2 dppe$, CH_2Cl_2 , $Pd(PPh_3)_4$, $Pd(OAc)/PPh_3$, $Cl_2Pd[(Pet)_3]_2$, $Pd(DIPHOS)_2$, $Cl_2Pd(Bipy)$, $[PdCl(Ph_2PCH_2PPh_2)]_2$, $Cl_2Pd[P(o-tolyl)_3]_2$, $Pd_2(dba)_3/P(o-tolyl)_3$, $Pd_2(dba)_3/P(furyl)_3$, $Cl_2Pd[P(furyl)_3]_2$, $Cl_2Pd(PMePh_2)_2$, $Cl_2Pd[P(4-F-Ph)_3]_2$, $Cl_2Pd[P(C_6F_5)_3]_2$, $Cl_2Pd[P(2-COOH-Ph)(Ph)_2]_2$, $Pd[P(t-Bu)_3]_2$, $PdCl_2(dppe)$, $PdCl_2(dppp)$, $PdCl_2[PCy_3]_2$, and $Cl_2Pd[P(4-COOH-Ph)(Ph)_2]_2$.

133. The process of claim 131, wherein the palladium catalyst is $Pd(PPh_3)_4$.

134. The process of any one of claims 131-133, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (3), or a salt thereof, is about 0.01:1 to about 0.1:1.

135. The process of any one of claims 131-134, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (3), or a salt thereof, is about 0.03:1.


136. The process of any one of claims 131-135, wherein reacting the compound of Formula (3), or a salt thereof, with the compound of Formula (4), or a salt thereof, is performed in the presence of a base.

137. The process of claim 136, wherein the base is selected from the group consisting of Na_2CO_3 , $\text{Ba}(\text{OH})_2$, K_3PO_4 , Cs_2CO_3 , K_2CO_3 , TlOH , KF , CsF , KOtBu , NEt_3 , Bu_4F , and NaOH .


138. The process of claim 136, wherein the base is K_3PO_4 .

139. The process of any one of claims 136-138, wherein the ratio of base to the compound of Formula (4), or a salt thereof, is about 3:1.

140. A process for preparing a compound of Formula (7)




or a salt thereof, wherein R^1 is nitrogen protecting group,
comprising reacting a compound of Formula (5)



(5)


or a salt thereof, with a compound of Formula (6)



(6)

or a salt thereof, to prepare a compound of Formula (7), or the salt thereof.

141. A process for preparing a compound of Formula (3)



(3)

or a salt thereof,

wherein:

R^1 is an nitrogen protecting group, and

A is selected from the group consisting of a boronic acid, a boronic ester, a boronate, a borinate, a boranate, a boranamide, an N-coordinated boronate, and a trifluoroborate;

comprising reacting a compound of Formula (2)



(2)


or a salt thereof,

wherein X¹ is selected from the group consisting of -Cl, -Br, -I, and -OTf, with a boron reagent to prepare a compound of Formula (3), or the salt thereof.


142. The process of claim 141, wherein the nitrogen protecting group is selected from the group consisting of acetyl, benzyl, cumyl, benzhydryl, trityl, benzyloxycarbonyl (Cbz), 9-fluorenylmethyloxycarbonyl (Fmoc), benzyloxymethyl (BOM), pivaloyl-oxy-methyl (POM), trichloroethoxycarbonyl (Troc), 1-adamantyloxycarbonyl (Adoc), allyl, allyloxycarbonyl, trimethylsilyl, tert-butyl-dimethylsilyl, triethylsilyl (TES), triisopropylsilyl, trimethylsilyloxyethyl (SEM), t-butoxycarbonyl (BOC), t-butyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methyl benzene, pyrrolidinyl, and pivaloyl.

143. The process of claim 141 or 142, wherein the nitrogen protecting group is trityl.

144. The process of any one of claims 141-143, wherein A is selected from the group consisting of:




145. The process of any one of claims 141-144, wherein A is:




146. The process of any one of claims 141-145, wherein X¹ is -Br.

147. The process of any one of claims 141-145, wherein X^1 is -I.
148. The process of any one of claims 141-147, wherein reacting the compound of Formula (2), or a salt thereof, with the boron reagent to prepare the compound of Formula (3), or a salt thereof, is performed in the presence of a palladium catalyst.
149. The process of claim 148, wherein the palladium catalyst is selected from the group consisting of palladium(II)acetate, Pd(dppf)Cl_2 , Pd(dba)_2 , tetrakis(triphenylphosphine)palladium(0), $(\text{MeCN})_2\text{PdCl}_2$, and tris(dibenzylideneacetone)dipalladium(0).
150. The process of claim 149, wherein the palladium catalyst is Pd(dppf)Cl_2 .
151. The process of any one of claims 148-150, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (2), or a salt thereof, is about 0.01:1 to about 0.1:1.
152. The process of any one of claims 148-150, wherein the ratio of molar equivalents of the palladium catalyst to the compound of Formula (2), or a salt thereof, is about 0.03:1.
153. The process of any one of claims 141-152, further comprising isolating the compound of Formula (3), or a salt thereof.
154. A process for preparing a compound of Formula (1)



(1)


or a salt thereof, the process comprising deprotecting a compound of Formula (12)



(12)

or a salt thereof, to prepare the compound of Formula (1), or a salt thereof.

155. A process for preparing a pharmaceutical composition, including mixing (i) a compound of Formula (1)



(1)

or a salt thereof, prepared according to the process of any one of claims 1-154, and (ii) a pharmaceutically acceptable carrier (excipient), to form the composition.

156. The process of claim 111 or 112, wherein the TFA is neat TFA.

157. The process of any one of claims 111-112 or 156, wherein the ratio of mass equivalents of TFA to the compound of Formula (12), or a salt thereof, is about 2:1 to about 16:1.

158. The process of any one of claims 111-112 or 156, wherein the ratio of mass equivalents of TFA to the compound of Formula (12), or a salt thereof, is about 7:1 to about 9:1.

159. The process of any one of claims 111-112 or 156, wherein the ratio of mass equivalents of TFA to the compound of Formula (12), or a salt thereof, is about 8:1.
160. The process of any one of claims 111-112 or 156-159, wherein deprotecting the compound of Formula (12), or a salt thereof, is performed at a temperature of about 15°C to about 25°C.
161. The process of any one of claims 111-112 or 156-159, wherein deprotecting the compound of Formula (12), or a salt thereof, is performed at a temperature of about 20°C.
162. The process of any one of claims 111-112 or 156-161, wherein deprotecting the compound of Formula (12), or a salt thereof, is performed for a time of about 2 hours to about 7 hours.
163. The process of claim 162, wherein deprotecting the compound of Formula (12), or a salt thereof, is performed for a time of about 3 hours to about 7 hours.
164. The process of claim 162, wherein deprotecting the compound of Formula (12), or a salt thereof, is performed for a time of about 2 hours to about 4 hours.
165. The process of claim 162, wherein deprotecting the compound of Formula (12), or a salt thereof, is performed for a time of about 5 hours.
166. The process of claim 162, wherein deprotecting the compound of Formula (12), or a salt thereof, is performed for a time of about 3 hours.
167. The process of any one of claims 111-112 or 156-166, wherein deprotecting the compound of Formula (12), or a salt thereof, comprises:
 forming a first mixture; and
 adding water to the first mixture at a temperature of about 0°C to about 10°C to form a second mixture.
168. The process of claim 167, comprising adding water at a temperature of about 5°C to form the second mixture.

169. The process of claim 167 or 168, comprising reslurrying the second mixture for a time of about 0.5 hours to about 1 hour.

170. The process of claim 167 or 168, comprising reslurrying the second mixture for a time of about 0.75 hours.

171. The process of any one of claims 167-170, comprising reslurrying the second mixture at a temperature of about 0°C to about 10°C.

172. The process of claim 171, comprising reslurrying the second mixture at a temperature of about 5°C.

173. The process of any one of claims 167-172, further comprising filtering the second mixture to provide a filtrate.

174. The process of claim 173, further comprising adding water to the filtrate at a temperature of about 0°C to about 10°C to form a third mixture.

175. The process of claim 174, comprising adding water to the filtrate at a temperature of about 5°C to form the third mixture.

176. The process of claim 174 or 175, comprising reslurrying the third mixture at a temperature of about 5°C to about 15°C.

177. The process of claim 176, comprising reslurrying the third mixture at a temperature of about 10°C.

178. The process of any one of claims 174-177, comprising reslurrying the third mixture for a time of about 1 hour to about 2 hours.

179. The process of claim 178, comprising reslurrying the third mixture for a time of about 1.5 hours.

180. The process of any one of claims 174-179, further comprising filtering the third mixture to provide a first residual solid.

181. The process of claim 180, further comprising adding ethanol to the first residual solid to form a fourth mixture.

182. The process of claim 181, comprising reslurrying the fourth mixture at a temperature of about 25°C to about 35°C.

183. The process of claim 182, comprising reslurrying the fourth mixture at a temperature of about 30°C.

184. The process of any one of claims 181-183, comprising reslurrying the fourth mixture for a time of about 2 hours to about 4 hours.

185. The process of claim 184, comprising reslurrying the fourth mixture for a time of about 3 hours.

186. The process of any one of claims 181-185, further comprising filtering the fourth mixture to provide a second residual solid.

187. The process of claim 186, further comprising adding water to the second residual solid to form a fifth mixture.

188. The process of claim 187, comprising reslurrying the fifth mixture at a temperature of about 20°C to about 30°C.

189. The process of claim 188, comprising reslurrying the fifth mixture at a temperature of about 25°C.

190. The process of any one of claims 187-189, comprising reslurrying the fifth mixture for a time of about 0.5 hour to about 1.5 hours.

191. The process of claim 190, comprising reslurrying the fifth mixture for a time of about 1 hour.

192. The process of any one of claims 187-191, further comprising adding a base to the fifth mixture to form a sixth mixture.

193. The process of claim 192, wherein the base is selected from the group consisting of: lithium carbonate, sodium carbonate, potassium carbonate, cesium carbonate, sodium hydrogen carbonate, potassium hydrogen carbonate, sodium sulfate, potassium sulfate,

cesium sulfate, lithium phosphate, sodium phosphate, potassium phosphate, and cesium phosphate.

194. The process of claim 192 or 193, wherein the base is sodium carbonate.

195. The process of any one of claims 192-194, comprising reslurrying the sixth mixture at a temperature of about 20°C to about 30°C.

196. The process of claim 195, comprising reslurrying the sixth mixture at a temperature of about 25°C.

197. The process of any one of claims 192-196, comprising reslurrying the sixth mixture for a time of about 5 hours to about 7 hours.

198. The process of claim 197, comprising reslurrying the sixth mixture for a time of about 6 hours.

199. The process of any one of claims 192-198, further comprising filtering the sixth mixture to provide a third residual solid.

200. The process of claim 199, further comprising adding water to the third residual solid to form a seventh mixture.

201. The process of claim 200, comprising reslurrying the seventh mixture at a temperature of about 20°C to about 30°C.


202. The process of claim 201, comprising reslurrying the seventh mixture at a temperature of about 25°C.

203. The process of any one of claims 200-202, comprising reslurrying the seventh mixture for a time of about 5 hours to about 8 hours.

204. The process of claim 203, comprising reslurrying the seventh mixture for a time of about 6.5 hours.

205. The process of any one of claims 200-204, further comprising filtering the seventh mixture to provide a fourth residual solid.

206. The process of claim 205, further comprising:
- (a) adding water to the fourth residual solid to provide an eighth mixture;
 - (b) reslurrying the eighth mixture; and
 - (c) filtering the eighth mixture to provide a fifth residual solid.
207. The process of claim 206, wherein steps (a)-(c) are performed an additional one or more times.
208. The process of claim 207, further comprising adding isopropanol to the fifth residual solid to form a ninth mixture.
209. The process of claim 208, comprising reslurrying the ninth mixture at a temperature of about 20°C to about 30°C.
210. The process of claim 209, comprising reslurrying the ninth mixture at a temperature of about 25°C.
211. The process of any one of claims 208-210, comprising reslurrying the ninth mixture for a time of about 1 hour to about 3 hours.
212. The process of claim 211, comprising reslurrying the ninth mixture for a time of about 2 hours.
213. The process of any one of claims 208-212, further comprising filtering the ninth mixture to provide a sixth residual solid.
214. A process for preparing a polymorph form of a compound of Formula (1)



(1),

the process comprising:

(a) preparing the compound of Formula (1) according to the process of any one of claims 1, 2, 54, 108, 113, 115, 154, or 155; and

(b) converting the compound of Formula (1) to the polymorph form.

215. The process of claim 214, wherein step (b) comprises reslurrying the compound of Formula (1) or a composition comprising the compound of Formula (1) in a solvent or mixture of solvents to generate the polymorph form.

216. The process of claim 215, wherein reslurrying is performed at room temperature.

217. The process of claim 215, wherein reslurrying is performed at a temperature of about 50°C.

218. The process of claim 215, wherein reslurrying is performed at a temperature of about 30°C to about 35°C.

219. The process of claim 215, wherein reslurrying is performed for a time of about 10 hours to about 80 hours.


220. The process of claim 215, wherein reslurrying is performed for a time of about 58 hours to about 80 hours.

221. The process of any one of claims 215-220, further comprising a filtering step to provide the polymorph form as a residual solid.

222. The process of any one of claims 215-221, wherein the solvent or mixture of solvents is selected from methanol, water, or a mixture thereof.

223. The process of claim 215, wherein the polymorph form of the compound of Formula (1) is a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water.

224. A process for preparing a polymorph form of a compound of Formula (1)



(1),

the process comprising converting the compound of Formula (1) to the polymorph form.

225. The process of claim 224, comprising reslurrying the compound of Formula (1) or a composition comprising the compound of Formula (1) in a solvent or mixture of solvents to generate the polymorph form.

226. The process of claim 225, wherein reslurrying is performed at room temperature.

227. The process of claim 225, wherein reslurrying is performed at a temperature of about 50°C.

228. The process of claim 225, wherein reslurrying is performed at a temperature of about 30°C to about 35°C.

229. The process of claim 225, wherein reslurrying is performed for a time of about 10 hours to about 80 hours.

230. The process of claim 225, wherein reslurrying is performed for a time of about 58 hours to about 80 hours.

231. The process of any one of claims 225-230, further comprising a filtering step to provide the polymorph form as a residual solid.

232. The process of any one of claims 225-231, wherein the solvent or mixture of solvents is selected from methanol, water, or a mixture thereof.

233. The process of claim 225-232, wherein the polymorph form of the compound of Formula (1) is a non-stoichiometric hydrate of polymorph Form 1 having between 1% and about 20% by weight water.