

85719pct.214


**NOVEL CRYSTALLINE ANHYDRATE WITH ANTICHOLINERGIC
EFFECT**

- 5 The invention relates to a new crystalline anhydrate of tiotropium bromide, processes for preparing it and its use for preparing a pharmaceutical composition for the treatment of respiratory complaints, particularly for the treatment of COPD (chronic obstructive pulmonary disease) and asthma.

10

Background to the invention

Tiotropium bromide is known from European Patent Application EP 418 716 A1 and has the following chemical structure:



- 15 Tiotropium bromide is a highly effective anticholinergic with a long-lasting effect, which may be used to treat respiratory complaints, particularly COPD (chronic obstructive pulmonary disease) and asthma. By tiotropium is meant the free ammonium cation.
- 20 Tiotropium bromide is preferably administered by inhalation. Suitable inhalable powders packed into appropriate capsules (inhalettes) may be used. Alternatively, it may be administered by the use of suitable inhalable aerosols. These also include powdered inhalable aerosols which contain, for example, HFA134a, HFA227 or mixtures thereof as propellant gas.
- 25 The correct manufacture of the abovementioned compositions which are suitable for use for the administration of a pharmaceutically active substance by inhalation is based on various parameters which are connected with the nature of the active

substance itself. In pharmaceutical compositions which are used like tiotropium bromide in the form of inhalable powders or inhalable aerosols, the crystalline active substance is used in ground (micronised) form for preparing the formulation. Since the pharmaceutical quality of a pharmaceutical formulation requires that the active substance should always have the same crystalline modification, the stability and properties of the crystalline active substance are subject to stringent requirements from this point of view as well. It is particularly desirable that the active substance should be prepared in the form of a uniform and clearly defined crystalline modification. It is also particularly desirable that the active substance be prepared in a crystalline form which is characterised by a high degree of stability even over long storage periods. The lower the tendency of a crystalline modification to absorb moisture, for example, the greater the physical stability of its crystal structure.

The aim of the invention is therefore to provide a new stable crystal form of the compound tiotropium bromide which meets the high demands mentioned above that are made of any pharmaceutically active substance. The present invention sets out in particular to provide a crystalline modification of tiotropium bromide which has only limited hygroscopic characteristics.

Detailed description of the invention

It has been found that, depending on the choice of the conditions which may be used during the purification of the crude product obtained after industrial production, tiotropium bromide may be obtained in different crystalline modifications.

It has been found that these different modifications can be decisively obtained by the choice of solvents used for the crystallisation and by the choice of the operating conditions selected during the crystallisation process.

It has surprisingly been found that, starting from the monohydrate of tiotropium bromide, which can be obtained in crystalline form by choosing specific reaction conditions and which was described in the prior art for the first time in WO 02/30928, an anhydrous crystal modification of tiotropium bromide may be obtained which meets the high requirements set out above and thereby solves the problem underlying the present invention. Accordingly, the present invention

relates to this crystalline anhydrous tiotropium bromide. Any reference made within the scope of the present invention to the term tiotropium bromide anhydrate is to be regarded as a reference to the crystalline anhydrous tiotropium bromide according to the invention.

5

In another aspect the present invention relates to a method of preparing the new crystalline form of anhydrous tiotropium bromide which is explained by way of example in the experimental section that follows.

10 The anhydrous tiotropium bromide according to the invention is particularly characterised by slightly hygroscopic characteristics, which ensure a high degree of stability of the crystal modification. The crystalline tiotropium bromide anhydrate according to the invention is highly crystalline and is therefore particularly well suited to the preparation of pharmaceutical formulations for administration by
15 inhalation.

The present invention also relates to the use of the crystalline tiotropium bromide anhydrate according to the invention for preparing a pharmaceutical composition for the treatment of respiratory complaints, particularly for the treatment of COPD
20 and/or asthma.

The Examples that follow serve to illustrate the present invention still further, without restricting the scope of the invention to the embodiments by way of example that follow.

25

A. I. Starting materials

Tiotropium bromide monohydrate:

5 Tiotropium bromide, which was obtained for example using the method described in European Patent Application EP 418 716, was converted into crystalline tiotropium bromide monohydrate according to the disclosure of WO 02/30928. This is used as the starting compound for preparing the tiotropium bromide anhydrate according to the invention.

A. II. Examples of synthesis according to the invention

Example 1:

10.0g of tiotropium bromide monohydrate were taken up in 100 ml of water and dissolved at boiling temperature. Then 80.0g of ammonium fluoride were added.
15 After 18 hours' stirring at about 20-25°C the product that crystallises out is isolated and dried at 70°C. The crude product obtained is dissolved in 25 ml of methanol at boiling temperature, filtered hot and cooled to ambient temperature (approx. 20-25°C). The product that crystallises out immediately after the wall of the flask has been scratched (with a glass rod) is isolated and dried at 50°C .
20 Yield: 7.35 g tiotropium bromide anhydrate, white solid.

Example 2:

4.39g tiotropium bromide monohydrate were taken up in 25 ml of methanol and dissolved at boiling temperature. The clear solution is inoculated with a few crystals
25 of the crystalline tiotropium bromide anhydrate obtained according to Example 1. The product crystallises out in the warm. After slowly cooling to ambient temperature (approx. 20-25°C) the precipitate obtained is filtered off and dried at 50°C . Yield: 3,47 g tiotropium bromide anhydrate, white solid.

A. III. Characterisation of the tiotropium bromide anhydrate according to the invention

The tiotropium bromide anhydrate obtained by the above method is highly crystalline. It was investigated further by X-ray powder diffraction. The following
5 procedure was used to record the X-ray powder diagram detailed below.

The X-ray powder diagram was recorded within the scope of the present invention using a Bruker D8 Advanced with a location-sensitive detector (CuK_α - radiation, $\lambda = 1.5418 \text{ \AA}$, 30 kV, 40 mA).
10

The X-ray powder diagram obtained for the tiotropium bromide anhydrate according to the invention is shown in Figure 1.

The following Table 1 lists the characteristic peaks and standardised intensities.
15

Table 1:

2θ [°]	d_{hkl} [Å]	intensity [%]
8.39	10.53	4
11.33	7.80	27
13.50	6.55	53
14.13	6.26	65
14.70	6.02	73
15.28	5.79	31
15.72	5.63	7
15.98	5.54	33
16.32	5.43	11
16.95	5.23	45
17.90	4.95	76
18.55	4.78	100

19.07	4.65	62
19.46	4.56	36
20.32	4.37	10
20.40	4.35	12
21.75	4.08	23
22.60	3.93	77
22.83	3.89	20
23.21	3.83	86
24.01	3.70	17
24.40	3.64	13
24.93	3.57	8
25.27	3.52	60
25.55	3.48	28
25.95	3.43	7
26.21	3.40	30
26.38	3.38	18
27.26	3.27	24
27.60	3.23	8
28.07	3.18	44
28.38	3.14	67
28.58	3.12	35
29.93	2.98	19
30.18	2.96	21
30.52	2.93	10

30.91	2.89	17
31.82	2.81	46
32.32	2.77	13
32.94	2.72	7
33.49	2.67	13
34.80	2.58	31

In the above Table the value " 2Θ [$^{\circ}$]" represents the diffraction angle in degrees and the value " d_{hkl} [$\text{\AA}$]" represents the specified lattice plane intervals in $\text{\AA}$.

- 5 The present invention therefore relates to crystalline tiotropium bromide anhydrate which is characterised in that in the X-ray powder diagram it has the characteristic values $d = 6.02 \text{ \AA}$; $4.95 \text{ \AA}$; $4.78 \text{ \AA}$; $3.93 \text{ \AA}$ and $3.83 \text{ \AA}$, *inter alia*.

10 **B. Formulations containing the tiotropium bromide anhydrate according to the invention**

- The crystalline tiotropium bromide anhydrate according to the invention is highly crystalline and is therefore particularly well suited to the preparation of, for example, pharmaceutical formulations for administration by inhalation such as inhalable powders or for example propellant-containing aerosol formulations,
15 particularly inhalable powders and propellant-containing aerosol suspensions.

B.1. Inhalable powders

- The present invention also relates to inhalable powder containing 0.001 to 3 % tiotropium in the form of the crystalline tiotropium bromide anhydrate according to
20 the invention combined with a physiologically acceptable excipient. By tiotropium is meant the ammonium cation.

- Inhalable powders which contain 0.01 to 2 % tiotropium are preferred according to the invention. Particularly preferred inhalable powders contain tiotropium in an amount from about 0.03 to 1 %, preferably 0.05 to 0.6 %, particularly preferably
25 0.06 to 0.3 %. Of particular importance according to the invention, finally, are inhalable powders which contain about 0.08 to 0.22 % tiotropium.

The amounts of tiotropium specified above are based on the amount of tiotropium cation contained.

5 The excipients that are used for the purposes of the present invention are prepared by suitable grinding and/or screening using current methods known in the art. The excipients used according to the invention may also be mixtures of excipients which are obtained by mixing excipient fractions of different mean particle sizes.

10 Examples of physiologically acceptable excipients which may be used to prepare the inhalable powders for use in the inhalettes according to the invention include monosaccharides (e.g. glucose, fructose or arabinose), disaccharides (e.g. lactose, saccharose, maltose, trehalose), oligo- and polysaccharides (e.g. dextrans, dextrans, maltodextrin, starch, cellulose), polyalcohols (e.g. sorbitol, mannitol, xylitol), cyclodextrins (e.g. α -cyclodextrin, β -cyclodextrin, χ -cyclodextrin, methyl- β -
15 cyclodextrin, hydroxypropyl- β -cyclodextrin), amino acids (e.g. arginine hydrochloride) or salts (e.g. sodium chloride, calcium carbonate), or mixtures thereof. Preferably, mono- or disaccharides are used, while the use of lactose or glucose is preferred, particularly, but not exclusively, in the form of their hydrates. For the purposes of the invention, lactose is the particularly preferred excipient,
20 while lactose monohydrate is most particularly preferred.

Within the scope of the inhalable powders according to the invention the excipients have a maximum average particle size of up to 250 μ m, preferably between 10 and 150 μ m, most preferably between 15 and 80 μ m. It may sometimes seem appropriate to add finer excipient fractions with an average particle size of 1 to 9 μ m to the
25 excipients mentioned above. These finer excipients are also selected from the group of possible excipients listed hereinbefore. The average particle size may be determined using methods known in the art (cf. for example WO 02/30389, paragraphs A and C). Finally, in order to prepare the inhalable powders according to the invention, micronised crystalline tiotropium bromide anhydrate, which is
30 preferably characterised by an average particle size of 0.5 to 10 μ m, particularly preferably from 1 to 5 μ m, is added to the excipient mixture (cf. for example WO 02/30389, paragraph B). Processes for grinding and micronising active substances are known from the prior art.

If no specifically prepared excipient mixture is used as the excipient, it is particularly preferable to use excipients which have a mean particle size of 10 - 50 μm and a 10 % fine content of 0.5 to 6 μm .

5 By average particle size is meant here the 50 % value of the volume distribution measured with a laser diffractometer using the dry dispersion method . The average particle size may be determined using methods known in the art (cf. for example WO 02/30389, paragraphs A and C). Analogously, the 10% fine content in this
10 diffractometer. In other words, for the purposes of the present invention, the 10% fine content denotes the particle size below which 10% of the quantity of particles is found (based on the volume distribution).

The percentages given within the scope of the present invention are always percent
15 by weight, unless specifically stated to the contrary.

In particularly preferred inhalable powders the excipient is characterised by a mean particle size of 12 to 35 μm , particularly preferably from 13 to 30 μm .

Also particularly preferred are those inhalable powders wherein the 10 % fine
20 content is about 1 to 4 μm , preferably about 1.5 to 3 μm .

The inhalable powders according to the invention are characterised, in accordance with the problem on which the invention is based, by a high degree of homogeneity in the sense of the accuracy of single doses. This is in the region of < 8 % ,
25 preferably < 6 % , most preferably < 4 %.

After the starting materials have been weighed out the inhalable powders are prepared from the excipient and the active substance using methods known in the art. Reference may be made to the disclosure of WO 02/30390, for example. The
30 inhalable powders according to the invention may accordingly be obtained by the method described below, for example. In the preparation methods described hereinafter the components are used in the proportions by weight described in the above-mentioned compositions of the inhalable powders.

First, the excipient and the active substance are placed in a suitable mixing container. The active substance used has an average particle size of 0.5 to 10 μm , preferably 1 to 6 μm , most preferably 2 to 5 μm . The excipient and the active substance are preferably added using a sieve or a granulating sieve with a mesh size of 0.1 to 2 mm, preferably 0.3 to 1 mm, most preferably 0.3 to 0.6 mm. Preferably, the excipient is put in first and then the active substance is added to the mixing container. During this mixing process the two components are preferably added in batches. It is particularly preferred to sieve in the two components in alternate layers. The mixing of the excipient with the active substance may take place while the two components are still being added. Preferably, however, mixing is only done once the two components have been sieved in layer by layer.

The present invention also relates to the use of the inhalable powders according to the invention for preparing a pharmaceutical composition for the treatment of respiratory complaints, particularly for the treatment of COPD and/or asthma.

The inhalable powders according to the invention may for example be administered using inhalers which meter a single dose from a reservoir by means of a measuring chamber (e.g. according to US 4570630A) or by other means (e.g. according to DE 36 25 685 A). Preferably, however, the inhalable powders according to the invention are packed into capsules (to make so-called inhalettes), which are used in inhalers such as those described in WO 94/28958, for example.

Most preferably, the capsules containing the inhalable powder according to the invention are administered using an inhaler as shown in Figure 2. This inhaler is characterised by a housing 1 containing two windows 2, a deck 3 in which there are air inlet ports and which is provided with a screen 5 secured via a screen housing 4, an inhalation chamber 6 connected to the deck 3 on which there is a push button 9 provided with two sharpened pins 7 and movable counter to a spring 8, and a mouthpiece 12 which is connected to the housing 1, the deck 3 and a cover 11 via a spindle 10 to enable it to be flipped open or shut and airholes 13 for adjusting the flow resistance.

The present invention further relates to the use of the inhalable powders containing the crystalline tiotropium bromide anhydrate according to the invention for preparing a pharmaceutical composition for treating respiratory complaints, particularly for the treatment of COPD and/or asthma, characterised in that the
5 inhaler described above and shown in Figure 2 is used.

For administering the inhalable powders containing the crystalline tiotropium bromide anhydrate according to the invention using powder-filled capsules it is particularly preferred to use capsules the material of which is selected from among
10 the synthetic plastics, most preferably selected from among polyethylene, polycarbonate, polyester, polypropylene and polyethylene terephthalate. Particularly preferred synthetic plastic materials are polyethylene, polycarbonate or polyethylene terephthalate. If polyethylene is used as one of the capsule materials which is particularly preferred according to the invention, it is preferable to use polyethylene
15 with a density of between 900 and 1000 kg/m³, preferably 940 - 980 kg/m³, more preferably about 960 - 970 kg/m³ (high density polyethylene). The synthetic plastics according to the invention may be processed in various ways using manufacturing methods known in the art. Injection moulding of the plastics is preferred according to the invention. Injection moulding without the use of mould
20 release agents is particularly preferred. This method of production is well defined and is characterised by being particularly reproducible.

In another aspect the present invention relates to the abovementioned capsules which contain the abovementioned inhalable powder according to the invention.
25 These capsules may contain about 1 to 20 mg, preferably about 3 to 15 mg, most preferably about 4 to 12 mg of inhalable powder. Preferred formulations according to the invention contain 4 to 6 mg of inhalable powder. Of equivalent importance according to the invention are capsules for inhalation which contain the formulations according to the invention in an amount of from 8 to 12 mg.

30 The present invention also relates to an inhalation kit consisting of one or more of the above capsules characterised by a content of inhalable powder according to the invention in conjunction with the inhaler according to Figure 2.

The present invention also relates to the use of the abovementioned capsules characterised by a content of inhalable powder according to the invention, for preparing a pharmaceutical composition for treating respiratory complaints, especially for treating COPD and/or asthma.

Filled capsules which contain the inhalable powders according to the invention are produced by methods known in the art, by filling the empty capsules with the inhalable powders according to the invention.

B.1.1. Examples of inhalable powders according to the invention

The following Examples serve to illustrate the present invention in more detail without restricting the scope of the invention to the exemplifying embodiments that follow.

B.1.1.1. Starting materials

Active substance

The crystalline tiotropium bromide anhydrate according to the invention is used to produce the inhalable powders according to the invention. The micronisation of this active substance is carried out analogously to methods known in the art (cf for example WO 03/078429 A1). Where reference is made within the scope of the present invention to the mean particle size of the crystalline tiotropium bromide anhydrate according to the invention, this is determined using methods of measurement known in the art (cf for example WO 03/078429 A1, para. D.2).

Excipient:

In the Examples that follow lactose-monohydrate is used as excipient. It may be obtained for example from Borculo Domo Ingredients, Borculo/NL under the product name *Lactochem Extra Fine Powder*. The specifications according to the invention for the particle size and specific surface area are met by this grade of lactose. For example, in the Examples that follow, batches of lactose were used having the following specifications:

B.1.1.2. Preparation of the powder formulations according to the invention:**I) Apparatus**

- 5 The following machines and equipment, for example, may be used to prepare the inhalable powders:

Mixing container or powder mixer: Turbulamischer 2 L, Type 2C; made by Willy A. Bachofen AG, CH-4500 Basel

10

Hand-held screen: 0.135 mm mesh size

The empty inhalation capsules may be filled with inhalable powders containing tiotropium by hand or mechanically. The following equipment may be used.

15

Capsule filling machine:

MG2, Type G100, manufacturer: MG2 S.r.l, I-40065 Pian di Macina di Pianoro (BO), Italy

20 **Formulation Example 1:**

Powder mixture :

To prepare the powder mixture, 299.39 g of excipient and 0.61 g of micronised crystalline tiotropium bromide anhydrate are used.

25

About 40-45 g of excipient are placed in a suitable mixing container through a hand-held screen with a mesh size of 0.315 mm. Then crystalline tiotropium bromide anhydrate in batches of about 90-110 mg and excipient in batches of about 40-45 g are screened in in alternate layers. The excipient and active substance are added in 7 and 6 layers, respectively.

30

Having been screened in, the ingredients are then mixed (mixing speed 900 rpm). The final mixture is passed twice more through a hand-held screen and then mixed again at 900 rpm.

35

Using the method described in Example 1 it is possible to obtain inhalable powders which when packed into suitable plastic capsules may be used to produce the following capsules for inhalation, for example:

5 **Formulation Example 2:**

	tiotropium bromide anhydrate:	0.0113 mg
	lactose monohydrate:	5.4887 mg
	capsule:	100.0 mg
10	Total:	105.5 mg

Formulation Example 3:

	tiotropium bromide anhydrate:	0.0225 mg
15	lactose monohydrate:	5.4775 mg
	polyethylene capsules:	100.0 mg
	Total:	105.5 mg

Formulation Example 4:

20	tiotropium bromide anhydrate:	0.0056 mg
	lactose monohydrate:	5.4944 mg
	polyethylene capsules:	100.0 mg
	Total:	105.5 mg

25 **Formulation Example 5:**

	tiotropium bromide anhydrate:	0.0113 mg
	lactose monohydrate:*	5.4887 mg
30	capsule:	100.0 mg
	Total:	105.5 mg

*) the lactose contains 5% specifically added fine content of micronised lactose monohydrate with a mean particle size of about 4µm.

Formulation Example 6:

	tiotropium bromide anhydrate:	0.0225 mg
	lactose monohydrate:*	5.4775 mg
5	<u>polyethylene capsules:</u>	<u>100.0 mg</u>
	Total:	105.5 mg

*) the lactose contains 5% specifically added fine content of micronised lactose monohydrate with a mean particle size of about 4µm.

10 Formulation Example 7:

	tiotropium bromide anhydrate:	0.0056 mg
	lactose monohydrate:*	5.4944 mg
	<u>polyethylene capsules:</u>	<u>100.0 mg</u>
15	Total:	105.5 mg

*) the lactose contains 5% specifically added fine content of micronised lactose monohydrate with a mean particle size of about 4µm.

20 B.2. Propellant-containing aerosol suspensions containing crystalline tiotropium bromide anhydrate

The crystalline tiotropium bromide anhydrate according to the invention may optionally also be administered in the form of propellant-containing inhalable
25 aerosols. Aerosol suspensions are particularly suitable for this.

The present invention therefore also relates to suspensions of the crystalline tiotropium bromide anhydrate according to the invention in the propellant gases HFA 227 and/or HFA 134a, optionally combined with one or more other
30 propellant gases, preferably selected from the group consisting of propane, butane, pentane, dimethylether, CHClF_2 , CH_2F_2 , CF_3CH_3 , isobutane, isopentane and neopentane.

According to the invention those suspensions which contain as propellant gas only
35 HFA 227, a mixture of HFA 227 and HFA 134a or only HFA 134a are preferred.

If a mixture of the propellant gases HFA 227 and HFA 134a is used in the suspension formulations according to the invention, the weight ratios in which these two propellant gas components are used are freely variable.

If one or more other propellant gases, selected from the group consisting of propane, butane, pentane, dimethylether, CHClF_2 , CH_2F_2 , CF_3CH_3 , isobutane, isopentane and neopentane are used in addition to the propellant gases HFA 227 and/or HFA 134a in the suspension formulations according to the invention, the amount of this additional propellant gas component is preferably less than 50 %, preferably less than 40%, particularly preferably less than 30%.

10

The suspensions according to the invention preferably contain an amount of tiotropium bromide anhydrate such that the amount of tiotropium cation is between 0.001 and 0.8%, preferably between 0.08 and 0.5%, and particularly preferably between 0.2 and 0.4% according to the invention.

Unless stated to the contrary, the percentages given within the scope of the present invention are always percent by weight.

In some cases, the term suspension formulation is used within the scope of the present invention instead of the term suspension. The two terms are to be regarded as equivalent within the scope of the present invention.

20

The propellant-containing inhalable aerosols or suspension formulations according to the invention may also contain other constituents such as surface-active agents (surfactants), adjuvants, antioxidants or flavourings.

25

The surface-active agents (surfactants) optionally present in the suspensions according to the invention are preferably selected from the group consisting of Polysorbate 20, Polysorbate 80, Myvacet 9-45, Myvacet 9-08, isopropyl myristate, oleic acid, propyleneglycol, polyethyleneglycol, Brij, ethyl oleate, glyceryl trioleate, glyceryl monolaurate, glyceryl monooleate, glyceryl monostearate, glyceryl monoricinoleate, cetylalcohol, sterylalcohol, cetylpyridinium chloride, block polymers, natural oil, ethanol and isopropanol. Of the above-mentioned suspension adjuvants Polysorbate 20, Polysorbate 80, Myvacet 9-45, Myvacet 9-08 or isopropyl myristate are preferably used. Myvacet 9-45 or isopropyl myristate are most preferably used.

35

If the suspensions according to the invention contain surfactants these are preferably used in an amount of 0.0005 - 1 %, particularly preferably 0.005 - 0.5 %.

5 The adjuvants optionally contained in the suspensions according to the invention are preferably selected from the group consisting of alanine, albumin, ascorbic acid, aspartame, betaine, cysteine, phosphoric acid, nitric acid, hydrochloric acid, sulphuric acid and citric acid. Ascorbic acid, phosphoric acid, hydrochloric acid or citric acid are preferably used, while hydrochloric acid or citric acid is most
10 preferably used.

If adjuvants are present in the suspensions according to the invention, these are preferably used in an amount of 0.0001-1.0 %, preferably 0.0005-0.1 %, particularly preferably 0.001-0.01 %, while an amount of 0.001-0.005 % is
15 particularly important according to the invention.

The antioxidants optionally contained in the suspensions according to the invention are preferably selected from the group consisting of ascorbic acid, citric acid, sodium edetate, editic acid, tocopherols, butylhydroxytoluene, butylhydroxyanisol
20 and ascorbylpalmitate, while tocopherols, butylhydroxytoluene, butylhydroxyanisol or ascorbylpalmitate are preferably used.

The flavourings optionally contained in the suspensions according to the invention are preferably selected from the group consisting of peppermint, saccharine,
25 Dentomint, aspartame and ethereal oils (for example cinnamon, aniseed, menthol, camphor), of which peppermint or Dentomint® are particularly preferred.

With a view to administration by inhalation it is essential to provide the active substances in finely divided form. For this purpose, the crystalline tiotropium
30 bromide anhydrate according to the invention is obtained in finely divided form using methods known in the prior art. Methods of micronising active substances are known in the art. Preferably after micronising the active substance has a mean particle size of 0.5 to 10µm, preferably 1 to 6µm, particularly preferably 1.5 to 5µm. Preferably at least 50%, preferably at least 60%, particularly preferably at least 70%
35 of the particles of active substance have a particle size which is within the size ranges

mentioned above. Particularly preferably at least 80%, most preferably at least 90% of the particles of active substance have a particle size which is within the size ranges mentioned above.

- 5 In another aspect the present invention relates to suspensions which contain only one of the two active substances according to the invention without any other additives.

10 The suspensions according to the invention may be prepared using methods known in the art. For this, the constituents of the formulation are mixed with the propellant gas or gases (optionally at low temperatures) and filled into suitable containers.

15 The above-mentioned propellant-containing suspensions according to the invention may be administered using inhalers known in the art (pMDIs = pressurized metered dose inhalers). Accordingly, in another aspect, the present invention relates to pharmaceutical compositions in the form of suspensions as hereinbefore described combined with one or more inhalers suitable for administering these suspensions. Moreover the present invention relates to inhalers,
20 characterised in that they contain the propellant-containing suspensions according to the invention described hereinbefore.

The present invention also relates to containers (cartridges) which when fitted with a suitable valve can be used in a suitable inhaler and which contain one of the
25 above-mentioned propellant-containing suspensions according to the invention. Suitable containers (cartridges) and processes for filling these cartridges with the propellant-containing suspensions according to the invention are known in the art.

30 In view of the pharmaceutical activity of tiotropium the present invention also relates to the use of the suspensions according to the invention for preparing a pharmaceutical composition for inhalation or nasal administration, preferably for preparing a pharmaceutical composition for inhalative or nasal treatment of diseases in which anticholinergics may develop a therapeutic benefit.

Particularly preferably the present invention also relates to the use of the suspensions according to the invention for preparing a pharmaceutical composition for the inhalative treatment of respiratory complaints, preferably asthma or COPD.

- 5 The Examples that follow serve to illustrate the present invention in more detail, by way of example, without restricting it to their contents.

B.2.1 Examples of aerosol suspension formulations

- 10 Suspensions containing other ingredients in addition to active substance and propellant gas:

Formulation Example 8:

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.04
oleic acid	0.005
HFA-227	99.955

15

Formulation Example 9:

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.02
oleic acid	0.01
HFA-227	60.00
HFA-134a	39.97

Formulation Example 10:

20

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.02
isopropylmyristate	1.00
HFA-227	98.98

Formulation Example 11:

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.02
Myvacet 9-45	0.3
HFA-227	99.68

Formulation Example 12:

5

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.02
Myvacet 9-45	0.1
HFA-227	60.00
HFA-134a	39.88

Formulation Example 13:

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.04
Polysorbate 80	0.04
HFA-227	99.92

10 **Formulation Example 14:**

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.01
Polysorbate 20	0.20
HFA-227	99.78

Formulation Example 15:

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.04
Myvacet 9-08	01.00
HFA-227	98.96

Formulation Example 16:

5

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.02
isopropylmyristate	0.30
HFA-227	20.00
HFA-134a	79.68

Suspensions containing only active substance and propellant gas:

Formulation Example 17:

10

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.02
HFA-227	60.00
HFA-134a	39.98

Formulation Example 18:

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.02
HFA-227	99.98

Formulation Example 19:

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.02
HFA-134a	99.98

Formulation Example 20:

5

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.02
HFA-227	99.98

Formulation Example 21:

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.02
HFA-134a	99.98

10 **Formulation Example 22:**

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.02
HFA-227	20.00
HFA-134a	79.98

Formulation Example 23:

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.04
HFA-227	40.00
HFA-134a	59.96

15

Formulation Example 24:

constituents	concentration [% w/w]
tiotropium bromide anhydrate	0.04
HFA-227	80.00
HFA-134a	19.96

Patent Claims

1) Anhydrous crystalline tiotropium bromide which is characterised in that in the X-ray powder diagram it has the characteristic values $d = 6.02 \text{ \AA}$; $4.95 \text{ \AA}$; $4.78 \text{ \AA}$; $3.93 \text{ \AA}$ and $3.83 \text{ \AA}$, *inter alia*.

2) Pharmaceutical composition, characterised in that it contains crystalline anhydrous tiotropium bromide according to claim 1.

3) Pharmaceutical composition according to claim 2, characterised in that it is an inhalable powder.

4) Pharmaceutical composition according to claim 3, characterised in that it is an inhalable powder which contains in addition to the anhydrous crystalline tiotropium bromide one or more suitable physiologically acceptable excipients selected from among the monosaccharides, the disaccharides, the oligo- and polysaccharides, the polyalcohols, the cyclodextrins, the amino acids or the salts or mixtures of these excipients with one another.

5) Pharmaceutical composition according to claim 4, characterised in that the excipient is selected from the group consisting of glucose, fructose, arabinose, lactose, saccharose, maltose, trehalose, dextrans, dextrans, maltodextrin, starch, cellulose, sorbitol, mannitol, xylitol, α -cyclodextrin, β -cyclodextrin, χ -cyclodextrin, methyl- β -cyclodextrin, hydroxypropyl- β -cyclodextrin, arginine hydrochloride, sodium chloride or calcium carbonate, or mixtures thereof.

6) Pharmaceutical composition according to claim 4 or 5, characterised in that it contains between 0.01 and 2 % tiotropium.

7) Capsules, characterised in that in that they contain an inhalable powder according to one of claims 3 to 6.

8) Pharmaceutical composition according to claim 2, characterised in that it is a propellant-containing aerosol suspension formulation.

9) Use of crystalline anhydrous tiotropium bromide according to claim 1 for preparing a pharmaceutical composition for the treatment of respiratory complaints.

10) Use according to claim 9, characterised in that the respiratory complaints are
5 asthma or COPD.