USE OF HDAC INHIBITORS FOR THE TREATMENT OF MELANOMA

Field of the Invention

The invention relates to the use of an histone deacetylase (HDAC) inhibitor or a pharmaceutically acceptable salt thereof for the manufacture of pharmaceutical compositions for the treatment of melanoma; the use of an HDAC inhibitor or a pharmaceutically acceptable salt thereof in the treatment of melanoma; a method of treating warm-blooded animals including mammals, especially humans, suffering from melanoma by administering to a said animal in need of such treatment a dose effective against said disease of an HDAC inhibitor or a pharmaceutically acceptable salt thereof.

Background of the Invention

Melanoma is the most serious type of skin cancer. It begins in skin cells called melanocytes. Melanocytes are the cells that make melanin, which gives skin its color. Melanin also protects the deeper layers of the skin from the sun's harmful ultraviolet (UV) rays. When people spend time in the sunlight, the melanocytes make more melanin and cause the skin to tan. This also happens when skin is exposed to other forms of ultraviolet light, such as in a tanning booth. If the skin receives too much ultraviolet light, the melanocytes may begin to grow abnormally and become cancerous. This condition is called melanoma. Therefore, there is a need to develop novel treatment methods.

Summary of the Invention

The compounds of formula (I), as defined herein, are HDAC inhibitors. Reversible acetylation of histones is a major regulator of gene expression that acts by altering accessibility of transcription factors to DNA. In normal cells, histone deacetylase (HDA) and histone acetyltrasferase together control the level of acetylation of histones to maintain a balance. Inhibition of HDA results in the accumulation of hyperacetylated histones, which results in a variety of cellular responses.


Surprisingly, it was now found that HDAC inhibitors, especially the compounds of formula (I), as defined herein, treat melanoma. Hence, the invention relates to the use of an HDAC inhibitor for the preparation of a medicament for the treatment of melanoma. The

invention also relates to the use of an HDAC inhibitor or a pharmaceutically acceptable salt thereof in the treatment of melanoma. The invention relates to a method of treating warm-blooded animals including mammals, especially humans, suffering from melanoma by administering to a said animal in need of such treatment a dose effective against said disease of an HDAC inhibitor or a pharmaceutically acceptable salt thereof.

Detailed Description of the Invention

HDAC inhibitor compounds

HDAC inhibitor compounds of particular interest for use in the inventive combination are hydroxamate compounds described by the formula (I):



(I)

wherein

R_1 is H; halo; or a straight-chain C_1-C_6alkyl, especially methyl, ethyl or *n*-propyl, which methyl, ethyl and *n*-propyl substituents are unsubstituted or substituted by one or more substituents described below for alkyl substituents;

R_2 is selected from H; C_1-C_{10}alkyl, preferably C_1-C_6alkyl, e.g., methyl, ethyl or $-CH_2CH_2$-OH; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; C_4-C_9heterocycloalkylalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; $-(CH_2)_nC(O)R_6$; $-(CH_2)_nOC(O)R_6$; amino acyl; HON-C(O)-CH=C(R_1)-aryl-alkyl-; and $-(CH_2)_nR_7$;

R_3 and R_4 are the same or different and independently H, C_1-C_6alkyl, acyl or acylamino, or

R_3 and R_4, together with the carbon to which they are bound, represent C=O, C=S or C=NR_8, or

R_2, together with the nitrogen to which it is bound, and R_3, together with the carbon to which it is bound, can form a C_4-C_9heterocycloalkyl, a heteroaryl, a polyheteroaryl, a non-aromatic polyheterocycle, or a mixed aryl and non-aryl polyheterocycle ring;

R_5 is selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; acyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; aromatic polycycles; non-aromatic polycycles; mixed aryl and non-aryl polycycles; polyheteroaryl; non-aromatic polyheterocycles; and mixed aryl and non-aryl polyheterocycles;

n, n_1, n_2 and n_3 are the same or different and independently selected from 0-6, when n_1 is 1-6, each carbon atom can be optionally and independently substituted with R_3 and/or R_4;

X and Y are the same or different and independently selected from H; halo; C_1-C_4alkyl, such as CH_3 and CF_3; NO_2; $C(O)R_1$; OR_9; SR_9; CN; and $NR_{10}R_{11}$;

R_6 is selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl and 2-phenylethenyl; heteroarylalkyl, e.g., pyridylmethyl; OR_{12}; and $NR_{13}R_{14}$;

R_7 is selected from OR_{15}, SR_{15}, $S(O)R_{16}$, SO_2R_{17}, $NR_{13}R_{14}$ and $NR_{12}SO_2R_6$;

R_8 is selected from H; OR_{15}; $NR_{13}R_{14}$; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

R_9 is selected from C_1-C_4alkyl, e.g., CH_3 and CF_3; C(O)-alkyl, e.g., $C(O)CH_3$; and $C(O)CF_3$;

R_{10} and R_{11} are the same or different and independently selected from H, C_1-C_4alkyl and -C(O)-alkyl;

R_{12} is selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; C_4-C_9heterocycloalkylalkyl; aryl; mixed aryl and non-aryl polycycle; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

R_{13} and R_{14} are the same or different and independently selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; amino acyl, or

R_{13} and R_{14}, together with the nitrogen to which they are bound, are C_4-C_9heterocycloalkyl, heteroaryl, polyheteroaryl, non-aromatic polyheterocycle or mixed aryl and non-aryl polyheterocycle;

R_{15} is selected from H, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl and $(CH_2)_mZR_{12}$;

R_{16} is selected from C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, polyheteroaryl, arylalkyl, heteroarylalkyl and $(CH_2)_mZR_{12}$;

R_{17} is selected from C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, aromatic polycycles, heteroaryl, arylalkyl, heteroarylalkyl, polyheteroaryl and $NR_{13}R_{14}$;

m is an integer selected from 0-6; and

Z is selected from O; NR_{13}; S; and S(O),

or a pharmaceutically acceptable salt thereof.

As appropriate, "unsubstituted" means that there is no substituent or that the only substituents are hydrogen.

Halo substituents are selected from fluoro, chloro, bromo and iodo, preferably fluoro or chloro.

Alkyl substituents include straight- and branched-C_1-C_6alkyl, unless otherwise noted. Examples of suitable straight- and branched-C_1-C_6alkyl substituents include methyl, ethyl, *n*-propyl, 2-propyl, *n*-butyl, *sec*-butyl, *t*-butyl and the like. Unless otherwise noted, the alkyl substituents include both unsubstituted alkyl groups and alkyl groups that are substituted by one or more suitable substituents, including unsaturation, i.e., there are one or more double or triple C-C bonds; acyl; cycloalkyl; halo; oxyalkyl; alkylamino; aminoalkyl; acylamino; and OR_{15}, e.g., alkoxy. Preferred substituents for alkyl groups include halo, hydroxy, alkoxy, oxyalkyl, alkylamino and aminoalkyl.

Cycloalkyl substituents include C_3-C_9cycloalkyl groups, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like, unless otherwise specified. Unless otherwise noted, cycloalkyl substituents include both unsubstituted cycloalkyl groups and cycloalkyl groups that are substituted by one or more suitable substituents, including C_1-C_6alkyl, halo, hydroxy, aminoalkyl, oxyalkyl, alkylamino and OR_{15}, such as alkoxy. Preferred substituents for cycloalkyl groups include halo, hydroxy, alkoxy, oxyalkyl, alkylamino and aminoalkyl.

The above discussion of alkyl and cycloalkyl substituents also applies to the alkyl portions of other substituents, such as, without limitation, alkoxy, alkyl amines, alkyl ketones, arylalkyl, heteroarylalkyl, alkylsulfonyl and alkyl ester substituents and the like.

Heterocycloalkyl substituents include 3- to 9-membered aliphatic rings, such as 4- to 7-membered aliphatic rings, containing from 1-3 heteroatoms selected from nitrogen, sulfur,

oxygen. Examples of suitable heterocycloalkyl substituents include pyrrolidyl, tetrahydrofuryl, tetrahydrothiofuranyl, piperidyl, piperazyl, tetrahydropyranyl, morphilino, 1,3-diazapane, 1,4-diazapane, 1,4-oxazepane and 1,4-oxathiapane. Unless otherwise noted, the rings are unsubstituted or substituted on the carbon atoms by one or more suitable substituents, including C_1-C_6alkyl; C_4-C_9cycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; halo; amino; alkyl amino and OR_{15}, e.g., alkoxy. Unless otherwise noted, nitrogen heteroatoms are unsubstituted or substituted by H, C_1-C_4alkyl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; acyl; aminoacyl; alkylsulfonyl; and arylsulfonyl.

Cycloalkylalkyl substituents include compounds of the formula -$(CH_2)_{n5}$-cycloalkyl, wherein n5 is a number from 1-6. Suitable alkylcycloalkyl substituents include cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl and the like. Such substituents are unsubstituted or substituted in the alkyl portion or in the cycloalkyl portion by a suitable substituent, including those listed above for alkyl and cycloalkyl.

Aryl substituents include unsubstituted phenyl and phenyl substituted by one or more suitable substituents including C_1-C_6alkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; O(CO)alkyl; oxyalkyl; halo; nitro; amino; alkylamino; aminoalkyl; alkyl ketones; nitrile; carboxyalkyl; alkylsulfonyl; aminosulfonyl; arylsulfonyl and OR_{15}, such as alkoxy. Preferred substituents include including C_1-C_6alkyl; cycloalkyl, e.g., cyclopropylmethyl; alkoxy; oxyalkyl; halo; nitro; amino; alkylamino; aminoalkyl; alkyl ketones; nitrile; carboxyalkyl; alkylsulfonyl; arylsulfonyl and aminosulfonyl. Examples of suitable aryl groups include C_1-C_4alkylphenyl, C_1-C_4alkoxyphenyl, trifluoromethylphenyl, methoxyphenyl, hydroxyethylphenyl, dimethylaminophenyl, aminopropylphenyl, carbethoxyphenyl, methanesulfonylphenyl and tolylsulfonylphenyl.

Aromatic polycycles include naphthyl, and naphthyl substituted by one or more suitable substituents including C_1-C_6alkyl; alkylcycloalkyl, e.g., cyclopropylmethyl; oxyalkyl; halo; nitro; amino; alkylamino; aminoalkyl; alkyl ketones; nitrile; carboxyalkyl; alkylsulfonyl; arylsulfonyl; aminosulfonyl and OR_{15}, such as alkoxy.

Heteroaryl substituents include compounds with a 5- to 7-membered aromatic ring containing one or more heteroatoms, e.g., from 1-4 heteroatoms, selected from N, O and S. Typical heteroaryl substituents include furyl, thienyl, pyrrole, pyrazole, triazole, thiazole, oxazole, pyridine, pyrimidine, isoxazolyl, pyrazine and the like. Unless otherwise noted,

heteroaryl substituents are unsubstituted or substituted on a carbon atom by one or more suitable substituents, including alkyl, the alkyl substituents identified above, and another heteroaryl substituent. Nitrogen atoms are unsubstituted or substituted, e.g., by R_{13}; especially useful N substituents include H, C_1-C_4alkyl, acyl, aminoacyl and sulfonyl.

Arylalkyl substituents include groups of the formula -$(CH_2)_{n5}$-aryl, -$(CH_2)_{n5-1}$-(CH-aryl)-$(CH_2)_{n5}$-aryl or -$(CH_2)_{n5-1}$CH(aryl)(aryl), wherein aryl and n5 are defined above. Such arylalkyl substituents include benzyl, 2-phenylethyl, 1-phenylethyl, tolyl-3-propyl, 2-phenylpropyl, diphenylmethyl, 2-diphenylethyl, 5,5-dimethyl-3-phenylpentyl and the like. Arylalkyl substituents are unsubstituted or substituted in the alkyl moiety or the aryl moiety or both as described above for alkyl and aryl substituents.

Heteroarylalkyl substituents include groups of the formula -$(CH_2)_{n5}$-heteroaryl, wherein heteroaryl and n5 are defined above and the bridging group is linked to a carbon or a nitrogen of the heteroaryl portion, such as 2-, 3- or 4-pyridylmethyl, imidazolylmethyl, quinolylethyl and pyrrolylbutyl. Heteroaryl substituents are unsubstituted or substituted as discussed above for heteroaryl and alkyl substituents.

Amino acyl substituents include groups of the formula -C(O)-$(CH_2)_n$-C(H)($NR_{13}R_{14}$)-$(CH_2)_n$-R_5, wherein n, R_{13}, R_{14} and R_5 are described above. Suitable aminoacyl substituents include natural and non-natural amino acids, such as glycinyl, D-tryptophanyl, L-lysinyl, D- or L-homoserinyl, 4-aminobutryic acyl and ±-3-amin-4-hexenoyl.

Non-aromatic polycycle substituents include bicyclic and tricyclic fused ring systems where each ring can be 4- to 9-membered and each ring can contain zero, one or more double and/or triple bonds. Suitable examples of non-aromatic polycycles include decalin, octahydroindene, perhydrobenzocycloheptene and perhydrobenzo-[*f*]-azulene. Such substituents are unsubstituted or substituted as described above for cycloalkyl groups.

Mixed aryl and non-aryl polycycle substituents include bicyclic and tricyclic fused ring systems where each ring can be 4- to 9-membered and at least one ring is aromatic. Suitable examples of mixed aryl and non-aryl polycycles include methylenedioxyphenyl, *bis*-methylenedioxyphenyl, 1,2,3,4-tetrahydronaphthalene, dibenzosuberane, dihydroanthracene and 9*H*-fluorene. Such substituents are unsubstituted or substituted by nitro or as described above for cycloalkyl groups.

Polyheteroaryl substituents include bicyclic and tricyclic fused ring systems where each ring can independently be 5- or 6-membered and contain one or more heteroatom, e.g., 1, 2, 3 or 4 heteroatoms, chosen from O, N or S such that the fused ring system is aromatic. Suitable examples of polyheteroaryl ring systems include quinoline, isoquinoline, pyridopyrazine, pyrrolopyridine, furopyridine, indole, benzofuran, benzothiofuran, benzindole, benzoxazole, pyrroloquinoline and the like. Unless otherwise noted, polyheteroaryl substituents are unsubstituted or substituted on a carbon atom by one or more suitable substituents, including alkyl, the alkyl substituents identified above and a substituent of the formula $-O-(CH_2CH=CH(CH_3)(CH_2))_{1-3}H$. Nitrogen atoms are unsubstituted or substituted, e.g., by R_{13}, especially useful N substituents include H, C_1-C_4alkyl, acyl, aminoacyl and sulfonyl.

Non-aromatic polyheterocyclic substituents include bicyclic and tricyclic fused ring systems where each ring can be 4- to 9-membered, contain one or more heteroatom, e.g., 1, 2, 3 or 4 heteroatoms, chosen from O, N or S and contain zero or one or more C-C double or triple bonds. Suitable examples of non-aromatic polyheterocycles include hexitol, *cis*-perhydro-cyclohepta[*b*]pyridinyl, decahydro-benzo[*f*][1,4]oxazepinyl, 2,8-dioxabicyclo[3.3.0]octane, hexahydro-thieno[3,2-*b*]thiophene, perhydropyrrolo[3,2-*b*]pyrrole, perhydronaphthyridine, perhydro-1*H*-dicyclopenta[*b,e*]pyran. Unless otherwise noted, non-aromatic polyheterocyclic substituents are unsubstituted or substituted on a carbon atom by one or more substituents, including alkyl and the alkyl substituents identified above. Nitrogen atoms are unsubstituted or substituted, e.g., by R_{13}, especially useful N substituents include H, C_1-C_4alkyl, acyl, aminoacyl and sulfonyl.

Mixed aryl and non-aryl polyheterocycles substituents include bicyclic and tricyclic fused ring systems where each ring can be 4- to 9-membered, contain one or more heteroatom chosen from O, N or S, and at least one of the rings must be aromatic. Suitable examples of mixed aryl and non-aryl polyheterocycles include 2,3-dihydroindole, 1,2,3,4-tetrahydroquinoline, 5,11-dihydro-10*H*-dibenz[*b,e*][1,4]diazepine, 5*H*-dibenzo[*b,e*][1,4]diazepine, 1,2-dihydropyrrolo[3,4-*b*][1,5]benzodiazepine, 1,5-dihydro-pyrido[2,3-*b*][1,4]diazepin-4-one, 1,2,3,4,6,11-hexahydro-benzo[*b*]pyrido[2,3-*e*][1,4]diazepin-5-one. Unless otherwise noted, mixed aryl and non-aryl polyheterocyclic substituents are unsubstituted or substituted on a carbon atom by one or more suitable substituents including -N-OH, =N-OH, alkyl and the alkyl substituents identified above. Nitrogen atoms are

unsubstituted or substituted, e.g., by R_{13}; especially useful N substituents include H, C_1-C_4alkyl, acyl, aminoacyl and sulfonyl.


Amino substituents include primary, secondary and tertiary amines and in salt form, quaternary amines. Examples of amino substituents include mono- and di-alkylamino, mono- and di-aryl amino, mono- and di-arylalkyl amino, aryl-arylalkylamino, alkyl-arylamino, alkyl-arylalkylamino and the like.

Sulfonyl substituents include alkylsulfonyl and arylsulfonyl, e.g., methane sulfonyl, benzene sulfonyl, tosyl and the like.

Acyl substituents include groups of formula -C(O)-W, -OC(O)-W, -C(O)-O-W or $-C(O)NR_{13}R_{14}$, where W is R_{16}, H or cycloalkylalkyl.

Acylamino substituents include substituents of the formula $-N(R_{12})C(O)-W$, $-N(R_{12})C(O)-O-W$ and $-N(R_{12})C(O)-NHOH$ and R_{12} and W are defined above.

The R_2 substituent HON-C(O)-CH=C(R_1)-aryl-alkyl- is a group of the formula:



Preferences for each of the substituents include the following:

R_1 is H, halo or a straight-chain C_1-C_4alkyl;

R_2 is selected from H, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, cycloalkylalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, $-(CH_2)_nC(O)R_6$, amino acyl and $-(CH_2)_nR_7$;

R_3 and R_4 are the same or different and independently selected from H and C_1-C_6alkyl, or

R_3 and R_4, together with the carbon to which they are bound, represent C=O, C=S or $C=NR_8$;

R_5 is selected from H, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, a aromatic polycycle, a non-aromatic polycycle, a mixed aryl and non-aryl polycycle, polyheteroaryl, a non-aromatic polyheterocycle, and a mixed aryl and non-aryl polyheterocycle;

n, n_1, n_2 and n_3 are the same or different and independently selected from 0-6, when n_1 is 1-6, each carbon atom is unsubstituted or independently substituted with R_3 and/or R_4;

X and Y are the same or different and independently selected from H, halo, C_1-C_4alkyl, CF_3, NO_2, $C(O)R_1$, OR_9, SR_9, CN and $NR_{10}R_{11}$;

R_6 is selected from H, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, alkylcycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, OR_{12} and $NR_{13}R_{14}$;

R_7 is selected from OR_{15}, SR_{15}, $S(O)R_{16}$, SO_2R_{17}, $NR_{13}R_{14}$ and $NR_{12}SO_2R_6$;

R_8 is selected from H, OR_{15}, $NR_{13}R_{14}$, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, arylalkyl and heteroarylalkyl;

R_9 is selected from C_1-C_4alkyl and C(O)-alkyl;

R_{10} and R_{11} are the same or different and independently selected from H, C_1-C_4alkyl and -C(O)-alkyl;

R_{12} is selected from H, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, arylalkyl and heteroarylalkyl;

R_{13} and R_{14} are the same or different and independently selected from H, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl and amino acyl;

R_{15} is selected from H, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl and $(CH_2)_mZR_{12}$;

R_{16} is selected from C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl and $(CH_2)_mZR_{12}$;

R_{17} is selected from C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl and $NR_{13}R_{14}$;


m is an integer selected from 0-6; and

Z is selected from O, NR13, S and S(O);

or a pharmaceutically acceptable salt thereof.

Useful compounds of the formula (I), include those wherein each of R_1, X, Y, R_3 and R_4 is H, including those wherein one of n_2 and n_3 is 0 and the other is 1, especially those wherein R_2 is H or $-CH_2-CH_2-OH$.

One suitable genus of hydroxamate compounds are those of formula (Ia):



(Ia)

,

wherein


n_4 is 0-3;

R_2 is selected from H, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, cycloalkylalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, -$(CH_2)_nC(O)R_6$, amino acyl and -$(CH_2)_nR_7$; and

R_5' is heteroaryl; heteroarylalkyl, e.g., pyridylmethyl; aromatic polycycles; non-aromatic polycycles; mixed aryl and non-aryl polycycles; polyheteroaryl or mixed aryl; and non-aryl polyheterocycles;

or a pharmaceutically acceptable salt thereof.

Another suitable genus of hydroxamate compounds are those of formula (Ia):



(Ia)

,

wherein


n_4 is 0-3;

R_2 is selected from H, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, cycloalkylalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, -$(CH_2)_nC(O)R_6$, amino acyl and -$(CH_2)_nR_7$;

R_5' is aryl; arylalkyl; aromatic polycycles; non-aromatic polycycles and mixed aryl; and non-aryl polycycles, especially aryl, such as *p*-fluorophenyl, *p*-chlorophenyl, *p*-*O*-C_1-C_4alkylphenyl, such as *p*-methoxyphenyl, and *p*-C_1-C_4alkylphenyl; and arylalkyl, such as benzyl, *ortho*-, *meta*- *or para*-fluorobenzyl, *ortho*-, *meta*- *or para*-chlorobenzyl, *ortho*-, *meta*- *or para*-mono, di- or tri-*O*-C_1-C_4alkylbenzyl, such as *ortho*-, *meta*- *or para*-methoxybenzyl, *m,p*-diethoxybenzyl, *o,m,p*-triimethoxybenzyl and *ortho*-, *meta*- *or para*-mono, di- or tri-C_1-C_4alkylphenyl, such as *p*-methyl, *m,m*-diethylphenyl;

or a pharmaceutically acceptable salt thereof.

Another interesting genus is the compounds of formula (Ib):



(Ib)


wherein

R_2' is selected from H; C_1-C_6alkyl; C_4-C_6cycloalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; $(CH_2)_{2-4}OR_{21}$, where R_{21} is H, methyl, ethyl, propyl and *i*-propyl; " and

R_5 is unsubstituted 1*H*-indol-3-yl, benzofuran-3-yl or quinolin-3-yl, or substituted 1*H*-indol-3-yl, such as 5-fluoro-1*H*-indol-3-yl or 5-methoxy-1*H*-indol-3-yl, benzofuran-3-yl or quinolin-3-yl;

or a pharmaceutically acceptable salt thereof.

Another interesting genus of hydroxamate compounds are the compounds of formula (Ic):



(Ic)

wherein

the ring containing Z_1 is aromatic or non-aromatic, which non-aromatic rings are saturated or unsaturated,

Z_1 is O, S or N-R_{20};

R_{18} is H; halo; C_1-C_6alkyl (methyl, ethyl, *t*-butyl); C_3-C_7cycloalkyl; aryl, e.g., unsubstituted phenyl or phenyl substituted by 4-OCH_3 or 4-CF_3; or heteroaryl, such as 2-furanyl, 2-thiophenyl or 2-, 3- or 4-pyridyl;

R_{20} is H; C_1-C_6alkyl; C_1-C_6alkyl-C_3-C_9cycloalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; acyl, e.g., acetyl, propionyl and benzoyl; or sulfonyl, e.g., methanesulfonyl, ethanesulfonyl, benzenesulfonyl and toluenesulfonyl;

A_1 is 1, 2 or 3 substituents which are independently H; C_1-C_6alkyl; -OR_{19}; halo; alkylamino; aminoalkyl; halo; or heteroarylalkyl, e.g., pyridylmethyl;

R_{19} is selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl and -$(CH_2CH=CH(CH_3)(CH_2))_{1\text{-}3}H$;

R_2 is selected from H, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, cycloalkylalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, -$(CH_2)_nC(O)R_6$, amino acyl and -$(CH_2)_nR_7$;

v is 0, 1 or 2;

p is 0-3; and


q is 1-5 and r is 0; or

q is 0 and r is 1-5;

or a pharmaceutically acceptable salt thereof. The other variable substituents are as defined above.

Especially useful compounds of formula (Ic), are those wherein R_2 is H, or -$(CH_2)_pCH_2OH$, wherein p is 1-3, especially those wherein R_1 is H; such as those wherein R_1 is H and X and Y are each H, and wherein q is 1-3 and r is 0 or wherein q is 0 and r is 1-3, especially those wherein Z_1 is N-R_{20}. Among these compounds R_2 is preferably H or -CH_2-CH_2-OH and the sum of q and r is preferably 1.

Another interesting genus of hydroxamate compounds are the compounds of formula (Id):



(Id)

wherein

Z_1 is O, S or N-R_{20};

R_{18} is H; halo; C_1-C_6alkyl (methyl, ethyl, *t*-butyl); C_3-C_7cycloalkyl; aryl, e.g., unsubstituted phenyl or phenyl substituted by 4-OCH_3 or 4-CF_3; or heteroaryl;

R_{20} is H; C_1-C_6alkyl, C_1-C_6alkyl-C_3-C_9cycloalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; acyl, e.g., acetyl, propionyl

and benzoyl; or sulfonyl, e.g., methanesulfonyl, ethanesulfonyl, benzenesulfonyl, toluenesulfonyl);

A_1 is 1, 2 or 3 substituents which are independently H, C_1-C_6alkyl, -OR_{19} or halo;

R_{19} is selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

p is 0-3; and


q is 1-5 and r is 0; or

q is 0 and r is 1-5;

or a pharmaceutically acceptable salt thereof. The other variable substituents are as defined above.

Especially useful compounds of formula (Id), are those wherein R_2 is H or -$(CH_2)_pCH_2OH$, wherein p is 1-3, especially those wherein R_1 is H; such as those wherein R_1 is H and X and Y are each H, and wherein q is 1-3 and r is 0 or wherein q is 0 and r is 1-3. Among these compounds R_2 is preferably H or -CH_2-CH_2-OH and the sum of q and r is preferably 1.

The present invention further relates to compounds of the formula (Ie):



(Ie)

or a pharmaceutically acceptable salt thereof. The variable substituents are as defined above.

Especially useful compounds of formula (Ie), are those wherein R_{18} is H, fluoro, chloro, bromo, a C_1-C_4alkyl group, a substituted C_1-C_4alkyl group, a C_3-C_7cycloalkyl group, unsubstituted phenyl, phenyl substituted in the para position, or a heteroaryl, e.g., pyridyl, ring.

Another group of useful compounds of formula (Ie), are those wherein R_2 is H or -$(CH_2)_pCH_2OH$, wherein p is 1-3, especially those wherein R_1 is H; such as those wherein R_1 is H and X and Y are each H, and wherein q is 1-3 and r is 0 or wherein q is 0 and r is 1-3.


Among these compounds R_2 is preferably H or $-CH_2-CH_2-OH$ and the sum of q and r is preferably 1. Among these compounds p is preferably 1 and R_3 and R_4 are preferably H.

Another group of useful compounds of formula (Ie), are those wherein R_{18} is H, methyl, ethyl, *t*-butyl, trifluoromethyl, cyclohexyl, phenyl, 4-methoxyphenyl, 4-trifluoromethylphenyl, 2-furanyl, 2-thiophenyl, or 2-, 3- or 4-pyridyl wherein the 2-furanyl, 2-thiophenyl and 2-, 3- or 4-pyridyl substituents are unsubstituted or substituted as described above for heteroaryl rings; R_2 is H or $-(CH_2)_pCH_2OH$, wherein p is 1-3; especially those wherein R_1 is H and X and Y are each H, and wherein q is 1-3 and r is 0 or wherein q is 0 and r is 1-3. Among these compounds R_2 is preferably H or $-CH_2-CH_2-OH$ and the sum of q and r is preferably 1.

Those compounds of formula (Ie), wherein R_{20} is H or C_1-C_6alkyl, especially H, are important members of each of the subgenuses of compounds of formula (Ie) described above.

N-hydroxy-3-[4-[[(2-hydroxyethyl)[2-(1*H*-indol-3-yl)ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide, *N*-hydroxy-3-[4-[[[2-(1*H*-indol-3-yl)ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide and *N*-hydroxy-3-[4-[[[2-(2-methyl-1*H*-indol-3-yl)-ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide or a pharmaceutically acceptable salt thereof, are important compounds of formula (Ie).

The present invention further relates to the compounds of the formula (If):



(If)

or a pharmaceutically acceptable salt thereof. The variable substituents are as defined above.


Useful compounds of formula (If), are include those wherein R_2 is H or $-(CH_2)_pCH_2OH$, wherein p is 1-3, especially those wherein R_1 is H; such as those wherein R_1 is H and X and Y are each H, and wherein q is 1-3 and r is 0 or wherein q is 0 and r is 1-3.

Among these compounds R_2 is preferably H or $-CH_2-CH_2-OH$ and the sum of q and r is preferably 1.

N-hydroxy-3-[4-[[[2-(benzofur-3-yl)-ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide or a pharmaceutically acceptable salt thereof, is an important compound of formula (If).


The compounds described above are often used in the form of a pharmaceutically acceptable salt. Pharmaceutically acceptable salts include, when appropriate, pharmaceutically acceptable base addition salts and acid addition salts, e.g., metal salts, such as alkali and alkaline earth metal salts, ammonium salts, organic amine addition salts and amino acid addition salts and sulfonate salts. Acid addition salts include inorganic acid addition salts, such as hydrochloride, sulfate and phosphate; and organic acid addition salts, such as alkyl sulfonate, arylsulfonate, acetate, maleate, fumarate, tartrate, citrate and lactate. Examples of metal salts are alkali metal salts, such as lithium salt, sodium salt and potassium salt; alkaline earth metal salts, such as magnesium salt and calcium salt, aluminum salt and zinc salt. Examples of ammonium salts are ammonium salt and tetramethylammonium salt. Examples of organic amine addition salts are salts with morpholine and piperidine. Examples of amino acid addition salts are salts with glycine, phenylalanine, glutamic acid and lysine. Sulfonate salts include mesylate, tosylate and benzene sulfonic acid salts.

Additional HDAI compounds within the scope of formula (I), and their synthesis, are disclosed in WO 02/22577. Two preferred compounds within the scope of WO 02/22577 are:



(II)

- *N*-hydroxy-3-[4-[(2-hydroxyethyl){2-(1*H*-indol-3-yl)ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide, or a pharmaceutically acceptable salt thereof; and



(III)

- *N*-hydroxy-3-[4-[[[2-(2-methyl-1*H*-indol-3-yl)-ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide, or a pharmaceutically acceptable salt thereof.

An HDAC inhibitor as used for the present invention displays in the assay described above preferably an IC_{50} value between 50 and 2500 nM, more preferably between 250 and 2000 nM, and most preferably between 500 and 1250 nM.

The term "treatment", as used herein, comprises the treatment of patients having melanoma or being in a pre-stage of said cancer which effects the delay of progression of the disease in said patients.

The invention relates to a method of treating a warm-blooded animal having melanoma comprising administering to said animal in need for such a treatment *N*-hydroxy-3-[4-[[[2-(2-methyl-1*H*-indol-3-yl)-ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide or a pharmaceutically acceptable salt thereof, in a quantity which is therapeutically effective against melanoma.

The invention relates to a method for administering to a human subject suffering from melanoma an acid addition salt of *N*-hydroxy-3-[4-[[[2-(2-methyl-1*H*-indol-3-yl)-ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide and preferably the lactic acid addition salt.

The person skilled in the pertinent art is fully enabled to select relevant test models to prove the beneficial effects mentioned herein on melanoma. The pharmacological activity of such a compound may, e.g., be demonstrated by means of the Examples described below, by *in vitro* tests and *in vivo* tests or in suitable clinical studies. Suitable clinical studies are, for example, open label non-randomized, dose escalation studies in patients with melanoma. The efficacy of the treatment is determined in these studies, e.g., by evaluation of the tumor sizes every 4 weeks, with the control achieved on placebo.

The effective dosage of the HDAC inhibitor may vary depending on the particular compound or pharmaceutical composition employed, on the mode of administration, the type

of the melanoma being treated or its severity. The dosage regimen is selected in accordance with a variety of further factors including the renal and hepatic function of the patient. A physician, clinician or veterinarian of ordinary skill can readily determine and prescribe the effective amount of compounds required to prevent, counter or arrest the progress of the condition.

Example 1 **Monolayer Growth Inhibition Assay on Melanoma Cell Lines**

The MTT is a colorimetric assay to determine the cell proliferation rate. The yellow tetrazolium MTT (3-(4, 5-dimethylthiazolyl-2)-2, 5-diphenyltetrazolium bromide) is reduced by metabolically active cells, in part by the action of dehydrogenase enzymes, to generate reducing equivalents, such as NADH and NADPH. The resulting intracellular purple formazan can be solubilized and quantified by spectrophotometric means. The signals produced is directly proportional to the cell numbers. Describing the MTT assay in detail, experiments were done using six-point or nine-point drug titrations in multi-well tissue culture dishes, with outer rows left empty. Cells were suspended in complete media at densities of between 10^3 and 10^4 cell/mL, respectively, and added per well. The appropriate medium (200 µL) was then added. Twenty-four hours later, 10 µL of MTS solution , were added to one plates to determine the activity at the time of compound addition (T_0). This plate was incubated at 37°C for 4 hours and the optical density was measured on a Molecular Devices Thermomax at 490 nm using the Softmax program. The T_0 plate served as a reference for initial activity at the beginning of the experiment.

Compound addition began 24 hours after seeding, the same time as the T_0 determination. Serial dilutions at 4-fold, 2-fold, 1-fold, 0.5-fold, 0.25-fold and 0.125-fold of previously determined IC_{50} values of each compound were made in a 96-deep well plate with the highest concentrations on the edge of the plate. Each of the six dilutions were added in triplicate and complete medium was added to the empty outer rows without cells. The compounds were added to the plates singly or in combination with *N*-hydroxy-3-[4-[[[2-(2-methyl-1*H*-indol-3-yl)-ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide (LBH589). The plates were incubated at 37°C for 72 hours from seeding. The MTS solution was added (as for the T_0 plate) and read 4 hours later. In order to analyze the data, the average value of media alone (background) was subtracted from each experimental well and the triplicate values were averaged for each compound dilution. The following formulas were used to calculate percent growth.

If $X > T_0$, % Growth = 100 x $((X-T_0)/(GC-T_0))$

If $X < T_0$, % Growth = 100 x $(X-T_0)/T_0$)

T_0 = average value of T_0 minus background

GC = average value of untreated cells (in triplicate) minus background

X = average value of compound treated cells (in triplicate) minus background

IC_{50} the concentration of LBH589 required to inhibit cell growth by 50% and LD_{50}s the concentration required to reduce cell number (kill cells) to 50% the original innoculum were determined. The "% Growth" was plotted against compound concentration and used to calculate IC_{50}s and LD_{50}s, employing the user-defined spline function in Microsoft Excel. The attached Table shows the Anti-proliferation and cytotoxic effects of LBH589 in a panel of Melanoma cell lines:


Melanoma Cell Lines	IC_{50} [nM] Measure of Cell Growth Inhibition	LD_{50} (nM) Measure of Cell Kill
A375	32.66	77.83
A2058	17.81	58.96
SK-MEL-28	43.1	129.33
WM-266-4	26.36	73.05
RPMI-7951	27.6	71.63
SK-MEL-5	6.15	54.95
MeWo	34.5	157.64
G-361	8.09	51.44

A panel of Melanoma cell lines were treated with DMSO vehicle control or varying concentrations of LBH589 for 3 days. Cell proliferation was measured on the day of cell plating and on the third day post-treatment. IC_{50} and LD_{50} values were calculated as described above. LBH589 exhibits potent anti-proliferative effect on all 8 melanoma cell lines examined, as demonstrated by the low nanomolar concentrations of IC_{50} values. LBH589 also exhibits potent cytotoxic effect in the cell lines tested with LD_{50} <160 nM

What is Claimed Is:

1. The use of an HDAC inhibitor for the preparation of a medicament for the treatment of melanoma.

2. Use according to Claim 1, wherein the HDAC inhibitor is a compound of the formula (I):



(I)

wherein

R_1 is H; halo; or a straight-chain C_1-C_6alkyl, especially methyl, ethyl or *n*-propyl, which methyl, ethyl and *n*-propyl substituents are unsubstituted or substituted by one or more substituents described below for alkyl substituents;

R_2 is selected from H; C_1-C_{10}alkyl, preferably C_1-C_6alkyl, e.g., methyl, ethyl or $-CH_2CH_2$-OH; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; C_4-C_9heterocycloalkylalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; $-(CH_2)_nC(O)R_6$; $-(CH_2)_nOC(O)R_6$; amino acyl; HON-C(O)-CH=C(R_1)-aryl-alkyl-; and $-(CH_2)_nR_7$;

R_3 and R_4 are the same or different and, independently, H, C_1-C_6alkyl, acyl, or acylamino, or

R_3 and R_4, together with the carbon to which they are bound, represent C=O, C=S or C=NR_8, or

R_2, together with the nitrogen to which it is bound, and R_3, together with the carbon to which it is bound, can form a C_4-C_9heterocycloalkyl, a heteroaryl, a polyheteroaryl, a non-aromatic polyheterocycle, or a mixed aryl and non-aryl polyheterocycle ring;

R_5 is selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; acyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; aromatic polycycles; non-aromatic polycycles; mixed aryl and non-aryl polycycles; polyheteroaryl; non-aromatic polyheterocycles; and mixed aryl and non-aryl polyheterocycles;

n, n_1, n_2 and n_3 are the same or different and independently selected from 0-6, when n_1 is 1-6, each carbon atom can be optionally and independently substituted with R_3 and/or R_4;

X and Y are the same or different and independently selected from H; halo; C_1-C_4alkyl, such as CH_3 and CF_3; NO_2; $C(O)R_1$; OR_9; SR_9; CN; and $NR_{10}R_{11}$;

R_6 is selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl and 2-phenylethenyl; heteroarylalkyl, e.g., pyridylmethyl; OR_{12}; and $NR_{13}R_{14}$;

R_7 is selected from OR_{15}, SR_{15}, $S(O)R_{16}$, SO_2R_{17}, $NR_{13}R_{14}$ and $NR_{12}SO_2R_6$;

R_8 is selected from H; OR_{15}; $NR_{13}R_{14}$; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

R_9 is selected from C_1-C_4alkyl, e.g., CH_3 and CF_3; C(O)-alkyl, e.g., $C(O)CH_3$; and $C(O)CF_3$;

R_{10} and R_{11} are the same or different and independently selected from H, C_1-C_4alkyl and -C(O)-alkyl;

R_{12} is selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; C_4-C_9heterocycloalkylalkyl; aryl; mixed aryl and non-aryl polycycle; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

R_{13} and R_{14} are the same or different and independently selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; amino acyl; or

R_{13} and R_{14}, together with the nitrogen to which they are bound, are C_4-C_9heterocycloalkyl, heteroaryl, polyheteroaryl, non-aromatic polyheterocycle, or mixed aryl and non-aryl polyheterocycle;

R_{15} is selected from H, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl and $(CH_2)_mZR_{12}$;

R_{16} is selected from C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, polyheteroaryl, arylalkyl, heteroarylalkyl and $(CH_2)_mZR_{12}$;


R_{17} is selected from C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, aromatic polycycles, heteroaryl, arylalkyl, heteroarylalkyl, polyheteroaryl and $NR_{13}R_{14}$;

m is an integer selected from 0-6; and

Z is selected from O, NR_{13}, S and S(O),

or a pharmaceutically acceptable salt thereof.

3. Use according to Claim 2, wherein the compound of formula (I) is *N*-hydroxy-3-[4-[[[2-(2-methyl-1*H*-indol-3-yl)-ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide having the formula (III):



(III)


,

or a pharmaceutically acceptable salt thereof.

4. Use according to any one of Claims 1 to 3, wherein the warm-blooded animal is a human.

5. A method of treating melanoma comprising administering a therapeutically effective amount of an HDAC inhibitor to a warm-blooded animal in need thereof.

6. A method according to Claim 5, comprising administering a therapeutically effective amount of a compound of formula (I):



(I)

,

wherein

R_1 is H; halo; or a straight-chain C_1-C_6alkyl, especially methyl, ethyl or *n*-propyl, which methyl, ethyl and *n*-propyl substituents are unsubstituted or substituted by one or more substituents described below for alkyl substituents;

R_2 is selected from H; C_1-C_{10}alkyl, preferably C_1-C_6alkyl, e.g., methyl, ethyl or -CH_2CH_2-OH; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; C_4-C_9heterocycloalkylalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl;

heteroarylalkyl, e.g., pyridylmethyl; $-(CH_2)_nC(O)R_6$; $-(CH_2)_nOC(O)R_6$; amino acyl; HON-C(O)-CH=C(R_1)-aryl-alkyl-; and $-(CH_2)_nR_7$;

R_3 and R_4 are the same or different and, independently, H, C_1-C_6alkyl, acyl, or acylamino, or

R_3 and R_4, together with the carbon to which they are bound, represent C=O, C=S or C=NR_8, or

R_2, together with the nitrogen to which it is bound, and R_3, together with the carbon to which it is bound, can form a C_4-C_9heterocycloalkyl, a heteroaryl, a polyheteroaryl, a non-aromatic polyheterocycle, or a mixed aryl and non-aryl polyheterocycle ring;

R_5 is selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; acyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; aromatic polycycles; non-aromatic polycycles; mixed aryl and non-aryl polycycles; polyheteroaryl; non-aromatic polyheterocycles; and mixed aryl and non-aryl polyheterocycles;

n, n_1, n_2 and n_3 are the same or different and independently selected from 0-6, when n_1 is 1-6, each carbon atom can be optionally and independently substituted with R_3 and/or R_4;

X and Y are the same or different and independently selected from H; halo; C_1-C_4alkyl, such as CH_3 and CF_3; NO_2; $C(O)R_1$; OR_9; SR_9; CN; and $NR_{10}R_{11}$;

R_6 is selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl and 2-phenylethenyl; heteroarylalkyl, e.g., pyridylmethyl; OR_{12}; and $NR_{13}R_{14}$;

R_7 is selected from OR_{15}, SR_{15}, $S(O)R_{16}$, SO_2R_{17}, $NR_{13}R_{14}$ and $NR_{12}SO_2R_6$;

R_8 is selected from H; OR_{15}; $NR_{13}R_{14}$; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

R_9 is selected from C_1-C_4alkyl, e.g., CH_3 and CF_3; C(O)-alkyl, e.g., $C(O)CH_3$; and $C(O)CF_3$;

R_{10} and R_{11} are the same or different and independently selected from H, C_1-C_4alkyl and -C(O)-alkyl;

R_{12} is selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; C_4-C_9heterocycloalkylalkyl; aryl; mixed aryl and non-aryl polycycle; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

R_{13} and R_{14} are the same or different and independently selected from H; C_1-C_6alkyl; C_4-C_9cycloalkyl; C_4-C_9heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; amino acyl; or

R_{13} and R_{14}, together with the nitrogen to which they are bound, are C_4-C_9heterocycloalkyl, heteroaryl, polyheteroaryl, non-aromatic polyheterocycle, or mixed aryl and non-aryl polyheterocycle;

R_{15} is selected from H, C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl and $(CH_2)_mZR_{12}$;

R_{16} is selected from C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, heteroaryl, polyheteroaryl, arylalkyl, heteroarylalkyl and $(CH_2)_mZR_{12}$;


R_{17} is selected from C_1-C_6alkyl, C_4-C_9cycloalkyl, C_4-C_9heterocycloalkyl, aryl, aromatic polycycles, heteroaryl, arylalkyl, heteroarylalkyl, polyheteroaryl and $NR_{13}R_{14}$;

m is an integer selected from 0-6; and

Z is selected from O; NR_{13}; S; and S(O),

or a pharmaceutically acceptable salt thereof to a warm-blooded animal in need thereof.

7. The method according to Claim 5, wherein the compound of formula (I) is *N*-hydroxy-3-[4-[[[2-(2-methyl-1*H*-indol-3-yl)-ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide having the formula (III):



(III)

or a pharmaceutically acceptable salt thereof.

8. The method according to Claim 5, wherein the warm-blooded animal is a human.